

STRAIGHT PEPTIDES

TRUE IN HEALTH. UNLOCKING HUMAN POTENTIAL

PRODUCT GUIDE

Cagrilintide 5mg — Mixing & Injection Instructions

Reconstitution and Subcutaneous Administration Protocol
For Straight Peptides Clients

Confidential | Internal Use Only | 2026

Stage 1: Reconstitution

1

Mixing Your Cagrilintide Vial

Time: 5 minutes | Location: Clean, flat surface

Reconstitution prepares the lyophilized Cagrilintide powder for subcutaneous injection. Follow each step precisely to preserve peptide integrity.

Products Needed

Equipment

Bacteriostatic Water	BAC water vial (supplied)
100-unit insulin syringe	For drawing and transferring BAC water
30-35-unit insulin syringe	For injection (used in Stage 2)
Alcohol swabs	For sterilizing vial stoppers and injection site
5mg Cagrilintide vial	Lyophilized peptide, vacuum-sealed

Prepare the Vials

Process

1. Remove the protective seals from both the BAC water and Cagrilintide vials.
2. Clean the rubber stoppers on both vials using an alcohol swab. Allow to air dry.

Mixing Instructions

Protocol

1. Using the 100-unit insulin syringe, insert the needle into the center of the BAC water vial.
2. Pull back to 100 units (1mL) of BAC water.
3. Insert the needle into the Cagrilintide vial. Allow the vacuum to pull the liquid in slowly.

Critical Warning

Do not force the liquid in. The vial is vacuum-sealed and draws the liquid in on its own. Forcing the plunger compromises the vacuum and may cause inaccurate dosing.

4. Gently roll the vial between your fingers. Do not shake. Allow the peptide to fully dissolve for 30–60 seconds.
5. Place the vial in the refrigerator. After 1–2 hours refrigerated, the solution is ready for use.

Storage Requirement

Reconstituted Cagrilintide must remain refrigerated at all times. Do not freeze. Do not expose to direct sunlight or heat.

Stage 2: Injection

2

Subcutaneous Administration

Time: 2 minutes | Location: Stomach area (preferred)

Administer your dose subcutaneously using the 30–35-unit insulin syringe. Rotate injection sites to prevent tissue irritation.

Products Needed

Equipment

Cagrilintide vial	Reconstituted and refrigerated
Alcohol swab	For sterilizing vial stopper and injection site
30–35-unit insulin syringe	For drawing and injecting the dose

Draw Your Dose

Protocol

1. Wipe the Cagrilintide vial stopper and your injection site with an alcohol swab.
2. Insert the insulin syringe into the center of the vial.
3. Pull back to your prescribed unit mark using the conversion table below.

Inject

Process

1. Insert the needle fully into the injection site at a 90-degree angle. Stomach area is preferred.
2. Inject slowly and steadily.
3. Lightly rub the injection site afterward in a circular motion to aid absorption.

Stage 3: Dosing Protocol

3

Titration Schedule, Conversion, and Vial Duration

Time: Ongoing | Once weekly

Cagrilintide is administered once weekly. Slow titration is essential due to the compound's long half-life, which causes it to build up in the body over time.

Recommended Titration Schedule**Protocol**

Week 1-4	0.5mg weekly (10 units)
Week 5-8	0.75mg weekly (15 units)
Week 9+	1mg weekly (20 units)

Increase slowly depending on tolerance. Do not skip titration phases.

Protocol Note

Starting at 0.5mg reduces nausea, stomach discomfort, and fatigue. Cagrilintide builds up in the body due to its long half-life, so slow titration produces better tolerance and more consistent appetite control. Rushing the dose escalation increases side effects without improving outcomes.

Dose Conversion

100 units total — each 1 unit = 50mcg (0.05mg)

10 units	0.5mg
15 units	0.75mg
20 units	1mg
25 units	1.25mg
30 units	1.5mg

Vial Duration (5mg vial)

0.5mg weekly (10 units)	10 weeks
0.75mg weekly (15 units)	≈6-7 weeks
1mg weekly (20 units)	5 weeks
1.25mg weekly (25 units)	4 weeks
1.5mg weekly (30 units)	≈3 weeks

Important Reminders**Protocol**

Refrigeration	Always keep the reconstituted vial refrigerated
Sterile syringes	Use a fresh syringe for each injection

Injection sites	Rotate sites to prevent tissue irritation
Frequency	Inject once weekly, same day each week

Stage 4: What to Expect

4

Response Timeline and Outcomes
Timeline: Weeks 1-12+

Cagrilintide is an amylin analogue used in metabolic research. It mimics the satiety-signaling hormone amylin, reducing appetite and slowing gastric emptying through central and peripheral pathways.

Common Observations

Weeks 1-12+

Appetite	Reduced appetite and decreased food cravings
Satiety	Improved feeling of fullness after meals
Gastric emptying	Slower gastric emptying supports portion control
Weight management	Supports structured weight management protocols

Important Notice

This guide is intended for research and educational purposes only. Not intended to diagnose, treat, cure, or prevent any disease.

Questions or Concerns

Contact Straight Peptides directly for dosage guidance, product questions, or support.

Phone: 915-637-6443
Email: straightpeptides@gmail.com