

STRAIGHT PEPTIDES

TRUE IN HEALTH. UNLOCKING HUMAN POTENTIAL

PRODUCT GUIDE

Glow 70mg — Mixing & Injection Instructions

Reconstitution and Subcutaneous Administration Protocol
For Straight Peptides Clients

Confidential | Internal Use Only | 2026

GHK-Cu 50mg | BPC-157 10mg | TB-500 10mg

Stage 1: Reconstitution

1

Mixing Your Glow Vial

Time: 5 minutes | Location: Clean, flat surface

Glow is a pre-blended vial containing GHK-Cu (50mg), BPC-157 (10mg), and TB-500 (10mg) for a total of 70mg active peptide. Reconstitution prepares the blend for subcutaneous injection.

Products Needed

Equipment

Bacteriostatic Water	BAC water vial (supplied)
100-unit insulin syringe	For drawing and transferring BAC water
30-35-unit insulin syringe	For injection (used in Stage 2)
Alcohol swabs	For sterilizing vial stoppers and injection site
Glow 70mg blend vial	Lyophilized peptide blend, vacuum-sealed

Prepare the Vials

Process

1. Remove the protective seals from both the BAC water and Glow vials.
2. Clean the rubber stoppers on both vials using an alcohol swab. Allow to air dry.

Mixing Instructions

Protocol

1. Using the 100-unit insulin syringe, insert the needle into the center of the BAC water vial.
2. Pull back to 100 units (1mL) of BAC water.
3. Insert the needle into the Glow vial. Allow the vacuum to pull the liquid in slowly.

Critical Warning

Do not push the syringe forcefully. The vial is vacuum-sealed and draws the liquid in on its own. GHK-Cu is a delicate copper peptide — aggressive handling degrades the compound.

4. Gently roll the vial between your fingers. Do not shake. Allow the blend to fully dissolve for 30–60 seconds.
5. Place the vial in the refrigerator. After 1–2 hours refrigerated, the solution is ready for use.

Storage Requirement

Reconstituted Glow must remain refrigerated at all times. GHK-Cu is light-sensitive. Store in a dark location. Do not freeze.

Dose Conversion

70mg vial mixed with 1mL BAC water = 0.7mg total peptide per unit

Per-Unit Breakdown

GHK-Cu	500mcg per unit (50mg / 100 units)
BPC-157	100mcg per unit (10mg / 100 units)
TB-500	100mcg per unit (10mg / 100 units)

Common Research Doses

2 units	GHK-Cu 1mg + BPC-157 200mcg + TB-500 200mcg
4 units	GHK-Cu 2mg + BPC-157 400mcg + TB-500 400mcg
5 units	GHK-Cu 2.5mg + BPC-157 500mcg + TB-500 500mcg

Vial Duration

2 units daily	50 days
4 units daily	25 days
5 units daily	20 days

Stage 2: Injection

2

Subcutaneous Administration

Time: 2 minutes | Location: Abdomen, thigh, or target area

Administer your dose subcutaneously using the 30–35-unit insulin syringe. For localized recovery benefits, inject near the target area. For systemic skin and tissue benefits, rotate between abdomen and thigh.

Products Needed

Equipment

Glow vial	Reconstituted and refrigerated
Alcohol swab	For sterilizing vial stopper and injection site
30–35-unit insulin syringe	For drawing and injecting the dose

Draw Your Dose

Protocol

1. Wipe the Glow vial stopper and your injection site with an alcohol swab.
2. Insert the needle into the center of the vial and pull back to your prescribed unit mark.

Inject

Process

1. Insert the needle fully into the injection site at a 90-degree angle.
2. Inject slowly and steadily.
3. After injecting, lightly rub the area in a circular motion to aid absorption.

Stage 3: Dosing Protocol

3

Frequency, Timing, and Injection Strategy

Time: Ongoing | Daily injection

Glow is administered once daily via subcutaneous injection. Each component targets a different recovery and rejuvenation pathway, and daily consistency maximizes the compounding effect.

Blend Components

Objective

GHK-Cu (50mg)	Copper peptide — stimulates collagen synthesis, skin remodeling, wound healing, and anti-inflammatory signaling
BPC-157 (10mg)	Body Protection Compound — accelerates tissue repair, gut healing, tendon and ligament recovery
TB-500 (10mg)	Thymosin Beta-4 fragment — promotes cell migration, reduces inflammation, supports tissue regeneration

Important Reminders

Protocol

Refrigeration	Always keep reconstituted Glow refrigerated and away from light
Sterilization	Always wipe the vial stopper and injection site before each use
Frequency	Once daily, subcutaneous
Injection site	Rotate for systemic benefits; inject near target area for localized recovery

Protocol Note

For skin quality and anti-aging benefits, rotate standard injection sites (abdomen, thigh). For targeted recovery — a healing joint, surgical site, or injured area — inject subcutaneously as close to the target tissue as possible. BPC-157 and TB-500 both exhibit local tissue affinity.

Stage 4: What to Expect

4

Response Timeline and Outcomes

Timeline: Weeks 1-12+

The Glow blend combines three peptides that work through complementary pathways: collagen synthesis, tissue repair, and inflammatory modulation. Results are progressive and compound with daily use.

Early Changes

Weeks 1-4

Skin texture	Smoother, more hydrated skin — often noticeable within 2-3 weeks
Recovery	Faster healing of minor injuries, soreness, and inflammation
Hair and nails	Improved growth rate and quality
Inflammation	Reduced joint stiffness and general inflammatory markers

Progressive Results

Weeks 4-12+

Collagen density	Visibly firmer, more elastic skin
Scar remodeling	Improved appearance of existing scars and marks
Gut health	BPC-157 supports gut lining integrity and digestive function

Tissue repair	Enhanced tendon, ligament, and connective tissue recovery
Overall vitality	Compounding anti-aging and regenerative effects

Important Notice

This guide is intended for research and educational purposes only. Not intended to diagnose, treat, cure, or prevent any disease.

Questions or Concerns

Contact Straight Peptides directly for dosage guidance, product questions, or support.

Phone: 915-637-6443

Email: straightpeptides@gmail.com