

STRAIGHT PEPTIDES

TRUE IN HEALTH. UNLOCKING HUMAN POTENTIAL

PRODUCT GUIDE

IGF-1 LR3 1mg — Mixing & Injection Instructions

Reconstitution and Subcutaneous Administration Protocol
For Straight Peptides Clients

Confidential | Internal Use Only | 2026

Stage 1: Reconstitution

1

Mixing Your IGF-1 LR3 Vial

Time: 5 minutes | Location: Clean, flat surface

Reconstitution prepares the lyophilized IGF-1 LR3 powder for subcutaneous injection. IGF-1 LR3 is a modified form of insulin-like growth factor with an extended half-life, requiring careful handling.

Products Needed

Equipment

Bacteriostatic Water	BAC water vial (supplied)
100-unit insulin syringe	For drawing and transferring BAC water
30-35-unit insulin syringe	For injection (used in Stage 2)
Alcohol swabs	For sterilizing vial stoppers and injection site
1mg IGF-1 LR3 vial	Lyophilized peptide, vacuum-sealed

Prepare the Vials

Process

1. Remove the protective seals from both the BAC water and IGF-1 LR3 vials.
2. Clean the rubber stoppers on both vials using an alcohol swab. Allow to air dry.

Mixing Instructions

Protocol

1. Using the 100-unit insulin syringe, insert the needle into the center of the BAC water vial.
2. Pull back to 100 units (1mL) of BAC water.

3. Insert the needle into the IGF-1 LR3 vial. Allow the vacuum to pull the liquid in slowly.

Critical Warning

Do not push the syringe forcefully. IGF-1 LR3 is a fragile peptide. Inject BAC water slowly along the inside wall of the vial. Do not spray directly onto the powder.

4. Gently roll the vial between your fingers. Do not shake. Allow the peptide to fully dissolve for 30–60 seconds.
5. Place the vial in the refrigerator. After 1–2 hours refrigerated, the solution is ready for use.

Storage Requirement

Reconstituted IGF-1 LR3 must remain refrigerated at all times. Use within 28 days of reconstitution. Do not freeze. Do not expose to direct sunlight or heat.

Dose Conversion

1mg vial mixed with 1mL BAC water = 10mcg per unit

2 units	20mcg
3 units	30mcg
4 units	40mcg
5 units	50mcg
6 units	60mcg
8 units	80mcg
10 units	100mcg

Vial Duration

20mcg daily (2 units)	50 days
40mcg daily (4 units)	25 days
50mcg daily (5 units)	20 days
80mcg daily (8 units)	≈12 days

Stage 2: Injection

2

Subcutaneous Administration

Time: 2 minutes | Location: Abdomen, thigh, or target muscle

Administer your dose subcutaneously using the 30–35-unit insulin syringe. For muscle-specific benefits, inject near the target muscle group post-workout.

Products Needed

Equipment

IGF-1 LR3 vial	Reconstituted and refrigerated
Alcohol swab	For sterilizing vial stopper and injection site
30-35-unit insulin syringe	For drawing and injecting the dose

Draw Your Dose

Protocol

1. Wipe the IGF-1 LR3 vial stopper and your injection site with an alcohol swab.
2. Insert the needle into the center of the vial and pull back to your prescribed unit mark using the conversion table above.

Dosage Reference

Common starting dose is 20–40mcg daily (2–4 units). Do not exceed recommended research doses. Start low and assess tolerance before increasing.

Inject

Process

1. Insert the needle fully into the injection site at a 90-degree angle.
2. Inject slowly and steadily.
3. After injecting, lightly rub the area in a circular motion to aid absorption.

Stage 3: Dosing Protocol

3

Frequency, Timing, and Cycling

Time: 4 weeks on / 4 weeks off

IGF-1 LR3 is administered once daily. Due to receptor desensitization, cycling is recommended to maintain efficacy over time.

Recommended Protocol

Protocol

Starting dose	20–40mcg daily (2–4 units)
Advanced dose	50–80mcg daily (5–8 units)
Cycle pattern	4 weeks on, 4 weeks off
Frequency	Once daily

Injection Timing

Objective

Timing determines whether IGF-1 LR3 supports systemic growth or localized muscle recovery.

Post-workout	Inject within 30 minutes after training for localized muscle benefit
Morning (non-training days)	Inject fasted in the morning for systemic benefit

Protocol Note

IGF-1 LR3 has an extended half-life of approximately 20–30 hours, meaning it remains active throughout the day. Cycling 4 weeks on, 4 weeks off prevents receptor downregulation and maintains sensitivity. Do not run continuously without off periods.

Important Reminders

Protocol

Refrigeration	Always keep reconstituted IGF-1 LR3 refrigerated
Shelf life	Use within 28 days of reconstitution
Sterilization	Always wipe the vial stopper and injection site before each use
Cycling	4 weeks on, 4 weeks off to maintain receptor sensitivity
Blood sugar	Monitor for signs of hypoglycemia — IGF-1 can lower blood glucose

Safety Note

IGF-1 LR3 can lower blood sugar. If you experience dizziness, shakiness, or lightheadedness after injection, consume a small carbohydrate source. Clients using this compound alongside insulin or GLP-1 medications should consult their provider about dose timing.

Stage 4: What to Expect

4

Response Timeline and Outcomes

Timeline: Weeks 1–4 per cycle

IGF-1 LR3 is a modified insulin-like growth factor with an extended half-life. It promotes muscle hyperplasia (new muscle cell formation), enhances recovery, and supports fat metabolism through direct tissue-level signaling.

Early Changes

Weeks 1–2

Muscle fullness	Increased pump and muscle density during training
Recovery	Faster recovery between training sessions
Soreness	Reduced delayed-onset muscle soreness (DOMS)
Energy	Improved training capacity and endurance

Progressive Results

Weeks 2-4

Muscle growth	Hyperplasia — new muscle cell formation, not just hypertrophy
Fat metabolism	Enhanced lipolysis and improved body composition
Connective tissue	Improved tendon and ligament integrity
Overall composition	Leaner, denser physique with improved vascularity

Important Notice

This guide is intended for research and educational purposes only. Not intended to diagnose, treat, cure, or prevent any disease.

Questions or Concerns

Contact Straight Peptides directly for dosage guidance, product questions, or support.

Phone: 915-637-6443

Email: straightpeptides@gmail.com