

STRAIGHT PEPTIDES

TRUE IN HEALTH. UNLOCKING HUMAN POTENTIAL

PRODUCT GUIDE

NAD+ 1000mg — Mixing & Injection Instructions

Reconstitution and Subcutaneous Administration Protocol
For Straight Peptides Clients

Confidential | Internal Use Only | 2026

Stage 1: Reconstitution

1

Mixing Your NAD+ Vial

Time: 5 minutes | Location: Clean, flat surface

Reconstitution prepares the lyophilized NAD+ powder for subcutaneous injection. NAD+ dissolves readily but requires gentle handling to preserve molecular integrity.

Products Needed

Equipment

Bacteriostatic Water	BAC water vial (supplied)
100-unit insulin syringe	For drawing and transferring BAC water
30-35-unit insulin syringe	For injection (used in Stage 2)
Alcohol swabs	For sterilizing vial stoppers and injection site
1000mg NAD+ vial	Lyophilized powder, vacuum-sealed

Prepare the Vials

Process

1. Remove the protective seals from both the BAC water and NAD+ vials.
2. Clean the rubber stoppers on both vials using an alcohol swab. Allow to air dry.

Mixing Instructions

Protocol

1. Using the 100-unit insulin syringe, insert the needle into the center of the BAC water vial.
2. Push air into the vial, then pull back to 100 units (1mL) of BAC water.
3. Insert the needle into the NAD+ vial.

- Repeat for a second draw of 100 units, for a total of 200 units (2mL) of BAC water.

Critical Warning

Do not push the syringe forcefully. Allow the vacuum to assist. Inject the BAC water slowly along the inside wall of the vial to minimize foaming.

- Gently roll the vial between your fingers. Do not shake. Allow the powder to fully dissolve for 30–60 seconds.
- Place the vial in the refrigerator. After 1–2 hours refrigerated, the solution is ready for injection.

Storage Requirement

Reconstituted NAD+ must remain refrigerated at all times. NAD+ is sensitive to heat and light. Store in a dark, cool location. Do not freeze.

Dose Conversion

1000mg vial mixed with 2mL BAC water = 5mg per unit

10 units	50mg
20 units	100mg
30 units	150mg
40 units	200mg
50 units	250mg

Vial Duration

100mg daily (20 units)	10 days
150mg daily (30 units)	≈7 days
200mg daily (40 units)	5 days
250mg daily (50 units)	4 days

Stage 2: Injection

2

Subcutaneous Administration

Time: 2 minutes | Location: Abdomen or thigh

Administer your dose subcutaneously using the 30–35-unit insulin syringe. NAD+ injections commonly produce a mild stinging or warming sensation at the injection site. This is normal and subsides within a few minutes.

Products Needed

Equipment

NAD+ vial	Reconstituted and refrigerated
------------------	--------------------------------

Alcohol swab	For sterilizing vial stopper and injection site
30-35-unit insulin syringe	For drawing and injecting the dose

Draw Your Dose

Protocol

1. Wipe the NAD+ vial stopper and your injection site with an alcohol swab.
2. Insert the needle into the center of the vial and pull back to your prescribed unit mark.

Safety Note
NAD+ SubQ injections commonly cause a temporary stinging, warming, or flushing sensation at the injection site. This is a normal reaction to the compound and resolves within 2-5 minutes. Injecting slowly reduces discomfort.

Inject

Process

1. Insert the needle fully into the injection site at a 90-degree angle. Abdomen or outer thigh preferred.
2. Inject slowly and steadily. A slower injection reduces the stinging sensation.
3. After injecting, lightly rub the area in a circular motion to aid absorption.

Stage 3: Dosing Protocol

3

Frequency, Timing, and Loading Phase
Time: Ongoing | Daily injection

NAD+ is administered once daily via subcutaneous injection. Many protocols include a loading phase at a higher dose for the first 1-2 weeks before transitioning to a maintenance dose.

Recommended Protocol

Protocol

Loading phase (Week 1-2)	200mg daily (40 units)
Maintenance (Week 3+)	100-150mg daily (20-30 units)

Adjust based on individual response and tolerance. Some protocols call for 5 days on, 2 days off.

Important Reminders

Protocol

Refrigeration	Always keep reconstituted NAD+ refrigerated and away
----------------------	--

	from light
Sterilization	Always wipe the vial stopper and injection site before each use
Frequency	Once daily, subcutaneous
Timing	Morning preferred for energy benefits; evening acceptable
Sensation	Mild stinging at injection site is normal and temporary

Protocol Note

NAD+ levels decline naturally with age. Subcutaneous administration bypasses the GI tract and delivers the molecule directly into circulation, avoiding the degradation that occurs with oral supplementation. Consistency is more important than dose — daily administration maintains elevated cellular NAD+ levels.

Stage 4: What to Expect

4

Response Timeline and Outcomes

Timeline: Weeks 1-12+

NAD+ (nicotinamide adenine dinucleotide) is a coenzyme essential to cellular energy production, DNA repair, and metabolic regulation. Supplementation supports mitochondrial function and activates sirtuins, the enzymes linked to longevity and cellular resilience.

Early Changes

Weeks 1-4

Energy	Improved sustained energy throughout the day
Mental clarity	Enhanced focus and reduced brain fog
Sleep quality	Deeper, more restorative sleep
Recovery	Faster recovery from exercise and daily activity

Progressive Results

Weeks 4-12+

Cellular repair	Enhanced DNA repair and mitochondrial function
Anti-aging markers	Improved skin quality and overall vitality
Metabolic function	More efficient energy metabolism
Cognitive performance	Sustained improvements in focus and memory
Exercise capacity	Improved endurance and workout performance

Important Notice

This guide is intended for research and educational purposes only. Not intended to diagnose, treat, cure, or prevent any disease.

Questions or Concerns

Contact Straight Peptides directly for dosage guidance, product questions, or support.

Phone: 915-637-6443

Email: straightpeptides@gmail.com