

STRAIGHT PEPTIDES

TRUE IN HEALTH. UNLOCKING HUMAN POTENTIAL

PRODUCT GUIDE

Tirzepatide 20mg — Mixing & Injection Instructions

Reconstitution and Subcutaneous Administration Protocol
For Straight Peptides Clients

Confidential | Internal Use Only | 2026

Stage 1: Reconstitution

1

Mixing Your Tirzepatide Vial

Time: 10 minutes | Location: Clean, flat surface

The 20mg vial requires two separate draws of bacteriostatic water. This double-draw process ensures proper dilution for accurate dosing across the full vial.

Products Needed

Equipment

Bacteriostatic Water	BAC water vial (supplied)
100-unit insulin syringe	For drawing and transferring BAC water
35-unit insulin syringe	For injection (used in Stage 2)
Alcohol swabs	For sterilizing vial stoppers and injection site
20mg Tirzepatide vial	Lyophilized peptide, vacuum-sealed

Prepare the Vials

Process

1. Remove the protective seals from both the BAC water and Tirzepatide vials.
2. Clean the rubber stoppers on both vials using an alcohol swab. Allow to air dry.

Mixing Instructions

Protocol

1. Using the 100-unit insulin syringe, insert the needle into the center of the BAC water vial.
2. Push air into the vial, then pull back to approximately 105 units. This extra amount compensates for air bubbles.
3. Insert the needle into the 20mg Tirzepatide vial.

Critical Warning

Do not push the syringe. The vial is vacuum-sealed and pulls the liquid in on its own. Forcing the plunger compromises the vacuum and may cause inaccurate dosing.

4. Repeat steps 1–3 to add a second 105 units of BAC water to the same 20mg Tirzepatide vial, for a total of 210 units of BAC water.

Protocol Note

The 20mg vial requires two full draws of 105 units each (210 units total). Do not attempt to draw the full volume in a single syringe. Each draw uses a fresh insertion into the BAC water vial.

5. Once mixed, gently roll the vial between your fingers. Do not shake.
6. Slowly rotate the vial for approximately 1 minute until the powder fully dissolves.
7. Place the vial in the refrigerator. After 2 hours refrigerated, the solution is ready for injection.

Storage Requirement

Reconstituted Tirzepatide must remain refrigerated at all times. Do not freeze. Do not expose to direct sunlight or heat.

Stage 2: Injection

2

Subcutaneous Administration

Time: 2 minutes | Location: Abdomen, thigh, or upper arm

Administer your dose subcutaneously using the 35-unit insulin syringe. Rotate injection sites weekly to prevent tissue irritation.

Products Needed

Equipment

20mg Tirzepatide vial	Reconstituted and refrigerated
Alcohol swab	For sterilizing vial stopper and injection site
35-unit insulin syringe	For drawing and injecting the dose

Draw Your Dose

Protocol

1. Wipe the Tirzepatide vial stopper and your injection site with an alcohol swab.
2. Insert the needle into the center of the vial and pull back to 25 units.

Dosage Reference

25 units = 2.5mg Tirzepatide. This is the standard starting dose. Do not adjust dosage without guidance from your provider.

Inject

Process

1. Insert the needle fully into the injection site at a 90-degree angle.
2. Slowly push the plunger to inject the medication. Insulin needles are very small and typically painless.
3. After injecting, gently rub the area in a circular motion. This aids absorption and prevents liquid from pooling under the skin.

Stage 3: Daily Protocol

3

Hydration, Nutrition, and Movement

Time: Ongoing | Begins Week 1

Tirzepatide delivers results progressively. The medication works best as part of a structured protocol that includes consistent movement, hydration, and fiber intake.

Important Reminders

Protocol

Refrigeration	Always keep the reconstituted Tirzepatide vial refrigerated
Sterilization	Always wipe the vial stopper and injection site before each injection
Onset timeline	Most people do not feel effects until Week 2 or Week 3
Early changes	Drop in inflammation, loss of water weight
Progressive effects	Feel full faster, stay full longer, eat smaller portions naturally

Hydration and Electrolytes

Process

1. Incorporate daily walking. Start with 10–15 minutes and gradually increase.
2. Drink significantly more water than usual. Aim for nearly double your pre-treatment intake. GLP-1 medications naturally cause dehydration because appetite and fluid intake decrease.
3. Use electrolytes daily. Electrolytes are essential for maintaining energy and preventing fatigue.

Nutrition Focus

Objective

Preserve muscle mass and support digestive function throughout treatment.

Fiber

Eat fruits and vegetables daily. Tirzepatide slows gastric emptying, meaning food stays in the stomach longer. Without adequate fiber, constipation occurs.

Protein

Aim for 0.5g of protein per pound of body weight. Protein preserves muscle mass, maintains energy levels, and supports recovery.

Protocol Note
Fiber and protein are non-negotiable. Clients who skip fiber consistently report constipation and digestive discomfort. Clients who under-eat protein lose muscle mass alongside fat, which undermines long-term results.

Stage 4: What to Expect

4

Physical, Mental, and Long-Term Outcomes
Timeline: Weeks 2-12+

Tirzepatide produces measurable changes across metabolic, physical, and psychological dimensions. These improvements compound over time with consistent protocol adherence.

Physical Improvements

Metabolic Health

Blood sugar control	More stable glucose levels throughout the day
Lipid panel	Decreased LDL cholesterol and triglycerides
Blood pressure	Reduced cardiovascular strain
Inflammation markers	Overall improvement in metabolic health

Energy and Vitality

Sustained energy	No more afternoon crashes
Sleep quality	Improved rest and recovery
Physical performance	Easier movement and exercise
Joint pain	Less stress on the body as weight decreases

Mental and Emotional Benefits

Relationship with Food

Food noise	The constant mental chatter about food quiets down
Appetite regulation	Natural satisfaction with appropriate portions
Mindful eating	Greater awareness of hunger and fullness cues

Binge cycle	Reduced emotional eating and food obsession
--------------------	---

Confidence and Well-Being

Self-image	Real, sustainable progress you can see and feel
Social confidence	Comfort in your own skin in any setting
Food anxiety	Decisions without guilt or overthinking
Sense of control	Taking charge of your health trajectory

Long-Term Success

This protocol is not only about losing weight. It is about transforming your relationship with food, reclaiming your energy, and optimizing your health from the inside out. When combined with consistent daily movement, proper hydration, and nutritious food choices, Tirzepatide becomes a powerful tool for sustainable, life-changing results.

Questions or Concerns

Contact Straight Peptides directly for dosage guidance, product questions, or support.

Phone: 915-637-6443

Email: straightpeptides@gmail.com