

STRAIGHT PEPTIDES

TRUE IN HEALTH. UNLOCKING HUMAN POTENTIAL

PRODUCT GUIDE

Tirzepatide 30mg — Mixing & Injection Instructions

Reconstitution and Subcutaneous Administration Protocol
For Straight Peptides Clients

Confidential | Internal Use Only | 2026

Stage 1: Reconstitution

1

Mixing Your Tirzepatide Vial

Time: 15 minutes | Location: Clean, flat surface

The 30mg vial requires three separate draws of bacteriostatic water. This triple-draw process ensures proper dilution for accurate dosing across the full vial.

Products Needed

Equipment

Bacteriostatic Water	BAC water vial (supplied)
100-unit insulin syringe	For drawing and transferring BAC water
35-unit insulin syringe	For injection (used in Stage 2)
Alcohol swabs	For sterilizing vial stoppers and injection site
30mg Tirzepatide vial	Lyophilized peptide, vacuum-sealed

Prepare the Vials

Process

1. Remove the protective seals from both the BAC water and Tirzepatide vials.
2. Clean the rubber stoppers on both vials using an alcohol swab. Allow to air dry.

Mixing Instructions

Protocol

1. Using the 100-unit insulin syringe, insert the needle into the center of the BAC water vial.
2. Push air into the vial, then pull back to approximately 105 units. This extra amount compensates for air bubbles.
3. Insert the needle into the 30mg Tirzepatide vial.

Critical Warning

Do not push the syringe. The vial is vacuum-sealed and pulls the liquid in on its own. Forcing the plunger compromises the vacuum and may cause inaccurate dosing.

4. Repeat steps 1–3 two more times, for a total of three draws (approximately 315 units of BAC water).

Protocol Note

The 30mg vial requires three full draws of 105 units each (315 units total). Do not attempt to shortcut this process. Each draw uses a fresh insertion into the BAC water vial. The final concentration is approximately 0.1mg per unit, matching the 10mg and 20mg vials.

5. Once mixed, gently roll the vial between your fingers. Do not shake.
6. Slowly rotate the vial for approximately 1 minute until the powder fully dissolves.
7. Place the vial in the refrigerator. After 2 hours refrigerated, the solution is ready for injection.

Storage Requirement

Reconstituted Tirzepatide must remain refrigerated at all times. Do not freeze. Do not expose to direct sunlight or heat.

Stage 2: Injection

2

Subcutaneous Administration

Time: 2 minutes | Location: Abdomen, thigh, or upper arm

Administer your dose subcutaneously using the 35-unit insulin syringe. Rotate injection sites weekly to prevent tissue irritation.

Products Needed

Equipment

30mg Tirzepatide vial	Reconstituted and refrigerated
Alcohol swab	For sterilizing vial stopper and injection site
35-unit insulin syringe	For drawing and injecting the dose

Draw Your Dose

Protocol

1. Wipe the Tirzepatide vial stopper and your injection site with an alcohol swab.
2. Insert the needle into the center of the vial and pull back to 25 units.

Dosage Reference

25 units = 2.5mg Tirzepatide. This is the standard starting dose. Do not adjust

dosage without guidance from your provider.

Inject

Process

1. Insert the needle fully into the injection site at a 90-degree angle.
2. Slowly push the plunger to inject the medication. Insulin needles are very small and typically painless.
3. After injecting, gently rub the area in a circular motion. This aids absorption and prevents liquid from pooling under the skin.

Stage 3: Ongoing Guidance

3

Reminders, Hydration, and Nutrition

Time: Ongoing | Begins Week 1

Tirzepatide delivers results progressively. The medication works best as part of a structured protocol that includes consistent movement, hydration, and fiber intake.

Important Reminders

Protocol

Refrigeration	Always keep the reconstituted Tirzepatide vial refrigerated
Sterilization	Always wipe the vial stopper and injection site before each injection
Onset timeline	Most people do not feel effects until Week 2 or Week 3
Early changes	Drop in inflammation, loss of water weight
Progressive effects	Feel full faster, stay full longer, eat smaller portions naturally

Lifestyle Integration

What Not to Do

Do not rely on Tirzepatide alone. The medication works best as part of a structured weight-loss protocol that includes movement, hydration, and fiber intake.

Process

1. Incorporate daily walking. Consistent low-impact movement accelerates metabolic adaptation.
2. Drink significantly more water than usual. Aim for nearly double your pre-treatment intake.
3. Use electrolytes daily. GLP-1 medications naturally cause dehydration because appetite and fluid intake decrease.
4. Eat fruits and vegetables daily. Tirzepatide slows gastric emptying, meaning food stays in the stomach longer. Without adequate fiber, constipation occurs.

Protocol Note

Fiber is critical. Tirzepatide slows gastric motility. Clients who skip fiber consistently report constipation and digestive discomfort. Prioritize whole fruits, leafy greens, and vegetables at every meal.

Important Notice

This guide is intended for research and educational purposes only. Not intended to diagnose, treat, cure, or prevent any disease.

Questions or Concerns

Contact Straight Peptides directly for dosage guidance, product questions, or support.

Phone: 915-637-6443

Email: straightpeptides@gmail.com