

Subject Index

- 100% cosine data taper, 189
- absolute deviations sinusoidal amplitude estimator
(Laplace method), 529
- absolutely
 - continuous, 120
 - summable, 165
- acausal
 - autoregressive (AR) process, 446, 506
 - filter, 147–8, 152, 155
 - purely
 - autoregressive (AR) process, 447–8
- ACF (*see* autocorrelation function)
- acoustic properties of human speech
 - AR model, 447
- ACLS (*see* approximate conditional least squares)
- ACS (*see* autocorrelation sequence)
- ACVF (*see* autocovariance function)
- ACVS (*see* autocovariance sequence)
- adaptive multitaper spectral estimator, 389
 - ACVS issue, 615
 - distribution of, 390
 - EDOFs, 390
 - ocean wave data, 427–8
 - Parseval's theorem in expectation, 390, 439
- aggregated processes, 42
- AIC (*see* Akaike's information criterion)
- AICC (*see* Akaike's information criterion corrected for bias)
- Akaike's information criterion, 494
 - corrected for bias (AICC), 495
 - order selection, 494
 - centered time series, 495
- Akaike's information criterion corrected for bias (AICC), 495
 - order selection, 495
 - centered time series, 495
 - Willamette River data, 575–6
- aliases, 82
- aliasing, 83, 108, 113, 122–3, 569
 - and averaging, 161
 - antialiasing filter, 57, 224
 - different sampling intervals example, 131
 - discrete time/continuous frequency, 81
 - sea level residuals, 522
- amplitude spectrum, 54
- analytic series, 114, 561, 578–9
 - imaginary part of, 579
 - Willamette River data, 578, 592
- angular frequency, 8
- antialiasing filter, 57, 224
- applications of spectral analysis, 16–7
- approximate conditional least squares (ACLS)
 - conditions needed for ACLS to become ECLS
 - complex-valued case, 517
 - real-valued case, 515
 - sinusoidal amplitude estimator, 515, 525, 549, 569–70, 572–3, 585, 589
 - white noise variance estimator, 549, 570, 589
- approximate least squares, 516, 585
- approximate least squares sinusoidal amplitude estimator
 - expected value of, 516, 585
- AR (autoregressive) process (*see* autoregressive (AR) process)
- AR(1) process
 - asymptotic stationarity and causality, 448–9
 - autocovariance sequence, 44
 - backward least squares autoregressive parameter estimator, 510
 - Burg autoregressive spectral estimator, 510
 - forward least squares autoregressive parameter estimator, 510
 - from sampling a continuous parameter/time process, 507
 - infinite-order moving average form, 44
 - maximum likelihood estimator, 482

- model for colored noise, 335
- parameter estimation from estimated ACVS, 335–6
- simulation of Gaussian
 - via circulant embedding, 639–40
 - via Gaussian spectral synthesis method (GSSM), 639–40
 - via Kay's method, 639
- Yule–Walker autoregressive parameter estimator, 510
- AR(2) process, 34, 168, 172–3
 - alternative models, 440
 - asymptotic time series bandwidth, 347
 - bias in periodogram, 177
 - Burg autoregressive spectral estimator, 509
 - cumulative periodogram test, 216
 - direct spectral estimator, 186
 - and variability, 208
 - discretely smoothed periodogram, 309–10, 348–9
 - form of FBLS parameter estimator, 510
 - innovation variance estimators, 406–7
 - lag window spectral estimator, 294, 348–9
 - log SDF estimator, 304–5
 - maximum likelihood autoregressive spectral estimator, 489
 - numerical examples
 - Burg's algorithm, 509
 - periodogram, 172–3, 510
 - Yule–Walker estimator, 507–8
 - simulated examples, 34
 - direct spectral estimator properties, 242
 - periodogram properties, 242
 - simulation of Gaussian
 - via circulant embedding, 602–3
 - via Gaussian spectral synthesis method (GSSM), 608–9
 - via Kay's method, 597
 - tapering, 195
 - unimodality of spectrum and bandwidth of time series, 300
 - WOSA estimator, 416–7
 - Yule–Walker autoregressive spectral estimator, 509
- AR(4) process, 35, 168, 172–3
 - and end-point matching, 239
 - asymptotic time series bandwidth, 347
 - bias in periodogram, 178
 - Burg autoregressive spectral estimator, 468–9, 509
 - DCT-based periodograms, 218
 - direct spectral estimator, 186
 - discretely smoothed direct spectral estimator, 310–11, 348–9
 - eigenspectra, 361, 363, 368, 395, 397, 437
 - FBLS autoregressive spectral estimator, 478, 509
 - innovation variance estimators, 406–7
 - lag window spectral estimator, 295, 301, 349
 - log SDF estimator, 305–6
 - low-rank weighted multitaper spectral estimator, 422–3
 - maximum likelihood autoregressive spectral estimator, 482–3, 490
 - multitaper spectral estimator, 358–9, 364–5, 398–9, 400, 419
 - multitaper versus WOSA spectral estimation, 418–9
 - nonunimodality of spectrum and bandwidth of time series, 300–1
 - numerical examples
 - Burg's algorithm, 468–9, 509
 - periodogram, 172–3
 - Yule–Walker estimator, 458–9, 461–2, 509
 - prewhitening, 198–201
 - SDF and expectation of direct spectral estimator, 193
 - simulated examples, 34
 - direct spectral estimator properties, 242
 - periodogram properties, 242
 - simulation of Gaussian
 - via circulant embedding, 640–1
 - via Gaussian spectral synthesis method (GSSM), 606, 641
 - via Kay's method, 638–9
 - smoothing window
 - leakage, 289
 - parameter choice, 292
 - tapering, 185
 - WOSA estimator, 417–8
 - Yule–Walker autoregressive spectral estimator, 458–9, 461–2, 509
 - Yule–Walker versus periodogram spectral estimators, 508
- ARIMA (autoregressive, integrated, moving average) process, 16
- ARMA (*see* autoregressive moving average (ARMA) process)
- ARMA(1,1) process, 44
 - autocovariance sequence (ACVS), 44
 - infinite-order moving average form, 44
 - kurtosis, 632
 - skewness, 632
 - inequality, 633, 641
 - variance, 632
- asymptotic equivalence of autoregressive estimators, 484
- asymptotic relative efficiency, 165
- asymptotic stationarity and causality, 448–9
- asymptotically efficient estimator, 166, 522–3
- atmospheric CO₂ data, 332–3
- atomic clock data, 1–2
 - ACS for
 - sample, 4
 - theoretical, 10
 - lag 1 scatter plot for, 3
 - simulation of, 11
 - spectrum for
 - estimated, 12, 14
 - theoretical, 10, 12, 14
- atomic clock fractional frequency deviates, 326, 430
 - associated time differences, 325–6
 - direct spectral estimator, 327
 - periodogram, 327
 - Burg autoregressive spectral estimator, 501
 - innovation variance estimation, 431–2, 499–500

- simulation, 628–9
- lag window spectral estimator, 328, 501
- multitaper spectral estimator, 501
- periodogram, 328
- atomic clock timekeeping, 162, 593
- augmented Yule–Walker equations, 452
- autocorrelated cosine lag window, 284, 345
- autocorrelation, 72, 96–8, 100, 102, 112
- autocorrelation function (ACF), 27
- autocorrelation method, 480
- autocorrelation sequence (ACS), 4, 9, 27
 - conjugate symmetry, 30
 - estimation, 5, 19–20
 - MA(2) process, 159
 - sample, 3
 - white noise, 19
- autocorrelation width, 73, 97–8, 100, 102
 - smoothing window, 251
 - spectral window, 191–2, 194
- autocovariance function (ACVF), 27, 38
 - and the integrated spectrum, 114
 - and the SDF, 114
 - conjugate symmetry, 30
 - Gaussian-shaped, 131
 - positive semidefinite, 30
 - symmetry, 27
- autocovariance sequence (ACVS), 27, 29
 - and integrated spectrum, 111
 - ARMA process, 44, 600
 - AR process, 44, 460, 508
 - band-limited white noise, 379, 604
 - biased estimator of, 2–7, 166, 168–9, 232, 295, 505, 561, 591
 - for complex-valued process, 231, 562
 - positive definite, 451
 - classification examples, 121
 - complex-valued process, 29, 42
 - corresponding to
 - direct spectral estimator, 188, 611–12
 - lag window spectral estimator, 248, 612
 - multitaper estimator, 433
 - periodogram, 170–1
 - WOSA estimator, 420, 435
 - Yule–Walker autoregressive spectral estimator, 504, 508
- decay to zero, 120
- estimation of, 166, 231
- for process squared, 43
- fractionally differenced (FD) process, 440
- GSSM-based approximation, 606–7, 640
- harmonic process, 36–7, 117, 591
 - complex-valued, 519, 562
- inverse Fourier transform of SDF, 111, 126
- moving average process, 32
 - MA(2), 159
- piecewise-constant SDF, 348
- positive semidefinite, 28, 30, 43
- power-law process, 391, 605
- randomly phased sinusoid, 35, 511, 553, 598
- subsampling, 130
- symmetry and conjugate symmetry, 27, 29
- unbiased estimator of, 166, 168–9, 236–7, 271
 - for complex-valued process, 562
- versus spectral density function, 124
- white noise, 31, 117
- Wold’s theorem, 117, 126
- autoregressive (AR) process, 33, 445–6
 - acausal, 446, 506
 - acoustic properties of human speech, 447
 - ACVS, 460, 508
 - asymptotic stationarity and causality, 448–9
 - causality, 446, 448, 506, 618, 638
 - and parameter estimators, 448, 617
 - assessment, 463
 - conditions for, 34, 446
 - characterization, 509
 - order, 446
 - order selection (*see* order selection for autoregressive process)
 - plus white noise is an ARMA process, 554, 591
 - pseudo-, 553
 - purely acausal, 447
 - simulation, 595
 - zero initial conditions, 595
 - simulation of Gaussian
 - stationary initial conditions, 595, 599, 624, 637
 - via Kay’s method, 595–7
 - via McLeod and Hipel’s method, 599
 - sinusoidal variations, 447
 - spectral density function (SDF), 144–5, 446, 485, 609, 617–8
 - stationarity condition, 34, 145, 446, 448, 557
- autoregressive estimator
 - and prewhitening, 491, 506
 - asymptotic equivalences, 484
 - Burg’s algorithm, 466, 505
 - complex-valued process, 502
 - least squares estimators, 475, 478, 505
 - maximum entropy method, 471
 - maximum likelihood estimators, 480, 506
 - pure parametric approach, 506
 - variability assessment
 - via large-sample theory, 485
 - via simulation, 626
- Yule–Walker method, 449–51, 505
 - complex-valued process, 502
- autoregressive moving average (ARMA) process, 35, 492, 597–8
- ACVS, 600
- AR process plus white noise, 159, 554, 591
- causal, 600, 631
- maximum entropy, 473–4
- pseudo-AR process plus white noise, 554
- SDF, 145, 159
- simulation of Gaussian, 593–4, 597–8
 - stationary initial conditions, 599–600, 637, 639
 - via Kay’s method, 598, 639
 - via McLeod and Hipel’s method, 600, 639
- simulation of non-Gaussian, 631, 638
- simulation via step-down scheme, 461
- with distinct AR and MA coefficients, 554, 591
- with identical AR and MA coefficients, 554

- autoregressive moving average (ARMA) spectral estimator, 492, 504
- autorelation sequence, 29, 31
- average value of a function, 71

- background continuum, 511, 513, 523, 544, 547, 549, 551, 553, 583, 588
- backward least squares (BLS)
 - autoregressive parameter estimator, 477, 484
 - AR(1) example, 510
- backward linear prediction, 455
 - complex-valued process, 502, 564
- backward linear prediction error, 455
- band-limited function, 57–8, 93, 139
 - extrapolation of, 66
 - recovering the continuous from the sampled, 84
- band-limited white noise, 128, 131, 379, 604
- band-pass filter, 146
- band-pass process, 299, 347
- bandwidth
 - of AR(2) and AR(4) time series, 347
 - of spectrum, 292, 296, 346
 - estimation of, 297
 - of time series, 300–1, 321–2, 347
 - estimation of, 300
 - regularization, 357, 377, 425, 428, 434
 - smoothing window, 245, 251, 268, 281, 292, 296, 300–1, 318, 322, 341, 346–7
 - Bartlett–Priestley design, 344
 - choice of, 252
 - finer frequency grid, 255
 - for sea level residuals, 522
 - Grenander, 251, 268, 279, 281
 - lag window spectral estimator, 279
 - Parzen, 251
 - repeated lag window estimator, 349–50
 - spectral window, 206, 226–7, 229, 235, 268, 294, 296, 317–8, 323, 325, 328, 331, 335, 341, 347, 568, 623–4, 628
 - approximation for sinusoidal multitaper spectral estimator, 400, 409
 - basic multitaper spectral estimator, 437, 575
 - definition for basic multitaper spectral estimator, 353
 - definition for direct spectral estimator, 194
 - definition for eigenspectrum, 353
 - definition for lag window estimator, 256
 - definition for weighted multitaper spectral estimator, 353
 - definition for WOSA, 414–5
 - for sea level residuals, 521–2
 - frequency range for statistical results, 213, 244
 - versus equivalent width, 214, 355, 369, 402
 - versus regularization bandwidth, 402
 - WOSA, 429
 - spectral window versus smoothing window, 257
- Bartlett
 - lag window, 268–9, 281, 344, 412
 - smoothing window, 269–70
 - spectral window, 269–70
- Bartlett–Priestley
 - design window, 273–4, 344
 - lag window, 268, 274, 281, 320, 344
 - smoothing window, 274–5
 - spectral window, 274–5
- basic multitaper spectral estimator, 352–5, 432–4, 611, 613
 - ACVS estimator for, 433
 - and Gaussian white noise, 370–4, 402–3
 - and improved estimation of
 - innovation variance, 403–7
 - exponent of power-law process, 407–12, 440
 - log SDF, 412
 - and periodicity testing, 544–8
 - approximated by WOSA spectral estimator, 424
 - AR(4) data, 358–65, 368–70, 393–400, 405–7, 419
 - atomic clock fractional frequency deviates, 430–2, 501, 628–30
 - bias of, 359, 392
 - distribution of, 354, 356–7, 381–2, 409, 434
 - energy preservation of, 366–7
 - equivalent degrees of freedom (EDOFs) of, 354–5, 434
 - equivalent width, 355
 - expected value of, 353, 433
 - in circulant embedding simulation method, 611
 - jackknifing of, 356–7, 369–70
 - ocean wave data, 425–9
 - spectral window for, 353, 433
 - bandwidth of, 353, 359, 433
 - approximation (sinusoidal), 400–1, 434
 - sinusoidal, 398–400
 - Slepian, 358–9, 364–5
 - variance of, 353–4, 359
 - Gaussian white noise, 372–4, 402–3
 - via quadratic spectral estimator, 374
- basis for the class of band-limited functions, 66
- Bessel function, 196, 234
- best (backward) linear predictor, 455
 - complex-valued process, 502, 564
- best (forward) linear predictor, 452–3
 - complex-valued process, 502, 564
- best linear predictor, 404, 455, 460–1, 463, 472
- best predictor, 463
- bias
 - and dynamic range, 177
 - broad-band, 359, 380–2, 387–8, 390
 - defined, 378
 - indicator, 379–81, 439
 - minimizing indicator of, 380
 - due to leakage, 185, 192, 351
 - due to smoothing window, 256, 290
 - indicator for minimum-bias multitaper spectral estimator, 392, 441
 - local, 290, 359, 381–2
 - defined, 378
 - magnitude indicator, 378–9, 381
 - of direct spectral estimator, 192
 - of multitaper spectral estimator, 359
 - of periodogram, 163

- of weighted multitaper spectral estimator, 391, 441
- bivariate standard Gaussian PDF, 634
 - Mehler's expansion, 635
- Blackman–Tukey estimator, 252
- block averaging, 270–1, 412–14
- Bluestein's substitution, 94
- blurring function, 93
- bootstrapping, 593, 611, 618, 631, 638
 - confidence intervals for innovation variance, 630
 - second approach in AR case, 630
 - confidence intervals for log of SDF, 623–5
 - second approach in AR case, 626
- boundary conditions (*see* circular or reflecting)
- Brent's method, 528, 558
- broad-band bias, 359, 380–2, 387–8, 390
 - defined, 378
 - indicator, 379–81, 439
 - minimizing indicator of, 380
- Burg autoregressive estimator, 466–8, 505
 - AR(1) process, 510
 - AR(2) data, 509–10
 - AR(4) data, 468–9, 509
 - assessing variability (SDF), 485, 624–6
- atomic clock fractional frequency deviates, 501
- causality, 466, 505
- complex-valued process, 502, 562
- FPE criterion, 494
- ocean wave data, 496–9
 - order selection via PACS, 498–9
- of innovation variance, 629
- peak in SDF
 - height, 576
 - shifting and splitting, 560, 562, 576, 591
 - width, 576
- Willamette River data, 575–7
- Burg's algorithm, 466, 468, 470, 473, 478, 482, 491, 493, 505, 560, 575–6
 - analytic series, 562
 - and Yule–Walker method, 471
 - AR(2) numerical examples, 509–10, 556
 - observed prediction error, 466
 - two-step, 560
- carbon dioxide and global temperature, 366
- cascaded filter, 145
- Cauchy inequality, 257, 262, 371, 442
- causal process
 - autoregressive (AR), 34, 446, 448, 506, 618, 638
 - autoregressive moving average (ARMA), 600, 631
- causal filter, 147, 161
- causality, 446, 448, 506, 618, 638
 - and parameter estimators, 448, 617
 - assessment, 463
 - conditions for, 34, 446
- centering
 - AIC and AICC, 494–5
 - and direct spectral estimator, 195, 241, 244, 548
 - and periodogram, 184, 213, 240
 - and simulations, 619, 627
 - and tapering of a time series, 195
 - and WOSA spectral estimator, 419–20
 - of data for harmonic analysis, 583
- central peak (lobe), 175
- cepstrum, 301, 473, 503
 - estimator, 302–3
- Cesàro summability and sums, 78, 165
- chaotic beam data, 228
 - direct spectral estimator, 228–30
 - periodogram, 228–9
- Chebyshev's inequality, 164
- chi-square distribution, 37, 204, 209–10, 539
 - noncentral, 542, 552
- chi-square RV, 37, 203, 209–10, 235, 408, 539, 546
 - and confidence intervals, 438, 486, 490
 - logarithm of, 209, 412
 - expected value and variance, 302
 - scaled, 264, 298, 343
 - sum of chi-square RVs, 299
 - weighted sum of chi-square RVs, 264
- chirp transform algorithm, 94, 216
- Cholesky decomposition, 599, 604, 637, 639
 - modified, 464, 480, 599
- circular embedding simulation method, 440, 601–4, 611, 620, 627, 629, 631
 - ACVS properties, 602, 640
 - and direct SDF estimator, 611
 - and lag window SDF estimator, 611–12
 - and multitaper SDF estimator, 611, 613
 - and periodogram, 615, 641
 - and WOSA SDF estimator, 611, 613
 - compared to method of surrogate time series (MSTS), 616
 - computational efficiency, 603
 - issues with ACVS, 615
 - overcoming negative weights, 604, 637, 640
 - weights, 601–2, 605, 611, 629, 637, 640
 - AR(1) process, 640–1
 - direct SDF estimator, 612
 - lag window SDF estimator, 612
 - multitaper SDF estimator, 613
 - WOSA SDF estimator, 613
- circular boundary conditions, 424
- circularized sequence, 601–2, 640
- classification of spectra, 120–2
- CO₂ data, 332
 - lag window spectral estimator for filtered, 335
 - periodogram for filtered, 335
- coefficient of determination, 570, 572, 621
- colored noise, 120, 126, 219, 241, 377–8, 381, 389, 511, 513, 518, 520, 523, 539, 544, 550, 574, 584
 - proper complex-valued, 519, 585
- complementary covariance, 31
- complete stationarity, 215
- complex demodulation, 160, 232, 592
- complex exponentials, 17, 48, 134
 - linear combinations of, 132, 137
- complex-valued colored noise (proper), 519, 585
- complex-valued stochastic process
 - continuous parameter/time, 23

- discrete parameter/time, 23
- complex-valued Gaussian or normal RV (proper), 30, 546–7
- complex-valued process
 - by complex demodulation, 160, 374
 - resulting ACVS, 160
 - resulting SDF, 160, 374
 - example of, 127
- complex-valued roots, 566–7, 580, 592
- complex-valued stationary process, 29, 231, 374, 501–2, 519, 585
 - time reverse of conjugate, 502
- complex-valued white noise (proper), 517–8, 564
- compound periodicity, 540–1
- computational pathways, 221, 224, 315
- concentration measure
 - continuous time/continuous frequency, 62, 65
 - discrete time/continuous frequency, 85, 105, 155, 386
 - discrete time/discrete frequency, 94
 - for eigenspectrum spectral window, 368
- condition number of covariance matrix, 185
 - and dynamic range, 185
- confidence interval for
 - innovation variance, 629
 - bootstrapping, 630
 - via simulation, 630
- jump in the integrated spectrum, 590
- log of SDF, 205, 227, 229, 266–7, 323, 328, 331, 335, 342, 354, 359, 425, 521
 - bootstrapping, 623–5
- SDF, 205, 265, 267, 342, 357, 485, 488–90
- chi-square versus jackknifing, 438
- conjugate symmetry, 29–30, 48, 54, 75, 92
- consistent estimator of
 - mean, 164
 - SDF
 - discretely smoothed direct spectral estimator, 246
 - lag window spectral estimator, 260, 342
- constructed multitaper spectral estimator, 355, 376
- continuous parameter deterministic sinusoid, 558
- continuous parameter white noise, 38
- continuous parameter/time stochastic process, 22, 38, 113–4, 131, 507
 - sampling and aliasing, 122–3
 - sampling and averaging, 161
- convergence
 - factors, 80, 152–3, 157, 579, 592
 - in mean square, 49, 53
 - pointwise, 50, 53, 84
- convolution, 67, 96, 98–9, 101
 - as an LTI filter, 137, 157
 - continuous parameter, 142, 247
 - discrete parameter, 143, 247
 - as smoothing operation, 69–70, 247, 385
- cyclic, 101, 106
- Fourier transform of, 67, 96, 98–9, 101
- illustration of, 68–9, 104
- in frequency domain, 96, 98–9, 101, 105
- of periodic functions, 96, 105
- copper wires, 57
- correlation distortion due to zero-memory nonlinearity (ZMNL), 635–6
- correlation length, 131
- correlation of direct spectral estimator, 212, 243
- correlation time, 214
- correlogram method power spectral density estimator, 252
- cosine data taper, 189–90, 260
- cosine lag window, 290
- covariance
 - complex-valued case, 25, 41
 - real-valued case, 24
- covariance matrix, 28, 42, 44, 184, 376, 379, 382, 391, 440, 450, 457, 465, 485
 - Cholesky decomposition, 599, 604, 637, 639
 - modified, 464, 480, 599
 - condition number, 185
- covariance method, 480
- covariances of
 - direct spectral estimator, 205, 209–13, 243
 - in WOSA blocks, 415, 442
- eigenspectra, 354, 372, 436, 438
- lag window spectral estimator, 260–1, 342
- periodogram, 203–4, 235
 - subtle issues, 214–5
- weighted multitaper spectral estimator, 437
- weighted sums of time series values, 242
- CPDF (*see* cumulative probability distribution function)
- cross-correlation (complex), 72, 96, 98–9, 101–2
- cumulant, 215, 632
 - inequality for fourth, 632–3
- cumulative periodogram test, 215–6
 - examples
 - AR(2) data, 216
 - ocean noise, 230–1
 - Willamette River data, 550, 573–4
- cumulative probability distribution function (CPDF), 23
 - N -dimensional, 23
- cyclic
 - autocorrelation, 220
 - convolution, 220, 273
 - convolution theorem, 101, 106
 - convolutions via FFTs, 106
- cyclical frequency, 8
- Daniell
 - design window, 272
 - lag window, 268, 273, 281, 290
 - smoothing window, 250, 272–3, 320
 - spectral window, 272–3
- data taper (*see* taper)
- data window, 186
- DC (direct current), 518
- DCT-based periodogram, 217, 244
 - AR(4) process, 218
 - asymptotic unbiasedness of, 218
 - bias of, 218–9
 - statistical properties, 219
- decay
 - of envelope of sidelobes, 269
 - rate of sidelobes

- Bartlett smoothing window, 269, 276
- Bartlett–Priestley smoothing window, 275
- Daniell smoothing window, 273
- Gaussian smoothing window, 277
- Papoulis smoothing window, 278
- Parzen smoothing window, 276
- spectral windows, 281, 328
- decibel (dB) scale, 13, 209, 225–7, 266
- deconstructed multitaper spectral estimator, 355, 376, 420–3
- decorrelation time, 214
- default data taper, 188, 190
- degenerate kernel, 86
- degrees of freedom
 - for chi-square RV, 37, 202
 - one, 13, 203
 - two, 13, 203
 - zero, 542
- equivalent (EDOFs), 264, 266
 - discretely smoothed direct spectral estimator, 343
 - discretely smoothed periodogram, 347
 - lag window spectral estimator, 264, 279, 342
 - multitaper spectral estimator, 354–5, 390, 434
 - WOSA spectral estimator, 415–6, 442–3
- in a time series, 298, 507
- of order 2, 298
- derivative as an LTI filter, 157
- design window, 247
 - for Bartlett–Priestley, 273–4, 319, 344
 - for Daniell, 272, 319
- determinant of a matrix, 480
- deterministic
 - function, 48
 - sequence, 48
 - sinusoid, 553
- detrending (*see* trend removal and quadratic trend removal)
- deviance, 308
 - generalized cross-validated (GCV), 309, 313, 324, 339
- DFT (*see* discrete Fourier transform)
- diagonal matrix, 464, 565–6, 599
- differencing
 - for linear trend removal, 40
 - for seasonality/periodicity removal, 40, 42, 333
 - higher-order, 340
 - to achieve stationarity, 340, 431
- digamma function, 210, 296, 302, 345, 405, 409
- Dirac delta function, 74, 113, 120, 157, 250, 524
- direct current (DC), 518
- direct spectral estimator, 164, 186, 201, 233, 340, 375, 535–6, 545, 611
 - and centering a time series, 195–6, 241, 244, 548
 - atomic clock data, 327–9
 - average value of, 241
 - averaging of uncorrelated, 443
 - bias, 192
 - chaotic beam data, 228–9
 - computation of, 219–24
 - correlations, 212, 243
 - corresponding ACVS estimator, 188, 452, 611
 - covariances, 205, 209–13, 243
 - discretely smoothed, 246–7
 - equivalent degrees of freedom (EDOFs), 347
 - spectral window bandwidth, 347
 - distribution of, 204, 235
 - for complex-valued process, 232
 - eigenspectrum, 352
 - expected value of, 186, 234
 - for real-valued harmonic process, 535, 585
 - exponential averaging, 444
 - for blocks in WOSA, 414
 - harmonic process examples, 536, 588
 - in circulant embedding simulation method, 611–12, 627–8
 - inconsistent estimator of SDF, 207, 235
 - linear combination of independent, 441–2
 - ocean wave data, 225–6, 496
 - relationship to quadratic spectral estimator, 376
 - ship altitude data, 330–1
 - simulating properties of, 242
 - smoothing of, 245
 - spectrum of, 213
 - use with AR order selection, 493
 - use with Yule–Walker autoregressive estimator, 452, 461–2, 505
 - variability, 208
 - variance of, 204, 235, 624
 - for complex-valued process, 232
 - white noise, 189, 371
- Dirichlet's kernel, 17–8, 76–7, 80, 85, 93, 106, 127, 233, 383, 537,
- discrete cosine transform, 184, 217
- discrete Fourier transform (DFT), 74–5, 92, 219, 594, 602, 608
 - as a convolution, 93–4
 - chirp algorithm, 94
 - conjugate symmetry, 75, 92
- discrete parameter/time stochastic process, 22
- discrete power spectrum, 50, 139–40
- discrete prolate spheroidal sequence (DPSS), 87, 155, 189, 357
- discrete prolate spheroidal wave function (DP-SWF), 87, 105–6, 383–5, 439
- discrete spectrum, 120, 511–13, 518, 538, 584–6
- discrete time sequence, 74
 - autocorrelation width, 100
 - convolution theorem, 99
 - equivalent width, 100
 - Fourier representation, 75, 99
 - Parseval's theorem, 75, 99
- segment of, 91
 - autocorrelation width, 102
 - convolution theorem, 101
 - equivalent width, 102
 - Fourier representation, 92, 100
 - Parseval's theorem, 92, 101
 - spectral properties, 101
 - spectral properties, 99
- discretely smoothed direct spectral estimator, 246–7
 - computation of, 316
 - consistent estimator of SDF, 246

- equivalence to lag window spectral estimator, 249–50
- finer frequency grid, 253
- discretely smoothed periodogram, 307
 - bias/variance trade-off, 309
 - computational scheme, 347
 - corresponding ACVS estimator, 312–13, 348, 615
 - equivalent degrees of freedom (EDOFs), 347
 - evaluating SDF estimates, 349
 - ice profile data, 324
 - parameter choice, 307, 309–10, 312, 324, 339, 348
 - spectral window bandwidth, 347
- distribution of
 - adaptive multitaper spectral estimator, 390
 - direct spectral estimator, 204, 235
 - complex-valued process, 232
 - lag window spectral estimator, 264, 342, 490
 - maximum likelihood AR parameter estimator, 483
 - other AR parameter estimators, 484
 - multitaper spectral estimator, 354, 356, 381, 409, 434
 - periodogram, 203, 235, 408
 - WOSA spectral estimator, 436
- Doppler measurements of blood flow, 131
- double orthogonality
 - of eigenfunctions, 65, 87
 - of Slepian sequences (DPSSs), 106
- double-sided versus single-sided SDF, 118
- DPSS (*see* discrete prolate spheroidal sequence)
- DPSWF (*see* discrete prolate spheroidal wave function)
- duration–bandwidth product, 66, 357
- dynamic range, 177, 184, 197, 492, 641
 - and condition number, 185
- echoes, 227
- ECLS (*see* exact conditional least squares)
- EDOFs (*see* equivalent degrees of freedom)
- effective bandwidth (*see* spectral window bandwidth)
- eigenfunction, 64
 - double orthogonality, 65, 87
 - for a linear filter, 135
- eigenspectra, 352, 385, 432, 545
 - bandwidth of spectral windows, 353, 358
 - covariances of, 354, 372, 436, 438
 - plots of for AR(4) data, 361, 363, 368, 395, 397
 - regularization bandwidth, 357
 - weights associated with, 352, 388–9
- eigenvalue, 64, 86–7, 95, 185, 375, 380, 392, 565, 567, 580, 591
 - for a linear filter, 135
- eigenvector, 87, 95, 380, 392, 591
 - left and right, 565, 567
- end-point mismatch, 181, 183
- energy
 - of a nonperiodic function, 54–5
 - of a periodic function, 49
- energy spectral density function, 55, 97, 99, 132
- ensemble, 21–2
- entire function, 57
- entropy, 471–3
- envelope of sidelobes, 269, 273
- Epanechnikov lag window (*see* Bartlett–Priestley lag window)
- equal
 - by definition, 8, 23
 - in distribution, 203, 485, 599
 - in the mean square sense, 49, 53
- equivalence of time and frequency domains, 55
- equivalent degrees of freedom (EDOFs), 264, 266
 - adaptive multitaper spectral estimator, 390
 - example (ocean wave data), 427–8
 - basic multitaper spectral estimator, 354–5, 434
 - example (ocean wave data), 427
 - discretely smoothed direct spectral estimator, 343
 - discretely smoothed periodogram, 347
 - lag window spectral estimator, 264, 279, 342
 - example (ocean wave data), 318–20
 - via eigenvalues, 423–4
 - weighted multitaper spectral estimator, 354
 - WOSA spectral estimator, 415–6, 442–3
 - example (ocean wave data), 429–30
- equivalent width, 58–60, 73, 97–8, 100, 102, 213–4
 - multitaper spectral estimator, 355, 437
 - sinusoidal multitaper spectral estimator, 401, 441
 - smoothing window, 251
 - versus spectral window bandwidth, 214, 355, 369
 - sinusoidal multitapers, 402
 - Slepian multitapers, 369
- ergodic theorems, 165
- Euler relationship, 17
- Euler's constant, 210, 302, 404, 408
- exact conditional least squares (ECLS)
 - including mean, 552
 - sinusoidal amplitude estimator, 549, 569–70, 572, 589–90
 - white noise variance estimator, 549, 552, 570, 589–90
- exact unconditional least squares (EULS)
 - including mean, 552
 - sinusoidal amplitude estimator, 549, 569–70, 572
 - sinusoidal frequency estimator, 549
 - white noise variance estimator, 550, 552, 570
- expected value (expectation), 4
- expected value of
 - direct spectral estimator, 186, 234
 - real-valued harmonic process, 535, 585
 - lag window spectral estimator, 255, 341, 343
 - multitaper spectral estimator
 - adaptive, 389–90, 439
 - basic, 353, 433
 - weighted, 352
 - periodogram, 174, 233, 240
 - complex-valued harmonic process, 524, 586
 - real-valued harmonic process, 524, 585
 - conditional on phase value, 537–8
 - quadratic spectral estimator, 376–7, 439

- sinusoidal frequency estimator using periodogram maximum, 526
- WOSA spectral estimator, 414, 435
- exponential averaging, 444
- exponential distribution, 202, 209, 215, 598
- exponential RV, 209
- extrapolation of a band-limited function, 66
- fader, 186
- fast Fourier transform (FFT), 92, 94, 237, 602, 608
- FBLS (*see* forward and backward least squares)
- F -distribution, 547, 574, 582, 587
 - non-central, 552
 - percentage points (special case), 544
- Fejér's kernel, 80, 174–6, 180, 233, 240, 524, 532, 534, 537
 - different expressions for, 236
 - sampling of, 524
 - sidelobes, 178–80, 328, 527, 532, 535–6
- FFT (*see* fast Fourier transform)
- filter
 - acausal, 147–8, 152, 155
 - analog, 133
 - antialias, 57, 224
 - band-pass, 146
 - cascaded, 145
 - causal, 147
 - demodulation, 160
 - digital, 140
 - effect on integrated spectrum (filtering theorem)
 - analog, 136, 157
 - digital, 141
 - example of an LTI digital, 148
 - finite impulse response (FIR), 147–8, 152, 155, 161, 503
 - first difference, 333
 - for pilot analysis, 160
 - high-pass, 146
 - as trend remover, 151
 - formed by subtracting output of low-pass filter from its input, 149
 - infinite impulse response (IIR), 147, 503
 - Kalman, 482
 - kurtosis norm of, 632
 - linear time-invariant (LTI), 132, 631, 633
 - analog, 133
 - and convolution, 137
 - digital, 140
 - low-pass, 146, 592
 - cutoff frequency, 151
 - design by least squares, 152
 - design using Slepian sequences, 155
 - prediction error, 198
 - prewhitening, 197, 234, 241, 447, 482, 491–3, 496, 510, 615
 - seasonal differencing, 333
 - transfer function, 136, 141
 - transformation or operator, 133
- final prediction error (FPE), 493
- finite impulse response (FIR) filter, 147–8, 152, 155, 161, 503
- FIR (*see* finite impulse response)
- first difference filter, 40, 333
 - effect of filter on SDF, 158, 326
- Fisher's test for simple periodicity, 539–40, 550, 586
 - critical value, 581
 - approximation, 540, 589
 - ocean noise data, 581
 - Willamette River data residuals, 571, 573
- flicker frequency noise, 431
- folding frequency (Nyquist frequency), 82
- forward and backward least squares (FBLS)
 - and SVD approach, 565
- autoregressive parameter estimator, 477, 484
 - AR(2) examples, 510, 555–6
- autoregressive spectral estimator
 - AR(4) example, 478, 482
 - Willamette River data, 592
- same as modified covariance method, 480
- forward least squares (FLS)
 - autoregressive parameter estimator, 477, 484
 - AR(1) example, 510
 - same as covariance method, 480
- forward linear prediction, 452–3
 - complex-valued process, 502, 564
- forward linear prediction error, 453, 596
- Fourier coefficient, 49
- Fourier frequencies, 8, 91, 171, 515, 523–4
 - fundamental, 357
 - grid of, 171
 - alternative grid for uncorrelated SDF estimators, 206, 208, 212
- Fourier integral representation, 54
- Fourier representation of a periodic function, 49, 95
- Fourier synthesis, 54
- Fourier theory
 - continuous time/continuous frequency, 53, 97
 - continuous time/discrete frequency, 48, 95
 - discrete time/continuous frequency, 74, 99
 - discrete time/discrete frequency, 91, 100
- Fourier transform pair, 54, 96–7, 99–100
 - examples of, 171, 181, 188, 212, 220–2, 237, 247–8, 301–2
 - for convolution of two functions, 67
 - for derivative of a function, 61
- FPE (*see* final prediction error)
- fractionally differenced (FD) process, 166, 407–8, 440
 - autocovariance sequence (ACVS), 440
 - SDF, 407–8
 - simulation by circulant embedding, 440
- Fredholm integral equation
 - of the first kind, 383
 - of the second kind, 63, 86
- frequency grid, 171, 527, 531, 558, 578, 605
 - for circulant embedding, 601
 - for uncorrelated SDF estimators, 206, 208, 212
- frequency response function, 136
 - Hilbert transform, 579
- function, special type of
 - blurring, 93
 - digamma, 210

- Dirac delta, 74
- gamma, 440
- modified Bessel, 196
- periodic, 48
- positive-part, 541
- trigamma, 296
- function width
 - autocorrelation, 73, 97–8, 105, 191, 194, 288
 - equivalent, 58, 73, 97–8, 105, 213
 - half-power, 191–2, 288
 - variance, 60, 73, 105, 191–2, 194, 288
- fundamental Fourier frequency, 357
- fundamental frequency, 569
- gain function, 136
 - squared
 - for differencing, 334
 - for Hilbert transform approximation, 592
 - for ideal low-pass, high-pass and band-pass filters, 146
- gamma function, 297, 345, 440, 542, 633
- Gaussian distribution, 265
 - generalized, 633
- Gaussian kernel, 69
- Gaussian lag window, 268, 276, 278, 281, 320, 330, 430
- Gaussian or normal process, 29, 445, 463, 480, 490, 544
 - autoregressive, 34, 445, 595
 - complex-valued, 30
 - entropy, 472
 - moving average, 33, 595
 - white noise, 31, 201, 210, 539, 594
- Gaussian or normal RV
 - complex-valued, 30, 210
 - PDF and its Fourier transform, 55
- Gaussian smoothing window, 276–8
- Gaussian spectral synthesis method (GSSM), 605–8, 637, 640–1
 - ACVS approximation, 607
 - adjustment for power-law process, 610
 - and adaptive multitaper spectral estimator, 615
 - and prewhitening, 615
 - role of sampling interval, 609
 - weights for AR(1) process, 640–1
- Gaussian spectral window, 278
- Gaussian white noise, 31, 201, 210, 539, 594
 - multitaper spectral estimator, 370
- Gaussian-shaped ACVF and SDF, 131
- Gegenbauer process, 610
- generalized cross-validated (GCV) deviance, 309, 313, 324, 339
- generalized Gaussian (GG) distribution, 633
- generalized inverse, 566
- generalized least squares, 412, 522
- Gibbs phenomenon, 77, 79, 81, 152
- goodness-of-fit test
 - Kolmogorov, 215
- grid size, 179, 527, 531, 558, 568, 578, 605
- group delay, 136
- GSSM (*see* Gaussian spectral synthesis method)
 - half-power width of function, 191–2, 288
 - Hamming lag window, 343
 - Hanning
 - data taper, 189–90, 222, 230, 234, 260, 305, 327, 331, 345, 415–8, 535
 - alternative form, 240, 285
 - lag window, 343, 579
 - harmonic analysis, 511
 - ocean noise data, 581
 - ocean wave data, 582
 - parametric approach, 553
 - AR examples, 555–6
 - problems with, 558
 - singular value decomposition (SVD) approach, 563
 - summary, 584
 - Willamette River data, 567
 - centering of time series, 583
 - harmonic process, 35, 108, 511
 - ACVS for, 36–7, 117
 - expected value of, 36
 - periodicity, 603–4
 - random phase, 37, 45
 - simulation, 593–4, 598–9, 616, 637, 641
 - examples, 36
 - variance of, 36, 108, 110
 - variance spectrum of, 108
 - harmonic process with additive noise
 - complex-valued, 519, 585–6
 - ACVS for, 519, 562, 585–7
 - expected value of, 519, 585–6
 - known frequencies, 517, 586
 - unknown frequencies, 586
 - variance of, 519, 585–6
 - real-valued, 512, 544, 584–5
 - ACVS for, 519, 584, 591
 - expected value of, 516, 584
 - known frequencies, 513, 515, 585
 - random phase, 516
 - signal-to-noise ratio, 526, 532
 - tapered, 535–6, 588
 - unknown frequencies, 523, 585
 - variance of, 519, 526, 532, 584
 - harmonic tidal prediction, 512
 - at Southampton, England, 513
 - harmonics
 - in tidal prediction, 522
 - of fundamental frequency, 569
- Heisenberg's uncertainty principle, 61–2, 104
- Hermite polynomials, 634–5
- Hermitian Toeplitz matrix, 30
- high-cut frequency, 57
- high-pass
 - filter, 146
 - process, 299, 347
 - creation of from a low-pass filter, 160
- Hilbert transform, 114, 562, 579
 - frequency response function of, 579, 592
 - impulse response sequence of, 579, 592
 - transfer function of, 579, 592
- ice profile data, 321
 - lag window spectral estimator, 322–3
 - periodogram, 321, 323

- plot of, 322
- IID (independent and identically distributed)
 - noise, 32, 631
 - process, 32
- IIR (*see* infinite impulse response)
- imaginary part of
 - analytic series, 579
 - complex-valued number, 551
 - complex-valued process, 602, 606, 640
- impulse response
 - function, 142
 - sequence, 143
 - for Hilbert transform, 579, 592
 - normalization of, 150
- inconsistency of
 - direct spectral estimator, 235
 - periodogram, 163–4, 207–8
- index-limited sequence, 85
- indirect nonparametric estimator, 252
- infinite impulse response (IIR) filter, 147, 503
- initial conditions
 - AR process simulation, 595
 - Gaussian AR process simulation, 595, 599
 - Gaussian ARMA process simulation, 600, 639
- inner product, 470
- innovation process, 446
 - estimation of, 17
- innovation variance, 403–4, 446, 449, 455, 468, 471–2, 510
 - Burg autoregressive estimator, 499–500, 629
 - confidence interval for, 629
 - different estimators compared, 406, 440
 - direct spectral estimator, 405
 - mean square linear prediction error for $AR(p)$, 455
 - multitaper spectral estimator, 405, 432, 629
 - variability assessment via simulation, 630
 - periodogram-based estimator, 404
 - simulation, 628–9
 - variability assessment via, 629
 - smoothed periodogram estimator, 404
 - Yule–Walker autoregressive estimator, 499–500
- integrable (in square sense), 49, 53–4
- integral equation, 63, 86
- integral time scale, 131, 214
- integrated magnitude square error, 52, 76
- integrated spectrum (spectral distribution function), 110, 117, 163, 511
 - absolutely continuous, 120
 - and AR SDF (parametric sinusoidal power estimation), 578
 - and LTI filters, 135–6, 141
 - cascade of filters, 146
 - basic properties, 116
 - classification examples, 121
 - continuous parameter stationary process, 114
 - continuous singular function, 120
 - differentiability, 120
 - discrete parameter stationary process, 110
 - jumps (steps), 120, 511, 549, 551, 589–90
 - confidence interval for, 590
 - distribution of estimator, 552, 590
 - Willamette River example, 570, 572, 575
- intrinsically stationary process
 - order d , 340
 - order unity, 431
- inverse Fourier transform (*see* Fourier transform pair)
- invertibility of moving average process, 43
- irregularly sampled observations, 355, 528, 543
- Isserlis theorem, 30, 43, 210, 236, 241, 298, 371
- jackknifing
 - and multitaper spectral estimator, 356, 369–70
 - for confidence interval calculation, 438
- Johnson system of distributions, 633
- joint moment, 27
- joint stationarity, 29
- jump process, 109
- jumps in the integrated spectrum, 120, 511, 549, 551, 589–90
 - confidence interval for, 590
 - distribution of estimator, 552, 590
 - Willamette River example, 570, 572, 575
- Kalman filter, 482
- kernel
 - degenerate, 86
 - Dirichlet's, 17–8, 76, 93
 - Fejér's, 80, 174–5
 - Gaussian, 69, 290
 - of an integral equation, 63, 86
 - rectangular, 71, 290
- Kolmogorov goodness-of-fit test, 215
- Kronecker's delta function, 44
- k th-order Slepian data taper, 357, 382, 384
 - plots of, 360, 362
- Kullback–Leibler (KL) discrepancy, 297, 312, 346, 349
 - expected value, 297
 - for scaled chi-square RVs, 345
 - symmetrized form, 345–6, 349
 - with swapped arguments, 345–6, 349
- kurtosis, 631
 - ARMA(1,1) process, 632
 - norm of a filter, 632
- lag, 2, 27
 - lag 1 scatter plot, 2
 - lag p differencing, 40
 - lag τ scatter plot, 2
- lag window, 248, 341
 - choice of, 287
 - compared to data taper, 282–3, 285
 - constructed from data taper, 283, 344–5
 - parameter m (controls smoothing), 247, 287
 - choice of, 291, 304, 313–4
 - examples of, 318, 322, 328, 330, 339
 - properties of, 250
 - reshaped, 257, 281–2, 375
 - type
 - autocorrelated 100% cosine, 283–5, 344–5
 - Bartlett, 268–9, 281, 344, 412
 - Bartlett–Priestley, 268, 273, 281, 320, 344
 - cosine (100%), 283–4, 290

- Daniell, 268, 271, 281, 290, 320
- Gaussian, 268, 276, 278, 281, 320, 331, 430
- Hamming, 343
- Hanning, 343, 579
- Papoulis, 268, 278–9, 281, 345
- Parzen, 248, 268, 275, 281, 301, 320–1, 323, 331, 344, 346, 496, 521–2
- Slepian, 285, 345
- lag window spectral estimator, 245, 248, 340–1, 492, 612
 - alternative names for, 252
 - as a quadratic spectral estimator, 375
 - as deconstructed multitaper estimator, 420–2
 - asymptotic distribution of, 264–5, 297, 342, 490
 - asymptotic unbiasedness of, 256, 341
 - asymptotic variance of, 258–9, 262, 265, 279, 341
 - verification of validity by simulation, 622–3
 - atomic clock data, 328–9, 501
 - computation of, 314–7
 - consistent estimator of SDF, 260, 342
 - corresponding ACVS estimator, 248, 341, 612
 - covariances, 260, 342
 - equivalence to discretely smoothed direct spectral estimator, 249–50
 - equivalent degrees of freedom (EDOFs), 264, 279, 342
 - expected value of, 255, 341, 343
 - filtered CO₂ data, 335
 - gamma PDF, 297
 - ice profile data, 321–4
 - in circulant embedding simulation method, 611–12, 619–20, 627
 - ocean wave data, 316–9, 349–50, 430, 496
 - parameter m (controls smoothing), 247, 287
 - choice of, 291, 304, 313–4
 - examples of, 318, 322, 328, 330, 339
 - repeated, 349–50
 - sea level residuals, 521–2
 - ship altitude data, 330–2
 - smoothing window bandwidth B_W , 251, 279
 - alternative measure β_W , 251, 279
 - use in prewhitening, 491
- Laplace periodogram, 529–30
- law of total expectation, 588
- leakage, 77–8, 178, 180, 196, 233, 287, 358, 532, 592
 - and end-point mismatch, 181, 239
 - bias in spectra due to, 225, 227, 233, 351
 - reduction of, 78, 185
 - smoothing window, 268, 287–9, 294–5, 316, 318–9, 498
- least squares (LS), 40, 50, 332, 409, 430–1, 475–8, 513–6
 - approximate, 516, 585
 - approximate conditional (*see* approximate conditional least squares)
 - complex-valued, 517, 546, 565
 - exact conditional (*see* exact conditional least squares)
 - exact unconditional (*see* exact unconditional least squares)
 - filter design, 152, 155
 - generalized (GLS), 412, 522
 - ordinary (OLS), 409, 430, 440, 522
- least squares (LS) autoregressive estimators, 475–9, 505–6
 - AR(1) process, 510
 - assessing variability (SDF), 485
 - backward, 477, 484, 510
 - causality not guaranteed, 479, 506, 510, 617
 - forward, 475–7, 484, 510
 - forward and backward, 477–8, 484, 509, 555–6, 565
 - relation to Yule–Walker scheme, 479–80
- Lebesgue decomposition theorem, 120, 518
- Levinson–Durbin recursions, 452, 456–7, 461–2, 464–6, 509, 556, 604, 639
 - for complex-valued time series, 502
 - linear algebra derivation, 463–4
 - step-down scheme, 460–1, 507–8, 597
 - causality assessment, 461, 463, 509
 - generation of ACVS for an AR process, 460, 508
 - simulation of ARMA process, 461, 595–8
- likelihood function, 308, 480
 - and order selection criteria, 494–5
 - log of, 480–1
 - reduced (profile), 482–4
 - Whittle, 307
- line spectrum, 120, 511, 522
- linear and log scales for SDFs, 336–9
- linear model, 2, 39, 46, 621
- linear regression, 151, 408–10, 412, 440, 446, 501
 - complex-valued, 517, 545
 - real-valued, 513–5, 529
 - robust, 530
- linear taper, 186
- linear time-invariant (LTI) filter, 132
 - analog, 133
 - digital, 140
 - linearity of, 133, 140
- linear trend removal
 - by differencing, 40
 - by linear least squares estimation, 40
- linear window, 186
- local bias, 290, 359, 381–2
 - defined, 378
 - indicator, 378–9, 381
 - minimizing indicator of, 381
- log and linear scales for SDFs, 336–9
- log spectral density function, 301
 - estimation of
 - log direct spectral estimator, 301–2
 - with smoothing, 303–6, 324
- Lomb–Scargle periodogram, 528–9, 543–4, 588
- long memory process, 407
- long-range dependence, 407–8, 411
- Lorenzian SDF, 131, 507
- loss of resolution, 76, 78, 271, 290, 357, 359, 400, 434, 535
- low-cut frequency, 57
- low-pass
 - filter, 146, 592
 - process, 299, 347
 - use of to create a high-pass filter, 160

- low-rank approximation to SDF estimator, 420–3
lower triangular matrix, 464, 484, 509, 599, 639
LTI (*see* linear time-invariant)
- MA (moving average) process (*see* moving average (MA) process)
male speech, 57
Mann–Wald theorem, 486
marginal distribution, 45, 631, 634
matching condition, 135–6, 141–3
matrix
 Cholesky decomposition of, 599, 604, 637, 639
 modified, 464, 480, 599
 covariance, 28, 42, 44, 184, 376, 379, 382, 391, 440, 450, 457, 465, 485
 determinant of, 480
 diagonal, 464, 565–6, 599
 generalized inverse of, 566
 lower triangular, 464, 484, 509, 599, 639
 rank of, 375–6, 380–1, 420, 422–3, 490, 565–6
 Schur, 484
 singular value decomposition (SVD) of, 425, 564–5
 and short time series, 567
 approach to harmonic analysis, 563
 Willamette River data, 578–80, 592
 singular value, 565
 singular vector, 565
Toeplitz, 29, 156, 457, 462, 480, 484
 Hermitian, 30
 trace of, 376, 378, 439
 tridiagonal, 88, 95
maximum confusion spectral analysis, 473
maximum entropy
 principle of, 471, 473
 for ARMA process, 473–4
 spectral analysis (MESA), 194, 447, 466, 471–4, 503
 example for MA processes, 475–6
maximum likelihood autoregressive estimator (MLE), 480–4, 506
 approximations to, 484, 510
 AR(1) process, 482
 AR(2) data, 489
 AR(4) data, 482–3, 490
 assessing variability (SDF), 485
 distribution of, 483
 recursive, 485
mean, 4, 24
 complex-valued case, 25
 estimation of, 3, 164–6, 184, 195–6, 213, 548
mean correction (*see* centering)
mean integrated square error (MISE), 303
mean square convergence, 49, 52–3
mean square error (MSE), 167, 236, 293, 346, 349, 386–8, 404, 463, 526, 537
 normalized (NMSE), 296, 314, 346, 349
mean square linear prediction error, 453–5, 457, 460, 468, 493, 509
 same as AR(p) innovation variance, 455
mean square log error (MSLE), 296, 346, 349
mean square prediction error, 463
Mehler's expansion, 635
method of surrogate time series (MSTS), 615–6, 641
 and harmonic processes, 641
 compared to circulant embedding, 616
minimum-bias multitapers, 392–3, 441
 bias indicator of, 392, 441
 plots of, 393
 relationship to sinusoidal multitapers, 392–3
mixed spectrum, 120–1, 359, 511, 513, 518–9, 521–2, 539, 584–5
modified
 Cholesky decomposition, 464, 480, 599
 covariance method, 480
 Bessel function, 196
 Daniell spectral estimator and windows, 280
 periodogram, 186
moment
 coefficient of kurtosis, 631
 coefficient of skewness, 631
 first and second, 27
 joint, 27
moving average (MA) process, 32, 43, 503–4, 594–5
 autocovariance sequence (ACVS), 32, 128
 invertibility, 43
 noninvertibility, 474
MA(1)
 ACVS, 504
 as infinite-order AR process, 43
 simulated examples, 33
MA(2), 159–60
 simulation of, 594–5, 637, 639
 spectral density function (SDF), 130, 144, 503
 textile slivers, 503
moving average (MA) spectral estimator, 503
 and truncated periodogram, 504
MSTS (*see* method of surrogate time series)
multitaper spectral estimator, 351–5, 432–4
 adaptive, 389–90
 ACVS (issue of being ill-defined), 615
 distribution of, 390
 versus weighted multitaper estimator, 390
 and improved estimation of
 innovation variance, 403–7
 exponent of power-law process, 407–12, 440
 log SDF, 412
 and periodicity testing, 544–8
 approximated by WOSA spectral estimator, 424
AR(4) data, 358–65, 368–70, 393–400, 405–7, 419
 atomic clock fractional frequency deviates, 430–2, 501, 628–30
basic, 352–5, 432–4, 611, 613
 ACVS estimator for, 433
 and Gaussian white noise, 370–4, 402–3
 bias of, 359, 392
 distribution of, 354, 356–7, 381–2, 409, 434
 energy preservation of, 366–7
 EDOFs of, 354–5, 434
 equivalent width, 355

- expected value of, 353, 433
 - jackknifing of, 356–7, 369–70
 - spectral window for, 353, 433
 - bandwidth of, 353, 359, 433
 - approximation (sinusoidal), 400–1, 434
 - sinusoidal, 398–400
 - Slepian, 358–9, 364–5
 - variance of, 353–4, 359
 - Gaussian white noise, 372–4, 402–3
- constructed, 355, 376
- deconstructed, 355, 376
- eigenspectra (*see basic entry for eigenspectra*)
- in circulant embedding simulation method, 611–13
- information recovery, 195, 366, 372
- minimum-bias, 391–3
- ocean wave data, 425–9
- peak-matched multitapers, 382
- sinusoidal multitapers, 355, 391–4, 396, 434
- Slepian multitapers, 355, 357–8, 360, 362, 368, 434
- versus prewhitening, 424
- versus WOSA spectral estimator, 414, 417–9, 424
- via quadratic spectral estimator, 374
- via regularization, 382
- weighted, 352–4, 388, 433, 611–13
 - and quadratic spectral estimator, 376
 - bias of, 391
 - covariances of, 437
 - distribution of, 354–5
 - EDOFs of, 354–5
 - equivalent width, 355, 437
 - expected value of, 352, 389
 - low-rank approximation, 422–3
 - Parseval's theorem in expectation, 390
 - spectral window for, 352–3, 391
 - bandwidth of, 353
 - variance of, 354
- multitaper-based test for periodicity, 544–8
 - sinusoidal amplitude estimator (complex-valued), 546, 551, 590
 - Willamette River data, 574–5
 - white noise variance estimator, 546, 551, 590
- multivariate Gaussian (normal), 4, 29–30, 483, 485, 510
- mutually (pairwise) uncorrelated random variables, 8, 13, 108, 203–4, 206, 235, 246, 258, 302, 306, 381, 410, 443, 465, 546, 595–6
- Newton–Raphson method, 528, 558
- noise
 - colored, (*see entry for colored noise*)
 - flicker frequency, 431
 - IID, 32, 631
 - in signal-to-noise ratio, 526
 - random walk frequency, 431
 - white, (*see entry for white noise*)
 - with heavy-tailed distribution, 530
- noncentral
 - F -distribution, 552
 - chi-square distribution, 542, 552
- nonergodic stationary process, 128
- non-Gaussian (non-normal) process, 36, 210, 412, 463, 530, 631–6
 - plots with examples of, 31
- nonindependent but uncorrelated random variables, 45–6, 108
- nonlinear lag relationship, 5–6
- nonnegative definite (*see positive semi-definite*)
- nonperiodic function, 53–4, 107, 143
- nonstationary process
 - complex-valued, 42
 - differencing for stationarity, 333, 340, 431
 - examples of, 39–41, 44–6, 332
- nonwhite noise (*see colored noise*)
- normal equations, 514, 517
- normal process (*see Gaussian process*)
- normalized mean square error (NMSE), 296, 314, 346, 349, 607
- normalized observed prediction error
 - backward, 618
 - forward, 618, 624, 626, 630, 638
- normalized root mean-square error, 406, 440
- Nyquist
 - frequency, 82, 84, 108, 122–3, 142
 - rate, 84, 123
- observed prediction error
 - backward, 466, 492, 618
 - normalized, 618
 - forward, 466, 491, 496, 617–8
 - normalized, 618, 624, 626, 630, 638
- ocean noise data, 1–2
 - ACS for
 - sample, 4
 - theoretical, 10
 - lag 1 scatter plot for, 3
 - periodogram for, 231
 - simulation of, 11
 - spectrum for
 - estimated, 12, 14
 - theoretical, 10, 12, 14
 - testing for white noise, 11, 230–1, 581–2
- ocean tide prediction, 512
- ocean wave data, 224–8, 316–21, 425–30, 496–9, 582–3, 619–26
 - adaptive multitaper spectral estimator, 428
 - equivalent degrees of freedom (EDOFs), 427–8
 - Burg autoregressive spectral estimator, 496, 500, 625
 - direct spectral estimator, 226
 - estimated PACS, 498–9
 - harmonic analysis of, 582–3
 - lag window spectral estimator, 318–9, 349–50, 430, 496, 500, 623
 - multitaper spectral estimator, 426, 582
 - periodogram, 226–7
 - plot of, 225, 620
 - prewhitening, 496–7
 - reshaped spectrum, 582
 - simulated series, 620
 - uses for, 626
 - Thomson's F -test for periodicity, 582

- WOSA spectral estimator, 430
- Yule–Walker autoregressive spectral estimator, 496
- octave, 269, 273, 276–7, 282, 288–9
- one-sided versus two-sided SDF, 118
- orchestral music, 57
- order
 - of autoregressive process, 33, 446
 - of autoregressive moving average process, 35
 - of autoregressive spectral estimator, 458
 - of linear predictor, 452–3, 455
 - of moving average process, 32, 503
 - of taper in multitapering scheme, 357, 392–3
- order selection for autoregressive process, 492–5
 - AIC method, 494
 - centered time series, 495
 - AICC method, 495
 - centered time series, 495, 575–6
 - direct spectral estimator criterion, 493
 - examples of use of, 496, 498–500
 - FPE method, 493
 - PACS method, 493, 498
- ordinary least squares (OLS), 409, 430, 440, 522
- orthogonal
 - data tapers, 354
 - increments, 109
 - process, 109
- orthogonality
 - of Hermite polynomials, 635
 - principle, 454
 - corollary to, 508
 - relationships (trigonometric), 19, 202, 516
- orthonormality
 - basis exhibiting, 138, 384
 - nonuniqueness of, 138
 - of data tapers, 432
 - of eigenvectors, 565
- oscilloscope, 21–2
- outliers, 196, 227
- oversmoothing, 71
- oxygen isotope ratios, 366
- PACS (*see* partial autocorrelation sequence)
- pairwise uncorrelated random variables (*see* mutually (pairwise) uncorrelated random variables)
- Papoulis
 - lag window, 268, 278–9, 281, 290, 345
 - smoothing window, 278–9
 - spectral window, 279
- parametric approach to harmonic analysis, 553–8
 - AR(2) example, 555–6
 - AR(16) example, 556–7
 - problems with, 558–62
 - Willamette River data, 575–8
- parametric spectral estimation, 14, 445, 482, 491–2, 503, 506
 - AR versus MA, 503
 - summary, 505
- Parseval's theorem, 49, 55, 75, 92
 - in expectation
 - adaptive multitaper spectral estimator, 390, 439
 - weighted multitaper spectral estimator, 390
 - one discrete time sequence, 99
 - example of usage, 85, 111, 115, 188, 194, 251, 260, 303, 385, 390
 - one nonperiodic function, 97
 - example of usage, 61
 - one periodic function, 96
 - example of usage, 49, 52, 137–8
 - one segment of discrete time sequence, 101
 - two discrete time sequences, 99
 - two nonperiodic functions, 97
 - two periodic functions, 96
 - two segments of discrete time sequences, 101
- partial autocorrelation
 - coefficient, 457, 466, 469, 509
 - sequence (PACS), 462–3
 - complex-valued process, 502
 - estimated for ocean wave data, 498–9
 - for AR order selection, 493, 498
- Parzen
 - lag window, 248, 268, 275–6, 281–2, 292, 316, 320–1, 331, 344, 346, 496, 521–2
 - smoothing window, 275, 280–1, 344
 - spectral window, 275
- PDF (*see* probability density function)
- peak-matched multitapering, 382
- peak shifting, 559–60, 576
 - and splitting, 560
 - analytic series, 561
 - multiple sinusoids, 562
 - with Burg's algorithm, 560, 562, 591
 - with Yule–Walker approach, 560–2, 591
- peak width, 576
- Pearson product moment correlation coefficient, 3
- percentage of block overlap in WOSA, 415
- periodic extension, 54, 92
- periodic function, 48–52, 74, 95–7
 - convolution theorems, 96–7
 - Fourier representation, 49, 95–6
 - Parseval's theorem, 49, 96
 - spectral properties, 50, 96
- periodicity
 - compound, 540–1, 587
 - elimination by differencing, 40
 - simple, 540–1, 586
 - testing with white noise, 538–44, 586–7
 - Fisher's test for (simple periodicity), 539, 586
 - example of use, 571, 573, 581
 - irregular sampling (Lomb–Scargle), 543–4
 - Schuster's test for (simple periodicity), 539
 - Siegel's test for (compound periodicity), 541, 587
 - example of use, 581
 - Thomson's F -test for (colored noise), 544–8, 587
 - example of use, 574–5, 582–3, 592
- periodogram, 170–185, 232–3
 - and DFT of time series, 116, 171
 - and real-valued harmonic process plus white noise, 523

- average value of, 241
- averaging of independent, 441
- bias, 163, 176–7, 218
 - AR(2) process (low dynamic range), 177
 - AR(4) process (high dynamic range), 178–80
- asymptotic (unbiasedness), 175
- average value of, 239
 - due to use of log periodogram, 304–5, 307
- bootstrapping and, 615–6, 631
- computation of, 219–24
- covariances, 203–4, 235
 - verification via simulation, 242
 - subtle issues, 214
- DCT-based, 217–9, 244
- deficiency of power after removal of lines, 550–1
- definition, 170, 232
 - for complex-valued time series, 231
- discretely smoothed (*see* discretely smoothed periodogram)
- distribution of, 203, 235, 307–8, 408
 - verification via simulation, 242
- effect on periodogram of
 - centering of time series, 184, 213, 240
 - at nonzero Fourier frequencies, 184
 - at zero frequency, 184
 - circular shifting of time series, 238–9
 - scaling of time series, 238
 - time indexing of time series, 184
- examples of
 - AR(2) data, 172–3, 182, 208, 216, 246, 249, 294, 305, 310
 - AR(4) data, 172–3, 182, 199, 508
 - atmospheric CO₂ data, 335
 - atomic clock data, 327–8
 - chaotic beam data, 228
 - ice profile data, 323
 - ocean noise data, 231
 - ocean wave data, 226–7, 497
 - rotation of earth data, 224
 - harmonic process plus white noise, 533–4, 536, 551
 - ship altitude data, 331
 - three sinusoids (Kay and Marple), 531
 - white noise data, 207, 210
 - Willamette River data, 568, 571, 573
- expected value of, 174–5, 203, 233, 240
 - complex-valued harmonic process plus white noise, 524, 586
 - real-valued harmonic process plus white noise, 524, 536, 585
 - conditional on phase value, 537–8
 - verification via simulation, 242
- in circulant embedding simulation method, 615
- inconsistent estimator of SDF, 163–4, 207, 235
- Laplace, 529
- log likelihood, 308
- Lomb–Scargle, 528–9, 543–4, 588
- modified, 186
- power-law exponent estimation, 408
- reflecting boundary conditions, 244
- rescaled, 541, 581
- spurious peaks, 527
- sum over
 - Fourier frequencies, 237, 540
 - non-Fourier frequencies, 233, 237
- symmetry, 170
- three-point example to verify properties, 238
- truncated, 343, 504
- utility compared to multitaper spectral estimator, 403–12
- variance of, 203, 235, 241
 - for AR(1) process, 243
 - verification via simulation, 242
- phase function (phase shift function), 136
- phase-shifted Fourier transform, 383
- phases (*see* random phases)
- pilot analysis (filters for), 160
- pointwise convergence, 50, 53, 84
- Poisson's formula, 84
- polynomial regression model, 522
- positive
 - definite, 28, 170, 375, 380, 422, 450–2, 454, 457, 461, 464, 480
 - semidefinite, 28, 30, 167, 170, 374–5, 457
- positive-part function, 541, 636, 640
- postcoloring, 198–201, 234, 496, 615
 - AR model approach, 491–2, 510
 - atomic clock data, 329, 339–40
 - ocean wave data, 497–500
- power
 - of a nonperiodic function, 55
 - of a periodic function, 50
- power spectral density function (PSDF), 113, 116
- power-law exponent, 403, 407–12
 - estimation using linear regression and multitaper estimator, 409
 - atomic clock data, 430
 - periodogram, 408
 - periodogram versus multitaper estimation, 410–11
- power-law process, 327–30, 336–7, 403, 407, 430–1
 - ACVS, 391, 434, 605
 - spectral density function (SDF), 391, 434, 605, 610
- predictability of a time series, 17, 403–4, 471–2
- prediction error
 - backward, 455
 - forward, 453, 596
- prediction error filter, 198
- prewhitening, 179, 197–201, 447, 491–2, 503, 506, 615
 - versus multitaper spectral estimator, 424
 - ocean wave data, 496–8, 510
- prewhitening filter, 197, 234, 241, 447, 482, 491–3, 496, 510, 615
- principle of maximum entropy, 447, 471, 473
- probability density function (PDF), 24
 - bivariate standard Gaussian, 634–5
 - chi-square with two degrees of freedom, 37
 - exponential (scaled chi-square with two degrees of freedom), 37, 209
 - gamma (scaled chi-square with arbitrary degrees of freedom), 297

- generalized Gaussian (GG), 633
- Gaussian (normal) with zero mean, 55
- modeling, 631
- spectral window as an example of, 192
- uniform
 - over interval $(-\pi, \pi]$, 37, 45, 598
 - over interval $(-\sqrt{3}, \sqrt{3}]$, 634
- probability integral transform, 634
- probability of an event, 23
- process
 - band-pass, 299, 347
 - high-pass, 299, 347
 - low-pass, 299, 347
 - residual, 550
 - with stationary differences of order d , 340
- profile likelihood, 482–4
- projections, 384, 439
- prolate spheroidal wave functions (PSWFs), 64
- proper
 - complex-valued Gaussian or normal RV, 546–7
 - complex-valued white noise, 517–8, 564
 - process, 30, 32
- PSDF (*see* power spectral density function)
- pseudo-AR process, 553
 - complex-valued, 563
 - finding the coefficients, 591
 - plus white noise is an ARMA process, 554
 - solution to defining equation, 554, 591
 - stationarity condition, 557
- SVD
 - parameter estimator, 566
 - sinusoidal frequency estimator, 567
- purely
 - acausal, 447–8
 - continuous spectrum, 120, 163, 165, 167, 511
 - continuous stationary process, 457, 532
 - discrete spectrum, 120, 511, 513
 - nondeterministic process, 404, 471
- QQ plot
 - Gaussian stationary process assumption, 621–2
 - i.i.d. Gaussian assumption, 621
- quadratic lag window (*see* Bartlett–Priestley lag window)
- quadratic spectral estimator, 374–82, 439
 - bias of
 - broad-band, 378
 - indicator, 379–81
 - minimizing indicator of, 380
 - for white noise, 377–8
 - local, 378
 - magnitude indicator, 378–9, 381
 - deconstructed multitaper spectral estimator, 376
 - examples of
 - direct spectral estimator, 375
 - lag window spectral estimator, 375
 - weighted multitaper spectral estimator, 376
 - justification for, 351
 - WOSA spectral estimator, 423
 - expected value of, 376–7, 439
 - regularization bandwidth, 377
 - variance of, 380
 - quadratic taper, 248
 - quadratic trend removal
 - by ordinary least squares (OLS), 332, 443
 - by differencing, 46, 333–4
 - quadratic window, 248
 - estimator, 252
- radar clutter, 131
- random amplitudes, 35–7, 111
- random phases, 37–8, 111
 - nonuniformly distributed, 38, 45
 - uniformly distributed, 37, 511, 553, 598, 616
- random variable, 3, 21–3
 - complex-valued, 21, 23
 - vector-valued, 21–2
- random walk, 43
 - frequency noise, 431
- randomly phased sinusoid, 35, 511, 553, 598
 - ACVS for, 553, 591
 - plus white noise, 554
- rank of matrix, 375–6, 380–1, 420, 422–3, 490, 565–6
- Rayleigh
 - distribution, 37, 598
 - quotient, 89, 105
 - theorem, 49
- real part of complex-valued
 - process, 602, 606, 640
 - variable, 61, 486
- real-valued stochastic process
 - continuous parameter/time, 22
 - stationary, 26
 - discrete parameter/time, 22
 - stationary, 26
- realizations, 3, 21–2
- reciprocity relationships, 58–62
 - equivalent width, 58–9
 - fundamental uncertainty relationship, 60–2
 - similarity theorem, 58–60
- rectangular
 - data taper, 188–90, 535
 - kernel, 71, 290
- recursive maximum likelihood autoregressive
 - parameter estimator (RMLE), 485
- reduced (profile) likelihood, 482–4
- reflecting boundary conditions, 181, 183, 216, 219, 244, 424
- reflection coefficient (*see* partial autocorrelation coefficient)
- regularization, 377, 386
 - bandwidth, 357, 377, 425, 428, 434
 - versus spectral window bandwidth, 402
- rejection filtration, 201
- relation function, 31
- relative efficiency, 165
- remodulated filtered series, 592
- repeated lag window spectral estimator, 349–50
- rescaled periodogram, 541, 581
- reshaped
 - lag window, 257, 281–2, 375
 - spectrum, 547, 551, 575, 582, 592
 - Willamette River data, 575

- residual process (EULS), 550
 - periodogram of
 - harmonic process plus white noise, 551
 - Willamette River data, 570
- resistor, 21, 40–1
- resolution
 - loss of, 76, 78, 271, 290, 357, 359, 400, 434, 535
- restoration of power, 195
- Riemann integral, 24, 26, 112–13, 258, 261
 - approximation by a Riemann sum, 174, 254, 258–9, 404, 605–6
- Riemann sum, 174, 254, 258–9, 404, 605–6
- Riemann–Lebesgue lemma, 120, 167
- Riemann–Stieltjes integral
 - ordinary, 24–6, 109, 112, 127
 - stochastic, 110, 112, 127
- ringing, 196, 225, 227, 277, 320, 331
- ripples, 72, 77, 79, 152–3, 189, 253, 275, 290, 294
- robust estimation of SDF via WOSA, 419
- rotation of the earth, 148, 222, 366, 407
- rule of thumb for significance levels, 548, 574, 582
- sample autocorrelation, 3
- sample autocorrelation sequence (sample ACS), 3
 - examples, 4
 - correlation of, 4
 - variance of, 4
- sample mean, 3, 164–6, 184, 195–6, 213
 - variance for AR(1) process, 507
- sampled function, 74
- sampling and aliasing, 122–3, 569
- sampling interval, 84–5, 113, 119, 122–5, 142–3, 163, 207, 232, 352, 374, 432, 445, 505, 511
 - computational details, 220–4, 314–5
 - irregular, 382, 528–9, 543
 - jitter, 124
 - effect on oxygen isotope time series, 124
 - oversampling, 123
 - role in simulation of time series, 594, 601, 609, 613
 - subsampling, 124
 - example, 130–1
- scale preservation (LTI filters), 133, 140
- scatter plot
 - lag 1 and four examples, 2–3
 - lag τ , 2, 5
 - $\tau = 6$ and 9 examples, 6
 - locations of peak frequencies from estimated spectra, 537–8, 556–7, 589, 591
- Schur matrix, 484
- Schuster's test for simple periodicity, 539
- Schwarz inequality, 55, 57, 61, 104, 257
- SDF (*see* spectral density function)
- sea levels, 520
- seasonality/periodicity, 40, 332
 - removal by differencing, 40, 42, 333
- second differencing for quadratic trend removal, 46
- seismic trace of earthquake, 58
- semiperiodogram, 217
- semivariance, 32, 546–7
- sequency (Walsh functions), 138
- sequency-limited and time-limited, 140
- shading sequence, 186
- Shannon number, 66–7, 90, 357, 372–4, 381
- ship altitude data, 330–2, 626–8
 - analysis of longer series, 350
 - direct spectral estimator, 331
 - lag window spectral estimator, 331
 - periodogram, 331
 - plot of, 330, 627
 - simulation of, 626–8
- sidelobe leakage, 77, 185, 386, 388, 390
- sidelobes
 - of smoothing windows, 268–9, 273, 275–8
 - of spectral windows, 281–2
 - Fejér's kernel, 178–80, 328, 527, 532, 535–6
- Siegel's test for compound periodicity, 540–3, 581, 587
 - asymptotic critical value, 542–3, 581
 - exact critical value, 541, 581
 - interpolated critical value, 542, 581
- sign of i convention, 55
- signal-to-noise ratio, 526, 532, 543, 552, 560–1, 567, 578, 587
 - signal in, 526
- similarity theorem, 58–60
- simple periodicity, 540–1, 586
- simulation of
 - atomic clock data, 628–30
 - autoregressive (AR) process, 595
 - zero initial conditions, 595
 - fractionally differenced (FD) process, 440
 - Gaussian AR(1) process
 - via Kay's method, 639
 - Gaussian AR(2) process, 597, 602
 - via circulant embedding, 603
 - via Gaussian spectral synthesis method (GSSM), 608
 - via Kay's method, 597
 - examples, 34
 - Gaussian AR(4) process, 597
 - via circulant embedding, 640–1
 - via Gaussian spectral synthesis method (GSSM), 606, 641
 - via Kay's method, 597, 638
 - examples, 34
 - Gaussian ARMA process, 593–4
 - via Kay's method, 598, 639
 - via McLeod and Hipel's method, 600, 639
 - Gaussian autoregressive (AR) process
 - stationary initial conditions, 595, 599, 624, 637
 - via Kay's method, 597
 - via McLeod and Hipel's method, 599
 - harmonic process, 593–4, 637
 - moving average (MA) process, 594, 637, 639
 - MA(1) examples, 33
 - non-Gaussian time series
 - ARMA, 631, 636, 638
 - with specified ACVS and PDF, 631, 633, 638

- ocean wave data, 619–26
 - uses, 626
- reduced-length time series
 - based on nonparametric SDF estimate, 614
- ship altitude data, 626–8
- time series, 593
 - based on
 - nonparametric SDF estimate, 611, 619–20, 627, 629, 637
 - with non-unity sampling interval, 613, 638
 - method of surrogate time series (MSTS), 615–6
 - parametric SDF estimate, 617, 638
 - periodogram, 615
 - centering, 619, 627–8
 - Gaussianity testing, 621
 - given a portion of its ACVS, 593, 601, 637
 - given its fully specified SDF, 593, 604, 637
 - role of sampling interval, 594, 609, 613
 - summary, 637–8
- white noise, 594, 637
 - examples, 31
- simulation via
 - circulant embedding, 601–4, 619–20, 627, 629, 631
 - ACVS properties, 602, 640
 - computational efficiency, 603
 - issues with ACVS, 615
 - overcoming negative weights, 604, 640
 - weights, 601–2, 605, 629, 640–1
 - Gaussian spectral synthesis method (GSSM), 605–6, 608
 - adjustment for power-law process, 610
 - and adaptive multitaper spectral estimator, 615
 - and prewhitening, 615
 - role of sampling interval, 609
 - weights, 640–1
 - method of surrogate time series (MSTS), 615, 641
 - relation of simulated process to harmonic process, 641
- sinc function, 63, 559
- single-sided versus double-sided SDF, 118
- singular value decomposition (SVD), 425, 564–5
 - and short time series, 567
 - approach to harmonic analysis, 563
 - Willamette River data, 578–80, 592
- singular value, 565
- singular vector, 565
- sinusoidal amplitude estimator
 - approximate conditional least squares (ACLS), 515, 525, 549, 569–70, 572–3, 585, 589
 - approximate least squares, 516, 585
 - exact conditional least squares (ECLS), 549, 569–70, 572, 589–90
 - including mean, 552
 - exact least squares
 - complex-valued, 517, 586
 - real-valued, 515, 525, 585
 - exact unconditional least squares (EULS), 549, 569–70, 572
 - including mean, 552
 - Laplace method, 529
 - Lomb–Scargle method, 528
 - multitaper (complex-valued), 546, 551, 574
 - sinusoidal frequency estimator
 - exact unconditional least squares (EULS), 549, 569–70, 572
 - including mean, 552
 - location of maximum (global or local) of AR SDF, 556–8, 591
 - forward and backward least squares, 556
 - direct spectral estimator, 537–8, 556–7, 585, 589
 - rationale for tapering, 535–7
 - periodogram, 526–8, 537–8, 589
 - at Fourier frequencies, 523–4, 585–6
 - at general frequencies, 524–5, 585
 - expected value of estimator, 526
 - convergence of estimator, 527
 - MSE of estimator, 526, 537
 - variance of estimator, 526, 534
 - Thomson's F -test, 574, 582–3
 - parametric approach, 553–8
 - problems with, 558–63
 - three-sinusoids example
 - noise-free (Kay and Marple), 530
 - effect of grid size, 530–2
 - with noise, 532
 - effect of noise variance, 533–4
 - effect of series length, 532, 534
- sinusoidal multitapers, 355, 372, 391, 393–4, 396, 434, 544, 574, 583, 590, 592
 - compared to Slepian multitapers, 393
 - defined, 392
 - equivalent width, 401, 441
 - versus spectral window bandwidth, 402
 - relationship to minimum-bias multitapers, 392
- sinusoidal power estimation (AR SDF), 578
- sinusoidal variations
 - deterministic, 553–4, 558, 590
 - expressible as AR model, 447
 - inexpressible as MA model, 503
 - randomly phased, 35, 511, 553, 561
 - ACVS for, 37, 553, 591
 - examples of realizations, 36
 - plus white noise, 512, 554
 - as ARMA process, 554
 - sum of, 6–7
- skewness, 631
 - ARMA(1,1) process, 632
 - inequality, 633, 641
- Slepian
 - data taper, 190–1, 206, 208, 225, 228, 234, 260, 263, 282, 287, 416, 461, 496, 535–6, 569, 591
 - approximation of zeroth-order, 196
 - lag window, 285, 345
 - multitapers, 355, 360, 362, 372, 391, 434, 544, 574, 582, 590, 592
 - compared to sinusoidal, 393
 - decomposition of energy, 367
 - equivalent width versus spectral window bandwidth, 369
 - sequence, 87, 105, 145, 189, 357
 - double orthogonality, 106
 - zeroth-order as a low-pass filter, 155

- smoothing of
 - direct spectral estimator, 245
 - log direct spectral estimator, 303
 - periodogram estimator, 246, 307
- smoothing window, 245, 248, 341
 - autocorrelation width of, 251
 - bandwidth, 245, 251, 268, 281, 292, 296, 300–1, 318, 322, 341, 346–7
 - Bartlett–Priestley design, 344
 - choice of, 252
 - finer frequency grid, 255
 - for sea level residuals, 522
 - Grenander, 251, 268, 279, 281
 - lag window spectral estimator, 279
 - Parzen, 251
 - repeated lag window estimator, 349–50
- bias due to, 256, 290
- equivalent width of, 251
- leakage, 268, 287–9, 294, 316, 318, 331, 498
- parameter m (controls smoothing), 247, 287
 - choice of, 291, 304, 313–4
 - examples of, 318, 322, 328, 330, 339
- properties of, 250
- type
 - Bartlett, 269–70
 - Bartlett–Priestley, 274–5
 - Daniell (rectangular), 250, 272–3
 - Gaussian, 277–8
 - modified Daniell, 280
 - Papoulis, 278–9
 - Parzen, 275–6, 280–2, 344
 - rectangular, 271
- spectral bandwidth, 292, 296, 346
- spectral density function (SDF), 111
 - after first difference and seasonal filtering, 333–4
 - aliasing effect, 122–3
 - alternative definitions, 114–6
 - and LTI digital filters, 143
 - and subsampling, 130
 - asymmetry for complex-valued process, 118
 - basic properties, 116–7
 - classification examples, 121
 - confidence interval for, 205, 265–7, 342
 - for log of, 205, 227, 229, 266–7, 342
 - double-sided versus single-sided, 118
 - dynamic range, 177, 184, 197, 492, 641
 - for first difference of a process, 158, 326
 - for particular stationary processes
 - AR, 145, 446, 485, 609, 617–8
 - AR(1), 336
 - AR(2), 172–3, 609
 - AR(4), 172–3
 - computation using zero padding, 449
 - ARMA, 145, 159
 - band-limited white noise, 379
 - fractionally differenced (FD), 166, 407–8
 - MA, 130, 144
 - MA(1), 130
 - MA(2), 159
 - power-law, 605, 610
 - white noise, 117
- Fourier transform of ACVS, 111, 114, 126
- imputed from noise model, 336
- linear and log scales, 336–8
- log of, 301
- one-sided versus two-sided, 118
- nonparametric estimators of
 - direct spectral estimator, 186
 - discretely smoothed, 246–7
 - lag window estimator, 248
 - multitaper, 352
 - periodogram, 170
 - discretely smoothed, 307
 - WOSA, 414
 - robust estimation via, 419
- parametric estimators of
 - AR, 446–7
 - Burg, 466
 - least squares, 475
 - maximum likelihood, 480
 - Yule–Walker, 451
 - MA, 503
- single-sided versus double-sided, 118
- symmetry for real-valued process, 111–12, 118, 126
- two-sided versus one-sided, 118
- type
 - Gaussian-shaped, 131
 - Lorenzian, 131
 - piecewise constant, 348
 - rectangular, 128
 - triangular, 128
- unimodal, 297
- units of (physically meaningful), 374
- variance-preserving, 337–9
- versus autocovariance sequence, 124–5
- weighted sum of uncorrelated processes with SDFs, 130
- spectral distribution function, 117
- spectral estimator of the Grenander–Rosenblatt type, 252
- spectral factorization, 636
- spectral lines, 518, 538–40, 543–4, 548–9
 - splitting of, 558, 560
- spectral representation
 - interpretation, 110–11
 - of stationary process, 108
 - complex-valued, 114
 - continuous parameter, 113–4, 135
 - discrete parameter, 110, 125
 - nonzero mean, 113
 - with discrete spectra, 518–9
 - with mixed spectra, 518–9
 - theorem, 7, 108–10, 382, 518–9
 - uniqueness, 110
- spectral reshaping, 547, 551, 575, 582
- spectral window, 175, 186–8, 255, 353, 414
 - bandwidth, 192, 194, 213–4, 355
 - definitions of
 - autocorrelation width (standard), 194
 - correlation time, 213–4, 355
 - half-power width, 192
 - standard (autocorrelation width), 194
 - variance width, 192, 194
 - definitions (standard) for spectral estimators
 - direct, 194, 235
 - discretely smoothed periodogram, 347

- lag window, 256, 341
- multitaper, 353
- WOSA, 415
- frequency range for statistical results, 206, 213, 244, 268
- standard measure of, 194
 - examples of use, 226–7, 229, 294, 317–8, 323, 325, 328, 331, 335, 568, 623–4
 - versus regularization bandwidth, 358, 402
 - versus smoothing window bandwidth, 296
 - versus correlation time, 214, 368–9, 401–2
- bias, 192, 391
 - and minimum-bias multitapers, 391–2
 - due to central lobe (spectral window bias), 194, 535
 - due to sidelobes (leakage), 194, 535
- decay rate, 281, 328
- definitions of for spectral estimators
 - direct, 186–8, 234
 - lag window, 255, 341
 - multitaper, 353
 - periodogram, 175
 - WOSA, 414
- type
 - autocorrelated 100% cosine lag window, 285
 - Bartlett lag window, 268, 270
 - Bartlett–Priestley lag window, 268, 274
 - Bartlett–Priestley taper, 288
 - Daniell (rectangular) lag window, 268, 272
 - Daniell taper, 288
 - Gaussian lag window, 268, 278
 - Gaussian taper, 288
 - Hanning taper, 191
 - modified Daniell window, 280
 - 100% cosine lag window, 284
 - Papoulis lag window, 268, 279
 - Papoulis taper, 288
 - Parzen lag window, 268, 276
 - $p \times 100\%$ taper, 191
 - periodogram, 176
 - Slepian lag window, 286
 - Slepian multitapers, 358, 364–5
 - k th-order taper, 361, 363, 368
 - Slepian taper (zeroth-order), 191, 288
 - sinusoidal multitapers, 398–401
 - k th-order taper, 395, 397
 - standard bandwidth approximation, 400–1
- spectrograph estimator, 252
- spectrum, 107–8
 - classification of, 120–2
 - discrete, 120, 511, 513, 518, 538, 584, 586
 - for simple time series model, 5–11
 - constant, 11
 - estimated
 - nonparametric, 11–13
 - parametric, 14–5
 - high frequency, 9–10
 - low frequency, 9–10
 - theoretical, 9–11
 - integrated, 120
 - interpretation via band-pass filtering, 147–8
 - line, 120, 511, 522
 - mixed, 120, 511, 513, 518, 521–2, 539, 584–5
 - purely
 - continuous, 120, 165, 167
 - discrete, 120, 511
 - spectrum analyzers, 419, 444
 - spinning rotor, 39
 - spontaneous line splitting (*see* spectral lines)
 - spurious peaks in
 - FBLs pseudo-AR sinusoidal frequency estimator, 567
 - periodogram, 527
 - square integrable, 48, 53–4, 75
 - squared norm, 470
 - St. Paul temperature data, 5–6
 - standard frequencies (Fourier frequencies), 8, 91, 515
 - stationarity
 - conditions for
 - autoregressive (AR) process, 145, 446–9, 557
 - pseudo-AR process, 557
 - second-order/weak/covariance, 26–7
 - strict/complete/strong, 26, 631, 634
 - stationary differences of order d
 - process with, 340
 - stationary process, 4, 7, 21, 26–31, 107
 - and modeling data, 38–41, 45
 - examples of
 - autoregressive (AR), 33
 - autoregressive moving average (ARMA), 35
 - bandlimited white noise, 379
 - harmonic, 35
 - fractionally differenced, 166
 - moving average (MA), 32
 - white noise, 31
 - intrinsically of order d , 340
 - nonergodic, 128
 - reverse of conjugate of complex-valued process, 502
 - reverse of real-valued process, 455, 477, 484
 - step function, 24, 26, 120
 - step-down Levinson–Durbin recursions, 460–1
 - and causality assessment, 453, 509
 - and generation of ACVS for an AR process, 460–1, 508
 - and simulation of ARMA process, 594–8
 - stochastic continuity, 114, 118
 - stochastic process, 21–2, 23–5
 - jump, 109
 - orthogonal, 109
 - storm surges, 522
 - strict stationarity, 26, 631, 634
 - sum of squares (SS), 467, 476, 514
 - summability
 - absolute, 165
 - Cesàro, 78, 165
 - superposition (LTI filters), 133, 140
 - SVD (*see* singular value decomposition)
 - symmetry, 27
 - conjugate, 29–30, 48, 54
 - Szegő–Kolmogorov theorem, 404
- tables
 - AR-based estimates of innovation variance, 500

- Burg's algorithm and sinusoidal frequency estimation, 559
- characteristics of AR SDFs and periodogram for Willamette River data, 578
- comparison of
 - spectral bandwidth measures for direct spectral estimator, 214
 - spectral bandwidth measures for sinusoidal multitaper spectral estimator, 402
 - spectral bandwidth measures for Slepian multitaper spectral estimator, 369
 - window bandwidth measures for discretely smoothed direct spectral estimator, 255
 - Yule-Walker AR(4) and AR(8) coefficient estimates, 459
- exact and approximate critical values for Siegel's test, 542
- harmonic analysis and centering for Willamette River data, 583
- horizontal and vertical components of crisscrosses (ice profile data), 325
- multitaper-based estimates of innovation variance, 432
- NMSE criteria for GSSM-based approximations for AR(4) process, 607
- parameter estimates in harmonic analysis for Willamette River data, 570
- properties of lag windows, 279
- ratio of variances for estimators of power-law exponent, 411
- reduced likelihoods for estimated AR(4) coefficients, 483
- true and estimated sinusoidal frequencies, 533
- variance inflation factor for tapers, 260
- taper, 186
 - compared to lag window, 282–3
 - advantages/disadvantages in use of
 - better performance of Yule-Walker autoregressive spectral estimator, 452, 461
 - decrease in bias due to sidelobes, 179, 185, 189, 234, 535–6
 - decrease in effective sample size, 194–5
 - decrease in pairwise correlation, 207
 - increase in bias due to central lobe, 192, 535
- converting
 - taper into lag window, 283
 - lag window into taper, 285
- linear, 186
- normalization
 - unit energy, 188
 - data-dependent, 195
- orthogonality, 354, 357, 392, 432
- plethora of types, 196–7
- quadratic, 248
- type
 - Bartlett–Priestley, 287
 - Daniell, 287
 - default, 188, 190
 - Gaussian, 287
 - Hanning, 189–90, 230, 234, 327, 331, 415–7
 - Hanning (alternative definition), 240
 - k th-order minimum-bias, 392–3
 - k th-order sinusoidal, 392–4, 396
 - k th-order Slepian, 357, 360, 362, 368
 - minimum-bias multitapers, 392–3
 - $p \times 100\%$ cosine, 189–90, 260
 - Papoulis, 287
 - Parzen, 287
 - rectangular, 188–90, 535–6
 - sinusoidal multitapers, 355, 392–3, 402–3
 - Slepian, 190, 196, 212, 225, 228, 230, 234, 260, 263, 287, 416, 461, 496, 535–6
 - Slepian multitapers, 355, 357, 372–3
 - variance inflation factor, 260, 262, 342, 417
- temporal window, 186
- terrestrial free oscillations, 359, 366
- test for
 - periodicity
 - Fisher's, 540–1, 586–7
 - in colored noise, 544, 587
 - in white noise, 538, 586–7
 - Schuster's, 539
 - Siegel's, 541, 587
 - Thomson's F -test (multitaper approach), 228, 359, 544–8, 574–5, 582–3, 587, 592
 - with irregular sampling, 543–4
 - white noise (cumulative periodogram), 215–6, 230–1, 573–4
- textile slivers, 33, 503
- Thomson's F -test for periodicity, 228, 359, 544–8, 574–5, 582–3, 587, 592
- tidal
 - analysis, 40, 520–3
 - frequencies, 521
- time indexing of a series
 - and periodogram, 184
- time invariance (LTI filters), 133, 140
- time series simulation, 593
 - based on
 - nonparametric SDF estimate, 611, 619–20, 627, 629, 637
 - with non-unity sampling interval, 613, 638
 - method of surrogate time series (MSTS), 615–6
 - parametric SDF estimate, 617, 638
 - periodogram, 615
 - centering, 619, 627–8
 - examples of,
 - atomic clock data, 628–30
 - ocean wave data, 619–26
 - uses, 626
 - ship altitude data, 626–8
 - for testing of Gaussianity, 621
 - given a portion of its ACVS, 593, 601, 637
 - given its fully specified SDF, 593, 604, 637
 - non-Gaussian
 - ARMA process, 631, 636, 638
 - with specified ACVS and PDF, 631, 633, 638
 - reduced-length
 - based on nonparametric SDF estimate, 614
 - role of sampling interval, 594, 609, 613
 - summary, 637–8
- time-limited function, 58
- time reverse of
 - conjugate of complex-valued stationary process
 - is stationary with same ACVS, 502

- real-valued stationary process is stationary with same ACVS, 455, 477, 484
- time-reversibility, 484
- Toeplitz
 - matrix, 29, 156, 457, 462, 480, 484
 - Hermitian, 30
 - system of equations, 452, 461
- trace of a matrix, 376, 378, 439
- trade-off
 - between bias and variance, 80, 296, 309
 - between fidelity and resolution, 80
- transfer function, 136, 142–3, 290
 - and
 - gain function, 136
 - phase function, 136
 - as Fourier transform of impulse response sequence, 143
 - for
 - AR filter, 144–5, 487
 - Hilbert transform, 579, 592
 - MA filter, 144
- transpose of vector, 28
- trend, 40, 151, 332
 - removal, 39, 334, 443
 - as high-pass filtering, 151
- tridiagonal matrix, 88, 95
- trigamma function, 296, 302, 405, 410
- truncated periodogram, 343, 504
- twin peaks, 76–8, 178–9, 192, 200–1, 218–9, 291, 294, 300–1, 306, 337, 382, 458–9, 489
- two-sided versus one-sided SDF, 118
- uncorrelated random variables
 - definition, 31
 - relationship to independent random variables, 31–2
- undersmoothing, 71
- uniform distribution
 - over interval $(-\pi, \pi]$, 37, 45, 598
 - over interval $(-\sqrt{3}, \sqrt{3}]$, 634
- uniqueness of
 - Fourier transform, 55, 82
 - integrated spectrum, 135
 - spectral representation, 110
 - stationary solution to AR equation, 144–5, 446–8
- unit circle, 446, 448–9, 463, 553, 557–8, 566–7, 580, 591–2, 600, 618
- variability of SDF estimators
 - assessment via
 - asymptotic theory, 204–5, 265–7, 342, 354, 485–91
 - simulation, 622–6
- variance
 - definition of
 - complex-valued case, 25
 - real-valued case, 24
- variance inflation factor due to tapering, 260, 262, 342, 417
- variance of
 - linear combination of random variables, 41
 - sample mean for AR(1) process, 507
- sinusoidal frequency estimator using periodogram maximum, 526–7, 534, 592
- spectral estimators
 - basic multitaper, 353–4, 359
 - for Gaussian white noise, 372–4, 402–3
 - direct, 204, 235, 624
 - for complex-valued process, 232
 - verification via simulation, 242
 - lag window, 258–9, 262, 265, 279, 341
 - periodogram, 203, 235, 241
 - for AR(1) process, 243
 - verification via simulation, 242
 - quadratic, 380
 - weighted multitaper, 354
 - WOSA, 415, 435
- stationary process, 27–8
 - bias in estimator (unknown mean), 168, 237
- variance-preserving SDF, 337–9
- variance spectrum for
 - harmonic process, 108
 - simple time series model, 8, 47
- variance-stabilizing transform, 13, 209
- variance width of function, 60, 73, 105, 191–2, 194, 288
- vector transposition, 28
- Walsh functions, 138–40
- weighted covariance estimator, 252
- weighted multitaper spectral estimator, 352–4, 388, 433, 611–13
 - and quadratic spectral estimator, 376
 - bias of, 391
 - covariances of, 437
 - distribution of, 354–5
 - equivalent degrees of freedom (EDOFs) of, 354–5
 - equivalent width, 355, 437
 - expected value of, 352, 389
 - in circulant embedding simulation method, 611–13
 - low-rank approximation, 422–3
 - Parseval's theorem in expectation, 390
 - spectral window for, 352–3, 391
 - bandwidth of, 353
 - variance of, 354
 - via regularization, 382
- weighted overlapped segment averaging (*see* WOSA spectral estimator)
- weights for circulant embedding simulation method (*see* circulant embedding simulation method)
- Welch's overlapped segment averaging spectral estimator (*see* WOSA spectral estimator)
- white noise, 31–2, 120, 169, 184, 197, 236, 381, 386–8, 446, 450, 511, 513, 524, 532, 534, 538, 586
 - and direct spectral estimator, 189, 371
 - and periodogram, 177, 207, 371
 - and quadratic spectral estimator, 377
 - and tapering, 195
 - autocovariance sequence (ACVS), 31
 - band-limited, 379
 - complex-valued, 32

- and proper, 517–8, 564
- continuous parameter, 38
- examples of, 31
- Gaussian, 201, 539
- spectral density function (SDF), 117
- simulation of, 594, 637
- test for (cumulative periodogram), 215, 230–1
- versus colored noise, 120, 126, 241, 372
- white noise variance estimator (harmonic analysis)
 - approximate conditional least squares (ACLS), 549, 570
 - exact conditional least squares (ECLS), 549, 552, 570
 - exact least squares
 - complex-valued, 518
 - real-valued, 516
 - exact unconditional least squares (EULS), 550, 552, 570
 - multitaper, 551
- Whittle likelihood, 307
- Wiener–Khinchine theorem, 118
- Willamette River data, 1–2, 567–8
 - AICC order selection, 575–6
 - analytic series version, 578–81, 592
 - Burg autoregressive spectral estimator, 575–7
 - cumulative periodogram test of residuals, 573–4
 - direct spectral estimator, 569, 591–2
 - FBLS autoregressive spectral estimator, 592
 - fitted values, 571–2
 - harmonic analysis of, 567–81
 - effect of centering, 583
 - integrated autoregressive spectrum, 577
 - multitaper spectral estimator, 575
 - parametric approach to harmonic analysis of, 575–8
 - periodogram, 568
 - of residuals, 570–1, 573
 - plot of
 - complex roots for, 580
 - fitted model, 571–2
 - residuals, 571–2
 - remodulated filtered series, 592
 - reshaped spectrum, 575, 592
 - residuals, 571–2
 - small portion of, 1–2
 - ACS for
 - sample, 4
 - theoretical, 10
 - lag 1 scatter plot for, 3
 - sample ACVS for, 169
 - simulation of, 11
 - spectrum for
 - estimated, 12, 14
 - theoretical, 10, 12, 14
 - SVD approach to harmonic analysis of, 578, 592
 - table of characteristics of AR SDFs and periodogram, 578
 - Thomson's F -test for periodicity, 575, 592
 - Yule–Walker autoregressive spectral estimator, 592
- wind speed data, 1–2
 - ACS for
 - sample, 4
 - theoretical, 10
 - lag 1 scatter plot for, 3
 - sample ACVS for, 169
 - simulation of, 11
 - spectrum for
 - estimated, 12, 14
 - theoretical, 10, 12, 14
- window
 - data, 186
 - design, 247
 - for Bartlett–Priestley, 273–4, 319, 344
 - for Daniell, 272, 319
 - lag (*see entry for* lag window)
 - linear, 186
 - quadratic, 248
 - smoothing (*see entry for* smoothing window)
 - spectral (*see entry for* spectral window)
 - temporal, 186
- window closing, 293
 - and WOSA, 416
 - examples, 293–5
- windowed periodogram estimator, 252
- windows and convergence factors, 80
- Wold's decomposition theorem, 404, 471
- Wold's theorem, 117–8, 126, 185
- WOSA spectral estimator, 351, 412–19, 432, 434–6
 - ACVS estimator for, 420, 435
 - AR(2) data, 416–7
 - AR(4) data, 417–8
 - at zero frequency, 420
 - boundary conditions
 - circular, 424
 - reflecting, 424
 - centering for unknown mean, 419–20
 - computational efficiency, 419
 - connection to Bartlett lag window spectral estimator, 270
 - deconstructed multitaper spectral estimator, 420–1, 423
 - EDOFs of, 415–6, 423–4, 442–3
 - expected value of, 414, 435
 - for approximating a multitaper spectral estimator, 424–5
 - in circulant embedding simulation method, 611, 613
 - ocean wave data, 425, 429–30
 - robust SDF estimation, 419
 - shift factor restriction, 420
 - variance of, 415, 435
 - versus multitaper spectral estimator, 414, 417–9, 424
- Yule–Walker autoregressive estimator, 449–52, 505
 - and ACVS for direct spectral estimator, 452, 461–2, 505
 - AR(1) process, 510
 - AR(2) numerical example, 507–8
 - AR(4) data, 458–9, 461–2, 508–9
 - assessing variability (SDF), 485
 - causality, 448, 505

- complex-valued process, 502
- corresponding ACVS, 451, 504, 508
- FPE criterion, 494
- ocean wave data, 496–9
 - order selection via PACS, 498–9
- order of, 458–9
- peak shifting and splitting, 560–2, 591
- relation to
 - autocorrelation method, 480
 - Burg's algorithm, 471
 - forward least squares, 479
- Willamette River data, 592
- Yule–Walker equations, 450, 561
 - arising from two related problems, 454–5
 - augmented, 452
 - Levinson–Durbin recursions and, 457
- zero padding, 93–4
 - use in computing AR SDF, 449
 - finer frequency grid, 179, 527, 558, 568
 - for noncyclic convolutions, 106, 169–70, 220
- zero-memory nonlinearity (ZMNL), 634
 - correlation distortion, 635–6

