

HIGHLIGHTS OF PRESCRIBING INFORMATION: AMOXICILLIN CAPSULES, USP

1 INDICATIONS AND USAGE

Amoxicillin is a penicillin-class antibacterial indicated for treatment of infections due to susceptible isolates of designated microorganisms:

- Infections of the ear, nose, and throat (e.g., otitis media, pharyngitis, tonsillitis due to *Streptococcus* species).
- Infections of the genitourinary tract (e.g., *Escherichia coli*, *Proteus mirabilis*, *Enterococcus faecalis*).
- Infections of the skin and skin structure (e.g., *Streptococcus* spp., *staphylococci*, *E. coli*).
- Infections of the lower respiratory tract (e.g., *Streptococcus pneumoniae*, *Haemophilus influenzae*).
- *Helicobacter pylori* eradication to reduce the risk of duodenal ulcer recurrence in triple or dual therapy regimens.

2 DOSAGE AND ADMINISTRATION

- Adults: 250 mg to 500 mg every 8 hours or 500 mg to 875 mg every 12 hours depending on infection severity.
- Pediatric Patients (aged > 3 months): 20 to 45 mg/kg/day in divided doses every 8 or 12 hours.
- Neonates and Infants (aged ≤ 12 weeks): Maximum dose 30 mg/kg/day divided every 12 hours.
- Renal Impairment: For GFR 10 to 30 mL/min, administer 250 mg or 500 mg every 12 hours. For GFR < 10 mL/min, administer 250 mg or 500 mg every 24 hours.

4 CONTRAINDICATIONS

Amoxicillin is contraindicated in patients who have experienced a serious hypersensitivity reaction (e.g., anaphylaxis or Stevens-Johnson syndrome) to amoxicillin or to other beta-lactam antibacterial drugs (e.g., penicillins and cephalosporins).

5 WARNINGS AND PRECAUTIONS

- Anaphylactic Reactions: Serious and occasionally fatal hypersensitivity (anaphylactic) reactions have been reported in patients on penicillin therapy. Carefully inquire regarding previous hypersensitivity before initiating therapy.
- *Clostridioides difficile*-Associated Diarrhea (CDAD): Reported with nearly all antibacterial agents, ranging in severity from mild diarrhea to fatal colitis. If CDAD is suspected or confirmed, ongoing antibiotic use not directed against *C. difficile* may need to be discontinued.
- Development of Drug-Resistant Bacteria: Prescribing amoxicillin in the absence of a proven or strongly suspected bacterial infection is unlikely to provide benefit and increases the risk of drug-resistant bacteria.

6 ADVERSE REACTIONS

The most frequent adverse effects are:

- Gastrointestinal: Nausea, vomiting, diarrhea, and black hairy tongue.
- Hypersensitivity Reactions: Erythematous maculopapular rashes, urticaria, serum sickness-like reactions.
- Hematologic: Anemia, thrombocytopenia, leukopenia, agranulocytosis (usually reversible upon cessation).
- Central Nervous System: Hyperactivity, agitation, anxiety, insomnia, confusion