Music Arrangement via Quantum Annealing

Lucas Kirby 19th January 2025

Durham University

Overview

Theory

Music arrangement

Quantum annealing

Methods

Results

Conclusions

Theory

Music arrangement

- Adaptation of previously composed pieces for practical or artistic reasons
- Traditionally complex and time-consuming
- This study focuses on **reduction**



Beethoven's String Quartet No. 10

Quantum annealing

- Materials heating and cooling a material to alter its physical properties
- Quantum changing a quantum system from one Hamiltonian to another
- Done slowly and adiabatically to remain in the ground state

$$H(t) = \left(1 - \frac{t}{T}\right)H_0 + \frac{t}{T}H_p$$

QUBO

Encodes problem solution into Hamiltonian's ground state

Ising model

$$H(s) = -\sum_{i < j} J_{ij} s_i s_j - \sum_{i=1}^{N} h_i s_i$$

QUBO

Quadratic Unconstrained Binary Optimisation

$$f(x) = \sum_{i < j} Q_{i,j} x_i x_j + \sum_i Q_{i,i} x_i$$

5

How to combine them?

Methods

Problem formulation

- 1. Split parts into phrases
- 2. Arrange phrases into a graph
- 3. Solve graph problem using QPU
- 4. Construct arrangement from solution

1. Split parts


Local boundary detection model (LBDM)

$$S_i = x_i \times (r_{i-1,i} + r_{i,i+1})$$



2. Create graph







3. Solve graph

Maximal independent set (MIS)

Largest subset of nodes such that no nodes within the subset are connected by an edge.

$$f(x) = A \sum_{ij \in E} x_i x_j - B \sum_i x_i$$






Problem graph

Solution graph

4. Construct arrangement



Results


Excerpt



String Quartet No. 10 by Ludwig van Beethoven


Phrase detection

Local boundary detection model (LBDM)




Boundary strengths for the Violin I part

Problem graph



Solutions



Solutions returned by the $\ensuremath{\mathsf{QPU}}$

Example solution

Blocks

$$\iint_A E \cdot dA = \frac{Q}{\varepsilon_0}$$

The *net electric flux* through any closed surface is proportional to the **enclosed charge**.

Alert


This is an alert.

Example

This is an example.

Apperance sync

- Volume rate of flow equal to divergence
- Summed over entire volume
- Equal to net flow across the boundary



Source: Wikimedia Commons

$$\iiint_{V} \nabla \cdot \mathbf{F} \, dV = \oiint_{A} \mathbf{F} \cdot d\mathbf{A}$$

Conclusions

Equation gather

$$\nabla \cdot \mathbf{E} = \frac{\rho}{\varepsilon_0}$$

$$\nabla \cdot \mathbf{B} = 0$$

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}$$

$$\nabla \times \mathbf{B} = \frac{1}{c^2} \frac{\partial \mathbf{E}}{\partial t} + \mu_0 I$$