Music Arrangement via Quantum Annealing

Lucas Kirby

30 January 2025

Department of Physics, Durham University

Overview

Theory

Music arrangement

Quantum annealing

Methods

Results

Conclusions

Theory



Beethoven's String Quartet No. 10

 Adaptation of previously composed pieces for practical or artistic reasons



Beethoven's String Quartet No. 10

- Adaptation of previously composed pieces for practical or artistic reasons
- Traditionally complex and time-consuming



Beethoven's String Quartet No. 10

- Adaptation of previously composed pieces for practical or artistic reasons
- Traditionally complex and time-consuming
- This study focuses on **reduction**



Beethoven's String Quartet No. 10

 Materials — heating and cooling a material to alter its physical properties

- Materials heating and cooling a material to alter its physical properties
- Quantum changing a quantum system from one Hamiltonian to another

- Materials heating and cooling a material to alter its physical properties
- Quantum changing a quantum system from one Hamiltonian to another
- Done slowly and adiabatically to remain in the ground state

- Materials heating and cooling a material to alter its physical properties
- Quantum changing a quantum system from one Hamiltonian to another
- Done slowly and adiabatically to remain in the ground state

$$H(t) = \left(1 - \frac{t}{T}\right)H_0 + \frac{t}{T}H_p$$

[Lucas, 2014]

Ising model

$$H_p(\sigma^z) = \sum_{i < j}^N J_{ij} \sigma_i^z \sigma_j^z + \sum_{i=1}^N h_i \sigma_i^z$$


Ising model

$$H_p(\sigma^z) = \sum_{i < j}^N J_{ij} \sigma_i^z \sigma_j^z + \sum_{i=1}^N h_i \sigma_i^z$$

Initial state

$$H_0 = h_0 \sum_{i=1}^{N} \sigma_i^x$$

[Lucas, 2014]



[Johnson et al., 2011]

$$f(x) = \sum_{i < j}^{N} Q_{i,j} x_i x_j + \sum_{i}^{N} Q_{i,i} x_i$$

$$f(x) = \sum_{i < j}^{N} Q_{i,j} x_i x_j + \sum_{i}^{N} Q_{i,i} x_i$$

$$f(x) = \sum_{i < j}^{N} Q_{i,j} x_i x_j + \sum_{i}^{N} Q_{i,i} x_i$$

$$f(x) = \sum_{i < j}^{N} Q_{i,j} x_i x_j + \sum_{i}^{N} Q_{i,i} x_i$$

- Encodes problem solution into Hamiltonian's ground state
- Remains in low-energy state via quantum tunneling

How to combine them?

Methods

1. Split parts into phrases

- 1. Split parts into phrases
- 2. Arrange phrases into a graph

- 1. Split parts into phrases
- 2. Arrange phrases into a graph
- 3. Solve graph problem using QPU

- 1. Split parts into phrases
- 2. Arrange phrases into a graph
- 3. Solve graph problem using QPU
- 4. Construct arrangement from solution

1. Split parts

1. Split parts

Local boundary detection model (LBDM)

$$S_i = x_i \times (r_{i-1,i} + r_{i,i+1})$$

[Cambouropoulos, 2011]

1. Split parts

Local boundary detection model (LBDM)

$$S_i = x_i \times (r_{i-1,i} + r_{i,i+1})$$

[Cambouropoulos, 2011]




2. Create graph




2. Create graph





3. Solve graph




3. Solve graph

Maximal independent set (MIS)

Largest subset of nodes such that no nodes within the subset are connected by an edge

$$f(x) = A \sum_{ij \in E} x_i x_j - B \sum_i x_i$$

[Lucas, 2014]




3. Solve graph


Maximal independent set (MIS)

Largest subset of nodes such that no nodes within the subset are connected by an edge

$$f(x) = A \sum_{ij \in E} x_i x_j - B \sum_i x_i$$


[Lucas, 2014]






4. Construct arrangement






4. Construct arrangement




Results

Excerpt




String Quartet No. 10 by Ludwig van Beethoven

Phrase detection




Boundary strengths for the Violin I part

Problem graph




Problem graph with 33 nodes and 70 edges

Solutions



Returned solutions for 1000 reads

Example solution



Solution graph returning a subset of 11 nodes

Final arrangement





Selected phrases

Final arrangement



• Successful in creating a valid single-part reduction



- Successful in creating a valid single-part reduction
- Advantage over classical algorithms [Huang et al., 2012]



- Successful in creating a valid single-part reduction
- Advantage over classical algorithms [Huang et al., 2012]
- Removes skill barrier for music arrangement



• Increased problem size

- Increased problem size
- Parametric variation of LBDM

- Increased problem size
- Parametric variation of LBDM
- Physical limitations of instruments

- Increased problem size
- Parametric variation of LBDM
- Physical limitations of instruments
- Reduction to more than one part

- Increased problem size
- Parametric variation of LBDM
- Physical limitations of instruments
- Reduction to more than one part
- Quality comparison of computer arrangements [Pearce and Wiggins, 2001]



Music Arrangement via Quantum Annealing

Lucas Kirby

30 January 2025

Department of Physics, Durham University

References i

- Cambouropoulos, E. (2011).

 The Local Boundary Detection Model (LBDM) and its Application in the Study of Expressive Timing.

 International Computer Music Association.
- Huang, J.-L., Chiu, S.-C., and Shan, M.-K. (2012).

 Towards an automatic music arrangement framework using score reduction.

ACM Trans. Multimedia Comput. Commun. Appl., 8(1):8:1–8:23.

References ii

Johnson, M. W., Amin, M. H. S., Gildert, S., Lanting, T., Hamze, F., Dickson, N., Harris, R., Berkley, A. J., Johansson, J., Bunyk, P., Chapple, E. M., Enderud, C., Hilton, J. P., Karimi, K., Ladizinsky, E., Ladizinsky, N., Oh, T., Perminov, I., Rich, C., Thom, M. C., Tolkacheva, E., Truncik, C. J. S., Uchaikin, S., Wang, J., Wilson, B., and Rose, G. (2011). Quantum annealing with manufactured spins. Nature, 473(7346):194–198.

Publisher: Nature Publishing Group.

References iii

Li, Y., Wilk, C. M., Hori, T., and Sagayama, S. (2019).

Automatic Piano Reduction of Orchestral Music Based on Musical Entropy.

In 2019 53rd Annual Conference on Information Sciences and Systems (CISS), pages 1–5.

Lucas, A. (2014).

Ising formulations of many NP problems.

Frontiers in Physics, 2.

Publisher: Frontiers.

References iv

Pearce, M. and Wiggins, G. A. (2001).

Towards A Framework for the Evaluation of Machine

Compositions.

In Proceedings of the AISB'01 Symposium on Artificial Intelligence and Creativity in the Arts and Sciences.

LBDM

Boundary strength

$$S_i = x_i \times (r_{i-1,i} + r_{i,i+1})$$
$$r_{i,i+1} = \frac{|x_i - x_{i+1}|}{x_i + x_{i+1}}$$

Normalisation

$$S_i' = \frac{S_i - \min(S_i)}{\max(S_i) - \min(S_i)}$$

Weighting

$$S = rac{1}{3} \left(S'_{
m pitch} + 2 S'_{
m IOI}
ight)$$

[Cambouropoulos, 2011]

Phrase entropy

Shannon entropy

$$H(X) := -\sum_{i} P(x_i) \log_2 P(x_i)$$

Probability distribution

$$P(x_i) = \frac{n_i}{N}$$

[Li et al., 2019]