

Quantum Annealing for Music Arrangement

Lucas Kirby
4 March 2025

Department of Physics, Durham University

Quantum Annealing for Music Arrangement

2025-03-04

└ Overview

- AQC, more general umbrella term for the technique
- Quantum annealing as a subset of AQC and what that involves
- Music arrangement and why we're looking at this problem
- How the problem is solved, and the following results
- Conclusions and future work

Theory
Adiabatic quantum computing
Quantum annealing
Motivations
Music arrangement
Method
Results
Conclusions

Overview

Theory

Adiabatic quantum computing

Quantum annealing

Motivations

Music arrangement

Method

Results

Conclusions

2025-03-04


Theory

Adiabatic principle

A system remains in its instantaneous eigenstate if a given perturbation is acting on it slowly enough.¹

¹Born and Fock, 'Beweis des Adiabatensatzes'.

2025-03-04



- Adiabatic principle — system remains in the same eigenstate if perturbed slowly enough (without transferring heat)
- Equation shows evolution from initial Hamiltonian H_0 to final H_p over time T
- Importantly, if the system starts in the ground state, it will end in the ground state
- Impossible in practice as true adiabatic evolution would take infinite time, infinitely many steps

Adiabatic principle

A system remains in its instantaneous eigenstate if a given perturbation is acting on it slowly enough.¹

$$H(t) = \left(1 - \frac{t}{T}\right) H_0 + \frac{t}{T} H_p$$

¹Born and Fock, 'Beweis des Adiabatensatzes'.

Theory

- └ Adiabatic quantum computing
 - └ Adiabatic quantum computing

2025-03-04

Adiabatic quantum computing

Adiabatic principle
A system remains in its instantaneous eigenstate if a given perturbation is acting on it slowly enough.¹

$$H(t) = \left(1 - \frac{t}{T}\right) H_0 + \frac{t}{T} H_p$$

¹Born and Fock, 'Beweis des Adiabatensatzes'

- Adiabatic principle — system remains in the same eigenstate if perturbed slowly enough (without transferring heat)
- Equation shows evolution from initial Hamiltonian H_0 to final H_p over time T
- Importantly, if the system starts in the ground state, it will end in the ground state
- Impossible in practice as true adiabatic evolution would take infinite time, infinitely many steps

Adiabatic quantum computing

Adiabatic principle

A system remains in its instantaneous eigenstate if a given perturbation is acting on it slowly enough.¹

$$H(t) = \left(1 - \frac{t}{T}\right) H_0 + \frac{t}{T} H_p$$

- Universal and guaranteed

¹Born and Fock, 'Beweis des Adiabatensatzes'.

Quantum Annealing for Music Arrangement

Theory

- Adiabatic quantum computing
 - Adiabatic quantum computing

2025-03-04

Adiabatic quantum computing

Adiabatic principle
A system remains in its instantaneous eigenstate if a given perturbation is acting on it slowly enough.¹

$$H(t) = \left(1 - \frac{t}{T}\right) H_0 + \frac{t}{T} H_p$$

- Universal and guaranteed

¹Born and Fock, 'Beweis des Adiabatensatzes'

Adiabatic quantum computing

Adiabatic principle

A system remains in its instantaneous eigenstate if a given perturbation is acting on it slowly enough.¹

$$H(t) = \left(1 - \frac{t}{T}\right) H_0 + \frac{t}{T} H_p$$

- Universal and guaranteed
- A system that starts in a ground state, ends in a ground state

¹Born and Fock, 'Beweis des Adiabatensatzes'.

Quantum Annealing for Music Arrangement

Theory

- └ Adiabatic quantum computing
 - └ Adiabatic quantum computing

2025-03-04

Adiabatic quantum computing

Adiabatic principle
A system remains in its instantaneous eigenstate if a given perturbation is acting on it slowly enough.¹

$$H(t) = \left(1 - \frac{t}{T}\right) H_0 + \frac{t}{T} H_p$$

- Universal and guaranteed
- A system that starts in a ground state, ends in a ground state

¹Born and Fock, 'Beweis des Adiabatensatzes'

Adiabatic quantum computing

Adiabatic principle

A system remains in its instantaneous eigenstate if a given perturbation is acting on it slowly enough.¹

$$H(t) = \left(1 - \frac{t}{T}\right) H_0 + \frac{t}{T} H_p$$

- Universal and guaranteed
- A system that starts in a ground state, ends in a ground state
- Not possible in practice

¹Born and Fock, 'Beweis des Adiabatensatzes'.

Quantum Annealing for Music Arrangement

Theory

Adiabatic quantum computing

Adiabatic quantum computing

2025-03-04

Adiabatic quantum computing

Adiabatic principle

A system remains in its instantaneous eigenstate if a given perturbation is acting on it slowly enough.¹


$$H(t) = \left(1 - \frac{t}{T}\right) H_0 + \frac{t}{T} H_p$$

- Universal and guaranteed
- A system that starts in a ground state, ends in a ground state
- Not possible in practice

Born and Fock, 'Beweis des Adiabatensatzes'

- Adiabatic principle — system remains in the same eigenstate if perturbed slowly enough (without transferring heat)
- Equation shows evolution from initial Hamiltonian H_0 to final H_p over time T
- Importantly, if the system starts in the ground state, it will end in the ground state
- Impossible in practice as true adiabatic evolution would take infinite time, infinitely many steps

Quantum annealing




Quantum Annealing for Music Arrangement

2025-03-04

Theory

Quantum annealing


Quantum annealing



- Subset of AQC, relaxes the adiabaticity condition
- Annealing — slow heating of a material to change its properties
- Evolution time shortened (order of a few μs)
- End state no longer guaranteed, if started in ground state could end in excited state
- Able to run the evolution many times
- Probabilistic distribution of outcomes, sometimes will get lucky

Quantum annealing

- Relaxes the adiabaticity




Quantum Annealing for Music Arrangement

2025-03-04

Theory


- └ Quantum annealing
- └ Quantum annealing



- Subset of AQC, relaxes the adiabaticity condition
- Annealing — slow heating of a material to change its properties
- Evolution time shortened (order of a few μs)
- End state no longer guaranteed, if started in ground state could end in excited state
- Able to run the evolution many times
- Probabilistic distribution of outcomes, sometimes will get lucky

Quantum annealing

- Relaxes the adiabaticity
- Rate of change determined heuristically




Quantum Annealing for Music Arrangement

2025-03-04

Theory

Quantum annealing


Quantum annealing



- Subset of AQC, relaxes the adiabaticity condition
- Annealing — slow heating of a material to change its properties
- Evolution time shortened (order of a few μs)
- End state no longer guaranteed, if started in ground state could end in excited state
- Able to run the evolution many times
- Probabilistic distribution of outcomes, sometimes will get lucky

Quantum annealing

- Relaxes the adiabaticity
- Rate of change determined heuristically
- Final state is probabilistic, not deterministic



4

Quantum Annealing for Music Arrangement

2025-03-04


Theory

Quantum annealing

Quantum annealing


Quantum annealing

- Relaxes the adiabaticity
- Rate of change determined heuristically
- Final state is probabilistic, not deterministic



- Subset of AQC, relaxes the adiabaticity condition
- Annealing — slow heating of a material to change its properties
- Evolution time shortened (order of a few μs)
- End state no longer guaranteed, if started in ground state could end in excited state
- Able to run the evolution many times
- Probabilistic distribution of outcomes, sometimes will get lucky

Advantages




Quantum Annealing for Music Arrangement

2025-03-04

Theory


- Quantum annealing
- Advantages



- Why is this technique useful?
- Allows us to find the ground state of complicated Hamiltonians by starting from an easy one
- Diagram — energy against configuration space, simulated annealing (classical) traverses the "energy landscape" whereas quantum annealing tunnels through it
- As opposed to classical methods, does not get affected by local minima
- Technique very good for solving optimisation problems e.g. travelling salesman, with complicated energy landscapes

Advantages

- Find the ground state of complicated Hamiltonians




By Brianlechthaler - Own work, CC BY-SA 4.0,
<https://commons.wikimedia.org/w/index.php?curid=112382195>

Quantum Annealing for Music Arrangement

2025-03-04

Theory


- Quantum annealing
- Advantages



- Why is this technique useful?
- Allows us to find the ground state of complicated Hamiltonians by starting from an easy one
- Diagram — energy against configuration space, simulated annealing (classical) traverses the "energy landscape" whereas quantum annealing tunnels through it
- As opposed to classical methods, does not get affected by local minima
- Technique very good for solving optimisation problems e.g. travelling salesman, with complicated energy landscapes

Advantages

- Find the ground state of complicated Hamiltonians
- Quantum tunneling avoids local minima




By Brianlechthaler - Own work, CC BY-SA 4.0,
<https://commons.wikimedia.org/w/index.php?curid=112382195>

Quantum Annealing for Music Arrangement

2025-03-04

Theory

- └ Quantum annealing
- └ Advantages



- Why is this technique useful?
- Allows us to find the ground state of complicated Hamiltonians by starting from an easy one
- Diagram — energy against configuration space, simulated annealing (classical) traverses the "energy landscape" whereas quantum annealing tunnels through it
- As opposed to classical methods, does not get affected by local minima
- Technique very good for solving optimisation problems e.g. travelling salesman, with complicated energy landscapes

Lattice of variables with two discrete values

2025-03-04

Ising model

Lattice of variables with two discrete values

Ising model

Lattice of variables with two discrete values

Initial Hamiltonian


$$H_0 = h_0 \sum_{i=1}^N \sigma_i^x$$

6

Quantum Annealing for Music Arrangement

2025-03-04

- └ Theory
 - └ Quantum annealing
 - └ Ising model



Ising model

Lattice of variables with two discrete values

Initial Hamiltonian

$$H_0 = h_0 \sum_{i=1}^N \sigma_i^x$$

Problem Hamiltonian

$$H_p = \sum_{i < j}^N J_{ij} \sigma_i^z \sigma_j^z + \sum_{i=1}^N h_i \sigma_i^z$$


Quantum Annealing for Music Arrangement

2025-03-04

Theory

Quantum annealing

Ising model



- How can we model the Hamiltonians?
- Ising model, a lattice of variables with two discrete values (+1/-1), acted on by spin operators σ
- Start with initial Hamiltonian, superposition of all possible states, easy to prepare and find the ground state
- Problem Hamiltonian, coupling strengths J_{ij} and field strengths h_i , describe interactions (biases) of the spins
- Want to encode the problem solution into the ground state of this Hamiltonian so that the system will give the solution after evolution

2025-03-04

Theory

Quantum annealing

QUBO

QUBO

- How to encode a problem into a Hamiltonian?
- Similar form to the Ising model, but with binary variables (0/1)
- Minimisation of this function should be the problem solution
- Set of binary variables x , matrix Q of real weights that describes interactions between variables
- After evolution, can read out the values of x to give solution

Quadratic Unconstrained Binary Optimisation

Vector x of qubits, matrix Q of weights

$$f(x) = \sum_{i < j}^N Q_{i,j}x_i x_j + \sum_i^N Q_{i,i}x_i$$

Quantum Annealing for Music Arrangement

2025-03-04

- └ Theory
 - └ Quantum annealing
 - └ QUBO

QUBO

Quadratic Unconstrained Binary Optimisation
Vector x of qubits, matrix Q of weights

$$f(x) = \sum_{i < j}^N Q_{i,j}x_i x_j + \sum_i^N Q_{i,i}x_i$$

Quadratic Unconstrained Binary Optimisation

Vector x of qubits, matrix Q of weights

$$f(x) = \sum_{i < j}^N Q_{i,j} x_i x_j + \sum_i^N Q_{i,i} x_i$$

Quantum Annealing for Music Arrangement

2025-03-04

Theory

- └ Quantum annealing
- └ QUBO

QUBO

Quadratic Unconstrained Binary Optimisation
Vector x of qubits, matrix Q of weights

$$f(x) = \sum_{i < j}^N Q_{i,j} x_i x_j + \sum_i^N Q_{i,i} x_i$$

Quadratic Unconstrained Binary Optimisation

Vector x of qubits, matrix Q of weights

$$f(x) = \sum_{i < j}^N Q_{i,j}x_i x_j + \sum_i^N Q_{i,i}x_i$$

Quantum Annealing for Music Arrangement

2025-03-04

Theory

- └ Quantum annealing
- └ QUBO

QUBO

Quadratic Unconstrained Binary Optimisation
Vector x of qubits, matrix Q of weights
$$f(x) = \sum_{i < j}^N Q_{i,j}x_i x_j + \sum_i^N Q_{i,i}x_i$$

Quadratic Unconstrained Binary Optimisation

Vector x of qubits, matrix Q of weights

$$f(x) = \sum_{i < j}^N Q_{i,j}x_i x_j + \sum_i^N Q_{i,i}x_i$$

- Aim to minimise this function
- Difficult to solve analytically
- Mapped to H_p using simple change of variable
- Encodes problem solution into Hamiltonian's ground state

Quantum Annealing for Music Arrangement

2025-03-04

Theory

Quantum annealing

QUBO

QUBO

Quadratic Unconstrained Binary Optimisation
Vector x of qubits, matrix Q of weights

$$f(x) = \sum_{i < j}^N Q_{i,j}x_i x_j + \sum_i^N Q_{i,i}x_i$$

- Aim to minimise this function
- Difficult to solve analytically
- Mapped to H_p using simple change of variable
- Encodes problem solution into Hamiltonian's ground state

- How to encode a problem into a Hamiltonian?
- Similar form to the Ising model, but with binary variables (0/1)
- Minimisation of this function should be the problem solution
- Set of binary variables x , matrix Q of real weights that describes interactions between variables
- After evolution, can read out the values of x to give solution

Motivations

What problems can we solve?

Music arrangement



www.freepik.com

²Moses and Demaine, 'Computational Complexity of Arranging Music'.

Quantum Annealing for Music Arrangement

2025-03-04

- └ Motivations
- └ Music arrangement
- └ Music arrangement

Music arrangement

www.freepik.com

©Moses and Demaine, 'Computational Complexity of Arranging Music'.

- Adaptation of music in terms of instrumentation, medium, or style
- Traditionally a complex process that requires a deep understanding of musical theory and structure
- Reduction is the rewriting of music for a smaller number of instruments (for example string quartet)
- Very large configuration space, many different combinations of notes that could produce the final arrangement
- For those interested, NP-hard in computational complexity theory, cannot be solved in polynomial time
- NB: all scores shown are own reproductions from public domain files

Music arrangement

- Adaptation of previously composed pieces for practical or artistic reasons



www.freepik.com

²Moses and Demaine, 'Computational Complexity of Arranging Music'.

Quantum Annealing for Music Arrangement

2025-03-04

- └ Motivations
- └ Music arrangement
- └ Music arrangement

Music arrangement

- Adaptation of previously composed pieces for practical or artistic reasons

www.freepik.com

© Moses and Demaine, 'Computational Complexity of Arranging Music'.

- Adaptation of music in terms of instrumentation, medium, or style
- Traditionally a complex process that requires a deep understanding of musical theory and structure
- Reduction is the rewriting of music for a smaller number of instruments (for example string quartet)
- Very large configuration space, many different combinations of notes that could produce the final arrangement
- For those interested, NP-hard in computational complexity theory, cannot be solved in polynomial time
- NB: all scores shown are own reproductions from public domain files

Music arrangement

- Adaptation of previously composed pieces for practical or artistic reasons
- Traditionally difficult and time-consuming



www.freepik.com

²Moses and Demaine, 'Computational Complexity of Arranging Music'.

Quantum Annealing for Music Arrangement

Motivations

- └ Music arrangement
- └ Music arrangement

2025-03-04

Music arrangement

- Adaptation of previously composed pieces for practical or artistic reasons
- Traditionally difficult and time-consuming

©Moses and Demaine, 'Computational Complexity of Arranging Music'.

www.freepik.com

- Adaptation of music in terms of instrumentation, medium, or style
- Traditionally a complex process that requires a deep understanding of musical theory and structure
- Reduction is the rewriting of music for a smaller number of instruments (for example string quartet)
- Very large configuration space, many different combinations of notes that could produce the final arrangement
- For those interested, NP-hard in computational complexity theory, cannot be solved in polynomial time
- NB: all scores shown are own reproductions from public domain files

Music arrangement

- Adaptation of previously composed pieces for practical or artistic reasons
- Traditionally difficult and time-consuming
- *Reduction* can be shown to be computationally complex²



www.freepik.com

²Moses and Demaine, 'Computational Complexity of Arranging Music'.


Quantum Annealing for Music Arrangement

Motivations

- └ Music arrangement
- └ Music arrangement

2025-03-04

- Adaptation of music in terms of instrumentation, medium, or style
- Traditionally a complex process that requires a deep understanding of musical theory and structure
- Reduction is the rewriting of music for a smaller number of instruments (for example string quartet)
- Very large configuration space, many different combinations of notes that could produce the final arrangement
- For those interested, NP-hard in computational complexity theory, cannot be solved in polynomial time
- NB: all scores shown are own reproductions from public domain files



³Huang, Chiu and Shan, 'Towards an automatic music arrangement framework using score reduction'; Nakamura and Yoshii, 'Statistical piano reduction controlling performance difficulty'; Li et al., 'Automatic Piano Reduction of Orchestral Music Based on Musical Entropy'.

⁴Freedline, 'Algorhythms'; Arya et al., 'Music Composition Using Quantum Annealing'.

⁵Miranda, *Quantum Computer Music*.

2025-03-04



- Context of previous work
- Classical methods — machine learning, statistical analysis, rule-based systems, iterative and slow
- Applying quantum computing to music in the last five years, still a very young technology with limitations
- Has been used to generate music, not arrange it
- Methods shown here have not been found in the literature

Motivations

- Already exist classical methods of automatic arrangement³

³Huang, Chiu and Shan, 'Towards an automatic music arrangement framework using score reduction'; Nakamura and Yoshii, 'Statistical piano reduction controlling performance difficulty'; Li et al., 'Automatic Piano Reduction of Orchestral Music Based on Musical Entropy'.

⁴Freedline, 'Algorhythms'; Arya et al., 'Music Composition Using Quantum Annealing'.

⁵Miranda, *Quantum Computer Music*.

Quantum Annealing for Music Arrangement

Motivations

Music arrangement

Motivations

2025-03-04

Motivations

- Already exist classical methods of automatic arrangement³

Yi Huang, Chiu and Shan, 'Towards an automatic music arrangement framework using score reduction'; Nakamura and Yoshii, 'Statistical piano reduction controlling performance difficulty'; Li et al., 'Automatic Piano Reduction of Orchestral Music Based on Musical Entropy'.

*Freedline, 'Algorhythms'; Arya et al., 'Music Composition Using Quantum Annealing'.

⁵Miranda, *Quantum Computer Music*.

- Context of previous work
- Classical methods — machine learning, statistical analysis, rule-based systems, iterative and slow
- Applying quantum computing to music in the last five years, still a very young technology with limitations
- Has been used to generate music, not arrange it
- Methods shown here have not been found in the literature

Motivations

- Already exist classical methods of automatic arrangement³
- Quantum annealing used to generate music⁴

³Huang, Chiu and Shan, 'Towards an automatic music arrangement framework using score reduction'; Nakamura and Yoshii, 'Statistical piano reduction controlling performance difficulty'; Li et al., 'Automatic Piano Reduction of Orchestral Music Based on Musical Entropy'.

⁴Freedline, 'Algorhythms'; Arya et al., 'Music Composition Using Quantum Annealing'.

⁵Miranda, *Quantum Computer Music*.

Quantum Annealing for Music Arrangement

Motivations

Music arrangement

Motivations

2025-03-04

Motivations

- Already exist classical methods of automatic arrangement³
- Quantum annealing used to generate music⁴

Huang, Chiu and Shan, 'Towards an automatic music arrangement framework using score reduction'; Nakamura and Yoshii, 'Statistical piano reduction controlling performance difficulty'; Li et al., 'Automatic Piano Reduction of Orchestral Music Based on Musical Entropy'.

Freedline, 'Algorhythms'; Arya et al., 'Music Composition Using Quantum Annealing'.

Miranda, *Quantum Computer Music*.

- Context of previous work
- Classical methods — machine learning, statistical analysis, rule-based systems, iterative and slow
- Applying quantum computing to music in the last five years, still a very young technology with limitations
- Has been used to generate music, not arrange it
- Methods shown here have not been found in the literature

Motivations

- Already exist classical methods of automatic arrangement³
- Quantum annealing used to generate music⁴
- Field of quantum computer music is very new⁵

³Huang, Chiu and Shan, 'Towards an automatic music arrangement framework using score reduction'; Nakamura and Yoshii, 'Statistical piano reduction controlling performance difficulty'; Li et al., 'Automatic Piano Reduction of Orchestral Music Based on Musical Entropy'.

⁴Freedline, 'Algorhythms'; Arya et al., 'Music Composition Using Quantum Annealing'.

⁵Miranda, *Quantum Computer Music*.

Quantum Annealing for Music Arrangement


2025-03-04

Motivations

Music arrangement

Motivations

- Context of previous work
- Classical methods — machine learning, statistical analysis, rule-based systems, iterative and slow
- Applying quantum computing to music in the last five years, still a very young technology with limitations
- Has been used to generate music, not arrange it
- Methods shown here have not been found in the literature



Motivations

- Already exist classical methods of automatic arrangement³
- Quantum annealing used to generate music⁴
- Field of quantum computer music is very new⁵
- Novel adaption of this method to a new problem

³Huang, Chiu and Shan, 'Towards an automatic music arrangement framework using score reduction'; Nakamura and Yoshii, 'Statistical piano reduction controlling performance difficulty'; Li et al., 'Automatic Piano Reduction of Orchestral Music Based on Musical Entropy'.

⁴Freedline, 'Algorhythms'; Arya et al., 'Music Composition Using Quantum Annealing'.

⁵Miranda, *Quantum Computer Music*.

Quantum Annealing for Music Arrangement

Motivations

Music arrangement

Motivations

2025-03-04

Motivations

- Already exist classical methods of automatic arrangement³
- Quantum annealing used to generate music⁴
- Field of quantum computer music is very new⁵
- Novel adaption of this method to a new problem

³Huang, Chiu and Shan, 'Towards an automatic music arrangement framework using score reduction'; Nakamura and Yoshii, 'Statistical piano reduction controlling performance difficulty'; Li et al., 'Automatic Piano Reduction of Orchestral Music Based on Musical Entropy'.

⁴Freedline, 'Algorhythms'; Arya et al., 'Music Composition Using Quantum Annealing'.

Nakamura, Quantum Computer Music

- Context of previous work
- Classical methods — machine learning, statistical analysis, rule-based systems, iterative and slow
- Applying quantum computing to music in the last five years, still a very young technology with limitations
- Has been used to generate music, not arrange it
- Methods shown here have not been found in the literature

Motivations

- Already exist classical methods of automatic arrangement³
- Quantum annealing used to generate music⁴
- Field of quantum computer music is very new⁵
- Novel adaption of this method to a new problem
- *This has never been done before!*

³Huang, Chiu and Shan, 'Towards an automatic music arrangement framework using score reduction'; Nakamura and Yoshii, 'Statistical piano reduction controlling performance difficulty'; Li et al., 'Automatic Piano Reduction of Orchestral Music Based on Musical Entropy'.

⁴Freedline, 'Algorhythms'; Arya et al., 'Music Composition Using Quantum Annealing'.

⁵Miranda, *Quantum Computer Music*.

Quantum Annealing for Music Arrangement

Motivations

Music arrangement

Motivations

2025-03-04

Motivations

- Already exist classical methods of automatic arrangement³
- Quantum annealing used to generate music⁴
- Field of quantum computer music is very new⁵
- Novel adaption of this method to a new problem
- *This has never been done before!*

³Huang, Chiu and Shan, 'Towards an automatic music arrangement framework using score reduction'; Nakamura and Yoshii, 'Statistical piano reduction controlling performance difficulty'; Li et al., 'Automatic Piano Reduction of Orchestral Music Based on Musical Entropy'.

⁴Freedline, 'Algorhythms'; Arya et al., 'Music Composition Using Quantum Annealing'.

⁵Miranda, *Quantum Computer Music*.

- Context of previous work
- Classical methods — machine learning, statistical analysis, rule-based systems, iterative and slow
- Applying quantum computing to music in the last five years, still a very young technology with limitations
- Has been used to generate music, not arrange it
- Methods shown here have not been found in the literature

2025-03-04

Method



Joseph Haydn playing in a string quartet,
painting from the StaatsMuseum,
Vienna

Quantum Annealing for Music Arrangement

Method

Aims

- What are we trying to do? What are the constraints to our problem?
- Take a musical score and reduce it to a smaller ensemble
- All notes must be taken from the original score, no new notes can be added
- Each instrument can only take notes from one part at a time

2025-03-04

Aims



Joseph Haydn playing in a string quartet,
painting from the StaatsMuseum,
Vienna

Aims

- Arrange a musical score for a smaller ensemble



Joseph Haydn playing in a string quartet,
painting from the StaatsMuseum,
Vienna

Quantum Annealing for Music Arrangement

Method

Aims

- What are we trying to do? What are the constraints to our problem?
- Take a musical score and reduce it to a smaller ensemble
- All notes must be taken from the original score, no new notes can be added
- Each instrument can only take notes from one part at a time

2025-03-04

Aims

- Arrange a musical score for a smaller ensemble



Joseph Haydn playing in a string quartet,
painting from the StaatsMuseum,
Vienna

Aims

- Arrange a musical score for a smaller ensemble
- All notes are taken from the original score



Joseph Haydn playing in a string quartet,
painting from the StaatsMuseum,
Vienna

Quantum Annealing for Music Arrangement

Method

Aims

2025-03-04

- What are we trying to do? What are the constraints to our problem?
- Take a musical score and reduce it to a smaller ensemble
- All notes must be taken from the original score, no new notes can be added
- Each instrument can only take notes from one part at a time

Aims

- Arrange a musical score for a smaller ensemble
- All notes are taken from the original score



Joseph Haydn playing in a string quartet
painting from the StaatsMuseum,
Vienna

Aims

- Arrange a musical score for a smaller ensemble
- All notes are taken from the original score
- Each instrument can only play one note at a time



Joseph Haydn playing in a string quartet,
painting from the StaatsMuseum,
Vienna

Quantum Annealing for Music Arrangement

Method

Aims

2025-03-04

- What are we trying to do? What are the constraints to our problem?
- Take a musical score and reduce it to a smaller ensemble
- All notes must be taken from the original score, no new notes can be added
- Each instrument can only take notes from one part at a time

Aims

- Arrange a musical score for a smaller ensemble
- All notes are taken from the original score
- Each instrument can only play one note at a time



Joseph Haydn playing in a string quartet
painting from the StaatsMuseum,
Vienna

2025-03-04

- Formulating arrangement as a problem to be solved via annealing, five-step process
- Split parts into musical phrases
- Arrange phrases into a graph (will explain later)
- Formulate the optimisation problem
- Solve corresponding graph problem using a quantum computer
- Construct final arrangement from the solution returned

Method

1. Split score into musical phrases

Quantum Annealing for Music Arrangement

└ Method

└ Method

- Formulating arrangement as a problem to be solved via annealing, five-step process
- Split parts into musical phrases
- Arrange phrases into a graph (will explain later)
- Formulate the optimisation problem
- Solve corresponding graph problem using a quantum computer
- Construct final arrangement from the solution returned

1. Split score into musical phrases

Method

Method

1. Split score into musical phrases
2. Arrange phrases into a graph

Quantum Annealing for Music Arrangement

Method

Method

- Formulating arrangement as a problem to be solved via annealing, five-step process
- Split parts into musical phrases
- Arrange phrases into a graph (will explain later)
- Formulate the optimisation problem
- Solve corresponding graph problem using a quantum computer
- Construct final arrangement from the solution returned

Method

1. Split score into musical phrases
2. Arrange phrases into a graph

Method

1. Split score into musical phrases
2. Arrange phrases into a graph
3. Formulate optimisation problem

11

Quantum Annealing for Music Arrangement

Method

Method

2025-03-04

- Formulating arrangement as a problem to be solved via annealing, five-step process
 - Split parts into musical phrases
 - Arrange phrases into a graph (will explain later)
 - Formulate the optimisation problem
 - Solve corresponding graph problem using a quantum computer
 - Construct final arrangement from the solution returned

- 1. Split score into musical phrases
- 2. Arrange phrases into a graph
- 3. Formulate optimisation problem

Method

Method

1. Split score into musical phrases
2. Arrange phrases into a graph
3. Formulate optimisation problem
4. Solve problem using QPU

Quantum Annealing for Music Arrangement

Method

Method

2025-03-04

Method

1. Split score into musical phrases
2. Arrange phrases into a graph
3. Formulate optimisation problem
4. Solve problem using QPU

- Formulating arrangement as a problem to be solved via annealing, five-step process
 - Split parts into musical phrases
 - Arrange phrases into a graph (will explain later)
 - Formulate the optimisation problem
 - Solve corresponding graph problem using a quantum computer
 - Construct final arrangement from the solution returned

Method

1. Split score into musical phrases
2. Arrange phrases into a graph
3. Formulate optimisation problem
4. Solve problem using QPU
5. Construct arrangement from solution

Quantum Annealing for Music Arrangement

Method

Method

2025-03-04

Method

1. Split score into musical phrases
2. Arrange phrases into a graph
3. Formulate optimisation problem
4. Solve problem using QPU
5. Construct arrangement from solution

- Formulating arrangement as a problem to be solved via annealing, five-step process
 - Split parts into musical phrases
 - Arrange phrases into a graph (will explain later)
 - Formulate the optimisation problem
 - Solve corresponding graph problem using a quantum computer
 - Construct final arrangement from the solution returned

1. Split score


⁶Cambouropoulos, 'The Local Boundary Detection Model (LBDM) and its Application in the Study of Expressive Timing'.

2025-03-04

Quantum Annealing for Music Arrangement

└ Method

└ 1. Split score



- First stage to separate each part of original score into phrases
- Phrases — smallest unit of music that preserves melody and structure
- Boundaries between phrases found using LBDM
- Measures the degree of change of a certain parameter (x) between notes (i) (explain equation)
- Strength calculated for both pitch and IOI, weighted and summed to give the final strength
- Strengths above a threshold value are considered phrases

1. Split score

- Musical phrases chosen as smallest unit of music
- Preserve melody and structure when rearranged


⁶Cambouropoulos, 'The Local Boundary Detection Model (LBDM) and its Application in the Study of Expressive Timing'.

Quantum Annealing for Music Arrangement

Method

1. Split score

- First stage to separate each part of original score into phrases
- Phrases — smallest unit of music that preserves melody and structure
- Boundaries between phrases found using LBDM
- Measures the degree of change of a certain parameter (x) between notes (i) (explain equation)
- Strength calculated for both pitch and IOI, weighted and summed to give the final strength
- Strengths above a threshold value are considered phrases



1. Split score

- Musical phrases chosen as smallest unit of music
- Preserve melody and structure when rearranged

Local boundary detection model (LBDM)⁶

$$S_i = x_i \times (r_{i-1,i} + r_{i,i+1})$$


⁶Cambouropoulos, 'The Local Boundary Detection Model (LBDM) and its Application in the Study of Expressive Timing'.

Quantum Annealing for Music Arrangement

Method

1. Split score

- First stage to separate each part of original score into phrases
- Phrases — smallest unit of music that preserves melody and structure
- Boundaries between phrases found using LBDM
- Measures the degree of change of a certain parameter (x) between notes (i) (explain equation)
- Strength calculated for both pitch and IOI, weighted and summed to give the final strength
- Strengths above a threshold value are considered phrases




1. Split score

- Musical phrases chosen as smallest unit of music
- Preserve melody and structure when rearranged

Local boundary detection model (LBDM)⁶

$$S_i = x_i \times (r_{i-1,i} + r_{i,i+1})$$




⁶Cambouropoulos, 'The Local Boundary Detection Model (LBDM) and its Application in the Study of Expressive Timing'.

Quantum Annealing for Music Arrangement

Method

1. Split score

- First stage to separate each part of original score into phrases
- Phrases — smallest unit of music that preserves melody and structure
- Boundaries between phrases found using LBDM
- Measures the degree of change of a certain parameter (x) between notes (i) (explain equation)
- Strength calculated for both pitch and IOI, weighted and summed to give the final strength
- Strengths above a threshold value are considered phrases



2025-03-04

- What is a graph? Nodes connected by edges, useful to model pairwise relations between objects
- Each phrase becomes a node, edges between nodes if phrases overlap (play at the same time)

2. Create graph

- Vertices (nodes) connected by edges
- Models pairwise relations between objects

Quantum Annealing for Music Arrangement

└ Method

└ 2. Create graph


- What is a graph? Nodes connected by edges, useful to model pairwise relations between objects
- Each phrase becomes a node, edges between nodes if phrases overlap (play at the same time)

2. Create graph

- Vertices (nodes) connected by edges
- Models pairwise relations between objects

2. Create graph

- Vertices (nodes) connected by edges
- Models pairwise relations between objects




- Nodes — phrases
- Edges — overlap between phrases

Quantum Annealing for Music Arrangement

Method

2. Create graph

- What is a graph? Nodes connected by edges, useful to model pairwise relations between objects
- Each phrase becomes a node, edges between nodes if phrases overlap (play at the same time)






Quantum Annealing for Music Arrangement

Method


2. Create graph

- Score on top becomes graph on bottom

2025-03-04



2. Create graph




Quantum Annealing for Music Arrangement

└ Method

└ 2. Create graph

- Score on top becomes graph on bottom

2025-03-04



3. Create optimisation problem

Quantum Annealing for Music Arrangement

Method

└ 3. Create optimisation problem

2025-03-04

- Use a graph theory problem to create the optimisation problem that matches our constraints
- Here each colour represents an instrument we are arranging for
- QUBO, set of n colours, $x_{v,i}$ is 1 if node v is colour i
- A — each node is only coloured once, sum over colours is one
- B — penalise adjacent nodes with the same colour
- C — weight of each node, preference for certain nodes
- D — weight of each edge, preference for certain edges
- Weights here are musical entropy i.e. how interesting the phrase is musically

3. Create optimisation problem

3. Create optimisation problem

Proper vertex colouring

Colour each vertex such that no edge connects two vertices of the same colour

Quantum Annealing for Music Arrangement

Method

3. Create optimisation problem

- Use a graph theory problem to create the optimisation problem that matches our constraints
- Here each colour represents an instrument we are arranging for
- QUBO, set of n colours, $x_{v,i}$ is 1 if node v is colour i
- A — each node is only coloured once, sum over colours is one
- B — penalise adjacent nodes with the same colour
- C — weight of each node, preference for certain nodes
- D — weight of each edge, preference for certain edges
- Weights here are musical entropy i.e. how interesting the phrase is musically

3. Create optimisation problem

Proper vertex colouring

Colour each vertex such that no edge connects two vertices of the same colour

$$x_{v,i} = \begin{cases} 1 & \text{if vertex } v \text{ is colour } i \\ 0 & \text{otherwise} \end{cases}$$

Quantum Annealing for Music Arrangement

Method

3. Create optimisation problem

- Use a graph theory problem to create the optimisation problem that matches our constraints
- Here each colour represents an instrument we are arranging for
- QUBO, set of n colours, $x_{v,i}$ is 1 if node v is colour i
- A — each node is only coloured once, sum over colours is one
- B — penalise adjacent nodes with the same colour
- C — weight of each node, preference for certain nodes
- D — weight of each edge, preference for certain edges
- Weights here are musical entropy i.e. how interesting the phrase is musically

3. Create optimisation problem

Proper vertex colouring
Colour each vertex such that no edge connects two vertices of the same colour

$$x_{v,i} = \begin{cases} 1 & \text{if vertex } v \text{ is colour } i \\ 0 & \text{otherwise} \end{cases}$$

3. Create optimisation problem

Proper vertex colouring

Colour each vertex such that no edge connects two vertices of the same colour

$$x_{v,i} = \begin{cases} 1 & \text{if vertex } v \text{ is colour } i \\ 0 & \text{otherwise} \end{cases}$$

$$\begin{aligned} f(x) = & +A \sum_{v \in V} \left(1 - \sum_{i=1}^n x_{v,i} \right)^2 + B \sum_{(u,v) \in E} \sum_{i=1}^n x_{u,i} x_{v,i} \\ & - C \sum_{v \in V} \sum_{i=1}^n W_v x_{v,i} \quad \quad \quad - D \sum_{(u,v) \in E} W_{uv} \sum_{i=1}^n \sum_{j=1}^n x_{u,i} x_{v,j} \end{aligned}$$

Quantum Annealing for Music Arrangement

Method

3. Create optimisation problem

3. Create optimisation problem

Proper vertex colouring
Colour each vertex such that no edge connects two vertices of the same colour

$$x_{v,i} = \begin{cases} 1 & \text{if vertex } v \text{ is colour } i \\ 0 & \text{otherwise} \end{cases}$$
$$f(x) = +A \sum_{v \in V} \left(1 - \sum_{i=1}^n x_{v,i} \right)^2 + B \sum_{(u,v) \in E} \sum_{i=1}^n x_{u,i} x_{v,i} - C \sum_{v \in V} \sum_{i=1}^n W_v x_{v,i} - D \sum_{(u,v) \in E} W_{uv} \sum_{i=1}^n \sum_{j=1}^n x_{u,i} x_{v,j}$$

3. Create optimisation problem

Proper vertex colouring

Colour each vertex such that no edge connects two vertices of the same colour

$$x_{v,i} = \begin{cases} 1 & \text{if vertex } v \text{ is colour } i \\ 0 & \text{otherwise} \end{cases}$$

$$\begin{aligned} f(x) = & +A \sum_{v \in V} \left(1 - \sum_{i=1}^n x_{v,i} \right)^2 + B \sum_{(u,v) \in E} \sum_{i=1}^n x_{u,i} x_{v,i} \\ & - C \sum_{v \in V} \sum_{i=1}^n W_v x_{v,i} \quad - D \sum_{(u,v) \in E} W_{uv} \sum_{i=1}^n \sum_{j=1}^n x_{u,i} x_{v,j} \end{aligned}$$

Quantum Annealing for Music Arrangement

Method

3. Create optimisation problem

3. Create optimisation problem

Proper vertex colouring
Colour each vertex such that no edge connects two vertices of the same colour

$$x_{v,i} = \begin{cases} 1 & \text{if vertex } v \text{ is colour } i \\ 0 & \text{otherwise} \end{cases}$$
$$f(x) = +A \sum_{v \in V} \left(1 - \sum_{i=1}^n x_{v,i} \right)^2 + B \sum_{(u,v) \in E} \sum_{i=1}^n x_{u,i} x_{v,i} - C \sum_{v \in V} \sum_{i=1}^n W_v x_{v,i} - D \sum_{(u,v) \in E} W_{uv} \sum_{i=1}^n \sum_{j=1}^n x_{u,i} x_{v,j}$$

3. Create optimisation problem

Proper vertex colouring

Colour each vertex such that no edge connects two vertices of the same colour

$$x_{v,i} = \begin{cases} 1 & \text{if vertex } v \text{ is colour } i \\ 0 & \text{otherwise} \end{cases}$$

$$\begin{aligned} f(x) = & +A \sum_{v \in V} \left(1 - \sum_{i=1}^n x_{v,i} \right)^2 + B \sum_{(u,v) \in E} \sum_{i=1}^n x_{u,i} x_{v,i} \\ & - C \sum_{v \in V} \sum_{i=1}^n W_v x_{v,i} - D \sum_{(u,v) \in E} W_{uv} \sum_{i=1}^n \sum_{j=1}^n x_{u,i} x_{v,j} \end{aligned}$$

Quantum Annealing for Music Arrangement

Method

3. Create optimisation problem

3. Create optimisation problem

Proper vertex colouring
Colour each vertex such that no edge connects two vertices of the same colour

$$x_{v,i} = \begin{cases} 1 & \text{if vertex } v \text{ is colour } i \\ 0 & \text{otherwise} \end{cases}$$
$$f(x) = +A \sum_{v \in V} \left(1 - \sum_{i=1}^n x_{v,i} \right)^2 + B \sum_{(u,v) \in E} \sum_{i=1}^n x_{u,i} x_{v,i} - C \sum_{v \in V} \sum_{i=1}^n W_v x_{v,i} - D \sum_{(u,v) \in E} W_{uv} \sum_{i=1}^n \sum_{j=1}^n x_{u,i} x_{v,j}$$

3. Create optimisation problem

Proper vertex colouring

Colour each vertex such that no edge connects two vertices of the same colour

$$x_{v,i} = \begin{cases} 1 & \text{if vertex } v \text{ is colour } i \\ 0 & \text{otherwise} \end{cases}$$

$$\begin{aligned} f(x) = & +A \sum_{v \in V} \left(1 - \sum_{i=1}^n x_{v,i} \right)^2 + B \sum_{(u,v) \in E} \sum_{i=1}^n x_{u,i} x_{v,i} \\ & - C \sum_{v \in V} \sum_{i=1}^n W_v x_{v,i} \quad - D \sum_{(u,v) \in E} W_{uv} \sum_{i=1}^n \sum_{j=1}^n x_{u,i} x_{v,j} \end{aligned}$$

Quantum Annealing for Music Arrangement

Method

3. Create optimisation problem

3. Create optimisation problem

Proper vertex colouring
Colour each vertex such that no edge connects two vertices of the same colour

$$x_{v,i} = \begin{cases} 1 & \text{if vertex } v \text{ is colour } i \\ 0 & \text{otherwise} \end{cases}$$
$$f(x) = +A \sum_{v \in V} \left(1 - \sum_{i=1}^n x_{v,i} \right)^2 + B \sum_{(u,v) \in E} \sum_{i=1}^n x_{u,i} x_{v,i} - C \sum_{v \in V} \sum_{i=1}^n W_v x_{v,i} - D \sum_{(u,v) \in E} W_{uv} \sum_{i=1}^n \sum_{j=1}^n x_{u,i} x_{v,j}$$

3. Create optimisation problem

Proper vertex colouring

Colour each vertex such that no edge connects two vertices of the same colour

$$x_{v,i} = \begin{cases} 1 & \text{if vertex } v \text{ is colour } i \\ 0 & \text{otherwise} \end{cases}$$

$$\begin{aligned} f(x) = & +A \sum_{v \in V} \left(1 - \sum_{i=1}^n x_{v,i} \right)^2 + B \sum_{(u,v) \in E} \sum_{i=1}^n x_{u,i} x_{v,i} \\ & - C \sum_{v \in V} \sum_{i=1}^n W_v x_{v,i} \quad - D \sum_{(u,v) \in E} W_{uv} \sum_{i=1}^n \sum_{j=1}^n x_{u,i} x_{v,j} \end{aligned}$$

Quantum Annealing for Music Arrangement

Method


3. Create optimisation problem

3. Create optimisation problem

Proper vertex colouring
Colour each vertex such that no edge connects two vertices of the same colour

$$x_{v,i} = \begin{cases} 1 & \text{if vertex } v \text{ is colour } i \\ 0 & \text{otherwise} \end{cases}$$
$$f(x) = +A \sum_{v \in V} \left(1 - \sum_{i=1}^n x_{v,i} \right)^2 + B \sum_{(u,v) \in E} \sum_{i=1}^n x_{u,i} x_{v,i} - C \sum_{v \in V} \sum_{i=1}^n W_v x_{v,i} - D \sum_{(u,v) \in E} W_{uv} \sum_{i=1}^n \sum_{j=1}^n x_{u,i} x_{v,j}$$

3. Create optimisation problem



Quantum Annealing for Music Arrangement


└ Method

└ 3. Create optimisation problem

3. Create optimisation problem



3. Create optimisation problem




Quantum Annealing for Music Arrangement


Method

3. Create optimisation problem

- $n = 1$
- One of many possible solutions



4. Solve problem



D-Wave Advantage QPU topology. Own work.

18

Quantum Annealing for Music Arrangement

Method


4. Solve problem

- D-Wave Systems is a company that gives access to true quantum annealers, normally for business applications
- Interact via a Python SDK, submit problems to the QPU
- Returns a distribution of results, each with an associated energy
- Run the problem thousands of times to find the lowest-energy solutions



4. Solve problem

- Problem embedded onto D-Wave quantum hardware



D-Wave Advantage QPU topology. Own work.

Quantum Annealing for Music Arrangement

Method

4. Solve problem

- D-Wave Systems is a company that gives access to true quantum annealers, normally for business applications
- Interact via a Python SDK, submit problems to the QPU
- Returns a distribution of results, each with an associated energy
- Run the problem thousands of times to find the lowest-energy solutions

2025-03-04

4. Solve problem


• Problem embedded onto D-Wave quantum hardware



D-Wave Advantage QPU topology. Own work.

4. Solve problem

- Problem embedded onto D-Wave quantum hardware
- Quantum annealer optimises QUBO formulation



D-Wave Advantage QPU topology. Own work.

18

Quantum Annealing for Music Arrangement

Method

4. Solve problem

- D-Wave Systems is a company that gives access to true quantum annealers, normally for business applications
- Interact via a Python SDK, submit problems to the QPU
- Returns a distribution of results, each with an associated energy
- Run the problem thousands of times to find the lowest-energy solutions

4. Solve problem


- Problem embedded onto D-Wave quantum hardware
- Quantum annealer optimises QUBO formulation



D-Wave Advantage QPU topology. Own work.

4. Solve problem

- Problem embedded onto D-Wave quantum hardware
- Quantum annealer optimises QUBO formulation
- Returns a sampleset of results



D-Wave Advantage QPU topology. Own work.

Quantum Annealing for Music Arrangement

Method

4. Solve problem

- D-Wave Systems is a company that gives access to true quantum annealers, normally for business applications
- Interact via a Python SDK, submit problems to the QPU
- Returns a distribution of results, each with an associated energy
- Run the problem thousands of times to find the lowest-energy solutions

4. Solve problem


- Problem embedded onto D-Wave quantum hardware
- Quantum annealer optimises QUBO formulation
- Returns a sampleset of results



D-Wave Advantage QPU topology. Own work.

4. Solve problem

- Problem embedded onto D-Wave quantum hardware
- Quantum annealer optimises QUBO formulation
- Returns a sampleset of results
- Run many times to find lowest-energy solution



D-Wave Advantage QPU topology. Own work.

Quantum Annealing for Music Arrangement

Method

4. Solve problem

- D-Wave Systems is a company that gives access to true quantum annealers, normally for business applications
- Interact via a Python SDK, submit problems to the QPU
- Returns a distribution of results, each with an associated energy
- Run the problem thousands of times to find the lowest-energy solutions


4. Solve problem

- Problem embedded onto D-Wave quantum hardware
- Quantum annealer optimises QUBO formulation
- Returns a sampleset of results
- Run many times to find lowest-energy solution



D-Wave Advantage QPU topology. Own work.

5. Construct arrangement



Quantum Annealing for Music Arrangement

Method



5. Construct arrangement

- Take chosen low-energy solution and construct the final arrangement
- Map each node back to its phrase, with colour corresponding to the instrument

5. Construct arrangement



5. Construct arrangement




Quantum Annealing for Music Arrangement

Method

5. Construct arrangement

- Take chosen low-energy solution and construct the final arrangement
- Map each node back to its phrase, with colour corresponding to the instrument



2025-03-04

Results

Score

The image shows a musical score for 'Quartet No. 1 in B_b major' by Joseph Haydn. The title is at the top center. Below it, the instrumentation is listed: Violin I, Violin II, Viola, and Cello. The first page starts with a 'Presto' tempo marking. The score consists of three staves of music, each with four measures. The violins play eighth-note patterns, the viola provides harmonic support, and the cello provides bass lines. The second page continues with similar patterns, maintaining the 'Presto' tempo.

Quartet No. 1 in Bb major by
Joseph Haydn

Quantum Annealing for Music Arrangement

Results

Score

- Quartet No. 1 in Bb major by Joseph Haydn
- Smaller instrumentation and length (about 3 min), keeping the problem graph small and manageable
- Musical style has well-defined structure and phrases

2025-03-04

The screenshot shows a software interface titled 'Score'. On the left, there's a navigation tree with 'Score' at the root, followed by 'Quartet No. 1 in Bb major by Joseph Haydn'. The main area displays a musical score for a string quartet. The score includes four staves: Violin I, Violin II, Viola, and Cello. The music is presented in a standard musical notation format with notes, rests, and dynamic markings like 'Presto' and 'f' (fortissimo). The interface has a clean, modern design with dark mode selected.

Score

- Smaller ensemble chosen for problem size



Quartet No. 1 in Bb major by
Joseph Haydn

Quantum Annealing for Music Arrangement

Results

Score

2025-03-04

Score

- Smaller ensemble chosen for problem size

Quartet No. 1 in Bb major by Joseph Haydn

A screenshot of a software interface titled "Score". It shows a visualization of a musical score with four staves. A legend on the right indicates that darker colors represent a "Smaller ensemble chosen for problem size". The interface has a dark theme with orange and white highlights.

- Quartet No. 1 in Bb major by Joseph Haydn
- Smaller instrumentation and length (about 3 min), keeping the problem graph small and manageable
- Musical style has well-defined structure and phrases

Score

- Smaller ensemble chosen for problem size
- Well-defined musical structure



Quartet No. 1 in Bb major by
Joseph Haydn

Quantum Annealing for Music Arrangement

Results


Score

2025-03-04

Score

- Smaller ensemble chosen for problem size
- Well-defined musical structure

Quartet No. 1 in Bb major by Joseph Haydn



Score

- Smaller ensemble chosen for problem size
- Well-defined musical structure
- Reduction to three instruments



Quartet No. 1 in Bb major by
Joseph Haydn

Quantum Annealing for Music Arrangement

Results

Score

2025-03-04


Score

- Smaller ensemble chosen for problem size
- Well-defined musical structure
- Reduction to three instruments

Quartet No. 1 in Bb major by Joseph Haydn



Phrase detection




Quantum Annealing for Music Arrangement


Results

Phrase detection

- Example of the LBDM finding suitable boundaries for phrases
- Threshold value of 0.4 chosen manually



QUBO parameter variation




Variation of the edge constraint Lagrange parameter B

Quantum Annealing for Music Arrangement


Results

QUBO parameter variation

- Each QUBO problem submitted five times with different edge constraint Lagrange parameter
- Checking against fulfillment of the desired constraint
- Lagrange parameters taken as multipliers of the maximum node weight for normalisation
- 12.0 chosen as the best parameter, with all others equal to one



Optimisation




Variation of the number of QPU reads, with the lowest-energy solution found

Quantum Annealing for Music Arrangement

Results

Optimisation

2025-03-04



- Once Lagrange parameters chosen, can check how well the annealer optimises the problem
- In general, more reads is more likely to find lower-energy solutions
- Sometimes the annealer gets lucky (see 2000 reads)
- Each number of reads repeated five times, exponential decay fitted

2025-03-04

Conclusions

2025-03-04

- Successful application of this method on a new problem
- QPU returns samples that fulfill the constraints of the problem, creating a valid arrangement
- New technology, limited in power
- What would it take for quantum to show advantage?

Conclusions

- Successful novel application of quantum annealing

Quantum Annealing for Music Arrangement

└ Conclusions

└ Conclusions

- Successful application of this method on a new problem
- QPU returns samples that fulfill the constraints of the problem, creating a valid arrangement
- New technology, limited in power
- What would it take for quantum to show advantage?

• Successful novel application of quantum annealing

Conclusions

Conclusions

- Successful novel application of quantum annealing
- QPU returns low-energy samples

Quantum Annealing for Music Arrangement

└ Conclusions

└ Conclusions

- Successful application of this method on a new problem
- QPU returns samples that fulfill the constraints of the problem, creating a valid arrangement
- New technology, limited in power
- What would it take for quantum to show advantage?

- Successful novel application of quantum annealing
- QPU returns low-energy samples

Conclusions

- Successful novel application of quantum annealing
- QPU returns low-energy samples
- Necessary constraints for a valid arrangement fulfilled

Quantum Annealing for Music Arrangement

└ Conclusions

└ Conclusions

- Successful application of this method on a new problem
- QPU returns samples that fulfill the constraints of the problem, creating a valid arrangement
- New technology, limited in power
- What would it take for quantum to show advantage?

Conclusions

- Successful novel application of quantum annealing
- QPU returns low-energy samples
- Necessary constraints for a valid arrangement fulfilled

Conclusions

- Successful novel application of quantum annealing
- QPU returns low-energy samples
- Necessary constraints for a valid arrangement fulfilled
- Still very new technology, does not show quantum advantage (yet)

Quantum Annealing for Music Arrangement

└ Conclusions

└ Conclusions

- Successful application of this method on a new problem
- QPU returns samples that fulfill the constraints of the problem, creating a valid arrangement
- New technology, limited in power
- What would it take for quantum to show advantage?

- Successful novel application of quantum annealing
- QPU returns low-energy samples
- Necessary constraints for a valid arrangement fulfilled
- Still very new technology, does not show quantum advantage (yet)

Conclusions

2025-03-04

Future work

Quantum Annealing for Music Arrangement

└ Conclusions

└ Future work

2025-03-04

Future work

⁷Pearce and Wiggins, 'Towards A Framework for the Evaluation of Machine Compositions'.

- How well does the method scale with larger scores? How well can it find low energies with smaller problems?
- Compare to classical optimisation methods, time to solution, energy of solutions
- Only tuned one parameter by hand, could use a more systematic approach to find lower-energy solutions
- Quality judgement — Turing-like test, present subjects with human-/computer-generated scores

⁷Pearce and Wiggins, 'Towards A Framework for the Evaluation of Machine Compositions'.

Future work

- Variation in problem size

⁷Pearce and Wiggins, 'Towards A Framework for the Evaluation of Machine Compositions'.

Quantum Annealing for Music Arrangement

Conclusions

Future work

2025-03-04



- How well does the method scale with larger scores? How well can it find low energies with smaller problems?
- Compare to classical optimisation methods, time to solution, energy of solutions
- Only tuned one parameter by hand, could use a more systematic approach to find lower-energy solutions
- Quality judgement — Turing-like test, present subjects with human-/computer-generated scores

Future work

- Variation in problem size
- Comparison to classical methods

⁷Pearce and Wiggins, 'Towards A Framework for the Evaluation of Machine Compositions'.

Quantum Annealing for Music Arrangement

Conclusions

Future work

2025-03-04

Future work

- Variation in problem size
- Comparison to classical methods

⁷Pearce and Wiggins, 'Towards A Framework for the Evaluation of Machine Compositions'

- How well does the method scale with larger scores? How well can it find low energies with smaller problems?
- Compare to classical optimisation methods, time to solution, energy of solutions
- Only tuned one parameter by hand, could use a more systematic approach to find lower-energy solutions
- Quality judgement — Turing-like test, present subjects with human-/computer-generated scores

Future work

- Variation in problem size
- Comparison to classical methods
- Lagrange parameter tuning

⁷Pearce and Wiggins, 'Towards A Framework for the Evaluation of Machine Compositions'.

Quantum Annealing for Music Arrangement

Conclusions

Future work

2025-03-04

- Variation in problem size
- Comparison to classical methods
- Lagrange parameter tuning

⁷Pearce and Wiggins, 'Towards A Framework for the Evaluation of Machine Compositions'

- How well does the method scale with larger scores? How well can it find low energies with smaller problems?
- Compare to classical optimisation methods, time to solution, energy of solutions
- Only tuned one parameter by hand, could use a more systematic approach to find lower-energy solutions
- Quality judgement — Turing-like test, present subjects with human-/computer-generated scores

Future work

- Variation in problem size
- Comparison to classical methods
- Lagrange parameter tuning
- Qualitative judgement of computer arrangements⁷


⁷Pearce and Wiggins, 'Towards A Framework for the Evaluation of Machine Compositions'.

Quantum Annealing for Music Arrangement

Conclusions

Future work

2025-03-04



- How well does the method scale with larger scores? How well can it find low energies with smaller problems?
- Compare to classical optimisation methods, time to solution, energy of solutions
- Only tuned one parameter by hand, could use a more systematic approach to find lower-energy solutions
- Quality judgement — Turing-like test, present subjects with human-/computer-generated scores

Quantum Annealing for Music Arrangement

└ Conclusions

2025-03-04

Thank you!

Quantum Annealing for Music Arrangement

Lucas Kirby
4 March 2025

Department of Physics, Durham University

Quantum Annealing for Music Arrangement

2025-03-04

Quantum Annealing
for Music Arrangement

Lucas Kirby
4 March 2025
Department of Physics, Durham University

Quantum Annealing for Music Arrangement

2025-03-04

└ References

-  Arya, Ashish et al. '**Music Composition Using Quantum Annealing**'. In: *arXiv* (Jan. 2022). DOI: 10.48550/arXiv.2201.10557. (Visited on 26/10/2024).
-  Born, M. and V. Fock. '**Beweis des Adiabatensatzes**'. de. In: *Zeitschrift für Physik* 51.3 (Mar. 1928), pp. 165–180. ISSN: 0044-3328. DOI: 10.1007/BF01343193. URL: <https://doi.org/10.1007/BF01343193> (visited on 01/03/2025).
-  Cambouropoulos, Emilios. '**The Local Boundary Detection Model (LBDM) and its Application in the Study of Expressive Timing**'. In: *International Computer Music Association* (2011). ISSN: 2223-3881.

-  Arya, Ashish et al. '**Music Composition Using Quantum Annealing**'. In: *arXiv* (Jan. 2022). DOI: 10.48550/arXiv.2201.10557. (Visited on 26/10/2024).
-  Born, M. and V. Fock. '**Beweis des Adiabatensatzes**'. de. In: *Zeitschrift für Physik* 51.3 (Mar. 1928), pp. 165–180. ISSN: 0044-3328. DOI: 10.1007/BF01343193. URL: <https://doi.org/10.1007/BF01343193> (visited on 01/03/2025).
-  Cambouropoulos, Emilios. '**The Local Boundary Detection Model (LBDM) and its Application in the Study of Expressive Timing**'. In: *International Computer Music Association* (2011). ISSN: 2223-3881.

References ii

-  Freedline, Alex. ‘**Algorhythms: Generating Music with D-Wave’s Quantum Annealer**’. en. In: *MIT 6.s089—Intro to Quantum Computing* (Feb. 2021).
-  Huang, Jiun-Long, Shih-Chuan Chiu and Man-Kwan Shan. ‘**Towards an automatic music arrangement framework using score reduction**’. In: *ACM Trans. Multimedia Comput. Commun. Appl.* 8.1 (Feb. 2012), 8:1–8:23. ISSN: 1551-6857. DOI: 10.1145/2071396.2071404. (Visited on 05/12/2024).

Quantum Annealing for Music Arrangement

2025-03-04

└ References

-  Freedline, Alex. ‘**Algorhythms: Generating Music with D-Wave’s Quantum Annealer**’. en. In: *MIT 6.s089—Intro to Quantum Computing* (Feb. 2021).
-  Huang, Jiun-Long, Shih-Chuan Chiu and Man-Kwan Shan. ‘**Towards an automatic music arrangement framework using score reduction**’. In: *ACM Trans. Multimedia Comput. Commun. Appl.* 8.1 (Feb. 2012), 8:1–8:23. ISSN: 1551-6857. DOI: 10.1145/2071396.2071404. (Visited on 05/12/2024).

References iii

 Li, You et al. '**Automatic Piano Reduction of Orchestral Music Based on Musical Entropy**'. In: *2019 53rd Annual Conference on Information Sciences and Systems (CISS)*. Mar. 2019, pp. 1–5. DOI: 10.1109/CISS.2019.8693036. URL: <https://ieeexplore.ieee.org/document/8693036> (visited on 27/12/2024).

 Miranda, Eduardo Reck, ed. **Quantum Computer Music: Foundations, Methods and Advanced Concepts**. en. Springer International Publishing, 2022. ISBN: 978-3-031-13908-6 978-3-031-13909-3. DOI: 10.1007/978-3-031-13909-3. (Visited on 28/12/2024).

Quantum Annealing for Music Arrangement

2025-03-04

└ References

-  Li, You et al. '**Automatic Piano Reduction of Orchestral Music Based on Musical Entropy**', In: *2019 53rd Annual Conference on Information Sciences and Systems (CISS)*. Mar. 2019, pp. 1–5. doi: 10.1109/CISS.2019.8693036. URL: <https://ieeexplore.ieee.org/document/8693036> (visited on 27/12/2024).
-  Miranda, Eduardo Reck, ed. **Quantum Computer Music: Foundations, Methods and Advanced Concepts**. en. Springer International Publishing, 2022. isbn: 978-3-031-13908-6 978-3-031-13909-3. doi: 10.1007/978-3-031-13909-3. (Visited on 28/12/2024).

References iv

-  Moses, William S. and Erik D. Demaine. ‘**Computational Complexity of Arranging Music**’. In: *arXiv* (July 2016). arXiv:1607.04220. DOI: 10.48550/arXiv.1607.04220. (Visited on 09/11/2024).
-  Nakamura, Eita and Kazuyoshi Yoshii. ‘**Statistical piano reduction controlling performance difficulty**’. en. In: *APSIPA Transactions on Signal and Information Processing* 7 (Jan. 2018), e13. ISSN: 2048-7703. DOI: 10.1017/ATSIIP.2018.18. (Visited on 17/12/2024).
-  Pearce, M. and Geraint A. Wiggins. ‘**Towards A Framework for the Evaluation of Machine Compositions**’. In: *Proceedings of the AISB'01 Symposium on Artificial Intelligence and Creativity in the Arts and Sciences*. 2001.

Quantum Annealing for Music Arrangement

2025-03-04

└ References

-  Moses, William S. and Erik D. Demaine. ‘**Computational Complexity of Arranging Music**’. In: *arXiv* (July 2016). arXiv:1607.04220. DOI: 10.48550/arXiv.1607.04220. (Visited on 09/11/2024).
-  Nakamura, Eita and Kazuyoshi Yoshii. ‘**Statistical piano reduction controlling performance difficulty**’. en. In: *APSIPA Transactions on Signal and Information Processing* 7 (Jan. 2018), e13. ISSN: 2048-7703. DOI: 10.1017/ATSIIP.2018.18. (Visited on 17/12/2024).
-  Pearce, M. and Geraint A. Wiggins. ‘**Towards A Framework for the Evaluation of Machine Compositions**’. In: *Proceedings of the AISB'01 Symposium on Artificial Intelligence and Creativity in the Arts and Sciences*. 2001.

Boundary strength

$$S_i = x_i \times (r_{i-1,i} + r_{i,i+1})$$

$$r_{i,i+1} = \frac{|x_i - x_{i+1}|}{x_i + x_{i+1}}$$

Normalisation

$$S'_i = \frac{S_i - \min(S_i)}{\max(S_i) - \min(S_i)}$$

Weighting


$$S = \frac{1}{3} (S'_{\text{pitch}} + 2S'_{\text{IOI}})$$

⁸Cambouropoulos, 'The Local Boundary Detection Model (LBDM) and its Application in the Study of Expressive Timing'.

2025-03-04

└ LBDM

- Boundaries always taken at beginning/end of piece
- Weightings derived by trial and error



Phrase entropy

x_i — parameter x of note i

Shannon entropy

$$H(X) := - \sum_i P(x_i) \log_2 P(x_i)$$

Probability distribution

$$P(x_i) = \frac{n_i}{N}$$

⁹Li et al., 'Automatic Piano Reduction of Orchestral Music Based on Musical Entropy'.

Quantum Annealing for Music Arrangement

2025-03-04

└ Phrase entropy

Phrase entropy

x_i — parameter x of note i

Shannon entropy


$$H(X) := - \sum_i P(x_i) \log_2 P(x_i)$$

Probability distribution

$$P(x_i) = \frac{n_i}{N}$$

Li et al., 'Automatic Piano Reduction of Orchestral Music Based on Musical Entropy'.


Solution graph



Quantum Annealing for Music Arrangement

2025-03-04

└ Solution graph



Solution score

A musical score for three woodwind instruments: Flute, Oboe, and Bassoon. The score consists of four staves, each with a treble clef and a key signature of one flat. The Flute staff starts with a measure of eighth notes followed by a measure of sixteenth-note patterns. The Oboe staff follows with eighth-note patterns. The Bassoon staff has eighth-note patterns. Measures 5 through 14 show the instruments playing eighth-note patterns in various rhythmic patterns. Measure 15 shows the Flute and Oboe playing eighth-note patterns, while the Bassoon rests. Measure 16 shows the Flute and Oboe continuing their eighth-note patterns.

Quantum Annealing for Music Arrangement

2025-03-04

Solution score

Solution score

