

Lecture 5: Generalized Linear Models, MLP, and Back-Propagation

Tao LIN

March 16, 2023



Reading materials

- Chapter 3, Stanford CS 229 Lecture Notes,
https://cs229.stanford.edu/notes2022fall/main_notes.pdf
- Lecture 4, Stanford CS 231n, <http://cs231n.stanford.edu/schedule.html>

Reference

- EPFL, CS-433 Machine Learning, https://github.com/epfml/ML_course

Table of Contents

1 Review of Last Week

- Generalization Gap and Model Selection
- Bias-Variance Decomposition
- Logistic Regression

2 Exponential Families and Generalized Linear Models

3 Multi-Layer Perceptron (MLP) and Back-Propagation (BP)

Table of Contents

- 1 Review of Last Week
 - Generalization Gap and Model Selection
 - Bias-Variance Decomposition
 - Logistic Regression
- 2 Exponential Families and Generalized Linear Models
 - Motivation
 - Exponential family
 - Application in ML (Generalized Linear Models)
- 3 Multi-Layer Perceptron (MLP) and Back-Propagation (BP)
 - The Basic Structure of MLP
 - Training of NNs and BP

Generalization gap: How far is the test from the true error?

- **True Error:**

$$L_{\mathcal{D}}(f) = \mathbb{E}_{(y, \mathbf{x}) \sim \mathcal{D}} [\ell(y, f(\mathbf{x}))]. \quad (1)$$

- **Test/Empirical Error:**

$$L_{S_{\text{test}}}(f) = \frac{1}{|S_{\text{test}}|} \sum_{(\mathbf{x}_n, y_n) \in S_{\text{test}}} \ell(y_n, f(\mathbf{x}_n)). \quad (2)$$

- **Generalization Error:**

$$|L_{\mathcal{D}}(f) - L_{S_{\text{test}}}(f)|. \quad (3)$$

- **In Expectation:**

$$L_{\mathcal{D}}(f) = \mathbb{E}_{S_{\text{test}} \sim \mathcal{D}} [L_{S_{\text{test}}}(f)], \quad (4)$$

Generalization gap: How far is the test from the true error?

- **True Error:**

$$L_{\mathcal{D}}(f) = \mathbb{E}_{(y, \mathbf{x}) \sim \mathcal{D}} [\ell(y, f(\mathbf{x}))]. \quad (1)$$

- **Test/Empirical Error:**

$$L_{S_{\text{test}}}(f) = \frac{1}{|S_{\text{test}}|} \sum_{(\mathbf{x}_n, y_n) \in S_{\text{test}}} \ell(y_n, f(\mathbf{x}_n)). \quad (2)$$

- **Generalization Error:**

$$|L_{\mathcal{D}}(f) - L_{S_{\text{test}}}(f)|. \quad (3)$$

- **In Expectation:**

$$L_{\mathcal{D}}(f) = \mathbb{E}_{S_{\text{test}} \sim \mathcal{D}} [L_{S_{\text{test}}}(f)], \quad (4)$$

The variation of the $|L_{\mathcal{D}}(f) - L_{S_{\text{test}}}(f)|$ matters.

Theorem 1

Given a model f and a test set $S_{test} \sim \mathcal{D}$ i.i.d. (not used to learn f) and a loss $\ell(\cdot, \cdot) \in [a, b]$:

$$\Pr \left[|L_{\mathcal{D}}(f) - L_{S_{test}}(f)| \geq \sqrt{\frac{(b-a)^2 \ln(2/\delta)}{2|S_{test}|}} \right] \leq \delta \quad (5)$$

Theorem 1

Given a model f and a test set $S_{test} \sim \mathcal{D}$ i.i.d. (not used to learn f) and a loss $\ell(\cdot, \cdot) \in [a, b]$:

$$\Pr \left[|L_{\mathcal{D}}(f) - L_{S_{test}}(f)| \geq \sqrt{\frac{(b-a)^2 \ln(2/\delta)}{2|S_{test}|}} \right] \leq \delta \quad (5)$$

- The error decreases as $\mathcal{O}(1/\sqrt{|S_{test}|})$ with the number test points.

Theorem 1

Given a model f and a test set $S_{test} \sim \mathcal{D}$ i.i.d. (not used to learn f) and a loss $\ell(\cdot, \cdot) \in [a, b]$:

$$\Pr \left[|L_{\mathcal{D}}(f) - L_{S_{test}}(f)| \geq \sqrt{\frac{(b-a)^2 \ln(2/\delta)}{2|S_{test}|}} \right] \leq \delta \quad (5)$$

- The error decreases as $\mathcal{O}(1/\sqrt{|S_{test}|})$ with the number test points.
- \Rightarrow The more data points we have, the more confident we are that the empirical loss we measure is close to the true loss.

Theorem 1

Given a model f and a test set $S_{\text{test}} \sim \mathcal{D}$ i.i.d. (not used to learn f) and a loss $\ell(\cdot, \cdot) \in [a, b]$:

$$\Pr \left[|L_{\mathcal{D}}(f) - L_{S_{\text{test}}}(f)| \geq \sqrt{\frac{(b-a)^2 \ln(2/\delta)}{2 |S_{\text{test}}|}} \right] \leq \delta \quad (5)$$

- The error decreases as $\mathcal{O}(1/\sqrt{|S_{\text{test}}|})$ with the number test points.
- \Rightarrow The more data points we have, the more confident we are that the empirical loss we measure is close to the true loss.

Given a predictor f and a dataset S , we can control the expected risk:

$$\Pr \left[\underbrace{L_{\mathcal{D}}(f)}_{\text{not computable}} \geq \underbrace{L_{S_{\text{test}}}(f)}_{\text{computable}} + \underbrace{\sqrt{\frac{(b-a)^2 \ln(2/\delta)}{2 |S_{\text{test}}|}}}_{\text{deviation}} \right] \leq \delta. \quad (6)$$

How far is each of the K test errors $L_{S_{\text{test}}}(f_k)$ from the true $L_{\mathcal{D}}(f_k)$?

Theorem 2

We can bound the maximum deviation for all K candidates, by

$$\Pr \left[\max_k |L_{\mathcal{D}}(f_k) - L_{S_{\text{test}}}(f_k)| \geq \sqrt{\frac{(b-a)^2 \ln(2K/\delta)}{2|S_{\text{test}}|}} \right] \leq \delta \quad (7)$$

How far is each of the K test errors $L_{S_{\text{test}}}(f_k)$ from the true $L_{\mathcal{D}}(f_k)$?

Theorem 2

We can bound the maximum deviation for all K candidates, by

$$\Pr \left[\max_k |L_{\mathcal{D}}(f_k) - L_{S_{\text{test}}}(f_k)| \geq \sqrt{\frac{(b-a)^2 \ln(2K/\delta)}{2|S_{\text{test}}|}} \right] \leq \delta \quad (7)$$

- The error decreases as $\mathcal{O}(1/\sqrt{|S_{\text{test}}|})$ with the number test points.

How far is each of the K test errors $L_{S_{\text{test}}}(f_k)$ from the true $L_{\mathcal{D}}(f_k)$?

Theorem 2

We can bound the maximum deviation for all K candidates, by

$$\Pr \left[\max_k |L_{\mathcal{D}}(f_k) - L_{S_{\text{test}}}(f_k)| \geq \sqrt{\frac{(b-a)^2 \ln(2K/\delta)}{2|S_{\text{test}}|}} \right] \leq \delta \quad (7)$$

- The error decreases as $\mathcal{O}(1/\sqrt{|S_{\text{test}}|})$ with the number test points.
- When testing K hyper-parameters, the error only goes up by $\sqrt{\ln(K)}$.

How far is each of the K test errors $L_{S_{\text{test}}}(f_k)$ from the true $L_{\mathcal{D}}(f_k)$?

Theorem 2

We can bound the maximum deviation for all K candidates, by


$$\Pr \left[\max_k |L_{\mathcal{D}}(f_k) - L_{S_{\text{test}}}(f_k)| \geq \sqrt{\frac{(b-a)^2 \ln(2K/\delta)}{2|S_{\text{test}}|}} \right] \leq \delta \quad (7)$$

- The error decreases as $\mathcal{O}(1/\sqrt{|S_{\text{test}}|})$ with the number test points.
 - When testing K hyper-parameters, the error only goes up by $\sqrt{\ln(K)}$.
- ⇒ So we can test many different models without incurring a large penalty.

Issues: Splitting the data once into two parts (one for training and one for testing)
is not the most efficient way to use the data!

K-fold cross-validation:

- 1 Randomly partition the data into K groups
- 2 Train K times. Each time leave out exactly one of the K groups for testing and use the remaining $K - 1$ groups for training.
- 3 Average the K results




Benefits:

Issues: Splitting the data once into two parts (one for training and one for testing)
is not the most efficient way to use the data!

K-fold cross-validation:

- 1 Randomly partition the data into K groups
- 2 Train K times. Each time leave out exactly one of the K groups for testing and use the remaining $K - 1$ groups for training.
- 3 Average the K results




Benefits:

- We have used all data for training, and all data for testing, and used each data point the same number of times.

Issues: Splitting the data once into two parts (one for training and one for testing)
is not the most efficient way to use the data!

K-fold cross-validation:

- 1 Randomly partition the data into K groups
- 2 Train K times. Each time leave out exactly one of the K groups for testing and use the remaining $K - 1$ groups for training.
- 3 Average the K results



Benefits:

- We have used all data for training, and all data for testing, and used each data point the same number of times.
- Cross-validation returns an unbiased estimate of the generalization error and its variance.

Table of Contents

- 1 Review of Last Week
 - Generalization Gap and Model Selection
 - **Bias-Variance Decomposition**
 - Logistic Regression
- 2 Exponential Families and Generalized Linear Models
 - Motivation
 - Exponential family
 - Application in ML (Generalized Linear Models)
- 3 Multi-Layer Perceptron (MLP) and Back-Propagation (BP)
 - The Basic Structure of MLP
 - Training of NNs and BP

Bias-Variance Decomposition

$$\begin{aligned}\mathbb{E}_{S_{\text{train}} \in \mathcal{D}} [L(f_{S_{\text{train}}})] &= \text{Var}_{\epsilon \sim \mathcal{D}_\epsilon} [\epsilon] && \text{(noise variance)} \\ &+ (f(\mathbf{x}_0) - \mathbb{E}_{S_{\text{train}}} [f_{S_{\text{train}}}(\mathbf{x}_0)])^2 && \text{(Bias)} \\ &+ \mathbb{E}_{S_{\text{train}} \sim \mathcal{D}} \left[(\mathbb{E}_{S_{\text{train}}} [f_{S_{\text{train}}}(\mathbf{x}_0)] - f_{S_{\text{train}}}(\mathbf{x}_0))^2 \right], && \text{(Variance)}\end{aligned}$$

which always lower-bounds the true error.

Bias-Variance Decomposition

$$\begin{aligned}\mathbb{E}_{S_{\text{train}} \in \mathcal{D}} [L(f_{S_{\text{train}}})] &= \text{Var}_{\epsilon \sim \mathcal{D}_{\epsilon}} [\epsilon] && \text{(noise variance)} \\ &+ (f(\mathbf{x}_0) - \mathbb{E}_{S_{\text{train}}} [f_{S_{\text{train}}}(\mathbf{x}_0)])^2 && \text{(Bias)} \\ &+ \mathbb{E}_{S_{\text{train}} \sim \mathcal{D}} \left[(\mathbb{E}_{S_{\text{train}}} [f_{S_{\text{train}}}(\mathbf{x}_0)] - f_{S_{\text{train}}}(\mathbf{x}_0))^2 \right], && \text{(Variance)}\end{aligned}$$

which always lower-bounds the true error.

⇒ To minimize the true error, we need to select a method that **simultaneously achieves low bias and low variance**.

Double descent curve in Deep Learning




Table of Contents

1 Review of Last Week

- Generalization Gap and Model Selection
- Bias-Variance Decomposition
- Logistic Regression

2 Exponential Families and Generalized Linear Models

- Motivation
- Exponential family
- Application in ML (Generalized Linear Models)

3 Multi-Layer Perceptron (MLP) and Back-Propagation (BP)

- The Basic Structure of MLP
- Training of NNs and BP


The logistic function

Consider first of all the case of two classes.

The posterior probability for class \mathcal{C}_1 :

$$p(\mathcal{C}_1|\mathbf{x}) = \frac{p(\mathbf{x}|\mathcal{C}_1)p(\mathcal{C}_1)}{p(\mathbf{x}|\mathcal{C}_1)p(\mathcal{C}_1) + p(\mathbf{x}|\mathcal{C}_2)p(\mathcal{C}_2)} \quad (8)$$

$$= \frac{1}{1 + \exp(-\eta)} = \sigma(\eta) \quad (9)$$



Properties of the logistic function:

- $1 - \sigma(\eta) = \sigma(-\eta)$
- $\sigma'(\eta) = \sigma(\eta) (1 - \sigma(\eta))$

The logistic function

Consider first of all the case of two classes.


The posterior probability for class \mathcal{C}_1 :

$$p(\mathcal{C}_1|\mathbf{x}) = \frac{p(\mathbf{x}|\mathcal{C}_1)p(\mathcal{C}_1)}{p(\mathbf{x}|\mathcal{C}_1)p(\mathcal{C}_1) + p(\mathbf{x}|\mathcal{C}_2)p(\mathcal{C}_2)} \quad (8)$$

$$= \frac{1}{1 + \exp(-\eta)} = \sigma(\eta) \quad (9)$$

where we have defined

$$\eta = \ln \frac{p(\mathbf{x}|\mathcal{C}_1)p(\mathcal{C}_1)}{p(\mathbf{x}|\mathcal{C}_2)p(\mathcal{C}_2)} \text{ and } \sigma(\eta) := \frac{e^\eta}{1 + e^\eta} \quad (10)$$



Properties of the logistic function:

- $1 - \sigma(\eta) = \sigma(-\eta)$
- $\sigma'(\eta) = \sigma(\eta) (1 - \sigma(\eta))$

The logistic function

Consider first of all the case of two classes.

The posterior probability for class \mathcal{C}_1 :

$$p(\mathcal{C}_1|\mathbf{x}) = \frac{p(\mathbf{x}|\mathcal{C}_1)p(\mathcal{C}_1)}{p(\mathbf{x}|\mathcal{C}_1)p(\mathcal{C}_1) + p(\mathbf{x}|\mathcal{C}_2)p(\mathcal{C}_2)} \quad (8)$$


$$= \frac{1}{1 + \exp(-\eta)} = \sigma(\eta) \quad (9)$$

where we have defined

$$\eta = \ln \frac{p(\mathbf{x}|\mathcal{C}_1)p(\mathcal{C}_1)}{p(\mathbf{x}|\mathcal{C}_2)p(\mathcal{C}_2)} \text{ and } \sigma(\eta) := \frac{e^\eta}{1 + e^\eta} \quad (10)$$

For the case of $K > 2$ classes, we have

$$p(\mathcal{C}_k|\mathbf{x}) = \frac{p(\mathbf{x}|\mathcal{C}_k)p(\mathcal{C}_k)}{\sum_j p(\mathbf{x}|\mathcal{C}_j)p(\mathcal{C}_j)} = \frac{\exp(\eta_k)}{\sum_j \exp(\eta_j)} \quad (11)$$



Properties of the logistic function:

- $1 - \sigma(\eta) = \sigma(-\eta)$
- $\sigma'(\eta) = \sigma(\eta) (1 - \sigma(\eta))$

Logistic Regression

Given a “new” feature vector \mathbf{x} , we predict the (posterior) probability of the two class labels given \mathbf{x} by means of

$$p(1|\mathbf{x}) := \Pr[Y = 1|\mathbf{X} = \mathbf{x}] = \sigma(\mathbf{x}^\top \mathbf{w} + w_0) \quad (12)$$

$$p(0|\mathbf{x}) := \Pr[Y = 0|\mathbf{X} = \mathbf{x}] = 1 - \sigma(\mathbf{x}^\top \mathbf{w} + w_0) , \quad (13)$$

where we predict a real value (a probability) and not a label.

MLE is a method of estimating the parameters of a statistical model

The MLE finds the parameters \mathbf{w}^* under which $\{\mathbf{y}, \mathbf{X}\}$ are the most likely:

$$\mathbf{w}^* = \arg \max_{\mathbf{w}} \left(\mathcal{L}(\mathbf{w}) := \prod_{n=1}^N p(\{\mathbf{x}_n, y_n\} | \mathbf{w}) \right) = \arg \min_{\mathbf{w}} [-\log \mathcal{L}(\mathbf{w})] . \quad (14)$$

MLE is a method of estimating the parameters of a statistical model

The MLE finds the parameters \mathbf{w}^* under which $\{\mathbf{y}, \mathbf{X}\}$ are the most likely:

$$\mathbf{w}^* = \arg \max_{\mathbf{w}} \left(\mathcal{L}(\mathbf{w}) := \prod_{n=1}^N p(\{\mathbf{x}_n, y_n\} | \mathbf{w}) \right) = \arg \min_{\mathbf{w}} [-\log \mathcal{L}(\mathbf{w})] . \quad (14)$$

The likelihood of the data $\{\mathbf{y}, \mathbf{X}\}$ given the parameter \mathbf{w} , i.e., $p(\mathbf{y}, \mathbf{X} | \mathbf{w})$.

$$p(\mathbf{y}, \mathbf{X} | \mathbf{w}) = p(\mathbf{X} | \mathbf{w}) p(\mathbf{y} | \mathbf{X}, \mathbf{w}) = p(\mathbf{X}) p(\mathbf{y} | \mathbf{X}, \mathbf{w}) , \quad (15)$$

where \mathbf{X} does not depend on \mathbf{w} .

MLE for Logistic Regression

For Logistic Regression, we have:

$$p(\mathbf{y}|\mathbf{X}, \mathbf{w}) = \prod_{n=1}^N p(y_n|\mathbf{x}_n) = \prod_{n:y_n=1} p(y_n = 1|\mathbf{x}_n) \prod_{n:y_n=0} p(y_n = 0|\mathbf{x}_n) \quad (16)$$

$$= \prod_{n=1}^N \sigma(\mathbf{x}_n^\top \mathbf{w})^{y_n} [1 - \sigma(\mathbf{x}_n^\top \mathbf{w})]^{1-y_n} \quad (17)$$

MLE for Logistic Regression

For Logistic Regression, we have:

$$p(\mathbf{y}|\mathbf{X}, \mathbf{w}) = \prod_{n=1}^N p(y_n|\mathbf{x}_n) = \prod_{n:y_n=1} p(y_n = 1|\mathbf{x}_n) \prod_{n:y_n=0} p(y_n = 0|\mathbf{x}_n) \quad (16)$$

$$= \prod_{n=1}^N \sigma(\mathbf{x}_n^\top \mathbf{w})^{y_n} [1 - \sigma(\mathbf{x}_n^\top \mathbf{w})]^{1-y_n} \quad (17)$$

Minimizing $\mathcal{L}(\mathbf{w})$ through the property of stationary points.

$$\nabla \mathcal{L}(\mathbf{w}) = \frac{1}{N} \sum_{n=1}^N \mathbf{x}_n (\sigma(\mathbf{x}_n^\top \mathbf{w}) - y_n) = \frac{1}{N} \mathbf{X}^\top [\sigma(\mathbf{X}\mathbf{w}) - \mathbf{y}] , \quad (18)$$

where $\mathbf{X} \in \mathbb{R}^{N \times d}$. It has no closed-form solution to $\nabla \mathcal{L}(\mathbf{w}) = 0$.

Last lecture:

- Generalization Gap and Model Selection
- Bias-Variance Decomposition
- Before Introducing Multilayer Perceptron: Logistic Regression

Last lecture:

- Generalization Gap and Model Selection
- Bias-Variance Decomposition
- Before Introducing Multilayer Perceptron: Logistic Regression

This lecture:

- Exponential Families and Generalized Linear Models
- Multi-Layer Perceptron
- Back-Propagation

Table of Contents

- 1 Review of Last Week
- 2 Exponential Families and Generalized Linear Models**
 - Motivation
 - Exponential family
 - Application in ML (Generalized Linear Models)
- 3 Multi-Layer Perceptron (MLP) and Back-Propagation (BP)

Table of Contents


- 1 Review of Last Week
 - Generalization Gap and Model Selection
 - Bias-Variance Decomposition
 - Logistic Regression
- 2 Exponential Families and Generalized Linear Models
 - **Motivation**
 - Exponential family
 - Application in ML (Generalized Linear Models)
- 3 Multi-Layer Perceptron (MLP) and Back-Propagation (BP)
 - The Basic Structure of MLP
 - Training of NNs and BP

The Least-Squares can be defined in two different ways

- **Geometric way:**

Minimizing the sum of the squares of the residuals:

$$\hat{\mathbf{w}} = \arg \min_{\mathbf{w}} \frac{1}{2N} \sum_{n=1}^N (y_n - \mathbf{x}_n^\top \mathbf{w})^2 \quad (19)$$



The Least-Squares can be defined in two different ways

- **Geometric way:**


Minimizing the sum of the squares of the residuals:

$$\hat{\mathbf{w}} = \arg \min_{\mathbf{w}} \frac{1}{2N} \sum_{n=1}^N (y_n - \mathbf{x}_n^\top \mathbf{w})^2 \quad (19)$$

- **Probabilistic way:**

Assume the data follow a linear Gaussian model:

$$\mathbf{y} = \mathbf{x}^\top \mathbf{w} + \epsilon \text{ where } \epsilon \sim \mathcal{N}(0, \sigma^2) \quad (20)$$



The Least-Squares can be defined in two different ways

- **Geometric way:**

Minimizing the sum of the squares of the residuals:


$$\hat{\mathbf{w}} = \arg \min_{\mathbf{w}} \frac{1}{2N} \sum_{n=1}^N (y_n - \mathbf{x}_n^\top \mathbf{w})^2 \quad (19)$$

- **Probabilistic way:**

Assume the data follow a linear Gaussian model:


$$\mathbf{y} = \mathbf{x}^\top \mathbf{w} + \epsilon \text{ where } \epsilon \sim \mathcal{N}(0, \sigma^2) \quad (20)$$

Doing MLE recovers the LS estimator $\hat{\mathbf{w}}$.




How to get non-linear models?

- **Features augmentations:** add non-linear features
(x, x^2, x^3, \dots)




How to get non-linear models?

- **Features augmentations:** add non-linear features
(x, x^2, x^3, \dots)
- **Different probabilistic models:**




How to get non-linear models?

- **Features augmentations:** add non-linear features
(x, x^2, x^3, \dots)
- **Different probabilistic models:**
 - Least Squares: $y \sim \mathcal{N}(x^\top \mathbf{w}, \sigma^2)$




How to get non-linear models?

- **Features augmentations:** add non-linear features
(x, x^2, x^3, \dots)
- **Different probabilistic models:**
 - Least Squares: $y \sim \mathcal{N}(x^\top \mathbf{w}, \sigma^2)$
 \Rightarrow The linear model predicts the mean of a distribution μ
(from which the data are sampled).




How to get non-linear models?

- **Features augmentations:** add non-linear features
(x, x^2, x^3, \dots)
- **Different probabilistic models:**
 - Least Squares: $y \sim \mathcal{N}(x^\top \mathbf{w}, \sigma^2)$
 \Rightarrow The linear model predicts the mean of a distribution μ
(from which the data are sampled).
 - Logistic Regression: $y \sim \mathcal{B}(\sigma(x^\top \mathbf{w}))$



How to get non-linear models?

- **Features augmentations:** add non-linear features
($\mathbf{x}, \mathbf{x}^2, \mathbf{x}^3, \dots$)
- **Different probabilistic models:**
 - Least Squares: $y \sim \mathcal{N}(\mathbf{x}^\top \mathbf{w}, \sigma^2)$
 \Rightarrow The linear model predicts the mean of a distribution μ
(from which the data are sampled).
 - Logistic Regression: $y \sim \mathcal{B}(\sigma(\mathbf{x}^\top \mathbf{w}))$
 \Rightarrow The linear model predicts another quantity $\eta := \mathbf{x}^\top \mathbf{w}$.



Recall the definition of Logistic Regression

Logistic Regression models the probability of the two classes $\{0, 1\}$ by

$$p(1|\eta) = \sigma(\eta) \text{ and } p(0|\eta) = 1 - \sigma(\eta), \quad (21)$$

where $\eta = \mathbf{x}^\top \mathbf{w}$.

Recall the definition of Logistic Regression

Logistic Regression models the probability of the two classes $\{0, 1\}$ by

$$p(1|\eta) = \sigma(\eta) \text{ and } p(0|\eta) = 1 - \sigma(\eta), \quad (21)$$

where $\eta = \mathbf{x}^\top \mathbf{w}$. This can be compactly written as

$$p(y|\eta) = \frac{e^{\eta y}}{1 + e^\eta} = \exp(\eta y - \ln(1 + e^\eta)) \quad (22)$$

Recall the definition of Logistic Regression

Logistic Regression models the probability of the two classes $\{0, 1\}$ by

$$p(1|\eta) = \sigma(\eta) \text{ and } p(0|\eta) = 1 - \sigma(\eta), \quad (21)$$

where $\eta = \mathbf{x}^\top \mathbf{w}$. This can be compactly written as

$$p(y|\eta) = \frac{e^{\eta y}}{1 + e^\eta} = \exp(\eta y - \ln(1 + e^\eta)) \quad (22)$$

- The linear model predicts $\eta := \sigma(\eta)$ which is not the mean of the distribution.

Recall the definition of Logistic Regression

Logistic Regression models the probability of the two classes $\{0, 1\}$ by

$$p(1|\eta) = \sigma(\eta) \text{ and } p(0|\eta) = 1 - \sigma(\eta), \quad (21)$$

where $\eta = \mathbf{x}^\top \mathbf{w}$. This can be compactly written as

$$p(y|\eta) = \frac{e^{\eta y}}{1 + e^\eta} = \exp(\eta y - \ln(1 + e^\eta)) \quad (22)$$

- The linear model predicts $\eta := \sigma(\eta)$ which is not the mean of the distribution.
- η is related to the mean μ by the non-linear relation

Recall the definition of Logistic Regression

Logistic Regression models the probability of the two classes $\{0, 1\}$ by

$$p(1|\eta) = \sigma(\eta) \text{ and } p(0|\eta) = 1 - \sigma(\eta), \quad (21)$$

where $\eta = \mathbf{x}^\top \mathbf{w}$. This can be compactly written as

$$p(y|\eta) = \frac{e^{\eta y}}{1 + e^\eta} = \exp(\eta y - \ln(1 + e^\eta)) \quad (22)$$

- The linear model predicts $\eta := \sigma(\eta)$ which is not the mean of the distribution.
- η is related to the mean μ by the non-linear relation

\Rightarrow The *link function*:

Recall the definition of Logistic Regression

Logistic Regression models the probability of the two classes $\{0, 1\}$ by

$$p(1|\eta) = \sigma(\eta) \text{ and } p(0|\eta) = 1 - \sigma(\eta), \quad (21)$$

where $\eta = \mathbf{x}^\top \mathbf{w}$. This can be compactly written as

$$p(y|\eta) = \frac{e^{\eta y}}{1 + e^\eta} = \exp(\eta y - \ln(1 + e^\eta)) \quad (22)$$

- The linear model predicts $\eta := \sigma(\eta)$ which is not the mean of the distribution.
- η is related to the mean μ by the non-linear relation

⇒ The *link function*:
the relation between (1) the value η we predict by the linear model and (2) the mean μ .

A unified framework

The distribution used in Logistic Regression can be written in a very specific form:

$$p(y|\eta) = \frac{e^{\eta y}}{1 + e^{\eta}} = \exp(\eta y - \ln(1 + e^{\eta})) \quad (23)$$

A unified framework

The distribution used in Logistic Regression can be written in a very specific form:

$$p(y|\eta) = \frac{e^{\eta y}}{1 + e^{\eta}} = \exp(\eta y - \ln(1 + e^{\eta})) \quad (23)$$

Goals: a unified framework to generalize other forms of distributions.

A unified framework

The distribution used in Logistic Regression can be written in a very specific form:

$$p(y|\eta) = \frac{e^{\eta y}}{1 + e^{\eta}} = \exp(\eta y - \ln(1 + e^{\eta})) \quad (23)$$

Goals: a unified framework to generalize other forms of distributions.

- The discussion on a class of distributions, known as *exponential families*.

A unified framework

The distribution used in Logistic Regression can be written in a very specific form:

$$p(y|\eta) = \frac{e^{\eta y}}{1 + e^{\eta}} = \exp(\eta y - \ln(1 + e^{\eta})) \quad (23)$$

Goals: a unified framework to generalize other forms of distributions.

- The discussion on a class of distributions, known as *exponential families*.
- Many distributions (but not all) fit into this framework and that distributions in this family have many nice properties.

Table of Contents

- 1 Review of Last Week
 - Generalization Gap and Model Selection
 - Bias-Variance Decomposition
 - Logistic Regression
- 2 Exponential Families and Generalized Linear Models
 - Motivation
 - Exponential family
 - Application in ML (Generalized Linear Models)
- 3 Multi-Layer Perceptron (MLP) and Back-Propagation (BP)
 - The Basic Structure of MLP
 - Training of NNs and BP

Exponential family — Definition

A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = \underbrace{h(y)}_{\geq 0} \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (24)$$

¹Assume that we are given independent samples from this distribution. We do know $\boldsymbol{\phi}(y)$ and $h(y)$ but not $\boldsymbol{\eta}$. In order to optimally estimate $\boldsymbol{\eta}$ given these samples, all we need is the empirical average of the $\boldsymbol{\phi}(y)$.

Exponential family — Definition

A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = \underbrace{h(y)}_{\geq 0} \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (24)$$

- natural or canonical parameter of the distribution $\boldsymbol{\eta}$

¹Assume that we are given independent samples from this distribution. We do know $\boldsymbol{\phi}(y)$ and $h(y)$ but not $\boldsymbol{\eta}$. In order to optimally estimate $\boldsymbol{\eta}$ given these samples, all we need is the empirical average of the $\boldsymbol{\phi}(y)$.

Exponential family — Definition

A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = \underbrace{h(y)}_{\geq 0} \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (24)$$

- natural or canonical parameter of the distribution $\boldsymbol{\eta}$
- sufficient statistics¹ $\boldsymbol{\phi}(y)$ contains all the relevant information

¹Assume that we are given independent samples from this distribution. We do know $\boldsymbol{\phi}(y)$ and $h(y)$ but not $\boldsymbol{\eta}$. In order to optimally estimate $\boldsymbol{\eta}$ given these samples, all we need is the empirical average of the $\boldsymbol{\phi}(y)$.

Exponential family — Definition

A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = \underbrace{h(y)}_{\geq 0} \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (24)$$

- natural or canonical parameter of the distribution $\boldsymbol{\eta}$
- sufficient statistics¹ $\boldsymbol{\phi}(y)$ contains all the relevant information
- $A(\boldsymbol{\eta})$: log partition, the quantity $e^{-A(\boldsymbol{\eta})}$ is used as a normalization constant:

¹Assume that we are given independent samples from this distribution. We do know $\boldsymbol{\phi}(y)$ and $h(y)$ but not $\boldsymbol{\eta}$. In order to optimally estimate $\boldsymbol{\eta}$ given these samples, all we need is the empirical average of the $\boldsymbol{\phi}(y)$.

Exponential family — Definition

A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = \underbrace{h(y)}_{\geq 0} \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (24)$$

- natural or canonical parameter of the distribution $\boldsymbol{\eta}$
- sufficient statistics¹ $\boldsymbol{\phi}(y)$ contains all the relevant information
- $A(\boldsymbol{\eta})$: log partition, the quantity $e^{-A(\boldsymbol{\eta})}$ is used as a normalization constant:

$$\int p(y|\boldsymbol{\eta}) dy = \int h(y) \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] dy = 1 \quad (25)$$

¹Assume that we are given independent samples from this distribution. We do know $\boldsymbol{\phi}(y)$ and $h(y)$ but not $\boldsymbol{\eta}$. In order to optimally estimate $\boldsymbol{\eta}$ given these samples, all we need is the empirical average of the $\boldsymbol{\phi}(y)$.

Exponential family — Definition

A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = \underbrace{h(y)}_{\geq 0} \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (24)$$

- natural or canonical parameter of the distribution $\boldsymbol{\eta}$
- sufficient statistics¹ $\boldsymbol{\phi}(y)$ contains all the relevant information
- $A(\boldsymbol{\eta})$: log partition, the quantity $e^{-A(\boldsymbol{\eta})}$ is used as a normalization constant:

$$\int p(y|\boldsymbol{\eta}) dy = \int h(y) \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] dy = 1 \quad (25)$$

$$\implies \int h(y) \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y)] dy = \int h(y) \exp [A(\boldsymbol{\eta})] dy = \exp [A(\boldsymbol{\eta})].$$

¹Assume that we are given independent samples from this distribution. We do know $\boldsymbol{\phi}(y)$ and $h(y)$ but not $\boldsymbol{\eta}$. In order to optimally estimate $\boldsymbol{\eta}$ given these samples, all we need is the empirical average of the $\boldsymbol{\phi}(y)$.

A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = \underbrace{h(y)}_{\geq 0} \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (26)$$

A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = \underbrace{h(y)}_{\geq 0} \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (26)$$

- A fixed choice of $\boldsymbol{\phi}(y)$, $A(\boldsymbol{\eta})$ and $h(y)$ defines a family of distributions (parameterized by $\boldsymbol{\eta}$).

A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = \underbrace{h(y)}_{\geq 0} \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (26)$$

- A fixed choice of $\boldsymbol{\phi}(y)$, $A(\boldsymbol{\eta})$ and $h(y)$ defines a family of distributions (parameterized by $\boldsymbol{\eta}$).
- As we vary $\boldsymbol{\eta}$, we then get different distribution within this family.

A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = \underbrace{h(y)}_{\geq 0} \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (26)$$

- A fixed choice of $\boldsymbol{\phi}(y)$, $A(\boldsymbol{\eta})$ and $h(y)$ defines a family of distributions (parameterized by $\boldsymbol{\eta}$).
- As we vary $\boldsymbol{\eta}$, we then get different distribution within this family.
- For some parameters $\boldsymbol{\eta}$, there exists some $h(y) \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y)]$ that cannot be normalized.

A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = \underbrace{h(y)}_{\geq 0} \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (26)$$

- A fixed choice of $\boldsymbol{\phi}(y)$, $A(\boldsymbol{\eta})$ and $h(y)$ defines a family of distributions (parameterized by $\boldsymbol{\eta}$).
- As we vary $\boldsymbol{\eta}$, we then get different distribution within this family.
- For some parameters $\boldsymbol{\eta}$, there exists some $h(y) \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y)]$ that cannot be normalized.

For example, $h(y) = 1$, $\boldsymbol{\phi}(y) = y^2$ and $\boldsymbol{\eta} = 1$.

A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = \underbrace{h(y)}_{\geq 0} \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (26)$$

- A fixed choice of $\boldsymbol{\phi}(y)$, $A(\boldsymbol{\eta})$ and $h(y)$ defines a family of distributions (parameterized by $\boldsymbol{\eta}$).
- As we vary $\boldsymbol{\eta}$, we then get different distribution within this family.
- For some parameters $\boldsymbol{\eta}$, there exists some $h(y) \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y)]$ that cannot be normalized.

For example, $h(y) = 1$, $\boldsymbol{\phi}(y) = y^2$ and $\boldsymbol{\eta} = 1$.

We will exclude such parameters by only looking at the set of parameters

$$M := \left\{ \boldsymbol{\eta} : \int_y h(y) \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y)] dy < \infty \right\} \quad (27)$$

Why?

Bernoulli distributions belong to the exponential family

Recall: A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = h(y) \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (28)$$

Bernoulli distributions belong to the exponential family

Recall: A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = h(y) \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (28)$$

The Bernoulli distribution is the binary random variable such that for $\mu \in [0, 1]$:

$$\Pr(Y = 1) = \mu \quad \text{and} \quad \Pr(Y = 0) = 1 - \mu \quad (29)$$

Bernoulli distributions belong to the exponential family

Recall: A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = h(y) \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (28)$$

The Bernoulli distribution is the binary random variable such that for $\mu \in [0, 1]$:

$$\Pr(Y = 1) = \mu \quad \text{and} \quad \Pr(Y = 0) = 1 - \mu \quad (29)$$

Claim: the Bernoulli distribution is a member of the exponential family.

$$p(y|\mu) = \mu^y (1 - \mu)^{1-y}, \text{ where } \mu \in (0, 1) \quad (30)$$

$$= \exp \left\{ \left(\ln \frac{\mu}{1 - \mu} \right) y + \ln(1 - \mu) \right\} = \exp \{ \eta \phi(y) - A(\eta) \}. \quad (31)$$

Bernoulli distributions belong to the exponential family

Recall: A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = h(y) \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (28)$$

The Bernoulli distribution is the binary random variable such that for $\mu \in [0, 1]$:

$$\Pr(Y = 1) = \mu \quad \text{and} \quad \Pr(Y = 0) = 1 - \mu \quad (29)$$

Claim: the Bernoulli distribution is a member of the exponential family.

$$p(y|\mu) = \mu^y (1 - \mu)^{1-y}, \text{ where } \mu \in (0, 1) \quad (30)$$

$$= \exp \left\{ \left(\ln \frac{\mu}{1 - \mu} \right) y + \ln(1 - \mu) \right\} = \exp \{ \eta \phi(y) - A(\eta) \}. \quad (31)$$

where we can identify:

$$\phi(y) = y, \quad \eta = \ln \frac{\mu}{1 - \mu}, \quad A(\eta) = -\ln(1 - \mu) = \ln(1 + e^\eta), \quad h(y) = 1. \quad (32)$$

Bernoulli distributions belong to the exponential family

Recall: A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = h(y) \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (28)$$

The Bernoulli distribution is the binary random variable such that for $\mu \in [0, 1]$:

$$\Pr(Y = 1) = \mu \quad \text{and} \quad \Pr(Y = 0) = 1 - \mu \quad (29)$$

Claim: the Bernoulli distribution is a member of the exponential family.

$$p(y|\mu) = \mu^y (1 - \mu)^{1-y}, \text{ where } \mu \in (0, 1) \quad (30)$$

$$= \exp \left\{ \left(\ln \frac{\mu}{1 - \mu} \right) y + \ln(1 - \mu) \right\} = \exp \{ \eta \phi(y) - A(\eta) \}. \quad (31)$$

where we can identify:

$$\boldsymbol{\phi}(y) = y, \quad \eta = \ln \frac{\mu}{1 - \mu}, \quad A(\eta) = -\ln(1 - \mu) = \ln(1 + e^\eta), \quad h(y) = 1. \quad (32)$$

$$\Rightarrow \eta = g(\mu) = \ln \frac{\mu}{1 - \mu} \iff \mu = g^{-1}(\eta) = \frac{e^\eta}{1 + e^\eta}, \text{ and } g(\mu) \text{ links the mean of } \boldsymbol{\phi}(y) \text{ to } \boldsymbol{\eta}.$$

Gaussian distributions belong to the exponential family

Recall: A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = h(y) \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (33)$$

Gaussian distributions belong to the exponential family

Recall: A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = h(y) \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (33)$$

Claim: the Gaussian distribution with mean μ and variance σ^2 is also a member of the exponential family.

Gaussian distributions belong to the exponential family

Recall: A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = h(y) \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (33)$$

Claim: the Gaussian distribution with mean μ and variance σ^2 is also a member of the exponential family.

$$p(y|\mu) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(y-\mu)^2}{2\sigma^2}}, \quad \mu \in \mathbb{R}, \sigma^2 \in \mathbb{R}^+ \quad (34)$$

$$= \exp \left[(\mu/\sigma^2, -1/(2\sigma^2))(y, y^2)^\top - \frac{\mu^2}{2\sigma^2} - \frac{1}{2} \ln(2\pi\sigma^2) \right]. \quad (35)$$

Gaussian distributions belong to the exponential family

Recall: A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = h(y) \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (33)$$

Claim: the Gaussian distribution with mean μ and variance σ^2 is also a member of the exponential family.

$$p(y|\mu) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(y-\mu)^2}{2\sigma^2}}, \quad \mu \in \mathbb{R}, \sigma^2 \in \mathbb{R}^+ \quad (34)$$

$$= \exp \left[(\mu/\sigma^2, -1/(2\sigma^2))(y, y^2)^\top - \frac{\mu^2}{2\sigma^2} - \frac{1}{2} \ln(2\pi\sigma^2) \right]. \quad (35)$$

$$\boldsymbol{\phi}(y) = (y, y^2)^\top, \quad \boldsymbol{\eta} = (\eta_1 = \mu/\sigma^2, \eta_2 = -1/(2\sigma^2))^\top, \quad (36)$$

$$A(\boldsymbol{\eta}) = -\frac{\eta_1^2}{4\eta_2} - \frac{1}{2} \ln(-\eta_2/\pi), \quad h(y) = 1. \quad (37)$$

Gaussian distributions belong to the exponential family

Recall: A distribution belongs to the exponential family if it can be written in the form

$$p(y|\boldsymbol{\eta}) = h(y) \exp [\boldsymbol{\eta}^\top \boldsymbol{\phi}(y) - A(\boldsymbol{\eta})] \quad (33)$$

Claim: the Gaussian distribution with mean μ and variance σ^2 is also a member of the exponential family.

$$p(y|\mu) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(y-\mu)^2}{2\sigma^2}}, \quad \mu \in \mathbb{R}, \sigma^2 \in \mathbb{R}^+ \quad (34)$$

$$= \exp \left[(\mu/\sigma^2, -1/(2\sigma^2))(y, y^2)^\top - \frac{\mu^2}{2\sigma^2} - \frac{1}{2} \ln(2\pi\sigma^2) \right]. \quad (35)$$

$$\boldsymbol{\phi}(y) = (y, y^2)^\top, \quad \boldsymbol{\eta} = (\eta_1 = \mu/\sigma^2, \eta_2 = -1/(2\sigma^2))^\top, \quad (36)$$

$$A(\boldsymbol{\eta}) = -\frac{\eta_1^2}{4\eta_2} - \frac{1}{2} \ln(-\eta_2/\pi), \quad h(y) = 1. \quad (37)$$

Link function ($\eta := g(\mu)$): $\eta_1 = \frac{\mu}{\sigma^2}, \eta_2 = -\frac{1}{2\sigma^2} \iff \mu = -\frac{\eta_1}{2\eta_2}, \sigma^2 = -\frac{1}{2\eta_2}.$

Basic properties

- Cumulant $A(\boldsymbol{\eta})$ is convex.

Basic properties

- Cumulant $A(\boldsymbol{\eta})$ is convex.
- $\nabla A(\boldsymbol{\eta}) = \mathbb{E}[\boldsymbol{\phi}(y)]$

Basic properties

- Cumulant $A(\boldsymbol{\eta})$ is convex.
- $\nabla A(\boldsymbol{\eta}) = \mathbb{E} [\boldsymbol{\phi}(\boldsymbol{y})]$
- $\nabla^2 A(\boldsymbol{\eta}) = \mathbb{E} [\boldsymbol{\phi}(\boldsymbol{y})\boldsymbol{\phi}(\boldsymbol{y})^\top] - \mathbb{E} [\boldsymbol{\phi}(\boldsymbol{y})] \mathbb{E} [\boldsymbol{\phi}(\boldsymbol{y})]^\top$

Basic properties

- Cumulant $A(\boldsymbol{\eta})$ is convex.
- $\nabla A(\boldsymbol{\eta}) = \mathbb{E} [\boldsymbol{\phi}(y)]$
- $\nabla^2 A(\boldsymbol{\eta}) = \mathbb{E} [\boldsymbol{\phi}(y) \boldsymbol{\phi}(y)^\top] - \mathbb{E} [\boldsymbol{\phi}(y)] \mathbb{E} [\boldsymbol{\phi}(y)]^\top$
- There is a 1 – 1 relationship between the “mean” $\boldsymbol{\mu} := \mathbb{E} [\boldsymbol{\phi}(y)]$ and natural parameter $\boldsymbol{\eta}$, defined using a so-called *link function* \mathbf{g} :

Basic properties

- Cumulant $A(\boldsymbol{\eta})$ is convex.
- $\nabla A(\boldsymbol{\eta}) = \mathbb{E} [\boldsymbol{\phi}(y)]$
- $\nabla^2 A(\boldsymbol{\eta}) = \mathbb{E} [\boldsymbol{\phi}(y)\boldsymbol{\phi}(y)^\top] - \mathbb{E} [\boldsymbol{\phi}(y)] \mathbb{E} [\boldsymbol{\phi}(y)]^\top$
- There is a 1 – 1 relationship between the “mean” $\boldsymbol{\mu} := \mathbb{E} [\boldsymbol{\phi}(y)]$ and natural parameter $\boldsymbol{\eta}$, defined using a so-called *link function* \mathbf{g} :

$$\boldsymbol{\eta} = \mathbf{g}(\boldsymbol{\mu} := \mathbb{E} [\boldsymbol{\phi}(y)]) \iff \boldsymbol{\mu} = \mathbf{g}^{-1}(\boldsymbol{\eta}) = \nabla A(\boldsymbol{\eta}) \quad (38)$$

Table of Contents

- 1 Review of Last Week
 - Generalization Gap and Model Selection
 - Bias-Variance Decomposition
 - Logistic Regression
- 2 Exponential Families and Generalized Linear Models
 - Motivation
 - Exponential family
 - Application in ML (Generalized Linear Models)
- 3 Multi-Layer Perceptron (MLP) and Back-Propagation (BP)
 - The Basic Structure of MLP
 - Training of NNs and BP

Maximum Likelihood Estimation (MLE)

Assume a set of iid samples $\{y_n\}_{n=1}^N$, sampled from a member of the exponential family with given $h(y)$, sufficient statistics $\phi(y)$, but unknown parameter η .

Maximum Likelihood Estimation (MLE)

Assume a set of iid samples $\{y_n\}_{n=1}^N$, sampled from a member of the exponential family with given $h(y)$, sufficient statistics $\phi(y)$, but unknown parameter η .

Goal: Estimate the natural parameter η .

Maximum Likelihood Estimation (MLE)

Assume a set of iid samples $\{y_n\}_{n=1}^N$, sampled from a member of the exponential family with given $h(y)$, sufficient statistics $\phi(y)$, but unknown parameter η .

Goal: Estimate the natural parameter η .

How: MLE for $p(y|\eta) = h(y) \exp [\eta^\top \phi(y) - A(\eta)]$.

Maximum Likelihood Estimation (MLE)

Assume a set of iid samples $\{y_n\}_{n=1}^N$, sampled from a member of the exponential family with given $h(y)$, sufficient statistics $\phi(y)$, but unknown parameter η .

Goal: Estimate the natural parameter η .

How: MLE for $p(y|\eta) = h(y) \exp [\eta^\top \phi(y) - A(\eta)]$.

$$\mathcal{L}(\eta) = -\frac{1}{N} \ln (p(y|\eta)) \tag{39}$$

$$= \frac{1}{N} \sum_{n=1}^N [-\ln (h(y_n)) - \eta^\top \phi(y_n) + A(\eta)] . \tag{40}$$

Maximum Likelihood Estimation (MLE)

Assume a set of iid samples $\{y_n\}_{n=1}^N$, sampled from a member of the exponential family with given $h(y)$, sufficient statistics $\phi(y)$, but unknown parameter η .

Goal: Estimate the natural parameter η .

How: MLE for $p(y|\eta) = h(y) \exp [\eta^\top \phi(y) - A(\eta)]$.

$$\mathcal{L}(\eta) = -\frac{1}{N} \ln (p(y|\eta)) \quad (39)$$

$$= \frac{1}{N} \sum_{n=1}^N [-\ln (h(y_n)) - \eta^\top \phi(y_n) + A(\eta)] . \quad (40)$$

\Rightarrow The cost function \mathcal{L} is a convex function in η since $A(\eta)$ is convex.

Given the definition

$$\mathcal{L}(\boldsymbol{\eta}) = \frac{1}{N} \sum_{n=1}^N [-\ln(h(y_n)) - \boldsymbol{\eta}^\top \boldsymbol{\phi}(y_n) + A(\boldsymbol{\eta})] \quad (41)$$

²It says that we should pick $\boldsymbol{\eta}$ s.t. the expected value of the sufficient statistics is equal to its empirical value!

Given the definition

$$\mathcal{L}(\boldsymbol{\eta}) = \frac{1}{N} \sum_{n=1}^N [-\ln(h(y_n)) - \boldsymbol{\eta}^\top \boldsymbol{\phi}(y_n) + A(\boldsymbol{\eta})] \quad (41)$$

Gradient:

$$\nabla_{\boldsymbol{\eta}} \mathcal{L}(\boldsymbol{\eta}) = -\frac{1}{N} \sum_{n=1}^N \boldsymbol{\phi}(y_n) + \mathbb{E}[\boldsymbol{\phi}(y)] , \quad (42)$$

²It says that we should pick $\boldsymbol{\eta}$ s.t. the expected value of the sufficient statistics is equal to its empirical value!

Given the definition

$$\mathcal{L}(\boldsymbol{\eta}) = \frac{1}{N} \sum_{n=1}^N [-\ln(h(y_n)) - \boldsymbol{\eta}^\top \boldsymbol{\phi}(y_n) + A(\boldsymbol{\eta})] \quad (41)$$

Gradient:

$$\nabla_{\boldsymbol{\eta}} \mathcal{L}(\boldsymbol{\eta}) = -\frac{1}{N} \sum_{n=1}^N \boldsymbol{\phi}(y_n) + \mathbb{E}[\boldsymbol{\phi}(y)] , \quad (42)$$

Stationary point:²

²It says that we should pick $\boldsymbol{\eta}$ s.t. the expected value of the sufficient statistics is equal to its empirical value!

Given the definition

$$\mathcal{L}(\boldsymbol{\eta}) = \frac{1}{N} \sum_{n=1}^N [-\ln(h(y_n)) - \boldsymbol{\eta}^\top \boldsymbol{\phi}(y_n) + A(\boldsymbol{\eta})] \quad (41)$$

Gradient:

$$\nabla_{\boldsymbol{\eta}} \mathcal{L}(\boldsymbol{\eta}) = -\frac{1}{N} \sum_{n=1}^N \boldsymbol{\phi}(y_n) + \mathbb{E}[\boldsymbol{\phi}(y)] , \quad (42)$$

Stationary point:²

$$\boldsymbol{\mu} := \mathbb{E}[\boldsymbol{\phi}(y)] = \frac{1}{N} \sum_{n=1}^N \boldsymbol{\phi}(y_n) , \quad (43)$$

²It says that we should pick $\boldsymbol{\eta}$ s.t. the expected value of the sufficient statistics is equal to its empirical value!

Given the definition

$$\mathcal{L}(\boldsymbol{\eta}) = \frac{1}{N} \sum_{n=1}^N [-\ln(h(y_n)) - \boldsymbol{\eta}^\top \boldsymbol{\phi}(y_n) + A(\boldsymbol{\eta})] \quad (41)$$

Gradient:

$$\nabla_{\boldsymbol{\eta}} \mathcal{L}(\boldsymbol{\eta}) = -\frac{1}{N} \sum_{n=1}^N \boldsymbol{\phi}(y_n) + \mathbb{E}[\boldsymbol{\phi}(y)] , \quad (42)$$

Stationary point:²

$$\boldsymbol{\mu} := \mathbb{E}[\boldsymbol{\phi}(y)] = \frac{1}{N} \sum_{n=1}^N \boldsymbol{\phi}(y_n) , \quad (43)$$

Closed-form: assume we have determined the link function $\mathbf{g}(\boldsymbol{\mu}) = \boldsymbol{\eta}$

$$\boldsymbol{\eta} = \mathbf{g} \left(\frac{1}{N} \sum_{n=1}^N \boldsymbol{\phi}(y_n) \right) , \quad (44)$$

and justify why we called $\boldsymbol{\phi}(y)$ a sufficient statistics.

²It says that we should pick $\boldsymbol{\eta}$ s.t. the expected value of the sufficient statistics is equal to its empirical value!

Generalized Linear Models (GLM)

Scenario:

Generalized Linear Models (GLM)

Scenario:

- We would like to build a model to estimate the number y of customers arriving in a store.

Generalized Linear Models (GLM)

Scenario:

- We would like to build a model to estimate the number y of customers arriving in a store.
- The Poisson distribution usually gives a good model for the number of visitors.

Generalized Linear Models (GLM)

Scenario:

- We would like to build a model to estimate the number y of customers arriving in a store.
- The Poisson distribution usually gives a good model for the number of visitors.
- How can we come up with a model for our problem?

Generalized Linear Models (GLM)

Scenario:

- We would like to build a model to estimate the number y of customers arriving in a store.
- The Poisson distribution usually gives a good model for the number of visitors.
- How can we come up with a model for our problem?
- Fortunately the Poisson is an exponential family distribution.

Generalized Linear Models (GLM)

Scenario:

- We would like to build a model to estimate the number y of customers arriving in a store.
- The Poisson distribution usually gives a good model for the number of visitors.
- How can we come up with a model for our problem?
- Fortunately the Poisson is an exponential family distribution.

We can apply a Generalized Linear Model (GLM)!

Constructing GLMs

To derive a GLM for a classification/regression problem (the conditional dist. of y given x):

Constructing GLMs

To derive a GLM for a classification/regression problem (the conditional dist. of y given \mathbf{x}):

- 1 The natural parameter η and the observed inputs \mathbf{x} are related linearly: $\eta = \mathbf{x}^\top \mathbf{w}$

Constructing GLMs

To derive a GLM for a classification/regression problem (the conditional dist. of y given \mathbf{x}):

- 1 The natural parameter η and the observed inputs \mathbf{x} are related linearly: $\eta = \mathbf{x}^\top \mathbf{w}$
- 2 The conditional mean μ is represented as a function $f(\eta)$ of the linear combination η

Constructing GLMs

To derive a GLM for a classification/regression problem (the conditional dist. of y given \mathbf{x}):

- 1 The natural parameter η and the observed inputs \mathbf{x} are related linearly: $\eta = \mathbf{x}^\top \mathbf{w}$
- 2 The conditional mean μ is represented as a function $f(\eta)$ of the linear combination η
- 3 The observed output y is assumed to be characterized by an exponential family distribution with conditional mean μ , i.e.,

Constructing GLMs

To derive a GLM for a classification/regression problem (the conditional dist. of y given \mathbf{x}):

- 1 The natural parameter η and the observed inputs \mathbf{x} are related linearly: $\eta = \mathbf{x}^\top \mathbf{w}$
- 2 The conditional mean μ is represented as a function $f(\eta)$ of the linear combination η
- 3 The observed output y is assumed to be characterized by an exponential family distribution with conditional mean μ , i.e.,

The condition probability is thus modeled as:

$$p(y|\mathbf{x}; \mathbf{w}) = h(y_n) \exp(\eta \phi(y) - A(\eta)) \quad \text{for } \eta = g \circ f(\mathbf{x}^\top \mathbf{w}) \quad (45)$$

Negative log-likelihood estimation:

$$\mathcal{L}(\mathbf{w}) = -\frac{1}{N} \sum_{n=1}^N \ln p(y_n | \mathbf{x}_n^\top \mathbf{w}) \quad (46)$$

$$= -\frac{1}{N} \sum_{n=1}^N (\ln(h(y_n)) + \eta_n \phi(y_n) - A(\eta_n)) \quad (47)$$

Negative log-likelihood estimation:

$$\mathcal{L}(\mathbf{w}) = -\frac{1}{N} \sum_{n=1}^N \ln p(y_n | \mathbf{x}_n^\top \mathbf{w}) \quad (46)$$

$$= -\frac{1}{N} \sum_{n=1}^N (\ln(h(y_n)) + \eta_n \phi(y_n) - A(\eta_n)) \quad (47)$$

If we rewrite this sum by using the matrix notation, we get

$$\nabla_{\mathbf{w}} \mathcal{L}(\mathbf{w}) = -\frac{1}{N} \sum_{n=1}^N \mathbf{x}_n \phi(y_n) - \nabla_{\mathbf{w}} A(\eta_n) \quad (48)$$

Negative log-likelihood estimation:

$$\mathcal{L}(\mathbf{w}) = -\frac{1}{N} \sum_{n=1}^N \ln p(y_n | \mathbf{x}_n^\top \mathbf{w}) \quad (46)$$

$$= -\frac{1}{N} \sum_{n=1}^N (\ln(h(y_n)) + \eta_n \phi(y_n) - A(\eta_n)) \quad (47)$$

If we rewrite this sum by using the matrix notation, we get

$$\nabla_{\mathbf{w}} \mathcal{L}(\mathbf{w}) = -\frac{1}{N} \sum_{n=1}^N \mathbf{x}_n \phi(y_n) - \nabla_{\mathbf{w}} A(\eta_n) \quad (48)$$

In the case of Logistic Regression:

$$\nabla_{\mathbf{w}} \mathcal{L}(\mathbf{w}) = \frac{1}{N} \mathbf{X}^\top [\sigma(\mathbf{X}\mathbf{w}) - \mathbf{y}] \quad (49)$$

Some examples

- Gaussian distribution



Least Squares

Some examples

- Gaussian distribution



Least Squares

- Bernoulli distribution



Logistic Regression

Some examples

• Gaussian distribution	\Rightarrow	Least Squares
• Bernoulli distribution	\Rightarrow	Logistic Regression
• Multi-nomial distribution	\Rightarrow	Softmax Regression

Table of Contents

- 1 Review of Last Week
- 2 Exponential Families and Generalized Linear Models
- 3 Multi-Layer Perceptron (MLP) and Back-Propagation (BP)**
 - The Basic Structure of MLP
 - Training of NNs and BP

Table of Contents

- 1 Review of Last Week
 - Generalization Gap and Model Selection
 - Bias-Variance Decomposition
 - Logistic Regression
- 2 Exponential Families and Generalized Linear Models
 - Motivation
 - Exponential family
 - Application in ML (Generalized Linear Models)
- 3 Multi-Layer Perceptron (MLP) and Back-Propagation (BP)**
 - The Basic Structure of MLP**
 - Training of NNs and BP

MLP from a single neuron view




Figure: A simple MLP.

The output at the node j in layer l is denoted by $x_j^{(l)}$ and it is given by

$$x_j^{(l)} = \phi\left(\sum w_{i,j}^{(l)} x_i^{(l-1)} + b_j^{(l)}\right). \quad (50)$$

NNs extract suitable features from the input

A NN can be decomposed into a feature extractor and the output layer.

NNs extract suitable features from the input

A NN can be decomposed into a feature extractor and the output layer.

- Feature extractor $\mathbb{R}^d \rightarrow \mathbb{R}^K$: It transforms data into a suitable representation.

NNs extract suitable features from the input

A NN can be decomposed into a feature extractor and the output layer.

- Feature extractor $\mathbb{R}^d \rightarrow \mathbb{R}^K$: It transforms data into a suitable representation.

This function is defined by

NNs extract suitable features from the input

A NN can be decomposed into a feature extractor and the output layer.

- Feature extractor $\mathbb{R}^d \rightarrow \mathbb{R}^K$: It transforms data into a suitable representation.

This function is defined by

- The biases $\{\mathbf{b}^{(l)}\}_{l \in [L]}$ and weights $\{\mathbf{W}^{(l)}\}_{l \in [L]}$

NNs extract suitable features from the input

A NN can be decomposed into a feature extractor and the output layer.

- Feature extractor $\mathbb{R}^d \rightarrow \mathbb{R}^K$: It transforms data into a suitable representation.

This function is defined by

- The biases $\{\mathbf{b}^{(l)}\}_{l \in [L]}$ and weights $\{\mathbf{W}^{(l)}\}_{l \in [L]}$
- The activation function σ we pick

NNs extract suitable features from the input

A NN can be decomposed into a feature extractor and the output layer.

- Feature extractor $\mathbb{R}^d \rightarrow \mathbb{R}^K$: It transforms data into a suitable representation.

This function is defined by

- The biases $\{\mathbf{b}^{(l)}\}_{l \in [L]}$ and weights $\{\mathbf{W}^{(l)}\}_{l \in [L]}$
- The activation function σ we pick

In practice: both L and K are large — over-parameterized NNs.

NNs extract suitable features from the input

A NN can be decomposed into a feature extractor and the output layer.

- Feature extractor $\mathbb{R}^d \rightarrow \mathbb{R}^K$: It transforms data into a suitable representation.

This function is defined by

- The biases $\{\mathbf{b}^{(l)}\}_{l \in [L]}$ and weights $\{\mathbf{W}^{(l)}\}_{l \in [L]}$
- The activation function σ we pick

In practice: both L and K are large — over-parameterized NNs.

- The last layer $\mathbb{R}^K \rightarrow \mathbb{R}$: It performs the desired ML task, either linear regression or classification.

Table of Contents

- 1 Review of Last Week
 - Generalization Gap and Model Selection
 - Bias-Variance Decomposition
 - Logistic Regression
- 2 Exponential Families and Generalized Linear Models
 - Motivation
 - Exponential family
 - Application in ML (Generalized Linear Models)
- 3 Multi-Layer Perceptron (MLP) and Back-Propagation (BP)
 - The Basic Structure of MLP
 - Training of NNs and BP

Training loss for a regression problem with $S_{\text{train}} = \{(\mathbf{x}_n, y_n)\}_{n=1}^N$:

$$\mathcal{L}(f) = \frac{1}{2N} \sum_{n=1}^N (y_n - f(\mathbf{x}_n))^2, \quad (51)$$

where

Training loss for a regression problem with $S_{\text{train}} = \{(\mathbf{x}_n, y_n)\}_{n=1}^N$:

$$\mathcal{L}(f) = \frac{1}{2N} \sum_{n=1}^N (y_n - f(\mathbf{x}_n))^2, \quad (51)$$

where

- f is the function represented by a NN.

Training loss for a regression problem with $S_{\text{train}} = \{(\mathbf{x}_n, y_n)\}_{n=1}^N$:

$$\mathcal{L}(f) = \frac{1}{2N} \sum_{n=1}^N (y_n - f(\mathbf{x}_n))^2, \quad (51)$$

where

- f is the function represented by a NN.
- The overall function $y = f(\mathbf{x}^{(0)})$ can then be written as the composition:

$$f(\mathbf{x}^{(0)}) = f^{(L+1)} \circ \dots \circ f^{(2)} \circ f^{(1)}(\mathbf{x}^{(0)}).$$

Compact description of output

Compact description of output

- The function that is implemented by each layer in the form

$$\mathbf{x}^{(l)} = f^{(l)}(\mathbf{x}^{(l-1)}) = \phi((\mathbf{W}^{(l)})^\top \mathbf{x}^{(l-1)} + \mathbf{b}^{(l)}) . \quad (52)$$

Compact description of output

- The function that is implemented by each layer in the form

$$\mathbf{x}^{(l)} = f^{(l)}(\mathbf{x}^{(l-1)}) = \phi((\mathbf{W}^{(l)})^\top \mathbf{x}^{(l-1)} + \mathbf{b}^{(l)}) . \quad (52)$$

- Let $\mathbf{W}^{(l)}$ denote the *weight* matrix that connects layer $l - 1$ to layer l .

Compact description of output

- The function that is implemented by each layer in the form

$$\mathbf{x}^{(l)} = f^{(l)}(\mathbf{x}^{(l-1)}) = \phi((\mathbf{W}^{(l)})^\top \mathbf{x}^{(l-1)} + \mathbf{b}^{(l)}) . \quad (52)$$

- Let $\mathbf{W}^{(l)}$ denote the *weight* matrix that connects layer $l - 1$ to layer l .
- The matrix $\mathbf{W}^{(1)}$ is of dimension $D \times K$, the matrices $\mathbf{W}^{(l)}$, $2 \leq l \leq L$, are of dimension $K \times K$, and the matrix $\mathbf{W}^{(L+1)}$ is of dimension $K \times 1$.

Compact description of output

- The function that is implemented by each layer in the form

$$\mathbf{x}^{(l)} = f^{(l)}(\mathbf{x}^{(l-1)}) = \phi((\mathbf{W}^{(l)})^\top \mathbf{x}^{(l-1)} + \mathbf{b}^{(l)}). \quad (52)$$

- Let $\mathbf{W}^{(l)}$ denote the *weight* matrix that connects layer $l - 1$ to layer l .
- The matrix $\mathbf{W}^{(1)}$ is of dimension $D \times K$, the matrices $\mathbf{W}^{(l)}$, $2 \leq l \leq L$, are of dimension $K \times K$, and the matrix $\mathbf{W}^{(L+1)}$ is of dimension $K \times 1$.
- The entries of each matrix \mathbf{W} are given by

$$\mathbf{W}_{i,j}^{(l)} = w_{i,j}^{(l)}, \quad (53)$$

Compact description of output

- The function that is implemented by each layer in the form

$$\mathbf{x}^{(l)} = f^{(l)}(\mathbf{x}^{(l-1)}) = \phi((\mathbf{W}^{(l)})^\top \mathbf{x}^{(l-1)} + \mathbf{b}^{(l)}). \quad (52)$$

- Let $\mathbf{W}^{(l)}$ denote the *weight* matrix that connects layer $l - 1$ to layer l .
- The matrix $\mathbf{W}^{(1)}$ is of dimension $D \times K$, the matrices $\mathbf{W}^{(l)}$, $2 \leq l \leq L$, are of dimension $K \times K$, and the matrix $\mathbf{W}^{(L+1)}$ is of dimension $K \times 1$.
- The entries of each matrix \mathbf{W} are given by

$$\mathbf{W}_{i,j}^{(l)} = w_{i,j}^{(l)}, \quad (53)$$

where $w_{i,j}^{(l)}$ is the edge weight that connects node i on layer $l - 1$ to node j on layer l .

The back-propagation algorithm

Cost function:

$$\mathcal{L}_n = \left(y_n - f^{(L+1)} \circ \dots \circ f^{(2)} \circ f^{(1)}(\mathbf{x}_n^{(0)}) \right)^2,$$

where $\mathbf{x}_n^{(l)} = f^{(l)}(\mathbf{x}_n^{(l-1)}) = \phi((\mathbf{W}^{(l)})^\top \mathbf{x}_n^{(l-1)} + \mathbf{b}^{(l)})$.

The back-propagation algorithm

Cost function:

$$\mathcal{L}_n = \left(y_n - f^{(L+1)} \circ \dots \circ f^{(2)} \circ f^{(1)}(\mathbf{x}_n^{(0)}) \right)^2,$$

where $\mathbf{x}_n^{(l)} = f^{(l)}(\mathbf{x}_n^{(l-1)}) = \phi((\mathbf{W}^{(l)})^\top \mathbf{x}_n^{(l-1)} + \mathbf{b}^{(l)})$.

Recall that we aim to compute:

$$\frac{\partial \mathcal{L}_n}{\partial w_{i,j}^{(l)}}, \quad l = 1, \dots, L+1,$$
$$\frac{\partial \mathcal{L}_n}{\partial b_j^{(l)}}, \quad l = 1, \dots, L+1.$$

Let's use two quantities (i.e., $\mathbf{z}^{(l)}$ and $\delta^{(l)}$) to aid the computation:

Let's use two quantities (i.e., $\mathbf{z}^{(l)}$ and $\delta^{(l)}$) to aid the computation:

- Quantity computed in the **forward pass**:

$$\mathbf{z}^{(l)} = (\mathbf{W}^{(l)})^\top \mathbf{x}^{(l-1)} + \mathbf{b}^{(l)} \quad (54)$$

be the input at the l -th layer before applying the activation function, where $\mathbf{x}^{(l)} = \phi(\mathbf{z}^{(l)})$.

Let's use two quantities (i.e., $\mathbf{z}^{(l)}$ and $\delta^{(l)}$) to aid the computation:

- Quantity computed in the **forward pass**:

$$\mathbf{z}^{(l)} = (\mathbf{W}^{(l)})^\top \mathbf{x}^{(l-1)} + \mathbf{b}^{(l)} \quad (54)$$

be the input at the l -th layer before applying the activation function, where $\mathbf{x}^{(l)} = \phi(\mathbf{z}^{(l)})$.

- Quantity computed in the **backward pass**:

$$\delta_j^{(l)} = \frac{\partial \mathcal{L}_n}{\partial z_j^{(l)}} \quad (55)$$

$$= \sum_k \frac{\partial \mathcal{L}_n}{\partial z_k^{(l+1)}} \frac{\partial z_k^{(l+1)}}{\partial z_j^{(l)}} \quad (56)$$

$$= \sum_k \delta_k^{(l+1)} \mathbf{w}_{j,k}^{(l+1)} \phi'(z_j^{(l)}), \quad (57)$$

Let's use two quantities (i.e., $\mathbf{z}^{(l)}$ and $\delta^{(l)}$) to aid the computation:

- Quantity computed in the **forward pass**:

$$\mathbf{z}^{(l)} = (\mathbf{W}^{(l)})^\top \mathbf{x}^{(l-1)} + \mathbf{b}^{(l)} \quad (54)$$

be the input at the l -th layer before applying the activation function, where $\mathbf{x}^{(l)} = \phi(\mathbf{z}^{(l)})$.

- Quantity computed in the **backward pass**:

$$\delta_j^{(l)} = \frac{\partial \mathcal{L}_n}{\partial z_j^{(l)}} \quad (55)$$

$$= \sum_k \frac{\partial \mathcal{L}_n}{\partial z_k^{(l+1)}} \frac{\partial z_k^{(l+1)}}{\partial z_j^{(l)}} \quad (56)$$

$$= \sum_k \delta_k^{(l+1)} \mathbf{w}_{j,k}^{(l+1)} \phi'(z_j^{(l)}), \quad (57)$$

In vector form, we can write this as

$$\delta^{(l)} = (\mathbf{W}^{(l+1)} \delta^{(l+1)}) \odot \phi'(\mathbf{z}^{(l)}), \quad (58)$$

where \odot denotes the Hadamard product (the point-wise multiplication of vectors).

Now that we have both $\mathbf{z}^{(l)}$ and $\delta^{(l)}$ let us get back to our initial goal.

$$\frac{\partial \mathcal{L}_n}{\partial w_{i,j}^{(l)}} = \sum_k \frac{\partial \mathcal{L}_n}{\partial z_k^{(l)}} \frac{\partial z_k^{(l)}}{\partial w_{i,j}^{(l)}} = \underbrace{\frac{\partial \mathcal{L}_n}{\partial z_j^{(l)}}}_{\delta_j^{(l)}} \underbrace{\frac{\partial z_j^{(l)}}{\partial w_{i,j}^{(l)}}}_{\mathbf{x}_i^{(l-1)}} = \delta_j^{(l)} \mathbf{x}_i^{(l-1)}$$

$$\frac{\partial \mathcal{L}_n}{\partial b_j^{(l)}} = \sum_k \frac{\partial \mathcal{L}_n}{\partial z_k^{(l)}} \frac{\partial z_k^{(l)}}{\partial b_j^{(l)}} = \underbrace{\frac{\partial \mathcal{L}_n}{\partial z_j^{(l)}}}_{\delta_j^{(l)}} \underbrace{\frac{\partial z_j^{(l)}}{\partial b_j^{(l)}}}_1 = \delta_j^{(l)} \cdot 1 = \delta_j^{(l)}.$$

Summary: Backpropagation Algorithm for Computing the Derivatives

Settings: We are given a NN with L hidden layers

Summary: Backpropagation Algorithm for Computing the Derivatives

Settings: We are given a NN with L hidden layers

- All weight matrices $\mathbf{W}^{(l)}$ and bias vectors $\mathbf{b}^{(l)}$, $l = 1, \dots, L + 1$, are fixed.

Summary: Backpropagation Algorithm for Computing the Derivatives

Settings: We are given a NN with L hidden layers

- All weight matrices $\mathbf{W}^{(l)}$ and bias vectors $\mathbf{b}^{(l)}$, $l = 1, \dots, L + 1$, are fixed.
- We are given in addition a sample (\mathbf{x}_n, y_n) .

Summary: Backpropagation Algorithm for Computing the Derivatives

Settings: We are given a NN with L hidden layers

- All weight matrices $\mathbf{W}^{(l)}$ and bias vectors $\mathbf{b}^{(l)}$, $l = 1, \dots, L + 1$, are fixed.
- We are given in addition a sample (\mathbf{x}_n, y_n) .
- We want to compute the derivatives

$$\frac{\partial \mathcal{L}_n}{\partial w_{i,j}^{(l)}}, \quad \frac{\partial \mathcal{L}_n}{\partial b_j^{(l)}}, \quad l = 1, \dots, L + 1,$$

where

$$\mathcal{L}_n = (y_n - f^{(L+1)} \circ \dots \circ f^{(2)} \circ f^{(1)}(\mathbf{x}_n))^2.$$

Summary: Backpropagation Algorithm for Computing the Derivatives

Settings: We are given a NN with L hidden layers

- All weight matrices $\mathbf{W}^{(l)}$ and bias vectors $\mathbf{b}^{(l)}$, $l = 1, \dots, L + 1$, are fixed.
- We are given in addition a sample (\mathbf{x}_n, y_n) .
- We want to compute the derivatives

$$\frac{\partial \mathcal{L}_n}{\partial w_{i,j}^{(l)}}, \quad \frac{\partial \mathcal{L}_n}{\partial b_j^{(l)}}, \quad l = 1, \dots, L + 1,$$

where

$$\mathcal{L}_n = (y_n - f^{(L+1)} \circ \dots \circ f^{(2)} \circ f^{(1)}(\mathbf{x}_n))^2.$$

Forward pass: Set $\mathbf{x}^{(0)} = \mathbf{x}_n$. Compute for $l = 1, \dots, L + 1$,

$$\mathbf{z}^{(l)} = (\mathbf{W}^{(l)})^\top \mathbf{x}^{(l-1)} + \mathbf{b}^{(l)}, \quad \mathbf{x}^{(l)} = \phi(\mathbf{z}^{(l)}).$$

Summary: Backpropagation Algorithm for Computing the Derivatives

Settings: We are given a NN with L hidden layers

- All weight matrices $\mathbf{W}^{(l)}$ and bias vectors $\mathbf{b}^{(l)}$, $l = 1, \dots, L + 1$, are fixed.
- We are given in addition a sample (\mathbf{x}_n, y_n) .
- We want to compute the derivatives

$$\frac{\partial \mathcal{L}_n}{\partial w_{i,j}^{(l)}}, \quad \frac{\partial \mathcal{L}_n}{\partial b_j^{(l)}}, \quad l = 1, \dots, L + 1,$$

where

$$\mathcal{L}_n = (y_n - f^{(L+1)} \circ \dots \circ f^{(2)} \circ f^{(1)}(\mathbf{x}_n))^2.$$

Forward pass: Set $\mathbf{x}^{(0)} = \mathbf{x}_n$. Compute for $l = 1, \dots, L + 1$,

$$\mathbf{z}^{(l)} = (\mathbf{W}^{(l)})^\top \mathbf{x}^{(l-1)} + \mathbf{b}^{(l)}, \quad \mathbf{x}^{(l)} = \phi(\mathbf{z}^{(l)}).$$

Backward pass: Set $\delta^{(L+1)} = -2(y_n - \mathbf{x}^{(L+1)})\phi'(\mathbf{z}^{(L+1)})$. Compute for $l = L, \dots, 1$,

$$\delta^{(l)} = (\mathbf{W}^{(l+1)} \delta^{(l+1)}) \odot \phi'(\mathbf{z}^{(l)}).$$

Summary: Backpropagation Algorithm for Computing the Derivatives

Settings: We are given a NN with L hidden layers

- All weight matrices $\mathbf{W}^{(l)}$ and bias vectors $\mathbf{b}^{(l)}$, $l = 1, \dots, L + 1$, are fixed.
- We are given in addition a sample (\mathbf{x}_n, y_n) .
- We want to compute the derivatives

$$\frac{\partial \mathcal{L}_n}{\partial w_{i,j}^{(l)}}, \quad \frac{\partial \mathcal{L}_n}{\partial b_j^{(l)}}, \quad l = 1, \dots, L + 1,$$

where

$$\mathcal{L}_n = (y_n - f^{(L+1)} \circ \dots \circ f^{(2)} \circ f^{(1)}(\mathbf{x}_n))^2.$$

Forward pass: Set $\mathbf{x}^{(0)} = \mathbf{x}_n$. Compute for $l = 1, \dots, L + 1$,

$$\mathbf{z}^{(l)} = (\mathbf{W}^{(l)})^\top \mathbf{x}^{(l-1)} + \mathbf{b}^{(l)}, \quad \mathbf{x}^{(l)} = \phi(\mathbf{z}^{(l)}).$$

Backward pass: Set $\delta^{(L+1)} = -2(y_n - \mathbf{x}^{(L+1)})\phi'(\mathbf{z}^{(L+1)})$. Compute for $l = L, \dots, 1$,

$$\delta^{(l)} = (\mathbf{W}^{(l+1)}\delta^{(l+1)}) \odot \phi'(\mathbf{z}^{(l)}).$$

Final computation: For all parameters compute

This lecture:

- Exponential Families and Generalized Linear Models
- Multi-Layer Perceptron
- Back-Propagation

This lecture:

- Exponential Families and Generalized Linear Models
- Multi-Layer Perceptron
- Back-Propagation

Next lecture ?