

# Unit 2: Polynomials and Rational Functions

## Honors Precalculus

Student

### Contents of this unit:

- 2.1 Quadratic Functions
- 2.2 Graphs of Polynomial Functions
- 2.3 Zeros of Polynomial Functions
- 2.4 Dividing Polynomials
- 2.5 Complex Numbers
- 2.6 Fundamental Theorem of Algebra
- 2.7 Rational Functions and Asymptotes
- 2.8 Graphs of Rational Functions

## Contents

|     |                                   |    |
|-----|-----------------------------------|----|
| 2.1 | Quadratic Functions               | 2  |
| 2.2 | Graphs of Polynomial Functions    | 4  |
| 2.3 | Zeros of Polynomial Functions     | 6  |
| 2.4 | Dividing Polynomials              | 8  |
| 2.5 | Complex Numbers                   | 11 |
| 2.6 | Fundamental Theorem of Algebra    | 13 |
| 2.7 | Rational Functions and Asymptotes | 15 |
| 2.8 | Graphs of Rational Functions      | 17 |

## 2.1 Quadratic Functions

### Objectives

- Classify polynomial functions by degree.
- Identify the key features of a parabola (vertex, axis of symmetry, intercepts).
- Convert between vertex form and standard form by completing the square.
- Solve quadratic-type equations using substitution.

### Polynomial Functions

A polynomial function is classified by its **degree** — the largest exponent on the variable.

| Equation                             | Degree | Type  |
|--------------------------------------|--------|-------|
| $f(x) = c$                           | _____  | _____ |
| $f(x) = mx + b$                      | _____  | _____ |
| $f(x) = ax^2 + bx + c$               | _____  | _____ |
| $f(x) = ax^3 + bx^2 + cx + d$        | _____  | _____ |
| $f(x) = ax^4 + bx^3 + cx^2 + dx + e$ | _____  | _____ |

- The **leading coefficient** is the coefficient on the highest-degree term. It controls end behavior and (for quadratics) whether the parabola opens up or down.
- The **constant term** is the term with no variable. It is always the  $y$ -intercept:  $f(0)$ .

The graph of every quadratic function is a **parabola**. It has five features worth naming.

### Anatomy of a Parabola

- **Vertex** — the highest or lowest point on the parabola.
- **Axis of symmetry** — the vertical line through the vertex; the parabola is a mirror image on either side.
- **$x$ -intercepts** — the points where the parabola crosses the  $x$ -axis (may have 0, 1, or 2).
- **$y$ -intercept** — the point where the parabola crosses the  $y$ -axis.
- If  $a > 0$ , the parabola opens \_\_\_\_\_ and the vertex is a \_\_\_\_\_.
- If  $a < 0$ , the parabola opens \_\_\_\_\_ and the vertex is a \_\_\_\_\_.

### Forms of a Quadratic Function

| Feature        | Vertex Form $a(x - h)^2 + k$ | Standard Form $ax^2 + bx + c$ |
|----------------|------------------------------|-------------------------------|
| Vertex         | _____                        | _____                         |
| Axis of sym.   | _____                        | _____                         |
| $y$ -intercept | _____                        | _____                         |

To convert from standard form to vertex form, \_\_\_\_\_.

### Note

#### Standard Form → Vertex Form (Completing the Square):

1. Factor out  $a$  from the  $x$ -terms:  $a\left(x^2 + \frac{b}{a}x\right) + c$ .

2. Take half the inner coefficient, square it:  $\left(\frac{b}{2a}\right)^2$ .
3. Add and subtract that value inside the parentheses.
4. Factor the perfect-square trinomial and simplify.

**Example 1**

Given  $f(x) = 2x^2 + 8x + 7$ , write the function in vertex form, identify the vertex, and sketch the graph.

**Example 2**

Write the standard form of the equation of the parabola whose vertex is  $(1, 2)$  and that passes through the point  $(3, -6)$ .

**Example 3**

Sophia has a food truck and sells tacos. Her daily costs are approximated by  $C(x) = x^2 - 40x + 610$ , where  $C(x)$  is the cost in dollars to sell  $x$  tacos.

- (a) Find the number of tacos she must sell to minimize her cost.
- (b) What is the minimum cost?

**Quadratic-Type Equations**

A polynomial that *isn't* degree 2 can still factor like a quadratic. Look for three clues:

1. Exactly **three terms** (a trinomial).
2. The exponent on the first term is \_\_\_\_\_ the exponent on the second.
3. The third term is a \_\_\_\_\_.

**Strategy:** Substitute  $u$  for the middle-term variable expression (e.g.,  $u = x^5$ ). The equation becomes

$au^2 + bu + c = 0$  — a familiar quadratic. Solve for  $u$ , then back-substitute to solve for  $x$ .

#### Example 4

Solve  $x^6 + 7x^3 = 8$ .

*Homework:* Quadratic Functions Worksheet

## 2.2 Graphs of Polynomial Functions

### Objectives

- Identify the features that make a graph polynomial (continuous, smooth).
- Predict end behavior from the leading term.
- Use multiplicity to determine whether the graph crosses or is tangent to the  $x$ -axis.
- Sketch polynomials from factored form.

### What Makes a Graph Polynomial

- The graph of every polynomial function is \_\_\_\_\_ — no breaks, no holes, no jumps.
- The graph has only \_\_\_\_\_ — no sharp corners or cusps.
- \_\_\_\_\_ properties must hold for the graph to be polynomial. Either one alone is \_\_\_\_\_.

### Zeros and Turning Points

For a polynomial function  $f$  of degree  $n$ :

- $f$  has at most \_\_\_\_\_ real zeros (i.e.,  $x$ -intercepts).
- The graph of  $f$  has at most \_\_\_\_\_ turning points.

A **zero** of  $f$  is a number  $x$  for which \_\_\_\_\_. Equivalently, it is an  $x$ -intercept of the graph.

### End Behavior and the Leading Coefficient Test

**End behavior** describes what happens as  $x$  becomes large in the \_\_\_\_\_ direction. It is determined by the function's \_\_\_\_\_ and the \_\_\_\_\_ of its leading coefficient.

| Case | Degree | Leading Coef. | End Behavior                         |
|------|--------|---------------|--------------------------------------|
| 1    | even   | positive      | both ends $\uparrow$                 |
| 2    | even   | negative      | both ends $\downarrow$               |
| 3    | odd    | positive      | left $\downarrow$ , right $\uparrow$ |
| 4    | odd    | negative      | left $\uparrow$ , right $\downarrow$ |

**Example 1**

Which of the following could be the graph of a polynomial whose leading term is  $-3x^4$ ?

**Example 2**

Describe the end behavior of each polynomial.

a)  $f(x) = -x^3 + 4x$

b)  $f(x) = x^4 - 5x^2 + 4$

**Repeated Zeros and Multiplicity**

A factor of  $(x - a)^k$  with  $k > 1$  means  $x = a$  is a **zero of multiplicity  $k$** . The multiplicity controls how the graph meets the  $x$ -axis at that zero:

- If  $k$  is \_\_\_\_\_, the graph \_\_\_\_\_ the  $x$ -axis at  $x = a$ .
- If  $k$  is \_\_\_\_\_, the graph is \_\_\_\_\_ the  $x$ -axis at  $x = a$  (touches and bounces back).
- The higher the multiplicity, the \_\_\_\_\_ the graph passes through (or past) the  $x$ -axis.

**Example 3**

For  $f(x) = -(x - 1)(x + 2)(x - 3)$ :

(a) Identify the degree.    (b) Identify the  $x$ -intercepts.    (c) Cross or tangent at each?    (d) Sketch.

**Example 4**

For  $f(x) = (x + 2)^2(x - 3)^3$ :

- (a) Identify the degree.    (b) Identify the  $x$ -intercepts.    (c) Cross or tangent at each?    (d) Sketch.

**Example 5**

How many turning points can each polynomial have at most?

a)  $f(x) = -(x + 1)^2(x - 3)$

b)  $f(x) = (x + 1)^2(x - 3)^3$

**Example 6**

Write an equation for a polynomial graph with  $x$ -intercepts at  $x = -3, -1, 1, 2$  whose left and right arms both point **down**. Assume the leading coefficient is 1 or  $-1$ .

*Homework:* Graphs of Polynomial Functions Worksheet

## 2.3 Zeros of Polynomial Functions

### Objectives

- Translate fluently among zero, root, solution, factor, and  $x$ -intercept.
- Sketch polynomial graphs by factoring and using multiplicities.
- Build a polynomial from a given set of zeros.
- Apply the Intermediate Value Theorem to guarantee zeros.

### Five Names, One Idea

If  $f(x)$  is a polynomial and  $a$  is a real number, the following five statements are equivalent — they all mean the same thing:

- $x = a$  is a \_\_\_\_\_ of the **function**  $f(x)$ .
- $x = a$  is a \_\_\_\_\_ of the polynomial **equation**  $f(x) = 0$ .
- $(x - a)$  is a \_\_\_\_\_ of the **polynomial**  $f(x)$ .
- $(a, 0)$  is an \_\_\_\_\_ of the **graph** of  $f$ .

- Sometimes zeros are called \_\_\_\_\_.

**Example 1**

For  $f(x) = x^3 - x^2 - 2x$ , use the zeros,  $y$ -intercept, and end behavior to sketch the graph.

**Example 2**

For  $g(x) = x^3 - 2x^2 - 4x + 8$ , factor and sketch.

**Example 3**

For  $f(x) = -2x^4 - x^3 + 3x^2$ , factor and sketch.

**Example 4**

For  $f(x) = x^4 - 12x^2 + 27$ , find all real zeros and sketch.

**Building a Polynomial from Its Zeros**

If a polynomial has zeros  $r_1, r_2, \dots, r_n$ , then:  $f(x) = a(x - r_1)(x - r_2) \cdots (x - r_n)$

- A zero listed twice gives a factor with exponent 2 (its multiplicity).



- For a fractional zero like  $-\frac{1}{2}$ , use the factor  $(2x + 1)$  to keep coefficients as integers.
- If  $p + \sqrt{q}$  is a zero, then  $p - \sqrt{q}$  is also a zero (conjugate radicals).

**Example 5**

Write an equation of a polynomial with zeros  $-\frac{1}{2}$ , 3, 3.

**Example 6**

Write an equation of a polynomial with zeros 3,  $2 + \sqrt{3}$ ,  $2 - \sqrt{3}$ .

**Intermediate Value Theorem (IVT)**

If  $f$  is a polynomial (continuous) and \_\_\_\_\_ while \_\_\_\_\_ (or vice versa), then  $f$  has **at least one zero** between  $a$  and  $b$ .

- A continuous function cannot jump over the  $x$ -axis.
- IVT proves a zero *exists* but does not tell you *where*.
- If  $f(a)$  and  $f(b)$  have the same sign, the test is inconclusive.

**Example 7**

Show that  $f(x) = x^4 - 4x + 1$  has at least one zero between  $x = 0$  and  $x = 1$ .

*Homework:* Zeros of Polynomial Functions Worksheet

## 2.4 Dividing Polynomials

**Objectives**

- Divide polynomials using long division.
- Divide polynomials using synthetic division (linear divisors only).
- Apply the Remainder Theorem and the Factor Theorem.
- Use the Rational Zero Test to list possible rational zeros.

### Polynomial Long Division

Long division of polynomials works exactly like long division of whole numbers: **divide, multiply, subtract, bring down, repeat**.

The result always has the form:

$$\frac{f(x)}{d(x)} = q(x) + \frac{r(x)}{d(x)}$$

where  $q(x)$  is the \_\_\_\_\_ and  $r(x)$  is the \_\_\_\_\_.

- The degree of the remainder is always \_\_\_\_\_ than the degree of the divisor.
- Always write polynomials in \_\_\_\_\_ of degree before dividing.
- Use 0 as a placeholder for any missing powers.

#### Example 1

Divide  $2x^3 + 3x^2 - 5x + 1$  by  $x + 2$  using long division.

#### Example 2

Divide  $6x^3 + 10x^2 + x + 8$  by  $2x^2 + 1$ .

### Synthetic Division

Synthetic division is a shortcut for dividing by a **linear** divisor of the form \_\_\_\_\_.

#### Steps:

1. Write  $c$  on the left and the coefficients of the dividend across the top.
2. Bring down the first coefficient.
3. Multiply by  $c$ , write the product under the next coefficient, and add.
4. Repeat until you reach the last column — that entry is the \_\_\_\_\_.

The bottom row (excluding the remainder) gives the coefficients of the quotient, whose degree is **one less** than the dividend's.

**Example 3**

Use synthetic division to divide  $x^3 - 7x + 6$  by  $x - 2$ .

**Example 4**

Use synthetic division to divide  $3x^4 - 5x^3 + 4x - 6$  by  $x - 1$ .

**The Remainder Theorem**

If a polynomial  $f(x)$  is divided by  $(x - c)$ , then the remainder equals \_\_\_\_\_.

In other words: evaluating  $f(c)$  and dividing by  $(x - c)$  give the same number — the remainder.

**The Factor Theorem**

$(x - c)$  is a factor of  $f(x)$  if and only if \_\_\_\_\_.

This is just the Remainder Theorem with the remainder equal to zero. It connects factoring and root-finding.

**Example 5**

Use the Remainder Theorem to evaluate  $f(3)$  for  $f(x) = 2x^3 - 5x^2 + x - 3$ .

**The Rational Zero Test**

If a polynomial  $f(x) = a_nx^n + \cdots + a_1x + a_0$  has integer coefficients, then every rational zero has the form:

$$\frac{p}{q}$$

where  $p$  is a factor of the \_\_\_\_\_  $a_0$  and  $q$  is a factor of the \_\_\_\_\_  $a_n$ .

**Strategy:** List all  $\frac{p}{q}$  combinations, then test each candidate using synthetic division or direct evaluation.

**Example 6**

List all possible rational zeros of  $f(x) = 2x^3 + 3x^2 - 8x + 3$ , then find all actual zeros.

*Homework:* Dividing Polynomials Worksheet

## 2.5 Complex Numbers

### Objectives

- Define the imaginary unit  $i$  and simplify powers of  $i$ .
- Write complex numbers in standard form  $a + bi$ .
- Add, subtract, multiply, and divide complex numbers.
- Solve quadratic equations with complex solutions.

### The Imaginary Unit

The imaginary unit  $i$  is defined by:

$$i = \text{_____}, \quad i^2 = \text{_____}$$

For any positive real number  $a$ :

$$\sqrt{-a} = \text{_____}$$

**Standard form** of a complex number: \_\_\_\_\_, where  $a$  is the \_\_\_\_\_ part and  $b$  is the \_\_\_\_\_ part.

**Example 1**

Write each number in standard form.

a)  $\sqrt{-9}$

b)  $\sqrt{-48}$

c)  $3 + \sqrt{-25}$

### Operations on Complex Numbers

Let  $a + bi$  and  $c + di$  be complex numbers.

- **Addition:**  $(a + bi) + (c + di) = \text{_____}$
- **Subtraction:**  $(a + bi) - (c + di) = \text{_____}$
- **Multiplication:** Use FOIL, then replace  $i^2$  with  $-1$ .

**Example 2**

Simplify. Write answers in standard form.

a)  $(3 + 2i) + (5 - 4i)$

b)  $(7 - i) - (3 + 2i)$

**Example 3**

Multiply. Write answers in standard form.

a)  $(2 + 3i)(4 - i)$

b)  $(3 + 2i)^2$

**Complex Conjugates**

The **complex conjugate** of  $a + bi$  is \_\_\_\_\_.

Key property:  $(a + bi)(a - bi) = a^2 + b^2$  — always a \_\_\_\_\_.

To **divide** complex numbers, multiply the numerator and denominator by the \_\_\_\_\_ of the denominator.

**Example 4**

Divide. Write in standard form:  $\frac{2 + 3i}{4 - i}$

**Powers of  $i$** 

The powers of  $i$  cycle with period \_\_\_\_\_:

$$i^1 = i, \quad i^2 = -1, \quad i^3 = -i, \quad i^4 = 1, \quad i^5 = i, \dots$$

To simplify  $i^n$ : divide  $n$  by 4 and use the \_\_\_\_\_.

**Example 5**Simplify: a)  $i^{23}$ b)  $i^{42}$ c)  $i^{100}$ **Example 6**Solve  $x^2 + 4x + 7 = 0$ .*Homework:* Complex Numbers Circuit Worksheet**2.6 Fundamental Theorem of Algebra****Objectives**

- Use the Fundamental Theorem of Algebra to count zeros.
- Factor any polynomial completely over the complex numbers.
- Apply the Conjugate-Zero Theorems to recover missing zeros.
- Build polynomials with given zeros and an additional condition.

**The Fundamental Theorem of Algebra (FTA)**

Every polynomial equation with degree greater than zero has \_\_\_\_\_ in the set of complex numbers (counting multiplicity).

The polynomial therefore factors completely as a product of \_\_\_\_\_ over  $\mathbb{C}$ :

$$f(x) = a(x - r_1)(x - r_2) \cdots (x - r_n)$$

- A degree- $n$  polynomial has  $n$  zeros in  $\mathbb{C}$ .
- A repeated zero of multiplicity  $k$  counts as  $k$  of those zeros.
- Over the *real* numbers, a polynomial can have fewer than  $n$  zeros. Over  $\mathbb{C}$ , the count is exact.

**Example 1**Factor  $P(x) = x^3 + x^2 - 4x + 6$  completely over  $\mathbb{C}$ .

**Example 2**

Find all zeros of  $f(x) = x^5 + 6x^3 + 2x^2 - 27x + 18$ .

**Conjugate-Zero Theorems**

When a polynomial has **real coefficients**, non-real and irrational zeros come paired with their conjugates:

- **Complex Conjugate Zeros:** If  $a + bi$  is a zero, then \_\_\_\_\_ is also a zero.
- **Irrational Conjugates:** If  $a + \sqrt{b}$  is a zero, then \_\_\_\_\_ is also a zero.

Non-real zeros always pair up — a degree- $n$  polynomial with real coefficients has an **even** number of non-real zeros.

Multiplying a conjugate pair gives a quadratic with real coefficients:

$$(x - (a + bi))(x - (a - bi)) = (x - a)^2 + b^2$$

**Example 3**

Write a cubic polynomial  $g$  with real coefficients that has zeros 2 and  $\sqrt{3}$ , and where  $g(4) = 13$ .

**Example 4**

Find a cubic polynomial  $f$  with real coefficients that has 2 and  $1 - i$  as zeros, and  $f(1) = 3$ .

**Example 5**

Given that  $1 + 3i$  is a zero of  $f(x) = x^4 - 6x^3 + 47x^2 - 98x + 290$ , find all zeros.

*Homework:* Fundamental Theorem of Algebra Worksheet

**2.7 Rational Functions and Asymptotes****Objectives**

- Define rational functions and their domains.
- Identify vertical and horizontal asymptotes.
- Distinguish between holes and vertical asymptotes.
- Compare degrees to determine horizontal asymptotes.

**Rational Functions**

A **rational function** is a quotient of two polynomials:

$$f(x) = \frac{g(x)}{h(x)}, \quad h(x) \neq 0$$

The **domain** excludes every  $x$ -value that makes the denominator equal to \_\_\_\_\_.

The parent rational function is  $f(x) = \frac{1}{x}$ , a \_\_\_\_\_ with domain  $x \neq 0$  and range  $y \neq 0$ .

**Vertical and Horizontal Asymptotes**

- If  $f(x) \rightarrow \pm\infty$  as  $x \rightarrow a$ , then  $x = a$  is a \_\_\_\_\_.
- If  $f(x) \rightarrow a$  as  $x \rightarrow \pm\infty$ , then  $y = a$  is a \_\_\_\_\_.

**Asymptotes and Holes — The Rules**

**Factor first**, cancel common factors to expose holes, then read off the features.



| Feature            | Condition                                   | How to Find                                                             |
|--------------------|---------------------------------------------|-------------------------------------------------------------------------|
| Vertical Asymptote | Values that make _____ the denominator zero | Set remaining denom. = 0.                                               |
| Horiz. Asymptote   | deg(num) _____ deg(denom)                   | $y = 0$                                                                 |
|                    | deg(num) _____ deg(denom)                   | $y = \frac{\text{leading coef. of num}}{\text{leading coef. of denom}}$ |
| Hole               | Common factors in _____ num. and denom.     | Set the common factor = 0.                                              |
| Slant Asymptote    | deg(num) _____ deg(denom) (by exactly 1)    | Long/synthetic division; quotient is the slant line.                    |

**Example 1**

Find the domain of  $f(x) = \frac{1}{x-1}$  and describe the behavior near any excluded  $x$ -values.

**Example 2**

Find the asymptotes and/or holes.

a)  $f(x) = \frac{1}{2x}$

b)  $f(x) = \frac{2x^2}{x^2 - 1}$

**Example 3**

Find the asymptotes and/or holes.

a)  $f(x) = \frac{3x}{2x^2 + 1}$

b)  $y = \frac{(x+4)(x-2)}{(x-3)(x+4)}$

**Example 4**

For  $y = \frac{x^2 + x - 6}{x^2 - x - 2}$ : (a) Identify asymptotes and/or holes. (b) Write the domain. (c) Find the intercepts.

**Example 5**

For  $f(x) = \frac{3x^3 + 7x^2 + 2}{-2x^3 + 16}$ : (a) Identify asymptotes and/or holes. (b) Write the domain.

**Example 6**

For  $f(x) = \frac{x^2 - 16}{x^2 + 8x}$ : (a) Asymptotes/holes. (b) Domain. (c) Intercepts.

*Homework:* Rational Functions Worksheet

## 2.8 Graphs of Rational Functions

### Objectives

- Follow a four-step procedure to sketch any rational function by hand.
- Identify and draw slant (oblique) asymptotes.
- Use test points to determine the shape of each branch.

### Guidelines for Graphing a Rational Function

Every rational graph follows the same four-step recipe:

1. **Find & sketch asymptotes.** Factor num. and denom. Cancel common factors (holes). Read VA, HA, or slant asymptote. Draw each as a dashed line.
2. **Find & plot intercepts.**  $x$ -int from zeros of the (simplified) numerator.  $y$ -int is  $f(0)$  (if in the domain).
3. **Plot test points.** At least one point between and one beyond each  $x$ -intercept and VA.
4. **Connect smoothly.** Approach but never cross VAs. May cross HAs in the middle.

**Note**

Don't connect across a vertical asymptote. The graph splits into separate branches — sketch each branch independently.

**Example 1**

Sketch  $g(x) = \frac{3}{x-2}$ .

**Example 2**

Sketch  $f(x) = \frac{x^2}{x^2 - x - 2}$ .

**Example 3**

Sketch  $f(x) = \frac{x^2 - 9}{x^2 - 2x - 3}$ .

**Slant (Oblique) Asymptotes**

When the degree of the numerator is *exactly one more* than the denominator's, the long-run behavior is a tilted line.

**Condition:**  $\deg(\text{num}) = \deg(\text{denom}) + 1$

**How to find:** Long-divide the numerator by the denominator. The \_\_\_\_\_ (ignore the remainder) is the slant asymptote.

- A slant asymptote replaces — does **not coexist with** — a \_\_\_\_\_ asymptote.
- If the numerator degree exceeds the denominator's by more than 1, there is \_\_\_\_\_.

**Example 4**

Sketch  $f(x) = \frac{x^2 + x}{x - 1}$ .

**Example 5**

Sketch  $y = \frac{x^3}{x^2 - 4}$ .

*Homework:* Graphs of Rational Functions Worksheet