

## Problem Set 2

Byron Ralph Boter

Nhemuel Robles


Giovanni Ross Sucalit

2024-02-25

### Problem Set 2

1. Use the bisection method with a hand calculator or computer to find the real root of  $x^3 - x - 3 = 0$ . Use an error tolerance of  $\epsilon = 0.0001$ . Graph the function  $f(x) = x^3 - x - 3$  and label the root.

- **Step 1:** The graph below is generated using R, and illustrates the function  $f(x) = x^3 - x - 3$



- **Step 2:** Based on the graph, we start with  $[a, b] = [1.5, 2.0]$
- **Step 3:** To determine the no. of iterations, let  $|a - c_n| = \epsilon = 0.0001$ , then solve for  $n$ .

$$|a - c_n| \leq \left[\frac{1}{2}\right]^n (b - a)$$

$$0.0001 \leq \left[\frac{1}{2}\right]^n (2.0 - 1.5)$$

$$0.0001 \leq \left[\frac{1}{2}\right]^n (0.5)$$

$$\frac{0.0001}{0.5} \leq \left[\frac{1}{2}\right]^n$$

$$0.0002 \leq \left[\frac{1}{2}\right]^n$$

$$\ln 0.0002 \leq n \ln \left[\frac{1}{2}\right]$$

$$\frac{\ln 0.0002}{\ln \frac{1}{2}} \leq n$$

$$\frac{-8.5171931914162374266547336972793}{-0.69314718055994530941723212145818} \leq n$$

$$12.28771238 \leq n$$

$$n \geq 13$$

- **Step 4:** The following table is generated starting with  $Bisect(f(x) = x^3 - x - 3, 1.5, 2.0, c_1, \epsilon = 0.0001)$ .

n	a	b	c	f(b)	f(c)	f(b)f(c)	b - c  ≤ 0.0001?
1	1.5	2.0	1.75	3.0	0.609375	1.828125	No
2	1.5	1.75	1.625	0.609375	-0.333984375	-0.203521728515625	No
3	1.625	1.75	1.6875	0.609375	0.117919921875	0.0718574523925781	No
4	1.625	1.6875	1.65625	0.117919921875	-0.112884521484375	-0.0133113339543343	No
5	1.65625	1.6875	1.671875	0.117919921875	0.00129318237304687	0.000152491964399815	No
6	1.65625	1.671875	1.6640625	0.00129318237304687	-0.0561003684997559	-0.0000725480076653184	No
7	1.6640625	1.671875	1.66796875	0.00129318237304687	-0.0274799466133118	-0.0000355365825726039	No
8	1.66796875	1.671875	1.669921875	0.00129318237304687	-0.0131124928593636	-0.000016956844632432	No
9	1.669921875	1.671875	1.6708984375	0.00129318237304687	-0.00591443572193384	-0.00000764844402212361	No
10	1.6708984375	1.671875	1.67138671875	0.00129318237304687	-0.00231182214338332	-0.00000298960764544276	No
11	1.67138671875	1.671875	1.671630859375	0.00129318237304687	-0.00050961879605893	-0.000000659030044036779	No
12	1.671630859375	1.671875	1.6717529296875	0.00129318237304687	0.000391707055314328	0.000000506548659330586	No
13	1.671630859375	1.6717529296875	1.67169189453125	0.000391707055314328	-0.0000589745529850916	-0.000000023100748488269	Yes
14	1.67169189453125	1.6717529296875	<b>1.67172241210937</b>	0.000391707055314328	0.000166361580426155	0.000000065165004786167	Yes

- **To conclude**, the real root of the function  $f(x) = x^3 - x - 3$  is **1.67172241210937**.

2. The function  $f(x) = -3x^3 + 2e^{\frac{x^2}{2}} - 1$  has values of zero near  $x = -0.5$  and  $x = 0.5$ .

a. What is the derivative of  $f$ ?

b. If you begin Newton's method at  $x = 0$ , which root is reached? How many iterations to achieve an error less than  $10^{-5}$ ?

c. Begin Newton's method at another starting point to get the other zero.

3. Use the function from no.2 and find the root using the secant method where  $x_0 = 0$  and  $x_1 = 1$ . Use an error tolerance of  $\epsilon = 0.001$ .

4. Consider the system

$$10.2x + 2.4y - 4.5z = 14.067,$$

$$-2.3x - 7.7y + 11.1z = -0.996,$$

$$-5.5x - 3.2x + 0.9z = -12.645.$$

a. Present the augmented matrix of the system.

$$\left( \begin{array}{ccc|c} 10.2 & 2.4 & -4.5 & 14.067 \\ -2.3 & -7.7 & 11.1 & -0.996 \\ -5.5 & -3.2 & 0.9 & -12.645 \end{array} \right)$$

b. Solve the system using  $Ax = LUx = Ly = b$  and round the final answer to 4 decimal digits.

c. Find the residual vector if the correct solution is  $x = 1.4531001$ ,  $y = -1.5891949$ ,  $z = -0.2748947$ .

- **Step 1:** The following are given values of the variables  $b$ ,  $A$ , and  $\bar{x}$ :

$$A = \begin{pmatrix} 10.2 & 2.4 & -4.5 \\ -2.3 & -7.7 & 11.1 \\ -5.5 & -3.2 & 0.9 \end{pmatrix}$$

$$b = \begin{pmatrix} 14.067 \\ -0.996 \\ -12.645 \end{pmatrix}$$

$$\bar{x} = \begin{pmatrix} 1.4531001 \\ -1.5891949 \\ -0.2748947 \end{pmatrix}$$

- **Step 2:** Use the equation  $r = b - A\bar{x}$  to calculate the residual error:

$$r = b - A\bar{x}$$

$$r = \begin{pmatrix} 14.067 \\ -0.996 \\ -12.645 \end{pmatrix} - \begin{pmatrix} 10.2 & 2.4 & -4.5 \\ -2.3 & -7.7 & 11.1 \\ -5.5 & -3.2 & 0.9 \end{pmatrix} \begin{pmatrix} 1.4531001 \\ -1.5891949 \\ -0.2748947 \end{pmatrix}$$

$$r = \begin{pmatrix} 14.067 \\ -0.996 \\ -12.645 \end{pmatrix} - \begin{pmatrix} 10.2(1.4531001) + & 2.4(-1.5891949) - & 4.5(-0.2748947) \\ -2.3(1.4531001) + & -7.7(-1.5891949) + & 11.1(-0.2748947) \\ -5.5(1.4531001) + & -3.2(-1.5891949) + & 0.9(-0.2748947) \end{pmatrix}$$

$$r = \begin{pmatrix} 14.067 \\ -0.996 \\ -12.645 \end{pmatrix} - \begin{pmatrix} 12.24457941 \\ 5.84333933 \\ -3.1540321 \end{pmatrix}$$

$$r = \begin{pmatrix} 14.067 - 12.24457941 \\ -0.996 - 5.84333933 \\ -12.645 + 3.1540321 \end{pmatrix}$$

$$r = \begin{pmatrix} 1.82242059 \\ -6.83933933 \\ -9.4909679 \end{pmatrix}$$

- **To conclude**, the residual vector of the solution  $x = 1.4531001, y = -1.5891949, z = -0.2748947$  is  $r = (1.82242059, -6.83933933, -9.4909679)^T$ .

5. Compute the Frobenius norm, maximum column sum, and maximum row sum of the matrix:

$$\begin{pmatrix} 10.2 & 2.4 & 4.5 \\ -2.3 & 7.7 & 11.1 \\ -5.5 & -3.2 & 0.9 \end{pmatrix}$$

6. Solve the system of equations given in no. 4, starting with the initial vector of  $[0, 0, 0]$ :

- Solve using the Jacobi method with 2-digit precision.
- Solve using Gauss-Seidel method with 2-digit precision.
- Solve for  $e$  if the true solution is  $x = (1.5, 0.33, 0.45)^T$ .