Observation, Memory, Control, and Time in Finite Models

Kirtus Justus

Theoretical Physicist

Port Coquitlam, British Columbia, Canada

10 November 2026 · baseline edition 12 November 2026

Abstract

For finite conventional models — probability kernels, hidden-state models, quantum states and continuous-time Markov generators — the paper determines which features are fixed by a stated menu of observations and interventions. Results include an exact criterion for controlled reduction and a complete-history bound under uniform row error; a parity-shift process with exact innovation coordinates; classical realization rank with finite-sample certificates; anytime-valid likelihood-mixture tests for complete record laws; nonidentification of fragment factorization from a final channel; sharp control criteria for positive hidden realizations; calibration-induced identification intervals; and, for sampled generators, two positive generators with all integer-time samples equal, exact identification from two incommensurate times, and the absence of uniform inverse stability over unbounded rates. Identifiability within a menu does not establish that the chosen states, controls or time coordinate describe a particular physical system.

Research direction and scholarly responsibility: Kirtus Justus.

Mathematical, computational and editorial assistance: ChatGPT (OpenAI). Editorial preparation of this edition: Claude (Anthropic). Assistance does not imply provider endorsement.

This manuscript is prepared for scholarly assessment. Such preparation does not establish independent peer review, experimental replication or institutional endorsement.

6 Observation, Memory, Control, and Time in Finite Models

This section studies what a specified collection of observations determines about a finite mathematical model. Its objects are conventional probability kernels, finite hidden-state models, quantum states, and continuous-time Markov generators. An observation law, an intervention, and a calibration premise are distinct inputs. Equality of observable laws will always refer to an explicitly stated experimental menu. Identifiability within such a menu does not establish that the chosen states, controls, environmental factors, or time coordinate describe a particular physical system.

We use row probability vectors. For probability measures on a finite set, $\mathrm{TV}(p,q) = \frac{1}{2}\sum_x |p_x - q_x|$. Matrix 1-norms below are entrywise unless specified otherwise; the Frobenius norm is denoted by $\|\cdot\|_F$. The vector $\mathbf{1}$ has all entries one. All logarithms in likelihood calculations are natural; binary entropy is explicitly measured in bits. Population probabilities are exact mathematical inputs unless a finite-sample statement is given.

6.1 Controlled reduction and the limits of short records

Let X, Z be finite sets, $q : X \to Z$ a surjection, and P_a a row-stochastic transition matrix on X for each allowed action a. The full process is Markov conditional on the selected action. A policy may randomize and may depend on the complete visible history, but it has no extra model-specific hidden-state information. Write

$$M_a(x, z') = \sum_{y : q(y) = z'} P_a(x, y).$$

The familiar strong-lumpability condition is the action-free version of the following statement; see Wolfer and Watanabe [1].

Theorem 6.1 (Exact controlled reduction)**.** *There are kernels Q_a on Z that reproduce the complete visible-history law for every initial distribution on X and every common visible-history policy if and only if, for every action a, $M_a(x,\cdot)$ is constant on each fiber of q. In that event the common row is $Q_a(q(x),\cdot)$.*

Proof. For necessity, choose point initial distributions at two states x, x' in one fiber and apply the same fixed action. Both reduced initial states are $q(x)$; their next-observation laws must therefore be equal. For sufficiency, condition on any visible history of positive probability and the next selected action. The conditional law of the hidden state is supported in the fiber of the current observation. Mixing identical pushed rows over that posterior gives the stated Q_a row. The policy uses the same conditional distribution of actions in both descriptions. Induction over the finite history proves equality, including the recorded actions. □

If only specified starting states are admitted, the fiber condition need only be imposed on the appropriately reachable domain. Reduction for one special initial law is a weaker property and does not imply the all-start statement.

Example 6.2 (Fair pairs with predictive memory)**.** On states $00, 01, 10, 11$, observe the final bit and use

$$P = \begin{pmatrix} 3/4 & 1/4 & 0 & 0 \\ 0 & 0 & 1/4 & 3/4 \\ 1/4 & 3/4 & 0 & 0 \\ 0 & 0 & 3/4 & 1/4 \end{pmatrix}.$$

The uniform distribution is stationary because each column sums to one. Both append choices have positive probability; the shift graph is strongly connected and has self-loops, so the chain is irreducible and aperiodic. In stationarity every adjacent observed pair has probability $1/4$. Nevertheless,

$$P(Y_1 = a, Y_2 = b, Y_3 = c) = \frac{1}{8}\left[1 + \frac{1}{2}(-1)^{a+b+c}\right], \qquad \mathrm{TV}(P_{Y_{1:3}}, \mathrm{Unif}\{0,1\}^3) = \frac{1}{4}.$$

For example, the next-zero probabilities after 00 and 10 are $3/4$ and $1/4$, respectively. The visible process is not first-order Markov. The best uniform row approximation by a kernel depending on the last bit has error $1/4$: the two relevant Bernoulli rows have distance $1/2$, and their fair midpoint attains half that distance.

Hidden state is not by itself an obstruction. For example,

$$P((z,h),(z',h')) = Q(z,z')\,\mathbf{1}_{\{h'=1-h\}}$$

has a perpetually alternating hidden bit and an exactly autonomous visible kernel Q.

Theorem 6.3 (Uniform row error controls complete histories)**.** *Suppose, uniformly over all admitted x, a,*

$$\mathrm{TV}(M_a(x,\cdot), Q_a(q(x),\cdot)) \le \delta, \qquad 0 \le \delta \le 1.$$

Start with the same visible law and use the same visible-history policy. After N transitions the total variation distance between complete visible histories, including actions, is at most

$$1 - (1-\delta)^N \le N\delta.$$

The first bound is sharp.

Proof. Couple the initial observations identically. As long as the histories agree, use the same policy randomness. The next visible law of the full model is a posterior mixture of pushed rows, hence is within δ of the reduced row by convexity of total variation. A maximal coupling keeps the next observations equal with probability at least $1-\delta$. Consequently the coupled histories agree with probability at least $(1-\delta)^N$. The coupling inequality proves the claim. For sharpness, let the reference chain remain at state 0, with state 1 absorbing if ever entered. Let the comparison chain escape from 0 to absorbing 1 with probability δ on each step. The row premise holds at both states. The reference history is all zero, and its probability under the comparison model is exactly $(1-\delta)^N$. □

Thus a test of size α for the reduced history law has size at most $\min\{1, \alpha + 1 - (1-\delta)^N\}$ for the full model. An average stationary row error supplies no such uniform conditional premise.

An intervention on fine states need not define an intervention on coarse states. As a simple exact example, let $q(1) = q(2) = 0$, $q(3) = q(4) = 1$. The point masses at $1, 2$ have identical coarse images. A permutation interchanging $2, 3$ while fixing $1, 4$ sends them to distinct coarse point masses. No map of the original coarse distribution can represent that operation on both inputs. The quantum counterpart is equally direct: Z-basis dephasing sends both $|+\rangle\langle+|$ and $|-\rangle\langle-|$ to $I/2$, while a Hadamard operation followed by that dephasing distinguishes them perfectly. Specifying the admitted interventions is therefore part of specifying a reduced model.

Example 6.4 (A full-support non-descent witness)**.** For the same partition $\{1,2\},\{3,4\}$, take the strictly positive fine laws

$$p = (3/5, 3/20, 1/5, 1/20), \qquad p' = (3/20, 3/5, 1/5, 1/20).$$

Both have coarse law $(3/4, 1/4)$. Interchanging the second and third fine labels sends them to coarse laws $(4/5, 1/5)$ and $(7/20, 13/20)$, of total variation $9/20$. Thus the failure is not confined to boundary point masses. Exactly eight of the twenty-four fine permutations descend for all fine distributions: the permutation must send each whole input block to one output block, giving two block permutations and two internal permutations in each block, hence $2(2!)^2 = 8$.

6.2 A parity-shift process and its exact innovation coordinates

Fix $L \geq 1$ and $-1 < \eta < 1$. The hidden state is the last L bits, initially uniform. From $x = (x_1, \dots, x_L)$, append b with probability

$$P_\eta(x, (x_2, \dots, x_L, b)) = \frac{1 + \eta(-1)^{x_1 + \cdots + x_L + b}}{2}. \tag{6.1}$$

This family is a parity form of the two-faced chains studied by Ryabko [2], with parameter $p = (1+\eta)/2$.

Proposition 6.5 (Finite-depth agreement)**.** *The uniform law on L-bit states is stationary. Every contiguous observed block of length at most L is uniform, whereas a block of length $L+1$ has law*

$$2^{-(L+1)} \left[1 + \eta(-1)^{\sum_{j=1}^{L+1} y_j}\right].$$

Its distance from the uniform law is $|\eta|/2$. For $\eta \neq 0$, the observed process has Markov order exactly L.

Proof. Each destination state has two predecessors, differing in their dropped bit. Their transition probabilities sum to one. Thus a uniform state distribution remains uniform. Marginalization gives every shorter contiguous block law. Multiplying the initial 2^{-L} probability by the append probability gives the displayed $L+1$ law, whose 2^{L+1} absolute discrepancies are all $2^{-(L+1)}|\eta|$. The transition rule depends on no more than the last L bits. For $\eta \neq 0$, changing the oldest of those bits while keeping the last $L-1$ fixed changes the next-bit probability. All such histories have positive probability, proving that smaller order is impossible. □

The statement concerns contiguous blocks. It does not assert independence of every collection of L widely separated observations. Since L can be arbitrarily large, no common finite validation depth distinguishes the whole family from fair independent bits.

For a record of length $L+m$, define parity innovations

$$E_t = Y_t \oplus \cdots \oplus Y_{t+L}, \quad 1 \leq t \leq m,$$

where $\oplus$ is addition modulo two.

Theorem 6.6 (Innovation factorization)**.** *The map from $Y_{1:L+m}$ to $(Y_{1:L}, E_{1:m})$ is bijective and*

$$P(Y_{1:L} = x, E_{1:m} = e) = 2^{-L} \prod_{t=1}^{m} \frac{1 + \eta(-1)^{e_t}}{2}.$$

Consequently the innovations are independent and identically distributed, with $P(E_t = 0) = (1+\eta)/2$, and are independent of the initial block.

Proof. The recursion $Y_{t+L} = E_t \oplus Y_t \oplus \cdots \oplus Y_{t+L-1}$ recovers the record uniquely. Its probability is the initial block probability times the m append probabilities in (6.1). Each append factor depends only on e_t, giving the asserted product law. Bijection alone would not establish independence; the factorization of probabilities is essential. □

Writing h_2 for binary entropy in bits and d_{Ber} for Bernoulli relative entropy in nats, this gives

$$H(Y_{1:L+m}) = L + mh_2((1+\eta)/2), \qquad D(P_{Y_{1:L+m}} \| \,\mathrm{Unif}) = md_{\mathrm{Ber}}((1+\eta)/2 \| 1/2).$$

These are entropy and divergence per indexed observation, not a physical communication capacity or a rate per unit of laboratory time. Cancellation of overlapping parities also gives

$$(-1)^{Y_t + Y_{t+L+1}} = (-1)^{E_t + E_{t+1}}, \qquad P(Y_t = a, Y_{t+L+1} = b) = \tfrac{1}{4}[1 + \eta^2(-1)^{a+b}].$$

The two retained outcomes therefore have distance $\eta^2/2$ from a fair independent pair, although their acquisition spans $L+2$ process positions. Repeated overlapping sparse pairs are not automatically independent trials. For $L \geq 2$, pairs at lags $1, \ldots, L$ are independent and fair: for lag L, marginalizing any other coordinate of the $L+1$ parity law removes its perturbation, and smaller lags follow from the uniform block laws. The qualifier $L \geq 2$ is necessary.

6.2.1 Ideal fixed-horizon testing

Lemma 6.7 (Finite likelihood-ratio ordering)**.** *For two strictly positive probability laws p_0, p_1 on a finite sample space, a randomized test of prescribed size α has maximum power by rejecting in decreasing order of p_1/p_0, randomizing only on a boundary level.*

Proof. If the threshold is c and the proposed test is ϕ_*, then $(\phi_* - \phi)(p_1 - cp_0) \geq 0$ pointwise for every other test $0 \leq \phi \leq 1$. Summation and equal size imply $\sum \phi_* p_1 \geq \sum \phi p_1$. A boundary randomization fills the required size. This is the finite Neyman–Pearson argument [3]. □

With known L, error-free readout, and alternatives $\eta = 0$ versus $\eta = 1/2$, let $C_m = \sum_{t=1}^{m} \mathbf{1}_{\{E_t = 0\}}$. The likelihood ratio of the complete record is $3^{C_m}/2^m$; the initial block is ancillary. Thus

$$C_m \sim \mathrm{Bin}(m, 1/2) \quad \text{or} \quad \mathrm{Bin}(m, 3/4).$$

At size $1/20$, the optimum powers for $m = 39, 40$ are respectively $0.9498706279714081\ldots$ and $0.9530027582141789\ldots$. At $m = 40$, reject for $C_m \geq 26$ and, when $C_m = 25$, reject with probability $4423136507/16760560440$. The exact power is

$$\frac{7200685253556740462047 29}{7555786372591432341913 60}.$$

Optimal full-record power cannot decrease with the horizon, because an extra observation can be discarded. The failure at 39 and success at 40 therefore establish the minimum within this randomized fixed-horizon experiment. It uses $L + 40$ raw outputs. The nonrandomized rule $m = 42, C_m \geq 27$ is a sufficient alternative, with size $0.04421477349\ldots$ and power $0.95837130194\ldots$; no global nonrandomized minimum is asserted.

6.2.2 Independent readout errors create dependent innovations

Now observe $\widetilde{Y}_t = Y_t \oplus N_t$, where the N_t are iid Bernoulli(ϵ), independent of the entire source, and $0 \le \epsilon < 1/2$. Put $a = 1 - 2\epsilon$, $W_t = (-1)^{E_t}$, $V_j = (-1)^{N_j}$, and

$$Z_t = (-1)^{\widetilde{E}_t} = W_t \prod_{j=t}^{t+L} V_j.$$

Proposition 6.8 (Noisy innovation law)**.** *The mean and covariance function are*

$$\mu = \eta a^{L+1}, \quad \Gamma_0 = 1 - \mu^2, \quad \Gamma_h = \begin{cases} \eta^2(a^{2h} - a^{2L+2}), & 1 \le h \le L, \\ 0, & h \ge L+1. \end{cases}$$

Collections of Z's separated by more than L indices are independent. For $C_m = \sum_{t=1}^{m}(1+Z_t)/2$,

$$\mathrm{Var}(C_m) = \frac{1}{4}\left[m\Gamma_0 + 2\sum_{h=1}^{\min\{L,m-1\}}(m-h)\Gamma_h\right].$$

Proof. The W_t are iid of mean η, while the independent V_j have mean a. In $Z_t Z_{t+h}$, the $L+1-h$ common detector factors square to one when $h \le L$, leaving $2h$ factors and expectation $\eta^2 a^{2h}$. For larger h, the defining independent variables are disjoint, giving expectation μ^2 and the stronger setwise independence statement. Summing the covariances gives the variance. □

The observed initial L-bit block remains uniform and independent of the complete observed innovation sequence: condition on all innovations and all detector bits, and use the uniform independent source initial block. Thus the ordered noisy innovations are sufficient within this specified family. Their count generally is not. For $L = 2, \eta = 1/2, \epsilon = 1/10$,

$$P(\widetilde{E}_{1:3} = 001) = 681/5000, \qquad P(\widetilde{E}_{1:3} = 010) = 609/5000,$$

although both words have the same count. At $m = 40$, the count mean is $628/25$ and its variance $185736/15625 = 11.887104$, rather than the attenuated-binomial value 9.34464.

Proposition 6.9 (Exact noisy count recursion)**.** *Let $A_t(h,k)$ be the joint probability of the most recent L detector bits h and k even observed innovations after t steps. Initialize*

$$A_0(h,0) = \epsilon^{|h|}(1-\epsilon)^{L-|h|}.$$

For $b \in \{0,1\}$, put $r_b = \epsilon^b(1-\epsilon)^{1-b}$, $h' = (h_2, \dots, h_L, b)$, and $q(h,b) = [1 + \eta(-1)^{|h|+b}]/2$. For each current mass, add

$$A_t(h,k)r_b q(h,b) \text{ to } A_{t+1}(h', k+1), \qquad A_t(h,k)r_b[1-q(h,b)] \text{ to } A_{t+1}(h',k).$$

Then $P(C_m = k) = \sum_h A_m(h,k)$.

Proof. The next detector bit is independent of the current detector history. The next source innovation is independent of all preceding innovations and all detector bits. Conditional on the detector-window parity, its observed even probability is exactly $q(h,b)$. The displayed update is the law of total probability on these two new independent inputs. Induction proves the result. □

For rational parameters this is an integer recurrence after keeping one common denominator at each step. Applying Lemma 6.7 to the count laws, sorting their actual likelihood ratios rather than presuming their order, gives the following finite certificate for $L = 2, \eta = 1/2, \epsilon = 1/10$ against the fair iid null. Every $m = 1, \dots, 182$ has optimal count-only power below 0.95 at size 0.05; $m = 183$ has power $0.95047965645\dots$. At $m = 182$ it is $0.94920331752\dots$. An easily specified sufficient rule at 183 rejects for $C_m \geq 104$ and with probability $43/50$ at $C_m = 103$; its size and power are $0.04986347138\dots$ and $0.95033569239\dots$. The exact recurrence and rational comparisons are supplied with the computational materials. This minimum is for tests of C_m only. Unlike the complete record, C_{m+1} need not recover C_m; monotonicity of the count experiments is not used. Here 183 innovations require 185 raw outputs. A nonrandomized sufficient rule uses 188 innovations and rejects for $C_m \geq 106$; its size is $0.04659560696\dots$ and power $0.95203135112\dots$. No nonrandomized minimum is claimed.

The ideal 40-innovation rule has power only $0.49258836198\dots$ under this noisy alternative, while retaining its null size. Replacing its count law by a binomial with the attenuated mean is incorrect. If instead parity windows start $L + 1$ indices apart, their defining source innovations and detector windows are disjoint, so the selected parities are iid. That thinning consumes additional raw outputs.

At the population level, known L, $\eta \neq 0$, and $a > 0$ allow recovery from $\mu = EZ_t$ and $\nu = EZ_tZ_{t+1}$:

$$a = (\mu^2/\nu)^{1/(2L)}, \qquad \epsilon = (1-a)/2, \qquad \eta = \mu/a^{L+1}.$$

Indeed $\mu^2/\nu = a^{2L}$ and $\nu = \eta^2 a^2$. The cases $\eta = 0$ and $a = 0$ are degenerate, and the inversion is poorly conditioned near them. These identities do not supply a finite-data confidence interval or validate the iid detector model.

6.3 Classical realization rank and finite-sample certificates

Let $H_L(x, y) = P(Y_{1:L} = x, Y_{L+1:2L} = y)$. Any D-state classical hidden realization with a sufficient state at this boundary factors the matrix as

$$H_L(x, y) = \sum_{s=1}^{D} P(Y_{1:L} = x, S_L = s) P(Y_{L+1:2L} = y \mid S_L = s).$$

Hence $\operatorname{rank} H_L \leq D$. Every source, detector, or controller variable that carries predictive information across the cut belongs in that sufficient state. General hidden-model realization theory distinguishes this necessary rank condition from the existence of a positive realization [4, 5].

Theorem 6.10 (Parity-block spectrum and exact state count)**.** *In the ideal parity-shift model the singular values of H_L are*

$$2^{-L}|\eta|^j, \qquad j = 0, \dots, L,$$

with multiplicity $\binom{L}{j}$. For $\eta \neq 0$, the minimum number of states of an exact classical transition-emitting realization is 2^L. It is one when $\eta = 0$.

Proof. Define separate bijections of the row and column bit strings by $u_t = x_t \oplus \cdots \oplus x_L$ and $v_t = y_1 \oplus \cdots \oplus y_t$. The tth innovation following the cut is $u_t \oplus v_t$. The innovation factorization therefore gives, up to separate row and column permutations,

$$H_L = 2^{-L} B_\eta^{\otimes L}, \qquad B_\eta = \frac{1}{2}\begin{pmatrix} 1+\eta & 1-\eta \\ 1-\eta & 1+\eta \end{pmatrix}.$$

The singular values of B_η are $1, |\eta|$; tensor products give the asserted list. Separate row and column permutations preserve singular values, though they need not be a similarity transformation. For nonzero η the rank is 2^L, providing the lower bound. The shift-state construction (6.1) has exactly that many states and provides the matching positive upper realization. For $\eta = 0$ the outputs are iid fair bits. □

This 2^L result belongs to the parity-shift model. A stationary random-phase concatenation of perturbed k-bit blocks is a different process, with a different $2k-1$ realization result. The parameters L, k, their record laws, and their state counts are not interchangeable. Neither count is a bound on physical constituents or on quantum memory.

Theorem 6.11 (Readout and the noisy Walsh spectrum)**.** *Independent symmetric readout with $a = 1 - 2\epsilon > 0$ preserves the rank 2^L for $\eta \neq 0$. For $s \in \{0,1\}^L$, define*

$$r_i(s) = \bigoplus_{j=1}^{i} s_j, \qquad \ell_i(s) = \bigoplus_{j=i}^{L} s_j.$$

The noisy matrix has singular values, indexed by all s,

$$\sigma_s = 2^{-L} |\eta|^{|s|} a^{|r(s)|+|\ell(s)|}.$$

In particular $\sigma_{\min} \geq (|\eta| a^2/2)^L > 0$.

Proof. Let $D_\epsilon = \left(\begin{smallmatrix} 1-\epsilon & \epsilon \\ \epsilon & 1-\epsilon \end{smallmatrix}\right)$. Independent noise on both blocks gives $\widetilde{H}_L = D_\epsilon^{\otimes L} H_L (D_\epsilon^{\otimes L})^T$. Its two eigenvalues are $1, a$, so it is invertible for $a > 0$. For the full spectrum, expand each transition factor to obtain

$$H_L(x,y) = 2^{-2L} \sum_s \eta^{|s|} (-1)^{r(s)\cdot x + \ell(s)\cdot y}.$$

The maps $s \mapsto r(s)$, $s \mapsto \ell(s)$ are bijections. The normalized Walsh vectors $2^{-L/2}(-1)^{v\cdot x}$ form orthonormal bases. In these bases, H_L is a weighted permutation matrix with weights $2^{-L}\eta^{|s|}$. Readout multiplies its row and column Walsh coefficients by $a^{|r(s)|}$ and $a^{|\ell(s)|}$. The absolute weights are the singular values. Since $|s| \leq L$ and $|r(s)|+|\ell(s)| \leq 2L$, the lower bound follows. □

The same 2^L-state realization with memoryless noisy emission gives the upper bound after readout. At $\epsilon = 1/2$ the observed process is iid fair and the rank collapses. Exact invertibility does not imply a well-conditioned statistical inversion.

Lemma 6.12 (Singular-tail distance)**.** *For a real matrix H with singular values in decreasing order, define $T_D(H) = \sum_{j>D} \sigma_j(H)^2$. Then*

$$d_D(H) := \inf_{\operatorname{rank} Q \leq D} \|H - Q\|_F = \sqrt{T_D(H)}, \qquad |d_D(H) - d_D(G)| \leq \|H - G\|_F.$$

Proof. If Q has rank at most D, its nullspace contains a subspace of dimension equal to the number of columns minus D. Let P be the orthogonal projection onto such a subspace. Then $\|H - Q\|_F^2 \geq \|(H-Q)P\|_F^2 = \|HP\|_F^2$. In an eigenbasis of $H^T H$, the latter is a weighted sum of its eigenvalues with weights in $[0,1]$ summing to the rank of P. It is at least the sum of the smallest required eigenvalues, which is $T_D(H)$. Keeping the largest D terms of a singular-value decomposition attains this lower bound. This is the Eckart–Young approximation theorem [6]. Finally, distance to any fixed nonempty subset is 1-Lipschitz by the triangle inequality; taking infima proves the last statement. □

The rank-D approximation just used is allowed to be any real matrix. It need not be nonnegative, normalized, or realizable by a hidden process.

Theorem 6.13 (Total variation bounds with marginal information)**.** *Let H, Q be normalized nonnegative joint matrices, with* $\operatorname{rank} Q \le D$*. Write*

$$\delta = \mathrm{TV}(H, Q), \quad r = \mathrm{TV}(H_X, Q_X), \quad c = \mathrm{TV}(H_Y, Q_Y), \quad T = T_D(H).$$

Then

$$T \le \|H - Q\|_F^2 \le \min\{2\delta^2, \delta(\delta + r + c)\},$$

and therefore

$$\delta \ge \max\left\{ \sqrt{T/2}, \frac{\sqrt{(r+c)^2 + 4T} - (r+c)}{2} \right\}.$$

When both marginals match, the second bound becomes $\delta \ge \sqrt{T}$.

Proof. Put $A = H - Q$. Its positive and negative entry masses both equal δ. The sum of squares within each sign is at most the square of its total mass, proving $\|A\|_F^2 \le 2\delta^2$. For a positive entry $A_{ij} = x$, the negative mass in row i is at least $x - r$, since the positive part of any row sum is at most r. The negative mass in column j is at least $x - c$. These sets of negative entries are disjoint, because their only possible intersection is the positive entry (i, j). Consequently $\delta \ge 2x - r - c$. Applying the same argument to $-A$ gives $\|A\|_{\max} \le (\delta + r + c)/2$. Thus $\|A\|_F^2 \le \|A\|_{\max}\|A\|_1 \le \delta(\delta + r + c)$. Lemma 6.12 supplies $T \le \|A\|_F^2$; solving the two quadratic inequalities proves the result. □

Known upper bounds on $r + c$ may be substituted, because the second lower bound decreases with that sum. For zero row and column sums, $\|A\|_F \le \delta$ is sharp, as the signed 2-by-2 cycle $\left(\begin{smallmatrix} x & -x \\ -x & x \end{smallmatrix}\right)$ shows. That norm sharpness does not show that a rank-constrained probability model attains the combined bound.

For $L = 2, \eta = 1/2, \epsilon = 1/10$, the exact noisy matrix is

$$\widetilde{H}_2 = \frac{1}{2000} \begin{pmatrix} 209 & 105 & 81 & 105 \\ 81 & 105 & 209 & 105 \\ 105 & 81 & 105 & 209 \\ 105 & 209 & 105 & 81 \end{pmatrix}. \tag{6.2}$$

Its singular values are $1/4, 8/125, 8/125, 1/25$, and

$$T_1 = 153/15625, \qquad T_2 = 89/15625, \qquad T_3 = 1/625.$$

Both marginals are uniform on four outcomes. A normalized rank-one nonnegative matrix with those same marginals is uniquely the matrix whose entries are $1/16$, and its total variation distance is exactly $21/125$. The iid fair-bit process attains this restricted rank-one competitor. The number does not apply when competitor marginals may vary; the bounds for two or three states have not been asserted attainable.

6.3.1 From population rank to finite samples

Lemma 6.14 (Bounded-differences inequality)**.** *Let $X_1, \dots, X_n$ be independent, and suppose changing only X_i changes f by at most c_i. Then*

$$P(f - Ef \ge t) \le \exp\left(-\frac{2t^2}{\sum_i c_i^2} \right).$$

Proof. The conditional Doob martingale of f has each martingale difference in an interval of length at most c_i: conditional expectations for two possible values of X_i differ by at most c_i, after coupling the still independent remaining coordinates identically. For a centered variable U in an interval of length c, $\log Ee^{\lambda U} \le \lambda^2 c^2/8$. Indeed the second derivative of its log moment-generating function is the variance under exponential tilting, which is at most $c^2/4$; the function and its first derivative vanish at zero, so integration twice gives the bound. The variance bound follows from $E[(U-a)(b-U)] \ge 0$, which gives $\mathrm{Var}(U) \le (EU-a)(b-EU) \le (b-a)^2/4$. Apply the exponential bound conditionally to the martingale differences, iterate, use Markov's inequality, and minimize over $\lambda = 4t/\sum c_i^2$. This is the standard bounded-differences method [7]. □

Theorem 6.15 (Empirical matrix radius and rank test)**.** *Fix a matrix shape with K cells, an observation map, and a sample size n. Observe n independent records having the same categorical joint law H, and let $\widehat{H}$ be their frequency matrix. Dependence within a record is allowed. For $0 < \alpha < 1$, set*

$$\varepsilon_\alpha(n,K) = \frac{\sqrt{1-1/K} + \sqrt{\log(1/\alpha)}}{\sqrt{n}}.$$

Then $P(\|\widehat{H}-H\|_F \le \varepsilon_\alpha) \ge 1-\alpha$. The test rejecting rank at most D when $d_D(\widehat{H}) > \varepsilon_\alpha$ has size at most α. If $d_D(H) > \varepsilon_\alpha + \varepsilon_\beta$, its power is at least $1-\beta$. On the same confidence event, simultaneously for every D,

$$\inf_{\substack{Q\ge 0,\ \sum Q=1 \\ \operatorname{rank} Q \le D}} \mathrm{TV}(H,Q) \ge \frac{[d_D(\widehat{H}) - \varepsilon_\alpha]_+}{\sqrt{2}}.$$

Proof. The multinomial second-moment identity gives

$$E\|\widehat{H}-H\|_F^2 = \frac{1-\|H\|_F^2}{n} \le \frac{1-1/K}{n}.$$

Jensen's inequality bounds the mean norm by the square root. Replacing one record changes $\widehat{H}$ by Frobenius norm at most $\sqrt{2}/n$. The reverse triangle inequality gives the same sensitivity for its error norm. Lemma 6.14 therefore gives an upper tail $\exp(-nt^2)$; use $t = \sqrt{\log(1/\alpha)/n}$. Under the rank null, $d_D(H) = 0$; Lemma 6.12 proves the size bound. On the radius event with probability at least $1-\beta$, $d_D(\widehat{H}) \ge d_D(H) - \varepsilon_\beta > \varepsilon_\alpha$, which proves power. No additional $\alpha+\beta$ penalty is needed for that power statement. Finally, $d_D(H) \ge [d_D(\widehat{H}) - \varepsilon_\alpha]_+$, and Theorem 6.13's unrestricted bound applies to every normalized rank-D competitor. □

At the single predeclared matrix and horizon, the confidence event simultaneously controls all D; repeated looks at changing horizons are a different experiment. Complete categorical outcomes, including any failure outcome required by the experiment, must be represented. A numerically computed singular tail requires an error allowance if rounding can affect the threshold.

For (6.2), $K = 16$ and $\alpha = \beta = 1/20$. Let $C = \sqrt{15/16} + \sqrt{\log 20}$. The condition is $n > 4C^2/T_D$. Its first integer solutions are

D	1	2	3
n	2976	5116	18213
raw outputs $4n$	11904	20464	72852

These are sufficient numbers of iid four-output records for this concentration bound, not optimal testing budgets. At $n = 18213$ the single common power event suffices for all three rank rejections. Experiments separately stopped at the three listed horizons do not automatically constitute one test of size 0.05.

6.4 Sequential evidence for complete record laws

The null in this subsection is the complete iid fair-bit law, $P_0(Y_{1:n} = y) = 2^{-n}$. Fair one-bit marginals alone do not define this null. Fix coherent candidate laws P_θ before seeing data and nonnegative mixing weights totaling one.

Theorem 6.16 (Likelihood mixtures and an anytime threshold)**.** *Each* $R_{\theta,n} = 2^n P_\theta(Y_{1:n})$ *is a nonnegative* P_0*-martingale starting at one. Their fixed mixture* $\mathcal{E}_n = \int R_{\theta,n}\, dW(\theta)$ *is also such a martingale, and*

$$P_0\{\exists n \geq 0 : \mathcal{E}_n \geq 1/\alpha\} \leq \alpha.$$

Proof. Conditional on a prefix under P_0, the next bit is fair. Coherence gives

$$E_0[R_{\theta,n+1} \mid Y_{1:n}] = 2^n \sum_{b=0}^{1} P_\theta(Y_{1:n}, b) = R_{\theta,n}.$$

Nonnegative integration permits conditional Tonelli, proving the mixture claim. Stop at the first threshold crossing or at a fixed N. The expected stopped value is one by finite optional stopping, so $P_0(\text{crossing by } N) \leq \alpha$. Let N increase. The same argument holds for nonnegative supermartingales and is Ville's inequality; see Howard et al. [8]. □

For the noisy parity family with fixed $\eta = 1/2, \epsilon = 1/10$ and unknown L, take $w_L = 1/[L(L+1)]$. When $L \geq n$, every n-bit observed law is uniform. Since $\sum_{L=n}^{\infty} w_L = 1/n$, the mixture has the finite expression

$$\mathcal{E}_n = \frac{1}{n} + \sum_{L=1}^{n-1} \frac{2^n P_L(Y_{1:n})}{L(L+1)}, \qquad n \geq 1, \quad \mathcal{E}_0 = 1. \tag{6.3}$$

This identity does not guarantee inexpensive evaluation: a component may require 2^L hidden states. Replacing the fixed mixture by a post-data maximum does not preserve its martingale proof.

Proposition 6.17 (A sufficient 1950-output horizon)**.** *At the fixed true alternative* $L = 2, \eta = 1/2, \epsilon = 1/10$*, the test crossing* 20 *in* (6.3) *has power by* 1950 *raw outputs at least* $0.9519745696463602\ldots$*.*

Proof. The 650 disjoint three-bit parity windows have independent even-parity probability $p = 157/250$. Let K be their even count and let

$$S = (157/125)^K (93/125)^{650-K}$$

be their likelihood ratio against fair parity. For the full-record component $R = 2^{1950} P_2(Y_{1:1950})$, strict positivity gives $E_2(S/R) = E_0 S = 1$. The mixture contains $R/6$. Failure to cross by this horizon implies in particular $R < 120$. For any $c > 0$,

$$P_2(R < 120) \leq P_2(S < 120c) + P_2(S/R > c) \leq P_2(S < 120c) + 1/c.$$

With $c = 40$, exact integer comparisons show $S \geq 4800$ if and only if $K \geq 384$. Therefore the miss probability is at most

$$\sum_{k=0}^{383} \binom{650}{k} (157/250)^k (93/250)^{650-k} + \frac{1}{40} = 0.04802543035363984\ldots < \frac{1}{20}.$$

All quantities in the last inequality are rational. The calculation does not require evaluating all nonconstant mixture components. □

This is a sufficient horizon for one specified alternative, not a minimum, an expected stopping time, or a bound uniform in L. It is a different experiment from the fixed-horizon 183-innovation count test.

Theorem 6.18 (Pointwise eventual detection)**.** *Fix a nondegenerate parity candidate with $L < \infty$, $\eta \neq 0$, $0 \leq \epsilon < 1/2$, and positive mixture weight w_L. If that candidate generates the data, its likelihood mixture eventually crosses every fixed finite threshold almost surely. More precisely,*

$$\liminf_{n\to\infty} \frac{\log \mathcal{E}_n}{n} \geq \frac{d_{\mathrm{Ber}}(p_L \| 1/2)}{L+1} > 0, \qquad p_L = \frac{1 + \eta(1-2\epsilon)^{L+1}}{2}.$$

Proof. Use $m = \lfloor n/(L+1) \rfloor$ disjoint parity windows, and let S_m be their likelihood ratio. The iid strong law gives $\log S_m/n \to d_{\mathrm{Ber}}(p_L\|1/2)/(L+1)$. For the full-record ratio $R_{L,n}$, the same change-of-measure identity as above gives $E_L[S_m/R_{L,n}] = 1$. For every $b > 0$,

$$P_L(R_{L,n} < S_m e^{-bn}) \leq e^{-bn}.$$

The sum over n is finite, so the first Borel–Cantelli lemma implies that this event occurs only finitely often. Apply the assertion to a countable sequence of positive rational b decreasing to zero and use $\mathcal{E}_n \geq w_L R_{L,n}$. The fixed multiplicative weight has vanishing logarithm per n. □

No uniform finite horizon follows: for $L \geq n$ the full length-n law equals the null exactly. Alternatives approaching $\eta = 0$ or $\epsilon = 1/2$ also approach the null on each fixed horizon. For a concrete warning about the null, the process $Y_t = B$ for one fair bit B has fair marginals. In the ideal-readout $\eta = 1/2$ mixture, its $L = 1$ contribution is $\frac{1}{2}(3/2)^{n-1}$, exceeding 20 at $n = 11$. It is not an iid-null process; rejecting it does not violate the theorem.

6.4.1 Conditional null envelopes and useful growth

Suppose instead that the null supplies a predictable envelope

$$p_t = P(Y_t = 1 \mid \mathcal{F}_{t-1}) \in [1/2 - \delta_t, 1/2 + \delta_t], \quad 0 \leq \delta_t \leq 1/2.$$

For a predictable Bernoulli forecast q_t, put $q_t(1) = q_t$ and $q_t(0) = 1 - q_t$. The factor

$$F_t = \frac{2q_t(Y_t)}{1 + 2\delta_t|2q_t - 1|}$$

has conditional null expectation at most one: its numerator has expectation $1 + 2(p_t - 1/2)(2q_t - 1)$, maximized exactly at the appropriate endpoint of the interval. Its products obey the crossing guarantee of Theorem 6.16. This sharp normalization does not ensure favorable evidence growth.

Theorem 6.19 (Log-optimal binary factor for an interval null)**.** *For $0 < \ell \le u < 1$ and an alternative $0 < q < 1$, let $p_* = \min\{u, \max\{\ell, q\}\}$. Define*

$$G_q(1) = q/p_*, \qquad G_q(0) = (1-q)/(1-p_*).$$

It has expectation at most one for every Bernoulli parameter in $[\ell, u]$, and maximizes $E_q \log g(Y)$ over all nonnegative factors g valid throughout that interval.

Proof. If $q > u$, the expectation of G_q increases with the actual Bernoulli parameter and equals one at $u = p_*$. If $q < \ell$, it decreases and equals one at $\ell = p_*$. If $q \in [\ell, u]$, it is identically one. For every valid g, Jensen's inequality gives

$$E_q \log \frac{g(Y)}{G_q(Y)} \le \log E_q \frac{g(Y)}{G_q(Y)} = \log E_{p_*} g(Y) \le 0.$$

Zero factors are handled by the extended value $-\infty$ for the log. □

This is a binary specialization of the established growth-optimal e-variable framework [9]. Predictable choices of ℓ_t, u_t, q_t again yield a nonnegative null supermartingale. The criterion is expected one-step log growth, not finite-horizon Neyman–Pearson power.

For $q = 13/20$ and null interval $[2/5, 3/5]$, the two factors are

$$(F(0), F(1)) = (35/53, 65/53), \qquad (G(0), G(1)) = (7/8, 13/12).$$

Their expected log growth under iid Bernoulli$(13/20)$ is negative and positive, respectively. The exact sign certificates are

$$65^{13}35^7 < 53^{20}, \qquad 13^{13}7^7 > 12^{13}8^7.$$

The strong law makes the first product tend to zero and the second diverge almost surely. The first can still cross a threshold early.

A conditional-bias envelope is not a memorylessness null. In the noisy parity model, conditioning first on the full source history and then averaging over the observed past yields

$$\left| P(\widetilde{Y}_t = 1 \mid \widetilde{Y}_{<t}) - \frac{1}{2} \right| \le \frac{|\eta|(1-2\epsilon)}{2}.$$

For the numerical alternative this is $1/5$. A null envelope at least that wide contains the dependent process, so a test valid for every law in it must retain its null error guarantee on that process. The envelope itself needs justification for the actual observation filtration; marginal calibration does not supply it.

6.5 Visibility, environmental records, and access to fragments

Consider an ordinary qubit and an environment with a fixed tensor decomposition into N fragments. A stipulated controlled interaction maps the two qubit basis states to branch records

$$|0\rangle \longmapsto |0\rangle|e_0\rangle^{\otimes N}, \qquad |1\rangle \longmapsto |1\rangle|e_1\rangle^{\otimes N}, \quad \langle e_0|e_1\rangle = \gamma, \quad |\gamma| \le 1.$$

For a balanced input, tracing out the fragments leaves coherence of magnitude $|\gamma|^N/2$. A phase setting with the convention $p_0(\phi) = \frac{1}{2}[1 + \mathrm{Re}(e^{i\phi}\gamma^N)]$ has visibility $|\gamma|^N$ and qubit purity $(1 + |\gamma|^{2N})/2$. These formulas assume the stated state preparation, tensor partition, controlled interaction, and Born-rule measurement. They derive no macroscopic amplification mechanism.

Proposition 6.20 (A classical realization of the binary visibility law)**.** *For real* $0 \le \gamma \le 1$*, all final binary laws*

$$p_0(N,\phi) = \tfrac{1}{2}(1+\gamma^N \cos\phi), \qquad N \in \{0,1,...\},\ \phi \in \mathbb{R},$$

are reproduced by a two-state classical Markov chain and a setting-dependent binary response.

Proof. Let $Z_0 = +1$. Flip the sign independently at each step with probability $(1-\gamma)/2$, so that the transition matrix is

$$K_\gamma = \frac{1}{2}\begin{pmatrix} 1+\gamma & 1-\gamma \\ 1-\gamma & 1+\gamma \end{pmatrix}.$$

Then $EZ_N = \gamma^N$, by the one-step conditional expectation $E(Z_{n+1} \mid Z_n) = \gamma Z_n$. Given $Z_N = z$ and the calibrated setting ϕ, report outcome zero with probability $(1+z\cos\phi)/2$. Averaging proves the formula. At $N = 0$ the chain has made no transition, including when $\gamma = 0$. □

For complex $\gamma = re^{i\beta}$, one may instead choose independent classical phases $\Theta_j = \beta$ or $\beta + \pi$ with probabilities $(1+r)/2, (1-r)/2$. Since $Ee^{i\Theta_j} = \gamma$, the response $[1 + \cos(\phi + \sum_j \Theta_j)]/2$ has the required averaged law. This construction uses accumulated phase; it is not the same assertion of a homogeneous two-state realization with one fixed response for all complex γ. Related random-unitary descriptions of quantum noise are established [10]; their scope differs from this explicitly classical simulation of a specified final binary family. Neither construction simulates arbitrary quantum preparations, joint measurements, Bell experiments, or every environmental observable.

Theorem 6.21 (A final channel does not identify fragment factorization)**.** *Fix the physical fragment partition and let* $0 < \gamma < 1$*,* $s = \sqrt{1-\gamma^2}$*. For* $N \ge 2$*, the product branches*

$$E_0 = |0\rangle^{\otimes N}, \qquad E_1 = (\gamma|0\rangle + s|1\rangle)^{\otimes N}$$

and the branches $F_i = U_N E_i$*, where* $U_N = \mathrm{CZ}_{12} \otimes I_{3:N}$*, induce exactly the same complete final qubit channel, including its action with a retained ancilla. Nevertheless* F_1 *is not a product across the first two fragments. If the transformed old fragments are never accessed or recoupled, the two descriptions also give the same complete histories for common operations on the retained system, ancillas, and fresh fragments.*

Proof. For any joint state ρ of a retained system and environment,

$$\mathrm{Tr}_E[(I \otimes U_N)\rho(I \otimes U_N^\dagger)] = \mathrm{Tr}_E\,\rho.$$

This identity is valid with an arbitrary retained ancilla, proving the complete-channel statement. It also preserves the branch overlap $\langle E_0|E_1\rangle = \gamma^N$. For the first two fragments, the coefficient matrix of F_1 is

$$\begin{pmatrix} \gamma^2 & \gamma s \\ \gamma s & -s^2 \end{pmatrix}, \qquad \det = -2\gamma^2 s^2 \neq 0.$$

A pure two-fragment state is a product precisely when its coefficient matrix has rank one; thus this branch is not a product. Appending the remaining product factors does not remove that fact. For a later history satisfying the stated access restriction, the common environmental unitary commutes with every later retained-system or

fresh-fragment operation. Move it through the product of the corresponding instrument maps and remove it under the final trace. This proves equality for each fixed outcome/action history; common history-dependent randomized policies preserve it by conditioning. □

The history assertion requires absence of later coupling to the transformed fragments. Equality of final reduced channels by itself does not prove equality under arbitrary later access. Also, the distinguished property here is tensor-product factorization of conditional quantum fragment states. It is not a claim that every chosen classical fragment readout becomes statistically dependent. For example, the common controlled phase leaves the branch's Z-basis probability distribution unchanged. If arbitrary entangling changes of tensor factorization are declared to be a gauge freedom, the target factorization property itself changes. At $\gamma = 0$ or 1, or with fewer than two fragments, the stated nonproduct witness is absent; common-unitary invariance of the reduced channel still holds.

Example 6.22 (A calibrated fragment measurement separates the pair)**.** For $\gamma = 3/5, s = 4/5, N = 2$, the branch-one vectors are

$$E_1 = (9, 12, 12, 16)^T/25, \qquad F_1 = (9, 12, 12, -16)^T/25.$$

Both are normalized and have overlap $9/25$ with $|00\rangle$. Their coefficient determinants are zero and $-288/625$. The first-fragment reduced off-diagonal entries are $12/25$ and $-84/625$, so a local X measurement has

$$P_{E_1}(X = +) = 49/50, \qquad P_{F_1}(X = +) = 457/1250, \qquad \text{gap} = 384/625.$$

This requires anchored preparation of branch one, access to the specified fragment, and a calibrated X readout. If each reported binary law is within total variation ε of its stated ideal law, the gap is at least $384/625 - 2\varepsilon$; strict positivity is a sufficient separation condition. This is pair separation, not identification of an arbitrary environmental model.

6.6 Positive hidden realizations and exact intervention contracts

A finite transition-emitting, or Mealy, model has nonnegative matrices T_y, indexed by a finite output alphabet, whose sum $P = \sum_y T_y$ is row-stochastic. From an initial row distribution π,

$$P(Y_{1:m} = y_1 \cdots y_m) = \pi T_{y_1} \cdots T_{y_m} \mathbf{1}.$$

A silent control is a row-stochastic matrix inserted between output steps, producing no extra recorded outcome. Linear reachability means that the rows $\{\pi T_w\}$ span the whole row space; linear observability means that $\{T_w \mathbf{1}\}$ spans the whole column space, where w includes the empty word. We use *linearly minimal* when both properties hold. It is stronger than merely displaying no obvious redundant state.

If S is invertible with $S\mathbf{1} = \mathbf{1}$, put

$$T'_y = S^{-1} T_y S, \qquad \pi' = \pi S.$$

Whenever these transformed quantities are valid probabilities, every passive word law is unchanged by telescoping. If each common fixed control commutes with S, all controlled word laws are unchanged as well. If instead controls are themselves allowed to vary, simultaneous transformation $K'_j = S^{-1} K_j S$ gives the same telescoping identity. The distinction between exact fixed controls and uncertain controls is substantive. Similarity equivalence for Mealy models and its distinction from other hidden-model classes are established [5].

Example 6.23 (A strictly positive minimal passive pair)**.** Take

$$T_0 = \frac{1}{20}\begin{pmatrix} 6 & 2 \\ 1 & 3 \end{pmatrix}, \quad T_1 = \frac{1}{20}\begin{pmatrix} 4 & 8 \\ 7 & 9 \end{pmatrix}, \quad \pi = (4/9, 5/9), \quad S = \frac{1}{10}\begin{pmatrix} 9 & 1 \\ 2 & 8 \end{pmatrix}.$$

The transformed matrices are

$$T_0' = \frac{1}{1400}\begin{pmatrix} 449 & 151 \\ 19 & 181 \end{pmatrix}, \qquad T_1' = \frac{1}{280}\begin{pmatrix} 67 & 93 \\ 125 & 115 \end{pmatrix}, \qquad \pi' = \pi S.$$

Both representations are strictly positive and stationary: $\pi P = \pi$ and $\pi' P' = \pi'$. Their common two-output matrix is

$$J = \frac{1}{900}\begin{pmatrix} 81 & 179 \\ 179 & 461 \end{pmatrix}, \qquad \det J = 53/8100.$$

The factorization of J through a hidden boundary state implies minimal dimension at least two; the displayed models attain it. Their total-transition diagonal multisets are $\{1/2, 3/5\}$ and $\{14/25, 27/50\}$, respectively. They therefore are not related by a permutation of hidden-state labels.

Proposition 6.24 (Exactly which fixed controls separate that pair)**.** *Let the same anchored control*

$$K(a,b) = \begin{pmatrix} 1-a & a \\ b & 1-b \end{pmatrix}, \qquad 0 \le a, b \le 1,$$

act in both models of Example 6.23. All controlled output-history laws agree if and only if $b = 2a$*. The next-zero contrast after one such control is*

$$\pi K T_0 \mathbf{1} - \pi' K T_0' \mathbf{1} = (2a - b)/35.$$

Proof. Direct multiplication gives the contrast and

$$SK - KS = \frac{1}{10}\begin{pmatrix} b-2a & 2a-b \\ b-2a & 2a-b \end{pmatrix}.$$

If $b = 2a$, the similarity telescopes through every interleaving of outputs and controls. If $b \neq 2a$, the displayed single-output experiment already distinguishes the pair. Common randomized or adaptive control policies preserve the equality in the first case by conditioning on their observed histories. □

The reset $R_2 = \left(\begin{smallmatrix} 0 & 1 \\ 0 & 1 \end{smallmatrix}\right)$ maximizes the absolute single-output contrast over this control square: it gives $2/35$, with zero probabilities $1/5$ and $1/7$. This is an optimum for that contrast and that fixed pair. It does not identify the entire model class. For example,

$$U = \begin{pmatrix} 9/10 & 1/10 \\ 0 & 1 \end{pmatrix} \quad \text{satisfies} \quad UR_2 = R_2 U = R_2.$$

The further transformed model has

$$\pi U = (2/5, 3/5), \quad U^{-1} T_0 U = \frac{1}{1800}\begin{pmatrix} 531 & 229 \\ 81 & 279 \end{pmatrix}, \quad U^{-1} T_1 U = \frac{1}{1800}\begin{pmatrix} 297 & 743 \\ 567 & 873 \end{pmatrix}.$$

It is strictly positive, remains minimal, and has total-transition diagonal multiset $\{23/50, 16/25\}$. It is a further nonpermutation partner agreeing with the original on all passive and exact-R_2 controlled laws.

Theorem 6.25 (One silent control does not identify the unrestricted positive minimal class)**.** *Let $n \geq 2$, let $(\pi, \{T_y\})$ be a strictly positive linearly minimal finite-alphabet realization, and assume its unanchored preparation and output matrices may vary in a neighborhood of their current values within the model class. A single exact common row-stochastic silent control K leaves a continuum of distinct positive realizations with the same complete controlled laws. Stationarity, if imposed on π, is preserved.*

Proof. For $K \neq I$, choose $S_t = I + t(K - I)$. It satisfies $S_t\mathbf{1} = \mathbf{1}$, commutes with K, and is invertible for sufficiently small t. At $t = 0$, all entries of T_y and π are positive. Continuity and finiteness preserve positivity for a sufficiently small interval of t. Normalization follows from $S_t\mathbf{1} = \mathbf{1}$. The transformed total transition matrix and preparation satisfy $\pi' P' = \pi P S_t = \pi S_t = \pi'$ when the original is stationary. The telescoping identity proves equality for all controlled words. If $K = I$, use any nonzero matrix A with $A\mathbf{1} = 0$ and $S_t = I + tA$, again sufficiently near the identity. To see that distinct similarities give distinct numerical realizations, suppose a similarity V fixes both π and every T_y. Then $\pi T_w V = \pi T_w$ for every word. Reachability forces $V = I$. The ratio of two similarities yielding identical realizations would be such a V, hence the similarities would be equal. A continuum of distinct realizations cannot be removed by the finite set of state permutations. □

The theorem concerns a local open positive class, an unanchored preparation, and silent stochastic controls. Structural zeros, fixed preparations, recorded control outcomes, or narrower parametrizations can change the conclusion. It is not a claim about all hidden-model classes. Adding invisible hidden labels is an easier ambiguity: one may append arbitrary hidden transitions whose pushed visible rows stay fixed, as in Theorem 6.1. Such redundant lifts are not counterexamples to minimal-realization theorems and do not establish distinct physical ontologies.

Theorem 6.26 (A sharp two-control criterion in two states)**.** *Consider stationary strictly positive binary Mealy models with two states, passive linear dimension two, and an unanchored initial distribution. Suppose two labelled controls are known exactly and have the same matrices in every admissible model:*

$$K_j = \begin{pmatrix} 1 - a_j & a_j \\ b_j & 1 - b_j \end{pmatrix}, \quad j = 1, 2.$$

Complete controlled population laws identify every numerical output matrix and the labelled preparation in this class if and only if

$$\Delta = a_1 b_2 - a_2 b_1 \neq 0.$$

When $\Delta \neq 0$, experiments with at most one control followed by at most two outputs suffice for the reconstruction.

Proof. Write $\pi = (r, 1 - r)$, $e = T_0\mathbf{1} = (e_1, e_2)^T$, $d = e_1 - e_2$, $k = rd$, $q = P(0)$, and $q_j = P(K_j \text{ then } 0)$. If $d = 0$, both $T_0\mathbf{1} = q\mathbf{1}$ and $T_1\mathbf{1} = (1 - q)\mathbf{1}$. Repeated multiplication makes every word iid Bernoulli, contradicting passive dimension two. Thus $d \neq 0$. The first-state mass after K_j is $r + b_j - (a_j + b_j)r$, hence

$$\begin{pmatrix} q_1 - q \\ q_2 - q \end{pmatrix} = B \begin{pmatrix} d \\ k \end{pmatrix}, \qquad B = \begin{pmatrix} b_1 & -(a_1 + b_1) \\ b_2 & -(a_2 + b_2) \end{pmatrix}, \quad \det B = \Delta.$$

If $\Delta \neq 0$, solve for d, k, then recover

$$r = k/d, \qquad e_2 = q - k, \qquad e_1 = e_2 + d.$$

At least one $q_j - q$ is nonzero, because otherwise B would map the nonzero vector $(d,k)^T$ to zero. For such j, define

$$A = \begin{pmatrix} \pi \\ \pi K_j \end{pmatrix}, \quad F = (\mathbf{1} \quad e), \quad C_y = \begin{pmatrix} P(y) & P(y0) \\ P(K_j \text{ then } y) & P(K_j \text{ then } y0) \end{pmatrix}.$$

Then $C_y = AT_yF$, $\det A = -(q_j - q)/d$, and $\det F = -d$. Both are invertible, so $T_y = A^{-1}C_yF^{-1}$. All quantities are obtained from the stated short experiments.

Conversely, direct multiplication gives

$$[K_1, K_2] = \Delta \begin{pmatrix} 1 & -1 \\ 1 & -1 \end{pmatrix}.$$

If $\Delta = 0$ and a control $K \neq I$ exists, the nonconstant similarities $I + t(K - I)$ commute with both controls. If both controls are identity, use any nontrivial near-identity row-normalized similarity. The positivity, stationarity, and minimality argument of Theorem 6.25 provides a continuum of admissible distinct realizations with all controlled laws equal. Identification is impossible. □

For a positive non-reset example,

$$K_1 = \begin{pmatrix} 3/4 & 1/4 \\ 1/2 & 1/2 \end{pmatrix}, \qquad K_2 = \begin{pmatrix} 1/2 & 1/2 \\ 1/4 & 3/4 \end{pmatrix}, \qquad \Delta = -3/16.$$

For Example 6.23,

$$q = 13/45, \quad q_1 = 29/90, \quad q_2 = 49/180, \quad d = 1/5, \quad k = 4/45,$$

and the short-word matrices for $j = 1$ are

$$C_0 = \begin{pmatrix} 13/45 & 9/100 \\ 29/90 & 21/200 \end{pmatrix}, \qquad C_1 = \begin{pmatrix} 32/45 & 179/900 \\ 61/90 & 337/1800 \end{pmatrix}.$$

The formulas recover the displayed π, T_0, T_1 exactly. Small $|\Delta|$, small $|d|$, or a small preparation contrast can make the inverse numerically unstable; this population theorem contains no finite-sample stability bound.

Proposition 6.27 (Pure preparations and a finite-state positive control)**.** *For a two-state model with two exactly anchored labelled pure preparations, put $e_i = P_i(0)$, $p_i(y) = P_i(y)$, $z_i(y) = P_i(y0)$. If $d = e_1 - e_2 \neq 0$, then*

$$(T_y)_{i1} = \frac{z_i(y) - e_2 p_i(y)}{d}, \qquad (T_y)_{i2} = \frac{e_1 p_i(y) - z_i(y)}{d}.$$

More generally, an observable n-state realization is identified by an exact labelled cyclic permutation control and an exact reset to one labelled state, using experiments with at most n output symbols.

Proof. For two states, the equations $(T_y)_{i1} + (T_y)_{i2} = p_i(y)$ and $e_1(T_y)_{i1} + e_2(T_y)_{i2} = z_i(y)$ give the formulas. For n states, a reset followed by powers of the cycle prepares every pure labelled state. Let $V_k = \operatorname{span}\{T_v\mathbf{1} : |v| \leq k\}$. The initial space V_0 has dimension one. If $V_k = V_{k+1}$, each T_y maps V_k into itself, so all later spaces are identical. Observability therefore implies that full dimension is reached by $k = n - 1$. Choose n spanning suffix columns to form an invertible matrix F. Their probabilities under the pure preparations give every entry of F. Probabilities of yv under the same preparations give T_yF, so recover T_y. Probabilities of those suffixes from the original preparation give πF, so recover π. □

The cycle and reset contain structural zeros and permit repeated uses. The bound concerns the number of output symbols per setting; it does not count control uses, distinct settings, repeated trials, duration, or calibration cost. Together with the one-control obstruction, two types are minimal only for the stated universal ideal control-design question over positive minimal models. Restricted model classes may require fewer.

6.7 Calibration intervals and statistical separation

Exact controls and approximately known controls define different inverse problems. This can be shown quantitatively without assigning a prior to the uncertain control.

Theorem 6.28 (An explicit calibration-induced identification interval)**.** *Use the positive model of Example 6.23 and the two controls displayed after Theorem 6.26. Keep K_1 fixed exactly, but allow K_2 to lie in its full maximum-row total-variation ball of radius ε. Keep the preparation unanchored. Among strictly positive stationary models with the same complete controlled output laws, the first preparation coordinate includes the interval*

$$\left[\frac{4}{9}, \frac{4}{9} + \min\left\{\frac{1}{15}, \frac{8\varepsilon}{9}\right\}\right].$$

This is a contained interval, not the complete identified set.

Proof. For $0 \le t \le 1/10$, use

$$S_t = \begin{pmatrix} 1-t & t \\ 2t & 1-2t \end{pmatrix}, \qquad \det S_t = 1 - 3t \ge 7/10.$$

Direct multiplication gives

$$S_t^{-1} K_1 S_t = K_1, \qquad S_t^{-1} K_2 S_t - K_2 = \begin{pmatrix} 3t/4 & -3t/4 \\ 3t/4 & -3t/4 \end{pmatrix}.$$

Thus the maximum-row total variation error is $3t/4$. The transformed preparation is $\pi S_t = ((4+6t)/9, (5-6t)/9)$. For completeness, the eight numerators of the two transformed output matrices, over the positive common denominator $20(1-3t)$, are

$$\begin{matrix} 6 - 15t - t^2 & 2 - 5t + t^2 \\ 1 - 8t - t^2 & 3 - 12t + t^2 \\ 4 - 3t - 35t^2 & 8 - 37t + 35t^2 \\ 7 - 4t - 35t^2 & 9 - 36t + 35t^2 \end{matrix}.$$

On the stated interval they are bounded below, in the same order, by

$$449/100,\ 3/2,\ 19/100,\ 9/5,\ 67/20,\ 43/10,\ 25/4,\ 27/5,$$

all strictly positive. Normalization and stationarity are preserved by similarity. Simultaneously transforming T_y, π, K_2, while K_1 stays fixed, makes every controlled word telescope to its original probability. The admissible segment is $0 \le t \le \min\{1/10, 4\varepsilon/3\}$. Its first preparation coordinate is $4/9 + 2t/3$, giving the interval. □

For example, $\varepsilon = 1/100$ gives $[4/9, 34/75]$, of width $2/225$. This is uncertainty in a chosen latent coordinate, not uncertainty in the common predictions. An independently fixed preparation or a model for extra calibration outcomes can break the equivalence.

Proposition 6.29 (Finite positive menus with open calibration classes)**.** *Let $n \geq 2$, with finitely many strictly positive output matrices, a positive unanchored preparation, and finitely many strictly positive silent controls. Suppose their admissible classes contain local neighborhoods of the output/preparation parameters and relative-open neighborhoods of each control within row-stochastic matrices. If the realization is linearly minimal, there is a local continuum of distinct admissible simultaneous-similarity realizations with the same complete controlled laws.*

Proof. Choose a nonzero matrix A with $A\mathbf{1} = 0$, and $S_t = I + tA$ sufficiently near I. Transform every output matrix and every control by $S_t^{-1}(\cdot)S_t$, and the preparation by πS_t. The sum of the transformed output matrices and each transformed control are row-stochastic, and the transformed preparation sums to one, because $S_t\mathbf{1} = \mathbf{1}$. Strict positivity and the finitely many open-neighborhood conditions persist for sufficiently small t. All probabilities telescope. The trivial stabilizer argument in Theorem 6.25 proves distinctness. □

The open-neighborhood premise excludes exact structural zeros and other constraints that obstruct the path. This qualitative result provides no universal numerical interval width.

6.7.1 The reset experiment with bounded nuisance

For the fixed pair in Example 6.23, suppose the intended reset to state two may place a candidate-specific mass $u \in [0, \delta]$ on state one. The next-zero laws range over

$$p_A = (1 + u)/5, \qquad p_B = (1 + 2u)/7.$$

For $0 \leq \delta < 1/5$, the two probability intervals have strictly positive separation

$$g_\delta = \frac{1}{5} - \frac{1 + 2\delta}{7} = \frac{2 - 10\delta}{35}.$$

At $\delta = 1/5$, take the exact R_2 in the first model and

$$S^{-1}R_2S = \begin{pmatrix} 1/5 & 4/5 \\ 1/5 & 4/5 \end{pmatrix}$$

in the second. The complete controlled laws, not only the next output, then agree by simultaneous similarity. The same witness is admissible for every larger δ. Thus this fixed-pair nuisance class has a sharp 20% threshold for uniform separation by arbitrary records. With an additional total-variation readout allowance ϵ on each binary law, $g_\delta - 2\epsilon > 0$ is a sufficient one-output separation condition. The full-history sharpness claim above concerns the explicitly specified reset nuisance, not every possible history-dependent readout-error class.

With perfect resets, perfect readout, and independent repeated reset trials, the simple hypotheses are Bernoulli$(1/7)$ versus Bernoulli$(1/5)$ for the zero indicator. Lemma 6.7 yields optimal size-$1/20$ randomized powers

n	467	468
power	$0.9498952948087144\ldots$	$0.9505519541817742\ldots$

At $n = 468$, reject for at least 81 zeros and randomize at 80 with probability $0.998625929855047\ldots$, defined exactly by

$$\gamma = \frac{1/20 - \sum_{k=81}^{468} \binom{468}{k}(1/7)^k(6/7)^{468-k}}{\binom{468}{80}(1/7)^{80}(6/7)^{388}}.$$

Discarding an extra iid trial cannot reduce attainable full-record power, so the $467/468$ boundary proves the minimum for this specified randomized fixed-horizon test class. It does not include calibration trials or apply to the two-control reconstruction theorem.

For a different, robust experiment suppose the conditional zero probability after every observed past is at most p_- under one class and at least p_+ under the other, with $g = p_+ - p_- > 0$. Threshold the mean at $(p_- + p_+)/2$. The conditional exponential bound used in Lemma 6.14, applied to the bounded martingale differences of the zero indicators, gives each error at most $\exp(-ng^2/2)$. It suffices that $n \geq \lceil 2\log(20)/g^2 \rceil$. For $g = 2/35$, this is 1835 trials. Independence is unnecessary for this conditional-mean argument, but the interval must hold under the actual filtration at every trial. An unconditional mean gap does not supply it.

6.8 Sampled generators and the information supplied by a clock

A finite continuous-time Markov generator Q has nonnegative off-diagonal entries and zero row sums, and produces transition matrices e^{tQ} for $t \geq 0$. Recovering Q from sampled transitions is the classical embedding and matrix-exponential inverse problem [11, 12]. It presupposes a state basis and time coordinate.

Theorem 6.30 (Two positive generators with all integer samples equal)**.** *Let*

$$C = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \quad \Pi = \tfrac{1}{3}\mathbf{1}\mathbf{1}^T, \quad R = I - \Pi, \quad A = C - C^2,$$

and define

$$Q_0 = -12R, \qquad Q_1 = -12R + \frac{2\pi}{\sqrt{3}}A.$$

Both are irreducible generators with uniform stationary distribution and strictly positive off-diagonal rates. They obey $e^{nQ_0} = e^{nQ_1}$ for every integer $n \geq 0$, but are neither similar nor related by a positive scalar time rescaling, even when a similarity is also allowed. One is reversible and the other has nonzero stationary circulation.

Proof. The off-diagonal rates are 4 for Q_0 and $4 \pm 2\pi/\sqrt{3} > 0$ for Q_1. The inequality follows, for example, from $\pi < 22/7 < 2\sqrt{3}$. Rows and columns sum to zero, so the uniform distribution is stationary; strict positive rates imply irreducibility and uniqueness. Direct algebra gives

$$A\Pi = \Pi A = 0, \quad A^2 = -3R, \quad AR = RA = A.$$

Summing the exponential series on the invariant subspaces yields

$$e^{tQ_0} = \Pi + e^{-12t}R, \qquad e^{tQ_1} = \Pi + e^{-12t}\left[\cos(2\pi t)R + \frac{\sin(2\pi t)}{\sqrt{3}}A\right].$$

At integer t the kernels agree. The spectra are $0, -12, -12$ and $0, -12 \pm 2\pi i$, respectively. Similarity preserves spectra, and a positive scalar cannot turn real nonzero eigenvalues into nonreal ones. The symmetric generator Q_0 is reversible for the uniform law. For Q_1, the oriented stationary flux difference along a cycle edge is $\frac{1}{3}[(4 + 2\pi/\sqrt{3}) - (4 - 2\pi/\sqrt{3})] = 4\pi/(3\sqrt{3}) > 0$. □

Proposition 6.31 (Exactly the integer-grid intervention menu)**.** *Use either generator in Theorem 6.30, a common initial law, and the same instantaneous labelled boundary controls and instruments. Suppose every control and every readout, including silent state maps, occurs on the integer time grid, the between-boundary generator stays fixed, and the policy depends only on the common observed history and private randomness. Then the complete finite controlled observation-history laws agree.*

Proof. A fixed outcome-history probability is a row product of the initial law, transition matrices for nonnegative integer durations, and the corresponding boundary instrument matrices, followed by **1**. Every factor is common to the two descriptions, so the probability is common. For an adaptive randomized policy, condition successively on each history; its next action, scheduling choice, and stop decision have the same conditional law. Induction gives the result. □

Integer-grid readouts alone are insufficient if silent controls are allowed at off-grid times. Persistent drives that change the generator, continuous monitoring, and observations of jump or waiting times also lie outside this menu and may supply distinguishing information.

An anchored preparation of state one and a readout distinguishing states two and three at time $t = 1/4$ give the contrast

$$(e^{Q_1/4})_{12} - (e^{Q_1/4})_{13} = \frac{2e^{-3}}{\sqrt{3}} = 0.0574891546486971\ldots,$$

whereas the contrast is zero for Q_0. An unconditioned stationary uniform preparation has identical uniform marginals under both models and does not provide this witness. The contrast assumes a calibrated quarter-time setting; it is not claimed optimal and carries no trial-count guarantee.

Theorem 6.32 (Exact identification from two incommensurate times)**.** *Let Q, B be finite real matrices represented in the same state basis. Let $t_1, t_2 > 0$ have irrational ratio. If*

$$e^{t_1 Q} = e^{t_1 B}, \qquad e^{t_2 Q} = e^{t_2 B},$$

then $Q = B$.

Proof. Denote the two common exponential matrices by M_1, M_2. They are invertible and commute, since they are both functions of Q. For every pair of integers m, n,

$$e^{(mt_1+nt_2)Q} = M_1^m M_2^n = e^{(mt_1+nt_2)B}.$$

The subgroup $\{mt_1 + nt_2 : m, n \in \mathbb{Z}\}$ is dense in $\mathbb{R}$. One proof is that irrational rotation modulo t_2 has arbitrarily small positive return differences by the pigeonhole principle; integer multiples of those differences approximate any real number, and all such multiples remain in the subgroup. Continuity gives equality of the exponentials for every real time. Differentiating at zero yields $Q = B$. □

The proof does not assume Q and B commute with one another. Negative powers are algebraic inverses, not proposed physical operations at negative time. The result requires full exact transition matrices, a common basis, homogeneous dynamics, and exact times. Finite-precision measurements cannot establish exact irrationality of a time ratio, and this uniqueness theorem is not a stability theorem.

Theorem 6.33 (No uniform inverse stability over unbounded rates)**.** *Fix a nonzero real c and any finite collection of strictly positive observation times. With the matrices R, A above, put*

$$Q_a = -aR, \qquad B_a = -aR + cA, \qquad a > 3|c|.$$

These are irreducible Markov generators. Their difference is the fixed nonzero matrix cA, while $\|e^{tQ_a} - e^{tB_a}\| \to 0$ as $a \to \infty$ at every specified time, for any matrix norm. Thus no uniform inverse error modulus holds over the unrestricted-rate class. In particular, even when the menu is injective, as for two incommensurate times, its inverse is not uniformly continuous.

Proof. Off-diagonal entries are $a/3$ or $a/3 \pm c$, all positive. The same invariant-subspace calculation gives

$$e^{tB_a} - e^{tQ_a} = e^{-at}\left[(\cos(\sqrt{3}ct) - 1)R + \frac{\sin(\sqrt{3}ct)}{\sqrt{3}}A\right].$$

For fixed positive t, the bracket is bounded independently of a and the prefactor tends to zero. A finite maximum over observation times tends to zero as well, while $\|B_a - Q_a\| = |c|\|A\| > 0$. □

On a specified compact generator class, the exact two-time map of Theorem 6.32 is continuous and injective, so its inverse on its image is uniformly continuous by compactness. That qualitative exception provides no quantitative modulus, and no joint finite-data result for uncertain timing and uncertain transition matrices is obtained here.

6.9 Scope of the mathematical conclusions

The positive results identify precisely what their declared experiments determine: autonomous coarse dynamics under a fiber condition; a classical predictive-state lower bound from a joint-law matrix; valid finite-sample or sequential tests under explicit sampling assumptions; finite-state reconstruction under exact anchored controls; and a generator under full exact observations at two incommensurate times. The counterexamples identify specific missing information: short-block agreement, a final channel without fragment access, a single silent control, open calibration uncertainty, or sampling on a coarse time grid can leave distinct descriptions observationally equivalent.

All these conclusions are conditional on conventional mathematical primitives and the stated observation maps. They do not select a physical tensor partition, demonstrate that a detector is independent, derive a probability rule from a prephysical proposal, or provide an emergent physical clock. A physical correspondence must identify which preparations, interventions, outcomes, reset procedures, and timings realize the premises and must confront the appropriate competing observation laws.

References

[1] G. Wolfer and S. Watanabe. *Geometric Aspects of Data-Processing of Markov Chains*. Author manuscript, arXiv:2203.04575v3 (2023). https://arxiv.org/abs/2203.04575v3.

[2] B. Ryabko. Low-Entropy Stochastic Processes for Generating k-Distributed and Normal Sequences, and the Relationship of These Processes with Random Number Generators. *Mathematics* **7**(9), 838 (2019). https://doi.org/10.3390/math7090838.

[3] J. Neyman and E. S. Pearson. On the Problem of the Most Efficient Tests of Statistical Hypotheses. *Philosophical Transactions of the Royal Society of London A* **231**, 289–337 (1933). https://doi.org/10.1098/rsta.1933.0009.

[4] Y. Ohta. *On the Realization of Hidden Markov Models and Tensor Decomposition*. Author manuscript, arXiv:2008.11487 (2020). https://arxiv.org/abs/2008.11487.

[5] B. Vanluyten, J. C. Willems, and B. De Moor. Equivalence of state representations for hidden Markov models. *Systems & Control Letters* **57**, 410–419 (2008). https://doi.org/10.1016/j.sysconle.2007.10.004.

[6] C. Eckart and G. Young. The Approximation of One Matrix by Another of Lower Rank. *Psychometrika* **1**, 211–218 (1936). https://doi.org/10.1007/BF02288367.

[7] C. McDiarmid. On the method of bounded differences. In *Surveys in Combinatorics, 1989*, pp. 148–188. Cambridge University Press (1989). https://doi.org/10.1017/CBO9781107359949.008.

[8] S. R. Howard, A. Ramdas, J. McAuliffe, and J. Sekhon. Time-uniform Chernoff bounds via nonnegative supermartingales. *Probability Surveys* **17**, 257–317 (2020). https://doi.org/10.1214/18-PS321.

[9] M. Larsson, A. Ramdas, and J. Ruf. *The numeraire e-variable and reverse information projection*. Author manuscript, arXiv:2402.18810v2 (2024). https://arxiv.org/abs/2402.18810v2.

[10] D. Crow and R. Joynt. Classical simulation of quantum dephasing and depolarizing noise. *Physical Review A* **89**, 042123 (2014). https://doi.org/10.1103/PhysRevA.89.042123.

[11] M. Baake and J. Sumner. Notes on Markov embedding. *Linear Algebra and its Applications* **594**, 262–299 (2020). https://arxiv.org/abs/1903.08736v2.

[12] Z. Yue, J. Thunberg, and J. Goncalves. *Inverse Problems for Matrix Exponential in System Identification: System Aliasing*. Author manuscript, arXiv:1605.06973 (2016). https://arxiv.org/abs/1605.06973.