

Study of (α, n) reactions in Spain

Ariel Tarifeño-Saldivia

Instituto de Física Corpuscular
C.S.I.C – Universidad de Valencia
Spain
atarisal@ific.uv.es

On behalf of the
MANY collaboration

Scientific motivation

Nuclear data on (alpha,n) reactions are important for:

- **Nuclear astrophysics.** Source of neutrons for the s-process, “light” r-process.
- **Rare-event experiments.** Neutron-induced background in underground experiments (dark matter, neutrinos, neutrinoless double beta decay).
- **Nuclear technologies.** Fission and fusion reactors, spent fuel management and nonproliferation. Neutron-induced background in particle accelerators.

MANY collaboration: Measurement of **Alpha Neutron Yields**

A joint effort with the aim to carry out measurements of (α, xn) reactions using existing facilities in Spain

Detection techniques:


- Direct neutron counting
- Neutron Time-of-Flight Spectroscopy
- Activation
- Gamma spectroscopy

Physical magnitudes:


- Reaction Yields
- Cross sections
- Neutron spectra
- Angular distributions

The MANY Collaboration


Two Spanish facilities




Three Spanish detectors



miniBELEN



MONSTER




GARY


Facilities: Centre for Micro Analysis of Materials (CMAM)

Maximum of $(Z+1) \times 5$ MeV beams, minimum energy below 300 keV with decent ripple.

Terminal voltage was originally calibrated using 11 different nuclear reactions. Recalibration was required after replacement of faulty diodes (2013), about 0.3% deviation found.




HVee tandem
5 MV maximum terminal voltage
Cockroft-Walton acceleration




Courtesy of LM Fraile

The HiSPANoS neutron source @CNA


- HiSPANoS is the first Accelerator-based neutron source in Spain and it is installed at the the 3 MV Tandem Accelerator.
- Operates since:
 - 2013 in continuous mode
 - 2018 in pulsed mode



Courtesy of C Guerrero



MiniBELEN commissioning 45° beamline @ CMAM (Madrid)



UPC
UNIVERSITAT POLITÈCNICA DE CATALUNYA
BARCELONATECH
Institut de Tècniques Energètiques

IFIC
INSTITUT DE FÍSICA
CORPUSCULAR



UNIVERSIDAD
COMPLUTENSE
MADRID



Ciemat
Centro de Investigaciones
Energéticas, Medioambientales
y Tecnológicas

CNA
Centro Nacional de Aceleradores
Infraestructuras
Científicas y
Técnicas
Singulares

UNIVERSIDAD DE
SEVILLA
U

IEM



IFIC
INSTITUT DE FÍSICA
CORPUSCULAR

CSIC
CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS

UNIVERSITAT
DE VALÈNCIA

DDR2 Collaboration Meeting/WP4 Scaling-up
challenges. CERN, 11-13 Feb. 2025

A. Tarifeño-Saldivia

MONSTER

MOdular Neutron time-of-flight Spectrometer is a detection system designed for DESPEC

MONSTER TDR, (2013)


It's the result of an international collaboration between CIEMAT, JYFL-ACCLAB, VECC, IFIC, and UPC

Main characteristics:

- Low neutron energy threshold
- High intrinsic neutron detection efficiency
- Discriminates between detected neutrons and γ -rays by their pulse shape
- Good time resolution
- The energy of the neutrons is determined with the TOF technique

A. R. Garcia *et al.*, JINST, **7**, (2012) C05012



T. Martinez *et al.*, Nuclear Data Sheets, **120**, (2014) 78



Courtesy of A Perez de la Rada



Detection systems: MONSTER



Courtesy of A Perez de la Rada

Thick (300 μm) 27Al (99 % purity) target
 $E\alpha = 5.5, 7, \text{ and } 8.25 \text{ MeV}$
(Buncher not optimized for α -particles)



Uncertainties

Jacobs and Liskien:

- Target stability, charge measurement: 2.0 %
- Neutron detection efficiency: 3.2 - 5.2 %
- Integration procedure: 2.6 %
- Statistics: 2.0 %
- Neutron energy determination:
 - 0.5 % @ 200 keV
 - 1.7 % @ 7 MeV

This work:

- Statistical
- Systematic (only):
 - Efficiency
 - Flight path
 - TOF resolution

Neutron yields

Threshold (keV)	Y_n^{JL}/Y_n
500	1.05
900	1.09
1750	1.01

Detection systems: GARY



Gamma-detector array
for Alpha-induced
Reaction Yield
measurements

LaBr₃(Ce) based array

HPGe detectors

Monitoring neutron detectors

Courtesy of LM Fraile





UNIVERSIDAD
COMPLUTENSE
MADRID

Activation measurements at CMAM

thick-target yields in $^{27}\text{Al}(\alpha, \text{ng})$

Activation measurements



- Using decay of 511 keV when possible
- Using total rate (activation + decay)
- Using activation time

HPGe seems to work fine
 $\text{LaBr}_3(\text{Ce})$ array

Courtesy of LM Fraile



The MANY collaboration:

V. Alcayne⁶, A. Algora², O. Alonso-Sañudo³, J. Balibrea-Correa², J. Benito³, M. J. García-Borge⁵, J. A. Briz³, F. Calviño¹, D. Cano-Ott⁶, G. Cortés¹, A. De Blas¹, C. Domingo-Pardo², A. Espinosa³, B. Fernández⁷, L. M. Fraile³, G. García⁴, R. García¹, V. García-Tavora⁴, J. Gómez-Camacho⁷, E.M. González-Romero⁶, C. Guerrero⁷, A. Illana³, J. Lerendegui-Marco², M. Llanos³, T. Martínez⁶, V. Martínez-Nouvilas³, E. Mendoza⁶, N. Mont-Geli¹, JR. Murias³, E. Nácher², A. Nerio-Aguirre⁵, V. V. O. Onecha³, S. Orrigo², M. Pallàs¹, A. Perea⁵, A. Pérez de Rada⁶, V. Pesudo⁶, J. Plaza⁶, J.M. Quesada⁷, A. Sánchez⁶, V. Sánchez-Tembleque³, R. Santorelli⁶, J.L. Tain², A. Tarifeño-Saldivia², O. Tengblad⁵, J.M. Udías³, D. Villamarín⁶ and S. Viñals⁴

¹ Institut de Tècniques Energètiques (INTE), Universitat Politècnica de Catalunya (UPC), E-08028, Barcelona, Spain

² Instituto de Física Corpuscular (IFIC), CSIC – Univ. Valencia (UV), E-46071, Valencia, Spain

³ Grupo de Física Nuclear (GFN) and IPARCOS, Universidad Complutense de Madrid (UCM), E-28040, Madrid, Spain

⁴ Centro de Micro-Análisis de Materiales (CMAM), Universidad Autónoma de Madrid (UAM), E-28049, Madrid, Spain

⁵ Instituto de Estructura de la Materia (IEM), CSIC, E-28006 Madrid, Spain

⁶ Centro de Investigaciones Energéticas, Medioambientales y Tecnológicas (CIEMAT), E-28040, Madrid, Spain

⁷ Departamento de Física Atómica, Molecular y Nuclear, Universidad de Sevilla (US), E-41012 Sevilla, Spain

⁸ Centro Nacional de Aceleradores (CNA) (Universidad de Sevilla - Junta de Andalucía - CSIC), E-41092, Sevilla, Spain

THANKS!