
(a) coupling disorder (chiral)
std of $\delta b$ (units of $\kappa_2$)
D: 1st order, 0.51σ
T: Monte Carlo
D: Monte Carlo
T: round-off level
(exactly pinned)
disorder strength $\sigma$
(b) on-site disorder
T: 1st order, 0.81σ
D: 1st order, 0.81σ
T: Monte Carlo
D: Monte Carlo
disorder strength $\sigma$
(c) width model ($\sigma_\beta = \sigma_\kappa$)
T: 1st order, 0.81σ
D: 1st order, 1.23σ
T: Monte Carlo
D: Monte Carlo
disorder strength $\sigma$
(d) mode SHAPE change
mean overlap $|\langle\psi_0|\psi\rangle|^2$
T coupling
T onsite
T width
D coupling
D onsite
D width
$\sigma$
(e) H2: T localisation
$\ln|\psi_n/\psi_0|$ (T zero mode)
mean ± 1 std (2000 samples)
$\sigma_\kappa = 0.0$
$\sigma_\kappa = 0.1$
$\sigma_\kappa = 0.3$
$\sigma_\kappa = 0.6$
cells from the centre
(f) H5: width model, unknown ratio
std $\delta b$ at $\sigma_\kappa = 0.05$
T
D
T / D (dotted)
$\sigma_\beta/\sigma_\kappa$ (units of $\kappa_2$)