

QUANTITATIVE APTITUDE FORMULAS



FOR VARIOUS GOVT EXAMS

Quantitative Aptitude section is one of the difficult sections of any Bank and Government Exam. However, with right tricks and practice it can also turn out to be a high scoring section. To help you prepare most effectively for **Quantitative Aptitude**, we are providing you with Free E-books about tips, tricks and formulas for the most important topics from the Quantitative Aptitude syllabus for the exams.

In this E-book we are providing you with all the **Important Quantitative Aptitude formulas** for **Geometry (Mensuration), Simplification and Trigonometry** to help you solve the questions in various Bank and Government Exams.

CRACK SSC CGL EXAM

One - stop solution for complete Tier I & II Prep

Attempt FREE mock test



oliveboard

	Mock Tests	Questions	Marks	Hours
Tier I	GI/ GA/ English/ Quant	100	200	1
Tier II	Paper I - Quantitative Aptitude	100	200	2
	Paper II - English	200	200	2

Tests can be taken in **HINDI & ENGLISH**








Take Your FREE Mock Test Now & Find Out Where You Stand

ATTEMPT NOW








Attempt Test ▶ Analyze Performance ▶ Improve

Visit <https://www.oliveboard.in/ssc-cgl/>







Quantitative Aptitude Formulas: AREA

Figure	Formula of Area
Circle :- 	$= \pi * R^2$ or $= (\pi * D^2) / 4$
Square :- 	$= s \times s$
Rectangle :- 	$= l * w$
Triangle :- 	$= (b * h) / 2$
Parallelogram :- 	$= b \times h$
Rhombus :- 	$= b \times h$
Trapezoid :- 	$= (a + b) / 2 \times h$

Quantitative Aptitude Formulas: Perimeter

Figure	Formula of Perimeter
Circle -: 	$= 2 \times \pi \times r$ or $= \pi \times d$
Square -: 	$= s + s + s + s$ $= 4 \times s$
Rectangle -: 	$= l + l + w + w$ or $= 2 \times l + 2 \times w$
Triangle -: 	$= a + b + c$
Parallelogram -: 	$= a + a + b + b$ or $= 2 \times a + 2 \times b$
Rhombus -: 	$= b + b + b + b$ $= 4 \times b$
Trapezoid -: 	$= a + b + c + d$

Quantitative Aptitude Formulas: Volume

Figure	Formula of Volume
<p>Cube -:</p> 	$= a^3$ $= a \times a \times a$
<p>Cylinder -:</p> 	$= \pi \times r^2 \times h$
<p>Rectangular Solid -:</p> 	$= l \times w \times h$
<p>Sphere -:</p> 	$= (4 \times \pi \times r^3)/3$
<p>Cone -:</p> 	$= (\pi \times r^2 \times h)/3$
<p>Pyramid -:</p> 	$= (B \times h)/3$

Quantitative Aptitude Formulas: Simplification

$$\diamond (a + b)(a - b) = (a^2 - b^2)$$

$$\diamond (a + b)^2 = (a^2 + b^2 + 2ab)$$

$$\diamond (a - b)^2 = (a^2 + b^2 - 2ab)$$

$$\diamond (a - b)^2 = (a^2 + b^2 - 2ab)$$

$$\diamond (a + b + c)^2 = a^2 + b^2 + c^2 + 2(ab + bc + ca)$$

$$\diamond (a^3 + b^3) = (a + b)(a^2 - ab + b^2)$$

$$\diamond (a^3 - b^3) = (a - b)(a^2 + ab + b^2)$$

$$\diamond (a^3 + b^3 + c^3 - 3abc) = (a + b + c)(a^2 + b^2 + c^2 - ab - bc - ac)$$

$$\diamond \text{ If } a + b + c = 0, \\ \text{then } a^3 + b^3 + c^3 = 3abc.$$

BODMAS Rule:

The rule is basically a sequence of operations to follow for solving and finding the correct value of the given arithmetic expression.

Here, BODMAS is an acronym depicting the correct sequence of operations to follow. It stands for -:

B – Bracket

O – Of

D – Division

M – Multiplication

A – Addition

S – Subtraction

Quantitative Aptitude Formulas: Trigonometry

$$\diamond \cos^2 x + \sin^2 x = 1$$

$$\diamond \sin^2 \theta = 1 - \cos^2 \theta$$

$$\diamond \cos^2 \theta = 1 - \sin^2 \theta$$

$$\diamond 1 + \tan^2 x = \sec^2 x$$

$$\diamond 1 + \cot^2 x = \operatorname{cosec}^2 x$$

$$\diamond \sin \theta \times \operatorname{cosec} \theta = 1$$

$$\diamond \cos(x \pm y) = \cos(x) \cdot \cos(y) \pm \sin(x) \cdot \sin(y)$$

$$\diamond \sin(x \pm y) = \sin(x) \cdot \cos(y) \pm \cos(x) \cdot \sin(y)$$

$$\diamond \sin(2x) = 2\sin(x) \cdot \cos(x)$$

$$\diamond \cos(2x) = 2\cos^2(x) - 1$$

$$\diamond \sin(2x) = \cos^2(x) - \sin^2(x)$$

$$\diamond \cos(2x) = 1 - 2\sin^2(x)$$

$$\diamond \tan(x \pm y) = [\tan(x) \pm \tan(y)] / [1 \pm \tan(x) \cdot \tan(y)]$$

$$\diamond \sin(x) \times \sin(y) = 1/2 [\cos(x-y) - \cos(x+y)]$$

$$\diamond \cos(x) \times \cos(y) = 1/2 [\cos(x-y) + \cos(x+y)]$$

$$\diamond \sin(x) \times \cos(y) = 1/2 [\sin(x+y) + \sin(x-y)]$$


$$\diamond \cos(x) \times \sin(y) = 1/2 [\sin(x+y) - \sin(x-y)]$$

Quantitative Aptitude Formulas: Trigonometry

Important Trigonometric Values:

	0	$30^\circ = \frac{\pi}{6}$	$45^\circ = \frac{\pi}{4}$	$60^\circ = \frac{\pi}{3}$	$90^\circ = \frac{\pi}{2}$
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0
tan	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	undefined
csc	undefined	2	$\sqrt{2}$	$\frac{2}{\sqrt{3}}$	1
sec	1	$\frac{2}{\sqrt{3}}$	$\sqrt{2}$	2	undefined
cot	undefined	$\sqrt{3}$	1	$\frac{1}{\sqrt{3}}$	0

Four Quadrant Rule :- All – Sine – Tan – Cosine. (Also known as All Students Take Calculus Rule) to find the values of any trigonometric angle.



FREE Ebooks

[Download Now](#)

Current Affairs

[Explore Now](#)

FREE MOCK TESTS + TOPIC TESTS + SECTIONAL TESTS

For Banking, Insurance, SSC & Railways Exams

[Web](#)

[APP](#)

BLOG

Your one-stop destination
for all exam related
information & preparation
resources.

[Explore Now](#)

FORUM

Interact with peers & experts,
exchange scores
& improve your preparation.

[Explore Now](#)



www.OliveBoard.in