Exercises in Type Theory

Exercise 1

For each of the following types, write down an element of that type, if it has one. If it does not have any elements, you should establish that this is the case. Recall the lecture:

to show that A has no elements, we construct an element of $A \rightarrow \text{empty}$.

- 1) $A \times (B + C) \rightarrow A \times B + A \times C$, given types A, B, C
- 2) $(A \rightarrow A) \rightarrow (A \rightarrow A)$, given type A (for extra credit, write down *five* elements of this type)
- 3) Id_nat (0, succ 0)
- 4) \sum (A: Universe) (A \rightarrow empty) \rightarrow empty
- 5) \prod (n : nat), \sum (m : nat), (n = 2 * m) + (n = 2 * m + 1), assuming you have got arithmetic.
- 6) $(\Sigma (x : A) B \times P x) \rightarrow B \times \Sigma (x : A) P x$, given types A, B, and P : A \rightarrow Universe
- 7) $B \rightarrow (B \rightarrow A) \rightarrow A$, given types A and B
- 8) B $\rightarrow \prod$ (A: Universe) (B \rightarrow A) \rightarrow A, given type B
- 9) (\prod (A : Universe) (B \rightarrow A) \rightarrow A) \rightarrow B, given type B

Exercise 2

- Using the basic rules for natural numbers, construct addition on natural numbers.
- 2) State associativity and commutativity of addition in a type-theoretic way.

Exercise 3

Write down the following types and elements:

- 1) even natural numbers, and 4 as an element of this type
- 2) prime numbers, and 7 as an element of this type
- 3) functions $A \rightarrow$ nat with a given argument at which they are zero (settheoretically this would be $\{(x, f) \in A \times (A \rightarrow nat) \mid f(x) = 0\}$).
- 4) pairs (m, n) of natural numbers such that $m \le n$, and (2,4) as an element of this type.

Exercise 4

We say that a dependent type $P:A \to Universe$ is **decidable** if there is an element of

$$\prod$$
 (x:A), P x + (P x \rightarrow Empty).

Which of the following are decidable?

- 1) P: bool → Universe, defined by P false = unit and P false = empty
- 2) P: nat → Universe, defined by P n = Id_nat (n, 42)