

# Sheaves and their cohomology

Date: 2023-09-01 8:00-10:00 PM

Lecturer: Nithi Rungtanapirom

Transcriber: Kittapat Ratanaphupa

*Recall* (in short)

$$M \otimes_R N = \left\{ \sum_{i=1}^k m_i \otimes n_i : k \in \mathbb{N}, m_i \in M, n_i \in N \right\}$$

subject to the following rules: ...

*Basic properties*

1.  $R \otimes_R M \cong M \cong M \otimes_R R$
2.  $(M \otimes_R N) \otimes_R L \cong M \otimes_R (N \otimes_R L)$
3.  $M \otimes_R N \cong N \otimes_R M$
4.  $(\bigoplus_{i \in I} M_i) \otimes_R N \cong \bigoplus_{i \in I} (M_i \otimes_R N)$

*Remark.*

(2) can be generalized to finitely many modules, i.e,

$$M_1 \otimes_R M_2 \otimes_R \cdots \otimes_R M_n \cong (M_1 \otimes_R (M_2 \otimes_R (\cdots))) \otimes_R M_n \cong M_1 \otimes_R (M_2 \otimes_R ((\cdots) \otimes_R M_n))$$

This also satisfies the universal property with  $R$ -multilinear maps  $M_1 \times \cdots \times M_n \rightarrow N$  instead of  $R$ -bilinear maps.

(4) the tensor product of free module is free.

$$M = R^{(I)} = \bigoplus_{i \in I} R e_i, N = R^{(J)} = \bigoplus_{j \in J} R f_j$$

$$M \otimes_R N \cong \bigoplus_{(i,j) \in I \times J} R(e_i \otimes f_j)$$

*Functoriality*

Let  $M_j, N_j$  be  $R$ -modules,  $\varphi_j : M_j \rightarrow N_j$  for  $j = 1, 2$ . There exists uniquely  $R$ -linear map such that

$$\varphi_1 \otimes \varphi_2 : M_1 \otimes_R M_2 \rightarrow N_1 \otimes_R N_2$$

that is  $(\varphi_1 \otimes \varphi_2)(m_1 \otimes m_2) = \varphi_1(m_1) \otimes \varphi_2(m_2)$  for all  $m_1 \in M_1, m_2 \in M_2$ .

Key step: apply the universal property to the  $R$ -bilinear map

$$M_1 \times M_2 \rightarrow N_1 \otimes_R N_2, (m_1, m_2) \mapsto \varphi_1(m_1) \otimes \varphi_2(m_2).$$

*Extension of scalars* Let  $\rho : R \rightarrow S$  be a ring homomorphism,  $M$  be an  $R$ -module.

$\rho^* M = M \otimes_R S$  is an  $S$ -module via  $t(m \otimes_R s) := m \otimes_R ts$  that  $m \in M, s, t \in S$ .

(The multiplication with  $t \in S$  is  $\text{id}_M \otimes_R (s \mapsto ts)$ )

$S$  is an  $R$ -module via  $r \cdot s := \rho(r)s, r \in R, s \in S$ .

### Examples

- Let  $M = \bigoplus_{i \in I} Re_i$  be a free  $R$ -module. Then,  $M \otimes_R S \cong \bigoplus_{i \in I} Se_i$ .
- Let  $V$  be an  $\mathbb{R}$ -vector space,  $\rho : \mathbb{R} \hookrightarrow \mathbb{C}$ . Then  $V_{\mathbb{C}} := V \otimes_{\mathbb{R}} \mathbb{C}$  “complexification of  $V$ .”  $V_{\mathbb{C}} = \{v + iw \mid v, w \in V\}$  where  $(\alpha + i\beta)(v + iw) = (\alpha v - \beta w) + i(\alpha w + \beta v)$ .

*Remark.* By this way, we obtain a functor  $\rho^* : (R\text{-Mod}) \rightarrow (S\text{-Mod})$

*Exercise* Let  $\rho : R \rightarrow S$  be a ring homomorphism,  $M, N$  be  $R$ -modules. Show that  $(\rho^*M) \otimes_S (\rho^*N) \cong \rho^*(M \otimes_R N)$ .

## Tensor Algebra & co.

Let  $M$  be an  $R$ -module.

### Def

- Let  $r \in \mathbb{N}_0$ . The  $r$ -th tensor power of  $M$  is

$$T_R^r(M) := T^r(M) = M^{\otimes r} = \underbrace{M \otimes_R \cdots \otimes_R M}_{r \text{ times}}$$

As  $T^0(M) = R$  and  $T^1(M) = M$ .

- The tensor algebra of  $M$  is  $T_R(M) := T(M) = \bigoplus_{r \geq 0} T^r(M)$  with the multiplication law obtained from  $T^r(M) \times T^s(M) \rightarrow (T^r(M) \otimes_R T^s(M)) = T^{r+s}(M)$  such that  $(m_1 \otimes \cdots \otimes m_r, n_1 \otimes \cdots \otimes n_s) \mapsto (m_1 \otimes \cdots \otimes m_r \otimes n_1 \otimes \cdots \otimes n_s)$ . This is an associative  $R$ -algebra (not necessary commutative.)

---

The exterior algebra of  $M$  is

$$\Lambda_R(M) := \Lambda(M) := T(M) / (m \otimes m \mid m \in M)$$

where  $(m \otimes m \mid m \in M)$  is two-sided ideal of  $T(M)$ . Then for this case  $[m \otimes n] = -[n \otimes m]$ .

This implies that  $[m_1 \otimes \cdots \otimes m_n] = 0$  if  $\exists i \neq j : m_i = m_j$ .

The multiplication on  $\Lambda(M)$  is typically denoted by  $\wedge$  (wedge product.)

Therefore, the equivalence class of  $m_1 \otimes m_2 \otimes \cdots \otimes m_n$  is  $m_1 \wedge m_2 \wedge \cdots \wedge m_n$ .

For  $r \in \mathbb{N}_0$ , we denoted by  $\Lambda^r(M)$  the submodule of  $\Lambda(M)$  generated by  $m_1 \wedge \cdots \wedge m_r$  “ $r$ -th exterior product.”  $\Lambda(M) = \bigoplus \Lambda^r(M)$ .

- The symmetric algebra of  $M$  is

$$\text{Sym}_R(M) := \text{Sym}(M) := T(M) / (m \otimes n - n \otimes m \mid m, n \in M)$$

Similarly to  $\Lambda(M)$ , we have

$$\text{Sym}(M) = \bigoplus_{r \in \mathbb{N}_0} \text{Sym}^r(M)$$

where  $\text{Sym}^r(M)$  is generated by  $[m_1 \otimes m_2 \otimes \cdots \otimes m_r]$  for  $M_i \in M$ .

*Example* Let  $M := R^n$  (free  $R$ -module with basis  $\{e_1, \dots, e_n\}$ )

- $T^r(M)$  is free with basis  $\{e_{i_1} \otimes e_{i_2} \otimes \cdots \otimes e_{i_r} \mid i_1, \dots, i_r \in \{1, 2, \dots, n\}\}$
- $\Lambda^r(M)$  is free with basis  $\{e_{i_1} \wedge e_{i_2} \wedge \cdots \wedge e_{i_r} \mid i_1, \dots, i_r \in \{1, 2, \dots, n\}\}$  The rank of  $\Lambda^r(M)$  is  $\binom{n}{r}$ .
- $\text{Sym}(M) \cong R[x_1, \dots, x_n]$  as  $R$ -algebra.

---

*Remark*  $\Lambda(M) = T(M) / (m \otimes m \mid m \in M)$

- $\Lambda^r(M)$  satisfies the following universal property

$$\begin{array}{ccc} M \times \cdots \times M & \xrightarrow{\quad\quad\quad} & \Lambda^r M \\ & \searrow & \swarrow \exists! \\ & N & \end{array}$$

$\forall R$ -module  $N, \varphi : M \times \cdots \times M \rightarrow N$  alternating  $r$ -multilinear map.

$\exists! \psi : \Lambda^r(M) \rightarrow N$  that is  $R$ -linear:  $\psi(m_1 \wedge \cdots \wedge m_r) = \varphi(m_1, m_2, \dots, m_r)$  for all  $m_i \in M$ .

- $\Lambda(M)$  is “graded commutative.”  $\forall \omega \in \Lambda^r(M), \eta \in \Lambda^s(M) : \eta \wedge \omega = (-1)^{rs} \omega \wedge \eta_j$ .

(Note:  $(-1)^{rs} = \text{sgn} \left( \begin{cases} i+r & 1 \leq i \leq s \\ i-s & s < i \leq r \end{cases} \right)$ )

**Exercise** Let  $M$  be an  $R$ -module,  $r \in \mathbb{N}$ .

1. Let  $\alpha_1, \dots, \alpha_r \in M^\vee = \text{Hom}_R(M, R)$ . Show that there is an  $R$ -linear map

$$\alpha_1 \wedge \cdots \wedge \alpha_r : \Lambda^r(M) \rightarrow R, m_1 \wedge \cdots \wedge m_r \mapsto \det(\alpha_i(m_j))_{i \times j}$$

2. Derive from (1) the  $R$ -linear map

$$\Phi : \Lambda^r(M^\vee) \rightarrow \Lambda^r(M)^\vee$$

Show that  $\Phi$  is an isomorphism if  $M$  is free and finitely operated.

Back to sheaves of modules:

Let  $(X, \mathcal{O}_X)$  be a ringed space.

*Exercise* Show that if  $\mathcal{F}$  is a locally free  $\mathcal{O}_X$ -module of rank  $n$ , then so is  $\mathcal{F}^\vee := \mathfrak{Hom}_{\mathcal{O}_X}(\mathcal{F}, \mathcal{O}_X)$ . (locally free of rank  $n$ : has open cover  $\{U_i\} : \mathcal{F}_{U_i} \cong (\mathcal{O}_X)|_{U_i}$ )

---

**Def** Let  $\mathcal{F}, \mathcal{G}$  be  $\mathcal{O}_X$ -modules. The tensor product  $\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{G}$  is a sheaf on  $X$  associated to the presheaf

$$(U \subseteq X) \mapsto \mathcal{F}(U) \otimes_{\mathcal{O}_X(U)} \mathcal{G}(U)$$

This is an  $\mathcal{O}_X$ -module.

*Some interesting properties.* Let  $\mathcal{F}, \mathcal{G}$  be  $\mathcal{O}_X$ -modules.

- $\forall x \in X : (\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{G})_a \cong \mathcal{F}_a \otimes_{\mathcal{O}_{X,a}} \mathcal{G}_a$  ( $\mathcal{F}_a$  is an  $\mathcal{O}_{X,a}$ -module via  $[(U, f)][(V, s)] := [(U \cap V, (f|_{U \cap V})(s|_{U \cap V}))]$ )
- If  $\mathcal{F}, \mathcal{G}$  are locally free of rank  $m, n$  respectively, then  $\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{G}$  is locally free of rank  $mn$ .

*Proof.* Exercise. (Wedhorn, Problem 8.7)

*Exercise* (Wedhorn) Problems 8.7-8.8

**Prop** Let  $\mathcal{L}$  be locally free  $\mathcal{O}_X$ -module of rank 1. Then,  $\mathcal{L}^\vee = \mathfrak{H}om_{\mathcal{O}_X}(\mathcal{L}, \mathcal{O}_X)$  is locally free of rank 1 (Exercise: !) and  $\mathcal{L} \otimes \mathcal{L}^\vee \cong \mathcal{O}_X$ .

*Proof.* For  $U \subseteq X$ , consider  $\mathcal{L}(U) \otimes_{\mathcal{O}_X(U)} \mathcal{L}^\vee(U) \rightarrow \mathcal{O}_X(U)$  arising from the  $\mathcal{O}_X(U)$ -bilinear map.

$$\mathcal{L}(U) \times \text{Hom}_{\mathcal{O}_X|_U}(\mathcal{L}|_U, \mathcal{O}_X|_U) \rightarrow \mathcal{O}_X(U), (s, \varphi) \mapsto \varphi_U(s)$$

Sheafifying this yields  $\mathcal{L} \otimes \mathcal{L}^\vee \rightarrow \mathcal{O}_X$ . This is an isomorphism since it is an isomorphism on stalks (Check!!)  $\square$

*Remark* Therefore, the isomorphism classes of locally free  $\mathcal{O}_X$ -modules of rank 1 together with the tensor product form an abelian group. This is called the Picard group and denoted by  $\text{Pic}(X)$ .

*Also a remark.* We can also define the tensor products of finitely many  $\mathcal{O}_X$ -modules in a similar manner.

Let  $M$  be a  $C^\infty$ -manifold.

$(r, s)$ -tensors are sections of  $(\mathcal{T}_M^1)^{\otimes r} \otimes (\mathcal{E}_M^1)^{\otimes s}$ .

$$F_{j_1, \dots, j_n}^{i_1, \dots, i_r} \partial_{i_1} \otimes \dots \otimes \partial_{i_r} \otimes dx^{j_1} \otimes \dots \otimes dx^{j_s}.$$

**Def** Let  $\mathcal{F}$  be an  $\mathcal{O}_X$ -module,  $r \in \mathbb{N}_0$ .

The  $r$ -th exterior power of  $\mathcal{F}$  is the sheaf associated to  $U \mapsto \Lambda_{\mathcal{O}_X(U)}^r(\mathcal{F}(U))$ . We call them  $\Lambda^r \mathcal{F}$ .

The  $r$ -symmetric power of  $\mathcal{F}$  is the sheaf associated to  $U \mapsto \text{Sym}_{\mathcal{O}_X(U)}^r(\mathcal{F}(U))$ . We call them  $\text{Sym}^r(\mathcal{F})$ .

*Example.* Let  $M$  be a  $C^\infty$ -manifold. Then,  $\mathcal{E}_M^p := \Lambda^p(\mathcal{E}_M^1)$  ( $p$ -th differential forms:  $dx^{i_1} \wedge \dots \wedge dx^{i_p}$ .)

*Remark.* If  $\mathcal{F}$  is a locally free  $\mathcal{O}_X$ -module of rank  $n$ , then  $\Lambda^r \mathcal{F}$  is locally free of rank  $\binom{n}{r}$ . If  $r = n$ ,  $\det(\mathcal{F}) := \Lambda^n \mathcal{F}$  is locally free of rank 1 (the determinant of  $\mathcal{F}$ )

( $T : V \rightarrow V$  of dimension  $n$ ,  $\dim V = n$ :  $\Lambda^n T : \Lambda^n V \rightarrow \Lambda^n V$  is given by multiplication with  $\det T$ )

Furthermore, there is an isomorphism (!)

$$\Lambda^r(\mathcal{F}^\vee) \xrightarrow{\sim} \Lambda^r(\mathcal{F})^\vee$$

---

Now let  $f : (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$  be a morphism of ringed spaces.

**Def**

- Let  $\mathcal{F}$  be an  $\mathcal{O}_X$ -module, then  $i_*\mathcal{F}$  is an  $\mathcal{O}_Y$ -module (a pushforward.) For  $V \subseteq Y$  :  $i_*(V) = \mathcal{F}(f^{-1}(V))$  is an  $\mathcal{O}_X(f^{-1}(V))$ -module hence also an  $\mathcal{O}_Y(V)$ -module via  $f_V^\flat : \mathcal{O}_Y(V) \rightarrow \mathcal{O}_X(f^{-1}(V))$ .
- Let  $\mathcal{G}$  be an  $\mathcal{O}_Y$ -module.  $f^{-1}\mathcal{G} :=$  a sheafification of  $\left( U \mapsto \varinjlim_{f(U) \subseteq V \subseteq Y} \mathcal{G}(V) \right)$  is an  $f^{-1}\mathcal{O}_Y$ -module.  $f^\sharp : f^{-1}\mathcal{O}_Y \rightarrow \mathcal{O}_X$ . Then  $f^*\mathcal{G} := f^{-1}\mathcal{G} \otimes_{f^{-1}\mathcal{O}_Y} \mathcal{O}_X$ . This is an  $\mathcal{O}_X$ -module. (a pullback)

*Remark* for  $x \in X$ :

- $(f^*(\mathcal{G}))_x \cong \mathcal{G}_{f(x)} \otimes_{\mathcal{O}_{Y,f(x)}} \mathcal{O}_{X,x}$
- If  $\mathcal{G}$  is locally free of rank  $n$ , so is  $f^*\mathcal{G}$ .

*Exercise.* Show that

$$f_* : (\mathcal{O}_X\text{-Mod}) \rightarrow (\mathcal{O}_Y\text{-Mod})$$

is a right adjoint of  $f^* : (\mathcal{O}_Y\text{-Mod}) \rightarrow (\mathcal{O}_X\text{-Mod})$ , i.e.,  $\text{Hom}_{\mathcal{O}_X}(f^*\mathcal{G}, \mathcal{F}) \cong \text{Hom}_{\mathcal{O}_Y}(\mathcal{G}, f_*\mathcal{F})$  for all ...