

## Q1

- (a) .
- (b) .
- (c)

$$\begin{aligned}0 &\leq x \leq 1 \\0 &\leq y \leq 1 \\x^2 &\leq z \leq x \iff \sqrt{x^2} \leq \sqrt{z} \wedge z \leq x\end{aligned}$$

Thus we can write the bounds as:

$$\begin{aligned}0 &\leq z \leq 1 \\0 &\leq y \leq 1 \\z &\leq x \leq \sqrt{z}\end{aligned}$$

As our triple integral is "nice", we can rewrite it with these new bounds:

$$\int_0^1 \int_0^1 \int_z^{\sqrt{z}} f(x, y, z) dx dy dz$$

## Q2

- (a) First we find integrals for the surface area of each component body:

- The box has 6 sides, each with area:

$$\int_0^4 \int_0^4 1 du dv$$

- For the tube, we parametrize the surface as:

$$r(u, v) = \begin{bmatrix} e^u \cos v \\ e^u \sin v \\ u \end{bmatrix}$$

Where  $u \in [0, 1]$  and  $v \in [0, 2\pi]$ . Giving a surface integral:

$$\begin{aligned}
 & \int_0^{2\pi} \int_0^1 \left\| \frac{\partial r(u, v)}{\partial u} \times \frac{\partial r(u, v)}{\partial v} \right\| du dv \\
 &= \int_0^{2\pi} \int_0^1 \left\| \begin{bmatrix} e^u \cos v \\ e^u \sin v \\ 1 \end{bmatrix} \times \begin{bmatrix} -e^u \sin v \\ e^u \cos v \\ 0 \end{bmatrix} \right\| du dv \\
 &= \int_0^{2\pi} \int_0^1 \left\| \begin{bmatrix} -e^u \cos v \\ -e^u \sin v \\ e^{2u} \cos^2 v + e^{2u} \sin^2 v \end{bmatrix} \right\| du dv \\
 &= \int_0^{2\pi} \int_0^1 \sqrt{e^{2u} + e^{4u}} du dv \\
 &= \int_0^{2\pi} \int_0^1 e^u \sqrt{1 + e^{2u}} du dv
 \end{aligned}$$

- The top of the box must have a hole for the tube, therefore we must remove an area correspond to the bottom of the tube, a disk with radius  $e^0 = 1$ :

$$- \int_0^1 \int_0^{2\pi} r d\theta dr$$

- The bottom of the half sphere connects the top of the tube to the sphere, thus it is a ring with inner radius  $e^1$  and outer radius 4. Again using polar coordinates:

$$\int_e^4 \int_0^{2\pi} r d\theta dr$$

- We parametrize the surface of the sphere in spherical coordinates,  $[\rho, \theta, \phi]^T$  for  $\rho = 4$ ,  $\theta \in [0, 2\pi]$ ,  $\phi \in [0, \frac{\pi}{2}]$ , the area element of the sphere is given by  $\rho^2 \sin \phi d\theta d\phi$ . Hence we have an integral for the surface area of:

$$\int_0^{\frac{\pi}{2}} \int_0^{2\pi} 4^2 \sin \phi d\theta d\phi$$

Thus the surface area of  $R$  is given by:

$$A = 6 \int_0^4 \int_0^4 1 du dv + \int_0^{2\pi} \int_0^1 e^u \sqrt{1 + e^{2u}} du dv - \int_0^1 \int_0^{2\pi} r d\theta dr + \int_e^4 \int_0^{2\pi} r d\theta dr + \int_0^{\frac{\pi}{2}} \int_0^{2\pi} 4^2 \sin \phi d\theta d\phi$$

(b) We find three separate integrals for the mass of each region of  $R$ :

- For the box we simply integrate  $\rho$  over its bound in Cartesian coordinates:

$$\int_{-4}^0 \int_{-2}^2 \int_{-2}^2 \rho(x, y, z) dx dy dz$$

- For the tube, we integrate in cylindrical coordinates where the volume element is given by  $dV = r dr d\theta dz$ , giving an integral of:

$$\int_0^1 \int_0^{2\pi} \int_0^{e^z} \rho(r \cos \theta, r \sin \theta, z) r dr d\theta dz$$

- Lastly for the sphere, we integrate in Cartesian coordinates:

$$\int_{-4}^4 \int_{-\sqrt{4^2-x^2}}^{\sqrt{4^2-x^2}} \int_1^{1+\sqrt{4^2-x^2-y^2}} \rho(x, y, z) dz dy dx$$

Thus we have a total mass of:

$$\begin{aligned}
 m &= \int_{-4}^0 \int_{-2}^2 \int_{-2}^2 \rho(x, y, z) dx dy dz \\
 &+ \int_0^1 \int_0^{2\pi} \int_0^{e^z} \rho(r \cos \theta, r \sin \theta, z) r dr d\theta dz \\
 &+ \int_{-4}^4 \int_{-\sqrt{4^2-x^2}}^{\sqrt{4^2-x^2}} \int_1^{1+\sqrt{4^2-x^2-y^2}} \rho(x, y, z) dz dy dx
 \end{aligned}$$

### Q3

- (a) Looking at the gradient field, we see that a factor of  $2x$  and  $2y$  indicating the inside was differentiated using the chain rule, and a reciprocal indicating that the outer function could have been  $\ln z$ , with derivative  $z^{-1}$ . Thus, as a guess we consider  $\phi(x, y) = \log(x^2 + y^2)$  defined on  $\mathbb{R}^2 \setminus \{0\}$ , which is continuous and differentiable over its domain. Checking its gradient:

$$\nabla \phi(x, y) = \begin{bmatrix} \frac{\partial \phi(x, y)}{\partial x} \\ \frac{\partial \phi(x, y)}{\partial y} \end{bmatrix} = \frac{2}{x^2 + y^2} \begin{bmatrix} x \\ y \end{bmatrix}$$

As  $\mathbb{R}^2 \setminus \{0\} \subset \{(x, y) | y = \alpha x \alpha \in \mathbb{R}\}$ ,  $\phi$  is defined here and one has  $\nabla \phi = F$ . Hence,  $\phi$  is a potential which satisfies the question.

- (b) In (a), we have already found a potential that satisfies the question, thus it exists.