

Model-derived PK target threshold profile of ustekinumab therapeutic thresholds in inflammatory bowel diseases to guide precision dosing

E.S.K. Widigson (1,2), C. Steenholdt (3), C. Frimor (4), M.A. Ainsworth (4), W. Huisinga (2,5), C. Kloft (1,2)

(1) Dept. of Clinical Pharmacy and Biochemistry, Institute of Pharmacy, Freie Universitaet Berlin, Germany, (2) Graduate Research Training program PharMetriX, Germany, (3) Dept. of Gastroenterology, Herlev and Gentofte Hospital, Denmark, (4) Dept. of Gastroenterology, Odense University Hospital, Denmark, (5) Institute of Mathematics, Universität Potsdam, Germany



Background and Objectives

Inflammatory bowel diseases (IBD) are a group of chronic immune-mediated diseases affecting the gastrointestinal tract, causing debilitating symptoms such as persistent abdominal pain, diarrhoea, and fatigue and with negative influence on quality of life [1].

Ustekinumab (UST) is a mAb drug used to treat IBD. Since for UST an exposure-response relationship has been demonstrated, the lack of response in some patients might be due to suboptimal drug concentrations. Subsequently, TDM has been proposed [2].

TDM is performed by measuring the drug serum concentration (typically trough levels) at a specific timepoint and comparing to a pre-defined PK concentration threshold linking drug concentration to beneficial PD treatment outcome. Given the long dosing intervals of mAb drugs, this means that there is usually a **time lag of eight to twelve weeks** between TDM measurements and subsequent individual treatment optimisation.

This work aimed to provide the framework for a more **rapid UST treatment individualisation**, leveraging the knowledge from nonlinear mixed-effects (NLME) modelling to account for and leverage PK/PD targets from several studies at different timepoints simultaneously.

Methods

A two-compartment **NLME model** with 1st-order absorption and elimination was identified from literature [3].




Table 1. PK/PD targets of UST concentration associated with endoscopic remission in IBD.

Week	Concentration [mg/L]	N patients	Reference
4	23.3	41	[5]
8	11.1	41	[5]
8	3.47	960	[4]
16	1.12	19	[6]
24	3.75	337	[7]
24	6.00	135	[8]
24	2.00	28	[9]
24	2.11	108	[10]
Maintenance (allocated: 24)	2.30	71	[11]
Maintenance (allocated: 24)	1.08	960	[4]

PK/PD targets for reaching endoscopic remission with UST treatment for IBD were collected from literature (Table 1) [4-11].

Threshold concentrations was **weighted** based on number of patients involved in deriving them.

Maximum a posteriori (MAP) parameter value for CL was estimated fitting the PK/PD targets using the published model, for deriving the PK threshold profile.

Results cont.

Therapeutic threshold profile of UST PK thresholds




Figure 1. Simulated target concentration-time threshold profile based on 10 therapeutic thresholds plotted together with the thresholds and stochastic simulations using model published by O.J. Adedokun et al. (2022) [3, 4-11].

Results

- Using the identified **NLME model** together with the **10 PK/PD targets** for reaching endoscopic remission, the **MAP** parameter value for CL was estimated to **0.182 L/day** (Table 2), for an UST clinical praxis dosing regimen of an intravenous induction loading dose of 390 mg followed by 90 mg subcutaneously every 8 weeks.
- From the **simulated UST concentration-time profile** (Figure 1), a **model-informed concentration threshold table** was generated for induction weeks 1-16 and maintenance therapy weeks 1-8 after dosing, increasing availability of the PK target profile for physicians without modelling experience (Table 3).

Table 2. Model parameters for UST therapeutic target profile

Parameter [unit]	Value
Ka [d ⁻¹]	0.181
F [%]	78.2
CL _{MAP} threshold profile [L/d]	0.182
Vc [L]	2.75
Vp [L]	1.88
Q [L/day]	0.287

Table 3. Model-informed therapeutic thresholds for the clinical praxis dosing regimen of 390 mg intravenous loading dose followed by 90 mg subcutaneously every 8 weeks.

Induction therapy		Maintenance (q8w)	
Week	Target concentration [mg/L]	Week	Target concentration [mg/L]
2	37.2	1	10.8
4	21.2	2	9.56
6	12.5	3	7.58
8 (pre-dose)	7.37	4	5.87
10	12.7	5	4.52
12	7.70	6	3.47
14	4.55	7	2.67
16 (pre-dose)	2.69	8 (pre-dose)	2.06

Conclusions

This work leverages the knowledge from NLME modelling to **account for** and **average** PK/PD targets from several studies at **different timepoints**. A complete concentration-time profile for induction and maintenance dosing every 8 weeks was derived from 10 published PK/PD targets and a published NLME model.

A framework is provided for more **rapid treatment individualisation** by **enabling TDM sampling** of UST treatment against IBD **at any point** during the dosing interval. By using the model-based threshold time profile, **dosing optimisation** to secure drug levels associated with endoscopic remission is facilitated.

[1] L.P. Bruner *et al.* Inflamm. Bowel Dis. (2023).
[2] S. Restellini, W. Afif. J Clin Med. (2021).
[3] O.J. Adedokun *et al.* Clin Ther. (2022).
[4] O.J. Adedokun *et al.* Clin. Gastroenterol. Hepatol. (2020).
[5] J. Hanžel *et al.* Clin. Gastroenterol. Hepatol. (2021).

[6] J.-Y. Yao *et al.* BMC Gastroenterol. (2021).
[7] M. Walshe *et al.* Crohns Colitis. (2021).
[8] J. Hanžel *et al.* Eur. J. Gastroenterol. Hepatol. (2021).
[9] H. Hirayama *et al.* BMC Gastroenterol (2022).
[10] J. Yao *et al.* J. Clin. Med. (2023).
[11] J. Petrov *et al.* J Clin Gastroenterol (2024).

IBD: Inflammatory bowel diseases
mAb: Monoclonal antibody
MAP: Maximum a posteriori
NLME: Nonlinear mixed-effects
PK/PD: Pharmacokinetic/pharmacodynamic
TDM: Therapeutic Drug Monitoring
UST: Ustekinumab



For more information:
Ella Signe Kassandra Widigson,
ella.widigson@fu-berlin.de
www.clinical-pharmacy.eu

{Population Approach Group Europe – PAGE}, {Rome 2024}

