

# Quantifying the impact of CYP2D6 allele activity on Z-endoxifen formation leveraging the multi-study CEPAM database: Towards treatment optimisation of tamoxifen

F. Klima (1,2), T. Helland (3,4,5), R. Michelet (1), W. Huisinga (6), D. Hertz (3), C. Kloft (1,2)  
for the CYP2D6 Endoxifen Percentage Activity Model in Breast Cancer (CEPAM) consortium

(1) Dept. of Clinical Pharmacy and Biochemistry, Institute of Pharmacy, Freie Universitaet Berlin, Germany, (2) Graduate Research Training program PharMetRX, Germany, (3) Dept. of Clinical Pharmacy, University of Michigan College of Pharmacy, Ann Arbor, USA, (4) Hormone Laboratory, Dept. of Medical Biochemistry and Pharmacology, Haukeland University Hospital, Bergen, Norway, (5) Dept. of Clinical Science, University of Bergen, Norway, (6) Institute of Mathematics, University of Potsdam, Germany




## Background and Objectives


### Tamoxifen treatment individualisation?

**Prodrug TAM:** Breast cancer recurrence ( $\leq 25\%$ ) at 20 mg QD standard dosing<sup>1</sup>  
**Active Z-ENDX:** Clinical studies found heterogeneous PK efficacy thresholds<sup>2-5</sup>  
**Large prospective study for TAM treatment individualisation unrealistic<sup>6</sup>**

**New approach needed: Modelling + simulation of multi-study data<sup>7,8</sup>**



### Role of varying CYP2D6 allele activities



**Develop PK model for TAM and Z-ENDX from large multi-study dataset**

**Investigate quantitative impact of CYP2D6 alleles on Z-ENDX formation**

## Methods

### CEPAM analysis dataset

36 studies n=8451 n=10574  
TAM <1 month  
No PK measurements  
19.4% Unexplained outliers  
No CYP2D6 genotype  
31 studies n=6841 n=8791

### PK samples


94% one sample  
6% multiple samples [2-20]  
**Z-ENDX 66.5%**  
**TAM + Z-ENDX and Z4'-ENDX 19.8%**  
**Total ENDX 13.8%** (Z-ENDX + Z4'-ENDX)

### Developing CYP2D6 percentage allele activity



**Modelling CYP2D6 activity**  
NONMEM<sup>®</sup> 7.5.1, FOCE-I  
Single CYP2D6 allele activity:  
 $\text{Activity} = \text{Activity} \cdot N \cdot \text{CNV}$   
Combined allele activity:  
 $\text{Activity} = \frac{\text{Activity}_1 + \text{Activity}_2}{2}$   
CYP2D6-dependent Z-ENDX formation:  
 $CL_{\text{CYP2D6}} = CL_{\text{CYP2D6}} \cdot \text{Activity} \cdot e^{\text{CYP2D6 inhibitor}}$

CYP2D6 alleles CEPAM,%		
xN	CNV	2.1
*1 (WT)	44	
*2	13	
*35	1.8	
*9	2.2	
*10	7.3	
*17	1.2	
*41	7.6	
*3	1.0	
*4	15	
*5	3.8	

## Results




**Figure 1** Parent-metabolite model of TAM and ENDX. All exponential and power covariate relationships were median-normalised. All RSEs:  $\leq 30\%$ . \*: Parameter values fixed.  
**Structural parameters:**  $t_{\text{lag}}$ : 0.385 h;  $k_a$ : 1.09 h<sup>-1</sup>;  $CL_{\text{TAM}}/F$ : 5.78 L/h; CYP2D6-dependent apparent Z-ENDX formation  $CL_{\text{CYP2D6}}/F$ : 0.442 L/h; CYP2D6-independent apparent Z-ENDX formation  $CL_{\text{other}}/F$ : 0.163 L/h;  $CL_{\text{ZENDX}}/F$ : 5.1 L/h;  $CL_{\text{Z4'MET}}/F$ : 0.318 L/h;  $CL_{\text{Z4'ENDX}}/F$ : 5.1 L/h;  $V_{\text{TAM,C}}/F$ : 743 L;  $V_{\text{ZENDX,C}}/F$ : 400 L;  $V_{\text{Z4'ENDX,C}}/F$ : 400 L.  
**Covariates on F:** Non-adherence (-0.697) as relative change; on  $CL_{\text{TAM}}/F$ : Age (-0.282) and body weight (0.177) as power functions; CYP3A4 and CYP2C19 phenotype (0.133 and 0.001) as exponential models; on  $CL_{\text{CYP2D6}}/F$ : CYP2D6 inhibitor (-0.565) as exponential model; CYP2D6 alleles as relative change, Fig. 4; on  $CL_{\text{other}}/F$ : CYP2C9 phenotype (0.380) as exponential model; on  $CL_{\text{ZENDX}}/F$  and  $CL_{\text{Z4'ENDX}}/F$ : Body weight (0.385 and 0.226) as power functions.  
**IIV:** On  $CL_{\text{TAM}}/F$  (51.0% CV) and on Z-ENDX formation  $CL_{\text{CYP2D6}}/F + CL_{\text{other}}/F$  (28.1% CV) as exponential model. **RUV:** Proportional model separated for patients with one sample (29.9-58.6% CV) and multiple samples ( $\leq 17.3\%$  CV).




CYP2D6 allele activity		
Alleles	Activity	
Category Percentage		
xN	CNV	Nx 0.70
*1 (WT)	1	1 fix
*35	1	0.74
*2	1	0.72
*9	0.25	0.58
*17	0.5	0.30
*10	0.25	0.25
*41	0.25	0.17
*3,*4,*5	0	0 fix

**Figure 4** CYP2D6 categorical and percentage allele activities and impact of gene duplication (CNV).


## Conclusions



**PK model was successfully developed:**



**Application of PK model to studies with survival data:**



### References

- [1] Davies et al. Lancet 381: 805-816 (2013).
- [2] Madlensky et al. Clin. Pharmacol. Ther. 89: 718-725 (2011).
- [3] Saladores et al. Pharmacogenomics J. 15: 84-94 (2015).
- [4] Helland et al. Breast Cancer Res. 19 (2011).
- [5] Love et al. Springerplus 2: 1-5 (2013).
- [6] Helland et al. J. Pers. Med. 11 (2021).
- [7] de Vries Schultink et al. PAGE 28 (2019).
- [8] Mc Laughlin et al. Clin. Pharmacol. Ther. (2024).
- [9] Caudle et al. Clin. Transl. Sci. 13: 116-124 (2020).

### Abbreviations

- |        |   |                  |                         |
|--------|---|------------------|-------------------------|
| CNV    | Copy number variation                               | PK               | Pharmacokinetic(s)      |
| CL     | Clearance   | QD               | Once daily              |
| ENDX   | Endoxifen   | RSE              | Relative standard error |
| F      | Bioavailability                                     | RUV              | Residual standard error |
| FOCE-I | First-order conditional estimation with interaction | TAM              | Tamoxifen               |
| IIV    | Interindividual variability                         | $t_{\text{lag}}$ | Absorption lag time     |
| $k_a$  | Absorption rate constant                            | V                | Volume of distribution  |
| N      | Number of gene duplications                         | WT               | wildtype                |



For more information:  
Fenja Klima,  
[fenja.klima@fu-berlin.de](mailto:fenja.klima@fu-berlin.de)  
[www.clinical-pharmacy.eu](http://www.clinical-pharmacy.eu)

32<sup>nd</sup> Population Approach Group Europe meeting – PAGE, Rome, Italy, 2024

