

X-2 ION-MOTION EXPERIMENT (ARD Nov 12–15)

Carl A. Lindstrøm & Severin Diederichs

FLASHFORWARD ▶ Research Fellow | Research Group for Plasma Wakefield Accelerators
Deutsches Elektronen-Synchrotron DESY, Particle Physics Division, Hamburg, Germany




Accelerator Research and Development, Matter and Technologies
Helmholtz Association of German Research Centres, Berlin, Germany



HELMHOLTZ
RESEARCH FOR GRAND CHALLENGES

PLANNED EXPERIMENT FOR THE ARD SHIFT Nov 12–15

- > What: **Observe emittance growth due to ion motion inside the plasma cavity.**
- > Why?
 - > An on-axis ion spike gives nonlinear focusing fields, resulting in emittance growth.
- > How?
 - > Using a tightly focused beam (semi-matched)
 - > Hydrogen plasma
 - > Argon plasma (as a control; expecting no ion motion)
 - > Observable: emittance vs. energy gain (a proxy for the z-location)
 - > *(object plane scan for different energies on a spectrometer)*
- > Big question: Can it be done at FLASHForward?
 - > Using Rosenzweig's ion-phase-advance formula: maybe...
 - > Severin performed HiPACE++ simulations to check.



$$\Delta\phi_i \simeq \sqrt{\frac{2\pi r_p Z_i N \sigma_z}{A \epsilon_n}} (r_e n_0 \gamma)^{1/4}$$

SIMULATIONS USING DIFFERENT BEAM SIZES (SIMULATED BY SEVERIN USING HIPACE++)

> The answer is yes, if the beam size is small enough.

> *Matched: 2 μm rms*

> Simulation:

> 1 GeV, 1 nC charge

> ~1 kA peak current


> $7 \times 10^{15} \text{ cm}^{-3}$, 40 mm long

> 2 mm mrad emittance


> 0.4% energy spread

> Shown:


> Slice emittance at the tail




3 μm rms




5 μm rms




8 μm rms




10 μm rms



20 μm rms




CROSS SECTION OF THE PLASMA CAVITY



GROWING EMITTANCE TOWARDS THE BEAM TAIL



> Main observable:



NON-GAUSSIAN TRANSVERSE PHASE SPACE IN HYDROGEN DUE TO ION MOTION

- > Looking at a slice towards the tail, we see clear evidence of nonlinear focusing:
 - > A galaxy-shaped phase space (hydrogen) shows that focusing is stronger closer to the axis (like in a galaxy).
 - > In argon, a regular bivariate Gaussian distribution is observed.

Simulations with a 5 μm rms beam size:



> In conclusion—we should be able to observe emittance growth due to ion motion!