

YSU ASDS, Statistics, Fall 2019

Lecture 26

Michael Poghosyan

25 Nov 2019

Contents

- ▶ Tests for the Normal Model Variance
- ▶ Large Sample Hypothesis Testing

Last Lecture ReCap

- ▶ What are we testing when using a (one-sample) Z - or t -Test?

Last Lecture ReCap

- ▶ What are we testing when using a (one-sample) Z - or t -Test?
- ▶ Describe the Z -test.

Last Lecture ReCap

- ▶ What are we testing when using a (one-sample) Z - or t -Test?
- ▶ Describe the Z -test.
- ▶ Describe the T -test.

Last Lecture ReCap

- ▶ What are we testing when using a (one-sample) Z - or t -Test?
- ▶ Describe the Z -test.
- ▶ Describe the T -test.
- ▶ Describe the relation between Testing and CIs.

Test for the Normal Variance, μ is known

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is known**, the Parameter (our unknown) is σ^2 ;

Test for the Normal Variance, μ is known

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is known**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Test for the Normal Variance, μ is known

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is known**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test for the Normal Variance, μ is known

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is known**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics:

Test for the Normal Variance, μ is known

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is known**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 =$

Test for the Normal Variance, μ is known

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is known**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \mu}{\sigma_0} \right)^2$

Test for the Normal Variance, μ is known

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is known**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \mu}{\sigma_0} \right)^2$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $\chi^2 \sim$

Test for the Normal Variance, μ is known

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is known**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \mu}{\sigma_0} \right)^2$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $\chi^2 \sim \chi^2(n)$;

Test for the Normal Variance, μ is known

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ is known, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \mu}{\sigma_0} \right)^2$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $\chi^2 \sim \chi^2(n)$;

\mathcal{H}_1 is	RR is
$\sigma^2 \neq \sigma_0^2$	

Test for the Normal Variance, μ is known

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is known**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \mu}{\sigma_0} \right)^2$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $\chi^2 \sim \chi^2(n)$;

\mathcal{H}_1 is	RR is
$\sigma^2 \neq \sigma_0^2$	$\chi^2 \notin \left[\chi_{n, \frac{\alpha}{2}}^2, \chi_{n, 1 - \frac{\alpha}{2}}^2 \right]$

Test for the Normal Variance, μ is known

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is known**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \mu}{\sigma_0} \right)^2$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $\chi^2 \sim \chi^2(n)$;

\mathcal{H}_1 is	RR is
$\sigma^2 \neq \sigma_0^2$ $\sigma^2 > \sigma_0^2$	$\chi^2 \notin \left[\chi_{n, \frac{\alpha}{2}}^2, \chi_{n, 1 - \frac{\alpha}{2}}^2 \right]$

Test for the Normal Variance, μ is known

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is known**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \mu}{\sigma_0} \right)^2$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $\chi^2 \sim \chi^2(n)$;

\mathcal{H}_1 is	RR is
$\sigma^2 \neq \sigma_0^2$	$\chi^2 \notin \left[\chi_{n, \frac{\alpha}{2}}^2, \chi_{n, 1 - \frac{\alpha}{2}}^2 \right]$
$\sigma^2 > \sigma_0^2$	$\chi^2 > \chi_{n, 1 - \alpha}^2$

Test for the Normal Variance, μ is known

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is known**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \mu}{\sigma_0} \right)^2$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $\chi^2 \sim \chi^2(n)$;

\mathcal{H}_1 is	RR is
$\sigma^2 \neq \sigma_0^2$	$\chi^2 \notin \left[\chi_{n, \frac{\alpha}{2}}^2, \chi_{n, 1 - \frac{\alpha}{2}}^2 \right]$
$\sigma^2 > \sigma_0^2$	$\chi^2 > \chi_{n, 1 - \alpha}^2$
$\sigma^2 < \sigma_0^2$	

Test for the Normal Variance, μ is known

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is known**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \mu}{\sigma_0} \right)^2$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $\chi^2 \sim \chi^2(n)$;

\mathcal{H}_1 is	RR is
$\sigma^2 \neq \sigma_0^2$	$\chi^2 \notin \left[\chi_{n, \frac{\alpha}{2}}^2, \chi_{n, 1 - \frac{\alpha}{2}}^2 \right]$
$\sigma^2 > \sigma_0^2$	$\chi^2 > \chi_{n, 1 - \alpha}^2$
$\sigma^2 < \sigma_0^2$	$\chi^2 < \chi_{n, \alpha}^2$

Test for the Normal Variance, μ is unknown

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is unknown**, the Parameter (our unknown) is σ^2 ;

Test for the Normal Variance, μ is unknown

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is unknown**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Test for the Normal Variance, μ is unknown

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is unknown**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test for the Normal Variance, μ is unknown

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is unknown**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics:

Test for the Normal Variance, μ is unknown

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is unknown**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 =$

Test for the Normal Variance, μ is unknown

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is unknown**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \bar{X}}{\sigma_0} \right)^2 = \frac{(n-1) \cdot S^2}{\sigma_0^2}$

Test for the Normal Variance, μ is unknown

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is unknown**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \bar{X}}{\sigma_0} \right)^2 = \frac{(n-1) \cdot S^2}{\sigma_0^2}$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $\chi^2 \sim$

Test for the Normal Variance, μ is unknown

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is unknown**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \bar{X}}{\sigma_0} \right)^2 = \frac{(n-1) \cdot S^2}{\sigma_0^2}$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $\chi^2 \sim \chi^2(n-1)$;

Test for the Normal Variance, μ is unknown

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ is unknown, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \bar{X}}{\sigma_0} \right)^2 = \frac{(n-1) \cdot S^2}{\sigma_0^2}$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $\chi^2 \sim \chi^2(n-1)$;

\mathcal{H}_1 is	RR is
$\sigma^2 \neq \sigma_0^2$	

Test for the Normal Variance, μ is unknown

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is unknown**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \bar{X}}{\sigma_0} \right)^2 = \frac{(n-1) \cdot S^2}{\sigma_0^2}$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $\chi^2 \sim \chi^2(n-1)$;

\mathcal{H}_1 is	RR is
$\sigma^2 \neq \sigma_0^2$	$\chi^2 \notin \left[\chi_{n-1, \frac{\alpha}{2}}^2, \chi_{n-1, 1-\frac{\alpha}{2}}^2 \right]$

Test for the Normal Variance, μ is unknown

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is unknown**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \bar{X}}{\sigma_0} \right)^2 = \frac{(n-1) \cdot S^2}{\sigma_0^2}$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $\chi^2 \sim \chi^2(n-1)$;

\mathcal{H}_1 is	RR is
$\sigma^2 \neq \sigma_0^2$	$\chi^2 \notin \left[\chi_{n-1, \frac{\alpha}{2}}^2, \chi_{n-1, 1-\frac{\alpha}{2}}^2 \right]$
$\sigma^2 > \sigma_0^2$	

Test for the Normal Variance, μ is unknown

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is unknown**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \bar{X}}{\sigma_0} \right)^2 = \frac{(n-1) \cdot S^2}{\sigma_0^2}$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $\chi^2 \sim \chi^2(n-1)$;

\mathcal{H}_1 is	RR is
$\sigma^2 \neq \sigma_0^2$	$\chi^2 \notin \left[\chi_{n-1, \frac{\alpha}{2}}^2, \chi_{n-1, 1-\frac{\alpha}{2}}^2 \right]$
$\sigma^2 > \sigma_0^2$	$\chi^2 > \chi_{n-1, 1-\alpha}^2$

Test for the Normal Variance, μ is unknown

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is unknown**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \bar{X}}{\sigma_0} \right)^2 = \frac{(n-1) \cdot S^2}{\sigma_0^2}$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $\chi^2 \sim \chi^2(n-1)$;

\mathcal{H}_1 is	RR is
$\sigma^2 \neq \sigma_0^2$	$\chi^2 \notin \left[\chi_{n-1, \frac{\alpha}{2}}^2, \chi_{n-1, 1-\frac{\alpha}{2}}^2 \right]$
$\sigma^2 > \sigma_0^2$	$\chi^2 > \chi_{n-1, 1-\alpha}^2$
$\sigma^2 < \sigma_0^2$	

Test for the Normal Variance, μ is unknown

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, μ **is unknown**, the Parameter (our unknown) is σ^2 ;

Null Hypothesis: $\mathcal{H}_0 : \sigma^2 = \sigma_0^2$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $\chi^2 = \sum_{k=1}^n \left(\frac{X_k - \bar{X}}{\sigma_0} \right)^2 = \frac{(n-1) \cdot S^2}{\sigma_0^2}$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $\chi^2 \sim \chi^2(n-1)$;

\mathcal{H}_1 is	RR is
$\sigma^2 \neq \sigma_0^2$	$\chi^2 \notin \left[\chi_{n-1, \frac{\alpha}{2}}^2, \chi_{n-1, 1-\frac{\alpha}{2}}^2 \right]$
$\sigma^2 > \sigma_0^2$	$\chi^2 > \chi_{n-1, 1-\alpha}^2$
$\sigma^2 < \sigma_0^2$	$\chi^2 < \chi_{n-1, \alpha}^2$

Large Sample Hypothesis Testing

aka

Asymptotic Testing

Asymptotic Test for the Mean of General Distribution

Model: X_1, X_2, \dots, X_n are IID from some Distribution with Mean μ and Variance σ^2 . σ^2 is not known. The Parameter is μ ;

Asymptotic Test for the Mean of General Distribution

Model: X_1, X_2, \dots, X_n are IID from some Distribution with Mean μ and Variance σ^2 . σ^2 is not known. The Parameter is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Asymptotic Test for the Mean of General Distribution

Model: X_1, X_2, \dots, X_n are IID from some Distribution with Mean μ and Variance σ^2 . σ^2 is not known. The Parameter is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$;

Asymptotic Test for the Mean of General Distribution

Model: X_1, X_2, \dots, X_n are IID from some Distribution with Mean μ and Variance σ^2 . σ^2 is not known. The Parameter is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$; This means that we want to have

$$\mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}) \rightarrow \alpha, \quad \text{as } n \rightarrow +\infty$$

Asymptotic Test for the Mean of General Distribution

Model: X_1, X_2, \dots, X_n are IID from some Distribution with Mean μ and Variance σ^2 . σ^2 is not known. The Parameter is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$; This means that we want to have

$$\mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}) \rightarrow \alpha, \quad \text{as } n \rightarrow +\infty$$

Test Statistics: $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$, where $S^2 =$

Asymptotic Test for the Mean of General Distribution

Model: X_1, X_2, \dots, X_n are IID from some Distribution with Mean μ and Variance σ^2 . σ^2 is not known. The Parameter is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$; This means that we want to have

$$\mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}) \rightarrow \alpha, \quad \text{as } n \rightarrow +\infty$$

Test Statistics: $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$, where $S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}$.

Asymptotic Test for the Mean of General Distribution

Model: X_1, X_2, \dots, X_n are IID from some Distribution with Mean μ and Variance σ^2 . σ^2 is not known. The Parameter is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$; This means that we want to have

$$\mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}) \rightarrow \alpha, \quad \text{as } n \rightarrow +\infty$$

Test Statistics: $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$, where $S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}$.

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $t \xrightarrow{D}$

Asymptotic Test for the Mean of General Distribution

Model: X_1, X_2, \dots, X_n are IID from some Distribution with Mean μ and Variance σ^2 . σ^2 is not known. The Parameter is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$; This means that we want to have

$$\mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}) \rightarrow \alpha, \quad \text{as } n \rightarrow +\infty$$

Test Statistics: $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$, where $S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}$.

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $t \xrightarrow{D} \mathcal{N}(0, 1)$;

Asymptotic Test for the Mean of General Distribution

Model: X_1, X_2, \dots, X_n are IID from some Distribution with Mean μ and Variance σ^2 . σ^2 is not known. The Parameter is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$; This means that we want to have

$$\mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}) \rightarrow \alpha, \quad \text{as } n \rightarrow +\infty$$

Test Statistics: $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$, where $S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}$.

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $t \xrightarrow{D} \mathcal{N}(0, 1)$;

\mathcal{H}_1 is	RR is
$\mu \neq \mu_0$	

Asymptotic Test for the Mean of General Distribution

Model: X_1, X_2, \dots, X_n are IID from some Distribution with Mean μ and Variance σ^2 . σ^2 is not known. The Parameter is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$; This means that we want to have

$$\mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}) \rightarrow \alpha, \quad \text{as } n \rightarrow +\infty$$

Test Statistics: $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$, where $S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}$.

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $t \xrightarrow{D} \mathcal{N}(0, 1)$;

\mathcal{H}_1 is	RR is
$\mu \neq \mu_0$	$ t > z_{1-\frac{\alpha}{2}}$

Asymptotic Test for the Mean of General Distribution

Model: X_1, X_2, \dots, X_n are IID from some Distribution with Mean μ and Variance σ^2 . σ^2 is not known. The Parameter is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$; This means that we want to have

$$\mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}) \rightarrow \alpha, \quad \text{as } n \rightarrow +\infty$$

Test Statistics: $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$, where $S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}$.

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $t \xrightarrow{D} \mathcal{N}(0, 1)$;

\mathcal{H}_1 is	RR is
$\mu \neq \mu_0$	$ t > z_{1-\frac{\alpha}{2}}$
$\mu > \mu_0$	

Asymptotic Test for the Mean of General Distribution

Model: X_1, X_2, \dots, X_n are IID from some Distribution with Mean μ and Variance σ^2 . σ^2 is not known. The Parameter is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$; This means that we want to have

$$\mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}) \rightarrow \alpha, \quad \text{as } n \rightarrow +\infty$$

Test Statistics: $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$, where $S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}$.

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $t \xrightarrow{D} \mathcal{N}(0, 1)$;

\mathcal{H}_1 is	RR is
$\mu \neq \mu_0$	$ t > z_{1-\frac{\alpha}{2}}$
$\mu > \mu_0$	$t > z_{1-\alpha}$

Asymptotic Test for the Mean of General Distribution

Model: X_1, X_2, \dots, X_n are IID from some Distribution with Mean μ and Variance σ^2 . σ^2 is not known. The Parameter is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$; This means that we want to have

$$\mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}) \rightarrow \alpha, \quad \text{as } n \rightarrow +\infty$$

Test Statistics: $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$, where $S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}$.

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $t \xrightarrow{D} \mathcal{N}(0, 1)$;

\mathcal{H}_1 is	RR is
$\mu \neq \mu_0$	$ t > z_{1-\frac{\alpha}{2}}$
$\mu > \mu_0$	$t > z_{1-\alpha}$
$\mu < \mu_0$	

Asymptotic Test for the Mean of General Distribution

Model: X_1, X_2, \dots, X_n are IID from some Distribution with Mean μ and Variance σ^2 . σ^2 is not known. The Parameter is μ ;

Null Hypothesis: $\mathcal{H}_0 : \mu = \mu_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$; This means that we want to have

$$\mathbb{P}(\text{Type I Error}) = \mathbb{P}(\text{Reject } \mathcal{H}_0 \mid \mathcal{H}_0 \text{ is True}) \rightarrow \alpha, \quad \text{as } n \rightarrow +\infty$$

Test Statistics: $t = \frac{\bar{X} - \mu_0}{S/\sqrt{n}}$, where $S^2 = \frac{\sum_{k=1}^n (X_k - \bar{X})^2}{n-1}$.

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $t \xrightarrow{D} \mathcal{N}(0, 1)$;

\mathcal{H}_1 is	RR is
$\mu \neq \mu_0$	$ t > z_{1-\frac{\alpha}{2}}$
$\mu > \mu_0$	$t > z_{1-\alpha}$
$\mu < \mu_0$	$t < z_{\alpha}$

Asymptotic Tests based on MLEs

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{F}_\theta;$

Asymptotic Tests based on MLEs

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{F}_\theta;$

Null Hypothesis: $\mathcal{H}_0 : \theta = \theta_0$

Asymptotic Tests based on MLEs

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{F}_\theta;$

Null Hypothesis: $\mathcal{H}_0 : \theta = \theta_0$

Asymptotic Significance Level: $\alpha \in (0, 1);$

Asymptotic Tests based on MLEs

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{F}_\theta$;

Null Hypothesis: $\mathcal{H}_0 : \theta = \theta_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $W = \frac{\hat{\theta}^{MLE} - \theta_0}{1/\sqrt{n \cdot \mathcal{I}(\theta_0)}}$ or $W = \frac{\hat{\theta}^{MLE} - \theta_0}{1/\sqrt{n \cdot \mathcal{I}(\hat{\theta}^{MLE})}}$

Asymptotic Tests based on MLEs

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{F}_\theta;$

Null Hypothesis: $\mathcal{H}_0 : \theta = \theta_0$

Asymptotic Significance Level: $\alpha \in (0, 1);$

Test Statistics: $W = \frac{\hat{\theta}^{MLE} - \theta_0}{1/\sqrt{n \cdot \mathcal{I}(\theta_0)}} \quad \text{or} \quad W = \frac{\hat{\theta}^{MLE} - \theta_0}{1/\sqrt{n \cdot \mathcal{I}(\hat{\theta}^{MLE})}}$

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $W \xrightarrow{D}$

Asymptotic Tests based on MLEs

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{F}_\theta$;

Null Hypothesis: $\mathcal{H}_0 : \theta = \theta_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $W = \frac{\hat{\theta}^{MLE} - \theta_0}{1/\sqrt{n \cdot \mathcal{I}(\theta_0)}}$ or $W = \frac{\hat{\theta}^{MLE} - \theta_0}{1/\sqrt{n \cdot \mathcal{I}(\hat{\theta}^{MLE})}}$

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $W \xrightarrow{D} \mathcal{N}(0, 1)$;

Asymptotic Tests based on MLEs

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{F}_\theta$;

Null Hypothesis: $\mathcal{H}_0 : \theta = \theta_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $W = \frac{\hat{\theta}^{MLE} - \theta_0}{1/\sqrt{n \cdot \mathcal{I}(\theta_0)}}$ or $W = \frac{\hat{\theta}^{MLE} - \theta_0}{1/\sqrt{n \cdot \mathcal{I}(\hat{\theta}^{MLE})}}$

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $W \xrightarrow{D} \mathcal{N}(0, 1)$;

\mathcal{H}_1 is	RR is
$\theta \neq \theta_0$	$ W > z_{1-\frac{\alpha}{2}}$
$\theta > \theta_0$	$W > z_{1-\alpha}$
$\theta < \theta_0$	$W < z_\alpha$

Example: Asymptotic Test for Bernoulli Model

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \text{Bernoulli}(p);$

Example: Asymptotic Test for Bernoulli Model

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \text{Bernoulli}(p)$; In this case, $\hat{p}^{MLE} =$

Example: Asymptotic Test for Bernoulli Model

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \text{Bernoulli}(p)$; In this case, $\hat{p}^{MLE} = \bar{X}$ and $\mathcal{I}(p) =$

Example: Asymptotic Test for Bernoulli Model

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \text{Bernoulli}(p)$; In this case, $\hat{p}^{MLE} = \bar{X}$ and

$$\mathcal{I}(p) = \frac{1}{p(1-p)};$$

Example: Asymptotic Test for Bernoulli Model

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \text{Bernoulli}(p)$; In this case, $\hat{p}^{MLE} = \bar{X}$ and

$$\mathcal{I}(p) = \frac{1}{p(1-p)};$$

Null Hypothesis: $\mathcal{H}_0 : p = p_0$

Example: Asymptotic Test for Bernoulli Model

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \text{Bernoulli}(p)$; In this case, $\hat{p}^{MLE} = \bar{X}$ and

$$\mathcal{I}(p) = \frac{1}{p(1-p)};$$

Null Hypothesis: $\mathcal{H}_0 : p = p_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$;

Example: Asymptotic Test for Bernoulli Model

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \text{Bernoulli}(p)$; In this case, $\hat{p}^{MLE} = \bar{X}$ and $\mathcal{I}(p) = \frac{1}{p(1-p)}$;

Null Hypothesis: $\mathcal{H}_0 : p = p_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $W = \frac{\bar{X} - p_0}{\sqrt{\frac{p_0(1-p_0)}{n}}}$ or $W = \frac{\bar{X} - p_0}{\sqrt{\frac{\bar{X}(1-\bar{X})}{n}}}$

Example: Asymptotic Test for Bernoulli Model

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \text{Bernoulli}(p)$; In this case, $\hat{p}^{MLE} = \bar{X}$ and $\mathcal{I}(p) = \frac{1}{p(1-p)}$;

Null Hypothesis: $\mathcal{H}_0 : p = p_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $W = \frac{\bar{X} - p_0}{\sqrt{\frac{p_0(1-p_0)}{n}}}$ or $W = \frac{\bar{X} - p_0}{\sqrt{\frac{\bar{X}(1-\bar{X})}{n}}}$

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $W \xrightarrow{D}$

Example: Asymptotic Test for Bernoulli Model

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \text{Bernoulli}(p)$; In this case, $\hat{p}^{MLE} = \bar{X}$ and $\mathcal{I}(p) = \frac{1}{p(1-p)}$;

Null Hypothesis: $\mathcal{H}_0 : p = p_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $W = \frac{\bar{X} - p_0}{\sqrt{\frac{p_0(1-p_0)}{n}}}$ or $W = \frac{\bar{X} - p_0}{\sqrt{\frac{\bar{X}(1-\bar{X})}{n}}}$

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $W \xrightarrow{D} \mathcal{N}(0, 1)$;

Example: Asymptotic Test for Bernoulli Model

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \text{Bernoulli}(p)$; In this case, $\hat{p}^{MLE} = \bar{X}$ and $\mathcal{I}(p) = \frac{1}{p(1-p)}$;

Null Hypothesis: $\mathcal{H}_0 : p = p_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $W = \frac{\bar{X} - p_0}{\sqrt{\frac{p_0(1-p_0)}{n}}}$ or $W = \frac{\bar{X} - p_0}{\sqrt{\frac{\bar{X}(1-\bar{X})}{n}}}$

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $W \xrightarrow{D} \mathcal{N}(0, 1)$;

\mathcal{H}_1 is	RR is
$p \neq p_0$	$ W > z_{1-\frac{\alpha}{2}}$
$p > p_0$	$W > z_{1-\alpha}$
$p < p_0$	$W < z_{\alpha}$

Example: Asymptotic Test for Bernoulli Model

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \text{Bernoulli}(p)$; In this case, $\hat{p}^{MLE} = \bar{X}$ and $\mathcal{I}(p) = \frac{1}{p(1-p)}$;

Null Hypothesis: $\mathcal{H}_0 : p = p_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $W = \frac{\bar{X} - p_0}{\sqrt{\frac{p_0(1-p_0)}{n}}}$ or $W = \frac{\bar{X} - p_0}{\sqrt{\frac{\bar{X}(1-\bar{X})}{n}}}$

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $W \xrightarrow{D} \mathcal{N}(0, 1)$;

\mathcal{H}_1 is	RR is
$p \neq p_0$	$ W > z_{1-\frac{\alpha}{2}}$
$p > p_0$	$W > z_{1-\alpha}$
$p < p_0$	$W < z_{\alpha}$

Note: People use this Test only if $n \cdot p_0 > 5$ and $n \cdot (1 - p_0) > 5$.

Wald Test

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{F}_\theta;$

Wald Test

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{F}_\theta$; We take any Estimator $\hat{\theta}$ which is Asymptotically Normal:

$$W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})} \xrightarrow{D} \mathcal{N}(0, 1)$$

Wald Test

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{F}_\theta$; We take any Estimator $\hat{\theta}$ which is Asymptotically Normal:

$$W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})} \xrightarrow{D} \mathcal{N}(0, 1)$$

Null Hypothesis: $\mathcal{H}_0 : \theta = \theta_0$

Wald Test

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{F}_\theta$; We take any Estimator $\hat{\theta}$ which is Asymptotically Normal:

$$W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})} \xrightarrow{D} \mathcal{N}(0, 1)$$

Null Hypothesis: $\mathcal{H}_0 : \theta = \theta_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$;

Wald Test

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{F}_\theta$; We take any Estimator $\hat{\theta}$ which is Asymptotically Normal:

$$W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})} \xrightarrow{D} \mathcal{N}(0, 1)$$

Null Hypothesis: $\mathcal{H}_0 : \theta = \theta_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})}$ or $W = \frac{\hat{\theta} - \theta_0}{SE(\theta_0)}$

Wald Test

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{F}_\theta$; We take any Estimator $\hat{\theta}$ which is Asymptotically Normal:

$$W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})} \xrightarrow{D} \mathcal{N}(0, 1)$$

Null Hypothesis: $\mathcal{H}_0 : \theta = \theta_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})}$ or $W = \frac{\hat{\theta} - \theta_0}{SE(\theta_0)}$

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $W \xrightarrow{D}$

Wald Test

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{F}_\theta$; We take any Estimator $\hat{\theta}$ which is Asymptotically Normal:

$$W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})} \xrightarrow{D} \mathcal{N}(0, 1)$$

Null Hypothesis: $\mathcal{H}_0 : \theta = \theta_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})}$ or $W = \frac{\hat{\theta} - \theta_0}{SE(\theta_0)}$

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $W \xrightarrow{D} \mathcal{N}(0, 1)$;

Wald Test

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{F}_\theta$; We take any Estimator $\hat{\theta}$ which is Asymptotically Normal:

$$W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})} \xrightarrow{D} \mathcal{N}(0, 1)$$

Null Hypothesis: $\mathcal{H}_0 : \theta = \theta_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $W = \frac{\hat{\theta} - \theta_0}{\widehat{SE}(\hat{\theta})}$ or $W = \frac{\hat{\theta} - \theta_0}{SE(\theta_0)}$

Asymptotic Distrib of the TS Under \mathcal{H}_0 : $W \xrightarrow{D} \mathcal{N}(0, 1)$;

\mathcal{H}_1 is	RR is
$\theta \neq \theta_0$	$ W > z_{1-\frac{\alpha}{2}}$
$\theta > \theta_0$	$W > z_{1-\alpha}$
$\theta < \theta_0$	$W < z_\alpha$

Note

Note: In all above Asymptotic Tests, one can replace the quantiles z_p of the Standard Normal by the quantiles $t_{n-1,p}$ of $t(n-1)$, since, for large n ,

$$t_{n-1,p} \approx z_p.$$

Note

Note: In all above Asymptotic Tests, one can replace the quantiles z_p of the Standard Normal by the quantiles $t_{n-1,p}$ of $t(n-1)$, since, for large n ,

$$t_{n-1,p} \approx z_p.$$

```
a <- 0.05  
qnorm(1-a/2)
```

```
## [1] 1.959964
```

```
n <- seq(50,250,50)  
qt(1-a/2, df = n)
```

```
## [1] 2.008559 1.983972 1.975905 1.971896 1.969498
```

Note

Note: In all above Asymptotic Tests, one can replace the quantiles z_p of the Standard Normal by the quantiles $t_{n-1,p}$ of $t(n-1)$, since, for large n ,

$$t_{n-1,p} \approx z_p.$$

```
a <- 0.05  
qnorm(1-a/2)
```

```
## [1] 1.959964
```

```
n <- seq(50,250,50)  
qt(1-a/2, df = n)
```

```
## [1] 2.008559 1.983972 1.975905 1.971896 1.969498
```

Another point is that, since $|t_{n-1,p}| > |z_p|$, $p \neq 0.5$, it is safer to have a little bit smaller Rejection Region:

Note

Note: In all above Asymptotic Tests, one can replace the quantiles z_p of the Standard Normal by the quantiles $t_{n-1,p}$ of $t(n-1)$, since, for large n ,

$$t_{n-1,p} \approx z_p.$$

```
a <- 0.05  
qnorm(1-a/2)
```

```
## [1] 1.959964
```

```
n <- seq(50,250,50)  
qt(1-a/2, df = n)
```

```
## [1] 2.008559 1.983972 1.975905 1.971896 1.969498
```

Another point is that, since $|t_{n-1,p}| > |z_p|$, $p \neq 0.5$, it is safer to have a little bit smaller Rejection Region: say, for the Two-Sided Tests, if $|W| > t_{n-1,1-\alpha/2}$, then, for sure, also $|W| > z_{1-\alpha/2}$.

Two Sample Tests

Z-Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2),$

Z-Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2), Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2),$

Z-Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$,
 σ_X, σ_Y are known,

Z-Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$,
 σ_X, σ_Y **are known**, and X_k -s and Y_j -s are all Independent.

Z-Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are known**, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

Z-Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are known**, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

Null Hypothesis: $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

Z-Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{iid}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are known**, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

Null Hypothesis: $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Z-Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are known**, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

Null Hypothesis: $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $Z =$

Z-Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are known**, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

Null Hypothesis: $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $Z = \frac{(\bar{X} - \bar{Y}) - \mu_0}{\sigma_{\bar{X} - \bar{Y}}}$

Z-Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are known**, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

Null Hypothesis: $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics:
$$Z = \frac{(\bar{X} - \bar{Y}) - \mu_0}{\sqrt{\frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}}}$$

Z-Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are known**, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

Null Hypothesis: $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics:
$$Z = \frac{(\bar{X} - \bar{Y}) - \mu_0}{\sqrt{\frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}}}$$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $Z \sim$

Z-Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are known**, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

Null Hypothesis: $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics:
$$Z = \frac{(\bar{X} - \bar{Y}) - \mu_0}{\sqrt{\frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}}}$$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $Z \sim \mathcal{N}(0, 1)$;

Z-Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{iid}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y are known, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

Null Hypothesis: $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics:
$$Z = \frac{(\bar{X} - \bar{Y}) - \mu_0}{\sqrt{\frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}}}$$

Distrib of the Test-Statistics Under \mathcal{H}_0 : $Z \sim \mathcal{N}(0, 1)$;

\mathcal{H}_1 is	RR is
$\mu_X - \mu_Y \neq \mu_0$	$ Z > z_{1-\frac{\alpha}{2}}$
$\mu_X - \mu_Y > \mu_0$	$Z > z_{1-\alpha}$
$\mu_X - \mu_Y < \mu_0$	$Z < z_\alpha$

t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu_X, \sigma_X^2),$

t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu_X, \sigma_X^2), Y_1, Y_2, \dots, Y_m \stackrel{iid}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2),$

t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{iid}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$,
 σ_X, σ_Y are unknown,

t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$,
 σ_X, σ_Y **are unknown**, and X_k -s and Y_j -s are all Independent.

t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{iid}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are unknown**, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are unknown**, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

Additional Assumption: $\sigma_X = \sigma_Y$

t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are unknown**, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

Additional Assumption: $\sigma_X = \sigma_Y$ (can be Tested separately, by F -Test)

t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are unknown**, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

Additional Assumption: $\sigma_X = \sigma_Y$ (can be Tested separately, by F -Test)

Null Hypothesis: $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are unknown**, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

Additional Assumption: $\sigma_X = \sigma_Y$ (can be Tested separately, by F -Test)

Null Hypothesis: $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

Significance Level: $\alpha \in (0, 1)$;

t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are unknown**, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

Additional Assumption: $\sigma_X = \sigma_Y$ (can be Tested separately, by F -Test)

Null Hypothesis: $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $t = \frac{(\bar{X} - \bar{Y}) - \mu_0}{S_p \cdot \sqrt{\frac{1}{n} + \frac{1}{m}}}$, where S_p is the **Pooled**

Sample Deviation:

t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are unknown**, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

Additional Assumption: $\sigma_X = \sigma_Y$ (can be Tested separately, by F -Test)

Null Hypothesis: $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics: $t = \frac{(\bar{X} - \bar{Y}) - \mu_0}{S_p \cdot \sqrt{\frac{1}{n} + \frac{1}{m}}}$, where S_p is the **Pooled**

Sample Deviation:

$$S_p^2 = \frac{(n-1)S_X^2 + (m-1)S_Y^2}{n+m-2} = \frac{\sum_{k=1}^n (X_k - \bar{X})^2 + \sum_{k=1}^m (Y_k - \bar{Y})^2}{n+m-2}.$$

t -Test for the Difference of two Normals Means, Cont'd

Distrib of the Test-Statistics Under \mathcal{H}_0 : $t \sim$

t -Test for the Difference of two Normals Means, Cont'd

Distrib of the Test-Statistics Under \mathcal{H}_0 : $t \sim t(n + m - 2)$;

t -Test for the Difference of two Normals Means, Cont'd

Distrib of the Test-Statistics Under \mathcal{H}_0 : $t \sim t(n + m - 2)$;

Rejection Region:

\mathcal{H}_1 is	RR is
$\mu_X - \mu_Y \neq \mu_0$	$ t > t_{n+m-2, 1-\frac{\alpha}{2}}$
$\mu_X - \mu_Y > \mu_0$	$t > t_{n+m-2, 1-\alpha}$
$\mu_X - \mu_Y < \mu_0$	$t < t_{n+m-2, \alpha}$

t -Test for the Difference of two Normals Means, Cont'd

Distrib of the Test-Statistics Under \mathcal{H}_0 : $t \sim t(n + m - 2)$;

Rejection Region:

\mathcal{H}_1 is	RR is
$\mu_X - \mu_Y \neq \mu_0$	$ t > t_{n+m-2, 1-\frac{\alpha}{2}}$
$\mu_X - \mu_Y > \mu_0$	$t > t_{n+m-2, 1-\alpha}$
$\mu_X - \mu_Y < \mu_0$	$t < t_{n+m-2, \alpha}$

Note: This Test is called the **Pooled t -Test**

t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu_X, \sigma_X^2),$

t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2), Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2),$

t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$,
 σ_X, σ_Y are unknown,

t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$,
 σ_X, σ_Y **are unknown**, and X_k -s and Y_j -s are all Independent.

t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are unknown**, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

t-Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are unknown**, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

Additional Assumption: $\sigma_X \neq \sigma_Y$ (can be Tested separately, by *F*-Test)

t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are unknown**, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

Additional Assumption: $\sigma_X \neq \sigma_Y$ (can be Tested separately, by F -Test)

Null Hypothesis: $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are unknown**, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

Additional Assumption: $\sigma_X \neq \sigma_Y$ (can be Tested separately, by F -Test)

Null Hypothesis: $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

Significance Level: $\alpha \in (0, 1)$;

t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are unknown**, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

Additional Assumption: $\sigma_X \neq \sigma_Y$ (can be Tested separately, by F -Test)

Null Hypothesis: $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics:

$$t = \frac{(\bar{X} - \bar{Y}) - \mu_0}{\sqrt{\frac{S_X^2}{n} + \frac{S_Y^2}{m}}},$$

t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y are unknown, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $\mu_X - \mu_Y$;

Additional Assumption: $\sigma_X \neq \sigma_Y$ (can be Tested separately, by F -Test)

Null Hypothesis: $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics:

$$t = \frac{(\bar{X} - \bar{Y}) - \mu_0}{\sqrt{\frac{S_X^2}{n} + \frac{S_Y^2}{m}}},$$

where S_X and S_Y are the Sample SDs for X and Y , respectively.

t -Test for the Difference of two Normals Means, Cont'd

Distrib of the Test-Statistics Under \mathcal{H}_0 : Approximately,

$$t \approx$$

t -Test for the Difference of two Normals Means, Cont'd

Distrib of the Test-Statistics Under \mathcal{H}_0 : Approximately,
 $t \approx t(\nu)$, where ν is

t -Test for the Difference of two Normals Means, Cont'd

Distrib of the Test-Statistics Under \mathcal{H}_0 : Approximately,
 $t \approx t(\nu)$, where ν is some scary thing...

t -Test for the Difference of two Normals Means, Cont'd

Distrib of the Test-Statistics Under \mathcal{H}_0 : Approximately,
 $t \approx t(\nu)$, where ν is some scary thing... given by

$$\nu = \left\lfloor \frac{\left(\frac{S_X^2}{n} + \frac{S_Y^2}{m} \right)^2}{\frac{(S_X^2/n)^2}{n-1} + \frac{(S_Y^2/m)^2}{m-1}} \right\rfloor$$

t -Test for the Difference of two Normals Means, Cont'd

Distrib of the Test-Statistics Under \mathcal{H}_0 : Approximately,
 $t \approx t(\nu)$, where ν is some scary thing... given by

$$\nu = \left\lfloor \frac{\left(\frac{S_X^2}{n} + \frac{S_Y^2}{m} \right)^2}{\frac{(S_X^2/n)^2}{n-1} + \frac{(S_Y^2/m)^2}{m-1}} \right\rfloor$$

Rejection Region:

\mathcal{H}_1 is	RR is
$\mu_X - \mu_Y \neq \mu_0$	$ t > t_{\nu, 1-\frac{\alpha}{2}}$
$\mu_X - \mu_Y > \mu_0$	$t > t_{\nu, 1-\alpha}$
$\mu_X - \mu_Y < \mu_0$	$t < t_{\nu, \alpha}$

t -Test for the Difference of two Normals Means, Cont'd

Distrib of the Test-Statistics Under \mathcal{H}_0 : Approximately,
 $t \approx t(\nu)$, where ν is some scary thing... given by

$$\nu = \left\lfloor \frac{\left(\frac{S_X^2}{n} + \frac{S_Y^2}{m} \right)^2}{\frac{(S_X^2/n)^2}{n-1} + \frac{(S_Y^2/m)^2}{m-1}} \right\rfloor$$

Rejection Region:

\mathcal{H}_1 is	RR is
$\mu_X - \mu_Y \neq \mu_0$	$ t > t_{\nu, 1-\frac{\alpha}{2}}$
$\mu_X - \mu_Y > \mu_0$	$t > t_{\nu, 1-\alpha}$
$\mu_X - \mu_Y < \mu_0$	$t < t_{\nu, \alpha}$

Note: The formula above for the DF ν is called **Welch – Satterthwaite Equation**, and the Tests is called the **Welch Test**.

Paired t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2),$

Paired t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2), Y_1, Y_2, \dots, Y_n \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2),$

Paired t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_n \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$,
 σ_X, σ_Y are unknown.

Paired t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_n \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$,
 σ_X, σ_Y **are unknown**. The Parameter of interest is $\mu_X - \mu_Y$;

Paired t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_n \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$,
 σ_X, σ_Y **are unknown**. The Parameter of interest is $\mu_X - \mu_Y$;

Note: Here we have the same number of X_k and Y_k ;

Paired t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_n \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are unknown**. The Parameter of interest is $\mu_X - \mu_Y$;

Note: Here we have the same number of X_k and Y_k ; also, importantly, X_k and Y_k can be dependent!

Paired t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_n \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are unknown**. The Parameter of interest is $\mu_X - \mu_Y$;

Note: Here we have the same number of X_k and Y_k ; also, importantly, X_k and Y_k can be dependent!

Notation: $D_k = X_k - Y_k$;

Paired t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_n \stackrel{iid}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are unknown**. The Parameter of interest is $\mu_X - \mu_Y$;

Note: Here we have the same number of X_k and Y_k ; also, importantly, X_k and Y_k can be dependent!

Notation: $D_k = X_k - Y_k$; clearly,

$$\mathbb{E}(D_k) = \mu_X - \mu_Y.$$

Paired t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_n \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are unknown**. The Parameter of interest is $\mu_X - \mu_Y$;

Note: Here we have the same number of X_k and Y_k ; also, importantly, X_k and Y_k can be dependent!

Notation: $D_k = X_k - Y_k$; clearly,

$$\mathbb{E}(D_k) = \mu_X - \mu_Y.$$

The Variance of D_k , although the same, $\sigma_D^2 = \text{Var}(X_k - Y_k)$, cannot be calculated, since X_k and Y_k can be dependent. But that's OK, we do not need it.

¹The Test will work also in the case when the Differences are nor Normally Distributed, but the Sample Size n is large. We jut need to use the CLT.

Paired t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_n \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are unknown**. The Parameter of interest is $\mu_X - \mu_Y$;

Note: Here we have the same number of X_k and Y_k ; also, importantly, X_k and Y_k can be dependent!

Notation: $D_k = X_k - Y_k$; clearly,

$$\mathbb{E}(D_k) = \mu_X - \mu_Y.$$

The Variance of D_k , although the same, $\sigma_D^2 = \text{Var}(X_k - Y_k)$, cannot be calculated, since X_k and Y_k can be dependent. But that's OK, we do not need it.

Additional Assumption: We will assume that the Differences D_k are Normally Distributed¹.

¹The Test will work also in the case when the Differences are nor Normally Distributed, but the Sample Size n is large. We jut need to use the CLT.

Paired t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_n \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are unknown**. The Parameter of interest is $\mu_X - \mu_Y$;

Note: Here we have the same number of X_k and Y_k ; also, importantly, X_k and Y_k can be dependent!

Notation: $D_k = X_k - Y_k$; clearly,

$$\mathbb{E}(D_k) = \mu_X - \mu_Y.$$

The Variance of D_k , although the same, $\sigma_D^2 = \text{Var}(X_k - Y_k)$, cannot be calculated, since X_k and Y_k can be dependent. But that's OK, we do not need it.

Additional Assumption: We will assume that the Differences D_k are Normally Distributed¹.

Null Hypothesis: $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

¹The Test will work also in the case when the Differences are nor Normally Distributed, but the Sample Size n is large. We jut need to use the CLT.

Paired t -Test for the Difference of two Normals Means

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_1, Y_2, \dots, Y_n \stackrel{IID}{\sim} \mathcal{N}(\mu_Y, \sigma_Y^2)$, σ_X, σ_Y **are unknown**. The Parameter of interest is $\mu_X - \mu_Y$;

Note: Here we have the same number of X_k and Y_k ; also, importantly, X_k and Y_k can be dependent!

Notation: $D_k = X_k - Y_k$; clearly,

$$\mathbb{E}(D_k) = \mu_X - \mu_Y.$$

The Variance of D_k , although the same, $\sigma_D^2 = \text{Var}(X_k - Y_k)$, cannot be calculated, since X_k and Y_k can be dependent. But that's OK, we do not need it.

Additional Assumption: We will assume that the Differences D_k are Normally Distributed¹.

Null Hypothesis: $\mathcal{H}_0 : \mu_X - \mu_Y = \mu_0$

Asymptotic Significance Level: $\alpha \in (0, 1)$;

¹The Test will work also in the case when the Differences are nor Normally Distributed, but the Sample Size n is large. We jut need to use the CLT.

Paired t -Test for the Difference of two Normals Means, Cont'd

Test Statistics: $t = \frac{\bar{D} - \mu_0}{S_D/\sqrt{n}}$, where S_D is the Sample Deviation of D .

²Or, Asymptotically, $t \approx t(n-1)$ or $t \approx \mathcal{N}(0,1)$, if D_k -s are not Normal, but n is large.

Paired t -Test for the Difference of two Normals Means, Cont'd

Test Statistics: $t = \frac{\bar{D} - \mu_0}{S_D/\sqrt{n}}$, where S_D is the Sample Deviation of D .

Distrib of the Test-Statistics Under \mathcal{H}_0 :² $t \sim t(n-1)$;

²Or, Asymptotically, $t \approx t(n-1)$ or $t \approx \mathcal{N}(0,1)$, if D_k -s are not Normal, but n is large.

Paired t -Test for the Difference of two Normals Means, Cont'd

Test Statistics: $t = \frac{\bar{D} - \mu_0}{S_D/\sqrt{n}}$, where S_D is the Sample Deviation of D .

Distrib of the Test-Statistics Under \mathcal{H}_0 :² $t \sim t(n-1)$;

Rejection Region:

\mathcal{H}_1 is	RR is
$\mu_X - \mu_Y \neq \mu_0$	$ t > t_{n-1, 1-\frac{\alpha}{2}}$
$\mu_X - \mu_Y > \mu_0$	$t > t_{n-1, 1-\alpha}$
$\mu_X - \mu_Y < \mu_0$	$t < t_{n-1, \alpha}$

²Or, Asymptotically, $t \approx t(n-1)$ or $t \approx \mathcal{N}(0, 1)$, if D_k -s are not Normal, but n is large.

Paired t -Test for the Difference of two Normals Means, Cont'd

Test Statistics: $t = \frac{\bar{D} - \mu_0}{S_D/\sqrt{n}}$, where S_D is the Sample Deviation of D .

Distrib of the Test-Statistics Under \mathcal{H}_0 :² $t \sim t(n-1)$;

Rejection Region:

\mathcal{H}_1 is	RR is
$\mu_X - \mu_Y \neq \mu_0$	$ t > t_{n-1, 1-\frac{\alpha}{2}}$
$\mu_X - \mu_Y > \mu_0$	$t > t_{n-1, 1-\alpha}$
$\mu_X - \mu_Y < \mu_0$	$t < t_{n-1, \alpha}$

Note: This Test is called the **Paired t -Test**

²Or, Asymptotically, $t \approx t(n-1)$ or $t \approx \mathcal{N}(0, 1)$, if D_k -s are not Normal, but n is large.

Two Sample test for Proportions

Model: $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \text{Bernoulli}(p_X),$

Two Sample test for Proportions

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \text{Bernoulli}(p_X),$
 $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \text{Bernoulli}(p_Y),$

Two Sample test for Proportions

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \text{Bernoulli}(p_X)$,
 $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \text{Bernoulli}(p_Y)$, and X_k -s and Y_j -s are all Independent.

Two Sample test for Proportions

Model: $X_1, X_2, \dots, X_n \overset{IID}{\sim} \text{Bernoulli}(p_X)$,
 $Y_1, Y_2, \dots, Y_m \overset{IID}{\sim} \text{Bernoulli}(p_Y)$, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $p_X - p_Y$;

Two Sample test for Proportions

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \text{Bernoulli}(p_X)$,
 $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \text{Bernoulli}(p_Y)$, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $p_X - p_Y$;

Null Hypothesis: $\mathcal{H}_0 : p_X - p_Y = p_0$

Two Sample test for Proportions

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \text{Bernoulli}(p_X)$,
 $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \text{Bernoulli}(p_Y)$, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $p_X - p_Y$;

Null Hypothesis: $\mathcal{H}_0 : p_X - p_Y = p_0$

Significance Level: $\alpha \in (0, 1)$;

Two Sample test for Proportions

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \text{Bernoulli}(p_X)$,
 $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \text{Bernoulli}(p_Y)$, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $p_X - p_Y$;

Null Hypothesis: $\mathcal{H}_0 : p_X - p_Y = p_0$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics:

$$Z = \frac{(\bar{X} - \bar{Y}) - p_0}{\sqrt{\hat{p}(1 - \hat{p}) \left(\frac{1}{n} + \frac{1}{m} \right)}} \quad \text{or} \quad Z = \frac{(\bar{X} - \bar{Y}) - p_0}{\sqrt{\frac{\bar{X}(1 - \bar{X})}{n} + \frac{\bar{Y}(1 - \bar{Y})}{m}}}$$

Two Sample test for Proportions

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \text{Bernoulli}(p_X)$,
 $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \text{Bernoulli}(p_Y)$, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $p_X - p_Y$;

Null Hypothesis: $\mathcal{H}_0 : p_X - p_Y = p_0$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics:

$$Z = \frac{(\bar{X} - \bar{Y}) - p_0}{\sqrt{\hat{p}(1 - \hat{p}) \left(\frac{1}{n} + \frac{1}{m} \right)}} \quad \text{or} \quad Z = \frac{(\bar{X} - \bar{Y}) - p_0}{\sqrt{\frac{\bar{X}(1 - \bar{X})}{n} + \frac{\bar{Y}(1 - \bar{Y})}{m}}}$$

where \hat{p} is the **Pooled Sample Proportion**:

$$\hat{p} = \frac{n}{n + m} \cdot \bar{X} + \frac{m}{n + m} \cdot \bar{Y}$$

Two Sample test for Proportions

Model: $X_1, X_2, \dots, X_n \stackrel{IID}{\sim} \text{Bernoulli}(p_X)$,
 $Y_1, Y_2, \dots, Y_m \stackrel{IID}{\sim} \text{Bernoulli}(p_Y)$, and X_k -s and Y_j -s are all Independent. The Parameter of interest is $p_X - p_Y$;

Null Hypothesis: $\mathcal{H}_0 : p_X - p_Y = p_0$

Significance Level: $\alpha \in (0, 1)$;

Test Statistics:

$$Z = \frac{(\bar{X} - \bar{Y}) - p_0}{\sqrt{\hat{p}(1 - \hat{p}) \left(\frac{1}{n} + \frac{1}{m} \right)}} \quad \text{or} \quad Z = \frac{(\bar{X} - \bar{Y}) - p_0}{\sqrt{\frac{\bar{X}(1 - \bar{X})}{n} + \frac{\bar{Y}(1 - \bar{Y})}{m}}}$$

where \hat{p} is the **Pooled Sample Proportion**:

$$\hat{p} = \frac{n}{n+m} \cdot \bar{X} + \frac{m}{n+m} \cdot \bar{Y} = \frac{X_1 + \dots + X_n + Y_1 + \dots + Y_m}{n+m}.$$

Two Sample test for Proportions, Cont'd

Distrib of the Test-Statistics Under \mathcal{H}_0 : Approximately,
 $Z \approx$

Two Sample test for Proportions, Cont'd

Distrib of the Test-Statistics Under \mathcal{H}_0 : Approximately,
 $Z \approx \mathcal{N}(0, 1)$

Two Sample test for Proportions, Cont'd

Distrib of the Test-Statistics Under \mathcal{H}_0 : Approximately,
 $Z \approx \mathcal{N}(0, 1)$

Rejection Region:

\mathcal{H}_1 is	RR is
$p_X - p_Y \neq p_0$	$ Z > z_{1-\frac{\alpha}{2}}$
$p_X - p_Y > p_0$	$Z > z_{1-\alpha}$
$p_X - p_Y < p_0$	$Z < z_\alpha$

Notes

Note: An important note is that you can perform 2 Sample t -Test even the DataSets are not Normally Distributed, but n and m are large (say, $n, m \geq 30$).

Notes

Note: An important note is that you can perform 2 Sample t -Test even the DataSets are not Normally Distributed, but n and m are large (say, $n, m \geq 30$). This is because the t -Statistics will have, approximately (in the limit), a Standard Normal Distribution, so we can use Asymptotic Tests, and replace the quantiles of Standard Normal by t -quantiles (as we have talked in the previous lecture).

Notes

Note: To perform a two-Sample t -Test in **R**, we can use the same `t.test`.

Notes

Note: To perform a two-Sample t -Test in **R**, we can use the same `t.test`. When the input of `t.test` consists of two Datasets, it performs a two-Sample Test.

Notes

Note: To perform a two-Sample t -Test in **R**, we can use the same `t.test`. When the input of `t.test` consists of two Datasets, it performs a two-Sample Test. Some appropriate parameters are:

- ▶ `paired` : if TRUE, then it is doing a Paired t -Test. The default value is FALSE;

Notes

Note: To perform a two-Sample t -Test in **R**, we can use the same `t.test`. When the input of `t.test` consists of two Datasets, it performs a two-Sample Test. Some appropriate parameters are:

- ▶ `paired` : if TRUE, then it is doing a Paired t -Test. The default value is FALSE;
- ▶ `mu` : our μ_0 , in two-Sample Test, the value of $\mu_X - \mu_Y$ from the Null.

Notes

Note: To perform a two-Sample t -Test in **R**, we can use the same `t.test`. When the input of `t.test` consists of two Datasets, it performs a two-Sample Test. Some appropriate parameters are:

- ▶ `paired` : if TRUE, then it is doing a Paired t -Test. The default value is FALSE;
- ▶ `mu` : our μ_0 , in two-Sample Test, the value of $\mu_X - \mu_Y$ from the Null. The default value is 0, so by default,

$$\mathcal{H}_0 : \mu_X = \mu_Y$$

Notes

Note: To perform a two-Sample t -Test in **R**, we can use the same `t.test`. When the input of `t.test` consists of two Datasets, it performs a two-Sample Test. Some appropriate parameters are:

- ▶ `paired` : if TRUE, then it is doing a Paired t -Test. The default value is FALSE;
- ▶ `mu` : our μ_0 , in two-Sample Test, the value of $\mu_X - \mu_Y$ from the Null. The default value is 0, so by default,

$$\mathcal{H}_0 : \mu_X = \mu_Y$$

- ▶ `var.equal`: if TRUE, then we are dealing with Equal Variances case, so Pooled Standard Deviation is calculated. Otherwise, **R** is doing non-equal Variance Test.

Notes

Note: To perform a two-Sample t -Test in **R**, we can use the same `t.test`. When the input of `t.test` consists of two Datasets, it performs a two-Sample Test. Some appropriate parameters are:

- ▶ `paired` : if TRUE, then it is doing a Paired t -Test. The default value is FALSE;
- ▶ `mu` : our μ_0 , in two-Sample Test, the value of $\mu_X - \mu_Y$ from the Null. The default value is 0, so by default,

$$\mathcal{H}_0 : \mu_X = \mu_Y$$

- ▶ `var.equal`: if TRUE, then we are dealing with Equal Variances case, so Pooled Standard Deviation is calculated. Otherwise, **R** is doing non-equal Variance Test. By default, the value is FALSE.

Note: To perform a two-Sample Proportion Test in **R**, we can use `prop.test` command.