

Assignment 12

Chiral vs Quantum Spin
Hall insulators.

RISHI PARESH JOSHI

2111093

Problems from Topological Quantum Materials Concepts, Models, and Phenomena by Grigory Tchakov

Section 2.6

1) Given Hamiltonian for chern insulator:

$$\vec{H}_{\vec{k}} = \vec{\sigma} \vec{I}_{2 \times 2} \cdot \hat{\vec{\alpha}} \quad ①, \text{ where } \vec{I}_{2 \times 2} \text{ is a } 2 \times 2 \text{ identity matrix and } \vec{\sigma} \text{ are the Pauli matrices.}$$

$$\vec{\alpha} = (\hbar v k_x - \hbar v k_y, m) \quad ②, \text{ where } v \text{ is the fermi velocity and } R_{x/y} \text{ is the canonical momentum.}$$

We need to show that,

(a) Edge modes are given by: $E(k) = -\text{sgn}(m) \pi v \quad -③$

(b) Berry connection components are:

$$A_{\vec{k}, s}^{(k)} = 0 \quad A_{\vec{k}, s}^{(\phi)} = -\left(\frac{1 - s u_3}{2k}\right) \quad ④, \text{ where } u_3 = \frac{\partial \vec{R}}{|\partial \vec{k}|}$$

(c) Berry phase is :

$$\gamma_s = \oint \vec{A}_{\vec{k},s} \cdot d\vec{k} = -\pi(1 - s\mu_3) \quad (5)$$

(d) Berry curvature is:

$$(\vec{\nabla} \times \vec{A}_{\vec{k},s})_3 = \frac{s}{2k} \frac{\partial \mu_3}{\partial k} \quad (6)$$

(e) Relation to the Chern insulator hamiltonian with:

$$\vec{d}_k = (\hbar v k_x, \hbar e k_y, m) - \tilde{\epsilon} \quad (7)$$

Solution:

Writing Hamiltonian in eqn ① in polar coordinates:

$$k_x = k \cos\phi, \quad k_y = k \sin\phi,$$

$$\vec{u}_k = \frac{\vec{d}_k}{1 d_k} = \frac{1}{\sqrt{(\hbar v k)^2 + m^2}} (\hbar v k \cos\phi, -\hbar v k \sin\phi, m) \quad (\text{From ②})$$

Define:

$$u_3 = \frac{m}{\sqrt{(\hbar v k)^2 + m^2}}, \quad u_{||} = \sqrt{1 - u_3^2} = \frac{\hbar v k}{\sqrt{(\hbar v k)^2 + m^2}}$$

$$\vec{u}_k = (u_{||} \cos\phi, -u_{||} \sin\phi, u_3).$$

This mean our Hamiltonian in eqn ① can be written as:

$$\hat{H}_{\vec{k}} = |\vec{d}_{\vec{k}}| \left(\vec{u}_{\vec{k}} \hat{I}_{2 \times 2} \right) \cdot \vec{\sigma}$$

$$\left(\vec{u}_{\vec{k}} \hat{I}_{2 \times 2} \cdot \vec{\sigma} \right) = u_{11} \cos \phi \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} + (-u_{11} \sin \phi) \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} + u_3 \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

$\underbrace{\hspace{10em}}$
 x

$\underbrace{\hspace{10em}}$
 y

$\underbrace{\hspace{10em}}$
 z

$$\Rightarrow \left(\vec{u}_{\vec{k}} \hat{I}_{2 \times 2} \cdot \vec{\sigma} \right) = \begin{bmatrix} u_3 & u_{11} \cos \phi + i u_{11} \sin \phi \\ u_{11} \cos \phi - i u_{11} \sin \phi & -u_3 \end{bmatrix} = \begin{bmatrix} u_3 & u_{11} e^{i\phi} \\ u_{11} e^{-i\phi} & -u_3 \end{bmatrix} \quad \text{--- ⑧}$$

We now find the eigenvalues & hence the eigen vectors:

$$\Rightarrow (\lambda - u_3)(\lambda + u_3) - u_{11}^2 = 0 \quad (\text{From characteristic eqn})$$

$$\Rightarrow \lambda^2 - (u_3^2 + u_{11}^2) = 0$$

$$\lambda = \pm \sqrt{u_3^2 + u_{11}^2} = \pm \sqrt{u_3^2 + u_{11}^2}$$

$$\Rightarrow E_{\vec{k}, \text{bulk}} = |\vec{d}_{\vec{k}}| / \lambda = \sqrt{\hbar v k^2 + m^2} \times \sqrt{\frac{m^2 + (\hbar v k)^2}{m^2 + (\hbar v k)^2}} = \sqrt{(\hbar v k)^2 + m^2}$$

(a) This is the bulk band structure. To find the edge modes, we write eqn ① sub $k_x \rightarrow -i\partial_x, k_y \rightarrow -i\partial_y$:

$$E \Psi(x, y) = [-i\hbar v (\hat{\sigma}_x \partial_y - \hat{\sigma}_y \partial_x) + m \hat{\sigma}_3] \Psi(x, y)$$

Taking the ansatz: $\vec{\Psi}(x, y) = \vec{\phi}(x, y) e^{ikx + iy/\lambda}$, $\vec{\phi}$ is 2×2 with real entries.

$$E \vec{\phi}(x, y) = \hbar v k \hat{\sigma}_x \vec{\phi} + \left(i \frac{\hbar v \omega}{\lambda} \hat{\sigma}_y + m \hat{\sigma}_3 \right) \vec{\phi}$$

Separating the real & complex parts:

$$\Rightarrow E \vec{\phi} = \text{tr} \vartheta k \hat{\sigma}_x \vec{\phi}$$

- I (Energy is real)

$$\left(i \frac{\text{tr} \vartheta}{\lambda} \hat{\sigma}_y + m \hat{\sigma}_z \right) \vec{\phi} = 0$$

Since $\hat{\sigma}_y \cdot \hat{\sigma}_x = i \hat{\sigma}_y$,

$$\hat{\sigma}_z \left(\frac{\text{tr} \vartheta}{\lambda} \hat{\sigma}_x + m \right) \vec{\phi} = 0$$

If $\vec{\phi}$ is an eigenstate of σ_x ,

$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \rightarrow$ eigenvalues $\in \{\pm 1\}$, eigenvectors $\in \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$

$$\hat{\sigma}_x \vec{\phi} = s_x \vec{\phi} \Rightarrow \left(\frac{\text{tr} \vartheta}{\lambda} s_x + m \right) \vec{\phi} = 0 \Rightarrow s_x = -\frac{m \lambda}{\text{tr} \vartheta}$$

Since $s_x = \pm 1$, $\lambda > 0$, $s_x = -\text{sgn}(m)$

$$\Rightarrow -\operatorname{sgn}(m) = -\frac{m\lambda}{\pi e} \Rightarrow \lambda = \frac{\pi e}{(m)} > 0.$$

This ensures as $y \rightarrow -\infty$, the wavefunction decays exponentially.

$$E \vec{\phi} = \text{tr} e K \hat{\sigma}_x \vec{\phi} = -\text{tr} e K \operatorname{sgn}(m) \vec{\phi}$$

$$\Rightarrow E_{k, \text{edge}} = -\operatorname{sgn}(m) \text{tr} e K$$

matching Eq ③.

(b) The eigenvectors of the hamiltonian (Eq ①) of helicity operator (Eq ⑧) are the same.

$$\begin{bmatrix} u_3 e^{i\phi} \\ u_1 e^{-i\phi} \\ -u_2 \end{bmatrix} \rightarrow \text{Eigenvalues } \in \{\pm 1\}$$

(S)

$$\begin{bmatrix} u_3 \\ u_{||} e^{-i\phi} \end{bmatrix} \begin{bmatrix} u_{||} e^{i\phi} & -u_3 \\ -u_3 & \end{bmatrix} \begin{bmatrix} q \\ b \end{bmatrix} = s \begin{bmatrix} q \\ b \end{bmatrix} \Rightarrow u_3 a + u_{||} e^{i\phi} b = sa$$

$$\Rightarrow u_{||} e^{-i\phi} a - u_3 b = sb$$

$$\Rightarrow u_{||} e^{-i\phi} a = \frac{(s+u_3)}{u_{||} e^{i\phi}} b$$

$$\frac{a}{b} = \frac{(s+u_3)}{u_{||} e^{i\phi}}$$

$\Rightarrow \vec{v} = \begin{pmatrix} s + u_3 \\ u_{||} e^{-i\phi} \end{pmatrix}$, where $u_3 = \frac{m}{\sqrt{(\hbar e k)^2 + m^2}}$

On normalizing,

$$c^2 (u_{||}^2 + (s+u_3)^2) = 1$$

$$c^2 = \frac{1}{u_{||}^2 + u_3^2 + 1 + 2su_3} = \frac{1}{2(1+su_3)}$$

$$c = \frac{1}{\sqrt{2(1+su_3)}}$$

$$\Rightarrow \Psi_{k,s} = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{1+su_3} e^{-i\phi} \\ \frac{\sqrt{1-u_3^2}}{\sqrt{1+su_3}} e^{-i\phi} \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{1+su_3} \\ \frac{(1-su_3)(1+su_3)}{(1+su_3)} \end{pmatrix} e^{i\phi} = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{1+su_3} \\ \sqrt{1-su_3} e^{i\phi} \end{pmatrix}$$

$$\Rightarrow \vec{A}_{k,s} = -i \langle \Psi_{k,s} | \vec{\nabla}_k | \Psi_{k,s} \rangle$$

$$\Rightarrow A_{k,s}^{(R)} = -i \langle \Psi_{k,s} | \partial_k | \Psi_{k,s} \rangle, \quad A_{k,s}^{(\phi)} = -\frac{i}{R} \langle \Psi_{k,s} | \partial_\phi | \Psi_{k,s} \rangle$$

$$\partial_\phi | \Psi_{k,s} \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ -i\sqrt{1-su_3} e^{-i\phi} \end{pmatrix}$$

$$\langle \Psi_{k,s} | = \frac{1}{\sqrt{2}} \left(\sqrt{1+su_3}, \sqrt{1-su_3} e^{i\phi} \right)$$

$$\Rightarrow A_{k,s}^{(\phi)} = -\frac{i}{R} \langle \Psi_{k,s} | \partial_\phi | \Psi_{k,s} \rangle = -\frac{1}{2} \frac{(1-su_3)}{R}$$

$$\partial_k \sqrt{1+s u_3} = \frac{1}{2} \frac{\partial_k u_k}{\sqrt{1+s u_3}}, \quad \partial_k (\sqrt{1-s u_3}) = -\frac{1}{2} \frac{\partial_k u_k}{\sqrt{1-s u_3}}$$

$$\Rightarrow \langle \psi_{k,s} | \partial_k | \psi_{k,s} \rangle = \frac{1}{2} \left[\sqrt{1+s u_3} \frac{\partial_k u_k}{\sqrt{1+s u_3}} + \sqrt{1-s u_3} \left(-\frac{\partial_k u_k}{\sqrt{1-s u_3}} \right) \right] = 0$$

$$\Rightarrow A_{k,s}^{(k)} = 0.$$

(c) Berry phase:

$$P_s = \oint \vec{A}_{k,k} \cdot d\vec{k} = 2\pi k A_{ks}^{(\phi)} = -\pi (1-s u_3)$$

(d) Berry curvature :

$$(\vec{\nabla} \times \vec{A}_{k,s})_z = \frac{1}{\hbar} \frac{\partial}{\partial k} (k A_{k,s}^{(\phi)}) = \frac{1}{\hbar} \left(\frac{\partial}{\partial k} \left(-\frac{k}{2\pi} (1 - s u_z) \right) \right)$$

$$(\vec{\nabla} \times \vec{A}_{k,s})_z = + \frac{s}{2\pi} \frac{\partial u_z}{\partial k}$$

(e) The hamiltonian in Eq. ① is the time reversed hamiltonian of the chern insulator hamiltonian (Eq. ⑦).

$$\vec{d}_k = (\hbar v k_x, \hbar v k_y, m) \xrightarrow{\text{TR}} \vec{d}_k = (\hbar v k_x, -\hbar v k_y, m)$$

TR \rightarrow $i\omega_y \vec{k}$ \rightarrow Modern QM - Sakurai
 Complex conjugation

4)

$$\hat{H} \rightarrow BH_2 = \begin{bmatrix} \hat{n}_k & 0 \\ 0 & \hat{n}_{-k}^* \end{bmatrix} - \textcircled{1}$$

Bernevig, Hughes & Zhang
 Hamiltonian for HgTe/CdTe
 quantum wells

$$\hat{n}_k = A(\sigma_x k_x - \sigma_y k_y) + M_k \sigma_z + D \vec{k}^2 \sigma_0 - \textcircled{2}$$

$$M_k = M + B \vec{k}^2 - \textcircled{3}$$

where A, B, ρ and M are constants characterizing the band structure. Using the unitary transformation:

$$U = \begin{bmatrix} 0 & \sigma_3 \\ -i\sigma_3 & 0 \end{bmatrix} \quad \text{--- (4)}$$

Obtain the Hamiltonian,

$H'_{\text{BH2}} = U H_{\text{BH2}} U^+ \quad \text{--- (5)}$ and show that for

$$\rho = \beta = 0, \quad H'_{\text{BH2}} = \begin{bmatrix} -i\hbar v(\sigma_x \partial_x + \sigma_y \partial_y) + \Delta \sigma_3 & 0 \\ 0 & i\hbar v(\sigma_x \partial_x + \sigma_y \partial_y) - \Delta \sigma_3 \end{bmatrix}$$

Eq. (6) ←

Solution:

Making $k \rightarrow -k$ and taking complex conjugate of Eq. ②,

$$\hat{h}_{-k}^* = [A(-\sigma_n k_n + \sigma_y k_y) + (M + B k^2) \tilde{\gamma}_3 + D k^2 \sigma_0]^*$$

Since $\sigma_n^* = \sigma_n$, $\sigma_y^* = -\sigma_y$, $\tilde{\gamma}_3^* = \tilde{\gamma}_3$ & $\sigma_0^* = \sigma_0$,

$$\hat{h}_{-k}^* = [A^*(-\sigma_n k_n - \sigma_y k_y) + (M^* + B^* k^2) \tilde{\gamma}_3 + D^* k^2 \sigma_0]$$

Thus, Eq. ① is:

$$\hat{H}_{BH2} = \begin{bmatrix} \alpha(\sigma_x k_x - \sigma_y k_y) + (\mu^* \beta^* k^2) \tilde{\gamma}_g + \delta^* k^2 \sigma_0 & 0 \\ 0 & \alpha^* (-\sigma_x k_x - \sigma_y k_y) + (\mu^* \beta^* k^2) \tilde{\gamma}_g + \delta^* k^2 \sigma_0 \end{bmatrix}$$

Now we define the unitary transformation :

$$\hat{U} = \begin{pmatrix} 0 & \tilde{\gamma}_g \\ -i\sigma_y & 0 \end{pmatrix}, \quad \hat{U}^+ = \begin{pmatrix} 0 & (-i\sigma_y)^+ \\ \sigma_g^{*+} & 0 \end{pmatrix} = \begin{pmatrix} 0 & i\sigma_y \\ \tilde{\gamma}_g & 0 \end{pmatrix}$$

$$\hat{U}^+ \hat{U} = \begin{pmatrix} \tilde{\gamma}_g^2 & 0 \\ 0 & (-i\sigma_y)^2 \end{pmatrix}$$

$$(-i\sigma_y)^2 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = I$$

$$(\tilde{\sigma}_y)^2 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = I$$

$$\hat{U}^+ U = \begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix}$$

Now we compute the transformed matrix:

$$H_{BH2}^I = \hat{U}^+ H_{BH2} \hat{U} = \begin{pmatrix} 0 & U_{12} \\ U_{21} & 0 \end{pmatrix} \begin{pmatrix} \hat{n}_k & 0 \\ 0 & \hat{n}_k^* \end{pmatrix} \begin{pmatrix} 0 & U_2^+ \\ U_{21}^+ & 0 \end{pmatrix}$$

$$= \begin{pmatrix} 0 & U_{12} \\ U_{21} & 0 \end{pmatrix} \begin{pmatrix} 0 & \hat{h}_k U_{12}^+ \\ \hat{h}_{-k}^* U_{21}^+ & 0 \end{pmatrix}$$

$$= \begin{pmatrix} U_{12} \hat{h}_{-k}^* U_{21}^+ & 0 \\ 0 & U_{21} \hat{h}_k U_{12}^+ \end{pmatrix}$$

$$U_{12} \hat{h}_{-k}^* U_{21}^+ = \sigma_3 \hat{h}_{-k}^* \sigma_3$$

$$= \tilde{\sigma}_3 \left[\alpha^* (-\sigma_n k_x - \alpha_y k_y) + (\mu^* + \beta^* k^2) \tilde{\sigma}_3 + \delta^* k^2 \alpha_0 \right] \sigma_3$$

Using properties of σ matrices,

$$\sigma_n \tilde{\sigma}_3 = -i \sigma_y, \quad \alpha_y \tilde{\sigma}_3 = i \sigma_x$$

$$\left(\begin{matrix} H \\ BH2 \end{matrix}\right)_{11} = \sigma_3 \left[A^* (i\sigma_y k_x - i\sigma_x k_y) + (M^* + \beta^* k^2) \sigma_0 + Q^* k^2 \sigma_3 \right]$$

$$= A^* \left[\begin{pmatrix} \sigma_x k_x \\ \sigma_y k_y \end{pmatrix} + \begin{pmatrix} \sigma_3 & k_y \\ \sigma_3 & k_y \end{pmatrix} \right] + (M^* + \beta^* k^2) \sigma_3 + Q^* k^2 \sigma_3$$

$$\left(\begin{matrix} H \\ BH2 \end{matrix}\right)_{22} = U_{21} \tilde{U}_K U_{12}^+$$

$$= (-i\sigma_y) \left[A (\sigma_x k_x - \sigma_y k_y) + (M + \beta k^2) \sigma_3 + Q k^2 \sigma_0 \right] (i\sigma_y)$$

$$= (-i\sigma_y) \left[A [(-\sigma_y) k_x - i k_y] + (M + \beta k^2) \sigma_0 + Q k^2 (i\sigma_y) \right]$$

$$= A \left[-\sigma_x k_x - \sigma_y k_y \right] - (M + \beta k^2) \sigma_3 + Q k^2 \sigma_0$$

$$\Rightarrow H'_{BH2} = \begin{bmatrix} A[(\sigma_x k_x) + (\sigma_y k_y)] + (M + \beta^* k^2) \sigma_3 + \phi^* k^2 \sigma_0 & 0 \\ 0 & A[(-\sigma_x k_x) - (\sigma_y k_y)] - (M + \beta k^2) \sigma_3 + \phi k^2 \sigma_0 \end{bmatrix}$$

Setting $\alpha = \hbar v$, $\beta = 0$, $\phi = 0$, $k_x = -i\partial_x$, $k_y = -i\partial_y$:

$$H'_{BH2} = \begin{bmatrix} -i\hbar v(\sigma_x \partial_x + \sigma_y \partial_y) + \Delta \sigma_3 & 0 \\ 0 & i\hbar v(\sigma_x \partial_x + \sigma_y \partial_y) - \Delta \sigma_3 \end{bmatrix}$$

Thus, we get back Eq. 6.