

Universidad Nacional Autónoma de México

OAN–SPM Catalog of Galactic Stellar Clusters

Doctoral Thesis Candidacy

STUDENT:
Ramón Caballero Villegas

TUTOR:
Dr. Raúl Michel Murillo (IA–Ensenada)
Dr. Miguel Ángel Aragón Calvo (IA–Ensenada)

Baja California, Ensenada. (November) 2026

Abstract

The study of stellar clusters (SCs) within the Galaxy serves an important role in understanding the Galaxy's physical and chemical characteristics. SCs are defined as gravitationally-bound groups of stars born from a single, homogeneous molecular cloud, typically composed of a single stellar population. This allows us to measure physical parameters such as a cluster's age, metallicity, distance, and extinction, which can be determined through the photometric data of resolved cluster members. Studies of Galactic clusters have seen a significant increase in recent years with the advent of Gaia, whose unmatched astrometric precision has allowed for the identification of thousands of new stellar clusters, as well as refining membership lists for previously identified clusters.

This doctoral thesis project makes use of Gaia DR3 (and DR4 in the near future) data to update and expand the membership lists from the Hunt & Reffert (2024) and the Unified Cluster Catalog (UCC). By using HDBSCAN we identify an additional population of stars for each cluster, retrieving stars beyond the main clusters' cores and at fainter magnitudes than previously identified. We determine each cluster's physical parameters using isochrone fitting methods using *ASteCA* and `simul_fit` on each clusters' Gaia DR3 data. We compare the derived fundamental cluster parameters between both methods as well as the reported literature values. With these preliminary results we have validated the efficacy of the membership determination pipeline and cluster parameter inference for future use with the rest of the cataloged observations made from the Observatorio Astronómico Universitario – San Pedro Mártir (SPM).

1 Introduction

Given that stars make up the fundamental blocks of the Galaxy, SCs therefore act as important astrophysical laboratories ideal for understanding the Galaxy's dynamics and structure. Stars are well known to be born in cohorts within molecular clouds, dissolving into the interstellar medium to form the field stars that can be seen today. From current star formation theory we know that the youngest of these clusters were formed from a singular burst of star formation, and are therefore referred to as *simple stellar populations* (SSPs), groups of chemically homogeneous stars (Huang et al., 2024). These groups are known as *open clusters* (OCs) due to their loosely-bound and irregular appearance. OCs also tend to exhibit lower total stellar masses, and are primarily located in the Galactic disk (Zwart et al., 2010). OCs are used as tracers for studying the Galactic spiral structure (Dias et al., 2022).

Globular clusters (GCs), on the other hand, are high-density collections of stars, distributed throughout the Galactic halo. GCs are found to be in a more stable virial state, with a much older age distribution as well as higher dynamical timescales, which contributes to their spherically symmetrical appearance. As a result of their age, GC populations tend to be more complex, to the point that they sometimes cannot be properly described as a SSP; chemical abundance variations between stars in a single GC indicates *multiple stellar populations* (MPs) with later-generation stars showing

distinct chemical compositions (Huang et al., 2024). Although they have historically been studied as a separate category from OCs, current models do not discriminate between either OCs or GCs for describing their dynamics and compositions (Krumholz et al., 2019).

The characterization of SCs typically involves two processes: the identification of member stars in order to separate them from the surrounding field stars, and the determination of the SC's fundamental parameters. Of particular relevance to this project is the study of SCs through theoretical isochrone fitting procedures using multiband observations.

1.1 Membership Determination

Many different approaches to discriminate SCs from a population of field stars have been proposed and employed in the literature with varying degrees of success. While early methodologies generally relied on spatial coherence, later surveys and multi-epoch photometric observations allowed for the consideration of kinematic coherence between cluster stars, eventually giving way to the clustering techniques used in more recent works on stars' astrometric solutions (Ramezani, Mondal, et al., 2026). While this project utilizes HDBSCAN as discussed in Section 4, other statistical approaches include a forest fire inspired model (Wei et al., 2025), Gaussian mixture models, K-means, etc. (Pera et al., 2021).

The Gaia mission (Gaia Collaboration et al., 2016) brought about a revolution in the study of Galactic SCs. Its unparalleled astrometric precision of over 1 billion sources in Data Release 1, and over 2 billion sources in the upcoming Data Release 4, allows for cluster identification at unprecedented scales. Systematic searches have revealed hundreds of previously unidentified clusters throughout the Galaxy (Castro-Ginard et al., 2022; Hunt & Reffert, 2021). Additionally, Gaia data has been extensively used to refine cluster membership lists (Hunt & Reffert, 2024; Perren et al., 2023), allowing for studies on clusters' structure (Mendes et al., 2026; Tarricq et al., 2022).

1.2 Cluster Characterization

Photometric studies of SCs involve fitting synthetic isochrones to color–magnitude diagrams (CMDs), and are a common approach to determining a cluster's fundamental parameters (Dias et al., 2021). While spectroscopic data provides a much stronger constraint on stellar chemical abundances (Kos et al., 2025), it is rare to have a significant portion of a cluster's member stars with a reliable measurement of their spectral energy distribution, and as such studies are restricted to use spectro-photometric measurements (in spite of this, narrow-band observations can help, for example, in measuring the blanketing of metallic lines (Paunzen, 2026)). Theoretical isochrone morphology depends on the given age and metallicity, while the cluster's distance and total reddening from interstellar extinction displaces the isochrone's sequence. While traditional isochrone fitting methods seek to minimize a likelihood function such as the distance of each star to the nearest isochrone point (Monteiro et al., 2010), models such as the

Mixture Model for Open Clusters (MiMO; Li et al., 2025) explicitly account for field star contamination, eliminating the need for member star discrimination.

Forward modeling of a cluster can also be performed using codes such as *ASteCA* (further described in Section 5.1), which when combined with tooling designed for solving the inverse problem can yield a cluster's fundamental parameters alongside an estimate of the posterior density function. More recent efforts have also been made with the implementation of machine learning tools on observed CMDs, allowing for quick parameter inference (Cantat-Gaudin et al., 2020; Cavallo et al., 2023; Hunt & Reffert, 2023).

2 Objectives

Although a variety of studies have been done using CMDs from other catalogs, few studies include observations in near-ultraviolet filters due to the demanding photometric conditions (as a result of atmospheric sensitivity) required to obtain reliable measurements for a cluster's metallicity and age. For this project, we will exploit over 20 years of UBVRI observations of over 1700 resolved Galactic clusters (both GCs and OCs), which can be used to provide reliable measurements of the clusters' fundamental parameters with stronger model constraints than is otherwise possible without UBV observations (Kaviraj et al., 2007). The observations from the SPM are complemented with photometry obtained from all-sky surveys when available, crossmatching the Gaia observations with GALEX, 2MASS, PAN-STARRS, SDSS, and WISE.

The project's key objectives are outlined as follows:

- Determine and expand on stellar cluster membership catalogs using high quality astrometric data, focusing on retrieving member stars that were otherwise discarded in previous studies.
- Implement existing tools for determining clusters' physical parameters based on cleaned CMDs using revised membership lists.
- Develop new tools for the efficient and accurate characterization of cluster CMDs.
- Use newly developed tools for determining fundamental and structural parameters of SCs and SC candidates using observations from SPM, supplemented with observations from external catalogs: 2MASS, WISE, GALEX, Gaia, PAN-STARRS, and SDSS when available.

3 Catalog of Observations

For this project we have observations of over 1700 clusters, compiling the results from 298 observation campaigns dating back to the year 2001. Each observational campaign focuses either on the suspected cluster members, or the foreground or background stars. Doing a preliminary cross-match with the *Unified Cluster Catalog* (UCC;

Perren et al., 2023) shows over 300 clusters not identified in the UCC, with some having been identified as asterisms (physically unrelated groups of observed stars). As part of the initial data exploration step of the methodology, a more in-depth review of the published literature will be needed to determine the characterization already performed for each cluster in the sample.

As can be seen in Figure 1, our observations encompass a dense sample of the Solar neighborhood, both spatially as well as characterization quality from the UCC UTI metric, providing a good opportunity to improve the worst-scoring clusters and provide a uniform study of the stellar parameter spatial distributions. Thanks to the photometric conditions at SPM, stars up to a magnitude of $V \sim 21$ can be recorded, allowing for a deeper high quality measurement of SC members. This will allow us to obtain quality photometric measurements for a higher amount of low luminosity members, thus reducing selection bias effects as a result of incomplete sampling of low mass members.




Figure 1: Spatial distribution of observed SCs in the Galaxy. Heliocentric coordinates were calculated using distances determined through an initial manual photometric analysis of each cluster.

3.1 Cluster Sample

As a first test of the membership determination pipeline and preliminary evaluation of the physical characterization of each OC's CMD, we selected four well-studied, well-populated clusters all located within a similar parallax range: *NGC 1245*, *NGC 2311*, *NGC 2420*, and *NGC 2506*. An initial literature distance determination was reported for all but NGC 2311 (Alter et al., 1970), while subsequent studies have been made in an effort to understand each cluster's fundamental parameters (Cavallo et al., 2023; Hunt & Reffert, 2024; Li et al., 2025) and use them—as part of a larger sample of Galactic SCs—to understand and trace the Galactic structure (Cantat-Gaudin et al., 2020; Kharchenko et al., 2016).

Due to the technical limitations of the cluster characterization tooling available at this time we opted to only use each cluster's Gaia DR3 observations, as this would have the most amount of information available and whose broadband passbands cover a portion of the UV spectrum, providing some sensitivity to metallic lines and thus increasing sensitivity to a cluster's global metallicity, albeit without the sensitivity a narrow-band passband might otherwise provide (Paunzen, 2026).

4 Membership Determination

In order to properly characterize a SC, its member stars must be properly identified to avoid including field stars not bound to the cluster in question for the ensuing CMD analysis. As a result, cluster membership determination algorithms have received a large amount of attention in recent years (Ramezani, Balciráková, et al., 2026). For this project we implement a membership determination pipeline using HDBSCAN, an improvement over the DBSCAN clustering algorithm by assembling minimum spanning trees and subsequently compacting them into a set of "groupings," from which stellar clusters are identified. HDBSCAN has been used in recent works with a high rate of success in both identifying previously unknown SCs as well as expanding on known clusters' membership lists (Hunt & Reffert, 2021; Sánchez-Sanjuán et al., 2024; Tarricq et al., 2022).

4.1 Gaia DR3 Data Selection

Our HDBSCAN pipeline operates on the Gaia DR3 (GDR3) (Gaia Collaboration et al., 2023) 5D astrometric phase space $(\alpha, \delta, \omega, \mu_\alpha \cos\delta, \mu_\delta)$; radial velocities are not included in order to maximize our sample size, as only a few stars within the GDR3 catalog include radial velocity measurements. For each of the four clusters analyzed for this work we selected all stars within a conical field centered on the cluster reported center. The search radius was taken as a multiple of the literature reported cluster radius: $\texttt{max}\left(10 \cdot R_{50,\ \mathrm{UCC}},\ 2 \cdot R_{\mathrm{tot},\ \mathrm{HR24}}\right)$, where $R_{50,\ \mathrm{UCC}}$ is the half-member radius reported in the UCC, and $R_{\mathrm{tot},\ \mathrm{HR24}}$ is the tidal radius reported by Hunt & Reffert, 2024.

Each astrometric parameter was normalized to avoid the different scales of each parameter to overpower the other four dimensions, in particular to avoid issues with each




Figure 2: Gaia DR3 fields obtained using each OC's reported coordinates, with concentric rings denoting the search fields radii, along with the corresponding Gaia DR3 CMD. The upper panels of each OC overlays the reported cluster members in the UCC in blue, while the lower panels show the HR24 members overlaid in red.

field's parallax range, as its scale is order of magnitudes smaller than the other parameters. This was done using the `RobustScaler` class from `scikit-learn` (Pedregosa et al., 2011), which removes the data median and scales the data with the quantile range to avoid scaling problems due to outliers. We performed a quality cut on our initial field sample by retrieving all stars in the field that satisfy `fidelity_v2` ≥ 0.5, as published and recommended by Rybizki et al., 2022, where the `fidelity_v2` flag quantifies the reliability of the Gaia EDR3 astrometric solution. It has been previously used with success to discriminate cluster and star formation groups from a population

of field stars without performing a hard cut on either magnitude or RUWE (Hunt & Reffert, 2023; Rottensteiner et al., 2026).

4.2 HDBSCAN Clustering Pipeline

Field star discrimination is performed using HDBSCAN (Campello et al., 2013), a hierarchical density-based machine learning algorithm adept at finding clusters of data, whose hierarchical approach improves upon its predecessor DBSCAN's performance by not being biased towards a prescribed density.

HDBSCAN takes two input parameters: a minimum number of neighboring points for a given cluster center (m_{pts}) and a minimum cluster size (m_{cls}) for a given *data cluster* (for the purposes of this paper, the term *data cluster* is used to denote a grouping of data, but not necessarily a physical stellar cluster) is considered significant and not labeled as noise in the data. The membership determination pipeline in its current state runs the HDBSCAN clustering algorithm for a grid of $10 \leq m_{cls} \leq 5000$ in steps of 20. m_{pts} was fixed to a value of 2 to reduce compute time, while the choice of $m_{pts} = 2$ was motivated to allow HDBSCAN to remain sensitive to more sparse clusters; when applied to more populated clusters such as NGC 1245, NGC 2311, NGC 2420, and NGC 2506, the hierarchical nature of HDBSCAN eventually collapses smaller groupings to a single cluster.

4.3 Membership Determination Results

The HDBSCAN data clusters are subsequently processed to find the cluster of interest, as HDBSCAN will almost always return more than one singular cluster per field. The most correct cluster is identified by its *average centroid linkage distance* (D_c) to the literature cluster:

$$D_c(H, L) = \frac{1}{|H| + |L|} \left\{ \sum_{\vec{x} \in H} d(\vec{x}, \vec{c}_L) + \sum_{\vec{y} \in L} d(\vec{y}, \vec{c}_H) \right\} \tag{1}$$

Where H is the data cluster identified from the HDBSCAN pipeline, L is the cluster composed of the members reported in the literature (taken from Hunt & Reffert, 2024 (henceforth HR24) when available, else from the UCC (Perren et al., 2023)), $\vec{x}$ and $\vec{y}$ are the normalized 5D astrometric parameters of the members in clusters H and L respectively, the function d returns the Euclidean distance between two points, and $\vec{c}_H$ and $\vec{c}_L$ are the center positions of clusters H and L respectively. The data cluster H with the lowest D_c to the literature cluster L is taken as the new solution.

Figure 3 shows the results of the HDBSCAN pipeline on each of the four clusters evaluated, having accepted only the members with a membership probability greater than 50%. In 3 of the 4 clusters the pipeline retrieves both an adequate overlap with the membership lists in the HR24 catalog, while identifying a significant amount of new member stars, both in the spatial outskirt regions of each cluster as well as an appreciable locus of new stars (as is the case of NGC 2311) of faint stars. NGC 2420 presents a case where the HDBSCAN pipeline did not retrieve an adequate amount




Figure 3: Matched clusters obtained by HDBSCAN for each of the evaluated OCs. Their spatial distribution can be seen in columns 1 and 3, where column 3 shows the KDE distribution of the cluster members in the UCC catalog (cyan), HR24 (red), and the recovered HDBSCAN cluster (magenta). Column 2 shows the CMD in the Gaia bands, and column 4 shows the clusters' distribution in the proper motion space. In all but one of the clusters we manage to retrieve members further than the previously reported furthest members from the core. The average centroid linkage distance (D_c) is computed on the scaled astrometric parameters of each member star.

of stars when compared with the literature, and, perhaps more significantly, did not identify any new member stars, likely due to the search field size when compared to the search radius used by HR24, processing multiple Gaia DR3 HEALPix pixels at once in search of OCs. As a result, for the subsequent photometric analysis, we used the

HR24 membership catalog for NGC 2420, while for the rest of the clusters we used the list obtained from the HDBSCAN pipeline.

5 Photometric Analysis

Using the membership list from the HDBSCAN pipeline, we performed a photometric analysis of each OC to determine their physical parameters: metallicity (Z; calculated using [Fe/H]), age ($\log t$), distance given as the distance modulus (d_m), visual extinction (A_V), and Galactic extinction curve (R_V). This was done using each cluster's computed Gaia CMDs as a first approximation and fitting PARSEC theoretical isochrones (Bressan et al., 2012), where the fit was done using two different approaches to explore the existing tooling available in the literature.

5.1 ASteCA

The *Automated Stellar Cluster Analysis* (ASteCA) package (Perren et al., 2015) is an open-source package for cluster analysis. ASteCA is organized into several modules for handling cluster membership probabilities, isochrone handling, synthetic cluster generation from theoretical isochrones, observation calibrations for comparing synthetic clusters with real observations, and likelihood calculations for comparing synthetic CMDs with a synthetic cluster's CMD. Since each clusters' stellar members were already determined from the HDBSCAN pipeline, we only made use of the `Synthetic` module to generate synthetic clusters calibrated on the Gaia member observations.

An *approximate Bayesian computation* (ABC) combined with *sequential Monte-Carlo* (ABC-SMC) approach was employed for parameter inference with ASteCA using the `pyABC` package (Schälte et al., 2022). Unlike other methods of parameter inference, ABC-SMC does not depend on the evaluation of a likelihood function, but rather only on a measurement of a distance metric of the generated model (in this case, the synthetic cluster created from a parameter vector sampled from a given target distribution) and the observed data (the observational Gaia DR3 CMD from the member stars identified from the HDBSCAN pipeline), denoted as ϵ.

In addition to the main intrinsic parameters that dictate the general shape of a theoretical isochrone, unresolved binary systems can have an impact on a cluster's CMD, appearing as an excessively red star and manifesting as a broadening of the cluster's main sequence (Li et al., 2020). When generating a synthetic cluster, ASteCA generates binary systems with the given probability ($P_b(m_1)$) as a function of the primary's mass (m_1) with free parameters α and β:

$$P_b(m_1) = \alpha + \beta \left(\frac{m_1}{1.4 + m_1} \right) \tag{2}$$

5.1.1 ABC Sampling

Parameter inference with `pyABC` was performed by sampling from a uniform distribution for all cluster intrinsic and extrinsic parameters. Parameter ranges can be seen

in Table 1. Although this imposes an initial restriction on the parameter space initially explored by `pyABC`, these are not hard limits, as proposal parameter vectors carry the sampler beyond the limits defined by the prior distributions (Sisson et al., 2007).

Parameter	**Minimum**	**Maximum**
$\log t$	7.0	9.5
Z	0.005	0.0197
d_m	7.0	10.5
A_V	0.0	3.0
R_V	2.0	4.0
α	0.0	0.2
β	0.8	0.99
$E(B-V)$	0	4.0

Table 1: Prior parameter sample ranges.

The logarithmic Poisson likelihood ratio (PLR) ($\log p$) defined within ASteCA is used as the distance metric for `pyABC`:

$$\log p \approx \sum_{i}^{B} \left[\log \Gamma \left(n_i + m_i + \frac{1}{2} \right) - \log \Gamma \left(m_i + \frac{1}{2} \right) \right] \tag{3}$$

The PLR is calculated on a 2D histogram of the Gaia DR3 CMD composed of B bins, where n_i and m_i are the number of observed and synthetic stars in bin i, respectively. Each cluster was sampled until either $\epsilon \leq 0.01$, $\Delta\epsilon \leq 10^{-9}$, or 3000 iterations were reached.

5.2 `simul_fit`

As an additional comparison, each cluster's Gaia CMD was also processed using `simul_fit` (speletry, 2026) to simultaneously fit theoretical PARSEC isochrones to the cluster Gaia CMDs using a χ^2 optimization technique in a Bayesian grid-based sampling scheme (Al et al., 2026), where χ^2 is the sum of the distances of each cluster member from the nearest point on the isochrone. The $\log t$, A_V, and d_m parameter ranges for the sample grid were obtained from the HR24 catalog, taking $R_V = 3.1$. Figure 4 shows the sampling results for NGC_2506, showing an estimation of the posterior distribution function.

5.3 CMD analysis

We compare the parameters obtained from both ASteCA and `simul_fit` with the reported parameters in the literature, as well as with a visual isochrone fit performed on the members' Gaia DR3, Johnson-Cousins *UBVRI*, Strömgren *ubv*, and Pan-STARRS *gri* observations. A detailed report with each cluster's fundamental parameters can be seen in Table 2. The resulting CMDs with overlaid isochrone sequences can be seen in




Figure 4: Corner plot for NGC_2506 from the weighted sampling done by `simul_fit` from the parameter grid exploration. `simul_fit` assigns a likelihood weight of $P \propto -\Delta\chi^2/2$ to each sample. The median is marked by the blue line, followed by the 16th and 84th percentile uncertainties.

Figure 5, where the parameters determined in this work result in an appreciably better fit to the observed data.

Using ASteCA we were able to assess the impact of binary systems on the cluster CMD; while the resulting ASteCA isochrone lies towards the blue edge of each of the main sequence, the binary probability of each star as reported by ASteCA reveals the extent that member binarity broadens the main sequence, as seen in Figure 6.




Figure 5: Gaia CMDs for each of the four clusters analyzed in this work, showing the best fit PARSEC isochrones using the parameters reported in the literature (UCC, HR24), best fits from ASteCA and `simul_fit`, and a visual isochrone fit using SPM observations.

Cluster	Parameter	UCC	HR24	Manual SPM	ASteCA	`simul_fit`
NGC 1245	$\log t$	9.0298	$8.9914^{+0.1659}_{-0.1839}$	9.2400	9.0896 ± 0.5626	$8.9000^{+0.0019}_{-0.0043}$
	Z	0.0181	0.0207 ($Z_\odot$)	0.0088	0.0153 ± 0.0031	$0.0191^{+0.0002}_{-0.0002}$
	d_m	12.3856	$12.6432^{+0.2072}_{-0.1601}$	12.0000	12.3966 ± 0.7244	$12.6724^{+0.0097}_{-0.0175}$
	$E(B-V)$	0.2613	$0.1757^{+0.0574}_{-0.0566}$	0.2800	2.1026 ± 1.1672	$0.3292^{+0.0036}_{-0.0083}$
NGC 2311	$\log t$	8.3010	$7.9845^{+0.2190}_{-0.1901}$	7.9000	8.3497 ± 0.6713	$7.5000^{+0.0017}_{-0.0038}$
	Z	0.0179	0.0207 ($Z_\odot$)	0.0125	0.0154 ± 0.0044	$0.0208^{+0.0002}_{-0.0003}$
	d_m	11.5799	$11.5365^{+0.1470}_{-0.1616}$	11.1000	11.5886 ± 0.8691	$11.9330^{+0.0112}_{-0.0178}$
	$E(B-V)$	0.3161	$0.2726^{+0.0544}_{-0.0673}$	0.3100	1.9979 ± 1.1505	$0.4867^{+0.0040}_{-0.0071}$
NGC 2420	$\log t$	9.3049	$9.3523^{+0.1731}_{-0.1656}$	9.3700	9.4016 ± 0.6724	$9.4000^{+0.0019}_{-0.0040}$
	Z	0.0130	0.0207 ($Z_\odot$)	0.0143	0.0102 ± 0.0029	$0.0174^{+0.0002}_{-0.0004}$
	d_m	12.0327	$11.7873^{+0.2266}_{-0.1924}$	12.0000	12.0648 ± 0.7509	$11.9806^{+0.0089}_{-0.0238}$
	$E(B-V)$	0.0290	$0.0098^{+0.0121}_{-0.0098}$	0.0300	2.0974 ± 1.1330	$0.0000^{+0.0033}_{-0.0069}$
NGC 2506	$\log t$	9.2101	$9.3819^{+0.1623}_{-0.1647}$	9.3900	9.3000 ± 0.4787	$9.3001^{+0.0019}_{-0.0038}$
	Z	0.0113	0.0207 ($Z_\odot$)	0.0095	0.0065 ± 0.0027	$0.0101^{+0.0002}_{-0.0004}$
	d_m	12.5460	$12.5030^{+0.2218}_{-0.2199}$	12.4500	12.5887 ± 0.5474	$12.5393^{+0.0101}_{-0.0176}$
	$E(B-V)$	0.0806	$0.0104^{+0.0116}_{-0.0104}$	0.0600	1.9323 ± 1.1549	$0.0501^{+0.0048}_{-0.0075}$

Table 2: Fundamental parameters for each cluster, comparing the literature values with the ASteCA and `simul_fit` derived parameters. The resulting isochrones can be seen in Figure 5.




Figure 6: The color map on the Gaia CMD for each cluster, representing the probability of each member star of being an unresolved binary system. This probability was derived by ASteCA from the best fit α and β parameters as described in Section 5.1.

References

Al, Z., Karataş, Y., Murillo, R. M., Bonatto, C., Güneş, O., Oralhan, İ. A., & Ülgen, E. K. (2026, May 18). *CCD () photometry of the open cluster NGC 6793 and its dynamical evolution* [Comment: 24 pages and 17 figures, accepted for publishing at Journal of Astrophysics and Astronomy JAA]. arXiv: 2605.18665 [astro-ph.GA].

Alter, G., Balazs, B., Ruprecht, J., & Vanysek, J. (1970, January 1). *Catalogue of star clusters and associations*. Retrieved November 12, 2026, from https://ui.adsabs.harvard.edu/abs/1970csca.book.....A

Bressan, A., Marigo, P., Girardi, Léo., Salasnich, B., Dal Cero, C., Rubele, S., & Nanni, A. (2012). PARSEC: Stellar tracks and isochrones with the PAdova and TRieste Stellar Evolution Code. *Monthly Notices of the Royal Astronomical Society*, *427*, 127–145.

Campello, R. J. G. B., Moulavi, D., & Sander, J. (2013). Density-Based Clustering Based on Hierarchical Density Estimates. In J. Pei, V. S. Tseng, L. Cao, H. Motoda, & G. Xu (Eds.), *Advances in Knowledge Discovery and Data Mining* (pp. 160–172). Springer.

Cantat-Gaudin, T., Anders, F., Castro-Ginard, A., Jordi, C., Romero-Gómez, M., Soubiran, C., Casamiquela, L., Tarricq, Y., Moitinho, A., Vallenari, A., Bragaglia, A., Krone-Martins, A., & Kounkel, M. (2020). Painting a portrait of the Galactic disc with its stellar clusters. *Astronomy and Astrophysics*, *640*, A1.

Castro-Ginard, A., Jordi, C., Luri, X., Cantat-Gaudin, T., Carrasco, J. M., Casamiquela, L., Anders, F., Balaguer-Núñez, L., & Badia, R. M. (2022). Hunting for open clusters in Gaia EDR3: 628 new open clusters found with OCfinder. *Astronomy & Astrophysics*, *661*, A118.

Cavallo, L., Spina, L., Carraro, G., Magrini, L., Poggio, E., Cantat-Gaudin, T., Pasquato, M., Lucatello, S., Ortolani, S., & Schiappacasse-Ulloa, J. (2023). Parameter Estimation for Open Clusters using an Artificial Neural Network with a QuadTree-based Feature Extractor. *The Astronomical Journal*, *167*(1), 12.

Dias, W. S., Monteiro, H., Caetano, T., Hickel, G., Alves, G., Sacchi, M., Lépine, J. R. D., & Amarinho, N. (2022). 58 open clusters investigated with UBVRI and Gaia eDR3 data. *Monthly Notices of the Royal Astronomical Society*, *512*, 4464–4483.

Dias, W. S., Monteiro, H., Moitinho, A., Lépine, J. R. D., Carraro, G., Paunzen, E., Alessi, B., & Villela, L. (2021). Updated parameters of 1743 open clusters based on Gaia DR2. *Monthly Notices of the Royal Astronomical Society*, *504*, 356–371.

Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., Brown, A. G. A., Vallenari, A., Babusiaux, C., Bailer-Jones, C. A. L., Bastian, U., Biermann, M., Evans, D. W., Eyer, L., Jansen, F., Jordi, C., Klioner, S. A., Lammers, U., Lindegren, L., Luri, X., Mignard, F., Milligan, D. J., ... Costigan, G. (2016). The Gaia mission. *Astronomy and Astrophysics*, *595*, A1.

Gaia Collaboration, Vallenari, A., Brown, A. G. A., Prusti, T., de Bruijne, J. H. J., Arenou, F., Babusiaux, C., Biermann, M., Creevey, O. L., Ducourant, C., Evans, D. W., Eyer, L., Guerra, R., Hutton, A., Jordi, C., Klioner, S. A., Lammers,

U. L., Lindegren, L., Luri, X., ... De March, R. (2023). Gaia Data Release 3. Summary of the content and survey properties. *Astronomy and Astrophysics*, *674*, A1.

Huang, R., Tang, B., Li, C., Geisler, D., Mateo, M., Song, Y.-Y., Baumgardt, H., Carballo-Bello, J. A., Wang, Y., Nie, J., Dias, B., & Fernández-Trincado, J. G. (2024). Driving factors behind multiple populations. *Science China Physics, Mechanics & Astronomy*, *67*(5), 259513.

Hunt, E. L., & Reffert, S. (2021). Improving the open cluster census. I. Comparison of clustering algorithms applied to Gaia DR2 data. *Astronomy and Astrophysics*, *646*, A104.

Hunt, E. L., & Reffert, S. (2023). Improving the open cluster census - II. An all-sky cluster catalogue with Gaia DR3. *Astronomy & Astrophysics*, *673*, A114.

Hunt, E. L., & Reffert, S. (2024). Improving the open cluster census. III. Using cluster masses, radii, and dynamics to create a cleaned open cluster catalogue. *Astronomy and Astrophysics*, *686*, A42.

Kaviraj, S., Rey, S.-C., Rich, R. M., Yoon, S.-J., & Yi, S. K. (2007). Better age estimation using ultraviolet–optical colours: Breaking the age–metallicity degeneracy. *Monthly Notices of the Royal Astronomical Society: Letters*, *381*(1), L74–L78.

Kharchenko, N. V., Piskunov, A. E., Schilbach, E., Röser, S., & Scholz, R.-D. (2016). Global survey of star clusters in the Milky Way. V. Integrated JHKS magnitudes and luminosity functions. *Astronomy and Astrophysics*, *585*, A101.

Kos, J., Buder, S., Beeson, K. L., Bland-Hawthorn, J., De Silva, G. M., D'Orazi, V., Freeman, K., Hayden, M., Lewis, G. F., Lind, K., Martell, S. L., Sharma, S., Zucker, D. B., Zwitter, T., Da Costa, G. S., de Grijs, R., Howell, M., McKenzie, M., Nordlander, T., ... Traven, G. (2025). The GALAH survey: Improving chemical abundances using star clusters. *Astronomy and Astrophysics*, *703*, A104.

Krumholz, M. R., McKee, C. F., & Bland-Hawthorn, J. (2019). Star Clusters Across Cosmic Time. *Annual Review of Astronomy and Astrophysics*, *57*, 227–303.

Li, L., Shao, Z., Li, Z., & Fu, X. (2025, October 27). *The MiMO Catalog: Physical Parameters and Stellar Mass Functions of 1,232 Open Clusters from Gaia DR3* [Comment: Accepted in AJ]. arXiv: 2510.23374 [astro-ph.GA].

Li, L., Shao, Z., Li, Z.-Z., Yu, J., Zhong, J., & Chen, L. (2020). Modeling Unresolved Binaries of Open Clusters in the Color-Magnitude Diagram. I. Method and Application of NGC 3532. *The Astrophysical Journal*, *901*, 49.

Mendes, D. C., Santiago, V. A., & Angelo, M. d. S. (2026). BATEIA-1: An Approach for determining the structural parameters of open star clusters via machine and deep learning. *Astronomy and Computing*, *56*, 101119.

Monteiro, H., Dias, W. S., & Caetano, T. C. (2010). Fitting isochrones to open cluster photometric data: A new global optimization tool. *Astronomy and Astrophysics*, *516*, A2.

Paunzen, E. (2026). The calibration of Gaia DR3 synthetic Strömgren-Crawford photometry. *Monthly Notices of the Royal Astronomical Society*, *548*, stag544.

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A.,

Cournapeau, D., Brucher, M., Perrot, M., & Duchesnay, É. (2011). Scikit-learn: Machine Learning in Python. *Journal of Machine Learning Research*, *12*(85), 2825–2830. Retrieved November 6, 2026, from http://jmlr.org/papers/v12/pedregosa11a.html

Pera, M. S., Perren, G. I., Moitinho, A., Navone, H. D., & Vazquez, R. A. (2021). pyUPMASK: An improved unsupervised clustering algorithm. *Astronomy and Astrophysics*, *650*, A109.

Perren, G. I., Vázquez, R. A., & Piatti, A. E. (2015). ASteCA: Automated Stellar Cluster Analysis. *Astronomy & Astrophysics*, *576*, A6.

Perren, G. I., Pera, M. S., Navone, H. D., & Vázquez, R. A. (2023). The Unified Cluster Catalogue: Towards a comprehensive and homogeneous data base of stellar clusters. *Monthly Notices of the Royal Astronomical Society*, *526*, 4107–4119.

Ramezani, T., Balciráková, J., Mondal, P., Neumannova, K., Paunzen, E., & Szász, G. (2026, September 1). *Cluster Membership Probabilities: A Review of Methods and Gaia Applications*.

Ramezani, T., Mondal, P., Neumannova, K., Paunzen, E., Supikova, J., & Szasz, G. (2026, July 1). *The different methods to calculate cluster membership probabilities*.

Rottensteiner, A., Ratzenböck, S., Alves, J., Meingast, S., & Hutschenreuter, S. (2026, August 1). *A comprehensive cluster census of Orion. An application of the Significance Mode Analysis (SigMA) algorithm*.

Rybizki, J., Green, G. M., Rix, H.-W., El-Badry, K., Demleitner, M., Zari, E., Udalski, A., Smart, R. L., & Gould, A. (2022). A classifier for spurious astrometric solutions in Gaia eDR3. *Monthly Notices of the Royal Astronomical Society*, *510*, 2597–2616.

Sánchez-Sanjuán, S., Hernández, J., Pérez-Villegas, Á., Román-Zúñiga, C., Aguilar, L., Ballesteros-Paredes, J., & Bonilla-Barroso, A. (2024). Kinematic study of the Orion Complex: Analysing the young stellar clusters from big and small structures. *Monthly Notices of the Royal Astronomical Society*, *534*(3), 2566–2584.

Schälte, Y., Klinger, E., Alamoudi, E., & Hasenauer, J. (2022). pyABC: Efficient and robust easy-to-use approximate Bayesian computation. *Journal of Open Source Software*, *7*(74), 4304.

Sisson, S. A., Fan, Y., & Tanaka, M. M. (2007). Sequential Monte Carlo without likelihoods. *Proceedings of the National Academy of Sciences*, *104*(6), 1760–1765.

speletry. (2026). Speletry/simul_fit.py: Simul_fit v1.0.0 - Initial Stable Release.

Tarricq, Y., Soubiran, C., Casamiquela, L., Castro-Ginard, A., Olivares, J., Miret-Roig, N., & Galli, P. a. B. (2022). Structural parameters of 389 local open clusters. *Astronomy & Astrophysics*, *659*, A59.

Wei, X., Chen, J., Zhang, S., He, F., Zhao, Y., He, X., Fang, Y., Chen, X., & Yang, H. (2025). Forest Fire Clustering: A Novel Tool for Identifying Star Members of Clusters. *The Astronomical Journal*, *169*, 115.

Zwart, S. F. P., McMillan, S. L. W., & Gieles, M. (2010). Young Massive Star Clusters. *Annual Review of Astronomy and Astrophysics*, *48*, 431–493.