

# M374M

## Homework #9


Logan.

Sections 3.3, 3.4: p.189 print [p.328 ut ebook] (#1<sup>1</sup>g). p.199 print [p.348 ut ebook] (#3<sup>1</sup>).

<sup>1</sup>Assume boundary layer is on left side of interval and find leading-order composite approximation.

Mini-project.

Due to surface tension, a liquid-gas interface will rise up or dip down at a solid boundary to form a meniscus, which is responsible for the so-called capillary effect in narrow tubes. In the planar case, the shape of the interface or meniscus curve  $y(x)$  is described by



$$\frac{\sigma}{\rho g} \frac{d^2 y}{dx^2} = y \left[ 1 + \left( \frac{dy}{dx} \right)^2 \right]^{3/2}, \quad 0 \leq x \leq L \quad (1)$$

$$\frac{dy}{dx}(0) = 0, \quad \frac{dy}{dx}(L) = \tan \gamma$$

where  $\rho$  is the density of the liquid,  $g$  is gravitational acceleration, and  $\sigma$  and  $\gamma$  are the surface tension and wetting angle constants associated with the interface. Here we develop an approximate solution of (1) under the assumption that  $0 < \frac{\sigma}{\rho g} \ll L^2$ . For this problem we expect a boundary layer at  $x = L$ , and we will see that  $\frac{dy}{dx}$  is regular, but  $\frac{d^2 y}{dx^2}$  behaves singularly.

- (a) Introduce the dimensionless variables  $h = y/L$  and  $s = x/L$  and show that (1) can be written in the following form for an appropriate parameter  $0 < \varepsilon \ll 1$ .

$$\varepsilon \frac{d^2 h}{ds^2} = h \left[ 1 + \left( \frac{dh}{ds} \right)^2 \right]^{3/2}, \quad 0 \leq s \leq 1. \quad \frac{dh}{ds}(0) = 0, \quad \frac{dh}{ds}(1) = \tan \gamma. \quad (2)$$

- (b) Find the leading-order outer approximation.
- (c) For the inner approximation, consider the change of variables  $\tau = \varepsilon^{-\alpha}(s - 1)$  and  $u = \varepsilon^{-\beta}h$ . Show that  $\alpha = \frac{1}{2}$  and  $\beta = \frac{1}{2}$  leads to the regular system

$$\frac{d^2 u}{d\tau^2} = u \left[ 1 + \left( \frac{du}{d\tau} \right)^2 \right]^{3/2}, \quad \tau \leq 0. \quad \frac{du}{d\tau}(0) = \tan \gamma. \quad (3)$$

This system is difficult to solve. To simplify it, we expect that  $\frac{du}{d\tau}$  varies between zero and  $\tan \gamma$  in the boundary layer; hence using an average of  $\frac{1}{2} \tan \gamma$ , we replace  $1 + \left( \frac{du}{d\tau} \right)^2$  by  $1 + \frac{1}{4} \tan^2 \gamma$  in the differential equation. Use this simplification to find an inner approximation.

- (d) By matching the results from (b) and (c) construct a leading-order composite approximation for  $h(s)$ , and hence for the meniscus curve  $y(x)$ . Use the approximation to find an expression for the meniscus height  $\ell = y(L)$ . Show that  $\frac{dy}{dx}$  remains bounded at all points as  $\varepsilon \downarrow 0$ , but that  $\frac{d^2 y}{dx^2}$  grows unbounded at  $x = L$  as  $\varepsilon \downarrow 0$ .
- (e) Download the Matlab files `program9.m`, `meniscusODE.m` and `meniscusBC.m` from the course webpage and simulate the equations in (1) for  $\sigma = 0.05$ ,  $\rho = 1000$ ,  $g = 10$ ,  $L = 0.07$  and  $\gamma = \frac{\pi}{3}$  in kg-m-s units. Superimpose plots of the curve  $y(x)$  produced by the simulation and by your approximation from (d). Is your prediction of the meniscus height  $\ell$  in (approximate) agreement with the simulation?

Turn in: responses to (a)–(e).