

current-current two-point functions

$$\begin{aligned}\Pi_{\mu\nu}(q) &= i \int d^4x e^{iqx} \langle 0 | T \left\{ \mathcal{J}_{ij}^\mu(x) \mathcal{J}_{ij}^{\nu\dagger}(0) \right\} \rangle \\ &= \left(q_\mu q_\nu - q^2 g_{\mu\nu} \right) \Pi_{ij,\mathcal{J}}^{(1)}(q^2) + q^\mu q^\nu \Pi_{ij,\mathcal{J}}^{(0)}(q^2)\end{aligned}\quad (1)$$

Inclusive ratio:

$$R_\tau = \frac{\Gamma[\tau^- \rightarrow \nu_\tau \text{ hadrons}]}{\Gamma[\tau^- \rightarrow \nu_\tau e^- \bar{\nu}_e]}\quad (2)$$

$$R_\tau = 12\pi S_{EW} \int_0^{m_\tau} \frac{ds}{m_\tau^2} \left(1 - \frac{s}{m_\tau^2} \right) \left[\left(1 + 2 \frac{s}{m_\tau^2} \right) \text{Im} \Pi^{(1)}(s) + \text{Im} \Pi^{(0)}(s) \right] \quad (3)$$

$$\Pi^{(J)}(s) \equiv |V_{uq}|^2 \left(\Pi_{ud,V}^{(J)} + \Pi_{ud,A}^{(J)}(s) \right) \quad (4)$$

OPE:

$$\Pi_{OPE}^{(1+0)}(s) = \sum_{k=0}^{\infty} \frac{C_{2k}(s)}{(-s)^k} \quad (5)$$

$$\mathcal{O}_{4,V/A} = \frac{1}{12} \left[1 - \frac{11}{18} a_s \right] \langle a_s G G \rangle + \left[1 + \frac{\pm 36 - 23}{27} a_s \right] \langle (m_u + m_d) \bar{q} q \rangle \quad (6)$$

D=6: Anomalous Dimension Matrix (V-A):

$$\hat{\gamma}_{Q-}^{(1)} = \begin{pmatrix} -\frac{3N_C}{2} + \frac{3}{N_C} & -\frac{3C_F}{2N_C} \\ -3 & 0 \end{pmatrix} \quad (7)$$

Anomalous Dimension Matrix (V+A):

$$\gamma_{Q_+}^{(1)} = \begin{pmatrix} -\frac{3}{N_C} & \frac{3C_F}{2N_C} & -\frac{1}{3N_C} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 3 & 0 & \frac{2}{3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{N_f}{3} - \frac{3N_c}{4} - \frac{1}{3N_c} & \frac{3N_c}{4} - \frac{3}{N_c} & 0 & 0 & 0 & 0 & 0 & 0 \\ \frac{3}{2} + \frac{3}{2N_c} & -\frac{3C_F}{2N_c} & \frac{3N_c}{4} + \frac{3}{2} - \frac{11}{6N_c} & -\frac{3N_c}{4} + \frac{3}{2} + \frac{3}{2N_c} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{11}{3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

non-diagonal V/A current

$$j_\mu^V(x) = (\bar{u}\gamma_\mu d)(x) \quad (9)$$

$$j_\mu^A(x) = (\bar{u}\gamma_\mu\gamma_5 d)(x) \quad (10)$$

Contour integral:

$$\int_{s_{th}}^{s_0} \frac{ds}{s_0} \omega(s) \text{Im} \Pi_{V/A}(s) = \frac{i}{2} \oint_{|s|=s_0} \frac{ds}{s_0} \omega(s) \Pi_{V/A}(s) \quad (11)$$

adler function

$$D(s) \equiv -s \frac{d\Pi^{PT}}{ds} = \frac{1}{12\pi^2} \sum_{n=0}^{\infty} a_{\mu}^n \sum_{k=1}^{n+1} k c_{nk} \ln \left(\frac{-s}{\mu^2} \right) \quad (12)$$

contour integral in terms of adler

$$A^{\omega,PT} = \frac{i}{2s_0} \oint_{|s|=s_0} \frac{ds}{s} [W(s) - W(s_0)] D(s) \quad (13)$$

spectral function:

$$\rho(s) \equiv \frac{1}{\pi} \text{Im } \Pi(s) \quad (14)$$

normalized invariant mass-squared distribution

$$\left(\frac{1}{N_{V/A}} \right) \left(\frac{dN_{V/A}}{ds} \right) \quad (15)$$

aleph:

$$\begin{aligned} v_1(s) &\equiv \frac{m_\tau^2}{6|V_{ud}|^2 S_{EW}} \frac{B(\tau^- \rightarrow V^- \nu_\tau)}{B(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau)} \frac{dN_V}{N_V ds} \left[\left(1 - \frac{s}{m_\tau^2} \right)^2 \left(1 + \frac{2s}{m_\tau^2} \right) \right]^{-1} \\ a_1(s) &\equiv \frac{m_\tau^2}{6|V_{ud}|^2 S_{EW}} \frac{B(\tau^- \rightarrow A^- \nu_\tau)}{B(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau)} \frac{dN_A}{N_A ds} \left[\left(1 - \frac{s}{m_\tau^2} \right)^2 \left(1 + \frac{2s}{m_\tau^2} \right) \right]^{-1} \\ a_0(s) &\equiv \frac{m_\tau^2}{6|V_{ud}|^2 S_{EW}} \frac{B(\tau^- \rightarrow \pi^- \nu_\tau)}{B(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau)} \frac{dN_A}{N_A ds} \left(1 - \frac{s}{m_\tau^2} \right)^2 \end{aligned} \quad (16)$$

$$\begin{aligned} \text{Im } \Pi_{\bar{u}d,V}^{(1)}(s) &= \frac{1}{2\pi} v_1(s) \\ \text{Im } \Pi_{\bar{u}d,A}^{(1)}(s) &= \frac{1}{2\pi} a_1(s) \\ \text{Im } \Pi_{\bar{u}d,A}^{(0)}(s) &= \frac{1}{2\pi} a_0(s) \end{aligned} \quad (17)$$

chisquared:

$$\chi^2(\alpha) = (I_i^{exp} - I_i^{th}(\alpha)) C_{ij}^{-1} (I_j^{exp} - I_j^{th}(\alpha)) \quad (18)$$

$$\begin{aligned} I_{i=kl}^{exp}(s_k, \omega_l) &= \int_{s_{th}}^{s_k} \frac{ds}{s_k} \omega_l(s) \text{Im } \Pi_{V/A}(s) \\ I_{i=kl}^{th}(s_k, \omega_l) &= \frac{i}{2s_k} \oint_{|s|=s_k} \frac{ds}{s} [W_l(s) - W_l(s_k)] D(s) \end{aligned} \quad (19)$$

used weights

$$\omega_{kl}(s) = \left(1 - \frac{s}{m_\tau^2} \right)^{2+k} \left(\frac{s}{m_\tau^2} \right)^l \left(1 + \frac{2s}{m_\tau^2} \right) \quad (20)$$

$$(k, l) = (0, 0), (1, 0), (1, 1), (1, 2), (1, 3)$$

weight to OPE:

$$\frac{1}{2\pi i s_0} \oint_{|s|=s_0} ds \left(\frac{s}{s_0} \right)^n \frac{C_{2k}}{(-s)^k} = (-1)^{n+1} \frac{C_{2(n+1)}}{s_0^{n+1}} \delta_{k,n+1} \quad (21)$$

implying that an n -th degree monomial in the weight $\omega(s/s_0)$ selects the $D = 2k = 2(n+1)$ term in the OPE.

$$\frac{s}{s_0} + \left(\frac{s}{s_0} \right)^2 + \left(\frac{s}{s_0} \right)^3 + \dots \quad (22)$$

$$\begin{aligned}
A_{00,V/A}^{ALEPH} &= A_{00,V/A}^{ALEPH}(a_s, \mathcal{O}_{6,V/A}) \\
A_{10,V/A}^{ALEPH} &= A_{10,V/A}^{ALEPH}(a_s, \langle a_s GG \rangle, \mathcal{O}_{6,V/A}, \mathcal{O}_{10,V/A}) \\
A_{11,V/A}^{ALEPH} &= A_{11,V/A}^{ALEPH}(a_s, \langle a_s GG \rangle, \mathcal{O}_{6,V/A}, \mathcal{O}_{10,V/A}, \mathcal{O}_{12,V/A}) \\
A_{12,V/A}^{ALEPH} &= A_{12,V/A}^{ALEPH}(a_s, \mathcal{O}_{6,V/A}, \mathcal{O}_{10,V/A}, \mathcal{O}_{12,V/A}, \mathcal{O}_{14,V/A}) \\
A_{13,V/A}^{ALEPH} &= A_{13,V/A}^{ALEPH}(a_s, \mathcal{O}_{10,V/A}, \mathcal{O}_{12,V/A}, \mathcal{O}_{14,V/A}, \mathcal{O}_{16,V/A})
\end{aligned} \tag{23}$$

duality violations

$$\Delta\rho_{V/A}^{DV}(s) = e^{-\delta_{V/A} + \gamma_{V/A}s} \sin(\alpha_{V/A} + \beta_{V/A}s) \tag{24}$$

$$\Delta A_{V/A}^{\omega, DV}(s_0) \equiv \frac{i}{2} \oint_{|s_0|=s_0} \frac{ds}{s_0} \omega(s) \left\{ \Pi_{V/A}(s) - \Pi_{V/A}^{OPE}(s) \right\} = -\pi \int_{s_0}^{\infty} \frac{ds}{s_0} \omega(s) \Delta\rho_{V/A}^{DV}(s) \tag{25}$$

Pich 2016 α_s V+A from five weights

Channel	$\alpha_s(m_\tau^2)$	$\langle a_s GG \rangle$ ($10^{-3} GeV^4$)	\mathcal{O}_6 ($10^{-3} GeV^6$)	\mathcal{O}_8 ($10^{-3} GeV^8$)
V+A (FOPT)	$0.319^{+0.010}_{-0.006}$	-3^{+6}_{-11}	$1.3^{+1.4}_{-0.8}$	$-0.8^{+0.4}_{-0.7}$
V+A (CIPT)	$0.339^{+0.011}_{-0.009}$	-16^{+5}_{-5}	$0.9^{+0.3}_{-0.4}$	$-1.0^{0.5}_{-0.7}$

Boito 2015 α_s V+A from three weights, multiple s_0 starting by s_{min}

Channel	$s_{min}(GeV^2)$	$\alpha_s(m_\tau^2)$	$\mathcal{O}_{6V,A}$	$\mathcal{O}_{8V,A}$	$\delta_{V,A}$	$\gamma_{V,A}$	$\alpha_{V,A}$	$\beta_{V,A}$
V+A (FOPT)	1.550	0.292(9)	-0.90(13)	1.57(22)	3.19(51)	0.80(30)	-2.65(79)	4.42(41)
			-0.63(61)	3.0(2.2)	1.53(56)	1.42(24)	5.73(84)	1.84(43)
V+A (CIPT)	1.550	0.312(13)	-0.90(13)	1.48(25)	3.35(49)	0.70(29)	-2.28(81)	4.23(42)
			1.59(55)	1.44(25)	5.37(89)	2.03(46)	-0.33(56)	2.0(1.8)