Inclusive ratio:

$$R_{\tau} = 12\pi S_{EW} \int_{0}^{m_{\tau}} \frac{\mathrm{d}s}{m_{\tau}^{2}} \left(1 - \frac{s}{m_{\tau}^{2}} \right) \left[\left(1 + 2 \frac{s}{m_{\tau}^{2}} \right) \operatorname{Im} \Pi^{(1)}(s) + \operatorname{Im} \Pi^{(0)}(s) \right]$$
(1)

OPE:

$$\Pi_{OPE}^{(1+0)}(s) = \sum_{k=0}^{\infty} \frac{C_{2k}(s)}{(-s)^k}$$
 (2)

Countour integral:

$$\int_{s_{th}}^{s_0} \frac{\mathrm{d}s}{s_0} \omega(s) \operatorname{Im} \Pi_{V/A}(s) = \frac{i}{2} \oint_{|s|=s_0} \frac{\mathrm{d}s}{s_0} \omega(s) \Pi_{V/A}(s)$$
 (3)

chisquared:

$$\chi^{2}(\alpha) = (I_{i}^{exp} - I_{i}^{th}(\alpha))C_{ij}^{-1}(I_{j}^{exp} - I_{j}^{th}(\alpha))$$
(4)

used weights

$$\omega_{wk}(s) = \left(1 - \frac{s}{m_{\tau}^2}\right)^{2+k} \left(\frac{s}{m_{\tau}^2}\right)^l \left(1 + \frac{2s}{m_{\tau}^2}\right) \tag{5}$$

$$(k, l) = (0, 0), (1, 0), (1, 1), (1, 2), (1, 3)$$

weight to OPE:

$$\frac{1}{2\pi i s_0} \oint_{|s|=s_0} ds \left(\frac{s}{s_0}\right)^n \frac{C_{2k}}{(-s)^k} = (-1)^{n+1} \frac{C_{2(n+1)}}{s_0^{n+1}} \delta_{k,n+1} \tag{6}$$

implying that an *n*-th degree monomial in the weight $\omega(s/s_0)$ selects the D=2k=2(n+1) term in the OPE.

$$\begin{split} A_{00,V/A}^{ALEPH} &= A_{00,V/A}^{ALEPH}(a_s, \mathcal{O}_{6,V/A}) \\ A_{10,V/A}^{ALEPH} &= A_{10,V/A}^{ALEPH}(a_s, \langle a_s GG \rangle, \mathcal{O}_{6,V/A}, \mathcal{O}_{10,V/A}) \\ A_{11,V/A}^{ALEPH} &= A_{11,V/A}^{ALEPH}(a_s, \langle a_s GG \rangle, \mathcal{O}_{6,V/A}, \mathcal{O}_{10,V/A} \mathcal{O}_{12,V/A}) \\ A_{12,V/A}^{ALEPH} &= A_{12,V/A}^{ALEPH}(a_s, \mathcal{O}_{6,V/A}, \mathcal{O}_{10,V/A}, \mathcal{O}_{12,V/A}, \mathcal{O}_{14,V/A}) \\ A_{13,V/A}^{ALEPH} &= A_{13,V/A}^{ALEPH}(a_s, \mathcal{O}_{10,V/A}, \mathcal{O}_{12,V/A}, \mathcal{O}_{14,V/A}, \mathcal{O}_{16,V/A}) \end{split}$$
 (7)

 α_s

Channel	$\alpha_s(m_{\tau}^2)$	$\langle a_s GG \rangle$
V+A (FOPT)	$0.319^{+0.010}_{-0.009}$	-3
V+A (CIPT)	0.339	-16