


AP Calculus AB Practice: Analyzing Graph of f'

AP[®] CALCULUS AB 2004 SCORING GUIDELINES (Form B)

Question 4

The figure above shows the graph of f' , the derivative of the function f , on the closed interval $-1 \leq x \leq 5$. The graph of f' has horizontal tangent lines at $x = 1$ and $x = 3$. The function f is twice differentiable with $f(2) = 6$.

- (a) Find the x -coordinate of each of the points of inflection of the graph of f . Give a reason for your answer.
- (b) At what value of x does f attain its absolute minimum value on the closed interval $-1 \leq x \leq 5$? At what value of x does f attain its absolute maximum value on the closed interval $-1 \leq x \leq 5$? Show the analysis that leads to your answers.
- (c) Let g be the function defined by $g(x) = xf(x)$. Find an equation for the line tangent to the graph of g at $x = 2$.




NO CALCULATOR

AP[®] CALCULUS AB
2006 SCORING GUIDELINES (Form B)

Question 2

Let f be the function defined for $x \geq 0$ with $f(0) = 5$ and f' , the first derivative of f , given by $f'(x) = e^{(-x/4)} \sin(x^2)$. The graph of $y = f'(x)$ is shown above.


- (a) Use the graph of f' to determine whether the graph of f is concave up, concave down, or neither on the interval $1.7 < x < 1.9$. Explain your reasoning.
- (b) On the interval $0 \leq x \leq 3$, find the value of x at which f has an absolute maximum. Justify your answer.
- (c) Write an equation for the line tangent to the graph of f at $x = 2$.



CALCULATOR

AP[®] CALCULUS AB
2008 SCORING GUIDELINES

Question 4



Graph of v

A particle moves along the x -axis so that its velocity at time t , for $0 \leq t \leq 6$, is given by a differentiable function v whose graph is shown above. The velocity is 0 at $t = 0$, $t = 3$, and $t = 5$, and the graph has horizontal tangents at $t = 1$ and $t = 4$. The areas of the regions bounded by the t -axis and the graph of v on the intervals $[0, 3]$, $[3, 5]$, and $[5, 6]$ are 8, 3, and 2, respectively. At time $t = 0$, the particle is at $x = -2$.


- (a) For $0 \leq t \leq 6$, find both the time and the position of the particle when the particle is farthest to the left. Justify your answer.
- (b) For how many values of t , where $0 \leq t \leq 6$, is the particle at $x = -8$? Explain your reasoning.
- (c) On the interval $2 < t < 3$, is the speed of the particle increasing or decreasing? Give a reason for your answer.
- (d) During what time intervals, if any, is the acceleration of the particle negative? Justify your answer.

NO CALCULATOR

AP[®] CALCULUS AB
2004 SCORING GUIDELINES (Form B)

Question 4

The figure above shows the graph of f' , the derivative of the function f , on the closed interval $-1 \leq x \leq 5$. The graph of f' has horizontal tangent lines at $x = 1$ and $x = 3$. The function f is twice differentiable with $f(2) = 6$.



- (a) Find the x -coordinate of each of the points of inflection of the graph of f . Give a reason for your answer.
- (b) At what value of x does f attain its absolute minimum value on the closed interval $-1 \leq x \leq 5$? At what value of x does f attain its absolute maximum value on the closed interval $-1 \leq x \leq 5$? Show the analysis that leads to your answers.
- (c) Let g be the function defined by $g(x) = xf(x)$. Find an equation for the line tangent to the graph of g at $x = 2$.

- (a) $x = 1$ and $x = 3$ because the graph of f' changes from increasing to decreasing at $x = 1$, and changes from decreasing to increasing at $x = 3$.

$$2 : \begin{cases} 1 : x = 1, x = 3 \\ 1 : \text{reason} \end{cases}$$

- (b) The function f decreases from $x = -1$ to $x = 4$, then increases from $x = 4$ to $x = 5$.
 Therefore, the absolute minimum value for f is at $x = 4$.
 The absolute maximum value must occur at $x = -1$ or at $x = 5$.

$$f(5) - f(-1) = \int_{-1}^5 f'(t) dt < 0$$

Since $f(5) < f(-1)$, the absolute maximum value occurs at $x = -1$.

$$4 : \begin{cases} 1 : \text{indicates } f \text{ decreases then increases} \\ 1 : \text{eliminates } x = 5 \text{ for maximum} \\ 1 : \text{absolute minimum at } x = 4 \\ 1 : \text{absolute maximum at } x = -1 \end{cases}$$

- (c) $g'(x) = f(x) + xf'(x)$
 $g'(2) = f(2) + 2f'(2) = 6 + 2(-1) = 4$
 $g(2) = 2f(2) = 12$


Tangent line is $y = 4(x - 2) + 12$

$$3 : \begin{cases} 2 : g'(x) \\ 1 : \text{tangent line} \end{cases}$$

AP[®] CALCULUS AB
2006 SCORING GUIDELINES (Form B)

Question 2

Let f be the function defined for $x \geq 0$ with $f(0) = 5$ and f' , the first derivative of f , given by $f'(x) = e^{(-x/4)} \sin(x^2)$. The graph of $y = f'(x)$ is shown above.



- (a) Use the graph of f' to determine whether the graph of f is concave up, concave down, or neither on the interval $1.7 < x < 1.9$. Explain your reasoning.
- (b) On the interval $0 \leq x \leq 3$, find the value of x at which f has an absolute maximum. Justify your answer.
- (c) Write an equation for the line tangent to the graph of f at $x = 2$.

- (a) On the interval $1.7 < x < 1.9$, f' is decreasing and thus f is concave down on this interval.

2 : $\begin{cases} 1 : \text{answer} \\ 1 : \text{reason} \end{cases}$

- (b) $f'(x) = 0$ when $x = 0, \sqrt{\pi}, \sqrt{2\pi}, \sqrt{3\pi}, \dots$
 On $[0, 3]$ f' changes from positive to negative only at $\sqrt{\pi}$. The absolute maximum must occur at $x = \sqrt{\pi}$ or at an endpoint.

3 : $\begin{cases} 1 : \text{identifies } \sqrt{\pi} \text{ and } 3 \text{ as candidates} \\ \quad - \text{ or } - \\ \text{indicates that the graph of } f \\ \text{increases, decreases, then increases} \\ 1 : \text{justifies } f(\sqrt{\pi}) > f(3) \\ 1 : \text{answer} \end{cases}$

$$f(0) = 5$$

$$f(\sqrt{\pi}) = f(0) + \int_0^{\sqrt{\pi}} f'(x) dx = 5.67911$$

$$f(3) = f(0) + \int_0^3 f'(x) dx = 5.57893$$

This shows that f has an absolute maximum at $x = \sqrt{\pi}$.

- (c) $f(2) = f(0) + \int_0^2 f'(x) dx = 5.62342$


$$f'(2) = e^{-0.5} \sin(4) = -0.45902$$

$$y - 5.623 = (-0.459)(x - 2)$$

4 : $\begin{cases} 2 : f(2) \text{ expression} \\ \quad 1 : \text{integral} \\ \quad 1 : \text{including } f(0) \text{ term} \\ 1 : f'(2) \\ 1 : \text{equation} \end{cases}$

AP[®] CALCULUS AB
2008 SCORING GUIDELINES

Question 4



Graph of v

A particle moves along the x -axis so that its velocity at time t , for $0 \leq t \leq 6$, is given by a differentiable function v whose graph is shown above. The velocity is 0 at $t = 0$, $t = 3$, and $t = 5$, and the graph has horizontal tangents at $t = 1$ and $t = 4$. The areas of the regions bounded by the t -axis and the graph of v on the intervals $[0, 3]$, $[3, 5]$, and $[5, 6]$ are 8, 3, and 2, respectively. At time $t = 0$, the particle is at $x = -2$.

- For $0 \leq t \leq 6$, find both the time and the position of the particle when the particle is farthest to the left. Justify your answer.
- For how many values of t , where $0 \leq t \leq 6$, is the particle at $x = -8$? Explain your reasoning.
- On the interval $2 < t < 3$, is the speed of the particle increasing or decreasing? Give a reason for your answer.
- During what time intervals, if any, is the acceleration of the particle negative? Justify your answer.

- (a) Since $v(t) < 0$ for $0 < t < 3$ and $5 < t < 6$, and $v(t) > 0$ for $3 < t < 5$, we consider $t = 3$ and $t = 6$.

$$x(3) = -2 + \int_0^3 v(t) \, dt = -2 - 8 = -10$$

$$x(6) = -2 + \int_0^6 v(t) \, dt = -2 - 8 + 3 - 2 = -9$$

Therefore, the particle is farthest left at time $t = 3$ when its position is $x(3) = -10$.

- (b) The particle moves continuously and monotonically from $x(0) = -2$ to $x(3) = -10$. Similarly, the particle moves continuously and monotonically from $x(3) = -10$ to $x(5) = -7$ and also from $x(5) = -7$ to $x(6) = -9$.

By the Intermediate Value Theorem, there are three values of t for which the particle is at $x(t) = -8$.

- (c) The speed is decreasing on the interval $2 < t < 3$ since on this interval $v < 0$ and v is increasing.
- (d) The acceleration is negative on the intervals $0 < t < 1$ and $4 < t < 6$ since velocity is decreasing on these intervals.

$$3 : \begin{cases} 1 : \text{identifies } t = 3 \text{ as a candidate} \\ 1 : \text{considers } \int_0^6 v(t) \, dt \\ 1 : \text{conclusion} \end{cases}$$

$$3 : \begin{cases} 1 : \text{positions at } t = 3, t = 5, \\ \quad \text{and } t = 6 \\ 1 : \text{description of motion} \\ 1 : \text{conclusion} \end{cases}$$

1 : answer with reason

$$2 : \begin{cases} 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$$