

# AP Calculus AB Practice: Functions/Miscellaneous

## AP<sup>®</sup> CALCULUS AB 2001 SCORING GUIDELINES

### Question 5

A cubic polynomial function  $f$  is defined by

$$f(x) = 4x^3 + ax^2 + bx + k$$

where  $a$ ,  $b$ , and  $k$  are constants. The function  $f$  has a local minimum at  $x = -1$ , and the graph of  $f$  has a point of inflection at  $x = -2$ .

(a) Find the values of  $a$  and  $b$ .

(b) If  $\int_0^1 f(x) dx = 32$ , what is the value of  $k$ ?

NO CALCULATOR

**AP<sup>®</sup> CALCULUS AB**  
**2001 SCORING GUIDELINES**

**Question 4**

Let  $h$  be a function defined for all  $x \neq 0$  such that  $h(4) = -3$  and the derivative of  $h$  is given by  $h'(x) = \frac{x^2 - 2}{x}$  for all  $x \neq 0$ .

- (a) Find all values of  $x$  for which the graph of  $h$  has a horizontal tangent, and determine whether  $h$  has a local maximum, a local minimum, or neither at each of these values. Justify your answers.
- (b) On what intervals, if any, is the graph of  $h$  concave up? Justify your answer.
- (c) Write an equation for the line tangent to the graph of  $h$  at  $x = 4$ .
- (d) Does the line tangent to the graph of  $h$  at  $x = 4$  lie above or below the graph of  $h$  for  $x > 4$ ? Why?

NO CALCULATOR

**AP<sup>®</sup> CALCULUS AB**  
**2003 SCORING GUIDELINES**

**Question 6**

Let  $f$  be the function defined by

$$f(x) = \begin{cases} \sqrt{x+1} & \text{for } 0 \leq x \leq 3 \\ 5-x & \text{for } 3 < x \leq 5. \end{cases}$$

- (a) Is  $f$  continuous at  $x = 3$ ? Explain why or why not.
- (b) Find the average value of  $f(x)$  on the closed interval  $0 \leq x \leq 5$ .
- (c) Suppose the function  $g$  is defined by

$$g(x) = \begin{cases} k\sqrt{x+1} & \text{for } 0 \leq x \leq 3 \\ mx+2 & \text{for } 3 < x \leq 5, \end{cases}$$

where  $k$  and  $m$  are constants. If  $g$  is differentiable at  $x = 3$ , what are the values of  $k$  and  $m$ ?

NO CALCULATOR

**AP<sup>®</sup> CALCULUS AB**  
**2007 SCORING GUIDELINES**

**Question 6**

Let  $f$  be the function defined by  $f(x) = k\sqrt{x} - \ln x$  for  $x > 0$ , where  $k$  is a positive constant.

- (a) Find  $f'(x)$  and  $f''(x)$ .
- (b) For what value of the constant  $k$  does  $f$  have a critical point at  $x = 1$ ? For this value of  $k$ , determine whether  $f$  has a relative minimum, relative maximum, or neither at  $x = 1$ . Justify your answer.
- (c) For a certain value of the constant  $k$ , the graph of  $f$  has a point of inflection on the  $x$ -axis. Find this value of  $k$ .

NO CALCULATOR

**AP<sup>®</sup> CALCULUS AB**  
**2006 SCORING GUIDELINES**

**Question 6**

The twice-differentiable function  $f$  is defined for all real numbers and satisfies the following conditions:

$$f(0) = 2, \quad f'(0) = -4, \quad \text{and} \quad f''(0) = 3.$$

- (a) The function  $g$  is given by  $g(x) = e^{ax} + f(x)$  for all real numbers, where  $a$  is a constant. Find  $g'(0)$  and  $g''(0)$  in terms of  $a$ . Show the work that leads to your answers.
- (b) The function  $h$  is given by  $h(x) = \cos(kx)f(x)$  for all real numbers, where  $k$  is a constant. Find  $h'(x)$  and write an equation for the line tangent to the graph of  $h$  at  $x = 0$ .

NO CALCULATOR

**AP<sup>®</sup> CALCULUS AB**  
**2001 SCORING GUIDELINES**

**Question 5**

A cubic polynomial function  $f$  is defined by

$$f(x) = 4x^3 + ax^2 + bx + k$$

where  $a$ ,  $b$ , and  $k$  are constants. The function  $f$  has a local minimum at  $x = -1$ , and the graph of  $f$  has a point of inflection at  $x = -2$ .

(a) Find the values of  $a$  and  $b$ .

(b) If  $\int_0^1 f(x) dx = 32$ , what is the value of  $k$ ?

(a)  $f'(x) = 12x^2 + 2ax + b$   
 $f''(x) = 24x + 2a$

$$f'(-1) = 12 - 2a + b = 0$$

$$f''(-2) = -48 + 2a = 0$$

$$a = 24$$

$$b = -12 + 2a = 36$$

$$5 : \left\{ \begin{array}{l} 1 : f'(x) \\ 1 : f''(x) \\ 1 : f'(-1) = 0 \\ 1 : f''(-2) = 0 \\ 1 : a, b \end{array} \right.$$

(b)  $\int_0^1 (4x^3 + 24x^2 + 36x + k) dx$   
 $= x^4 + 8x^3 + 18x^2 + kx \Big|_{x=0}^{x=1} = 27 + k$

$$27 + k = 32$$

$$k = 5$$

$$4 : \left\{ \begin{array}{l} 2 : \text{antidifferentiation} \\ \quad < -1 > \text{ each error} \\ 1 : \text{expression in } k \\ 1 : k \end{array} \right.$$


**AP<sup>®</sup> CALCULUS AB**  
**2001 SCORING GUIDELINES**

**Question 4**

Let  $h$  be a function defined for all  $x \neq 0$  such that  $h(4) = -3$  and the derivative of  $h$  is given by  $h'(x) = \frac{x^2 - 2}{x}$  for all  $x \neq 0$ .

- (a) Find all values of  $x$  for which the graph of  $h$  has a horizontal tangent, and determine whether  $h$  has a local maximum, a local minimum, or neither at each of these values. Justify your answers.
- (b) On what intervals, if any, is the graph of  $h$  concave up? Justify your answer.
- (c) Write an equation for the line tangent to the graph of  $h$  at  $x = 4$ .
- (d) Does the line tangent to the graph of  $h$  at  $x = 4$  lie above or below the graph of  $h$  for  $x > 4$ ? Why?

(a)  $h'(x) = 0$  at  $x = \pm\sqrt{2}$



Local minima at  $x = -\sqrt{2}$  and at  $x = \sqrt{2}$

(b)  $h''(x) = 1 + \frac{2}{x^2} > 0$  for all  $x \neq 0$ . Therefore, the graph of  $h$  is concave up for all  $x \neq 0$ .

(c)  $h'(4) = \frac{16 - 2}{4} = \frac{7}{2}$

$$y + 3 = \frac{7}{2}(x - 4)$$

- (d) The tangent line is below the graph because the graph of  $h$  is concave up for  $x > 4$ .

$$4 : \begin{cases} 1 : x = \pm\sqrt{2} \\ 1 : \text{analysis} \\ 2 : \text{conclusions} \\ < -1 > \text{not dealing with} \\ & \text{discontinuity at } 0 \end{cases}$$

$$3 : \begin{cases} 1 : h''(x) \\ 1 : h''(x) > 0 \\ 1 : \text{answer} \end{cases}$$

1 : tangent line equation

1 : answer with reason

**AP<sup>®</sup> CALCULUS AB**  
**2003 SCORING GUIDELINES**

**Question 6**

Let  $f$  be the function defined by

$$f(x) = \begin{cases} \sqrt{x+1} & \text{for } 0 \leq x \leq 3 \\ 5-x & \text{for } 3 < x \leq 5. \end{cases}$$

- (a) Is  $f$  continuous at  $x = 3$ ? Explain why or why not.  
 (b) Find the average value of  $f(x)$  on the closed interval  $0 \leq x \leq 5$ .  
 (c) Suppose the function  $g$  is defined by

$$g(x) = \begin{cases} k\sqrt{x+1} & \text{for } 0 \leq x \leq 3 \\ mx + 2 & \text{for } 3 < x \leq 5, \end{cases}$$

where  $k$  and  $m$  are constants. If  $g$  is differentiable at  $x = 3$ , what are the values of  $k$  and  $m$ ?

- (a)  $f$  is continuous at  $x = 3$  because

$$\lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^+} f(x) = 2.$$

$$\text{Therefore, } \lim_{x \rightarrow 3} f(x) = 2 = f(3).$$

$$2 : \begin{cases} 1 : \text{answers "yes" and equates the} \\ \text{values of the left- and right-hand} \\ \text{limits} \\ 1 : \text{explanation involving limits} \end{cases}$$

$$\begin{aligned} \text{(b) } \int_0^5 f(x) dx &= \int_0^3 f(x) dx + \int_3^5 f(x) dx \\ &= \frac{2}{3}(x+1)^{3/2} \Big|_0^3 + \left( 5x - \frac{1}{2}x^2 \right) \Big|_3^5 \\ &= \left( \frac{16}{3} - \frac{2}{3} \right) + \left( \frac{25}{2} - \frac{21}{2} \right) = \frac{20}{3} \end{aligned}$$

$$4 : \begin{cases} 1 : k \int_0^3 f(x) dx + k \int_3^5 f(x) dx \\ \text{(where } k \neq 0) \\ 1 : \text{antiderivative of } \sqrt{x+1} \\ 1 : \text{antiderivative of } 5-x \\ 1 : \text{evaluation and answer} \end{cases}$$

$$\text{Average value: } \frac{1}{5} \int_0^5 f(x) dx = \frac{4}{3}$$

- (c) Since  $g$  is continuous at  $x = 3$ ,  $2k = 3m + 2$ .

$$g'(x) = \begin{cases} \frac{k}{2\sqrt{x+1}} & \text{for } 0 < x < 3 \\ m & \text{for } 3 < x < 5 \end{cases}$$

$$\lim_{x \rightarrow 3^-} g'(x) = \frac{k}{4} \text{ and } \lim_{x \rightarrow 3^+} g'(x) = m$$

Since these two limits exist and  $g$  is differentiable at  $x = 3$ , the two limits are

equal. Thus  $\frac{k}{4} = m$ .

$$3 : \begin{cases} 1 : 2k = 3m + 2 \\ 1 : \frac{k}{4} = m \\ 1 : \text{values for } k \text{ and } m \end{cases}$$

$$8m = 3m + 2; m = \frac{2}{5} \text{ and } k = \frac{8}{5}$$



**AP<sup>®</sup> CALCULUS AB**  
**2007 SCORING GUIDELINES**

**Question 6**

Let  $f$  be the function defined by  $f(x) = k\sqrt{x} - \ln x$  for  $x > 0$ , where  $k$  is a positive constant.

- (a) Find  $f'(x)$  and  $f''(x)$ .
- (b) For what value of the constant  $k$  does  $f$  have a critical point at  $x = 1$ ? For this value of  $k$ , determine whether  $f$  has a relative minimum, relative maximum, or neither at  $x = 1$ . Justify your answer.
- (c) For a certain value of the constant  $k$ , the graph of  $f$  has a point of inflection on the  $x$ -axis. Find this value of  $k$ .

(a)  $f'(x) = \frac{k}{2\sqrt{x}} - \frac{1}{x}$

$$f''(x) = -\frac{1}{4}kx^{-3/2} + x^{-2}$$

$$2 : \begin{cases} 1 : f'(x) \\ 1 : f''(x) \end{cases}$$

(b)  $f'(1) = \frac{1}{2}k - 1 = 0 \Rightarrow k = 2$

When  $k = 2$ ,  $f'(1) = 0$  and  $f''(1) = -\frac{1}{2} + 1 > 0$ .  
 $f$  has a relative minimum value at  $x = 1$  by the Second Derivative Test.

$$4 : \begin{cases} 1 : \text{sets } f'(1) = 0 \text{ or } f'(x) = 0 \\ 1 : \text{solves for } k \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$$

(c) At this inflection point,  $f''(x) = 0$  and  $f(x) = 0$ .

$$f''(x) = 0 \Rightarrow \frac{-k}{4x^{3/2}} + \frac{1}{x^2} = 0 \Rightarrow k = \frac{4}{\sqrt{x}}$$

$$f(x) = 0 \Rightarrow k\sqrt{x} - \ln x = 0 \Rightarrow k = \frac{\ln x}{\sqrt{x}}$$

$$3 : \begin{cases} 1 : f''(x) = 0 \text{ or } f(x) = 0 \\ 1 : \text{equation in one variable} \\ 1 : \text{answer} \end{cases}$$

Therefore,  $\frac{4}{\sqrt{x}} = \frac{\ln x}{\sqrt{x}}$   
 $\Rightarrow 4 = \ln x$   
 $\Rightarrow x = e^4$   
 $\Rightarrow k = \frac{4}{e^2}$

**AP<sup>®</sup> CALCULUS AB**  
**2006 SCORING GUIDELINES**

**Question 6**

The twice-differentiable function  $f$  is defined for all real numbers and satisfies the following conditions:

$$f(0) = 2, \quad f'(0) = -4, \quad \text{and} \quad f''(0) = 3.$$

- (a) The function  $g$  is given by  $g(x) = e^{ax} + f(x)$  for all real numbers, where  $a$  is a constant. Find  $g'(0)$  and  $g''(0)$  in terms of  $a$ . Show the work that leads to your answers.
- (b) The function  $h$  is given by  $h(x) = \cos(kx)f(x)$  for all real numbers, where  $k$  is a constant. Find  $h'(x)$  and write an equation for the line tangent to the graph of  $h$  at  $x = 0$ .

(a)  $g'(x) = ae^{ax} + f'(x)$   
 $g'(0) = a - 4$

$g''(x) = a^2e^{ax} + f''(x)$   
 $g''(0) = a^2 + 3$

$$4 : \begin{cases} 1 : g'(x) \\ 1 : g'(0) \\ 1 : g''(x) \\ 1 : g''(0) \end{cases}$$

(b)  $h'(x) = f'(x)\cos(kx) - k\sin(kx)f(x)$   
 $h'(0) = f'(0)\cos(0) - k\sin(0)f(0) = f'(0) = -4$   
 $h(0) = \cos(0)f(0) = 2$   
 The equation of the tangent line is  $y = -4x + 2$ .

$$5 : \begin{cases} 2 : h'(x) \\ 3 : \begin{cases} 1 : h'(0) \\ 1 : h(0) \\ 1 : \text{equation of tangent line} \end{cases} \end{cases}$$