

Symmetries in Classical and Quantum Field Theory

E. Arnold & K. Okyere

October 2025

Undergraduate Summer Research Project

Durham University

Abstract

Abstract here

Contents

1	Introduction	3
2	Symmetries in Classical field theory	3
2.1	Classical field theory	3
2.2	Noether's first theorem	4
2.3	Noether's second theorem	5
3	The Dirac Lagrangian	6
3.1	Gauging the vector symmetry	6
3.2	Gauging the axial symmetry	7
4	Quantum Field Theory	8
4.1	Second Quantisation	8
4.2	Fock Space	9
5	Symmetries in QFT	10
5.1	Slavnov-Taylor identities	10
5.2	Ward-Takahashi identities	11
6	Anomalies in QFT	12
6.1	The Path Integral measure & Fujikawa's method	12
6.2	Atiyah-Singer index Theorem	14
6.3	Anomalies as obstructions to gauging	15
	References	19

1 Introduction

2 Symmetries in Classical field theory

2.1 Classical field theory

In classical mechanics we consider a countable set of particles each with finitely many degrees of freedom and generalised coordinates $q_i(t)$. These generalised coordinates $\{q_i(t)\}_i$ specify the system's configuration (position in configuration space) and together with the generalised conjugate momenta $\left\{p_i(t) := \frac{\partial L}{\partial \dot{q}_i}\right\}_i$ define the system's position in phase space [1]. In classical field theory we generalise this notion of configuration space to a continuum with infinite degrees of freedom. The scalar field $\phi(t)$ can be seen as the generalised coordinates of a continuum $\left(q_i(t) \xrightarrow{i \rightarrow \vec{x}} \phi(\vec{x}, t)\right)$ and for a system of continua the set of scalar fields $\{\phi_k(\vec{x}, t)\}_k$ specifies the system's configuration [2]. Subsequently, we generalise the classical Lagrangian $L(q_i(t), \dot{q}_i(t), t)$ via

$$L(t) = \int d^3\vec{x} \mathcal{L}(\phi_k(\vec{x}, t), \partial_\mu \phi_k(\vec{x}, t), \vec{x}, t), \quad (2.1)$$

where $\mathcal{L}$ is the Lagrangian density. With respect to a path in configuration space, $\vec{\Phi}(\vec{x}, t)$, the classical action becomes

$$S[\vec{\Phi}(\vec{x}, t)] = \int dt L = \int d^4x \mathcal{L}(\phi_k(\vec{x}, t), \partial_\mu \phi_k(\vec{x}, t), \vec{x}, t). \quad (2.2)$$

Using Hamilton's principle [3] ($\delta S[\vec{\Phi}; \vec{\rho}] = 0, \forall \vec{\rho} \in C_0^\infty$)¹ we obtain the field-theoretic Euler-Lagrange equations

$$\frac{\partial \mathcal{L}}{\partial \phi_k} \approx \partial_\mu \left[\frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi_k)} \right], \quad (2.3)$$

where we use $\approx$ to denote an on-shell equality. In Lagrangian systems the Hamiltonian, $H(p_i(t), q_i(t), t)$, is characterised by the Legendre transform

$$H = \sum_i p_i(t) \dot{q}_i(t) - L, \quad (2.4)$$

to generalise this to field theory we introduce the momentum field conjugate to $\phi_k(\vec{x}, t)$

$$\pi_k(\vec{x}, t) := \frac{\partial \mathcal{L}}{\partial \dot{\phi}_k}, \quad (2.5)$$

$$p_i(t) := \frac{\partial L}{\partial \dot{q}_i} \xrightarrow{i \rightarrow \vec{x}} \frac{\partial}{\partial \dot{\phi}_k} \left[\int d^3\vec{x} \mathcal{L} \right] = \int d^3\vec{x} \pi_k(\vec{x}, t). \quad (2.6)$$

Thus the classical Hamiltonian is generalised to

$$H(t) = \int d^3\vec{x} \mathcal{H}(\pi_k(\vec{x}, t), \phi_k(\vec{x}, t), \vec{x}, t), \quad (2.7)$$

¹ δS is a Gateaux derivative, and $\vec{\rho}$ vanishes at the endpoints of $\vec{\Phi}$

$$\mathcal{H} = \sum_k \pi_k(\vec{x}, t) \dot{\phi}_k(\vec{x}, t) - \mathcal{L}, \quad (2.8)$$

where $\mathcal{H}$ is the Hamiltonian density, and the field-theoretic analogue of Hamilton's equations [3] is the Hamiltonian field equations [2]

$$\dot{\phi}_k = \frac{\delta \mathcal{H}}{\delta \pi_k}, \quad \dot{\pi}_k = -\frac{\delta \mathcal{H}}{\delta \phi_k}, \quad (2.9)$$

where we use the functional derivative identify

$$\frac{\delta \mathcal{F}}{\delta g} = \frac{\partial \mathcal{F}}{\partial g} - \partial_\mu \left[\frac{\partial \mathcal{F}}{\partial (\partial_\mu g)} \right]. \quad (2.10)$$

To begin second quantisation we will also require the notion of a field-theoretic poisson bracket. Given two functionals, F and G , of the dynamical fields given by²

$$F = \int d^3 \vec{x} \mathcal{F}(\pi_k, \phi_k, \vec{x}, t), \quad G = \int d^3 \vec{x} \mathcal{G}(\pi_k, \phi_k, \vec{x}, t), \quad (2.11)$$

we define the poisson bracket in field theory via

$$\{F, G\}_f = \int d^3 \vec{x} \sum_k \left[\frac{\delta \mathcal{F}}{\delta \phi_k} \frac{\delta \mathcal{G}}{\delta \pi_k} - \frac{\delta \mathcal{G}}{\delta \phi_k} \frac{\delta \mathcal{F}}{\delta \pi_k} \right] \quad (2.12)$$

2.2 Noether's first theorem

In classical mechanics Noether's first theorem shows the correspondence between global symmetries of the lagrangian and conserved quantities called Noether charges. This generalises to global symmetries of the lagrangian density corresponding to conserved Noether currents in classical field theory. Consider an infinitesimal field transformation $\vartheta(\epsilon)$ such that

$$\phi_n \mapsto \phi_n + \delta_\epsilon \phi_n + \mathcal{O}(\epsilon^2); \quad \delta_\epsilon \phi_n = \epsilon \vartheta_n(\phi_k), \quad (2.13)$$

where the generators, ϑ_k , are independent of spacetime. A transformation is called a global symmetry if its effect on the Lagrangian density is

$$\mathcal{L} \mapsto \mathcal{L} + \epsilon \partial_\mu \Lambda^\mu(\phi_k, x) + \mathcal{O}(\epsilon^2). \quad (2.14)$$

This transformation changes the action by a surface term and the Euler-Lagrange equations are invariant. Using a Taylor expansion of the Lagrangian density under $\vartheta(\epsilon)$ we find that

$$\delta_\epsilon \mathcal{L} = \sum_k \delta_\epsilon \phi_k E_k + \partial_\mu \theta^\mu, \quad (2.15)$$

$$\theta^\mu = \sum_k \delta_\epsilon \phi_k \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi_k)}. \quad (2.16)$$

²n.b. $K(x) = \int dy [K(y) \delta(x-y)]$

Where E_k are the Euler-Lagrange derivatives and θ^μ is the symplectic potential density [4]. By considering the shift in the action we arrive at Noether's first theorem [5, 6]; an infinitesimal transformation $\vartheta(\epsilon)$ is a symmetry of the Lagrangian if and only if

$$\epsilon j^\mu := \theta^\mu - \epsilon \Lambda^\mu, \quad \epsilon \partial_\mu j^\mu = -\delta_\epsilon \phi_k E_k \approx 0. \quad (2.17)$$

Where we use the summation convention for the field indices hereafter and the on-shell divergence-free quantity j^μ is called the conserved Noether current³. This easily generalises to local symmetries: spacetime dependent generators $\vartheta_k(\phi, x)$.

2.3 Noether's second theorem

Noether's second theorem shows that a local symmetry introduces off-shell constraints on the equations of motion and leads to Noether current identities which are crucial for gauge theories. Under local infinitesimal symmetries (gauge symmetries) R_ξ of the form

$$\phi_n \mapsto \phi_n + \delta_\xi \phi_n + \mathcal{O}(\xi^2); \quad \delta_\xi \phi_n = \sum_{m=0} R_n^{\mu_1 \dots \mu_m}(\phi_k) \partial_{\mu_1} \dots \partial_{\mu_m} \xi \lambda(x), \quad (2.18)$$

the action shifts by

$$\delta_\xi S = \int d^4x \delta_\xi \phi_n E_n + \partial_\mu \theta^\mu. \quad (2.19)$$

If $\lambda(x)$ is a compactly supported smooth function, the boundary term vanishes and we have

$$\int d^4x \delta_\xi \phi_n E_n = 0, \quad (2.20)$$

and integration by parts leads to

$$\int d^4x \lambda(x) \Delta(E) = 0; \quad \Delta(E) := \sum_{m=0} (-1)^m \partial_{\mu_1} \dots \partial_{\mu_m} [R_n^{\mu_1 \dots \mu_m}(\phi_k) E_n]. \quad (2.21)$$

By the fundamental lemma of the calculus of variations [7] we obtain the off-shell constraint $\Delta(E) = 0$. Thus the equations of motion are not all independent and the Euler-Lagrange equations ($E_n = 0$) are under-determined. However, observables must be uniquely determined by the theory hence we conclude some degrees of freedom are gauge. Using the surface term S^μ from integration by parts we find an expression similar to [eq2.17](#)

$$\delta_\xi \phi_n E_n = \xi \partial_\mu S^\mu(E_k, \lambda), \quad (2.22)$$

with the important difference that S^μ vanishes on-shell. The quantity $j^\mu + S^\mu$ is divergence-free off-shell, subsequently by the Poincaré lemma we obtain Noether's second theorem [5, 6, 8, 9]; on-shell the canonical Noether current j^μ is the divergence of a two-form

$$j^\mu = -S^\mu(E_k, \lambda) + \partial_\nu \kappa^{\mu\nu} \approx \partial_\nu \kappa^{\mu\nu}; \quad \kappa^{\mu\nu} = -\kappa^{\nu\mu}, \quad (2.23)$$

³n.b. j^μ is ϵ -independent

the antisymmetric two-form $\kappa^{\mu\nu}$ is called a superpotential. Although the superpotential is arbitrary ($\kappa^{\mu\nu}$ is determined up to addition by a divergence-free quantity) it can be uniquely determined if one demands the superpotential obeys additional constraints (such as asymptotic conservation) [9]. In the case of electrodynamics the superpotential is electromagnetic field tensor and eq2.23 becomes Maxwell's equations

$$J^\mu = \partial_\nu F^{\nu\mu}, \quad F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu \quad (2.24)$$

3 The Dirac Lagrangian

3.1 Gauging the vector symmetry

In quantum mechanics the Schrodinger equation is not Lorentz invariant and is thus incompatible with special relativity. A generalisation of the relativistic dispersion relation using Hamiltonian and momentum operators for free particles leads to the Klein-Gordon equation⁴ [10, 11]. However, the Klein-Gordon equation is a second order partial differential equation and subsequently does not uniquely determine time-evolution of the wavefunction. Alternatively, by finding the half-iterate of d'Alembert operator⁵ the Klein-Gordon equation simplifies to a first order partial differential equation; the Dirac equation [12]

$$(i\gamma^\mu \partial_\mu - m)\psi(x) = 0. \quad (3.1)$$

We call the corresponding Lagrangian density the Dirac Lagrangian

$$\mathcal{L}_D = \bar{\psi}(i\gamma^\mu \partial_\mu - m)\psi, \quad (3.2)$$

where the adjoint bispinor is defined as $\bar{\psi} := \psi^\dagger \gamma^0$. This Lagrangian admits a global U(1) symmetry called the vector symmetry

$$\psi \mapsto e^{i\vartheta} \psi, \quad \bar{\psi} \mapsto \bar{\psi} e^{-i\vartheta}. \quad (3.3)$$

This symmetry corresponds to a conserved current, called the vector current, via Noether's first theorem

$$j^\mu = \bar{\psi} \gamma^\mu \psi. \quad (3.4)$$

We now attempt to 'gauge' this symmetry by adding spacetime dependence to the infinitesimal parameter

$$\psi \mapsto e^{i\vartheta(x)} \psi, \quad \bar{\psi} \mapsto \bar{\psi} e^{-i\vartheta(x)}. \quad (3.5)$$

However, the Lagrangian is not invariant under such a transformation so we introduce a derivative operator which transforms covariantly

$$D_\mu := \partial_\mu + ie\Pi_\mu, \quad (3.6)$$

⁴ $(\square + m^2)\psi = 0$

⁵ $\square := \partial_t^2 - \nabla^2$

$$D_\mu \psi \mapsto D'_\mu \left[e^{i\vartheta(x)} \psi \right] = e^{i\vartheta(x)} D_\mu \psi. \quad (3.7)$$

To satisfy this the gauge field must transform as

$$\Pi_\mu \mapsto \Pi_\mu - \frac{1}{e} \partial_\mu \vartheta(x), \quad (3.8)$$

Thus we have a modified Lagrangian which admits a local U(1) gauge symmetry

$$\mathcal{L} = \bar{\psi}(i\rlap{\not{D}} - m)\psi = \bar{\psi}(i\gamma^\mu \partial_\mu - m)\psi - e\Pi_\mu j^\mu, \quad (3.9)$$

where $\rlap{\not{D}} := \gamma^\mu D_\mu$ is the Dirac operator. This is strikingly similar to the electromagnetic Lagrangian density which also has a U(1) gauge symmetry

$$\mathcal{L}_{\text{EM}} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - A_\mu J^\mu. \quad (3.10)$$

Thus we identify $\Pi_\mu = A_\mu$ and $e j^\mu = J^\mu$ and arrive at the Lagrangian for quantum electrodynamics.

$$\mathcal{L}_{\text{QED}} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - e A_\mu j^\mu + \bar{\psi}(i\gamma^\mu \partial_\mu - m)\psi. \quad (3.11)$$

Schwinger arrived at the covariant formulation of quantum electrodynamics similarly by constructing a Lagrangian invariant under Lorentz transformations, gauge transformations, and charge conjugation [13]. However, this is not the end of the story, the massless Dirac equation admits another global U(1) symmetry. In the next sections we construct a gauge-invariant Lagrangian and investigate the resultant gauge theory.

3.2 Gauging the axial symmetry

The massless Dirac Lagrangian admits another symmetry better seen in the Weyl basis

$$\psi = \begin{pmatrix} \psi_L \\ \psi_R \end{pmatrix}, \quad \gamma^\mu \partial_\mu = \begin{pmatrix} 0 & \sigma^\mu \partial_\mu \\ \bar{\sigma}^\mu \partial_\mu & 0 \end{pmatrix}. \quad (3.12)$$

In this basis the Dirac Lagrangian can be seen as the interaction between two Weyl fermions of opposite chirality [14]

$$\mathcal{L}_D = \psi_L^\dagger (i\sigma^\mu \partial_\mu) \psi_L + \psi_R^\dagger (i\bar{\sigma}^\mu \partial_\mu) \psi_R - m(\psi_L^\dagger \psi_R + \psi_R^\dagger \psi_L). \quad (3.13)$$

In the massless case the Lagrangian gains a global U(1) symmetry known as the axial symmetry

$$\psi_L \mapsto e^{i\vartheta} \psi_L, \quad \psi_R \mapsto e^{-i\vartheta} \psi_R, \quad (3.14)$$

which is equivalent to

$$\psi \mapsto e^{i\vartheta \gamma^5} \psi, \quad (3.15)$$

and a Noether current

$$j_5^\mu = \bar{\psi} \gamma^\mu \gamma^5 \psi, \quad \partial_\mu j_5^\mu = 2im \bar{\psi} \gamma^5 \psi. \quad (3.16)$$

We now proceed à la Schwinger, gauging the axial symmetry by promoting ∂_μ to a covariant derivative $\tilde{D}_\mu$ as in eq3.6 and determining the required gauge transformation

$$\mathcal{L}_D|_{m=0} = \bar{\psi}(i\gamma^\mu\partial_\mu)\psi - q\bar{\psi}\gamma^\mu\Pi_\mu\psi, \quad (3.17)$$

$$\Pi_\mu \mapsto e^{i\vartheta\gamma^5}\Pi_\mu e^{-i\vartheta\gamma^5} - \frac{1}{q}\partial_\mu\vartheta\gamma^5. \quad (3.18)$$

Now we must add a kinetic term to the Lagrangian to determine the dynamics of the matrix-valued gauge field. Such a term must be gauge-invariant to preserve the U(1) symmetry. We find a Lagrangian analogous to the Maxwell Lagrangian

$$\mathcal{L}_\Pi = -\frac{1}{4}\text{Tr}(\Gamma_{\mu\nu}\Gamma^{\mu\nu}), \quad (3.19)$$

with field strength

$$\Gamma_{\mu\nu} = \partial_\mu\Pi_\nu - \partial_\nu\Pi_\mu + iq[\Pi_\mu, \Pi_\nu], \quad (3.20)$$

has the required gauge-invariance and is Lorentz invariant. Thus we have constructed a classical theory describing the interaction between massless dirac fermions and an axial gauge boson

$$\mathcal{L}_A = -\frac{1}{4}\text{Tr}(\Gamma_{\mu\nu}\Gamma^{\mu\nu}) + \bar{\psi}i\tilde{D}\psi, \quad (3.21)$$

$$\mathcal{L}_A = -\frac{1}{4}\text{Tr}(\Gamma_{\mu\nu}\Gamma^{\mu\nu}) - q\bar{\psi}\gamma^\mu\Pi_\mu\psi + \bar{\psi}(i\gamma^\mu\partial_\mu)\psi. \quad (3.22)$$

4 Quantum Field Theory

So far we have studied relativistic quantum mechanics, making use of classical fields to describe the dynamics of quantum states. However, such a quantum theory is inconsistent and predicts particles with negative energy which require peculiar interpretations to reconcile like Dirac's infamous hole theory [15]. To proceed to a quantum theory of particle interactions we must promote the dynamical fields to field operators; second quantisation. We shall see how this shift to a quantised field theory affects our calculation of expectation values and probabilities.

4.1 Second Quantisation

In classical mechanics, the canonical transformations are exactly those which preserve the symplectic structure; invariance of poisson brackets of the dynamical variables (p_i, q_i) characterises transformations which leave Hamilton's equations unchanged [3]. A system's position in phase space specifies its classical state. However, in quantum mechanics all properties of a system are included in a quantum state, $|\psi\rangle$, inside of a Hilbert space upon which operators corresponding to observables act.

Dirac's famous canonical quantisation rule ($\{A, B\} \rightarrow \frac{1}{i\hbar}[\hat{A}, \hat{B}]$) [12] allows us quantise the canonical structure of classical mechanics (although Groenewold's theorem shows us that such a rule cannot hold for all functions of the dynamical variables [16]). To obtain a quantised field theory we consider the field-theoretic poisson brackets

$$\{\phi_n(z), \phi_m(w)\}_f = 0 = \{\pi_n(z), \pi_m(w)\}_f, \quad (4.1)$$

$$\{\phi_n(z), \pi_m(w)\}_f = \delta_{nm} \delta^{(3)}(\vec{z} - \vec{w}). \quad (4.2)$$

Which are quantised to the canonical commutation relations for the dynamical field operators acting on a Fock space [17]

$$[\hat{\phi}_n(z), \hat{\phi}_m(w)] = 0 = [\hat{\pi}_n(z), \hat{\pi}_m(w)], \quad (4.3)$$

$$[\hat{\phi}_n(z), \hat{\pi}_m(w)] = i\hbar \delta_{nm} \delta^{(3)}(\vec{z} - \vec{w}). \quad (4.4)$$

However, just like Dirac's rule this procedure does not always produce a consistent quantum theory. In the case of fermionic fields canonical quantisation produces negative energy states [18]; we require anti-commutation relations because of the spin-statistics theorem [19].

4.2 Fock Space

After second quantisation in general we obtain an expression for the Hamiltonian operator in terms of various creation and annihilation operators (e.g. $a_{\vec{p}_1}^{r\dagger}, b_{\vec{p}_2}^{s\dagger}, \dots$ and $a_{\vec{p}_1}^r, b_{\vec{p}_2}^s, \dots$ respectively). Subsequently we define the vacuum state to be annihilated by all particle annihilation operators

$$a_{\vec{p}_1}^r |0\rangle = b_{\vec{p}_1}^r |0\rangle = \dots = 0 \quad (4.5)$$

The indistinguishable n -particle state is given by

$$a_{\vec{p}_1}^{r_1\dagger} a_{\vec{p}_2}^{r_2\dagger} \dots a_{\vec{p}_n}^{r_n\dagger} |0\rangle = |r_1, \vec{p}_1; r_2, \vec{p}_2; \dots; r_n, \vec{p}_n\rangle \in H^{\otimes n} \quad (4.6)$$

$$H^{\otimes n} := \bigotimes_{k=1}^n H \quad (4.7)$$

Where $H^{\otimes n}$ is the n -particle Hilbert space ($H^0 = \mathbb{C}$ corresponds to the vacuum), and an arbitrary quantum state is an element of the Fock space

$$F(H) := \bigoplus_{n=0}^{\infty} H^{\otimes n} \quad (4.8)$$

Thus the probability amplitude of a state transitioning from one particle to another is

$$\langle r_2, \vec{p}_2 | r_1, \vec{p}_1 \rangle = \left\langle 0 \left| a_{\vec{p}_2}^{r_2} a_{\vec{p}_1}^{r_1\dagger} \right| 0 \right\rangle := \left\langle a_{\vec{p}_2}^{r_2} a_{\vec{p}_1}^{r_1\dagger} \right\rangle \quad (4.9)$$

which generalises to transitions between arbitrary quantum states. Hence in quantum field theory the study of probabilities is characterised by vacuum expectation values of operators and their compositions.

5 Symmetries in QFT

5.1 Slavnov-Taylor identities

We now extend our investigation of symmetries from classical fields to a quantised theory. Suppose the classical action admits a local symmetry such that the path integral measure is invariant

$$\phi_\mu \mapsto \phi_\mu + \epsilon \vartheta_\mu(\phi, x), \quad (5.1)$$

$$\mathcal{D}\phi \mapsto \mathcal{D}\phi \det\left(\frac{\partial\phi'_\mu(x)}{\partial\phi_\nu(y)}\right) = \mathcal{D}\phi \quad (5.2)$$

where $K_{\mu\nu} = \frac{\partial\phi'_\mu(x)}{\partial\phi_\nu(y)}$ is the transformation Jacobian. Thus the partition function $Z[J]$ is invariant and for a source J^μ corresponding to ϕ_μ we can write

$$\int d^4x J^\mu(x) \langle \vartheta_\mu(\phi, x) \rangle_J = 0. \quad (5.3)$$

The quantum action is $W[J] = -i \ln Z[J]$, and we define the effective quantum action as the Legendre transform

$$\Gamma[\varphi] = W[J] - \int d^4x J^\mu(x) \varphi_\mu(x); \quad \varphi_\mu := \langle \phi_\mu \rangle_J. \quad (5.4)$$

Taking the functional derivative we obtain

$$J^\mu(x) = -\frac{\delta\Gamma[\varphi]}{\delta\varphi_\mu(x)}, \quad (5.5)$$

after using [eqn7.3](#) and the definition of a functional derivative we have

$$\delta\Gamma[\varphi_\mu; \langle \vartheta_\mu(\phi, x) \rangle_J] = 0. \quad (5.6)$$

Thus the effective quantum action is invariant under the transformations

$$\varphi_\mu \mapsto \varphi_\mu + \epsilon \langle \vartheta_\mu(\phi, x) \rangle_J, \quad (5.7)$$

such a relation is called a Slavnov-Taylor identity [\[20–22\]](#). If the transformation is linear in the fields⁶ such that

$$\vartheta_\mu = \Theta_\mu[\phi, x] = \alpha_\mu(x) + \int d^4y \beta_\mu^\nu(x, y) \phi_\nu(y), \quad (5.8)$$

the transformation Jacobian becomes field independent

$$K_{\mu\nu} = \frac{\delta}{\delta\phi_\nu(y)} (\phi_\mu + \epsilon \Theta_\mu[\phi, x]) = \delta_{\mu\nu} \delta^{(4)}(x - y) + \epsilon \beta_\mu^\nu(x, y). \quad (5.9)$$

⁶i.e. linear $R_n^\mu(\phi_k)$ in [eqn2.18](#)

Moreover, we have $\langle \Theta_\mu[\phi, x] \rangle_J = \Theta_\mu[\varphi, x]$ which implies the effective quantum action is invariant under the same transformation as the classical action, namely,

$$\varphi_\mu \mapsto \varphi_\mu + \epsilon \Theta_\mu[\varphi, x], \quad (5.10)$$

is a symmetry of $\Gamma[\varphi]$ so the normalised one-particle irreducible correlators are invariant. In this case, the quantum theory inherits the classical symmetry and it is impossible to violate the symmetry via quantum effects like loop diagrams because the corresponding correlators are invariant [22].

5.2 Ward-Takahashi identities

The classical conservation laws for Noether currents present in quantum mechanics have a quantum field theoretic analogue, the Ward-Takahashi identities. The conservation of classical fields induces analogous relations for the correlation functions of the corresponding field operators. Consider a bounded functional of the fields $F[\phi]$ and its VEV

$$\langle F[\phi] \rangle = \frac{1}{Z} \int \mathcal{D}\phi' F[\phi'] e^{iS[\phi]}, \quad (5.11)$$

$$\phi' = \phi + \delta_\epsilon \phi + \mathcal{O}(\epsilon^2); \quad \delta_\epsilon \phi = \epsilon \lambda(\phi, x) \quad (5.12)$$

which follows from relabeling the variable of integration. Suppose the field transformation preserves the path integral measure, then taking the functional Taylor expansion leads to

$$\langle F[\phi] \rangle = \frac{1}{Z} \int \mathcal{D}\phi \left(F[\phi] + \frac{\delta F[\phi]}{\delta \phi} \epsilon \lambda + \mathcal{O}(\epsilon^2) \right) \exp \left[iS[\phi] + i \frac{\delta S[\phi]}{\delta \phi} \epsilon \lambda + \mathcal{O}(\epsilon^2) \right], \quad (5.13)$$

$$\langle F[\phi] \rangle = \frac{1}{Z} \int \mathcal{D}\phi \left(F[\phi] + \frac{\delta F[\phi]}{\delta \phi} \epsilon \lambda \right) \left(1 + i \frac{\delta S[\phi]}{\delta \phi} \epsilon \lambda \right) e^{iS[\phi]} + \mathcal{O}(\epsilon^2), \quad (5.14)$$

and to first-order in ϵ we have

$$\int \mathcal{D}\phi \lambda(\phi, x) \left(\frac{\delta F[\phi]}{\delta \phi} + i F[\phi] \frac{\delta S[\phi]}{\delta \phi} \right) e^{iS[\phi]} = 0. \quad (5.15)$$

Since $\lambda(\phi, x)$ is smooth and arbitrary this implies the Dyson-Schwinger equation

$$\left\langle \frac{\delta F[\phi]}{\delta \phi} \right\rangle = -i \left\langle F[\phi] \frac{\delta S[\phi]}{\delta \phi} \right\rangle. \quad (5.16)$$

Suppose the classical Noether current $j^\mu(x)$ has an analogue $\hat{j}^\mu(x)$ in the operator formalism and the operators $\{\hat{\mathcal{V}}_i(z_i)\}$ are functions of spacetime. The derivative of their correlator is [WE SUPPRESS THE HATS IN THE CORRELATOR TO INDICATE A TIME-ORDERED PROD][ADD PROOF L8R]

$$\partial_\mu^x \left\langle j^\mu(x) \prod_{i=1}^n \mathcal{V}_i(z_i) \right\rangle = \left\langle \partial_\mu^x j^\mu(x) \prod_{i=1}^n \mathcal{V}_i(z_i) \right\rangle + \sum_{k=1}^n \delta(x_0 - z_{k0}) \left\langle [j^0(x), \mathcal{V}_k(z_k)] \prod_{i \neq k}^n \mathcal{V}_i(z_i) \right\rangle. \quad (5.17)$$

where we define the commutator in the time-ordered product as

$$[j(x), \mathcal{A}_m(y_m)] := \Theta(y_{\alpha_0} - x_0) j(x) \mathcal{A}_m(y_m) - \Theta(x_0 - y_{\beta_0}) \mathcal{A}_m(y_m) j(x); \quad (5.18)$$

$$\mathcal{T} \prod_{i=1}^n \hat{\mathcal{A}}_i = \cdots \hat{\mathcal{A}}_\alpha(y_\alpha) \hat{\mathcal{A}}_m(y_m) \hat{\mathcal{A}}_\beta(y_\beta) \cdots. \quad (5.19)$$

See appendix for proof. The Noether current is only conserved on-shell so we cannot apply current conservation to the second term (the path integral is taken over all field configurations). Assuming that the symmetry corresponding to j^μ leaves the path integral measure invariant and transforms $\mathcal{V}_i$ as

$$\mathcal{V}_i(\phi) \mapsto \mathcal{V}_i(\phi) + \delta_\epsilon \mathcal{V}_i(\phi) + \mathcal{O}(\epsilon^2); \quad \delta_\epsilon \mathcal{V}_i = \frac{\partial \mathcal{V}_i}{\partial \phi} \delta_\epsilon \phi \quad (5.20)$$

using the Dyson-Schwinger equation with $F[\phi] = \prod_i \mathcal{V}_i$ and [eq2.19](#) we have to first-order in ϵ

$$\left\langle \int d^4x \epsilon \partial_\mu^x j^\mu(x) \prod_{i=1}^n \mathcal{V}_i(z_i) \right\rangle = -i \sum_{k=1}^n \left\langle \delta_\epsilon \mathcal{V}_k(z_k) \prod_{i \neq k}^n \mathcal{V}_i(z_i) \right\rangle. \quad (5.21)$$

Demanding the generator $\lambda(\phi, x)$ is smooth and compactly supported and interchanging the spacetime and path integrals leads to⁷

$$\left\langle \partial_\mu^x j^\mu(x) \prod_{i=1}^n \mathcal{V}_i(z_i) \right\rangle = -i \sum_{k=1}^n \delta^{(4)}(x - z_k) \left\langle \lambda(\phi, x) \frac{\partial \mathcal{V}_k}{\partial \phi}(z_k) \prod_{i \neq k}^n \mathcal{V}_i(z_i) \right\rangle, \quad (5.22)$$

via the fundamental lemma of the calculus of variations. These delta function terms are called Schwinger/contact terms and in the case that none of the insertion times coincide we have the Ward-Takahashi identity

$$\partial_\mu^x \left\langle j^\mu(x) \prod_{i=1}^n \mathcal{V}_i(z_i) \right\rangle = 0; \quad x_0 \neq z_{i_0}. \quad (5.23)$$

6 Anomalies in QFT

6.1 The Path Integral measure & Fujikawa's method

We shall now study transformations which are not necessarily symmetries of the path integral measure, to this end we must introduce a more precise notion of the path integral for fermionic fields. We decompose the Dirac spinor and its adjoint using an orthonormal basis $\{\phi_n(x)\}$ of the Dirac operator's eigenfunctions and the independent Grassmann variables θ_n and $\bar{\xi}_m$

$$\psi = \sum_n \theta_n \phi_n(x) = \sum_n \theta_n \langle x | n \rangle, \quad (6.1)$$

⁷n.b. [eqn2.17](#) $\implies \partial_\mu^x j^\mu = -\lambda E$

$$\bar{\psi} = \sum_m \bar{\xi}_m \phi_m^\dagger(x) = \sum_m \bar{\xi}_m \langle m|x \rangle. \quad (6.2)$$

Hence the path integral measure can be seen as a transformation of a Grassmann measure under a change of variables

$$\theta \mapsto \theta_n \langle x|n \rangle, \quad \bar{\xi} \mapsto \bar{\xi}_m \langle m|x \rangle, \quad (6.3)$$

$$\mathcal{D}\psi \mathcal{D}\bar{\psi} = [\det \langle x|n \rangle \det \langle m|x \rangle]^{-1} \lim_{\substack{N \rightarrow \infty \\ M \rightarrow \infty}} \left\{ \prod_n^N d\theta_n \prod_m^M d\bar{\xi}_m \right\} = \lim_{N \rightarrow \infty} \prod_n^N d\theta_n d\bar{\xi}_n. \quad (6.4)$$

Under the infinitesimal vector transformation the Dirac spinor becomes

$$\psi' = \sum_n \theta'_n \phi_n(x); \quad \theta'_n = e^{i\vartheta(x)} \theta_n, \quad (6.5)$$

and similarly for the adjoint spinor. Thus the path integral measure transforms as

$$\mathcal{D}\psi \mathcal{D}\bar{\psi} \mapsto \left[\det \left(e^{i\vartheta(x)\mathbb{I}} \right) \det \left(e^{-i\vartheta(x)\mathbb{I}} \right) \right]^{-1} \mathcal{D}\psi \mathcal{D}\bar{\psi}, \quad (6.6)$$

and we see the measure is invariant; the vector symmetry is retained by the quantum theory and the correlators are gauge invariant by the Slavnov–Taylor identities. However, this is not the case for the axial transformation; the axial symmetry is ‘anomalous’. The following is an outline of Fujikawa’s method for the existence of the axial anomaly [23]. Under the axial transformation where the Grassmann variables transform as

$$\theta'_n = \sum_m C_{nm} \theta_m, \quad \bar{\xi}'_n = \sum_m C_{nm} \bar{\xi}_m, \quad (6.7)$$

$$C_{nm} = \delta_{nm} + i \int d^4x \vartheta(x) \phi_n^\dagger(x) \gamma^5 \phi_m(x). \quad (6.8)$$

Using the identities

$$\ln(\mathbb{I} + \epsilon A) = \epsilon A + \mathcal{O}(\epsilon^2), \quad (6.9)$$

$$\det(e^A) = e^{\text{Tr}(A)}, \quad (6.10)$$

we find that the path integral measure changes by the Jacobian

$$J[\vartheta] = (\det C)^{-2} = \exp[-2\text{Tr}(\ln C)], \quad (6.11)$$

$$J[\vartheta] = \exp \left[-2\text{Tr} \left(i \int d^4x \vartheta(x) \phi_n^\dagger(x) \gamma^5 \phi_m(x) \right) \right], \quad (6.12)$$

$$J[\vartheta] = \exp \left[-2i \int d^4x \vartheta(x) \sum_n \phi_n^\dagger(x) \gamma^5 \phi_n(x) \right]. \quad (6.13)$$

The sum appearing in the Jacobian is clearly divergent (by orthonormality) and requires the introduction of a regulator or ultraviolet cut-off to obtain a non-singular Jacobian. However, there is a much more elegant approach available involving the Atiyah–Singer index theorem but first we show the relation between the axial anomaly and the axial

current. Henceforth, we define the axial anomaly $\mathcal{A}$ via the Jacobian of the path integral measure

$$J[\vartheta] = \exp \left[- \int d^4x \vartheta(x) \mathcal{A} \right], \quad (6.14)$$

under the axial transformation the partition function becomes

$$Z'[A_\mu, \vartheta] = \int \mathcal{D}\psi' \mathcal{D}\bar{\psi}' \exp \left[\int d^4x \bar{\psi}' (i \not{D} - m) \psi' - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} \right], \quad (6.15)$$

but this is just a relabeling of the integration variables and thus the partition function is invariant. Considering the change in the QED lagrangian we have

$$Z[A_\mu, \vartheta] = \int \mathcal{D}\psi \mathcal{D}\bar{\psi} J[\vartheta] \exp \left[\int d^4x \mathcal{L}_{\text{QED}} + \vartheta(x) (\partial_\mu j_5^\mu - 2im\bar{\psi}\gamma^5\psi) \right]. \quad (6.16)$$

Thus to first order in $\vartheta(x)$ we have

$$Z[A_\mu, \vartheta] = \int \mathcal{D}\psi \mathcal{D}\bar{\psi} \left\{ 1 + \int d^4x \vartheta(x) (\partial_\mu j_5^\mu - 2im\bar{\psi}\gamma^5\psi - \mathcal{A}) \right\} \exp \left[\int d^4x \mathcal{L}_{\text{QED}} \right], \quad (6.17)$$

by the fundamental lemma of the calculus of variations [7] we attain the anomalous divergence for the axial current

$$\partial_\mu j_5^\mu = 2im\bar{\psi}\gamma^5\psi + \mathcal{A}. \quad (6.18)$$

6.2 Atiyah–Singer index Theorem

Given an eigenfunction of the Dirac operator $\not{D}$, ϕ_n , with an eigenvalue $\lambda_n \neq 0$, $\gamma^5\phi_n$ is also an eigenfunction

$$\not{D}(\gamma^5\phi_n) = -\gamma^5\not{D}(\phi_n) = -\lambda_n\gamma^5\phi_n, \quad (6.19)$$

and the two are orthogonal as they have different eigenvalues. However, when the eigenvalue is zero ϕ_n and $\gamma^5\phi_n$ correspond to $\lambda_n = 0$ and are called zero modes of $\not{D}$. We also have that γ^5 is hermitian thus by the spectral theorem there is a basis that diagonalises γ^5 and spans $\text{Ker}(\not{D})$. The eigenvalues of γ^5 are ± 1 (because $(\gamma^5)^2 = I_4$) and we use the projection operators $P_\pm = \frac{1}{2}(I_4 \pm \gamma^5)$ to decompose spinors into their positive and negative eigenvalue (positive and negative chirality respectively) components. Ignoring the infinitesimal parameter for now, only the zero modes contribute to the integral in the Jacobian because of orthonormality and we have

$$\int d^4x \sum_n \phi_n^\dagger \gamma^5 \phi_n = \sum_n \int d^4x \phi_{n+}^{0\dagger} \phi_{n+}^0 - \sum_n \int d^4x \phi_{n-}^{0\dagger} \phi_{n-}^0 = n_+ - n_-, \quad (6.20)$$

$$\int d^4x \mathcal{A} = 2i(n_+ - n_-), \quad (6.21)$$

where $n_{\pm}$ are the number of positive and negative chirality zero modes $\phi_{n_{\pm}}^0$ respectively. This is precisely the analytical index of the Dirac operator projected onto the positive chirality subspace

$$\text{index}(\mathcal{D}_+) = n_+ - n_-, \quad (6.22)$$

$$\mathcal{D}_{\pm} := \mathcal{D}P_{\pm} = \mathcal{D}|_{\{\pm\}}. \quad (6.23)$$

By the Atiyah–Singer index theorem this coincides with the Dirac operator’s topological index on a compact manifold [24]. On a 4-dimensional compact manifold Ω without boundary and Riemannian curvature R this index is [25, 26]

$$\text{index}(\mathcal{D}_+) = \int_{\Omega} \hat{A}(\Omega) \text{ch}(F), \quad (6.24)$$

where the integrand, called the index density, only includes 4-form terms and the curvature form is given by $F = e dA = \frac{e}{2} F_{\mu\nu} dx^{\mu} \wedge dx^{\nu}$. The characteristic classes $\text{ch}(F)$ and $\hat{A}(\Omega)$ are the Chern character of the curvature and Dirac genus of the manifold respectively

$$\text{ch}(F) = \text{Tr} \left(\exp \left[\frac{i}{2\pi} F \right] \right), \quad (6.25)$$

$$\hat{A}(\Omega) = \sqrt{\det \left(\frac{\frac{i}{4\pi} R}{\sinh \frac{i}{4\pi} R} \right)}. \quad (6.26)$$

Now we pick the manifold $S^4 = \Omega$ so the Riemannian curvature is constant and $\hat{A}(\Omega) = 1$. Thus the index becomes

$$\text{index}(\mathcal{D}_+) = \int_{S^4} \text{ch}(F) = \frac{1}{2!} \left(\frac{i}{2\pi} \right)^2 \int_{S^4} \text{Tr}(F^2). \quad (6.27)$$

Assuming that the action due to the Maxwell Lagrangian is finite

$$\int_{S^4} d^4x F_{\mu\nu} F^{\mu\nu} < \infty \iff A_{\mu}|_{S^4} \xrightarrow{x \rightarrow \infty} -\frac{i}{e} g^{-1} \partial_{\mu} g|_{S^4}; \quad g(x) \in \text{U}(1), \quad (6.28)$$

we can map our problem from S^4 to $\mathbb{R}^4$ and $\text{ch}(F)$ coincides with the Dirac operator’s index density in Euclidean space if the gauge fields A_{μ} obey eqn7.38 [27, 28]. Using antisymmetry of the wedge product we identify the axial anomaly in Euclidean space as

$$\partial_{\mu} j_5^{\mu} - 2im\bar{\psi}\gamma^5\psi = \mathcal{A}[A_{\mu}] = \frac{-ie^2}{16\pi^2} \text{Tr}(\epsilon^{\mu\nu\rho\sigma} F_{\mu\nu} F_{\rho\sigma}). \quad (6.29)$$

6.3 Anomalies as obstructions to gauging

We now study the effect of gauging an anomalous symmetries in the case of the Dirac Lagrangian’s axial symmetry. Using Fujikawa’s method [23] we shall calculate the anomaly for a quantum theory with Lagrangian eq3.22 although a topological approach analogous to the previous section is possible using the Atiyah–Singer Families index theorem [29–31].

For the theory of massless Dirac fermions coupling to a U(1) axial gauge field in [eq3.22](#), the path integral in Euclidean space is

$$\int \mathcal{D}\psi \mathcal{D}\bar{\psi} \exp \left[- \int d^4x \bar{\psi} i \tilde{D} \psi - \frac{1}{4} \text{Tr}(\Gamma_{\mu\nu} \Gamma^{\mu\nu}) \right]. \quad (6.30)$$

The classical action is gauge-invariant by construction, thus the only change in the path integral under a gauge transformation is from the Jacobian

$$\int \mathcal{D}\psi' \mathcal{D}\bar{\psi}' e^{-S[\psi', \bar{\psi}']} = \int \mathcal{D}\psi \mathcal{D}\bar{\psi} J[\vartheta] e^{-S[\psi, \bar{\psi}]}, \quad (6.31)$$

where the Jacobian takes the same form as in [eq9.23](#) for an orthonormal basis of eigenfunctions $\{\phi_n(x)\}$ of the operator $\tilde{D}$. We now proceed via Fujikawa's method of gauge-invariant mode cut-off to calculate the Jacobian by introducing a smooth regulator $f(x)$ with all derivatives vanishing at infinity

$$J[\vartheta] = \exp \left[-2i \lim_{N \rightarrow \infty} \left\{ \sum_{n=1}^N \int d^4x \vartheta(x) \phi_n^\dagger \gamma^5 \phi_n \right\} \right], \quad (6.32)$$

$$J[\vartheta] = \exp \left[-2i \lim_{M \rightarrow \infty} \left\{ \sum_{n=1}^{\infty} \int d^4x \vartheta(x) \phi_n^\dagger \gamma^5 f\left(\frac{\lambda_n^2}{M^2}\right) \phi_n \right\} \right], \quad (6.33)$$

$$J[\vartheta] = \exp \left[-2i \lim_{M \rightarrow \infty} \text{Tr} \left\{ \vartheta(x) \gamma^5 f\left(\frac{\tilde{D}^2}{M^2}\right) \right\} \right], \quad (6.34)$$

where the functional trace is taken over spacetime using the basis $\{\phi_n(x)\}$. Now we compute the series using momentum space

$$\phi_n(x) = \int \frac{d^4k}{(2\pi)^2} \tilde{\phi}_n(k) e^{ikx}, \quad (6.35)$$

$$\sum_{n=1}^{\infty} \phi_n^\dagger \gamma^5 f\left(\frac{\tilde{D}^2}{M^2}\right) \phi_n = \int \frac{d^4k d^4p}{(2\pi)^4} \sum_{n=1}^{\infty} e^{-ipx} \tilde{\phi}_n^\dagger(p) \gamma^5 f\left(\frac{\tilde{D}^2}{M^2}\right) \tilde{\phi}_n(k) e^{ikx}, \quad (6.36)$$

$$= \int \frac{d^4k}{(2\pi)^4} \text{Tr} \left\{ \gamma^5 f\left(\frac{(\tilde{D}_\mu + ik_\mu)(\tilde{D}^\mu + ik^\mu)}{M^2} + \frac{iq[\gamma^\mu, \gamma^\nu] \Gamma_{\mu\nu}}{4M^2}\right) \right\}, \quad (6.37)$$

where we have used the identities

$$\sum_{n=1}^{\infty} \tilde{\phi}_n^\dagger(p) A \tilde{\phi}_n(k) = \delta(k-p) \text{Tr}(A), \quad (6.38)$$

$$\tilde{D}^2 = \tilde{D}_\mu \tilde{D}^\mu + \frac{iq}{4} [\gamma^\mu, \gamma^\nu] \Gamma_{\mu\nu}, \quad (6.39)$$

$$e^{-ikx} f(\partial_\mu) e^{ikx} = f(\partial_\mu + ik_\mu). \quad (6.40)$$

Now we rescale the momenta via $k_\mu \rightarrow Mk_\mu$ and the series becomes

$$M^4 \int \frac{d^4 k}{(2\pi)^4} \text{Tr} \left\{ \gamma^5 f \left(\frac{\tilde{D}_\mu \tilde{D}^\mu + 2iMk_\mu \tilde{D}^\mu}{M^2} - k^2 + \frac{iq[\gamma^\mu, \gamma^\nu] \Gamma_{\mu\nu}}{4M^2} \right) \right\}, \quad (6.41)$$

expanding the Taylor series of $f(x)$ about $-k^2 = -k_\mu k^\mu$, and using the trace properties of the gamma matrices the only term non-vanishing in the limit $M \rightarrow \infty$ is the term quadratic in $\Gamma_{\mu\nu}$

$$\frac{M^4 q^2}{2!M^4} \text{Tr}(\epsilon^{\mu\nu\rho\sigma} \Gamma_{\mu\nu} \Gamma_{\rho\sigma}) \int \frac{d^4 k}{(2\pi)^4} f''(-k^2), \quad (6.42)$$

and we have

$$J[\vartheta] = \exp \left[\int d^4 x \frac{-iq^2}{16\pi^2} \text{Tr}(\epsilon^{\mu\nu\rho\sigma} \Gamma_{\mu\nu} \Gamma_{\rho\sigma}) \vartheta(x) \right], \quad (6.43)$$

thus there is gauge-variance in the path integral measure and the quantum theory is inconsistent. In the general case, gauging an anomalous symmetry will produce an inconsistent quantum theory as constructing a gauge invariant classical action ensures the shift in the measure makes the path integral change under a gauge transformation.

References

- [1] L. Landau and E. Lifshitz, *The Equations of Motion*, in *Mechanics*, vol. 1, (Oxford), pp. 1–12, Butterworth-Heinemann (1976).
- [2] W. Greiner and J. Reinhardt, *Classical Field Theory*, in *Field Quantization*, (Heidelberg), pp. 31–54, Springer Berlin Heidelberg (1996).
- [3] L. Landau and E. Lifshitz, *The Canonical Equations*, in *Mechanics*, vol. 1, (Oxford), pp. 131–167, Butterworth-Heinemann (1976).
- [4] J. Lee and R.M. Wald, *Local symmetries and constraints*, *J. Math. Phys.* **31** (1990) 725.
- [5] E. Noether, *Invariante variationsprobleme*, *Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse* **1918** (1918) 235.
- [6] E. Noether, *Invariant variation problems*, *Transport Theor. Stat. Phys.* **1** (1971) 186.
- [7] J. Jost and X. Li-Jost, *The classical theory*, in *Calculus of Variations*, Cambridge Studies in Advanced Mathematics, (Cambridge), pp. 4–8, Cambridge University Press (1998).
- [8] S.G. Avery and B.U.W. Schwab, *Noether’s second theorem and ward identities for gauge symmetries*, *J. High Energy Phys.* **2016** (2016) 31.
- [9] G. Barnich and F. Brandt, *Covariant theory of asymptotic symmetries, conservation laws and central charges*, *Nucl. Phys. B* **633** (2002) 3.
- [10] O. Klein, *Quantentheorie und fünfdimensionale relativitätstheorie*, *Zeitschrift für Physik* **37** (1926) 895.
- [11] W. Gordon, *Der comptoneffekt nach der schrödingerschen theorie*, *Zeitschrift für Physik* **40** (1926) 117.
- [12] P.A.M. Dirac, *The Principles of Quantum Mechanics*, Clarendon Press, Oxford, 4 ed. (1958).
- [13] J. Schwinger, *Quantum electrodynamics. i. a covariant formulation*, *Phys. Rev.* **74** (1948) 1439.
- [14] H. Weyl, *Elektron und gravitation*, *Z. Phys.* **56** (1929) 330.
- [15] P.A.M. Dirac, *A theory of electrons and protons*, *Proc. Roy. Soc. Lond. A* **126** (1930) 360.
- [16] H. Groenewold, *On the principles of elementary quantum mechanics*, *Physica* **12** (1946) 405.
- [17] J.R. Klauder, *Exponential hilbert space: Fock space revisited*, *J. Math. Phys.* **11** (1970) 609.
- [18] W. Pauli, *On dirac’s new method of field quantization*, *Rev. Mod. Phys.* **15** (1943) 175.
- [19] W. Pauli, *On the connection between spin and statistics*, *Prog. Theor. Exp. Phys.* **5** (1950) 526.
- [20] A.A. Slavnov, *Ward identities in gauge theories*, *Theor. Math. Phys.* **10** (1972) 99.
- [21] J. Taylor, *Ward identities and charge renormalization of the yang-mills field*, *Nucl. Phys. B* **33** (1971) 436.
- [22] M. Dütsch, *Slavnov–taylor identities from the causal point of view*, *Int. J. Mod. Phys. A* **12** (1997) 3205.
- [23] K. Fujikawa and H. Suzuki, *The Jacobian in path integrals and quantum anomalies*, in *Path Integrals and Quantum Anomalies*, (Oxford), Clarendon Press (2004).
- [24] M.F. Atiyah and I.M. Singer, *The index of elliptic operators: I*, *Ann. Math.* **87** (1968) 484.

- [25] L. Alvarez-Gaume, *A note on the atiyah-singer index theorem*, *J. Phys. A: Math. Gen.* **16** (1983) 4177.
- [26] T. Eguchi, P.B. Gilkey and A.J. Hanson, *Gravitation, gauge theories and differential geometry*, *Phys. Rep.* **66** (1980) 213.
- [27] R. Jackiw and C. Rebbi, *Spinor analysis of yang-mills theory*, *Phys. Rev. D* **16** (1977) 1052.
- [28] A. Belavin, A. Polyakov, A. Schwartz and Y. Tyupkin, *Pseudoparticle solutions of the yang-mills equations*, *Phys. Lett. B* **59** (1975) 85.
- [29] L. Alvarez-Gaumé and P. Ginsparg, *The topological meaning of non-abelian anomalies*, *Nucl. Phys. B* **243** (1984) 449.
- [30] T. Sumitani, *Chiral anomalies and the generalised index theorem*, *J. Phys. A: Math. Gen.* **17** (1984) L811.
- [31] M.F. Atiyah and I.M. Singer, *The index of elliptic operators: Iv*, *Ann. Math.* **93** (1971) 119.