

Norwegian University of Science and Technology Department of Mathematical Sciences TMA4165 Differential equations and dynamical systems Spring 2017

Solutions exercise 10

 $\boxed{\textbf{3.7}}$ A limit cycle encloses N nodes, F spirals, C centres, and S saddle points. Show that

$$N + F + C - S = 1.$$

Let e be the limit cycle in question. Then $I_e = 1$. By theorem 3.2 in [JS] we have

$$1 = I_e = \sum_{i=1}^{n} I_i = N + F + C - S$$

since the index for a saddle point is -1 and for nodes, spirals and centres we have I = 1.

3.12 We are to find the index at infinity for the system

$$\dot{x} = x - y$$

$$\dot{y} = x - y^2.$$

Here, there are two equilibrium points, namely (0,0) and (1,1). The matrix of linearization is given by

$$A = \begin{bmatrix} 1 & -1 \\ 1 & -2y \end{bmatrix}$$

at the point (x, y).

The eigenvalues of A at the origin can be found by solving the system $-\lambda(1-\lambda)+1=0$ which gives $\lambda=\frac{1}{2}\pm\frac{\sqrt{3}}{2}i$. This is an unstable spiral also in the original system and so the index at the origin is I(0,0)=1.

The eigenvalues of A at (1,1) can be found by solving $(1-\lambda)(-2-\lambda)+1=0$. This gives $\lambda=-\frac{1}{2}\pm\frac{\sqrt{5}}{2}$. This is a saddle point, so I(1,1)=-1.

The index at infinity is now given by J.S. theorem 3.4,

$$I_{\infty} = 2 - (1 - 1) = 2.$$

3.25 Show that the second order system $\dot{\mathbf{x}} = \mathbf{X}(\mathbf{x})$ and $\operatorname{curl}(\mathbf{X}) = 0$ in a simply connected region D cannot have closed paths.

First note that

$$\operatorname{curl}(\mathbf{X}) = 0 \quad \Leftrightarrow \quad \frac{\partial Y}{\partial x} = \frac{\partial X}{\partial y}.$$

We solve the problem using Green's theorem. Suppose we have a closed curve Γ and call the interior region R. Then we have

$$0 = \iint_{R} \left(\frac{\partial Y}{\partial x} - \frac{\partial X}{\partial y} \right) dA = \oint_{\Gamma} \mathbf{X} \cdot \mathbf{T} ds$$

where **T** is the unit tangent vector for the curve Γ . However, since Γ is a phase path, **T** will always be parallel to the vector **X**. Hence, the right hand side in the previous equation cannot be zero and we have a contradiction.

The system

$$\dot{x} = y + 2xy,$$

$$\dot{y} = x + x^2 - y^2$$

satisfy the conditions, so it follows that the system does not have any closed paths.

8.3 Find the limit cycles of the system

$$\begin{split} \dot{x} &= -y + x \sin \sqrt{x^2 + y^2}, \\ \dot{y} &= x + y \sin \sqrt{x^2 + y^2}. \end{split}$$

Determine which of them are Poincaré stable.

Introduce polar coordinates to find

$$\dot{r} = r \sin r,$$

$$\dot{\theta} = 1.$$

Since \dot{r} is only a function of r, we have limit cycles where $\dot{r} = 0$. This gives us limit cycles when $r = n\pi$ where $n \in \mathbb{N}$. For limit cycles of the form $r = (2n + 1)\pi$, \dot{r} is positive just inside the limit cycle and negative just outside the limit cycle. This is a stable limit cycle. The other limit cycles are unstable. See figure 1 for a sketch of the phase diagram.




Figure 1: Phase diagram of $\dot{x} = -y + x \sin \sqrt{x^2 + y^2}$, $\dot{y} = x + y \sin \sqrt{x^2 + y^2}$