TMA4255 Applied Statistics Solution to Exercise 6

Problem 1

a) Main effect of z_1 = expected average response when z_1 is on the high level minus the expected average response when z_1 is on the low level.

$$\frac{y_4 + y_2}{2} - \frac{y_3 + y_1}{2} = \frac{(\beta_0 + \beta_1 + \beta_2 + \beta_{12}) + (\beta_0 + \beta_1 - \beta_2 - \beta_{12})}{2} - \frac{(\beta_0 - \beta_1 + \beta_2 - \beta_{12}) + (\beta_0 - \beta_1 - \beta_2 + \beta_{12})}{2} = 2\beta_1$$

Main effect of z_2 = expected average response when z_2 is on the high level minus the expected average response when z_2 is on the low level.

$$\frac{y_4 + y_3}{2} - \frac{y_2 + y_1}{2} = \frac{(\beta_0 + \beta_1 + \beta_2 + \beta_{12}) + (\beta_0 - \beta_1 + \beta_2 - \beta_{12})}{2} - \frac{(\beta_0 + \beta_1 - \beta_2 - \beta_{12}) + (\beta_0 - \beta_1 - \beta_2 + \beta_{12})}{2} = 2\beta_2$$

The interaction between z_1 and z_2 : 1) half the main effect of z_1 when z_2 is on the high level minus 2) half the main effect of z_1 when z_2 is on the low level.

1) the main effect of z_1 when z_2 is on the high level:

$$\frac{(\beta_0 + \beta_1 + \beta_2 + \beta_{12}) - (\beta_0 - \beta_1 + \beta_2 - \beta_{12})}{1} = 2\beta_1 + 2\beta_{12}$$

2) the main effect of z_1 when z_2 is on the low level:

$$\frac{(\beta_0 + \beta_1 - \beta_2 - \beta_{12}) - (\beta_0 - \beta_1 - \beta_2 + \beta_{12})}{1} = 2\beta_1 - 2\beta_{12}$$

Then, the interaction between z_1 and z_2 :

$$\frac{2\beta_1 + 2\beta_{12}}{2} - \frac{2\beta_1 - 2\beta_{12}}{2} = 2\beta_{12}$$

b) Main effect of z_1 while keeping z_2 at low level (we have already calculated above):

$$\frac{(\beta_0 + \beta_1 - \beta_2 - \beta_{12}) - (\beta_0 - \beta_1 - \beta_2 + \beta_{12})}{1} = 2\beta_1 - 2\beta_{12}$$

Main effect of z_1 while keeping z_2 at high level (we have already calculated above):

$$\frac{(\beta_0 + \beta_1 + \beta_2 + \beta_{12}) - (\beta_0 - \beta_1 + \beta_2 - \beta_{12})}{1} = 2\beta_1 + 2\beta_{12}$$

c) Based on the results in **b** we see that main effect of z_1 when z_2 is at its low level is $2\beta_1 - 2\beta_{12}$, and main effect of z_1 when z_2 is at its high level is $2\beta_1 + 2\beta_{12}$. If the interaction between z_1 and z_2 is zero we can find the main effect of z_1 by fixing the other factors at a given level. But, when there are interactions present, just fixing one factor at a given level will not give us estimate of the main effect, but the main effect and the interaction effect (as shown above). Therefore, we do not fix one factor at one level and vary the other factor in DOE!

Problem 2

a) We do the analysis in MINITAB:

Estimated Effects and Coefficients for C9 (coded units)

Term	Effect	Coef
Constant		17,544
A	8,837	4,419
В		
Ь	-2,512	-1,256
C	-1,087	-0,544
D	0,112	0,056
A*B	-0,762	-0,381
A*C	1,013	0,506
A*D	0,212	0,106
B*C	1,012	0,506
B*D	0,262	0,131
C*D	-0,162	-0,081
A*B*C	0,213	0,106
A*B*D	-0,038	-0,019
A*C*D	1,387	0,694
B*C*D	0,288	0,144
A*B*C*D	-0,263	-0,131

S = * PRESS = *

Analysis of Variance for C9 (coded units)

```
Source
                    DF
                         Seq SS
                                  Adj SS
                                            Adj MS
Main Effects
                     4
                        342,437
                                 342,437
                                           85,6094
2-Way Interactions
                     6
                         11,089
                                  11,089
                                            1,8481
3-Way Interactions
                     4
                          8,217
                                   8,217
                                            2,0544
4-Way Interactions
                     1
                          0,276
                                   0,276
                                            0,2756
Residual Error
                     0
                    15 362,019
Total
```

$$\hat{A} = 8.84$$

$$\hat{B} = -2.51$$

$$\hat{C} = -1.09$$

$$\hat{D} = 0.11$$

$$\vdots \quad \vdots$$

$$\hat{ABCD} = -0.262$$

From the normal plot in figure (1) it looks like A and B are the most important factors.




Figure 1: Normal plot a)

b) The corresponding regression model is

$$Y = \beta_0 + \beta_1 z_1 + \beta_2 z_2 + \beta_3 z_3 + \beta_4 z_4 \tag{1}$$

$$+\beta_{12}z_1z_2 + \beta_{13}z_1z_3 + \beta_{14}z_1z_4 \tag{2}$$

$$+\beta_{23}z_2z_3 + \beta_{24}z_2z_4 + \beta_{34}z_3z_4 \tag{3}$$

$$+\beta_{123}z_1z_2z_3 + \beta_{124}z_1z_2z_4 + \beta_{134}z_1z_3z_4 \tag{4}$$

$$+\beta_{234}z_2z_3z_4 + \beta_{1234}z_1z_2z_3z_4 + \epsilon \tag{5}$$

And the estimated effects are of the kind

$$\hat{A} = 2b_1 \tag{6}$$

where b_1 is the least squares estimator of β_1 . Same goes for the other effects.

c) In the analysis in a) we have 16 equations and 16 coefficients to estimate. Therefore there are no degrees of freedom left to estimate the variance. If we assume that the variance is known it is possible to make inference about the effects. For factor A we have:

$$\begin{vmatrix}
\hat{A} = \frac{1}{8}(-Y_1 + Y_2 - \dots - Y_{15} + Y_{16}) \\
\operatorname{Var}(\hat{A}) = \frac{1}{64}16\sigma^2 = \frac{\sigma^2}{4}
\end{vmatrix} \Rightarrow (\hat{A} \sim N(\mu_A, \frac{\sigma^2}{4})$$

95 % confidence interval for μ_A :

$$\hat{A} \pm z_{0.025} \frac{\sigma}{2} = (6.88, 10.80)$$

95 % confidence interval for μ_B :

$$\hat{B} \pm z_{0.025} \frac{\sigma}{2} = (-4.47, -0.5)$$

d) If there are good reasons to assume that the 3- and 4-factor interactions are 0, we have enough degrees of freedom to estimate the variance.

From MINITAB we get:

Fractional Factorial Fit

Estimated Effects and Coefficients for Response (coded units)

Term	Effect	Coef	StDev Coef	Т	P
Constant		17,544	0,3258	53,84	0,000
A	8,837	4,419	0,3258	13,56	0,000
В	-2,512	-1,256	0,3258	-3,86	0,012
C	-1,087	-0,544	0,3258	-1,67	0,156
D	0,112	0,056	0,3258	0,17	0,870
A*B	-0,762	-0,381	0,3258	-1,17	0,295
A*C	1,012	0,506	0,3258	1,55	0,181
A*D	0,212	0,106	0,3258	0,33	0,758
B*C	1,012	0,506	0,3258	1,55	0,181
B*D	0,262	0,131	0,3258	0,40	0,704
C*D	-0,162	-0,081	0,3258	-0,25	0,813

Analysis of Variance for Response (coded units)

Source	DF	Seq SS	Adj SS	Adj MS	F	P
Main Effects	4	342,437	342,437	85,609	50,40	0,000
2-Way Interactions	6	11,089	11,089	1,848	1,09	0,473
Residual Error	5	8,493	8,493	1,699		
Total	15	362,019				

We see that the estimator for σ^2 is now:

$$s^2 = MS_E = \frac{s_{ABC} + \dots + s_{BCD} + s_{ABCD}}{5} = \frac{8.22 + 0.28}{5} = 1.70$$

where 8.22 is 3-way Seq SS, and 0.28 is 4-way Seq SS from the full analysis in section a. The variance of the effects is thus estimated by

$$s_{effect}^2 = \frac{4s^2}{n} = 0.425$$

We can also obtain this estimate of σ_{effect}^2 directly by using the estimated effects

$$s_{\textit{effect}}^2 = \frac{\widehat{ABC}^2 + \dots + \widehat{BCD}^2 + \widehat{ABCD}^2}{5} = \frac{0.213^2 + 0.038^2 + 1.387^2 + 0.288^2 + 0.263^2}{5} = 0.425$$

Now we can do a T-test or an equivalent F-test to decide which of the effects are significant. We use the results and do an F-test:

$$F_A = \frac{MS_A}{MS_E} = \frac{s_A^2}{1.7} = \frac{(\hat{A}2^k/2)^2/2^k}{1.7} = \frac{312.37}{1.7} = 183.74$$

or alternatively, with $n = 2^k$,

$$F_A = \frac{MS_A}{MS_E} = \frac{\beta_A^2 n}{1.7} = \frac{(\hat{A}/2)^2 n}{1.7} = \frac{312.37}{1.7} = 183.74$$

$$F_B = \frac{MS_B}{MS_E} = \frac{s_B^2}{1.7} = \frac{25.26}{1.7} = 14.86,$$

and get the p-values:

$$p = P(F_{1,5} > 183.74) = 2P(T_5 > 13.56) \approx 0$$

 $p = P(F_{1,5} > 14.86) = 2P(T_5 > 3.85) = 0.012$

Use that

$$F_{1,\nu} = T_{\nu}^2$$

): A has effect and B is significant at all levels > 0.012.

e) From MINITAB we get

Full Factorial Design

Factors: 4 Base Design: 4; 16 Resolution with blocks: V

Runs: 16 Replicates: 18 Replicates: 2 Center pts (total): 19

Block Generators: ABCD

Alias Structure

Ι

Blk = ABCD

Α

B C

D

AB

AC AD

BC

BD

CD

ABC

ABD ACD

BCD

Design Table

Run Block A B C D
1 1 + - - 2 1 - + - 3 1 - - + 4 1 + + + 5 1 - - +

We see that ABCD is the only effect confounded with the block effect

f) To perform the experiment in two blocks, we need two generators. Choosing ABC and AD as generators gives

$$ABC \cdot AD = BCD \tag{7}$$

$$ABC \cdot BCD = AD \tag{8}$$

$$AD \cdot BCD = ABC \tag{9}$$

And we see that the effects confounded with the blocks are ABC, BCD and AD. This design avoids main effects being confounded with the block effect.