

TMA4125 Matematikk

4N

Spring 2017

Norwegian University of Science and Technology Institutt for matematiske fag

Exercise set 7

1 Use the Laplace transform to solve the Initial Value Problem

$$y'' - 9y = \delta(t - 1),$$

with initial conditions y(0) = 2, y'(0) = 0. (See Kreyszig, Chapter 6.4 for details on how to handle the delta function $\delta(t-1)$)

2 In this exercise we will solve the integral equation

$$y(t) + \int_0^t y(\tau) \cosh(t - \tau) d\tau = t + e^t$$

- a) Rewrite the integral appearing in the equation as a convolution (see Kreyszig, Chapter 6.5)
- b) By taking Laplace transforms of both sides and using the convolution theorem, show that the Laplace transform of y(t) obeys the equation

$$Y(s) = \frac{s^2 - 1}{s^2(s - 1)}$$

c) Solve the integral equation for y(t) by simplifying the above expression and then taking the inverse Laplace transform.

 $\boxed{\bf 3}$ The oscillations of a spring under a forcing term r(t) are described by the equation

$$y'' + 4y = r(t),$$

where the system is initially at rest, i.e. y(0) = y'(0) = 0.

a) Find an expression for the Laplace transform of y(t) in the form Y(s) = Q(s)R(s), where R(s) is the Laplace transform of r(t), and Q(s) is a function you should find (this Q(s) is the transfer function of the system, see Kreyszig 6.5). Conclude that

$$y(t) = \frac{1}{2}\sin 2t * r(t),$$

b) By computing the above convolution, solve for y in the case that $r(t) = 4\cos 2t$. You may use the identity

$$\sin \alpha \cos \beta = \frac{1}{2} (\sin(\alpha + \beta) + \sin(\alpha - \beta))$$

What is happening as t becomes large? What physical phenomenon does this correspond to?

4 We investigate the variable coefficient ODE,

$$ty'' - 2ty' + 2y = 2$$

with initial condition y(0) = 1.

a) Show that the Laplace transform of y(t) obeys the first order linear ODE

$$s\frac{dY}{ds} + 2Y = \frac{1}{s}$$

You will require the formula for $\mathcal{L}(tf(t))$ given in Chapter 6.6 of Kreyszig.

- b) (Optional) Solve the equation for Y and hence find y(t). (Hint: consider solutions of the form $Y = s^k$. It may be helpful to solve the homogeneous equation sY' + 2Y = 0 separately. Alternatively, use an integrating factor)
- 5 A system of two connected springs subject to an external force is described by the equations

$$y_1'' = -8y_1 + 4y_2 + 11\sin t$$

$$y_2'' = 4y_1 - 8y_2 - 11\sin t$$

with initial conditions $y_1(0) = 1, y'_1(0) = 1, y_2(0) = 1, y'_2(0) = -1$. (See Kreyszig, Chapter 6.7 for a discussion of such systems)

a) Show that the Laplace transforms of y_1 and y_2 satisfy the linear system

$$\begin{pmatrix} s^2 + 8 & -4 \\ -4 & s^2 + 8 \end{pmatrix} \begin{pmatrix} Y_1 \\ Y_2 \end{pmatrix} = \begin{pmatrix} s + 1 + \frac{11}{s^2 + 1} \\ s - 1 - \frac{11}{s^2 + 1} \end{pmatrix}$$

b) (Optional, for those who wish to practise their algebra, or can solve linear systems by computer!) By solving for Y and then inverting, show that the solution is given by

$$y(t) = \begin{pmatrix} \cos 2t + \sin t \\ \cos 2t - \sin t \end{pmatrix}$$