Statement E.74 (Proved). The complete matrix-distribution and Gaussian central-layer fibres. *Keep the literal* 24 *E.17 words f_i retained in E.71, and put*

$$N = N_B, \qquad K_6 = \overline{N'N^6}, \qquad J = K_6 \cap \widehat{F}', \qquad \rho : J \longrightarrow P \tag{E.74.1}$$

for the entire source and the actual 7501*-point correction map of E.71. Thus* $|P| = 2^{105}3^{17}$*, and its specified matrix, coefficient, distribution, Ferrari and orbifold conditions have* 18 432 *finite values, with all their source words retained.*

Let $\widetilde{\Theta}$ be the actual twelve-evaluation homomorphism of E.69, restricted to J. Write

$$L = \widetilde{\Theta}(J) = \mathcal{M} \times \widetilde{\mathcal{G}} \simeq C_3^3 \times H_3^3, \qquad M = \mathcal{M} \times \mathcal{G} \simeq C_3^9, \tag{E.74.2}$$

where H_3 is the exponent-three Heisenberg group with the exact coordinates and graph maps of E.69. The homomorphism $\varepsilon : L \to M$ forgets its Gaussian central coordinates; its kernel is $L' \simeq C_3^3$. The old orbifold coordinate of E.71 gives an actual epimorphism $\pi : P \to M$. On the identical source J these maps satisfy

$$\pi\rho = \varepsilon\widetilde{\Theta}. \tag{E.74.3}$$

The complete common correction image is

$$(\rho, \widetilde{\Theta})(J) = P \times_M L, \qquad |P \times_M L| = 2^{105}3^{20}. \tag{E.74.4}$$

Every one of the 18 432 *E.71 values has exactly three lifts to the specified Gaussian central-frame targets of E.69. Consequently the complete common finite locus has* 55 296 *values, all with saved actual J-word preimages. For a retained joint correction u over f_i, its entire word fibre inside f_iJ is*

$$f_i u\big(\ker\rho \cap \ker(\widetilde{\Theta}|_J)\big). \tag{E.74.5}$$

The frame choices and the absolute twelve-group values use the literal E.17 origin in each row. This statement concerns precisely the displayed finite maps and their whole word fibres.

Proof. We exhibit three actual words in the kernel of the entire E.71 source which supply all Gaussian central corrections.

First, $N' \subset J \subset K_6$, and E.69 proves $\widetilde{\Theta}(N') = \widetilde{\Theta}(K_6) = L$. This proves the equality defining L in (E.74.2) on the stated domain. The coordinate-forgetting homomorphism is the literal map from the Heisenberg law

$$(v, c)(w, e) = (v + w, c + e + 2\det(v, w)) \longmapsto v + w \tag{E.74.6}$$

over $\mathbf{F}_3$. E.69.5 identifies its restriction to L with the particular E.67 abelianization map. E.71 retains this same old coordinate, so (E.74.3) holds for every source word. This already places the common image in the fibre product in (E.74.4).

Literal common-kernel generators. The E.69 certificate supplies three words z_0, z_1, z_2, each the commutator of two actual words in N'. In the graph

$$\widetilde{\mathcal{G}} = \{(u, v, t, A(t), B(v), B(u)) : u, v, t \in H_3\}, \tag{E.74.7}$$

their values are the three successive positive central basis vectors, with trivial Mordell values. In particular $z_j \in N'' \subset J$, and these three values form a basis of $\ker\varepsilon$ over $\mathbf{F}_3$.

Substitute the saved word DAGs into the same 97 literal N-generators in the original 11 691-point E.71 source, before its retained-coordinate projection. Their exact permutation orders and the powers used are

word	z_0	z_1	z_2
order on the full E.71 source	4	1	1
retained power	4	1	1.

(E.74.8)

The order is obtained from every cycle of each full permutation. The finite certificate separately evaluates these words in the actual twelve groups and compares their values with the saved E.69 central basis. Its derived-depth check follows the word operations, so their membership in N'' is retained as well.

Put $k_0 = z_0^4$, $k_1 = z_1$, $k_2 = z_2$. Their full E.71 permutations are identity. Since their Gaussian central values have order three and $4 \equiv 1 \pmod 3$, they keep exactly the same three central basis values. Thus

$$\rho(k_j) = 1, \qquad \langle \widetilde{\Theta}(k_0), \widetilde{\Theta}(k_1), \widetilde{\Theta}(k_2) \rangle = \ker \varepsilon. \tag{E.74.9}$$

These are images of the displayed actual words, rather than generators assigned only in an abstract central group.

For any $(p, l) \in P \times_M L$, choose $u \in J$ with $\rho(u) = p$. Equation (E.74.3) gives $\widetilde{\Theta}(u)^{-1} l \in \ker \varepsilon$. A product of the k_j realizes this difference. Multiplying u on the right by that product retains p and gives exactly l. This proves (E.74.4), including its source map. Its order is $|P|\,|\ker \varepsilon| = 2^{105} 3^{20}$.

The entire finite locus and its word preimages. The Gaussian frame enumeration of E.69 has exactly three allowed central-frame values over each of the four E.67 Gaussian values. In applying this enumeration, the required correction is computed from the absolute target t as $\widetilde{\Theta}(f_i)^{-1} t$ in the actual twelve-group product. Its image under the central-coordinate-forgetting map is the E.67 correction already attached to the E.71 row. Thus it is compatible with that row's P coordinate in (E.74.4). Equations (E.74.8)–(E.74.9) realize all three choices while preserving every E.71 coordinate. Conversely, every permitted joint value projects to one of the exhaustive 18 432 E.71 values and to one of those three frame targets. This proves both completeness and the count 55 296.

The second program substitutes the central words into the unchanged 97-generator E.71 word DAG. It first replays every ancestor needed for the 18 432 preceding witnesses in the twelve actual groups. For each absolute target, it then computes the central difference by the complete group law (E.74.6), chooses the corresponding product of the k_j, and checks the resulting word value. It also independently replays the three powered words on all 11 691 original E.71 points. Thus retention of the preceding finite values is checked in the same full source used for (E.74.8). Every added word lies in N'', so the exact domain J of the old witnesses is preserved. Equality of two joint word values is exactly membership of their quotient in the two displayed kernels; this proves (E.74.5).

The two mathematical programs in the code directory share the prefix `a_cm_matrix_gaussian_`; their suffixes are `join.py` and `witnesses.py`. Their JSON and NPZ outputs retain all source indices, all three powered words, every joint target and the complete 55 296-word DAG family. The initial unsigned residue-conversion failure is preserved separately; its failed run produced no accepted mathematical output. No further Gaussian Frattini coordinate or full-GT lift is assigned by this finite-layer calculation. □