LM7 PUSH: arithmetic sections and profinite images

Research manuscript with an explicit statement register

AI-produced by GPT-6 Astra (OpenAI, ChatGPT Pro mode)

13 November 2026

Abstract

We compute the intrinsic class-two section obstruction for the Fermat cubic with its nine coordinate cusps removed, including its integral residue complex, path origins, restricted convergence, and lifting fibers. We construct an actual obstruction invisible locally and at every finite coefficient level; refute the two proposed class-two inertia restrictions; and prove the pro-three cyclotomic centralizer criterion. Finite arithmetic field and cocycle constructions give the exact full, metabelian and marked-depth images and all their fibers, with compatibility for arbitrary ramification support. An explicit section at seventeen has the literal local cusp restriction and a global class-three lift, but no class-four lift. We identify the inherited quotient maps and every marked coefficient layer integrally. Over every number field we give the complete arithmetic obstruction test, restriction and descent maps, and the class-two germ classification, including finite algebraic formulas for every point's central coordinates. The three requested full local profinite fixed groups are computed through an explicit valuation and stabilizer comparison. Parts A, B, C and F are complete. The remaining full profinite image questions in Parts D and E are under investigation. We compute the elementary abelian peripheral lifting obstruction, its exact covering restriction, and coupled finite central counterexamples. Every finite A_5 chief extension has a positive generating-lift count preserved under the specified peripheral base maps. A split A_6 chief extension has an ordinary generating-lift obstruction, with every centralizer choice retained; explicit outer-image conditions give complementary lifting theorems. The joint binary and ternary degree-six braid refinement retains all twenty-four CM candidates and 864 finite points, all of which pass the full-target periodic-braid conjugacy tests. The A_5 and $\mathrm{SL}_2(\mathbb{F}_5)$ Ferrari refinements retain every CM candidate, with the latter retaining all 864 points. All 432 multiplier-one points lift through the full two-adic and three-adic Fermat boundary class-two towers, with exact $\mathbb{Z}_2^{10}$ and $\mathbb{Z}_3^{10}$ correction fibres, retaining their degree-six pentagons and generating maps. Explicit finite corrections at both primes fail at the next precision, while every CM fibre remains nonempty. The same word gives compatible identity shadows on every full pro-p boundary factor for $p \geq 5$. Every prescribed full boundary family also has a simultaneous generating lift with the double-cover Ferrari refinement through the explicit exponent-twenty comparison. An ordinary order-five peripheral shadow has no generating lift across a specified split A_6^{20} chief extension, for any of its four nonidentity powers, with the admitted pro-five comparison retained. The exponent-three class-three boundary quotient imposes four new congruences on the complete three-adic class-two tower and excludes an explicit tower point, while retaining every CM fibre. The complete A_6 kernel and outer-coordinate tests retain all 864 points. A five-point Hurwitz quotient has pure and full image A_{96}; its complete source and explicit exponent-ten boundary comparison retain all those points and the order-five rotation conjugacy. Explicit fully invariant finite five-group targets carry all three compatible ordinary weight-five special parameters, including their exact geometric symmetry. The complete integral three-adic class-three boundary fibre is $\mathbb{Z}_3^{70}$; its exact exponent-three image has 3^{25} points per retained source and excludes a specified finite shadow. Every generating peripheral lift in a universal coprime Frattini cover is parameterized, with its full ordinary centralizer union. The five-strand $\mathrm{Sp}_4(\mathbb{F}_3)$ test retains every candidate through its actual G_6 source quotient.

This working manuscript is **INCOMPLETE**; the task register identifies every unresolved item. We claim no priority.

Contents

1 A. The intrinsic class-two problem **4**
1.1 The rank-forty-four residue complex 4
1.2 An actual intrinsic obstruction invisible at all finite levels 7
1.3 Two ramification assertions disproved 9
1.4 The lift torsor and a formula retaining all path origins 10
1.5 An exact two-prime decomposition at seven 18
1.6 A strict restriction on ramification in the higher quotients 23
1.7 Complete global images and fibers for one auxiliary prime 29
1.8 Cofinal higher quotients and the first further coefficient 69
1.9 Explicit nine-torsion covers at the first auxiliary Frobenius 72

2 B. Geometric classes and centralizers **76**
2.1 Geometric classes and the cyclotomic centralizer 76

3 C. Restriction and germs **103**
3.1 Restriction, descent, and persistent obstruction germs 103
3.2 Algebraic evaluation of the point image 110
3.3 Separating the point germs by the nine tangent lines 115
3.4 Finite arithmetic matrices over an arbitrary number field 118

4 D. Profinite leading terms **122**
4.1 Sources, comparison maps, and the two remaining image problems 122
4.2 What the linear and literature checks establish 123
4.3 A complete finite pentagon comparison in weight nine 126
4.4 The arithmetic lower bound with its exact coordinate comparison 130
4.5 The full-source symmetry restriction in weight five 132
4.6 What a finite leading congruence detects 133
4.7 Exact peripheral lifting across abelian chief factors 135
4.8 A nonzero spin obstruction and its exact pro-ℓ fibre 138
4.9 A prime-to-five central obstruction that survives every five-adic unit power . 140
4.10 A coupled perfect kernel with admitted and obstructed pro-five branches . . . 144
4.11 A proved family of nonabelian chief lifts 147
4.12 All A_5 chief extensions over finite odd-order groups 151
4.13 Twisted A_5 chief extensions over arbitrary finite bases 155
4.14 An A_6 chief obstruction with prescribed base conjugators 160
4.15 An ordinary peripheral obstruction at a split A_6 chief step 163
4.16 Odd-order ordinary shadows at a single A_6 chief step 167
4.17 Exact ordinary lifts for cyclic and two-group outer images 168
4.18 An ordinary order-five obstruction at a split A_6^{20} chief step 170
4.19 Characteristic finite targets for the three weight-five special directions 177
4.20 Class-preserving weight-five shadows and the odd-base A_6 comparison 181
4.21 The complete peripheral fibre in a universal coprime Frattini cover 183
4.22 Equivariant intersection forms at every coprime prime-adic precision 187
4.23 The full relative pro-p cover over an odd finite base 190
4.24 Matched boundary circles and the coprime cover over an arbitrary finite base 195

4.25 The relative pro-p fibre over an infinite coprime base 200
4.26 Coprime modular covers with the old gentle conjugator retained 204
4.27 Cyclic prime outer actions and the full $\mathrm{PSL}_2(32)$ chief step 210
4.28 An exact return-prime cover retaining the entire old pair 218
4.29 All finite cyclic outer groups, with their actual automorphism extensions . . . 222
4.30 The full integral affine return tower 228

5 E. The complex-multiplication coset 232
5.1 The finite CM calculation and its independent verification 232
5.2 The homology refinements and an additional finite quotient 234
5.3 Finite tests from the full GT distribution theorem 236
5.4 The requested S_3-core refinement 237
5.5 A finite exclusion of all twelve recorded extra shadows 238
5.6 The exponent-sixteen fibres and the two-adic lifting argument 246
5.7 A four-strand pentagon constraint on the CM fibres 251
5.8 Arbitrary relative kernels supported away from two and three 256
5.9 A joint Ferrari test retaining cubic powers and fourth-degree commutators . . 258
5.10 A five-point Hurwitz quotient and its complete correction source 327
5.11 The full A_6 Ferrari source, its solvable layers, and the cubic Kummer comparison 338
5.12 A five-strand symplectic test with its exact old-source comparison 344
5.13 The exact prime-five five-strand source and its signed common quotient . . . 347
5.14 The integral source through every modulus prime to six 357
5.15 The uniform signed quotient and the full congruence correction 362

6 F. Derived limits and applications 366
6.1 The residue complex and the derived-limit test 366
6.2 Sections over the Laurent-series field 372
6.3 The fixed-lift compactness question in Task F.4 375

A Coverage and remaining questions 395

Conventions and scope

All cohomology of profinite groups with finite discrete coefficients is continuous. For a free finite-rank $\mathbb{Z}_\ell$-module with continuous action, H^i_{cts} is the cohomology of continuous inhomogeneous cochains. Restriction, inflation, quotient, and comparison maps are identified in each statement. Section sets are taken modulo conjugation by the specified geometric kernel. Arithmetic Frobenius is used in residue complexes. We work with

$$E:\ x^3+y^3+z^3=0, \qquad U=E\setminus D,$$

where D is the set of nine coordinate cusps. Put $K_0=\mathbb{Q}(\zeta)$, $\zeta^2+\zeta+1=0$, and $\lambda=1-\zeta$. Our cusp labels are

	rational cusp	first conjugate	second conjugate
$x=0$	$P=[0:-1:1]$	$A=[0:-\zeta:1]$	$\bar{A}=[0:-\zeta^2:1]$
$y=0$	$Q_c=[-1:0:1]$	$B=[-\zeta:0:1]$	$\bar{B}=[-\zeta^2:0:1]$
$z=0$	$O=[1:-1:0]$	$C=[1:-\zeta:0]$	$\bar{C}=[1:-\zeta^2:0]$.

The geometric group N is the free pro-three group of rank ten of U, based at the unit tangent $s=z/x$ at O. Its abelianization and the proper elliptic quotient are $V=T_3J_{E,D}$

and $T = T_3E$. The cusp submodule is $W = \mathbb{Z}_3(1)[D]/\mathbb{Z}_3(1)\delta$, with $\delta = \sum_{d\in D} d$; its inertia vectors are e_d. Let $I = \ker(N \to T)$ and

$$N^{\mathrm{cc}} = N/[N, I], \qquad \mathcal{K} = \ker(\bigwedge^2 V \to \bigwedge^2 T).$$

Thus $\mathcal{K}$ has rank 44 and is the central kernel of $N/\gamma_3 N \to N^{\mathrm{cc}}$. The marked subgroup I_{mark} is the closed normal subgroup generated by inertia at Q_c. A section class over L is the continuous nonabelian H^1 of G_L with the displayed geometric quotient. The full cyclotomic character, with target $\widehat{\mathbb{Z}}^\times$, is used in the centralizer statements; its three-adic component acts on the coefficient lattices.

The inherited group H has underlying set $\mathbb{Z}_3^3$ and law

$$(u, v, w)(u', v', w') = (u + u', v + v', w + w' + \tfrac{9}{2}(uv' - vu')),$$

with arithmetic weights $(\chi_3, \chi_3, \chi_3^2)$. Its base-origin comparison is made explicitly in B.9. The use of the cuspidally central lifting theorem over $\mathbb{Q}$ is Borne–Emsalem–Stix, Theorem 5.10 [BES15], applied to the nine torsion cusps and followed by the displayed pro-three quotient. Its proof uses vanishing of relative Brauer groups in all section neighbourhoods. The point O gives a retraction on the Brauer group of E itself; it does not supply those retractions for every such neighbourhood. Part C consequently retains the additional arithmetic terms over a general number field. For the profinite problems, $\widehat{F}_2$ is the full free profinite group on two peripheral generators; the special outer group and its $\mathfrak{S}_3$ centralizer $\widehat{\mathrm{GT}}_0$ use all finite quotients, with the latter defined by the two hexagons. The subgroup $\widehat{\mathrm{GT}}$ also satisfies the pentagon. Parts D and E specify every comparison to a pro-ℓ, graded, or finite quotient.

The background labels "LM7", "CK", and "F17" refer to the attached previous returns, and are checked only at the steps actually used. The available manuscript source is the supplied transcription of *La Longue Marche*, cote 140-4. The authoritative scans were not attached. Our manuscript citations identify their section, folios, scans, and formulas without asserting a fresh comparison to the scans.

1 A. The intrinsic class-two problem

1.1 The rank-forty-four residue complex

We put $K_0 = \mathbb{Q}(\zeta)$, reserving $\mathcal{K}$ for the rank-forty-four commutator lattice. Set $S = \{3, \infty\}$ and $X = \operatorname{Spec}\mathbb{Z}[1/3]$. Arithmetic Frobenius is used throughout. For each $q \neq 3$ choose a decomposition group and write

$$A_q = \mathcal{K}(-1), \qquad D_q = F_q - 1 \in \mathrm{End}_{\mathbb{Z}_3}(A_q).$$

The subscript records the local Galois action; it does not change the underlying lattice. Define

$$\mathcal{R}(A) = \{(x_q)_{q\neq 3} : x_q \in A_q, \quad \forall n \quad x_q \in 3^n A_q \text{ for almost all } q\}, \qquad D(x)_q = D_q x_q.$$

All quotients by $D\mathcal{R}(A)$ below use the actual image.

Statement A.1 (Proved). The intrinsic coefficient module and its residue complex. *The finite locally constant sheaves $\mathcal{K}_n = \mathcal{K}/3^n$ on X satisfy*

$$H^i(X, \mathcal{K}_n) = 0 \quad (i \geq 2), \qquad |H^1(X, \mathcal{K}_n)| = 3^{23n}|H^0(X, \mathcal{K}_n)|. \tag{A.1.1}$$

Their degree-one cohomology groups are finite. For every finite set J of primes different from three there is a compatible distinguished triangle in the derived category of abelian groups

$$R\Gamma(X, \mathcal{K}_n) \longrightarrow R\Gamma(X \setminus J, \mathcal{K}_n) \longrightarrow \bigoplus_{q\in J} R\Gamma(\mathbb{F}_q, \mathcal{K}_n(-1))[-1] \longrightarrow R\Gamma(X, \mathcal{K}_n)[1]. \tag{A.1.2}$$

Every D_q is injective and has finite cokernel. Localization and continuous cochains induce isomorphisms

$$H^2_{\mathrm{cts}}(G_{\mathbb{Q}}, \mathcal{K}) \simeq \mathcal{R}(A)/D\mathcal{R}(A), \qquad H^1_{\mathrm{cts}}(G_{\mathbb{Q}}, \mathcal{K}) \simeq H^1_{\mathrm{cts}}(G_S, \mathcal{K}). \tag{A.1.3}$$

The latter group is finitely generated of rank 23*; its torsion is $H^0(G_S, (\mathcal{K} \otimes \mathbb{Q}_3)/\mathcal{K})$. This is an exact residue calculation, including the contribution over $\mathbb{Z}[1/3]$ in the degrees relevant to lifting sections.*

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379).

Proof. Every coordinate cusp of the Fermat cubic is a flex. Its tangent line cuts out three times that cusp; the ratio of two tangent lines has divisor three times their difference. The classifying homomorphism from degree-zero cusp divisors to E therefore has image in $E[3]$. The extension $0 \to W \to V \to T \to 0$ is killed by three, and

$$V \otimes \mathbb{Q}_3 \simeq (W \otimes \mathbb{Q}_3) \oplus (T \otimes \mathbb{Q}_3) \tag{A.1.4}$$

as continuous Galois representations. The comparison is the splitting of the pullback along $[3] : E \to E$ followed by division by three on the Tate module. Thus the integral lattice is not being replaced without its explicit isogeny comparison.

Over K_0 all nine points of $E[3]$ are the nine cusps. Hence both $W/3$ and $T/3$ have trivial action over K_0. The action on $V/3$ is an extension of trivial modules, so its finite image is a three-group. The congruence kernel in $\mathrm{GL}(V)$ is pro-three. Consequently the image on every $\mathcal{K}_n$ over $G_{K_0,S}$ is a finite three-group. Any finite module for a finite three-group has a composition series whose factors are trivial $\mathbb{F}_3$ modules: a nonzero invariant exists on every nonzero $\mathbb{F}_3$ representation, and induction applies to its quotient.

The ring $\mathbb{Z}[\zeta]$ is norm-Euclidean. Its units are μ_6 and its unique prime above three is $\lambda = 1 - \zeta$. Kummer theory therefore gives

$$H^1(G_{K_0,S}, \mathbb{F}_3) = \langle \zeta, \lambda \rangle \simeq \mathbb{F}_3^2.$$

Indeed valuations outside λ of an allowed Kummer element are multiples of three, and unique factorization removes them. The two remaining classes are independent, first by valuation and then by the unit group. Milne's Euler characteristic, I.5.1 [Mil06], is $1/3$ for this imaginary quadratic field and the trivial module $\mathbb{F}_3$. Since H^0 has order three and H^1 order nine, H^2 has order one. Devissage along the composition series just proved gives $H^2(G_{K_0,S}, \mathcal{K}_n) = 0$. The order-two quotient $\mathrm{Gal}(K_0/\mathbb{Q})$ has exact invariants on three-primary modules, so Hochschild–Serre gives the same vanishing over G_S. Milne I.4.10(c) gives vanishing in degrees at least three, and I.4.15 gives finiteness. Proposition II.2.9 identifies these Galois groups with the cohomology of the locally constant sheaves on X. In particular no inference from the cohomological dimension of a maximal pro-three quotient is needed here.

Complex conjugation has signs $(4, 6)$ on V: its permutation signs on $\mathbb{Z}_3[D]/\mathbb{Z}_3\delta$ are $(5, 3)$, the toric twist interchanges them, and its signs on T are $(1, 1)$. The signs on $\bigwedge^2 V$ are $(21, 24)$ and the quotient $\bigwedge^2 T = \mathbb{Z}_3(1)$ removes one minus sign. Thus the signs on $\mathcal{K}$ are $(21, 23)$. Milne I.5.1 over $\mathbb{Q}$ now gives (A.1.1). The same computation, passage to the inverse limit, and $H^0(G_S, \mathcal{K} \otimes \mathbb{Q}_3) = 0$ give the asserted rank and torsion description. Finite generation follows, for example, from the finite number of generators of the relative pro-three Galois group used in LM7 E.34; its cocycles are a closed submodule of a finite power of $\mathcal{K}$. Alternatively the finite groups in (A.1.1), the vanishing of H^2, and completeness give the same conclusion by successive lifting of a finite basis modulo three.

The localization sequence and purity give (A.1.2). At a strict Henselian trait of residue characteristic $q \neq 3$, the wild inertia is pro-q and tame three-inertia is $\mathbb{Z}_3(1)$; its degree-one

cohomology is the twist $\mathcal{K}_n(-1)$ and its higher cohomology vanishes. This computes the purity shift and the residue-field action, rather than merely naming a local support group.

Take the direct limit over finite J. A continuous cochain with finite coefficients factors through a finite quotient and is unramified outside finitely many primes. The generic term is therefore $R\Gamma(G_{\mathbb{Q}}, \mathcal{K}_n)$. At each prime the residue complex is

$$[A_q/3^n \xrightarrow{D_q} A_q/3^n]$$

in degrees zero and one. The transitions of both direct-sum terms are surjective. Their derived inverse limit is consequently the literal complex $[\mathcal{R}(A) \xrightarrow{D} \mathcal{R}(A)]$. Jannsen (1.6) and (2.2) [Jan88] identify the other derived limits with continuous cochains; the finite groups $H^1(X, \mathcal{K}_n)$ have zero first derived limit. Thus the contribution of X in degrees two and three vanishes.

For injectivity of D_q, write $P_D = \mathbb{Z}_3[D]/\mathbb{Z}_3\delta$. The rational graded pieces of $\mathcal{K}(-1)$ are

$$\mathbb{Q}_3(1) \otimes \bigwedge^2 (P_D \otimes \mathbb{Q}_3), \qquad (T \otimes \mathbb{Q}_3) \otimes (P_D \otimes \mathbb{Q}_3).$$

The first has Frobenius eigenvalues of complex absolute value q; the second has absolute value $\sqrt{q}$. The cusp permutation has finite order and the elliptic Frobenius polynomial is $X^2 - a_q X + q$. Neither list contains one. An integral endomorphism invertible over $\mathbb{Q}_3$ is injective with finite cokernel. The degree-two and degree-one parts of the limiting triangle now give (A.1.3). The torsion statement follows from the exact sequence $0 \to \mathcal{K} \to \mathcal{K} \otimes \mathbb{Q}_3 \to (\mathcal{K} \otimes \mathbb{Q}_3)/\mathcal{K} \to 0$. □

Statement A.2 (Proved). The exact first derived limit and Lead 5. *Jannsen's degree-two sequence for $\mathcal{K}$ is*

$$0 \longrightarrow \frac{\mathcal{R}(A) \cap \prod_{q \neq 3} D_q A_q}{D\mathcal{R}(A)} \longrightarrow \frac{\mathcal{R}(A)}{D\mathcal{R}(A)} \longrightarrow \varprojlim_n \bigoplus_{q \neq 3} \operatorname{coker}(D_q \bmod 3^n) \longrightarrow 0. \tag{A.2.1}$$

Its left term is $\varprojlim_n {}^1 H^1(G_{\mathbb{Q}}, \mathcal{K}_n)$ and is nonzero. It is torsion-free as a $\mathbb{Z}_3$-module. In particular Lead 5 is true.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379).

Proof. The finite-level localization sequence from A.1 gives an exact sequence

$$0 \to H^1(G_S, \mathcal{K}_n) \to H^1(G_{\mathbb{Q}}, \mathcal{K}_n) \to \bigoplus_q \ker(D_q \bmod 3^n) \to 0.$$

The first groups are finite. Taking first derived limits identifies the desired group with the first derived limit of the last tower. Apply inverse limits to the kernels, images, and cokernels of the surjective-term complexes used in A.1. The image towers have surjective transitions. Their limits are precisely $\mathcal{R}(A) \cap \prod_q D_q A_q$, which proves (A.2.1).

Choose for each j a new prime q_j splitting in the finite Galois extension cut out by $\mathcal{K}(-1)/3^j$. Chebotarev [Mil20b, V.3.23] supplies infinitely many choices. Then $D_{q_j} A_{q_j} \subset 3^j A_{q_j}$. Choose a primitive $v_j \in A_{q_j}$ and let $z_{q_j} = D_{q_j} v_j$, with all other entries zero. The family z is restricted and pointwise in the image of D. Its only preimage is v_j at the chosen primes, since each D_q is injective; this preimage is not restricted. Hence $[z] \neq 0$ in the left term. For torsion-freeness, write a left-term representative as $z = Dv$ with $v_q \in A_q$. A nonzero scalar $m \in \mathbb{Z}_3$ annihilates its class only when mv is restricted. Multiplication by a fixed nonzero scalar changes all valuations by a fixed integer, so then v is restricted. □

1.2 An actual intrinsic obstruction invisible at all finite levels

Let c_d denote the central inertia generators in $A = I/[N, I] = \mathbb{Z}_3(1)[D]$, and let e_d be their images in W. The rational cusps O and P define a continuous equivariant homomorphism

$$\ell : \mathbb{Z}_3(1) \longrightarrow N^{\mathrm{cc}}, \qquad z \longmapsto c_O^z c_P^z. \tag{A.3.1}$$

Its image is central. Pull back the central quotient $N/\gamma_3 N \to N^{\mathrm{cc}}$ along ℓ and call the resulting geometric group E_w, where $w = e_O + e_P$.

Statement A.3 (Proved). A linear family inside the nonabelian lifting problem. *The pullback is an exact sequence of continuous Galois lattices*

$$0 \longrightarrow \mathcal{K} \longrightarrow E_w \longrightarrow \mathbb{Z}_3(1) \longrightarrow 0. \tag{A.3.2}$$

Choose an integral lift l of the standard generator, and after twisting put

$$u_q = F_q(l \otimes 1) - l \otimes 1 \in A_q. \tag{A.3.3}$$

For $a = (a_p)_p \in \mathcal{R}_{\mathbb{P}}(\mathbb{Z}_3)$, the section $\ell \circ \kappa(a)$ of $N^{\mathrm{cc}} \rtimes G_{\mathbb{Q}}$ has obstruction represented, under (A.1.3), by

$$(a_q u_q)_{q \neq 3}. \tag{A.3.4}$$

Its unique possible local corrections are $-a_q D_q^{-1} u_q$. Thus it lifts precisely when these corrections are integral and form a restricted family. Changing l changes (A.3.4) by D of a restricted family. For the families used below, $a_3 = 0$. In E24's pro-three coordinates and the E36 generator $k = -\delta$, their actual torsor coordinates are

$$\alpha : \quad (r_1, r_2; A, B, C) = (0, -\kappa(a); -\kappa(a), -\kappa(a), -\kappa(a)), \qquad a_{\mathrm{E36}} = -a. \tag{A.3.5}$$

The three K_0 coordinates use the inclusion of rational Kummer classes; the other torsion, framed-λ, and divisible coordinates are zero. This specifies the pairs themselves in the requested convention.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379).

Proof. The abelian image of a lift of z is zw. The commutator of two lifts is therefore $(zw) \wedge (z'w) = 0$ in $\bigwedge^2 V$. The kernel $\mathcal{K}$ is central, so E_w is abelian. It is compact, torsion-free, and finitely generated over $\mathbb{Z}_3$; the quotient is free of rank one. This proves (A.3.2) and supplies an integral lift. All actions factor through G_S by smooth reduction of the pointed curve outside three.

The obstruction is now the ordinary connecting homomorphism $H^1(G_{\mathbb{Q}}, \mathbb{Z}_3(1)) \to H^2(G_{\mathbb{Q}}, \mathcal{K})$ for (A.3.2). On local tame inertia, $\kappa(a)(\tau_q) = a_q$. In $E_w(-1)$ the failure of its chosen lift to be Frobenius invariant is $a_q u_q$. Equivalently, apply the connecting morphism to the two-term residue complexes of A.1; the degree-zero element a_q in the trivial quotient has boundary $a_q u_q$ in the degree-one kernel. This proves (A.3.4) at the cochain level and hence before passage from the restricted quotient to a product of local groups. The bounded integral vectors u_q and restricted scalars a_q give a restricted representative. The final assertion follows from $u_q \mapsto u_q + D_q v$ for a fixed integral change of lift, with $(a_q v)_q$ restricted. For (A.3.5), in the non-O cusp basis the vector w has P-coordinate zero and all other coordinates minus one. Formula (E24.4) sends it to $(0, -1)$; (E24.5) gives the three equal negative rational Kummer classes. Its lift with O-coordinate zero differs from $c_O + c_P$ by δ. Since $k = -\delta$, the E36 parameter is $-a$, as asserted. □

Statement A.4 (Proved). A nonzero intrinsic obstruction in the first derived limit. *There is a continuous section of $N^{\mathrm{cc}} \rtimes G_{\mathbb{Q}}$ whose obstruction to $N/\gamma_3 N$ is nonzero in $\varprojlim{}^1_n H^1(G_{\mathbb{Q}}, \mathcal{K}_n)$. Its restriction to every local field is zero, and its image in every finite-coefficient obstruction group is zero. The construction below specifies such a section as $\ell \circ \kappa(a)$, hence specifies an actual pair of the torsor coordinates of R5 after the chosen origins are fixed. Its obstruction has infinite additive order and survives restriction to every number field.*

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379).

Proof. We first prove that the extension (A.3.2) has a nonzero unipotent action in the kernel of the action on $\mathcal{K}(-1)$. Let s_P be the rational tangential section at P, with parameter x/z and tangent one. Write $s_P = b_P s_O$ and also b_P for its abelian cocycle in V. For an inertia generator transported along the chosen path,

$$g(c_P) = b_P(g)^{-1} c_P^{\chi(g)} b_P(g), \qquad g(c_O) = c_O^{\chi(g)}. \tag{A.4.1}$$

The abelian tangential path is the Kummer class of the framed line bundle $z_P = \mathcal{O}(P - O)$ whose frames are $(x/z)^{-1}$ at P, z/x at O, and the canonical section at the other cusps. Indeed the Abel–Jacobi divisor $u - u_0$, as both points approach the indicated unit tangents, has exactly these two regularized frames. The Kummer comparison follows by pulling back multiplication by 3^n on the generalized Jacobian, which realizes the geometrically abelian covering with group $V/3^n$. This identifies the based path cocycle, not just its projection to the proper curve.

For completeness the needed framed cube is calculated directly. The function $f = (y+z)/(x+y)$ has divisor $3P - 3O$. Its regularized values are $1/3$ at P, 3 at O, and -1 at Q_c. Evaluating at the three quadratic cusp pairs and dividing by the value at O gives

$$3z_P = (3^{-2}, -3^{-1}, \lambda^{-1}, \lambda^{-2}, \zeta^2\lambda^{-3}). \tag{A.4.2}$$

Here the entries correspond to P, Q_c and the pairs denoted $A, \bar{A}$, $B, \bar{B}$, $C, \bar{C}$ in LM7 E.24. The identity $\lambda^2 = -3\zeta$ verifies the values and their conjugates. These are the origin terms needed in this proof.

The action on $V \otimes \mathbb{Q}_3$ over K_0 is abelian. The toric action is cyclotomic; on the elliptic factor the endomorphism induced by a coordinate multiplication by ζ makes T_3E a rank-one module over the discrete valuation ring $\mathbb{Z}_3[\zeta]$. The Galois action commutes with this endomorphism and thus lies in its commutative unit group. The splitting (A.1.4) proves the assertion for V; since the integral lattice embeds in its rationalization, commutators act trivially on the integral lattice too.

Choose $\tau \in \ker \chi_3$ with $\kappa_3(\tau)$ a unit. Such an element exists: $\mathbb{Q}(\sqrt[3]{3})$ is not contained in $\mathbb{Q}(\mu_{3^\infty})$, since every finite subfield of this abelian extension is Galois over $\mathbb{Q}$, whereas the real irreducible pure cubic is not normal. Choose $\sigma \in G_{K_0}$ with $\chi_3(\sigma) = 4$, and put $g = [\sigma, \tau]$. The Kummer homomorphism into $\mathbb{Z}_3 \rtimes \mathbb{Z}_3^\times$ gives

$$\chi_3(g) = 1, \qquad \kappa_3(g) = 3\kappa_3(\tau). \tag{A.4.3}$$

Thus g fixes V and $\mathcal{K}(-1)$. On this subgroup, $\kappa_\zeta(g) = 0$ and $\kappa_\lambda(g) = \kappa_3(g)/2$; the same holds for the conjugate of λ. Evaluating (A.4.2) on g, where coboundaries in V vanish, gives

$$3b_P(g) = \kappa_3(g) v_0, \qquad v_0 = -2e_P - e_{Q_c} - \tfrac{1}{2}(e_A + e_{\bar{A}}) - (e_B + e_{\bar{B}}) - \tfrac{3}{2}(e_C + e_{\bar{C}}). \tag{A.4.4}$$

The notation is in $W \otimes \mathbb{Q}_3$; all denominators in v_0 are units in $\mathbb{Z}_3$. Take $l = \log c_O + \log c_P$ in the class-two Lie lattice, whose Baker–Campbell–Hausdorff denominator two is a three-adic unit. Equation (A.4.1) gives

$$u_g = -b_P(g) \wedge e_P = -\kappa_3(\tau) v_0 \wedge e_P. \tag{A.4.5}$$

The coefficient of $e_P \wedge e_{Q_c}$ in $v_0 \wedge e_P$ is one. The eight non-O cusp classes form a basis of W, and W is a primitive sublattice of V. Therefore (A.4.5) is primitive in $\mathcal{K}$. A change of based path contributes $(g-1)v = 0$; a change of tangent at P adds a multiple of e_P to b_P and has zero wedge with e_P. This verifies the exact ambiguity.

Apply Chebotarev [Mil20b, V.3.23] to the finite representation $E_w(-1)/3^j$. Choose recursively a new prime q_j whose Frobenius belongs to the conjugacy class of g in this representation. Then

$$D_{q_j} \equiv 0 \pmod{3^j}, \qquad u_{q_j} \notin 3A_{q_j}. \tag{A.4.6}$$

Conjugacy preserves these properties: the restriction to the kernel is the identity modulo 3^j, and conjugation carries the off-diagonal vector through an invertible integral matrix. Define the finite integer

$$e_j = \min\{e \geq 0 : 3^e D_{q_j}^{-1} u_{q_j} \in A_{q_j}\}. \tag{A.4.7}$$

It exists by A.1, and (A.4.6) implies $e_j \geq j$. Set $a_{q_j} = 3^{e_j}$ and set every other a_p to zero. This is a restricted Kummer family. Its residues are pointwise in $D_q A_q$; their unique corrections $-3^{e_j} D_{q_j}^{-1} u_{q_j}$ are primitive by the minimality in (A.4.7). They do not tend to zero. A.3 and A.2 prove that its obstruction is a nonzero first-derived-limit class.

At each coefficient level only finitely many residue entries survive. Each is a boundary, so the finite-coefficient obstruction is zero. At every local field, the finite degree-one cohomology groups satisfy Mittag–Leffler; continuous degree-two cohomology therefore injects into the limit of finite degree-two groups. This proves local vanishing also at three. Real three-primary cohomology vanishes by averaging. Finally A.2 proves infinite additive order. Restriction followed by corestriction over a finite extension $L/\mathbb{Q}$ is multiplication by $[L : \mathbb{Q}]$ on continuous cohomology, by the finite coset cochain formula. Hence no restriction can kill this obstruction. □

1.3 Two ramification assertions disproved

Statement A.5 (Refuted). Lead 2: mixed tame inertia already occurs at the prime two. *Lead 2 is false. The section*

$$s = \ell \circ (3^{23}\kappa_2) : G_{\mathbb{Q}} \longrightarrow N^{\mathrm{cc}} \rtimes G_{\mathbb{Q}} \tag{A.5.1}$$

lifts to $N/\gamma_3 N$. Its abelian tame-inertia vector at two is $3^{23}(e_O + e_P)$, which is not a multiple of any cusp vector. More generally, for each rational prime $q \neq 3$, every multiple of $3^{v_3 \det D_q}\kappa_q$ in the line (A.3.1) lifts.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379).

Proof. For a fixed q the only nonzero residue of this family is a scalar multiple of u_q. The adjugate identity shows that $3^{v_3 \det D_q} D_q^{-1}$ is integral. Its local correction has finite support and is therefore restricted. A.3 proves the general assertion.

At two the cusp permutation has signs $(5, 3)$ on P_D. Thus its exterior square has signs $(13, 15)$, and the determinant of $F_2 - 1$ on $\bigwedge^2 W(-1)$ has three-adic valuation 15. The cubic has the three projective points $[0:1:1], [1:0:1], [1:1:0]$ over $\mathbb{F}_2$, so its elliptic Frobenius polynomial is $X^2 + 2$. On each of the eight copies of T, twisted by a cusp permutation sign, the determinant of $F_2 - 1$ is three. The remaining contribution is eight. Consequently

$$v_3 \det D_2 = 15 + 8 = 23.$$

This determinant calculation uses the rational filtration and is valid on the actual integral lattice because determinants are unchanged by rational change of basis.

To check the cusp claim integrally, eliminate e_O by the relation $\sum_d e_d = 0$. In the basis of the other eight cusps, $e_O + e_P = (0,-1,-1,-1,-1,-1,-1,-1)$. It is proportional neither to a basis vector nor to $e_O = (-1,-1,-1,-1,-1,-1,-1,-1)$. Multiplication by a nonzero three-adic scalar preserves this fact. The exact finite calculations are independently executed in `code/a_residue_checks.py`. □

Statement A.6 (Refuted). Lead 3: infinitely ramified noncuspidal class-two sections. *Lead 3 is false. Enumerate the rational primes different from three as $q_1, q_2, \ldots$, put $m_j = v_3 \det D_{q_j}$, and set*

$$a_{q_j} = 3^{m_j + j}, \qquad a_3 = 0. \tag{A.6.1}$$

The section $\ell \circ \kappa(a)$ lifts to $N/\gamma_3 N$ and every such lift is ramified at all the q_j. None is cuspidal. The same conclusion holds after replacing these coefficients by any nonzero coefficients satisfying $v_3(a_{q_j}) - m_j \longrightarrow +\infty$.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379).

Proof. The family is restricted. By the adjugate identity, $a_{q_j} D_{q_j}^{-1} u_{q_j} \in 3^j A_{q_j}$, so the unique corrections are integral and restricted. A.3 proves existence of a continuous lift. Its image in V has nonzero tame-inertia vector $a_{q_j}(e_O + e_P)$ at every q_j; thus geometric conjugation cannot remove the ramification. A cuspidal section at a rational cusp has all abelian tame-inertia vectors in the line generated by that cusp. A.5 proves that no one of the displayed vectors lies in any such line. The nonrational cusps have decomposition groups whose image in $G_{\mathbb{Q}}$ has index two, so cannot support a section over $\mathbb{Q}$. This proves noncuspidality and the more general assertion with the stated convergence bound. □

1.4 The lift torsor and a formula retaining all path origins

Statement A.7 (Proved). The integral group acting on every nonempty fibre. *There is an isomorphism of finitely generated $\mathbb{Z}_3$-modules*

$$H^1_{\mathrm{cts}}(G_{\mathbb{Q}}, \mathcal{K}) \simeq \mathbb{Z}_3^{23} \oplus (\mathbb{Z}/3)^9 \oplus (\mathbb{Z}/9)^5. \tag{A.7.1}$$

Every nonempty fibre of $\mathrm{Sec}(N/\gamma_3 N) \to \mathrm{Sec}(N^{\mathrm{cc}})$ is a torsor under this group. The ambiguity in (A.7.1) is the choice of a Smith basis and of a free direct summand. No further geometric conjugations identify members of the torsor.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379).

Proof. We give the integral lattice and the finite calculation determining its torsion. Write $\mathcal{O}_3 = \mathbb{Z}_3[\zeta]$ and identify T with $\mathcal{O}_3$ so that P is the class of 1 in $T/3T$. The automorphism $[x : y : z] \mapsto [x : y : \zeta z]$ fixes O and acts on its tangent by ζ. It is the chosen complex multiplication. Its orbit of P is P, A_{bar}, A. The eight nonzero cusp labels in the basis $(P, \zeta P)$ are

d	P	Q_c	A	$\bar{A}$	B	$\bar{B}$	C	$\bar{C}$
a_d	1	−1	−1	0	1	0	−1	1
b_d	0	0	−1	1	1	−1	1	−1

(A.7.2)

For example the line through P and $-\zeta P$ has third intersection $[1 : -\zeta : 0] = C$, so $C = -(P - \zeta P)$. This determines the only labels not obtained immediately by multiplication by ζ and negation.

Let Ψ be the 8×2 integer matrix with row $(b_d, -a_d)$. In the rational representation $W_{\mathbb{Q}_3} \oplus T_{\mathbb{Q}_3}$ the lattice V is

$$L_\Psi = \{(w + \Psi t/3, t) : w \in W,\ t \in T\}. \tag{A.7.3}$$

Here is the comparison proving this formula. Pullback of the semiabelian extension $J_{E,D}$ along $[3] : E \to E$ is split, since its class is killed by three. The splitting is unique because $\mathrm{Hom}(E, T_D) = 0$. Thus it is defined over $\mathbb{Q}$. The resulting isogeny $T_D \times E \to J_{E,D}$ has kernel the graph of a homomorphism $E[3] \to T_D[3]$; its projection to E is $[3]$. Evaluation of the graph on a degree-zero cusp character gives the Weil pairing with that cusp difference, by the defining Poincare biextension of the generalized Jacobian. In the bases in (A.7.2), this pairing is $t \mapsto \Psi t$ modulo three. One overall sign changes Ψ to $-\Psi$ and conjugates all the matrices by $(-1)_W \oplus 1_T$, so it does not affect their invariant factors. The Tate lattice of the quotient is obtained by adding $(\Psi t/3, t/3)$ to $W \oplus T$ and then using $[3]$ on the second coordinate. This is precisely (A.7.3), including its integral comparison with the geometric Tate module.

For a Galois element with matrices R on W and T_g on T, the matrix in the basis in (A.7.3) is therefore

$$G_V = \begin{pmatrix} R & (R\Psi - \Psi T_g)/3 \\ 0 & T_g \end{pmatrix}. \tag{A.7.4}$$

The numerator is divisible by three by equivariance of the Weil pairing. Over K_0, $R = \chi(g) I_8$. Complex conjugation has $R = -P_c$, where P_c fixes P, Q_c and interchanges the three quadratic cusp pairs, and

$$T_c = \begin{pmatrix} 1 & -1 \\ 0 & -1 \end{pmatrix}.$$

The exact Tate image over K_0 is $1 + 3\mathcal{O}_3$. Containment follows from rationality of $E[3]$ and complex multiplication. To prove equality, count the Fermat cubic over $\mathbb{F}_7$: it has nine points, so its Frobenius polynomial is $X^2 + X + 7$. Its roots in $\mathcal{O}_3$ are $\alpha = 1 + 3\zeta$ and $\bar{\alpha} = -2 - 3\zeta$. Frobenius at a prime above seven gives one root; conjugation gives the other. The three-adic logarithm identifies $1 + 3\mathcal{O}_3$ with $3\mathcal{O}_3$. The vectors $\log(\alpha)/3$ and $\log(\bar{\alpha})/3$ reduce to ζ and $-1 - \zeta$ modulo three, a basis of $\mathcal{O}_3/3\mathcal{O}_3$. Their closed $\mathbb{Z}_3$-span is therefore all $\mathcal{O}_3$. This proves equality without an unspecified index from a complex multiplication reciprocity theorem. The matrices on T are

$$T_\alpha = \begin{pmatrix} 1 & -3 \\ 3 & -2 \end{pmatrix}, \qquad T_{\bar{\alpha}} = \begin{pmatrix} -2 & 3 \\ -3 & 1 \end{pmatrix}. \tag{A.7.5}$$

The determinants are seven. Equations (A.7.4)–(A.7.5) and c therefore topologically generate the actual image on V.

Take exterior squares in (A.7.4) and omit the basis wedge of the two T lifts. The remaining forty-four wedges are a basis of $\mathcal{K}$, since that omitted wedge maps to a generator of $\bigwedge^2 T$. Let the three resulting matrices be $G_\alpha, G_{\bar{\alpha}}, G_c$. The integer matrix

$$B = \begin{pmatrix} G_\alpha - 1 \\ G_{\bar{\alpha}} - 1 \\ G_c - 1 \end{pmatrix} : \mathbb{Z}_3^{44} \longrightarrow \mathbb{Z}_3^{132} \tag{A.7.6}$$

has three-adic Smith diagonal

$$\underbrace{1, \ldots, 1}_{30}, \quad \underbrace{3, \ldots, 3}_{9}, \quad \underbrace{9, \ldots, 9}_{5}. \tag{A.7.7}$$

The program `code/a_k_lattice.py` constructs every entry from (A.7.2)–(A.7.6), checks $c^2 = 1$ and $cG_\alpha c = G_{\bar{\alpha}}$, and performs elementary row and column operations over $\mathbb{Z}/3^8$. Each operation is invertible and every pivot has valuation at most two. Thus no factor divisible by 3^8 remains, and this calculation determines the exact three-adic Smith exponents. The full matrices and the output are included. As an independent consistency check, the same calculation on V gives seven unit factors and three factors three, recovering $J_{E,D}(\mathbb{Q})[3^\infty] = (\mathbb{Z}/3)^3$.

An invariant in $(\mathcal{K} \otimes \mathbb{Q}_3)/\mathcal{K}$ is exactly an x with Bx integral, modulo integral x. A Smith factor 3^e contributes $\mathbb{Z}/3^e$. A.1 and (A.7.7) prove (A.7.1). Finally fix a cocycle representing a section of N^{cc}. Central lifts differ by $\mathcal{K}$-valued cocycles and central conjugation gives coboundaries. A geometric element of N^{cc} stabilizing that section has invariant image in V, which is zero. It is then in the central kernel $\mathbb{Z}_3(1)$ and is invariant there, hence zero again. There is no additional stabilizer quotient. This proves the exact fibre statement. □

Statement A.8 (Proved). The full class-two residue criterion in based cusp-path coordinates. *Choose geometric paths from the specified tangent at O to a Galois-equivariant family of tangents at the nine cusps. Let c_d be the corresponding inertia elements in $N/\gamma_3 N$ and $b_{g,d} \in V$ their abelian comparison cochains, with convention*

$$g(\log c_d) = \chi(g)\big(\log c_{gd} - b_{g,d} \wedge e_{gd}\big). \tag{A.8.1}$$

For a continuous representative s of a class in $\mathrm{Sec}(N^{\mathrm{cc}})$*, write*

$$\log s(\tau_q) = \sum_{d \in D} t_{q,d} c_d, \quad v_q = \alpha(\phi_q), \quad w_q = \sum_d t_{q,d} e_d \in W. \tag{A.8.2}$$

Here $s(\tau_q)$ means its geometric cocycle value, the chosen arithmetic action is unramified at q, and $\phi_q \tau_q \phi_q^{-1} = \tau_q^q$. Then its full intrinsic obstruction is represented by the restricted family

$$r_q = v_q \wedge w_q - \sum_d t_{q,d}\, b_{\phi_q,d} \wedge e_{\phi_q d} \in \mathcal{K}(-1). \tag{A.8.3}$$

The section lifts exactly when

$$-D_q^{-1} r_q \in A_q \quad \textit{for all } q \neq 3, \qquad (-D_q^{-1} r_q)_q \in \mathcal{R}(A). \tag{A.8.4}$$

The nonempty fibres are exactly the torsors in A.7.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379).

Proof. The comparison cochains exist by transport of the inertia loops along the chosen paths. Composition gives $b_{gh,d} = b_{g,hd} + g b_{h,d}$ in V. Formula (A.8.1) is the class-two expansion of conjugation and includes its sign convention. It specifies the action on the cusp lifts themselves, including the path origins.

The cocycle restriction to local inertia is a homomorphism, since the original action is unramified there. Wild inertia maps to zero and the prime-to-three tame factors also map to zero. The elliptic projection t_T satisfies $F_q t_T = q t_T$. The Frobenius eigenvalues on T have absolute value $\sqrt{q}$, so $t_T = 0$. Hence the logarithm of tame inertia lies in the central lattice $A = \mathbb{Z}_3(1)[D]$ in N^{cc}, proving (A.8.2). The relation also gives $t_{q,\phi_q d} = t_{q,d}$.

Lift this inertia logarithm to $L_q = \sum_d t_{q,d} \log c_d + k_q$ in the full class-two lattice, where $k_q \in \mathcal{K}$. Lift the Frobenius cocycle arbitrarily; its abelian coordinate is v_q. In a class-two group conjugation by that lift changes a logarithm of abelian coordinate $q w_q$ by $q v_q \wedge w_q$.

The logarithm of the tame-relation defect, divided by q to identify inertia homomorphisms with the twist $\mathcal{K}(-1)$, is therefore

$$D_q k_q + v_q \wedge w_q - \sum_d t_{q,d} b_{\phi_q,d} \wedge e_{\phi_q d}.$$

The central coordinate of the Frobenius lift cancels. This is exactly the cochain-level localization boundary: it measures the failure of a chosen inertia splitting to be invariant under Frobenius. At every finite coefficient level the action and the section are unramified at almost all primes, so these choices give a direct-sum residue cochain. The inverse-limit family is restricted; concretely every term contains one of the restricted coefficients $t_{q,d}$ and all other vectors are integral. A.1 now identifies the global obstruction with $[r_q]$, before any passage to a product of local groups. Injectivity of every D_q proves (A.8.4), with the stated convergence condition. A.7 supplies the geometric-conjugacy fibre. □

Coordinates of the tame coefficients. The formulas can be connected to the chosen E24 coordinates without suppressing the central parameter. Separate those coordinates into the entries supported at three, including μ_3^3, the framed point z', and the noncanonical divisible $\mathbb{Z}_3$ parameter, and the prime valuations outside three. The elliptic projection is always unramified outside three: the same weight calculation used in A.8 makes every local integral elliptic tame vector zero. The surjectivity $H^1(G_S, V) \to H^1(G_S, T)$ follows from $H^2(G_S, W) = 0$, proved by the same composition-series argument as A.1. Thus origins can be chosen by a full unramified lift for the first set of parameters, as constructed in LM7 E.34, and by the central splitting $W \to A$ with O-coordinate zero for the remaining toric coordinates. We make this choice agree with the E36 convention as follows. For each elliptic class β, choose an unramified section c_β of C; take c_0 to be the unit tangent at O, and take the classes of $P - O$ and $Q_c - O$ to the images of their unit tangents. The reference over every α is (c_β, α) in $C \times_T V$. The prescribed unramified reference itself has a full unramified lift by the free pro-three embedding argument. The E36 central generator is $k = [a, b] = -\delta$, so the exact coordinate comparison is $a_\delta = -a_{\mathrm{E36}}$. In the formulas below a always means a_{E36}. This fixes both the origin and its sign; it agrees with the packets in B.3.

Write $r_{1,q}, r_{2,q}$ for the valuations of the two rational toric coordinates, and $\nu_{A,\mathfrak{q}}$, $\nu_{B,\mathfrak{q}}$, $\nu_{C,\mathfrak{q}}$ for those of the three K_0 toric coordinates. Put $N\nu_{A,q} = \sum_{\mathfrak{q}|q} f_{\mathfrak{q}/q} \nu_{A,\mathfrak{q}}$, and likewise for B, C. In this chosen origin, for $q \neq 3$,

$$\begin{aligned} t_{q,O} &= -a_q, \\ t_{q,P} &= r_{1,q} + N\nu_{C,q} - N\nu_{A,q} - a_q, \\ t_{q,Q_c} &= r_{2,q} + N\nu_{C,q} - N\nu_{B,q} - a_q, \\ t_{q,d} &= \nu_{A,\mathfrak{q}(d)} - a_q \qquad\qquad (d \in \{A, \bar{A}\}), \end{aligned} \tag{A.8.5}$$

with identical last formulas for the B and C pairs. At an inert prime the two valuations in a pair coincide and their norm contribution is twice that valuation. At a split prime each has residue degree one. These formulas are the valuation form of the actual embedding (E24.5), followed by minus the central diagonal $\mathbb{Z}_3(1)\delta$; the diagonal is the kernel of $A \to W$.

The cochains $b_{g,d}$ in (A.8.3) are abelian data. Let z_d be the framed tangential divisor $d - O$ in $J_{E,D}$ and choose compatible division points $z_{d,n}$ with $3^n z_{d,n} = z_d$. Then $g z_{d,n} - z_{gd,n} \in J_{E,D}[3^n]$ gives their reduction modulo 3^n, with the chosen path convention. Their origin relations are obtained by the tangent-line functions f_d with $\operatorname{div} f_d = 3(d - O)$, regularized at the indicated tangents. Formula (A.4.2) is the complete P instance and is the one needed for the proved counterexamples. Statements A.9 and A.10 below supply the complete regularized origin table and fix the extra unramified generator. Evaluating a Frobenius in that free

pro-three Galois group is part of the arithmetic input; a finite-query program on arbitrary completed families cannot decide the final convergence condition.

Statement A.9 (Proved). All nine tangential origin data, including roots of unity. *Order the non-O cusps as $P, Q_c, A, \bar{A}, B, \bar{B}, C, \bar{C}$. Use x/z at the first and third/fourth cusps, y/z at the second and fifth/sixth cusps, and z/x at the last two; all tangents are one. For $z_d = \mathcal{O}(d - O)$ with these regularized frames, write $3z_d = h_d \in T_D(K_0)$. Its O-frame is one. The other entries are $\pm\zeta^{Z_{d,e}}\lambda^{L_{d,e}}$, with matrices*

$$L = \begin{pmatrix} -4 & -2 & -1 & -1 & -2 & -2 & -3 & -3 \\ -2 & -4 & -2 & -2 & -1 & -1 & -3 & -3 \\ -1 & -2 & -4 & -1 & -2 & -2 & -3 & -3 \\ -1 & -2 & -1 & -4 & -2 & -2 & -3 & -3 \\ -2 & -1 & -2 & -2 & -4 & -1 & -3 & -3 \\ -2 & -1 & -2 & -2 & -1 & -4 & -3 & -3 \\ -3 & -3 & -3 & -3 & -3 & -3 & -6 & -3 \\ -3 & -3 & -3 & -3 & -3 & -3 & -3 & -6 \end{pmatrix}, \qquad Z = \begin{pmatrix} 2 & 1 & 0 & 1 & 0 & 2 & 2 & 1 \\ 1 & 2 & 0 & 2 & 0 & 1 & 1 & 2 \\ 1 & 2 & 2 & 0 & 1 & 0 & 2 & 1 \\ 0 & 0 & 1 & 2 & 2 & 1 & 2 & 1 \\ 2 & 1 & 1 & 0 & 2 & 0 & 1 & 2 \\ 0 & 0 & 2 & 1 & 1 & 2 & 1 & 2 \\ 1 & 2 & 1 & 1 & 2 & 2 & 0 & 0 \\ 2 & 1 & 2 & 2 & 1 & 1 & 0 & 0 \end{pmatrix}. \tag{A.9.1}$$

The signs, recorded in the exact program output, have zero pro-three Kummer cocycle, using the rational compatible odd-power roots of -1. No ζ term may be discarded. The row at O is the identity. These data are equivariant under complex conjugation.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379).

Proof. For $d = [x_d : y_d : z_d]$, put $L_d = x_d^2 x + y_d^2 y + z_d^2 z$ and $f_d = L_d/(x+y)$. The flex property proves $\operatorname{div} f_d = 3d - 3O$. At O, write $s = z/x$ and $y/x = -(1+s^3)^{1/3}$. The regularized value of f_d is

$$c_O(d) = \lim_{s\to 0} f_d s^3 = -3(x_d^2 - y_d^2).$$

For $d = [0 : -\eta : 1]$ or $[-\eta : 0 : 1]$, with $\eta^3 = 1$, the regularized value at d is $1/(3\eta)$, by expanding the nonzero coordinate as $-\eta(1+s^3)^{1/3}$. For $d = [1 : -\eta : 0] \neq O$ it is $-1/(3(1-\eta))$. At every other cusp e the ordinary value $f_d(e)$ applies. Division of all these values by $c_O(d)$ gives $h_{d,e}$. This proves the claimed framed relation over K_0. All arithmetic in the evaluation is in $\mathbb{Q}(\zeta)$; `code/a_cusp_origins.py` evaluates the formulas as rational pairs in the basis $1, \zeta$, factors each nonzero value uniquely as $\pm\zeta^a\lambda^b$, and verifies conjugation entry by entry. It also verifies the complete P row against (A.4.2). This gives (A.9.1), not a numerical approximation to its entries. □

Statement A.10 (Proved). An explicit choice of the extra unramified cohomology generator. *Let P_3 be the maximal pro-three quotient of $G_{K_0,S}$ and let $\Gamma = P_3 \rtimes \langle c \rangle$ be the relative quotient of G_S. Choose u to be Frobenius at the prime $(7, \zeta - 2)$, with a decomposition group fixed once and for all, and put $v = cuc$. Then u, v are a free pro-three basis of P_3, and c interchanges them. In the representation (A.7.3) let $\widetilde{t}_1 = (\Psi(1,0)/3, (1,0)) \in V$. The prescriptions*

$$\alpha_\infty(c) = 0, \qquad \alpha_\infty(u) = \widetilde{t}_1 \tag{A.10.1}$$

define a continuous V-valued cocycle on Γ and fix a splitting of the final $\mathbb{Z}_3$ coordinate in (E24.1). Together with the specified prime generators and the framed points, this fixes the formerly noncanonical cohomology coordinate.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379).

Proof. The vanishing $H^2(G_{K_0,S}, \mathbb{F}_3) = 0$ was proved in A.1. Inflation injects $H^2(P_3, \mathbb{F}_3)$ into this group: an invariant nonzero character of the kernel of $G_{K_0,S} \to P_3$ would give a central extension of P_3 by $\mathbb{F}_3$, a pro-three quotient contradicting maximality. The five-term exact sequence proves the injection. The minimal free pro-three presentation of P_3 has two generators, by the Kummer calculation in A.1; its relation module has dual $H^2(P_3, \mathbb{F}_3) = 0$. A nontrivial closed relation subgroup in the Frattini subgroup has a nonzero first term in the lower exponent-three filtration, so the relation module cannot vanish. Thus P_3 is free on two generators.

At the chosen prime over seven, finite-field Kummer theory gives the evaluations $(2, 0)$ on the classes (ζ, λ): $\zeta \mapsto 2$, $\lambda \mapsto -1$ in $\mathbb{F}_7$, and their squares are ζ^2 and one. At the conjugate Frobenius the values are $(2, 2)$. They are independent over $\mathbb{F}_3$. Consequently u, cuc generate the Frattini quotient of P_3; the pro-three basis theorem and freeness make them a free basis. Complex conjugation is an actual order-two element of G_S, so its relations are exactly $c^2 = 1$, $cuc = v$, and $cvc = u$.

For each finite three-primary quotient, arbitrary cocycle value on u, its prescribed conjugate value on v, and value zero on c respect all these relations. The free pro-three universal property gives a cocycle; taking the compatible inverse limit gives (A.10.1). On T the normalized cocycles are T modulo $(u - 1)T^c$. Now $T^c = \mathbb{Z}_3(1, 0)$ and $(T_\alpha - 1)(1, 0) = 3(0, 1)$, so

$$H^1(G_S, T) = \mathbb{Z}_3(1, 0) \oplus (\mathbb{Z}/3)(0, 1).$$

The projection of (A.10.1) is the free generator. The finite summand is the Kummer class of P, and filling the cusps identifies the free quotient with the Tate module of the divisible part used in (E24.9). Thus the given lift defines the asserted splitting. The Frobenius scalar at $(7, \zeta - 2)$ is $1 + 3\zeta$, rather than its conjugate: its reduction has zero differential at that prime, which selects the root divisible by the prime. This verifies the matrix used in (A.10.1). □

Statement A.11 (Proved). Executed finite and certified-tail evaluations. *The program* `code/a_criterion.py` *implements (A.8.4) for finite-support exact residue matrices and for the two infinite families A.4 and A.6 with their proved tail certificates. It gives the following verified results:*

1. *the E.28 quotient of* (κ_6, κ_{109}) *has correction* $-1/12$ *at* 109*, of three-adic valuation* -1*; all* 729 *central lift pairs modulo* 27 *fail the tame relation;*

2. *the E.31 origin retains* $\kappa_3(\zeta)$*, its local Frobenius and inertia values are* 2 *and* 1*, and all nine central lift pairs fail; the entire coset* $\alpha + 3\mathcal{K}_0$ *has the same excluded image;*

3. *all the off-three residues vanish for every section unramified outside three;*

4. *the E.35 norm polynomial is* $X^3 + AY^3 + A^2Z^3 - 3AXYZ$*, its example* $A = 2, b = 3$ *has witness* $(1, 1, 0)$*, and its finite conjugation calculation has span-rank histogram* $(1, 32, 48)$ *and* 153 *orbits over all* 81 *abelian assignments and* 729 *generator assignments.*

The infinite family A.6 is certified by correction valuations at least j*, whereas A.4 is certified by primitive corrections at every selected prime. These two conclusions use their complete tail proofs, with no inference from a finite sample.*

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379).

Proof. For exact finite-support data, the program calculates $-D_q^{-1} r_q$ by rational linear algebra and tests the three-adic valuation of every coordinate. All unlisted corrections are zero, so this is exactly (A.8.4). The two family constructors apply the adjugate bound of A.6

and the minimal-denominator proof of A.4, respectively. Their output records these bounds explicitly.

The comparison for E.28 is the actual covering homomorphism $N \to \widehat{F}_2^{(3)}$ followed by its class-two quotient. The two coordinate inertia exponents multiply by three, so the bracket multiplier is nine. At 109, choose the residue generator 6^4 of the three-primary roots of unity; the Kummer value of six is one. The resulting differential is $109 - 1 = 108$ and the residue is nine, giving $-9/108 = -1/12$. The program also enumerates both central coordinates modulo 27 and checks the same nonzero defect for all 27^2 possibilities.

For E.31 use the actual function $h = (x+y)/(x+\zeta y)$ on the cubic, with the same tangent at O. Writing $s = z/x$, its leading constant is $-1/(3\lambda) = \zeta\lambda^{-3}$. At $\mathfrak{q} = (3+\zeta)$ over seven, $\zeta = 4$, $\lambda = 4$ and $\bar{\pi} = 2 - \zeta = 5$. At the specified torus point the total h unit reduces to three, whose square is two, the square of $\zeta = 4$. Thus the Kummer Frobenius exponent is two and the other coordinate has inertia exponent one. The commutator is nonzero in the order-27 Heisenberg quotient. The morphism on abelian cohomology to these two $\mathbb{F}_3$ coordinates is additive, with the fixed affine origin just computed; adding $3\mathcal{K}_0$ leaves it unchanged. This proves the whole coset assertion, rather than a list of sampled coset elements.

For an unramified section $t_{q,d} = 0$ for every $q \neq 3$; every term of (A.8.3) is zero. The program verifies the corresponding polynomial identity. The existence of lifts to the full N uses the free pro-three argument, not this finite computation.

Finally the inner action of E.31 is removed by the explicit arithmetic-group isomorphism $(n, g) \mapsto (nq_g, g)$, $q_g = (\kappa_3(\zeta)(g), 0, 0)$. It sends the abelian pair (a, b) to $(A, b) = (\zeta a, b)$. The central extension then has cup obstruction $\kappa_3(A) \cup \kappa_3(b)$. The crossed-product comparison [Mil20b, IV.3.14] represents this class by the cyclic algebra $L_A \oplus L_A U \oplus L_A U^2$, with $U\ell = \sigma(\ell)U$ and $U^3 = b$. It splits exactly when $b = N_{L_A/K_0}(w)$: a norm witness gives the representation $U = w\sigma$ on the three-dimensional K_0-space L_A; a splitting conversely gives that form for U on a one-dimensional L_A-module. Its norm multiplication matrix on $1, t, t^2$ is

$$\begin{pmatrix} X & AZ & AY \\ Y & X & AZ \\ Z & Y & X \end{pmatrix}.$$

The determinant is the displayed polynomial. The finite group law $\frac{1}{2}(uv' - vu')$ gives the central change $rv - su$ under conjugation. Thus the complete nonempty fibre is $K_0^\times / (K_0^{\times 3} \langle A, b \rangle)$. The program checks the determinant identity, the norm witnesses, and every finite conjugation orbit. This confirms the exact quotient calculation used in E.35; it does not assert that a lift in this finite quotient already lifts to $N/\gamma_3 N$. □

Statement A.12 (Refuted). A universal finite-precision decision program. *There is no program which, for every completed Kummer input in A.3 supplied through finite-coefficient queries, terminates and decides whether its continuous intrinsic obstruction vanishes. The same failure holds for the full inputs of A.8. Finite-support inputs and inputs furnished with proved tail certificates have the exact evaluator in A.11. There is also a concrete family of inputs described by finite programs, each returning the actual R4/E36 finite-coefficient cocycles and each with the uniform known tail bound $v_3(a_j) \geq j$, on which the vanishing decision is undecidable. Thus replacing an oracle input by a finite program for its coefficients does not provide a total evaluator on this explicitly specified class. The task's phrase "given (α, a)" does not itself specify an input encoding; these two precise input classes state the scope of the refutation, while A.11 states the implemented domain and proves all four requested reproductions.*

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379).

Proof. Run such a program on the zero input. Its terminating computation uses finitely many coefficient levels, with maximum 3^n. Discard the first n terms of the nonzero obstruction constructed in A.4. The resulting input is zero modulo 3^n, since its remaining coefficients have valuations at least their indices. Every query in the alleged computation therefore receives exactly the same answer. Nevertheless its corrections remain primitive at infinitely many primes, so A.3 proves that its obstruction is nonzero. The program gives the same answer on two inputs requiring different answers, a contradiction. The counterexample is a tail of the actual intrinsic family, rather than an arbitrary formal sequence outside the section problem. This establishes the stated precise computational limit and explains the role of the tail certificates in A.11.

For the finite-program assertion, first make the family A.4 effective. At level 3^r, the representation $E_w(-1)/3^r$ is determined by the 3^r-division points of the explicit generalized Jacobian and the 3^r-division torsor of the framed point $P - O$. The latter point and its torus multiple are the actual framed point of A.9 and (E24.6). Multiplication by 3^r is a finite algebraic morphism on this semiabelian variety in characteristic zero. Its fiber equations, the elliptic division polynomials, and the torus root equations give a finite collection of algebraic numbers and their exact Galois action. These are effective finite algebraic operations: roots are represented by polynomials and isolating data, and their equalities, embeddings, and permutations are decided by exact algebraic arithmetic. The exterior-square map and the cochain $b_{g,P} \wedge e_P$ then give the finite representation used in A.4, in the fixed existing coordinates.

We can consequently enumerate primes and test the finite conditions $D_q \equiv 0 \pmod{3^j}$ and $u_q \not\equiv 0 \pmod 3$ by reduction in that explicitly constructed finite Galois extension. Take q_j to be the first unused larger prime passing these tests. A.4 and Chebotarev prove termination of each search. Put $m_j = v_3(\det D_{q_j})$. The matrix D_{q_j} is effective from the integral Frobenius matrices, and is invertible over $\mathbb{Q}_3$ by the weight calculation. Its inverse has denominators dividing 3^{m_j}. The minimal exponent e_j such that $3^{e_j} D_{q_j}^{-1} u_{q_j}$ is integral is therefore determined by the finite representation modulo 3^{m_j+1}; it satisfies $j \le e_j \le m_j$. This constructs uniformly computable (q_j, e_j) and all finite cocycles of the actual A.4 input, rather than a formal sequence of unrelated residue classes.

Now fix an ordinary finite Turing-machine description T. Define its input coefficients by the bounded calculation

$$a_{T,q_j} = \begin{cases} 3^{e_j}, & T \text{ has not halted within } j \text{ steps}, \\ 0, & T \text{ has halted within } j \text{ steps}, \end{cases} \qquad a_{T,q} = 0 \quad (q \notin \{q_j\}). \tag{A.12.1}$$

Every coefficient query requires only a bounded simulation. To return the entire input modulo 3^n, the program constructs q_j, e_j for $j < n$, performs those finitely many bounded simulations, and returns the finite Kummer product with the surviving exponents 3^{e_j}. Every omitted coefficient is divisible by 3^n. Thus $j \ge n$ is one uniform convergence modulus, independent of whether T halts.

The input section is the actual equivariant line $\ell\kappa(a_T)$ of A.3. Its R4 coordinates are $(0, -\kappa(a_T); -\kappa(a_T), -\kappa(a_T), -\kappa(a_T))$ and its E36 parameter is $-\kappa(a_T)$, with the same fixed origins as A.8. The finite Kummer product and its algebraic roots supply the corresponding finite cocycle; the preceding division-point construction supplies every required origin cochain at the requested level. Hence these programs give legitimate inputs in the existing coordinate system and include their finite cochain data.

If T halts, (A.12.1) has finite support. All of its local corrections are integral by the definition of e_j, so its obstruction vanishes. If T does not halt, (A.12.1) is the A.4 phantom input and its obstruction is nonzero. A total vanishing evaluator for these finite program descriptions would therefore decide whether an arbitrary Turing machine halts. The halting

theorem proves the second asserted impossibility. This conclusion concerns the precise uniform family (A.12.1); it does not assign an unstated encoding to all possible symbolic descriptions of completed inputs. □

Statement A.13 (Proved). The full finite-S complex, with its integral differential. *With the basis and the Frobenius u in A.7 and A.10, put $M = \mathcal{K}$, $M^+ = M^{c=1}$ and $d = (G_\alpha - 1)|_{M^+}$. There are compatible explicit quasi-isomorphisms*

$$R\Gamma(\mathbb{Z}[1/3], M/3^n) \simeq [M^+/3^n M^+ \xrightarrow{d} M/3^n M]$$

in degrees zero and one, and

$$R\Gamma_{\mathrm{cts}}(G_S, M) \simeq [M^+ \xrightarrow{d} M]. \tag{A.13.1}$$

The first lattice has rank 21 *and the second rank* 44. *The twenty-one nonzero Smith factors of d are seven units, nine factors three, and five factors nine. In particular*

$$\begin{aligned} H^0(G_S, M/3^n) &\simeq (\mathbb{Z}/3)^9 \oplus (\mathbb{Z}/3^{\min(n,2)})^5, \\ H^1(G_S, M/3^n) &\simeq (\mathbb{Z}/3^n)^{23} \oplus (\mathbb{Z}/3)^9 \oplus (\mathbb{Z}/3^{\min(n,2)})^5. \end{aligned} \tag{A.13.2}$$

The full integral matrices are printed by `code/a_k_lattice.py`*; thus (A.13.1) specifies the complex and not only its Euler characteristic.*

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379).

Proof. Every finite-coefficient cocycle factors through Γ. On $G_{K_0,S}$ its graph in the semidirect product of the coefficient three-group and the action image is a homomorphism to a three-group, so it kills the kernel defining P_3. This gives the required factorization; it is compatible with c. Inflation is therefore an isomorphism in degrees zero and one. Both groups have vanishing higher three-primary cohomology for these coefficients, by A.1 and the free pro-three kernel of Γ with its order-two quotient. Thus inflation is a quasi-isomorphism.

Normalize a one-cocycle on Γ to vanish on c, using averaging over its group of order two. Its value on the free generator u is arbitrary in $M/3^n$, and uniquely determines its value on cuc and on the entire group. The remaining coboundaries come precisely from $M^{c=1}/3^n$ and have value $(G_\alpha - 1)m$ at u. This construction is a cochain map from the displayed two-term complex to continuous cochains: in degree zero it is inclusion of the c-invariants; in degree one it sends a vector to the unique normalized cocycle. It induces the stated cohomology isomorphisms, hence is the claimed quasi-isomorphism. Reduction and inverse limits commute with these maps. The invariant lattice commutes with reduction because two is a unit.

For the matrix calculation, the columns of $1 + G_c$ span M^+. The program selects the twenty-one columns independent modulo three; Nakayama's lemma makes them a $\mathbb{Z}_3$-basis. It applies $G_\alpha - 1$ and performs the same certified Smith elimination as in A.7. The resulting valuations are seven zeros, nine ones, and five twos. Kernel and cokernel modulo 3^n give (A.13.2). The matrix has trivial integral kernel, consistent with $H^0(G_S, M) = 0$, and its integral cokernel is (A.7.1). □

1.5 An exact two-prime decomposition at seven

Put $\mathfrak{q} = (2 - \zeta)$ and $\overline{\mathfrak{q}} = (3 + \zeta)$ in $\mathbb{Z}[\zeta]$. Their residue fields are $\mathbb{F}_7$, with $\zeta = 2$ at $\mathfrak{q}$ and $\zeta = 4$ at $\overline{\mathfrak{q}}$. Write

$$S_7 = \{\lambda, \mathfrak{q}, \overline{\mathfrak{q}}, \infty\}, \quad G = G_{K_0,S_7}, \quad P_7 = G(3), \quad D_{\mathfrak{p}} = G_{K_{0,\mathfrak{p}}}(3).$$

Here (3) denotes the maximal pro-three quotient. Choose an embedding of the decomposition group at $\mathfrak{q}$ into G, and choose the embedding at $\bar{\mathfrak{q}}$ by conjugating it with the fixed complex conjugation c. The star in the following statement is the free product in the category of pro-three groups.

Statement A.14 (Proved). The two local factors generate the entire global pro-three group. *The natural morphism*

$$D_{\mathfrak{q}} *_3 D_{\bar{\mathfrak{q}}} \longrightarrow P_7 \tag{A.14.1}$$

is an isomorphism. Each factor has the explicit pro-three presentation

$$D_{\mathfrak{q}} = \langle \phi, \tau \mid \phi\tau\phi^{-1} = \tau^7 \rangle_3 \simeq \mathbb{Z}_3(1) \rtimes \mathbb{Z}_3, \tag{A.14.2}$$

where ϕ is arithmetic Frobenius and τ is tame inertia. The relative pro-three quotient of $G_{\mathbb{Q},\{3,7,\infty\}}$ over $\mathbb{Q}(\zeta)$ is therefore

$$\Gamma_7 \simeq (D_{\mathfrak{q}} *_3 D_{\bar{\mathfrak{q}}}) \rtimes \langle c \rangle, \qquad c^2 = 1, \tag{A.14.3}$$

with c exchanging the conjugately embedded factors. In particular the local morphisms in (A.14.1) are injective; this is part of the conclusion, not an assumption about realization of local extensions.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379); the one-prime higher-lifting problem in B.4.

Proof. We first prove exactly the comparison with a maximal pro-three quotient that will be used. Let H be any profinite group, $H \twoheadrightarrow H(3)$ its maximal pro-three quotient, and J its kernel. A nonzero invariant character $\eta : J \to \mathbb{F}_3$ would have a closed kernel normal in H. The resulting quotient $H/\ker\eta$ would be an extension of the pro-three group $H(3)$ by the group $\mathbb{F}_3$, hence a pro-three group. The defining maximality would make its map from H kill J, a contradiction. Thus

$$H^1(J, \mathbb{F}_3)^{H(3)} = 0, \qquad H^2(H(3), \mathbb{F}_3) \hookrightarrow H^2(H, \mathbb{F}_3). \tag{A.14.4}$$

The second assertion is the inflation map in the five-term Hochschild–Serre sequence. No assertion about all higher cohomology of this quotient is needed.

Let $U = \operatorname{Spec}\mathbb{Z}[\zeta, 1/(3 \cdot 7)]$. The ring is a localization of a principal ideal domain, so $\operatorname{Pic}(U) = 0$. The Kummer sequence, the fixed isomorphism $\mathbb{F}_3 \simeq \mu_3$ sending 1 to ζ, and Milne II.2.9 [Mil06] identify

$$H^1(G, \mathbb{F}_3) = \langle \zeta, \lambda, \pi, \bar{\pi} \rangle_{\mathbb{F}_3}, \qquad \pi = 2 - \zeta, \quad \bar{\pi} = 3 + \zeta, \tag{A.14.5}$$

and identify $H^2(G, \mathbb{F}_3)$ with $\operatorname{Br}(U)[3]$. Milne II.2.1 and Remark II.2.2(a) give its explicit invariant description

$$\operatorname{Br}(U)[3] = \{(x_\lambda, x_{\mathfrak{q}}, x_{\bar{\mathfrak{q}}}) \in (\tfrac{1}{3}\mathbb{Z}/\mathbb{Z})^3 : x_\lambda + x_{\mathfrak{q}} + x_{\bar{\mathfrak{q}}} = 0\}. \tag{A.14.6}$$

Consequently restriction from the full group G to the two local groups at seven is an isomorphism in degree two with these coefficients. Its inverse on invariants is $(x, y) \mapsto (-x - y, x, y)$. In particular it is injective. Combined with (A.14.4) and the natural local inflation maps, this proves the injection

$$H^2(P_7, \mathbb{F}_3) \hookrightarrow H^2(D_{\mathfrak{q}}, \mathbb{F}_3) \oplus H^2(D_{\bar{\mathfrak{q}}}, \mathbb{F}_3). \tag{A.14.7}$$

Indeed a class restricting to zero in both pro-three local groups inflates to a full global class with zero restrictions, and is therefore zero by (A.14.6) and (A.14.4).

For degree one, the two local Kummer groups each have dimension two: one uniformizer class and one residue-unit class. The unramified Kummer values of ζ and λ are

$$\begin{array}{c|cc} & \kappa_\zeta & \kappa_\lambda \\ \hline \phi_{\mathfrak{q}} & 2 & 0 \\ \phi_{\bar{\mathfrak{q}}} & 2 & 2 \end{array} \quad \text{in } \mathbb{F}_3. \tag{A.14.8}$$

These values follow by taking the square of the residue and expressing it as a power of the residue of ζ. For example $\lambda = -1$ at $\mathfrak{q}$, while $\lambda = 4 = \zeta$ at $\bar{\mathfrak{q}}$. The determinant of (A.14.8) is $4 = 1$ modulo three. The valuations of π and $\bar{\pi}$ at the two places form the identity matrix. Thus the restriction

$$H^1(P_7, \mathbb{F}_3) \longrightarrow H^1(D_{\mathfrak{q}}, \mathbb{F}_3) \oplus H^1(D_{\bar{\mathfrak{q}}}, \mathbb{F}_3) \tag{A.14.9}$$

is an isomorphism. With Frobenius lifts fixing the respective uniformizer roots, the complete matrix, in row order $(\phi_{\mathfrak{q}}, \tau_{\mathfrak{q}}, \phi_{\bar{\mathfrak{q}}}, \tau_{\bar{\mathfrak{q}}})$ and column order $(\zeta, \lambda, \pi, \bar{\pi})$, is

$$\begin{pmatrix} 2 & 0 & 0 & 2 \\ 0 & 0 & 1 & 0 \\ 2 & 2 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \tag{A.14.10}$$

Here both tame generators are oriented using the same global ζ. Hence $c\tau_{\mathfrak{q}}c^{-1} = \tau_{\bar{\mathfrak{q}}}^{-1}$. Using the conjugately transported second generator instead negates the last row; it does not change invertibility. This specifies the orientation in the factor exchange in (A.14.3).

For completeness we give the group-theoretic detection argument. Let $B = D_{\mathfrak{q}} *_3 D_{\bar{\mathfrak{q}}}$ and $f : B \to P_7$ the natural morphism. Its image is compact and therefore closed. A proper closed subgroup of a pro-three group is contained in an open subgroup of index three after passing to a finite quotient. Such a subgroup would yield a nonzero character of P_7 trivial on the image of f, contrary to (A.14.9). Thus f is surjective.

Restriction $H^2(B, \mathbb{F}_3) \to \bigoplus_{\mathfrak{p}|7} H^2(D_{\mathfrak{p}}, \mathbb{F}_3)$ is injective. To see this directly, a degree-two class is represented by a central extension of B by $\mathbb{F}_3$. Zero restriction provides homomorphic splittings on the two factors. The extension is a pro-three group, so the universal property of the free pro-three product combines these into a splitting on B. Equation (A.14.7) in particular shows that $f^* : H^2(P_7, \mathbb{F}_3) \to H^2(B, \mathbb{F}_3)$ is injective. Put $R = \ker f$. Its five-term sequence and (A.14.9) now give $H^1(R, \mathbb{F}_3)^{P_7} = 0$.

A nontrivial closed normal subgroup R of a pro-three group B always has a nonzero invariant character to $\mathbb{F}_3$. For a proof, choose a finite three-group quotient B_0 in which its image R_0 is nontrivial. The nonzero vector space $R_0/\Phi(R_0)$ is a finite module for the three-group B_0. Its coinvariants are nonzero: the augmentation ideal of $\mathbb{F}_3[B_0]$ is nilpotent, so equality of that ideal times the module with the whole module would force the module to be zero. A nonzero linear functional on the coinvariants gives an invariant character of R, a contradiction. Thus $R = 1$ and (A.14.1) is proved.

At residue characteristic seven every finite extension of three-power degree is tame. The maximal pro-three extension is obtained from the unramified three-power tower and compatible three-power roots of a uniformizer. Its tame inertia is $\mathbb{Z}_3(1)$, its unramified quotient is $\mathbb{Z}_3$, and arithmetic Frobenius acts on inertia by seventh powers. Choosing Frobenius to fix the uniformizer roots gives exactly (A.14.2). Finally the kernel of $G \to P_7$ is characteristic in G and stable under c. The chosen order-two element c therefore splits the relative quotient over $\mathbb{Q}$ and exchanges the conjugately chosen local factors, proving (A.14.3).

The elementary residue and matrix computations, including the orientation change, are reproduced in `code/a_two_prime.py` and its output file. □

Statement A.15 (Proved). The exact local equation for every section supported at three and seven. *Let Q be any of the geometric pro-three quotients of N under consideration, with the action defined by the unit tangential section at O. Write F for the action of $\phi_{\mathfrak{q}}$ on Q. Sections unramified outside $\{3,7,\infty\}$ have the exact parameterization*

$$\mathrm{Sec}_{\{3,7\}}(Q) \simeq \{(a,b) \in Q^2 : aF(b)a^{-1} = b^7\}/Q^c, \tag{A.15.1}$$

where

$$h \cdot (a,b) = (h^{-1}aF(h),\ h^{-1}bh) \quad (h \in Q^c). \tag{A.15.2}$$

The parameterization is natural under all the quotient maps. For a surjective equivariant morphism $R \twoheadrightarrow Q$ among these geometric pro-three groups, a class represented by (a,b) lifts exactly when there are lifts $A, B \in R$ satisfying

$$AF(B)A^{-1} = B^7. \tag{A.15.3}$$

The same statement holds for $R = N$. Moreover a fixed class-two section supported at these primes lifts to N exactly when it lifts to every lower-central quotient $N/\gamma_r N$. The equivalence includes compatible full lifts; no compatibility of separately chosen representatives is assumed.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379); the one-prime higher-lifting problem in B.4.

Proof. The action on N defined by the tangential basepoint is unramified away from three. Over K_0, its action on $N/\Phi(N) = V/3$ has three-group image, as proved in A.1. The kernel of $\mathrm{Aut}(N) \to \mathrm{Aut}(N/\Phi(N))$ is pro-three. One can see this on the finite lower-exponent-three quotients: successive kernels embed into homomorphism groups between finite $\mathbb{F}_3$-vector spaces, and are three-groups; their inverse limit is the stated kernel. Hence the action image over K_0 on N, and on each Q, is pro-three. The graph of any Q-valued cocycle on G_{K_0,S_7} is a homomorphism into the pro-three semidirect product of Q with this action image. It factors through P_7 by its defining maximality. Thus the sections in question factor through the exact group Γ_7 of A.14.

Every cocycle on $\langle c \rangle$ with values in a pro-three group is a coboundary. For finite three-groups this follows inductively along c-stable central elementary abelian subgroups: the induction leaves an abelian cocycle, and division by two solves it. For a pro-three group apply the finite assertion to c-stable finite quotients; the nonempty closed sets of conjugating elements have the finite-intersection property, so compactness supplies a common solution. We may therefore represent any section by a cocycle β with $\beta(c) = 1$. The remaining conjugacies are precisely by Q^c.

The restriction of a normalized cocycle to the first local factor is determined by $a = \beta(\phi_{\mathfrak{q}})$ and $b = \beta(\tau_{\mathfrak{q}})$. Tame inertia acts trivially on Q, because the reference action is unramified at seven. The single relation in (A.14.2) therefore becomes $aF(b)a^{-1} = b^7$. Conversely any pair satisfying this relation gives a continuous splitting over the local pro-three group: send the two generators to (a,ϕ) and (b,τ) in $Q \rtimes D_{\mathfrak{q}}$ and use its pro-three presentation. On the conjugate factor define

$$\beta(cdc^{-1}) = c(\beta(d)) \quad (d \in D_{\mathfrak{q}}).$$

The free-product property extends these local splittings uniquely over P_7, and the displayed identity extends them with $\beta(c) = 1$ over Γ_7. Conjugation of the two generator values by $h \in Q^c$ gives exactly (A.15.2). This proves (A.15.1) and its naturality.

The fixed-point map $R^c \to Q^c$ is surjective. Indeed, for a lift r of a fixed element of Q, the discrepancy $r^{-1}c(r)$ is a cocycle in the pro-three kernel of $R \to Q$. The preceding coprime argument removes it by an element of that kernel. Thus changes of normalized representatives

can be lifted, and (A.15.3) gives the stated criterion for the conjugacy class, not only for an individually chosen pair.

Finally fix the pair (a, b) in the class-two quotient. In the compact space of its lifts to N^2, the condition

$$AF(B)A^{-1}B^{-7} \in \gamma_r N$$

defines a closed subset. Lifting at level r makes this set nonempty, by lifting the two elements from $N/\gamma_r N$ to N; the representative issue has just been resolved. These sets are nested as r increases. Compactness provides a pair belonging to all of them. The intersection $\bigcap_r \gamma_r N$ is trivial: any nontrivial element survives in a finite three-group quotient, and every such quotient is nilpotent. The resulting pair satisfies the full equation in N, proving the last assertion. □

Equations (A.15.1)–(A.15.3) remove the additional global embedding problem at the specified prime seven. They do not evaluate the full local equation for every ramified mixed inertia vector. In particular A.14–A.15 alone do not settle the all-nilpotent assertion in B.4 or the general ramified metabelian classification in A.6.

Statement A.16 (Proved). A positive-density set with the same exact one-prime reduction. *Define the following explicit set of rational primes:*

$$\Sigma = \{q : q \equiv 4 \text{ or } 7 \pmod 9, \quad 3^{(q-1)/3} \not\equiv 1 \pmod q\}. \tag{A.16.1}$$

It has natural density $2/9$. *For every* $q \in \Sigma$, *write* $(q) = \mathfrak{q}\overline{\mathfrak{q}}$ *in* $\mathbb{Z}[\zeta]$. *The natural local morphisms give an isomorphism*

$$G_{K_0,\{\lambda,\mathfrak{q},\overline{\mathfrak{q}},\infty\}}(3) \simeq D_{\mathfrak{q}} *_3 D_{\overline{\mathfrak{q}}}, \quad D_{\mathfrak{q}} = \langle \phi, \tau \mid \phi\tau\phi^{-1} = \tau^q \rangle_3. \tag{A.16.2}$$

Its relative quotient over $\mathbb{Q}$ *is the semidirect product by complex conjugation exchanging the factors. The exact section parameterization and lifting assertions of A.15 hold with* 7 *replaced throughout by* q. *In particular the defining equation is* $aF_q(b)a^{-1} = b^q$, *with the same action of* Q^c *on pairs. The primes in* Σ *up to* 1000 *are*

$$\begin{gathered} 7, 13, 31, 43, 79, 97, 139, 157, 211, 223, 229, 241, \\ 277, 283, 313, 331, 337, 349, 373, 409, 421, 457, 463, \\ 571, 601, 607, 673, 691, 709, 733, 751, 769, 823, 859, 877, 907. \end{gathered} \tag{A.16.3}$$

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379); the one-prime higher-lifting problem in B.4.

Proof. Let $q \equiv 1 \pmod 3$ be a rational prime different from three, choose one residue root of ζ at $\mathfrak{q}$, and write

$$a = \kappa_\zeta(\phi_{\mathfrak{q}}) = \frac{q-1}{3} \bmod 3, \qquad b = \kappa_\lambda(\phi_{\mathfrak{q}}) \in \mathbb{F}_3.$$

Conjugation sends ζ to ζ^{-1} and λ to $-\zeta^2\lambda$. Its action on the Kummer coefficients μ_3 is multiplication by minus one. Thus the conjugately chosen Frobenius at $\overline{\mathfrak{q}}$ has Kummer vector $(a, a-b)$, with both characters measured using the same global ζ. In particular the determinant of the two unramified-unit rows is

$$\det \begin{pmatrix} a & b \\ a & a-b \end{pmatrix} = a(a+b) \quad \text{in } \mathbb{F}_3. \tag{A.16.4}$$

The identity $\lambda^2 = -3\zeta$ gives $\kappa_3 = 2b - a = -a - b$. Hence (A.16.4) is nonzero precisely when $a \neq 0$ and the residue of 3 is not a cube. These are exactly the two elementary conditions in (A.16.1).

Since $\mathbb{Z}[\zeta]$ is principal, choose a generator π of $\mathfrak{q}$ and its conjugate for $\bar{\mathfrak{q}}$. The four S-unit Kummer classes are $\zeta, \lambda, \pi, \bar{\pi}$. Their inertia submatrix is the identity, while (A.16.4) is their unramified-unit submatrix. For $q \in \Sigma$ this proves the degree-one isomorphism used in A.14. The Brauer invariant proof is unchanged, since exactly three finite places are inverted. All group-theoretic maps and kernel arguments in A.14 consequently apply to these actual local morphisms and prove (A.16.2). The local tame presentation has exponent q; $q \equiv 1 \pmod 3$ ensures that its Frobenius action on three-inertia factors through $\mathbb{Z}_3$. The construction of normalized cocycles and the compactness argument in A.15 now give all the stated section assertions.

For the density assertion put

$$L = K_0(\zeta_9, \sqrt[3]{\lambda}).$$

The Kummer classes of ζ and λ are independent: valuation at λ first forces the latter exponent to be zero modulo three, and ζ is not a cube in K_0, as its unit group is μ_6. Hence $[L : K_0] = 9$ and its Galois group is $\mathbb{F}_3^2$, with the two displayed Kummer coordinates. The relation $\bar{\lambda} = -\zeta^2\lambda$ makes L stable under complex conjugation: a conjugate cube root is $-\zeta_9^2\sqrt[3]{\lambda}$. Thus $L/\mathbb{Q}$ is Galois of degree eighteen. The chosen complex conjugation acts on the normal subgroup by

$$(a, b) \longmapsto (a, a - b).$$

The admissible vectors are exactly $(1, 0), (1, 1), (2, 0), (2, 2)$; this four-element subset is stable under conjugation in the full group. A rational prime outside three belongs to Σ exactly when its Frobenius class belongs to this subset. Chebotarev [Mil20b, V.3.23] therefore gives natural density $4/18 = 2/9$. The exhaustive elementary check through 1000, including both residue roots, the conjugation identity, the κ_3 identity and the determinant test, is in `code/a_two_prime.py` and its output. □

1.6 A strict restriction on ramification in the higher quotients

Statement A.17 (Proved). Tame abelian inertia in a metabelian section lies on one cusp line. *Let k be a finite extension of $\mathbb{Q}_q$ with $q \neq 3$, and let*

$$s : G_k \longrightarrow (N/N'') \rtimes G_k$$

be a continuous section. For a generator of tame three-inertia, write t for its geometric value and w for its image in $V = N^{\mathrm{ab}}$. Then

$$w = 0 \quad \text{or} \quad w \in \mathbb{Z}_3 e_d \setminus \{0\} \text{ for a unique cusp } d \in D(k). \tag{A.17.1}$$

In particular the assertion T1 of Lead 2, refuted for class two by A.5, holds for the tame abelian inertia of every metabelian section through the actual quotient $N/N'' \twoheadrightarrow N/\gamma_3 N$.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379).

Proof. Put $Q = N/N''$ and let Q_k denote the size of the residue field of k. The good marked model and its integral tangent make the reference action of inertia on N trivial. Consequently the restriction of the cocycle of s to inertia is a homomorphism; wild inertia and the tame factors prime to three have trivial image in Q. For an arithmetic Frobenius lift $F = s(\phi_k)$ the exact relation in the arithmetic quotient is

$$FtF^{-1} = t^{Q_k}. \tag{A.17.2}$$

It follows on abelianization that w is in W: its image in T_3E is a Q_k-eigenvector, and that Tate module has Frobenius eigenvalues of complex absolute value $\sqrt{Q_k}$ and is torsion-free.

For $n \geq 1$ define the actual subgroups of the full geometric group

$$M_n = \ker(N \to V/3^n) = N'N^{3^n}, \qquad H_n = \{g \in N : g \bmod M_n \in \langle w \bmod 3^n \rangle\}. \quad \text{(A.17.3)}$$

Here products and powers denote closed subgroups. The group H_n is open and normal in N, contains N', and is stable under the action induced by the local section. Indeed inertia acts trivially on V, while Frobenius sends the generating vector w to $Q_k w$, and Q_k is a three-adic unit. Crucially, the inclusion $N' \subset H_n$ gives

$$N'' = [N', N'] \subset [H_n, H_n], \qquad H_n^{\mathrm{ab}} \xrightarrow{\sim} (H_n/N'')^{\mathrm{ab}}. \quad \text{(A.17.4)}$$

The displayed isomorphism is induced by the quotient map; its kernel is explicitly $N''[H_n, H_n]/[H_n, H_n] = 0$.

The subgroup $(H_n/N'')s(G_k)$ of $Q \rtimes G_k$ is open. Pull it back through the actual morphism $N \rtimes G_k \to Q \rtimes G_k$. The resulting arithmetic subgroup defines a finite etale cover $Y_n \to U_k$, with geometric group H_n. This cover has the smooth proper tame compactification needed for the weight argument in B.7. We spell out why that argument applies to the present metabelian section, without assuming a full section lift. The cocycle values on arithmetic inertia are powers of t and therefore lie in H_n/N''. Thus the pulled-back subgroup contains the original arithmetic inertia. Original inertia centralizes N, so its action on every coset in N/H_n is trivial. Prime-to-q specialization therefore descends the cover to the good marked model. At each boundary point its compactification is the tame normalization $z^e = x$ after an unramified extension, with e prime to q; it is smooth there and etale away from the boundary. This is the specialization comparison used in B.7, with the same hypotheses.

Choose any lift $\widetilde{t} \in H_n$ of t, and any lift $\widetilde{F}$ of the Frobenius value in this arithmetic cover subgroup. Relation (A.17.2) says

$$\widetilde{F}\widetilde{t}\widetilde{F}^{-1}\widetilde{t}^{-Q_k} \in N''.$$

By (A.17.4), the class of $\widetilde{t}$ in H_n^{ab} is therefore an exact Q_k-eigenvector. Let X_n denote the smooth proper compactification of Y_n. Its boundary sequence is the integral exact sequence

$$0 \longrightarrow \mathbb{Z}_3(1)[X_n \setminus Y_n]/\mathbb{Z}_3(1)\delta_n \longrightarrow H_n^{\mathrm{ab}} \longrightarrow T_3 J_{X_n} \longrightarrow 0. \quad \text{(A.17.5)}$$

Frobenius on the last term has complex absolute values $\sqrt{Q_k}$, by the smooth proper finite-field weight theorem used in B.7. It has no Q_k-eigenvector even integrally, because the lattice is torsion-free. Thus the class of $\widetilde{t}$ is in the actual boundary submodule in (A.17.5).

The finite cyclic three-group H_n/M_n is generated by the image of $\widetilde{t}$. The boundary submodule therefore maps onto this group. At least one boundary inertia subgroup maps onto it: otherwise all its images would be contained in the unique maximal subgroup of this cyclic group. Such a boundary generator is conjugate in N to c_d^e for some cusp $d \in D$ and some positive integer e. Its image in $V/3^n$ is ee_d. The assertion that this element generates H_n/M_n means that w mod 3^n is a scalar multiple of e_d mod 3^n. If the cyclic group is trivial, the same conclusion holds with scalar zero.

There are only nine cusps. One cusp occurs for an unbounded subsequence of n, and its line $\mathbb{Z}_3 e_d$ is closed in V. Equivalently, e_d is primitive and the quotient of V by that line is a free $\mathbb{Z}_3$-module, so congruence membership for unbounded n gives exact membership. This proves $w \in \mathbb{Z}_3 e_d$. Distinct cusp lines have zero intersection: in $\mathbb{Z}_3[D]/\mathbb{Z}_3\delta$ an equality $ae_d = be_{d'}$ with $d \neq d'$ has zero coefficients at the other seven cusps, forcing the diagonal coefficient and then a, b to be zero. Thus the cusp is unique when $w \neq 0$. Frobenius sends e_d to $Q_k e_{\phi_k d}$. Together with $F(w) = Q_k w$, this forces $\phi_k d = d$. The cusp scheme is unramified at q, so this is precisely $d \in D(k)$, proving (A.17.1). $\square$

Statement A.18 (Proved). Class-two mixed examples have no metabelian or marked-depth lift. *Let I_{mark} be the closed normal subgroup generated by the inertia at Q_c, so that the marked depth quotient in the task is $N/\gamma_3(I_{\mathrm{mark}})$. The class-two sections of A.5 and A.6 have no lift to either*

$$N/N'' \quad or \quad N/\gamma_3(I_{\mathrm{mark}}). \tag{A.18.1}$$

This holds for every class-two lift of their specified N^{cc} sections, and continues to hold after every finite number-field extension. Thus their section germs also have no lift to these two quotients.

More generally, a local section of the marked depth quotient over a finite extension $k/\mathbb{Q}_q$, $q \neq 3$, has tame abelian inertia w whose image in $V/\mathbb{Z}_3 e_{Q_c}$ is zero or lies on a single cusp line belonging to $D \setminus \{Q_c\}$ in that quotient. The images of both section maps in (A.18.1) to class two are therefore proper; the entire unramified family of A.20 still belongs to both images.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379).

Proof. The two comparison maps to class two are genuine quotient maps. Indeed $N'' \subset \gamma_4 N \subset \gamma_3 N$, and $\gamma_3(I_{\mathrm{mark}}) \subset \gamma_3 N$ because $I_{\mathrm{mark}} \subset N$. At each prime used in A.5 or A.6, the specified abelian tame inertia is a nonzero scalar multiple of

$$w_0 = e_O + e_P.$$

It is on no cusp line. In the basis of W given by the eight cusps other than O, its coefficient at P is zero and its coefficients at the other seven cusps are all minus one. This vector is neither a multiple of a basis vector nor of e_O, whose eight coefficients are all minus one. Multiplication by a nonzero three-adic scalar does not change this conclusion. A.17 consequently rules out every metabelian lift.

For the marked quotient use the explicit filling morphisms

$$N/\gamma_3(I_{\mathrm{mark}}) \twoheadrightarrow N/I_{\mathrm{mark}} \xrightarrow{\sim} \pi_1(E_{\overline{\mathbb{Q}}} setminus(D \setminus \{Q_c\}))^{(3)}. \tag{A.18.2}$$

The second map is induced by filling the puncture Q_c. It is an isomorphism because a finite three-group etale cover of U extends etale across that puncture exactly when its inertia there is trivial. Normal closure kills precisely that inertia condition, and the finite covers on the two sides are thereby identified. The same map on abelianizations is

$$V \longrightarrow V/\mathbb{Z}_3 e_{Q_c},$$

as follows from the boundary exact sequence, with the single summand at Q_c removed. These morphisms preserve the arithmetic action and the tangent at O.

The resulting open elliptic curve has eight cusps and good marked reduction at $q \neq 3$. The proof of B.7 applies verbatim to its full geometric pro-three group: it uses only the finite-cover boundary sequence, smooth proper weights, and compactness of the lifted cusp set. Hence every nonzero tame inertia of the projected full section lies in one cusp inertia subgroup. Its abelian image is zero or on that cusp line, which proves the more general assertion of the statement.

In the filled curve the images of the seven cusps other than O, Q_c form a basis of its boundary lattice. The image of w_0 has coefficient zero at P and minus one at the other six basis cusps. It is not zero and is on no cusp line by the same coordinate comparison. Thus (A.18.2) rules out the marked-depth lifts of A.5 and A.6 as well.

Under a finite extension of local fields the tame inertia generator maps to a nonzero integer multiple of the old one, with that integer equal to the ramification index after compatible choices. The new vector is consequently a nonzero scalar multiple of the same mixed vector; changes of decomposition group merely conjugate and permute its two distinct cusps. The two line exclusions remain valid over every such extension, including those of degree divisible by three. A hypothetical lift over a finite number field would give a forbidden local lift at a place above one of the specified ramified primes. This proves the finite-extension and germ assertions. Finally the full unramified lifts of A.20 project through both quotient maps, while A.5 supplies a class outside both images. The images are therefore nonempty and proper. The integral line exclusions before and after filling are independently certified by unit two-by-two minors in `code/a_higher_checks.py` and its output. □

Statement A.19 (Proved). The complete local metabelian image at every good prime. *Let $k/\mathbb{Q}_q$ be finite with $q \neq 3$, put $Q = N/N''$, and retain the unit tangential origin at O. Every continuous section of $Q \rtimes G_k$ is either unramified or contained in the decomposition group of a unique lifted k-rational cusp. It is unramified exactly when its abelianization is unramified.*

More precisely, with F denoting the reference arithmetic Frobenius action and Q_k the residue-field cardinality, the local section set is the disjoint union

$$\mathrm{Sec}_k(Q) = Q/\{a \sim h^{-1}aF(h) : h \in Q\} \;\sqcup \coprod_{d \in D(k)} \{u \in k^{\times \wedge 3} : v_k(u) \neq 0\}. \tag{A.19.1}$$

The first term consists of unramified sections, with a their geometric Frobenius value. The term at d uses its actual unit tangent section $s_{d,1}$ and the map $g \mapsto c_d^{\kappa(u)(g)} s_{d,1}(g)$, followed by the quotient to Q. Its parameter group is

$$k^{\times \wedge 3} \simeq \mathbb{Z}_3 \oplus \mu_{3^{v_3(Q_k - 1)}}(k).$$

The morphism $\mathrm{Sec}_k(N) \to \mathrm{Sec}_k(Q)$ is surjective and is bijective on the ramified parts. The ramified image of $\mathrm{Sec}_k(Q) \to \mathrm{Sec}_k(N/\gamma_3 N)$ is exactly the full cuspidal packet of B.8, with the same parameters, including the central inertia and Frobenius coordinates. In particular no extra local metabelian section remains over a class-two section with a noncuspidal central coordinate.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379).

Proof. We give the inverse-limit and integral arguments that strengthen the line condition of A.17 to an actual cusp subgroup. Put $M = Q' = N'/N''$. Let $t \in Q$ be the tame inertia value, $w \in V$ its abelianization, and $\widetilde{t} \in N$ a lift. Use the groups M_n, H_n of (A.17.3), and set

$$H_\infty = \bigcap_n H_n = \{g \in N : g^{\mathrm{ab}} \in \mathbb{Z}_3 w\}. \tag{A.19.2}$$

All these groups contain N'. The same good-reduction cover argument as in A.17 makes the image of $\widetilde{t}$ in every H_n^{ab} lie in its integral boundary submodule B_n.

First we justify the precise inverse limits. For any descending sequence of closed subgroups of a profinite group, their images in a fixed finite quotient eventually stabilize to the image of their intersection. This follows by compactness: a compatible nonempty sequence of fibers over the fixed finite quotient has nonempty intersection. The images of their closed commutator subgroups therefore stabilize to the commutator of that intersection's image. Applying this to the H_n and testing in every finite quotient of N proves

$$\bigcap_n H_n' = H_\infty', \qquad H_\infty^{\mathrm{ab}} \xrightarrow{\sim} \varprojlim_n H_n^{\mathrm{ab}}. \tag{A.19.3}$$

For surjectivity of the second map, representatives of a compatible family define nested compact cosets of H_n'; their intersection gives a representative in H_∞. Thus the image of $\widetilde{t}$ lies in $\varprojlim B_n \subset H_\infty^{\mathrm{ab}}$.

Let $C_n(e)$ be the geometric cusps of the cover Y_n above a base cusp e. Since H_n contains N', these finite sets and their inertia exponents are explicitly

$$C_n(e) = V/(\mathbb{Z}_3 w + \mathbb{Z}_3 e_e + 3^n V), \qquad j_n(e) = \min\{j \geq 1 : j e_e \in \mathbb{Z}_3 w + 3^n V\}. \tag{A.19.4}$$

The minimum is a positive power of three. A basis vector at a cusp in $C_n(e)$ represents a conjugate of $c_e^{j_n(e)}$. For the map from level $n+1$ to level n, the boundary-module transition is the pushforward on cusp sets multiplied by $j_{n+1}(e)/j_n(e)$. This follows directly from the map on tame inertia generators, so the transition is specified integrally, including ramification. There is an exact sequence of compact modules

$$0 \to \mathbb{Z}_3(1)\delta_n \to \bigoplus_{e \in D} \mathbb{Z}_3(1)[C_n(e)] \to B_n \to 0. \tag{A.19.5}$$

The diagonal transition multiplies by $[H_n : H_{n+1}]$: over each cusp, the number of its preimages times their ramification index equals the cover degree.

For every inverse sequence of compact abelian groups the first derived limit is zero. Indeed, in the standard difference equation $x_n - f_n(x_{n+1}) = y_n$, any finite set of equations can be solved backwards. Their solution sets are closed in the compact product of the groups and have the finite-intersection property, which solves all equations simultaneously. This applies to the diagonal groups in (A.19.5), without asserting a Mittag–Leffler property for their underlying discrete groups.

Suppose first that $w = 0$. Then $H_\infty = N'$, $j_n(e) = 3^n$ for every cusp, and every cusp-module transition in (A.19.5) has a factor three. Its inverse limit is zero: each coefficient at a fixed level is divisible by arbitrarily high powers of three. The diagonal transitions have the same vanishing inverse limit, and the preceding derived-limit argument gives $\varprojlim B_n = 0$. By (A.19.3) the image of $\widetilde{t}$ in $N'/N'' = M$ is zero. Hence

$$w = 0 \quad \Longrightarrow \quad t = 1. \tag{A.19.6}$$

Now suppose $w \neq 0$. By A.17 there is a unique $d \in D(k)$ and $a \in \mathbb{Z}_3 \setminus \{0\}$ with $w = a e_d$. Write $a = 3^s u$, $u \in \mathbb{Z}_3^\times$. For $n > s$ the exponents in (A.19.4) are

$$j_n(d) = 3^s, \qquad j_n(e) = 3^n \quad (e \neq d). \tag{A.19.7}$$

For the second equality, distinct cusp vectors form a primitive independent pair in V: this is immediate in $\mathbb{Z}_3[D]/\mathbb{Z}_3\delta$ by choosing a third cusp to eliminate. Thus no smaller power of e_e can belong to the line generated by $a e_d$ modulo 3^n. The transition degree is 3^9 for $n > s$. Consequently the diagonal inverse limit is zero, all summands at $e \neq d$ have zero inverse limit, and the remaining summand gives the exact isomorphism

$$\varprojlim_n B_n \simeq \mathbb{Z}_3(1)[[A_d]], \qquad A_d = V/\mathbb{Z}_3 e_d. \tag{A.19.8}$$

Here the right side means the completed permutation module $\varprojlim_n \mathbb{Z}_3(1)[A_d/3^n A_d]$, with pushforward transition maps. Its basis measure at a coset represented by $g \in N$ maps to the class of $g c_d^{3^s} g^{-1}$ in H_∞^{ab}. The isomorphism follows from (A.19.5), including its diagonal term and the vanishing of that term's first derived limit; no closure or rationalization replaces the integral boundary module.

The actual Frobenius action on this module is Q_k times an affine permutation of A_d. To specify its origin, let $v_F \in V$ be the geometric Frobenius value of the given section and use

the cusp-path cocycle of A.8, so that reference Frobenius sends c_d to $b_{F,d}^{-1}c_d^{Q_k}b_{F,d}$. The affine map on the boundary cosets is

$$\Psi(x) = F_V x + \overline{v_F - b_{F,d}} \quad (x \in A_d). \tag{A.19.9}$$

The bar means the quotient by $\mathbb{Z}_3 e_d$. This proves the sign and the tangential origin of the translation: the section contributes left translation by v_F, and the reference path contributes $-b_{F,d}$. Changing the based path conjugates this affine description by a translation and leaves the underlying boundary action unchanged.

We use the following elementary integral-measure fact. For a finite free $\mathbb{Z}_3$-module A, let $\Psi(x) = Bx + v$ be an affine automorphism with $B^m - 1$ invertible over $\mathbb{Q}_3$ for every positive integer m. Then

$$\mathbb{Z}_3[[A]]^{\Psi} = \begin{cases} \mathbb{Z}_3\delta_{x_0}, & x_0 = (1-B)^{-1}v \in A, \\ 0, & x_0 \notin A. \end{cases} \tag{A.19.10}$$

Here is a proof with the necessary uniformity. The sole possible periodic point in $A \otimes \mathbb{Q}_3$ is x_0, since translation by x_0 turns Ψ into B and $B^m - 1$ is invertible. A positive power $\Theta = \Psi^e$ has pro-three closure in the affine automorphism group: take e to kill its finite action on $A/3A$. Thus every orbit of Θ on a finite quotient $A/3^nA$ has three-power length. Fix a residue cell in A not containing x_0, and let the corresponding orbit at level n have length 3^r. For each h, there is a level $m \geq n$ at which no point of the cell is fixed modulo 3^m by $\Theta^{3^{r+h}}$. Otherwise the nested nonempty compact solution sets would give an actual periodic point in that cell. Every orbit at level m meeting this cell therefore has length divisible by 3^{r+h+1}. An invariant coefficient is constant on such an orbit, and its pushforward to the chosen level-n cell is multiplied by the ratio of orbit lengths, divisible by 3^{h+1}. Hence that level-n coefficient is divisible by every power of three and is zero. All cells away from x_0 have zero coefficient. If $x_0 \notin A$ every coefficient is zero. If $x_0 \in A$, compatibility leaves exactly one constant mass on its residue cells, giving (A.19.10).

Apply (A.19.10) to (A.19.9). The eigenvalues of F_V on $A_d \otimes \mathbb{Q}_3$ have complex absolute values Q_k or $\sqrt{Q_k}$; the rational isogeny comparison is (A.1.4), followed by the quotient of its boundary part by e_d. Hence no positive power has eigenvalue one, as required. The class of $\widetilde{t}$ is a Q_k-eigenvector in H_∞^{ab}, so its measure in (A.19.8), after removing the Tate twist, is invariant under Ψ. Its total mass is u, because every boundary basis measure has abelian image $3^s e_d$ whereas $\widetilde{t}$ has image ae_d. Thus this mass is nonzero. Formula (A.19.10) proves that $x_0 \in A_d$ and that the measure is exactly $u\delta_{x_0}$. Choose $g \in N$ representing x_0. Then

$$\widetilde{t} \equiv g c_d^a g^{-1} \pmod{H'_\infty}. \tag{A.19.11}$$

This congruence gives actual conjugacy in the metabelian group. The image J of H_∞ in Q fits into $1 \to M \to J \to \mathbb{Z}_3 ae_d \to 1$ and splits by the subgroup generated by t, or by c_d^a. Since M is abelian, its commutator subgroup is

$$J' = (T-1)M, \tag{A.19.12}$$

where T is conjugation by any element with abelian image ae_d. The image is already closed, as the continuous image of the compact group M. Write $h = gc_d^a g^{-1}$ in Q. Equations (A.19.11)–(A.19.12) give $th^{-1} = (T-1)m$ for some $m \in M$, in additive notation for M. The right side equals $m^{-1}hmh^{-1}$. Hence $t = m^{-1}hm$ in Q: the tame inertia lies in one actual cusp inertia subgroup of the metabelian group.

We recall and justify the centralizer calculation needed to conclude for the whole section. For every nonzero integer b,

$$C_Q(c_d^b) = \overline{\langle c_d \rangle}. \tag{A.19.13}$$

This is Yamaguchi, Proposition 1.1.6 [Yam23], with the full class of finite three-groups and $m = 2$. Each c_d is a free generator in some basis of N: in the surface presentation one eliminates a different cusp, which is available because there are nine cusps. For a direct proof in this case, commutation in class two first makes the abelian image of any centralizer element a multiple of e_d, since $\bigwedge^2 V$ is torsion-free. Remove that power of c_d to obtain an element of M. The completed abelian-cover chain complex of a graph with ten free generators embeds

$$M \hookrightarrow \mathbb{Z}_3[[V]]^{10}, \qquad \partial(f_i) = \sum_i (T_i - 1) f_i.$$

One obtains this injection by taking the inverse limit of the degree-one chain sequences of its finite abelian covers; (A.19.3) with $H_\infty = N'$ identifies the limiting first homology with M. This is also Proposition 1.1.2 of the cited paper. The ring $\mathbb{Z}_3[[V]] \simeq \mathbb{Z}_3[[X_1, \ldots, X_{10}]]$ is an integral domain, and $T_d^b - 1$ is nonzero. Its multiplication is therefore injective on M, which proves (A.19.13).

After a geometric conjugation take $t = c_d^a$. Its closed generated subgroup equals that generated by $c_d^{3^s}$. Compare the actual Frobenius value with the Frobenius value of the unit tangent section at d. Both conjugate t to t^{Q_k}, so their geometric ratio centralizes a nonzero integral power of c_d. By (A.19.13) that ratio lies in its inertia group. Frobenius and inertia generate the local arithmetic image topologically; hence the whole section is contained in the cusp decomposition group. The cusp lift is unique: distinct base cusps have distinct abelian lines, and for two lifts of the same cusp any conjugator identifying a nonzero inertia element belongs to (A.19.13).

Sections in this decomposition group are exactly $H^1(G_k, \mathbb{Z}_3(1)) = k^{\times \wedge_3}$ with the specified tangential origin. A nonzero valuation is exactly nontrivial tame inertia. Their conjugacies in Q are exactly their conjugacies inside the cusp decomposition group, by the same uniqueness and centralizer argument. These packets therefore give the second, disjoint term in (A.19.1). For unramified sections the cocycle factors through $G_k/I_k = \widehat{\mathbb{Z}}$. A geometric value $a \in Q$ at arithmetic Frobenius determines it and always defines a continuous section: the closed procyclic subgroup generated by $(a, 1)$ in $Q \rtimes \widehat{\mathbb{Z}}$ maps isomorphically onto $\widehat{\mathbb{Z}}$. Indeed the map from $\widehat{\mathbb{Z}}$ sending 1 to this element composes with projection to the identity. Conjugation is precisely the relation in the first term of (A.19.1).

Finally choose any lift of an unramified Frobenius value to N and repeat this procyclic construction. Every cuspidal packet already comes from the actual full cusp decomposition group. Thus all local metabelian sections lift to N. B.7 and the injective packet parameterization give bijectivity on the ramified parts. Projecting the same actual packets to class two gives exactly B.8; its injective parameters also show that both the central inertia and the Frobenius coordinate are retained. This proves all claims. □

1.7 Complete global images and fibers for one auxiliary prime

Put $A = N/\gamma_3 N$ and, for a geometric quotient B mapping to A, put $J_B = \ker(B \to A)$. For $q \neq 3$ let $\mathscr{L}_q \subset \mathrm{Sec}_{\mathbb{Q}_q}(A)$ be the union of the unramified local section classes and the ramified cuspidal packets in B.8. Thus this subset uses the actual class-two inertia and Frobenius values and their unit-tangent origins, including all central coordinates.

Statement A.24 (Proved). The complete one-prime metabelian image and its fibers for $q \in \Sigma$. *For each $q \in \Sigma$ of A.16, put $S_q = \{3, q, \infty\}$. Then*

$$\mathrm{Sec}_{S_q}(N) \longrightarrow \mathrm{Sec}_{S_q}(N/N'') \quad \textit{is surjective.} \tag{A.24.1}$$

For either $B = N$ or $B = N/N''$, the exact image in class two is

$$\mathrm{im}\big(\mathrm{Sec}_{S_q}(B) \to \mathrm{Sec}_{S_q}(A)\big) = \mathrm{res}_q^{-1}(\mathscr{L}_q). \tag{A.24.2}$$

All fibers are as follows. A target outside the displayed subset has empty fiber. A target unramified at q has precisely the unramified fiber of (A.20.2); every lift in the present fiber is unramified outside three. For a target whose restriction at q is ramified and belongs to $\mathscr{L}_q$, choosing one normalized lift gives a bijection of its fiber with the set of right cosets

$$J_B/J_B^c. \tag{A.24.3}$$

Here J_B^c is the fixed subgroup of the actual complex conjugation, and it is not asserted to be normal. The same description for the ramified fibers of (A.24.1) is $N''/(N'')^c$.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379); Tasks A.6 and B.4 for this explicit infinite set of primes.

Proof. Use the exact relative free product of A.16 and normalize all cocycles to have value one on c. A normalized target section is determined by its pair (a, b) on the chosen local factor at q. A local full lift of its conjugacy class can be made a lift of this exact pair: lift a conjugator through the surjection of geometric groups and conjugate the local lift. A.15 then extends that pair, together with its c-conjugate pair and value one at c, to a global section over $G_{\mathbb{Q},S_q}$. A.19 supplies such a local full lift for every local metabelian section. This proves (A.24.1). B.7–B.8 and A.19 give exactly the local subset $\mathscr{L}_q$, so the same construction proves both directions of (A.24.2).

If the target is unramified at q, its abelian inertia is zero. The inertia of every full or metabelian lift is then trivial, by B.7 or (A.19.6), respectively. Such a lift factors through $G_{\mathbb{Q},\{3,\infty\}}$: the kernel of the map from $G_{\mathbb{Q},S_q}$ to this group is the closed normal subgroup generated by the inertia groups above q, and the reference action is also trivial on these groups. Thus its entire fiber is exactly the one already determined in A.20.

Now fix a ramified target in the image and fix a full or metabelian global lift $\widetilde{\beta}$ with $\widetilde{\beta}(c) = 1$ and exact projection β. Such an exact representative exists because the fixed-point map $B^c \to A^c$ is surjective, as proved in A.15. The ramified local packet injects into class two by B.8 and A.19. Consequently all local B-section lifts of the target pair are conjugate in B.

The target local section has trivial centralizer in A. Indeed a centralizer element has abelian image fixed by the twisted Frobenius, whose action on V has no eigenvalue one. It therefore lies in $\bigwedge^2 V$; the Frobenius eigenvalues there also have absolute value strictly greater than one, so that central element is zero. The local section centralizer in B is likewise trivial: any such element centralizes its nonzero cusp inertia and so lies in that cusp group, by B.4 or (A.19.13); Frobenius multiplies this group by q, with no nonzero fixed element. Hence every local pair lifting the exact target pair is uniquely of the form

$$\widetilde{\beta}|_{D_q}^{j} \quad (j \in J_B),$$

where the superscript means geometric conjugation by j, with cocycle convention $\beta^j(g) = j^{-1}\beta(g)g(j)$. The conjugator belongs to J_B, since its projection centralizes the target local section.

Extend each such local pair to a global normalized cocycle by A.15. Every element of the global fiber is obtained this way. A conjugation between two resulting global normalized cocycles belongs to B^c and its projection centralizes β; the latter centralizer is already trivial after restriction at q. Thus the conjugator lies in J_B^c. Composition of conjugations is $(\widetilde{\beta}^{\cdot j})^h = \widetilde{\beta}^{\cdot jh}$, so the identifications are exactly right multiplication by J_B^c. This proves (A.24.3). For the map to the metabelian quotient the same argument uses its trivial ramified local section centralizer and kernel N'', giving the last assertion. □

Statement A.25 (Proved). The relative arithmetic presentation at every inert prime $q \equiv 2, 5 \pmod 9$. *Put*

$$\Xi = \{q \text{ prime} : q \equiv 2 \text{ or } 5 \pmod 9\}. \tag{A.25.1}$$

This set has natural density $1/3$ and contains 2. For $q \in \Xi$ let $\mathfrak{q} = (q)$ be the unique prime of K_0 above q, and write $P_q = G_{K_0,\{\lambda,\mathfrak{q},\infty\}}(3)$. After conjugating the local embedding compatibly with the fixed global complex conjugation, there is an isomorphism

$$P_q \simeq D_{q^2} *_3 \mathbb{Z}_3^{\langle z \rangle}, \qquad D_{q^2} = \langle \phi, \tau \mid \phi\tau\phi^{-1} = \tau^{q^2} \rangle_3, \tag{A.25.2}$$

and its relative arithmetic quotient is

$$\Gamma_q \simeq (D_{q^2} *_3 \mathbb{Z}_3^{\langle z \rangle}) \rtimes \langle c \rangle, \qquad c\phi c^{-1} = \phi, \quad c\tau c^{-1} = \tau^{-1}, \quad czc^{-1} = z^{-1}. \tag{A.25.3}$$

Here $c^2 = 1$. Thus $\sigma = zc$ is another involution, with no further relation beyond (A.25.3). The source category in these presentations consists of profinite groups whose kernel over $\langle c \rangle$ is pro-three; the kernel free product is precisely the pro-three free product, as displayed.

For an inert prime $q \equiv 8 \pmod 9$, the two unit Frobenius values used in this construction are both zero modulo three. In particular $q = 17$ is the smallest prime not covered by $\Sigma \cup \Xi$; (A.25.2) is not asserted there. The union $\Sigma \cup \Xi$ has natural density $5/9$.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379).

Proof. The three S-unit Kummer characters over K_0 are $\kappa_\zeta, \kappa_\lambda, \kappa_q$. At the inert prime the residue field is $\mathbb{F}_{q^2}$. Write $a = (q^2 - 1)/3 \bmod 3$. The two unramified values are

$$(\kappa_\zeta(\phi), \kappa_\lambda(\phi)) = (a, 2a). \tag{A.25.4}$$

Indeed $\lambda^{q-1} = -\zeta^2$ in this residue field, because Frobenius sends ζ to ζ^2 and $\bar{\lambda} = -\zeta^2\lambda$. Consequently $\lambda^{(q^2-1)/3} = \zeta^{2(q+1)/3}$. The sign is one for odd q, since $(q+1)/3$ is even, and is one in characteristic two as well. Since $q - 1 = 1$ modulo three, this exponent is $2a$. The value a is nonzero exactly for $q \equiv 2, 5 \pmod 9$. The character κ_q gives the independent tame-inertia valuation.

Thus restriction in degree one from P_q to the local group D_{q^2} has rank two, with a one-dimensional kernel. In degree two the argument of A.14 applies with just two finite places in S: the Brauer invariants at λ and $\mathfrak{q}$ sum to zero, so restriction to $\mathfrak{q}$ is injective. Inflation from the maximal pro-three quotient is injective by (A.14.4). Choose one element complementing the local image in the global Frattini quotient. The resulting map $D_{q^2} *_3 \mathbb{Z}_3 \to P_q$ is an isomorphism by the five-term kernel proof of A.14. In particular the local morphism is injective before any assertion about an equivariant complement is made.

The relative local group over $\mathbb{Q}_q$ is $D_{q^2} \rtimes C_2$. It is explicitly obtained from the tower of unramified extensions with degrees $2 \cdot 3^n$ and compatible three-power roots of the uniformizer q. Its order-two unramified element acts on tame three-inertia by inversion and commutes with the pro-three unramified generator ϕ; the latter acts on inertia by q^2. Its morphism to the global relative quotient is injective, since it is injective on the pro-three kernel just proved and on the quotient C_2. Its order-two complement is conjugate to the fixed global c by an element of P_q, using the coprime complement argument of A.15. Conjugate the local embedding accordingly. This yields the first two action formulas in (A.25.3) with the actual chosen c.

It remains to produce the equivariant free generator. On the three global Kummer characters, conjugation is

$$c(\kappa_\zeta) = \kappa_\zeta, \quad c(\kappa_\lambda) = \kappa_\zeta - \kappa_\lambda, \quad c(\kappa_q) = -\kappa_q. \tag{A.25.5}$$

This uses conjugation on the fixed coefficient μ_3 as well as on the S-units. The Frattini quotient therefore has signs $(1,2)$, whereas the local subgroup has signs $(1,1)$ by its explicit ϕ,τ action. Over $\mathbb{F}_3$, division by two gives an invariant complement, and that one-dimensional complement has minus sign. Choose an element $x \in P_q$ lifting a nonzero vector in it and put

$$z = x\,c(x)^{-1}.$$

Then $c(z) = z^{-1}$ exactly, while its Frattini class is twice that chosen vector. It and the local Frattini image span the whole quotient. The map with this new generator in place of the initial complement is again an isomorphism by the same degree-one and degree-two argument. This proves both (A.25.2) and the full equivariant presentation (A.25.3). In terms of $\sigma = zc$, the last relation is exactly $\sigma^2 = 1$; substitution $z = \sigma c$ gives the reverse morphism of presentations, proving that no additional relation was omitted.

Chebotarev in $\mathbb{Q}(\zeta_9)/\mathbb{Q}$ assigns density $1/6$ to each invertible residue class modulo nine, so (A.25.1) has density $1/3$. It consists of inert primes, whereas Σ consists of split primes. Their densities add to $1/3+2/9 = 5/9$. Formula (A.25.4) gives zero for both units at $q \equiv 8 \pmod 9$; thus the local degree-one restriction at 17 has rank one, not two. This identifies the precise failure of the present free-product construction there. The exact residue-field computations, involution matrices, and all primes through 1000 are reproduced in `code/a_inert_prime.py` and its output. The list has 58 primes in Ξ in that range. □

Statement A.26 (Proved). The complete one-prime metabelian image and fibers for $q \in \Xi$. *For every $q \in \Xi$ and $S_q = \{3, q, \infty\}$, the conclusions (A.24.1)–(A.24.2) hold. Thus every metabelian section supported at these primes lifts to N, and for $B = N$ or N/N'' the exact image in class two is*

$$\operatorname{im}\bigl(\mathrm{Sec}_{S_q}(B) \to \mathrm{Sec}_{S_q}(A)\bigr) = \mathrm{res}_q^{-1}(\mathscr{L}_q). \tag{A.26.1}$$

For targets unramified at q, the fiber is exactly the unramified fiber of A.20. For a ramified target in this image, choose a normalized lift and let X_0 be its cocycle value at the involution $\sigma = zc$ of A.25. On J_B put

$$\Theta = \mathrm{Ad}(X_0) \circ \sigma.$$

This is an involution. The fiber is in bijection with either of the following explicitly identified sets:

$$\{j \in J_B : j\Theta(j) = 1\} \simeq J_B^{\Theta}\backslash J_B, \qquad h \longmapsto h^{-1}\Theta(h). \tag{A.26.2}$$

The quotient in the right member is a set of left cosets, without an assertion of normality. The same ramified-fiber formula for the surjection from full to metabelian sections uses $J_B = N''$ and the involution supplied by the chosen full lift.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379); Tasks A.6 and B.4, including the prime two.

Proof. Normalize the cocycle at c to one. Presentation (A.25.3), with z replaced by σc, gives the following exact data for a section with values in a geometric quotient B:

$$\begin{gathered} a \in B^c, \qquad c(b) = b^{-1}, \qquad aF_{q^2}(b)a^{-1} = b^{q^2}, \\ X \in B, \qquad X\sigma(X) = 1. \end{gathered} \tag{A.26.3}$$

The first row is its section on the relative local group at q, with values at ϕ,τ. The second is its value at the additional involution. The remaining conjugations are by B^c, acting on the first row as in (A.15.2) and on the second row by $X \mapsto h^{-1}X\sigma(h)$. The pro-three

universal property in (A.25.2) proves that these data define continuous sections, and the given presentation proves that they exhaust them.

For every equivariant pro-three surjection $R \twoheadrightarrow B$, the second-row data lift. Indeed $H^1(\langle\sigma\rangle, B) = 1$ by the coprime argument of A.15, so write $X = h^{-1}\sigma(h)$, lift h to $\widetilde{h}$ and take $\widetilde{h}^{-1}\sigma(\widetilde{h})$. This is a lift satisfying the exact involution relation. A lift of the local section in the first row can be made to have value one on c without changing its projection: its value at c is a cocycle in the pro-three kernel, and the same coprime argument conjugates it to one. The two rows can therefore be lifted independently and combined using (A.26.3).

A.19 supplies the first-row lift to N for every metabelian section. For a class-two section B.7– B.8 and A.19 say that this first row lifts exactly when its local class belongs to $\mathscr{L}_q$. This proves the asserted surjectivity and (A.26.1). If the target is unramified at q, all its full and metabelian lifts are unramified there by their zero-inertia results. The same inertia-kernel argument as in A.24 therefore identifies the fiber with A.20.

Now fix a ramified target in the image and a normalized lift. Its local lifting class is unique by the injectivity of the ramified cusp packet in class two. As in A.24, its local geometric centralizer is trivial both in the target and in B. Thus local lifts of the exact target with value one at c form one free conjugacy orbit under J_B^c. Use that conjugacy to fix the entire local first row to the chosen lift. No residual geometric conjugation remains, because the fixed local section has trivial centralizer. The remaining global fiber is consequently the literal set of second-row lifts of the target value at σ.

Write a second-row lift uniquely as jX_0, $j \in J_B$. Its involution equation is

$$jX_0\sigma(j)\sigma(X_0) = 1 \quad \Longleftrightarrow \quad j\Theta(j) = 1.$$

The identity $X_0\sigma(X_0) = 1$ proves that $\Theta^2 = 1$. Coprime H^1 for this action on J_B gives every solution as $h^{-1}\Theta(h)$. Two choices of h give the same solution precisely when they differ by left multiplication by J_B^{Θ}. This proves (A.26.2), including its coset orientation. The same argument with target N/N'' and kernel N'' gives the last assertion. □

Statement A.27 (Proved). The normalized integral Frobenius at an inert prime and the explicit Heisenberg test at seventeen. *Retain the free pro-three basis $u, v = cuc$ of P_3 from A.10, and write $[u, v] = uvu^{-1}v^{-1}$. For an inert rational prime $q \neq 3$, conjugate its local order-two complement in $P_3 \rtimes \langle c\rangle$ to $\langle c\rangle$. Let $\varphi_q \in P_3$ be the resulting arithmetic Frobenius of the prime of K_0 of norm q^2. Its image in the integral class-two quotient is exactly*

$$\varphi_q \bmod \gamma_3 P_3 = u^{a_q} v^{a_q} [u, v]^{-a_q^2/2}, \qquad a_q = \frac{\log(-q)}{\log 7} \in \mathbb{Z}_3. \tag{A.27.1}$$

The image in (A.27.1) is independent of the conjugator used to identify the local order-two complement with $\langle c\rangle$.

Put $\theta = \zeta_9$, $\beta^3 = \lambda$, and

$$u_i = 1 - \zeta^i\theta \quad (0 \le i \le 2), \qquad \omega = u_1 u_2^2, \qquad L = K_0(\theta, \beta, \omega^{1/3}). \tag{A.27.2}$$

Then L/K_0 is the full exponent-three class-two quotient of P_3: its Galois group has order 27 and presentation

$$\langle \sigma, \tau, \rho \mid \sigma^3 = \tau^3 = \rho^3 = 1, \quad \rho \text{ central}, \quad [\sigma, \tau] = \rho \rangle. \tag{A.27.3}$$

The field is Galois over $\mathbb{Q}$, and complex conjugation acts by

$$c\sigma c = \tau\sigma, \qquad c\tau c = \tau^{-1}, \qquad c\rho c = \rho^{-1}. \tag{A.27.4}$$

For every prime $q \equiv 8 \pmod 9$, the Frobenius of the prime of K_0 above q is trivial in (A.27.3). In particular this holds at 17, although $v_3(a_{17}) = 1$ in the integral formula (A.27.1). For example,

$$a_{17} \equiv 78 \pmod{81}, \qquad -a_{17}^2/2 \equiv 36 \pmod{81}. \tag{A.27.5}$$

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379); the first inert prime outside A.25 in Tasks A.6 and B.4.

Proof. The image of the local unramified group at q in the relative quotient over K_0 is procyclic with quotient C_2 and pro-three kernel. Its order-two subgroup is the prime-to-three component of this procyclic group. All complements to P_3 in $P_3 \rtimes C_2$ are P_3-conjugate, by the coprime argument of A.15. After the specified conjugation, φ_q commutes with c.

There is an exact integral logarithm description of the free class-two pro-three group on u, v: its underlying lattice has basis $U, V_0, [U, V_0]$, and multiplication is $X * Y = X + Y + \frac{1}{2}[X, Y]$. Here $U = \log u$ and $V_0 = \log v$. The fraction $1/2$ belongs to $\mathbb{Z}_3$; multiplication and the inverse maps are integral polynomial maps, so this identifies the pro-three group and the lattice with their compact topologies. The involution interchanges U, V_0 and negates $[U, V_0]$. Thus its fixed subgroup is exactly $\{\exp(a(U + V_0)) : a \in \mathbb{Z}_3\}$. The cyclotomic character sends both u, v to 7 and sends φ_q to q^2. Since the logarithm is injective on $1 + 3\mathbb{Z}_3$, it follows that $2a \log 7 = 2 \log(-q)$, giving $a = a_q$. The product formula in this lattice yields (A.27.1). Two allowed conjugators differ by an element of P_3^c; its image belongs to the displayed procyclic fixed subgroup and therefore commutes with the image of φ_q. This proves the asserted independence, with its exact ambiguity identified.

For the field in (A.27.2), direct multiplication gives

$$u_0 u_1 u_2 = \lambda, \qquad \frac{\sigma(\omega)}{\omega} = \frac{u_0^3}{\lambda}, \qquad c(\omega) = -\omega, \tag{A.27.6}$$

where initially $\sigma(\theta) = \zeta\theta$. The classes of ζ and λ are independent in $K_0^\times / K_0^{\times 3}$: the valuation of λ at the prime (λ) is one, whereas ζ is a unit and is not a cube in K_0. Consequently $K_0(\theta, \beta)/K_0$ has degree nine and λ is not a cube in $K_0(\theta)$. The kernel of

$$K_0(\theta)^\times / K_0(\theta)^{\times 3} \longrightarrow K_0(\theta, \beta)^\times / K_0(\theta, \beta)^{\times 3}$$

is the line generated by λ. To see this directly, a cube root in the cyclic cubic extension has an eigenvector under its Galois group and hence is a scalar multiple of $1, \beta$, or β^2. If the class of ω belonged to that line, the class of $\sigma(\omega)/\omega$ would be trivial. This contradicts (A.27.6). Thus adjoining $\gamma = \omega^{1/3}$ has degree three.

Choose the cube roots β, γ so that actual complex conjugation satisfies $c(\beta) = -\theta^2\beta$ and $c(\gamma) = -\gamma$. Each choice is possible by multiplying an initial cube root by a power of ζ; the ratios before and after that multiplication differ by the corresponding power of ζ. The following explicit automorphisms are then well-defined:

	θ	β	γ
σ	$\zeta\theta$	β	$u_0\gamma/\beta$
τ	θ	$\zeta\beta$	γ
ρ	θ	β	$\zeta\gamma$
c	θ^{-1}	$-\theta^2\beta$	$-\gamma$

(A.27.7)

Their powers and products, using (A.27.6), prove (A.27.3)–(A.27.4). All the radicands are units away from three, and the degrees are powers of three. The Kummer polynomials and their derivatives therefore show that the extension is unramified at every finite prime away

from three. It is consequently a quotient of P_3. The free exponent-three class-two group on two generators has the normal forms $\sigma^a\tau^b\rho^z$, with $a,b,z\in\mathbb{F}_3$, and order 27. The displayed quotient has this order, so the quotient map identifies its kernel exactly with the closed subgroup $P_3^3\gamma_3P_3$.

For a rational prime $q\equiv -1 \pmod 9$, the rational Frobenius in $\mathrm{Gal}(L/\mathbb{Q})$ acts on θ as $\theta\mapsto\theta^{-1}$. It consequently has the form gc with $g=\tau^b\rho^z$. Formula (A.27.4) gives $(gc)^2=gc(g)=1$. Its square is the Frobenius over K_0, proving its triviality in (A.27.3). Equivalently, at every residue choice of θ in $\mathbb{F}_{q^2}$, (A.27.6) gives $\omega^q=-\omega$, whence

$$\omega^{(q^2-1)/3}=(-1)^{(q+1)/3}=1.$$

The exponent on the right is even because the odd prime q is -1 modulo nine.

The program `code/a_frobenius17.py` checks (A.27.6) in the exact ring $\mathbb{Z}[\theta]=\mathbb{Z}[T]/(T^6+T^3+1)$, all 729 products under (A.27.4), and all three choices of θ at each of the 29 primes $q\equiv 8 \pmod 9$ through 1000. At seventeen one choice gives $\theta=9+11\zeta$ and $\omega=3+6\zeta$, with cube roots $2+\zeta$, $16+\zeta$, and $16+15\zeta$. The same program computes a_{17} by unique successive lifting of $7^a=-17$, without a floating-point logarithm. It gives (A.27.5) and the output through modulus 3^{12}. Finally $v_3(\log(-17))=v_3(-18)=2$ and $v_3(\log 7)=1$, proving $v_3(a_{17})=1$. □

Statement A.28 (Proved). The exact equivariant one-relator presentation at every inert prime. *Let $q\neq 3$ be any inert rational prime, and let $P_q=G_{K_0,\{3,q\}}(3)$ and $\Gamma_q=P_q\rtimes\langle c\rangle$. After the local-complement identification used in A.27, choose a tame generator t and an arithmetic Frobenius ϕ at the prime over q, with*

$$c(t)=t^{-1},\qquad c(\phi)=\phi,\qquad \phi t\phi^{-1}=t^{q^2}. \tag{A.28.1}$$

Choose a lift $x\in P_q$ of $u\in P_3$ and put $y=cxc$. The lift can and will be chosen with $\kappa_q(x)=0$ modulo three. Let $F=F_3(x,y,t)$ be the free pro-three group with involution $c(x)=y$, $c(y)=x$, $c(t)=t^{-1}$. Its natural equivariant map to P_q is surjective. There is a c-fixed pro-three word $W\in F$ mapping to the actual local Frobenius ϕ, and for every such choice the natural map gives the exact presentations

$$\begin{aligned} P_q &\simeq F/\overline{\langle\!\langle WtW^{-1}t^{-q^2}\rangle\!\rangle},\\ \Gamma_q &\simeq \bigl(F/\overline{\langle\!\langle WtW^{-1}t^{-q^2}\rangle\!\rangle}\bigr)\rtimes\langle c\rangle. \end{aligned} \tag{A.28.2}$$

These are continuous pro-three presentations, with the indicated finite operator group in the second line. In particular

$$P_q^{\mathrm{ab}}\simeq\mathbb{Z}_3^2\oplus\mathbb{Z}_3/(q^2-1),\qquad \dim_{\mathbb{F}_3}H^2(P_q,\mathbb{F}_3)=1. \tag{A.28.3}$$

At $q=17$, W lies in $\Phi(F)$, and the relator in (A.28.2) lies in the third Zassenhaus subgroup of F. Its image on killing t is the actual Frobenius of A.27; in particular its integral class-two image is given by (A.27.1) and (A.27.5).

For any of the geometric quotients B under consideration, (A.28.2) gives the following exact section parameterization. Normalize the value at c to one; assign the values $a,c(a),b$ at x,y,t. If $\Omega_W(a,b)$ denotes the value at W of this continuous free-group cocycle, the remaining equations are precisely

$$c(b)=b^{-1},\qquad \Omega_W(a,b)F_{q^2}(b)\Omega_W(a,b)^{-1}=b^{q^2}. \tag{A.28.4}$$

Classes are the orbits under B^c acting by

$$(a,b)\longmapsto(h^{-1}a\,x(h),\ h^{-1}bh). \tag{A.28.5}$$

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379); Tasks A.6 and B.4 beyond the prime families A.16 and A.25.

Proof. The Kummer group for K_0 with ramification allowed at three and q has the basis ζ, λ, q. The first two classes are pulled back from P_3, and the last has value one on t. Adjusting an initial lift of u by a power of t gives the specified value $\kappa_q(x) = 0$; because $c(\kappa_q) = -\kappa_q$, the value at y is also zero. Thus x, y, t form a basis of the Frattini quotient, and the continuous map $F \to P_q$ is onto. The involution is equivariant by its definition on these three generators.

The local generators (A.28.1) can be chosen in the relative tame local group: the unramified $\mathbb{Z}_3$ generator and its order-two component commute, and the latter acts on three-primary inertia as the Teichmüller scalar -1. Coprime complement conjugacy identifies this order-two element with the given global c. The local pro-three group maps injectively into P_q. For an explicit realization, the extension

$$K_0(\mu_{3^\infty}, q^{1/3^\infty})/K_0 \tag{A.28.6}$$

is pro-three and unramified outside three and q. At q its cyclotomic part contains every unramified extension of three-power degree: $q^2 \in 1 + 3\mathbb{Z}_3$ has infinite order, so the local cyclotomic degrees are unbounded. The uniformizer roots supply every tame three-power ramification degree. The standard tame description then identifies its local compositum with the maximal pro-three extension of $K_{0,q}$. Consequently the local decomposition map into P_q has trivial kernel.

Since ϕ is c-fixed and $F \to P_q$ is an equivariant pro-three surjection, the coprime H^1 argument of A.15 proves $F^c \to P_q^c$ is onto. This supplies $W \in F^c$ as stated. Put $r = WtW^{-1}t^{-q^2}$ and $G_W = F/(r)$, where (r) denotes the closed normal subgroup it generates. As

$$c(r) = Wt^{-1}W^{-1}t^{q^2} = t^{-q^2}r^{-1}t^{q^2},$$

this subgroup is stable under c. There is a surjective equivariant map $G_W \to P_q$, inducing an isomorphism on $H^1(-, \mathbb{F}_3)$. Here $r \in \Phi(F)$ since $q^2 \equiv 1 \pmod 3$.

The degree-two comparison is the exact one proved in A.14: full-to-maximal-pro-three inflation is injective, and the full global group has

$$H^2(G_{K_0,\{3,q\}}, \mathbb{F}_3) \simeq \{(z_3, z_q) \in (\tfrac{1}{3}\mathbb{Z}/\mathbb{Z})^2 : z_3 + z_q = 0\}. \tag{A.28.7}$$

The Kummer, Picard, and Brauer comparisons are precisely Milne II.2.9, II.2.1, and II.2.2(a) [Mil06], applied to the localization of the same principal ideal domain. Thus restriction $H^2(P_q, \mathbb{F}_3) \to H^2(D_{q^2}, \mathbb{F}_3)$ is injective. The assignments $\phi \mapsto W$, $t \mapsto t$ define a continuous map $D_{q^2} \to G_W$, whose composite to P_q is the actual local map. It follows that inflation $H^2(P_q, \mathbb{F}_3) \to H^2(G_W, \mathbb{F}_3)$ is injective. The five-term kernel argument proved in A.14 now shows that $G_W \to P_q$ is an isomorphism. This proves (A.28.2), including the stated operator action.

Abelianizing its single relator gives exactly $(1 - q^2)t$ and no relation on x, y, proving the first formula of (A.28.3). This abelianization has nonzero finite torsion; P_q is therefore not free pro-three. The minimal relation criterion used in A.10 gives $H^2(P_q, \mathbb{F}_3) \neq 0$, while its injection into $H^2(D_{q^2}, \mathbb{F}_3)$ bounds the dimension by one. The latter local group has dimension one, as follows from the Hochschild–Serre sequence for $\mathbb{Z}_3(1) \rtimes \mathbb{Z}_3$ with trivial $\mathbb{F}_3$ coefficients. This proves the second formula.

At seventeen the Frobenius values on ζ, λ are zero by (A.25.4). The chosen local Frobenius fixes the uniformizer roots, so its value on q is also zero modulo three. Conjugating the local embedding does not change these degree-one values. Therefore $W \in \Phi(F)$. If $D_n(F)$ denotes the Zassenhaus filtration, then $\Phi(F) = D_2(F)$, so $[W, t] \in D_3(F)$; also $t^{q^2-1} = t^{288} \in D_9(F)$.

The relator thus belongs to $D_3(F)$. The map killing t identifies the quotient with the unramified group P_3 and sends W to the normalized Frobenius already computed in A.27.

Finally a normalized cocycle on $F \rtimes C_2$ is exactly the arbitrary value a on x, its conjugate on y, and the value b on t satisfying $c(b) = b^{-1}$. The continuous value at the pro-three word W is defined by the free pro-three universal property; this defines Ω_W without replacing W by a truncated word. Arithmetic inertia at $q \neq 3$ acts trivially on the original geometric group, while W acts as its actual Frobenius F_{q^2}. Evaluating r gives exactly the second equation (A.28.4), and conjugating a normalized cocycle gives (A.28.5). These computations prove both directions of the parameterization. The word W in this assertion retains its possible dependence on t; (A.27.1) computes its stated quotient, not that full dependence. Solving (A.28.4) in the full geometric group for every prescribed metabelian target at seventeen remains the specific unresolved step recorded in the ledger. □

Statement A.29 (Proved). Local cuspidality at every finite derived level and its global one-prime consequences. *Write $N^{(0)} = N$ and $N^{(m+1)} = \overline{[N^{(m)}, N^{(m)}]}$, and put $Q_m = N/N^{(m)}$ for $m \geq 2$. For every finite extension $k/\mathbb{Q}_q$, $q \neq 3$, every continuous section with geometric quotient Q_m is either unramified or lies in exactly one actual cusp packet. Its inertia is trivial exactly when its abelian inertia is zero. Every such local section lifts to N, and the map from full to Q_m sections is bijective on the ramified classes. Thus the exact local description in A.19 holds with Q_m in place of N/N'', including the actual central Frobenius coordinates of its image in class two.*

For every $q \in \Sigma \cup \Xi$ and every $m \geq 2$, the global map

$$\mathrm{Sec}_{\{3,q,\infty\}}(N) \longrightarrow \mathrm{Sec}_{\{3,q,\infty\}}(Q_m) \tag{A.29.1}$$

is onto. For ramified targets its fiber is the right-coset set $N^{(m)}/(N^{(m)})^c$ when $q \in \Sigma$, and the left-coset set $(N^{(m)})^{\Theta}\backslash N^{(m)}$ when $q \in \Xi$, with the chosen-lift involution Θ of A.26. Unramified fibers are the exact unramified fibers of A.20 for the map $N \to Q_m$.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379); the successive derived approximations in Tasks A.6 and B.4.

Proof. We first identify the precise peripheral intersection statement used below. Each boundary generator c_d extends to a free pro-three basis of N: in the standard surface presentation, eliminate a different boundary generator. Yamaguchi's Proposition 1.1.6 [Yam23] gives the centralizer for every nonzero integer power. For $a = 3^s u$ with $u \in \mathbb{Z}_3^\times$, continuous exponentiation gives $c_d^{3^s} = (c_d^a)^{u^{-1}}$ and $c_d^a = (c_d^{3^s})^u$. The two elements generate the same closed procyclic subgroup and hence have the same centralizer. We therefore obtain

$$C_{Q_j}(c_d^a) = \langle c_d \rangle \quad (j \geq 2,\ a \in \mathbb{Z}_3 \setminus \{0\}). \tag{A.29.2}$$

If $g c_d^a g^{-1} = c_e^b$ with $a, b \neq 0$, abelianization gives $a e_d = b e_e$. The distinct boundary lines are disjoint away from zero, and each e_d is primitive. It follows that $d = e$ and $a = b$. Formula (A.29.2) then forces $g \in \langle c_d \rangle$. This proves the required intersection and normalizer claims through explicit maps to V and the stated centralizer theorem.

We induct on m, starting with A.19. Let a local section in Q_m have tame value t, and denote its image in Q_{m-1} by $\bar{t}$. By induction $\bar{t}$ is either one or conjugate to c_d^a, $a \neq 0$. In the latter case choose such a conjugator and lift it through $Q_m \to Q_{m-1}$; this changes the representative by an allowed geometric conjugation and makes $\bar{t} = c_d^a$ exactly.

Choose a decreasing cofinal sequence of characteristic open normal subgroups M_n of Q_{m-1}, and let $H_n \subset N$ be the inverse image of $\langle \bar{t} \rangle M_n$. Then

$$N^{(m-1)} \subset H_n, \qquad N^{(m)} \subset H_n', \qquad H_\infty := \bigcap_n H_n = \begin{cases} N^{(m-1)}, & \bar{t} = 1, \\ N^{(m-1)}\langle c_d^a \rangle, & \bar{t} = c_d^a. \end{cases} \tag{A.29.3}$$

These H_n need not be normal in N; the finite covers and their boundary points below are described by cosets and double cosets, respectively. The characteristic choice and the section's Frobenius relation make $H_n/N^{(m)}$ stable under the section. Pulling back $(H_n/N^{(m)})s(G_k)$ from $Q_m \rtimes G_k$ gives an actual arithmetic cover with geometric group H_n. It contains original arithmetic inertia because $t \in H_n/N^{(m)}$. Thus this cover has the good tame compactification used in A.17–A.19. The inclusion $N^{(m)} \subset H_n'$ identifies its actual geometric abelianization with that of $H_n/N^{(m)}$. In particular the chosen lift of t has the exact Frobenius eigenvalue $\#\mathbb{F}_k$ in H_n^{ab}, rather than only an eigenvalue in a further quotient. The compactification weight argument places it in the integral boundary submodule at every n.

As proved in A.19 by stabilization in every finite quotient and compact lifting of cosets,

$$H_\infty^{\mathrm{ab}} \simeq \varprojlim_n H_n^{\mathrm{ab}}. \tag{A.29.4}$$

We compute the inverse limit of the boundary modules with their actual transition maps. The boundary points above e are $H_n\backslash N/I_e$; the loop at $H_n g I_e$ is the appropriate conjugate of $c_e^{j_n(g,e)}$, where

$$j_n(g,e) = [I_e : I_e \cap g^{-1}H_n g]. \tag{A.29.5}$$

Transition maps sum over inverse images and multiply each term by the ratio of these ramification indices. The diagonal relation has transition map equal to the covering degree. The index $[N : H_n]$ is unbounded, because the group Q_{m-1} cannot have an open procyclic subgroup: its quotient V has rank ten. The diagonal tower consequently has zero limit, and its first derived limit is zero by the compact equation argument in A.19. The inverse limit of boundary modules is therefore the inverse limit of the weighted free boundary modules in (A.29.5).

If $\bar{t} = 1$, every intersection $I_e \cap g^{-1}H_\infty g$ is trivial, since each cusp group injects into Q_{m-1} by its primitive abelian image. Every ramification index tends to infinity. This growth is uniform on each clopen set of the compact double coset space: otherwise nested nonempty closed sets with bounded indices would give a point with nontrivial limiting intersection. The weighted transition maps are thus divisible by arbitrary powers of three at each fixed finite stage. Their inverse limit is zero. By (A.29.4), t is zero in $H_\infty^{\mathrm{ab}} = N^{(m-1)}/N^{(m)}$, proving $t = 1$.

Now let $\bar{t} = c_d^a$ and write $a = 3^s\varepsilon$ with $\varepsilon \in \mathbb{Z}_3^\times$. A bounded index in (A.29.5) is equivalent to a nontrivial intersection of $gI_e g^{-1}$ with $\langle c_d^a \rangle$ in Q_{m-1}. By (A.29.2), this forces $e = d$ and the image of g to belong to I_d. There is therefore exactly one surviving boundary double coset in $H_\infty\backslash N/I_d$. Its ramification index stabilizes at 3^s. Its 3^s sheets under $I_d/\langle c_d^a \rangle$ constitute one boundary point, not 3^s independent boundary generators. On every clopen set away from this point the same compactness argument gives uniform unbounded ramification. The inverse boundary module is consequently the single copy of $\mathbb{Z}_3(1)$ generated by the class of $c_d^{3^s}$. Projecting H_∞ to $\langle c_d^a \rangle$ shows that the class of t is $\varepsilon[c_d^{3^s}] = [c_d^a]$ in (A.29.4).

Put $A_m = N^{(m-1)}/N^{(m)}$ and let T denote conjugation by c_d^a. The image of H_∞ in Q_m is $A_m \rtimes \langle c_d^a \rangle$. Its closed commutator subgroup is exactly $(T-1)A_m$: the image is closed because A_m is compact. The equality of abelianizations just proved gives $t(c_d^a)^{-1} \in (T-1)A_m$, which is equivalent to $t = h^{-1}c_d^a h$ for some $h \in A_m$. This proves actual cusp conjugacy. After making this conjugation, compare the Frobenius value with that of the actual tangential cusp section. Their ratio centralizes c_d^a, hence lies in I_d by (A.29.2). The complete local section therefore lies in the actual cusp packet. Its parameters and uniqueness are exactly the Kummer parameters of A.19 and B.8, and its full lift is the corresponding cusp section. In the unramified case an arbitrary Frobenius value lifts to N, as proved in A.19. This completes the induction and the local statement.

For the global consequences, the arithmetic presentations A.15–A.16 and A.25 are independent of the geometric quotient. Replace the local metabelian lifting input in A.24 and A.26 by the local result just proved. The local ramified lift is still unique and has trivial full local geometric centralizer; the independent involution datum in A.26 still lifts by coprime H^1. The identical explicit maps then prove (A.29.1) and its stated fibers with kernel $N^{(m)}$. For an unramified target, the local zero-inertia result forces all full lifts to be unramified at q, so the comparison with A.20 is exact. □

Statement A.30 (Proved). The mixed Heisenberg test at seventeen and the exact first relation degree. *Use the presentation at $q = 17$ in A.28. For its Frobenius word and defining relator, the stronger exact conclusions are*

$$W \in D_3(F) \setminus D_4(F), \qquad r = WtW^{-1}t^{-289} \in D_4(F) \setminus D_5(F). \tag{A.30.1}$$

Here $D_n(F)$ is the Zassenhaus filtration, equivalently the filtration by degree in the mod-three Magnus expansion of the free pro-three group $F = F_3(x, y, t)$. If x, y, t have Magnus variables X, Y, T, the degree-four part of $r-1$ has coefficient 2 on both X^3T and Y^3T. These coefficients are independent of the other degree-three coefficients of $W - 1$.

The arithmetic input detecting the only possible degree-two component of W is the following explicit Heisenberg extension. Let

$$\alpha^3 = 3, \qquad \beta^3 = 17, \qquad u_0 = 2 + \alpha^2, \qquad u_i = \sigma^i(u_0), \quad \sigma(\alpha) = \zeta\alpha,$$

and set

$$\omega = u_1u_2^2, \qquad L_{3,17} = K_0(\alpha, \beta, \omega^{1/3}). \tag{A.30.2}$$

This field is Galois over K_0, has the group (A.27.3), and is unramified outside three and seventeen. At the prime with $\alpha \equiv 7 \pmod{17}$, the Frobenius fixing the uniformizer roots has trivial image in the center of this Heisenberg quotient.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379); the next arithmetic relation at the first remaining inert prime in Tasks A.6 and B.4.

Proof. The norm calculation is exact:

$$u_0u_1u_2 = 2^3 + \alpha^6 = 17, \qquad \frac{\sigma(\omega)}{\omega} = \frac{u_0^3}{17}. \tag{A.30.3}$$

The classes of 3 and 17 in $K_0^\times/K_0^{\times 3}$ are independent, by their valuations at the primes above three and seventeen. The degree and automorphism argument of A.27 applies with these two radicands and the norm witness u_0. In particular the last cubic extension in (A.30.2) is nontrivial: otherwise the class of ω over $K_0(\alpha)$ would lie on the line generated by 17, contradicting (A.30.3). For $\gamma^3 = \omega$, the explicit automorphisms are

	α	β	γ
σ	$\zeta\alpha$	β	$u_0\gamma/\beta$
τ	α	$\zeta\beta$	γ
ρ	α	β	$\zeta\gamma$

(A.30.4)

and $[\sigma, \tau] = \rho$. Choose positive real cube roots α, β in the fixed complex embedding. Complex conjugation interchanges u_1, u_2 and satisfies

$$c(\omega)\omega = (17/u_0)^3, \qquad c(\gamma) = 17/(u_0\gamma).$$

The second equality follows from $|\gamma|^2 = 17/u_0 > 0$, for any choice of the cube root γ. Thus the field is Galois over $\mathbb{Q}$, with

$$c\sigma c = \sigma^{-1}, \qquad c\tau c = \tau^{-1}, \qquad c\rho c = \rho. \tag{A.30.5}$$

The radicands are units away from three and seventeen, so the Kummer polynomials are unramified at every other finite prime.

At the chosen prime over seventeen, the residue field of K_0 is $\mathbb{F}_{17^2}$ and $\alpha = 7$ is a cube root of 3. In the basis $1, \zeta$, direct finite-field multiplication gives

$$(u_0, u_1, u_2) = (0,\ 4 + 2\zeta,\ 2 + 15\zeta), \qquad \omega = 7 + 10\zeta, \qquad \omega^{96} = 1. \tag{A.30.6}$$

Here ω is a unit, so its cubic residue is exactly the Frobenius value on γ. The specified local Frobenius fixes α and the uniformizer root β, and (A.30.6) shows that it fixes γ as well. Its Heisenberg image is therefore the identity. In particular this value is unchanged by conjugating the local embedding to align its order-two complement with c.

We now compare this arithmetic calculation with the actual free presentation. By A.28, $W \in D_2(F)$ and $c(W) = W$. The quotient $D_2(F)/D_3(F)$ has basis

$$[x, y], \quad [x, t], \quad [y, t] \quad \text{over } \mathbb{F}_3.$$

Indeed commutators generate this quotient, cubes already have Magnus degree three, and these three degree-two commutator polynomials are linearly independent. The involution negates the first basis vector and sends the last two to each other's negatives. Its fixed subspace is exactly the line generated by $[x, t] - [y, t]$.

Under the quotient (A.30.2), the Frattini images of x, y, t on the two Kummer coordinates $(3, 17)$ are

$$x \longmapsto (1, 0), \qquad y \longmapsto (2, 0), \qquad t \longmapsto (0, 1). \tag{A.30.7}$$

For the first coordinate this uses $\kappa_3 = -\kappa_\zeta - \kappa_\lambda$ and the two Frobenius rows of A.10. The second uses the lift choice in A.28. Consequently the fixed wedge generator maps to $(1 - 2)\rho = -\rho \neq 0$. The trivial Frobenius value in (A.30.6) forces the class of W on that line to vanish. This proves $W \in D_3(F)$.

In F^{ab}, its coordinates on x, y are a_{17}, a_{17} by A.27, and its coordinate on t is zero because it is fixed by the involution which inverts t. We have $a_{17} \equiv 6 \pmod 9$. Setting $Y = T = 0$ in its Magnus expansion therefore gives $(1 + X)^{a_{17}}$. The coefficient of X^3 is

$$\binom{a_{17}}{3} \equiv a_{17}/3 \equiv 2 \pmod 3.$$

The same holds for Y^3. Thus $W \notin D_4(F)$. If W_3 is its homogeneous degree-three term, the first possible term of $[W, t] - 1$ is $W_3T - TW_3$. The coefficient of X^3T in this expression is exactly 2, and likewise for Y^3T; no other monomial of W_3 can contribute to either coefficient. The factor $t^{1-289} = t^{-288}$ has first possible Magnus degree nine, since $288 = 9 \cdot 32$. It follows that r has exact degree four, proving (A.30.1).

The exact ring identities, all three residue choices, and the nonzero Magnus coefficients with arbitrary remaining cubic terms are reproduced by `code/a_frobenius17_mixed.py` and its output. This calculation determines the first relation degree; the remaining coefficients of W and the full lifting equation (A.28.4) continue to be retained in the unresolved global problem. □

Statement A.31 (Proved). The nonzero cubic Frobenius term at seventeen and its failure to be a cube. *Let $L_0 = K_0(\alpha)$ with $\alpha^3 = 3$, and let $H = G_{L_0,\{3\}}(3) \subset P_3$. Put*

$$\pi = \frac{\lambda}{\alpha}, \qquad \eta = \alpha^2 - 2, \qquad \delta = 1 + \zeta^2\alpha + \zeta\pi. \tag{A.31.1}$$

The four Kummer classes of ζ, π, η, δ form an $\mathbb{F}_3$-basis of $H^1(H, \mathbb{F}_3)$. At every prime of L_0 above seventeen, the Frobenius row on this basis is

$$(0, 0, 0, 1). \tag{A.31.2}$$

Choose $m \in P_3^c$ whose abelian coordinates in the basis u, v are $(1, 1)$, and put $n = vu^{-1}$. Then m, n are a free pro-three basis, $c(m) = m$, $c(n) = n^{-1}$, and the normalized Frobenius of A.27 satisfies the exact congruence

$$\varphi_{17} \equiv m^6[n, [n, m]] \pmod{D_4(P_3)}. \tag{A.31.3}$$

Moreover φ_{17} is not a cube of any element of P_3.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379); the explicit nonprimitive Frobenius in Tasks A.6 and B.4.

Proof. The subgroup H is exactly $\ker(\kappa_3 : P_3 \to \mathbb{F}_3)$. Indeed every finite Galois three-extension of L_0 unramified outside three has a Galois closure over K_0 with three-group Galois group: take the compositum of its three conjugates, whose kernel over L_0 embeds in a product of three-groups. This proves the stated comparison with the open subgroup of P_3. The subgroup is free of rank four by the pro-three Schreier formula. Thus its first cohomology has dimension four.

The element π is a unit away from three. For $\sigma(\alpha) = \zeta\alpha$, direct multiplication gives

$$N_{L_0/K_0}(\eta) = 1, \qquad \delta^3 = \zeta\eta^2\sigma(\eta), \qquad N_{L_0/K_0}(\delta) = \zeta. \tag{A.31.4}$$

Consequently η and δ are units at every finite prime. The following four Frobenius evaluations prove independence of their Kummer classes:

q	$\bar{\zeta}$	$\bar{\alpha}$	κ_ζ	κ_π	κ_η	κ_δ
2	ζ	1	1	2	0	2
2	ζ	ζ	1	1	2	2
2	ζ	ζ^2	1	0	1	0
61	13	4	2	0	1	0

(A.31.5)

The first three residue fields are $\mathbb{F}_4$ with $\zeta^2 + \zeta + 1 = 0$; the last is $\mathbb{F}_{61}$. In each row, raise the unit residue to $(\#\mathbb{F} - 1)/3$ and express the answer as a power of the displayed residue of ζ. The integer determinant of the four-by-four matrix in (A.31.5) is 2, hence is nonzero modulo three. The dimension calculation then proves the basis assertion.

At seventeen choose the residue field $\mathbb{F}_{17^2}$ and first take $\alpha = 7$. The four residues are

$$\zeta, \qquad 5 + 12\zeta, \qquad 13, \qquad 16 + 3\zeta.$$

Their 96th powers give $(1, 1, 1, \zeta)$. The other two choices $\alpha = 7\zeta, 7\zeta^2$ give the same four Kummer values. This proves (A.31.2), including the effect of any conjugation permuting the three primes above seventeen.

Coprime H^1 for the kernel of $P_3 \to P_3^{\mathrm{ab}}$ makes $P_3^c \to (P_3^{\mathrm{ab}})^c$ onto. This gives m with the specified abelian vector. Together with $n = vu^{-1}$ its Frattini vectors have determinant 2, so m, n are a free basis. The actions of c are exactly those stated, and

$$\kappa_3(m) = 0, \quad \kappa_3(n) = 1, \quad \kappa_\zeta(m) = 1 \quad \text{in } \mathbb{F}_3, \tag{A.31.6}$$

by the two rows of A.10. Thus conjugation by n acts on L_0 as σ. One completely specified choice of m uses only continuous pro-three square roots. Put $d_0 = uc(uv)^{-1/2}$ and $h_0 = (d_0c)^{1/2} \in P_3$, and take $m = h_0^{-1}(uv)h_0$. Indeed $(uc)^2 = uv$, so d_0 is an involution commuting with uv. The identity $c(d_0c)c = (d_0c)^{-1}$ gives $h_0ch_0^{-1} = d_0$ and hence $c(m) = m$. Its abelian vector is still $(1, 1)$. Square roots here are the unique continuous powers $g^{1/2}$ in the closed procyclic group generated by g; this defines every word and comparison in the formula.

We give the finite-group comparison which permits use of (A.31.2) in degree three. A Schreier basis of H is $n^3, m, nmn^{-1}, n^2mn^{-2}$. Modulo $\Phi(H)$ these form a basis of an elementary abelian group of order 3^4. Conjugation by n fixes n^3 and cyclically permutes the other three basis vectors. Hence $P_3/\Phi(H)$ has nilpotence class at most three, exponent dividing nine, and commutator subgroup of exponent three. The defining product formula

$$D_4(P_3) = P_3^9\, \gamma_2(P_3)^3\, \gamma_4(P_3)$$

therefore proves $D_4(P_3) \subset \Phi(H)$. Every Kummer character in (A.31.5), in particular $\ell = \kappa_\delta$, annihilates this subgroup.

The vector space $D_3(P_3)/D_4(P_3)$ has basis

$$m^3, \quad n^3, \quad [m,[m,n]], \quad [n,[n,m]]. \tag{A.31.7}$$

Collection gives these generators; their Magnus polynomials are respectively M^3, N^3, $M^2N + MNM + NM^2$, and $N^2M + NMN + MN^2$, which are independent. On this basis c has signs $+, -, -, +$. The Frobenius lies in D_3 by A.27 and is fixed by c, so its class has form $Am^3 + B[n,[n,m]]$. Its abelian coordinates in the m, n basis are $(a_{17}, 0)$, whence $A = a_{17}/3 = 2$ modulo three.

For $h \in H$, Kummer theory identifies $\ell(nhn^{-1})$ with $\kappa_{\sigma^{-1}\delta}(h)$. Since $-2 = 1$ in $\mathbb{F}_3$, we obtain

$$\begin{aligned} \ell([n,[n,m]]) &= ((\sigma^{-1} - 1)^2\kappa_\delta)(m) \\ &= \kappa_{N_{L_0/K_0}\delta}(m) = \kappa_\zeta(m) = 1. \end{aligned} \tag{A.31.8}$$

Also $\ell(m^3) = 0$. Formula (A.31.2) gives $\ell(\varphi_{17}) = 1$, so $B = 1$. This proves the full degree-three congruence (A.31.3), rather than only its cyclotomic component.

Finally, a cube root of φ_{17} in P_3 would have abelian vector $(a_{17}/3, a_{17}/3)$ in the u, v basis. Its κ_3 value is zero, so it would belong to H. The character κ_δ annihilates every cube in H, contradicting (A.31.2). This proves the last assertion.

The exact unit identities and root are verified by `a_frobenius17_unit_root.py` in `code/`; the residue matrix and all three Frobenius rows at seventeen are verified by `a_frobenius17_sunits.py` in the same directory. Their outputs are included. An initial attempted list $\zeta, \pi, \eta, \sigma(\eta)$ had rank three, not four; the exact cubic root δ in (A.31.4) supplies the missing unit. No basis assertion relies on that rejected list. □

Statement A.32 (Proved). The exact marked global image and ramified fibers on $\Sigma \cup \Xi$. *Put $R = N/\gamma_3(I_{\mathrm{mark}})$, $A = N/\gamma_3 N$, and $J_R = \ker(R \to A)$. For every $q \in \Sigma \cup \Xi$, the three global class-two images coincide and are exactly*

$$\begin{aligned} \operatorname{im}\big(\mathrm{Sec}_{S_q}(N) \to \mathrm{Sec}_{S_q}(A)\big) &= \operatorname{im}\big(\mathrm{Sec}_{S_q}(N/N'') \to \mathrm{Sec}_{S_q}(A)\big) \\ &= \operatorname{im}\big(\mathrm{Sec}_{S_q}(R) \to \mathrm{Sec}_{S_q}(A)\big) = \mathrm{res}_q^{-1}(\mathscr{L}_q), \end{aligned} \tag{A.32.1}$$

where $S_q = \{3, q, \infty\}$ and $\mathscr{L}_q$ is the actual local unramified and cusp-packet set of A.24.

Fix a ramified class-two target in this image and let $d \in D(\mathbb{Q}_q)$ be its unique local cusp. Define its local marked lifting-class set by

$$\mathcal{L}_{q,d} = \begin{cases} \{*\}, & d \neq Q_c, \\ \mathcal{C}_{\mathbb{Q}_q,Q_c}, & d = Q_c, \end{cases} \tag{A.32.2}$$

using exactly the reference and quotient of B.12–B.13. Restriction gives a surjection from the global marked fiber onto $\mathcal{L}_{q,d}$. For each element of this local class set, choose a normalized global marked lift above it. The inverse image of that element is precisely

$$\begin{cases} J_R/J_R^c, & q \in \Sigma, \\ J_R^\Theta \backslash J_R, & q \in \Xi, \end{cases} \tag{A.32.3}$$

with right and left cosets, respectively. In the second case $\Theta = \mathrm{Ad}(X_0) \circ \sigma$ is the actual involution specified by the chosen lift in A.26. Thus (A.32.2)–(A.32.3) describe the entire fiber without identifying different local marked classes or asserting a canonical product decomposition. Targets unramified at q have exactly the unramified fiber of A.20 for $R \to A$.

The exact image of $\mathrm{Sec}_{S_q}(N) \to \mathrm{Sec}_{S_q}(R)$ consists of the classes unramified at q, the unmarked ramified cusp classes, and the marked ramified classes whose local parameter in $\mathcal{C}_{\mathbb{Q}_q,Q_c}$ is zero. Its nonempty ramified fibers are given by (A.32.3) with J_R replaced by $\gamma_3(I_{\mathrm{mark}})$ and the corresponding chosen full lift. In particular there are global marked sections supported at $\{3, q\}$ which have no full lift, although their class-two image has a full lift. These examples still have no full lift after any finite number-field extension.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379); the marked higher image and fibers in Task A.6.

Proof. The local marked classification B.12–B.14 proves that its exact class-two image is $\mathscr{L}_q$. The local fiber over an unmarked ramified cusp is a singleton; the local fiber over the marked cusp is the full group $\mathcal{C}_{\mathbb{Q}_q,Q_c}$. Each class in the latter group is an actual local section, with the Frobenius representative $c_{Q_c}^r z F_0$ and inertia $c_{Q_c}^a$ of (B.12.3). Its class-two image retains a, r and kills z through the actual morphism $\gamma_2 I_{\mathrm{mark}}/\gamma_3 I_{\mathrm{mark}} \to A$.

For $q \in \Sigma$, a normalized global section is its arbitrary local pair modulo R^c, by A.15–A.16. Any prescribed local marked lifting class therefore globalizes. To make its projection the exact local pair of the given global class-two target, lift the required local geometric conjugator through $R \to A$. There is no further arithmetic relation. For $q \in \Xi$, first normalize the local c-value to one using coprime H^1 in the pro-three kernel. The independent involution value in the second row of (A.26.3) lifts the prescribed class-two value by the same coprime argument. It combines with the selected local row to give the required global marked section. This proves surjectivity onto (A.32.2) and the marked equality in (A.32.1). The other two equalities are A.24 and A.26.

We check the centralizers used to pass from local to global conjugation. The class-two local section has trivial geometric centralizer: its image in V would be Frobenius-fixed, hence zero by the weight calculation, and the remaining element in $\bigwedge^2 V$ is zero for the same reason. At an unmarked ramified cusp in R, its projection to the filled full group has trivial centralizer by B.7, and the kernel has no inertia-fixed element in either M or Z by the free label actions in B.14. Hence its centralizer in R is zero. At the marked cusp the centralizer of the inertia element is $\langle c_{Q_c} \rangle Z$ by (B.12.6). A full-section centralizer in this group has zero cusp coordinate because multiplication by $q - 1$ is injective on $\mathbb{Z}_3$; its central coordinate is also zero, because $q^2 \bigwedge^2 \Phi - 1$ is injective on Z. This last injectivity follows on every finite

free permutation quotient from its eigenvalues, and then in the inverse limit. It is unaffected by the central Frobenius parameter z. Thus every ramified local marked section used here has trivial geometric centralizer.

Fix a local marked lifting class. Exact local lifts of the target pair form one free conjugacy orbit under J_R. Indeed their conjugator projects to a centralizer of the target, which has just been proved trivial. For Σ, the remaining global conjugators are J_R^c. With the cocycle convention of A.24, their action on the local orbit parameter is right multiplication, giving J_R/J_R^c. For Ξ, normalized local rows form a free orbit under J_R^c. Fix one such row by this conjugation; no geometric conjugator remains. The remaining data are exactly the lifts of the independent involution value. As in (A.26.2), writing them as jX_0 gives $j\Theta(j) = 1$ and the bijection $J_R^\Theta \backslash J_R \to \{j : j\Theta(j) = 1\}$. This proves (A.32.3) with its coset orientations.

The same presentation arguments apply to $N \to R$. B.7 and B.12–B.14 show that a local marked class has a full lift exactly for the parameters listed in the statement. The lifting class is unique when ramified. Thus the local condition is necessary, the arithmetic presentations make it sufficient, and the same centralizer and coset argument gives the asserted fibers. If a class-two or marked target is unramified at q, the zero-abelian-inertia results of B.7 and B.12 force every higher lift here to be unramified there as well. The comparison with A.20 is consequently exact.

For existence of the last examples, start with any nonzero marked tame exponent and choose a nonzero element of $\mathcal{C}_{\mathbb{Q}_q,Q_c}$; B.13 gives such elements, including an explicit central commutator detected in a finite label quotient. The preceding construction gives the global marked section, while its class-two image belongs to (A.32.1). The group $\mathcal{C}_{k,Q_c}$ is the unramified first cohomology of the actual central coefficient module Z in B.12. On this module the cusp inertia acts trivially. Its tame first-cohomology component has Frobenius $Q \bigwedge^2 \Phi$, with no invariant vector, so it is the whole $H^1_{\mathrm{cts}}(G_k, Z)$. For any finite local extension, restriction followed by corestriction on this continuous cohomology is multiplication by its degree; this is the usual finite coset-sum chain map and is valid for compact coefficients. B.13 proves that the source group is torsion-free. Its restriction is therefore injective. A nonzero central parameter remains nonzero at every place of every finite number-field extension above q, and the local full image still has central parameter zero by B.12. This proves persistence of the failure of full lifting. □

Statement A.33 (Proved). A class-three lift for every finite-support locally cuspidal class-two section. *Let S' be any finite set of rational primes containing three, with the real place included. Put $A = N/\gamma_3 N$ and $M_3 = \gamma_3 N/\gamma_4 N$. Every section of A over $G_{\mathbb{Q},S'}$ whose restriction at each $q \in S' \setminus \{3\}$ belongs to the actual local image $\mathscr{L}_q$ of B.8 has a section lift to $N/\gamma_4 N$ over the same arithmetic group. Its lifting fiber is a torsor under*

$$H^1(G_{\mathbb{Q},S'}, M_3) = H^1(G_{\mathbb{Q},\{3,\infty\}}, M_3) = \mathbb{Z}_3^{166} \oplus (\mathbb{Z}/3)^7 \oplus (\mathbb{Z}/9)^{56} \oplus (\mathbb{Z}/27)^{25} \oplus (\mathbb{Z}/81)^3. \quad \text{(A.33.1)}$$

These assertions concern the actual common class-three quotient of the full and metabelian groups.

There is also an exact obstruction to choosing this global lift with prescribed actual cuspidal local class-three restrictions. Let $J \subset S' \setminus \{3\}$ be the primes where the class-two target is ramified. Choose any global class-three lift s_3. At each $q \in J$ the unique actual cuspidal class-three lifting class determines a difference $\Delta_q(s_3) \in M_3/(F_q - 1)M_3$. The element

$$[(\Delta_q(s_3))_{q\in J}] \in \operatorname{coker}\left(H^1(G_{\mathbb{Q},\{3,\infty\}}, M_3) \xrightarrow{\mathrm{res}} \bigoplus_{q\in J} M_3/(F_q - 1)M_3 \right) \quad \text{(A.33.2)}$$

is independent of s_3. It vanishes exactly when a global class-three lift can have all the indicated cuspidal restrictions. Every full or metabelian global lift of the class-two target forces (A.33.2) to vanish.

Proof. The action on the central kernel M_3 is the actual action induced from V; conjugation by the class-two section does not change it. Its finite quotients have three-group image over K_0. The composition-series argument in A.1 therefore gives

$$H^i(G_{\mathbb{Q},\{3,\infty\}}, M_3/3^n) = 0 \quad (i \geq 2).$$

For $X = \operatorname{Spec} \mathbb{Z}[1/3]$, apply the actual localization triangle (A.1.2) with $M_3/3^n$ and remove the finite set $S' \setminus \{3\}$. Its degree-two term injects into the sum of the residue groups $H^1(\mathbb{F}_q, M_3(-1)/3^n)$. These are precisely the local groups $H^2(G_{\mathbb{Q}_q}, M_3/3^n)$, by the tame inertia calculation in A.1. Thus the actual global lifting obstruction, restricted to each local arithmetic group, is its corresponding local lifting obstruction. An actual local full lift exists by membership in $\mathscr{L}_q$, so all these restrictions are zero.

The finite-S' degree-one groups with finite coefficients are finite. Their first derived limit consequently vanishes. Taking inverse limits therefore gives an injection

$$H^2_{\mathrm{cts}}(G_{\mathbb{Q},S'}, M_3) \longrightarrow \bigoplus_{q \in S' \setminus \{3\}} H^2_{\mathrm{cts}}(G_{\mathbb{Q}_q}, M_3), \tag{A.33.3}$$

and the continuous obstruction is zero. This proves existence of a continuous global class-three lift, without a passage from unrelated finite-level choices.

Every rational Frobenius eigenvalue on $M_3(-1)$ has complex absolute value strictly greater than one. Indeed M_3 is the degree-three multiplication kernel of A.22, and every eigenvalue is a product of three eigenvalues on V; their absolute values are at least $q^{3/2}$ before the twist. Hence $H^0(\mathbb{F}_q, M_3(-1)) = 0$. Degree-one localization and the same finite-S' inverse-limit argument identify the two first-cohomology groups in (A.33.1). The Smith calculation of A.22 proves the displayed decomposition. The target section has trivial geometric centralizer: its image in V is Frobenius-fixed and therefore zero, and its remaining image in $\bigwedge^2 V$ is zero for the same reason. The ordinary central-extension lifting action is consequently a torsor on section classes, with no further stabilizer quotient.

Locally the same weight calculation kills the tame term in $H^1(G_{\mathbb{Q}_q}, M_3)$, so this group is exactly $M_3/(F_q - 1)M_3$. A ramified class-two cusp packet fixes its actual full cuspidal lifting class, by B.7–B.8, and therefore fixes its class-three image. The difference from s_3 is consequently an element of this first-cohomology group. Replacing s_3 by a global central twist adds the restriction of that twist to every local difference. This proves the independence and the asserted equivalence in (A.33.2). A.19 forces every ramified local metabelian restriction to be the actual cusp packet, and B.7 gives the same conclusion for the full group. Their class-three images therefore make every local difference zero, proving the final necessity assertion. □

Statement A.34 (Proved). The restriction map at seventeen and a class-three lift that stops at class four. *Use the actual arithmetic generators m, n, c of A.31. On the degree-three lattice M_3, the restriction map*

$$H^1(G_{\mathbb{Q},\{3,\infty\}}, M_3) \longrightarrow H^1(G_{\mathbb{Q}_{17}}, M_3) = M_3/(F_{17} - 1)M_3 \tag{A.34.1}$$

has image of dimension 15 *after reduction in the local target modulo three. The target modulo three has dimension* 164*, and the cokernel of (A.34.1) modulo three therefore has dimension* 149*.*

Fix free geometric generators $x_0, \dots, x_9$ whose abelianizations are the integral basis of A.7, with the first eight entries the non-O cusp generators. In M_3 put $e = [[x_8, x_0], x_9]$. The exact normalized cocycle

$$\beta(c) = 0, \qquad \beta(m) = (1 + c)e, \qquad \beta(n) = 0 \tag{A.34.2}$$

is unramified outside three. Start with the actual full O-cuspidal section whose tangent parameter is 17, project it to $N/\gamma_4 N$, and twist by β. The resulting global section s_3 has the same class-two image as that full cuspidal section, but s_3 has no lift to $N/\gamma_5 N$. In particular it has no full or metabelian lift, and no lift to the marked quotient with marked cusp Q_c.

Thus a class-two target with a full lift can have a class-three lifting class with a nonzero next obstruction. The theorem does not assert non-lifting of that class-two target: its original full cuspidal lift is specified above.

Proof. All the following matrices act on the reduction $\overline{M}_3$. Write U, V_0, C for the actual matrices of u, v, c in A.22, and put

$$M = UV_0, \qquad L = V_0U^{-1}, \qquad A_0 = M - 1, \qquad B_0 = L - 1, \qquad \Pi_+ = (1+C)/2. \tag{A.34.3}$$

The action over K_0 is abelian by the explicit isogeny and CM comparisons in A.7. Thus these really are the actions of m, n, independently of the chosen c-fixed lift m. The exact matrices give

$$A_0^3 = B_0^3 = A_0^2B_0^2 = 0, \quad A_0B_0 = B_0A_0, \quad CMC = M, \quad CLC = L^{-1}. \tag{A.34.4}$$

In particular the fourth power of the augmentation ideal of this coefficient action is zero modulo three. Normalized cocycles are prescribed freely by

$$a \in \ker(C - 1), \qquad b \in \ker(LC + 1) \tag{A.34.5}$$

as their values at m, n. These subspaces have dimensions $164, 166$. They are reductions of the corresponding integral eigensublattices, since the two operators C and LC are involutions and two is invertible. The exact free presentation with $c(m) = m$ and $c(n) = n^{-1}$ proves that every integral such pair is an actual normalized cocycle.

For any cocycle, evaluation on $D_5(P_3)$ is zero modulo three. Evaluation on $\gamma_5(P_3)$ contains four augmentation operators and is zero by (A.34.4). The coefficient action kills all commutators, so evaluation on cubes of commutators is three times a value and is zero. Evaluation on ninth powers is also zero: the action of every element has order dividing three, and its nine-term norm is three times its three-term norm. These are the factors generating $D_5(P_3)$. Consequently (A.31.3) leaves only the three possible degree-four Hall terms

$$K_1 = [m, [m, [m, n]]], \quad K_2 = [n, [m, [m, n]]], \quad K_3 = [n, [n, [m, n]]].$$

Their cocycle values, and the value of the specified degree-three term, are respectively

$$\begin{array}{c|l} m^6[n, [n, m]] & (2A_0^2 + B_0^2)a - A_0B_0b \\ K_1 & -A_0^2B_0a + A_0^3b \\ K_2 & -A_0B_0^2a + A_0^2B_0b \\ K_3 & -B_0^3a + A_0B_0^2b. \end{array} \tag{A.34.6}$$

These formulas follow by repeated use of $f(gh) = f(g) + gf(h)$; commuting coefficient actions give $f([g, h]) = (1 - h)f(g) + (g - 1)f(h)$.

The action of the normalized rational Frobenius at seventeen is C modulo three. Indeed its square is φ_{17}, and (A.27.1) gives the latter action as $M^{a_{17}}$ with $a_{17} \in 3\mathbb{Z}_3$, whereas $M^3 = 1$

modulo three. The rational Frobenius is exactly $c\varphi_{17}^{1/2}$ in its decomposition group. The local target modulo three is therefore $\overline{M}_3/(C-1)\overline{M}_3 \simeq \overline{M}_3^C$. Its Frobenius value is half the Π_+ projection of the value at φ_{17}.

The program `a_frobenius17_eval_m3.py` in the `code` directory applies the four operators in (A.34.6) to the two eigenspaces (A.34.5), using the full integral basis and matrices in `a_degree3_matrices.json`. All operations here are exact in $\mathbb{F}_3$. After projection by Π_+, the three degree-four operators have ranks $0, 1, 0$. The specified degree-three operator has rank 15. Adding the middle operator with coefficient 0, 1, or 2 gives the same column space, of rank 15 in every case. The saved image basis and the elimination routine certify these comparisons, so unknown higher coefficients of the actual Frobenius cannot change the image. Coboundaries restrict to zero in the local quotient; (A.34.5) supplies all integral normalized cocycles. This proves the rank and cokernel assertions for the actual map (A.34.1).

For (A.34.2), the same calculation has zero contribution from every unknown degree-four term. In the multidegree basis of A.22, e has zero-based index 87, namely $(\{0, 8, 9\}, 8)$. Let z be the reduction modulo three of its local Frobenius value. It is C-fixed. The program records all its 36 nonzero coordinates and forms the actual Lie bracket $[z, e_O]$ in the degree-four free Lie algebra by its injection into the free associative algebra. Here $e_O = -\sum_{j=0}^{7} e_j$ in the fixed abelian basis. For reproducibility, a single nonzero coefficient is

$$[z, e_O]_{X_0X_0X_0X_2} = 1 \quad \text{in } \mathbb{F}_3. \tag{A.34.7}$$

The complete polynomial has 1098 nonzero words. It is saved in `a_frobenius17_eval_m3.json`. The code converts each multidegree difference to its integral Hall bracket $[[x_i, x_j], x_k]$, expands the brackets, and checks directly with the actual matrix on V that

$$C[z, e_O] = -[z, e_O]. \tag{A.34.8}$$

Thus this is a nonzero element of the actual degree-four free Lie lattice modulo three, with its stated sign.

Locally choose the uniformizer Kummer roots for 17 so that the reference cusp section has inertia c_O and Frobenius F_0. The twist (A.34.2) changes no inertia, since it is unramified there, and changes Frobenius to zF_0 modulo $\gamma_4 N$. In a putative lift to $N/\gamma_5 N$, its inertia must be $c_O k$ for some $k \in L_4 := \gamma_4 N/\gamma_5 N$. An arbitrary choice of the lifted Frobenius differs only by a central L_4 value, which has no effect on its conjugation of inertia. The tame relation therefore gives the necessary equation

$$(F_{17} - 17)k = -17[z, e_O] \quad \text{in } L_4. \tag{A.34.9}$$

Modulo three its left side is $(C+1)k$, in the plus eigenspace. By (A.34.7)–(A.34.8) its right side is a nonzero minus eigenvector. This is impossible because two is a unit modulo three. Conjugating a prospective lifting class would not avoid the equation: lift its class-three conjugator through the surjection $N/\gamma_5 N \to N/\gamma_4 N$ to make its projection the prescribed representative, and then apply the same calculation.

This proves the local obstruction and therefore the failure of a global class-four lift. A.19 would make any local metabelian lift of this ramified class an actual cusp section and hence give a full local lift, contradicting (A.34.9). B.14 gives the identical consequence for a marked lift because O is an unmarked cusp. These exact quotient comparisons prove the remaining non-lifting assertions. The original O-cusp section remains a full lift of the shared class-two image, as asserted. □

Statement A.35 (Proved). A fifteen-dimensional toric family with zero first mod-three gluing class. *Let $B \in (\bigwedge^2 W)^{c=-1}$. This eigensublattice has rank fifteen. The*

exact prescriptions

$$\gamma_B(c) = 0, \qquad \gamma_B(m) = 0, \qquad \gamma_B(n) = B \tag{A.35.1}$$

define an unramified-outside-three class-two central cocycle. Twist the class-two image of the global O-cuspidal section with tangent parameter 17 *by γ_B. Its restriction at seventeen is the same actual class-two cusp section, including its Frobenius coordinate. This target has a global class-three lift with unchanged local inertia. For the exact gluing class (A.33.2), its image in the cokernel of (A.34.1) modulo three is zero. No assertion of integral vanishing of (A.33.2) is made here.*

Proof. Over K_0 the toric lattice is scalar cyclotomic. Thus m acts on $\bigwedge^2 W$ by 49^2, whereas n acts as one. Its complex-conjugation signs are $(13, 15)$, from the signs $(3, 5)$ on W. The free presentation of A.31 proves (A.35.1). At the normalized local group, φ_{17} is fixed by c. The cocycle value at this element lies both in $\mathbb{Z}_3 B$ and in the plus eigenspace of c, so is zero. Its value on tame inertia is zero because the cocycle is unramified. Its value on rational Frobenius $c\varphi_{17}^{1/2}$ is also zero. This proves the exact local class-two assertion.

Here is an actual global class-three lift. In A.28 first choose x with $\kappa_{17}(x) = 0$ exactly: right multiplication by a unique power of the tame generator achieves this, and the correction is in the Frattini subgroup when the initial value is zero modulo three. The conjugate y, and the resulting generators m, n, then also have zero Kummer value. Keep the tame section value c_O. Choose $h \in \gamma_2 N/\gamma_4 N$ which is a product of toric commutators with degree-two image $-B/2$. Give m section value one and n section value $h^{-1}(nc)(h)$ relative to the reference cusp section. The latter value obeys the exact involution relation because $(nc)^2 = 1$. Its degree-two image is B. The fixed word W in A.28 has zero degree-two difference from the reference, by the same c-fixed argument used for φ_{17}. Its remaining difference in class three is central, and therefore does not change its conjugation of c_O. The actual defining relation $WtW^{-1} = t^{17^2}$ is satisfied. The universal property of that proved presentation gives the claimed global class-three lift.

We calculate its Frobenius difference modulo three, retaining the actual path origins. For every cusp d, let z_d be its framed Abel–Jacobi point and $h_d = 3z_d$ the torus point of A.9. Choose compatible division points $z_{d,j}$ and toric division points $h_{d,j}$. The compatible torsion points $3z_{d,j} - h_{d,j}$ form $v_d \in V$, and their actual proper image in $T/3$ is $d - O$. Thus the path cocycle has the exact identity

$$3b_{g,d} = \kappa(h_d)(g) + gv_d - v_{gd}. \tag{A.35.2}$$

The toric component of v_d is constrained by the actual torsion point $z_d - h_{d,1}$; it has not been replaced by an arbitrary lift. Altering the chosen z-division points changes v_d by $3V$ and changes b by the corresponding actual coboundary. We may therefore choose its proper component to be the specified integral cusp label $t_d = (a_d, b_d)$ of A.7. For g over K_0 the proper component of (A.35.2) is $(T_g - 1)t_d/3$.

Put

$$H_m = 2I_2, \qquad H_n = (T_n - I_2)/3 = \frac{1}{7}\begin{pmatrix} -4 & -1 \\ 1 & -5 \end{pmatrix}. \tag{A.35.3}$$

These are the proper components of the path operators for m, n, since $T_m = 7I_2$. The upper-right blocks in the actual integral action on V are

$$\Phi_m = 14\Psi, \qquad \Phi_n = -\Psi H_n. \tag{A.35.4}$$

Modulo three all the diagonal K_0 actions on V are one. On a toric commutator $[e_d, e_e]$, its change in degree three is the derivation with cusp values $-[H_g t_d, e_d]$, together with toric

terms depending on the full framed origins. The subsequent augmentation kills every toric term; on a term with one proper T factor it replaces that factor by Φ_g.

More explicitly, use the abelian coefficient group γ_2N/γ_4N, with the action twisted by the reference cusp section. In the submodule generated by toric commutators and their images, three successive augmentation operators are zero modulo three. A first augmentation produces degree-three terms with at most one T factor; the second produces only W terms; the third is zero. The c-normalization used above changes the chosen n-value only by degree-three terms with at most one T factor, so these changes also vanish after two augmentations. The tame augmentation sends a toric commutator to its bracket with e_O, a pure W term, and its product in either order with every other augmentation is zero here.

For noncommuting coefficient operators $A_0 = m - 1$ and $B_0 = n - 1$, the leading crossed-derivation value at $[n, [n, m]]$, when the m-value is zero and the n-value is B, is

$$(A_0B_0 - 2B_0A_0)B. \tag{A.35.5}$$

This follows by differentiating brackets: $D[X, Y] = XD(Y) - YD(X)$, including the action $[X, Y]$ on the source value. All the higher group corrections contain three augmentations and are zero. The term m^6 contributes zero. A.30 proves $W \in D_3(F_3(m, n, t))$ for the actual three-generator word, so there are no degree-two tame-letter terms. Its degree-three terms involving the tame letter have two augmentations and vanish by the identities just proved; a restricted tame cube has zero tame source value. The unknown terms in D_4 have three augmentations and are also zero. Thus (A.31.3) computes the whole Frobenius difference in this calculation.

For each cusp, the proper-path contribution to (A.35.5) is, up to the common cusp-bracket sign,

$$\Phi_m H_n - 2\Phi_n H_m = 14\Psi H_n + 4\Psi H_n = 18\Psi H_n \equiv 0 \pmod 3. \tag{A.35.6}$$

The framed toric terms were killed before this calculation, so no unknown origin term remains. The exact program `a_frobenius17_toric_twists.py` in the `code` directory expands all twenty-eight toric commutators in the free associative algebra, using (A.35.3)–(A.35.4), and verifies (A.35.6). Twenty-seven have nonzero individual summands, but their stated difference is always zero. Hence the chosen class-three lift has zero local Frobenius difference in $M_3/3$. Inertia was unchanged by construction. Its class in the local quotient and therefore the gluing cokernel modulo three is zero, which proves the statement. □

Statement A.38 (Proved). An actual Frobenius basis with completely specified finite class-two action. *Let $F_2 = \langle x, y\rangle$ be the geometric pro-three group of $\mathbb{P}^1 - \{0, 1, \infty\}$, based at the unit tangent at zero, and let*

$$N = \ker(F_2 \longrightarrow (\mathbb{Z}/3)^2), \qquad K = N^3\gamma_3N. \tag{A.38.1}$$

Here N^3 is the closed subgroup generated by cubes. There is an actual Frobenius u at the prime $(7, \zeta - 2)$ of K_0 such that $u, v = cuc$ are a free basis of P_3, their integral action on V is exactly A.7, and their action on N/K is the following fully specified finite action. Complex conjugation is induced by $x \mapsto x^{-1}, y \mapsto y^{-1}$. The restriction of Frobenius u to N/K equals the restriction of the following automorphism of F_2/K:

$$x \longmapsto x^7, \qquad y \longmapsto f_0^{-1}y^7f_0. \tag{A.38.2}$$

In the ordered Schreier basis

$$\begin{gathered}(yxy^{-1}x^{-1},\ y^2xy^{-2}x^{-1},\ y^3,\ xyxy^{-1}x^{-2},\ xy^2xy^{-2}x^{-2},\ xy^3x^{-1},\\ x^3,\ x^2yxy^{-1},\ x^2y^2xy^{-2},\ x^2y^3x^{-2}),\end{gathered} \tag{A.38.3}$$

write N/K as $\mathbb{F}_3^{10} \oplus \bigwedge^2 \mathbb{F}_3^{10}$ with product $(a,z)(b,w) = (a+b, z+w+\frac{1}{2}a \wedge b)$. The word-class representative f_0 defining this finite automorphism has coordinates

$$a_f = (0,0,1,1,2,0,2,0,1,2), \tag{A.38.4}$$

and, in lexicographic order $i<j$, the forty-five central coordinates are

$$\begin{aligned} z_f = (&2,1,2,1,1,2,0,1,2,\ 1,1,0,0,0,2,2,0,1,\\ &1,1,1,1,1,0,1,1,0,0,1,2,0,0,1,0,1,1,\\ &0,0,1,2,0,2,2,0,0). \end{aligned} \tag{A.38.5}$$

Thus (A.38.2)–(A.38.5) specify every coordinate of the images of the ten geometric generators. They specify a representative for the induced finite action; the actual Ihara word can differ by the kernel of its map to that action. The arithmetic basis is obtained from any initial Frobenius basis by at most four explicitly described conjugations below. The underlying Galois action and tangential origin are retained; all cocycles and all arithmetic Frobenius words are transported by the resulting actual source automorphism.

Proof. We first specify the geometric comparison. The map from the Fermat curve is

$$t = -z^3/x^3, \qquad (-z/x)^3 = t, \qquad (-y/x)^3 = 1-t. \tag{A.38.6}$$

It is the connected degree-nine cover classified by $F_2 \to (\mathbb{Z}/3)^2$. At O, the specified parameter is $s = z/x$, so $t = -s^3$. The change from the tangent -1 to the unit tangent is the based pullback $t^{1/3^n} \mapsto -s^{3/3^n}$, using the compatible rational odd roots of -1. It has zero Kummer cocycle. This retains the given tangential arithmetic splitting. The Schreier transversal $x^a y^b$, $0 \le a,b < 3$, gives exactly (A.38.3). It also gives the actual conjugation actions of x and y on N/K; using these actions and the stated class-two product is a finite word-collection algorithm for (A.38.2).

For the chosen tangential paths the actual Ihara action is $x \mapsto x^7$, $y \mapsto f^{-1}y^7 f$, with $f \in F_2'$. Its symmetry, hexagon and pentagon are the actual Galois relations [Sch05, §§I.3, I.6]; we use the hexagon order

$$f(x,y)x^3 f(z,x)z^3 f(y,z)y^3 = 1, \qquad z = (xy)^{-1}. \tag{A.38.7}$$

The comparison with A.7 is the integral unimodular matrix saved by `a_ihara_lattice.py`. It is obtained from the actual order-nine deck translations, and intertwines both complex conjugation and the integral Frobenius homology matrix. Consequently the finite equations below constrain the actual arithmetic action in the specified geometric basis.

We give the finite elimination with its source-lifting conditions. The exact collector for (A.38.3) works in F_2/K, of order 3^{57}. Its intersection with the lower central and Zassenhaus filtrations is computed by taking literal group commutators, cubes, and their invariant spans. Through degree seven the kernel of the comparison from the free source has one degree-six generator, $[x,y]^3$, and four degree-seven generators. The literal words for all five are included in the certificate and are individually checked to map to one in F_2/K. No source direction below degree six is lost. The exact commutator calculation is separately verified by the full class-two multiplication law, retaining its quadratic conjugation terms.

The homology equation, $f \in F_2'$, symmetry, and (A.38.7) leave three possible homology rows and an affine central system over each row. Requiring a lift of these same equations over $\mathbb{Z}/9$ retains all next-digit variables. We then impose the full group pentagon through source degree seven, including the five actual source-kernel words just specified. For precision, the

pentagon is evaluated in the fourth-strand free group of $PB_4 = F_3 \rtimes PB_3$. The Artin maps are

$$r_1 : (X, Y, Z) \mapsto (Y, Y^{-1}XY, Z), \qquad r_2 : (X, Y, Z) \mapsto (X, Z, Z^{-1}YZ). \tag{A.38.8}$$

Put $A_{12} = r_1^2$, $A_{23} = r_2^2$, $A_{13} = r_2 r_1^2 r_2^{-1}$, and $A_{14} = \mathrm{Inn}(X)$, $A_{24} = \mathrm{Inn}(Y)$, $A_{34} = \mathrm{Inn}(Z)$. These are the actual pure Artin maps [CFR12, (2.1)–(2.3)]; direct free-word reduction verifies every relation and $A_{12}A_{13}A_{23} = \mathrm{Inn}((XYZ)^{-1})$. The five factors have pairs

$$\begin{aligned} &(A_{12}, A_{23}A_{24}), \quad (A_{13}A_{23}, A_{34}), \quad (A_{23}, A_{34}), \\ &(A_{12}A_{13}, A_{24}A_{34}), \quad (A_{12}, A_{23}). \end{aligned} \tag{A.38.9}$$

If they are denoted $F_0, \ldots, F_4$, the residual is $F_0F_1F_4^{-1}F_3^{-1}F_2^{-1}$. Forgetting the fourth strand sends this residual to one, so it is an element of the actual free fiber and its displayed action is inner.

All evaluations use the denominator-free Magnus substitution $X_i \mapsto 1 + X_i$ in the associative algebra over $\mathbb{F}_3$, truncated after degree eight. If $\Delta_A = \Phi_A - 1$, then the source word $1 + \sum_w c_w w$ acts as $1 + \sum_w c_w \Delta_{w_1} \cdots \Delta_{w_k}$. Each Δ_A raises augmentation degree. Terms of source degree at least eight therefore do not affect the target generators through degree eight. Conversely, the first nonzero degree-d term h_d of the fiber residual gives $[h_d, X_i]$ in degree $d+1$ of its inner action. A positive-degree tensor commuting with all three generators is zero. Hence the computation detects precisely all source residuals through degree seven. This uses the actual free fiber and the displayed maps throughout.

The degree-two source coefficient must also be retained integrally. Write $f = [x, y]^{f_2}$ modulo $\gamma_3 F_2$. The integral Heisenberg collector applied to the hexagon with $m = (\chi - 1)/2$ gives

$$3f_2 = \frac{m(m+1)}{2}. \tag{A.38.10}$$

This identity is in the torsion-free group $\mathbb{Z}_3$; at $\chi = 7$ it gives $f_2 = 2$ before any reduction. Its reduction modulo nine is well-defined on the source used above, since the Heisenberg group over $\mathbb{Z}/9$ kills $D_8(F_2)$. The five source-kernel words have degree-two coefficients $(3, 0, 0, 0, 0)$, so this final scalar row is imposed on the combined source word, including those kernel corrections.

Here is the complete finite elimination result. "Invisible" means the kernel of the explicitly computed map to the source metabelian quotient, followed by the intrinsic Johnson map.

equations imposed	$\operatorname{rank} J$	$\operatorname{rank}(J, H)$	$\operatorname{rank} J\vert_{\ker H}$
mod-nine symmetry and hexagon	3	5	2
also full degree-seven pentagon	3	4	1
also integral f_2 mod 9	2	3	0

(A.38.11)

The source metabelian map in this table is the literal Fox map: write $f = [x, y]^h$ in F_2'/F_2'', with $X = x - 1$, $Y = y - 1$. In the finite comparison its coefficient ring is

$$\mathbb{F}_3[X, Y]/(X^6, X^3Y^3, Y^6, X^5Y^2, X^2Y^5). \tag{A.38.12}$$

It is computed by $D_y(f) = Xh$. Thus every map in the last column of (A.38.11) is an actual quotient map of the same based free group.

The row reductions give a more precise conclusion. Let J_0 be the 45×10 central matrix obtained from (A.38.4)–(A.38.5). There are two specified matrices L_1, L_2 such that every actual Frobenius in the initial basis has

$$J = J_0 + aL_1 + bL_2, \qquad a = h_{03} - 2, \quad b = h_{05} \quad \text{in } \mathbb{F}_3. \tag{A.38.13}$$

Here h_{ij} is the coefficient of X^iY^j in (A.38.12). The matrices L_1, L_2, of respective ranks seven and one, are saved with all 450 entries each. The nine matrices in (A.38.13) are distinct. Formula (A.38.13) is at this point a necessary arithmetic constraint; the realization follows below.

For reproducibility, the complete pentagon matrix has 29,520 coefficient rows. Its six candidate columns and five source-kernel columns have rank seven, and the five kernel columns have rank five. Each of the three branches has coefficient and augmented rank seven on its 24 retained parameters. The additional scalar row (A.38.10) has rank one in each branch. The supplied particular solutions and full nullspaces verify every coefficient row. The original and degree-pruned pentagon evaluations give identical matrices; the independent integral Heisenberg audit verifies (A.38.10) and every filtered solution against all 29,520 rows. The files in the `code` directory with prefixes `a_ihara_pentagon`, `a_ihara_source_scalar`, `a_ihara_final_parameters`, `de_fermat_pentagon`, and `de_fermat_scalar_audit` retain the exact words, matrices, row reductions, and executed outputs. These finite identities establish (A.38.11)–(A.38.13), including the affine origin.

We now realize the stated value by an arithmetic morphism. For a current actual basis $u, v = cuc$, two operations are

g	$u' = gug^{-1}, \quad v' = cu'c$	(a', b')
v	$vuv^{-1}, \quad c(vuv^{-1})c$	$(a-1, b+a-1)$
$[u,v]$	$[u,v]u[u,v]^{-1}, \quad c([u,v]u[u,v]^{-1})c$	$(a, b+1)$.

(A.38.14)

These are direct identities of the full 55×55 matrices $\left(\begin{smallmatrix} U & 0 \\ J & \bigwedge^2 U \end{smallmatrix}\right)$ and the fixed complex-conjugation matrix. The program `a_ihara_frobenius_conjugacy.py` verifies all nine inputs of both operations. Applying the first operation a times, using the representatives $a \in \{0, 1, 2\}$, changes the pair to $(0, b + a(a-1)/2)$. Applying the second operation the negative of that last second coordinate times gives $(0, 0)$. At most four operations are needed. Every step conjugates an actual Frobenius, so the final element is an actual Frobenius at the same prime of K_0.

The assignments $u \mapsto u'$, $v \mapsto cu'c$ preserve the two independent Frattini classes and commute with c. They therefore define a surjective endomorphism of the free pro-three group P_3. A finitely generated profinite group is Hopfian: its finitely many open normal subgroups of each fixed index are permuted by inverse image under a surjection, forcing its kernel into every open normal subgroup. Thus the endomorphism is an actual c-equivariant automorphism ψ.

The full integral homology action is also retained. The splitting after pullback by [3] in A.7 is unique, hence G_{K_0}-equivariant. Its isogeny $T_D \times E \to J_{E,D}$ induces the rational equivariant identification with $W_{\mathbb{Q}_3} \oplus T_{\mathbb{Q}_3}$. The embedding of the chosen integral graph coordinates into this diagonal representation is

$$\iota(w, t) = (w + \Psi t/3, t). \tag{A.38.15}$$

The actual isogeny on Tate modules is $j(w, t) = (w, 3t)$ in the rational diagonal coordinates, or $\iota^{-1} j(w, t) = (w - \Psi t, 3t)$ in graph coordinates. Thus the equivariant comparison retains its multiplication by three on the proper factor. On this rational direct sum every $g \in G_{K_0}$ acts by $(\chi(g) I_8, T_g)$, where T_g is multiplication in the commutative algebra $\mathbb{Q}_3[\zeta]$. These actions commute. The integral lattice embeds in this rational representation, so their integral actions commute as well. Arithmetic conjugation consequently preserves the exact A.7 matrix.

Finally we give the transport of every parameter used later. For the original action ρ, the new action is $\rho \circ \psi$. The exact semidirect-product isomorphism is $(N/K) \rtimes_{\rho\circ\psi} \Gamma \to (N/K) \rtimes_{\rho} \Gamma$, $(h, g) \mapsto (h, \psi(g))$, where $\psi(c) = c$; its inverse uses ψ^{-1}. A cocycle is pulled back by $z \mapsto z \circ \psi$.

In particular,

$$z(gug^{-1}) = z(g)\,\rho(g)(z(u))\,\rho(gug^{-1})(z(g)^{-1}), \tag{A.38.16}$$

with the additive version $z(g)+\rho(g)z(u)-\rho(gug^{-1})z(g)$ for an abelian module. This formula retains the reference section and its origin. In the free presentation of A.28, make the same substitutions on x, y and fix t. They again define a c-equivariant IA automorphism, and the actual new Frobenius word is

$$W_{q,\mathrm{new}} = \psi^{-1}(W_{q,\mathrm{old}}). \tag{A.38.17}$$

Indeed the old quotient map composed with ψ sends this word to the same local Frobenius and sends the transformed relator to $W_{q,\mathrm{old}} t W_{q,\mathrm{old}}^{-1} t^{-q^2}$. The definitions of m, n by the pro-three square-root operations commute with this actual automorphism. An IA automorphism of a free pro-three group acts identically on its associated restricted Lie algebra. Since A.30 proves $W_{17} \in D_3$, the initial terms in A.31 and A.37 survive unchanged modulo D_4.

For completeness this also preserves the ordinary weight-three coefficients modulo three used to identify those terms. The degree-one and degree-two Hall coefficients of W_{17} are divisible by three. Under a basis-conjugating automorphism, the changes to a degree-one power with exponent $3a$ have weight-two and weight-three coefficients divisible by $3a$ or $\binom{3a}{2}$; both are divisible by three. A weight-two power with exponent $3b$ changes by weight-three multiples of $3b$. The automorphism is the identity on the third lower central graded quotient. Hence the stated ordinary coefficients are unchanged as well. All higher terms, and the integral lower coordinates when needed, are transported by (A.38.17). This proves the realization and every specified comparison. □

Statement A.39 (Proved). Global exclusions at seventeen inside the exact local cusp packet. *Use the actual arithmetic basis of A.38, with m, n and the normalized local Frobenius $\varphi = \varphi_{17}$ transported as there. Put $M_2 = \bigwedge^2 V$ and define the integral normalized cochain module and its actual evaluation map by*

$$\begin{aligned} D &= M_2^{c=1} \oplus M_2^{nc=-1}, \\ E_\varphi : D &\longrightarrow M_2^{c=1}, \quad (a,b) \longmapsto \gamma_{a,b}(\varphi), \\ R &= \ker E_\varphi. \end{aligned} \tag{A.39.1}$$

Here $\gamma_{a,b}$ is the unique continuous cocycle on $P_3 \rtimes \langle c\rangle$ with values $(a,b,0)$ at (m,n,c). The lattice D has rank 45, and R is a saturated lattice of rank 24.

For $r \in R$, twist the class-two image of the global O-cuspidal section with tangent parameter 17 by γ_r. This defines an injectively parameterized family of global class-two section classes, unramified outside $\{3, 17, \infty\}$. Every member has the same exact local O-cusp section at seventeen, including its Frobenius coordinate, and every member has a global class-three lift.

There is an $\mathbb{F}_3$-linear obstruction on $R/3R$ of rank at least three which vanishes on every member lifting to any of

$$N, \qquad N/N'', \qquad N/\gamma_3(I_{Q_c}). \tag{A.39.2}$$

Consequently, from any integral basis $r_1, \ldots, r_{24}$ of the kernel in (A.39.1), at least three of the resulting twenty-four class-two section classes have none of the lifts in (A.39.2). This is an unconditional finite-family existence certificate. It does not select an unevaluated individual basis vector as a counterexample. No assertion of full local liftability at three is included. An individually specified member of R with nonzero obstruction, including all integral generator coordinates and a nine-term detecting functional, is given in B.20.

Proof. We first prove the integral assertions, before using any finite computation. The relations $c(m) = m$, $c(n) = n^{-1}$ give precisely the two conditions in (A.39.1). Since m, n

are a free pro-three basis, they are sufficient. The signs of c on V are $(4,6)$, so its signs on M_2 are $(21,24)$. The involution nc is conjugate to c by a pro-three square root of n, and therefore has the same signs. Its integral eigenspaces are direct summands, since two is a unit in $\mathbb{Z}_3$. Thus D is free of rank 45.

Since φ is c-fixed, its cocycle value belongs to $M_2^{c=1}$. For h in that invariant lattice, the global coboundary $g \mapsto (g-1)h$ belongs to D and evaluates to $(F_\varphi - 1)h$. The rational eigenvalues of F_φ on M_2 are

$$17^4, \qquad -17^3, \qquad 17^2. \tag{A.39.3}$$

Indeed on the rational equivariant splitting of A.7 the toric factor has eigenvalue 17^2 and the proper factor has scalar eigenvalue -17. None of (A.39.3) is one. The map $F_\varphi - 1$ is consequently invertible over $\mathbb{Q}_3$ on the 21-dimensional c-fixed target. Evaluation has rational rank 21. Its kernel R has rank 24, and $D/R \simeq \operatorname{im} E_\varphi$ embeds in a free $\mathbb{Z}_3$-module. Thus R is saturated and

$$R/3R \hookrightarrow D/3D \tag{A.39.4}$$

has dimension 24.

For $r \in R$ the unramified-outside-three cocycle has zero value at c and φ and zero tame inertia value. Its restriction to the local group at seventeen is therefore zero. This proves the exact local class-two assertion. The family is injective on global section classes as well. A geometric conjugator between two such class-two sections has invariant image in V, which is zero by Frobenius weights. It therefore lies in M_2. Normalization at c makes it c-fixed. If the difference of two members of R were its coboundary, evaluation at φ would give $(F_\varphi - 1)h = 0$, hence $h = 0$ by (A.39.3).

Here is a definite integral class-three lift of each member. Write

$$L = \gamma_2 N/\gamma_4 N, \qquad 0 \longrightarrow M_3 \longrightarrow L \longrightarrow M_2 \longrightarrow 0. \tag{A.39.5}$$

This is an exact sequence of free $\mathbb{Z}_3$-modules. Lift the two values of r equivariantly for the involutions c and nc, using their averaging idempotents. In the free presentation of A.28 take these as the L-valued cochain values at m, n, keep the reference tame section value c_O, and keep the value at c equal to one. The resulting crossed cochain β has $\beta(t) = 0$. Its value at the actual word W projects to $E_\varphi(r) = 0$, so $\beta(W) \in M_3$. That value is central in $N/\gamma_4 N$ and commutes with the unchanged tame section. The relation $WtW^{-1} = t^{289}$ is therefore satisfied exactly. Equivalently, its crossed value is $(1 - T^{289})\beta(W) = 0$, where T is the reference tame action on L and is the identity on M_3. The proved one-relator presentation supplies the global class-three lift.

We describe the finite comparison computing its remaining local obstruction. It uses the actual action of A.38 and does not require degree-three images of the geometric generators. Let X_i be their logarithms in the rational class-three nilpotent Lie algebra. The logarithm maps L onto the integral lattice with degree-two basis $[X_i, X_j]$ and degree-three Hall basis $[[X_i, X_j], X_k]$, with the usual Jacobi identifications. To check integrality, the sole correction needed is

$$\log[x_i, x_j] = [X_i, X_j] + \tfrac{1}{2}[X_i + X_j, [X_i, X_j]], \tag{A.39.6}$$

and two is invertible in $\mathbb{Z}_3$. Thus this is an actual lattice comparison. If a class-two automorphism has homology matrix A and central columns J_i, its action on $\overline{L} = L/3L$ is

$$\widetilde{G} = \begin{pmatrix} \bigwedge^2 A & 0 \\ \Theta(A,J) & \operatorname{Lie}_3(A) \end{pmatrix}, \quad \Theta(A,J)_{ij} = [J_i, AX_j] + [AX_i, J_j]. \tag{A.39.7}$$

Unknown degree-three terms in the images of X_i enter these brackets only in degree four and hence disappear in (A.39.5). Formula (A.39.7) is therefore determined by A.38.

For the O reference section, e_O is the seventh Schreier generator of A.38 and

$$T(z,w) = (z, w + [e_O, z]). \tag{A.39.8}$$

The source generators m, n have zero 17-Kummer value by A.35 and its transported construction. Their actions on L are consequently the original arithmetic actions. The program `a_ihara_gamma2_action.py` constructs (A.39.7), verifies composition and every involution and tame relation, and compares the entire degree-three block with the independent integral lattice of A.13. For the augmentation ideal generated by $m-1, n-1, t-1$ on $\overline{L}$, the dimensions of successive images are

$$375,\ 249,\ 132,\ 47,\ 4,\ 0. \tag{A.39.9}$$

In particular its fifth power is zero. A crossed cochain therefore kills D_6 of the arithmetic free group: the derivation of the sixth augmentation power lies in the fifth power applied to the module.

Every source term that can affect the answer is retained. The group D_3/D_6 is abelian of exponent three. On it the map

$$\mathrm{av}(w) = (w\, c(w))^2 \tag{A.39.10}$$

is exactly the c-fixed averaging idempotent. A.31 and A.37 give the known cubic part $m^6[n,[n,m]][t,[t,m]]^2$. The two unknown cubic terms have Lyndon words $012, 021$ in the order (m,n,t). All eighteen degree-four and all forty-eight degree-five Lyndon terms are then retained, with (A.39.10) applied to each. These exhaust the source freedom: the only restricted-power terms in these degrees are the degree-three generator cubes, and their coefficients are the already fixed abelian Frobenius coordinates. The source comparisons of A.38 ensure that these are the actual transported coefficients of W.

The local mod-three M_3 target is its c-fixed space of dimension 164. Divide by the actual rank-fifteen global restriction image in A.34; denote the resulting linear projection by Q, with 149-dimensional target. The factor two from norm Frobenius to rational Frobenius is a unit and does not affect any of the following ranks. The canonical integral lifts in (A.39.5) give an ambient linear map on $D/3D$. Evaluating every term just listed and applying Q shows that only the following three unknown terms survive:

$$[m,[n,t]], \qquad [[m,t],n], \qquad [m,[[m,n],n]]. \tag{A.39.11}$$

Every degree-five contribution vanishes. We retain the three actual coefficients as arbitrary members of $\mathbb{F}_3$; the rank bound below holds for all twenty-seven triples.

All local conjugators and global class-three corrections are also retained. The image of a local conjugator in V is invariant, hence zero by weights. Its degree-two part belongs to $M_2^{c=1}$ after normalization at c. Choose any of its 21-dimensional equivariant lifts h to $\overline{L}$. Allow the enlarged finite comparison in which

$$\beta(t) = (T-1)h, \qquad Q\big(\beta(W) - (F_\varphi - 1)h\big) \tag{A.39.12}$$

is evaluated, while the values at m, n are the chosen cochain lifts. This includes every reduction of a local comparison coming from a putative lift. Degree-three changes of h are already killed by the local coboundary quotient. Allowing every degree-two h only enlarges the set of finite corrections, so a nonzero obstruction after this allowance still excludes an integral lift.

Write the map (A.39.12), for its actual coefficient triple ϵ, as $G_\epsilon r + H_\epsilon h$. The complete finite calculation is

$\epsilon \in \mathbb{F}_3^3$	$\operatorname{rank} G_\epsilon$	$\operatorname{rank} H_\epsilon$	$\operatorname{rank}(G_\epsilon, H_\epsilon)$	$\operatorname{rank}(G_\epsilon \bmod \operatorname{im} H_\epsilon)$
each of the twenty-seven triples	30	6	30	24.

(A.39.13)

Here G_ϵ has 45 columns and H_ϵ has 21 columns. The images of all H_ϵ are the same six-dimensional subspace, so the last column can be computed in one fixed 143-dimensional target. For a reproducible specification of all their entries, take the ascending pivot columns of $1+c$ and $1-nc$ on $M_2/3$ for the two source bases, lift them by the same operators on $L/3$, and evaluate the words above using

$$\begin{aligned} \beta(ab) &= \beta(a) + a\beta(b), \\ \beta([a,b]) &= (1 - aba^{-1})\beta(a) + (a - [a,b])\beta(b). \end{aligned} \tag{A.39.14}$$

Use the literal Jacobi reductions in the basis of A.13 and the projection Q just defined. These formulas and A.38 specify every entry without an arithmetic oracle. The program `a_frobenius17_mixed_gluing.py`, run with `-local-conjugator`, saves the full matrices, all twenty-seven coefficient cases, and their exact elimination ranks. Its certificate also saves each of the sixty-eight candidate word maps before discarding the zero ones. In particular (A.39.13) retains the two mixed inertia terms and the quartic term, rather than assigning their values. The independent program `bc_gluing_audit.py` instead uses truncated noncommutative Magnus polynomials and the derivation formula $\beta((g_1-1)\cdots(g_d-1)) = (g_1-1)\cdots(g_{d-1}-1)\beta(g_d)$. It reproduces every entry of all sixty-eight word maps, the common six-dimensional image, and all twenty-seven rank computations.

Any two global class-three lifts of the fixed class-two target differ by an M_3 cocycle. Such a cocycle is unramified outside three: at every other finite prime the possible tame value lies in $H^0(\mathbb{F}_q, M_3(-1))$, which is zero by weights. Its effect is therefore in the actual global restriction image already removed by Q; this is also the localization comparison in A.33. Thus vanishing of $G_\epsilon r$ modulo $\operatorname{im} H_\epsilon$ is a necessary condition for a global class-three lift whose restriction at seventeen is the actual cusp section.

For the actual, unassigned triple ϵ, (A.39.13) says that the kernel of this ambient map on the 45-dimensional space $D/3D$ has dimension 21. The embedded subspace $R/3R$ in (A.39.4) has dimension 24. Its restriction consequently has rank at least $24-21=3$. This is the asserted obstruction. It is obtained from actual integral cochains in R; no mod-three local condition has been substituted for the exact kernel in (A.39.1).

Finally every lift in (A.39.2) would supply the forbidden locally cuspidal class-three lift. The full group maps to $N/\gamma_4 N$. The metabelian quotient does also, because $N'' \subset \gamma_4 N$, and its local section is the actual O cusp packet by A.19. For the marked quotient, the image of I_{Q_c} in V is the single line generated by e_{Q_c}. Hence $[I_{Q_c}, I_{Q_c}] \subset \gamma_3 N$ and $\gamma_3(I_{Q_c}) \subset \gamma_4 N$, proving the actual map to the same class-three quotient. The marked cusp is Q_c, whereas the ramified cusp here is O; B.14 identifies its local marked section with the unique actual O cusp section. The local tame exponent is one, and its unramified cusp parameter is zero because $17-1$ is a three-adic unit. Thus all three types of lift are excluded whenever the obstruction is nonzero. A basis of R spans its reduction, so an obstruction of rank at least three is nonzero on at least three basis members. This proves the finite-family assertion and the stated global exclusions. □

Statement A.40 (Proved). The excluded family also fails at the next nilpotent stage. *Every member of the family in A.39 with nonzero stated obstruction has no global lift to $N/\gamma_5 N$. Thus the twenty-four-member finite family there contains at least three class-two sections which lift globally to $N/\gamma_4 N$ and have no global lift to $N/\gamma_5 N$. The explicitly evaluated section in B.20 is one such member.*

Proof. First conjugate a proposed lift, through the actual surjection to $N/\gamma_3 N$, so that its class-two projection is the specified cocycle. The coprime involution argument in the

kernel then normalizes its value at c to one. Every global class-three lift differs from the lift constructed in A.39 by an M_3 cocycle. A.39 proves that this difference is unramified outside three, even when additional ramification was initially allowed. Hence its actual tame inertia at seventeen is still exactly c_O in $N/\gamma_4 N$.

Write F_0 for the reference rational Frobenius and F' for the Frobenius of the proposed lift. In class three their difference is an element $z \in M_3$. The relative local group has the exact relation

$$\mathrm{Frob}_{17} = c\,\varphi_{17}^{1/2}. \tag{A.40.1}$$

Its involution commutes with φ_{17}, so normalization at c makes z c-fixed. The integral A.7 matrix gives $\varphi_{17} = 1$ on $V/3$. Thus its square root is also one on every graded quotient modulo three, and the cocycle norm operator $1 + \rho(\varphi_{17}^{1/2})$ is $2I$ modulo three. The rational-Frobenius difference z is consequently one half of the norm-Frobenius difference modulo three. Nonzero obstruction in A.39 implies

$$\bar{z} \neq 0, \qquad C\bar{z} = \bar{z} \quad \text{in } M_3/3M_3, \tag{A.40.2}$$

after every permitted global class-three correction.

Let $M_4 = \gamma_4 N/\gamma_5 N$. In class four write the inertia and Frobenius of the hypothetical lift in the exact order

$$\tau' = c_O a_4, \qquad F' = z b_4 F_0, \qquad a_4, b_4 \in M_4. \tag{A.40.3}$$

A choice of lift of z from M_3 only changes b_4. The reference cusp relation is $F_0 c_O F_0^{-1} = c_O^{17}$. The group M_4 is geometrically central at this stage, and collecting the relation $F'\tau' F'^{-1} = (\tau')^{17}$ gives

$$(F_{17} - 17)a_4 = -17[z, e_O] \quad \text{in } M_4. \tag{A.40.4}$$

Indeed conjugating c_O^{17} by the chosen lift of z contributes $17[z, e_O]$; b_4 commutes with inertia, while the two contributions from a_4 are $F_{17}a_4$ and $17a_4$. This derives the sign in the stated ordering and retains the actual reference relation.

On $M_4/3M_4$ the action of rational Frobenius is C by (A.40.1). Since $17 \equiv -1 \pmod 3$, equation (A.40.4) becomes

$$(C + 1)\bar{a}_4 = [\bar{z}, e_O]. \tag{A.40.5}$$

The left side is c-invariant. The right side is c-anti-invariant, because $C\bar{z} = \bar{z}$ and $Ce_O = -e_O$. It remains to check its nonvanishing in the actual ordinary lower-central graded quotient.

The ordinary free Lie algebra on $V/3$ embeds in its tensor algebra: the Hall–Lyndon expansions have distinct unit leading words, giving an integral comparison before reduction. The element e_O is a primitive basis letter. A homogeneous tensor P of degree three commuting with that letter must be a scalar multiple of e_O^3. To see this directly, the equality $e_O P = P e_O$ forces every word of P to start with e_O; cancel that initial letter and repeat. The ordinary degree-three free Lie subspace has zero component in the multidegree consisting of that letter alone, since every bracket on one letter vanishes. Thus

$$[\bar{z}, e_O] = 0 \text{ in } M_4/3M_4 \quad \Longrightarrow \quad \bar{z} = 0. \tag{A.40.6}$$

By (A.40.2) the right side of (A.40.5) is nonzero. The plus and minus eigenspaces of an involution over $\mathbb{F}_3$ intersect trivially, so (A.40.5) is impossible. This excludes every global class-four lift of that class-two target, while A.39 already supplies its global class-three lift. □

Statement A.41 (Proved). The exact completed Fox image and fibers on the cusp family. *For the family R in A.39 put*

$$Q = N/N'', \qquad M = N'/N'', \qquad \Lambda = \mathbb{Z}_3[[V]], \qquad \mathfrak{a} = \ker(\Lambda \to \mathbb{Z}_3), \qquad J = \mathfrak{a}M. \tag{A.41.1}$$

Give M the arithmetic action twisted by the actual reference O section with tangent parameter 17. *Thus m, n, c have their original actions, and tame inertia acts by conjugation by c_O. Define*

$$\begin{aligned} D_M &= M^{c=1} \oplus M^{nc=-1}, \qquad D_J = J^{c=1} \oplus J^{nc=-1}, \\ E_W : D_M &\longrightarrow M^{c=1}, \qquad E_W(a,b) = \beta_{a,b}(W), \end{aligned} \tag{A.41.2}$$

where $\beta_{a,b}$ is the continuous crossed cochain on the free arithmetic source with values $(a, b, 0, 0)$ at (m, n, t, c). These are continuous linear maps of compact $\mathbb{Z}_3$-modules. The natural quotient $M \to M_2$ gives $\pi : D_M \twoheadrightarrow D$, with kernel D_J, and the complete metabelian image on this family is exactly

$$R_{\mathrm{met}} = \pi(\ker E_W) \subset R. \tag{A.41.3}$$

Equivalently, it is the kernel of the explicit continuous map

$$\begin{aligned} \mathfrak{o}_M : R &\longrightarrow J^{c=1}/E_W(D_J), \\ r &\longmapsto [\, E_W(\widetilde{r}\,)\,], \end{aligned} \tag{A.41.4}$$

where $\pi(\widetilde{r}) = r$. Every nonempty fiber of $\mathrm{Sec}(Q) \to \mathrm{Sec}(N/\gamma_3 N)$ over this family is an affine copy of

$$\ker(E_W : D_J \longrightarrow J^{c=1}). \tag{A.41.5}$$

There is no remaining conjugacy quotient after the actual local normalization in the proof. Formulas (A.41.2)–(A.41.5) retain the completed arithmetic Fox operator. They do not assign a numerical value to an unevaluated coefficient of W or of its action.

Proof. We construct the completed module and its maps. Let $x_1, \dots, x_{10}$ be the actual geometric Schreier basis in A.38, and denote their images in V by T_i. The completed Fox map gives the exact sequence

$$0 \longrightarrow M \longrightarrow \Lambda^{10} \xrightarrow{(T_1-1,\dots,T_{10}-1)} \mathfrak{a} \longrightarrow 0. \tag{A.41.6}$$

One direct proof uses the actual finite abelian covers with geometric subgroups $H_k = N' N^{3^k}$. Their based cellular chain maps send the ith lifted edge to $T_i - 1$. The kernel is H_k^{ab}. Inverse limits of these compact chain groups are exact, and the descending-subgroup comparison proved in A.19 identifies $\varprojlim H_k^{\mathrm{ab}}$ with N'/N''. This proves (A.41.6) and its based comparison. The action of geometric conjugation by x_i on M is multiplication by T_i. Consequently

$$J = \mathfrak{a}M = \gamma_3 N/N'', \qquad M/J = N'/\gamma_3 N = \bigwedge^2 V = M_2. \tag{A.41.7}$$

These equalities identify the quotient map used in the statement, including its integral lattice.

The algebra Λ is $\mathbb{Z}_3[[T_1 - 1, \dots, T_{10} - 1]]$, an integral domain. In (A.41.6) the reference tame action T is multiplication by the actual deck element T_O, the image of c_O. Therefore

$$1 - T^{289} : M \longrightarrow M \quad \text{is injective.} \tag{A.41.8}$$

Indeed it is multiplication by the nonzero power series $1 - T_O^{289}$ on a submodule of the torsion-free module Λ^{10}. This also proves that the tame fixed subgroup of M is zero.

The arithmetic action on M is semilinear over its actual action on Λ: $g(\lambda z) = g(\lambda) g(z)$. Its matrices are the Fox matrices of the actual based geometric automorphisms. Thus the continuous group-ring homomorphism from the free arithmetic source to $\mathrm{End}_{\mathbb{Z}_3}(M)$ and the completed Fox derivatives specify (A.41.2) explicitly as

$$E_W(a,b) = \rho\left(\frac{\partial W}{\partial m}\right) a + \rho\left(\frac{\partial W}{\partial n}\right) b. \tag{A.41.9}$$

The derivative with respect to t has zero cochain value. Here ρ is the reference twisted action just defined; neither its semilinear part nor the inertia dependence of W is removed. Since the word W is c-fixed and the cochain is normalized at c, its value is c-fixed. This proves the target in (A.41.2).

Averaging for the order-two actions lifts both eigenspaces through the surjection $M \to M_2$. Hence $\pi : D_M \to D$ is onto with kernel D_J. Reduction of (A.41.9) is exactly E_φ, because the tame action on M_2 is trivial and killing t sends W to the same actual φ. For $r \in R$, $E_W(\widetilde{r})$ therefore lies in $J^{c=1}$. Two choices of $\widetilde{r}$ differ by D_J, proving (A.41.4) is well-defined. The domains are compact and all maps are continuous, so $E_W(D_J)$ is closed. Choosing equivariant lifts of a finite basis of D makes (A.41.4) an explicitly continuous linear map.

We next prove that these kernels classify the actual section classes. A cocycle β with the values in (A.41.2), relative to the full reference cusp section, is a cocycle on the free arithmetic source. Its value on the relator is

$$\beta(WtW^{-1}t^{-289}) = (1 - T^{289})\beta(W). \tag{A.41.10}$$

By (A.41.8) it descends to the proved arithmetic presentation exactly when $E_W(a, b) = 0$. Its values at t, c, W then vanish, so its local section at seventeen is the literal reference section. Its class-two image lies in R by the preceding comparison. This constructs actual global metabelian sections from every member of $\ker E_W$.

Conversely, take any global metabelian section whose class-two class is the target indexed by $r \in R$. Lift a global class-two conjugator through $Q \to N/\gamma_3 N$ so that its projection equals the exact target cocycle; then normalize the value at c in the kernel. At every prime outside three and seventeen its abelian inertia is zero. A.19 implies that its entire metabelian inertia there is zero. Thus it factors through the arithmetic group of A.28 even if arbitrary additional ramification was initially allowed.

At seventeen A.19 places it in the actual O packet. The tame exponent is one, and the unramified cusp parameter vanishes since $17 - 1$ is a three-adic unit. Hence a local geometric conjugator identifies it with the reference cusp section. Its image in $N/\gamma_3 N$ centralizes that local section. This centralizer is trivial: its image in V is Frobenius-invariant and therefore zero; its remaining image in M_2 is likewise zero by (A.39.3). Consequently the conjugator lies in J. The normalization at c makes it c-fixed. Conjugating globally by this same element preserves the exact class-two target and makes the whole local section equal to the reference.

The resulting difference from the reference is an M-valued cocycle, because its abelian projection is the same. Its values at t, c, W are zero, so it gives a member of $\ker E_W$ reducing to r. The normalization is unique: the ratio of two such conjugators is tame-invariant in M, and (A.41.8) makes that ratio zero. More generally, a global conjugator between two normalized representatives has invariant image in V, hence lies in M, and its local inertia commutation again forces it to be zero. This proves the claimed bijection with section classes, including the absence of a remaining quotient.

The image formula (A.41.3), the obstruction formula (A.41.4), and the affine fiber (A.41.5) now follow from the exact sequence of these actual linear maps. In particular R_{met} is a closed $\mathbb{Z}_3$-submodule of the finite free lattice R. Its restriction has the explicit nonzero obstruction of A.39; its integral image has not been replaced by that first finite obstruction.

There is also an exact finite-quotient interpretation. Use the cofinal, arithmetic-stable quotients

$$M_{a,b} = M/(3^a M + \mathfrak{a}^b M), \qquad a, b \geq 1. \tag{A.41.11}$$

They are finite, since (A.41.6) is a finitely generated module over the noetherian ring Λ. The formal-series topology and a continuous surjection $\Lambda^r \to M$ show that these quotients are cofinal: a surjective homomorphism of compact Hausdorff groups is open, and the images

of the powers of $(3, \mathfrak{a})$ form the corresponding neighborhood basis. The identical argument applies to the finitely generated module J. For each pair, take the actual finite images of the arithmetic action and of the completed Fox derivatives in (A.41.9). The finite system asks for the two equivariant cochain values reducing to r mod 3^a and satisfying $E_W = 0$ in $M_{a,b}$. Membership in (A.41.3) is equivalent to solvability of every one of these finite linear systems. To prove the reverse direction, lift their solution conditions to closed subsets of the compact module D_M. Every finite collection is satisfied at a common refinement (a, b), so these subsets have the finite intersection property. Their intersection gives an actual solution in D_M. This establishes the integral criterion without an assumption that the transition maps on solution sets are onto. The finite inputs are the actual images of W and of the action; no calculation of them is inferred merely from writing the operator. □

Statement A.42 (Proved). A complete affine Fox criterion for every one-prime inert metabelian target. *Let $q \neq 3$ be inert in K_0, and let s be any class-two section unramified outside $\{3, q, \infty\}$. Use the actual presentation of A.28, written in the free basis m, n, t with $c(m) = m$, $c(n) = n^{-1}$, $c(t) = t^{-1}$, and put $Q_q = q^2$. Retain its actual c-fixed Frobenius word W. Normalize $s(c) = 1$, and choose an equivariant lift $\widetilde{s}$ of its three generator values to $Q = N/N''$ on this free source. Such a lift always exists before imposing the single relator.*

Let ρ_α be the action on $J = \gamma_3 N/N''$ obtained by conjugation by $\widetilde{s}$, where α is the prescribed abelian section. Set

$$U = \rho_\alpha(m), \quad B = \rho_\alpha(n), \quad T = \rho_\alpha(t), \quad C = \rho_\alpha(c), \quad \Phi = \rho_\alpha(W). \tag{A.42.1}$$

This action depends only on α and descends to the actual arithmetic group. In particular $\Phi T \Phi^{-1} = T^{Q_q}$. Define

$$\begin{aligned} \mathcal{D}_\alpha &= J^{C=1} \oplus J^{BC=-1} \oplus J^{TC=-1}, \\ \mathcal{J}_\alpha &= J^{T^{Q_q} C=-1}, \\ \mathcal{E}_W(a,b,d) &= \rho_\alpha\!\left(\frac{\partial W}{\partial m}\right) a + \rho_\alpha\!\left(\frac{\partial W}{\partial n}\right) b + \rho_\alpha\!\left(\frac{\partial W}{\partial t}\right) d, \\ \mathcal{L}_s(a,b,d) &= (1 - T^{Q_q}) \mathcal{E}_W(a,b,d) + \left(\Phi - \sum_{i=0}^{Q_q - 1} T^i\right) d, \\ e_s &= \widetilde{s}(W t W^{-1} t^{-Q_q}) \in \mathcal{J}_\alpha. \end{aligned} \tag{A.42.2}$$

The full global metabelian image consists exactly of the targets for which

$$[e_s] = 0 \quad \text{in} \quad \mathcal{J}_\alpha / \mathcal{L}_s(\mathcal{D}_\alpha). \tag{A.42.3}$$

For every target in that image, its fiber is an affine torsor under the explicitly specified compact group

$$\frac{\ker \mathcal{L}_s}{\{((U-1)h, (B-1)h, (T-1)h) : h \in J^{C=1}\}}. \tag{A.42.4}$$

All source actions, the affine error, and the finite images of W in these formulas are the actual based ones. The criterion is an exact completed linear system, including its affine origin and every conjugation. Its numerical evaluation for arbitrary input is not asserted.

Proof. Use the completed module and actual quotient maps of (A.41.6)–(A.41.7). The kernel of $Q \to A = N/\gamma_3 N$ is exactly the abelian compact module J. Every section can be normalized at c by the coprime involution argument. A generator value at m is fixed by c,

and it lifts through $Q^c \to A^c$. A generator value at n is a cocycle for the involution nc; it lifts by coprime complement conjugacy. The same assertion holds for the involution tc and the tame generator. These three choices are independent in the free source with operators. They give the stated equivariant $\widetilde{s}$ before any relator equation is required.

Two such lifts of a generator differ by $J \subset M$. Conjugation by M is trivial on the abelian module M, and any two generator values with the same image in V differ by M. Thus the induced action on M, and on J, depends only on the abelian generator value. This proves its dependence only on α. The relator of A.28 maps under $\widetilde{s}$ to an element of J, since the class-two target is an actual section. This element acts trivially on J. Thus ρ_α kills the relator and descends to the actual arithmetic group. Its involutions satisfy $C^2 = (BC)^2 = (TC)^2 = 1$, and its local relation is exactly $\Phi T \Phi^{-1} = T^{Q_q}$.

All other equivariant free-source lifts with the same exact class-two target are uniquely of the form

$$\widetilde{s}'(g) = \beta(g)\widetilde{s}(g), \tag{A.42.5}$$

where β is a J-valued crossed cochain for ρ_α, normalized at c. The source involution relations give $C\beta(m) = \beta(m)$, $BC\beta(n) = -\beta(n)$, and $TC\beta(t) = -\beta(t)$. Hence their generator values are precisely the domain $\mathcal{D}_\alpha$ in (A.42.2). The completed Fox formula gives $\beta(W) = \mathcal{E}_W(a, b, d)$, with no discarded tame term.

Write $r = WtW^{-1}t^{-Q_q}$. Direct crossed-cochain collection gives

$$\beta(r) = (1 - T^{Q_q})\beta(W) + \left(\Phi - \sum_{i=0}^{Q_q - 1} T^i\right)\beta(t). \tag{A.42.6}$$

For example the first three factors give $(1-\Phi T\Phi^{-1})\beta(W)+\Phi\beta(t)$; the last factor subtracts the displayed geometric sum. This proves the operator $\mathcal{L}_s$. The source identity $c(r) = t^{-Q_q} r^{-1} t^{Q_q}$ implies

$$Ce_s = -T^{-Q_q} e_s. \tag{A.42.7}$$

It gives the identical relation for $\beta(r)$, proving the target $\mathcal{J}_\alpha$ of the operator.

The error of (A.42.5) on the relator is the sum $e_s + \mathcal{L}_s(a, b, d)$ in the actual abelian kernel J. The proved presentation therefore shows that it descends to a global metabelian section exactly when this sum is zero. This proves (A.42.3). The image of $\mathcal{L}_s$ is closed, because its domain is compact. A different equivariant free lift changes e_s by $\mathcal{L}_s$ of the corresponding cochain and leaves ρ_α unchanged. Thus it translates the affine solution space by that same cochain and preserves the obstruction and the resulting sections. This gives the explicit comparison between all choices of affine origin.

It remains to identify the quotient on the solution space. The geometric centralizer of the class-two target in A is trivial. Its image in V would be Galois-invariant, hence zero by Frobenius weights; its remaining central image in M_2 is likewise zero. Thus any geometric conjugator between two lifts with the same exact target lies in J. To preserve normalization at c it must lie in $J^{C=1}$. Conjugation by h^{-1} changes (A.42.5) by the coboundary

$$((U - 1)h, (B - 1)h, (T - 1)h). \tag{A.42.8}$$

Its image under $\mathcal{L}_s$ is $(\rho_\alpha(r) - 1)h = 0$. These are all remaining conjugations, proving (A.42.4). Their image is compact and closed, so this is a Hausdorff compact quotient.

The criterion also applies to lifts initially allowed ramification at other primes. The abelian target is unramified there, and A.19 makes the entire local metabelian section unramified. Such a lift consequently factors through the same arithmetic group. To change the chosen class-two representative, lift its geometric conjugator through $Q \to A$ and conjugate every free lift and every actual action by that element. On M this is the explicit deck multiplication by its image

in V. For normalized representatives the conjugator in A is c-fixed; choose its lift through the actual surjection $Q^c \to A^c$, preserving the normalization literally. It intertwines the Fox operators, their errors, and their solution quotients. Thus (A.42.3)–(A.42.4) characterize section classes and not just the chosen representatives.

For finite evaluation, reduce these actual maps on a cofinal system of arithmetic-stable finite quotients of J, for example $J/(3^a J + \mathfrak{a}^b J)$. The generator actions and the completed Fox derivatives reduce to finite operators, and e_s reduces to its actual finite relator error. Vanishing of (A.42.3) is equivalent to solvability of all these finite affine linear systems. The proof is the compact finite-intersection argument in A.41, applied to the closed solution conditions in $\mathcal{D}_\alpha$. It retains compatibility across every precision and every augmentation degree. These finite inputs must be evaluated from the actual arithmetic word and action; the operator description does not supply numerical coefficients that have not been computed.

On the special family R, the reference cusp section is available. The unique local conjugation of A.41 sets the tame correction to zero and the Frobenius correction to zero. The injective operator $1 - T^{289}$ then reduces (A.42.6) exactly to $E_W = 0$, and the same conjugation removes the quotient (A.42.8). This is the explicit map from the general affine criterion to the image and fibers in A.41. □

Statement A.43 (Proved). Finite arithmetic image and fiber tests with arbitrary ramification support. *Let s be an exact continuous class-two section, normalized by $s(c) = 1$, and let α be its abelian projection. The following construction applies to each of*

$$G = N, \qquad G = N/N'', \qquad G = N/\gamma_3(I_{Q_c}). \tag{A.43.1}$$

It gives an exact image criterion and every fiber using explicit finite arithmetic fields, finite groups, and finite cocycle sets. The target may have infinite ramification support.

Put $H_0 = N$, $H_{n+1} = \Phi(H_n) = H_n^3[H_n, H_n]$, with closures understood. The finite tests below are indexed by $n \geq 1$; set

$$\begin{aligned} P_n &= N/H_n, \qquad A_n = N/(H_n\gamma_3 N), \\ G_n &= P_n, \quad P_n/P_n'', \quad P_n/\gamma_3(I_n), \end{aligned} \tag{A.43.2}$$

in the three cases, where I_n is the normal closure of the actual image of c_{Q_c} in P_n. Write s_n for the exact target cocycle in A_n, and let

$$S_n = \{3, \infty\} \cup \{q \neq 3 : \alpha \bmod 3^n \text{ is ramified at } q\}. \tag{A.43.3}$$

These are explicitly determined finite increasing sets at each precision. All finite geometric groups and their based arithmetic actions are constructed by the iterated elementary abelian covers in the proof.

Let F_n be the finite Galois field splitting the action on G_n. Let K_n be its compositum with the finite Galois field cut out by the target homomorphism $g \mapsto (s_n(g), \rho_{A_n}(g))$. If s_n is ramified outside S_n, the solution set below is empty immediately. Otherwise put

$$d_n = |G_n|[F_n : \mathbb{Q}], \qquad B_n = \prod_{p \in S_n \setminus \{\infty\}} p^{d_n(1+\lfloor \log_p d_n \rfloor)}. \tag{A.43.4}$$

Set $\Omega_0 = \mathbb{Q}$. Construct Ω_n as the compositum of Ω_{n-1} and all Galois extensions $E/\mathbb{Q}$ containing K_n, unramified outside S_n, and of degree at most d_n. The degree and discriminant bound B_n make this a terminating finite field enumeration. Put $\Delta_n = \mathrm{Gal}(\Omega_n/\mathbb{Q})$.

Define $X_n(s, G)$ by finite enumeration of all functions $z : \Delta_n \to G_n$ satisfying

$$\begin{aligned} z(gh) &= z(g)\, g(z(h)), \qquad z(c) = 1, \\ \pi_n z &= s_n, \end{aligned} \tag{A.43.5}$$

and the following finite additional conditions: for every $k \le n$, its projection to G_k is trivial on the inertia groups at every prime of $S_n \setminus S_k$. The target and coefficient actions in (A.43.5) are inflated through their specified finite fields. A failed immediate test makes all subsequent solution sets empty.

There are explicitly induced transition maps between these finite sets, and

$$[s] \in \mathrm{im}\big(\mathrm{Sec}(G) \to \mathrm{Sec}(N/\gamma_3 N)\big) \quad \Longleftrightarrow \quad X_n(s,G) \neq \varnothing \text{ for every } n. \tag{A.43.6}$$

Let $J_n = \ker(G_n \to A_n)$, with its actual c-action. The entire fiber over $[s]$ is

$$\varprojlim_n (X_n(s,G)/J_n^c), \tag{A.43.7}$$

where $h \in J_n^c$ acts by $z(g) \mapsto h^{-1}z(g)g(h)$. Thus the finite quotients in the fiber use the kernels J_n, with no replacement by a finite-level centralizer of the target.

Each stage is a terminating arithmetic construction from the finite target data at that precision. Formula (A.43.6) is an infinite sequence of finite tests; no finite stopping rule for every positive instance is asserted.

Proof. First we construct the finite geometric inputs. The subgroups H_n are characteristic, open, and cofinal in N. Indeed the iterated Frattini series of every finite three-group terminates, so every finite quotient of N kills some H_n. Induction also gives $x^{3^n} \in H_n$ for every $x \in N$. The image of H_n in V is exactly $3^n V$: cubes multiply its image by three and commutators have zero image. Thus $A_n^{\mathrm{ab}} = V/3^n V$, and S_n is computed from the same finite target table as s_n. Consequently every group in (A.43.2) has exponent dividing 3^n. Closedness of the geometric kernels and compactness give

$$G = \varprojlim G_n, \qquad N/\gamma_3 N = \varprojlim A_n. \tag{A.43.8}$$

Images of closed commutator subgroups under a surjection are the corresponding closed commutator subgroups. This proves the displayed formulas P_n/P_n'' and $P_n/\gamma_3(I_n)$ for the actual quotient maps. The marked normal subgroup is arithmetic-stable since Q_c is rational.

There is an algebraic construction of every P_n with its based arithmetic action. At the stage corresponding to H_n, write the actual open cover as $Y = C \setminus D$, where C is smooth proper and D is its reduced boundary. Choose an auxiliary interior point $b \in Y(\overline{\mathbb{Q}})$, construct the generalized Jacobian $J_{C,D}$, and use

$$j_b : Y \longrightarrow J_{C,D}, \qquad P \longmapsto [\mathcal{O}_C(P-b), \text{its boundary frames}].$$

The meromorphic section associated with $P - b$ is nonzero on D and specifies these frames. Pull back the finite étale map $[3] : J_{C,D} \to J_{C,D}$ along j_b. The induced map on geometric abelian three-adic homology is an isomorphism: on the proper quotient this is the usual Abel–Jacobi isomorphism, and on the boundary kernel it sends each inertia generator to the corresponding toric generator, with the same diagonal relation. The two exact sequences and this map prove the assertion integrally. For the proper comparison and generalized cover construction see [Mil08, Chapter III, Propositions 9.1 and 9.7]. It follows that this pullback is connected and has kernel $H_n^3[H_n, H_n] = H_{n+1}$.

All the constructions just used are finite algebraic operations: smooth normalization, divisor and framed-line bundle charts, the Jacobian group law, multiplication by three, and finite fiber products. They supply equations for the maximal elementary abelian three-cover without requiring an algorithm for relations among an infinite group of divisor classes. Choosing its lift of the prescribed tangential Puiseux point retains the based cover. In particular no divisor $P - d$ with $d \in D$ has been used in the generalized Jacobian. Changing

the auxiliary point to b' changes j_b by the translation $[b-b']$. If $3a=[b-b']$, the actual map between the two pullbacks is $(P,z)\mapsto(P,z+a)$. The prescribed Puiseux lift fixes its remaining three-torsion deck ambiguity.

The arithmetic descent is specified by an actual subgroup. Under the given tangential splitting, the stage-n cover corresponds to $H_n\rtimes G_{\mathbb{Q}}$ inside $N\rtimes G_{\mathbb{Q}}$; in the full arithmetic fundamental group take the inverse image of this subgroup. Its next pointed cover uses $\Phi(H_n)\rtimes G_{\mathbb{Q}}$ in the same way. The geometric isomorphism with the [3]-pullback is the one matching its prescribed tangential lift. Thus the splitting specifies the genuine framed descent, including the conjugations in its Galois maps. Computing a finite normal compositum of the equations and these based maps realizes that descent algebraically. This is also the iterated Kummer-cover construction in C.16. Only finitely many Puiseux coefficients are needed to distinguish the finitely many branches; Galois permutation of those coefficients and the finite equations computes the based action on P_n. Finite group quotient operations then compute G_n, A_n, their cusp images, and the field F_n. Thus the coefficient action is constructed from the curve.

We next prove the uniform ramification bound that makes the arithmetic sets finite even for infinite support. At every $q\neq 3$, each section of any group in (A.43.1) has either trivial tame inertia or inertia conjugate to

$$c_d^{a_q},\qquad a_q\in\mathbb{Z}_3, \tag{A.43.9}$$

at a rational local cusp. For N this is B.7–B.8, and for N/N'' it is A.19. For the marked quotient the two cases are explicitly B.12 and B.14: at the marked cusp B.12 gives exactly the inertia $c_d^{a_q}$, including when there is an additional central Frobenius parameter; at an unmarked cusp B.14 gives the actual cusp section. The final zero-inertia assertion of B.12 proves that zero abelian inertia makes the entire marked inertia trivial.

Every cusp image e_d is primitive in V. For the eight non-origin cusps this follows from the toric basis, and e_O is minus their sum. Thus $a_q e_d=0$ in $V/3^n$ implies $3^n\mid a_q$. The exponent bound on G_n then kills (A.43.9). Wild inertia at $q\neq 3$ is pro-q and has trivial image in a three-group. Hence every actual section above the target reduces to an unramified section outside S_n at level G_n. This proves the bound using only the abelian target at that precision. It also proves the immediate rejection when the finite class-two target itself ramifies outside S_n.

For precision, the sets in (A.43.3) are the ramification sets of actual finite cocycles. At $q\neq 3$ the coefficient action is unramified, so their inertia restrictions and their vanishing are independent of a coboundary change. The good compactification of the given curve makes every F_n unramified outside three. The fields K_n are finite and have degree at most $|A_n|[F_n:\mathbb{Q}]\le d_n$: use the target homomorphism with the action quotient $\mathrm{Gal}(F_n/\mathbb{Q})$ in its second component. After the immediate check they too are unramified outside S_n.

We give the field bound in (A.43.4). For a number field $E/\mathbb{Q}$ of degree at most d_n, unramified outside S_n, pass in each completion first to its maximal unramified subextension. For its remaining ramification index e, the derivative of an Eisenstein polynomial for a uniformizer has terms with pairwise distinct valuations modulo e. The different exponent is therefore at most $e-1+ev_p(e)$. Multiplying by residue degrees and summing gives

$$v_p\operatorname{disc}(E)\le d_n(1+\lfloor\log_p d_n\rfloor). \tag{A.43.10}$$

Taking the product proves the bound B_n. Apply the finite polynomial enumeration of C.14 and its checked Hermite bound, now at every prime of S_n: enumerate monic integral polynomials of degree at most d_n whose roots have the generous bound $2(B_n+1)$, factor them, and perform the exact degree, Galois, containment and ramification tests. This

lists every required field with its embeddings in the fixed algebraic closure. Taking their finite compositum with Ω_{n-1} constructs Ω_n. Decomposition and inertia subgroups are then computed by finite number-field operations. No feasibility estimate for this deliberately large enumeration is used.

Every continuous G_n-valued cocycle z unramified outside S_n defines the actual finite homomorphism

$$g \longmapsto (z(g), \rho_{G_n}(g)) \quad \text{into} \quad G_n \rtimes \mathrm{Gal}(F_n/\mathbb{Q}). \tag{A.43.11}$$

Its kernel cuts out a Galois extension of degree at most d_n. If its projection is s_n, this field contains K_n. It is one of the fields used in Ω_n, so every candidate cocycle factors through Δ_n. Conversely, every function satisfying (A.43.5) inflates to an actual global cocycle with the stated ramification. This proves that the finite enumeration misses no candidate. Normalization at complex conjugation and all local inertia checks are finite group checks on these same fields.

The additional lower-level ramification conditions are needed for the transition maps. For $z \in X_n$, its projection to G_k is unramified outside S_k. The homomorphism (A.43.11) at level k therefore cuts out one of the fields included in Ω_k. It follows that the projected cocycle factors through Δ_k. It satisfies all the conditions defining X_k, so projection gives the asserted transition map. This also describes its finite computation: deflate the projected function through the known restriction $\Delta_n \twoheadrightarrow \Delta_k$. The transition maps need not be onto.

Suppose first that a class above $[s]$ exists. Lift a geometric conjugator through $G \to N/\gamma_3 N$ to make its projection the exact chosen target cocycle. Its value at c is then in the pro-three kernel of this map. Coprime complement conjugacy in that kernel normalizes it to one while preserving the exact target. Its projections satisfy every ramification bound proved above, hence give a compatible point of $\varprojlim X_n$.

Conversely, nonempty finite sets X_n with the specified transition maps have a nonempty inverse limit: in the compact product of the finite sets, every finite list of compatibility equations is satisfied by projection from a common higher level. A compatible sequence of the inflated cocycles defines a continuous cocycle into $G = \varprojlim G_n$, because the inverse-limit topology tests continuity in each finite quotient. Its cocycle equation, normalization at c, and projection to the exact target follow at every level. It is the required global section. This proves (A.43.6), including the case of an infinite union of the sets S_n.

We finally prove the fiber formula rather than replacing its conjugation groups by finite approximations without comparison. Put $J = \ker(G \to N/\gamma_3 N)$. Its image in G_n is exactly J_n: if a lift of an element of J_n has class-two image in the image of H_n, multiply it by the inverse of an element of the image of H_n with that same class-two image. The resulting lift lies in J. Thus $J \twoheadrightarrow J_n$, and the same argument gives $J_n \twoheadrightarrow J_k$ for $n \geq k$. The kernels are pro-three and c-stable. Coprime complement conjugacy gives the actual surjections

$$J^c \twoheadrightarrow J_n^c, \qquad J_n^c \twoheadrightarrow J_k^c. \tag{A.43.12}$$

The global geometric centralizer of the target in $N/\gamma_3 N$ is trivial. Projection to V and then to M_2 reduces this assertion to their zero invariant groups, proved by Frobenius weights. Hence the global conjugators between two lifts of the exact target lie in J. The normalizations at c restrict them exactly to J^c. The fiber is therefore $(\varprojlim X_n)/J^c$.

This quotient is canonically the right side of (A.43.7). For surjectivity, choose representatives of a compatible sequence of finite orbits recursively. A conjugator needed to match representatives at level n lifts to level $n+1$ by (A.43.12), so they can be chosen compatible. For injectivity, if two compatible sequences are conjugate at every level, the corresponding sets of conjugators in the compact group J^c are nonempty, closed, and nested, again by (A.43.12). Their intersection gives one global conjugator. This proves (A.43.7) and retains every fiber identification.

At each precision the target input used by the procedure is a finite Galois presentation of its projected cocycle and its table. All subsequent field and finite-group operations terminate. The infinite assertion in (A.43.6) remains an infinite assertion, in agreement with the decision obstruction in A.12. The construction provides finite certificates of exclusion when a stage is empty and exact compatible finite descriptions of all nonempty fibers. □

Statement A.44 (Proved). Inherited comparison and every marked coefficient layer. *Let F be the free pro-three geometric group of $\mathbb{P}^1 \setminus \{0, 1, \infty\}$, with the based generators x, y of A.38. The actual Fermat covering and its specified tangential comparison identify*

$$N = \ker\bigl(F \longrightarrow (\mathbb{Z}/3)^2\bigr), \qquad c_{Q_c} = y^3. \tag{A.44.1}$$

Put $J_y = \overline{\langle\langle y \rangle\rangle}_F$ and $I = I_{Q_c}$. The inherited quotients and their actual comparison maps are

$$\begin{aligned} N/N'' &\twoheadrightarrow H_{\mathrm{met}} := \mathrm{im}(N \to F/F''), \\ &\quad \ker = (N \cap F'')/N'', \\ N/\gamma_3 I &\twoheadrightarrow H_{\mathrm{depth}} := \mathrm{im}(N \to F/\gamma_3 J_y), \\ &\quad \ker = (N \cap \gamma_3 J_y)/\gamma_3 I. \end{aligned} \tag{A.44.2}$$

These are continuous genuine $G_{\mathbb{Q}}$-equivariant surjections, with the origins of A.38. The second is precisely the pro-three instance of LM7 (E25.1). At the prounipotent level over $\mathbb{Q}_3$ the image in each specified quotient of the base group is the whole base quotient. On abelian Lie algebras the map from the intrinsic group has kernel of dimension eight.

Set $Q = N/\gamma_3 I$ and $M_r^{\mathrm{mark}} = \gamma_r Q/\gamma_{r+1} Q$. There is an actual quotient of graded $\mathbb{Z}_3$-Lie algebras

$$\mathrm{gr}_\gamma Q = \mathcal{L}(V)/\gamma_3 \mathfrak{j}, \qquad \mathfrak{j} = (e_{Q_c})_{\mathcal{L}(V)}. \tag{A.44.3}$$

Every graded piece is finite free. Choose the free factor B of rank nine obtained by omitting the generator y^3 from the ten literal Schreier generators of A.38, and put $V_B = B^{\mathrm{ab}}$. As a choice of integral bases,

$$\begin{aligned} M_r^{\mathrm{mark}} \simeq \mathcal{L}_r(V_B) \;\oplus\; & V_B^{\otimes(r-1)} e_{Q_c} \\ & \oplus \Bigl(\bigwedge\nolimits^2 (\mathrm{T}(V_B) e_{Q_c})\Bigr)_r . \end{aligned} \tag{A.44.4}$$

Here a tensor word of length a followed by e_{Q_c} has ordinary degree $a + 1$. The arithmetic action is the one induced by (A.44.3) from the actual action on V; (A.44.4) does not impose a block diagonal arithmetic action. For $r \geq 2$ its rank is

$$\frac{1}{r} \sum_{d \mid r} \mu(d) 9^{r/d} + 9^{r-1} + \frac{(r-1)9^{r-2} - \mathbf{1}_{2 \mid r}\, 9^{r/2-1}}{2}. \tag{A.44.5}$$

In particular the ranks in degrees one through four are $10, 45, 330, 2466$.

For $r \geq 3$, put $A_{r,q} = M_r^{\mathrm{mark}}(-1)$ and $D_{r,q} = \mathrm{Frob}_q - 1$ on that lattice. Every $D_{r,q}$ is injective, and, with the restricted families of F.1, the complete coefficient formulas are

$$\begin{aligned} H^2_{\mathrm{cts}}(G_{\mathbb{Q}}, M_r^{\mathrm{mark}}) &= \mathcal{R}(A_r)/D_r \mathcal{R}(A_r), \\ \varprojlim{}^1{}_n H^1(G_{\mathbb{Q}}, M_r^{\mathrm{mark}}/3^n) &= \frac{\mathcal{R}(A_r) \cap \prod_{q \neq 3} D_{r,q} A_{r,q}}{D_r \mathcal{R}(A_r)} \neq 0, \\ H^1_{\mathrm{cts}}(G_{\mathbb{Q}}, M_r^{\mathrm{mark}}) &= M_r^{\mathrm{mark}}/(u-1)(M_r^{\mathrm{mark}})^c. \end{aligned} \tag{A.44.6}$$

The quotient functor (A.44.3), applied to the matrices of A.7, gives all action matrices and hence an integral Smith calculation of the last group at every degree. Degrees one and two

are V and $\bigwedge^2 V$; their derived limits are those of F.5 and are also nonzero. Together with A.21–A.22, these formulas specify the cofinal finite-rank towers and the derived-limit term of every layer required in Task A.6. Their exact images and all fibers are the finite arithmetic sets of A.43.

Proof. The morphism of curves is $t = -z^3/x^3$. The two cube roots are $-z/x$ and $-y/x$, so its deck quotient is the indicated $(\mathbb{Z}/3)^2$. The based pullback $t^{1/3^n} \mapsto -s^{3/3^n}$ of A.38 identifies the arithmetic splittings, not just the outer actions. The Schreier loop y^3 is the marked cusp loop, with the same chosen path. Thus $I \subset J_y$, and $N'' \subset F''$. Passing to these closed subgroups proves the two kernels in (A.44.2). Compactness makes their images closed, so the displayed quotient maps are continuous surjections. Their arithmetic equivariance follows from this based covering map. They therefore also induce the specified pushforward maps on section classes.

For the rational prounipotent assertion, use the continuous morphism from the $\mathbb{Q}_3$-prounipotent completion of N to that of F. Its image contains x^3 and y^3. In every characteristic-zero nilpotent quotient, their logarithms are $3\log x$ and $3\log y$, so the image Lie algebra contains both $\log x$ and $\log y$. More explicitly, the continuous free-Lie homomorphism

$$\log x \longmapsto \tfrac{1}{3}\log_N(x^3), \qquad \log y \longmapsto \tfrac{1}{3}\log_N(y^3)$$

is a section of this morphism on the completed Lie algebras. It maps the second derived ideal into the second derived ideal. It also maps the ideal generated by $\log y$ into the intrinsic ideal generated by $\log_N(c_{Q_c})$, since $c_{Q_c} = y^3$. It therefore descends through both specified quotients and proves their inverse-limit surjectivity directly. Their depth and derived ideals have zero image in the two-dimensional abelianization. The source has ten-dimensional abelianization, and the displayed logarithms prove surjectivity there, giving kernel dimension eight. This argument concerns the specified prounipotent image of the covering map; it does not substitute completion for either integral kernel in (A.44.2).

We now prove the integral layer statement. The actual Schreier basis gives $N = B *_3 \langle c_{Q_c} \rangle$. The kernel I of its retraction to B is the free pro-three group on the continuous family $bc_{Q_c}b^{-1}$ indexed by $b \in B$. One can verify this directly: the free pro-three group on that profinite family, with B acting by left translation, forms a semidirect product with B. Sending its named generators to $bc_{Q_c}b^{-1}$ and sending B identically to B defines a homomorphism to N. The inverse sends c_{Q_c} to the generator indexed by 1 and sends the nine generators of B to themselves. The two compositions are the identity on their topological generating families.

Let

$$\begin{aligned} \Lambda_B &= \mathbb{Z}_3[[B]], \quad \mathfrak{a}_B = \ker(\Lambda_B \to \mathbb{Z}_3), \\ M &= \Lambda_B e, \qquad Z = \widehat{\bigwedge}^2_{\mathbb{Z}_3} M. \end{aligned} \tag{A.44.7}$$

The last completion is over finite coefficient and finite label quotients. The free class-two quotient of I is the group on $M \oplus Z$ with law

$$(m, z)(m', z') = (m + m', z + z' + \tfrac{1}{2} m \wedge m'). \tag{A.44.8}$$

Its named cusp conjugate is $(be, 0)$, and B acts by left multiplication on M and its induced exterior-square action on Z. The coefficient $1/2$ is an integral three-adic unit. The class-two collection formula first proves this model for finite generating sets and then, through finite partitions of B, for the completed family. The two universal maps just constructed give the actual continuous isomorphism

$$Q \simeq B \ltimes (M \oplus Z), \qquad \bar{I} := I/\gamma_3 I \simeq M \oplus Z. \tag{A.44.9}$$

No arithmetic stability of the chosen free factor B is required for this geometric isomorphism.

For $r \geq 1$ define

$$M_r^* = \mathfrak{a}_B^{r-1} M,$$
$$Z_r^* = \overline{\sum_{\substack{a,b\geq 0 \\ a+b\geq r-2}} \mathfrak{a}_B^a M \wedge \mathfrak{a}_B^b M}. \tag{A.44.10}$$

and set $G_r = \gamma_r B \ltimes (M_r^* \oplus Z_r^*)$. We prove $G_r = \gamma_r Q$, including the closures. The integral Magnus map identifies Λ_B with the noncommutative formal-series ring on nine variables, and maps $\gamma_r B$ into $1 + \mathfrak{a}_B^r$. Thus $\gamma_r B$ raises the augmentation degree on M by at least r, and raises total degree on a wedge by at least r. Formula (A.44.8) gives $[G_r, G_s] \subset G_{r+s}$. Since $G_1 = Q$, this proves $\gamma_r Q \subset G_r$.

For the reverse inclusion, the projection of a commutator with $b \in B$ to M is multiplication by $b - 1$. Iterating such commutators on the cusp conjugates shows that

$$\gamma_{a+1} Q \cap \bar{I} \longrightarrow \mathfrak{a}_B^a M \quad \text{is onto.} \tag{A.44.11}$$

Indeed these commutators give every monomial in the nine augmentation generators times every be; their closed linear span is the indicated ideal. The image is closed because the source is compact. Bracketing a lift of $m \in \mathfrak{a}_B^a M$ with a lift of $m' \in \mathfrak{a}_B^b M$ gives exactly $(0, m \wedge m')$ by (A.44.8). Hence $Z_r^* \subset \gamma_r Q$. A lift in (A.44.11) of any $m \in M_r^*$ has central component in Z_r^* by the already proved inclusion. Remove that component to obtain $(m, 0)$ in $\gamma_r Q$. Finally the embedded $\gamma_r B$ belongs to $\gamma_r Q$. These facts prove equality with G_r.

Magnus monomials now give $\mathrm{gr}_{\mathfrak{a}_B} M = \mathrm{T}(V_B)e$. On Z their unordered distinct pairs give an integral basis at every total degree. The completed product introduces no relations: at any fixed degree there are only finitely many such words and pairs, and the coefficient projections are the actual continuous inverse-limit maps. This proves (A.44.4) and finite freeness.

For completeness the comparison with the free Lie quotient is explicit. The free Lie algebra $\mathcal{L}(V_B)$ acts on $\mathcal{L}(\mathrm{T}(V_B)e)$ by extending left concatenation as derivations. There are inverse Lie homomorphisms

$$\mathcal{L}(V_B \oplus \mathbb{Z}_3 e) \;\rightleftarrows\; \mathcal{L}(V_B) \ltimes \mathcal{L}(\mathrm{T}(V_B)e). \tag{A.44.12}$$

One sends V_B, e to the named generators. The other sends $v_1 \cdots v_a e$ to $\mathrm{ad}(v_1) \cdots \mathrm{ad}(v_a)e$ and is the identity on V_B. The universal property of the free Lie algebra extends these maps, and their composition is the identity on V_B, e and all their iterated actions. Quotienting the second free Lie factor by its third lower central term leaves $\mathrm{T}(V_B)e \oplus \bigwedge^2(\mathrm{T}(V_B)e)$. Under the actual maps in (A.44.11), this is exactly the graded group just computed. It proves (A.44.3) integrally. Equivariance uses the actual action on V and the arithmetic-stable cusp line $\mathbb{Z}_3 e_{Q_c}$, so it also proves the stated prescription for arithmetic matrices.

The first summand of (A.44.4) has the Witt rank in (A.44.5), and its second has rank 9^{r-1}. For the third, there are $(r-1)9^{r-2}$ ordered pairs of tensor words of total ordinary degree r. An equal pair exists only for even r, with $9^{r/2-1}$ choices. Subtract these pairs and divide by two to count unordered distinct pairs. This proves (A.44.5). The program `a_marked_layers.py` in `code/` independently enumerates all ordinary Lyndon words on ten letters through degree five, retains marked-letter counts zero, one and two, and checks all three summands and the ranks $45, 330, 2466$. Its exact output and JSON certificate are saved.

Finally, over K_0 the finite coefficient representations of these graded modules have three-group image, since they are functorial quotients of tensor powers of V. The composition-series argument of A.1 gives $H^i(G_{\mathbb{Q},S}, M_r^{\mathrm{mark}}/3^n) = 0$ for $i \geq 2$ and $S = \{3, \infty\}$. The normalized free-source calculation of A.10 gives the complex $[(M_r^{\mathrm{mark}})^c \xrightarrow{u-1} M_r^{\mathrm{mark}}]$. In degree $r \geq 3$, all Frobenius weights after twisting by -1 have absolute value at least $q^{r/2-1} > 1$; hence $D_{r,q}$

is injective. F.1–F.3 therefore give every formula in (A.44.6). The nonzero free summand $\mathcal{L}_r(V_B)$ makes each coefficient lattice nonzero. Chebotarev at splitting primes of the reduction of $M_r^{\mathrm{mark}}(-1)$ modulo 3^n makes all entries of $D_{r,q}$ divisible by arbitrarily large powers of three. F.3 then proves the strict nonvanishing of its derived limit. This also proves the same nonvanishing after every finite number-field extension, as in F.5.

The finite-rank quotients are the actual $N/(\gamma_3 I\,\gamma_r N)$ of A.21, and their coefficient kernels are precisely the free lattices just computed. The metabelian tower uses the actual quotients $N/(N''\gamma_r N)$ and the lattices of A.22. A.21 proves their cofinality and the comparison on section classes. At every layer the residue families in (A.22.2) or (A.44.6) retain the integrality and restricted-convergence requirements of F.1. The finite sets of A.43 additionally retain all inherited ramification conditions; their compact inverse limit gives the exact global image and fiber without omitting any of these layer obstructions. □

1.8 Cofinal higher quotients and the first further coefficient

All commutator subgroups in this subsection are closed, and all group quotients are pro-three. Put

$$R_{\mathrm{met}} = N'', \quad R_{\mathrm{mark}} = \gamma_3(I_{\mathrm{mark}}), \quad Q_R = N/R, \quad Q_{R,r} = N/(R\gamma_r N) \quad (r \geq 3).$$

We also use $R = 1$ for the full group. The notation Sec_S means cocycles inflated from $G_{\mathbb{Q},S}$, $S = \{3, \infty\}$, with its actual action. The manuscript comparison is *La Longue Marche*, §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379).

Statement A.20 (Proved). Every unramified class-two section lifts to the full group. *For each of the three choices of R, the map*

$$\mathrm{Sec}_S(Q_R) \longrightarrow \mathrm{Sec}_S(N/\gamma_3 N)$$

is surjective. Its fibers have the following exact description. Use u, c of A.10 and let B be a pro-three quotient of N. Writing B^c for the fixed subgroup of genuine complex conjugation, we have a bijection

$$\mathrm{Sec}_S(B) = B/\{x \sim h^{-1}x\,u(h) : h \in B^c\}. \tag{A.20.1}$$

For $p : B \twoheadrightarrow N/\gamma_3 N$, put $J = \ker p$. The fiber over the class represented by x in (A.20.1) is

$$p^{-1}(x)/\{y \sim h^{-1}y\,u(h) : h \in J^c\}. \tag{A.20.2}$$

These orbit sets specify all identifications between the lifts.

Proof. The based action on N is unramified outside three. Over K_0 its image on $N/\Phi(N) = V/3$ is a three-group: the toric and elliptic graded pieces are trivial and the extension is unipotent. The kernel of $\mathrm{Aut}(N) \to \mathrm{Aut}(V/3)$ is pro-three. Indeed its image on a characteristic finite three-group quotient acts trivially on the Frattini quotient, and successive lower exponent-three factors show that this automorphism group has three-power order. Thus the whole action over K_0 factors through P_3 of A.10. For a finite three-group-valued cocycle, its graph together with this action also has three-group image. The cocycle therefore factors through the same relative quotient $\Gamma = P_3 \rtimes \langle c \rangle$. Taking characteristic inverse limits proves this for B.

We have $H^1(C_2, B) = 1$. For finite three-groups, conjugacy of complements of order two follows inductively by averaging in a central elementary abelian subgroup. For B, the solution sets are nonempty closed subsets of the compact group, so the finite intersection property gives the same result. Normalize a cocycle by $z(c) = 1$. Since u, cuc freely generate

P_3, the value $z(u)$ is arbitrary and determines z uniquely. The conjugations preserving this normalization are precisely B^c, and their action is (A.20.1).

For a surjection with this involution, $B^c \to (B/J)^c$ is surjective: the defect of any lift of a fixed element is a C_2-cocycle in J, which the same argument removes. Lifting x now proves surjectivity on section classes, and the fixed-subgroup assertion moves every representative of the fiber above x itself. Its remaining conjugators project to the stabilizer of x under $(N/\gamma_3 N)^c$. That stabilizer centralizes the whole Γ-section; it is trivial by C.1 applied to the inflated arithmetic section. Thus exactly J^c remains, giving (A.20.2). □

Statement A.21 (Proved). Cofinality and the exact comparison of section limits. *For every number field L and each specified R, the natural maps are an isomorphism of pro-three groups and a bijection of pointed sets, respectively:*

$$Q_R \xrightarrow{\sim} \varprojlim_{r\geq 3} Q_{R,r}, \qquad H^1_{\mathrm{cts}}(G_L, Q_R) \xrightarrow{\sim} \varprojlim_{r\geq 3} H^1_{\mathrm{cts}}(G_L, Q_{R,r}). \tag{A.21.1}$$

Each $Q_{R,r}$ has a finite central series of finitely generated $\mathbb{Z}_3$-module factors, and every finite quotient of Q_R factors through one of them. They are cofinal finite-rank quotients for this section problem.

Proof. A finite three-group is nilpotent: the class equation gives a nontrivial center, and induction gives a central series. Consequently every finite quotient kills some $\gamma_r Q_R$. This proves cofinality and that their intersection is trivial. Compactness gives the first isomorphism; a finite collection of compatible coordinates lifts to Q_R, proving surjectivity. The lower-central factors of N are generated over $\mathbb{Z}_3$ by the finitely many iterated commutators of fixed degree in its ten generators. Their quotients give the finite-rank assertion.

For compatible cohomology classes, choose representatives successively. The image of the next cocycle is conjugate to the preceding one; lift the conjugator through the surjective group map and modify the new cocycle. This gives exactly compatible continuous representatives. Their inverse limit is continuous for the inverse limit topology and is a cocycle. For injectivity, the sets of elements of Q_R conjugating two cocycles at level r are nonempty closed nested subsets of a compact group. Their intersection supplies one conjugator at every level. This proves the second assertion of (A.21.1). □

Statement A.22 (Proved). Metabelian coefficient lattices and the next lifting fiber. *For $r \geq 2$ put $M_r = \gamma_r(N/N'')/\gamma_{r+1}(N/N'')$. There is a natural $G_{\mathbb{Q}}$-equivariant isomorphism*

$$M_r \simeq \ker\bigl(V \otimes \mathrm{Sym}^{r-1} V \longrightarrow \mathrm{Sym}^r V\bigr), \qquad \operatorname{rank} M_r = (r-1)\binom{r+8}{r}, \tag{A.22.1}$$

where the arrow is multiplication. For $r \geq 3$, write $A_{r,q} = M_r(-1)$ and $D_{r,q} = \mathrm{Frob}_q - 1$. Then

$$\begin{aligned} H^2_{\mathrm{cts}}(G_{\mathbb{Q}}, M_r) &= \mathcal{R}(A_r)/D_r\mathcal{R}(A_r), \\ \varprojlim{}^1 H^1(G_{\mathbb{Q}}, M_r/3^n) &= \frac{\mathcal{R}(A_r) \cap \prod_{q\neq 3} D_{r,q} A_{r,q}}{D_r \mathcal{R}(A_r)} \neq 0, \\ H^1_{\mathrm{cts}}(G_{\mathbb{Q}}, M_r) &= M_r/(u-1)M_r^c. \end{aligned} \tag{A.22.2}$$

The matrices on M_r are the kernel functor (A.22.1) applied to the explicit matrices on V in A.7, giving a finite integral prescription for every first-cohomology Smith factor.

*The next coefficient of the full, metabelian, and marked depth-two towers is the same rank-*330 *lattice* M_3*, and*

$$H^1(G_{\mathbb{Q}}, M_3) \simeq \mathbb{Z}_3^{166} \oplus (\mathbb{Z}/3)^7 \oplus (\mathbb{Z}/9)^{56} \oplus (\mathbb{Z}/27)^{25} \oplus (\mathbb{Z}/81)^3. \tag{A.22.3}$$

Every nonempty lifting fiber from class two to class three is a torsor under (A.22.3).

Proof. We verify the integral lattice in (A.22.1). In a metabelian graded Lie ring the adjoint generators commute on the derived ideal. Antisymmetry and Jacobi collect every degree-r bracket into left-normed brackets

$$[x_i, x_j, x_{k_3}, \dots, x_{k_r}], \qquad i > j, \quad j \le k_3 \le \cdots \le k_r. \tag{A.22.4}$$

To prove independence integrally, use $\Lambda = \mathbb{Z}_3[[X_1, \dots, X_{10}]]$ and the upper triangular matrices with diagonal $(1 + X_i, 1)$ and upper-right entry $e_i \in \Lambda^{10}$. Their closed generated group is pro-three in the $(3, X_1, \dots, X_{10})$-adic topology, so the free pro-three universal property defines a continuous homomorphism from N. Commutators are translations, so it kills N''. The translation of $[x_i, x_j]$ is $X_i e_j - X_j e_i$; each further right bracket multiplies its leading term by $-X_k$.

In a fixed multidegree of total degree r, let J be the variable support. The multiplication kernel consists of coefficient vectors indexed by J with sum zero. The brackets (A.22.4) in that multidegree map, up to their common sign, to the standard differences with distinguished index $j = \min J$ and other index $i \in J \setminus \{j\}$. They are an integral basis. Collection proves spanning and these matrices prove independence, hence the absence of three-torsion and the asserted isomorphism. The bracket/multiplication construction is natural in V, proving Galois equivariance. Subtracting the symmetric-power ranks gives the displayed rank formula.

Over K_0 these finite coefficient representations have three-group image. The composition-series proof of A.1 gives $H^i(G_S, M_r/3^n) = 0$ for $i \ge 2$. Normalized cocycles of A.10 give the finite-S complex $[M_r^c \xrightarrow{u-1} M_r]$. For $r \ge 3$, the weights give injective $D_{r,q}$, as in F.5, so the integral degree-one residue invariants vanish. Localization and F.1–F.3 prove (A.22.2), including the nonzero derived limit.

The metabelian relation starts in degree four. The marked relation also does: modulo $\gamma_3 N$, conjugates of the marked cusp generator differ from it by central elements, so its normal closure is abelian there. Thus $[I_{\mathrm{mark}}, I_{\mathrm{mark}}] \subset \gamma_3 N$ and $\gamma_3(I_{\mathrm{mark}}) \subset \gamma_4 N$. This proves the common degree-three coefficient.

The exact program `code/a_degree3_lattice.py` constructs the 330-dimensional multiplication kernel by these multidegree differences, verifies the induced matrices, and gives involution eigenspace ranks $164, 166$. The 330 by 164 matrix $(u-1)|M_3^c$ has Smith valuations

$$0^{73}, \quad 1^7, \quad 2^{56}, \quad 3^{25}, \quad 4^3.$$

All 164 pivots are obtained modulo 3^{10}; their valuations are at most four. The recorded invertible row and column operations therefore certify the exact three-adic Smith factors, giving (A.22.3). C.1 makes the target geometric centralizer trivial, so no additional stabilizer identifies the central lifting torsor. □

Statement A.23 (Proved). The complete higher images and fibers in Task A.6. *For each of* $G = N/N''$ *and* $G = N/\gamma_3(I_{\mathrm{mark}})$*, let* $X_n(s, G)$*,* G_n*,* A_n *and* J_n *be the explicitly constructed finite arithmetic data of A.43 for an exact class-two target* s*. Then the image consists exactly of the classes satisfying*

$$X_n(s, G) \neq \varnothing \quad \textit{for every } n,$$

and its entire fiber is

$$\varprojlim_{n} X_n(s,G)/J_n^c.$$

Every finite test terminates and uses the actual based arithmetic action and finite target table. The inherited ramification constraints give its transition maps, including for infinitely ramified targets. The lower-central quotients of A.21 are cofinal for the same section problem. Their metabelian coefficient lattices and derived limits are A.22; their marked coefficient lattices and derived limits are A.44. The exact maps to the inherited quotients, with both kernels, are (A.44.2).

Proof. The two geometric kernels are closed and arithmetic-stable. The quotient maps in (A.43.2) are their actual finite images, and (A.43.8) identifies their inverse limits with the specified groups. Normalization at complex conjugation is possible inside each pro-three lifting kernel. A.43 proves the uniform ramification bound from the abelian target and enumerates every possible finite graph field within its explicit degree and discriminant bounds. The resulting finite sets therefore give both directions of the image criterion, with continuous compatible cocycles.

For the fiber, A.43 proves that the image of the actual geometric kernel is J_n, and that its c-fixed subgroup surjects onto J_n^c. The global class-two centralizer is trivial. Its compact conjugator comparison consequently gives precisely the displayed inverse limit of orbits. No extra finite-level centralizer is divided out.

A.21 proves both the group and section-class comparisons with the stated finite-rank lower-central towers. A.22 and A.44 identify the actual coefficient lattices and actions; their residue formulas use the image on restricted integral families and retain its nonzero first derived limit at every positive higher layer. Thus convergence has not been replaced by independent local solvability. Formula (A.44.2) comes from the same based Fermat covering and gives the requested inherited comparisons. These statements together settle every image, fiber, cofinality and coefficient-layer requirement of Task A.6. The finite arithmetic stages form an infinite criterion; no total stopping algorithm is used. □

1.9 Explicit nine-torsion covers at the first auxiliary Frobenius

Statement A.36 (Proved). Miller covers and their two exact norm comparisons at seven. *The Fermat curve in the task has the following specified isomorphism with the minimal Weierstrass curve $E_m : y_m^2 + y_m = x_m^3 - 7$:*

$$x_m = \frac{-3z}{x+y}, \quad y_m = \frac{4x-5y}{x+y}, \qquad [x:y:z] = [y_m+5 : 4-y_m : -3x_m]. \tag{A.36.1}$$

It carries O to the Weierstrass origin and carries P, Q_c to $(3,-5),(3,4)$, respectively. All of $E_m[9]$ is defined over the unramified degree-three extension of $\mathbb{Q}_7$. Its reduction over $k = \mathbb{F}_7[\theta]/(\theta^3-2)$ has exactly 81 points. One point above P under multiplication by three is

$$Q = (2+\theta^2,\ 1+\theta+6\theta^2), \qquad \varphi Q - Q = (6,2), \tag{A.36.2}$$

where φ raises the residue coordinates to their seventh powers. The unique nine-torsion lifts give the corresponding identities over the unramified local extension.

For each of the nine points Q above either $d = P$ or $d = Q_c$, let $v_R = x_m - x_m(R)$ and let $\ell_{R,S}$ be the line through R, S, tangent at R when $R = S$, normalized to have y_m coefficient one. Then the actual function

$$a_Q = \frac{\ell_{Q,Q}\ell_{2Q,Q}}{v_{2Q}v_d}, \qquad \operatorname{div}(a_Q) = 3(Q) - (d) - 2(O), \tag{A.36.3}$$

defines a connected cubic cover $w^3 = a_Q$ which is finite étale over U. Its geometric character is an explicit Kummer class with nonzero cusp valuations at d, O. With $\iota(R) = -R$, the following function identities hold:

$$a_Q\, \iota^* a_Q = \frac{(x_m - x_m(Q))^3}{x_m - x_m(d)}, \qquad \prod_{i=0}^{2} a_{\varphi^i Q} = g_Q^3. \tag{A.36.4}$$

In the second identity, put $Q_i = \varphi^i Q$, $S = Q_0 + Q_1$, and

$$g_Q = \frac{\ell_{Q_0,Q_1}\ell_{S,Q_2}}{v_S v_d}. \tag{A.36.5}$$

This function is defined over $\mathbb{Q}_7$, and its reduction is defined over $\mathbb{F}_7$.

For the exact Fermat parameter $s = z/x$ and its unit tangent, the regularized constants are

$$(s^2 a_Q)(O) = 9, \qquad (s^2(x_m - 3))(O) = (s^2(x_m - 3\zeta^2))(O) = 9. \tag{A.36.6}$$

In the large model $Y^2 = X^3 - 432$, $X = 4x_m$, $Y = 8y_m + 4$, the corresponding normalized Miller function is $4a_Q$, with regularized constant 36. For the residue embedding $\zeta \mapsto 2 \in \mathbb{F}_7$, the cubic residue of 9 is $4 = \zeta^2$, whereas that of 36 is 1. These are arithmetic cover data; no central Galois action on an intrinsic Heisenberg quotient is inferred from them.

Proof. Substitution in (A.36.1) gives the identity

$$(x + y)^3(y_m^2 + y_m - x_m^3 + 7) = 27(x^3 + y^3 + z^3).$$

The displayed inverse gives the reverse identity. The rational maps extend across their omitted points because both projective curves are smooth. Their composites on the displayed dense affine opens are the identity, proving the claimed isomorphism and its stated cusp images. The discriminant of the minimal equation is -3^9, so it has good reduction at seven. Multiplication by nine is étale over $\mathbb{Z}_7$, and reduction identifies its nine-torsion over an unramified extension with the nine-torsion of the residue curve.

The exact program `a_miller_f343_root.py` constructs k, lists all points of $y_m^2 + y_m = x_m^3$, and applies the stated Weierstrass group law. There are 324 points, 81 killed by nine, and nine killed by three. Every three-torsion point has coordinates in $\mathbb{F}_7$. It lists all nine preimages of each specified cusp and verifies (A.36.2) and each Frobenius orbit. The geometric nine-torsion has order 81 because nine is invertible in characteristic seven; the enumerated points therefore exhaust it. Finite étale lifting proves the local assertion, rather than merely an assertion about an unrelated finite-field group.

The divisors of the normalized line and vertical functions are

$$\begin{aligned} \operatorname{div}(\ell_{R,S}) &= (R) + (S) + (-R - S) - 3(O), \\ \operatorname{div}(v_R) &= (R) + (-R) - 2(O). \end{aligned}$$

Their substitution proves (A.36.3). At every point of U the valuation is a multiple of three, so the normalized cubic cover is unramified there. Characteristic zero then makes it étale. Its valuation -1 at d proves that the function is not a cube and that the cover is connected. This gives an actual finite cover whose Kummer character is defined by the displayed function.

Writing $t = -x_m/y_m$ at the origin gives $x_m = t^{-2} + O(t)$, $y_m = -t^{-3} + O(1)$. Each normalized line has leading term $-t^{-3}$ and each vertical function has leading term t^{-2}. Consequently a_Q and g_Q have leading term t^{-2}. The divisors of the two sides of the first identity in (A.36.4) agree, and their leading coefficients at O are both one. Their ratio is therefore the constant one on the smooth proper curve, proving that identity.

For the second identity, write $D_Q = \varphi Q - Q$. The finite computation and finite étale comparison give $D_Q \in E_m[3](\mathbb{Q}_7)$. Thus $Q_i = Q + iD_Q$, and $Q_0 + Q_1 + Q_2 = 3Q + 3D_Q = d$. The line-divisor calculation in (A.36.5) gives

$$\operatorname{div}(g_Q) = (Q_0) + (Q_1) + (Q_2) - (d) - 2(O).$$

This divisor is fixed by φ. Its normalized leading coefficient is one, so $\varphi(g_Q) = g_Q$. The function descends to $\mathbb{Q}_7$. Comparing divisors and the same leading coefficient proves the cube identity. The finite-field program also multiplies the function numerators in the exact ring $k[x_m, y_m]/(y_m^2 + y_m - x_m^3)$ and verifies both identities by cross-multiplication, for all 18 points.

Finally on the Fermat chart $x = 1$, $y = -(1 + s^3)^{1/3}$ gives $x_m = 9s^{-2} + O(s)$ and $y_m = -27s^{-3} + O(1)$. Hence $s = 3t + O(t^4)$, which proves (A.36.6). The large-model line and vertical functions are eight and four times the corresponding minimal-model functions, so their two-line quotient is $4a_Q$. Reduction of the retained constants gives $9 \equiv 2$, $36 \equiv 1 \pmod 7$; the cubic residue map on $\mathbb{F}_7^\times$ is $u \mapsto u^2$. This proves the stated tangent phases without dropping the origin term. The program retains the characteristic-zero coordinate identity, tangent expansions, complete field coordinates, line coefficients, and all exact outputs. □

Statement A.37 (Proved). An order-eighty-one arithmetic quotient and a tame-letter Frobenius coefficient at seventeen. *Put $K_0 = \mathbb{Q}(\zeta)$, $\zeta^2 + \zeta + 1 = 0$, and $L_{17} = K_0(\beta)$ with $\beta^3 = 17$. Let $\sigma(\beta) = \zeta\beta$ and set*

$$\delta = \frac{-7 - \zeta - 3\beta - \beta^2}{3}, \qquad \delta_i = \sigma^i(\delta), \qquad r_i^3 = \delta_i \quad (i \in \mathbb{Z}/3). \tag{A.37.1}$$

The extension

$$M_{17} = L_{17}(r_0, r_1, r_2)/K_0 \tag{A.37.2}$$

is Galois, unramified outside three and seventeen, with group

$$G = (\mathbb{F}_3)^3 \rtimes \langle s \rangle, \qquad s^3 = 1, \tag{A.37.3}$$

where $s(r_i) = r_{i+1}$ and $s(\beta) = \zeta\beta$. For $a = (a_0, a_1, a_2)$ write $\tau_a(r_i) = \zeta^{a_i} r_i$, $\tau_a(\beta) = \beta$. Then $s\tau_a s^{-1} = \tau_{Sa}$, where $(Sa)_i = a_{i-1}$. The group has order 81, lower-central orders $81, 9, 3, 1$, exponent nine, and trivial fourth three-Zassenhaus subgroup.

Use the arithmetic generators m, n, t of A.28 and A.31, with the exact Kummer normalization of A.35. Their modulo-three characters are

	m	n	t
κ_ζ	1	0	0
κ_{17}	0	0	1

. (A.37.4)

The restriction map to (A.37.2), composed with the actual free pro-three presentation, is a continuous epimorphism

$$F_3(m, n, t) \longrightarrow G_{K_0,\{3,17\}}(3) \longrightarrow G. \tag{A.37.5}$$

Let $W \in D_3(F_3(m, n, t))$ be the actual arithmetic Frobenius word of A.30. Under (A.37.5),

$$W \longmapsto \tau_{(2,2,2)}, \qquad [t, [t, m]] \longmapsto \tau_{(1,1,1)}. \tag{A.37.6}$$

In the restricted degree-three Lie algebra $D_3(F_3)/D_4(F_3)$, this map annihilates the three generator cubes and every ordinary degree-three component except the multidegree $(1, 0, 2)$

component generated by $[t,[t,m]]$. Consequently the coefficient of this bracket in the full W is exactly $2 \in \mathbb{F}_3$. This records a coefficient involving tame inertia; the map killing t in A.31 does not retain it.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379); Tasks A.6 and B.4.

Proof. For $a,b,c \in K_0$ direct multiplication of the three conjugates gives

$$N_{L_{17}/K_0}(a+b\beta+c\beta^2) = a^3+17b^3+289c^3-51abc. \tag{A.37.7}$$

At $(a,b,c)=(-7-\zeta,-3,-1)$ the right side is 27ζ. Thus

$$\delta_0\delta_1\delta_2 = \zeta. \tag{A.37.8}$$

Each δ_i is integral away from three. Since its norm is a unit, it is a unit at every finite prime away from three. Kummer extensions by its cube root are unramified at such primes, because three is invertible there and the radical is a unit. The extension L_{17}/K_0 is ramified only at three and seventeen. This proves the asserted ramification bound once the degree and Galois group are established.

The three classes of δ_i are independent in $L_{17}^{\times}/L_{17}^{\times 3}$. Indeed the map

$$\mathbb{F}_3[\langle\sigma\rangle] \longrightarrow L_{17}^{\times}/L_{17}^{\times 3}, \qquad \sigma^i \longmapsto [\delta_i] \tag{A.37.9}$$

is a module homomorphism. Every nonzero submodule of the regular module $\mathbb{F}_3[u]/(u-1)^3$ contains its socle, generated by $1+u+u^2$. By (A.37.8) that element maps to $[\zeta]$. The latter is nonzero: adjoining a cube root of ζ gives $K_0(\mu_9)$, a cubic extension unramified at seventeen, whereas L_{17}/K_0 is Eisenstein at the unique prime over seventeen. These two cubic extensions therefore have different ramification indices and cannot coincide. Thus the kernel in (A.37.9) is zero.

Kummer theory now gives $\mathrm{Gal}(M_{17}/L_{17}) = (\mathbb{F}_3)^3$ with the displayed diagonal actions. Independence also proves that the defining radical algebra is a field of degree 27 over L_{17}. Cyclic permutation of its generators defines the specified extension s of σ, with $s^3=1$. This proves the semidirect presentation (A.37.3). Its commutators are

$$\gamma_2 G = (S-1)(\mathbb{F}_3)^3, \quad \gamma_3 G = (S-1)^2(\mathbb{F}_3)^3 = \langle \tau_{(1,1,1)} \rangle, \quad \gamma_4 G = 1. \tag{A.37.10}$$

The three dimensions are $2,1,0$. Every cube in G is diagonal central, while $G^9=1$ and $(\gamma_2 G)^3 = 1$. The defining formula $D_j(G) = \prod_{i3^a \geq j} (\gamma_i G)^{3^a}$ therefore gives $D_4(G)=1$.

We next compute the actual local Frobenius. The residue field of K_0 at seventeen is $k = \mathbb{F}_{289} = \mathbb{F}_{17}[\zeta]$. At its unique extension to L_{17} the radical β reduces to zero, so every δ_i has the same residue

$$d_0 = (-7-\zeta)/3 = 9+11\zeta, \qquad d_0^3 = \zeta, \qquad d_0^{96} = \zeta^2. \tag{A.37.11}$$

In particular d_0 is not a cube in k. The polynomial $R^3 - d_0$ defines the unramified residue extension of degree three. Its roots obey

$$R^{289} = d_0^{96} R = \zeta^2 R. \tag{A.37.12}$$

An arithmetic Frobenius fixing the chosen uniformizer roots fixes β and acts by $r_i \mapsto \zeta^2 r_i$ for all three radicals. To see that the residue calculation determines this exact automorphism, the ratio of any two possible images of r_i is in μ_3, whose reduction is injective at seventeen. This

proves the first formula in (A.37.6). It is independent of conjugating the chosen decomposition subgroup because this image is central.

Finally put $v = r_0 r_1 r_2$, so $v^3 = \zeta$. For an element $\tau_a s^k$, the two characters $\kappa_\zeta, \kappa_{17}$ are respectively $a_0 + a_1 + a_2$ and k. Thus the actual images of the normalized generators have the form

$$m \mapsto \tau_a, \qquad n \mapsto \tau_b, \qquad t \mapsto \tau_c s, \qquad \sum a_i = 1, \quad \sum b_i = \sum c_i = 0. \tag{A.37.13}$$

We have made no choice of their individual coordinates. All three images have cube one. Conjugation by $\tau_c s$ on the abelian kernel is S, and

$$(S-1)^2 a = (1 + S + S^2)a = (1, 1, 1), \qquad (S-1)^2 b = 0. \tag{A.37.14}$$

Every bracket with two entries from the abelian kernel vanishes. The only surviving ordinary degree-three bracket is therefore $[t, [t, m]]$; its image is the second formula in (A.37.6). The standard free restricted degree-three basis consists of the ordinary degree-three free Lie part and the three generator cubes. The preceding calculations identify the entire map on that basis, so the Frobenius image proves that its indicated coefficient is two. This argument uses the full three-generator word and the actual arithmetic restriction (A.37.5).

For comparison with the lower Heisenberg test, the fixed field of the diagonal subgroup in (A.37.10) is

$$L_{17}(v, w), \qquad v^3 = \zeta, \quad w = r_0^2 r_1, \quad w^3 = \delta_0^2 \delta_1. \tag{A.37.15}$$

The two character rows $(1, 1, 1)$ and $(2, 1, 0)$ are independent, so this field has degree nine over L_{17}, hence degree 27 over K_0. The action satisfies $s(w) = (v/\delta_0)w$. Its weighted radical has residue $d_0^3 = \zeta$, which already has the cube root d_0 in k. The Frobenius image computed above becomes the identity in this quotient, exactly as follows from its diagonal kernel. This specifies both the lower quotient map and the coefficient it forgets.

The exact program `a_wreath17_root.py` verifies (A.37.7), the residue field and its actual degree-three radical extension, all 81 group elements, the lower-central orders, and every one of the 729 triples in (A.37.13). The preliminary norm search and its exact algebraic certification are retained in `a_norm_zeta_17.py`. □

2 B. Geometric classes and centralizers

2.1 Geometric classes and the cyclotomic centralizer

We retain the tangent at O with parameter z/x and tangent one. All inertia groups in this section are their images in the geometric pro-3 group N. We write κ for continuous pro-3 Kummer classes. A cuspidal section means a section contained in the decomposition group of a lifted cusp, with geometric conjugacy taken in the group actually displayed. The manuscript comparison is §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379).

Statement B.1 (Proved). The three abelian cusp packets, with their framed origins. *Put $J = J_{E,D}$ and use the frame coordinates of LM7 (E24.4). Let z_P be the framed line bundle $\mathcal{O}(P - O)$ with frames $(x/z)^{-1}$ at P, z/x at O, and the canonical section at the other cusps. Define z_Q in the same way from $Q_c - O$, using y/z at Q_c. Put $\alpha_P = \kappa(z_P)$, $\alpha_Q = \kappa(z_Q)$ and $\alpha_O = 0$. The three packets in $H^1(G_\mathbb{Q}, V)$ are*

$$\mathscr{P}_d^V = \alpha_d + e_d\, \mathcal{R}_\mathbb{P}(\mathbb{Z}_3), \qquad d = O, P, Q_c. \tag{B.1}$$

Here $e_d\, \mathcal{R}_\mathbb{P}(\mathbb{Z}_3)$ denotes the image of the continuous homomorphism on cohomology induced by the $G_\mathbb{Q}$-equivariant inclusion $\mathbb{Z}_3(1) \to V$, $1 \mapsto e_d$.

The torus-point relation specifying the second framed origin is

$$z_P + z_Q = (3^{-1}, 3^{-1}, -\lambda^{-1}, -\lambda^{-1}, \zeta\lambda^{-2}). \tag{B.2}$$

In the presentation (E24.7), eliminate the first λ-generator and retain z'. Write g_2, g_3 for the remaining two λ-generators and τ_3 for ζ in the third quadratic factor. After pro-3 completion the origins have toric coordinates zero, divisible-cohomology coordinate zero, and remaining coordinates

$$\alpha_P = \kappa(z'), \qquad \alpha_Q = \kappa(2z' + g_2 + g_3 - \tau_3). \tag{B.3}$$

The inertia maps in the original frame coordinates are

$$\begin{aligned} i_O(u) &= (u^{-1}, u^{-1}, u^{-1}, u^{-1}, u^{-1}), \\ i_P(u) &= (u, 1, 1, 1, 1), \qquad i_Q(u) = (1, u, 1, 1, 1). \end{aligned} \tag{B.4}$$

Their images under the two coordinate-unit characters are respectively $(-\kappa u, -\kappa u)$, $(\kappa u, 0)$ and $(0, \kappa u)$. The three sets in (B.1) are pairwise disjoint, and each displayed parameter map is injective.

Proof. The tangential Albanese class at d, relative to the tangent at O, is represented in the generalized Picard group by $\mathcal{O}(d - O)$ with its local parameter frames. This description can be obtained directly from the divisor construction: replace d and O by points with local parameters s_d and s_O, take the line bundle of their difference, and divide its local frames by the leading powers s_d^{-1} and s_O. The resulting fibers at the reduced boundary are the displayed frames. Applying multiplication by 3^n identifies the finite abelian path torsor with the torsor of 3^n-division points of this framed line bundle. The inverse limit therefore gives its Kummer class in $T_3 J = V$. Changing the nonzero tangent at d multiplies its local root system by the root system of a scalar u and changes the abelian cocycle by $e_d\kappa(u)$. The same description applies to an arbitrary continuous section of the cusp decomposition extension, because that extension is $\mathbb{Z}_3(1) \rtimes G_{\mathbb{Q}}$ and its section classes are $H^1(G_{\mathbb{Q}}, \mathbb{Z}_3(1))$.

We verify the origin calculation. The rational function

$$h = \frac{x + y + z}{x + y}$$

has divisor $P + Q_c - 2O$. Its leading values are -1 after division by x/z at P, -1 after division by y/z at Q_c, and -3 after multiplication by $(z/x)^2$ at O. On the representatives $[0 : -\zeta : 1]$, $[-\zeta : 0 : 1]$ and $[1 : -\zeta : 0]$ its values, normalized by -3, are $-\lambda^{-1}, -\lambda^{-1}, \zeta\lambda^{-2}$. These are exactly (B.2). The expansions follow, for example, from $y/z = -(1 + (x/z)^3)^{1/3}$ near P. The relation $3 = -\zeta^2\lambda^2$ verifies the three quadratic values without choosing complex approximations.

For comparison, multiplication by $(x + z)/(x + y)$ trivializes $\mathcal{O}(3Q_c - 3O)$ and gives

$$3z_Q = (3^{-1}, -3^{-2}, -\lambda^{-2}, -\lambda^{-1}, \zeta\lambda^{-3}). \tag{B.5}$$

Together with (E24.6), this checks the cube of (B.2) in all five coordinates. The program `code/bc_checks.py` verifies all these identities over the exact field $\mathbb{Q}[\zeta]$.

Signs disappear in pro-3 Kummer theory: -1 has compatible rational 3^n-th roots. Relation (E24.7) then reads $3z' = -g_1 - 2g_2 - 3g_3 + 2\tau_3$. The quadratic coordinates in (B.2) are $-g_1 - g_2 - 2g_3 + \tau_3$. Subtracting z' and eliminating g_1 gives $2z' + g_2 + g_3 - \tau_3$, proving (B.3). The last factor of (E24.1) is zero because these are Kummer classes of actual rational generalized-Jacobian points; we use the splitting there with its rational-point completion as kernel.

Normalizing the frame at O to one gives (B.4). Applying (E24.4) gives the three asserted toric images; the quadratic norm of a rational scalar is its square. These images prove injectivity of each parameter map. Filling all cusps gives the continuous $G_{\mathbb{Q}}$-equivariant quotient $V \twoheadrightarrow T_3E$. It sends the packets to the point classes of O, P, Q_c, respectively. Their pairwise differences have exact order three. Reduction at two embeds rational odd torsion into $E(\mathbb{F}_2)$, which has order three, as checked by the program. Consequently none of these nonzero differences is three-divisible in $E(\mathbb{Q})$, and its modulo-three Kummer class is nonzero. This proves disjointness. □

Statement B.7 (Proved). Every nontrivial tame inertia in a full section is cuspidal. *Let $q \neq 3$ be a rational prime, put $k = \mathbb{Q}_q$, and use the actual pro-three fundamental group N of the marked Fermat cubic with the integral unit tangent at O as origin. For a continuous section $s : G_k \to N \rtimes G_k$, its restriction to inertia is either trivial or has image in one cuspidal inertia subgroup. In the latter case the whole section is contained in that cusp's decomposition group. The cusp is k-rational and the lifted cusp is unique. Moreover, a full section is unramified at q exactly when its image in V is unramified there.*

Proof. The marked curve and the tangent have good reduction at q. Thus the original based action of inertia on N is trivial, by prime-to-q specialization. Its cocycle restriction is a homomorphism. Wild inertia, being pro-q, maps to one; the only tame factor which can map to N is $\mathbb{Z}_3(1)$. Let $t = s(\tau_q)$ denote the geometric value on a generator of that factor, and assume $t \neq 1$. If $F = s(\phi_q)$ is the full Frobenius lift, the section relation is the exact group equality

$$FtF^{-1} = t^q. \tag{B.7.1}$$

Take a characteristic open normal subgroup M of N and put $H = M\overline{\langle t \rangle}$. Such M form a cofinal family, since N is finitely generated. Equation (B.7.1) and the characteristic property of M show that H is stable under $s(G_k)$. The open arithmetic subgroup $H\, s(G_k)$ is therefore the subgroup of the pro-three arithmetic fundamental group defining a finite etale cover $Y_H \to U_k$. Its geometric group is H, by the exact finite-three-group cover comparison proved in B.4.

We justify the reduction property needed for weights. The subgroup $H\, s(G_k)$ contains the original arithmetic inertia: its geometric cocycle values are powers of t, which lie in H. Original inertia centralizes N, so it acts trivially on every coset in N/H. The cover consequently descends to the smooth marked integral model by prime-to-q specialization. Its compactification is smooth and proper over the valuation ring. Locally at a boundary point this last assertion is the tame normalization $z^e = x$ after an unramified extension, with e prime to q; away from the boundary it is etale. These charts are exactly the tame-cover specialization comparison, also obtained from the logarithmic comparison in Stix, Theorem 3.4.6 [Sti02].

Let X_H be this compactification. The boundary sequence (B.8), with $\mathbb{Z}_3$ coefficients, is equivariant for the actual arithmetic Frobenius F of this descended cover. The image of t in $T_3J_{X_H}$ satisfies $Fv = qv$ by (B.7.1). All Frobenius eigenvalues on this Tate module have complex absolute value $\sqrt{q}$, by Deligne, Theorem (1.6) [Del74], applied in degree one to the smooth proper special fiber. The Tate module of its Jacobian has these same eigenvalues for arithmetic Frobenius, by the Abel–Jacobi comparison and the dual Weil pairing. Thus $v = 0$. The image of t in H^{ab} is in the boundary submodule.

The cyclic finite three-group H/M is generated by the image of t. Its homomorphism from H^{ab} consequently maps the boundary submodule onto all of H/M. At least one individual cusp inertia maps onto this cyclic group: otherwise every such image would be in its unique maximal subgroup. It follows that t fixes a cusp in the boundary set of the normal cover

associated to M. The inverse images of these finite fixed-cusp sets in $\coprod_d N/I_d$ are nonempty closed subsets with the finite-intersection property. Compactness gives a lifted cusp fixed by t at every level, so t belongs to its actual inertia group $I_{\widetilde{d}}$.

Distinct cusp inertia groups have trivial intersection by the boundary and cover argument in B.4. Hence this lifted cusp is unique. Equation (B.7.1) shows that F normalizes $I_{\widetilde{d}}$, because its image has the nontrivial element t^q in common with that group. Its normalizer is the cusp decomposition group, again by the explicit stabilizer argument in B.4. The inertia image and F topologically generate the image of the local section; therefore that whole image lies in the decomposition group. Its projection is all G_k, forcing $d \in D(k)$.

Finally the inclusion $I_{\widetilde{d}} \to V$ sends its generator to the primitive nonzero boundary vector e_d. It is injective. Thus $t \neq 1$ has nonzero abelian image, while a trivial full inertia image certainly has trivial abelian image. This proves the last assertion. □

Statement B.8 (Proved). The exact local image of ramified full sections in class two. *With $k = \mathbb{Q}_q$, $q \neq 3$, let $A = N/\gamma_3 N$. For each $d \in D(k)$ choose the coordinate unit tangent used in C.10, its actual section $s_{d,1}$, and a peripheral generator c_d based at O. The ramified part of the image of full sections in* $\mathrm{Sec}_k(A)$ *is precisely the disjoint union*

$$\coprod_{d \in D(k)} \left\{ \left[g \longmapsto c_d^{\kappa_q(u)(g)} s_{d,1}(g) \right] : u \in k^{\times \wedge_3},\ v_q(u) \neq 0 \right\}. \tag{B.8.1}$$

Here the expression is multiplication in the arithmetic group, followed by its actual class-two quotient. Its parameter space is explicitly

$$k^{\times \wedge_3} = \mathbb{Z}_3 \kappa(q) \oplus \mu_{3^{v_3(q-1)}}(k). \tag{B.8.2}$$

There are nine packets for $q \equiv 1 \pmod 3$ and three for $q \equiv 2 \pmod 3$. In the central cuspidal lattice $I/[N,I] = \mathbb{Z}_3(1)[D]$ the tame vector of a packet is $v_q(u)c_d$, with exactly this one cusp coordinate nonzero. In particular the mixed-inertia counterexamples in A.5 and A.6 cannot lift to full N.

Proof. The decomposition extension for a rational cusp is $\mathbb{Z}_3(1) \rtimes G_k$ using its unit tangent. Its sections modulo inertia conjugacy are the Kummer group (B.8.2). The decomposition inclusion is a continuous homomorphism of actual arithmetic groups; after quotienting the geometric group it gives exactly (B.8.1). The Kummer cocycle equation and $s_{d,1}(g) c_d s_{d,1}(g)^{-1} = c_d^{\chi_3(g)}$ verify directly that the displayed product is a section. The unit-tangent section is unramified, and the tame Kummer exponent is $v_q(u)$, giving the stated ramification condition. Every displayed class has its full cuspidal lift. Conversely B.7 puts every ramified full section in one of these decomposition extensions.

To check that the displayed parametrization remains injective in class two, first pass to a finite unramified extension splitting all cusps. The boundary inclusion $e_d : \mathbb{Z}_3(1) \to W$ is then split, and $H^1(k,W) \to H^1(k,V)$ is injective because $T_3 E^{G_k} = 0$. Restriction on the Kummer group is injective by the same invariant argument as C.10. Thus two parameters at the same cusp cannot coincide. Different cusps map in the proper elliptic quotient to the Kummer classes of different three-torsion points. Reduction at $q \neq 3$ injects three-primary torsion into the finite group $E(\mathbb{F}_q)$; its nonzero elements are not divisible by every power of three. The formal group is pro-q, on which multiplication by three is invertible. Hence a nonzero difference of two cusp points has nonzero integral local Kummer class, proving disjointness.

The decomposition of $k^\times$ into a uniformizer, the roots of unity of order $q-1$, and principal units gives (B.8.2), since the principal units are pro-q. The two possible cusp counts follow

from the actual field of the six nonrational cusps, $\mathbb{Q}(\zeta_3)$. Projection to $I/[N,I]$ sends the peripheral generator to its indicated basis element, proving the tame-vector formula. The vectors in A.5 and A.6 involve $c_O + c_P$ and therefore do not have this form. □

Statement B.9 (Proved). All inherited class-two cusp origins. *Use exactly the diagonal coordinates of LM7 E.29 for H, based at the lifted cusp P over $t = 0$, and put*

$$f(g) = \frac{\chi_3(g)^2 - 1}{24} \in \mathbb{Z}_3(2). \tag{B.9.1}$$

The three rational cusp packets in $\mathrm{Sec}(H)$ *have the following continuous cocycle representatives:*

	$u(g)$	$v(g)$	$w(g)$
P	$\kappa(a)(g)$	0	0
Q_c	0	$\kappa(a)(g)$	$f(g)$
O	$-\kappa(a)(g)$	$-\kappa(a)(g)$	$f(g)/2$

$$a \in \mathbb{Q}^{\times \wedge 3}. \tag{B.9.2}$$

Each row is injectively parametrized and the rows are pairwise disjoint. Above inherited abelian class $(0,0)$ *the three unit tangents give, respectively, the central labels* $0, 2, 1 \in \mathbb{Z}/3$ *relative to* $g \mapsto (\chi_3(g)^2 - 1)/3$. *The actual arithmetic map from the intrinsic group based at* O *to this* H*-extension is*

$$(n, g) \longmapsto \big(q_O(n)(0, 0, f(g)/2), g\big), \tag{B.9.3}$$

after the geometric path choices in the proof. Thus (B.9.2) retains the change of section origin between the O *and* P *base points.*

Proof. Write $t = -u_0^3$, $1 - t = -v_0^3$ for the cubic cover of the tripod, with $u_0 = x/z$, $v_0 = y/z$. The standard path from the unit tangent at zero to the unit tangent at one lifts from P to Q_c: along the real interval its cube roots are $u_0 = -t^{1/3}$ and $v_0 = -(1-t)^{1/3}$. It reaches exactly the displayed rational cusps. At either end the lifted local parameter has tangent minus one relative to u_0 or v_0. Changing that sign has zero pro-three Kummer class, with the explicit compatible rational roots $(-1)^{1/3^n} = -1$. The local decomposition-section comparison therefore changes no section class.

Let X, Y be the logarithms of the standard tripod loops and $C = [X, Y]$. The H lattice is $\mathbb{Z}_3(3X) \oplus \mathbb{Z}_3(3Y) \oplus \mathbb{Z}_3 C$. Wickelgren's evaluation of the standard path cocycle in the subsection "Evaluating the 2-nilpotent quotient of $\mathfrak{f}$", equation $f(\sigma) = (\chi(\sigma)^2 - 1)/24$ [Wic12], says that the section at Q_c has central cocycle fC relative to P. The sign follows from the actual action convention $g(y) = \mathfrak{f}_g^{-1} y^{\chi(g)} \mathfrak{f}_g$: the lift $\mathfrak{f}_g s_P(g)$ conjugates y to its $\chi(g)$-th power. Thus the P, Q_c origins and their peripheral generators are $0, fC$ and $3X, 3Y$.

Consider the rational coordinate interchange $(x, y, z) \mapsto (y, x, z)$ on the cubic. It covers $t \mapsto 1 - t$, interchanges P, Q_c, and fixes O. Transporting the fundamental group along the same standard path gives an actual automorphism of the arithmetic H-extension. Its geometric map swaps the two abelian coordinates and negates C, and its change of arithmetic origin is fC. There are no additional linear central terms in its geometric map: because fC is central, the geometric map commutes with the diagonal $G_{\mathbb{Q}}$-action; a term from weight one to weight two would satisfy $(\chi - \chi^2)c = 0$ and therefore is zero in the torsion-free lattice. Consequently the exact automorphism is

$$((u, v, w), g) \longmapsto ((v, u, -w + f(g)), g). \tag{B.9.4}$$

Its formula also checks directly that it is a continuous group homomorphism and that its square is the identity. It preserves H because the covering subgroup is the kernel of the two abelian coordinates modulo three.

At O, use the local parameter $s = z/x$. The regularized leading terms of u_0, v_0 are $1, -1$; thus the inherited abelian class of the unit tangent is zero. Conjugating by an element of H makes its abelian cocycle identically zero. Its remaining class is some $h \in H^1(G_{\mathbb{Q}}, \mathbb{Z}_3(2))$. The coordinate interchange takes s to z/y, whose leading coefficient at O is -1. The rational root comparison above makes these two pro-three tangent sections conjugate. Since (B.9.4) acts on central classes by $h \mapsto -h + [f]$, we obtain $2h = [f]$. Multiplication by two is invertible on this group, so $h = [f/2]$. A central geometric conjugation then chooses the representative $f/2$.

We also check the inertia generator in this gauge. Its abelian coordinates are $(-1, -1)$, since both u_0 and v_0 have a simple pole in s. Write its central coordinate as w_0. The section just chosen is central, so its cyclotomic normalization of the inertia generator and the diagonal action give $\chi(g)^2 w_0 = \chi(g) w_0$ for every g. The lattice is torsion-free and the character is nontrivial, hence $w_0 = 0$. This determines the actual inertia image and avoids an unsupported choice of a conjugate of $(xy)^{-3}$. Twisting each of the three sections by its Kummer parameter now gives (B.9.2), using the displayed group law of E.29. For each row the two abelian entries are proportional, so that its BCH cross term is zero. Equation (B.9.3) is the arithmetic map with exactly the calculated image of the intrinsic reference section; centrality of that image makes its geometric part q_O equivariant for the stated diagonal action.

Finally $H^1(G_{\mathbb{Q}}, \mathbb{Z}_3(2)) = \mathbb{Z}/3$ with generator $(\chi_3^2 - 1)/3$ by E.29m. The coefficients $1/8$ and $1/16$ reduce to 2 and 1 modulo three. The three abelian packet lines meet only at zero, and their central classes there are distinct. This proves all injectivity and disjointness assertions. The exact rational-arithmetic program `code/bc_h_packets.py` verifies the BCH automorphism, the three cocycle identities and the three distinct central labels; its output is included alongside the code. □

Statement B.10 (Proved). The full global image at the specified prime seven. *Let $A = N/\gamma_3 N$, and use the normalized pair coordinates (a, b) and the A^c-action of A.15 for sections supported at $\{3, 7, \infty\}$. Such a class has a full N-lift exactly in the following cases:*

$$b = 1, \quad \text{or its restriction to } G_{\mathbb{Q}_7} \text{ belongs to one of the nine ramified packets (B.8.1).} \tag{B.10.1}$$

Equivalently these are precisely the classes which lift to every lower-central quotient. A full lift can be chosen unramified outside $\{3, 7, \infty\}$.

Here is an exact packet parametrization in the pair coordinates. Fix the embedding $\zeta_3 \equiv 2 \pmod 7$, a Frobenius ϕ fixing compatible roots of 7, and the actual unit-tangent values $g_d = s_{d,1}(\phi) \in A$. Use the same based inertia element $c_d \in A$ in each case. The ramified part of the global image is

$$\left\{ [(c_d^{2j} g_d, c_d^v) \cdot h] : d \in D(\mathbb{Q}_7), \quad v \in \mathbb{Z}_3 \setminus \{0\}, \quad j \in \{0, 1, 2\}, \quad h \in A/A^c \right\}. \tag{B.10.2}$$

Here $(a, b) \cdot h = (h^{-1} a F(h), h^{-1} b h)$, so A/A^c means right cosets, and the displayed parameters are injective after these explicit choices. The symbols g_d retain the actual tangential Frobenius origins; they are not replaced by arbitrary solutions of a graded normalization equation.

Proof. The exact isomorphism of arithmetic groups in A.14 and the normalized-cocycle construction in A.15 identify every global class with the asserted pair. If $b = 1$, lift a arbitrarily to $\widetilde{a} \in N$ and take $\widetilde{b} = 1$. The full tame equation holds, and A.15 extends this pair to the required global section. If $b \neq 1$ and a full section exists, B.7 forces its local restriction to be cuspidal. Conversely a local cuspidal class in (B.8.1) has its actual full local lift. Conjugate this lift by a lift in N of the geometric element comparing its class-two image to the specified pair (a, b). Its generator values then project to that exact pair. A.15

extends them, with their conjugate values on the second local factor, to a full normalized global section. This proves both directions of (B.10.1), including the ramification support. The nilpotent assertion follows either from A.15's compactness proof or from B.11 below.

For (B.10.2), the completed local Kummer group is $\mathbb{Z}_3\kappa(7) \oplus (\mathbb{Z}/3)\kappa(\zeta_3)$. Choose compatible roots of ζ_3. Its unramified cocycle is $(\chi_3 - 1)/3$, which has value two at ϕ and value zero on tame inertia. The uniformizer cocycle has values zero and one there. The full decomposition-group formula of B.8 therefore gives exactly the pair $(c_d^{2j} g_d, c_d^v)$. Taking $j + 3$ instead changes the pair by conjugation with c_d, so the three chosen representatives cover precisely the unit parameter group.

Different d, v, j have different local section classes by B.8. Within each local class the geometric centralizer in A is trivial: its abelian image would be fixed by Frobenius on V, and its remaining central image would be fixed on $\bigwedge^2 V$; the respective eigenvalue absolute values are $\sqrt{7}, 7$ and $7, 7^{3/2}, 49$, all greater than one. Thus the full A-conjugacy orbit of that local pair is freely parametrized by $h \in A$. The normalized global conjugacies are exactly A^c by A.15, acting on the right. This proves the stated right-coset parametrization and its injectivity. □

Statement B.11 (Proved). No new ramification above class two and compactness for a finite set of primes. *Let S be any finite set of rational places containing 3 and infinity, and let s_2 be a class-two section unramified outside S. Every lift of s_2 to $N/\gamma_r N$, $r \geq 3$, is unramified outside the same set S. Moreover the following assertions are equivalent:*

$$\begin{gathered} s_2 \text{ lifts to } N; \\ s_2 \text{ lifts to } N/\gamma_r N \text{ for every } r \geq 3. \end{gathered} \tag{B.11.1}$$

The separately given nilpotent lifts need not be compatible. The conclusion supplies a compatible continuous full lift, also unramified outside S.

Proof. For each r, iterated commutators give a continuous $G_{\mathbb{Q}}$-equivariant surjection

$$V^{\otimes r} \longrightarrow \gamma_r N/\gamma_{r+1} N. \tag{B.11.2}$$

The target is a finite free $\mathbb{Z}_3$-module. This integral assertion follows by Hall collection in the free pro-three group: basic commutators of weight r span the quotient, and their Magnus expansions have the linearly independent leading Hall Lie monomials. Collection at finite exponent quotients, followed by compactness, gives the same spanning assertion over $\mathbb{Z}_3$. The leading monomials prove independence and exclude torsion. Thus (B.11.2) is the actual integral graded comparison being used.

For $q \neq 3$, Frobenius eigenvalues on V have weights one and two: on the elliptic factor their absolute values are $\sqrt{q}$ and on the boundary factor they are q. The eigenvalues on (B.11.2) are therefore among products of r such eigenvalues. For $r \geq 3$ their absolute values are at least $q^{r/2} > q$. This eigenvalue conclusion is valid for a quotient of the tensor representation and does not assume semisimplicity of the extension defining V.

Suppose a lift from class $r-1$ to class r acquires inertia at a prime outside S. Its geometric inertia values lie in the central kernel $\gamma_r N/\gamma_{r+1} N$ and, as in B.7, factor through the tame three-part. The exact tame relation requires its value t to satisfy $F_q t = qt$; conjugation by the section's Frobenius value acts trivially on this central kernel. For $r \geq 3$ the preceding eigenvalue calculation and torsion-freeness give $t = 0$. Induction starting at the fixed class-two section proves that every higher nilpotent lift has the asserted support.

Let $K_0 = \mathbb{Q}(\zeta_3)$ and let S' be its places over S. The relative pro-three quotient Γ_S of $G_{\mathbb{Q},S}$ has kernel $G_{K_0,S'}(3)$ and quotient of order two. That pro-three kernel is finitely generated:

its $\mathbb{F}_3$-characters form the finite Kummer group of S'-units and the finite three-primary ideal class group. The finite-dimensional Frattini quotient generates a pro-three group, by the same finite-quotient maximal-subgroup argument used in A.14. Thus Γ_S is finitely generated. The actual geometric action factors through it, as proved in A.15 from the three-group action on the Frattini quotient of N. Every cocycle under consideration also factors through Γ_S: over K_0 its graph is a homomorphism to a pro-three semidirect product.

Fix generators $g_1, \dots, g_m$ of Γ_S and a cocycle representative of s_2 on it. Consider the compact space of tuples in N^m whose projections are its fixed generator values. Choose a free profinite presentation on these generators, with its closed subgroup of relators. For each $r \geq 3$, impose all those relators on the lifted tuples in $(N/\gamma_r N) \rtimes \Gamma_S$. These are closed conditions: at any finite quotient each profinite word is a continuous function of the generator values. A lift at level r makes the resulting closed subset nonempty, after conjugating its projection to the fixed representative; the required geometric conjugator lifts because $N/\gamma_r N \to N/\gamma_3 N$ is surjective. The closed sets are nested. Compactness supplies one tuple satisfying every level of every relator.

The intersection $\bigcap_r \gamma_r N$ is trivial: a nontrivial element survives in a finite three-group, and that group is nilpotent. Thus the tuple satisfies all relators in $N \rtimes \Gamma_S$ itself and defines a continuous section. Composing with $G_{\mathbb{Q},S} \to \Gamma_S$ gives the desired full cocycle with the prescribed class-two projection. This proves the nontrivial implication in (B.11.1); the other is induced by the continuous quotient maps. □

Statement B.2 (Refuted). The inherited class-two image need not be a point or cusp image. *The exclusion proposed in Lead 1 is false. There are sections of $N \rtimes G_{\mathbb{Q}}$ whose image in* $\mathrm{Sec}(H)$ *has toric coordinates $(\kappa(3), 2\kappa(3))$. Every such image lies outside the three rational cusp packets, and $U(\mathbb{Q})$ has no point producing it. It therefore supplies the asserted counterexample already with a full geometric pro-3 lift.*

Proof. We use exactly the onto map in LM7 E.34: the coordinate-unit homomorphism on the unramified group $H^1(G_{\mathbb{Q},\{3,\infty\}}, V)$ has image $\mathbb{Z}_3\kappa(3)^2$, and every class in this source lifts to the full N. Its projectivity step is checked as follows. Over $\mathbb{Q}(\zeta)$, the maximal pro-3 Galois group unramified away from three is free on two generators: Kummer theory gives the two classes $\zeta, 1-\zeta$; the class group of $\mathbb{Z}[\zeta, 1/3]$ is zero, and the three-primary Brauer group supported at its single omitted finite place is zero by the sum of local invariants. Thus degree-two cohomology with $\mathbb{F}_3$ vanishes. The free-pro-3 criterion follows by solving finite central embedding problems. Complex conjugation has order two, so averaging gives the same degree-two vanishing for the relative semidirect product. The geometric action and every abelian cocycle factor through this relative group: their restrictions over $\mathbb{Q}(\zeta)$ take values in pro-3 groups. Successive finite geometric embedding problems therefore have solutions; finite generation of the relative group and compactness make them compatible. This is the full-group lifting step used from E.34. The direct-summand coordinate-unit map is the explicit algebraic section $(a, b) \mapsto (a, b, 1, 1, 1)$ of (E24.4), so its onto property requires no further lifting assertion. These arguments also agree with the free-Galois-group mechanism in Hoshi, Theorem 3.5 [Hos10]; that theorem concerns the specified proper Jacobian sections, while the preceding relative-group argument is the needed one for all ten coordinates here.

By (B.4), the cusp projections are $(c, 0)$, $(0, c)$ and $(-c, -c)$, with $c \in H^1(G_{\mathbb{Q}}, \mathbb{Z}_3(1))$. The valuation at three sends the proposed vector to $(1, 2)$, which lies on none of the three lines, already over $\mathbb{F}_3$. The code verifies this finite separation. Conjugation in H does not alter the abelian cohomology class, so no choice of central lift can change this conclusion. LM7's computation $E(\mathbb{Q}) = \{O, P, Q_c\}$ gives $U(\mathbb{Q}) = \varnothing$.

The drafter's suggested vector $\kappa(3)(e_O + e_P)$ does have an intrinsic abelian image outside the packets (B.1): its elliptic image is zero and its toric image is not diagonal. Its inherited toric image is $(0, -\kappa(3))$, on the Q_c line. That particular toric calculation therefore gives no exclusion in $\mathrm{Sec}(H)$. The vector used above supplies the required exclusion there. □

Statement B.3 (Proved). The cusp packets in the cuspidally central quotient. *In $N^{\mathrm{cc}} = C \times_T V$, let the generator of the central $\mathbb{Z}_3(1)$ of C be the elliptic commutator $[a, b]$. Choose the reference C-section above the elliptic class zero to be the tangent at O; above $\kappa(P - O)$ and $\kappa(Q_c - O)$ choose the images of the respective unit tangents. Use these references for every abelian class in the corresponding three elliptic fibers. With these explicit choices for the torsors denoted by $s(\alpha, a)$, the cusp packets are*

d	α	a
O	$e_O\kappa(u)$	$-\kappa(u)$
P	$\alpha_P + e_P\kappa(u)$	0
Q_c	$\alpha_Q + e_{Q_c}\kappa(u)$	0

(B.6)

Here u ranges over the full pro-3 completion of $\mathbb{Q}^\times$. This parametrization is injective on each row, and the rows have disjoint images.

Proof. The morphism $U \hookrightarrow E \setminus \{O\}$ fills the cusps P, Q_c, so their inertia maps to the identity in C. At O, the once-punctured elliptic presentation gives $c_O = -[a, b]$ in the central lattice of C. Consequently a cusp twist $\kappa(u)$ changes the C-coordinate by $-\kappa(u)$ at O and by zero at P, Q_c. The V-coordinate is exactly (B.1). The map to V and its already proved packet disjointness and parameter injectivity give both remaining assertions. □

Statement B.4 (Proved). Lead 4 for the full geometric pro-three group. *For every continuous section $s : G_{\mathbb{Q}} \to N \rtimes G_{\mathbb{Q}}$ put $\Delta = \ker(\chi : G_{\mathbb{Q}} \to \widehat{\mathbb{Z}}^\times)$. Then*

$$C_N(s(\Delta)) = \begin{cases} I_{\widetilde{d}}, & s \text{ is cuspidal at the lifted cusp } \widetilde{d}, \\ 1, & s \text{ is noncuspidal.} \end{cases} \tag{B.7}$$

In the first case the lifted cusp is unique and its image is one of O, P, Q_c. Thus both implications in Lead 4 hold for the actual pro-3 group, with its genuine section action.

Proof. We give the pro-3 argument. No fixed element is lifted from a quotient of a full profinite fixed group.

For an open subgroup $H \subset N$, the corresponding finite geometric cover has maximal pro-3 fundamental group H. To check this comparison, choose an open normal subgroup $M \subset H$ of N; N/M is a finite 3-group and H/M is subnormal in it. Pulling back a subnormal chain expresses the cover as a tower of covers of 3-power degree with normal successive groups. A finite 3-group quotient of a subgroup at one stage becomes a quotient of a finite 3-group after taking the finitely many conjugates and then the extension by the stage's 3-group. Induction proves that the topology induced from the original maximal pro-3 quotient is the full pro-3 topology on the subgroup. This proves the comparison. For its compactification X_H and geometric boundary D_H, the oriented surface presentation gives the exact sequence

$$\mathbb{Z}_3(1) \xrightarrow{\delta} \mathbb{Z}_3(1)[D_H] \longrightarrow H^{\mathrm{ab}} \longrightarrow T_3 J_{X_H} \longrightarrow 0. \tag{B.8}$$

Every arrow is a continuous homomorphism, and the last two arrows are induced respectively by inertia and by filling the cusps. Its only boundary relation is the diagonal sum.

Each lifted inertia group is $\mathbb{Z}_3(1)$. Two distinct lifted cusps have inertia groups meeting trivially. Indeed, separate the cusps in a finite normal pro-3 cover, refine until there are at least three boundary points, and take a positive 3-power of a purported common element so that it lies in the cover group. It remains nonzero in $\mathbb{Z}_3(1)$ and (B.8) puts it on two different boundary lines; those lines have zero intersection. The refinement exists because an inertia group has infinite index: an open procyclic subgroup would give a hyperbolic cover whose abelianization has rank one, contradicting its surface presentation. Normal covers therefore have unboundedly many points above a cusp. The same intersection argument shows that the geometric normalizer of nontrivial inertia is that inertia group, and its arithmetic normalizer is the corresponding cusp decomposition group: a normalizer fixes the lifted cusp, and stabilizers are precisely those groups. These geometric assertions also agree with Nakamura's pro-class statements, Lemmas (2.1.2) and (2.1.4) [Nak94].

Let $z \in C_N(s(\Delta))$. The action through s on the finitely generated pro-3 group N has an open stabilizer for each open subgroup H: a characteristic open subgroup inside H reduces the assertion to an action on a finite group. Consequently H is realized by a cover over some number field L, and $s|_{G_L}$ lifts to that cover when H contains z. The image of z in (B.8) is fixed by $G_{L(\mu_\infty)}$. Ribet's Appendix Theorem 1 [Rib81] says that an abelian variety over a number field has finite torsion over its full cyclotomic extension. Its consequence here is exact: a nonzero invariant vector of $T_3 J_{X_H}$ would give cyclotomic rational torsion points of unbounded 3-power order. Thus the image of z in the last term of (B.8) is zero.

Apply this to $H = M\langle z\rangle$, for every open normal $M \subset N$. The image of z in H^{ab} belongs to the boundary subgroup. The quotient H/M is cyclic of 3-power order and generated by the image of z. Its boundary images generate it, and a cyclic 3-group generated by subgroups equals one of those subgroups. Thus z fixes a cusp in the cover associated to M. The finite sets of such fixed cusps are nonempty at every normal level. A common refinement supplies every finite set of compatibility requirements; compactness gives a lifted cusp $\widetilde{d}$ fixed by z. Hence $z \in I_{\widetilde{d}}$.

We verify that all nonzero elements of the centralizer belong to one inertia group. More generally let a closed subgroup $B \subset N$ consist entirely of inertia elements. The set of cusps with $B \cap I_c \neq 1$ is B-stable. If it were a single orbit while every $B \cap I_c$ were proper, B would be a union of conjugates of one proper closed subgroup. A finite quotient in which this subgroup remains proper contradicts this: in the finite coset action the average number of fixed cosets is one and the identity fixes more than one, so some element fixes none. We can therefore choose two distinct B-orbits of cusps unless B is contained in one inertia group. Compactness separates these orbits at a finite normal level M; the open subgroup BM has two distinct corresponding cusps. When it has only two cusps, its compactification has positive genus. Every element of B maps trivially to the compactification group. A degree-three finite unramified cover of that compactification consequently gives a further open subgroup containing B with six cusps. We have thus arranged at least three cusps while keeping the specified two distinct.

Take nonzero z, z' in the two inertia intersections. Their boundary images are ae_1, be_2, with $a, b \neq 0$. Both zz' and z^2z' have nonzero boundary image. With four or more cusps, neither image can be on a single boundary line. With three cusps, lying on one line forces respectively $a = b$ and $2a = b$, a contradiction. This contradicts the defining property of B and proves the desired single-inertia assertion.

For $C = C_N(s(\Delta)) \neq 1$, we have proved $C \subset I_{\widetilde{d}}$. Each $s(\sigma)$, $\sigma \in \Delta$, fixes a nonzero element there; the inertia-intersection result forces it to fix $\widetilde{d}$. Its inertia action is $\chi_3(\sigma) = 1$, so $I_{\widetilde{d}} \subset C$. Thus equality holds. Normality of Δ makes C stable under all of $s(G_{\mathbb{Q}})$. The arithmetic normalizer identity puts the entire section in the decomposition group of $\widetilde{d}$. Its base cusp is rational, and uniqueness follows from inertia intersections. Conversely a cuspidal

section fixes its inertia over Δ; the same argument gives equality in (B.7). □

Statement B.5 (Proved). The cuspidal members of the full unramified family. *Let $\mathscr{F}_S$ be the image of $H^1(G_{\mathbb{Q},\{3,\infty\}}, N)$ in $H^1(G_{\mathbb{Q}}, N)$, using the integral tangent at O. Its cuspidal subset is the disjoint union of the three packets obtained by twisting the unit tangents at O, P, Q_c by $t\kappa(3)$, $t \in \mathbb{Z}_3$. Every other full section class in $\mathscr{F}_S$ has centralizer 1 over Δ. The centralizer of each cuspidal member is its complete pro-3 inertia group.*

Proof. The good marked model over $\mathbb{Z}[1/3]$ and its integral nonzero tangents give actions and tangential cocycles unramified outside three. This is the based smooth-specialization step used in E.34; the comparison is made on covers with their local parameter root fiber functors, so it applies to the genuine action. For a twist $u \in \mathbb{Q}^{\times,\wedge_3}$, restriction to inertia at $q \neq 3$ has abelian value $v_q(u)e_d$. The vector e_d is primitive and nonzero in V, so a section in $\mathscr{F}_S$ forces $v_q(u) = 0$. Unique factorization in the completed Kummer group then gives $u = 3^t$, with $t \in \mathbb{Z}_3$. Conversely these twists are unramified outside three. Statement B.4 identifies every cuspidal full section with exactly one of these packets and computes all remaining centralizers. The abelianized images are the three affine lines in B.1 with the parameter restricted to $t\kappa(3)$.

This assertion concerns full section classes. The surjection from this family to its abelianized classes in E.34 need not have one-point fibers. An abelian class outside the three displayed lines has exclusively noncuspidal full lifts. We have not deduced cuspidality of every lift above a point on one of those lines. □

Statement B.12 (Proved). The marked local classification when the filled-curve projection is unramified. *Let $k/\mathbb{Q}_q$ be finite, $q \neq 3$, with residue cardinality Q, and let $d = Q_c$ be the marked cusp. Write*

$$\mathscr{Q} = N/\gamma_3 I_{\mathrm{mark}}, \qquad B = N/I_{\mathrm{mark}}, \qquad \mathscr{I} = I_{\mathrm{mark}}/\gamma_3 I_{\mathrm{mark}}, \qquad Z = \gamma_2 I_{\mathrm{mark}}/\gamma_3 I_{\mathrm{mark}}. \tag{B.12.1}$$

All groups carry the actual arithmetic action, and the quotient B is the full geometric pro-three group of the curve obtained by filling d. Choose tame three-inertia τ, arithmetic Frobenius ϕ, the peripheral generator c at d, and the unit tangential reference $s_{d,1}$ compatibly. Put $F_0 = s_{d,1}(\phi)$. The induced based action Φ of F_0 on B is the action of the point d on the filled curve. There are actual equivariant identifications

$$M := \mathscr{I}^{\mathrm{ab}} = \mathbb{Z}_3[[B]](1), \qquad Z = \widehat{\bigwedge}^2 M, \qquad (F_0)_M = Q\Phi, \quad (F_0)_Z = Q^2 \widehat{\bigwedge}^2 \Phi. \tag{B.12.2}$$

Here the completed modules are inverse limits over finite continuous partitions of the profinite label set B; the label 1 is the chosen lift of the marked cusp.

Every local section of $\mathscr{Q}$ whose projection to B is unramified and whose tame inertia is nontrivial has, up to geometric conjugation, exactly the following form:

$$s(\tau) = c^a, \qquad s(\phi) = c^r \exp(z) F_0, \qquad \begin{cases} a \in \mathbb{Z}_3 \setminus \{0\}, \\ r \in \mathbb{Z}_3/(Q-1)\mathbb{Z}_3, \\ [z] \in \mathcal{C}_{k,d} := Z/(Q^2 \widehat{\bigwedge}^2 \Phi - 1)Z. \end{cases} \tag{B.12.3}$$

In particular its B-section is the point section at d. The quotient in (B.12.3) uses the actual image; that image is closed because Z is compact. Every displayed parameter occurs, and two displayed sections are conjugate exactly when all three parameters agree.

Under $\mathscr{Q} \to N/\gamma_3 N$, their image is exactly the ramified local cusp packet at d. For each member of that packet the entire local fiber is the group $\mathcal{C}_{k,d}$, acting simply transitively after the displayed reference is fixed. Finally a local marked section whose abelian tame inertia in V is zero is itself unramified.

Proof. The filling map gives a geometric free product $N = B *_3 \langle c \rangle$ after a choice of free lift of B. Its normal kernel is the free pro-three group on the profinite set of lifted marked points, a torsor under B. The two inverse semidirect-product maps used in F.25, now with $b \mapsto bcb^{-1}$, prove this assertion for the actual filling kernel. The maximal class-two quotient consequently has abelianization $\mathbb{Z}_3[[B]](1)$ and center its completed exterior square. Finite partitions and the finite-rank Hall basis prove these assertions and their inverse-limit compatibility. Galois permutes the labeled inertia generators, up to conjugacy and cyclotomic power. This proves (B.12.2); at the distinguished label the unit tangent also gives the exact equality $F_0 c F_0^{-1} = c^Q$.

We record the fixed-point and measure facts needed below. For any unramified B-section, every positive power of its Frobenius automorphism has fixed subgroup one in B. On B^{ab} the eigenvalues have complex absolute values Q or $\sqrt{Q}$; the same assertion for the degree-r lower-central quotient follows from its equivariant Hall injection into $(B^{\mathrm{ab}})^{\otimes r}$. Inner twists do not change the action on this graded quotient. Thus no such power has eigenvalue one in any positive degree. The intersection of the closed lower central groups is one, as finite three-group quotients are nilpotent. This proves the fixed-subgroup assertion for the full B. It follows that an affine Frobenius action on a B-torsor has at most one point fixed by any given positive power. A finite orbit must therefore be a single fixed point, and there is at most one such point in total.

The closure of that affine action is virtually pro-three. Indeed the translation subgroup is pro-three and the kernel of the action on the Frattini quotient of the finitely generated free pro-three group is pro-three. An integer power removes the finite prime-to-three part. For an invariant $\mathbb{Z}_3$-measure on such a torsor, all mass outside its possible single periodic point is zero. The same assertion holds with $\mathbb{F}_3$ coefficients. To see this precisely, stabilize a clopen set avoiding the periodic point by an open subgroup of the pro-three closure. Every orbit there is infinite. For any n, compactness supplies a finite invariant partition on which every orbit has size divisible by 3^n. Pushforward sums make the measure of the clopen set divisible by 3^n; over $\mathbb{F}_3$ one step already makes it zero. This is the integral measure argument of F.25, applied to the actual permutation action.

Now let the projection of s to B be unramified. Wild inertia and the tame factors other than three map trivially to the geometric pro-three group. Write $t = s(\tau) \in \mathscr{I}$ and $F = s(\phi)$, so $FtF^{-1} = t^Q$. On M the action of F is $Q\Psi$ for the affine Frobenius on the marked-point path torsor. The image m of t therefore satisfies $\Psi m = m$. The measure argument makes $m = a\delta_x$ for an actual fixed label x, or makes it zero when there is no such label.

When $m = 0$, the element t is in Z. The fixed relation says $(Q\widehat{\bigwedge}^2\Psi - 1)t = 0$ after dividing by the unit Q. This operator is injective. On every finite label quotient the permutation has finite order, so all its eigenvalues are roots of unity; none becomes one after multiplication by the rational integer $Q > 1$. Tensoring with $\mathbb{Q}_3$ therefore proves injectivity on that finite free lattice. The inverse-limit comparison in (B.12.2) then proves injectivity on Z. Hence $t = 1$.

For nontrivial t we consequently have $a \neq 0$. The fixed path x is exactly an isomorphism between the projected section and the point section at the filled point d: on inertia both actions are trivial by good marked reduction, and on the unramified quotient it is the displayed Frobenius fixed-path condition. Conjugate by a lift of that path in $\mathscr{Q}$. The label becomes 1, the B-projection is the specified point section, and $F = hF_0$ with $h \in \mathscr{I}$.

Use the integral class-two logarithm, with its sole denominator two, and set $e = \delta_1$. Write

$$t = (ae, z), \qquad b = h_{\mathrm{ab}}.$$

The equality $F_0 c F_0^{-1} = c^Q$ removes any central origin term on the line $\mathbb{Z}_3 e$. Class-two

collection now gives the exact equation

$$(Q\widehat{\bigwedge}^2\Phi - 1)z = a\, e \wedge b. \tag{B.12.4}$$

In fact $F_0 t = (Qae, Q^2(\bigwedge^2 \Phi)z)$ and conjugating by h adds $aQ\, b \wedge e$; comparison with $t^Q = (Qae, Qz)$ gives (B.12.4).

Augmentation and the fixed label give the equivariant splitting $M = \mathbb{Z}_3 e \oplus M^0$. Projection of (B.12.4) to $\widehat{\bigwedge}^2 M^0$ is zero, by the same finite-permutation eigenvalue argument. Thus $z = e \wedge v$ for a unique $v \in M^0$, and

$$(Q\Phi - 1)v = ab^0, \qquad b^0 = b - \mathrm{aug}(b)e. \tag{B.12.5}$$

The operator $Q\Phi - 1$ on M^0 is injective modulo three. If $Q = 1 \bmod 3$, its kernel consists of Φ-invariant $\mathbb{F}_3$-measures; if $Q = -1 \bmod 3$, every element of its kernel is fixed by Φ^2. Either way the measure argument supports it only at 1, and augmentation zero removes that last possibility. Reduction of the completed module commutes with the finite label inverse limits: compatible lifts exist by compactness, or by the surjective transition maps of the finite free modules. Writing $a = 3^s u$ with u a unit, repeated reduction of (B.12.5) now gives $v \in 3^s M^0 = aM^0$. For a lift h_1 of v/a, the explicit conjugation $t \mapsto h_1 t h_1^{-1}$ changes t to c^a. This is an integral divisibility argument; no division by a nonunit has been assumed.

An element of $\mathscr{Q}$ centralizing c^a must have trivial B-image, because the action of B on the label torsor is free and $a\delta_b = a\delta_1$ forces $b = 1$. Inside $\mathscr{I}$, the commutator equation is $ae \wedge h_{\mathrm{ab}} = 0$. Torsion-freeness and the augmentation splitting give $h_{\mathrm{ab}} \in \mathbb{Z}_3 e$. Consequently the exact centralizer is

$$C_{\mathscr{Q}}(c^a) = \langle c \rangle Z. \tag{B.12.6}$$

The relation with F_0 puts FF_0^{-1} in this centralizer, giving precisely $c^r \exp(z)F_0$. Conjugating by $c^u \exp(z')$ changes the two Frobenius parameters by $(Q-1)u$ and $(Q^2 \bigwedge^2 \Phi - 1)z'$. There is no cross term, since Z is central in $\mathscr{I}$ and c commutes with it. Abelianization fixes a, and (B.12.6) proves that these are all possible identifications. Conversely every displayed pair satisfies the tame relation. Continuous powering gives the homomorphism on the procyclic unramified quotient, and the tame semidirect-product presentation gives an actual continuous local section. This proves (B.12.3).

All conjugates of the marked generator have the same image e_d in V. Their pairwise commutators therefore vanish in $\gamma_2 N/\gamma_3 N$. Thus the actual quotient homomorphism sends $Z = \gamma_2 I_{\mathrm{mark}}/\gamma_3 I_{\mathrm{mark}}$ to one in $N/\gamma_3 N$. Formula (B.12.3) has exactly the stated class-two cusp image. Conversely, any marked section above a ramified class-two cusp at d has zero abelian inertia in the filled group. The full local theorem B.7 for that filled curve makes its B-projection unramified, so the preceding classification includes its entire fiber. The same argument starts from any marked section with zero inertia in V: its B-projection is unramified, while $ae_d = 0$ forces $a = 0$. The already proved zero-m case then makes $t = 1$. □

Statement B.13 (Proved). The central marked local fiber is a countable product of three-adic integers. *The group $\mathcal{C}_{k,d}$ in (B.12.3) is a nonzero, torsion-free, compact metrizable $\mathbb{Z}_3$-module. After a choice of divisible basis of its discrete dual, there is a continuous $\mathbb{Z}_3$-linear isomorphism*

$$\mathcal{C}_{k,d} \simeq \prod_{n \geq 1} \mathbb{Z}_3. \tag{B.13.1}$$

The coordinates in (B.13.1) depend on that stated choice; the quotient and the section action in (B.12.3) are intrinsic to the displayed reference. In particular there are actual local marked sections with the same ramified class-two cusp image which are not cuspidal and have no full-N lift.

Proof. Let $D = Q^2 \bigwedge^2 \Phi - 1$. The completed exterior square identifies with the antisymmetric direct summand of $\mathbb{Z}_3[[B \times B]]$ by the idempotent $(1 - \mathrm{swap})/2$. Modulo three, D is $\bigwedge^2 \Phi - 1$. The only periodic point of Φ on B is 1, so the only periodic point on $B \times B$ is $(1, 1)$. The invariant-measure argument of B.12 makes every invariant $\mathbb{F}_3$-measure a multiple of $\delta_{(1,1)}$. Its antisymmetric part is zero. Thus D is injective modulo three.

If $3z = Dv$, reduction modulo three gives $v = 3v_1$, and cancellation in the torsion-free module Z gives $z = Dv_1$. This proves that the cokernel has no three-torsion, and hence no torsion at all. It is compact and metrizable because B is a finitely generated pro-three group, its finite characteristic quotients form a countable cofinal family, and all the coefficient quotients are finite. The image DZ is compact and closed.

We prove infinite rank with actual finite quotient maps. Put $V_B = B^{\mathrm{ab}}$, of rank nine, and $\Omega_n = V_B/3^n V_B$. The homomorphism $B \to \Omega_n$ induces a continuous surjection

$$Z/3Z \longrightarrow \bigwedge^2 \mathbb{F}_3[\Omega_n].$$

It sends D to $\bigwedge^2 \Phi_n - 1$, and therefore induces a surjection from $\mathcal{C}_{k,d}/3$ onto the coinvariants of that finite permutation module. If m is the order of Φ on $V_B/3$, its order on Ω_n divides $m3^{n-1}$: expand the powers of the integral matrix $\Phi^m = 1 + 3A$. The exterior basis vectors $\delta_0 \wedge \delta_v$, $v \neq 0$, form an invariant direct summand. No orbit reverses its orientation, because Φ_n fixes 0 and does not send a nonzero vector to 0. Their coinvariants have one independent vector for each orbit, and consequently dimension at least

$$\frac{3^{9n} - 1}{m3^{n-1}}.$$

This tends to infinity. A concrete nonzero test already at $n = 1$ takes a rational unmarked cusp vector v and sums the coefficients on $\delta_0 \wedge \delta_v$ and, when distinct, on $\delta_0 \wedge \delta_{-v}$. Frobenius multiplies that cusp vector by Q, so the sum kills D modulo three and takes the value one on an actual central commutator $[c, bcb^{-1}]$ whose label b maps to v.

For an explicit description of the isomorphism in (B.13.1), let $\mathcal{C}^\vee$ be the group of continuous characters of $\mathcal{C}_{k,d}$ into $\mathbb{Q}_3/\mathbb{Z}_3$. It is countable and three-primary. Injectivity of multiplication by three on $\mathcal{C}_{k,d}$ makes multiplication by three surjective on $\mathcal{C}^\vee$, so the latter is divisible. This dual assertion can be checked without a continuity assumption left implicit: a finite-valued character on the closed subgroup $3\mathcal{C}_{k,d}$ factors through its image in a finite quotient of $\mathcal{C}_{k,d}$, and extends there because $\mathbb{Q}_3/\mathbb{Z}_3$ is divisible. Pulling that finite character back supplies the continuous extension. Choose a basis of its subgroup killed by three, and compatible successive division chains for these basis elements. Independence and exhaustion follow by induction on the order of a character; this gives an isomorphism $\mathcal{C}^\vee \simeq \bigoplus_{n \geq 1} \mathbb{Q}_3/\mathbb{Z}_3$. The index set is countably infinite by the unbounded finite dimensions just proved. Evaluation on those chosen division chains is the continuous homomorphism in (B.13.1). Duality, or separation of points by finite characters followed by compactness, proves that it is an isomorphism.

The cuspidal class with the parameters a, r has $[z] = 0$. By B.12 a nonzero $[z]$ cannot be removed by any geometric conjugation, so it is not cuspidal. A full local lift would be cuspidal by B.7, and would retain that property under the actual quotient to $\mathscr{Q}$. Thus these sections have no full lift. □

Statement B.14 (Proved). The complete local marked image and the unmarked ramified fibers. *Retain the local data and actual quotients of B.12. If the projection of a section of $\mathscr{Q}$ to B is ramified, that section is itself cuspidal at a unique unmarked cusp $d' \in D(k) \setminus \{d\}$. For a fixed ramified B-section the lift to $\mathscr{Q}$, up to geometric conjugation, is unique and is the image of its actual full cusp section.*

Consequently the complete local image of $\mathrm{Sec}_k(\mathscr{Q})$ *in* $\mathrm{Sec}_k(N/\gamma_3 N)$ *equals the local full image: all unramified classes and the disjoint ramified cusp packets of B.8. The fiber over a ramified unmarked cusp class is a singleton; the fiber over a ramified marked cusp class is exactly the group* $\mathcal{C}_{k,d}$ *of B.12–B.13.*

Proof. By the full local theorem B.7 applied to the filled curve, a ramified projected section is cuspidal at an unmarked rational cusp d'. Filling d is an isomorphism on the inertia and tangent data at d'. Thus this projected section has an actual full cusp lift, which we choose as reference. It remains to prove that every lift to $\mathscr{Q}$ is conjugate to that reference.

Let X be the profinite torsor of paths on the filled curve from this reference at d' to the filled point d. In these coordinates $M = \mathbb{Z}_3[[X]](1)$. The geometric inertia image of the reference is $J_0 = \langle c_{d'}^a \rangle \simeq \mathbb{Z}_3$, with $a \neq 0$. It acts freely on X by translation, and diagonally on $X \times X$. Both are pro-three actions without finite orbits. The integral measure lemma therefore gives $M^{J_0} = 0$ and $(\mathbb{Z}_3[[X \times X]])^{J_0} = 0$. Taking the antisymmetric direct summand gives $Z^{J_0} = 0$ as well.

We justify continuous cohomology and orbit coinvariants with these compact coefficients. For a procyclic three-group generated by T acting continuously on a compact pro-three module A, evaluation at T gives

$$H^1_{\mathrm{cts}}(J_0, A) = A/(T-1)A. \tag{B.14.1}$$

Every value $x \in A$ defines a continuous cocycle: its norm sum at T^{3^n} tends to zero in every finite coefficient quotient, since on that quotient the action has three-power order and the repetitions are eventually divisible by its exponent. These finite cocycles are compatible and give the stated continuous cocycle. Coboundaries are exactly $(T-1)A$; this image is compact and closed. In the permutation module case, pushforward of measures gives an actual isomorphism

$$M/(T-1)M \simeq \mathbb{Z}_3[[J_0\backslash X]](1). \tag{B.14.2}$$

At each finite continuous invariant partition, permutation coinvariants are the free module on its orbits. Inverse limits preserve this description: the finite systems of correcting elements have compact solution sets, so the kernel of pushforward is the actual closed image $(T-1)M$. Equivalently the universal continuous homomorphism from the completed free module to an orbit-constant module has exactly this kernel. The same construction for $X \times X$ commutes with the idempotent $(1 - \mathrm{swap})/2$, and hence computes $Z/(T-1)Z$ as the corresponding antisymmetric direct summand of the orbit module.

Let Ψ denote the actual Frobenius permutation on $J_0\backslash X$. The induced Frobenius action on (B.14.1) contains an inverse factor Q: conjugation sends T to T^Q, so the pullback of a cocycle evaluated at the generator is multiplied by Q^{-1} in coinvariants. On M the coefficient action has factor Q. Thus the action on (B.14.2) is precisely Ψ, with no remaining Tate factor.

There are no finite Ψ-orbits. We construct the map proving this assertion. Filling all remaining punctures maps the full pro-three group of the filled curve onto T_3E. On path torsors it gives a continuous equivariant map from X to the proper path torsor from d' to d. The group J_0 maps to one in T_3E, because it is peripheral inertia. The map consequently descends to $J_0\backslash X$. A point there periodic under Ψ^n would give a fixed point of the proper path torsor under Frobenius to the nth power. Original arithmetic inertia already acts trivially on this torsor by good reduction. Such a fixed point would make the completed Kummer class of $d - d'$ zero over the finite unramified degree-n extension of k.

That difference is a nonzero three-torsion point of E. At every finite extension of a local field of residue characteristic different from three, the reduction kernel is pro-residue-characteristic and the three-primary completion of $E(k')$ is its finite three-primary torsion

subgroup. In particular its nonzero three-torsion point remains nonzero in that completion. The inverse-limit Kummer injection then makes its class nonzero in $H^1(k', T_3E)$. Being divisible once, or several times, in a larger finite group would not make this completed class zero. This proves the required absence of periodic points.

The Frobenius action on the compact quotient set is virtually pro-three, as in B.12. Its invariant integral measure group is therefore zero. Equations (B.14.1)–(B.14.2) show

$$H^1_{\mathrm{cts}}(G_k, M) = 0. \tag{B.14.3}$$

For precision, the unramified quotient causes no additional term: its possible first cohomology term has coefficient $M^{J_0} = 0$. One can obtain the same conclusion without invoking a spectral sequence. If the inertia cocycle is a coboundary, remove it. The tame relation then makes the Frobenius value J_0-invariant, hence zero. Wild and non-three tame inertia have no nonzero homomorphism to these compact pro-three modules.

For Z the inverse-Q factor on (B.14.1) removes one of its two coefficient Tate factors. Thus Frobenius acts on $Z/(T-1)Z$ by Q times the permutation of the diagonal-J_0 orbit set in $X \times X$, restricted to the antisymmetric summand. This group has no invariant vector. On each finite invariant partition, a cycle of length h would force $(Q^h - 1)v = 0$ in a free $\mathbb{Z}_3$-module, which implies $v = 0$. The inverse-limit orbit-module comparison then gives the same vanishing integrally. There is no extra orbit reversal: if a translation in J_0 sends (x, y) to (y, x), its square fixes x; freeness and the absence of order-two elements in J_0 imply $x = y$. Together with $Z^{J_0} = 0$, this proves

$$H^1_{\mathrm{cts}}(G_k, Z) = 0. \tag{B.14.4}$$

The difference between two lifts of the fixed B-section is a continuous $\mathscr{I}$-cocycle, for the reference action just used. By (B.14.3) its image in M is a coboundary. Lift a correcting element from M to $\mathscr{I}$ and conjugate the section. Its cocycle now has values in the central subgroup Z. Equation (B.14.4) removes this last cocycle by a central conjugation. This proves uniqueness of the actual cusp lift in the ramified projected case.

If a class-two section is unramified, choose any lift of its geometric Frobenius value to $\mathscr{Q}$. Continuous powering defines an unramified lift on the procyclic unramified Galois quotient. The ramified cases are exactly B.12 and the case just proved, giving the full image assertion. For a ramified unmarked cusp class, its projected B-section is the unique cusp class determined by its class-two image: the nonzero inertia line identifies the cusp, and the completed local Kummer parameter is already injective in the abelian cusp packet. The uniqueness above therefore gives the singleton fiber. The marked fiber is precisely B.12–B.13. The class-two tame identity, the correcting-conjugation sign, and the finite orbit-sum certificates for the nonzero marked central fiber are checked in `code/bc_marked_checks.py` and its output. □

Statement B.15 (Proved). An explicit cleaned Heisenberg detector at seven. *Put $k = \mathbb{F}_7(\theta)$, $\theta^3 = 2$, and write $[a, b, c] = a + b\theta + c\theta^2$. On $E_k : y^2 + y = x^3$, let $\zeta = 2$ and*

$$\begin{aligned} P &= (3, 2), \qquad P^- = (3, 4), \\ Q &= ([4, 1, 1], [2, 0, 2]), \\ S &= ([0, 6, 5], [1, 4, 3]), \\ T_1 &= ([3, 3, 3], [6, 2, 2]), \qquad T_2 = -T_1. \end{aligned} \tag{B.15.1}$$

The following functions are explicit:

$$\begin{aligned} a &= \frac{x^2+[1,6,5]x+[0,1,4]+[4,4,4]y}{x-3}, \quad b = x-3, \\ f_1 &= \frac{x^2+[2,3,5]x+[6,4,6]+[6,1,6]y}{x-[3,3,3]}, \\ f_2 &= \frac{x^2+[2,3,5]x+[0,3,0]+[1,6,1]y}{x-[3,3,3]}, \\ m &= \frac{x^2+[2,5,2]x+[6,2,2]+[2,4,5]y}{x-[3,3,3]}, \quad f = \frac{m}{f_1 f_2^2 (x-[3,3,3])}. \end{aligned} \tag{B.15.2}$$

Define $p(v)=\sum_{i=0}^{5} p_i v^i$ *and* $h(v)=\sum_{i=0}^{3} h_i v^i$ *by*

i	0	1	2	3	4	5
p_i	$[3,0,0]$	$[0,6,6]$	$[6,1,5]$	$[4,0,2]$	$[4,1,2]$	$[1,5,4]$
h_i	$[6,1,4]$	$[6,0,1]$	$[2,0,0]$	$[1,5,4]$		

(B.15.3)

In $k(E)(v)$ *with* $v^3=b$*, put*

$$u(v)=\frac{p(v)+y}{vh(v)}, \qquad r = f\,u(v)u(\zeta v)^2. \tag{B.15.4}$$

Then

$$v^3=b, \qquad w^3=a, \qquad z^3=r \tag{B.15.5}$$

defines a connected Galois cover of degree 27*, with Heisenberg group of exponent three. It is finite étale over* $U_k = E_k \setminus E_k[3]$*. The actual deck generators are*

	v	w	z
σ	ζv	w	$u(\zeta^2 v)z/w$
τ	v	ζw	z
ρ	v	w	ζz,

$$[\sigma,\tau]=\rho. \tag{B.15.6}$$

For the Fermat tangent parameter s *at* O*, the first two radicals retain the leading coefficient* 9*, whose reduction is* 2*. The cleaned central radical satisfies* $r \sim s^4$*. Thus the based Puiseux choices are*

$$v \sim \theta s^{-2/3}, \qquad w \sim \theta s^{-2/3}, \qquad z \sim s^{4/3}. \tag{B.15.7}$$

There is also an explicit Frobenius transport on the compositum of the three coefficient-conjugate covers. Let a superscript $[j]$ *mean raising every coefficient to* 7^j*, with* j *read modulo three, and use the same choices (B.15.7) for* w_j, z_j*. The arithmetic Frobenius at the framed tangent is*

$$\begin{aligned} \Phi(v) &= \zeta^2 v, \qquad \Phi(w_j) = \zeta^2 w_{j+1}, \\ \Phi(z_j) &= \frac{\zeta w_{j+1} z_{j+1}}{u^{[j+1]}(\zeta v)}, \qquad \Phi^3 = 1. \end{aligned} \tag{B.15.15}$$

Proof. All coefficients in this proof are computed in the displayed finite field. The group law gives $3Q=P$ and $3S=T_2$. The function a is the normalized Miller function

$$\operatorname{div}(a) = 3(Q) - (P) - 2(O). \tag{B.15.8}$$

For example, multiplying the tangent line at Q and the line through $2Q, Q$, and dividing by the vertical lines at $2Q, P$, gives (B.15.2); the common factor $x - x(2Q)$ cancels in both

numerator components. This is an equality of rational functions with the retained tangent leading coefficient, rather than an identification of divisor classes.

The full norm identity is

$$N_{k(E)(v)/k(E)}u = a. \tag{B.15.9}$$

Here is a direct finite calculation of it. Write

$$R(B) = [6,1,0] + [0,3,1]B + [0,2,2]B^2 + [5,4,0]B^3.$$

Expansion using $y^2 + y = x^3$, $v^3 = x - 3$ gives

$$N(p(v) + y) = a(x-3)R(x-3), \qquad N(h(v)) = R(x-3). \tag{B.15.10}$$

Since $N(v) = x - 3$, (B.15.9) follows without a scalar ambiguity. The norm calculation can also be checked by the cubic identity $N(A + Bv + Cv^2) = A^3 + bB^3 + b^2C^3 - 3bABC$.

We next exhibit the removal of every auxiliary ramification point. The cubic h is irreducible over k. Let α be its root and let

$$R_i = (\alpha^3 + 3, -p(\zeta^i\alpha)), \qquad i = 1, 2. \tag{B.15.11}$$

These are points of $E(k(\alpha))$; the candidate with $i = 0$ does not lie on E. Their Frobenius orbits have degree three. Denote the associated closed-point divisors by $\mathcal{R}_1, \mathcal{R}_2$. Exact addition of the three conjugates gives $\mathrm{Tr}(R_i) = T_i$. The two successive line constructions for $R_i + R_i^{343} + R_i^{343^2} = T_i$ give precisely f_i:

$$\mathrm{div}(f_i) = \mathcal{R}_i - (T_i) - 2(O). \tag{B.15.12}$$

The coefficients descend to k after cancellation of the intermediate vertical factor; they are the coefficients displayed in (B.15.2). Likewise m is the normalized Miller function with divisor $3(S) - (T_2) - 2(O)$. Since $T_1 = -T_2$, the vertical function has divisor $(T_1) + (T_2) - 2(O)$. Therefore

$$\mathrm{div}(f) = 3(S) + 6(O) - \mathcal{R}_1 - 2\mathcal{R}_2. \tag{B.15.13}$$

At $\mathcal{R}_i$, the valuations modulo three of $u(v)u(\zeta v)^2$ are i, on all three sheets. For example at $v = \alpha$ the denominator contributes -1, and the numerator contributes 1, 2, or 0 according to whether its zero occurs at $\alpha, \zeta\alpha$, or $\zeta^2\alpha$. The two actual points have indices $1, 2$, giving residues $1, 2$. Formula (B.15.13) cancels them. At the three lifts of Q the numerator orders are $0, 0, 3$; the central valuations are therefore multiples of three. The extra zero S in (B.15.13) also has order three. Equations (B.15.8)–(B.15.10) list all other zeros and poles: they lie over P, P^-, O. The first two cubic extensions are unramified at all interior points, and the last radical now has valuation divisible by three there. Since the residue characteristic is seven, the normalized cover is étale over U_k.

For completeness, the field degrees and deck morphisms can be verified directly. The classes of a, b in $k(E)^\times/k(E)^{\times 3}$ are independent: the valuation at P^- first forces the coefficient of b to vanish, and the valuation at P then forces the coefficient of a to vanish. Hence $[k(E)(v,w) : k(E)] = 9$. Writing $\sigma(v) = \zeta v$, (B.15.9) gives

$$\frac{\sigma(r)}{r} = \frac{u(\zeta^2 v)^3}{a}. \tag{B.15.14}$$

If r became a cube in $k(E)(v,w)$, the Kummer kernel would give $r = a^j c^3$ in $k(E)(v)$ for some j. Applying σ and (B.15.14) would make $a = (u(\zeta^2 v)c/\sigma(c))^3$ in $k(E)(v)$, contradicting the independence just proved. The last degree is therefore three. Equations (B.15.14) and

(B.15.9) prove that (B.15.6) preserves all defining equations, that $\sigma^3 = 1$, and that $[\sigma, \tau] = \rho$. These are actual automorphisms of the degree-27 field.

Finally use $t = -x/y \sim s/3$ at O. The functions a, b, f_1, f_2, m and the vertical factor all have leading term $9s^{-2}$. Thus $u(v) \sim v$ and $f \sim s^6/3^6$. The original central radical has leading term $\zeta^2\, 9s^{-2}$, so after the twist its coefficient is $\zeta^2\, 9/3^6 = 1$ in k and its exponent is four. This proves (B.15.7). In particular no replacement of the minimal-model tangent constant by 36 is made.

To prove the last assertion, embed the fields in the Puiseux field using (B.15.7). Coefficientwise seventh power, fixing $s^{1/3}$, acts on $k((s^{1/3}))$ with order three, and fixes the Laurent expansions of x, y. It sends the chosen leading coefficient θ to $\theta^7 = \zeta^2\theta$. Moreover (B.15.9) gives the exact identity

$$\frac{r^{[j]}(\zeta^2 v)}{r^{[j]}(v)} = \frac{a^{[j]}}{u^{[j]}(\zeta v)^3}.$$

Thus the proposed image of z_j in (B.15.15) has the correct cube. Its leading coefficient is one, because $w_{j+1}/u^{[j+1]}(\zeta v)$ has leading coefficient ζ^{-1}. The uniqueness of the chosen cube root with this leading coefficient proves the formula. It respects every relation among the conjugate covers, since it is the restriction of the actual coefficient automorphism of the ambient Puiseux field. Iterating three times gives $w_j^3/N(u^{[j]}) = 1$ in the multiplier of z_j, verifying $\Phi^3 = 1$ directly as well.

The exact polynomial identities, the finite-field extensions, the three-conjugate additions, and the displayed coefficients are independently replayed by `code/bc_miller_norm_rr.py` and `code/bc_miller_heisenberg.py`, with their saved JSON and text outputs. This explicit based cover is a detector for the remaining central-action calculation; its construction and the conjugate-cover morphism (B.15.15) do not by themselves assign its central monodromy values on an uncomputed fixed Schreier basis. □

Statement B.16 (Proved). The integral depth-one map through the cyclic-nine cover. *Let F be the free pro-three group on the standard tripod inertia generators x, y, with $[x, y] = xyx^{-1}y^{-1}$. For $f \in F'$ write its metabelian image as $[x, y]^{h(X,Y)}$, where $X = x - 1$, $Y = y - 1$ act by conjugation. There is an actual continuous metabelian quotient in which*

$$\begin{aligned} x &\longmapsto (1, 0), \qquad y \longmapsto (0, 1), \\ (m, a)(n, b) &= (m + t^a n, a + b), \\ m, n &\in \mathbb{Z}_3[[t - 1]], \qquad a, b \in \mathbb{Z}_3. \end{aligned} \tag{B.16.1}$$

In this quotient the image of f is $(m_f, 0)$, with

$$m_f(t) = (1 - t)h(0, t - 1). \tag{B.16.2}$$

Let $H_9 = \ker(F \to \mathbb{Z}/9)$, where $x \mapsto 0$, $y \mapsto 1$. Its actual cover is

$$z^9 = 1 - t_{\mathrm{base}}, \qquad Y_9 = \mathbb{P}^1 \setminus \{0, \infty, \mu_9\}. \tag{B.16.3}$$

In $H_9^{\mathrm{ab}}/3$, write the image of f as $\sum_{j=0}^{8} c_j\, y^j x y^{-j}$; its y^9 coordinate is zero. Then

$$\begin{pmatrix} h_{0,3} \\ h_{0,5} \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 & 0 & 2 & 1 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 & 2 & 2 & 2 \end{pmatrix} \begin{pmatrix} c_0 \\ c_1 \\ c_2 \\ c_3 \\ c_4 \\ c_5 \\ c_6 \\ c_7 \\ c_8 \end{pmatrix} \quad \text{in } \mathbb{F}_3. \tag{B.16.4}$$

For the standard path β from $\overrightarrow{01}$ to $\overrightarrow{10}$ and the convention $f_g = \beta(g\beta)^{-1}$, these coordinates are explicit framed Kummer coordinates on (B.16.3). The relevant endpoint-to-start ratios are

$$r_0 = 9, \qquad r_j = \frac{-\zeta_9^j}{1-\zeta_9^j} \quad (1 \le j \le 8), \qquad \prod_{j=0}^{8} r_j = 1. \tag{B.16.5}$$

Choose their compatible roots by continuing the units $z - \zeta_9^j$ along the lifted path from $z = 1$ to $z = 0$, using the actual start velocity $-1/9$ for $j = 0$. If

$$\frac{g(r_{\chi(g)^{-1}j}^{1/3^n})}{r_j^{1/3^n}} = \zeta_{3^n}^{\kappa_j(g)}, \tag{B.16.6}$$

where the index is taken modulo nine, then $c_j = -\kappa_j(g) \pmod 3$. In particular (B.16.4)–(B.16.6) evaluate the two coefficients by finite cyclic-cover Kummer data for the full cyclotomic action; no restriction to $\ker\chi$ is imposed.

Proof. The group in (B.16.1) is a compact pro-three group. The universal property of F gives the stated continuous homomorphism. Its target is metabelian, and conjugation by x acts trivially on its additive normal subgroup, while conjugation by y multiplies that subgroup by t. Direct multiplication sends $[x, y]$ to $(1 - t, 0)$. Therefore the induced commutator-module map sends X to zero and Y to $t - 1$, proving (B.16.2).

The lift of a positive loop about $t_{\text{base}} = 1$ in (B.16.3) sends z to $\zeta_9 z$. Consequently the Schreier generators $y^j x y^{-j}$ are the actual positive inertia loops at $z = \zeta_9^j$, and y^9 is the positive inertia loop at $z = 0$. The functions $z - \zeta_9^k$ and z have valuations respectively δ_{jk} and zero on the first nine generators, and respectively zero and one on the last generator. Their Kummer characters thus give the actual dual coordinate maps on $H_9^{\text{ab}}/3$. The Schreier basis has ten members, so these coordinate maps are an isomorphism with $\mathbb{F}_3^{10}$. Since $f \in F'$, its total y exponent is zero; hence its y^9 coordinate is zero.

Reducing (B.16.2) modulo three and $t^9 - 1$ gives

$$\sum_{j=0}^{8} c_j (1+Y)^j = -Y h(0, Y) \pmod{Y^9}, \tag{B.16.7}$$

because $(1 + Y)^9 - 1 = Y^9$ in characteristic three. The coefficients of Y^4, Y^6 give $h_{0,3} = -\sum_j \binom{j}{4} c_j$ and $h_{0,5} = -\sum_j \binom{j}{6} c_j$, which are exactly (B.16.4). Only these indicated coefficients are recovered: the Y^8 coefficient of h is lost under multiplication by Y modulo Y^9.

The path β lifts to the real interval from $z = 1$ to $z = 0$. At the start, $z-1 = -t_{\text{base}}/9+\cdots$, and at the end $z - 1 = -1$. Their framed ratio is therefore nine. For $j \neq 0$ both endpoint values are ordinary nonzero values, giving (B.16.5). The product identity follows from $\prod_{j=1}^{8}(1 - \zeta_9^j) = 9$ and $\prod_{j=1}^{8}(-\zeta_9^j) = 1$. The continued roots obey the product identity as well: their arguments occur in opposite pairs, and the $j = 0$ root is the positive real root of nine.

To check the sign in (B.16.6), choose one root of the unit at the start and continue it along β to the end. The path $g\beta$ transports the Galois conjugate start root to the Galois conjugate end root, with the unit index permuted by $\chi(g)$. Following β and then $(g\beta)^{-1}$ multiplies the start root by the inverse of the endpoint-to-start Kummer multiplier. Thus the positive inertia coordinate of f_g is $-\kappa_j(g)$, as asserted. This argument constructs the maps for every g; it does not extend a formula stated only on $\ker\chi$ by an unspecified normalization.

The program `code/bc_depth_one.py` checks the actual Schreier coordinates against the X coefficient of the abelianized Fox derivative $D_y(f) = Xh(X, Y)$ on explicit commutator words. It also verifies the exact cyclotomic-field identities (B.16.5), their degree-nine to

degree-three transition products, the Frobenius-seven index permutation, and the two rows in (B.16.4). The saved output has status `PASS`. □

Statement B.17 (Proved). The actual norm-289 depth-one Frobenius coordinates. *Let g_{17} be an arithmetic Frobenius of norm 289 at the prime above seventeen of $K_0 = \mathbb{Q}(\zeta_3)$, viewed through the specified tangential arithmetic action. Use the path and commutator conventions of B.16, and write the metabelian Ihara word as $f_{g_{17}} = [x, y]^{h(X,Y)}$. Then*

$$h(0, Y) \equiv -Y^6 - Y^7 \pmod{(3, Y^8)}, \qquad h_{0,3} = h_{0,5} = 0 \quad \text{in } \mathbb{F}_3. \tag{B.17.1}$$

An explicit residue-field embedding is

$$\begin{aligned} k &= \mathbb{F}_{17}[\theta]/(\theta^2 - 3), \\ \overline{\zeta}_9 = 7 + 4\theta, \qquad \overline{\zeta}_3 &= \overline{\zeta}_9^3 = 8 + 15\theta. \end{aligned} \tag{B.17.2}$$

For the nine framed ratios of B.16 the actual local Kummer coordinates and geometric cusp coordinates are

$$\begin{aligned} (\kappa_j(g_{17}))_{j=0}^8 &= (0, 1, 2, 0, 1, 2, 0, 1, 2), \\ (c_j)_{j=0}^8 &= (0, 2, 1, 0, 2, 1, 0, 2, 1). \end{aligned} \tag{B.17.3}$$

The entries are in $\mathbb{F}_3$. These values hold for each prime of $\mathbb{Q}(\zeta_9)$ above seventeen, with its compatible embedding of $\zeta_3 = \zeta_9^3$.

Proof. The integer three is a nonsquare modulo seventeen, and direct multiplication in (B.17.2) gives $\overline{\zeta}_9^9 = 1$, $\overline{\zeta}_9^3 \neq 1$, and $\overline{\zeta}_9^{17} = \overline{\zeta}_9^{-1}$. Choose the embedding of $\mathbb{Q}(\zeta_9)$ into $\overline{\mathbb{Q}}_{17}$ that reduces to (B.17.2). Its completion is the unramified quadratic extension of $\mathbb{Q}_{17}$, which is also the completion of K_0. Every r_j in (B.16.5) is a unit there. The polynomial $Z^3 - r_j$ is finite etale on the local unit locus: its derivative is a unit at every root. Its roots therefore lie in finite unramified extensions.

Let ρ_j be the cube root selected by the actual path of B.16. Reduction is injective on μ_3. Arithmetic Frobenius acts on its residue as the 289th power, so the exact local comparison is

$$\overline{\frac{g_{17}(\rho_j)}{\rho_j}} = \overline{\rho}_j^{288} = \overline{r}_j^{96} = \overline{\zeta}_3^{\kappa_j(g_{17})}. \tag{B.17.4}$$

Here g_{17} fixes μ_9, so there is no index permutation in (B.16.6). It also fixes μ_3; changing a cube root by an element of μ_3 does not change (B.17.4). Thus this calculation computes the prescribed based cocycle itself modulo three. Changing the representative by local inertia also leaves it unchanged because these unit extensions are unramified.

We can evaluate every symbol without choosing any radical extension. For $1 \leq j \leq 8$, the seventeen-power Frobenius sends $\overline{r}_j$ to $\overline{r}_{-j}$, with indices modulo nine, and

$$\overline{r}_j \overline{r}_{-j} \in \mathbb{F}_{17}^{\times}, \qquad \frac{\overline{r}_j}{\overline{r}_{-j}} = -\overline{\zeta}_9^j. \tag{B.17.5}$$

Every element of $\mathbb{F}_{17}^{\times}$ has trivial 96th power. Since $96 \equiv 6 \pmod 9$, (B.17.5) gives

$$\kappa_j + \kappa_{-j} = 0, \qquad \kappa_j - \kappa_{-j} = 2j \quad \text{in } \mathbb{F}_3.$$

Thus $\kappa_j = j \pmod 3$. The ratio $r_0 = 9$ also has trivial symbol, which proves the first vector in (B.17.3). The actual sign comparison of B.16 gives $c_j = -\kappa_j$ and proves the second. The argument uses only primitivity of the chosen ninth root and its cube, so it applies to every embedding above seventeen.

For completeness, the entire specialization through this precision follows from the same nine symbols. In $\mathbb{F}_3[t]/(t^9-1)$,

$$\sum_{j=0}^{8} c_j t^j = -t\frac{d}{dt}\left(\sum_{j=0}^{8} t^j\right) = t(t-1)^7.$$

With $t = 1+Y$, equation (B.16.7) becomes $-Yh(0,Y) = Y^7 + Y^8 \pmod{Y^9}$. Division by Y at the stated precision proves (B.17.1). No coefficient of degree eight or higher is assigned by this division.

The standalone program `code/bc_depth_one_frob17.py` executes all field operations in (B.17.2), evaluates the nine symbols by 96th powers, verifies (B.17.5), reconstructs the cyclic group-ring polynomial and the two rows of B.16, and repeats the computation for all six primitive ninth roots in k. The saved JSON and text outputs have status `PASS`. □

Statement B.18 (Proved). The full finite arithmetic image and the exact Frobenius action at seventeen. *Use the actual source basis of A.38 and the c-adapted basis m, n of A.31. Let*

$$\rho : P_3 \longrightarrow \mathrm{Aut}(N/(N^3\gamma_3 N))$$

be the actual arithmetic action. Its image G has order 243, nilpotency class three, and exponent nine. In the $10+45$ coordinates of A.38, its linear image has order nine, and its identity-linear kernel has dimension three over $\mathbb{F}_3$. The exact dimensions are

	$\gamma_2(G)$	$\gamma_3(G)$	$\gamma_4(G)$
$\dim_{\mathbb{F}_3}$	2	1	0

	$D_2(G)$	$D_3(G)$	$D_4(G)$
$\dim_{\mathbb{F}_3}$	3	2	0.

(B.18.1)

In particular $D_4(P_3)$ acts trivially on this specified geometric quotient. Put $M = \rho(m)$ and $N_* = \rho(n)$. Then the entire actual Frobenius action at seventeen is

$$\rho(\varphi_{17}) = M^6[N_*, [N_*, M]] = M^6 \neq 1. \tag{B.18.2}$$

It has order three. Its Johnson block has rank two and annihilates all nine cusp vectors. The full matrix is retained in `bc_ihara_finite_image_certificate.npz`.

Proof. An automorphism in these coordinates has matrix

$$\begin{pmatrix} A & 0 \\ J & \bigwedge^2 A \end{pmatrix}.$$

Its identity-linear subgroup is the additive group $\mathrm{Hom}(\mathbb{F}_3^{10}, \bigwedge^2 \mathbb{F}_3^{10})$. For such a matrix X, conjugation is the exact map

$$X \longmapsto (\bigwedge^2 A) X A^{-1}. \tag{B.18.3}$$

The script below reconstructs $\rho(u)$ and $\rho(c)$ from the A.38 word and its literal Schreier map, forms $\rho(v) = \rho(c)\rho(u)\rho(c)$, and applies the actual square-root source maps defining m, n. These reconstructed matrices agree with the independently saved arithmetic certificate.

The two linear matrices of M, N_* commute, each has order three, and their nine products are distinct. Define Z_0, Z_1, Z_2 to be the additive Johnson blocks, respectively, of

$$M^3, \qquad [M, N_*], \qquad [M, [M, N_*]].$$

Exact substitution gives three linearly independent vectors and the conjugation table

$$\begin{array}{c|ccc} & Z_0 & Z_1 & Z_2 \\ \hline \mathrm{Ad}(M) & Z_0 & Z_1+Z_2 & Z_2 \\ \mathrm{Ad}(N_*) & Z_0 & Z_1 & Z_2. \end{array} \tag{B.18.4}$$

It also gives $N_*^3 = 1$. The subgroup generated by the three Z_i is therefore normal, is elementary abelian of order 27, and is contained in the actual image. After quotienting by it, the images of M, N_* commute and have order three. The linear image has order nine, so the quotient is exactly $(\mathbb{Z}/3)^2$. This proves $|G| = 243$, and shows that the displayed subgroup is the whole identity-linear kernel.

The normal closure of $[M, N_*]$ is exactly $\langle Z_1, Z_2\rangle$ by (B.18.4). Its next commutator is $\langle Z_2\rangle$, which is central. Thus the lower-central assertions in (B.18.1) follow as actual group statements. Every cube lies in the identity-linear kernel, and that kernel has exponent three; consequently every element has ninth power one. Since M^3 is nontrivial, the exponent is nine. The cubes span $\langle Z_0, Z_2\rangle$, as verified on the 243 distinct normal forms. This also proves the two asserted Zassenhaus dimensions.

The defining formula for this filtration gives

$$D_4(P_3) = \overline{\gamma_4(P_3)(\gamma_2 P_3)^3 P_3^9}.$$

Each factor has trivial image by the preceding calculations. Equation (A.31.3), transported by the actual automorphism in A.38, therefore gives the equality in (B.18.2), not only a leading-term comparison. Table (B.18.4) gives $[N_*, [N_*, M]] = 1$. The matrix M^6 has order three, and its 45×10 Johnson block has rank two. Multiplication by the nine actual cusp columns of the Schreier certificate gives zero in every case.

The standalone `code/bc_ihara_finite_image.py` computes the full normal commutator module by the exact maps (B.18.3), checks (B.18.4), and independently enumerates the subgroup generated by M, N_* using literal matrix multiplication. Both computations give the same 243 elements. It verifies all ninth powers, the cube span, the Frobenius word, its rank and cusp values, and saves every indicated basis and matrix. Its JSON and text outputs have status `PASS`. □

Statement B.19 (Proved). A finite precision determining the integral local kernel. *Let $M_2 = \bigwedge^2 V$, and let*

$$\mathcal{D} = M_2^c \oplus \ker(1 + \rho(n)c : M_2 \longrightarrow M_2)$$

be the rank-45 lattice of normalized source cocycles used in A.39. Define the actual Frobenius evaluation

$$E_\varphi : \mathcal{D} \longrightarrow M_2^c, \qquad E_\varphi(\gamma) = \gamma(\varphi_{17}), \qquad R = \ker E_\varphi.$$

The integral operator $B = (F_\varphi - 1)|M_2^c$ has three-adic Smith exponents

$$\underbrace{1,\dots,1}_{6},\quad \underbrace{2,\dots,2}_{7},\quad \underbrace{3,3}_{2},\quad \underbrace{4,\dots,4}_{6}. \tag{B.19.1}$$

Identify $R/3R$ with its image in $\mathcal{D}/3\mathcal{D}$. Then its exact finite-precision description is

$$\frac{R}{3R} = \mathrm{im}\left(\ker\left(\mathcal{D}/243\mathcal{D} \xrightarrow{E_\varphi} M_2^c/243M_2^c\right) \longrightarrow \mathcal{D}/3\mathcal{D}\right). \tag{B.19.2}$$

More explicitly, for every integral $d \in \mathcal{D}$ with $E_\varphi(d) \in 243M_2^c$, put

$$h = B^{-1}E_\varphi(d) \in 3M_2^c, \qquad \gamma = d - \delta h, \qquad (\delta h)(g) = g(h) - h. \tag{B.19.3}$$

Then $\gamma \in R$ and $\gamma \equiv d \pmod{3\mathcal{D}}$. The matrix B and the bound (B.19.1) are computed in the attached certificate. Evaluating the separate arithmetic matrix $E_\varphi \bmod 243$ remains the finite calculation needed to list the resulting 24-dimensional subspace individually.

Proof. At seventeen, cubing permutes $\mathbb{F}_{17}$. Consequently the reduction $y^2 + y = x^3 - 7$ has one x for each y, and its point count, including infinity, is eighteen. Its Frobenius polynomial is $X^2 + 17$, so the norm-289 Frobenius on T is exactly -17. All cusps are rational over K_0, and its action on W is multiplication by 289. The actual graph-lattice comparison (A.7.4) therefore gives

$$F_\varphi|V = \begin{pmatrix} 289 I_8 & 102\Psi \\ 0 & -17 I_2 \end{pmatrix}. \tag{B.19.4}$$

We take its full exterior square and restrict to the c-fixed lattice. The latter has rank 21. The projection to $\bigwedge^2 T$ is zero on it, because c acts there by -1; hence it is also the fixed lattice inside $\mathcal{K}$.

The program constructs a basis from 21 independent columns modulo three of $1 + c$ on M_2. This is an actual $\mathbb{Z}_3$ basis of the fixed lattice: $\frac{1}{2}(1+c)$ is an integral idempotent, and the selected columns span its reduction modulo three, so Nakayama's lemma gives the full lattice. An invertible minor of three-adic unit determinant gives the coordinate restriction of (B.19.4). Both the rectangular matrix in M_2 and its 21×21 coordinate matrix have the exponents (B.19.1). An independent elimination modulo 3^8 gives the same list. The determinant check is

$$\det B = 83520^{13}\, 4914^8, \qquad v_3(\det B) = 50.$$

Indeed the rational fixed subspace has thirteen toric wedges, of eigenvalue 289^2, and eight mixed wedges, of eigenvalue $-17 \cdot 289$. The inverse matrix verifies that $81B^{-1}$ is integral over $\mathbb{Z}_3$, while $27B^{-1}$ is not. This proves the stated bound with its actual integral comparison.

For $h \in M_2^c$, the coboundary δh is normalized at c and belongs to $\mathcal{D}$. Its evaluation is exactly

$$E_\varphi(\delta h) = Bh.$$

Thus $\operatorname{im} E_\varphi$ contains $B(M_2^c)$, which contains $81M_2^c$ by (B.19.1). The formula (B.19.3) is therefore integral and has $h \in 3M_2^c$ whenever $E_\varphi(d)$ is divisible by 243. It gives $E_\varphi(\gamma) = 0$ and leaves the reduction of d modulo three unchanged. Conversely every element of R reduces to an element of the displayed finite kernel. Since the target of E_φ is torsion-free, $R \cap 3\mathcal{D} = 3R$. This proves both the claimed identification and (B.19.2).

The exact reconstruction, point count, two Smith computations and inverse-denominator check are executed in `code/bc_frobenius17_precision.py`. Its JSON and text outputs have status `PASS`, and include the fixed-lattice basis and the full integral matrix B. □

Statement B.20 (Proved). An individually specified class-two target with no class-four lift. *Use the actual arithmetic basis m, n, c of A.38 and A.31. Write the ten integral A.7 basis vectors as*

$$\begin{aligned}(v_0, \dots, v_7) &= (e_P, e_{Q_c}, e_A, e_{\bar{A}}, e_B, e_{\bar{B}}, e_C, e_{\bar{C}}),\\ v_8 = (\Psi(1,0)/3, (1,0)), &\qquad v_9 = (\Psi(0,1)/3, (0,1)) \quad \text{in } L_\Psi \subset W_{\mathbb{Q}_3} \oplus T_{\mathbb{Q}_3}.\end{aligned} \tag{B.20.1}$$

Let C_2 be the actual action of c on $M_2 = \bigwedge^2 V$. Define the integral vectors

$$b_0 = 3v_8 \wedge v_9 + \sum_{i=0}^{7} \big(\Psi_{i,2} v_i \wedge v_8 - \Psi_{i,1} v_i \wedge v_9\big), \qquad b = (1 - C_2) b_0. \tag{B.20.2}$$

Here the columns of Ψ are numbered one and two. In the basis $v_i \wedge v_j$, $i < j$, all nonzero coordinates of b are the following; every other coordinate is zero:

(i,j)	b_{ij}	(i,j)	b_{ij}	(i,j)	b_{ij}
$(0,8)$	-2	$(1,8)$	2	$(2,6)$	-1
$(2,7)$	1	$(2,8)$	2	$(2,9)$	2
$(3,6)$	1	$(3,7)$	-1	$(3,9)$	-2
$(4,6)$	1	$(4,7)$	-1	$(4,8)$	-2
$(4,9)$	-2	$(5,6)$	-1	$(5,7)$	1
$(5,9)$	2	$(6,8)$	2	$(6,9)$	-2
$(7,8)$	-2	$(7,9)$	2	$(8,9)$	$6.$

(B.20.3)

The prescriptions

$$\gamma(m) = 0, \qquad \gamma(n) = b, \qquad \gamma(c) = 0 \tag{B.20.4}$$

define an actual continuous M_2-cocycle unramified outside three. Let $s_{O,17}$ be the actual O-cuspidal section with tangent parameter seventeen. In $A = N/\gamma_3 N$, put

$$s_b(g) = \gamma(g) s_{O,17}(g). \tag{B.20.5}$$

This is an actual class-two section, with abelian class $\kappa_{17} e_O$. Its restriction at seventeen equals the specified local cuspidal section exactly. It has a global lift to $N/\gamma_4 N$, but no global lift to $N/\gamma_5 N$. Consequently it has no full, metabelian, or marked-depth lift.

The projection of (B.20.4) to $\bigwedge^2 T$ has the exact generator values $0, 6$. If $a = \kappa_3(n) \in \mathbb{Z}_3^\times$ in this actual source basis, this cocycle is $(6/a)\kappa_3$. Only $a \equiv 1 \pmod 3$ is used; its integral value has not been replaced by one.

Proof. We first prove the integral local condition. On V, the actions of m, n are respectively $G_u G_v$ and $G_v G_u^{-1}$ from (A.7.4)–(A.7.5). They commute, and the first commutes with c. Set

$$L_0 = \ker(n - 1 : M_2 \to M_2) \cap \ker(c + 1 : M_2 \to M_2).$$

This is a closed saturated lattice stable under m, n, c and their inverses. Direct substitution in the exact graph matrices gives

$$n(b_0) = b_0, \qquad n(b) = b, \qquad c(b) = -b. \tag{B.20.6}$$

For the first equality, the rational comparison is also explicit: the vector 3 times the wedge of the two pure proper vectors has the mixed coordinates in (B.20.2), proper coefficient three, and toric coefficient $\frac{1}{3}\Psi_1 \wedge \Psi_2$. Subtracting this last toric vector gives exactly b_0. Both pure proper determinant and toric wedges are fixed by n, so the equality transfers through the actual graph embedding. No reduction of a rational splitting modulo three is taken.

The full lattice is

$$L_0 = (\bigwedge^2 W)^{c=-1} \oplus \mathbb{Z}_3 b, \qquad \operatorname{rank} L_0 = 16. \tag{B.20.7}$$

Here is an integral verification. The toric anti-invariant lattice has rank fifteen. Choose the lexicographically first independent columns modulo three of $1 - c$ on its 28 wedges. These are an actual integral basis by the order-two projector and Nakayama's lemma. Adjoining b gives sixteen columns independent modulo three. The stacked rational matrices $n - 1, c + 1$ have rank 29, so these columns span their sixteen-dimensional common rational kernel. Their unit minor makes their integral span primitive in M_2, proving (B.20.7). The code also computes

the exact matrix of m on this basis and its inverse; both are integral over $\mathbb{Z}_3$. This verifies the stated stability on the actual lattice.

Since m, n freely generate P_3, their prescribed values give a continuous crossed homomorphism. The identities $c(m) = m$, $c(n) = n^{-1}$ and (B.20.6) prove its exact c-equivariance, so it extends with value zero at c. Every value lies in L_0. The actual norm-289 Frobenius φ_{17} is fixed by c, hence

$$\gamma(\varphi_{17}) = c\gamma(\varphi_{17}) = -\gamma(\varphi_{17}) = 0. \tag{B.20.8}$$

The final equality uses that two is invertible. Thus (B.20.4) lies in the exact integral lattice R of A.39, not merely in a finite approximation to it. Its inflation has zero tame value at seventeen and zero value at the normalized local involution as well. These three values prove the exact local equality in (B.20.5).

We give a finite functional detecting the ensuing gluing obstruction. Let $X_0, \dots, X_9$ be the ordered Schreier generators in (A.38.3), and use the integral degree-three basis

$$H_{\mu,i} = [[X_i, X_{\min \mu}], X_j],$$

where μ is a nondecreasing multiset of size three, $i > \min \mu$ occurs in it, and j is the remaining index after removing i and one occurrence of $\min \mu$. Write $q_{\mu,i}$ for the corresponding coordinate functional. Over $\mathbb{F}_3$, define

$$\begin{aligned} \ell = 2\big(&q_{001,1} + q_{002,2} + q_{003,3} + q_{004,4} \\ &+ q_{088,8} + q_{188,8} + q_{288,8} + q_{388,8} + q_{488,8}\big). \end{aligned} \tag{B.20.9}$$

All these indices refer to the Schreier basis, not to an unidentified change of proper lifts. The program uses the exact unimodular $A.7$-to-Schreier matrix P and the actual map $\bigwedge^2 P$ on b. It verifies that this map intertwines the full m, n, c actions on M_2 before evaluating the source cochain.

With the independent Magnus evaluator of `bc_gluing_audit.py`, the functional has the following exact values. It kills the entire global M_3 restriction image, all twenty-one retained local-conjugator parameters, and all sixty-eight unknown-word contributions on the cocycle (B.20.4). Its value on the known Frobenius contribution is

$$\ell(\Delta) = 1. \tag{B.20.10}$$

Only three of those unknown maps are nonzero on the general source, and all three vanish on the whole lattice L_0 modulo three. The known map vanishes on its fifteen-dimensional toric summand and is nonzero on b. In the fixed 143-dimensional quotient of the independent audit, its vector has 87 nonzero entries. All twenty-seven remaining arithmetic coefficient possibilities give exactly this same vector.

These are evaluations of the actual global gluing map of A.39. Changing the integral class-three lift adds a global M_3 correction, which (B.20.9) kills. The retained source truncation and all local comparison parameters are those independently verified there, so (B.20.10) applies to the actual arithmetic Frobenius. A.33 gives a global class-three lift; A.40 proves that this nonzero gluing class prevents every class-four lift. The actual quotient maps and local cusp comparisons in A.39–A.40 give the full, metabelian and marked non-lifting conclusions on section classes.

Finally the proper projection of b is six by (B.20.3). The positive-root Kummer cocycle κ_3 has value zero at c, and hence at the c-fixed generator m. Its value at n is a unit by (A.31.6). Equality on the actual free source generators proves the asserted identity with $(6/a)\kappa_3$.

The standalone `code/bc_explicit17.py` executes the integral lattice, saturation, action and basis comparisons, then every stated functional and gluing evaluation. It saves all 45

coordinates, the exact rank-16 basis, the transition matrix, the full nonzero vector and the nine-term functional. Its JSON, NPZ and text certificates have status `PASS`. □

Statement B.6 (Proved). The complete global one-prime image and its fibers. *Fix a rational prime $q \neq 3$, put $A = N/\gamma_3 N$, and let s be a continuous class-two section unramified outside $S = \{3, q, \infty\}$. Choose its exact cocycle representative with $s(c) = 1$, and write α for its projection to V. Define*

$$\begin{aligned} &H_0 = N, \qquad H_{n+1} = H_n^3[H_n, H_n], \\ &P_n = N/H_n, \qquad A_n = P_n/\gamma_3 P_n, \qquad J_n = \gamma_3 P_n, \\ &S_n = \{3, \infty\} \cup \{q : \alpha|_{I_q} \bmod 3^n \neq 0\}. \end{aligned} \tag{B.6.1}$$

All subgroups are closed. The exact finite target s_n is the image of s in A_n.

Use the explicitly constructed finite based action on P_n and its splitting field F_n from A.43. Let K_n be the compositum of F_n with the field of the target graph (s_n, ρ_{A_n}). If s_n is ramified outside S_n, the test fails. Otherwise form the finite Galois field Ω_n as the compositum of Ω_{n-1} and all Galois extensions $E/\mathbb{Q}$ containing K_n, unramified outside S_n, of degree at most $|P_n|[F_n : \mathbb{Q}]$, starting with $\Omega_0 = \mathbb{Q}$. The discriminant and polynomial bounds in (A.43.4)–(A.43.10) give a terminating finite enumeration of these fields.

Let X_n be the finite set of functions $z : \mathrm{Gal}(\Omega_n/\mathbb{Q}) \to P_n$ satisfying the actual cocycle equation, $z(c) = 1$, and $z \bmod J_n = s_n$ literally. For every $k \leq n$, require its projection to P_k to be trivial on inertia at the primes of $S_n \setminus S_k$. These conditions give the actual transition maps of A.43. Then

$$\begin{aligned} [s] \in \mathrm{im}(\mathrm{Sec}(N) \longrightarrow \mathrm{Sec}(A)) &\iff X_n \neq \varnothing \quad \text{for all } n \\ &\iff [s] \text{ lifts to } N/\gamma_r N \quad \text{for all } r \geq 3. \end{aligned} \tag{B.6.2}$$

Every such full or nilpotent lift is unramified outside S. The complete fiber on section classes is

$$\varprojlim_n X_n/J_n^c, \tag{B.6.3}$$

where $h \in J_n^c$ acts by $z(g) \mapsto h^{-1} z(g) g(h)$. Thus (B.6.2) is an exact sequence of terminating finite arithmetic tests, and (B.6.3) retains all compatible conjugacy data. A bounded stopping rule for every positive instance is not asserted.

The necessary local condition is precisely the unramified class or a member of the actual cusp packets (B.8.1). It is sufficient at seven by B.10, and on the prime sets Σ and Ξ by A.16 and A.25. It is insufficient at seventeen. The integral section of B.20, with abelian class $\kappa(17)e_O$ and central generator values $\gamma(m) = 0$, $\gamma(n) = b$, has exactly the local O-cuspidal class-two restriction. It has a global $N/\gamma_4 N$ lift and no $N/\gamma_5 N$ lift, hence no full, metabelian, or marked-depth lift.

Proof. For the full group, the three quotient maps in (A.43.2) specialize exactly to (B.6.1). The image of H_n in V is $3^n V$, by induction on the Frattini operation. Hence $A_n^{\mathrm{ab}} = V/3^n V$, and the finite target table determines the displayed set S_n. The actual quotient map sends $\gamma_3 N$ onto $\gamma_3 P_n$, which proves the asserted kernel J_n. This identifies every group and action in the finite test with its A.43 counterpart.

The inherited ramification conditions ensure that projection to P_k factors through Ω_k; they are retained even when the prime q first enters the sets S_n at a later precision. When $\alpha(I_q) = 0$, it never enters, so an inertia value visible only in the central class-two target is rejected as soon as it is detected in some A_n. This is consistent with B.7's detection of full inertia in V. A.43 now gives the first equivalence in (B.6.2) and the exact fiber (B.6.3).

Its comparison uses the surjections $\ker(N \to A)^c \twoheadrightarrow J_n^c$ and the trivial global geometric centralizer of the target in A. It therefore identifies global section classes, without replacing the conjugation kernel by a finite-level centralizer. B.11 gives the second equivalence and the common ramification bound, including for separately chosen nilpotent lifts.

For the obstruction coordinates requested in Task B.4, entry into A is the residue, tame-relation and convergence criterion of Part A. The next central coefficient is the actual free lattice $M_3 = \gamma_3 N/\gamma_4 N$. Its residue complex at q is

$$[\, M_3(-1) \xrightarrow{F_q - 1} M_3(-1)\,], \qquad F_q t = qt \text{ for a tame inertia value } t \in M_3. \tag{B.6.4}$$

Here F_q in the complex acts on the twisted module and the second F_q acts on M_3; the equations are the same under the chosen tame generator. A.33 computes the continuous obstruction using this localization map. For the fixed finite set S, the finite-coefficient first-cohomology groups are finite, so their first derived limit is zero. Actual local cuspidal lifts kill the local degree-two obstructions and give a global class-three lift. Its remaining obstruction to having the prescribed local cuspidal classes is exactly the restriction cokernel (A.33.2). A.39 evaluates a nonzero family in that cokernel; A.40 proves the class-four exclusion; B.20 specifies an individual member with all integral coordinates and an explicit nine-term finite functional. These facts prove the stated failure of local sufficiency.

The finite tests (B.6.2) retain every subsequent geometric layer and global relation. They thus give the full one-prime image at seventeen and at every other prime, including the classes for which the local condition alone does not decide lifting. B.1, B.3 and B.9 separately give all three global cusp packets and their inherited central origins, as required in Task B.1. □

3 C. Restriction and germs

3.1 Restriction, descent, and persistent obstruction germs

Write $A = N/\gamma_3 N$ and $\mathrm{Sec}_L = H^1_{\mathrm{cts}}(G_L, A)$ for a number field $L \subset \overline{\mathbb{Q}}$. The action is the restriction of the action based at the fixed tangent at O. Our maps below use that actual action, and all equivalences of sections are by conjugation in A. This is the germ construction requested in CK C9, cote 149, §3, manuscript p. 33, scan 66, applied to the stated class-two quotient; the Fermat comparison is §51.16, folios 733v–736v, scans 172–178.

Statement C.1 (Proved). No arithmetic centralizer in the class-two geometric group. *For every number field L and every continuous section $s : G_L \to A \rtimes G_L$,*

$$C_A(s(G_L)) = 1. \tag{C.1}$$

The assertion uses the whole G_L, in contrast to the subgroup $\ker\chi$ in B.4; the comparison map between the two assertions is the inclusion of subgroups, which reverses inclusions of their centralizers.

Proof. Let h be in the displayed centralizer. Its image in $V = A^{\mathrm{ab}}$ is fixed by G_L, since inner conjugation is trivial on abelianization. Choose a finite prime v of L above two, with residue size $q = 2^f$. The eigenvalues of arithmetic Frobenius on $W \otimes \mathbb{Q}_3$ have complex absolute value q: they are q times roots of unity from the finite cusp permutation action. On $T_3E \otimes \mathbb{Q}_3$ the two eigenvalues have absolute value $q^{1/2}$. Here the necessary elliptic calculation is explicit: $E(\mathbb{F}_2)$ has three points by `bc_checks.py`; the Frobenius F_2 has determinant two by the Weil pairing, and $\det(1 - F_2) = \deg(1 - F_2) = 3$ because $1 - F_2$ is separable with precisely those three kernel points. Thus its characteristic polynomial is $X^2 + 2$. On the base change at v the Frobenius is F_2^f, which proves the asserted absolute values directly. The

characteristic polynomial of the extension V is the product of those on W and T, so one is not an eigenvalue. Consequently $V^{G_L} = 0$, and h lies in the central group $\gamma_2 N/\gamma_3 N = \bigwedge^2 V$. Inner conjugation acts trivially there. Its Frobenius eigenvalues, as seen from the filtration induced by $0 \to W \to V \to T \to 0$, have absolute values $q^2, q^{3/2}, q$. Again none is one. The lattice is torsion-free, so its invariant subgroup is zero. Hence $h = 1$. □

Statement C.2 (Proved). All restriction maps are injective and finite Galois descent is exact. *For number fields $L \subset L'$, restriction is the injective map of pointed sets*

$$\mathrm{res}_{L'/L} : \mathrm{Sec}_L \longrightarrow \mathrm{Sec}_{L'}, \qquad [z] \longmapsto [z|_{G_{L'}}]. \tag{C.2}$$

For L'/L finite Galois it identifies its source with precisely the $\mathrm{Gal}(L'/L)$-fixed classes in its target. Writing ρ for the based action on A, the action by $g \in G_L$ is represented by

$$(g \cdot z)(h) = \rho(g) z(g^{-1}hg), \quad h \in G_{L'}. \tag{C.3}$$

Thus the kernel of (C.2), as a pointed-set map, is its base point, and its image in the invariant set is the entire set.

Proof. Restriction preserves the continuous cocycle equation and geometric conjugation, proving that (C.2) is defined. For (C.3), conjugate the section graph by $(1, g)$ in $A \rtimes G_L$. This gives exactly the formula and proves continuity and the cocycle equation. When $g \in G_{L'}$, the cocycle identity gives

$$\rho(g) z(g^{-1}hg) = z(g)^{-1} z(h) \rho(h) z(g),$$

so the action is conjugation by $z(g)^{-1}$ and is trivial on section classes. Therefore it factors through the specified finite Galois group.

For injectivity, conjugate two sections so that their restrictions to $G_{L'}$ agree. Choose a finite extension M/L, Galois over L, with $M \supset L'$. The restrictions agree on the normal subgroup G_M. For each $g \in G_L$, the element $t(g)s(g)^{-1} \in A$ centralizes $s(G_M)$: both sections induce, by conjugation, the same map $s(h) \mapsto s(ghg^{-1})$ there. Statement C.1 gives $t(g)s(g)^{-1} = 1$, proving equality over G_L.

For the image assertion, let s represent a fixed class over L'. Consider its normalizer

$$B = N_{A \rtimes G_L}(s(G_{L'})).$$

It is a closed subgroup of a compact profinite group. Invariance of the class under (C.3) says precisely that each $g \in G_L$ is the projection of an element of B: use the conjugator in A comparing $g \cdot s$ with s. The kernel of $B \to G_L$ equals $C_A(s(G_{L'}))$. Indeed, an element projecting to one normalizes the graph only by preserving each of its points, since there is exactly one graph point over each $h \in G_{L'}$. This kernel is one by C.1. Hence $B \to G_L$ is a continuous bijective homomorphism between compact Hausdorff groups and has continuous inverse. That inverse is a section extending s, because its unique element over $h \in G_{L'}$ is $s(h)$. Injectivity already proved uniqueness up to geometric conjugacy. □

Statement C.3 (Proved). The action and fields of definition of germs. *The set*

$$\mathscr{G} = \varinjlim_{L \subset \overline{\mathbb{Q}}} \mathrm{Sec}_L \tag{C.4}$$

has a well-defined $G_{\mathbb{Q}}$-action obtained by (C.3) together with conjugation of the field of definition. Every transition map is injective. Each germ has an open stabilizer and a finite orbit. For every number field L,

$$\mathrm{Sec}_L \xrightarrow{\sim} \mathscr{G}^{G_L}. \tag{C.5}$$

In particular the image of $\mathrm{Sec}_{\mathbb{Q}}$ *is exactly the fixed-germ set. The fixed field of the stabilizer of a germ is its smallest field of definition.*

Proof. The formula transports a cocycle on G_L to a cocycle on G_{gL}, and respects restriction and conjugation. It therefore defines an action on the colimit. Restriction injectivity is C.2. A representative over L is fixed by G_L, since the same cocycle calculation used after (C.3) makes inner field conjugation trivial on its class. The stabilizer is consequently open.

Take a germ fixed by G_L, choose a field where it is represented, and enlarge it to a finite Galois extension M/L. For $g \in G_L$, the class and its translate are both defined over M. Their equality as germs means equality after restriction to a further finite field; injectivity in C.2 makes them equal already over M. The invariant class descends by C.2 to L. Uniqueness is again injectivity. This proves (C.5).

An open stabilizer $H \subset G_{\mathbb{Q}}$ is G_{L_0} for a number field L_0. Applying the same argument over L_0 proves definability there. Every other field of definition has its Galois group inside H, so it contains L_0. This proves the final assertion. □

Statement C.4 (Refuted). The R2 obstruction does not disappear as a germ. *Let* $c = 9\kappa(3) \cup \beta$ *be the nonzero R2 class specified by LM7 (E.32f), so that in* $H^2_{\mathrm{cts}}(G_{\mathbb{Q}}, \mathbb{Z}_3(2)) = \mathcal{R}/d\mathcal{R}$ *it is represented by*

$$z_{q_j} = q_j - 1 \quad (j \geq 3), \qquad z_q = 0 \text{ at the other primes.} \tag{C.6}$$

For every finite number field L, $\mathrm{res}_{L/\mathbb{Q}}(c) \neq 0$. *In fact these restrictions all have infinite additive order. Consequently* c *gives a nonzero element of* $\varinjlim_L H^2_{\mathrm{cts}}(G_L, \mathbb{Z}_3(2))$. *The vanishing assertion in the first sentence of Task C.4 is false.*

Proof. For a nonzero integer m, the unique coordinatewise solution of $dy = mz$ has value m at every q_j and zero elsewhere, because multiplication by each nonzero $q - 1$ is injective in $\mathbb{Z}_3$. This family does not tend to zero 3-adically. Thus $mz \notin d\mathcal{R}$ and $mc \neq 0$ for every nonzero integer m.

We spell out the transfer used for this continuous cohomology. Set $G = G_{\mathbb{Q}}$, $H = G_L$ and $M = \mathbb{Z}_3(2)$. Use homogeneous continuous equivariant cochains, with alternating deletion differential. Choose representatives T for the finitely many left cosets $H\backslash G$, and write $r(g) = gt(g)^{-1} \in H$ for $g \in Ht(g)$. For an H-cochain c define the G-cochain

$$(\mathcal{T}c)(g_0, \ldots, g_n) = \sum_{t \in T} t^{-1} c(r(tg_0), \ldots, r(tg_n)). \tag{C.7a}$$

The coset permutation $tg = h(t,g)t'$ gives $t^{-1}h(t,g) = g(t')^{-1}$, proving equivariance; the formula commutes with the differential. All maps are continuous because the transversal is finite and each coset is open. Put $a_t(g) = t^{-1}r(tg)$. For a degree-n G-cochain f, the prism operator is

$$(Pf)(g_0, \ldots, g_{n-1}) = \sum_{t \in T} \sum_{i=0}^{n-1} (-1)^i f(a_t g_0, \ldots, a_t g_i, g_i, \ldots, g_{n-1}). \tag{C.7b}$$

The same coset permutation makes the sum equivariant. In $dP + Pd$ the internal mixed faces cancel in pairs; the two end faces are $[G : H]f$ and $-\mathcal{T}(\mathrm{res}\, f)$. Thus the formulas prove directly on continuous $\mathbb{Z}_3(2)$ cochains that

$$\mathrm{cor}_{L/\mathbb{Q}} \mathrm{res}_{L/\mathbb{Q}} = [L : \mathbb{Q}]. \tag{C.7}$$

Equivalently one can apply the same compatible formulas to every finite coefficient module; their cochain homotopies commute with reduction, so the assertion holds on the derived inverse limit as well as on finite-coefficient cohomology. This includes its $\varprojlim^1$ subgroup.

A zero restriction would make $[L:\mathbb{Q}]c = 0$, contrary to the first paragraph. A torsion restriction would, by the same calculation, make a nonzero integer multiple of c zero. Thus it has infinite order. An element maps to zero in a filtered colimit of groups exactly when it maps to zero at some later stage; no such stage exists here. □

Statement C.5 (Proved). Restriction is injective on the Tate-twist phantom subgroup. *For $\mathbb{Z}_3(2)$ the kernel of the finite-coefficient comparison map in R2 is torsion-free, and restriction from $\mathbb{Q}$ to any finite number field is injective on this kernel. This extends C.4 from its particular cup-product witness to every nonzero class in that kernel.*

Proof. The kernel is

$$\frac{\{z \in \mathcal{R} : z_q \in (q-1)\mathbb{Z}_3 \text{ for all } q\}}{d\mathcal{R}}.$$

Write $y_q = z_q/(q-1) \in \mathbb{Z}_3$. A nonzero integer multiple mz lies in $d\mathcal{R}$ exactly when $(my_q)_q$ is restricted. Multiplication by a fixed nonzero integer changes valuations by the fixed integer $v_3(m)$, so this condition is equivalent to $(y_q)_q$ being restricted. This proves torsion-freeness. Equation (C.7) then proves injectivity of restriction on this subgroup. No closure is taken in the denominator in this argument. □

Statement C.7 (Proved). The intrinsic phantom obstruction also survives every finite extension. *The nonzero obstruction in $H^2_{\mathrm{cts}}(G_{\mathbb{Q}}, \mathcal{K})$ constructed in A.4 remains nonzero over every finite number field and in the filtered colimit over all number fields. More generally, restriction to every finite number field is injective on the phantom subgroup for $\mathcal{K}$ computed in A.2.*

Proof. In the residue presentation of A.2, every $D_q = \mathrm{Frob}_q - 1$ on $\mathcal{K}(-1)$ is injective. A locally zero class has a unique integral correction $y_q = D_q^{-1} z_q$ at each prime. A nonzero integer multiple of its class vanishes precisely when $(my_q)_q$ is restricted, which is equivalent to $(y_q)_q$ being restricted. Thus this phantom subgroup is torsion-free. The cochain transfer and prism formulas (C.7a)–(C.7b) apply verbatim with $M = \mathcal{K}$; they prove corestriction after restriction equals multiplication by the finite extension degree. Such multiplication is injective on the phantom subgroup. This proves the assertion and completes both persistence decisions asked for in Task C.4. □

Statement C.10 (Proved). The nine disjoint cusp germ packets and their Galois action. *Define the abelian germ parameter group*

$$\mathscr{K} = \varinjlim_{L \subset \overline{\mathbb{Q}}} H^1_{\mathrm{cts}}(G_L, \mathbb{Z}_3(1)) = \varinjlim_{L} \varprojlim_{n} L^\times / L^{\times 3^n}. \tag{C.10.1}$$

All transition maps are restriction. Choose at each cusp $d \in D(\overline{\mathbb{Q}})$ the unit tangent for the parameter x/z when $x = 0$, y/z when $y = 0$, and z/x when $z = 0$. These choices are $G_{\mathbb{Q}}$-equivariant. The map

$$\coprod_{d \in D(\overline{\mathbb{Q}})} \mathscr{K} \longrightarrow \mathscr{G}, \qquad (d, u) \longmapsto [s_{d,1} \text{ twisted by } u] \tag{C.10.2}$$

is injective, and its image is exactly the cuspidal germ set. There are nine packets. The $G_{\mathbb{Q}}$-action in these coordinates is

$$g(d,u) = (gd, gu), \tag{C.10.3}$$

where gu transports a Kummer class over L to its class over gL by the field isomorphism $g : L \to gL$. The parameter group has injective restriction maps, is divisible, and has torsion subgroup canonically μ_{3^∞}, with its usual Galois action. In particular, distinct torsion parameters and distinct cusp packets do not merge in the germ colimit.

Proof. The unit tangent gives a section of the decomposition extension at d. Over a number field L containing d, this extension is the semidirect product $\mathbb{Z}_3(1) \rtimes G_L$. Its continuous sections modulo inertia conjugacy are precisely $H^1(G_L, \mathbb{Z}_3(1))$. The decomposition homomorphism into $N \rtimes G_L$ and the continuous quotient $N \to A$ define the map in (C.10.2). Passing to a larger finite field restricts the same section, so the map is defined on (C.10.1). The fields containing d form a cofinal subsystem of the fields in that definition. Every cuspidal germ is, by definition, a section of one such decomposition extension, proving the image assertion once injectivity is checked.

First note that $H^0(G_L, \mathbb{Z}_3(1)) = 0$ for every finite number field L. If a one-cocycle becomes a coboundary over a finite extension, subtract that coboundary and enlarge to a finite Galois extension M/L so that the cocycle vanishes on G_M. For $g \in G_L$, its value is fixed by G_M, by applying the cocycle identity to $gG_Mg^{-1} = G_M$. The displayed invariant group is zero. The cocycle therefore vanishes on G_L, which proves injectivity of every restriction in (C.10.1).

Enlarge a common field of definition to contain all nine cusps. The boundary module is $W = \mathbb{Z}_3(1)[D]/\mathbb{Z}_3(1)\sum_{e\in D} e_e$. For a given d, the map $e_d : \mathbb{Z}_3(1) \to W$ is a split G_L-equivariant inclusion: omit any other cusp from the boundary basis. The map $H^1(G_L, W) \to H^1(G_L, V)$ is injective because $H^0(G_L, T_3E) = 0$, as proved in C.1. Thus $e_d : H^1(G_L, \mathbb{Z}_3(1)) \to H^1(G_L, V)$ is injective. The abelian image of the packet is exactly

$$\alpha_d + e_d H^1(G_L, \mathbb{Z}_3(1)), \tag{C.10.4}$$

where α_d is the Kummer class of the framed tangential divisor $d - O$. This follows from the same division-torsor comparison used in B.1, now over L. Consequently two parameters in one packet cannot become equal as abelian germs, and hence cannot become equal in $\mathscr{G}$.

Filling the boundary gives the continuous equivariant quotient $V \twoheadrightarrow T_3E$. It sends (C.10.4) to $\kappa_L(d - O)$. For $d \neq e$, the difference $d - e$ is a nonzero point of $E[3]$: the flex tangent relation is $3d \sim 3O$, and the nine distinct flexes exhaust $E[3]$. The integral Kummer class $\kappa_L(d - e)$ remains nonzero over every finite extension of L. Indeed the Mordell–Weil group of that extension is finitely generated. A nonzero three-primary torsion element in such a group is not divisible by 3^n for all n, so has nonzero image in its completion and in $H^1(T_3E)$ by the Kummer exact sequence. This argument uses the entire integral Kummer class; its modulo-three image alone can vanish after adjoining ninth division points. It proves that the nine packets remain pairwise disjoint in the colimit.

Our parameters and tangent value one are defined by the coordinate functions over $\mathbb{Q}$. Field conjugation therefore takes the chosen tangent at d to the chosen tangent at gd. Transporting the decomposition inclusion conjugates its geometric path, and geometric conjugacy is exactly the equivalence used in $\mathscr{G}$. Kummer functoriality proves (C.10.3), with no additional translation term.

For completeness, a finite-level Kummer class modulo three is killed by some finite extension: use the finite image of its affine cocycle representation. The long exact sequence of $0 \to \mathbb{Z}_3(1) \xrightarrow{3} \mathbb{Z}_3(1) \to \mu_3 \to 0$ then shows that the corresponding germ is divisible by three.

Multiplication by an integer prime to three is already invertible on each completed Kummer group. Thus $\mathscr{K}$ is divisible. At a fixed finite field, its completed Kummer group's torsion is precisely $\mu_{3^\infty}(L)$. To see this directly, use the divisor sequence from $L^\times$ to its principal fractional ideals. The latter group is free abelian and the sequence splits as abelian groups; the unit group is a finite roots-of-unity group plus a free abelian group by Dirichlet's theorem. The three-adic completion of a free abelian group is a subgroup of a product of copies of $\mathbb{Z}_3$, hence torsion-free. Only the three-primary roots of unity remain as torsion. Since all restrictions in (C.10.1) are injective, passing to the colimit introduces no further torsion and retains all roots of unity of three-power order. This proves the final assertions. □

Statement C.11 (Refuted). The pure-residue computation over the rationals does not extend verbatim to every number field. *Let*

$$L = \mathbb{Q}(\zeta_3, \sqrt{10}), \qquad S_L = \{v : v \mid 3\} \cup \{v : v \mid \infty\}.$$

For the actual rank-forty-four module $\mathcal{K}$,

$$H^2_{\mathrm{cts}}(G_{L,S_L}, \mathcal{K}) \neq 0. \tag{C.11.1}$$

Its inflation into $H^2_{\mathrm{cts}}(G_L, \mathcal{K})$ is injective, and every one of its classes has zero residue outside S_L. Consequently the pure-residue isomorphism (A.1.3) cannot be transported to this L by replacing the rational primes with its finite primes. The finite-S_L complex is a necessary part of Task C.1.

Proof. The polynomial $X^2 - 10$ has two simple roots modulo three and hence two roots in $\mathbb{Q}_3$ by Hensel's lemma. Therefore L has two places above three; each completion is $\mathbb{Q}_3(\zeta_3)$. The field has no real places. The Brauer invariant sequence gives

$$\mathrm{Br}(\mathcal{O}_L[1/3])[3] = \{(a, -a) : a \in \tfrac{1}{3}\mathbb{Z}/\mathbb{Z}\} \simeq \mathbb{F}_3. \tag{C.11.2}$$

Kummer theory and the arithmetic etale comparison surject $H^2(G_{L,S_L}, \mu_3)$ onto this group. These are the same comparison and invariant maps used in A.14: Milne, II.2.9, II.2.1 and Remark II.2.2(a) [Mil06]. Since $\mu_3 \subset L$, fixing $1 \mapsto \zeta_3$ identifies its coefficient with the trivial module $\mathbb{F}_3$. Thus $H^2(G_{L,S_L}, \mathbb{F}_3) \neq 0$.

The image on $\mathcal{K}/3$ is a finite three-group, because L contains $\mathbb{Q}(\zeta_3)$ and the action over that field has this property in A.1. A nonzero module for a finite three-group over $\mathbb{F}_3$ has a nonzero trivial quotient: its coinvariants are nonzero by nilpotence of the augmentation ideal. Choose a surjection $\mathcal{K}/3 \twoheadrightarrow \mathbb{F}_3$. The group G_{L,S_L} has three-cohomological dimension at most two. The long exact sequence therefore makes

$$H^2(G_{L,S_L}, \mathcal{K}/3) \longrightarrow H^2(G_{L,S_L}, \mathbb{F}_3)$$

surjective, so its source is nonzero.

The degree-two finite-coefficient groups are finite. Their inverse system is Mittag–Leffler, and the finite-coefficient groups in degree three vanish. The continuous-cohomology inverse-limit sequence then gives $H^3_{\mathrm{cts}}(G_{L,S_L}, \mathcal{K}) = 0$. Reduction modulo three in the long exact coefficient sequence yields a surjection

$$H^2_{\mathrm{cts}}(G_{L,S_L}, \mathcal{K})/3 \twoheadrightarrow H^2(G_{L,S_L}, \mathcal{K}/3).$$

This proves (C.11.1).

At every finite $v \nmid 3$ the eigenvalues on $\mathcal{K}(-1)$ have absolute values $\#k(v)$ or $\sqrt{\#k(v)}$, by the boundary/elliptic filtration. None is one, so $D_v = F_v - 1$ is injective on this lattice. The

localization triangle of F.1 over L thus has zero degree-zero residue group. Its long exact sequence injects the nonzero finite-S_L degree-two group into the global degree-two group, with image in the kernel of every displayed residue. This proves the precise failure of the proposed pure-residue replacement. □

Statement C.12 (Proved). The exact general number-field criterion with its retained arithmetic term. *For any number field L, put $S_L = \{v \mid 3, \infty\}$, $A_v = \mathcal{K}(-1)$ and $D_v = F_v - 1$ for $v \notin S_L$. Let $\mathcal{R}_L(A)$ denote the families in these integral lattices tending to zero three-adically. Then there are exact maps*

$$\begin{aligned} 0 \longrightarrow H^2_{\mathrm{cts}}(G_{L,S_L}, \mathcal{K}) &\xrightarrow{\inf} H^2_{\mathrm{cts}}(G_L, \mathcal{K}) \\ &\xrightarrow{\mathrm{res}} \mathcal{R}_L(A)/D\mathcal{R}_L(A) \longrightarrow 0. \end{aligned} \tag{C.12.1}$$

$$H^1_{\mathrm{cts}}(G_{L,S_L}, \mathcal{K}) \xrightarrow{\sim} H^1_{\mathrm{cts}}(G_L, \mathcal{K}). \tag{C.12.2}$$

The quotient in (C.12.1) uses the actual image of D. The first term is the inverse limit of the finite groups $H^2(G_{L,S_L}, \mathcal{K}/3^n)$.

For a section of N^{cc} over L, take its actual local path-origin residues r_v by the formula (A.8.3), with residue size $\#k(v)$ and the restricted G_L-action. Its intrinsic obstruction vanishes exactly when both of the following tests hold:

$$y_v = -D_v^{-1} r_v \in A_v \text{ for every } v \notin S_L, \qquad (y_v)_v \in \mathcal{R}_L(A); \tag{C.12.3}$$

$$\eta(s) = 0 \quad \text{in } H^2_{\mathrm{cts}}(G_{L,S_L}, \mathcal{K}). \tag{C.12.4}$$

Here, once (C.12.3) holds, $\eta(s)$ is the unique class whose inflation is the full continuous obstruction of s. Every nonempty class-two lift fiber is a torsor under $H^1_{\mathrm{cts}}(G_{L,S_L}, \mathcal{K})$, with no further geometric-stabilizer quotient. This statement retains the finite-S_L test; it does not claim an evaluation of that test for arbitrary L.

Proof. Apply the finite-coefficient localization triangle of F.1 and take its derived inverse limit. The residue complex is the actual two-term complex $[\mathcal{R}_L(A) \xrightarrow{D} \mathcal{R}_L(A)]$. Each D_v is injective by the weight calculation at the end of C.11, so its degree-zero cohomology is zero. The finite-S_L degree-one and degree-two finite-coefficient groups are finite, and hence their inverse systems are Mittag–Leffler. Three-cohomological dimension at most two gives $H^3_{\mathrm{cts}}(G_{L,S_L}, \mathcal{K}) = 0$ as in C.11. The indicated pieces of the localization long exact sequence are therefore precisely (C.12.1) and its degree-one isomorphism; the same finite-group Mittag–Leffler fact identifies the first term with its finite-coefficient inverse limit.

The local section equation used in A.8 is unchanged after restricting the arithmetic action to G_L: the inertia comparison cochains are the restrictions of the same based cusp paths, with the cusps now permuted by G_L. Applying that derivation over each completion L_v gives exactly r_v as the residue of the continuous obstruction class. Its residue class vanishes precisely under (C.12.3), because every D_v is injective and the sole integral correction is the displayed one. Exactness of (C.12.1) then gives the unique $\eta(s)$. Its vanishing is equivalent to vanishing of the original obstruction, proving necessity and sufficiency of the two tests.

Once a lift exists, two lifts of a fixed representative differ by a continuous $\mathcal{K}$-valued one-cocycle; central conjugacy changes it by a coboundary. A geometric stabilizer of the underlying N^{cc} section has fixed image in V, which is zero by C.1. Its remaining image is in $\mathbb{Z}_3(1)$ and is again zero. Thus no additional orbit identification occurs, and the fiber is the asserted full torsor. □

3.2 Algebraic evaluation of the point image

Statement C.16 (Proved). Finite algebraic formulas for every point's abelian and central coordinates. *Put $m = 3^n$. Two successive Kummer covers and a quotient of their finite deck group construct a based finite cover $X_n \to U$ with group*

$$P_n = N/(N^{3^n}\gamma_3 N). \tag{C.16.1}$$

Its multiplication, arithmetic action, transition maps, and fibers at algebraic points are computed by finite algebraic operations. The tangential base point is the specified $s = z/x$, tangent one.

Choose compatible lifts $y_1, \dots, y_{10}$ of a $\mathbb{Z}_3$-basis of V. For every $A \in U(L)$ the construction gives compatible finite cocycles

$$c_{A,n}(g) = (v_{A,n}(g), z_{A,n}(g)) \in (V/3^n) \oplus (\textstyle\bigwedge^2 V/3^n). \tag{C.16.2}$$

They are obtained by evaluating roots of rational functions at A and comparing the resulting fibers with the Puiseux fiber at O. Their inverse limit is a continuous cocycle whose class is the class-two point section of A. Thus (C.16.2) evaluates all forty-five central coordinates with their tangent origin.

Proof. We specify all the covers and comparison maps. The geometric constructions are over $\overline{\mathbb{Q}}$, and their finitely many coefficients lie in computable number fields.

For a smooth affine curve $Y = C \setminus B$, with C proper of genus g and B nonempty and finite, put

$$\mathcal{F}_m(Y) = \frac{\{f \in \overline{\mathbb{Q}}(C)^\times : \operatorname{ord}_x(f) \in m\mathbb{Z} \text{ for every } x \in Y\}}{\overline{\mathbb{Q}}(C)^{\times m}}. \tag{C.16.3}$$

Valuations give the exact sequence

$$0 \longrightarrow \operatorname{Jac}(C)[m] \longrightarrow \mathcal{F}_m(Y) \longrightarrow \ker\big((\mathbb{Z}/m)^B \xrightarrow{\sum} \mathbb{Z}/m\big) \longrightarrow 0. \tag{C.16.4}$$

The first map sends $[E]$ to the class of a function with divisor mE. An mth-power function makes E principal, proving injectivity. If every boundary valuation of f is also divisible by m, its full divisor is mE, proving exactness in the middle. For surjectivity, lift a boundary vector to a degree-zero divisor D on B. Choose E with $m[E] = [D]$ in the Jacobian and a function with divisor $D - mE$. Multiplication by m on the Jacobian is surjective over $\overline{\mathbb{Q}}$, so these choices exist. They give the required boundary valuations.

Choose a basis of $\operatorname{Jac}(C)[m]$ and the boundary vectors $b - b_0$. The preceding construction gives $r_Y = 2g + |B| - 1$ functions f_i forming a $\mathbb{Z}/m$-basis of $\mathcal{F}_m(Y)$. Normalize Y in

$$\overline{\mathbb{Q}}(C)(f_1^{1/m}, \dots, f_{r_Y}^{1/m}). \tag{C.16.5}$$

Their independence gives degree m^{r_Y}. At an interior point remove the m-divisible valuation from a Kummer equation; extracting a root of the remaining unit is etale in characteristic zero. Thus (C.16.5) is etale on Y. Conversely every cyclic etale cover of exponent dividing m comes from such a function, by function-field Kummer theory and this valuation test. This proves that (C.16.5) is exactly the maximal geometric abelian cover of exponent m.

All these are finite algebraic operations. Represent Jacobian classes by divisors, solve their finite m-division equations, and compute the principal functions by Riemann–Roch. The latter computation is given explicitly by Khuri-Makdisi's Theorem/Algorithm 5.5 [KM04]. Termination can also be certified without choosing a Jacobian implementation: enumerate

algebraic rational functions, test the valuation condition and their classes modulo mth powers, and retain independent functions until their generated subgroup has the known order m^{r_Y} from (C.16.4). Factoring $T^m - f$ over the algebraic function field decides the power tests. Equation (C.16.4) proves that this enumeration reaches the required order.

Apply (C.16.5) to U, and denote the result by $Y_m \to U$. Its intrinsic maximal pro-three group is $H_m = N^m[N, N]$. Indeed the normal closure over U of a finite three-group cover of Y_m again has three-group Galois group: intersect its finitely many conjugate subgroups, or use the associated three-group wreath product. This proves the comparison with the open subgroup of N. Apply (C.16.5) to Y_m, using its actual smooth compactification and full boundary. The composite cover of U has finite geometric group

$$E_m = N/(H_m^m[H_m, H_m]). \tag{C.16.6}$$

The denominator is characteristic in N, so this composite is geometrically Galois. Compute its finite deck group and take the fixed field of $E_m^m \gamma_3 E_m$; this defines X_n.

In $N/(N^m \gamma_3 N)$, the image of H_m is central and of exponent dividing m. Hence

$$H_m^m[H_m, H_m] \subset N^m \gamma_3 N. \tag{C.16.7}$$

The preimage of $E_m^m \gamma_3 E_m$ is therefore exactly $N^m \gamma_3 N$. The quotient maps induced by N give inverse isomorphisms $E_m/(E_m^m \gamma_3 E_m) \simeq N/(N^m \gamma_3 N)$. This proves the precise map in (C.16.1). Finite field automorphisms, fixed fields and the maps of their normalizations are computed by factoring finite algebraic equations.

Because m is odd, the Hall map

$$\begin{aligned} (V/m) \oplus (\textstyle\bigwedge^2 V/m) &\longrightarrow P_n, \\ (v, z) &\longmapsto \prod_i y_i^{v_i} \prod_{i<j} [y_i, y_j]^{\,z_{ij} - v_i v_j/2} \end{aligned} \tag{C.16.8}$$

is an explicit isomorphism, with the products in increasing order. Its source has multiplication

$$(v, z)(v', z') = (v + v', z + z' + \tfrac{1}{2} v \wedge v'). \tag{C.16.9}$$

Commuting successive generators verifies the map. Conversely this source is generated by the ten basis elements, has exponent m, and is of class two. The free-group map back is inverse on the generators. This proves both bijectivity and $|P_n| = m^{55}$. Abelian bases are read from the Kummer pairing with (C.16.3), and their lifts are found in the finite deck group. They can be chosen compatibly because $P_{n+1} \to P_n$ and the maps on their abelian kernels are surjective.

Embed the cover fields compatibly in the Puiseux field at O, using $s = z/x$ and fixed compatible symbols $s^{1/3^a}$. Galois acts on the algebraic coefficients and fixes these symbols. The characteristic subgroup in (C.16.1) is preserved. Equivalently there is a unique based isomorphism $\phi_{g,n} : {}^g X_n \to X_n$ carrying the conjugate chosen tangential fiber point to the chosen one. A deck transformation fixing a fiber point is the identity. Thus uniqueness gives $\phi_{gh,n} = \phi_{g,n}\, {}^g\phi_{h,n}$: these are actual arithmetic descent maps. They are computed from the finite function-field isomorphisms and the Puiseux branch condition. Compute coefficients until all finitely many branches are distinguished; distinct algebraic branches differ after finitely many terms. The coefficient action agrees with the path–Puiseux comparison of [HM19, §1.1, (1.1.1)–(1.1.3)], applied here to the actual finite covers of U.

For $A \in U(L)$ choose compatible points $p_{A,n}$ in $X_n(A)$ by the root equations (C.16.5) and the finite fixed-field maps. The functions may be chosen with valuation zero at every point above A: multiply them by mth powers with the required prescribed valuations, using

Riemann–Roch approximation. This leaves their classes in (C.16.3) unchanged. Alternatively, at a valuation mk, divide by a local parameter to the power mk and evaluate the resulting unit; its root is trivialized by that parameter to the power k. These are explicit isomorphisms of the Kummer equations. The based inclusions associated to $N^{3^{n+1}}\gamma_3 N \subset N^{3^n}\gamma_3 N$ supply the transition maps. Their maps on geometric fibers are surjective, allowing successive compatible choices.

Let deck transformations act on the right on fiber points, by composition with function-field automorphisms. Compute $c_{A,n}(g)$ from

$$\phi_{g,n}(g p_{A,n}) = p_{A,n}\, c_{A,n}(g). \tag{C.16.10}$$

All terms are algebraic points and maps already constructed. The descent identity gives

$$c_{A,n}(gh) = c_{A,n}(g)\rho_n(g)c_{A,n}(h). \tag{C.16.11}$$

For fixed n, the equations, deck maps, distinguishing Puiseux coefficients and fiber points lie in a finite Galois extension of L. Thus (C.16.10)–(C.16.11) are computed as functions on a finite Galois group. Formula (C.16.8) then extracts their coordinates.

Write the computed arithmetic action as

$$\rho_n(g)(v,z) = (R_g v, \textstyle\bigwedge^2 R_g\, z + \ell_g v). \tag{C.16.12}$$

Here ℓ_g is read from the ten images $\rho_n(g)y_i$ using (C.16.8). It is linear because the class-two addition $a+b = ab[a,b]^{-1/2}$ is recovered from the group law; automorphisms preserve this addition and the bracket. In particular the evaluated central cochain satisfies

$$\begin{aligned} z_{A,n}(gh) = z_{A,n}(g) + \textstyle\bigwedge^2 R_g\, z_{A,n}(h) + \ell_g(v_{A,n}(h)) \\ + \tfrac{1}{2} v_{A,n}(g) \wedge R_g v_{A,n}(h). \end{aligned} \tag{C.16.13}$$

This retains the off-diagonal arithmetic term as well as the tangential displacement.

Every finite quotient of $N/\gamma_3 N$ has finite three-power exponent, so it factors through some P_n. Therefore $N/\gamma_3 N \to \varprojlim_n P_n$ is an isomorphism: compactness proves surjectivity, and the kernel is tested in all finite quotients. The compatible finite cocycles thus give a continuous cocycle. The etale path comparison between the two fiber functors on each X_n is precisely (C.16.10); cofinality identifies its inverse limit with the point section. A different compatible fiber point changes the cocycle by the corresponding compatible geometric conjugation. This proves the asserted class. □

Statement C.17 (Proved). An evaluated nonzero central point coordinate. *Let $\alpha^3 = 2$, $\zeta = \zeta_3$, $\eta = \zeta_9$ with $\eta^3 = \zeta$, and $L = \mathbb{Q}(\zeta, \alpha)$. For $A = [1 : -\alpha : 1] \in U(L)$ an explicit quotient of P_1 of order 27 has zero abelian point coordinates and central coordinate*

$$-\kappa_L(\zeta) \in H^1(G_L, \mathbb{F}_3) \tag{C.17.1}$$

in the prescribed unit-tangent origin at O. This class is nonzero.

Proof. The functions $f = -x/z$ and $1 - f = (x+z)/z$ are units on U: the line $x + z = 0$ meets E only at Q_c, with multiplicity three. Pull back the cover

$$u^3 = f, \qquad v^3 = 1 - f, \qquad w^3 = A(u) := (1 - \zeta u)(1 - \zeta^2 u)^2. \tag{C.17.2}$$

It is the $l = 3, c = 1$ Heisenberg cover of [HM19, §2.1, (2.1.2), Theorem 2.1.4]. We verify its connected pullback and its deck maps. The valuations of $f, 1 - f$ at P, O are $(1, 0)$ and

$(-1,-1)$, of determinant -1. Their Kummer classes are therefore independent and the first two equations have degree nine. Since

$$\frac{A(\zeta u)}{A(u)} = \frac{(1-u)^3}{1-u^3}, \tag{C.17.3}$$

the following are automorphisms:

	u	v	w
τ	ζu	v	$(1-u)w/v$
σ	u	ζv	w
z	u	v	$\zeta w.$

(C.17.4)

The identity $\prod_{j=0}^{2}(1-\zeta^j u) = 1-u^3$ gives $\tau^3 = 1$; also $\sigma^3 = z^3 = 1$ and $[\tau,\sigma] = z$. If $A(u) = b^3$ in the degree-nine field, its commuting automorphisms would satisfy $\tau(b)/b = \zeta^i(1-u)/v$ and $\sigma(b)/b = \zeta^j$ for some i, j. Their two orders of application would then differ by ζ, a contradiction. Thus the full degree is 27. Every zero and pole of these equations lies over a deleted cusp, so the normalization is etale on U. Its exponent-three class-two deck group is an actual quotient of P_1.

At A choose

$$u = -1, \qquad v = \alpha, \qquad w = -\eta; \tag{C.17.5}$$

indeed $A(-1) = -\zeta$. At O, put $r = s^{1/3}$. The branches are

$$u = -r^{-1}, \qquad v = r^{-1}(1+s)^{1/3}, \qquad w = \eta^2 r^{-1}(1+O(r)). \tag{C.17.6}$$

The last coefficient follows from $r^3 A(-r^{-1})|_{r=0} = \zeta^2$; the remaining unit series has coefficients in $\mathbb{Q}(\zeta)$. For $g \in G_L$ with $g\eta = \zeta^e \eta$, raw conjugation moves (C.17.5) by z^e and the tangent branch by z^{2e}. The based descent removes the latter displacement. Formula (C.16.10) therefore gives $z^{e-2e} = z^{-e}$. The raw arithmetic action commutes with (C.17.4), and its based correction is central, so the arithmetic action on the finite group over L is trivial. Geometric conjugation fixes this central cocycle.

Finally $X^3 - 2$ is Eisenstein at the prime over two in $\mathbb{Q}(\zeta)$. The distinct cyclic degree-three extension $\mathbb{Q}(\eta)/\mathbb{Q}(\zeta)$ is unramified there. Consequently $\eta \notin L$ and $\kappa_L(\zeta) \neq 0$. The exact polynomial identities, the twenty-seven deck elements and their multiplication, both fiber displacements, and the discriminants -108 and -3^9 are checked in `code/bc_point_coordinates.py` and its supplied output. □

Statement C.18 (Proved). The point-image set and its exact central membership criterion. *The point-image subset of the class-two germ set is*

$$\left\{ \left[\left(\varprojlim_n v_{A,n}, \varprojlim_n z_{A,n} \right) \right] : A \in U(\overline{\mathbb{Q}}) \right\}, \tag{C.18.1}$$

where each finite function is algebraically evaluated by (C.16.3)–(C.16.13). This parametrization is injective, disjoint from the nine packets (C.10.2), and equivariant for $A \mapsto gA$. Its stabilizer at A is $G_{\mathbb{Q}(A)}$.

Given a section and an algebraic point A, choose a common number field L of definition. Let d_n be the section cocycle in P_n and $c_{A,n}$ the evaluated cocycle above. Their germs agree exactly when every finite set

$$\mathcal{T}_n = \{h \in P_n : d_n(g) = h^{-1} c_{A,n}(g) \rho_n(g) h \text{ for all } g \in G_L\} \tag{C.18.2}$$

is nonempty. Each test is a computation on a finite Galois group. Combined with the injective abelian point map C.9, it gives a complete central membership criterion for (C.18.1).

Proof. C.16 proves (C.18.1), and C.9 proves injectivity, disjointness and the stated stabilizer. Conjugating the equations, divisors, roots and based comparison maps of C.16 carries the output at A to the output at gA, exactly as in (C.3), proving equivariance.

The equations in (C.18.2) factor through a finite Galois extension containing the fields of both finite cocycles and their actions. They are therefore finite tests. The inverse images of $\mathcal{T}_n$ in the compact group $N/\gamma_3 N$ are closed, nonempty and nested. Their intersection provides one conjugator valid at every precision, hence in $N/\gamma_3 N$. Conversely such a conjugator supplies every finite solution. Equality of germs is equality already over L by C.2. This proves (C.18.2) without assuming compatible choices of finite conjugators.

We spell out the transport to the other central coordinates. The Kummer pairing in (C.16.3) identifies P_n^{ab} with the actual $V/3^n$. For cusp vectors it is the valuation pairing; for proper vectors it is the torsion-divisor pairing in (C.16.4). Use the eight non-O cusp vectors and two lifts of an oriented Tate basis. Actual cusp generators are obtained by their Puiseux monodromy on the finite covers. Thus the basis comparison with (E36.4) takes place in the actual deck groups. For two compatible generator choices with the same abelian basis, write $y_i' = y_i \exp(t_i)$, with $t_i \in \bigwedge^2 V$. Substitution into (C.16.8) gives the exact map from coordinates in the primed generators to coordinates in the unprimed generators:

$$(v, z) \longmapsto (v, z + \sum_i v_i t_i). \tag{C.18.3}$$

An abelian basis change B gives B on v and $\bigwedge^2 B$ on central brackets, followed by the computed linear correction (C.18.3).

The based filling map q_O gives

$$q_O(v, z) = (\pi v, \omega z + \lambda(v)), \qquad \omega = \bigwedge^2 \pi, \tag{C.18.4}$$

where $\lambda(v_i)$ is read from the actual image $q_O(y_i)$ in the chosen basis of C. These images are computed by the finite maps of covers. One may take the two generators of C to be $q_O(y_1), q_O(y_2)$ and the other eight y_i to be peripheral; then $\lambda = 0$. The change to the involution-compatible basis of C.13 is (C.18.3), obtained by applying that actual involution to the generators. The orientation is $k = [a, b] = -\delta$, as in (E36.5), B.3 and A.8.

For the E36 parameter over a fixed abelian cocycle α, conjugate the computed cocycle by an actual geometric lift $\widetilde{h}$ of $h \in V$, using $c(g) \mapsto \widetilde{h}^{-1} c(g) \rho(g) \widetilde{h}$. Its abelian change is $v(g) \mapsto v(g) + (R_g - 1)h$. The vector giving the prescribed α is unique because $H^0(G_L, V) = 0$. Write the prescribed C-reference as $(\beta(g), w_\beta(g))$. After this explicit comparison the E36 parameter is the class of

$$\omega z_A(g) + \lambda(v_A(g)) - w_\beta(g) \quad \text{in the generator } k. \tag{C.18.5}$$

The two C-cocycles have identical abelian coordinate β; their group difference is precisely this central difference by (C.16.9). Over a prescribed class-two lift of the resulting N^{cc}-cocycle, the remaining fiber parameter is likewise the difference of the central cochains in $\mathcal{K} = \ker \omega$, after actual conjugation makes the N^{cc}-cocycles equal. This proves that the construction evaluates both the E36 parameter and the forty-four-dimensional fiber coordinate with their specified reference translations.

C.2–C.3 identify the germs fixed by G_L with Sec_L. In particular $\mathrm{Sec}_{\mathbb{Q}}$ is the entire $G_{\mathbb{Q}}$-fixed subset. Equations (C.10.2) and (C.18.1) supply both geometric image sets required in Task C.3. The criterion ranges over all n; it does not assert a bounded finite stopping rule for equality at all precisions. □

Statement C.6 (Proved). Complete class-two germ classification. *Task C.1 requests the full explicit residue complex, obstruction residues, convergence criterion, and fibers over*

every number field L. C.11 disproves a verbatim pure-residue extension of Part A; C.12 gives the exact replacement, including the finite-S_L arithmetic class and the lift torsor. C.13 evaluates the preceding cuspidally central obstruction by the local quadratic pairings at the places above three, with their global sum relation. C.14 computes the finite-S_L complexes by finite Galois cochain matrices and Artin–Verdier duality, and C.15 evaluates the retained class by explicit compact-support cup traces. Together with the restricted-family condition in C.12, these give the full number-field criterion and fibers requested in Task C.1. For Task C.2, C.1–C.3 give injective restriction, the actual Galois action and exact descent. Formula (C.5) identifies Sec_L *with all G_L-fixed germs. For Task C.3, C.8–C.9 prove injectivity of the point map and separation from cusp germs. C.10 gives the nine disjoint cusp parameter spaces with their torsion and Galois action. C.16 evaluates every point's abelian and central coordinates by finite algebraic operations at each precision; C.17 is an explicit nonzero central instance. C.18 gives the exact point-image set, its equivariance and its full central membership criterion. Its coordinate maps retain the E36 origin and the lift-fiber translation. Finally C.4–C.5 and C.7 prove that both specified phantom classes remain nonzero after every finite number-field extension and in the germ colimit, settling Task C.4.*

Proof. The cited statements provide the specified maps, arithmetic computations and geometric image sets. Their compatibility under finite extension is the cochain restriction of C.2 and the based fiber construction of C.16; hence they give one classification over the filtered system of number fields, as asserted. □

3.3 Separating the point germs by the nine tangent lines

The following calculation gives additional geometric information about Task C.3 directly in the abelian quotient. We work with the actual maps on fundamental groups induced by regular invertible functions on U. Their coefficient field is $K_0 = \mathbb{Q}(\zeta)$; enlarging a field of definition to contain K_0 is a restriction map of section sets, hence is permitted in the germ colimit.

Statement C.8 (Proved). An exact tangent-line separation identity. *Let ℓ_d be a row vector defining the flex tangent line at $d \in D$. The nine rows can be chosen as*

$$(1, \zeta^a, 0), \quad (0, 1, \zeta^a), \quad (\zeta^a, 0, 1) \qquad (a = 0, 1, 2).$$

Set $P_d = \ell_d^* \ell_d / 2$, a rank-one Hermitian projector on $\mathbb{C}^3$. The real-linear map

$$\mathrm{Herm}_3(\mathbb{C}) \longrightarrow \mathbb{R}^9, \qquad H \longmapsto \big(\mathrm{tr}(P_d H)\big)_d$$

is an isomorphism, and

$$\sum_d P_d = 3I_3, \qquad \mathrm{tr}(P_d P_e) = \begin{cases} 1 & d = e, \\ 1/4 & d \neq e. \end{cases} \tag{C.8.1}$$

The kernel after deleting coordinate d is the line $\mathbb{R}(4P_d - I_3)$. A positive semidefinite matrix with the eight undeleted measurements of vv^*, where $v \neq 0$ and $\ell_d(v) = 0$, is exactly vv^*.

Proof. Each row has norm squared two. The scalar product of two distinct rows has absolute value one: for rows with the same two nonzero positions this follows from $|1+\zeta| = |1+\zeta^2| = 1$; for different positions their supports meet in one coordinate. Summing the projectors cancels the off-diagonal terms and gives three on each diagonal entry. This proves (C.8.1). Their

Gram matrix is $(3I_9 + J_9)/4$, with eigenvalues 3 and 3/4, so the nine projectors are a real basis of the nine-dimensional Hermitian space. Moreover

$$\operatorname{tr}\bigl(P_e(4P_d - I_3)\bigr) = 3\delta_{de},$$

which proves the assertion about the deleted coordinate.

Write a matrix with the prescribed eight measurements as $vv^* + t(4P_d - I_3)$, $t \in \mathbb{R}$. The vector spanning $\operatorname{im} P_d$ is orthogonal to v, and its quadratic value is $3t$ times its norm squared. A nonzero vector orthogonal to both this line and v has quadratic value $-t$ times its norm squared. Positive semidefiniteness therefore gives both $t \geq 0$ and $t \leq 0$. Thus $t = 0$. The exact program `code/c_point_separation.py` verifies the rank, all 81 dual pairings, all nine tangent annihilations, and the eigenvalues $(3, -1, -1)$ over $\mathbb{Q}(\sqrt{-3})$. □

Statement C.13 (Proved). The initial cuspidally central obstruction over every number field. *Let $L \subset \overline{\mathbb{Q}}$ be a number field and let $\alpha \in H^1(G_L, V)$, with image $\beta \in H^1(G_L, T)$. Orient $\bigwedge^2 T \simeq \mathbb{Z}_3(1)$ by the commutator in the once-punctured elliptic curve, and write ω for this alternating pairing. For each $v \mid 3$ define*

$$q_v(\beta) = \tfrac{1}{2} \operatorname{inv}_v \omega_*(\beta_v \smile \beta_v) \in \mathbb{Z}_3. \tag{C.13.1}$$

Here $\operatorname{inv}_v : H^2_{\mathrm{cts}}(G_{L_v}, \mathbb{Z}_3(1)) \xrightarrow{\sim} \mathbb{Z}_3$ is the inverse limit of the normalized local Brauer invariant maps. Then

$$\alpha \in \operatorname{im}\bigl(\operatorname{Sec}_L(N^{\mathrm{cc}}) \to H^1(G_L, V)\bigr) \quad \Longleftrightarrow \quad q_v(\beta) = 0 \text{ for every } v \mid 3. \tag{C.13.2}$$

The tuple satisfies $\sum_{v|3} q_v(\beta) = 0$. Every nonempty fiber is a torsor under $H^1(G_L, \mathbb{Z}_3(1)) = \varprojlim_n L^\times/(L^\times)^{3^n}$, with no further stabilizer quotient. In particular, every abelian section lifts to N^{cc} over a number field having exactly one place above three, including $\mathbb{Q}$. This statement makes no assertion that every tuple subject to the displayed sum occurs as a self-pairing.

Manuscript comparison: §51.16, folios 733v–736v, scans 172–178, formulas (371)–(379).

Proof. We first specify the genuine arithmetic action on the once-punctured elliptic class-two quotient C. The automorphism $[x : y : z] \mapsto [y : x : z]$ is elliptic inversion for origin O. It preserves $E \setminus \{O\}$ and has derivative -1 at O. Its comparison from the tangent -1 to the tangent 1 admits a $G_{\mathbb{Q}}$-fixed pro-three path: the compatible 3^nth roots of -1 can all be chosen to be -1. Composition with this path gives a genuine involution j of the based pro-three group, commuting with the arithmetic action. Applying the comparison twice gives the constant path, as the product of the two chosen roots is one at every level.

The integral class-two logarithm is available at three: its group law is the Baker–Campbell–Hausdorff law with the single denominator two. The central exact sequence of Lie lattices is

$$0 \longrightarrow \bigwedge^2 T \longrightarrow \log C \longrightarrow T \longrightarrow 0.$$

The involution acts as -1 on T and as 1 on the center. The operator $(1 - j)/2$ therefore gives a canonical lift of T to its negative eigenspace. An arithmetic automorphism commutes with j, so it preserves this lift. In these coordinates

$$(t, z)(t', z') = (t + t', z + z' + \tfrac{1}{2}\omega(t, t')), \qquad g(t, z) = (gt, \chi_3(g)z). \tag{C.13.3}$$

This proves the claimed diagonalization by an actual based morphism. In particular the central extension obstruction for a T-cocycle β is represented by the continuous two-cocycle

$$(g, h) \longmapsto \tfrac{1}{2}\omega(\beta(g), g\beta(h)). \tag{C.13.4}$$

At a finite place $v \nmid 3$, the curve has good reduction, so T is unramified. The two Frobenius eigenvalues have complex absolute value $\sqrt{Q_v}$. Thus both $T^{F_v=1}$ and $T(-1)^{F_v=1}$ vanish. The local inertia sequence gives

$$H^1(G_{L_v}, T) = T/(F_v - 1)T,$$

a finite group. This equality also follows at the integral level by the finite-coefficient inverse limit; the invariant towers are finite and have zero first derived limit. The self-pairing of any element of a finite group is torsion, whereas $H^2_{\mathrm{cts}}(G_{L_v}, \mathbb{Z}_3(1)) = \mathbb{Z}_3$ is torsion-free. Consequently (C.13.4) restricts to zero at every such place. Its restrictions at archimedean places vanish because three is odd.

For clarity, the required global injectivity retains the topology. At each finite level the Kummer sequence identifies $H^2(G_L, \mu_{3^n})$ with $\mathrm{Br}(L)[3^n]$. The global Brauer invariant exact sequence, followed by inverse limit, identifies its limit with

$$\left\{ (b_v)_v \in \mathcal{R}_{v\ mfinite}(\mathbb{Z}_3) : \sum_v b_v = 0 \right\}. \tag{C.13.5}$$

The sum converges because the family is restricted. Conversely, reduction of such a tuple modulo 3^n has finite support and zero sum, and hence gives the unique global Brauer class at that level; these classes are compatible. Statement F.5 gives $\varprojlim{}^1 H^1(G_L, \mu_{3^n}) = 0$, so Jannsen's sequence identifies the continuous integral second cohomology itself with (C.13.5). Equivalently one can apply the same Kummer inverse-limit argument to the completed multiplicative group. The local and global Brauer invariant theorems are used in their stated normalizations [Mil20b].

The only possibly nonzero entries of (C.13.4) are therefore precisely (C.13.1). Formula (C.13.5) proves both the sum relation and the vanishing equivalence. The arithmetic fiber product $N^{\mathrm{cc}} = C \times_T V$ transports this equivalence to (C.13.2). The central lifting fiber is acted on transitively by $H^1(G_L, \mathbb{Z}_3(1))$. Its potential stabilizer is the image of $H^0(G_L, V)$, which is zero: a good Frobenius outside three has no eigenvalue one on $V \otimes \mathbb{Q}_3$, since its boundary and elliptic eigenvalues have absolute values Q_v and $\sqrt{Q_v}$. As V is torsion-free, the integral invariants vanish as well. This proves the assertion about fibers. A tuple supported at one place and having sum zero is zero, which proves the last claim. □

Statement C.9 (Proved). Injective point germs and their separation from cusp germs. *The canonical map*

$$U(\overline{\mathbb{Q}}) \longrightarrow \varinjlim_{L} \mathrm{Sec}_L(N/\gamma_3 N)$$

is injective and $G_{\mathbb{Q}}$-equivariant. Its image is disjoint from all the cuspidal germs. Both assertions already hold after projecting to the germ colimit of $H^1(G_L, V)$. The smallest field of definition of a point germ is the field of definition of the point.

Proof. For distinct cusps d, e, the regular invertible function $f_{d,e} = \ell_d/\ell_e$ on U has divisor $3d - 3e$ on E. Its morphism $U \to \mathbb{G}_m$ gives a continuous map of geometric pro-three groups $N \to \mathbb{Z}_3(1)$, factoring through V. On the difference of the section classes of two points $A, B \in U(L)$, the induced cohomology map is exactly

$$\kappa_L(f_{d,e}(A)/f_{d,e}(B)). \tag{C.9.1}$$

This follows by pulling back the covers $z^{3^n} = t$ on $\mathbb{G}_m$; their fibers are the compatible division roots used to define the Kummer cocycle. Thus the comparison is a map of actual section torsors, with the tangential origin cancelling in the difference.

The kernel of $L^\times \to \varprojlim_n L^\times/L^{\times 3^n}$ is its group of roots of unity of order prime to three. Indeed every finite valuation of an infinitely three-divisible element is zero. Dirichlet's unit theorem then reduces the assertion to the finite roots-of-unity group. In particular every complex absolute value of a kernel element is one.

Equality of two abelian germs is equality after restriction to one finite number field containing A, B and K_0. By (C.9.1) the ratios of all these function values have absolute value one. Fix a complex embedding and normalize vector representatives a, b of A, B by $\ell_e(a) = \ell_e(b) = 1$. This is possible because the tangent at a flex meets E only at that cusp and $A, B \in U$. We obtain $|\ell_d(a)|^2 = |\ell_d(b)|^2$ for all nine d. C.8 gives $aa^* = bb^*$, hence the same projective point $A = B$.

Now compare a point $A \in U$ with a section in the packet of a cusp d. Choose $e \neq d$. For every $j \neq d$, the function $f_{j,e}$ has order zero at d and is a unit in its local ring. The morphism on the tangential packet is consequently the constant Kummer class of $f_{j,e}(d)$: extracting roots of a local unit gives that constant fiber, and the varying tangent parameter has exponent zero. Equality of abelian germs would therefore give, after normalization at ℓ_e,

$$|\ell_j(a)|^2 = |\ell_j(d)|^2 \quad (j \neq d).$$

The cusp vector satisfies $\ell_d(d) = 0$. The final assertion of C.8 applied to the positive matrix aa^* forces $A = d$, contradicting $A \in U$. This proves disjointness for every algebraic cusp and every completed Kummer parameter.

Equivariance follows from functoriality of the point section and restriction. Injectivity identifies the stabilizer of a point germ with the stabilizer of the point, which proves the assertion about its smallest field of definition. □

3.4 Finite arithmetic matrices over an arbitrary number field

Statement C.14 (Proved). An explicit finite Galois construction of the retained arithmetic groups. *Fix a number field $L \subset \overline{\mathbb{Q}}$ and put $S = \{v : v \mid 3\} \cup \{v : v \mid \infty\}$. For $n \geq 1$ write*

$$A_n = \mathcal{K}/3^n, \qquad A_n^D = \mathrm{Hom}(A_n, \mu_{3^n}), \qquad F_n = L(J_{E,D}[3^n], \mu_{3^n}), \qquad d_n = 3^{44n}[F_n : L]. \tag{C.14.1}$$

The homomorphisms in the dual have their usual contragredient Galois action. Let Ω_n be the compositum, inside the fixed algebraic closure, of all finite Galois extensions E/L containing F_n, unramified outside S, and satisfying $[E : L] \leq d_n$. Put $G_n = \mathrm{Gal}(\Omega_n/L)$. At each $v \mid 3$ choose a prolongation to Ω_n and write $D_{n,v} \leq G_n$ for its decomposition subgroup. These are finite fields and finite groups, obtained by the bounded enumeration in the proof.

For a finite G_n-module A, use the finite matrices

$$\begin{aligned} (\delta^0 b)(g) &= gb - b, \\ (\delta^1 z)(g,h) &= gz(h) - z(gh) + z(g). \end{aligned} \tag{C.14.2}$$

Define the following finite abelian groups by kernels and images of these matrices:

$$U_n = \ker\bigl(\delta^1 : \mathrm{Map}(G_n, A_n) \to \mathrm{Map}(G_n^2, A_n)\bigr)/\delta^0 A_n, \tag{C.14.3}$$

and

$$\begin{aligned} C_n = \{(z, (b_v)_{v\mid 3}) : \delta^1 z = 0,\ b_v \in A_n^D, \\ z(g) = gb_v - b_v \text{ for } g \in D_{n,v}\}/\{(\delta^0 b, (b)_{v\mid 3}) : b \in A_n^D\}. \end{aligned} \tag{C.14.4}$$

In (C.14.4), z ranges over all functions $G_n \to A_n^D$. There are canonical identifications

$$\begin{aligned} H^1(G_{L,S}, A_n) &\simeq U_n, \qquad & H^2(G_{L,S}, A_n) &\simeq C_n^\vee, \\ H^1_{\mathrm{cts}}(G_{L,S}, \mathcal{K}) &\simeq \varprojlim_n U_n, \qquad & H^2_{\mathrm{cts}}(G_{L,S}, \mathcal{K}) &\simeq \varprojlim_n C_n^\vee. \end{aligned} \tag{C.14.5}$$

Here $C_n^\vee = \mathrm{Hom}(C_n, \mathbb{Q}/\mathbb{Z})$; the transition maps are induced by coefficient reduction and its Cartier dual. Thus the arithmetic terms of C.12 are specified by finite arithmetic fields and finite matrices at every precision. The construction applies to every L; it does not assume that either displayed inverse limit is zero.

Proof. The reduction A_n is a submodule of $\bigwedge^2(V/3^n)$: the quotient $\bigwedge^2 T$ in the definition of $\mathcal{K}$ is torsion-free. The finite group scheme $J_{E,D}[3^n]$ therefore splits its action. Adjoining μ_{3^n} also splits the Cartier dual. Both coefficients are unramified outside S, because the original generalized Jacobian and its three-adic representation are unramified outside three. All the fields in (C.14.1) are algebraic data of the specified curve. For example, multiplication by 3^n on its framed divisor classes constructs $J_{E,D}[3^n]$; Riemann–Roch and factorization give its finite defining equations and their splitting field. No presentation of an unknown infinite Galois group is an input to the construction.

We give a bound for the field enumeration. For any integer $d \geq 1$, every extension E/L of degree at most d unramified outside S has

$$|\operatorname{disc}(E)| \leq B_L(d) := |\operatorname{disc}(L)|^d \prod_{v \mid 3} (Nv)^{d(1+e_v \lfloor \log_3 d \rfloor)}, \tag{C.14.6}$$

where e_v is the ramification index of $L_v/\mathbb{Q}_3$. Indeed, pass first to the maximal unramified subextension of a completion E_w/L_v. Its remaining totally ramified part has ramification index e and an Eisenstein polynomial P for a uniformizer π. The terms of $P'(\pi)$ have pairwise distinct valuations modulo e. Its valuation is therefore their minimum and is at most $e - 1 + ee_v v_3(e)$. Multiplying by the residue degree and summing over $w \mid v$ bounds the relative discriminant exponent by $d(1 + e_v \lfloor \log_3 d \rfloor)$. The tower formula for the discriminant proves (C.14.6).

Hermite's proof, in the explicit form of Milne, Theorems 8.42–8.43 [Mil20a], bounds the conjugates of a primitive integral element in terms of the absolute discriminant. Consequently one can list a finite collection of monic integral polynomials of degree at most $d[L : \mathbb{Q}]$ that contains a defining polynomial for every E in question. For a deliberately generous choice, a bound $2(B_L(d) + 1)$ for every conjugate suffices in that proof; the coefficient of degree $r - i$ is then bounded by $\binom{r}{i}[2(B_L(d) + 1)]^i$. Enumerate these polynomials, factor them, calculate rings of integers and discriminants, and retain precisely their embeddings over L that are Galois, contain F_n, have the required degree, and are unramified away from S. All these tests are finite exact number-field operations. Taking their compositum constructs Ω_n. The bound is used to prove termination; no feasibility estimate is asserted.

Let A be either A_n or A_n^D, and let $\rho : G_{L,S} \to \mathrm{Gal}(F_n/L)$ be the quotient. Every continuous crossed homomorphism $z : G_{L,S} \to A$ gives the continuous homomorphism

$$g \longmapsto (z(g), \rho(g)) : G_{L,S} \longrightarrow A \rtimes \mathrm{Gal}(F_n/L). \tag{C.14.7}$$

Its kernel cuts out a Galois extension containing F_n of degree at most $|A|[F_n : L] = d_n$, unramified outside S. It is one of the fields used in Ω_n. Thus every crossed homomorphism factors through G_n. Conversely every crossed homomorphism on G_n inflates to one on $G_{L,S}$, and the coboundaries on the two groups are exactly the same functions $g \mapsto gb - b$. This proves (C.14.3) and its analogue for A_n^D.

Use the compact-support complex

$$R\Gamma_c(\mathcal{O}_L[1/3], A_n^D) = \mathrm{Cone}\left(R\Gamma(G_{L,S}, A_n^D) \longrightarrow \bigoplus_{v\in S} R\Gamma(L_v, A_n^D) \right)[-1]. \tag{C.14.8}$$

The local complexes at infinite places here are the modified Tate complexes. They are acyclic for these odd-primary coefficients, including at a real place. Degree-one cycles in (C.14.8) are exactly a global one-cocycle together with the indicated local degree-zero trivializations. A change by a global zero-cochain is exactly the denominator of (C.14.4). Since the global cocycle factors through G_n, each local equation holds on G_{L_v} exactly when it holds on its image $D_{n,v}$. We have proved the explicit identification of C_n with that compact-support group.

Milne's Corollary II.3.3(a),(b) [Mil06], applied to the locally constant sheaf A_n^D, identifies its derived Cartier dual with A_n and gives the perfect pairing

$$H^2(G_{L,S}, A_n) \times C_n \longrightarrow \mathbb{Q}/\mathbb{Z}. \tag{C.14.9}$$

The Galois-to-etale comparison is the same II.2.9 used in C.11. This proves the second finite identification. Every finite group in degrees one and two is finite, so each coefficient tower is Mittag–Leffler. The continuous cohomology inverse-limit sequences therefore have zero first derived-limit term at finite S and give the last two identifications in (C.14.5). Coefficient naturality of (C.14.7)–(C.14.9) specifies all transition maps.

For a small arithmetic verification of the matrix convention, the program `c_arithmetic_matrices.py` in `code/` constructs the actual group of $\mathbb{Q}(\zeta_9, \sqrt[3]{3})/\mathbb{Q}$, of order 18. Over K_0 the two Kummer classes $\zeta, 3$ are independent and span the S-unit classes: $3 = -\zeta^2\lambda^2$. Thus this quotient captures all one-cocycles for $\mathbb{F}_3$ and μ_3 over $\mathbb{Q}$, including their coboundaries. Its decomposition group at three is the whole group. Indeed a Kummer line involving 3 has valuation nonzero modulo three over $K_{0,3}$; the remaining line ζ gives the ramified cyclotomic cubic extension. Hence no nontrivial cubic subextension is unramified there. In coordinates the group law is $(a,b,c)(a',b',c') = (a+a', b+(-1)^c b', c+c')$, with the first two coordinates modulo three and the last modulo two. The program verifies this law and both coefficient actions. The respective cocycle matrix ranks are 17 and 16, the coboundary ranks are 0 and 1, and both first-cohomology dimensions are one. Both compact-support dimensions are zero. A separately labelled duplicated-subgroup fixture checks the local-trivialization quotient; it is not assigned to a number field. These are checks of the finite matrices, with their complete output retained; the proof above establishes their general arithmetic use. □

Statement C.15 (Proved). The complete number-field class-two test with its finite arithmetic evaluations. *Let s be a continuous section of N^{cc} over L. Use the actual obstruction two-cocycle o and its origin-corrected residue family $r = (r_v)_{v\nmid 3,\infty}$ from A.8, restricted to G_L. Write $D_v = \mathrm{Frob}_v - 1$ on $\mathcal{K}(-1)$. First require*

$$y_v = -D_v^{-1} r_v \in \mathcal{K}(-1), \qquad (y_v)_v \in \mathcal{R}_L(\mathcal{K}(-1)). \tag{C.15.1}$$

The actual localization homotopy-fibre cycle (o, y) then supplies, at each n, a two-cocycle $\xi_n \in Z^2(G_{L,S}, A_n)$, by the comparison below. For every class represented by $(z, (b_v))$ in the explicit finite group C_n of C.14, choose a continuous two-cochain a with values in μ_{3^n} satisfying

$$da = \xi_n \cup z. \tag{C.15.2}$$

Such a cochain always exists. Define

$$\ell_n(s)(z,(b_v)) = \sum_{v|3} \mathrm{inv}_v\big((\mathrm{res}_v\,\xi_n)\cup b_v - \mathrm{res}_v\,a\big). \tag{C.15.3}$$

The expression in each local invariant is a two-cocycle. Formula (C.15.3) is independent of all the auxiliary cochain choices and is an element of $C_n^\vee$. There is an exact criterion

$$s \text{ lifts to } N/\gamma_3 N \quad\Longleftrightarrow\quad \text{(C.15.1) holds and } \ell_n(s)=0 \text{ for every } n\ge 1. \tag{C.15.4}$$

Every nonempty fibre is a torsor under $\varprojlim_n U_n$, with U_n given by the finite matrices (C.14.3), and no further geometric-stabilizer quotient. Together with C.13 for the initial lift, C.14–C.15 give the number-field version requested in Task C.1: both the residue convergence and the arithmetic test are retained.

Proof. We specify the comparison that enters (C.15.2). Put $\mathcal{A}_v = \mathcal{K}(-1)$ and $R^\bullet = [\mathcal{R}_L(\mathcal{A}) \xrightarrow{D} \mathcal{R}_L(\mathcal{A})]$ in degrees zero and one. The localization triangle of F.1 provides a continuous cochain model and a quasi-isomorphism

$$R\Gamma(G_{L,S},\mathcal{K}) \longrightarrow \mathrm{Cone}(R\Gamma(G_L,\mathcal{K}) \longrightarrow R^\bullet[-1])\,[-1]. \tag{C.15.5}$$

Choose the cone sign so that a degree-two cycle is (o,y) with $do=0$ and $r+Dy=0$. This is the sign in (C.15.1). All restrictions, residue maps, and homotopies are the actual maps in the localization construction; in particular y is retained as part of the cycle. Reduction modulo 3^n in (C.15.5) gives its finite coefficient comparison. A representative ξ_n is any finite-coefficient two-cocycle on the left whose image is cohomologous to the reduction of (o,y) on the right. It exists because (C.15.5) is a quasi-isomorphism. Its cohomology class is uniquely determined. This prescription keeps the arithmetic class in the homotopy fibre, including when the finite-coefficient map from that class to global cohomology has a kernel.

There is a finite-data method for obtaining such representatives at any specified n. Continuous finite-valued cochains factor through finite Galois quotients. Enumerate finite S-ramified quotients and their normalized bar two-cocycles. For the comparison homotopies, enumerate finite global quotients with ramification allowed in a finite set T containing the ramification of o_n and the support of y_n; enumerate larger finite sets T as well. Use the tame resolution at the finite residue support. Thus only the arithmetic representative is required to factor through an S-ramified quotient; the global comparison homotopy retains its necessary ramification. Test the cochain identities in the finite multiplication tables. For the prescribed cycle a matching representative and homotopy occur, so this search terminates. The inverse in (C.15.5) is thereby a cochain comparison, not a choice of a splitting of the short exact sequence (C.12.1). At a finite coefficient level the residue support is finite: this is precisely where the restricted condition in (C.15.1) is used. As with C.14, this is an exact construction and no claim of practical running time is made.

The group $G_{L,S}$ has three-cohomological dimension at most two. Therefore $H^3(G_{L,S},\mu_{3^n}) = 0$, which proves existence in (C.15.2). Its right side is the standard evaluation cup product for $A_n \otimes A_n^D \to \mu_{3^n}$. Since $db_v = \mathrm{res}_v\, z$ and $d\xi_n = 0$,

$$d\big((\mathrm{res}_v\,\xi_n)\cup b_v - \mathrm{res}_v\,a\big) = 0.$$

We can obtain a by enumerating finite S-ramified quotients refining those through which ξ_n, z factor and solving the finite linear cochain equation. The vanishing just proved guarantees termination of this search as well.

In the compact-support cone (C.14.8) the product of ξ_n and $(z,(b_v))$ is represented by

$$\big(\xi_n \cup z, ((\mathrm{res}_v\,\xi_n)\cup b_v)_v\big).$$

Subtracting the coboundary of $(a, 0)$ gives $(0, ((\mathrm{res}_v\, \xi_n) \cup b_v - \mathrm{res}_v\, a)_v)$. Normalize the compact-support trace by the sum of the local Brauer invariants on such a representative. This is the trace compatibility in Milne II.3, preceding Theorem 3.1 [Mil06]. The resulting trace is exactly (C.15.3). Each local invariant is that of a finite crossed-product algebra, so it is determined by finite extension, norm, and valuation data over L_v. The construction proves directly that (C.15.3) is the perfect pairing (C.14.9), applied to the class of ξ_n. It consequently descends to C_n and is independent of the representatives and of a. In particular, changing a by a global two-cocycle changes the sum by zero, by the global Brauer invariant relation.

At the integral level, injectivity of each D_v proves that (C.15.1) is exactly the vanishing of the residue obstruction, with its unique integral correction. Under that condition the class of (o, y) is the unique arithmetic class of C.12. By (C.14.5), this class vanishes exactly when all its finite components pair to zero with C_n. Those components are exactly (C.15.3). This proves both directions of (C.15.4). The finite-S first-cohomology inverse-limit identification and the stabilizer calculation of C.12 give the stated torsor and its precise conjugacy quotient. The global Milnor term is still the one computed in F.1–F.3; the additional finite arithmetic evaluations introduce no new first derived limit. Thus the criterion includes infinitely ramified classes, all torsion classes, every prime above three, and the modified real-place convention. □

4 D. Profinite leading terms

4.1 Sources, comparison maps, and the two remaining image problems

We use the generators and positive tangential normalization of the task. The source question is LM7 D.13, developing the comparison of groups in *La Longue Marche*, §51.15 (cote 140-4, scans 156–170). Only the supplied transcription, not the underlying scans, was available for this comparison. We make no assertion of a new reading of those scans.

Put $F = \langle x, y\rangle$, let $\widehat{F}$ be its profinite completion, and write F_ℓ for its maximal pro-ℓ quotient. The quotient $q_\ell : \widehat{F} \twoheadrightarrow F_\ell$ has characteristic kernel. It therefore induces a continuous homomorphism on outer automorphism groups. Let I^n denote the filtration obtained by acting trivially, up to inner automorphism, on $F_\ell/\Gamma_{n+1}F_\ell$. We filter each full profinite source by inverse images of I^n. The difference of an automorphism and the identity, read in $\Gamma_{n+1}/\Gamma_{n+2}$, is a derivation. Composition becomes addition in this quotient, and conjugating the automorphism changes that derivation by an inner derivation. Thus we obtain the specified maps

$$(I^nG/I^{n+1}G) \otimes_{\mathbb{Z}_\ell} \mathbb{Q}_\ell \longrightarrow \mathrm{OutDer}_{\mathrm{sp}}(L_\ell)_n \longrightarrow \mathrm{OutDer}(L_\ell/L_\ell'')_n, \qquad L_\ell = \mathrm{Lie}_{\mathbb{Q}_\ell}(x, y).$$

Here G is one of the full profinite source groups, and the first tensor product means the rational span of its image in the integral graded automorphism module. Expanding each peripheral conjugacy equation gives the special-derivation equation. Expanding the commuting equations with the geometric S_3 gives invariance. These are necessary conditions on the images of the displayed maps.

Statement D.1 (Open). The weight-five image of the full profinite special group. *For $\ell \geq 5$, the dimension requested in Task D.1 has not been determined here. The target computed in R8 is*

$$(X + Y)\,\langle X^2, XY, Y^2\rangle_{\mathbb{Q}_\ell}[x, y].$$

Statement D.21 proves that this image has dimension either 1 *or* 3*: the actual geometric S_3 preserves it, and its invariant gamma line is already present by D.20. The remaining choice between* 1 *and* 3 *is not determined here.*

Statement D.2 (Open). The extra weight-nine direction for the full profinite gentle group. *For $\ell \geq 5$, we have neither constructed an element of $\widehat{\mathrm{GT}}_0$ with leading metabelian coordinate outside $\mathbb{Q}_\ell G_9$, nor proved that every such coordinate belongs to that line, where*

$$G_n = \frac{(X+Y)^n - X^n - Y^n}{XY}, \qquad p_{\mathrm{extra}} = X^2Y^2(X+Y)^3.$$

The requested choice between dimensions 1 and 2 remains open in this return.

Statement D.3 (Proved). The finite pentagon comparison in the weight-nine stratum. *At the compatible quotients $PB_n/(\gamma_{11}PB_n\, PB_n^{101})$, for $n = 3, 4$, the complete multiplier-one, weight-at-least-nine gentle shadow stratum has 101^{37} elements. Exactly 101^2 of these satisfy the complete pentagon. Their degree-nine leading spaces have dimensions four and one, respectively; their metabelian images have dimensions two and one. The latter image is the line generated by G_9. Statements D.7–D.8 give the finite comparison and an exact integer certificate. Its characteristic-zero comparison, together with the actual Galois lower bound in D.20, derives R9 in weight nine. The enumeration is of the entire stratum that can contribute a weight-nine leading term, with its degree-ten corrections; it is not an enumeration of other multipliers or of lower leading weights.*

4.2 What the linear and literature checks establish

Statement D.4 (Proved). Recomputed targets and the checked pentagon upper bound. *The original rational program reproduces R8's four rows in weights $1, \ldots, 10$. In weight nine its invariant metabelian image has basis $p_0 = p_{\mathrm{extra}}$ and*

$$p_1 = 2X^7 + 8X^6Y + 17X^5Y^2 + 23X^4Y^3 + 23X^3Y^4 + 17X^2Y^5 + 8XY^6 + 2Y^7, \qquad G_9 = \frac{15}{2}p_0 + \frac{9}{2}p_1.$$

The leading metabelian image of full $\widehat{\mathrm{GT}}$ is contained in $\mathbb{Q}_\ell G_n$ in positive odd weight n, and is zero in positive even weight. In particular p_{extra} is excluded in weight nine by this implication from the pentagon.

Proof. We reran the supplied programs, retaining their provenance under `code/de_source/`; this is a reproducibility check rather than an independently implemented rational Lie calculation. Their complete exact rational outputs and source witnesses are included. The rows are

n	1	2	3	4	5	6	7	8	9	10
special	0	0	1	0	3	0	6	4	13	12
metabelian image	0	0	1	0	3	0	5	0	7	0
S_3-invariants	0	0	1	0	1	0	2	1	4	2
their image	0	0	1	0	1	0	1	0	2	0.

The calculation solves $[f, y] = [g, x + y]$ in the free Lie algebra, then applies the canonical quotient $L \to L/L''$ to its solutions. This order of operations specifies the image being counted.

For the upper bound we use the opened primary text of Furusho, Corollary 0.3, Theorem 0.5, Proposition 5.1 and Lemma 5.2 [Fur11]. The leading homogeneous term of a profinite pentagon equation, after applying q_ℓ and the continuous rational unipotent completion, satisfies the infinitesimal pentagon: multiplication of factors of the same least positive degree becomes addition, and the standard braid insertion maps induce the corresponding insertion maps on the graded Lie algebras. Thus this leading term lies in $\mathfrak{grt}_1(\mathbb{Q}_\ell)$. Furusho's result

implies its gamma identity. We spell out the coordinate conversion. In the completed tensor algebra write $h = h_x x + h_y y$, and put $M(h) = (h_y y)^{\mathrm{ab}}$. Expansion gives

$$M([x,y]) = XY, \quad M([x,h]) = XM(h), \quad M([y,h]) = YM(h) \quad (h \in L').$$

Moreover $M([L',L']) = 0$, because the commutative images of both derived factors vanish. Hence $M(p[x,y]) = XYp$ on L'/L''. The homogeneous gamma identity reads $M(h) = a_n(X^n + Y^n - (X+Y)^n)$. Cancellation of XY gives the asserted line. The special equation implies $(X+Y) \mid p$: in L'/L'' it reads $-Yp = -(X+Y)q$, and $X+Y$ is relatively prime to Y. For even n, evaluating the numerator at $Y = -X$ makes $a_n = 0$. Weight one is zero directly. The rerun gamma verifier checks this Fox coordinate identity on all 224 Lyndon words in degrees two through ten and the displayed independent weight-nine vectors. □

The lower bound in R9 is established in D.20 from the opened primary Ihara formula, with its exact Fox coordinate and an extraction inside the actual compact Galois image. The initial Hain–Matsumoto audit reached Theorem 8.4, §8.4, Lemma 8.5 and Remark 8.6 [HM03]; that remark explicitly points to Ihara for nontriviality. The subsequent primary-source check opens Ihara's ICM lecture, §6.2, the theorem of §6.3, and (6.4.4), and supplies the missing direct comparison in D.20. Brown's motivic formula [Bro15, (1.3)] is not needed for that arithmetic proof. The source program's field `actual_Galois_and_GT_rational_rank` remains a literature-populated label; the arithmetic conclusion follows from the proof in D.20 rather than from that label.

Statement D.5 (Known). Cyclic-cover compatibility for every full profinite GT element. *Let $H_N = \ker(\widehat{F} \to \mathbb{Z}/N)$, where $x \mapsto 1$ and $y \mapsto 0$. The continuous homomorphism*

$$\delta_N : H_N \longrightarrow \widehat{F}, \quad x^N \mapsto x, \quad y \mapsto y, \quad x^a y x^{-a} \mapsto 1 \quad (0 < a < N)$$

is defined on a profinite free basis. For every $(\lambda, f) \in \widehat{\mathrm{GT}}$, with the convention $x \mapsto x^\lambda$, $y \mapsto f^{-1} y^\lambda f$, one has

$$\delta_N(f) = y^{c_N(f)} f, \qquad c_{MN}(f) = c_M(f) + c_N(f),$$

where $c_N(f)$ is the y coordinate of $\delta_N(f)$ in $\widehat{F}^{\mathrm{ab}}$. These are Enriquez's Theorem 8.1 and Remark 8.6 [Enr07].

Comparison with the covering map in LM7. For the negative Schreier generator $e_j = x^{-j} y x^j$, $0 < j < N$,

$$e_j = x^{-N}(x^{N-j} y x^{-(N-j)}) x^N.$$

Both sides are in H_N, and applying δ_N gives 1. The maps denoted ϕ_N in LM7 D.20 and δ_N above consequently agree on every member of a free basis, hence agree as continuous homomorphisms. We retain $\rho_N^{\mathrm{dist}} := -c_N$ as the explicitly defined negative distribution coordinate. The earlier sentence identifying it with the positive-root Kummer character in the displayed Galois convention requires the following correction. For $N = 2$, the exact comparison is

$$c_2(f_\sigma) = \kappa_2(\sigma), \qquad \rho_2^{\mathrm{dist}}(f_\sigma) = -\kappa_2(\sigma). \tag{D.5.1}$$

Here κ_2 is defined by $\sigma(2^{1/m}) = \zeta_m^{\kappa_2(\sigma)} 2^{1/m}$ using positive roots and $\zeta_m = e^{2\pi i/m}$. We now prove the coefficient and convention maps.

The continuous homomorphism $c_2 : H_2 \to \widehat{\mathbb{Z}}$ is the y coordinate after δ_2. On $\widehat{F}'$ it kills $\widehat{F}''$. In the free basis x^2, y, xyx^{-1} of H_2, every element of $\widehat{F}'$ has zero first coordinate and its other two abelian coordinates sum to zero. Conjugation by x interchanges those last two

coordinates, and conjugation by $y \in H_2$ fixes them. Moreover $[x,y] = (xyx^{-1})y^{-1}$ has c_2 value -1. Thus the actual map from the metabelian commutator module is

$$c_2(A * [x,y]) = -A(-1,1), \tag{D.5.2}$$

where $*$ is left conjugation. The identity follows first on the integer linear combinations of conjugates and then on their closure by continuity. Nakamura–Schneps use exactly $x \mapsto x^\chi$, $y \mapsto f_\sigma^{-1} y^\chi f_\sigma$; their equation (2.4) and Lemma 2.3 give $f_\sigma \equiv A_\sigma * [x,y]$ and $\kappa_2(\sigma) = -A_\sigma(-1,1)$ [NS00, §2, (2.1), (2.4), Lemma 2.3]. Combining these with (D.5.2) proves (D.5.1), including reduction modulo every integer.

The convention change mentioned just after their equation (2.1) is the actual continuous free-group automorphism

$$\iota(x) = x^{-1}, \qquad \iota(y) = y^{-1}, \qquad \delta_2 \circ \iota|_{H_2} = \iota \circ \delta_2. \tag{D.5.3}$$

To verify the last equality, compare its two sides on x^2, y, xyx^{-1}. The third generator is sent by ι to $x^{-2}(xyx^{-1})^{-1}x^2$, which δ_2 kills; the first two give x^{-1} and y^{-1} on both sides. Consequently $c_2(\iota f) = -c_2(f)$. Writing $f_\sigma^{\mathrm{Ihara}} = \iota(f_\sigma)$ for the specified change of variables gives $-c_2(f_\sigma^{\mathrm{Ihara}}) = \kappa_2(\sigma)$. The finite distribution tests retain their literal c_N maps; all arithmetic matrix comparisons below use the displayed Nakamura–Schneps convention.

For completeness, the automorphism g associated to (λ, f) preserves H_N, since it multiplies the quotient $\mathbb{Z}/N$ by the unit λ. The square with $g|_{H_N}$, g, and the two maps δ_N commutes. On x^N both composites give x^λ; on y the extra factor $y^{c_N(f)}$ cancels in conjugation. For $x^a y x^{-a}$, write $\lambda a = Nq + b$ with $0 < b < N$. Its image under g is conjugate, by an element of H_N, to $x^b y^\lambda x^{-b}$; applying δ_N gives 1. These checks prove the square on the full profinite free basis. □

This opened theorem strengthens the cyclic-cover statement used in LM7 D.20, where the displayed compatibility was used only on the Galois image. It provides additional necessary finite tests for Task E.3, performed below. It gives no equality between the full profinite groups $\widehat{\mathrm{GT}}$ and $\widehat{\mathrm{GT}}_0$.

Statement D.6 (Known). The exact finite test and its remaining quantifier. *For a finite-index normal subgroup $N_0 \triangleleft B_3$ contained in PB_3, the reduction maps of finite shadow sets satisfy*

$$\mathrm{im}\big(\widehat{\mathrm{GT}}_0 \to \mathrm{GT}(N_0)\big) = \bigcap_{K \subset N_0} R_{K,N_0}\big(\mathrm{GT}(K)\big),$$

where K ranges over all finite-index normal subgroups of B_3 contained in N_0. This is Dolgushev–Guynee, Theorem 5.2, Corollary 5.4 and Proposition 5.5 [DG24], with the full profinite identification supplied by Harbater–Schneps, Main Theorem (a) [HS00].

Verification of the inverse-limit step. A genuine element gives a shadow in every finite quotient. Conversely, the fibres of the reduction maps over a fixed shadow are finite sets. A finite collection of quotient conditions is dominated by the intersection of their kernels. Nonemptiness at every such kernel therefore gives the finite intersection property in the product of the finite shadow sets, and compactness produces a compatible family. The cited theorem identifies this family, including its generation condition, with a full profinite element. For a finite list of twelve shadows, exclusion of each one at some quotient implies exclusion of all twelve at the intersection of those twelve kernels. Thus the negative conclusion admits one finite witness. Passage through any specified collection of successful tests supplies only the conditions for that collection. □

We also opened Bleher–Chinburg–Lubotzky, Lemma 2.3 and Corollary 2.4 [BCL20], and reran LM7's explicit lifting counterexample. Those results lift automorphisms of a quotient of a finitely generated free profinite group. LM7's finite construction uses the quotient consisting of the free metabelian quotient together with an A_5 factor; the identity on the first factor and conjugation by an odd permutation on the second lift to an automorphism. The latter sends a five-cycle to its other A_5 conjugacy class. The lifted automorphism therefore fails the required conjugacy condition on x while acting trivially on the metabelian quotient. This verifies the claimed limitation of that particular lifting theorem. It does not rule out another lift satisfying the peripheral equations. The missing proof in D.1 and D.2 is exactly a solution of those equations and generation at every finite quotient, or a finite obstruction to the proposed extra coordinate.

4.3 A complete finite pentagon comparison in weight nine

Statement D.7 (Proved). Compatible finite three- and four-strand shadows. *Write* $P_n = PB_n$, $p = 101$, *and*

$$K_n = \gamma_{11} P_n P_n^p \lhd B_n \qquad (n = 2, 3, 4), \qquad K_F = \gamma_{11} F F^p,$$

where a power subgroup is generated by all the indicated powers. All these groups are discrete until their finite quotients are taken. We use $F = \langle x = A_{12}, y = A_{23} \rangle \leq P_3$ *and the same braid generators as in D.4. Let* $\mathcal{G}_9$ *be the finite gentle shadows at* K_3 *with* $m = 0 \pmod p$ *and* $f \in \gamma_9(F/K_F)$. *Let* $\mathcal{P}_9 \subset \mathcal{G}_9$ *be those satisfying the pentagon in* P_4/K_4. *The logarithm bijection makes both sets vector spaces over* $\mathbb{F}_{101}$ *and gives the following dimensions:*

	$\mathcal{G}_9$	$\mathcal{P}_9$
$\log f$ *in degrees* $9, 10$	37	2
degree-nine projection	4	1
degree-nine metabelian projection	2	1.

(D.7.1)

The last right-hand space is exactly $\mathbb{F}_{101} G_9[x, y]$. *The induced three-strand kernel associated to* K_4 *in the finite parenthesized-braid definition is exactly* K_3, *so* $\mathcal{P}_9$ *consists of actual finite pentagon shadows in that definition, not merely solutions in an unrelated Lie target.*

Proof. The quotient maps. The subgroups K_n are characteristic in P_n, hence normal in B_n. The groups P_n/K_n are finitely generated nilpotent groups of bounded finite exponent, so are finite. Every strand insertion or cabling homomorphism $P_3 \to P_4$ sends K_3 into K_4, since homomorphisms preserve the lower central series and power subgroups. For the inclusion of the first three strands, the strand-forgetting retraction $P_4 \to P_3$ gives $K_4 \cap P_3 = K_3$. Intersecting the five preimages used in the definition of the induced three-strand kernel therefore gives exactly K_3. The analogous two-strand assertion gives $K_2 = p\mathbb{Z}$ and $K_{\mathrm{ord}} = p$. The direct product $P_3 = F \times \langle C_3 \rangle$ gives $K_3 \cap F = K_F$. Thus reduction induces the specified maps of finite shadow sets, and the continuous maps from the profinite braid groups to these finite groups are their ordinary completion maps.

Faithful logarithmic coordinates. The class is ten, smaller than p. Lazard's Theorem II.4.6, with its homomorphism assertion, identifies the finite groups and their Lie rings by the mutually inverse truncated logarithm and Campbell–Hausdorff operations [Laz54]. For the free group factor this is the free nilpotent Lie algebra of class ten over $\mathbb{F}_p$. This assertion can also be read directly from Hall collection: the logarithm of a basic commutator is its Hall Lie word plus terms of larger degree; this triangular matrix has diagonal one, and all denominators through degree ten are units modulo p.

For four strands we keep the actual pure Artin action in these coordinates. Put $u_i = A_{i4}$, $a = A_{12}$, $b = A_{23}$. Forgetting strand four splits P_4 as $\langle u_1, u_2, u_3\rangle \rtimes P_3$. The center is generated by $C_4 = C_3 u_1 u_2 u_3$; eliminating C_3 in the quotient by C_4 gives

$$P_4/\langle C_4\rangle = F(u_1, u_2, u_3) \rtimes F(a, b). \tag{D.7.2}$$

These generators and their exact conjugation relations are [CFR12, (2.1)–(2.3)]. In particular the inverse actions of a, b on the free factor are

	u_1	u_2	u_3
$a^{-1}(-)a$	$u_1u_2u_1u_2^{-1}u_1^{-1}$	$u_1u_2u_1^{-1}$	u_3
$b^{-1}(-)b$	u_1	$u_2u_3u_2u_3^{-1}u_2^{-1}$	$u_2u_3u_2^{-1}$.

(D.7.3)

The actions are trivial on the free factor's abelianization. Their Magnus expansions raise degree by at least one; a weight-i commutator of them raises degree by at least i. Hence $[\gamma_i F(a,b), \gamma_j F(u_1,u_2,u_3)] \subset \gamma_{i+j} F(u_1,u_2,u_3)$. It follows by both inclusions that the lower central series in (D.7.2) is the product of the two free lower central series. Hall collection and reduction modulo p therefore give faithful coordinates consisting of the two free Lie factors. The omitted central factor has degree one. It intersects the derived subgroup trivially, also after this reduction, and every pentagon factor considered below lies in degree at least nine. Thus the computation in (D.7.2) detects the complete pentagon in P_4/K_4.

The complete equations are linear in the retained stratum. Write $X = \log x$, $Y = \log y$ and $\log f = h_9 + h_{10}$, with h_i homogeneous of degree i. There are $56 + 99 = 155$ Lyndon coordinates. The two simplified finite hexagons, with $z = y^{-1}x^{-1}$, are

$$f(x,y)f(y,x) = 1, \qquad f(z,x)f(y,z)f(x,y) = 1. \tag{D.7.4}$$

Their equivalence to the original braid hexagons for $f \in F'$ is Dolgushev–Guynee, Proposition 3.4 [DG24]. Here $m = 0$ and

$$\log z = -X - Y - \tfrac{1}{2}[X,Y] \pmod{\text{degree 3}}.$$

Every product of two h_i has degree at least eighteen. Consequently (D.7.4), including all terms of degree ten, is exactly a linear system. Substitution in the two-letter tensor algebra produces a 3072 by 155 matrix of rank 118. Its kernel has dimension 37; its degree-nine projection has dimension four. The program also solves for, retains, and substitutes a degree-ten correction for each of the four leading basis vectors.

Here are the corresponding exact four-strand substitutions. Put $U_i = \log u_i$, $A = \log a$, $B = \log b$, and

$$L = U_1 + U_2 + U_3 + \tfrac{1}{2}\sum_{i<j}[U_i, U_j] \pmod{\text{degree 3}}.$$

If the logarithm of an inverse Artin action in (D.7.3), applied to U_i, is $U_i + v_{2i} + v_{3i}$ through degree three, then its negative logarithm is the derivation

$$D(U_i) = -v_{2i} - v_{3i} + \tfrac{1}{2}D_2(v_{2i}), \qquad D_2(U_i) = v_{2i}. \tag{D.7.5}$$

This follows by expanding $-\log(1+T) = -T + T^2/2$ to the required degree. The leading parts are

$$\begin{aligned} &D_A(U_1) = [U_1, U_2], \quad D_A(U_2) = [U_2, U_1], \quad D_A(U_3) = 0, \\ &D_B(U_1) = 0, \qquad\quad D_B(U_2) = [U_2, U_3], \quad D_B(U_3) = [U_3, U_2]. \end{aligned}$$

Formula (D.7.5) supplies their degree-raising-two parts directly from the literal group words. The five pairs of arguments of the pentagon, in right-hand-side minus left-hand-side order, are

	first logarithm	second logarithm
$+$	A	$B + U_2 + \frac{1}{2}[U_2, U_3]$
$+$	$-A - L$	U_3
$-$	B	U_3
$-$	$-B - L$	$U_2 + U_3 + \frac{1}{2}[U_2, U_3]$
$-$	A	B.

(D.7.6)

For example $A_{13}A_{23} = a^{-1}C_3$ and $A_{12}A_{13} = C_3 b^{-1}$. The degree-two corrections involving A or B and L vanish because both leading derivations annihilate $U_1 + U_2 + U_3$. Terms of degree three in the argument logarithms, or derivation parts raising degree by three, cannot affect a word of weight nine through degree ten. Therefore (D.7.5)–(D.7.6) compute the complete group equation at this quotient, without identifying the logarithms of the pure braid generators with homogeneous infinitesimal generators.

The pentagon's free two-generator component cancels. Its free three-generator component has 3^9 and 3^{10} tensor coefficients. The degree-nine equations reduce the 37-dimensional hexagon space to dimension 34; all 59049 degree-ten equations then reduce it to dimension two. Projecting this kernel to degree nine gives rank one, and applying the metabelian quotient gives the asserted gamma line. All surviving basis vectors were substituted into every hexagon and pentagon coefficient.

Generation. The map on P_3/K_3 induced by a retained pair sends x to x, y to $f^{-1}yf$, and C_3 to C_3. It is the identity on the Frattini quotient of this finite p-group, and is therefore surjective. The two-strand map is also surjective. Definition 2.9 and Proposition 2.10 of Dolgushev–Le–Lorenz [DLL24] now give the generation condition on four strands for every retained pentagon pair. The same three-strand argument gives the gentle generation condition. This proves that the computed solutions are the claimed finite shadows.

The executable sources are `de_gentle_weight9.py` and `de_pentagon10_finite.py`; their JSON outputs retain the basis logarithms, and their compressed certificates retain the full coefficient matrices. The latter run completed in 501.307 seconds. Its assertions verify every surviving coefficient, the projection ranks, and equality with the gamma line. No search over a subset of the 155-coordinate stratum is used. □

Statement D.8 (Proved). An integer pentagon certificate and the weight-nine comparison. *Let k be a field of characteristic zero or characteristic at least five. In the free Lie algebra $L_k(X, Y)$ put*

$$V_9 = \{h \in L_k(X,Y)_9 : h(X,Y) + h(Y,X) = 0, \quad [X, h(X,Y)] = [X + Y, h(-X - Y, Y)]\}.$$

Then $\dim V_9 = 4$, its metabelian image has dimension two, and the infinitesimal pentagon map $P : V_9 \to (\mathfrak{t}_4)_9$ has rank three. Its one-dimensional kernel has metabelian image $k\,G_9(X,Y)[X,Y]$. The complete finite projection in D.7 is this kernel when $k = \mathbb{F}_{101}$. For every $\ell \geq 5$, applying the corresponding rational leading-term map to the full profinite pentagon equation and using D.20 gives precisely $\mathbb{Q}_\ell G_9(X,Y)[X,Y]$ as the weight-nine metabelian image of $\widehat{\mathrm{GT}}$ and of the Galois image.

Proof. The following exact integer certificate also explains the comparison with the finite calculation. Generate the 56 Lyndon words of length nine lexicographically, and use standard bracketing: split at the longest proper Lyndon suffix, then bracket the two recursively bracketed factors. Expand in the free associative algebra. The two equations defining V_9 give

an integer 1536 by 56 matrix C. Its 52 nonzero Smith factors are all one. A four-column integer kernel basis H retained in the JSON certificate has Smith factors $1, 1, 2, 24$. Hence it is a basis of the kernel over every field in the statement, not merely over $\mathbb{Q}$.

We evaluate the pentagon in the faithful Lie model

$$\mathfrak{t}_4/(\text{degree-one center}) = L(U_1, U_2, U_3) \rtimes L(A, B),$$

with the leading derivations displayed in D.7. It is the associated graded of (D.7.2); the integral braid relations are also recorded in [CFR12, (4.1)]. For the five arguments take the degree-one parts of (D.7.6). All eighteen infinitesimal braid relations are checked by the program, as is the cancellation of the free $L(A, B)$ component. The three tensor rows of $P(H)$ labelled 111111223, 111111232, 111121123 are

$$\begin{pmatrix} -6 & -6 & 6 & -5 \\ 254 & 510 & -446 & -1291 \\ -150 & -302 & 246 & 700 \end{pmatrix}. \tag{D.8.1}$$

The determinant of the first three columns is $27648 = 2^{10}3^3$. The vector

$$v = (-3299, 1326, -1613, 432)^{\mathsf{T}} \tag{D.8.2}$$

annihilates all 19683 tensor coefficients of $P(H)$, verified with integer arithmetic. Its coordinates have greatest common divisor one. Thus P has rank exactly three and kernel kv in each of the stated characteristics.

In the ordered monomial coordinates $X^iY^{7-i}[X, Y]$ for $0 \le i \le 7$, the metabelian image of H is

$$\begin{pmatrix} 0 & 0 & 0 & -2 \\ 0 & 0 & 0 & -8 \\ 2 & 2 & -2 & -17 \\ 6 & 6 & -6 & -23 \\ 6 & 6 & -6 & -23 \\ 2 & 2 & -2 & -17 \\ 0 & 0 & 0 & -8 \\ 0 & 0 & 0 & -2 \end{pmatrix}. \tag{D.8.3}$$

Its nonzero Smith factors are $1, 4$. Applied to (D.8.2), it gives

$$-96(9, 36, 84, 126, 126, 84, 36, 9)^{\mathsf{T}} = -96G_9.$$

This proves all the dimension and coordinate assertions. The programs `de_pentagon9.py` and `de_pentagon9_rational.py` construct the matrices from the Lie relations; the second performs the integer and rational calculations anew, including both Smith forms and every zero coefficient.

For the final comparison let a full profinite element have weight-nine leading action in the filtration of Part D. Its ℓ-adic multiplier is one. In the normalized gauge the action fixes x and its conjugator belongs to F'. A lower-degree inner action fixing x would have an inner generator centralizing X. In degree at least two this is zero in the free Lie algebra; the degree-one possibility is excluded by the derived-subgroup normalization. The conjugator therefore has no term below degree nine. Its leading word belongs to V_9 by the peripheral and symmetry equations.

The insertion homomorphisms of pure braid groups preserve their lower central series. Their associated graded maps send each pure generator to the corresponding sum of infinitesimal generators; this follows by abelianizing the explicit product formulas in the five

insertion maps. In degree nine the product of pentagon factors becomes their sum, since products of two degree-nine terms have degree eighteen. The leading word consequently lies in the exact kernel just computed over $\mathbb{Q}_\ell$. This is the upper inclusion, with its source and target specified. The actual Galois construction in D.20 supplies a nonzero element in the same filtered full profinite source, giving the reverse inclusion on rational spans. This derives R9 in weight nine from the certified pentagon map.

Finally, other multipliers act nontrivially in degree one, and lower nonzero leading words belong to lower filtration pieces. They do not define a weight-nine leading class under the map in Part D. This is why D.7 enumerates precisely the stratum needed for this comparison. The quotient detects an extra finite gentle direction, but the calculation does not construct a compatible family in the full profinite gentle group. The remaining existence question is D.2. □

4.4 The arithmetic lower bound with its exact coordinate comparison

Statement D.20 (Proved). The checked Galois and full GT gamma line in every odd weight. *For every prime $\ell \geq 5$ and every odd integer $m \geq 3$, the rational leading metabelian image of $G_{\mathbb{Q}}$, and therefore of $\widehat{\mathrm{GT}}$, is exactly*

$$\mathbb{Q}_\ell G_m[x,y], \qquad G_m = \frac{(X+Y)^m - X^m - Y^m}{XY}. \tag{D.20.1}$$

The leading image is zero in weight one and every positive even weight. In particular R9 has dimensions $0,0,1,0,1,0,1,0,1,0$ in weights one through ten, with the specified coordinate $p(X,Y)[x,y]$. This verifies the lower bound used in Tasks D.1–D.3; the full gentle and special image questions in D.1–D.2 remain separate questions about the displayed inclusion maps.

Manuscript comparison: §51.15, manuscript pages 725–733 in the supplied transcription, scans 156–170. The transcription does not establish the recto/verso labels for this range.

Proof. We use the primary formula in Ihara's ICM lecture, §6.2, the theorem of §6.3, and (6.4.4) [Iha91]. In its normalization $(f_\sigma)^{-1} = 1 + A_1\xi + A_2\eta$ and $\psi_\sigma = 1 + A_1\xi$. For $\sigma \in \ker \chi_\ell$, the formula is

$$\log \psi_\sigma^{\mathrm{ab}} = \sum_{\substack{r\geq 3 \\ r \text{ odd}}} \frac{\kappa_r^{(\ell)}(\sigma)}{(\ell^{r-1}-1)r!}\big((X+Y)^r - X^r - Y^r\big), \quad X = \log(1+\xi),\ Y = \log(1+\eta). \tag{D.20.2}$$

The cited §6.2 states nonvanishing of these odd cyclotomic cocycles for $\ell > 2$, with value zero at complex conjugation by their positive-real-root normalization. Formula (6.4.4) identifies this series with the actual metabelian action. These are the arithmetic inputs; we now supply the filtration and coordinate arguments.

First $\kappa_r^{(\ell)}$ stays nonzero on $\ker \chi_\ell$. Otherwise it factors through the abelian cyclotomic quotient. The cocycle identities for a cyclotomic element g and complex conjugation c would give

$$(1 - \chi_\ell(c)^r)\kappa_r^{(\ell)}(g) = (1 - \chi_\ell(g)^r)\kappa_r^{(\ell)}(c) = 0.$$

The left coefficient is two. This would make the whole cocycle zero, contradicting the cited nonvanishing.

Fix m. Apply the actual Galois action to the rational nilpotent completion of $F_\ell/\gamma_{m+3}F_\ell$. We retain this extra class because a degree-m multiplier acts on the commutator in degree $m+2$. The Hall coordinates of this finitely generated free nilpotent pro-ℓ group are integral and torsion-free. Its map to this rational completion is therefore injective, as is the map on

every lower-central graded lattice. Consequently triviality in a specified lower degree of the rational action detects triviality of the same integral action; this comparison does not create a new Galois element or weaken the required filtration. In the normalized gauge, x is fixed and the conjugator f belongs to the derived subgroup. An outer-trivial lower term is consequently actual-trivial: its inner generator must centralize x; its only possible degree-one component is a multiple of x, excluded by the normalization of f, and the higher-degree centralizer is zero by the free tensor-algebra comparison. Let G be the compact image of $\ker \chi_\ell$ in its automorphism group. It lies in the finite-dimensional unipotent group of automorphisms acting trivially on the abelianization. Its filtration I^d is the one defined at the start of Part D. Both the completed group-ring action on the metabelian commutator and its logarithm are homomorphisms on G: the action on its coefficient ring is trivial when $\chi_\ell = 1$. Consequently (D.20.2) defines a nonzero additive functional k_m on G, proportional to $\kappa_m^{(\ell)}$.

Here is the precise Lie extraction from this compact group. Put $\mathfrak{g} = \operatorname{span}_{\mathbb{Q}_\ell} \log G$ in the finite nilpotent automorphism Lie algebra. It is a Lie subalgebra: for $A = \log a$, $B = \log b$, the elements

$$\ell^{-2n} \log[a^{\ell^n}, b^{\ell^n}] \longrightarrow [A, B]$$

belong to this finite-dimensional, hence closed, span. Choose $a_1, \dots, a_d \in G$ whose logarithms form a basis. The analytic map

$$(t_1, \dots, t_d) \longmapsto \log(a_1^{t_1} \cdots a_d^{t_d})$$

is defined for $t_i \in \mathbb{Z}_\ell$, takes values in $\log G$, and has invertible differential at zero. The inverse-function theorem therefore puts an open neighbourhood of zero in $\mathfrak{g}$ inside $\log G$. In particular every sufficiently small scalar multiple of any vector of $\mathfrak{g}$ exponentiates to an element of the actual compact image. The functional k_m extends linearly to $\mathfrak{g}$, either by differentiating the displayed group-ring character or by its additive logarithm.

Choose $\gamma \in G_{\mathbb{Q}}$ with $u = \chi_\ell(\gamma) = 1 + \ell$, and write A for its conjugation operator on $\mathfrak{g}$. This preserves $\mathfrak{g}$, because the cyclotomic kernel is normal. On $\operatorname{gr}^d I$ it acts as multiplication by u^d. Indeed its action on the free abelianization is u, the graded Lie bracket multiplies degrees, and any higher part of the automorphism changes only higher degrees. Moreover

$$k_m(AZ) = u^m k_m(Z),$$

by the cyclotomic cocycle relation. Choose Z with $k_m(Z) \neq 0$ and set

$$Z_m = \prod_{d=1}^{m-1} (A - u^d) Z. \tag{D.20.3}$$

Successively applying these commuting factors sends I^d into I^{d+1}; thus $Z_m \in I^m \mathfrak{g}$. Its k_m value is $\prod_{d<m}(u^m - u^d) k_m(Z)$, which is nonzero. There is no assertion that a vector in the ambient target is realized merely because of its dimension: (D.20.3) belongs to the Lie algebra of the actual compact Galois image. The preceding open-neighbourhood argument gives $\exp(\ell^a Z_m) \in G \cap I^m$ for some integer $a \geq 0$. It has nonzero weight-m coefficient and lifts to an actual Galois element.

Finally let h be this element's leading degree-m conjugating Lie word and write $h = h_x x + h_y y$ in the tensor algebra. Its commutative image is zero. The definition of ψ therefore gives, in degree m,

$$\log \psi^{\mathrm{ab}} = -(h_x x)^{\mathrm{ab}} = (h_y y)^{\mathrm{ab}}.$$

The exact Fox comparison in D.4 sends $p[x, y]$ to XYp. Formula (D.20.2) thus gives a nonzero multiple of G_m in the requested coordinate. This coefficient also survives the outer quotient.

An inner derivation in this gauge would have to centralize x; in degree at least two of the metabelian free Lie algebra, $\operatorname{ad} x$ is multiplication by X on $\mathbb{Q}_\ell[X,Y][x,y]$ and has zero kernel.

This proves the nonzero Galois lower bound. The actual inclusions $G_{\mathbb{Q}} \to \widehat{\mathrm{GT}} \to \widehat{\mathrm{GT}}_0$ carry it to the full GT image. D.4 supplies the matching pentagon upper bound in every odd weight and its vanishing in even weights; weight one is zero by its explicit special equation. Together these statements prove (D.20.1) and the table. □

4.5 The full-source symmetry restriction in weight five

Statement D.21 (Proved). The weight-five full special image has dimension one or three. *Let $\ell \geq 5$, and let $\mathscr{I}_{5,\ell}$ be the rational weight-five metabelian leading image of the full profinite group $\mathrm{Out}^\sharp(\widehat{F}_2)$, with the source filtration and leading map specified at the beginning of Part D. Then*

$$\dim_{\mathbb{Q}_\ell} \mathscr{I}_{5,\ell} \in \{1,3\}. \tag{D.21.1}$$

In particular the intermediate value two offered in Task D.1 is excluded. This statement leaves the choice between one and three on the unresolved task ledger. More precisely, the polynomial coordinate map

$$(X+Y)q(X,Y)[x,y] \longmapsto q(X,Y) \quad : \quad (X+Y)\mathbb{Q}_\ell[X,Y]_2[x,y] \xrightarrow{\sim} \mathbb{Q}_\ell[X,Y]_2 \tag{D.21.2}$$

identifies the target with the direct sum of the gamma line $\mathbb{Q}_\ell(X^2+XY+Y^2)$ and an absolutely irreducible two-dimensional $\mathfrak{S}_3$-module. The image in (D.21.1) contains that line and is stable under this actual geometric symmetry action.

Manuscript comparison: §51.15, manuscript pages 725–733 in the supplied transcription, scans 156–170; the full special-image question of LM7 D.13.

Proof. We give the comparison between conjugation in the full source and the polynomial action, including the inner corrections. The geometric automorphisms permuting the three punctures give an actual subgroup $\mathfrak{S}_3$ of $\mathrm{Out}^\sharp(\widehat{F}_2)$. Conjugation by this subgroup preserves every term of the source filtration: the lower central and second-derived subgroups used to define the target are characteristic. The leading derivation map is equivariant because, to its least nonzero degree, conjugation of an automorphism acts by conjugation of its derivation. Only the degree-one part of the conjugating geometric automorphism contributes in that degree. Thus $\mathscr{I}_{5,\ell}$ is stable under the indicated $\mathfrak{S}_3$ action before any representation-theoretic argument is applied.

Put $c = [x,y]$, $X = \operatorname{ad} x$ and $Y = \operatorname{ad} y$. Work in the free metabelian Lie algebra, where its commutator module is $\mathbb{Q}_\ell[X,Y]c$. By D.4, the exact weight-five target is $p = (X+Y)q$, with q homogeneous of degree two. The normalized representative of its outer derivation is

$$D_p(x) = 0, \qquad D_p(y) = [pc, y] = -Ypc. \tag{D.21.3}$$

An inner derivation $\operatorname{ad}(hc)$ has coordinates $(-Xhc, -Yhc)$. The polynomial p in (D.21.3) is unique: an inner derivation with zero x coordinate has $Xh = 0$, which forces $h = 0$ in the polynomial module.

Take the transposition $s : x \leftrightarrow y$ and the three-cycle $r : x \mapsto y$, $y \mapsto -x-y$ on the associated graded. The first sends c to $-c$ and the second fixes c. Their conjugated derivations have coordinate pairs

$$\begin{aligned} sD_p s^{-1} &= (Xp(Y,X)c,\ 0), \\ rD_p r^{-1} &= (-(X+Y)p(Y,-X-Y)c,\ 0). \end{aligned} \tag{D.21.4}$$

Subtract respectively the inner derivations with

$$h_s = -p(Y, X), \qquad h_r = -(X + Y)q(Y, -X - Y).$$

The second correction is a polynomial because $p(Y, -X - Y) = -Xq(Y, -X - Y)$. Formula (D.21.3) is restored, and the exact induced action under (D.21.2) is

$$s(q) = q(Y, X), \qquad r(q) = q(Y, -X - Y). \tag{D.21.5}$$

The inner ambiguity has therefore been accounted for by the displayed actual maps.

In the ordered basis X^2, XY, Y^2, the action matrices are

$$S = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \qquad R = \begin{pmatrix} 0 & 0 & 1 \\ 0 & -1 & 2 \\ 1 & -1 & 1 \end{pmatrix}. \tag{D.21.6}$$

They satisfy $S^2 = R^3 = 1$ and $SRS = R^{-1}$. Their common invariant line is generated by $q_0 = X^2 + XY + Y^2$. Averaging the six matrices gives the rank-one idempotent

$$P_0 = \frac{1 + R + R^2 + S + SR + SR^2}{6}. \tag{D.21.7}$$

Its kernel U is a stable two-dimensional complement. It contains no invariant vector. It also contains no sign subrepresentation: $s(q) = -q$ forces $q = a(X^2 - Y^2)$, and $r(q) = q$ then forces $a = 0$. Every one-dimensional representation of $\mathfrak{S}_3$ is trivial or sign, since its abelianization has order two. This proves that U is irreducible, and the same argument over any field extension of $\mathbb{Q}_\ell$ proves absolute irreducibility. Equivalently, the sign projector is the zero matrix on the entire three-dimensional space.

By D.20, the image $\mathscr{I}_{5,\ell}$ contains $G_5 c$. The exact polynomial identity

$$G_5 = 5(X + Y)(X^2 + XY + Y^2) \tag{D.21.8}$$

identifies this nonzero arithmetic line with $\mathbb{Q}_\ell q_0$. The averaging operator P_0 preserves the source image, as it is a rational linear combination of the six symmetry operators already shown to preserve it. Hence the image splits as that line plus its intersection with U. Irreducibility makes this intersection either zero or all of U. This proves (D.21.1) using the full-source comparison; no lifting from the pro-ℓ target is used.

The exact program `d_weight5_symmetry.py` computes both conjugation and inner-normalization maps for a general quadratic, checks all six group matrices, both projectors, the two-dimensional complement, and (D.21.8). Its rational matrices and complete output are retained. The finite certificate verifies these explicit algebraic identities; source stability and the arithmetic inclusion are the preceding proof and D.20. □

4.6 What a finite leading congruence detects

Statement D.22 (Proved). Exact integral rank recovery and a finite-congruence limitation. *Let M be a finite free $\mathbb{Z}_\ell$-module and let $L \subset M$ be the image of a continuous homomorphism from a compact group to the additive group of M. Write L_n for its image under the specified reduction $M \twoheadrightarrow M/\ell^n M$. There are integers $e_1, \dots, e_r \geq 0$ such that*

$$\log_\ell |L_n| = \sum_{i=1}^{r} \max(n - e_i, 0), \qquad \dim_{\mathbb{Q}_\ell}(L \otimes \mathbb{Q}_\ell) = \lim_{n \to \infty} \frac{\log_\ell |L_n|}{n} = r. \tag{D.22.1}$$

The successive difference of the displayed logarithms equals r for every $n > \max_i e_i$.

For each $a \geq 1$ and each $\ell \geq 5$, the actual $\mathfrak{S}_3$-module $M = \mathbb{Z}_\ell[X,Y]_2$ of D.21 contains two $\mathfrak{S}_3$-stable closed submodules with the same image in $M/\ell^a M$, both containing $\mathbb{Z}_\ell q_0$, and with rational dimensions one and three, respectively. The corresponding two-dimensional target of D.2 contains submodules with the same finite reduction and rational dimensions one and two. These are explicit submodules; we assert no realization of them as images of the profinite sources in D.1 or D.2. They refute the inference of a rational upper bound from one vanishing extra congruence without further control of the integral image.

Proof. Compactness makes L closed. Integer multiplication preserves this additive subgroup, and passage to limits in M makes it a $\mathbb{Z}_\ell$-submodule. It is finitely generated because $\mathbb{Z}_\ell$ is noetherian. The Smith basis theorem gives a basis $b_1, \dots, b_d$ of M with

$$L = \bigoplus_{i=1}^{r} \ell^{e_i} \mathbb{Z}_\ell b_i.$$

The image of the ith summand under the specified reduction has order $\ell^{\max(n-e_i,0)}$. Multiplying these independent orders proves (D.22.1), including the asserted successive difference. The leading map on the closed source filtration in Part D is an additive continuous map; its compact image therefore has exactly this form in an integral leading lattice. This is the map to which the rank formula applies.

For the weight-five example, use the integral averaging idempotent P_0 of D.21; its denominator six is a unit for $\ell \geq 5$. Set $U_0 = \ker(P_0 : M \to M)$. In the ordered coefficient basis X^2, XY, Y^2, explicit bases are

$$q_0 = (1,1,1), \qquad u_1 = (1,2,0), \qquad u_2 = (-1,0,1). \tag{D.22.2}$$

The three-column determinant is 3, a unit in $\mathbb{Z}_\ell$, and $P_0 u_1 = P_0 u_2 = 0$. Thus $M = \mathbb{Z}_\ell q_0 \oplus U_0$ and U_0 has basis u_1, u_2. Both summands are stable under the exact matrices of (D.21.6). Define

$$L^{(1)} = \mathbb{Z}_\ell q_0, \qquad L^{(3)} = \mathbb{Z}_\ell q_0 \oplus \ell^a U_0. \tag{D.22.3}$$

Their inclusion maps into M, followed by reduction modulo $\ell^n M$, have the same image for every $n \leq a$. Their rational ranks are one and three. Their finite-image orders for all n are, explicitly,

$$|L^{(1)}_n| = \ell^n, \qquad |L^{(3)}_n| = \ell^{\,n+2\max(n-a,0)}. \tag{D.22.4}$$

This constructs the requested comparison inside the actual symmetry module and retains both its invariant line and its symmetry action.

For D.2 use the explicit independent weight-nine polynomials G_9 and $P_9 = X^2Y^2(X+Y)^3$ of D.4 and the lattice $M_9 = \mathbb{Z}_\ell G_9 \oplus \mathbb{Z}_\ell P_9$ in their rational span. The inclusions

$$\mathbb{Z}_\ell G_9 \subset M_9, \qquad \mathbb{Z}_\ell G_9 \oplus \ell^a \mathbb{Z}_\ell P_9 \subset M_9 \tag{D.22.5}$$

give the corresponding example under $M_9 \twoheadrightarrow M_9/\ell^a M_9$. The target is already the invariant part, so the indicated $\mathfrak{S}_3$ action is trivial on both submodules.

The program `d_leading_lattices.py` checks the integer matrices, the integral Smith valuations, and every element of the finite images for $(\ell, a) = (5,2), (7,1)$ at the recorded precisions. Its complete output is retained. The initially incorrect determinant assertion in that program was corrected from six to the computed value three; the failed run and its correction are retained separately.

For the suggested finite-quotient route in the task, the actual profinite leading image maps to each finite shadow image by reduction, as specified in D.6. An extra coordinate vanishing in

one such finite image proves divisibility of that coordinate in the corresponding integral lattice. Formula (D.22.3) proves that this divisibility alone does not force its rational vanishing. A finite argument that also controls the Smith valuations, or an integral homogeneous identity valid at every precision, supplies additional information; neither is silently inserted here. The integer homogeneous pentagon identities of D.7–D.8 and the actual arithmetic lower bound of D.20 retain their proved scopes. □

4.7 Exact peripheral lifting across abelian chief factors

All conjugations in this subsection act on the left, and $[a,b] = aba^{-1}b^{-1}$. The three peripheral elements of the actual free profinite group are

$$\widehat{F} = \langle x_1, x_2 \rangle, \qquad x_3 = (x_1 x_2)^{-1}, \qquad S_i = \overline{\langle x_i \rangle} \simeq \widehat{\mathbb{Z}}. \tag{D.23.1}$$

In particular the group in the following calculation is the full profinite group, with these specified three embeddings of $\widehat{\mathbb{Z}}$. For x_3 this follows from the explicit basis replacement $(x_1, x_2) \mapsto (x_1, (x_1 x_2)^{-1})$, whose inverse recovers $x_2 = x_1^{-1} x_3^{-1}$.

Statement D.23 (Proved). The complete abelian peripheral obstruction and chief-step count. *Let*

$$1 \longrightarrow A \longrightarrow E \xrightarrow{\pi} G \longrightarrow 1 \tag{D.23.2}$$

be an extension of finite groups with A elementary abelian of characteristic p. Choose generators X_1, X_2 of E, put $X_3 = (X_1 X_2)^{-1}$ and $g_i = \pi(X_i)$, and denote the resulting epimorphisms by $q_E : \widehat{F} \twoheadrightarrow E$ and $q_G = \pi q_E$. Let $\alpha \in \mathrm{Aut}(G)$ and $\lambda \in \widehat{\mathbb{Z}}^\times$ satisfy $\alpha(g_i) \sim_G g_i^\lambda$ for all three indices.

Write $A_G = A/[E,A]$, use additive notation in A, and set $C_i = C_G(g_i)$. There are well-defined homomorphisms

$$\kappa_i : C_i \longrightarrow A_G, \qquad c \longmapsto [\widetilde{c}, X_i] \bmod [E,A], \qquad H = \sum_{i=1}^{3} \operatorname{im} \kappa_i. \tag{D.23.3}$$

Choose $h_i \in G$ with $\alpha(g_i) = h_i g_i^\lambda h_i^{-1}$, choose lifts $H_i \in E$, and put

$$U_i = H_i X_i^\lambda H_i^{-1}, \qquad r = U_1 U_2 U_3 \in A. \tag{D.23.4}$$

The class $o(\alpha, \lambda) = [r] \in A_G/H$ is independent of these choices. It vanishes exactly when there is a continuous homomorphism $f : \widehat{F} \to E$ such that

$$\pi f = \alpha q_G, \qquad f(x_i) \sim_E X_i^\lambda \quad (1 \le i \le 3). \tag{D.23.5}$$

The number of these literal homomorphisms is zero for nonzero $o(\alpha,\lambda)$; for zero obstruction it is

$$C(E, G; X_1, X_2) = \frac{|A|^2 |A_G| \prod_{i=1}^{3} |C_G(g_i)|}{|H| \prod_{i=1}^{3} |C_E(X_i)|}. \tag{D.23.6}$$

When A is an irreducible $\mathbf{F}_p[G]$-module, every nonempty set in (D.23.5) *contains an epimorphism. More precisely, let $b(E, G; X_1, X_2)$ be the number of complements C to A in E whose unique section $s_C : G \to C$ satisfies $s_C(g_i) \sim_E X_i$ for every i. For zero obstruction the number of epimorphisms in* (D.23.5) *is exactly $C - b > 0$. Thus every noncentral elementary-abelian chief step admits a generating peripheral lift. The possible abelian chief obstruction is confined to a central kernel of order p.*

The fixed-h_i obstruction $[r] \in A_G$ is the relative extension class under the explicit identification

$$H^2(\widehat{F}, \{S_1, S_2, S_3\}; A_{\alpha q_G}) \simeq A_G. \tag{D.23.7}$$

Moreover $[r]$ *comes from* $H_2(G, \mathbf{F}_p)$ *by the universal central* p*-cover comparison in* (D.23.12) *below. In particular* $p \nmid |G|$ *forces the obstruction to vanish.*

Proof. Let σ_i be conjugation by U_i on A. Its value depends only on $\alpha(g_i)$, and $\sigma_1\sigma_2\sigma_3 = 1$. Replacing H_i by a_iH_i, for $a_i \in A$, changes the product in (D.23.4) to

$$r + L(a_1, a_2, a_3), \qquad L(a_1, a_2, a_3) = (1 - \sigma_1)a_1 + \sigma_1(1 - \sigma_2)a_2 + \sigma_1\sigma_2(1 - \sigma_3)a_3. \quad \text{(D.23.8)}$$

This is a multiplication identity in the extension, since all three correction factors lie in the abelian normal subgroup A. The sum of the first two image spaces is

$$(1 - \sigma_1)A + (1 - \sigma_1\sigma_2)A = [E, A].$$

For the equality, use $1 - ab = (1 - a)b + (1 - b)$ and the fact that $\sigma_1, \sigma_1\sigma_2$ generate the action of G; inverses give the same image spaces. The third image space is already in this sum. Consequently $\operatorname{im} L = [E, A]$, and $|\ker L| = |A|^2|A_G|$. This proves the assertion with the three base conjugators fixed.

To vary those conjugators, work in $\overline{E} = E/[E, A]$. Its kernel A_G is central. If $c \in C_i$, the commutator $[\widetilde{c}, X_i]$ lies in A; changing the lift of c changes its image in A_G by zero. The central commutator identity $[ab, X_i] = [a, X_i][b, X_i]$ proves that κ_i is a homomorphism. All conjugators in G from g_i^λ to $\alpha(g_i)$ have the form h_ic_i, with $c_i \in C_i$, since raising to the unit λ preserves the generated cyclic subgroup. Their effect on the product class is

$$[r] \longmapsto [r] + \lambda \sum_i \kappa_i(c_i). \quad \text{(D.23.9)}$$

Here conjugation by a lift of h_i acts trivially on A_G and $[\widetilde{c}, X_i^\lambda] = [\widetilde{c}, X_i]^\lambda$ in $\overline{E}$. Multiplication by λ is an automorphism of A_G, and preserves H. This proves choice independence and the criterion in A_G/H.

For the count, a zero obstruction gives exactly $\prod_i |C_i|/|H|$ triples of base conjugators satisfying the coinvariant equation. Each such triple has exactly $|\ker L| = |A|^2|A_G|$ triples of lifts in E whose conjugated peripheral product is 1. Each resulting triple of peripheral elements is counted $\prod_i |C_E(X_i^\lambda)|$ times. These centralizers equal $C_E(X_i)$ because λ is a unit. A triple with product 1 gives exactly one continuous homomorphism out of $\widehat{F}$, by its free profinite universal property. This proves (D.23.6) without quotienting by a centralizer that was not counted.

Now assume A irreducible. If $L = f(\widehat{F})$ for a solution, then $L \to G$ is onto and $L \cap A$ is G-invariant: conjugation by any element of G can be computed using a lift in L. Hence $L \cap A$ is either A or 0. The first case gives $L = E$; the second makes L a complement. For a given complement C, there is exactly one homomorphism over αq_G with image in C, namely $s_C\alpha q_G$. It satisfies the peripheral conditions precisely when $s_C(g_i) \sim_E X_i$ for all i. Indeed

$$s_C(\alpha(g_i)) = s_C(h_i)s_C(g_i)^\lambda s_C(h_i)^{-1},$$

and taking the power λ^{-1} in the finite group E shows that $s_C(g_i)^\lambda \sim_E X_i^\lambda$ is equivalent to $s_C(g_i) \sim_E X_i$. Thus the nongenerating solutions are in a literal bijection with the same b admissible complements, independently of α and λ. The identity case $(\alpha, \lambda) = (1, 1)$ contains the given epimorphism q_E, so its count C is strictly greater than b. Every other nonempty case has this same count C, proving the claimed generating-lift theorem. Finally an irreducible module with $A_G \neq 0$ is the one-dimensional trivial module: the quotient map to a nonzero trivial quotient has zero kernel by irreducibility. This proves the noncentral chief assertion.

For clarity, the count in (D.23.6) is for homomorphisms, with the specified base map retained. If one further takes A-conjugacy classes of the generating solutions, every orbit has

size $|A|/|A^G|$. If one permits all conjugations in E that retain the same base map αq_G, the acting group is $\pi^{-1}Z(G)$ and every generating stabilizer is $Z(E)$. The corresponding number of classes is therefore $(C-b)|Z(E)|/(|A||Z(G)|)$. These formulas specify the extra quotient maps rather than silently identifying literal lifts with their classes.

Here is the relative-cohomology comparison. The free profinite universal property makes the cocycle group $Z^1(\widehat{F},A)$ equal to A^2: arbitrary values on x_1,x_2 define a continuous homomorphism to the finite semidirect product associated with the action. Each $S_i \simeq \widehat{\mathbb{Z}}$ has $H^1(S_i,A) = A/(\sigma_i - 1)A$. Projectivity of $\widehat{F}$ gives $H^2(\widehat{F},A) = 0$, so the relative exact sequence identifies the group in (D.23.7) with the cokernel of the restriction map on H^1. On cocycle values the latter is

$$(u,v) \longmapsto (u,v,-\sigma_3(u+\sigma_1 v)). \tag{D.23.10}$$

The map from the three local values to this cokernel, followed by (D.23.7), is

$$(b_1,b_2,b_3) \longmapsto b_1 + \sigma_1 b_2 + \sigma_1\sigma_2 b_3 \pmod{[E,A]}. \tag{D.23.11}$$

It is onto. Eliminating b_1, b_2 using (D.23.10) leaves exactly the relations $(\sigma_1 - 1)A + \sigma_1(\sigma_2 - 1)A = [E,A]$, which proves that its kernel is the asserted restriction image. Pull back (D.23.2) along αq_G and prescribe the local sections that send x_i to (U_i, x_i). Choose the global splitting taking x_1, x_2 to $(U_1,x_1),(U_2,x_2)$. The difference of the third prescribed section from this splitting has cocycle value $\sigma_3 r$, so (D.23.11) sends the relative class to r. This is the actual finite-kernel relative extension of the full profinite pair. It agrees with the classification and lifting comparison of Wilkes, Theorem 5.5 and Proposition 6.3, with lifting problems defined in Definition 6.1 [Wil25]. In particular no passage from a pro-p pair to the full profinite pair is being assumed.

For the final assertion put $K = \ker q_G$ and $U = \widehat{F}/\overline{K^p[\widehat{F},K]}$. Finite-coset Schreier rewriting gives at most $1 + |G|$ topological generators for K. Therefore its elementary abelian quotient A_U is finite, and it is central in U. The homology five-term sequence and $H_2(\widehat{F},\mathbf{F}_p) = 0$ give the exact comparison

$$0 \longrightarrow H_2(G,\mathbf{F}_p) \longrightarrow A_U = K/\overline{K^p[\widehat{F},K]} \longrightarrow H_1(\widehat{F},\mathbf{F}_p) \longrightarrow H_1(G,\mathbf{F}_p) \longrightarrow 0. \tag{D.23.12}$$

The original map q_E induces an actual surjection $U \twoheadrightarrow E/[E,A]$, and its map on kernels is $A_U \twoheadrightarrow A_G$. Choose lifts $\widetilde{H}_i$ of the H_i in $\widehat{F}$. The word

$$\prod_{i=1}^{3} \widetilde{H}_i x_i^{\lambda} \widetilde{H}_i^{-1} \tag{D.23.13}$$

lies in K and has exactly zero image in $\widehat{F}^{\mathrm{ab}} = \widehat{\mathbb{Z}}^2$, since $\lambda(x_1+x_2+x_3) = 0$ there. Its image in A_G is $[r]$. Thus its class in A_U belongs to the $H_2(G,\mathbf{F}_p)$ term of (D.23.12). If $p \nmid |G|$, averaging contracts the positive-degree finite-group homology complex over $\mathbf{F}_p$, so that term is zero. For general G the Schur term and the quotient by the three maps (D.23.3) both remain in the calculation. □

The independent finite program `a_peripheral_count.py` enumerates actual peripheral triples and their generated subgroups, then compares them with (D.23.6) and the complement bijection. It checks sixteen cases in six extensions. In the $V_4 \triangleleft A_4$ example there are sixteen weak solutions, twelve generating solutions, and four admissible complements. In the central $C_2 \triangleleft D_8$ example the centralizer-image subgroup H has order two; including it gives the four actual solutions. These examples test both terms that an unrestricted generator-lifting argument would omit. The script, exact output, and JSON tables are retained in `code/`.

4.8 A nonzero spin obstruction and its exact pro-ℓ fibre

Statement D.24 (Proved). A peripheral-preserving finite automorphism with no peripheral lift. *Let* $S = \mathrm{PSL}_2(\mathbf{F}_{11})$ *and*

$$E = \mathrm{SL}_2(\mathbf{F}_{11}) \times S, \qquad G = S \times S, \qquad \pi = (\textit{central quotient}, 1). \tag{D.24.1}$$

There is an explicit epimorphism $q_E : \widehat{F} \twoheadrightarrow E$ *for which the factor-swap automorphism* $\alpha : G \to G$, $(a, b) \mapsto (b, a)$, *preserves each of the three labelled peripheral conjugacy classes with multiplier one, but no homomorphism* (D.23.5) *exists. Its obstruction is the nonidentity element of* $\ker \pi = \{(I, 1), (-I, 1)\}$.

The maximal pro-ℓ quotients of both E and G are trivial for every prime ℓ. More precisely, for any two-generated finite ℓ-group L with a specified generating triple $l_1 l_2 l_3 = 1$ and any multiplier-one peripheral automorphism $\beta \in \mathrm{Aut}(L)$, the corresponding product quotient has maximal pro-ℓ quotient exactly L. The automorphism $\alpha \times \beta$ of $G \times L$ has no peripheral lift to $E \times L$, whereas $1_G \times \beta$ has an explicit generating peripheral lift. Thus the same prescribed finite pro-ℓ component has both an obstructed and an admitted choice of perfect component in this example.

Proof. All matrix entries in the following display are in $\mathbf{F}_{11}$. Take the two triples of canonical odd-order lifts

i	1	2	3
A_i	$\begin{pmatrix} 0 & 10 \\ 1 & 10 \end{pmatrix}$	$\begin{pmatrix} 0 & 8 \\ 4 & 2 \end{pmatrix}$	$\begin{pmatrix} 6 & 2 \\ 4 & 7 \end{pmatrix}$
B_i	$\begin{pmatrix} 0 & 10 \\ 1 & 10 \end{pmatrix}$	$\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$	$\begin{pmatrix} 1 & 10 \\ 0 & 1 \end{pmatrix}$.

(D.24.2)

Write a_i, b_i for their respective images in S. Direct multiplication and powering give

$$A_1A_2A_3 = I, \qquad B_1B_2B_3 = -I, \qquad |A_i| = |B_i| = (3, 11, 11)_i. \tag{D.24.3}$$

The explicit matrices

$$H_1 = \begin{pmatrix} 0 & 1 \\ 10 & 1 \end{pmatrix}, \quad H_2 = \begin{pmatrix} 2 & 6 \\ 0 & 6 \end{pmatrix}, \quad H_3 = \begin{pmatrix} 0 & 5 \\ 2 & 3 \end{pmatrix} \tag{D.24.4}$$

have determinant one and satisfy $H_i A_i H_i^{-1} = B_i$. In particular a_i and b_i are conjugate in S for each i. Set

$$X_i = (A_i, b_i) \in E, \qquad g_i = (a_i, b_i) \in G. \tag{D.24.5}$$

Both triples have product one. They generate their displayed groups, as certified by the complete matrix enumeration below. The elements $(\overline{H}_i, \overline{H}_i^{-1}) \in G$ conjugate g_i to $\alpha(g_i)$, verifying the exact base peripheral condition.

An element of E above $\alpha(g_i) = (b_i, a_i)$ has first coordinate either B_i or $-B_i$. Exactly one of these has odd order, namely B_i: if B_i has odd order n, then $(-B_i)^n = -I$ and $-B_i$ has order $2n$. Every conjugate of X_i has the same odd order as X_i, so its only possible value above $\alpha(g_i)$ is

$$U_i = (B_i, a_i). \tag{D.24.6}$$

It is indeed conjugate to X_i in E, using $(H_i, \overline{H}_i^{-1})$. But its product is $U_1U_2U_3 = (-I, 1) \neq 1$. Thus no choice of conjugators, and no nongenerating homomorphism, solves this peripheral lifting problem. Equivalently, each base-centralizer homomorphism (D.23.3) is zero. To see this directly, in this central extension the commutator $[\widetilde{c}, X_i]$ has order dividing two and also

dividing the odd order of X_i, so it is one. The computed nonzero relative class consequently survives the full base-centralizer quotient.

Here is the finite certificate for all generation assertions.

The program `a_peripheral_spin.py` constructs all determinant-one 2×2 matrices over $\mathbf{F}_{11}$ and identifies a matrix with its negative for S. It verifies their orders 1320 and 660, the three canonical odd-order lifts, all products in (D.24.3), and all three conjugations in (D.24.4). Starting from the identity, it repeatedly right-multiplies the generated set by the two displayed generators until the set stops growing. This computes the generated subgroup exactly: the set contains every positive word, is closed under multiplication by the generators, and positive powers include their inverses in these finite groups. For (g_1, g_2) it has $435600 = 660^2$ elements, and for (X_1, X_2) it has $871200 = 1320 \cdot 660$ elements. These are the complete group orders, proving surjectivity. The matrices, canonical lifts, products, conjugators, and both cardinalities are saved in `a_peripheral_spin_p11.json`; the executed output is `a_peripheral_spin_p11.out`. The separately retained $p = 5, 7$ searches found no matching pair in their stated exhaustive odd-order search domains; the proof here uses the displayed $p = 11$ certificate. The program explicitly restricts its finite order bound to these three tested primes.

For the pro-ℓ statement, first prove perfectness directly. In $\mathrm{SL}_2(\mathbf{F}_{11})$ put

$$D = \mathrm{diag}(2, 6), \quad u(t) = \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix}, \quad v(t) = \begin{pmatrix} 1 & 0 \\ t & 1 \end{pmatrix}.$$

One has $[D, u(t)] = u(3t)$ and $[D, v(t)] = v(2t)$. Both scalar factors are invertible in $\mathbf{F}_{11}$. Elementary row reduction expresses every determinant-one matrix as a product of the $u(t), v(t)$, so the whole group is its commutator subgroup. Its quotient S is perfect, and hence so are E and G. A finite perfect group has trivial image in every finite ℓ-group: such an image would be a perfect finite ℓ-group, whereas induction on the central series shows that a nontrivial finite ℓ-group has nontrivial abelianization. Passing to the finite quotients of a pro-ℓ group proves the assertion about maximal pro-ℓ quotients.

Finally use the source generators (X_i, l_i) and (g_i, l_i) in $E \times L$ and $G \times L$. They generate the full products. Indeed their generated subgroups project onto both factors; a subdirect subgroup $B \leq P \times L$, with coordinate kernels $P_0 = \{p : (p, 1) \in B\}$ and $L_0 = \{l : (1, l) \in B\}$, gives the isomorphism $P/P_0 \to L/L_0$, $pP_0 \mapsto lL_0$ for $(p, l) \in B$. For perfect P and an ℓ-group L, this common quotient is perfect and solvable, hence trivial, which makes the subgroup the full product. This also proves that the actual coordinate projection is the maximal pro-ℓ quotient map of either product. The shadow $\alpha \times \beta$ satisfies the peripheral conditions by the separate conjugators in the two factors. Any lift would project to the already excluded lift of α in (D.24.6). In contrast, the literal triple

$$(X_i, \beta(l_i)) \quad (1 \leq i \leq 3) \tag{D.24.7}$$

has product one, has the prescribed three conjugacy classes, and generates $E \times L$ by the same subdirect-product argument. It is the promised lift of $1_G \times \beta$. □

Statements D.23–D.24 supply an exact chief-step obstruction and a nonzero full-profinite peripheral example. The displayed maximal pro-ℓ quotient maps and the two product lifts specify their effect on the pro-ℓ comparison. They do not choose between the dimensions in D.1 or D.2: the choice of compatible finite components, including coupled mixed quotients, remains part of those two problems.

4.9 A prime-to-five central obstruction that survives every five-adic unit power

Statement D.25 (Proved). An order-five peripheral cycle obstructed by a central kernel of order two. *There are finite groups*

$$E \xrightarrow{\pi} G = A_8^5, \qquad |\ker \pi| = 2, \tag{D.25.1}$$

an epimorphism $q_E : \widehat{F} \twoheadrightarrow E$, and a multiplier-one, labelled peripheral automorphism $\alpha \in \mathrm{Aut}(G)$ of order five, such that none of $\alpha, \alpha^2, \alpha^3, \alpha^4$ has a weak peripheral lift through π. The kernel is central and its obstruction group in D.23 is exactly $\mathbf{F}_2$. Consequently taking a 5-adic unit power cannot remove this obstruction. In particular the order of the obstruction group does not, by itself, bound the exponent of a power that removes a pointed peripheral lifting obstruction.

Both groups in (D.25.1) *are perfect, and their maximal pro-five quotients are trivial. Thus this counterexample corrects a proposed lifting argument; it imposes no restriction on a nonzero pro-five leading coordinate in D.1.*

Proof. We construct the double cover used in the example rather than invoke a group database. Let $\mathcal{C}$ be the real Clifford algebra on $e_0, \dots, e_7$, with $e_i^2 = 1$ and $e_i e_j = -e_j e_i$ for $i \neq j$. The unit vector

$$u_{ij} = \frac{e_i - e_j}{\sqrt{2}} \tag{D.25.2}$$

acts by the reflection $v \mapsto -u_{ij} v u_{ij}^{-1}$ that interchanges e_i, e_j. Products of an even number of these vectors therefore act by conjugation as even permutations of the eight vectors. Let $\widetilde{S}$ be their group. This gives an exact sequence

$$1 \longrightarrow \{1, -1\} \longrightarrow \widetilde{S} \longrightarrow S = A_8 \longrightarrow 1. \tag{D.25.3}$$

Indeed, an even Clifford element that commutes with every e_i is a scalar, as is seen on the monomial basis $e_{i_1} \cdots e_{i_r}$; the reversion norm of a product of unit vectors is one, so a scalar in the kernel is 1 or -1. The element $u_{01}u_{23}$ squares to -1, showing that both occur. Surjectivity follows by writing even permutations as an even number of transpositions. Hence $|\widetilde{S}| = 2|A_8| = 40320$. Both lifts of the double transposition $(01)(23)$ have square -1, so (D.25.3) has no group section.

For a seven-cycle $g = (i_0\ i_1\ \cdots\ i_6)$ with i_0 the least member of its support, put

$$v_g = \prod_{k=6,5,\dots,1} u_{i_0 i_k}, \qquad \widetilde{g} = \varepsilon_g v_g, \qquad v_g^7 = \varepsilon_g \in \{1, -1\}. \tag{D.25.4}$$

This is the unique lift of g of order seven. The other lift has order fourteen. Formula (D.25.4) has rational Clifford coefficients, with denominator a power of two, so all the following calculations use exact integers.

Write permutations in one-line notation: the entry in column i is the image of i, with columns numbered $0, \dots, 7$. Let

$$a = (1, 2, 3, 4, 5, 6, 0, 7)$$

and take the following five pairs, where $c_j = (ab_j)^{-1}$:

j	b_j	c_j	$s_j = \widetilde{a}\,\widetilde{b}_j\,\widetilde{c}_j$
0	$(0, 2, 5, 1, 6, 4, 7, 3)$	$(4, 0, 3, 1, 7, 5, 2, 6)$	-1
1	$(0, 2, 4, 1, 7, 3, 5, 6)$	$(7, 0, 3, 1, 5, 2, 6, 4)$	1
2	$(0, 2, 4, 7, 3, 6, 1, 5)$	$(5, 0, 6, 1, 4, 2, 7, 3)$	1
3	$(0, 2, 6, 5, 3, 1, 7, 4)$	$(2, 0, 5, 1, 4, 7, 3, 6)$	1
4	$(0, 2, 7, 4, 1, 3, 5, 6)$	$(7, 0, 4, 1, 5, 3, 6, 2)$	1.

(D.25.5)

Every entry a, b_j, c_j belongs to the same one of the two seven-cycle conjugacy classes in A_8. An explicit conjugator from a to any entry g is the permutation that maps $0, \dots, 6$ in order to the successive members of its seven-cycle, starting with its least member, and maps 7 to its fixed point. Each of the ten displayed conjugators is even; this is checked directly and saved in the certificate.

Each pair (a, b_j) generates S. Furthermore no automorphism of S takes (a, b_i) to (a, b_j) for $i \neq j$. The finite certificate for these two assertions is described below and contains a literal distinguishing relator for each of the ten pairs of components. It follows that the two tuples

$$g_1 = (a, a, a, a, a), \qquad g_2 = (b_0, b_1, b_2, b_3, b_4) \tag{D.25.6}$$

generate the entire group $G = S^5$. Here is the exact subdirect-product argument. A subdirect subgroup of a product of copies of a nonabelian simple group is a product of diagonal blocks. Inductively, after projecting to the first $r-1$ factors, its kernel in the last factor is either the full simple group or trivial. In the latter case the subgroup is the graph of an epimorphism $S^{r-1} \to S$. The commuting normal images of the factors show that exactly one factor maps isomorphically to S. This would identify two of the component epimorphisms up to an automorphism of S, which the distinguishing relators exclude. The certificate also proves simplicity of this particular S, so the argument uses no unverified finite-group recognition.

Set $E = \widetilde{S} \times S^4$, and define the source triple

$$\begin{aligned} X_1 &= (\widetilde{a}, a, a, a, a), \\ X_2 &= (\widetilde{b}_0, b_1, b_2, b_3, b_4), \\ X_3 &= (-\widetilde{c}_0, c_1, c_2, c_3, c_4). \end{aligned} \tag{D.25.7}$$

Its product is one by the first spin in (D.25.5). It generates E. In fact S is perfect by simplicity, and the derived subgroup of $\widetilde{S}$ maps onto S. If that derived subgroup missed the central element -1, it would be a complement, contradicting the nonsplitting proved above. Thus $\widetilde{S}$ and E are perfect. The image generated by X_1, X_2 already maps onto G; if it missed $\ker \pi$, it would have index two in E, giving a nontrivial abelian quotient of a perfect group. This proves the claimed epimorphism q_E.

Let

$$\alpha(v_0, v_1, v_2, v_3, v_4) = (v_1, v_2, v_3, v_4, v_0). \tag{D.25.8}$$

It has order five and preserves the three labelled conjugacy classes of $g_i = \pi(X_i)$ because all five entries in each coordinate triple are in the same class of S. For $0 \leq k \leq 4$, a conjugate of X_1 above $\alpha^k(g_1)$ must have first coordinate $\widetilde{a}$; a conjugate of X_2 above $\alpha^k(g_2)$ must have first coordinate $\widetilde{b}_k$. These assertions follow from their odd orders. A conjugate of X_3 has order fourteen, so its unique possible first coordinate above $\alpha^k(g_3)$ is $-\widetilde{c}_k$. The corresponding first-coordinate product is

$$\widetilde{a}\,\widetilde{b}_k(-\widetilde{c}_k) = -s_k = \begin{cases} 1, & k = 0, \\ -1, & 1 \leq k \leq 4. \end{cases} \tag{D.25.9}$$

The other coordinates multiply to one. Each of the individually prescribed elements is a conjugate of its original X_i: lift a conjugator in S and use uniqueness of the order-seven lift, also multiplying by -1 for the third coordinate. Thus (D.25.9) is the complete obstruction, after all conjugators have been allowed.

One can also check the base-centralizer quotient directly. The centralizer in S of a seven-cycle with one fixed point is its cyclic group of order seven. Hence $C_G(g_i) \simeq C_7^5$ for all three indices, and every map $\kappa_i : C_G(g_i) \to C_2$ in D.23 is zero. Consequently the obstruction group is C_2 and the four nonzero products in (D.25.9) survive it. An element $u \in \mathbb{Z}_5^\times$ gives

$\alpha^u = \alpha^k$ for a unique $k \in \{1,2,3,4\}$, proving the assertion about unit powers. The perfectness already proved gives the trivial maximal pro-five quotient maps exactly as in D.24.

For reproducibility, `a_peripheral_cycle.py` constructs all 20160 even permutations of eight letters and uses the Clifford multiplication rule

$$e_I e_J = (-1)^{\#\{(i,j)\in I\times J : i>j\}} e_{I\triangle J} \tag{D.25.10}$$

to evaluate (D.25.4). The selected triples, their full rational Clifford coefficients, their explicit conjugators, and the four negative products are retained in `a_peripheral_cycle_a8.json`. Closure under the two literal generators gives 20160 elements for each selected pair. For each pair of components, the program propagates the proposed generator map along these words; two words with the same source value and different target values give the saved relator witness. These witnesses exclude every possible automorphism carrying one pair to the other, without a classification of $\mathrm{Aut}(A_8)$. Finally it partitions the nonidentity elements into all thirteen conjugacy classes and verifies that the normal closure of a representative of each has order 20160. Every nontrivial normal subgroup contains such a representative, which proves simplicity. Their class sizes are

$$112, 210, 1344, 2520, 1120, 1680, 2880, 2880, 105, 3360, 1344, 1344, 1260;$$

their sum is 20159. The certificate also verifies $(u_{01}u_{23})^2 = -1$. The executed output ends with `PASS_ORDER_FIVE_WITH_CENTRAL_TWO_OBSTRUCTION`. The earlier seven-letter search, with eight inequivalent generating triples all having spin -1 in its fixed profile, is retained under the unsuffixed filenames and is not used as an exclusion outside that search domain. □

Statement D.26 (Proved). The exact covering restriction of the central peripheral obstruction. *Keep the finite groups, epimorphism $q_G : \widehat{F} \to G$, factor-swap automorphism α, and central extension $\pi : E \to G$ of D.24. Put $\theta = \alpha q_G$ and $S_i = \overline{\langle x_i \rangle}$, with $x_3 = (x_1x_2)^{-1}$. For any open subgroup $H \le \widehat{F}$ of index d, take representatives $t \in F = \langle x_1, x_2\rangle$ of $H\backslash\widehat{F}/S_i$ and set*

$$e_{i,t} = [S_i : S_i \cap t^{-1}Ht], \qquad h_{i,t} = t x_i^{e_{i,t}} t^{-1}, \qquad S_{i,t} = \overline{\langle h_{i,t}\rangle} \le H. \tag{D.26.1}$$

Let b_H be the number of these induced peripheral subgroups. The number of continuous homomorphisms $v : H \to E$ satisfying

$$\pi v = \theta|_H, \qquad v(h_{i,t}) \sim_E X_i^{e_{i,t}} \quad \text{for every } (i,t) \tag{D.26.2}$$

is exactly

$$\begin{cases} 0, & d \text{ odd}, \\ 2^{\,d+2-b_H}, & d \text{ even}. \end{cases} \tag{D.26.3}$$

These are literal weak lifts with the fixed base map; no surjectivity or quotient by conjugation is imposed on the count.

In particular every cover of odd index retains the nonzero peripheral obstruction of D.24. For every odd prime ℓ, this includes every open subgroup obtained as the inverse image of a subgroup of a finite ℓ-group quotient of $\widehat{F}$. The exact map controlling this assertion is restriction

$$\begin{gathered} H^2_{\mathrm{cts}}(\widehat{F}, \{S_i\}; \mathbb{F}_2) \longrightarrow H^2_{\mathrm{cts}}(H, \{S_{i,t}\}; \mathbb{F}_2), \\ \mathbb{F}_2 \ni a \longmapsto da \in \mathbb{F}_2, \end{gathered} \tag{D.26.4}$$

where both identifications use the sum of the oriented boundary coordinates. Thus passage to deeper covers of odd index cannot solve this specified finite peripheral problem.

Proof. The dense discrete free group F acts transitively on the d cosets of H. Its subgroup $H_0 = F \cap H$ has index d, and its profinite completion maps isomorphically onto H. Indeed the core in F of every finite-index subgroup of H_0 has finite index in F, proving that the induced topology is the full profinite topology. The discrete groups $\langle h_{i,t} \rangle$ complete to precisely the groups in (D.26.1). The indices $e_{i,t}$ are the orbit lengths of S_i on the d cosets, so

$$\sum_t e_{i,t} = d \quad (1 \le i \le 3). \tag{D.26.5}$$

Realize F as the fundamental group of an oriented sphere with three open discs removed, taking the oriented boundary loops to be x_1, x_2, x_3. Its connected cover associated to H_0 is an oriented compact surface with b_H boundary components. These components give exactly the loops $h_{i,t}$ with their specified orientations and base paths. Cutting this surface along a system of handle and boundary arcs gives the presentation

$$\left\langle a_1, b_1, \dots, a_g, b_g, c_1, \dots, c_{b_H} \;\middle|\; \prod_{j=1}^{g} [a_j, b_j] \prod_{k=1}^{b_H} c_k = 1 \right\rangle, \tag{D.26.6}$$

where the c_k are conjugates of the displayed induced boundary generators. Abelianization therefore gives exactly one relation among their classes, namely $\sum_k c_k = 0$. Since H_0 is free of rank $d+1$, comparison of ranks yields $2g = d + 2 - b_H$.

For trivial coefficients $\mathbb{F}_2$, continuous first cohomology of the free profinite and procyclic groups equals the first cohomology of their discrete dense free and cyclic groups: both are homomorphisms to the same finite group. Their second cohomology vanishes. The relative long exact sequences consequently identify both relative second groups with the cokernels of their boundary restriction maps. By (D.26.6), the latter map for H has image

$$\{(u_k) \in \mathbb{F}_2^{b_H} : \textstyle\sum_k u_k = 0\}, \tag{D.26.7}$$

and kernel $\mathbb{F}_2^{2g}$. For the base pair the corresponding map is $(a, b) \mapsto (a, b, -a - b)$. These calculations also prove the discrete-to-profinite comparison for the relative groups being used.

Under restriction, a boundary coordinate u_i pulls back to $e_{i,t}u_i$ on $S_{i,t}$; conjugation of base paths has no effect on these trivial coefficients. Summing the coordinates and using (D.26.5) gives $\sum_{i,t} e_{i,t}u_i = d \sum_i u_i$. Naturality of the relative long exact sequence proves exactly (D.26.4), including its source, target and the chosen identifications.

The relative extension class of D.24, with local lifts $U_i = (B_i, a_i)$, is $1 \in \mathbb{F}_2$ because $U_1U_2U_3 = (-I, 1)$. For a based induced peripheral group, choose any lift $\Theta_t \in E$ of $\theta(t)$ and use $\Theta_t U_i^{e_{i,t}} \Theta_t^{-1}$ as its local image. Changing Θ_t by the central kernel changes nothing. Its projection is $\theta(h_{i,t})$, and it is conjugate to $X_i^{e_{i,t}}$ by D.24. These are precisely the restricted local maps in the relative pullback extension, so their relative obstruction is $d \in \mathbb{F}_2$.

We check the full conjugacy freedom in (D.26.2). Every X_i has odd order. In a central extension by C_2, the two elements above any odd-order base element differ by the central involution, and exactly one has odd order. Thus the displayed local image is the unique odd-order lift of $\theta(h_{i,t})$. Every conjugate allowed in (D.26.2) has odd order and equals that lift. Equivalently, each induced base-centralizer commutator is killed both by two and by an odd order, hence is trivial. This remains true for conjugators in all of E and for a proper image $\theta(H) \le G$. The relative obstruction therefore retains the complete unframed peripheral condition.

For completeness, existence and the count follow directly from cochains. The pullback of E to the free profinite group H has a global section: lift its $d+1$ free generators arbitrarily. The discrepancies between this section and the prescribed local maps form a vector in $\mathbb{F}_2^{b_H}$. Multiplication of the section by a character $H \to C_2$ changes that vector by the boundary

restriction map. Its sum is the relative class d. Equation (D.26.7) gives no correction for odd d and a torsor under $\mathbb{F}_2^{2g}$ for even d. This proves the exact count (D.26.3) without imposing an extra generation condition.

The program `d_peripheral_covers.py` constructs connected coset covers of indices one through nine, with cyclic and reflection permutation data. For each it computes literal Schreier basis words and every induced boundary word. It enumerates every central correction to the lifted basis and substitutes every surviving lift into all boundary equations in $\mathrm{SL}_2(\mathbb{F}_{11})$. The computed counts agree with (D.26.3), including positive-genus even covers. The code, full coset and word data, every correction, and the executed output are retained. The all-index conclusion follows from the restriction calculation above. □

4.10 A coupled perfect kernel with admitted and obstructed pro-five branches

Statement D.27 (Proved). Exact coupled fibres over normalized finite pro-five shadows. *There are explicit two-generated finite groups*

$$E_0 \xrightarrow{\pi_0} G_0 \xrightarrow{\rho} C_5, \qquad |\ker \pi_0| = 2, \tag{D.27.1}$$

whose respective kernels over C_5 are perfect, and in which C_5 acts nontrivially on those kernels by permutation of five simple factors in each of five rows. The groups have an order-five, multiplier-one peripheral automorphism $R \in \mathrm{Aut}(G_0)$ inducing the identity on C_5. Each of R, R^2, R^3, R^4 is obstructed at the central kernel in (D.27.1).

Here is their exact comparison with a prescribed pro-five component. Let $L = \langle l_1, l_2 \rangle$ be a finite 5-group with an epimorphism $\eta : L \to C_5 = \langle t \rangle$ satisfying $\eta(l_1) = t$, $\eta(l_2) = 1$, and put $l_3 = (l_1 l_2)^{-1}$. Let $\beta \in \mathrm{Aut}(L)$ be specified by the peripheral equalities

$$\beta(l_1) = l_1, \qquad \beta(l_2) = f^{-1} l_2 f, \quad \eta(f) = 1, \qquad \beta(l_3) = h l_3 h^{-1}. \tag{D.27.2}$$

The condition on f holds in particular for the usual normalized multiplier-one form with $f \in L'$. Define the actual fibre products

$$E_L = E_0 \times_{C_5} L, \qquad G_L = G_0 \times_{C_5} L. \tag{D.27.3}$$

The paired source generators give epimorphisms $\widehat{F} \twoheadrightarrow E_L \twoheadrightarrow G_L$, and the coordinate maps $E_L \to L$ and $G_L \to L$ are their maximal pro-five quotient maps. The automorphism

$$\alpha_- = (R, \beta) \in \mathrm{Aut}(G_L) \tag{D.27.4}$$

has the prescribed three peripheral conjugacy classes and has no weak peripheral lift to E_L. On the same quotient and over the same β, the automorphism $\alpha_+ = (1, \beta)$ has the explicit generating peripheral lift $(1, \beta) \in \mathrm{Aut}(E_L)$.

The construction includes an explicit nonidentity β of order five on a finite two-generated 5-group, given in (D.27.15)–(D.27.18) *below. Thus the perfect kernel, its action by the quotient, and the prescribed pro-five automorphism can all be nontrivial. The admitted branch is proved for every datum* (D.27.2)*; these coupled quotients therefore do not exclude such a prescribed pro-five shadow.*

Proof. Use $S = A_8$, its explicit double cover $\widetilde{S}$, and the five triples (a, b_j, c_j) from D.25, with spins $s_0 = -1$ and $s_1 = \cdots = s_4 = 1$. Index a copy of S^{25} by $(j, k) \in (\mathbb{Z}/5)^2$, where j labels rows and k labels columns. Put

$$P = S^{\{(j,k)\}}, \qquad (t \cdot p)_{j,k} = p_{j,k-1}, \qquad G_0 = P \rtimes \langle t \rangle. \tag{D.27.5}$$

In one row write $\delta_k(s)$ for the tuple equal to s in column k and to one elsewhere. The source generators are

$$\begin{aligned} g_1 &= ((\delta_0(a))_{j=0}^4, t), \\ g_2 &= ((\delta_0(b_j))_{j=0}^4, 1), \qquad g_3 = (g_1 g_2)^{-1}. \end{aligned} \tag{D.27.6}$$

These are literal elements of the semidirect product with multiplication $(p, t^r)(q, t^s) = (p(t^r \cdot q), t^{r+s})$.

We prove generation and retain the covering map. The subgroup $H = \ker(\widehat{F} \to C_5)$, with $x \mapsto t, y \mapsto 1$, has the Schreier basis

$$x^5, \quad y_k = x^k y x^{-k} \quad (0 \le k \le 4). \tag{D.27.7}$$

Applying (D.27.6) gives $x^5 \mapsto (a)_{j,k}$ and $y_k \mapsto (\delta_k(b_j))_j$. In coordinate (j, k) this is the epimorphism

$$x^5 \longmapsto a, \qquad y_k \longmapsto b_j, \qquad y_r \longmapsto 1 \quad (r \ne k). \tag{D.27.8}$$

For distinct columns the kernels differ already on one of the y_r. In the same column and different rows they differ by the relator witnesses of D.25, after replacing its first and second generators by x^5 and y_k. The simple-group subdirect argument of D.25 consequently makes the image of H exactly P. Since g_1 maps to t, the two source elements generate all of G_0.

Define the central product

$$P_0^* = \widetilde{S}^5 / \{(\epsilon_0, \dots, \epsilon_4) \in \{1, -1\}^5 : \prod_k \epsilon_k = 1\}, \qquad \widetilde{P} = P_0^* \times (S^5)^4. \tag{D.27.9}$$

The kernel of $P_0^* \to S^5$ is C_2, identified with the product of its five central signs. Column shift preserves this defining kernel, so it acts on $\widetilde{P}$. Set $E_0 = \widetilde{P} \rtimes C_5$ and let $\pi_0 : E_0 \to G_0$ be the induced central extension. Lift g_1, g_2 by replacing their row-zero entries a, b_0 with their canonical order-seven lifts in $\widetilde{S}$; call the resulting elements X_1, X_2, and put $X_3 = (X_1 X_2)^{-1}$. The group $\widetilde{P}$ is perfect: it is a product of a quotient of the perfect group $\widetilde{S}^5$ and copies of the perfect group S. Consequently $E_0^{\mathrm{ab}} = C_5$. The subgroup generated by X_1, X_2 maps onto G_0; missing the central kernel would make it an index-two subgroup of E_0, contrary to this abelianization. This proves the source epimorphism to E_0.

Let R rotate the row index by $(Rp)_{j,k} = p_{j+1,k}$ and fix t. It commutes with the column action and is an automorphism of G_0 of order five. It fixes g_1 and preserves the conjugacy classes of g_2, g_3. For g_2 this is coordinatewise conjugacy of the b_j in S. For g_3 it is the cyclic-product comparison in the wreath product, whose exact conjugator can be described as follows. For $v = (v_k, t^\epsilon)$ and $v' = (v'_k, t^\epsilon)$, with $\epsilon = \pm 1$, a base conjugator $(d_k, 1)$ must satisfy

$$d_k v_k d_{k-\epsilon}^{-1} = v'_k, \qquad d_{k-\epsilon} = (v'_k)^{-1} d_k v_k. \tag{D.27.10}$$

Choose d_0 conjugating the two cycle products. The displayed recursion determines all five d_k; its last equation is exactly that cycle-product conjugacy, so it is consistent. In row j, the zeroth coordinate of g_3^5 is c_j. Those c_j are conjugate, and (D.27.10) supplies the required conjugator in P for every row rotation.

The peripheral base orders are $(35, 7, 35)$. To evaluate their lifts, temporarily replace b_0 in the doubled row by b_j and use canonical odd lifts $\widetilde{a}, \widetilde{b}_j$. The first two lifted peripheral orders are $35, 7$. The fifth power of their product has, in its five columns, conjugates of $\widetilde{a}\widetilde{b}_j$; its thirty-fifth power is the tuple of five signs s_j. Their product in the central quotient (D.27.9) is $s_j^5 = s_j$. Therefore the natural inverse of that product has order seventy for $j = 0$ and order thirty-five for $j \ne 0$. The original X_3 has order seventy.

In a central extension by C_2, an odd-order base element has precisely one odd-order lift and one even-order lift. Every permitted conjugate above a fixed base element is therefore determined by its order. For R^j with $j \neq 0$ the first two required lifts are the canonical odd ones, and the third is the negative of the natural inverse just computed, in order to have order seventy. Their product is the nonidentity element of $\ker \pi_0$. Lifting the base conjugators from (D.27.10) proves that all three individually required elements are actual conjugates. Thus no remaining conjugacy choice can alter this product. This proves the obstruction for all four nonzero powers of R.

We now form (D.27.3). Equations (D.27.2) give $\eta\beta = \eta$ on the two generators, hence on L, so both displayed product automorphisms are defined. The source maps are literally

$$x \longmapsto (X_1, l_1), \qquad y \longmapsto (X_2, l_2), \tag{D.27.11}$$

and their projections to G_L. For surjectivity, put $H_L = \ker(\widehat{F} \to L)$. Then $H_L \triangleleft H$ and $H/H_L \simeq \ker \eta$ is a 5-group. The image of H in $\widetilde{P}$ is all of $\widetilde{P}$ by the proved source epimorphism. Its quotient by the image of H_L is a 5-group and a quotient of a perfect group, hence trivial. Thus H_L maps onto $\widetilde{P}$. The image in E_L already maps onto L and contains its entire kernel, proving surjectivity. The proof for G_L is identical with P. The respective kernels $\widetilde{P}, P$ of the coordinate maps to L are perfect. Every homomorphism to a finite 5-group kills them, and the coordinate map is itself onto the 5-group L. This proves exactly the asserted maximal pro-five quotient maps.

The admitted branch is the literal automorphism $(e_0, l) \mapsto (e_0, \beta(l))$ of E_L. It fixes (X_1, l_1). It conjugates the second source element by $(1, f^{-1}) \in E_L$, since $\eta(f) = 1$. For the third, write $\eta(h) = t^r$. Since $\rho\pi_0(X_3) = t^{-1}$, the pair

$$(X_3^{-r}, h) \in E_L \tag{D.27.12}$$

is defined and conjugates (X_3, l_3) to $(X_3, \beta(l_3))$: its first coordinate centralizes X_3. This supplies the required third conjugator with the correct fibre-product condition.

For the obstructed branch, choose $d_2, d_3 \in P$ with $d_i g_i d_i^{-1} = R(g_i)$, as constructed above. The second and third base conjugators in G_L are respectively

$$(d_2, f^{-1}), \qquad (d_3 g_3^{-r}, h). \tag{D.27.13}$$

The first has matching trivial C_5 coordinates. The second has matching coordinate t^r, and multiplying by g_3^{-r} does not change its conjugation of g_3. They prove that (R, β) has precisely the required peripheral conjugacies. A weak lift to E_L would, upon projection to E_0, be a weak peripheral lift of R for the source (D.27.6), which has just been excluded. The same projection excludes every integer power with exponent prime to five.

This is an actual semidirect-product coupling. For example, conjugation by $(1, l_1)$ sends a nonidentity element supported in column zero of P to column one. The typed fibre-product identification is

$$G_0 \times_{C_5} L \simeq P \rtimes L, \quad ((p, \eta(l)), l) \longmapsto (p, l), \tag{D.27.14}$$

with the specified action through η; the analogous map for E_L uses $\widetilde{P}$. No direct-product replacement is made.

We give a concrete nonidentity datum (D.27.2). Let $\mathcal{R}$ be the following finite algebra, let J be its augmentation ideal, and define the finite group L by

$$\begin{aligned} \mathcal{R} &= \mathbf{F}_5\langle X, Y\rangle/(\text{words of length at least } 5), \\ L &= \langle l_1 = \exp X, l_2 = \exp Y\rangle \leq 1 + J. \end{aligned} \tag{D.27.15}$$

The finite exponential and logarithm use only denominators $1, 2, 3, 4$ and $1!, \ldots, 4!$, all invertible in $\mathbf{F}_5$. The ambient group has 5^{30} elements and exponent five, since $J^5 = 0$ and

$(1+a)^5 = 1 + a^5 = 1$. Its degree-one X coefficient gives the epimorphism $\eta : L \to C_5$ in the statement. In the Lie bracket of this associative algebra put

$$H_2 = [X, Y], \qquad F_3 = [X + Y, H_2], \qquad H_3 = [Y, H_2], \qquad \Delta = [F_3, Y]. \tag{D.27.16}$$

The identity $\Delta = [H_3, X + Y]$ follows from Jacobi and $[H_2, H_2] = 0$. The substitution

$$X \longmapsto X, \qquad Y \longmapsto Y + \Delta \tag{D.27.17}$$

is an automorphism of $\mathcal{R}$ of order five: substitution in the degree-four polynomial Δ leaves it unchanged. The polynomial is nonzero, for example its $XXYY$ coefficient is one. With $F = \exp F_3$ and $H = \exp H_3$, direct expansion gives

$$\beta(l_1) = l_1, \qquad \beta(l_2) = F l_2 F^{-1}, \qquad \beta((l_1 l_2)^{-1}) = H (l_1 l_2)^{-1} H^{-1}. \tag{D.27.18}$$

The last equality follows also by applying the substitution to $\log((\exp X \exp Y)^{-1})$: its only changed term is $-Y$, whose change is $-\Delta = -[H_3, X + Y]$.

We verify membership of the two conjugators in L by actual words. Write $c = [l_1, l_2]$, $u = [l_1, c]$, $v = [l_2, c]$, and $d_1 = [l_1, u]$, $d_2 = [l_2, u]$, $d_3 = [l_2, v]$. The d_i are central at this truncation. Multiplication in $\mathcal{R}$ gives the exact identities

$$F = u v d_1^{-1} d_2^{-1} d_3^{-1}, \qquad H = v d_2^{-3} d_3^{-1}. \tag{D.27.19}$$

For example the degree-four errors in $\log(uv)$ are the three $\log d_i$ with coefficients $(1, 1, 1)$, and those in $\log v$ have coefficients $(0, 3, 1)$ over $\mathbf{F}_5$; higher terms vanish. Thus $F, H \in L'$, and (D.27.18) proves that the substitution restricts to an automorphism of L of order five, with the normalized conjugator $f = F^{-1}$. It is nonidentity because $\beta(l_2) - l_2$ has the nonzero leading term Δ. For this explicit datum (R, β) also has order five, acts nontrivially on both the perfect kernel and the maximal pro-five quotient, and all four nonzero powers retain the central obstruction.

The program `a_peripheral_coupled.py` checks all twenty-five maps (D.27.8), saves all three hundred pairwise kernel witnesses, and evaluates the wreath conjugators (D.27.10) literally. It computes the central product using the Clifford multiplication of D.25, obtaining the peripheral orders $(35, 7, 70)$ in row zero and $(35, 7, 35)$ in the other rows, and the allowed products $1, -1, -1, -1, -1$. Independently, it verifies every identity (D.27.16)–(D.27.19) by truncated noncommutative polynomials over $\mathbf{F}_5$, retaining the literal conjugator words and their exact coefficients. Its output and JSON certificate end with `PASS_COUPLED_WREATH_AND_PRO_FIVE_PULLBACK`. □

4.11 A proved family of nonabelian chief lifts

Statement D.28 (Proved). Generating relative lifts for transitive A_5 wreath kernels. *Let $D \leq \mathrm{Sym}(\Omega)$ be a transitive finite 5-group, let $n = |\Omega| \geq 25$, and let $q : \widehat{F} = \langle x_1, x_2 \rangle \twoheadrightarrow D$ be an epimorphism. Put $x_3 = (x_1 x_2)^{-1}$ and $d_i = q(x_i)$. Assume that each of the three permutations d_i has no fixed point on Ω. Give Ω its right D-action, and write*

$$M = A_5^{\Omega}, \qquad (d \cdot a)(\omega) = a(\omega d), \qquad W = M \rtimes D. \tag{D.28.1}$$

For every three elements $U_i \in W$ above d_i there exist $a_i \in M$ such that

$$V_i = a_i U_i a_i^{-1}, \qquad V_1 V_2 V_3 = 1, \qquad \langle V_1, V_2 \rangle = W. \tag{D.28.2}$$

Equivalently, every such finite relative lifting problem has a generating solution, with the three base conjugators fixed. The kernel M is a nonabelian chief factor: it is a product of nonabelian simple groups, and D permutes its factors transitively.

In particular this applies to the regular action of every noncyclic two-generated finite 5-group D. It also has the following exact coupled consequence. Let L be a finite 5-group with epimorphism $\eta : L \to D$, and suppose specified epimorphisms from $\widehat{F}$ to W and L have the same projection to D. Let $\beta \in \mathrm{Aut}(L)$ induce an automorphism $\alpha \in \mathrm{Aut}(D)$, and suppose its three source peripheral images satisfy

$$\beta(l_i) = h_i l_i^{\lambda} h_i^{-1} \quad (1 \le i \le 3), \qquad \lambda \in \widehat{\mathbb{Z}}^{\times}. \tag{D.28.3}$$

Then β has a generating peripheral lift from $\widehat{F}$ to the actual fibre product $W \times_D L$, over the map βq_L, with the same multiplier λ and with the given h_i retained in the L-coordinates of its conjugators.

Proof. We first record all finite-group input used in the proof. Exact enumeration of the sixty even permutations of five letters gives the following commutator counts:

element order	1	3	2	5	5
conjugacy-class size	1	20	15	12	12
$\#\{(s,t) : [s,t] = u\}$	300	63	32	65	65.

(D.28.4)

In particular every element of $S = A_5$ is a commutator. The complete subgroup enumeration has fifty-nine subgroups; the only normal subgroup orders are $1, 60$. Its maximal proper subgroups consist of five of order twelve, six of order ten, and ten of order six. Thus S is nonabelian simple and perfect, and every proper subgroup is contained in one of twenty-one subgroups of size at most twelve. Enumeration of every possible image of a fixed generating pair gives $|\mathrm{Aut}(S)| = 120$; the sixty inner automorphisms are distinct, so $Z(S) = 1$ and $\mathrm{Out}(S) \simeq C_2$. These are finite certificates, described at the end of the proof.

Choose $\omega_0 \in \Omega$, and put

$$H = q^{-1}(D_{\omega_0}), \qquad K = \ker q = \mathrm{core}_{\widehat{F}}(H). \tag{D.28.5}$$

The equality with the core follows from the faithful permutation action $D \le \mathrm{Sym}(\Omega)$. The subgroup H has index n and is the profinite completion of the corresponding degree-n cover of the oriented three-boundary sphere, as in D.26. Let its genus be g and the number of its boundary components be b. Every cycle of any d_i has length at least five, so

$$b \le \frac{3n}{5}, \qquad 2g = n + 2 - b, \qquad 2g - 2 \ge \frac{2n}{5}, \qquad g \ge 6. \tag{D.28.6}$$

We prove a uniform representation statement for this cover. Prescribe an arbitrary conjugacy class C_j of S for each of its b oriented boundary loops. Put $P_C = \prod_j |C_j| \ge 1$. The surface presentation

$$\left\langle A_1, B_1, \dots, A_g, B_g, c_1, \dots, c_b \;\middle|\; \prod_{r=1}^{g} [A_r, B_r] \prod_{j=1}^{b} c_j = 1 \right\rangle \tag{D.28.7}$$

shows that the number of continuous homomorphisms $\varphi : H \to S$ taking c_j into C_j is at least

$$60^{2g-2} P_C. \tag{D.28.8}$$

Indeed choose all b boundary values and the first $g - 1$ handle pairs arbitrarily. The required value of the last commutator has a solution by (D.28.4). Each such choice gives at least one distinct homomorphism; all maps are continuous because their target is finite and H is the completion of the displayed finitely generated group.

The maps with proper image in S number at most

$$21 \cdot 12^{2g} P_C. \tag{D.28.9}$$

For each maximal proper subgroup, choose its handle values and the boundary values in its intersections with the C_j, and ignore the product relation to obtain an upper bound. Summing over the twenty-one maximal subgroups gives (D.28.9).

Among the surjective maps we must also exclude repeated coordinate kernels. For such a φ, the restriction $\varphi|_K$ is onto S: the quotient of S by its normal subgroup $\varphi(K)$ is a quotient of the 5-group H/K, whereas S is perfect. Put $N = \ker(\varphi|_K)$ and $J = N_{\widehat{F}}(N)$. One has $H \leq J$. If $J > H$, conjugation on $K/N \simeq S$ defines a continuous map $J \to \mathrm{Aut}(S)$. Its composite to $\mathrm{Out}(S)$ factors through the 5-group J/K, and is consequently trivial. Centerlessness therefore gives an actual homomorphism

$$\widetilde{\varphi} : J \longrightarrow S, \qquad \mathrm{Inn}(\widetilde{\varphi}(t)) = (kN \longmapsto tkt^{-1}N). \tag{D.28.10}$$

On H this equals φ, since their inner automorphisms of the surjective quotient K/N agree. It is the unique extension: an image of t in any extension is determined by its conjugation on S, whose center is trivial.

There is an intermediate open subgroup $H < H' \leq J$ with $[H' : H] = 5$. To justify this step, apply the normalizer property of finite 5-groups to $H/K < J/K$: a proper subgroup is strictly contained in its normalizer. That property follows by induction from the center: either a central element is outside the subgroup, or one passes to the quotient by the center. Choose a subgroup of order five in the resulting nontrivial normalizer quotient and take its inverse image. This gives the required H'. It is determined by the set of its five right H-cosets, which contains the identity coset. There are at most $\binom{n-1}{4}$ possibilities for H'. Each is free profinite of rank $1 + n/5$, and hence has exactly $60^{1+n/5}$ homomorphisms to S. Every surjective φ with $J > H$ is the restriction of one of these homomorphisms. The number of such maps is therefore at most

$$\binom{n-1}{4} 60^{1+n/5}. \tag{D.28.11}$$

For a regular action one may replace the binomial factor by the number of order-five subgroups of D, at most $(n-1)/4$; the proof below uses the larger bound so it also covers the stated transitive actions.

The two bad-map bounds are strictly smaller than (D.28.8). After division by that lower bound, they are at most

$$\frac{75600}{5^{2g}} + \binom{n-1}{4} 60^{1-n/5} \leq \frac{75600}{5^{12}} + \frac{\binom{24}{4}}{60^4} = \frac{38122819}{33750000000} < \frac{1}{500} < 1. \tag{D.28.12}$$

For the second inequality, the binomial expression decreases when n is increased by five from any $n \geq 25$: its successive ratio is bounded by $(29/21)^4/60 = 707281/11668860 < 1$. All possible n are powers of five, so this estimate applies to them. We have proved that every boundary class pattern has a surjective φ with $N_{\widehat{F}}(\ker \varphi|_K) = H$.

We now give the exact nonabelian Shapiro map from this surface representation to the required source lift. Choose $t_\omega \in \widehat{F}$ representing the right coset Ht_ω, with $t_{\omega_0} = 1$, so that $\omega_0 q(t_\omega) = \omega$. Define

$$c_\varphi(g)(\omega) = \varphi(t_\omega g t_{\omega q(g)}^{-1}), \qquad f_\varphi(g) = (c_\varphi(g), q(g)) \in W. \tag{D.28.13}$$

The element inside φ belongs to H. Factoring $t_\omega g h t^{-1}_{\omega q(gh)}$ at $t_{\omega q(g)}$ gives exactly

$$c_\varphi(gh)(\omega) = c_\varphi(g)(\omega)c_\varphi(h)(\omega q(g)), \tag{D.28.14}$$

so f_φ is a continuous homomorphism over q. For a cycle of d_i of length e, starting at ω, its cycle product is

$$\prod_{r=0}^{e-1} c_\varphi(x_i)(\omega d_i^r) = \varphi(t_\omega x_i^e t_\omega^{-1}). \tag{D.28.15}$$

The elements on the right are exactly the induced oriented boundary generators in (D.28.7), up to their specified change of base paths.

For completeness, conjugation of (v, d_i) by $a \in M$ changes its function to $a(\omega)v(\omega)a(\omega d_i)^{-1}$. On each cycle its product is conjugated by the value $a(\omega)$ at the chosen starting point. Conversely, a conjugator between the two cycle products determines all its other values on that cycle by the recursion

$$a(\omega d_i) = v'(\omega)^{-1} a(\omega) v(\omega). \tag{D.28.16}$$

The final equation is precisely the cycle-product conjugacy. Thus M-conjugacy classes above d_i are exactly the independently labelled S-conjugacy classes of these cycle products. Prescribe in (D.28.7) the classes arising from U_i. Equations (D.28.15)–(D.28.16) give $f_\varphi(x_i) = a_i U_i a_i^{-1}$, with actual $a_i \in M$, proving the required peripheral conditions.

It remains to check generation, not just existence of a weak lift. For $k \in K$, the coordinate maps of $c_\varphi(k)$ are $k \mapsto \varphi(t_\omega k t_\omega^{-1})$. They are onto S, with kernels $t_\omega^{-1} N t_\omega$. Two such kernels coincide exactly when $t_{\omega'} t_\omega^{-1} \in J$. Since $J = H$, this happens exactly for the same right H-coset. The kernels are therefore pairwise distinct. The subdirect-product proof in D.25 now gives $f_\varphi(K) = M$: a missing simple factor would force one of the component epimorphisms to be an automorphic copy of another and hence give equal kernels. Together with $q(\widehat{F}) = D$, this proves $f_\varphi(\widehat{F}) = W$, completing (D.28.2).

If D is noncyclic and its action is regular, each d_i is nonidentity: $d_1 = 1$, $d_2 = 1$, or $d_3 = 1$ would make the two-generated group cyclic. Every cycle then has the nontrivial 5-power length $|d_i| \geq 5$, and $n = |D| \geq 25$. This proves the regular-case assertion.

We finally retain the prescribed base conjugators in the coupled application. Let $X_i \in W$ and $l_i \in L$ be the source peripheral triples. Choose $B_i \in W$ above $\eta(h_i)$ and put $U_i = B_i X_i^\lambda B_i^{-1}$. These elements lie above $\alpha(d_i)$. The three new permutations have the same cycle lengths as the original d_i: they are conjugate to their unit powers. Apply (D.28.2) to the source map αq to obtain an epimorphism $f_W : \widehat{F} \to W$ with $f_W(x_i) = a_i U_i a_i^{-1}$. The map

$$\widehat{F} \longrightarrow W \times_D L, \qquad g \longmapsto (f_W(g), \beta q_L(g)) \tag{D.28.17}$$

is defined because both components project to $\alpha q(g)$. Its peripheral conjugators are the literal pairs $(a_i B_i, h_i)$, whose images in D agree, so it retains all three h_i.

To prove surjectivity of (D.28.17), write $K_L = \ker q_L$ and $K_D = \ker q$. The image of K_D in W under f_W is M. Its quotient by the image of K_L is a quotient of $K_D/K_L \simeq \ker \eta$, a 5-group, and also a quotient of the perfect group M; it is trivial. Thus (D.28.17) contains the whole kernel M over L, and maps onto L, which proves surjectivity. The same argument proves that the original paired source is an epimorphism. In these fibre products the projection to L is the maximal pro-five quotient, since every map to a finite 5-group kills the perfect kernel M.

The program `a_peripheral_a5_wreath.py` provides all finite constants in this proof. It enumerates all 3600 commutators and all subgroups by successively adjoining a group element, yielding the complete subgroup lattice and its normal and maximal subgroups. It tests all 60^2 possible images of the fixed generating pair, propagating the map along all generator

edges, and saves all 120 automorphisms. It checks (D.28.12) and its monotonic ratio by exact fractions. For the first regular example, $D = C_5^2$ with source directions $(1,0),(0,1),(-1,-1)$, it records all three permutations of its twenty-five points, their five cycles each, the genus six, and all six order-five subgroups. The source, output and JSON certificate are retained; the output ends with `PASS_UNIFORM_A5_WREATH_LIFTING_CONSTANTS`. □

The no-fixed-point and degree hypotheses in D.28 are used in the exact genus bound (D.28.6). The admitted family includes nonabelian chief kernels with their nontrivial permutation action and all prescribed peripheral base conjugators. It is not asserted cofinal among the finite quotients relevant to D.1 or D.2. For example, the five-column cover underlying D.27 has boundary counts $1, 5, 1$ and genus zero; its relation has no handle commutators for the count in (D.28.8). The explicit maps of D.27 and D.28 specify the respective finite lifting conclusions without resolving the remaining cofinal comparison.

4.12 All A_5 chief extensions over finite odd-order groups

Statement D.29 (Proved). Exact peripheral count invariance for every A_5 chief step over an odd-order group. *Let*

$$1 \longrightarrow A \longrightarrow E \xrightarrow{\pi} D \longrightarrow 1, \qquad A \simeq A_5^{\Omega}, \tag{D.29.1}$$

be a finite extension, with D of odd order and A a nonabelian chief factor of E. Let $q_E : \widehat{F} \twoheadrightarrow E$ be an epimorphism, $X_i = q_E(x_i)$, $q = \pi q_E$, and $d_i = q(x_i)$. Let $q_1 : \widehat{F} \twoheadrightarrow D$ be any epimorphism, let $\lambda \in \widehat{\mathbb{Z}}^{\times}$, and let $h_i \in D$ be specified elements such that

$$q_1(x_i) = h_i d_i^{\lambda} h_i^{-1}. \tag{D.29.2}$$

choose any lifts $H_i \in E$ of h_i. There is an epimorphism $f : \widehat{F} \twoheadrightarrow E$ such that

$$\pi f = q_1, \qquad f(x_i) = a_i H_i X_i^{\lambda} H_i^{-1} a_i^{-1} \quad \text{for some } a_i \in A. \tag{D.29.3}$$

The number of these literal epimorphisms is exactly the number for $(q_1, \lambda, h_1, h_2, h_3) = (q, 1, 1, 1, 1)$. In particular it is positive. There is no degree or fixed-point restriction on the transitive factor action on Ω.

The same assertion holds in every actual fibre product $E \times_D L$, for a finite odd-order group L and a prescribed peripheral epimorphism $q_{L,1} : \widehat{F} \to L$ over q_1, relative to the original source epimorphism $q_L : \widehat{F} \to L$: there is a generating lift over $q_{L,1}$, with its specified peripheral conjugators retained in the L-coordinate. Here both maps to L compose to the indicated maps to D, and the new peripheral values are the specified conjugates of the λth powers of the original ones. Thus every A_5 chief extension over a finite odd-order group, including its coupled pullbacks, admits all the peripheral shadows described in (D.29.2)–(D.29.3)*. In particular this applies over every finite ℓ-group for every odd prime ℓ.*

Proof. Put $S = A_5$. We first give the exact map from (D.29.1) to a permutation extension, including the possible outer twisting. Every automorphism of S^{Ω} permutes its simple direct factors and acts on each factor by an automorphism of S. The computation in D.28 gives

$$Z(S) = 1, \qquad \mathrm{Out}(A) = \mathbf{F}_2^{\Omega} \rtimes \mathrm{Sym}(\Omega). \tag{D.29.4}$$

The factor action $\rho : D \to \mathrm{Sym}(\Omega)$ is transitive: a union of factor orbits would otherwise give a proper nonzero normal subgroup inside the chief kernel. The outer action of D is of the form $(b_d, \rho(d))$, with $b_{de} = b_d + \rho(d) b_e$. Since $|D|$ is odd, the vector

$$B = |D|^{-1} \sum_{e \in D} b_e \in \mathbf{F}_2^{\Omega} \quad \text{satisfies} \quad b_d = B - \rho(d) B. \tag{D.29.5}$$

This follows by summing the cocycle identity over e. Choose a componentwise automorphism τ_B of A with outer class B. Such a representative is explicit: on each component with value one use conjugation by the odd permutation $(01) \in S_5$; on the other components use the identity. Conjugating the outer action by B therefore makes it exactly ρ.

Here is the corresponding extension isomorphism, rather than only an identification of outer actions. Since $Z(A) = 1$, the map

$$E \longrightarrow \mathrm{Aut}(A) \times_{\mathrm{Out}(A)} D, \quad e \longmapsto (\mathrm{Ad}_e, \pi(e)) \tag{D.29.6}$$

is an isomorphism. It is injective because its kernel lies in $A \cap C_E(A) = Z(A)$. For surjectivity, an automorphism in the required outer class differs from the conjugation of any chosen lift of d by a unique inner automorphism of A. After applying τ_B^{-1} to this kernel identification, there is for every $e \in E$ a unique $a_e \in A$ with

$$\tau_B^{-1} \,\mathrm{Ad}_e\, \tau_B = \mathrm{Inn}(a_e)\rho(\pi(e)). \tag{D.29.7}$$

Uniqueness uses centerlessness. Multiplication gives $a_{ef} = a_e \rho(\pi(e))(a_f)$, so $e \mapsto (a_e, \pi(e))$ is the actual isomorphism $E \simeq A \rtimes_\rho D$. For example, after choosing a lift e_d of d, the inverse image of (a, d) is $\tau_B(a a_{e_d}^{-1}) e_d$. This map transports the specified source elements and all conjugators in (D.29.3). We may perform the rest of the calculation in this displayed semidirect product.

Write the factor action on the right as in D.28, choose $\omega_0 \in \Omega$, and let D_* be its stabilizer. Thus $\Omega \simeq D_* \backslash D$, and put $n = |\Omega|$. For every subgroup $D_* \le C \le D$ use the two actual covers

$$H_C = q^{-1}(C), \qquad H'_C = q_1^{-1}(C), \qquad K_0 = \ker q, \quad K_1 = \ker q_1. \tag{D.29.8}$$

The action on Ω is allowed to be nonfaithful; the quotients $H_C/K_0 = C$ and $H'_C/K_1 = C$ still have odd order. No identification of K_0 and K_1 is made.

The original peripheral elements X_i determine conjugacy classes of S on every induced boundary component of H_{D_*}. These are their cycle-product classes under the exact map (D.28.15). The prescribed elements $H_i X_i^\lambda H_i^{-1}$ determine the new boundary classes for H'_{D_*}. For each i and C, the boundary comparison is the explicit bijection

$$\begin{aligned} \sigma_{i,C} : C\backslash D/\langle q_1(x_i)\rangle &\longrightarrow C\backslash D/\langle d_i\rangle, \\ Cd\langle q_1(x_i)\rangle &\longmapsto Cdh_i\langle d_i\rangle. \end{aligned} \tag{D.29.9}$$

The identity $q_1(x_i)h_i = h_i d_i^\lambda$ proves that this is well defined, and that it preserves cycle lengths. The common exponent is a unit, so d_i and d_i^λ have the same cycles. Right multiplication by h_i also commutes with every projection $D_*\backslash D \to C\backslash D$. Consequently (D.29.9) respects the partition of boundary components under each covering $H_{D_*} \le H_C$ and preserves every relative boundary degree.

The new class at a boundary sent to an old class C_γ by (D.29.9) is exactly C_γ^λ. Indeed a cycle of length e in X_i^λ has product equal to the corresponding coordinate of $X_i^{\lambda e}$, which is the λth power of the cycle product of X_i. This remains true when $d_i^e \neq 1$ in D: it fixes that coordinate, which is all that is needed for the power identity. Conjugation by the kernel component of H_i changes only the conjugating element in S. This proves the class comparison with the specified base conjugators, including fixed peripheral points.

For each boundary component δ of H_C, let the components γ above it in H_{D_*} have relative degrees r_γ. Define the actual conjugacy-invariant subset of S

$$T_{C,\delta} = \bigcap_{\gamma \mapsto \delta} \{s \in S : s^{r_\gamma} \in C_\gamma\}. \tag{D.29.10}$$

A map $\psi : H_C \to S$ restricts to one with the specified classes on H_{D_*} exactly when its δ-boundary value belongs to $T_{C,\delta}$. This uses the actual covering identity that a boundary generator upstairs is conjugate, inside H_C, to the r_γth power of the generator downstairs. No condition on an unspecified path value remains because C_γ is a conjugacy class. The transported subset for H'_C is

$$T'_{C,\delta'} = T^{\lambda}_{C,\sigma_{i,C}(\delta')}. \tag{D.29.11}$$

The unit-power map on a finite group is a bijection and commutes with every power r_γ, which proves this equality even when that latter power map is not injective.

For A_5 all the powers in (D.29.11) are realized by one actual target automorphism. Namely let τ_λ be the identity for $\lambda \equiv 1, 4 \pmod 5$ and conjugation by (01) in S_5 for $\lambda \equiv 2, 3 \pmod 5$. Then

$$\tau_\lambda(T) = T^\lambda \quad \text{for every conjugacy-invariant } T \subseteq A_5. \tag{D.29.12}$$

The identity, double-transposition and three-cycle classes are each preserved by every unit power. The two five-cycle classes are exchanged precisely by the nonsquare residues modulo five, as they are by the indicated odd conjugation. All eight units modulo $30 = \exp(A_5)$ are checked in the finite certificate below.

The two covers in (D.29.8) have the same degree and the same boundary cycle lengths under (D.29.9), hence the same genus and number of boundary components. Their boundary permutation is realized by an explicit surface-group isomorphism. In the presentation (D.28.7), the adjacent boundary transformation

$$(c_j, c_{j+1}) \longmapsto (c_{j+1}, c_{j+1}^{-1} c_j c_{j+1}) \tag{D.29.13}$$

fixes all handles and other boundaries and preserves the boundary product literally. Its inverse sends (c_j, c_{j+1}) to $(c_j c_{j+1} c_j^{-1}, c_j)$. These actual group automorphisms realize every permutation of the oriented boundary conjugacy classes; the same formulas give the inverse maps on the profinite completions. Choosing matching surface presentations and composing these maps yields an isomorphism $u_C : H'_C \to H_C$ with the boundary correspondence (D.29.9). Thus

$$\psi \longmapsto \tau_\lambda \circ \psi \circ u_C \tag{D.29.14}$$

is an actual bijection between the epimorphisms $H_C \twoheadrightarrow S$ with the conditions (D.29.10) and those on H'_C with the conditions (D.29.11). Let their common finite cardinality be N_C. These isomorphisms are chosen separately for each C; compatibility among them is not assumed or needed for the count below.

We now impose full generation of the simple-factor kernel. In the original case put $K = K_0$; the following argument is repeated for the new case with $K = K_1$ and H_C replaced by H'_C. For any epimorphism $\varphi : H_{D_*} \twoheadrightarrow S$, its restriction to K is onto. Indeed $\varphi(K)$ is normal in the simple group S, so it is either 1 or S. The former alternative would make the even-order group S a quotient of the odd-order group H_{D_*}/K, which is impossible. Put $N = \ker(\varphi|_K)$. For $D_* \le C \le D$, the condition $H_C \le N_{\widehat{F}}(N)$ is equivalent to extension of φ to H_C, and the extension is unique. Indeed conjugation on $K/N \simeq S$ gives $H_C \to \mathrm{Aut}(S)$; its outer part factors through the odd-order group C and is trivial because $\mathrm{Out}(S) = C_2$. The inverse of the injective map $S \xrightarrow{\mathrm{Inn}} \mathrm{Inn}(S)$ is the required homomorphism to S. It agrees with φ on H_{D_*} by centerlessness, and is uniquely determined by conjugation on the surjective image of K.

Conversely an epimorphism $H_C \to S$ restricts surjectively to H_{D_*}, because its restriction to K is already onto by the same odd-order quotient argument. Its boundary conditions on that restriction are exactly (D.29.10). Hence N_C is also the number of the prescribed

epimorphisms from H_{D_*} whose coordinate-kernel stabilizer contains H_C. Let e_B count those with stabilizer exactly H_B, for $D_* \le B \le D$. On this finite subgroup poset we have

$$N_C = \sum_{B \ge C} e_B, \qquad e_{D_*} = \sum_{C \ge D_*} \mu(D_*, C) N_C, \tag{D.29.15}$$

where μ is the inverse of the incidence matrix of inclusion. Formula (D.29.14) proves equality of every N_C for the original and the new problem, so (D.29.15) proves equality of their e_{D_*} as well.

The original source epimorphism q_E gives $e_{D_*} > 0$. In fact the explicit nonabelian Shapiro map of (D.28.13) identifies the coordinate maps on K with $k \mapsto \varphi(t_\omega k t_\omega^{-1})$. Their kernels are pairwise distinct precisely when the stabilizer of N is H_{D_*}. For the original epimorphism their joint image is all of S^Ω, so these kernels are distinct. Conversely, for the new problem distinct kernels and surjectivity of the coordinate maps force the full product by the subdirect argument of D.25. The Shapiro formula therefore turns each new map counted by e_{D_*} into an epimorphism in (D.29.3), including the prescribed h_i.

The literal count can also be retained exactly. For fixed representatives t_ω, every homomorphism to $S^\Omega \rtimes D$ over the base map has a unique kernel conjugation by $v \in S^\Omega$ with $v(\omega_0) = 1$ that makes its cocycle satisfy $c(t_\omega)(\omega_0) = 1$. Explicitly take $v(\omega) = c(t_\omega)(\omega_0)$ and use

$$c^v(g)(\omega) = v(\omega) c(g)(\omega) v(\omega q(g))^{-1}. \tag{D.29.16}$$

Here q denotes the base map in the case being counted; in the new case it is replaced by q_1. The resulting cocycle is exactly the Shapiro cocycle of $\varphi = c|_{H_{D_*}}(\omega_0)$. Thus each surface map has exactly 60^{n-1} literal cocycles. Both generation and the specified kernel-conjugacy conditions are preserved by these conjugations, so the literal epimorphism count is

$$60^{n-1} \sum_{C \ge D_*} \mu(D_*, C) N_C. \tag{D.29.17}$$

Its proved invariance gives the asserted exact count, as well as existence.

For the coupled assertion, choose the specified L-conjugators of its new peripheral epimorphism and use their images in D as the h_i in (D.29.2). The lift just constructed pairs with $q_{L,1}$, and the peripheral conjugators are the literal pairs of the chosen lifted conjugators. For surjectivity, let K_L be the kernel of the new map from $\widehat{F}$ to L. Its image in A is normal in A, and the quotient of A by this image is a quotient of the odd-order group $\ker(L \to D)$. Every normal subgroup of a direct power of the nonabelian simple group S is a product of its simple factors: commuting such a subgroup with each factor first includes every factor on which its projection is nontrivial. Thus every nontrivial quotient of A has a factor of order 60 and has even order. The quotient in question is therefore trivial, which proves surjectivity of the paired map. When L is an actual ℓ-group, its coordinate projection is the maximal pro-ℓ quotient: the perfect kernel A has no nontrivial finite ℓ-group quotient. These arguments use neither a solvability theorem for odd-order groups nor an inference from perfectness to absence of all odd-order quotients.

The program `a_peripheral_a5_invariance.py` checks the actual outer involution of A_5, all eight unit powers modulo thirty, all thirty-two conjugacy-invariant subsets, and their inverse images under every power modulo thirty. It verifies (D.29.13) and its inverse as literal free-word identities, and checks the averaging identity on a basis of the outer permutation module. It also treats the degree-five regular cover with $x \mapsto t, y \mapsto 1$, which has seven boundaries and genus zero. For a generating triple of class orders $(3, 5, 5)$ it checks all eight multipliers and all five possible y-conjugator cosets. In every one of the forty cases there are sixty proper surface maps; the corresponding literal count is 60^5. The

script evaluates their wreath cycle products and the coordinate Schreier maps, verifying surjectivity and distinct kernels. The code, executed output and JSON certificate end with `PASS_ALL_DEGREE_A5_COUNT_COMPARISONS`. □

D.29 proves the complete A_5-chief comparison over finite odd-order bases. It does not iterate automatically over an arbitrary mixed finite base: after adjoining the first A_5 factor, the base need no longer have odd order. The specific uses of that hypothesis are the outer-cocycle averaging in (D.29.5) and the unique inner extension in (D.29.15). Those exact maps identify the step needed to extend this particular proof. D.30 supplies that extension over every finite base by the actual twisted S_5 model, its character count, and its finite-base subtraction. The full cofinal problem in D.1 and D.2 still includes the other chief factors.

4.13 Twisted A_5 chief extensions over arbitrary finite bases

Statement D.30 (Proved). Exact peripheral count invariance for every finite A_5 chief extension. *Let*

$$1 \longrightarrow A \longrightarrow E \xrightarrow{\pi} D \longrightarrow 1, \qquad A \simeq A_5^{\Omega}, \tag{D.30.1}$$

be a finite extension in which A is a nonabelian chief factor. Let $q_E : \widehat{F} \twoheadrightarrow E$ be an epimorphism, put $X_i = q_E(x_i)$, $q_0 = \pi q_E$, and $d_i = q_0(x_i)$. Let $q_1 : \widehat{F} \twoheadrightarrow D$ be any epimorphism, and suppose its three specified peripheral values are

$$q_1(x_i) = h_i d_i^{\lambda} h_i^{-1}, \qquad h_i \in D, \quad \lambda \in \widehat{\mathbb{Z}}^{\times}. \tag{D.30.2}$$

For arbitrary chosen lifts $H_i \in E$ of h_i, there is an epimorphism $f : \widehat{F} \twoheadrightarrow E$ with

$$\pi f = q_1, \qquad f(x_i) = a_i H_i X_i^{\lambda} H_i^{-1} a_i^{-1} \quad (a_i \in A). \tag{D.30.3}$$

The number of these literal epimorphisms equals the number for $(q_1, \lambda, h_1, h_2, h_3) = (q_0, 1, 1, 1, 1)$. In particular it is positive. There is no restriction on the order of D, the degree of its factor action, its fixed peripheral points, or its outer action on A. The result applies at every successive A_5 chief step of a finite tower with its given original source epimorphism and its prescribed new peripheral base epimorphism.

Proof. Write $S = A_5$, $\widetilde{S} = S_5$, let $\sigma = (01) \in \widetilde{S}$, and write $\tau(s) = \sigma s \sigma^{-1}$. The exact finite calculation in D.28 identifies $\widetilde{S} \xrightarrow{\mathrm{Ad}} \mathrm{Aut}(S)$ as an isomorphism and $\mathrm{Out}(S)$ with $\mathbf{F}_2$ by parity. In particular $C_{\widetilde{S}}(S) = 1$.

The actual twisted permutation extension. The factor action of D on Ω is transitive because the kernel is chief. Choose a factor ω_0, let D_* be its stabilizer, and identify $\Omega = D_* \backslash D$ with the right action. Choose representatives $t_\omega \in D$ with $t_{\omega_0} = 1$. The outer action of D is a homomorphism to $\mathbf{F}_2^{\Omega} \rtimes \mathrm{Sym}(\Omega)$, so its component b satisfies

$$b_{de}(\omega) = b_d(\omega) + b_e(\omega d). \tag{D.30.4}$$

Set

$$\epsilon(c) = b_c(\omega_0) \quad (c \in D_*), \qquad s_\omega = b_{t_\omega}(\omega_0), \qquad k(\omega, d) = t_\omega d t_{\omega d}^{-1}. \tag{D.30.5}$$

Then $\epsilon : D_* \to \mathbf{F}_2$ is a character, $k(\omega, d) \in D_*$, and the cocycle identity gives

$$b_d(\omega) = s_\omega + \epsilon(k(\omega, d)) + s_{\omega d}. \tag{D.30.6}$$

This formula supplies the required change of coordinates, without averaging. More explicitly, centerlessness gives the isomorphism

$$E \longrightarrow \mathrm{Aut}(A) \times_{\mathrm{Out}(A)} D, \qquad e \longmapsto (\mathrm{Ad}_e, \pi(e)) : \tag{D.30.7}$$

injectivity and surjectivity follow exactly by the kernel and inner-automorphism calculation in (D.29.6). Represent each factor automorphism by its unique element of $\widetilde{S}$. Conjugate the resulting wreath coordinates by the vector $(\sigma^{s_\omega})_\omega$. In coordinates this sends (z, d) to $(\sigma^{-s_\omega} z(\omega) \sigma^{s_{\omega d}}, d)$. Equations (D.30.6)–(D.30.7) identify the given extension, by this actual isomorphism, with

$$\begin{aligned} E_* &= \{(z, d) \in \widetilde{S}^\Omega \rtimes D : \operatorname{sgn} z(\omega) = \epsilon(k(\omega, d))\}, \\ (z, d)(w, e) &= (z(\omega) w(\omega d), de). \end{aligned} \tag{D.30.8}$$

Its kernel is precisely S^Ω. All the X_i, H_i are transported by the displayed isomorphism; these transported elements are used below.

For either source epimorphism $q = q_j$, put $K_j = \ker q_j$ and $H_{j,C} = q_j^{-1}(C)$ for $D_* \le C \le D$. Choose actual words $T_\omega \in \widehat{F}$ with $q(T_\omega) = t_\omega$ and $T_{\omega_0} = 1$. The twisted nonabelian Shapiro map is the literal formula

$$\begin{aligned} \varphi : H_{j,D_*} &\longrightarrow \widetilde{S}, \quad \operatorname{sgn} \varphi = \epsilon q_j, \\ z_\varphi(g)(\omega) &= \varphi(T_\omega g T^{-1}_{\omega q_j(g)}), \qquad f_\varphi(g) = (z_\varphi(g), q_j(g)). \end{aligned} \tag{D.30.9}$$

The arguments of φ lie in H_{j,D_*}; multiplication and the parity condition in (D.30.8) follow by cancellation of the middle representative. Conversely, evaluate a cocycle at ω_0 on this subgroup. Conjugating by the unique vector $v \in S^\Omega$ with $v(\omega_0) = 1$ and $v(\omega) = z(T_\omega)(\omega_0)$ makes it exactly (D.30.9). The entries of v are even: the transition element for (ω_0, t_ω) in (D.30.5) is 1. Thus there are exactly $60^{|\Omega|-1}$ literal cocycles for each normalized surface map.

All intermediate extensions and their exact embeddings. For $D_* \le C \le D$, consider each character $\epsilon_C : C \to \mathbf{F}_2$ extending ϵ. There may be none. Choose representatives R_ν for $\nu \in C\backslash D$, with the identity at the base coset, and define

$$k_C(\nu, d) = R_\nu d R_{\nu d}^{-1}, \qquad b_C(d, \nu) = \epsilon_C(k_C(\nu, d)). \tag{D.30.10}$$

Let E_C be the analogue of (D.30.8) with these data. Write $p(\omega) = C t_\omega$ and

$$g_\omega = \epsilon_C(t_\omega R_{p(\omega)}^{-1}). \tag{D.30.11}$$

The argument belongs to C. The following is an injective homomorphism over D:

$$j_C : E_C \longrightarrow E_*, \qquad j_C(z, d)(\omega) = \sigma^{g_\omega} z(p(\omega)) \sigma^{-g_{\omega d}}. \tag{D.30.12}$$

The base coordinate d is retained. Multiplication is verified by cancelling the two adjacent powers of σ; injectivity follows by reading one coordinate above every ν. Its parity is correct because

$$\epsilon(k(\omega, d)) = g_\omega + b_C(d, p(\omega)) + g_{\omega d}. \tag{D.30.13}$$

This proves the embedding with its outer twisting. At the base coordinate $g_{\omega_0} = 0$; consequently the Shapiro map of a homomorphism $H_{j,C} \to \widetilde{S}$, followed by j_C, restricts at H_{j,D_*} to that very homomorphism's restriction.

The boundary constraint sets. Fix a peripheral i. An element of E_* over $q_j(x_i)$ is conjugate under $A = S^\Omega$ to another such element exactly when all their cycle products are conjugate under S. Indeed the usual successive conjugator equation along a cycle determines every conjugator after its first value; all these values are even because the two component parity patterns agree. The last equation is exactly the stated S-conjugacy of the cycle product.

This gives finite subsets $T_{j,C,\delta}$ of $\widetilde{S}$, one for every boundary component δ of $H_{j,C}$, that express the condition on $j_C f_\varphi(x_i)$ literally. Here is an explicit construction. Let δ be a cycle on $C\backslash D$, choose its initial coordinate ν, and choose the initial coordinate ω of each cycle γ above it so that $p(\omega) = \nu$. If their lengths are e and re, respectively, the cycle product under (D.30.12) is $\sigma^{g_\omega} a^r \sigma^{-g_\omega}$, where a is the product at δ. Intersect the conditions

$$\sigma^{g_\omega} a^r \sigma^{-g_\omega} \in \mathcal{C}_{j,\gamma} \tag{D.30.14}$$

over all such γ, and impose the parity $\operatorname{sgn} a = \epsilon_C(q_j(c_\delta))$. Here $\mathcal{C}_{0,\gamma}$ is the actual S-class of the cycle product of X_i, and $\mathcal{C}_{1,\gamma}$ comes from $H_i X_i^\lambda H_i^{-1}$. The result is $T_{j,C,\delta}$. It is a union of S-conjugacy classes, all of its prescribed parity. Formula (D.30.14) is also valid when the corresponding subset is empty.

The comparison of these sets retains the prescribed h_i. Match the boundary cycles by

$$Cd\langle q_1(x_i)\rangle \longmapsto Cdh_i\langle d_i\rangle. \tag{D.30.15}$$

It preserves lengths and every relative boundary degree, by (D.30.2). Choose an initial coordinate ν' of the new cycle and use $\nu' h_i$ as the initial coordinate of its old match. Let $L_i \in E_C$ be the canonical element over h_i, with components $\sigma^{b_C(h_i,\nu)}$. Its image under j_C differs from H_i by an element of A. Therefore conjugating a local element by L_i after taking its λth power transports precisely the old allowable set to the new one. Its cycle product is conjugated by the actual component $\sigma^{b_C(h_i,\nu')}$ of L_i. Consequently

$$T_{1,C,\delta'} = \tau^{b_C(h_i,\nu')}(T_{0,C,\delta}^\lambda). \tag{D.30.16}$$

The inverse power and inverse conjugation prove the reverse inclusion, as λ is a unit on every finite group involved. This argument does not require X_i itself to belong to $j_C(E_C)$.

Actual surface maps with a specified parity character. The covers $H_{0,C}$ and $H_{1,C}$ have the same index, genus, and boundary count, and (D.30.15) matches their boundary parities for the characters $\chi_j = \epsilon_C q_j$. Both characters are trivial when $\epsilon_C = 0$; both are nontrivial otherwise, since q_j maps $H_{j,C}$ onto C. There is an isomorphism of their surface groups, and hence of their profinite completions, that matches these characters and the indicated oriented boundary classes. The following gives actual homeomorphisms realizing the comparison. First use the adjacent Hurwitz maps (D.29.13) for the boundary permutation. If all boundary character values vanish, the remaining nonzero characters are covectors on the handle symplectic space over $\mathbf{F}_2$. A Dehn twist around a simple curve of class v acts by the transvection $w \mapsto w + \langle w, v\rangle v$. Every nonzero mod-two handle class has such a curve: in each handle take the indicated one of the three slopes $(1,0), (0,1), (1,1)$, and join the selected disjoint curves by bands away from the boundary. These transvections are transitive on nonzero vectors. Explicitly, for distinct a, b with $\langle a, b\rangle = 1$ use the transvection at $a+b$; when this pairing is zero choose c pairing to one with both and use the two transvections through c. This gives a product of boundary-fixed Dehn twists matching the two nonzero covectors.

If a boundary class d has character value one, take two disjoint curves of classes a and $a + d$ bounding a pair of pants with that boundary. The bounding-pair map $T_{a+d} T_a^{-1}$ fixes the boundary and acts on homology by

$$w \longmapsto w + \langle w, a\rangle d. \tag{D.30.17}$$

Thus it adds the handle covector $\langle -, a\rangle$ to the character. Choosing the handle basis curves supplies every required translation. For genus zero the boundary values already determine the character. Base these homeomorphisms outside their supports. They induce actual inverse maps on the surface groups and their profinite completions. A boundary generator may be conjugated by a based path, whose parity is specified by χ_j; this can apply one further power of τ to its target class. This is the only additional boundary change in this comparison.

We now prove the exact target-count fact that accounts for all those possible changes. For a fixed nontrivial character χ of a surface group, the number of epimorphisms to $\widetilde{S}$ with parity χ and boundary values in specified S-invariant subsets is unchanged by applying τ independently to any of the boundary subsets.

Here is an integral finite proof. In the group algebra of $\widetilde{S}$, write $[T] = \sum_{t\in T} t$ and define

$$K_{ab} = \sum_{\substack{u,v\in\widetilde{S}\\ \operatorname{sgn} u=a,\operatorname{sgn} v=b}} [u, v], \qquad a, b \in \mathbf{F}_2. \tag{D.30.18}$$

Here each group element in the sum is its basis element, so multiplicities are retained. Let $\Delta = [5A] - [5B]$, using the two S-classes of five-cycles. Exact multiplication of the 120 permutations gives

$$\begin{gathered} K_{01} = K_{10} = K_{11} = L, \\ L\Delta = 0, \qquad L(K_{00} - L) = 0, \\ [O]\Delta = 0, \qquad [O](K_{00} - L) = 0 \quad \text{for each odd } S\text{-class } O. \end{gathered} \tag{D.30.19}$$

The coefficients of K_{00} and L on the even classes, per element, are

class	1	2^2	3	$5A$	$5B$
K_{00}	300	32	63	65	65
L	180	72	63	45	45.

(D.30.20)

They vanish on odd permutations. All eight S-class sums in $\widetilde{S}$ commute. The complete integer coefficient vectors and all products in (D.30.19) are checked by the exhaustive permutation certificate specified below.

For genus g and boundaries $c_1, \dots, c_b$, the number of all homomorphisms with character χ and boundary subsets T_i is exactly the coefficient of 1 in

$$\prod_{j=1}^{g} K_{\chi(a_j),\chi(b_j)} \prod_{i=1}^{b} [T_i]. \tag{D.30.21}$$

This is the tuple count for the literal surface relation $\prod[a_j, b_j] \prod c_i = 1$. Changing one even boundary subset by τ changes its class sum by 0, Δ, or $-\Delta$; odd classes are unchanged. Nontriviality of χ provides either a factor L or another odd boundary factor. Thus (D.30.19) makes the difference in (D.30.21) zero.

The same conclusion holds after excluding nonsurjective maps. The exhaustive subgroup lattice of S_5 has 156 subgroups. Exactly six proper subgroups contain both an odd permutation and a five-cycle, and each is the normalizer $F_{20} = C_5 \rtimes C_4$ of its unique subgroup of order five. In the difference from one changed boundary, every contributing nonsurjective map has precisely one of these six subgroups as its image. It has a five-cycle because the changed factor is supported there, and an odd image because χ is nontrivial. For each embedded F_{20} put $R = \sum_{c\in C_5} c$. Its corresponding restricted distributions are

$$K_{00}^{F_{20}} = 25 \cdot 1 + 15R, \qquad K_{01}^{F_{20}} = K_{10}^{F_{20}} = K_{11}^{F_{20}} = 20R. \tag{D.30.22}$$

The restriction of Δ to C_5 has augmentation zero. Every nonempty odd intersection of an S-class with F_{20} is the union of its two odd C_5-cosets. Consequently both $20R$ and every such odd class sum annihilate that restricted Δ. The relevant restricted class sums commute; this also follows immediately by multiplying in $C_5 \rtimes C_4$ and is checked for all six embeddings. The difference of the homomorphism counts into each F_{20} is therefore zero by the same formula (D.30.21). Every map in this difference is already onto that F_{20}, by the proved proper subgroup classification. There is no overcount between the six image groups. Subtracting their zero differences proves the asserted equality of epimorphism counts to S_5.

Finally every unit power on the eight S-classes in S_5 is realized by the actual automorphism τ_λ: it is the identity for $\lambda \equiv 1, 4 \pmod 5$ and τ for $\lambda \equiv 2, 3 \pmod 5$. The three odd classes of orders $2, 4, 6$ are each preserved by all unit powers; the even classes have the transformation in (D.29.12). Thus the surface isomorphism above, τ_λ, and the independent-flip count prove equality of the following two finite numbers, for each C, ϵ_C:

$$P_{j,C,\epsilon_C} = \#\{\varphi : H_{j,C} \twoheadrightarrow G_C : \operatorname{sgn}\varphi = \epsilon_C q_j,\ \varphi(c_\delta) \in T_{j,C,\delta}\}, \quad G_C = \begin{cases} A_5 & \epsilon_C = 0, \\ S_5 & \epsilon_C \neq 0. \end{cases} \tag{D.30.23}$$

For $\epsilon_C = 0$, the character and all flips in (D.30.16) and in the based surface comparison are trivial; the actual target automorphism and source isomorphism give the equality directly. For $\epsilon_C \neq 0$, the just-proved fixed-character count applies. No claim that the independent class flips themselves are group automorphisms is needed.

The exact finite-base subtraction. An epimorphism counted in (D.30.23) has $\varphi(K_j) = 1$ or A_5. Indeed this image is contained in A_5 by the parity condition and is normal in G_C, hence normal in the simple group A_5. The first alternative consists precisely of maps

$$\varphi = \Phi q_j|_{H_{j,C}}, \qquad \Phi : C \twoheadrightarrow G_C, \operatorname{sgn}\Phi = \epsilon_C. \tag{D.30.24}$$

Their admissible counts on the two sides agree by an actual finite-group comparison. For such a Φ, form the induced section

$$s_\Phi : D \longrightarrow E_C, \qquad s_\Phi(d)(\nu) = \Phi(R_\nu d R_{\nu d}^{-1}), \tag{D.30.25}$$

with base coordinate d. It is a homomorphism by cancellation of representatives, and its parity is (D.30.10). The two maps in (D.30.24) give exactly the Shapiro maps $s_\Phi q_0$ and $s_\Phi q_1$. Their compositions with j_C have the literal equality

$$j_C s_\Phi(q_1(x_i)) = j_C s_\Phi(h_i)\, j_C s_\Phi(d_i)^\lambda\, j_C s_\Phi(h_i)^{-1}. \tag{D.30.26}$$

The element $j_C s_\Phi(h_i)$ and the prescribed H_i have the same base coordinate; their difference is in A. It follows in both directions that the same Φ is admissible for $j = 0$ and $j = 1$. The inverse unit power proves the reverse implication. This is a bijection of the maps that factor through the finite base; it also retains their surjectivity and prescribed character.

Subtract this common number from (D.30.23). We obtain equal numbers N_{j,C,ϵ_C} of maps with the same boundary conditions and $\varphi(K_j) = A_5$. Put

$$N_{j,C} = \sum_{\epsilon_C|_{D_*} = \epsilon} N_{j,C,\epsilon_C}. \tag{D.30.27}$$

The empty sum is zero. We have proved $N_{0,C} = N_{1,C}$ for every C, without assuming that a finite-base quotient of A_5 is impossible.

Normalizer, generation, and literal counts. Fix j and a map $\varphi : H_{j,D_*} \to S_5$ of prescribed character and boundary classes with $\varphi(K_j) = A_5$. Let $N = \ker(\varphi|_{K_j})$. Its normalizer in $\widehat{F}$ contains H_{j,D_*} and K_j, so it equals $H_{j,B}$ for a unique $B \geq D_*$. For $C \geq D_*$, the condition that $H_{j,C}$ normalize N is equivalent to extension of φ to a map $H_{j,C} \to S_5$ with image containing A_5. The extension is unique. For existence, conjugation on $K_j/N \xrightarrow[\varphi]{\sim} A_5$ gives the actual homomorphism $H_{j,C} \to \mathrm{Aut}(A_5) = S_5$. Its restriction agrees with φ because $C_{S_5}(A_5) = 1$. Its parity factors through a unique character $\epsilon_C : C \to C_2$ extending ϵ. Conversely every extension with K_j-image A_5 normalizes N, and centerlessness gives uniqueness. Thus each map whose normalizer contains $H_{j,C}$ is counted exactly once in (D.30.27).

Let $e_{j,B}$ count those whose normalizer is exactly $H_{j,B}$. On the same finite subgroup poset on both sides,

$$N_{j,C} = \sum_{B \geq C} e_{j,B}, \qquad e_{j,D_*} = \sum_{C \geq D_*} \mu(D_*, C) N_{j,C}. \tag{D.30.28}$$

The already proved equalities of $N_{j,C}$ therefore give $e_{0,D_*} = e_{1,D_*}$. Under (D.30.9), the coordinate maps on K_j are $k \mapsto \varphi(T_\omega k T_\omega^{-1})$. They are onto A_5, and their kernels are pairwise distinct exactly when the normalizer is H_{j,D_*}. The simple-factor subdirect argument of D.25 then gives the whole product A_5^Ω. Together with the epimorphism to D this is precisely surjectivity onto E_*. The original epimorphism q_E shows $e_{0,D_*} > 0$. Finally the unique even gauges in (D.30.9) show that the literal proper-lift count on side j is

$$60^{|\Omega|-1} \sum_{C \geq D_*} \mu(D_*, C) N_{j,C}. \tag{D.30.29}$$

The gauges preserve the required A-conjugacy classes, so no further quotient or centralizer factor is implicit in this formula. Its equality and positivity prove (D.30.3).

Finite certificates and the scope of iteration. The script `a_peripheral_s5_twist.py` constructs all 120 permutations, enumerates all 156 subgroups, and verifies every integer coefficient in (D.30.19)–(D.30.22). It verifies all sixteen units modulo 60 on the eight A_5-classes in S_5. Its output and JSON certificate end with `PASS_TWISTED_S5_CLASS_ALGEBRA`. The independent map script `a_peripheral_s5_maps.py` uses the actual bases $D = S_3, S_4$, the stabilizer generated by a transposition, and every extending character on every intermediate subgroup. It verifies 4680 literal multiplication comparisons for (D.30.12) and 1473 nonempty boundary comparisons for (D.30.16), with all permutation coordinates retained. Its output and JSON end with `PASS_TWISTED_WREATH_EXACT_MAPS`. The general maps and count argument above prove the statement beyond those finite test bases.

At a successive A_5 chief step, use the epimorphism already obtained at the previous base as q_1; its peripheral values have exactly the required form. The given original source epimorphism at that step supplies q_E. This proves the asserted finite iteration. It makes no assertion that towers with these chief factors are cofinal among all finite quotients, and it does not remove the central obstruction in D.24 or other chief factors in D.1–D.2. □

4.14 An A_6 chief obstruction with prescribed base conjugators

Statement D.31 (Proved). A proper-lift obstruction and its complete forgetful fibre. *There is an actual chief extension*

$$1 \longrightarrow \mathrm{Inn}(A_6) \longrightarrow \mathrm{Aut}(A_6) \xrightarrow{\pi} \mathbf{F}_2^2 \longrightarrow 1 \tag{D.31.1}$$

and a generating peripheral triple (X, Y, Z) *of orders* $(2, 8, 8)$ *such that the following holds. Put* $d_1 = \pi(X)$*,* $d_2 = \pi(Y)$*, and* $d_3 = d_1 + d_2$*. With the same base epimorphism* $q_1 = q_0$*, multiplier* $\lambda = 1$*, and prescribed base conjugators*

$$(h_1, h_2, h_3) = (0, 0, d_1), \tag{D.31.2}$$

there are exactly ninety weak lifts and no epimorphic lift. Every weak lift has image isomorphic to

$$C_8 \rtimes_5 C_2 = \langle r, s : r^8 = s^2 = 1,\ srs^{-1} = r^5 \rangle. \tag{D.31.3}$$

The original choice $(h_1, h_2, h_3) = (0, 0, 0)$ *has exactly 360 lifts, all epimorphic.*

More precisely, write $h_i = p_i d_1 + q_i d_2$*, with* $p_i, q_i \in \mathbf{F}_2$*. The complete prescribed-conjugator count is*

$q_1 + p_2 + p_3 + q_3$	*weak lifts*	*epimorphic lifts*
0	360	360
1	90	0.

(D.31.4)

Thus each row occurs for 32 of the 64 possible base-centralizer triples. Forgetting the prescribed base conjugators gives a union of 1800 weak lifts and 1440 epimorphic lifts. These form exactly two simultaneous $\mathrm{Aut}(A_6)$*-conjugacy orbits, of sizes 1440 and 360, respectively. The first is the orbit of the original generating triple. In particular the ordinary peripheral lift fibre is nonempty.*

Both groups in (D.31.1) *have trivial maximal pro-five quotient. For every finite five-group* L *and every specified multiplier-one peripheral automorphism* β *of its given two-generated source quotient, the actual products of* (D.31.1) *with* L *have maximal pro-five quotient* L*. Over the same base shadow* $1_{\mathbf{F}_2^2} \times \beta$ *they have both the obstructed prescribed-conjugator branch above and an explicit admitted branch.*

Proof. All permutations below act on $\{0, 1, 2, 3, 4, 5\}$. In $S = A_6$, take

$$a = (0123)(45), \qquad b = (01234). \tag{D.31.5}$$

They generate S. An automorphism is specified by the images of this actual generating pair. The following table defines X, Y, Z; a row $[u_0, \ldots, u_5]$ denotes the permutation $i \mapsto u_i$:

	a image	b image
X	$[1, 0, 4, 2, 5, 3]$	$[4, 1, 0, 2, 5, 3]$
Y	$[1, 2, 3, 0, 5, 4]$	$[5, 1, 0, 4, 2, 3]$
Z	$[1, 4, 5, 0, 3, 2]$	$[0, 3, 4, 2, 5, 1]$.

(D.31.6)

The exact permutation certificate verifies that these assignments extend to automorphisms, that $XYZ = 1$, and that X, Y generate all 1440 automorphisms. It enumerates every automorphism, rather than taking an order formula as input: any automorphism must send a, b to elements of orders $4, 5$ with the same product and commutator orders. For each such pair it propagates the proposed map over every edge of the Cayley graph of the actual 360-element group. Consistency on these edges and bijectivity are necessary and sufficient for an automorphism. This exhaustive calculation gives 1440 maps. Composition is computed from their two generator images. The inner maps form the actual 360-element normal subgroup, and its computed quotient is $\mathbf{F}_2^2$, with d_1, d_2 as the indicated basis. The same certificate computes all seven conjugacy classes of A_6 and the normal closure of each. Every nonidentity class normally generates all 360 elements. This proves simplicity, so the displayed kernel is indeed chief. A nontrivial commutator normally generates it as well, which proves perfectness.

Let $\mathcal{X}_0, \mathcal{Y}_0, \mathcal{Z}_0$ be the conjugacy classes of X, Y, Z under $A = \mathrm{Inn}(S)$. Their sizes are $15, 90, 90$. Put

$$\mathcal{X}_1 = Y\mathcal{X}_0Y^{-1}, \qquad \mathcal{Y}_1 = X\mathcal{Y}_0X^{-1}, \qquad \mathcal{Z}_1 = X\mathcal{Z}_0X^{-1}. \tag{D.31.7}$$

For each letter, the two displayed classes are distinct and of equal size. Conjugation by an element above $pd_1 + qd_2$ changes their indices as follows:

$$\mathcal{X}_i \longmapsto \mathcal{X}_{i+q}, \qquad \mathcal{Y}_i \longmapsto \mathcal{Y}_{i+p}, \qquad \mathcal{Z}_i \longmapsto \mathcal{Z}_{i+p+q}. \tag{D.31.8}$$

These are actions of the actual outer quotient on the actual inner classes. They are verified by conjugation of the saved automorphisms, and do not replace that action by an abstract isomorphic group.

The tuple map $q_E : \widehat{F} \to E = \mathrm{Aut}(S)$, $x_1 \mapsto X$, $x_2 \mapsto Y$, $x_3 \mapsto Z$, is an epimorphism. For the base conjugators (D.31.2), take the actual lifts $(1, 1, X)$. The required images belong to

$$\mathcal{X}_0, \quad \mathcal{Y}_0, \quad \mathcal{Z}_1, \tag{D.31.9}$$

respectively. Their base values are still d_1, d_2, d_3, because the quotient is abelian. Every other choice of lifts of (D.31.2) gives the same three A-conjugacy classes. The complete enumeration of $u \in \mathcal{X}_0$, $v \in \mathcal{Y}_0$ with $(uv)^{-1} \in \mathcal{Z}_1$ gives exactly ninety triples. They are one simultaneous A-conjugacy orbit. For one representative $(s, r, (sr)^{-1})$, the literal products give $|r| = 8$, $|s| = 2$, and $srs^{-1} = r^5$; its generated subgroup has sixteen elements and its intersection with A is $\langle r^2 \rangle$ of order four. These equalities prove (D.31.3) for every member of the orbit.

There is also a direct count proof of nonexistence of a proper lift. The centralizer of A in $\mathrm{Aut}(A)$ is trivial: an automorphism commuting with every inner automorphism fixes every element because $Z(A) = 1$. Thus a generating triple in E has trivial simultaneous centralizer in A. Its A-conjugacy orbit has 360 elements and remains inside the required three classes. It cannot fit inside the ninety triples in (D.31.9). The count with $\mathcal{Z}_0$ is 360, and its triples are exactly the A-conjugates of (X, Y, Z); all of them generate E.

We retain every base-centralizer choice. For arbitrary h_i the profile indices in (D.31.7) are

$$(q_1,\ p_2,\ p_3 + q_3). \tag{D.31.10}$$

Simultaneous conjugation by an element above $pd_1 + qd_2$ adds $(q, p, p + q)$ to these three indices. These four vectors are exactly the even-parity vectors in $\mathbf{F}_2^3$. They send the original 360-element orbit to every even-parity profile, and send the ninety-element orbit to every odd-parity profile. This proves (D.31.4) from the two exact counts above. Each of the eight profiles has eight preimages in the 64 conjugator triples.

The forgetful map is the literal set map

$$\coprod_{(h_1,h_2,h_3)\in D^3} \mathrm{Lift}(h_1, h_2, h_3) \longrightarrow \{(U_1, U_2, U_3) : U_1U_2U_3 = 1,\ U_i \in X_i^E,\ \pi(U_i) = d_i\}, \tag{D.31.11}$$

which discards the conjugator labels and retains the triple. It is onto: any E-conjugator has some base image h_i, and changing its lift by A is precisely the inner-class freedom already counted. Its image is the union of eight disjoint profiles, not the sum of 64 labelled counts. There are $4 \cdot 360 + 4 \cdot 90 = 1800$ weak triples and $4 \cdot 360 = 1440$ generating triples. The simultaneous E-orbits are exactly the 1440 conjugates of the generating source and the 360 conjugates of the order-sixteen source. Their centralizers have orders one and four, respectively. In particular (D.31.11) retains the original epimorphism itself.

For the pro-five comparison, every homomorphism from E to a finite five-group kills its perfect normal subgroup A. It then factors through D of order four and is trivial. The same holds for D. Thus their actual universal maximal pro-five quotients are trivial. Given a finite five-group source $\widehat{F} \twoheadrightarrow L$, write its peripheral values as l_i and suppose $\beta(l_i) = k_i l_i k_i^{-1}$. The paired original source into $E \times L$ is onto: a subdirect subgroup would otherwise give a nontrivial common quotient of E and L, contrary to the just-proved maximal-pro-five calculation. The projection

$$E \times L \longrightarrow D \times L \tag{D.31.12}$$

has the simple kernel $A \times 1$. Take new base values $(d_i, \beta(l_i))$ and prescribed base conjugators $(0, k_1), (0, k_2), (d_1, k_3)$. Any epimorphic lift would project to an epimorphic lift of (D.31.9), which is impossible. On the other hand the same base values, with base conjugators $(0, k_1), (0, k_2), (0, k_3)$, have the explicit generating lift $(X_i, \beta(l_i))$. The coordinate projections $E \times L \to L$ and $D \times L \to L$ are the universal maximal pro-five maps, with sections $l \mapsto (1, l)$. Both branches therefore lie over the same specified pro-five shadow.

The two certificate programs in `code` have the common prefix `a_peripheral_a6_` and suffixes `outer.py` and `obstruction.py`. They save the complete automorphism maps, all inner classes, all ninety bad triples, their generated subgroup, and all 64 conjugator cases in matching output and JSON files. The latter independently checks every generator edge of every saved automorphism and verifies both simultaneous-conjugacy orbit equalities. Its final status is `PASS_A6_PROPER_OBSTRUCTION_AND_FORGETFUL_UNION`. An initial, incorrect guess that the bad image was a Sylow subgroup of order 32 is explicitly retained in the files with suffix `failed_sylow32`; the exact order-sixteen calculation above supersedes that guess.

D.31 disproves the extension of D.30's arbitrary-prescribed-conjugator assertion to all nonabelian simple kernels. The forgetful map (D.31.11) proves that this example does not obstruct its underlying ordinary base shadow. No restriction on the rational pro-five leading image in D.1 or D.2 is inferred from it. □

4.15 An ordinary peripheral obstruction at a split A_6 chief step

Statement D.32 (Proved). An exhausted Nielsen family obstructs the ordinary generating lift. *There is a split finite chief extension*

$$1 \longrightarrow A_6 \longrightarrow \mathcal{E} \xrightarrow{\pi} \mathcal{D} \longrightarrow 1, \qquad |\mathcal{D}| = 4 \cdot 360^{12}, \quad |\mathcal{E}| = 4 \cdot 360^{13}, \tag{D.32.1}$$

an actual source epimorphism $q_{\mathcal{E}} : \widehat{F} \twoheadrightarrow \mathcal{E}$, and an order-two automorphism α of $\mathcal{D}$ preserving all three labelled peripheral conjugacy classes, such that no epimorphism $f : \widehat{F} \twoheadrightarrow \mathcal{E}$ satisfies

$$\pi f = \alpha \pi q_{\mathcal{E}}, \qquad f(x_i) \in q_{\mathcal{E}}(x_i)^{\mathcal{E}}. \tag{D.32.2}$$

All base conjugators are allowed in (D.32.2)*. There are exactly 2160 weak lifts; each is the graph of an explicit coordinate projection of $\mathcal{D}$, conjugated in the added A_6 factor, and has image a complement to that factor. For the original base map, the corresponding ordinary peripheral fibre has 3240 weak lifts, of which 1080 are epimorphic and 2160 have complementary image.*

The maximal pro-five quotients of $\mathcal{E}$ and $\mathcal{D}$ are trivial. For every prescribed finite-five-group multiplier-one peripheral automorphism β, taking the actual products with its source quotient gives both an ordinary obstructed base shadow $\alpha \times \beta$ and an explicitly admitted base shadow $1 \times \beta$, with the same induced pro-five shadow β.

Proof. Use the actual group $G = \mathrm{Aut}(A_6)$ of D.31, its inner subgroup $S = \mathrm{Inn}(A_6)$, and its quotient $O = G/S = \mathbf{F}_2^2$. The elements of G below are specified by their images of the same

generators $a = (0123)(45)$ and $b = (01234)$. Write $[i_0 i_1 i_2 i_3 i_4 i_5]$ for the permutation $j \mapsto i_j$. Define $U \in G$ by

$$U(a) = [103452], \qquad U(b) = [023451]. \tag{D.32.3}$$

It has order six. The following table defines fifteen actual automorphisms V_{H_j} and V_{L_j}:

	a image	b image		a image	b image
H_0	[103524]	[325140]	L_0	[123054]	[510423]
H_1	[104532]	[423015]	L_1	[123054]	[542103]
H_2	[130254]	[025413]	L_2	[140523]	[513402]
H_3	[130254]	[352410]	L_3	[150432]	[314250]
H_4	[153204]	[214530]	L_4	[203154]	[524301]
H_5	[230514]	[153042]	L_5	[320154]	[054213]
H_6	[240531]	[134205]			
H_7	[241503]	[054132]			
H_8	[401523]	[035214]			

(D.32.4)

For each label T, also use T to denote the actual homomorphism

$$T : \widehat{F} \twoheadrightarrow G, \qquad (T(x_1), T(x_2), T(x_3)) = (U, V_T, (UV_T)^{-1}). \tag{D.32.5}$$

All fifteen maps are onto, and all have the same outer map

$$\theta : \widehat{F} \twoheadrightarrow O, \qquad (x_1, x_2, x_3) \longmapsto (e_1, e_2, e_1 + e_2). \tag{D.32.6}$$

These assertions are verified by literal automorphism multiplication and enumeration of the generated 1440-element subgroup, using the complete automorphism construction in D.31.

The two complete Nielsen families. Let $\mathcal{U}$ be the S-class of U, let $\mathcal{V}$ be the S-class of V_{H_0}, and put

$$\mathcal{W}_H = ((UV_{H_0})^{-1})^S, \qquad \mathcal{W}_L = U\mathcal{W}_H U^{-1}. \tag{D.32.7}$$

Their sizes are $120, 90, 90, 90$. The classes $\mathcal{W}_H$ and $\mathcal{W}_L$ are distinct. The exact tuple counts and orbit decompositions are

profile	literal triples	S-conjugacy orbits	generated order
$(\mathcal{U}, \mathcal{V}, \mathcal{W}_H)$	3240	9	1440
$(\mathcal{U}, \mathcal{V}, \mathcal{W}_L)$	2160	6	1440.

(D.32.8)

The representatives are precisely the nine H_j and the six L_j in (D.32.4)–(D.32.5). For completeness, the enumeration fixes the first element U and uses its inner centralizer of order three on the remaining admissible second elements. There are 27 admissible second elements for $\mathcal{W}_H$ and eighteen for $\mathcal{W}_L$; their orbits all have size three. Every representative generates G. Conjugating the first element through its 120 values gives the literal counts in (D.32.8). Thus these counts retain properness and the complete orbit decomposition.

Put $K = \ker \theta$. Each of the fifteen maps sends K onto S. Their fifteen kernels on K are pairwise distinct. This last statement is verified by actual relators, not inferred only from the orbit labels: for every pair T, T' the certificate saves a word $w_{T,T'} \in K$ with $T(w_{T,T'}) = 1$ and $T'(w_{T,T'}) \neq 1$. There are 105 such words. For example

$$w = x_2 x_1 x_2^2 x_1^2 x_2^{-1} x_1^{-1} \tag{D.32.9}$$

has $H_0(w) = 1$, whereas $H_1(w)$ sends a to [320154] and b to [324105]. Its two exponent parities are zero, so it lies in K. All 105 saved words are checked both in their two actual images and in (D.32.6).

This also gives the precise comparison between kernel orbits and the relevant automorphisms. If two of the K-kernels agreed, their onto maps $K \to S$ would differ by some $\eta \in \mathrm{Aut}(S) = G$. For any $g \in \widehat{F}$, the conjugation actions of $T'(g)$ and $\eta T(g)\eta^{-1}$ on that onto image would then agree. Their difference centralizes S in G and is trivial. Hence $T' = \eta T \eta^{-1}$ as actual maps on all of $\widehat{F}$. The literal relators rule this out, including automorphisms outside the inner subgroup S.

The actual common-outer products and their source. Define

$$\begin{aligned} \mathcal{D} &= \{(g_1, \dots, g_{12}) \in G^{12} : \overline{g}_1 = \cdots = \overline{g}_{12} \in O\}, \\ \mathcal{E} &= \{(g_0, g_1, \dots, g_{12}) \in G^{13} : \overline{g}_0 = \cdots = \overline{g}_{12} \in O\}, \end{aligned} \tag{D.32.10}$$

with coordinatewise multiplication. Projection forgetting g_0 is π in (D.32.1). Its kernel is the actual first copy of S; the map $d \mapsto (d_1, d)$ is a splitting, where d_1 means the first coordinate of $d \in \mathcal{D}$. Thus the extension is split and its kernel is simple.

The source maps, in the exact coordinate order used below, are

$$\begin{aligned} q_{\mathcal{D}} &= (H_0, L_0, H_1, L_1, \dots, H_5, L_5), \\ q_{\mathcal{E}} &= (H_6, H_0, L_0, H_1, L_1, \dots, H_5, L_5). \end{aligned} \tag{D.32.11}$$

They land in (D.32.10) because of their common outer map. They are epimorphisms. Indeed their coordinate maps on K are onto the nonabelian simple group S and have pairwise distinct kernels by the relators above. The simple subdirect-product argument therefore gives all of S^{12} or S^{13}. Here is the exact implication: induct on the number of coordinates; if the last coordinate were constrained after projection to the preceding full product, it would be an onto homomorphism from that product to S. The commuting normal images of its simple factors force this homomorphism to factor through one coordinate, repeating a kernel. This contradicts the saved witness. Together with the common onto map to O, full generation and the two orders in (D.32.1) follow.

Let α swap each adjacent pair of coordinates of $\mathcal{D}$. This is an actual automorphism of the displayed group, of order two, and fixes its common quotient O. It preserves each labelled peripheral conjugacy class. For the first peripheral all coordinates equal U and no conjugator is needed. For the second, choose $c_j \in S$ with

$$c_j V_{H_j} c_j^{-1} = V_{L_j}. \tag{D.32.12}$$

Then the tuple $(c_0, c_0^{-1}, \dots, c_5, c_5^{-1})$ belongs to $\mathcal{D}$ and performs all six swaps. For the third peripheral, choose $b_j \in G$ of outer class e_1 sending the H_j third value to the L_j third value. These elements exist by (D.32.7), and the tuple $(b_0, b_0^{-1}, \dots, b_5, b_5^{-1})$ again belongs to $\mathcal{D}$: all twelve outer coordinates are e_1. The certificate supplies every c_j, b_j as an actual saved automorphism and verifies each conjugacy equation. Denote the three resulting base conjugators by h_i. Their common outer coordinates are respectively

$$0, \quad 0, \quad e_1. \tag{D.32.13}$$

The complete ordinary centralizer union. Write $d_i = q_{\mathcal{D}}(x_i)$ and let $\rho : \mathcal{D} \to O$ be the common-outer projection. The exact centralizer images are

$$\rho(C_{\mathcal{D}}(d_1)) = \langle e_1 \rangle, \qquad \rho(C_{\mathcal{D}}(d_2)) = \langle e_2 \rangle, \qquad \rho(C_{\mathcal{D}}(d_3)) = \langle e_1 + e_2 \rangle. \tag{D.32.14}$$

To prove this, the image of the centralizer of any relevant coordinate in G is exactly the indicated line. Equivalently, that line is the stabilizer in O of its S-conjugacy class. This is

computed for the actual classes in (D.32.7). The image of a centralizer in the common-outer product is the intersection of these coordinate images. Equality with the intersection follows by choosing in each coordinate a centralizing element with the same prescribed outer value; the resulting tuple belongs to (D.32.10). The certificate saves such a tuple for both values of every line in (D.32.14).

Every base conjugator sending d_i to $\alpha(d_i)$ belongs to the actual coset $h_i C_{\mathcal{D}}(d_i)$. Thus all their possible outer values, allowing the full ordinary conjugacy condition, are

$$\langle e_1\rangle, \qquad \langle e_2\rangle, \qquad e_1 + \langle e_1 + e_2\rangle = \{e_1, e_2\}. \tag{D.32.15}$$

The first line stabilizes $\mathcal{U}$, the second stabilizes $\mathcal{V}$, and both elements of the third coset send $\mathcal{W}_H$ to $\mathcal{W}_L$. Consequently the entire ordinary union of allowable first-coordinate triples over $\alpha q_{\mathcal{D}}$ is exactly

$$(\mathcal{U}, \mathcal{V}, \mathcal{W}_L). \tag{D.32.16}$$

This is also sufficient, not just necessary. For any triple in that profile, choose first-coordinate conjugators of outer values $0, 0, e_1$. Pair them with the actual twelve-coordinate h_i above; the pairs belong to $\mathcal{E}$ and realize the three required conjugacies. The product relation is coordinatewise. Hence the 2160 triples in the second row of (D.32.8) give exactly all ordinary weak lifts in (D.32.2).

Exhaustion and the exact two fibres. Each triple in (D.32.16) is uniquely an S-conjugate of one of the six L_j representatives. All six are already present in the base source. More exactly, under the swapped base map $\alpha q_{\mathcal{D}}$, the coordinate in the position formerly occupied by H_j equals L_j. If the added first-coordinate map is cL_jc^{-1}, its image in $\mathcal{E}$ is therefore the graph of the actual homomorphism

$$\mathcal{D} \longrightarrow \mathcal{E}, \qquad d \longmapsto (c\ \mathrm{pr}_{H_j}(d)c^{-1}, d). \tag{D.32.17}$$

Here $c \in S$ and pr_{H_j} means that fixed coordinate position, not the old source map. This graph is a complement to the first S factor and has order $|\mathcal{D}|$. It is a proper subgroup of $\mathcal{E}$ of index 360. The six indices and 360 distinct inner conjugates give the 2160 weak lifts, all complementary and none epimorphic. This proves the ordinary obstruction.

For the original base map, the same calculation using (D.32.14) gives the entire first-coordinate profile $(\mathcal{U}, \mathcal{V}, \mathcal{W}_H)$. Of its nine complete kernel classes, six $H_0, \ldots, H_5$ are present in the base and give 2160 graph lifts. The three remaining classes H_6, H_7, H_8 have distinct K-kernels from every base coordinate by the actual relators. Each therefore gives a full subdirect product and 360 epimorphisms. This proves the original counts $3240, 2160, 1080$ and also verifies the original source choice H_6 directly.

The exact pro-five comparison. Both groups in (D.32.10) have a perfect normal subgroup, S^{12} or S^{13}, and quotient O of order four. Every map to a finite five-group is therefore trivial, by the same explicit quotient argument as in D.31. For any finite five-group source quotient L with a multiplier-one peripheral automorphism β, the paired sources generate $\mathcal{E} \times L$ and $\mathcal{D} \times L$: the subdirect-product comparison would otherwise give a nontrivial common five-group quotient. Their maximal pro-five maps are the actual coordinate projections to L. The automorphism $\alpha \times \beta$ is peripheral on the base, using the pairs of the conjugators above and the prescribed conjugators for β. Any ordinary epimorphic lift would project to a lift in (D.32.2), which has just been excluded. For $1 \times \beta$ the explicit paired map $(q_{\mathcal{E}}, \beta q_L)$ is an epimorphic peripheral lift. Both base automorphisms induce the same β on the maximal pro-five quotient.

The computation files in the `code` directory are `a_peripheral_a6_nielsen.py` and `a_peripheral_a6_ordinary.py`, with executed outputs and complete JSON certificates. The first retains every fixed-first orbit and its generated subgroup. The second retains all 105 literal kernel witnesses, all twelve-coordinate conjugators, the complete centralizer-image vectors, and the six exact graph coordinate maps. The common-outer product orders are derived from the proved simple-factor generation argument; they are not reported as an enumeration of those large groups. The final status is `PASS_A6_ORDINARY_CHIEF_OBSTRUCTION`.

This is an ordinary finite chief-step obstruction, with all base-centralizer choices included. The explicit products and pro-five projections show its precise limitation for D.1–D.2: it excludes a finite branch while the same prescribed pro-five shadow has the admitted branch displayed above. □

4.16 Odd-order ordinary shadows at a single A_6 chief step

Statement D.33 (Proved). Every odd-order multiplier-one ordinary shadow lifts across a single A_6 kernel. *Let*

$$1 \longrightarrow S \longrightarrow E \xrightarrow{\pi} D \longrightarrow 1, \qquad S \simeq A_6, \tag{D.33.1}$$

be a finite extension, with a given epimorphism $q_E : \widehat{F} \twoheadrightarrow E$ and peripheral values X_i, whose images in D are d_i. Let $\alpha \in \mathrm{Aut}(D)$ have odd order m and satisfy $\alpha(d_i) \in d_i^D$ for the three labelled peripherals. There is an actual automorphism $\widetilde{\alpha}$ of E, of order m, covering α, fixing S pointwise, and satisfying

$$\widetilde{\alpha}(X_i) \in X_i^E. \tag{D.33.2}$$

Consequently $\widetilde{\alpha} q_E$ is an ordinary peripheral epimorphic lift of $\alpha\pi q_E$. In particular a multiplier-one ordinary obstruction of order five cannot occur at a single A_6 kernel.

Proof. Use the actual identification $\mathrm{Aut}(S) = \mathrm{Aut}(A_6)$ and $\mathrm{Out}(S) = O = \mathbf{F}_2^2$ proved by the complete automorphism calculation in D.31. The extension determines a homomorphism $\epsilon : D \to O$. Centerlessness gives the actual isomorphism

$$E \xrightarrow{\sim} \mathrm{Aut}(S) \times_{O,\epsilon} D, \qquad e \longmapsto (\mathrm{Ad}_e, \pi(e)). \tag{D.33.3}$$

Its kernel is $Z(S) = 1$, and any compatible automorphism over a given d differs from the conjugation action of a chosen lift of d by an inner automorphism of S. This proves surjectivity as well.

Choose $h_i \in D$ with $\alpha(d_i) = h_i d_i h_i^{-1}$. Since O is abelian, $\epsilon(\alpha(d_i)) = \epsilon(d_i)$. The elements d_1, d_2 generate D, so the two homomorphisms agree:

$$\epsilon\alpha = \epsilon. \tag{D.33.4}$$

Define the actual cumulative conjugator

$$c_i = \alpha^{m-1}(h_i)\, \alpha^{m-2}(h_i) \cdots \alpha(h_i) h_i. \tag{D.33.5}$$

Iterating the conjugacy equation gives $\alpha^m(d_i) = c_i d_i c_i^{-1}$. Thus $c_i \in C_D(d_i)$. Equation (D.33.4) and the oddness of m give

$$\epsilon(c_i) = m\epsilon(h_i) = \epsilon(h_i). \tag{D.33.6}$$

Put $h_i' = h_i c_i^{-1}$. The centralizer membership proves $h_i' d_i (h_i')^{-1} = \alpha(d_i)$, and (D.33.6) proves $\epsilon(h_i') = 0$. This is the exact adjustment allowed by ordinary peripheral conjugacy.

In the pullback (D.33.3), define

$$\widetilde{\alpha} : (g, d) \longmapsto (g, \alpha(d)). \tag{D.33.7}$$

Equation (D.33.4) makes it a homomorphism of that very group, with inverse $(g, d) \mapsto (g, \alpha^{-1}(d))$. It fixes its kernel pointwise. It has order exactly m: its mth power is the identity, and its projection to the base has order m. If $X_i = (g_i, d_i)$, then $(1, h_i')$ belongs to the pullback because $\epsilon(h_i') = 0$, and

$$(1, h_i')\, X_i\, (1, h_i')^{-1} = (g_i, \alpha(d_i)) = \widetilde{\alpha}(X_i). \tag{D.33.8}$$

Transporting (D.33.7)–(D.33.8) back through the actual isomorphism (D.33.3) proves the statement. □

D.33 concerns the ordinary fibre: the h_i' need not equal the initially prescribed h_i. Their difference is exactly the centralizer element (D.33.5), so the result does not erase the labelled obstruction in D.31. It also gives a precise limitation on changing the order-two example of D.32 into an order-five example while retaining a single A_6 kernel. For an A_6 chief kernel, an ordinary order-five obstruction would have to involve more than one simple factor. In that case the actual outer group is

$$\mathrm{Out}(A_6^{\Omega}) = (\mathbf{F}_2^2)^{\Omega} \rtimes \mathrm{Sym}(\Omega), \tag{D.33.9}$$

with its transitive factor action. The argument proving (D.33.4) used the abelian single-factor outer quotient; the displayed permutation component must be retained in an analysis of that remaining case.

4.17 Exact ordinary lifts for cyclic and two-group outer images

Statement D.34 (Proved). Cyclic outer image admits every multiplier-one peripheral base automorphism. *Let*

$$1 \longrightarrow A \longrightarrow E \xrightarrow{\pi} D \longrightarrow 1 \tag{D.34.1}$$

be a finite extension with $Z(A) = 1$ and a given source epimorphism $q_E : \widehat{F} \twoheadrightarrow E$. Let $X_i = q_E(x_i)$ and $d_i = \pi(X_i)$. Write $\epsilon : D \to \mathrm{Out}(A)$ for its actual outer action, and suppose its image is cyclic. Every $\alpha \in \mathrm{Aut}(D)$ with $\alpha(d_i) \in d_i^D$ has an actual automorphism lift $\widetilde{\alpha} \in \mathrm{Aut}(E)$ satisfying $\widetilde{\alpha}(X_i) \in X_i^E$. The full group $\mathrm{Out}(A)$ need not be abelian.

Proof. As in (D.33.3), use the actual centerless pullback isomorphism

$$E \xrightarrow{\sim} \mathrm{Aut}(A) \times_{\mathrm{Out}(A),\epsilon} D, \qquad e \longmapsto (\mathrm{Ad}_e, \pi(e)). \tag{D.34.2}$$

Choose $h_i \in D$ with $\alpha(d_i) = h_i d_i h_i^{-1}$. All $\epsilon(h_i)$ and $\epsilon(d_i)$ lie in the cyclic image C and commute. Since d_1, d_2 generate D, this proves

$$\epsilon\alpha = \epsilon. \tag{D.34.3}$$

Choose a generator t of C, of order n, and write $\epsilon(d_i) = t^{a_i}$ and $\epsilon(h_i) = t^{b_i}$. The actual source relation gives $a_1 + a_2 + a_3 = 0 \pmod{n}$, and a_1, a_2 generate $\mathbf{Z}/n$.

There is an $r \in \mathbf{Z}/n$ with

$$r - b_i \in a_i(\mathbf{Z}/n) \qquad (i = 1, 2, 3). \tag{D.34.4}$$

Here is a constructive proof. For every p^e exactly dividing n, at least one of a_1, a_2 is a unit modulo p, and the sum relation makes at least two of a_1, a_2, a_3 units modulo p. Thus at most one index imposes a congruence on r in that primary component. For that index, take $r = b_i \pmod{p^e}$; if none does, take $r = 0 \pmod{p^e}$. The Chinese remainder isomorphism combines these choices and proves (D.34.4). Solving the three resulting linear congruences gives integers k_i with $a_i k_i = r - b_i \pmod{n}$.

Put $h_i' = h_i d_i^{k_i}$. Right multiplication by a power of d_i retains its conjugacy equation, and

$$\epsilon(h_i') = t^r \qquad (i = 1, 2, 3). \tag{D.34.5}$$

Choose an actual $c \in \mathrm{Aut}(A)$ representing the outer class t^r. In (D.34.2) define

$$\widetilde{\alpha}(g, d) = (cgc^{-1}, \alpha(d)). \tag{D.34.6}$$

Conjugation by the outer class of c fixes every element of $\epsilon(D) = C$, and (D.34.3) holds. Thus (D.34.6) preserves the actual pullback condition. Its inverse is $(g, d) \mapsto (c^{-1}gc, \alpha^{-1}(d))$, so it is an automorphism over α. If $X_i = (g_i, d_i)$, then (c, h_i') belongs to the pullback by (D.34.5), and

$$(c, h_i')X_i(c, h_i')^{-1} = (cg_ic^{-1}, \alpha(d_i)) = \widetilde{\alpha}(X_i). \tag{D.34.7}$$

This is the required ordinary peripheral lift, transported through (D.34.2). □

Statement D.35 (Proved). A two-group outer image admits every order-five multiplier-one peripheral shadow. *In* (D.34.1)*, retain $Z(A) = 1$ and the given two-generated source. Suppose the actual image $\epsilon(D)$ is a two-group. Every order-five automorphism α of D preserving the three labelled peripheral conjugacy classes has an ordinary peripheral automorphism lift of order five to E, fixing A pointwise. In particular an order-five ordinary obstruction with kernel A_6^{Ω} requires its transitive factor-permutation image to have non-two-power order. It cannot occur with two simple factors or with any purely two-group factor-permutation action.*

Proof. Let P be the characteristic maximal two-group quotient of the finite group D. It is the quotient by the intersection of kernels of all maps to finite two-groups. There are only finitely many distinct such kernels in D; the resulting quotient is a subgroup of their finite product and is itself a two-group. The given source shows that P is generated by at most two elements.

There is no nontrivial automorphism of order five of a two-generated finite two-group P. To prove this directly, let $\Phi(P)$ be its Frattini subgroup, and put $d = \dim_{\mathbf{F}_2} P/\Phi(P) \le 2$. The action on this quotient has image in $\mathrm{GL}_d(\mathbf{F}_2)$, whose order divides six. The kernel of this action is a two-group. Indeed fix one ordered basis of $P/\Phi(P)$ and consider all ordered generating tuples lifting it. Every lift is generating: a proper generated subgroup would lie in a maximal subgroup of index two, contradicting its full image modulo $\Phi(P)$. There are exactly $|\Phi(P)|^d$ such tuples. An automorphism fixing the Frattini quotient acts freely on this set, because an automorphism fixing a generating tuple is the identity. Its group order therefore divides the two-power $|\Phi(P)|^d$. This proves the claim, including the trivial quotient cases.

The order-five automorphism α induces the identity on P. Since the actual map ϵ factors through P, it follows, before any conjugator adjustment, that

$$\epsilon\alpha = \epsilon. \tag{D.35.1}$$

Choose h_i with $\alpha(d_i) = h_i d_i h_i^{-1}$ and form

$$c_i = \alpha^4(h_i)\alpha^3(h_i)\alpha^2(h_i)\alpha(h_i)h_i. \tag{D.35.2}$$

Then $c_i \in C_D(d_i)$, and (D.35.1) gives the literal power identity

$$\epsilon(c_i) = \epsilon(h_i)^5. \tag{D.35.3}$$

No commutativity of the two-group image is assumed: the five factors on the right are the same element. Choose an integer k with $5k \equiv 1$ modulo the two-power exponent of $\epsilon(D)$ and put $h_i' = h_i c_i^{-k}$. It still conjugates d_i to $\alpha(d_i)$ and now satisfies $\epsilon(h_i') = 1$.

In the actual centerless pullback (D.34.2), the formula

$$(g, d) \longmapsto (g, \alpha(d)) \tag{D.35.4}$$

is an automorphism by (D.35.1). It has order five, fixes A pointwise, and its peripheral conjugators are the actual elements $(1, h_i')$ of the pullback. This proves the lift.

For $A = A_6^{\Omega}$, the exact outer group is (D.33.9). If its factor-permutation image is a two-group, then $\epsilon(D)$ is an extension of a subgroup of $(\mathbf{F}_2^2)^{\Omega}$ by a two-group and is itself a two-group. The result therefore applies. For two simple factors, the transitive permutation image is the group of order two, so that case is included. □

Both results retain the ordinary centralizer freedom by explicit maps. Neither asserts the corresponding lift with every initially prescribed base conjugator unchanged.

4.18 An ordinary order-five obstruction at a split A_6^{20} chief step

Statement D.36 (Proved). Every nonidentity power of an ordinary peripheral order-five shadow is obstructed. *Put*

$$\begin{aligned} m &= 61478762617650840213814367698461663428608, \\ h &= 61478762617650840528585632300833961934848. \end{aligned} \tag{D.36.1}$$

There are explicitly specified finite groups, a split chief extension, and an actual source epimorphism

$$1 \longrightarrow A_6^{20} \longrightarrow \mathcal{E} \xrightarrow{\pi} \mathcal{B} \longrightarrow 1, \qquad q_{\mathcal{E}} : \widehat{F} \twoheadrightarrow \mathcal{E}, \tag{D.36.2}$$

with

$$|\mathcal{B}| = 5662310400 \cdot 360^{100m}, \qquad |\mathcal{E}| = 5662310400 \cdot 360^{100m+20}. \tag{D.36.3}$$

An actual automorphism γ of $\mathcal{B}$ has order five and preserves the three labelled peripheral conjugacy classes of $q_{\mathcal{B}} = \pi q_{\mathcal{E}}$, with multiplier one. For every $k = 1, 2, 3, 4$, the ordinary fibre

$$\left\{ f : \widehat{F} \longrightarrow \mathcal{E} : \pi f = \gamma^k q_{\mathcal{B}},\ f(x_i) \in q_{\mathcal{E}}(x_i)^{\mathcal{E}}\ (i = 1, 2, 3) \right\} \tag{D.36.4}$$

has weak solutions and has no epimorphic solution. All base conjugators, including their full centralizer ambiguity, are allowed in this assertion.

The maximal pro-five quotients of $\mathcal{B}$ and $\mathcal{E}$ are trivial. Taking their actual products with a prescribed finite five-group source L retains an obstructed branch $\gamma^k \times \beta^k$, for each $k = 1, 2, 3, 4$, and an admitted branch $1 \times \beta^k$, over the same multiplier-one pro-five shadow β^k. In particular, every 5-adic unit power of the specified order-five finite component remains obstructed. These maps concern a specified finite component; the simultaneous admitted branch is part of the asserted comparison with D.1.

Proof. We give the finite groups and maps before the exhaustion argument. The large integers in (D.36.1) count explicitly defined finite sets; selecting elements from these sets means taking their lexicographically first generator tuples in the fixed enumeration below. Thus the construction makes no choice of an unevaluated infinite lifting problem.

The small base with its actual order-five automorphism. Let $H = A_5$ act on

$$M = \{(a_0, \dots, a_4) \in \mathbf{F}_2^5 : \textstyle\sum a_i = 0\}, \qquad b_i = e_i + e_4 \quad (0 \le i < 4). \tag{D.36.5}$$

Use the actual permutations

$$u = (01234), \qquad v = (01342), \qquad w = (uv)^{-1} = (03412), \qquad C = \langle (012) \rangle. \tag{D.36.6}$$

All three peripheral permutations have order five and u, v generate the sixty-element group. Put $V = M^4$, with diagonal H-action, and give $V \rtimes H$ the multiplication $(a, s)(b, t) = (a + sb, st)$. The actual source map is

$$q_0^{\mathrm{mod}} : \widehat{F} \longrightarrow D_0 = V \rtimes H, \qquad x_1 \longmapsto ((b_0, b_1, b_2, b_3), u), \quad x_2 \longmapsto (0, v). \tag{D.36.7}$$

It is onto. One direct certificate follows every Cayley edge of $\langle u, v \rangle$: the resulting 61 actual source relators, evaluated in V, and their H-translates span all sixteen coordinates. Therefore the source image contains V and surjects onto H. The saved relator words and sixteen-dimensional span check prove the assertion without a generation assumption about a split extension. The same finite calculation verifies that every nonzero vector of M has an orbit spanning M, and that $\mathrm{End}_H(M) = \mathbf{F}_2$. Its crossed-homomorphism constraint matrix has rank four on the eight generator coordinates: $Z^1(H, M) = B^1(H, M)$ has dimension four. These give a second, compatible verification of (D.36.7).

Define the multiplicity operator

$$T(a_0, a_1, a_2, a_3) = (a_3, a_0 + a_3, a_1 + a_3, a_2 + a_3). \tag{D.36.8}$$

Its four columns are $2, 4, 8, 15$ in binary coordinates. It commutes with the actual H-action, $T^5 = 1$, and $T \neq 1$. Let $G = \mathrm{Aut}(A_6)$, $S = \mathrm{Inn}(A_6)$, and $O = G/S = \mathbf{F}_2^2$, with the actual multiplication and outer coordinates of D.31. Take as $q_Q : \widehat{F} \twoheadrightarrow G$ the map H_0 of (D.32.4)–(D.32.5). Its peripheral indices in the complete generator-image enumeration of D.31 are $0, 24, 1380$, and its outer values are $1, 2, 3$. Its complete peripheral centralizer images are

$$\overline{C_G(q_Q(x_1))} = \{0, 1\}, \quad \overline{C_G(q_Q(x_2))} = \{0, 2\}, \quad \overline{C_G(q_Q(x_3))} = \{0, 3\}. \tag{D.36.9}$$

The certificate enumerates all members of these three centralizers.

Set

$$D = D_0 \times G, \qquad q_0 = (q_0^{\mathrm{mod}}, q_Q), \qquad \alpha(a, s, g) = (Ta, s, g), \qquad d_i = q_0(x_i). \tag{D.36.10}$$

The map q_0 is onto. Indeed, D_0 is perfect: $[V, H] = V$ by the displayed orbit-span calculation, and A_5 is perfect. The group G has no nontrivial perfect quotient. For if $N \lhd G$, simplicity of S gives either $S \subset N$ or $N \cap S = 1$; in the latter case $[N, S] = 1$, whence $N \subset C_G(S) = 1$. Its proper quotients therefore factor through O, and G itself has nontrivial abelian quotient O. Goursat's actual common-quotient map for the two onto projections of q_0 must consequently be trivial. This proves the claimed product generation. In particular $|D| = 60 \cdot 2^{16} \cdot 1440 = 5662310400$.

Write $q_0^{\mathrm{mod}}(x_i) = (v_i, h_i)$. The operator $1 - h_i$ is invertible on each copy of M. For $j = 0, \dots, 4$, define the literal module element

$$c_i^{(j)} = (1 - h_i)^{-1}(T^j - 1)v_i \in V \subset D. \tag{D.36.11}$$

It satisfies

$$\alpha^j(d_i) = c_i^{(j)} d_i (c_i^{(j)})^{-1}, \qquad c_i^{(0)} = c_i^{(5)} = 0, \qquad c_i^{(j+1)} - c_i^{(j)} = \alpha^j(c_i^{(1)}). \tag{D.36.12}$$

These identities use the displayed multiplication in $V \rtimes H$; they are also checked on every saved vector. Thus α is an ordinary peripheral automorphism of exact order five.

A specified twenty-factor chief extension. For a vector in the basis $b_0, \dots, b_3$, put $\ell(a) = a_0 + a_1 + a_2$ and $r(a) = a_3$. The homomorphism

$$\epsilon : D_* = (V \rtimes C) \times G \longrightarrow O, \qquad \epsilon((a_0, a_1, a_2, a_3), s, g) = (\ell(a_0), r(a_1)) + \overline{g} \tag{D.36.13}$$

is well defined: the two module functionals are C-invariant. They are encoded by the sixteen-bit masks $7, 128$. Let $\Omega = D_* \backslash D = C \backslash H$, of size twenty, and choose representatives $t_\omega = (0, s_\omega, 1)$. Here s_ω is given by the signed-edge breadth-first tree in the u, v Schreier graph, with edge order $x_1, x_2, x_1^{-1}, x_2^{-1}$. All group permutations, cosets, and tree words are retained in the certificate. Put

$$a(\omega, d) = \epsilon(t_\omega d t_{\omega d}^{-1}), \qquad E_* = \{(z, d) \in G^\Omega \rtimes D : \overline{z(\omega)} = a(\omega, d)\}. \tag{D.36.14}$$

The multiplication is literally

$$(z, d)(z', d') = \big((z(\omega) z'(\omega d))_\omega, dd'\big). \tag{D.36.15}$$

The cocycle identity for a follows by cancelling the intermediate representative. Projection $E_* \to D$ is onto: in each coordinate choose an actual G-representative of the specified outer value. Its kernel is $A = S^\Omega$, and D permutes the twenty factors transitively. A normal subgroup of E_* contained in A is a product of simple direct factors and is therefore 1 or A. Thus this is an actual chief extension.

Put $K = \ker q_0$ and $\mathcal{H} = q_0^{-1}(D_*)$. Since α preserves D_*, these two source subgroups are the same for all five maps $q_j = \alpha^j q_0$. The index-twenty cover of the three-punctured sphere has four cycles of length five over each peripheral. It has twelve boundaries, genus five, and a free Schreier basis of rank 21. The twenty tree words will be denoted T_ω. They are actual words in x_1, x_2, whereas t_ω in (D.36.14) is the chosen element of D. This difference is retained below.

For a homomorphism $\varphi : \mathcal{H} \to G$ with $\overline{\varphi} = \epsilon q_j|_{\mathcal{H}}$, choose actual $b_{\omega,j} \in G$ satisfying

$$b_{*,j} = 1, \qquad \overline{b}_{\omega,j} = \epsilon(q_j(T_\omega) t_\omega^{-1}). \tag{D.36.16}$$

For example, take the least indexed G-element of that outer value away from the identity coset. The actual induced source map $p_\varphi : \widehat{F} \to E_*$, over q_j, is

$$z_\varphi(g)(\omega) = b_{\omega,j}^{-1} \varphi(T_\omega g T_{\omega g}^{-1}) b_{\omega g, j}. \tag{D.36.17}$$

Multiplying two right sides cancels the intermediate b and gives (D.36.15). Taking outer coordinates gives $a(\omega, q_j(g))$, by (D.36.16). Conversely restriction to the identity coordinate on $\mathcal{H}$, with the actual path values $z(T_\omega)(*)$, recovers this formula. This is the needed nonabelian Shapiro comparison, including its forced outer path coordinates.

The fixed peripheral classes and their full based transport. Let

$$\mathcal{C}_1 = X^S, \qquad \mathcal{C}_2 = q_Q(x_2)^S, \qquad \mathcal{C}_3 = q_Q(x_3)^S, \tag{D.36.18}$$

where X is the actual involution of (D.31.6). Their indices in the relative class table are $12, 7, 5$; their sizes are $15, 90, 90$, and their outer values are $1, 2, 3$. The action of O on S-classes means conjugation by any actual representative in G; it is independent of that representative.

Choose the least coset in each peripheral cycle as its start. Write T_ν for its tree word and

$$q_0^{\mathrm{mod}}(T_\nu) = (v_\nu, s_\nu), \qquad \delta_\nu = T_\nu x_i^5 T_\nu^{-1} \in \mathcal{H} \quad (\nu \text{ lies over } x_i). \tag{D.36.19}$$

There are four such ν for each i. Define $\eta_\nu = 2$ at the first x_1-cycle, which contains the identity coset, and $\eta_\nu = 0$ elsewhere. Specify the cycle-product class of a reference element $Z_i \in E_*$, over d_i, to be

$$\big(\epsilon_V(v_\nu) + \eta_\nu\big) \cdot \mathcal{C}_i, \qquad \epsilon_V(a) = (\ell(a_0), r(a_1)). \tag{D.36.20}$$

These actual reference elements exist coordinate by coordinate. Along a length-five cycle choose the first four coordinate elements with their required outer values, and choose the fifth so that the ordered product is the least element of the class in (D.36.20). The required outer product is $i \in O$, so the last coordinate has exactly the required outer value. The certificate gives all sixty coordinate elements.

For a source over q_j whose i-th peripheral is conjugate to Z_i by an element over $c_i^{(j)}$, the precise class of $\varphi(\delta_\nu)$ is

$$\mathcal{T}_{\nu,j} = \Big(\overline{q_Q(T_\nu)} + \epsilon_V((T^j - 1)v_\nu) + \epsilon_V(s_\nu c_i^{(j)}) + \eta_\nu\Big) \cdot \mathcal{C}_i. \tag{D.36.21}$$

Here $s_\nu c_i^{(j)}$ is the specified module action. To verify the formula, the cycle product changes under a lift of $c_i^{(j)}$ by the outer value $\epsilon_V(s_\nu c_i^{(j)})$. The based boundary value is that cycle product conjugated by $b_{\nu,j}$ from (D.36.16), whose outer value is $\epsilon_V(T^j v_\nu) + \overline{q_Q(T_\nu)}$. Adding the initial adjustment $\epsilon_V(v_\nu) + \eta_\nu$ in (D.36.20) proves (D.36.21). In particular the module term $\epsilon_V((T^j - 1)v_\nu)$ is required even though all five source maps have the same permutation action on the cosets.

For completeness the conjugacy criterion just used has a literal inverse. For two arrays over the same d_i, let P, P' be their ordered products around a cycle. Choose $a_0 \in S$ with $P' = a_0 P a_0^{-1}$. If the old and new coordinates are z_ω, z'_ω, successively put

$$a_{\omega d_i} = (z'_\omega)^{-1} a_\omega z_\omega. \tag{D.36.22}$$

All a_ω lie in S, and the return to the starting coordinate is a_0 exactly because of the product equation. This produces the actual kernel conjugator. Thus (D.36.21) is both necessary and sufficient for the specified local conjugator class. The five saved source maps verify this recurrence on every coordinate.

The complete finite count, with proper images removed. The twelve boundary words in the rank-21 Schreier basis are literal words. Across them every Schreier letter occurs once positively and once negatively. Introduce twelve boundary generators δ_ν. Merging faces along eleven tree edges gives a single quadratic relation with ten paired letters and twelve boundary letters. All these eliminations are actual Tietze maps: from relations $aU = 1$ and $a^{-1}V = 1$, substitute $a = U^{-1}$ and retain $UV = 1$. The remaining handles are extracted by the exact identity

$$\begin{gathered} R = aUbVa^{-1}Wb^{-1}Z, \qquad C = WVU, \\ A = C^{-1}aU, \qquad B = bVU, \qquad C^{-1}RC = [A, B]ZWVU. \end{gathered} \tag{D.36.23}$$

The substitutions are invertible: $a = CAU^{-1}$ and $b = BU^{-1}V^{-1}$. Cyclic changes of the remaining polygon conjugate previously extracted handles explicitly and leave each boundary generator unchanged. The certificate checks every free-word identity, the resulting relation, and the determinant -1 of the 21-element integral basis comparison. It also inverts these maps in G, producing actual homomorphisms on the original Schreier generators.

For each j, let $\mathscr{X}_j$ consist of the onto maps $\varphi : \mathcal{H} \to G$ with the exact outer character ϵq_j and all twelve conditions $\varphi(\delta_\nu) \in \mathcal{T}_{\nu,j}$. The complete count is

j	0	1	2	3	4
$\lvert\mathscr{X}_j\rvert$	$360h$	$360m$	$360m$	$360m$	$360m$.

(D.36.24)

We describe the entire finite calculation so that generation and framed transport are part of its proof.

First enumerate every subgroup $L \le G$ mapping onto O. Such a subgroup contains a pair with outer values $1, 2$. The first member can be moved, by S-conjugation, into one of its five inner conjugacy-class representatives. The 1800 resulting pairs give all starting subgroups up to conjugacy. Adjoining every inner element to each new subgroup exhausts all possibilities: a subgroup surjecting onto O is generated by such a starting pair and its intersection with S. This gives ten conjugacy classes and exactly 452 actual subgroups, with order histogram

$\lvert L\rvert$	8	16	32	40	144	1440
$\#L$	180	180	45	36	10	1.

(D.36.25)

For these subgroups G-conjugacy and S-conjugacy coincide: if L surjects onto O and $g \in G$, choose $l \in L$ with $\bar{l} = \bar{g}$; then $gLg^{-1} = (gl^{-1})L(gl^{-1})^{-1}$ and $gl^{-1} \in S$. Thus all prescribed boundary sets are preserved in using the ten representatives.

For each L, put $L_0 = L \cap S$. In $\mathbf{Z}[L]^{L_0}$ compute the actual elements

$$K^L_{a,b} = \sum_{\substack{u,v \in L \\ \bar{u}=a,\ \bar{v}=b}} [u, v], \qquad B^L_{\mathcal{T}} = \sum_{t \in \mathcal{T} \cap L} t. \tag{D.36.26}$$

The handle outer values are those obtained from the literal words in (D.36.23), and the boundary sets are exactly (D.36.21). The coefficient of 1 in the ordered product of the five appropriate $K^L_{a,b}$ and the twelve inverse-boundary sums counts all such maps into L. This is just expansion of the actual surface relation; it uses no inference from a rational character formula. The relative class-algebra structure constants are computed by counting x in one class with $x^{-1}g$ in the other. Every coefficient is an integer.

In increasing subgroup order subtract the counts of maps whose image is each actual proper subgroup:

$$E_L = N_L - \sum_{J < L,\ \bar{J} = O} E_J. \tag{D.36.27}$$

Because the prescribed outer character is onto O, this list includes every proper image. The count for $L = G$ is (D.36.24). The positive excess of kernel classes is

$$h - m = 314771264602372298506240. \tag{D.36.28}$$

As a separately executed check, omitting the one fixed flip $\eta_\nu = 2$ gives $360(m, h, h, h, h)$. An earlier exploratory calculation omitted the module-path term in (D.36.21); its programs and outputs are preserved with the suffix `incomplete_framing`. They are not used in (D.36.24). The corrected calculation includes both the term and the two opposite patterns.

The exact comparison with source kernels. Every $\varphi \in \mathscr{X}_j$ sends K onto S. Its image on K is normal in G, lies in S, and is therefore 1 or S. If it were 1, φ would factor through

$$\mathcal{H}/K = D_* = (V \rtimes C) \times G. \tag{D.36.29}$$

The first factor is solvable. Its image would be a solvable normal subgroup of G, hence trivial: a normal subgroup meeting S nontrivially contains the nonsolvable simple group S, while a normal subgroup disjoint from S centralizes S and is trivial. The factored map would then kill V, contradicting its prescribed nonzero outer character $\epsilon_V T^j$ on V. This proves $\varphi(K) = S$.

For the five ordered module-character pairs, the exact stabilizer and transporter calculation is

$$\{s \in H : \epsilon_V T^j s = \epsilon_V T^k \text{ on } V\} = \begin{cases} C, & j = k, \\ \varnothing, & j \neq k. \end{cases} \tag{D.36.30}$$

It is verified on all sixteen basis vectors for all sixty actual permutations and all 25 ordered pairs. Equivalently, the two masks $7,128$ on the specified module copies and the matrix (D.36.8) give the displayed finite calculation.

Let $R_\varphi = \ker(\varphi|_K)$. Its normalizer in $\widehat{F}$ is exactly $\mathcal{H}$. Indeed, the normalizer contains K, and its conjugation action on $K/R_\varphi = S$ gives an actual map to $\mathrm{Aut}(S) = G$ extending φ. Taking outer classes gives an extension of $\epsilon q_j|_{\mathcal{H}}$. Every subgroup of D containing D_* is the preimage of a subgroup of H containing C, since D_* already contains $V \times G$. Its action must fix the character pair on V; (D.36.30) forces that subgroup to be C. Thus the normalizer equals $\mathcal{H}$. In particular the twenty coordinate kernels in (D.36.17) are distinct.

If two restrictions to K have equal kernels, their onto maps $K \to S$ differ by an automorphism of S, that is, by an actual element $g \in G$. The two extended maps on $\mathcal{H}$ then differ by conjugation by g: their conjugation actions agree on the onto subgroup S, and $C_G(S) = 1$. Conversely such conjugation preserves the kernel. Within $\mathscr{X}_j$, exactly 360 elements of this G-orbit remain in the twelve prescribed classes. Indeed the common stabilizer in O of the three classes in (D.36.18) is 0; the independent class transports in (D.36.21) do not change that stabilizer because O is abelian. The S-action on onto maps is free. Thus $\mathscr{X}_j$ represents exactly

$$h, m, m, m, m \tag{D.36.31}$$

different kernels on K.

These are also distinct $\widehat{F}$-orbits of kernels. Any conjugator between two such kernels must normalize $\mathcal{H}$, by the normalizer equality just proved. Its image in H belongs to $N_H(C)$. The V-part acts trivially on the C-invariant characters, and the G-part acts trivially on O. Consequently the required character comparison is precisely (D.36.30). For equal colours it forces the conjugator into $\mathcal{H}$, where each kernel is already normal; for different colours no conjugator exists.

The simple-product generation argument is exact. An onto map to each of finitely many copies of a nonabelian simple group, with pairwise distinct kernels, maps onto their product. Inductively, a proper subdirect subgroup of $S^{n-1} \times S$ would be the graph of an onto map $S^{n-1} \to S$. The images of the direct factors are commuting normal subgroups of S; simplicity and centerlessness force this map to factor through one coordinate. Its kernel would coincide with an earlier one. Applying this argument to the twenty coordinate kernels proves that every map in $\mathscr{X}_j$ induces an epimorphism in (D.36.17), and applying it across colours proves all product-generation assertions used below.

The entire ordinary centralizer union. This step is independent of the numerical count. Since $1 - h_i$ is invertible on V, the element (v_i, h_i) is conjugate by a unique module translation to $(0, h_i)$. Its centralizer is conjugate to $C_H(h_i)$, which is the cyclic group of order five. Hence

$$C_D(d_i) = \langle q_0^{\mathrm{mod}}(x_i) \rangle \times C_G(q_Q(x_i)). \tag{D.36.32}$$

This group is generated by d_i and the elements $(1, c)$ with $c \in C_G(q_Q(x_i))$. Conjugation by the actual lift Z_i of d_i fixes Z_i. For $(1, c)$, the permutation of Ω is the identity and the outer value in every coordinate is $\bar{c}$. By (D.36.9), this is 0 or i. Every S-class in outer coset i is fixed by the outer action of i: use a representative of that very class as the conjugating element. Thus conjugation by any lift of $(1, c)$ preserves each cycle-product S-class of Z_i. The explicit recurrence (D.36.22) shows that it preserves its full A-conjugacy class. Changing

the chosen lift only adds an A-conjugation. It follows that every element of the complete base centralizer (D.36.32) preserves the A-class of Z_i.

Consequently, for every j, allowing all ordinary base conjugators from d_i to $\alpha^j(d_i)$ gives exactly the class condition (D.36.21). Multiplying a specified $c_i^{(j)}$ by any centralizer element cannot add another cycle-product profile. This proves the full ordinary comparison used by the exhaustion argument, rather than only a comparison for one chosen tuple of conjugators.

The five-colour finite product and its source. For $j \in \mathbf{Z}/5\mathbf{Z}$, let

$$E_j = E_* \times_{D,\alpha^j} D = \{(e,d) : \mathrm{pr}(e) = \alpha^j(d)\}. \tag{D.36.33}$$

Order the onto maps in $\mathscr{X}_j$ by their 21 generator values in the actual enumeration of G. In each equivalence class under G-conjugation retain its least member still satisfying (D.36.21). The preceding kernel comparison shows that these lists have lengths h, m, m, m, m and have disjoint kernel orbits across colours. Use the first m members for colour zero and all m members for each other colour. Denote the induced source maps by $p_{j,r} : \widehat{F} \to E_*$, over q_j, for $1 \le r \le m$. Use the next colour-zero member for a map p_*. This is a completely specified finite procedure on tuples in G^{21}; its required positive excess is (D.36.28).

Define the actual fibre product

$$\mathcal{B} = \left\{((e_{j,r}), d) : \mathrm{pr}(e_{j,r}) = \alpha^j(d) \text{ for every } j, r\right\}, \qquad q_{\mathcal{B}}(g) = ((p_{j,r}(g)), q_0(g)). \tag{D.36.34}$$

The comparison of all coordinate kernels proves that $q_{\mathcal{B}}$ is onto. Now define

$$\begin{aligned} \mathcal{E} &= \mathcal{B} \times_D E_*, \\ q_{\mathcal{E}}(g) &= (q_{\mathcal{B}}(g), p_*(g)), \\ \gamma((e_{j,r}), d) &= ((e_{j+1,r}), \alpha(d)). \end{aligned} \tag{D.36.35}$$

The same distinct-kernel argument proves that $q_{\mathcal{E}}$ is onto. The shift in (D.36.35) has the required type because $\mathrm{pr}(e_{j+1,r}) = \alpha^j(\alpha(d))$, including $j = 4$. It is an automorphism of exact order five. Forgetting the last E_*-coordinate gives (D.36.2), with kernel S^{20}. It splits by the actual map taking that coordinate to $e_{0,1}$. Its factor action is transitive, so the kernel is chief, as for (D.36.14). Counting the $5m$ kernels of size 360^{20} proves (D.36.3).

The automorphism γ is ordinary peripheral. For each i, j, r, choose an actual conjugator from Z_i to $p_{j,r}(x_i)$ over $c_i^{(j)}$. These exist by (D.36.21)–(D.36.22). The quotient of two successive such conjugators lies over $c_i^{(j+1)} - c_i^{(j)} = \alpha^j(c_i^{(1)})$. These $5m$ coordinates therefore assemble into one actual element of $\mathcal{B}$ over $c_i^{(1)}$ conjugating $q_{\mathcal{B}}(x_i)$ to $\gamma q_{\mathcal{B}}(x_i)$. The cyclic closing coordinate uses $c_i^{(5)} = 0$ and gives the same equality. This establishes all three peripheral conjugacies in the actual product.

Exhaustion for every nonidentity power. Suppose $f : \widehat{F} \to \mathcal{E}$ belongs to (D.36.4) and is onto. Its extra E_*-coordinate e is an epimorphism over $q_k = \alpha^k q_0$. Project an arbitrary peripheral conjugator in $\mathcal{E}$ to D. Its value belongs to $c_i^{(k)} C_D(d_i)$. The full centralizer comparison (D.36.32) proves that the corresponding map on $\mathcal{H}$ belongs to $\mathscr{X}_k$. Its K-kernel orbit therefore occurs among the m already selected colour-k maps.

In the base map $\gamma^k q_{\mathcal{B}}$, these same m maps occur as the $j = 0$ coordinates: the shift has put $p_{k,r}$ in those positions. Thus some simple-coordinate kernel of the extra E_*-map coincides with a simple-coordinate kernel already present in the base. The image of K in the kernel of $\mathcal{E} \to D$ is contained in the corresponding proper diagonal graph. An epimorphism f would send K onto that whole kernel, a contradiction. This proves the emptiness of the generating fibre for all four values of k.

Weak lifts nevertheless exist. For any r, compose $\gamma^k q_{\mathcal{B}}$ with the split section that uses $e_{0,r}$ as the extra coordinate. The extra source map is then $p_{k,r}$. Its peripheral classes agree with those of p_* after conjugators over $c_i^{(k)}$; these may be paired with the already constructed base conjugators, giving actual elements of $\mathcal{E}$. This supplies an element of (D.36.4) with complementary image.

The exact pro-five comparison and reproducible inputs. Every homomorphism from A_6 or A_5 to a finite five-group is trivial, and a homomorphism from the module V to a five-group is trivial. The quotient $G/S = O$ has order four. Applying these facts to the displayed normal sequences shows directly that $D, E_*, \mathcal{B}$, and $\mathcal{E}$ all have trivial maximal pro-five quotient. For a finite five-group source L, the maps $(q_{\mathcal{E}}, q_L)$ and $(q_{\mathcal{B}}, q_L)$ are onto the actual products: Goursat's common quotient would be a five-group quotient of the first factor and is therefore trivial. Their maximal pro-five quotients are exactly L. Projection to (D.36.4) obstructs every branch $\gamma^k \times \beta^k$; the actual automorphism $1_{\mathcal{E}} \times \beta^k$ gives the admitted branch $1_{\mathcal{B}} \times \beta^k$. This proves the stated comparison without discarding the specified finite component.

The finite dependencies are the actual A_6 model and source table from D.31–D.32, followed by the programs in the `code` directory with common prefix `a_peripheral_` and suffixes `a5_deleted.py`, `a6_outer_subgroups.cpp`, `a6_cup_epi.py`, `surface20.py`, `a6_boundary_search.py`, and `order5_certificate.py`. The authoritative surface data have suffix `surface20_framed`; the authoritative count data have suffix `boundary_framed_flip`. The last certificate constructs one actual onto surface map in each colour, inverts the Tietze maps to its 21 original generator values, computes the two induced twenty-coordinate source arrays using (D.36.17), and checks all twelve cycle products and all sixty peripheral conjugator coordinates for every colour. It also checks all 25 transporter sets in (D.36.30) and the complete base centralizer images. The five-colour exhaustion uses the full counts (D.36.24), with the exact finite lexicographic selection in (D.36.34), rather than claiming to enumerate the enormous product group. □

4.19 Characteristic finite targets for the three weight-five special directions

Statement D.37 (Proved). An exact characteristic Magnus target and all three ordinary outer parameters. *For every $n \geq 1$ there is an explicitly specified characteristic finite quotient*

$$\rho_n : \widehat{F} \twoheadrightarrow P_n, \qquad |P_n| = 5^{17n+2}, \tag{D.37.1}$$

which is metabelian of nilpotency class six. It has $P_n^{\mathrm{ab}} \simeq (\mathbf{Z}/5^{n+1})^2$, derived subgroup $(\mathbf{Z}/5^n)^{15}$, and centre $\gamma_6 P_n \simeq (\mathbf{Z}/5^n)^5$. Its kernel is fully invariant, by an explicit Fox substitution map. Coefficient reduction gives onto maps $P_{n+1} \to P_n$.

There is a specified injective homomorphism

$$(\mathbf{Z}/5^n)^3 \longrightarrow \mathrm{Out}^\sharp(P_n), \qquad (A, B, C) \longmapsto [\beta_{A,B,C}], \tag{D.37.2}$$

whose representatives fix x_1 and have actual multiplier-one conjugators at x_2 and x_3. The parameters have exactly the requested leading coordinate

$$(X + Y)(AX^2 + BXY + CY^2)[x, y]. \tag{D.37.3}$$

The homomorphism is injective in the ordinary outer group, including all inner automorphisms of the finite quotient. At $n = 1$, every nonzero parameter gives an actual automorphism of order five of the characteristic group of order 5^{19}. The parameter maps and their geometric $\mathfrak{S}_3$ action are compatible with coefficient reduction.

These are fixed finite source targets for the full profinite lifting question. No assertion that a non-gamma parameter in (D.37.2) *has a full profinite peripheral lift is included.*

Proof. Put $R_n = \mathbf{Z}/5^n$, let $I = (X, Y)$, and define

$$B_n = R_n[X, Y]/I^7, \qquad U_n = (R_n[X, Y]/I^6)^2. \tag{D.37.4}$$

The ambient finite group consists of pairs (a, u) with $a \in 1 + IB_n$ and $u \in U_n$, with the exact multiplication

$$(a, u)(b, v) = (ab, u + av). \tag{D.37.5}$$

Multiplication by a on U_n uses its reduction modulo I^6. This is a finite five-group. Define P_n to be the actual subgroup generated by

$$x = (1 + X, (1, 0)), \qquad y = (1 + Y, (0, 1)). \tag{D.37.6}$$

Thus ρ_n is an actual continuous epimorphism from the free profinite group, rather than a presentation whose realization is left implicit.

Every element $(a, (u_1, u_2))$ in this source image satisfies

$$a - 1 = Xu_1 + Yu_2 \quad \text{in } B_n. \tag{D.37.7}$$

The identity holds for the two generators and is preserved by (D.37.5) and inversion. The two diagonal generators have order 5^{n+1} and are independent. To see the order exactly, for $k \geq 1$ the coefficients of $(1 + X)^{5^k} - 1$ through degree six have valuations $k - v_5(j)$ at degree j, when that coefficient is present. The degree-five coefficient is nonzero modulo 5^n for $k = n$, and all coefficients vanish for $k = n + 1$. Specializing respectively $Y = 0$ and $X = 0$ proves independence. The same binomial calculation applied to $\sum_{r=0}^{5^{n+1}-1}(1 + X)^r$, through degree five, shows that the full group element x, and likewise y, has order 5^{n+1}.

Write $c = [x, y]$. Direct multiplication gives

$$c = (1, (-Y, X)), \qquad c^p := (1, p(-Y, X)) \quad \big(p \in R_n[X, Y]_{\leq 4}\big). \tag{D.37.8}$$

Here the notation c^p denotes the displayed module element, not an undefined exponent by a polynomial. It is realized by actual group operations: conjugation by x multiplies its coefficient by $1 + X$, and $[x, c^p] = c^{Xp}$; similarly $[y, c^p] = c^{Yp}$. Integer powers and products realize every coefficient in R_n. Thus all fifteen monomials of total degree at most four in (D.37.8) belong to the derived subgroup.

Conversely, if the diagonal of a source element is one, (D.37.7) is the actual polynomial identity $Xu_1 + Yu_2 = 0$, of degree at most six. It implies $u_1 = -Yp,\ u_2 = Xp$ for a unique polynomial of degree at most four. This uses only that multiplication by a polynomial variable is injective in a polynomial ring over R_n: setting $X = 0$ first shows that X divides u_2, and the identity then gives the claim. No division by a nonunit of R_n occurs. Therefore the kernel of the diagonal map is precisely the fifteen-coordinate module in (D.37.8). It is the derived subgroup. Every source element has a unique normal form

$$x^r y^s c^p, \qquad 0 \leq r, s < 5^{n+1}, \quad p \in R_n[X, Y]_{\leq 4}. \tag{D.37.9}$$

The product of the diagonal and module orders proves (D.37.1) and the asserted abelianization. Furthermore

$$\gamma_j P_n = \{c^p : p \in I^{j-2}/I^5\} \quad (2 \leq j \leq 6), \qquad \gamma_7 P_n = 1. \tag{D.37.10}$$

Both inclusions follow from the exact commutators $[x, c^p] = c^{Xp}$, $[y, c^p] = c^{Yp}$. In particular the class is exactly six.

The actual fully invariant kernel map. Let $g = (a_g, u_g)$ and $h = (a_h, u_h)$ be any two elements of P_n. The substitutions

$$\sigma_{g,h}(X) = a_g - 1, \qquad \sigma_{g,h}(Y) = a_h - 1 \tag{D.37.11}$$

preserve the augmentation ideal and its powers. Define

$$\Phi_{g,h}(a, (u_1, u_2)) = (\sigma_{g,h}(a),\ \sigma_{g,h}(u_1)u_g + \sigma_{g,h}(u_2)u_h)\,. \tag{D.37.12}$$

This is a homomorphism for (D.37.5): the module formula is $\sigma(u + av)$ followed by the displayed linear map, and it equals the module part of the product of the two images. The truncations have the required types because $\sigma(I^j) \subset I^j$. The map sends x to g and y to h. Consequently, for every actual word w,

$$w(g, h) = \Phi_{g,h}(\rho_n(w)). \tag{D.37.13}$$

The same equality holds for profinite words by continuity into the finite groups. If $\rho_n(w) = 1$, its image under every endomorphism of $\widehat{F}$ is therefore again in $\ker \rho_n$. This proves full invariance, and hence characteristicness, with the exact quotient map. It also proves that the geometric permutations of the three peripherals act on this same target. Reducing the coefficients in (D.37.4)–(D.37.6) maps the two generators onto those of P_n, which proves the stated surjectivity of $P_{n+1} \to P_n$.

All three special automorphisms. For

$$Q = AX^2 + BXY + CY^2, \quad p = (X + Y)Q, \quad f_Q = c^p, \quad h_Q = c^{YQ}, \quad \Delta_Q = c^{-Y(X+Y)Q}, \tag{D.37.14}$$

all four elements on the right are actual group elements by (D.37.8). The last belongs to $\gamma_6 P_n$ and is central. The linear coefficient of Y in a diagonal element is an additive homomorphism

$$s : P_n \longrightarrow R_n, \qquad s(x) = 0, \quad s(y) = 1. \tag{D.37.15}$$

Define the actual automorphism

$$\beta_Q(g) = g\Delta_Q^{s(g)}. \tag{D.37.16}$$

Centrality proves the homomorphism identity. It fixes Δ_Q, since $s(\Delta_Q) = 0$; the inverse is obtained by replacing Q by $-Q$. The same formula proves $\beta_Q\beta_{Q'} = \beta_{Q+Q'}$. It fixes the whole derived subgroup and fixes x. The exact relation

$$[f_Q, y] = c^{-Yp} = \Delta_Q \tag{D.37.17}$$

shows that $\beta_Q(y) = f_Q y f_Q^{-1}$.

Put $z = (xy)^{-1}$. Then $\beta_Q(z) = z\Delta_Q^{-1}$. Conjugation of a module element by z multiplies its coefficient by $((1 + X)(1 + Y))^{-1}$. The identity

$$\Big(1 - ((1 + X)(1 + Y))^{-1}\Big)\, YQ = Y(X + Y)Q \pmod{I^5} \tag{D.37.18}$$

is exact at the required degree: the difference between the first parenthesis and $X + Y$ has degree at least two, whereas YQ has degree three. It follows that $[h_Q, z] = \Delta_Q^{-1}$, and hence

$$\beta_Q(z) = h_Q z h_Q^{-1}. \tag{D.37.19}$$

Thus all three peripheral conditions hold as literal identities in P_n. The image of the normalized conjugator in $\gamma_5 P_n/\gamma_6 P_n$ is precisely $(X+Y)Q[x, y]$, because the group conjugation differences X, Y induce the actual ordinary graded brackets. This proves the stated leading coordinate comparison. The polynomial $-Y(X + Y)Q$ has degree four; multiplication by $Y(X + Y)$ is injective on the quadratic coefficient module. Thus the order of β_Q is exactly the additive order of Q, before taking outer classes.

The ordinary outer quotient, including restricted inner powers. We compute the actual centralizer of x, rather than using a centralizer theorem for a different free group. Let $g = x^r y^s c^p$ be the normal form (D.37.9), and write

$$S_s(Y) = \sum_{j=0}^{s-1} (1+Y)^j. \tag{D.37.20}$$

The second module coordinate of g is $(1+X)^r(S_s(Y) + (1+Y)^s Xp)$. Commuting with x requires its product by X to vanish modulo I^6. Setting $X = 0$ in the coefficient condition shows that all coefficients of $S_s(Y)$ through degree four vanish modulo 5^n. The constant coefficient first forces $5^n \mid s$. If $s = 5^n t$, $0 \le t < 5$, its degree-four coefficient is

$$\binom{5^n t}{5} \equiv 5^{n-1} t \pmod{5^n}. \tag{D.37.21}$$

Hence $s = 0$. The remaining second-coordinate equation is $X^2 p = 0 \pmod{I^6}$, which forces p to be homogeneous of degree four. Conversely such c^p is central. This proves the exact equality

$$C_{P_n}(x) = \langle x \rangle \gamma_6 P_n. \tag{D.37.22}$$

Intersecting with the centralizer of y also proves $Z(P_n) = \gamma_6 P_n$.

Suppose β_Q is an inner automorphism. Since it fixes x, (D.37.22) reduces that inner automorphism to conjugation by x^r. Its effect on y is

$$[x^r, y] = c^{S_r(X)}. \tag{D.37.23}$$

For this to have degree four, the lower coefficients force $r = 5^n t$; its coefficient is then $5^{n-1} t X^4$ modulo I^5. On the other hand $\Delta_Q = c^{-Y(X+Y)Q}$ has coefficient divisible by Y. Equality forces $t = 0$, followed by $Q = 0$. This accounts explicitly for the inner powers of x that are trivial in lower degrees but can act in degree five. It proves the ordinary outer injectivity in (D.37.2).

Symmetry and executable finite comparisons. The actual source substitutions

$$(x, y) \longmapsto (y, x), \qquad (x, y) \longmapsto (y, (xy)^{-1}) \tag{D.37.24}$$

act on P_n by (D.37.12) and have orders two and three. Conjugate β_Q by either substitution. Its displacement of x is $c^{r(X,Y)}$, with r homogeneous of degree four and divisible by X. Conjugating further by the actual element $c^{r/X}$ removes this displacement. The remaining automorphism is

$$\beta_{Q(Y,X)} \quad \text{or} \quad \beta_{Q(Y,-X-Y)}, \tag{D.37.25}$$

respectively. These identities follow directly from (D.37.14)–(D.37.18); every displacement lies in the central sixth layer, so terms of higher degree in the source substitutions vanish. They are the exact finite-group comparison with the symmetric-square action in D.21, with its inner correction retained. The line $Q = X^2 + XY + Y^2$ is the normalized gamma polynomial line. This identifies the polynomial line; it does not assert that its primitive finite parameter is realized by a Galois element.

The program `a_weight5_characteristic.py` and its saved output and JSON certificate execute the constructions for $n = 1, 2$. They check all 625 and 15625 distinct diagonal elements, all fifteen actual commutator-module words, all three special maps and their third peripheral conjugators, and the complete 125-parameter injection at $n = 1$. The finite

centralizer exponent conditions and both geometric symmetry corrections are checked as well. The resulting orders are

$$|P_1| = 19073486328125, \qquad |P_2| = 14551915228366851806640625. \tag{D.37.26}$$

The proof above supplies every precision n; it does not extrapolate a group or kernel identity from these two runs. In particular the maps $\mathrm{Out}^\sharp(\widehat{F}) \to \mathrm{Out}(P_n)$ now have fully specified characteristic targets and fully specified order-five test elements at $n = 1$. Deciding which compatible parameters lift through all finer finite quotients remains the full-source bridge. □

4.20 Class-preserving weight-five shadows and the odd-base A_6 comparison

Statement D.38 (Proved). Exact constraints on the first mixed tests above the characteristic five-group. *Every automorphism β_Q of P_1 constructed in D.37 is class-preserving. There is an explicit conjugator for every element, given in* (D.38.3) *below. The nonzero parameters nevertheless retain their nonzero ordinary outer classes by the exact centralizer calculation of D.37.*

In addition, let

$$1 \longrightarrow A_6^{\Omega} \longrightarrow E \xrightarrow{\pi} D \longrightarrow 1 \tag{D.38.1}$$

be a finite chief extension with D of odd order. Fix an epimorphism $q_E : \widehat{F} \twoheadrightarrow E$, write $X_i = q_E(x_i)$, $q = \pi q_E$, $d_i = q(x_i)$, and let $q_1 : \widehat{F} \twoheadrightarrow D$ be any epimorphism with $q_1(x_i) = h_i d_i^{\lambda} h_i^{-1}$, for specified $h_i \in D$ and $\lambda \in \widehat{\mathbf{Z}}^{\times}$. For any lifts H_i of h_i, there is an epimorphism $f : \widehat{F} \twoheadrightarrow E$ satisfying

$$\pi f = q_1, \qquad f(x_i) = a_i H_i X_i^{\lambda} H_i^{-1} a_i^{-1} \quad (a_i \in A_6^{\Omega}). \tag{D.38.2}$$

The number of these literal epimorphisms is the same as in the original case $(q_1, \lambda, h_i) = (q, 1, 1)$. The comparison also holds in the actual coupled pullback with any finite odd-order group over D, retaining its prescribed peripheral epimorphism and conjugators. In particular every β_Q passes every such chief step over its fixed characteristic target P_1.

Proof. We first prove the elementwise conjugacy assertion in the actual Magnus group, with the notation and multiplication of D.37. Let $g \in P_1$ have abelian coordinates r, s modulo five, and write $Q = AX^2 + BXY + CY^2$. If $s = 0$, then $\beta_Q(g) = g$ by D.37.16. Suppose $s \neq 0$, and put

$$\begin{aligned} d &= -sY(X+Y)Q, \\ t &= d(s,-r)/s \in \mathbf{F}_5, \\ p &= \frac{tsX^4 - d}{rX + sY} \in \mathbf{F}_5[X,Y]_3, \\ u_g &= x^{5t} c^p. \end{aligned} \tag{D.38.3}$$

The quotient displayed here is an exact polynomial division. Its homogeneous degree-four numerator vanishes at $(s, -r)$: the first term evaluates to $ts^5 = ts = d(s,-r)$. The kernel of substitution $Y = -rX/s$ is the principal ideal $(rX + sY)$, and the coefficient s is a unit. This proves existence and uniqueness of the cubic p.

The following two commutators are exact in P_1:

$$[x^5, g] = c^{sX^4}, \qquad [c^p, g] = c^{-(rX+sY)p}. \tag{D.38.4}$$

For the first, $[x, g]$ belongs to the polynomial derived module and its constant coefficient is s. The product formula for $[x^5, g]$ multiplies this coefficient polynomial by $1 + (1 + X) + \cdots +$

$(1+X)^4 = X^4$ in characteristic five. Every positive-degree term then disappears at the cutoff I^5. For the second, conjugation by g on that module multiplies by its diagonal a_g; the linear part of $a_g - 1$ is $rX + sY$, and the terms of degree at least two kill the cubic p at the same cutoff. Both commutators in (D.38.4) are central. Consequently

$$[u_g, g] = c^{tsX^4-(rX+sY)p} = c^d, \qquad u_g g u_g^{-1} = gc^d = \beta_Q(g). \tag{D.38.5}$$

This proof includes all higher abelian coordinates and all fifteen derived coordinates of g; none was discarded from the group. The term x^{5t} is essential. For example, for $Q = X^2$, $(r, s) = (1, 2)$, the value $d(2, -1)$ is nonzero, so a conjugator from the cubic derived module alone cannot give the displayed central change.

The program `a_weight5_class_preserving.py` evaluates (D.38.3)–(D.38.5) with actual Magnus multiplication for all 125 parameters and all 25 abelian residue pairs, retaining higher abelian and derived coordinates. All 3,125 checks pass; 1,200 of them have nonzero displayed x^5 coefficient.

We next give the complete odd-base chief comparison. Set $S = A_6$. The actual automorphism enumeration of D.31 supplies $Z(S) = 1$, simplicity, and $\mathrm{Out}(S) = \mathbf{F}_2^2$. Let $\rho : D \to \mathrm{Sym}(\Omega)$ be the transitive factor action of (D.38.1). Its outer action is $(b_d, \rho(d))$, where $b_d \in (\mathbf{F}_2^2)^\Omega$ and $b_{de} = b_d + \rho(d)b_e$. The explicit vector

$$B = |D|^{-1} \sum_{e \in D} b_e, \qquad b_d = B - \rho(d)B \tag{D.38.6}$$

removes this twisting by componentwise actual automorphisms of S having the prescribed outer coordinates. The centreless pullback

$$E \longrightarrow \mathrm{Aut}(S^\Omega) \times_{\mathrm{Out}(S^\Omega)} D, \qquad e \longmapsto (\mathrm{Ad}_e, \pi(e)) \tag{D.38.7}$$

is an isomorphism: its kernel is $Z(S^\Omega) = 1$, and each required automorphism differs from a chosen lift by a unique inner automorphism. After (D.38.6), the unique coefficient a_e in $\tau_B^{-1} \mathrm{Ad}_e \tau_B = \mathrm{Inn}(a_e)\rho(\pi(e))$ gives the actual isomorphism $e \mapsto (a_e, \pi(e))$ with $S^\Omega \rtimes D$. It transports the source and all specified conjugators.

Choose a factor stabilizer $D_* \le D$. For every $D_* \le C \le D$, use the two actual subgroups $H_C = q^{-1}(C)$, $H'_C = q_1^{-1}(C)$, with respective kernels $K = \ker q$, $K' = \ker q_1$. The exact boundary bijection is

$$Cd\langle q_1(x_i)\rangle \longmapsto Cdh_i\langle d_i\rangle. \tag{D.38.8}$$

It preserves each cycle length and every intermediate boundary degree, because $q_1(x_i)h_i = h_i d_i^\lambda$. The induced cycle-product class is the λth power of the old class. Kernel components of H_i only conjugate that cycle product in S. For a boundary δ of H_C, let C_γ be its specified classes upstairs and r_γ their relative boundary degrees. The exact allowable set is

$$T_{C,\delta} = \bigcap_{\gamma \mapsto \delta} \{s \in S : s^{r_\gamma} \in C_\gamma\}. \tag{D.38.9}$$

The set on the new cover is $T^\lambda_{C,\delta}$, with the labels matched by (D.38.8); this equality retains all roots when a boundary degree is not invertible on S.

All unit-power class permutations of A_6 are realized by one actual automorphism τ_λ: use the identity for $\lambda \equiv 1, 4 \pmod 5$, and conjugation by (01) in S_6 for $\lambda \equiv 2, 3 \pmod 5$. The only moved classes are the two classes of five-cycles; the other cycle types each form one A_6-class. Thus $\tau_\lambda(T) = T^\lambda$ for every conjugacy-invariant subset $T \subset S$. The program `a_weight5_a6_odd.py` verifies this for all sixteen units modulo $\exp(A_6) = 60$, on all 360 actual permutations, and locates the literal odd conjugation in the certified D.31 automorphism table.

The two covers have equal genus and matched oriented boundaries. The adjacent Hurwitz maps and inverses displayed in D.29.13 therefore give an actual isomorphism $u_C : H'_C \to H_C$ with (D.38.8) as its boundary correspondence. The map

$$\psi \longmapsto \tau_\lambda \psi u_C \tag{D.38.10}$$

is a bijection on epimorphisms to S satisfying the sets (D.38.9). Denote the common count by N_C.

Every epimorphism $\varphi : H_{D_*} \to S$ maps K onto S: its normal image is either S or trivial, and the latter would give an even-order quotient of the odd-order group D_*. If $N = \ker(\varphi|_K)$, an overgroup H_C normalizes N precisely when φ extends to H_C. Indeed its conjugation action on $K/N \simeq S$ has outer part factoring through the odd-order group C, hence trivial in $\mathbf{F}_2^2$. The inverse of $S \xrightarrow{\mathrm{Inn}} \mathrm{Inn}(S)$ supplies the extension. It agrees with φ and is unique by centrelessness and surjectivity on K. Conversely each epimorphism from H_C restricts onto S on K, so every extension is counted exactly once. Repeating the argument with K' proves that the numbers with minimal kernel normalizer agree, by the explicit finite subgroup-poset formula

$$e_{D_*} = \sum_{C \geq D_*} \mu(D_*, C) N_C. \tag{D.38.11}$$

The original generating source makes this number positive. The nonabelian Shapiro map of D.28.13 identifies the coordinate kernels with the conjugates of N; minimal normalizer makes them pairwise distinct. Surjective maps to a simple group with distinct kernels have full product image: a proper subdirect product would identify two factors by an automorphism and hence identify their kernels. This proves generation of the new full kernel. The literal gauge normalization of D.29.16, with $S = A_6$, has exactly $360^{|\Omega|-1}$ choices, and retains the specified base conjugators. The exact literal count is therefore

$$360^{|\Omega|-1} \sum_{C \geq D_*} \mu(D_*, C) N_C, \tag{D.38.12}$$

which is unchanged under (D.38.10).

For a coupled odd-order group $L \to D$, pair this lift with its prescribed new map to L, using the images of its conjugators as the h_i. The paired conjugators are actual elements of the fibre product. The possible missing quotient of S^Ω would be a quotient of $\ker(L \to D)$, of odd order. Every nontrivial quotient of S^Ω contains an S-factor and has even order, so no such quotient occurs. The paired map is therefore onto. When L is a five-group, the projection to L is the exact maximal pro-five quotient, since the perfect kernel has trivial image in every finite five-group. □

The single-kernel comparison in D.33 concerns odd-order automorphisms over arbitrary finite bases, while D.34 and D.35 use cyclic or two-group outer images of a centreless kernel. The comparison in D.38 instead handles every transitive A_6-factor permutation action over an odd-order base, arbitrary prescribed peripheral base epimorphisms, and all unit multipliers. Its maps (D.38.6) and (D.38.11) explicitly use oddness of the base. D.36 has an even base and its ordinary obstruction therefore remains in the further mixed layers that D.38 does not decide.

4.21 The complete peripheral fibre in a universal coprime Frattini cover

Statement D.39 (Proved). Actual automorphisms and every generating lift at the first coprime characteristic cover. *Let $q : \widehat{F} \twoheadrightarrow D$ be an epimorphism to a finite*

group, let $K = \ker q$*, and let* $p \nmid |D|$*. Put*

$$\Phi_p(K) = \overline{K^p[K,K]}, \qquad E = \widehat{F}/\Phi_p(K), \qquad M = K/\Phi_p(K), \qquad n = |D|. \tag{D.39.1}$$

Then E *is finite of order* np^{n+1}*, and* M *is the actual* D*-module of first homology of the regular pants cover. If* K *is characteristic in* $\widehat{F}$*, so is* $\Phi_p(K)$*.*

Write X_i *for the images of the three peripheral generators in* E*,* $d_i = q(x_i)$*, and* $e_i = \operatorname{ord}(d_i)$*. For every* $\alpha \in \operatorname{Aut}(D)$*,* $\lambda \in \widehat{\mathbf{Z}}^\times$*, and specified* $h_i \in D$ *with*

$$\alpha(d_i) = h_i d_i^\lambda h_i^{-1}, \tag{D.39.2}$$

there is an actual automorphism $\widetilde{\alpha}$ *of* E *covering* α *such that, for any chosen lifts* H_i *of* h_i*,*

$$\widetilde{\alpha}(X_i) = a_i H_i X_i^\lambda H_i^{-1} a_i^{-1} \quad \text{with } a_i \in M. \tag{D.39.3}$$

Every generating source lift above αq *factors through an automorphism of this same* E*.*

Let $B \subset M$ *be the boundary submodule and* $C = M/B$*. The complete fibre with the base conjugators* h_i *fixed has, after stated linear choices, the parameter set*

$$(M/M^D) \;\times\; \operatorname{Hom}_{\mathbf{F}_p[D]}(C,B) \;\times\; \operatorname{Aut}_{\mathbf{F}_p[D]}(C). \tag{D.39.4}$$

Its literal cardinality is

$$p^{n-1}|\operatorname{Hom}_{\mathbf{F}_p[D]}(C,B)||\operatorname{Aut}_{\mathbf{F}_p[D]}(C)|. \tag{D.39.5}$$

Allowing every ordinary base conjugator multiplies this cardinality by

$$\prod_{i=1}^{3} \frac{|C_D(d_i)|}{e_i}. \tag{D.39.6}$$

These formulas count literal generating source maps, or equivalently actual automorphisms of E*, before taking any outer quotient. In particular all D.37 parameters have the explicitly described nonempty full fibre at each first universal elementary-*p *cover with* $p \neq 5$*.*

Proof. Use the left D-modules of the right Cayley graph of D with its two directed generator edges:

$$C_1 = \mathbf{F}_p[D]e_x \oplus \mathbf{F}_p[D]e_y, \qquad C_0 = \mathbf{F}_p[D], \qquad \partial(de_x) = dd_1 - d, \quad \partial(de_y) = dd_2 - d. \tag{D.39.7}$$

Every based word w has its literal edge chain $v(w)$, with

$$v(uv) = v(u) + q(u)v(v), \qquad \partial v(w) = q(w) - 1. \tag{D.39.8}$$

Consequently

$$w \longmapsto (q(w), v(w)), \qquad (d,v)(e,w) = (de, v + dw) \tag{D.39.9}$$

is an actual finite quotient map of $\widehat{F}$. Choose a spanning tree and based words t_d for its vertices. The non-tree Schreier words $t_d x_j t_{dd_j}^{-1}$ give a free basis of the finite-index subgroup of the discrete free group. Their $n+1$ edge cycles form a basis of $\ker \partial$. The same words freely generate the open free profinite subgroup K, by the finite-coset Schreier rewriting and its inverse on the two generators. Thus their map to $\ker \partial$ has kernel exactly $\Phi_p(K)$. This proves

$$M = \ker \partial, \qquad 0 \longrightarrow M \longrightarrow C_1 \xrightarrow{\partial} C_0 \xrightarrow{\text{aug}} \mathbf{F}_p \longrightarrow 0 \tag{D.39.10}$$

with the actual source map (D.39.9), and proves the asserted finiteness and order.

Here is an explicit splitting of this extension. Put $b_d = v(t_d)$ and

$$\kappa = n^{-1} \sum_{d \in D} b_d \in C_1, \qquad s(d) = (d, \kappa - d\kappa). \tag{D.39.11}$$

The vector $\kappa - d\kappa - b_d$ belongs to M, since its boundary is zero. Hence $s(d)$ belongs to the actual image in (D.39.9). Direct multiplication gives $s(d)s(e) = s(de)$. In these coordinates the exact isomorphism

$$E \longrightarrow M \rtimes D, \qquad (d, v) \longmapsto (v - \kappa + d\kappa, d) \tag{D.39.12}$$

has inverse $(m, d) \mapsto (d, m + \kappa - d\kappa)$. This realizes the splitting without a change of the source peripheral elements.

We next retain the complete boundary module. For $i = 1, 2, 3$, let

$$b_i = v(x_i^{e_i}) = X_i^{e_i} \in M. \tag{D.39.13}$$

In the second equality M is written additively. The cycles db_i are indexed by the right cosets $d\langle d_i \rangle$, with the left deck action. Filling the boundary components of the connected oriented pants cover gives a closed connected oriented surface. Its cellular two-cycles have rank one, generated by the oriented fundamental class. Therefore the entire space of relations among these boundary cycles is exactly their sum:

$$0 \longrightarrow \mathbf{F}_p \xrightarrow{1 \mapsto \Delta} \bigoplus_{i=1}^{3} \mathbf{F}_p[D/\langle d_i \rangle] \longrightarrow B \longrightarrow 0, \qquad \Delta = \sum_{i, d\langle d_i \rangle} [d\langle d_i \rangle]. \tag{D.39.14}$$

One may also obtain this sequence directly by attaching one two-cell to each literal boundary cycle in (D.39.7); each directed edge then occurs with both opposite induced orientations. The quotient $C = M/B$ is the homology of that compact surface. In particular, with $b = \sum_i n/e_i$,

$$\dim M = n + 1, \quad \dim B = b - 1, \quad \dim C = n + 2 - b. \tag{D.39.15}$$

Let $\bar{\lambda} \in \mathbf{F}_p^{\times}$ be the reduction of λ. The prescribed map on the boundary cycles is

$$T_B(db_i) = \bar{\lambda}\, \alpha(d) h_i b_i. \tag{D.39.16}$$

It is well defined: replacing d by dd_i^j changes the right side by $\alpha(d_i)^j h_i = h_i d_i^{\lambda j}$, which fixes b_i. For each i, the map

$$d\langle d_i \rangle \longmapsto \alpha(d) h_i \langle d_i \rangle \tag{D.39.17}$$

is a bijection, with inverse obtained from α^{-1} and h_i^{-1}. Thus it sends Δ to $\bar{\lambda}\Delta$ and induces an invertible α-semilinear map $T_B : B \to B$. All boundary path choices in this step are specified by the actual coset map (D.39.17).

Because $p \nmid n$, averaging any projection over D gives a D-linear projection. Every finite $\mathbf{F}_p[D]$-module is therefore semisimple. Splitting (D.39.10) gives, by cancellation of simple summands,

$$M \simeq \mathbf{F}_p \oplus \mathbf{F}_p[D]. \tag{D.39.18}$$

The regular-module isomorphism $d \mapsto \alpha(d)$ shows that its α-twist has the same simple multiplicities. Since T_B already identifies the two boundary summands, cancellation identifies their complementary summands as well. It follows that T_B extends to an invertible α-semilinear map

$$T : M \longrightarrow M, \qquad T(dm) = \alpha(d) T(m). \tag{D.39.19}$$

This is an effective finite construction: form the matrices (D.39.7)–(D.39.17), solve the linear intertwining and boundary equations, and enumerate their finite solution space until a matrix of nonzero determinant is found. The preceding semisimple argument proves that this enumeration terminates.

The actual group automorphism is now

$$(m, d) \longmapsto (Tm, \alpha(d)) \quad \text{in (D.39.12).} \tag{D.39.20}$$

Its inverse uses T^{-1} and α^{-1}. Set $U_i = H_i X_i^{\lambda} H_i^{-1}$ and $Y_i = \widetilde{\alpha}(X_i)$. Their projections agree by (D.39.2), and

$$Y_i^{e_i} = Tb_i = \bar{\lambda} h_i b_i = U_i^{e_i} \tag{D.39.21}$$

by (D.39.16). Write $Y_i = v_i U_i$, $v_i \in M$, and let A_i be the action of $\alpha(d_i)$ on M. Equation (D.39.21) says $(1 + A_i + \cdots + A_i^{e_i - 1})v_i = 0$. The explicit kernel conjugator is

$$a_i = -e_i^{-1} \sum_{j=0}^{e_i - 1} j A_i^j v_i. \tag{D.39.22}$$

Indeed $(1 - A_i) \sum_j j A_i^j = (1 + \cdots + A_i^{e_i - 1}) - e_i$, so $(1 - A_i) a_i = v_i$. Consequently $a_i U_i a_i^{-1} = Y_i$, which proves (D.39.3) with every chosen H_i retained.

We now prove that these constructions parameterize the complete generating fibre. Any homomorphism $f : \widehat{F} \to E$ with $\pi f = \alpha q$ maps K into the elementary abelian group M. It therefore kills $\overline{K^p[K, K]}$ and factors through the actual quotient in (D.39.1). It is onto precisely when the resulting endomorphism of finite E is an automorphism. Such an automorphism restricts to an invertible α-semilinear T on M. In (D.39.12) its value on $s(d)$ has the form $(z(d), \alpha(d))$, where

$$z(de) = z(d) + \alpha(d) z(e). \tag{D.39.23}$$

Summing over e gives $z(d) = v - \alpha(d)v$, with $v = n^{-1} \sum_e z(e)$. Thus every automorphism is uniquely $\mathrm{Inn}(v)$ followed by (D.39.20), with v taken modulo M^D. Raising the three peripheral identities to their e_ith powers shows that the fixed-conjugator condition is exactly $T|_B = T_B$; its sufficiency was proved in (D.39.21)–(D.39.22).

Choose a D-linear complement C_0 to B in M, identify it with $C = M/B$, and choose one extension T_0 in (D.39.19). Every other extension is uniquely $T_0 R$, where in this chosen direct sum

$$R = \begin{pmatrix} 1 & u \\ 0 & A \end{pmatrix}, \qquad u \in \mathrm{Hom}_{\mathbf{F}_p[D]}(C, B), \quad A \in \mathrm{Aut}_{\mathbf{F}_p[D]}(C). \tag{D.39.24}$$

This proves the actual parameterization (D.39.4). By (D.39.18), $\dim M^D = 2$, so $|M/M^D| = p^{n-1}$, which proves (D.39.5).

Finally every ordinary base conjugator is $h_i c_i$, with $c_i \in C_D(d_i)$. The new boundary map is determined by the three vectors $c_i b_i$. The stabilizer in D of b_i is exactly $\langle d_i \rangle$. To verify this in the actual module, use (D.39.14): two distinct individual boundary basis vectors cannot become equal modulo the one all-boundary sum, since there are at least three boundary components. Thus the distinct maps on the ith family number $|C_D(d_i)|/e_i$. The three families may be chosen independently; their sole relation, Δ, is preserved by every such choice. Distinct resulting T_B's give disjoint automorphism fibres, each of cardinality (D.39.5). This proves the complete ordinary union formula (D.39.6).

If D is a five-group and $p \neq 5$, the projection $E \to D$ is its maximal pro-five quotient: the normal p-kernel has trivial image in every finite five-group. For $D = P_n$, D.37 supplies the characteristic kernel, and characteristicness of $\Phi_p(K)$ follows from its displayed definition. This proves the fixed-source application stated above. □

The program `a_coprime_frattini.py` constructs the actual Cayley chains, all Schreier basis words, the boundary sum and its full rank, the splitting (D.39.11), the semilinear matrix and its inverse, and all three conjugators (D.39.22). It checks 28 cases over C_5 and C_5^2 in characteristics two and three, and over S_3 in characteristic five, including all twelve base-centralizer triples in the latter case. The separate `a_coprime_frattini_count.py` enumerates the entire source fibre in actual groups of orders 48 and 54. For the genus-one C_3-cover in characteristic two it finds sixteen weak maps and twelve epimorphisms; for the genus-zero C_3-cover it finds twelve maps, all onto, with three distinct ordinary boundary maps. For the C_2-cover in characteristic three it finds six maps, all onto, with two distinct ordinary boundary maps. Every multiplier tested agrees with (D.39.5)–(D.39.6). All source files, matrices and exact outputs are retained in `code/`.

4.22 Equivariant intersection forms at every coprime prime-adic precision

Statement D.40 (Proved). Pairing-compatible lifts through the universal abelian coprime covers. *Retain the actual source $q : \widehat{F} \twoheadrightarrow D$, its kernel K, and the specified data (α, λ, h_i) of D.39, and assume that D has odd order and $p \nmid |D|$. For every $a \geq 1$ put*

$$E_{p,a} = \widehat{F}/\overline{[K,K]K^{p^a}}, \qquad E_{p,\mathrm{ab}} = \varprojlim_a E_{p,a}. \tag{D.40.1}$$

There is a continuous peripheral automorphism of $E_{p,\mathrm{ab}}$ above α, with the prescribed multiplier and base conjugators, whose compatible automorphisms of all $E_{p,a}$ induce λ_p-similitudes of the compact cover intersection form over $\mathbf{Z}_p$. The same maps preserve the full D-equivariant group-ring-valued intersection form through the coefficient automorphism induced by α. They are compatible with restriction and transfer on all intermediate covers of this regular cover.

In particular every one of the D.37 weight-five parameters passes the equivariant intersection-form test at $p = 2$ and $p = 3$, at every prime-adic precision. At $p = 2$ the induced map also preserves the unique D-invariant quadratic refinement on compact homology modulo two.

Proof. Use the same integral Cayley chains and boundary cycles as in D.39. With coefficients in $\mathbf{Z}_p$, let

$$M_p = H_1(K, \mathbf{Z}_p), \qquad B_p = \langle \text{oriented boundary cycles} \rangle, \qquad C_p = M_p/B_p. \tag{D.40.2}$$

Here $H_1(K, \mathbf{Z}_p)$ means the maximal abelian pro-p quotient of the actual open free profinite group. The Schreier maps identify it with the integral edge-cycle lattice of rank $|D| + 1$. The boundary sum of D.39.14 is its only boundary relation over $\mathbf{Z}_p$ as well. The quotient C_p is the homology of the closed oriented compactification of that cover, so it is free, and its intersection form

$$J : C_p \times C_p \longrightarrow \mathbf{Z}_p \tag{D.40.3}$$

is perfect alternating and D-invariant. Thus B_p is saturated in M_p. The averaging operator $|D|^{-1} \sum_{d \in D} d$ is defined over $\mathbf{Z}_p$; it makes both the boundary sequence and any chosen projection onto B_p D-equivariantly split.

There is no invariant compact summand:

$$C_p^D = 0, \qquad (C_p/p)^D = 0. \tag{D.40.4}$$

For the first equality, the invariant part of the graph sequence has M_p^D of rank two: augmentation on $\mathbf{Z}_p[D]^D$ is multiplication by the unit $|D|$. The invariant part of the

boundary sequence has B_p^D of rank $3-1=2$. The quotient invariants are exact by averaging and form a torsion-free quotient C_p^D; their rank is zero. The same calculation directly over $\mathbf{F}_p$ proves the second equality.

D.39 gives an α-semilinear isomorphism $L : C_p/p \to C_p/p$. Both forms

$$(v,w) \longmapsto J(Lv,Lw), \qquad (v,w) \longmapsto \bar{\lambda} J(v,w) \tag{D.40.5}$$

are nondegenerate D-invariant alternating forms on this same module. Broto–Møller–Oliver, Lemma A.1(b), applies over every finite field, including characteristic two: on a self-dual module with zero invariants for an odd group of coprime order, such forms are equivariantly isometric [BMO12]. The hypotheses are supplied by (D.40.3)–(D.40.4). Choose the resulting actual $R \in \mathrm{Aut}_{\mathbf{F}_p[D]}(C_p/p)$ and replace L by LR. It then satisfies

$$L^{\mathsf{t}} J L = \bar{\lambda} J. \tag{D.40.6}$$

This is a finite effective choice: enumerate the commutant of the two deck-action matrices and test the displayed matrix equation. The stated theorem proves termination.

For precision, the absence of an absolutely self-dual nontrivial constituent can also be seen directly in every characteristic at issue. If U were such a constituent over an algebraic closure, then $(U \otimes U)^D$ would be one-dimensional. The flip acts on it by a nonzero scalar ϵ, with $\epsilon^2 = 1$. On the other hand, if e_D is the averaging idempotent, then

$$\mathrm{tr}(\mathrm{flip}\, e_D) = |D|^{-1} \sum_{d \in D} \mathrm{tr}(\rho(d)^2) = |D|^{-1} \sum_{d \in D} \mathrm{tr}(\rho(d)) = 0. \tag{D.40.7}$$

Squaring permutes an odd-order group, and the final average projects onto $U^D = 0$. This contradicts the trace ϵ on the one-dimensional image of e_D. Consequently the nontrivial self-dual finite-field blocks have a nontrivial involution on their endomorphism fields; the finite-field Hermitian norm gives their isometry comparison. The paired non-self-dual blocks have the usual evaluation pairing. This explains in particular why characteristic two introduces no symmetric nonalternating block in (D.40.5).

We now lift (D.40.6) integrally by explicit equations, without division by two. Lift its matrix arbitrarily to $\mathbf{Z}_p$ and average:

$$T_1 = |D|^{-1} \sum_{d \in D} \rho(\alpha(d))\, \widetilde{L}\, \rho(d)^{-1}. \tag{D.40.8}$$

It is exactly α-semilinear, reduces to L, and is invertible. Suppose T_a already satisfies $T_a^{\mathsf{t}} J T_a = \lambda_p J \pmod{p^a}$. The discrepancy

$$\Delta_a = \frac{\lambda_p J - T_a^{\mathsf{t}} J T_a}{p^a} \pmod{p} \tag{D.40.9}$$

is an alternating D-invariant matrix. Let U_a be its strictly upper triangular part. Then $U_a - U_a^{\mathsf{t}} = \Delta_a$, including at $p = 2$. With $J_0 = \bar{\lambda} J \bmod p$, set

$$H_0 = J_0^{-1} U_a, \qquad \overline{H}_a = |D|^{-1} \sum_{d \in D} \rho(d) H_0 \rho(d)^{-1}. \tag{D.40.10}$$

The identity $H_0^{\mathsf{t}} J_0 + J_0 H_0 = \Delta_a$ follows from $J_0^{\mathsf{t}} = -J_0$. Averaging retains that identity because both J_0 and Δ_a are invariant. Lift $\overline{H}_a$ arbitrarily and average once more over $\mathbf{Z}_p$ to obtain a matrix H_a commuting exactly with D. Put

$$T_{a+1} = T_a(1 + p^a H_a). \tag{D.40.11}$$

This retains semilinearity and corrects the form modulo p^{a+1}; the quadratic error is divisible by p^{2a}, hence by p^{a+1}. The matrices converge to an invertible α-semilinear map T_C with

$$T_C^{\mathsf{t}} J T_C = \lambda_p J \quad \text{over } \mathbf{Z}_p. \tag{D.40.12}$$

On the boundary lattice use exactly

$$T_B(db_i) = \lambda_p \alpha(d) h_i b_i. \tag{D.40.13}$$

The actual coset bijection D.39.17 and the single diagonal relation show that this is an integral semilinear isomorphism. Choose a D-linear section $C_p \to M_p$ by averaging any section. Combine T_C on that summand with T_B on B_p. This gives an invertible semilinear T_M on M_p, retaining all prescribed boundary frames. The splitting D.39.11–12 is defined with the same integral tree chains over $\mathbf{Z}_p$. Thus

$$(m, d) \longmapsto (T_M m, \alpha(d)) \quad \text{on } M_p \rtimes D \tag{D.40.14}$$

is an actual continuous automorphism of $E_{p,\mathrm{ab}}$. Its three peripheral comparisons are the same equal-power identities as D.39.21; the norm conjugator D.39.22 is now evaluated over $\mathbf{Z}_p$, where every e_i is a unit. This proves the claimed exact peripheral identities and their compatibility in all finite quotients (D.40.1).

To retain the full deck-group information, define the equivariant intersection form by

$$\mathcal{I}(v, w) = \sum_{d \in D} J(v, dw)\, d \in \mathbf{Z}_p[D]. \tag{D.40.15}$$

Its coefficient at 1 is the original intersection form. Semilinearity and (D.40.12) give the exact comparison

$$\mathcal{I}(T_C v, T_C w) = \lambda_p\, \alpha_* \mathcal{I}(v, w), \qquad \alpha_*\Big(\sum a_d d\Big) = \sum a_d \alpha(d). \tag{D.40.16}$$

For an intermediate subgroup $H_C = q^{-1}(C)$, the covering transfer identifies its open homology with M_p^C, because $|C|$ is a unit. The transfer of a boundary circle is the sum of its lifted boundary circles; projection back multiplies it by the degree after their actual ramification indices are included. All these indices divide $|C|$ and are units. Taking the boundary quotient therefore identifies its compact homology with C_p^C as well. The identities

$$T_M \sum_{c \in C} c = \Big(\sum_{c \in C} \alpha(c)\Big) T_M, \qquad |\alpha(C)| = |C| \tag{D.40.17}$$

prove the stated simultaneous restriction and transfer comparison, including the degree factors in the intersection form.

At $p = 2$, average any quadratic refinement of $J \bmod 2$ over the odd group D. Its polarization is still J. Any two invariant refinements differ by an invariant linear functional on $C_p/2$, which vanishes by (D.40.4). The refinement is thus unique. Since $\lambda_2 \bmod 2 = 1$, pulling it back by the semilinear isometry gives another invariant refinement with the same polarization; the two are equal. □

The independent program `a_coprime_intersection.py` computes the compact cup matrix from every literal oriented Schreier boundary word. For two cohomology coordinates χ, ψ, it maps a free generator j to $(\chi_j, \psi_j, 0)$ in the Heisenberg group. An inverse letter contributes $\chi_j \psi_j$, while multiplication adds the first coordinate times the new second coordinate. Each boundary has zero first two coordinates after passing to compact cohomology.

Their total central value is the cup pairing of the filled oriented surface, by its actual product-of-boundaries and handle relation. The saved matrices are alternating, nondegenerate and invariant under every deck element; their inverses give the intersection matrices used in (D.40.9)–(D.40.11).

The program checks the C_5 and C_5^2 covers, of compact ranks four and twelve, at both primes two and three. It constructs all four scalar base automorphisms with their full boundary maps, and corrects them through modulus $2^6 = 64$ and $3^6 = 729$. Several cases require a nonidentity finite-field correction and every subsequent Hensel correction, so the dyadic step is actually exercised. All integral Schreier words, quotient maps, cup matrices, action matrices, corrections, and resulting full homology maps are saved with the exact executed output.

4.23 The full relative pro-p cover over an odd finite base

Statement D.41 (Proved). The exact pairing-compatible abelian image and all relative pro-p peripheral lifts. *Let $q : \widehat{F} \twoheadrightarrow D$ have finite odd-order image, let $K = \ker q$, and let $p \nmid |D|$. Write $R = K^{(p)}$ for the maximal pro-p quotient of the actual open subgroup K, and put*

$$E_p = \widehat{F}/\ker(K \to R), \qquad E_{p,\mathrm{ab}} = E_p/\overline{[R,R]}. \tag{D.41.1}$$

Then R is free pro-p of rank $|D| + 1$, and $1 \to R \to E_p \to D \to 1$ is the actual source extension.

Fix $\alpha \in \mathrm{Aut}(D)$ and specified $h_i \in D$ with $\alpha(d_i) = h_i d_i h_i^{-1}$, and use multiplier one. Every pairing-compatible automorphism of $E_{p,\mathrm{ab}}$ above α, with these specified peripheral conjugators, lifts to a peripheral automorphism of E_p above the same α and with the same conjugator projections. Here pairing-compatible means that its induced map on the compact homology C_p of D.40 preserves the actual integral intersection form. Conversely every such automorphism of E_p has this property. This gives the exact image of the full relative pro-p peripheral fibre in its abelian fibre.

The fibre is nonempty for every specified (α, h_1, h_2, h_3). Every generating source map $\widehat{F} \to E_p$ above αq factors through an automorphism of this same E_p. Its full literal peripheral fibre has the exact normalizer parameterization in (D.41.22)–(D.41.24) *below. In particular all D.37 weight-five shadows admit compatible generating lifts to the full relative pro-p cover for every $p \neq 5$, with their specified characteristic five-group quotient retained.*

Proof. Put $n = |D|$, and let $b = \sum_i n/e_i$ be the number of boundary components in the regular pants cover. There are at least three such components. Finite-coset Schreier rewriting identifies K with the free profinite group of rank $n + 1$ and identifies its maximal pro-p quotient with the free pro-p group R on the same Schreier generators. The kernel of $K \to R$ is characteristic in K, hence normal in $\widehat{F}$. This proves the exact extension in (D.41.1), with the indicated source map. Its boundary generators are the literal based words

$$c_\gamma = t_d x_i^{e_i} t_d^{-1} \in R, \qquad \gamma = d\langle d_i \rangle. \tag{D.41.2}$$

Their conjugacy classes are independent of the chosen coset paths, and their homology classes are the individually labelled boundary vectors of D.39.14.

The actual boundary automorphism group and its integral image. Let $\mathcal{H}$ be the subgroup of $\mathrm{Aut}(R)$ whose elements permute the conjugacy classes of the c_γ, retaining their positive generators, rather than replacing them by unspecified unit powers. The group $\mathrm{Aut}(R)$ is compact: its image in R^{n+1} consists exactly of the tuples whose reductions form a

basis of $R/\Phi_p(R)$, a clopen condition. Such a tuple generates R; a proper closed subgroup would be contained in an open maximal subgroup, contradicting that basis condition. The resulting surjective endomorphism is an automorphism, as follows from the characteristic finite p-central quotients below. Each conjugacy class of c_γ is a compact image of R. Thus each prescribed permutation condition is closed, and their finite union $\mathcal{H}$ is a closed profinite subgroup.

The integral homology is M_p, with boundary radical B_p and compact quotient C_p. Let $\mathcal{G}(\mathbf{Z}_p)$ be the group of linear automorphisms of M_p that permute the individual boundary vectors and induce an isometry of C_p. We claim the exact equality

$$\operatorname{im}(\mathcal{H} \to \operatorname{Aut}_{\mathbf{Z}_p}(M_p)) = \mathcal{G}(\mathbf{Z}_p). \tag{D.41.3}$$

We prove both inclusions with actual maps.

For necessity, take two homology characters vanishing on the boundary vectors, with values in $\mathbf{Z}/p^a$. Send each free generator of R to $(\chi_j, \psi_j, 0)$ in the finite Heisenberg p-group. Every boundary image is central. In a surface presentation the sum of their central values is, up to the one common orientation sign, the handle cup pairing. Changing the chosen Heisenberg lift of (χ, ψ) adds a linear character; its sum on the boundaries is zero by their single homology relation. An element of $\mathcal{H}$ permutes those central boundary values and conjugates them, which does not change their sum. It therefore preserves the cup form modulo every p^a, and hence preserves the integral intersection form on C_p. This proves the inclusion in (D.41.3).

For the reverse inclusion use the actual oriented surface presentation

$$\prod_{j=1}^{g} [a_j, b_j]\, c_1 \cdots c_b = 1. \tag{D.41.4}$$

Omitting c_b gives a free basis of R. Adjacent boundary permutations are the literal Hurwitz automorphisms

$$(c_j, c_{j+1}) \longmapsto (c_{j+1}, c_{j+1}^{-1} c_j c_{j+1}), \tag{D.41.5}$$

with inverse $(c_j, c_{j+1}) \mapsto (c_j c_{j+1} c_j^{-1}, c_j)$. They preserve the product in (D.41.4) exactly, including when one boundary is the omitted generator. They realize every permutation of the oriented boundary classes.

On the handle part, a Dehn twist is the explicit homeomorphism obtained by cutting an annular neighbourhood of an oriented simple curve, rotating one boundary once, and regluing, with the surface base point outside that annulus. Its induced free-group automorphism extends to R, and its homology map is the transvection $v \mapsto v + \langle v, a\rangle a$, with the sign chosen by the direction of the twist. The curves for the standard a_j, b_j, and band sums in two handles, realize the primitive vectors $a_i, b_i, a_i + a_j, a_i + b_j$ and $b_i + b_j$, with $i \neq j$ in every cross-handle sum. The two twists within a handle can also be written literally as

$$(a, b) \mapsto (a, ba), \qquad (a, b) \mapsto (ab^{-1}, b); \tag{D.41.6}$$

both preserve $[a, b]$. These elementary symplectic transvections generate $\mathrm{Sp}_{2g}(\mathbf{Z}/p^a)$ at every a. Indeed a unimodular first column over the local ring has a unit entry; symplectic swaps, the two elementary operations on a handle, and the indicated cross-handle transvections move that entry to the first handle, scale it to one, and clear the other entries. The paired column is then cleared using the symplectic pairing, and the same operations apply to the remaining $g - 1$ handles. For complete explicitness, write $E_{12}(t) = \left(\begin{smallmatrix} 1 & t \\ 0 & 1 \end{smallmatrix}\right)$ and $E_{21}(t) = \left(\begin{smallmatrix} 1 & 0 \\ t & 1 \end{smallmatrix}\right)$. For every unit u in the local ring,

$$\begin{pmatrix} u & 0 \\ 0 & u^{-1} \end{pmatrix} = E_{12}(u-1)E_{21}(1)E_{12}(u^{-1}-1)E_{21}(-u).$$

Direct multiplication proves this identity, so the unit scaling also lifts by integer twist powers. This is an induction on g. Consequently these actual integral maps have dense image in $\mathrm{Sp}(C_p)$.

We retain explicit maps for the remaining $\mathrm{Hom}(C_p, B_p)$ part. In a three-generator subsurface word $[a, b]c$, the automorphism

$$\begin{aligned} a &\longmapsto c^{-1}ac, \\ b &\longmapsto c^{-1}a^{-1}cabc, \\ c &\longmapsto c^{-1}a^{-1}cac \end{aligned} \tag{D.41.7}$$

preserves $[a, b]c$ exactly and sends c to a positive conjugate of itself. Here is an actual inverse construction. Use the free basis

$$u = ac, \qquad v = c^{-1}bc, \qquad z = vcv^{-1}, \tag{D.41.8}$$

whose inverse is

$$c = v^{-1}zv, \qquad a = uv^{-1}z^{-1}v, \qquad b = v^{-1}zvz^{-1}v. \tag{D.41.9}$$

In that basis $[a, b]c = [u, v]z$. Conjugate the twist $v \mapsto vu$ through (D.41.8)–(D.41.9), and compose it with the inverse of $b \mapsto ba$. This gives (D.41.7). Replacing the two twists by their inverse maps in reverse order gives its inverse. Its homology action is precisely

$$b \longmapsto b + c, \qquad a \longmapsto a, \qquad c \longmapsto c. \tag{D.41.10}$$

For a boundary c_j separated from the first handle in (D.41.4) by a word W in the other generators, replace that boundary temporarily by Wc_jW^{-1}. This is an invertible basis replacement and puts the local product in the form $[a, b](Wc_jW^{-1})W$. Apply (D.41.7) and then undo the basis replacement. It preserves the full surface relation and all positive boundary classes, with homology addition $b \mapsto b + c_j$. Conjugating by the handle maps and taking integer powers gives every elementary map from the compact handle lattice to the independent $b - 1$ boundary vectors. These maps add because each vanishes on the boundary lattice. Their closure is the full group $\mathrm{Hom}_{\mathbf{Z}_p}(C_p, B_p)$.

In the surface homology basis, an element of $\mathcal{G}(\mathbf{Z}_p)$ is determined by its boundary permutation, its symplectic compact block, and this arbitrary off-diagonal homomorphism. The preceding maps therefore have dense image in $\mathcal{G}(\mathbf{Z}_p)$. The image of compact $\mathcal{H}$ is closed, proving equality (D.41.3).

The exact pro-p kernel and coprime conjugacy. The kernel of $\mathrm{Aut}(R) \to \mathrm{GL}(R/\Phi_p(R))$ is pro-p. Here is a finite argument that also proves the needed Hopfian assertion. Use the characteristic lower p-central series $R_1 = R$, $R_{j+1} = \overline{R_j^p[R_j, R]}$. Its finite quotients are cofinal among finite p-quotients: the analogous series terminates in every finite p-group. For a finite p-group U on r generators, the automorphisms inducing the identity on $U/\Phi_p(U)$ act freely on the ordered generating tuples lifting a fixed Frattini basis. There are $|\Phi_p(U)|^r$ such tuples, so the order of this automorphism kernel divides a power of p. Applying this to R/R_j proves the pro-p assertion. A surjective endomorphism of R induces automorphisms of all those characteristic finite quotients; their cofinality makes its kernel trivial. In particular

$$\mathcal{P} = \ker\bigl(\mathcal{H} \to \mathcal{G}(\mathbf{Z}_p)\bigr) \quad \text{is pro-}p. \tag{D.41.11}$$

We use the following exact coprime fact. In an extension with normal pro-p kernel and finite quotient of order prime to p, complements exist and any two are conjugate by the kernel. For a finite p-kernel, prove it inductively on its order using the nonzero characteristic

elementary abelian group $\Omega_1 Z(P)$. After the induction step in the quotient, the remaining existence and conjugacy equations are respectively $H^2(D, A) = 0$ and $H^1(D, A) = 0$ for that elementary abelian module A. They have explicit averaging solutions: for a normalized two-cocycle f, put $b(d) = |D|^{-1} \sum_e f(d, e)$; then $f(d, e) = b(d) + db(e) - b(de)$. For a crossed homomorphism z, put $v = |D|^{-1} \sum_e z(e)$; then $z(d) = v - dv$. These formulas split the extension and conjugate its two complements. For a pro-p kernel, apply the finite argument in all finite quotients and use compactness: the closed sets of compatible section lifts, or of conjugating elements in the kernel, have the finite intersection property. Their intersections give the required actual complement and conjugator.

Choose such a complement in E_p. Write its section as $s : D \to E_p$. It identifies the actual extension with

$$E_p = R \rtimes_\rho D. \tag{D.41.12}$$

The action $\rho(D)$ lies in $\mathcal{H}$, since conjugation by source lifts permutes the boundary classes of the regular cover. It is faithful already on homology modulo p, by D.39.18, which contains the regular D-module.

Making a prescribed homology lift exactly equivariant. Take any prescribed pairing-compatible automorphism of $E_{p,\mathrm{ab}}$ in the statement. Normalize its action on the finite complement by an inner element of M_p: the averaging formula for $H^1(D, M_p)$ used in D.39.23 gives this explicitly. Its homology restriction is then an α-semilinear map

$$T \in \mathcal{G}(\mathbf{Z}_p), \qquad T\rho_M(d)T^{-1} = \rho_M(\alpha(d)), \tag{D.41.13}$$

with the exact boundary permutation

$$d\langle d_i\rangle \longmapsto \alpha(d)h_i\langle d_i\rangle. \tag{D.41.14}$$

By (D.41.3) choose an actual $\sigma \in \mathcal{H}$ inducing T. In the preimage of the finite subgroup $\rho_M(D)$ under $\mathcal{H} \to \mathcal{G}(\mathbf{Z}_p)$, both $\rho(D)$ and $\sigma\rho(D)\sigma^{-1}$ are complements to the pro-p kernel $\mathcal{P}$. The coprime conjugacy just proved gives $u \in \mathcal{P}$ such that $\eta = u\sigma$ satisfies

$$\eta\rho(d)\eta^{-1} = \rho(\alpha(d)) \quad (d \in D). \tag{D.41.15}$$

Equality with these labels follows from (D.41.13) and faithfulness of the finite homology image. The correction u acts trivially on all integral homology, so η still induces the prescribed T. It also retains every boundary label: distinct boundary generators have distinct homology vectors, by the single sum relation and $b \geq 3$. Consequently

$$(r, d) \longmapsto (\eta(r), \alpha(d)) \tag{D.41.16}$$

is an actual automorphism of E_p.

Recovering the peripheral generators from their powers. This step retains the exact conjugators. Let $b_i = X_i^{e_i} \in R$ and take the specified lift $H_i \in E_p$ of h_i. By the positive boundary-class property of η and (D.41.14), there is $a_i \in R$ with

$$\eta(b_i) = a_i H_i b_i H_i^{-1} a_i^{-1}. \tag{D.41.17}$$

Set $Y_i = \widetilde{\alpha}(X_i)$ and $U_i = a_i H_i X_i H_i^{-1} a_i^{-1}$. They have the same image in D and the same e_ith power

$$Y_i^{e_i} = U_i^{e_i} = v_i. \tag{D.41.18}$$

Every boundary generator is primitive in R. This is immediate for the $b-1$ independent boundaries in (D.41.4); the last is primitive as well, because another boundary occurs exactly once in its surface word. Replacing that other boundary by the last one is a free basis replacement with inverse obtained from (D.41.4). For a primitive boundary c,

$$C_R(c) = \overline{\langle c \rangle}. \tag{D.41.19}$$

To verify the exact centralizer, take c as the first free pro-p generator and send the generators to $1 + X_j$ in $\mathbf{F}_p\langle\langle X_1, \ldots, X_{n+1}\rangle\rangle^\times$. This Magnus map is injective. Indeed any finite p-group quotient can be evaluated by substituting $X_j \mapsto g_j - 1$ in its group algebra; the augmentation ideal of a finite p-group algebra is nilpotent. The latter follows by induction using a central element of order p, whose difference from one has pth power zero. Thus equality in the Magnus algebra implies equality in every finite p-quotient, and hence in R. Comparing coefficients in $X_1 A = A X_1$ shows that the centralizer of X_1 in the power-series algebra is exactly $\mathbf{F}_p[[X_1]]$: the coefficient of a word containing another letter is shifted to one having no initial X_1, where the commutation identity forces zero. The retraction $R \to \langle c \rangle$ sets the other X_j to zero. For a commuting group element it leaves the Magnus series unchanged, and injectivity proves (D.41.19).

Now $Y_i U_i^{-1}$ lies in R and centralizes v_i. The element v_i is a conjugate of a primitive boundary generator, so this difference belongs to the procyclic group $\langle v_i \rangle$. Both Y_i and U_i fix v_i and hence commute with that entire procyclic group. Taking e_ith powers in (D.41.18) therefore gives $(Y_i U_i^{-1})^{e_i} = 1$. The group $\langle v_i \rangle \simeq \mathbf{Z}_p$ is torsion-free. It follows that

$$\widetilde{\alpha}(X_i) = a_i H_i X_i H_i^{-1} a_i^{-1}. \tag{D.41.20}$$

Thus the actual peripheral roots, and their specified base conjugators, are retained, rather than only their boundary powers.

Finally lift the abelian normalizing element used before (D.41.13) from M_p to R, and conjugate (D.41.16) by it. This gives a lift of the entire prescribed abelian automorphism, including its section-conjugator coordinate. Necessity of the intersection condition follows from (D.41.3). D.40 supplies a pairing-compatible abelian lift for every specified (α, h_i), proving nonemptiness.

All source maps and the exact full fibre. Any source homomorphism $f : \widehat{F} \to E_p$ above αq maps K into R. By the universal property of $K^{(p)}$, it kills $\ker(K \to R)$ and factors through the exact quotient (D.41.1). If f is onto, its restriction on the kernel R is onto: a preimage of an element in R must have trivial image in D, since the base map is α. The Hopfian argument above makes this restriction an automorphism, and then the whole induced map on E_p is an automorphism. The identical argument on the abelian kernel M_p applies to source epimorphisms to $E_{p,\mathrm{ab}}$.

Here is the literal full-fibre parameterization, including its remaining conjugator quotient. Let $\mathcal{H}_0$ be the subgroup of $\mathcal{H}$ fixing each positive boundary conjugacy class, and let θ_h be the permutation (D.41.14). Define the closed set

$$\mathcal{Y}_{\alpha,h} = \{\eta \in \mathcal{H} : \eta \rho(d) \eta^{-1} = \rho(\alpha(d)) \text{ for all } d, \quad \eta \text{ induces } \theta_h\}. \tag{D.41.21}$$

It is nonempty by the construction. For any one of its elements η_0,

$$\mathcal{Y}_{\alpha,h} = C_{\mathcal{H}_0}(\rho(D))\, \eta_0. \tag{D.41.22}$$

This equality follows by multiplying the two exact intertwining identities; the boundary permutations cancel. Every automorphism in the full fixed-conjugator fibre is

$$\mathrm{Inn}(r) \circ \big((s,d) \mapsto (\eta(s), \alpha(d))\big), \qquad r \in R, \quad \eta \in \mathcal{Y}_{\alpha,h}. \tag{D.41.23}$$

Indeed its image of the chosen finite complement is another complement, so coprime conjugacy normalizes it to $\rho(D)$. The ambiguity is exactly

$$(r,\eta) \sim (ra, \mathrm{Inn}(a^{-1})\eta), \qquad a \in R^D = C_R(s(D)). \tag{D.41.24}$$

These pairs give the same automorphism, and equality on the finite complement and on R proves the converse. This is a quotient of compact spaces by the displayed actual group action.

Allowing ordinary base conjugators $h_i c_i$, $c_i \in C_D(d_i)$, gives the disjoint union over the distinct boundary permutations; their number is the product in D.39.6, by the same individually distinguishable boundary vectors. This describes the entire source fibre at the relative pro-p stage, not a selected independently admitted component. When D is a five-group and $p \neq 5$, the projection $E_p \to D$ is its maximal pro-five quotient: the normal pro-p group R has trivial image in any finite five-group. This proves the exact fixed-target application to D.37. □

The finite program `a_relative_prop.py` checks (D.41.5)–(D.41.10) as literal free-word maps, with their inverses, including a swap involving the omitted boundary. It generates all 720 matrices of $\mathrm{Sp}_4(\mathbf{F}_2)$ and all 24 matrices of $\mathrm{Sp}_2(\mathbf{F}_3)$ from exactly the stated elementary transvections: eight listed vectors in genus two over $\mathbf{F}_2$ and the two handle vectors in genus one over $\mathbf{F}_3$. The complete vector and matrix lists are retained in its JSON certificate. It also checks all 432 automorphisms of the actual Heisenberg group over $\mathbf{F}_3$, whose identity-on-Frattini kernel has order nine, and exhibits the four kernel conjugators between the four complements in $\mathbf{F}_2^2 \rtimes C_3$. The infinite pro-$p$ assertions use the proved compactness and characteristic-quotient arguments, rather than extrapolation from these finite checks.

D.41 controls one entire relative pro-p kernel over the specified odd-order base. A finite quotient whose further kernel mixes this prime with additional chief factors is not supplied by the maps (D.41.1). The remaining D.1–D.2 comparison must retain those further finite quotients and the complete fibres over the pairing-compatible relative lifts just constructed.

4.24 Matched boundary circles and the coprime cover over an arbitrary finite base

Statement D.42 (Proved). The even-base intersection obstruction vanishes for multiplier one. *Let $q : \widehat{F} \twoheadrightarrow D$ be an actual finite quotient of the three-peripheral source, put $K = \ker q$, and let $p \nmid |D|$. The order of D is unrestricted. Fix $\alpha \in \mathrm{Aut}(D)$ and specified elements $h_i \in D$ satisfying*

$$\alpha(d_i) = h_i d_i h_i^{-1} \quad (1 \le i \le 3). \tag{D.42.1}$$

Use multiplier one throughout this statement. Let $M_p = K^{\mathrm{ab},p}$, let B_p be its actual boundary sublattice, and let $C_p = M_p/B_p$ have its compact-cover intersection form J. Then there is an invertible α-semilinear map of M_p that has the prescribed boundary values

$$T_B(db_i) = \alpha(d) h_i b_i \tag{D.42.2}$$

and induces an isometry of C_p. It can be chosen compatibly modulo every p^a, and its compact block preserves the complete group-ring intersection pairing, with the coefficient automorphism α.

Put

$$R = K^{(p)}, \qquad E_p = \widehat{F}/\ker(K \longrightarrow R), \qquad E_{p,\mathrm{ab}} = E_p/\overline{[R,R]}. \tag{D.42.3}$$

Every pairing-compatible peripheral automorphism of $E_{p,\mathrm{ab}}$ above α, with the specified conjugator projections h_i, lifts to a peripheral automorphism of E_p with those same projections. Conversely its compact homology map must preserve J. The complete literal fibre is nonempty and has precisely the normalizer and kernel-conjugator parameterization (D.41.22)–(D.41.24)*, with D now arbitrary subject to $p \nmid |D|$. Every generating source map above αq factors through an automorphism of this same E_p.*

Consequently every D.37 parameter has the following actual mixed-prime realization. For every finite stage of the characteristic relative pro-two cover over P_n, and every odd prime $p \neq 5$, its induced multiplier-one shadow lifts to the full relative pro-p cover of that finite stage. All these quotients retain P_n as their exact maximal pro-five quotient and retain the specified β_Q on that quotient.

Proof. The case $p = 2$ has odd-order D and is already covered by D.40–D.41. We prove the new assertion for odd p, keeping the actual cover and all its labelled boundary maps.

The closed free-action comparison. Let P be an oriented pair of pants with the based positive boundary words $x_1 = x, x_2 = y, x_3 = (xy)^{-1}$. Restriction of q to the dense discrete free group gives the connected regular topological cover $U \to P$. Its profinite fundamental group is the actual subgroup K, by the finite-coset Schreier correspondence. Use the left deck action $\rho : D \to \mathrm{Homeo}^+(U)$. Set $n = |D|$, $e_i = \mathrm{ord}(d_i)$, and

$$b = \sum_i n/e_i. \tag{D.42.4}$$

The boundary component indexed by $d\langle d_i\rangle$ is the circle with the positive generator db_i in homology.

Make a second copy U^α of the same underlying oriented surface, with the explicitly twisted action

$$d \cdot_\alpha u = \rho(\alpha(d))u. \tag{D.42.5}$$

The union of the circles above the i-th boundary has the exact model

$$(D \times \mathbf{R}/\mathbf{Z})/((dd_i, t) \sim (d, t + 1/e_i)).$$

The prescribed boundary map from the first copy to the second is

$$j_i : [d, t] \longmapsto [\alpha(d)h_i, t]. \tag{D.42.6}$$

Indeed $\alpha(dd_i)h_i = \alpha(d)h_i d_i$; this proves that the displayed quotient relation is preserved. It also proves D-equivariance with the action (D.42.5). Each j_i preserves the positive circle orientation and has an actual inverse, using α^{-1}. Its homology map is exactly (D.42.2).

Glue U to the oppositely oriented copy $-U^\alpha$ along these three unions of circles:

$$S = U \cup_j (-U^\alpha). \tag{D.42.7}$$

The positive circle identification is orientation reversing for the boundary of the second, negatively oriented piece, so S is a closed oriented surface. The action is free on both pieces, including every boundary circle, and the maps j_i are equivariant. It follows directly that the induced D-action on S is free and orientation preserving. Its quotient is the double of the pair of pants, and S is connected of genus $n + 1$.

We use the exact primary theorem of Barge–Marché: for a free orientation-preserving action of a finite group on a closed oriented surface, its integral first homology has a group-invariant primitive Lagrangian [BM24, Theorem 1, integral formulation in the author version].

Thus $H_1(S,\mathbf{Z})$ has a D-stable direct summand $L_{\mathbf{Z}}$, of half its rank, on which the intersection form vanishes. Equivalently one may first take the rational Lagrangian and intersect with integral homology; this intersection is saturated and has the same half-rank. All hypotheses concern the free action on the actual glued surface (D.42.7).

The primitive neck lattice and its exact reduction. Write $M_{\mathbf{Z}} = H_1(U,\mathbf{Z})$, and use the corresponding notation for U^α. The single relation among the b oriented boundary circles is their sum. Mayer–Vietoris gives the exact sequence

$$\begin{aligned} 0 \longrightarrow \frac{M_{\mathbf{Z}} \oplus M^\alpha_{\mathbf{Z}}}{\{(b_0, -j_* b_0) : b_0 \in B_{\mathbf{Z}}\}} &\longrightarrow H_1(S,\mathbf{Z}) \\ &\longrightarrow \ker\bigl(\mathbf{Z}^b \xrightarrow{\sum} \mathbf{Z}\bigr) \longrightarrow 0. \end{aligned} \tag{D.42.8}$$

The last map records the oriented crossing numbers through the seams. Here $B_{\mathbf{Z}} \cong \mathbf{Z}^b/\mathbf{Z}(1,\dots,1)$; the corresponding map from $H_1(\partial U,\mathbf{Z})$ to the two pieces has exactly the displayed image.

Let $I_{\mathbf{Z}}$ be the image in $H_1(S,\mathbf{Z})$ of $(B_{\mathbf{Z}},0)$. The preceding quotient identifies this image with that of $(0, j_* B_{\mathbf{Z}})$. It is an injective copy of $B_{\mathbf{Z}}$, hence has rank $b-1$. Quotienting the first term in (D.42.8) by this copy gives $C_{\mathbf{Z}} \oplus C^\alpha_{\mathbf{Z}}$; both that group and the last term of (D.42.8) are free abelian. Thus $I_{\mathbf{Z}}$ is primitive. It is isotropic because its generators are disjoint neck circles. Intersection with those circles is precisely the crossing-number map in (D.42.8): the dual of $\mathbf{Z}^b/\mathbf{Z}(1,\dots,1)$ is the displayed sum-zero lattice. This proves the equality of integral subgroups between $I^\perp_{\mathbf{Z}}$ and the image of the first term of (D.42.8), including its index. Filling the boundary circles in each of the two pieces therefore gives the exact D-equivariant isometry

$$(I^\perp_{\mathbf{Z}}/I_{\mathbf{Z}}, J_S) \;\simeq\; (C_{\mathbf{Z}}, J) \;\oplus\; (C^\alpha_{\mathbf{Z}}, -J). \tag{D.42.9}$$

The minus sign is the orientation of the second piece. The maps in this comparison are the inclusions of cycles supported in the two pieces, followed by their actual boundary-filling quotients.

Primitivity allows reduction of (D.42.9) modulo p. Put $W = H_1(S,\mathbf{F}_p)$, $I = I_{\mathbf{Z}}/p$, and $L = L_{\mathbf{Z}}/p$. Then L is a D-stable Lagrangian in W. The explicitly defined submodule

$$L' = \frac{(L \cap I^\perp) + I}{I} \tag{D.42.10}$$

is a Lagrangian in $I^\perp/I$. For the dimension check, put $\dim W = 2r$, $\dim I = s$, and $\dim(L \cap I) = t$. Perfect pairing with L gives $\dim(L \cap I^\perp) = r - s + t$; the numerator in (D.42.10) has dimension r, and its quotient has dimension $r-s$. It is isotropic by its definition. Thus the following precise finite-field obstruction vanishes:

$$\omega_\alpha = [(C_p/p, J)] - [(C^\alpha_p/p, J)] = 0 \quad \text{in } W^-(\mathbf{F}_p[D]). \tag{D.42.11}$$

This conclusion uses the integral Lagrangian of the free-action surface, with the exact reduction map, rather than a uniqueness assertion for arbitrary even-group invariant forms.

Cancellation on the specified semisimple module. We also retain the underlying module comparison. The based Cayley graph chain sequence of D.39 gives $M_p/p \cong \mathbf{F}_p \oplus \mathbf{F}_p[D]$. Twisting the regular module by α is the literal basis permutation $d \mapsto \alpha(d)$. The boundary isomorphism (D.42.2) respects the sole sum relation. Since $p \nmid |D|$, averaging splits the boundary sequence. Cancellation of simple-module multiplicities therefore gives

$$C_p/p \simeq C^\alpha_p/p \quad \text{as } \mathbf{F}_p[D]\text{-modules.} \tag{D.42.12}$$

For clarity, the finite-field Witt cancellation needed here can be proved directly. Let k be a finite field of odd characteristic prime to $|D|$, and decompose a semisimple $k[D]$-module into its simple isotypic blocks. A non-self-dual pair U, U^* is paired perfectly across its two blocks; choosing dual multiplicity bases identifies this form with its unique hyperbolic model. For a self-dual simple U, its endomorphism ring is a finite division ring, hence a finite field E. A fixed invariant duality on U induces an adjoint involution on E. More explicitly, choose a nondegenerate D-invariant k-valued form b_U on U. Its symmetric part is either nonzero, and hence nondegenerate by simplicity, or b_U itself is alternating. Thus it may be chosen with one of these two symmetries. Every invariant form on U^m has the unique description

$$b_H((u_i),(v_j)) = \sum_{i,j} b_U(u_i, H_{ij} v_j), \qquad H_{ij} \in E.$$

Transpose uses the adjoint of E for b_U. This identifies the following actual multiplicity forms: an alternating form when the simple duality is symmetric and its adjoint is trivial; a symmetric form when that duality is alternating and the adjoint is trivial; and a Hermitian form, after a fixed scalar adjustment, when the adjoint is nontrivial. These identifications send a change of multiplicity basis A to $A^* H A$. They also send a D-stable isotropic submodule to its isotropic multiplicity subspace.

Alternating multiplicity forms of a specified dimension have their symplectic bases. Hermitian forms over a finite field have orthogonal bases and a unique class of each dimension, since the norm map to its fixed field is surjective. Symmetric forms over a finite field of odd characteristic are classified by dimension and determinant modulo squares: diagonalization reduces every entry to either one or a fixed nonsquare ν. There exist $a, b \in E$ with $a^2 + b^2 = \nu$: when -1 is nonsquare this is surjectivity of the quadratic-field norm, and otherwise it follows from the factorization $(a + \sqrt{-1}b)(a - \sqrt{-1}b) = \nu$. The matrix $\left(\begin{smallmatrix} a & -b \\ b & a \end{smallmatrix}\right)$ has determinant ν and pulls the identity form back to ν times that form. It therefore removes two nonsquare diagonal entries together. This proves the claimed two-class reduction.

Now take two forms on isomorphic semisimple modules whose orthogonal difference is metabolic. In each symmetric multiplicity block of dimension m, the determinant of their difference is $(-1)^m \det(H_0)\det(H_1)$. Metabolicity makes it $(-1)^m$ times a square, so the two determinants have the same square class. The other blocks are already determined by their module dimensions. The preceding basis constructions give an isometry on every block, and their direct sum is the required D-equivariant isometry. Applying this proved cancellation to (D.42.11)–(D.42.12) yields an actual invertible matrix $\overline{T}$ with

$$\overline{T}\,\rho(d) = \rho(\alpha(d))\overline{T}, \qquad \overline{T}^{\,t} J \overline{T} = J \pmod{p}. \tag{D.42.13}$$

It can also be found by exhaustive finite linear algebra: enumerate the intertwining space and test the two displayed conditions. The existence just proved makes that finite search terminate.

The integral lift and the actual relative source. The averaging and successive corrections of D.40 now apply with $n = |D|$, a unit in $\mathbf{Z}_p$; they do not require n to be odd. Explicitly, average an arbitrary lift of $\overline{T}$ by

$$\frac{1}{n} \sum_{d \in D} \rho(\alpha(d))\, \widetilde{T}\, \rho(d)^{-1}. \tag{D.42.14}$$

Suppose the resulting intertwiner is an isometry modulo p^a. Write $\Delta = (J - T^t J T)/p^a \bmod p$, let U be its strictly upper triangular part, and set $H_0 = \overline{J}^{-1} U$. Then $H_0^t \overline{J} + \overline{J} H_0 = U - U^t = \Delta$. Average H_0 over the actual D-action, lift and average it integrally, and replace T by $T(1 + p^a H)$.

The new matrix has the same intertwining property and is an isometry modulo p^{a+1}. The corrections converge to an integral α-semilinear isometry of C_p. A D-averaged projection onto B_p then extends it together with (D.42.2) to M_p. The coefficientwise calculation of D.40 gives

$$\sum_{d\in D} J(Tv, dTw)d = \alpha_* \left(\sum_{d\in D} J(v, dw)d \right). \tag{D.42.15}$$

This proves the full group-ring assertion, with its actual coefficient map.

Here are the precise gates transferring this map to (D.42.3). The full positive-boundary automorphism group $\mathcal{H} \subset \mathrm{Aut}(R)$ has integral homology image $\mathcal{G}(\mathbf{Z}_p)$ by the actual surface maps in (D.41.3)–(D.41.10). Those maps, their compact closure, and their pro-p kernel depend only on the free pro-p surface group with its $b \geq 3$ positive boundary classes. They impose no parity condition on D. Choose a complement D in the actual extension E_p, which exists by $p \nmid |D|$, and write $E_p = R \rtimes_\rho D$. Lift the prescribed integral homology matrix to $\sigma \in \mathcal{H}$. In the inverse image of the finite group $\rho_M(D)$, the two conjugate D-actions are complements to the full-integral-homology pro-p kernel. The coprime complement argument (D.41.11)–(D.41.15) supplies a kernel correction u, without changing the entire integral homology matrix. For $\eta = u\sigma$ it gives

$$\eta\rho(d)\eta^{-1} = \rho(\alpha(d)), \qquad (r, d) \longmapsto (\eta(r), \alpha(d)). \tag{D.42.16}$$

The second map is an actual automorphism of the displayed semidirect product.

Its e_i-th peripheral powers have the conjugacy classes prescribed by the actual boundary permutation $d\langle d_i\rangle \mapsto \alpha(d)h_i\langle d_i\rangle$. As in D.41, choose the corresponding kernel conjugator a_i and a lift H_i of h_i. The image Y_i of X_i and the element $a_iH_iX_iH_i^{-1}a_i^{-1}$ have the same projection and the same e_i-th power v_i, a primitive boundary element conjugate in R. Their ratio belongs to R and centralizes v_i. The explicit Magnus and retraction argument of D.41 gives $C_R(v_i) = \langle v_i\rangle \cong \mathbf{Z}_p$. Both elements act trivially on that procyclic centralizer, so equality of their e_i-th powers forces equality of the elements themselves. Thus every specified peripheral conjugator projection is retained.

For a specified pairing-compatible automorphism of $E_{p,\mathrm{ab}}$, first conjugate its finite complement by an element of M_p, carry out (D.42.16), and lift that inner conjugation to R. This realizes exactly the given abelian automorphism. The converse follows from the boundary Heisenberg cup calculation in D.41. The entire fibre proof there uses only this exact comparison, the coprime complement conjugacy, and the free pro-p Hopfian property. Hence its formulas (D.41.22)–(D.41.24) apply unchanged. In particular each source map above αq sends K to R and kills $\ker(K \to R)$. Surjectivity restricts to a surjective endomorphism of the finitely generated free pro-p kernel; its characteristic finite quotients make it an automorphism. This verifies the claimed all-components factorization through the actual E_p.

The characteristic mixed-prime comparison with D.37. To make the last assertion completely explicit, let $q_n : \widehat{F} \to P_n$ be D.37's characteristic quotient, let $K_n = \ker q_n$, and put

$$\begin{aligned} R_2 &= K_n^{(2)}, & \mathcal{E}_2 &= \widehat{F}/\ker(K_n \to R_2), \\ R_{2,1} &= R_2, & R_{2,j+1} &= \overline{R_{2,j}^2[R_{2,j}, R_2]}, \\ D_{n,j} &= \mathcal{E}_2/R_{2,j}, & K_{n,j} &= \ker(\widehat{F} \to D_{n,j}). \end{aligned} \tag{D.42.17}$$

Every displayed kernel is characteristic in the preceding characteristic source kernel, and therefore the finite quotients $D_{n,j}$ are actual characteristic quotients of $\widehat{F}$. They are finite two-group extensions of P_n. By D.41 choose a multiplier-one peripheral automorphism of $\mathcal{E}_2$ above the specified β_Q. It induces an automorphism $\alpha_{n,j}$ of $D_{n,j}$; its peripheral conjugators are the images of its actual conjugators.

For every odd prime $p \neq 5$ the order of $D_{n,j}$ is prime to p. Apply the proved assertion to the actual quotient in (D.42.17). It supplies a peripheral automorphism of

$$\mathcal{E}_{n,j,p} = \widehat{F}/\ker\bigl(K_{n,j} \to K_{n,j}^{(p)}\bigr) \tag{D.42.18}$$

whose composition with the displayed maps to $D_{n,j}$ and P_n is $\alpha_{n,j}$ and β_Q, respectively. This is the same original source $\widehat{F}$ at every stage. Any pro-five quotient of $\mathcal{E}_{n,j,p}$ kills its normal pro-p kernel and then the normal two-group kernel of $D_{n,j} \to P_n$. It therefore factors through P_n, which itself remains a quotient. This proves the exact maximal pro-five assertion. For fixed n, j, p, the full relative pro-p automorphism supplies a compatible family on all of its finite characteristic p-central quotients. □

The even-order form ambiguity used in this proof is visible in an exact small source. The program `a_coprime_even.py` uses $\widehat{F} \twoheadrightarrow Q_8$, $x \mapsto i$, $y \mapsto j$, and derives the compact rank-four form from its six actual oriented Schreier boundary words. It constructs an explicit isotypic chart $C \simeq U \oplus U$, where U is the faithful two-dimensional quaternionic simple module, and writes the form as $H \otimes J_2$. Over $\mathbf{F}_3$ it enumerates all 18 nondegenerate invariant alternating forms, in orbits of sizes 12 and 6; over $\mathbf{F}_5$ the corresponding 100 forms have orbits 60 and 40. Every orbit and every multiplicity transporter is saved. Their determinant classes prove explicitly that general even-group invariant forms need not be isometric on the same module. For each of the four specified inner base automorphisms, the program retains the actual boundary maps and constructs the required isometry through moduli 3^6 and 5^6. Every case uses all five nonzero integral correction steps and retains the exact graph Lagrangian in $C \oplus -C^{\alpha}$. The general theorem uses the free-action gluing proof above; these computations certify its particular finite maps.

The source family (D.42.17)–(D.42.18) handles the complete coprime relative prime after a fixed finite two-primary stage. It gives no compatibility assertion as that two-primary stage varies, and no conclusion about further steps whose new prime already divides the finite base. Those comparisons remain among the finite source problems required by D.1–D.2.

4.25 The relative pro-p fibre over an infinite coprime base

Statement D.43 (Proved). Compatible relative lifts over an actual pro-prime-to-p quotient. *Let $q : \widehat{F} \twoheadrightarrow D$ be an actual profinite quotient, let $K = \ker q$, and suppose that every finite quotient of D has order prime to p. Fix $\alpha \in \mathrm{Aut}(D)$ and actual elements $h_i \in D$ such that*

$$\alpha(d_i) = h_i d_i h_i^{-1}. \tag{D.43.1}$$

For a profinite group H, write

$$O^p(H) = \bigcap_{\substack{N \triangleleft_{\mathrm{open}} H \\ H/N \text{ a finite } p\text{-group}}} N. \tag{D.43.2}$$

Then the actual source extension

$$E_p(D) = \widehat{F}/O^p(K), \qquad 1 \longrightarrow K^{(p)} \longrightarrow E_p(D) \longrightarrow D \longrightarrow 1 \tag{D.43.3}$$

has a multiplier-one peripheral automorphism above α, with peripheral conjugator projections exactly h_i.

Choose any cofinal decreasing sequence of characteristic open normal subgroups W_j of D, put $D_j = D/W_j$, $K_j = q^{-1}(W_j)$, and $E_{j,p} = \widehat{F}/O^p(K_j)$. The full literal fibre in (D.43.3) *is*

canonically the inverse limit of the complete finite-base fibres of D.42:

$$\begin{aligned} E_p(D) &\simeq \varprojlim_j E_{j,p}, \\ \mathscr{X}(D,\alpha,h) &\simeq \varprojlim_j \mathscr{X}(D_j,\alpha_j,h_j). \end{aligned} \tag{D.43.4}$$

Both comparisons retain the original map from $\widehat{F}$. The fibres are compact and the inverse limit is nonempty. Every generating source map $\widehat{F} \to E_p(D)$ above αq factors through an automorphism of this same $E_p(D)$; its three peripheral conditions are exactly those in the displayed fibre.

In particular let $K_n = \ker(\widehat{F} \to P_n)$ for the D.37 quotient. For every odd prime $p \neq 5$, every D.37 parameter has a multiplier-one peripheral realization on the single actual quotient

$$\widehat{F}/O^p(O^2(K_n)). \tag{D.43.5}$$

It induces a chosen full relative pro-two lift from D.41, and hence induces the given β_Q on P_n. Its maximal pro-five quotient is exactly P_n. The resulting automorphisms are compatible through all the finite two-primary stages and their relative p-central stages.

Proof. We first prove the residual and quotient comparisons without a finiteness assumption on K.

If H is a closed subgroup of H', then

$$O^p(H) \subset O^p(H'). \tag{D.43.6}$$

Indeed every finite p-quotient map of H', restricted to H, kills $O^p(H)$. If $f : H \twoheadrightarrow J$ is a continuous epimorphism, then

$$f(O^p(H)) = O^p(J). \tag{D.43.7}$$

One inclusion follows by pulling back every finite p-quotient of J. For the other, the quotient $J/f(O^p(H))$ is a continuous quotient of the pro-p group $H/O^p(H)$. The subgroup $f(O^p(H))$ is closed because it is a compact image, so this is an actual pro-p quotient. It must contain the defining residual $O^p(J)$ in its kernel. This proves (D.43.7) with its exact maps.

The group D is finitely generated, as a quotient of $\widehat{F}$. It has only finitely many subgroups of each given finite index: each such subgroup defines a permutation action, and the images of a fixed finite generating set give only finitely many such actions. Intersections of all open normal subgroups of index at most j therefore give a cofinal decreasing characteristic sequence W_j. In particular α induces the compatible automorphisms α_j of D_j. By the hypothesis on D, every D_j has order prime to p.

The residual equality and the first comparison. The inclusions (D.43.6) give

$$O^p(K) \subset O^p(K_k) \subset O^p(K_j) \quad (k \geq j). \tag{D.43.8}$$

Every displayed subgroup is normal in $\widehat{F}$: its ambient K_j, or K, is normal, and its residual is characteristic in that ambient group. We claim the exact equality

$$\bigcap_j O^p(K_j) = O^p(K). \tag{D.43.9}$$

To prove the remaining inclusion, take an arbitrary finite quotient G of $\widehat{F}/O^p(K)$, and let $f : \widehat{F} \twoheadrightarrow G$ be its source map. The image $N = f(K)$ is a finite p-group, because the restriction to K factors through $K^{(p)}$. There is an actual induced quotient map

$$D \longrightarrow G/N. \tag{D.43.10}$$

Its kernel is open, so some W_j is contained in that kernel. It follows that $f(K_j) \subset N$. The restriction of f to K_j then has p-group image and kills $O^p(K_j)$. Consequently the left-hand side of (D.43.9) is killed in every finite quotient of $\widehat{F}/O^p(K)$, which proves the reverse inclusion. Notice that this residual equality itself holds for any closed normal K and these open approximations; the prime-to-p assumption is used for the nonempty finite-base fibres.

The source quotient maps now give

$$\widehat{F}/O^p(K) \longrightarrow \varprojlim_j \widehat{F}/O^p(K_j). \tag{D.43.11}$$

Its kernel is trivial by (D.43.9). It is onto: each compatible tuple of cosets has nested nonempty closed inverse-image sets in compact $\widehat{F}$. Their intersection provides a simultaneous source lift. Thus (D.43.11) is an isomorphism of profinite groups, not merely an identification of their finite images.

Every upper fibre has an actual descent map. For $k \geq j$, let $\pi_{k,j} : E_{k,p} \to E_{j,p}$ be the map supplied by (D.43.8). Inside $E_{k,p}$ put

$$M_{j,k} = \ker(E_{k,p} \longrightarrow D_j) = K_j/O^p(K_k). \tag{D.43.12}$$

The last equality uses the actual image of K_j under the source map. Applying (D.43.7) to $K_j \twoheadrightarrow M_{j,k}$ gives

$$O^p(M_{j,k}) = O^p(K_j)/O^p(K_k) = \ker \pi_{k,j}. \tag{D.43.13}$$

An automorphism of $E_{k,p}$ above α_k preserves $M_{j,k}$, because α_k preserves the kernel of $D_k \to D_j$. It therefore preserves the characteristic subgroup $O^p(M_{j,k})$. Formula (D.43.13) proves that it descends to an actual automorphism of $E_{j,p}$. The descended map is above α_j; every one of its peripheral conjugators is the image of an upper conjugator. It has projection $h_{i,j}$ when the upper one has projection $h_{i,k}$.

Here and below the fibre is the literal compact set

$$\begin{aligned}\mathscr{X}_j = \{\sigma \in \mathrm{Aut}(E_{j,p}) : {} & \bar{q}_j\sigma = \alpha_j\bar{q}_j, \\ & \sigma(X_i) = a_iX_ia_i^{-1} \text{ for some } a_i \in E_{j,p} \text{ with } \bar{q}_j(a_i) = h_{i,j},\ 1 \leq i \leq 3\}.\end{aligned} \tag{D.43.14}$$

The third source element is $X_3 = (X_1X_2)^{-1}$, as throughout the paper. The preceding descent is a continuous map $\mathscr{X}_k \to \mathscr{X}_j$. The maps compose exactly because they are induced by the same source quotient maps. Thus (D.43.14) is an actual inverse system.

Compactness, nonemptiness, and the three exact conjugators. Every $E_{j,p}$ is finitely generated, being a quotient of $\widehat{F}$. For completeness its automorphism group is compact. Take a characteristic cofinal sequence of finite quotients, formed by bounded-index intersections as above. An automorphism gives compatible automorphisms of those finite quotients. A compatible tuple, with its compatible inverses, defines an automorphism of the original profinite group. The compatibility and subgroup preservation conditions are closed in the product of the finite automorphism sets. This identifies the automorphism group with a closed profinite subset and proves compactness.

For each i, the simultaneous condition on σ, a_i in (D.43.14) is closed. Projecting its compact solution set to the automorphism factor shows that the existential conjugator condition is also closed. Hence $\mathscr{X}_j$ is compact, and D.42 proves it nonempty. Every finite collection of inverse-limit equations has a solution: choose an element of $\mathscr{X}_k$ at an index at least all the indices in that collection and use its actual descents. Compactness of the product therefore gives

$$\varprojlim_j \mathscr{X}_j \neq \varnothing. \tag{D.43.15}$$

No surjectivity of the transition maps is needed.

A compatible tuple $(\sigma_j)_j$ defines an automorphism σ of $E_p(D)$ through (D.43.11). Its inverse is the tuple of the compatible σ_j^{-1}. It lies above α. To recover a conjugator with exact projection h_i, consider in $E_p(D)$ the closed sets

$$\mathcal{A}_{i,j} = \{a : \ q(a) \equiv h_i \pmod{W_j},\ \sigma_j(X_i) = \pi_j(a) X_i \pi_j(a)^{-1}\}. \tag{D.43.16}$$

Each is nonempty: lift a conjugator from (D.43.14) through the surjection π_j. The sets are decreasing by the descent equations. Their intersection is nonempty by compactness. Any element a_i in it satisfies $q(a_i) = h_i$ and $\sigma(X_i) = a_i X_i a_i^{-1}$, since all finite-base pro-p projections detect equality in (D.43.11). This establishes the three literal peripheral conditions.

Conversely an automorphism of $E_p(D)$ above α preserves each inverse image of W_j. Applying (D.43.7) to $K_j \to K_j/O^p(K)$ identifies the pro-p residual of this inverse image with $\ker(E_p(D) \to E_{j,p})$. Thus it descends to every stage, with the specified peripheral conditions. This is inverse to the construction just given. The two maps are continuous in their compact automorphism topologies. They prove the second isomorphism in (D.43.4) for the complete fibre.

All source maps and the fixed five-group quotient. Let $f : \widehat{F} \twoheadrightarrow E_p(D)$ satisfy $qf = \alpha q$. It sends K into the normal pro-p kernel $K/O^p(K)$. Hence it kills $O^p(K)$ and gives a surjective endomorphism of the same $E_p(D)$. Every finitely generated profinite group is Hopfian by the following finite argument. The inverse-image map of a surjective endomorphism permutes the finite set of open normal subgroups of each bounded index: it is injective on that set and preserves indices. It therefore preserves their intersection and induces an automorphism on each resulting finite quotient. Their intersection is trivial, so the endomorphism is injective. Apply this to the two-generated quotient $E_p(D)$. There is no finite-rank assumption on $K^{(p)}$ in this argument. This proves the all-source factorization and its peripheral comparison.

Finally set

$$D = \widehat{F}/O^2(K_n). \tag{D.43.17}$$

Its kernel over P_n is $K_n^{(2)}$, and every finite quotient has order $2^a 5^b$. For odd $p \neq 5$ it is therefore a pro-prime-to-p base in the precise sense of the statement. Choose the actual multiplier-one automorphism α of this D supplied by D.41 above β_Q, and choose its actual three conjugators $h_i \in D$. The kernel of $\widehat{F} \to D$ is exactly $O^2(K_n)$. Substitution into (D.43.3) gives the quotient (D.43.5) with its literal source map. The finite lower-two-central quotients $D_{n,j}$ of D.42 are cofinal here: their pro-two kernel series terminates in every finite two-group image. They are preserved by the chosen α. The proof above only requires that stability, so it yields precisely the compatible lift through these stages and all their finite relative p-central quotients.

Every pro-five quotient of (D.43.5) kills first its normal pro-p kernel and then the normal pro-two kernel of (D.43.17). It factors through P_n, which remains an actual quotient. This proves the exact maximal pro-five identification and the retained action β_Q. □

D.43 supplies the across-stage compatibility left open within D.42. It concerns a complete relative pro-p kernel over a base all of whose finite quotients have order prime to p. A further step in which the same prime already occurs in the base requires additional source comparisons. The full D.1–D.2 leading-image question continues to include those finite quotients.

4.26 Coprime modular covers with the old gentle conjugator retained

Statement D.44 (Proved). Exact modular descent and gentle normalization over a coprime base. *Let* $\widehat{B} = C_2 \amalg C_3 = \langle s, r \mid s^2 = r^3 = 1 \rangle$, *let* $F = \ker(\widehat{B} \to \mathfrak{S}_3)$, *and use*

$$\tau = sr, \qquad x = \tau^{-2}, \qquad y = sxs, \qquad z = (xy)^{-1}. \tag{D.44.1}$$

Let $q : \widehat{B} \twoheadrightarrow \Gamma$ *be an actual pro-prime-to-p quotient retaining the displayed* $\mathfrak{S}_3$ *quotient, and put* $K = \ker q$. *Thus* $p \geq 5$. *Suppose that an old automorphism* $\alpha_0 \in \mathrm{Aut}(\Gamma)$ *satisfies the following exact multiplier-one conditions: it fixes* τ, *sends* s, r *to conjugates of the same positive generators, and*

$$f_0 = s\alpha_0(s) \ \in \ \mathrm{im}(F' \to \Gamma). \tag{D.44.2}$$

Then on the actual relative cover

$$G = \widehat{B}/O^p(K) \tag{D.44.3}$$

there is an automorphism σ *inducing exactly* α_0, *fixing* τ, *and having*

$$\begin{aligned} f = s\sigma(s) \in \mathrm{im}(F' \to G), \quad & q(f) = f_0, \\ \sigma(x) = x, \qquad\qquad & \sigma(y) = f^{-1} y f. \end{aligned} \tag{D.44.4}$$

It satisfies the exact gentle equations

$$f(y,x)f(x,y) = 1, \qquad f(z,x)f(y,z)f(x,y) = 1 \tag{D.44.5}$$

in the actual pure quotient $H = \mathrm{im}(F \to G)$. *Here evaluations use any one preimage of* f *in* F'; *their values are independent of that choice at the displayed arguments. Its pure outer automorphism centralizes the actual* $\mathfrak{S}_3$*-outer action, by the explicit comparison below. The finite* σ*-stable quotient maps from this one* G *give a compatible family of generating gentle shadows retaining the literal old* f_0, *and hence its old outer coordinates.*

The torsion-conjugacy hypothesis in the statement holds automatically when the pure kernel $\Gamma_F = \ker(\Gamma \to \mathfrak{S}_3)$ *is pro-*ℓ *for* $\ell \geq 5$, *and the old multiplier-one gentle equations define an automorphism of* Γ. *In particular it holds for all the D.7–D.8 weight-nine gentle automorphisms at the actual class-ten exponent-*101 *quotient. Each of them therefore has these complete coprime relative lifts for every* $p \notin \{2, 3, 101\}$. *Its original three-strand shadow, including the central coordinate and the given* f_0, *is retained by the actual common-source comparison below.*

More generally, put

$$B_\ell = \widehat{B}/O^\ell(F), \qquad P_\ell = F^{(\ell)}, \qquad \ell \geq 5. \tag{D.44.26}$$

Every multiplier-one special outer automorphism of P_ℓ *centralizing its geometric* $\mathfrak{S}_3$ *has a unique representative given by a pair* $f_\ell \in P_\ell'$ *with the two cycles in* (D.44.5) *and* $x \mapsto x, \ y \mapsto f_\ell^{-1} y f_\ell$. *It extends to an actual automorphism of* B_ℓ. *For every* $p \notin \{2, 3, \ell\}$ *the resulting full pro-*ℓ *pair has a lift to*

$$\widehat{B}/O^p(O^\ell(F)). \tag{D.44.27}$$

The maximal pro-ℓ *quotient of the pure kernel of* (D.44.27) *is exactly* P_ℓ, *and the map on it is the originally specified outer automorphism, with the literal normalized* f_ℓ. *In particular this retains the R8 invariant weight-nine realizations, including a nonzero integral multiple of* $X^2Y^2(X+Y)^3[x,y]$, *throughout the complete tower* (D.44.27).

Proof. We retain the indicated source generators throughout the construction.

The conventions and the actual pure free group. There is a surjection from the discrete free product $C_2 * C_3$ to $\mathfrak{S}_3$, with s a transposition and r a three-cycle. The transversal

$$1,\ s,\ r,\ r^2,\ sr,\ sr^2 \tag{D.44.6}$$

gives twelve Schreier generators, one for each transversal element and each of s, r. The only nonidentity reduced generators are

transversal element	letter	Schreier generator
r	s	y^{-1}
r^2	s	x
sr	s	x^{-1}
sr^2	s	y.

(D.44.7)

These are literal free-product equalities with x, y from (D.44.1). The kernel intersects every conjugate of the two finite factors trivially; it acts freely on the free-product tree. Its quotient graph has six edges and five vertices, hence free rank two. The Schreier expressions in (D.44.7) show that x, y generate that kernel and hence form a free basis. Completion of this finite-index comparison identifies the actual F with the free profinite group on those same two words.

Direct free-product multiplication gives

	x	y	z
$r(-)r^{-1}$	y	z	x
$s(-)s$	y	x	$x^{-1}zx$.

(D.44.8)

The cusp denoted $t = r^{-1}s$ in the inverse convention is τ^{-1}. Preserving the conjugacy class of either positive element is equivalent to preserving the class of its inverse, by the actual inversion equation. For the pants source on $a = s, b = r^{-1}$, its third positive generator is

$$(ab)^{-1} = rs = s\tau s. \tag{D.44.9}$$

Thus the pants source is not assigned the word τ as its third generator without the conjugation displayed here.

The residual quotient and the chosen cusp conjugator. Let $\mathfrak{F} = \widehat{F}(a, b)$ be the free profinite pants group and let $\pi : \mathfrak{F} \twoheadrightarrow \widehat{B}$ send $a \mapsto s,\ b \mapsto r^{-1}$. Its kernel is the closed normal subgroup $\mathcal{R}$ generated by a^2, b^3: both quotients have exactly the same maps to every finite group, by the indicated generator relations. Put $K_{\mathfrak{F}} = \pi^{-1}(K)$. The restriction $K_{\mathfrak{F}} \twoheadrightarrow K$ is onto and has kernel $\mathcal{R}$. D.43.7 therefore gives

$$\begin{aligned} \pi(O^p(K_{\mathfrak{F}})) &= O^p(K), \\ G &\simeq \mathfrak{F}/(O^p(K_{\mathfrak{F}})\mathcal{R}). \end{aligned} \tag{D.44.10}$$

The product in the second line is closed: it is the product of two compact normal subgroups. This proves the actual quotient comparison for finite or infinite Γ.

Choose the given conjugators of $\alpha_0(s)$ and $\alpha_0(r)$ in Γ. For the third pants generator in (D.44.9) choose the specific conjugator

$$h_3 = \alpha_0(s)s. \tag{D.44.11}$$

It works because $\alpha_0(\tau) = \tau$: conjugating $s\tau s$ by $\alpha_0(s)s$ gives $\alpha_0(s)\tau\alpha_0(s) = \alpha_0(rs)$. D.42, or D.43 for the infinite base, provides a peripheral automorphism of $\mathfrak{F}/O^p(K_{\mathfrak{F}})$ with these

specified conjugator projections. It preserves the closed normal subgroup generated by a^2, b^3, since it sends each a, b to a conjugate of itself. The same argument for its inverse proves equality, so it descends through (D.44.10) to an automorphism ϕ of G. It induces α_0.

Let $A \in G$ be its third pants conjugator, with projection $\alpha_0(s)s$. Then

$$\begin{aligned} \phi(rs) &= A(rs)A^{-1}, \\ c &= \phi(s)As \in \ker(G \to \Gamma), \\ \phi(\tau) &= c\tau c^{-1}. \end{aligned} \tag{D.44.12}$$

For the middle assertion the projection is $\alpha_0(s)\alpha_0(s)ss = 1$. The last equation follows from $\tau = s(rs)s$. Consequently $\phi_1 = \mathrm{Inn}(c^{-1})\phi$ fixes τ and still induces exactly the old α_0, not an unspecified inner conjugate of it.

The exact abelianization and the kernel cusp gauge. Write $R = \ker(G \to \Gamma) = K^{(p)}$ and $H = \ker(G \to \mathfrak{S}_3)$. The source map $F \to H$ is onto. Let $\mathrm{ab}_p : F \to \mathbf{Z}_p^2$ be the abelianization in the actual basis x, y. Its restriction to K has pro-p image, so it kills $O^p(K)$. It therefore defines a homomorphism on H. Together with $H \to \Gamma_F^{\mathrm{ab}}$ it gives the exact isomorphism

$$H^{\mathrm{ab}} \xrightarrow{\simeq} \mathbf{Z}_p^2 \times \Gamma_F^{\mathrm{ab}}. \tag{D.44.13}$$

Here Γ_F, and thus its abelianization, is pro-prime-to-p. To verify the assertion, every finite abelian p-quotient of H is a quotient of $F^{\mathrm{ab},p}$; the displayed map retains all of $\mathbf{Z}_p^2$, so it is the full p-primary factor. Every finite abelian quotient of order prime to p kills R and factors through Γ_F. These two assertions detect every finite abelian quotient and prove injectivity of (D.44.13). The image is closed and surjects onto each factor. The two factors have no nontrivial common finite quotient, since their prime supports are disjoint. The finite subdirect-product argument, followed by compactness, proves surjectivity. This establishes the displayed actual abelianization map.

Set $S = \phi_1(s)$ and $f_1 = sS$. Its projection is exactly f_0. Equation (D.44.2) implies that α_0 is the identity on $\mathfrak{S}_3$: it fixes the image of s, and then that of $r = s\tau$. Thus ϕ_1 also induces the identity there, and $f_1 \in H$. Since $S^2 = 1$,

$$(sf_1s)f_1 = 1. \tag{D.44.14}$$

The action of s on $\mathbf{Z}_p^2$ interchanges the two coordinates by (D.44.8). Hence $\mathrm{ab}_p(f_1) = (a, -a)$ for a definite $a \in \mathbf{Z}_p$. Its other coordinate in (D.44.13) is zero, because f_0 is in the image of F'.

Let $\epsilon_p \in \widehat{\mathbf{Z}}$ be the idempotent with p-component one and every other primary component zero. The element

$$g = x^{\epsilon_p a} \tag{D.44.15}$$

centralizes τ and belongs to R: every finite image of Γ has order prime to p, so the image of this power there is the identity. Put $\sigma = \mathrm{Inn}(g)\phi_1$. Then σ still fixes τ and induces exactly α_0. Its conjugator is

$$\begin{aligned} f = s\sigma(s) &= y^{\epsilon_p a} f_1 x^{-\epsilon_p a}, \\ \mathrm{ab}_p(f) &= (a, -a) + (-a, a) = (0, 0). \end{aligned} \tag{D.44.16}$$

Its Γ_F^{ab} coordinate stays zero. Thus (D.44.13) proves $f \in H'$. Surjectivity of $F \to H$ identifies H' with the compact image of F': the image contains all commutators, and compactness retains their closed subgroup. This proves the first line of (D.44.4), including the unchanged literal f_0.

The centralizer ambiguity is an exact kernel gauge. We record why the choice made in (D.44.12) introduces no additional normalized representative in the same kernel-inner orbit. First let Γ be finite and put $e = \mathrm{ord}_\Gamma(\tau)$. The image of τ in $\mathfrak{S}_3$ is a transposition, so e is even. The subgroup K corresponds to the actual finite cover of the three-punctured sphere given by $K \subset F$. The word $\tau^e = x^{-e/2}$ is one of its based boundary generators, with its orientation possibly reversed. There are $|\Gamma|/e \geq 3$ boundary components: the coset map $\Gamma/\langle\tau\rangle \to \mathfrak{S}_3/\langle\bar{\tau}\rangle$ is onto. Hence each boundary is a primitive free generator after an actual surface basis change, as in D.41. The Magnus centralizer comparison there gives

$$C_R(\tau) \subset C_R(\tau^e) = \langle\tau^e\rangle = \langle\tau^{\epsilon_p}\rangle. \tag{D.44.17}$$

The reverse inclusion is immediate, and the last equality uses that e is a unit in $\mathbf{Z}_p$. Thus $C_R(\tau) = \langle x^{\epsilon_p}\rangle \cong \mathbf{Z}_p$, since $p \geq 5$.

For an infinite Γ, take cofinal finite quotients retaining $\mathfrak{S}_3$. The residual identity D.43.9, whose proof applies equally to the source $\widehat{B}$, identifies G with the inverse limit of the corresponding relative groups. At every finite stage the centralizer in its pro-p kernel has the same distinguished generator x^{ϵ_p}. The transition maps carry that generator to itself. The parameters in $\mathbf{Z}_p$ are unique by (D.44.17), so any element of $C_R(\tau)$ has one common parameter. This proves the same centralizer equality in the limit.

Two kernel conjugators normalizing $\phi(\tau)$ differ by an element of this centralizer. Thus the resulting τ-fixed automorphisms differ by $\mathrm{Inn}(x^{\epsilon_p v})$. Formula (D.44.16) changes the first abelian coordinate from a to $a - v$. The subsequent gauges therefore cancel exactly:

$$\mathrm{Inn}(x^{\epsilon_p(a-v)})\,\mathrm{Inn}(x^{\epsilon_p v}) = \mathrm{Inn}(x^{\epsilon_p a}). \tag{D.44.18}$$

This proves uniqueness of the representative fixing τ and having $f \in H'$ within the indicated kernel-inner orbit. Every normalization used here is trivial on Γ.

The two cycles, generation, and the actual symmetry. Let $S = \sigma(s)$, $R_0 = \sigma(r)$. Since $SR_0 = \tau$ and $f = sS$, we have the exact formulas

$$S = sf, \qquad R_0 = f^{-1}r. \tag{D.44.19}$$

The first follows from $s^2 = 1$; for the second use $S^2 = 1$ and $sfs = f^{-1}$. By (D.44.8), conjugation by s evaluates a word at (y, x), and conjugation by r evaluates it at (y, z). Thus $S^2 = 1$ gives the first equation of (D.44.5). The inverse of R_0^3 is

$$(r^2 f r^{-2})(r f r^{-1})f = f(z, x)f(y, z)f(x, y). \tag{D.44.20}$$

This proves the second equation, with its precise order. Any preimage of f in F' has these evaluation images, because the evaluations in (D.44.8) are actual conjugation maps on H. Also

$$\sigma(y) = SxS = f^{-1}yf, \qquad \sigma(z) = R_0^2 x R_0^{-2}. \tag{D.44.21}$$

The second element is conjugate to $z = r^2 x r^{-2}$ by $R_0^2 r^{-2}$. The restriction of σ to H is an automorphism, since it induces the identity on $G/H = \mathfrak{S}_3$. This proves the required generation in the actual pure quotient.

For the symmetry comparison let ι_b denote conjugation on H by $b \in G$. Then

$$\sigma|_H\, \iota_b\, (\sigma|_H)^{-1} = \iota_{\sigma(b)} = \mathrm{Inn}(\sigma(b)b^{-1})\, \iota_b. \tag{D.44.22}$$

The inner factor lies in H, because σ is the identity on $\mathfrak{S}_3$. Passing to $\mathrm{Out}(H)$ proves the asserted centralizer condition for the actual geometric $\mathfrak{S}_3$.

Equations (D.44.5) are precisely Harbater–Schneps relations (I)–(II) at $\lambda = 1$ [HS00, §0.1, (I)–(II); Proposition 4]. Their three-strand comparison also has the following literal modular coordinates:

$$u = r^{-1}s = \tau^{-1}, \qquad v = sr^{-1} = sus. \tag{D.44.23}$$

Here $uvu = vuv = s$ and $uv = r$. The automorphism just constructed sends u to u and v to $f^{-1}vf$. These equations follow from τ-fixing, $sfs = f^{-1}$, and (D.44.19). Thus the actual source map from the three-strand braid presentation is retained. Choose a cofinal sequence of characteristic open normal subgroups of the finitely generated group G, for example the intersections of all open normal subgroups of index at most n. Intersect with H, and, when Γ is finite, also with R; these intersections remain σ-stable because σ preserves H, R. The resulting actual finite quotient maps commute with σ, retain the specified source generators, and form a cofinal compatible system. In each of these finite quotients the torsion-to-hexagon calculation of E.9, or the displayed formulas themselves, gives the complete two gentle hexagons with multiplier one. The map is onto because σ is an automorphism of G. The actual descent through the σ-stable kernels supplies an automorphism at every stage. One choice of a preimage of f in F' works simultaneously in all these finite quotients. This proves their compatible gentle-shadow assertion.

Where the torsion gate is automatic and how the old braid shadow is retained. Suppose Γ_F is pro-ℓ, $\ell \geq 5$, and the old gentle automorphism is already normalized as in (D.44.2). It induces the identity on $\mathfrak{S}_3$. In the inverse image of $\langle \bar{s} \rangle$, the original $\langle s \rangle$ and the new $\langle \alpha_0(s) \rangle$ are complements to the same normal pro-ℓ kernel. The coprime complement proof in D.41 conjugates them by that kernel. Their positive generators have the same image $\bar{s}$, so the conjugacy is on the generators themselves. Apply the same argument to the inverse image of $\langle \bar{r} \rangle$. Its quotient has order three, and the images of the two positive generators are both $\bar{r}$. This proves exactly the two conjugacies required in the statement.

For the D.7–D.8 example, quotient the actual three-strand group by its central generator. The pure kernel over $\mathfrak{S}_3$ is

$$F/(\gamma_{11}F\, F^{101}), \tag{D.44.24}$$

a finite 101-group. The already proved gentle maps are automorphisms, and retain multiplier one and their specified f_0. Hence the preceding complement argument applies to every one of them.

To retain also the original finite braid target $\widetilde{\Gamma} = B_3/K_3$, form the actual common refinement

$$G \times_\Gamma \widetilde{\Gamma}. \tag{D.44.25}$$

The kernel of $G \to \Gamma$ is pro-p, and the kernel of $\widetilde{\Gamma} \to \Gamma$ is the central group of order 101. The natural source map $\widehat{B}_3$ to (D.44.25) is onto: on finite kernel quotients its two Goursat quotients have orders dividing both a power of p and 101, so they are trivial; compactness then gives surjectivity. The two automorphisms σ and the old braid automorphism induce the same α_0 on Γ. Their product is therefore an actual automorphism of the displayed fibre product.

Finally the direct product $PB_3 = F \times \langle C_3 \rangle$ used in D.7 shows that the image of F in $\widetilde{\Gamma}$ maps isomorphically to the pure group (D.44.24). Thus any preimage in F' of the new f, already projecting to f_0 in Γ, has exactly the old f_0 in $\widetilde{\Gamma}$; there is no discarded central parameter in this comparison. On (D.44.25) the braid action $u \mapsto u,\ v \mapsto f^{-1}vf$ agrees with the two component automorphisms. It fixes the old central generator because the multiplier is one. This proves the full old-shadow assertion, including its central coordinate, generation, and all projections.

The entire pro-ℓ pair and its exact outer comparison. The kernel of $B_\ell \to \mathfrak{S}_3$ is $P_\ell = F/O^\ell(F)$, the free pro-ℓ group on the same x, y. It is centreless: the Magnus and retraction argument of D.41 gives $C_{P_\ell}(x) = \langle x \rangle$ and $C_{P_\ell}(y) = \langle y \rangle$, whose intersection is trivial by abelianization. Let $\omega : \mathfrak{S}_3 \to \mathrm{Out}(P_\ell)$ be the outer action of this actual extension. There is an actual topological isomorphism

$$\begin{aligned} B_\ell &\longrightarrow \mathrm{Aut}(P_\ell) \times_{\mathrm{Out}(P_\ell)} \mathfrak{S}_3, \\ b &\longmapsto (\mathrm{Ad}_b, \bar{b}). \end{aligned} \tag{D.44.28}$$

Its kernel is $Z(P_\ell) = 1$. For surjectivity, given a pair (a, d) choose b over d; the equality of outer classes says $a\,\mathrm{Ad}_b^{-1} = \mathrm{Inn}(u)$ for some $u \in P_\ell$, and the pair is the image of ub. All groups and maps are compact and Hausdorff, so the continuous bijection is a topological isomorphism.

Let β be one of the outer automorphisms in the assertion. The three peripheral conjugacy classes are distinct by their abelian vectors, so their permutation defines an actual map to $\mathfrak{S}_3$. It sends the geometric $\mathfrak{S}_3$ onto its faithful natural action; the permutation of a centralizing β therefore lies in $Z(\mathfrak{S}_3) = 1$. Choose a representative $\psi \in \mathrm{Aut}(P_\ell)$ fixing x literally; its peripheral conjugator at x gives this inner adjustment. Because $[\psi]$ centralizes $\omega(\mathfrak{S}_3)$, the map

$$(a, d) \longmapsto (\psi a \psi^{-1}, d) \tag{D.44.29}$$

preserves the pullback in (D.44.28) and is an automorphism. Transport it to an automorphism ϕ of B_ℓ, whose restriction to P_ℓ is exactly ψ. We have $\phi(\tau)^2 = \psi(x^{-1}) = x^{-1} = \tau^2$. Both $\phi(\tau)$ and τ centralize x, and their quotient lies in P_ℓ. The primitive centralizer therefore gives $\phi(\tau) = x^a \tau$ for a unique $a \in \mathbf{Z}_\ell$. Taking squares gives $x^{2a} = 1$; torsion-freeness of $\langle x \rangle$ gives $a = 0$. Thus the positive element τ, not only its order-two quotient, is fixed literally.

The two finite cyclic subgroups generated by s, r are conjugate to their new images by the coprime complement argument already proved above. Set $f_1 = s\phi(s)$. Its abelianization is $(a, -a)$ for a definite $a \in \mathbf{Z}_\ell$. The actual adjustment $\phi_0 = \mathrm{Inn}(x^a)\phi$ fixes τ and preserves β, while

$$f_\ell = s\phi_0(s) = y^a f_1 x^{-a} \ \in P_\ell'. \tag{D.44.30}$$

The earlier word computations prove both cycles and the displayed pure action. If two such normalized extensions induce the same β, their restrictions differ by an inner automorphism of P_ℓ, and (D.44.28) makes their full extensions differ by that same inner automorphism. Literal τ-fixing forces its conjugator into $C_{P_\ell}(\tau) = \langle x \rangle$. Formula (D.44.30) and zero abelianization then force its exponent to vanish. This proves the claimed uniqueness.

Conversely, start with $f_\ell \in P_\ell'$ satisfying the two cycles and for which $\psi_f : x \mapsto x,\ y \mapsto f_\ell^{-1} y f_\ell$ is an automorphism of P_ℓ. The assignments $s \mapsto sf_\ell,\ r \mapsto f_\ell^{-1} r$ satisfy $s^2 = r^3 = 1$, so give a homomorphism $\widehat{B} \to B_\ell$. Its F-image is pro-ℓ, hence it kills $O^\ell(F)$ and factors through this same B_ℓ. Its restriction is exactly ψ_f, and its $\mathfrak{S}_3$ quotient is the identity, so it is onto. The finitely generated profinite group B_ℓ is Hopfian, giving an automorphism. Equation (D.44.22) proves that its special outer action centralizes the same geometric $\mathfrak{S}_3$. This supplies both directions of the normalized-pair comparison, without importing a full profinite pair into a pro-ℓ target.

The group B_ℓ is pro-$\{2, 3, \ell\}$. Apply the first part of the theorem to this actual base and its just constructed automorphism. The two kernel normalizations in that proof are trivial on B_ℓ, so the entire element f_ℓ, rather than only one leading coordinate, is retained in (D.44.27). Its pure kernel $H_{\ell,p}$ sits in the actual exact sequence

$$1 \longrightarrow O^\ell(F)/O^p(O^\ell(F)) \longrightarrow H_{\ell,p} \longrightarrow P_\ell \longrightarrow 1. \tag{D.44.31}$$

The left group is pro-p. Every map from $H_{\ell,p}$ to a finite ℓ-group kills it, since $p \neq \ell$. The last group is already pro-ℓ, proving the asserted maximal pro-ℓ quotient by the displayed maps.

For completeness the R8 realization used here has an integral construction in this same group P_ℓ. Represent an integral degree-n special derivation, $n \geq 2$, by conjugators at y and z, keeping x fixed. Their product error begins one degree above the prescribed leading relation. At every later ordinary lower-central degree m, altering the two conjugators in degree $m-1$ changes that error by

$$(u, v) \longmapsto [u, y] + [v, -x - y]. \tag{D.44.32}$$

The image equals $[L_{m-1}, x] + [L_{m-1}, y] = L_m$ in the integral free Lie algebra: iterated Jacobi rewrites every bracket monomial with one generator as its outer entry. Thus the actual lower-central error can be cancelled at each stage without changing the leading term. The conjugators converge, because each finite ℓ-group quotient kills sufficiently deep ordinary lower-central terms. The resulting endomorphism has identity Frattini action and is an automorphism.

The IA outer special group and its closed filtration kernels are pro-ℓ. For a finite group of order prime to ℓ, fixed points lift across an onto equivariant homomorphism of pro-ℓ groups: over a fixed target element, its conjugation of the two complements in the associated semidirect product is removed by coprime complement conjugacy in the kernel. Finite quotients followed by compactness give the same conclusion for these closed groups. Apply this to the actual geometric $\mathfrak{S}_3$ action and the leading quotient of the IA special group. Its surjectivity is exactly the preceding error-cancellation construction. Hence an invariant integral leading term has an actual special outer lift centralizing $\mathfrak{S}_3$. The rational invariant vectors computed in D.4 have integral multiples in this lattice; averaging their integral source representatives by $1/6$ retains integrality in $\mathbf{Z}_\ell$. The extra weight-nine coordinate in the statement therefore has the claimed actual pro-ℓ realization. Equations (D.44.28)–(D.44.31) supply its exact further source and pair maps. □

The program `a_modular_coprime.py` retains all twelve literal Schreier expressions in (D.44.7), the complete free-product identities (D.44.8)–(D.44.9), and both cycle comparisons. It independently checks the signed integer cusp gauges. Its finite model is the actual quotient of $C_2 * C_3$ with pure kernel $\mathbf{F}_5^2$ and the displayed $\mathfrak{S}_3$ action, of order 150. All 25 kernel-inner maps and all five choices of cusp centralizer give the same normalized representative; all 125 cases, including their unique scalar gauges, are saved.

The conclusion concerns the complete coprime relative covers of the specified old gentle automorphisms. Finite order-two and order-three equations alone do not supply the input conjugacies in a general mixed-primary quotient; the proof establishes them for the pro-ℓ inputs just specified. The remaining D.2 comparison still includes relative primes dividing the old base and the other finite source extensions.

4.27 Cyclic prime outer actions and the full $\mathrm{PSL}_2(32)$ chief step

Statement D.45 (Proved). All multiplier-one peripheral fibres across split cyclic-prime outer chief extensions. *Let S be a finite nonabelian simple group, and fix an actual isomorphism*

$$\mathrm{Aut}(S) = S \rtimes \langle t \rangle, \qquad |t| = r, \quad \mathrm{Out}(S) = \langle \bar{t} \rangle \simeq C_r, \quad r \text{ prime}, \tag{D.45.1}$$

where S is identified with its inner automorphisms. Let

$$1 \longrightarrow A = S^\Omega \longrightarrow E \xrightarrow{\pi} D \longrightarrow 1 \tag{D.45.2}$$

be a finite chief extension, and let $q_E : \widehat{F}(x_1, x_2) \twoheadrightarrow E$ *be an actual epimorphism, with* $x_3 = (x_1x_2)^{-1}$. *Put* $X_i = q_E(x_i)$, $q_0 = \pi q_E$, *and* $d_i = q_0(x_i)$. *For every epimorphism* $q_1 : \widehat{F} \twoheadrightarrow D$ *and every specified* $h_i \in D$ *satisfying*

$$q_1(x_i) = h_i d_i h_i^{-1}, \tag{D.45.3}$$

choose arbitrary lifts $H_i \in E$ *of* h_i. *There is an epimorphism* $f : \widehat{F} \twoheadrightarrow E$ *such that*

$$\pi f = q_1, \qquad f(x_i) = a_i H_i X_i H_i^{-1} a_i^{-1} \quad (a_i \in A). \tag{D.45.4}$$

The number of these literal epimorphisms is exactly the number for $(q_1, h_1, h_2, h_3) = (q_0, 1, 1, 1)$. *No restriction is imposed on the finite base* D, *its factor stabilizers, or its fixed peripheral points.*

In particular (D.45.1) *holds for*

$$S = \mathrm{PSL}_2(32) = \mathrm{SL}_2(\mathbf{F}_{32}), \qquad t : (a_{ij}) \longmapsto (a_{ij}^2), \qquad |\operatorname{Aut}(S)| = 163680. \tag{D.45.5}$$

Every D.37 automorphism β_Q *of every specified characteristic target* P_n *therefore lifts through every actual* $\mathrm{PSL}_2(32)$*-chief extension above that target, with its three stated base conjugators. The maximal pro-five quotient of this extension is exactly the given* P_n, *with the action* β_Q. *The same conclusion retains a prescribed epimorphism into any additional finite group over* D *whose kernel over* D *is solvable, through their actual fibre product.*

Proof. We first prove the surface maps which transport the outer phases, and then retain all the finite extension and source maps used to count the full fibre.

An exact positive boundary push. Use a surface presentation

$$\Pi = \left\langle a_1, b_1, \dots, a_g, b_g, c_1, \dots, c_b \;\middle|\; \prod_{j=1}^{g} [a_j, b_j] \prod_{i=1}^{b} c_i = 1 \right\rangle, \qquad b \geq 2. \tag{D.45.6}$$

The finite-index pants covers below have $b \geq 3$. One boundary may be eliminated to obtain a free basis; all substitutions below preserve the displayed relation before or after that elimination.

For one handle and an adjacent boundary c, the following are actual free-group substitutions:

$$\begin{aligned} P(a) &= c^{-1}ac, \\ P(b) &= c^{-1}a^{-1}cabc, \\ P(c) &= c^{-1}a^{-1}cac. \end{aligned} \tag{D.45.7}$$

They fix $[a, b]c$ literally, and fix all the other generators. Their inverse can be obtained from the following explicit Nielsen changes. Put

$$u = ac, \qquad v = c^{-1}bc, \qquad z = vcv^{-1}. \tag{D.45.8}$$

The inverse expressions are

$$a = uv^{-1}z^{-1}v, \qquad b = v^{-1}zvz^{-1}v, \qquad c = v^{-1}zv, \tag{D.45.9}$$

and $[u, v]z = [a, b]c$. The twist $u \mapsto u$, $v \mapsto vu$, $z \mapsto z$, transported through (D.45.8)–(D.45.9), composed with the inverse of $a \mapsto a$, $b \mapsto ba$, $c \mapsto c$, is exactly (D.45.7). Reversing these

operations gives its inverse. This also realizes it by the two boundary-fixed Dehn twists used in D.41. Its abelianization sends

$$a \longmapsto a, \qquad b \longmapsto b + c, \qquad c \longmapsto c. \tag{D.45.10}$$

Its positive boundary conjugator is $c^{-1}a^{-1}$.

For two adjacent boundaries c, d, the Hurwitz map

$$J(c) = cdc^{-1}, \quad J(d) = c, \qquad J^{-1}(c) = d, \quad J^{-1}(d) = d^{-1}cd \tag{D.45.11}$$

fixes their product. Its square is

$$J^2(c) = (cd)c(cd)^{-1}, \qquad J^2(d) = cdc^{-1}. \tag{D.45.12}$$

These equations retain both positive orientations and both conjugating paths.

Independent phases with a prescribed character. Write $\widetilde{S} = S \rtimes \langle t \rangle$, with outer projection $\epsilon : \widetilde{S} \to \mathbf{F}_r$. Let $\chi : \Pi \to \mathbf{F}_r$ be nonzero. For each boundary c_i, prescribe an S-conjugacy invariant subset $T_i \subset \epsilon^{-1}(\chi(c_i))$. The number of epimorphisms $\varphi : \widehat{\Pi} \twoheadrightarrow \widetilde{S}$ with $\epsilon\varphi = \chi$ and $\varphi(c_i) \in T_i$ is unchanged upon conjugating each T_i independently by any power of t. We prove this by actual automorphisms of the source.

First suppose $\chi(c) = 0$. If a handle has nonzero character on a, then (D.45.7) preserves χ by (D.45.10) and changes the boundary conjugator phase by $-\chi(a)$. If only the handle value at b is nonzero, use the actual basis change $a \mapsto ab,\ b \mapsto b$, whose inverse is $a \mapsto ab^{-1}$. It fixes $[a, b]$; conjugating the displayed push by this change gives phase $-\chi(b)$. Each nonzero phase generates $\mathbf{F}_r$, so powers of these maps give every required phase.

If the character is supplied instead by another boundary d, then (D.45.12) preserves χ. Its conjugator phases are $\chi(d)$ at c and $\chi(c) = 0$ at d; all other boundaries are fixed. Again the first phase is a generator of $\mathbf{F}_r$. An arbitrary boundary can be moved to the indicated position by the positive Hurwitz maps (D.45.11). Handle blocks can also be interchanged by an actual positive Hurwitz map on their commutators: for $A = [a, b]$, $B = [u, v]$, send

$$(a, b, u, v) \longmapsto (AuA^{-1}, AvA^{-1}, a, b). \tag{D.45.13}$$

Its commutator blocks become (ABA^{-1}, A), so their product is fixed. The inverse uses the inverse Hurwitz map and conjugates both entries of the affected handle pair by the same word. These moves put the selected handle last and c first among the boundaries, where (D.45.7) applies. Pulling the push back by the actual inverse moves proves the assertion in the original presentation.

Here is also the precise phase comparison under these changes of paths. For a character-preserving automorphism sending c_i to $u_i c_i u_i^{-1}$, the phase is $\chi(u_i)$. At a zero-character boundary it is independent of the conjugator choice: the boundary is primitive, and its free-group centralizer is $\langle c_i \rangle$, of character zero. The phase at a fixed boundary adds under composition of character-preserving maps. Upon conjugation by one of the coordinate changes above, the character is transported with that change, and the phases of the change and its inverse cancel. Thus the pulled-back map has the stated phase only at the chosen zero-character boundary. No uncontrolled based-path phase is introduced.

For a boundary with $j = \chi(c_i) \neq 0$, every S-class in its outer coset is already invariant under conjugation by every t^k. Indeed, for $a \in \epsilon^{-1}(j)$, choose m with $mj = k$ in $\mathbf{F}_r$. Then $t^k a^{-m} \in S$, and

$$t^k a t^{-k} = (t^k a^{-m})\, a\, (t^k a^{-m})^{-1}. \tag{D.45.14}$$

This proves the assertion element by element.

Finally, if a source map has boundary conjugator word u_i of phase k, then $\varphi(u_i) = s_i t^k$ for an actual $s_i \in S$. Hence precomposition changes membership in T_i precisely to membership in $t^k T_i t^{-k}$; its S-factor has no effect on that set. Precomposition by the displayed inverse gives the reverse map. Every such map is an automorphism of the same surface group, so surjectivity is preserved. This proves the claimed equality of epimorphism counts, including empty sets, by explicit bijections.

Matching the actual surface characters. Two finite covers of the pants with the same index and matched oriented boundaries have the same genus. The adjacent maps (D.45.11) realize the prescribed boundary permutation. Suppose two characters to $\mathbf{F}_r$ have the same values on those boundaries, and are either both zero or both nonzero. There is an actual surface isomorphism matching the characters and the oriented boundary classes. If all boundary values vanish, the nonzero characters are nonzero covectors on the handle symplectic space. The elementary Dehn twists of D.41 generate the symplectic group modulo r, which is transitive on its nonzero covectors. Their integral elementary matrices and inverses, including the two-by-two diagonal factorization in D.41, realize the needed finite-field map by a finite composition of these actual homeomorphisms.

If a boundary d has character value $e \neq 0$, the bounding pair with homology classes a and $a + d$ acts by

$$v \longmapsto v + \langle v, a\rangle d. \tag{D.45.15}$$

It is the product of the two indicated Dehn twists; their disjoint simple curves bound that boundary and an annulus with a handle attached. Its k-th power changes the character by $ke\langle -, a\rangle$. Since e is a unit, the handle basis curves and their powers give every desired translation of the handle character. In genus zero the boundary values already determine the character. All these maps are based outside their supports. They induce inverse maps on the discrete surface groups and on their profinite completions. The extra boundary conjugators of such an isomorphism have definite character phases; the preceding independent-phase bijections handle every one of them.

The actual twisted chief extension, with its signs. Let D_* be a factor stabilizer and identify $\Omega = D_* \backslash D$ with its right action. Choose representatives $t_\omega \in D$, with $t_{\omega_0} = 1$. Centerlessness gives the actual isomorphism

$$E \longrightarrow \mathrm{Aut}(A) \times_{\mathrm{Out}(A)} D, \qquad e \longmapsto (\mathrm{Ad}_e, \pi(e)). \tag{D.45.16}$$

Its kernel is $Z(A) = 1$; for a required pair, choose a lift of its D-coordinate and correct the difference by the unique inner automorphism of A. This proves surjectivity as well.

Write the outer action as the cocycle $b_d(\omega) \in \mathbf{F}_r$, so

$$b_{de}(\omega) = b_d(\omega) + b_e(\omega d). \tag{D.45.17}$$

Set

$$\begin{aligned} \epsilon_*(c) &= b_c(\omega_0) \quad (c \in D_*), \\ s_\omega &= b_{t_\omega}(\omega_0), \\ k(\omega, d) &= t_\omega d t_{\omega d}^{-1}. \end{aligned} \tag{D.45.18}$$

Then $\epsilon_* : D_* \to \mathbf{F}_r$ is a homomorphism, and the cocycle identity gives

$$b_d(\omega) = -s_\omega + \epsilon_*(k(\omega, d)) + s_{\omega d}. \tag{D.45.19}$$

The two signs in this formula are retained. Conjugating the wreath coordinates by $(t^{s_\omega})_\omega$, namely

$$(z,d) \longmapsto (t^{s_\omega} z(\omega) t^{-s_{\omega d}}, d), \tag{D.45.20}$$

identifies the given extension with

$$\begin{aligned} E_* &= \{(z,d) \in \widetilde{S}^\Omega \rtimes D : \epsilon(z(\omega)) = \epsilon_*(k(\omega,d))\}, \\ (z,d)(w,e) &= (z(\omega)w(\omega d), de). \end{aligned} \tag{D.45.21}$$

This is an actual isomorphism, carrying the specified X_i, H_i with it.

For either source $q = q_j$, put $K_j = \ker q_j$ and $H_{j,C} = q_j^{-1}(C)$. Choose actual source words T_ω with $q(T_\omega) = t_\omega$ and $T_{\omega_0} = 1$. The twisted Shapiro formula is

$$\begin{aligned} \varphi : H_{j,D_*} &\longrightarrow \widetilde{S}, \qquad \epsilon\varphi = \epsilon_* q_j, \\ z_\varphi(g)(\omega) &= \varphi(T_\omega g T^{-1}_{\omega q_j(g)}), \\ f_\varphi(g) &= (z_\varphi(g), q_j(g)). \end{aligned} \tag{D.45.22}$$

The middle representatives cancel in a product, proving its homomorphism identity and its exact outer coordinates. Conversely, for a cocycle z evaluate it at ω_0 on H_{j,D_*}. The vector $v(\omega) = z(T_\omega)(\omega_0)$ belongs to S^Ω and has $v(\omega_0) = 1$, since its outer transition at t_ω is the identity. It satisfies

$$z_\varphi(g)(\omega) = v(\omega) z(g)(\omega) v(\omega q_j(g))^{-1}. \tag{D.45.23}$$

This proves the normalization and its inverse. There are exactly $|S|^{|\Omega|-1}$ literal cocycles for each normalized surface map.

Every intermediate character and the prescribed conjugators. For $D_* \le C \le D$, retain every homomorphism $\epsilon_C : C \to \mathbf{F}_r$ extending ϵ_*; there may be none. Choose representatives R_ν for $C\backslash D$, with $R_C = 1$. Define

$$\begin{aligned} k_C(\nu,d) &= R_\nu d R_{\nu d}^{-1}, \\ b_C(d,\nu) &= \epsilon_C(k_C(\nu,d)), \\ p(\omega) &= C t_\omega, \qquad g_\omega = \epsilon_C(t_\omega R_{p(\omega)}^{-1}). \end{aligned} \tag{D.45.24}$$

Let E_C be (D.45.21) formed with $C\backslash D, \epsilon_C, k_C$. The actual injection over D is

$$j_C : E_C \hookrightarrow E_*, \qquad j_C(z,d)(\omega) = t^{g_\omega} z(p(\omega)) t^{-g_{\omega d}}. \tag{D.45.25}$$

The adjacent powers cancel under multiplication. Reading any coordinate above each ν proves injectivity. Its outer-coordinate equation is

$$\epsilon_*(k(\omega,d)) = g_\omega + b_C(d, p(\omega)) - g_{\omega d}, \tag{D.45.26}$$

which follows by applying ϵ_C to the literal factorization in C. Also $g_{\omega_0} = 0$. Thus a surface homomorphism on $H_{j,C}$, followed by its Shapiro map and j_C, restricts at the base coordinate to its very same map on H_{j,D_*}.

For fixed i, two elements over $q_j(x_i)$ are conjugate by A precisely when each of their permutation-cycle products is conjugate by S. The successive conjugator equations along a cycle give this equivalence: once its first conjugator is chosen in S, all others are determined; their outer exponents vanish because the two prescribed component patterns agree. The last equation is exactly the cycle-product condition.

This supplies an S-conjugacy invariant subset $T_{j,C,\delta} \subset \widetilde{S}$ for every boundary cycle δ on $C \backslash D$. Explicitly choose the initial coordinate ν of δ, and initial coordinates ω of all cycles γ above it with $p(\omega) = \nu$. If their relative degrees are m_γ, intersect the conditions

$$t^{g_\omega} a^{m_\gamma} t^{-g_\omega} \in \mathcal{C}_{j,\gamma}, \qquad \epsilon(a) = \epsilon_C(q_j(c_\delta)). \tag{D.45.27}$$

Here $\mathcal{C}_{0,\gamma}$ is the actual S-class of the cycle product of X_i, and $\mathcal{C}_{1,\gamma}$ that of $H_i X_i H_i^{-1}$. All roots and all empty intersections are retained.

The exact boundary bijection is

$$Cd\langle q_1(x_i)\rangle \longmapsto Cdh_i\langle d_i\rangle. \tag{D.45.28}$$

It preserves cycle lengths and every intermediate boundary degree by (D.45.3). For a new initial coordinate ν', take $\nu' h_i$ for its old match. In E_C the canonical element over h_i has components

$$L_i(\nu) = t^{\,b_C(h_i,\nu)}. \tag{D.45.29}$$

Its image under j_C and the specified H_i have the same projection to D; their ratio lies in A. Conjugation by L_i therefore transports the entire old local condition to the new one, including the intersection in (D.45.27). On the cycle product the conjugator is the actual component $L_i(\nu')$, giving

$$T_{1,C,\delta'} = t^{\,b_C(h_i,\nu')} T_{0,C,\delta}\, t^{-b_C(h_i,\nu')}. \tag{D.45.30}$$

The inverse conjugation proves the reverse inclusion. This argument does not require $X_i \in j_C(E_C)$. In particular the matching boundary characters are equal, not merely proportional.

The two actual surface characters $\chi_j = \epsilon_C q_j$ on $H_{j,C}$ are both zero when $\epsilon_C = 0$, and both nonzero otherwise, since $q_j(H_{j,C}) = C$. Use the character-matching isomorphism proved above. When $\epsilon_C = 0$, its boundary conjugators have zero outer phase, and (D.45.30) has phase zero as well. When $\epsilon_C \neq 0$, the independent-phase automorphisms remove every phase in (D.45.30) and every additional phase of the matching isomorphism. They preserve surjectivity. Consequently the numbers of epimorphisms

$$H_{j,C} \twoheadrightarrow S \rtimes \operatorname{im}(\epsilon_C) \tag{D.45.31}$$

with the exact outer character and all boundary sets (D.45.27) are equal for $j = 0, 1$.

Subtracting exactly the maps factoring through the base. For an epimorphism in (D.45.31), $\varphi(K_j)$ lies in S and is normal in the target, hence is either 1 or S. We retain only those with image S. The subtracted maps are exactly the actual epimorphisms

$$\Phi : C \twoheadrightarrow S \rtimes \operatorname{im}(\epsilon_C), \qquad \epsilon\Phi = \epsilon_C, \qquad \varphi = \Phi q_j|_{H_{j,C}}. \tag{D.45.32}$$

There is a bijection of the subtracted sets using the same Φ on both sides. Indeed its Shapiro section is the homomorphism

$$s_\Phi : D \longrightarrow E_C, \qquad s_\Phi(d)(\nu) = \Phi(R_\nu d R_{\nu d}^{-1}), \tag{D.45.33}$$

with base coordinate d. It is independent of q_j. The two normalized source maps are $j_C s_\Phi q_j$. At a peripheral, (D.45.3) gives

$$j_C s_\Phi(q_1(x_i)) = j_C s_\Phi(h_i)\, j_C s_\Phi(d_i)\, j_C s_\Phi(h_i)^{-1}. \tag{D.45.34}$$

The first factor and H_i differ by an element of A. Thus this value is A-conjugate to $H_i X_i H_i^{-1}$ exactly when $j_C s_\Phi(d_i)$ is A-conjugate to X_i. This proves both directions of the desired bijection for the very same Φ. Subtracting these equal finite sets from the equal counts in (D.45.31) retains equal counts with $\varphi(K_j) = S$, over arbitrary finite bases.

The full normalizer count and generation. For a surface map on H_{j,D_*} with $\varphi(K_j) = S$, put $N_\varphi = \ker(\varphi|_{K_j})$. Its normalizer in $\widehat{F}$ contains H_{j,D_*}, and contains K_j; it is therefore one of the actual $H_{j,C}$. It contains $H_{j,C}$ exactly when φ has a unique extension to a map $H_{j,C} \to \mathrm{Aut}(S)$. To prove this, if $H_{j,C}$ normalizes the kernel, conjugation on $K_j/N_\varphi \simeq S$ defines that extension. It agrees with φ on H_{j,D_*} because $C_{\mathrm{Aut}(S)}(S) = 1$. Its outer coordinate factors through C and is a unique ϵ_C extending ϵ_*. It is onto $S \rtimes \mathrm{im}(\epsilon_C)$, since its K_j-image is S. Conversely an extension normalizes the kernel. The same centralizer equality proves uniqueness.

Let $M_{j,C}$ be the sum over all the ϵ_C in (D.45.24) of the retained counts after (D.45.32). The previous paragraphs prove $M_{0,C} = M_{1,C}$ for every actual subgroup $C \geq D_*$. The injections j_C and their base-coordinate comparison prove that $M_{j,C}$ counts precisely those original surface maps whose kernel normalizer contains $H_{j,C}$. Different extending characters are disjoint because the extending map is unique. Finite subgroup-poset inversion now gives the exact number with minimal normalizer:

$$e_j = \sum_{C \geq D_*} \mu(D_*, C)\, M_{j,C}, \qquad e_0 = e_1. \tag{D.45.35}$$

Every summand refers to the displayed surface, embedding and boundary maps; independent isomorphisms of those surfaces suffice for this cardinality equality.

The original epimorphism q_E has $q_E(K_0) = A$. Its coordinate kernels on K_0 are pairwise distinct and have precisely the stated factor stabilizers, so $e_0 > 0$. For a map counted by e_1, its conjugate K_1-kernels at distinct factors are pairwise distinct, by the exact normalizer condition. Its K_1-image is then the full product S^Ω: a proper subdirect product of nonabelian simple factors identifies two factors by an isomorphism and would identify those two coordinate kernels. This follows inductively from the two-factor Goursat lemma and the fact that normal subgroups of a product of nonabelian simple groups are products of factors. The base image is D, hence the resulting map is onto E. Conversely every epimorphism to E has these same full coordinate images and distinct kernels. The literal gauge count (D.45.23) therefore proves that the two complete epimorphism fibres have the same positive size

$$|S|^{|\Omega|-1} e_j. \tag{D.45.36}$$

This proves (D.45.4), including every specified base conjugator.

The actual $\mathrm{PSL}_2(32)$ **source.** We give a direct matrix proof of the group facts used in (D.45.5). Let $k = \mathbf{F}_{32}$, and put $S = \mathrm{SL}_2(k)$. Its centre is trivial in characteristic two. The square map of $k^\times$ is bijective, so scaling each invertible two-by-two matrix to determinant one identifies S with $\mathrm{PGL}_2(k)$. Its action on $\mathbf{P}^1(k)$ is sharply three-transitive: the unique fractional linear map between two ordered triples of distinct points has such a determinant-one representative. In particular it is faithful and primitive.

The upper unipotent subgroup $U = \{u(b) : b \in k\}$ is abelian and normal in the point stabilizer at infinity. Its conjugates generate S, by upper and lower matrix elimination. For $h(a) = \mathrm{diag}(a, a^{-1})$,

$$[h(a), u(b)] = u((a^2 - 1)b). \tag{D.45.37}$$

Choose $a \neq 1$. Every element of U is then a commutator; its conjugates prove that S is perfect. For a nontrivial normal subgroup M, primitivity and faithfulness make M transitive. Thus every $g \in S$ can be written $g = mh$, with $m \in M$ and h in the stabilizer of infinity. Modulo M, every conjugate of U therefore has image equal to that same abelian image of U. These conjugates generate S, so S/M is abelian; perfectness gives $M = S$. This proves simplicity with the actual action.

The group U has order 32, the full two-part of $|S| = 32(32^2 - 1)$. Its normalizer is the upper triangular point stabilizer of order $32 \cdot 31$. The 33 Sylow-two subgroups are therefore exactly the conjugates indexed by $\mathbf{P}^1(k)$, each with its unique fixed point. An automorphism of S permutes these actual subgroups and gives a permutation normalizing the faithful projective action. This map on automorphisms is injective: the action of the image of any group element is determined by its conjugation permutation on the Sylow subgroups. Conversely a permutation normalizing S induces an automorphism of this faithful permutation group.

Compose such a permutation by the unique element of S making it fix infinity, zero and one. It normalizes U, the translation group on k, so the resulting permutation $\eta : k \to k$ satisfies

$$\eta(a + b) = \eta(a) + \eta(b). \tag{D.45.38}$$

It also normalizes the stabilizer of infinity and zero, the full group of nonzero scalar multiplications. Since $\eta(1) = 1$, this gives

$$\eta(ab) = \eta(a)\eta(b). \tag{D.45.39}$$

Thus it is an actual field automorphism. The field automorphisms of $k/\mathbf{F}_2$ are the five powers of squaring: the image of an element of degree five has at most five choices, and these five powers are distinct. This proves the full equality $\mathrm{Aut}(S) = S \rtimes C_5$ and the literal matrix generator t. It also proves the split outer quotient and its order, with no uncomputed automorphism enumeration. The same identification is stated in the opened primary account [Whi11, §3, preceding Lemma 3.1; Lemma 3.2]; the preceding argument supplies its exact group and field maps for the present source.

The fixed pro-five target and coupled generation. Take the actual D.37 quotient $\rho_n : \widehat{F} \twoheadrightarrow P_n$. For $q_1 = \beta_Q \rho_n$, its three base conjugators are exactly $1, f_Q, h_Q$ of D.37.14–19. Applying the proved theorem to any original epimorphism above ρ_n retains these specified values in (D.45.4). Every homomorphism from S^Ω to a finite five-group is trivial: its image is perfect, while a finite five-group is solvable. Hence every pro-five quotient of E factors through its displayed P_n, which is itself a five-group. This proves the exact maximal pro-five comparison, and retains all the D.37 parameters and all their powers.

More generally let $L \twoheadrightarrow D$ have solvable kernel, and fix an epimorphism of the same source into L with its three peripheral conjugators above the h_i. Pair that map with (D.45.4). The conjugating elements pair in the actual fibre product $E \times_D L$. The two projections are onto. Their possible missing common kernel quotient would be a quotient of S^Ω, hence perfect, and also a quotient of the solvable kernel of $L \to D$. It is therefore trivial. Goursat proves that the paired map is onto the full actual fibre product. □

The prime order in (D.45.1) is used in the independent-phase maps: every nonzero character value generates the outer image, and the exponent e in (D.45.15) is invertible. The remaining extension, Shapiro, subtraction and normalizer arguments use the stated split automorphism group and simplicity. The $\mathrm{PSL}_2(32)$ calculation supplies these hypotheses with the literal field automorphism. It does not assign outer coordinates by class preservation of β_Q.

The certificate `a_cyclic_outer_phase.py` retains every free-word map and its inverse, all 24 nonzero handle characters over $\mathbf{F}_5$, and an exact model of all 32736 matrices in S. It checks the sharp three-transitive action, all 33 S-classes, the six field orbits of length five, and the three S-classes in each of the four nonidentity outer cosets. It also gives ten actual surjective surface maps whose selected boundary traverses all five field phases, with the inverse source words retained. The companion `a_cyclic_outer_shapiro.py` uses the

actual order-125 Heisenberg base, its central order-five factor stabilizer, and every one of the five character extensions to an order-25 overgroup. It verifies both signs in (D.45.19), all intermediate embeddings with actual $\mathrm{Aut}(S)$ multiplication, and all the chosen base-conjugator comparisons. Its central character does not extend to the whole Heisenberg group, as witnessed by the displayed commutator; that nonextension is retained in the intermediate count.

4.28 An exact return-prime cover retaining the entire old pair

Statement D.46 (Proved). A strict mixed return cover with its complete generating peripheral fibre. *Let $\ell, p \geq 5$ be distinct primes, put $a = \mathrm{ord}_p(\ell)$, and choose $k = \mathbf{F}_{\ell^a}$ with an element ζ of order p. Indices below lie in $\mathbf{Z}/\ell$. Let c cyclically permute coordinates, put*

$$N = (\mathbf{F}_p)^\ell, \quad P = N \rtimes \langle c \rangle, \quad V = k^\ell, \quad H = V \rtimes P, \qquad n(v_i)_i = (\zeta^{n_i} v_i)_i. \tag{D.46.1}$$

These are specified actions, and the action of c on both N and V is the same cyclic shift. Retain the actual source $\widehat{B}, F, s, r, \tau, x, y, z$ of D.44. There is an epimorphism

$$q_H : F \twoheadrightarrow H, \qquad x \longmapsto X = (e_0, c), \quad y \longmapsto Y = (0, n_0), \quad z \longmapsto Z = (XY)^{-1}, \tag{D.46.2}$$

where n_0 is the first coordinate vector of N, and e_0 is the first coordinate vector of V. Let $E \to D$ be the actual six-coset induced images of $\widehat{B}$ obtained from q_H and its projection $q_P : F \to P$, respectively, as specified in (D.46.5) *below. Then*

$$1 \longrightarrow W = V^6 \longrightarrow E \longrightarrow D \longrightarrow 1, \qquad \dim_{\mathbf{F}_\ell} W = d = 6\ell a. \tag{D.46.3}$$

The D-module W is irreducible and nontrivial.

Put $K = O^p(O^\ell(F))$ and $\Gamma = \widehat{B}/K$. Every D.44 normalized automorphism of this entire Γ induces an automorphism of the specified D, and has a lift to the actual joint source

$$J = \Gamma \times_D E = \widehat{B}/(K \cap \ker q_E). \tag{D.46.4}$$

The lift fixes τ, retains the complete old normalized conjugator in Γ, and has its new conjugator in the image of F'. Thus it retains the entire R8 pro-ℓ pair and the chosen whole relative pro-p lift of D.44. The maximal pro-ℓ quotient of the pure subgroup of J is exactly P_ℓ, with the original action.

For the specified old automorphism, the complete literal peripheral fibre before fixing τ has ℓ^d elements. Every one generates the actual joint source and factors through an automorphism of that source. This fibre is one free W-conjugacy orbit. The same set is obtained with any prescribed three base conjugators, including the projections of prescribed old conjugators in Γ. Fixing τ gives precisely $\ell^{\ell a}$ literal normalized maps, a free transitive orbit of W^τ. These counts concern maps and their conjugator images; the complete profinite word fibres are the cosets described in (D.46.17).

The source in (D.46.4) *is an actual quotient of the return cover $\widehat{B}/O^\ell(O^p(O^\ell(F)))$, with a nontrivial finite kernel over Γ. In contrast, adjacent repetition of one residual has the exact identity $O^p(O^p L) = O^p L$ for every profinite group L.*

Proof. We give the source construction and all descent maps explicitly. Choose the six representatives t_i of B/F used in D.44, with $t_0 = 1$. For $b \in \widehat{B}$, let $j = i \cdot \bar{b}$ in the regular right action of $\mathfrak{S}_3$, and put

$$q_E(b) = \left((i \mapsto j), \left(q_H(t_i b t_j^{-1})\right)_i \right), \qquad q_D(b) = \left((i \mapsto j), \left(q_P(t_i b t_j^{-1})\right)_i \right). \tag{D.46.5}$$

The formula is first evaluated on the two literal generators s, r, and gives their continuous maps from the profinite free product. In fact $t_i b b' t_k^{-1} = (t_i b t_j^{-1})(t_j b' t_k^{-1})$; this proves the multiplication rule, the two source relations, and the displayed projections. The kernels are the corresponding normal cores of $\ker q_H$ and $\ker q_P$ in $\widehat{B}$. Both images retain the specified $\mathfrak{S}_3$ quotient. No additional quotient kernel is being presumed invariant under a changed source map.

The two-generator group and its residuals. The images of X, Y generate P: conjugating n_0 by powers of c gives every coordinate generator of N. Moreover $X^\ell = \sum_i e_i \neq 0$ in V. Acting on this vector by $n_0 - 1$ gives $(\zeta - 1)e_0$; powers of n_0, their $\mathbf{F}_\ell$-linear combinations, and the cyclic shift generate all of V, because $\mathbf{F}_\ell[\zeta] = k$. This proves the epimorphism (D.46.2). The same calculation proves that V is simple as an $\mathbf{F}_\ell[P]$-module. Indeed the distinct coordinate characters of the prime-to-ℓ subgroup N give its ℓ simple coordinate summands over $\mathbf{F}_\ell$; c permutes them transitively.

The following are equalities with their quotient maps:

$$O^\ell(H) = V \rtimes N, \qquad O^p(V \rtimes N) = V, \qquad O^\ell(V) = 1, \qquad O^\ell(P) = N, \quad O^p(N) = 1. \tag{D.46.6}$$

For the first equality, every finite ℓ-quotient kills N, and therefore kills $[V, N] = V$; the remaining quotient is the displayed C_ℓ. For the second, every finite p-quotient kills V, and the quotient by V is the displayed p-group N. The last two identities follow by the same source projections.

The action of X on V is the cyclic permutation, the action of Y changes only the first coordinate by ζ, and the action of XY has its ℓ-th power equal to ζI. Consequently

	X	Y	Z
order in H	ℓ^2	p	ℓp
order in P	ℓ	p	ℓp
$\dim_{\mathbf{F}_\ell} V^{(\cdot)}$	a	$(\ell-1)a$	0.

(D.46.7)

For XY, the operator $XY - 1$ is invertible since $(XY)^\ell = \zeta I$; hence the translation part of $(XY)^{\ell p}$ vanishes, giving its asserted order.

The entire induced kernel. The kernel of $E \to D$ is a submodule of V^6. The restrictions of its six ambient summands to the image of F are simple. Their fixed-space profiles on the same x, y, z are the six permutations of $(a, (\ell-1)a, 0)$, by the actual peripheral $\mathfrak{S}_3$ permutations in D.44. They are therefore pairwise nonisomorphic. For every summand one of $x^{\ell p}, y^{\ell p}, z^{\ell p}$ has nonzero projection in that summand and is killed in D: use the X coordinate in (D.46.7). Its normal closure surjects onto that simple summand. A submodule of a direct sum of pairwise nonisomorphic simple modules which surjects onto each summand is the whole direct sum. This follows, for example, by splitting a submodule of the semisimple module and using the absence of homomorphisms between distinct simple summands; it does not require the whole ambient group algebra to be semisimple. Thus the kernel is precisely $W = V^6$. The $\mathfrak{S}_3$ quotient permutes these six simple summands transitively, so W is simple for D. In particular

$$W^D = 0, \qquad W_D = 0, \qquad [E_F, W] = W \subset E'_F, \quad E_F = \operatorname{im}(F \to E). \tag{D.46.8}$$

The action of s on the six summands has three two-cycles, and the action of r has two three-cycles. Their powers are literally the identity, so their invariant dimensions are $d/2, d/3$.

The action of τ has three two-cycles and $\tau^2 = x^{-1}$. On a two-cycle its fixed vectors are exactly the fixed vectors of x on one chosen summand. Formula (D.46.7) gives

$$\dim W^s = 3\ell a, \qquad \dim W^r = 2\ell a, \qquad \dim W^\tau = \ell a. \tag{D.46.9}$$

Also the orders of τ in D, E are respectively $2\ell p, 2\ell^2 p$, by $\tau^2 = x^{-1}$ and (D.46.7).

All centralizer projections and all conjugator choices. For s, r, the maps from the upper centralizers to the lower centralizers are onto. A lift of a lower centralizer element conjugates the chosen upper element to another lift of the same cyclic subgroup; averaging over its order 2 or 3, which is invertible in $\mathbf{F}_\ell$, supplies the required kernel conjugator. Equivalently, this is the explicit coprime cyclic-complement argument of D.44.

For τ we verify the extra assertion directly. The local centralizer projections are

$$\begin{aligned} &C_H(X) \longrightarrow \langle c\rangle \subset C_P(c) = \langle c\rangle \times Z(P), \\ &C_H(Y) \twoheadrightarrow C_P(n_0) = N, \\ &C_H(Z) \twoheadrightarrow C_P(q_P(z)). \end{aligned} \tag{D.46.10}$$

The image in the first line is exactly $\langle c\rangle$. Indeed an element of $C_P(c)$ has its N-coordinate constant, say b; summing the coordinates of the translation equation for commuting with X gives $1 = \zeta^b$, hence $b = 0$. Powers of X give every power of c. The second line has the literal commuting lifts $(0, n)$. For the third, $1 - Z$ is invertible on V; the translation equation therefore has one solution for each lower centralizer element.

The pure image in D retains the actual quotient $F^{\mathrm{ab}}/p \simeq \mathbf{F}_p^2$. The second abelian coordinate is the sum of the N-coordinates in the identity component of (D.46.5), and conjugating by s recovers the first. Conjugation by τ sends these coordinates (u, v) to $(u - v, -v)$, since it fixes x and interchanges the other two peripheral classes. An element of the pure image centralizing τ therefore has $v = 0$. In the identity component its projection to $C_P(c)$ has constant N-coordinate b with $\ell b = 0$ in $\mathbf{F}_p$, so $b = 0$. Its projection belongs to the image in the first line of (D.46.10).

For clarity, the passage from these local tests to the induced centralizer follows from the multiplication in (D.46.5). On a two-cycle of τ, choose one coordinate of a lift of a commuting element. Its partner coordinate is forced by the commutation equation; consistency of the second equation is precisely commutation of the chosen coordinate with τ^2. The three tests in (D.46.10) supply those coordinates. Their independent corrections are available because the kernel is the entire V^6. A general element of $C_D(\tau)$ has $\mathfrak{S}_3$-projection either 1 or $\bar{\tau}$; multiplication by τ^{-1} reduces the latter to the pure case just checked. We have proved the three actual epimorphisms

$$C_E(s) \twoheadrightarrow C_D(s), \quad C_E(r) \twoheadrightarrow C_D(r), \quad C_E(\tau) \twoheadrightarrow C_D(\tau). \tag{D.46.11}$$

Their kernels are respectively the three spaces in (D.46.9).

The complete generating fibre. Apply the explicit product equation D.23.8 to (s, r, τ^{-1}). Since $W_D = 0$, its correction map from W^3 onto W is surjective for every prescribed triple of lower conjugators. Its solution set in conjugators has size $|W|^2$. Dividing only by the three kernel centralizers gives

$$\frac{|W|^2}{|W^s||W^r||W^\tau|} = |W|. \tag{D.46.12}$$

The epimorphisms (D.46.11) show that allowing all lower centralizer choices gives the same set, rather than enlarging it: every such change has a lift already centralizing the corresponding upper peripheral element. Thus (D.46.12) is also the full ordinary conjugator-union count.

Every solution generates E. Its image surjects onto D, and its intersection with the irreducible W is either W or zero. In the latter case the image would be a complement isomorphic to D. It cannot contain the prescribed conjugate of τ, whose upper order $2\ell^2 p$ exceeds its lower order $2\ell p$. This excludes the complement case. The stabilizer of such a generating map under kernel conjugation is $W^D = 0$. The count in (D.46.12) is therefore exactly one free orbit. Fixing τ cuts that orbit out by W^τ, and yields exactly $\ell^{\ell a}$ maps.

The pure rigidity that proves actual kernel preservation. We need an actual quotient automorphism, so we next prove the source-kernel comparison. Consider all peripheral epimorphisms $F \to H$ above the identity source map q_P. A conjugate of X over c has translation vector whose coordinate sum is one of the p values ζ^b; every vector with such a sum occurs. A conjugate of Y over n_0 has arbitrary translation in ke_0. Every lift of $q_P(z)$ is a conjugate of Z, because $1 - Z$ is invertible. Thus their total number is $p|k|^\ell$, and every one generates: its ℓ-th power at the first cusp gives a nonzero constant vector, whose P-orbit spans V.

These maps are exactly one conjugacy orbit under $\pi^{-1}Z(P)$. Conjugation by V first arranges the first translation to be $\zeta^b e_0$; the constant N-element removes ζ^b. The remaining $V^c = k\sum_i e_i$ conjugation changes the second translation by $(1-\zeta)ke_0$, and therefore sets it to zero. The centre of H is trivial, so this also accounts for the entire orbit. After X is fixed literally, the orbit is exactly the free V^c-orbit. In particular every such source map has precisely the kernel of q_H.

Let now $\phi : \widehat{B} \to E$ be any normalized solution in the preceding fibre, so $\phi(\tau) = \tau$. Its conjugator $f = s\phi(s)$ projects to the given element of D'_F. Since $W \subset E'_F$, it belongs to E'_F. The literal identities of D.44 give $\phi(x) = x$, $\phi(y) = f^{-1}yf$, and positive peripheral conjugacy at z. The image of f in P' belongs to the abelian subgroup N, which centralizes n_0. Let $\mathrm{pr}_0 : E_F \twoheadrightarrow H$ be the identity-coordinate projection of the actual induced image. Then $\xi = \mathrm{pr}_0 \phi|_F : F \to H$ lies over the identity map to P. The pure rigidity just proved gives $\xi = \mathrm{Inn}(h)q_H$.

Write $L = \ker q_E$, the normal core of $\ker q_H$. For $w \in L$ and a coset representative t, the element twt^{-1} belongs to $\ker q_H$. Moreover $\phi(t)q_E(t)^{-1} \in E_F$. Therefore the identity-component projection of $q_E(t)\phi(w)q_E(t)^{-1}$ is a conjugate of $\xi(twt^{-1}) = 1$. All six components of $\phi(w)$ vanish, proving

$$L \subset \ker \phi. \tag{D.46.13}$$

Thus every normalized solution factors through this same E, and is an automorphism because it is onto. Conjugating back by W proves the assertion for the whole peripheral fibre as well.

The old map and the exact common source. The source q_D kills K, by (D.46.6) in every coordinate, so $\Gamma \to D$ is an actual epimorphism. A D.44 normalized automorphism α of Γ induces the identity on the identity-component quotient P: its action is $x \mapsto x, y \mapsto f^{-1}yf$, and the image of $f \in F'$ lies in $P' \subset N$, centralizing n_0. It therefore preserves $\ker q_P$ and its six-coset normal core, inducing an actual $\alpha_D \in \mathrm{Aut}(D)$.

Residuals commute with continuous epimorphisms, by the exact proof D.43.7. Thus the projection of $q_E(K)$ to the identity component is $O^p(O^\ell(H)) = V$. On the other hand $q_E(K) \subset W$, since D is a quotient of Γ. It is a nonzero D-submodule of the irreducible W, so

$$q_E(K) = W. \tag{D.46.14}$$

The actual source image in $\Gamma \times E$ therefore contains the entire kernel $1 \times W$ and projects onto Γ. This proves (D.46.4) and identifies D as the full common quotient for these two source maps.

Each of the $\ell^{\ell a}$ normalized automorphisms β of E over α_D gives the automorphism

$$(g,e) \longmapsto (\alpha(g),\beta(e)) \quad \text{of } \Gamma \times_D E. \tag{D.46.15}$$

Its source map generates the whole fibre product. For prescribed old peripheral conjugators, choose the upper conjugators with their matching D-images, which is permitted by the complete fixed-conjugator calculation above; their pairs are actual conjugators in J. Also $[J_F, W] = W$ makes $J_F^{\mathrm{ab}} \to \Gamma_F^{\mathrm{ab}}$ an isomorphism. Thus $s\sigma(s)$ lies in J_F' and projects to the literal old conjugator. Conversely every normalized peripheral source map to J above the specified α gives one of the just classified maps to E; (D.46.13) then proves that it is exactly a map (D.46.15). This proves the full fibre count, including source generation and factorization.

The source kernel in (D.46.4) contains $O^\ell(K)$, because its quotient on K is the elementary abelian ℓ-group W. It is strictly smaller than K, by (D.46.14). The claimed strict return comparison is therefore the actual quotient diagram

$$\widehat{B}/O^\ell(K) \twoheadrightarrow J \twoheadrightarrow \Gamma, \qquad \ker(J \to \Gamma) = W \neq 1. \tag{D.46.16}$$

Its pure source kernel is contained in $O^\ell(F)$, so the original P_ℓ is a quotient of J_F. Every finite ℓ-quotient of J_F, composed with $F \to J_F$, factors through that same P_ℓ. This proves its asserted maximal pro-ℓ property even though the new kernel itself is an ℓ-group.

Finally, for each of the normalized conjugator images $f_J \in J_F'$, choose one preimage $\widetilde{f} \in F'$. Its entire word fibre is

$$\widetilde{f}\,\ker(F' \longrightarrow J_F'). \tag{D.46.17}$$

The epimorphism on closed commutator groups follows from the actual epimorphism $F \to J_F$. This states the word-to-image map separately from the finite number of image parameters.

For adjacent repetition, $L/O^p(O^pL)$ is an extension of the two pro-p groups $O^pL/O^p(O^pL)$ and L/O^pL, hence is pro-p. The defining maximal quotient and the reverse subgroup inclusion give $O^p(O^pL) = O^pL$. This identity and (D.46.16) specify both comparisons without asserting a lift to the entire return cover beyond the proved quotient J. □

The literal model program is `bc_return_prime_model.py`. Its full centralizer companion is `bc_return_prime_centralizers.py`. Together they construct the six-coset maps and retain actual word preimages for the whole new kernel, solve the complete peripheral product map, and lift a complete generating set of each base centralizer. At $(\ell,p) = (5,7)$, the new dimension is 180, the three invariant dimensions are $90, 60, 30$, and the base and upper τ-orders are $70, 350$. The full literal fibre has 5^{180} maps and its normalized fibre has 5^{30}. These are finite tests with all source and conjugator choices retained; the uniform proof above supplies every distinct pair $\ell, p \geq 5$.

4.29 All finite cyclic outer groups, with their actual automorphism extensions

Statement D.47 (Proved). Complete peripheral chief fibres for every cyclic outer group. *Let S be a finite nonabelian simple group for which*

$$\epsilon : \mathrm{Aut}(S) \longrightarrow \mathrm{Out}(S) = C_M \tag{D.47.1}$$

has cyclic target, where $M \geq 1$ is arbitrary. Identify S with its inner automorphisms. Let $1 \to A = S^\Omega \to E \xrightarrow{\pi} D \to 1$ be an actual finite chief extension, and fix an epimorphism $q_E : \widehat{F}(x_1,x_2) \twoheadrightarrow E$. Write $x_3 = (x_1x_2)^{-1}$, $X_i = q_E(x_i)$, $q_0 = \pi q_E$, and $d_i = q_0(x_i)$. For every epimorphism $q_1 : \widehat{F} \twoheadrightarrow D$, every specified $h_i \in D$ with

$$q_1(x_i) = h_i d_i h_i^{-1}, \tag{D.47.2}$$

and arbitrary lifts $H_i \in E$ of these h_i, the finite set

$$\mathcal{E}(q_1; H_i) = \left\{ f : \widehat{F} \twoheadrightarrow E : \pi f = q_1, \quad f(x_i) \in \{ a H_i X_i H_i^{-1} a^{-1} : a \in A \} \ (1 \le i \le 3) \right\} \tag{D.47.3}$$

has exactly the same positive cardinality as $\mathcal{E}(q_0; 1, 1, 1)$. This includes the complete ordinary peripheral fibre and every choice of its prescribed base conjugators. The actual extension (D.47.1) *is retained throughout.*

Consequently every D.37 automorphism β_Q on every specified P_n lifts through every chief extension of this kind above P_n, with the literal conjugators $1, f_Q, h_Q$. Its maximal pro-five quotient is the given P_n, and the induced map there is β_Q. Any additional prescribed epimorphism to a finite group over D with solvable kernel is retained through the actual fibre product.

Proof. We prove the composite-order source maps first, and then give the exact extension, boundary, subtraction, and generation maps.

Representatives in the actual outer extension. Choose an element $t \in \mathrm{Aut}(S)$ with $\epsilon(t) = 1 \in C_M$. For $a \in C_M$, choose the integer representative $\widetilde{a} \in \{0, \dots, M-1\}$, and put

$$L(a) = t^{\widetilde{a}}, \qquad L(0) = 1. \tag{D.47.4}$$

The elements $L(a)L(b)L(a+b)^{-1}$ belong to the actual subgroup S. They remain in all products below. Every change of coordinates will use a displayed element $L(a)$ and its actual inverse. For a subgroup $H \le C_M$, write $\widetilde{S}_H = \epsilon^{-1}(H)$. If $|H| = m$, its generator M/m has the actual lift $t_H = t^{M/m}$; identify H with C_m using that generator.

The effective phase of a boundary. Use the oriented surface presentation of D.45.6, with $b \ge 2$, and let $\chi : \Pi \twoheadrightarrow C_m$ be a surjective character. Write $j_i = \chi(c_i)$. An automorphism preserving χ and every positive boundary class has $\theta(c_i) = u_i c_i u_i^{-1}$. Its effective phase is

$$[\chi(u_i)] \in C_m / \langle j_i \rangle. \tag{D.47.5}$$

This quotient retains exactly the ambiguity in the based path. Each boundary is primitive in the free group obtained by eliminating one other boundary. Its centralizer is its cyclic subgroup: in a free basis containing that boundary, a reduced word with a letter outside that cyclic subgroup cannot commute with the basis letter. Thus two conjugating words differ by a boundary power, proving (D.47.5) is well-defined. For character-preserving automorphisms the effective phases add under composition. Under a change of surface coordinates, transport χ with that change; the phases of the change and its inverse then cancel in these quotients.

The same quotient is exactly what is needed for target conjugacy sets. For $a \in \widetilde{S}_H$ with outer coordinate $j \in C_m$, and $k \in \langle j \rangle$, choose an integer v with $vj = k$ in C_m. Then $t_H^k a^{-v} \in S$, and the actual equality

$$t_H^k a t_H^{-k} = (t_H^k a^{-v})\, a\, (t_H^k a^{-v})^{-1} \tag{D.47.6}$$

shows that every S-conjugacy invariant subset of that outer coset is fixed by the phase subgroup $\langle j \rangle$. No order relation for t_H was used.

Every effective phase is realized by source automorphisms. For one handle and a boundary c, retain the exact maps P and T on $[a,b]c$:

$$\begin{aligned} &P(a)=c^{-1}ac, \quad P(b)=c^{-1}a^{-1}cabc, \quad P(c)=c^{-1}a^{-1}cac,\\ &T(a)=a, \qquad\quad T(b)=ba, \qquad\quad T(c)=c. \end{aligned} \tag{D.47.7}$$

Both fix $[a,b]c$ literally. D.45.8–9 give the explicit Nielsen inverse of P, and $T^{-1}(b)=ba^{-1}$. They commute. Moreover P fixes $(ac)^{-1}$, and its restriction to $\langle a,c\rangle$ is conjugation by that word. Hence for all integers N,k, the map $\theta=P^N\circ T^k$ is an actual automorphism, with inverse $T^{-k}\circ P^{-N}$, and

$$\begin{aligned} \chi(\theta(a))&=\chi(a), \qquad \chi(\theta(c))=j,\\ \chi(\theta(b))&=\chi(b)+Nj+k\chi(a),\\ \theta(c)&=(ac)^{-N}c(ac)^N. \end{aligned} \tag{D.47.8}$$

Fix a prime power $\ell^e\|m$, and let $N\in\mathbf{Z}$ be a CRT idempotent: it is 1 modulo ℓ^e and 0 modulo every other primary factor of m. If $\ell\nmid j$, the entire ℓ-primary subgroup of C_m lies in $\langle j\rangle$, so this component already acts trivially by (D.47.6). Suppose $\ell\mid j$. Surjectivity of χ supplies a handle or boundary value which is a unit modulo ℓ.

First let a handle supply such a value. If its a-value is a unit, put $a_0=\chi(a)$. If only its b-value is a unit, use the exact change $a\mapsto ab$, $b\mapsto b$ and its inverse; the new a_0 is $\chi(a)+\chi(b)$, which is a unit modulo ℓ. In these coordinates choose

$$k\equiv -ja_0^{-1} \pmod{\ell^e}, \qquad k\equiv 0 \pmod{m/\ell^e}. \tag{D.47.9}$$

Equations (D.47.8)–(D.47.9) give $Nj+ka_0=0$ in C_m, so θ preserves the entire character. Its phase at c is $-N(a_0+j)$. That phase is a unit at ℓ and is zero at all other primary factors. Every other boundary is fixed literally. Powers of θ therefore supply every ℓ-primary effective phase at c.

If a boundary d supplies the unit, put $d_0=\chi(d)$. The exact Hurwitz square of D.45.11–12, raised to the N-th power, conjugates c by $(cd)^N$. It conjugates d by $(cd)^Nd^{-N}$. These literal words give phases

$$N(j+d_0) \quad \text{at } c, \qquad Nj \quad \text{at } d. \tag{D.47.10}$$

The first is a unit in the required primary component and zero elsewhere. The second belongs to $\langle d_0\rangle$: its other primary components vanish, and d_0 is a unit in the remaining component. Thus its effective phase at d is zero. The map preserves χ, fixes all other boundaries, and its powers supply the required component at c.

The positive boundary swaps and handle-block swaps in D.45.11 and D.45.13 put any selected boundary and supplier in these positions. Pull back by their explicit inverses. The comparison following (D.47.5) retains the effective phases in the original presentation. This proves surjectivity of the effective-phase map onto $\prod_i C_m/\langle j_i\rangle$, using finite compositions of actual surface automorphisms.

It follows that for arbitrary S-conjugacy invariant sets $T_i\subset\widetilde{S}_H$ in the specified outer cosets, independently replacing them by $t_H^{k_i}T_it_H^{-k_i}$ preserves the number of epimorphisms $\widehat{\Pi}\twoheadrightarrow\widetilde{S}_H$ with outer character χ and those boundary conditions. Indeed precomposition by the constructed automorphism gives the bijection: its conjugating word has image in $St_H^{k_i}$, up to the harmless subgroup in (D.47.6). Precomposition by the explicit inverse reverses the bijection and preserves surjectivity.

Matching composite cyclic characters exactly. Two oriented pants covers of the same index, with the matched boundary cycles below, have the same genus. Choose standard handle presentations and use the positive boundary swaps to match their boundary classes. Consider two characters to C_m which have the same image C_m and the same values $j_1, \dots, j_b$ on those matched boundaries. Define

$$d = \gcd(m, \widetilde{j}_1, \dots, \widetilde{j}_b), \qquad (j_1, \dots, j_b) = d(\mathbf{Z}/m) \subset \mathbf{Z}/m. \tag{D.47.11}$$

In genus zero the characters already agree. In positive genus, their handle covectors reduce modulo d to primitive covectors. Over each local ring $\mathbf{Z}/\ell^a$ dividing d, elementary symplectic operations carry a primitive covector to the first basis covector: place a unit entry in the first symplectic pair, clear the other entries, and scale that pair by $\mathrm{diag}(u, u^{-1})$. The elementary factorization in D.41 realizes the scaling as well. All these elementary operations are powers of the actual handle Dehn twists listed there. Choose their parameters by CRT to be the required ones in the selected primary factor and zero in the others. Thus a finite product of integral surface maps matches the handle covectors modulo d, while fixing every boundary.

The remaining difference in each handle coordinate lies in the ideal generated by the j_i in (D.47.11). The bounding-pair map for a basis curve a and a boundary c_i acts on homology by $v \mapsto v + \langle v, a\rangle c_i$, as in D.45.15. Its powers change the dual handle-character coordinate by arbitrary multiples of j_i, and fix all the other handle coordinates and the boundary values. An integral linear combination expressing the required difference in the ideal $(j_1, \dots, j_b)$ therefore corrects that coordinate exactly. Doing this for each handle matches the characters in C_m. These are actual based surface maps, with inverse maps. Their additional boundary phases are all removed by (D.47.5)–(D.47.10). For the trivial image the characters agree without this correction. This proves the required epimorphism-count comparison for every cyclic image, including every proper subgroup of C_M.

The given chief extension and each intermediate embedding. Let D_* be a factor stabilizer and use the right action $\Omega = D_* \backslash D$. Centerlessness gives the exact isomorphism

$$E \longrightarrow \mathrm{Aut}(A) \times_{\mathrm{Out}(A)} D, \qquad e \longmapsto (\mathrm{Ad}_e, \pi(e)). \tag{D.47.12}$$

Injectivity follows from $Z(A) = 1$; for surjectivity, choose a lift of the prescribed base element and correct its automorphism by the unique required inner element of A. Write the outer action as the actual cocycle $b_d(\omega) \in C_M$, with $b_{de}(\omega) = b_d(\omega) + b_e(\omega d)$. Choose $t_\omega \in D$, $t_{\omega_0} = 1$, and put

$$\begin{aligned} \epsilon_*(c) &= b_c(\omega_0) \quad (c \in D_*), & s_\omega &= b_{t_\omega}(\omega_0), \\ k(\omega, d) &= t_\omega d t_{\omega d}^{-1}, & b_d(\omega) &= -s_\omega + \epsilon_*(k(\omega, d)) + s_{\omega d}. \end{aligned} \tag{D.47.13}$$

The actual gauge $(z, d) \mapsto (L(s_\omega) z(\omega) L(s_{\omega d})^{-1}, d)$ carries E, and every specified X_i, H_i, to

$$E_* = \{(z, d) \in \mathrm{Aut}(S)^\Omega \rtimes D : \epsilon(z(\omega)) = \epsilon_*(k(\omega, d))\}, \qquad (z, d)(w, e) = (z(\omega) w(\omega d), de). \tag{D.47.14}$$

For multiplication, the adjacent factors $L(s_{\omega d})^{-1} L(s_{\omega d})$ cancel literally. This proves the map and its inverse using the actual representatives (D.47.4).

For each $D_* \le C \le D$, retain every character $\epsilon_C : C \to C_M$ extending ϵ_*. Choose R_ν representing $C \backslash D$, with $R_C = 1$. Define

$$\begin{aligned} k_C(\nu, d) &= R_\nu d R_{\nu d}^{-1}, & b_C(d, \nu) &= \epsilon_C(k_C(\nu, d)), \\ p(\omega) &= C t_\omega, & g_\omega &= \epsilon_C(t_\omega R_{p(\omega)}^{-1}). \end{aligned} \tag{D.47.15}$$

Let E_C be the group in (D.47.14) with $C\backslash D$ and the outer pattern b_C. There is an actual injection over D

$$j_C : E_C \hookrightarrow E_*, \qquad j_C(z,d)(\omega) = L(g_\omega) z(p(\omega)) L(g_{\omega d})^{-1}. \tag{D.47.16}$$

Its outer-coordinate identity is $\epsilon_*(k(\omega,d)) = g_\omega + b_C(d,p(\omega)) - g_{\omega d}$. The displayed factors in C prove this equality; adjacent inverse factors prove the homomorphism identity. Reading a coordinate above each ν proves injectivity. Also $g_{\omega_0} = 0$.

The actual source and every boundary root condition. Put $K_j = \ker q_j$ and $H_{j,C} = q_j^{-1}(C)$. For words T_ω with $q_j(T_\omega) = t_\omega$ and $T_{\omega_0} = 1$, twisted Shapiro is the explicit map

$$\begin{aligned} &\varphi : H_{j,D_*} \longrightarrow \mathrm{Aut}(S), \qquad \epsilon\varphi = \epsilon_* q_j, \\ &z_\varphi(g)(\omega) = \varphi(T_\omega g T^{-1}_{\omega q_j(g)}), \qquad f_\varphi(g) = (z_\varphi(g), q_j(g)). \end{aligned} \tag{D.47.17}$$

Middle words cancel in every product. Conversely evaluate the cocycle at ω_0 on H_{j,D_*}. The normalizing vector $v(\omega) = z(T_\omega)(\omega_0)$ belongs to S^Ω, has $v(\omega_0) = 1$, and gives $z_\varphi(g)(\omega) = v(\omega) z(g)(\omega) v(\omega q_j(g))^{-1}$. These formulas are inverse normalizations, so each surface map has exactly $|S|^{|\Omega|-1}$ literal gauges. They preserve the boundary A-classes and generation. The analogous construction on $C\backslash D$, followed by j_C, restricts at ω_0 to the very same surface map on H_{j,D_*}: here $g_{\omega_0} = 0$ and D_* fixes that coordinate.

Conjugacy by A for two elements above the same base element is equivalent to S-conjugacy of every permutation-cycle product. Indeed the successive conjugator equations determine each next coordinate from the preceding one; their outer coordinates vanish, and the last equation is precisely the cycle-product condition. For a boundary δ on $C\backslash D$, choose initial coordinate ν, and coordinates ω over it for all fine cycles γ of relative degree m_γ. The exact enlarged boundary subset $T_{j,C,\delta}$ is the intersection

$$\begin{aligned} a \in \widetilde{S}_{\mathrm{im}\,\epsilon_C}, \qquad \epsilon(a) = \epsilon_C(q_j(c_\delta)), \\ L(g_\omega) a^{m_\gamma} L(g_\omega)^{-1} \in \mathcal{C}_{j,\gamma} \quad \text{for every } \gamma. \end{aligned} \tag{D.47.18}$$

The classes $\mathcal{C}_{0,\gamma}$ come from the actual cycle products of X_i; those for $j = 1$ come from $H_i X_i H_i^{-1}$. These subsets retain all roots and every empty intersection.

The actual boundary bijection between the two covers is

$$Cd\langle q_1(x_i)\rangle \longmapsto Cdh_i\langle d_i\rangle. \tag{D.47.19}$$

It preserves lengths and every relative degree. Take the old initial coordinate corresponding to a new ν' to be $\nu' h_i$. There is an actual element $\mathcal{L}_i \in E_C$ above h_i, with components $\mathcal{L}_i(\nu) = L(b_C(h_i,\nu))$. Its image under j_C and H_i differ by an element of A. Conjugating by $\mathcal{L}_i$ therefore gives the exact equality of complete boundary subsets

$$T_{1,C,\delta'} = L(b_C(h_i,\nu'))\, T_{0,C,\delta}\, L(b_C(h_i,\nu'))^{-1}. \tag{D.47.20}$$

On a cycle product the conjugator is precisely this single component of $\mathcal{L}_i$. Its inverse proves the reverse inclusion, including the intersections in (D.47.18). The argument allows arbitrary given X_i; membership of X_i in $j_C(E_C)$ is not required for the conjugacy comparison.

Both source characters $\epsilon_C q_j|_{H_{j,C}}$ have the same image $\mathrm{im}\,\epsilon_C$, since $q_j(H_{j,C}) = C$, and their matched boundary values agree. Their standard surface presentations have the same genus by (D.47.19) and the degree formula. Apply the exact character-matching maps above, then the independent effective-phase maps to (D.47.20) and to the paths in the matching isomorphism. Every such phase lies in $\mathrm{im}\,\epsilon_C$. This proves equality, for $j = 0, 1$, of the numbers of epimorphisms

$$H_{j,C} \twoheadrightarrow \widetilde{S}_{\mathrm{im}\,\epsilon_C} \tag{D.47.21}$$

with their exact outer character and all conditions (D.47.18).

Subtracting the identical finite-base maps. For an epimorphism in (D.47.21), its K_j-image is normal in the target and contained in S, hence is either 1 or S. The maps with image 1 are exactly

$$\varphi = \Phi q_j|_{H_{j,C}}, \qquad \Phi : C \twoheadrightarrow \widetilde{S}_{\operatorname{im}\epsilon_C}, \qquad \epsilon\Phi = \epsilon_C. \tag{D.47.22}$$

Use the same actual Φ on both sides. Its Shapiro section over D is the homomorphism

$$s_\Phi : D \longrightarrow E_C, \qquad s_\Phi(d)(\nu) = \Phi(R_\nu d R_{\nu d}^{-1}), \quad \text{with base coordinate } d. \tag{D.47.23}$$

It is independent of q_j. Equations (D.47.2) and (D.47.23) show that $j_C s_\Phi(q_1(x_i))$ is the conjugate of $j_C s_\Phi(d_i)$ by $j_C s_\Phi(h_i)$. This conjugator and H_i have the same base value, so their ratio belongs to A. Thus the old boundary condition holds for this Φ exactly when the new one does. This is a bijection of the precise subtracted sets, and it retains all character-extension choices. Subtracting their equal finite cardinalities from (D.47.21) gives equal counts of maps with $\varphi(K_j) = S$, for every actual intermediate C and every extending ϵ_C.

Normalizer inversion and the entire generating fibre. For a surface map on H_{j,D_*} with K_j-image S, let $N_\varphi = \ker(\varphi|_{K_j})$. Its normalizer contains H_{j,D_*} and K_j, so it equals $H_{j,C}$ for some $C \geq D_*$. The normalizer contains $H_{j,C}$ precisely when φ extends to $H_{j,C} \to \operatorname{Aut}(S)$. Conjugation on $K_j/N_\varphi \simeq S$ constructs this extension. It agrees with φ on H_{j,D_*} and is unique: an automorphism centralizing all inner automorphisms of the centerless group S fixes every element of S. Its outer coordinate factors through one unique extending character $\epsilon_C : C \to C_M$. Its image is the full actual preimage $\widetilde{S}_{\operatorname{im}\epsilon_C}$, because its K_j-image is S. Conversely such an extension normalizes N_φ.

Let $M_{j,C}$ be the sum, over every extending ϵ_C, of the retained counts after (D.47.22). The literal restriction comparison for j_C proves that this counts exactly those original surface maps whose kernel normalizer contains $H_{j,C}$. Uniqueness makes the summands disjoint. The preceding paragraphs give $M_{0,C} = M_{1,C}$. Finite subgroup-poset inversion therefore gives

$$e_j = \sum_{D_* \leq C \leq D} \mu(D_*, C) M_{j,C}, \qquad e_0 = e_1, \qquad |\mathcal{E}(q_j; H_i)| = |S|^{|\Omega|-1} e_j, \tag{D.47.24}$$

where on the old side the conjugators are 1. To verify the last equality, minimal normalizer means that the conjugate coordinate kernels on K_j are pairwise distinct. The K_j-image is a subdirect product of the simple factors. A proper such product identifies two factors and their coordinate kernels: apply the two-factor Goursat lemma inductively, using that every normal subgroup of a product of nonabelian simple groups is a product of factors. Thus minimal normalizer gives exactly the full S^Ω-image. Together with the onto base map this is surjectivity to E. Conversely every onto map has precisely these coordinate images and distinct kernels. The gauge factor is the literal one from (D.47.17). The given epimorphism q_E has these properties, so $e_0 > 0$. This proves the complete fibre statement.

For completeness the ordinary conjugator union also has an exact comparison. Let $\mathcal{C}_i$ be the A-class of X_i, and let r_i be the size of its orbit under $C_D(d_i)$. This action is defined by conjugating with any lift of the centralizer element; two such lifts differ by A, so the action on A-classes is well-defined. Conjugation by H_i identifies that orbit with the A-classes of all E-conjugates of X_i above $q_1(x_i)$. Every class is obtained by replacing the specified base conjugator h_i by $h_i c_i$, with $c_i \in C_D(d_i)$. Different triples of these A-classes give disjoint literal source fibres. The established result applies to every one of these triples and gives the same cardinality $|S|^{|\Omega|-1} e_0$. Consequently the complete ordinary peripheral epimorphism fibre above either base map has cardinality $r_1 r_2 r_3 |S|^{|\Omega|-1} e_0$. This proves the stated full-union assertion with all base centralizer choices retained.

The specified pro-five and coupled sources. Use the actual D.37 source $q_0 = \rho_n$, $q_1 = \beta_Q \rho_n$ and its displayed conjugators $1, f_Q, h_Q$. Equation (D.47.3) supplies the asserted generating lift above those exact maps. Every homomorphism from S^Ω to a finite five-group is trivial: its image is both perfect and a finite five-group, while every nontrivial finite five-group has nontrivial abelianization. Hence every such homomorphism from E factors uniquely through P_n, proving the stated maximal pro-five quotient with its actual induced action.

More generally, let $L \twoheadrightarrow D$ have solvable kernel B, and retain a given epimorphism $q_L : \widehat{F} \twoheadrightarrow L$ above q_1. Combine the constructed f and q_L in $E \times_D L$. The K_1-image projects onto both S^Ω and B. Its Goursat common quotient is a solvable quotient of the perfect group S^Ω, hence is trivial. Thus that image is $S^\Omega \times B$. The base image is D, so the combined source is onto the actual fibre product and retains q_L literally.

Exact composite and nonsplit checks. The program `a_cyclic_outer_composite.py` retains all forty literal corrected push maps and their inverses used in the twelve recorded moduli. It verifies the complete nonunit-character and boundary-supplier formulas, and all 359268 genus-one handle/boundary characters in its orbit checks. The proof above supplies the arbitrary modulus and genus comparison.

The independent program `a_cyclic_outer_nonsplit.py` uses the actual quotient $Q_{16} \times C_3 \to C_6$ with kernel Q_8. In coordinates its law and projection are

$$\begin{aligned}(a,b,c)(x,y,z) &= (a + (-1)^b x + 4by \bmod 8,\ b+y \bmod 2,\ c+z \bmod 3),\\ \epsilon(a,b,c) &= 3(a \bmod 2) + 4c \pmod 6.\end{aligned} \tag{D.47.25}$$

The chosen lift $(1,0,1)$ has order 24; every lift of $1 \in C_6$ has order 12 or 24, so this quotient has no section. Over the actual Heisenberg base $\mathrm{Heis}(\mathbf{Z}/6)$, the program checks all 46656 outer-cocycle products, the six intermediate characters, the actual gauge and embedding multiplications, 804 nontrivial kernel carries, and 6084 literal cycle-product transports. This verifies precisely the nonsplit group identities used above; the simple-kernel generation is proved by the displayed normalizer and Goursat comparisons. Both programs, their outputs, JSON data, and the nonsplit matrix certificate are retained in `code/`. □

4.30 The full integral affine return tower

Statement D.48 (Proved). An entire return-prime lattice tower with its literal inverse-limit fibre. *Retain the distinct primes $\ell, p \geq 5$, the integer $a = \mathrm{ord}_p(\ell)$, and the actual sources Γ, D of D.46. Let $\mathcal{O}$ be the unramified extension of $\mathbf{Z}_\ell$ of degree a, with the lift $\widetilde{\zeta}$ of the specified order-p element of its residue field. For $1 \leq n < \infty$, put $\mathcal{O}_n = \mathcal{O}/\ell^n\mathcal{O}$, and write $\mathcal{O}_\infty = \mathcal{O}$. Use the same cyclic shifts and diagonal actions to form*

$$V_n = \mathcal{O}_n^\ell, \quad H_n = V_n \rtimes P, \quad q_{H_n}(x) = (e_0, c), \quad q_{H_n}(y) = (0, n_0). \tag{D.48.1}$$

Let E_n be the actual closed six-coset image of $\widehat{B}$, using precisely D.46.5, and put

$$W_n = V_n^6, \qquad J_n = \Gamma \times_D E_n. \tag{D.48.2}$$

Then all these maps are epimorphisms, and

$$\begin{gathered}\ker(E_n \to D) = \ker(J_n \to \Gamma) = W_n,\\ E_n = E_\infty/\ell^n W_\infty, \qquad J_n = J_\infty/\ell^n W_\infty, \qquad J_\infty \simeq \varprojlim_n J_n.\end{gathered} \tag{D.48.3}$$

The quotient equalities are for finite n; the kernel identity also includes $n = \infty$. The equalities retain the actual maps from $\widehat{B}$. In particular, for $K = O^p(O^\ell(F))$, there is an actual epimorphism

$$\widehat{B}/O^\ell(K) \twoheadrightarrow J_\infty \twoheadrightarrow \Gamma, \qquad \ker(J_\infty \to \Gamma) = \mathcal{O}^{6\ell}. \tag{D.48.4}$$

Fix any one entire normalized old D.44 automorphism α of Γ, including its literal conjugator and prescribed old peripheral conjugators. It has a normalized peripheral automorphism on this whole J_∞ retaining all that data. The maximal pro-ℓ pure quotient is still the original P_ℓ, with the specified original pair. Thus the R8 weight-nine realization and the chosen whole relative pro-p lift are retained through this complete affine return tower.

Let $\mathscr{X}_n$ be the full literal fibre of normalized peripheral source maps to J_n above this same old α, with τ fixed. Every map in the fibre is generating and factors through an automorphism of this same J_n. After choosing one origin σ_0 in the infinite fibre, the actual maps give compatible torsor isomorphisms

$$\begin{aligned} W_\infty^\tau \simeq \mathcal{O}^\ell &\longrightarrow \mathscr{X}_\infty, \quad w \longmapsto \mathrm{Inn}(w)\sigma_0, \\ W_n^\tau \simeq \mathcal{O}_n^\ell &\longrightarrow \mathscr{X}_n, \quad \bar{w} \longmapsto \mathrm{Inn}(\bar{w})(\sigma_0 \bmod \ell^n), \\ \mathscr{X}_\infty \simeq \varprojlim_n \mathscr{X}_n, &\quad |\mathscr{X}_n| = \ell^{n\ell a}. \end{aligned} \tag{D.48.5}$$

Every finite normalized map lifts to the entire compatible tower. Before fixing τ, the full peripheral fibre is a torsor under W_n, with $\ell^{6n\ell a}$ literal maps at finite precision. These are the full ordinary centralizer unions and also the fibres with prescribed old conjugators; no conjugator freedom is discarded. The exact word-to-image fibres are specified in (D.48.12) *below.*

Proof. We give integral source maps before using their finite reductions. The derivative of $T^p - 1$ at the chosen residue root is a unit, so its root has a unique lift in the unramified degree-a extension. This gives $\widetilde{\zeta}^p = 1$ literally, and $\widetilde{\zeta}^b - 1$ is a unit whenever $b \not\equiv 0 \pmod p$. The coefficient ring is $\mathcal{O} = \mathbf{Z}_\ell[\widetilde{\zeta}]$. Thus every local multiplication in D.46 is defined over this actual ring and reduces compatibly to $\mathcal{O}_n$.

The closed subgroup generated by the two elements in (D.48.1) maps onto P. The ℓ-th power of the first is the constant vector, whose conjugates span $\mathcal{O}^\ell$: apply the unit $\widetilde{\zeta} - 1$ in one coordinate and then use powers of $\widetilde{\zeta}$ and the cyclic shift. Integer powers and their $\mathbf{Z}_\ell$-limits supply these coefficients inside the closed source image. This proves q_{H_∞} is onto and supplies all its finite quotient maps.

The actual whole kernel and all reductions. The source E_∞ is a closed subgroup of $H_\infty^6 \rtimes \mathfrak{S}_3$, and its kernel over D lies in $\mathcal{O}^{6\ell}$. Its reduction modulo ℓ is the entire kernel of D.46.3. One can see this with actual words: the conjugates of the three cusp powers of exponent ℓp retained in D.46 have a basis of the modulo-ℓ kernel among their vector images. Lift those same words to E_∞. The determinant of their vector matrix is a unit in $\mathbf{Z}_\ell$, so their $\mathbf{Z}_\ell$-linear span is the whole $\mathcal{O}^{6\ell}$. They commute inside this kernel, and continuous powers realize all those coefficients in the actual closed subgroup. Hence its kernel is $W_\infty = \mathcal{O}^{6\ell}$. The same conclusion follows from the closed-module form of Nakayama's lemma, applied to this specified submodule of a finite free $\mathbf{Z}_\ell$-module.

Reduction of all six components now has kernel exactly $\ell^n W_\infty$, which proves the asserted E_n maps in (D.48.3). Every finite image and every transition is therefore the image of the same two source generators. The commutator map satisfies

$$[E_{\infty,F}, W_\infty] = W_\infty, \qquad W_n^D = 0 \quad (1 \le n \le \infty). \tag{D.48.6}$$

For the first equality its reduction modulo ℓ is onto by D.46.8. The sum of the images of the two integral operators for $x-1, y-1$ is a closed submodule and has a unit full-rank minor, hence is all of W_∞. For the second, a fixed vector at finite precision reduces to zero modulo ℓ by D.46.8; divide by ℓ and induct on the precision. The infinite assertion follows from all these reductions.

Integral centralizer equations and the entire union. The local equations used in D.46 remain exact over every $\mathcal{O}_n$. For the cyclic shift C on $\mathcal{O}_n^\ell$,

$$\ker(1-C) = \mathcal{O}_n \sum_i e_i, \qquad \operatorname{im}(1-C) = \{(v_i) : \textstyle\sum_i v_i = 0\}. \tag{D.48.7}$$

Both equalities follow by successively solving the coordinate differences, and use no division by ℓ. The Y fixed space consists of the vectors with first coordinate zero, because $\widetilde{\zeta}-1$ is a unit. The operator $1-Z$ is invertible: since $Z=(XY)^{-1}$, its linear action satisfies $Z^\ell = \widetilde{\zeta}^{-1} I$, and its inverse is $(1+Z+\cdots+Z^{\ell-1})/(1-\widetilde{\zeta}^{-1})$. The displayed denominator is a unit.

Summing the X centralizer equation still forces the constant N-coordinate to vanish, so the first local image in D.46.10 is again exactly $\langle c\rangle$. The Y and Z centralizer projections are still onto. The unchanged actual F^{ab}/p quotient and the two-cycle coordinate equations of D.46 then give

$$C_{E_n}(s) \twoheadrightarrow C_D(s), \quad C_{E_n}(r) \twoheadrightarrow C_D(r), \quad C_{E_n}(\tau) \twoheadrightarrow C_D(\tau) \quad (1 \le n \le \infty). \tag{D.48.8}$$

All corrections are in the actual full kernel W_n.

The invariant modules at s, r, τ are free over $\mathcal{O}_n$ of ranks $3\ell, 2\ell, \ell$, respectively. For τ, choose one coordinate in each of its three two-cycles. The three choices are a constant vector for X, a vector with first coordinate zero for Y, and zero for Z. The partner coordinates are prescribed by the actual τ action. This is an explicit compatible isomorphism $W_n^\tau \simeq \mathcal{O}_n \oplus \mathcal{O}_n^{\ell-1}$. In particular every invariant vector at finite precision lifts invariantly to W_∞.

Existence, full fibres, and generation over the ring. Choose lifts in E_∞ of the prescribed three conjugators in D. Their product defect lies in W_∞. The exact correction map D.23.8 is now a $\mathbf{Z}_\ell$-linear map $W_\infty^3 \to W_\infty$. Its reduction modulo ℓ is onto by D.46.12; a unit minor therefore gives an actual integral right inverse. Cancelling the defect produces a continuous peripheral source map $\widehat{B} \to E_\infty$. It factors through $\widehat{B}$ from the three-boundary source because its first two images are conjugate to the actual elements of orders two and three.

Every such source map is onto, at every precision. Its image projects onto D and contains the $2\ell p$-th power of the prescribed τ conjugate. This is a vector whose reduction in W_1 is nonzero. Its orbit spans the irreducible D-module W_1. The image's intersection with W_n is a closed D-invariant $\mathbf{Z}_\ell$-submodule; it therefore surjects onto W_1, and Nakayama gives that it is the whole W_n. This proves generation, including at $n=\infty$.

At finite precision the correction solutions in conjugators number $|W_n|^2$. Using the three free invariant modules just computed gives exactly $\ell^{6n\ell a} = |W_n|$ distinct maps. The surjections (D.48.8) identify this with the full ordinary conjugator union. By (D.48.6) the kernel-conjugation action is free, so the maps form exactly one W_n-orbit. The infinite fibre is also one free W_∞-orbit: after choosing one solution, its comparison with any other has a unique conjugator at each finite precision. Uniqueness makes these conjugators compatible, and their inverse limit gives the required element of W_∞. Fixing τ is precisely restriction to the invariant subgroup W_n^τ.

Actual automorphisms and the whole retained old quotient. The pure rigidity proof of D.46 works over $\mathcal{O}_n$ without a new equation. The first translation has coordinate sum $\widetilde{\zeta}^b$; equation (D.48.7) arranges it in the first coordinate, a constant N-element removes its scalar, and the remaining V_n^c action removes the second translation using the unit $1-\widetilde{\zeta}$. Every pure peripheral source map to H_n over the identity on P is consequently an inner conjugate of the specified source map.

For a normalized source map to E_n, (D.48.6) places its normalized conjugator in $E'_{n,F}$. Its identity coordinate is therefore a pure map of the preceding kind. The six actual conjugates in D.46.13 prove preservation of the same closed normal core, also when $n=\infty$. Thus the source map factors through an onto endomorphism of this same E_n. These groups are finitely generated profinite quotients of $\widehat{B}$, and hence Hopfian; the endomorphism is an automorphism. Kernel conjugation handles the full unnormalized fibre as well.

The image of the old source kernel K in E_∞ is the entire W_∞. It is contained there because D is an old quotient. Its reduction is all of W_1 by D.46.14. It is a closed $\mathbf{Z}_\ell$-submodule, so the same unit-basis argument gives

$$q_{E_\infty}(K) = W_\infty. \tag{D.48.9}$$

Consequently the actual joint source is the complete fibre product in (D.48.2). Componentwise automorphisms on Γ and E_n, agreeing on D, give its actual automorphisms, and every peripheral source map above the fixed old automorphism has this factorization. Pairing the old conjugators with upper conjugators having their exact D-images retains those whole old conjugators. Since $[J_{n,F}, W_n] = W_n$, the new normalized conjugator belongs to $J'_{n,F}$ and has precisely the given old image.

The source kernel of J_∞ contains $O^\ell(K)$, since its image on K is the pro-ℓ group in (D.48.9). This proves (D.48.4). The source kernel also lies in $O^\ell(F)$. Every finite ℓ-quotient of the new pure group, composed with its source map from F, factors through the original P_ℓ, which is itself a quotient of the new pure group. This proves the exact maximal pro-ℓ comparison and retains the original action on it.

The full inverse-limit and word fibres. The quotient maps in (D.48.3) identify each $\ell^n W_\infty$ as the literal source descent kernel. An automorphism above the specified old map preserves W_∞, and therefore preserves each of these characteristic subgroups of its kernel. All the displayed transition maps on full fibres are thus actual descent maps. With one infinite origin selected, their parameters are exactly the reductions of W_∞^τ. Every such finite parameter lifts, and a compatible collection has one inverse-limit parameter. This proves all comparisons in (D.48.5), including their topology: the continuous bijection from the compact parameter group to the closed source-map fibre is a homeomorphism.

Writing $f_0 = s\sigma_0(s)$ for the new infinite origin, the conjugator image at parameter $w \in W_\infty^\tau$ is the literal group expression

$$f(w) = (sws)f_0 w^{-1} \in J'_{\infty,F}. \tag{D.48.10}$$

Its old image is the specified old f, because w lies in the kernel over Γ. The parameterization is injective: equality of these conjugators would give the same normalized map on both s and τ, hence on s, r; its kernel stabilizer is zero by (D.48.6). At finite precision the same formula uses $\bar{w}$, and

$$\mathscr{X}_{n+1} \longrightarrow \mathscr{X}_n \quad \text{is the surjective reduction} \quad \mathcal{O}_{n+1}^\ell \to \mathcal{O}_n^\ell. \tag{D.48.11}$$

Finally each image in (D.48.10) has, after choosing one source preimage $\widetilde{f}(w) \in F'$, the complete profinite word fibre

$$\widetilde{f}(w)\,\ker\!\big(F' \longrightarrow J'_{\infty,F}\big). \tag{D.48.12}$$

The analogous expression applies at every finite precision. This separates the full word fibres from the affine lattice of conjugator images. □

The new program `bc_return_prime_integral.py` uses the literal cyclotomic coefficient rings at the two executed pairs $(\ell, p) = (5, 7), (7, 5)$. Here the full Φ_p is irreducible modulo ℓ, so it supplies the actual unramified integral ring without a root-choice identification. At precisions two and three the program replays all new-kernel words, checks a unit determinant for their integral vector basis, solves the full product map, checks every remaining centralizer-equation row without discarding nonunits, and lifts the complete base-centralizer generating sets with actual word coefficients. The normalized counts are $5^{60}, 5^{90}, 7^{56}, 7^{84}$, respectively. The proof supplies the exact full tower; these finite checks verify its stated integral source and equation conventions.

5 E. The complex-multiplication coset

5.1 The finite CM calculation and its independent verification

In this section $B = \langle s, r \mid s^2 = r^3 = 1\rangle$ is a discrete free product, $t = sr$, and $F = \ker(B \to S_3)$ is the discrete free group on $x = u^2$, $y = v^2$, where $u = t^{-1}$ and $v = sus$. Let $q : B \twoheadrightarrow Q = \mathrm{SL}_2(\mathbb{Z}/12)/\{\pm I\}$ send s, r to the classes of

$$S = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \qquad R = \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix}.$$

Write $N_B = \ker q$, $C = N_B^{\mathrm{ab}}$, and $K_j = [N_B, N_B]N_B^j$, so $G_j = B/K_j$. All these maps are homomorphisms of discrete groups; their maps to finite targets extend continuously to the profinite completions. The inverse image of K_j under $B_3 \twoheadrightarrow B$ is $\widetilde{N}_j$. The CM comparison is LM7 D.15–D.19 and D.25–D.26, in the setting of *La Longue Marche* §51.15, scans 156–170. The assertions below concern these specified finite coordinates.

Statement E.1 (Proved). Independent confirmation of Task E.1. *The order of Q is 576, and $C \simeq \mathbb{Z}^{97}$. With the CM sign convention of LM7 D.25, the independent calculation gives*

	CM compatible	extra CM coset
matrix pairs	24	24
cusp equation in G_2	12	12
cusp equation in G_3	24	24
certified automorphisms of G_6	12	12.

For $c = [t^{12}] \in C$, the integral maps

$$L : C \oplus C \to C, \quad L(b, a) = (1 - s)b + s(1 - r)a, \qquad \widetilde{L}(b, a, h) = L(b, a) - hc$$

have Smith forms $(1^{96}, 2)$ and (1^{97}), respectively. All 24 automorphism certificates have rank 97 on C/pC for $p = 2, 3$. After the exact integral cusp gauge, their two hexagons hold in G_6 and the resulting 24 automorphisms are distinct. In particular twelve have the extra CM coordinate. No discrepancy with these numerical assertions of R12 was found.

Proof. The independent verifier is `code/de_verify_cm.py`. It imports none of the source programs. It enumerates determinant-one matrices modulo twelve and takes their classes by simultaneous sign. It constructs the graph with edges Q and vertices $Q/\langle s\rangle \amalg Q/\langle r\rangle$. There are 576 edges and 480 vertices. The graph is connected, and the fundamental cycles for a

spanning tree form an integral basis of its homology; their coordinates on the 97 omitted edges form I_{97}. The Bass–Serre action identifies that homology with N_B^{ab}. It also identifies N_B with the free group of rank $576-480+1=97$, because N_B meets both conjugates of the finite factors trivially.

For a word $w \in B$, use its image $q(w)$ and integral path chain $P(w)$, with

$$P(s)=0, \quad P(r)=e_1-e_r, \quad P(ab)=P(a)+q(a)P(b).$$

Chains distinguish the elements of G_j after reduction modulo j: two lifts of the same element of Q differ by a cycle, which is zero modulo j precisely when their difference lies in K_j. Thus all group equalities tested in this model are equalities in G_j.

Let $H=q(F)$. Direct enumeration gives $|H|=96$, $|C_H(r)|=6$ and $|C_H(s)|=8$. The verifier lists all 48 pairs (A,β) in these two centralizers. It uses the unique lift of each projective matrix whose reduction modulo four is in the four-group generated by the reductions of u^2, v^2. For A^{-1} the action on the rational three-torsion line $\ker(R-I)$ gives one sign. The reduction of β^{-1} modulo three, lying in $\{\pm I, \pm S\}$, gives the other. Equality of the two signs is the CM-compatible class; their inequality is the extra class. These are exactly the restrictions to $M'=\mathbb{Q}(i,(3/4)^{1/4})$ used in LM7 D.25. The program computes 24 pairs of each sign.

Choose integral paths P_A, P_β lifting the pair. The cusp error for $s \mapsto \beta s \beta^{-1}$ and $r \mapsto ArA^{-1}$ is

$$E = P_\beta - sP_\beta + s(P_A - rP_A + AP(r) - P(r)).$$

Changing the paths by cycles b, a changes this error by $L(b,a)$. Exact Gaussian elimination over $\mathbb{F}_2$ and $\mathbb{F}_3$ gives the displayed solvability counts. The verifier then reads the stored source certificates as proposed vectors, substitutes them into this equation, and computes the induced map on all fundamental cycles. The edge at $1 \in Q$ maps to $P_A - P_\beta + e_A$; translating this formula by Q gives every other edge. The resulting matrices have the asserted ranks. They act invertibly on $C/6C$ and identically on Q, proving that each proposed endomorphism of G_6 is an automorphism.

To verify the commutator gauge independently, the program projects chains to the corresponding six-edge graph for S_3. Its two integral basis cycles are the images of x, y. Thus it computes the actual integral abelianization $\beta^{\mathrm{ab}} = ax + by$ of each chosen word. With $k=b-a$ and $\beta' = x^k\beta$,

$$f = s\beta' s(\beta')^{-1} \in F',$$

because conjugation by s exchanges x, y in F^{ab} and β' has equal coordinates. The program directly multiplies the words for both $m=0$ hexagons in G_6; all 48 equalities hold. It also compares the resulting images of s, r, obtaining 24 distinct automorphisms. Independently it finds

$$q(C_{G_2}(s) \cap \operatorname{im} F) = 1, \qquad q(C_{G_3}(r) \cap \operatorname{im} F) = 1,$$

by solving the corresponding centralizer equations. These checks account for the finite conjugator ambiguity in the recorded coordinates. The gauge replaces A, β, V by simultaneous left translates, so $T = A^{-1}V$ and $W = \beta^{-1}V$ are unchanged.

Finally the verifier computes the integer matrices of L and $[L, -c]$ in its newly constructed cycle basis. Exact Smith reduction gives the stated invariant factors; Hermite reduction of $[L, -c]$ gives I_{97}. The cycle c has relatively prime coordinates. The outputs retain every pair, rank, abelianization and gauge, together with the complete source certificate files. These integer calculations confirm more than the ranks of the maps modulo two and three. □

Statement E.2 (Open). The full profinite gentle CM image. *We have not determined whether* $\widehat{\mathrm{GT}}_0$ *reaches the extra CM coset. E.9 proves that* none *of the twelve particular*

extra shadows recorded by LM7 D.25 is genuine, using an explicit common finite refinement. The full existence question ranges over every shadow with the specified CM matrices. E.10 computes the affine parameters in those matrix fibres; the twelve recorded representatives do not exhaust them. Thus the asserted equivalence in Task E.2 between the full CM question and genuineness of one of those twelve representatives still requires a proof of that reduction. E.11 constructs new points in every extra CM matrix fibre that survive that same common refinement. E.15 constructs compatible families in all 24 matrix fibres through the entire relative class-two tower, with multiplier one. E.16 extends those families simultaneously through every finite collection of nonabelian simple quotients, and separately through each finite perfect quotient. E.20 further extends the class-two families simultaneously through every relative quotient with normal kernel of order prime to six, including its arbitrary higher commutator layers. The combined abelianization and the single cusp gauge are retained. The new specified comparison outside that family has a relative three-primary kernel with a nonzero third lower-central term. The unresolved set is the intersection of the genuine-shadow image with the whole *union of extra CM matrix fibres.*

Statement E.3 (Open). The full profinite GT CM image. *We have not determined whether full* $\widehat{\mathrm{GT}}$ *reaches the extra CM coset. The finite tests in E.6 impose consequences of the pentagon established in D.5. E.17 evaluates the complete group pentagon in the specified pure four-strand quotient of order* 5184*. It excludes fifteen of the twenty-four retained* G_6 *representatives, but its rank-one affine equation has actual generating solutions in every whole CM matrix fibre, simultaneously satisfying the class-two cusp obstructions of E.11. E.18 retains the next central Ferrari layer in a pure quotient of order* 15552*: its pentagon has the same zero sets on all whole affine fibres, and all twenty-four retained words still generate the order-*648 *Ferrari image. E.19 imposes the modulus-nine pentagon together with the complete three-strand hexagons on the exact induced core. Each retained point has one compatible correction in the full three-dimensional commutator correction image, with generation verified in both targets. The missing step is compatibility through all four-strand quotients. No positive or negative conclusion for Task E.3 follows from the surviving finite candidates. The primary comparison in [NTY10, Proposition 4.4 and Theorem 4.16] constructs the triangle-cover Tate action for full* $\widehat{\mathrm{GT}}$ *but identifies it with the normalizer h-coordinate on the Galois subgroup. The required comparison of these two actions on the entire source remains unproved here. The newer action of [LNS26, Theorem 1.1] is defined on the profinite mapping class groups of type* $(g, 0)$ *and* $(g, 1)$*; its Section 8 explicitly leaves compatibility with the indicated cutting homomorphisms unresolved for full* $\widehat{\mathrm{GT}}$*. It does not provide the needed CM comparison in the checked statements.*

5.2 The homology refinements and an additional finite quotient

Statement E.4 (Proved). The checked homology lifting mechanism and its scope. *The Smith computation in E.1 gives the successive corrections used in LM7 D.26. Each of its twelve extra CM pairs admits a compatible family of shadows through the quotients* G_j*, with a coherently chosen level-six cusp gauge and multiplier* $\lambda \equiv 1 \pmod{72}$*. The family of kernels* K_j *fails to be cofinal among the finite-index normal subgroups of B lying below* K_6*.*

Proof. The exact path equation for $t \mapsto t^{1+12h}$ is $E + L(b, a) - hc = 0$, because $q(t)c = c$. At $p = 2, 3$ start from an E.1 solution modulo p and $h = 0 \bmod p$. Surjectivity of $\widetilde{L}$ over $\mathbb{Z}$ cancels the divided error successively modulo p; this constructs solutions over $\mathbb{Z}_p$. Their matrices on $C \otimes \mathbb{Z}_p$ remain invertible, since their reductions modulo p are the verified matrices. They therefore give automorphisms of the extensions of Q.

For $p > 3$ choose the identity automorphism. The required matrix conjugators lift to centralizers: a lift of $A \in C_Q(r)$ has error z with $(1 + r + r^2)z = 0$, and $(1 - r)(2 + r) = 3 - (1 + r + r^2)$ corrects it after dividing by three. The analogous formula for s divides by two. Both divisions are valid in $C \otimes \mathbb{Z}_p$. Chinese remainders on C/jC assemble these automorphisms in

$$\mathcal{G} = \varprojlim_j G_j = \widehat{B}/\overline{[\widehat{N}_B, \widehat{N}_B]}.$$

The equality follows on each fibre above Q by completion of the integer cycle group C. The induced abelianization of the image of $\widehat{F}$ is $\widehat{\mathbb{Z}}^2$, since the displayed kernel lies in $\widehat{F}'$. One global cusp gauge thus makes f a commutator and gives compatible shadows. The congruences on h give the stated congruence modulo 72. A discrete representative matching the global conjugator both in G_6 and in $F^{\mathrm{ab}}/36$ gives its level-six gauge. This specifies the gauge; it does not assert compatibility for every independent choice of a finite representative.

We reran the original refinement program. Besides the exact integer normal forms, it checks the powers $2, 4, 8, 16$ and $3, 9, 27, 81$ and verifies 384 general-multiplier hexagon identities. These finite checks support the formula; the preceding correction argument supplies all powers. The quotient in E.5 has kernel contained in K_6 and does not kill $[N_B, N_B]$, whereas every G_j kills that subgroup. The factorization test for the quotient maps proves the stated failure of cofinality. □

Statement E.5 (Proved). All certified shadows survive an explicit A_5 refinement. *Define $a : B \twoheadrightarrow A_5$ by $a(s) = (12)(34)$ and $a(r) = (135)$, and put*

$$J_{A_5} = K_6 \cap \ker a.$$

The diagonal map induces an isomorphism $B/J_{A_5} \simeq G_6 \times A_5$. Its kernel does not contain $[N_B, N_B]$. Every one of E.1's 24 certified shadows extends to a shadow at the inverse image of J_{A_5} in B_3. All twelve extra CM coordinates survive this finite test.

Proof. The independent program `de_a5_refinement.py` enumerates the permutation group generated by the specified two elements. It has order 60, is perfect and simple, and its cusp $a(t)$ has order five. The program also verifies the derived-series orders $576, 96, 8, 1$ for Q. Thus G_6, an extension of Q by the abelian group $C/6C$, is solvable. The image of the diagonal map is a subdirect product. Goursat's lemma expresses a proper subdirect product by a common nontrivial quotient of the two factors. Every quotient of G_6 is solvable, whereas the only nontrivial quotient of A_5 is A_5 itself and is perfect. Hence the image is the whole product.

The subgroup $a(N_B)$ is normal in A_5, and $A_5/a(N_B)$ is a quotient of Q. The same argument gives $a(N_B) = A_5$, and therefore $a([N_B, N_B]) = A_5$. This proves $[N_B, N_B] \not\subset J_{A_5}$.

Let α_6 and f_6 be any certified, gauged automorphism and commutator from E.1. On the product choose $(\alpha_6, \mathrm{id}_{A_5})$, multiplier $\lambda = 1$, and $f = (f_6, 1)$. The image of F is $\mathrm{im}(F \to G_6) \times A_5$: its projection to A_5 is surjective because its quotient has order dividing six, and the same Goursat argument applies to its solvable first factor. Taking derived groups shows that f is the image of an element of F'. Both hexagons hold componentwise, and the product automorphism verifies generation. This proves the shadow claim by the finite definition.

The refined cusp has order $\mathrm{lcm}(72, 5) = 360$ and the shadow order parameter is 180; $m = 0$ and $\lambda = 1$ satisfy its unit condition. The projection $G_6 \times A_5 \to G_6$ gives the exact reduction map that retains the original CM coordinate. This quotient is additional to, and has not been identified with, LM7 D.26's S_3-core quotient B/J. □

5.3 Finite tests from the full GT distribution theorem

Statement E.6 (Proved). Five simultaneous finite cyclic-cover tests preserve all 24 candidates. *Let $f_0 \in F'$ be the explicit gauged word of any E.1 certificate, and let $q_F : F \twoheadrightarrow H = q(F)$ be the matrix quotient. There is one word $k \in [N_B, N_B]$ such that $f_* = f_0 k$ has the same image as f_0 in G_6 and, simultaneously for all $N \in \{2, 3, 4, 6, 12\}$, satisfies*

$$q_F \delta_N(f_*) = q_F(y)^{c_N(f_*)} q_F(f_*), \qquad c_N(f_*) \text{ read modulo } 6.$$

It can be chosen also to satisfy, modulo six, $c_4 = 2c_2$, $c_6 = c_2 + c_3$, and $c_{12} = 2c_2 + c_3$. These are assertions in the listed finite targets.

Proof. The programs `de_distribution_test.py` and `de_distribution_more.py` reconstruct words for the 97 fundamental cycles of the graph used in E.1, rewrite them in x, y, and evaluate the maps below. Put $A = H \times \mathbb{Z}/6$, and define

$$\psi_N : H_N \cap F \to A, \qquad w \longmapsto (q_F \delta_N(w), c_N(w) \bmod 6).$$

The x exponent of every member of N_B is even. Consequently $K_6 \subset H_N$ for every $N \mid 12$; also $N_B' \subset F'$. Induce ψ_N to a homomorphism from F into $A^N \rtimes \mathbb{Z}/N$ by the transversal $1, x, \ldots, x^{N-1}$. Its coordinates are the Schreier words $x^i w x^{-(i+s)}$, with residues taken modulo N and s the x exponent of w modulo N. If an element is written (v_i, s), multiplication is $v_i(w_1 w_2) = v_i(w_1) v_{i+s(w_1)}(w_2)$.

The normal subgroup generated by commutators of the 97 images is the image of N_B'. Every such commutator has shift zero. Conjugation of a shift-zero vector by (v_i, s) sends its ith coordinate to $v_i k_{i+s} v_i^{-1}$. Starting from the commutators, the program repeatedly takes coordinate subgroup closures and these conjugations until stable. This computes the coordinate projections exactly: every generator added is a coordinate of an actual product of conjugates, and the final projections contain all such products. The coordinate-zero orders obtained are

N	2	3	4	6	12
$\lvert\psi_N(N_B')\rvert$	8	48	48	48	48
$\lvert\psi_N(K_6)\rvert$	8	144	96	144	288
passing compatible candidates	12	12	12	12	12
passing extra candidates	12	12	12	12	12.

For each certificate it intersects the exact coset $\psi_N(f_0)\psi_N(N_B')$ with

$$\{(q_F(y)^c q_F(f_0), c) : c \in \mathbb{Z}/6\}.$$

Every intersection is nonempty. The order of $q_F(y)$ is six, so the exponent modulo six suffices for this target.

To impose these tests simultaneously, `de_distribution_joint.py` uses a faithful degree-17 permutation representation of A: $H \simeq C_2^2 \times \mathrm{SL}_2(\mathbb{F}_3)$ acts regularly on four points and naturally on the eight nonzero vectors of $\mathbb{F}_3^2$, while $\mathbb{Z}/6$ acts on disjoint sets of two and three points. The product of the five induced representations has degree $17(2+3+4+6+12) = 459$. The program takes the derived subgroup of the image of the 97 N_B generators and projects its shift-zero elements to coordinate zero in each factor. Exact permutation-group membership gives the joint image of N_B' in A^5, of order 42 467 328.

For each certificate we test the finite target tuples with $(c_2, c_3, c_4, c_6, c_{12}) = (a, b, 2a, a + b, 2a + b)$ modulo six. All 24 required coset intersections are nonempty. A tuple in this joint

image is attained by a single word of N'_B, which gives the asserted common correction. The complete output includes the chosen coefficient tuple, the derived-image generators, its order, and the membership results. Thus no inference from five independent choices is used for the simultaneous statement. □

These computations concern the explicit projections ψ_N. They do not enumerate all shadows at a refinement carrying these projections, and they do not impose generation there. In particular they show that these necessary distribution projections do not exclude an extra CM candidate. D.6 still requires the full collection of finite shadow fibres. The theorem in D.5 provides a concrete additional relation to test, but E.2 and E.3 remain the precise unsettled questions.

5.4 The requested S_3-core refinement

Statement E.7 (Proved). All 24 candidates lift through the specified S_3 core. *Fix the free basis $n_1, \dots, n_{97}$ of N_B given by the fundamental cycles in E.1, ordered by the omitted-edge order in its program. Let $\rho : N_B \twoheadrightarrow S_3$ send n_1, n_2 to two different transpositions and the other basis elements to 1. Put*

$$J = \mathrm{core}_B(\ker \rho \cap K_6), \qquad D_J = B/J.$$

There is an exact sequence of finite groups

$$1 \longrightarrow (\mathbb{Z}/3)^{288} \longrightarrow D_J \longrightarrow G_6 \longrightarrow 1.$$

The cusp t has order 72 in D_J. Each of E.1's 24 certified shadows is the reduction of a shadow at the inverse image of J in B_3, with multiplier one. All twelve extra CM shadows consequently survive this first requested nonabelian refinement.

Proof. The complete finite calculation is `de_s3_core.py`; it was run to completion. We describe its faithful model and the exact surjectivity test. Identify S_3 with the affine group of $\mathbb{F}_3$, whose elements are pairs (ϵ, b) acting by $z \mapsto \epsilon z + b$, with $\epsilon \in \{1, -1\}$. The two transpositions are $(-1, 0)$ and $(-1, 1)$. On the graph of E.1 label the first and second omitted edges by these elements, and every other edge by 1. Denote these labels by a_q for $q \in Q$. The induced homomorphism

$$\omega : B \longrightarrow S_3^Q \rtimes Q$$

has s component identically 1, right permutation $q \mapsto qs$, and r component $a_q a_{qr}^{-1}$ with right permutation $q \mapsto qr$. The identities $s^2 = r^3 = 1$ follow by cancellation. For $n \in N_B$, the component at 1 is $\rho(n)$; the other components are its conjugate Schreier representations. Hence $\ker \omega$ on N_B is the normal core of $\ker \rho$. The combined map $(B \to G_6, \omega)$ has kernel exactly J and is a faithful model of D_J.

Since every element of S_3 has sixth power one, $\omega(K_6) = \omega(N'_B)$. Each of its components lies in $A_3 \simeq \mathbb{F}_3$. This image is therefore a vector subspace $M \subset \mathbb{F}_3^Q$. The program forms the commutators of the images of the 97 basis elements and closes their span under conjugation by s, r. Conjugation by $(\epsilon_q, b_q; g)$ is the linear map

$$(z_q)_{q \in Q} \longmapsto (\epsilon_q z_{qg})_{q \in Q}.$$

The iteration stabilizes at dimension 288. Every inserted vector is the image of a product of conjugates of the basis commutators; conversely a stable span contains the image of their normal closure. Thus the computed subspace is exactly $\omega(N'_B)$. The kernel of $D_J \to G_6$ is

M by the combined faithful model. Direct evaluation gives $t^{72} = 1$; its order is at least 72 from its image in G_6.

For each original pair choose the already recorded words A, β and write $S' = \beta s \beta^{-1}$, $R' = ArA^{-1}$ in this model. The error $S'R't^{-1}$ lies in M. Replacing the conjugators by $b\beta, aA$, with $b, a \in M$, changes the error by

$$(1 - S')b + S'(1 - R')a.$$

The conjugation actions in this equation are the just specified linear maps on M. For every one of the 24 pairs the matrix of this map has rank 288. The program solves the equation and records its 576 correction coordinates. Each correction is represented by an actual word of N'_B, by the construction of M.

The proposed images of s, r satisfy their finite-order relations, so define a homomorphism $B \to D_J$, whose composition with $D_J \to G_6$ is the original automorphism. To check surjectivity on the kernel, evaluate all 97 n_i on the proposed images, take their pairwise commutators, and close their vector span under the two proposed generator actions. Their sixth powers are all trivial in the wreath component. These operations give exactly the image of K_6. In every case the resulting span has dimension 288. The map is therefore onto D_J: it is onto G_6 and onto the entire kernel. The first deterministic solution already has this property for all 24 pairs; no probabilistic inference is used.

Finally the correction words lie in $N'_B \subset F'$, so the integral abelianization of β and the cusp gauge from E.1 are unchanged. After that gauge the program directly verifies both hexagons in the wreath component; they hold in G_6 by E.1 and hence in the faithful combined model. The commutator condition holds for the same abelianization reason. With $m = 0$ and $\lambda = 1$, the unit condition and the verified surjectivity give a finite GT shadow. Reduction by $D_J \to G_6$ is the specified original shadow. This proves the claim for all 24 recorded fibres. □

For reproducibility, the first basis word in this instance is $(sr^{-1})^{12}$; the complete first two words and the deterministic basis construction are in the JSON and source. LM7 D.26 left the free basis in this construction to be chosen. We fix it here rather than identifying cores arising from different choices. The next calculation excludes all twelve representatives that survived this particular core.

5.5 A finite exclusion of all twelve recorded extra shadows

Statement E.8 (Known). Finite-factor conjugacy for the full profinite gentle action. *Every $g = (\lambda, f) \in \widehat{\mathrm{GT}}_0$ induces a continuous automorphism of $\widehat{B} = C_2 \amalg C_3$ sending s to a conjugate of s and r to a conjugate of $r^{\lambda \bmod 3}$. For $\lambda \equiv 1 \pmod 3$, the conjugators can be chosen in $\widehat{F} = \ker(\widehat{B} \to S_3)$.*

Proof. Harbater–Schneps, Proposition 4, proof of $(i) \Rightarrow (iii)$, explicitly extends the full profinite gentle action by $u \mapsto u^\lambda$, $v \mapsto f^{-1}v^\lambda f$ on $\widehat{B}_3$, and proves that its central generator goes to its λth power [HS00]. It descends to the quotient by the closed center. The universal property of profinite completion identifies this quotient with the free profinite product $C_2 \amalg C_3$: continuous homomorphisms from either to a finite group are precisely the pairs of elements satisfying $s^2 = r^3 = 1$.

Scheiderer's Appendix to Lochak–Schneps, unnumbered Corollary on printed p.598, places every nontrivial finite subgroup of a free profinite product in a conjugate of a factor [LS97]. Applying it to the images of $\langle s \rangle$ and $\langle r \rangle$ gives the stated conjugacy possibilities. The induced automorphism of $B^{\mathrm{ab}} = C_2 \times C_3$ is multiplication by λ, as follows from the abelianization of the displayed braid action. This distinguishes r from r^{-1} and selects the stated power. For

$\lambda \equiv 1 \pmod 3$, reduction of the braid action to S_3 fixes both braid generators, because λ is odd and $f \in \widehat{F}'$. A conjugator for s consequently projects into $C_{S_3}(s) = \langle s \rangle$, and a conjugator for r projects into $C_{S_3}(r) = \langle r \rangle$. Right multiplication by the appropriate power of s or r leaves the conjugate unchanged and makes that projection trivial. This gives conjugators in $\widehat{F}$. □

Put $N = N_B$ for the rest of this subsection, and define characteristic subgroups of N, hence normal subgroups of B, by

$$J_3 = (\gamma_3 N\, N^3) \cap (N' N^2), \qquad J_2 = (\gamma_3 N\, N^4) \cap (N' N^3), \qquad J_* = J_2 \cap J_3.$$

All are contained in $K_6 = N' N^6$. Write $D_i = B/J_i$ and let $\widetilde{J}_i$ be the inverse image of J_i under $B_3 \twoheadrightarrow B$. These are finite-index normal subgroups of B_3 contained in PB_3. The projections $D_* \to D_i \to G_6$ induce the reduction maps on gentle shadows used below. Their pure parts surject onto $H = q(F) \simeq C_2^2 \times \mathrm{SL}_2(\mathbb{F}_3)$ by restriction of q.

Statement E.9 (Proved). An explicit common refinement removes all twelve recorded extra shadows. *Number the 24 E.1 certificates from zero in their stored source order, retaining exactly the records containing* `mod6_beta_cycles`*. The exact gentle-shadow fibres of the two projections have the following nonempty instances among these certificates:*

refinement	*surviving certificate indices*	*compatible*	*extra*
$D_3 \to G_6$	$0, 1, 2, 3$	2	2
$D_2 \to G_6$	$0, 16$	1	1
$D_* \to G_6$	0	1	0.

The extra indices are $1, 2, 4, 7, 9, 10, 12, 15, 16, 19, 21, 22$*. Consequently none of the twelve recorded extra shadows is in the image of* $\widehat{\mathrm{GT}}_0$*, and none is in the image of* $\widehat{\mathrm{GT}}$*. This excludes those specified finite shadows, while E.2 asks about all shadows in their CM matrix fibres.*

Proof. We give the exact linear equations, including the multiplier and commutator conditions. Since N is free of rank $n = 97$, its image in D_3 is the product of the free class-two exponent-three group on n generators with $C/2C$. The kernel of $D_3 \to G_6$ is

$$M_3 = \Lambda^2(\mathbb{F}_3^{97}), \qquad \dim M_3 = 4656.$$

For D_2 the corresponding product is the free class-two exponent-four group on n generators with $C/3C$. Squares and commutators have order two and generate the central kernel

$$M_2 \simeq (\mathbb{Z}/2)^{97+\binom{97}{2}}, \qquad \dim M_2 = 4753.$$

These descriptions follow directly from the normal forms

$$(a, z)(b, w) = (a + b, z + w + \tfrac{1}{2} a \wedge b) \quad (p = 3)$$

and $n_1^{a_1} \cdots n_{97}^{a_{97}} \prod_{i<j} [n_i, n_j]^{z_{ij}}$ with $a_i \bmod 4$, $z_{ij} \bmod 2$ at two. Collecting these normal forms verifies the defining exponent and class-two relations, and their independent generator and commutator coordinates prove that no further relations have been imposed. The two prime parts form the stated products since their orders are coprime.

Let $\mathcal{S}, \mathcal{R}$ denote conjugation by s, r on M_i. At three these are the exterior-square actions induced from $C/3C$. At two their duals act on the quadratic functions on $C/2C$ by $q(v) \mapsto q(Sv)$ and $q(v) \mapsto q(Rv)$; here S, R are the verified 97-dimensional cycle actions. This formula follows by squaring the collected normal form, whose square is the quadratic function of its parity vector. It includes both square and commutator coordinates.

For one recorded certificate, undo its explicit E.1 cusp gauge and take its stored word β. Set

$$S_0 = \beta s \beta^{-1}, \qquad R_0(\lambda) = S_0 t^{\lambda}.$$

The first element is an involution. For every possible lift of the same finite shadow, the new images have the unique form

$$S' = S_0 k, \qquad R' = k^{-1} R_0(\lambda), \qquad k \in M_i.$$

The action of S_0 on M_i is $\mathcal{S}$, and that of $R_0(\lambda)$ is $\mathcal{R}$: their images in Q are s, r, and N acts trivially on M_i. The exact order equations are therefore

$$(1 + \mathcal{S})k = 0, \qquad (1 + \mathcal{R} + \mathcal{R}^2)k = R_0(\lambda)^3. \tag{1}$$

We explain why these test all finite gentle lifts, without assuming that torsion in a finite group is conjugate to a specified factor. Put $f = sS'$ after undoing the gauge, $u = t^{-1}$, and $v = sus$. For the two finite hexagons, multiplying the first on the right by u^{λ} gives $f^{-1}s$, while multiplying the second on the left by u^{λ} gives sf. Their left sides are identical. Thus they imply $(sf)^2 = 1$. After this equality the first hexagon reads $(S' t^{\lambda})^3 = 1$. Reversing these algebraic steps proves the converse. In particular (1) is the complete pair of hexagon equations at the indicated quotient. The gauge is conjugation by a fixed power of $x = t^{-2}$ and commutes with t^{λ}, so undoing it and reinstating it are inverse operations throughout this calculation.

In D_3 the cusp has order 72. Its order is at least 72 from G_6 and divides 72 from the exponent-three and exponent-two factors. The finite order parameter is therefore 36, and the multiplier over a recorded shadow has the unique residue $\lambda = 1 \bmod 72$. Every $k \in M_3$ is in the image of F'. As two is invertible, the first equation in (1) is parametrized by $k = (1 - \mathcal{S})w$. The program `de_class2_gentle.py` constructs the complete matrix

$$(1 + \mathcal{R} + \mathcal{R}^2)(1 - \mathcal{S}) : M_3 \to M_3$$

and all 24 right sides by word collection. Its rank is 1542. Exact elimination of the augmented matrix, whose rank is 1547, gives exactly indices $0, 1, 2, 3$. Each retained correction is substituted into both original equations.

In D_2 the cusp has order 144. Indeed t^{12} has the primitive cycle from E.1, so its sixth power has a nonzero square coordinate modulo two, while its twelfth power is trivial in both factors. The finite order parameter is 72. All multiplier lifts are $\lambda = 1, 73 \bmod 144$; both are tested. For the commutator condition let L_N be the image of N in $F^{\mathrm{ab}} = \mathbb{Z}^2$. Exact word abelianization gives the column lattice

$$L_N = \begin{pmatrix} 6 & 4 \\ 0 & 2 \end{pmatrix} \mathbb{Z}^2.$$

The image of J_2 in this abelianization is exactly $12L_N$: membership gives divisibility by four and three, and twelfth powers of elements of N give the reverse inclusion. A square coordinate in M_2 is represented by n_i^6 and a commutator coordinate by a commutator. Hence $k \in M_2$ lies in the image of F' exactly when the two coordinates of its square contribution to $6L_N/12L_N$ vanish. The program `de_class2_two_gentle.py` appends these two rows to (1). The coefficient matrix has rank 3927; its augmentation by all 48 right sides has rank 3932. Its complete solution test gives exactly indices $0, 16$. Every retained solution is substituted into the two order equations and the two abelianization rows. The fixed E.1 gauge made the original f a commutator, so this is precisely the condition that the corrected, regauged f be a commutator as well.

All passing solutions satisfy generation. The homomorphism $B \to D_i$ defined by S', R' preserves the image of N, because its map to Q is the recorded identity before gauge. The defining class and power laws show that it kills J_i. On each prime part of the image of N, its Frattini quotient map is exactly the invertible E.1 map on C/pC. The elementary Frattini argument for a finite p-group therefore makes it surjective on both prime parts, and hence on D_i. The multiplier, commutator, hexagon and generation conditions of a finite gentle shadow have all been checked.

The bit-packed elimination routines in `de_gf3.c` are independently compared with the earlier Python elimination on four rectangular matrices for each prime, including multiple machine word boundaries. An additional independent calculation `de_class2_direct_check.py` uses the full 9312 by 4656 matrix of the peripheral correction map at three, agrees with the separate Jordan-block computation, and gives the same four survivors after all conjugator-centralizer freedoms. The full hexagon calculation above supplies the exact finite-shadow scope.

Every shadow over $\widetilde{J}_*$ reduces to shadows over both $\widetilde{J}_2$ and $\widetilde{J}_3$. Intersecting the two displayed index lists leaves only index zero. That record has $A = \beta = 1$ and zero cycle corrections; its identity shadow lifts to every quotient, including D_*. Thus the common-refinement image among the 24 recorded points is exactly the claimed singleton. The all-finite-quotient criterion D.6 excludes the other 23 from the full profinite gentle image. The inclusion $\widehat{\mathrm{GT}} \subset \widehat{\mathrm{GT}}_0$ gives the stated full GT exclusion as well. □

Statement E.10 (Proved). The affine sizes of the CM matrix fibres. *For each of the 24 admissible CM matrix pairs and multiplier one modulo 72, the cusp-preserving changes of its two homology conjugators form affine spaces of dimensions 98 over $\mathbb{F}_2$ and 97 over $\mathbb{F}_3$. Quotienting by the changes that leave both generator images fixed gives dimensions 17 and 18. Quotienting further by simultaneous conjugation by cusp powers gives dimensions 16 and 17. The finite classification modulo all inner conjugations by the centralizer of t in C/pC has dimensions 8 and 9. Invertibility of the resulting action on C/pC is an additional condition in each of these spaces.*

Proof. The exact program `de_affine_fibres.py` computes the kernel of L from E.1 and the following subspaces. Over $\mathbb{F}_2$ the dimensions of C^s, C^r, C^t, C^Q are respectively $52, 29, 9, 0$; over $\mathbb{F}_3$ they are $45, 34, 9, 0$. The changes that fix both generator images are $C^s \oplus C^r \subset \ker L$. Simultaneous inner conjugation adds the diagonal copy of C^t; its intersection with that first subspace is the diagonal $C^Q = 0$. The cusp powers give the one-dimensional diagonal subspace generated by the primitive cycle $c = [t^{12}]$. Taking the indicated dimensions gives all assertions. The output stores explicit bases and complements, and directly substitutes every column into L.

Conjugation by an actual power of t centralizes t in every quotient, so that operation can be transported through every refinement. The larger finite space C^t is classified here only at the homology quotient: transporting one of its elements as a centralizer would require a lift that centralizes t. Consequently the last quotient does not by itself replace the whole finer lifting problem. In particular the twelve original representatives cannot be treated as exhaustive matrix fibres. R11 supplies a genuine arithmetic point in each compatible CM matrix fibre, while E.9 excludes the recorded points in every compatible fibre except the identity fibre. These give concrete distinct points with the same CM matrices. □

Statement E.11 (Proved). Every admissible CM matrix fibre has a replacement at the common refinement. *Each of the 24 admissible CM matrix pairs of E.1 is attained by a gentle shadow at $\widetilde{J}_*$, where $J_* = J_2 \cap J_3$ is the explicit refinement of E.9. In particular*

all twelve extra CM matrix fibres *have nonempty image from this refinement, although their twelve original recorded representatives are excluded by E.9. The new witnesses are finite shadows; their membership in the full profinite image is still to be determined.*

Proof. The programs `de_fibre_three.py` and `de_fibre_two.py` use the complete 18- and 17-dimensional spaces after conjugator centralizers in E.10. They construct exact five-component quadratic obstruction maps on these spaces. The three-primary map projects the eight alternating coinvariants modulo their three centralizer directions. The two-primary map projects the sixteen quadratic coinvariants modulo their eleven centralizer and multiplier directions. Both maps are checked against direct collection of words for every matrix pair. Searching their finite parameter spaces gives a zero in each of the 24 fibres, and the induced action has rank 97 in every case. Only these produced and verified witnesses are used; the search is not used to infer any negative assertion.

Write the two pairs of changes of the conjugator cycles as (b_2, a_2) and (b_3, a_3). The program `de_fibre_crt_verify.py` uses

$$b = 3b_2 + 4b_3 \pmod 6, \qquad a = 3a_2 + 4a_3 \pmod 6,$$

and prefixes the original words β, A by $\prod n_i^{b_i}, \prod n_i^{a_i}$. These are literal words in s, r stored in its output. Their graph cusp equation and both 97-dimensional generation ranks are checked again. Their images in Q are the original matrices, so their CM coordinates agree with the prescribed pair, also after the common cusp gauge.

The complete norm systems of E.9 are then solved anew for these CRT words. Every one of the 24 points has a solution at both primes with $\lambda = 1 \bmod 144$. The two-primary system includes the two exact commutator-image rows. Let k_2 be its square and commutator coordinates and k_3 the three-primary commutator coordinates. The common correction is the explicit word

$$k = \prod_i n_i^{6(k_2)_i} \prod_{i<j} [n_i, n_j]^{3(k_2)_{ij} + 4(k_3)_{ij}}.$$

Its two-primary component is k_2 and its three-primary component is k_3. For the integral cusp gauge g computed from the new word β, set

$$f = sx^g(\beta s \beta^{-1}) k x^{-g} j.$$

Here j is an explicitly recorded product of twelfth powers of two basis words of N, chosen to cancel the abelianization of k. The two commutator rows ensure that this abelianization belongs to $12L_N$; an exact unimodular pair in the displayed basis of L_N constructs j. Thus $j \in J_*$ and the displayed word f lies in F' itself. This gives one common commutator word, with a single integral gauge, for the two primary components.

Finally both hexagons are substituted directly, as words, into a faithful collector for the product of the exponent-four and exponent-three class-two images of N. There are 48 successful hexagon checks. For efficiency a factored central block is evaluated through its exact action under the current Q coset: at three its alternating matrix changes by $Z \mapsto AZA^t$; at two a square vector u and symmetric commutator matrix Z change to Au and the upper entries of $A(Z + \operatorname{diag} u)A^t$. These formulas follow by collecting the explicit square and commutator factors above, so this evaluation is precisely substitution of those actual words. The other letters are collected individually by the omitted-edge basis. A word is trivial in D_* exactly when its Q image and both collected primary normal forms vanish.

The checked Frattini ranks give generation on both primary kernels, as in E.9. The word, multiplier, commutator and generation conditions therefore give a shadow at $\widetilde{J}_*$ in every asserted fibre. The JSON retains the literal CRT conjugators, all factored correction words, both ranks, the common gauge and multiplier, and every hexagon result. □

Statement E.12 (Proved). All admissible CM pairs lift through the whole class-two three-primary tower. *For every admissible CM matrix pair in E.1 there is a compatible family of gentle shadows through all quotients*

$$B/\big((\gamma_3 N\, N^{3^k}) \cap (N' N^2)\big), \qquad k \geq 1,$$

with multiplier one and that CM pair. Thus all twelve extra matrix fibres survive this entire tower. This assertion concerns matrix fibres with a coherent cusp gauge; it does not assert that an independently chosen normalized finite representative extends.

Proof. Let $\mathcal{P}$ be the free class-two pro-three group on 97 generators. Its logarithmic coordinates form $V \oplus \Lambda^2 V$ for $V = \mathbb{Z}_3^{97}$, with product $(a, z)(b, w) = (a + b, z + w + \frac{1}{2} a \wedge b)$. The relevant completed kernel of the extension of Q is $\mathcal{P} \times (C/2C)$. The odd denominator two is a unit in $\mathbb{Z}_3$. Consequently the cusp equation in two conjugators has polynomial coordinate functions over $\mathbb{Z}_3$, of degree at most two.

We verify that its full derivative is onto at each three-primary witness of E.11. Its homology part is L, of rank 97 modulo three. The derivative in the central coordinates has rank 4648, with the eight-dimensional alternating coinvariant quotient computed in E.9. Changes by the order-three conjugator centralizer give rank three in this quotient. The remaining five components are exactly the quadratic map used in `de_fibre_three.py`. Here is a coordinate description of that map and its derivative. For an invariant alternating form B, write τ_{S_0}, τ_t for the central parts of conjugation by $S_0 = \beta s \beta^{-1}$ and by t, evaluated by B. A homology change $(b, a) \in \ker L$ changes the central cusp coordinate by

$$\tau_{S_0}(a - b) - \tau_t(a) + \tfrac{1}{2} B\big(b, (T - 1)a\big), \qquad T = SR.$$

This follows by multiplying $n(b)\, S_0 (n(b)^{-1} n(a)) S_0^{-1}$, the old cusp error, and $t n(a)^{-1} t^{-1}$ in the displayed group law. The central coordinates chosen for $n(a), n(b)$ cancel in the coinvariant quotient. The program checks the formula against direct word collection.

The independent derivative calculation `de_fibre_three_smooth.py` evaluates this full eight-component derivative at each of the 24 zeros, adjoins the three centralizer directions, and obtains rank eight in every case. Thus the full coordinate derivative has rank $97 + 4648 + 8 = 4753$, equal to the dimension of the cusp target. The centralizer corrections and central corrections used to make the initial error zero do not change that rank: central corrections do not change the homology coordinates, and the five-component quadratic map is constant along the discarded centralizer directions. All these equalities are checked over $\mathbb{F}_3$.

Successive lifting is now explicit. At a solution modulo 3^k, divide its error by 3^k, reduce modulo three, and solve the surjective derivative equation for a correction of the two conjugators by 3^k times their coordinates. The polynomial expansion shows that the corrected error is zero modulo 3^{k+1}; its terms of degree at least two in the correction are divisible by 3^{2k}. The corrections preserve all preceding levels, and the derivative retains the same reduction modulo three. Induction gives convergent conjugators over $\mathbb{Z}_3$. Their component in $C/2C$ is kept fixed throughout.

The resulting map preserves Q and is invertible on the three-primary Frattini quotient and on $C/2C$, by the verified ranks in E.11. The pro-three Frattini argument makes it onto $\mathcal{P}$, and its induced finite maps are automorphisms. Choose profinite lifts of the limiting conjugators to $\widehat{F}$ and one cusp gauge using their actual abelianizations. The formula $f = s\beta' s(\beta')^{-1}$ then gives a commutator and compatible hexagons at every finite level. The common multiplier is one. The simultaneous gauge leaves the CM coordinates unchanged.

This tower kills $\gamma_3 N$. The map to A_5 in E.5 sends N onto A_5, and hence also sends $\gamma_3 N$ onto A_5. Its refinement therefore does not factor through any member of the displayed tower.

This explicit factorization obstruction records why these compatible class-two lifts do not complete E.2. □

Statement E.13 (Proved). Every CM matrix fibre survives the common exponent-eight refinement. *Put*

$$J_{8,3} = (\gamma_3 N\, N^8) \cap (\gamma_3 N\, N^3).$$

This is a normal finite-index subgroup of B, contained in J_* of E.9. Its inverse image in B_3 is normal, has finite index, and is contained in PB_3. Each of the 24 admissible CM matrix pairs of E.1 is attained by a gentle shadow at this refinement, with multiplier one modulo 288. *In particular every one of the twelve extra CM matrix fibres remains nonempty here.*

Proof. We compute entire central correction fibres above the 24 affine homology points of E.11. The image of N in the two-primary factor is the free class-two exponent-eight group, with ordered normal form

$$\prod_i n_i^{a_i} \prod_{i<j} [n_i, n_j]^{z_{ij}}, \qquad a_i \in \mathbb{Z}/8, \quad z_{ij} \in \mathbb{Z}/4. \tag{E.13.1}$$

The product in these coordinates is

$$(a, z)(b, w) = (a + b, z + w - (a_j b_i)_{i<j}). \tag{E.13.2}$$

Indeed moving $n_i^{b_i}$ past $n_j^{a_j}$ for $j > i$ introduces $[n_i, n_j]^{-a_j b_i}$. Commutators have eighth power one, and the eighth power of a product gives their twenty-eighth power one. Since $\gcd(8, 28) = 4$, their fourth powers vanish. Conversely the ordered normal form with these relations has exponent eight. This proves the stated free quotient and the faithful collector used below.

The kernel above the fixed G_6 homology image, retaining $N^{\mathrm{ab}}/3$, has a particularly useful description:

$$A \cong (\mathbb{Z}/4)^{4753}, \qquad e_i = n_i^6, \quad e_{ij} = [n_i, n_j]^3. \tag{E.13.3}$$

The two-primary abelian coordinates of an element of this kernel are even, and multiplication by three is invertible modulo four, so these elements span the whole kernel. They are independent by (E.13.1). Moreover their commutators are $[n_i, n_j]^{36} = 1$, so the displayed group is abelian. This is the exact reason that this entire finite correction problem is linear over $\mathbb{Z}/4$.

Let β be one of the literal affine conjugators from E.11, before its cusp gauge, and put $S_0 = \beta s \beta^{-1}$, $R_0 = S_0 t$. Every lift of the fixed homology shadow has

$$S = S_0 k, \qquad R = k^{-1} S_0 t^{1+72h}, \quad k \in A, \quad h \in \mathbb{Z}/4.$$

The complete equations, with the two commutator-image rows, are consequently one linear system over $\mathbb{Z}/4$:

$$(1 + S_0)k = 0, \qquad (1 + R_0 + R_0^2)k = (S_0 t^{1+72h})^3, \qquad B_N^{-1} \operatorname{ab}_F(k) = 0 \pmod{24}. \tag{E.13.4}$$

Here $B_N = \left(\begin{smallmatrix} 6 & 4 \\ 0 & 2 \end{smallmatrix}\right)$ is the chosen basis matrix for the lattice L_N in E.9. Define the integer columns $H_i = B_N^{-1} \operatorname{ab}_F(n_i)$. Then the two equations are $H(k_i)_i = 0 \pmod 4$, because $\operatorname{ab}_F(k) = 6 B_N H(k_i)_i$ and the new kernel has abelian image $24 L_N$. The right side of the second equation is affine in h: $t^{72} \in A$, and A is abelian. The common multiplier range is $1 + 72h \pmod{288}$.

For clarity, the action matrices in (E.13.4) use the ungauged S_0, R_0. The normalized pair is obtained by conjugation by $x^g = t^{-2g}$, which fixes t. Thus it preserves both order

equations, and its conjugation of $k \in F$ preserves the exact F^{ab} coordinates. This supplies the comparison used when restoring the cusp gauge.

Here is the coefficient construction and the exhaustive linear solution. If conjugation by a word q sends n_i to normal coordinates $(A_q e_i, \tau_i)$, its action on (E.13.3) has blocks

$$\begin{pmatrix} A_q & 0 \\ T_q & \Lambda^2 A_q \end{pmatrix}, \qquad (T_q)_{(a,b),i} = 2(\tau_i)_{ab} - (A_q)_{ai}(A_q)_{bi} \pmod 4, \tag{E.13.5}$$

where $a < b$ indexes a normal commutator coordinate. The formula follows by taking the sixth power of the image of n_i in (E.13.2) and then dividing its central coordinate by three. Conjugation by an additional normal word with abelian coordinate c adds $2(c_{i_1}(A_q)_{i_2 i} - c_{i_2}(A_q)_{i_1 i})$ to that block. These formulas compute every coefficient.

The program `de_class2_eight_fibres.py` reduces the 9508 by 4754 system modulo two. Its rank is 3927 and its nullspace has dimension 827, including the multiplier variable. For a particular solution x_0 and a binary nullspace matrix K, write every possible lift as $x = x_0 + Kz + 2y$. The second-layer equation is exactly

$$A_0 y + \frac{A_4 K}{2} z = \frac{b_4 - A_4 x_0}{2} \pmod 2, \tag{E.13.6}$$

where the numerators are even and are reduced modulo four before division. This system has coefficient rank 3953 and is consistent in each of the 24 cases. It retains all 827 nullspace parameters, so it tests the complete fibre above each fixed affine homology point. A solution with $h = 0$ is retained in every case. Direct substitution verifies both order equations and the two commutator-image rows.

We then form actual words at the common refinement. Let k_2 be the retained $\mathbb{Z}/4$ correction, and let k_3 be the three-primary correction already retained in E.11. Set

$$k = \prod_i n_i^{6(k_2)_i} \prod_{i<j} [n_i, n_j]^{3(k_2)_{ij} + 4(k_3)_{ij}}.$$

Its exponent-eight image is k_2, and its exponent-three image is k_3. Since $H((k_2)_i)$ is divisible by four, the same unimodular two-column calculation as in E.11 gives an explicit product j of twenty-fourth powers of basis words satisfying $\mathrm{ab}_F(kj) = 0$. It is trivial in both factors of $J_{8,3}$. Thus

$$f = s x^g S_0 k x^{-g} j \in F' \tag{E.13.7}$$

is a single actual word for the common shadow.

The final program `de_fibre_eight_crt_verify.py` substitutes both complete braid hexagons for (E.13.7) in both faithful collectors: there are 96 successful component checks. The exponent-eight collector also checks sixteen block actions against literal expanded conjugate words. It keeps the section comparison in the Schreier calculation: if the short representative is $n(c)$ times the spanning-tree representative, the latter's central action is $\tau_{\mathrm{short}} - c \wedge A_q$. This is the inner correction required by (E.13.2).

The new corrections preserve the verified G_6 homology shadow. Both 97-dimensional Frattini ranks remain maximal, so generation follows on the two primary kernels. The actual commutator word, unit multiplier, complete hexagons and generation prove the assertion. The JSON retains all literal base words and the factored correction and adjustment words. The quotient still kills $\gamma_3 N$; the explicit A_5 comparison at the end of E.12 therefore applies to this refinement as well. No cofinality conclusion follows from these positive finite fibres. □

5.6 The exponent-sixteen fibres and the two-adic lifting argument

Statement E.14 (Proved). Every CM matrix fibre survives the common exponent-sixteen refinement. *Let $J_{16,3} = (\gamma_3 N\, N^{16}) \cap (\gamma_3 N\, N^3)$. Its inverse image in B_3 is a finite-index normal subgroup contained in PB_3. Every one of the 24 admissible CM matrix pairs of E.1 occurs as a gentle shadow at this refinement, with multiplier one modulo* 576.

Proof. The exponent-sixteen collector has coordinates $a \in (\mathbb{Z}/16)^{97}$ and $z \in (\mathbb{Z}/8)^{4656}$, with the product in (E.13.2). The image of $N'N^6$, retaining $N^{\mathrm{ab}}/3$, has normal coordinates

$$k = \prod_i n_i^{6u_i} \prod_{i<j} [n_i, n_j]^{3v_{ij}}, \qquad (u, v) \in (\mathbb{Z}/8)^{4753},$$

and exact product

$$(u, v)(u', v') = (u + u',\, v + v' - 12(u_j u'_i)_{i<j}). \tag{E.14.1}$$

The correction group is therefore nonabelian at this level; the program retains its full product and inverse.

For each affine homology point of E.11 use the same ungauged $S_0 = \beta s \beta^{-1}$ and $R_0 = S_0 t$. With $\lambda = 1 + 72h$, the complete equations are

$$(S_0 k)^2 = 1, \qquad (k^{-1} S_0 t^{1+72h})^3 = 1, \qquad Hu = 0 \pmod 8. \tag{E.14.2}$$

Here H is the two-row integer matrix of E.13. The last condition is exact: the common kernel has F-abelian image $48L_N$, and the correction has image $6B_N Hu$.

Write $x = (u, v, h)$ and let $\mathcal{F}(x)$ be the two normal-coordinate order errors followed by Hu. These are integer polynomials of degree at most two. Their diagonal quadratic coefficients are divisible by six and their mixed quadratic coefficients by twelve. Indeed the sixth-power contribution of an automorphism with abelian matrix A and generator cochain τ is

$$2u_i\tau_i - (6u_i^2 - u_i)A_{ai}A_{bi}$$

in a central coordinate; reordering contributes multiples of $12u_iu_j$. Cusp powers have the same divisibility, since $12\binom{h}{2} = 6h^2 - 6h$. Only linear homology expressions are substituted into these formulas. Thus $\mathcal{F}$ modulo four is exactly the affine system of E.13.

Its homogeneous solutions are represented by

$$Vt + 2Kw \pmod 4, \qquad t \in \mathbb{F}_2^{801}, \quad w \in \mathbb{F}_2^{827}, \tag{E.14.3}$$

where K is the entire binary kernel and V lifts the kernel of the second-layer obstruction. Every column is verified modulo four. The projection of V mod 2 to the 97 homology coordinates and the multiplier has rank 15. Select 15 columns spanning that projection and subtract their combinations from the other columns. The remaining 786 columns have even homology and multiplier coordinates. Twice a selected column is already represented by $2K$, so this change preserves all solutions in (E.14.3).

Every direction with even homology and multiplier coordinates changes $\mathcal{F}$ modulo eight linearly and independently of the base point: the stated divisibilities kill its quadratic cross terms. Hence all 786 remaining lower parameters, all 827 upper parameters, and the arbitrary correction $4y$ enter linearly; only the 15 selected bits enter quadratically. The program `de_class2_sixteen_fibres.py` retains the full linear system and evaluates all 32768 choices of those bits. In each of the 24 cases the linear correction rank is 3953, and every constant, linear, and quadratic coefficient vanishes modulo its image. All 32768 choices survive. A

solution with $h = 0$ is retained in every case and substituted into both literal order equations and Hu.

For these two-primary coordinates and the three-primary central correction k_3 of E.11, the common actual word is

$$k = \prod_i n_i^{6u_i} \prod_{i<j} [n_i, n_j]^{3v_{ij}+16(k_3)_{ij}}. \tag{E.14.4}$$

The unimodular two-column calculation of E.11 supplies a product j of forty-eighth powers with $\mathrm{ab}_F(kj) = 0$. It is trivial in both factors. Restoring the recorded cusp gauge gives $f = sx^g S_0 k x^{-g} j \in F'$. The program `de_fibre_sixteen_crt_verify.py` substitutes both full braid hexagons in both faithful collectors, with 96 successful component checks. Normal-block inversion retains the central term $-a_i a_j$. Twelve additional literal conjugation checks include a nontrivial change of section. Both Frattini ranks remain 97 because the corrections preserve the G_6 homology point. This proves the full shadow and generation conditions.

The separate `de_class2_sixteen.py` finds no lift of the 24 particular words retained in E.13. Its projection fixes their upper corrections. Calculation (E.14.3) permits all corrections over the homology point and supplies the replacements above. These exact projection maps explain the scopes of the two calculations. □

Statement E.15 (Proved). Every admissible CM pair survives the entire relative class-two tower. *For each of the 24 admissible CM matrix pairs there is a compatible family of gentle shadows at the inverse images in B_3 of*

$$J_j = \gamma_3 N\, N^j \qquad (6 \mid j), \tag{E.15.1}$$

ordered by divisibility, with multiplier $\lambda = 1$. All twelve extra matrix fibres therefore survive this entire tower. The assertion specifies matrix fibres; it does not assert extension of the particular normalized words retained in E.13 or E.14.

Proof. We construct the two-primary component first. Let $\mathcal{N}_2$ be the free class-two pro-two group on the 97 basis elements, in the extension of Q obtained by completing N in this quotient. Let $\mathcal{U} \subset \mathcal{N}_2$ have even abelian coordinates. Multiplication by three is invertible over $\mathbb{Z}_2$, so the elements $n_i^6, [n_i, n_j]^3$ give its normal coordinates $(u, v) \in \mathbb{Z}_2^{4753}$, with product (E.14.1). The polynomial bijection

$$Z = (u, z), \qquad z_{ij} = v_{ij} + 6u_i u_j$$

puts its operations in the form

$$Z * Z' = Z + Z' + \tfrac{1}{2}[Z, Z'], \qquad [Z, Z'] = (0, 12u \wedge u'). \tag{E.15.2}$$

Every coefficient is integral. We use $\exp Z$ and $\log$ for this explicit group-coordinate correspondence; (E.15.2) justifies that notation at the prime two.

Conjugation by S_0 is an integral Lie automorphism M_S with $M_S^2 = 1$. The first order equation is equivalent to

$$G_S(Z) = (1 + M_S)Z = 0. \tag{E.15.3}$$

In fact its logarithm is $G_S(Z) + \frac{1}{2}[G_S(Z), Z]$; the map $Y \mapsto Y + \frac{1}{2}[Y, Z]$ is an integral unipotent automorphism. The target of G_S is the saturated submodule $\mathrm{Fix}(M_S)$.

Put $E = \log(R_0^3)$ and $r_* = \exp(-E/3)R_0$. Here $R_0^3 \in \mathcal{U}$, division by three is integral, and R_0 commutes with its cube; thus $r_*^3 = 1$. Write M for its Lie action and $P = (1 + M + M^2)/3$

for the integral projection onto $\mathrm{Fix}(M)$. For $k = \exp Z$, keep the multiplier equal to one and write

$$R = k^{-1} S_0 t = \exp(W) r_*, \qquad C = \log(R^3).$$

The coordinates of W and C are integral polynomials of degree at most two in Z, by (E.15.2). The second order equation is equivalent to

$$G_R(Z) = PC = 0. \tag{E.15.4}$$

To prove this integrally, the homology of C is the norm of the homology of W, so it is fixed by M and equals that of PC. Since R^3 commutes with R, $(M-1)C = -[W, C]$. On $\ker P$ the inverse of $M - 1$ is $-(M+2)/3$. With $Q_0 = 1 - P$ this gives

$$C = PC + \frac{M+2}{3} Q_0 [W, PC]. \tag{E.15.5}$$

The bracket uses only homology. Thus $PC = 0$ is equivalent to $C = 0$, and (E.15.5) is an integral linear embedding of the error PC, with left inverse P.

There is only one further independent commutator equation. The integral matrix H satisfies

$$H A_S = \begin{pmatrix} -2 & -1 \\ 3 & 2 \end{pmatrix} H, \tag{E.15.6}$$

where A_S acts on N^{ab}. Equivalently, apply B_N^{-1} to the interchange of the two coordinates of F^{ab}. The first row of $H(1 + A_S)u$ is $-(H_1 + H_2)u$. Hence (E.15.3) and $H_2 u = 0$ imply both commutator rows, and the converse holds. The exact compressed system is the integral polynomial map

$$G : \mathbb{Z}_2^{4753} \longrightarrow \mathrm{Fix}(M_S) \oplus \mathrm{Fix}(M) \oplus \mathbb{Z}_2, \qquad G = (G_S, G_R, H_2 u). \tag{E.15.7}$$

The target lattices are primitively embedded. The normal-coordinate changes and (E.15.5)–(E.15.6) recover the full error vector from G, with primitive linear part at a zero. At a point whose errors are divisible by eight the same derivative comparison holds modulo four, since the additional derivative terms multiply those errors.

Here are the exact target and derivative ranks. The cycle actions have traces $-7, -5, -5$ for s, r, r^2. The exterior-square traces are $-24, 15, 15$, respectively, from $\mathrm{tr}(\Lambda^2 A) = ((\mathrm{tr}\, A)^2 - \mathrm{tr}(A^2))/2$. Central cochains do not change a trace. It follows that

$$\mathrm{rank}\, \mathrm{Fix}(M_S) = 2361, \quad \mathrm{rank}\, \mathrm{Fix}(M) = 1591, \quad \mathrm{rank}\, \mathrm{target}(G) = 3953. \tag{E.15.8}$$

The program `de_two_hensel_tangent.py` checks these traces and (E.15.6). At each of the 24 points from E.14 it computes the full integral Jacobian modulo four. The program `de_two_hensel_fixed.py` then deletes the multiplier column, recomputes the entire binary kernel, and verifies all the ranks below for the resulting 4753-variable system. Its binary kernel has dimension 826. The divisibilities in the proof of E.14 give

$$\mathcal{F}(x + 2e_i) - \mathcal{F}(x) = 2J_i \pmod 8$$

and make $J \bmod 4$ independent of x. The central columns are the exterior-square norm matrices. All homology columns are retained, with five additional complete direction checks per point. For the full binary kernel K, the exact ranks are

$$\mathrm{rank}_{\mathbb{F}_2}(J \bmod 2) = 3927, \qquad \mathrm{rank}_{\mathbb{F}_2} \left(J \bmod 2 \;\middle|\; \frac{JK}{2} \bmod 2 \right) = 3953. \tag{E.15.9}$$

The primitive comparison with (E.15.7) identifies these with the ranks for dG. Its target has rank 3953, so its elementary divisors consist of 3927 units and 26 elements of valuation one. Its image contains twice the target lattice.

The lifting can now be given explicitly. At the integral point x_0 retained in E.14, $G(x_0)$ is divisible by eight. Choose a right inverse of dG_{x_0} over $\mathbb{Q}_2$ carrying the integral target into $\frac{1}{2}\mathbb{Z}_2^{4753}$; the elementary divisors give such an inverse. The Newton correction belongs to $4\mathbb{Z}_2^{4753}$. The new error is divisible by sixteen, because G is integral of degree at most two, and its derivative is unchanged modulo four. Repeating, an error of valuation $v \geq 3$ is replaced by one of valuation at least $2v-2$. The corrections converge to an exact zero of G, with multiplier still equal to one. Equations (E.15.3)–(E.15.6) give both exact orders and zero F-abelianization for the correction k. The retained cusp gauge commutes with t and makes the normalized f have zero abelianization, by the formula (E.15.12) below. The reduction on $N^{\mathrm{ab}}/2$ has rank 97, so the pro-two Frattini argument gives an automorphism at every finite level.

We give the gluing before applying the cusp gauge. For the same CM matrix pair, take the ungauged three-primary action of E.12, and take the identity action at every other prime. In each component the images of s, r, t in Q are s, r, t: the matrix representatives β, A centralize s, r in Q, and all their corrections belong to the normal kernel. Thus these ungauged actions have the same identity action on Q. The precise comparison of their target groups is

$$\varprojlim_{j} B/(\gamma_3 N\, N^j) \simeq \prod_{p,Q} \varprojlim_{a} B/(\gamma_3 N\, N^{p^a}), \tag{E.15.10}$$

where the product is the fibre product over Q. On the normal kernel this is the Sylow decomposition of the class-two completion of N: every finite nilpotent quotient is the product of its Sylow groups. For surjectivity at finitely many levels, lift their common Q coordinate and adjust independently in these normal Sylow groups. Compactness gives (E.15.10).

We now specify one gauge in this common group. Let $g_2 \in \mathbb{Z}_2$ be the integral gauge retained in E.14. For the limiting three-primary conjugator β_3, let $g_3 \in \mathbb{Z}_3$ be the difference of its second and first F-abelian coordinates. The E.12 corrections preserve these coordinates modulo three, so the reduction of g_3 is that of the recorded integral gauge. The image of N in F^{ab} has basis matrix $B_N = \left(\begin{smallmatrix} 6 & 4 \\ 0 & 2 \end{smallmatrix}\right)$; therefore every integral change in this difference is even. The two-primary residue of the three-primary gauge can accordingly be kept equal to that of g_2. Choose $g \in \widehat{\mathbb{Z}}$ with these two primary components and with $g_p = 0$ for $p \neq 2, 3$. The element $x = t^{-2}$ has order six in Q. Consequently x^g restricts to the specified cusp gauge in the two- and three-primary extensions. Apply its inner conjugation to the glued ungauged action, obtaining S', R' with $S'R' = t$.

Put $f = sS'$. Its common Q coordinate is

$$q(f) = s x^{\bar{g}} s x^{-\bar{g}}, \qquad \bar{g} = g \bmod 6. \tag{E.15.11}$$

In particular f need not lie in the image of N. The independent matrix program `de_adelic_gauge.py` verifies the identity ungauged Q action and (E.15.11) for all 24 pairs. For $\bar{g} = 0, 4, 2$, respectively, its canonical matrix representatives in $\mathrm{SL}_2(\mathbb{Z}/12)/\{\pm 1\}$ are

$$I, \qquad \begin{pmatrix} 1 & 4 \\ 4 & 5 \end{pmatrix}, \qquad \begin{pmatrix} 1 & 8 \\ 8 & 5 \end{pmatrix}.$$

The exact abelianization formula is

$$\mathrm{ab}_F(s x^g S x^{-g}) = \mathrm{ab}_F(sS) + (-g, g). \tag{E.15.12}$$

At two and three the selected gauges make this zero. At every other prime the ungauged action is identity, and $g_p = 0$, so it is zero there also. In those components the final action is

inner by x^g; its image can have nontrivial prime-to-p order dividing six, exactly as (E.15.11) requires. Hence f has zero full F-abelianization. Since $\gamma_3 N \subset F'$, the kernel of the resulting map to $\widehat{F}^{\mathrm{ab}} = \widehat{\mathbb{Z}}^2$ is the image of $\widehat{F}'$, and f belongs to that image. Both order equations hold in every component. The exact hexagon equivalence of E.9 now supplies compatible gentle shadows. Generation holds on each normal Sylow kernel by its Frattini map at two and three and by the specified inner maps at the other primes; it also holds on Q. Each finite map is therefore an automorphism. The two- and three-primary reductions recover the prescribed CM matrix pair, proving the family.

Every subgroup in (E.15.1) contains $\gamma_3 N$. The map to A_5 in E.5 sends that subgroup onto A_5, so its common refinement factors through none of these quotients. This explicit comparison leaves the cofinal lifting problem of E.2 unresolved. □

Statement E.16 (Proved). Simultaneous extension through every nonabelian simple quotient. *Let Σ be any finite collection of epimorphisms $\theta : B \twoheadrightarrow S_\theta$ to nonabelian finite simple groups, and put*

$$K_{j,\Sigma} = \gamma_3 N\, N^j \cap \bigcap_{\theta \in \Sigma} \ker \theta, \qquad 6 \mid j. \tag{E.16.1}$$

Every one of the 24 CM matrix pairs has a compatible family of gentle shadows at the inverse images of these subgroups in B_3, with multiplier one, as j increases by divisibility and Σ increases by inclusion. Thus the class-two families of E.15 extend simultaneously through all these nonabelian simple quotients. For each fixed epimorphism $B \twoheadrightarrow R$ to an arbitrary finite perfect group, they also extend compatibly in j through $\gamma_3 N\, N^j \cap \ker(B \to R)$.

The finite solvable quotient $B/(\gamma_4 N\, N^5)$ is not a quotient, through the natural map from B, of any $B/K_{j,\Sigma}$. This gives a precise remaining finite comparison outside this enlarged family.

Proof. Write $E_j = B/(\gamma_3 N\, N^j)$ and let R_Σ be the image of B in $\prod_{\theta \in \Sigma} S_\theta$. The latter is a product of nonabelian simple groups, with repeated diagonal factors identified. To prove this, adjoin the simple factors one at a time. The kernel of projection to the previous factors, viewed in the new simple group, is either the whole group or trivial. In the first case the new group is a direct product. In the second, the new factor is a simple quotient of the preceding product and is the graph of an isomorphism from one of its simple factors. This proves the claim by induction. In particular R_Σ is perfect.

The group Q is solvable. Indeed, $\mathrm{SL}_2(\mathbb{Z}/12)$ is the product of its reductions modulo four and three. The kernel of reduction $\mathrm{SL}_2(\mathbb{Z}/4) \to \mathrm{SL}_2(\mathbb{F}_2)$ is an elementary abelian group of order eight, and the target is $\mathfrak{S}_3$. The other factor has a normal quaternion group of order eight with cyclic quotient of order three. Passing to the quotient by $\{\pm I\}$ preserves solvability. Since the kernel of $E_j \to Q$ is nilpotent, E_j is solvable as well.

The image of B in $E_j \times R_\Sigma$ projects onto both factors. For a subdirect product $H \subset U \times V$, put $U_0 = \{u : (u,1) \in H\}$ and $V_0 = \{v : (1,v) \in H\}$. Sending one coordinate of a pair in H to the other defines an isomorphism $U/U_0 \simeq V/V_0$. Here that common quotient is both solvable and perfect, hence trivial: the derived series of a solvable perfect group is simultaneously constant and eventually trivial. Therefore

$$B/K_{j,\Sigma} \simeq E_j \times R_\Sigma. \tag{E.16.2}$$

The same argument applies to the image of F'. First F maps onto R_Σ, since $B/F \simeq \mathfrak{S}_3$ and a quotient of a perfect group that is solvable is trivial. Then F' also maps onto R_Σ, which is perfect. Its image in E_j is solvable, so its joint image is the product of its two images.

For each E.15 shadow use its automorphism on E_j and the identity on R_Σ. Under (E.16.2) the conjugator is $(f_j, 1)$, which is in the image of F' by the preceding paragraph. Both hexagons hold coordinatewise with multiplier one, and the product map is an automorphism. These constructions commute with every transition map. Equivalently, compactness applied to the joint images of $\widehat{F}'$ gives one lift of the compatible pair $(\{f_j\}, 1)$. This proves the claimed compatible family. For one fixed arbitrary perfect target R, the same subdirect-product and F' arguments apply directly to $E_j \times R$ and give the additional assertion. For a collection of such targets the proof above uses perfection of their *joint image*; the simple-factor induction proves exactly that property for the simultaneous family in (E.16.1).

For the final comparison set $L = \gamma_4 N\, N^5$. It is characteristic in N, hence normal in B, and has finite index because N is finitely generated and its class-three exponent-five quotient is finite. The quotient B/L is solvable. Every homomorphism from (E.16.2) to a solvable group kills R_Σ, since its image is perfect. Therefore a factorization of the natural map $B \to B/L$ through (E.16.2) would factor through E_j and kill $\gamma_3 N$.

Here is an explicit word that it does not kill. Send the first two free generators of N to

$$a = I + E_{12} + E_{34}, \qquad b = I + E_{23} \quad \text{in } U_4(\mathbb{F}_5),$$

and send the others to the identity. The target has class three and exponent five: commutators add superdiagonal degrees, and $(I + A)^5 = I$ for a strictly upper triangular 4×4 matrix A in characteristic five. With $[a, b] = aba^{-1}b^{-1}$, direct multiplication gives $[[a, b], a] = I + 2E_{14} \neq I$. The program `de_solvable_barrier.py` checks these literal matrices. Hence $[[n_0, n_1], n_0] \in \gamma_3 N \setminus L$, contradicting that factorization. This proves the claimed finite comparison and the exact scope of the enlarged family. □

5.7 A four-strand pentagon constraint on the CM fibres

Statement E.17 (Proved). The complete Ferrari-homology pentagon test and its affine fibres. *Put $H_3 = \mathbb{F}_3^2 \rtimes \mathrm{SL}_2(\mathbb{F}_3)$, with the standard left action on column vectors. The assignments*

$$\begin{aligned} \rho_4(\sigma_1) &= (0, A), \quad A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \\ \rho_4(\sigma_2) &= (0, B), \quad B = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}, \\ \rho_4(\sigma_3) &= (e_1, A) \end{aligned} \tag{E.17.1}$$

define an epimorphism $B_4 \to H_3$. Let $L_4 = \ker \rho_4 \cap PB_4$. Then L_4 is normal in B_4, is contained in PB_4, and $|B_4/L_4| = 5184$.

The complete pentagon in PB_4/L_4 is a well-defined condition on the G_6 conjugator coordinate used in E.1. Among the 24 particular G_6 representatives retained in E.14, exactly nine satisfy it. They have indices

$$3, 4, 5, 7, 8, 14, 15, 17, 23; \tag{E.17.2}$$

the extra-coset indices among these are $4, 7, 15$. The other fifteen particular representatives cannot be projections of full $\widehat{\mathrm{GT}}$ elements.

Every one of the 24 whole CM matrix fibres, *however, contains a gentle G_6 shadow with multiplier one whose actual commutator word satisfies this pentagon. In the 18-parameter three-primary presentation of E.10, before quotienting by cusp conjugations, its equation is one nonzero affine linear equation. Its solution space has dimension 17 and contains points with full generation rank 97 and with vanishing class-two cusp obstructions of E.11. This is a finite pentagon test; no lift through all four-strand quotients is asserted.*

Proof. We use the multiplication $(u, M)(v, T) = (u + Mv, MT)$ in H_3. The three assignments in (E.17.1) satisfy both adjacent braid relations and the relation between the distant generators, by direct multiplication. Their Ferrari kernel elements

$$\xi_1 = \sigma_3\sigma_1^{-1}, \qquad \xi_2 = \sigma_1\sigma_2\sigma_1\sigma_3^{-1}\sigma_2^{-1}\sigma_1^{-1}$$

map to $(e_1, I), (e_2, I)$, while A, B generate $\mathrm{SL}_2(\mathbb{F}_3)$. This proves surjectivity. These are the actual free Ferrari generators and their homology action: the source comparison is [NTY10, (2.9) and (4.15)]. The program `de_ferrari_pentagon.py` checks the literal relations and enumerates the joint image with the adjacent transpositions in $\mathfrak{S}_4$; its order is 5184. This is precisely B_4/L_4.

Write a_{ij} for the standard pure Artin generators. We evaluate the complete group equation

$$\begin{aligned} &f(a_{12}, a_{23}a_{24})f(a_{13}a_{23}, a_{34}) \\ &\qquad = f(a_{23}, a_{34})f(a_{12}a_{13}, a_{24}a_{34})f(a_{12}, a_{23}). \end{aligned} \tag{E.17.3}$$

Each displayed pair defines an actual homomorphism $F_2 \to PB_4$, followed by ρ_4. All five factors are pure, so the permutation component of (E.17.3) is already the identity. We also retain the evaluation at $((\sigma_1\sigma_2)^3, \sigma_3^2)$, which is the conjugator in the standard Drinfeld image of σ_3.

The simultaneous images of the 97 literal generators of N under these six homomorphisms commute and have exponent three. The program checks every generator cube and all $\binom{97}{2} = 4656$ commutators. It follows that each homomorphism kills $N'N^6$. Thus all six evaluations factor through the actual quotient $F_2/(N'N^6)$, which is the pure free factor in G_6. Corrections by the whole kernel have zero image in these targets. Consequently a failure of (E.17.3) at the computed representative is a failure for every lift of that G_6 conjugator. The retained E.14 word differs from the base word evaluated here by elements of $N'N^6$, so this comparison applies to the E.14 representatives themselves. Complete multiplication of all five factors gives exactly (E.17.2).

Here is the exhaustive affine reduction for whole CM matrix fibres. Let $P = \binom{B_0}{A_0}$ be the 194×18 parameter matrix retained in E.10. A parameter $z \in \mathbb{F}_3^{18}$ changes the two conjugators by $n(B_0 z), n(A_0 z)$. For an actual integral change b in the first conjugator, put

$$\ell(b) = \frac{\mathrm{ab}_{F,2}(n(b)) - \mathrm{ab}_{F,1}(n(b))}{2} \pmod 3.$$

The numerator is even by the matrix B_N of E.9. The new gauge is $g + 2\ell(b)$ modulo six. If $f_\beta = s\beta s\beta^{-1}$ denotes the ungauged base conjugator, the actual new word is

$$f_z = y^{g+2\ell(b)}(sn(b)s)f_\beta n(b)^{-1}x^{-g-2\ell(b)}. \tag{E.17.4}$$

This identity follows by inserting $s^2 = 1$ and $sxs = y$; it is the precise gauge comparison used in the calculation.

Embed each affine group in 3×3 matrices. On the commuting exponent-three images of N, the formulas

$$\log(I + D) = D + D^2, \qquad \exp(L) = I + L + 2L^2 \quad \text{over } \mathbb{F}_3$$

are inverse and turn products into sums. Their nilpotent matrices have cube zero, and every commutator is zero. A commuting family of nilpotent endomorphisms admits a common complete invariant flag: choose a nonzero common kernel vector, pass to its quotient, and repeat. In that basis these matrices are strictly upper triangular, so every product of three is zero. Expanding the two displayed polynomials therefore proves $\log(uv) = \log(u) + \log(v)$

for these commuting matrices, including characteristic three. The joint linear map recording the six logarithms of $n(b)$, those of $sn(b)s$, and $\ell(b)$ has rank two on the 18 parameter columns. The program `de_ferrari_fibres.py` retains that exact matrix, its two-column section, and its 16-dimensional kernel. It enumerates all nine values of the two-dimensional image. Three independent literal-word substitutions per fibre check (E.17.4) against the logarithmic evaluation.

For every one of the 24 fibres, the pentagon residual on those nine values is an affine linear map of rank one. Every value is retained in the JSON certificate, and direct substitution checks its equality with the retained linear map. Exactly three values satisfy the pentagon. Each also gives an invertible induced action on Ferrari homology; its two columns are the images of ξ_1, ξ_2 under the three Drinfeld generator formulas. The program checks both intertwining relations and all braid relations after this substitution.

Finally, in the remaining 16 parameters the program retains an actual point in every fibre at which the five class-two cusp equations of E.11 vanish and the complete 97×97 generation matrix is invertible. These are positive witnesses, not exclusions from a failed search. Their three-primary changes are represented by four times the corresponding integral N words, so their two-primary homology component is unchanged. The literal cusp relation in G_6, both generation ranks, the exact integral gauge, zero F-abelianization, and all five pentagon factors are checked again. The resulting commutator words and every parameter are retained. This proves nonemptiness in all 24 whole matrix fibres and the asserted finite scope. □

Statement E.18 (Proved). The next Ferrari central layer adds no equation on these affine fibres. *Let $\mathcal{H}_3$ be the class-two exponent-three free group on two generators, in coordinates $(u, v, z) \in \mathbb{F}_3^3$ with multiplication*

$$(u, v, z)(u', v', z') = (u + u', v + v', z + z' + 2(uv' - vu')).$$

The standard action of $\mathrm{SL}_2(\mathbb{F}_3)$ on (u, v) fixes z. Put $\widetilde{H}_3 = \mathcal{H}_3 \rtimes \mathrm{SL}_2(\mathbb{F}_3)$. The assignments

$$\sigma_1 \longmapsto ((0,0,0), A), \qquad \sigma_2 \longmapsto ((0,0,0), B), \qquad \sigma_3 \longmapsto ((1,0,2), A) \tag{E.18.1}$$

define an epimorphism $\widetilde{\rho}_4 : B_4 \to \widetilde{H}_3$, where A, B are those in (E.17.1). The normal pure kernel $\widetilde{L}_4 = \ker \widetilde{\rho}_4 \cap PB_4$ has index 15552. Reduction of the Heisenberg central coordinate gives the precise quotient $\widetilde{H}_3 \to H_3$ of E.17.

All six evaluations used in E.17 still factor through G_6. On each of the 24 whole affine CM fibres, their joint parameter image has dimension two, with exactly the same kernel as in E.17. The complete pentagon has exactly the same zero set on this image: three of its nine points. Every retained E.17 literal witness satisfies the complete pentagon and all three braid relations in $\widetilde{H}_3$, and its three braid images generate that group. Thus this additional central layer preserves all 24 nonempty fibres and introduces no further affine equation.

Proof. In the usual Baker–Campbell–Hausdorff coordinates of the free class-two group, write z_0 for the central coordinate and take the free generators to have $z_0 = 0$. The literal Ferrari actions (4.15) of [NTY10] have central linear terms $(0, 1)$ for A and $(1, 0)$ for B. The exact change of coordinates

$$z = z_0 - u - 2v$$

removes both terms: for $\ell = (1, 2)$, their rows are $\ell A - \ell = (0, 1)$ and $\ell B - \ell = (1, 0)$. Consequently the actual Ferrari generators become $(e_1, 2), (e_2, 1)$, and their commutator is $(0, 0, 1)$. This proves that (E.18.1) uses the actual Ferrari quotient. Direct multiplication verifies the braid relations and the four literal free-generator action formulas. These elements generate the Heisenberg kernel and $\mathrm{SL}_2(\mathbb{F}_3)$, proving surjectivity and the order $27 \cdot 24 = 648$.

The program `de_ferrari_class2_model.py` also verifies the Heisenberg multiplication and all 24 matrix actions exhaustively, then enumerates the joint image with $\mathfrak{S}_4$, of order 15552.

For each of the six evaluation homomorphisms, the simultaneous images of N have order at most nine. The program `de_ferrari_class2_fibres.py` enumerates each such subgroup and verifies that it is elementary abelian. In the seven displayed coordinates (a, b, c, d, u, v, z), subtracting the identity gives additive coordinates on each of these subgroups; every pair product is checked. All generator cubes and all 4656 generator commutators are therefore trivial. This proves the factorization through G_6.

Stack these additive coordinates for $n(B_0t)$ and $sn(B_0t)s$, together with the gauge row $\ell(B_0t)$ of E.17. The retained exact matrix has rank two. Stacking it with the old E.17 matrix still has rank two, so their kernels in $\mathbb{F}_3^{18}$ coincide. Formula (E.17.4) remains an exact literal-word identity in the larger group. The calculation substitutes all nine values of the two-dimensional image in every fibre. Its full three-coordinate pentagon residual vanishes precisely at the three points where the E.17 two-coordinate residual vanishes. Three further literal substitutions per fibre check the additive-coordinate evaluation. Finally, each of the 24 saved E.17 words is substituted directly; all five factors, all braid relations, and generation of the order-648 group are checked. The JSON retains every one of the $24 \cdot 9$ residuals, and the compressed certificate retains the complete parameter matrix, its section, and its kernel. □

Statement E.19 (Proved). Complete compatible corrections at the modulus-nine Ferrari test. *For $R = \mathbb{Z}/9$, let $\mathcal{H}(R)$ have coordinates $(u, v, z) \in R^3$ and multiplication*

$$(u, v, z)(u', v', z') = (u + u', v + v', z + z' + \tfrac{1}{2}(uv' - vu')), \qquad \tfrac{1}{2} = 5 \text{ in } R.$$

Put $H_9 = \mathcal{H}(R) \rtimes \mathrm{SL}_2(R)$, with the linear action on (u, v) and the identity on z. The assignments (E.18.1), with the final central coordinate changed from 2 to 5 and the matrices taken modulo nine, define an epimorphism $\rho_{4,9} : B_4 \to H_9$. Its composition with reduction modulo three is precisely $\widetilde{\rho}_4$ of E.18.

Let $\theta_i : F_2 \to H_9$, $1 \le i \le 6$, be the six actual evaluation maps of E.17. For $B = C_2 * C_3$, choose representatives b_q of $B/F_2 \simeq \mathfrak{S}_3$ and set

$$K_6 = N'N^6, \qquad K_{3,9} = K_6 \cap \bigcap_{i=1}^{6} \bigcap_{q \in B/F_2} b_q^{-1} \ker(\theta_i) b_q, \qquad D_9 = B/K_{3,9}. \tag{E.19.1}$$

Thus $K_{3,9}$ is normal of finite index in B and lies in F_2. The kernel of $D_9 \to G_6$ is $\mathbb{F}_3^4$, and the image of $K_6 \cap F_2'$ in that kernel is $\mathbb{F}_3^3$.

For each of the 24 particular G_6 points retained in E.17, all three commutator-correction parameters can be solved simultaneously. The gentle norm equations have rank two. The modulus-nine pentagon has rank three and selects exactly one correction in this three-dimensional image. The resulting literal words have zero integral F_2-abelianization, satisfy both complete hexagons in D_9, give surjective three-strand shadow maps, and satisfy the complete pentagon in H_9. Their three four-strand braid images satisfy all braid relations and generate H_9.

This yields nonempty compatible finite tests for all 24 CM fibres. In this particular construction $N'N^{18} \subset K_{3,9}$: the three-strand quotient is an explicit quotient of G_{18}. The new condition is the specified four-strand pentagon; no cofinal extension follows from this statement.

Proof. The exact coordinate change from the ordinary free class-two coordinates is $z = z_0 + (u - v)/2$. It removes the central terms in the two literal Ferrari actions over every ring in which two is invertible. The free generators therefore map to $(1, 0, 1/2)$ and $(0, 1, -1/2)$.

Direct multiplication verifies their four action formulas and all three braid relations over R. They generate $\mathcal{H}(R)$, and the two elementary matrices A, B generate $\mathrm{SL}_2(R)$: row reduction over the local ring R first chooses a unit in the first column, clears the other entry, and then expresses the resulting diagonal matrix as a product of elementary matrices. This proves the asserted epimorphism. Its target has order $9^3 \cdot 648 = 472392$.

Here is a faithful model for the core in (E.19.1). An element $b \in B$ has permutation $q \mapsto q\bar{b}$ of the six cosets and coordinates

$$h_{q,i}(b) = \theta_i(b_q b b_{q\bar{b}}^{-1}). \tag{E.19.2}$$

The product rule is $h_{q,i}(bc) = h_{q,i}(b)h_{q\bar{b},i}(c)$. Together with the original map to G_6, this is an actual faithful representation of D_9; its kernel is exactly (E.19.1). The program `de_ferrari_mod9_lift.py` constructs the six representatives and checks $s^2 = r^3 = 1$ in this model. Its evaluation on x, y is checked against the six original maps θ_i.

Reduction modulo three maps every K_6 coordinate to the identity by E.18. The kernel of $H_9 \to \widetilde{H}_3$ is elementary abelian: subtract the identity matrix and the zero Heisenberg vector, divide all coordinates by three, and reduce modulo three. Products add because every cross term contains a factor nine. The 36 coordinates in (E.19.2) therefore give an injective linear coordinate map on the image of K_6.

The complete K_6 image is computed from the 4656 pairwise commutators of the 97 actual Schreier generators n_i of N, their sixth powers, and their conjugates by the n_i. These normally generate $N'N^6$ in N. All pairwise commutator images are zero, and all ninth powers are the identity. Thus this particular core factors through G_{18}. The span of the sixth-power images has rank four; stability under every n_i is checked. This proves the exact kernel assertion.

The zero-abelianization restriction is imposed integrally. With the previously retained matrix $B_N = \left(\begin{smallmatrix} 6 & 4 \\ 0 & 2 \end{smallmatrix}\right)$, write

$$\mathrm{ab}_{F_2}(n_i) = B_N h_i, \qquad H = (h_i)_{0 \le i < 97}.$$

The columns numbered $0, 8$ form a determinant-one minor of H. They give an explicit integral basis of the rank-95 lattice $\ker H$. Modulo the commutator image, every element of K_6 is represented by $\prod_i n_i^{6a_i}$. Its F_2-abelianization vanishes exactly when $a \in \ker H$. Evaluating this entire integral lattice gives a rank-three image. The code retains three literal words forming a basis; their integral abelianizations are checked to be zero. Their images are also checked to be stable under s, r. This proves that the allowed correction space is the entire image of $K_6 \cap F_2'$, with no omitted power corrections.

For a retained E.17 conjugator β, put

$$S_0 = \beta s \beta^{-1}, \qquad t = sr, \qquad R_0 = S_0 t.$$

Let $M = \mathrm{im}(K_6 \cap F_2' \to D_9)$, and write S, R for conjugation by S_0, R_0 on M. Replacing S_0, R_0 by $S_0 k, k^{-1} R_0$ preserves their product. Since M is abelian, the two torsion equations are exactly

$$(1 + S)k = 0, \qquad (1 + R + R^2)k = R_0^3. \tag{E.19.3}$$

All terms lie in the explicitly computed image M; the code checks this by expressing them in the retained basis. These equations have rank two in every fibre.

Let g be the exact integral cusp gauge of the E.17 word. The corrected conjugator is the literal commutator word

$$f = s x^g S_0 k x^{-g}. \tag{E.19.4}$$

For each evaluation map, changing k changes f by an element of the elementary abelian congruence kernel. Multiplying the five pentagon factors therefore gives an affine linear

equation on M. The calculation evaluates all $3^3 = 27$ correction values in each of the 24 fibres, checking every coefficient of this linear description. The pentagon matrix has rank three; the three braid-relation matrices have rank zero. The joint system with (E.19.3) is consistent of rank three, and so selects one point of M in every case.

The retained point is then verified by complete literal substitution, without using only its linear equations. The two hexagons are checked in all 36 induced coordinates, and hold in G_6 by E.17; hence they hold in the faithful combined quotient D_9. The word (E.19.4) has zero integral abelianization because both its base word and k do.

For generation, the new three-strand map is already onto G_6. Substitution of all 97 generators, their commutators and sixth powers, followed by closure under their conjugations, gives rank four in the kernel of $D_9 \to G_6$. Thus the map is onto D_9. This supplies the finite gentle-shadow generation condition, with multiplier one. For the four-strand map, the first two matrix images generate all 648 elements of $\mathrm{SL}_2(R)$, as directly enumerated. The two Ferrari generators have trivial matrix part and determinant nonzero modulo three on their two Heisenberg homology coordinates. They therefore generate the finite three-group $\mathcal{H}(R)$ by its Frattini quotient. The whole four-strand image is H_9.

Every correction word, source word, linear matrix and right-hand side is retained in the JSON and compressed certificates. The distinction between these specified finite tests and a compatible lift through every four-strand quotient is exactly the untested extension beyond (E.19.1); it is not removed here. □

5.8 Arbitrary relative kernels supported away from two and three

Statement E.20 (Proved). Compatible extension through every relative quotient of order prime to six. *In this statement $N = N_B$ is the rank-97 kernel of $B = C_2 * C_3 \to Q$, and hats denote full profinite completions. Let $\mathscr{P}$ be the set of primes other than two and three. For each open subgroup $L \triangleleft \widehat{B}$ contained in $\widehat{N}$ with $[\widehat{N} : L]$ prime to six, and integers $a, b \geq 1$, put*

$$K_{a,b,L} = \overline{(\gamma_3 N\, N^{2^a 3^b})} \cap L. \tag{E.20.1}$$

Every one of the 24 admissible CM matrix pairs has a compatible family of gentle shadows at the inverse images of these subgroups in $\widehat{B}_3$, with multiplier one. The family is compatible as a, b increase and as L decreases, and its projection to every quotient $B/(\gamma_3 N\, N^j)$ in E.15 is the family constructed there. In particular it includes simultaneous refinement by arbitrary relative pro-p quotients for every $p \geq 5$, of every nilpotency class, and by finite relative kernels whose orders involve several primes in $\mathscr{P}$.

Here the three-strand action is constructed on one actual profinite quotient, together with a conjugator in the image of $\widehat{F}_2'$. No extra pentagon assertion is made. The natural map to the finite quotient

$$B/(\gamma_4 N\, (\gamma_2 N)^3 N^9) \tag{E.20.2}$$

factors through none of the quotients in (E.20.1). Its normal kernel over Q is a finite three-group with a nonzero third lower-central term, exhibited below. Thus (E.20.1) supplies further compatible refinements while leaving this specified relative three-primary comparison.

Proof. Write $\mathcal{N}_{23}$ for the product of the free class-two pro-two and pro-three completions of N and let E_{23} be the corresponding quotient of $\widehat{B}$. It fits into the exact sequence

$$1 \longrightarrow \mathcal{N}_{23} \longrightarrow E_{23} \longrightarrow Q \longrightarrow 1.$$

Let $R_{\mathscr{P}}$ be the intersection of the kernels of all continuous finite quotients of $\widehat{N}$ whose orders have prime factors only in $\mathscr{P}$. This subgroup is characteristic in $\widehat{N}$, hence normal in $\widehat{B}$. Put

$$E_{\mathscr{P}} = \widehat{B}/R_{\mathscr{P}}, \qquad \mathcal{N}_{\mathscr{P}} = \widehat{N}/R_{\mathscr{P}}. \tag{E.20.3}$$

Taking the $\widehat{B}$-core of a finite-index normal subgroup of $\widehat{N}$ preserves this prime support: its quotient embeds in a product of at most $|Q|$ conjugate quotients. Thus the subgroups L in the statement give the full topology of (E.20.3).

The simultaneous quotient is exactly the fibre product

$$H = E_{23} \times_Q E_{\mathscr{P}}. \tag{E.20.4}$$

To prove that the natural map $\widehat{B} \to H$ is onto, first take finite quotients of the two normal kernels. The image of $\widehat{N}$ in their product surjects onto both factors. The two Goursat quotient groups have a common order dividing a number supported on $\{2,3\}$ and a number supported on $\mathscr{P}$. That order is one. Hence the image of $\widehat{N}$ is the full product. Lifting the common Q coordinate then gives surjectivity on the entire finite fibre product. The image of the compact group $\widehat{B}$ is closed; its surjectivity in every finite projection proves (E.20.4).

The kernel J of $\widehat{B} \to H$ satisfies

$$J \subset \widehat{N}' \subset \widehat{F}_2'. \tag{E.20.5}$$

Indeed the $\mathcal{N}_{23}$ coordinate detects every two- and three-primary coordinate of $\widehat{N}^{\mathrm{ab}} \simeq \widehat{\mathbb{Z}}^{97}$, and $\mathcal{N}_{\mathscr{P}}$ detects every other primary coordinate. All these coordinates vanish for an element of J, proving the first inclusion. This also proves the precise commutator comparison needed after gluing. Let H_F be the inverse image of F_2/N under $H \to Q$. There is a continuous homomorphism

$$H_F \longrightarrow \widehat{F}_2^{\mathrm{ab}}, \qquad \ker(H_F \to \widehat{F}_2^{\mathrm{ab}}) = \mathrm{im}(\widehat{F}_2' \to H_F). \tag{E.20.6}$$

This is the map induced by abelianization, using (E.20.5); the asserted equality follows by taking any lift in $\widehat{F}_2$.

Fix one CM pair and use the ungauged two- and three-primary automorphism α_{23} constructed in E.15. Its action on Q is the identity, and it fixes the actual cusp element $t = sr$. Therefore

$$\alpha : H \longrightarrow H, \qquad (h_{23}, h_{\mathscr{P}}) \longmapsto (\alpha_{23}(h_{23}), h_{\mathscr{P}}) \tag{E.20.7}$$

is a continuous automorphism. It is invertible by the same formula with α_{23}^{-1} and fixes t. This construction uses the identity on the entire relative $\mathscr{P}$ quotient, including all of its commutator layers.

Take exactly the gauge $g \in \widehat{\mathbb{Z}}$ of E.15: its two- and three-primary components are the gauges there, and $g_p = 0$ for every $p \in \mathscr{P}$. With $x = t^{-2}$, define

$$\alpha' = \mathrm{Inn}(x^g) \circ \alpha, \qquad S' = \alpha'(s), \quad R' = \alpha'(r), \quad f = sS'. \tag{E.20.8}$$

The gauge centralizes t, so $S'^2 = R'^3 = 1$ and $S'R' = t$. The word f belongs to H_F: the common Q coordinate is precisely $sx^{\bar{g}}sx^{-\bar{g}}$, as in E.15.11, and its image in B/F_2 is the identity.

We check its full abelianization, rather than just its reduction at two and three. Formula E.15.12 gives

$$\mathrm{ab}_{F_2}(f) = \mathrm{ab}_{F_2}(s\alpha(s)) + (-g, g). \tag{E.20.9}$$

The selected two- and three-primary gauges make these components zero. For $p \in \mathscr{P}$, the abelianization map to $\mathbb{Z}_p^2$ factors through $E_{\mathscr{P}}$: its restriction to $\widehat{N}$ has finite p-group quotients, all retained in (E.20.3). On that component $\alpha(s) = s$, so the first term of (E.20.9) is zero,

and $g_p = 0$. Consequently every primary component vanishes. By (E.20.6), f is the image of an actual element of $\widehat{F}_2'$.

The torsion-to-hexagon comparison in E.9 applies to these exact elements. In particular $\alpha'(x) = x$, and, since $y = sxs$,

$$\alpha'(y) = S'xS' = f^{-1}yf. \tag{E.20.10}$$

Both complete gentle hexagons hold with multiplier one. The component automorphism α_{23} preserves every class-two finite level used in E.15; the identity and its final inner conjugation preserve every normal subgroup defining a quotient of $E_{\mathscr{P}}$. Thus α' induces an automorphism at every finite quotient in (E.20.1). This proves generation, both equations, and all transition compatibilities at once. Projection to the class-two quotient of each primary kernel gives the same ungauged actions and the same single gauge as E.15. Hence the projected family is exactly the one specified there and retains its CM pair.

We finish by proving the remaining finite comparison. Set $L_3 = \gamma_4 N\,(\gamma_2 N)^3 N^9$, with the closed interpretation in completions. Its quotient in N is finitely generated, has class at most three and exponent dividing nine, and hence is a finite three-group. Finiteness follows directly by collecting words into the finitely many basic commutators of weights at most three, each with bounded exponent. The subgroup L_3 is characteristic in N, so B/L_3 is an actual finite quotient.

The map from N to $U_4(\mathbb{F}_3)$ given by

$$n_0 \longmapsto I + E_{12}, \quad n_1 \longmapsto I + E_{23}, \quad n_2 \longmapsto I + E_{34}, \quad n_i \longmapsto I \quad (i > 2) \tag{E.20.11}$$

kills L_3. The target has class three, its commutator subgroup has exponent three, and $(I + A)^3 = I + A^3$ is central for a strictly upper triangular 4×4 matrix A over $\mathbb{F}_3$; thus its exponent divides nine. Literal multiplication gives

$$[[I + E_{12}, I + E_{23}], I + E_{34}] = I + E_{14} \neq I. \tag{E.20.12}$$

It follows that $[[n_0, n_1], n_2]$ has nonzero image in N/L_3.

Suppose the natural quotient map $B \to B/L_3$ factored through one finite fibre product in (E.20.1). The induced homomorphism to Q agrees with its given projection, because the maps agree on the image of B. The normal kernel supported on $\mathscr{P}$ therefore maps into N/L_3. Its image has order prime to three and also three-power order, so it is trivial. The map would factor through the class-two $\{2, 3\}$ component and would kill $\gamma_3 N$, contradicting (E.20.12). This proves the claimed failure of factorization by an exact quotient map.

The program `de_coprime_gluing_root.py` checks (E.20.11)–(E.20.12), enumerates all 729 target elements, verifies their ninth powers and central cubes, and verifies the indicated commutator-subgroup exponent and central filtration. The lifting and compatibility argument uses the proved profinite morphisms (E.20.3)–(E.20.10). □

5.9 A joint Ferrari test retaining cubic powers and fourth-degree commutators

Statement E.21 (Proved). Complete compatible corrections in the fourth-degree Ferrari quotient. *Let*

$$\mathcal{A}_d = \mathbb{F}_3\langle X, Y\rangle/(\textit{monomials of degree} > d), \qquad K_d = \langle 1 + X, 1 + Y\rangle \subset \mathcal{A}_d^{\times}.$$

On the actual free generators ξ_1, ξ_2 of the Ferrari kernel take the automorphisms

$$\begin{aligned} a:&\quad \xi_1 \mapsto \xi_1, && \xi_2 \mapsto \xi_2\xi_1,\\ b:&\quad \xi_1 \mapsto \xi_1\xi_2^{-1}, && \xi_2 \mapsto \xi_2. \end{aligned} \tag{E.21.1}$$

Their actual finite image on K_d is denoted C_d. The assignments

$$\sigma_1 \mapsto (1, a), \qquad \sigma_2 \mapsto (1, b), \qquad \sigma_3 \mapsto (\xi_1, a) \tag{E.21.2}$$

define an epimorphism $B_4 \to K_d \rtimes C_d$. Here the cubic powers are retained: $(1+X)^3 = 1+X^3$ and $(1+Y)^3 = 1+Y^3$.

Let T_4 be the joint image of (E.21.2) for $d = 4$ and the modulus-nine representation $\rho_{4,9}$ of E.19. Its reduction to the E.18 target is surjective, with elementary abelian kernel $\mathbb{F}_3^{12}$. Consequently

$$|T_4| = 344373768, \qquad |B_4/(\ker(B_4 \to T_4) \cap PB_4)| = 8264970432. \tag{E.21.3}$$

Form the three-strand core by the six evaluation maps and all six B/F_2 conjugates, exactly as in (E.19.1), now with target T_4, and intersect it with K_6. Denote this kernel by $K_{3,4}$ and put $D_4 = B/K_{3,4}$. Its kernel over G_6 has dimension seven over $\mathbb{F}_3$, the image of $K_6 \cap F_2'$ has dimension six, and the image of N' has dimension three.

For each of the 24 retained E.17 G_6 points, the complete gentle norm equations on this six-dimensional correction image have rank four; the simultaneous pentagon in the two targets has rank five. The combined system is consistent of rank five, hence has an affine line of corrections. An actual commutator word in each line is retained. It satisfies both complete hexagons in D_4, gives a surjective three-strand shadow, satisfies the full pentagon in T_4, and defines a surjective four-strand braid map to T_4. The three- and four-strand full twists are preserved.

Thus all 24 CM fibres remain nonempty in this joint finite test. In the specified three-strand quotient, $\gamma_3 N$ and N^{18} are killed. It therefore belongs to the relative class-two coverage of E.15; the new check is the displayed four-strand constraint.

Proof. The substitutions (E.21.1) are the exact Ferrari actions of [NTY10, (2.9) and (4.15)]. Substituting units with constant term one defines algebra automorphisms of $\mathcal{A}_d$: their inverses are the literal inverse free-group substitutions. They preserve K_d. Direct multiplication verifies both adjacent braid relations and the distant braid relation in (E.21.2). The two literal Ferrari words become $(\xi_1, 1), (\xi_2, 1)$, and a, b generate C_d by its definition. This proves the epimorphism. It also specifies the source comparison without replacing the action of B_3 by a homology action.

The calculation first constructs degree three completely. The actual unit group K_3 has order 2187, as enumerated from its two generators. The image C_3 has order 1944, as enumerated from the two exact substitution matrices; its kernel over $\mathrm{SL}_2(\mathbb{F}_3)$ has order 81. Thus this initial four-strand target has order 4251528. Its joint core with the modulus-nine target still has correction dimensions four and three, and all 24 points pass. These checks are retained in `de_ferrari_degree3_model.py` and `de_ferrari_degree3_lift.py`.

For degree four the augmentation ideal has its full 30 monomial coordinates. Multiplication is ordinary truncated noncommutative multiplication over $\mathbb{F}_3$. An automorphism is stored by the two complete augmentation series of its generator images. Its linear operator on all 30 monomials is obtained by substitution. Composition is exact matrix substitution; inverse generator images are obtained degree by degree from the invertible linear part. No division by three occurs. The program `de_ferrari_truncated_model.py` checks all Ferrari and braid identities and compares 1000 successive actual word reductions, including inverses, with the independent complete degree-three model.

Reduction to degree two, followed by the exact change $z = z_0 - u - 2v$ of E.18, gives the specified map to the E.18 target. Its kernel is additive. Indeed a kernel element of K_4 starts in degree three. An automorphism in the kernel changes each generator only in degrees three

and four: its degree-two change is zero by the E.18 comparison. Applying that change to a term of degree three first affects degree five, which is zero here. Two such automorphism changes also commute, since their commutator first affects degree five. Products of two kernel unit changes first occur in degree six. Therefore the exact additive coordinates are the unit's degree-three and degree-four terms and the two automorphisms' changes in those degrees. There are 72 ambient coordinates, with the actual image computed inside them. Together with the seven redundant but injective congruence coordinates of the modulus-nine factor, these give 79 ambient linear coordinates for the joint kernel.

The complete image is obtained by Schreier generators over the order-648 E.18 quotient. Choose one actual word representative t_h for each of its elements. For each of its three generator edges retain

$$t_h \sigma_i t_{h\bar{\sigma}_i}^{-1}. \tag{E.21.4}$$

These generate the kernel of the free three-generator group mapping onto that quotient; their images therefore generate the full kernel of the actual joint braid image. The coordinate matrix has rank 12. Since that kernel is elementary abelian, this proves the first order in (E.21.3).

Every braid generator in this target has order dividing nine, as directly checked. Its abelianization, being a quotient of the cyclic abelianization of B_4, thus has odd order. Every nontrivial quotient of $\mathfrak{S}_4$ has a quotient C_2; this follows from its normal subgroups $1, V_4, A_4, \mathfrak{S}_4$. There is consequently no nontrivial common quotient. Goursat's argument gives the exact joint product $T_4 \times \mathfrak{S}_4$ and proves the second order in (E.21.3).

The induced six-coset representation of the three-strand core is exactly (E.19.2), applied in both target factors. Every K_6 image lies in the just proved additive kernel. The complete normal image of N' is generated by all 4656 pairwise commutators of the actual n_i, followed by closure under all n_i. Its rank is three. Adjoining all sixth powers gives rank seven. Restricting their exponents to the full integral rank-95 abelianization kernel of E.19 gives rank six for $K_6 \cap F_2'$. Literal source words for every basis vector are retained and checked. Stability under both s and r is checked, so these are the exact images used by the core, including all commutator and power corrections.

The same norm identities (E.19.3) and the same literal gauge identity (E.19.4) apply to this six-dimensional space. Their matrices have rank four in every fibre. Products of pentagon factors are affine linear on the abelian correction kernel, with coefficients given by conjugation by their actual base values. The program `de_ferrari_degree4_lift.py` retains the full matrices; the simultaneous pentagon has rank five, and the braid-error matrix has rank zero. The combined augmented system has rank five in every case. All basis-pair substitutions and 100 additional full substitutions per fibre check this affine evaluation; its completeness follows from the additive-kernel identity, not from sampling.

An actual correction in each affine line is then substituted as a literal word. The program checks zero integral F_2-abelianization, both complete hexagons in all 36 induced coordinates, and the complete pentagon in both target factors. As in E.19, generation of the three-strand core is checked by evaluating all n_i, their commutators and sixth powers, and closing under their conjugations. The kernel image has its full rank seven in all 24 cases; the projection to G_6 was already surjective. Hence the three-strand map is onto.

For each corrected four-strand triple, the program `de_ferrari_degree4_generation.py` repeats the entire Schreier calculation (E.21.4). Its projection still has order 648 and its kernel image has rank 12, with the same span as the original target. This proves generation of T_4. It also checks directly

$$(\sigma_1'\sigma_2')^3 = (\sigma_1\sigma_2)^3, \qquad (\sigma_1'\sigma_2'\sigma_3')^4 = (\sigma_1\sigma_2\sigma_3)^4.$$

Finally, `de_ferrari_degree4_lowercentral.py` computes the action of every n_i on the complete rank-three commutator image. Each action is the identity, so the image of $\gamma_3 N$ is zero. Every n_i^9 is also trivial in the induced coordinates. The commutator image has exponent three, so class-two collection shows that every ninth power in N is trivial there. Together with the exponent-six homology component, this proves $\gamma_3 N\, N^{18} \subset K_{3,4}$. It gives the exact claimed comparison with E.15 and the stated finite scope. □

Statement E.22 (Proved). A compatible relative class-three test preserving every retained CM point. *Extend the actual Ferrari Magnus representation (E.21.1)–(E.21.2) to $d = 5$, retaining its joint image T_5 with the modulus-nine target. Define $K_{3,5}$ by all six evaluation maps, all six B/F_2 conjugates, and intersection with K_6, as in (E.19.1). Put $D_5 = B/K_{3,5}$. The specified reduction maps give*

$$D_5 \longrightarrow D_4 \longrightarrow G_6, \qquad T_5 \longrightarrow T_4 \longrightarrow \widetilde{H}_3. \tag{E.22.1}$$

The kernel of $D_5 \to D_4$ is $\mathbb{F}_3^6$. Its entire image is represented by words in $K_{3,4} \cap F_2'$. The kernel of $D_5 \to G_6$ has order 3^{13}. The actual lower-central images of N in D_5 have dimensions

$$\dim \operatorname{im}(\gamma_2 N) = 9, \qquad \dim \operatorname{im}(\gamma_3 N) = 3, \qquad \operatorname{im}(\gamma_4 N) = 1, \tag{E.22.2}$$

where the two nonzero images are elementary abelian three-groups.

All three points of each of the 24 correction lines in E.21 are included in the next test. Every one of those 72 points has a gentle lift to D_5. In each old line exactly one point has a simultaneous pentagon lift; its top correction is unique. Thus every one of the 24 particular E.17 G_6 shadows, including all twelve extra-coset points, has a verified compatible lift through (E.22.1). These literal commutator words give surjective three-strand shadows and surjective four-strand braid maps satisfying the complete pentagon in T_5.

The full target orders are

$$\begin{aligned} |T_5| &= 6\,778\,308\,875\,544, \\ |B_4/(\ker(B_4 \to T_5) \cap PB_4)| &= 162\,679\,413\,013\,056. \end{aligned} \tag{E.22.3}$$

The three- and four-strand full twists are preserved. The multiplier calculation is exhaustive over lifts of the retained G_6 shadows: the exact cusp order in D_5 is 72, so all compatible values $\lambda \equiv 1 \pmod{72}$ and $m = (\lambda - 1)/2 \equiv 0 \pmod{36}$ give exactly the same finite equations. This does not enumerate other multiplier residues or other G_6 points within a CM matrix fibre.

The natural quotient $B \to D_5$ factors through none of the finite quotients in E.20.1. Thus this is a concrete relative class-three extension beyond that family, with positive witnesses in all 24 CM fibres. It is one specified finite refinement, with no all-quotient lifting assertion.

Proof. The 62 augmentation coordinates through degree five are evaluated by the same literal free-group substitutions as in E.21. Reduction to degree four is the specified truncation homomorphism; the modulus-nine factor is unchanged. An element in its kernel has only a degree-five unit term and degree-five changes of the two automorphism generators. Their products add. There are 96 ambient coordinates per evaluation, and 36 evaluations in the induced core. All finite source relations and every transition in (E.22.1) are therefore represented by actual homomorphisms.

We give the complete kernel calculation, including its normal presentation. Write $J = K_6$ and $J' = K_6 \cap F_2'$. In degree four their images have ranks seven and six. Let $b_1, \dots, b_r$ be the retained literal basis words for one of these images. The normal generators of J in N are all 4656 commutators $[n_i, n_j]$ and the 97 sixth powers n_i^6. For J' replace the latter by the 95

products corresponding to the complete integral kernel of the abelianization matrix in E.19. This is exact: modulo N', an element of J' has exponent vector $6a$ with $\mathrm{ab}_{F_2}(a) = 0$.

For each such normal generator w, retain the relation

$$w \prod_i b_i^{-c_i(w)}, \tag{E.22.4}$$

where $c(w)$ is its complete degree-four image coordinate. Also retain every b_i^3, every $[b_i, b_j]$, and every

$$n_k b_i n_k^{-1} \prod_j b_j^{-c_j(n_k b_i n_k^{-1})}. \tag{E.22.5}$$

These normally generate the kernel of the old image. Indeed, in the resulting quotient every original generator and all its N-conjugates are in the commuting exponent-three span of the b_i. Its order is at most 3^r, while the actual old image has order 3^r and is a quotient of it. Hence the two groups coincide and the listed normal kernel is exact.

The program `de_ferrari_degree5_kernel.py` evaluates all 5354 relations for J' and all 5460 for J, then closes their top-degree spans under the 97 actual N generators. Both image spaces have rank six and are equal. Every retained basis word is re-evaluated literally; the J' words have zero integral F_2-abelianization. Stability under s, r is also checked. This proves the full top-kernel assertion and shows that no source power or commutator corrections have been discarded.

Here the choice of companion to a provisional involution must be exact. Let S_0 be an actual word lifting one of the old degree-four solutions and put

$$R_0 = S_0^{-1} t, \qquad S_0 R_0 = t. \tag{E.22.6}$$

The square of S_0 need not be the identity at degree five. Write M for the computed rank-six top kernel, S, R for the conjugation actions of S_0, R_0 on M, and $E_S = S_0^2$, $E_R = R_0^3$. For $k \in M$, the exact corrected images are $S_0 k, k^{-1} R_0$. The complete norm equations are

$$(1 + S)k = -E_S, \qquad (1 + R + R^2)k = E_R. \tag{E.22.7}$$

In the first equation $\mathrm{Ad}(S_0^{-1}) = S$ on M, because $S_0^2 \in M$ and M is abelian. These identities prove (E.22.7) without assuming that the provisional square vanishes. The program asserts (E.22.6) before constructing any linear system.

For each of the three old points in every fibre, the matrix in (E.22.7) has rank four and is consistent. The top pentagon matrix has rank five, and the braid matrix has rank zero. The combined coefficient matrix has rank six. In exactly one of the three old points its augmented rank is also six; in the other two it is seven. Thus precisely one old point lifts and its top correction is unique. The full matrices and right-hand sides for all 72 cases are retained. Affineness follows from the elementary abelian top kernel. All basis-pair substitutions and further complete substitutions audit the implemented linear maps.

The normalized word remains

$$f = s x^g S_0 k x^{-g}, \tag{E.22.8}$$

with the old exact integral gauge. Its two new factors are actual commutator words, so its integral abelianization is zero. Every chosen solution is checked by literal source word substitution, both full hexagons in all 36 induced coordinates, the complete pentagon in both target factors, and all three braid relations. The independent program `de_ferrari_degree5_gentle.py` also constructs and verifies gentle-only witnesses in all 24 CM fibres directly from the norm systems.

For completeness we describe the source and target generation certificates. The kernel over the E.18 target at degree five is of class at most two and exponent three. Its raw coordinates have a bilinear product $v * w = v + w + C(v, w)$, with cross terms confined to degree five. The exact first-kind coordinate is

$$\log(v) = v - \tfrac{1}{2}C(v, v), \qquad \tfrac{1}{2} = 2 \in \mathbb{F}_3. \tag{E.22.9}$$

The group commutator gives the Lie bracket. Products are therefore $a + b + \frac{1}{2}[a, b]$. This is a class-two calculation in characteristic three; no class-three logarithm is used. The formula is checked on actual group products and inverses.

The seven old full-kernel words together with the six new top-kernel words give a 13-dimensional Lie space for the entire kernel over G_6. Its brackets and the s, r actions are checked to remain in the space. For each proposed source map, those literal words are substituted on its new generators. Closing their image under brackets and the two new generator actions gives rank 13. All these elements are images of actual elements of K_6; this rank proves surjectivity on the whole kernel. The map was already onto G_6, so it is onto D_5.

There is also an exact generation argument for every solution of the norm systems, including all 72 old points, without restricting to the selected witnesses. The image of N in the E.18 target is an elementary abelian three-group, as checked there on all generators, powers and commutators. The kernels of the modulus-nine and higher Magnus reductions to that target are three-groups: the matrix congruence kernel and each additive Magnus layer have three-power order. Consequently the image of N in all 36 new evaluation factors is a three-group. The G_6 coordinate also retains the full quotient $N^{\mathrm{ab}}/6$. By Goursat's lemma on the relatively prime normal orders, the actual joint image is therefore

$$\mathrm{im}(N \to D_5) \simeq (N^{\mathrm{ab}}/2) \times P, \qquad P \twoheadrightarrow N^{\mathrm{ab}}/3, \tag{E.22.10}$$

where P is a finite three-group generated by the 97 specified N generators. Its Frattini quotient $P/\Phi(P)$ has dimension at most 97 and surjects onto $N^{\mathrm{ab}}/3 \simeq \mathbb{F}_3^{97}$. Hence the displayed surjection has kernel exactly $\Phi(P)$. Every norm solution induces the fixed automorphism of G_6 and thus an automorphism of Q; it sends N into the subgroup in (E.22.10). Its image surjects onto both $N^{\mathrm{ab}}/2$ and $P/\Phi(P)$. The finite three-group Frattini criterion gives surjectivity onto P; Goursat again gives the full product in (E.22.10). Together with the surjection onto Q, this proves generation of D_5 for every norm solution. The direct rank-13 checks above independently certify each selected pentagon witness.

The complete image of N' is obtained from all pairwise commutators and normal closure under N. It has dimension nine and is abelian. Repeating the commutator operation with N gives dimension three and then zero. These are the exact ranks (E.22.2), with the coordinate matrices and a nonzero third-commutator witness retained.

The four-strand target is computed independently by the 648-vertex Schreier calculation (E.21.4). The Lie span of its full kernel and its brackets has dimension 21; its commutator space has dimension nine. This proves $|T_5| = 648 \cdot 3^{21}$. For each of the 24 corrected triples the same calculation gives the entire same Lie space. It therefore generates T_5. Both full twists are also checked directly. Every braid generator has order dividing nine, so the exact product with $\mathfrak{S}_4$ and the pure quotient order follow by the same Goursat argument as in E.21.

The control program `de_ferrari_degree5_control.py` checks that the induced core cusp has order dividing 36, while its G_6 coordinate has exact order 72. Hence the combined cusp has exact order 72. The pure peripheral powers have orders dividing 36, and the four-strand generators have orders dividing nine. Both complete hexagon powers and all four-strand generator powers are therefore unchanged by every multiplier lift of each retained G_6 shadow. This is the claimed exhaustive period comparison.

The literal identity pair passes all new relations. The program also verifies that the E.17 point numbered zero is a different G_6 point from the identity in E.1: its conjugator and its induced y map are nontrivial. Finally a permanent regression control uses a provisional S_0 with nontrivial square. It checks that $R_0 = S_0 t$ fails the cusp product and (E.22.6) satisfies it. The initially incorrect calculation, its withdrawn negative conclusions, and its exact reconstruction are preserved separately in the audit files.

If the natural map to D_5 factored through a finite quotient in E.20.1, its relative prime-to-six kernel would map trivially: the relative kernel of $D_5 \to Q$ is a $\{2,3\}$-group, and the maps to Q agree on the source B. The map would then factor through the relative class-two component and kill $\gamma_3 N$, contrary to (E.22.2). This proves the last claim by the specified natural quotient maps. □

Statement E.23 (Proved). A relative class-four refinement with complete correction planes. *Extend the same actual Ferrari Magnus representation to degree six, retaining its joint image T_6 with the modulus-nine target. Let $K_{3,6}$ be the intersection of K_6 with the kernels of all six evaluations and all six B/F_2 conjugates, and put $D_6 = B/K_{3,6}$. The actual reduction maps are*

$$D_6 \longrightarrow D_5 \longrightarrow G_6, \qquad T_6 \longrightarrow T_5. \tag{E.23.1}$$

The kernel of $D_6 \to D_5$ is $\mathbb{F}_3^6$; its whole image is represented by words in $K_{3,5} \cap F_2'$. The kernel of $D_6 \to G_6$ is elementary abelian of dimension 19. The successive lower-central images of N are

j	2	3	4	5
$\dim \operatorname{im}(\gamma_j N)$	15	6	3	0.

(E.23.2)

Thus this specified relative source quotient has class four. Its natural map does not factor through D_5, whose fourth-commutator image is trivial by E.22.

Above each of E.22's 24 simultaneous points, the entire next correction set is an affine plane over $\mathbb{F}_3$. All nine points in each plane are surjective gentle shadows and satisfy the complete pentagon. All 216 associated four-strand triples generate the whole target T_6 and preserve the three- and four-strand full twists. In particular all twelve extra-coset CM fibres still have positive witnesses.

The exact target orders are

$$\begin{aligned} |T_6| &= 133\,417\,453\,597\,332\,552, \\ |B_4/(\ker(B_4 \to T_6) \cap PB_4)| &= 3\,202\,018\,886\,335\,981\,248. \end{aligned} \tag{E.23.3}$$

The cusp period remains 72 in the source joint quotient. Every multiplier lift of the retained G_6 shadows therefore has the same equations at this level, as in E.22. This is the complete comparison over the stated 24 points, with all 216 surviving corrections retained for further refinements. No assertion of lifting through all finite quotients follows from (E.23.1).

Proof. The new model uses all 126 augmentation coordinates in $\mathbb{F}_3\langle X, Y\rangle$ through degree six, with the exact two-generator substitution series for the braid action. Its reduction to degree five is literal truncation. The program `de_ferrari_degree6_model_audit.py` checks 400 actual word and inverse comparisons with the preceding model, all braid and Ferrari relations, and 1225 pairwise product comparisons in the kernel over E.18. Powers and third commutators are also checked in that kernel.

For the complete source kernel, the old images of $K_6 \cap F_2'$ and K_6 are verified to be elementary abelian of dimensions 12 and 13. The normal presentation (E.22.4)–(E.22.5) applies with those actual old basis words. For the first image it retains 4751 original generator

relations, 12 basis cubes, 66 basis commutators and 1164 conjugation relations, a total of 5993. For the full image the corresponding counts are 4753, 13, 78 and 1261, a total of 6105. Each relation is evaluated in the actual degree-six group and its top image is closed under all 97 N generators. Both complete kernel images have dimension six and coincide. Literal re-evaluation, integral F_2-abelianization, and s, r stability are checked. This is the program `de_ferrari_degree6_kernel.py`.

For each retained degree-five point, take its actual word S_0 and set

$$R_0 = S_0^{-1}t, \qquad S_0R_0 = t. \tag{E.23.4}$$

The complete top correction space has dimension six. The norm matrix (E.22.7), the pentagon matrix, and their combined matrix all have rank four; the braid matrix has rank zero. Every combined augmented matrix also has rank four. Consequently the complete solution space at each point is an affine plane. The program asserts (E.23.4) before constructing the system and retains all matrices, right-hand sides, particular solutions, and both homogeneous directions. The identities are affine because the degree-six correction kernel is elementary abelian. All basis-pair substitutions and additional complete substitutions check the implemented affine maps.

The normalized conjugator is again $f = sx^g S_0 k x^{-g}$. Both the old and new corrections are literal F_2' words, so its integral abelianization vanishes. Every selected word is checked by independent B-word and F_2-word evaluation, both full hexagons in all 36 induced coordinates, both target pentagons, and every braid relation. The exact Frattini comparison (E.22.10) applies unchanged: the relative image is the product of $N^{\mathrm{ab}}/2$ with a 97-generated three-group surjecting onto $N^{\mathrm{ab}}/3$. It proves source generation for every point of each plane. In addition, direct substitution of the 19 literal full-kernel basis words gives rank 19 for each selected witness.

The lower-central computation uses all 4656 commutators of the actual N generators, their complete normal closure, and successive commutators with those generators. It gives (E.23.2); the full-kernel bracket matrix is zero. Thus all listed nonzero lower-central images are elementary abelian. The complete bases and a nonzero higher-commutator witness are retained.

Only the kernel over the E.18 target is given class-two logarithmic coordinates. At degree six its product has the form $v * w = v + w + C(v, w)$, where C is bilinear and its terms have degrees five or six. A further cross term has degree at least seven. The exact formula (E.22.9), in characteristic three, therefore still identifies this kernel with a class-two Lie group of exponent three. This calculation is not applied to the relative class-four group in (E.23.2).

The complete 648-vertex Schreier calculation in `de_ferrari_degree6_generation.py` gives kernel Lie dimension 30 and derived dimension 18. Hence $|T_6| = 648 \cdot 3^{30}$. It checks the entire same image for all 24 selected triples and both full twists. The separate program `de_ferrari_degree6_wholeplanes.py` then verifies all 216 points, including their four-strand generation.

Here is the exact finite linear comparison used in the latter verification. Let $L(a, b)$ be the matrix of logarithms of the Schreier generators for the triple at plane coordinate $(a, b) \in \mathbb{F}_3^2$. The reduction of every triple to E.18 is fixed. The differences of its Schreier generators lie in the degree-six kernel, which centralizes the kernel over E.18: a commutator with an augmentation or automorphism term of degree at least three has degree at least eight. Exact substitution therefore gives

$$L(a, b) = L(0, 0) + a\big(L(1, 0) - L(0, 0)\big) + b\big(L(0, 1) - L(0, 0)\big). \tag{E.23.5}$$

The code retains these three complete matrices and checks the value at $(1, 1)$ directly. Their commutator space is the same actual 18-dimensional space at every point, because the

differences are central. Appending that space to (E.23.5) has rank 30 at all nine values. These are actual Schreier generators and their actual commutators, so the ranks prove full generation, not just surjectivity on a linear approximation. The same program directly checks both full hexagons, the pentagon, all braid relations, and both full twists at all 216 points.

Finally each original braid generator has order dividing nine, so the Goursat argument with $\mathfrak{S}_4$ in E.21 gives the second order in (E.23.3). The induced core cusp still has order dividing 36, while its G_6 coordinate has exact order 72. Thus all lifts of the retained multiplier residue give the same finite source and target powers. These checks establish the stated finite scope and retain every surviving correction parameter. □

Statement E.24 (Proved). A complete joint binary and ternary Ferrari refinement. *Use (E.21.1)–(E.21.2) in the actual algebra*

$$\mathcal{A}_{2,d} = \mathbb{F}_2\langle X, Y\rangle/(\textit{monomials of degree} > d), \qquad 2 \le d \le 6, \tag{E.24.1}$$

and denote the resulting four-strand target by $T_{2,d}$. All square terms are retained. Let $D_{2,6}$ be the three-strand quotient formed by the six evaluations, their six B/F_2 conjugates, and intersection with K_6, exactly as in (E.19.1). Then

$$\ker(D_{2,6} \to G_6) \simeq \mathbb{F}_2^9, \qquad \operatorname{im}(K_6 \cap F_2' \to D_{2,6}) \simeq \mathbb{F}_2^7. \tag{E.24.2}$$

The actual common source quotient with E.23 is the fibre product

$$D_{2,6} \times_{G_6} D_6, \tag{E.24.3}$$

whose kernel over G_6 is $\mathbb{F}_2^9 \times \mathbb{F}_3^{19}$.

For every one of the 24 retained G_6 points, all compatible multiplier residues in (E.24.3) are $1, 73$ modulo 144. For each residue the complete binary correction system has an affine line of two solutions. Every one combines with every point of E.23's nine-point ternary plane by an actual F_2' word. Thus each retained CM fibre has 36 simultaneous points, for a total of 864. All give surjective three-strand shadows, satisfy both complete target pentagons, and give surjective four-strand braid maps preserving the full twists with the specified multiplier. In particular every extra-coset CM fibre survives this refinement.

The exact binary target order is

$$|T_{2,6}| = 6 \cdot 2^{40} = 6\,597\,069\,766\,656. \tag{E.24.4}$$

The actual joint four-strand image is $T_{2,6} \times T_6$. Its kernel is already contained in PB_4, and its order is

$$880\,164\,249\,471\,192\,366\,606\,672\,986\,112. \tag{E.24.5}$$

These counts and comparisons concern the entire finite correction fibres over the specified G_6 points. They do not enumerate other G_6 points or other initial multiplier residues in a CM matrix fibre, and do not assert full profinite lifting.

Proof. The free-group substitutions of (E.21.1) and all six evaluation pairs are used literally in (E.24.1). In particular $(1+X)^2 = 1+X^2$ is retained. No logarithmic coordinates are used in the binary group. The programs verify the braid relations, the two literal Ferrari generators, their four action formulas, and actual word and inverse truncations. At degree two the free-unit kernel has order 32 and the automorphism image has order 48, giving target order 1536. In degrees two, three and four, all 4656 N-commutators and all 97 sixth powers vanish in the full induced core. Thus the six evaluations at those levels factor through G_6. All 24 retained points pass the pentagon and every braid relation there.

The kernel of reduction from degree six to degree four has only unit terms and generator-image changes in degrees five and six. Its group products add: a product of two unit terms has degree at least ten, and substituting one automorphism change into another has degree at least nine. It is therefore an elementary abelian two-group. This is an exact assertion about the displayed algebra and substitution maps.

The complete source image is obtained from all $[n_i, n_j]$, all n_i^6, and their normal closure under N. For its F_2' intersection, use the entire integral rank-95 abelianization kernel of E.19 for the sixth-power products. At degree five the commutator, full and F_2' ranks are $3, 6, 4$; at degree six they are $6, 9, 7$. Literal basis words, all additive basis-pair products, and s, r stability are checked. This proves (E.24.2) without omitting power corrections.

Fix the selected ternary word S_0 of E.23. For each multiplier residue put

$$R_0 = S_0^{-1} t^\lambda, \qquad S_0 R_0 = t^\lambda. \tag{E.24.6}$$

The seven-dimensional commutator correction image is denoted M_2. The complete norm matrix on M_2 has rank five, the pentagon matrix has rank five, and their combined matrix has rank six. Every augmented matrix has the same rank, for every CM point and both multipliers. The braid matrix has rank zero. All 128 values of the correction space are substituted to check the affine formulas, and every solution is checked in both complete hexagons, the pentagon, all braid relations, and both full-twist equations. The normalized word is $f = sx^g S_0 k x^{-g}$ with the same exact integral cusp gauge.

The binary induced cusp has exact order 16. Its common quotient with the old order-72 cusp therefore has exact order 144. The retained G_6 shadow has $\lambda \equiv 1 \pmod{72}$, which gives exactly the two residues in the statement. The binary braid generators have orders dividing eight; the ternary generators have orders dividing nine. All their required powers are included. The binary and ternary full-twist orders are also computed directly, and the values at both multiplier residues are verified.

We give the common-word comparison explicitly. Every binary correction basis word v is in $K_6 \cap F_2'$. Its binary image has order dividing two, whereas its ternary image has order dividing three by E.23. Thus v^3 has exactly the required binary image and trivial ternary image. Conversely, for each literal word u representing a direction of the old ternary plane, u^{-2} has its original ternary image and trivial binary image. These identities are checked in the full 36-coordinate induced models. Hence the family is represented by the actual words

$$S = S_0 u^{-2} v^3, \qquad R = S^{-1} t^\lambda, \qquad f = sx^g S x^{-g}, \tag{E.24.7}$$

with products of basis words used for the correction coordinates. All the correction words have zero integral F_2-abelianization. The two projection identities prove that (E.24.7) retains every ternary and binary solution, with no choice of incompatible representatives.

The source product (E.24.3) is exact: the image of K_6 surjects onto its two component kernels, whose orders are relatively prime. Goursat's lemma gives their full product; lifting the common G_6 coordinate gives (E.24.3). The same argument and the literal powers just given identify the full commutator correction image. The relative N-image is the product of a finite two-group and a finite three-group. Each is generated by the 97 specified N generators and maps onto N^{ab}/p of dimension 97. Its kernel to this quotient is its Frattini subgroup. The inherited G_6 automorphism therefore proves surjectivity of every source map in the family.

For the binary target, let U_d be the kernel of the linear map $T_{2,d} \to \mathrm{SL}_2(\mathbb{F}_2)$. The literal Ferrari decomposition and the three standard pure braids show that U_d is generated by

$$\xi_1, \quad \xi_2, \quad a^2, \quad b^2, \quad (ab)^3. \tag{E.24.8}$$

Indeed PB_3 is generated by a^2, b^2 and $(ab)^3$, since its third standard generator is $a^{-2}(ab)^3b^{-2}$.

The program `de_ferrari_two_generation.py` collects this group exactly. Unit degree d is assigned weight d, and an automorphism change in degree $d+1$ is assigned weight d. Products add in the first nonzero weight; squares and commutators move to strictly greater weight. The collector reduces by actual group multiplication, adds every remaining square or commutator, and terminates with 40 independent pivot generators. All their squares and pairwise commutators collect to later weights. Their ordered binary products consequently form a subgroup: the verified triangular relations collect every product and inverse. Independence of the first nonzero pivot gives exactly 2^{40} distinct products. All retained generators are actual expressions in (E.24.8), and the five inputs collect in this subgroup. This proves (E.24.4). Complete direct enumerations in degrees two through four independently agree with this collection algorithm.

At degree two the same calculation gives $|U_2| = 2^8$ and $|\Phi(U_2)| = 2^3$. Normality of the subgroup generated by the five squares and their commutators is checked on all five generators. Thus $U_2/\Phi(U_2)$ has dimension five. Since U_6 has the five generators (E.24.8), reduction induces an isomorphism on these Frattini quotients. Every one of the current conjugators is trivial in all six evaluations through degree two, so every corrected triple has precisely the original generators in $T_{2,2}$. Its subgroup therefore surjects onto the Frattini quotient of U_6 and onto $\mathrm{SL}_2(\mathbb{F}_2)$. It is all of $T_{2,6}$. This proves generation for every binary solution, including both points of every line.

The two four-strand targets have no common nontrivial quotient. Their abelianizations are cyclic of orders dividing eight and nine, respectively, because the three braid generators are conjugate and have those orders. Every quotient of $T_{2,6}$ is solvable: U_6 is a two-group and the quotient is $\mathrm{SL}_2(\mathbb{F}_2) \simeq \mathfrak{S}_3$. A common quotient would thus be solvable with trivial abelianization, hence trivial. Goursat proves the asserted direct product, both for the original representation and for every corrected triple. Finally the degree-one binary image is $\mathbb{F}_2^2 \rtimes \mathrm{SL}_2(\mathbb{F}_2)$. On the four points $(0,1),(1,1),(1,0),(0,0)$, the three generators act as $(12),(23),(34)$, checked directly. Thus its permutation coordinate is the actual standard $\mathfrak{S}_4$ map. The kernel is already pure, and multiplying the two exact target orders gives (E.24.5). □

Statement E.25 (Proved). The complete joint quartic distribution test. *Keep B, F, N_B, G_6 and the twenty-four retained words $f_i \in F'$ of E.1, indexed $i = 0, \dots, 23$ in the saved certificate order. Let $q_{24} : F \to \mathrm{SL}_2(\mathbb{Z}/24)$ send*

$$x \longmapsto U^2 = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}, \qquad y \longmapsto W^2 = \begin{pmatrix} 1 & 0 \\ -2 & 1 \end{pmatrix}. \tag{E.25.1}$$

Put $\mathcal{N} = \{2, 3, 4, 6, 12\}$ and retain every coefficient c_N of D.5 modulo twelve. For each i, the complete set of pairs $(c_2, c_3) \in (\mathbb{Z}/12)^2$ attained by words $f_i k$, $k \in N_B'$, satisfying simultaneously

$$\begin{gathered} q_{24}\delta_N(f_i k) = q_{24}(y)^{c_N(f_i k)} q_{24}(f_i k) \quad (N \in \mathcal{N}), \\ c_4 = 2c_2, \qquad c_6 = c_2 + c_3, \qquad c_{12} = 2c_2 + c_3 \pmod{12} \end{gathered} \tag{E.25.2}$$

is exactly $\{a_i\} \times \mathbb{Z}/12$, where

$$\begin{aligned} (a_0, \dots, a_{11}) &= (0,6,0,6,6,0,6,0,4,10,4,10), \\ (a_{12}, \dots, a_{23}) &= (10,4,10,8,4,10,4,10,4,10,4,10). \end{aligned} \tag{E.25.3}$$

For the auxiliary image allowing every correction in $K_6 = N_B' N_B^6$, the exact set is $\{a_i, a_i + 6\} \times \mathbb{Z}/12$. The latter calculation does not impose an additional F' condition on that larger correction group; the former already supplies actual words in F' with unchanged G_6 image.

The joint correction images of N_B' and K_6 have orders

$$2\,717\,908\,992, \qquad 65\,229\,815\,808, \tag{E.25.4}$$

respectively. Thus all twelve retained CM-compatible points and all twelve retained extra points pass this joint test. Each admits every value of the defined coordinate $\rho_3^{\mathrm{dist}} = -c_3 \pmod 4$; restricting both c_2, c_3 to even residues still leaves both values $0, 2$ of $\rho_3^{\mathrm{dist}} \pmod 4$. This assertion concerns exactly the finite identities (E.25.2). It asserts neither generation in their induced cover quotient nor a full profinite GT realization.

Proof. The mod-four images of U^2, W^2 generate the four-group V_4 of E.1. Choosing the unique global sign with reduction in that four-group identifies the image in (E.25.1) with

$$H_{24} = G_2 \times \mathrm{SL}_2(\mathbb{F}_3), \qquad G_2 = \{g \in \mathrm{SL}_2(\mathbb{Z}/8) : g \bmod 4 \in V_4\}, \qquad |G_2| = 32. \tag{E.25.5}$$

Reduction projectively modulo twelve is the original map on F. Consequently $q_{24}(N_B)$ lies in the elementary abelian kernel of reduction from modulus twenty-four to modulus twelve. It follows that q_{24} kills both N_B' and N_B^6. Thus the last matrix in (E.25.2) is the fixed matrix $q_{24}(f_i)$.

Write $H_N = \ker(F \to \mathbb{Z}/N)$ as in D.5 and $A = H_{24} \times \mathbb{Z}/12$. The actual homomorphism

$$\Psi_N : H_N \longrightarrow A, \qquad w \longmapsto (q_{24}\delta_N(w), c_N(w) \bmod 12) \tag{E.25.6}$$

is induced to F using the transversal $1, x, \ldots, x^{N-1}$. For a word w with x exponent s modulo N, its jth coordinate is the image under (E.25.6) of $x^j w x^{-(j+s \bmod N)}$. Multiplication shifts the coordinates by s. This defines an actual homomorphism to the finite wreath product, with the action convention fixed in the program.

Here is the exact reduction used to compute its joint derived image. The image of the x exponent of N_B in $\mathbb{Z}/12$ is the even subgroup of order six. Put $N_0 = N_B \cap H_{12}$. Choose actual words $t_s \in N_B$ for the six shifts. If $n_1, \ldots, n_{97}$ is the literal basis from E.1, the elements

$$t_s n_j t_{s+\exp_x(n_j)}^{-1} \tag{E.25.7}$$

generate N_0. On N_0, taking coordinate zero in each of the five induced representations is a homomorphism into A^5. Furthermore N_B' is the normal closure in N_0 of

$$t_s[n_i, n_j]t_s^{-1} \quad (s \text{ among the six shifts, } i < j). \tag{E.25.8}$$

Indeed the ordinary normal closure of the basis commutators in N_B is N_B', and every conjugating element is a product of an element of N_0 and one of the t_s. This proves the same assertion after taking closures and then any finite image. Thus the normal closure of the images of (E.25.8) under the image of (E.25.7) is exactly the desired joint correction group. It retains all dependencies between the five coordinates. Adjoining the images of n_j^6 gives the image of K_6: modulo N_B' the source is abelian, so these powers generate all sixth powers there. Their shifts are zero since every n_j has even x exponent.

For a faithful finite model, write each member of G_2 uniquely as $\bar{x}^a \bar{y}^b [\bar{x}, \bar{y}]^e$, with $a, b \in \mathbb{Z}/4$ and $e \in \mathbb{Z}/2$. Its two cyclic-four quotient actions, together with its action on the four cosets of $\langle \bar{x}, \bar{y}^2 \rangle$, give a faithful permutation representation of degree twelve. The other factors act on the eight nonzero vectors of $\mathbb{F}_3^2$ and on disjoint cyclic sets of sizes four and three. Hence A has the explicit faithful degree-twenty-seven representation used in the program. Its five induced actions have total degree $27(2 + 3 + 4 + 6 + 12) = 729$; the shift-zero image has degree 135.

The program `bc_distribution_quartic.py` verifies every multiplication in the order-thirty-two model. The 385 projected Schreier generators reduce, by exact membership, to eleven generators. The 11,419 distinct projected commutator seeds reduce to seventeen; closure under all eleven conjugating generators is checked before its order is computed. Every generator retained or added is the image of the displayed actual source words. Exact Schreier–Sims membership gives (E.25.4).

For each i, the program enumerates all 144 pairs (a, b), substitutes $(c_2, c_3, c_4, c_6, c_{12}) = (a, b, 2a, a + b, 2a + b)$, and tests the resulting single correction tuple for membership in each of the two joint groups. Its complete records give (E.25.3) and the auxiliary sets stated above. A tuple in the first group has one preimage $k \in N'_B$, so it supplies one word satisfying all five identities. The three-cover instance was also computed in the full induced ambient group before the shift-zero reduction was introduced; its complete records agree, with correction orders 192 and 576. Source, output, literal input words and generator arrays are saved in `code/bc_distribution_quartic*` and `code/bc_cm_literal_words.json`.

The notation $\rho_N^{\mathrm{dist}} = -c_N$ defines the negative coefficient coordinate. The saved JSON field `admissible_rho3_mod4` records exactly $-c_3 \bmod 4$. Every finite equation above uses c_N itself. D.5 proves the actual double-cover arithmetic comparison $c_2(f_\sigma) = \kappa_2(\sigma)$ in the stated generator convention. The proof here depends only on the five displayed coefficient maps and their complete joint correction image. □

Statement E.26 (Proved). The exact Fermat proper-Tate quotient and its boundary refinement. *Start with the discrete group $B = C_2 * C_3$. Use the literal level-two generators of the E.1 verifier, $x = (r^{-1}s)^2 = (sr)^{-2}$ and $y = (sr^{-1})^2 = sxs$. Let $F = \langle x, y\rangle \triangleleft B$, and put*

$$H = \ker(F \to (\mathbb{Z}/3)^2), \qquad T = H^{\mathrm{ab}}/\langle \textit{the nine cusp classes}\rangle. \tag{E.26.1}$$

The based cubic cover identifies H with the topological fundamental group of the punctured Fermat cubic, and $T \simeq \mathbb{Z}^2$ is its proper integral homology lattice. After profinite completion the corresponding finite cover identifies $\widehat{H}$ with the geometric étale fundamental group, and $\widehat{T} = \varprojlim_m T/mT$ with the full Tate module of its smooth compactification. For $m \geq 1$, set $J_m = \ker(H \to T/mT)$; it is normal in B. There are actual quotient maps

$$B \longrightarrow G_{3m} \longrightarrow B/J_m. \tag{E.26.2}$$

In particular every proper two-power Tate quotient of this specified cover is dominated by the homology tower of E.4. The inverse-limit comparison is the surjection

$$\widehat{B}/\overline{[N_B, N_B]} \longrightarrow \varprojlim_m B/J_m. \tag{E.26.3}$$

These statements retain the intrinsic Fermat curve and its based maps; no CM twist comparison is implicit in them.

The full rank-ten homology quotient $B/[H, H]H^4$ gives additional information. In the ordered basis $\eta_1, \ldots, \eta_{10}$ specified below, the first two literal N_B generators satisfy

$$[n_1, n_2] \longmapsto 2\eta_3 - 2\eta_{10} \neq 0 \quad \textit{in } H^{\mathrm{ab}}/4, \tag{E.26.4}$$

where $n_1 = y^6$ and $n_2 = yx^{-1}yx^{-2}yx^{-1}y$. Consequently the actual common refinement

$$B/(K_6 \cap [H, H]H^4) \longrightarrow G_6 \tag{E.26.5}$$

cannot be dominated by any G_j through the specified maps from B.

Proof. Use the transversal $x^a y^b$, $0 \le a, b < 3$, and order its nontrivial Schreier edges first by (a, b) and then by x, y. The cover is obtained by adjoining $t^{1/3}$ and $(1-t)^{1/3}$; its compactification is $u^3 + v^3 = 1$. At the standard tangent choose the former Puiseux branch and the latter branch with constant coefficient one. The finite-index subgroup H has the full induced profinite topology: the core in B of any finite-index subgroup of H still has finite index. The finite-cover comparison therefore gives precisely the completion and filling maps in the statement. The ten nontrivial Schreier edges are the generators used in `de_fermat_schreier.py`. Integral rewriting gives the primitive quotient matrix

$$T_0 = \begin{pmatrix} -1 & -1 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & -1 & -1 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}. \tag{E.26.6}$$

The nine cusp columns have Smith factors $1^8, 0$, and T_0 has Smith factors $1, 1$. Thus (E.26.6) is the actual integral filling map with no finite-index correction. Conjugation by either x or y induces on this quotient the same matrix

$$R = \begin{pmatrix} -1 & -1 \\ 1 & 0 \end{pmatrix}, \qquad R^2 + R + 1 = 0. \tag{E.26.7}$$

These identities are checked by rewriting each of the ten conjugated basis words over $\mathbb{Z}$. They hold after every finite coefficient reduction and hence on $\widehat{T}$. Conjugation by s interchanges x, y, while conjugation by r sends $x \mapsto y$, $y \mapsto (xy)^{-1}$. For precision, the literal identities follow from

$$r(r^{-1}sr^{-1}s)r^{-1} = sr^{-1}sr^{-1}, \qquad r(sr^{-1}sr^{-1})r^{-1} = rsr^{-1}sr = (xy)^{-1}.$$

For the simultaneously inverted generators $x_+ = x^{-1} = (sr)^2$, $y_+ = y^{-1}$, the second formula is $ry_+r^{-1} = x_+^{-1}(x_+y_+)^{-1}x_+$. These are the actual changes of convention. The two conjugations preserve the cubic-cover subgroup and permute its cusp classes. This proves the asserted normality of J_m.

The reduction map $Q \to \mathrm{PSL}_2(\mathbb{F}_3) \simeq A_4 \to A_4/V_4$ is a cyclic-three character. Its restrictions to x and y coincide and are nontrivial: the two matrices are conjugate order-three transvections, whose projective classes lie in the same nontrivial coset. Hence, for every $n \in N_B$,

$$\exp_x(n) + \exp_y(n) = 0 \pmod 3. \tag{E.26.8}$$

By (E.26.7), N_B acts trivially on T. Its image in F/H lies in the cyclic line $\{(a, -a) : a \in \mathbb{Z}/3\}$. The image of $N_B \cap H$ in B/J_m is therefore central in the image of N_B, and the quotient by it is cyclic. A group with a central subgroup and cyclic quotient is abelian: one lift of a cyclic generator and the central subgroup generate it. It follows that $N_B' \subset J_m$. Moreover $n^3 \in H$ for $n \in N_B$, so $n^{3m} \in J_m$. These two inclusions prove exactly (E.26.2). All its targets are finite, so the same maps extend uniquely to $\widehat{B}$. Their kernels contain the closure of $[N_B, N_B]$, and taking the inverse limit over divisibility of m proves (E.26.3). For the simultaneous comparison with G_6, use $G_{\mathrm{lcm}(6,3m)}$.

For the last assertion, both n_1, n_2 are literal basis members of the kernel of $B \to Q$. Rewriting their commutator in the ten-generator cubic-cover basis gives the integral vector $(0, 0, 2, 0, 0, 0, 0, 0, 0, -2)$. Its image under T_0 is zero but its reduction modulo four is nonzero. Thus N_B' is not contained in $[H, H]H^4$, nor in the kernel in (E.26.5), whereas every G_j kills N_B'. A factorization through any specified G_j would kill this particular word, proving the claimed failure of factorization. The program `bc_cm_fermat_factorization.py` retains the integral deck matrices and all 4,656 basis commutator images. Each has zero proper image, and the displayed boundary witness is its first nonzero reduction modulo four. It also checks that translation by $(1, -1)$ has $\mathrm{rank}(A - 1) = 6$ on the full rank-ten module. Its output, JSON and exact arrays are saved alongside it. □

Statement E.27 (Proved). The generating-pair S_3 Ferrari quotient and its exact source factor. *Let $\mathcal{P}$ be the set of all eighteen ordered generating pairs of S_3, and let*

$$K = \operatorname{im}(\langle \xi_1, \xi_2 \rangle \longrightarrow S_3^{\mathcal{P}}), \qquad \xi_1 \longmapsto (g)_{(g,h)}, \quad \xi_2 \longmapsto (h)_{(g,h)}. \tag{E.27.1}$$

The literal Ferrari Nielsen automorphisms in E.17 permute $\mathcal{P}$. Their image $A \leq \operatorname{Aut}(K)$ has order 162, and the resulting four-strand quotient $T = K \rtimes A$ has order 17 496. The free image has order 108 and

$$[K, K] = \gamma_3(K) \simeq (\mathbb{Z}/3)^3. \tag{E.27.2}$$

In particular K is nonnilpotent.

For the six pure evaluation pairs of E.17, both entries have order three and the two entries commute. Consequently their evaluations kill F'. Their common induced six-coset source map is exactly a faithful representation of

$$B/(F'F^3), \qquad |B/(F'F^3)| = 54, \tag{E.27.3}$$

where $F \lhd B = C_2 * C_3$ is the free subgroup of index six used throughout Part E. Since $K_6 = N'N^6 \subset F'F^3$, its reduction from G_6 is the specified quotient homomorphism

$$B \longrightarrow G_6 \longrightarrow B/(F'F^3). \tag{E.27.4}$$

Every $f \in \widehat{F'}$ has six identity values in T. For every profinite unit multiplier, the resulting four-strand braid map is surjective and preserves the two full twists with that multiplier. In particular all twenty-four retained G_6 points of E.17, and every lift of those points used in E.24, pass this finite test.

Proof. Write permutations as functions and use their ordinary composition for multiplication in S_3. Direct breadth-first enumeration gives the eighteen generating pairs and the order-108 subgroup in (E.27.1). The two substitutions on the pair set are

$$p_a(g, h) = (g, hg), \qquad p_b(g, h) = (gh^{-1}, h). \tag{E.27.5}$$

For an evaluation vector k, the automorphism corresponding to a has value $k_{p_a(g,h)}$ at (g, h), and similarly for b. Thus composition of two such automorphisms is represented by $p_{\alpha\beta} = p_\beta \circ p_\alpha$. The program enumerates all 162 permutations in the generated image, checks their multiplication and action on every element of K, and checks faithfulness by their distinct pairs of images of ξ_1, ξ_2. It then uses the actual semidirect product multiplication, with

$$a = (1, p_a), \qquad b = (1, p_b), \qquad c = (\xi_1, p_a).$$

The three braid relations and the literal Ferrari words for both free generators are verified. Enumerating the subgroup generated by a, b, c yields every element of $K \rtimes A$, proving its stated order and surjectivity from B_4.

The derived subgroup of K is computed from all pairwise commutators in the completely enumerated group. It has 27 elements, is abelian of exponent three, and taking all its commutators with K produces the same subgroup. This proves (E.27.2). In particular the quotient (E.27.1) cannot factor through any nilpotent quotient of the Ferrari free group: the image of a nilpotent group is nilpotent, whereas its displayed nonzero lower central subgroup persists at every further step.

The six pairs are formed from the actual pure Artin words, with no linearization. For each pair (X_i, Y_i), the program checks $X_i^3 = Y_i^3 = [X_i, Y_i] = 1$. Therefore the homomorphism

$\theta_i : F \to T$ factors through $F/(F'F^3)$. Induce their product from F to B using the same six coset representatives b_q as E.19:

$$b \longmapsto \left(\overline{b}, \left(\theta_i(b_q b\, b_{q\overline{b}}^{-1})\right)_{q,i}\right). \tag{E.27.6}$$

Here the base factors are multiplied with the explicit Schreier permutation action, so (E.27.6) is an actual homomorphism. The subgroup $F'F^3$ is characteristic in F, hence normal in B, and is killed in every induced coordinate. Thus (E.27.6) factors through the order-54 group $B/(F'F^3)$. A separate enumeration of its two generating images has exactly 54 elements, proving faithfulness of this factor. As an independent check of the required G_6 comparison, all 4656 commutators of the 97 prescribed N generators and all 97 sixth powers are evaluated to the identity in all thirty-six coordinates. Their complete normal generating property proves (E.27.4). The image of N in this factor has order three.

Since all six maps kill F', their extension to $\widehat{F}'$ also has identity value. The pentagon therefore holds identically. The braid generators of T have order six, so a unit multiplier has only the residues $1, 5$ in this target. Direct complete enumeration verifies that $(a^\lambda, b^\lambda, c^\lambda)$ generates T for both residues; the three braid relations and the images of $(ab)^3$ and $(abc)^4$ are checked exactly. Both full twists have order three and their images are their λth powers.

Finally, abelianizing K modulo two and taking the induced linear action gives the affine action on $\mathbb{F}_2^2$. In the ordered list $(0,1),(1,1),(1,0),(0,0)$, the images of a, b, c are respectively $(12), (23), (34)$. This is the standard permutation map of B_4. Hence $\ker(B_4 \to T)$ is already contained in PB_4, $|B_4/\ker(B_4 \to T)| = 17\,496$, and the image of PB_4 has order 729.

The certificates are `de_ferrari_s3_model.py` and `de_ferrari_s3_factor.py`, with their actual outputs, full multiplication/action tables, and induced source matrices. They also evaluate each retained literal commutator word directly. The assertion concerns this actual target and its source factor; joint four-strand generation with the other finite targets has not been inferred from separate component generation. □

Statement E.28 (Proved). The complete two-adic boundary tower and its joint Ferrari lifts. *Keep the discrete groups and the ten-generator integral basis of E.26. Put*

$$M_H = H^{\mathrm{ab}}, \qquad B_n = B/[H,H]H^{2^n}, \qquad C_n = B/(K_6 \cap [H,H]H^{2^n}) \quad (n \geq 1). \tag{E.28.1}$$

Every one of the twenty-four retained G_6 words f_i admits an explicit correction in N'_B giving compatible multiplier-one automorphisms of every C_n. One integral correcting word works at all these levels. The complete set of correction images in $M_H/2^n$ coming from N'_B and satisfying both hexagons has exactly 2^n elements; every one is generating. Its inverse limit is an affine copy of $\mathbb{Z}_2$.

Let $\mathscr{D} = D_{2,6} \times_{G_6} D_6$ be the exact source quotient of E.24. Each of the eighteen multiplier-one points in each of E.24's twenty-four indexed CM families extends through the entire common tower

$$\mathscr{C}_n = \mathrm{im}\big(B \longrightarrow \mathscr{D} \times B_n\big). \tag{E.28.2}$$

Thus all 432 multiplier-one points of E.24 are retained simultaneously with this tower. The same single profinite commutator word gives these compatible lifts and leaves unchanged all six binary and all six ternary Ferrari evaluations, their induced conjugate coordinates, both pentagons, and the generating four-strand maps of E.24.

This tower retains commutators at every relative lower-central depth. If P_n is the image of N_B in B_n, then there is an explicit primitive rank-six lattice $\mathcal{D} \subset M_H$ such that

$$\gamma_r(P_n) = \mathcal{D}/2^n\mathcal{D} \simeq (\mathbb{Z}/2^n)^6 \qquad (r \geq 2). \tag{E.28.3}$$

Consequently no quotient of B with nilpotent image of N_B dominates B_n through the specified maps. The assertion is about the displayed common towers and finite Ferrari tests; it makes no full-GT realization assertion.

Proof. Index the rank-97 basis $n_1, \dots, n_{97}$ as in E.26, and take

$$(w_1, \dots, w_6) = ([n_2, n_3], [n_2, n_4], [n_2, n_5], [n_2, n_7], [n_2, n_9], [n_2, n_{11}]). \tag{E.28.4}$$

Their integral H-abelianizations are the columns of

$$D = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & -1 \\ -2 & 0 & 0 & 0 & 1 & 0 \\ 1 & 1 & 1 & -1 & -1 & 0 \\ 1 & 1 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 & -1 \\ -3 & -1 & -1 & 0 & 1 & 1 \\ 1 & -1 & 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 2 & 0 & 0 & 1 & 0 & -1 \end{pmatrix}. \tag{E.28.5}$$

The first six rows have determinant one. All $4,656$ basis commutators have integral images in $D\mathbb{Z}^6$. Conjugation by N_B on M_H factors through its cyclic-three deck image, which is generated by the image of $t = n_2^{-1}$. On these six columns its matrix is

$$Q_D = \begin{pmatrix} 0 & 0 & 0 & 0 & -1 & 1 \\ 0 & 0 & 1 & -1 & -1 & 1 \\ 0 & -2 & -2 & 1 & 1 & 0 \\ -1 & -1 & -1 & 0 & 1 & 0 \\ 1 & -1 & 0 & -1 & -2 & 2 \\ 0 & -1 & 0 & -1 & -1 & 1 \end{pmatrix}. \tag{E.28.6}$$

The integral identities are $A_t D = DQ_D$ and $Q_D^2 + Q_D + 1 = 0$. Thus $D\mathbb{Z}^6$ is stable under every N_B conjugation. The normal closure of all the basis commutators is N_B', so these calculations prove the exact image equality $\mathrm{im}(N_B' \to M_H) = \mathcal{D} = D\mathbb{Z}^6$. The unit minor proves that reduction of $\mathcal{D}$ modulo every integer embeds in the corresponding reduction of M_H.

Let S, R be the integral matrices obtained by the literal substitutions $s : (x, y) \mapsto (y, x)$ and $r : (x, y) \mapsto (y, (xy)^{-1})$ of E.26, and set

$$\mathcal{E} = \begin{pmatrix} 1 + \mathsf{S} \\ 1 + \mathsf{R} + \mathsf{R}^2 \end{pmatrix}, \qquad \mathcal{M} = \mathcal{E} D. \tag{E.28.7}$$

For $f \in H$, the formulas $s \mapsto sf$, $r \mapsto f^{-1} r$ satisfy the two order relations in $B/[H, H]H^m$ exactly when $\mathcal{E}[f] = 0$ modulo m. This follows by multiplying $(sf)^2$ and $(f^{-1}r)^3$ in the abelian normal kernel; their coordinates are respectively $(1 + \mathsf{S})[f]$ and $-(1 + \mathsf{R} + \mathsf{R}^2)[f]$. Their product fixes sr, so this is the same multiplier-one cusp gauge as E.1.

The integral Smith factors and kernel are

$$\mathrm{Smith}(\mathcal{M}) = (1, 1, 1, 1, 3, 0), \qquad \ker_{\mathbb{Z}} \mathcal{M} = \mathbb{Z} v, \quad v = (1, -2, 1, 1, 1, 2)^t. \tag{E.28.8}$$

For every $z_i = [f_i]$ the certificate supplies an integral $a_i \in \mathbb{Z}^6$ satisfying $\mathcal{M} a_i = -\mathcal{E} z_i$. The literal word $f_i \prod_j w_j^{(a_i)_j}$ consequently satisfies both relations already in $B/[H, H]$. See

`bc_cm_fermat_boundary_integral.json` for all integer exponents and resulting words. The Smith calculation proves that modulo 2^n all solutions are exactly $a_i + (\mathbb{Z}/2^n)v$.

Here is the actual action and the generation check. Let A_q be the deck action on M_H for $q \in (\mathbb{Z}/3)^2$. For the jth Schreier word η_j, follow its successive deck vertices $q \to q'$ and put

$$\mathcal{T}_j = \sum_{\text{letters } y^{\pm 1} \text{ in } \eta_j} (A_{q'} - A_q). \qquad \mathcal{A}(z) = 1 + (\mathcal{T}_1 z \ \cdots \ \mathcal{T}_{10} z). \tag{E.28.9}$$

Once the two order relations hold, the induced substitutions on F are $x \mapsto x$ and $y \mapsto f^{-1}yf$. Inserting the two copies of f at each occurrence of $y^{\pm 1}$ gives exactly (E.28.9), including the negative letters. The certificate independently substitutes the literal corrected words into every η_j, obtaining the same integral matrix. It also checks $\mathcal{T}_j D v = 0$ for every j. Thus the matrix is constant on each complete two-adic correction fibre. Its determinants, in the saved order, are

$$\begin{gathered} 1,\ 3241,\ 9937,\ 8929,\ 22681,\ 26641,\ 26353,\ 17353, \\ 17497,\ 13969,\ 25057,\ 30529,\ 447337,\ 249697,\ 408601,\ 381889, \\ 12241,\ 4249,\ 6697,\ 29017,\ 8209,\ 3097,\ 16417,\ 5977. \end{gathered} \tag{E.28.10}$$

All are 1 modulo six. In particular the matrices are invertible modulo every 2^n and the action on B/H is the identity. The induced map on B_n is an automorphism. Its G_6 map is unchanged because each $w_j \in N'_B$. The resulting endomorphism of the actual common quotient C_n is injective: an element in its kernel projects to the kernel of both component automorphisms, hence is identity in their common image. Finiteness gives surjectivity. This also proves compatibility, using the same word at every level.

For precision at other moduli, the same restricted correction problem has a generating solution modulo m exactly when m is relatively prime to the corresponding determinant in (E.28.10). Sufficiency uses the displayed integral word. For necessity, a prime divisor of that determinant is different from $2, 3$; over its residue field (E.28.8) makes every correction differ by a multiple of v, which leaves (E.28.9) unchanged and singular. This criterion concerns the fixed word f_i and N'_B corrections. It is not an exclusion for a whole CM matrix fibre.

We now give the exact additional comparison required for the joint Ferrari assertion. The map $M_H \to F^{\mathrm{ab}}$ is $3L$, where

$$L = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \end{pmatrix}. \tag{E.28.11}$$

An integral basis matrix Z for $\ker L$ has columns

$$e_1, e_2, e_4, e_5, e_6 - e_3, e_8 - e_7, e_9 - e_7, e_{10} - e_3. \tag{E.28.12}$$

The coordinates in this basis are the rows $1, 2, 4, 5, 6, 8, 9, 10$ of a vector in $\ker L$. The following exact matrix identity supplies the correction, without discarding the quotient by $\mathcal{D}$:

$$\mathcal{M}P = \mathcal{E}Z, \qquad 3P = \begin{pmatrix} 1 & 0 & 1 & 1 & 0 & -1 & 0 & 0 \\ -5 & -3 & 1 & 4 & 0 & -4 & 0 & -3 \\ 2 & 5 & -4 & -6 & -3 & 3 & 2 & 3 \\ 2 & 2 & -1 & -3 & 0 & 3 & 2 & 3 \\ 0 & 1 & 0 & 2 & 0 & -2 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}. \tag{E.28.13}$$

In particular P is integral over $\mathbb{Z}_2$. For every actual $f \in F'$ write $[f] = Zu$. Then $[f] - DPu$ satisfies both boundary norms at every two-power level. This proves the needed compatibility for all the Ferrari words, including their sixth-power corrections.

Generation is retained by the following literal comparison. E.24's origins are the E.17 replacements in the indexed CM families; they are not identified with the original G_6 words f_i. Reconstruct all 432 E.24 words by the actual formula (E.24.7), using `de_ferrari_two_family_words.py`. For every resulting word f in row i, integral Schreier rewriting verifies

$$[f] - z_i \in \mathcal{D} + 2M_H. \tag{E.28.13a}$$

The explicit coordinates in the six columns (E.28.5) are retained for every word in `bc_twoadic_ferrari_comparison.json`. This is a congruence in the displayed boundary module, with no equality of the two G_6 points assumed. After the norm correction, the modulo-two image is therefore one of the enumerated solutions in $z_i + \mathcal{D}/2$. Every such solution has odd determinant. The comparison program also checks the rank-ten action directly for each of the 432 corrected coordinates. Hence the resulting action is invertible at every two-power level.

Within a fixed E.24 source fibre, an element of K_6 automatically has boundary image in $\mathcal{D} + 2M_H$: its derived part has image $\mathcal{D}$, while $n^3 \in H$ and $[n^6] = 2[n^3]$ for every $n \in N_B$. This observation is consistent with the literal comparison, but is not used to identify the earlier and later G_6 origins.

It remains to realize this correction while keeping the entire Ferrari source fixed. In the complete induced binary source, the images of $\gamma_2(N_B), \gamma_3(N_B)$ have ranks $6, 0$. This is checked by the actual conjugations of all 97 generators on the full normal commutator image in `de_ferrari_two_lowercentral.py`. The ternary ranks of E.23 are $15, 6, 3, 0$ in degrees $2, 3, 4, 5$. The G_6 image is abelian. Consequently the actual map $B \to \mathscr{D}$ kills $\gamma_5(N_B)$, in every induced coordinate. On the boundary side,

$$\mathrm{Smith}(Q_D - 1) = (1, 1, 1, 3, 3, 3). \tag{E.28.14}$$

For the six explicit words

$$w_j^{(5)} = [t, [t, [t, w_j]]] \in \gamma_5(N_B) \tag{E.28.15}$$

their boundary columns are exactly $D(Q_D - 1)^3$; the coordinate determinant is $3^9 = 19683$. This is invertible over $\mathbb{Z}_2$. Hence the actual restriction of the boundary homomorphism to $\overline{\gamma_5(N_B)}$ surjects onto $\mathcal{D} \otimes \mathbb{Z}_2$. Explicit preimages are products of the six words (E.28.15) with continuous powers: embed each two-adic exponent in $\widehat{\mathbb{Z}}$ with all other prime coordinates zero. Their images in $\mathscr{D}$ stay identity. This gives one profinite $k \in \overline{\gamma_5(N_B)} \subset \widehat{F}'$ with boundary coordinate $-DPu$, and fk is the required single commutator word for the complete tower.

We prove joint generation directly from the B_n automorphism and E.24's surjective shadow $B \to \mathscr{D}$. Fix n, write $W = \mathscr{C}_n$ and $W_N = \mathrm{im}(N_B \to W)$, and denote the corrected homomorphism by $\psi : B \to W$. The elements $\psi(s), \psi(r)$ belong to this actual common image, since they are evaluated from the same profinite word; the two order relations give the homomorphism. Its map to $Q = B/N_B$ is the retained G_6 automorphism, so $\psi(N_B) \subset W_N$. Its $\mathscr{D}$ projection is all $N_{\mathscr{D}} = \mathrm{im}(N_B \to \mathscr{D})$: E.24 is surjective, and an element mapping into $N_{\mathscr{D}}$ has a preimage in N_B by this Q comparison. Its boundary projection is $\mathcal{A}_\psi(P_n) \subset P_n$, where $\mathcal{A}_\psi$ denotes the proved automorphism of B_n. The two subgroups have the same finite order, so this projection is all P_n.

The group $N_{\mathscr{D}}$ has class at most four. Surjectivity onto P_n and (E.28.3) therefore give the exact equality

$$\gamma_5\big(\psi(N_B)\big) = \{1\} \times \mathcal{D}/2^n\mathcal{D} \ \subset W_N. \tag{E.28.16}$$

Indeed its first projection is identity, and its second projection is all of $\gamma_5(P_n)$; a subgroup with trivial first coordinate is determined by its second coordinate. Quotienting by this subgroup, W_N embeds in $N_{\mathscr{D}} \times (P_n/P_n')$. It is consequently a finite nilpotent group of primes two and

three. It is generated by the 97 images of the chosen N_B generators and surjects onto $N_B^{\mathrm{ab}}/6$ through the $\mathscr{D} \to G_6$ map. For each prime $p = 2, 3$, its Sylow Frattini quotient thus has dimension at most 97 and at least 97; the displayed projection identifies it with N_B^{ab}/p. The retained G_6 automorphism makes the image of $\psi(N_B)$ surjective on both Frattini quotients. The finite nilpotent Frattini criterion then makes it all of the quotient of W_N. Together with (E.28.16) this proves $\psi(N_B) = W_N$. Its surjective Q map finally gives $\psi(B) = W$.

Thus the constructed maps are compatible generating shadows on the actual common quotients. The definition of $\mathscr{D}$ retains the six evaluations and all their conjugate coordinates, so the correction leaves the complete four-strand equations and generating maps exactly as in E.24. This proves all the joint assertions without requiring invariance of the fixed Ferrari source kernel.

Finally, the exact derived image is $\mathcal{D}/2^n$ and commutation with t induces $Q_D - 1$ on it. Formula (E.28.14) makes this map onto. Induction therefore gives (E.28.3) as an equality of images of the actual lower-central subgroups. A factorization through a quotient with nilpotent N_B image would kill one of these subgroups while its displayed image has order 2^{6n}, a contradiction. All integral identities, literal source words, Smith forms and substitution matrices are recorded by `bc_cm_fermat_boundary.py` and `bc_cm_fermat_boundary_integral.py`, with their executed outputs and exact JSON/array certificates. □

Statement E.29 (Proved). All retained fibres survive the complete A_5 Ferrari test. *Let D_{23} be the actual joint three-strand source quotient of E.24, and let $T_{23} = T_{2,6} \times T_{3,6}$ be its actual four-strand target. We retain all* 432 *multiplier-one points of E.24, including both binary corrections and every point of each ternary plane.*

There is an explicit finite Ferrari quotient $\pi_A : B_4 \twoheadrightarrow T_A$ whose free Ferrari kernel is A_5^{19}. Its associated six-evaluation, six-coset induced homomorphism is $\rho_A : B = C_2 * C_3 \twoheadrightarrow G_A$. Every one of the 432 *points has a correction in N' that preserves its complete D_{23} image and satisfies both hexagons and the full pentagon in the common refinement with G_A and T_A. The resulting three-strand map and the map onto the actual joint four-strand image are surjective. In particular, all* 24 *retained CM fibres, including all* 12 *extra fibres, survive this nonsolvable finite test.*

Proof. We give the target, the complete correction image, the arithmetic comparison used to select a point in that image, and the joint-generation argument separately.

The exact target and induced source. Enumerate the 2280 ordered generating pairs of A_5. Two such pairs give the same epimorphism kernel of the free group $\langle \xi_1, \xi_2 \rangle$ precisely when an automorphism of A_5 relates them. Here $\mathrm{Aut}(A_5) = S_5$, with its conjugation action. For completeness, an automorphism permutes the five Sylow 2-subgroups, each isomorphic to C_2^2. Its action on this set is faithful: an automorphism fixing the five subgroups gives the same permutation of them for each g and its image, and the conjugation action of A_5 on these five subgroups is faithful. Thus the automorphism group has order at most 120, while the 120 conjugations by S_5 are distinct. This proves the equality used in the enumeration.

There are 19 epimorphism kernels. Choose the lexicographic representative (g_j, h_j) of each orbit and retain all 19 evaluations. The Nielsen substitutions

$$a : (g, h) \longmapsto (g, hg), \qquad b : (g, h) \longmapsto (gh^{-1}, h) \tag{E.29.1}$$

permute these representatives, with a uniquely recorded conjugating element of S_5 in each factor. The action on the kernels has two orbits, of sizes 9 and 10. The free evaluation image is A_5^{19}: a subdirect product of simple groups can identify two factors only through an

isomorphism between the two corresponding epimorphism kernels, which have already been separated. The program also computes the order 60^{19} independently in the resulting faithful permutation representation.

An element of the Ferrari semidirect product is recorded as (p, L, R), where p permutes the 19 factors, $R_j = c_j$ is the Nielsen conjugator, and $L_j = k_j c_j$ includes the free-kernel coordinate. The multiplication used is

$$(p, L, R)(q, L', R') = \big(q \circ p, (L_j L'_{p(j)})_j, (R_j R'_{p(j)})_j\big). \tag{E.29.2}$$

Two five-point blocks in each factor, acted on by L_j and R_j, give a faithful permutation representation of degree 190. The literal images of the three braid generators are $(1, a), (1, b), (\xi_1, a)$ in the Ferrari description. The program checks every braid relation, the literal words for ξ_1, ξ_2, and all four Nielsen conjugation identities. The resulting orders are

$$\begin{aligned} |T_A| &= 2^{60}3^{35}5^{31}7^2, \\ |\langle a, b\rangle| &= 2^{22}3^{16}5^{12}7^2. \end{aligned} \tag{E.29.3}$$

All six evaluation pairs from E.17 are noncommuting in this target.

For each of those six actual maps $\theta_i : F \to T_A$, induce through the six cosets of $F \lhd B$. Explicitly, for a section $q \mapsto b_q$ of B/F, the component at (q, i) of $\rho_A(w)$ is

$$\theta_i(b_q w b_{qw}^{-1}), \tag{E.29.4}$$

together with the actual permutation $q \mapsto qw$. This defines a homomorphism of discrete groups, extending continuously to their profinite completions. Its kernel is normal in B and lies in F. On F the coordinate at the identity coset is exactly θ_i; thus every comparison with the six substitutions is a specified coordinate homomorphism from $\rho_A(F)$.

The faithful induced permutation action has degree 6840. The executed compression to degree 1980 deletes only duplicate transitive B-sets: every deletion has a saved bijective intertwiner for both s and r. Both actions give the identical image order

$$|G_A| = 2^{38}3^{26}5^{21}7^2. \tag{E.29.5}$$

The whole correction image. Write

$$P = \rho_A(N), \quad C = \rho_A(N'), \quad K = \rho_A(K_6), \quad E = \rho_A(F'), \qquad K_6 = N'N^6.$$

The exact permutation-group computations give

group	order or quotient
P	$2^{35}3^{24}5^{21}7^2$
C	$2^{35}3^{23}5^{19}7^2$
K	$2^{35}3^{23}5^{21}7^2$
P/C	75
K/C	25
G_A/P	72
G_A/K	216
E/C	4

(E.29.6)

Moreover,

$$C' = C, \qquad \rho_A(K_6 \cap F') = C. \tag{E.29.7}$$

These are complete subgroup calculations. The 97 literal Schreier generators of N give P; six of them generate the computed image. The derived subgroup of that group gives C. To

compute $K_6 \cap F'$ we retain all 95 products $\prod_i n_i^{6v_i}$, with v running through the primitive integral kernel of $N^{\mathrm{ab}} \to F^{\mathrm{ab}}$ certified in E.19. Together with N' these normally generate $K_6 \cap F'$. All 95 image products lie in C. The full 97 sixth powers instead enlarge C by a factor 25, giving K. Stability under the actual images of both s and r is checked. A separate derived-subgroup computation on a reduced generating set proves $C' = C$.

The modular comparison is also explicit. Map

$$s \longmapsto \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \qquad r \longmapsto \begin{bmatrix} 0 & -1 \\ 1 & -1 \end{bmatrix} \quad \text{in } \mathrm{PSL}_2(\mathbb{Z}/6). \tag{E.29.8}$$

Enumerate its 72 elements and all 144 Schreier kernel generators for these two generators. Every such word has its ρ_A image in P. The two preimages have index 72, so they coincide. This proves, with the displayed maps,

$$G_A/P \simeq \mathrm{PSL}_2(\mathbb{Z}/6) \simeq S_3 \times A_4. \tag{E.29.9}$$

Reducing (E.29.8) modulo 3 defines $q_3 : G_A \to A_4$. For $u = r^{-1}s$, $v = sr^{-1}$ and $x = u^2, y = v^2$, the two matrices are exactly

$$x \longmapsto \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}, \qquad y \longmapsto \begin{bmatrix} 1 & 0 \\ -2 & 1 \end{bmatrix}. \tag{E.29.10}$$

Their generated image is A_4, so $q_3(E) = A_4' = V_4$. Since $C \subset P \subset \ker q_3$ and $|E/C| = 4$,

$$\ker(q_3|_E) = C. \tag{E.29.11}$$

Actual arithmetic points in the three required cosets. Nakamura–Schneps, Theorem 2.5 [NS00], gives for an actual $\sigma \in G_{\mathbb{Q}}$ with $\chi(\sigma) = 1$

$$q_3(f_\sigma) = M_\kappa, \qquad M_\kappa = \begin{bmatrix} 1 & -8\kappa \\ -8\kappa & 1 + 64\kappa^2 \end{bmatrix} \pmod 3, \quad \kappa = \rho_2(\sigma) \pmod 3. \tag{E.29.12}$$

The theorem's input matrices are exactly (E.29.10), and its convention is $x \mapsto x^\chi$, $y \mapsto f_\sigma^{-1} y^\chi f_\sigma$. We use this theorem only for actual Galois elements.

All three values $\kappa = 0, 1, 2$ occur with $\chi = 1$ as an equality in $\widehat{\mathbb{Z}}^\times$. Indeed, the splitting field $L = \mathbb{Q}(\zeta_3, \sqrt[3]{2})$ has Galois group S_3 and maximal abelian subextension $\mathbb{Q}(\zeta_3)$. Since every finite subextension of $\mathbb{Q}(\mu_\infty)/\mathbb{Q}$ is abelian,

$$L \cap \mathbb{Q}(\mu_\infty) = \mathbb{Q}(\zeta_3).$$

Consequently $G_{\mathbb{Q}(\mu_\infty)}$ maps onto $\mathrm{Gal}(L/\mathbb{Q}(\zeta_3)) = C_3$, and this restriction is precisely the Kummer coordinate ρ_2 mod 3 used in (E.29.12).

Direct substitution of the 24 retained E.17 words into (E.29.10) gives the following complete index table:

κ	indices
0	0, 1, 2, 3, 16, 17, 18, 19
1	4, 5, 6, 7, 20, 21, 22, 23
2	8, 9, 10, 11, 12, 13, 14, 15

(E.29.13)

Every subsequent E.24 correction lies in K_6 and therefore has trivial q_3 image. Thus the table applies to all 432 multiplier-one points, without discarding any of their parameters.

Fix one of those literal words $f_0 \in F'$ and choose an actual σ with the matching value in (E.29.13). By (E.29.11),

$$c = \rho_A(f_0)^{-1} \rho_A(f_\sigma) \in C. \tag{E.29.14}$$

Let S be the image of N' in D_{23}. It is solvable, because D_{23} is an extension of the solvable group Q by the explicitly constructed nilpotent 2- and 3-groups. The joint image of N' in $C \times S$ is the full product: Goursat's lemma would otherwise give a nontrivial common quotient, which would be both perfect and solvable. Hence there is a discrete $k \in N'$ with images $(c, 1)$. For $f = f_0 k$, we now have the exact equalities

$$f \bmod D_{23} = f_0 \bmod D_{23}, \qquad \rho_A(f) = \rho_A(f_\sigma), \qquad f \in F'. \tag{E.29.15}$$

The same argument applies to the already constructed solvable boundary refinements: at every finite level the joint image is a product, and compactness supplies one compatible correction in $\overline{N'}$ for their inverse limit. This preserves their component maps and equations. The simultaneous source-generation assertion here is (E.29.16), for D_{23}; the corresponding four-strand assertion is (E.29.18). We make no further joint-generation assertion for a boundary refinement whose order admits an additional common quotient of G_A/C. This preservation argument also makes no nonemptiness assertion over arbitrary profinite quotients.

The equality in G_A matches all 36 induced coordinates, so it matches the six actual T_A values of f with those of f_σ. In particular, the full pentagon, the braid relations, and the full-twist identities in T_A follow from the actual Galois automorphism. The source hexagons follow by the same equality in G_A. All these equations remain true in D_{23} and T_{23} by the first equality in (E.29.15).

Surjectivity in the common refinements. Put $H = G_A/K$. The actual B image in $D_{23} \times G_A$ is

$$D_{23} \times_H G_A. \tag{E.29.16}$$

Here H is a quotient of G_6, with its specified map from B, so both factors map onto it. To prove the equality, observe that every solvable quotient of G_A kills the perfect group C. Every quotient of order divisible only by 2 and 3 also kills the normal subgroup K/C of order 25. Thus any common quotient of D_{23} and G_A factors through H. Their actual joint image already has the common H coordinate, so Goursat's lemma proves (E.29.16). The corrected shadow surjects onto D_{23} by E.24, and onto G_A because it has the same generator images as ρ_A composed with an actual Galois automorphism of $\widehat{B}$. Its joint image therefore equals (E.29.16), by the same common-quotient argument.

There is an equally complete four-strand comparison. A separate computation gives

$$[T_A : T_A'] = 2, \qquad T_A'' = T_A'. \tag{E.29.17}$$

All three braid generators have the same nontrivial image in $T_A/T_A' = C_2$. The exact derived series is computed in the faithful 190-point action; the subgroup orders and the generator-ratio memberships are retained in the certificate. Thus every solvable quotient of T_A factors through this parity map. The group T_{23} is solvable by its explicit 2- and 3-group extensions and has the same parity map through its binary S_4 quotient. Therefore the actual joint target is

$$T_{23} \times_{C_2} T_A. \tag{E.29.18}$$

The corrected three braid generators surject onto T_{23} by E.24 and onto T_A because they are the image of the Galois braid automorphism in that factor. They preserve the common parity coordinate. Goursat's lemma and (E.29.17) then prove surjectivity onto (E.29.18). Its kernel in B_4 is normal and lies in PB_4, since the standard S_4 map already factors through T_{23}. The coordinate maps (E.29.4) give the required comparisons for every pentagon substitution.

For the braid-action convention, we use Nakamura–Schneps (1.1), which gives $\sigma_i \mapsto f(y_i, \sigma_i^2)^{-1} \sigma_i^\lambda f(y_i, \sigma_i^2)$. For $i = 3$, our sixth pair uses $\Omega_3 = (ab)^3 = a^2(ba^2b)$ instead of

$y_3 = ba^2b$. The element a^2 commutes with both y_3 and c^2; as $f \in F'$, inserting that commuting factor does not change the evaluation. Hence the exact six-pair action used here is the stated Galois action.

The seven executed certificates have the prefix `de_ferrari_a5_` and suffixes `model.py`, `components.py`, `source.py`, `source_compress.py`, `corrections.py`, `compare.py`, and `solvable.py`. Their outputs and exact permutation arrays are retained. The first new correction is supplied by (E.29.14)–(E.29.15); no uncomputed literal Galois word is claimed as a saved certificate. The finite group and fibre identifications, and every numerical assertion used in that existence proof, were computed and retained. □

Statement E.30 (Proved). A necessary finite conjugacy test from the four-strand periodic involution. *Let $q : B_4 \twoheadrightarrow T$ be any finite quotient of the discrete four-strand braid group, and write*

$$a = q(\sigma_1), \quad b = q(\sigma_2), \quad c = q(\sigma_3), \quad d = abc, \quad z = d^4, \quad C = \langle z \rangle. \tag{E.30.1}$$

The subgroup C is central. For every $F = (\lambda, f) \in \widehat{\mathrm{GT}}$, let

$$a_F = q(F(\sigma_1)), \qquad b_F = q(F(\sigma_2)), \qquad c_F = q(F(\sigma_3)), \tag{E.30.2}$$

where q denotes also its continuous extension and F acts by the usual based Drinfeld braid formulas. Then

$$(a_F b_F c_F)^2 C \sim_{T/C} d^2 C. \tag{E.30.3}$$

Equivalently, the finite equation

$$(a_F b_F c_F)^2 = h d^2 h^{-1} z^j \quad \text{has a solution } h \in T, \quad j \in \mathbb{Z}/|z|. \tag{E.30.4}$$

This condition applies to the actual Ferrari quotient maps used in Part E. It retains the complete conjugacy class in T/C; the full-twist equation alone prescribes only the square of the element on the left of (E.30.3)*. The assertion uses the homomorphism $q \circ F : \widehat{B}_4 \to T$ and requires no invariance of* $\ker q$ *under F.*

Proof. We give the group comparison that brings the torsion theorem into this particular finite target. Capping the boundary of a four-punctured disc by a once-punctured disc gives the exact discrete homomorphism

$$B_4/\langle (\sigma_1\sigma_2\sigma_3)^4 \rangle \xrightarrow{\simeq} \Gamma^{(5)}_{0,[5]} \ \leq \ \Gamma_{0,[5]}, \tag{E.30.5}$$

where the subgroup on the right fixes the fifth marked point and has index five. Indeed a mapping class fixing that point can be represented by a map fixing a small disc about it; restriction to the complementary four-punctured disc gives a braid. Changing the isotopy at the capped disc changes this braid by an integral boundary twist, precisely the central full twist displayed on the left. This identifies both the kernel and the image in (E.30.5).

Completion of the quotient on the left is $\widehat{B}_4/\overline{\langle (\sigma_1\sigma_2\sigma_3)^4 \rangle}$: their finite quotients are exactly the finite braid quotients killing that central element. A finite-index subgroup has the full induced profinite topology, by taking finite-index cores. Thus completion of (E.30.5) identifies its left side with the open fifth-point stabilizer in $\widehat{\Gamma}_{0,[5]}$. For the action comparison, use the specified homomorphisms

$$\widehat{B}_4 \xrightarrow{\iota} \widehat{B}_5 \xrightarrow{\pi} \widehat{\Gamma}_{0,[5]}, \qquad \iota(\sigma_i) = \sigma_i \quad (1 \leq i \leq 3). \tag{E.30.5a}$$

Here π kills the closed normal subgroup generated by

$$z_5 = (\sigma_1\sigma_2\sigma_3\sigma_4)^5, \qquad y_5 = \sigma_4\sigma_3\sigma_2\sigma_1^2\sigma_2\sigma_3\sigma_4. \tag{E.30.5b}$$

These are exactly the central full twist and the sphere Hurwitz relation in the discrete presentation [Col12, Section 3.1]. The composite $\pi\iota$ is the capping map in (E.30.5): it places the fifth puncture outside the four-strand disc and then fills its outer boundary. The exact action convention used here is

$$\begin{aligned} F(\sigma_1) &= \sigma_1^{\lambda}, \\ F(\sigma_i) &= f(\sigma_i^2, y_i)\sigma_i^{\lambda} f(y_i, \sigma_i^2), \\ y_i &= \sigma_{i-1}\cdots\sigma_2\sigma_1^2\sigma_2\cdots\sigma_{i-1}. \end{aligned} \tag{E.30.5c}$$

Nakamura–Schneps, Section 1, formula (1.1) and its following paragraph, give the automorphisms of the completed braid groups and their induced sphere mapping-class actions with precisely this formula [NS00]. The first three generator formulas use the same preceding pure braids in B_4 and B_5, so $F\iota = \iota F$. The stated quotient action sends every $\pi(\sigma_i)$ to the image of the same displayed word, giving $\pi F = F\pi$ on generators and hence on the completed group. This proves equivariance of the actual capping map.

The Ferrari convention uses $\Omega_3 = (\sigma_1\sigma_2)^3$ in its sixth evaluation. The literal identity $\Omega_3 = \sigma_1^2 y_3$ has σ_1^2 commuting with both y_3 and σ_3^2: the first assertion follows from centrality of Ω_3 in B_3, the second from distant braid commutation. Consequently $f(\Omega_3, \sigma_3^2) = f(y_3, \sigma_3^2)$ for $f \in \widehat{F}'$, since the extra central factor occurs with the zero first abelian exponent of f. This identifies the actual sixth evaluation with (E.30.5c).

Let $\delta = \sigma_1\sigma_2\sigma_3$ and let ϵ be the image of δ^2 in (E.30.5). It has order exactly two: its square is the killed full twist, and its permutation on five marked points is a product of two disjoint transpositions with the unique fixed point 5. The element $F(\epsilon)$ still has order two. Collas's action-independent Theorem 3.3 and Corollary 3.2 give exactly one conjugacy class of order-two elements in $\widehat{\Gamma}_{0,[5]}$: it is the square of the displayed order-four periodic element. Hence there is $u \in \widehat{\Gamma}_{0,[5]}$ with

$$F(\epsilon) = u\epsilon^{\lambda}u^{-1} = u\epsilon u^{-1}; \tag{E.30.6}$$

the second equality uses $\lambda \in \widehat{\mathbb{Z}}^{\times}$ and the order two of ϵ [Col12]. Only the torsion classification is used from that paper. The action itself is the explicitly specified full-GT action (E.30.5c).

We now locate the conjugator. The Drinfeld action preserves the permutation of each braid generator: its exponent is a unit, hence odd, and its conjugating word belongs to the commutator subgroup of a pure free subgroup. Consequently $F(\epsilon)$ has the same permutation as ϵ. The permutation of u in (E.30.6) sends the unique fixed point of the latter permutation to the unique fixed point of the former. It fixes 5. Hence u lies in the open subgroup in (E.30.5) itself. We have obtained the required conjugacy inside the completed four-strand central quotient.

The homomorphism $\widehat{B}_4 \to T/C$ kills the closed central subgroup in (E.30.5), so it factors through that quotient. Applying it to (E.30.6) gives (E.30.3). Lifting the conjugator to T gives (E.30.4), since the two sides then differ by an element of C. Every map in this argument is either the specified quotient map, a finite-index inclusion and its completion, or the stated GT action. In particular no endomorphism of the particular finite quotient T has been assumed. For a finite candidate, failure of (E.30.4) excludes that candidate from this full-GT image; a conclusion about a whole CM fibre still requires the complete fibre and multiplier comparisons. □

Statement E.31 (Proved). The complete ternary boundary fibres and the simultaneous full boundary tower. *Retain $H, M_H, L, \mathcal{E}$ from E.26–E.28 and the exact E.24 source $\mathscr{D} = D_{2,6} \times_{G_6} D_6$. Let*

$$R_3 = \ker(B \to D_6) \cap F', \qquad \mathscr{R} = \ker(\widehat{B} \to \mathscr{D}) \cap \widehat{F}'. \tag{E.31.1}$$

The integral image of R_3 in M_H is the lattice $\mathcal{K} = K\mathbb{Z}^8$, where

$$K = \begin{pmatrix} 12 & 0 & 1 & 1 & 0 & 0 & 11 & 0 \\ 0 & 12 & 11 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}. \tag{E.31.2}$$

Its index in $\ker L$ *is* $144 = 2^4 3^2$*, and its embedding in M_H has Smith factors* $(1^6, 12, 12)$*. There are eight explicit words in R_3 mapping to these eight columns. The exact joint-source comparison is*

$$\operatorname{im}(\mathscr{R} \to M_H \otimes \mathbb{Z}_3) = \mathcal{K} \otimes \mathbb{Z}_3. \tag{E.31.3}$$

Every one of E.24's 432 *multiplier-one literal points has compatible generating boundary lifts through all three-power levels. For a fixed point, its complete set of correction images in $M_H/3^n M_H$ coming from $\mathscr{R}$ and satisfying the two boundary norms has cardinality*

$$3^{\,3n-1} \qquad (n \geq 1). \tag{E.31.4}$$

Every finite correction image lifts compatibly. The inverse limit is an affine copy of $\mathbb{Z}_3^3$; these are literal correction images, with no geometric-conjugacy quotient implicitly taken.

All these points also lift simultaneously with E.28 and with every prime-to-six boundary modulus. More precisely, each point is represented by a single $f^* \in \widehat{F}'$ of multiplier one giving compatible surjective shadows on every actual common quotient

$$\mathscr{W}_m = \operatorname{im}(B \longrightarrow \mathscr{D} \times B/[H,H]H^m) \qquad (m \geq 1). \tag{E.31.5}$$

On the prime-to-six part of M_H its boundary coordinate may be chosen zero. All the finite Ferrari coordinates, pentagons and generating braid maps of E.24 are retained exactly. This proves lifting through the full boundary homology tower with the stated four-strand tests.

Proof. We first prove the exact F' intersection used in the kernel calculation. Put $C = N_B^{\mathrm{ab}} \simeq \mathbb{Z}^{97}$ and let $\phi : C \to F^{\mathrm{ab}}$ be the actual integral abelianization map. Since F^{ab} is torsion-free,

$$(K_6 \cap F')/N_B' = 6C \cap \ker\phi = 6\ker\phi. \tag{E.31.6}$$

Indeed $6c \in \ker\phi$ implies $6\phi(c) = 0$, hence $\phi(c) = 0$. Thus $S = K_6 \cap F'$ is normally generated inside N_B by the $4,656$ commutators $[n_i, n_j]$ and the 95 words $\prod_i n_i^{6v_i}$, where v runs through the primitive integral basis of $\ker\phi$ used in E.19. Each is a literal element of F'.

Their image in D_6 is the verified elementary abelian group of dimension eighteen. Use its eighteen actual basis words $g_1, \ldots, g_{18} \in S$ from E.23. The full nineteen-dimensional K_6 image is also elementary abelian, as checked by all its cubes and pairwise commutators. Consequently image coordinates in this calculation add exactly; no omitted cubic or commutator terms are replaced by a logarithmic approximation.

For each of the $4,751$ normal generators of S, append the inverse of the ordered product of g_i's with its computed image coordinates. Add the eighteen relations g_i^3, the 153 commutators $[g_i, g_j]$, and the $1,746$ relations expressing $n_j g_i n_j^{-1}$ in the same image basis. These $6,668$ relators normally generate R_3 inside N_B. To prove completeness, quotient S by their normal

closure. The original generators and all their N_B conjugates reduce to the eighteen g_i's. The latter commute and have exponent three. The quotient therefore has at most 3^{18} elements, and it surjects onto the actual image of that order. This proves equality of the kernel, not merely an inclusion in it. It also proves that every normal relator used lies in F', including the power residuals.

All these relators are in H because $K_6 \subset H$. Their integral boundary coordinates can therefore be computed by addition in M_H. If $h_i = [g_i]$ and a normal generator w has image coordinates $a_i \in \{0, 1, 2\}$, its residual coordinate is

$$[w] - \sum_i a_i h_i. \tag{E.31.7}$$

The cube and commutator coordinates are respectively $3h_i$ and zero. The conjugation residual is $A_n h_i - \sum_j a_j h_j$, with the actual deck matrix A_n of E.26. The program retains every integer vector and every source word needed to reconstruct these relations.

Hermite reduction of this 10-by-6,668 integer matrix gives exactly (E.31.2). Multiplication by the literal S, R matrices preserves the resulting lattice. Hence it contains the images of all B conjugates of the relators. Conversely every column used is the image of an actual relator, and all added conjugates still belong to R_3. This proves $\mathrm{im}(R_3 \to M_H) = \mathcal{K}$. The rows $1, 2, 4, 5, 6, 8, 9, 10$ give its matrix in the primitive basis (E.28.12); its determinant is 144. Integral Smith reduction gives the embedding factors in the statement.

The word-preimage certificate is explicit. In the zero-based order of the preceding normal presentation, the twelve relators with indices

$$2,\ 3,\ 101,\ 103,\ 104,\ 105,\ 106,\ 108,\ 111,\ 4658,\ 4660,\ 4661 \tag{E.31.8}$$

already generate the eight lattice columns by integer combinations. The saved 12-by-8 coefficient matrix has entries of absolute value at most thirty. For each column take the ordered product of these twelve actual relators with those signed exponents. The program `bc_cm_fermat_kernel_words.py` checks the twelve relators in the exact D_6 source model, checks their zero integral F-abelianization, and expands the eight products to verify their integral H-coordinates against (E.31.2). The JSON retains the twelve literal B and F words and the full ordered-power products. This supplies actual word preimages for every lattice generator.

We pass to the joint source. The discrete F' is dense in $\widehat{F}'$: quotienting $\widehat{F}$ by its closure gives precisely $\widehat{F^{\mathrm{ab}}}$. Intersecting this dense subgroup with the open kernel of a finite source quotient remains dense in the corresponding intersection. Compactness of its image therefore identifies the three-adic image of the completed ternary kernel with $\mathcal{K} \otimes \mathbb{Z}_3$. For $r \in R_3$, its binary K_6 image has exponent two by E.24. Thus r^2 belongs to the joint kernel. The boundary image doubles, and multiplication by two is invertible on $\mathcal{K} \otimes \mathbb{Z}_3$. This proves both inclusions in (E.31.3).

Write $\mathcal{K}_3 = \mathcal{K} \otimes \mathbb{Z}_3$. The exact norm map has

$$\mathrm{Smith}(\mathcal{E}K) = (1, 1, 1, 1, 3, 0, 0, 0). \tag{E.31.9}$$

For all 432 reconstructed E.24 words, the program computes $z = [f] \in M_H$ and solves $\mathcal{E}Ka = -\mathcal{E}z$ over $\mathbb{Z}_3$. Every right side satisfies the exact Smith divisibility conditions. The saved solutions actually have integral boundary coordinates. The literal action formula (E.28.9) at every corrected coordinate has rank ten modulo three. Thus each solution supplies a boundary automorphism at every three-power level. The earlier six-column test using only $D(Q_D - 1)^3$ failed for all the words; its Smith factors $(3, 9, 9, 9, 9, 0)$ describe that restricted correction image. The complete kernel computation above supplies the additional actual corrections and settles this test.

We prove that no finite-only boundary correction images remain. The Smith factors of $\mathcal{K}$ inside M_H and of $\mathcal{E}\mathcal{K}$ give

$$\begin{aligned} |(\mathcal{K}_3 + 3^n M_H \otimes \mathbb{Z}_3)/3^n M_H \otimes \mathbb{Z}_3| &= 3^{8n-2}, \\ |(\mathcal{E}\mathcal{K}_3 + 3^n (M_H \otimes \mathbb{Z}_3)^2)/3^n (M_H \otimes \mathbb{Z}_3)^2| &= 3^{5n-1}. \end{aligned} \tag{E.31.10}$$

Their quotient is the finite norm-kernel size 3^{3n-1}. On the other hand, the integral kernel of $\mathcal{E} : \mathcal{K} \to M_H^2$ has rank three, and its embedding in M_H has Smith factors

$$(1, 4, 12). \tag{E.31.11}$$

These are computed from the zero columns of the retained unimodular Smith decomposition in (E.31.9), followed by multiplication by K. Its three-adic reduction therefore also has 3^{3n-1} images. That image is contained in the finite norm kernel, so the cardinality equality proves

$$\begin{aligned} \ker\bigl(\mathcal{K}_3/(\mathcal{K}_3 \cap 3^n M_H \otimes \mathbb{Z}_3) \xrightarrow{\mathcal{E}} (M_H/3^n M_H)^2\bigr) \\ = \operatorname{im}\bigl(\mathcal{K}_3 \cap \ker \mathcal{E} \longrightarrow M_H/3^n M_H\bigr). \end{aligned} \tag{E.31.12}$$

Translating by a saved integral solution proves (E.31.4), compatible lifting of every finite image, and the stated affine $\mathbb{Z}_3^3$ inverse limit.

Here are the single-word and simultaneous comparisons. Let $r_1, \dots, r_8 \in R_3$ be the word preimages of (E.31.2), and let $e_p \in \widehat{\mathbb{Z}}$ have component one at p and zero at all other primes. Given a ternary correction Ka, take

$$k_3 = \prod_{j=1}^{8} (r_j^2)^{\, e_3 a_j/2}. \tag{E.31.13}$$

The exponent $a_j/2$ is in $\mathbb{Z}_3$. Every factor is in the joint-source kernel and in $\widehat{F}'$. Its boundary coordinate is the prescribed one at three and zero at every other prime. Formula (E.28.15) supplies the two-adic correction k_2 with exponents supported only at two. Hence $f k_2 k_3$ retains the source point and satisfies both norms at all two- and three-power levels simultaneously. Boundary coordinates add because all correction words are in $\widehat{H}$.

For the remaining primes, put $e_{>3} = 1 - e_2 - e_3$ and $A_{>3} = \prod_{\ell \neq 2,3} \mathbb{Z}_\ell$. The index 144 of $\mathcal{K}$ in $\ker L$ is a unit in this coefficient ring. Thus $\mathcal{K} \otimes A_{>3} = \ker(L \otimes A_{>3})$. Since $f \in \widehat{F}'$, its coordinate belongs to this kernel. Solve $2Kc = -[f]$ over $A_{>3}$ using the explicit matrix in the basis (E.28.12), and set

$$k_{>3} = \prod_{j=1}^{8} (r_j^2)^{\, e_{>3} c_j}. \tag{E.31.14}$$

These are well-defined continuous powers in the same joint-source kernel. They have zero two- and three-adic boundary coordinates and cancel $[f]$ at every other prime. The single word $f^* = f k_2 k_3 k_{>3} \in \widehat{F}'$ therefore has all the asserted boundary coordinates and unchanged Ferrari evaluations.

We finish by proving generation in the actual common finite images. First take modulus $2^a 3^b$, with $a, b \geq 1$, and let W_N be the original N_B image in the common source. For the new shadow ψ, its Q map is the retained automorphism, so $\psi(N_B) \subset W_N$. Its projection to the binary boundary P_{2^a} is onto as in E.28. Choose a lower-central degree larger than the class of the finite three-group P_{3^b} and at least five. The other source $N_{\mathscr{D}}$ has class at most four, whereas every binary boundary lower-central image from degree two is $\mathcal{D}/2^a\mathcal{D}$. It follows that $\psi(N_B)$ contains the exact subgroup

$$\{1\} \times \mathcal{D}/2^a\mathcal{D} \times \{1\}. \tag{E.31.15}$$

After quotienting by this subgroup, W_N embeds in the product of $N_{\mathscr{D}}$, $P_{2^a}^{\mathrm{ab}}$ and P_{3^b}. It is finite nilpotent of primes two and three, generated by the 97 source generators and surjecting onto $N_B^{\mathrm{ab}}/6$. Its two Sylow Frattini quotients are therefore exactly the rank-97 quotients already controlled by the G_6 automorphism. This proves surjectivity after quotienting by (E.31.15), and then on W_N itself. The Q projection proves surjectivity of the complete shadow. The same argument with only the ternary boundary factor also proves that every finite correction image counted in (E.31.4) is generating.

Now append any modulus q prime to six. The actual Chinese remainder maps identify

$$B/[H,H]H^{\,2^a3^bq} \simeq B/[H,H]H^{\,2^a3^b} \times_{B/H} B/[H,H]H^q. \tag{E.31.16}$$

On the normal H coordinates this is the ordinary isomorphism $M_H/(2^a3^bq) \simeq M_H/(2^a3^b) \times M_H/q$; lifting a common B/H coordinate proves the displayed group isomorphism. The old common source has order involving only two and three. Its H image and M_H/q consequently have coprime orders. The original and the corrected H images project onto both: for the corrected map, the first follows from the just-proved surjectivity and its identity B/H map, and the second follows because the boundary map at q is literally the identity. A subdirect subgroup of these two groups is their full product: their common quotient has order dividing two coprime integers. This proves generation in the actual common source at 2^a3^bq. These moduli are cofinal, and the one word f^* makes all reductions compatible. Thus (E.31.5) follows.

The executed complete-kernel calculation is `bc_cm_fermat_kernel_boundary.py`, with all normal-relation metadata, integer matrices, Smith transformations and the 432 evaluated solutions. Its companion `bc_cm_fermat_kernel_words.py` retains the actual lattice-generator preimages and the embedding check (E.31.11). The earlier restricted test and the literal two-adic comparison are also retained, with their exact scopes. □

Statement E.32 (Proved). Every retained binary fibre passes the order-two torsion-class test. *Let $T_{2,6}$ be the exact characteristic-two Ferrari target of E.24, and put $\delta = abc$, $z_4 = \delta^4$. For all 96 branch/line inputs of E.24, including both multiplier residues $1, 73$ and both binary correction points in each of the 24 retained fibres, the resulting triple satisfies*

$$(a'b'c')^2 = h\delta^2h^{-1} \qquad \text{for an explicitly certified } h \in T_{2,6}. \tag{E.32.1}$$

Thus the necessary conjugacy condition of E.30 holds already in $T_{2,6}$ itself. It holds for every one of the 864 joint points of E.24, since its nine ternary parameters leave the binary coordinates unchanged. This assertion retains the specified finite fibres and makes no full profinite realization assertion.

Proof. All group operations take place in the exact tensor-algebra semidirect product of E.24. Let $U = \ker(T_{2,6} \to \mathrm{SL}_2(\mathbb{F}_2))$. The complete collection certificate gives $|U| = 2^{40}$. Give a unit increment of degree d weight d, and an automorphism increment of degree $d+1$ weight d. Let U_d consist of the elements having zero coordinates of weight less than d. Multiplication and substitution in the actual truncated tensor algebra show

$$[U_i, U_j] \subset U_{i+j}, \qquad U_d/U_{d+1} \text{ has exponent two}, \qquad U_7 = 1. \tag{E.32.2}$$

These assertions use the positive-weight filtration of the collected group, including all square terms.

Here is the complete conjugacy algorithm, including its centralizer fibres. Enumerate six actual representatives of $T_{2,6}/U$. For each representative g, put $x_g = g\delta^2g^{-1}$, which belongs

to U. At stage d, let C_d be the full subgroup of U centralizing x_g modulo U_d. Initially $C_1 = U$. The actual commutator gives a homomorphism

$$L_d : C_d \longrightarrow U_d/U_{d+1}, \qquad c \longmapsto [c, x_g] \bmod U_{d+1}. \tag{E.32.3}$$

It is a homomorphism because the target is central in U/U_{d+1}. Its matrix is computed on an exact polycyclic basis of C_d. Products corresponding to the binary nullspace give actual elements of $\ker L_d$; exact group collection of these products has order $|C_d|/2^{\operatorname{rank} L_d}$. Containment in the kernel is checked by substitution. This order equality proves that the entire kernel, rather than a subset of its lifts, has been retained. The next centralizer is exactly $C_{d+1} = \ker L_d$.

Suppose hx_gh^{-1} agrees with a specified target y modulo U_d. Its discrepancy in U_d/U_{d+1} is an explicit vector. Solving (E.32.3) for that vector replaces h by hc and gives agreement modulo U_{d+1}. All solutions at that stage form the coset hC_{d+1}. An inconsistent linear system rules out that entire coset. Running through all six outer representatives therefore decides conjugacy in the full target. The exact centralizer dimensions and successive image ranks are, in every outer coset,

	C_1	C_2	C_3	C_4	C_5	C_6	C_7
$\log_2 \lvert C_d \rvert$	40	40	38	37	34	31	27

, $\qquad (\operatorname{rank} L_d)_{d=1}^{6} = (0, 2, 1, 3, 3, 4).$ (E.32.4)

The central element z_4 has order four. The program tests all four possible translates $(a'b'c')^2 z_4^j$ for each input. There are exactly two distinct values of $(a'b'c')^2$, hence eight distinct translated targets. In every one of the 96 inputs, precisely the translate $j = 0$ is conjugate to δ^2. The resulting conjugators are substituted into the complete group multiplication to verify (E.32.1).

The 96 inputs are reconstructed directly from the complete seven-dimensional binary correction module, its two affine branches, and the retained cusp gauge. If f changes to fk, its two corrected braid generators change to $k_5^{-1}b'k_5$ and $k_6^{-1}c'k_6$ in the corresponding evaluation factors. This exact formula retains both binary points at both multiplier residues. The ternary homogeneous corrections are represented by the powers u^{-2} of E.24, whose binary image is identity, proving the assertion about all 864 joint points.

Two independent controls accompany the computation. First, the same conjugacy algorithm is run in the degree-two target of order 1536 and compared with direct enumeration of every target element and every conjugate. All 1536 decisions agree; the conjugacy class has size 12. Second, a separate program imports no conjugacy solver. It reconstructs all 40 collected basis elements from their literal words in the original braid generators, collects each certified conjugator, and obtains explicit B_4 words. Substitution of these words verifies all 96 equalities (E.32.1). The certificates are `de_ferrari_two_conjugacy.py`, `de_ferrari_two_torsion.py`, and `de_ferrari_two_torsion_words.py`, together with the executed outputs, all centralizer matrices, literal conjugators and indexed input map. □

Statement E.33 (Proved). The full rank-ten class-two refinement at three. *Retain the actual groups $H \triangleleft F \triangleleft B$, the ten signed Schreier generators $\eta_1, \dots, \eta_{10}$ of E.26, and the common Ferrari source $\mathscr{D}$ of E.24. Put*

$$P_3 = B/(\gamma_3(H)H^3), \qquad \mathscr{J}_3 = \operatorname{im}(B \longrightarrow \mathscr{D} \times P_3). \tag{E.33.1}$$

The geometric kernel $H/(\gamma_3(H)H^3)$ has order 3^{55}, with ten abelian and forty-five commutator coordinates. Every one of the 432 multiplier-one points of E.24 has a generating lift to the actual common source $\mathscr{J}_3$. For each such point the complete set of correction images coming from $\ker(\widehat{B} \to \mathscr{D}) \cap \widehat{F}'$ and satisfying both order relations has exactly

$$3^{11} \tag{E.33.2}$$

elements.

This refinement imposes a strict additional condition on the abelian boundary fibre. For a commutator word f, write its H-abelianization modulo three as v. The abelian order relations and the exact F-abelianization condition give

$$v = v(a,b,c) = (c,b,a,b,a+2c,2a+c,2a,a+2c,c,2c)^t, \qquad a,b,c \in \mathbf{F}_3. \tag{E.33.3}$$

The class-two order relations admit a central coordinate over v exactly when

$$a-b-c=0, \qquad \textit{equivalently} \qquad v_3-v_2-v_1=0. \tag{E.33.4}$$

Each admissible v has an affine ten-dimensional space of central coordinates. In each complete E.24 correction fibre, exactly three of the nine abelian solutions of E.31 satisfy (E.33.4)*. Thus six actual abelian norm solutions in each such fibre do not extend to* (E.33.1)*, while all* 432 *source points retain generating extensions. Every one of the* 3^{11} *finite solutions over a given source point extends compatibly and simultaneously through*

$$\mathscr{J}_{3,m} = \operatorname{im}\bigl(B \longrightarrow \mathscr{D} \times P_3 \times B/[H,H]H^m\bigr), \qquad m \geq 1. \tag{E.33.17}$$

One profinite commutator word realizes each such compatible family, retaining all the Ferrari coordinates.

Proof. The normality of the subgroup in (E.33.1) follows from characteristicness inside the actual normal subgroup H. For $q=(a,b)\in(\mathbb{Z}/3)^2$ use the section $c(q)=x^ay^b$, with $0\leq a,b<3$. The nontrivial signed Schreier edges $c(q)\ell c(q+\ell)^{-1}$, in the fixed order, are

$$\begin{gathered}((0,1),x),\ ((0,2),x),\ ((0,2),y),\ ((1,1),x),\ ((1,2),x),\ ((1,2),y),\\ ((2,0),x),\ ((2,1),x),\ ((2,2),x),\ ((2,2),y).\end{gathered} \tag{E.33.5}$$

This specifies every generator and every based comparison without a choice of a further conjugating path.

Let $V=\mathbf{F}_3^{10}$, with this ordered basis, and $Z=\Lambda^2V$. The explicit isomorphism

$$H/\bigl(\gamma_3(H)H^3\bigr) \longrightarrow V \oplus Z, \qquad \eta_i \longmapsto (e_i,0), \tag{E.33.6}$$

uses the multiplication

$$(v,z)(w,t) = \bigl(v+w, z+t+2v\wedge w\bigr). \tag{E.33.7}$$

Hall collection gives the inverse map. The ordinary commutators are the independent coordinates of Z; all cubes in (E.33.7) are one. Conversely the class-two collection identities and $\binom{3}{2}=3$ show that the relations $\gamma_3(H)=H^3=1$ introduce no further relation among these coordinates. This proves (E.33.6), rather than only a graded dimension count. Since $[B:H]=54$, the order of P_3 is $54\cdot 3^{55}$.

Here are the exact action and norm calculations. On the free group F the actual substitutions are

$$s:(x,y)\longmapsto(y,x), \qquad r:(x,y)\longmapsto\bigl(y,(xy)^{-1}\bigr). \tag{E.33.8}$$

They are the negative-square convention verified in E.26. Substitute each of the ten words (E.33.5), perform the signed Schreier rewrite, and collect by (E.33.7). This gives matrices

$$\mathsf{S}=\begin{pmatrix} S_1 & 0 \\ J_s & \Lambda^2 S_1 \end{pmatrix}, \qquad \mathsf{R}=\begin{pmatrix} R_1 & 0 \\ J_r & \Lambda^2 R_1 \end{pmatrix}. \tag{E.33.9}$$

The computed matrices satisfy $\mathsf{S}^2=\mathsf{R}^3=1$ and preserve the displayed bracket. Their Johnson blocks have ranks seven and nine. The actual words $x=(r^{-1}s)^2$ and $y=(sr^{-1})^2$

give precisely conjugation by x and y on all fifty-five coordinates. Thus no diagonal action has been substituted for the based one.

For completeness, the finite group operations used in the certificate also have an explicit source comparison. Write an arbitrary F word as $h\,c(q)$ by signed Schreier rewriting. Multiplication is

$$(h,q)(k,t) = \bigl(h^{\,c(q)}k\,c(q)c(t)c(q+t)^{-1},\; q+t\bigr). \tag{E.33.10}$$

Both the conjugation and each of the eighty-one carries in this formula are collected from their literal words. This evaluates every relator used below in the actual quotient, including all power terms.

For $g=(v,z)$ let multiplication in the following expression mean (E.33.7), and let the displayed product be its logarithmic coordinate:

$$\mathcal{N}(g) = \begin{pmatrix} (\mathsf{S}g)\,g \\ (-g)\,(\mathsf{R}(-g))\,(\mathsf{R}^2(-g)) \end{pmatrix}. \tag{E.33.11}$$

These are exactly $(sf)^2$ and $(f^{-1}r)^3$. Their abelian parts are $(1+S_1)v$ and $-(1+R_1+R_1^2)v$. The map $H^{\mathrm{ab}} \to F^{\mathrm{ab}}$ is the integral map $3L$ of E.28. A word in F' therefore has $L[f]=0$ before reduction, and hence $Lv=0$. Solving these three literal linear conditions gives (E.33.3). The $Lv=0$ condition is retained; the two norm conditions alone have a four-dimensional kernel modulo three.

The central part of (E.33.11) has the form

$$Mz + e(v), \qquad M = \begin{pmatrix} 1+\Lambda^2 S_1 \\ -1-\Lambda^2 R_1 - (\Lambda^2 R_1)^2 \end{pmatrix}. \tag{E.33.12}$$

Its rank is 35, so its kernel has dimension ten. There is a small certificate for the extra obstruction. For the central coordinates t_{ij} of the second norm component put

$$\begin{aligned} \omega(t) = {} & t_{13}+t_{14}-t_{15}+t_{16}+t_{17}-t_{18} \\ & +t_{23}+t_{24}+t_{27}-t_{28}. \end{aligned} \tag{E.33.13}$$

Literal matrix multiplication gives $\omega M = 0$ and

$$\omega e(v(a,b,c)) = a-b-c. \tag{E.33.14}$$

The certificate verifies this as a polynomial identity: the three linear coefficients are $(1,-1,-1)$ and every square and cross coefficient is zero. It also solves (E.33.12) for each of the nine vectors $v(b+c,b,c)$. These forty-five-coordinate particular solutions, together with a basis of $\ker M$, are saved in the exact certificate. This proves both directions of (E.33.4), and not only a necessary linear test.

We next prove that the correction image is complete. Put $R_3 = \ker(B \to D_6) \cap F'$. Use the complete $6,668$ normal relators and their actual word metadata from E.31: the $4,656$ rank-97 commutators, the 95 primitive integral-F^{ab} kernel power products, the eighteen basis cubes, the 153 basis commutators and the $1,746$ conjugation relations, each corrected by its literal eighteen-coordinate representative. Evaluate every one by (E.33.10). Their logarithms span a 51-dimensional subspace, already stable under S,R and every bracket. Its abelian projection is exactly the six-dimensional reduction $\overline{K}$ of the integral lattice K in E.31, and its central intersection is all of Z. Thus

$$\operatorname{im}(R_3 \to V \oplus Z) = \overline{K} \oplus Z. \tag{E.33.15}$$

The logarithmic span equals the group image: in (E.33.7), addition is obtained from group multiplication by multiplying by the appropriate half-power of the commutator, and scalar

multiplication is ordinary group powering. Stability under both actual B generators and brackets therefore proves equality with the full normal image. Every chosen spanning vector retains a relator or conjugation/commutator preimage in the saved certificate.

The same image is obtained from $\ker(B \to \mathscr{D}) \cap F'$. Indeed the binary K_6 image has exponent two. For each $u \in R_3$, the word u^4 kills that binary image, kills the ternary image, and has the same image as u in the exponent-three group (E.33.6). This is an actual word map retaining all induced Ferrari coordinates. Density of the discrete commutator subgroup in the full profinite commutator subgroup, intersected with the specified open source kernel, gives the same finite image for the profinite notation in the statement.

For each retained literal word f, the allowed targets therefore form exactly its coset by $\overline{K} \oplus Z$. On $\overline{K}$ the abelian norm map has rank four and kernel dimension two, giving the nine abelian options. The literal 432-word reconstruction computes these affine planes and verifies that each meets (E.33.4) in a line of three elements. Each point of that line has 3^{10} central solutions by (E.33.12). Multiplication by the fixed value of f is a bijection on correction images, proving the exact count (E.33.2). For example the actual E.24 point with indexed CM row 2, ternary parameters $(0, 0)$ and binary parameter 0 has the abelian solution

$$(0, 0, 1, 0, 1, 2, 2, 1, 0, 0)^t. \tag{E.33.16}$$

Its obstruction in (E.33.14) is one. This supplies a specified boundary solution that does not lift, while the other three abelian solutions for that source point do lift.

There is an exact kernel intersection that supplies the simultaneous assertion. Let $g_1, \ldots, g_8 \in R_3$ be the actual words of E.31 whose integral H-abelianizations are the basis columns of K. For an original normal relator u write, uniquely, $[u]_H = Kc_u$ with $c_u \in \mathbb{Z}^8$, and form

$$u^\circ = u \left(\prod_{j=1}^{8} g_j^{(c_u)_j} \right)^{-1}. \tag{E.33.18}$$

This is an actual element of $R_3 \cap [H, H]$.

The program `bc_fermat_class_two_integral_kernel.py` computes the exact integral c_u, using the same eight literal words, and evaluates (E.33.18). Forty-five of these adjusted relators have linearly independent central images modulo three. Their indices and all eight integral exponents are saved. Consequently

$$\operatorname{im}(R_3 \cap [H, H] \longrightarrow Z) = Z. \tag{E.33.19}$$

This equality follows already from these forty-five word preimages; it requires no assertion that a finite kernel automatically lifts integrally. Taking fourth powers again retains their central images and puts them in $\ker(B \to \mathscr{D}) \cap [H, H]$.

Now prescribe any one of the finite solutions counted in (E.33.2). E.31's proved finite-fibre comparison lifts its abelian correction modulo three through the entire three-adic norm fibre. The two-adic and prime-to-six supported corrections of that statement give one word realizing the abelian tower at every modulus with that specified reduction. The central discrepancy from the prescribed P_3 solution belongs to Z. Correct it by a product of the actual fourth powers in (E.33.18)–(E.33.19). These words have zero integral H-abelianization, so every abelian modulus and every Ferrari coordinate is retained. The resulting single word has the prescribed P_3 image and both norm equations in every factor of (E.33.17).

Finally all these finite norm solutions generate the common source. Let W be the image of N_B in $\mathscr{J}_3$. Its image in $\mathscr{D}$ is finite nilpotent with prime divisors two and three, and its image in P_3 is a three-group: the latter maps to the cyclic-three image in F/H with a three-group kernel. Consequently W is finite nilpotent, is generated by the 97 specified

generators, and maps onto $(\mathbb{Z}/6)^{97}$ through the retained G_6. Its two Sylow Frattini quotients therefore both have dimension exactly 97. The corrected map is a homomorphism into the actual common image, because both norm equations hold in both factors. It preserves N_B through its retained Q map and induces the known G_6 automorphism on these Frattini quotients. The finite Frattini criterion gives surjectivity on W, and then on $\mathscr{J}_3$. This proves generation without imposing invariance of the fixed Ferrari source kernel.

Generation in (E.33.17) follows through the same actual groups. First take $m = 2^a 3^b$ with $a, b \geq 1$. Choose a lower-central index larger than the nilpotence classes of the images of N_B in $\mathscr{D}$, in P_3 and in the three-power abelian boundary quotient. Its image in the two-power boundary quotient is exactly $\mathcal{D}/2^a\mathcal{D}$ by E.28. The corrected image projects onto that two-power boundary group, since the already generating abelian action has not changed. Thus this deep commutator subgroup of the corrected joint image is exactly the pure two-primary factor $\mathcal{D}/2^a\mathcal{D}$. After quotienting it out, the joint N_B image is finite nilpotent with primes two and three and has the same two rank-97 Frattini quotients through G_6. The preceding Frattini argument therefore proves joint surjectivity. For a further factor of m prime to six, the actual H image in all existing factors is a two-three group, whereas the additional abelian H image has order prime to six. The exact CRT map and coprime subgroup argument of E.31 give their full product over B/H. The added P_3 factor changes neither coprimality nor the prescribed identity action on this prime-to-six boundary factor. This proves surjectivity for every m; the same constructed word proves compatibility.

The programs `bc_fermat_class_two.py`, `bc_fermat_class_two_kernel.py`, `bc_fermat_class_two_fibres.py`, and `bc_fermat_class_two_obstruction.py` save the full matrices, literal preimage metadata, all finite solutions and the polynomial obstruction certificate. Each executed output ends in `PASS`. The additional integral-kernel certificate supplies the actual preimages needed for the simultaneous all-moduli comparison. □

Statement E.34 (Proved). The four-strand order-three torsion class gives a second finite test. *Keep the finite quotient $q : B_4 \twoheadrightarrow T$ and the elements a, b, c, z, C of E.30. Put*

$$e = a^2bc, \qquad e_F = a_F^2 b_F c_F \quad \text{for } F = (\lambda, f) \in \widehat{\mathrm{GT}}. \tag{E.34.1}$$

Then $e^3 = z$, and every full-GT element satisfies

$$e_F C \sim_{T/C} e^{\lambda} C. \tag{E.34.2}$$

Equivalently there exist $h \in T$ and $j \in \mathbb{Z}/|z|$ with

$$e_F = h e^{\lambda} h^{-1} z^j. \tag{E.34.3}$$

Only the residue of λ modulo three enters the class on the right of (E.34.2)*. In particular both E.24 multiplier residues $1, 73$ give the class eC there. The test uses the actual homomorphism $q \circ F$ and the capping map of E.30.*

Proof. Write $\delta = \sigma_1\sigma_2\sigma_3$ and $\epsilon = \sigma_1^2\sigma_2\sigma_3$ in B_4. There is a short proof of the exact braid identity $\epsilon^3 = \delta^4$. Using a digit i for σ_i, its successive positive words are

$$\begin{aligned} 112311231123 &= 112131231123 = 112131213123 \\ &= 121231213123 = 121232123123 \\ &= 121323123123 = 123123123123. \end{aligned} \tag{E.34.4}$$

The six changes use respectively distant commutation, distant commutation, the $121 = 212$ relation twice, the $232 = 323$ relation, and distant commutation. The program

`e_periodic_braid_root.py` independently verifies every location and substitution. Applying q proves $e^3 = z$.

Let $\bar{\epsilon}$ be the image in $B_4/\langle\delta^4\rangle$, identified with the fifth-point stabilizer in $\Gamma_{0,[5]}$ by (E.30.5). Formula (E.34.4) and its permutation (234) show that its order is exactly three, with fixed marked points $1, 5$. The actual Nakamura–Schneps action comparison (E.30.5a)–(E.30.5c) makes $F(\bar{\epsilon})$ another element of order three. The action-independent torsion classification in Collas's Theorem 3.3 and Corollary 3.2 makes it conjugate to $\bar{\epsilon}$ or $\bar{\epsilon}^2$. The explicit normalizer below conjugates these two powers. Since λ is a unit modulo three, the resulting conjugacy can be written

$$F(\bar{\epsilon}) = u\bar{\epsilon}^{\lambda}u^{-1}, \qquad u \in \widehat{\Gamma}_{0,[5]}. \tag{E.34.5}$$

We must locate this conjugator in the actual fifth-point stabilizer before passing to T/C.

There is an explicit geometric normalizer reversing $\bar{\epsilon}$ and interchanging its two fixed points. The periodic braid in (E.34.4) is the order-three representative ϵ_{n-2} for $n = 5$ in Collas's Corollary 3.2 [Col12]. Its rotation model has the five marked points

$$\{0, \infty, 1, \zeta_3, \zeta_3^2\}, \qquad R(t) = \zeta_3 t. \tag{E.34.6}$$

The order-two Möbius map $w_0(t) = 1/t$ preserves this set, exchanges 0 and ∞, and satisfies $w_0 R w_0^{-1} = R^{-1}$ by literal substitution. Transporting this map by the marked-sphere identification with the periodic braid gives $w \in \Gamma_{0,[5]}$ such that

$$w\bar{\epsilon}w^{-1} = \bar{\epsilon}^{-1}, \qquad w(1) = 5, \quad w(5) = 1. \tag{E.34.7}$$

This is an actual discrete mapping class and hence also gives an element of its profinite completion. The two additional equatorial points are exchanged and the third is fixed, as recorded by the literal permutation check in the program.

The Drinfeld action preserves the permutation of $\bar{\epsilon}$. Formula (E.34.5) therefore makes u preserve its fixed-point set $\{1, 5\}$. If its permutation exchanges these points, then $v = uw$ fixes 5 and (E.34.7) gives

$$F(\bar{\epsilon}) = v\bar{\epsilon}^{-\lambda}v^{-1}. \tag{E.34.8}$$

Both sides and the conjugator now lie in the completed fifth-point stabilizer. The abelianization of that group is $\mathbb{Z}/12$: the braid relations identify all three generator classes, and the killed full twist has exponent sum twelve. This also computes its profinite abelianization, since every finite abelian quotient has exactly those relations. In this quotient $[\bar{\epsilon}] = 4$, and the based GT formulas multiply every braid abelianization by λ. Equation (E.34.8) would imply

$$4\lambda = -4\lambda \pmod{12}, \quad \text{hence } 3 \mid \lambda. \tag{E.34.9}$$

The latter contradicts $\lambda \in \widehat{\mathbb{Z}}^{\times}$. Thus u fixes 5 itself, proving the conjugacy (E.34.5) inside $\widehat{B}_4/\overline{\langle\delta^4\rangle}$.

The actual map to T/C factors through this central quotient. Its application to (E.34.5) proves (E.34.2), and lifting the conjugator to T proves (E.34.3). Since $e^3 = z$, its class in T/C has order dividing three, which proves the claimed multiplier scope. This argument requires no invariance of the finite kernel of q under the GT action. The earlier order-two test and this order-three test use their respective explicit periodic elements and the same equivariant capping homomorphism. □

Statement E.35 (Proved). All retained joint fibres pass the order-three torsion-class test. *Let $T_{23} = T_{2,6} \times T_{3,6}$ be the exact joint Ferrari target of E.24. Write $e = a^2bc$, $z_4 = (abc)^4$, and $e' = a'^2b'c'$ for the image of the same word under a retained braid triple.*

For every one of the 864 *points of E.24, and every compatible multiplier lift* $\lambda \equiv 1 \pmod{72}$*, there is* $h \in T_{23}$ *with*

$$e' = he^{\lambda}h^{-1}. \tag{E.35.1}$$

Thus every retained fibre passes E.34 already in the actual joint target. On the ternary factor, all 216 *parameter points have been evaluated. There are* 15 *distinct values of* e'*; for every such value,*

$$e'z_4^j \sim_{T_{3,6}} e \quad \Longleftrightarrow \quad j \equiv 0 \pmod 3, \qquad j \in \mathbb{Z}/9. \tag{E.35.2}$$

All 45 *positive comparisons have saved literal* B_4 *conjugators. The assertion concerns these exact finite fibres and does not assert a full profinite lift.*

Proof. **Complete conjugacy in the ternary target.** Use the actual projection from E.23,

$$1 \longrightarrow J \longrightarrow T_{3,6} \longrightarrow T_{3,2} \longrightarrow 1, \qquad |T_{3,2}| = 648. \tag{E.35.3}$$

The complete four-strand Schreier image, retained in E.23, gives a class-two exponent-three group J with

$$\dim_{\mathbb{F}_3} \log J = 30, \qquad \dim_{\mathbb{F}_3} \log[J,J] = 18. \tag{E.35.4}$$

The logarithm and exponential here are the exact class-two maps on J certified in E.23. We do not apply a logarithm to the whole degree-six group. On this kernel the BCH identity is $u * v = u + v + \frac{1}{2}[u,v]$, with $\frac{1}{2} = 2$ in $\mathbb{F}_3$.

The new program independently enumerates all 648 base elements using actual words in a, b, c, and reconstructs the whole 30-dimensional kernel from the resulting Schreier relations and their commutators. It verifies equality with the saved kernel of E.23 and records a literal B_4 word for every basis element. Let $Z = \log[J,J]$, let $V = \log J$, and put $A = \mathrm{Ad}(e)$ on V. All these maps are computed by substitution in the original finite group.

For a prescribed target y, enumerate every base conjugator q satisfying $q\bar{e}q^{-1} = \bar{y}$ in $T_{3,2}$, and choose its recorded lift g_q. There are 72 base conjugates of $\bar{e}$ and nine possible base conjugators for each of them. Conjugacy above a given q is equivalent to conjugating e to $g_q^{-1}yg_q$ by an element of J. Its first equation is

$$(1-A)u \equiv \log\bigl(g_q^{-1}yg_qe^{-1}\bigr) \pmod{Z}. \tag{E.35.5}$$

This is the exact equation modulo the central subgroup $[J,J]$. Its homogeneous solution space is

$$V_1 = \ker\bigl(V \xrightarrow{1-A} V/Z\bigr).$$

The corresponding group $\exp(V_1)$ is the full centralizer of e modulo $[J,J]$. For $v \in V_1$, $Av - v$ is central, so the exact class-two formula gives

$$[\exp(v), e] = \exp((1-A)v). \tag{E.35.6}$$

In particular the second equation, after any solution u of (E.35.5), is the linear equation on *all* of V_1 given by the remaining central discrepancy. Multiplying $\exp(u)$ on the right by $\exp(v)$ corrects that discrepancy. Its full homogeneous fibre is $\exp(V_2)$, where

$$V_2 = \ker\bigl((1-A)|_{V_1}\bigr). \tag{E.35.7}$$

The first solution set is an entire coset of $\exp(V_1)$; the second is an entire coset of $\exp(V_2)$. Thus both kernel spaces retain every possible conjugator lift. An inconsistent equation

excludes that complete coset, and enumeration of all base conjugators decides conjugacy in $T_{3,6}$. The computed ranks and dimensions are

$$\begin{array}{c|ccc} & V & V_1 & V_2 \\ \hline \dim & 30 & 22 & 10 \end{array}, \qquad \operatorname{rank}\overline{(1-A)} = 8, \quad \operatorname{rank}((1-A)|_{V_1}) = 12. \tag{E.35.8}$$

Every ternary input is reconstructed from its retained E.23 braid triple and its complete two-dimensional homogeneous correction plane. For each of the 24 indices, the program evaluates all nine parameters and all nine central translates of the resulting e'. The following exact facts are checked:

$$e^3 = z_4, \quad |z_4| = 9, \quad |e| = 27, \qquad (e')^3 = z_4. \tag{E.35.9}$$

All nine parameters in a given indexed plane have the same value of e'. Across the 24 indices there are 15 values, giving 135 distinct translated targets. The complete two-stage algorithm gives exactly the three positive translates $j = 0, 3, 6$ for each value. The other six translates fail after the complete base and kernel search. This proves (E.35.2).

The resulting 45 conjugators have literal braid words of lengths between 6 and 912. They are obtained by multiplying the Schreier words for a logarithm's basis coordinates and then correcting the resulting central BCH term with the recorded central commutator words. Every reconstructed word is evaluated in the actual group and checked against the computed conjugator. A separate program imports no conjugacy solver: it evaluates all 45 words, verifies the conjugacy identities, and reconstructs all 216 original parameter inputs independently. The identity GT element, whose triple is (a, b, c), is retained as a literal control.

For the multiplier scope, the binary CRT corrections of E.24 have identity image in the entire ternary source, so this test covers both binary parameters of each point. Both multiplier branches have the same ternary triple; the exact braid-generator orders divide nine. Moreover

$$e^{73} = ez_4^6, \qquad e^\lambda = ez_4^{(\lambda-1)/3} \quad (\lambda \equiv 1 \pmod{72}). \tag{E.35.10}$$

The exponent $(\lambda - 1)/3$ is a multiple of three. All its residues modulo nine are among the three positive translates in (E.35.2). Hence e' is conjugate to e^λ in the ternary group itself for every compatible higher multiplier lift.

An exhaustive auxiliary control tests the same two-stage method on the class-two exponent-three group with generators a, b, c and central relation $[b, c] = 1$, of order $3^5 = 243$. The order-three automorphism fixes a, sends b to $b + c$ in logarithmic coordinates, and fixes c; it sends $[a, b]$ to $[a, b] + [a, c]$. All 243 target values are compared with the complete enumeration of all 243 conjugators. All decisions agree; the class has nine elements and the two successive ranks are one and one.

The binary factor and the actual product. Let $U = \ker(T_{2,6} \to \mathrm{SL}_2(\mathbb{F}_2))$. It is the finite 2-group of E.24, and its center contains $\langle z_4 \rangle$ of order four. In $T_{2,6}/\langle z_4 \rangle$, both e^λ and e' have order three. Their images in $\mathrm{SL}_2(\mathbb{F}_2) = S_3$ are the same specified generator of C_3: the three retained braid generators have the original degree-two images, and, writing A, B for their linear transvections,

$$\overline{e} = A^2BA = BA, \qquad \overline{e'} = BA, \qquad (BA)^\lambda = BA. \tag{E.35.11}$$

Here $A^2 = 1$ and $\lambda \equiv 1 \pmod 3$.

We recall an elementary proof of the needed coprime conjugacy fact. Two order-three complements to a normal finite 2-group P are conjugate by P. Induct on $|P|$ and use the nonzero characteristic elementary abelian subgroup $D = \Omega_1 Z(P)$. The induction hypothesis conjugates the two complements to agree modulo D. For their generators with the same

specified image in C_3, write the remaining difference as $t = vs$, $v \in D$, and let α be conjugation by s. The equation $t^3 = 1$ is

$$v + \alpha v + \alpha^2 v = 0.$$

In the $\mathbb{F}_2$-vector space D, take $w = \alpha v$. Then $(1 - \alpha)w = v$, so conjugation by w removes the difference. This completes the induction and keeps the generator orientation fixed.

Apply this argument to the preimage of the displayed C_3 in $T_{2,6}/\langle z_4 \rangle$, whose normal 2-group is $U/\langle z_4 \rangle$. Lifting its conjugator gives

$$e' = h_2 e^{\lambda} h_2^{-1} z_4^j.$$

Both cubes are z_4^{λ}: this follows from the literal braid identity of E.34 and E.24's full-twist calculation. Consequently $3j = 0 \pmod 4$, so $j = 0$. This proves the binary equality in the group itself.

Finally E.24 proves that the actual joint target is the full product $T_{2,6} \times T_{3,6}$. Pairing the binary conjugator with the ternary one therefore gives an element of that actual target and proves (E.35.1), with no further joint-generation assumption. The new executed files are `de_ferrari_three_conjugacy.py`, `de_ferrari_three_torsion.py`, and `de_ferrari_three_torsion_words.py`, with the complete centralizer matrices, translated target decisions, indexed parameter map, and literal conjugators. □

Statement E.36 (Proved). The complete exponent-four refinement and its joint exponent-twelve fibres. *With $H \triangleleft F \triangleleft B$ and $\mathscr{D}$ as in E.33, put*

$$P_4 = B/(\gamma_3(H)H^4), \qquad P_{12} = B/(\gamma_3(H)H^{12}). \tag{E.36.1}$$

Every one of the 432 *multiplier-one points of E.24 has exactly* 2^{10} *complete correction images giving generating lifts to* $\mathrm{im}(B \to \mathscr{D} \times P_4)$*. All four abelian boundary solutions modulo four lift; each has an affine eight-dimensional space of central corrections over* $\mathbf{F}_2$*.*

Every pair consisting of one of these solutions and one of the 3^{11} *solutions in E.33 is realized by one commutator word. The complete joint correction fibre in* $\mathrm{im}(B \to \mathscr{D} \times P_{12})$ *therefore has*

$$2^{10}3^{11} = 181398528 \tag{E.36.2}$$

images over each retained source point. Every such solution extends compatibly and simultaneously through

$$\mathrm{im}(B \longrightarrow \mathscr{D} \times P_{12} \times B/[H,H]H^m), \qquad m \geq 1. \tag{E.36.3}$$

All these finite shadows are generating and retain the complete binary and ternary Ferrari coordinates.

Proof. We keep the exact ordered Schreier basis (E.33.5). At exponent four the correct Hall coordinates are

$$H/(\gamma_3(H)H^4) = (\mathbb{Z}/4)^{10} \times (\mathbb{Z}/2)^{45}, \qquad \prod_i \eta_i^{v_i} \prod_{i<j} [\eta_i, \eta_j]^{z_{ij}}. \tag{E.36.4}$$

The multiplication and powers are

$$\begin{aligned} (v,z)(w,t) &= (v+w,\ z+t-(v_j w_i)_{i<j}), \\ (v,z)^n &= \big(nv,\ nz-\tbinom{n}{2}(v_i v_j)_{i<j}\big), \end{aligned} \tag{E.36.5}$$

with the first coordinate reduced modulo four and the second modulo two. In any class-two exponent-four group, the identities for $(ab)^4$ and $[a,b]^4$ give $[a,b]^2 = 1$, since $\gcd(6,4) = 2$.

Conversely (E.36.5) has exponent four, and its independent collected coordinates give the inverse to the generator map from the free class-two exponent-four group. This proves the isomorphism (E.36.4). Its order is 2^{65}, and its central Frattini subgroup consists of the 55 binary coordinates $(v/2, z)$ with v even. In particular the square of (v, z) contains the term $(v_i v_j)_{i<j}$; it is retained throughout. At modulus two there is no corresponding new commutator layer: $(ab)^2 = a^2 = b^2 = 1$ gives $ab = ba$, so $\gamma_3(H)H^2 = H^2$.

Use the actual substitutions (E.33.8) and the same nine-coset carries (E.33.10), now collected by (E.36.5). If the ten images under an action have abelian columns A_i and central columns J_i, the image of (v, z) is the ordered product of their v_ith powers and their pairwise commutators to the powers z_{ij}. Equivalently its central part is

$$\sum_i v_i J_i - \sum_i \tbinom{v_i}{2}(A_{\ell i}A_{ki})_{k<\ell} - \sum_{i<j} v_i v_j (A_{\ell i}A_{kj})_{k<\ell} + (\Lambda^2 A)z. \tag{E.36.6}$$

This formula, reduced modulo two, preserves all the square terms and gives the exact action on every Hall coordinate. The literal calculations verify $s^2 = r^3 = 1$ and verify that the negative-square words for x, y induce their actual inner actions.

Here is the complete source-kernel computation. Let $R_2 = \ker(B \to D_{2,6}) \cap F'$. The binary image of $K_6 \cap F'$ is the actual seven-dimensional elementary-abelian group of E.24. Its complete normal presentation has

$$4656 + 95 + 7 + 21 + 7 \cdot 97 = 5458 \tag{E.36.7}$$

relators: the original commutators and primitive integral-F^{ab} kernel sixth-power products, the seven basis squares, the twenty-one basis commutators and all N_B conjugation relations. For each original generator or conjugate, subtract its actual seven-basis representative by group multiplication. The proof that these normally generate the kernel is the same exact presentation argument as in E.31: after imposing them the quotient is generated by seven commuting elements of order two and still maps onto the actual group of order 2^7.

Evaluate all 5458 relators by (E.36.5)–(E.36.6). Their complete normal image is computed using the parity quotient $(\mathbb{Z}/2)^{10}$ and the central Frattini group. For a dependent parity vector multiply by the inverse of the ordered product of its chosen parity representatives; the resulting even abelian vector and central coordinate belong to the central Frattini group. Retain each parity representative's square and every pairwise commutator, then close under the actual s, r actions. These operations give precisely the generated subgroup, since they include the central collection corrections between all parity representatives.

The resulting image has parity dimension six, central Frattini intersection dimension 45, and high-bit abelian dimension six. Its order is therefore 2^{51}. Its abelian image is exactly $\mathcal{D}/4\mathcal{D} \simeq (\mathbb{Z}/4)^6$, where $\mathcal{D}$ is the primitive lattice of E.28; all six columns are explicitly lifted in the certificate. The intersection with the commutator centre has dimension $45 - 6 = 39$. If $U = \mathcal{D}/2\mathcal{D} \subset \mathbf{F}_2^{10}$, that intersection is exactly

$$C_2 = U \wedge \mathbf{F}_2^{10}, \qquad \dim C_2 = 39. \tag{E.36.8}$$

This follows both by the complete calculation and by the explicit preimages below.

The joint source kernel has this same image. Every $u \in R_2$ belongs to K_6, whose full ternary source image has exponent three. Thus u^9 kills that ternary image, retains the binary source kernel, and has the same image as u in the exponent-four group (E.36.4). This proves the exact comparison with $\ker(\widehat{B} \to \mathscr{D}) \cap \widehat{F}'$. The discrete-to-profinite finite-image comparison is the open-kernel density argument of E.33.

The abelian norm equation on $\mathcal{D}/4$ has exactly four solutions for each retained word, by the Smith factors $(1, 1, 1, 1, 3, 0)$ in E.28. For a chosen abelian correction, first multiply

the selected parity representatives, then solve for the required even abelian difference in the central Frattini group. This supplies an actual element of the complete source-kernel image with that abelian coordinate. All other such elements differ by C_2. The two central norm equations on C_2 form a binary linear map of rank 31. The literal 432-word calculation checks all four abelian choices for each word: every right-hand side lies in that map's image. Its kernel has dimension $39 - 31 = 8$. This proves the exact count $4 \cdot 2^8 = 2^{10}$. The induced action on the ten-dimensional Frattini quotient of the geometric two-group has rank ten in all 1728 abelian cases. The central choices do not change that action.

There are actual zero-abelianization preimages for the entire space (E.36.8). The six parity representatives can be taken to be the normal relators with zero-based indices

$$96,\ 97,\ 98,\ 100,\ 103,\ 104 \tag{E.36.9}$$

in the saved complete presentation. Write them as $u_1, \dots, u_6$. Their parity vectors form a basis of U. The sixty literal words $[u_i, \eta_j]$ have zero integral H-abelianization, and their central images span $U \wedge \mathbf{F}_2^{10}$. The independent preimage checker constructs all sixty words, verifies these zero abelianizations and their Hall coordinates, and retains thirty-nine independent columns. Their ninth powers preserve these central values and kill both the full ternary source and the exponent-three quotient P_3. Consequently

$$\operatorname{im}\bigl(R_2 \cap [H,H] \longrightarrow [H,H]/(\gamma_3(H)H^4)\bigr) = C_2. \tag{E.36.10}$$

This supplies the integral kernel comparison required to prescribe a central value while retaining every abelian modulus.

We spell out the exponent-twelve comparison, including its coordinate change. The Hall coordinates on the geometric kernel of P_{12} are $(\mathbb{Z}/12)^{10} \times (\mathbb{Z}/6)^{45}$ with the first multiplication formula in (E.36.5). The commutator exponent six follows from $\gcd(12, \binom{12}{2}) = 6$, and the same collected-coordinate construction proves independence. The exact maps to the two previous coordinate systems are

$$\begin{aligned} (v, z) &\longmapsto (v \bmod 4,\ z \bmod 2), \\ (v, z) &\longmapsto \bigl(v \bmod 3,\ z + 2(v_i v_j)_{i<j} \bmod 3\bigr). \end{aligned} \tag{E.36.11}$$

The second line is the actual change from ordered Hall coordinates to the BCH coordinates of E.33. Its inverse, together with the first line, is the explicit CRT formula

$$\begin{aligned} v &= 9v_4 + 4v_3 \pmod{12}, \\ z &= 3z_4 + 4(z_3 - 2(v_i v_j)_{i<j}) \pmod 6. \end{aligned} \tag{E.36.12}$$

Both lines respect multiplication. The literal B actions respect these reductions, so they give the actual isomorphism

$$P_{12} \simeq P_4 \times_{B/H} P_3. \tag{E.36.13}$$

The common source-kernel image also splits as the product of its two displayed geometric images. A fourth power retains an exponent-three image and kills an exponent-four image; a ninth power does the converse. Apply these powers to actual preimages in the common source kernel and multiply the resulting words. This proves surjectivity onto the product, and injectivity of the image into that product is immediate from its definition. The two norm equations are tested coordinatewise under (E.36.13), giving (E.36.2).

For simultaneous lifting through (E.36.3), prescribe the desired abelian values modulo four and modulo three. E.28's full two-adic fibre and E.31's full three-adic fibre lift those values using their disjoint supported exponents; their prime-to-six correction supplies all remaining abelian moduli. Correct the central exponent-three discrepancy by the actual

fourth-power words in (E.33.18). They have zero integral H-abelianization and are one in the exponent-four quotient. Then correct the exponent-four discrepancy by the ninth powers from (E.36.9)–(E.36.10). These leave P_3 and every abelian modulus unchanged. This constructs the claimed single profinite commutator word.

Finally we prove joint generation without assuming a fixed-kernel automorphism on $\mathscr{D}$. Each of the boundary maps descends to its displayed quotient, since it preserves H and kills the appropriate characteristic class-two or abelian subgroup. The rank-ten Frattini test makes the P_4 map an automorphism; E.33 gives the same conclusion for P_3, and E.31 for the retained abelian factors. Their actual common boundary quotient therefore also has an automorphism: the intersection of these kernels is preserved, and the induced map has trivial kernel in every factor.

First retain only moduli 2^a3^b. Let T denote this finite common boundary quotient, and let T_N be its N_B image. Choose $r \geq 5$ after the lower-central series of T_N has stabilized, and write $L = \gamma_r(T_N)$. The retained G_6 map places the corrected N_B image inside T_N; its automorphism of the finite group T then makes this inclusion an equality. The corrected map is therefore onto T_N, whereas γ_r of the N_B image in $\mathscr{D}$ is one. Hence the corrected joint image contains exactly the pure factor $\{1\} \times L$ at this lower-central stage. The quotient of the full joint N_B image by that factor is finite nilpotent with prime divisors two and three. It is generated by the 97 source generators and still maps onto $(\mathbb{Z}/6)^{97}$ through G_6. Both Sylow Frattini dimensions are therefore 97, and the retained G_6 automorphism forces joint surjectivity. The quotient by N_B then gives the whole common group. For additional moduli prime to six, the old geometric H image has order divisible only by two and three, whereas the new abelian H image has order prime to six. The actual CRT map and the coprime subgroup argument of E.31 give their full product. This proves generation at every level of (E.36.3).

The exact Hall model, complete source presentation, all 432 fibres, sixty central preimages and the coordinate CRT map are recorded respectively in `bc_fermat_class_two_four.py`, `bc_fermat_class_two_four_kernel.py`, `bc_fermat_class_two_four_fibres.py`, `bc_fermat_class_two_four_central.py`, and `bc_fermat_class_two_crt.py`. Their executed outputs all end in `PASS`; the matrices and literal word certificates accompany them. □

Statement E.37 (Proved). The complete three-adic class-two fibre and the first nonlifting boundary shadow. *Put*

$$P_{3^n} = B/(\gamma_3(H)H^{3^n}), \qquad \mathscr{J}_{3^n} = \mathrm{im}(B \longrightarrow \mathscr{D} \times P_{3^n}). \tag{E.37.1}$$

Fix any one of the 432 *multiplier-one source points of E.24, retaining its literal commutator word. The complete set of correction images satisfying the two order relations in $\mathscr{J}_{3^n}$ has*

$$3^{10n+1} \tag{E.37.2}$$

elements. Its subset lifting to $\mathscr{J}_{3^{n+1}}$ is exactly its subset lifting through every level, and has 3^{10n-1} elements. An explicit two-coordinate criterion for this subset is given in (E.37.15) *below.*

The full inverse-limit fibre of correction images has the explicit coordinate parametrization $\mathbb{Z}_3^{10}$ in (E.37.13)*. Every such compatible family is generating in all the actual common finite sources and is realized by one profinite commutator word. It can be prescribed simultaneously with any of the 2^{10} exponent-four solutions of E.36 and with the whole abelian boundary tower.*

There is an individually specified multiplier-one commutator shadow in $\mathscr{J}_3$ with no lift to $\mathscr{J}_9$, displayed in (E.37.17)–(E.37.18)*. It still has the abelian-tower compatibility proved in E.33 and E.36. The failed projection here is the additional class-two map $P_9 \to P_3$.*

Proof. We use exact integral doubled-log coordinates

$$(v, Z) \in \mathbb{Z}_3^{10} \oplus \Lambda^2 \mathbb{Z}_3^{10}, \qquad Z = 2z, \quad (v, Z)(w, T) = (v + w, Z + T + v \wedge w). \tag{E.37.3}$$

Thus the change from E.33 is the actual invertible map $(v, z) \mapsto (v, 2z)$. Hall collection over $\mathbb{Z}_3$ gives the class-two pro-three quotient of H, and reduction modulo 3^n gives precisely the geometric kernel of P_{3^n}. All entries of the based s, r actions are integral in these coordinates. They are obtained by the same literal substitutions and signed Schreier rewriting as E.33. The program verifies their order relations over $\mathbb{Z}$ and the exact doubled-log comparison with the previously saved modulo-three matrices.

We first retain the full source image. Let $K \subset H^{\mathrm{ab}}$ be E.31's actual eight-column integral kernel lattice. The forty-five words in (E.33.18)–(E.33.19) have exactly zero integral H-abelianization and form a basis of the central reduction modulo three. Their integral class-two central matrix therefore has determinant a three-adic unit. Continuous three-adic powers of these actual words supply every element of the entire central lattice, not just its first reduction. Squaring the words clears the binary source; two is a unit in $\mathbb{Z}_3$, so this does not reduce their three-adic image. The eight actual K-basis words give every element of $K \otimes \mathbb{Z}_3$ in abelianization; the central words then correct their central values arbitrarily. Consequently the exact common source-kernel image is

$$\left\{(v, Z) : v \in K \otimes \mathbb{Z}_3, \quad Z \in \mathbb{Z}_3^{45}\right\}. \tag{E.37.4}$$

This equality is a statement about the image with multiplication (E.37.3); no arithmetic splitting is assumed. The upper inclusion follows from the integral abelian image of E.31. The explicit word preimages prove the reverse inclusion. The same argument proves equality after reduction at every finite level.

The abelian norm equations together with $Lv = 0$ have a primitive integral rank-three kernel. A basis parametrization is

$$v = Bt = (-t_1, t_2, -t_1 + t_3, t_2, t_3, -t_3, t_1 - t_3, t_3, -t_1, t_1)^t. \tag{E.37.5}$$

The nonzero Smith factors of these abelian equations are units at three, so this same parametrization is exact modulo every 3^n. The affine source condition has not been discarded. For a vector in $\ker L$, membership in $K \otimes \mathbb{Z}_3$ is exactly divisibility by three of the two functionals

$$\begin{aligned} \lambda_1(v) &= v_1 - v_4 - v_5 - 11v_9, \\ \lambda_2(v) &= v_2 - 11v_4 - v_8 - v_9. \end{aligned} \tag{E.37.6}$$

They vanish on K and are independent modulo three on $\ker L$; the index of K there has three-part 3^2, proving equality of these conditions with that lattice. On Bt the two functionals both reduce to $t_1 - t_2 - t_3$. For each retained source word f, the literal calculation verifies $\lambda_1([f]) = \lambda_2([f])$ modulo three. Denote this common residue by $c_f \in \{0, 1, 2\}$. The complete affine condition is therefore

$$t_1 - t_2 - t_3 \equiv c_f \pmod 3. \tag{E.37.7}$$

The saved rank-three source lattice has determinant 48 in the coordinates t, so its three-part is exactly the index three in (E.37.7). All 432 residues are saved; their multiplicities are $90, 144, 198$ for $c_f = 0, 1, 2$ respectively.

In the doubled-log coordinates the central norm equations are $MZ + e(t) = 0$. The 90-by-45 integral matrix has a saved unimodular Smith decomposition

$$S_M = U_M M V_M, \qquad \operatorname{diag}(S_M) = (1^{35}, 3, 3, 0^8). \tag{E.37.8}$$

The certificate checks the full matrices over $\mathbb{Z}$, including the unimodular inverses. Write $Y = V_M^{-1} Z$ and $E(t) = U_M e(t)$. The nonzero entries before the two factors three may have the saved unit signs; these are retained in the formula below.

Define the integral polynomials

$$\begin{aligned} Q(t) &= t_1^2 + t_2^2 + t_3^2 + t_1 t_2 + t_1 t_3 - t_2 t_3, \\ F(t) &= t_3 - t_2 + 3Q(t), \\ G(t) &= t_2 + t_3 - 3t_1^2 - 4t_2^2 - 2t_3^2 - 3t_1 t_2 - 5t_1 t_3 + 5t_2 t_3 - (t_3 - t_2) Q(t). \end{aligned} \tag{E.37.9}$$

Index the Smith coordinates from 0 to 44, as in the certificate. The exact polynomial identities are

$$\begin{aligned} E_{35}(t) &= (1 + t_3 - t_2) F(t) + 3G(t), \\ E_{36}(t) &= 3(-t_1 + t_3), \\ E_{63}(t) &= 2F(t). \end{aligned} \tag{E.37.10}$$

Every other zero-Smith-row polynomial is an integer multiple of $F(t)$. These statements are verified coefficient by coefficient, including every quadratic term. The row indexed 63 is among the 90 equations, not among the 45 central unknowns. Since two is a three-adic unit, the complete central-solvability obstruction is

$$F(t) = 0. \tag{E.37.11}$$

When it holds, the two factors three in (E.37.8) are divisible by construction. All the central solutions are exactly

$$\begin{cases} Y_i = -E_i(t)/(S_M)_{ii}, & 0 \le i < 35, \\ Y_{35} = -G(t), \quad Y_{36} = t_1 - t_3, & Z = V_M Y. \\ Y_{37}, \dots, Y_{44} \in \mathbb{Z}_3 \text{ arbitrary}, \end{cases} \tag{E.37.12}$$

Thus no unnamed rational cokernel or divisibility condition remains in the description.

The equation has an explicit global three-adic solution graph. Its derivative with respect to t_3 is $1 + 3(2t_3 + t_1 - t_2)$, a unit. For any t_1, t_2, its reduction has the unique root $t_3 = t_2$ modulo three. Starting with that root, at precision 3^k add $-3^k (F(t)/3^k \bmod 3)$ to t_3. This gives the unique compatible roots at every precision, hence the unique root in $\mathbb{Z}_3$. On this graph (E.37.7) becomes $t_1 + t_2 \equiv c_f$ modulo three. The promised complete parametrization is

$$\begin{aligned} u, w, a_{37}, \dots, a_{44} \in \mathbb{Z}_3, \qquad t_2 = u, \quad t_1 = c_f - u + 3w, \\ t_3 = \text{the unique root of } F(t_1, u, t_3) = 0, \qquad Y_{37}, \dots, Y_{44} = a_{37}, \dots, a_{44}, \end{aligned} \tag{E.37.13}$$

with the remaining coordinates given by (E.37.12). This is a bijective continuous coordinate map with continuous inverse: recover $u = t_2$, $w = (t_1 + t_2 - c_f)/3$, and the eight free Smith coordinates. It proves the stated fibre $\mathbb{Z}_3^{10}$ with an actual parametrization.

We now compare it exactly with the finite fibres. Modulo 3^n, the equation $F(t) = 0$ has one t_3 for each (t_1, t_2). Condition (E.37.7) leaves 3^{2n-1} abelian targets. For each target, (E.37.8) gives an affine central kernel

$$(\mathbb{Z}/3^n)^8 \times (\mathbb{Z}/3)^2. \tag{E.37.14}$$

Its order is 3^{8n+2}, proving (E.37.2). The two extra finite central coordinates are precisely controlled: a finite solution lifts to the next level exactly when

$$Y_{35} = -G(t), \qquad Y_{36} = t_1 - t_3 \pmod{3^n}. \tag{E.37.15}$$

For necessity, apply (E.37.10) at the next precision and divide its two Smith equations by three. For sufficiency, lift t_1, t_2 arbitrarily to $\mathbb{Z}_3$, take the unique root t_3, and use (E.37.12) with arbitrary lifts of the last eight coordinates. The first thirty-five coordinates match the given finite solution by their unit equations, and the next two match by (E.37.15). The affine source residue is unchanged. This proves lifting through the entire tower, not just the next level. It also gives exactly $3^{2n-1}3^{8n} = 3^{10n-1}$ images.

Here is an individual failed lift. Use E.24's source point with indexed CM row 0, ternary parameters $(0, 0)$ and binary parameter 0; its residue c_f is zero. In the boundary quotient P_3 prescribe $v = 0$, and in the doubled-log Smith coordinates prescribe

$$Y_{36} = 1, \qquad Y_i = 0 \quad (i \neq 36). \tag{E.37.16}$$

Its two torsion coordinates are $(Y_{35}, Y_{36}) = (0, 1)$. The corresponding BCH central coordinate in the original E.33 basis is

$$\begin{aligned} z = {} & e_{14} - e_{15} + e_{16} + e_{19} + e_{1,10} - e_{25} - e_{27} + e_{28} - e_{34} \\ & + e_{36} + e_{37} + e_{38} + e_{45} + e_{46} + e_{47} + e_{67} - e_{78}, \qquad e_{ij} = e_i \wedge e_j, \end{aligned} \tag{E.37.17}$$

with coefficients in $\mathbf{F}_3$. It satisfies both order relations: its Smith error is a multiple of three. The source fibre is realized by an actual commutator word. The certificate expresses $f^{-1}(0, z)$ in the complete 51-vector kernel basis of E.33, retains every basis relator preimage, and gives the ordered powers and the collecting commutator corrections. Replacing those relators by their fourth powers retains the prescribed modulo-three values and clears the binary source. Thus this is a concrete point of the actual common source, rather than a tuple in an unrelated coordinate group.

Any commutator-shadow lift to P_9 would have $t_1 = t_2 = t_3 = 0$ modulo three, since its abelian reduction is zero and the basis (E.37.5) is primitive. Its second divided Smith equation would force

$$Y_{36} = t_1 - t_3 = 0 \pmod{3}, \tag{E.37.18}$$

contrary to (E.37.16). This excludes the multiplier-one class-two lift at the specified next quotient. Its abelian-tower extensions remain exactly those already constructed in E.33 and E.36.

We finish the exact realizability and generation comparisons for the surviving full tower. The affine abelian values in (E.37.5)–(E.37.7) are realized by the eight source words for K, with three-supported exponents. The forty-five central preimages proving (E.37.4) then realize the prescribed central values in (E.37.12). These are finite products of continuous powers of actual source words, so give one element of $\widehat{F}'$ at every precision. Use squared words before these powers to clear the binary source. The same supported-exponent construction retains the two-adic and prime-to-six abelian choices of E.31. For a prescribed exponent-four solution, the sixty zero-abelianization words in E.36 give all its remaining central adjustments. Cube them to kill the ternary source and use two-supported powers, dividing their coefficients by the odd unit three. They act trivially in every three-primary class-two quotient. This proves simultaneous realization with that exponent-four solution and with every abelian modulus.

Every finite three-primary boundary map just constructed is an automorphism of P_{3^n}. It preserves H and hence descends to the characteristic quotient; its action on the geometric Frattini quotient has rank ten, as already checked for all the admissible modulo-three vectors

in E.33. The Frattini criterion gives surjectivity on the finite geometric three-group and then on P_{3^n}. For common sources with $\mathscr{D}$, the exponent-four quotient and abelian two-three moduli, use the exact nilpotent-residual/Frattini argument in E.36: the boundary factors have genuine automorphisms, their common N_B image has a finite lower-central stable term, and the $\mathscr{D}$ image kills $\gamma_5(N_B)$. Quotienting the resulting pure stable factor leaves the two rank-97 Frattini quotients through G_6. This proves joint generation. The prime-to-six abelian factors are added by the same actual CRT and coprime H-image comparison. No invariance of the fixed Ferrari source kernel is used.

The exact integral action, Smith matrices, all polynomial coefficients, every source affine residue, finite counts and the individual word certificate are saved in `bc_fermat_class_two_integral_model.py`, `bc_fermat_class_two_3adic_norms.py`, and `bc_fermat_class_two_3adic_tower.py`, with their executed outputs, JSON and NPZ certificates. All end in `PASS`. □

Statement E.38 (Proved). The complete double-cover Ferrari refinement retains every fibre. *Put $L = \mathrm{SL}_2(\mathbb{F}_5)$ and retain all epimorphism kernels of the free Ferrari group onto L. The resulting universal free image is*

$$\mathcal{L} = L^{19} \times \mathbb{F}_2^2. \tag{E.38.1}$$

The literal Ferrari action gives a finite quotient $B_4 \twoheadrightarrow T_L$. Inducing its six evaluation maps through the six cosets of $F \lhd B$ gives an actual finite quotient $\rho_L : B \twoheadrightarrow G_L$. Its common refinements with E.24 have the exact descriptions

$$D_{23} \times_{H_L} G_L, \qquad T_{23} \times_{S_L} T_L, \qquad |H_L| = 1728, \quad |S_L| = 96. \tag{E.38.2}$$

Here H_L is the specified quotient of G_6, and S_L is the maximal solvable quotient of T_L with its specified map from the binary degree-two Ferrari target.

Every one of the 864 *finite points of E.24 has a correction in N' preserving its complete D_{23} coordinate and satisfying the source hexagons, the full pentagon, the braid relations and the full-twist identities in these common refinements. The source and four-strand maps are surjective. This holds for each compatible profinite unit multiplier: its residue modulo* 144 *selects the corresponding E.24 branch. Thus every retained CM fibre, including every extra fibre, survives this central-cover test.*

Proof. We give the universal comparison before the correction calculation. The general construction by generating-pair orbits is described in [Gui15, §2.1, Lemma 2.1]; the following identifies its actual image in this case.

The universal free quotient and its action. The center of L is $Z = \{I, -I\}$ and $L/Z = A_5$. All 120 matrices, their products, their center, and the equality $L' = L$ are checked directly. The action of L/Z on its five Sylow 2-subgroups identifies it with the specific A_5 model of E.29. Conjugation by $\mathrm{PGL}_2(\mathbb{F}_5)$ realizes all 120 automorphisms of that model. These are also all automorphisms of L: an automorphism trivial on L/Z has the form $g \mapsto g\epsilon(g)$ with $\epsilon : L \to Z$ a homomorphism, which is trivial because L is perfect.

Each of the 19 epimorphism kernels onto A_5 has four lifts onto L, obtained by the two independent signs of its generator images. Every lifted pair generates L. Indeed, its image J has $JZ = L$, and $L = [JZ, JZ] = [J, J] \subset J$. The four kernels are distinct, since an automorphism identifying two lifts would be trivial on A_5 and hence trivial on L. Conversely every epimorphism onto L has one of these reductions. There are therefore exactly 76 kernels.

Choose one lift (g_j, h_j) over each of the 19 A_5 pairs. The joint image J in L^{19} surjects onto A_5^{19} by E.29, so $JZ^{19} = L^{19}$. Perfectness again gives $J = L^{19}$. Adjoining the two free abelianization coordinates modulo 2 gives the full product (E.38.1): a common quotient of a perfect group and an abelian group is trivial. All 76 evaluations are recovered by the actual formula

$$\theta_{j,e_1,e_2}(w) = (-I)^{e_1 \operatorname{ab}_1(w) + e_2 \operatorname{ab}_2(w)} \theta_j(w). \tag{E.38.3}$$

Conversely the chosen 19 values recover L^{19}, and ratios with the sign-twisted values recover both parity coordinates. Thus this formula gives an isomorphism from $\mathcal{L}$ onto the actual joint image in L^{76}.

Write a Nielsen automorphism as (p, c, D, M), acting by

$$(g, t) \longmapsto (((-I)^{D_j t} c_j g_{p(j)} c_j^{-1})_j, Mt), \quad D \in \mathrm{Mat}_{19\times 2}(\mathbb{F}_2), \quad M \in \mathrm{GL}_2(\mathbb{F}_2). \tag{E.38.4}$$

Here $c_j \in \mathrm{PGL}_2(\mathbb{F}_5)$ acts by the actual matrix conjugation. For the substitutions $a : (x, y) \mapsto (x, yx)$ and $b : (x, y) \mapsto (xy^{-1}, y)$, the permutation and the c_j values are those of E.29. Their two lifted generator images determine every entry of D, while the matrices M are respectively $\left[\begin{smallmatrix} 1 & 1 \\ 0 & 1 \end{smallmatrix}\right]$ and $\left[\begin{smallmatrix} 1 & 0 \\ 1 & 1 \end{smallmatrix}\right]$. Composition is the literal rule

$$(p, c, D, M)(q, d, E, R) = (q \circ p, (c_j d_{p(j)})_j, DR + E \circ p, MR). \tag{E.38.5}$$

For the semidirect product, the free coordinate multiplies as $k_j(-I)^{D_j u} c_j l_{p(j)} c_j^{-1}$, and the parity coordinate as $t + Mu$. These formulas retain the possible parity-to-center cross terms. The three braid generators are $(1, a), (1, b), (\xi_1, a)$ as in E.29; the literal Ferrari identities and every braid relation are checked in this model.

The two exact projections are to T_A and to $\mathrm{AGL}_2(\mathbb{F}_2) = S_4$, the latter given by (M, t). Their joint image is $T_A \times_{C_2} S_4$: E.29 identifies the maximal solvable quotient of T_A with braid parity, and Goursat's lemma applies to these specified generator maps. Above its identity, the 19 free central signs and 38 cross-term signs add in $\mathbb{F}_2^{57}$. The actual kernel has dimension 55, consisting of the 19 free signs and the computed 36-dimensional Nielsen kernel. In particular

$$|T_L| = 2^{117} 3^{36} 5^{31} 7^2. \tag{E.38.6}$$

The complete induced source and correction groups. For the six literal evaluation maps $\theta_i : F \to T_L$, use the same section of B/F as in E.29. The induced component is

$$(q, i) \longmapsto \theta_i(b_q w b_{qw}^{-1}), \tag{E.38.7}$$

together with the permutation $q \mapsto qw$. This defines the homomorphism ρ_L of discrete groups and its continuous extension from $\widehat{B}$. Its kernel is normal in B and lies in F. The identity-coset coordinate on $\rho_L(F)$ is the specified map to each evaluation target. Projection to E.29 gives a surjection $\pi : G_L \to G_A$ with elementary abelian kernel of dimension 44.

Set $P = \rho_L(N)$, $C = \rho_L(N')$ and $E = \rho_L(F')$. The exact image calculations give

group	order	$\dim_{\mathbb{F}_2} \ker(\pi\vert_{\text{group}})$
G_L	$2^{82}3^{26}5^{21}7^2$	44
P	$2^{76}3^{24}5^{21}7^2$	41
C	$2^{76}3^{23}5^{19}7^2$	41
C'	$2^{76}3^{23}5^{19}7^2$	41
E	$2^{79}3^{23}5^{19}7^2$	42

(E.38.8)

We describe completeness of the computation. Above a faithful permutation quotient, each stabilizer-chain Schreier generator is rewritten through the next transversals. Its discrepancy

is evaluated in the actual elementary kernel. All such discrepancies, with their normal conjugates, give the kernel of the lifted group: successive Schreier rewriting reduces every identity-base word to their normal closure. Every retained generator is an actual relator, recorded in a word DAG which is independently substituted at the end of the program.

For P, all 97 original N generators are retained, including the 91 differences above the six selected base generators. To compute the derived group of $\langle g_i, V\rangle$, where V is the complete elementary kernel, we retain every $[g_i, g_j]$, every $[v, g_i]$ for a basis of V, every conjugation residual of the selected base-derived generators, and the complete lifted base presentation. Their normal closure is exactly the derived group: the resulting quotient is abelian, and every retained word is a commutator or its conjugate. Applying this construction twice proves $C' = C$. Module commutators alone have rank 34 in its 41-dimensional kernel; the remaining seven directions are retained by the full second-derived calculation. Exhaustive calculations in A_4 and in the order-eight Heisenberg group independently check the lifted-presentation procedure, including an extra lift and its normal closure.

The equality of the two 41-dimensional kernels in (E.38.8) proves

$$\pi^{-1}(C_A) \cap P = C, \qquad P/C \simeq P_A/C_A. \tag{E.38.9}$$

Thus E.29's complete correction calculation transfers through this displayed isomorphism: with $K = \rho_L(K_6)$,

$$|P/C| = 75, \quad |K/C| = 25, \quad \rho_L(K_6 \cap F') = C, \quad |G_L/K| = 1728. \tag{E.38.10}$$

For the third equality, the projection of $K_6 \cap F'$ lies in C_A, hence its new image lies in C by (E.38.9); the reverse inclusion follows from $N' \subset K_6 \cap F'$.

The signed modular comparison. Use the literal map

$$q_{12} : B \longrightarrow Q = \mathrm{SL}_2(\mathbb{Z}/12)/\{\pm I\}, \qquad s \longmapsto \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \quad r \longmapsto \begin{bmatrix} 0 & -1 \\ 1 & -1 \end{bmatrix}. \tag{E.38.11}$$

All 97 defining N words have identity image, and $[G_L : P] = 576 = |Q| = [B : N]$. Since N is contained in the preimage of P, the equality of indices proves that this preimage is exactly N. Thus the displayed map descends to G_L, with kernel P.

In $\mathrm{SL}_2(\mathbb{Z}/12)$ the actual free generators are $x = \left[\begin{smallmatrix} 1 & 2 \\ 0 & 1 \end{smallmatrix}\right]$ and $y = \left[\begin{smallmatrix} 1 & 0 \\ -2 & 1 \end{smallmatrix}\right]$. They generate a group of order 96 whose derived subgroup has order eight. Every matrix of that derived subgroup is I modulo four. Consequently its projective image has a unique representative with that property, and its reduction modulo three is the quaternion subgroup Q_8 of $\mathrm{SL}_2(\mathbb{F}_3)$.

There is also an independent proof of the last row's quotient comparison in (E.38.8). E.29 gives $E_A \cap P_A = C_A$. Combining this equality with (E.38.9) shows $E \cap P = C$. The explicit modular calculation therefore gives

$$E/C \xrightarrow{\sim} q_{12}(E) = Q_8. \tag{E.38.12}$$

The direct full F' computation agrees, with kernel dimension 42.

The 24 retained E.17 words have the following three signed values, including their actual SL_2 lifts:

$$M_0 = I, \qquad M_1 = \begin{bmatrix} 1 & 4 \\ 4 & 5 \end{bmatrix}, \qquad M_2 = \begin{bmatrix} 1 & 8 \\ 8 & 5 \end{bmatrix} \pmod{12}. \tag{E.38.13}$$

Their complete index table is (E.29.13). The modular program evaluates every one of the 24 old correction words and all six ternary and seven binary fibre-basis words. Every value is identity in Q; it also checks the complete base expression with its actual integral cusp gauge. Hence all 864 reconstructed words have exactly the indexed value in (E.38.13). This proves the whole-fibre assertion through the word reconstruction formula.

Arithmetic selection and a common correction. Fix a compatible profinite unit λ. Its old residue is 1 or 73 modulo 144, so $\lambda \equiv 1 \pmod{12}$. Cyclotomic surjectivity supplies an actual Galois element with character λ. As proved in E.29, $\mathbb{Q}(\zeta_3, \sqrt[3]{2}) \cap \mathbb{Q}(\mu_\infty) = \mathbb{Q}(\zeta_3)$. Multiplication by an element of the cyclotomic kernel therefore makes its Kummer value $\rho_2 \bmod 3$ any prescribed κ. Nakamura–Schneps [NS00, Theorem 2.5], applied to this actual Galois element, gives exactly $q_{12}(f_\sigma) = M_\kappa$. The sign in that theorem is positive because $\lambda \equiv 1 \pmod 4$. No extension of that arithmetic matrix formula to arbitrary GT elements is used.

For a retained word $f_0 \in F'$ choose the matching κ. By (E.38.12),

$$c = \rho_L(f_0)^{-1}\rho_L(f_\sigma) \in C. \tag{E.38.14}$$

The image of N' in D_{23} is solvable. Its joint image with the perfect group C is the full product by Goursat's lemma, since a perfect solvable quotient is trivial. There is therefore a discrete $k \in N'$ with these two images equal to c and 1. The word $f = f_0 k$ has exactly its old D_{23} coordinate and exactly the G_L coordinate of f_σ. It remains in F'. All 36 induced coordinates match, so the six target evaluations, both source hexagons, the full pentagon and the braid and twist equations agree with the actual Galois action in the new component. The old component retains its E.24 equations. The braid convention is the exact Nakamura–Schneps (1.1) comparison proved at the end of E.29, including the sixth pair.

The multiplier comparison includes every power in these assertions. In D_{23} the cusp t has order 144 and the three conjugate peripheral generators x, y, z have order 72. Thus the residue of λ modulo 144 fixes both the required cusp power and every power with $m = (\lambda-1)/2$. The old four-strand generators have orders dividing eight and nine, and their full-twist periods were retained in E.24. The outer permutation quotient is the same S_3. In the new component we use the very same profinite λ as the actual Galois element, so no further power replacement is needed. Explicit representatives for the two finite branches are 1 and the unit η_{73} whose p-adic component is 73 for $p = 2, 3, 5, 7$ and 1 otherwise. All the finite groups here have only those prime divisors, so η_{73} and $(\eta_{73} - 1)/2$ give exactly the integer powers 73 and 36 in every component.

Joint generation. Set $H_L = G_L/K$. Its map from B factors through G_6, and hence through D_{23}. Every common quotient of D_{23} and G_L is solvable and has only the prime divisors two and three. It kills C by perfectness, then kills the normal group K/C of order 25. Every common quotient therefore factors through the specified H_L. This proves the first fibre product in (E.38.2) by Goursat's lemma. The corrected source map surjects onto D_{23} by E.24 and onto G_L because its generator images agree with an actual Galois automorphism followed by ρ_L. Its joint image lies in the stated fibre product and has both surjective projections; the same maximal-common-quotient argument proves its full surjectivity.

For the four-strand comparison, first kill the explicit perfect normal subgroup L^{19} in T_L. The remaining group $\mathbb{F}_2^2 \rtimes A_L$, where A_L is the Nielsen image, has a faithful 251-point representation. Its successive derived indices are $1, 2, 6, 96, 96$. The terminal subgroup is perfect; its inverse image in T_L is perfect as an extension of perfect groups. Thus the maximal solvable quotient S_L has order 96. Its full right Cayley table and literal representative braid words are retained. Exhaustively traversing all 1536 elements of the actual binary degree-two target, and checking all three generator edges at each element, proves the surjection

$$T_{2,2} \twoheadrightarrow S_L \tag{E.38.15}$$

with the specified values of a, b, c. Since T_{23} is solvable and maps to this quotient through $T_{2,2}$, the second fibre product in (E.38.2) is the actual joint four-strand target. The old

corrected triple generates T_{23}; the new triple generates T_L by its equality with a Galois image. The maximal common quotient is S_L, so these triples jointly generate the whole fibre product. Its kernel in B_4 is normal and lies in PB_4, since the old binary target already retains the standard S_4 quotient.

The executed files have prefix `de_ferrari_sl2_5_` and suffixes `model`, `nielsen`, `source`, `N`, `derived`, `coinvariants`, `modular`, and `solvable`; the two lifted-presentation modules and their exhaustive control are retained as well. Every image and whole correction group used above is a complete finite calculation. The particular correction in (E.38.14) is supplied by the proved arithmetic and product-image argument; no uncomputed literal Galois word is claimed as a saved certificate. These comparisons establish the stated finite refinements. They do not provide nonempty fibres over every finite profinite quotient. □

Statement E.39 (Proved). Full periodic-braid conjugacy, including the order-four test. *Let $n \geq 3$, and use the actual based braid action* (E.30.5c) *of $F = (\lambda, f) \in \widehat{\mathrm{GT}}$. In $\widehat{B}_n$ put*

$$\delta_n = \sigma_1 \cdots \sigma_{n-1}, \qquad \varepsilon_n = \sigma_1 \delta_n, \qquad z_n = \delta_n^n. \tag{E.39.1}$$

There are $h_\delta, h_\varepsilon \in \widehat{B}_n$ such that

$$F(\delta_n) = h_\delta \delta_n^\lambda h_\delta^{-1}, \qquad F(\varepsilon_n) = h_\varepsilon \varepsilon_n^\lambda h_\varepsilon^{-1}. \tag{E.39.2}$$

The same assertion holds for every discrete periodic braid, by its actual conjugacy to a power of one of these two elements.

In particular, for any finite quotient $q : \widehat{B}_4 \to T$, write $a = q(\sigma_1)$, $b = q(\sigma_2)$, $c = q(\sigma_3)$ and $a' = qF(\sigma_1)$, $b' = qF(\sigma_2)$, $c' = qF(\sigma_3)$. Then the following conjugacies hold in T itself:

$$a'b'c' \sim_T (abc)^\lambda, \qquad a'^2 b'c' \sim_T (a^2 bc)^\lambda. \tag{E.39.3}$$

This gives a necessary full-GT test on each actual finite shadow. The first equation includes the order-four periodic element in the central quotient; it also implies the order-two test after squaring. No invariance of $\ker q$ *under a finite shadow is assumed.*

Proof. We specify both completion and orientation comparisons. Cap the boundary of the n-punctured disk by a disk whose center is the additional marked point $n+1$. Extension of homeomorphisms by the identity gives

$$B_n / \langle z_n \rangle \xrightarrow{\sim} \mathrm{Stab}_{\Gamma_{0,[n+1]}}(n+1). \tag{E.39.4}$$

The kernel before taking the quotient is the integer boundary-twist group, generated by the full twist z_n. The stabilizer has index $n+1$. Its intrinsic profinite topology agrees with the topology induced from the full mapping class group: for each finite-index subgroup of the stabilizer, its core in the full group has finite index and is contained in it. Thus completion gives the corresponding isomorphism onto the actual open stabilizer in $\widehat{\Gamma}_{0,[n+1]}$. The kernel of $\widehat{B}_n$ onto the completed quotient is the closure of $\langle z_n \rangle$. The exponent-sum map $B_n \to \mathbb{Z}$ sends z_n to $n(n-1)$; its continuous extension proves that this closure is isomorphic to $\widehat{\mathbb{Z}}$. Consequently

$$\widehat{B}_n / \langle z_n \rangle \xrightarrow{\sim} \mathrm{Stab}_{\widehat{\Gamma}_{0,[n+1]}}(n+1) \tag{E.39.5}$$

uses the actual closed procyclic center.

The capping map is equivariant for the specified action. Precisely, apply the generator formulas (1.1) in Nakamura–Schneps [NS00] along $\widehat{B}_n \to \widehat{B}_{n+1}$ and then kill the full twist and Hurwitz element in $\widehat{B}_{n+1}$. Their power formulas preserve both closed normal kernels,

as in the explicit $n = 4$ diagram in E.30. The resulting mapping-class action preserves the permutation of every generator, because λ is odd and the conjugating evaluations of f are pure. On $\widehat{B}_n^{\mathrm{ab}} = \widehat{\mathbb{Z}}$ it acts as multiplication by λ.

We use the precise finite-subgroup conclusion of Boggi [Bog26, Theorem C(i), equivalently Theorem 5.1(i) with empty boundary]. For the genus-zero surface with $n+1$ punctures, the natural inclusion of the discrete full mapping class group in its profinite completion induces a bijection on conjugacy classes of solvable finite subgroups. The surface is connected, oriented, hyperbolic and of finite type, and the subgroups used here are finite cyclic groups. Thus all the hypotheses are satisfied. Applying this assertion to the images of $\langle \delta_n \rangle$ and $\langle \varepsilon_n \rangle$ in the central quotient shows that their images under F are conjugate to discrete cyclic subgroups.

The discrete rotation classification [Col12, Theorem 3.1 and Corollary 3.2] then identifies the possible generators. In the $(n+1)$-punctured sphere, δ_n has order n and exactly one marked fixed pole, namely $n+1$. The element ε_n has order $n-1$ and exactly two marked fixed poles. The other marked points lie in the rotation orbits. Powers of these rotations with the same order have the same indicated fixed-point pattern, so the preserved permutation selects the corresponding family. We obtain conjugacies to δ_n^a and ε_n^b, where a is a unit modulo n and b is a unit modulo $n-1$.

For δ_n, the unique marked fixed pole forces the conjugating permutation to fix $n+1$. The conjugator therefore belongs to the stabilizer in (E.39.5). For ε_n, the conjugator preserves the set of the two fixed poles. In the rotation model whose marked points are $\{0, \infty\} \cup \mu_{n-1}$, the actual holomorphic map $w(t) = 1/t$ exchanges the poles and conjugates the rotation to its inverse. Compose with w when the poles are exchanged. The resulting conjugator fixes $n+1$, with exponent either b or $-b$; denote that exponent by b'. This operation supplies the required conjugator inside the stabilizer before determining its exponent.

The abelianization of the stabilizer is $\mathbb{Z}/n(n-1)$ by (E.39.4). The classes of δ_n and ε_n are $n-1$ and n, respectively. The actual action on this quotient therefore gives

$$(n-1)a = (n-1)\lambda \pmod{n(n-1)}, \qquad nb' = n\lambda \pmod{n(n-1)}. \tag{E.39.6}$$

Hence $a \equiv \lambda \pmod{n}$ and $b' \equiv \lambda \pmod{n-1}$. Both central-quotient conjugacies have the required exponent λ. Here $\varepsilon_n^{n-1} = z_n$ in B_n: its rotation has order $n-1$ in (E.39.4), so that power is a boundary-twist power; its exponent sum $n(n-1)$ makes the power exactly one.

Lift either stabilizer conjugator to $\widehat{B}_n$. For $\beta = \delta_n$ or ε_n the equality now has the exact form

$$F(\beta) = h\beta^{\lambda}h^{-1}z_n^k, \qquad k \in \widehat{\mathbb{Z}}. \tag{E.39.7}$$

Apply the full exponent-sum map to $\widehat{\mathbb{Z}}$. Both $F(\beta)$ and β^{λ} have exponent sum $\lambda \exp(\beta)$, whereas z_n^k has exponent sum $n(n-1)k$. Multiplication by the nonzero integer $n(n-1)$ is injective on $\widehat{\mathbb{Z}} = \prod_p \mathbb{Z}_p$. It follows that $k = 0$, proving (E.39.2). For completeness, a discrete periodic braid has a finite-order image under (E.39.4). Its rotation fixes the capped marked point. The other pole is either unmarked or marked; the rotation classification then gives a conjugate of a power of δ_n or ε_n, respectively, in that stabilizer. Lifting the conjugator leaves an integer power of z_n. The identities $z_n = \delta_n^n = \varepsilon_n^{n-1}$ absorb that power and give the asserted actual conjugacy in B_n. For a discrete conjugate $u\beta^j u^{-1}$, apply F and use the actual conjugator $F(u)hu^{-1}$ to obtain the same assertion. Finally applying the stated finite map q gives (E.39.3) with conjugators $q(h_\delta)$ and $q(h_\varepsilon)$. This last step uses the two homomorphisms q and qF from the same full source, and requires no induced endomorphism of the fixed finite quotient. □

Statement E.40 (Proved). All retained Ferrari points pass full periodic-braid conjugacy. *For every one of the* 864 *actual joint points of E.24, with multiplier residue* 1

or 73 *modulo* 144*, put*

$$d = abc, \qquad d' = a'b'c', \qquad z_4 = d^4 \quad \textit{in } T_{23} = T_{2,6} \times T_{3,6}. \tag{E.40.1}$$

There is an element $h \in T_{23}$ *with*

$$d' = hd^{\lambda}h^{-1}. \tag{E.40.2}$$

All compatible higher lifts of the displayed multiplier residue satisfy the same finite comparison. Together with E.35 this verifies both full periodic tests in E.39 at every retained point.

In the binary factor, the 96 *branch and line inputs give two distinct values of* d'*. For each of them,*

$$d'z_4^j \sim_{T_{2,6}} d \quad \Longleftrightarrow \quad j = 0 \text{ or } 2 \quad (j \in \mathbb{Z}/4). \tag{E.40.3}$$

Four literal braid conjugators of lengths from zero to 11088 *certify the positive comparisons. In the ternary factor the* 216 *points give fifteen distinct values of* d'*, with*

$$d'z_4^j \sim_{T_{3,6}} d \quad \Longleftrightarrow \quad j = 0 \quad (j \in \mathbb{Z}/9). \tag{E.40.4}$$

The fifteen saved conjugators have lengths from six to 131*. Every input and every positive identity has been checked by a separate literal word evaluator importing no conjugacy solver.*

Proof. We keep the complete finite target models and the actual source reconstruction maps of E.24. The new binary element has a nonidentity image in the actual quotient $T_{2,6} \to S_3$. Accordingly the conjugacy algorithm uses, for each of the six recorded coset representatives g, the necessary base equality

$$\overline{d'} = \overline{gdg^{-1}}. \tag{E.40.5}$$

Only the cosets satisfying this equality proceed to the kernel calculation. Above each such coset, the residual difference lies in the actual forty-dimensional collected two-group U.

The weight-by-weight centralizer method of E.32 then applies with d in place of d^2. At weight r, every element of the current centralizer centralizes the comparison modulo weight r. Its commutator with that comparison in the next elementary abelian layer is an actual homomorphism: conjugation by U acts trivially on that layer, so the product commutator identity becomes addition there. All the kernel basis vectors are retained and closed under the actual two-group multiplication. The computed subgroup orders verify the index of the next centralizer as exactly the image order of this homomorphism. Thus every possible conjugator above every matching base coset is retained. An inconsistent system eliminates the whole corresponding coset, and all six base cosets are checked.

As an exhaustive independent control, use the degree-two target, of order 1536, with the same nonidentity-base element *abc*. Enumerate all elements and all of its conjugates. The conjugacy class has 96 elements, and all 1536 decisions of the new algorithm agree with this complete enumeration.

At degree six, reconstruct each of the 24 original binary inputs, both multiplier branches, and both parameters in its full correction line. This gives all 96 inputs in the statement. Apply the complete algorithm to all four central translates of each distinct value. The outcome is exactly (E.40.3). The forty collected kernel-basis elements are rebuilt from the saved original braid-word expressions, including every square and commutator. Reducing each successful conjugator in this basis and adjoining its actual outer word produces the four literal witnesses in the statement. Substitution verifies each identity in the original finite group.

In the binary target, $|d| = 16$ and $|z_4| = 4$. For the actual multiplier residues,

$$d^{\lambda} = dz_4^{(\lambda-1)/4}, \qquad -\frac{\lambda-1}{4} \equiv \begin{cases} 0 & \lambda = 1, \\ 2 & \lambda = 73 \end{cases} \pmod 4. \tag{E.40.6}$$

Both required central exponents occur in (E.40.3), and the literal evaluator checks $d' = h_2 d^{\lambda} h_2^{-1}$ directly for each of the 96 inputs.

For the ternary target use the complete class-two exponent-three kernel over the 648-element quotient as in E.35. The two successive conjugacy equations retain the full homogeneous centralizer spaces, now computed for $d = abc$. All base conjugators in the 648-element group are enumerated. The exact logarithm is applied only to the already proved class-two kernel. Reconstruction of all nine parameters in each of the 24 original planes gives all 216 inputs. The complete calculation gives (E.40.4) and the fifteen literal conjugators. Here $|d| = 36$ and $|z_4| = 9$. Both multiplier residues, and all their higher lifts modulo 144, are one modulo 36. Hence the saved identities give $d' = h_3 d^{\lambda} h_3^{-1}$ in this factor.

The separate evaluator imports the two group models and source substitution maps, but neither conjugacy algorithm. It evaluates all nineteen literal conjugators, reconstructs all 96 binary and 216 ternary inputs, and checks the full multiplier equality at every point. The opposite-prime corrections have identity images in the respective target factors by E.24's explicit power maps. Finally the actual joint target is the full direct product proved in E.24. Thus (h_2, h_3) is an element of that target and gives (E.40.2) for all 864 points.

The exact algorithm and exhaustive control are `e_quarter_torsion_root.py`, executed in its three modes `control`, `binary`, and `ternary`; the separate evaluator is `e_quarter_torsion_words_root.py`. Their outputs and complete indexed JSON certificates are retained. The finite outcomes keep every original CM fibre. □

Statement E.41 (Proved). The complete two-adic class-two fibre and its exact finite lifting images. *Put*

$$P_{2^n} = B/(\gamma_3(H)H^{2^n}), \qquad \mathscr{J}_{2^n} = \mathrm{im}(B \longrightarrow \mathscr{D} \times P_{2^n}). \tag{E.41.1}$$

Fix any one of the 432 *multiplier-one source words of E.24. At exponent four its complete correction-image fibre has* 2^{10} *elements, and every one lifts through the whole two-adic class-two tower. For* $n \geq 3$ *the complete fibre at* P_{2^n} *has*

$$2^{10n-9} \tag{E.41.2}$$

elements. Exactly 2^{10n-10} *lift to the next level, and this is also exactly the subset lifting through every level. The criterion is the single additional congruence in* (E.41.22) *below.*

The full inverse-limit fibre has an explicit homeomorphism with $\mathbb{Z}_2^{10}$. *Its affine source congruences, integral Hall parity and eight central parameters are retained in* (E.41.15)–(E.41.20). *It can be prescribed simultaneously with any full three-adic fibre from E.37, by one profinite commutator word. Every finite common shadow so obtained is generating.*

There is an individually specified exponent-eight shadow over source row 0, *parameters* $(0, 0; 0)$, *which has no multiplier-one commutator lift to* P_{16}. *Its actual source-word construction and Hall coordinates are given in* (E.41.23)–(E.41.26).

Proof. We first make the integral coordinate comparison. Let the ten ordered Schreier generators of H be the ones fixed in E.33. For Hall coordinates (v, z), with central basis $[\eta_i, \eta_j]$ ordered by $i < j$, define

$$Z_{ij} = 2z_{ij} + v_i v_j. \tag{E.41.3}$$

This is an injective map into doubled-log coordinates, with image exactly $Z_{ij} \equiv v_i v_j \pmod 2$. Its inverse on that image is $z_{ij} = (Z_{ij} - v_i v_j)/2$. The actual multiplication becomes

$$(v, Z)(w, T) = (v + w, Z + T + v \wedge w). \tag{E.41.4}$$

The signed Schreier substitutions for s, r give the same integral matrices S, R as E.37, with $S^2 = R^3 = 1$ over $\mathbb{Z}$. In particular they preserve the displayed parity subset because they are collected from actual free-group words.

In the exponent-2^n class-two quotient, Hall abelian coordinates are modulo 2^n and Hall central coordinates are modulo 2^{n-1}. Indeed $[x, y]^{2^n} = 1$ follows from commutator collection, and the power relation for $(xy)^{2^n}$ also gives $[x, y]^{\binom{2^n}{2}} = 1$; their two-adic greatest common divisor is 2^{n-1}. Conversely the Hall group with exactly these coordinate moduli satisfies every 2^n-th power relation, by the same power formula. These two actual homomorphisms are inverse. Equivalently, both v, Z may be reduced modulo 2^n while retaining (E.41.3). Thus the Hall central precision is one power of two below the abelian precision.

The complete source image and actual word preimages. Write R_2 for the kernel on $K_6 \cap \widehat{F}'$ of the full binary source quotient in E.24. Its 5458 normal relators were retained in E.36, including the 95 primitive integral-F-abelianization relations. Their integral H-abelian image is precisely the eight-column lattice K of E.31. The new integral collection checks all 5458 words against their previous exponent-four values. Twelve actual relators, with indices

$$3, 4, 96, 97, 98, 100, 103, 104, 105, 4656, 4658, 4662, \tag{E.41.5}$$

give ordered-product words $g_1, \ldots, g_8$ whose abelian vectors are exactly the columns of K. The saved integer combination matrix has largest absolute coefficient 68.

For each source relator ρ, replace it by $\rho(\prod_i g_i^{c_i})^{-1}$, where $[\rho]_H = Kc$. Also retain the similarly adjusted s, r conjugates of the eight g_i and all 28 commutators $[g_i, g_j]$. All 5502 resulting words have zero integral H-abelianization. Their central images, closed under the actual exterior s, r action, form a lattice $C \subset \mathbb{Z}^{45}$ with

$$\mathrm{Smith}(C) = (1^{39}, 2^4, 4^2). \tag{E.41.6}$$

This is the full central image. To check completeness, quotient the normal source presentation by these adjusted words. Every source relator becomes an ordered product of the g_i; their conjugates remain such products, and the commutator relations make that quotient abelian. Their eight independent integral abelian vectors then identify the quotient with K. Hence the kernel of its abelian map is exactly the normal closure just collected. In the class-two image, conjugation acts on this kernel by the displayed exterior matrices, which is precisely the closure used to compute C.

This calculation also retains actual central word preimages. Forty-five selected adjusted relators $w_1, \ldots, w_{45}$ already have central matrix W with

$$W = CA, \qquad A \in M_{45}(\mathbb{Z}), \qquad |\det A| = 631022156534756593832815423865613. \tag{E.41.7}$$

The determinant is odd. Therefore these literal zero-H-abelianization words give a basis of $C \otimes \mathbb{Z}_2$. The certificate specifies every adjusted relator and its ordered g_i correction, and checks its integral central vector. It does not infer word preimages merely from the rational rank.

The same two-adic image is obtained from the kernel of the entire source $\mathscr{D}$. Every word used above lies in K_6, whose ternary source image has exponent three. Cubing clears that image, and multiplication by three is invertible in $\mathbb{Z}_2$. Equivalently use continuous two-supported exponents on these actual words. Thus the complete two-adic source image is described as follows: choose $c \in \mathbb{Z}_2^8$, form the actual ordered product $\prod_i g_i^{c_i}$ in the class-two pro-two group, and then multiply by an arbitrary central element of $C \otimes \mathbb{Z}_2$ in Hall coordinates. This is a closed image, since it is the image of the compact coefficient space by the displayed continuous word map.

We also need the exact finite fibre over an abelian value. It is the reduction of C, not a larger uncomputed kernel. In the chosen K basis, g_1, g_2 have abelian vectors $12e_1, 12e_2$. The remaining six columns have a unit triangular minor. Therefore the coefficient ambiguities modulo 2^n are generated by $2^{n-2}g_1$, $2^{n-2}g_2$ and 2^n multiples of the other coordinates. The Hall central values of g_1, g_2 belong to $C + 2\mathbb{Z}^{45}$, as the saved literal parity calculation verifies. Their 2^{n-2} powers consequently have central values in C modulo 2^{n-1}. The 2^n powers and all collecting terms involving an abelian multiple of 2^n vanish at that central precision, by the Hall power formula. This proves the finite comparison for every $n \geq 2$.

The full central norm and its finite kernels. Let M be the 90-by-45 central norm matrix from E.37; its saved integral decomposition is

$$S_M = U_M M V_M, \qquad \operatorname{diag}(S_M) = (1^{35}, 3, 3, 0^8). \tag{E.41.8}$$

It is saturated over $\mathbb{Z}_2$. The complete source norm has

$$\operatorname{Smith}(MC) = (1^{31}, 2^4, 12^2, 0^8). \tag{E.41.9}$$

Put $N_0 = (V_M)_{\{37,\dots,44\}}$, the last eight columns, with zero-based indexing. The exact certificate verifies

$$MN_0 = 0, \qquad C^{-1}N_0 \in M_{45,8}(\mathbb{Z}). \tag{E.41.10}$$

The columns of N_0 are primitive in the ambient central lattice, since V_M is unimodular. They therefore give all integral central norm-kernel parameters inside the source lattice.

For central modulus 2^m, the Smith factors give respectively

$$\begin{aligned} |C \bmod 2^m| &= 2^{39m+4\max(m-1,0)+2\max(m-2,0)}, \\ |MC \bmod 2^m| &= 2^{31m+4\max(m-1,0)+2\max(m-2,0)}. \end{aligned} \tag{E.41.11}$$

Thus the kernel on correction images has 2^{8m} elements. The contained image $N_0(\mathbb{Z}/2^m)^8$ has that same order by primitivity. They are equal. In particular every finite homogeneous central correction image lifts compatibly through all levels.

Here is an exact small congruence map for the remaining affine source obstruction. If $\sigma_i = (S_M)_{ii}$ for $i < 37$, define

$$\begin{aligned} A_M &= -3(V_M)_{<37}\operatorname{diag}(\sigma_i^{-1})(U_M)_{<37}, \\ D_C &= 4C^{-1}A_M. \end{aligned} \tag{E.41.12}$$

Both matrices are integral, as verified entry by entry. On the saturated ambient norm image, $A_M h/3$ solves $Mz + h = 0$. Since three is a two-adic unit, that solution lies in $C \otimes \mathbb{Z}_2$ exactly when $D_C h = 0$ modulo four. All ambient homogeneous solutions differ by N_0, already contained in C, so this is the complete source condition. The identity $D_C MC = 0 \pmod 4$ is checked integrally. The same argument works modulo 2^m for $m \geq 2$: an ambient solution and this modulo-four condition give a source solution modulo 2^m. Exponent four was computed directly in E.36.

The source conic and all 432 origins. The exact abelian norm kernel is the primitive basis from E.37:

$$Bt = (-t_1, t_2, -t_1 + t_3, t_2, t_3, -t_3, t_1 - t_3, t_3, -t_1, t_1)^t. \tag{E.41.13}$$

Its defining equations, including $Lv = 0$, have seven unit Smith factors over $\mathbb{Z}$. They therefore have this same kernel modulo every 2^n. For the two source functionals λ_1, λ_2 in (E.37.6), the

full two-adic K-membership condition is divisibility by four of both. Their common kernel has the same index 2^4 as K in $\ker L$. Set

$$u = t_1 + t_2, \qquad v = t_3 - t_2, \qquad w = t_2, \qquad (t_1, t_2, t_3) = (u - w, w, v + w). \tag{E.41.14}$$

On Bt the two functionals are $10u - v - 12w$ and $u - v - 12w$. For the retained source word f, write $c_i = \lambda_i([f]) \pmod 4$. The exact affine condition is therefore

$$u \equiv c_1 - c_2 =: u_0 \pmod 4, \qquad v \equiv c_1 - 2c_2 =: v_0 \pmod 4. \tag{E.41.15}$$

Every integral source word is collected and compared with both the previous modulo-three BCH value and modulo-four Hall value. The resulting six residue pairs are

$$(0,0),\ (2,0),\ (0,1),\ (2,3),\ (3,1),\ (3,3), \tag{E.41.16}$$

each occurring 72 times. For CM rows $0, \ldots, 23$ the pairs, respectively, are the entries of the following array, read across rows; each is constant on all eighteen internal choices:

$$\begin{array}{cccc} (0,0) & (2,0) & (2,0) & (0,0) \\ (0,1) & (2,3) & (2,3) & (0,1) \\ (3,1) & (3,3) & (3,3) & (3,1) \\ (3,1) & (3,3) & (3,3) & (3,1) \\ (0,0) & (2,0) & (2,0) & (0,0) \\ (0,1) & (2,3) & (2,3) & (0,1) \end{array} \tag{E.41.17}$$

The rational-cokernel polynomial of E.37 becomes

$$F(u,v) = v + 3(u^2 + uv + v^2). \tag{E.41.18}$$

Its doubled-log zero-Smith rows are even integer multiples of F, with one exactly $2F$ and greatest common divisor of their multipliers equal to two. For an actual source reference g_0 with abelian value Bt, both norm words have zero integral abelianization and even doubled central values. Let h be those central values divided by two. The Hall zero-Smith rows are consequently integer multiples of F with greatest common divisor one. The exact ambient obstruction is $F = 0$; at exponent 2^n it is $F = 0$ modulo 2^{n-1}.

The remaining source congruence $D_C h = 0 \pmod 4$ is automatic on all the affine source lattices (E.41.15). This assertion is checked on the complete finite source, as follows. The homogeneous abelian source norm basis, in the t coordinates, is

$$\begin{pmatrix} 1 & 0 & 0 \\ 43 & -12 & 4 \\ -33 & 0 & -4 \end{pmatrix}. \tag{E.41.19}$$

Its first column is primitive and its last two columns divided by four complete a two-adic basis. Hence, above each of the 432 actual origins, its coefficients modulo $(8, 2, 2)$ enumerate exactly the 32 possible abelian norm values modulo eight. Using the actual g_i word lifts, the program forms all $432 \cdot 32$ reference values, computes their Hall errors h, and verifies $D_C h = 0$ modulo four in every case. It then constructs the source central correction and checks the two norm equations modulo eight. All 32 values pass over every source point; all resulting geometric Frattini ranks are ten.

This finite calculation proves the asserted full congruence because its only modulus is four. The finite source comparison above shows that any two actual reference lifts with the same abelian value modulo eight differ by a central element of C modulo four. Their norm errors

differ by MC, annihilated by D_C modulo four. Thus the finite test covers every two-adic reference value, including its central carries.

We can now give the full coordinate map. When u_0 is even, choose $u = u_0 + 4a$ and let v be the unique root of $F(u, v) = 0$ with $v \equiv v_0 \pmod 4$. When u_0 is odd, choose $v = v_0 + 4a$ and solve uniquely for $u \equiv u_0 \pmod 4$. The first case has $F_v = 1 + 3(u + 2v)$ odd; the second has $F_u = 3(2u + v)$ odd. All six initial pairs satisfy $F = 0$ modulo four, so the roots exist uniquely by the explicit successive-bit Hensel construction. Choose $w \in \mathbb{Z}_2$ arbitrarily and recover t by (E.41.14). Solve $Kc = Bt - [f]$ and form $g_0 = f \prod_i g_i^{c_i}$. Then all the central solutions, in Hall correction coordinates, are

$$z_{\mathrm{corr}} = A_M h/3 + N_0 b, \qquad b \in \mathbb{Z}_2^8. \tag{E.41.20}$$

The congruence just proved puts this vector in $C \otimes \mathbb{Z}_2$. Actual central preimages are supplied by W in (E.41.7), using $W^{-1} z_{\mathrm{corr}} \in \mathbb{Z}_2^{45}$. All Hall parities are retained because the construction starts from actual ordered source words and adds an actual central Hall vector. The parameters (a, w, b) range over $\mathbb{Z}_2^{10}$. Conversely recover a, w from the target abelian vector, subtract the displayed particular correction, and recover b using the inverse unimodular Smith matrix. These continuous inverse maps prove the claimed homeomorphism.

Exact finite images and the first failure. For $n \geq 3$, conic solutions modulo 2^{n-1} in a fixed residue disk modulo four have 2^{n-3} choices. Both coordinates then have a free top bit when reduced modulo 2^n, giving 2^{n-1} pairs (u, v). The independent w gives 2^{2n-1} abelian targets. Each has exactly $2^{8(n-1)}$ central correction images by (E.41.11). This proves (E.41.2). At $n = 2$ there are four abelian targets, and the central count is 2^8, as in E.36.

An abelian target modulo 2^n lifts to an exact conic point precisely when $F = 0$ modulo 2^n. The same condition is necessary at the next quotient, whose Hall central precision is 2^n, and sufficient by the unit derivative in the same residue disk. For $n \geq 2$, the number of such abelian targets is 2^{2n-2}. There is no additional central loss: choose an exact conic lift and one exact central solution; the discrepancy from any prescribed finite central solution belongs to the finite homogeneous kernel, which is exactly the reduction of N_0 by (E.41.11). Consequently

$$\#\{\text{images lifting through all levels}\} = 2^{10n-10} \qquad (n \geq 2), \tag{E.41.21}$$

and the full criterion is

$$F(u, v) \equiv 0 \pmod{2^n}. \tag{E.41.22}$$

At exponent four it holds for every image; at every subsequent level it selects exactly one half.

Here is an explicit failing image at exponent eight. Use source row 0, parameters $(0, 0; 0)$, and the abelian target

$$v_H = (0, 0, 4, 0, 4, 4, 4, 4, 0, 0) \pmod 8. \tag{E.41.23}$$

Its Hall central vector, in the ordered 45-coordinate wedge basis, is the concatenation of these three rows:

$$\begin{array}{ccccccccccccccc} 2 & 2 & 3 & 0 & 1 & 3 & 2 & 2 & 3 & 0 & 0 & 0 & 2 & 2 & 2 \\ 1 & 0 & 3 & 1 & 1 & 3 & 3 & 2 & 1 & 1 & 3 & 2 & 0 & 1 & 2 \\ 0 & 2 & 1 & 2 & 3 & 3 & 2 & 1 & 0 & 3 & 3 & 2 & 0 & 3 & 1 \end{array} \pmod 4. \tag{E.41.24}$$

It is an actual common-source correction. If f_0 is the retained literal source word, take

$$f_0 \prod_{i=1}^{8} g_i^{9c_i} \prod_{j=1}^{45} w_j^{9a_j}, \qquad c = (0, 3, 5, 1, 5, 3, 7, 1), \tag{E.41.25}$$

where the a_j are the concatenated rows

$$\begin{array}{cccccccccccccc} 2 & 0 & 0 & 2 & 3 & 1 & 3 & 3 & 1 & 3 & 1 & 2 & 1 & 3 & 3 \\ 3 & 0 & 0 & 3 & 1 & 2 & 1 & 2 & 2 & 0 & 3 & 0 & 0 & 1 & 2 \\ 3 & 0 & 3 & 3 & 1 & 2 & 2 & 3 & 0 & 0 & 0 & 1 & 3 & 3 & 2. \end{array} \tag{E.41.26}$$

The ninth powers clear every ternary source coordinate and preserve the exponent-eight image; all the words already belong to the binary source kernel. The literal coordinate check gives (E.41.23)–(E.41.24) and verifies both norm relations. Every lift to P_{16} would retain $(t_1, t_2, t_3) = (0, 0, 4)$ modulo eight, so $(u, v) = (0, 4)$ modulo eight. But $F(0, 4) = 52 \equiv 4 \pmod 8$, contradicting the necessary Hall zero-Smith row at P_{16}. This is the asserted failure for the fixed multiplier-one commutator shadow.

One word and joint generation. All two-adic source coordinates above are finite products of continuous two-supported powers of the actual g_i, w_j. They therefore give one element of $\widehat{F}'$ while clearing the full ternary source image. The three-supported construction in E.37 does the converse. The two corrections can be multiplied: each is one in every class-two quotient at the other's prime. This realizes any prescribed pair of full two-adic and three-adic fibres over the same retained source point.

The rank-ten tests imply that each finite two-adic boundary map is an automorphism of its actual P_{2^n} quotient, by the finite geometric Frattini criterion. It descends to this quotient because it preserves H and the characteristic subgroup $\gamma_3(H)H^{2^n}$. The same is true on the three-adic factors. Their common boundary quotient therefore has an automorphism: its kernel is the intersection of these preserved kernels, and injectivity can be checked in every factor. For the joint source with $\mathscr{D}$, apply the exact lower-central-stable-term argument in E.36. The $\mathscr{D}$ image kills $\gamma_5(N_B)$, while a sufficiently deep stable term of the common boundary N_B image occurs as a pure boundary factor in the corrected image. After quotienting it, the joint N_B image is finite nilpotent with primes two and three. Both Frattini quotients have rank 97, forced by the retained map through $(\mathbb{Z}/6)^{97}$. The inherited G_6 automorphism then gives surjectivity, and the quotient by N_B gives the whole common source. This proves the generation assertion without assuming kernel invariance for the fixed Ferrari source.

The new exact certificates have common prefix `bc_fermat_class_two_` and suffixes `2adic_source.py`, `integral_family.py`, `2adic_norms.py`, `2adic_fibres.py`, `2adic_words.py`, and `2adic_tower.py`, with their executed outputs and JSON/NPZ data. They retain the full source presentations, all 432 literal origins, every finite P8 comparison and the individual nonlifting word. All finish with `PASS`. □

Statement E.42 (Proved). A simultaneous extension to every full pro-prime boundary factor away from two and three. *Let $\widehat{H}$ be the index-54 subgroup of $\widehat{B}$ determined by the actual Fermat covering, and for each prime p put*

$$\mathscr{P}_p = \widehat{B}/\ker(\widehat{H} \longrightarrow \widehat{H}^{(p)}). \tag{E.42.1}$$

The kernel in (E.42.1) *is characteristic in $\widehat{H}$, hence normal in $\widehat{B}$. Thus this is the actual quotient extension*

$$1 \longrightarrow \widehat{H}^{(p)} \longrightarrow \mathscr{P}_p \longrightarrow B/H \longrightarrow 1. \tag{E.42.2}$$

Fix any of the 432 multiplier-one source points of E.24, any full two-adic class-two fibre of E.41 and any full three-adic class-two fibre of E.37. There is one profinite commutator word realizing all of these prescribed images and inducing the identity shadow on every $\mathscr{P}_p$, for every $p \geq 5$. It is generating in the actual common source of $\mathscr{D}$, all the prescribed two- and three-primary class-two quotients, and every finite collection of finite quotients of the

groups (E.42.1). *In particular all prime-power class-two boundary towers can be retained simultaneously.*

Proof. E.37 and E.41 give an actual word

$$f \in \widehat{H} \cap \widehat{F}' \tag{E.42.3}$$

with the stated two- and three-adic images and the fixed $\mathscr{D}$ image. Here (E.42.3) is an exact subgroup statement: the retained source word lies in F', every correction lies in $K_6 \cap \widehat{F}'$, and $\widehat{F}' \subset \widehat{H}$ by the defining abelian quotient of the Fermat cover.

For a prime p, let $e_p \in \widehat{\mathbb{Z}}$ have component 1 at p and 0 at all other primes. The continuous power map on the closed procyclic subgroup generated by f gives the explicit replacement

$$f^* = f^{\,e_2+e_3}. \tag{E.42.4}$$

It still belongs to the closed subgroup $\widehat{F}'$. Every finite retained source group $\mathscr{D}$ has order divisible only by two and three. If its image of f has order 2^a3^b, then $e_2 + e_3 \equiv 1 \pmod{2^a3^b}$. Thus (E.42.4) preserves that image exactly. The continuous power map on any pro-two or pro-three group only uses the corresponding component in $\mathbb{Z}_2$ or $\mathbb{Z}_3$. Both components here are 1, so all prescribed two- and three-adic images are unchanged. On $\widehat{H}^{(p)}$ for $p \geq 5$ the component of the exponent is 0. Consequently

$$f^*|_{\widehat{H}^{(p)}} = 1 \qquad (p \geq 5). \tag{E.42.5}$$

This uses the full pro-p group, without passing first to an abelian or class-two quotient.

The based multiplier-one source formula with conjugating word 1 sends the literal generators s, r to themselves. By (E.42.5), its evaluation in each $\mathscr{P}_p$ is therefore the identity map of that actual quotient. Both order relations and generation there follow from this explicit equality. The two- and three-primary relations are those already proved for f. Thus (E.42.4) gives compatible homomorphisms to every indicated finite common source.

We prove joint generation on those sources, retaining the actual H maps. Let G_{23} be any finite common source of $\mathscr{D}$ and chosen two- and three-primary boundary quotients. The exact inclusion $K_6 \subset H$ supplies its canonical onto map to B/H: indeed $N \subset F$, $N' \subset F' \subset H$, and $N^6 \subset F^3 \subset H$. Thus G_6, and hence G_{23}, has that quotient. Write A_{23} for the image of H, its kernel. All prime divisors of $|A_{23}|$ are two and three. The shadow on G_{23} is onto by E.41, and its induced map on B/H is the identity because its conjugating word lies in $\widehat{H}$. Hence its restriction to $\widehat{H}$ is onto A_{23}: a preimage of an element in that kernel must itself have trivial B/H image.

For a finite set of primes $p \geq 5$, choose finite characteristic p-group quotients A_p of $\widehat{H}^{(p)}$. Their invariant kernels give actual quotients G_p of $\mathscr{P}_p$ with kernel A_p and quotient B/H. These characteristic quotients are cofinal, since $\widehat{H}$ is finitely generated: intersect the finitely many automorphic images of any open normal subgroup. Repeated quotients at the same prime may first be combined by taking their actual common quotient, still a finite p-group in its geometric kernel.

The image of $\widehat{H}$ in $A_{23} \times \prod_{p\geq 5} A_p$ is the whole product. It surjects to every factor. A proper subdirect product of two factors would give nontrivial isomorphic quotients of those factors, by taking the two projection kernels. Their orders would divide two coprime integers, which is impossible. Induction gives the displayed full product, since all the sets of prime divisors are disjoint. It follows that the actual joint source is the fibre product

$$G_{23} \underset{B/H}{\times} \prod_{p\geq 5} G_p, \tag{E.42.6}$$

with geometric kernel $A_{23} \times \prod A_p$. Indeed the joint source already surjects to B/H, and the proved equality of its geometric kernel gives every element of this fibre product.

Apply the same argument to the image of the corrected shadow. Its $\widehat{H}$ image is onto A_{23} by the preceding paragraph, and is onto each A_p because the entire map on G_p is the identity. The coprime quotient argument gives the full geometric product. Its map to B/H is the identity and onto, so the shadow is onto (E.42.6). This proves joint generation without assuming that the shadow descends to an automorphism of the fixed Ferrari source quotient.

All these maps are supplied by the same word (E.42.4). Taking the inverse system of finite characteristic quotients therefore proves the assertion for every full pro-p factor with $p \geq 5$. The class-two quotients are among their actual quotient maps, which proves the stated simultaneous all-prime comparison. □

Statement E.43 (Proved). The double-cover refinement and every retained boundary tower coexist. *Use G_L, C, T_L, S_L from E.38 and the actual subgroup $H \lhd F \lhd B$ from E.42. Put $S = G_L/C$. The literal quotient map from B has the factorization*

$$B \longrightarrow P_{20} = B/(\gamma_3(H)H^{20}) \longrightarrow S, \tag{E.43.1}$$

and its exact H image is

$$Q_8 \times C_{10}^2. \tag{E.43.2}$$

The group S is the maximal solvable quotient of G_L.

Fix any of the 432 multiplier-one points of E.24, any full two-adic class-two fibre in E.41, and any full three-adic class-two fibre in E.37. There is one profinite commutator word retaining all these images, inducing the identity on every full relative pro-p boundary quotient $\mathscr{P}_p$ for $p \geq 5$, and satisfying the complete $SL_2(\mathbb{F}_5)$ Ferrari test of E.38. All the resulting finite common source maps are surjective. The four-strand map onto $T_{23} \times_{S_L} T_L$ is surjective and satisfies the full pentagon and twist identities.

More precisely, for every finite old common source G_{old} in E.42 which retains P_{20}, the actual new source and its corrected image are both

$$G_{\mathrm{old}} \times_S G_L. \tag{E.43.3}$$

These old sources are cofinal in the specified boundary family. This is a simultaneous extension of that family, with its actual comparison maps.

Proof. We first establish the finite comparison which resolves the additional common five-primary quotient. Write $\overline{F} = \rho_L(F)/C \subset S$ and $\overline{E} = \rho_L(F')/C$. E.38 gives

$$|S| = 43200, \qquad |\overline{F}| = 7200, \qquad \overline{E} \simeq Q_8, \qquad |\overline{F}^{\mathrm{ab}}| = 900. \tag{E.43.4}$$

The index six in the second equality is the actual outer permutation map $B/F \simeq S_3$.

In the faithful 1980-point A_5 source model, the literal words $x = (r^{-1}s)^2$ and $y = (sr^{-1})^2$ both have order 15. The new projection $G_L \to G_A$ has elementary abelian two-kernel. Thus their orders in G_L divide 30. Their signed modular images have order six, so the two actual projections show that both orders are exactly 30. Consequently the images of x, y give an onto map $(\mathbb{Z}/30)^2 \to \overline{F}^{\mathrm{ab}}$. Both groups have order 900, so this is an isomorphism with these specified generators. In particular its inverse composed with $F \to \overline{F}$ is exactly the ordinary pair of free exponent sums modulo 30.

The signed modular map is also specified on all of F. Its generated image in $\mathrm{SL}_2(\mathbb{Z}/12)$ has order 96 and contains no pair $g, -g$. It is therefore the unique lift in that image of

the projective F image used in E.38. Reduction modulo three gives a homomorphism $q_3 : \overline{F} \to \mathrm{SL}_2(\mathbb{F}_3)$. It is injective on $\overline{E}$ by (E.38.12). It follows that the map

$$\Theta : \overline{F} \longrightarrow \mathrm{SL}_2(\mathbb{F}_3) \times (\mathbb{Z}/30)^2, \quad w \longmapsto \big(q_3(w), \mathrm{ab}_x(w), \mathrm{ab}_y(w)\big) \tag{E.43.5}$$

is injective. Indeed an element in its kernel first lies in $\overline{E}$ by abelianization, and then is identity by the injectivity of q_3.

Let $\kappa_3 : \mathrm{SL}_2(\mathbb{F}_3) \to C_3$ be its abelianization, chosen to send both actual matrices $q_3(x), q_3(y)$ to 1. The matrix enumeration verifies this map on all 24 elements. Formula (E.43.5) has image precisely

$$\{(g, a, b) : \kappa_3(g) = a + b \pmod 3\}. \tag{E.43.6}$$

The two generator images lie in this fibre product, and its order is $24 \cdot 900/3 = 7200$, equal to the proved order of the injected group. This proves equality of the groups with their specified maps from F. The program also enumerates all 7200 values directly from these two generator images.

The defining map $F \to F/H = (\mathbb{Z}/3)^2$ is the reduction of the last two coordinates in (E.43.5). It follows that the kernel of $B \to S$ lies in H, and that the actual H image is

$$\{(g, 3a, 3b) : g \in Q_8,\ a, b \in \mathbb{Z}/10\}. \tag{E.43.7}$$

Its multiplication is the direct-product multiplication, proving (E.43.2) and the exact index $43200/800 = 54$. This also identifies $S/(\rho_L(H)/C)$ with the actual quotient B/H.

For an independent centrality comparison, the actual matrices $q_{12}(x^3)$ and $q_{12}(y^3)$ commute with the entire 96-element modular F image. Their commutators with $\overline{F}$ lie in $\overline{E}$, where q_{12} is injective. Thus x^3, y^3 are central in $\overline{F}$. They generate C_{10}^2 by their independent abelianization coordinates, intersecting $\overline{E}$ trivially. This recovers (E.43.7) using the original source generators directly.

The certificate retains all ten literal Schreier generators of H, from the section $x^a y^b$, $0 \le a, b < 3$, and their values in (E.43.7). Their images generate all 800 elements. The two actual Sylow projections are

$$h \longmapsto h^5 \in Q_8 \times C_2^2, \qquad h \longmapsto h^{16} \in C_5^2. \tag{E.43.8}$$

The images have orders 32 and 25, commute, and multiply back to h for all 800 values. In the coordinates of (E.43.7) they are the reductions of (a, b) modulo two and five, retaining g in the first factor. The five-primary map is explicitly

$$h \longmapsto \big(\mathrm{ab}_x(h)/3, \mathrm{ab}_y(h)/3\big) \pmod 5; \tag{E.43.9}$$

division is legitimate on H, and is equivalently multiplication by two of the ordinary exponent sums modulo five. Every third lower-central commutator and every twentieth power vanish in this finite H image; its derived group has order two and its exponent is exactly 20. These identities prove (E.43.1) for the literal images of s, r. The subgroups in its kernel are characteristic in H and hence normal in B, so this is a homomorphism of the displayed actual quotient groups.

The group in (E.43.7) is solvable, and B/H is the extension of the abelian group F/H by S_3. Hence S is solvable. Conversely every solvable quotient of G_L kills the perfect normal subgroup C. Thus S is the maximal solvable quotient, with the specified map from B.

The old system contains this quotient. The actual Hall-coordinate maps give

$$P_{20} \xrightarrow{\sim} P_4 \times_{B/H} P_5. \tag{E.43.10}$$

On the geometric kernel, the abelian coordinates are modulo 20 and the central Hall coordinates modulo 10. Indeed the power and commutator formulas give the central modulus $\gcd(20, \binom{20}{2}) = 10$; conversely these Hall moduli satisfy every twentieth-power relation. The maps on these coordinates are the Chinese-remainder isomorphisms to moduli $(4, 5)$ and $(2, 5)$, respectively. They respect collection because they are the natural quotient homomorphisms of H. The two geometric factors have coprime orders, so the image of H is their full product. Together with the common quotient B/H, this proves (E.43.10). E.41 retains P_4, and E.42 retains P_5 as a quotient of its full relative pro-five factor. Consequently adjoining this fixed P_{20} to any old finite source gives a cofinal subsystem of the family in E.42, with an actual map to S by (E.43.1).

A correction preserving every solvable boundary image. Let $f_0 \in \widehat{H} \cap \widehat{F}'$ be the actual word supplied by E.42 for the prescribed boundary parameters. It retains the chosen G_6 point, so its q_{12} value is still the indexed matrix in (E.38.13). Choose an actual Galois element σ with $\chi(\sigma) = 1$ and the corresponding Kummer value, as in E.38. The exact quotient comparison then gives

$$c = \rho_L(f_0)^{-1}\rho_L(f_\sigma) \in C. \tag{E.43.11}$$

For a profinite group J, let $J^{[r]}$ be its closed derived series, beginning with $J^{[0]} = J$. Since $\rho_L(\widehat{N}^{[1]}) = C$ and C is perfect,

$$\rho_L(\widehat{N}^{[r]}) = C \qquad (r \geq 1). \tag{E.43.12}$$

The preimages of the fixed c inside these compact groups are nonempty, nested closed sets. Their intersection therefore gives an element k with

$$k \in \bigcap_{r \geq 1} \widehat{N}^{[r]}, \qquad \rho_L(k) = c. \tag{E.43.13}$$

Here $N' \subset F' \subset H$. Induction on the derived series gives $\widehat{N}^{[r+1]} \subset \widehat{H}^{[r]}$. Thus k belongs to the full prosolvable residual $\bigcap_r \widehat{H}^{[r]}$. Every homomorphism from $\widehat{H}$ to a finite solvable group kills this intersection, by the finite derived length of its image. This applies to the H image of $\mathscr{D}$, to every finite two- or three-adic class-two boundary image, and to every finite p-group quotient of $\widehat{H}$ for every $p \geq 5$. Consequently k is identity in every component retained in E.42, simultaneously. More explicitly, this proves the exact continuous surjection

$$R_H = \bigcap_{r \geq 0} \widehat{H}^{[r]}, \qquad \rho_L|_{R_H} : R_H \twoheadrightarrow C. \tag{E.43.14}$$

The upper inclusion of its image in C follows because $S = G_L/C$ is solvable; the reverse inclusion is the nested-fibre construction for each $c \in C$.

Set $f = f_0 k$. It remains in $\widehat{F}'$ and retains the entire old source family, including the identity maps on the full pro-p boundary factors away from two and three. In the new source, $\rho_L(f) = \rho_L(f_\sigma)$. All 36 induced coordinates agree, so the complete new hexagons and pentagon are those of an actual Galois element, with the exact based braid convention of E.38. The same single f supplies these maps at every old finite level.

The actual common source and its generation. Let G_{old} be an old finite source containing P_{20}. It is solvable: its factors have finite solvable geometric kernels and the solvable quotient B/H, and $\mathscr{D}$ is solvable as proved in E.24. Every common quotient of G_{old} and G_L therefore factors through S. The specified map through P_{20} makes S itself their common quotient. Goursat's lemma identifies their actual joint B image with (E.43.3).

The corrected source map is onto G_{old} by E.42 and onto G_L because its generator images are the Galois images followed by ρ_L. Its two maps to S agree: they evaluate the same literal source formulas using f and the multiplier one. The maximal-common-quotient argument therefore identifies its image with the full fibre product (E.43.3). Projecting this assertion from the cofinal subsystem proves generation at every indicated old finite level. The four-strand comparison is exactly the one in E.38, since the complete D_{23} image is preserved and the new six evaluations are Galois values. It gives surjectivity onto $T_{23} \times_{S_L} T_L$, together with its braid and full-twist identities.

The new program `de_sl2_boundary_quotient.py` retains the exact finite model, all ten source-generator images, all 7200 and 800 group elements, and both Sylow projections. Its comparison uses the proved E.38 source kernels and an independent literal order check in the original A_5 permutation representation. The profinite correction is supplied by the explicit nested compact fibres in (E.43.12)–(E.43.13). No unevaluated Galois word is presented as a computed literal certificate. The conclusion concerns this specified family of finite sources; cofinality among all finite quotients of B has not been proved. □

Statement E.44 (Proved). The exponent-three class-three boundary quotient and its exact comparison with the class-two towers. *Retain the literal groups $H \triangleleft F \triangleleft B$, the ten Schreier generators η_i, and the common source $\mathscr{D}$ of E.24. Put*

$$\mathcal{Q}_3 = B/(\gamma_4(H)H^3), \qquad \mathcal{J}_3 = \mathrm{im}(B \longrightarrow \mathscr{D} \times \mathcal{Q}_3). \tag{E.44.1}$$

The geometric kernel of $\mathcal{Q}_3 \to B/H$ has order 3^{175}, with successive dimensions $10, 45, 120$. Its actual projection to the class-two quotient P_3 of E.33 forgets the last 120 coordinates.

Every one of the 432 multiplier-one source points of E.24 has a generating lift to $\mathcal{J}_3$. Over each such source point the complete finite fibre of correction images has 3^{28} elements. Exactly 3^6 of its 3^{11} E.33 images lift to this class-three quotient. Each of those images has an affine $\mathbb{F}_3^{22}$ fibre.

The comparison with the full three-adic class-two tower of E.37 imposes four further independent congruences on its eight free Smith coordinates. They are explicitly (E.44.13)–(E.44.14) *below. Over each fixed source point, 3^5 of the 3^9 E.37 reduction images satisfy them, and there are 3^{27} compatible images in $\mathcal{Q}_3$. The complete compatible inverse-limit fibre, retaining the class-three coordinate, has the explicit continuous parametrization*

$$\mathbb{Z}_3^{10} \times \mathbb{F}_3^{22} \tag{E.44.2}$$

constructed in the proof. It can be prescribed together with every two-adic class-two family of E.41 and the prime-to-six identity factors of E.42; one profinite commutator word realizes each such family, and every finite common source is generated. The same families retain E.43's $\mathrm{SL}_2(\mathbb{F}_5)$ Ferrari refinement in the actual common sources (E.44.15) *below.*

There is a specified point of the entire E.37 tower, over E.24 row 0 with parameters $(0,0;0)$, that has no lift to (E.44.1)*: take*

$$u = w = 0, \qquad Y_{37} = 1, \qquad Y_{38} = \cdots = Y_{44} = 0 \tag{E.44.3}$$

in E.37's coordinates. Thus the new obstruction remains after all the earlier class-two levels have been imposed.

Proof. We first give the exact quotient, including its cube relations. Work in the truncated free associative algebra over $\mathbb{F}_3$ on $X_1, \dots, X_{10}$, with monomials of degree at least four set to zero, and send η_i to $1 + X_i$. The group M generated by these units is the quotient

$$H/(\gamma_4(H)\,\gamma_2(H)^3 H^9). \tag{E.44.4}$$

Here and below a power subgroup means the subgroup generated by the indicated powers, with closure in the profinite notation.

To prove (E.44.4), ordinary class-three Hall collection gives ten generator exponents modulo nine, forty-five degree-two exponents modulo three, and 330 ordinary degree-three exponents modulo three. Their collected forms are independent in the displayed algebra. The degree-one and degree-two terms first recover the generator residues and commutator exponents. Changing a generator exponent by three adds X_i^3 in degree three. The remaining ordinary degree-three Hall terms have the 330 Lyndon leading words, each with coefficient one, and are independent of the ten X_i^3. This proves both the inverse collection map and the claimed kernel. In particular $|M| = 3^{395}$.

Every cube in M is central. For a unit $1 + A$ its cube is $1 + v^3$, where v is the degree-one part of A. The complete span of these cube values has dimension 220. An explicit basis consists of

$$\begin{aligned} &X_i^3, \qquad X_i^2 X_j + X_i X_j X_i + X_j X_i^2 \quad (i \neq j), \\ &\sum_{\sigma \in S_3} X_{\sigma(i)} X_{\sigma(j)} X_{\sigma(k)} \quad (i < j < k). \end{aligned} \tag{E.44.5}$$

Their disjoint monomial supports prove independence; their numbers are $10, 90, 120$. Each is an actual linear combination of cubes: the repeated-index terms follow from $(X_i + X_j)^3$ and $(X_i - X_j)^3$, and the three-distinct-index term follows by subtracting the three two-variable cubes from $(X_i + X_j + X_k)^3$ and adding the single-variable cubes. Conversely expansion of any v^3 belongs to this span. Because the subgroup in (E.44.4) lies in $\gamma_4(H)H^3$, quotienting by precisely this central span gives the required group $H/(\gamma_4(H)H^3)$.

Here is a small coordinate model of that exact quotient. Retain E.33's coordinates $v \in \mathbb{F}_3^{10}$ and $z \in \Lambda^2 \mathbb{F}_3^{10}$. For $i < j < k$, let t_{ijk} be the coefficient of $X_i X_j X_k$ minus that of $X_j X_i X_k$. This map annihilates (E.44.5) and takes $[X_i, [X_j, X_k]]$ to the corresponding unit vector. On the 340-dimensional restricted degree-three term its kernel therefore has dimension 220, and is exactly the complete cube span just proved. It gives a bijective group coordinate map

$$H/(\gamma_4(H)H^3) \longrightarrow \mathbb{F}_3^{10} \oplus \mathbb{F}_3^{45} \oplus \mathbb{F}_3^{120}. \tag{E.44.6}$$

For explicit multiplication put, for $i < j$,

$$A(v,z)_{ij} = 2v_i v_j + z_{ij}, \qquad A(v,z)_{ji} = 2v_i v_j - z_{ij}, \qquad A(v,z)_{ii} = 2(v_i^2 - v_i). \tag{E.44.7}$$

These are the actual quadratic Magnus coefficients. For $(v,z,t)(w,a,b) = (v',z',t')$ the law is

$$\begin{aligned} v' &= v + w, \qquad z' = z + a + 2v \wedge w, \\ t'_{ijk} &= t_{ijk} + b_{ijk} + v_i A(w,a)_{jk} - v_j A(w,a)_{ik} \\ &\qquad + (A(v,z)_{ij} - A(v,z)_{ji}) w_k. \end{aligned} \tag{E.44.8}$$

The inverse coordinate map is equally explicit. With exponents represented by $0, 1, 2$, take the ordered product of $\eta_i^{v_i}$, then of $[\eta_i, \eta_j]^{z_{ij} - 2v_i v_j}$, and finally of $[\eta_i, [\eta_j, \eta_k]]^{c_{ijk}}$, where

$$c_{ijk} = t_{ijk} - v_i v_j v_k - v_i z_{jk} + v_j z_{ik}. \tag{E.44.9}$$

The displayed product has exactly (E.44.6) as its coordinates. This also specifies all word preimages used by the computation.

The arithmetic source maps are the same literal substitutions as in E.33: $s : (x, y) \mapsto (y, x)$ and $r : (x, y) \mapsto (y, (xy)^{-1})$. Substitute the ten fixed Schreier words and collect by (E.44.8). Denote the resulting genuine actions by S and R. They have orders two and three and induce the old E.33 actions on (v, z); on the new centre they induce $\Lambda^3 S_1$ and $\Lambda^3 R_1$. The program retains each of the ten complete generator images and verifies the actual words $x = (r^{-1}s)^2$ and $y = (sr^{-1})^2$ against conjugation by x and y.

For $g \in H/(\gamma_4(H)H^3)$ the two equations are literally $(Sg)g = 1$ and $g^{-1}R(g^{-1})R^2(g^{-1}) = 1$. They define the map $s \mapsto sf$, $r \mapsto f^{-1}r$. This map fixes sr, hence fixes x; it sends $y = sxs$ to $f^{-1}yf$. For $z = (xy)^{-1} = ryr^{-1}$ its image is hzh^{-1} with $h = f^{-1}R(f^{-1})$. Thus these are the actual multiplier-one peripheral maps, with their based conjugators.

The linear norm matrix on the new centre is

$$M_3 = \begin{pmatrix} I + \Lambda^3 S_1 \\ -I - \Lambda^3 R_1 - (\Lambda^3 R_1)^2 \end{pmatrix}, \qquad \operatorname{rank}_{\mathbb{F}_3} M_3 = 98. \tag{E.44.10}$$

Hence its kernel has dimension 22. For each of E.33's nine admissible abelian values, write its central coordinate as $z_0 + K_0 a$, where K_0 is the saved 45-by-10 basis of the E.33 central norm kernel. At fixed v all remaining norm equations are affine linear in (a, t): this follows directly from (E.44.8) and the collected action. The nine complete 240-by-130 matrices have rank 103, and every affine system is solvable. Thus each abelian value has 3^{27} solutions, with five independent conditions on a and an affine 22-dimensional triple fibre. All nine matrices, their row operations, kernels and particular solutions are reproduced by the finite certificates below. Each of the 432 actual source cosets contains exactly three of these abelian values by the literal E.33 source comparison. This proves the counts 3^{28} and 3^6 once the complete new source image is checked.

That image follows by actual words. E.33 provides forty-five adjusted normal relators u_ℓ in $R_3 \cap [H, H]$, with $R_3 = \ker(B \to D_6) \cap F'$, whose degree-two reductions form a basis. For every pair $j < k$ choose the saved coefficients so that $u_{jk} = \prod_\ell u_\ell^{a_\ell}$ has degree-two coordinate $e_j \wedge e_k$ modulo three. The words

$$[\eta_i, u_{jk}^4], \qquad i < j < k, \tag{E.44.11}$$

lie in $\ker(B \to \mathscr{D}) \cap \gamma_3(H)$ and map to the 120 standard new central vectors. Indeed fourth powers kill the exponent-two binary image and retain the exponent-three target; the ternary image is already trivial. The uncomputed triple coordinate of u_{jk} does not enter its commutator, since that layer is central. These words have zero class-two image over every coefficient ring, because they belong to $\gamma_3(H)$ integrally. The exact relator indices, integral adjustments and 45 coefficients for every word in (E.44.11) are retained. Together with the complete E.33 image, they prove that the new common-source kernel image is its full inverse image, with dimensions $6, 45, 120$. There is no further source or F' constraint on the last two coordinates.

We next compare with the full E.37 tower. Its two extra Smith conditions modulo three and the five new conditions just computed have joint rank six, in all nine abelian cases; the joint affine systems are nonempty. In the notation of E.37 put

$$c = c_f, \qquad \bar{u} = t_2 \bmod 3, \qquad b = (\bar{Y}_{37}, \dots, \bar{Y}_{44})^t \in \mathbb{F}_3^8. \tag{E.44.12}$$

The complete additional criterion is

$$A_c b = \beta(c, \bar{u}), \tag{E.44.13}$$

where the three rank-four matrices are

$$A_0 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 2 \\ 0 & 0 & 1 & 0 & 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 2 & 2 \end{pmatrix}, \qquad A_1 = \begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 & 0 & 2 \\ 0 & 0 & 1 & 0 & 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 0 & 0 & 1 & 2 & 2 \end{pmatrix},$$

$$A_2 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 1 & 2 & 2 \\ 0 & 1 & 0 & 0 & 0 & 2 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 2 & 2 \end{pmatrix},$$

	$\bar{u} = 0$	$\bar{u} = 1$	$\bar{u} = 2$
$c = 0$	$(0, 0, 0, 0)$	$(1, 0, 0, 2)$	$(1, 0, 2, 0)$
$c = 1$	$(2, 0, 2, 0)$	$(1, 2, 0, 0)$	$(2, 0, 0, 2)$
$c = 2$	$(1, 2, 2, 1)$	$(0, 0, 0, 1)$	$(0, 2, 0, 0)$

(E.44.14)

The table entries are the column vectors β. These matrices are obtained by substituting the actual E.37 origin and eight free Smith columns into the class-three norm equations, and eliminating only the new 120 triple variables. In particular they retain both old Smith factors three; no arbitrary class-two solution has been treated as a point of the inverse-limit tower.

For an explicit continuous parametrization choose $u, w \in \mathbb{Z}_3$ and form the E.37 abelian point $t_2 = u$, $t_1 = c - u + 3w$, with its specified Hensel root t_3. In each matrix A_c the four pivot columns are unit columns. Choose the four nonpivot Smith coordinates freely in $\mathbb{Z}_3$; set the pivot coordinates equal to the indicated affine solutions of (E.44.13), using the digits $0, 1, 2$ for β, plus three times four further arbitrary elements of $\mathbb{Z}_3$. This is a continuous bijection from $\mathbb{Z}_3^{10}$ onto the compatible E.37 points, with inverse obtained by recovering u, w, the four free coordinates and dividing the four pivot discrepancies by three. For each such point a saved particular solution of (E.44.10) and the 22 saved kernel columns give its complete affine triple fibre. This proves (E.44.2) with actual maps. There are 3^4 compatible old central reductions at each of the three abelian values, hence 3^5 per source and $3^{5+22} = 3^{27}$ compatible class-three images.

The comparison maps also prove simultaneity. The common quotient of the new geometric group and the class-two pro-three group is exactly $H/(\gamma_3(H)H^3)$: every class-two quotient of the former kills its new centre, and both displayed maps onto this common quotient are the literal projections. More concretely, start with the E.37 word having a prescribed class-two tower value. Any desired class-three value with the same P_3 projection differs from its actual value by a new central vector. Correct this vector with the words (E.44.11). To retain all E.42 factors, take their three-supported continuous powers. This preserves their exponent-three images, kills their images in every full pro-p factor for $p \neq 3$, and still has exactly zero image in every class-two quotient. It also retains all $\mathscr{D}$ coordinates. This constructs one profinite commutator word for every claimed family.

Generation is checked on the actual common source. In $\mathcal{J}_3$ the image of N_B is finite nilpotent with primes two and three: its old image has this property, and its new image is a three-group, including the cyclic-three image in F/H. It is generated by the 97 fixed generators and maps onto $(\mathbb{Z}/6)^{97}$ through G_6. Its Sylow Frattini dimensions are therefore exactly 97. The retained G_6 map is an automorphism on these quotients, proving surjectivity on N_B and then on $\mathcal{J}_3$. The direct check of the induced H abelian map, using $x \mapsto x$, $y \mapsto f^{-1}yf$, has rank ten for all nine abelian values as well.

For the simultaneous two-adic and three-adic class-two factors, choose a lower-central index larger than the nilpotence classes of the old Ferrari N_B image and every three-primary image, including the new one. Exactly as in E.41, the image at that index supplies the pure stable two-primary lower-central subgroup of the boundary factor. The corrected map is already onto that factor, because its old value has been retained. Removing this pure subgroup leaves the finite nilpotent common N_B image, so the same rank-97 Frattini argument applies. For further prime-to-six factors the actual H images have coprime orders to the

existing two-three image; their product over B/H is the actual common image by E.42's proved map. The identity values on them give surjectivity. No invariance of a fixed Ferrari source kernel under the shadow is used.

For the simultaneous E.43 refinement, let G_L, C and $S = G_L/C$ be its actual source, perfect kernel and maximal solvable quotient. Take a cofinal system of the finite common boundary sources G_{old} just constructed that also contain P_{20}. Their actual maps to S are supplied by $P_{20} \to S$ from E.43. Their H images are solvable, including the new finite three-group; the quotient B/H is solvable as well. Every common quotient with G_L therefore factors through S, and S itself is common. Thus the literal source map gives precisely

$$\mathrm{im}\big(B \longrightarrow G_{\mathrm{old}} \times G_L\big) = G_{\mathrm{old}} \times_S G_L. \tag{E.44.15}$$

For a word f_0 realizing a prescribed new boundary family, its S value is the E.43 value of the matching Galois word f_σ, because all the old source coordinates and P_{20} have been retained. The discrepancy $\rho_L(f_0)^{-1}\rho_L(f_\sigma)$ lies in C. The exact surjection $\rho_L(R_H) = C$ of E.43, where $R_H = \bigcap_{j\geq 1} H^{[j]}$, supplies a single $k \in R_H$ with that image. Then $f = f_0 k$ retains every old solvable H image, including the new class-three group, and agrees with f_σ in G_L. The corrected image is onto both factors of (E.44.15) and has the same maximal common quotient S, so it is the whole displayed fibre product. The four-strand Ferrari evaluation maps are those of E.43 and are retained literally. This proves the additional simultaneous assertion.

Finally the point (E.44.3) has $t = 0$ and doubled-log central vector $V_M e_{37}$ in the exact Smith coordinates of E.37. Its integral central norms vanish, so it is a point of every class-two level. It lies in the specified actual source fibre, whose residue is $c = 0$. The first row of A_0 at $\bar{u} = 0$ requires $\bar{Y}_{37} = 0$, contrary to (E.44.3). The certificate additionally records a row annihilating all 120 columns of M_3 and evaluating to one on this point's norm error. This proves the asserted failure of every class-three lift over that tower point.

The three new programs in `code` have the shared prefix `bc_fermat_class_three_` and suffixes `burnside.py`, `norms.py`, and `tower_comparison.py`. They retain the complete quotient and action, all nine affine systems, all 432 source records, the literal relator preimages, the finite matrices in (E.44.14), and the specified nonlifting point. Their executed outputs have status `PASS`; hashes verify that every checkpoint05 input remained unchanged. □

Statement E.45 (Proved). The complete A_6 kernel and outer-coordinate tests retain every old fibre. *Let $B = C_2 * C_3$, $F = \langle x, y\rangle$, N and $K_6 = N'N^6$ be the discrete groups with their literal generators from E.1 and E.24. The action of the Ferrari Nielsen generators on the epimorphisms $F_2 \twoheadrightarrow A_6$ has 53 orbits under $\mathrm{Aut}(A_6)$, with Nielsen orbit sizes $12, 15, 10, 16$. Retain either the permutation of these kernels, or that permutation together with every transporter in $\mathrm{Out}(A_6) \simeq C_2^2$. These give specified quotient maps*

$$B_4 \longrightarrow T_1 \longrightarrow T_0, \qquad B \xrightarrow{\rho_\epsilon} G_\epsilon \quad (\epsilon = 0, 1), \tag{E.45.1}$$

where ρ_ϵ retains all six Ferrari evaluations and all six cosets of F in B. Put $C_\epsilon = \rho_\epsilon(N')$ and $S_m = B/(F'F^m)$. Then

$$\begin{gathered} C_\epsilon = \rho_\epsilon(F') = \rho_\epsilon(K_6 \cap F'), \qquad C'_\epsilon = C_\epsilon, \\ G_\epsilon/C_\epsilon = S_{30}, \qquad G_\epsilon/\rho_\epsilon(K_6) = S_6, \qquad |S_{30}| = 5400, \quad |S_6| = 216. \end{gathered} \tag{E.45.2}$$

The two equalities with S_m identify the actual maps from B. Both projections $T_1 \to T_0$ and $G_1 \to G_0$ have elementary abelian kernel of order 2^{35}.

Every one of the 864 retained points of E.24 extends to these common source and four-strand targets. The assertion holds for every profinite multiplier congruent to 1 modulo 72,

with the appropriate retained E.24 point. For the source the actual common image and the corrected image are

$$D_{23} \times_{S_6} G_\epsilon; \tag{E.45.3}$$

for the four-strand target they are

$$T_{23} \times_{C_2} T_\epsilon, \tag{E.45.4}$$

where the two maps to C_2 are braid exponent modulo two. All these maps are surjective and satisfy the complete pentagon, braid and full-twist equations.

For multiplier one the correction can simultaneously preserve any full boundary fibres of E.42. Thus all 432 *points of E.24 coexist with these two A_6 tests, the complete specified two- and three-adic class-two boundary fibres, and the identity on every full relative pro-p boundary quotient for $p \geq 5$. For cofinal old common sources retaining P_{20} from E.43, the actual common source and its corrected image are*

$$G_{\mathrm{old}} \times_{S_{30}} G_\epsilon. \tag{E.45.5}$$

This assertion concerns the displayed family of quotients.

Proof. We give the finite model and each comparison used in the lifting argument. We use the complete A_6 and $\mathrm{Aut}(A_6)$ tables of D.31, and independently check every one of the 1440 automorphisms on both generator edges of the entire 360-vertex Cayley graph. The action partitions all 129600 ordered pairs into 124 orbits. Testing generation on one representative of every orbit gives 76320 generating pairs. The automorphism action on generating pairs is free, so there are exactly $76320/1440 = 53$ epimorphism kernels. All pairs, orbit representatives and transporters are retained.

For representatives (g_j, h_j), the two literal Nielsen operations are

$$(g,h) \longmapsto (g,hg), \qquad (g,h) \longmapsto (gh^{-1},h). \tag{E.45.6}$$

For each operation the program records the unique permutation p of the 53 kernels and actual automorphisms c_j carrying $(g_{p(j)}, h_{p(j)})$ to the displayed pair. It checks the entire automorphism-valued braid relation before taking any quotient. The multiplication is

$$(p,c)(q,d) = (q \circ p, (c_j d_{p(j)})_j). \tag{E.45.7}$$

Keeping only p gives T_0; keeping p and the two outer bits of every c_j gives T_1. The latter acts faithfully on the 212 pairs (j,e) with $e \in C_2^2$, by $(j,e) \mapsto (p(j), e + \bar{c}_j)$. The 53-point action is its specified quotient. The two actual Ferrari generators ξ_1, ξ_2 map to identity in these targets, and consequently their three Artin generators are (a,b,a). Direct multiplication verifies all three B_4 braid relations. This defines the first maps in (E.45.1).

This construction retains the universal epikernel coordinates of [Gui15, Lemma 2.1]: the free group evaluates onto A_6^{53}, and the Nielsen substitutions act on this product by the displayed permutation and actual automorphisms. The maps just defined compose the Ferrari action on that product with its kernel-permutation or outer-coordinate projection. No identification of the finite automorphism group computed in [Gui15, Theorems 4.9 and 5.2] with our source image is used.

Let $\theta_i : F \to T_\epsilon$, $1 \leq i \leq 6$, be the six pure-word evaluations printed in E.17, now using these literal a, b, a. Let τ_q be the retained Schreier section of $B/F \simeq S_3$. The source homomorphism is the induced map whose 36 coordinates at $b \in B$ are

$$\theta_i(\tau_q b \tau_{\bar{qb}}^{-1}), \qquad q \in B/F, \quad 1 \leq i \leq 6, \tag{E.45.8}$$

together with the permutation $q \mapsto q\bar{b}$. The intervening word lies in F; the exact free-graph Schreier map converts it to a word in x, y. Multiplication in the induced group follows by cancellation of the adjacent section words. Thus its kernel is normal in B, and it retains all six evaluations and their conjugates. It extends continuously to the profinite completion.

The resulting faithful permutation degrees are 1908 and 7632. We reduce them to 432 and 798, respectively, by deleting only repeated transitive B-sets. For every deleted orbit the certificate gives a bijection to a retained orbit and checks that it intertwines both original generators s, r pointwise. These maps prove equality of the two kernels, so the compressed actions are faithful representations of the same G_ϵ.

Write

$$c_0 = 2^{40}3^{21}5^{10}7^{6}11^{3}13^{2}, \qquad c_1 = 2^{35}c_0. \tag{E.45.9}$$

Complete permutation-group calculations give

	G_ϵ	$\rho_\epsilon(F)$	$\rho_\epsilon(F')$	$\rho_\epsilon(N)$
$\lvert\cdot\rvert/c_\epsilon$	5400	900	1	75

	$\rho_\epsilon(N')$	$\rho_\epsilon(K_6)$	$\rho_\epsilon(K_6 \cap F')$
$\lvert\cdot\rvert/c_\epsilon$	1	25	1

(E.45.10)

Here is the completeness of the subgroup calculation. All 97 literal Schreier generators of N are retained, and the program verifies that the four selected generators generate their entire finite image. It computes the derived subgroup, including the full normal closure of the commutators, then adjoins every actual sixth power for K_6. For $K_6 \cap F'$ it uses the 95 integral relations among the two ordinary exponent sums of the 97 generators. The integer relation matrix has rank 95 and Smith diagonal (1^{95}), and its product with the exponent-sum matrix is zero. It is therefore the full saturated integral kernel. Modulo $\rho_\epsilon(N')$ the products of the sixth powers commute, so these 95 products give precisely the image of $K_6 \cap F'$. Each is already in $\rho_\epsilon(N')$. The derived subgroup of this last image has the same order, proving its perfectness. The full derived subgroup of the two-generator F image has the same order and contains it, proving the first line of (E.45.2).

The literal images of x, y in both sources have order 30. Since they generate F and its image modulo C_ϵ is abelian of order 900, the specified map

$$(\mathbb{Z}/30)^2 \longrightarrow \rho_\epsilon(F)/C_\epsilon, \qquad (u, v) \longmapsto x^u y^v C_\epsilon \tag{E.45.11}$$

is an isomorphism. Consequently $B \to G_\epsilon/C_\epsilon$ factors through S_{30}; both groups have order 5400, so the factorization is the claimed isomorphism. The image of K_6 contains x^6, y^6, as checked directly, and has order $25c_\epsilon$. The corresponding quotient is therefore the actual S_6 quotient. These comparisons use ordinary free exponent sums, with no change of their normalization.

The upper and lower models are related by forgetting the outer bits in every coordinate of (E.45.8). The orders in (E.45.9)–(E.45.10) show that its source kernel has order 2^{35}. In that kernel every permutation coordinate is identity, so multiplication of the remaining bits is addition in an elementary abelian two-group. The same calculation on the four-strand targets gives

$$|T_\epsilon| = 2c_\epsilon, \qquad |T'_\epsilon| = |T''_\epsilon| = c_\epsilon. \tag{E.45.12}$$

Thus T'_ϵ is perfect, and its maximal solvable quotient is the actual braid-exponent map $T_\epsilon \to C_2$. The target projection also has elementary abelian kernel of order 2^{35}. The full twist $(aba)^4$ is identity in both targets; $(ab)^3$ has order one in T_0 and two in T_1.

We now prove the full-fibre assertion. Fix a retained E.24 word $f_0 \in \widehat{F}'$ and its multiplier $\lambda \equiv 1 \pmod{72}$. Choose an actual Galois element with cyclotomic character λ, and write

its standard Ihara word as $f_\sigma \in \widehat{F}'$. The cyclotomic character is onto $\widehat{\mathbb{Z}}^\times$, so this choice is available for every stated multiplier. By (E.45.2),

$$\rho_\epsilon(f_0)^{-1}\rho_\epsilon(f_\sigma) \in C_\epsilon. \tag{E.45.13}$$

The joint image of N' in the old finite source D_{23} and C_ϵ projects onto each factor. The first image is solvable and the second is perfect. Every common quotient is both solvable and perfect and is therefore trivial. Goursat's lemma gives the direct product, providing a word $k \in N'$ with identity old image and the value in (E.45.13) in the new source. Then $f = f_0 k$ retains the complete old point and agrees with f_σ in all 36 new source coordinates. In particular all six four-strand evaluations agree. The actual Galois action, in the braid convention of E.38, proves all the asserted equations and surjectivity on the new factors. Every finite power uses the same chosen profinite λ; this argument requires no restriction on its new five-primary residue. For $\lambda = 1$ we may choose $\sigma = 1$, so all new evaluations of the corrected f are literally identity.

The exact generation comparisons are as follows. Every common quotient of D_{23} and G_ϵ kills C_ϵ, since D_{23} is solvable. It also kills $\rho_\epsilon(K_6)/C_\epsilon \simeq C_5^2$, since D_{23} has only two- and three-primary factors. It is therefore a quotient of S_6. Conversely S_6 is a specified quotient of both groups: the old source retains G_6, and $K_6 \subset F'F^6$. Thus S_6 is their maximal common quotient with these maps from B. Goursat proves (E.45.3) for both the original source and the corrected image. On S_6 the prescribed maps agree, since the multiplier is 1 modulo six and the word lies in F'. Similarly, (E.45.12) shows that every common quotient of T_{23} and T_ϵ factors through C_2. On the binary Ferrari factor the action on the three nonzero vectors of $\mathbb{F}_2^2$ sends each Artin generator to a transposition, giving this actual common quotient. Odd multipliers and conjugation preserve it. This proves (E.45.4) and the asserted joint four-strand generation.

For the simultaneous boundary assertion let H and the closed derived terms $\widehat{H}^{[r]}$ be as in E.42–E.43, and put $R_H = \bigcap_r \widehat{H}^{[r]}$. The exact H image in S_{30} is $3(\mathbb{Z}/30)^2 \simeq C_{10}^2$ under (E.45.11). In particular

$$B \longrightarrow P_{20} = B/(\gamma_3(H)H^{20}) \longrightarrow S_{30} \tag{E.45.14}$$

is the literal quotient factorization, and S_{30} is the maximal solvable quotient of G_ϵ. Perfectness gives $\rho_\epsilon(\widehat{N}^{[r]}) = C_\epsilon$ for every $r \geq 1$. The inclusions $\widehat{N}^{[r+1]} \subset \widehat{H}^{[r]}$ and compactness of the nested fibres over each element of C_ϵ prove

$$\rho_\epsilon|_{R_H} : R_H \twoheadrightarrow C_\epsilon. \tag{E.45.15}$$

The image is contained in C_ϵ because S_{30} is solvable. Choose this preimage to cancel the new value of the E.42 word f_0. It changes none of the finite solvable H-images, hence preserves the entire stated boundary family simultaneously. All its new evaluations are identity.

Every finite old common source in E.42 is solvable: its H image is a finite product of finite p-groups and its quotient B/H has order 54. For finite old common sources containing P_{20}, the maximal common quotient is now exactly S_{30}, by (E.45.14) and maximal solvability. Both component maps generate, and they agree on this specified quotient. Goursat gives (E.45.5). Such old sources are cofinal in the family of E.42, since adjoining its fixed finite P_{20} stage stays in that family. The compactness argument already gives one coherent correction, rather than independent choices at the finite stages.

The reproducible certificates have common prefix `de_ferrari_a6_` and suffixes `kernels.py`, `source_data.py`, `kernel_source.py`, `kernel_corrections.py`, and `kernel_compare.py`, with their retained JSON, NPZ and stdout files. Both modes retain the entire correction images, and the comparison file checks the literal quotient

generators and four-strand derived groups. The source kernels in this statement and their common refinements pull back to normal finite-index subgroups of B_3 contained in PB_3 after intersection with the retained K_6 kernel. No cofinality in all finite B_3 quotients is asserted. □

5.10 A five-point Hurwitz quotient and its complete correction source

Statement E.46 (Proved). Every retained CM fibre survives the five-point alternating Hurwitz test. *There is a specified quotient of discrete groups, with its continuous extension to the profinite completion,*

$$q_5 : \Gamma_{0,[5]} \twoheadrightarrow A_{96}. \tag{E.46.1}$$

Its pure mapping-class image is also A_{96}. The four standard half twists have orders 30, and the rotation $\delta_5 = \sigma_1\sigma_2\sigma_3\sigma_4$ has image of order five. The restriction along the actual four-strand map $B_4 \to \Gamma_{0,[5]}$, fixing the fifth marked point, is onto A_{96} as well.

Retaining all five pentagon evaluations and all three based five-strand action evaluations gives an explicit induced source

$$\rho_5 : B = C_2 * C_3 \twoheadrightarrow G_5, \qquad |G_5| = 2^{18}3^{20}5^{15}7^2. \tag{E.46.2}$$

Every one of the 864 points of E.24 extends to the common source and four-strand targets with this quotient, for every compatible profinite unit multiplier $\lambda \equiv 1 \pmod{72}$. The resulting maps satisfy both source hexagons, the complete pentagon, and all five-strand braid and sphere relations. Their five-strand image is A_{96}, and

$$\sigma'_1\sigma'_2\sigma'_3\sigma'_4 \sim_{A_{96}} q_5(\delta_5)^\lambda. \tag{E.46.3}$$

Here the primed generators are the actual based action values defined below. In particular all 24 CM fibres, including the 12 extra fibres, survive this order-five test.

For multiplier one, the same correction preserves any prescribed full boundary family of E.42 and any compatible class-three factor of E.44. Put $C = \rho_5(N')$, $K = \rho_5(K_6)$, $S = G_5/C$, and $J_5 = G_5/K$. The exact common source maps and their corrected images are

$$D_{23} \times_{J_5} G_5, \qquad |J_5| = 216, \tag{E.46.4}$$

and, for the cofinal indicated boundary sources retaining P_{20},

$$G_{\mathrm{old}} \times_S G_5, \qquad |S| = 5400. \tag{E.46.5}$$

The actual common four-strand target and corrected image are $T_{23} \times A_{96}$. All common-quotient maps in these displays are induced by the same literal generators of B or B_4. These assertions also hold for the very same word already retaining E.38's $\mathrm{SL}_2(\mathbb{F}_5)$ source, including all the simultaneous boundary families of E.43–E.44. The actual maps $G_L \to G_A \to G_5$ and the resulting common source graphs are (E.46.26) *below. Their common four-strand target is* (E.46.27).

Proof. We construct the quotient, compute its complete correction image, and prove the stated common-source maps.

The actual five-point action. Let $\Pi_5 = \langle c_1, \dots, c_5 \mid c_1 \cdots c_5 = 1 \rangle$ be the discrete fundamental group of the sphere with five marked points removed, with the chosen positively oriented peripheral loops. Let $\mathcal{X}$ be the set of epimorphisms $\Pi_5 \twoheadrightarrow A_5$ sending each c_i to an involution, modulo postcomposition by $\mathrm{Aut}(A_5)$. Precomposition with the peripheral outer mapping-class action defines a permutation of this set. The ambiguity from a based inner automorphism is absorbed by an inner automorphism of A_5. Thus this construction defines an actual action of $\Gamma_{0,[5]}$ on $\mathcal{X}$.

We use $\mathrm{Aut}(A_5) = S_5$ with its actual conjugation action, proved through its faithful permutation of the five Sylow C_2^2 subgroups in E.29. Enumerating the fifteen involutions gives 11820 ordered product-one five-tuples. Exactly 300 generate a C_2^2 subgroup; the remaining 11520 generate A_5. The automorphism action on generating tuples is free, because an automorphism fixing the five entries fixes their generated group. Hence

$$|\mathcal{X}| = 11520/120 = 96. \tag{E.46.6}$$

Every tuple, orbit representative, and transporter is retained. With right actions on the index set, the positive half twist is

$$(g_i, g_{i+1}) \longmapsto (g_i g_{i+1} g_i^{-1}, g_i). \tag{E.46.7}$$

The constructed permutations satisfy all Artin relations and the two literal sphere relations

$$\sigma_1\sigma_2\sigma_3\sigma_4^2\sigma_3\sigma_2\sigma_1 = 1, \qquad (\sigma_1\sigma_2\sigma_3\sigma_4)^5 = 1. \tag{E.46.8}$$

Each generator has cycle type $2^6 3^9 5^{10} 1^7$ and is even. The image of the second word before taking its fifth power has order exactly five. These word checks accompany the topological construction of the map; they fix its conventions.

We prove the full image order by a short exact permutation certificate. Put $u_i = \sigma_i^2$ for $1 \le i \le 4$. The saved word W of length 120 in the signed u_i has

$$W^{410} = (17\ 26\ 43). \tag{E.46.9}$$

The certificate gives 72 explicit conjugates of this three-cycle by signed u_i-words. In the retained order, each new support meets the preceding union, and their union is the entire 96-element set. The deterministic verifier multiplies every printed word and checks each support and permutation. Array composition $g[c[g^{-1}]]$ in that certificate represents the word $g^{-1}cg$ under our right action convention.

Here is the group-theoretic implication of the support check. Start with the alternating group on the first three vertices. A three-cycle with two vertices in an already covered set U and one new vertex, together with A_U, generates $A_{U\cup\{a\}}$: A_U is transitive on unordered pairs for $|U| \ge 3$, and conjugation and inversion give every three-cycle involving the new vertex and two old vertices. These and the three-cycles on U generate that alternating group. A new cycle meeting U in one vertex x has the form $(x\ a\ b)$, after an orientation choice. Choose distinct $y, z \in U \setminus \{x\}$. Conjugating $(x\ y\ z)$ by a suitable orientation of the new cycle gives $(a\ y\ z)$; the preceding argument first adjoins a, and the original cycle then adjoins b. Induction along the saved supports proves $\langle u_1, u_2, u_3, u_4 \rangle = A_{96}$. All these squares are pure. The evenness upper bound therefore proves both image assertions in (E.46.1). This proof uses the explicit three-cycles and does not rely on the initially interrupted generic order calculation.

The reversed sphere word is checked on the same arrays:

$$y_4 = \sigma_3\sigma_2\sigma_1^2\sigma_2\sigma_3, \qquad y_4\sigma_4^2 = 1. \tag{E.46.10}$$

Thus the subgroup generated by the first three half twists contains all four u_i, and equals A_{96}. It is precisely the image of the stated B_4 map.

All evaluation maps and the actual induced source. For $1 \le i < j \le 5$ let

$$A_{ij} = (\sigma_{j-1} \cdots \sigma_{i+1})\sigma_i^2(\sigma_{j-1} \cdots \sigma_{i+1})^{-1} \tag{E.46.11}$$

in this finite target. The first five pairs are

$$\begin{gathered}(A_{12}, A_{23}A_{24}), \quad (A_{13}A_{23}, A_{34}), \quad (A_{23}, A_{34}), \\ (A_{12}A_{13}, A_{24}A_{34}), \quad (A_{12}, A_{23}).\end{gathered} \tag{E.46.12}$$

They are the five literal pentagon substitutions of E.17. For $i = 2, 3, 4$ put

$$y_i = \sigma_{i-1} \cdots \sigma_1^2 \cdots \sigma_{i-1}; \qquad \text{the last three pairs are } (y_i, \sigma_i^2). \tag{E.46.13}$$

These are exactly the evaluations in the based action [NS00, (1.1)]:

$$\sigma_1' = \sigma_1^\lambda, \qquad \sigma_i' = f(y_i, \sigma_i^2)^{-1}\sigma_i^\lambda f(y_i, \sigma_i^2) \quad (2 \le i \le 4). \tag{E.46.14}$$

The order of the two inputs and the conjugation orientation are the exact comparison established in E.29 and E.38.

Each pair gives a homomorphism $\theta_i : F \to A_{96}$, $1 \le i \le 8$, taking x, y to its entries. Choose the retained Schreier section b_q of $B/F \simeq S_3$. For $b \in B$, the induced map has permutation $q \mapsto q\bar{b}$ and coordinate

$$\theta_i(b_q b b_{q\bar{b}}^{-1}) \quad (q \in B/F,\ 1 \le i \le 8). \tag{E.46.15}$$

Cancellation of adjacent section words proves multiplication, so this is a homomorphism of discrete groups and extends continuously to the profinite completion. Its kernel lies in F and is normal in B. On F, its coordinate at the identity coset is the specified θ_i.

The original action has degree 4608. Its 87 transitive orbits comprise nine isomorphism classes of B-sets, of sizes

$$72, 24, 300, 162, 6, 162, 300, 30, 18. \tag{E.46.16}$$

For every deleted orbit a saved bijection intertwines both s and r pointwise with a retained orbit. Deleting only these repeated actions therefore gives a faithful action of the same source image on 1074 points. The exact permutation-group calculation gives (E.46.2).

The complete correction image and its modular map. Put $P = \rho_5(N)$, $E = \rho_5(F')$, and retain C, K from the statement. The full subgroup computations give

$$\begin{gathered}|C| = 2^{15}3^{17}5^{13}7^2, \qquad C' = C, \qquad \rho_5(K_6 \cap F') = C, \\ |G_5/P| = 72, \quad |P/C| = 75, \quad |K/C| = 25, \quad |E/C| = 4, \quad |G_5/C| = 5400.\end{gathered} \tag{E.46.17}$$

All 97 actual N generators are evaluated. The four selected generators, with indices $1, 7, 14, 27$, generate their entire image. Its full derived subgroup is C. The 97 sixth powers give K; the 95 products prescribed by the primitive integral kernel of $N^{\mathrm{ab}} \to F^{\mathrm{ab}}$ give the entire $\rho_5(K_6 \cap F')$. The rank and Smith diagonal (1^{95}) of this kernel are checked, as is its zero product with the actual two-row exponent-sum matrix. All 95 image products lie in C. Normality under both s, r, the second derived image of C, and the complete derived image of $\rho_5(F)$ are recomputed. Thus (E.46.17) concerns full subgroup images.

Map B to $\mathrm{PSL}_2(\mathbb{Z}/6)$ by

$$s \longmapsto \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \qquad r \longmapsto \begin{bmatrix} 0 & -1 \\ 1 & -1 \end{bmatrix}. \tag{E.46.18}$$

Its actual image has 72 elements. Enumerating its Cayley graph gives 144 Schreier words generating the entire kernel. Each of their ρ_5 images lies in P. Both inverse images have index 72 in B, proving equality of the kernels. Reduction modulo three therefore gives the specified homomorphism $q_3 : G_5 \to A_4$, with

$$x \longmapsto \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}, \qquad y \longmapsto \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}. \tag{E.46.19}$$

These generate A_4, so $q_3(E) = V_4$. Since $C \subset P \subset \ker q_3$ and $|E/C| = 4$, we obtain

$$\ker(q_3|_E) = C. \tag{E.46.20}$$

The retained 24 base words have the three matrices and the exact index table of E.29.12–E.29.13. Every old correction word and all six ternary and seven binary fibre-basis words are checked to be identity under (E.46.18). Checking the actual cusp-gauged reconstruction proves that all 864 points have their specified value M_κ under q_3.

Arithmetic selection and finite generation. Fix such a point and its profinite unit λ. The arithmetic construction in E.38 chooses an actual $\sigma \in G_{\mathbb{Q}}$ with $\chi(\sigma) = \lambda$ and the required $\rho_2(\sigma)$ mod 3. The intersection $\mathbb{Q}(\zeta_3, \sqrt[3]{2}) \cap \mathbb{Q}(\mu_\infty) = \mathbb{Q}(\zeta_3)$ proves this freedom in the cyclotomic kernel. Applying Nakamura–Schneps, Theorem 2.5, to this actual element gives $q_3(f_\sigma) = M_\kappa$, with precisely the inputs (E.46.19). For the retained word $f_0 \in \widehat{F}'$,

$$c = \rho_5(f_0)^{-1}\rho_5(f_\sigma) \in C. \tag{E.46.21}$$

The joint image of N' in C and its solvable image in D_{23} is their direct product. Indeed, Goursat's common quotient would be both perfect and solvable, and its derived series forces it to be trivial. There is a discrete $k \in N'$ with these two images equal to c and 1. Thus $f = f_0 k$ retains the entire old point and satisfies $\rho_5(f) = \rho_5(f_\sigma)$.

This equality matches all 48 induced coordinates. It gives all five pentagon values and all three values in (E.46.14) of the actual Galois action. The resulting $B_5 \to A_{96}$ and $B_4 \to A_{96}$ maps are onto: they are the original onto maps precomposed with the corresponding profinite Galois automorphisms. Their relations, including both sphere words, are therefore satisfied. E.39 applied with $n = 5$ and then with q_5 proves (E.46.3). This uses two actual maps from the full braid group; it imposes no unproved invariance of a finite quotient kernel.

Every common quotient of D_{23} and G_5 kills C by perfectness, and then kills K/C of order 25 because D_{23} has only two- and three-primary factors. It factors through the actual J_5. Conversely $B \to J_5$ kills K_6, so it factors through G_6, which D_{23} retains. This proves the maximal common-quotient assertion in (E.46.4). The corrected maps agree there, since their values are the same literal source formulas. Their onto projections and Goursat's lemma give the asserted fibre product.

The group A_{96} is perfect: every three-cycle is a commutator of two double transpositions, using two extra letters, and three-cycles generate A_{96}. Its common quotient with the solvable T_{23} is therefore trivial. Both original four-strand projections are onto, as are both corrected projections. Goursat's lemma proves the actual product $T_{23} \times A_{96}$ in both cases.

The exact solvable quotient and every retained boundary stage. The literal images of x, y in G_5 have order 15. By (E.46.17), the image of F in S has order 900, its derived group has order four, and its abelianization has order 225. The map from $(\mathbb{Z}/15)^2$ taking its two basis vectors to x, y in that abelianization is onto between groups of equal order. It is

consequently the actual abelianization map. Together with (E.46.20) this gives an injection of $\rho_5(F)/C$ into $A_4 \times (\mathbb{Z}/15)^2$. Its exact image is

$$\{(g, a, b) : \kappa_3(g) = a + b \pmod 3\}, \tag{E.46.22}$$

where $\kappa_3 : A_4 \to C_3$ takes both matrices in (E.46.19) to one. The complete 900-element enumeration checks this equality with the generator maps.

For the actual Fermat subgroup $H = \ker(F \to (\mathbb{Z}/3)^2)$, its ten Schreier words give exactly

$$\rho_5(H)C/C = V_4 \times C_5^2. \tag{E.46.23}$$

Here V_4 is the zero-abelianization part of (E.46.22), and the two central order-five generators are the images of x^3, y^3. All 100 elements and all ten generator images are retained; commutativity and exponent ten are checked on the exact model. In particular the literal source map has the factorization

$$B \longrightarrow B/([H, H]H^{10}) \longrightarrow S. \tag{E.46.24}$$

The quotient S is solvable, being an extension of the solvable group in (E.46.22) by S_3; every solvable quotient of G_5 kills the perfect C. Thus S is its maximal solvable quotient. Since $\gamma_3(H)H^{20} \subset [H, H]H^{10}$, the same map factors through the retained P_{20}.

Let $R_H = \bigcap_r \widehat{H}^{[r]}$ be the closed prosolvable residual used in E.43. Perfectness gives $\rho_5(\widehat{N}^{[r]}) = C$ for every $r \geq 1$. The inclusions $\widehat{N}^{[r+1]} \subset \widehat{H}^{[r]}$ and compactness of the nested nonempty fibres over each element of C therefore prove the actual continuous epimorphism

$$\rho_5|_{R_H} : R_H \twoheadrightarrow C. \tag{E.46.25}$$

Its image is contained in C because the H image in S is abelian.

Now fix any compatible multiplier-one boundary word f_0 of E.42, with any E.44 factor included. It has the same D_{23} value as its retained point, so the modular value used in (E.46.21) is unchanged. Choose a preimage $k \in R_H$ of c by (E.46.25). Every finite geometric image in the specified old family is solvable. Hence this single k has identity in each of them and preserves the full compatible family, including every full pro-p boundary component for $p \geq 5$. The corrected word still matches f_σ in G_5.

Every finite old source is solvable, including the class-three factor, since its geometric kernel is solvable and B/H has order 54. For old sources retaining P_{20}, formula (E.46.24) makes S a specified common quotient. Maximal solvability makes it the maximal common quotient. The component source maps are onto by E.42 and E.44 on the old side and by the actual Galois automorphism on the new side. They agree on S by their literal source formulas. This proves (E.46.5) for the original and corrected images. Adjoining P_{20} stays in the displayed boundary family, so these old sources are cofinal within that family.

The same word retains the central-cover Ferrari source. There is a further actual quotient comparison. Let G_A be E.29's full A_5 Ferrari source, with its specified map ρ_A. Form the disjoint union of its 1980-point action and the 1074-point action of G_5, using exactly the two images of s, r in both factors. This represents the actual joint image $J \leq G_A \times G_5$ with its two onto projections. The complete intertwiner compression retains 2538 points. Four new transitive B-set classes remain, so their deletion alone does not prove a factorization. The exact joint group calculation instead gives

$$|J| = |G_A| = 2^{38}3^{26}5^{21}7^2.$$

Thus $J \to G_A$ is bijective, and its inverse followed by the second projection defines an epimorphism

$$\begin{gathered} G_L \longrightarrow G_A \xrightarrow{\psi} G_5, \qquad \psi(\rho_A(b)) = \rho_5(b) \quad (b \in B), \\ j \longmapsto (j, \psi(\pi_A \pi_L(j))). \end{gathered} \tag{E.46.26}$$

Here $G_L \to G_A$ is the actual E.38 central-cover projection, denoted π_A; π_L is the projection from any of E.38, E.43 or E.44's common sources to G_L. The second line is an isomorphism from that common source onto its actual joint image with G_5. Its inverse is the first-coordinate projection. Both maps are continuous in the profinite inverse-limit family. The separate root replay reconstructs the disjoint action, every intertwiner, the exact joint order and the identical strong-generator arrays; its original inputs stay unchanged.

For each simultaneous word supplied by E.38 or E.43–E.44, the equality $\rho_L(f) = \rho_L(f_\sigma)$ holds for the same actual Galois element used in its construction. Applying (E.46.26) gives $\rho_5(f) = \rho_5(f_\sigma)$ with no new correction. It therefore retains every five-point assertion already proved, and its source image is the complete graph in (E.46.26).

Write $T_{\mathrm{old}} = T_{23} \times_{S_L} T_L$ for the four-strand target of E.38. Its order has only the prime divisors $2, 3, 5, 7$, since $|T_L| = 2^{117}3^{36}5^{31}7^2$ and T_{23} has only the prime divisors two and three. Every common quotient of T_{old} and A_{96} therefore kills an eleven-cycle $c = (1\ 2\ \cdots\ 11)$ in A_{96}. Its normal closure is the entire A_{96}: in our right action convention, with $h = (1\ 12\ 13)$, $chc^{-1}h^{-1} = (1\ 13\ 11)$. All three-cycles are conjugate in A_{96} and generate it. Conjugacy here can be checked by extending any prescribed map on the three letters to an even permutation, using two of the unused letters to adjust parity. Thus the common quotient is trivial. The original and corrected four-strand maps both have onto projections, so Goursat gives exactly

$$(T_{23} \times_{S_L} T_L) \times A_{96} \tag{E.46.27}$$

for their joint images. This proves the additional simultaneous assertion with the specified quotient maps.

Certificates and scope. The files with prefix `e_hurwitz_five_a5_` retain the tuple model, pure three-cycle word and conjugates, and deterministic order verification. The files with prefix `e_hurwitz_five_` and suffixes `source_root`, `corrections_root`, `compare_root`, and `boundary_root` retain the entire induced source, all subgroup images, the 144 modular Schreier words, and every element of the two small comparison models. The independent `bc_hurwitz_five_a5_audit` reconstructs the tuples, all transporters, and all pure conjugates. The additional `a5_common_root` certificate with the same `e_hurwitz_five_` prefix gives the entire joint action and the quotient (E.46.26); its independent replay is `audit_hurwitz_a5_common_root`. The interrupted generic order calculation and its recorded interruption are preserved. The assertion proves nonemptiness for these specified sources and their stated boundary products; the full cofinality questions E.2 and E.3 remain open. □

Statement E.47 (Proved). The complete three-adic class-three correction-image fibre and its exact exponent-three image. *Let*

$$\begin{aligned} C_3 &= H^{(3)}/\overline{\gamma_4(H^{(3)})}, \qquad K_3 = \ker(\widehat{H} \longrightarrow C_3), \\ \mathcal{B}_3 &= \widehat{B}/K_3, \qquad \mathscr{J}_3^{\infty} = \mathrm{im}(\widehat{B} \longrightarrow \mathscr{D} \times \mathcal{B}_3). \end{aligned} \tag{E.47.1}$$

Here $H^{(3)}$ is the maximal pro-three quotient, $\mathscr{D}$ is the actual E.24 common source, and $H \triangleleft F \triangleleft B$ and all based maps are those of E.26. Fix any one of E.24's 432 *multiplier-one source values, retaining its literal commutator word.*

The complete set of boundary correction images in $\mathscr{J}_3^\infty$ satisfying the two order relations has an explicit continuous parametrization

$$\mathbb{Z}_3^{70}. \tag{E.47.2}$$

Its projection to the full E.37 class-two fibre is given by the four exact equations (E.47.10) *and the two residue equations* (E.47.15) *below. Every point of this projection has a complete central fibre affine under a free $\mathbb{Z}_3$ module of rank* 64*.*

The image in E.44's actual exponent-three target has exactly 3^{25} elements per source value. Its complete criterion consists of the E.37 reduction conditions, E.44's equations, and the two additional equations (E.47.22)*. All* 24 *retained CM rows, with all* 18 *source points in each row, survive. There is a specified E.44 shadow, compatible with the entire class-two tower, which has no lift to this full class-three tower:*

$$t = 0, \qquad Y_{41} = 1, \qquad Y_j = 0 \quad (37 \le j \le 44,\ j \ne 41). \tag{E.47.3}$$

Its exact finite central value is printed in (E.47.24)*. The failure persists after changing every higher class-two digit.*

All the stated correction images are generating in the actual common finite sources. They are realized by single profinite commutator words, simultaneously with E.41's full two-adic class-two family, E.42's full pro-p factors for $p \ge 5$, the E.43 refinement, and the five-point Hurwitz factor through the exact maps of E.46.26.

Proof. We first give the actual class-three group and the exact source comparison. Let $V = \mathbb{Z}_3^{10}$ with ordered basis $e_1, \dots, e_{10}$ and let $L_3(V)$ be the ordinary degree-three free Lie lattice. Its 330 ordered Lyndon brackets are expanded in the 1000 tensor words of length three; each has leading coefficient one. Use scaled logarithmic coordinates

$$\log g = (v, Z/2, T/12), \qquad v \in V, \quad Z \in \Lambda^2 V, \quad T \in L_3(V). \tag{E.47.4}$$

Multiplication is the following integral polynomial:

$$\begin{aligned}(v, Z, T)(w, W, U) = (&v + w,\ Z + W + v \wedge w,\\ &T + U + 3([v, W] - [w, Z]) + [v - w, v \wedge w]).\end{aligned} \tag{E.47.5}$$

The inverse negates all three coordinates. This formula follows directly by collecting the degree-one, degree-two and degree-three terms of the Baker–Campbell–Hausdorff identity; the displayed factors retain its denominator three.

Here is the precise group lattice in these coordinates. Write the product of the ordered generator powers $\prod_i \eta_i^{v_i}$ as $(v, Z_0(v), T_b(v))$, so $(Z_0)_{ij} = v_i v_j$. Define the integral linear map

$$\begin{aligned}\mathcal{C}(e_i \wedge e_j) &= [e_i, [e_i, e_j]] + [e_j, [e_i, e_j]],\\ T_0(v, Z) &= T_b(v) + 3\mathcal{C}(Z - Z_0) + 3[v, Z - Z_0].\end{aligned} \tag{E.47.6}$$

The actual Hall normal form gives exactly

$$C_3 = \{(v, Z, T) : T - T_0(v, Z) \in 3L_3(V)\}, \qquad \xi = \frac{T - T_0(v, Z)}{12} \in L_3(V). \tag{E.47.7}$$

Indeed the ordered commutator exponents are $(Z - Z_0)/2$; their logarithms and their products with $\prod_i \eta_i^{v_i}$ give (E.47.6). The remaining ordinary Hall commutators of weight three have actual logarithms equal to the Lyndon basis, and hence scaled T coordinates equal to 12 times that basis. Since 2 and 4 are units in $\mathbb{Z}_3$, these expressions give inverse continuous coordinate

maps between the Hall normal form and (E.47.7). They prove the comparison with the actual pro-three quotient, including the asserted congruence. Reduction of the integral coordinates at increasing three-power precision gives a cofinal system of finite continuous images of this group.

The based actions S, R are obtained by substituting the literal words $x \mapsto y, y \mapsto x$ and $x \mapsto y, y \mapsto (xy)^{-1}$ in all ten Schreier words. Their full 385 by 385 matrices in (E.47.4) are integral and satisfy $S^2 = R^3 = 1$ over $\mathbb{Z}$. Their first 55 rows are exactly the E.37 doubled-log action. The last diagonal blocks S_3, R_3 are the ordinary free Lie actions on $L_3(V)$; all off-diagonal blocks from the literal substitutions are retained.

The source image is also exact. The 45 adjusted relators $u_j \in H'$ of E.33 have central images forming a three-adic basis. The 450 literal words

$$[\eta_i, u_j^4] \in \ker(B \longrightarrow \mathscr{D}) \cap \gamma_3(H) \tag{E.47.8}$$

map to a spanning set of $L_3(V)$ modulo three. The retained bracket matrix has rank 330 and specifies a unit 330-column minor. This rank also follows from the free Lie brackets $[V, \Lambda^2 V] = L_3(V)$ and the unit degree-two matrix of the u_j. The fourth powers clear the binary source image and preserve its ternary kernel. Their three-supported continuous powers therefore supply every element of $L_3(V)$ while retaining all class-two factors and all pro-p factors for $p \neq 3$. Together with the exact E.37 source image, this proves that the new source image is its full inverse image under $C_3 \to C_3/\gamma_3(C_3)$. Thus the only affine source restriction is the one already proved in E.37 on v; there is no uncomputed central source condition.

We solve the two order equations with this entire source image. Over an E.37 point (v, Z) put $g_0 = (v, Z, T_0(v, Z))$, and let E_s, E_r be the scaled third coordinates of $(Sg_0)g_0$ and $g_0^{-1}R(g_0^{-1})R^2(g_0^{-1})$. Their lower coordinates vanish. The equations for $g = (v, Z, T_0 + 12\xi)$ are

$$M\xi = b, \qquad M = \begin{pmatrix} 1 + S_3 \\ -1 - R_3 - R_3^2 \end{pmatrix}, \qquad b = -\frac{1}{12}\begin{pmatrix} E_s \\ E_r \end{pmatrix}. \tag{E.47.9}$$

The exact 660 by 330 matrix has Smith factors $1^{264}, 3, 3, 0^{64}$. In particular its rational rank is 266, its integral kernel has rank 64, and its image has index 9 in its three-saturated rational image. The errors lie in $L_3(V)^{S_3=1} \oplus L_3(V)^{R_3=1}$: cyclic permutation of each central norm word gives the same word. The two invariant dimensions are 164 and 106; the traces are -2 and -6. The common dual invariant space has rank four. For its four saved primitive rows Λ, the identities $(3\Lambda, 2\Lambda)M = 0$ give four independent conditions on this 270-dimensional error space. They therefore characterize rational membership in the image of M.

We give those four conditions as small explicit equations on the full E.37 chart. Let $c \in \{0, 1, 2\}$ be the actual affine source residue computed in E.37 and put

$$U = t_1 + t_2 = c + 3w, \quad u = t_2, \quad t_3 = u + 3e, \qquad \Phi = e + U^2 + 3Ue + 9e^2 = 0. \tag{E.47.10a}$$

For every $U \in \mathbb{Z}_3$ there is a unique $e \in \mathbb{Z}_3$ satisfying this equation: $e \equiv -U^2 \pmod 3$ and $\partial\Phi/\partial e \equiv 1 \pmod 3$. The substitution sends the E.37 equation $t_3 - t_2 + 3Q(t) = 0$ exactly to $3\Phi = 0$. Set $a = (Y_{37}, \ldots, Y_{44})^t$. The complete rational criterion is

$$A(U, e)a + h(U, u, e) = 0, \qquad A = A_0 + UA_U + eA_e, \tag{E.47.10}$$

where

$$A_0 = \begin{pmatrix} -1 & -1 & -2 & -1 & -1 & 0 & 0 & 1 \\ 3 & 3 & 6 & 3 & 5 & 0 & 0 & -1 \\ -7 & -7 & -10 & -7 & -13 & 2 & 0 & -1 \\ -4 & -6 & -8 & -4 & -6 & -2 & 2 & 0 \end{pmatrix},$$

$$A_U = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 16 & 24 & 8 & 12 & 0 & -4 & -10 & -8 \\ -30 & -36 & -12 & -24 & 0 & 12 & 12 & 24 \end{pmatrix}, \tag{E.47.11}$$

$$A_e = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 6 & 36 & 12 & 0 & 0 & 12 & -24 & 24 \\ -54 & -108 & -36 & -36 & 0 & 0 & 54 & 0 \end{pmatrix}. \tag{E.47.12}$$

The four constant polynomials, reduced by the displayed monic relation Φ in the variable U, are

$$\begin{aligned} h_1 &= -U + 6e^2 + 6eu + 2u^2 + u, \\ h_2 &= U(12e + 8u + 1) - 6e^2 - 30eu - 14u^2 - u, \\ h_3 &= U(12e^2 - 24eu - 150e - 14u^2 - 76u - 1) \\ &\quad - 306e^3 - 360e^2u - 372e^2 - 102eu^2 + 12eu - 28e + 38u^2 + u, \\ h_4 &= 2\big[U(-63e^2 - 36eu + 39e - 3u^2 + 31u - 1) \\ &\qquad + 135e^3 + 216e^2u + 333e^2 + 72eu^2 + 120eu + 30e + 12u^2 + 3u\big]. \end{aligned} \tag{E.47.13}$$

These equations are obtained by applying the four common dual rows to the exact integral norms, substituting the E.37 Smith origin, and performing the saved invertible rational row change. All row divisions and polynomial multiples of Φ are retained. This gives an equality of zero loci over $\mathbb{Q}_3$, not a discarded integral condition.

Number the columns of A from zero. For $P = (0, 1, 2, 4)$ and $F = (3, 5, 6, 7)$, its pivot minor satisfies the exact identity

$$\det A_P = -16(30\Phi + 1 - 3U - 3e). \tag{E.47.14}$$

It is a three-adic unit at every point of the chart. Thus $a_F = b = (b_0, b_1, b_2, b_3)^t$ is free and $a_P = -A_P^{-1}(h + A_F b)$ gives every rationally soluble integral point.

The two remaining integral Smith tests, after this substitution, are exactly

$$\begin{pmatrix} 2 & 0 & 1 & 1 \\ c+2 & 1 & -c & -c \end{pmatrix} \bar{b} = \rho(c, \bar{u}) \quad \text{in } \mathbb{F}_3^2, \qquad \begin{array}{c|ccc} & \bar{u}=0 & \bar{u}=1 & \bar{u}=2 \\ \hline c=0 & (0,0) & (1,1) & (0,2) \\ c=1 & (1,0) & (0,0) & (0,2) \\ c=2 & (2,0) & (2,0) & (0,1) \end{array}. \tag{E.47.15}$$

For completeness, the two functionals are constructed on the three-saturated image of M, rather than extended from a chosen rational complement. Starting with 266 independent columns of M, two recorded divisions by three give a matrix N whose reduction has rank 266. Hence its columns are a basis of $(\operatorname{im} M_{\mathbb{Q}_3}) \cap \mathbb{Z}_3^{660}$. Choose the recorded 266 rows I for which N_I is invertible over $\mathbb{Z}_3$. The annihilator of the image of $N_I^{-1} M_I$ modulo three has dimension two. Pulling its rows back by N_I^{-1} gives the two saved tests. The Smith factors show that their vanishing is both necessary and sufficient for integral membership. In their polynomial evaluation the substitution $t_3 = u + 3e$ is made before division by three. The resulting two weighted error polynomials have integral coefficients after that division. Their

residues depend only on $(c, \bar{u}, \bar{b})$; the exact coefficient matrices give (E.47.15) in all nine charts.

Every chart is soluble, since the first two columns of the matrix in (E.47.15) have determinant two. More explicitly choose $u, w, b_2, b_3, \ell_0, \ell_1 \in \mathbb{Z}_3$. Use the digits $d_0, d_1 \in \{0, 1, 2\}$ defined by

$$\begin{aligned} d_0 = 2(\rho_1 - \bar{b}_2 - \bar{b}_3), \qquad d_1 = \rho_2 - (c+2)d_0 + c(\bar{b}_2 + \bar{b}_3) \quad \text{in } \mathbb{F}_3, \\ b_0 = d_0 + 3\ell_0, \qquad b_1 = d_1 + 3\ell_1. \end{aligned} \tag{E.47.16}$$

Equations (E.47.10a), (E.47.14), (E.47.16) give a continuous bijection from $\mathbb{Z}_3^6$ onto the entire projected E.37 locus. Its inverse recovers u, w, the four entries of a_F, and divides the last two digit discrepancies by three.

We next supply the central solution, including its integral right inverse. The same two saturation operations give an integral matrix P_0 with

$$MP_0 = 3N. \tag{E.47.17}$$

The saved numerator initially has denominator nine; every numerator entry is divisible by three, giving exactly (E.47.17). Each saturation step enlarges the selected column lattice by index three: the coefficient of the replaced column in the saved relation is a three-adic unit. Thus the original selected-column lattice has index nine in $N\mathbb{Z}_3^{266}$. The full image of M has that same index by its Smith factors and contains the selected columns. The two lattices are therefore equal. This proves that $P_0/3$, on the specified N coordinates, is an integral right inverse exactly on the full image of M. For the right side b in (E.47.9) set

$$\xi_0 = \frac{1}{3} P_0 N_I^{-1} b_I, \qquad \xi = \xi_0 + Kz, \quad z \in \mathbb{Z}_3^{64}, \qquad T = T_0 + 12\xi. \tag{E.47.18}$$

Here K is the saved integral kernel matrix. It satisfies $MK = 0$ and has a unit 64-row minor. Its rational columns span $\ker M$, so these properties prove that its columns form the complete three-adic kernel lattice. On the rational image, $NN_I^{-1}b_I = b$. The tests (E.47.15) state exactly that the numerator of ξ_0 is divisible by three. Consequently (E.47.18) is an actual integral solution of (E.47.9), and subtracting it from any solution gives a unique Kz. This proves the 64-dimensional central fibre and the complete 70-parameter parametrization (E.47.2).

We now determine the precise finite projection. The map to E.44 is the actual Hall/Magnus map

$$\begin{aligned} (v, Z, T) \longmapsto \left(\bar{v},\, 2\bar{Z},\, t_{\mathrm{can}}(\bar{v}, 2\bar{Z}) + \Pi\bar{\xi}\right), \\ \Pi(L)_{ijk} = [X_i X_j X_k]L - [X_j X_i X_k]L \quad (i < j < k). \end{aligned} \tag{E.47.19}$$

It is obtained by expanding the literal Hall normal form. The tensor-coordinate definition is the cube-span quotient proved in E.44, so it respects multiplication and both based actions. The matrix ΠK has rank 21 over $\mathbb{F}_3$; E.44's central norm kernel has rank 22. Thus over each fixed exact E.37 point in the locus just proved, the finite central image is the explicit affine space

$$t_{\mathrm{can}} + \Pi\bar{\xi}_0 + \mathrm{im}(\Pi K), \tag{E.47.20}$$

of size 3^{21}. This is the complete fixed-lower-point comparison.

Changing a higher class-two digit supplies the remaining direction when only the finite E.44 point is prescribed. The central error is affine in Z; write its coefficient after division by three as $C(t)$. The exact rational graph has Z-derivative $D = V_{M[:,37:44]}\, \partial a/\partial b$. Changing b to $b + 3d$ keeps all integral tests and all class-two residues. Formula (E.47.18) gives its central derivative modulo three as

$$-P_0 N_I^{-1} (C(t)D)_I. \tag{E.47.21}$$

The factor $1/4$ from (E.47.9) has residue one. A saved linear functional that annihilates ΠK evaluates the four projected columns of (E.47.21) as $(2,0,1,1)$ in each of the nine charts. In particular changing ℓ_0 by one supplies a complement to $\mathrm{im}(\Pi K)$. The full finite central kernel is therefore obtained, using this actual change of lower parameters and then the actual central kernel corrections.

It remains to identify which finite lower points occur. On E.44's locus compatible with E.37, the four rational residue equations and the two integral tests add exactly two independent equations. With $a = (\bar{Y}_{37}, \dots, \bar{Y}_{44})^t$, they are

$$\begin{aligned} &\begin{pmatrix} 2 & 2 & 1 & 2 & 2 & 0 & 0 & 1 \\ c+2 & 2 & 2c+2 & 2 & 2 & 2c+2 & 2c & c+2 \end{pmatrix} a \\ &= -\begin{pmatrix} 2c + 2\bar{u}^2 + \bar{u} \\ (c+2)\bar{u}^2 + (2c+1)\bar{u} + c^2 + 2c \end{pmatrix} \quad \text{in } \mathbb{F}_3^2. \end{aligned} \tag{E.47.22}$$

These are rows one and three of (E.47.10) reduced modulo three. For each of the three values of $\bar{u}$, their joint rank with E.44.14 is six, and the joint affine system has nine solutions for a. Formula (E.47.16) realizes all of them. There are consequently 3^3 lower images per source, each with the entire 3^{22} finite central fibre by (E.47.21). This proves the exact count 3^{25} and the complete image criterion. For comparison, E.44 has 3^{28} points per source, of which 3^{27} are compatible with the full E.37 tower.

The new nonlifting example has the integral E.37 central coordinate

$$Z_* = -e_1 \wedge e_7 - e_2 \wedge e_7 - e_3 \wedge e_6 - e_3 \wedge e_{10} - e_4 \wedge e_7 - e_5 \wedge e_7 + e_7 \wedge e_8 + e_7 \wedge e_9. \tag{E.47.23}$$

It is the column $V_M e_{41}$, so it has exactly the free Smith coordinates in (E.47.3). Use the actual source (row 0; ternary $(0,0)$; binary 0), whose affine residue is zero. In the E.44 coordinates the finite point is $(0, 2\bar{Z}_*, t_*)$, where the following array lists t_* consecutively in the lexicographic order of the triples $1 \le i < j < k \le 10$:

$$\begin{pmatrix} 0 & 2 & 2 & 0 & 2 & 2 & 0 & 2 & 2 & 0 \\ 1 & 2 & 2 & 0 & 1 & 1 & 0 & 1 & 2 & 2 \\ 1 & 1 & 1 & 1 & 0 & 2 & 1 & 0 & 2 & 0 \\ 1 & 1 & 0 & 0 & 2 & 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 & 2 & 2 & 1 & 0 & 0 & 2 \\ 2 & 0 & 2 & 2 & 0 & 0 & 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 0 & 2 & 1 & 1 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 2 & 1 & 0 & 0 & 0 & 2 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}. \tag{E.47.24}$$

Both E.44 norms vanish exactly. Its full class-two tower point satisfies E.44.14, so E.44's literal central corrections realize it in the actual common source, retaining that whole tower. In (E.47.22) its left side is $(2,2)^t$ and its right side is zero. Since these are conditions on the finite lower coordinates, no change of any higher digit can give a full class-three lift. This proves the stated failure for an actual peripheral generating shadow.

We finish with the simultaneous source and generation assertions. The new source-kernel equality proved using (E.47.8) lifts every prescribed E.37 point to every permitted class-three central value. The corresponding three-supported powers preserve the whole two-adic class-two family and all full pro-p factors with $p \neq 3$. Compactness of the closed source-kernel

image, or the explicit products of the 330 basis words with continuous exponents, supplies a single profinite word for every compatible family.

At every finite three-primary level the image of N_B in the new factor is a three-group: its image in F/H is cyclic of order three and its intersection with the geometric kernel is a finite three-group. Together with $\mathscr{D}$ its image is finite nilpotent of primes two and three, generated by the 97 fixed generators and mapping onto $(\mathbb{Z}/6)^{97}$ through G_6. The retained G_6 shadow is an isomorphism on both Sylow Frattini quotients, so the corrected map is onto this actual common N_B image. It is then onto the common source. With the two-adic boundary factor, take a lower-central index larger than the nilpotence classes of all the finite three-primary images and the old Ferrari image. E.41's exact stable two-primary lower-central subgroup is supplied as a pure factor. The quotient is the nilpotent common image just considered. This proves joint generation with the full E.41 tower at every finite level. E.42's actual coprime H-image product gives the remaining full pro-p factors and joint generation.

For E.43 take cofinal old common sources containing P_{20}. Their new H images are solvable, and B/H is solvable. Their actual common quotient with G_L is therefore exactly $S = G_L/C$, with the specified $P_{20} \to S$ map, and the actual joint source is $G_{\mathrm{old}} \times_S G_L$. The surjection $\rho_L(R_H) = C$ of E.43 supplies a correction in $R_H = \bigcap_j H^{[j]}$ matching the Galois word in G_L. It preserves every new class-three image. The exact common-quotient argument gives generation of this fibre product, as in E.44.15. The further maps are the proved maps

$$\begin{gathered} G_L \longrightarrow G_A \xrightarrow{\psi} G_5, \qquad \psi\rho_A = \rho_5, \\ j \longmapsto (j, \psi\pi_A\pi_L(j)). \end{gathered} \tag{E.47.25}$$

of E.46.26. They identify the old joint source with its actual joint image after adjoining G_5. The same corrected word therefore retains the five-point Hurwitz value and the full common four-strand target of E.46.27.

Precisely, (E.47.2) parametrizes the images under the actual map $\rho : \widehat{F}' \to \mathscr{D} \times \mathcal{B}_3$ with its prescribed first coordinate. For each such image and any one word f realizing it, the set of actual profinite words realizing it is exactly

$$f\ \ker(\rho|_{\widehat{F}'}). \tag{E.47.26}$$

This follows by multiplying two preimages and is the complete comparison between correction-image fibres and word fibres. Adjoining the simultaneous target maps replaces the kernel in this same formula by their joint kernel.

The eight new programs in `code` have shared prefix `bc_fermat_class_three_` and suffixes `integral.py`, `norm_smith_probe.py`, `rational_obstruction.py`, `curve_obstruction.py`, `integral_tests.py`, `graph_tests.py`, `fibres.py`, and `finite_image.py`. Their executed certificates retain the tensor/Hall comparison, the actual source word preimages, the Smith matrix and saturation operations, every polynomial row change, the central right inverse, all nine residue systems, the full 432-point coverage, and the exact nonlifting value. All have status `PASS`. □

5.11 The full A_6 Ferrari source, its solvable layers, and the cubic Kummer comparison

Statement E.48 (Proved). Every retained D_{23} fibre survives the full A_6 Ferrari test. *Use the actual groups $B = C_2 * C_3 = \langle s, r\rangle$, F, N, $K_6 = N'N^6$, and D_{23} of E.24. The 53 epikernels of E.45 give a finite Ferrari quotient*

$$\widehat{B}_4 \twoheadrightarrow T_6, \qquad |T_6| = 2^{287}3^{127}5^{86}7^6 11^3 13^2. \tag{E.48.1}$$

Its full free Ferrari image is A_6^{53}. Retaining the six specified evaluation pairs and all six B/F cosets gives an actual epimorphism

$$\rho : B \twoheadrightarrow G, \qquad |G| = 2^{218}3^{84}5^{62}7^{6}11^{3}13^{2}. \tag{E.48.2}$$

Put $P = \rho(N)$, $E = \rho(F')$, and $E_1 = \rho(F'')$. There is an explicitly certified perfect normal subgroup C of G such that the following are the actual quotient maps:

$$\begin{gathered} G/\rho(F) \simeq S_3, \qquad \rho(F)/E \simeq (\mathbf{Z}/120)^2, \qquad E/E_1 \simeq \mathbf{F}_3^9, \\ E_1/C \simeq \mathbf{F}_2^6, \qquad |C| = 2^{205}3^{72}5^{60}7^{6}11^{3}13^{2}, \\ |G/C| = 108839116800, \qquad E/P' \simeq \mathbf{F}_3^3, \qquad P' = \rho(N') = \rho(\gamma_j N) \quad (j \geq 2). \end{gathered} \tag{E.48.3}$$

In the first abelian quotient the coordinates are the ordinary x, y exponent sums. The subgroup C is the kernel of the maximum solvable quotient of this specified source.

Every one of the 864 *retained E.24 points, and every compatible profinite multiplier $\lambda \equiv 1 \pmod{72}$, admits a commutator word f with the same D_{23} coordinate and with*

$$\rho(f) = \rho(f_\sigma) \tag{E.48.4}$$

for an actual Galois element σ with $\chi(\sigma) = \lambda$. The correction from the retained word can be chosen in the actual subgroup $\gamma_5 N$. Consequently all six Ferrari evaluations, the pentagon, the Artin relations, the full twists, and the periodic-conjugacy tests in T_6 agree with the corresponding Galois action. The new four-strand map is onto.

The source and target comparisons in this assertion are

$$\begin{gathered} B/(\ker \rho_{D_{23}} \cap \ker \rho) \;\longrightarrow\; D_{23} \times G, \\ T_{23} \times_{C_2} T_6, \end{gathered} \tag{E.48.5}$$

with the first group identified with the image of its displayed map, and the second fibre product taken over actual braid parity. Both corrected source and target maps are onto these same groups. For each fixed old point and multiplier, the complete fibre in G/C contains 18 *norm solutions, exactly* 16 *of which generate the source. All these values and their correction preimages are retained in the certificates.*

Proof. We retain the literal source maps at every finite layer and then lift the complete old fibres through their exact common correction image.

The full finite model and the induced source. The A_6 and $\mathrm{Aut}(A_6)$ tables, the 53 epikernels, and all Nielsen transporters are those of E.45. The action on the ten actual Sylow-three subgroups of A_6 is faithful on $\mathrm{Aut}(A_6)$; all 1440^2 composition pairs are checked. For each factor the holomorph is embedded by the literal map

$$(k, \alpha) \longmapsto (\mathrm{Inn}(k)\alpha, \alpha). \tag{E.48.6}$$

It acts on two ten-point blocks. Taking all 53 factors gives the faithful 1060-point model. The six nonidentity A_6 conjugacy-class normal closures are the full group. The distinct epikernels and the simple-factor Goursat argument therefore give the full image A_6^{53} of the free Ferrari group. The retained Nielsen image has order $2^{128}3^{21}5^{33}7^{6}11^{3}13^{2}$. These exact factors prove (E.48.1). The three standard braid generators have order 60; $(ab)^3$ has order 40, and the central element $(abc)^4$ has order 20.

Use the six pairs of E.29, including $(A_{12}, A_{23}A_{24})$, $(A_{13}A_{23}, A_{34})$, (A_{23}, A_{34}), $(A_{12}A_{13}, A_{24}A_{34})$, (A_{12}, A_{23}), and $((ab)^3, A_{34})$. The six-coset induction formula is

the same actual formula as in E.38; it makes each evaluation a homomorphism on the pure subgroup and produces a homomorphism from B after induction. Its original permutation degree is $36 \cdot 1060 = 38160$. The 348 transitive orbits reduce to 55 distinct B-sets, of total degree 9462. Every removed orbit has a saved bijection intertwining both s and r with its retained copy. The final reordering of the retained points is a saved permutation with its inverse. Thus this compression preserves the kernel of the source map.

The source order in (E.48.2) is verified by a complete Schreier stabilizer chain. Every strong generator has an actual original-generator word in a saved directed acyclic graph. The final check sifts all 865049 Schreier relations. A fresh replay reconstructs the chain, checks every word, and repeats the complete final relation check. A separately implemented NumPy word replay and the reconstruction of all 36 induced components give independent comparisons with the supplied literal source arrays.

The successive actual subgroup images. The same method is applied to all 97 literal generators of N and to the two literal free generators x, y. Derived-subgroup certificates retain the input commutators, normal conjugation checks, and word membership in the derived subgroup. They give the following exact orders:

actual image	order
$P = \rho(N)$	$2^{215}3^{82}5^{62}7^{6}11^{3}13^{2}$
$E = \rho(F')$	$2^{211}3^{81}5^{60}7^{6}11^{3}13^{2}$
$P' = \rho(N')$	$2^{211}3^{78}5^{60}7^{6}11^{3}13^{2}$
$E_1 = \rho(F'')$	$2^{211}3^{72}5^{60}7^{6}11^{3}13^{2}$
C	$2^{205}3^{72}5^{60}7^{6}11^{3}13^{2}$.

(E.48.7)

These identifications include their source words. For example, the six selected N words are checked to generate the entire image of all 97 words before their derived image is used. The nine selected E words are checked to generate the entire image E before computing E_1.

The subgroup C is first obtained as the derived group of ten specified words in E_1. All 664 conjugates of its strong generators by the two actual source generators lie in C. A second derived computation, on ten specified words in C, has exactly the same order as C. The recorded word inclusions give $[\langle w_1, \dots, w_{10}\rangle, \langle w_1, \dots, w_{10}\rangle] \subset C' \subset C$; equality of the certified orders proves $C' = C$. The second final relation check has 720615 relations and has also been freshly replayed. Thus C is an actual perfect normal subgroup.

The quotient G/P is obtained by exact canonical-coset comparison. It has order 72. The map from the specified 576-element modular quotient $Q = B/N$ is checked on all 1152 Cayley-generator edges. Its pure image is abelian of order 12, with the actual exponent lattice

$$L = \begin{pmatrix} 6 & 4 \\ 0 & 2 \end{pmatrix} \mathbf{Z}^2. \tag{E.48.8}$$

This proves $G/P = B/(F'N)$, with the indicated quotient map, and in particular $E \subset P$. The actual elements x, y in G have order 120. Together with $[G : E] = 6 \cdot 120^2$, this identifies $\rho(F)/E$ with the ordinary exponent quotient $(\mathbf{Z}/120)^2$. It also identifies P as the inverse image of $L/120\mathbf{Z}^2$ under that exponent map.

Nine explicit basis cubes are killed in E_1, and the exact index $[E : E_1] = 3^9$ identifies E/E_1 with $\mathbf{F}_3^9$. The actual conjugation matrices of s, r, x, y and the actual coordinate of $[x, y]$ are obtained by permutation sifting. The 64 cosets of C in E_1 are independently enumerated; all quotient generators have square one and commute. Six saved words give a basis of $E_1/C = \mathbf{F}_2^6$. The program checks their squares, commutators, all 64 distinct cosets, and their conjugates by s, r, x, y and the nine E words. It also finds nine commuting order-three lifts of the E/E_1

basis, by testing all 64 lifts of each required basis value. These data give the exact collected normal forms

$$x^a y^b e_v w, \qquad a, b \in \mathbf{Z}/120, \quad v \in \mathbf{F}_3^9, \quad w \in \mathbf{F}_2^6 \tag{E.48.9}$$

in the pure part of G/C. The displayed commutation and power relations come from actual permutation equalities modulo C; their normal-form count equals the actual quotient order. They therefore present this quotient. The six B/F representatives and their literal extension cocycle give the complete B quotient, including all 216 cocycle checks. Since this quotient is solvable and C is perfect, C is exactly the kernel of the maximum solvable quotient.

The complete correction image over the old source. Write $V = E/E_1$ and $W = E_1/C$. In the retained V basis, let X, Y, S, R be conjugation by x, y, s, r, respectively, and let c represent $[x, y]$. The matrices X, Y commute and have order four. The cyclic module generated by c has dimension nine; its seven relations among the 16 monomials $X^i Y^j c$ $(0 \le i, j < 4)$ are retained exactly.

The complete 6668-relator presentation of E.31 for $\ker(B \to D_6) \cap F'$ is evaluated in V and closed under the literal S, R actions. Its image has dimension six. Squaring each actual relator kills its entire binary K_6 image and multiplies its ternary value by the invertible scalar two. This gives the complete image

$$V_6 = \rho_V(\ker \rho_{D_{23}} \cap F') = \rho_V(N'), \qquad \dim V_6 = 6. \tag{E.48.10}$$

The six independent squared relators are retained as literal B words and as literal F words. Their two old source images have been independently replayed in all 36 components of each characteristic. Their lengths are $628, 672, 612, 1032, 10160, 13696$.

The 18 actual $K_6 \cap F'$ basis words in the old ternary presentation also have image in V_6. Together with its complete kernel presentation this proves that $K_6 \cap F'$ has image V_6 in V. The full N' image modulo C has order $3^6 2^6$ by (E.48.7). Actual N-commutator words and their retained products give all six binary coordinates of W inside that image. Successive commutators with the actual N words still span all six coordinates at lower-central degrees three, four, and five. The six degree-five preimages have lengths $4746, 4754, 4682, 4682, 6440, 6438$. Both their nested word expressions and their expanded literal words have identity image in both old source components. Their lower-central membership is checked on the word graph: a commutator adds the two source degrees, and products and inverses retain their minimum degree.

The remaining ternary commutator calculation is

$$\operatorname{span}_{\mathbf{F}_3}((X^6 - 1)V_6 + (X^4 Y^2 - 1)V_6) = V_6. \tag{E.48.11}$$

The exponents in this formula are exactly the two generators of the lattice (E.48.8). The 4656 literal N commutators give the same V_6. The perfect subgroup C lies in P' because $C \subset E_1 = [E, E]$ and $E \subset P$. It therefore lies in $[P, P']$, since $C = [C, C]$. The full binary degree-five image and (E.48.11) show that $[P, P']$ maps onto P'/C. Hence $[P, P'] = P'$, and induction proves the stable lower-central equality in (E.48.3).

E.24 proves $\gamma_5 N \subset \ker \rho_{D_{23}}$. The reverse bound in (E.48.10), including the 18-word comparison, now gives the exact full-source formula

$$\rho(\ker \rho_{D_{23}} \cap F') = P' = \rho(\gamma_5 N). \tag{E.48.12}$$

This identity concerns the entire correction fibre. It is not obtained by restricting to a selected representative.

The quotient coefficient and its sign. The following matrix, acting on the retained V coordinates, is the actual quotient map $E \to E/P'$:

$$Q_3 = \begin{pmatrix} 2 & 0 & 0 & 2 & 0 & 1 & 0 & 0 & 0 \\ 2 & 1 & 1 & 2 & 0 & 0 & 0 & 1 & 0 \\ 2 & 0 & 0 & 2 & 1 & 0 & 1 & 0 & 1 \end{pmatrix} \pmod 3. \tag{E.48.13}$$

Its kernel is V_6. Define c_2 by the actual index-two cover $\ker(F \to C_2)$, where x maps to the nonidentity element and y maps to the identity. The Schreier generators are x^2, y, xyx^{-1}. The coefficient c_2 is the ordinary coefficient of y in its abelianization; the other y-coefficient is its negative on F'. The literal two-coset collector has multiplication

$$(\epsilon, u_0, u_1)(\eta, v_0, v_1) = \big(\epsilon + \eta, (u_0, u_1) + \mathrm{swap}^{\epsilon}(v_0, v_1)\big), \tag{E.48.14}$$

with the last two coordinates modulo three. It sends x to $(1, 0, 0)$ and y to $(0, 1, 0)$. Evaluation of the nine actual E words gives

$$c_2(f) = h\rho_V(f), \qquad h = (1, 2, 2, 1, 2, 1, 2, 2, 2) \pmod 3. \tag{E.48.15}$$

In particular $hc = -1$, $hX = -h$, $hY = h$, and $hV_6 = 0$. These identities, the actual cyclic-module relations, and the powers $x^{120} = y^{120} = 1$ give the homomorphism from the collected quotient to (E.48.14). Thus (E.48.15) holds for every source word, with the stated cover and coefficient map.

In the convention $x \mapsto x^{\chi}$, $y \mapsto f_\sigma^{-1} y^{\chi} f_\sigma$ and $[x, y] = xyx^{-1}y^{-1}$, Nakamura–Schneps write $f_\sigma \equiv A_\sigma(\bar{x}, \bar{y}) * [x, y]$ in the metabelian commutator module. Their equation (2.4) and Lemma 2.3 give $\kappa_2(\sigma) = -A_\sigma(-1, 1)$ for the positive-root Kummer character [NS00, §2, (2.4), Lemma 2.3]. Our explicit cover has $c_2([x, y]) = -1$; x-conjugation negates this coefficient and y-conjugation fixes it. Consequently the comparison, with its sign, is

$$c_2(f_\sigma) = \kappa_2(\sigma) \pmod 3. \tag{E.48.16}$$

Only this Galois comparison is used here.

Every norm fibre, including the final binary layer. In the quotient with derived module V, the two complete source norm equations are

$$(1 + S)v = 0, \qquad (1 + R + R^2)v = 0. \tag{E.48.17}$$

Their combined rank is six, so there are 27 values. The normalized action on the cyclic derived module sends

$$c \longmapsto c + (X - 1)(Y - 1)v. \tag{E.48.18}$$

Exactly 24 of these values give rank nine; the other three give rank three. On the complete norm kernel the exact quotient relation is

$$Q_3 v = \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix} hv. \tag{E.48.19}$$

Thus its quotient value is determined by c_2, and every value of c_2 mod 3 occurs.

The full collected quotient (E.48.9) retains every binary lift $w \in W$ of every one of these 27 values. For $f = (v, w)$ put $S_f = sf$ and $R_f = S_f^{-1}(sr)^{\lambda}$. The program keeps the companion-product identity $S_f R_f = (sr)^{\lambda}$ and solves both order equations in the entire six-dimensional binary kernel. The exact binary norm matrix has two solutions over each

of the 27 ternary points. For all 48 values above the rank-nine ternary points, nine literal F' words have rank nine in V, and six literal F'' words have rank six in W. Their images therefore contain all of E/C. The ordinary exponent map is multiplication by the unit λ, so these maps generate all of G/C. The remaining six norm values have derived image of rank three in V, by the cyclic calculation (E.48.18), and do not generate it.

All 48 generating maps are also checked on the complete finite presentation of (E.48.9). The 8496 checks include complement cubes and commutators, all complement actions on W, the x, y actions on both layers, the actual value of $[x, y]$, and the two order-120 relations. Thus these particular maps are automorphisms of G/C. This conclusion is made for this explicitly presented quotient.

The actual cusp order is 240; the old cusp period is 144. All compatible multiplier classes in the common period 720 are retained:

$$1, 73, 217, 289, 361, 433, 577, 649 \pmod{720}. \tag{E.48.20}$$

The actual elements x^{12}, y^{12} are central in the pure quotient, and $(xy)^{12} = x^{12}y^{12}$ there. This verifies the halved-exponent comparison as well as the ordinary powers. All eight classes have the same 54 norm values and the same 48 generating values. The three new braid generators have order 60, so this period also covers every power in their standard GT formulas.

Every old retained point has one fixed quotient value in (E.48.19). The complete image (E.48.12) gives all 18 norm values above it, of which exactly 16 generate. Among the 864 old points, the three scalar values occur 288 times each. Only the 288 points with scalar zero permit new value $f = 1$; the complete fibres retain the other 576 points as well. The literal comparison with E.38's positive-root labels gives the following common index sequence for all 24 old fibres:

$$\begin{aligned}(&0, 0, 0, 0, 1, 1, 1, 1, 2, 2, 2, 2,\\ &2, 2, 2, 2, 0, 0, 0, 0, 1, 1, 1, 1).\end{aligned} \tag{E.48.21}$$

This comparison evaluates the same literal F word in the double cover and in $\mathrm{SL}_2(\mathbf{Z}/12)$. All six ternary and seven binary old fibre-basis words have zero coefficient and identity matrix, so the comparison holds for every one of the 864 points.

The Galois selection and the full correction. The field $\mathbf{Q}(\zeta_3, \sqrt[3]{2})$ has group S_3 over $\mathbf{Q}$ and maximum abelian subfield $\mathbf{Q}(\zeta_3)$. Since the cyclotomic extension is abelian, its intersection with this field is exactly $\mathbf{Q}(\zeta_3)$. Therefore $\kappa_2 \bmod 3$ maps the cyclotomic kernel onto $\mathbf{Z}/3$. Cyclotomic surjectivity first supplies σ_0 with the specified full character λ; multiplication by an element of its kernel then gives any required value of $\kappa_2 \bmod 3$. This is the same positive-root normalization used in E.38.

For the old word f_0 choose that value to be $c_2(f_0)$. Both f_0 and f_σ have the norm quotient value in (E.48.19). Equations (E.48.16) and (E.48.19) therefore give

$$\rho(f_0)^{-1}\rho(f_\sigma) \in P'. \tag{E.48.22}$$

By (E.48.12) it has an actual preimage $k \in \gamma_5 N$, killed by D_{23}. Taking $f = f_0 k$ proves (E.48.4) and retains every old coordinate. All 36 induced coordinates agree with the same Galois word. The new braid map is consequently the original quotient of $\widehat{B}_4$ precomposed with the actual Galois automorphism given by Nakamura–Schneps (1.1). This proves its relations and surjectivity without an invariance assertion about the kernel of ρ.

Generation on the actual common source. First the new Galois source map takes N onto P. In G/C its norm value is one of the 48 generating values above: the actual Galois map is onto, while the other six values have rank-three derived image. The full-presentation check shows that this map is an automorphism of G/C. It preserves the image of N, since that image is the inverse image of $L/120\mathbf{Z}^2$ under ordinary abelianization and the multiplier is a unit. Its full image of N in G is normal, and the quotient is an image of the solvable group $B/N = Q$. The perfect subgroup C is therefore killed in this quotient. Surjectivity modulo C now proves that the full N image is exactly P.

Let J be the actual first source in (E.48.5), let J_N be its actual N image, and let P_{23} be the N image in D_{23}. The group P_{23} is finite nilpotent, with primes only two and three, and retains $N^{\mathrm{ab}}/6$. Both the original and corrected images of N are subdirect in $P_{23} \times P$. Every nilpotent quotient of P kills its stable lower-central subgroup P'. Goursat therefore puts $\{1\} \times P'$ in both images. The quotient $J_N/(\{1\} \times P')$ is finite nilpotent, being a subgroup of $P_{23} \times P/P'$.

For primes two and three its Frattini quotient is exactly N^{ab}/p: it is a quotient of that rank-97 vector space, and projects onto the full rank-97 quotient retained by P_{23}. The old corrected source maps are invertible on both of these vector spaces, by E.24. For the prime five the old factor is trivial, and the remaining quotient is exactly the ordinary exponent C_5^2; the lattice (E.48.8) has determinant 12, a unit modulo five. The corrected map on this factor is multiplication by λ, hence is invertible. The finite nilpotent Frattini criterion gives the entire quotient $J_N/(\{1\} \times P')$ in the corrected image. It already contains $\{1\} \times P'$, so its N image is all of J_N. Finally the old map on $J/J_N = Q$ is the actual cusp-gauge conjugation of E.24 and is onto. The corrected source map is therefore onto J. Only the specified epimorphisms from B have been used for the old and the full new source.

The common four-strand target. The Nielsen image in (E.48.6) has a perfect derived subgroup of index two, by its two complete derived-word certificates. The kernel A_6^{53} is perfect, so T_6' is perfect and the maximum solvable quotient of T_6 is the actual braid parity C_2. The old target T_{23} is solvable and retains that same parity quotient. The original joint braid image is therefore exactly $T_{23} \times_{C_2} T_6$ by Goursat. The corrected projections are both onto and have the same parity values, so the same argument proves generation of this target.

The replay guide in the research directory lists the complete source-word and quotient certificates: `DE_a6_replay.md`. Their program prefixes are `de_ferrari_a6_` and `de_a6_`; the source model and both successive kernel layers are independently replayable. This statement retains the full old D_{23} fibre. The scalar equality (E.48.21) also permits the same arithmetic choice as E.38; a simultaneous assertion with that earlier nonsolvable source additionally requires its actual combined-source comparison. □

5.12 A five-strand symplectic test with its exact old-source comparison

Statement E.49 (Proved). Every retained CM fibre passes the complete five-strand symplectic test at three. *There is an explicitly specified epimorphism*

$$q_3 : B_5 \twoheadrightarrow \mathrm{Sp}_4(\mathbb{F}_3), \qquad |\mathrm{Sp}_4(\mathbb{F}_3)| = 51840, \tag{E.49.1}$$

*whose four braid generators are the chain transvections below. Its restriction to the first four strands factors through an explicit isomorphism from the actual order-*648 *target of E.18. The complete induced $B = C_2 * C_3$ source for the five pentagon evaluations and the three braid-action evaluations has order* 46656*. Its specified source map factors through G_6.*

All 864 *E.24 points satisfy the pentagon and every five-strand braid relation in this target. Each resulting five-strand map is onto the full group in* (E.49.1)*, preserves the full twist, and*

has the two periodic conjugacies of E.39 in that group itself. This holds for the two profinite multiplier representatives of E.38, and for its whole multiplier range $\lambda \equiv 1 \pmod{72}$.

Every later retained word with these specified G_6 *values has the same eight evaluations. Thus all the simultaneous families of E.43, E.44 and E.47 retain this test without an additional correction. On the first four strands the joint target with each old target retaining E.18 is exactly the graph of the displayed quotient map.*

Proof. We specify the action convention as well as the finite maps. Put

$$J = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & -1 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}, \qquad A_i = I + e_i(e_i^t J) \quad (1 \le i \le 4). \tag{E.49.2}$$

The determinant of J is one. The matrices satisfy $A_i^t J A_i = J$ and $A_i^3 = I$ over $\mathbb{F}_3$. Use the 80 nonzero row vectors, with right action $v \mapsto vA_i^t$, and denote the resulting permutations by a_i. Right permutation multiplication therefore agrees with right multiplication by the matrices A_i^t. These preserve the alternating pairing vJw^t. All adjacent Artin relations and the distant commutations are evaluated as literal permutations; equivalently they follow by multiplying the indicated transvections with adjacent pairing one. Consequently $\sigma_i \mapsto a_i$ is an actual homomorphism from the discrete braid group, with its unique continuous extension from $\widehat{B}_5$.

The complete permutation-group calculation gives order 51840. An independent enumeration of all generated four-by-four matrices gives the same order and retains a word preimage of every matrix. The symplectic-basis count is

$$(3^4 - 1)3^3(3^2 - 1)3 = 51840. \tag{E.49.3}$$

Indeed choose a nonzero first vector, then its partner with pairing one, and then a symplectic basis of their two-dimensional orthogonal complement. A symplectic matrix is determined by such an ordered basis. The faithful vector action and equality of these orders prove the epimorphism (E.49.1). The same arrays give

$$\begin{gathered} d = a_1a_2a_3a_4, \quad e = a_1d, \qquad d^5 = e^4 = -I, \\ |d| = 10, \qquad |e| = 8. \end{gathered} \tag{E.49.4}$$

Here is the comparison with the old four-strand target, retaining its generators. Let $\widetilde{H}_3$ have E.18's actual Heisenberg and matrix coordinates, and its specified generating triple (a, b, c). Starting at the identity, enumerate $\widetilde{H}_3$ by right multiplication by these three elements. In parallel multiply (a_1, a_2, a_3) along the same words. The 1944 generator edges all have the same value at each repeated vertex. Exactly 648 vertices and 648 distinct permutations occur. Thus

$$\theta : \widetilde{H}_3 \xrightarrow{\sim} \langle a_1, a_2, a_3 \rangle, \qquad (a, b, c) \longmapsto (a_1, a_2, a_3) \tag{E.49.5}$$

is a well-defined onto homomorphism with trivial kernel. Every vertex, image and edge is retained in the certificate. For any old four-strand target T with its proved map $\pi : T \to \widetilde{H}_3$, the actual joint image is therefore $\{(t, \theta\pi(t)) : t \in T\}$. Both the original and corrected generating maps have this graph image, since their evaluations commute with (E.49.5).

For the full five-strand map set

$$A_{ij} = a_{j-1} \cdots a_{i+1} a_i^2 a_{i+1}^{-1} \cdots a_{j-1}^{-1}.$$

The five pentagon pairs are

$$\begin{gathered} (A_{12}, A_{23}A_{24}), \quad (A_{13}A_{23}, A_{34}), \quad (A_{23}, A_{34}), \\ (A_{12}A_{13}, A_{24}A_{34}), \quad (A_{12}, A_{23}). \end{gathered} \tag{E.49.6}$$

For $i = 2, 3, 4$ append

$$(y_i, a_i^2), \qquad y_i = a_{i-1} \cdots a_2 a_1^2 a_2 \cdots a_{i-1}. \tag{E.49.7}$$

Every pair defines an actual evaluation homomorphism from $\widehat{F}_2$. Induce all eight maps over all six B/F_2 cosets, with the literal based Schreier maps of E.46. This defines $\rho_3 : B \to G_3$ as an actual permutation representation of degree 3840. Explicit transitive B-set intertwiners compress it faithfully to degree 558; the source images of s, r satisfy $s^2 = r^3 = 1$ and have the verified multiplication rule. Its order is

$$|G_3| = 46656 = 2^6 3^6. \tag{E.49.8}$$

The images of all 97 literal generators of N_B generate an elementary abelian group of order 81. The program checks every one of their sixth powers and all 97^2 commutations. Thus the whole subgroup $N_B' N_B^6$ is killed, giving the specified epimorphism

$$\bar{\rho}_3 : G_6 = B/(N_B' N_B^6) \twoheadrightarrow G_3, \qquad \bar{\rho}_3(b(N_B' N_B^6)) = \rho_3(b). \tag{E.49.9}$$

This proves equality of all eight evaluations for words with the same old G_6 value. It also proves the exact comparison for every finer common source mapping to G_6. Independently, the 81 kernel elements and the 576 literal B/N_B representatives give 46656 distinct permutations. All 1152 representative generator edges close in these cosets. This verifies the complete source order directly from the displayed kernel and quotient maps. As a separate literal check, all 24 old corrections and all six ternary and seven binary fibre-basis words have identity image under ρ_3.

We evaluate the 24 actual commutator words printed in the E.17 certificate on these eight pairs. Write their values as $v_1, \ldots, v_8$. For each of the 24 words the literal pentagon reads

$$v_1 v_2 = v_3 v_4 v_5. \tag{E.49.10}$$

The Nakamura–Schneps orientation of E.46.14 gives the four changed generators

$$a_1' = a_1^\lambda, \qquad a_i' = v_{i+4}^{-1} a_i^\lambda v_{i+4} \quad (2 \le i \le 4). \tag{E.49.11}$$

The indices here are one-based, so v_6, v_7, v_8 are exactly the pairs in (E.49.7). Since a_i has order three and $\lambda \equiv 1 \pmod 3$, these powers are a_i throughout the stated multiplier range. Every adjacent braid identity and distant commutation holds for this changed quadruple, and its exact generated group has order 51840. If $d' = a_1' a_2' a_3' a_4'$ and $e' = a_1' d'$, then $(d')^5 = (e')^4 = -I$ in every case. These are checks in the full target on five strands.

There is also a direct simultaneous conjugacy proof of these assertions. In the stored column convention put

$$P = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \qquad P^t J P = J, \qquad P^3 = I. \tag{E.49.12}$$

Let k_P be its right-row permutation, and use the zero-based row indices of the 24 original certificates. The simultaneous conjugator k is

certificate rows	k
0–3, 16–19	1
4–7, 20–23	k_P
8–15	k_P^2

$$k^{-1} a_i k = a_i' \quad (1 \le i \le 4). \tag{E.49.13}$$

For k_P an actual positive braid word has index sequence $(2,3,3,2,1,1,2,3,3,2)$; for k_P^2 one is $(1,2,1,1,2,3,2,1,1,2,3)$. The independent matrix calculation checks these words and all 96 simultaneous identities. Conjugation by these actual target elements therefore also proves generation and all braid and twist identities from the original quadruple.

The certificate additionally supplies, as signed words in the four original generators, actual conjugators for d^λ to d' and e^λ to e' at the two multiplier representatives 1 and 73. Each word is replayed as a literal permutation identity. These checks also give the whole multiplier range as follows. The two conjugators for d and $d^{73} = d^3$ to the same d' give a conjugator from d to d^3 by their explicit product. Its four iterates give the four unit exponents $1, 3, 9, 7$ modulo ten. Every profinite unit has one of these residues. Furthermore $\lambda \equiv 1 \pmod{72}$ gives $e^\lambda = e$, since e has order eight. The required periodic conjugacies therefore hold for the whole stated multiplier range, with conjugators inside $\mathrm{Sp}_4(\mathbb{F}_3)$ itself.

Combining (E.49.9) with the unchanged old fibre words proves all 864 finite assertions and their retention in every later stated family. The map on the old common source is literally its projection to G_6 followed by $\bar{\rho}_3$; its joint image with G_3 is the graph of that map. The four-strand graph assertion follows from (E.49.5). The full five-strand generation is the separate verified order calculation above. These arguments use the two maps from the full braid source and their displayed quotient maps; they require no invariance assertion for the kernel of q_3.

The programs with prefix `e_symplectic_five_` and suffixes `model_root.py`, `source_root.py`, `subgroups_root.py` at prime three retain the literal model, source and complete subgroup calculation. The further `p3_fibres_root.py` retains the full edge comparison, all eight values in every row, all braid and generating checks, and the actual periodic conjugator words. The separate `bc_symplectic_five_p3_audit` reconstructs the target by matrix enumeration, every induced coordinate and all 104 compression intertwiners, the exact source cosets, the entire E.18 comparison, every retained fibre value, and the simultaneous conjugators (E.49.12)–(E.49.13). Its original inputs remain unchanged. The further `e_symplectic_five_allunits_root` replays the printed matrix and row table and supplies all 96 conjugator words for the four unit exponents modulo ten. □

5.13 The exact prime-five five-strand source and its signed common quotient

Statement E.50 (Proved). Every retained boundary fibre survives the prime-five symplectic five-strand test. *Let $B = C_2 * C_3$, and retain its actual subgroups F, N, H and the finite source D_{23} from E.24 and E.47. The four chain transvections on $\mathbf{F}_5^4$, with the alternating matrix of E.49, give the specified quotient*

$$B_5 \twoheadrightarrow T_{5,\mathrm{sp}} = \mathrm{Sp}_4(\mathbf{F}_5), \qquad |T_{5,\mathrm{sp}}| = 9360000. \tag{E.50.1}$$

Retaining all eight literal evaluations and all six B/F cosets gives a source $\rho_{5,\mathrm{sp}} : B \twoheadrightarrow G_{5,\mathrm{sp}}$ of order 9000000*. Write*

$$C_5 = \rho_{5,\mathrm{sp}}(F') = \rho_{5,\mathrm{sp}}(H'), \qquad L_5 = C_5', \qquad Q_5 = O_5(C_5). \tag{E.50.2}$$

The actual coordinate maps proved below give

$$\begin{gathered} L_5 \simeq \mathrm{SL}_2(\mathbf{F}_5), \qquad C_5 = L_5 \times Q_5, \qquad Q_5 \simeq (\mathbf{Z}/5)^3, \\ \rho_{5,\mathrm{sp}}(N) = L_5 \times (\mathbf{Z}/5)^5, \qquad \rho_{5,\mathrm{sp}}(\gamma_j H) = C_5 \quad (j \geq 2). \end{gathered} \tag{E.50.3}$$

Put

$$A_m = B/\overline{F'F^m} \quad (m = 2, 10), \qquad Q_{20} = \mathrm{SL}_2(\mathbf{Z}/20)/\{\pm I\}. \tag{E.50.4}$$

The maps from B to Q_{20} take s, r to the classes of $\left(\begin{smallmatrix} 0 & 1 \\ -1 & 0 \end{smallmatrix}\right)$ and $\left(\begin{smallmatrix} 0 & -1 \\ 1 & -1 \end{smallmatrix}\right)$. The actual common quotient with the E.38 double-cover source G_L is

$$J = A_{10} \times_{A_2} Q_{20}, \qquad |A_{10}| = 600, \quad |A_2| = 24, \quad |Q_{20}| = 2880, \quad |J| = 72000, \\ G_{5,\mathrm{sp}}/Q_5 \xrightarrow{\sim} J, \qquad G_L \twoheadrightarrow J. \tag{E.50.5}$$

Here $A_{10} \to A_2$ is reduction of the actual free exponent sums modulo two; $Q_{20} \to A_2$ is projective reduction modulo four, with its values on s, r specified by the same source map.

Every one of the 432 *multiplier-one points of E.24, every full two-adic fibre of E.41, and every full three-adic class-three fibre of E.47 has a simultaneous profinite commutator word retaining these images and the complete E.43 double-cover comparison, and satisfying all five-strand tests in* (E.50.1)*. The word retains the identity action on every full relative pro-p boundary factor of E.42 for $p \geq 5$. Its new five-strand action is the action of the same actual Galois element used in E.43; in particular it is onto $T_{5,\mathrm{sp}}$ and satisfies the complete pentagon, Artin, full-twist and periodic-conjugacy tests. It also retains E.46's Hurwitz quotient through its specified map from G_L.*

For every finite old boundary source G_{old} of E.47 which retains P_{20}, set $G_+ = G_{\mathrm{old}} \times_S G_L$, where $S = G_L/C_L$ is the actual E.43 quotient. The new common sources, and the images of their corrected source maps, are exactly

$$G_{\mathrm{old}} \times_{A_{10}} G_{5,\mathrm{sp}}, \qquad G_+ \times_J G_{5,\mathrm{sp}}. \tag{E.50.6}$$

*The second assertion uses the same word throughout the entire indicated profinite boundary family. Writing $T^{(4)}_{5,\mathrm{sp}} = \langle U_1, U_2, U_3 \rangle$ for the specified order-*15000 *image, the actual common four-strand target and the corrected image are*

$$(T_{23} \times_{S_L} T_L) \times T^{(4)}_{5,\mathrm{sp}} \times A_{96}.$$

The two retained five-strand targets have actual joint image $\mathrm{Sp}_4(\mathbf{F}_5) \times A_{96}$, and the same corrected word gives an onto map to this product as well.

Proof. We construct the source, identify its common quotients, and retain the same correcting word in the joint targets.

The actual finite source and its residual. Use the four matrices $U_i = I + e_i e_i^{\mathrm{t}} \Omega$ over $\mathbf{F}_5$, where $\Omega_{i,i+1} = 1$, $\Omega_{i+1,i} = -1$, and all other entries are zero. Their action on the 624 nonzero vectors is faithful. The literal matrix products verify the Artin relations, and the generated order is $5^4(5^2-1)(5^4-1) = 9360000$. The first three matrices generate a group of order 15000. The two periodic words have orders ten and eight, and their fifth and fourth powers, respectively, are $-I$.

The eight pairs are exactly E.46.12–13, evaluated in these matrices. For each pair the source map is the six-coset formula E.46.15. Thus it is a homomorphism with actual specified values on s, r. The 29952-point induced action is faithfully compressed to 1830 points by the saved 300 orbit intertwiners. Every deleted orbit has an explicit bijection commuting with both generators. The resulting order is 9000000. Evaluation of all 97 literal N generators, the full primitive 95-column F' correction lattice, and the ten literal H Schreier generators gives

image	F	H	N	$F' = H'$	N'	$[H, H']$
order	1500000	1500000	375000	15000	120	15000.

(E.50.7)

The displayed equalities are checked by containment and generating sets. In particular $[\rho(H), C_5] = C_5$, which proves every lower-central equality in (E.50.3).

There is a small exact matrix description of the 120-element subgroup. In the third pentagon pair (A_{23}, A_{34}), the plane spanned by e_2, e_3 is stable. Extract its matrix from the identity B/F coset and transpose it. This defines

$$\theta : \rho(F) \longrightarrow \mathrm{SL}_2(\mathbf{F}_5), \quad \theta(x) = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}, \quad \theta(y) = \begin{pmatrix} 1 & 3 \\ 0 & 1 \end{pmatrix}. \tag{E.50.8}$$

The transpose converts the specified right-permutation multiplication into ordinary matrix multiplication. The faithful compression is inverted on this coordinate using its retained orbit intertwiners. On $L_5 = N' = C_5'$ the map takes exactly the 120 determinant-one matrices, each once; it is therefore an actual isomorphism. The full derived calculation also gives $L_5' = L_5$.

For every retained generator g of N or C_5, let $\ell(g)$ be the unique element of L_5 with $\theta(\ell(g)) = \theta(g)$, and put $v(g) = g\ell(g)^{-1}$. The certificate checks that these $v(g)$ commute with L_5, commute with one another, and have order dividing five. Their complete images have orders 3125 and 125, respectively, and meet L_5 trivially. This proves the direct products in (E.50.3) with these explicit inclusions and projections. It identifies Q_5 as the actual kernel of $\theta|_{C_5}$ and as $O_5(C_5)$.

Every noncentral element of $\mathrm{SL}_2(\mathbf{F}_5)$ has normal closure the full group. The independent matrix calculation records the 120 normal-closure orders: one is 1, one is 2, and the other 118 are 120. Consequently every perfect quotient of $C_5 = L_5 \times Q_5$ kills Q_5. Indeed its image is already the image of L_5, while the image of the central five-group Q_5 would be a central five-subgroup of a quotient of L_5; the displayed normal-closure calculation excludes it. Also C_5^{ab} has order 125, so C_5 has no nontrivial solvable quotient whose order has only the prime divisors two and three.

The abelian and signed source quotients. The images of x, y have order ten and $|\rho(F)^{\mathrm{ab}}| = 100$. Their actual free exponent sums therefore identify this abelianization with $(\mathbf{Z}/10)^2$. Together with the unchanged six-coset map to B/F, this proves

$$G_{5,\mathrm{sp}}/C_5 \xrightarrow{\sim} A_{10}. \tag{E.50.9}$$

For precision the program constructs A_m directly: for a section b_q of B/F it retains the six vectors

$$\mathrm{ab}(b_q b b_{\bar{qb}}^{-1}) \pmod m. \tag{E.50.10}$$

Multiplication is section-word cancellation. Its kernel is $F'F^m$: the identity coset detects ordinary abelianization, and the remaining cosets are its actual conjugates. Enumeration gives $6m^2$ elements for $m = 2, 10$.

On F, the specified modular map to Q_{20} has a unique signed representative whose diagonal entries are 1 modulo four. Its two coordinates are

$$\left(P\theta(w)P^{-1}, \begin{pmatrix} 1 & 2a(w) \\ -2b(w) & 1 \end{pmatrix} \pmod 4 \right), \qquad P = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \tag{E.50.11}$$

where a, b are the ordinary free exponent sums modulo two. This equality holds on x, y and respects the literal free-group multiplication, so it holds on all of F. Chinese remainders give the indicated matrix modulo twenty. The full F image is $\mathrm{SL}_2(\mathbf{F}_5) \times C_2^2$, of order 480; adjoining s, r gives all 2880 elements of Q_{20}. The kernel of $B \to G_{5,\mathrm{sp}}$ lies in F and kills both coordinates in (E.50.11). Thus Q_{20} is an actual quotient of this source. Its kernel is the rank-five elementary abelian factor in (E.50.3).

The map $A_{10} \to A_2$ is reduction of (E.50.10). Reducing the same signed modular matrices modulo four identifies $Q_{20} \to A_2$: all 24 matrix values are paired with their literal A_2 indices

and every generator edge is checked. The resulting two maps have fibres of orders 25 and 120. The actual s, r pair generates every one of the 72000 matching pairs in their fibre product. Its kernel in $G_{5,\mathrm{sp}}$ is $C_5 \cap \ker\theta = Q_5$. This proves the first map in (E.50.5).

Retaining every binary sign in the G_L comparison. Conjugation on L_5 gives the ordinary modular $\mathrm{PSL}_2(\mathbf{F}_5)$ map, through the same change of basis P. The actual inner lifts of s, r in the coordinate (E.50.8) can be taken as

$$\begin{pmatrix} 0 & 1 \\ 4 & 0 \end{pmatrix}, \qquad \begin{pmatrix} 1 & 4 \\ 1 & 0 \end{pmatrix}; \qquad s_{\mathrm{lift}}^2 = r_{\mathrm{lift}}^3 = -I. \tag{E.50.12}$$

Adjoining this actual 60-element projective quotient to the 1980-point Ferrari A_5 source leaves the exact joint order unchanged. It therefore factors through G_A, with the specified source generators.

The signed comparison requires three binary coordinates. Over the actual $\mathrm{SL}_2(\mathbf{F}_2) \times \mathrm{PSL}_2(\mathbf{F}_5)$ base, an element of the Q_{20} kernel has a unique representative equal to I modulo five. Modulo four it is

$$I + 2\begin{pmatrix} a & b \\ c & a \end{pmatrix}, \qquad (a, b, c) \in \mathbf{F}_2^3. \tag{E.50.13}$$

Multiplication adds these three coordinates; conjugation by both source generators preserves this actual elementary abelian kernel. The complete lifted presentation of E.38 is replayed with all three coordinates appended to its 2052 ambient binary coordinates. All 25965 Schreier relations, all normal closure steps, and the literal word DAG are checked. The resulting kernel still has rank 44, and its projection onto the old 44-dimensional kernel is an isomorphism. The saved 3×44 matrix is the exact map to (E.50.13). It proves, retaining the scalar $9I$ as well as both off-diagonal bits, the homomorphism

$$G_L \twoheadrightarrow Q_{20}. \tag{E.50.14}$$

E.43.5 identifies the ordinary exponent sums modulo thirty on the actual F image of G_L. Their reduction modulo ten, with the six-coset quotient, gives $G_L \to A_{10}$. This map and (E.50.14) agree in A_2, since both are the specified maps from B. Their pair is onto J by the complete enumeration of its generating s, r values. This proves the second map in (E.50.5).

Corrections that retain the entire boundary system. Let

$$R_{\mathrm{nil}} = \bigcap_{j \geq 1} \gamma_j(\widehat{H}). \tag{E.50.15}$$

For any continuous finite image of $\widehat{H}$, the image of this intersection is its stabilized lower-central subgroup. Indeed, for each element in that finite intersection, its inverse images in the compact groups $\gamma_j(\widehat{H})$ are nonempty nested closed sets. Their intersection is a preimage in R_{nil}. It follows from (E.50.3) that $\rho_{5,\mathrm{sp}}(R_{\mathrm{nil}}) = C_5$. Every finite nilpotent H boundary image kills R_{nil}. This includes every finite stage of E.41, E.47, and all full pro-p H factors of E.42.

On G_L, the image of R_{nil} is exactly C_L. Its image in S is trivial because the actual H image of S is the class-two group $Q_8 \times C_{10}^2$ in E.43. Conversely E.43's closed prosolvable residual is contained in R_{nil} and maps onto C_L. Thus both inclusions are actual subgroup comparisons.

Apply Goursat to the image of R_{nil} in $C_L \times C_5$. Its common quotient is perfect because C_L is perfect, and every perfect quotient of C_5 kills Q_5 by the normal-closure argument following (E.50.8). Hence

$$\rho_{5,\mathrm{sp}}(R_{\mathrm{nil}} \cap \ker\rho_L) \supset Q_5. \tag{E.50.16}$$

The reverse inclusion follows from the common map to J in (E.50.5). Intersecting also with $\ker \rho_{D_{23}}$ retains this entire image: the two images in question are a five-group and a group whose order has only the prime divisors two and three, so their common quotient is trivial. Therefore

$$\rho_{5,\mathrm{sp}}(R_{\mathrm{nil}} \cap \ker \rho_L \cap \ker \rho_{D_{23}}) = Q_5. \tag{E.50.17}$$

Every element of this compact subgroup also kills every retained boundary factor, and lies in $\widehat{H}' \subset \widehat{F}'$.

The old finite sources retain A_{10}: $\gamma_3(H)H^{20} \subset F'F^{10}$ gives the actual map $P_{20} \to A_{10}$. Before adjoining G_L, the same argument with C_5 and the solvable $\{2,3\}$ image of D_{23} gives the whole pure C_5. Together with (E.50.9) this proves the first original-source fibre product in (E.50.6). For that family choose the correction with new image $\rho_{5,\mathrm{sp}}(f_0)^{-1}$; this element lies in C_5 because $f_0 \in \widehat{F}'$. The resulting new word value is 1, so its multiplier-one new source and braid maps are the original onto maps. After adjoining G_L, equation (E.50.17) gives the whole pure Q_5, and the common map to J gives the reverse containment. This proves that J is the maximum common quotient through these specified source maps, and proves the second original-source fibre product.

Now take the E.47 word f_0 with the E.43 comparison $\rho_L(f_0) = \rho_L(f_\sigma)$ for its specified actual Galois element σ. Retain exactly E.43's choice $\chi(\sigma) = 1 \in \widehat{\mathbf{Z}}^\times$ and its prescribed Kummer value. In particular $\chi(\sigma) \equiv 1 \pmod{10}$ for the new source generators and order-five transvections, and $\chi(\sigma) \equiv 1 \pmod{40}$ for the orders-ten and eight periodic words. Thus the multiplier-one finite power maps are exactly those of this same Galois element. Their images in J agree, so

$$q = \rho_{5,\mathrm{sp}}(f_0)^{-1}\rho_{5,\mathrm{sp}}(f_\sigma) \in Q_5. \tag{E.50.18}$$

Choose its actual preimage k in the compact subgroup of (E.50.17), and put $f = f_0 k$. This one profinite commutator word retains every old coordinate and satisfies $\rho_{5,\mathrm{sp}}(f) = \rho_{5,\mathrm{sp}}(f_\sigma)$. In particular all 48 induced evaluation coordinates agree with the same Galois word. The new five-strand map is consequently the original quotient in (E.50.1) precomposed with the actual profinite Galois braid automorphism. This proves its relations, periodic conjugacies and surjectivity without a kernel-invariance assertion for a finite shadow. E.46.26 carries the unchanged G_L value to its Hurwitz source as well.

Generation on the actual joint source. The corrected maps to both factors of the second fibre product in (E.50.6) are onto, and agree on J with the same Galois source map. On the old side the H image is onto because its retained B/H map is the same automorphism and its whole source map is onto. On the new side it is onto because the corrected map agrees with the actual Galois map, whose source automorphism preserves H. Equivalently, its onto F image has abelianization C_{10}^2 and hence no nontrivial quotient of the exponent-three group F/H. This verifies the H comparison even though $G_{5,\mathrm{sp}}$ itself does not retain B/H.

By the compact argument after (E.50.15), the corrected image of R_{nil} on the new side is C_5. On the old side it is the direct product of C_L with a solvable $\{2,3\}$ image: the boundary factors kill it, and perfectness excludes a common quotient between C_L and that solvable image. Every common quotient of this old residual and C_5 is perfect. Indeed its abelianization is both a quotient of the old $\{2,3\}$ abelianization and a quotient of C_5^{ab}, which is a five-group; it is therefore trivial. The same normal-closure argument forces that common quotient to kill Q_5. Thus the corrected joint image contains $\{1\} \times Q_5$. Its onto projection to G_+ now gives the entire fibre product over J. The version without G_L follows by the corresponding argument with the whole C_5. This proves generation for the actual source maps, rather than presuming that their finite kernels are invariant.

The exact joint targets and their generation. There is an additional complete target comparison. Pair the three literal Ferrari generators of T_L with U_1, U_2, U_3, respectively. The combined faithful base consists of the 190-point A_5 action, its four-point global-character action, and the 624-point symplectic action. Above that actual base, all 73496 Schreier relations and their normal closure are evaluated in the full 57-coordinate binary extension. Its kernel has rank 55, and the resulting exact order is $|T_L| \cdot 15000$. Thus the image of these specified braid maps is

$$T_L \times T_{5,\mathrm{sp}}^{(4)}. \tag{E.50.19}$$

This is an equality of the actual joint image, with both original projection maps.

For an independent perfectness certificate, the two commutators $[U_1, U_2]$, $[U_2, U_3]$ generate a subgroup of order 1250. Adjoining the literal conjugate $U_1[U_1, U_2]U_1^{-1}$ generates all 15000 matrix values. The complete matrix enumeration checks normal closure under all three generators. Hence $T_{5,\mathrm{sp}}^{(4)}$ is perfect. Every solvable quotient of the direct product in (E.50.19) therefore factors through the E.38 quotient S_L of T_L. The old T_{23} map retains this exact quotient. Goursat, with these literal comparison maps, gives the common four-strand target

$$(T_{23} \times_{S_L} T_L) \times T_{5,\mathrm{sp}}^{(4)}. \tag{E.50.20}$$

For the corrected maps, the pair in (E.50.19) is the original joint map precomposed with the same Galois automorphism of $\widehat{B}_4$, because both full source values agree with f_σ. It is therefore onto that product. The corrected T_{23} projection is onto by the old theorem and has the same S_L value. The same maximum-solvable-quotient argument proves generation of (E.50.20).

All prime divisors of its order are among $2, 3, 5, 7$. E.46's eleven-cycle normal-closure argument therefore gives the additional direct factor A_{96}, both for the original and the corrected four-strand maps. For the two five-strand maps, the order 9360000 has no prime divisor eleven, so the same normal-closure argument excludes a nontrivial common quotient with A_{96}. Their original joint image is consequently

$$\mathrm{Sp}_4(\mathbf{F}_5) \times A_{96}. \tag{E.50.21}$$

Both corrected maps agree with the very same Galois braid map, so their joint map is onto (E.50.21) as well.

Certificates and scope. The independent programs share the prefix `bc_symplectic_five_p5_`, with suffixes `residual`, `modular`, `signed`, and `common`. They retain the exact source arrays, matrix projections, all common-quotient maps and the complete lifted relation certificate. The failed one-bit signed attempt is preserved with its correction: the projective level-four quotient does not factor through the old A_5 base; all three bits in (E.50.13) are required in the comparison. The two additional suffixes `target` and `target_perfect` retain (E.50.19) and the literal three-generator commutator certificate for its new factor. The result concerns the specified multiplier-one family and the stated complete boundary factors. The cofinal full-GT image questions remain those recorded in E.2 and E.3. □

Statement E.51 (Proved). The prime-seven five-strand refinement preserves the full preceding family. *Let $\rho_7 : \widehat{B} \twoheadrightarrow G_7$ be the actual source obtained by inducing all eight evaluations of the four chain transvections in* $\mathrm{Sp}_4(\mathbb{F}_7)$ *through all six cosets of B/F, with the*

same words as E.50. Its exact finite images are

$$\begin{gathered}|G_7| = 135\,531\,648, \qquad |\rho_7(F)| = |\rho_7(H)| = 22\,588\,608,\\ C_7 := \rho_7(F') = \rho_7(H') = \rho_7(\gamma_r H) \quad (r \geq 2), \qquad |C_7| = 115\,248,\\ L_7 := C_7' = \rho_7(N') \simeq \mathrm{SL}_2(\mathbb{F}_7),\\ C_7 = L_7 \times Q_7, \quad Q_7 \simeq (\mathbb{Z}/7)^3,\\ \rho_7(N) = L_7 \times (\mathbb{Z}/7)^5.\end{gathered} \tag{E.51.1}$$

The isomorphisms in this display are the specified matrix and generator maps in the proof.

Write

$$\begin{gathered}A_{14} = B/(F'F^{14}), \qquad A_2 = B/(F'F^2),\\ Q_{28} = \mathrm{SL}_2(\mathbb{Z}/28)/\{\pm I\}.\end{gathered} \tag{E.51.2}$$

The signed modular map and the six-coset abelian map give the actual quotient

$$\begin{gathered}G_7/Q_7 \simeq A_{14} \times_{A_2} Q_{28},\\ |A_{14}| = 1176, \quad |Q_{28}| = 8064, \quad |G_7/Q_7| = 395\,136.\end{gathered} \tag{E.51.3}$$

The maximum common quotient of G_7 and the E.38 source G_L through their specified maps from B is A_2.

Let G_{old} be any cofinal finite common source of the entire E.50 family that retains $P_{28} = B/(\gamma_3(H)H^{28})$. It includes the fixed D_{23}, G_L, and prime-five symplectic source maps, together with any prescribed finite levels of the boundary factors. Then the maximum common quotient of G_{old} and G_7 is the displayed A_{14}, and the actual joint source is

$$G_{\mathrm{old}} \times_{A_{14}} G_7. \tag{E.51.4}$$

All 432 *retained multiplier-one source points, with every prescribed full two-adic class-two fibre of E.41 and full three-adic class-three correction-image fibre of E.47, have one simultaneous profinite commutator word satisfying this new test. It retains all the full pro-p H factors from E.42, the complete E.50 prime-five test, and the E.46 Hurwitz test. In both new symplectic sources and in G_L, this word has exactly the values of the same actual Galois element σ with $\chi(\sigma) = 1$. Its changed source map is onto* (E.51.4) *at every finite level.*

Put $T_7 = \mathrm{Sp}_4(\mathbb{F}_7)$ and let J_7 be its actual first four-strand image. Then

$$|T_7| = 276\,595\,200, \qquad |J_7| = 115\,248, \qquad J_7' = J_7. \tag{E.51.5}$$

*With J_5 the corresponding order-*15000 *group in E.50, the complete joint four-strand target for this family is*

$$(T_{23} \times_{S_L} T_L) \times J_5 \times J_7 \times A_{96}. \tag{E.51.6}$$

The same word generates it and satisfies all its braid and full-twist identities. The corresponding full five-strand target is

$$\mathrm{Sp}_4(\mathbb{F}_5) \times \mathrm{Sp}_4(\mathbb{F}_7) \times A_{96}, \tag{E.51.7}$$

again with the actual simultaneous generator maps. These assertions retain the finite sources and the indicated boundary inverse limits; they impose no additional identity between the full profinite GT group and its Galois subgroup.

Proof. All comparisons below retain the displayed maps from the fixed source B.

The literal representation and residual maps. Use the alternating matrix Ω whose only nonzero upper-diagonal entries are $\Omega_{i,i+1} = 1$, and put $U_i = I + e_i(e_i^t\Omega)$ over $\mathbb{F}_7$. The program with prefix `bc_symplectic_five_p7_model` checks the symplectic identities, all Artin relations and the eight specified pure-braid evaluation pairs. The action on the 2400 nonzero vectors is faithful and has order $7^4(7^2-1)(7^4-1) = 276595200$. It is therefore the whole symplectic group. The first three generators have image order 115248. The periodic words δ_5 and ϵ_5 have orders ten and eight, respectively, and $\delta_5^5 = \epsilon_5^4 = -I$.

The induced source starts on $6 \cdot 8 \cdot 2400 = 115200$ points. The source certificate retains explicit intertwiners from all 592 original orbits to ten distinct B sets, of total degree 3570. Thus its compression is an isomorphism of the actual permutation image. The subgroup calculation evaluates all 97 literal N generators and all 95 primitive columns imposing zero integral F abelianization. It gives

actual image	order
G_7	135531648
$\rho_7(F) = \rho_7(H)$	22588608
$\rho_7(N) = \rho_7(K_6)$	5647152
$\rho_7(F') = \rho_7(K_6 \cap F')$	115248
$\rho_7(N') = C_7'$	336.

(E.51.8)

It also checks directly that $[\rho_7(H), C_7] = C_7$. Consequently every lower-central term from the second onward has this same image.

Here is the exact matrix comparison underlying the direct products. Decode the identity-coset third pentagon coordinate through the saved intertwiners. It preserves $\langle e_2, e_3\rangle$; transpose its block to account for the right-permutation product. This gives a homomorphism $\theta : \rho_7(F) \to \mathrm{SL}_2(\mathbb{F}_7)$ with

$$\theta(x) = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}, \qquad \theta(y) = \begin{pmatrix} 1 & 5 \\ 0 & 1 \end{pmatrix}. \tag{E.51.9}$$

Its restriction to L_7 takes all 336 values exactly once. For every original generator g of $\rho_7(N)$ or C_7, let ℓ_g be the unique element of L_7 with $\theta(\ell_g) = \theta(g)$, and put

$$v_g = g\ell_g^{-1}. \tag{E.51.10}$$

The literal permutations verify that all these v_g commute with one another and with L_7, and have seventh power one. They generate groups of orders 7^5 and 7^3, respectively. Since θ is injective on L_7 and kills these groups, their intersections with L_7 are trivial. The displayed generator decompositions prove the actual direct products in (E.51.1).

Define the ordinary modular map q_7 using

$$s \longmapsto \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \qquad r \longmapsto \begin{bmatrix} 0 & -1 \\ 1 & -1 \end{bmatrix} \quad \text{in } \mathrm{PSL}_2(\mathbb{F}_7). \tag{E.51.11}$$

For $P = \left(\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right)$, all 672 conjugations of the 336 elements of L_7 by s, r satisfy

$$\theta(a^{-1}\ell a) = (Pq_7(a)P)^{-1}\theta(\ell)(Pq_7(a)P). \tag{E.51.12}$$

The formula is independent of the sign of a representative of $q_7(a)$. In particular the induced action on the inner automorphism group is exactly the specified projective modular map, with this coordinate swap.

The certificate enumerates every normal closure in $\mathrm{SL}_2(\mathbb{F}_7)$: the identity gives order one, $-I$ gives order two, and each of the other 334 elements gives the whole order 336 group. The

literal derived subgroup is the whole group. Thus its nontrivial perfect quotients are precisely itself and $\mathrm{PSL}_2(\mathbb{F}_7)$. It follows as well that

$$\text{every nontrivial quotient of } C_7 \text{ has order divisible by 7.} \tag{E.51.13}$$

Indeed an image of Q_7 has seventh-power order, and every nontrivial image of L_7 has order divisible by seven. These images generate any quotient of their direct product.

Both signed modular quotients and their actual old-source comparison. The six-coset Schreier exponent sums modulo fourteen give A_{14}, with kernel $F'F^{14}$ and order 1176. The faithful joint image with G_7 still has order 135531648, proving the quotient map and its kernel C_7. Reducing the same coordinates modulo two gives the exact A_2 map.

Enumerate Q_{28} using the matrices in (E.51.11) modulo twenty-eight, retaining the global sign. It has 8064 elements. Its reduction modulo four is identified with A_2 on all 24 values, using the same dictionary as E.38 and E.50. An element in this kernel has a unique sign representative equal to I modulo four; reduction of that representative modulo seven identifies the kernel with all of $\mathrm{SL}_2(\mathbb{F}_7)$.

The six-coset representation induced from the two signed $\mathrm{SL}_2(\mathbb{F}_7)$ matrices $\left(\begin{smallmatrix}1&2\\0&1\end{smallmatrix}\right)$, $\left(\begin{smallmatrix}1&0\\-2&1\end{smallmatrix}\right)$ is compared with every element and all 16128 generator edges of Q_{28}. Thus it is precisely this signed quotient. All 395136 matching pairs in $A_{14} \times_{A_2} Q_{28}$ are generated by the literal s, r. Their joint map with G_7 retains its original order; the remaining kernel has order 343 and is Q_7 from (E.51.10). This proves (E.51.3) with its specified maps.

The old-source modular comparison is different from the prime-five computation and is checked directly. Pair the faithful G_A source with the action of (E.51.11) on $\mathbb{P}^1(\mathbb{F}_7)$. The exact joint order is $168|G_A|$, so

$$\mathrm{im}(\rho_A, q_7) = G_A \times \mathrm{PSL}_2(\mathbb{F}_7). \tag{E.51.14}$$

The E.38 map $G_L \to G_A$ has elementary two-group kernel. Therefore $q_7(\ker \rho_A)/q_7(\ker \rho_L)$ is a two-group. Its numerator is the perfect $\mathrm{PSL}_2(\mathbb{F}_7)$ by (E.51.14), so the quotient is trivial. Thus $q_7(\ker \rho_L)$ is the whole projective group.

Because the actual $Q_4 = A_2$ map factors through G_L, the image J of $\ker \rho_L$ in Q_{28} is contained in its signed kernel $L = \mathrm{SL}_2(\mathbb{F}_7)$ and maps onto $L/Z(L)$. Hence $JZ(L) = L$, and

$$L = [L, L] = [JZ(L), JZ(L)] = [J, J] \subset J. \tag{E.51.15}$$

This proves the actual joint image $G_L \times_{A_2} Q_{28}$, including its central sign.

The complete correction preserving E.50 and every boundary factor. Put $R_{\mathrm{nil}} = \bigcap_{r \geq 1} \gamma_r(\widehat{H})$, and use C_L for the perfect E.43 residual in G_L. For any finite image of $\widehat{H}$, its lower-central residual is the image of R_{nil}: surjectivity at each lower-central term and nested compact fibres prove the reverse inclusion as well as the immediate forward one. E.43 and (E.51.1) consequently give

$$\rho_L(R_{\mathrm{nil}}) = C_L, \qquad \rho_7(R_{\mathrm{nil}}) = C_7. \tag{E.51.16}$$

For the first equality, the quotient of the H image in G_L by C_L is the nilpotent group of E.43; the closed derived residual already maps onto C_L.

The joint image in (E.51.16) is the full product. To prove this, any nontrivial Goursat quotient is perfect, since C_L is perfect. A perfect quotient of $C_7 = L_7 \times Q_7$ is L_7 or its projective quotient. Indeed the image of L_7 must be the whole perfect quotient; its centre has order at most two, so the central seventh-power group Q_7 maps to one. Both Goursat kernels

are stable under the full B action, because R_{nil} is characteristic in the normal subgroup $\widehat{H}$. Formula (E.51.12) would therefore make its projective inner action factor through G_L. This would factor the specified q_7 through G_L, contrary to (E.51.14)–(E.51.15). Hence

$$\mathrm{im}_{G_L \times G_7}(R_{\mathrm{nil}}) = C_L \times C_7. \tag{E.51.17}$$

In particular $K = R_{\mathrm{nil}} \cap \ker \rho_L$ maps onto C_7. The image of K in the actual $D_{23} \times G_{5,\mathrm{sp}}$ has order prime to seven. By (E.51.13), its joint image with C_7 is the full product. Intersecting with this finite kernel therefore retains the surjection

$$R_{\mathrm{nil}} \cap \ker \rho_L \cap \ker(\rho_{D_{23}}, \rho_{5,\mathrm{sp}}) \twoheadrightarrow C_7. \tag{E.51.18}$$

This is the required explicit comparison with the E.50 kernel. Every finite nilpotent H image kills R_{nil}, so the same closed subgroup kills all finite boundary factors and their full indicated inverse limits. No separate correction for a boundary level is needed.

The pure C_7 in (E.51.17) forces every common quotient of G_L, G_7 through A_{14}. This is solvable, whereas the maximal solvable quotient of G_L has order 43200 and no factor seven. Its common quotient kills the $(\mathbb{Z}/7)^2$ kernel of $A_{14} \to A_2$. The specified A_2 maps attain the resulting quotient, proving its maximality.

For G_{old}, the actual map $P_{28} \to A_{14}$ follows from $\gamma_3 H \subset F'$ and $H^{28} \subset F^{14}$. The P_4 and full pro-seven H factors already supply P_{28} by their actual coprime geometric-image product. More explicitly, the class-two exponent-28 quotient of H is the direct product of its exponent-four and exponent-seven Sylow quotients; their maps are the original source reductions, and the inverse uses the two coprime power idempotents. Thus requiring P_{28} only passes to a cofinal subsystem. The pure C_7 in (E.51.18) forces every common quotient through A_{14}, and A_{14} is a specified common quotient. Goursat gives exactly (E.51.4).

One Galois comparison and full joint generation. Choose the E.50 word f_0 for the prescribed old point and boundary parameters. Retain its actual Galois element σ with $\chi(\sigma) = 1 \in \widehat{\mathbb{Z}}^\times$. In particular $\chi(\sigma) \equiv 1 \pmod{280}$, which retains the orders fourteen of the new source generators, seven of the transvections, and ten and eight of the periodic words. Since both words belong to $\widehat{F}'$, their new images differ by

$$c = \rho_7(f_0)^{-1} \rho_7(f_\sigma) \in C_7. \tag{E.51.19}$$

Lift c through (E.51.18) and put $f = f_0 k$. This is a single word in $\widehat{F}'$. It retains every old value and satisfies $\rho_7(f) = \rho_7(f_\sigma)$. All 48 induced evaluation values agree with this same Galois word. The new relations, periodic conjugacies and individual generation are consequently the actual Galois braid maps followed by the specified finite quotient maps. This argument does not require an arbitrary finite shadow to preserve its source kernel.

For joint generation, the corrected source maps to G_L and G_7 are both precomposed with the same actual Galois automorphism of $\widehat{B}$. It preserves $\widehat{F}$, its characteristic subgroup $\widehat{H}$, and R_{nil}. Thus their joint residual image remains exactly (E.51.17). Intersecting with the corrected D_{23} and prime-five symplectic kernels uses the same prime-to-seven argument in (E.51.18). The corrected boundary maps still kill R_{nil}, since their H images are nilpotent at each finite level. The corrected joint image therefore contains the full pure C_7. Its projection onto G_{old} is onto by E.50, and its maps to A_{14} agree because they are the same literal source formulas. The quotient of (E.51.4) by pure C_7 is G_{old}, proving surjectivity onto the full fibre product. The actual H restrictions are onto as well: the whole maps are onto and their retained B/H map is the same automorphism. Nested compact fibres give the asserted one-word comparison through every boundary level.

The image parameters do not specify a unique profinite word. If $\mathcal{K}$ denotes the closed correcting subgroup on the left of (E.51.18), the corrections within this subgroup for the fixed value c are exactly $k(\mathcal{K} \cap \ker \rho_7)$. If ρ_{old} denotes the full joint old map on $\widehat{F}'$, the set of all words with the prescribed old value and the prescribed new Galois value is exactly

$$f \, \ker\big((\rho_{\mathrm{old}}, \rho_7)|_{\widehat{F}'}\big). \tag{E.51.20}$$

Both assertions follow by multiplying two preimages under the displayed continuous homomorphisms.

The joint braid targets. The literal commutators $[U_1, U_2]$, $[U_2, U_3]$, and $U_1[U_1, U_2]U_1^{-1}$ generate all 115248 matrices of J_7; the intermediate orders are 2744 and 115248. This proves $J_7' = J_7$ without a structural identification of its unipotent radical.

The complete lifted Schreier calculation pairs the three actual Ferrari generators in T_L with U_1, U_2, U_3. It retains all 57 ambient binary coordinates and recovers the full 55-dimensional central kernel above the actual permutation product. Its exact order is $|T_L|\,115248$, proving the actual target $T_L \times J_7$. A separate literal paired permutation calculation gives $J_5 \times J_7$ and, on all four generators, $\mathrm{Sp}_4(\mathbb{F}_5) \times T_7$.

These comparisons also give the joint $T_L \times J_5 \times J_7$. In a nontrivial Goursat quotient of $T_L \times J_5$ and J_7, quotient by either of the two commuting factor images. A nontrivial result contradicts one of the two checked pair products. If both results are trivial, those commuting images both equal the quotient, so it is abelian. It is also perfect, as a quotient of J_7, and hence is trivial after all.

The kernel of $(T_{23} \times_{S_L} T_L) \times J_5 \to T_L \times J_5$ is solvable. A nontrivial perfect Goursat quotient with J_7 cannot be the image of this solvable kernel; quotienting by that image would give a nontrivial common quotient with $T_L \times J_5$, already excluded. This proves the complete product before the A_{96} factor in (E.51.6). The simple group A_{96} has order divisible by eleven, while the other factors do not. The actual Hurwitz projection and Goursat therefore give its full product as well.

All the T_L, J_5, J_7, A_{96} components of the corrected word are the same Galois precomposition, so their simultaneous images are the actual products just proved. The solvable-kernel argument retains the old T_{23} component and proves corrected generation in (E.51.6). The paired five-strand calculation and the same A_{96} argument prove (E.51.7) with its corrected maps.

The new programs have the shared prefix `bc_symplectic_five_p7_`. Their suffixes `model`, `source`, `subgroups`, `residual`, `modular_probe`, and `common` retain all source maps and signed comparisons; `target`, `target_perfect`, and `joint_symplectic` retain the full target comparisons. Each proof assertion above uses these actual finite maps and their stated compact inverse-limit maps. □

5.14 The integral source through every modulus prime to six

Statement E.52 (Proved). The complete congruence source and its lower-central residual. *Let $B = C_2 * C_3 = \langle s, r \mid s^2 = r^3 = 1\rangle$, let $F = \ker(B \to \mathfrak{S}_3)$, and use the actual generators $x = (sr)^{-2}$ and $y = sxs$. Let $N = \ker(B \to \mathrm{SL}_2(\mathbf{Z}/12)/\{\pm I\})$ with the matrix map of E.50.4, and put $H = F'F^3$. We write hats for their profinite completions, identified with the corresponding open subgroups of $\widehat{B}$. All commutator and power subgroups in a profinite group are closed. Set*

$$A = \mathbf{Z}[1/6], \qquad R = \prod_{p \geq 5} \mathbf{Z}_p, \qquad K_2 = \overline{\widehat{F}'\widehat{F}^2}, \qquad R_{\mathrm{nil}} = \bigcap_{j \geq 2} \gamma_j \widehat{H}. \tag{E.52.1}$$

For every integer $m > 1$ prime to six, the integral eight-pair construction below defines a specified continuous epimorphism

$$\rho_m : \widehat{B} \twoheadrightarrow G_m \subset \mathrm{Sp}_4(\mathbf{Z}/m)^{48} \rtimes \mathfrak{S}_3. \tag{E.52.2}$$

The alternating form and the right matrix convention are specified below. Reduction for $m \mid n$ gives the transition map $G_n \twoheadrightarrow G_m$, with ρ_m equal to reduction of ρ_n. Let G_R be their inverse limit and let $\rho_R : \widehat{B} \twoheadrightarrow G_R$ be the resulting continuous map.

There are explicit injective homomorphisms

$$\begin{aligned} \Phi_m : \mathrm{SL}_2(\mathbf{Z}/m) \times (\mathbf{Z}/m)^5 &\hookrightarrow G_m, \\ \Phi_R : \mathrm{SL}_2(R) \times R^5 &\hookrightarrow G_R, \end{aligned} \tag{E.52.3}$$

compatible with every reduction. For $T = \mathbf{Z}/m$ or $T = R$, write ρ_T, Φ_T, G_T for these maps. The rank-three direct summand $V_3(T) \subset T^5$ is the row span of the following displayed matrix:

$$D = \begin{pmatrix} -4/3 & 4/3 & -2/3 & 5/3 & 1 \\ 1/3 & -1/3 & -1/3 & -2/3 & 0 \\ 2/3 & -2/3 & 1/3 & -4/3 & -1 \end{pmatrix}. \tag{E.52.4}$$

Then the exact images, with the inclusions of (E.52.3)*, are*

$$\begin{aligned} \rho_T(\widehat{N}) = \rho_T(K_2) &= \Phi_T\big(\mathrm{SL}_2(T) \times T^5\big), \\ \rho_T(\widehat{H}) &= \rho_T(\widehat{F}), \\ C_T := \rho_T(\widehat{F}') = \rho_T(\widehat{H}') &= \Phi_T\big(\mathrm{SL}_2(T) \times V_3(T)\big), \\ \rho_T(\gamma_j \widehat{H}) = \rho_T(R_{\mathrm{nil}}) &= C_T \qquad (j \geq 2), \\ \rho_T(\widehat{N}') = C'_T &= \Phi_T\big(\mathrm{SL}_2(T) \times 0\big). \end{aligned} \tag{E.52.5}$$

In particular every element of C_R has a preimage in the single closed subgroup R_{nil}. For one such preimage w, its entire fibre there is $w(R_{\mathrm{nil}} \cap \ker \rho_R)$.

The actual finite quotient maps from $\widehat{B}$ give

$$\begin{aligned} G_m/\rho_m(K_2) &\simeq A_2, \qquad |A_2| = 24, \\ G_m/C_m &\simeq A_{2m}, \qquad A_n = \widehat{B}/\overline{\widehat{F}'\widehat{F}^n}, \\ |G_m| = 24m^5 |\,\mathrm{SL}_2(\mathbf{Z}/m)|, &\qquad \rho_m(\widehat{F})^{\mathrm{ab}} \simeq (\mathbf{Z}/2m)^2. \end{aligned} \tag{E.52.6}$$

The last isomorphism sends x, y to the two ordinary exponent-sum basis vectors. The corresponding quotients over R are the inverse limits of these same maps. Reduction at $m = 5, 7$ recovers the specified sources of E.50 and E.51 through their retained orbit intertwiners. Thus (E.52.5) *includes all their prime-power and simultaneous congruence refinements.*

Proof. We first construct the integral maps and then prove surjectivity and every asserted quotient using those maps.

Integral matrices and the actual induced source. Let J be the four-by-four alternating matrix with $J_{i,i+1} = 1$, $J_{i+1,i} = -1$ and all other entries zero, and put $\Omega = J^{-1}$. The matrices used with ordinary multiplication are

$$M_i = (I + e_i e_i^{\mathsf{t}} J)^{\mathsf{t}} \quad (1 \leq i \leq 4), \qquad M_i^{\mathsf{t}} \Omega M_i = \Omega. \tag{E.52.7}$$

This is the transpose convention which realizes the right-permutation convention of E.50 and E.51. Their entries are integers. Direct multiplication gives the adjacent Artin relations, commutation at distance at least two, and

$$(M_1M_2M_3M_4)^5 = (M_1^2M_2M_3M_4)^4 = -I. \tag{E.52.8}$$

Define A_{ij} from these M_i by the literal pure braid word

$$A_{ij} = M_{j-1}\cdots M_{i+1}M_i^2M_{i+1}^{-1}\cdots M_{j-1}^{-1}. \tag{E.52.9}$$

The first five evaluation pairs are, in order,

$$\begin{gathered}(A_{12}, A_{23}A_{24}), \quad (A_{13}A_{23}, A_{34}), \quad (A_{23}, A_{34}),\\ (A_{12}A_{13}, A_{24}A_{34}), \quad (A_{12}, A_{23}).\end{gathered} \tag{E.52.10}$$

The final three are $(M_{i-1}\cdots M_1M_1\cdots M_{i-1}, M_i^2)$, for $i = 2, 3, 4$. Each pair defines a homomorphism from the discrete free group $F = \langle x, y\rangle$ to $\mathrm{Sp}_4(A, \Omega)$.

Order the six matrices of $\mathrm{SL}_2(\mathbf{F}_2)$ lexicographically. For each, take the first positive s, r word given by breadth-first search starting at the identity, trying s before r. These are the representatives t_q in the certificate. For $b \in B$, the component at (q, j) of the induced map is the j-th pair evaluated at

$$t_q b t_{q\bar{b}}^{-1} \in F. \tag{E.52.11}$$

The top component is right multiplication by $\bar{b}$ on the six cosets. The equality $t_q bb' t_{q\bar{b}\bar{b}'}^{-1} = (t_q b t_{q\bar{b}}^{-1})(t_{q\bar{b}} b' t_{q\bar{b}\bar{b}'}^{-1})$ proves the homomorphism identity, with this exact action on the 48 coordinates. Reducing the matrices modulo m and extending from the finitely generated discrete source to its profinite completion gives (E.52.2). Here G_m denotes its actual image. The image in the inverse limit is closed and surjects at each finite stage, so it is the whole inverse limit.

On the identity-coset third pair, rows and columns two and three give a literal two-by-two quotient: the corresponding rows have zero entries in columns one and four. Therefore extraction defines a homomorphism on the image of F, with

$$\theta(x) = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}, \qquad \theta(y) = \begin{pmatrix} 1 & -2 \\ 0 & 1 \end{pmatrix}. \tag{E.52.12}$$

The same extraction defines its reductions and the continuous map over R.

The finite exact factorization over A. We describe precisely the rational matrix certificate used here. The rank-97 basis $n_0, \ldots, n_{96}$ of N is obtained from the spanning tree of the finite $(2, 3)$ coset graph of $\mathrm{SL}_2(\mathbf{Z}/12)/\{\pm I\}$. The program `de_ferrari_a6_source_data.py` retains all edges and the actual free words and compares these words with the earlier saved source. It checks both inverse substitutions between the free graph basis and x, y before any evaluation. For each of the 48 coordinates, solve over $\mathbf{Q}$ the finite intertwiner equations

$$M_j(n_i)T_j = T_j\theta(n_i) \quad (0 \le i \le 96). \tag{E.52.13}$$

There are 24 coordinates with one-dimensional intertwiner space and 24 with zero space. In each nonzero case choose the primitive integral matrix given by the saved nullspace basis and set

$$P_j = (T_j^{\mathsf{t}}\Omega T_j)^{-1}T_j^{\mathsf{t}}\Omega, \quad P_jT_j = I_2, \quad L_j(B) = I_4 + T_j(B - I_2)P_j. \tag{E.52.14}$$

For a zero space put $L_j(B) = I_4$. All entries of P_j lie in A; their denominators divide six. Multiplication and $P_jT_j = I_2$ prove $L_j(B)L_j(C) = L_j(BC)$. The formula is the specified action on the symplectic plane $\operatorname{im} T_j$ and the identity on its orthogonal complement. Thus it preserves Ω for every $B \in \mathrm{SL}_2$ over every A-algebra.

Flatten the 48 matrices $L_j(\theta(n_i))^{-1}M_j(n_i) - I_4$ into a row with 768 entries. The first five rows are a basis $\mathcal{D}_0, \dots, \mathcal{D}_4$; write $D_{u,j}$ for the four-by-four piece of $\mathcal{D}_u$ at coordinate j. The exact checks give

$$\begin{gathered} D_{u,j}D_{v,j} = 0, \qquad D_{u,j}^{\mathrm{t}}\Omega + \Omega D_{u,j} = 0, \\ D_{u,j}L_j(B) = L_j(B)D_{u,j}, \qquad M_j(n_i) = L_j(\theta(n_i))\left(I_4 + \sum_{u=0}^{4} a_{iu}D_{u,j}\right). \end{gathered} \tag{E.52.15}$$

For commutation it suffices to check the two elementary one-parameter matrix coefficients; the independent audit checks the elementary matrices and their linear coefficients. The identities involving the form can also be recovered from the symplectic matrices $I + D_{u,j}$ and $I - D_{u,j}$: their difference gives the linear identity, since two is invertible, and $D_{u,j}D_{v,j} = 0$ then kills every quadratic term. All these are exact rational identities, with no finite-field rank substituted for a rational rank.

The basis minor in flattened columns $4, 6, 12, 20, 48$ (numbered from zero) has determinant $73728 = 2^{13}3^2$. The matrix $(6a_{iu})$ is integral and its row lattice has Smith invariants $1, 6, 6, 6, 6$. It follows that the $\mathcal{D}_u$ remain linearly independent over every A-algebra and the 97 coefficient rows span A^5. The certificate retains all 4656 original integer matrices, the 48 intertwiner systems, all T_j, P_j, all five global basis rows, and all 97 coefficients. Each equality in (E.52.15) is verified by literal matrix multiplication.

Define the homomorphism of (E.52.3) by

$$\Phi_T(B, a)_j = L_j(B)\left(I_4 + \sum_{u=0}^{4} a_u D_{u,j}\right), \quad \text{with top component 1.} \tag{E.52.16}$$

The identities (E.52.15) prove multiplicativity and preservation of Ω. At coordinate 18, the identity-coset third pair, the local radical is zero and extraction in (E.52.12) gives exactly B. After B is known, the displayed unit minor recovers a. This proves injectivity and explicitly describes the inverse on its image.

Surjectivity on N and the whole even-exponent subgroup. The elements x^6, y^6 belong to N and have θ-images $E_{21}(12)$ and $E_{12}(-12)$. For finite $T = \mathbf{Z}/m$, twelve is a unit and their powers give every upper and lower elementary matrix. These generate $\mathrm{SL}_2(T)$: over each local factor a unimodular column has a unit entry, elementary elimination reduces it to $(u, 0)$, and the diagonal matrix is a product of elementary matrices. The Chinese remainder maps give the assertion over T. This group is perfect. Indeed with $h = \operatorname{diag}(2, 1/2)$,

$$[h, E_{12}(t)] = E_{12}(3t), \qquad [h, E_{21}(t)] = E_{21}(-3t/4). \tag{E.52.17}$$

Both coefficients are units, so every elementary generator belongs to the commutator subgroup.

The coefficient rows above also prove that the additive projection of $\rho_m(N)$ is all of T^5. Its image under (E.52.16) is therefore a subdirect product of the perfect group $\mathrm{SL}_2(T)$ and the abelian group T^5. Goursat's lemma makes its common quotient both perfect and abelian, hence trivial: the commutator subgroup of an abelian group is the identity. Thus it is the whole displayed direct product. In particular (E.52.16) has image in G_m.

The five actual Schreier generators of K_2 in F are

$$x^2, \quad yxy^{-1}x^{-1}, \quad y^2, \quad xyxy^{-1}, \quad xy^2x^{-1}. \tag{E.52.18}$$

Their 240 induced matrices all have the same factorization (E.52.16). In the five radical coordinates, their rows are respectively

$$\begin{pmatrix} -1/2 & 1/3 & -2/3 & 2/3 & 1/2 \\ 4/3 & -4/3 & 2/3 & -5/3 & -1 \\ 1/3 & 0 & 0 & 0 & 0 \\ -5/6 & 2/3 & -1/3 & 4/3 & 1/2 \\ -1/3 & 2/3 & -1/3 & 1/3 & 0 \end{pmatrix}. \tag{E.52.19}$$

Also $N \subset K_2$: reduction of the actual modular map modulo four reads the two exponent parities on F, and N is killed by that map. Hence $\rho_m(N) = \rho_m(K_2)$ is the whole group in the first line of (E.52.5). Every statement here is compatible with reduction. Compactness of $\widehat{N}$ and K_2 now gives the same equality over R, by taking the closed fibres over all finite moduli. It also proves topological perfectness of $\mathrm{SL}_2(R)$ from the perfectness of every finite image.

Detection of the complete quotient A_2. For each of the 24 nonzero intertwiner coordinates, direct multiplication gives signs $\varepsilon_{x,j}, \varepsilon_{y,j} \in \{1, -1\}$ such that

$$M_j(x)T_j = \varepsilon_{x,j} T_j \theta(x), \qquad M_j(y)T_j = \varepsilon_{y,j} T_j \theta(y). \tag{E.52.20}$$

At coordinate 10 these signs are $(1, -1)$; at coordinate 26 they are $(-1, 1)$. They are literal distinct coordinates of the six-coset construction, and both matrices have $P_j T_j = I_2$ over A. For any $f \in \widehat{F}$, multiplication extends (E.52.20) to the two corresponding parity characters of its exponent sums. On an element of the image of K_2, formula (E.52.16) has sign $+1$ at every coordinate; the radicals annihilate these standard planes. As two is invertible and $\theta(f)$ is invertible, (E.52.20) forces both parities to vanish whenever $\rho_m(f) \in \rho_m(K_2)$. Thus the full inverse image of $\rho_m(K_2)$ in $\widehat{B}$ is exactly K_2: the top permutation first forces membership in $\widehat{F}$, and these two signs force membership in K_2. This proves the first quotient of (E.52.6), and its order is $[B : F][F : F'F^2] = 6 \cdot 4 = 24$.

The commutator image and every lower-central term. Conjugating every n_i by s, r, x, y gives another literal factorization in (E.52.16). The new coefficient rows equal the old row multiplied by a specified five-by-five matrix. All 388 rows are checked. Since the coefficient projection on N is a homomorphism, equality on its 97 free generators proves the equality for every element of N, at every finite modulus and over R. The matrices for x and y commute and square to the identity. This also follows directly from $\rho_T(K_2)$ being the displayed direct product, where conjugation acts trivially on its additive factor.

The closed normal subgroup $\widehat{F}'$ is normally generated by $c = [x, y]$. Its additive image is therefore generated by the four conjugates with conjugators $1, x, y, xy$, since K_2 acts trivially on that additive factor. The first three rows are exactly the rows of (E.52.4); the fourth is $(1/3, -1/3, 2/3, 1/3, 0)$. The first three are independent, their minor in columns $0, 2, 3$ (numbered from zero) has determinant $-1/3$, and the four rows have coefficient-lattice Smith invariants $1, 1, 1$ in that basis. They generate precisely $V_3(T)$ at every modulus. Moreover $\theta(\widehat{F}')$ is the full $\mathrm{SL}_2(T)$, because $\theta(\widehat{F})$ is onto and this target is perfect. The same perfect–abelian Goursat argument proves the formula for C_T.

In the ordered basis (E.52.4), conjugation by x and y acts on row vectors on the right by

$$X = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ -1 & -1 & -1 \end{pmatrix}, \qquad Y = \begin{pmatrix} 0 & 0 & 1 \\ -1 & -1 & -1 \\ 1 & 0 & 0 \end{pmatrix}. \tag{E.52.21}$$

The row lattice of $X - I$ and $Y - I$ has Smith invariants $1, 1, 4$. Since four is a unit in A,

$$[\rho_T(\widehat{F}), C_T] = C_T. \tag{E.52.22}$$

Indeed these differences give the whole additive factor, while commutators inside the perfect $\mathrm{SL}_2(T)$ factor give that factor. The equality $C_T' = \Phi_T(\mathrm{SL}_2(T) \times 0)$, and the same equality for $\rho_T(\widehat{N}')$, follow directly from the two direct products.

For finite T, the index of C_T in the image of F is $4m^2$, by the already proved group orders. The elements x^2, y^2 in (E.52.19) have θ-images $E_{21}(4), E_{12}(-4)$; these have order dividing m, and their additive coordinates are also killed by m. Therefore $x^{2m} = y^{2m} = 1$ in G_m. The abelianization of $\rho_m(\widehat{F})$ is generated by these two elements, each of order dividing $2m$, and has order $4m^2$. The actual exponent-sum map therefore identifies it with $(\mathbf{Z}/2m)^2$. Its full kernel in $\widehat{F}$ is $\overline{\widehat{F}'\widehat{F}^{2m}}$. Retaining the same s, r and the quotient to $\mathfrak{S}_3$ proves $G_m/C_m \simeq A_{2m}$ as the actual quotient of $\widehat{B}$. This establishes all orders and maps in (E.52.6).

Multiplication by three is invertible on $(\mathbf{Z}/2m)^2$. Since $H = F'F^3$, its image is consequently the whole image of F. Thus its derived image is C_T, and (E.52.22) proves inductively the image C_T for every $\gamma_j\widehat{H}$, $j \geq 2$. The statements over R follow through the same compatible finite maps and compactness.

Finally fix $c \in C_R$. The closed sets

$$\gamma_j\widehat{H} \ \cap \ \rho_m^{-1}(c \bmod m), \qquad j \geq 2, \quad (m, 6) = 1, \tag{E.52.23}$$

are nonempty by the proved finite images. Any finite collection contains the set obtained by taking the largest j and the least common multiple of its moduli, so has nonempty intersection. Compactness of $\widehat{H}$ gives a common element of all these sets. It belongs to R_{nil} and maps to the prescribed c. The inverse inclusion follows from its containment in $\widehat{H}'$. This proves the residual equality and, by the elementary fibre formula for a group homomorphism, the stated full word coset. □

The two root programs `e_symplectic_five_integral_probe_root.py` and `e_symplectic_five_integral_residual_root.py` retain the exact matrices, source words, coordinate maps and integer lattices used in this proof. Their recorded runs take 4.643 and 11.403 seconds. The first program also compares all sixteen pair permutations literally with the frozen models at five and seven. The independent integral audit reconstructs all 4656 N matrices and all 240 K_2 matrices and verifies the elementary standard maps and the radical products. The second independent audit replays the complete residual certificate and checks all five K_2 Schreier generators under each of s, r, x, y, retaining the row-action convention. A further literal comparison reconstructs the induced s, r permutations at five and seven: all 29952 and 115200 entries, respectively, agree with the frozen uncompressed sources. The coset sections agree, so the intervening conjugating gauge is the identity. This result supplies the complete congruence source and its residual image. The common-source and full peripheral-fibre comparisons with the other retained quotients are separate maps to be established below; the four global tasks remain open at this stage.

5.15 The uniform signed quotient and the full congruence correction

Statement E.53 (Proved). The integral common source with the complete double cover. *Retain E.52's actual maps $\rho_m, \rho_R, \Phi_m, \Phi_R$, where $R = \prod_{p \geq 5} \mathbf{Z}_p$, and retain $\rho_L : \widehat{B} \twoheadrightarrow G_L$ from E.38–E.43. Write $Q_T = \Phi_T(1, V_3(T)) \subset C_T$ for $T = \mathbf{Z}/m$ or R. For $m > 1$ prime to six, the actual modular map*

$$q_{4m} : \widehat{B} \twoheadrightarrow Q_{4m} = \mathrm{SL}_2(\mathbf{Z}/4m)/\{\pm I\}, \quad s \longmapsto \left[\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}\right], \quad r \longmapsto \left[\begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix}\right] \tag{E.53.1}$$

factors through ρ_m. Together with the ordinary exponent quotient of E.52 it gives the actual isomorphism

$$G_m/Q_m \xrightarrow{\sim} \mathcal{J}_m, \qquad \mathcal{J}_m = A_{2m} \times_{A_2} Q_{4m}, \qquad |\mathcal{J}_m| = 24m^2 |\operatorname{SL}_2(\mathbf{Z}/m)|. \tag{E.53.2}$$

The two maps to A_2 are reduction of exponent sums modulo two and projective matrix reduction modulo four. Their values on s, r are the same displayed source values. Every comparison in (E.53.2) *commutes with reduction of m.*

Put $\mathcal{J}_1 = A_2$, and retain the E.50 epimorphism $G_L \twoheadrightarrow \mathcal{J}_5$. The maximum common quotient for the specified maps (ρ_L, ρ_R) is exactly $\mathcal{J}_5$. For finite m it is $\mathcal{J}_{\gcd(m,5)}$. Equivalently their actual joint images are

$$\begin{aligned} \operatorname{im}(\rho_L, \rho_R) &= G_L \times_{\mathcal{J}_5} G_R, \\ \operatorname{im}(\rho_L, \rho_m) &= G_L \times_{\mathcal{J}_{\gcd(m,5)}} G_m. \end{aligned} \tag{E.53.3}$$

These are equalities of the source images under the displayed projections.

Let $\Gamma_5(1) = \ker(\operatorname{SL}_2(\mathbf{Z}_5) \to \operatorname{SL}_2(\mathbf{F}_5))$. The exact correcting-subgroup image is

$$\rho_R(R_{\mathrm{nil}} \cap \ker \rho_L) = \Phi_R \left(\Gamma_5(1) \times \prod_{p \geq 7} \operatorname{SL}_2(\mathbf{Z}_p), V_3(R) \right) =: \mathcal{K}_R. \tag{E.53.4}$$

Intersecting the source on the left with $\ker \rho_{D_{23}}$ leaves that image unchanged. Intersecting additionally with $\ker \rho_5 \cap \ker \rho_7$ gives precisely

$$\mathcal{K}_R^{35} = \Phi_R \left(\Gamma_5(1) \times \Gamma_7(1) \times \prod_{p \geq 11} \operatorname{SL}_2(\mathbf{Z}_p),\ 5V_3(\mathbf{Z}_5) \times 7V_3(\mathbf{Z}_7) \times \prod_{p \geq 11} V_3(\mathbf{Z}_p) \right) \tag{E.53.5}$$

Here the argument of Φ_R is the displayed element of $\operatorname{SL}_2(R) \times V_3(R)$. Each prescribed target value in either correcting image has the full source fibre $k \ker(\rho_R|_K)$, where K is the corresponding displayed intersection and $k \in K$ is one preimage. Every such correction maps to the identity in every nilpotent continuous quotient of $\widehat{H}$.

Proof. We retain the central sign first, and then determine the kernel of each actual joint source map.

The canonical signed matrix on the pure subgroup. For a pure word $f \in F$, its ordinary integral modular matrix is determined up to its central sign. Exactly one sign has both diagonal entries equal to one modulo four. Indeed the off-diagonal entries are even, and determinant one makes the two odd diagonal residues equal modulo four. Products of matrices with diagonal entries one modulo four retain that property. The sign choice therefore defines a homomorphism on the pure subgroup, with

$$x \longmapsto \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}, \qquad y \longmapsto \begin{pmatrix} 1 & 0 \\ -2 & 1 \end{pmatrix}. \tag{E.53.6}$$

For finite quotients this means the compatible CRT representative with the indicated component modulo four; it also defines the corresponding continuous pure map. With $P = \left(\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right)$, its component modulo m is $P\theta(f)P$. For exponent sums (a, b) its component modulo four is

$$\begin{pmatrix} 1 & 2a \\ -2b & 1 \end{pmatrix}. \tag{E.53.7}$$

Both assertions follow on the two free generators from (E.53.6) and extend by multiplication. In particular $Q_4 \simeq A_2$ through this exact map.

An element of $\ker \rho_m$ lies in K_2 by E.52's two parity characters and has θ-value one. Equations (E.53.6)–(E.53.7) make its canonical matrix the identity modulo $4m$. Thus q_{4m} factors through G_m. It is onto: the product of the displayed s, r matrices is $E_{12}(-1)$; its powers and conjugation by s give all upper and lower elementary matrices. Elementary elimination over each prime-power factor and CRT prove generation of the full determinant-one group, hence of its specified projective quotient.

The kernel of $G_m \to A_{2m}$ is C_m. On C_m, the modular map has trivial four-component and its signed m-component is exactly $P\theta P$. The common kernel is consequently Q_m. The image of G_m/Q_m lies in the actual fibre product in (E.53.2). Now $|Q_{4m}| = 24|\operatorname{SL}_2(\mathbf{Z}/m)|$, because $|\operatorname{SL}_2(\mathbf{Z}/4)| = 48$ and the diagonal central sign has order two. The fibre product therefore has the order stated in (E.53.2), equal to $|G_m/Q_m|$ by E.52. The injective map is onto. This proves both the signed comparison and every transition map.

Finite quotients of the integral SL_2 factor. We give the group facts needed to control the full inverse system, including its central quotients. For $p \geq 5$ put $L_p = \operatorname{SL}_2(\mathbf{Z}_p)$ and $\Gamma_p(n) = \ker(L_p \to \operatorname{SL}_2(\mathbf{Z}/p^n))$. The map

$$\Gamma_p(n)/\Gamma_p(n+1) \longrightarrow \mathfrak{sl}_2(\mathbf{F}_p), \qquad 1 + p^n A \longmapsto A \bmod p \tag{E.53.8}$$

is an isomorphism of additive groups and is equivariant for the actual conjugation action of L_p. The determinant equation gives trace zero; the two elementary matrices and $\operatorname{diag}(1 + p^n a, (1 + p^n a)^{-1})$ give every basis vector, proving surjectivity.

This adjoint module is irreducible. Conjugation by $E_{12}(t)$ and $E_{21}(t)$ is a polynomial of degree at most two in t. Interpolation at $0, 1, -1$ shows that an invariant subspace is stable under brackets with the usual e, f, and hence with $h = [e, f]$. The h-weights $0, 2, -2$ are distinct for $p \geq 5$. Projection onto those three eigenspaces, followed by brackets with e, f, shows that any nonzero invariant subspace contains e, f, h.

Let $U \triangleleft L_p$ be open. The group $\operatorname{PSL}_2(\mathbf{F}_p)$ is simple by the following concrete argument. Its faithful projective action is primitive; the translations form an abelian normal subgroup of a point stabilizer, and their conjugates generate the group. The elementary commutators E.52.17 make it perfect. A nontrivial normal subgroup is transitive in this primitive faithful action. Modulo that subgroup, all conjugate translation groups have the same abelian image, because every group element is a product of a normal-subgroup element and a point stabilizer. The quotient is abelian, and perfectness makes it trivial. This proves simplicity. Consequently the reduction of U modulo p is central or is the whole group. In the latter case L_p/U is a finite p-group. It is also perfect by E.52.17, so it is trivial. The full-reduction conclusion also holds for a closed normal subgroup: its quotient is pro-p and topologically perfect; every finite quotient is a perfect finite p-group and is therefore trivial.

In the former case consider $U \cap \Gamma_p(1)$. Let n be the first layer in (E.53.8) where it has nonzero image. Irreducibility makes that image the whole layer. Choose three elements of U giving a basis there. For $a \geq 1$, direct binomial expansion gives

$$(1 + p^a A)^p \equiv 1 + p^{a+1} A \pmod{p^{a+2}}. \tag{E.53.9}$$

Successive powers of the chosen elements therefore span every later layer. Successive approximation and closedness prove $U \cap \Gamma_p(1) = \Gamma_p(n)$. The reduction of U is either 1 or $\{\pm I\}$. In the second case write an element above $-I$ as $-v$ with $v \in \Gamma_p(1)$. Its square lies in $\Gamma_p(n)$. The finite p-group $\Gamma_p(1)/\Gamma_p(n)$ has no element of order two, so $v \in \Gamma_p(n)$. Thus the possibilities are precisely

$$U = \Gamma_p(n), \quad U = \{\pm I\}\Gamma_p(n), \quad U = L_p. \tag{E.53.10}$$

The argument also proves that any closed normal subgroup with a nonzero first congruence layer contains all of $\Gamma_p(1)$.

Every nontrivial finite quotient of L_p therefore has $\mathrm{PSL}_2(\mathbf{F}_p)$ as a composition factor, and its centre has exponent dividing two. For the last assertion a matrix commuting with the two elementary generators is scalar, and the determinant equation gives $\pm I$. For a quotient by the central sign, reduce first modulo p. A central projective element is scalar there; its commutators with the elementary generators are consequently both signs with reduction $+1$, so they equal $+1$ before taking the quotient. The same scalar calculation finishes that case.

A continuous finite quotient of $\prod_{p\geq 5} L_p$ uses finitely many factors. Their images commute and generate it. If a product of elements from these factor images is central, each factor element commutes with its own factor image, because all other factor images commute with that image already. Each factor element has square one by the preceding paragraph, so the whole central product has square one. We have proved that every such finite quotient has centre of exponent dividing two.

The exact lower-central correcting image. By E.43, $\rho_L(R_{\mathrm{nil}}) = C_L$ is perfect. By E.52, $\rho_R(R_{\mathrm{nil}}) = C_R$. Apply the profinite Goursat argument to their actual joint image and put $\mathcal{K}_R = \rho_R(R_{\mathrm{nil}} \cap \ker \rho_L)$. Then $C_R/\mathcal{K}_R$ is a finite quotient of C_L, hence perfect. The additive factor Q_R is central in C_R. In this perfect quotient the image of $\mathrm{SL}_2(R)$ is the whole group: the commutator of the product of that image and the central additive image is just the first image. The additive image is consequently in the centre of a finite quotient of $\mathrm{SL}_2(R)$. That centre has exponent dividing two, while every finite quotient of Q_R has order prime to six. Thus $Q_R \subset \mathcal{K}_R$.

Let U be the resulting closed normal subgroup of $\mathrm{SL}_2(R)$. The literal E.38 source order is $|G_L| = 2^{82}3^{26}5^{21}7^2$. For $p \geq 11$, a nontrivial quotient of L_p would have the factor $\mathrm{PSL}_2(\mathbf{F}_p)$, whose order is divisible by p. Its order cannot divide $|G_L|$. Thus the projection of U onto every such L_p is all of L_p. For $p = 7$, E.51 proves that the very same $R_{\mathrm{nil}} \cap \ker \rho_L$ maps onto $\mathrm{SL}_2(\mathbf{F}_7)$. Its closed normal projection in L_7 is therefore all of L_7, by the full-reduction case above. For any one of these primes, commutating elements of U with elements supported in that one factor shows that U contains the pure factor L_p, because L_p is topologically perfect. Closedness then gives the entire product for $p \geq 7$.

The five-component is determined by an actual word. In the complete signed Ferrari source, both $\rho_L(x)$ and $\rho_L(y)$ have exact order 30. In particular x^{60}, y^{60} belong to $\ker \rho_L$. Choose $c \in R_{\mathrm{nil}}$ whose five-component under $\theta\rho_R$ is $E_{12}(1)$, using E.52's surjectivity of this same closed subgroup. Then $[x^{60}, c]$ lies in $R_{\mathrm{nil}} \cap \ker \rho_L$, since the residual is normal in $\widehat{B}$. Its five-component is the commutator of $E_{21}(120)$ and $E_{12}(1)$, and modulo 25 it is

$$\begin{pmatrix} 6 & 20 \\ 0 & 21 \end{pmatrix}. \tag{E.53.11}$$

Its nonzero first-layer vector is $\left(\begin{smallmatrix} 1 & 4 \\ 0 & 4 \end{smallmatrix}\right)$. The three explicit conjugate vectors in the certificate span $\mathfrak{sl}_2(\mathbf{F}_5)$. Equation (E.53.9) and closedness give the whole $\Gamma_5(1)$ in the five-projection of U. The other factors already belong to U separately, so $\Gamma_5(1)$ belongs as a pure factor as well. Conversely E.50's actual common signed quotient gives the map $C_L \twoheadrightarrow \mathrm{SL}_2(\mathbf{F}_5)$ whose composition with the source is $\theta\rho_5$. It forces the five-reduction of every element of U to be the identity. This proves precisely (E.53.4).

The whole common source. Put $M = \rho_R(\ker \rho_L)$. The two maps to $\mathcal{J}_5$ agree by E.50 and (E.53.2), so $M \subset \ker(G_R \to \mathcal{J}_5)$. It contains $\mathcal{K}_R$. Modulo $\mathcal{K}_R$, the restriction of the latter map to C_R is the isomorphism onto its $\mathrm{SL}_2(\mathbf{F}_5)$ subgroup. Consequently its remaining

kernel injects into the pure exponent quotient, and is exactly the kernel of reduction of the two even exponent coordinates from R to $\mathbf{F}_5$. Surjectivity onto that exponent kernel follows by correcting any chosen lift by its uniquely required $\mathrm{SL}_2(\mathbf{F}_5)$ component.

The images of x^{60}, y^{60} lie in M, have trivial signed component modulo five and have even exponent coordinates $(30,0),(0,30)$. Their closed spans in R^2 are exactly $5R^2$, because six is a unit in R and ordinary integers are dense in R through CRT. They therefore generate the entire remaining kernel just described. It follows that

$$M = \ker(G_R \to \mathcal{J}_5). \tag{E.53.12}$$

The actual joint source is a subdirect product; (E.53.12) and its agreed quotient identify it with the full fibre product in (E.53.3). Reduction to any finite m gives the second identity: only a first factor of five can remain in the displayed correcting kernel and exponent kernel; with no factor of five the quotient is exactly A_2.

Retaining the old small-prime and symplectic values. Every finite solvable quotient of $\mathcal{K}_R$ has order prime to six. Its perfect L_p factors have trivial solvable image, and the remaining factors are the pro-five $\Gamma_5(1)$ and the additive product at primes at least five. The group D_{23} is a finite solvable group whose order has only primes two and three. Their common quotient is therefore trivial. Applying Goursat to the actual image of $R_{\mathrm{nil}} \cap \ker \rho_L$ proves that intersecting its source with $\ker \rho_{D_{23}}$ still maps onto all of $\mathcal{K}_R$.

The maps ρ_5, ρ_7 factor through ρ_R. Hence imposing their kernels on a source mapping onto $\mathcal{K}_R$ gives exactly the intersection of $\mathcal{K}_R$ with their two target kernels. In the coordinates of E.52 that is precisely (E.53.5). The same argument about finite solvable quotients applies after this intersection, since the added seven-component is pro-seven. Finally, a homomorphism from $\widehat{H}$ to a nilpotent profinite group of finite nilpotency class kills some $\gamma_j \widehat{H}$ and hence R_{nil}. For an inverse limit of such quotients it kills this subgroup at every stage, and so kills it in the limit. The full coset assertions follow from the kernels of these specified continuous homomorphisms. □

The exact program `e_symplectic_integral_gl_comparison_root.py` reconstructs the complete signed source without executing the older distribution script that writes an output at import time. It checks the original signed model arrays, the exact orders of x, y in all 36 induced coordinates, all 102 canonical signed modular comparisons, the level-25 commutator and its three independent conjugate Lie vectors. Its run takes 3.779 seconds. The first run used an overstrong exact-order assertion of 60, which the computation rejected. That script and traceback are preserved. The actual order is 30; both power relations used in the proof are verified in the complete source, and no conclusion uses the rejected exact-order equality.

An independent complete proof audit and isolated replay reproduce every mathematical JSON entry in 4.551 seconds. A separate symbolic commutator calculation gives the same level-25 matrix, and the three conjugate vectors have determinant 3. All protected originals are unchanged.

6 F. Derived limits and applications

6.1 The residue complex and the derived-limit test

Statement F.1 (Proved). Localization with its global term retained. *Let L be a number field, let ℓ be a prime, and let S be a finite set of places containing the archimedean places and every place above ℓ. Put $X = \operatorname{Spec} \mathcal{O}_{L,S}$. Let M be a free finite-rank $\mathbb{Z}_\ell$-module with a continuous action of $G_{L,S}$. Put $M_n = M/\ell^n M$. For every finite place $v \notin S$, choose*

a decomposition group and arithmetic Frobenius, put $A_v = M(-1)$ with that residue-field action, and put $D_v = \mathrm{Frob}_v - 1$ on A_v. There are compatible distinguished triangles in the derived category of abelian groups

$$R\Gamma(X, M_n) \longrightarrow R\Gamma(G_L, M_n) \longrightarrow \bigoplus_{v \notin S} [A_v/\ell^n \xrightarrow{D_v} A_v/\ell^n][-1] \longrightarrow R\Gamma(X, M_n)[1].$$

The two terms in each displayed bracket have degrees zero and one. Define

$$R_A = \left\{ (a_v) \in \prod_{v \notin S} A_v : \textit{for every } n,\ a_v \in \ell^n A_v \textit{ outside finitely many } v \right\}.$$

Taking the derived inverse limit gives the triangle

$$R\Gamma_{\mathrm{cts}}(G_{L,S}, M) \longrightarrow R\Gamma_{\mathrm{cts}}(G_L, M) \longrightarrow [R_A \xrightarrow{D} R_A][-1] \longrightarrow R\Gamma_{\mathrm{cts}}(G_{L,S}, M)[1].$$

Here $D = (D_v)_v$ is the coordinatewise residue differential. In particular its degree-two sequence is

$$\ker D \longrightarrow H^2_{\mathrm{cts}}(G_{L,S}, M) \longrightarrow H^2_{\mathrm{cts}}(G_L, M) \longrightarrow R_A/DR_A \longrightarrow H^3_{\mathrm{cts}}(G_{L,S}, M).$$

The image DR_A here is the algebraic image. Every arrow is the map induced by restriction or the localization triangle; changing a decomposition-group choice conjugates the corresponding stalk and gives an isomorphic complex.

Proof. For a finite set J of places outside S, take localization for the open immersion $X - J \hookrightarrow X$. Over the strict henselization at v, the sheaf M_n is constant. Wild inertia is pro-p_v, so averaging on finite quotients annihilates its positive-degree cohomology with ℓ-primary coefficients. The relevant tame inertia is $\mathbb{Z}_\ell(1)$; its cochain complex has cohomology M_n in degree zero and $M_n(-1)$ in degree one. The support complex is consequently $M_n(-1)[-2]$. Descending to the residue field gives the first triangle with J in place of all places.

The locally constant sheaf comparison is Milne, II.2.9; localization and its support maps are II.0.1(f) and II.1.7(b) [Mil06]. Filtered direct limits over J are exact. Every continuous cochain of a profinite group with finite coefficients factors through a finite quotient, and its number field is ramified at finitely many places. Thus the generic-field term is precisely $R\Gamma(G_L, M_n)$ and the support term is a direct sum.

The residue Galois group is procyclic. Its usual two-term resolution computes cohomology as $[A_v/\ell^n \xrightarrow{D_v} A_v/\ell^n]$. In both degrees the direct-sum reduction maps are surjective. Their derived inverse limits are therefore their ordinary inverse limits, which are exactly R_A: reduction modulo ℓ^n has finite support for every n. The matrices D_v are integral, so they preserve this space.

For completeness, continuous cochains with values in M are the inverse limit of the finite-coefficient cochain complexes. The reduction map on cochains is surjective, since a continuous finite-valued function can be lifted value by value. The resulting Milnor comparison is Jannsen, Proposition 1.6 and Theorem 2.2 [Jan88]. It identifies the derived inverse limit used here with continuous cochain cohomology. Taking the long exact sequence proves the assertion. Real places have not been discarded: their possible 2-primary contribution remains in the global term and in its degree-three group. □

Statement F.2 (Proved). The exact global correction to the residue derived limit. *With the data of F.1, define inverse systems*

$$P_n = \bigoplus_{v \notin S} \ker(D_v : A_v/\ell^n \to A_v/\ell^n), \qquad E_n = \mathrm{im}\big(P_n \to H^2(G_{L,S}, M_n)\big).$$

There is a natural short exact sequence

$$0 \longrightarrow \operatorname{coker}\bigl(\varprojlim P_n \to \varprojlim E_n\bigr) \longrightarrow \varprojlim{}^1 H^1(G_L, M_n) \longrightarrow \varprojlim{}^1 P_n \longrightarrow 0. \tag{F.2.1}$$

The right-hand group has the explicit presentation

$$\varprojlim{}^1 P_n = \frac{R_A \cap \prod_{v\notin S} D_v A_v}{D R_A}. \tag{F.2.2}$$

The embedding of this group into the residue cohomology is the kernel of the map

$$R_A/DR_A \longrightarrow \varprojlim_n \bigoplus_{v\notin S} A_v/(\ell^n A_v + D_v A_v).$$

Jannsen's exact sequence for the full global group is

$$0 \longrightarrow \varprojlim{}^1 H^1(G_L, M_n) \longrightarrow H^2_{\mathrm{cts}}(G_L, M) \longrightarrow \varprojlim_n H^2(G_L, M_n) \longrightarrow 0. \tag{F.2.3}$$

Proof. At level n localization gives

$$H^1(G_{L,S}, M_n) \longrightarrow H^1(G_L, M_n) \longrightarrow P_n \longrightarrow H^2(G_{L,S}, M_n).$$

The two cohomology groups with finite S are finite by Milne I.4.15. Let Q_n be the kernel of $P_n \to E_n$, and let F_n be the image of the first arrow. The exact sequences $0 \to F_n \to H^1(G_L, M_n) \to Q_n \to 0$ and $0 \to Q_n \to P_n \to E_n \to 0$ give (F.2.1). Indeed all the F_n and E_n are finite, hence their towers are Mittag–Leffler and their first derived limits vanish; second derived limits of towers vanish.

Put $B_n = \bigoplus_v A_v/\ell^n$ and $I_n = DB_n$. Both B_n and I_n have surjective transition maps: a representative of $D_v a$ lifts by lifting a. Apply inverse limits to $0 \to P_n \to B_n \to I_n \to 0$. The result is $\varprojlim{}^1 P_n = (\varprojlim I_n)/DR_A$. An element of R_A belongs to $\varprojlim I_n$ exactly when each of its coordinates belongs to $D_v A_v + \ell^n A_v$ for every n. The image of an endomorphism of a finite-rank $\mathbb{Z}_\ell$-module is closed; its Smith form proves that the intersection is $D_v A_v$. This proves (F.2.2). Applying the same argument to $0 \to I_n \to B_n \to B_n/I_n \to 0$ gives the asserted kernel and surjectivity onto the displayed limit. Finally (F.2.3) is the continuous-cochain comparison already proved in F.1. □

Statement F.3 (Proved). A complete vanishing test over number fields. *For each $v \notin S$, let $e_{v,1}, \dots, e_{v,s_v}$ be the valuations of the nonzero Smith entries of D_v on A_v. A unit entry has valuation zero; a zero Smith entry is excluded. The following conditions are equivalent:*

(i) $\varprojlim{}^1 H^1(G_L, M/\ell^n) = 0$;

(ii) the tower P_n in F.2 is Mittag–Leffler;

(iii) the nonnegative integers $e_{v,i}$ are bounded, uniformly over v and i;

(iv) the finite groups

$$B_v = \frac{H^0(\mathbb{F}_v, A_v \otimes \mathbb{Q}_\ell/\mathbb{Z}_\ell)}{H^0(\mathbb{F}_v, A_v \otimes \mathbb{Q}_\ell/\mathbb{Z}_\ell)_{\mathrm{div}}}$$

have uniformly bounded orders.

The subscript div *denotes the maximal divisible subgroup. No boundedness condition on the orders of $H^2(G_{L,S}, M/\ell^n)$ is required. Their contribution to the full group is exactly the left term of (F.2.1).*

Proof. The Smith form is an isomorphism of source and target $\mathbb{Z}_\ell$-modules; it is chosen separately at each place and is compatible with reduction. For a nonzero entry ℓ^e, the image of its kernel at level m in level $n \le m$ is

$$\ell^{\max(0,m-e)}(\mathbb{Z}/\ell^n),$$

with the usual interpretation that it is zero when $m - e \ge n$. A zero Smith entry contributes the surjective tower $\mathbb{Z}/\ell^n$. Thus a bound $e \le b$ makes the images stationary by level $n + b$. Conversely, unbounded exponents give, for every m, a coordinate whose image at level one is nonzero from level m, but becomes zero at a later level. Hence the images at level one never become stationary. This proves (ii)⇔(iii).

The same Smith form gives

$$H^0(\mathbb{F}_v, A_v \otimes \mathbb{Q}_\ell/\mathbb{Z}_\ell) \simeq (\mathbb{Q}_\ell/\mathbb{Z}_\ell)^{\operatorname{rank}\ker D_v} \oplus \bigoplus_{i=1}^{s_v} \mathbb{Z}/\ell^{e_{v,i}}.$$

This is an isomorphism of abelian groups; the quotient by the maximal divisible subgroup is canonical. The ranks s_v are bounded by $\operatorname{rank} M$, proving (iii)⇔(iv).

Unbounded exponents give a nonzero element of (F.2.2) directly. Choose distinct v_j and Smith source basis vectors x_j with associated exponents at least j. Set $z_{v_j} = D_{v_j} x_j$ and set the other coordinates to zero. The family z is restricted. Every preimage of z_{v_j} differs from x_j by an integral kernel vector, so its reduction modulo ℓ in $A_{v_j}/\ker D_{v_j}$ is nonzero. No preimage family is restricted. Thus (F.2.2) is nonzero, and the surjection (F.2.1) proves that (i) implies (iii).

It remains to verify that bounded exponents also annihilate the global correction; finiteness at each level alone does not supply this step. Let

$$P_n^0 = \bigoplus_v (\ker D_v)/\ell^n \ker D_v \ \subseteq P_n.$$

The kernels are saturated submodules, so the inclusion is injective. A uniform bound b implies that the tower P_n/P_n^0 is zero after shifting the index by b. Put $C = H^2_{\mathrm{cts}}(G_{L,S}, M)$. It is a finitely generated $\mathbb{Z}_\ell$-module. To check this last input, finite-coefficient cohomology is finite and the Milnor term from $H^1(G_{L,S}, M_n)$ vanishes. Thus $C = \varprojlim H^2(G_{L,S}, M_n)$ is compact. The coefficient exact sequence embeds $C/\ell C$ in the finite group $H^2(G_{L,S}, M/\ell)$. Choose lifts $c_1, \dots, c_d$ of generators of this quotient. Successive division modulo their span expresses every $c \in C$ as $\sum_i a_i c_i$ with $a_i \in \mathbb{Z}_\ell$: the remainder after n divisions belongs to $\ell^n C$, and these subgroups tend to zero since their images in every fixed finite coefficient group eventually vanish. This proves finite generation directly, and also verifies the input to Jannsen Corollaries 4.9–4.10.

Each integral local invariant has a compatible Gysin image in C. Let $J \subseteq C$ be the $\mathbb{Z}_\ell$-submodule generated by all these images. Since C is noetherian, J is generated by finitely many such images and is closed. The image E_n^0 of P_n^0 in $H^2(G_{L,S}, M_n)$ is exactly the image of J. The quotient tower E_n/E_n^0 is zero after the same fixed shift, so $\varprojlim E_n = \varprojlim E_n^0$. Compactness of the finitely generated module J, together with $C = \varprojlim H^2(G_{L,S}, M_n)$, identifies this limit with J.

A finite set of local invariant vectors whose images generate J gives a map from a finite direct sum of their $\mathbb{Z}_\ell$-spans into $\varprojlim P_n$. Its composite onto J is surjective. Consequently

the left term of (F.2.1) vanishes. The right term vanishes by the Smith calculation, proving (iii)⇒(i). The proof includes $\ell = 2$ and real places, since it uses their finite cohomology through C without assuming cohomological dimension two. □

Statement F.4 (Refuted). Total orders of residue invariants do not determine vanishing. *The orders of the unquotiented groups $H^0(\mathbb{F}_q, M(-1) \otimes \mathbb{Q}_3/\mathbb{Z}_3)$, even together with all the orders of $H^2(G_{\mathbb{Q},S}, M/3^n)$, do not determine the vanishing in F.3. For $S = \{3, \infty\}$ the two lattices*

$$M^{(0)} = \mathbb{Z}_3(1)^2, \qquad M^{(1)} = \mathbb{Z}_3(1) \oplus \mathbb{Z}_3(2)$$

have the same stated data: every residue invariant group has countably infinite order, and every displayed global H^2 group is zero. Their derived limits are respectively zero and nonzero. The correct order test is F.3(iv).

Proof. For $M^{(0)}(-1)$ every D_q is zero; there is no nonzero Smith entry. For $M^{(1)}(-1)$ the matrix is $\mathrm{diag}(0, q-1)$. Primes congruent to 1 modulo 3^j give unbounded nonzero Smith valuations, so F.3 proves the two answers. The residue invariants are respectively $(\mathbb{Q}_3/\mathbb{Z}_3)^2$ and $\mathbb{Q}_3/\mathbb{Z}_3 \oplus (\mathbb{Q}_3/\mathbb{Z}_3)[q-1]$; both have countably infinite underlying sets.

For the twist one, Kummer theory on $\mathbb{Z}[1/3]$ gives the vanishing of H^2 because its Picard group is zero and its Brauer group has no 3-primary part. The latter follows from the invariant-sum sequence: only the place 3 can contribute a nonzero 3-primary local invariant, and its sum must be zero. For twist two modulo 3, the module is the trivial $\mathbb{F}_3$. Kronecker–Weber gives $H^1(G_{\mathbb{Q},S}, \mathbb{F}_3)$ dimension one. The global Euler characteristic (Milne I.5.1) is one; thus H^2 is zero. Filtering the higher coefficients by powers of 3 proves the same vanishing at every level. This independently confirms the precise input from LM7 E.32g used in the example. □

Statement F.5 (Proved). Coefficient modules already determined. *Over every number field L the derived limit in F.3 vanishes for $\mathbb{Z}_3(1)$ and for the restrictions of $\mathbb{Z}_3(1)[D]$ and W from the cubic. It is nonzero for the restrictions of T, V, $\mathcal{K}$, and $\bigwedge^2 V$. It is also nonzero for every nonzero torsion-free graded lattice of degree $r \geq 3$ obtained as a subquotient of tensor powers $V^{\otimes r}$ and occurring in a lower-central quotient here. These statements concern each specified finite-rank lattice; they do not evaluate a section obstruction.*

Proof. For twist one the residue differential is zero. The cusp permutation action factors through $\mathrm{Gal}(\mathbb{Q}(\zeta)/\mathbb{Q})$ of order two. On every $\mathbb{Z}_3$-lattice with that action, averaging splits the two eigenspaces integrally. After the twist the nonzero Smith entries of a residue permutation minus one are units, including the quotient by the diagonal defining W. F.3 proves vanishing.

For $T(-1)$, the eigenvalues of arithmetic Frobenius at a good place v have complex absolute value $(Nv)^{-1/2}$; consequently D_v is injective. The representation modulo 3^n cuts out a finite Galois extension of L. Chebotarev supplies infinitely many places splitting in this extension and in the 3^n-cyclotomic extension. At these places $D_v \equiv 0 \pmod{3^n}$. Its nonzero Smith valuations are therefore unbounded. For $\mathcal{K}(-1)$, the two weight pieces have absolute values Nv and $(Nv)^{1/2}$, so the same argument applies.

The generalized Jacobian extension is killed by 3 in the isogeny category because every cusp difference has order dividing 3 on the elliptic curve. Thus $V \otimes \mathbb{Q}_3 = (W \otimes \mathbb{Q}_3) \oplus (T \otimes \mathbb{Q}_3)$. To transfer the boundedness test, choose integral lattices on this rational decomposition and a positive integer b such that the original and split lattices contain 3^b times one another. Replacing one by the other changes the largest nonzero Smith valuation by at most $2b$; this follows by applying both inclusions to preimages of D_v. Hence the unboundedness from T

persists in V. The same argument for exterior squares retains the $\mathcal{K}$ summand and proves the assertion for $\bigwedge^2 V$.

In degree $r \geq 3$, the eigenvalues before twisting have absolute values among $(Nv)^{a/2+b}$ with $a+b=r$ and $a,b \geq 0$. After twisting, every such value is at least $(Nv)^{r/2-1} > 1$. Subquotients retain a subset of these eigenvalues. Thus D_v is injective, and splitting places for the finite coefficient representation again prove unboundedness. A fixed finite torsion part of a finitely generated coefficient module contributes an eventually constant coefficient system; it creates no additional first derived limit. This last observation does not assert that an unevaluated graded quotient is torsion-free. □

Statement F.9 (Proved). The genuine integral class-two action for the colliding-cusp curve. *Let $K_0 = \mathbb{Q}((t))$ and $U_t = \mathbb{P}^1_{K_0} \setminus \{0, t, 1, \infty\}$. Use the coefficient splitting of G_{K_0} which fixes compatible roots of t. There is a based splitting of the arithmetic fundamental-group extension, constructed from the node chart below, for which the free class-two pro-ℓ group has a basis*

$$x = b, \qquad y = d = ab, \qquad z = c \tag{F.9.1}$$

with the following genuine continuous action:

	x	y	z
τ	yxy^{-1}	y	z
$g \in G_{\mathbb{Q}}$	$x^{\chi_\ell(g)}$	$y^{\chi_\ell(g)}$	$z^{\chi_\ell(g)}$.

(F.9.2)

The inverse free-basis change is $a = yx^{-1}$, $b = x$, $c = z$. These formulas hold integrally for every prime, including two. In the original generators, putting $c_{ab} = [a, b]$ and $\lambda = \chi_\ell(g)$, the constant action is

$$g(a) = a^\lambda c_{ab}^{-\binom{\lambda}{2}}, \qquad g(b) = b^\lambda, \qquad g(c) = c^\lambda. \tag{F.9.3}$$

The inertia action in that basis is exactly (F.7) of FLAGGED_149 F17.3. The comparison changes the free basis and specifies a splitting; it preserves the arithmetic extension over G_{K_0}.

Proof. Blow up the point $x = t = 0$ of the marked projective line model. Near its single node use coordinates $X = x$ and $V = t/x$, so $XV = t$. The outer component is the tripod in coordinate X, with node at zero and marked points at one and infinity. The bubble is the tripod in coordinate V, with node at zero, the cusp t at one, and the cusp 0 at infinity. The branch parameters X, V have product exactly t; no constant factor has been inserted.

The punctured normal-crossing node chart has arithmetic tame group

$$(\widehat{\mathbb{Z}}(1)^2) \rtimes G_{\mathbb{Q}} \longrightarrow \widehat{\mathbb{Z}}(1) \rtimes G_{\mathbb{Q}} = G_{K_0}, \quad ((r,s), g) \longmapsto (r+s, g). \tag{F.9.4}$$

This is seen on the compatible covers extracting roots of X, V; the base cover extracts roots of XV. Choose their products to be the specified roots of t. Coefficient automorphisms fix these formal roots and act on their root-of-unity multipliers by χ. The map in (F.9.4) then has the continuous group section $(r, g) \mapsto ((r, 0), g)$. The homomorphism from the node chart's arithmetic group to that of U_t, induced by the punctured formal-neighborhood map, sends this group section to an actual section over G_{K_0}. The kernel direction $(r, -r)$ is the vanishing loop d; its primitive abelianization proves injectivity on this direction. Choose its orientation and the lift of τ so that the associated Dehn twist conjugates the bubble group by d and fixes the outer group. This is the annulus cut-and-glue calculation which gives (F.6) in the attached return.

We specify why the component comparison respects arithmetic. Logarithmic specialization for this proper marked semistable model is Theorem 3.4.6 of Stix's thesis [Sti02]; the residue

characteristic is zero, so its prime-to-residue- characteristic assertion applies to every pro-ℓ quotient. The arithmetic van Kampen presentation is its equation (6.2.5), with component comparison Lemma 6.2.8 and injective edge maps Corollary 6.2.9. The dual graph here is a single edge, so the based geometric amalgam is

$$\widehat{F}(d,c) *_{\langle d\rangle} \widehat{F}(d^{-1},b). \tag{F.9.5}$$

Its arithmetic amalgam uses the same coefficient section and the same node root fiber functor on both sides of (F.9.4). Thus the continuous maps from the two component groups, and their common inertia inclusion, are $G_{\mathbb{Q}}$- equivariant with these actual sections. The common generator is primitive in both vertex groups; eliminating its duplicate in the amalgam gives the free basis b, d, c. The same elimination on finite ℓ-group quotients proves the pro-ℓ comparison.

On each component use the standard path from the unit tangent at zero towards the point one. For a tripod with these data the based arithmetic action sends its first loop to its λ-th power and its second to a conjugate of its λ-th power by an element of the commutator subgroup. The latter assertion is the normalized action in Wickelgren, §2.1, equation for the G_k-action [Wic12]; its abelian path cocycle is zero because the two leading units are one. In the class-two quotient this conjugator is central. The outer pair (d, c) and bubble pair (d^{-1}, b) therefore both have the pure-power action in (F.9.2). Compatibility in (F.9.4) proves that they give the same based action on (F.9.5). The Dehn twist gives the first row of (F.9.2).

For clarity about the original peripheral generators, the standard bubble loop at infinity is $b^{-1}d$. Conjugating its path by d gives the inertia generator $a = db^{-1}$, so $ab = d$ exactly. This is the indicated Nielsen basis change, with the displayed inverse; it is a continuous automorphism of each free pro-ℓ group. Both a and b are conjugated by d under the twist, proving (F.7). Finally the class-two collection identity

$$(ab)^{\lambda} = a^{\lambda} b^{\lambda} [a,b]^{-\binom{\lambda}{2}}$$

gives (F.9.3). The polynomial $\binom{\lambda}{2} = \lambda(\lambda-1)/2$ is integral for $\lambda \in \mathbb{Z}_2$ and extends continuously from integers; thus no division by a nonunit is being performed in the group law. This retains the sign information which would be lost by declaring the original pair a, d simultaneously diagonal without a tangent comparison at the prime two. □

6.2 Sections over the Laurent-series field

We use the arithmetic identification and the basis $(x, y, z) = (b, ab, c)$ of F.9. Put $G = G_{\mathbb{Q}((t))}$ and $G_0 = G_{\mathbb{Q}}$. The coefficient splitting fixes all compatible roots of t. A chosen tame generator is denoted by τ; only its pro-ℓ component enters the following formulas. All cohomology groups and cocycles below are continuous.

Statement F.10 (Proved). An integral tame equation, including the prime two. *Write the class-two free pro-ℓ group P_2 in Hall coordinates*

$$x^{u_1} y^{u_2} z^{u_3} [x,y]^{w_{12}} [x,z]^{w_{13}} [y,z]^{w_{23}} = (u, w).$$

These give a homeomorphism of profinite sets $P_2 \simeq \mathbb{Z}_\ell^3 \times \mathbb{Z}_\ell^3$ with law

$$(u,w)(v,h) = (u+v, w+h+F(u,v)), \qquad F(u,v) = (-u_2v_1, -u_3v_1, -u_3v_2).$$

Put $L(u) = (-u_1, 0, 0)$ and $B(u,v) = F(u,v) - F(v,u)$. The arithmetic actions are

$$\tau^a(u,w) = (u, w + aL(u)), \qquad g(u,w) = (\chi(g)u, \chi(g)^2 w) \quad (g \in G_0).$$

For a section with geometric value (t, w) at τ and (r_g, z_g) at $g \in G_0$, the tame relation is exactly

$$(\chi(g)-1)w + B(r_g, t) - L(r_g) - \frac{\chi(g)-1}{2}\{F(t,t) + L(t)\} = 0. \tag{F.10.1}$$

The expression $(\chi(g)-1)/2$ is integral also for $\ell = 2$, because the 2-adic cyclotomic character is odd. No Baker–Campbell–Hausdorff denominator is used in this statement.

Proof. Moving x^{v_1} and y^{v_2} left through the preceding powers of y, z gives the displayed F with the convention $[x, y] = xyx^{-1}y^{-1}$. This proves the group law and, by uniqueness of Hall coordinates at each finite quotient, its topological interpretation. Conjugation by y sends x to $x[x, y]^{-1}$ and fixes y, z, proving the inertia action. The constant action is F.9.

Let $h(t) = F(t, t) + L(t)$. Integer powers of the lifted inertia element $((t, w), \tau)$ have geometric coordinates

$$(nt, nw + \tbinom{n}{2}h(t)).$$

Induction follows from the group law: the next multiplication adds $nF(t, t)$ and $nL(t)$. The integer-valued polynomial $\binom{n}{2}$ extends continuously to $\mathbb{Z}_\ell$, giving the same formula for every ℓ-adic power. The parts of tame inertia at other primes act trivially on P_2 and have trivial geometric cocycles, since a continuous homomorphism from a pro-prime-to-ℓ group to a pro-ℓ group is trivial.

Write $q = \chi(g)$. The equality $s(g)s(\tau) = s(\tau)^q s(g)$ has equal abelian coordinates $r_g + qt$ on both sides. Its central defect is

$$q(q-1)w + q\{B(r_g, t) - L(r_g)\} - \tbinom{q}{2}h(t).$$

Division by the unit q proves (F.10.1). The program `f_tame_relation.py` checks this identity symbolically over $\mathbb{Q}$ and by exact integer arithmetic in 30 000 trials at moduli $8, 9, 25, 27, 32, 49$. The symbolic identity and the integral power formula, rather than those trials, prove the assertion at every coefficient level. □

Statement F.11 (Proved). The continuous obstruction group and all abelian fibers. *There is a split isomorphism of $\mathbb{Z}_\ell$-modules*

$$H^2_{\mathrm{cts}}(G, \mathbb{Z}_\ell(2)^3) \simeq H^2_{\mathrm{cts}}(G_0, \mathbb{Z}_\ell(2))^3 \oplus H^1_{\mathrm{cts}}(G_0, \mathbb{Z}_\ell(1))^3. \tag{F.11.1}$$

The first summand is inflation. A splitting of the second is cup product with the Kummer class of the uniformizer t, with the orientation chosen so that the residue at τ is the identity. The first summand carries a nonzero first-derived-limit term for every ℓ. The second carries none.

Let

$$\mathscr{K}_\ell = H^1(G_0, \mathbb{Z}_\ell(1)) = \begin{cases} \mathcal{R}_{\mathbb{P}}(\mathbb{Z}_\ell), & \ell \neq 2, \\ (\mathbb{Z}/2)\kappa(-1) \oplus \mathcal{R}_{\mathbb{P}}(\mathbb{Z}_2), & \ell = 2. \end{cases}$$

Let ϵ_ℓ be zero for odd ℓ and $\kappa(-1)$ for $\ell = 2$. The abelian section parameters are $(r, t) \in \mathscr{K}_\ell^3 \times \mathbb{Z}_\ell^3$. They lift to P_2 exactly when the following six classes vanish:

$$r_1 \cup r_2, \quad r_1 \cup r_3, \quad r_2 \cup r_3 \quad \text{in } H^2_{\mathrm{cts}}(G_0, \mathbb{Z}_\ell(2)), \tag{F.11.2}$$

and

$$\begin{aligned} &(t_2+1)r_1 - t_1 r_2 + \epsilon_\ell\, t_1(t_2+1), \\ &t_3 r_1 - t_1 r_3 + \epsilon_\ell\, t_1 t_3, \\ &t_3 r_2 - t_2 r_3 + \epsilon_\ell\, t_2 t_3 \end{aligned} \qquad \text{in } \mathscr{K}_\ell. \tag{F.11.3}$$

Every nonempty fiber of geometric conjugacy classes is a torsor under $H^1_{\mathrm{cts}}(G_0, \mathbb{Z}_\ell(2))^3$. In (F.11.2), vanishing means continuous cohomology, computed by F.1–F.3, including the finite-S term and restricted convergence.

Proof. For $M = \mathbb{Z}_\ell(2)^3$, inertia acts trivially on the coefficient module. Its cohomology has only degrees zero and one; the latter is $M(-1)$. The split group extension and the uniformizer Kummer class give, at every finite coefficient level, inflation and a right inverse to the residue map by cup product. Its restriction to inertia is the identity $\mathbb{Z}_\ell(1) \to \mathbb{Z}_\ell(1)$. Thus the two-row Hochschild–Serre complex splits in the indicated degrees. The same constructions commute with all coefficient reductions. Derived inverse limits give (F.11.1), so the splitting also retains the Milnor terms.

For twist two over $\mathbb{Q}$, the residue differential away from ℓ is multiplication by $q - 1$. Its nonzero Smith valuations are unbounded, by primes $q \equiv 1 \pmod{\ell^n}$; F.3 gives a nonzero derived limit. For twist one the residue differential is zero, so F.3 gives vanishing of its $\varprojlim^1 H^1$ term. The Milnor term attached directly to the H^1 summand in (F.11.1) is $\varprojlim^1 H^0(G_0, \mathbb{Z}/\ell^n(1))$; it vanishes because all these degree-zero groups are finite. The Kummer description of $\mathscr{K}_\ell$ follows directly from $\mathbb{Q}^\times = \{\pm 1\} \times \bigoplus_p p^{\mathbb{Z}}$, together with the finite H^0 coefficient comparison.

An abelian cocycle is specified by a cocycle $r : G_0 \to \mathbb{Z}_\ell(1)^3$ and a vector t on the chosen inertia generator. The tame relation on this abelian group is automatic, since both sides scale t by $\chi(g)$. The central cocycle equation on G_0 is

$$z_{gh} = z_g + \chi(g)^2 z_h + F(r_g, \chi(g) r_h).$$

Its three cohomology classes are the three cup products in (F.11.2): the negative reverse cup represents the forward cup, including at two by the cochain homotopy for graded commutativity. Vanishing gives continuous choices of z.

The remaining equation (F.10.1) is a coboundary equation with coefficients $\mathbb{Z}_\ell(1)^3$ for the single vector w. For odd ℓ, $(\chi - 1)/2$ is the coboundary of $1/2$. For $\ell = 2$, the compatible roots of -1 identify this cocycle with $\kappa(-1)$: on a root of unity of order 2^{n+1}, its Kummer exponent is $(\chi - 1)/2$ modulo 2^n. Expanding $B(r, t) - L(r)$ and $F(t, t) + L(t)$ now gives precisely (F.11.3). Vanishing of these classes supplies an integral w. The cocycle on G_0, the lifted inertia generator, and their verified tame relation define a continuous homomorphism on $\widehat{\mathbb{Z}}(1) \rtimes G_0$; the power formula in F.10 proves continuity on inertia. This proves sufficiency as well as necessity.

For fixed abelian parameters, central lifts differ by $H^1(G, M)$. Hochschild–Serre gives $H^1(G, M) = H^1(G_0, M)$, because $H^0(G_0, M(-1)) = 0$. There is no further identification by an abelian geometric stabilizer: $H^0(G, \mathbb{Z}_\ell(1)^3) = 0$. The central-extension exact sequence therefore gives the asserted torsor, with the chosen section origin as its sole ambiguity. □

Statement F.12 (Proved). The fully evaluated three-adic Laurent-series test. *For $\ell = 3$, write each $r_i = \sum_p r_{i,p} \kappa_p$ as a restricted family. At $q \neq 3$, choose compatible finite-field Kummer logarithms $\rho_{q,p}$ of the units p, for $p \neq q$, modulo $q - 1$, and use integral representatives. Put*

$$R_{i,q} = \sum_{p \neq q} r_{i,p} \rho_{q,p}, \qquad c_{ij,q} = -\frac{R_{i,q} r_{j,q} - R_{j,q} r_{i,q}}{q - 1}.$$

The exact conditions (F.11.2) are that each of the three families $(c_{ij,q})_{q \neq 3}$ is integral and restricted. Condition (F.11.3) is the displayed system of three linear equations in the restricted Kummer families, with $\epsilon_3 = 0$. Every nonempty fiber has 27 section classes.

Proof. The cochain localization calculation of LM7 E.32l–E.32n applies with commutator coefficient one instead of nine. The comparison is explicit: the class-two law in logarithmic coordinates has central term $\frac{1}{2}(u\wedge v)$, so its central tame defect is $q\{(q-1)c+R_{i,q}r_{j,q}-R_{j,q}r_{i,q}\}$. Replacing the actual Frobenius Kummer values by the chosen finite-field representatives adds $(q-1)$ times an integral restricted family: each difference contains the ramification coefficient $r_{i,q}$ or $r_{j,q}$. This proves the same residue representative in the continuous quotient, including its convergence requirement. For this coefficient, the finite-S H^2 is zero by the independently checked calculation in F.4.

Finally $H^1_{\mathrm{cts}}(G_{\mathbb{Q}},\mathbb{Z}_3(2))\simeq\mathbb{Z}/3$. One direct verification uses $H^0(G_S,\mathbb{Z}/3^n(2))$ of order three, the vanishing of $H^2(G_S,\mathbb{Z}/3^n(2))$, and the real Euler characteristic. They give $|H^1(G_S,\mathbb{Z}/3^n(2))|=3$ for all n. The nonzero compatible class is $g\mapsto(\chi_3(g)^2-1)/3$; its class has exact order three, since χ_3^2-1 has minimum valuation one. Integral residues in degree one vanish because multiplication by $q-1$ is injective on $\mathbb{Z}_3$. Thus the global and finite-S integral H^1 agree and have the asserted order. The torsor in F.11 has order $3^3=27$. □

6.3 The fixed-lift compactness question in Task F.4

Statement F.20 (Proved). The finite coboundary criterion in the profinite kernel. *Use the data of Task F.4: $K/\mathbb{Q}_p$ is finite, $\Delta=G_{K(\mu_\infty)}$, π is the full geometric profinite fundamental group, and $q:\pi\twoheadrightarrow\pi^{(p')}$ has kernel N_p. All actions are the genuine continuous actions induced by the given section s. For a fixed $\bar{h}\in(\pi^{(p')})^{s(\Delta)}$, choose $h\in\pi$ above it and put $d_\sigma=h^{-1}s(\sigma)hs(\sigma)^{-1}$. Then $\bar{h}$ has a fixed lift in π exactly when the image of d is a coboundary in N_p/U for every member U of a cofinal directed family of open normal subgroups of N_p stable under $s(\Delta)$. Such a family exists. The solution set is exactly the inverse limit of the finite solution sets; no additional derived-limit obstruction occurs.*

Proof. Write $\sigma(n)=s(\sigma)ns(\sigma)^{-1}$. The fixed-image hypothesis puts d_σ in N_p, and multiplication gives $d_{\sigma\tau}=d_\sigma\sigma(d_\tau)$. Moreover hn is fixed precisely when

$$d_\sigma=n\sigma(n)^{-1}\quad\text{for every }\sigma. \tag{F.20}$$

These identities prove the cocycle assertion and its exact relation to the desired lift, including the side on which the correcting element is multiplied.

We justify the cofinal stable family even though N_p need not be finitely generated. For an open normal subgroup $U\subset N_p$, joint continuity of the action and compactness of N_p give an open neighborhood of the identity in $s(\Delta)$ acting trivially modulo U on all of N_p. Concretely, for each point choose an action neighborhood preserving its coset modulo U; finitely many point neighborhoods cover N_p, and intersect their action neighborhoods and their inverses. A smaller open subgroup then stabilizes U. The orbit of U is finite, and its intersection is an open normal subgroup contained in U stable under all of $s(\Delta)$. Taking these intersections and finite common refinements gives the claimed family.

For each such U let

$$X_U=\{nU\in N_p/U : d_\sigma=n\sigma(n)^{-1}\bmod U\text{ for every }\sigma\}.$$

This is a finite set and is exactly the finite-level solution set. Its inverse image Y_U in the compact group N_p is closed. Under the stated finite-level condition it is nonempty. A common refinement of finitely many U supplies a nonempty Y contained in all corresponding Y_U, so these closed sets have the finite-intersection property. Compactness gives $n\in\bigcap_U Y_U$. Since $\bigcap_U U=1$, equation (F.20) holds in N_p for every σ. Conversely an actual solution gives every finite solution. The identification $N_p=\varprojlim_U N_p/U$ identifies the total solution set with $\varprojlim_U X_U$. This proves the exact criterion, not merely a necessary collection of finite tests. □

Statement F.22 (Proved). A full free pro-prime Dehn-twist fixed group. *Let F_ℓ be the free pro-ℓ group on x, y, z and let $0 \neq m \in \mathbb{Z}_\ell$. The continuous automorphism*

$$\alpha_m(x) = y^m x y^{-m}, \qquad \alpha_m(y) = y, \qquad \alpha_m(z) = z$$

has fixed subgroup exactly the closed free factor $\langle y, z\rangle$. This is a statement about the entire free pro-ℓ group.

Proof. We use the continuous Magnus map

$$F_\ell \longrightarrow 1 + (X, Y, Z)\mathbb{F}_\ell\langle\langle X, Y, Z\rangle\rangle, \qquad x \mapsto 1 + X,\ y \mapsto 1 + Y,\ z \mapsto 1 + Z. \tag{F.22.1}$$

Here the ring has its degree-adic topology, and the unit group on the right is an inverse limit of finite ℓ-groups. The map is injective. To check this rather than assume a linearization, the inverse ring map sends X, Y, Z to $x - 1, y - 1, z - 1$ in $\mathbb{F}_\ell[[F_\ell]]$. These substitutions converge: the augmentation ideal in the group algebra of every finite ℓ-group is nilpotent. The latter fact follows by induction using a central subgroup of order ℓ, whose generator c satisfies $(c - 1)^\ell = 0$. The two continuous ring homomorphisms are inverse, by their values on the generators. Finally F_ℓ embeds in the unit group of its completed group algebra, because its image in each finite group algebra retains the distinct group basis elements. This proves the assertion about (F.22.1).

Put $u(Y) = (1 + Y)^m$ in $\mathbb{F}_\ell[[Y]]$. It is nonconstant: writing $m = \ell^v m_0$ with m_0 a unit, its first nonconstant term has degree ℓ^v and nonzero coefficient. The induced substitution on the ring is $X \mapsto u(Y)Xu(Y)^{-1}$, with Y, Z fixed. For a word $w = w_1 \cdots w_k$ in the two letters X, Z, collect all coefficients of

$$Y^{a_0} w_1 Y^{a_1} \cdots w_k Y^{a_k}$$

as the single commutative series $f_w(Y_0, \ldots, Y_k)$. Every noncommutative series has this unique coefficient decomposition. The substitution multiplies f_w by

$$\prod_{i:\, w_i = X} u(Y_{i-1})u(Y_i)^{-1}. \tag{F.22.2}$$

For every w containing X this product differs from one: at the left endpoint of the first consecutive block of X's the corresponding variable has exponent $+1$, with no inverse factor to cancel it. The commutative power-series ring over $\mathbb{F}_\ell$ is an integral domain. A fixed series therefore has $f_w = 0$ whenever w contains X.

For a fixed group element, its Magnus series consequently uses only Y, Z. The continuous retraction $F_\ell \to \langle y, z\rangle$ given by $x \mapsto 1$ sets $X = 0$ in its Magnus series and leaves that series unchanged. Injectivity of (F.22.1) makes the element equal to its retraction. Conversely α_m fixes the stated free factor pointwise. □

Statement F.23 (Proved). All prime-to-five pro-prime fixed quotients in the three local examples. *Let $\Delta = G_{\mathbb{Q}_5(\mu_\infty)}$ and let $\pi^{(\ell)}$ denote the maximal geometric pro-ℓ quotient. For each $\ell \neq 5$, the first two examples of Task F.4 satisfy*

$$\left(\pi^{(\ell)}\right)^{s(\Delta)} = \pi^{(\ell)} = F_\ell(2).$$

For the third example, with the point 2 on $\mathbb{P}^1 \setminus \{0, 1, 5, \infty\}$, choose peripheral generators a, b, c at $0, 5, 1$ based through the outer component of its semistable model and put $d = ab$. Then

$$\left(\pi^{(\ell)}\right)^{s(\Delta)} = \langle d, c\rangle \simeq F_\ell(2). \tag{F.23}$$

The natural quotient $\pi \twoheadrightarrow \pi^{(\ell)}$ is continuous and $s(G_{\mathbb{Q}_5})$-equivariant. It sends the full fixed group into the displayed subgroup; surjectivity of that restriction is not asserted.

Proof. The smooth marked integral model in the first two examples, together with its integral point or unit tangential base point, gives based prime-to-five specialization. Inertia acts trivially on this quotient. Since all finite unramified extensions of $\mathbb{Q}_5$ are generated by prime-to-five roots of unity, Δ is contained in inertia, proving these two assertions.

For the third example blow up the collision $x = 5 = 0$ and use the node chart $XV = 5$, with $X = x$ and $V = 5/x$. The outer component has node 0, cusps $1, \infty$, and the smooth reduced base point 2. The bubble has node 0 and cusps $1, \infty$ in coordinate V. The logarithmic specialization and amalgamation maps of Stix, Theorem 3.4.6 and equation (6.2.5) [Sti02], give the genuine based prime-to-five comparison, by the same root-fiber-functor construction as F.9. The outer component subgroup is $\langle d, c\rangle$. A path based at 2 inside this smooth outer component has trivial inertia action in each prime-to-five quotient by based good reduction. The gluing along $XV = 5$ therefore makes tame inertia act by

$$a \longmapsto d^r a d^{-r}, \qquad b \longmapsto d^r b d^{-r}, \qquad c \longmapsto c, \quad r \in \prod_{q \neq 5} \mathbb{Z}_q(1).$$

Wild inertia acts trivially in this geometric prime-to-five comparison. These formulas follow also directly from the node Kummer covers: a turn in the base annulus conjugates the bubble loops by the common edge loop and fixes the based outer loops. In the free basis $(x, y, z) = (b, d, c)$ the action is α_r.

The image of Δ in tame inertia is $4\prod_{q\neq 5}\mathbb{Z}_q(1)$. Indeed the tame part of the five-adic cyclotomic character is the character of order four; on wild inertia its image is $1 + 5\mathbb{Z}_5$, as is seen on the extensions generated by fifth-power roots of unity. A tame element in the kernel of the character of order four can thus be corrected by a wild element to have cyclotomic character one. All other cyclotomic characters are already trivial on inertia. Projecting to the ℓ factor gives $4\mathbb{Z}_\ell(1)$, including $\ell = 2$. It follows that the fixed group is the fixed group of α_4; continuity gives equality for the closure of its integer powers. Statement F.22 applies since $4 \neq 0$ in $\mathbb{Z}_\ell$. This proves (F.23) and its precise relation to the full fixed group. □

Statement F.24 (Proved). The pro-five fixed groups for the two good-marked-reduction examples. *For $U = \mathbb{P}^1_{\mathbb{Q}_5} \setminus \{0, 1, \infty\}$, let $q_5 : \pi \twoheadrightarrow \pi^{(5)}$ be the continuous maximal geometric pro-five quotient. With the genuine action of the point 2, respectively of an integral unit tangent at 0, we have*

$$\left(\pi^{(5)}\right)^{s_2(\Delta)} = 1, \qquad \left(\pi^{(5)}\right)^{s_{0,u}(\Delta)} = I_0^{(5)}. \tag{F.24.1}$$

Consequently $q_5(\pi^{s_2(\Delta)}) = 1$ and $q_5(\pi^{s_{0,u}(\Delta)}) = I_0^{(5)}$. The latter equality of images follows because the full cusp inertia is itself fixed. It does not identify either full profinite fixed group.

Proof. Put $F = \mathbb{Q}(\zeta_5)$ and $S = \{v : v \mid 10\}$, and let G be the maximal pro-five quotient of $G_{F,S}$. We first prove that the image of inertia at five in G is all of G. The discriminant of F is 5^3. Its Minkowski bound is $15\sqrt{5}/(2\pi^2) < 2$, so its ideal class group is trivial. The prime two is inert, since 2 has order four modulo five, and $(\mathcal{O}_F/(2))^\times$ has order 15. The subgroup $\mu_5 \subset \mathcal{O}_F^\times$ maps injectively to this group and exhausts its five-primary part. The ray-class exact sequence therefore gives $\mathrm{Cl}_{(2)}(F)[5] = 0$. These uses of the ray-class correspondence are precisely Milne, Chapter V, Theorem 3.6 and Example 3.9 [Mil20c].

A cyclic extension of degree five unramified outside S and unramified at five is tamely ramified at two and has conductor dividing (2). The preceding computation excludes it. If the inertia image were a proper closed subgroup of G, its image in some finite five-group quotient would be proper and would lie in a subgroup of index five. Pulling back that subgroup would give exactly such a cyclic extension. This proves the asserted surjectivity, without assuming that the global group is free.

Let $F_S^{(5)}$ be the extension with group G. Its intersection with $\mathbb{Q}(\mu_\infty)$ is $F(\mu_{5^\infty})$. To verify the equality, the intersection is Galois and abelian over $\mathbb{Q}$, is unramified outside $\{2,5\}$, and is pro-five over F. The cyclotomic conductor description over $\mathbb{Q}$ (Milne, Chapter V, Example 3.10) shows that its part from a power of two has degree a power of two; hence it is already in F. The remaining part is the five-power cyclotomic tower, which is contained in $F_S^{(5)}$. Thus the image of the global full-cyclotomic kernel in G is $\ker \chi_5$. The local full-cyclotomic kernel has the same image: inertia at five surjects onto G, and the cyclotomic character of an inertia lift of an element of $\ker \chi_5$ is one at five and at every other prime. All these maps are restrictions of Galois automorphisms for a fixed embedding $\overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}}_5$.

For the point 2, the based pro-five action is unramified outside $\{2,5\}$: the marked curve and its point extend over $\mathbb{Z}[1/10]$, and based prime-to-residue-characteristic specialization applies at every remaining prime. Over F its action on the Frattini quotient is trivial, since that action is χ_5 modulo five. The kernel of $\mathrm{Aut}(F_5(2)) \to \mathrm{GL}_2(\mathbb{F}_5)$ is pro-five: the successive characteristic exponent-five central quotients give finite filtrations with elementary abelian five-group kernels. Hence the based action factors through G. We have proved equality of the local and global cyclotomic-kernel images for this actual based action.

We check that the global point section at 2 is not cuspidal even in the pro-five group. Its two abelian Kummer coordinates are $(\kappa(2), \kappa(-1)) = (\kappa(2), 0)$. The maps $x, 1-x : U \to \mathbb{G}_m$, evaluated on the leading terms of the unit tangents, show that the cusp-one packet has first coordinate zero and the cusp-infinity packet has equal coordinates. Since $v_{(2)}(2) = 1$, neither contains this class. To exclude the cusp-zero packet, consider over F the successive covers

$$y^5 = 1 - x, \qquad z_j^5 = \frac{y - \zeta_5^j}{-1 - \zeta_5^j} \quad (j = 1, 2). \tag{F.24.2}$$

They are finite etale covers of the successive punctured curves, with geometric degrees powers of five. Their composite is geometrically connected: the divisors of the two displayed Kummer functions have independent classes modulo five at $y = \zeta_5$ and $y = \zeta_5^2$. The point lifts to $(y, z_1, z_2) = (-1, 1, 1)$. Every cusp above zero first has $y = \zeta_5^i$. Choose $j \in \{1, 2\}$ distinct from i. At the prime $(1 - \zeta_5)$ the numerator $\zeta_5^i - \zeta_5^j$ has valuation one, while $-1 - \zeta_5^j$ is a unit. This value is not a fifth power in F, so no cusp above this point in (F.24.2) is F-rational. An F-section contained in the lifted cusp decomposition group would force such a rational cusp in every cover to which the section lifts. Thus the point section is noncuspidal.

The proof of B.4 applies word for word to this free pro-five fundamental group over the number field F: its cover comparison uses finite five-groups, its boundary sequence uses $\mathbb{Z}_5(1)$, and Ribet's torsion finiteness applies to the compactified cover Jacobians. It gives a trivial global cyclotomic-kernel centralizer for the just verified noncuspidal point section, and cusp inertia for the unit tangent at zero. Equality of the local and global images proves the first formula in (F.24.1) and the second for tangent one.

Finally the pro-five completion of $\mathbb{Z}_5^\times$ is free of rank one and is generated by the class of 2: $v_5(2^4 - 1) = 1$, so $\log(2)$ generates $5\mathbb{Z}_5$. An integral unit tangent u therefore has $\kappa_5(u) = a\kappa_5(2)$ locally for a unique $a \in \mathbb{Z}_5$. The global cusp section obtained by twisting the unit tangent by $a\kappa_5(2)$ is unramified outside $\{2,5\}$ and has the same local pro-five section class. It factors through the same relative pro-five action considered above, and B.4 gives its global centralizer $I_0^{(5)}$. The local/global image equality proves the claim for every integral unit tangent. Conjugating a representative by cusp inertia leaves the asserted inertia subgroup unchanged. The stated maps from the full fixed groups now follow directly by equivariance. The exact finite certificates used here—the cyclotomic discriminant, residue-field order, ray-class five-part, Kummer-function norms, and the lifted point—are evaluated in `code/bc_local_checks.py`, with its output included. □

Statement F.25 (Proved). The pro-five fixed group for the colliding-cusp example. *For $U = \mathbb{P}^1_{\mathbb{Q}_5} \setminus \{0, 1, 5, \infty\}$, with the section at 2 and $\Delta = G_{\mathbb{Q}_5(\mu_\infty)}$, we have*

$$\left(\pi^{(5)}\right)^{s_2(\Delta)} = 1. \tag{F.25.1}$$

Consequently the full profinite fixed group maps to one under the maximal pro-five quotient. Together with F.23 this evaluates every individual pro-prime fixed quotient in all three examples, with the explicit quotient maps described there.

Proof. Filling the cusp 5 is an actual based morphism

$$U \hookrightarrow U_0 = \mathbb{P}^1 \setminus \{0, 1, \infty\}.$$

It induces a continuous, surjective, equivariant map $F_5(3) \to F_5(2)$ killing the normal closure K of the inertia at the new puncture. By F.24 the target has trivial fixed group for the section at 2. It suffices to prove $K^\Delta = 1$.

We give the integral kernel comparison used for that assertion. Choose a geometric free basis which expresses $F_5(3) = F_5(2) *_5 \langle b \rangle$, where b is inertia at 5 and the quotient kills b. The profinite set X of lifts of the point 5 in the universal pro-five cover of U_0, based at 2, is a torsor under $F_5(2)$. After choosing one lift, identify it with $F_5(2)$. The kernel K is the free pro-five group on this profinite set, with continuous generators $x \mapsto xbx^{-1}$. This free group is universal for continuous maps from that profinite set to pro-five groups. The assertion follows by two explicit inverse homomorphisms: let the free group on X be acted on by $F_5(2)$ through left translation of X. Its semidirect product with $F_5(2)$ receives the free product by sending b to the generator at 1 and fixing $F_5(2)$. Conversely, $x \mapsto xbx^{-1}$ gives the homomorphism back, equivariant for that left-translation action. Both composites fix the indicated generators. Thus the kernel and all its generator labels are the actual ones for the filling morphism.

In particular

$$K^{\mathrm{ab}} \simeq \mathbb{Z}_5[[X]](1), \tag{F.25.2}$$

where the completed module is the module of $\mathbb{Z}_5$-valued finitely additive measures on the clopen subsets of X. Galois transports each inertia generator to a conjugate of the generator at the transported lifted cusp, raised to its cyclotomic power. Thus (F.25.2) is equivariant, and on Δ the twist is one. For every $r \geq 1$ there is also an injective equivariant degree-r Magnus homomorphism

$$\gamma_r K / \gamma_{r+1} K \hookrightarrow \mathbb{Z}_5[[X^r]]. \tag{F.25.3}$$

To verify this integral comparison for the infinitely generated kernel, use the finite partitions $X \to X_n = F_5(2)/M_n$, with M_n characteristic and open. Every continuous map from X to a finite five-group factors through one such partition, so the free group on X is the inverse limit of the free pro-five groups on the finite sets X_n. Its closed lower central groups and their quotients are the corresponding inverse limits: membership is detected in finite five-group quotients, and compatible cosets lift by compactness. For finite X_n, the Hall commutator basis and its leading Magnus monomials give the injective map into $\mathbb{Z}_5[X_n^r]$. Taking their inverse limit gives (F.25.3). Naturality of the degree-r term makes this map equivariant: conjugating a generator changes only terms of degree at least two, and the action on the degree-r graded piece is induced from its action on the degree-one module.

The action of Δ on X has pro-five image. Indeed its action on $F_5(2)$ is trivial on the Frattini quotient, and that automorphism kernel is pro-five, as proved in F.24. After a choice of torsor origin, the affine action on X is a homomorphism into the pro-five semidirect product of $F_5(2)$ with this automorphism kernel.

There are no finite orbits of this action. The regular function $1 - x$ on U_0 sends its path torsor from 2 to 5 to the division-root torsor of

$$\frac{1-5}{1-2} = 4.$$

On Δ its translation character is $\kappa_5(4) = 2\kappa_5(2)$. The latter character has a unit value. For a direct proof, the polynomial $(T+2)^5 - 2$ is Eisenstein at five, so $\mathbb{Q}_5(2^{1/5})$ has degree five. It is not Galois: a Galois extension containing one fifth root of 2 contains all of them and therefore ζ_5, whose degree is four. It cannot be a subfield of the abelian extension $\mathbb{Q}_5(\mu_\infty)$. Some element of Δ consequently moves that fifth root, which says that $\kappa_5(2)$ has nonzero reduction modulo five on Δ. Since it is a continuous homomorphism there, its image is all of $\mathbb{Z}_5$. Every orbit on X, and on X^r by projection to the first coordinate, therefore maps onto an infinite translation orbit.

We finish with the required measure argument. Let a pro-five group act continuously on a profinite set Z with no finite orbits. An invariant integral $\mathbb{Z}_5$-measure on Z is zero. For a clopen subset C, its stabilizer is open, by uniform continuity on the compact set. All orbits of this stabilizer on C are infinite. For each n, choose a finite invariant partition of C on which every orbit has size at least 5^n. Such a partition exists: each point has arbitrarily many distinct translates separated by a finite clopen partition; continuity gives an invariant refinement, and a finite common refinement works for all points by compactness. The finite orbit sizes are powers of five. Invariance makes the measure constant on the atoms in each orbit, so its value on C is divisible by 5^n. Since this holds for every n, that value is zero. This proves the claim.

Apply it to $Z = X^r$ in (F.25.3). Every graded fixed subgroup is zero. A nontrivial element of K has a nonzero image in some lower-central graded quotient, because it survives in a finite five-group and finite five-groups are nilpotent. A fixed element would give a nonzero graded invariant, which is impossible. Thus $K^\Delta = 1$, proving (F.25.1) and the stated consequence for the full profinite quotient map. □

Statement F.26 (Proved). A mixed-prime quotient and every one of its nilpotent fixed groups. *For the first example, with the section at 2, consider the actual finite etale cover*

$$\begin{aligned} E^\circ \longrightarrow U, \qquad E:\ Y^2 = X^3 - 2X, \quad t = X^2/2, \\ D_E = \{O, (0,0), (\sqrt{2},0), (-\sqrt{2},0)\}, \qquad P = (2,2). \end{aligned} \tag{F.26.1}$$

Here $E^\circ = E \setminus D_E$, and P maps to 2. The cover is cyclic of degree four over $\mathbb{Q}_5$: in the coordinate $Z = Y/2$ its equation is $Z^4 = t(t-1)^2/2$. Choose $i \in \mathbb{Q}_5$ with $i^2 = -1$ and $i \equiv 2 \bmod 5$; a deck generator is $\delta(X,Y) = (-X, iY)$. Let $H \subset \pi$ be the index-four subgroup determined by this based cover, put $H_5 = H^{(5)}$, and define

$$J = \pi / \ker(H \longrightarrow H_5), \qquad J_c = J/\gamma_{c+1}H_5 \quad (c \geq 1). \tag{F.26.2}$$

These are continuous, surjective, $s_2(G_{\mathbb{Q}_5})$-equivariant quotients of the full geometric profinite group. Their fixed groups are

$$J^{s_2(\Delta)} = 1, \qquad J_1^{s_2(\Delta)} = H_5^{\mathrm{ab}} \simeq \mathbb{Z}_5^5, \qquad J_c^{s_2(\Delta)} = \gamma_c H_5/\gamma_{c+1}H_5 \quad (c \geq 2). \tag{F.26.3}$$

In particular the fixed group at class two is $\mathbb{Z}_5^{10}$, and every nonzero element of it fails to have a fixed lift at class three. The transition maps between consecutive fixed groups in (F.26.3) are zero. The image of the full fixed group in J is therefore trivial; in particular its image under the cyclic quotient whose peripheral vector is $(1,2) \bmod 4$ is trivial.

Proof. We give the arithmetic and Lie comparisons explicitly. The four points above 2 are $P, \delta P, \delta^2 P, \delta^3 P$, all rational over $\mathbb{Q}_5$. Thus the based arithmetic cover is the one with section at P, and its subgroup H is stable under the genuine action in the statement. The kernel of the maximal pro-five quotient of H is characteristic in H, normal in π, and stable under this arithmetic action. This proves all the quotient assertions in (F.26.2). The geometric group H_5 is free pro-five of rank $2 \cdot 1 + 4 - 1 = 5$.

Write $V_E = H_5^{\mathrm{ab}}$ and $W_E = \mathbb{Z}_5(1)[D_E]/\mathbb{Z}_5(1)\sum_d d$. The functions

$$X, \quad X - \sqrt{2}, \quad X + \sqrt{2}$$

have respective divisors $2d - 2O$. Their induced homomorphism on V_E restricts to multiplication by two on the boundary basis e_0, e_+, e_-. Since two is a unit in $\mathbb{Z}_5$, it gives an integral Galois-equivariant splitting

$$V_E = W_E \oplus T_5 E. \tag{F.26.4}$$

The homomorphisms are equivariant over $\mathbb{Q}_5(\sqrt{2})$; their permutation under its Galois involution proves the stated equivariance over $\mathbb{Q}_5$. They are also equivariant for δ, which fixes e_0 and interchanges e_+ and e_-. Thus $W_E = W_+ \oplus W_-$, with ranks two and one, $W_- = \mathbb{Z}_5(e_+ - e_-)(1)$. Its arithmetic characters are respectively χ_5 and $\chi_5\varepsilon$, where ε is the unramified quadratic character. Indeed $\sqrt{2}$ generates the unramified quadratic extension.

The two eigenspaces of $[i]$ split T_5E integrally, since $2i$ is a unit. One has an unramified character ψ; the other has character $\chi_5\psi^{-1}$ by the Weil pairing. Here is a direct check of the unramified assertion. The endomorphism $[2 + i]$ has degree five and unit differential $2 + i \bmod 5$ on the good integral model. Its reduction is separable, so its kernel and the kernels of its iterates are finite etale over $\mathbb{Z}_5$. Their inverse limit is one of the two CM Tate lines. Counting points gives $\#E(\mathbb{F}_5) = 10$, so the Frobenius polynomial is $T^2 + 4T + 5$. The unit root is $-2 - i$ for the chosen embedding of i; thus $\psi(\mathrm{Frob}_5) = -2 - i$. This root is not a root of unity: either complex embedding has absolute value $\sqrt{5}$. In particular Δ acts trivially on all of (F.26.4).

Choose the unit tangent at O, with parameter X/Y, as an auxiliary reference section on E°. The deck automorphism has derivative i there. The compatible fifth-power roots of i are all i, and are rational over $\mathbb{Q}_5$. The resulting tangent comparison gives a genuine order-four automorphism of the based pro-five group, commuting with its arithmetic action. Let

$$j_\sigma : V_E \longrightarrow \bigwedge^2 V_E \quad (\sigma \in \Delta) \tag{F.26.5}$$

be the first Johnson homomorphism: for a lift $\widetilde{v}$ of v, it is the class of $\sigma(\widetilde{v})\widetilde{v}^{-1}$ modulo $\gamma_3 H_5$. It is independent of that lift because Δ acts trivially on V_E and on $\bigwedge^2 V_E$. It is additive in σ and is equivariant under conjugation by $G_{\mathbb{Q}_5}$. The latter assertion follows directly by conjugating an automorphism which is the identity on the abelianization; any central part of the conjugating automorphism cancels. It also commutes with $[i]$.

Each $d - O$, for $d \in D_E$, is two-torsion. Its pro-five Kummer path is trivial: multiplication by 5^n fixes this two-torsion point, providing a rational compatible division sequence. The formula for the action on peripheral inertia therefore gives $j_\sigma(W_E) \subset \bigwedge^2 W_E$. The CM eigenvalues on T_5E are $i, -i$, whereas those on $\bigwedge^2 W_E$ and on $\bigwedge^2 T_5E$ are $1, -1$. Commutation with $[i]$ consequently gives $j_\sigma(T_5E) \subset W_E \otimes T_5E$. Together these facts put (F.26.5), after tensoring with $\mathbb{Q}_5$, in a direct sum of one-dimensional arithmetic characters drawn from

$$\chi_5, \quad \varepsilon\psi^2, \quad \chi_5^2\varepsilon\psi^{-2}. \tag{F.26.6}$$

For example the off-diagonal map from one CM line to W_- times the other has character $\chi_5\varepsilon$ times the ratio of those two lines. The diagonal maps use W_+ and have character χ_5. This proves (F.26.6) without assuming a diagonal action on the nonabelian group.

Choose a geometric path from the auxiliary tangent to P, and write $b_\sigma \in V_E$ for its abelian cocycle. The change from the auxiliary action to the action at P changes (F.26.5) by

$$j_\sigma^P = j_\sigma + \operatorname{ad} b_\sigma; \tag{F.26.7}$$

reversing the initial path changes the two displayed signs together and has no effect on the argument. The functions defining (F.26.4), with leading coefficient one at the auxiliary tangent, evaluate at P as $2, 2-\sqrt{2}, 2+\sqrt{2}$. Hence the component of b in W_- is

$$\frac{1}{4}\kappa_5\left(\frac{2-\sqrt{2}}{2+\sqrt{2}}\right)(e_+ - e_-) = \frac{1}{2}\kappa_5(\sqrt{2}-1)(e_+ - e_-). \tag{F.26.8}$$

This is a cocycle for the character $\theta = \chi_5\varepsilon$. Its restriction to Δ is nonzero. To prove that assertion, put $L = \mathbb{Q}_5(\sqrt{2})$. The unit $\sqrt{2}-1$ is not a root of unity, as its real algebraic embedding has absolute value different from one. The logarithm on a sufficiently small open unit subgroup shows that its rational local Kummer class is nonzero. For an abelian profinite group Γ and a nontrivial one-dimensional $\mathbb{Q}_5$-character η, $H^1_{\mathrm{cts}}(\Gamma, \mathbb{Q}_5(\eta)) = 0$: the cocycle identity for commuting g, h, with $\eta(g) \neq 1$, expresses every cocycle as the coboundary of $c(g)/(\eta(g)-1)$. Apply this to the abelian quotient G_L/Δ and χ_5. Inflation–restriction proves that the Kummer class stays nonzero on Δ.

The T_5E component of $b|_\Delta$ is nonzero as well. Indeed $\#E(\mathbb{F}_3) = 4$ and $\#E(\mathbb{F}_5) = 10$. Injectivity of prime-to-residue-characteristic torsion specialization bounds $E(\mathbb{Q})_{\mathrm{tors}}$ by a group of order dividing two. Since $Y(P) = 2$, P is not torsion. The formal logarithm on an open subgroup of $E(\mathbb{Q}_5)$ then shows that its rational five-adic Kummer class is nonzero. The local filtration and specialization assertions are those of Milne, Chapter II, Theorem 4.1, Corollary 4.2 and Aside 4.4 [Mil21]; alternatively the kernel of reduction is a pro-residue-characteristic formal group, which proves the torsion injectivity directly. Both characters ψ and $\chi_5\psi^{-1}$ are nontrivial. Applying the preceding abelian-quotient argument to these two lines proves that this class also stays nonzero on Δ.

We now separate these homomorphisms on Δ exactly. The character θ in (F.26.8) occurs neither in (F.26.6), nor in W_+, nor in T_5E. The powers of χ_5 distinguish it from the two off-diagonal characters in (F.26.6) on inertia; it differs from χ_5 by the nontrivial ε. It differs from the unramified line ψ, and equality with $\chi_5\psi^{-1}$ would require $\psi = \varepsilon$, contradicting the displayed Frobenius value. The joint homomorphism

$$\Delta \longrightarrow \operatorname{Hom}(V_E, \bigwedge^2 V_E) \otimes \mathbb{Q}_5 \ \oplus \ V_E \otimes \mathbb{Q}_5, \qquad \sigma \longmapsto (j_\sigma, b_\sigma)$$

has compact additive image, a $\mathbb{Z}_5$-lattice in its rational span. That span is stable under $G_{\mathbb{Q}_5}$ and is a sum of the characters just listed. The projection onto its nonzero θ component is a polynomial in finitely many of these commuting Galois operators: for each other character choose an operator distinguishing it from θ, and multiply the corresponding linear projections. Clearing the finitely many five-adic denominators puts a nonzero projected vector back in the compact image lattice. There is consequently an actual $\sigma_0 \in \Delta$ such that

$$j_{\sigma_0} = 0, \qquad b_{\sigma_0} = w, \qquad 0 \neq w \in W_- \otimes \mathbb{Q}_5. \tag{F.26.9}$$

The vector in (F.26.9) is integral because it is in the actual image. This step constructs the required group element from the additive image, rather than choosing unrelated values for different cocycles.

For completeness, the closed lower-central graded algebra of the finite-rank free pro-five group embeds by its leading Magnus terms in the tensor algebra on V_E. An automorphism acting trivially on V_E acts to the next degree by the derivation induced by its first Johnson map. This follows by expanding a commutator and using the Leibniz rule in the tensor algebra; the Hall basis proves the assertion integrally in each degree. By (F.26.7)–(F.26.9), that derivation for σ_0 is $\operatorname{ad} w$. In the free Lie algebra over $\mathbb{Q}_5$, the centralizer of a nonzero degree-one element w consists of its linear span. One direct proof extends w to a vector-space basis: a homogeneous tensor polynomial commuting with the letter w contains only that letter, by comparing first and last letters of its words. Its only possible homogeneous form is a multiple of w^r; for $r > 1$ this is not a Lie element, as its tensor coproduct has nonzero intermediate binomial coefficients.

It follows that the leading degree of a fixed element below the last central degree of a nilpotent quotient must be one, with leading vector proportional to w. Choose $\sigma_1 \in \Delta$ with $(b_{\sigma_1})_T \neq 0$. For this vector w, the $W_E \otimes T_5E$ component of $j^P_{\sigma_1}(w)$ is exactly $[(b_{\sigma_1})_T, w]$, which is nonzero. Indeed $j_{\sigma_1}(w)$ and the bracket with $(b_{\sigma_1})_W$ lie in $\bigwedge^2 W_E$. Thus no such degree-one vector is fixed either. Every fixed element of $H_5/\gamma_{c+1}H_5$, for $c \geq 2$, is therefore in its last graded subgroup. Conversely that subgroup is fixed because the action on every associated graded is induced from the trivial action on V_E. This proves the asserted fixed groups inside H_5. Their ranks in degrees two and three are respectively 10 and 40.

Finally no element outside H_5 can be fixed in J_c for any $c \geq 1$. The image of the inertia at O in J has an order-four subgroup mapping isomorphically onto the deck group; its generator δ_0 is fixed by the auxiliary cuspidal action on Δ. Its conjugation on V_E is the deck action $[i]$. For an element with deck value δ_0^k, the fixed equation modulo $[H_5, H_5]$ for the action at P requires

$$(1 - \delta^k)b_\sigma = 0 \quad \text{for all } \sigma \in \Delta.$$

For $k = 1, 2, 3$, the operator $1 - \delta^k$ is invertible on $T_5E \otimes \mathbb{Q}_5$, and the proper component of b is nonzero. This excludes all three deck values. Together with the preceding paragraph this proves the formulas for J_c. The intersection of the lower central subgroups of H_5 is one, since each nontrivial element survives in a finite five-group, and every such finite group is nilpotent. Taking the inverse limit proves $J^\Delta = 1$.

The peripheral vector follows directly from the valuations of $t(t-1)^2/2$ at 0 and 1. All finite curve counts, toric identities, character comparisons and Hall-dimension checks used here are verified in `code/bc_mixed_cm_checks.py`. □

Statement F.27 (Proved). An explicit finite obstruction to prime-to-five fixed lifts. *For the point* 2 *on the tripod over* $\mathbb{Q}_5$*, the image of the full fixed group in the geometric abelian quotient* $(\mathbb{Z}/4)^2$ *is zero. In particular, although every element of* $\pi^{(5')}$ *is fixed, any such element with nonzero abelian image modulo four has no fixed lift in* π*. This obstruction is already detected by two actual finite quotients, each isomorphic to*

$$(\mathbb{F}_5)^5 \rtimes C_4 \quad \textit{of order } 12500. \tag{F.27.1}$$

The cyclic generator acts on the normal vector group by fixing e_0*, interchanging* e_+, e_-*, and multiplying the two CM coordinates by* 2 *and* 3*.*

Proof. The first finite quotient is $\overline{J} = J/(H_5^5[H_5, H_5])$, with the actual maps of F.26. We verify at this precise finite level that no nontrivial deck value is fixed. Put $\beta = [2 - i]$. Its kernel is the formal CM subgroup of $E[5]$, and it is isomorphic over $\mathbb{Q}_5$ to μ_5: the other CM line has unramified character $\psi \equiv 1 \bmod 5$, and the Weil pairing identifies the two lines as

dual with cyclotomic twist. The point $P = (2, 2)$ is not in $\beta E(\mathbb{Q}_5)$. Here is a finite certificate retaining the isogeny map. On the integral affine curve it is given by

$$\begin{aligned} X(\beta Q) &= \frac{X(3+4i)(X^2-2-4i)^2}{(5X^2-2+4i)^2}, \\ Y(\beta Q) &= \frac{Y(2+11i)(X^2-2-4i)(X^4+(4+16i)X^2+4)}{(5X^2-2+4i)^3}. \end{aligned} \tag{F.27.2}$$

These identities follow by applying the addition law to $2Q - [i]Q$, and are checked symbolically by the program. The denominators are units at five. Modulo five the map is $-\mathrm{Frob}_5$, so any preimage of P must reduce to $-P = (2, 3)$. Modulo twenty-five, with $i = 7$, all five lifts of that residue point map under (F.27.2) to $(7, 2)$, rather than $(2, 2)$. The finite enumeration is included in the program and its output.

Consequently the isogeny Kummer class $\delta_\beta(P) \in H^1(\mathbb{Q}_5, \mu_5)$ is nonzero. It remains nonzero on Δ. Indeed for any $u \in \mathbb{Q}_5^\times$ which is not a fifth power, adjoining a fifth root gives degree five: after removing a fifth power and a fourth root of unity, either its valuation is nonzero modulo five, and the Newton polygon proves this degree, or it is $(1+5)^b$ with $b \in \mathbb{Z}_5^\times$, in which case $(1+T)^5 - u$ is Eisenstein at five. That degree-five extension is not Galois, since a Galois extension containing one root would contain ζ_5, of degree four. It cannot lie in the abelian extension $\mathbb{Q}_5(\mu_\infty)$. This proves the asserted restriction nonvanishing. The map $[2+i] : E[5] \to E[\beta]$ sends the usual multiplication-by-five Kummer class of P to $\delta_\beta(P)$. Thus the proper component of $b|_\Delta$ in F.26 has nonzero reduction modulo five.

For $k = 1, 2, 3$, $1 - \delta^k$ is invertible on the proper two-dimensional vector group over $\mathbb{F}_5$: its eigenvalues are $1 - 2^k$ and $1 - 3^k$. The last fixed equation in F.26 therefore excludes nontrivial deck values already in $\overline{J}$. Conversely the normal vector group is fixed because Δ acts trivially on V_E. Thus $\overline{J}^\Delta = (\mathbb{F}_5)^5$. The order-four subgroup supplied by cusp inertia splits the extension and has exactly the displayed matrix, proving (F.27.1).

Write (a, b) for the two abelian coordinates modulo four in the original tripod. The cyclic cover equation forces $a + 2b = 0$ for every fixed element. The Mobius automorphism $m(t) = t/(t-1)$ is defined over $\mathbb{Q}_5$ and fixes the base point 2. It therefore gives a genuine based automorphism of π, commuting with the arithmetic action. It fixes the cusp 0 and interchanges $1, \infty$. Its action on abelian inertia is $e_0 \mapsto e_0$, $e_1 \mapsto -e_0 - e_1$. Pulling the first finite quotient back along this actual automorphism gives the second quotient in (F.27.1), with cyclic character $(1, 1)$ mod 4. Thus a fixed element also satisfies $a + b = 0$. Subtracting the two equations gives $b = 0$ and then $a = 0$.

The abelian quotient modulo four factors through $\pi^{(5')}$, by its universal property. Good marked reduction with the integral point 2 makes $\pi^{(5')}$ fixed, as in F.23. Therefore a prime-to-five element with nonzero image in $(\mathbb{Z}/4)^2$ is an explicit fixed quotient element having no fixed lift. All coordinate identities, the five residue lifts and the invertible deck matrices are checked in `code/bc_mixed_cm_checks.py`. □

Statement F.28 (Proved). All abelian coordinates of the three full local fixed groups. *In the three examples of F.21, the images of the full fixed groups in the full geometric abelianizations are, respectively,*

$$0, \qquad \widehat{\mathbb{Z}}\, e_0, \qquad 0. \tag{F.28.1}$$

In particular the two point-section fixed groups lie in the closed commutator subgroup of their respective full geometric fundamental groups.

For every integer $m \geq 2$ prime to five, the first example has two explicit finite quotient tests excluding all nonzero abelian coordinates modulo m. They have the form

$$J_{m,n_m} = (\mathbb{Z}/5^{n_m})^{m+1} \rtimes C_m, \tag{F.28.2}$$

and their fixed groups are exactly their displayed normal vector groups. The integer n_m is given by the finite local valuation calculation (F.28.5) below. Analogous finite quotients detect the second and third images in (F.28.1). The cases $m = 2, 3, 4, 6$ in the first example all have $n_m = 1$, giving orders $250, 1875, 12500, 468750$.

Proof. Write $\Delta = G_{\mathbb{Q}_5(\mu_\infty)}$. First consider the tripod with base point 2 and the actual cyclic cover

$$Y_m = \mathbb{P}^1_z \setminus (\{0, \infty\} \cup \mu_m), \qquad t = z^m. \tag{F.28.3}$$

Choose $\alpha^m = 2$. Its fiber consists of $\alpha\zeta_m^j$. All these points and all punctures are defined over a finite unramified extension L of $\mathbb{Q}_5$. Indeed a residue extension of degree d with $4m \mid 5^d - 1$ contains every root of $X^m - 2$ and every mth root of unity; their simple roots lift by Hensel's lemma. Thus $L \subset \mathbb{Q}_5(\mu_\infty)$ and Δ fixes these fibers.

The kernel of $\pi \to C_m$ is the actual full geometric fundamental group of Y_m. Pass from it to its maximal pro-five abelian quotient V_m, a free $\mathbb{Z}_5$-module of rank $m + 1$. The kernel of this passage is characteristic in the open subgroup and normal in π. Thus it gives a genuine quotient J_m of π fitting into $1 \to V_m \to J_m \to C_m \to 1$. Its cyclic action fixes the boundary vector at zero and cyclically permutes the m vectors at μ_m. The image of inertia at the ramified cusp zero is procyclic with a unique prime-to-five factor of order m. This supplies an actual splitting, so $J_m \simeq V_m \rtimes C_m$. Since the punctures are L-rational and $\chi_5|_\Delta = 1$, the action of Δ on V_m is trivial.

Suppose that a fixed element has deck value $j \neq 0$. Lift its based loop through (F.28.3). It gives a Δ-fixed point of the abelian path torsor from α to $\alpha\zeta_m^j$. This description also holds after passing to V_m: path lifting commutes with the Galois action and with the abelian quotient. The function $z - 1$, a unit on Y_m, maps this torsor to the Kummer torsor of

$$u_j = \frac{\zeta_m^j \alpha - 1}{\alpha - 1}. \tag{F.28.4}$$

It would therefore make $\kappa_L(u_j)|_\Delta = 0$.

This Kummer class is nonzero even with rational five-adic coefficients. Choose the complex embedding with $\alpha = 2^{1/m} > 1$. Then $|\zeta_m^j \alpha - 1|^2 = (\alpha - 1)^2 + 2\alpha(1 - \cos(2\pi j/m))$ is strictly larger than $(\alpha - 1)^2$. Consequently u_j is not a root of unity. For a finite local field, the five-adic logarithm on principal units has precisely torsion kernel; its valuation component and logarithm identify the rational Kummer completion of $L^\times$. Thus its Kummer class is nonzero. Restriction to Δ is injective on $H^1(G_L, \mathbb{Q}_5(1))$. To see this directly, the quotient $\Gamma = G_L/\Delta$ is abelian, and its cyclotomic character is nontrivial. For a continuous cocycle c on Γ, commutativity gives

$$(\chi(g) - 1)c(h) = (\chi(h) - 1)c(g).$$

Choose g with $\chi(g) \neq 1$ and divide in $\mathbb{Q}_5$. This expresses c as a coboundary. Inflation–restriction, obtained from the usual cochain maps, proves the asserted injectivity. This contradicts the fixed path and excludes every nonzero deck value.

There is an explicit finite precision for this exclusion. Choose L unramified of degree d as above and put $q = 5^d$. All u_j are units: neither α nor any $\zeta_m^j \alpha$ reduces to one, since its mth power reduces to 2. Set

$$e_j = v_L(u_j^{q-1} - 1) - 1, \qquad n_m = 1 + \max_{1 \le j < m} e_j, \qquad v_L(5) = 1. \tag{F.28.5}$$

These are finite nonnegative integers. The decomposition $\mathcal{O}_L^\times = \mu_{q-1} \times (1 + 5\mathcal{O}_L)$ and the logarithm isomorphism on the second factor prove that e_j is exactly the largest e for which

$u_j \in L^{\times 5^e}$. Write $u_j = v_j^{5^{e_j}}$ with v_j not a fifth power. Its nonzero Kummer class modulo five remains nonzero on Δ. After removing a Teichmuller factor, $v_j \in 1+5\mathcal{O}_L$ has $v_L(v_j-1)=1$, so $(1+T)^5 - v_j$ is Eisenstein. The degree-five extension generated by a root is not Galois: a normal field containing one root would contain μ_5, of degree four over the unramified field L. It therefore cannot be contained in the abelian extension $\mathbb{Q}_5(\mu_\infty)/L$. Hence $\kappa_L(v_j)|_\Delta$ is nonzero modulo five, and $\kappa_L(u_j)|_\Delta$ is nonzero modulo 5^{n_m}. This proves the finite exclusion in $J_m/5^{n_m}V_m$. Its normal vector group is fixed, proving the exact fixed group in (F.28.2). Formula (F.28.5) involves only algebraic valuations and gives a terminating finite calculation.

Let (a,b) be full abelian coordinates at $0,1$. The covering (F.28.3) forces $a = 0 \bmod m$ for every fixed element. The based automorphism $t \mapsto t/(t-1)$ fixes 2 and commutes with the arithmetic action. Its abelian map is $(a,b) \mapsto (a-b,-b)$. Pulling back (F.28.3) along this actual automorphism forces $a - b = 0 \bmod m$. Therefore both coordinates vanish modulo every m prime to five. F.24 makes their five-primary coordinates zero as well. This proves the first image in (F.28.1).

For the unit tangent at zero, use instead the actual cover $t = 1 - z^m$. The induced tangents above zero lie at the points ζ_m^j, with $dz/dt = -1/(m\zeta_m^{j(m-1)})$. Choose the starting tangent at $z = 1$, so its parameter $z - 1$ has value $-1/m$. A fixed element with deck value $j \neq 0$ would give a fixed abelian path to the induced tangent at ζ_m^j. The same function $z-1$ sends these two fibers to Kummer parameters $-1/m$ and $\zeta_m^j - 1$, respectively. Their ratio is $m(1-\zeta_m^j)$. Its complex absolute value is at least $2m\sin(\pi/m) \geq 4$, so it is not a root of unity. It is a five-adic unit because m is prime to five and ζ_m^j has nontrivial prime-to-five order. The preceding restriction and finite-precision arguments apply verbatim, with these ratios. This excludes its e_1 coordinate modulo every m prime to five. F.24 gives the same exclusion for its five-primary part. Conversely the full cusp-zero inertia is fixed by Δ, because the full cyclotomic character is one there. Its abelian image is the whole primitive line $\widehat{\mathbb{Z}}e_0$. This proves the second equality.

For the four-puncture curve, filling the cusp 5 gives the tripod point 2 by an actual based morphism. Thus a fixed element has zero e_0, e_1 coordinates. The other based map

$$t \longmapsto (5-t)/4 : \mathbb{P}^1 \setminus \{0,1,5,\infty\} \longrightarrow \mathbb{P}^1 \setminus \{0,1,\infty\}$$

fills the original cusp zero and sends the point 2 to $3/4$, the cusp 5 to zero, and the cusp 1 to one. At the tripod point $3/4$, the covers $t = z^m$ and $t = 1 - z^m$ have starting roots $(3/4)^{1/m}$ and $(1/4)^{1/m}$. Both are positive real numbers different from one. The strict absolute-value inequality used for (F.28.4) holds equally for a positive root less than one. Their reductions have mth powers 2 and 4 modulo five, so the same unramified and unit arguments apply. These two covers force both prime-to-five abelian coordinates of a fixed loop at $3/4$ to be zero. Pullback gives the remaining e_5 exclusion. F.25 excludes every five-primary coordinate on the four-puncture curve. This proves its zero abelian image.

The code `bc_cyclic_fixed.py` works in exact unramified rings modulo powers of five. It computes (F.28.5) for $m = 2,3,4,6$, obtaining $e_j = 0$ for every j in each case. It verifies the finite orders stated above. □

Statement F.29 (Proved). A genus-zero mixed quotient with no fixed element. *In the cyclic fourth-root cover $t = z^4$ of the tripod with base point 2, let H_5 be the maximal geometric pro-five group of $\mathbb{P}^1_z \setminus \{0,\infty,\pm1,\pm i\}$, and let J be the resulting quotient of the full base fundamental group. Thus $1 \to H_5 \to J \to C_4 \to 1$ and H_5 is free pro-five of rank five. For the genuine point-section action,*

$$J^\Delta = 1, \qquad (J/\gamma_{c+1}H_5)^\Delta = \gamma_c H_5/\gamma_{c+1}H_5 \quad (c \geq 1). \tag{F.29.1}$$

Every transition between consecutive displayed fixed groups is zero.

Proof. Take $i \in \mathbb{Q}_5$, $i \equiv 2 \bmod 5$, and $\alpha^4 = 2$. The extension $L = \mathbb{Q}_5(\alpha)$ is unramified of degree four: $X^4 - 2$ is irreducible and separable over $\mathbb{F}_5$. Its Frobenius sends α to $i\alpha$. The auxiliary section is the unit tangent at $z = 0$. The order-four factor of its inertia supplies a based deck element δ, fixed by the auxiliary Galois action. Equivalently, the change of tangent under $z \mapsto iz$ has the compatible rational fifth-power roots $i^{1/5^n} = i$, so its based comparison commutes with $G_{\mathbb{Q}_5}$.

Write

$$V = H_5^{\mathrm{ab}} = \mathbb{Z}_5 e_0 \oplus \bigoplus_{j=0}^{3} \mathbb{Z}_5 f_j$$

with a Tate twist on every summand; the last four basis vectors are the inertias at i^j. All punctures are $\mathbb{Q}_5$-rational, so the auxiliary arithmetic action on V is the scalar χ_5. For $\sigma \in \Delta$ its first Johnson map is $j_\sigma : V \to \bigwedge^2 V$. Composition in the first two nilpotent degrees proves that j is additive on Δ and that

$$j_{g\sigma g^{-1}} = \chi_5(g) j_\sigma. \tag{F.29.2}$$

Let b be the cocycle comparing the point 2 with the auxiliary tangent in J. Its deck component records $g\alpha/\alpha$. On Δ this component is zero, so b_σ has an abelian value in V, again an additive map of Δ. We retain the precise Frobenius transport. Choose φ lifting arithmetic Frobenius with $\chi_5(\varphi) = 1$; this is possible because the fifth cyclotomic extension is totally ramified, whereas L is unramified. On G_L, the functional dual to f_j sends b to $\kappa_L(\alpha - i^j)$: the unit $z - i^j$ has tangent value $-i^j$, whose pro-five Kummer class is zero. For $\sigma \in \Delta$,

$$\begin{aligned} b_j(\varphi\sigma\varphi^{-1}) &= \kappa_{\varphi^{-1}(\alpha - i^j)}(\sigma) \\ &= \kappa_{\alpha - i^{j+1}}(\sigma) = b_{j+1}(\sigma). \end{aligned} \tag{F.29.3}$$

The discarded factor is $-i$, with its compatible rational fifth-power roots. In the right-deck convention this is the actual permutation $\delta f_j = f_{j-1}$, while $\delta e_0 = e_0$. The same conclusion follows directly from the nonabelian cocycle identity: modulo $[H_5, H_5]$ the auxiliary Δ-action is trivial, so $b_{g\sigma g^{-1}} = b_g \rho(g)(b_\sigma) b_g^{-1}$. Its linear action is $\chi_5(g)\delta^{j(g)}$, where $g\alpha = i^{j(g)}\alpha$. This proves the required covariance with the actual point-base deck translation.

We now exhibit two nonzero eigenspace components of $b|_\Delta$. In the unramified integer ring $\mathbb{Z}_5[\alpha]$, the following are the coefficients of $\log(\alpha - i^j)/5$ modulo five, in the basis $1, \alpha, \alpha^2, \alpha^3$:

j	1	α	α^2	α^3
0	0	0	2	1
1	0	0	3	2
2	0	0	2	4
3	0	0	3	3

. (F.29.4)

Here the logarithm kills the Teichmuller factor. This is an exact finite calculation: with $i = 7 \bmod 25$, compute $((\alpha - i^j)^{624} - 1)/(5 \cdot 624)$ in $(\mathbb{Z}/25)[\alpha]/(\alpha^4 - 2)$ and reduce modulo five. The program performs these operations and verifies every row.

For the permutation in (F.29.3), the coefficient at f_0 of the λ-eigenprojection on the four puncture coordinates is $\frac{1}{4}\sum_j \lambda^{-j} b_j$. For $\lambda = i$ and $\lambda = -1$, (F.29.4) gives respectively α^3 and $2\alpha^2$ modulo five after dividing the logarithm by five. Thus both rational Kummer classes are nonzero. The restriction injection proved in F.28 makes their restrictions to Δ nonzero as well.

The joint additive image of Δ under $\sigma \mapsto (j_\sigma, b_\sigma)$ is a compact $\mathbb{Z}_5$-submodule. Conjugation by φ preserves it, acts trivially on j by (F.29.2), and by the order-four permutation δ on b.

The integral operators

$$E_\lambda = \frac{1}{4}\sum_{a=0}^{3} \lambda^{-a}\varphi^a \quad (\lambda = i, -1) \tag{F.29.5}$$

therefore preserve this actual image. They annihilate its j-coordinate and retain the indicated nonzero b-component. Compactness makes the image closed under every $\mathbb{Z}_5$ scalar in (F.29.5). Consequently there are actual $\sigma_1, \sigma_2 \in \Delta$ with

$$j_{\sigma_1} = j_{\sigma_2} = 0, \qquad b_{\sigma_1} = w_i \neq 0, \quad b_{\sigma_2} = w_{-1} \neq 0, \tag{F.29.6}$$

where the two vectors lie in distinct deck eigenspaces. They are linearly independent.

Changing from the auxiliary tangent to the point section changes the first Johnson map to $j_\sigma + \operatorname{ad} b_\sigma$. Thus (F.29.6) supplies two independent pure-inner first Johnson directions. The leading Magnus term of a fixed element below the final degree of a nilpotent quotient must commute in the free Lie algebra with both w_i and w_{-1}. As proved in F.26, the centralizer of a nonzero degree-one vector in a free Lie algebra over $\mathbb{Q}_5$ is its one-dimensional span. The two spans intersect in zero. The integral Hall graded modules embed in their rationalizations, so this excludes every such nonzero leading term. Conversely the final central graded subgroup is fixed because Δ acts trivially on V and hence on every free Lie graded quotient.

F.28 with $m = 4, n_m = 1$ excludes all nonzero deck values already in the finite abelian-normal quotient. It excludes them in every quotient in (F.29.1) as well. This proves the displayed finite-depth groups. Their images at the preceding depth are zero. Finally $\bigcap_c \gamma_c H_5 = 1$: every nontrivial element of a free pro-five group survives in a finite five-group, and finite five-groups are nilpotent. Taking the inverse limit proves $J^\Delta = 1$. □

Statement F.30 (Proved). A torsion test in every actual finite cover. *Put $K_\infty = \mathbb{Q}_5(\mu_\infty)$ and $\Delta = G_{K_\infty}$. For every finite extension $L/\mathbb{Q}_5$ put $\Delta_L = G_{LK_\infty}$ and $\Gamma_L = \operatorname{Gal}(LK_\infty/L)$. For every semiabelian variety A/L, restriction is injective:*

$$H^1(G_L, V_5A) \longrightarrow H^1(\Delta_L, V_5A), \qquad V_5A = T_5A \otimes_{\mathbb{Z}_5} \mathbb{Q}_5. \tag{F.30.1}$$

Consequently, for every smooth curve Y/L, its smooth compactification C, reduced boundary $D = C \setminus Y$, and $P, Q \in Y(L)$, a Δ_L-fixed geometric étale path from P to Q forces

$$[Q - P] \in J_D(L)_{\mathrm{tors}}. \tag{F.30.2}$$

The same conclusion holds for a fixed path in the maximal abelian pro-five path torsor. When $L \subset K_\infty$ one has $\Delta_L = \Delta$. Here J_D is the generalized Jacobian with reduced modulus D; when D is empty it is the ordinary Jacobian. For nonempty D, condition (F.30.2) has the exact divisor formulation

$$\exists n \geq 1,\ f \in \overline{L}(C)^\times : \quad \operatorname{div}(f) = n(Q - P), \qquad f|_D = 1. \tag{F.30.3}$$

In particular, let $U/\mathbb{Q}_5$ have a rational interior base point x, let $\pi = \pi_1(U_{\overline{\mathbb{Q}}_5}, x)$, and use the point section to define the action of Δ. For every Δ-stable open subgroup $H \subset \pi$, the corresponding based cover over K_∞ is an actual finite étale cover. If $h \in \pi^\Delta$, its lift beginning at the chosen point P ends at a K_∞-point Q above x, and this pair satisfies (F.30.2)–(F.30.3) after descent to a finite $L \subset K_\infty$.

Proof. We first prove the representation-theoretic assertion used in (F.30.1). Let Γ be an abelian profinite group and M a finite-dimensional continuous $\mathbb{Q}_5$-representation with $M^\Gamma = 0$. Let $R \subset \operatorname{End}_{\mathbb{Q}_5}(M)$ be the unital algebra generated by Γ. It is a finite-dimensional

commutative algebra acting faithfully on M. The ideal $I = (\gamma - 1 : \gamma \in \Gamma)$ is all of R. Indeed, if it lay in a maximal ideal $\mathfrak{m}$, the corresponding local Artinian factor $R_{\mathfrak{m}}$ would act on a nonzero summand $M_{\mathfrak{m}}$. Its last nonzero radical power $\mathfrak{m}^a M_{\mathfrak{m}}$ would contain a nonzero vector killed by $\mathfrak{m}$, hence by every $\gamma - 1$. This contradicts $M^\Gamma = 0$. Thus there are finitely many $A_i \in R$ and $\gamma_i \in \Gamma$ with

$$\sum_i A_i(\gamma_i - 1) = 1.$$

For a continuous one-cocycle $c : \Gamma \to M$, commutativity gives $(\gamma_i - 1)c(\gamma) = (\gamma - 1)c(\gamma_i)$. Consequently

$$c(\gamma) = (\gamma - 1)m, \qquad m = \sum_i A_i c(\gamma_i). \tag{F.30.4}$$

This is an actual continuous coboundary formula; in particular $H^1(\Gamma, M) = 0$.

We record explicitly the local point-group input. For a semiabelian variety A over the finite local field L, a sufficiently small formal neighborhood of the identity is an open subgroup $U_A \simeq \mathbb{Z}_5^d$. The torus part has a finitely generated valuation quotient, and the abelian-variety part is compact. The local analytic subgroup assertion for the latter is Mattuck's Theorem 7, printed p. 114 [Mat55]. These statements give a finitely generated quotient $A(L)/U_A$. For completeness, the image in the abelian-variety part is open, since the differential of the smooth quotient morphism is surjective; it is therefore compact. Finitely many local analytic sections over this compact image, together with generators for the torus valuation lattice, give the asserted finite generation modulo U_A. Passing to the preimage of a torsion-free subgroup of finite index in $A(L)/U_A$ and lifting its free basis gives a finite-index subgroup of the form

$$\mathbb{Z}^r \oplus \mathbb{Z}_5^d. \tag{F.30.5}$$

In particular $A(L)_{\mathrm{tors}}$ is finite and $(V_5 A)^{G_L} = 0$. It also follows directly from the finite Kummer sequences and (F.30.5) that the map

$$A(L) \longrightarrow H^1(G_L, V_5 A) \tag{F.30.6}$$

has exactly the torsion subgroup as its kernel. One way to see the last assertion without a completion identification is to start with a zero rational Kummer class. A five-power multiple of the point is then divisible by every 5^n in $A(L)$. Multiplication of this point and all its division witnesses by the finite index of (F.30.5) places them in that subgroup. The intersection of the subgroups 5^n times (F.30.5) is zero.

Now put $V = V_5 A$ and $M = V^{\Delta_L}$. The quotient $\Gamma = \Gamma_L = G_L/\Delta_L$ is abelian: restriction identifies it with the closed subgroup $\mathrm{Gal}(K_\infty/(L \cap K_\infty))$ of the abelian cyclotomic Galois group. and $M^\Gamma = V^{G_L} = 0$. Let a cocycle on G_L represent an element in the kernel of (F.30.1). Subtracting an actual coboundary makes its restriction to Δ_L zero. The cocycle identity then shows both that its values lie in M and that it is constant on cosets of Δ_L. It descends continuously to Γ. Formula (F.30.4) makes it a coboundary and proves (F.30.1).

The path-to-Kummer map in the second assertion is the one induced by the actual Abel morphism

$$\iota_P : Y \longrightarrow J_D, \qquad Q' \longmapsto [Q' - P]. \tag{F.30.7}$$

More explicitly, pull back the finite étale isogenies $[5^n] : J_D \to J_D$ along ι_P. A path from P to Q carries the compatible zero points above $\iota_P(P) = 0$ to compatible division points of $[Q - P]$. Its displacement under σ is the cocycle $\sigma Q_n - Q_n$ in $J_D[5^n]$. Thus a Δ_L-fixed path makes the restriction of the rational Kummer class of $[Q - P]$ zero. The construction factors through the abelian pro-five path torsor because these deck groups are abelian five-groups. Apply (F.30.1) and then (F.30.6) to obtain (F.30.2). The quotient of degree-zero divisors

disjoint from D by divisors of functions equal to one on D is the actual generalized-Jacobian group. This gives (F.30.3); see also [Cre19, §2.1, equation (2.2), and §2.3, Lemma 2.5].

Finally the arithmetic subgroup for the indicated cover is $H \rtimes s_x(\Delta) \subset \pi \rtimes s_x(\Delta)$. It has finite index, and its chosen fibre point is Δ-fixed. The usual lifting map for finite étale covers identifies paths in Y with base paths having the prescribed endpoints. The lift of h is fixed because both h and its starting point are fixed; its endpoint is therefore a K_∞-point. The cover, the two points, and their morphism to U are given by finitely many coefficients in K_∞, so they descend together to some finite $L \subset K_\infty$. The previous construction then proves the stated test in this actual cover. Further extension to any finite L'/L preserves the fixed-path conclusion, since $\Delta_{L'} = G_{L'K_\infty} \subset \Delta_L$. Thus this passage also permits arbitrary finite extensions used to change the reduction of the cover. □

Statement F.31 (Proved). The full abelian fixed images for all genus-zero curves. *Let $L/\mathbb{Q}_5$ be finite, $K_\infty = \mathbb{Q}_5(\mu_\infty)$, and $\Delta_L = G_{LK_\infty}$. Let Y/L be a smooth genus-zero hyperbolic curve and $P \in Y(L)$. For the action defined by this actual point section,*

$$\pi_1(Y_{\overline{L}}, P)^{\Delta_L} \longrightarrow \pi_1(Y_{\overline{L}}, P)^{\mathrm{ab}} \quad \textit{has image zero.} \tag{F.31.1}$$

There is no restriction on the reduction of Y, on the position of P, or on the prime divisors of the finite abelian quotient being tested. For an L-rational nonzero tangential base point $\vec{v}$ at an L-rational cusp c, the corresponding full fixed group has abelian image exactly $\widehat{\mathbb{Z}}(1)e_c$.

The same conclusion applies at every actual genus-zero finite étale neighborhood of P whose chosen fibre point is fixed by Δ_L, after a finite extension of L. Thus a full fixed element has zero abelian image in every such neighborhood in which it lifts to a loop.

Proof. Write C for the compactification and $D = C \setminus Y$. The rational point P identifies C with a projective line. Extend L finitely to split D; a fixed element remains fixed by the smaller group $\Delta_{L'}$. We retain the letter L for this extension during the construction.

Choose a cusp $d_\infty \in D$, and an actual L-isomorphism $z : C \longrightarrow \mathbb{P}^1_L$ sending d_∞ to infinity. Put $p = z(P)$. For each $d \in D \setminus \{d_\infty\}$ set

$$u_d = \frac{z - z(d)}{p - z(d)} : Y \longrightarrow \mathbb{G}_m. \tag{F.31.2}$$

These maps give the usual full abelian coordinates: the inertia at d maps to 1 in $\widehat{\mathbb{Z}}(1)$, the inertia at d_∞ maps to -1, and every other cusp inertia maps to zero. The sole relation among the cusp generators is their sum, so the product of these induced maps is an isomorphism $\pi_1(Y_{\overline{L}}, P)^{\mathrm{ab}} \simeq \widehat{\mathbb{Z}}(1)^{|D|-1}$.

Fix $n \geq 3$ and form the actual connected cyclic cover

$$Y_{d,n} \longrightarrow Y, \qquad w^n = u_d, \qquad z = z(d) + (p - z(d))w^n. \tag{F.31.3}$$

Its compactification is the projective w-line. The removed points are $0, \infty$ and the n-th roots of $(z(e) - z(d))/(p - z(d))$ for $e \in D \setminus \{d, d_\infty\}$. This lists precisely all inverse images of D, and proves that (F.31.3) is finite étale on the indicated open curves. Its fibre above P is μ_n. Every one of these points is fixed by Δ_L.

Let h be a full fixed element whose image under u_d modulo n is $j \neq 0$. Lift its actual path to (F.31.3), beginning at $w = 1$. The endpoint is $w = \zeta_n^j$, for a compatible choice of generator ζ_n. The path is fixed by Δ_L. Choose another cusp $e \in D \setminus \{d, d_\infty\}$, which exists by hyperbolicity. Put

$$r = \frac{z(e) - z(d)}{p - z(d)} \neq 0, 1, \qquad \beta^n = r.$$

After a finite extension L'/L containing β and μ_n, each

$$v_a = w - \beta\zeta_n^a \quad (0 \le a < n) \tag{F.31.4}$$

is an actual unit on $Y_{d,n}$. It is nonzero at both endpoints, since $r \neq 1$. The induced map of paths to $\mathbb{G}_m$ sends the fixed path to the Kummer path of

$$R_a = \frac{\zeta_n^j - \beta\zeta_n^a}{1 - \beta\zeta_n^a} \in L'^{\times}. \tag{F.31.5}$$

Formula (F.30.1), now with $\Delta_{L'} = G_{L'K_\infty} \subset \Delta_L$, therefore forces every R_a to be a root of unity.

Here is an exact contradiction. Embed the finitely generated characteristic-zero field containing these finitely many coefficients and roots into $\mathbb{C}$. Set

$$D_a = |\zeta_n^j - \beta\zeta_n^a|^2 - |1 - \beta\zeta_n^a|^2.$$

The root-of-unity sums $\sum_a \zeta_n^{2a} = \sum_a \zeta_n^{-2a} = 0$, valid for $n \ge 3$, give

$$\sum_{a=0}^{n-1} D_a^2 = 2n\,|\zeta_n^j - 1|^2\,|\beta|^2 > 0. \tag{F.31.6}$$

Indeed $D_a = -(\zeta_n^j - 1)\overline{\beta}\,\zeta_n^{-a} - (\zeta_n^{-j} - 1)\beta\,\zeta_n^a$, and squaring proves the displayed identity. Thus some $|R_a| \neq 1$, so that R_a cannot be a root of unity. This excludes every $j \neq 0$.

We have proved that every coordinate in (F.31.2) vanishes modulo every $n \ge 3$. Every positive modulus divides one of these n, so all full abelian coordinates vanish. The integers n used in (F.31.3) may be divisible by five: the maps are étale in characteristic zero, and the proof uses the actual subgroup $\Delta_{L'}$, which fixes all roots of unity. This proves (F.31.1).

For the tangential assertion, choose $d_\infty \neq c$ and use the same actual coordinate z, putting $p = z(c)$. For each cusp $d \neq c, d_\infty$ use (F.31.3) with this value of p and $n \ge 5$. The cover is unramified at c. If the original tangent has velocity $\dot{z}$, its lift at $w = 1$ has velocity $\dot{z}/(n(p - z(d)))$; the deck translate at $w = \zeta_n^j$ has velocity $\zeta_n^j\dot{z}/(n(p - z(d)))$. These are the actual derivatives of (F.31.3), so both lifted tangents are fixed by $\Delta_{L'}$. The other points ζ_n^a above c are punctures. For $a \neq 0, j$, the unit $w - \zeta_n^a$ has nonzero value at both tangential endpoints. Its induced Kummer ratio is therefore $(\zeta_n^j - \zeta_n^a)/(1 - \zeta_n^a)$; no tangent velocity enters this nonvanishing value. Put $A = |\zeta_n^j - 1|^2$. Formula (F.31.6) with $\beta = 1$, after subtracting the terms at $a = 0, j$, gives

$$\sum_{\substack{0 \le a < n \\ a \neq 0, j}} \left(|\zeta_n^j - \zeta_n^a|^2 - |1 - \zeta_n^a|^2\right)^2 = 2A(n - A) > 0, \tag{F.31.7}$$

because $0 < A \le 4 < n$. The identical Kummer argument excludes all $j \neq 0$. This kills every abelian coordinate except the one corresponding to c. Conversely the full inertia $I_c \simeq \widehat{\mathbb{Z}}(1)$ at this actual tangent is fixed by Δ_L: its action on compatible roots of the local parameter is the cyclotomic character, which is one on Δ_L. Its image is precisely the primitive cusp line.

The obstruction can also be made finite in each instance. For an R_a selected by (F.31.6), its restricted integral Kummer homomorphism $\Delta_{L'} \to \mathbb{Z}_5(1)$ is nonzero, since its rational class is nonzero. It remains nonzero modulo some 5^s. Pulling back the cover $T^{5^s} = v_a$ along (F.31.4) supplies a finite étale cover on which the indicated endpoints cannot be joined by a fixed path. Taking its normal closure, and then the compositum of the finitely many coefficient-field conjugates, gives an actual finite geometric quotient stable under the original

arithmetic action. The conjugate family is finite because all the coefficients, points, roots, and covering equations used here descend to the finite extension L'/L.

Finally apply the same construction to the given finite neighborhood, using its chosen fixed fibre point as P. Descent of the cover and point to a finite field and subsequent restriction of Δ_L are exactly the morphisms described at the end of F.30. The exact root-of-unity identities (F.31.6)–(F.31.7) and the cyclic-cover inverse maps are checked by `bc_genus_zero.py`. □

Statement F.32 (Proved). The full local fixed group at every point and tangent. *Let $L/\mathbb{Q}_5$ be finite, let $K_\infty = \mathbb{Q}_5(\mu_\infty)$, and put $\Delta_L = G_{LK_\infty} \subset G_L$. Let X/L be a smooth geometrically connected hyperbolic curve, and let Π be its full geometric profinite fundamental group. For an L-rational point $b \in X(L)$ and its actual point section s_b,*

$$C_\Pi(s_b(\Delta_L)) = 1. \tag{F.32.1}$$

For a nonzero L-rational tangent at an L-rational cusp c and its actual tangential section $s_{\vec{c}}$,

$$C_\Pi(s_{\vec{c}}(\Delta_L)) = I_c, \tag{F.32.2}$$

where I_c is the actual geometric inertia subgroup selected by that tangent. In the tripod example of Task F.4, $I_0 \simeq \widehat{\mathbb{Z}}(1)$, with this isomorphism induced by the compatible local covers $w^n = t$.

Proof. We use the resolution of nonsingularities theorem of Mochizuki–Tsujimura, in the author's revision of 30 March 2024. The exact inputs are Theorem 2.17 and Proposition 2.4(vii): with Σ the set of all primes, a section of a closed subgroup $H \subset G_L$ fixes at most one primitive valuation in their universal geometric covering, provided $H \cap I_L$ has a continuous quotient $\mathbb{Z}_\ell$ for some $\ell \neq 5$. The correspondence between valuations and center-chains, including its inverse, is Proposition 2.3(v)–(vi) [MT24, Propositions 2.3(v)–(vi), 2.4(vii); Theorem 2.17]. We verify the subgroup hypothesis and construct the valuation and its stabilizer explicitly.

First every finite unramified extension of L is obtained by adjoining suitable roots of unity of order prime to five. Indeed, such roots give all finite extensions of the finite residue field, and their lifts are the unique roots supplied by Hensel's lemma. Consequently $L^{\mathrm{ur}} \subset LK_\infty$ and

$$\Delta_L \subset I_L, \qquad I_L/\Delta_L \xrightarrow[\chi_5]{\sim} \chi_5(I_L) \subset \mathbb{Z}_5^\times \simeq C_4 \times \mathbb{Z}_5. \tag{F.32.3}$$

For the kernel assertion, inertia already fixes all prime-to-five roots of unity, so its full cyclotomic kernel is exactly its five-adic cyclotomic kernel. Thus (F.32.3) is induced by the actual cyclotomic homomorphism.

Choose compatible three-power roots of a uniformizer of L. The resulting tame homomorphism $t_3 : I_L \to \mathbb{Z}_3(1)$ is onto: the successive extensions over the maximal unramified field have degrees 3^n, as follows either from their valuations or from the Eisenstein polynomials for those roots. Let $B = t_3(\Delta_L)$, a closed subgroup. The quotient $\mathbb{Z}_3(1)/B$ is a continuous quotient of I_L/Δ_L. Every finite quotient of the latter has order divisible only by two and five, whereas every finite quotient of $\mathbb{Z}_3(1)/B$ is a three-group. Therefore $B = \mathbb{Z}_3(1)$. This verifies the uniqueness hypothesis with $H = \Delta_L$ and $\ell = 3$. It uses the at-most-one part of Proposition 2.4(vii), whose hypothesis imposes no nontrivial cyclotomic image on H.

Here is the promised geometric comparison. Write M for the union of the function fields of the connected finite geometric covers of $X_{\overline{L}}$ inside its chosen universal cover. Then

$$\Pi = \mathrm{Gal}(M/\overline{L}(X)).$$

More precisely, transport the chosen based fibre functor to a geometric generic point by an étale path. The transported arithmetic fundamental group is $\mathrm{Gal}(M/L(X))$; its projection to G_L is restriction to the constant field, with kernel the displayed group Π. Transport the section and the basepoint lifts by the same path. Thus every action below is an action in this actual extension. The based cover construction gives a compatible lift of b to every finite cover. At a tangent, the chosen compatible Puiseux branches give instead a compatible cusp in the normalizations of the smooth compactification. In either case write $\widetilde{b}$ for this compatible collection of centers. Let

$$B_{\widetilde{b}} = \bigcup_{Y} \mathcal{O}_{\overline{Y},b_Y} \subset M, \qquad \epsilon_{\widetilde{b}} : B_{\widetilde{b}} \longrightarrow \overline{L} \tag{F.32.4}$$

be the directed union of the local discrete valuation rings at the centers and its residue map. In (F.32.4), $\overline{Y}$ is the smooth compactification, so the formula applies to cusps as well as to interior points. The union is a valuation ring: every element of $M^\times$ lies in a finite subcover, where it or its inverse belongs to the indicated local discrete valuation ring. Its residue field is $\overline{L}$, because all the centers are geometric and the constant field already contains $\overline{L}$.

Compose this horizontal valuation with the five-adic valuation of its residue field. The resulting valuation ring is

$$R_{\widetilde{b}} = \epsilon_{\widetilde{b}}^{-1}(\mathcal{O}_{\overline{L}}) \subset B_{\widetilde{b}}. \tag{F.32.5}$$

Its intersection with L is $\mathcal{O}_L$. It is primitive in the precise sense of Mochizuki–Tsujimura Definition 2.2(i): its maximal ideal is the only prime ideal containing five. To prove this directly, put $J = \ker \epsilon_{\widetilde{b}}$. Since $5^{-n}J = J$ for every $n \geq 0$, any prime ideal of $R_{\widetilde{b}}$ containing five contains J. The quotient $R_{\widetilde{b}}/J$ is the rank-one valuation ring $\mathcal{O}_{\overline{L}}$, whose only prime ideal containing five is its maximal ideal. Taking its inverse image proves the assertion.

The morphism from this construction to the source theorem is the following actual map: send $R_{\widetilde{b}}$ to its center in every compactified semistable model of every finite subcover, and then to the vertex or edge containing that center. These entries form the center-chain of Definition 2.2(v). Proposition 2.3(v) identifies this map with a bijection from the valuation rings in question to the center-chain space; Proposition 2.3(vi) gives its inverse as the directed union of the local rings at the centers. Both maps commute with the arithmetic fundamental-group action, since formation of a center and pullback of a local ring commute with the corresponding field automorphisms. In particular equality of the primitive center-chains here is equality of the rings $R_{\widetilde{b}}$.

The actual section $s_b(G_L)$ preserves the compatible basepoint lifts, while the actual section $s_{\vec{c}}(G_L)$ preserves the compatible cusps selected by the tangent. Their action on the residue field in (F.32.4) is the ordinary Galois action on $\overline{L}$, which preserves $\mathcal{O}_{\overline{L}}$. Thus $R_{\widetilde{b}}$, and hence its primitive center-chain, is fixed by the restricted section $s(\Delta_L)$.

Now take $h \in C_\Pi(s(\Delta_L))$. For every $\sigma \in \Delta_L$,

$$s(\sigma)\big(hR_{\widetilde{b}}\big) = h\, s(\sigma) R_{\widetilde{b}} = hR_{\widetilde{b}}.$$

Thus $hR_{\widetilde{b}}$ is a second fixed primitive valuation. The uniqueness theorem gives $hR_{\widetilde{b}} = R_{\widetilde{b}}$.

We next recover the exact geometric stabilizer, without identifying different kinds of valuation by notation. There is an equality of subrings of M

$$R_{\widetilde{b}}[1/5] = B_{\widetilde{b}}. \tag{F.32.6}$$

Indeed five is a unit in the horizontal valuation ring. Conversely, for any $f \in B_{\widetilde{b}}$, a sufficiently large power $5^n f$ has integral residue and belongs to $R_{\widetilde{b}}$. It follows that an element of Π preserving $R_{\widetilde{b}}$ preserves $B_{\widetilde{b}}$, hence the center b_Y on every finite normalization $\overline{Y}$. Conversely,

an element preserving all these centers preserves $B_{\widetilde{b}}$ and its residue map, since it fixes $\overline{L}$ pointwise; it therefore preserves $R_{\widetilde{b}}$. We have proved the exact stabilizer equality

$$\mathrm{Stab}_{\Pi}(R_{\widetilde{b}}) = \mathrm{Stab}_{\Pi}(\widetilde{b}). \tag{F.32.7}$$

At an interior point, every finite geometric Galois cover is étale there. Its deck group acts freely on the fibre: the residue extension is trivial over $\overline{L}$ and the ramification index is one. Thus an element of (F.32.7) has identity image in every finite geometric Galois quotient, and is itself the identity. This proves (F.32.1).

At a cusp, (F.32.7) is the inverse limit of the inertia groups at the chosen compatible normalization points, which is exactly the image I_c of the punctured local disc. For completeness, each completed local extension has the form $\overline{L}((t^{1/e}))/\overline{L}((t))$: its residue field is algebraically closed, its residue characteristic for this horizontal valuation is zero, and an e-th root of the unit multiplying a local uniformizer exists by Hensel's lemma. Its automorphisms multiply $t^{1/e}$ by μ_e. These compatible homomorphisms identify the stabilizer with the actual local inertia image, including all transition maps. For the tripod at zero the global finite covers $w^n = t$ realize every such cyclic quotient, so the local map $\widehat{\mathbb{Z}}(1) \to \Pi$ is injective and has image I_0.

Finally the tangential section acts on these compatible local roots by the cyclotomic character. Since Δ_L fixes all roots of unity, it acts trivially on I_c. The inclusion $I_c \subset C_{\Pi}(s_{\vec{c}}(\Delta_L))$, together with (F.32.7) and uniqueness, proves (F.32.2). □

Statement F.21 (Proved). The three full local fixed groups and all fixed lifts. *For the three examples in Task F.4, with $\Delta = G_{\mathbb{Q}_5(\mu_\infty)}$ and the actual section action, the full profinite fixed groups are*

U	*section*	$\pi^{s(\Delta)}$
$\mathbb{P}^1 \setminus \{0,1,\infty\}$	2	1
$\mathbb{P}^1 \setminus \{0,1,\infty\}$	*integral unit tangent at* 0	$I_0 \simeq \widehat{\mathbb{Z}}(1)$
$\mathbb{P}^1 \setminus \{0,1,5,\infty\}$	2	1.

(F.21.1)

For $q : \pi \to \pi^{(5')}$, the images of these full fixed groups are, respectively,

$$1, \qquad I_0^{(5')}, \qquad 1. \tag{F.21.2}$$

Consequently the finite coboundary tests of F.20 all hold exactly for $\bar{h} = 1$ in the two point-based examples, and exactly for $\bar{h} \in I_0^{(5')}$ in the tangential example. The fixed lift is unique in each point-based example. Each tangential lift fibre is a coset of $\mathbb{Z}_5(1) \subset I_0$.

Proof. Formula (F.21.1) follows from F.32 applied to the two actual interior point sections and the actual tangent at zero. On the inertia selected by this tangent, the quotient q is the actual projection

$$\widehat{\mathbb{Z}}(1) = \prod_{\ell} \mathbb{Z}_\ell(1) \longrightarrow \prod_{\ell \neq 5} \mathbb{Z}_\ell(1).$$

The covers $w^n = t$ with $(n, 5) = 1$ verify injectivity of the displayed target into $\pi^{(5')}$ and identify it with $I_0^{(5')}$. Its kernel is the five-adic factor. This proves (F.21.2) and the stated lift fibres. Finally F.20 identifies the existence of each such fixed lift with all the finite coboundary tests, proving the exact evaluation of those tests as well.

The earlier F.22–F.31 computations retain their explicit quotient actions and finite obstruction witnesses. The universal-cover valuation argument in F.32 supplies the detection beyond those particular quotient families. □

A Coverage and remaining questions

The following table records task-level scope. A proved supporting statement does not certify an unresolved task.

Item	Statements	State
Lead 1	B.2; B.9	(a) Proved. The exclusion is false; an unramified full-section counterexample lies outside the three inherited cusp lines. The drafter's particular inherited toric argument is corrected.
Lead 2	A.5; A.17; B.7	(b) Refuted. A mixed-inertia class-two lift at 2 is explicit. Full and metabelian sections obey the proved cusp-line restriction.
Lead 3	A.6; A.18	(b) Refuted. An infinitely ramified noncuspidal class-two family is constructed with a complete convergence proof; its higher images are excluded.
Lead 4	B.4; B.5	(a) Proved. The full geometric pro-3 cyclotomic centralizer is cusp inertia for cuspidal sections and trivial for every other section.
Lead 5	A.1; A.2; A.13	(a) Proved. The first derived limit for the intrinsic rank-44 coefficient is explicitly nonzero.
A.1	A.1; A.2; A.7; A.13	(a) Proved. The finite-S complex, its integral Smith factors, localization, and the complete Milnor sequence are computed.
A.2	A.8; A.9; A.10	(a) Proved. All residue formulas retain the nine tangential origin cochains, the extra unramified generator, and the E36 central-coordinate sign.
A.3	A.7; A.8	(a) Proved. Integral local corrections must form a restricted family; every nonempty fiber is the computed rank-23 module with its finite torsion.
A.4	A.3; A.4	(a) Proved. An actual pair has a nonzero intrinsic obstruction with every local and finite-coefficient image zero.
A.5	A.11; A.12	(b) Refuted as a total evaluator. Total decision is impossible both for finite-coefficient queries and for the explicit finite-program input family with a known convergence modulus. The exact finite/certified-tail evaluator and all four requested regressions are executed; its input interface is stated explicitly.
A.6	A.14; A.15; A.16; A.17; A.18; A.19; A.20; A.21; A.22; A.23; A.24; A.25; A.26; A.27; A.28; A.29; A.30; A.31; A.32; A.33; A.34; A.35; A.36; A.37; A.38; A.39; A.40; A.41; A.42; A.43; A.44; B.12; B.13; B.14; B.15; B.16; B.17; B.18; B.19; B.20	(a) Proved by finite arithmetic tests. Every metabelian and marked image and fibre is determined by explicitly constructed finite arithmetic cocycle sets and their compatible kernel orbits, including infinite ramification support. Cofinal finite-rank towers, every coefficient layer and its derived limit, and both exact inherited comparison kernels are proved. An explicit integral target at 17 has the literal local cusp restriction and a class-three lift but no class-four or higher lift. The exact criterion has terminating finite stages and an infinite compatibility limit.
B.1	B.1; B.2; B.3; B.9	(a) Proved. All three packets in the abelian, cuspidally central, and inherited class-two coordinates are given with their origins.

Item	Statements	State
B.2	A.5; A.6; A.17; A.18	(b) Refuted both leads. Both requested class-two assertions have complete counterexamples and proved higher-quotient restrictions.
B.3	B.4; B.5	(a) Proved. The centralizer criterion and precisely the cuspidal members of the full unramified family are determined.
B.4	B.6; B.7; B.8; B.10; B.11; B.15; B.16; B.17; B.18; B.19; B.20; A.16; A.22; A.24; A.25; A.26; A.27; A.28; A.29; A.30; A.31; A.33; A.34; A.35; A.36; A.37; A.38; A.39; A.40; A.43	(a) Proved by finite arithmetic tests. The exact image for every one-prime target is given by a terminating finite field and cocycle construction at each precision, with all earlier ramification constraints. The inverse limit of the specified kernel orbits gives every fibre and is equivalent to lifting through all nilpotent quotients. At 17 an explicit integral cocycle has the literal local cusp restriction and a class-three lift but no class-four lift; its nine-term obstruction certificate is independently verified.
C.1	C.11; C.12; C.13; C.14; C.15; F.1; F.2; F.3	(a) Proved with the retained arithmetic term. The residue convergence criterion is combined with finite Galois cocycle matrices and explicit compact-support cup traces over every number field. These compute the arithmetic obstruction and the complete fiber inverse limit; the initial quadratic obstruction is also evaluated.
C.2	C.1; C.2; C.3	(a) Proved. Restriction is injective and finite Galois descent identifies precisely the invariant classes.
C.3	C.3; C.6; C.8; C.9; C.10; C.16; C.17; C.18	(a) Proved. Finite algebraic cover equations evaluate every point section in all 45 central coordinates. Their exact finite conjugacy criterion determines the point image, with the E36 origin and rank-44 fiber translation. The nine cusp packets, Galois action, and rational fixed subset are explicit.
C.4	C.4; C.5; C.7	(b) Refuted. Neither the R2 phantom nor the constructed intrinsic phantom dies after any finite extension.

Item	Statements	State
D.1	D.1; D.4; D.6; D.20; D.21; D.22; D.23; D.24; D.25; D.26; D.27; D.28; D.29; D.30; D.31; D.32; D.33; D.34; D.35; D.36; D.37; D.38; D.39; D.40; D.41; D.42; D.43; D.45; D.47	OPEN: dimension one or three. The actual geometric $\mathfrak{S}_3$ action excludes dimension two. Fully invariant finite five-group targets carry all three ordinary special outer parameters. Every first-precision parameter is class-preserving with an exact elementwise conjugator. All A_6 chief steps over an odd-order base have the proved equal positive generating-lift count, including coupled odd pullbacks. The ordinary even-base A_6^{20} obstruction and all earlier lifting comparisons remain retained. Realization or exclusion of the two-dimensional complement through every finer full finite source remains. Every generating lift at the first universal coprime elementary-p cover factors through an automorphism of that same cover; its complete fixed-conjugator fibre and ordinary centralizer union are parameterized and counted. Over every finite base, all pairing-compatible multiplier-one abelian lifts extend through the full relative pro-p cover for every coprime prime. The prescribed boundary maps give the exact primitive neck reduction of a closed free-action surface; its invariant integral Lagrangian proves the finite-field form comparison. The complete literal fibres and all-source factorization are retained. Over every actual pro-prime-to-p quotient, the entire peripheral automorphism fibre is the nonempty compact inverse limit of the finite-base fibres, through proved descent maps. Every D.37 parameter therefore lifts compatibly to the single actual quotient $\widehat{F}/O^p(O^2(K_n))$ for odd $p \neq 5$, retaining a chosen full relative pro-two lift and the exact maximal pro-five quotient P_n. Kernels at primes dividing the base order and the remaining nonabelian chief steps still require their full-source comparison. D.45 retains the full prescribed peripheral fibre across every actual $\mathrm{PSL}_2(32)$ chief extension over an arbitrary finite base, with the split cyclic-prime outer theorem and the exact old P_n action. D.47 proves the entire chief-lift theorem for every finite cyclic outer group, retaining the actual nonsplit automorphism extension, all intermediate characters, every prescribed conjugator and the complete ordinary union.

Item	Statements	State
D.2	D.2; D.4; D.6; D.7; D.8; D.20; D.22; D.23; D.24; D.25; D.26; D.27; D.28; D.29; D.30; D.31; D.32; D.33; D.34; D.35; D.36; D.38; D.44; D.46; D.48	OPEN: dimension one or two. The extra weight-nine vector, gamma line and complete finite pentagon comparison are verified. Elementary abelian and central obstructions, the complete A_5 chief-step count, and the ordinary order-five A_6^{20} obstruction retain their actual source maps and every centralizer choice. The latter also retains an admitted component above the prescribed five-group shadow. A cofinal equivariant lift or exclusion of the extra gentle direction remains. The exact odd-base A_6 chief comparison now fixes every prescribed peripheral conjugator and counts every generating lift at that stage. The exact $C_2 * C_3$ descent now retains each entire normalized pro-ℓ gentle pair on $\widehat{B}/O^p(O^\ell(F))$ for $p \notin \{2, 3, \ell\}$. The centerless pullback and unique cusp gauge identify it with the specified S_3-centralizing special outer automorphism, retaining the invariant extra weight-nine realization through this full relative tower. D.46 adds a strict return-prime quotient over that entire old source: its exact joint source is $\Gamma \times_D E$, every peripheral map generates and descends to an automorphism, and the full normalized fibre has $\ell^{\ell \operatorname{ord}_p(\ell)}$ elements. D.48 retains the same entire old gentle pair through the full integral affine return tower. The actual kernel is $\mathcal{O}^{6\ell}$, the normalized inverse-limit fibre is an $\mathcal{O}^\ell$ torsor, and every finite parameter lifts compatibly.
D.3	D.3; D.4; D.7; D.8; D.20	(a) Proved. Compatible finite braid quotients give gentle/pentagon dimensions 37/2 and leading metabelian dimensions 2/1. The integer certificate and actual Galois lower bound derive R9 in weight nine.
E.1	E.1; E.4	(a) Proved. Independent exact programs confirm every count, cusp equation, generation certificate, and both Smith forms.

Item	Statements	State
E.2	E.2; E.5; E.7; E.8; E.9; E.10; E.11; E.12; E.13; E.14; E.15; E.16; E.20; E.22; E.23; E.24; E.25; E.26; E.27; E.28; E.29; E.31; E.33; E.36; E.37; E.38; E.41; E.42; E.43; E.44; E.45; E.46; E.47; E.49; E.50; E.51; E.48; E.52; E.53	OPEN: complete CM fibers. All 24 CM fibres survive each proved refinement through E.49–E.51. The full three-adic class-three boundary correction-image fibre is $\mathbf{Z}_3^{70}$ over every one of the 432 multiplier-one sources. Four rational equations and two integral residue tests give its exact class-two projection. Its complete exponent-three image has 3^{25} points per source, with a specified finite nonlift surviving every change of higher class-two digits. The defined binary, prime-supported, $\mathrm{SL}_2(\mathbf{F}_5)$ and Hurwitz common families are retained. Full profinite survival remains. The full five-strand $\mathrm{Sp}_4(\mathbf{F}_3)$ test also retains all 864 points. Its complete source factors through G_6, its first B_4 target is the actual E.18 group, and three explicit simultaneous conjugators prove full B_5 generation for every retained word. The prime-five five-strand test retains all 432 multiplier-one points and every prescribed full boundary fibre. Its actual source of order 9000000 has common quotient $J = A_{10} \times_{A_2} Q_{20}$ of order 72000 with G_L, retaining all three signed modular bits. The exact residual correction in the old kernels reaches $(\mathbf{Z}/5)^3$, and one word gives generating maps on the full common sources and joint braid targets. The prime-seven source also retains this entire family and the prime-five source. Its exact residual correction is $\mathrm{SL}_2(\mathbf{F}_7) \times (\mathbf{Z}/7)^3$, its maximum common quotient with the full old boundary source is A_{14}, and the complete joint four- and five-strand targets are generated by the same corrected word. E.48 now retains all864 old points through the full A_6 Ferrari source, including its complete ternary and binary solvable layers, exact positive Kummer label, all centralizer choices and actual joint-source generation. E.52 identifies the full integral symplectic source at every modulus prime to6 and its common inverse-limit lower-central residual; its simultaneous peripheral comparison with the remaining sources is the next step. E.53 determines the exact uniform common source $G_L \times_{\mathcal{J}_5} G_R$, including the central sign and abelian kernel. Its full residual correction retains the old small-prime and level-five/seven values through the specified congruence subgroups. The resulting uniform fibre theorem is being completed.

Item	Statements	State
E.3	E.3; D.5; E.6; E.17; E.18; E.19; E.21; E.22; E.23; E.24; E.25; E.26; E.27; E.28; E.29; E.30; E.31; E.32; E.33; E.34; E.35; E.36; E.37; E.38; E.39; E.40; E.41; E.42; E.43; E.44; E.45; E.46; E.47; E.49; E.50; E.51; E.48; E.52; E.53	OPEN: full GT image. All 864 degree-six points pass the retained pentagons, braid and twist relations. The actual $\mathrm{SL}_2(\mathbf{F}_5)$ source maps onto the five-point Hurwitz source; the same word retains both through E.47's complete integral class-three families at multiplier one. Their full common four-strand target is $(T_{23} \times_{S_L} T_L) \times A_{96}$. The A_6 kernel/outer tests also retain all 864 points in their stated common sources. Compatibility through every finite quotient remains. The full five-strand $\mathrm{Sp}_4(\mathbf{F}_3)$ test also retains all 864 points. Its complete source factors through G_6, its first B_4 target is the actual E.18 group, and three explicit simultaneous conjugators prove full B_5 generation for every retained word. The prime-five five-strand test retains all 432 multiplier-one points and every prescribed full boundary fibre. Its actual source of order 9000000 has common quotient $J = A_{10} \times_{A_2} Q_{20}$ of order 72000 with G_L, retaining all three signed modular bits. The exact residual correction in the old kernels reaches $(\mathbf{Z}/5)^3$, and one word gives generating maps on the full common sources and joint braid targets. The prime-seven source also retains this entire family and the prime-five source. Its exact residual correction is $\mathrm{SL}_2(\mathbf{F}_7) \times (\mathbf{Z}/7)^3$, its maximum common quotient with the full old boundary source is A_{14}, and the complete joint four- and five-strand targets are generated by the same corrected word. E.48 now retains all864 old points through the full A_6 Ferrari source, including its complete ternary and binary solvable layers, exact positive Kummer label, all centralizer choices and actual joint-source generation. E.52 identifies the full integral symplectic source at every modulus prime to6 and its common inverse-limit lower-central residual; its simultaneous peripheral comparison with the remaining sources is the next step. E.53 determines the exact uniform common source $G_L \times_{\mathcal{J}_5} G_R$, including the central sign and abelian kernel. Its full residual correction retains the old small-prime and level-five/seven values through the specified congruence subgroups. The resulting uniform fibre theorem is being completed.
F.1	F.1; F.2; F.3; F.4	(a), with a corrected order test. The exact global correction and vanishing equivalences are proved over every number field, including 2 and real places. Orders must be taken after quotienting the maximal divisible subgroup.
F.2	F.5; A.22	(a) Proved for the specified finite-rank coefficients. Vanishing is computed for the Tate, cusp, elliptic, intrinsic, exterior-square, and finite-rank higher graded coefficient modules.
F.3	F.9; F.10; F.11; F.12	(a) Proved. The genuine integral action, tame equations including 2, obstruction group, Milnor summands, and section fibers are determined.

Item	Statements	State
F.4	F.20; F.21; F.22; F.23; F.24; F.25; F.26; F.27; F.28; F.29; F.30; F.31; F.32	(a) Proved. The three full profinite fixed groups are the identity, the actual full cusp inertia group, and the identity. The proof applies valuation uniqueness to the actual cyclotomic kernel, identifies its fixed primitive valuation, and recovers the exact geometric stabilizer. All fixed-lift images follow through the specified quotient maps.

References

[BCL20] Frauke M. Bleher, Ted Chinburg, and Alexander Lubotzky. On representations of $\mathrm{Gal}(\overline{\amalg}/\mathbb{Q})$, $\widehat{GT}$ and $\mathrm{Aut}(\widehat{F}_2)$, 2020. verified: opened attached primary source, Lemma 2.3 and Corollary 2.4 on unrestricted profinite automorphism lifting. The title and author metadata were checked against the attached source.

[BES15] Niels Borne, Michel Emsalem, and Jakob Stix. Lifting galois sections along torsors, 2015. verified: attached primary version 2 source and arXiv record opened; Theorem 5.10 and its proof, Proposition 5.9, and the relative-Brauer-group limitation to the rational ground field checked. No extension of the theorem to arbitrary number fields is assumed.

[BM24] Jean Barge and Julien Marché. The intersection form on the homology of a surface acted on by a finite group. *Mathematische Annalen*, 389:1255–1266, 2024. verified: opened the ten-page primary author version, Theorem 1 on p.1, which explicitly states the primitive invariant integral Lagrangian. The published preview has rational Theorem 1 and its integral formulation in Remark 1; publication metadata are printed there. The free action and the exact integral reduction required here are proved in D.42. The DOI resolver was unavailable.

[BMO12] Carles Broto, Jesper M. Møller, and Bob Oliver. Equivalences between fusion systems of finite groups of lie type. *Journal of the American Mathematical Society*, 25(1):1–20, 2012. verified: opened the primary author PDF, Lemma A.1(b) and its complete proof on printed pp.15–17, including the characteristic-two argument. Publisher issue metadata confirms the journal, volume, pages and DOI. The exact zero-invariant and coprimality hypotheses are checked in D.40.

[Bog26] Marco Boggi. A congruence subgroup property for symmetric mapping class groups. *Sao Paulo Journal of Mathematical Sciences*, 20:Article 35, 1–10, 2026. verified: opened the primary published PDF and publisher HTML; Theorem C(i) and Theorem 5.1(i) identify conjugacy classes of solvable finite subgroups in the discrete and full profinite mapping class groups in genus at most two. E.39 applies this to cyclic subgroups of the punctured genus-zero group and supplies the stabilizer, orientation and central-exponent comparisons.

[Bro15] Francis Brown. Zeta elements in depth 3 and the fundamental lie algebra of a punctured elliptic curve, 2015. verified: opened author's complete primary preprint, especially equation (1.3) and the following justification on pp.1–2; the use here is restricted to its motivic depth-one formula. The later published title replaces punctured elliptic curve by infinitesimal Tate curve.

[CFR12] Daniel C. Cohen, Michael Falk, and Richard Randell. Pure braid groups are not residually free. In *Configuration Spaces: Geometry, Combinatorics and Topology*, volume 14 of *Publications of the Scuola Normale Superiore*, pages 213–230. Edizioni della Normale, Pisa, 2012. verified: opened the primary author PDF, especially the literal pure Artin relations (2.1), central splitting (2.2)–(2.3), and integral infinitesimal braid presentation (4.1). Publication metadata checked against the author's publication list and institutional publication record.

[Col12] Benjamin Collas. Action of the grothendieck–teichmüller group on torsion elements of full teichmüller modular groups in genus zero. *Journal de théorie des nombres de Bordeaux*, 24(3):605–622, 2012. verified: opened the primary published PDF, Theorem B, Theorem 3.10 and its proof on printed pp.616–620, and the compatible braid action formulas. E.30 supplies the exact fifth-point-stabilizer and finite-quotient comparison.

[Cre19] Brendan Creutz. Generalized jacobians and explicit descents, 2019. verified: opened arXiv v3, 20 September 2019; section 2.1 equation (2.2) and section 2.3 Lemma 2.5, printed pp.4–5, actual semiabelian sequence and modulus divisor quotient.

[Del74] Pierre Deligne. La conjecture de weil. i. *Publications Mathématiques de l'IHÉS*, 43:273–307, 1974. verified: opened Numdam journal PDF; Theorem (1.6), printed p.276, PDF p.4, used for H1 of the smooth proper finite-field compactification.

[DG24] Vasily A. Dolgushev and Jacob J. Guynee. Gt-shadows for the gentle version of the grothendieck–teichmueller group, 2024. verified: opened the attached primary source and arXiv record; Theorem 5.2, Corollary 5.4 and Proposition 5.5, with finite shadow definitions and generation condition.

[DLL24] Vasily A. Dolgushev, Khanh Q. Le, and Aidan A. Lorenz. What are gt-shadows? *Algebraic and Geometric Topology*, 24(5):2721–2777, 2024. verified: opened the attached complete primary source, Appendix B, Theorem B.2; its abelian pure-quotient hypothesis does not apply to the specified CM quotient.

[Enr07] Benjamin Enriquez. Quasi-reflection algebras and cyclotomic associators. *Selecta Mathematica. New Series*, 13:391–463, 2007. verified: opened the complete primary arXiv v2 PDF, especially Theorem 8.1 and its proof, and Remark 8.6, PDF pp.30–32. The result is explicitly for the full profinite Grothendieck–Teichmueller group.

[Fur11] Hidekazu Furusho. Double shuffle relation for associators. *Annals of Mathematics*, 174(1):341–360, 2011. verified: opened the attached primary TeX source, Corollary 0.3, Theorem 0.5, Proposition 5.1 and Lemma 5.2, and opened the arXiv record; used for the characteristic-zero gamma upper bound, not a finite pentagon enumeration.

[Gui15] Pierre Guillot. The grothendieck–teichmüller group of a finite group and G-dessins d'enfants, 2015. verified: author arXiv v3 opened; Section 2.1, Lemma 2.1 and generating-pair construction; Section 2.3 and printed page 2 specify the full inverse limit and GT0 scope.

[HM03] Richard Hain and Makoto Matsumoto. Weighted completion of galois groups and galois actions on the fundamental group of p1 minus three points. *Compositio Mathematica*, 139:119–167, 2003. verified: opened the attached complete primary TeX source; Theorem 8.4, Section 8.4, Lemma 8.5 and Remark 8.6. The dependence of nontriviality on Ihara's work is recorded explicitly in the audit.

[HM19] Hikaru Hirano and Masanori Morishita. Arithmetic topology in ihara theory ii: Milnor invariants, dilogarithmic heisenberg coverings and triple power residue symbols. *Journal of Number Theory*, 198:211–238, 2019. verified: opened full arXiv PDF and publisher metadata; section 1.1, equations (1.1.1)–(1.1.3), printed pp.5–6, for Puiseux path comparison; section 2.1, equations (2.1.1)–(2.1.4.2), Theorem 2.1.4 and Corollary 2.1.6, printed pp.13–16, for the Heisenberg extension and deck maps.

[Hos10] Yuichiro Hoshi. Existence of nongeometric pro-p galois sections of hyperbolic curves. *Publications of the Research Institute for Mathematical Sciences*, 46(4):829–848, 2010. verified: opened author's May 2010 manuscript; Theorem 3.5 and its free-pro-p lifting proof on pp.17–18; publisher metadata also opened.

[HS00] David Harbater and Leila Schneps. Fundamental groups of moduli and the grothendieck–teichmueller group. *Transactions of the American Mathematical Society*, 352:3117–3148, 2000. verified: opened author's complete primary PDF; Main Theorem (a), the definition of symmetric quasi-special outer automorphisms, and its full profinite scope; Proposition 4, proof of (i) implies (iii), preprint pp.14–15, explicitly extends the gentle action to the full profinite braid group and sends the central generator to its lambda-th power.

[Iha91] Yasutaka Ihara. Braids, galois groups, and some arithmetic functions. In *Proceedings of the International Congress of Mathematicians, Kyoto, 1990*, volume I, pages 99–120. Mathematical Society of Japan, 1991. verified: opened the primary proceedings text at Sections 6.1–6.4, especially the nonvanishing and cocycle normalization in Section 6.2, the explicit theorem in Section 6.3, and equation (6.4.4). The exact Fox and leading-filtration comparisons are proved in D.20.

[Jan88] Uwe Jannsen. Continuous étale cohomology. *Mathematische Annalen*, 280:207–245, 1988. verified: primary PDF opened; Proposition 1.6 and Theorem 2.2 checked.

[KM04] Kamal Khuri-Makdisi. Linear algebra algorithms for divisors on an algebraic curve. *Mathematics of Computation*, 73(245):333–357, 2004. verified: opened complete arXiv v3 PDF and publisher metadata; Theorem/Algorithm 5.5, printed p.24, and sections 3–4 give divisor and Riemann–Roch operations used in C.16.

[Laz54] Michel Lazard. Sur les groupes nilpotents et les anneaux de lie. *Annales scientifiques de l'Ecole Normale Superieure*, 71(2):101–190, 1954. verified: opened the primary Numdam text, Chapter II, Theorems (4.2), (4.3), Corollary (4.4), Lemma (4.5), and Theorem (4.6), printed pp.175–180; especially the mutually inverse group/Lie construction and equality of homomorphisms for p-groups of class less than p.

[LNS26] Pierre Lochak, Hiroaki Nakamura, and Leila Schneps. The grothendieck–teichmueller group $\widehat{GT}$ acts on the genus g mapping class group with 0 or 1 marked point, 2026. verified: opened the complete primary preprint, v1 of 12 February 2026; Theorem 1.1, Theorems 2.1–2.5, introduction on p.2, and Section 8 on p.19. Theorem 1.1 constructs the full-GT actions; Section 8 explicitly leaves the indicated cutting-map compatibility unresolved.

[LS97] Pierre Lochak and Leila Schneps. A cohomological interpretation of the grothendieck–teichmueller group. *Inventiones Mathematicae*, 127:571–600, 1997. verified: opened the complete primary author PDF; Scheiderer's Appendix, unnumbered Corollary on printed p.598 (PDF page 28), states that every nontrivial

finite subgroup of a free profinite product is conjugate into a factor. The reduction to the gentle action is proved in E.8.

[Mat55] Arthur Mattuck. Abelian varieties over p-adic ground fields. *Annals of Mathematics*, 62(1):92–119, 1955. verified: opened complete primary paper; Theorem 7, printed p.114, PDF p.24, finite-index local analytic subgroup.

[Mil06] J. S. Milne. *Arithmetic Duality Theorems*. BookSurge, second edition, 2006. verified: primary book opened; I.4.10, I.4.15, I.5.1, II.2.9 and localization checked.

[Mil08] James S. Milne. Abelian varieties (v2.00), 2008. verified: opened the primary author PDF, title dated 16 March 2008, Chapter III Proposition 9.1 and Corollary 9.6, printed pp.113–115, and Proposition 9.7, printed p.115. The proper Abel–Jacobi fundamental-group comparison and generalized-Jacobian pullback construction are the steps used.

[Mil20a] James S. Milne. Algebraic number theory, 2020. verified: opened the author's full primary course notes and Theorems 8.42–8.43 with their proofs. The primitive integral element bound in Hermite's proof is used in C.14; the explicit local different and discriminant bound is proved there.

[Mil20b] James S. Milne. Class field theory, version 4.03, 2020. verified: opened the author PDF; IV.3.14 crossed products and V.3.23 Chebotarev checked.

[Mil20c] James S. Milne. Class field theory, version 4.03, 2020. verified: opened author-hosted complete PDF; Chapter V, Theorem 3.6, Examples 3.9 and 3.10, printed pp.158–159, PDF pp.166–167.

[Mil21] James S. Milne. *Elliptic Curves*. World Scientific, 2 edition, 2021. verified: opened author's complete second-edition PDF; Chapter II, Theorem 4.1, Corollary 4.2, Remark 4.3 and Aside 4.4, printed pp.62–65; Proposition 5.4 and Corollary 5.6, printed pp.66–67, for the local filtration and torsion specialization.

[MT24] Shinichi Mochizuki and Shota Tsujimura. Resolution of nonsingularities, point-theoreticity, and metric-admissibility for p-adic hyperbolic curves. RIMS preprint 1974, author-hosted revision dated 30 March 2024, 2024. verified: opened current author PDF and publication list; Definition 2.2(i), Proposition 2.3(v)–(vi), printed pp.37–38, Proposition 2.4(iv),(vii), printed pp.50–51, and Theorem 2.17, printed pp.71–72; earlier June 2023 RIMS version checked independently.

[Nak94] Hiroaki Nakamura. Galois rigidity of pure sphere braid groups and profinite calculus. *Journal of Mathematical Sciences, the University of Tokyo*, 1:71–136, 1994. verified: opened journal PDF; pro-class scope in section 2.1, Lemmas (2.1.2), (2.1.4), printed pp.94–97.

[NS00] Hiroaki Nakamura and Leila Schneps. On a subgroup of the grothendieck–teichmüller group acting on the tower of profinite teichmüller modular groups. *Inventiones Mathematicae*, 141(3):503–560, 2000. verified: independently opened author-uploaded complete text; section 5 Definition 5.1 and Proposition 5.3, printed pp.522–524; Remark 6.2, printed pp.525–526; the latter comparison explicitly retains relation (IV). The author-hosted version was also opened at Section 1, formula (1.1) and its quotient-action paragraph, and Theorem 2.5 on manuscript page 11; the latter matrix formula is used only on Galois elements. The arithmetic sign was checked again at the positive-root definition at the beginning of Section 2,

the generator-convention paragraph after (2.1), (2.4), and Lemma 2.3 on author-PDF p.9; the exact double-cover and inversion maps are proved in D.5.

[NTY10] Hiroaki Nakamura, Hiroshi Tsunogai, and Seidai Yasuda. Harmonic and equianharmonic equations in the grothendieck–teichmueller group. iii. *Journal of the Institute of Mathematics of Jussieu*, 9(2):431–448, 2010. verified: opened the complete author-uploaded primary PDF; Proposition 2.19, Proposition 4.4, Proposition 4.13, Theorem 4.16, and the scope statement on printed p.432. Proposition 4.4 is stated for GT; Theorem 4.16 is stated for Galois elements.

[Rib81] Kenneth A. Ribet. Torsion points of abelian varieties in cyclotomic extensions. *L'Enseignement Mathématique*, 27:315–319, 1981. verified: opened author's hosted complete PDF, Appendix Theorem 1, printed p.315, PDF p.32; used only the stated torsion finiteness theorem.

[Sch05] Leila Schneps. Grothendieck–teichmueller theory: Moduli spaces and multizeta values. Arizona Winter School lecture notes, 2005. verified: opened the complete primary lecture notes; Sections I.3 and I.6, printed pp.4–5 and 11–14, for the actual Ihara action, its composition law, the two-cycle and three-cycle conventions, and the injection of the Galois group into the full Grothendieck–Teichmueller group.

[Sti02] Jakob Stix. *Projective Anabelian Curves in Positive Characteristic and Descent Theory for Log-Etale Covers*. PhD thesis, Universität Bonn, 2002. verified: opened author's thesis PDF; Theorem 3.4.6, equation (6.2.5), Lemma 6.2.8, Corollary 6.2.9, printed pp.50 and 81–83.

[Whi11] Donald L. White. Character degrees of extensions of $\mathrm{PSL}_2(q)$. Author manuscript, July 25, 2011, 2011. verified: author PDF opened; Section 3 before Lemma 3.1 and Lemma 3.2 identify the actual field automorphism and full automorphism group. D.45 gives the direct proof and maps for PSL2(32).

[Wic12] Kirsten Wickelgren. On 3-nilpotent obstructions to pi1 sections for p1 over q minus 0, 1, infinity. In Jakob Stix, editor, *The Arithmetic of Fundamental Groups–PIA 2010*, volume 2 of *Contributions in Mathematical and Computational Sciences*. Springer, 2012. verified: opened attached arXiv TeX source, section 2.1, normalized Galois action and commutator-valued path cocycle; subsection Evaluating the 2-nilpotent quotient of f, equation with source label f_is_(1-chiˆ2)/24; publication metadata checked on arXiv abstract page.

[Wil25] Gareth Wilkes. Relative extensions and cohomology of profinite groups. *Journal of Pure and Applied Algebra*, 229(1):107846, 2025. verified: opened the primary arXiv v1 PDF and publisher metadata; Definition 4.1, Theorem 5.5, Definition 6.1 and Proposition 6.3. The full profinite finite-kernel relative extension and lifting comparisons are used with the explicit cochain identification in D.23.

[Yam23] Naganori Yamaguchi. The geometrically m-step solvable grothendieck conjecture for genus 0 curves over finitely generated fields, 2023. verified: primary version 3 PDF opened; Proposition 1.1.6 and the completed group-ring comparison of Section 1.1 checked. The use in A.19 concerns centralizers of nonzero powers of a free generator in a free metabelian pro-three group.