Preceding material.

Theorem If Under the preceding conditions
m1

Marginal note. m1

($\mathcal{G}$ a discrete metabelian group, G a group scheme locally of finite presentation over a base scheme S), the canonical map (61) is bijective if we assume moreover, so as not to tire myself|oneself, that injective, and has as image the set of homomorphisms quasi-unipotent from $\mathcal{G}$ into $G(S)$ — at least if we assume, to simplify life for ourselves, that S is noetherian and G affine over $S|S_0$.

Preceding material.

Theorem [d1] Under the preceding conditions
[m2]

Marginal note. [m2]

($\mathcal{G}$ a discrete metabelian group, G a group scheme locally of finite presentation over a base scheme S), the canonical map (61) is [d2] injective, and has as image[i1] the set of homomorphisms quasi-unipotent from $\mathcal{G}$ into $G(S)$ — at least if we assume, to simplify life for ourselves, that S is noetherian and G affine over $S|S_0$.

Preceding material.

Theorem † Under the preceding conditions
m3

Marginal note. m3

($\mathcal{G}$ a discrete metabelian group, G a group scheme locally of finite presentation over a base scheme S), the canonical map (61) is † injective, and has as image the set of homomorphisms quasi-unipotent from $\mathcal{G}$ into $G(S)$ — at least if we assume, to simplify life for ourselves, that S is noetherian and G affine over $S|S_0$.

Preceding material.

Theorem If Under the preceding conditions
m4

Marginal note. m4

($\mathcal{G}$ a discrete metabelian group, G a group scheme locally of finite presentation over a base scheme S), the canonical map (61) is bijective if we assume moreover, so as not to tire myself|oneself, that injective, and has as image the set of homomorphisms quasi-unipotent from $\mathcal{G}$ into $G(S)$ — at least if we assume, to simplify life for ourselves, that S is noetherian and G affine over $S|S_0$.

Preceding material.

Theorem [d1] Under the preceding conditions
[m5]

Marginal note. [m5]

($\mathcal{G}$ a discrete metabelian group, G a group scheme locally of finite presentation over a base scheme S), the canonical map (61) is [d2] injective, and has as image[i1] the set of homomorphisms quasi-unipotent from $\mathcal{G}$ into $G(S)$ — at least if we assume, to simplify life for ourselves, that S is noetherian and G affine over $S|S_0$.

Preceding material.

Theorem † Under the preceding conditions
m6

Marginal note. m6

($\mathcal{G}$ a discrete metabelian group, G a group scheme locally of finite presentation over a base scheme S), the canonical map (61) is † injective, and has as image the set of homomorphisms quasi-unipotent from $\mathcal{G}$ into $G(S)$ — at least if we assume, to simplify life for ourselves, that S is noetherian and G affine over $S|S_0$.

Preceding material.

Theorem If Under the preceding conditions
m7

Marginal note. m7

($\mathcal{G}$ a discrete metabelian group, G a group scheme locally of finite presentation over a base scheme S), the canonical map (61) is bijective if we assume moreover, so as not to tire myself|oneself, that injective, and has as image the set of homomorphisms quasi-unipotent from $\mathcal{G}$ into $G(S)$ — at least if we assume, to simplify life for ourselves, that S is noetherian and G affine over $S|S_0$.

Preceding material.

Theorem [d1] Under the preceding conditions
[m8]

Marginal note. [m8]

($\mathcal{G}$ a discrete metabelian group, G a group scheme locally of finite presentation over a base scheme S), the canonical map (61) is [d2] injective, and has as image[i1] the set of homomorphisms quasi-unipotent from $\mathcal{G}$ into $G(S)$ — at least if we assume, to simplify life for ourselves, that S is noetherian and G affine over $S|S_0$.

Preceding material.

Theorem † Under the preceding conditions
m9

Marginal note. m9

($\mathcal{G}$ a discrete metabelian group, G a group scheme locally of finite presentation over a base scheme S), the canonical map (61) is † injective, and has as image the set of homomorphisms quasi-unipotent from $\mathcal{G}$ into $G(S)$ — at least if we assume, to simplify life for ourselves, that S is noetherian and G affine over $S|S_0$.