Fixture en-0-90.

[scan 495]

[p. 550] 49. Homomorphisms of $M_{0,3}$, the general groups $M_{\rho,\sigma}$.
m2

Where we get our feet wet…

Marginal note. m2 All of this section I seems entirely useless for what follows, it is a preliminary groping to lead into section II page 553.

I) Consider a closed subgroup G of $M_{0,3}$, containing $\mathcal{C}^{+\wedge}_{0,3}$, hence the inverse image of a closed subgroup of $\tilde{\Gamma}_{0,3}$. We assume that the latter is "big enough" — say that it contains a subgroup of finite index in $\Gamma_{\mathbb{Q}} \subset \tilde{\Gamma}_{0,3}$. For the moment, we assume for simplicity that $G \subset \operatorname{Ker} \varepsilon_3$. In summary

$$(1) \qquad \mathcal{C}^{+\wedge}_{0,3} \subset G \subset \widetilde{M}_{0,3}, \qquad u \in G \Longrightarrow \mu(u) \equiv 1 \ (3).$$

End of opening definition.

Fixture en-0-170.

[scan 495]

[p. 550] 49. Homomorphisms of $M_{0,3}$, the general groups $M_{\rho,\sigma}$.
m4

Where we get our feet wet…

Marginal note. m4 All of this section I seems entirely useless for what follows, it is a preliminary groping to lead into section II page 553.

I) Consider a closed subgroup G of $M_{0,3}$, containing $\mathcal{C}_{0,3}^{+\wedge}$, hence the inverse image of a closed subgroup of $\tilde{\Gamma}_{0,3}$. We assume that the latter is "big enough" — say that it contains a subgroup of finite index in $\Gamma_{\mathbb{Q}} \subset \tilde{\Gamma}_{0,3}$. For the moment, we assume for simplicity that $G \subset \operatorname{Ker} \varepsilon_3$. In summary

$$(1) \qquad \mathcal{C}_{0,3}^{+\wedge} \subset G \subset \widetilde{M}_{0,3}, \qquad u \in G \Longrightarrow \mu(u) \equiv 1 \ (3).$$

End of opening definition.

Fixture en-0-260.

[scan 495]

[p. 550] m6 49. Homomorphisms of $M_{0,3}$, the general groups $M_{\rho,\sigma}$.

Where we get our feet wet…

Marginal note. m6 All of this section I seems entirely useless for what follows, it is a preliminary groping to lead into section II page 553.

I) Consider a closed subgroup G of $M_{0,3}$, containing $\mathcal{C}_{0,3}^{+\wedge}$, hence the inverse image of a closed subgroup of $\tilde{\Gamma}_{0,3}$. We assume that the latter is "big enough" — say that it contains a subgroup of finite index in $\Gamma_{\mathbb{Q}} \subset \tilde{\Gamma}_{0,3}$. For the moment, we assume for simplicity that $G \subset \operatorname{Ker} \varepsilon_3$. In summary

$$(1) \qquad \mathcal{C}_{0,3}^{+\wedge} \subset G \subset \widetilde{M}_{0,3}, \qquad u \in G \Longrightarrow \mu(u) \equiv 1 \ (3).$$

End of opening definition.

Fixture en-1-90.

[scan 495]

[p. 550] 49. Homomorphisms of $M_{0,3}$, the general groups $M_{\rho,\sigma}$.
m8

Where we get our feet wet…

Marginal note. m8 All of this section I[i1] seems entirely useless for what follows, it is a preliminary groping to lead into section II page 553.

I) Consider a closed[i2] subgroup G of $M_{0,3}$, containing $\mathcal{C}_{0,3}^{+\wedge}$, hence the inverse image of a closed subgroup of $\tilde{\Gamma}_{0,3}$. We assume that the latter is "big enough" — say that it contains a subgroup of finite index in $\Gamma_{\mathbb{Q}} \subset \tilde{\Gamma}_{0,3}$. For the moment, we assume for simplicity that $G \subset \operatorname{Ker} \varepsilon_3$. In summary

$$(1) \qquad \mathcal{C}_{0,3}^{+\wedge} \subset G \subset \widetilde{M}_{0,3}, \qquad u \in G \Longrightarrow \mu(u) \equiv 1 \ (3).$$

End of opening definition.

Fixture en-1-170.

[scan 495]

[p. 550] 49. Homomorphisms of $M_{0,3}$, the general groups $M_{\rho,\sigma}$.
m10

Where we get our feet wet…

Marginal note. m10 All of this section I[i1] seems entirely useless for what follows, it is a preliminary groping to lead into section II page 553.

I) Consider a closed[i2] subgroup G of $M_{0,3}$, containing $\mathcal{C}_{0,3}^{+\wedge}$, hence the inverse image of a closed subgroup of $\tilde{\Gamma}_{0,3}$. We assume that the latter is "big enough" — say that it contains a subgroup of finite index in $\Gamma_{\mathbb{Q}} \subset \tilde{\Gamma}_{0,3}$. For the moment, we assume for simplicity that $G \subset \operatorname{Ker} \varepsilon_3$. In summary

$$(1) \qquad \mathcal{C}_{0,3}^{+\wedge} \subset G \subset \widetilde{M}_{0,3}, \qquad u \in G \Longrightarrow \mu(u) \equiv 1 \ (3).$$

End of opening definition.

Fixture en-1-260.

[scan 495]

[p. 550] m12 49. Homomorphisms of $M_{0,3}$, the general groups $M_{\rho,\sigma}$.

Where we get our feet wet…

Marginal note. m12 All of this section I[i1] seems entirely useless for what follows, it is a preliminary groping to lead into section II page 553.

I) Consider a closed[i2] subgroup G of $M_{0,3}$, containing $\mathcal{C}_{0,3}^{+\wedge}$, hence the inverse image of a closed subgroup of $\tilde{\Gamma}_{0,3}$. We assume that the latter is "big enough" — say that it contains a subgroup of finite index in $\Gamma_{\mathbb{Q}} \subset \tilde{\Gamma}_{0,3}$. For the moment, we assume for simplicity that $G \subset \operatorname{Ker} \varepsilon_3$. In summary

$$(1) \qquad \mathcal{C}_{0,3}^{+\wedge} \subset G \subset \widetilde{M}_{0,3}, \qquad u \in G \Longrightarrow \mu(u) \equiv 1 \ (3).$$

End of opening definition.

Fixture en-2-90.

49. Homomorphisms of $M_{0,3}$, the general groups $M_{\rho,\sigma}$. m14

[scan 495]
[p. 550]
Where we get our feet wet…

Marginal note. m14 All of this section I seems entirely useless for what follows, it is a preliminary groping to lead into section II page 553.

I) Consider a closed subgroup G of $M_{0,3}$, containing $\mathcal{C}^{+\wedge}_{0,3}$, hence the inverse image of a closed

subgroup of $\tilde{\Gamma}_{0,3}$. We assume that the latter is "big enough" — say that it contains a subgroup of finite index in $\Gamma_{\mathbb{Q}} \subset \tilde{\Gamma}_{0,3}$. For the moment, we assume for simplicity that $G \subset \operatorname{Ker} \varepsilon_3$. In summary

$$(1) \qquad \mathcal{C}_{0,3}^{+\wedge} \subset G \subset \widetilde{M}_{0,3}, \qquad u \in G \Longrightarrow \mu(u) \equiv 1 \ (3).$$

End of opening definition.

Fixture en-2-170.

49. Homomorphisms of $M_{0,3}$, the general groups $M_{\rho,\sigma}$.
m16

[scan 495]
[p. 550]
Where we get our feet wet…

Marginal note. m16 All of this section I seems entirely useless for what follows, it is a preliminary groping to lead into section II page 553.

I) Consider a closed subgroup G of $M_{0,3}$, containing $\mathcal{C}^{+\wedge}_{0,3}$, hence the inverse image of a closed subgroup of $\tilde{\Gamma}_{0,3}$. We assume that the latter is "big enough" — say that it contains a subgroup of finite index in $\Gamma_{\mathbb{Q}} \subset \tilde{\Gamma}_{0,3}$. For the moment, we assume for simplicity that $G \subset \operatorname{Ker} \varepsilon_3$. In summary

$$(1) \qquad \mathcal{C}^{+\wedge}_{0,3} \subset G \subset \widetilde{M}_{0,3}, \qquad u \in G \Longrightarrow \mu(u) \equiv 1 \ (3).$$

End of opening definition.

Fixture en-2-260.

49. Homomorphisms of $M_{0,3}$, the general groups $M_{\rho,\sigma}$.
m18

[scan 495]
[p. 550]
Where we get our feet wet…

Marginal note. m18 All of this section I seems entirely useless for what follows, it is a preliminary groping to lead into section II page 553.

I) Consider a closed subgroup G of $M_{0,3}$, containing $\mathcal{C}^{+\wedge}_{0,3}$, hence the inverse image of a closed subgroup of $\tilde{\Gamma}_{0,3}$. We assume that the latter is "big enough" — say that it contains a subgroup of finite index in $\Gamma_{\mathbb{Q}} \subset \tilde{\Gamma}_{0,3}$. For the moment, we assume for simplicity that $G \subset \operatorname{Ker} \varepsilon_3$. In summary

$$(1) \qquad \mathcal{C}^{+\wedge}_{0,3} \subset G \subset \widetilde{M}_{0,3}, \qquad u \in G \Longrightarrow \mu(u) \equiv 1 \ (3).$$

End of opening definition.