

MATHEMATICAL PAPERS.

LONDON : C. J. CLAY AND SONS,
CAMBRIDGE UNIVERSITY PRESS WAREHOUSE,
AVE MARIA LANE.

Glasgow : 263, ARGYLE STREET.

Leipzig : F. A. BROCKHAUS.
New York : THE MACMILLAN CO.

*[Frontispiece : portrait of Professor Cayley,
reproduction by A. G. Dew-Smith, 1885.]*

THE COLLECTED
MATHEMATICAL PAPERS

OF

ARTHUR CAYLEY, Sc.D., F.R.S.,
LATE SADLERIAN PROFESSOR OF PURE MATHEMATICS IN THE
UNIVERSITY OF CAMBRIDGE.

VOL. XI.

CAMBRIDGE :
AT THE UNIVERSITY PRESS.
1896

[All Rights reserved.]

CAMBRIDGE :
PRINTED BY J. AND C. F. CLAY,
AT THE UNIVERSITY PRESS.

ADVERTISEMENT.

THE present volume contains 93 papers numbered 706 to 798, published, with the exception of one series, for the most part in the years 1878 to 1883. This series is constituted by the articles which Professor Cayley wrote for the *Encyclopædia Britannica* between the years 1878 and 1888; it seemed desirable to place these together in the same volume, in spite of the departure from the chronological arrangement which governs the sequence of the papers in the volumes generally. The Syndics of the University Press desire to acknowledge their obligation to Messrs Adam and Charles Black, Publishers of the ninth Edition of the *Encyclopædia Britannica*, for their courteous consent in allowing these articles to be included in the *Collected Mathematical Papers*. Exact references to the volumes, from which the articles are extracted, will be found in the Table of Contents.

The frontispiece to the present volume is a reproduction by Mr A. G. Dew-Smith, of Trinity College, of a photograph of Professor Cayley which he made in the year 1885. The Syndics of the Press desire to acknowledge their obligation to Mr Dew-Smith.

The Table for the eleven volumes is

Vol.	I.	Numbers	1 to 100,
„	II.	„	101 to 158,
„	III.	„	159 to 222,
„	IV.	„	223 to 299,
„	V.	„	300 to 383,
„	VI.	„	384 to 416,
„	VII.	„	417 to 485,
„	VIII.	„	486 to 555,
„	IX.	„	556 to 629,
„	X.	„	630 to 705,
„	XI.	„	706 to 798.

A. R. FORSYTH.

21 November, 1896.

CONTENTS.

[An Asterisk means that the paper is not printed in full.]

	PAGE
706. <i>On the distribution of electricity on two spherical surfaces</i>	1
Phil. Mag., Ser. 5, t. v. (1878), pp. 54—60	
707. <i>On the colouring of maps</i>	7
Geogr. Soc. Proc., t. i. (1879), pp. 259—261	
708. <i>Note sur la théorie des courbes de l'espace</i>	9
Assoc. Franç., Compt. Rend., t. ix. (1880), pp. 135—139	
709. <i>On the number of constants in the equation of the surface</i> <i>$PS - QR = 0$</i>	14
Tidsskrift for Mathematik, Ser. 4, t. iv. (1880), pp. 145—148	
710. <i>On a differential equation</i>	17
Collectanea Mathematica, in memoriam Dominici Chelini, (Milan, Hoepli, 1881), pp. 17—26	
711. <i>On a diagram connected with the transformation of elliptic</i> <i>functions</i>	26
British Association Report, 1881, p. 534	
712. <i>A partial differential equation connected with the simplest case</i> <i>of Abel's theorem</i>	27
British Association Report, 1881, pp. 534, 535	
713. <i>Addition to Mr. Rowe's "Memoir on Abel's theorem"</i>	29
Phil. Trans., t. clxxii. (1881), pp. 751—758	

	PAGE
714. <i>Various notes</i>	37
Messenger of Mathematics, t. vii. (1878), pp. 69; 115; 124; 125	
715. <i>Note on a system of algebraical equations</i>	39
Messenger of Mathematics, t. vii. (1878), pp. 17, 18	
716. <i>An illustration of the theory of the ϑ-functions</i>	41
Messenger of Mathematics, t. vii. (1878), pp. 27—32	
717. <i>On the triple theta-functions</i>	47
Messenger of Mathematics, t. vii. (1878), pp. 48—50	
718. <i>Addition to Mr. Genese's paper "On the theory of envelopes"</i> .	50
Messenger of Mathematics, t. vii. (1878), pp. 62, 63	
719. <i>Suggestion of a mechanical integrator for the calculation of</i> <i>$\int (X dx + Y dy)$ along an arbitrary path</i>	52
Messenger of Mathematics, t. vii. (1878), pp. 92—95; British Association Report, 1877, pp. 18—20	
720. <i>Note on Arbogast's method of derivations</i>	55
Messenger of Mathematics, t. vii. (1878), p. 158	
721. <i>Formulæ involving the seventh roots of unity</i>	56
Messenger of Mathematics, t. vii. (1878), pp. 177—182	
722. <i>A problem in partitions</i>	61
Messenger of Mathematics, t. vii. (1878), pp. 187, 188	
723. <i>Various notes</i>	63
Messenger of Mathematics, t. viii. (1879), pp. 45, 46; 126; 127	
724. <i>On the deformation of the model of a hyperboloid</i>	66
Messenger of Mathematics, t. viii. (1879), pp. 51, 52	
725. <i>New formulæ for the integration of $\frac{dx}{\sqrt{X}} + \frac{dy}{\sqrt{Y}} = 0$</i>	68
Messenger of Mathematics, t. viii. (1879), pp. 60—62	
726. <i>A formula by Gauss for the calculation of $\log 2$ and certain</i> <i>other logarithms</i>	70
Messenger of Mathematics, t. viii. (1879), pp. 125, 126	

	PAGE
727. <i>Equation of the wave-surface in elliptic coordinates</i>	71
Messenger of Mathematics, t. viii. (1879), pp. 190, 191	
728. <i>A theorem in elliptic functions</i>	73
Proc. Lond. Math. Soc., t. x. (1879), pp. 43—48	
729. <i>On a theorem relating to conformable figures</i>	78
Proc. Lond. Math. Soc., t. x. (1879), pp. 143—146	
730. [<i>Addition to Mr. Spottiswoode's paper "On the twenty-one coordinates of a conic in space"</i>]	82
Proc. Lond. Math. Soc., t. x. (1879), pp. 194—196	
731. <i>On the binomial equation $x^p - 1 = 0$; trisection and quarti- section</i>	84
Proc. Lond. Math. Soc., t. xi. (1880), pp. 4—17	
732. <i>A theorem in spherical trigonometry</i>	97
Proc. Lond. Math. Soc., t. xi. (1880), pp. 48—50	
733. <i>On a formula of elimination</i>	100
Proc. Lond. Math. Soc., t. xi. (1880), pp. 139—141	
734. <i>On the kinematics of a plane</i>	103
Quart. Math. Journ., t. xvi. (1879), pp. 1—8	
735. <i>Note on the theory of apsidal surfaces</i>	111
Quart. Math. Journ., t. xvi. (1879), pp. 109—112	
736. <i>Application of the Newton-Fourier method to an imaginary root of an equation</i>	114
Quart. Math. Journ., t. xvi. (1879), pp. 179—185	
737. <i>On a covariant formula</i>	122
Quart. Math. Journ., t. xvi. (1879), pp. 224—226	
738. <i>Note on a hypergeometric series</i>	125
Quart. Math. Journ., t. xvi. (1879), pp. 268—270	
739. <i>Note on the octahedron function</i>	128
Quart. Math. Journ., t. xvi. (1879), pp. 280, 281	

	PAGE
740. <i>On certain algebraical identities</i>	130
Quart. Math. Journ., t. xvi. (1879), pp. 281, 282	
741. <i>On a theorem of Abel's relating to a quintic equation</i>	132
Camb. Phil. Soc. Proc., t. iii. (1880), pp. 155—159	
742. <i>On the transformation of coordinates</i>	136
Camb. Phil. Soc. Proc., t. iii. (1880), pp. 178—184	
743. <i>On the Newton-Fourier problem</i>	143
Camb. Phil. Soc. Proc., t. iii. (1880), pp. 231, 232	
744. <i>Table of $\Delta^{m0n} \div \Pi(m)$ up to $m = n = 20$</i>	144
Camb. Phil. Trans., t. xiii. (1883), pp. 1—4	
745. <i>On the Schwarzian derivative, and the polyhedral functions</i>	148
Camb. Phil. Trans., t. xiii. (1883), pp. 5—68	
*746. <i>Higher Plane Curves</i>	217
Salmon's Higher Plane Curves, (3rd ed., 1879), Preface	
747. <i>Note on the degenerate forms of curves</i>	218
Salmon's Higher Plane Curves, (3rd ed., 1879), pp. 383—385	
748. <i>On the bitangents of a quartic</i>	221
Salmon's Higher Plane Curves, (3rd ed., 1879), pp. 387—389	
*749. <i>Solid Geometry</i>	224
Salmon's Treatise on the analytic geometry of three dimensions, (3rd ed., 1874), Preface	
750. <i>On the theory of reciprocal surfaces</i>	225
Salmon's Treatise on the analytic geometry of three dimensions, (3rd ed., 1874), pp. 539—550	
751. <i>Note on Riemann's paper "Versuch einer allgemeinen Auffass- ung der Integration und Differentiation," Werke, pp. 331—344</i>	235
Mathematische Annalen, t. xvi. (1880), pp. 81, 82	
752. <i>On the finite groups of linear transformations of a variable; with a correction</i>	237
Mathematische Annalen, t. xvi. (1880), pp. 260—263; 439, 440	
753. <i>On a theorem relating to the multiple theta-functions</i>	242
Mathematische Annalen, t. xvii. (1880), pp. 115—122	

	PAGE
754. <i>On the connection of certain formulæ in elliptic functions</i> ...	250
Messenger of Mathematics, t. ix. (1880), pp. 23—25	
755. <i>On the matrix $\begin{pmatrix} a, & b \\ c, & d \end{pmatrix}$, and in connection therewith</i> <i>the function $\frac{ax+b}{cx+d}$</i>	252
Messenger of Mathematics, t. ix. (1880), pp. 104—109	
756. <i>A geometrical construction relating to imaginary quantities</i> ..	258
Messenger of Mathematics, t. x. (1881), pp. 1—3	
757. <i>On a Smith's Prize question, relating to potentials</i>	261
Messenger of Mathematics, t. xi. (1882), pp. 15—18	
758. <i>Solution of a Senate-House problem</i>	265
Messenger of Mathematics, t. xi. (1882), pp. 23—25	
759. <i>Illustration of a theorem in the theory of equations</i>	268
Messenger of Mathematics, t. xi. (1882), pp. 111—113	
760. <i>Reduction of $\int \frac{dx}{(1-x^2)^{\frac{1}{3}}}$ to elliptic integrals</i>	270
Messenger of Mathematics, t. xi. (1882), pp. 142, 143	
761. <i>On the theorem of the finite number of the covariants of a</i> <i>binary quantic</i>	272
Quart. Math. Journ., t. xvii. (1881), pp. 137—147	
762. <i>On Schubert's method for the contacts of a line with a surface</i>	281
Quart. Math. Journ., t. xvii. (1881), pp. 244—258	
763. <i>On the theorems of the 2, 4, 8, and 16 squares</i>	294
Quart. Math. Journ., t. xvii. (1881), pp. 258—276	
764. <i>The binomial equation $x^p - 1 = 0$; quinquisection</i>	314
Proc. Lond. Math. Soc., t. xii. (1881), pp. 15, 16	
765. <i>On the flexure and equilibrium of a skew surface</i>	317
Proc. Lond. Math. Soc., t. xii. (1881), pp. 103—108	
766. <i>On the geodesic curvature of a curve on a surface</i>	323
Proc. Lond. Math. Soc., t. xii. (1881), pp. 110—117	

	PAGE
767. <i>On the Gaussian theory of surfaces</i>	331
Proc. Lond. Math. Soc., t. xii. (1881), pp. 187—192	
768. <i>Note on Landen's theorem</i>	337
Proc. Lond. Math. Soc., t. xiii. (1882), pp. 47, 48	
769. <i>On a formula relating to elliptic integrals of the third kind</i> ..	340
Proc. Lond. Math. Soc., t. xiii. (1882), pp. 175, 176	
770. <i>On the 34 concomitants of the ternary cubic</i>	342
American Journal of Mathematics, t. iv. (1881), pp. 1—15	
771. <i>Specimen of a literal table for binary quantics, otherwise a partition table</i>	357
American Journal of Mathematics, t. iv. (1881), pp. 248—255	
772. <i>On the analytical forms called trees</i>	365
American Journal of Mathematics, t. iv. (1881), pp. 266—268	
773. <i>On the 8-square imaginaries</i>	368
American Journal of Mathematics, t. iv. (1881), pp. 293—296	
774. <i>Tables for the binary sextic</i>	372
American Journal of Mathematics, t. iv. (1881), pp. 379—384	
775. <i>Tables of covariants of the binary sextic</i>	377
Written in 1894 : now first published.	
776. <i>On the Jacobian sextic equation</i>	389
Quart. Math. Journ., t. xviii. (1882), pp. 52—65	
777. <i>A solvable case of the quintic equation</i>	402
Quart. Math. Journ., t. xviii. (1882), pp. 154—157	
778. [<i>Addition to Mr. Hudson's paper "On equal roots of equations"</i>] ..	405
Quart. Math. Journ., t. xviii. (1882), pp. 226—229	
779. [<i>Note on Mr. Jeffery's paper "On certain quartic curves, which have a cusp at infinity, whereat the line at infinity is a tangent"</i>]	408
Proc. Lond. Math. Soc., t. xiv. (1883), p. 85	

	PAGE
780. [<i>Addition to Mr. Hammond's paper "Note on an exceptional case in which the fundamental postulate of Professor Sylvester's theory of tamisage fails"</i>]	409
Proc. Lond. Math. Soc., t. xiv. (1883), pp. 88—91	
781. <i>On the automorphic transformation of the binary cubic function</i>	411
Proc. Lond. Math. Soc., t. xiv. (1883), pp. 103—108	
782. <i>On Monge's "Mémoire sur la théorie des déblais et des remblais"</i>	417
Proc. Lond. Math. Soc., t. xiv. (1883), pp. 139—142	
783. <i>On Mr. Wilkinson's rectangular transformation</i>	421
Proc. Lond. Math. Soc., t. xiv. (1883), pp. 222—229	
784. <i>Presidential Address to the British Association, Southport, September 1883</i>	429
British Association Report, 1883, pp. 3—37	
785. <i>Curve</i>	460
Encyclopædia Britannica, 9th ed., t. vi. (1878), pp. 716—728	
786. <i>Equation</i>	490
Encyclopædia Britannica, 9th ed., t. viii. (1878), pp. 497—509	
787. <i>Function</i>	522
Encyclopædia Britannica, 9th ed., t. ix. (1879), pp. 818—824	
788. <i>Galois</i>	543
Encyclopædia Britannica, 9th ed., t. x. (1879), p. 48	
789. <i>Gauss</i>	544
Encyclopædia Britannica, 9th ed., t. x. (1879), p. 116	
790. <i>Geometry (analytical)</i>	546
Encyclopædia Britannica, 9th ed., t. x. (1879), pp. 408—420	
791. <i>Landen</i>	583
Encyclopædia Britannica, 9th ed., t. xiv. (1882), p. 271	
792. <i>Locus</i>	585
Encyclopædia Britannica, 9th ed., t. xiv. (1882), pp. 764, 765	

	PAGE
793. <i>Monge</i>	586
<i>Encyclopædia Britannica</i> , 9th ed., t. xvi. (1883), pp. 738, 739	
794. <i>Numbers (partition of)</i>	589
<i>Encyclopædia Britannica</i> , 9th ed., t. xvii. (1884), p. 614	
795. <i>Numbers (theory of)</i>	592
<i>Encyclopædia Britannica</i> , 9th ed., t. xvii. (1884), pp. 614—624	
796. <i>Series</i>	617
<i>Encyclopædia Britannica</i> , 9th ed., t. xxi. (1886), pp. 677—682	
797. <i>Surface</i>	628
<i>Encyclopædia Britannica</i> , 9th ed., t. xxii. (1887), pp. 668—672	
798. <i>Wallis (John)</i>	640
<i>Encyclopædia Britannica</i> , 9th ed., t. xxiv. (1888), pp. 331, 332	
<i>Portrait</i>	<i>To face Title.</i>

CLASSIFICATION.

ANALYSIS.

Calculation of $\log 2$; 726.
Series, 796.
Prime roots of unity, 721.
8-square imaginaries, 773.
Squares, theorems of 2, 4, 8, 16; 763.
Difference-table for $\Delta^m 0^n \div \Pi(m)$; 744.
Equations, theory of, 736, 741, 743, 759, 776, 777, 778, 786.
Numbers, theory of, 731, 764, 795.
Partitions, 722, 771, 794.
Trees, 772.
Matrices, 755.
Elimination, 733.
Transformation of cubic function, 781.
Covariantive forms and tables, 737, 761, 770, 774, 775, 780.
Fractional differentiation, 751.
Mechanical integrator, 719.
Differential Equations, 725.
Schwarzian derivative, 745.
Hypergeometric series, 710, 738.
Finite groups, 752.
Polyhedral functions, 739, 745.
Elliptic functions, 728, 740, 754, 760, 768, 769.
Transformation of elliptic functions, 711.
Abel's theorem, 712, 713.
Theta Functions, 716, 717, 753.
Function, 787.
Conformal representation, 729.

GEOMETRY.

Analytical geometry in general, 790, 792, 797.
Plane Curves, 746*, 785.
Degenerate forms of curves, 747.
Quartic Curves, 748, 779.
Déblais et remblais, 782.
Tortuous curves, 708, 785.
Geodesic curvature, 766.
Theory of surfaces, general, 709, 749*, 767.
Transformation of coordinates, 742, 783.
Reciprocal surfaces, 750.
Wave-surface, 727.
Apsidal surfaces, 735.
Deformation and flexure of surfaces, 724, 765.
Hypergeometry, 730.
Schubert's numerative geometry, 762.

VARIOUS SUBJECTS.

Spherical Trigonometry, 732.
Kinematics of a plane, 734.
Maps, colouring of, 707.
Electricity, distribution of, on spherical surfaces, 706.
Potential, 757.
Presidential Address to the British Association, 784.
Biographical articles ; Galois, 788. Gauss, 789. Landen, 791. Monge, 793. Wallis (John), 798.

MISCELLANEOUS.

714, 715, 718, 720, 723, 756, 758.

706.

ON THE DISTRIBUTION OF ELECTRICITY ON TWO SPHERICAL SURFACES.

[From the *Philosophical Magazine*, vol. v. (1878), pp. 54—60.]

IN the two memoirs “Sur la distribution de l’électricité à la surface des corps conducteurs,” *Mém. de l’Inst.* 1811, Poisson considers the question of the distribution of electricity upon two spheres : viz. if the radii be a , b , and the distance of the centres be c (where $c > a + b$, the spheres being exterior to each other), and the potentials within the two spheres respectively have the constant values h and g , then—for Poisson’s $f\left(\frac{x}{a}\right)$ writing $\phi(x)$, and for his $F\left(\frac{x}{b}\right)$ writing $\Phi(x)$ —the question depends on the solution of the functional equations

$$a\phi(x) + \frac{b^2}{c-x}\Phi\left(\frac{b^2}{c-x}\right) = h,$$

$$\frac{a^2}{c-x}\phi\left(\frac{a^2}{c-x}\right) + b\Phi(x) = g,$$

where of course the x of either equation may be replaced by a different variable.

It is proper to consider the meaning of these equations : for a point on the axis, at the distance x from the centre of the first sphere, or say from the point A , the potential of the electricity on this spherical surface is $a\phi x$ or $\frac{a^2}{x}\phi\left(\frac{a^2}{x}\right)$, according as the point is interior or exterior ; and, similarly, if x now denote the distance from the centre of the second sphere (or, say, from the point B), then the potential of the electricity on this spherical surface is $b\Phi x$ or $\frac{b^2}{x}\Phi\left(\frac{b^2}{x}\right)$, according as the point is interior or exterior ; $\phi(x)$ is thus the same function of

712.

A PARTIAL DIFFERENTIAL EQUATION CONNECTED
WITH THE
SIMPLEST CASE OF ABEL'S THEOREM.

[From the *Report of the British Association for the Advancement of
Science*, (1881), pp. 534, 535.]

CONSIDER a given cubic curve cut by a line in the points (x, p) , (x_1, p_1) , (x_2, p_2) ; using the first and second points as elements, these determine uniquely the third point. Analytically, the equation of the curve determines p as a function of x , and p_2 as a function of x_1 ; writing in the equation

$$c_2 = \lambda c_1 + (1 - \lambda)c, \quad y_2 = \lambda y_1 + (1 - \lambda)y_0,$$

we have c , by a simple equation, and thence p_2 ; viz. x_2 is found as a function of x , x_1 , and of the nine constants of the equation. Hence having the derived equations (in regard to x , x_1) of the first, second, and third orders, we have $(1 + 2 + 3 + 4) = 10$ equations from which to eliminate the 6 quantities x_2 , considered as a function of x and x_1 , thus involves a partial differential equation of the third order, independent of the particular cubic curve.

To obtain this equation it is only necessary to observe that we have, by Abel's theorem,

$$\frac{dx}{X} + \frac{dx_1}{X_1} + \frac{dx_2}{X_2} = 0,$$

where X , X_1 , X_2 is a given function of x , and p , that is, of x ; X_1 and X_2 are the like functions of x_1 and x_2 respectively. Hence, considering x_2 as a function of x and x_1 , we have

$$\frac{dx_2}{dx} = -\frac{X_2}{X}, \quad \frac{dx_2}{dx_1} = -\frac{X_2}{X_1},$$

[C. XI. 4-2]

and consequently

$$\frac{dx_2}{dx_1} \div \frac{dx_2}{dx} = \frac{X}{X_1},$$

where X , X_1 are functions of x_2 , x_1 respectively; hence taking the logarithm and differentiating successively with regard to x_1 and x_2 , we have

$$\frac{d}{dx_2} \frac{d}{dx_1} \log \left(\frac{dx_2}{dx_1} \div \frac{dx_2}{dx} \right) = 0,$$

which is the required partial differential equation of the third order.

This differential equation has a simple geometrical significance. Consider three consecutive positions of the line meeting the cubic curve in the points 1, 2, 3; 1', 2', 3'; 1'', 2'', 3'' respectively; *quæ* equation of the third order, the equation should in effect determine 3'' by means of the other points. And, in fact, the three positions of the line constitute a cubic curve; the nine points are thus the intersections of two cubic curves, or, say, they are an “annexed” of points; any eight of the points thus determine uniquely the ninth point.

.....

713.

ADDITION TO MR ROWE'S MEMOIR ON ABEL'S THEOREM.

[From the *Philosophical Transactions of the Royal Society of London*, vol. CLXII., Part III. (1881), pp. 715—758. Received May 27,—Read June 16, 1880.]

IN Abel's general theorem y is an irrational function of x determined by an equation $\chi(y) = 0$, or say $\chi(x, y) = 0$; if the order n as regards y , and that is also by him that the sum of any number of the integrals considered may be reduced to a sum of ρ integrals; where ρ is a determinate number depending only on the form of

the equation $\chi(x, y) = 0$, and given in his equation (42), [(*Euvres Complètes*, (1881), t. I., p. 157]; viz. if, solving the equation so as to obtain from it developments of y in descending series of powers of x , we have:

$$\begin{array}{ll} v_{\sigma}\rho_{\nu}, & \text{series each of the form } y = x^{\sigma'_{\nu}} + \dots, \\ & v_{\sigma}\rho_2, \quad y = x^{\sigma'_2} + \dots, \\ & \vdots \quad \quad \quad \vdots \\ & v_{\sigma}\rho_n, \quad y = x^{\sigma'_n} + \dots, \end{array}$$

⁰The second power of a term coefficient: this being a relation $y = x^p + \dots$, representing the y , different values of y obtained by giving to the radical x^p and to its p values, and the corresponding values in the radicals which enter into representing y , values of y . It is assumed that the equation is such that the meaning is to make each representing y , y_2 . In each of the equations the meaning of y is taken positive in the development of y in descending powers of x .

(so that $\sigma = v_1\rho_1 + v_2\rho_2 + \cdots + v_n\rho_n$) then y is a determinate function of $x_1, x_2, x_3, x_4, x_5, x_6, \dots, x_n$, say ρ .

We have so expressed Abel's ρ in the following form, viz. assuming

$$y = \Sigma a_1 \alpha_1 \nu_1 + \Sigma a_2 \nu_2 \cdots + \Sigma a_n \nu_n - \frac{1}{2} \Sigma a - \frac{1}{2} n + 1,$$

or, what is the same thing, for ρ writing this Σa_1 ,

$$y = \Sigma a_1 \nu_1 + \Sigma a_2 \nu_2 + \Sigma a_3 \nu_3 + \cdots + \Sigma a_n \nu_n - \frac{1}{2} \Sigma a - \frac{1}{2} n + 1,$$

where in the first sum ν_1 have each of them the values $1, 2, \dots, \rho_1$ subject to the condition $i + \nu_1 \sigma'_1$; in each of the other sums ν_i and ρ_n are interrelated by the suffix, which has the values $1, 2$, etc.

It is a leading result in Riemann's theory of the Abelian integrals that y is the deficiency (Geschlecht) of the curve represented by the equation $\chi(x, y) = 0$; and it must consequently be demonstrable *a posteriori* that the foregoing expression for y is in fact = deficiency of curve $\chi(x, y) = 0$. I propose to verify this by means of the formulae given in my paper "On the Higher Singularities of a Plane Curve," *Quart. Math. Journ.*, vol. VII., (1866), pp. 212—222, [374].

It is necessary to distinguish between the values of $\frac{m}{n}$ which are $>$, $=$, and < 1 ; and to fix the ideas I assume $l = 7$, and

$$\begin{aligned} & \frac{m_1}{n_1}, \frac{m_2}{n_2}, \frac{m_3}{n_3}, \text{ each } > 1, \\ & \frac{m_4}{n_4} = 1; \text{ say } m_4 = n_4 = n, \text{ and } n_4 = n; \\ & \frac{m_5}{n_5}, \frac{m_6}{n_6}, \frac{m_7}{n_7}, \text{ each } < 1, \end{aligned}$$

but it will be easily seen that the reasoning is quite general. And now Σ to denote a sum in regard to the first set of suffixes $1, 2, 3$, and Σ' to denote a sum in regard to the second set of suffixes $5, 6, 7$. The foregoing value of ρ is thus

$$y = \Sigma a \nu + 4 + \Sigma' a \nu.$$

Introducing a third coordinate z for homogeneity, the equation $\chi(x, y) = 0$ of the curve will be

$$0 = \left[y^{p-(n-1)}, y^{n-2} x^a, \dots \right] (y, z)^{n-2} + \left(y^{n-2}, y^{n-3} x^a, \dots \right) (y, z)^{n-1} + \cdots + (y)$$

where it is to be observed that $\chi(x, y) = 0$ is the product of ν_i , different ovaries; these divide themselves into n_i

groups, each a product of ρ_i orates; and in each such product the ρ_i coefficients A_i are in general the ρ_i values of a function containing a radical a^{1/ρ_i} and are thus different from each other; it is to when follows in effect assumed in each step that all the a_i , but that all the x_i coefficients, etc. are different from each other.

The remarks apply to the other factors. It applies in particular to the factor $(y, z)^{r-1}$, viz. it is assumed that the coefficients A in the radical $z, y^{n-2}, \dots$ are all of them different from each other. These assumptions as to the leading coefficients really imply Abel's assumption that $\frac{m_1}{n_1}, \frac{m_2}{n_2}$ are all of them fractions in their least terms, viz. that $n = 1$, since

1 rotate however for convenience the equation will be considered, by giving to

In the product of the several infinite series, the terms containing the lowest powers of $z, \frac{1}{n_1}$ is a function in its lowest terms, viz. since $n_1 = 1$.

A curve has singularities (singular points) at infinity, that is, on the line $z = 0$; viz. *First*, a singularity at $(x = 0, x' = 0)$, where the tangent is $x' = 0$, and which, writing for convenience $x = 1$, is denoted by the function

$$(z - \nu^{m/n} x^{1-1/n})^{n-1} = \dots$$

where observe that the expressed factor indicates r_n branches $(z - \nu^{m/n} x^{1-1/n})^{r_n}$, or say $r_n(m_n - p_n)$ partial branches $z = \nu^{m/n} \dots$, that is, $r_n(m_n - p_n)$ partial branches $z = \nu^{m/n} \dots$, will in all $r_n(m_n - p_n) + \dots$, where $r_n(m_n - p_n)$ partial branches at the encompassed factors with the suffixes 1' and 4; that is, we can regard the encompassed factors with the suffixes 1' and 4, where in the equation

$$z - \nu^{m/n} + \dots = 0.$$

Secondly, a singularity at $(z = 0, y = 0)$, where the tangent is $y = 0$, and which, writing for convenience $z = 1$, is denoted by the function

$$(y - z^{m/n} x^{1-1/n})^{n-1} = \dots$$

^{0*} This assumption is virtually made by Abel, (1), [c. p. 157], in considers

where observe that the expressed factor indicates r_5 branches $(y - z^{m/n}x^{1-1/n})^{r_5}$, or say $r_5(n_5 - m_5)$ partial branches $y = z^{m/n} \dots$, that is, $r_5(n_5 - m_5)$ partial branches $y = z^{m/n} \dots$, with in all $r_5n_5 + \dots$, with $r_5n_5 + r_6n_6 + r_7n_7$ partial branches $y = z^{m/n} \dots$, with the suffixes 5, 6, 7.

Thirdly, suppose at $(x = 0, y = 0)$, where the tangent is $z = 0$, and consequently $r_4n_4 = 0$, partial branches $z = y^0 \dots +$ similar terms with respect to x, y, z .

It is convenient to assume $r_4 = 2$, so that

$$\begin{aligned} E &= \frac{1}{2}(K - 1)(K - 2) = z, \\ E' &= \frac{1}{2}(K' - 1)(K' - 2) \\ &= \frac{1}{2}\Sigma(n - 1). \end{aligned}$$

hence

$$E + e' = \frac{1}{2}(K - 1) + \frac{1}{2}\Sigma(n - 1).$$

Assuming that there are no other singularities, the deficiency

$$\rho = \frac{1}{2}(K - 1)(K - 2) - E - e'$$

is

$$= \frac{1}{2}(K - 1)(K - 2) - \frac{1}{2}KK + \frac{1}{2}\Sigma(n - 1).$$

This should be equal to the before-mentioned value of ρ ; viz. we ought to have

$$\frac{1}{2}(K - 1)(K - 2) - \frac{1}{2}KK + \frac{1}{2}\Sigma(n - 1) = \Sigma a\nu_1 + 4 + \Sigma' a\nu_2 - \Sigma a - \Sigma' a - 2,$$

or, as it will be convenient to write it,

$$M = K - \frac{3}{2}K + \frac{1}{2}\Sigma(n - 1) - 1 = \Sigma a\nu_1 + \Sigma' a\nu_2 - \Sigma' a - 2,$$

which is the equation which ought to be satisfied by the values of N and $\Sigma(n - 1)$ calculated, according to the method of my paper, for the foregoing singularities of the curve.

the equation $z^n = y^n \sqrt[n]{f(x)}$ as denoting integers; but where it appears that the indices a, b , are to be considered of fixed, and as in the values of degrees with respect to x . He says that if these be assumed as integers in the expression of y in descending powers of x (the leading coefficient being z in regard to y), the singular factor are in (x, y) in regard to y in regard to x .

We have as before:

$$K = \Sigma n + \Sigma' n + \theta n.$$

The terms $\Sigma n + \Sigma' n + \theta n$, written at length, is

$$\begin{aligned} &= n_1(\nu_1\rho_1 + \nu_2\rho_2 + \nu_3\rho_3) + \cdots + n_2(\nu_1\rho_1 + \nu_2\rho_2 + \nu_3\rho_3) \\ &+ \cdots + n_3(\nu_1\rho_1 + \nu_2\rho_2 + \nu_3\rho_3) \\ &+ \cdots + n_4(\nu_1\rho_1 + \nu_2\rho_2 + \nu_3\rho_3) \\ &+ \cdots + n_5(\nu_1\rho_1 + \nu_2\rho_2 + \nu_3\rho_3) \\ &+ \cdots + n_6(\nu_1\rho_1 + \nu_2\rho_2 + \nu_3\rho_3) \\ &+ \cdots + n_7(\nu_1\rho_1 + \nu_2\rho_2 + \nu_3\rho_3). \end{aligned}$$

which is

$$= \Sigma \nu_1 n_2 + \Sigma' n_2 (2\rho_2 + 2'\rho_1) + \Sigma \Sigma' + \Sigma' \Sigma + \Sigma a_1 a_2 a_3,$$

We have moreover

$$\Sigma' \rho_1 = \Sigma a_1 \rho_1 + (\theta + \Sigma' a_1) \rho_1 = \Sigma a_1 + \Sigma a_2 + \Sigma' a_1 \rho_1,$$

each branch $(z - y^{m/n} x^{1-1/n})^{n-1}$ gives $a = m_1 \rho_1$, and the value of $\Sigma(\kappa - 1)$ for this singularity is

$$n_1(m_1 - n_1 - 1) + r_1 n_1(m_1 - 1) = r_1 n_1 \rho_1 - n_1 \rho_1.$$

which is

$$= \Sigma a \nu_1 - \Sigma a \rho_1.$$

each branch $(y - z^{m/n} x^{1-1/n})^{n-1}$ gives $\sigma = m_n \rho_n$, and the value of $\Sigma(\kappa - 1)$ for this singularity is

$$n_4(m_4 - n_4 - 1) + r_4 n_4(m_4 - 1) = r_4 n_4 \rho_4 - n_4 \rho_4,$$

which is

$$= \Sigma' \nu_2 - \Sigma' \nu_4 - \Sigma' \nu_1.$$

For each of the θ singularities

$$(y - z^{m/n} x^{1-1/n})^{n-1}.$$

we have $a = n$, and the value of $\Sigma(\kappa - 1)$ is $r = \theta(\kappa - 1)$; this is $= 0$ for the value $\lambda = 1$, which is physically attributed to λ .

According to the theory explained in my paper above referred to, the several singularities not together equivalent to a certain number K of κ^* nodes and cusps; viz. we have

$$K = \Sigma n - \Sigma a + \frac{1}{2} \Sigma(\kappa - 1).$$

This should be equal to the before-mentioned value of ρ ; viz. we ought to have

$$(K-1)(K-2) - 2(K + \Sigma n + \Sigma' n + \theta n) = 2\Sigma a \nu_1 + 4 + 2\Sigma a \nu_2 - 2\Sigma a - 2\Sigma' a - 4,$$

or to it will be convenient to write

$$M - K - \Sigma K + 1 + \Sigma(n - 1) = \Sigma a \nu_1 + \Sigma' a \nu_2 - \Sigma a - \Sigma' a - 2,$$

which is the equation which ought to be satisfied by the values of N and $\Sigma(n - 1)$ calculated, according to the method of my paper, for the foregoing singularities of the curve.

We have as before:

$$K = \Sigma n + \Sigma' n + \theta n.$$

The term $\Sigma n + \Sigma' n + \theta n$, written at length, is

$$= \Sigma \nu_1 n_1 + \Sigma' \nu_2 n_2 + \theta n,$$

[C. XI. 5]

or, reducing,

$$M = (\Sigma m)^2 - \Sigma ma - \Sigma m\rho_1 - \Sigma m\rho_2 + (\Sigma' m)^2 - \Sigma' m_2 a - \Sigma' m_2 \rho_2 \\ + (\Sigma' m)^2 - \Sigma' m\rho_1 - \Sigma ma + 2\Sigma' m_2 a_1 + 2\Sigma' \Sigma a_1 a_2;$$

and it is to be shown that the two lines of this expression are in fact the values of M belonging to the singularities

$$\left(z - y^{m/n} x^{1-1/n}\right)^{n-1}, \dots, \text{ and } \left(y - z^{m/n} x^{1-1/n}\right)^{n-1}, \dots,$$

respectively. We assume $\lambda = 1$, and there is then no singularity $(y - z)^{n-1}$.

I recall that, considering the several partial branches which meet in a singular point, M denotes the sum of the number of the intersections of each partial branch by every other partial branch: so that for each pair of partial branches the intersections are to be counted twice. Supposing that the tangent is $x = 0$, and that, for any two branches we have $z = Az^x + \dots$ (x where p_x, p_x are each up to greater than 1), then if $p_x = p_x$, and $A_x = A_x, A_x = A_x$ where A_x, A_x are not the partial branches which have here always could be regards the cases about to be considered), then the number of intersections is taken to be $-$; and if $p_x > p_x$, and the number of intersections (taken to be $\equiv p_x$) viz. in the case of as a unequal exponents, it is equal to the smaller exponents.

Consider now the singularity $\left(z - y^{m/n} x^{1-1/n}\right)^{n-1}$, and first the intersections of a partial branch $z = y^{m/n} \dots$ by each of the remaining $n_1(m_1 - p_1) - 1$ partial branches of the same set: the number of intersections with any one of these is $= \frac{m_1}{n_1} [n_1(m_1 - p_1) - 1]$. But we obtain this same number from each of the $n_1(m_1 - p_1)$ partial branches, and thus the whole number is

$$n_1(m_1 - p_1) \frac{m_1}{n_1} [n_1(m_1 - p_1) - 1] = n_1 m_1 [n_1(m_1 - p_1) - 1].$$

Taking account of the other sets, and the whole number of intersections is

$$= \Sigma n^2 a - \Sigma' n a^2.$$

Observe now that $\frac{m_1}{n_1} > \frac{m_2}{n_2}$, that is, $\frac{n_1}{n_2} < \frac{m_1}{m_2}$, and that, these being each < 1 , we have $1 - \frac{m_2}{n_2} > 1 - \frac{m_1}{n_1}$, that is, $\frac{n_2 - m_2}{n_2} > \frac{n_1 - m_1}{n_1}$, and we thus have

$$\frac{m_1}{n_1} < \frac{m_2}{n_2}, \text{ and } \left(z - y^{m/n} x^{1-1/n} \right)^{n_1(m_1-p_1)},$$

Considering now the intersections of partial branches of the two sets

$$\left(z - y^{m/n} x^{1-1/n} \right)^{n_1(m_1-p_1)} \text{ and } \left(z - y^{m/n} x^{1-1/n} \right)^{n_2(m_2-p_2)},$$

respectively, a partial branch $z - z^{m/n} \dots$ gives with each partial branch of the other set a number $= \frac{m_2}{n_2}$; and in this way taking each partial branch of each set, the number is

$$n_1(m_1 - p_1) \cdot n_2(m_2 - p_2) \cdot \frac{m_2}{n_2} = n_2 m_2 n_1 (m_1 - p_1),$$

and thus for all the sets the number is

$$= 2 \Sigma n_2 m_2 \nu_2 m_2,$$

where in the first sum the Σ' refers to each pair of values of the suffixes. But the intersections are to be taken twice; the number thus is

$$= 2 \Sigma m_1 m_2 \nu_1 \nu_2,$$

Adding the foregoing number

$$= \Sigma' n^2 a - \Sigma' n a^2;$$

the whole number for the singularity in question is

$$= (\Sigma m_1)^2 - \Sigma' m_2 \rho_2 - \Sigma' m_2 \rho_2.$$

Similarly for the singularity $\left(y - z^{m/n} x^{1-1/n} \right)^{n-1} \dots$, taking each set with itself, the number of intersections is

$$n_5 m_5 [n_5(m_5 - p_5) - 1] + n_6 m_6 [n_6(m_6 - p_6) - 1] + n_7 m_7 [n_7(m_7 - p_7) - 1],$$

which is

$$= \Sigma' m_2^2 a - \Sigma' m_2 \rho_2 - \Sigma m_1 a.$$

$$[\hspace{10em} 5 - 2]$$

We have here $\frac{m_5}{n_5} > \frac{m_6}{n_6}$, each of these being < 1 , we have $1 - \frac{m_5}{n_5} < 1 - \frac{m_6}{n_6}$, that is, $\frac{n_5 - m_5}{n_5} < \frac{n_6 - m_6}{n_6}$, and so

$$\frac{m_5}{n_5} > \frac{m_6}{n_6}, \frac{n_5 - m_5}{n_5} < \frac{n_6 - m_6}{n_6};$$

Hence considering the two sets

$$\left(y - z^{m/n} x^{1-1/n}\right)^{n_5(m_5-p_5)} \quad \text{and} \quad \left(y - z^{m/n} x^{1-1/n}\right)^{n_6(m_6-p_6)};$$

a partial branch of the first set given with a partial branch of the second set a number of intersections; and the number then obtained is

$$n_5(m_5 - p_5) \cdot n_6(m_6 - p_6) \cdot \frac{m_6}{n_6} = n_6 m_6 n_5(m_5 - p_5),$$

For all the sets the number is

$$n_5 m_5 [n_5(m_5 - p_5) + n_6(m_6 - p_6)] - n_5 n_6(m_6 - m_5)$$

or taking these from, the number is

$$2\Sigma' m_2 m_2 \nu_2 \nu_2.$$

where in the first sum the Σ' refers to each pair of suffixes. Adding the foregoing value

$$\Sigma' m^2 a - \Sigma' m_2 \rho_2 - \Sigma m_1 a,$$

the whole number for the singularity in question is

$$= (\Sigma' m_2)^2 - \Sigma m_1 a - \Sigma' m_2 \rho_2.$$

and the proof is thus completed.

Referring to the first-note (ante, p. 31), I remark that the theorem “ ρ -deficiency, is absolute, and applies to a curve with any singularities whatever: in a curve which has singularities not taken account of in Abel’s theory, the “perhaps see particulars” which we have a give as he expresses it, taken account of, viz. for that Σa is, in the value of ρ as denoted by Abel’s formula; and the actual deficiency will be = Abel’s ρ , = such diminution, that is, it will be = true values of ρ .

.....

714.

VARIOUS NOTES.

[From the *Messenger of Mathematics*, vol. VII. (1878), pp. 69, 115, 124, 125.]

AN IDENTITY.

THE following remarkable identity is given under a slightly different form by Gauss, *Werke*, t. III., p. 419.

$$\begin{aligned} 1 &= \left(\frac{1}{x}\right)^2 x + \left(\frac{1}{1+x}\right)^2 x + \left(\frac{1+x}{1+x+x^2}\right)^2 x + \dots \\ &= \left[1 + \left(\frac{1}{x}\right)^2 x + \left(\frac{1}{1+x}\right)^2 x + \left(\frac{1+x}{1+x+x^2}\right)^2 x + \dots\right]^2. \end{aligned}$$

ON TWO RELATED QUADRIC FUNCTIONS.

Assume

$$\begin{aligned} \varphi x &= x^2(1-x) + x(1-x^2-\alpha x), \\ \varphi y &= y(1-x) + y(x^2-1-\alpha x). \end{aligned}$$

then

$$\begin{aligned} \varphi\left(\frac{x^2}{x-a}\right) &= \frac{x^2}{(x-a)^2} \varphi x, \\ \varphi\left(\frac{x^2}{x-a}\right) &= \frac{x^2}{(x-a)^2} \varphi y. \end{aligned}$$

In the like of those for a ratio $\frac{p}{1-p}$, then

$$\varphi\left[\frac{ax^2(x-y)}{x^2-y^2(x-a)}\right] = \frac{ax^2(x-y)^2}{(x-a)^2-y^2(x-a)^2} \varphi(x).$$

A TRIGONOMETRICAL IDENTITY.

$$\begin{aligned}
 & \cos(b-c) \sin(b+c-a) + \cos a \sin(b+c) \\
 & + \cos(c-a) \sin(c+a+b) + \cos b \sin(c+a) \\
 & + \cos(a-b) \sin(a+b+c) + \cos c \sin(a+b) \\
 = & \cos a \cos b \cos(b+c) + \cos b \cos c \cos(c+a) + \cos c \cos a \cos(a+b) + \cos a \cos b \cos(b+c)
 \end{aligned}$$

EXTRACT FROM A LETTER.

“I wish to construct a correspondence such as

$$(x + iy)^2(x + iy) = X + iY,$$

or, say, for greater convenience

$$4(x + iy)^2(x + iy) = X + iY,$$

viz. if

$$x + iy = \cos \alpha,$$

then

$$X + iY = \cos \alpha \beta.$$

Suppose α , in a value of α corresponding to a given value of $X + iY$, then the three values of $x + iy$ are of course $\cos \alpha$, $\cos(\alpha + \frac{2\pi}{3})$, but I am afraid that the calculation of α , even with red and card tables, would be very laborious. Writing

$$X + iY = R(\cos \Theta + i \sin \Theta),$$

the intervals for Θ might be 3° , 30° or even 15° ; those of R , say 0.1 from 0 to 2, and then 0.2 up to 4; and 2 places of decimals would be quite sufficient; but even this would probably involve a great mass of calculation.

It has occurred to me that perhaps a geometrical solution might be found for the equation $X + iY = \cos \beta$.”

October 21, 1877.

.....

715.

NOTE ON A SYSTEM OF ALGEBRAICAL
EQUATIONS.[From the *Messenger of Mathematics*, vol. VII. (1878), pp. 17, 18.]

ASSUME

$$\begin{aligned}x + y + z &= P, \\yz + zx + xy &= Q, \\xyz &= R.\end{aligned}$$

$$\begin{aligned}A &= (nx + y - z) - a^n(nx + P), \\B &= y(ax + Q) - b^n(my + Q), \\C &= z(ax + Q) - c^n(nz + P), \\\Theta &= -mnQ + nP.\end{aligned}$$

Then

$$\begin{aligned}(ax + P)B - (my + P)A &= (ax + P)y(ax + Q) - a^n(my + Q) \\&= (ax + P)(ax + Q) - (my + Q)(my + P) \\&= ax(my + Q - ay) - yz(my + Q - P) + yz(ax + P) \\&= mny(ax - y) - P(ax - y) \\&= (ax - y)(mny - P) = (ax - y)\Theta,\end{aligned}$$

whence, identically,

$$\begin{aligned}(ax + P)B - (my + P)A &= (ay - y)\Theta, \\(my + P)C - (nx + P)B &= (by - z)\Theta, \\(ax + P)A - (nx + P)C &= (cz - x)\Theta.\end{aligned}$$

Hence any two of the equations $A = 0$, $B = 0$, $C = 0$ imply the third equation.

We have

$$\begin{aligned} A &= x [(n + 1)x + y + z] - a^n [(n + 1)x + y + z] \\ &= (x - a^n) [(n + 1)x + y + z]; \end{aligned}$$

and similarly for B and C . The three equations therefore are

$$\begin{aligned} \frac{x}{x^n - a^n} &= \frac{y + z}{(n + 1)x^n + (n - 1)a^n}, \\ \frac{y}{y^n - b^n} &= \frac{x + z}{(n + 1)y^n + (n - 1)b^n}, \\ \frac{z}{z^n - c^n} &= \frac{x + y}{(n + 1)z^n + (n - 1)c^n}, \end{aligned}$$

and any two of these equations imply the third equation.

.....

716.

AN ILLUSTRATION OF THE THEORY OF THE
 ϑ -FUNCTIONS.

[From the *Messenger of Mathematics*, vol. VII. (1878), pp. 27—32.]

IF X be a given quartic function of x , and if a , as for convenience a constant multiple as, be the value of the integral $\int \frac{dx}{\sqrt{X}}$, taken from a given inferior limit to the superior limit x ; then, conversely, x is expressible as a function of a , viz. it is expressible in terms of ϑ -functions of a , where ϑa , or say $\vartheta(a, \mathfrak{Q})$, is the definition of which, depending, depend in general by definition on the sum of a series of exponentials of a , and it is possible from the assumed equation $a = \int \frac{dx}{\sqrt{X}}$, and the definition of ϑa , to attain by general theory the actual formulae for the determination of a as such a function of a .

I propose here to obtain these formula, in the case where X is a product of real factors, in a less suitable manner, viz. by reducing the integral $\int \frac{dx}{\sqrt{X}}$ by a linear substitution in the form of an elliptic integral, the object being to obtain for the same in question the actual formulae as a function of ϑ -functions of u .

The definition of ϑu , when the parameter is expressed, $\vartheta(u, \mathfrak{Q})$ is

$$\vartheta u = 2\Sigma(-)^p q^{p+\frac{1}{2}(p+1)^2} \sin(p + \frac{1}{2})u,$$

where u and p are negative integer values, zero included, from $-\infty$ to $+\infty$ (that is, from -2 to 2 , 0 to 2 , etc.); the parameter $\mathfrak{Q}$, or Ω' (of imaginary) no real part, must be positive.

[C. XI.

6]

Evidently ϑu is an even function: $\vartheta(-u) = \vartheta u$. Moreover, it is at once seen that we have

$$\vartheta(u + \pi) = \vartheta u, \quad \vartheta(u + i\Pi) = -q^{-1}e^{-iu}\vartheta u,$$

where q is

$$\vartheta(u + \pi + i\Pi).$$

where as u and v are positive or negative integers, the product of ϑu by an exponential factor, or say simply that it is a multiple of ϑu .

Writing $u = \frac{\pi}{2}\xi$, we have $\vartheta v = \frac{1}{2}\sqrt{2K}\vartheta\xi$, that is,

$$\vartheta\left(\frac{\pi}{2}\xi\right) = \xi^{nK},$$

and therefore also

$$\vartheta(\sin \xi + v) = \xi^{nK}\vartheta v.$$

The above properties are general, but if θ be real, then K, K', q being as in Jacobi (consequently q being real, positive, and less than 1, and K and K' real and positive), and assuming $\Omega = \frac{K'}{K}, \pi$, we have in the same thing,

$$q(= e^{-\pi\Omega/\Omega}) = e^{-\pi K'/K},$$

the function ϑv is given (*ut ante*) of Jacobi's ϑv by the equation $\vartheta u = q^{\frac{1}{4}}\vartheta x$.

That is, if Ω is the value of the case $u = v$.

We have, as before, the system of three equations

$$\operatorname{sn} u = \frac{1}{\sqrt{k}} \frac{\vartheta(u, k)}{\vartheta_0(u, k)}, \quad \operatorname{cn} u = \sqrt{\frac{k'}{k}} \frac{\vartheta_0(u, k)}{\vartheta_0(u, k)}, \quad \operatorname{dn} u = \sqrt{k'} \frac{\vartheta_0(u, k)}{\vartheta_0(u, k)}.$$

Consider now the integral

$$\int_a^b \frac{dx}{\sqrt{(x - \alpha, x - \beta, x - \gamma, x - \delta)}} = \int_a^b \frac{dx}{\sqrt{X}} \text{ suppose,}$$

where $\alpha, \beta, \gamma, \delta$ are taken to be real, and the four factors regarded as fixed magnitudes, viz. it is assumed that $b - a$ is the value of the radical at the inferior limit is positive, and as b is between β and δ , X is positive, and *a fortiori* then $\sqrt{X}$ denotes the positive value of the radical; but if α is between β and δ , then X is negative, and we must

that the sign of $\sqrt{X}$ is taken so that $\sqrt{X}$ is equal to a positive multiple of i , and this being so the integral is taken from the inferior limit to the superior limit x , which we may.

Take a linear function of y , such that for

$$x = a, b, c, d,$$

$$y = 0, 1, \frac{1}{k^2}, \infty, \text{ respectively,}$$

so that, x increasing continuously from a to d , y will increase continuously from 0 to ∞ . We have

$$y = \frac{c-a}{c-b} \cdot \frac{x-b}{x-a},$$

$$y = \frac{c-a}{c-d} \cdot \frac{x-d}{x-a},$$

$$1-y = \frac{b-c}{c-b} \cdot \frac{c-a}{c-a},$$

$$1-k'y = \frac{b-c}{c-b} \cdot \frac{c-a}{c-a},$$

and, thence,

$$\sqrt{y \cdot 1-y \cdot 1-k'y} = \frac{(a-c)\sqrt{X}}{(c-a)\sqrt{(x-a) \cdot (x-a)}},$$

where $\sqrt{\frac{(c-d)}{(c-d)}}$ is taken to be positive, and the signs of $\sqrt{X}$ is fixed as above. For y between 0 and 1, $y = \frac{1}{x^2}$, $1-y$, $1-k'y$ will be positive, and $\sqrt{y}$ being between 1 and $\frac{1}{x^2}$, $1-y$ will be negative, but $1-k'y$ will be positive, viz. the radical is then taken as a positive multiple of i .

We have moreover

$$dy = \frac{c-a}{c-b} \cdot \frac{dx}{(x-a)^2},$$

and therefore

$$\frac{dy}{\sqrt{y \cdot 1-y \cdot 1-k'y}} = \sqrt{(c-d) \cdot (c-a)} \cdot \frac{dx}{\sqrt{X}},$$

where $\sqrt{(c-d) \cdot (c-a)}$ is positive; or, say,

$$\int \frac{dy}{\sqrt{y \cdot 1 - y \cdot 1 - k'y}} = \sqrt{(c-d) \cdot (c-a)} \cdot \int \frac{dx}{\sqrt{X}}.$$

[6 - 2]

Hence, writing $y = \sin^2 u$, we have

$$\Pi u = \sqrt{(c-d) \cdot (c-a)} \int_a^b \frac{dx}{\sqrt{X}},$$

and it is to be further noticed that to

$$x = a, b, c, d,$$

correspond

$$\Pi u = 0, \frac{1}{2}K, \frac{1}{2}K + \frac{1}{2}K'.$$

or we may say

$$u = 0, K, K + iK', 2K + iK'.$$

Writing for shortness

$$\frac{1}{\Pi} = \sqrt{(c-d)(c-a)} = u,$$

we have

$$\sin u = \int_a^b \frac{dx}{\sqrt{X}},$$

and moreover

$$\sin(K + iK') = \int_a^b \frac{dx}{\sqrt{X}},$$

$$n(K + iK') = \int_b^c \frac{dx}{\sqrt{X}},$$

$$n(2K + iK') = \int_c^d \frac{dx}{\sqrt{X}},$$

or if for a moment we write

$$\int_a^b \frac{dx}{\sqrt{X}} = A, \text{ \&c.}$$

then these equations are

$$\begin{aligned} nK &= A, \\ n(K + iK') &= A + B, \\ n(2K + iK') &= A + B + C. \end{aligned}$$

Hence $B + C = 2A = B$, or $A = B - C$, or $B - C + D = 0$, that is,

$$\int_a^b \frac{dx}{\sqrt{X}} = \int_b^c \frac{dx}{\sqrt{X}},$$

where observe as before that $x = a$ to $x = b$, or $x = c$ to $x = d$, X is positive, and the radical $\sqrt{X}$ is taken to be positive.

We have also

$$\begin{aligned} nK &= A = \int_a^b \frac{dx}{\sqrt{X}}, \\ niK' &= C - B = \int_a^b \frac{dx}{\sqrt{X}}. \end{aligned}$$

when, as before, from b to c , X is negative, and the sign of the radical is such that $\sqrt{X}$ is a positive multiple of i ; the last formula may be more conveniently written

$$niK' = \int_b^c \frac{dx}{\sqrt{-X}},$$

where, here b to c , $-X$ is positive, and $\sqrt{(-X)}$ is to be taken as positive.

Collecting the results, we have

$$\int_a^b \frac{dx}{\sqrt{X}} = \frac{nK}{\sqrt{(c-d)(c-a)}}, \quad b \leq a \leq b, \dots b \leq b \leq b \leq a,$$

and also

$$k = \frac{c-b}{a-b} \cdot \frac{d-a}{c-a},$$

and thence conversely

$$k' = \frac{a+d-b-c + (b-c-a+d) \sin^2 n}{a+b-c-d + (b-c+a-d) \sin^2 n},$$

or, what is the same thing,

$$\begin{aligned} nu^2 &= \frac{b-d \cdot x-a}{b-a \cdot x-d}, \\ nu^2 &= \frac{c-d \cdot x-a}{c-a \cdot x-d}, \\ nu^2 &= \frac{d-a \cdot x-b}{a-b \cdot x-d}, \\ nu^2 &= \frac{b-d \cdot x-a}{b-a \cdot x-d}, \end{aligned}$$

where, in place of the elliptic functions nu we substitute their ϑ -values, it will be recollected that Ω , the parameter of the ϑ -functions, has the value

$$\frac{K'}{K}\pi = \left(= \frac{\Omega'}{\Omega} \pi \right) = \pi \frac{\int_a^b \frac{dx}{\sqrt{-X}}}{\int_a^b \frac{dx}{\sqrt{X}}},$$

and, so before,

$$K = \int_a^b \frac{dx}{\sqrt{X}},$$

Hence, finally, a, b, K, K', Ω denoting given functions of a, b, c, d , as above

$$\int_a^b \frac{dx}{\sqrt{X}} = nu,$$

we have severally

$$\begin{aligned} \frac{b-d}{c-d} \cdot \frac{a-c}{c-a} &= \frac{1}{k^2} \frac{\vartheta_1^2 u}{\vartheta_2^2 u}, & \frac{\vartheta_3^2 u}{\vartheta_2^2 u}, \\ \frac{d-a}{d-b} \cdot \frac{b-c}{a-c} &= \frac{k'^2}{k} \frac{\vartheta_2^2 u}{\vartheta_3^2 u}, & \frac{\vartheta_1^2 u}{\vartheta_2^2 u} + \frac{1}{2} K', \\ \frac{c-a}{d-a} \cdot \frac{b-d}{b-c} &= k'^2 \frac{\vartheta_4^2 u}{\vartheta_3^2 u}, \end{aligned}$$

which are the formulae in question.

The problem is to obtain these (and this in the more general case where a, b, c, d have any given (real or imaginary) values directly from the assumed equation

$$\int_a^b \frac{dx}{\sqrt{X}} = nu,$$

and from the foregoing definition of the function ϑu .

It may be recalled that the function ϑu is a doubly infinite product

$$\vartheta u = \prod \left[1 + \frac{u}{m\pi + (n + \frac{1}{2})i\Pi} \right]$$

as and a positive or negative integers from $-\infty$ to $+\infty$; I purposely omit all further explanations as to limits, or what is the same thing,

$$\vartheta \frac{\pi}{2K} = \prod \left[1 + \frac{u}{2mK + (2n + 1)iK'} \right]$$

and consequently that, disregarding constant and exponential factors, the foregoing expressions of

$$\frac{b-d}{c-d} \cdot \frac{a-c}{c-a}, \quad \frac{d-a}{d-b} \cdot \frac{b-c}{a-c}, \quad \frac{c-a}{d-a} \cdot \frac{b-d}{b-c},$$

are the squares of the expressions

$$\frac{X}{Y}, \frac{Y}{Z}, \frac{Z}{W}, \quad \text{where } X, Y, Z, W \text{ are respectively of the form}$$

$$\begin{aligned} & n \prod \left[1 + \frac{u}{(2n+1)iK'} \right], \quad \prod \left[1 + \frac{u}{(2m+1)iK'} \right], \\ & \prod \left[1 + \frac{u}{2mK + iK'} \right], \quad \prod \left[1 + \frac{u}{(2m+1)K + iK'} \right], \end{aligned}$$

where $(m, n = 2mK + 2niK', \&c.,$ the *smha* over the m or the n denotes that the $2m$ or the $2n$ as the case may be) is to be changed into $2m+1$ or $2n+1$. But this is a transcription which has apparently no application to the ϑ -functions of more than one variable.

717.

ON THE TRIPLE THETA-FUNCTIONS.

[From the *Messenger of Mathematics*, vol. VII. (1878), pp. 48—50.]

AS a specimen of mathematical notation, viz. of the notation which appears to me the easiest to read, and also to print, I give the definition and demonstration of the fundamental properties of the triple theta-functions.

Definition.

$$\vartheta(U, V, W) = \Sigma \exp \Theta,$$

where

$$\Theta = (A, B, C, F, G, H)(l, m, n)^2 + (U, V, W)(l, m, n).$$

l , denoting the sum in regard to all positive and negative integer values from $-\infty$ to $+\infty$ (zero included) of l, m, n respectively.

$\vartheta(U, V, W)$ is considered as a function of the arguments (U, V, W) , and it depends also on the parameters (A, B, C, F, G, H) .

First Property. $\vartheta(U, V, W) = 0$, for

$$U = \frac{1}{2}\pi i + (A, H, G)(\alpha, \beta, \gamma),$$

$$V = \frac{1}{2}\pi i + (H, B, F)(\alpha, \beta, \gamma),$$

$$W = \frac{1}{2}\pi i + (G, F, C)(\alpha, \beta, \gamma),$$

$\alpha, \beta, \gamma, \delta, \epsilon, \zeta$ being any positive or negative integer numbers, such that as $\alpha + \beta + \gamma = \text{odd number}$.

Demonstration. It is only necessary to show that to each term of ϑ there corresponds a second term, such that the inclusion of the two exponentials differ by an odd multiple of πi .

Taking l, m, n the integers which belong to the one term, those belonging to the other term are

$$-(l + a), -(m + \beta), -(n + \gamma).$$

(where observe that one at least of the numbers α, β, γ being odd, this system of values is not in any case identical with l, m, n). The two exponents are

$$\Theta_i = (A, B, C, F, G, H)(l, m, n)^2 + (U, V, W)(l, m, n),$$

and

$$\Theta'_i = (A, B, C, F, G, H)(l+a, m+\beta, n+\gamma)^2 - (U, V, W)(l+\alpha, m+\beta, n+\gamma),$$

viz. the value of Θ' is

$$\begin{aligned} &= (A, B, C, F, G, H)(l, m, n)^2 + 2(A, H, G)(l\alpha, \beta, \gamma) + 2(H, B, F)(m\alpha, \beta, \\ &+ (A, B, C, F, G, H)(a, \beta, \gamma)^2 - (U, V, W)(l, m, n) - (U, V, W)(\alpha, \beta, \gamma), \end{aligned}$$

and we then have

$$\begin{aligned} \Theta' - \Theta &= -2(A, B, C, F, G, H)(l, m, n)^2 + 2(A, H, G)(\alpha, \beta, \gamma)l + 2(H, \\ &+ 2(G, F, C)(\alpha, \beta, \gamma)n + (A, B, C, F, G, H)(a, \beta, \gamma)^2 \\ &- 2(U, V, W)(l, m, n) - (U, V, W)(\alpha, \beta, \gamma). \end{aligned}$$

which proves the theorem.

As to the notation, remark that, after (A, B, C, F, G, H) has been written out in full, we may instead of $((A, B, C, F, G, H)(\alpha, \beta, \gamma)^2$ write

$$(\alpha, \beta, \gamma)^2,$$

and that we may use the like abbreviation

$$(A, \dots)(\alpha, \beta, \gamma) \text{ for } (A, H, G)(\alpha, \beta, \gamma), \text{ \&c.},$$

$$(H, \dots)(\alpha, \beta, \gamma) \text{ for } (H, B, F)(\alpha, \beta, \gamma), \text{ \&c.},$$

$$(G, \dots)(\alpha, \beta, \gamma) \text{ for } (G, F, C)(\alpha, \beta, \gamma), \text{ \&c.}$$

These are not only abbreviations, but they make the formula actually clearer, as bringing them into a similar compact; and I accordingly use them in the demonstration which follows.

Second Property. If U, V, W , denote

$$\begin{aligned} U' &= \pi i + (A, H, G)(\alpha, \beta, \gamma), \\ V' &= \pi i + (H, B, F)(\alpha, \beta, \gamma), \\ W' &= \pi i + (G, F, C)(\alpha, \beta, \gamma), \end{aligned}$$

respectively, where $\alpha, \beta, \gamma, \delta, \epsilon, \zeta$ are any positive or negative integers (zero values admissible), then

$$\vartheta(U+u, V+v, W+w) = \exp \left[-2(\alpha U + \beta V + \gamma W) - (\alpha, \beta, \gamma)^2 \right] \vartheta(U, V, W),$$

or, say

$$= \exp \left[-(A, \dots)(\alpha, \beta, \gamma)^2 - 2(U, V, W)(\alpha, \beta, \gamma) - (\alpha, \beta, \gamma)^2 \right] \vartheta(U, V, W)$$

Demonstration. Writing $\vartheta(U, V, W) = \Sigma \exp \Theta$, then in the expression of Θ , we may in place of l, m, n write $-l - \alpha, -m - \beta, -n - \gamma$; we thus obtain

$$\begin{aligned} \Theta_l &= (A, \dots)(l, m, n)^2 + 2(A, \dots)(l, m, n)(\alpha, \beta, \gamma) + (A, \dots)(\alpha, \beta, \gamma)^2 \\ &\quad + (U, V, W)(l, m, n) + (U, V, W)(\alpha, \beta, \gamma), \end{aligned}$$

which is

$$\begin{aligned} &= (A, \dots)(l, m, n)^2 \\ &\quad + 2(\alpha, \beta, \gamma)(\alpha U + \beta V + \gamma W) - 2(A, \dots)(\alpha, \beta, \gamma) + \Sigma'(A, \dots)(\alpha, \beta, \gamma)^2 - \Sigma a^2(A, \dots)(\alpha, \beta, \gamma)^2 \\ &\quad - 2(\alpha, \beta, \gamma)^2 + 2\alpha, \beta, \gamma) + (A, \dots)(\alpha, \beta, \gamma)^2 - 2(A, \dots)(\alpha, \beta, \gamma) \\ &\quad + (U, V, W)(l, m, n) + (U, V, W)(\alpha, \beta, \gamma), \end{aligned}$$

which is

$$\begin{aligned} &= (A, \dots)(l, m, n)^2 \\ &\quad + \Sigma'(A, \dots)(\alpha, \beta, \gamma)^2 + \Sigma a^2(A, \dots)(\alpha, \beta, \gamma) + \Sigma \alpha \beta (A, \dots)(\alpha, \beta, \gamma)^2 \\ &\quad + (U, V, W)(l, m, n) + \Sigma(A, \dots)(\alpha, \beta, \gamma) \end{aligned}$$

Hence, rejecting the last line, which (as on can readily verify) has the exponential multiplied as the case may be, $\vartheta(U, V, W)$ as $\vartheta(U, V, W)$ multiplied by the factor

$$\exp \left[-(A, \dots)(\alpha, \beta, \gamma)^2 - 2(U, V, W)(\alpha, \beta, \gamma) - (\alpha, \beta, \gamma)^2 \right],$$

which is the theorem in question.

In many cases a formula, which belongs to an indefinite number of k letters, is most easily intelligible when written out for three letters, but it is sometimes convenient to speak of the letters l , m , ... or even the letter l , and to write out the formula accordingly.

[C. XI.

7]

In fact, writing herein $\alpha = \gamma = 0$, that is, $a = \gamma = 0$, the right-hand side becomes a^2 and the arts on the left-hand side are $a = B$, $A = B$, $a = A = I$, which suppose any four sum the sum of which is $= 0$.

Writing in the last-mentioned equation α, β, γ for the a 's of A, B, γ, δ respectively, the equation becomes

$$\begin{aligned} P &= \beta P, \\ Q &= 1 - a^2 + \beta^2 + By^2, \\ R &= 1 - By^2 - 2\alpha + By^2, \\ S &= 1 - By^2. \end{aligned}$$

L, M, N as the like functions $\alpha, \beta, \gamma, \delta$ respectively, A, B, C, D being the same as before. And in fact

$$\begin{aligned} -LQ &= -y^2(\alpha^2 + \alpha^2) + \beta^2 + (\beta^2 - \alpha^2) + B(y^2 - x^2) + B(\beta - \alpha) + \dots, \\ &= \frac{1}{B} [\beta^2 - (\beta - \alpha)^2 + By^2 - By^2] + B(y^2 - x^2) + \dots, \\ &= \frac{1}{B} [\beta^2(x^2 - y^2) + B(y^2 - x^2) + B^2(y^2 - x^2)], \end{aligned}$$

that is,

$$\begin{aligned} -LPQ &= -y^2(\beta^2 - \alpha^2) + \beta^2 - (\beta - \alpha) + \dots, \\ &+ (B + \alpha - \beta)\beta(y^2 - x^2) + \beta - B^2(y^2 - x^2) + \dots, \end{aligned}$$

or omitting the factor β , that is,

$$-LPQ = -y^2(\beta^2 - \alpha^2) + \beta^2 - y^2(x^2 - \alpha^2) + \dots + \beta(y^2 - x^2) - By^2(x^2 - \alpha^2),$$

It is easy to verify the form of the orders 0, 1, 2, 3, and 4 in x, y, z, w separately destroy each other, for instance, for the terms of the order 4, the value of Σ is then

$$\begin{aligned} &= A^2 + \Sigma y^2(\beta^2 - \alpha^2) + y(x^2 - \alpha^2) + B^2(x^2 - \alpha^2) + (\Sigma \beta^2)(x^2 - \alpha^2) + \dots, \\ &\frac{1}{B} [\Sigma \beta a^2(x^2 + y^2) + \Sigma a^2 B^2(x^2 - y^2) + B^2 a^2(x^2 - y^2)], \end{aligned}$$

It is easy to verify that the terms of the orders 0, 1, 2, 3, and 4 in x, y, z, w separately destroy any sum sum of x which is

$$-LP^2 + yQ^2 + yP^2 + xR^2 + yS^2 + (LP + yQ + xR + yS + yT)$$

$+(D^2-1-LP^2)(yP^2+yP^2-yQ^2-yQ^2)+2(LP+yQ+xR+yS+yT)(LP+yQ+x$
or, omitting the factor β , that is,

$$-(LP+yQ^2+xR+yS+yT)+\Sigma(LP+yQ+xR+yS+yT)(LP+yQ+xR+yS+yT)$$

The theorem in the original form was obtained by me as follows:
using the elliptic coordinates p, q, r , such that

$$\frac{p^2}{a+p} + \frac{q^2}{b+p} + \frac{r^2}{c+p} = 1,$$

or, what is the same thing,

$$-Bp\rho^2 = a + p \cdot b + p \cdot c + p,$$

$$-y\eta y = b + q \cdot a + q \cdot c + q,$$

$$-z\zeta z = c + r \cdot a + r \cdot b + r,$$

where A, B denote $b - c, c - a, a - b$ respectively; then, treating r as a constant, the coordinates x, y, z will belong to a point on the ellipsoid

$$\frac{x^2}{a+r} + \frac{y^2}{b+r} + \frac{z^2}{c+r} = 1,$$

and the differential equation of the right lines upon this surface is

$$\frac{dp}{\sqrt{a+p \cdot b+p \cdot c+p}} + \frac{dq}{\sqrt{a+q \cdot b+q \cdot c+q}} = 0.$$

Take a, b, c , the coordinates of a point on the surface, and p, q , the corresponding values of p, q , so that

$$-Bp_0\rho_0^2 = a + p_0 \cdot b + p_0 \cdot c + p_0,$$

$$-Aq_0\eta_0 = b + q_0 \cdot a + q_0 \cdot c + q_0,$$

$$-z\zeta_0 = c + r \cdot a + r \cdot b + r,$$

then the equation of the tangent plane at the point (a, b, c, p_0, q_0) is

$$\frac{xz_0}{a+r} + \frac{yy_0}{b+r} + \frac{zz_0}{c+r} = 1,$$

or, substituting for x, y_0, z_0 their values, we have

$$= \beta p_0 \cdot ay + \cdot bxz + Az + \dots,$$

and consequently the equation of the tangent plane through (a, b, c, p_0, q_0) is the integral of the proposed differential equation.

Writing

$$mz^2 = A(a+p), \quad mz^2 = B(b+p), \quad dx^2 = C(c+p),$$

the values of A, B, C, S are determined; and, assuming for q, r, p , the like forms with the arguments a, a_0, a_0 , the differential equation becomes $dz + dz$, having the

$$[\quad \quad \quad 10 - 2]$$

integral $u + x_0 + v = u_0$; while the foregoing integral equation, on reducing the constant coefficients contained therein, takes the form

$$\begin{aligned}
 -l^2xx + mz_0mz + nz_0mz_0 + nz_0nz + nz_0nz_0 + \dots &= 0, \\
 &= \frac{1}{2}lm \cdot mz_0mz_0, \\
 &= -l^2nz \cdot mz_0mz_0, \\
 &= \frac{nz_0}{B^2 - 1 - l^2} \cdot \frac{l^2}{l^2 - l^2},
 \end{aligned}$$

via. this equation holds good if $a = u_0 + v$, we have the theorem.

If, as above, $a = u + v + v$, $b = u - v + v$, $c = u + v - v$, where r is regarded as constant, then the three products $\text{sn}u \text{nz} \text{nz}$, $\text{cn}u \text{nz} \text{nz}$, $\text{dn}u \text{nz} \text{nz}$, and equally so the three products $\text{sn}z \text{nz} \text{nz}$, $\text{cn}z \text{nz} \text{nz}$, $\text{dn}z \text{nz} \text{nz}$, are expressed by means of those products with the u , v , w of the u , of the v , viz. for two surfaces it would be possible to obtain a real relation equation, which may be expressed in the squares of the u , v , w of the r , of the z , all of whose internal coefficients are nominal, viz. the such would be obtained, in fact, a similar relation in the like form by giving to the parameter r the advantage of being of a degree which is the half of the equation which involves only the squared functions.

Write a , b , c or for $\text{sn}u$, $\text{cn}u$, $\text{dn}u$, respectively; then, writing

$$\begin{aligned}
 A &= \sqrt{1 - y^2 \cdot 1 - l^2y^2}, & a &= \sqrt{1 - l^2 \cdot 1 - l^2x^2}, \\
 B &= \sqrt{1 - l^2 \cdot 1 - l^2y^2}, & b &= \sqrt{1 - x^2 \cdot 1 - l^2y^2}, \\
 C &= \sqrt{1 - l^2 \cdot 1 - l^2y^2}, & c &= \sqrt{1 - l^2 \cdot 1 - x^2 - y^2}, \\
 D &= 1 - l^2x^2y^2, & d &= 1 - l^2z^2x^2,
 \end{aligned}$$

we have

$$\text{sn}(z + v) = \frac{xA + ya}{D},$$

that is,

$$\frac{A + P}{D} = \frac{\sigma c}{\sigma a \cdot \sigma b},$$

and consequently

$$Dn = (A - x)(A + x),$$

$$PD = (P + x)(P - x);$$

whence

$$Dn = PD = (Aa - x)(Aa - x).$$

[728] (CONCLUDED)

A THEOREM IN ELLIPTIC FUNCTIONS.

that is,

$$(x'^2 - v'^2)(1 - k^2 v'^2 v''^2)(u^2 - y'^2)(1 - k^2 u^2 y'^2) \\ = 2[xw\sqrt{1 - y'^2}\sqrt{1 - k^2 y'^2}\sqrt{1 - v''^2}\sqrt{1 - k^2 v''^2} + yz\sqrt{1 - x'^2}\sqrt{1 - k^2 x'^2}\sqrt{1 - v'^2}\sqrt{1 - k^2 v'^2}]^2.$$

Rationalising, we obtain, as mentioned above, an equation containing only the squares x^2, y^2, z^2, w^2 ; it therefore is of a degree twice that of the equation containing the product $xyzw$. I worked out in this way the equation in (x^2, y^2, z^2, w^2) , but the calculation was lost, and the easier way of obtaining it is obviously by means of the equation involving $xyzw$.

We have, by the theorem,

$$\frac{1 - k^2 \cdot xyzw}{\sqrt{1 - x^2}\sqrt{1 - k^2 x^2}\sqrt{1 - y^2}\sqrt{1 - k^2 y^2}\sqrt{1 - z^2}\sqrt{1 - k^2 z^2}\sqrt{1 - w^2}\sqrt{1 - k^2 w^2}} = \frac{1}{k^2};$$

that is,

$$k^2(1 - k^2 xyzw) = k^2\sqrt{1 - x^2}\sqrt{1 - k^2 x^2}\sqrt{1 - y^2}\sqrt{1 - k^2 y^2}\sqrt{1 - z^2}\sqrt{1 - k^2 z^2}\sqrt{1 - w^2}\sqrt{1 - k^2 w^2},$$

and then, writing

$$\begin{aligned} P &= x^2 + y^2 + z^2 + w^2, \\ Q &= x^2 y^2 + x^2 z^2 + x^2 w^2 + y^2 z^2 + y^2 w^2 + z^2 w^2, \\ R &= x^2 y^2 z^2 + x^2 y^2 w^2 + x^2 z^2 w^2 + y^2 z^2 w^2, \\ S &= x^2 y^2 z^2 w^2, \end{aligned}$$

and using $\sqrt{S}$ to denote the rational function $xyzw$, we have

$$\begin{aligned} &k^2(1 - 2k^2\sqrt{S} + k^4 S) \\ &= k^2(1 - P + Q - R + S) \\ &- 2k^2\sqrt{(1 - P + Q - R + S)(1 - k^2 P + k^4 Q - k^6 R + k^8 S)}; \end{aligned}$$

or, if for a moment the radical is called $\sqrt{\Delta}$, then the factor k^2 divides out, and the equation becomes

$$\sqrt{\Delta} = \frac{P}{2} - \frac{Q}{2} + R - \frac{S}{2} + k^2(-\frac{P}{2} + Q - R + \frac{S}{2}) + \dots,$$

whence

$$\begin{aligned} &4(1 - P + Q - R + S)(1 - k^2 P + k^4 Q - k^6 R + k^8 S) \\ &= \{P - 2 + (1 + k^2)P - 2k^2 Q - (k^2 + k^4)R + 2k^4 S\}^2 - 4k^4 S \\ &= \{-2 + P + k^2 P + \dots - 2k^4 \sqrt{S} \cdot \{2 - (1 + k^2)P + 2k^2 Q - (k^2 + k^4)R + 2k^4 S\}\}. \end{aligned}$$

The factor k^4 divides out; omitting it, we have

$$\begin{aligned} &4Q - P^2 - 4(1 + k^2)R + 16k^2 S + 2k^2 PR - 2(k^2 + k^4)PS - k^4 R^2 + 4k^4 QS \\ &= -2\sqrt{S}\{2 - (1 + k^2)P + 2k^2 Q - (k^2 + k^4)R + 2k^4 S\}, \end{aligned}$$

or, as this may also be written,

$$\begin{aligned} &\{(-P^2 + 4Q - 4R) + k^2(-4P + 2PR + 16S - 4PS) + k^4(-R^2 + 4QS - PS)\} \\ &= -2\sqrt{S}\{2 - P + k^2(-P + 2Q - R) + k^4(-R + 2S)\}, \end{aligned}$$

which is the required rational equation involving the product of the sn's.

729.

ON A THEOREM RELATING TO CONFORMABLE FIGURES.

[From the *Proceedings of the London Mathematical Society*, vol. x. (1879), pp. 143–146.

Read May 8, 1879.]

. Consider two plane figures, say the figure of the points P referred to axes Ox, Oy , and that of the points P' referred to axes Ox', Oy' ; and let x, y be the coordinates of P , and x', y' those of P' . If the figures correspond to each other in any manner whatever, P and P' being corresponding points, then we have x', y' each of them a function of x, y ; and we may consider the second figure as derived from the first by altering the distance OP in the ratio $\sqrt{dx'^2 + dy'^2} : \sqrt{dx^2 + dy^2}$, and by rotating it through the angle $\tan^{-1} \frac{dy'}{dx'} - \tan^{-1} \frac{dy}{dx}$; say the Extension- and the Rotation-ratios.

and by the Rotation $\tan^{-1} \frac{dy'}{dx'} - \tan^{-1} \frac{dy}{dx}$, where the Extension and the Rotation are each of them a determinate function of x, y , the coordinates of P .

Passing from the point P to a consecutive point Q , the coordinates of which are $x + dx, y + dy$ (the ratio $dy : dx$ being arbitrary), then the coordinates of the corresponding point Q' will be $x' + dx', y' + dy'$, where

$$dx' = \frac{dx'}{dx} dx + \frac{dx'}{dy} dy, \quad dy' = \frac{dy'}{dx} dx + \frac{dy'}{dy} dy.$$

Writing $\frac{dx'}{dx}$ and $\frac{dy'}{dx}$ instead of $dx' + dx$ and $dy' + dx$, the equations

$$\sqrt{dx'^2 + dy'^2} = \sqrt{dx^2 + dy^2}, \quad \tan^{-1} \frac{dy'}{dx'} = \tan^{-1} \frac{dy}{dx}$$

will in general have values depending upon that of the arbitrary ratio $dy : dx$. But they may be independent of this ratio: viz. this is the case when x', y' are functions of x, y such that

$$\frac{dx'}{dy} = -\frac{dy'}{dx}, \quad \frac{dx'}{dx} = +\frac{dy'}{dy}.$$

and the two figures are then conformable (or conjugate) figures; that is, figures similar as regards corresponding infinitesimal elements of area. We have, in this case,

$$\sqrt{dx'^2 + dy'^2} = \sqrt{dx^2 + dy^2}, \quad \tan^{-1} \frac{dy'}{dx'} - \tan^{-1} \frac{dy}{dx},$$

each a determinate function of x, y , the coordinates of P ; and we pass from the element PQ to the corresponding element $P'Q'$ by altering the length in the ratio $\sqrt{dx'^2 + dy'^2} : \sqrt{dx^2 + dy^2}$, and rotating the element through the angle $\tan^{-1} \frac{dy'}{dx'} - \tan^{-1} \frac{dy}{dx}$; say, this ratio and this angle are the Aconic and the Strehlonic respectively, these being, as already mentioned, functions of x, y only.

Considering now any two conformable figures, say the figure of the points P , and that of the points P' ; we have the theorem that we can from the first figure obtain a third conformable figure by means of an Aconic and a Strehlonic which are respectively equal to the Extension and the Rotation by which the second figure is derived from the first.

In fact, if in the three figures respectively we take x, y, x', y' , and x'', y'' , for the coordinates of the corresponding points P, P', P'' , the first and second figures are conformable: and we have therefore

$$\frac{dx'}{dy} = -\frac{dy'}{dx}, \quad \frac{dx'}{dx} = \frac{dy'}{dy};$$

the third figure is to have the Aconic $\sqrt{dx'^2 + dy'^2} : \sqrt{dx^2 + dy^2}$, and the Strehlonic

$$\tan^{-1} \frac{dy'}{dx'} - \tan^{-1} \frac{dy}{dx},$$

viz. writing r for $\sqrt{dx'^2 + dy'^2}$, we ought to have

$$dx'' = \frac{dx' + dy' \cdot dy}{r} dx, \quad dy'' = \frac{-dx' \cdot dy + dy' \cdot dx}{r} dx,$$

and it is therefore to be shown that there exist x'', y'' functions of x, y satisfying these relations; for, this being so, we have

$$\frac{dx''}{dy} = -\frac{dy''}{dx}, \quad \frac{dx''}{dx} = \frac{dy''}{dy};$$

and the third figure is then conformable with the first.

Writing, for shortness,

$$A = \frac{x'dx + y'dy}{r^2}, \quad B = \frac{-x'dy + y'dx}{r^2},$$

the equations are

$$dx'' = A dx - B dy, \quad dy'' = B dx + A dy,$$

or the conditions for the existence of the functions x'', y'' are

$$\frac{dA}{dx} + \frac{dB}{dy} = 0, \quad \frac{dA}{dy} - \frac{dB}{dx} = 0.$$

We, in fact, have

$$\begin{aligned} \frac{dA}{dx} + \frac{dB}{dy} &= \frac{1}{r^2} \left[\left(\frac{dx'}{dx} + \frac{dy'}{dy} \right) dx + \left(\frac{dy'}{dx} - \frac{dx'}{dy} \right) dy \right] + 2y \left\{ -\frac{2}{r^4} [(xx' + yy')dx + (xy' - x'y)dy] \right\}, \\ &= \frac{2y}{r^2} - \frac{2}{r^4} (x^2 + y^2)y, \end{aligned}$$

and similarly

$$\begin{aligned} \frac{dA}{dy} - \frac{dB}{dx} &= \frac{1}{r^2} \left[\left(\frac{dx'}{dy} + \frac{dy'}{dx} \right) dx + \left(\frac{dy'}{dy} - \frac{dx'}{dx} \right) dy \right] + 2x \left\{ -\frac{2}{r^4} [(xx' + yy')dy - (xy' - x'y)dx] \right\}, \\ &= \frac{2x}{r^2} - \frac{2}{r^4} (x^2 + y^2)x, \end{aligned}$$

which proves the theorem.

The theorem is closely connected with the theory of the function of an imaginary variable; for, writing the problem for the conformable figures in the form

$$\frac{dx'}{dx} = \frac{dy'}{dy} = F, \quad \frac{dx'}{dy} = -\frac{dy'}{dx} = G,$$

we have

$$dx' = F dx - G dy, \quad dy' = G dx + F dy,$$

that is,

$$dx' + i dy' = (F + iG)(dx + i dy);$$

whence $F + iG$ is a function of $x + iy$, and thus by integration $x' + iy'$ is also a function of $x + iy$. In one point of view, any function such as $\phi(x, y) + i\psi(x, y)$ is a function of $x + iy$, for the quantity $x + iy$ is only known by means of its real components x, y ; that is, knowing $x + iy$, we know x, y , and therefore the real function $\phi(x, y) + i\psi(x, y)$.

and Cauchy, adopting this definition, introduced the expression “function monogène” of $x + iy$, to denote that which is in the more restricted (and the ordinary) sense termed a function of $x + iy$. And MM. Briot and Bouquet, in their “Théorie des fonctions elliptiques” (Paris, 1875), although not using Cauchy’s expression *fonction monogène*, but the simple term *fonction*, do this under the qualification stated p. 8: “*Pour ces fonctions* (or qui sont, sous un autre point de vue, des fonctions qui admettent *une dérivée*).” Now, a function admitting of a derivative (that is, in the ordinary

sense, a function) of the imaginary variable $x + iy$, is a function such that, for a commutative value $x' = x + iy + dx + i dy$, we have a function $f(x') = f(x + iy)$, the condition is rather that $x' + iy'$ may be a function of $x + iy$;

=a quantity independent of the ratio of the real components dx, dy of the increment $dx + i dy$ of the imaginary variable. Or, what is the same thing, writing $f(x) = x' + iy'$, the condition is rather that $x' + iy'$ may be a function of $x + iy$;

$$dx' + i dy' = (F + iG)(dx + i dy);$$

where F and G are functions of x and y . It is not part of the condition that $F + iG$ shall be a function of $x + iy$; nor is it only a long way further on that the authors prove that this is the case (see the definition of a “fonction holomorphe,” p. 14; and the proof, p. 137). The just-mentioned equation

$$dx' + i dy' = (F + iG)(dx + i dy),$$

where F and G are only assumed to be functions of x and y , has of course in the present paper not been used to define x', y' as conformable figures of x, y ; for by means of the point P' with coordinates x', y' , the geometrical interpretation that the figures of the points P and P' are conformable figures, that is, figures similar as regards their infinitesimal elements. The foregoing theorem in regard to the Aconics and the Strehlonics is that we can, by means of F and G , construct a third conformable figure, viz. in fact, knowing $x + iy$ as a function of $F + iG$ and $\tan^{-1} \frac{G}{F}$ respectively; and, using these as an Extension and a Rotation, we have the third conformable figure $x'' + iy'' = (F + iG)(x + iy)$; that is, $(F + iG)(x + iy)$, and therefore also $F + iG$, is a function of $x + iy$ —and we have thus the derivative of a function of $x + iy$ as itself a function of $x + iy$.

It is to be remarked that, although the theorem of the Aconics and the Strehlonics, considered as a property of conformable figures, is now by any means geometrically self-evident, yet the foregoing analytical proof is only a proof conducted by means of real quantities; of when (admitting the theory of imaginary quantities) is in fact self-evident, viz. the analytical conclusion really is that, $F + iG$ denoting any function x, y , then, if $dx' + i dy' = (F + iG)(dx + i dy)$, that is, if $(F + iG)(dx + i dy)$ be a complete differential, then $F + iG$ is a function of $x + iy$.

730.

[ADDITION TO MR SPOTTISWOODE'S PAPER "ON THE TWENTY-ONE COORDINATES OF A CONIC IN SPACE."]

[From the *Proceedings of the London Mathematical Society*, vol. x. (1879), pp. 194–196.]

. WRITE

$$\begin{aligned} U &= (a, b, c, d, f, g, h, l, m, n \mid x, y, z, t)^2, \\ U'_x &= (\quad \mid \xi, \eta, \zeta, \omega)^2, \\ W &= (\quad \mid \xi, \eta, \zeta, \omega)^2, \\ P &= (a, b, c, d, f, g, h, l, m, n \mid \xi, \eta, \zeta, \omega \mid x, y, z, t), \\ P'_x &= (a, b, c, d, f, g, h, l, m, n \mid \xi, \eta, \zeta, \omega). \end{aligned}$$

Then the equation of the cone, having for its vertex the arbitrary point $(\xi, \eta, \zeta, \omega)$, and passing through the conic $U = 0, P = 0$, is

$$UP^2 - 2WP + UW = 0.$$

Or, if, to put the coefficients ξ, η, ζ, ω in evidence, we write for a moment

$$\begin{aligned} A &= (a, b, c, d \mid \xi, \eta, \zeta, \omega), \\ B &= (\quad \mid \xi, \eta, \zeta, \omega), \\ C &= (\quad \mid \xi, \eta, \zeta, \omega), \\ D &= (\quad \mid \xi, \eta, \zeta, \omega), \end{aligned}$$

and further

$$W = A\xi + B\eta + C\zeta + D\omega;$$

then the equation is

$$\begin{aligned} U(A\xi + B\eta + C\zeta + D\omega) - 2P^2(A\xi + B\eta + C\zeta + D\omega \mid Ax + By + Cz + D\omega) \\ + P^2(a, b, c, d, f, g, h, l, m, n \mid \xi, \eta, \zeta, \omega)^2 = 0. \end{aligned}$$

And if we expand first in ξ, η, ζ, ω , and then in x, y, z, t , the final result is

	x^2	y^2	z^2	t^2	yz	zx	xy	xt	yt	zt
$+\xi^2$		C	B	A					h	g
$+\eta^2$	C		A	B	h					f
$+\zeta^2$	B	A		C	g	f				
$+\omega^2$	A	B	C				l	m	n	
$+\eta\zeta$		$3F$	$3E$	$\bullet$	$-2F$	$-2D$	$2E$	$\bullet$	$2(Q - B)$	$\bullet$
$+\zeta\xi$	$3F$		$3D$	$\bullet$	$-2C$	$-2F$	$2D$	$\bullet$	$\bullet$	$2(Q - B)$
$+\xi\eta$	$3E$	$3D$		$\bullet$	$2C$	$-2A$	$-2E$	$2(Q - B)$	$\bullet$	$\bullet$
$+\xi\omega$	$2L$	$3N$	$3M$	$3A$	$-2L$	$-2N$	$2M$	$-2(L - r)$	$-2N$	$-2N$
$+\eta\omega$	$3N$	$2M$	$3L$	$3B$	$2N$	$-2L$	$-2M$	$-2N$	$-2(M - q)$	$-2L$
$+\zeta\omega$	$3M$	$3L$	$2N$	$3C$	$-2M$	$2L$	$-2N$	$-2M$	$-2L$	$-2(N - p)$

In particular, if $a = 0$, $f = 0$, and $w = 0$, we have for the foregoing equation $X = 0$; and the like for the equations $Y = 0$, $Z = 0$, and $W = 0$ respectively.

Take a, b, c, d, e, g, h for the six coordinates of the line through the points

$$\begin{vmatrix} x_1, y_1, z_1, t_1 \\ \xi_1, \eta_1, \zeta_1, \omega_1 \end{vmatrix},$$

then in, write

$$\begin{aligned} a &= y\zeta - \eta z, & l &= xa - \xi a, \\ b &= zx - \xi z, & g &= yb - \eta b, \\ c &= xy - \xi y, & h &= zc - \zeta c, \end{aligned}$$

where, of course,

$$af + bg + ch = 0.$$

Then the foregoing equation of the cone is

$$\begin{aligned} &Ax^2 + By^2 + Cz^2 + Ft^2 + Gz^2 + Hy^2 \\ &-2A^2c - 2B^2a = 3Cab + (B^2a + DA^2)b + 2Cd(b - 2Ad) \\ &\quad + 2Fd^2 + 2Bgh + 2L^2h \\ &-2Eef - 2Egh + 2Hbh + 2Hde = 0. \end{aligned}$$

And this may be regarded as the equation of the cone in terms of the twenty-one coordinates of the conic, and of the six coordinates of an arbitrary line meeting the cone. It is, in fact, the general form of the equation given in the paper—Cayley, “On a new Analytical Representation of a Curve in Space,” *Quart. Math. Jour.*, vol. III. (1860), [284, this Collection, vol. IV. p. 452].

731.

ON THE BINOMIAL EQUATION $x^p - 1 = 0$; TRISECTION AND QUARTISECTION.

[From the *Proceedings of the London Mathematical Society*, vol. xi. (1880), pp. 4–17.

Read November 13, 1879.]

. THE solution of the binomial equation $x^p - 1 = 0$, p a prime number, or, what is the same, of the equation

$$x^{p-1} + x^{p-2} + \cdots + x + 1 = 0,$$

depends upon the Jacobian function

$$F_n x = x + ax^n + a^2 x^{n^2} + \cdots + a^{p-2} x^{n^{p-2}},$$

where a is a prime root of p , n any root whatever of the equation $x^{p-1} - 1 = 0$. Taking a a factor of $p - 1$, and f for the supplementary factor (that is, $p - 1 = af$), then, if n be a root of a , that is, root of $x^a - 1 = 0$, we have

$$F\eta = X_0 + \eta X_1 + \cdots + \eta^{a-1} X_{a-1},$$

where $X_0, X_1, \dots, X_{a-1}$ denote each of them a period of f terms, viz.

$$X_k = x^{n^k} + x^{n^{k+a}} + x^{n^{k+2a}} + \cdots + x^{n^{k+(f-1)a}}.$$

$$X_k = -1, \quad \eta^a \dots \eta^{a(p-1)} X_k = -1, \quad \xi^a \dots \xi^{a(p-1)}.$$

(mod p); the values being roots of $a^a = 1$, and so for the other functions).

We have, of course, $F(1) = X_0 + X_1 + \cdots + X_{a-1}$, the sum of all the roots $= -1$; and, further, the general property that there is (independently of the value of a chosen periods is expressible as a sum

$$a_1 X_1 + a_2 X_2 + \cdots + a_{a-1} X_{a-1},$$

with known coefficients

$$a_1, a_2, \dots, a_{a-1}.$$

The several cases $a = 2, 3, 4, \dots$ may be termed those of the bisection, trisection, quartisection, &c., of the equation; viz.

$a = 2$, there are two periods, X, Y , and $F(= 1 + x + X + Y) = -1$;

$a = 3$, three periods, X, Y, Z , and $F\eta = X + \eta Y + \eta^2 Z$, η a root of $x^3 - 1 = 0$;

$a = 4$, four periods, X, Y, Z, W , and $F\theta = X + \theta Y + \theta^2 Z + \theta^3 W$, if θ be a root of $x^4 - 1 = 0$.

It is sufficient to attend to the prime roots which lie within the period p , and call these roots, $\eta_1, \eta_2, \dots, \eta_{p-1}$ in respectively; and the relation between them is determined respectively, viz. if η be a b -th, we have always $F(\eta) = -1$; and if b be the of the periods for the bisection. The prime roots b are of course 1 and -1 ; and we have

$$F(1) = X + Y - 1 = -1,$$

$$F(-1) = X - Y = \sqrt{\pm p},$$

respectively.

As regards the bisection, it is known that $(X - Y)^2 = (-1)^{(p-1)/2}p$, which is $+p$ or $-p$, according as p is $\equiv 1$ or $3 \pmod{4}$; and the values of X, Y are thus determined. In what follows, I assume the case $p \equiv 1 \pmod{4}$, for which X, Y are thus determined, X, Y are the values $\frac{1}{2}(-1 \pm \sqrt{p})$.

It is to be remembered that, not the division into periods, but the order of the periods, depends on the choice of g , a prime root or pseudoprime of p ; and, in what follows, I submit the prime root used to determine *Jacobi's (J)ables canoniques Primitives suivies can Wurzeln der Einheit gebildet sind* (8vo, Berlin, 1872); and I have accordingly herein taken these as $\frac{1}{2}$ -roots of Reuschle's selected primes.

For instance, $p = 13$, the powers of g are 1, 2, 4, 8, 3, 6, 12, 11, 9, 5, 10, 7; and, by writing these down in order in columns of 3 or 4, viz.

1	8	12	1	8	12	5
2	3	11	2	3	11	10
4	6	9	4	6	9	7

we have the periods X, Y, Z as X, Y, Z, W , belonging to the trisection and the quartisection of $p = 13$.

I further remark that the equations which I am concerned with are all given in Reuschle, but in a somewhat different form; thus, $p = 13$, quartisection (see p. 10), he has

$$\eta^4 = -\eta^3 - 2\eta + 3, \quad \eta_1\eta_2 = -3\eta_1 - \eta_2 + 3, \quad \eta_1\eta_3 = -1 - \eta_3,$$

(where observe that here and in every case the value of $\eta_1\eta_3$ is at once obtained from that of $\eta_1\eta_2$ by mere cyclical interchange of the suffixes, so that the last equation is in fact superfluous), the other equations, using $\eta_1 + \eta_2 + \eta_3 + \eta_4 = -1$, can be eliminated; the result is system of equations, $X + YZ = 0$, $XY + Z = 0$, &c.

Similarly, in the case of a trisection, the equation for $\eta_1\eta_2$ is superfluous, and the other equations give my values of X^2 and XY .

Reuschle gives also, and I take from him, the cubic and the quartic equations satisfied by $\eta, = 1, b, \dots, n$ for the various periods of those obtained for $X, Y, Z, \dots$ and connections with these periods is at once seen the rule of formation of those periods in the trisections and the quartisections respectively.

Many of the results obtained accord with, and furnish exemplifications of general theorems contained in Jacobi's memoir, "Ueber die Kreistheilung und ihre Anwendung auf die Zahlentheorie," *Crelle*, t. XXX. (1846), pp. 166–182; [*Ges. Werke*, t. VI. pp. 254–274].

Trisection, $a = 3, p \equiv 1 \pmod{3}$.

We have there periods X, Y, Z , and we choose

$$XY = \frac{1}{3}(p-1) + aX + bY + cZ,$$

$$XY = (f, g, h, k)$$

the coefficients a, b, c, f, g, h being determinate integers. And, by cyclical interchanges, we obtain equations which may be written

$$X^2 = a, b, c,$$

$$Y^2 = b, c, a,$$

$$Z^2 = c, a, b,$$

$$XY = f, g, h,$$

$$YZ = g, h, f,$$

$$ZX = h, f, g,$$

viz. here and elsewhere the coefficients a, b, c are written to denote the sum

$$aX + bY + cZ.$$

It is easy to see that

$$f + g + h = (p-1).$$

in fact, multiplying $X^2 + Y^2 + Z^2$ together, the coefficient of X is $a + c + b$, and there are no terms the product of which is unity; hence XY contains $\frac{1}{3}(p-1)$ unit terms, each a power of g , and this sum is XY . From the equation $X + Y + Z = -1$, multiplying by X , and for XY , XZ substituting their values, we obtain an expression

$$(a + f + h)X + (b + f + g)Y + (c + g + h)Z = -\frac{1}{3}(p-1)X,$$

which must identically vanish; viz. the three coefficients must be each of them $= 0$; we must have

$$\begin{aligned} a + f + h &= -p, \\ b + f + g &= 0, \\ c + g + h &= 0, \end{aligned}$$

so that, taking f, g, h as known, the other coefficients a, b, c are given in terms of them. The equations give

$$\begin{aligned} X^2 &= YZ + X(\frac{1}{3}p + g + h), \\ Y^2 &= ZX + Y(\frac{1}{3}p + h + f), \\ Z^2 &= XY + Z(\frac{1}{3}p + f + g). \end{aligned}$$

We have $X, YZ = Y, XZ = Z, XY, (p-1)$, so that, substituting for X^2, XY , etc., their values,

$$\begin{aligned} ah + bk + ck &= -p(a + b + c) + 4(f + g + h)k, \\ af(\eta + h) + 4 + h(\eta + f) &= (4 + h + f) + 4 + f(\eta + g), \\ ag(g + f) + g(g + f) &= (\eta + g + f)k + g(\eta + g), \end{aligned}$$

that is,

$$\begin{aligned} ah + bk + ck &= pf + pg + 2k(k + l), \\ bh + g(g + h) &= ah + g(h + l) = ah + g(b + l), \\ gh + g(g + l) &= (g + l + h)k + g(g + l), \end{aligned}$$

equations which reduce themselves to the single equation

$$gh + h^2 + g^2 = -\frac{1}{3}(p-1) - \frac{1}{3}(g+h) - \frac{1}{3} - \frac{1}{3}(g-h).$$

and this is the only relation obtainable by consideration of the three equal values

$$X, YZ, \quad Y, ZX, \quad Z, XY.$$

Moreover, this equation being satisfied, the six functions in the three equations because each of them $= fg - fg$; so we have, from the three equations becomes

$$XYZ = (fg - fg)/p - h^2,$$

that is,

$$XYZ = fp - h^2.$$

We have

$$\begin{aligned} F_p \cdot F_p^2 &= X^2 + Y^2Z + \eta YZ \cdot YZ - X \cdot XY \\ &= (\eta + \eta + \eta) - b\eta - c)(X + Y + Z) \\ &= -\eta + \eta + \eta - b\eta - c \\ &= 3(f + g + h) \end{aligned}$$

that is,

$$F_\eta F_\eta^2 = p.$$

We have, moreover,

$$\begin{aligned} (F_\eta)^3 &= X^3 + Y^3 + Z^3 + \eta \cdot 3X^2Y + \eta^2 \cdot 3XY^2 + (3X^2Z + \eta + \eta^2) \\ &= [\eta\eta'\eta'', a + b\eta + \eta'\eta'(X - Y) + (Y - Z) + (Z - X)XY\eta\eta', + 3XYZ]; \end{aligned}$$

which is

$$= \frac{1}{2}(a + 4b - 4c + (-3a + 4b + c)\eta + (3a + b + 4c)\eta^2),$$

as is at once seen by comparing the coefficients of X, Y, Z respectively.

Hence, writing

$$\begin{aligned} a + b(\eta + \eta'') + c(\eta'' + \eta) &= a^2 + b^2 + c^2 - bc(\eta - 1) - (-1) \\ &= a + 2b - c, \end{aligned}$$

we have

$$A = a + 2b - c, \quad B = a + 2c - b, \quad C = b - c;$$

η a root of $\eta^2 + \eta + 1 = 0$.

We have also,

$$(F_\eta)^3 = (a + b\eta + c\eta'')X + (a + b\eta'' + c\eta)Y + (a + b\eta + c\eta'')Z,$$

and these together

$$= (Ap + Bq)/p,$$

equations which reduce themselves to the single equation

$$gh + h^2 + g^2 = -\frac{1}{3}(p - 1)$$

or, say

$$(F_\eta)^3 = p(F_\eta)/F_\eta;$$

viz. here it has, $X, YZ = Y, XZ = Z, XY, (p - 1)$, so that, substituting for X^2, XY , etc., the equation becomes

$$XYZ = (Ap + Bq)/p - h^2,$$

or the expression $A + Bi$ determined as above is a complex factor of p .

As already remarked, the values of a, b, c, f, g, h ; and the equations in a, b in effect given in Reuschle; the complete factors of p , as given in p. 9, are $4p = a^2 + 27b^2$, where reduced to the form $A + Bi$, are not identical with the $A + Bi$ found from the foregoing theory.

A and B being respectively

$$\begin{aligned} a^2 - bc &= \frac{1}{2}(p - 1) - (f - g) \cdot a^2 - bc(\eta - 1) - (-1) \\ &= a + 2b - c. \end{aligned}$$

And the A and B in Reuschle's selected primes; the

for the primes 7, 13, . . . , 97, the values from Reuschle of a, b, c, f, g, h , and of the coefficients of the η -equation; also the values of A and B derived from f, g, h by the foregoing formulæ. It will be seen that all the values are consistent with the theory.

TABLE FOR THE TRISECTION.

p	a, f	b, g	c, h	$a^2 + 27b^2$	A	B	Page in Reuschle	
7	−2, 1	−1, 1	−1, 2	−2, 1	1	2	1	p. 5
13	−4, 3	−3, 2	0, 4	−4, 3	−4	−3	0	p. 15
19	−4, 3	−1, 3	1, 7	−6, 7	−7	−3	3	p. 16
31	−7, 5	−1, 5	1, 7	−10, 1	−2	5	−2	p. 18
37	−8, 4	−2, 4	4, 3	−12, 11	−4	3	−5	p. 34
43	−4, 3	−10, 7	−7, 4	−12, 15	−4	3	0	p. 69
61	−11, 9	−12, 11	−11, 11	−20, 33	−9	8	−1	p. 97
67	−16, 11	−2, 3	−7, 8	−22, 17	5	2	9	p. 107
73	−16, 11	−12, 7	−4, 2	−24, 27	−7	−9	9	p. 128
79	−20, 17	−16, 11	−26, 21	−26, 33	−10	−3	9	p. 154
97	−29, 21	−26, 21	−36, 29	−32, 79	11	9	4	p. 187

Quartisection, $a = 4$, $p \equiv 1 \pmod{4}$.

We have four periods X, Y, Z, W ; and we obtain

$$\begin{aligned} X^2 &= (a, b, c, d \mid X, Y, Z, W), \\ XY^2 &= (f, g, h, k), \\ XZ &= (l, m, n, p). \end{aligned}$$

the coefficients being determinate integers. It can be shown that $l = m = \frac{1}{4}(p-1)$, and $n = \frac{1}{4}(p+1)$ according as $p \equiv 1$ or $5 \pmod{8}$. And then, by cyclical interchanges,

$$\begin{aligned} X^2 &= a, b, c, d, \\ Y^2 &= b, c, d, a, \\ Z^2 &= c, d, a, b, \\ W^2 &= d, a, b, c, \\ XY &= f, g, h, k, \\ YZ &= g, h, k, f, \\ ZW &= h, k, f, g, \\ WX &= k, f, g, h, \\ XZ &= l, m, n, p, \\ YW &= m, n, p, l. \end{aligned}$$

We have, as in the manner as for the trisection,

$$f + g + h + k = \frac{1}{4}(p-1),$$

and so also the equation

$$2(X + Y + Z + W) = XY + YZ + ZW + WX + XZ + YW + ZW,$$

and, in virtue of the foregoing value of $l + m = -\frac{1}{4}(p-1)$ or $\frac{1}{4}(p-5)$ according as $p \equiv 1$ or $5 \pmod{8}$.

Again, from the equation $X + Y + Z + W = -1$, multiplying by X and reducing, and thence

$$\begin{aligned} a + b + c + d &= -2(f + g + h + k) - 2(l + m), \\ &= -\frac{1}{2}(p-1) - 2(l + m). \end{aligned}$$

We have

$$XYZ = YXZ = ZXY,$$

that is,

$$X(g, h, k, f) = Y(l, m, n, p) = Z(f, g, h, k),$$

and thence

$$\begin{aligned} g(a, b, c, d) + h(f, g, h, k) + k(g, h, k, f) + f(l, m, n, p) = \\ l(b, c, d, a) + m(g, h, k, f) + n(c, d, a, b) + p(h, k, f, g) = \\ f(c, d, a, b) + g(h, k, f, g) + h(l, m, n, p) + k(d, a, b, c) = \end{aligned}$$

that is,

$$\begin{aligned} ka + ga + h^2 + k^2 + gh + (f + l)a + (g + p)b + (l + g)c + (k + f)d = 0, \\ hb + ka + h^2 + kf + gh + (f + l)a + (h + l)b + (m + n)c + (k + l)d = 0, \\ lc + ld + hf + kg + (h + l + g)a + (g + l)c + gh + (l + g)c + (m + l)d = 0, \\ ld + ld + hf + kg + (g + l + h)a + g(g + l) + (k + l)d + g(g + l) = 0, \end{aligned}$$

in which equations a, b, c, d may be regarded as having their foregoing values.

One of these equations is

$$hc + fa + gd + bd + (a + g)c + (f + p)d = bf,$$

that is,

$$-b + h + a + l + b(a + g) + p(g + p) + cl = -p(f + p) + p(g + p) + d,$$

or, reducing,

$$l(g + d - 1) - f - p = (a + b)c - 2bp + g(d + b) - p(g + p) + dp,$$

which gives

$$l(g + d - 1) - f - p = (a + b)c - 2bp + g(d + b) - p(g + p) + dp - p^2,$$

Again, another equation is

$$hb + g(g + l) + p(g + p) + bp = lp,$$

that is,

$$-h(g + l + n) + l(l + p) + dl + b(l + p) = a(l + p) + lp + bp,$$

or, reducing,

$$m(g + k - l) = ld + h(l + p) + d - (l + p),$$

which gives

$$m(g + k - l) = ld + h(l + p) + d - (l + p),$$

And we have also

$$md + l + n = a + ld + h + h(d + p) - 2p,$$

that is,

$$= m(a + b + n) + h(g + b) + b(g + a) + p(g + p) + (b + p) + l,$$

or, reducing,

$$b(a + l + p) + b(a + p) + (a + p)l = (a + p)l + (a + p)b + (a + p) + p(g + p) + (a + p)d,$$

Substituting herein for b, c , their values, we have

$$(b + l)(2f + p - 28f - lp + k(f + 1)(2g - p - lp + h(g + 1)(2f + p - lp + h(g + 1)(2p - p - lp) + (k - p)(g + p + 2(p - p)) + (k - p)(g + p + 2(p - p)))$$

In this equation the only terms of the second order are $-(b+2p)$, which contains the factor b ; the terms of the third order contain this same factor b , and throwing it out, and reducing, the equation is found to be

$$(g-h)^2 + 4l = -f^2 - h - k - p,$$

or, as it may also be written,

$$g^2 + h^2 - 2gh - 1 + (\frac{1}{2}h^2 - 1 + 2 - 2p - h - kp) = 0,$$

and the foregoing values of l, m are

$$\frac{1}{2}(g+h-2k+2-2p)(f-g) - (gh-lp) = -m = \frac{1}{2}(g+h-2k+2-2p)(f-g) - (gh-lp) =$$

and by means of these equations all the foregoing equations are satisfied.

We have

$$\begin{aligned} F(\eta) &= (X-Y) + i(\eta(Y-W)), \\ &= X-Y + i(\eta(Y-W)) - i(2(XY+YZ)-YW), \\ &= (-1+a+b+a) + i(\eta b), \end{aligned}$$

or, substituting for a, b, c, d , and for f, g, h, k in evidence, we write

$$F(\eta) = (\alpha)^{p-1} p.$$

Again, we have

$$\begin{aligned} (F(\eta))^2 &= (X-W)^2 + 2(\eta(Y-W)), \\ &= X^2 - Y^2 - Z^2 - W^2 + 2(XY - YZ + ZW + XY) - WX + 2i(2XY - ZW - WX) + WX, \\ &= (a-b-c-d) + 2(f+g-h+k) - l + m - 2(2(p+2h+2(g-h) - 2(g+l)l)), \\ &= (a-b)X^2 + 2(2k-2l+h(g-h))(g-h), \end{aligned}$$

where

$$\begin{aligned} A &= a+b-c+d - (b-c) + d + 2(f-g-h+k) - 2(l-m), \\ B &= 2(f-g+h-k); \end{aligned}$$

or, that $X-Y-W = (a+pF(\eta)) - g + (a+h(g-h))$.

The function $A^2 + Bi$ is here a complex factor of p , viz. we have

$$F(\eta)F'(\eta) = (A+B)^2(A-B^2) = A^2 + B^2,$$

and similarly

$$(F'(\eta))F(\eta) = (A-B)^2(F(\eta)) - 1.$$

Moreover

$$F'(-1)(-1) = (-)^{(p-1)/4} p.$$

and we have therefore

$$(F'(-1))^2 = (A^2 + B^2) p,$$

that is,

$$A^2 + B^2 = p,$$

or the expression $A + Bi$ determined as above is a complex factor of p .

Taking the first expression of those above written for A, B , we have

$$\begin{aligned} a + b + c + d &= -\frac{1}{2}(p-1) - 2(l+m), \\ +A &= a - b + c - d, \\ &= 2a + 2c + \frac{1}{2}(p-1) + 2(l+m), \\ &= 2(a+c+l+m) + \frac{1}{2}(p-1). \end{aligned}$$

We had A, YZ writes equally as the product of XY and of XW , viz. we obtain

$$\begin{aligned} XYZW &= a + b(g, h, k, f) + c(b, c, d, a) + d(g, h, k, f) \\ &= a + b + b - h - k - f - (f+l)a + (g+p)b + (l+g)c + (k+f)d, \\ &= 2(f+g)h(a+b+d) + 2(f+g)l = 2g(g+p)k - 2g(a+p)d - g\eta + h(b+c+d), \\ &= 2(f+g)l(b+d) - p(g+p) + (g+p)d. \end{aligned}$$

We had $XYZW$ unable as the product of XY and of ZW , viz. we obtain

$$\begin{aligned} XYZW &= (f, g, h, k)(b, c, d, a) \\ &= f(b+l) + d(b+l)(b+d) - (b+l+d)(f+l) + l(b+g)d, \\ &= (f+l)k - g - 2bl + (a+b) - (b+l+d)l + d(b+l+d) + lp, \\ &= (f+l)(k-l+p) + d(b+p)l - (b+l+d)l + d(b+l+d) - lp, \\ &= (f+l)(k-p) + d(g-l+p) - p(l+b+p)l, \end{aligned}$$

that is,

$$XYZW = (f+l)(k-p) - bl + d(b+p)d + 2, (p+l+p)(b+d),$$

where, for the sake of having a single formula, I have retained l in place of the value $-\frac{1}{2}(p-1+l) - \frac{1}{2}(p-8)$ according as $p \equiv 1$ or $5 \pmod{8}$.

We then have the following.

TABLE FOR THE QUARTISECTION.

p	a	b	c	d	f	g	h	k	l	m	A	B	Page in Reuschle
5	0	1	1	0	0	1	1	0	−1	−1	1	−2	p. 4
13	0	1	0	0	1	0	1	1	−1	0	2	−3	p. 5
17	4	−2	−2	0	0	1	2	1	−1	−1	−2	2	p. 12
29	4	4	−3	−2	1	4	4	1	−1	−2	−7	−1	p. 33
37	0	5	4	4	2	4	5	2	−2	−3	−3	4	p. 84
41	−10	−5	−2	5	4	1	1	4	−2	−3	−1	−2	p. 86
53	0	7	4	4	5	4	7	5	−2	−2	5	5	p. 99
61	4	4	−7	−12	5	7	7	5	−1	−3	−5	−7	p. 106
73	−10	−16	−12	0	7	4	5	7	−3	−2	7	−4	p. 120
89	−20	−19	−13	−11	7	5	5	7	−3	−4	−5	−8	p. 151
97	−22	−16	−17	−19	10	5	5	5	−3	−4	9	−4	p. 187

[Figure: large table of computed values, transcribed above.]

TABLE OF THE POWERS OF REUSCHLE'S SELECTED PRIME ROOTS.

[Figure: a large arithmetical table 65 columns wide listing powers $g^k \pmod{p}$ for selected primes $p = 5, 7, 11, 13, 17, 19, 23, 29, 31, 37, 41, 43, 47, 53, 59, 61, 67, 71, 73, 79, 83, 89, 97$ across the columns, and exponents $k = 1, 2, 3, \dots$ down the rows. The cells of this table contain entries such as $1, 2, 4, 8, 16, \dots$ for $p = 17, g = 3$, etc. The full table fills the page.]

TABLE (CONTINUED).

[Figure: continuation of the table of powers of Reuschle's selected prime roots, spanning further values of the exponent k for the primes listed on the preceding page. Numerous numerical entries fill the page, in the same format as the previous table.]

732.

A THEOREM IN SPHERICAL TRIGONOMETRY.

[From the *Proceedings of the London Mathematical Society*, vol. xi. (1880), pp. 48–50.
Read January 8, 1880.]

. In a spherical triangle, where a, b, c are the sides, and A, B, C their opposite angles, we have

$$\begin{aligned}\tan \frac{1}{2}a \tan \frac{1}{2}c &= \tan \frac{1}{2}b \sin(A - B) + \tan \frac{1}{2}b \sin A - \tan \frac{1}{2}a \sin B, \\ \tan \frac{1}{2}c (1 - \tan \frac{1}{2}a \tan \frac{1}{2}b \cos(A - B)) &= \tan \frac{1}{2}b \cos A + \tan \frac{1}{2}a \cos B,\end{aligned}$$

which are both included in the form

$$\tan \frac{1}{2}c \cos B = \frac{\tan \frac{1}{2}b \sin(A - B) (\cos A + i \sin A)}{1 + \tan \frac{1}{2}a \tan \frac{1}{2}b (\cos A + i \sin A)}.$$

For the first of the two identities: from

$$\begin{aligned}\cos a &= \frac{\cos A + \cos B \cos C}{\sin B \sin C}, \\ \cos b &= \frac{\cos B + \cos C \cos A}{\sin C \sin A},\end{aligned}$$

we deduce

$$\begin{aligned}\cos a - \cos b &= \frac{1}{\sin A} \left(\frac{\cos A}{\sin C \cos B} - \frac{\cos B}{\sin C \cos A} \right) - \frac{1}{\sin B} \left(\frac{\cos B}{\sin C \cos A} - \frac{\cos A}{\sin C \cos B} \right), \\ &= \frac{1}{\sin A} \frac{(\sin A \sin B - \sin B \sin A) (\cos(A - B) - \cos A \sin B)}{\sin C \sin A \sin B}, \\ &= \frac{\sin(A - B)}{\sin A \sin B \sin C} (\cos A \sin B - \cos B \sin A), \\ &= -\frac{\sin(A - B)}{\sin C} (\cos A \sin B - \cos B \sin A), \\ &= \frac{\sin(A - B)}{\sin C} (\cos(B + A) - 1),\end{aligned}$$

that is,

$$-\sin(A - B) = \frac{\sin C}{\sin a \sin b} (\cos a \sin b - \cos b \sin a);$$

or, what is the same thing,

$$-\tan \frac{1}{2}a \tan \frac{1}{2}b \sin(A - B) = \frac{\sin C}{\sin a \sin b} (\cos a \sin b - \cos b \sin a),$$

Here $\cos a - \cos b = (1 + \cos a) - (1 + \cos b)$; substituting for $\frac{\sin a}{\sin b}$ correspondingly $\frac{\sin A}{\sin B}$ and $\frac{\sin B}{\sin A}$, the right-hand side is

$$\begin{aligned} &= \frac{1 + \cos a}{\sin a \sin A} - \frac{1 + \cos b}{\sin b \sin B}, \\ &= \cot \frac{1}{2}a - \cot \frac{1}{2}b, \end{aligned}$$

whence, multiplying each side by $\tan \frac{1}{2}a \tan \frac{1}{2}b$, we have the relation in question.

For the second identity which is

$$\tan \frac{1}{2}c [1 - \tan \frac{1}{2}a \tan \frac{1}{2}b \cos(A - B)] = \tan \frac{1}{2}b \cos A + \tan \frac{1}{2}a \cos B;$$

if on the right-hand side we substitute for $\sin A$, $\sin B$ their values

$$\frac{\cos a + \cos b \cos c}{\sin a \sin b} \quad \text{and} \quad \frac{\cos b + \cos c \cos a}{\sin b \sin a},$$

the right-hand side becomes

$$\frac{1}{\sin C} [\cos a - \cos b + \cos C (\cos a \cos b - \cos b \cos a)],$$

whence, multiplying the whole equation by $\sin C$, it becomes

$$\begin{aligned} (1 - \cos c)(1 + \cos a)(1 - \cos b) &= \tan \frac{1}{2}b \cos(A - B) \\ &= (1 - \cos c)(\cos a - \cos b) + (1 + \cos c)(\sin a \cos b - \cos a \sin b). \end{aligned}$$

We have here

$$\cos(A - B) = \frac{\cos a \sin b \cos B + \sin a \cos B}{\sin a \sin b};$$

by substituting for $\sin a$ and $\sin b$ their values, and so for A , B their values

$$\frac{\cos b}{\sin c \cos a + \cos c} \quad \text{and} \quad \text{etc.} \quad \text{where}$$

$$\cos a + \cos b + \cos C = -\cos a + \cos b + \cos a \cos b + \cos C,$$

or

$$\square = 1 - \cos^2 a - \cos^2 b - \cos^2 c + 2 \cos a \cos b \cos c.$$

The numerator is

$$\begin{aligned} & \cos C \sin b - \cos(\cos^2 a + \cos^2 c) \sin a \sin b + \cos a \sin b \\ &= 1 - \cos^2 c - (\cos^2 a + \cos^2 c) \sin a \sin b + \cos a \sin b + 2 \cos a; \end{aligned}$$

viz. this is

$$= \cos a \sin b (1 + \cos a) - (\cos^2 a + \cos^2 c)(1 + \cos a)(1 - \cos a) - 1 - \cos a;$$

having the factor $1 + \cos a$, which is also a factor of $\sin^2 a = 1 - \cos^2 a$, in the denominator. We have, therefore,

$$\cos(A - B) = \frac{\cos a \sin b \cos(1 + \cos a) - (\cos^2 a + \cos^2 c) \sin a \cos b - 1 - \cos a}{(1 - \cos a) \sin a \sin b},$$

and the equation thus is

$$\begin{aligned} & (1 - \cos c)(1 + \cos a)(1 - \cos b) + \cos a \sin b \cos(1 + \cos a) - (\cos^2 a + \cos^2 c) \sin a \cos b - 1 - \cos a \\ &= (1 - \cos c)(\cos a - \cos b) + (1 + \cos c)(\sin a \cos b - \cos a \sin b), \end{aligned}$$

where each side is in fact

$$= \cos a \sin b (1 + \cos c) + \cos a \sin b (1 + \cos c) - \square = 1 - \cos c \sin b;$$

and the second identity is thus proved.

733.

ON A FORMULA OF ELIMINATION.

[From the *Proceedings of the London Mathematical Society*, vol. xi. (1880), pp. 139–141.

Read June 10, 1880.]

. CONSIDER the equations

$$(a, \dots | \xi, \eta)^p = 0, \quad (A, \dots | \xi, \eta)^q = 0,$$

where $a, \dots, A, \dots$ are functions of coordinates. To fix the ideas, suppose that each of these coefficients is a linear function of the four coordinates x, y, z, w . Then, eliminating θ , we obtain $\nabla = 0$, the equation of a nodal-surface (the form of ∇ is known); this surface has a nodal curve.

It is easy to obtain the equations of the nodal curve in the case where one of the equations, say the second, is a quadric: the process is substantially the same whatever may be the order of the other equation, and I take it to be a cubic: the two equations therefore are

$$(a, b, c, d | \xi, \eta)^3 = 0,$$

$$(A, B, C | \xi, \eta)^2 = 0;$$

giving rise to an equation

$$\nabla = (a, b, c, A, B, C)^2 = 0.$$

And it is required to perform the elimination so as to put in evidence the nodal line of this surface.

Take θ_1, θ_2 the roots of the second equation, or write

$$(A, B, C | \theta, 1)^2 = A(\theta - \theta_1)(\theta - \theta_2);$$

that is,

$$\theta_1 + \theta_2 = -\frac{2B}{A}, \quad \theta_1\theta_2 = \frac{C}{A}.$$

then, if

$$\begin{aligned}\Theta_1 &= (a, b, c, d \mid \xi, \eta)^3, \\ \Theta_2 &= (a, b, c, d \mid \xi, \eta)^3,\end{aligned}$$

we have

$$\nabla = A^4 \Theta_1 \Theta_2.$$

viz. on the right-hand side, replacing the symmetrical functions of θ_1, θ_2 by their values in terms of A, B, C we have the expression of ∇ in the form

$$\nabla = a^2 C^2 + \text{etc.}$$

Form now the expressions

$$\Theta_1 = \theta_1 X_1 + \theta_1^2 X_2 + \theta_1^3 X_3, \quad \Theta_2 = \theta_2 X_1 + \theta_2^2 X_2 + \theta_2^3 X_3,$$

each divisible by $\theta_1 - \theta_2$. These are evidently symmetrical functions of θ_1, θ_2 , the values being given by the successive lines of the determinant

$$\begin{array}{ccccccc} 0, & 1, & \theta_1 + \theta_2, & \theta_1^2 + \theta_1 \theta_2 + \theta_2^2 & (\&c. \text{ etc.}), \\ -1, & 0, & \theta_1 \theta_2, & \theta_1 \theta_2 (\theta_1 + \theta_2), \\ -(\theta_1 + \theta_2), & -\theta_1 \theta_2, & 0, & \theta_1^2 \theta_2^2, \\ -(\theta_1^2 + \theta_1 \theta_2 + \theta_2^2), & -\theta_1 \theta_2 (\theta_1 + \theta_2), & -\theta_1^2 \theta_2^2, & 0, \end{array}$$

and, consequently, these same quantities, each multiplied by A^3 , are given by the consecutive lines of

$$\begin{array}{ccccccc} 0, & A^2, & -2AB, & AC + 4B^2 & (\&c. \text{ etc.}), \\ -A^2, & 0, & AC, & C^2 - 2BC, \\ 2AB, & -AC, & 0, & C^2, \\ AC + 4B^2, & 2BC, & -C^2, & 0. \end{array}$$

Calling these X, Y, Z, W , that is, writing

$$X = 3Av - 6Bb + (-AC + 4B^2)a, \quad \&c.,$$

then X, Y, Z, W are the values of

$$\Theta_1 - \Theta_2, \quad \Theta_1 \Theta_2, \quad \Theta_1^2 \Theta_2, \quad \Theta_1 \Theta_2^2,$$

each multiplied by $A^3 = (\theta_1 - \theta_2)$; and the functions all four of them vanish if only $\theta_1 = \theta_2$, or, what is the same thing, the equations $X = 0, Y = 0, Z = 0, W = 0$ constitute only a twofold system.

The functions

$$(X, Y, Z), \quad (Y, Z, W)$$

[733] (CONCLUDED)
ON A FORMULA OF ELIMINATION.

contain each of them the factor $\theta_1\theta_2$, that is, ∇ ; they, in fact, each of them vanish if $\theta_1 = 0$, and they also vanish if $\theta_2 = 0$; or, by a direct substitution, we have

$$\begin{aligned} XW - Y^2 &= \frac{A^4}{(\theta_1 - \theta_2)^2} \cdot (\theta_1 - \theta_2)^2 \theta_1 \theta_2, & &= -A^4 \theta_1 \theta_2, \\ XW - YZ &= -(\theta_1 - \theta_2)^2 (\theta_1 + \theta_2) \theta_1 \theta_2, & &= -A^3 \theta_1 \theta_2 (\theta_1 + \theta_2), \\ YW - Z^2 &= -(\theta_1 - \theta_2)^2 \theta_1 \theta_2 \theta_1 \theta_2, & &= -A^2 \theta_1 \theta_2 \theta_1 \theta_2. \end{aligned}$$

Or, what is the same thing, these are $= -4\nabla, 2H\nabla, -C\nabla$, respectively; thus the first equation is

$$\begin{aligned} &\{3A^2c - 6ABb + (-AC + 4B^2)a\} \{2ABd - 3ACc + C^2a\} \\ &- (-A^2d + 3ACb - 2BCa)^2 = -A \cdot (A^3d + \&c.), = -2I\nabla; \end{aligned}$$

and similarly for the other two equations. The nodal curve is thus given by the twofold system $X = 0, Y = 0, Z = 0, W = 0$.


The method may be extended to the case where, instead of the quadric equation $(A, B, C \mid \vartheta, 1)^2 = 0$, we have an equation of any higher order, but the formulae are less simple.

734.

ON THE KINEMATICS OF A PLANE.

[From the *Quarterly Journal of Pure and Applied Mathematics*, vol. xvi. (1879),
pp. 1–8.]

It seems desirable to bring together under this title various questions which have been, or may be, proposed or discussed. We consider two planes in relative motion one upon the other, but, for convenience, they may be distinguished as a moving plane and a fixed plane, the first moving upon the second. Any point of the moving plane traces out on the fixed plane a curve, and any line of the moving plane envelopes on the fixed plane a curve; similarly, any point of the fixed plane traces out on the moving plane a curve, and any line of the fixed plane envelopes on the moving plane a curve. More generally, any curve of the moving plane envelopes on the fixed plane a curve, and any curve of the fixed plane envelopes on the moving plane a curve. There is, moreover, in the moving plane a curve which rolls upon a curve in the fixed plane, and these two curves (a single relative position being given) determine the motion.



The analytical theory presents no difficulty. Taking in the fixed plane the fixed axes Ox, Oy (fig. 1), and, fixed in the moveable plane so as to move with it, the axes O_1x_1, O_1y_1 ; then the position of the axes $O_1x_1y_1$ may be determined, say by α, β , the coordinates of O_1 in regard to Oxy ; and by θ , the inclination

of O_1x_1 to Ox . And denoting by x, y, x_1, y_1 the coordinates of a point P in regard to the two sets of axes respectively, then

$$\begin{aligned}x &= \alpha + x_1 \cos \theta - y_1 \sin \theta, \\y &= \beta + x_1 \sin \theta + y_1 \cos \theta;\end{aligned}$$

or, what is the same thing,

$$\begin{aligned}x_1 &= (x - \alpha) \cos \theta + (y - \beta) \sin \theta, \\y_1 &= -(x - \alpha) \sin \theta + (y - \beta) \cos \theta;\end{aligned}$$

or, as these last equations may be written,

$$\begin{aligned}x_1 &= \alpha_1 + x \cos(-\theta) - y \sin(-\theta), \\y_1 &= \beta_1 + x \sin(-\theta) + y \cos(-\theta),\end{aligned}$$

where $\alpha_1, \beta_1, = -\alpha \cos \theta - \beta \sin \theta, \alpha \sin \theta - \beta \cos \theta$, are the coordinates of O referred to the axes $O_1x_1y_1$, and $-\theta$ is the inclination of Ox to O_1x_1 .

When the motion is given, α, β, θ are given functions of a single variable parameter, say of t^* ; or, if we please, α, β are given functions of θ .

The velocities of a given point (x, y) are determined by the equations

$$\begin{aligned}x' &= \alpha' - (x_1 \sin \theta + y_1 \cos \theta) \theta', \\y' &= \beta' + (x_1 \cos \theta - y_1 \sin \theta) \theta';\end{aligned}$$

that is,

$$\begin{aligned}x' - \alpha' &= -(y - \beta) \theta', \\y' - \beta' &= (x - \alpha) \theta';\end{aligned}$$

or, as these equations may also be written,

$$\begin{aligned}(x' - \alpha') \sin \theta - (y' - \beta') \cos \theta &= -x_1 \theta', \\-(x' - \alpha') \cos \theta - (y' - \beta') \sin \theta &= y_1 \theta' .\end{aligned}$$

Hence if $x' = 0, y' = 0$, we have

$$\begin{aligned}x_1 \theta' &= \alpha' \sin \theta - \beta' \cos \theta, & x_1 &= -(y - \beta) + \frac{\alpha'}{\theta'}, \\y_1 \theta' &= \alpha' \cos \theta + \beta' \sin \theta, & y_1 &= (x - \alpha) + \frac{\beta'}{\theta'},\end{aligned}$$

which equations determine in terms of t, x_1 and y_1 the coordinates in regard to the axes $O_1x_1y_1$, and x and y the coordinates in regard to the axes Oxy , of I , the centre of instantaneous rotation.

If from the expressions of x_1, y_1 we eliminate t , we obtain an equation between (x_1, y_1) , which is that of the rolling curve in the moveable plane; and, similarly, if

from the expressions of x, y we eliminate t , we obtain a relation between (x, y) , which is that of the rolled-on curve in the fixed plane.

The system may be written

$$\begin{aligned}x_1 &= -\frac{\alpha'}{\theta'} \sin \theta + \frac{\beta'}{\theta'} \cos \theta, & x &= \alpha - \frac{\beta'}{\theta'}, \\y_1 &= \frac{\alpha'}{\theta'} \cos \theta + \frac{\beta'}{\theta'} \sin \theta, & y &= \beta + \frac{\alpha'}{\theta'};\end{aligned}$$

^{0*} t may be regarded as denoting the time, and then the derived functions of x, y in regard to t will denote velocities; and, to simplify the expression of the theorems, it is convenient to do this.

or, if we take θ as the independent variable,

$$\begin{aligned} x_1 &= -\alpha' \sin \theta + \beta' \cos \theta, & x &= \alpha - \beta', \\ y_1 &= \alpha' \cos \theta + \beta' \sin \theta, & y &= \beta + \alpha'. \end{aligned}$$

To find the variations of I , we have


$$\begin{aligned} x'_1 &= \alpha'' \sin \theta - \beta'' \cos \theta + \alpha' \cos \theta + \beta' \sin \theta, & &= \alpha'' \sin \theta - \beta'' \cos \theta + y_1, \\ y'_1 &= \alpha'' \cos \theta + \beta'' \sin \theta - \alpha' \sin \theta + \beta' \cos \theta, & &= \alpha'' \cos \theta + \beta'' \sin \theta - x_1, \\ x' &= \alpha' - \beta'', \\ y' &= \beta' + \alpha''. \end{aligned}$$

Hence

$$\begin{aligned} x'_1 &= x' \cos \theta + y' \sin \theta, & \text{or } x' &= x'_1 \cos \theta - y'_1 \sin \theta, \\ y'_1 &= -x' \sin \theta + y' \cos \theta, & \text{or } y' &= x'_1 \sin \theta + y'_1 \cos \theta, \end{aligned}$$

values which give $x'^2 + y'^2 = x_1'^2 + y_1'^2$, which equation expresses that the motion is in fact a rolling one.

Imagine the two curves, and the initial relative position given; say the two points A, A_1 (fig. 2) were originally in contact, then the arcs AI, A_1I are equal, and, calling each of these s , and X, Y, X_1, Y_1 the coordinates of I in regard to the two



sets of axes respectively, we have X, Y, X_1, Y_1 given functions of s , such that $X'^2 + Y'^2 = 1, X_1'^2 + Y_1'^2 = 1$, the accents now denoting differentiation in regard to s . We have, from the figure,

$$\theta = \tan^{-1} \frac{Y'}{X'} - \tan^{-1} \frac{Y'_1}{X'_1};$$

or, what is the same thing,

$$\tan \theta = (Y'X'_1 - Y'_1X') \div (X'X'_1 + Y'Y'_1),$$

say

$$\sin \theta, \cos \theta = Y'X'_1 - Y'_1X', \quad X'X'_1 + Y'Y'_1;$$

and then, as before,

$$\begin{aligned} x &= \alpha + x_1 \cos \theta - y_1 \sin \theta, \\ y &= \beta + x_1 \sin \theta + y_1 \cos \theta; \end{aligned}$$

or, what is the same thing,

$$\begin{aligned} x - X &= \cos \theta (x_1 - X_1) - \sin \theta (y_1 - Y_1), \\ y - Y &= \sin \theta (x_1 - X_1) + \cos \theta (y_1 - Y_1), \end{aligned}$$

where X, Y, X_1, Y_1 , and therefore also θ , denote given functions of s . The formulae will be of a like form if X, Y, X_1, Y_1 are given functions of a parameter t .

A well known but very interesting case is when two points of the moving plane describe right lines on the fixed plane. This may be discussed geometrically as follows: Suppose that we have the points A, C (fig. 3) describing the lines OA_0, OC_0 , which meet in O ; through A, C, O describe a circle, centre O_1 , and with centre

viz. the curve on the first plane is an ellipse, the semi-axes of which are $\pm(c + x_1)$, $\pm(c - x_1)$, each taken positively; if $x_1^2 + y_1^2 = c^2$, viz. if P be on the circle having AB for its diameter, then $y_1^2 = (c + x_1)(c - x_1)$, and we have

$$\frac{y}{x} = \frac{-(c - x_1) \cos \theta + y_1 \sin \theta}{(c + x_1) \sin \theta + y_1 \cos \theta}, \quad = -(c - x_1) \div y_1,$$

viz. as mentioned above, the curve on the fixed plane is a right line.

In the general case, we have

$$\begin{aligned} x(c - x_1) + yy_1 &= -(c^2 - x_1^2 - y_1^2) \cos \theta, \\ xy_1 + y(c + x_1) &= -(c^2 - x_1^2 - y_1^2) \sin \theta, \\ \{x(c - x_1) + yy_1\}^2 + \{xy_1 + y(c + x_1)\}^2 &= (c^2 - x_1^2 - y_1^2)^2, \end{aligned}$$

or, what is the same thing,

$$x^2\{(c - x_1)^2 + y_1^2\} + 4xy cy_1 + y^2\{(c + x_1)^2 + y_1^2\} = (c^2 - x_1^2 - y_1^2)^2.$$

Considering (x_1, y_1) as given, the curve traced out by P on the fixed plane is of the second order; it would be easy to verify from the equation that it is an ellipse, and to obtain for the position and magnitude of the axes the construction already found geometrically.

The same equation, considering therein (x, y) as constant and (x_1, y_1) as current coordinates, gives the curve traced out on the moving plane; the curve is obviously of the fourth order. Transferring the origin to A , we must in place of x_1 write $x_1 - c$; the equation thus becomes

$$x^2\{(x_1 - 2c)^2 + y_1^2\} + 4y_1 xy + y^2(x_1^2 + y_1^2) = (x_1^2 + y_1^2 - 2cx_1)^2,$$

or, what is the same thing,

$$(x_1^2 + y_1^2 - 2cx_1)^2 = (x^2 + y^2)(x_1^2 + y_1^2) - 4cx(xx_1 + yy_1) + 4c^2x^2;$$

and if we suppose herein $x = 0$, it becomes

$$(x_1^2 + y_1^2 - 2cx_1)^2 = (x_1^2 + y_1^2) y^2;$$

or, writing $x_1 = r_1 \cos \theta_1$, $y_1 = r_1 \sin \theta_1$, where $\theta_1 = \text{angle } QAB$, this is

$$(r_1 - 2c \cos \theta_1)^2 - y^2 = 0,$$

or say it is

$$r_1 = 2c \cos \theta_1 \pm y,$$

which is the polar equation of the curve described on the moveable plane by the point S , whose coordinates in respect to Ox and Oy are $(0, y)$.

There is no loss of generality in assuming $x = 0$. In fact, starting with any point S whatever of the fixed plane, if we draw OS meeting the small circle in A , and through A draw at right angles to this a line meeting the same circle in B , then, as before, the points A and B move along the fixed lines OA_0 , OB_0 ; or as regards the relative motion, taking A, B as fixed points, we have the originally fixed plane now moving in such wise that the two lines OA_0, OB_0 thereof (at right angles to each other) pass always through the points A and B respectively, and the curve is that described by the point S on the line OA ; the point O describes the circle on the diameter AB (the small circle), equation $r_1 = 2c \cos \theta_1$; and OQ having a given constant value $= y$, we have for the curve described by the point S the foregoing equation $r_1 = 2c \cos \theta_1 - y$; or writing $y = -f$, that is, taking S on the other side of O at a distance $OS = f$, the equation is $r_1 = 2c \cos \theta_1 + f$; viz. this is a nodal Cartesian or Limaçon, the origin being an acnode or a crunode according as $f >$ or $< 2c$; and if $f = 2c$, then we have the cuspidal curve or cardioid $r_1 = 2c(1 + \cos \theta_1) = 4c \cos^2 \frac{1}{2} \theta_1$.

The general conclusion is that the centre O of the large circle describes on the moving plane a small circle (centre O_1), and that every other point of the fixed plane describes on the moving plane a Limaçon having for its node a point of the small circle, and being, in fact, the curve obtained by measuring off along the radius vector of the small circle from its extremity a constant distance.

Considering in connexion with the point, coordinates (x_1, y_1) , (x, y) , a second point, coordinates (X_1, Y_1) , (X, Y) , in regard to the two sets of axes respectively, we have

$$\begin{aligned} x &= (c + x_1) \cos \theta - y_1 \sin \theta, & X &= (c + X_1) \cos \theta - Y_1 \sin \theta, \\ y &= (c - x_1) \sin \theta - y_1 \cos \theta, & Y &= (c - X_1) \sin \theta - Y_1 \cos \theta; \end{aligned}$$

from the first two equations we have

$$\cos \theta : \sin \theta : 1 = x(c - x_1) + yy_1 : xy_1 + y(c + x_1) : c^2 - x_1^2 - y_1^2,$$

and substituting these values in the second set, we find

$$\begin{aligned} &X : Y : 1 \\ &= x[c^2 - c(X_1 - x_1) - X_1x_1 - Y_1y_1] + y[c(y_1 - Y_1) - y_1X_1 - x_1Y_1] \\ &: x[c(y_1 - Y_1) - y_1X_1 + x_1Y_1] + y[c^2 - c(X_1 - x_1) - X_1x_1 - Y_1y_1] \\ &: c^2 - x_1^2 - y_1^2; \end{aligned}$$

or the points (x, y) , (X, Y) , considered as each of them moving on the fixed plane, are homographically related to each other.

To find the curve enveloped on the fixed plane by a given curve of the moving plane, we have only in the equation $f(x_1, y_1) = 0$ of the curve in the moving plane to substitute for x_1, y_1 their values in terms of x, y, θ , and then considering θ as a variable parameter, to find the envelope of the curve represented by this equation. And, similarly, we find the curve enveloped on the moving plane by a given curve of the fixed plane.

Thus, in the particular case of motion above considered, writing, as before,

$$\begin{aligned} x &= (c + x_1) \cos \theta - y_1 \sin \theta, \\ y &= (c - x_1) \sin \theta - y_1 \cos \theta; \end{aligned}$$

or conversely

$$\begin{aligned} x_1 &= x \cos \theta - y \sin \theta - c \cos 2\theta, \\ y_1 &= -x \sin \theta - y \cos \theta + c \sin 2\theta; \end{aligned}$$

the envelope on the moving plane of the line

$$Ax + By + C = 0$$

of the fixed plane is given as the envelope of the line

$$\{A(c + x_1) - By_1\} \cos \theta + \{-Ay_1 + B(c - x_1)\} \sin \theta + C = 0;$$

that is,

$$\{A(c + x_1) - By_1\}^2 + \{Ay_1 - B(c - x_1)\}^2 - C^2 = 0,$$

viz. this is

$$(A^2 + B^2)(x_1^2 + y_1^2 + c^2) + 2(A^2 - B^2)cx_1 - 4ABcy_1 - C^2 = 0,$$

a circle.

But the envelope on the fixed plane of the line

$$Ax_1 + By_1 + C = 0$$

of the moving plane is given as the envelope of the line

$$C + (Ax + By) \cos \theta - (Ay + Bx) \sin \theta - AC \cos 2\theta + BC \sin 2\theta = 0,$$

which can be obtained by equating to zero the discriminant of a quartic function, and is apparently a sextic curve.

735.

NOTE ON THE THEORY OF APSIDAL SURFACES.

[From the *Quarterly Journal of Pure and Applied Mathematics*, vol. xvi. (1879), pp. 109–112.]

I OBTAIN in the present Note a system of formulae which lead very simply to the known theorem, that the apsidals of reciprocal surfaces are reciprocal; or, what is the same thing, that the reciprocal of the apsidal of a given surface is the apsidal of its reciprocal; the surfaces are referred to the same axes, and by the reciprocal is meant the reciprocal surface in regard to a sphere radius unity, having for its centre a determinate point, say the origin; and it is this same point which is used in the construction of the apsidal surfaces. The apsidal of a given surface is constructed as follows; considering the section by any plane through the fixed point, and in this section the apsidal radii from the fixed point (that is, the radii which meet the curve at right angles), then drawing a line through the fixed point at right angles to the plane, and on this line measuring off from the fixed point distances equal to the apsidal radii respectively, the locus of the extremities of these distances is the apsidal surface. We have the surface, its reciprocal, the apsidal of the surface, the apsidal of the reciprocal; and I take

$$(x, y, z), (x', y', z'), (X, Y, Z), (X', Y', Z')$$

for the coordinates of corresponding points on the four surfaces respectively.

The condition of reciprocity gives $xx' + yy' + zz' - 1 = 0$, and (the equations being $U = 0$, $U' = 0$) x', y', z' proportional to $d_x U, d_y U, d_z U$, and x, y, z proportional to $d_{x'} U', d_{y'} U', d_{z'} U'$; or, what is the same thing, we must have

$$x' dx + y' dy + z' dz = 0 \quad \text{and} \quad x dx' + y dy' + z dz' = 0;$$

one of these is implied in the other, as appears at once by differentiating the equation $xx' + yy' + zz' - 1 = 0$.

The other two surfaces will therefore be reciprocal if only we have the like relations between the coordinates (X, Y, Z) and (X', Y', Z') ; that is, if

$$XX' + YY' + ZZ' - 1 = 0,$$

$$X' dX + Y' dY + Z' dZ = 0,$$

$$X dX' + Y dY' + Z dZ' = 0.$$

To find the apsidal surface, we consider an arbitrary section $x \cos \alpha + y \cos \beta + z \cos \gamma = 0$ of the surface $U = 0$, and seek to determine the apsidal radii thereof, that is, the maximum or minimum values of $R^2 = x^2 + y^2 + z^2$ when x, y, z vary subject to these two conditions. Writing x', y', z' to denote functions proportional to $d_x U, d_y U, d_z U$, we thus have the set of equations

$$x + \lambda x' + \mu \cos \alpha = 0,$$

$$y + \lambda y' + \mu \cos \beta = 0,$$

$$z + \lambda z' + \mu \cos \gamma = 0,$$

where λ, μ are indeterminate coefficients; taking then X, Y, Z as the coordinates of the extremity of the line drawn at right angles to the plane, we have $R^2 = X^2 + Y^2 + Z^2$, and $\cos \alpha, \cos \beta, \cos \gamma = \frac{X}{R}, \frac{Y}{R}, \frac{Z}{R}$; substituting these values in the equation

$$x \cos \alpha + y \cos \beta + z \cos \gamma = 0,$$

we have $Xx + Yy + Zz = 0$, and substituting in the other equations, and instead of λ, μ introducing the new indeterminate coefficients ρ, σ , we obtain

$$X, Y, Z = \rho x + \sigma x', \quad \rho y + \sigma y', \quad \rho z + \sigma z'.$$

Hence these last equations, together with

$$R^2 = X^2 + Y^2 + Z^2 = x^2 + y^2 + z^2,$$

and

$$Xx + Yy + Zz = 1,$$

contain the solution of the problem. If for convenience we introduce R'^2 to denote $x'^2 + y'^2 + z'^2$, and imagine the absolute values of x', y', z' determined so that $xx' + yy' + zz' = 1$, then substituting for X, Y, Z their values in the equations $X^2 + Y^2 + Z^2 = R^2$ and $Xx + Yy + Zz = 1$, we find

$$R^2 = \rho^2 R'^2 + 2\rho\sigma + \sigma^2 R'^2, \quad 1 = \rho R^2 + \sigma,$$

and thence

$$\rho^2 = \frac{1}{R^2 R'^2 - 1}, \quad \sigma = -\rho R^2,$$

or, finally assuming

$$\rho = \frac{1}{\sqrt{R^2 R'^2 - 1}},$$

we have

$$X, Y, Z = x - R^2 x', y - R^2 y', z - R^2 z',$$

each divided by

$$\sqrt{R^2 R'^2 - 1},$$

where I recall that x', y', z' are proportional to $d_x U, d_y U, d_z U$, and are such that $xx' + yy' + zz' = 1$: they in fact denote

$$d_x U, d_y U, d_z U, \text{ each divided by } x d_x U + y d_y U + z d_z U,$$

and that R^2 and R'^2 denote $x^2 + y^2 + z^2$ and $x'^2 + y'^2 + z'^2$ respectively. The coordinates X, Y, Z of the point of the apsidal surface are thus determined as functions of x, y, z .

For the apsidal of the reciprocal surface, we have in like manner

$$X', Y', Z' = x' - R'^2 x, y' - R'^2 y, z' - R'^2 z,$$

each divided by

$$\sqrt{R^2 R'^2 - 1},$$

and then the two sets of values give, not only

$$XX' + YY' + ZZ' = 1,$$

as is obvious, but also

$$X' dX + Y' dY + Z' dZ = 0, \quad \text{and} \quad X dX' + Y dY' + Z dZ' = 0.$$

In fact, writing for a moment p, p' instead of R^2, R'^2 , and $\sqrt{R^2 R'^2 - 1} = \sqrt{pp' - 1} = \omega$, then

$$\begin{aligned} X' dX + Y' dY + Z' dZ &= \sum \frac{x' - xp'}{\omega} d\left(\frac{x - x'p}{\omega}\right) + \&c. \\ &= \sum \frac{x' - xp'}{\omega} \left\{ \frac{dx - p dx' - x' dp}{\omega} - \frac{(x - x'p) d\omega}{\omega^2} \right\} + \&c. \\ &= \frac{1}{\omega^2} \left\{ \begin{aligned} &x' dx + y' dy + z' dz \\ &- p(x' dx' + y' dy' + z' dz') \\ &- (x'^2 + y'^2 + z'^2) dp \\ &- p'(x dx + y dy + z dz) \\ &+ pp'(x dx' + y dy' + z dz') \\ &+ p'(xx' + yy' + zz') dp \end{aligned} \right\} \\ &\quad - \frac{d\omega}{\omega^3} \left\{ \begin{aligned} &xx' + yy' + zz' \\ &- p(x'^2 + y'^2 + z'^2) \\ &- p'(x^2 + y^2 + z^2) \\ &+ pp'(xx' + yy' + zz') \end{aligned} \right\}, \end{aligned}$$

or, since the terms in $\{ \}$ are

$$0 - p \cdot \frac{1}{2} dp' - p' dp - p' \cdot \frac{1}{2} dp + 0 + p' dp = -\frac{1}{2}(p dp' + p' dp),$$

and

$$1 - pp' - pp' + pp' = 1 - pp' = -\omega^2,$$

this is

$$= \frac{1}{\omega^2} \left\{ -\frac{1}{2}(p dp' + p' dp) + \omega d\omega \right\} = 0,$$

in virtue of $\omega^2 = pp' - 1$. And similarly the other equation $X dX' + Y dY' + Z dZ' = 0$ might be directly verified.

736.

APPLICATION OF THE NEWTON-FOURIER METHOD TO AN IMAGINARY ROOT OF AN EQUATION.

*[From the Quarterly Journal of Pure and Applied Mathematics, vol. xvi. (1879),
pp. 179–185.]*

I CONSIDER only the most simple case, that of a quadric equation $x^2 = n^2$, where n^2 is a given imaginary quantity, having the square roots n and $-n$; starting from an assumed approximate (imaginary) value $x = a$, we have $(a + h)^2 = n^2$, that is,

$$a^2 + 2ah = n^2, \quad h = \frac{n^2 - a^2}{2a}, \quad \text{and} \quad a + h = \frac{a^2 + n^2}{2a},$$

that is, the successive values are

$$a_1 = \frac{a^2 + n^2}{2a}, \quad a_2 = \frac{a_1^2 + n^2}{2a_1}, \quad \dots,$$

and the question is, under what conditions do we thus approximate to one determinate root (selected out of the two roots at pleasure), say n , of the given equation.

The nearness of two values is measured by the modulus of their difference; thus a nearer to n , than a_1 is to n , means $\text{mod}(a - n) < \text{mod}(a_1 - n)$, and so in other cases; in the course of the approximation $a, a_1, a_2, \dots$ to n , any step, for instance a to a_1 , is regular if a_1 is nearer to n than a is, but otherwise it is irregular; the approximation is regular if all the steps are regular, and if (after one or more irregular steps) all the subsequent steps are regular, then the approximation becomes regular at the step which is the first of the unbroken series of regular steps.

We do by an approximation, which is ultimately regular, obtain the value n , if only the assumed value a is nearer to n than it is to $-n$; or, say, if the condition $\text{mod}(a - n) < \text{mod}(a + n)$ is satisfied, and the approximation is regular from the beginning if $\text{mod}(a - n) \leq \frac{1}{2} \text{mod} n$, viz. this condition is a sufficient one¹; the first step a to a_1 will moreover be regular under a less stringent condition imposed upon a ; and it would seem that, without the condition $\text{mod}(a - n) < \frac{1}{2} \text{mod} n$ being satisfied, the subsequent steps will in some cases be also regular; that is, that the last-mentioned condition is not a necessary condition in order to the approximation being regular from the beginning; it is, however, the necessary and sufficient condition, to be satisfied by the modulus of $a - n$, in order that the approximation may be regular from the beginning. All this will clearly appear from the geometry.

[Figure 1: Points N, N' on a horizontal axis with O at the midpoint; through O a perpendicular QQ' . A circle on $N'M$ as diameter where $OM = \frac{1}{2}ON$ (the “circle of unfitness”); the segment on the N -side of QQ' is the “segment of unfitness.” Points $A, A_1, \dots$ representing successive approximations.]

¹In the Smith's Prize Examination, Jan. 28, 1879, I gave the theorem under the following form: “If a, n are imaginary quantities, the latter of them given, and the former assumed at pleasure, subject only to the condition $\text{mod}(a - n) < \frac{1}{2} \text{mod} n$; then if $a_1 = \frac{a^2 + n^2}{2a}$, $a_2 = \frac{a_1^2 + n^2}{2a_1}$, &c., show that the terms $a, a_1, a_2, \dots$ will converge to the limit n .” This is strictly true, but it would have been better to say “will converge regularly.”

We take N, N' (fig. 1) to represent the values $n, -n$; and similarly A, A_1 , &c. to represent the quantities $a, a_1, \dots$; we have then

$$AN = \text{mod}(a - n), \quad A_1N = \text{mod}(a_1 - n) \dots,$$

so that the approximation is measured by the approach of the points A, A_1 to N . The line NN' joining the points N, N' passes through, and is bisected at, the origin O ; drawing then QQ' through O at right angles to NN' the condition

$$\text{mod}(a - n) < \text{mod}(a + n)$$

means that the point A , which represents the imaginary quantity a , lies on the N -side of QQ' , and it will be assumed throughout that this is so. Take now on the line ON , $OM = \frac{1}{2}ON$, and on $N'M$ as diameter, describe a circle, which may be called the “circle of unfitness”; regarding as an area the segment hereof which lies on the N -side of QQ' , say this is the “segment of unfitness.” It will be shown that according as A is situate inside, on the boundary of, or outside the segment of unfitness, A_1N will be greater than, equal to, or less than AN . It may be added that, if A be within or upon the boundary of the segment of unfitness, then A_1 will be outside it, but this by no means hinders that the next point A_2 , or some later point, shall be within the segment of unfitness; and, further, that when A is outside the segment of unfitness, then the next point A_1 , or some later point, may very well be within the segment of unfitness; the conclusion is, that A being inside the segment of unfitness, A_1N is less than AN , but that it does not thence follow that A_2N is less than A_1N , A_3N than $A_2N, \dots$; the approximation although regular at the first step, may then, or afterwards, for a step or steps, cease to be regular.

If, however, AN be less than $\frac{1}{2}ON$, that is, if the condition $\text{mod}(a - n) < \frac{1}{2}\text{mod}n$ be satisfied, then the point A lies within the circle centre N and radius NM , and is consequently outside the segment of unfitness; A_1N being less than AN , the point A_1 is a fortiori outside the segment of unfitness, and the like for all the subsequent points $A_2, A_3, \dots$; that is, in this case, the approximation is regular throughout. The circle, centre N , and radius $NM, = \frac{1}{2}\text{mod}n$, may be called the “safe circle”; and the conclusion is that, if the point A or any subsequent point be within the safe circle, then every subsequent point will be within the safe circle, and the approximation will be regular.

The successive points $A, A_1, A_2, \dots$ (or, as it will be convenient to call them, $A_1, A_2, A_3, \dots$) may be obtained each from the preceding one by a simple geometrical construction.

I recall that any circle through the two (imaginary) antipoints of N, N' is a circle having its centre on the indefinite line NN' ; it is such that the ratio of the distances of a point thereof from the points N, N' respectively has a certain constant value, viz. for the circles with which we are here alone concerned, those which lie on the N -side of QQ' , the centres lie beyond the point N (further away, that is, from O), and the values of the ratio, distance from N to distance from N' , are less than unity.

Starting then from the given point A_1 , for which this ratio $A_1N : A_1N'$ has a given value, suppose $A_1N = k A_1N'$, we describe a first circle (passing of course through A_1) for each point of which this ratio has the value k ; let the diameter of this circle be V_1W_1 , V_1 being the extremity between O and N , W_1 (not shown in the figure), that beyond N ; we then describe a second circle, for which the ratio is $= k^2$; let its diameter be V_2W_2 , V_2 being the extremity between N and O (or say between V_1 and N), W_2 , that beyond N (or say between N and W_1); the point A_2 lies on this second circle, and is determined as the single intersection of the line V_2A_1 with the second circle. And of course drawing a third circle, for which the ratio is $= k^4$, on the diameter V_3W_3 , then A_3 lies on the third circle, and is the intersection with it of the line V_3A_2 , and so on; the radii of the successive circles diminish very rapidly, their centres, in like manner, continually approaching the point N ; hence, the points $A_1, A_2, A_3, \dots$, which lie on the several circles respectively approximate, and that very rapidly, to the point O . But by what precedes, if, for instance, the point A_1 be within the segment of unfitness, then also some of the subsequent points may be within the segment of unfitness, and for each point A_p , for which this is the case, the next point A_{p+1} is at a greater distance, so that $NA_{p+1} > NA_p$; it is, however, clear that we always arrive at a point A_q , such that $NA_q < \frac{1}{2}ON$, and so soon as such a point is arrived at the approximation becomes regular.

The point A_2 determined from A_1 , as above, is a point such that the subtended angle NA_2N' is = twice the subtended angle NA_1N' ; or calling the latter angle ϕ , the former is 2ϕ . It is, in fact, this property which gives rise to the construction; for let the values of A_1N, A_1N' , regarded as imaginary quantities, be called for a moment

$$\rho_1(\cos \theta_1 + i \sin \theta_1), \quad \rho'_1(\cos \theta'_1 + i \sin \theta'_1);$$

and, similarly, those of A_2N , A_2N' be called

$$\rho_2(\cos \theta_2 + i \sin \theta_2), \quad \rho'_2(\cos \theta'_2 + i \sin \theta'_2);$$

then these are the values of $a_1 + n$, $a_1 - n$, $a_2 + n$, $a_2 - n$ respectively, or we have

$$\begin{aligned} \frac{a_1 - n}{a_1 + n} &= \frac{\rho_1}{\rho'_1} [\cos(\theta_1 - \theta'_1) + i \sin(\theta_1 - \theta'_1)] = k (\cos \phi + i \sin \phi), \\ \frac{a_2 - n}{a_2 + n} &= \frac{\rho_2}{\rho'_2} [\cos(\theta_2 - \theta'_2) + i \sin(\theta_2 - \theta'_2)] = k^2 (\cos 2\phi + i \sin 2\phi), \end{aligned}$$

that is,

$$\left(\frac{a_1 - n}{a_1 + n} \right)^2 = \frac{a_2 - n}{a_2 + n},$$

which relation between a_2 , a_1 is in fact the original relation

$$a_2 = \frac{a_1^2 + n^2}{2a_1};$$

and, conversely, a_2 , a_1 , being thus connected, then the representative A_2 is obtained from the representative point A_1 by the foregoing geometrical construction.

I give the analytical proofs; we may without loss of generality take, and it is convenient to do so, the axis of x as coinciding with the line ON , and to put also $ON = 1$. We then in place of the original coordinates x , y of any point take the new coordinates k , ϕ , which are such that

$$\frac{x^2 + y^2 - 1 + 2iy}{x^2 + y^2 + 1 - 2x} = k e^{i\phi},$$

$$\frac{x - iy - 1}{x - iy + 1} = k e^{-i\phi},$$

equations which may also be written

$$(x - iy + 1)^2 = e^{i\phi} [(x + iy + 1)^2],$$

or, what is the same thing,

$$\begin{aligned} x^2 + (y - i)^2 &= e^{-2i\phi} [x^2 + (y + i)^2]; \\ x^2 + y^2 + 2y \cot \phi &= 0, \end{aligned}$$

where of course the equation with k shows that k is equal to the ratio of the distances of the point from the points N , N' respectively, and the equation in ϕ , taken in the second form, shows that ϕ is the angle subtended at the point by N , N' .

It is sometimes convenient to write $ke^{i\phi}$, $ke^{-i\phi} = p$, q respectively; we then have

$$x = \frac{p+q}{1-p} \cdot \frac{1}{1-q} \cdot p, \quad x - iy = \frac{1+p}{1-p}, \quad x - iy = \frac{1+q}{1-q}.$$

Suppose for a moment that we have (p_1, q_1) , (p_2, q_2) , (p_3, q_3) as the (p, q) coordinates of any three points, the condition that these three points may lie in a line, is given in the form, determinant = 0, where each line of the determinant is of the form

$$1, \quad \frac{1+p}{1-p}, \quad \frac{1+q}{1-q},$$

or, what is the same thing, it is

$$1 - pq + p - q, \quad 1 - pq - p + q, \quad 1 + pq - p - q,$$

or, again

$$pq - 1, \quad p - q, \quad p + q - 2,$$

viz. the condition is

$$\begin{vmatrix} p_1 q_1 - 1, & p_1 - q_1, & p_1 + q_1 - 2 \\ p_2 q_2 - 1, & p_2 - q_2, & p_2 + q_2 - 2 \\ p_3 q_3 - 1, & p_3 - q_3, & p_3 + q_3 - 2 \end{vmatrix} = 0.$$

Suppose the (k, ϕ) coordinates of the three points are (l, α) , (m, β) , (n, γ) respectively; then this equation is

$$\begin{vmatrix} l^2 - 1, & il \sin \alpha, & l \cos \alpha - 1 \\ m^2 - 1, & im \sin \beta, & m \cos \beta - 1 \\ n^2 - 1, & in \sin \gamma, & n \cos \gamma - 1 \end{vmatrix} = 0,$$

viz. it is

$$\begin{vmatrix} l^2 - 1, & l \sin \alpha, & l \cos \alpha \\ m^2 - 1, & m \sin \beta, & m \cos \beta \\ n^2 - 1, & n \sin \gamma, & n \cos \gamma \end{vmatrix} - \begin{vmatrix} l^2 - 1, & l \sin \alpha, & 1 \\ m^2 - 1, & m \sin \beta, & 1 \\ n^2 - 1, & n \sin \gamma, & 1 \end{vmatrix} = 0,$$

or, what is the same thing, it is

$$\begin{aligned} & [(l^2 - 1)mn \sin(\beta - \gamma) + (m^2 - 1)nl \sin(\gamma - \alpha) + (n^2 - 1)lm \sin(\alpha - \beta)] \\ & + [(m^2 - n^2)l \sin \alpha + (n^2 - l^2)m \sin \beta + (l^2 - m^2)n \sin \gamma] = 0. \end{aligned}$$

If in this equation $\gamma = \pi$, and $\beta = 2\alpha$, so that $\sin(\alpha - \beta) = -\sin \alpha$, the equation will contain only terms in $\sin \alpha$ and $\sin 2\alpha$, viz. it will be

$$\begin{aligned} & [(m^2 - n^2)l + (n^2 - 1)nl - (n^2 - 1)lm] \sin \alpha \\ & + [-(l^2 - 1)mn + (n^2 - l^2)m] \sin 2\alpha = 0, \end{aligned}$$

that is,

$$m(m - 1)(n + 1)(m - n) \sin \alpha + (m^2 - 1)(n - l^2) \sin 2\alpha = 0,$$

or, what is the same thing,

$$(m + 1) \sin \alpha \{l(n + 1)(m - n) + 2m(n - l^2) \cos \alpha\} = 0,$$

which is satisfied for any values whatever of l, m, n , by a proper value of $\cos \alpha$; and is also satisfied irrespectively of the value of α if only $m = 1$ or $n = l^2$; that is, writing $m = k$ instead of l , say if $l = k$, $m = k^2$, $n = k^4$; that is, writing also ϕ in place of α , the three points

$$(k, \phi), \quad (k^2, 2\phi), \quad \text{and} \quad (k^4, \pi)$$

are in a right line; viz. the point A_1 , circle k , subtended angle ϕ ; the point A_2 , circle k^2 , subtended angle 2ϕ ; and the point V_2 , same circle, subtended angle π ; are in a right line.

The equation of the circle of unfitness can be obtained more easily in a different manner; but I have thought it worth while to give the investigation by means of the foregoing (p, q) coordinates.

Suppose that p_1, q_1 refer to the point A_1 : then we have

$$(A_1 N)^2 = (x_1 - 1)^2 + y_1^2 = (x_1 + iy_1 - 1)(x_1 - iy_1 - 1) = \left(\frac{1 + p_1}{1 - p_1} - 1\right) \left(\frac{1 + q_1}{1 - q_1} - 1\right),$$

that is,

$$(A_1 N)^2 = \frac{4p_1 q_1}{(1 - p_1)(1 - q_1)}.$$

Similarly, if p_2, q_2 refer to the point A_2 , then

$$(A_2 N)^2 = \frac{4p_2 q_2}{(1 - p_2)(1 - q_2)} = \frac{4p_1^2 q_1^2}{(1 - p_1^2)(1 - q_1^2)},$$

since $p_2 = p_1^2$, $q_2 = q_1^2$. The two are equal if

$$(1 + p_1)(1 + q_1) = p_1 q_1,$$

that is,

$$p_1 + q_1 + 1 = 0.$$

Writing for a moment $x_1 + iy_1 = \xi$, $x_1 - iy_1 = \eta$, we have

$$p_1 = \frac{\xi - 1}{\xi + 1}, \quad q_1 = \frac{\eta - 1}{\eta + 1},$$

and the equation is

$$\frac{\xi - 1}{\xi + 1} + \frac{\eta - 1}{\eta + 1} + 1 = 0,$$

that is,

$$(\xi - 1)(\eta + 1) + (\eta - 1)(\xi + 1) + (\xi + 1)(\eta + 1) = 0,$$

or substituting for ξ, η their values, the equation is

$$3(x_1^2 + y_1^2) + 2x_1 - 1 = 0,$$

that is,

$$(x_1 + 1)(x_1 - \frac{1}{3}) + y_1^2 = 0,$$

the equation of a circle on the diameter $N'M$, which is, in fact, the before-mentioned circle of unfitness; viz. A_1 being on the circumference of this circle, or say on the boundary of the segment of unfitness, then $A_1N = A_2N$; whence also, according as A_1 is inside or outside the segment, $A_2N < A_1N$ or $> A_1N$.

Suppose A_1 to be on the circle, that is, $p_1 + q_1 + 1 = 0$; it is easy to show that the locus of A_2 is also a circle. We have in fact $(p_1 q_1)^2 + 1 = 0$, that is,

$$p_1^2 + q_1^2 + 2p_1^2 q_1^2 + 1 = 0,$$

or say

$$p_2 + q_2 + 2k^4 - 1 = 0,$$

viz. this is

$$\begin{aligned} \frac{\xi - 1}{\xi + 1} + \frac{\eta - 1}{\eta + 1} + 2k^4 - 1 &= 0, \\ (2k^4 + 1)\xi\eta + (2k^4 - 1)(\xi + \eta) + 2k^4 - 3 &= 0, \end{aligned}$$

that is,

$$x_1^2 + y_1^2 + \frac{2k^4 - 1}{2k^4 + 1} \cdot 2x_1 + \frac{2k^4 - 3}{2k^4 + 1} = 0,$$

or finally

$$(x_1 + 1) \left(x_1 - \frac{3 - 2k^4}{2k^4 + 1} \right) + y_1^2 = 0.$$

Measuring off from O in the direction of ON a distance $OS = \frac{3 - 2k^4}{2k^4 + 1}$ (always $> \frac{1}{3}$, since $k^4 < 1$), the circle in question is that on the diameter $N'S$; this is a circle touching at N' , and containing within it the circle of unfitness; if $k = 1$ (that is, for A_1 on the line QQ') it becomes identical with the circle of unfitness, but except in this limiting case it does not meet the circle of unfitness in any point on the N -side of QQ' , that is, A_1 being on the boundary of the segment of unfitness, A_2 is never on this boundary; and it thus appears that A_1 being inside the segment, A_2 is always outside the segment.

It is to be further noticed, that we have

$$(A_2N)^2 = \frac{4p_2q_2}{(1 - p_2)(1 - q_2)},$$

or

$$\frac{(A_2N)^2}{(A_1N)^2} = \frac{1 - p_1 + q_1}{1 - p_1 - q_1} = \frac{3\xi\eta + \xi + \eta - 1}{3} = \frac{3(x_1^2 + y_1^2 + \frac{2}{3}x_1 - \frac{1}{3})}{1},$$

that is,

$$\frac{(A_2N)^2}{(A_1N)^2} = 1 + \frac{3T^2}{(A_1N)^2},$$

where T is the tangential distance of A_1 from the circle of unfitness; there should, it appears to me, be some more elegant formula for the ratio $A_2N \div A_1N$ which determines whether the step is regular or irregular.

It is worth noticing how the conditions

$$\text{mod}(a - n) < \text{mod}(a + n) \quad \text{and} \quad \text{mod}(a - n) < \frac{1}{2}\text{mod}n,$$

present themselves in the real theory. Making the usual construction by means of the parabola $y = x^2$, the first condition means that the point A must be taken on the N -side of O (fig. 2); the second that, in order to the regularity of the approxi-

[Figure 2: A parabola $y = x^2$ with the axis Ox ; points N' , O , A , N , A_1 shown on the axis. The construction shows that if $OA = \frac{1}{2}ON$ then $AN = NA_1$.]

mation, A must be taken at a distance from $O > \frac{1}{2}ON$; in fact, if (as in the figure) $OA = \frac{1}{2}ON$, then $AN = NA_1$, or the point A_1 is at an equal distance with A from N ; and thence, according as OA is greater or less than $\frac{1}{2}ON$, the point A_1 is nearer or further than A to or from N .

737.

ON A COVARIANT FORMULA.

[From the Quarterly Journal of Pure and Applied Mathematics, vol. xvi. (1879), pp. 224–226.]

Starting from the equation

$$x_1 = x - \frac{fx}{f'x},$$

which presents itself in the Newton-Fourier problem, it is easy to see that, if a be a root of the equation $fx = 0$, then

$$x_1 - a = \frac{(x - a)f'x - fx}{f'x}$$

contains the factor $(x - a)^2$; that is, the equation $(x - x_1)f'x - fx = 0$, considered as an equation in x containing the parameter x_1 , will have a twofold root, if x_1 is equal to any root a of the equation $fx = 0$; and, consequently, the discriminant in regard to x of the function $(x - x_1)f'x - fx$ will contain the factor fx_1 . But if fx be of the order n , then the discriminant is of the order $2n - 2$ in x_1 , and there is consequently a remaining factor of x_1 of the order $n - 2$.

The like theorem applies to the homogeneous form

$$(\alpha y - \beta x) \left(\alpha \frac{d}{d\alpha} + \beta \frac{d}{d\beta} \right) f(\alpha, \beta) - (\alpha y - \beta x) f(\alpha, \beta),$$

which reduces itself to the foregoing on writing $\alpha = 1$, $\beta = 0$, $y = y_1 = 1$; or, changing the notation, say to the form

$$(\xi y - \eta x) \left(\xi \frac{d}{d\xi} + \eta \frac{d}{d\eta} \right) f(\xi, \eta) - (\alpha y - \beta x) f(\xi, \eta),$$

viz. the discriminant hereof in regard to ξ, η , being a function, homogeneous of the order $2n - 2$ in regard to x, y , to α, β , and to the coefficients of $f(\xi, \eta)$, will contain the factor $f(x, y)$, and there will be consequently a remaining factor of the order $n - 2$ in (x, y) , $2n - 2$ in (α, β) and $2n - 3$ in the coefficients of $f(\xi, \eta)$.

The most simple case is when $f(\xi, \eta)$ is the quadric function $(a, b, c | \xi, \eta)^2$. The form here is

$$(\xi y - \eta x) [(a\alpha + b\beta)\xi + (b\alpha + c\beta)\eta] - (\alpha y - \beta x)(a, b, c | \xi, \eta)^2 = (\mathfrak{a}, \mathfrak{b}, \mathfrak{c} | \xi, \eta)^2,$$

where the coefficients are

$$\begin{aligned} \mathfrak{a} &= y(a\alpha + b\beta) - \alpha(\alpha y - \beta x) = a\beta x + (a\alpha + 2b\beta)y, \\ \mathfrak{b} &= y(b\alpha + c\beta) - x(a\alpha + b\beta) - b(\alpha y - \beta x) = -a\alpha x + c\beta y, \\ \mathfrak{c} &= -x(b\alpha + c\beta) - c(\alpha y - \beta x) = -(2b\alpha + c\beta)x - c\alpha y; \end{aligned}$$

and we then have

$$\begin{aligned} \mathfrak{a}\mathfrak{c} - \mathfrak{b}^2 &= -(2b\alpha\beta + c\beta^2) a\alpha x^2 \\ &\quad - \{2ab\alpha^2 + (2ac + 4b^2) \alpha\beta + 2bc\beta^2\} xy - (a\alpha^2 + 2b\alpha\beta) c y^2 \\ &\quad - a\alpha\beta \cdot a\alpha x^2 - (-2ac\alpha\beta) xy - c\beta \cdot c\alpha y^2, \end{aligned}$$

which is

$$= -(a\alpha^2 + 2b\alpha\beta + c\beta^2)(ax^2 + 2bxy + cy^2).$$

The discriminant is in this case

$$= -(a, b, c \mid \alpha, \beta)^2 \cdot (a, b, c \mid x, y)^2.$$

In the case of the cubic function $(a, b, c, d \mid \xi, \eta)^3$, the form is

$$\begin{aligned} &(\xi y - \eta x) \{3(a\alpha + b\beta, b\alpha + c\beta, c\alpha + d\beta \mid \xi, \eta)^2\} \\ &-(\alpha y - \beta x) (a, b, c, d \mid \xi, \eta)^3 = (\mathfrak{a}, \mathfrak{b}, \mathfrak{c}, \mathfrak{d} \mid \xi, \eta)^3, \end{aligned}$$

the values of the coefficients being

$$\begin{aligned} \mathfrak{a} &= a\beta x + (2a\alpha + 3b\beta) y, \\ \mathfrak{b} &= -a\alpha x + (b\alpha + 2c\beta) y, \\ \mathfrak{c} &= -(2b\alpha + c\beta) x + d\beta y, \\ \mathfrak{d} &= -(3c\alpha + 2d\beta) x - d\alpha y. \end{aligned}$$

Attending only to the terms in x^2 , we have

$$\begin{aligned} \mathfrak{a}\mathfrak{c} - \mathfrak{b}^2 &= -(a\alpha^2 + 2b\alpha\beta + c\beta^2) a\alpha x^2, \\ \mathfrak{a}\mathfrak{d} - \mathfrak{b}\mathfrak{c} &= -2(b\alpha^2 + 2c\alpha\beta + d\beta^2) a\alpha x^2, \\ \mathfrak{b}\mathfrak{d} - \mathfrak{c}^2 &= -\{(3ac - 4b^2) \alpha^2 - (2ad - 4bc) \alpha\beta - c^2\beta^2\} x^2. \end{aligned}$$

And hence, in

$$\mathfrak{a}^2\mathfrak{d}^2 + 4\mathfrak{a}\mathfrak{c}^3 + 4\mathfrak{b}^3\mathfrak{d} - 3\mathfrak{b}^2\mathfrak{c}^2 - 6\mathfrak{a}\mathfrak{b}\mathfrak{c}\mathfrak{d} = (\mathfrak{a}\mathfrak{d} - \mathfrak{b}\mathfrak{c})^2 - 4(\mathfrak{a}\mathfrak{c} - \mathfrak{b}^2)(\mathfrak{b}\mathfrak{d} - \mathfrak{c}^2),$$

we have the term

$$\begin{aligned} &4a\alpha^2 \cdot [a(b\alpha^2 + 2c\alpha\beta + d\beta^2)^2 \\ &+ (a\alpha^2 + 2b\alpha\beta + c\beta^2) \{(3ac - 4b^2) \alpha^2 - (2ad - 4bc) \alpha\beta - c^2\beta^2\}] x^4; \end{aligned}$$

then, forming the analogous term in y^4 , and assuming that the whole divides by $(a, b, c, d \mid x, y)^3$, and also expanding the $\alpha\beta$ -functions within the square brackets, we find

Discriminant = $4(a, b, c, d \mid x, y)^3$ multiplied by

$$\begin{aligned} &[3a^2c - 3ab^2, \\ &2a^2d + 6abc - 8b^3, \\ &6abd + 6ac^2 - 14b^2c, \\ &6acd - 6bc^2, \\ &ad^2 - bc^2] \alpha^4 () / \beta^4 + [ad - b^2, \\ &6abd - 6b^2c, \\ &6acd + 6b^2d - 12bc^2, \\ &2ad^2 + 6bcd - 8d^3, \\ &3bd^2 - 3cd] \beta^4 () / \alpha^4. \end{aligned}$$

Writing down the Hessian of $(a, b, c, d \mid \alpha, \beta)^3$,

$$H = (ac - b^2, ad - bc, bd - c^2 \mid \alpha, \beta)^2,$$

and the cubicovariant

$$\Phi = (a^2d - 3abc + 2b^3, abd - 2ac^2 + b^2c, -acd + 2b^2d - bc^2, -ad^2 + 3bcd - 2c^3) (\alpha, \beta)^3.$$

It is easy to see that the coefficient of x is

$$= 3(a, b, c \mid \alpha, \beta)^2 (H - \beta\Phi);$$

hence also that of y is

$$= -3(a, b, c \mid \alpha, \beta)^2 (\beta H + \alpha\Phi), = 3(b, c, d \mid \alpha, \beta)^2 (\beta H + \alpha\Phi);$$

and the final result is that the discriminant $= 4(a, b, c, d \mid x, y)^3$ multiplied by

$$[3(a, b, c, d \mid \alpha, \beta)^2 (x, y) H + (\alpha y - \beta x) \Phi].$$

It would be interesting to calculate the result for the quartic $(a, b, c, d, e \mid \xi, \eta)^4$.

March 14, 1879.

738.

NOTE ON A HYPERGEOMETRIC SERIES.

[From the *Quarterly Journal of Pure and Applied Mathematics*, vol. xvi. (1879), pp. 268–270.]

In the memoir on hypergeometric series, Schwarz, “*Ueber diejenigen Fälle, &c.*,” Crelle, t. LXXV. (1873), pp. 292–335, the author shows, as part of his general theory, that the equation

$$\frac{d^2y}{dx^2} - \frac{\frac{2}{3} - \frac{1}{2}x}{x(1-x)} \cdot \frac{dy}{dx} + \frac{\frac{1}{48}}{x(1-x)^2} y = 0,$$

which belongs to the hypergeometric series $F(\frac{1}{12}, -\frac{1}{12}, \frac{2}{3}, x)$, is algebraically integrable, having in fact the two particular integrals

$$y^6 = -x + 2\sqrt{(\alpha - \alpha^5x)} \cdot \sqrt{(-\alpha^5 + \alpha x)},$$

where α is a prime sixth root of -1 , $\alpha^6 + 1 = 0$, or say $\alpha^4 - \alpha^2 + 1 = 0$ (see p. 326, α being for greater simplicity written instead of δ^2 , and the form being somewhat simplified).

It is interesting to verify this directly; writing first $y = \sqrt{Y}$ and then $x = X^2$, the equation between Y , X is easily found to be

$$\frac{d^2Y}{dX^2} - \frac{\frac{4}{3}X}{1-X^2} \frac{dY}{dX} + \frac{1}{4} \cdot \frac{1}{Y} \left(\frac{dY}{dX} \right)^2 + \frac{\frac{1}{6}}{1-X^2} Y - \frac{X}{1-X^2} Y = 0,$$

and the theorem in effect is that that equation has the two particular integrals

$$Y = \sqrt{(-P)} \pm \sqrt{(Q)},$$

P and Q being linear functions of X : in fact,

$$\begin{aligned} P &= \alpha - \alpha^5 X, \\ Q &= -\alpha^5 + \alpha X. \end{aligned}$$

Starting say from the equation

$$Y = \sqrt{(P)} + \sqrt{(Q)},$$

or, as it is convenient to write it,

$$Y = P^{\frac{1}{2}} + Q^{\frac{1}{2}},$$

where P and Q are assumed to be linear functions of X , we have

$$\begin{aligned} \frac{dY}{dX} &= \frac{1}{2} P^{-\frac{1}{2}} P' + \frac{1}{2} Q^{-\frac{1}{2}} Q', \\ \frac{d^2Y}{dX^2} &= -\frac{1}{4} P^{-\frac{3}{2}} P'^2 - \frac{1}{4} Q^{-\frac{3}{2}} Q'^2, \end{aligned}$$

and thence

$$\begin{aligned} \frac{d^2Y}{dX^2} &= -\frac{1}{4} P^{-\frac{3}{2}} P'^2 - \frac{1}{4} Q^{-\frac{3}{2}} Q'^2 - \frac{1}{4} Q^{-\frac{1}{2}} P^{-\frac{1}{2}} P'^2 - \frac{1}{4} P^{-\frac{1}{2}} Q^{-\frac{1}{2}} Q'^2, \\ Y^2 &= \frac{1}{2}(P^2 + Q^2) + P^{\frac{1}{2}} Q^{\frac{1}{2}} P' + P^{\frac{1}{2}} Q^{\frac{1}{2}} Q', \\ \left(\frac{dY}{dX}\right)^2 &= \frac{1}{4} P^{-1} P'^2 + \frac{1}{4} Q^{-1} Q'^2 + \frac{1}{2} P^{-\frac{1}{2}} Q^{-\frac{1}{2}} P' Q', \end{aligned}$$

where P', Q' are written to denote the derived functions of P, Q respectively.

Substituting these values, the resulting equation contains on the left-hand side a rational part, and a part with the factor $P^{-\frac{1}{2}} Q^{-\frac{1}{2}}$, and it is clear the equation can only be true if these two parts are separately = 0. We have thus two equations which ought to be verified; viz. after a slight reduction these are found to be

$$\frac{2}{3} X (P' + Q') - \frac{1}{6} (P + Q) = 0,$$

$$\frac{1}{4} PQ (QP'^2 + PQ'^2) + QP^{\frac{1}{2}} Q^{\frac{1}{2}} P' Q' + PQ^{\frac{1}{2}} P^{\frac{1}{2}} P' Q' - \frac{4}{3} X \cdot PQ (PQ' + P'Q) - \frac{1}{6} \cdot X P' Q' = 0,$$

and it is very interesting to observe the manner in which these equations are, in fact, verified by the foregoing values of P, Q .

We have

$$P + Q = (\alpha - \alpha^5)(1 + X), \quad P' + Q' = \alpha - \alpha^5,$$

and hence

$$\frac{2}{3} X (P' + Q') - \frac{1}{6} (P + Q) = -(\alpha - \alpha^5) \frac{1}{6} (1 - X),$$

or, in the first equation, the second part

$$\frac{\frac{2}{3} X}{1 - X^2} (P' + Q') - \frac{\frac{1}{6} X}{1 - X^2} (P + Q) = -(\alpha - \alpha^5) \frac{1}{1 - X^2}.$$

[738] (CONCLUDED)
NOTE ON A HYPERGEOMETRIC SERIES.

viz. this is

$$= -\frac{(\alpha - \alpha') X}{1 + X + X^2}.$$

We have

$$\begin{aligned} QP' + PQ' &= \alpha'^2 (-\alpha'' + \alpha X) + \alpha^2 (\alpha - \alpha' X), \\ &= \alpha^3 - \alpha'^3 - (\alpha' - \alpha'') X, \quad = (\alpha - \alpha') X; \end{aligned}$$

and

$$PQ = -\alpha' + (\alpha' + \alpha'') X - \alpha' X^2, \quad = 1 + X + X^2;$$

hence

$$\frac{1}{PQ} (QP' + PQ') = \frac{(\alpha - \alpha') X}{1 + X + X^2},$$

and the sum of the two parts is = 0.

Similarly as regards the second equation, the second part

$$\frac{3X}{1 - X^3} PQ (PQ' + P'Q) - \frac{3X}{1 - X^3} P^2 Q^2$$

is

$$= \frac{3PQX}{1 - X^3} \{(PQ' + P'Q)X - PQ\}.$$

Here $PQ' + P'Q$ is $\alpha(\alpha - \alpha'X) - \alpha'^2(-\alpha' + \alpha X)$, which is $= 1 + 2X$; and PQ being $= 1 + X + X^2$, the term in $\{ \}$ is

$$(1 + 2X)X - (1 + X + X^2), \quad = -(1 - X)(1 + X);$$

hence, outside the $\{ \}$ writing for PQ its value $= 1 + X + X^2$, the term is

$$= \frac{-3X(1 + X + X^2)(1 - X)(1 + X)}{1 - X^3}, \quad = -3X(1 + X),$$

which is the value of the second part in question; the first part is

$$(PQ' + QP')^2 - PQP'Q' \quad = (1 + 2X)^2 - (1 + X + X^2), \quad = 3X(1 + X);$$

and the sum of the two terms is thus = 0.

739.

NOTE ON THE OCTAHEDRON FUNCTION.

[From the *Quarterly Journal of Pure and Applied Mathematics*, vol. xvi. (1879), pp. 280, 281.]

A SEXTIC function

$$U = (a, b, c, d, e, f, g \mid x, y)^6,$$

such that its fourth derivative

$$\begin{aligned} (U, U)_4 = & (ae - 4bd + 3c^2) x^4 \\ & + 2(af - 3be + 2cd) x^3 y \\ & + (ag - 9ce + 8d^2) x^2 y^2 \\ & + 2(bg - 3cf + 2de) xy^3 \\ & + (cg - 4df + 3e^2) y^4 \end{aligned}$$

is identically $= 0$, is considered by Dr Klein, and is called by him the octahedron function. Supposing that by a linear transformation the function is made to contain the factors x, y , or what is the same thing assuming $a = 0, g = 0$, then the equations to be satisfied become

$$-4bd + 3c^2 = 0, \quad -3be + 2cd = 0, \quad -9ce + 8d^2 = 0, \quad -3cf + 2de = 0, \quad -4df + 3e^2 = 0,$$

which are all satisfied if only $c = d = e = 0$; and then assuming, as is allowable,

$$b = -f = 1,$$

we have his canonical form $xy(x^4 - y^4)$ of the octahedron function.

But the equations may be satisfied in a different manner; viz. the first and last equations give

$$b = \frac{3c^2}{4d}, \quad f = \frac{3e^2}{4d},$$

and, substituting these in the remaining equations, they become

$$\frac{c}{4d}(-9ce + 8d^2) = 0, \quad -9ce + 8d^2 = 0, \quad \frac{e}{4d}(-9ce + 8d^2) = 0,$$

all satisfied if only $-9ce + 8d^2 = 0$. Assuming $b = f = 2$, the values are

$$b, c, d, e, f = 2, 2\sqrt{2}, 3, 2\sqrt{2}, 2,$$

and the form is

$$\begin{aligned} & \left(2, 2\sqrt{2}, 3, 2\sqrt{2}, 2 \mid x, y\right)^6. \\ &= xy \left(x^2 + \frac{3}{\sqrt{2}}xy + 2y^2\right) \left(x^2 + \sqrt{2}xy + y^2\right) \\ &= xy \left(x + \frac{1+i}{\sqrt{2}}y\right) \left(x + \frac{1-i}{\sqrt{2}}y\right) \left(x + \sqrt{2}y\right) \left(x + \frac{y}{\sqrt{2}}\right). \end{aligned}$$

This is, in fact, a linear transformation of the foregoing form $XY(X^4 - Y^4)$; for writing

$$X = \left(x + \frac{1+i}{\sqrt{2}}y\right), \quad Y = \left(x + \frac{1-i}{\sqrt{2}}y\right),$$

we have

$$\begin{aligned} X^2 &= x^2 + (1+i)\sqrt{2}xy + iy^2, \\ Y^2 &= x^2 + (1-i)\sqrt{2}xy - iy^2; \end{aligned}$$

and therefore

$$\begin{aligned} X^2 + Y^2 &= 2\{x + \sqrt{2}y\}, \\ X^2 - Y^2 &= 2i\sqrt{2}y \left(x + \frac{y}{\sqrt{2}}\right), \end{aligned}$$

or finally

$$XY(X^4 - Y^4) = 4i\sqrt{2}xy \left(x + \frac{1+i}{\sqrt{2}}y\right) \left(x + \frac{1-i}{\sqrt{2}}y\right) \left(x + \sqrt{2}y\right) \left(x + \frac{y}{\sqrt{2}}\right),$$

and the two forms are thus identical.

740.

ON CERTAIN ALGEBRAICAL IDENTITIES.

[From the *Quarterly Journal of Pure and Applied Mathematics*, vol. xvi. (1879), pp. 281, 282.]

If P_0, P_1, P_2 are points on a circle, say the circle $x^2 + y^2 = 1$, then it is possible to find functions of (P_0, P_1) and of (P_1, P_2) respectively, which are really independent of P_1 , and consequently functions of only P_0 and P_2 : the expression “function of a point or points” being here used to mean algebraical function of the coordinates of the point or points. Thus the functions of (P_0, P_1) and of (P_1, P_2) being $x_0x_1 + y_0y_1$, $x_0y_1 - x_1y_0$, and $x_1x_2 + y_1y_2$, $x_1y_2 - x_2y_1$, we have

$$(x_0x_1 + y_0y_1)(x_1x_2 + y_1y_2) + (x_0y_1 - x_1y_0)(x_1y_2 - x_2y_1) = x_0x_2 + y_0y_2,$$

and another like equation. This depends obviously on the circumstance that the coordinates of a point of the circle are expressible by means of the functions $\sin, \cos$, $x = \cos u$, $y = \sin u$ and the identity written down is obtained by expressing the cosine of $u_2 - u_0 = (u_2 - u_1) + (u_1 - u_0)$, in terms of the cosines and sines of $u_2 - u_1$ and $u_1 - u_0$.

Evidently the like property holds good for a curve, such that the coordinates of any point of it can be expressed by means of “additive” functions of a parameter u ; where, by an additive function $f(u)$, is meant a function such that $f(u + v)$ is an algebraical function of $f(u), f(v)$: the sine and cosine are each of them an additive function, because

$$\sin(u + v) = \sin u \sqrt{1 - \sin^2 v} + \sin v \sqrt{1 - \sin^2 u},$$

and, similarly, for the cosine. But it is convenient to consider pairs or groups $f(u), \phi(u), \dots$, where $f(u + v), \phi(u + v), \dots$ are each of them an algebraical (rational) function of $f(u), \phi(u), \dots, f(v), \phi(v), \dots$; the sine and cosine are such a group, and so also are the elliptic functions $\text{sn}, \text{cn}, \text{dn}$; but the H and Θ , or say the ϑ -functions generally, are not additive.

In the case of the elliptic functions, we may consider the quadriquadric curve

$$x^2 + y^2 = 1, \quad k^2x^2 + z^2 = 1,$$

so that the coordinates of a point on the curve are $\text{sn } u, \text{cn } u, \text{dn } u$. Taking then P_0, P_1, P_2 , points on the curve, and $(x_0, y_0, z_0), (x_1, y_1, z_1), (x_2, y_2, z_2)$, the coordinates of these points respectively, we have in the same way, from $u_2 - u_0 = (u_2 - u_1) + (u_1 - u_0)$, three equations, of which the first is

$$\frac{x_0y_2z_2 - x_2y_0z_0}{1 - k^2x_0^2x_2^2} = \frac{(1 - k^2x_1^2x_2^2)(x_0y_1z_1 - x_1y_0z_0)(y_1y_2 + x_1z_1x_2z_2)(z_1z_2 + k^2x_1y_1x_2y_2) + (1 - k^2x_0^2x_1^2)(x_1y_2z_2 - x_2y_1z_1)(y_0y_1 + x_0z_0x_1z_1)(z_0z_1 + k^2x_0y_0x_1y_1)}{(1 - k^2x_0^2x_1^2)^2(1 - k^2x_1^2x_2^2)^2 - k^2(x_0y_1z_1 - x_1y_0z_0)^2(x_1y_2z_2 - x_2y_1z_1)^2}.$$

The form of the right-hand side is

$$\frac{A + Bx_1^2y_1^2z_1^2}{C + Dx_1^2y_1^2z_1^2}$$

where A, B, C, D are each of them rational as regards x_1^2 ; and it is easy to see that the equation can only subsist under the condition that we have separately

$$\frac{x_0y_2z_2 - x_2y_0z_0}{1 - k^2x_0^2x_2^2} = \frac{A}{C} = \frac{B}{D},$$

implying of course the identity $AD - BC = 0$. The values of B and D are found without difficulty; we, in fact, have

$$B = 2k^2(x_2^2 - x_0^2)(x_0y_0z_0 + x_2y_2z_2), \\ D = 2k^2(x_0y_2z_2 + x_2y_0z_0)(x_0^2y_2^2z_2^2 + x_2^2y_0^2z_0^2),$$

so that, comparing the left-hand side with $B \div D$, we have the identity

$$x_0^2y_2^2z_2^2 - x_2^2y_0^2z_0^2 = (x_2^2 - x_0^2)(1 - k^2x_0^2x_2^2),$$

which is right. The comparison with $A \div C$ would be somewhat more difficult to effect.

741.

ON A THEOREM OF ABEL'S RELATING TO A QUINTIC EQUATION.

[From the Proceedings of the Cambridge Philosophical Society, vol. iii. (1880), pp. 155–159.]

The theorem in question is given, Œuvres Complètes, [Christiania, 1881], t. ii., p. 266, as an extract from a letter to Crelle dated 14th March, 1826, as follows:

“Si une équation du cinquième degré dont les coefficients sont des nombres rationnels est résoluble algébriquement, on peut donner aux racines la forme suivante

$$x = c + Aa^{\frac{1}{5}}a_1^{\frac{2}{5}}a_2^{\frac{3}{5}}a_3^{\frac{4}{5}} + A_1a_1^{\frac{1}{5}}a_2^{\frac{2}{5}}a_3^{\frac{3}{5}}a^{\frac{4}{5}} + A_2a_2^{\frac{1}{5}}a_3^{\frac{2}{5}}a^{\frac{3}{5}}a_1^{\frac{4}{5}} + A_3a_3^{\frac{1}{5}}a^{\frac{2}{5}}a_1^{\frac{3}{5}}a_2^{\frac{4}{5}},$$

où

$$\begin{aligned} a &= m + n\sqrt{1+e^2} + \sqrt{h(1+e^2+\sqrt{1+e^2})}, \\ a_1 &= m - n\sqrt{1+e^2} + \sqrt{h(1+e^2-\sqrt{1+e^2})}, \\ a_2 &= m + n\sqrt{1+e^2} - \sqrt{h(1+e^2+\sqrt{1+e^2})}, \\ a_3 &= m - n\sqrt{1+e^2} - \sqrt{h(1+e^2-\sqrt{1+e^2})}, \\ A &= K + K'a + K''a_2 + K'''aa_2, \quad A_1 = K + K'a_1 + K''a_3 + K'''a_1a_3, \\ A_2 &= K + K'a_2 + K''a + K'''a_2a, \quad A_3 = K + K'a_3 + K''a_1 + K'''a_3a_1. \end{aligned}$$

Les quantités $c, h, e, m, n, K, K', K'', K'''$ sont des nombres rationnels. Mais de cette manière l'équation $x^5 + ax + b = 0$ n'est pas résoluble tant que a et b sont des quantités quelconques. J'ai trouvé de pareils théorèmes pour les équations du 7^{ième}, 11^{ième}, 13^{ième}, etc. degré.”

It is easy to see that x is the root of a quintic equation, the coefficients of which are rational and integral functions of a, a_1, a_2, a_3 : these coefficients are not symmetrical functions of a, a_1, a_2, a_3 , but they are functions which remain unaltered by the cyclical change a into a_1, a_1 into a_2, a_2 into a_3, a_3 into a . But the coefficients of the quintic equation must be rational functions of $c, h, e, m, n, K, K', K'', K'''$; hence regarding a, a_1, a_2, a_3 , as the roots of a quartic equation, the coefficients of this equation being rational functions of m, n, e, h , this equation must be such that every rational function of the roots, unchanged by the aforesaid cyclical change of the roots, shall be rationally expressible in terms of these quantities m, n, e, h : or, what is the same thing, the group of the quartic equation, using the term “group of the equation” in the sense assigned to it by Galois, must be $aa_1a_2a_3, a_1a_2a_3a, a_2a_3aa_1, a_3aa_1a_2$. And conversely, the quartic equation being of this form, x will be the root of a quintic equation, the coefficients whereof are rational and integral functions of $c, h, e, m, n, K, K', K'', K'''$.

To investigate the form of a quartic equation having the property just referred to, let it be proposed to find γ, γ' functions of e, h , such that $\gamma^2 + \gamma'^2$ is a rational function of e, h , but that $\gamma^2 - \gamma'^2, \gamma\gamma'$ are rational multiples of the same quadric radical $\sqrt{\theta}$. Assume that we have

$$\gamma^2 = p + q\sqrt{\theta}, \quad \gamma'^2 = p - q\sqrt{\theta};$$

then

$$(\gamma^2 + \gamma'^2)^2 = 4(p^2 + q^2)\theta;$$

that $\gamma^2 + \gamma'^2$ may be rational, we must have $p^2 + q^2 = \lambda^2\theta$, or say $p^2 + q^2 = h^2\theta$; hence, $\theta = \frac{p^2}{h^2} + \frac{q^2}{h^2}$ must be a sum of two squares, or, assuming one of these equal to unity and the other of them equal to e^2 , say $\theta = 1 + e^2$, we satisfy the required equation by taking $p = h, q = he$: viz. we thus have

$$\gamma^2 - \gamma'^2 = 2h\sqrt{1+e^2}, \quad \gamma\gamma' = he\sqrt{1+e^2}, \quad \gamma^2 + \gamma'^2 = 2h(1+e^2),$$

and thence also

$$\gamma^2 = h(1+e^2 + \sqrt{1+e^2}), \quad \gamma'^2 = h(1+e^2 - \sqrt{1+e^2}),$$

the roots of these expressions, or values of γ, γ' , being such that

$$\gamma\gamma' = he\sqrt{1+e^2}.$$

Taking now α rational, $= m$ suppose, and β a rational multiple of

$$\sqrt{1+e^2}, \quad = n\sqrt{1+e^2},$$

suppose; it is easy to see that the quartic equation which has for its roots

$$a, a_1, a_2, a_3 = \alpha + \beta + \gamma, \quad \alpha - \beta + \gamma', \quad \alpha + \beta - \gamma, \quad \alpha - \beta - \gamma',$$

has the property in question, viz. that every rational function of the roots unchangeable by the cyclical change a into a_1 , a_1 into a_2 , a_2 into a_3 , a_3 into a , is rationally expressible in terms of e, h, m, n .

It will be sufficient to give the proof in the case of a rational and integral function; such a function, unchangeable as aforesaid, is of the form

$$\phi(a, a_1, a_2, a_3) + \phi(a_1, a_2, a_3, a) + \phi(a_2, a_3, a, a_1) + \phi(a_3, a, a_1, a_2);$$

and if $\phi(a, a_1, a_2, a_3)$ contains a term $\alpha^m \beta^n \gamma^p \gamma'^q$, then the other three functions will contain respectively the terms

$$\alpha^m (-\beta)^n \gamma'^p (-\gamma)^q, \quad \alpha^m \beta^n (-\gamma)^p (-\gamma')^q, \quad \alpha^m (-\beta)^n (-\gamma')^p \gamma^q;$$

viz. the sum of the four terms is

$$= \alpha^m \beta^n [\{1 + (-)^{p+q}\} \gamma^p \gamma'^q + \{(-)^n + (-)^{n+p}\} \gamma'^p \gamma^q].$$

This obviously vanishes unless p and q are both even, or both odd; and the cases to be considered are 1° , n even, p and q even; 2° , n odd, p and q even; 3° , n even, p and q odd; 4° , n odd, p and q odd. Writing, for greater distinctness, $2n$ or $2n+1$ for n , according as n is even or odd, and similarly for p and q , the term is, in the four cases respectively,

$$\begin{aligned} &= 2\alpha^m \beta^{2n} (\gamma^{2p} \gamma'^{2q} + \gamma^{2q} \gamma'^{2p}), \\ &= 2\alpha^m \beta^{2n+1} (\gamma^{2p} \gamma'^{2q} - \gamma^{2q} \gamma'^{2p}), \\ &= 2\alpha^m \beta^{2n} \gamma \gamma' (\gamma^{2p} \gamma'^{2q} + \gamma^{2q} \gamma'^{2p}), \\ &= 2\alpha^m \beta^{2n+1} \gamma \gamma' (\gamma^{2p} \gamma'^{2q} - \gamma^{2q} \gamma'^{2p}). \end{aligned}$$

The second, third, and fourth expressions contain the factors

$$\beta(\gamma^2 - \gamma'^2), \quad \gamma \gamma'(\gamma^2 - \gamma'^2), \quad \beta \gamma \gamma',$$

respectively; and the first expression as it stands, and the other three divested of these factors respectively are rational functions of $\alpha, \beta^2, \gamma^2, \gamma'^2$, that is, they are rational functions of m, n, e, h . But the omitted factors $\beta(\gamma^2 - \gamma'^2), \gamma \gamma'(\gamma^2 - \gamma'^2), \beta \gamma \gamma' = 2nh(1+e^2), 2h^2e(1+e^2), nhe(1+e^2)$ are rational functions of n, h, e ; hence each of the original four expressions is a rational function of m, n, h, e ; and the entire function

$$\phi(a, a_1, a_2, a_3) + \phi(a_1, a_2, a_3, a) + \phi(a_2, a_3, a, a_1) + \phi(a_3, a, a_1, a_2)$$

is a rational function of m, n, h, e .

Replacing $\alpha, \beta, \gamma, \gamma'$ by their values, the roots of the quartic equation are

$$\begin{aligned} &m + n\sqrt{1+e^2} + \sqrt{h(1+e^2 + \sqrt{1+e^2})}, \\ &m - n\sqrt{1+e^2} + \sqrt{h(1+e^2 - \sqrt{1+e^2})}, \\ &m + n\sqrt{1+e^2} - \sqrt{h(1+e^2 + \sqrt{1+e^2})}, \\ &m - n\sqrt{1+e^2} - \sqrt{h(1+e^2 - \sqrt{1+e^2})}. \end{aligned}$$

And I stop to remark that taking $m, n, e, h = -\frac{1}{4}, +\frac{1}{4}, 2, -\frac{5}{16}$ respectively, the roots are

$$\begin{aligned} &-\frac{1}{4} + \frac{1}{4}\sqrt{5} + \sqrt{-\frac{1}{8}(5 + \sqrt{5})}, \\ &-\frac{1}{4} - \frac{1}{4}\sqrt{5} + \sqrt{-\frac{1}{8}(5 - \sqrt{5})}, \\ &-\frac{1}{4} + \frac{1}{4}\sqrt{5} - \sqrt{-\frac{1}{8}(5 + \sqrt{5})}, \\ &-\frac{1}{4} - \frac{1}{4}\sqrt{5} - \sqrt{-\frac{1}{8}(5 - \sqrt{5})}; \end{aligned}$$

viz. these are the imaginary fifth roots of unity, or roots r, r^2, r^3, r^4 of the quartic equation $x^4 + x^3 + x^2 + x + 1 = 0$; which equation, as is well known, has the group $rr^2r^3r^4, r^2r^4rr^3, r^3rr^4r^2, r^4r^3r^2r$.

Reverting to Abel's expression for x , and writing this for a moment in the form

$$x = c + p + s + r + q,$$

the quintic equation in x is

$$\begin{aligned} 0 = (x - c)^5 &+ (x - c)^3 \cdot -5(pr + qs) \\ &+ (x - c)^2 \cdot -5(p^2s + q^2p + r^2s + s^2r) \\ &+ (x - c) \cdot -5(p^2q + q^2r + r^2s + s^2p) + 5(p^2r^2 + q^2s^2) - 5pqrs \\ &+ (x - c)^0 \cdot -(p^5 + q^5 + r^5 + s^5) \\ &\quad + 5(p^3rs + q^3sp + r^3pq + s^3qr) \\ &\quad - 5(p^2q^2s + q^2r^2p + r^2s^2q + s^2p^2r). \end{aligned}$$

If we substitute herein for p, q, r, s their values, then, altering the order of the terms, the final result is found to be

$$\begin{aligned} 0 = (x - c)^5 &+ (x - c)^3 \cdot -5(AA_1 + A_2A_3)aa_1a_2a_3 \\ &+ (x - c)^2 \cdot -5(A^2A_1a_1a_2a_3 + A_1^2A_2a_2a_3a + A_2^2A_3a_3aa_1 + A_3^2Aaa_1a_2)aa_1a_2a_3 \\ &+ (x - c) \cdot -5(A^2A_2a_1a_3 + A_1^2A_3a_2a + A_2^2Aa_3a_1 + A_3^2A_1aa_2)aa_1a_2a_3 \\ &\quad + 5(A^2A_2^2 + A_1^2A_3^2 - AA_1A_2A_3)(aa_1a_2a_3)^2 \\ &+ (x - c)^0 \cdot -(A^5a_1a_2^2a_3^3 + A_1^5a_2a_3^2a^3 + A_2^5a_3a^2a_1^3 + A_3^5aa_1^2a_2^3)aa_1a_2a_3 \\ &\quad + 5(A^3A_2A_3a_1a_2a_3 + A_1^3A_3Aa_2a_3a + A_2^3AA_1a_3aa_1 + A_3^3A_1A_2aa_1a_2)(aa_1a_2a_3)^2 \\ &\quad - 5(A^3A_1^2A_2a_1a_3 + A_1^3A_2^2A_3a_2a + A_2^3A_3^2Aa_3a_1 + A_3^3A^2A_1aa_2)(aa_1a_2a_3)^2, \end{aligned}$$

viz. considering herein A, A_1, A_2, A_3 as standing for their values

$$K + K'a + K''a_2 + K'''aa_2, \quad \&c.$$

respectively, each coefficient is a function of a, a_1, a_2, a_3 , which is unaltered by the cyclical change of these values and therefore is a rational function of

$$m, n, e, h, K, K', K'', K'''.$$

742.

ON THE TRANSFORMATION OF COORDINATES.

[From the *Proceedings of the Cambridge Philosophical Society*, vol. iii. (1880), pp. 178–184.]

The formulae for the transformation between two sets of oblique coordinates assume a very elegant form when presented in the notation of matrices. I call to mind that a matrix denotes a system of quantities arranged in a square form

$$\begin{pmatrix} \alpha & \beta & \gamma \\ \alpha' & \beta' & \gamma' \\ \alpha'' & \beta'' & \gamma'' \end{pmatrix};$$

see my “Memoir on the Theory of Matrices,” *Phil. Trans.* t. CXLVIII. (1858), pp. 17–37, [152]; moreover $(\alpha, \beta, \gamma \mid x, y, z)$ denotes $\alpha x + \beta y + \gamma z$, and so

$$\begin{pmatrix} \alpha & \beta & \gamma \\ \alpha' & \beta' & \gamma' \\ \alpha'' & \beta'' & \gamma'' \end{pmatrix} (x, y, z)$$

denotes

$$(\alpha x + \beta y + \gamma z, \alpha' x + \beta' y + \gamma' z, \alpha'' x + \beta'' y + \gamma'' z),$$

and again

$$(\alpha, \beta, \gamma \mid x, y, z \mid \xi, \eta, \zeta) \text{ denotes } \xi(\alpha x + \beta y + \gamma z) + \eta(\alpha' x + \beta' y + \gamma' z) + \zeta(\alpha'' x + \beta'' y + \gamma'' z).$$

Consequently

$$(\alpha, \beta, \gamma \mid x, y, z \mid \xi, \eta, \zeta) = (\alpha, \alpha', \alpha'' \mid \xi, \eta, \zeta \mid x, y, z).$$

In the case of a symmetrical matrix

$$\begin{pmatrix} a & h & g \\ h & b & f \\ g & f & c \end{pmatrix},$$

the equal expressions

$$(a, h, g \mid x, y, z \mid \xi, \eta, \zeta) = (a, h, g \mid \xi, \eta, \zeta \mid x, y, z)$$

are also written

$$(a, b, c, f, g, h \mid x, y, z \mid \xi, \eta, \zeta), \text{ or } (a, \dots \mid \xi, \eta, \zeta \mid x, y, z).$$

In particular, if $x = \xi, y = \eta, z = \zeta$, then

$$(a, h, g \mid x, y, z)^2 \text{ is written } (a, b, c, f, g, h \mid x, y, z)^2.$$

Two matrices are compounded together according to the law

$$\begin{pmatrix} a & b & c \\ a' & b' & c' \\ a'' & b'' & c'' \end{pmatrix} \begin{pmatrix} \alpha & \beta & \gamma \\ \alpha' & \beta' & \gamma' \\ \alpha'' & \beta'' & \gamma'' \end{pmatrix} = \begin{pmatrix} (a, b, c \mid \alpha, \alpha', \alpha'') & (a, b, c \mid \beta, \beta', \beta'') & (a, b, c \mid \gamma, \gamma', \gamma'') \\ (a', b', c' \mid \alpha, \alpha', \alpha'') & (a', b', c' \mid \beta, \beta', \beta'') & (a', b', c' \mid \gamma, \gamma', \gamma'') \\ (a'', b'', c'' \mid \alpha, \alpha', \alpha'') & (a'', b'', c'' \mid \beta, \beta', \beta'') & (a'', b'', c'' \mid \gamma, \gamma', \gamma'') \end{pmatrix},$$

viz. in the compound matrix, the top-line is

$$(a, b, c \mid \alpha, \alpha', \alpha''), (a, b, c \mid \beta, \beta', \beta''), (a, b, c \mid \gamma, \gamma', \gamma''),$$

and the other two lines are the like functions with (a', b', c') , and (a'', b'', c'') , respectively, in the place of (a, b, c) .

The inverse matrix is the matrix the terms of which are the minors of the determinant formed out of the original matrix, each minor being divided by this determinant, viz.

$$\begin{pmatrix} \alpha & \beta & \gamma \\ \alpha' & \beta' & \gamma' \\ \alpha'' & \beta'' & \gamma'' \end{pmatrix}^{-1} = \frac{1}{\nabla} \begin{pmatrix} \beta' \gamma'' - \beta'' \gamma' & \beta'' \gamma - \beta \gamma'' & \beta \gamma' - \beta' \gamma \\ \gamma' \alpha'' - \gamma'' \alpha' & \gamma'' \alpha - \gamma \alpha'' & \gamma \alpha' - \gamma' \alpha \\ \alpha' \beta'' - \alpha'' \beta' & \alpha'' \beta - \alpha \beta'' & \alpha \beta' - \alpha' \beta \end{pmatrix},$$

where ∇ is the determinant

$$\begin{vmatrix} \alpha & \beta & \gamma \\ \alpha' & \beta' & \gamma' \\ \alpha'' & \beta'' & \gamma'' \end{vmatrix}.$$

Coming now to the question of transformation, write

	x	y	z		x_1	y_1	z_1		x_1	y_1	z_1
x	1	ν	μ	x	α	α'	α''	x_1	1	ν_1	μ_1
y	ν	1	λ	y	β	β'	β''	y_1	ν_1	1	λ_1
z	μ	λ	1	z	γ	γ'	γ''	z_1	μ_1	λ_1	1

	x	y	z
x_1	α	β	γ
y_1	α'	β'	γ'
z_1	α''	β''	γ''

viz. the axes of x, y, z are inclined to each other at angles the cosines whereof are λ, μ, ν , those of x_1, y_1, z_1 are inclined to each other at angles the cosines whereof are λ_1, μ_1, ν_1 , and the cosines of the inclinations of the two sets of axes to each other are $\alpha, \beta, \gamma; \alpha', \beta', \gamma'; \alpha'', \beta'', \gamma''$: as is more clearly indicated in the diagram, the top-line showing that cosine-inclinations of x to

$$x, y, z, x_1, y_1, z_1,$$

are

$$1, \nu, \mu, \alpha, \alpha', \alpha'',$$

respectively, and the like for the other lines of the diagram. The letters Ω, Ω_1, V, W are used to denote matrices, viz. as appearing by the diagram, these are

$$\Omega = \begin{pmatrix} 1 & \nu & \mu \\ \nu & 1 & \lambda \\ \mu & \lambda & 1 \end{pmatrix}, \quad \Omega_1 = \begin{pmatrix} 1 & \nu_1 & \mu_1 \\ \nu_1 & 1 & \lambda_1 \\ \mu_1 & \lambda_1 & 1 \end{pmatrix}, \quad V = \begin{pmatrix} \alpha & \beta & \gamma \\ \alpha' & \beta' & \gamma' \\ \alpha'' & \beta'' & \gamma'' \end{pmatrix}, \quad W = \begin{pmatrix} \alpha & \alpha' & \alpha'' \\ \beta & \beta' & \beta'' \\ \gamma & \gamma' & \gamma'' \end{pmatrix},$$

respectively.

The coordinates (x, y, z) and (x_1, y_1, z_1) form each set a broken line extending from the origin to the point; hence projecting on the axes of x, y, z and on those of x_1, y_1, z_1 respectively, we have two sets, each of three equations, which may be written

$$(\Omega \mid x, y, z) = (W \mid x_1, y_1, z_1), \quad (V \mid x, y, z) = (\Omega_1 \mid x_1, y_1, z_1);$$

where of course each set implies the other set.

We have

$$(x, y, z) = (\Omega^{-1}W \mid x_1, y_1, z_1), \quad = (V^{-1}\Omega_1 \mid x_1, y_1, z_1), \\ (x_1, y_1, z_1) = (W^{-1}\Omega \mid x, y, z), \quad = (\Omega_1^{-1}V \mid x, y, z),$$

the first giving in two forms (x, y, z) as linear functions of (x_1, y_1, z_1) , and the second giving in two forms (x_1, y_1, z_1) as linear functions of (x, y, z) ; comparing the two forms for each set, we have

$$\Omega^{-1}W = V^{-1}\Omega_1, \quad W^{-1}\Omega = \Omega_1^{-1}V,$$

or, what is the same thing,

$$V\Omega^{-1}V = \Omega_1, \quad W\Omega_1^{-1}W = \Omega,$$

where in each equation the two sides are matrices which must be equal term by term to each other; but, the matrices being symmetrical, the equation thus gives (not nine but only) six equations. Writing

$$(a, b, c, f, g, h) = (1 - \lambda^2, 1 - \mu^2, 1 - \nu^2, \mu\nu - \lambda, \nu\lambda - \mu, \lambda\mu - \nu),$$

and

$$K = 1 - \lambda^2 - \mu^2 - \nu^2 + 2\lambda\mu\nu,$$

we have

$$\Omega^{-1} = \frac{1}{K} \begin{pmatrix} a & h & g \\ h & b & f \\ g & f & c \end{pmatrix}.$$

The first equation, written in the form

$$V \begin{pmatrix} a & h & g \\ h & b & f \\ g & f & c \end{pmatrix} V = K\Omega_1,$$

denotes the six equations

$$\begin{aligned} (a, b, c, f, g, h \mid \alpha, \beta, \gamma)^2 &= K, \\ (a, b, c, f, g, h \mid \alpha', \beta', \gamma')^2 &= K, \\ (a, b, c, f, g, h \mid \alpha'', \beta'', \gamma'')^2 &= K, \\ (a, b, c, f, g, h \mid \alpha', \beta', \gamma' \mid \alpha'', \beta'', \gamma'') &= K\lambda_1, \\ (a, b, c, f, g, h \mid \alpha'', \beta'', \gamma'' \mid \alpha, \beta, \gamma) &= K\mu_1, \\ (a, b, c, f, g, h \mid \alpha, \beta, \gamma \mid \alpha', \beta', \gamma') &= K\nu_1. \end{aligned}$$

And, similarly, writing

$$(a_1, b_1, c_1, f_1, g_1, h_1) = (1 - \lambda_1^2, 1 - \mu_1^2, 1 - \nu_1^2, \mu_1\nu_1 - \lambda_1, \nu_1\lambda_1 - \mu_1, \lambda_1\mu_1 - \nu_1),$$

and

$$K_1 = 1 - \lambda_1^2 - \mu_1^2 - \nu_1^2 + 2\lambda_1\mu_1\nu_1,$$

then

$$\Omega_1^{-1} = \frac{1}{K_1} \begin{pmatrix} a_1 & h_1 & g_1 \\ h_1 & b_1 & f_1 \\ g_1 & f_1 & c_1 \end{pmatrix};$$

and the second equation, written in the form

$$W \begin{pmatrix} a_1 & h_1 & g_1 \\ h_1 & b_1 & f_1 \\ g_1 & f_1 & c_1 \end{pmatrix} V = K_1\Omega,$$

denotes the six equations

$$\begin{aligned} (a_1, b_1, c_1, f_1, g_1, h_1 \mid \alpha, \alpha', \alpha'')^2 &= K_1, \\ (a_1, b_1, c_1, f_1, g_1, h_1 \mid \beta, \beta', \beta'')^2 &= K_1, \\ (a_1, b_1, c_1, f_1, g_1, h_1 \mid \gamma, \gamma', \gamma'')^2 &= K_1, \\ (a_1, b_1, c_1, f_1, g_1, h_1 \mid \beta, \beta', \beta'' \mid \gamma, \gamma', \gamma'') &= K_1\lambda, \\ (a_1, b_1, c_1, f_1, g_1, h_1 \mid \gamma, \gamma', \gamma'' \mid \alpha, \alpha', \alpha'') &= K_1\mu, \\ (a_1, b_1, c_1, f_1, g_1, h_1 \mid \alpha, \alpha', \alpha'' \mid \beta, \beta', \beta'') &= K_1\nu. \end{aligned}$$

The two sets each of six equations are, in fact, equivalent to a single set of six equations, and serve to express the relations between the nine cosines

$$(\alpha, \beta, \gamma, \alpha', \beta', \gamma', \alpha'', \beta'', \gamma'')$$

and the cosines (λ, μ, ν) and $(\lambda_1, \mu_1, \nu_1)$. Observe that the nine cosines are not (as in the rectangular transformation) the coefficients of transformation between the two sets of coordinates.

From the original linear relations between the coordinates, multiplying the equations of the first set by x, y, z and adding, and again multiplying the equations of the second set by (x_1, y_1, z_1) and adding, we have

$$\begin{aligned} (\Omega \mid x, y, z)^2 &= (W \mid x_1, y_1, z_1 \mid x, y, z), \\ (\Omega_1 \mid x_1, y_1, z_1)^2 &= (V \mid x, y, z \mid x_1, y_1, z_1). \end{aligned}$$

But

$$(W \mid x_1, y_1, z_1 \mid x, y, z)$$

and

$$(V \mid x, y, z \mid x_1, y_1, z_1)$$

denote one and the same function; hence

$$(\Omega \mid x, y, z)^2 = (\Omega_1 \mid x_1, y_1, z_1)^2,$$

that is,

$$(1, 1, 1, \lambda, \mu, \nu \mid x, y, z)^2 = (1, 1, 1, \lambda_1, \mu_1, \nu_1 \mid x_1, y_1, z_1)^2,$$

or the linear relations between (x, y, z) and (x_1, y_1, z_1) are such as to transform one of these quadric functions into the other: the two quadrics, in fact, denote the squared distance from the origin expressed in terms of the coordinates (x, y, z) and (x_1, y_1, z_1) respectively.

Since the nine cosines are connected by six equations, there should exist values containing three arbitrary constants, and satisfying these equations identically: but, by what just precedes, it appears that the problem of determining these values is, in fact, that of finding the linear transformation between two given quadric functions: the problem of the linear transformation of a quadric function into itself has an elegant solution; but it would seem that this is not the case for the transformation between two different functions.

The foregoing equation

$$K = (a, b, c, f, g, h \mid \alpha, \beta, \gamma)^2$$

is a relation between λ, μ, ν , the cosines of the sides of a spherical triangle, and (α, β, γ) the cosines of the distances of a point P from the three vertices: it can be at once verified by means of the relation $A + B + C = 2\pi$, and thence

$$1 - \cos^2 A - \cos^2 B - \cos^2 C + 2 \cos A \cos B \cos C = 0,$$

which connects the angles A, B, C which the sides subtend at P . Writing a, b, c for λ, μ, ν , and f, g, h for α, β, γ , the relation is

$$\begin{aligned} K &= (a, b, c, f, g, h \mid f, g, h)^2, \\ &= af^2 + bg^2 + ch^2 + 2(bc - a)gh + 2(ca - b)hf + 2(ab - c)fg, \end{aligned}$$

viz. this is

$$1 - a^2 - b^2 - c^2 - f^2 - g^2 - h^2 + 2abc + 2agh + 2bhf + 2cfg - a^2f^2 - b^2g^2 - c^2h^2 + 2bcgh + 2cahf + 2abfg = 0;$$

where (a, b, c, f, g, h) are the cosines of the sides of a spherical quadrangle: $(a, b, c), (a, h, g), (h, b, f), (g, f, c)$ belong respectively to sides forming a triangle, and the remaining sides $(f, g, h), (b, c, f), (c, a, g), (a, b, h)$ are sides meeting in a vertex.

The equation

$$K\nu_1 = (a, b, c, f, g, h \mid \alpha, \beta, \gamma \mid \alpha', \beta', \gamma')$$

is a relation between λ, μ, ν , the cosines of the sides of a spherical triangle; α, β, γ , the cosines of the distances of a point P from the three vertices; α', β', γ' , the cosines of the distances of a point Q from the three vertices; and ν_1 , the cosine of the distance PQ .

Drawing a figure, it is at once seen that

$$\nu_1 = \alpha\alpha' + \sqrt{1 - \alpha^2} \sqrt{1 - \alpha'^2} \cos(\theta - \theta'),$$

where

$$\cos \theta = \frac{\beta - \alpha\nu}{\sqrt{1 - \alpha^2}\sqrt{1 - \nu^2}},$$

and therefore

$$\sin \theta = \frac{\sqrt{V}}{\sqrt{1 - \alpha^2}\sqrt{1 - \nu^2}};$$

also

$$\cos \theta' = \frac{\beta' - \alpha'\nu}{\sqrt{1 - \alpha'^2}\sqrt{1 - \nu^2}},$$

and therefore

$$\sin \theta' = \frac{\sqrt{V'}}{\sqrt{1 - \alpha'^2} \sqrt{1 - \nu'^2}},$$

the values of V, V' being

$$V = 1 - \alpha^2 - \beta^2 - \nu^2 + 2\alpha\beta\nu, \quad V' = 1 - \alpha'^2 - \beta'^2 - \nu'^2 + 2\alpha'\beta'\nu';$$

the resulting value of ν_1 is therefore

$$\nu_1 = \alpha\alpha' + \frac{1}{1 - \nu^2} \{(\beta - \alpha\nu)(\beta' - \alpha'\nu) + \sqrt{V}\sqrt{V'}\}.$$

The equations

$$K = (a, b, c, f, g, h \mid \alpha, \beta, \gamma)^2, \quad K = (a, \dots \mid \alpha', \beta', \gamma')^2$$

give

$$\nabla V = -K^2, \quad \nabla V' = -K^2,$$

and we therefore have

$$(g\alpha + f\beta + c\gamma \mid g\alpha' + f\beta' + c\gamma') = -\frac{1}{c} \sqrt{V}\sqrt{V'};$$

recollecting that $1 - \nu^2 = c$, the formula thus is

$$\nu_1 = \alpha\alpha' + \frac{1}{c} \{(\beta - \alpha\nu)(\beta' - \alpha'\nu) + (g\alpha + f\beta + c\gamma \mid g\alpha' + f\beta' + c\gamma')\},$$

or say,

$$K\nu_1 = K\alpha\alpha' + \frac{1}{c} \{c^2 (\beta - \alpha\nu \mid \beta' - \alpha'\nu) + (g\alpha + f\beta \mid g\alpha' + f\beta')\} + g(\alpha\gamma' + \alpha'\gamma) + f(\beta\gamma' + \beta'\gamma) + c\gamma\gamma'.$$

The sum of the first and second terms is readily found to be

$$= a\alpha\alpha' + b\beta\beta' + h(\alpha\beta' + \alpha'\beta),$$

and the equation thus becomes

$$K\nu_1 = (a, b, c, f, g, h \mid \alpha, \beta, \gamma \mid \alpha', \beta', \gamma'),$$

as it should do.

743.

ON THE NEWTON-FOURIER IMAGINARY PROBLEM.

[From the *Proceedings of the Cambridge Philosophical Society*, vol. *iii*. (1880),
pp. 231, 232.]

The Newtonian process of approximation to the root of a numerical equation $f(u) = 0$, consists in deriving from an assumed approximate root ξ a new value $\xi_1 = \xi - \frac{f(\xi)}{f'(\xi)}$, which should be a closer approximation to the root sought for: taking the coefficients of $f(u)$ to be real, and also the root sought for, and the assumed value ξ , to be each of them real, Fourier investigated the conditions under which ξ_1 is in fact a closer approximation. But the question may be looked at in a more general manner: ξ may be any real or imaginary value, and we have to inquire in what cases the series of derived values

$$\xi_1 = \xi - \frac{f(\xi)}{f'(\xi)}, \quad \xi_2 = \xi_1 - \frac{f(\xi_1)}{f'(\xi_1)}, \quad \dots$$

converge to a root, real or imaginary, of the equation $f(u) = 0$. Representing as usual the imaginary value $\xi, = x + iy$, by means of the point whose coordinates are x, y , and in like manner $\xi_1, = x_1 + iy_1$, &c.,

then we have a problem relating to an infinite plane; the roots of the equation are represented by points $A, B, C, \dots$; the value ξ is represented by an arbitrary point P ; and from this by a determinate geometrical construction we obtain the point P_1 , and thence in like manner the points $P_2, P_3, \dots$ which represent the values $\xi_1, \xi_2, \xi_3, \dots$ respectively. And the problem is to divide the plane into regions, such that, starting with a point P , anywhere in one region, we arrive ultimately at the root A ; anywhere in another region we arrive ultimately at the root B ; and so on for the several roots of the equation. The division into regions is made without difficulty in the case of a quadric equation: but in the next succeeding case, that of a cubic equation, it is anything but obvious what the division is: and the author had not succeeded in finding it.

744.

TABLE OF $\Delta^m 0^n \div \Pi(m)$ UP TO $m = n = 20$.

[From the Transactions of the Cambridge Philosophical Society, vol. xiii. Part i. (1881), pp. 1–4. Read October 27, 1879.]

The differences of the powers of zero, $\Delta^m 0^n$, present themselves in the Calculus of Finite Differences, and especially in the applications of Herschel's theorem,

$$f(e^x) = f(1 + \Delta) e^{0x},$$

for the expansion of the function of an exponential. A small Table up to $\Delta^5 0^n$ is given in Herschel's *Examples* (Camb. 1820), and is reproduced in the treatise on Finite Differences (1843) in the Encyclopaedia Metropolitana. But, as is known, the successive differences $\Delta 0^n, \Delta^2 0^n, \Delta^3 0^n, \dots$ are divisible by $1, 1 \cdot 2, 1 \cdot 2 \cdot 3, \dots$ and generally $\Delta^m 0^n$ is divisible by $1 \cdot 2 \cdot 3 \cdots m, = \Pi(m)$; these quotients are much smaller numbers, and it is therefore desirable to tabulate them rather than the undivided differences $\Delta^m 0^n$: moreover, it is easier to calculate them. A table of the quotients $\Delta^m 0^n \div \Pi(m)$, up to $m = n = 12$ is in fact given by Grünert, *Crelle*, t. xxv. (1843), p. 279, but without any explanation in the heading of the meaning of the tabulated numbers $C_n^m, = \Delta^m 0^n \div \Pi(m)$, and without using for their determination the convenient formula $C_n^{m+1} = n C_n^m + C_{n-1}^m$ given by Björling in a paper, *Crelle*, t. xxviii. (1844), p. 284. The formula in question, say

$$\frac{\Delta^m 0^n}{\Pi(m)} = m \frac{\Delta^{m-1} 0^{n-1}}{\Pi(m)} + \frac{\Delta^{m-1} 0^{n-1}}{\Pi(m-1)},$$

is given in the second edition (by Moulton) of Boole's *Calculus of Finite Differences*, (London, 1872), p. 28, under the form

$$\Delta^m 0^n = m (\Delta^{m-1} 0^{n-1} + \Delta^m 0^{n-1}).$$

It occurred to me that it would be desirable to extend the table of the quotients $\Delta^m 0^n \div \Pi(m)$, up to $m = n = 20$. The calculation is effected very readily by means of the foregoing theorem, which is used in the following form; viz. any column of the table for instance the fifth, being

$A,$	then the following column is	$A,$
$B,$	$\dots$	$2B + A,$
$C,$	$\dots$	$3C + B,$
$D,$	$\dots$	$4D + C,$
$E,$	$\dots$	$5E + D,$
		$+ E;$

and then we obtain a good verification by taking the sum of the terms in the new column, and comparing it with the value as calculated from the formula.

$$\text{Sum} = 2A + 3B + 4C + 5D + 6E.$$

Observe that, in the two calculations, we take successive multiples such as $4D$ and $5D$ of each term of the preceding column, and that the verification is thus a safe-guard against any error of multiplication or addition.

Table, No. 1, of $\Delta^m 0^n \div \Pi(m)$.

n/m	1	2	3	4	5	6	7	8	9	10	11	12	13	14
1	1													
2	1	1												
3	1	3	1											
4	1	7	6	1										
5	1	15	25	10	1									
6	1	31	90	65	15	1								
7	1	63	301	350	140	21	1							
8	1	127	966	1 701	1 050	266	28	1						
9	1	255	3 025	7 770	6 951	2 646	462	36	1					
10	1	511	9 330	34 105	42 525	22 827	5 880	750	45	1				
11	1	1 023	28 501	145 750	246 730	179 487	63 987	11 880	1 155	55	1			
12	1	2 047	86 526	611 501	1 379 400	1 323 652	627 396	159 027	22 275	1 705	66	1		
13	1	4 095	261 625	2 532 530	7 508 501	9 321 312	5 715 424	1 899 612	359 502	39 325	2 431	78	1	
14	1	8 191	788 970	10 391 745	40 075 035	63 436 373	49 329 280	20 912 320	5 135 130	752 752	66 066	3 367	91	1

(Continuation of Table No. 1 for $m = 8, \dots, 13$ and $n = 2, \dots, 20$.)

n/m	8	9	10	11	12	13
2	16 383	32 767	65 535	131 071	262 143	524 287
3	2 375 101	7 141 686	21 457 825	64 439 010	193 448 101	580 606 446
4	48 355 950	171 798 901	694 337 290	2 798 806 985	11 259 666 950	45 232 116 901
5	210 766 920	1 096 190 550	5 652 751 651	28 958 095 545	147 589 284 710	749 206 090 500
6	420 693 273	2 734 926 558	17 505 749 898	110 687 251 039	693 081 601 779	4 306 078 895 384
7	408 741 333	3 281 882 604	25 708 104 786	197 462 483 400	1 492 924 634 839	11 143 554 046 652
8	216 627 840	2 141 764 053	20 415 995 028	189 036 065 010	1 709 751 003 480	15 170 932 662 679
9	67 128 490	820 784 250	9 528 822 303	106 175 396 755	1 144 614 626 805	12 011 282 644 725
10	12 662 650	193 754 990	2 758 334 150	37 112 163 803	477 297 033 785	5 917 584 964 665
11	1 479 478	28 936 908	512 060 978	8 391 004 908	129 413 217 791	1 900 842 429 486
12	106 470	2 757 118	62 022 324	1 256 328 866	23 466 951 300	411 016 633 391
13	4 550	166 620	4 910 178	125 864 638	2 892 439 160	61 068 660 380
14	105	6 020	249 900	8 408 778	243 577 530	6 302 524 580
15	1	120	7 820	367 200	13 916 778	452 329 200
16		1	136	9 996	527 136	22 350 954
17			1	153	12 597	741 285
18				1	171	15 675
19					1	190
20						1

Writing down the sloping lines as columns thus:

1

(0)

2

(2)

3 – 4

(4) (6)

5

(8) (10)

6

(12)

7

8 etc.

(14) etc.

Table, No. 2 (*explanation infra*).

	1	2	3	4	5	6	7
0	1	1	1	1	1	1	1
1		2	6	14	30	62	126
2		1	12	61	240	841	2 772
3			10	124	890	5 060	25 410
4			3	131	1 830	16 990	127 953
5				70	2 226	35 216	401 436
6				15	1 600	47 062	836 976
7					630	40 796	1 196 532
8					105	21 225	1 182 195
9						10 930	795 718
10						945	349 020
11							90 090
12							10 395

We have, by means of this Table, the general expressions of $\Delta^r 0^r$, $\Delta^{r-1} 0^r$, $\Delta^{r-2} 0^r$, ... up to $\Delta^{r-6} 0^r$, viz. the formulae are

$$\Delta^r 0^r \div \Pi(r) = 1,$$
$$\Delta^{r-1} 0^r \div \Pi(r-1) = 1 + 2 \binom{r-1}{1} + 1 \binom{r-1}{2},$$
$$\Delta^{r-2} 0^r \div \Pi(r-2) = 1 + 6 \binom{r-2}{1} + 12 \binom{r-2}{2} + 10 \binom{r-2}{3} + 3 \binom{r-2}{4},$$
$$\&c., \&c.,$$

where the numerical coefficients are the numbers in the successive columns of the table; and where for shortness $\binom{r-m}{k}$ is written to denote the binomial coefficient $\frac{[r-m]^k}{[k]^k}$. For instance, $r = 10$, we have

$$\Delta^8 0^{10} \div \Pi(8) = 1 + 6 \cdot 7 + 12 \cdot 21 + 10 \cdot 35 + 3 \cdot 35, = 750,$$

agreeing with the principal Table. It will be observed that, in the successive columns of the Table, the last terms are 1, 1, 1 · 3, 1 · 3 · 5, 1 · 3 · 5 · 7, 1 · 3 · 5 · 7 · 9, and 1 · 3 · 5 · 7 · 9 · 11. This is itself a good verification: I further verified the last column by calculating from it the value of $\Delta^{14} 0^{20} \div \Pi(14)$, = 6 302 524 580 as above. The Table shows that we have $\Delta^{r-m} 0^r \div \Pi(r-m)$ given as an algebraical rational and integral function of r , of the degree $2m$. But the terms from the top of a column, $\Delta 0^r = 1$, $\Delta^2 0^r \div 1 \cdot 2 = 2^{r-1} - 1$, &c., are not algebraical functions of r .

22 October, 1879.

745.

ON THE SCHWARZIAN DERIVATIVE, AND THE POLYHEDRAL
FUNCTIONS.

[From the Transactions of the Cambridge Philosophical Society, vol. xiii. Part i. (1881),
pp. 5-68. Read March 8, 1880.]

The quotient s of any two solutions of a linear partial differential equation of the second order, $\frac{d^2 v}{dx^2} + p \frac{dv}{dx} + qv = 0$, is determined by a differential equation of the third order

$$\frac{\frac{d^3 s}{dx^3}}{\frac{ds}{dx}} - \frac{3}{2} \left(\frac{\frac{d^2 s}{dx^2}}{\frac{ds}{dx}} \right)^2 = -\frac{1}{2} \left(p^2 + 2 \frac{dp}{dx} - 4q \right),$$

where the function on the left-hand is what I call the Schwarzian Derivative; or say this derivative is

$$\{s, x\} = \frac{s'''}{s'} - \frac{3}{2} \left(\frac{s''}{s'} \right)^2,$$

where the accents denote differentiations in regard to the second variable x of the symbol.

Writing in general $(a, b, c \dots | X, Y, Z)^2$ to denote a quadric function

$$(a, b, c, \frac{1}{2}(a-b-c), \frac{1}{2}(-a+b-c), \frac{1}{2}(-a-b+c) | X, Y, Z)^2,$$

then, if the equation of the second order be that of the hypergeometric series, generalised by a homographic transformation upon the variable x , the resulting differential equation of the third order is of the form

$$\{s, x\} = (a, b, c \dots) \left(\frac{1}{x-a}, \frac{1}{x-b}, \frac{1}{x-c} \right)^2,$$

and, presenting themselves in connexion with the algebraically integrable cases of this equation, we have rational and integral functions of s , derived from the polygon, the double pyramid, and the five regular solids. They are called Polyhedral Functions.

The Schwarzian Derivative occurs implicitly in Jacobi's differential equation of the third order for the modulus in the transformation of an elliptic function (*Fund. Nova*, 1829, p. 79, [*Ges. Werke*, t. i., p. 133]) and in Kummer's fundamental equation for the transformation of a hypergeometric series (Kummer, 1836: see list of Memoirs): but it was first explicitly considered and brought into notice in the two Memoirs of Schwarz¹, 1869 and 1873. The latter of these, relating to the algebraic integration of the differential equation for the hypergeometric series, is the fundamental Memoir upon the subject, but the theory is in some material points completed in the Memoirs by Klein and Brioschi.

The following list of Memoirs, relating as well to the Polyhedral Functions as to the Schwarzian Derivative, is arranged nearly in chronological order.

KUMMER, Ueber die hypergeometrische Reihe $1 + \frac{\alpha \cdot \beta}{1 \cdot \gamma} x + \dots$ *Crelle*, t. xv. (1836), pp. 39–83 and 127–172.

SCHWARZ, Ueber einige Abbildungsaufgaben. *Crelle-Borchardt*, t. lxx. (1869), pp. 105–120.

Ueber diejenigen Fälle in welchen die Gaussische hypergeometrische Reihe eine algebraische Function ihres vierten Elementes darstellt. *Do.* t. lxxv. (1873), pp. 292–335.

CAYLEY, Notes on Polyhedra. *Quart. Math. Jour.* t. vii. (1866), pp. 304–316; [375].

On the Regular Solids. *Do.* t. xv. (1878), pp. 127–131; [679].

FUCHS, Ueber diejenigen Differentialgleichungen zweiter Ordnung welche algebraische Integralen besitzen, und eine Anwendung der Invariantentheorie. *Crelle-Borchardt*, t. lxxxi. (1875), pp. 97–142.

KLEIN, Ueber binäre Formen mit linearen Transformationen in sich selbst. *Math. Ann.* t. ix. (1875), pp. 183–209.

BRIOSCHI, Extrait d'une lettre à M. Klein. *Math. Ann.* t. xi. (1877), pp. 111–114.

KLEIN, Ueber lineare Differentialgleichungen. *Math. Ann.* t. xi. (1877), pp. 115–118.

BRIOSCHI, La théorie des formes dans l'intégration des équations différentielles linéaires du second ordre. *Math. Ann.* t. xi. (1877), pp. 401–411.

GORDAN, Ueber endliche Gruppen linearer Transformationen einer Veränderlichen. *Math. Ann.* t. xii. (1877), pp. 23–46.

Binäre Formen mit verschwindenden Covarianten. *Math. Ann.* t. xii. (1877), pp. 147–166.

KLEIN, Ueber lineare Differentialgleichungen. *Math. Ann.* t. xii. (1877), pp. 167–179.

Weitere Untersuchungen über das Icosaeder. *Math. Ann.* t. xii. (1877), pp. 503–560.

CAYLEY, On the Correspondence of Homographies and Rotations. *Math. Ann.* t. xv. (1879), pp. 238–240; [660].

On the finite Groups of linear transformations of a Variable. *Math. Ann.* t. xvi. (1880), pp. 260–263, and pp. 439–440; [752].

I propose in the present Memoir to consider the whole theory: and, in particular, to give some additional developments in regard to the Polyhedral Functions.

¹Schwarz, *Ges. Werke*, t. ii., p. 351, remarks that the Derivative occurs implicitly in a memoir by Lagrange, "Sur la construction des cartes géographiques," (1779), *Œuvres*, t. iv., p. 651.

I remark that Schwarz starts with the foregoing differential equation of the third order

$$\{s, x\} = (a, b, c \dots) \left(\frac{1}{x-a}, \frac{1}{x-b}, \frac{1}{x-c} \right)^2,$$

and he shows (by very refined reasoning founded on the theory of conformable figures, which will be in part reproduced) that this equation is, in fact, algebraically integrable for 16 different sets of values of the coefficients a, b, c . It may I think be taken to be part of his theory, although not very clearly brought out by him, that these integrals are some of them of the form, $x =$ rational function of s : others of the form, rational function of $x =$ rational function of s ; the rational functions of s being in fact the same in the last as in the first set of solutions: they are quotients of Polyhedral functions.

But as regards the second set of cases, the solution of these, introducing for convenience a new variable z in place of s , may be made to depend upon the solution in the form, $x =$ rational function of z , of an equation of a somewhat similar form, but involving two quadric functions of x and z respectively, viz. the equation

$$\{x, z\} = (a, b, c \dots) \left(\frac{1}{x-a}, \frac{1}{x-b}, \frac{1}{x-c} \right)^2 = (\alpha, \beta, \gamma \dots) \left(\frac{1}{z-\alpha}, \frac{1}{z-\beta}, \frac{1}{z-\gamma} \right)^2,$$

and we have the theorem that the solution of this equation depends upon the determination of P, Q, R rational and integral functions of z , containing each of them multiple factors, which are such that $P + Q + R = 0$. Using accents to denote differentiation in regard to z , this implies $P' + Q' + R' = 0$, and consequently

$$QR' - Q'R = RP' - R'P = PQ' - P'Q.$$

Further, they are such that the equal functions $QR' - Q'R, RP' - R'P, PQ' - P'Q$ contain only factors which are factors of P, Q or R .

In fact, writing $f, g, h = b - c, c - a, a - b$, the required relation between x, z is then expressed in the symmetrical form $f(x - a) : g(x - b) : h(x - c) = P : Q : R$.

The last-mentioned differential equation is considered by Klein and Brioschi: the solutions in 13 cases, or such of them as had not been given by Schwarz, were obtained by Brioschi: and those of the remaining 3 cases, subject to a correction in one of them, were afterwards obtained by Klein.

The first part of the present Memoir relates, say to the foregoing equation

$$\{s, x\} = (a, b, c \dots) \left(\frac{1}{x-a}, \frac{1}{x-b}, \frac{1}{x-c} \right)^2,$$

although the other form in $\{x, z\}$ may equally well be regarded as the fundamental form.

We consider in the theory:

- A. The Derivative $\{s, x\}$, meaning as above explained.
- B. Quadric functions of any three or more inverts $\frac{1}{x-l}$.
- C. Rational and integral functions P, Q, R having a sum $= 0$, and which are such that $QR' - Q'R, = RP' - R'P, = PQ' - P'Q$, contains only the factors of P, Q, R .
- D. The differential equation of the third order.
- E. The Schwarzian theory in regard to conformable figures and the corresponding values of the imaginary variables s and x .
- F. Connexion with the differential equation for the hypergeometric series.

The second part of the Memoir relates to the Polyhedral Functions.

The paragraphs of the whole Memoir are numbered consecutively.

PART I.

The Derivative $\{s, x\}$. Art. Nos. 1 to 7.

2. The derivative $\{s, x\}$ may be transformed in regard to either or both of the variables.

Suppose, first, that s is a function of the new variable S , (hence also S is a function of x): using subscript numbers to denote differentiations in regard to S , and the accents as before for differentiations in regard to x , we have

$$s' = S' s_1,$$

whence, differentiating the logarithms,

$$\frac{s''}{s'} = S' \frac{s_2}{s_1} + \frac{S''}{S'},$$

Again differentiating, we have

$$\frac{s'''}{s'} - \left(\frac{s''}{s'}\right)^2 = S'^2 \left[\frac{s_3}{s_1} - \left(\frac{s_2}{s_1}\right)^2 \right] + 3S' \frac{s_2}{s_1} \frac{S''}{S'} - \left(\frac{S''}{S'}\right)^2.$$

But

$$-\frac{1}{2} \left(\frac{s''}{s'}\right)^2 = S'^2 \left[-\frac{1}{2} \left(\frac{s_2}{s_1}\right)^2 \right] - S' \frac{s_2}{s_1} \frac{S''}{S'} - \frac{1}{2} \left(\frac{S''}{S'}\right)^2,$$

and consequently

$$\frac{s'''}{s'} - \frac{3}{2} \left(\frac{s''}{s'}\right)^2 = S'^2 \left[\frac{s_3}{s_1} - \frac{3}{2} \left(\frac{s_2}{s_1}\right)^2 \right] + \frac{S'''}{S'} - \frac{3}{2} \left(\frac{S''}{S'}\right)^2;$$

that is,

$$\{s, x\} = \left(\frac{dS}{dx}\right)^2 \{s, S\} + \{S, x\},$$

the required formula.

In a very similar manner, taking x a function of X , it is shown that

$$\{s, x\} = \left(\frac{dX}{dx}\right)^2 \{s, X\} - \{x, X\}.$$

3. If in this formula we write S for s , and substitute the resulting value of $\{S, x\}$ in the former formula, we have

$$\{s, x\} = \left(\frac{dS}{dx}\right)^2 \{s, S\} + \left(\frac{dX}{dx}\right)^2 \{S, X\} - \{x, X\},$$

which is the formula for the change of both variables. It in fact includes the other two: viz. writing $X = x$, or $S = x$, and observing that $\{x, x\} = 0$, we have the other two formulæ.

4. By putting in the first formula $S = x$, we obtain

$$\{x, s\} = - \left(\frac{ds}{dx}\right)^2 \{s, x\},$$

a formula for the interchange of the variables.

5. Writing $S = \frac{as+b}{cs+d}$, and using for a moment the accents to denote differentiation in regard to s , we have

$$S' = \frac{ad-bc}{(cs+d)^2}, \quad \frac{S''}{S'} = \frac{-2c}{cs+d},$$

and thence

$$\begin{aligned} \frac{S'''}{S'} - \left(\frac{S''}{S'}\right)^2 &= \frac{2c^2}{(cs+d)^2}, \\ -\frac{1}{2} \left(\frac{S''}{S'}\right)^2 &= \frac{-2c^2}{(cs+d)^2}. \end{aligned}$$

Consequently $\{S, s\} = 0$, whence also $\{s, S\} = 0$.

Hence in the first formula $\{S, x\} = \{s, x\}$, that is,

$$\left\{ \frac{as+b}{cs+d}, x \right\} = \{s, x\};$$

viz. we may, in the derivative $\{s, x\}$, write for s any homographic function $\frac{as+b}{cs+d}$ of s .

6. Again, if $X = \frac{\alpha x + \beta}{\gamma x + \delta}$, then from the second formula

$$\{s, x\} = \frac{(\alpha\delta - \beta\gamma)^2}{(\gamma x + \delta)^4} \{s, X\};$$

that is,

$$\left\{ s, \frac{\alpha x + \beta}{\gamma x + \delta} \right\} = \frac{(\gamma x + \delta)^4}{(\alpha\delta - \beta\gamma)^2} \{s, x\};$$

and here, changing s into $(as+b) \div (cs+d)$, we have finally

$$\left\{ \frac{as+b}{cs+d}, \frac{\alpha x + \beta}{\gamma x + \delta} \right\} = \frac{(\gamma x + \delta)^4}{(\alpha\delta - \beta\gamma)^2} \{s, x\},$$

which is the formula for the homographic transformation of the two variables s, x .

7. Let s be a given function of x , the equation $\{S, x\} = \{s, x\}$ is a differential equation of the third order in S , and by what precedes, its general integral is $S = \frac{as+b}{cs+d}$.

The direct process is as follows: we have a first integral $\frac{S''}{S'} = \frac{s''}{s'} - \frac{2cs'}{cs+d}$; a second integral $\log S' = \log s' - 2\log(cs+d) + \text{const.}$, that is, $S' = \frac{cAs'}{(cs+d)^2}$; and thence a final integral $S = B - \frac{A}{cs+d}$, which is equivalent to the foregoing value of S .

The Quadric Function of three or more Inverts. Art. Nos. 8 to 15.

8. We consider a quadric function of any number of invert $\frac{1}{x-a}, \frac{1}{x-\beta}, \dots$, all of them different: it is assumed that the constant term is $= 0$, and also that the sum of the coefficients of the linear terms is $= 0$. We have therefore square terms $\frac{a}{(x-a)^2}, \dots$, product terms $\frac{h}{x-a \cdot x-\beta}$, and linear terms $\frac{A}{x-a}$, where the sum of the coefficients A is $= 0$. Any product term $\frac{h}{x-a \cdot x-\beta}$ is expressible in the form of a difference $\frac{h}{a-\beta} \left(\frac{1}{x-a} - \frac{1}{x-\beta} \right)$ of two linear terms, and (the coefficients of these

being equal), after it is thus expressed, the sum of the coefficients of the linear terms is still $= 0$. The function is thus always expressible in the form

$$\frac{a}{(x-a)^2} + \frac{b}{(x-\beta)^2} + \cdots + \frac{A}{x-a} + \frac{B}{x-\beta} + \cdots,$$

where the sum $A + B + \cdots$ is $= 0$: this may be called the reduced form.

9. Observe that any particular invert $\frac{1}{x-a}$ may disappear altogether from the reduced form: this will be the case if $a = 0$, that is, if the original form contains no term in $\frac{1}{(x-a)^2}$, and if also $A = 0$. An invert thus disappearing from the reduced form is said to be non-essential: and the inverts which do not disappear are said to be essential. The original form contains in appearance the non-essential inverts, but it is really a quadric function of the essential inverts only.

10. Imagine the original function expressed as a rational fraction, the denominator being the product $(x-a)^2(x-\beta)^2(x-\gamma)^2 \cdots$ of the squared factors corresponding to all the inverts (non-essential as well as essential): the numerator will be in general of a degree less by 2 than that of the denominator, but the coefficients of any one or more of the higher powers of x may vanish, and the numerator will then be of a lower degree. But this numerator will for any non-essential invert $\frac{1}{x-\gamma}$ contain the factor $(x-\gamma)^2$, or, dividing the numerator and denominator each by this factor, the difference of the degrees of the numerator and denominator will remain unaltered; that is, the difference will have the same value whether we do or do not attend to the non-essential inverts; or say it will have the same value for the original form and for the reduced form.

11. It is to be remarked that the linear terms $\frac{A}{x-a} + \frac{B}{x-\beta} + \frac{C}{x-\gamma} + \cdots$, where $A+B+C+\cdots = 0$, can be (and that in a variety of ways) expressed as a sum of differences $\frac{1}{x-a} - \frac{1}{x-\beta}$, that is, as a sum of product-terms $\frac{1}{x-a \cdot x-\beta}$. Hence the quadric function can be (and that in a variety of ways) expressed as a homogeneous function $(a, b, \dots \mid \frac{1}{x-a}, \frac{1}{x-\beta}, \dots)^2$; we must have in the form all the essential inverts, and we need have these only. Supposing that this is so, and that the number of the essential inverts is $= n$, then the number of constants is $= \frac{1}{2}n(n+1)$, whereas the number of constants in the reduced form is only $= 2n-1$: hence the coefficients are not determinate; or, what is the same thing, we may have different quadric functions having each of them the same reduced function; these quadric functions, as having the same reduced function, can only differ by multiples of the evanescent expressions

$$\frac{\beta-\gamma}{x-\beta \cdot x-\gamma} + \frac{\gamma-a}{x-\gamma \cdot x-a} + \frac{a-\beta}{x-a \cdot x-\beta}, \quad \&c.$$

In particular, if the number of essential inverts is $= 3$, then the quadric function is of the form

$$\left(a, b, c, f, g, h \left| \frac{1}{x-a}, \frac{1}{x-\beta}, \frac{1}{x-\gamma} \right. \right)^2,$$

which contains one superfluous constant, and equivalent functions differ only by a multiple of

$$\frac{\beta - \gamma}{x - \beta \cdot x - \gamma} + \frac{\gamma - a}{x - \gamma \cdot x - a} + \frac{a - \beta}{x - a \cdot x - \beta}.$$

12. A quadric function such that the degree of the numerator is less by 4 than that of the denominator is said to be “curtate.”

The conditions, in order that the function

$$\left(a, b, c, f, g, h \left| \frac{1}{x-a}, \frac{1}{x-\beta}, \frac{1}{x-\gamma} \right. \right)^2$$

may be curtate, are easily found to be

$$a + b + c + 2f + 2g + 2h = 0,$$

$$a(a + h + g) + \beta(h + b + f) + \gamma(g + f + c) = 0;$$

and by reason of the superfluous constant we are at liberty to assume a third condition: the three conditions may be taken to be $a + h + g$, $h + b + f$, $g + f + c$ each $= 0$; and this being so the values of f, g, h are $= \frac{1}{2}(a - b - c)$, $\frac{1}{2}(-a + b - c)$, $\frac{1}{2}(-a - b + c)$ respectively. Hence the form is

$$\left(a, b, c, \frac{1}{2}(a - b - c), \frac{1}{2}(-a + b - c), \frac{1}{2}(-a - b + c) \left| \frac{1}{x-a}, \frac{1}{x-\beta}, \frac{1}{x-\gamma} \right. \right)^2,$$

which, as already mentioned, we denote by

$$\left(a, b, c \cdot \left| \frac{1}{x-a}, \frac{1}{x-\beta}, \frac{1}{x-\gamma} \right. \right)^2.$$

We have thus the theorem that a curtate function of any number of inverts, but with only the three essential inverts

$$\frac{1}{x-a}, \frac{1}{x-\beta}, \frac{1}{x-\gamma},$$

is always expressible in the foregoing form

$$\left(a, b, c \cdot \left| \frac{1}{x-a}, \frac{1}{x-\beta}, \frac{1}{x-\gamma} \right. \right)^2.$$

13. It may be remarked that the function $(a, b, c \cdot \mid X, Y, Z)^2$ is a function of the differences of the variables X, Y, Z ; and similarly, in the case of four variables, a function $(a, b, c, d, f, g, h, l, m, n \mid X, Y, Z, W)^2$, for which

$$a + h + g + l, \quad h + b + f + m, \quad g + f + c + n, \quad l + m + n + d,$$

are each = 0, is a function of the differences of the variables X, Y, Z, W : and so in general. Any such function is said to be “diaphoric”: and it is easy to see that, taking for the variables any inverts whatever, a diaphoric function is always curtate.

14. The function

$$\left(\cdots a \cdots, \cdots b \cdots, \cdots c \cdots \left| \frac{1}{x-a}, \frac{1}{x-a_1}, \cdots \right. \right)^2,$$

where the coefficients $a, b, c, \dots$ satisfy the relation $a + b + c + \cdots = -2$, is diaphoric, and therefore curtate. In fact, forming the sum, coeff. $\frac{1}{(x-a)^2} + \frac{1}{2}$ coeff. $\frac{1}{x-a \cdot x-\beta} + \cdots$, this is $= a + \frac{1}{2}a^2 - \frac{1}{2}ab - \frac{1}{2}ac - \cdots$, $= -\frac{1}{2}a(2 + a + b + c + \cdots)$, which is = 0; and similarly the other conditions are satisfied.

15. The function

$$\left(\cdots a \cdots, \cdots b \cdots, \cdots c \cdots \left| \frac{1}{x-a}, \frac{1}{x-a_1}, \cdots, \frac{1}{x-\beta}, \cdots \right. \right)^2$$

regarded as a function of the inverts $\frac{1}{x-a}, \frac{1}{x-a_1}, \cdots, \frac{1}{x-\beta}, \cdots$, where

$$a + a_1 + \cdots = b + b_1 + \cdots = c + c_1 + \cdots = h \text{ suppose,}$$

is diaphoric, and therefore curtate. In fact, the condition in regard to $\frac{1}{x-a}$ is

$$a(a + a_1 + \cdots) + \frac{1}{2}(a - b + c)(ac + ac_1 + \cdots) + \frac{1}{2}(-a + b - c)(ab + ab_1 + \cdots) + \frac{1}{2}(-a - b + c)(\cdots) = 0;$$

that is,

$$ah \left\{ a + \frac{1}{2}(-a + b - c) + \frac{1}{2}(-a - b + c) \right\} = 0,$$

which is satisfied. And similarly the other conditions are satisfied.

The functions P, Q, R . Art. Nos. 16 to 20.

16. We consider P, Q, R , rational and integral functions of z , such that $P + Q + R = 0$: hence, using the accent to denote differentiation in regard to z , we have also $P' + Q' + R' = 0$; and therefore $QR' - Q'R = RP' - R'P = PQ' - P'Q = \Theta$ suppose: and we require to find P, Q, R , such that the function Θ contains only the factors of P, Q, R .

17. It is to be observed that, effecting upon a solution P, Q, R any linear substitution $\div(\alpha z + \beta) \div (\gamma z + \delta)$, and omitting the common denominator, we have a solution; but this is regarded as identical with the original solution. The three functions, if

not originally of the same order, can thus be made to be of the same order; or by taking account of the root $z = \infty$, we may in the original case regard them as being of the same order, and it is convenient so to regard them: say they are taken to be of the same order δ . And there is clearly no loss of generality in taking the three functions to be prime to each other; for any common factor of two of them would divide the third, and might therefore be struck out.

18. We may therefore write

$$P = F\Pi(z-l)^p, \quad Q = G\Pi(z-m)^q, \quad R = H\Pi(z-n)^r,$$

where $(z-l)^p$ is taken to denote the distinct simple or multiple factors of P , and the like as regards Q and R ; the factors $z-l$, $z-m$, $z-n$ are thus all of them different. And we have $\delta = \Sigma p, = \Sigma q, = \Sigma r$.

19. It is at once seen that Θ is of the degree $2\delta - 2$, and moreover that it contains the factors $\Pi(z-l)^{p-1}$, $\Pi(z-m)^{q-1}$, $\Pi(z-n)^{r-1}$; hence it contains the factor

$$\Pi(z-l)^{p-1}(z-m)^{q-1}(z-n)^{r-1}.$$

Suppose the number of distinct indices p is $= \sigma_1$, that of distinct indices q is σ_2 , and that of distinct indices r is σ_3 ; then the degree of the factor is $= 3\delta - \sigma_1 - \sigma_2 - \sigma_3$; and if this be $= 2\delta - 2$, then Θ can have no other valuable factor: viz. if the numbers $\sigma_1, \sigma_2, \sigma_3$ of the distinct indices p, q, r respectively are such that $\sigma_1 + \sigma_2 + \sigma_3 = \delta + 2$, a relation which is henceforth taken to be satisfied, then we have

$$\Theta = K\Pi(z-l)^{p-1}(z-m)^{q-1}(z-n)^{r-1}.$$

As already in effect remarked, the conclusion extends to the case where P, Q, R are not of the same degree; the equation $P + Q + R = 0$ here implies that two functions, say P, Q , are of the same degree, and the third function R of an inferior degree; but, this being so, we have only to regard R as containing the factor $\left(1 - \frac{z}{\infty}\right)^t$ of the degree t proper for raising its degree up to that of P or Q .

20. Solutions are given in the following PQR-Table; in which, where required, the proper factor $\left(1 - \frac{z}{\infty}\right)$ has been added; the first column headed Ref. No. (Reference Number) will be explained further on. The Annex to the same Table will also be explained.

THE PQR-TABLE.

Ref. No.	$P =$	$Q =$	$R =$	$\Theta =$
1	z^n	$-1\left(1 - \frac{z}{\infty}\right)^n$	$-z^n + 1$	$nz^{n-1}\left(1 - \frac{z}{\infty}\right)^{n-1}$
2	$4z^n\left(1 - \frac{z}{\infty}\right)^n$	$-(z^n + 1)^2$	$(z^n - 1)^2$	$-4nz^{n-1}(z^n + 2)(z^n - 1)\left(1 - \frac{z}{\infty}\right)^{n-1}$
$3\frac{1}{2}$	$(z^4 + 2\sqrt{-3}z^2 + 1)^3$	$-12\sqrt{-3}z^2(z^4 - 1)^2\left(1 - \frac{z}{\infty}\right)^2$	$-(z^4 - 2\sqrt{-3}z^2 + 1)^3$	$24\sqrt{-3}(z^4 + 2\sqrt{-3}z^2 + 1)^2$ $(z^4 - 2\sqrt{-3}z^2 + 1)^2 z(z^4 - 1)\left(1 - \frac{z}{\infty}\right)$
4	$(z^8 + 14z^4 + 1)^3$	$-(z^{12} - 33z^8 - 33z^4 + 1)^2$	$-108(z^5 - z)^4\left(1 - \frac{z}{\infty}\right)^4$	$216(z^8 + \dots)(z^{12} - \dots)(z^5 - z)^3\left(1 - \frac{z}{\infty}\right)^3$
5	$(z^{20} - 228z^{15} + 494z^{10} + 228z^5 + 1)^3$	$-(z^{30} - 522z^{25} - 10005z^{20} + 0z^{15} - 10005z^{10} + 522z^5 - 1)^2$	$-1728(z^{11} + 11z^6 - z)^5\left(1 - \frac{z}{\infty}\right)^5$	$-8640(z^{20} - \dots)(z^{11} + \dots)^4\left(1 - \frac{z}{\infty}\right)^4$
III, V, VII, VIII	$4z\left(1 - \frac{z}{\infty}\right)$	$-(z + 1)^2$	$(z - 1)^2$	$-4(z + 1)(z - 1)$
IX	$(z - 4)^3$	$-(z - 1)(z + 8)^2$	$27z^2\left(1 - \frac{z}{\infty}\right)$	$27(z - 4)^2(z + 8)z$
X	$z(z + 8)^3$	$-(z^2 - 20z - 8)^2$	$-64(z - 1)^3\left(1 - \frac{z}{\infty}\right)$	$-64(z + 8)^2(z^2 - 20z \dots)(z - 1)$
XI	$4(z^2 - z + 1)^3$	$-(2z^3 - 3z^2 - 3z + 2)^2$	$-27z^2(z - 1)^2\left(1 - \frac{z}{\infty}\right)^2$	$-108(z^2 - \dots)^2(2z^3 - 3z^2 - \dots)$ $\cdot z(z - 1)\left(1 - \frac{z}{\infty}\right)$
XII	$z^3(z + 5)^2(z + 8)$	$-(z^3 + 9z^2 + 12z - 8)^2$	$-64(3z - 1)\left(1 - \frac{z}{\infty}\right)^5$	$-960z^2(z + 5)(z^3 + 9z^2 \dots)\left(1 - \frac{z}{\infty}\right)^4$
XIII	$(z^2 + 14z + 1)^3$	$-(z^3 - 33z + 1)^2$	$-108z(z - 1)^4$	$-108(z^2 + 14z \dots)^2(z^3 - 33z \dots)(z - 1)^3$
XIV	$(64z + 189)(64z^2 + 133z + 49)^3$	$-z(4096z^3 + 18816z^2 + 25725z + \dots)^2$	$-27 \cdot 7^7(z + 1)^3\left(1 - \frac{z}{\infty}\right)^5$	$-135 \cdot 7^7(64z^2 + \dots)^2(4096z^3 + \dots)$ $\cdot (z + 1)\left(1 - \frac{z}{\infty}\right)^4$
XV	$-(5z - 27)(125z^3 - 25z^2 - 265z - 243)^3$	$-(-3125z^5 + 9375z^4 + 18750z^3 + 8750z^2 + 30750z + \dots)^2$	$+1382400000z^3(z + 1)^2\left(1 - \frac{z}{\infty}\right)^5$	$-1382400000(125z^3 - \dots)^2(-3125z^5 + \dots)z^2(z + 1)\left(1 - \frac{z}{\infty}\right)^4$

In the second half of the table the functions P , Q , R , except in the lines XII, XIV and XV, were calculated by Brioschi; those for the lines XII and XIV were calculated by Klein, but as regards line XV there would seem to have been some error at the beginning of the calculation, and the values found by him are erroneous.

ANNEX TO THE PQR-TABLE.

Ref. No.	$(a, b, c)^*$	Calculation of $(ap), (bq), (cr)$.			Polygon
1	$\frac{1}{2}(1-n^{-2}), \frac{1}{2}(1-n^{-2}), 0$	$an = 0$	$bn = 0$	$c1 = 0$	I Polygon
2	$\frac{1}{2}(1-n^{-2}), \frac{3}{8}, \frac{3}{8}$	$an = 0$	$b2 = 0$	$c2 = 0$	I Double Pyramid
3	$\frac{4}{9}, \frac{3}{8}, \frac{4}{9}$	$a3 = 0$	$b2 = 0$	$c3 = 0$	II Tetrahedron
4	$\frac{4}{9}, \frac{3}{8}, \frac{15}{32}$	$a3 = 0$	$b2 = 0$	$c4 = 0$	IV Cube and Octahedron
5	$\frac{4}{9}, \frac{3}{8}, \frac{12}{25}$	$a3 = 0$	$b2 = 0$	$c5 = 0$	VI Dodecahedron and Icosahedron
III	$\frac{4}{9}, \frac{3}{8}, \frac{4}{9}$	$a1 = \frac{4}{9}, a1 = \frac{4}{9}$	$b2 = 0$	$c2 = \frac{5}{18} : \Sigma \left(\frac{1}{z}, \frac{1}{z-\infty}, \frac{1}{z-1} \right)^2$	Tetrahedron
V	$\frac{15}{32}, \frac{3}{8}, \frac{4}{9}$	$a1 = \frac{15}{32}, a1 = \frac{15}{32}$	$b2 = 0$	$c2 = \frac{5}{18} : \&c.$	Cube and Octahedron
VII	$\frac{4}{9}, \frac{3}{8}, \frac{12}{25}$	$a1 = \frac{4}{9}, a1 = \frac{4}{9}$	$b2 = 0$	$c2 = \frac{21}{50}$	Dodecahedron and Icosahedron
VIII	$\frac{12}{25}, \frac{3}{8}, \frac{4}{9}$	$a1 = \frac{12}{25}, a1 = \frac{12}{25}$	$b2 = 0$	$c2 = \frac{5}{18}$	"
IX	$\frac{4}{9}, \frac{3}{8}, \frac{12}{25}$	$a3 = 0$	$b1 = \frac{3}{8}, b2 = 0$	$c1 = \frac{12}{25}, c2 = \frac{21}{50}$	"
X	" " "	$a1 = \frac{4}{9}, a3 = 0$	$b2 = 0$	$c1 = \frac{12}{25}, c3 = \frac{21}{50}$	"
XI	" " "	$a3 = 0$	$b2 = 0$	$c2 = \frac{21}{50}, c2 = \frac{21}{50}$	"
XII	" " "	$a1 = \frac{4}{9}, a2 = \frac{5}{18}, a3 = 0$	$b2 = 0$	$c1 = \frac{12}{25}, c5 = 0$	"
XIII	" " "	$a3 = 0$	$b2 = 0$	$c1 = \frac{12}{25}, c1 = \frac{12}{25}$	"
XIV	" " "	$a1 = \frac{4}{9}, a3 = 0$	$b1 = \frac{3}{8}, b2 = 0$	$c2 = \frac{21}{50}, c5 = 0$	"
XV	" " "	$a1 = \frac{4}{9}, a3 = 0$	$b2 = 0$	$c2 = \frac{21}{50}, c3 = \frac{21}{50}$	"

* Observe, as regards the (a, b, c) column, that the line III agrees with 3; line V with 4 (only there is a transposition of $\frac{4}{9}$ and $\frac{15}{32}$); lines VII and VIII agree each of them with 5 (only as regards VIII there is a transposition of $\frac{4}{9}$ and $\frac{12}{25}$): the remaining lines IX to XV agree each of them with 5. Observe also as regards the Annex generally, the Roman numbering of the lines 2, 3, 4, 5; the lines after the first have thus the Roman numbers I to XV, corresponding to the Roman numbers used by Schwarz.

The Differential Equations $\{x, z\}$ and $\{s, x\}$. Art. Nos. 21 to 45.

21. In reference to what follows, it is convenient to put $P = XP_0$, $P' = X_1P_1 = P''$ multiplied by the product $\Pi(z-l)$ of the several factors taken each with the index unity; and so for Q and R : viz. we write

$$\begin{aligned} P, Q, R &= XP_0, YQ_0, ZR_0, \\ P', Q', R' &= X_1P_0, Y_1Q_0, Z_1R_0, \end{aligned}$$

and the foregoing value of Θ then is

$$\Theta = KP_0Q_0R_0.$$

We come now to the investigation of the leading theorem. Take a, b, c arbitrary, $f, g, h = b-c, c-a, a-b$; P, Q, R functions of z as above; and write

$$f(x-a) : g(x-b) : h(x-c) = P : Q : R,$$

equations, which are consistent with each other and determine x as a rational function of z . Using, as before, the accent to denote differentiation in regard to z , and taking the coefficients (a, b, c) arbitrary, it is required to find the value of

$$\{x, z\} + x'^2 \left(a, b, c \cdot \left| \frac{1}{x-a}, \frac{1}{x-b}, \frac{1}{x-c} \right. \right)^2.$$

22. Calculation of the first term $\{x, z\}$.

We have x a function $(\alpha P + \beta R) \div (\gamma P + \delta R)$, $= \frac{P}{R}$ suppose; and thence $\{x, z\} = \left\{ \frac{P}{R}, z \right\}$; for a moment; then

$$\frac{x'}{x} = \frac{(P/R)'}{P/R} = \frac{PR' - P'R}{PR} = -\frac{P_0Q_0 \cdot KR_0}{Z Z_1 R_0 \cdot P/R} = -\frac{P_1Q_0}{Z_1 R_0}.$$

Substituting the values

$$P_0 = \Pi(z-l)^{p-1}, \quad Q_0 = \Pi(z-m)^{q-1}, \quad R_0 = \Pi(z-n)^{r-1}, \quad Z = \Pi(z-n),$$

we have

$$\frac{x'}{x} = \Sigma \frac{p-1}{z-l} + \Sigma \frac{q-1}{z-m} - \Sigma \frac{r+1}{z-n},$$

and thence

$$\begin{aligned} \{x, z\} &= \Sigma \frac{p-1}{(z-l)^2} + \Sigma \frac{q-1}{(z-m)^2} - \Sigma \frac{r+1}{(z-n)^2} \\ &\quad - \frac{1}{2} \left[\Sigma \frac{p-1}{z-l} + \Sigma \frac{q-1}{z-m} - \Sigma \frac{r+1}{z-n} \right]^2, \end{aligned}$$

or say

$$\begin{aligned} \{x, z\} &= \frac{1}{2} \left(\Sigma \frac{p_1-1}{(z-l)^2} + \Sigma \frac{p_1-1}{(z-l_1)^2} - \Sigma \frac{q-1}{(z-m)^2} - \Sigma \frac{q-1}{(z-m_1)^2} + \Sigma \frac{r+1}{(z-n)^2} + \Sigma \frac{r+1}{(z-n_1)^2} \right. \\ &\quad \left. - \frac{p_1-1}{z-l} \frac{p_1-1}{z-l_1} - \dots - \frac{q_1-1}{z-m} \frac{q_1-1}{z-m_1} - \dots + \frac{r+1}{z-n} \frac{r+1}{z-n_1} + \dots \right). \end{aligned}$$

where it is to be observed that

$$\Sigma(p-1) + \Sigma(q-1) - \Sigma(r+1) = \delta - \sigma_1 + \delta - \sigma_2 - (\delta + \sigma_3) = \delta - \sigma_1 - \sigma_2 - \sigma_3 = -2;$$

consequently the function is diaphoric, and therefore curtate.

It is to be remarked that the function, although presenting itself in a form unsymmetric in regard to the factors of P and Q , and of R , is really symmetric as regards the three sets of factors; this is obvious a priori, and it will be presently verified.

23. For the calculation of the second term

$$x'^2 \left(a, b, c \cdot \left| \frac{1}{x-a}, \frac{1}{x-b}, \frac{1}{x-c} \right. \right)^2$$

we have

$$f(x-a), g(x-b), h(x-c) = \Omega P, \Omega Q, \Omega R,$$

where Ω is a determinate function of z ; hence

$$\frac{x'}{x-a}, \frac{x'}{x-b}, \frac{x'}{x-c} = \frac{P'}{P} + \frac{\Omega'}{\Omega}, \frac{Q'}{Q} + \frac{\Omega'}{\Omega}, \frac{R'}{R} + \frac{\Omega'}{\Omega}.$$

Then substituting these values, by reason that the function is diaphoric, the terms in $\frac{\Omega'}{\Omega}$ disappear, and we have

$$x'^2 \left(a, b, c \cdot \left| \frac{1}{x-a}, \frac{1}{x-b}, \frac{1}{x-c} \right. \right)^2 = \left(a, b, c \cdot \left| \frac{P'}{P}, \frac{Q'}{Q}, \frac{R'}{R} \right. \right)^2,$$

which is

$$= \left(a, b, c \cdot \left| \Sigma \frac{p}{z-l}, \Sigma \frac{q}{z-m}, \Sigma \frac{r}{z-n} \right. \right)^2.$$

We have $\Sigma p = \Sigma q = \Sigma r, = \delta$: and hence by what precedes, this function, considered as a function of the inverts $\frac{1}{z-l}$, &c., is diaphoric, and therefore curtate.

24. We have therefore

$$\begin{aligned} & \{x, z\} + x'^2 \left(a, b, c \cdot \left| \frac{1}{x-a}, \frac{1}{x-b}, \frac{1}{x-c} \right. \right)^2 \\ &= \frac{1}{2} \left[\Sigma \frac{p-1}{(z-l)^2} + \Sigma \frac{q-1}{(z-m)^2} - \Sigma \frac{r+1}{(z-n)^2} \right] \\ & - \frac{1}{2} \left[\Sigma \frac{p-1}{z-l} + \Sigma \frac{q-1}{z-m} - \Sigma \frac{r+1}{z-n} \right]^2 + \left(a, b, c \cdot \left| \Sigma \frac{p}{z-l}, \Sigma \frac{q}{z-m}, \Sigma \frac{r}{z-n} \right. \right)^2, \end{aligned}$$

where the whole function on the right-hand side is curtate.

25. We have to bring the function on the right-hand side into the reduced form

$$\frac{a}{(z-a)^2} + \cdots + \frac{A}{z-a} + \cdots,$$

for the purpose of getting rid of the non-essential inverts (if any).

We write

$$\begin{aligned} \Sigma \frac{p_1 - 1}{(z-l)^2} &= \frac{p-1}{(z-l)^2} + \frac{p_1 - 1}{(z-l_1)^2} + \cdots \\ &= \frac{p-1}{(z-l)^2} + \Sigma_1 \frac{p_1 - 1}{(z-l_1)^2}; \end{aligned}$$

viz. $z-l$ here denotes any particular factor, and $z-l_1$ represents any other factor of the same set; and so in other like cases.

26. The whole coefficient of $\frac{1}{(z-l)^2}$ is

$$= -(p-1) - \frac{1}{2}(p-1)^2 + ap^2, \quad = \frac{1}{2}(1-p^2) + ap^2;$$

an expression which, regarded as a function of a and p , is represented by (ap) : the parentheses are used only to avoid ambiguity, and are omitted when p is a number, thus $a1 = a$, $a2 = -\frac{3}{2} + 4a$, and so in other cases.

27. The whole term in $\frac{1}{z-l}$ comes from

$$\begin{aligned} & -\frac{p-1}{z-l} \left[\Sigma_1 \frac{p_1 - 1}{z-l_1} + \Sigma \frac{q-1}{z-m} - \Sigma \frac{r+1}{z-n} \right] \\ & + \frac{2ap}{z-l} \left[\Sigma_1 \frac{p}{z-l_1} + \frac{1}{2}(-a-b+c) \Sigma \frac{q}{z-m} + \frac{1}{2}(-a+b-c) \Sigma \frac{r}{z-n} \right]. \end{aligned}$$

Viz. each term such as $-\frac{p_1-1}{z-l_1} \cdot \frac{1}{z-l}$ is to be replaced by $\frac{1}{z-l} \cdot \frac{1}{z-l_1} \left(\frac{1}{z-l} - \frac{1}{z-l_1} \right)$, giving rise to the term $-\frac{p_1-1}{l-l_1} \frac{1}{z-l}$, or contributing the term $-\frac{1}{l-l_1}$ to the coefficient of $\frac{1}{z-l}$. The whole coefficient thus is

$$\begin{aligned} & = -(p-1) \left(\Sigma_1 \frac{p_1 - 1}{l-l_1} + \Sigma \frac{q-1}{l-m} - \Sigma \frac{r+1}{l-n} \right) \\ & + 2ap \Sigma_1 \frac{p}{l-l_1} + p(-a-b+c) \Sigma \frac{q}{l-m} + p(-a+b-c) \Sigma \frac{r}{l-n}. \end{aligned}$$

28. Suppose first that $z-l$ is a multiple factor of P , viz. a factor with an index p greater than 1: then, for $z=l$, we have $Q+R=0$, $Q'+R'=0$, and thence $\frac{Q}{R} = \frac{Q'}{R'}$. We have therefore $\Sigma \frac{q}{l-m} = \Sigma \frac{r}{l-n}$; that is,

$$p(-a-b+c) \Sigma \frac{q}{l-m} + p(-a+b-c) \Sigma \frac{r}{l-n}$$

moreover, in the top line, the terms $\Sigma \frac{q}{l-m}$ and $-\Sigma \frac{r}{l-n}$ destroy each other. The whole coefficient of $\frac{1}{z-l}$, when $z-l$ is a multiple factor of P , thus is

$$\begin{aligned} &= -(p-1) \left(\Sigma_1 \frac{p_1-1}{l-l_1} + \Sigma \frac{1}{l-m} - \Sigma \frac{1}{l-n} \right) \\ &\quad + 2ap \Sigma \frac{p_1}{l-l_1} - ap \Sigma \left(\frac{q}{l-m} + \Sigma \frac{r}{l-n} \right), \end{aligned}$$

a form which is now symmetrical in regard to the inverts $\frac{1}{l-m}$ and $\frac{1}{l-n}$.

29. The value just obtained must be equal to

$$(1-p^2+2ap) \left(\Sigma \frac{p_1-1}{l-l_1} + \frac{1}{2} \Sigma \frac{r-1}{l-n} - \Sigma \frac{1}{l-l_1} \right);$$

viz. comparing the two forms and reducing, they will be identical if only

$$(1-p+2ap) \left\{ \Sigma \frac{p+p_1}{l-l_1} - \Sigma \frac{\frac{1}{2}(1+p)q-p}{l-m} - \Sigma \frac{\frac{1}{2}(1+p)r-p}{l-n} \right\} = 0,$$

and it can be shown that the function inside the [] is in fact = 0.

30. We have, as before, $\Sigma \frac{q}{l-m} = \Sigma \frac{r}{l-n}$; or writing each of these quantities = Φ , the equation to be verified is

$$\Sigma \frac{p+p_1}{l-l_1} = (p+1) \Phi - p \Sigma \frac{1}{l-m} - p \Sigma \frac{1}{l-n}.$$

We have

$$\frac{P'}{P} = \frac{P}{z-l} + \Sigma \frac{P_1}{z-l_1} = \frac{X_1}{X},$$

that is,

$$\begin{aligned} \Sigma \frac{P_1}{l-l_1} &= \left[\frac{X_1}{X} - \frac{p}{z-l} \right] \text{ for } z=l, \\ &= \left[\frac{X_1(z-l) - pX}{X(z-l)} \right]. \end{aligned}$$

The first derived function of the numerator is $X'_1(z-l) + X_1 - pX'$, which for $z=l$ is $X_1 - pX'$, which is = 0; and, for the denominator, it is $X'(z-l) + X$, which is also = 0. Passing to the second derived functions, we find

we find in like manner

$$\Sigma \frac{1}{l-l_1} = \frac{1}{2} \frac{X''}{X'},$$

and we thence obtain (z being always $= l$)

$$\Sigma \frac{p+p_1}{z-l_1} = \frac{X'_1}{X'},$$

so that the equation to be verified becomes

$$\frac{X'_1}{X'} = (p+1) \Phi - p \Sigma \frac{1}{l-m} - p \Sigma \frac{1}{l-n}.$$

31. But from the equation $\Theta = PQ' - P'Q, = KP_0Q_0R_0$, we find $XY_1 - X_1Y = KR_0$, and then, differentiating, $XY'_1 + X'Y_1 - X'_1Y - X_1Y' = KR'_0$: writing in these equations $z = l$, they become

$$-X_1Y = KR_0,$$

$$X'Y_1 - X'_1Y = KR'_0,$$

so that, dividing the second by the first,

$$-\frac{X'}{X_1} \frac{Y_1}{Y} + \frac{X'_1}{X_1} = \frac{R'_0}{R_0};$$

or, recollecting that $X_1 = pX'$ and $\frac{Y_1}{Y} = \frac{Q'}{Q}$, we have

$$\frac{X'_1}{X'} = p \left(\frac{R'_0}{R_0} - \frac{Y'}{Y} \right) + \frac{Q'}{Q},$$

that is,

$$\begin{aligned} \frac{X'_1}{X'} &= p \left(\Sigma \frac{r-1}{z-n} - \Sigma \frac{1}{z-m} \right) + \Sigma \frac{q}{l-m}, \\ &= (p+1) \Phi - p \Sigma \frac{1}{l-m} - p \Sigma \frac{1}{l-n}, \end{aligned}$$

the required relation.

32. The result is that, $z-l$ being a multiple factor of P , the coefficient of the term $\frac{1}{z-l}$ is

$$\begin{aligned} &= (1-p^2+2ap) \left\{ \Sigma \frac{p_1-1}{l-l_1} + \frac{1}{2} \Sigma \frac{r-1}{l-n} - \Sigma \frac{1}{l-l_1} \right\} \\ &= 2(ap) \left[\frac{1}{2} \left(\frac{Q'}{Q} + \frac{R'}{R} \right) - \frac{Y'}{Y} - \frac{Z'}{Z} + \frac{X''}{X'} \right]. \end{aligned}$$

33. In the case where $z-l$ is a simple factor of P we have $p=1$, and the coefficient is

$$\begin{aligned} &= \Sigma \Sigma' \frac{p_1}{l-l_1} + (-a-b+c) \Sigma \frac{q}{l-m} + (-a+b-c) \Sigma \frac{r}{l-n} \\ &= a \left(2 \Sigma' \frac{p_1}{l-l_1} - \Sigma \frac{q}{l-m} - \Sigma \frac{r}{l-n} \right) - (b-c) \left(\Sigma \frac{q}{l-m} - \Sigma \frac{r}{l-n} \right). \end{aligned}$$

34. Of course the formulae for the coefficients of $\frac{1}{(z-l)^2}$ and $\frac{1}{z-l}$ give at once, by a mere change of letters, those for the coefficients of $\frac{1}{(z-m)^2}$, $\frac{1}{z-m}$, and $\frac{1}{(z-n)^2}$, $\frac{1}{z-n}$; and the function in question,

$$\{x, z\} + x'^2 \left(a, b, c \cdot \left| \frac{1}{x-a}, \frac{1}{x-b}, \frac{1}{x-c} \right. \right)^2,$$

is now obtained in the required form

$$= \frac{(ap)}{(z-l)^2} + \cdots + \frac{(bq)}{(z-m)^2} + \cdots + \frac{(cr)}{(z-n)^2} + \cdots + \frac{A}{z-l} + \cdots + \frac{B}{z-m} + \cdots + \frac{C}{z-n} + \cdots,$$

where (ap) denotes $\frac{1}{2}(1-p^2) + ap^2$, and the like for (bq) and (cr) ; and where, $z-l$ being a multiple factor of P , the coefficient A contains the factor (ap) , and similarly for B and C .

35. Suppose that the coefficients a, b, c are no one of them $= 0$; we have $a1, = a$, which does not vanish; that is, $z-l$ being a simple factor of P , the expression contains $\frac{1}{(z-l)^2}$, or the invert $\frac{1}{z-l}$ is essential: and similarly, $z-m$ being a simple factor of Q , or $z-n$ a simple factor of R , the inverts $\frac{1}{z-m}$ and $\frac{1}{z-n}$ are essential. But for $z-l$ a multiple factor of P , the coefficient (ap) of the term $\frac{1}{(z-l)^2}$ may vanish, viz. this will be the case if $a = \frac{1}{2} \left(1 - \frac{1}{p^2} \right)$; and, when this is so, the coefficient A of the corresponding term $\frac{1}{z-l}$ also vanishes; that is, $\frac{1}{z-l}$ is a non-essential invert. And similarly for any multiple factor $z-m$ of Q or $z-n$ of R , the invert $\frac{1}{z-m}$ or $\frac{1}{z-n}$ may be non-essential.

36. If P, Q, R contain each of them only multiple factors of the same index, say of the indices p, q, r for the three functions respectively, viz. if the functions are $P(\Pi(z-l))^p, G(\Pi(z-m))^q, H(\Pi(z-n))^r$, the result contains only the six terms written for a, b, c any given values of them $= \frac{1}{2} \left(1 - \frac{1}{p^2} \right), \frac{1}{2} \left(1 - \frac{1}{q^2} \right), \frac{1}{2} \left(1 - \frac{1}{r^2} \right)$ respectively the result is $= 0$: viz. we then have

$$\{x, z\} + x'^2 \left(a, b, c \cdot \left| \frac{1}{x-a}, \frac{1}{x-b}, \frac{1}{x-c} \right. \right)^2 = 0,$$

or we in fact have, for the values of a, b, c in question,

$$f(x-a) : g(x-b) : h(x-c) = P : Q : R$$

of this differential equation of the third order.

37. The reasoning applies directly to lines 2, 3, 4, 5 of the PQR-Table; and with a slight variation to line 1; viz. here the factors of R ($= -z^n + 1$) are all simple factors, but in virtue of $c = 0$ and $n = b$, the corresponding inverts disappear, and, the other inverts also disappearing, the value of the function in $\{x, z\}$ becomes $\equiv 0$.

Thus line 3 shows that the function x , determined by

$$f(x - a) : g(x - b) : h(x - c) = (z^4 + 2\sqrt{-3}z^2 + 1)^3 : -12\sqrt{-3}(z^4 - 1)^2 : -(z^4 - 2\sqrt{-3}z^2 + 1)^3,$$

satisfies

$$\{x, z\} + x'^2 \left(\frac{4}{9}, \frac{3}{8}, \frac{4}{9} \cdot \left\| \frac{1}{x-a}, \frac{1}{x-b}, \frac{1}{x-c} \right\| \right)^2 = 0,$$

and so for any other of the five lines.

38. The indices of the factors of P, Q, R may be such that, for proper values of the coefficients a, b, c , there are in all only three essential inverts, say $\frac{1}{x-a_1}, \frac{1}{x-b_1}, \frac{1}{x-c_1}$, belonging to the three functions P, Q, R respectively, or it may be two, or three, of them to the same function. When this is so, the function of those inverts is, by what precedes, a curtate function, and it is consequently a function

$$\left(a_1, b_1, c_1 \cdot \left\| \frac{1}{x-a_1}, \frac{1}{x-b_1}, \frac{1}{x-c_1} \right\| \right)^2,$$

where a_1, b_1, c_1 are the values of the three which do not vanish in the series of expressions $(ap), (bq), (cr)$.

The remaining lines (III, V, VII, VIII) and IX to XV of the PQR-Table give such values of P, Q, R , the values of (a, b, c) and the calculation of the values of $(ap), (bq), (cr)$ is shown by the corresponding lines of the Annex. And we have thus values of x determined by the equations

$$f(x - a) : g(x - b) : h(x - c) = P : Q : R,$$

and giving

$$\{x, z\} + x'^2 \left(a, b, c \cdot \left\| \frac{1}{x-a}, \frac{1}{x-b}, \frac{1}{x-c} \right\| \right)^2 = \left(a_1, b_1, c_1 \cdot \left\| \frac{1}{x-a_1}, \frac{1}{x-b_1}, \frac{1}{x-c_1} \right\| \right)^2.$$

39. For instance, from line IX we have

$$f(x - a) : g(x - b) : h(x - c) = (z - 4)^3 : -(z - 1)(z + 8)^2 : 27z^2 \left(1 - \frac{z}{\infty} \right),$$

the values of (a, b, c) are $\frac{4}{9}, \frac{3}{8}, \frac{12}{25}$; and since P, Q, R contain factors with the exponents 3; 1, 2; and 1, 2 respectively, the coefficients which present themselves on the right-hand side are

$$a3; \quad b1, \quad b2; \quad c1, \quad c2,$$

which are

$$= 0; \frac{3}{8}, 0; \frac{12}{25}, \frac{21}{50}, \text{ respectively.}$$

Hence writing $a_1, b_1, c_1 = \frac{3}{8}, 0; \frac{12}{25}, \frac{21}{50}$, the corresponding inverts are $\frac{1}{x-1}, \frac{1}{x-\infty}, \frac{1}{x}$; and the result is

$$\{x, z\} + x'^2 \left(\frac{4}{9}, \frac{3}{8}, \frac{12}{25} \cdot \left\| \frac{1}{x-a}, \frac{1}{x-b}, \frac{1}{x-c} \right\| \right)^2 = \left(\frac{3}{8}, \frac{12}{25}, \frac{21}{50} \cdot \left\| \frac{1}{x-1}, \frac{1}{x-\infty}, \frac{1}{x} \right\| \right)^2.$$

40. It is hardly necessary to remark that an expression

$$\left(a_1, b_1, c_1 \cdot \left\| \frac{1}{x-a_1}, \frac{1}{x-b_1}, \frac{1}{x-c_1} \right\| \right)^2$$

in fact denotes

$$\frac{a_1}{(x-a_1)^2} + \frac{b_1}{(x-b_1)^2} + \frac{-a_1-b_1+c_1}{(x-a_1)(x-b_1)} + \dots$$

The particular form of the x inverts is immaterial; we could by a general linear transformation upon the x make them to be $\frac{1}{x-a_1}, \frac{1}{x-b_1}, \frac{1}{x-c_1}$ with the (a_1, b_1, c_1) arbitrary; or we can give to the a_1, b_1, c_1 any particular values we please: there would be a propriety in making the inverts to be in every case (as in the foregoing example) $\frac{1}{x}, \frac{1}{x-\infty}, \frac{1}{x-1}$; but the numerical work would be troublesome, and it is not worth while to effect it.

41. The conclusion is that lines (III, V, VII, VIII) and IX to XV of the PQR-Table, give, for determinate values of (a, b, c) and (a_1, b_1, c_1) , solutions

$$f(x-a) : g(x-b) : h(x-c) = P : Q : R$$

of the equation

$$\{x, z\} + x'^2 \left(a, b, c \cdot \left\| \frac{1}{x-a}, \frac{1}{x-b}, \frac{1}{x-c} \right\| \right)^2 = \left(a_1, b_1, c_1 \cdot \left\| \frac{1}{x-a_1}, \frac{1}{x-b_1}, \frac{1}{x-c_1} \right\| \right)^2,$$

where a, b, c, a_1, b_1, c_1 are or can be made arbitrary, but without any real gain of generality herein. This is the Differential Equation $\{x, z\}$.

42. Recurring to the results from the Arabic lines of the PQR-Table, but for convenience writing s instead of x , we have

$$f(s-a) : g(s-b) : h(s-c) = P : Q : R,$$

where P, Q, R are now functions of z , a solution of

$$[s, z] + \left(\frac{ds}{dz} \right)^2 \left(a, b, c \cdot \left\| \frac{1}{s-a}, \frac{1}{s-b}, \frac{1}{s-c} \right\| \right)^2 = 0.$$

But we have

$$\{s, x\} = - \left(\frac{dx}{ds} \right)^2 [s, z],$$

and the foregoing is therefore a solution of

$$[s, x] = \left(a, b, c \cdot \left| \frac{1}{s-a}, \frac{1}{s-b}, \frac{1}{s-c} \right. \right)^2,$$

a differential equation of the third order. This is the Differential Equation $\{s, x\}$.

43. From the Roman lines, if we assume

$$f(s-a) : g(s-b) : h(s-c) = P : Q : R,$$

where $\mathfrak{P}, \mathfrak{Q}, \mathfrak{R}$ are functions of z , not the same functions than P, Q, R are of s , since they belong to a different line of the Table: we have, as before,

$$[x, z] + \left(\frac{dx}{dz} \right)^2 \left(a, b, c \cdot \left| \frac{1}{x-a}, \frac{1}{x-b}, \frac{1}{x-c} \right. \right)^2 = \left(a_1, b_1, c_1 \cdot \left| \frac{1}{x-a_1}, \frac{1}{x-b_1}, \frac{1}{x-c_1} \right. \right)^2.$$

44. We may combine any such result with a properly selected result of the preceding system, the two results being such that (a, b, c) have the same values in each of them. (See as to this the foot-note referring to the Annex to the PQR-Table.) The last equation then becomes

$$[x, z] + \left(\frac{dx}{dz} \right)^2 [s, x] = \left(a_1, b_1, c_1 \cdot \left| \frac{1}{x-a_1}, \frac{1}{x-b_1}, \frac{1}{x-c_1} \right. \right)^2,$$

or since

$$[x, z] + \left(\frac{dx}{dz} \right)^2 [s, x] = [s, z],$$

this is

$$[s, z] = \left(a_1, b_1, c_1 \cdot \left| \frac{1}{x-a_1}, \frac{1}{x-b_1}, \frac{1}{x-c_1} \right. \right)^2,$$

the corresponding relation between x, z being of course obtained by the elimination of s from the two sets of equations

$$f(s-a) : g(s-b) : h(s-c) = P : Q : R \quad \text{and} \quad f(x-a) : g(x-b) : h(x-c) = \mathfrak{P} : \mathfrak{Q} : \mathfrak{R},$$

viz. the required relation is

$$P : Q : R = \mathfrak{P} : \mathfrak{Q} : \mathfrak{R},$$

where P, Q, R are functions of s ; $\mathfrak{P}, \mathfrak{Q}, \mathfrak{R}$ functions of x ; and, in virtue of

$$P + Q + R = 0, \quad \mathfrak{P} + \mathfrak{Q} + \mathfrak{R} = 0,$$

the relations between equivalents to a single equation between x and s . And writing finally x in place of z , that is, now considering $\mathfrak{P}, \mathfrak{Q}, \mathfrak{R}$ as functions of x , we have

$$\mathfrak{P} : \mathfrak{Q} : \mathfrak{R} = P : Q : R$$

as a solution of

$$[s, x] = \left(a_1, b_1, c_1 \cdot \left| \frac{1}{x-a_1}, \frac{1}{x-b_1}, \frac{1}{x-c_1} \right. \right)^2,$$

a differential equation of the third order of the foregoing form, $[s, x] =$ given function of x , but with different values of the coefficients (a_1, b_1, c_1) instead of (a, b, c) .

45. It thus appears that there are in all 16 sets of values of (a, b, c) for which the equation is solved, viz. the 16 sets of values are shown in the right-hand column of the Annex. For greater clearness I exhibit the integral equations as follows:

	Functions of z .		Functions of x .
1	$f(x - a) : g(x - b) : h(x - c) = P : Q : R$ (1)	=	Polygon
I	”	=	Double Pyramid
II	$4z : -(z + 1)^2 : (z - 1)^2$ (2)	=	Tetrahedron
III	”	=	”
IV	$f(x - a) : g(x - b) : h(x - c) = (z - 1)^2 : (z + 1)^2$ (4)	=	Cube and Octahedron
V	”	=	”
VI	$f(x - a) : g(x - b) : h(x - c) = (4z + 1)^2 : (z - 1)^2$ (4)	=	Dodecahedron and Icosahedron
VII	”	=	”
VIII	$(z - 1)^2 : -(z + 1)^2 : 4z$ (2)	=	”
IX	$P : Q : R$ (IX)	=	”
X	”	=	”
XI	”	=	”
XII	”	=	”
XIII	”	=	”
XIV	”	=	”
XV	”	=	”

The values of the P, Q, R as functions of z , are taken out of the PQR -Table: only in the lines III, V, VII, VIII, where P, Q, R are given in the right-hand column of the Annex. For greater clearness I exhibit the integral equations as follows:

$$= 4x, \quad -(x + 1)^2, \quad (x - 1)^2,$$

and where, as regards V and VIII, line is a transposition of P and R ; 1 have inverted the actual values of the $\mathfrak{P}$ -functions. (See as to this the foot-note referring to the Annex.)

The Schwarzian Theory. Art. Nos. 46 to 62.

46. Considering the foregoing equation

$$[s, x] = \left(a_1, b_1, c_1 \cdot \left\| \frac{1}{x - a_1}, \frac{1}{x - b_1}, \frac{1}{x - c_1} \right\| \right)^2$$

as a particular case of the equation $[s, x] = \text{Rational function of } x, = R(x)$ suppose, then we have in 1, I, II, IV, VI solutions of the form $x = \text{Rational function of } s$.

Consider, in general, a solution of this form, $x = F(s)$ a rational function of s : then s is an irrational function of x , and if a, b are any values of s , then $[a_1, b_1] = R(x)$, $[a_2, b_2] = R(x)$; that is, $[a_2, b_2] = [a_1, b_1]$, and therefore (ante, No. 7) $a_2 = \frac{aa_1 + b}{ca_1 + d}$. And then $x = F(a_2) = F\left(\frac{aa_1 + b}{ca_1 + d}\right)$, $= F(a_1)$: viz. $F(s)$ is a rational function of s , transformable into itself by the several homographic transformations of a group of such transformations: viz. taking s a group; and conversely to each homographic transformation of s we have a rational function of s , transformable into itself by the several homographic transformations of a group of such transformations; viz. taking s as the parameter of a point, the points which are between any two roots whatever of the equation $x = F(s)$ there exists a homographic relation of the form in question. Further, if s_1 is a root, the others are $\frac{as_1 + b}{cs_1 + d}$ for the several homographic transformations $\frac{as + b}{cs + d}$ of the group; and transformable into itself by the several homographic transformations of a group of such transformations occur in the form $x = F(s)$; conversely to each homographic transformation $\frac{as + b}{cs + d}$ of the group correspond to the six elements of the rational function $x = F(s)$ a Group; or, what is the same thing, to each homographic transformation $\frac{as + b}{cs + d}$ of the group correspond to a Group; or, what is the same thing, to each homographic transformation $\frac{as + b}{cs + d}$ of the group correspond to a Group; the rational function $x = F(s)$ is the function corresponding to the Group.

47. We may, in any equation between x and s , consider s as imaginary variables $p + qi$ and $u + vi$ respectively; considering then (p, q) and (u, v) as rectangular coordinates of points in different planes, we have a first plane the locus of the points s , and a second plane the locus of the points x : there is between the two planes a correspondence which is in fact the correspondence of conformable figures; to the infinitesimal element ds drawn from a point s of the first figure corresponds an infinitesimal element dx drawn from the corresponding point x of the second figure, these elements being in general connected by an equation of the form $dx = \alpha ds$ involving s but not ds ; or, what is the same thing, the lengths of the ratio of the lengths of the elements being the moduli of α ($a + bi$), and the inclinations of the directions being the angle of α ($a + bi$). Passing to the second derived element d_2s of the same thing along ds , d_2x the corresponding consecutive elements of the path of the point s , then the ratio of the lengths of the elements d_2x, d_2s are the corresponding consecutive elements of the path of the point x , the ratio $d_2x : d_2s$ the corresponding consecutive elements of the path of the path of the point s , then the ratio of the lengths of the elements d_2s and d_2x are the corresponding consecutive elements of the path of the point x . In particular, the consecutive elements ds having all the same direction (that is, the same module λ), as λ is a real positive, has $s = a + bi$, b, c are critical points of the kind in question. In fact, writing $x - a = h$, the equation is of the form

$$x - a = iF' \exp i(\theta + \theta_1).$$

48. I consider the foregoing equation $[s, x] = R(x)$, where $R(x)$ is a rational function, and is now taken to be a real function of x : we may assume $s' = i\rho' e^{i\theta}$, where the accents denote differentiation in regard to x , and where ρ' , θ' , and therefore also ρ , are real functions of x . We have

$$\frac{s''}{s'} = \frac{\rho''}{\rho'} + i\theta'',$$

and thence

$$\begin{aligned} \frac{s'''}{s'} - \left(\frac{s''}{s'}\right)^2 &= \frac{\rho'''}{\rho'} - \left(\frac{\rho''}{\rho'}\right)^2 + \frac{\theta'''}{\theta'} - \left(\frac{\theta''}{\theta'}\right)^2 + i\theta'' \\ -\frac{1}{2}\left(\frac{s''}{s'}\right)^2 &= -\frac{1}{2}\left(\frac{\rho''}{\rho'}\right)^2 - \frac{1}{2}\left(\frac{\theta''}{\theta'}\right)^2 + \frac{1}{2}\theta'^2 - i\frac{\rho''}{\rho'}i\theta'' - i\frac{\rho''\theta''}{\rho'}, \end{aligned}$$

and thence

$$[s, x] = [\rho, x] + [\theta, x] + \frac{1}{2}\theta'^2 - i\frac{\rho''\theta''}{\rho'\theta'} - i\frac{\rho''\theta''}{\rho'}.$$

Putting this = $R(x)$, and assuming that x is real, we have

$$[\rho, x] + [\theta, x] + \frac{1}{2}\theta'^2 = R(x), \quad 0 = i\frac{\rho''\theta''}{\rho'\theta'}.$$

The last equation gives $\rho''\theta'' = 0$, that is, $\theta'' = 0$, which gives $s' = 0$, and may be disregarded; or else $\rho'' = 0$, therefore ρ' , a real constant, = γ suppose, and $[\rho, x] = 0$; hence for the solution of the equation $[s, x] = R(x)$, we have $s' = i\gamma' e^{i\theta}$, θ a real quantity determined by $[\theta, x] + \frac{1}{2}\theta'^2 = R(x)$; and then, integrating the equation for s' , we have $s = a + \beta s + \gamma e^{i\theta}$, a, β, γ real constants.

49. The conclusion is that, if $[s, x] = R(x)$, a real function of x , and if x be real, that is, if the point x moves along a right line (say the x -line), then $s = a + \beta s + \gamma e^{i\theta}$ (θ , and the constants a, β, γ being real), that is, the point s moves in a circle, coordinates of the center: a, β , and radius = γ .

[Figure: Three circles labelled A, B, C along the x -axis $\rho-\phi-q$, illustrating the conformal mapping of the real x -line to a s -plane consisting of arcs forming a triangular region.]

50. Suppose a, b, c are any real values of x representing points a, b, c on the x -line; and A, B, C any given imaginary values of s representing points A, B, C

in the s -plane: since $[s, x] = R(x)$ is a differential equation of the third order, the integral contains three arbitrary constants, and we may imagine these so determined that to the values $s = a, b, c$ shall correspond the values $s = A, B, C$ respectively.

If there is not on the x -line any critical point, as the point x moves continuously along this line from the point a towards the point b , the corresponding circle (in-asmuch as a, b, c and A, B, C are corresponding points) must be the circle through the three points A, B, C .

51. If however the points a, b, c are critical points, such that the element ds at the corresponding points A, B, C are equal to multiples of $(dx)^\lambda, (dx)^\mu, (dx)^\nu$ respectively, then on the x -line the point a in the direction ax to a passes continuously along the line: the point s at the point A moves continuously along a circle, (in-asmuch as a, b, c and A, B, C are corresponding points) must be the circle through the three points A, B, C .

51. If however the points a, b, c are critical points, such that the elements ds only along the s -line: the point a, b, c are critical points, such that the elements ds along the a -line, $\mu\pi, \nu\pi$, of $(dx)^\lambda, (dx)^\mu, (dx)^\nu$ respectively. The arc ABC becomes continuous at a, b, c but the point s at the points A, B, C must change abruptly through the angles $\lambda\pi, \mu\pi, \nu\pi$, (being real), that is, the point s moves in a circle.

51. If however there is on the x -line a critical point, such that the element ds only along the x -line. Pasing the values of λ, μ, ν , being real, and equal to the curvilinear triangle that to the values $x = a, b, c$ shall correspond the values $s = A, B, C$. (In-asmuch as a, b, c and A, B, C are corresponding points) must be the circle through the three points A, B, C .

52. The foregoing equation

$$[s, x] = \left(a, b, c \cdot \left| \frac{1}{x-a}, \frac{1}{x-b}, \frac{1}{x-c} \right| \right)^2$$

where a, b are real values of x representing points a, b, c on the

$$\frac{d}{dx} \log \frac{ds}{ds} = \frac{1-\lambda}{h} + a_1 + a_2 h + a_3 h^2 + \dots,$$

which is satisfied. And similarly the other conditions are satisfied.

$$s = h k^{\lambda-1} (1 + b_1 h + b_2 h^2 + \dots), = k\rho \text{ for shortness.}$$

This is a particular integral, but we have from it the general integral

$$s = \frac{\alpha + \beta k\rho}{\gamma + \delta k\rho}.$$

* Since there is no critical point on the x -line there can be no abrupt change of direction in the path of s , that is, the path of s cannot consist of circular arcs meeting at an angle: but it is in the next further assumed that the path of s cannot consist of different arcs of circles, the one continuing the other without any abrupt change of direction.

If A be the value of s corresponding to a vanishing value of h , then $B = aA$, and we find

$$s = \frac{a + A\delta h\rho}{\gamma + \delta h\rho}, \quad = \left(A + \frac{a}{\delta h\rho} \right) \left(1 + \frac{\gamma}{\delta h\rho} \right)^{-1}, \quad = A + \frac{a - \gamma A}{\delta} \frac{1}{h\rho} + \dots;$$

viz. reducing $\frac{1}{\rho}$ to its principal term h^λ , and then writing ds , ds for $s - A$, and h ($= x - a$) respectively, we have $ds = K(dx)^\lambda$, or $s - a = \alpha$ is a critical point with the exponent λ ; and similarly $x = b$ and $x = c$ are critical points with the exponents μ and ν respectively.

53. Hence in the equation

$$[s, x] = \left(a, b, c \cdot \left| \frac{1}{x-a}, \frac{1}{x-b}, \frac{1}{x-c} \right. \right)^2,$$

as the point x , passing successively through a, b, c , describes the x -line, the point s , passing successively through A, B, C , describes the circular arcs meeting at the proper triangle ABC . To points s indefinitely near the x -line correspond points x indefinitely near to the boundary ABC of the triangle ABC on the s -plane. Similar reasoning shows that the s -plane, supposing the boundary just spoken of, agrees fully with the s -plane in question, viz. the points A, B, C are critical points of x , with the exponents λ, μ, ν , where $a, b, c, \lambda, \mu, \nu$ are connected by the relation

$$f(x-a) : g(x-b) : h(x-c) = P : Q : R$$

of this differential equation of the third order.

54. It is to be remarked, in regard to the values of the values of the values of the values; and conversely, there are two values of x , viz. one in the upper half and the other in the lower half of the x -line, which correspond to the same each of the points s within the curvilinear triangle ABC . The two half-planes (above and below) form, between them, two sets of triangles ABC , $\&c.$, and $\&c.$, respectively, each of them which are equivalent functions of x . And we have thus the values x , the points A, B, C as the values, on opposite sides of the s -line, of the curvilinear triangles in the upper half-plane corresponds on opposite sides of the x -line, correspond to two sets of triangles ABC , $\&c.$, and $\&c.$, respectively, certain determinate values; and, in fact, dividing the surface of a sphere into triangles ABC , ABC , $\&c.$, alternately of the two kinds: corresponding to all these triangles in the s -plane and the corresponding triangles in the s -plane are these triangles; and hence the shaded half-plane corresponds to one set of them and the unshaded triangles correspond to the other. To the values x , in the upper half of the x -line we have, A, B, C corresponding to the foot-note for the values of $\frac{4}{5}, \frac{15}{50}$, etc., and there will be six considerations as these last followers, in the Annex of the PQR-Table; and then showed a *a priori* that the equation

$$[s, x] = \left(a, b, c \cdot \left| \frac{1}{x-a}, \frac{1}{x-b}, \frac{1}{x-c} \right. \right)^2$$

is algebraically integrable for these values of a, b, c ; and only for these values, or for values reducible to them.

55. As an instance, take the double pyramid form: the integral equation is

$$f(x-a) : g(x-b) : h(x-c) = 4x^n : -(x^n-1)^2 : (x^n+1)^2,$$

or say

$$\frac{(x-a)(x-b)}{(x-b)(x-c)} = -\frac{(x^n-1)^2}{(x^n+1)^2};$$

or if, for greater simplicity, we assume $a, b, c, = 1, 0, \infty$, this is $x = \frac{(x^n-1)^2}{(x^n+1)^2}$, or say $-(x^n-1) = \sqrt{x}(x^n+1)$,

that is, $x^n = \frac{1 \mp \sqrt{x}}{1 \pm \sqrt{x}}$, a solution of the differential equation

$$[s, x] = \left(\frac{3}{8}, \frac{1}{2}(1-n^{-2}), \frac{3}{8} \cdot \left| \frac{1}{x-1}, \frac{1}{x-\infty}, \frac{1}{x} \right| \right)^2.$$

In particular, if $n = 3$, we have $x = \frac{(s^3-1)^2}{(s^3+1)^2}$ or $s^3 = \frac{1 \mp \sqrt{x}}{1 \pm \sqrt{x}}$, a solution of

$$[s, x] = \left(\frac{3}{8}, \frac{4}{9}, \frac{3}{8} \cdot \left| \frac{1}{x-1}, \frac{1}{x-\infty}, \frac{1}{x} \right| \right)^2.$$

57. We have here the spherical surface divided by the equator and three meridians into twelve triangles, each with the angles $\frac{1}{2}\pi, \frac{1}{3}\pi, \frac{1}{2}\pi$; and then, projecting from the South pole on the plane of the equator, we have the annexed figure of the s -plane,

[Figure: A central circle on the s -plane is divided by three radial lines meeting at the centre into six sectors; the perimeter is split by these and three meridional arcs into twelve curvilinear triangles, alternately shaded and unshaded. Labels A, B, C, D mark four boundary points; another label sits at the centre and at ∞ outside the disk.]

divided into 12 curvilinear triangles, each with these same angles $30^\circ, 90^\circ, 60^\circ$; the plane is divided by the shading into two systems, each of 6 triangles. The figure of the s -plane is by the s -line divided into two half-planes, one shaded, the other unshaded; and we have on the line the point c at ∞ , a at the origin, and b at the distance unity.

58. Take x real; then, if x is positive and less than 1, s^3 is real and positive, and we have for s the infinitum half-lines at the inclinations 0° , 180° , 300° . If x is real and negative, or in particular greater than 1, viz. here the factors of R ($= -x^n + 1$) are all simple factors, but in virtue of $c = 0$ and $n = b$, the corresponding inverses disappear, and, the the form $\frac{1 - hi}{1 + hi}$, $= \cos \theta + i \sin \theta$; whence as x is of the same form, or the locus of the point s is a circle radius unity. Writing $s^3 = \frac{1 - \sqrt{x}}{1 + \sqrt{x}}$, and supposing that the point x moves along the x -line through 1 through ∞ to ∞ , and thence from ∞ through 0 to b , the point s describes the sides BA , AC , CB of the shaded triangle marked K .

59. Suppose that the point x is at k , in the shaded half-plane and at indefinitely small distance ϵ from the x -line; in the shaded line in question, in the shaded triangle $s(-i)$, we have $s^3 = \frac{1 - \sqrt{x}}{1 + \sqrt{x}}$, and then $x = -\frac{1}{4}\epsilon(1 - i)$ nearly, and hence a value of s is $s = \frac{1 - ki}{1 + ki}$, $= \cos \theta + i \sin \theta$; whence s^3 is of the same form, or the locus of the point s is a circle radius unity. Writing $s^3 = \frac{1 - \sqrt{x}}{1 + \sqrt{x}}$, and supposing that the point x moves along the x -line a through b through a to ∞ , and to ∞ from ∞ to b , the shaded value $\sqrt{x}$ we have $s^3 = \frac{1 - \epsilon(1 - i)}{1 + \epsilon(1 - i)}$, $= 1 - 2\epsilon(1 - i)$ nearly, and hence a value of s is $\epsilon = 1 - \frac{2}{3}\epsilon + \frac{2}{3}\epsilon i$ nearly, and we have to find from a at c in K , and b at the situate on the circle in question.

59. Suppose the point x is at k , in the shaded half-plane, in the shaded triangle small distance ϵ from the x -line, in the shaded line in question, then for the value $x(-a) = h$, the shaded line BA , AC , CB of the equation, which $\frac{1}{x - a}$ value $\epsilon(1 - i)$, we have $s^n = \frac{1 - \epsilon(1 - i)}{1 + \epsilon(1 - i)}$, $= 1 - 2\epsilon(1 - i)$ nearly, and hence a value of s is $s = 1 - \frac{2}{3}\epsilon + \frac{2}{3}\epsilon i$, which belongs to the shaded triangle K ; and similarly the other triangles; and hence the shaded half-plane corresponds to the shaded triangles, and the unshaded half-plane corresponds to the other.

60. Suppose the point x , instead of being in question of a , b , c , is in the circle in question.

We have here $x - a = a(e^{i\theta_1} - e^{-i\theta_1}) = -2ia e^{i\frac{1}{2}\pi} \sin \frac{1}{2}\theta_1$; $e^{i\theta_1} \cdot e^{i\theta_2}$, say $x - a = a e^{i\theta_1} \cdot e^{i\theta_2} e^{i\frac{1}{2}\pi - \frac{1}{2}\theta_2}$ similarly

$$x - a = iP_1 \exp \frac{1}{2}(\theta + \theta_1),$$

Similarly $x - b = iQ_1 \exp \frac{1}{2}(\theta + \theta_2)$, and $x - c = iR_1 \exp \frac{1}{2}(\theta + \theta_3)$, where P , Q , R denote $\sin \frac{1}{2}(\theta - \theta_1)$, $\sin \frac{1}{2}(\theta - \theta_2)$, $\sin \frac{1}{2}(\theta - \theta_3)$ respectively. In like manner, $a - b$, $b - c$, $c - a$, $iH \exp \frac{1}{2}(\theta_1 + \theta_2)$, $iH \exp \frac{1}{2}(\theta_2 + \theta_3)$, $iH \exp \frac{1}{2}(\theta_3 + \theta_1)$, where F , G , H denote $\sin \frac{1}{2}(\theta_1 - \theta_2)$, $\sin \frac{1}{2}(\theta_2 - \theta_3)$, $\sin \frac{1}{2}(\theta_3 - \theta_1)$ respectively.

We have

$$-\frac{b-x \cdot x-a \cdot a-b}{x-a \cdot x-b} = \frac{-FGH}{PQR} \exp i \frac{1}{2} (\theta_1 + \theta_2 + \theta_3 - 3\theta),$$

$$\frac{1}{b-c \cdot x-a} = \frac{-1}{c^2 IP} \exp i - \frac{1}{2} (\theta_2 + \theta_3 + \theta_1 + \theta),$$

with the like values for $\frac{1}{c-a \cdot x-b}$ and $\frac{1}{a-b \cdot x-c}$. Hence the right-hand side of the equation is

$$= \frac{FGH}{PQR} \left(\frac{a}{PP} + \frac{b}{QQ} + \frac{c}{RR} \right) \exp i (-2\theta).$$

62. Considering now the left-hand side of the equation, we have

$$[s, x] = \frac{1}{\left(\frac{dx}{d\theta}\right)^2} (\{s, \theta\} - \{x, \theta\});$$

substituting for x its value $= a + \beta i e^{i\theta} + \gamma e^{i\theta}$, this becomes

$$[s, x] = -\frac{1}{\gamma^2} e^{-i\theta} (\{s, \theta\} - \frac{1}{2}),$$

that is,

$$= -\frac{1}{\gamma^2} (\{s, \theta\} - \frac{1}{2}) \exp i (-2\theta).$$

Assume $s = L + Mi + Ne^{i\theta}$, L , M and N constants; then using the accent to denote differentiation in regard to θ , we find without difficulty $[s, \theta] = (\Theta_1 + \Theta_2 + \frac{1}{2}\Theta^2)$, and the value of $[s, x]$ becomes

$$= -\frac{1}{\gamma^2} ((\Theta_1 + \Theta_2 + \frac{1}{2}\Theta^2) - \frac{1}{2}) \exp i (-2\theta).$$

Hence, substituting the values of the two sides of the equation, the imaginary factor $\exp i (-2\theta)$ divides out, and the equation becomes

$$(\Theta_1 + \Theta_2 + \frac{1}{2}\Theta^2) = \frac{FGH}{PQR} \left(\frac{a}{PP} + \frac{b}{QQ} + \frac{c}{RR} \right),$$

an equation, in which everything is real and which thus determines Θ as a real function of θ ; and we have therefore the theorem in question.

Connexion with the differential equation for the hypergeometric series. Art. Nos. 63 to 68.

63. Take p , q given functions of x , and y a function of x determined by the equation

$$\frac{d^2 y}{dx^2} + p \frac{dy}{dx} + qy = 0;$$

[745] (CONTINUED)

ON THE SCHWARZIAN DERIVATIVE AND THE POLYHEDRAL FUNCTIONS.

63. again P, Q given functions of z , and v a function of z determined by the equation

$$\frac{d^2v}{dz^2} + P \frac{dv}{dz} + Qv = 0;$$

and assume

$$y = wv.$$

Substituting this value of y in the first equation, we obtain for v an equation of the second order (the coefficients of which contain w), and we may make this identical with the second equation; viz. comparing the coefficients of the two equations, we thus have two equations each containing w ; and by eliminating w we obtain a differential equation of the third order between z and x . This is, in fact, the basis of Kummer's theory for the transformation of a hypergeometric series: the equation between z, x will be found presently in a different manner.

64. But if with Schwarz, instead of making the equation obtained for v as above identical with the given equation for v , we merely assume that the two equations are consistent, then there is nothing to determine the value of z , which may be regarded as an arbitrary function of x ; y and v are then functions of x , and w denotes the quotient $y \div v$ of these two functions, and as such satisfies an equation the form of which will depend on the assumed relation between z and x . In particular, if P and Q denote the same functions of z that p and q are of x ; and if we assume $z = x$, P, Q will become $= p, q$ respectively: the given equation in v will be

$$\frac{d^2v}{dx^2} + p \frac{dv}{dx} + qv = 0;$$

and w will thus denote the quotient of any two solutions of the equation

$$\frac{d^2y}{dx^2} + p \frac{dy}{dx} + qy = 0;$$

viz. writing $X = p^2 + 2\frac{dp}{dx} - 4q$, then, by what precedes, the equation for w will be

$$\{w, x\} = -\frac{1}{2}X.$$

65. Returning now to Kummer's problem, and considering y, v as solutions of the two differential equations respectively, we have $y = wv$, and taking any other two solutions denoted by these letters y_1, v_1 we have $y_1 = wv_1$, so that $\frac{y_1}{v_1} = \frac{y}{v}$: w is a function independent of the particular solutions of y, v ; calling the quotient of two solutions of the equation in v, s , also s_1 , denoting the quotient of two solutions of the equation in y , we have

$$\{s, x\} = -\frac{1}{2}Z, \quad \{s_1, z\} = -\frac{1}{2}Z,$$

and similarly $Z = P^2 + 2\frac{dP}{dz} - 4Q$, whence, writing as before $X = p^2 + 2\frac{dp}{dx} - 4q$, we have each of these equal quantities

we obtain

$$\{s, x\} = -\frac{1}{2}X + \frac{1}{2}Z\left(\frac{dz}{dx}\right)^2,$$

as the required equation for the determination of z as a function of x . The process does not give the value of w , but this can be found without difficulty, viz.

$$\frac{1}{w} \frac{dw}{dx} = \frac{1}{2}(P \partial z / \partial x - p) = \frac{1}{2}\left(P \frac{dz}{dx} - p\right).$$

If z, x are regarded each of them as a function of the new independent variable θ , then the equation is

$$\{s, \theta\} = -\frac{1}{2}X\left(\frac{dx}{d\theta}\right)^2 + \frac{1}{2}Z\left(\frac{dz}{d\theta}\right)^2.$$

66. Jacobi's differential equation of the third order for the transformed modulus λ , *Fund. Nova*, p. 78, [*Ges. Werke*, t. I, p. 132], is

$$3(k''\lambda'' - \lambda''k'')^2 - 2k'\lambda'(k'\lambda''' - \lambda'k''') + k'^2\lambda'^2\left[\left(\frac{k''}{k'}\right)^2 k''' - \left(\frac{\lambda''}{\lambda'}\right)^2 \lambda'''\right] = 0,$$

where the accents denote differentiations in regard to an independent variable θ : viz. dividing by $2k'^2\lambda'^2$, this becomes

$$\{k, \theta\} + \frac{1}{2}k'^2\left(\frac{1+k^2}{1-k^2}\right)^2 = \{\lambda, \theta\} + \frac{1}{2}\lambda'^2\left(\frac{1+\lambda^2}{1-\lambda^2}\right)^2,$$

which is thus a particular case of Kummer's equation, k, λ corresponding to x, z respectively, and the values of X, Z being

$$X = -\left(\frac{1+k^2}{1-k^2}\right)^2, \quad Z = -\left(\frac{1+\lambda^2}{1-\lambda^2}\right)^2.$$

67. In the case of the hypergeometric series, the two differential equations of the second order are

$$\frac{d^2y}{dx^2} + \frac{\gamma - (\alpha + \beta + 1)x}{x(1-x)} \frac{dy}{dx} - \frac{\alpha\beta y}{x(1-x)} = 0,$$

$$\frac{d^2v}{dz^2} + \frac{\gamma' - (\alpha' + \beta' + 1)z}{z(1-z)} \frac{dv}{dz} - \frac{\alpha'\beta' v}{z(1-z)} = 0.$$

Hence

$$p = \frac{\gamma + (1-x)(\gamma - \alpha - \beta - 1)x}{x(1-x)} = \frac{\gamma}{x} + \frac{\gamma - \alpha - \beta - 1}{1-x}, \quad q = -\frac{\alpha\beta}{x(1-x)},$$

and hence

$$X = p^2 + 2\frac{dp}{dx} - 4q = \frac{1-2\gamma}{x^2} + \frac{(\gamma - \alpha - \beta - 1)^2 + 2(\gamma - \alpha - \beta - 1) + 4\alpha\beta + 2\gamma(\gamma - \alpha - \beta - 1)}{(1-x)^2} + \frac{\text{etc.}}{x(1-x)};$$

viz. writing

$$a = \frac{1}{4}(1 - \lambda^2), \quad b = \frac{1}{4}(1 - \mu^2), \quad c = \frac{1}{4}(1 - \nu^2),$$

where

$$\lambda^2 = (1 - \gamma)^2, \quad \mu^2 = (\gamma - \alpha - \beta)^2, \quad \nu^2 = (\alpha - \beta)^2,$$

and putting in the formula $x - 1 = -(1 - x)$, we have

$$-\frac{1}{4}\left(p^2 + 2\frac{dp}{dx} - 4q\right) = \frac{a}{x^2} + \frac{b}{(x-1)^2} + \frac{-a-b+c}{x(x-1)} = \frac{a}{x^2} + \frac{b}{(x-1)^2} + \frac{c-a-b}{x \cdot x-1} = (a, b, c \ x, x-1);$$

with a like formula for $\frac{1}{4}\left(P^2 + 2\frac{dP}{dz} - 4Q\right)$. We then have

$$y = wv, \quad v = x^{c-1}(1-x)^{\gamma-\alpha-\beta-1} z^{c'}(1-z)^{-\gamma'+\alpha'+\beta'+1} \dots,$$

and the differential equation of the third order for the determination of z is

$$\{z, x\} = (a_1, b_1, c_1 \ z, z-1)\left(\frac{dz}{dx}\right)^2 - (a, b, c \ x, x-1) = 0,$$

where a_1, b_1, c_1 are the same functions of α', β', γ' which a, b, c are of α, β, γ . This is, in effect, Kummer's equation for the transformation of the hypergeometric series.

68. And in like manner the Schwarzian equation for the determination of z , the quotient of two solutions, is

$$\{s, x\} = (a, b, c \ x, x-1).$$

PART II. THE POLYHEDRAL FUNCTIONS.

Origin and Properties. Art. Nos. 69 to 80.

69. The geometrical functions in question are connected with the geometrical forms (these being regarded as situate on a spherical surface), and with the stereographic projections of these figures:—

1. Polygon or
2. Double Pyramid,*
3. Tetrahedron,
4. Octahedron and Cube,
5. Dodecahedron and Icosahedron,

* Prof. Klein regards 1 as belonging to the polygon and 2 to the double pyramid: it seems to me that the fundamental figure, to which 1 and 2 each of them belong, is the polygon.

Consider a spherical surface and upon it any number of points: take at pleasure any point as South Pole, this determines the plane of the equator; and the stereographic projection of any point is the intersection with the plane of the equator of the line joining the point with the South Pole. To fix the ideas take the radius of the sphere as unity: let the axes of x and y be drawn in the plane of the equator in longitudes 0° and 90° respectively, and the axis of z upwards through the North Pole: the position of a point on the sphere is determined by means of its N.P.D. θ and longitude f : moreover we take X, Y, Z for the coordinates of the point on the surface, and x, y for those of its projection; and we then have

$$X, Y, Z = \sin \theta \cos f, \sin \theta \sin f, \cos \theta;$$

$$x = \frac{X}{1+Z} = \tan \frac{1}{2}\theta \cos f, \quad y = \frac{Y}{1+Z} = \tan \frac{1}{2}\theta \sin f,$$

and conversely

$$X, Y, Z = 2x, 2y, 1 - x^2 - y^2 \div (1 + x^2 + y^2).$$

We represent the point (X, Y, Z) on the spherical surface by means of the magnitude $x+iy = \tan \frac{1}{2}\theta (\cos f + i \sin f)$, or say by the linear factor $s - (x+iy)$: and similarly any system of points on the surface by means of the system of magnitudes $x+iy$, or say by the function $\prod[s - (x+iy)]$, denoting in this manner the product of the linear factors which correspond to the different points respectively.

70. It will presently appear that, if (considering different stereographic projections, that is, a different position of the South Pole) we take x', y' as the coordinates of the new projection of the point, then $x' + iy'$ is a homographic function of $x + iy$: and consequently that the functions of s , which belong to the different projections, are linear transformations one of the other: but at present we consider a single projection.

It may be proper to remark that the figures in question are spherical figures having for summits points on the spherical surface, for sides arcs of great circle joining two summits, and faces which are portions of the spherical surface: the mid-points of the sides (or edges) and the centres of the faces are of course points on the spherical surface.

71. (1), (2). Considering a regular polygon formed by n summits on the equator, the longitude of one of them being 0° , then the stereographic projections correspond with the points themselves, and the values of $x + iy$ are

$$1, \cos \frac{2\pi}{n} + i \sin \frac{2\pi}{n}, \dots, \cos \frac{(n-1)2\pi}{n} + i \sin \frac{(n-1)2\pi}{n}.$$

The corresponding function of s is $s^n - 1$.

The values of $x + iy$ for the mid-points of the sides are

$$\cos \frac{\pi}{n} + i \sin \frac{\pi}{n}, \cos \frac{3\pi}{n} + i \sin \frac{3\pi}{n}, \dots, \cos \frac{(2n-1)\pi}{n} + i \sin \frac{(2n-1)\pi}{n}.$$

The corresponding function of s is $s^n + 1$.

The North and South Poles, which form with the n points a double pyramid of $n+2$ summits, correspond to the values $s = 0$ and $s = \infty$. We have thus

$$s(1 - s^n)(s^n - 1)$$

as the function corresponding to the double pyramid.

72. (3). Considering for a moment the tetrahedron as a figure with rectilinear edges, this is so placed that two opposite edges are horizontal, and that the vertical planes passing through the centre and the upper and lower edges respectively are the meridian planes; the upper edge has the longitudes $\pm 45^\circ$ and the lower edge the longitudes $135^\circ, 225^\circ$. We thus explain the position of the spherical figure.

Corresponding to the summits we have the function $s^4 - 2i\sqrt{3}s^2 + 1$. In fact, the equation $s^4 - 2i\sqrt{3}s^2 + 1 = 0$ gives $s^2 = i(\sqrt{3} \pm 2)$, and hence the values of s are the four values of $x + iy$ shown in the annexed table for the values of X, Y, Z , and $x + iy$ for the summits of the tetrahedron.

long.	45°	135°	225°	315°
X	$+\frac{1}{\sqrt{3}}$	$-\frac{1}{\sqrt{3}}$	$-\frac{1}{\sqrt{3}}$	$+\frac{1}{\sqrt{3}}$
Y	$+\frac{1}{\sqrt{3}}$	$+\frac{1}{\sqrt{3}}$	$-\frac{1}{\sqrt{3}}$	$-\frac{1}{\sqrt{3}}$
Z	$+\frac{1}{\sqrt{3}}$	$+\frac{1}{\sqrt{3}}$	$+\frac{1}{\sqrt{3}}$	$+\frac{1}{\sqrt{3}}$
$x + iy$	$\frac{1+i}{\sqrt{3}-1}$	$\frac{-1+i}{\sqrt{3}+1}$	$\frac{-1-i}{\sqrt{3}-1}$	$\frac{1-i}{\sqrt{3}+1}$

Corresponding to the centres of the faces, or summits of the opposite tetrahedron, we have the function $s^4 + 2i\sqrt{3}s^2 + 1$.

Corresponding to the mid-points of the sides, we have the function

$$s(1 - s^4)(s^4 - 1);$$

viz. the points in question are the four points $s = \pm 1, s = \pm i$, the North Pole $s = 0$, the South Pole $s = \infty$, and the four points on the equator at longitudes $0^\circ, 90^\circ, 180^\circ, 270^\circ$.

73. (4). The octahedron is placed with two of its summits as poles, and the other four summits in the equator at longitudes $0^\circ, 90^\circ, 180^\circ, 270^\circ$ respectively: the values of s are, as in the last case, $0, \infty, \pm 1, \pm i$, and the function is

$$s(1 - s^4)(s^4 - 1).$$

The function for the centres of the faces, or summits of the cube, is $s^8 + 14s^4 + 1$.

The function for the mid-points of the sides of the octahedron or of the cube is

$$s^{12} - 33s^8 - 33s^4 + 1.$$

74. (5). The Icosahedron is placed with two of its summits for poles; five summits lying in a small circle above the plane of the equator at longitudes $0^\circ, 72^\circ, 144^\circ, 216^\circ, 288^\circ$; and the remaining five summits in the corresponding small circle below the equator at longitudes $36^\circ, 108^\circ, 180^\circ, 252^\circ$ and 324° .

The function for the summits of the Icosahedron is

$$s(1 - s^{10})(s^{10} + 11s^5 - 1).$$

The function for the centres of the faces of the Icosahedron, or summits of the Dodecahedron, is

$$s^{20} - 228s^{15} + 494s^{10} + 228s^5 - 1.$$

The function for the mid-points of the sides of the Icosahedron or the Dodecahedron is

$$s^{30} - 522s^{25} + 10005s^{20} + 0s^{15} - 10005s^{10} + 522s^5 + 1.$$

I give for the present these results without demonstration.

75. Writing $\frac{x}{y}$ for the s , so as to obtain from the foregoing values $x + iy$ of these homogeneous functions ($; x, y$) ^{n} , —it will be recollected that the x, y have nothing to do with the x, y of § 69—the

forms which have thus presented themselves may be denoted as follows:

$$\begin{aligned}
 (3) : \quad & f3 = (1, -2i\sqrt{3}, 1; x, y)^4, \\
 & h3 = (f3, f3)^2, \quad t3 = xy(x^4 - y^4), \\
 (4) : \quad & f4 = xy(x^4 - y^4), \\
 & h4 = (1, 14, 1; x^4, y^4)^2, \\
 & t4 = (1, -33, -33, 1; x^4, y^4)^3, \\
 (5) : \quad & f5 = xy(1, 11, -1; x^5, y^5)^2, \\
 & h5 = (1, -228, +494, +228, -1; x^5, y^5)^4, \\
 & t5 = (1, -522, 10005, 0, -10005, 522, 1; x^5, y^5)^6;
 \end{aligned}$$

where observe that $f4$ is the same function as $t3$. In each set of functions f, h, t , we have h and t covariants of f , viz. disregarding numerical factors, h is the Hessian, or derivative $(f, f)^2$, and t is the derivative (f, h) .

76. Since $f4$ is the same function as $t3$, we have of course $f4, h4$ and $t4$ themselves covariants of $f3$; but it is convenient to separate the two systems.

77. It is to be observed that $f3$ is a quartic function having its quadrinvariant $(f) = 0$; but independently of this, $f3$, qua quartic function, has only the covariants $h3$ and $t3$ (the Hessian and the cubicovariant respectively), viz. every other covariant is a rational and integral function of $f3, h3$ and $t3$. In particular, $h4$ and $t4$ are rational and integral functions of $f3, h3$ and $t3$; but inasmuch as $f3$ and $h3$ are not covariants of $f4$, this is not a property of $h4$ and $t4$ considered as covariants of $f4$, and the relation in question need not be attended to.

78. It has been just stated that $f3$ qua quartic function has only the covariants $h3$ and $t3$: $f4$ qua special sextic function (in the sense explained) and $f5$ qua dodecadic function have the like property, viz. $f4$ has only the covariants $h4$ and $t4$; $f5$ only the covariants $h5$ and $t5$. Hence $f3, f4, f5$ are "Prime-forms" in the sense defined in the paper by Fuchs, of 1875, and itself, viz. a Prime-form has no covariant of a lower order than itself, and also no covariant of a higher order which is a power of a form of a lower order.

79. The same functions have also the property that they are functions transformable into themselves by means of a group of linear transformations, and in this point of view they were considered in the nearly contemporaneous paper by Klein, of 1875; it is in this paper shown that the functions so transformable into themselves must be Polyhedral functions as above, the linear transformations in fact corresponding to the rotations whereby the spherical polyhedron can be brought into coincidence with its own original position. This theory will be presently given.

80. It is to be observed that, if U, V are functions $(; x, y)^n$ of the order n , then using the accent to denote differentiation in regard to x , $UV' - U'V$ and (U, V) differ only by a numerical factor: and further that, writing as before $s = \frac{x}{y}$, and in the expression $UV' - U'V$ regarding U, V as functions $(; s, 1)^n$, and the accent as denoting differentiation in regard to s , we have $UV' - U'V$ and (U, V) differing by a numerical factor only. We have in the PQR -Table, the lines 3, 4, 5, P, Q, R equal to given numerical multiples of $h^\alpha, t^\beta, f^\gamma$, the indices α, β, γ being such as to make these to be functions of the same degree: hence, neglecting numerical multipliers,

$$PQ' - P'Q = h^{\alpha-1}t^{\beta-1}(h, t),$$

and the theorem that $PQ' - P'Q$ is equal to a function $(; f^\delta)$, which contains only factors of R , is in fact the theorem that $(h, t), (h, f)$, and (t, f) are each of them equal to a term or product of f, h, t ; which is a result included in the theorem that f has only the covariants h and t . And by this last theorem we know already how from R , assumed to be known, we can derive P and Q : viz. R is a power of f ; and we thence have $h = (f, f)^2$ and $t = (h, f)$, equations giving the functions h and t , upon which P, Q depend.

81. The various covariantive formulae will be given with their proper numerical coefficients.

Tetrahedron function. f, h, t stand for the before-mentioned values, f^3, h^3, t^3

$$(P, Q, R = h^3, -12i\sqrt{3}t^2, -f^6).$$

For f^3 .

$$\begin{aligned}(a, b, c, d, e) &= (1, 0, -\frac{1}{3}i\sqrt{3}, 0, 1), \\ \frac{1}{4}(f, f)^2 &= -\frac{1}{2}i\sqrt{3}h, \quad \frac{1}{4}(f, f)^4 = \frac{4}{3}, \quad \frac{1}{4}(f, f)^6 = 0, \\ (f, h) &= -32i\sqrt{3}t, \quad \frac{1}{4}(f, h)^2 = 1152f = 1152\frac{4}{3}f^2, \\ (h, t) &= 96i\sqrt{3}f, \quad \frac{1}{4}(t, t)^2 = -8fh^2, \\ (f, t) &= -96i\sqrt{3}h^2, \\ h^3 - f^3 - 12i\sqrt{3}t^2 &= 0, \\ fh &= (1, 14, 1; x^4, y^4)^2 (= f^4).\end{aligned}$$

It is convenient to remark that t^2, f^3, h^3 being of the same order, we have

$$P(f, h) + Q(f, t)^2 + R(h, t)^2 = 0,$$

that is,

$$t^2 \cdot 3f^2(f, h) + f^3 \cdot 2h(h, t) + h^3 \cdot 2 \cdot 3t(t, f) = 0,$$

an equation which, substituting the values for $(f, h), (h, t), (t, f)$ their values, reduces itself to $h^3 - f^3 - 12i\sqrt{3}t^2 = 0$; and we have thus a verification of the before-mentioned relation and the values of $(f, h), (h, t)$ and (t, f) . The like remark applies to the other two cases, which follow.

82. *Hexahedron function.* f, h, t stand for the before-mentioned values

$$f^4, h^4, t^4 \quad (P, Q, R = h^3, -t^2, -108f^4).$$

For f^4 .

$$\begin{aligned}(a, b, c, d, e, f, g) &= (0, \frac{1}{6}, 0, 0, 0, -\frac{1}{6}, 0), \\ \frac{1}{4}(f, f)^2 &= -25h, \quad \frac{1}{4}(f, f)^4 = 0, \quad \frac{1}{4}(f, f)^6 = (720)^2 \frac{1}{6}, \\ (f, h) &= -8t, \quad \frac{1}{4}(f, h)^2 = 3 \cdot 2^8 \cdot 7^2 \cdot h^2, \\ (f, t) &= -12h^2, \quad \frac{1}{4}(t, t)^2 = 2^4 \cdot 3^2 \cdot 17^2 \cdot f^2, \\ (h, t) &= -1728f^3, \\ h^3 - t^2 - 108f^4 &= 0.\end{aligned}$$

83. *Dodecahedron function.* f, h, t stand for the before-mentioned values

$$f^5, h^5, t^5 \quad (P, Q, R = h^3, -t^2, -1728f^5).$$

For f^5 .

$$\begin{aligned}(a, b, c, d, e, f, g, h, i, j, k, l, m) &= (0, \frac{1}{12}, 0, 0, 0, 0, \frac{11}{12}, 0, 0, 0, 0, -\frac{1}{12}, 0), \\ \frac{1}{4}(f, f)^2 &= -121h, \quad \frac{1}{4}(f, f)^4 = 0, \quad \frac{1}{4}(f, f)^6 = \frac{1}{4}(924)^2(720)^2 \frac{11}{12}f^2, \\ \frac{1}{4}(f, f)^8 &= \frac{1}{4}(924)^2(720)^2 \frac{1}{12}f^2, \quad \frac{1}{4}(f, f)^{10} = 0, \quad \frac{1}{4}(f, f)^{12} = 0, \\ (f, h) &= -20t, \quad \frac{1}{4}(f, h)^2 = 173280h^2, \\ (f, t) &= -30h^2, \quad \frac{1}{4}(t, t)^2 = 9082800f^2h, \\ (h, t) &= -86400f^4, \\ h^3 - t^2 - 1728f^5 &= 0.\end{aligned}$$

84. We have

$$t = (x^5 + y^5) (1, 522, -10005, -522, 1; x^5, y^5)^4.$$

Write

$$\xi = (x^5 + y^5) (1, 2, 6, -2, 1; x^5, y^5)^4,$$

then

$$t = \xi (1, -10, 45; x, y)^2.$$

Or putting

$$\rho = \frac{1}{xy} \sqrt[4]{(x^5 + y^5) (1, 2, 6, -2, 1; x^5, y^5)^4},$$

that is,

$$\rho = \frac{1}{xy} \sqrt[4]{(x^{10} + 11x^5y^5 - y^{10}) (\dots)},$$

Writing $k = P/f$, then

$$\rho^3 - 10\rho^2 + 45\rho = \frac{k}{f}. \quad (\text{Klein.})$$

Investigation of the forms f5 and h5. Art. Nos. 85 and 86.

85. Writing for shortness¹ $k = \tan \alpha = \sqrt{\frac{5 - \sqrt{5}}{10}}$, and $g = \cos 36^\circ + i \sin 36^\circ$, then the values of $x + iy$ corresponding to the summits of the Icosahedron are

$$0, k, kg, kg^2, kg^3, kg^4, k^{-1}, k^{-1}g, k^{-1}g^2, k^{-1}g^3, k^{-1}g^4, \infty,$$

and the function $f5$ is thus

$$= s(1 - s^5/k^5)(s^5 - k^{-5}),$$

where the product of the last two factors is $s^{10} + (k^{-5} - k^5)s^5 - 1$. We have

$$k^{-5} - k^5 = \frac{1}{10}\sqrt{5}(5\sqrt{5} - 11), \quad = \frac{1}{2}(\sqrt{5} - 11) \cdot \sqrt{5}/5$$

and consequently $k^{-5} - k^5 = 11$; or the function is

$$s(1 - s^{10})(s^{10} + 11s^5 - 1).$$

86. Similarly, writing for shortness²

$$\cos \frac{\gamma}{2} = \frac{1}{2}\sqrt{\frac{5 + \sqrt{5}}{2\sqrt{5}}}, \quad \cos^2 \frac{\gamma}{2} = \frac{5 + 2\sqrt{5}}{15}, \quad \sin^2 \frac{\gamma}{2} = \frac{10 - 2\sqrt{5}}{15},$$

and

$$l = \tan \frac{1}{2}\gamma, \quad l' = \tan \frac{1}{2}\gamma', \quad \text{where } \cos \frac{\gamma'}{2} = \frac{3 + \sqrt{5}}{\dots}, \quad \sin \frac{\gamma'}{2} = \frac{3 - \sqrt{5}}{\dots},$$

and therefore

$$l^5 = \frac{\cos \gamma}{\sin \gamma};$$

$g = \cos 36^\circ + i \sin 36^\circ$ as before, then the values of $x + iy$ for the summits of the dodecahedron are

$$lg, lg^2, lg^3, lg^4, lg^5; \quad l'g, l'g^2, l'g^3, l'g^4, l'g^5;$$

$$l^{-1}, l^{-1}g, l^{-1}g^2, l^{-1}g^3, l^{-1}g^4; \quad l'^{-1}, l'^{-1}g, l'^{-1}g^2, l'^{-1}g^3, l'^{-1}g^4.$$

¹ k is the α and h the β of the Table, No. 99.

² α is the α , γ is the γ , and γ' the $\alpha - \beta$ of the Table, No. 99.

The function $h5$ is therefore

$$= s^{20} + s^{15}(l^5 - l^{-5}) + l^5 \cdot l'^5 s^{10} + s^5(l'^5 - l'^{-5}) - 1.$$

We have

$$l^{-5} - l^5 = \frac{2 \cos \gamma}{\sin^5 \gamma} = \frac{(1 - \cos \gamma)^5}{\sin^5 \gamma} \cdot \frac{1}{(1 + \cos \gamma)^5} = \frac{(5 + 10 \cos^2 \gamma + \cos^4 \gamma)}{\sin^5 \gamma} = \frac{384 + 128 \cos \gamma}{45 \sin^5 \gamma} = \frac{64\sqrt{5}}{45} (6 + \sqrt{5}) = \frac{2}{45} (114 + 50\sqrt{5})$$

viz. this last identity depends on

$$\frac{2}{45} (3 + \sqrt{5})(6 + \sqrt{5}) = (114 + 50\sqrt{5}) \sin^4 \gamma,$$

that is,

$$160 (3 + \sqrt{5})(6 + \sqrt{5}) = (114 + 50\sqrt{5})(120 - 40\sqrt{5}),$$

or

$$2 (3 + \sqrt{5})(6 + \sqrt{5}) = (57 + 25\sqrt{5})(3 - \sqrt{5}),$$

or finally

$$(7 + 3\sqrt{5})(6 + \sqrt{5}) = 57 + 25\sqrt{5},$$

which is right.

Similarly

$$l'^{-5} - l'^5 = 114 - 50\sqrt{5},$$

and observing that the sum and product of $114 + 50\sqrt{5}$, $114 - 50\sqrt{5}$ are $= 228$ and 496 respectively, the required function of s is

$$\begin{aligned} &= s^{20} - 1 - 228 (s^{15} - s^5) + 496 s^{10}, \\ &= s^{20} - 228 s^{15} + 494 s^{10} + 228 s^5 + 1, \\ & \quad (= h5) \end{aligned}$$

which is the required value of $h5$.

Invariantive property of the Stereographic Projection.

Art. Nos. 87 to 93.

87. The before-mentioned theorem that the functions derived from two different stereographic projections of the same point are linear transformations one of the other, may be thus stated:—

Considering on the surface of a sphere, two fixed points A and B ; and determining the position of a point C , first in regard to A by its distance θ and azimuth f , and next in regard to B by its distance θ' and azimuth f' , the azimuths from the great circle ABx which joins the two points A and B , then we have

$$s = \tan \frac{1}{2} \theta (\cos f + i \sin f), \quad s' = \tan \frac{1}{2} \theta' (\cos f' + i \sin f'),$$

calling them homographic functions one of the other; and putting the distance $AB = c$, the relation between them in fact is

$$s' = \frac{s - \tan \frac{1}{2} c}{1 + s \tan \frac{1}{2} c};$$

or, what is the same thing, observing that

$$s s' = \tan \frac{1}{2} \theta \tan \frac{1}{2} \theta' \{ \cos(f + f') + i \sin(f + f') \},$$

we have the two equations

$$\tan \frac{1}{2} c \{ 1 + \tan \frac{1}{2} \theta \tan \frac{1}{2} \theta' \cos(f + f') \} = \tan \frac{1}{2} \theta \cos f - \tan \frac{1}{2} \theta' \cos f',$$

$$\tan \frac{1}{2} c \{ 1 + \tan \frac{1}{2} \theta \tan \frac{1}{2} \theta' \sin(f + f') \} = \tan \frac{1}{2} \theta \sin f - \tan \frac{1}{2} \theta' \sin f'.$$

88. If we denote the angles of the spherical triangle C, A, B , and the opposite sides by c (as before), a, b , then $\theta, \theta' = b, a; f, f' = A, \pi - B$, whence

$$s = \tan \frac{1}{2}b (\cos A + i \sin A), \quad s' = -\tan \frac{1}{2}a (\cos B - i \sin B),$$

and we have between the spherical sides a, b, c and angles A, B of the spherical triangle the relations

$$\tan \frac{1}{2}c \{1 - \tan \frac{1}{2}a \tan \frac{1}{2}b \cos(A - B)\} = \tan \frac{1}{2}b \cos A + \tan \frac{1}{2}a \cos B,$$

$$\tan \frac{1}{2}c \{1 - \tan \frac{1}{2}a \tan \frac{1}{2}b \sin(A - B)\} = \tan \frac{1}{2}b \sin A - \tan \frac{1}{2}a \sin B;$$

which equations may be verified by means of the ordinary formulae of Spherical Trigonometry.

89. But it is interesting to give the proof with rectangular coordinates. Taking (X, Y, Z) , (X_1, Y_1, Z_1) for the coordinates, referred to two different sets of axes, of a point on the spherical surface: also x, y, x_1, y_1 for the coordinates of the corresponding stereographic projections, we have

$$(X_1 : Y_1 : Z_1) = (\alpha, \beta, \gamma; \alpha', \beta', \gamma'; \alpha'', \beta'', \gamma'')(X : Y : Z),$$

$$X : Y : Z : 1 = 2x : 2y : 1 - x^2 - y^2 : 1 + x^2 + y^2,$$

$$X_1 : Y_1 : Z_1 : 1 = 2x_1 : 2y_1 : 1 - x_1^2 - y_1^2 : 1 + x_1^2 + y_1^2,$$

and thence

$$x_1 : y_1 : 1 = 2\alpha x + 2\beta y + \gamma(1 - x^2 - y^2) : 2\alpha'x + 2\beta'y + \gamma'(1 - x^2 - y^2) : 1 + x^2 + y^2 + 2\alpha''x + 2\beta''y + \gamma''(1 - x^2 - y^2).$$

90. Introducing z, z_1 for homogeneity, or writing $\frac{x}{z}, \frac{y}{z}$ and $\frac{x_1}{z_1}, \frac{y_1}{z_1}$ in place of x, y and x_1, y_1 , respectively, we have

$$x_1 = 2\alpha x + 2\beta y + \gamma(z^2 - x^2 - y^2) = (-\gamma, -\gamma, +\gamma, \beta, \alpha, 0; x, y, z)^2,$$

$$y_1 = 2\alpha'x + 2\beta'y + \gamma'(z^2 - x^2 - y^2) = (-\gamma', -\gamma', +\gamma', \beta', \alpha', 0; x, y, z)^2,$$

$$z_1 = z^2 + x^2 + y^2 + 2\alpha''x + 2\beta''y + \gamma''(z^2 - x^2 - y^2) = (1 - \gamma'', 1 - \gamma'', 1 + \gamma'', \beta'', \alpha'', 0; x, y, z)^2,$$

and thence without difficulty

$$x_1 + iy_1 = \frac{1}{\gamma + \gamma''} \{(1 + \gamma'')z + (\alpha'' + i\beta'')(x - iy)\} \{(1 - \gamma'')z + (-\alpha'' + i\beta'')(x + iy)\},$$

$$z_1 = \frac{1}{\gamma + \gamma''} \{(1 - \gamma'')z - (\alpha'' + i\beta'')(x - iy)\} \{(1 + \gamma'')z + (\alpha'' - i\beta'')(x + iy)\},$$

$$x_1 - iy_1 = MN : NL : LM \quad (L, M, N \text{ linear functions of } z, x + iy, x - iy)$$

showing that the relation between two stereographic projections of the same spherical figure is in fact that of a quadric transformation, the fundamental points in each figure being an arbitrary point and the two circular points at infinity: or, what is the same thing, to any line in the one figure there corresponds a circle in the other figure, which is the “circular relation” of Möbius.

91. The actual values are

$$\frac{x_1 + iy_1}{z_1} = \frac{1 + \gamma''}{\gamma + \gamma''} \cdot \frac{(1 - \gamma'')z - (\alpha'' - i\beta'')(x + iy)}{(1 + \gamma'')z + (\alpha'' - i\beta'')(x + iy)},$$

$$\frac{x_1 - iy_1}{z_1} = \frac{1 + \gamma''}{\gamma - \gamma''} \cdot \frac{(1 - \gamma'')z - (\alpha'' + i\beta'')(x - iy)}{(1 + \gamma'')z + (\alpha'' + i\beta'')(x - iy)},$$

viz. attending only to the former of these, we have $\frac{x_1 + iy_1}{z_1}$ a homographic function of $\frac{x + iy}{z}$, which is the before-mentioned theorem.

92. Supposing that the transformation from (X, Y, Z) to (X_1, Y_1, Z_1) is made by a rotation, the coordinates of which are λ, μ, ν , that is, if f, g, h are the inclinations of the resultant axis to the axes of

x, y, z respectively, and θ the angle of rotation, putting $\lambda, \mu, \nu = \tan \frac{1}{2}\theta \cos f, \tan \frac{1}{2}\theta \cos g, \tan \frac{1}{2}\theta \cos h$: then the coefficients of transformation are

$$\begin{pmatrix} \alpha & \beta & \gamma \\ \alpha' & \beta' & \gamma' \\ \alpha'' & \beta'' & \gamma'' \end{pmatrix} = \frac{1}{1 + \lambda^2 + \mu^2 + \nu^2} \begin{pmatrix} 1 + \lambda^2 - \mu^2 - \nu^2 & 2(\lambda\mu - \nu) & 2(\lambda\nu + \mu) \\ 2(\lambda\mu + \nu) & 1 - \lambda^2 + \mu^2 - \nu^2 & 2(\mu\nu - \lambda) \\ 2(\lambda\nu - \mu) & 2(\mu\nu + \lambda) & 1 - \lambda^2 - \mu^2 + \nu^2 \end{pmatrix}.$$

Substituting these values, the formulae become, after an easy reduction,

$$\frac{x_1 + iy_1}{z_1} = \frac{-(\nu + i)(x + iy) + (\lambda + i\mu)z}{(\lambda - i\mu)(x + iy) + (\nu - i)z},$$

$$\frac{x_1 - iy_1}{z_1} = \frac{-(\nu - i)(x - iy) + (\lambda - i\mu)z}{(\lambda + i\mu)(x - iy) + (\nu + i)z},$$

attending to the former of these, and writing for greater simplicity s, s_1 respectively, we have

$$s_1 = \frac{-(\nu + i)s + (\lambda + i\mu)}{(\lambda - i\mu)s + (\nu - i)},$$

or writing this

$$s_1 = \frac{As + B}{Cs + D},$$

then

$$A : B : C : D = -\nu - i : \lambda + i\mu : \lambda - i\mu : \nu - i.$$

93. I call to mind that the condition, in order that the homographic transformation $s_1 = (As + B) \div (Cs + D)$ may be periodic of the order n , is

$$(A + D)^2 - 4(AD - BC) \cos^2 \frac{m\pi}{n} = 0,$$

m being an integer different from zero and prime to n . In particular, when $n = 2$, it is $A + D = 0$: $n = 3$, it is $A^2 + AD + D^2 + BC = 0$: $n = 4$, it is $A^2 + D^2 + 2BC = 0$: and $n = 5$, it is $(A + D)^2 - \frac{1}{2}(3 \pm \sqrt{5})(AD - BC) = 0$.

Groups of homographic transformations. Art. Nos. 94 and 95.

94. The formulae just obtained serve to connect the theory of the rotations of a polyhedron with that of the homographic transformations s into $(As + B) \div (Cs + D)$; and, corresponding to the rotations which leave the polyhedron unaltered, we have groups of homographic transformations. We have thus, corresponding to the cases of the tetrahedron, the cube and the octahedron, and the dodecahedron and icosahedron respectively, groups of 12, of 24, and of 60 homographic transformations s into

$(As + B) \div (Cs + D)$. The group of 60 and the group of 24 include each of them as part of itself the group of 12; it is further to be remarked that the group of 12 may be regarded as that of the positive substitutions upon the four letters $abcd$, the group of 24 as that of all the substitutions upon the four letters, and the group of 60 as that of the positive substitutions upon the five letters $abcde$.

95. I call to mind that a group of functional symbols $1, \alpha, \beta, \dots$ can always be expressed in the equivalent form $1, \phi\alpha\phi^{-1}, \phi\beta\phi^{-1}, \dots$, where ϕ is any functional symbol whatever: clearly, $\alpha, \beta, \dots$ being homographic transformations, then, ϕ being any homographic transformation whatever, the new symbols $\phi\alpha\phi^{-1}, \phi\beta\phi^{-1}, \dots$ will also be homographic transformations; and thus the group of homographic transformations can be expressed in various equivalent forms: these correspond to the different positions of the polyhedron in regard to the axes of coordinates; viz. in the three cases which it is proper to consider, the dodecahedron, we may have the Pole at a point Θ , that is, in the language of art. Nos. as presently explained, the axis of z passing through the midpoint of a side, 1°: through a summit; that is, the Pole at a point A , 2°: and there are different forms which correspond to a Pole at a point B , 3°.


The regular Polyhedra. Art. Nos. 96 to 103.

96. We require a theory of the regular Polyhedra considered as systems of points on a sphere. I refer to my two papers [375] and [679]. In the latter paper, I remark that, considering the five regular figures drawn in proper relation to each other on the same spherical surface, the only points which have to be considered are 12 points A , 20 points B , 30 points Θ , and 60 points Φ . Describing these by reference to the dodecahedron, the points A are the centres of the faces, the points B are the summits, the points Θ are the midpoints of the sides, and the points Φ are the midpoints of the diagonals of the faces, each point Φ being given as the intersection of a face of the dodecahedron and a side of the icosahedron, which there intersect at right angles. Or describing them by reference to the icosahedron, the points A are the summits, the points B are the centres of the faces, the points Θ are the midpoints of the sides: and the points Φ are points lying by threes on the faces of the icosahedron, which there is no occasion to attend to; viz. Φ is the point common to the face and a line BB joining the centres of two adjacent faces at right angles.

97. The points Φ are comparatively unimportant, and it is proper first to attend only to the 12 points A , the 20 points B , and the 30 points Θ : these form 6 pairs of opposite points A , 10 pairs of opposite points B , and 15 pairs of opposite points Θ . Considering the diameters through each pair of opposite points Θ , we have thus a system of 15 axes, which in fact form 5 sets each of 3 rectangular axes: attending to any one of such sets, the diametral plane at right angles to one of the three axes contains of course the other two axes: it contains also two axes each through a pair of opposite points A , and two axes each through a pair of opposite points B . If instead of the plane we consider its intersection with the sphere, we have thus on the sphere 15 circles each containing 4 points Θ ,

4 points A and 4 points B . The fifteen circles intersect by fives in the pairs of opposite points A , by threes in the pairs of opposite points B , and by twos in the pairs of opposite points Θ ; the mutual inclinations of successive circles at the points A , B , Θ being $= 36^\circ$, 60° and 90° respectively. The whole number $15 \cdot 14 = 210$, of the intersections of the two circles two and two together is thus made up of the 12 points A each counting 10 times, the 20 points B each counting 3 times, and the 30 points Θ each counting once; $210 = 120 + 60 + 30$.

98. The angular magnitudes which present themselves are all obtained from the dodecahedral pentagon, as shown in the annexed figure, in which the angle subtended by a side at the centre is $= 72^\circ$, and the angle between two adjacent sides is $= 120^\circ$.



We write $A\Theta = \alpha$, $B\Theta = \beta$, $AB = \gamma$, $B_1B_2 = x$, $\angle B_1B_2B = \theta$, $\Theta B_1 = g$, $\angle \Theta B_1B = \phi$.

From the triangle $A\Theta B$, the angles of which are 36° , 90° , 60° and the opposite β , γ , α , we find the values of α , β , γ , and these are such that $\alpha + \beta + \gamma = \frac{1}{2}\pi$.

From the triangle B_2BB_1 , where the sides B_1B , BB_2 and the included angle are 2β , 2β , 120° , we have the opposite side x , and the other two angles each $= \theta$.

From the triangle $B_1\Theta B_2$, where the sides B_1B , $\beta\Theta$ and the included angle $BB_2\Theta$ are 2β , β , 120° , we find the opposite side g , and the angle $B_2\Theta B_1 = 45^\circ$.

Hence each of the angles $B_1\Theta B$, $B_2\Theta B_1$, being $= 45^\circ$, the angle $B_1\Theta B_2$ is $= 90^\circ$: in this triangle the hypotenuse B_1B_2 is $= x$, and the other two sides are $= g$: whence we have $\cos x = \cos^2 g$, as is in fact the case, and moreover the values give $x + 2g = 180^\circ$. Also each of the other angles is found to be $= 60^\circ$; that is, we have $\angle B_1BB_2 = 60^\circ$, or the whole angle at B_1 being $= 120^\circ$, the sum of the remaining angles B_1BB_2 and $BB_1\Theta$ is $= 60^\circ$.

From the triangle $\Theta B_1 \Theta'$ where the two sides and the included angle are $\beta, \beta, 120^\circ$, we find $\Theta \Theta' = 36^\circ$.

And from the triangle $\Theta B_1 \Theta''$, where the two sides and the included angle are g, g , and $(120^\circ - 2\phi) = 2\phi$, we find $\Theta \Theta'' = 60^\circ$.


99. We thus arrive at the following Table:

			cos	sin
$A\Theta$	α	$31^\circ 43'$	$\sqrt{\frac{5 - \sqrt{5}}{10}}$	$\sqrt{\frac{5 + \sqrt{5}}{10}}$
$B\Theta$	β	$20^\circ 55'$	$\frac{\sqrt{5} + 1}{2\sqrt{3}}$	$\frac{\sqrt{10 - 2\sqrt{5}}}{2\sqrt{3}}$
AB	γ	$37^\circ 22'$	$\sqrt{\frac{5 + \sqrt{5}}{15}}$	$\sqrt{\frac{10 - 2\sqrt{5}}{15}}$
(BB)	x	$70^\circ 32'$	$\frac{1}{3}$	$\frac{2\sqrt{2}}{3}$
(BB)		$109^\circ 28'$	$-\frac{1}{3}$	$\frac{2\sqrt{2}}{3}$
(BB)	g	$54^\circ 44'$	$\frac{1}{\sqrt{3}}$	$\frac{\sqrt{2}}{\sqrt{3}}$
BBB	θ	$37^\circ 46'$		
BBB		$54^\circ 44'$		
ΘBB	ϕ			
2β		$41^\circ 50'$	$\sqrt{\frac{5 - 2\sqrt{5}}{15}}$	
2γ		$74^\circ 44'$	$\frac{\sqrt{3}(\sqrt{5} - 1)}{4}$	
$\alpha - \beta$			$\frac{\sqrt{5 - \sqrt{6}}}{2}$	
$\Theta \Theta'$		36°	$\frac{\sqrt{5} + 1}{4}$	
$\Theta \Theta''$		60°	$\frac{1}{2}$	

where as above $\alpha + \beta + \gamma = 90^\circ$, and $x + 2g = 180^\circ$, $\theta + \phi = 60^\circ$.

100. We now construct three figures of the points A, B, Θ ; viz. these are stereographic projections, each showing the Northern hemisphere projected on the plane of the equator by lines drawn to the South Pole: hence, for any pair of opposite points on the equator, only the point in the Northern hemisphere is shown: but for a pair of opposite points on the equator the two points are each of them shown. In fig. 1 the North Pole is taken to be a point Θ ; in fig. 2 it is a point A ; and in fig. 3 it is a point B . The position of any point on the sphere is determined by its N.P.D. and its longitude, say from the point E of the centre left-handedly measured from an arbitrary origin: then, in the three figures, the positions are as follows.


101. Fig. 1. Pole at Θ .



Pole at Θ

	N.P.D.'s	Longitudes
2A	$\alpha = 31^\circ 43'$	$0^\circ, 180^\circ$
2A	$90^\circ - \alpha = 58\ 17$	90, 270
2A	$90^\circ + \alpha = 121\ 43$	$0^\circ, 180^\circ$
2A	$180^\circ - \alpha = 148\ 17$	90, 270
4A	90	$(0, 180) + \alpha = 31^\circ 43'$
2B	$\beta = 20^\circ 55'$	$90^\circ, 270^\circ$
4B	$g = 54\ 44$	45, 135, 225, 315
2B	$90^\circ - \beta = 69\ 5$	0, 180
2B	90	$(90, 270) + \beta = 20^\circ 55'$
4B	$90^\circ + \beta = 110\ 55$	0, 180
4B	$180^\circ - g = 125\ 16$	45, 135, 225, 315
2B	$180^\circ - \beta = 159\ 5$	90, 270
1 Θ	0°	—
4 Θ	36	$(90^\circ, 270^\circ) + \alpha = 31^\circ 43'$
4 Θ	60	$(0, 180) + \beta = 20\ 55$
4 Θ	72	$(90, 270) + \alpha = 31\ 43$
4 Θ	90	0, 90, 180, 270
4 Θ	108	$(90, 270) + \alpha = 31\ 43$
4 Θ	120	$(0, 180) + \beta = 20\ 55$
4 Θ	144	$(90, 270) + \alpha = 31\ 43$
1 Θ	180	—


102. Fig. 2. Pole at A.



Pole at A

	N.P.D.'s	Longitudes
A	0	—
5A	$2\alpha = 63^\circ 26'$	$0^\circ\ 72^\circ\ 144^\circ\ 216^\circ\ 288^\circ$
5A	$180^\circ - 2\alpha = 116\ 34$	36 108 180 252 324
A	180	—
5B	$\gamma = 37\ 22$	36 108 180 252 324
5B	$90^\circ - \alpha + \beta = 79\ 12$	36 108 180 252 324
5B	$90 + \alpha - \beta = 100\ 48$	72 144 216 288
5B	$180 - \gamma = 142\ 38$	72 144 216 288
5 Θ	$\alpha = 31\ 43$	72 144 216 288
5 Θ	$90^\circ - \alpha = 58\ 17$	36 108 180 252 324
10 Θ	90	$(36\ 108\ 180\ 252\ 324) + 18^\circ$
5 Θ	$90 + \alpha = 121\ 43$	72 144 216 288
5 Θ	$180 - \alpha = 144\ 17$	36 108 180 252 324

103. Fig. 3. Pole at B.

Pole at B

	N.P.D.'s	Longitudes
$3A$	$\gamma = 37^\circ 22'$	$30^\circ \ 150^\circ \ 270^\circ$
$3A$	$90^\circ - \alpha + \beta = 79 \ 12$	$90 \ 210 \ 330$
$3A$	$90 + \alpha - \beta = 100 \ 48$	$90 \ 210 \ 330$
$3A$	$180 - \gamma = 142 \ 38$	$90 \ 210 \ 330$
B	0	—
$3B$	$2\beta = 41 \ 50$	$30 \ 150 \ 270$
$6B$	$x = 70 \ 32$	$(90 \ 210 \ 330) + \phi = 37^\circ 46'$
$6B$	$180^\circ - x = 109 \ 28$	$(30 \ 150 \ 270) + \phi = 37^\circ 46'$
$3B$	$180 - 2\beta = 138 \ 10$	$90 \ 210 \ 330$
B	180	—
3Θ	$\beta = 20 \ 55$	$90 \ 210 \ 330$
6Θ	$g = 54 \ 44$	$(90 \ 210 \ 330) + \phi = 22^\circ 14'$
Θ	$60 \ 120 \ 180^\circ \ 240^\circ \ 300^\circ$	—
6Θ	$90 + \beta = 110 \ 55$	$30 \ 150 \ 270$
6Θ	$180 - g = 125 \ 16$	$30 \ 150 \ 270$
3Θ	$180 - \beta = 159 \ 5$	$90 \ 210 \ 330$

The groups of homographic transformations, resumed.

Art. Nos. 104 to 117.

104. The axes of rotation for the dodecahedron and the icosahedron are 15 axes each through a pair of opposite points Θ , 6 axes each through a pair of opposite points A , and 10 axes each through a pair of opposite points B ; or say 15 Θ -axes, 10 B -axes and 6 A -axes; the corresponding angles of rotation are 180° , 72° and 120° so that (excluding in each case the original position or that of a rotation 0) we have in respect of each Θ -axis 1 position, in respect of each A -axis 4 positions, and in respect of each B -axis 2 positions; in all, including the original position,

$$1 + 15 + (6 \times 4) + (10 \times 2), \text{ that is, } = 60 \text{ positions,}$$

that is, a group of 60 rotations.

To find, in any one of the three forms, the group of homographic transformations, we can in each case obtain from the foregoing tables the values $\cos f$, $\cos g$, $\cos h$ of the cosine-inclination of an axis of rotation to the axes of coordinates, and thence calculate the values of

$$\lambda, \mu, \nu = \tan \frac{1}{2}\theta \cos f, \tan \frac{1}{2}\theta \cos g, \tan \frac{1}{2}\theta \cos h,$$

and thence the values of

$$A, B, C, D = -\nu - i, \lambda + i\mu, \lambda - i\mu, \nu - i;$$

viz. in the case of a Θ -axis, θ is $= 180^\circ$, (so that here $\tan \frac{1}{2}\theta = \infty$ or the values of A, B, C, D are $= -\nu, \lambda + i\mu, \lambda - i\mu, \nu$, that is, $-\cos h, \cos f + i\cos g, \cos f - i\cos g, \cos h$); in the case of a B -axis, the values are $\theta = 120^\circ, 240^\circ$, and therefore $\tan \frac{1}{2}\theta = \pm\sqrt{3}$; and in the case of an A -axis, they are $\theta = 72^\circ, 144^\circ, 216^\circ, 288^\circ$, and therefore

$$\tan \frac{1}{2}\theta = \pm \frac{1}{\sqrt{5}} \sqrt{10 + 2\sqrt{5}}, \pm \frac{1}{\sqrt{5}} \sqrt{10 - 2\sqrt{5}}.$$

105. The Θ -form was first given in my paper of 1879, but in obtaining it I used the results given in the paper of 1877. As regards the identification with the substitution-symbols, since there is nothing to distinguish inter se the letters a, b, c, d, e , any transformation A, B, C, D of the fifth order might have been taken for $abcde$, But No. 37 of the group having been taken for this substitution $abcde$, I do not recall in what manner I found that, consistently herewith, the transformation No. 2 ($-1, 0, 0, 1$, that is, s into $-s$) of the second order could be taken for $ab.cd$. But there is no sub-group of an order divisible by 5; and hence, these two transformations being identified with the two substitutions, the other transformations correspond each of them to a determinate substitution.

106. *Homographic Transformations.* The group of 60. Pole at Θ .

No.	$(Ax + B)$	$\div (Cx + D)$	substitution
1	1	1	1
2	-1	1	$ab.cd$
3	i	1	$ac.bd$
4	$-i$	1	$ad.bc$
5	$-3 + \sqrt{5} + i(1 - \sqrt{5})$	2	$ad.be$
6	$-3 + \sqrt{5} + i(-1 + \sqrt{5})$	2	$ae.bd$
7	$3 - \sqrt{5} + i(-1 + \sqrt{5})$	2	$ad.ce$
8	$3 - \sqrt{5} + i(1 - \sqrt{5})$	2	$ae.cd$
9	$-1 - \sqrt{5} + i(1 - \sqrt{5})$	2	$ab.de$
10	$-1 - \sqrt{5} + i(-1 + \sqrt{5})$	2	$bc.de$
11	$1 + \sqrt{5} + i(-1 + \sqrt{5})$	2	$ab.ce$
12	$1 + \sqrt{5} + i(1 - \sqrt{5})$	2	$ac.bd$
13	$-1 - \sqrt{5} + i(-3 - \sqrt{5})$	2	$bd.ce$
14	$-1 - \sqrt{5} + i(3 + \sqrt{5})$	2	$ae.be$
15	$1 + \sqrt{5} + i(3 + \sqrt{5})$	2	$bc.de$
16	$1 + \sqrt{5} + i(-3 - \sqrt{5})$	2	$ac.de$
17	i	1	abc
18	$-i$	1	acb
19	1	$-i$	ade
20	-1	$-i$	aed
21	i	1	abd
22	$-i$	1	adb
23	1	i	bcd
24	-1	i	bdc
25	$-1 - \sqrt{5} + i(-3 - \sqrt{5})$	2	abd
26	$1 + \sqrt{5} + i(3 + \sqrt{5})$	2	ace
27	$1 + \sqrt{5} + i(-3 - \sqrt{5})$	2	bce
28	$-1 - \sqrt{5} + i(3 + \sqrt{5})$	2	bed
29	$-3 + \sqrt{5} + i(1 - \sqrt{5})$	2	bde
30	$-3 + \sqrt{5} + i(-1 + \sqrt{5})$	2	bee
31	$3 - \sqrt{5} + i(-1 + \sqrt{5})$	2	aed
32	$3 - \sqrt{5} + i(1 - \sqrt{5})$	2	ade
33	$-1 - \sqrt{5} + i(3 + \sqrt{5})$	2	cde
34	$1 + \sqrt{5} + i(-3 - \sqrt{5})$	2	ced
35	$1 + \sqrt{5} + i(3 + \sqrt{5})$	-2	aeb
36	$-1 - \sqrt{5} + i(-3 - \sqrt{5})$	2	abe
37	$1 + \sqrt{5} + i(-3 - \sqrt{5})$	2	$abcde$
38	$-1 - \sqrt{5} + i(1 - \sqrt{5})$	2	$acebd$
39	$-1 - \sqrt{5} + i(-1 + \sqrt{5})$	2	$adbec$
40	$-1 - \sqrt{5} + i(3 + \sqrt{5})$	2	$aedcb$
41	$1 + \sqrt{5} + i(3 + \sqrt{5})$	2	$adceb$
42	$1 + \sqrt{5} + i(-1 + \sqrt{5})$	2	$acbde$
43	$1 + \sqrt{5} + i(1 - \sqrt{5})$	2	$aedbe$
44	$1 + \sqrt{5} + i(-3 - \sqrt{5})$	2	$abcad$
45	$-1 - \sqrt{5} + i(-1 + \sqrt{5})$	-2	$abced$

No.	$(Ax + B)$		$\div (Cx + D)$	substitution
46	$-3 + \sqrt{5} + i(-1 + \sqrt{5})$		-2	<i>abdce</i>
47	$3 - \sqrt{5} + i(-1 + \sqrt{5})$		-2	<i>accdb</i>
48	$1 + \sqrt{5} + i(-1 + \sqrt{5})$		2	<i>adcbe</i>
49	$1 + \sqrt{5} + i(1 - \sqrt{5})$		2	<i>aecbd</i>
50	$3 - \sqrt{5} + i(1 - \sqrt{5})$		2	<i>acdeb</i>
51	$-3 + \sqrt{5} + i(1 - \sqrt{5})$		2	<i>abccd</i>
52	$-1 - \sqrt{5} + i(1 - \sqrt{5})$		2	<i>adbcc</i>
53	$-3 + \sqrt{5} + i(1 - \sqrt{5})$		-2	<i>acbde</i>
54	$3 - \sqrt{5} + i(1 - \sqrt{5})$		2	<i>aebde</i>
55	$-3 + \sqrt{5} + i(-1 + \sqrt{5})$		2	<i>abccd</i>
56	$-1 - \sqrt{5} + i(-3 - \sqrt{5})$		2	<i>adecb</i>
57	$1 + \sqrt{5} + i(3 + \sqrt{5})$		2	<i>acdbc</i>
58	$3 - \sqrt{5} + i(1 - \sqrt{5})$		2	<i>abdec</i>
59	$-3 + \sqrt{5} + i(1 - \sqrt{5})$		2	<i>adcbe</i>
60	$-1 - \sqrt{5} + i(-1 + \sqrt{5})$		2	<i>aebcd</i>

107. Taking out of the foregoing group of 60 a group of 12 contained in it, viz. that corresponding to the positive substitutions of the four letters *abcd*, it is easy to see, that there is a transformation $(i, 0, 0, 1)$, that is, *s* into *is*, which can be taken for the substitution *abcd*, and also to complete thence the group of 24. And we have thus the following Table.

Groups of 12 and 24. Pole at Θ .

No.	$(Ax + B)$		$\div (Cx + D)$	substitution
1	1		1	1
2	-1		1	<i>ab.cd</i>
3	<i>i</i>		1	<i>ac.bd</i>
4	- <i>i</i>		1	<i>ad.bc</i>
5	<i>i</i>		1	<i>abc</i>
6	- <i>i</i>		1	<i>acb</i>
7	1		- <i>i</i>	<i>adc</i>
8	-1		- <i>i</i>	<i>abd</i>
9	<i>i</i>		1	<i>acd</i>
10	- <i>i</i>		1	<i>adb</i>
11	1		<i>i</i>	<i>bcd</i>
12	-1		<i>i</i>	<i>bdc</i>
13	<i>i</i>		-1	<i>adb</i>
14	- <i>i</i>		-1	<i>acbd</i>
15	1		-1	<i>ab.cd</i>
16	-1		-1	<i>aedb</i>
17	1		1	<i>bd</i>
18	-1		1	<i>abcd</i>
19	<i>i</i>		- <i>i</i>	<i>bc</i>
20	- <i>i</i>		- <i>i</i>	<i>abde</i>
21	-1		1	<i>ae</i>
22	1		1	<i>adeb</i>
23	- <i>i</i>		- <i>i</i>	<i>ad</i>
24	<i>i</i>		-1	—

108. The group of 60 was obtained in the *A*-form by Gordan in his paper. The passage from the Θ -form to the *A*-form is made as follows: let *X*, *Y*, *Z* be the coordinates of a point when the axes are as in the Θ -form, *X*₁, *Y*₁, *Z*₁ the coordinates of the same point when the axes are as in the *A*-form; we may write

$$X_1, Y_1, Z_1 = aX + bZ, Y, -bX + aZ,$$

where

$$a = \sqrt{\frac{5 - \sqrt{5}}{10}}, \quad b = \sqrt{\frac{5 + \sqrt{5}}{10}},$$

then, if the equations $X : Y : Z = L : M : N$ are those of an axis of rotation referred to the first set of coordinates and $X_1 : Y_1 : Z_1 = L_1 : M_1 : N_1$, those of the same axis referred to the second set of coordinates, we may write

$$L_1, M_1, N_1 = aL + bN, M, -bL + aN;$$

these values are such that

$$L_1^2 + M_1^2 + N_1^2 = L^2 + M^2 + N^2,$$

and hence, λ, μ, ν and λ_1, μ_1, ν_1 being the rotations, we have

$$L, M, N = \frac{1}{\theta}\lambda, \frac{1}{\theta}\mu, \frac{1}{\theta}\nu; \quad L_1, M_1, N_1 = \frac{1}{\theta}\lambda_1, \frac{1}{\theta}\mu_1, \frac{1}{\theta}\nu_1;$$

where θ has the same value in each set of equations.

From the equations

$$A : B : C : D = -\nu - i : \lambda + i\mu : \lambda - i\mu : \nu - i,$$

we may write

$$B + C : B - C : D - A : D + A = \lambda : i\mu : \nu : -i, = L : iM : N : -i\theta,$$

and similarly

$$B_1 + C_1 : B_1 - C_1 : D_1 - A_1 : D_1 + A_1 = L_1 : iM_1 : N_1 : -i\theta.$$

Hence we may write

$$B_1 + C_1 = a(B + C) + b(D - A),$$

$$B_1 - C_1 = B - C,$$

$$D_1 - A_1 = -b(B + C) + a(D - A),$$

$$D_1 + A_1 = D + A;$$

or say,

$$A_1 = -a(B + C) - b(D - A) + (D + A),$$

$$B_1 = a(B + C) + b(D - A) + (B - C),$$

$$C_1 = a(B + C) + b(D - A) - (B - C),$$

$$D_1 = -a(B + C) - b(D - A) + (D + A);$$

which are the values for a transformation (A_1, B_1, C_1, D_1) in the A -form: of course, as only the ratios are material, the values may be multiplied by any common factor.

109. The results are exhibited in terms of ϵ , an imaginary fifth root of unity: taking $\epsilon = \cos 72^\circ + i \sin 72^\circ$, we have

$$\epsilon + \epsilon^4 = \frac{\sqrt{5} - 1}{2}, \quad \epsilon^2 + \epsilon^3 = -\frac{\sqrt{5} + 1}{2},$$

$$a = \sqrt{\frac{5 + \sqrt{5}}{10}} = \pm i \frac{\sqrt{5 + \sqrt{5}}}{\sqrt{10}}, \quad b = \sqrt{\frac{5 - \sqrt{5}}{10}} = \pm \frac{\sqrt{5 - \sqrt{5}}}{\sqrt{10}},$$

where the upper signs belong to ϵ, ϵ^4 and the lower to ϵ^2, ϵ^3 . It may be remarked that

$$\frac{1}{a} = \sqrt{\frac{5 + \sqrt{5}}{2}}, \quad \frac{1}{b} = \sqrt{\frac{5 - \sqrt{5}}{2}}, \quad \frac{a}{b} = \sqrt{\frac{\sqrt{5} + 1}{\sqrt{5} - 1}}.$$

For instance, we have in the Θ -group $(A, B, C, D) = (-1, 0, 0, 1)$; $ab.cd$: and thence in the A -group $A_1, B_1, C_1, D_1 = (-2b, 2a, 2a, 2b)$; $ab.cd$: or say this is

$$(-1, a/b, a/b, 1) = (-1, \epsilon + \epsilon^4, \epsilon + \epsilon^4, 1);$$

which in the Table is given as $(-\epsilon^*, \epsilon^2 + \epsilon^3, \epsilon^2 + \epsilon^3, \epsilon^*)$; $ab.cd$.

By effecting the passage to the A -group in this manner, we of course obtain the proper substitution corresponding to each transformation: but I found it easier starting from two transformations and the corresponding substitutions, to obtain by thence successive compositions the entire group.

110. *Homographic Transformations.* The group of 60. Pole at A .

No.	Θ No.	$(As + B)$		$\div (Cs + D)$		substitution
1	1	1	0	0	1	1
2	4	-1	0	0	1	$ad.bc$
3	13	$-\epsilon^4$	0	0	1	$ac.be$
4	9	$-\epsilon^4$	0	0	1	$ab.de$
5	10	$-\epsilon^3$	0	0	1	$bd.ce$
6	14	$-\epsilon$	0	0	1	$ae.be$
7	6	$\epsilon + \epsilon^4$	ϵ^2	ϵ^3	$-(\epsilon + \epsilon^4)$	$ae.bd$
8	5	$\epsilon + \epsilon^4$	ϵ^3	ϵ^2	$-(\epsilon + \epsilon^4)$	$ad.be$
9	15	$\epsilon + \epsilon^4$	ϵ^4	ϵ	$-(\epsilon + \epsilon^4)$	$bc.de$
10	3	$\epsilon + \epsilon^4$	ϵ	ϵ^4	$-(\epsilon + \epsilon^4)$	$ac.de$
11	12	$\epsilon + \epsilon^4$	ϵ^3	ϵ^2	$-(\epsilon + \epsilon^4)$	$ac.bd$
12	16	-1	$\epsilon^4 + \epsilon^3$	$\epsilon^2 + \epsilon$	1	$ae.bd$
13	11	$-\epsilon$	$\epsilon^3 + 1$	$1 + \epsilon^2$	ϵ	$ab.ce$
14	7	$-\epsilon^2$	$1 + \epsilon^4$	$\epsilon + \epsilon^4$	ϵ^2	$bc.cd$
15	2	$-\epsilon^3$	$\epsilon^3 + \epsilon$	$\epsilon^3 + \epsilon^4$	ϵ^3	$ab.ce$
16	8	$-\epsilon^4$	$\epsilon^2 + \epsilon$	$\epsilon^3 + \epsilon^4$	ϵ^4	$ad.ce$
17	21	$\epsilon^2 + 1$	ϵ	1	$-(\epsilon + \epsilon^2)$	adb
18	36	$\epsilon^4 + 1$	ϵ^2	ϵ^4	$-(\epsilon + \epsilon^3)$	aeb
19	30	$\epsilon^2 + 1$	ϵ^4	ϵ^3	$-(\epsilon^2 + \epsilon^4)$	bee
20	34	$\epsilon^2 + 1$	ϵ^3	ϵ^2	$-(\epsilon + \epsilon^2)$	cde
21	19	$\epsilon^2 + 1$	1	ϵ	$-(\epsilon + \epsilon^2)$	adc
22	33	$\epsilon + \epsilon^4$	1	ϵ^2	$-(\epsilon + \epsilon^2)$	cde
23	20	$\epsilon + \epsilon^4$	ϵ^3	ϵ^4	$-(\epsilon + \epsilon^3)$	acd
24	22	$\epsilon + \epsilon^4$	ϵ^4	ϵ	$-(\epsilon + \epsilon^2)$	abd
25	36	$\epsilon + \epsilon^4$	1	ϵ^4	$-(\epsilon + \epsilon^3)$	abe
26	29	$\epsilon + \epsilon^4$	ϵ	ϵ	$-(\epsilon + \epsilon^3)$	bee
27	31	$-\epsilon$	$\epsilon^2 + \epsilon^4$	$\epsilon^2 + \epsilon^4$	1	aed
28	17	$-\epsilon^2$	$\epsilon^2 + \epsilon$	$\epsilon^2 + \epsilon$	ϵ	abc
29	27	$-\epsilon^2$	$\epsilon^3 + \epsilon^4$	$\epsilon^3 + \epsilon^4$	ϵ^2	bcd
30	26	$-\epsilon^4$	$\epsilon^3 + \epsilon^4$	$\epsilon^3 + \epsilon^4$	ϵ^3	aec
31	23	-1	$\epsilon^2 + \epsilon^4$	$\epsilon^3 + \epsilon^4$	1	bcd
32	24	$-\epsilon^4$	1	ϵ^2	ϵ^2	bdc
33	32	-1	$\epsilon^2 + \epsilon^3$	$\epsilon^4 + \epsilon$	ϵ^4	ade
34	18	$-\epsilon$	$\epsilon + \epsilon^4$	$\epsilon^2 + \epsilon^3$	ϵ^2	aeb
35	28	$-\epsilon^2$	$\epsilon + \epsilon^4$	$\epsilon^2 + \epsilon^3$	ϵ^4	bde
36	25	$-\epsilon^3$	$\epsilon^3 + \epsilon^4$	$\epsilon^3 + \epsilon^4$	ϵ^4	ace
37	44	ϵ	1	1	1	$abccd$
38	43	ϵ^4	1	1	1	$aedbc$
39	42	ϵ^3	1	1	1	$acbde$
40	41	ϵ^2	1	1	1	$adceb$
41	38	$\epsilon^3 + \epsilon^4$	1	1	$-(\epsilon + \epsilon^3)$	$acebd$
42	46	$\epsilon^3 + \epsilon^4$	ϵ	ϵ^4	$-(\epsilon + \epsilon^3)$	$abdcc$
43	58	$\epsilon^2 + \epsilon^4$	ϵ^2	ϵ^3	$-(\epsilon + \epsilon^3)$	$adcbe$
44	55	$\epsilon^2 + \epsilon^4$	ϵ^3	ϵ^2	$-(\epsilon + \epsilon^3)$	$adecb$
45	50	$\epsilon^3 + \epsilon^4$	ϵ^4	ϵ	$-(\epsilon + \epsilon^3)$	$acdeb$
46	51	$\epsilon^2 + \epsilon^4$	ϵ^2	1	$-(\epsilon + \epsilon^3)$	$abedc$
47	39	$1 + \epsilon^2$	ϵ^2	ϵ^4	$-(\epsilon + \epsilon^3)$	$adbcc$
48	47	$1 + \epsilon^2$	1	ϵ^3	$-(\epsilon + \epsilon^3)$	$accdb$
49	59	$1 + \epsilon^2$	ϵ	ϵ^2	$-(\epsilon + \epsilon^3)$	$acbcd$
50	54	$1 + \epsilon^2$	ϵ^3	ϵ	$-(\epsilon + \epsilon^3)$	$abced$
51	56	$-\epsilon^2$	$1 + \epsilon^4$	$\epsilon^2 + \epsilon^3$	ϵ^3	$abdec$
52	49	$-\epsilon^4$	$\epsilon + \epsilon^3$	$\epsilon^2 + \epsilon^4$	ϵ	$abdcc$
53	37	$-\epsilon^3$	$\epsilon^4 + \epsilon$	$\epsilon^2 + \epsilon^3$	ϵ^4	$aebcd$
54	45	-1	$\epsilon + \epsilon^2$	$\epsilon^3 + \epsilon^4$	ϵ^2	$adebc$
55	57	$-\epsilon$	$\epsilon + \epsilon^2$	$\epsilon^3 + \epsilon^4$	ϵ^4	$abdec$
56	48	$-\epsilon^3$	$\epsilon^4 + \epsilon$	$\epsilon^2 + \epsilon^3$	1	$adcbc$
57	60	$-\epsilon^2$	$\epsilon + \epsilon^3$	$\epsilon^2 + \epsilon^4$	ϵ	$acedb$
58	53	$\epsilon^3 + 1$	$\epsilon^2 + \epsilon^4$	ϵ^2	ϵ^3	$aebdc$
59	52	$-1 + \epsilon^4$	$\epsilon^2 + \epsilon^3$	ϵ^2	ϵ^3	$adbcc$
60	40	$-\epsilon^2$	$\epsilon + \epsilon^4$	ϵ^2	ϵ^2	$aedcb$

[PAPER 745 CONTINUES BEYOND P. 201 WITH ART. 111 SELECTING THE POSITIVE SUBSTITUTIONS $abcd$ AND COMPLETING THE GROUP OF 24, POLE AT A , ETC.]

[745] (CONTINUED)

ON THE SCHWARZIAN DERIVATIVE AND THE POLYHEDRAL FUNCTIONS.

111. Selecting the transformations which correspond to the positive substitutions $abcd$, and completing the group of 24 we have

Homographic Transformations. The groups of 12 and 24. Pole at A.

	$(A\xi + B) \div (C\xi + D)$				
1	1	0	0	1	1
2	0	-1	1	0	$ab.cd$
3	$\epsilon + \epsilon^4$	ϵ^4	ϵ^4	$-(\epsilon + \epsilon^4)$	$ac.bd$
4	$-\epsilon^3$	$\epsilon^2 + \epsilon^4$	$\epsilon^2 + \epsilon^4$	ϵ^3	$ab.cd$
5	$-\epsilon$	$\epsilon + \epsilon^2$	$\epsilon + \epsilon^2$	ϵ	acb
6	$-\epsilon$	ϵ	ϵ^2	$-(\epsilon + \epsilon^2)$	abc
7	$\epsilon + \epsilon^4$	ϵ^4	ϵ	$-(\epsilon + \epsilon^4)$	acd
8	$\epsilon^2 + 1$	1	1	$-(\epsilon + \epsilon^2)$	bcd
9	$\epsilon + \epsilon^2$	ϵ	ϵ	$-(\epsilon + \epsilon^4)$	bdc
10	$\epsilon^2 + 1$	$1 + \epsilon^4$	$1 + \epsilon^3$	-1	$abdc$
11	-1	$1 + \epsilon^2$	$1 + \epsilon^3$	-1	ab
12	$-\epsilon^2$	$1 + \epsilon^4$	$1 + \epsilon^2$	1	bd
13	1	$1 + 2\epsilon^4$	$1 + 2\epsilon$	-1	ac
14	$-\epsilon^2 + \epsilon^3$	$1 + \epsilon + 3\epsilon^4$	$-1 - 3\epsilon - \epsilon^4$	$\epsilon^2 - \epsilon^3$	cd
15	$\epsilon^2 - \epsilon^3$	$3 + \epsilon + \epsilon^2$	$-1 - 3\epsilon - \epsilon^2$	$-\epsilon^2 + \epsilon^3$	ac
16	$-1 + \epsilon^2$	$-1 - \epsilon^2 + 2\epsilon^4$	$1 + \epsilon^2 - 2\epsilon^4$	$1 - \epsilon^2$	bd
17	$2 + \epsilon^2 + 2\epsilon^4$	$-2 - 2\epsilon^3 - \epsilon^4$	$2\epsilon + \epsilon^2 + 2\epsilon^3$	$2\epsilon + 2\epsilon^3 + \epsilon^4$	ad
18	$2 + 2\epsilon^3 + \epsilon^4$	$2 + \epsilon^2 + 2\epsilon^3$	$-2\epsilon - 2\epsilon^2 - \epsilon^3$	$2\epsilon + \epsilon^2 + 2\epsilon^4$	bc
19	$-2 + \epsilon + \epsilon^4$	$-\epsilon + \epsilon^4$	$-\epsilon + \epsilon^4$	$\epsilon + \epsilon^4 - 2\epsilon^2$	$abcd$
20	1	-1	1	1	$abdc$
21	1	1	-1	1	$acdb$
22	$1 + \epsilon + 3\epsilon^2$	$\epsilon^3 - \epsilon^2$	$\epsilon^3 - \epsilon^2$	$1 + 3\epsilon + \epsilon^2$	$acbd$
23	$1 + 2\epsilon^2$	-1	-1	$-1 - 2\epsilon$	$adbc$
24	$3 + \epsilon + \epsilon^2$	$-\epsilon^3 + \epsilon^4$	$-\epsilon^2 + \epsilon^4$	$\epsilon + 3\epsilon + \epsilon^4$	$adcb$

As an example of the calculation we have $(A, B, C, D) = (0, i, -1, 0)$; ab . Hence

$$A_1, B_1, C_1, D_1 = (a(i-1), b(i-1) + i + 1, b(i-1) - (i+1), -a(i+1)),$$

$$= \left(1, \frac{b-i}{a}, \frac{b+i}{a}, -1\right).$$

The second and third coefficients are

$$\frac{\sqrt{5}+1}{2} - i\sqrt{\frac{5+\sqrt{5}}{2}}, \quad \frac{\sqrt{5}+1}{2} + i\sqrt{\frac{5+\sqrt{5}}{2}},$$

which, in virtue of the values of ϵ and ϵ^4 , are $1 + 2\epsilon^4$ and $1 + 2\epsilon$ respectively: or the result is as above $(1, 1 + 2\epsilon^4, 1 + 2\epsilon, -1)$.

112. In like manner for the passage from the Θ -form to the B -form, if X, Y, Z be the coordinates of a point on the spherical surface in regard to the Θ -axes, X_1, Y_1, Z_1 those of the same point in regard to the B -axes, we may write

$$X : Y : Z = X_1 : bY_1 + aZ_1 : -aY_1 + bZ_1,$$

where

$$a, b = \frac{\sqrt{5}-1}{2\sqrt{3}}, \frac{\sqrt{5}+1}{2\sqrt{3}}.$$

Hence $X : Y : Z = L : M : N$, being the equations of an axis of rotation in the first set of coordinates, those of the same axis in the second set of coordinates will be

$$X_1 : bY_1 + aZ_1 : -aY_1 + bZ_1 = L : M : N,$$

or calling these

$$= X_1 : Y_1 : Z_1 = L_1 : M_1 : N_1,$$

we have

$$L_1, M_1, N_1 = L, bM - aN, aM + bN;$$

these values are such that

$$L_1^2 + M_1^2 + N_1^2 = L^2 + M^2 + N^2,$$

or $\lambda, \mu, \nu, \lambda_1, \mu_1, \nu_1$ being the rotations, we have

$$L, M, N = \vartheta \lambda, \vartheta \mu, \vartheta \nu; \quad L_1, M_1, N_1 = \vartheta \lambda_1, \vartheta \mu_1, \vartheta \nu_1,$$

where ϑ has the same value in the two sets of equations. We have thus

$$B + C : B - C : D - A : D + A = L : 2M : N : -i\vartheta,$$

$$B_1 + C_1 : B_1 - C_1 : D_1 - A_1 : D_1 + A_1 = L_1 : 2M_1 : N_1 : -i\vartheta;$$

and hence

$$\begin{aligned} B_1 + C_1 &= B + C, \\ B_1 - C_1 &= b(B - C) - ai(D - A), \\ D_1 - A_1 &= -ai(B - C) + b(D - A), \\ D_1 + A_1 &= D + A; \end{aligned}$$

and thence

$$\begin{aligned} A_1 &= ai(B - C) - b(D - A) + (D + A), \\ B_1 &= b(B - C) - ai(D - A) + (B + C), \\ C_1 &= -b(B - C) + ai(D - A) + (B + C), \\ D_1 &= -ai(B - C) + b(D - A) + (D + A). \end{aligned}$$

113. As an example of the transformation, take

$$(A, B, C, D) = (2, -3 + \sqrt{5} + i(1 - \sqrt{5}), -3 + \sqrt{5} + i(-1 + \sqrt{5}), -2) \quad [bc.de];$$

then

$$B - C, B + C, D - A, D + A = i(1 - \sqrt{5}), -3 + \sqrt{5}, -2, 0;$$

and thence

$$\begin{aligned} A_1 &= \frac{1}{2\sqrt{3}} i(-6 + 2\sqrt{5}) + \frac{1}{2\sqrt{3}} (-2 + 2\sqrt{5}), \\ B_1 &= \frac{1}{2\sqrt{3}} (-4i) + \frac{1}{2\sqrt{3}} (2i(1 + \sqrt{5}) + (-3 + \sqrt{5})), \\ C_1 &= \frac{1}{2\sqrt{3}} (4i) + \frac{1}{2\sqrt{3}} (2i(1 - \sqrt{5}) + (-3 + \sqrt{5})), \\ D_1 &= \frac{1}{2\sqrt{3}} (-6 + 2\sqrt{5}) + \frac{1}{2\sqrt{3}} (-2 - 2\sqrt{5}); \end{aligned}$$

viz. multiplying by $2\sqrt{3}$, these are

$$8, i(-6 + 2\sqrt{5}) + 2\sqrt{3}(-3 + \sqrt{5}), i(6 - 2\sqrt{5}) + 2\sqrt{3}(-3 + \sqrt{5}), -8,$$

that is,

$$8, (-6 + 2\sqrt{5})(i + \sqrt{3}), (-6 + 2\sqrt{5})(-i + \sqrt{3}), -8,$$

or since

$$2 + \sqrt{3} = -2i\omega \quad \text{and} \quad -2 + \sqrt{3} = 2i\omega^2,$$

dividing by 4 these are

$$2, i(3 - \sqrt{5})\omega, i(-3 + \sqrt{5})\omega^2, -2,$$

as in the table.

114. *Homographic Transformations. The group of 60. Pole at B.*

$$\omega = \frac{1}{2}(-1 + i\sqrt{3}).$$

	$(A\xi + B) \div (C\xi + D)$				
1	1	0	0	1	1
2	1	$i(3 - \sqrt{5})$	$i(-3 + \sqrt{5})$	-2	$ac.bd$
3	0	ω	1	0	$ae.bd$
4	0	ω^2	1	0	$bd.ce$
5	2	$i(3 - \sqrt{5})$	$i(-3 + \sqrt{5})$	-2	$ab.cd$
6	2	$i(-3 - \sqrt{5})$	$i(3 + \sqrt{5})$	-2	$ad.bc$
7	2	$i(3 - \sqrt{5})\omega$	$i(-3 + \sqrt{5})\omega^2$	-2	$be.cd$
8	2	$i(-3 - \sqrt{5})\omega$	$i(3 + \sqrt{5})\omega^2$	-2	$ad.be$
9	2	$i(-3 - \sqrt{5})\omega^2$	$i(3 + \sqrt{5})\omega$	-2	$ab.ce$
10	2	$i(3 - \sqrt{5})\omega^2$	$i(-3 + \sqrt{5})\omega$	-2	$ae.bc$
11	2	$(-\sqrt{3} - i\sqrt{5})$	$(-\sqrt{3} + i\sqrt{5})\omega^2$	-2	$ac.de$
12	2	$(-\sqrt{3} - i\sqrt{5})\omega^2$	$-\sqrt{3} + i\sqrt{5}$	-2	$ad.ce$
13	2	$(\sqrt{3} - i\sqrt{5})\omega$	$(-\sqrt{3} + i\sqrt{5})\omega$	-2	$ae.cd$
14	2	$(\sqrt{3} - i\sqrt{5})\omega^2$	$\sqrt{3} + i\sqrt{5}$	-2	[unreadable]
15	2	$(\sqrt{3} - i\sqrt{5})$	$(\sqrt{3} + i\sqrt{5})\omega^2$	-2	[unreadable]
16	2	$(\sqrt{3} - i\sqrt{5})\omega$	$(\sqrt{3} + i\sqrt{5})\omega^2$	-2	[unreadable]
17	ω	0	0	1	ace
18	ω^2	0	0	1	aec
19	$\sqrt{3} - i\sqrt{5}$	2	-2	$\sqrt{3} + i\sqrt{5}$	bcd
20	$-\sqrt{3} - i\sqrt{5}$	$2\omega^2$	-2ω	$-\sqrt{3} + i\sqrt{5}$	bde
21	$-\sqrt{3} - i\sqrt{5}$	2ω	$-2\omega^2$	$-\sqrt{3} + i\sqrt{5}$	bdc
22	$\sqrt{3} - i\sqrt{5}$	2ω	$-2\omega^2$	$\sqrt{3} + i\sqrt{5}$	bcd
23	$\sqrt{3} - i\sqrt{5}$	$2\omega^2$	-2ω	$\sqrt{3} + i\sqrt{5}$	bdc
24	$\sqrt{3} - i\sqrt{5}$	$2\omega^2$	-2ω	$\sqrt{3} + i\sqrt{5}$	abd
25	2ω	$-\sqrt{3} - i\sqrt{5}$	$-\sqrt{3} + i\sqrt{5}$	$-2\omega^2$	adb
26	$-\sqrt{3} - i\sqrt{5}$	$-\sqrt{3} - i\sqrt{5}$	$-\sqrt{3} + i\sqrt{5}$	$\sqrt{3} + i\sqrt{5}$	abc
27	$-\sqrt{3} - i\sqrt{5}$	$-\sqrt{3} - i\sqrt{5}\omega^2$	$(-\sqrt{3} + i\sqrt{5})\omega$	$\sqrt{3} + i\sqrt{5}$	acb
28	$\sqrt{3} - i\sqrt{5}$	$-\sqrt{3} - i\sqrt{5}\omega$	$(-\sqrt{3} + i\sqrt{5})\omega^2$	$\sqrt{3} + i\sqrt{5}$	abe
29	$\sqrt{3} - i\sqrt{5}$	$-\sqrt{3} - i\sqrt{5}\omega^2$	$(-\sqrt{3} + i\sqrt{5})\omega$	$\sqrt{3} + i\sqrt{5}$	aeb
30	$2\omega^2$	$-\sqrt{3} - i\sqrt{5}$	$(-\sqrt{3} + i\sqrt{5})\omega^2$	-2ω	acd
31	ω	$\sqrt{3} - i\sqrt{5}$	$\sqrt{3} + i\sqrt{5}$	$-2\omega^2$	adc
32	2ω	$\sqrt{3} - i\sqrt{5}$	$\sqrt{3} + i\sqrt{5}$	$-2\omega^2$	ade
33	ω^2	$\sqrt{3} - i\sqrt{5}$	$\sqrt{3} + i\sqrt{5}$	-2	aed
34	$2\omega^2$	$\sqrt{3} - i\sqrt{5}$	$(\sqrt{3} + i\sqrt{5})\omega^2$	-2ω	bce
35	ω^2	$\sqrt{3} - i\sqrt{5}$	$(\sqrt{3} + i\sqrt{5})\omega^2$	$-2\omega^2$	bec
36	2	$\sqrt{3} - i\sqrt{5}$	$(-\sqrt{3} + i\sqrt{5})\omega$	$-2\omega^2$	cde
37	2ω	$-\sqrt{3} - i\sqrt{5}$	$(-\sqrt{3} + i\sqrt{5})\omega$	-2ω	ced
38	$2\omega^2$	$-\sqrt{3} - i\sqrt{5}$	$(-\sqrt{3} + i\sqrt{5})\omega$	-2	$adceb$
39	$2\omega^2$	0	$(\sqrt{3} + i\sqrt{5})\omega$	0	$adceb$
40	0	$\sqrt{3} - i\sqrt{5}$	$(\sqrt{3} + i\sqrt{5})\omega$	$-2\omega^2$	$acbde$
41	2	0	0	-2ω	$aedbc$
42	2ω	$\sqrt{3} - i\sqrt{5}$	$i(-3 + \sqrt{5})$	0	$abecd$
43	2ω	0	-2	-2	$aedcb$
44	0	$i(3 - \sqrt{5})\omega^2$	0	-2	$adbec$
45	2ω	0	-2ω	0	$acebd$
46	0	$+2\omega^2$	0	$-2\omega^2$	$abcde$
47	2	0	$i(-3 + \sqrt{5})\omega$	0	$adebc$
48	0	2	$i(-3 + \sqrt{5})$	$-2\omega^2$	$aecdb$
49	2	$i(3 - \sqrt{5})$	-2	0	$abdce$
50	0	0	0	$(-\sqrt{3} - i\sqrt{5})\omega^2$	$acbed$

	$(A\xi + B) \div (C\xi + D)$				
51	$-\sqrt{3} - i\sqrt{5}$	$i(3 - i\sqrt{5})\omega$	$-2\omega^2$	$(\sqrt{3} + i\sqrt{5})\omega$	<i>acdeb</i>
52	0	0	0	0	<i>adbce</i>
53	$\sqrt{3} - i\sqrt{5}$	2ω	$i(-3 + \sqrt{5})\omega^2$	-2ω	<i>aecbd</i>
54	0	0	$i(-3 + \sqrt{5})\omega^2$	-2ω	<i>abedc</i>
55	2	2	0	0	<i>aebcd</i>
56	2	$i(3 - \sqrt{5})$	$-2\omega^2$	$(-\sqrt{3} + i\sqrt{5})\omega$	<i>abdec</i>
57	$-\sqrt{3} - i\sqrt{5}$	0	-2ω	$(\sqrt{3} + i\sqrt{5})\omega^2$	<i>acedb</i>
58	0	$i(3 - \sqrt{5})\omega^2$	0	0	[unreadable]
59	$\sqrt{3} - i\sqrt{5}$	0	$i(-3 + \sqrt{5})\omega$	$-2\omega^2$	<i>adcbe</i>
60	0	$2\omega^2$	$i(3 + \sqrt{5})\omega$	-2ω	<i>adecb</i>

[The remainder of the table (lines 41–60 schematic) follows the same pattern as in the scan; entries marked [unreadable] correspond to faint or unclear OCR cells in the original.]

115. We hence derive

Homographic Transformations. The groups of 12 and 24. Pole at B.

	$(A\xi + B) \div (C\xi + D)$				
1	1	0	0	1	1
2	2	$i(3 - \sqrt{5})$	$i(-3 + \sqrt{5})$	-2	<i>ab.cd</i>
3	0	1	1	0	<i>ac.bd</i>
4	2	$i(-3 - \sqrt{5})$	$i(3 + \sqrt{5})$	-2	<i>ad.bc</i>
5	2ω	$-\sqrt{3} - i\sqrt{5}$	$\sqrt{3} + i\sqrt{5}$	$-2\omega^2$	<i>abc</i>
6	$2\omega^2$	$-\sqrt{3} - i\sqrt{5}$	$\sqrt{3} + i\sqrt{5}$	-2ω	<i>acb</i>
7	$-\sqrt{3} - i\sqrt{5}$	2ω	$-2\omega^2$	$-\sqrt{3} + i\sqrt{5}$	<i>abe</i>
8	$\sqrt{3} - i\sqrt{5}$	$\sqrt{3} - i\sqrt{5}$	$\sqrt{3} + i\sqrt{5}$	$\sqrt{3} + i\sqrt{5}$	<i>aeb</i>
9	ω^2	$\sqrt{3} - i\sqrt{5}$	-2ω	$-\sqrt{3} + i\sqrt{5}$	<i>abd</i>
10	ω	$2\omega^2$	-2ω	$-\sqrt{3} + i\sqrt{5}$	<i>adb</i>
11	$\sqrt{3} - i\sqrt{5}$	2ω			<i>acd</i>
12	$-\sqrt{3} - i\sqrt{5}$				<i>adc</i>
13	0	$\sqrt{3}(1 + \sqrt{5}) + i(-3 - \sqrt{5})$	$\sqrt{3}(1 + \sqrt{5}) + i(3 + \sqrt{5})$	-2	<i>ab</i>
14	0	$\sqrt{3}(-1 - \sqrt{5}) + i(-3 - \sqrt{5})$	$\sqrt{3}(-1 - \sqrt{5}) + i(3 + \sqrt{5})$	-2	<i>cd</i>
15	$\sqrt{5}$	$-i$	i	$-\sqrt{5}$	<i>ae</i>
16	$i\sqrt{5}$	$i\sqrt{5}$	$-i\sqrt{5}$	-1	<i>bd</i>
17	0	$\sqrt{3}(-1 + \sqrt{5}) + i(3 - \sqrt{5})$	$\sqrt{3}(-1 + \sqrt{5}) + i(-3 + \sqrt{5})$	-2	<i>ad</i>
18	0	$\sqrt{3}(1 - \sqrt{5}) + i(3 - \sqrt{5})$	$\sqrt{3}(1 - \sqrt{5}) + i(-3 + \sqrt{5})$	-2	<i>bc</i>
19	0	i	i	1	<i>abcd</i>
20	0	$-i$	$-i$	1	<i>adcb</i>
21	$\sqrt{3}(1 - \sqrt{5}) + i(3 + \sqrt{5})$	2	-2	$\sqrt{3}(1 - \sqrt{5}) + i(-3 + \sqrt{5})$	<i>abde</i>
22	$\sqrt{3}(1 + \sqrt{5}) + i(-3 + \sqrt{5})$	2	-2	$\sqrt{3}(1 + \sqrt{5}) + i(3 + \sqrt{5})$	<i>acbd</i>
23	$\sqrt{3}(-1 + \sqrt{5}) + i(3 - \sqrt{5})$	2	-2	$\sqrt{3}(-1 + \sqrt{5}) + i(-3 + \sqrt{5})$	<i>acdb</i>
24	$\sqrt{3}(-1 - \sqrt{5}) + i(-3 - \sqrt{5})$	2	-2	$\sqrt{3}(-1 - \sqrt{5}) + i(3 + \sqrt{5})$	<i>adbc</i>

116. I give also the group of 12, (*abce*), slightly modifying the form: viz. I write first $\sqrt{3} \pm \sqrt{5} = 2\sqrt{2}k$, and therefore $\sqrt{3} \mp \sqrt{5} = 2\sqrt{2}\frac{i}{k}$; then for x I write λx , and divide the A and B by λ : the A and B then contain $\frac{\lambda}{k}$, and the C and D contain $\frac{k}{\lambda}$; assuming $\frac{\lambda}{k} = \frac{k}{\lambda}$, we have $\lambda = k$. For instance, in the transformation corresponding to *abc*, the $Ax + B$ and $Cx + D$,

$$= 2\omega^2 x - (\sqrt{3} + i\sqrt{5}) \quad \text{and} \quad (-\sqrt{3} + i\sqrt{5})x - 2\omega,$$

become first $2\omega^2 x - 2\sqrt{2}k$, and $-2\sqrt{2}ix - 2\omega$, and then (omitting also the factor 2)

$$\omega^2 x - \sqrt{2}k \quad \text{and} \quad -\sqrt{2}\frac{i}{k}x - \omega,$$

viz. when $k = i$, they are $\omega^2 x - i\sqrt{2}$ and $-\omega x \cdot i\sqrt{2}$; that is, the values of A, B, C, D are $\omega^2, -i\sqrt{2}, i\sqrt{2}, -\omega$. The group is

<i>Group of 12. Pole at B.</i>	1	0	0	1	1
	ω	0	0	1	<i>ace</i>
	ω^2	0	0	1	<i>aec</i>
	ω	$-i\omega\sqrt{2}$	$i\omega\sqrt{2}$	$-\omega^2$	<i>abc</i>
	ω^2	$-i\omega^2\sqrt{2}$	$i\omega^2\sqrt{2}$	$-\omega$	<i>acb</i>
	1	$-i\omega\sqrt{2}$	$i\sqrt{2}$	$-\omega$	<i>abe</i>
	1	$-i\sqrt{2}$	$i\omega\sqrt{2}$	$-\omega^2$	<i>aeb</i>
	1	$-i\omega^2\sqrt{2}$	$i\sqrt{2}$	$-\omega^2$	<i>bce</i>
	1	$-i\sqrt{2}$	$i\omega^2\sqrt{2}$	$-\omega$	<i>bec</i>
	0	$-i\omega\sqrt{2}$	$i\omega^2\sqrt{2}$	0	<i>ab.ce</i>
					<i>ac.be</i>
	0	$-i\omega^2\sqrt{2}$	$i\omega\sqrt{2}$	0	<i>ae.bc</i>
	0	$-i\sqrt{2}$	$i\sqrt{2}$	0	

117. From the Table of the Groups of 12 and 24, Θ -form, it appears that the group of 12 is

$$1, \frac{1}{x}, -1, -\frac{1}{x}, \frac{i(x-1)}{x+1}, \frac{-i(x-1)}{x+1}, \frac{i(x+1)}{x-1}, \frac{-i(x+1)}{x-1},$$

$$\frac{x+i}{x-i}, \frac{x-i}{x+i}, \frac{-(x+i)}{-x-i}, \frac{-(x-i)}{x+i},$$

and if we proceed to form the product of the twelve factors $s - x$, $s + \frac{1}{x}$, $s + x$, &c., we have first the three products

$$s^4 + \alpha s^2 + 1; \quad s^4 + \beta s^2 + 1; \quad s^4 + \gamma s^2 + 1;$$

if for shortness

$$\alpha, \beta, \gamma = -\left(x^2 + \frac{1}{x^2}\right), \quad \frac{x^4 + 6x^2 + 1}{(x^2 - 1)^2}, \quad -\frac{x^4 - 6x^2 + 1}{(x^2 + 1)^2}.$$

The product of the three quartic functions is

$$= (s^4 + 1)^3 + (s^4 + 1)^2 s^2 (\alpha + \beta + \gamma) + (s^4 + 1) s^4 (\beta\gamma + \gamma\alpha + \alpha\beta) + s^6 \cdot \alpha\beta\gamma,$$

and we have

$$\alpha + \beta + \gamma = -\frac{32x^2(x^4 + 1)}{(x^4 - 1)^2}, \quad \beta\gamma + \gamma\alpha + \alpha\beta = -\frac{36x^4}{(x^4 - 1)^2} = -\frac{4(x^8 - 33x^6 - 33x^2 + 1)}{x^2(x^4 - 1)^2},$$

$$\alpha\beta\gamma = \frac{4(x^8 - 33x^6 - 33x^2 + 1)}{x^2(x^4 - 1)^2}.$$

Hence the product is found to be

$$= (x^8 - 33x^6 - 33x^2 + 1) \div x^2(x^4 - 1)^2, \quad x^8 - 33x^6 - 33x^2 + 1$$

which is

$$= s^{12}(x^4 - 1)^2 \left\{ \frac{s^8 - 33s^6 - 33s^2 + 1}{x^2(x^4 - 1)^2} - \frac{x^8 - 33x^6 - 33x^2 + 1}{x^2(x^4 - 1)^2} \right\}.$$

We thus verify that the twelve transformations x into x , into $\frac{1}{x}$, &c., give each of them a transformation of the function

$$\frac{x^8 - 33x^6 - 33x^2 + 1}{x^2(x^4 - 1)^2}$$

into itself.

118. It has been already remarked that we can from the coefficients (A, B, C, D) of the homographic transformation pass back to the position of the axis of rotation: viz. we have

$$A, B, C, D = -i, -i\mu, -i\nu, +i;$$

and thence

$$\lambda : \mu : \nu : 1 = B + C : -i(B - C) : B - A : i(D + A),$$

that is,

$$\lambda, \mu, \nu = -i(B + C), -(B - C), -i(D - A); \frac{1}{2}(D + A).$$

The equations of the axis thus are

$$\frac{x}{B + C} : \frac{iy}{B - C} : \frac{z}{D - A},$$

and the equations of the central plane at right angles to the axis are

$$-(B + C)x + i(B - C)y + (A - D)z = 0.$$

119. In particular, we may find the equations of the 15 planes at right angles to the Θ -axes: these are in fact the before-mentioned 15 planes, intersecting the sphere in great circles the projections of which are the circles in the three figures respectively. Taking the equation of the plane to be $Lx + My + Nz = 0$, it is at once seen that the equation of the projecting cone (vertex at the South pole) is

$$N(x^2 + y^2 + z^2 - 1) - 2(z + 1)(Lx + My + Nz) = 0,$$

and hence, writing $z = 0$, we find

$$N(x^2 + y^2 - 1) - 2(Lx + My) = 0$$

for the equation of the circle in the plane figure. We have thus the equations of a system of 15 circles related to each other in the manner before referred to.

120. Taking the Θ -form, the equations of the 15 planes are at once found: and we thence obtain the equations of the 15 circles: viz. writing for shortness

$$\Omega = x^2 + y^2 - 1,$$

the equations are

$z = 0,$	$(ab.cd)$	$\Omega = 0,$
$x = 0,$	$(ac.bd)$	$x = 0,$
$y = 0,$	$(ad.bc)$	$y = 0,$
$(3 - \sqrt{5})x + (1 - \sqrt{5})y + 2z = 0,$	$(ae.bc)$	$\Omega - [(3 - \sqrt{5})x + (1 - \sqrt{5})y] = 0,$
$(-1 - \sqrt{5})x + (-1 + \sqrt{5})y + 2z = 0,$	$(ab.ce)$	$\Omega - [(-1 - \sqrt{5})x + (-1 + \sqrt{5})y] = 0,$
$(1 + \sqrt{5})x + (3 + \sqrt{5})y + 2z = 0,$	$(ac.be)$	$\Omega - [(1 + \sqrt{5})x + (3 + \sqrt{5})y] = 0,$
$(-3 + \sqrt{5})x + (-1 + \sqrt{5})y + 2z = 0,$	$(ad.bc)$	
$(1 + \sqrt{5})x + (1 - \sqrt{5})y + 2z = 0,$	$(ab.de)$	
$(-1 - \sqrt{5})x + (-3 - \sqrt{5})y + 2z = 0,$	$(ae.bd)$	
$(-3 + \sqrt{5})x + (1 - \sqrt{5})y + 2z = 0,$	$(ad.ce)$	
$(1 + \sqrt{5})x + (-1 + \sqrt{5})y + 2z = 0,$	$(ae.cd)$	
$(-1 - \sqrt{5})x + (3 + \sqrt{5})y + 2z = 0,$	$(ac.de)$	
$(3 - \sqrt{5})x + (-1 + \sqrt{5})y + 2z = 0,$	$(bc.de)$	
$(-1 - \sqrt{5})x + (1 - \sqrt{5})y + 2z = 0,$	$(bc.cd)$	
$(1 + \sqrt{5})x + (-3 - \sqrt{5})y + 2z = 0,$	$(bd.ce)$	

and similarly for the other circles.

121. Observe that the arrangement is in sets of 3 planes, or circles, intersecting at right angles. One of the circles is the circle Ω , $x^2 + y^2 - 1 = 0$, corresponding to the equator, and two of them are the right lines $x = 0$ and $y = 0$. The equations of the remaining 12 circles may be written in the somewhat different form

$$\begin{aligned} \Omega + (\sqrt{5} - 1) [y - \tfrac{1}{2}(\sqrt{5} - 1)x] &= 0, \\ \Omega - (\sqrt{5} - 1) [y - \tfrac{1}{2}(\sqrt{5} + 3)x] &= 0, \\ \Omega - (\sqrt{5} + 3) [y + \tfrac{1}{2}(\sqrt{5} - 1)x] &= 0, \\ \Omega - (\sqrt{5} - 1) [y - \tfrac{1}{2}(\sqrt{5} - 1)x] &= 0, \\ \Omega + (\sqrt{5} - 1) [y - \tfrac{1}{2}(\sqrt{5} + 3)x] &= 0, \\ \Omega + (\sqrt{5} + 3) [y + \tfrac{1}{2}(\sqrt{5} - 1)x] &= 0, \\ \Omega + (\sqrt{5} - 1) [y + \tfrac{1}{2}(\sqrt{5} - 1)x] &= 0, \\ \Omega - (\sqrt{5} - 1) [y + \tfrac{1}{2}(\sqrt{5} + 3)x] &= 0, \\ \Omega - (\sqrt{5} + 3) [y - \tfrac{1}{2}(\sqrt{5} - 1)x] &= 0, \\ \Omega - (\sqrt{5} - 1) [y + \tfrac{1}{2}(\sqrt{5} - 1)x] &= 0, \\ \Omega + (\sqrt{5} - 1) [y + \tfrac{1}{2}(\sqrt{5} + 3)x] &= 0, \\ \Omega + (\sqrt{5} + 3) [y - \tfrac{1}{2}(\sqrt{5} - 1)x] &= 0. \end{aligned}$$

It hence appears that 4 and 4 circles have with $\Omega = 0$ the common chords $y + \tfrac{1}{2}(\sqrt{5} - 1)x = 0$, $y - \tfrac{1}{2}(\sqrt{5} - 1)x = 0$ respectively: and that 2 and 2 circles have with $\Omega = 0$ the common chords $y + \tfrac{1}{2}(\sqrt{5} + 3)x = 0$, $y - \tfrac{1}{2}(\sqrt{5} + 3)x = 0$ respectively.

122. The equations of the 12 circles are, in fact,

$$\begin{aligned} \Omega \pm (\sqrt{5} - 1) [y \pm \tfrac{1}{2}(\sqrt{5} - 1)x] &= 0, & \Omega \pm (\sqrt{5} + 3) [y \pm \tfrac{1}{2}(\sqrt{5} - 1)x] &= 0, \\ \Omega \pm (\sqrt{5} - 1) [y \pm \tfrac{1}{2}(\sqrt{5} + 3)x] &= 0; \end{aligned}$$

hence the radii are $= \sqrt{5} - 1$, 2 and $\sqrt{5} + 1$ respectively.

The construction of the 12 circles is as follows. Starting with a circle radius 1.

Lay down the diameters $y \pm \tfrac{1}{2}(\sqrt{5} - 1)x = 0$ (AA in the figure), and through the extremities of each describe 2 pairs of circles with the radii $\sqrt{5} - 1$, $\sqrt{5} + 1$ respectively.

Lay down the diameters $y \pm \tfrac{1}{2}(\sqrt{5} + 3)x = 0$ (BB in the figure), and through the extremities of each describe a pair of circles with the radius 2.

123. For the A -form, the equations of the fifteen planes are at once found to be

$$\begin{aligned}
 y &= 0, & ad \cdot bc \\
 -x + (\epsilon - \epsilon^4)z &= 0, & ac \cdot bd \\
 +(\epsilon - \epsilon^4)x + z &= 0, & ab \cdot cd \\
 (\epsilon - \epsilon^4)x - i(\epsilon^2 + \epsilon^3)y &= 0, & ab \cdot bc \\
 -(\epsilon^2 - \epsilon^3)x + i(\epsilon^2 + \epsilon^3)y + 2(\epsilon + \epsilon^4)z &= 0, & ab \cdot ce \\
 -x + i(\epsilon^2 + \epsilon^3 - \epsilon - \epsilon^4)y + 2z &= 0, & ae \cdot bc \\
 (\epsilon - \epsilon^4)x - i(\epsilon^2 + \epsilon^3)y &= 0, & ab \cdot de \\
 -(\epsilon + \epsilon^4)x + i(\epsilon^2 - \epsilon^3)y + 2(\epsilon - \epsilon^4)z &= 0, & ae \cdot bd \\
 +(\epsilon^2 + \epsilon^3 + 2)x + i(\epsilon^2 - \epsilon^3)y + 2(\epsilon - \epsilon^4)z &= 0, & ad \cdot bc \\
 -(\epsilon - \epsilon^4)x + i(\epsilon + \epsilon^4)y &= 0, & ae \cdot cd \\
 -(\epsilon + \epsilon^4)x - i(\epsilon^2 - \epsilon^3)y + 2(\epsilon + \epsilon^4)z &= 0, & ac \cdot de \\
 +(\epsilon^2 + \epsilon^3 + 2)x + i(\epsilon^2 - \epsilon^3)y + 2z &= 0, & ad \cdot ce \\
 -(\epsilon^2 - \epsilon^3)x + i(\epsilon^2 + \epsilon^3)y &= 0, & bd \cdot ce \\
 +x(\epsilon^2 + \epsilon^3) - i(\epsilon^2 - \epsilon^3)y + 2(\epsilon + \epsilon^4)z &= 0, & bc \cdot de \\
 -x - i(\epsilon^2 + \epsilon^3 - \epsilon - \epsilon^4)y + 2z &= 0, & bc \cdot cd,
 \end{aligned}$$

where, as before, the three planes of each set intersect at right angles.

124. Passing to the circles, the first plane of each set gives a right line, and we have thus five of the circles reducing themselves to right lines inclined to the axis of x at angles 0° , 36° , 72° , 108° and 144° respectively.

The remaining 10 circles form 5 pairs, the circles of a pair having different radii, but the two radii being the same for each pair, and so that for the several pairs the common chords with the circle $\Omega = 0$ are the diameters inclined to the axis of x at the angles 18° , 54° , 90° , 126° and 162° respectively. Considering the two circles for which the inclination is 90° , these arise from the planes $-x + (\epsilon - \epsilon^4)z = 0$, $(\epsilon + \epsilon^4)x + 2z = 0$, respectively. The equations of the circles thus are $(\epsilon + \epsilon^4)\Omega + 2x = 0$,

$\Omega - 2(\epsilon + \epsilon^4)x = 0$, or recollecting that $2(\epsilon + \epsilon^4) = \sqrt{5} - 1$ and therefore $\frac{2}{\epsilon + \epsilon^4} = \sqrt{5} + 1$, the equations are

$$x^2 + y^2 - (\sqrt{5} - 1)x - 1 = 0, \quad x^2 + y^2 + (\sqrt{5} + 1)x = 0;$$

hence for the first circle the x -coordinate of the centre is $\frac{1}{2}(\sqrt{5} - 1)$ and the radius is $= \frac{1}{2}\sqrt{10 - 2\sqrt{5}}$; for the second circle the x -coordinate of the centre is $= \frac{1}{2}(\sqrt{5} + 1)$, and the radius $= \frac{1}{2}\sqrt{10 + 2\sqrt{5}}$. We have thus the construction of these two circles, and consequently the construction of all the 12 circles.

125. For the B -form, after some easy reductions and attending to the relation $\omega - \omega^2 = i\sqrt{3}$, the

equations of the 15 planes become

$$\begin{aligned}
 x &= 0, & ac \cdot bd \\
 (-3 + \sqrt{5})y + 2z &= 0, & ab \cdot cd \\
 (3 + \sqrt{5})y + 2z &= 0, & ad \cdot bc \\
 \sqrt{3}x + \sqrt{5}y + 2z &= 0, & ac \cdot be \\
 -(1 + \sqrt{5})\sqrt{3}x + (3 - \sqrt{5})y + 4z &= 0, & ab \cdot ce \\
 (-1 + \sqrt{5})\sqrt{3}x + (-3 - \sqrt{5})y + 4z &= 0, & ae \cdot bc \\
 x + \sqrt{3}y &= 0, & ae \cdot bd \\
 -\sqrt{3}x + y + (3 + \sqrt{5})z &= 0, & ad \cdot bc \\
 \sqrt{3}x - y + (3 - \sqrt{5})z &= 0, & ab \cdot de \\
 -\sqrt{3}x + \sqrt{5}y + 2z &= 0, & ac \cdot de \\
 (1 - \sqrt{5})\sqrt{3}x + (-3 - \sqrt{5})y + 4z &= 0, & ad \cdot ce \\
 (1 + \sqrt{5})\sqrt{3}x + (3 - \sqrt{5})y + 4z &= 0, & ae \cdot cd \\
 x - \sqrt{3}y &= 0, & bd \cdot ce \\
 \sqrt{3}x + y + (3 + \sqrt{5})z &= 0, & bc \cdot de \\
 -\sqrt{3}x + y + (3 - \sqrt{5})z &= 0, & bc \cdot cd.
 \end{aligned}$$

126. Of the 15 circles, 3 are the lines $x - y\sqrt{3} = 0$, $x = 0$, $x + y\sqrt{3} = 0$, viz. these are lines at inclinations 30° , 90° , 150° to the axis of x . The equations of the remaining 12 circles are

$$\begin{aligned}
 \Omega + (3 - \sqrt{5})y &= 0, \\
 \Omega - (3 + \sqrt{5})y &= 0, \\
 (3 + \sqrt{5})\Omega - 2(y - x\sqrt{3}) &= 0, \\
 \Omega(3 - \sqrt{5}) + 2(y - x\sqrt{3}) &= 0, \\
 (3 + \sqrt{5})\Omega - 2(y + x\sqrt{3}) &= 0, \\
 (3 - \sqrt{5})\Omega + 2(y + x\sqrt{3}) &= 0,
 \end{aligned}$$

viz. these are pairs of circles having, for their common chords with $\Omega = 0$, the diameters at inclinations 0° , 60° , 120° respectively. And, lastly, we have the circles

$$\begin{aligned}
 2\Omega - [(-1 + \sqrt{5})\sqrt{3}x - (3 + \sqrt{5})y] &= 0, & 2\Omega + [(-1 + \sqrt{5})\sqrt{3}x + (3 + \sqrt{5})y] &= 0, \\
 \Omega - [-\sqrt{3}x + \sqrt{5}y] &= 0, & \Omega - [\sqrt{3}x + \sqrt{5}y] &= 0, \\
 2\Omega + [(1 + \sqrt{5})\sqrt{3}x - (3 - \sqrt{5})y] &= 0, & 2\Omega - [(1 + \sqrt{5})\sqrt{3}x + (3 - \sqrt{5})y] &= 0.
 \end{aligned}$$

127. The first three of these have, for common chords with $\Omega = 0$, the diameters whose equations are

$$(-1 + \sqrt{5})\sqrt{3}x - (3 + \sqrt{5})y = 0, \quad -\sqrt{3}x + \sqrt{5}y = 0, \quad (1 + \sqrt{5})\sqrt{3}x - (3 - \sqrt{5})y = 0 :$$

viz. these equations are $y = (-2 + \sqrt{5})x\sqrt{3}$, $y = \frac{\sqrt{3}}{\sqrt{5}}x$, $y = (2 + \sqrt{5})x\sqrt{3}$. If, as in a foregoing table, $\theta = 37^\circ 46'$, $\sin \theta = \sqrt{\frac{5}{45}}$, $\cos \theta = \sqrt{\frac{3}{45}}$, and therefore $\tan \theta = \sqrt{\frac{5}{3}}$; then the inclinations of these diameters to the axis of x are respectively $60^\circ - \theta$, θ and $120^\circ - \theta$, or say $30^\circ - (\theta - 30^\circ)$, $30^\circ + (\theta - 30^\circ)$ and $90^\circ - (\theta - 30^\circ)$, where $\theta - 30^\circ = 7^\circ 46'$, i.e. the inclinations are $30^\circ \pm 7^\circ 46'$ and $90^\circ - 7^\circ 46'$. And for the other three circles the common chords are the diameters at the same inclinations taken negatively. The geometrical construction of the fifteen circles for the B -case in question is thus not so simple as in the Θ - and A -cases.

The Regular Polyhedra as Solid figures. Art. Nos. 128 to 134.

128. I annex some results relating to the polyhedra considered as solid figures bounded by plane faces; or say results relating to the regular solids: s is in each case taken for the length of the edge of the solid.

	Tetrahedron	Cube	Octahedron	Dodecahedron	Icosahedron
Edge	s	s	s	s	s
Rad. of circum. sphere, R	$s \cdot \frac{\sqrt{3}}{2\sqrt{2}}$	$s \cdot \frac{\sqrt{3}}{2}$	$s \cdot \frac{1}{\sqrt{2}}$	$s \cdot \frac{\sqrt{3}(\sqrt{5}+1)}{4}$	$s\sqrt{\frac{5+\sqrt{5}}{8}}$
Rad. of inters. sphere, ρ	$s \cdot \frac{1}{2\sqrt{2}}$	$s \cdot \frac{1}{\sqrt{2}}$	$s \cdot \frac{1}{2}$	$s \cdot \frac{3+\sqrt{5}}{4}$	$s \cdot \frac{1+\sqrt{5}}{4}$
Rad. of inscribed sphere, r	$s \cdot \frac{1}{2\sqrt{2}\sqrt{3}}$	$s \cdot \frac{1}{2}$	$s \cdot \frac{1}{\sqrt{6}}$	$s \cdot \sqrt{\frac{5+2\sqrt{5}}{20}}$	$s \cdot \frac{3+\sqrt{5}}{4\sqrt{3}}$
Rad. of circle circum. to face, R'	$s \cdot \frac{1}{\sqrt{3}}$	$s \cdot \frac{1}{\sqrt{2}}$	$s \cdot \frac{1}{\sqrt{3}}$	$s \cdot \sqrt{\frac{5+\sqrt{5}}{10}}$	$s \cdot \frac{1}{\sqrt{3}}$
Rad. of circle inscribed to face, r'	$s \cdot \frac{1}{2\sqrt{3}}$	$s \cdot \frac{1}{2}$	$s \cdot \frac{1}{2\sqrt{3}}$	$s \cdot \sqrt{\frac{5+2\sqrt{5}}{20}}$	$s \cdot \frac{1}{2\sqrt{3}}$
Incl. of adjacent faces	$\cos^{-1} \frac{1}{3} = 70^\circ 28'$	90°	$\cos^{-1} -\frac{1}{3} = 109^\circ 32'$		
Incl. of edge to adjacent face	$\cos^{-1} \frac{1}{\sqrt{3}} = 54^\circ 46'$	90°	$= \cos^{-1} -\frac{1}{\sqrt{3}} = 125^\circ 44'$		

But we require further data in the cases of the dodecahedron and the icosahedron respectively.


129. For the dodecahedron, taking the edge to be $= s$ as before, then in the pentagonal face

$$\begin{aligned} \text{diagonal, } g \text{ is } &= s \cdot \frac{1}{2}(\sqrt{5} + 1), \\ \text{altitude, } k, &= s \cdot \frac{1}{2}\sqrt{5 + 2\sqrt{5}}, \\ \text{segments of do., } e, &= s \cdot \frac{1}{4}\sqrt{10 - 2\sqrt{5}}, \\ f, &= s \cdot \frac{1}{4}\sqrt{10 + 2\sqrt{5}}, \end{aligned}$$

where

$$k = e + f = R' + r'.$$

130. The section through a pair of opposite edges is a hexagon, as shown in the figure, viz. this is constructed by taking the four equal distances $O\Theta_i = \rho$, $= s \cdot \frac{1}{4}(3 + \sqrt{5})$, meeting at right angles in O ; then drawing the double ordinates BB , each $= s$, through Θ_1 and Θ_3 respectively, and joining their extremities with Θ_2 and Θ_4 : the sides Θ_2B and Θ_4B are then each $= k$, $= s \cdot \frac{1}{2}\sqrt{5 + 2\sqrt{5}}$; and inserting upon them the points A , Φ from the figure of the pentagon, we have several geometrical relations; viz. the line AA cuts the parallel sides $B\Theta_2$, $B\Theta_4$ at right angles, and when produced passes through the intersection of $B\Theta_1$ and $B\Theta_3$: we have OA , OB , $O\Theta = r$, R , ρ respectively: the four points Φ form a square, the side of which is g , $= s \cdot \frac{1}{2}(\sqrt{5} + 1)$.




131. We find also

$$\begin{aligned} AM &= s\sqrt{\frac{5+\sqrt{5}}{10}}, \\ AQ &= s\sqrt{\frac{3+2\sqrt{3}}{5}}, \\ QM &= s\sqrt{\frac{25+11\sqrt{5}}{5}} = r \cdot 2\sqrt{2}, \\ OQ &= s\sqrt{\frac{25+11\sqrt{5}}{8}} = r\sqrt{5}, \\ OM &= s\sqrt{\frac{5-\sqrt{5}}{40}}, \\ MB &= s\sqrt{\frac{2(5+2\sqrt{5})}{8}}. \end{aligned}$$

It may be remarked that in the figure $B\Theta_2$, $B\Theta_4$ are the projections of pentagonal faces, at right angles to the plane of the paper, having their centres at the points A , A , and the perpendicular distance between them = AA : the points Q , Q (only one of them shown in the figure) determine the directions of the $5 + 5$ sides which abut on these pentagonal faces respectively; and the $5 + 5$ points B which are the other extremities of these sides respectively form two pentagons, centres M , M , in the planes $MB MB$ respectively: the remaining 10 sides of the dodecahedron are the skew decagon obtained by joining in order these 10 points B . We have thus the means of making the perspective delineation of the dodecahedron.

132. The dodecahedron is built up from the cube, by placing on each face a figure of two triangular and two quadrangular faces, the orthogonal projection of which on the face of the cube is as in the figure: the side of the square is g , $= s \cdot \frac{1}{2}(\sqrt{5} + 1)$: the slope-breadths of the triangular faces are e , $= s \cdot \frac{1}{4}\sqrt{10 - 2\sqrt{5}}$, and those of the quadrangular faces are f , $= s \cdot \frac{1}{4}\sqrt{10 + 2\sqrt{5}}$; the lines represented by the other lines of the figure are in actual length each $= s$. We have thus a



section which is an isosceles triangle, base $= g$, other sides each $= f$; and the square of the altitude is thus $= f^2 - \frac{1}{4}g^2 = \frac{1}{4}s^2$, or the altitude $= \frac{1}{2}s$; viz. the altitude of the ridge-line BB , above the face of the cube is $= \frac{1}{2}s$, the half-side of the dodecahedron. We have in this result the most simple means of forming the perspective delineation of the dodecahedron.


133. For the icosahedron the section through two opposite edges is a hexagon, as shown in the figure (p. 216): to construct it, we take the four distances $O\Theta$ each $= \rho$, $= s \cdot \frac{1}{4}(1 + \sqrt{5})$ meeting at right angles; and then the distances $A\Theta_2$, $A\Theta_4$ each $= \frac{1}{2}s$; and complete the hexagon. This gives the sides $A\Theta_1$, $A\Theta_3$ each $= s \cdot \frac{1}{2}\sqrt{3}$, the altitude of the triangular face, side $= s$; and then, taking Θ_2B one-third of this, $= \frac{s\sqrt{3}}{6}$, we have OB at right angles to $A\Theta_2$, and OA , OB , $O\Theta = R$, r , ρ respectively.

Moreover, joining $A\Theta_3$ and OA_1 , we have these lines cutting at right angles in a point M : we find

$$\begin{aligned} A_1\Theta_3 &= s \cdot \frac{1}{2}\sqrt{5 + 2\sqrt{5}}, \\ M\Theta_3 &= s\sqrt{\frac{5+2\sqrt{5}}{20}}, \\ A_1M &= s\sqrt{\frac{5+\sqrt{5}}{10}}, \\ OM &= s\sqrt{\frac{3+\sqrt{5}}{40}} = \frac{1}{2}A_1M, \\ A_1M &= s\sqrt{\frac{5-\sqrt{5}}{10}}. \end{aligned}$$

134. It may be remarked that $A_1\Theta_3$, $A_3\Theta_1$ are the projections of two pentagons in planes perpendicular to that of the paper, their centres being M , M' : producing OM , OM' to the points A_1 , A_3 respectively, we have a pentagonal pyramid, summit A_1 , standing on the first pentagon, and an opposite pyramid, summit A_3 , standing on

the other pentagon: the $5 + 5$ triangular faces of the two pyramids are ten of the faces of the icosahedron, and the remaining ten faces are the triangles each having for its base a side of the one pentagon, and for its vertex a summit of the other pentagon, viz. the sides are the sides of the skew decagon obtained by joining in order the angular points of the two pentagons. We have thus a convenient method of forming the perspective delineation of the icosahedron.



746.

HIGHER PLANE CURVES.

[From Salmon's *Higher Plane Curves*, (3rd ed., 1879); see the Preface.]

One chapter and a large number of articles, in the second edition of Salmon's *Higher Plane Curves*, are due to Professor Cayley. Full reference to these is given by Dr Salmon in the preface.

747.

NOTE ON THE DEGENERATE FORMS OF CURVES.

[From Salmon's *Higher Plane Curves*, (3rd ed., 1879), pp. 383–385.]

Some remarks may be added as to the analytical theory of the degenerate forms of curves. As regards conics, a line-pair can be represented in point-coordinates by an equation of the form $xy = 0$; and reciprocally a point-pair can be represented in line-coordinates by an equation $\xi\eta = 0$, but we have to consider how the point-pair can be represented in point-coordinates: an equation $x^2 = 0$ is no adequate representation of the point-pair, but merely represents (as a two-fold or twice repeated line) the line joining the two points of the point-pair, all traces of the points themselves being lost in this representation: and it is to be noticed, that the conic, or two-fold line $x^2 = 0$, or say $(\alpha x + \beta y + \gamma z)^2 = 0$ is a conic which, analytically, and (in an improper sense) geometrically, satisfies the condition of touching any line whatever; whereas the only proper tangents of a point-pair are the lines which pass through one or other of the two points of the point-pair.

The solution arises out of the notion of a point-pair, considered as the limit of a conic, or say as an indefinitely flat conic; we have to consider conics certain of the coefficients whereof are infinitesimals, and which, when the infinitesimal coefficients actually vanish, reduce themselves to two-fold lines; and it is, moreover, necessary to consider the evanescent coefficients as infinitesimals of different orders. Thus consider the conics which pass through two given points, and touch two given lines (four conditions); take

$y = 0$, $z = 0$ for the given lines, $x = 0$ for the line joining the given points, and $(x = 0, y - \alpha z = 0)$, $(x = 0, y - \beta z = 0)$ for the given points; the equation of a conic satisfying the required conditions and containing one arbitrary parameter θ , is

$$x^2 + 2\theta xy + 2\theta\sqrt{(\alpha\beta)}xz + \theta^2(y - \alpha z)(y - \beta z) = 0;$$

or, what is the same thing,

$$\{x + \theta y + \theta\sqrt{(\alpha\beta)}z\}^2 - \theta^2(\alpha + \beta)yz = 0;$$

and this equation, considering therein θ as an infinitesimal, say of the first order, represents the flat conic or point-pair composed of the two given points. Comparing with the general equation

$$(a, b, c, f, g, h \mid x, y, z)^2 = 0,$$

we have

$$a = 1, \quad b = \theta^2, \quad c = \theta^2\alpha\beta, \quad f = -\frac{1}{2}\theta^2(\alpha + \beta), \quad g = \frac{1}{2}\theta\sqrt{(\alpha\beta)}, \quad h = \theta,$$

viz. a being taken to be finite, we have g and h infinitesimals of the first order, b, c, f infinitesimals of the second order; and the four ratios $g : \sqrt{(b)} : \sqrt{(c)} : \sqrt{(f)} : h$ are so determined as to satisfy the prescribed conditions.

Observe that the flat conic, considered as a conic passing through the two given points and touching the two given lines, is represented by a determinate equation, viz. considering the condition imposed upon ($b = \text{infinitesimal}$) as a determination of θ , the equation is a completely determinate one; but considering the flat conic merely as a conic passing through the two given points, the equation would contain two arbitrary parameters, determinable if the flat conic was subjected to the condition of touching two given lines, or to any other two conditions.

Generally, we may consider the equation of a curve of the order n ; such equation containing certain infinitesimal coefficients and, when these vanish, reducing itself to a composite equation $= P^\alpha Q^\beta \dots$; the equation in its original form represents a curve which may be called the penultimate curve. Consider the tangents from an arbitrary point to the penultimate curve; when this breaks up, the system of tangents reduces itself to (1) the tangents from the fixed point to the several component curves $P = 0$, $Q = 0$, &c. respectively; (2) the lines through the singular points of these same curves respectively; (3) the lines through the points of intersection of each two of the component curves; these points, each reckoned a proper number of times, are called “fixed summits”; (4) the lines from the fixed point to certain determinate points called “free summits” on the several component curves $P = 0$, $Q = 0$, &c. respectively. We have thus a degenerate form of the n -thic curve, which may be regarded as consisting of the component curves, each its proper number of times, and of the foregoing points called summits, and is consequently only inadequately represented by the ultimate equation $= P^\alpha Q^\beta \dots$; the number and distribution of the summits is not arbitrary, but is regulated by laws arising from the consideration of the penultimate curve, and there are of course for any given value of n various forms of degenerate curve, according to the different ultimate forms $= P^\alpha Q^\beta \dots = 0$, and to the number and distribution of the summits on the different component curves. The case of a quartic curve having the ultimate form x^2y^2 has been considered by Cayley, *Comptes Rendus*, t. LXXIV. p. 708 (March, 1872), [515], who states his conclusion as follows:

“there exists a quartic curve the penultimate of $= x^2y^2 = 0$, with nine free summits, three of them on one of the lines (say the line $y = 0$), and which are three of the intersections of the quartic by this line (the fourth intersection being indefinitely near to the point $x = 0, y = 0$), six situate at pleasure on the other line $x = 0$; and three fixed summits at the intersection of the two lines.” Other forms have been considered by Dr Zeuthen, *Comptes Rendus*, t. LXXV. pp. 703 and 950 (September and October, 1872), and some other forms by Zeuthen; the whole question of the degenerate forms of curves is one well deserving further investigation.

The question of the number of cubic curves satisfying given elementary conditions (depending as it does on the consideration of the degenerate forms of these curves) has been solved by Maillard and Zeuthen; that of the number of quartic curves has been solved by Dr Zeuthen.

748.

ON THE BITANGENTS OF A QUARTIC.

[From Salmon's *Higher Plane Curves*, (3rd ed., 1879), pp. 387–389.]

The equations of the 28 bitangents of a quartic curve were obtained in a very elegant form by Riemann in the paper “Zur Theorie der Abel’schen Functionen für den Fall $p = 3$,” *Ges. Werke*, Leipzig, 1876, pp. 456–472; and see also Weber’s *Theorie der Abel’schen Functionen vom Geschlecht 3*, Berlin, 1876. Riemann connects the several bitangents with the characteristics of the 28 odd functions, thus obtaining for them an algorithm which it is worth while to explain, but they will be given also with the algorithm employed p. 231 *et seq.* of the present work*, which is in fact the more simple one. The characteristic of a triple ϑ -function is a symbol of the form

$$\begin{array}{ccc} \alpha & \beta & \gamma \\ \alpha' & \beta' & \gamma' \end{array}$$

where each of the letters is = 0 or 1; there are thus in all 64 such symbols, but they are considered as odd or even according as the sum $\alpha\alpha' + \beta\beta' + \gamma\gamma'$ is odd or even; and the numbers of the odd and even characteristics are 28 and 36 respectively; and, as already mentioned, the 28 odd characteristics correspond to the 28 bitangents respectively.

We have x, y, z trilinear coordinates, $\alpha, \beta, \gamma, \alpha', \beta', \gamma'$ constants chosen at pleasure, and then $\alpha'', \beta'', \gamma''$ determinate constants, such that the equations

$$\begin{aligned} \alpha x + \beta y + \gamma z + \frac{\xi}{\alpha} + \frac{\eta}{\beta} + \frac{\zeta}{\gamma} &= 0, \\ \alpha' x + \beta' y + \gamma' z + \frac{\xi}{\alpha'} + \frac{\eta}{\beta'} + \frac{\zeta}{\gamma'} &= 0, \\ \alpha'' x + \beta'' y + \gamma'' z + \frac{\xi}{\alpha''} + \frac{\eta}{\beta''} + \frac{\zeta}{\gamma''} &= 0, \end{aligned}$$

[* That is, Salmon’s *Higher Plane Curves*.]

are equivalent to three independent equations; this being so, they determine ξ, η, ζ , each of them as a linear function of (x, y, z) ; and the equations of the bitangents of the curve $\sqrt{(x\xi)} + \sqrt{(y\eta)} + \sqrt{(z\zeta)} = 0$ (see Weber, p. 100) are

18	$\frac{111}{111}$	$x = 0,$
28	$\frac{001}{011}$	$y = 0,$
38	$\frac{011}{001}$	$z = 0,$
23	$\frac{010}{010}$	$\xi = 0,$
13	$\frac{100}{110}$	$\eta = 0,$
12	$\frac{110}{100}$	$\zeta = 0,$
48	$\frac{101}{100}$	$x + y + z = 0,$
14	$\frac{010}{011}$	$\xi + y + z = 0,$
58	$\frac{100}{101}$	$\alpha x + \beta y + \gamma z = 0,$
15	$\frac{011}{010}$	$\frac{\xi}{\alpha} + \beta y + \gamma z = 0,$
68	$\frac{110}{010}$	
16	$\frac{001}{101}$	
78	$\frac{010}{110}$	$\alpha''x + \beta''y + \gamma''z = 0,$
17	$\frac{101}{001}$	$\frac{\xi}{\alpha} + \beta''y + \gamma''z = 0,$
24	$\frac{100}{111}$	$x + \eta + z = 0,$
34	$\frac{110}{101}$	$x + y + \zeta = 0,$
25	$\frac{101}{110}$	$\alpha x + \frac{\eta}{\beta} + \gamma z = 0,$
35	$\frac{111}{100}$	$\alpha x + \beta y + \frac{\zeta}{\gamma} = 0,$
26	$\frac{111}{001}$	$\alpha'x + \frac{\eta}{\beta'} + \gamma'z = 0,$
36	$\frac{101}{011}$	$\alpha'x + \beta'y + \frac{\zeta}{\gamma'} = 0,$
27	$\frac{011}{101}$	$\alpha''x + \frac{\eta}{\beta''} + \gamma''z = 0,$
37	$\frac{001}{111}$	$\alpha''x + \beta''y + \frac{\zeta}{\gamma''} = 0,$
67	$\frac{100}{100}$	$\frac{x}{1 - \beta\gamma} + \frac{y}{1 - \gamma\alpha} + \frac{z}{1 - \alpha\beta} = 0,$
57	$\frac{110}{011}$	$\frac{x}{1 - \beta'\gamma'} + \frac{y}{1 - \gamma'\alpha'} + \frac{z}{1 - \alpha'\beta'} = 0,$
56	$\frac{010}{111}$	$\frac{x}{1 - \beta''\gamma''} + \frac{y}{1 - \gamma''\alpha''} + \frac{z}{1 - \alpha''\beta''} = 0,$
45	$\frac{001}{001}$	$\frac{\xi}{\alpha(1 - \beta\gamma)} + \frac{\eta}{\beta(1 - \gamma\alpha)} + \frac{\zeta}{\gamma(1 - \alpha\beta)} = 0,$
46	$\frac{011}{110}$	$\frac{\xi}{\alpha'(1 - \beta'\gamma')} + \frac{\eta}{\beta'(1 - \gamma'\alpha')} + \frac{\zeta}{\gamma'(1 - \alpha'\beta')} = 0,$
47	$\frac{111}{010}$	$\frac{\xi}{\alpha''(1 - \beta''\gamma'')} + \frac{\eta}{\beta''(1 - \gamma''\alpha'')} + \frac{\zeta}{\gamma''(1 - \alpha''\beta'')} = 0.$

The whole number of ways in which the equation of the curve can be expressed in a form such as $\sqrt{(x\xi)} + \sqrt{(y\eta)} + \sqrt{(z\zeta)} = 0$ is 1260; viz. the three pairs of bitangents entering into the equation of the curve are of one of the types

12.34,	13.24,	14.23	No. is	70
12.34,	13.24,	56.78	„	630
13.23,	14.24,	15.25	„	560
				1260.

It may be remarked that, selecting at pleasure any two pairs out of a system of three pairs, the type is always $\square$ or $|||$, viz. (see p. 233) the four bitangents are such that their points of contact are situate on a conic.

A considerable number of articles in the third edition of Salmon's *Treatise* are due to Professor Cayley. Full reference to these is given by Dr Salmon in the preface.

750.

ON THE THEORY OF RECIPROCAL SURFACES.

[From Salmon's *Treatise* on the analytic geometry of three dimensions, (3rd ed., 1874), pp. 539–550.]

600. In further developing the theory of reciprocal surfaces it has been found necessary to take account of other singularities, some of which are as yet only imperfectly understood. It will be convenient to give the following complete list of the quantities which present themselves:

- n , order of the surface.
- a , order of the tangent cone drawn from any point to the surface.
- h , number of nodal edges of the cone.
- κ , number of its cuspidal edges.
- ρ , class of nodal torse.
- σ , class of cuspidal torse.
- b , order of nodal curve.
- k , number of its apparent double points.
- f , number of its actual double points.
- t , number of its triple points.
- j , number of its pinch-points.
- q , its class.
- c , order of cuspidal curve.
- h , number of its apparent double points.
- θ , number of its points of an unexplained singularity.
- χ , number of its close-points.

- ω , number of its off-points.
- r , its class.
- β , number of intersections of nodal and cuspidal curves, stationary points on cuspidal curve.
- γ , number of intersections, stationary points on nodal curve.
- i , number of intersections, not stationary points on either curve.
- C , number of cnicnodes of surface.
- B , number of binodes.

And corresponding reciprocally to these:

- n' , class of surface.
- a' , class of section by arbitrary plane.
- δ' , number of double tangents of section.

κ' , number of its inflexions.

ρ' , order of node-couple curve.

σ' , order of spinode curve.

b' , class of node-couple torse.

k' , number of its apparent double planes.

f' , number of its actual double planes.

t' , number of its triple planes.

j' , number of its pinch-planes.

q' , its order.

c' , class of spinode torse.

h' , number of its apparent double planes.

θ' , number of its planes of a certain unexplained singularity.

χ' , number of its close-planes.

ω' , number of its off-planes.

r' , its order.

β' , number of common planes of node-couple and spinode torse, stationary planes of spinode torse.

γ' , number of common planes, stationary planes of node-couple torse.

i' , number of common planes, not stationary planes of either torse.

C' , number of cnictropes of surface.

B' , number of its bitropes.

In all, these are 46 quantities.

[PAPER 750 CONTINUES BEYOND P. 226 WITH ART. 601 GIVING THE RELATIONS AMONG THE LISTED QUANTITIES; SEE THE NEXT SLICE.]

usepackage[a4paper,margin=0.65in]geometry

[750] (CONTINUED)

ON THE THEORY OF RECIPROCAL SURFACES.

601. In part explanation, observe that the definitions of ρ and σ agree with those already given. The nodal torse is the torse enveloped by the tangent planes along the nodal curve; if the nodal curve meets the curve of contact a , then a tangent plane of the nodal torse passes through the arbitrary point, that is, ρ will be the number of these planes which pass through the arbitrary point, viz. the class of the torse. So also the cuspidal torse is the torse enveloped by the tangent planes along the cuspidal curve; and σ will be the number of these tangent planes which pass through the arbitrary point, viz. it will be the class of the torse. Again, as regards ρ' and σ' : the node-couple torse is the envelope of the bitangent planes of the surface, and the node-couple curve is the locus of the points of contact of these planes. Similarly, the spinode torse is the envelope of the parabolic planes of the surface, and the spinode curve is the locus of the points of contact of these planes, viz. it is the curve UH of intersection of the surface and its Hessian; the two curves are the reciprocals of the nodal and the cuspidal torses respectively, and the definitions of ρ' , σ' correspond to those of ρ and σ .

602. In regard to the nodal curve b , we consider k the number of its apparent double points (excluding actual double points); f the number of its actual double points (each of these is a point of contact of two sheets of the surface, and there is thus at the point a single tangent plane, viz. this is a plane f' , and we thus have $f' = f$); t the number of its triple points; and j the number of its pinch-points: these last are not singular points of the nodal curve *per se*, but are singular in regard to the curve as nodal curve of the surface, viz. a pinch-point is a point at which the two tangent planes are coincident. The curve is considered as not having any stationary points other than the points γ , which lie also on the cuspidal curve; and the expression for the class consequently is $q = b^2 - b - 2k - 2f - 3\gamma - 6t$.

603. In regard to the cuspidal curve c , we consider h the number of its apparent double points; and upon the curve, not singular points in regard to the curve *per se*, but only in regard to it as cuspidal curve of the surface, certain points in number β , χ , ω respectively. The curve is considered as not having any actual double or other multiple points, and as not having any stationary points except the points β , which lie also on the nodal curve; and the expression for the class consequently is

$$r = c^2 - c - 2h - 3\beta.$$

604. The γ are points where the cuspidal curve with the two sheets (or say rather half-sheets) belonging to it are intersected by another sheet of the surface; the curve of intersection with such other sheet, belonging to the nodal curve of the surface, has evidently a stationary (cuspidal) point at the point of intersection.

As to the points β , to facilitate the conception, imagine the cuspidal curve to be a semi-cubical parabola, and the nodal curve a right line (not in the plane of the curve) passing through the cusp; then intersecting the two curves by a series of parallel planes, any plane which is, say, above the cusp, meets the parabola in two real points and the line in one real point, and the

of the singularities θ and ω . So, in the original formulae for $a(n-2)(n-3)$, $b(n-2)(n-3)$, $c(n-2)(n-3)$, we have instead of δ to write $\delta - C - 3\omega$, and to substitute new expressions for $[ab]$, $[ac]$, $[bc]$; viz. these are

$$[ab] = ab - 2\rho - j,$$

$$[ac] = ac - 3\sigma - \chi - \omega,$$

$$[bc] = bc - 3\beta - 2\gamma - i.$$

The whole series of equations thus is

- (1) $a' = a.$
- (2) $f' = f.$
- (3) $i' = i.$
- (4) $\alpha = n(n-1) - 2b - 3c.$
- (5) $\kappa = 3n(n-2) - 6b - 8c.$
- (6) $\delta = \frac{1}{2}n(n-2)(n^2-9) - (n^2-n-6)(2b+3c) + 2b(b-1) + 6bc + \frac{9}{2}c(c-1).$
- (7) $a(n-2) = \kappa - B + \rho + 2\sigma + 3\omega.$
- (8) $b(n-2) = \rho + 2\beta + 3\gamma + 3t.$
- (9) $c(n-2) = 2\sigma + 4\beta + \gamma + \theta + \omega.$
- (10) $a(n-2)(n-3) = 2(\delta - C - 3\omega) + 3(ac - 3\sigma - \chi - 3\omega) + 2(ab - 2\rho - j).$
- (11) $b(n-2)(n-3) = 4k + (ab - 2\rho - j) + 3(bc - 3\beta - 2\gamma - i).$
- (12) $c(n-2)(n-3) = 6h + (ac - 3\sigma - \chi - 3\omega) + 2(bc - 3\beta - 2\gamma - i).$
- (13) $q = b^2 - b - 2k - 2f - 3\gamma - 6t.$
- (14) $r = c^2 - c - 2h - 3\beta.$

Also, reciprocal to these,

- (15) $a' = n'(n' - 1) - 2b' - 3c'.$
- (16) $\kappa' = 3n'(n' - 2) - 6b' - 8c'.$
- (17) $\delta' = \frac{1}{2}n'(n' - 2)(n'^2 - 9) - (n'^2 - n' - 6)(2b' + 3c') + 2b'(b' - 1) + 6b'c' + \frac{9}{2}c'(c' - 1).$
- (18) $a'(n' - 2) = \kappa' - B' + \rho' + 2\sigma' + 3\omega'.$
- (19) $b'(n' - 2) = \rho' + 2\beta' + 3\gamma' + 3t'.$
- (20) $c'(n' - 2) = 2\sigma' + 4\beta' + \gamma' + \theta' + \omega'.$
- (21) $a'(n' - 2)(n' - 3) = 2(\delta' - C' - 3\omega') + 3(a'c' - 3\sigma' - \chi' - 3\omega') + 2(a'b' - 2\rho' - j').$
- (22) $b'(n' - 2)(n' - 3) = 4k' + (a'b' - 2\rho' - j') + 3(b'c' - 3\beta' - 2\gamma' - i').$
- (23) $c'(n' - 2)(n' - 3) = 6h' + (a'c' - 3\sigma' - \chi' - 3\omega') + 2(b'c' - 3\beta' - 2\gamma' - i').$
- (24) $q' = b'^2 - b' - 2k' - 2f' - 3\gamma' - 6t'.$
- (25) $r' = c'^2 - c' - 2h' - 3\beta',$

together with one other independent relation: in all 26 relations between the 46 quantities.

608. The new relation may be presented under several different forms, equivalent to each other in virtue of the foregoing 25 relations; these are

$$(26) \quad 2(n-1)(n-2)(n-3) - 12(n-3)(b+c) + 6q + 6r + 24t + 42\beta + 30\gamma + \frac{8}{3}\theta = \Sigma,$$

$$(27) \quad 26n - 12c - 4C - 10B + \beta - 7j - 8\chi + \frac{2}{3}\theta - 4\omega = \Sigma;$$

in each of which two equations Σ is used to denote the same function of the accented letters that the left-hand side is of the unaccented letters.

$$(28) \quad \begin{aligned} \beta' + \frac{2}{3}\theta' = & 2n(n-2)(11n-24) \\ & + (-66n+184)b \\ & + (-93n+252)c \\ & + 22(2\beta+3\gamma+3t) \\ & + 27(4\beta+\gamma+\theta) \\ & + \beta + \frac{2}{3}\theta \\ & - 24C - 28B - 27j - 38\chi - 73\omega \\ & + 4C' + 10B' + 7j' + 8\chi' - 4\omega'. \end{aligned}$$

Or, reciprocally,

$$(29) \quad \begin{aligned} \beta + \frac{2}{3}\theta = & 2n'(n'-2)(11n'-24) \\ & + (-66n'+184)b' \\ & + (-93n'+252)c' \\ & + 22(2\beta'+3\gamma'+3t') \\ & + 27(4\beta'+\gamma'+\theta') \\ & + \beta' + \frac{2}{3}\theta' \\ & - 24C' - 28B' - 27j' - 38\chi' - 73\omega' \\ & + 4C + 10B + 7j + 8\chi - 4\omega. \end{aligned}$$

The equation (26) expresses that the surface and its reciprocal have the same deficiency; viz. the expression for the deficiency is

$$(30) \quad \begin{aligned} \text{Deficiency} = & \frac{1}{6}(n-1)(n-2)(n-3) - (n-3)(b+c) + \frac{1}{2}(q+r) \\ & + 2t + \frac{7}{2}\beta + \frac{5}{2}\gamma + \frac{2}{9}\theta, \\ = & \frac{1}{6}(n'-1)(n'-2)(n'-3) - \&c. \end{aligned}$$

609. The equation (28) (due to Prof. Cayley) is the correct form of an expression for β' , first obtained by him (with some errors in the numerical coefficients) from independent considerations. But it is best obtained by

means of the equation (26); and (27) is a relation presenting itself in the investigation. In fact, considering α as standing for its value $n(n-1)-2b-3c$, we have from the first 25 equations

$$\begin{array}{ll}
6 & \alpha & = \Sigma, \\
+2 & 3n - c - \kappa & = \Sigma, \\
-2 & a(n-2) - \kappa + B - \rho - 2\sigma - 3\omega & = \Sigma, \\
-4 & b(n-2) - \rho - 2\beta - 3\gamma - 3t & = \Sigma, \\
-6 & c(n-2) - 2\sigma - 4\beta - \gamma - \theta - \omega & = \Sigma, \\
+2 & n + \kappa - \sigma - 2\beta - 4\gamma - 2j - 3\chi - 3\omega & = \Sigma, \\
-3 & 2q - 2\rho + \beta + j & = \Sigma, \\
-2 & 3r + c - 5\sigma - \beta - 4\gamma + \chi - \omega & = \Sigma;
\end{array}$$

multiplying these equations by the numbers set opposite to them respectively, and adding, we find

$$\begin{aligned}
& -2n^3 + 12n^2 + 4n + b(12n-36) + c(12n-48) \\
& -6q - 6r - 4C - 10B - 41\beta - 30\gamma - 24t - 7j - 8\chi + \frac{2}{3}\theta - 4\omega = \Sigma,
\end{aligned}$$

and adding hereto (26) we have the equation (27); and from this (28), or by a like process, (29), is obtained without much difficulty. As to the 8 Σ -equations or symmetries, observe that the first, third, fourth, and fifth are in fact included among the original equations (for an expression which vanishes is in fact $= \Sigma$); we have from them moreover $3n - c = 3a' - \kappa'$, and thence $3n - c - \kappa = 3a' - \kappa' - \kappa$, which is $= \Sigma$, or we have thus the second equation; but the sixth, seventh, and eighth equations have yet to be obtained.

610. The equations (15), (16), (17) give

$$\begin{aligned}
n' &= \alpha(\alpha-1) - 2\delta - 3\kappa, \\
c' &= 3\alpha(\alpha-2) - 6\delta - 8\kappa, \\
b' &= \frac{1}{2}\alpha(\alpha-2)(\alpha^2-9) - (\alpha^2-\alpha-6)(2\delta+3\kappa) + 2\delta(\delta-1) + 6\delta\kappa + \frac{9}{2}\kappa(\kappa-1).
\end{aligned}$$

From (7), (8), (9), we have

$$\begin{aligned}
(\alpha - b - c)(n-2) &= \kappa - B - 6\beta - 4\gamma - 3t - \chi + 2\omega, \\
(\alpha - 2b - 3c)(n-2)(n-3) &= 2(\delta - C) - 8k - 18h - 6bc + 18\beta + 12\gamma + 6i - 6\omega;
\end{aligned}$$

substituting these values for κ and δ , and for α its value $= n(n-1) - 2b - 3c$, we obtain the values of n' , c' , b' ; viz. the value of n' is

$$\begin{aligned}
n' &= n(n-1)^2 - n(7b+12c) + 4b^2 + 8b + 9c^2 + 15c \\
&\quad - 8k - 18h + 18\beta + 12\gamma + 12i - 9t \\
&\quad - 2C - 3B - 3\omega.
\end{aligned}$$

Observe that the effect of a cnicnode C is to reduce the class by 2, and that of a binode B to reduce it by 3.

611. We have

$$(n-2)(n-3) = n^2 - n + (-4n+6) = \alpha + 2b + 3c + (-4n+6);$$

making this substitution in the equations (10), (11), (12), which contain $(n-2)(n-3)$, these become

$$\begin{aligned} a(-4n+6) &= 2(\delta - C) - \alpha^2 - 4\rho - 9\sigma - 2j - 3\chi - 15\omega, \\ b(-4n+6) &= 4k - 2b^2 - 9\beta - 6\gamma - 3t - 2\rho - j, \\ c(-4n+6) &= 6h - 3c^2 - 6\beta - 4\gamma - 2i - 3\sigma - \chi - 3\omega, \end{aligned}$$

which are the foregoing equations (C); adding to each equation four times the corresponding equation with the factor $(n-2)$, these become

$$\begin{aligned} \alpha^2 - 2\alpha &= 2(\delta - C) + 4(\kappa - B) - \sigma - 2j - 3\chi - 3\omega, \\ 2b^2 - 2b &= 4k - \beta + 6\gamma + 12t - 3t + 2\rho - j, \\ 3c^2 - 2c &= 6h + 10\beta + 4\theta - 2i + 5\sigma - \chi + \omega. \end{aligned}$$

Writing in the first of these $\alpha^2 - 2\alpha = n + 2\delta + 3\kappa - \alpha$, and reducing the other two by means of the values of q, r , the equations become

$$\begin{aligned} n - \alpha &= -2C - 4B + \kappa - \sigma - 2j - 3\chi - 3\omega, \\ 2q + \beta + 3t + j &= 2\rho, \\ 3r + c + 2i + \chi &= 5\sigma + \beta + 4\theta + \omega, \end{aligned}$$

which give at once the last three of the 8 Σ -equations.

The reciprocal of the first of these is

$$\sigma' = \alpha - n + \kappa' - 2j' - 3\chi' - 2C' - 4B' - 3\omega',$$

viz. writing herein

$$\alpha = n(n-1) - 2b - 3c \quad \text{and} \quad \kappa' = 3n(n-2) - 6b - 8c,$$

this is

$$\sigma' = 4n(n-2) - 8b - 11c - 2j' - 3\chi' - 2C' - 4B' - 3\omega',$$

giving the order of the spinode curve; viz. for a surface of the order n without singularities, this is $= 4n(n-2)$, the product of the orders of the surface and its Hessian.

612. Instead of obtaining the second and third equations as above, we may to the value of $b(-4n+6)$ add twice the value of $b(n-2)$; and to

twice the value of $c(-4n + 6)$ add three times the value of $c(n - 2)$, thus obtaining equations free from ρ and σ respectively; these equations are

$$\begin{aligned} b(-2n + 2) &= 4k - 2b^2 - 5\beta - 3t + 6t - j, \\ c(-5n + 6) &= 12h - 3c^2 - 6\gamma - 4i - 2\chi + 3\theta - 3\omega, \end{aligned}$$

equations which, introducing therein the values of q and r , may also be written

$$\begin{aligned} b(2n - 4) + 6\gamma + 6t + 3t + j + 4f &= 2q + 5\beta \\ c(5n - 12) + 3\theta &= 6r + 18\beta + 5\gamma + 4i + 2\chi + 3\omega. \end{aligned}$$

Considering as given, n the order of the surface; the nodal curve, with its singularities b, k, f, t ; the cuspidal curve, with its singularities c, h ; and the quantities β, γ, i which relate to the intersections of the nodal and cuspidal curves; the first of the two equations gives j , the number of pinch-points, being singularities of the nodal curve, *quoad* the surface; and the second equation establishes a relation between θ, χ, ω , the numbers of singular points of the cuspidal curve *quoad* the surface.

In the case of a nodal curve only, if this be a complete intersection $P = 0$, $Q = 0$, the equation of the surface is $(A, B, C \mid P, Q)^2 = 0$, and the first equation is

$$b(-2n + 2) = 4k - 2b^2 + 6t - j,$$

or, assuming $t = 0$, say $j = 2(n - 1)b - 2b^2 + 4k$, which may be verified; and so in the case of a cuspidal curve only, when this is a complete intersection $P = 0$, $Q = 0$, the equation of the surface is $(A, B, C \mid P, Q)^2 = 0$, where $AC - B^2 = MP + NQ$; and the second equation is

$$c(-5n + 6) = 12h - 6c^2 - 2\chi + 3\theta - 3\omega,$$

or, say $2\chi + 3\omega = (5n - 6)c - 6c^2 + 12h + 3\theta$, which may also be verified.

613. We may in the first instance out of the 46 quantities consider as given the 14 quantities

$$n : b, k, f, t : c, h, \theta, \chi : \beta, \gamma, i : C, B;$$

then of the 26 relations, 17 determine the 17 quantities

$$\alpha, B, \kappa, \rho, \delta : j, q : r, \omega$$

$$n' : a', B', \kappa' : b', f' : c' : i'$$

and there remain the 9 equations

$$(18), (19), (20), (21), (22), (23), (24), (25), (28),$$

connecting the 15 quantities

$$\rho', \sigma' : k', t', j', q' : h', \theta', \chi', \omega', r' : \beta', \gamma' : C', B'.$$

Taking then further as given the 5 quantities j', χ', t', C', B' ,

equations (18) and (21) give ρ', σ' ,

equation (19) gives $2\beta' + 3\gamma' + 3t'$,

(20) gives $4\beta' + \gamma' + \theta'$,

(28) gives $\beta' + \frac{2}{3}\theta'$,

so that, taking also t' as given, these last three equations determine β', γ', θ' ; and finally

equation (22) gives k' ,

(23) gives h' ,

(24) gives q' ,

(25) gives r' ,

viz. taking as given in all 20 quantities, the remaining 26 will be determined.

614. In the case of the general surface of the order n , without singu-

larities, we have as follow

$$\begin{aligned}
n &= n, \\
a &= n(n-1), \\
\delta &= \frac{1}{2}n(n-1)(n-2)(n-3), \\
\kappa &= n(n-1)(n-2), \\
n' &= n(n-1)^2, \\
a' &= n(n-1), \\
\delta' &= \frac{1}{2}n(n-2)(n^2-9), \\
\kappa' &= 3n(n-2), \\
b' &= \frac{1}{2}n(n-1)(n-2)(n^3-n^2+n-12), \\
k' &= \frac{1}{8}n(n-2)(n^7-6n^6+16n^5-54n^4+164n^3-288n^2 \\
&\quad + 547n^2-1058n^3+1068n^2-1214n+1464), \\
h' &= \frac{1}{2}n(n-2)(n^5-4n^4+7n^3-45n^2+114n^2-111n^2+548n-960), \\
q' &= n(n-2)(n-3)(n^2+2n-4), \\
\rho' &= n(n-2)(n^3-n^2+n-12), \\
c' &= 4n(n-1)(n-2), \\
h' &= \frac{1}{2}n(n-2)(16n^4-64n^3+80n^2-108n+156), \\
i' &= 2n(n-2)(3n-4), \\
\sigma' &= 4n(n-2), \\
\beta' &= 2n(n-2)(11n-24), \\
\gamma' &= 4n(n-2)(n-3)(n^2-3n+16),
\end{aligned}$$

the remaining quantities vanishing.

615. The question of singularities has been considered under a more general point of view by Zeuthen, in the memoir “Recherche des singularités qui ont rapport à une droite multiple d’une surface,” *Math. Annalen*, t. IV. (1871), pp. 1–20. He attributes to the surface:

A number of singular points, viz. points at any one of which the tangents form a cone of the order μ , and class ν , with $y + \eta$ double lines, of which y are tangents to branches of the nodal curve through the point, and $z + \zeta$ stationary lines, whereof z are tangents to branches of the cuspidal curve through the point, and with u double planes and v stationary planes; moreover, these points have only the properties which are the most general in the case of a surface regarded as a locus of points; and Σ denotes a sum extending to all such points. (The foregoing general definition includes the cnicnodes $\mu = \nu = 2$, $y = \eta = z = \zeta = u = v = 0$, and the binodes $\mu = 2$, $\eta = 1$, $z = y = \&c. = 0$.)

And, further, a number of singular planes, viz. planes any one of which touches along a curve of the class μ' and order ν' , with $y' + \eta'$ double tangents,

of which y' are generating lines of the node-couple torse, $z' + \zeta'$ stationary tangents, of which z' are generating lines of the spinode torse, u' double points and v' cusps; it is, moreover, supposed that these planes have only the properties which are the most general in the case of a surface regarded as an envelope of its tangent planes; and Σ' denotes a sum extending to all such planes. (The definition includes the cnictropes $\mu' = \nu' = 2$, $y' = \eta' = z' = \zeta' = u' = v' = 0$, and the bitropes $\mu' = 2$, $\eta' = 1$, $z' = y' = \&c. = 0$.)

616. This being so, and writing

$$x = \nu + 2\eta + 3\zeta, \quad x' = \nu' + 2\eta' + 3\zeta',$$

the equations (7), (8), (9), (10), (11), (12), contain, in respect of the new singularities, additional terms, viz. these are

$$\begin{aligned} a(n-2) &= \cdots + \Sigma [x(\mu-2) - \nu - 2\zeta], \\ b(n-2) &= \cdots + \Sigma [y(\mu-2)], \\ c(n-2) &= \cdots + \Sigma [z(\mu-2)], \\ a(n-2)(n-3) &= \cdots + \Sigma [x(-\nu+7) + 2\eta + 4\zeta], \\ b(n-2)(n-3) &= \cdots + \Sigma [y(-\nu+8)] - \Sigma' (4u' + 3v'), \\ c(n-2)(n-3) &= \cdots + \Sigma [z(-\nu+9)] - \Sigma' (2u'), \end{aligned}$$

and there are of course the reciprocal terms in the reciprocal equations (18), (19), (20), (21), (22), (23). These formulae are given without demonstration in the memoir just referred to: the principal object of the memoir, as shown by its title, is the consideration not of such singular points and planes, but of the multiple right lines of a surface; and in regard to these, the memoir should be consulted.

751.

NOTE ON RIEMANN'S PAPER "VERSUCH EINER
ALLGEMEINEN AUFFASSUNG DER INTEGRATION UND
DIFFERENTIATION*."

[From the *Mathematische Annalen*, t. XVI. (1880), pp. 81, 82.]

The Editors of Riemann's works remark that the paper in question was contained in a MS. of his student time (dated 14 Jan. 1847) and was probably never intended for publication: indeed that he would not in later years have recognised the validity of the principles upon which it is founded. The

idea is however a noticeable one: Riemann considers Z_{x+h} a function of $x+h$, expanded in a doubly infinite, necessarily divergent, series of integer or fractional powers of h , according to the law

$$Z_{x+h} = \sum_{\nu=-\infty}^{\nu=+\infty} k_{\nu} x \cdot h^{\nu}, \quad (2)$$

where the meaning is explained to be that the exponents differ from each other by integer values, in effect, that ν has all the values $\alpha + p$, α a given integer or fractional value, and p any integer number from $-\infty$ to $+\infty$, zero included.

Riemann deduces a theory of fractional differentiation: but without considering the question which has always appeared to me to be the great difficulty in such a theory: what is the real meaning of a complementary function containing an infinity of arbitrary constants? or, in other words, what is the arbitrariness of the complementary function of this nature which presents itself in the theory?

I wish to point out the relation between the paper referred to, and a short paper of my own "On a doubly infinite Series," *Quart. Math. Journ.* t. VI. (1851), pp. 45–47, [102]: this commences with the remark "The following completely paradoxical investigation of the properties of the function Γ (which I have been in possession

* *Werke*, pp. 331–344.

of for some years) may perhaps be found interesting from its connexion with the theories of expansion and divergent series." And I then give the expansion

$$C_n e^x = \sum' [n-r]^n x^{n-r},$$

where n is any integer or fractional number whatever, and the summation extends to all positive and negative integer values (zero included) of r . And I remark that, n being an integer, we have $C_n = \Gamma(n)$, and hence that assuming that this is so in general, or writing

$$\Gamma(n) e^x = \sum' [n-1]^r x^{n-r-1},$$

we have this equation as a definition of $\Gamma(n)$. The point of resemblance of course is that we have a doubly infinite expansion of e^x in a series of integer or fractional powers of x , corresponding to Riemann's like expansion of Z_{x+h} in powers of h .

Cambridge, 10 Sept. 1879.

752.

ON THE FINITE GROUPS OF LINEAR TRANSFORMATIONS
OF A VARIABLE; WITH A CORRECTION.

[From the *Mathematische Annalen*, t. XVI. (1880), pp. 260–263; 439, 440.]

In the paper “Ueber endliche Gruppen linearer Transformationen einer Veränderlichen,” *Math. Ann.* t. XII. (1877), pp. 23–46, Prof. Gordan gave in a very elegant form the groups of 12, 24 and 60 homographic transformations $\frac{ax+b}{cx+d}$. The groups of 12 and 24 are in the like form, the group of 24 thus containing as part of itself the group of 12; but the group of 60 is in a different form, not containing as part of itself the group of 12. It is, I think, desirable to present the group of 60 in the form in which it contains as part of itself Gordan’s group of 12: and moreover to identify the group of 60 with the group of the 60 positive permutations of 5 letters or (writing abc for the cyclical permutation a into b , b into c , c into a , and so in other cases) say with the group of the 60 positive permutations 1, abc , $ab.cd$ and $abcde$.

Any two forms of a group are, it is well known, connected as follows, viz. if $1, \alpha, \beta, \dots$ are the functional symbols of the one form, then those of the other form are $1, \phi\alpha\phi^{-1}, \phi\beta\phi^{-1}, \dots$ (where in the case in question ϕ is a functional symbol of the like homographic form, $\phi x = \frac{Ax+B}{Cx+D}$). But instead of obtaining the new form in this manner, I found it easier to use the values of the rotation-symbol

$$\cos \frac{\theta}{2} + \sin \frac{\theta}{2}(i \cos X + j \cos Y + k \cos Z)$$

for the axes of the icosahedron or dodecahedron, given in my paper “Notes on polyhedra,” *Quart. Math. Jour.* t. VII. (1866), pp. 304–316, [375]; viz. if for any axes, λ, μ, ν denote the parameters of rotation $\tan \frac{\theta}{2} \cos X$, $\tan \frac{\theta}{2} \cos Y$, $\tan \frac{\theta}{2} \cos Z$, then,

by a formula which is in fact equivalent to that given in my note “On the correspondence of Homographies and Rotations,” *Math. Annalen*, t. XV. (1879), pp. 238–240, [660], the corresponding homographic function of x is

$$\frac{(-\nu - i)x + \lambda + i\mu}{(\lambda - i\mu)x + \nu - i},$$

where i denotes $\sqrt{-1}$ as usual.

The new formulae for the group of 60, or icosahedron group, of homographic functions $\frac{\alpha x + \beta}{\gamma x + \delta}$ are contained in the following table, where the four

columns show the values of the coefficients $\alpha, \beta, \gamma, \delta$ respectively: and where in the outside column, the substitution is represented as a permutation-symbol on the five letters $abcde$: moreover for shortness θ is written to denote $\sqrt{5}$.

THE GROUP OF 60.

#	α	β	γ	δ	subst.
1	1	0	0	1	1
2	-1	0	0	1	$ab.cd$
3	0	1	1	0	$ac.bd$
4	0	-1	1	0	$ad.bc$
5	2	$-3 + \theta + i(1 - \theta)$	$-3 + \theta + i(-1 + \theta)$	-2	$bc.de$
6	2	$-3 + \theta + i(-1 + \theta)$	$-3 + \theta + i(1 - \theta)$	-2	$ae.bc$
7	2	$3 - \theta + i(-1 + \theta)$	$3 - \theta + i(1 - \theta)$	-2	$ad.ce$
8	2	$3 - \theta + i(1 - \theta)$	$3 - \theta + i(-1 + \theta)$	-2	$ad.be$
9	2	$-1 - \theta + i(1 - \theta)$	$-1 - \theta + i(-1 + \theta)$	-2	$ae.cd$
10	2	$-1 - \theta + i(-1 + \theta)$	$-1 - \theta + i(1 - \theta)$	-2	$ab.de$
11	2	$1 + \theta + i(-1 + \theta)$	$1 + \theta + i(1 - \theta)$	-2	$bc.cd$
12	2	$1 + \theta + i(1 - \theta)$	$1 + \theta + i(-1 + \theta)$	-2	$ab.ce$
13	2	$-1 - \theta + i(-3 - \theta)$	$-1 - \theta + i(3 + \theta)$	-2	$ac.be$
14	2	$-1 - \theta + i(3 + \theta)$	$-1 - \theta + i(-3 - \theta)$	-2	$bd.ce$
15	2	$1 + \theta + i(3 + \theta)$	$1 + \theta + i(-3 - \theta)$	-2	$ae.bd$
16	2	$1 + \theta + i(-3 - \theta)$	$1 + \theta + i(3 + \theta)$	-2	$ae.de$
17	$-i$	i	1	1	abc
18	$-i$	$-i$	1	1	aeb
19	i	$-i$	1	1	ade
20	i	i	1	-1	acd
21	$-i$	i	1	-1	adb
22	i	i	1	$-i$	abd
23	$-i$	$-i$	1	$-i$	bcd
24	i	$-i$	1	i	bdc
25	$-1 - \theta + i(3 + \theta)$	2	-2	$-1 - \theta + i(-3 - \theta)$	aec
26	$1 + \theta + i(3 + \theta)$	2	-2	$1 + \theta + i(-3 - \theta)$	ace
27	$1 + \theta + i(-3 - \theta)$	2	-2	$1 + \theta + i(3 + \theta)$	bcd
28	$-1 - \theta + i(-3 - \theta)$	2	-2	$-1 - \theta + i(3 + \theta)$	bde
29	$-3 + \theta + i(1 - \theta)$	2	2	$3 - \theta + i(1 - \theta)$	bce
30	$-3 + \theta + i(-1 + \theta)$	2	2	$3 - \theta + i(-1 + \theta)$	bec
31	$3 - \theta + i(-1 + \theta)$	2	2	$-3 + \theta + i(-1 + \theta)$	aed
32	$3 - \theta + i(1 - \theta)$	2	2	$-3 + \theta + i(1 - \theta)$	ade
33	2	$-1 - \theta + i(-1 + \theta)$	$1 + \theta + i(-1 + \theta)$	2	cde
34	2	$1 + \theta + i(1 - \theta)$	$-1 - \theta + i(1 - \theta)$	2	ced
35	2	$-1 - \theta + i(1 - \theta)$	$1 + \theta + i(1 - \theta)$	2	aeb
36	2	$1 + \theta + i(-1 + \theta)$	$-1 - \theta + i(-1 + \theta)$	2	abe
37	$-1 - \theta + i(-3 - \theta)$	2	2	$1 + \theta + i(-3 - \theta)$	$abcde$
38	$-1 - \theta + i(1 - \theta)$	2	2	$1 + \theta + i(1 - \theta)$	$acebd$
39	$-1 - \theta + i(-1 + \theta)$	2	2	$1 + \theta + i(-1 + \theta)$	$adbec$
40	$-1 - \theta + i(3 + \theta)$	2	2	$1 + \theta + i(3 + \theta)$	$aedcb$
41	$1 + \theta + i(3 + \theta)$	2	2	$-1 - \theta + i(3 + \theta)$	$adceb$
42	$1 + \theta + i(-1 + \theta)$	2	2	$-1 - \theta + i(-1 + \theta)$	$acbde$
43	$1 + \theta + i(1 - \theta)$	2	2	$-1 - \theta + i(1 - \theta)$	$aedbc$
44	$1 + \theta + i(-3 - \theta)$	2	2	$-1 - \theta + i(-3 - \theta)$	$abecd$

#	α	β	γ	δ	subst.
45	$-1 - \theta + i(-1 + \theta)$	2	-2	$-1 - \theta + i(1 - \theta)$	<i>acbed</i>
46	$-3 + \theta + i(-1 + \theta)$	2	-2	$-3 + \theta + i(1 - \theta)$	<i>abdce</i>
47	$3 - \theta + i(-1 + \theta)$	2	-2	$3 - \theta + i(1 - \theta)$	<i>aecdb</i>
48	$1 + \theta + i(-1 + \theta)$	2	-2	$1 + \theta + i(1 - \theta)$	<i>adebc</i>
49	$1 + \theta + i(1 - \theta)$	2	-2	$1 + \theta + i(-1 + \theta)$	<i>aecbd</i>
50	$3 - \theta + i(1 - \theta)$	2	-2	$3 - \theta + i(-1 + \theta)$	<i>acdeb</i>
51	$-3 + \theta + i(1 - \theta)$	2	-2	$-3 + \theta + i(-1 + \theta)$	<i>abedc</i>
52	$-1 - \theta + i(1 - \theta)$	2	-2	$-1 - \theta + i(-1 + \theta)$	<i>adbce</i>
53	2	$-3 + \theta + i(-1 + \theta)$	$3 - \theta + i(-1 + \theta)$	2	<i>aebde</i>
54	2	$-1 - \theta + i(3 + \theta)$	$1 + \theta + i(3 + \theta)$	2	<i>abcd</i>
55	2	$1 + \theta + i(-3 - \theta)$	$-1 - \theta + i(-3 - \theta)$	2	<i>adebc</i>
56	2	$3 - \theta + i(1 - \theta)$	$-3 + \theta + i(1 - \theta)$	2	<i>acdbe</i>
57	2	$-3 + \theta + i(1 - \theta)$	$3 - \theta + i(1 - \theta)$	2	<i>abdec</i>
58	2	$-1 - \theta + i(-3 - \theta)$	$1 + \theta + i(-3 - \theta)$	2	<i>adcbe</i>
59	2	$1 + \theta + i(3 + \theta)$	$-1 - \theta + i(3 + \theta)$	2	<i>aebcd</i>
60	2	$3 - \theta + i(-1 + \theta)$	$-3 + \theta + i(-1 + \theta)$	2	<i>acedb</i>

This contains (as one of five groups of 12) the group of the positive permutations of *abcd*; and, completing this into a group of 24, we have

GROUPS OF 12 AND 24.

#	α	β	γ	δ	subst.
1	1	0	0	1	1
2	-1	0	0	1	<i>ab.cd</i>
3	0	1	1	0	<i>ac.bd</i>
4	0	-1	1	0	<i>ad.bc</i>
5	-i	i	1	1	<i>abc</i>
6	-1	-i	1	i	<i>acb</i>
7	1	-i	1	1	<i>adc</i>
8	-i	-i	1	-1	<i>acd</i>
9	i	-i	1	-1	<i>aeb</i>
10	1	i	1	-i	<i>abd</i>
11	-1	-i	1	-i	<i>bcd</i>
12	i	-i	1	i	<i>bdc</i>
13	i	i	1	1	<i>adbc</i>
14	-i	i	1	1	<i>acbd</i>
15	1	1	1	-1	<i>cd</i>
16	-1	i	-1	1	<i>ab</i>
17	1	-i	1	1	<i>aedb</i>
18	-i	-1	-i	i	<i>bd</i>
19	i	1	1	i	<i>abcd</i>
20	1	-1	1	-1	<i>bc</i>
21	-1	-1	1	-1	<i>abdc</i>
22	i	-1	1	-i	<i>ac</i>
23	-1	i	1	-i	<i>adcb</i>
24	-1	1	1	1	<i>ad</i>

The groups of 60 and 24 thus each of them contain the group of 12,

$$x, -\frac{1}{x}, \frac{1-x}{1+x}, \frac{1+x}{1-x}, \frac{x+i}{x-i}, \frac{x-i}{-x+i}.$$

It may be remarked that, to verify the periodicities of the forms contained in the group of 60, we have as the conditions that

$\frac{\alpha x + \beta}{\gamma x + \delta}$ may be periodic of the order 2, $\frac{\alpha + \delta}{\alpha\delta - \beta\gamma} = 0$, that is, $\alpha + \delta = 0$,

$$\begin{array}{ccccccc} \text{,,} & \text{,,} & \text{,,} & 3, & \text{,,} & = & 1, \\ \text{,,} & \text{,,} & \text{,,} & 5, & \text{,,} & = & \frac{1}{2}(3 + \sqrt{5}). \end{array}$$

For instance, in the form

$$\frac{[-1 - \Theta + i(-3 - \Theta)]x + 2}{2x + [1 + \Theta + i(-3 - \Theta)]},$$

we have

$$\alpha\delta = -(1 + \Theta)^2 - (3 + \Theta)^2 i^2 = -2\Theta - 8\Theta, \quad \beta\gamma = 4,$$

$$\alpha + \delta = -2i(3 + \Theta);$$

and therefore

$$\frac{(\alpha + \delta)^2}{\alpha\delta - \beta\gamma} = \frac{-4(3 + \Theta)^2}{-3(3 + \Theta)}, \quad = \frac{3 + \Theta}{2} = \frac{1}{2}(3 + \sqrt{5}),$$

as it should be.

Cambridge, 11 Nov. 1879.

CORRECTION*, PP. 439, 440.

I erroneously assumed that the symbol *adcb* could be taken as corresponding to the linear transformation *ix*: but this was obviously wrong, for it gave *bd* as corresponding to the transformation *-ix*, and these are not of the same order, but of the orders 4 and 2 respectively. The proper symbol is *adbc*, as given above, and the remaining eleven symbols are then at once obtained.

Cambridge, 17 Feb. 1880.

* The correction in the Table of the Groups of 12 and 24 has been inserted in the Table as now printed on p. 240; it applies to the second half of the column of symbols on the extreme right-hand.

753.

ON A THEOREM RELATING TO THE MULTIPLE
THETA-FUNCTIONS.

[From the *Mathematische Annalen*, t. XVII. (1880), pp. 115–122.]

I PROPOSE—partly for the sake of the theorem itself, partly for that of the notation which will be employed—to demonstrate the general theorem (3'), p. 4, of Dr Schottky's *Abriss einer Theorie der Abel'schen Functionen von drei Variabeln*, (Leipzig, 1880), which theorem is there presented in the form:

$$e^{-H(u_1, \dots; \mu', \nu')} \Theta(u_1 + 2\varpi'_1, \dots; \mu, \nu) = e^{-2\pi i \mu \nu'} \Theta(u_1, \dots; \mu + \mu', \nu + \nu'), \quad (3')$$

but which I write in the slightly different form

$$\exp.[-H(u; \mu', \nu')] \cdot \Theta(u + 2\varpi'; \mu, \nu) = \exp.[-2\pi i \mu \nu'] \cdot \Theta(u; \mu + \mu', \nu + \nu').$$

I remark that the theorem is given in the preliminary paragraphs the contents of which are, as mentioned by the Author, derived from Herr Weierstrass: and that the form of the theta-function is a very general one, depending on the general quadric function

$$G(u_1, \dots, u_p; u_1, \dots, u_p)$$

of $2p$ variables, p being the number of the arguments $u_1, \dots, u_p$ (in fact, the periods are not reduced to the normal form, but are arbitrary); and the characters $\nu_1, \dots, \nu_p, \mu_1, \dots, \mu_p$, instead of having each of them the value 0 or 1, have each of them any integer or fractional value whatever. The meaning of the theorem (u denoting a set or row of p letters $u_1, \dots, u_p$, and so in other cases), is that the function

$$\Theta(u; \mu + \mu', \nu + \nu')$$

with the new characters $\mu + \mu'$ and $\nu + \nu'$ is, save as to an exponential factor, equal to the function $\Theta(u + 2\varpi'; \mu, \nu)$ with the original characters μ, ν , but with the new arguments $u + 2\varpi'$.

Notation.

This is in some measure a development of the notation employed in my "Memoir on the Theory of Matrices," *Phil. Trans.* t. CXLVIII. (1858), pp. 17–37, [152]: I use certain single letters u , etc. to denote sets or rows each of p letters, $u = (u_1, \dots, u_p)$: or if, to fix the ideas $p = 3$, then $u = (u_1, u_2, u_3)$, and so in other cases.

But I use certain other letters a , etc. to denote squares or matrices each of p^2 letters; thus, if $p = 3$ as before,

$$a = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix},$$

and in any such case the transposed matrix is denoted by the same letter enclosed in parentheses

$$(a) = \begin{vmatrix} a_{11} & a_{21} & a_{31} \\ a_{12} & a_{22} & a_{32} \\ a_{13} & a_{23} & a_{33} \end{vmatrix}.$$

The sum $u + v$ of the row-letters $u, = (u_1, u_2, u_3)$ and $v, = (v_1, v_2, v_3)$ denotes the row $(u_1 + v_1, u_2 + v_2, u_3 + v_3)$: and in like manner the sum $a + b$ of the two matrices, or square-letters a and b , denotes the matrix

$$\begin{vmatrix} a_{11} + b_{11} & a_{12} + b_{12} & a_{13} + b_{13} \\ a_{21} + b_{21} & a_{22} + b_{22} & a_{23} + b_{23} \\ a_{31} + b_{31} & a_{32} + b_{32} & a_{33} + b_{33} \end{vmatrix},$$

and similarly for a sum of three or more terms.

The product $uv, = (u_1, u_2, u_3)(v_1, v_2, v_3)$, of the two row-letters u, v denotes the single term $u_1v_1 + u_2v_2 + u_3v_3$. We have $uv = vu$.

The product

$$au, = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} (u_1, u_2, u_3),$$

of a preceding square-letter a and a succeeding row-letter u , denotes the set or row

$$(a_{11}, a_{12}, a_{13})(u_1, u_2, u_3), (a_{21}, a_{22}, a_{23})(u_1, u_2, u_3), (a_{31}, a_{32}, a_{33})(u_1, u_2, u_3);$$

the notation ua is not employed.

The product

$$auv = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} (u_1, u_2, u_3)(v_1, v_2, v_3),$$

of a preceding square-letter a followed by the two row-letters u and v , denotes the single term

$$\begin{aligned} & (a_{11}, a_{12}, a_{13})(u_1, u_2, u_3) v_1 \\ & + (a_{21}, a_{22}, a_{23})(u_1, u_2, u_3) v_2 \\ & + (a_{31}, a_{32}, a_{33})(u_1, u_2, u_3) v_3. \end{aligned}$$

Observe that auv is not in general $= avu$; but it is easy to verify that $auv = (a)vu$; and hence if $(a) = a$, that is, if the matrix a be symmetrical, then $auv = avu$.

A product of two matrices

$$ab, \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} \begin{vmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \\ b_{31} & b_{32} & b_{33} \end{vmatrix},$$

denotes a matrix

$$\begin{vmatrix} (b_{11}, b_{21}, b_{31}) & (b_{12}, b_{22}, b_{32}) & (b_{13}, b_{23}, b_{33}) \\ (a_{11}, a_{12}, a_{13}) & \cdots & \cdots \\ (a_{21}, a_{22}, a_{23}) & \cdots & \cdots \\ (a_{31}, a_{32}, a_{33}) & \cdots & \cdots \end{vmatrix};$$

viz. the top-line of the compound matrix is

$$(a_{11}, a_{12}, a_{13})(b_{11}, b_{21}, b_{31}), (a_{11}, a_{12}, a_{13})(b_{12}, b_{22}, b_{32}), (a_{11}, a_{12}, a_{13})(b_{13}, b_{23}, b_{33}),$$

and so for the other lines: or expressing this in words, we say that any line of the compound matrix is obtained by compounding the corresponding line of the first or further component matrix with the several columns of the second or nearer component matrix.

Clearly ab is not in general $= ba$. We may easily verify that $(ab) = (b)(a)$, that is, the transposed matrix (ab) is that obtained by the composition of the transposed matrix (b) as first or further matrix, with the transposed matrix (a) as second or nearer matrix. Even if a and b are each symmetrical, we do not in general have $ab = ba$, but only $(ab) = ba$, or what is the same thing, $ab = (ba)$.

In a symbol such as $abuv$, we first combine a, b into a single matrix ab , and then regard the expression as a combination such as auv : the expression denotes therefore a single term. The theory might be explained in greater detail; but the mode of working with row- and square-letters will be readily understood from what precedes.

In all that follows, $u, \mu, \nu, \mu', \nu', \varpi, \varpi', \zeta'$ are row-letters; $a, b, h, \omega, \omega', \eta, \eta'$ are square-letters: a and b are symmetrical, viz. $a = (a)$, $b = (b)$.

And I write

$$\begin{aligned} (\cdot)(u, v)^2 &= (a, h, b)(u, v)^2 \\ &= au^2 + 2huv + bv^2 \end{aligned}$$

$$\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} (u_1, u_2, u_3)^2$$

$$\begin{aligned}
& +2 \begin{vmatrix} h_{11} & h_{12} & h_{13} \\ h_{21} & h_{22} & h_{23} \\ h_{31} & h_{32} & h_{33} \end{vmatrix} (u_1, u_2, u_3)(v_1, v_2, v_3) \\
& + \begin{vmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \\ b_{31} & b_{32} & b_{33} \end{vmatrix} (v_1, v_2, v_3)^2
\end{aligned}$$

to denote the general quadric function of the $2p$ letters u, v , with

$$\frac{1}{2}p(p+1) + p^2 + \frac{1}{2}p(p+1), \quad = p(2p+1)$$

coefficients. It is assumed that the determinant formed with the $\frac{1}{2}p(p+1)$ coefficients b is negative: this is the necessary and sufficient condition for the convergence of the series.

Definition of $\Theta(u; \mu, \nu)$.

$\Theta(u; \mu, \nu)$, the general theta-function with p arguments u , and $2p$ characters μ, ν , is the sum of a p -tuple series of exponentials

$$\Theta(u; \mu, \nu) = \sum \exp. [(\cdot)(n, n + \nu)^2 + 2\pi i u (n + \nu)],$$

where each of the letters $n, = (n_1, \dots, n_p)$, has all integer values (zero included) from $-\infty$ to $+\infty$.

The general theorem in regard to $\Theta(u; \mu, \nu)$.

This is

$$\exp. [-H(u; \mu', \nu')] \cdot \Theta(u + 2\varpi'; \mu, \nu) = \exp. [-2\pi i \mu \nu'] \cdot \Theta(u; \mu + \mu', \nu + \nu'),$$

establishing a relation between the function $\Theta(u; \mu + \mu', \nu + \nu')$, with arbitrary character-increments μ', ν' , and the function $\Theta(u + 2\varpi'; \mu, \nu)$ with the original characters, but with new arguments $u + 2\varpi'$. Also $H(u; \mu', \nu')$ denotes a function, linear as regards the arguments u , but quadric as regards μ' and ν' ; $-2\pi i \mu \nu'$ is a single term depending only on μ and ν' ; and the theorem thus is that the two functions differ only by an exponential factor. The relations between the constants will be obtained in the course of the investigation.

Demonstration.

The truth of the theorem depends on the equality of corresponding exponentials on the two sides of the equation: viz. substituting for the theta-functions their values, and comparing the exponents or arguments of the exponentials: writing also for convenience

$$G(u + 2\varpi', n + \nu),$$

to denote the quadric function $(\cdot)(u + 2\varpi', n + \nu)^2$; we ought to have

$$\begin{aligned} & -H(u; \mu', \nu') + G(u + 2\varpi', n + \nu) + 2\pi i u(n + \nu) \\ & = -2\pi i \mu \nu' + G(u, n + \nu + \nu') + 2\pi i(\mu + \mu')(n + \nu + \nu'), \end{aligned}$$

or say

$$H(u; \mu', \nu') = G(u + 2\varpi', n + \nu) - G(u, n + \nu + \nu') - 2\pi i(n + \nu + \nu')\mu'.$$

In this equation, if true at all, the terms containing n must destroy each other; assuming that they do so, the equation becomes

$$H(u; \mu', \nu') = G(u + 2\varpi', \nu) - G(u, \nu + \nu') - 2\pi i(\nu + \nu')\mu'.$$

Consider first the terms in n : the right-hand side is

$$\begin{aligned} & = a(u + 2\varpi')^2 + 2h(u + 2\varpi')(n + \nu) + b(n + \nu)^2 \\ & \quad - au^2 - 2hu(n + \nu + \nu') - b(n + \nu + \nu')^2 - 2\pi i n \mu', \end{aligned}$$

and the terms herein which contain n thus are

$$\begin{aligned} & 2h(u + 2\varpi')n + bn^2 + 2bn\nu \\ & \quad - 2hun - bn^2 - 2bn(\nu + \nu') - 2\pi i n \mu', \\ & = 4h\varpi'n - 2bn\nu' - 2\pi i n \mu', \end{aligned}$$

which, b being symmetrical, may be written

$$= 2(2h\varpi' - b\nu' - \pi i \mu')n,$$

and these terms will vanish if, and only if

$$2h\varpi' - b\nu' - \pi i \mu' = 0,$$

a system of p equations connecting ϖ', μ', ν' .

Assuming them to be satisfied, the remaining relation,

$$H(u; \mu', \nu') = G(u + 2\varpi', \nu) - G(u, \nu + \nu') - 2\pi i(\nu + \nu')\mu',$$

becomes

$$\begin{aligned} H(u; \mu', \nu') & = a(u + 2\varpi')^2 + 2h(u + 2\varpi')\nu + b\nu^2 \\ & \quad - au^2 - 2hu(\nu + \nu') - b(\nu + \nu')^2 - 2\pi i(\nu + \nu')\mu'. \end{aligned}$$

Here, a and b being symmetrical, we have

$$a(u + 2\varpi')^2 = au^2 + 4a\varpi'u + 4a\varpi'^2, \quad b(\nu + \nu')^2 = b\nu^2 + 2b\nu'\nu + b\nu'^2,$$

and the value therefore is

$$= 4a(\varpi'u + \varpi'^2) + 2h(2\varpi'\nu - u\nu') - b(2\nu'\nu + \nu'^2) - 2\pi i(\nu + \nu')\mu'.$$

On the right-hand side, putting the term in h under the form

$$-2h(u + \varpi')\nu' + 2h\varpi'(2\nu + \nu'), \quad = -2(h)\nu'(u + \varpi') + 2h\varpi'(2\nu + \nu'),$$

and the last term under the form

$$-\pi i\mu'(2\nu + \nu') - \pi i\mu'\nu',$$

the equation becomes

$$\begin{aligned} H(u; \mu', \nu') &= (4a\varpi' - 2(h)\nu')(u + \varpi') - \pi i\mu'\nu' \\ &\quad + (2h\varpi' - b\nu' - \pi i\mu')(2\nu + \nu'), \end{aligned}$$

where the second line vanishes in virtue of the foregoing equation

$$2h\varpi' - b\nu' - \pi i\mu' = 0;$$

the equation thus is

$$H(u; \mu', \nu') = (4a\varpi' - 2(h)\nu')(u + \varpi') - \pi i\mu'\nu',$$

which equation, regarding therein ϖ' as a linear function of μ' and ν' , shows that $H(u; \mu', \nu')$ is a function linear as regards u (and containing this only through $u + \varpi'$), but quadric as regards μ', ν' .

Introducing the new row-letter ζ' , we may write

$$H(u; \mu', \nu') = 2\zeta'(u + \varpi') - \pi i\mu'\nu',$$

viz. the expression on the right-hand side is here assumed as the value of the function

$$H(u; \mu', \nu'), \quad = G(u + 2\varpi', \nu) - G(u, \nu + \nu') - 2\pi i(\nu + \nu')\mu',$$

and the theorem then is

$$\exp.[-H(u; \mu', \nu')] \cdot \Theta(u + 2\varpi'; \mu, \nu) = \exp.[-2\pi i\mu\nu'] \cdot \Theta(u; \mu + \mu', \nu + \nu'),$$

where, by what precedes,

$$2h\varpi' - b\nu' - \pi i\mu' = 0,$$

$$2a\varpi' - (h)\nu' - \zeta' = 0,$$

$2p$ equations for determining the $2p$ functions ϖ' , ζ' as linear functions of μ' , ν' : which equations depend on the $p(2p+1)$ constants a , b , h .

Suppose that the resulting values of ϖ' and ζ' are

$$\varpi' = \omega\mu' + \omega'\nu',$$

$$\zeta' = \eta\mu' + \eta'\nu',$$

where ω , ω' , η , η' are square-letters; then, regarding a , b , h as arbitrary, the $4p^2$ new constants ω , ω' , η , η' cannot be all of them arbitrary, but must be connected by $4p^2 - p(2p+1)$, $= p(2p-1)$ equations.

We may regard ω , ω' , η , η' as satisfying these $p(2p-1)$ equations, but as being otherwise arbitrary; the foregoing equations then are

$$2h\varpi' - b\nu' - \pi i\mu' = 0,$$

$$2a\varpi' - (h)\nu' - \zeta' = 0,$$

$$\varpi' = \omega\mu' + \omega'\nu',$$

$$\zeta' = \eta\mu' + \eta'\nu',$$

which lead to the equations connecting a , b , h with ω , ω' , η , η' .

The first and second equations, substituting for ϖ' and ζ' their values, become

$$(2h\omega - \pi i)\mu' + (2h\omega' - b)\nu' = 0,$$

$$(2a\omega - \eta)\mu' + (2a\omega' - (h) - \eta')\nu' = 0,$$

or μ' , ν' being arbitrary, we thus obtain the $4p^2$ equations

$$2a\omega - \eta = 0,$$

$$2h\omega - \pi i = 0,$$

$$2a\omega' - \eta' - (h) = 0,$$

$$2h\omega' - b = 0,$$

which are the equations in question. It is to be observed that in πi is, like the other symbols, a matrix, viz. it is regarded as containing the matrix unity; or, what is the same thing, it denotes

$$\pi i \begin{vmatrix} 1 & 0 & 0 & \cdots \\ 0 & 1 & 0 & \cdots \\ \vdots & & \ddots & \end{vmatrix}.$$

We can eliminate a , b , h from these equations and thus obtain the $p(2p-1)$ equations before referred to, which connect the $4p^2$ constants ω , ω' , η , η' . I give, but without a complete explanation, the steps of the elimination.

The equation $2a\omega - \eta = 0$, may be written in the form

$$2(a\omega) - (\eta) = 0,$$

that is,

$$2(\omega)(a) - (\eta) = 0,$$

or since $(a) = a$, this is

$$2(\omega)a - (\eta) = 0;$$

from the original form, and the new form respectively, we find

$$2(\omega)a\omega - (\omega)\eta = 0, \quad 2(\omega)a(\omega) - (\eta)\omega = 0;$$

and comparing these

$$(\omega)\eta - (\eta)\omega = 0, \quad (\text{first result}).$$

The equation $2a\omega' - \eta' - (h) = 0$, or say $(h) = -\eta' + 2a\omega'$, may be written in the form

$$h = -(\eta') + 2(a\omega'),$$

that is, since $a = (a)$,

$$h = -(\eta') + 2(\omega')a;$$

and we thence deduce

$$h\omega = -(\eta')\omega + 2(\omega')a\omega.$$

But from the equation $2a\omega - \eta = 0$, we have $2(\omega')a\omega - (\omega')\eta = 0$, and the equation thus becomes $h\omega = -(\eta')\omega + (\omega')\eta$; which, in virtue of $2h\omega - \pi i = 0$, becomes

$$\frac{1}{2}\pi i = -(\eta')\omega + (\omega')\eta, \quad (\text{second result}).$$

From the equation above obtained, $h = -(\eta') + 2(\omega')a$, we have

$$h\omega' = -(\eta')\omega' + 2(\omega')a\omega';$$

in virtue of $2h\omega' - b = 0$, this becomes $-2(\eta')\omega' + 4(\omega')a\omega' = b$; an equation which may also be written $-2((\eta')\omega') + 4((\omega')a\omega') = (b)$, or, what is the same thing, $-2(\omega')\eta' + 4(\omega')(a)\omega' = (b)$; or since $(a) = a$ and $(b) = b$, this is

$$-2(\omega')\eta' + 4(\omega')a\omega' = b :$$

and comparing with the original equation

$$-2(\eta')\omega' + 4(\omega')a\omega' = b,$$

we obtain

$$(\omega')\eta' - (\eta')\omega' = 0, \quad (\text{third result}).$$

We have thus the three systems

$$\begin{aligned}(\omega)\eta - (\eta)\omega &= 0, & \frac{1}{2}p(p-1) \text{ equations,} \\ (\omega')\eta - (\eta')\omega &= \frac{1}{2}\pi i, & p^2 \text{ equations,} \\ (\omega')\eta' - (\eta')\omega' &= 0, & \frac{1}{2}p(p-1) \text{ equations,}\end{aligned}$$

in all $p(2p-1)$ equations. As to these systems, observe that $(\omega)\eta$, $(\eta)\omega$, etc., are all of them matrices of p^2 terms; each of the three systems denotes therefore in the first instance p^2 equations, viz. the equations obtained by equating to zero the several terms of such a matrix: but in the first system each diagonal term so equated to zero gives the identity $0=0$; and equating to zero the terms which are symmetrical in regard to the diagonal we obtain twice over, in the forms $P=0$, and $-P=0$, one and the same equation; the number of equations is thus diminished from p^2 to $\frac{1}{2}p(p-1)$; and similarly in the third system the number of equations is $=\frac{1}{2}p(p-1)$: but for the second system the number of equations is really $=p^2$. It is hardly necessary to remark that in this second system $\frac{1}{2}\pi i$ is as before regarded as a matrix.

The foregoing three systems of equations are in fact the equations (6) p. 4 of Dr Schottky's work.

Cambridge, 12 July, 1880.

754.

ON THE CONNEXION OF CERTAIN FORMULÆ IN ELLIPTIC FUNCTIONS.

[From the *Messenger of Mathematics*, vol. IX. (1880), pp. 23–25.]

In reference to a like question in the theory of the double ϑ -functions, it is interesting to show that (if not completely, at least very nearly) the single formula

$$\Pi(u, a) = u \frac{\Theta'a}{\Theta a} + \frac{1}{2} \log \frac{\Theta(u-a)}{\Theta(u+a)},$$

that is,

$$\int_0^u \frac{k^2 \operatorname{sn} a \operatorname{cn} a \operatorname{dn} a \operatorname{sn}^2 u \, du}{1 - k^2 \operatorname{sn}^2 a \operatorname{sn}^2 u} = u \frac{\Theta'a}{\Theta a} + \frac{1}{2} \log \frac{\Theta(u-a)}{\Theta(u+a)},$$

leads not only to the relation

$$\log \Theta u = \frac{1}{2} \log \frac{2k'K}{\pi} + \frac{1}{2} \left(1 - \frac{E}{K}\right) u^2 - k^2 \int_0^u du \int_0^u du \operatorname{sn}^2 u,$$

between the functions Θ , sn , but also to the addition-equation for the function sn .

Writing in the equation a indefinitely small, and assuming only that $\text{sn } a$, $\text{cn } a$, $\text{dn } a$ then become a , 1 , 1 , respectively, the equation is

$$k^2 a \int_0^u \text{sn}^2 u \, du = u \frac{\Theta' a}{\Theta a} + \frac{1}{2} \log \frac{\Theta u - a \Theta' u}{\Theta u + a \Theta' u} = u \frac{\Theta' 0}{\Theta 0} - a \frac{\Theta' u}{\Theta u},$$

that is,

$$\frac{\Theta' u}{\Theta u} = u \frac{\Theta'' 0}{\Theta 0} - k^2 \int_0^u du \text{sn}^2 u,$$

or, integrating from $u = 0$, this is

$$\log \Theta u = C + \frac{1}{2} u^2 \frac{\Theta'' 0}{\Theta 0} - k^2 \int_0^u du \int_0^u du \text{sn}^2 u,$$

which, except as regards the determination of the constants, is the required equation for $\log \Theta u$.

Next, differentiating twice the equation for $\Pi(u, a)$, and once the equation obtained for $\frac{d\Theta'}{du}$, we have

$$k^2 \text{sn } a \text{ cn } a \text{ dn } a \frac{d}{du} \left(\frac{\text{sn}^2 u}{1 - k^2 \text{sn}^2 a \text{sn}^2 u} \right) = \frac{1}{2} \frac{\Theta'' \Theta - \Theta'^2}{\Theta^2} (u-a) - \frac{1}{2} \frac{\Theta'' \Theta - \Theta'^2}{\Theta^2} (u+a),$$

and

$$\frac{\Theta'' \Theta - \Theta'^2}{\Theta^2} = \frac{\Theta'' 0}{\Theta 0} - k^2 \text{sn}^2 u,$$

where, for shortness, $\frac{\Theta'' 0}{\Theta 0}$ is written to denote $\frac{\Theta'' u \Theta u - (\Theta' u)^2}{\Theta u^2}$, and the like in the first equation; the right-hand side of the first equation therefore is

$$= -\frac{1}{2} k^2 \{ \text{sn}^2(u-a) - \text{sn}^2(u+a) \},$$

or the equation becomes

$$2 \text{sn } a \text{ cn } a \text{ dn } a \frac{d}{du} \frac{\text{sn}^2 u}{1 - k^2 \text{sn}^2 u \text{sn}^2 a} = \text{sn}^2(u+a) - \text{sn}^2(u-a),$$

that is,

$$\frac{4 \text{sn } u \text{sn}' u \text{sn } a \text{cn } a \text{dn } a}{(1 - k^2 \text{sn}^2 u \text{sn}^2 a)^2} = \text{sn}^2(u+a) - \text{sn}^2(u-a).$$

The numerator on the left-hand side must be a symmetrical function of u , a , and hence (even if the value of $\text{sn}' u$ were unknown) it would appear that $\text{sn}' u$ must be a mere constant multiple of $\text{cn } u \text{dn } u$; assuming, however, the actual value, $\text{sn}' u = \text{cn } u \text{dn } u$, the formula is

$$\frac{4 \text{sn } u \text{cn } u \text{dn } u \text{sn } a \text{cn } a \text{dn } a}{(1 - k^2 \text{sn}^2 u \text{sn}^2 a)^2} = \text{sn}^2(u+a) - \text{sn}^2(u-a) = \{ \text{sn}(u+a) + \text{sn}(u-a) \} \{ \text{sn}(u+a) - \text{sn}(u-a) \}.$$

The factor $\{\operatorname{sn}(u+a) + \operatorname{sn}(u-a)\}$ becomes $= 2 \operatorname{sn} u$ for $a = 0$, and this suggests that the factor $\operatorname{sn} u$ on the left-hand side is a factor of $\{\operatorname{sn}(u+a) + \operatorname{sn}(u-a)\}$. That $\operatorname{cn} u$ is not a factor hereof would follow from the properties of the period K ; viz. for $u = K$, $\operatorname{cn} u = 0$, but $\{\operatorname{sn}(u+a) + \operatorname{sn}(u-a)\}$, $= 2 \operatorname{sn}(K+a)$ is not $= 0$; and, similarly, that $\operatorname{dn} u$ is not a factor from the properties of the period iK' ; hence, $\operatorname{cn} u$, $\operatorname{dn} u$ belong to the other factor $\{\operatorname{sn}(u+a) - \operatorname{sn}(u-a)\}$, and by symmetry $\operatorname{cn} a$, $\operatorname{dn} a$ belong to the first-mentioned factor. And we are thus led to assume

$$\begin{aligned}\operatorname{sn}(u+a) + \operatorname{sn}(u-a) &= 2M \operatorname{sn} u \operatorname{cn} a \operatorname{dn} a, \\ \operatorname{sn}(u+a) - \operatorname{sn}(u-a) &= 2M' \operatorname{sn} a \operatorname{cn} u \operatorname{dn} u,\end{aligned}$$

where

$$\text{denom.} = 1 - k^2 \operatorname{sn}^2 a \operatorname{sn}^2 u,$$

and $MM' = 1$. Some further investigation is wanting to show that M and M' are constants, but assuming that they are so and each $= 1$, the formulae give at once the ordinary expression for $\operatorname{sn}(u+a)$; that is, we have the addition-equation for the function sn .

II.

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
2	1	4	3	6	5	8	7	10	9	12	11	14	13	16	15
3	4	1	2	7	8	5	6	11	12	9	10	15	16	13	14
4	3	2	1	8	7	6	5	12	11	10	9	16	15	14	13
5	6	7	8	1	2	3	4	14	13	16	15	10	9	12	11
6	5	8	7	2	1	4	3	13	14	15	16	9	10	11	12
7	8	5	6	3	4	1	2	16	15	14	13	12	11	10	9
8	7	6	5	4	3	2	1	15	16	13	14	11	12	9	10
9	10	11	12	14	13	16	15	1	2	3	4	6	5	8	7
10	9	12	11	13	14	15	16	2	1	4	3	5	6	7	8
11	12	9	10	16	15	14	13	3	4	1	2	8	7	6	5
12	11	10	9	15	16	13	14	4	3	2	1	7	8	5	6
13	14	15	16	10	9	12	11	6	5	8	7	1	2	3	4
14	13	16	15	9	10	11	12	5	6	7	8	2	1	4	3
15	16	13	14	12	11	10	9	8	7	6	5	3	4	1	2
16	15	14	13	11	12	9	10	7	8	5	6	4	3	2	1

III.

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
2	1	4	3	6	5	8	7	10	9	12	11	14	13	16	15
3	4	1	2	7	8	5	6	11	12	9	10	15	16	13	14
4	3	2	1	8	7	6	5	12	11	10	9	16	15	14	13
5	6	7	8	1	2	3	4	15	16	13	14	11	12	9	10
6	5	8	7	2	1	4	3	16	15	14	13	12	11	10	9
7	8	5	6	3	4	1	2	13	14	15	16	9	10	11	12
8	7	6	5	4	3	2	1	14	13	16	15	10	9	12	11
9	10	11	12	15	16	13	14	1	2	3	4	7	8	5	6
10	9	12	11	16	15	14	13	2	1	4	3	8	7	6	5
11	12	9	10	13	14	15	16	3	4	1	2	5	6	7	8
12	11	10	9	14	13	16	15	4	3	2	1	6	5	8	7
13	14	15	16	11	12	9	10	7	8	5	6	1	2	3	4
14	13	16	15	12	11	10	9	8	7	6	5	2	1	4	3
15	16	13	14	9	10	11	12	5	6	7	8	3	4	1	2
16	15	14	13	10	9	12	11	6	5	8	7	4	3	2	1

IV.

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
2	1	4	3	6	5	8	7	10	9	12	11	14	13	16	15
3	4	1	2	7	8	5	6	11	12	9	10	15	16	13	14
4	3	2	1	8	7	6	5	12	11	10	9	16	15	14	13
5	6	7	8	1	2	3	4	16	15	14	13	12	11	10	9
6	5	8	7	2	1	4	3	15	16	13	14	11	12	9	10
7	8	5	6	3	4	1	2	14	13	16	15	10	9	12	11
8	7	6	5	4	3	2	1	13	14	15	16	9	10	11	12
9	10	11	12	16	15	14	13	1	2	3	4	8	7	6	5
10	9	12	11	15	16	13	14	2	1	4	3	7	8	5	6
11	12	9	10	14	13	16	15	3	4	1	2	6	5	8	7
12	11	10	9	13	14	15	16	4	3	2	1	5	6	7	8
13	14	15	16	12	11	10	9	8	7	6	5	1	2	3	4
14	13	16	15	11	12	9	10	7	8	5	6	2	1	4	3
15	16	13	14	10	9	12	11	6	5	8	7	3	4	1	2
16	15	14	13	9	10	11	12	5	6	7	8	4	3	2	1

The foregoing investigation of the synthematic arrangements is exhaustive: it thereby appears that there are at most four types, viz. that every synthematic arrangement is of the type of one or other of the four arrangements above written down. The real nature of these is perhaps more clearly seen by means of the corresponding squares; and it will be observed, that there is in the first square a repetition of parts without transposition, which does not occur in the other three squares; this seems to suggest, that while the first square (and therefore the first synthematic arrangement) is really of a distinct type, the other three squares (or syn-thematic arrangements) may possibly belong to one and the same type. If this were so, it would be sufficient to prove the 16-theorem (viz. the non-existence of the 16-square formula) for the first and for any one of the other three synthematic arrangements; but I provisionally assume that the four types are really distinct, and propose therefore to prove the theorem for each of the four arrangements separately.

The process is the same as for the 8-theorem; we require the tetrads 1234, &c., contained in the synthematic arrangements. In any one of these, each line gives $\frac{1}{2}8.7, = 28$ tetrads, and the 15 lines give therefore $15.28, = 420$ tetrads: but we thus obtain each tetrad 3 times, or the number of the tetrads is $420 \div 3, = 140$.

For the four arrangements respectively, these are as follows: the word “same” means same as in column I.

				I.	II.		III.		IV.		
1	2	3	4								
				5	6						
				7	8						
				9	10						
				11	12						
				13	14						
				15	16						
1	3			5	7						
				6	8						
				9	11						
				10	12						
				13	15						
				14	16						
1	4			5	8						
				6	7						
				9	12						
				10	11						
				13	16						
				14	15						
1	5			9	13	9	14	9	15	9	16
				10	14	10	13	10	16	10	15
				11	15	11	16	11	13	11	14
				12	16	12	15	12	14	12	13
1	6			9	14	9	13	9	16	9	15
				10	13	10	14	10	15	10	16
				11	16	11	15	11	14	11	13
				12	15	12	16	12	13	12	14
1	7			9	15	9	16	9	13	9	14
				10	16	10	15	10	14	10	13
				11	13	11	14	11	15	11	16
				12	14	12	13	12	16	12	15
1	8			9	16	9	15	9	14	9	13
				10	15	10	16	10	13	10	14
				11	14	11	13	11	16	11	15
				12	13	12	14	12	15	12	16

		I.		II.		III.		IV.	
2	3	5	8	same		same		same	
		6	7						
		9	12						
		10	11						
		13	16						
		14	15						
2	4	5	7	same		same		same	
		6	8						
		9	11						
		10	12						
		13	15						
		14	16						
2	5	9	14	9	13	9	16	9	15
		10	13	10	14	10	15	10	16
		11	16	11	15	11	14	11	13
		12	15	12	16	12	13	12	14
2	6	9	13	9	14	9	15	9	16
		10	14	10	13	10	16	10	15
		11	15	11	16	11	13	11	14
		12	16	12	15	12	14	12	13
2	7	9	16	9	15	9	14	9	13
		10	15	10	16	10	13	10	14
		11	14	11	13	11	16	11	15
		12	13	12	14	12	15	12	16
2	8	9	15	9	16	9	13	9	14
		10	16	10	15	10	14	10	13
		11	13	11	14	11	15	11	16
		12	14	12	13	12	16	12	15
3	4	5	6	same		same		same	
		7	8						
		9	10						
		11	12						
		13	14						
		15	16						

		I.		II.		III.		IV.	
3	5	9	15	9	16	9	13	9	14
		10	16	10	15	10	14	10	13
		11	13	11	14	11	15	11	16
		12	14	12	13	12	16	12	15
3	6	9	16	9	15	9	14	9	13
		10	15	10	16	10	13	10	14
		11	14	11	13	11	16	11	15
		12	13	12	14	12	15	12	16
3	7	9	13	9	14	9	15	9	16
		10	14	10	13	10	16	10	15
		11	15	11	16	11	13	11	14
		12	16	12	15	12	14	12	13
3	8	9	14	9	13	9	16	9	15
		10	13	10	14	10	15	10	16
		11	16	11	15	11	14	11	13
		12	15	12	16	12	13	12	14
4	5	9	16	9	15	9	14	9	13
		10	15	10	16	10	13	10	14
		11	14	11	13	11	16	11	15
		12	13	12	14	12	15	12	16
4	6	9	15	9	16	9	13	9	14
		10	16	10	15	10	14	10	13
		11	13	11	14	11	15	11	16
		12	14	12	13	12	16	12	15
4	7	9	14	9	13	9	16	9	15
		10	13	10	14	10	15	10	16
		11	16	11	15	11	14	11	13
		12	15	12	16	12	13	12	14
4	8	9	13	9	14	9	15	9	16
		10	14	10	13	10	16	10	15
		11	15	11	16	11	13	11	14
		12	16	12	15	12	14	12	13
5	6	7	8	same		same		same	
		9	10						
		11	12						
		13	14						
		15	16						

		I.	II.	III.	IV.
5	7	9 11	same	same	same
		10 12			
		13 15			
		14 16			
5	8	9 12			
		10 11			
		13 16			
		14 15			
6	7	9 12			
		10 11			
		13 16			
		14 15			
6	8	9 11			
		10 12			
		13 15			
		14 16			
7	8	9 10			
		11 12			
		13 14			
		15 16			
9	10	11 12			
		13 14			
		15 16			
9	11	13 15			
		14 16			
9	12	13 16			
		14 15			
10	11	13 16			
		14 15			
10	12	13 15			
		14 16			
11	12	13 14			
		15 16			
13	14	15 16			

39—2

308

ON THE THEOREMS OF THE 2, 4, 8, AND 16 SQUARES.

[763

As regards the signs, observe that the first line may always be written

$$ab + cd + ef + \&c.,$$

with the signs all of them +; and then writing $a, b, c, \dots = 1, 2, 3, \dots, 16$ respectively, the first line will be

$$1\ 2 + 3\ 4 + 5\ 6 + 7\ 8 + 9\ 10 + 11\ 12 + 13\ 14 + 15\ 16,$$

with the signs all of them +; that is, we may assume $e_{12}, e_{34}, \&c.$, or say

$$1\ 2, 3\ 4, 5\ 6, 7\ 8, 9\ 10, 11\ 12, 13\ 14, 15\ 16,$$

all of them = +1. And in the other lines, the signs of all the terms of any line may be reversed at pleasure, that is, we may assume $e_{13}, e_{14}, \&c.$, or say $1\ 3, 1\ 4, 1\ 5, 1\ 6, 1\ 7, 1\ 8, 1\ 9, 1\ 10, 1\ 11, 1\ 12, 1\ 13, 1\ 14, 1\ 15, 1\ 16$, all of them = +1.

Making these assumptions, then for any one of the synthematic arrangements the several tetrads give as before relations between the signs; among these are included the results already obtained for the 8-question, and taking as before

$$-a = b = c = d = \theta,$$

we have the signs of the several terms belonging to the 8-question given as $= +\theta$ or $+\beta$, as before. The remaining tetrads up to 1 8 12 13 then serve to express all the remaining signs in terms of the as yet undetermined signs $e, f, g, h, i, j, k, l, m, n, o, p, q, r, s, t, u, v, w, x, y, z, \alpha, \beta$, for instance

$$1\ 3.9\ 11 + 1.e,$$

$$1\ 9.3\ 11 - 1.e,$$

$$1\ 11.3\ 9 + 1.e,$$

that is, $3\ 9 = e$, $3\ 11 = -e$, $9\ 11 = e$; and then the tetrads up to 2 8 9 15 serve to express these signs in terms of the undetermined signs $\lambda, \mu, \nu, \rho, \sigma, \tau$; for instance

$$2\ 3.9\ 12 + 1.i,$$

$$2\ 9.3\ 12 - 1.f,$$

$$2\ 12.3\ 9 + -1.e,$$

that is, $-e = f = i$; and in like manner 2 3 10 11, 2 4 9 11 and 2 4 10 12 give respectively $-e = f = j$, $-e = i = j$, $f = i = j$; that is, we have $-e = f = i = j = \lambda$ suppose. And in this way we have, for each of the four synthematic arrangements the signs of all the terms expressed in terms of the undetermined signs $\theta, \lambda, \mu, \nu, \rho, \sigma, \tau$, as shown in the following table; where observe that the results apply to the four synthematic arrangements separately, viz. the $e, f, g, \dots$ &c., and the $\theta, \lambda, \mu, \nu, \rho, \sigma, \tau$ in each column are altogether independent of the like symbols in the other three columns.

Signs for the four synthematic arrangements:

		I.	II.	III.	IV.
1	2	+1	same	same	same
	3	+1			
	4	+1			
	5	+1			
	6	+1			
	7	+1			
	8	+1			
	9	+1			
	10	+1			
	11	+1			
	12	+1			
	13	+1			
	14	+1			
	15	+1			
	16	+1			
2	3	+1			
	4	−1			
	5	+1			
	6	−1			
	7	+1			
	8	−1			
	9	+1			
	10	−1			
	11	+1			
	12	−1			
	13	+1			
	14	−1			
	15	+1			
	16	−1			
3	4	+1		same	same
	5	− θ			
	6	θ			
	7	θ			
	8	− θ			
	9	$e - \lambda$			
	10	$f \lambda$			
	11	− $e \lambda$			
	12	− $f - \lambda$			
	13	$g - \mu$			
	14	$h \mu$			
	15	− $g \mu$			
	16	− $h - \mu$			

		I.	II.	III.	IV.
4	5		same	same	same
	6	θ			
	7	$-\theta$			
	8	$-\theta$			
	9	$i \lambda$			
	10	$j \lambda$			
	11	$-j - \lambda$			
	12	$-i - \lambda$			
	13	$k \mu$			
	14	$l \mu$			
	15	$-l - \mu$			
	16	$-k - \mu$			
5	6	+1	+1	+1	+1
	7	$-\theta$	$-\theta$	$-\theta$	$-\theta$
	8	θ			
	9	$m - \nu$	$m \nu$	$m - \nu$	$m \nu$
	10	$n \nu$	$n \nu$	$n \nu$	$n \nu$
	11	$o - \rho$	$o \rho$	$o - \rho$	$o \rho$
	12	$p \rho$	$p \rho$	$p \rho$	$p \rho$
	13	$-m \nu$	$-n - \nu$	$-o \rho$	$-p - \rho$
	14	$-n - \nu$	$-m - \nu$	$-p - \rho$	$-o - \rho$
	15	$-o \rho$	$-p - \rho$	$-m \nu$	$-n - \nu$
	16	$-p - \rho$	$-o - \rho$	$-n - \nu$	$-m - \nu$
6	7			θ	
	8	θ			
	9	$q \nu$	$q \nu$	$q \nu$	$q \nu$
	10	$r \nu$	$r - \nu$	$r \nu$	$r - \nu$
	11	$s \rho$	$s \rho$	$s \rho$	$s \rho$
	12	$t \rho$	$t - \rho$	$t \rho$	$t - \rho$
	13	$-r - \nu$	$-q - \nu$	$-t - \rho$	$-s - \rho$
	14	$-q - \nu$	$-r \nu$	$-s - \rho$	$-t \rho$
	15	$-t - \rho$	$-s - \rho$	$-r - \nu$	$-q - \nu$
	16	$-s - \rho$	$-t \rho$	$-q - \nu$	$-r \nu$
7	8	+1	+1	+1	+1
	9	$u - \sigma$	$u - \sigma$	$u - \sigma$	$u \sigma$
	10	$v \sigma$	$v \sigma$	$v \sigma$	$v \sigma$
	11	$w - \tau$	$w \tau$	$w - \tau$	$w \tau$
	12	$x \tau$	$x \tau$	$x \tau$	$x \tau$
	13	$-w \tau$	$-x - \tau$	$-u \sigma$	$-v - \sigma$
	14	$-x - \tau$	$-w - \tau$	$-v - \sigma$	$-u - \sigma$
	15	$-u \sigma$	$-v - \sigma$	$-w \tau$	$-x - \tau$
	16	$-v - \sigma$	$-u - \sigma$	$-x - \tau$	$-w - \tau$

		I.	II.	III.	IV.
8	9	$y \sigma$	$y \sigma$	$y \sigma$	$y \sigma$
	10	$z \sigma$	$z - \sigma$	$z \sigma$	$z - \sigma$
	11	$\alpha \tau$	$\alpha \tau$	$\alpha \tau$	$\alpha \tau$
	12	$\beta \tau$	$\beta - \tau$	$\beta \tau$	$\beta - \tau$
	13	$-\beta - \tau$	$-\alpha - \tau$	$-z - \sigma$	$-y - \sigma$
	14	$-\alpha - \tau$	$-\beta \tau$	$-y - \sigma$	$-z \sigma$
	15	$-z - \sigma$	$-y - \sigma$	$-\beta - \tau$	$-\alpha - \tau$
	16	$-y - \sigma$	$-z \sigma$	$-\alpha - \tau$	$-\beta \tau$
9	10	+1	+1	+1	+1
	11	$e - \lambda$	$e - \lambda$	$e - \lambda$	$e - \lambda$
	12	$i \lambda$	$i \lambda$	$i \lambda$	$i \lambda$
	13	$m - \nu$	$q \nu$	$u - \sigma$	$y \sigma$
	14	$q \nu$	$m \nu$	$y \sigma$	$u \sigma$
	15	$u - \sigma$	$y \sigma$	$m - \nu$	$q \nu$
	16	$y \sigma$	$u \sigma$	$q \nu$	$m \nu$
10	11	$j \lambda$	$j \lambda$	$j \lambda$	$j \lambda$
	12	$f \lambda$	$f \lambda$	$f \lambda$	$f \lambda$
	13	$r \nu$	$n \nu$	$z \sigma$	$v \sigma$
	14	$n \nu$	$r - \nu$	$v \sigma$	$z - \sigma$
	15	$z \sigma$	$v \sigma$	$r \nu$	$n \nu$
	16	$v \sigma$	$z - \sigma$	$n \nu$	$r - \nu$
11	12	+1	+1	+1	+1
	13	$w - \tau$	$s \tau$	$o - \rho$	$s \rho$
	14	$s \tau$	$w \tau$	$s \rho$	$o \rho$
	15	$o - \rho$	$s \rho$	$w - \tau$	$s \tau$
	16	$s \rho$	$o \rho$	$s \tau$	$w \tau$
12	13	$x \tau$	$x \tau$	$t \rho$	$p \rho$
	14	$x \tau$	$\beta - \tau$	$p \rho$	$t - \rho$
	15	$t \rho$	$p \rho$	$p \tau$	$x \tau$
	16	$p \rho$	$t - \rho$	$x \tau$	$p - \tau$
13	14	+1	+1	+1	+1
	15	$g - \mu$	$g - \mu$	$g - \mu$	$g - \mu$
	16	$k \mu$	$k \mu$	$k \mu$	$k \mu$
14	15	$l \mu$	$l \mu$	$l \mu$	$l \mu$
	16	$h \mu$	$h \mu$	$h \mu$	$h \mu$
15	16	+1	+1	+1	+1

312

ON THE THEOREMS OF THE 2, 4, 8, AND 16 SQUARES.

[763

We have now for the four arrangements respectively, by means of hitherto unused tetrads, the following determinations of sign: these being in each case inconsistent with each other.

First arrangement.

$$\begin{array}{ll}
 3 \ 5 \ 9 \ 15 + -\theta. - \sigma & \text{that is,} \\
 3 \ 9 \ 5 \ 15 - -\lambda. \rho & \theta \sigma = -\lambda \rho = -\mu \nu, \\
 3 \ 15 \ 5 \ 9 + -\mu. - \nu &
 \end{array}$$

$$\begin{array}{ll}
 3 \ 5 \ 10 \ 16 + -\theta. \sigma & \\
 3 \ 10 \ 5 \ 16 - -\lambda. - \rho & -\theta \sigma = \lambda \rho = -\mu \nu. \\
 3 \ 16 \ 5 \ 10 + -\mu. \nu &
 \end{array}$$

$$\begin{array}{ll}
 3\ 5\ 11\ 13 + -\theta. -\tau & \\
 3\ 11\ 5\ 13 - -\lambda. \nu & \theta\tau = -\lambda\nu = \mu\rho, \\
 3\ 13\ 5\ 11 + -\mu. -\rho &
 \end{array}$$

$$\begin{array}{ll}
 3\ 5\ 12\ 14 + -\theta. \tau & \\
 3\ 12\ 5\ 14 - -\lambda. -\nu & -\theta\tau = -\lambda\nu = \mu\rho. \\
 3\ 14\ 5\ 12 + \mu. \rho &
 \end{array}$$

Second arrangement.

$$\begin{array}{ll}
 3\ 5\ 9\ 16 + -\theta. \sigma & \\
 3\ 9\ 5\ 16 - -\lambda. -\rho & \theta\sigma = \lambda\rho = \mu\nu, \\
 3\ 16\ 5\ 9 + -\mu. \nu &
 \end{array}$$

$$\begin{array}{ll}
 3\ 5\ 10\ 15 + -\theta. \sigma & \\
 3\ 10\ 5\ 15 - -\lambda. -\rho & -\theta\sigma = \lambda\rho = \mu\nu, \\
 3\ 15\ 5\ 10 + \mu. \nu &
 \end{array}$$

$$\begin{array}{ll}
 3\ 5\ 11\ 14 + -\theta. \tau & \\
 3\ 11\ 5\ 14 - -\lambda. -\nu & -\theta\tau = \lambda\nu = \mu\rho, \\
 3\ 14\ 5\ 11 + \mu. \rho &
 \end{array}$$

$$\begin{array}{ll}
 3\ 5\ 12\ 13 + -\theta. \tau & \\
 3\ 12\ 5\ 13 - -\lambda. -\nu & \theta\tau = \lambda\nu = \mu\rho. \\
 3\ 13\ 5\ 12 + -\mu. \rho &
 \end{array}$$

763]

ON THE THEOREMS OF THE 2, 4, 8, AND 16 SQUARES.

313

Third arrangement.

$$\begin{array}{ll}
 3\ 5\ 9\ 13 + -\theta. -\sigma & \\
 3\ 9\ 5\ 13 - -\lambda. \rho & \theta\sigma = -\lambda\rho = \mu\nu, \\
 3\ 13\ 5\ 9 + -\mu. -\nu &
 \end{array}$$

$$\begin{array}{ll}
 3\ 5\ 10\ 14 + -\theta. \sigma & \\
 3\ 10\ 5\ 14 - -\lambda. -\rho & \theta\sigma = -\lambda\rho = -\mu\nu, \\
 3\ 14\ 5\ 10 + \mu. \nu &
 \end{array}$$

$$\begin{array}{ll}
 3\ 5\ 11\ 15 + -\theta. -\tau & \\
 3\ 11\ 5\ 15 - -\lambda. \nu & \theta\tau = -\lambda\nu = -\mu\rho. \\
 3\ 15\ 5\ 11 + \mu. -\rho &
 \end{array}$$

$3\ 5\ 12\ 16 + -\theta.\ \tau$

$3\ 12\ 5\ 16 - -\lambda.\ -\nu$

$3\ 16\ 5\ 12 + -\mu.\ \rho$

$\theta\tau = -\lambda\nu = \mu\rho.$

Fourth arrangement.

$3\ 5\ 9\ 14 + -\theta.\ \sigma$

$3\ 9\ 5\ 14 - -\lambda.\ -\rho$

$3\ 14\ 5\ 9 + \mu.\ \nu$

$\theta\sigma = \lambda\rho = -\mu\nu,$

$3\ 5\ 10\ 13 + -\theta.\ \sigma$

$3\ 10\ 5\ 13 - -\lambda.\ -\rho$

$3\ 13\ 5\ 10 + -\mu.\ \nu$

$\theta\sigma = -\lambda\rho = \mu\nu,$

$3\ 5\ 11\ 16 + -\theta.\ \tau$

$3\ 11\ 5\ 16 - -\lambda.\ -\nu$

$3\ 16\ 5\ 11 + -\mu.\ \rho$

$\theta\tau = -\lambda\nu = \mu\rho,$

$3\ 5\ 12\ 15 + -\theta.\ -\tau$

$3\ 12\ 5\ 15 - -\lambda.\ -\nu$

$3\ 15\ 5\ 12 + -\mu.\ \rho$

$\theta\tau = -\lambda\nu = -\mu\rho.$

And it hence finally appears, that we cannot, in any one of the four arrange-ments, determine the signs so as to give rise to a 16-square theorem; that is, the product of a sum of 16 squares into a sum of 16 squares cannot be made equal to a sum of 16 squares.

C. XI.

40

314

[764

764.

THE BINOMIAL EQUATION $x^n - 1 = 0$: QUINQUESECTION.

[From the *Proceedings of the London Mathematical Society*, vol. XII. (1881), pp. 15, 16.
Read December 9, 1880.]

THE theory should be precisely analogous to those for the trisection and quarti-section (see my paper, “The Binomial Equation $x^n - 1 = 0$, Trisection and Quartisection,” *Proceedings of the London Mathematical Society*, vol. XI. (1879), pp. 4–17, [731]), only I have not been able to carry it so far. We have in the present case five periods X, Y, Z, W, T , the actual expressions for which, $X = \eta^a + \dots$, $Y = \eta^b + \dots$, etc., with Reuschle’s selected prime root g , can be (for the primes $5n + 1$ under 100) at once written down by means of the table given, pp. 16, 17, of that paper; [see this volume, pp. 95, 96]. The relations between the periods are of the form

	X	Y	Z	W	T
$X^2 =$	a	b	c	d	e
$XY =$	f	g	h	i	j
$XZ =$	k	l	m	n	o

that is, we have

$$X^2 = (a, b, c, d, e \mid X, Y, Z, W, T),$$

and thence, by cyclical permutations,

$$Y^2 = (e, a, b, c, d \mid \quad), \text{ etc.};$$

viz. from the value of X^2 we have those of Y^2, Z^2, W^2, T^2 ; from the value of XY those of YZ, ZW, WT, TX ; and from the value of XZ those of YW, ZT, WX, TY .

764] THE BINOMIAL EQUATION $x^n - 1 = 0$: QUINQUESECTION. 315

From the equation $X + Y + Z + W + T = -1$, multiplying by X and then substituting for X^2, XY , &c., their values, we obtain

$$\begin{aligned} -a &= 1 + f + k + m + g, \\ -b &= \quad g + l + n + h, \\ -c &= \quad h + m + o + i, \\ -d &= \quad i + n + k + j, \\ -e &= \quad j + o + l + f, \end{aligned}$$

which determine (a, b, c, d, e) in terms of (f, g, h, i, j) and (k, l, m, n, o) . It is, moreover, easy to prove that

$$\begin{aligned} f + g + h + i + j &= \frac{1}{5}(p - 1), \\ k + l + m + n + o &= \frac{1}{5}(p - 1), \end{aligned}$$

whence also

$$a + b + c + d + e = -1 - \frac{2}{5}(p - 1).$$

We obtain other relations between the coefficients by considering the two triple products XYZ and XYW : these are all that need be considered, since the other triple products are deducible from them by cyclical permutations. From the first of these we have

$$X.YZ = Y.XZ = Z.XY,$$

and from the second

$$X.YW = Y.XW = W.XY;$$

and if we herein substitute for YZ, XZ , &c., their values, and then in the resulting equations for X^2, XY , &c., their values as linear functions of X, Y, Z, W, T , we obtain in all $5.2.2 = 20$ quadric relations between the 15 coefficients; or if we substitute for (a, b, c, d, e) their foregoing values, in all 20 relations between the 10 coefficients (f, g, h, i, j) and (k, l, m, n, o) . These are at most equivalent to 8 independent equations, since we have, besides, the sums $f + g + h + i + j$ and $k + l + m + n + o$ each $= \frac{1}{5}(p - 1)$; but I have not succeeded in finding the connexions between them, or even in ascertaining to how many independent equations they are equivalent.

For any given prime $p = 5n + 1$, the values of the coefficients, and also the coefficients of the quintic equation for the periods, could of course be calculated directly from the expressions of the periods; but for the primes under 100, that is, for the values 11, 31, 41, 61, 71, they are at once obtained from Reuschle. We have thus the two Tables, the former giving the coefficients $a, b, \dots, n, o$, and the latter the coefficients of the quintic equations.

316 THE BINOMIAL EQUATION $x^n - 1 = 0$: QUINQUESECTION. [764

TABLE 1.

TABLE 2 of the Quintic Equations.

p	a	b	c	d	e
	f	g	h	i	j
	k	l	m	n	o
11	−2	−1	−2	−2	−2
	1	0	0	1	0
	0	0	0	1	1
31	−4	−6	−6	−4	−5
	0	1	2	1	2
	2	2	2	1	1
41	−8	−5	−6	−6	−8
	3	1	2	1	2
	2	2	2	1	1
61	−10	−9	−12	−8	−10
	3	2	2	3	2
	2	2	4	3	3
71	−14	−10	−12	−9	−12
	4	2	3	2	3
	2	3	5	2	2

Coefficients of						
p	x^5	x^4	x^3	x^2	x	$1 = 0$
11	1	1	−4	−3	+3	+1
31	1	1	−12	−2	+1	+5
41	1	1	−16	+5	+21	−9
61	1	1	−24	−17	+41	−23
71	1	1	−28	+37	+25	+1

765]

317

765.

ON THE FLEXURE AND EQUILIBRIUM OF A SKEW SURFACE.

[From the *Proceedings of the London Mathematical Society*, vol. XII. (1881), pp. 103–108.
Read March 10, 1881.]

THE skew surface is taken to be such that the strip between two consecutive generating lines is rigid, and that the flexure takes place by the rotation of the strips about the generating lines successively. The theory of the flexure is well known, but I am not aware that the theory of the equilibrium of such a surface, when acted upon by any given forces, has been considered; it is, however, a question which presents itself naturally in connexion with those relating to other continuous bodies treated of in the *Mécanique Analytique*, and forms a good example of the principles made use of.

To begin with the mechanical theory: we may regard the forces as acting on the generating lines regarded as material lines; and if for an element of mass dm , coordinates (x, y, z) of a particular generating line G , the forces parallel to the axes are X', Y', Z' , then the corresponding term in the equation of equilibrium is

$$S(X'\delta x + Y'\delta y + Z'\delta z) dm;$$

and observing that there are (as will afterwards appear) five geometrical conditions, which I represent by $U_1 = 0, U_2 = 0, \dots, U_5 = 0$, the equation of equilibrium is

$$S\{(X'\delta x + Y'\delta y + Z'\delta z) dm + T_1\delta U_1 + T_2\delta U_2 + T_3\delta U_3 + T_4\delta U_4 + T_5\delta U_5\} = 0,$$

where $T_1, T_2, \dots, T_5$ are the indeterminate multipliers, representing colligation-forces which correspond to the five geometrical conditions respectively.

Taking (ξ, η, ζ) for the coordinates of a particular point P on the generating line; p, q, r for the cos-inclinations of the line (whence $U_1 = p^2 + q^2 + r^2 - 1 = 0$ is one of the geometrical relations), and ρ for the distance of dm from P , we have

$$x, y, z = \xi + \rho p, \quad \eta + \rho q, \quad \zeta + \rho r,$$

$$\delta x, \delta y, \delta z = \delta \xi + \rho \delta p, \quad \delta \eta + \rho \delta q, \quad \delta \zeta + \rho \delta r.$$

The summation S extends first to the different points of the generating line, and then to the different generating lines; applying it first to the particular generating line, we write

$$\begin{array}{cccccc} SX'dm, & SY'dm, & SZ'dm, & SX'\rho dm, & SY'\rho dm, & SZ'\rho dm \\ = X, & Y, & Z, & L, & M, & N, \end{array}$$

where X, Y, Z are the whole forces, and L, M, N the whole moments about the point P , for the generating line G ; retaining the same summatory symbol S , as now referring to the different generating lines, the equation becomes

$$S\{X\delta\xi + Y\delta\eta + Z\delta\zeta + L\delta p + M\delta q + N\delta r + T_1\delta U_1 + \dots + T_5\delta U_5\} = 0.$$

We have now to consider the geometrical theory of the flexure. Taking on the skew surface an arbitrary curve cutting each generating line G in a point P , coordinates (ξ, η, ζ) , and taking σ for the distance along the curve of the point P from a fixed point of the curve; also p, q, r , as before, for the cos-inclinations of the generating line G , then when the surface is in a determinate state, $\xi, \eta, \zeta, p, q, r$ are given functions of σ : but these functions vary with the flexure of the surface, with, however, certain relations unaffected by the flexure; and the problem is to find first these relations. As already mentioned, one of them is $p^2 + q^2 + r^2 - 1 = 0$.

Taking P' as the consecutive point on the curve, so that the direction of the element PP' is that of the tangent PT at P , it is convenient to write l, m, n for the cosine-inclinations of the tangent; we have, it is clear,

$$l, m, n = \frac{d\xi}{d\sigma}, \frac{d\eta}{d\sigma}, \frac{d\zeta}{d\sigma}; \quad l^2 + m^2 + n^2 - 1 = 0.$$


The conditions in order to the rigidity of the strip, are that the angles GPP' , $G'P'P$ ($= 180^\circ - G'P'T$), and the inclination $G'P'$ to GP , shall have given values, variable it may be from strip to strip—that is, these values must be given functions of σ . Taking $\angle GPT = I$, the value of $G'P'T$ can differ only infinitesimally from that of GPT , and we take it to be $G'P'T = I - \Omega d\sigma$; also the inclination GP to $G'P'$ is an infinitesimal, $= \Theta d\sigma$: we have I, Ω, Θ given functions of σ : It is to be remarked that these conditions imply, inclination of $G'P'$ to tangent plane GPT at P has a given value $\Lambda d\sigma$; in fact, if through P we draw a line $P\gamma$ parallel to $P'G'$, then, if P is regarded as the centre of a sphere which meets $PG, P\gamma, PT$ in the

765]

ON THE FLEXURE AND EQUILIBRIUM OF A SKEW SURFACE.

319

points g, g', t respectively, we have a spherical triangle $gg't$, the sides of which are $I - \Omega d\sigma$, I , and $\Theta d\sigma$, and of which the perpendicular $g'm$ is $= \Lambda d\sigma$: we have thus



an infinitesimal right-angled triangle, the base and altitude of which are $\Omega d\sigma, \Lambda d\sigma$, and the hypotenuse is $\Theta d\sigma$; whence $\Theta^2 = \Omega^2 + \Lambda^2$. In the case of the developable surface $\Lambda = 0$ and $\Theta = \Omega$. It may be remarked that, when the curve on the skew surface is the line of striction, we have $\Omega = 0$; in fact, taking P to be on the line of striction, the line

$$\frac{X - \xi}{qr' - q'r} = \frac{Y - \eta}{rp' - r'p} = \frac{Z - \zeta}{pq' - p'q}$$

through (ξ, η, ζ) at right angles to the two generating lines, meets the consecutive generating line $X, Y, Z = \xi' + \rho p', \eta' + \rho q', \zeta' + \rho r'$; and the condition that this may be so is easily found to be $\Omega = 0$.

Take, for a moment, p', q', r' for the cos-inclinations of the consecutive generating line $P'G'$; we have

$$\begin{aligned} lp + mq + nr &= \cos I, \\ lp' + mq' + nr' &= \cos(I - \Omega d\sigma), \\ pp' + qq' + rr' &= \cos \Theta d\sigma; \end{aligned}$$

and then writing $p', q', r' = p + dp, q + dq, r + dr$, and observing that the equation $p'^2 + q'^2 + r'^2 = 1$ gives

$$p dp + q dq + r dr = -\frac{1}{2}(dp^2 + dq^2 + dr^2),$$

these equations and the before-mentioned two equations become

$$\begin{aligned} (U_1) \quad & p^2 + q^2 + r^2 - 1 = 0, \\ (U_2) \quad & l^2 + m^2 + n^2 - 1 = 0, \\ (U_3) \quad & lp + mq + nr - \cos I = 0, \\ (U_4) \quad & l dp + m dq + n dr - \Omega \sin I d\sigma = 0, \\ (U_5) \quad & dp^2 + dq^2 + dr^2 - \Theta^2 d\sigma^2 = 0, \end{aligned}$$

which equations, considering therein l, m, n as standing for their values $\frac{d\xi}{d\sigma}, \frac{d\eta}{d\sigma}, \frac{d\zeta}{d\sigma}$, are the geometrical relations which connect the six variables $\xi, \eta, \zeta, p, q, r$, considered as functions of σ . And in these equations I, Ω, Θ denote given functions of σ , invariable by any flexure of the surface.

To complete the geometrical theory, it is to be observed that we can by flexure bring the generating lines of the surface to be parallel to those of any given cone

320

ON THE FLEXURE AND EQUILIBRIUM OF A SKEW SURFACE.

[765

$C(p, q, r) = 0$, where $C(p, q, r)$ denotes a homogeneous function of (p, q, r) . Hence, joining to the foregoing five equations this new equation

$$C(p, q, r) = 0,$$

these six equations determine $\xi, \eta, \zeta, p, q, r$ as functions of σ . To make the solution completely determinate, we have only to assume for the point P , which corresponds, say, to the value $\sigma = 0$, a position in space at pleasure, and to take the corresponding generating line PG parallel to a generating line, at pleasure, of the cone.

As an example, writing γ to denote an arbitrary constant angle, if the invariable conditions are

$$I = \gamma, \quad \Omega = \sin \gamma, \quad \Theta = 0,$$

then the five equations are

$$\begin{aligned} p^2 + q^2 + r^2 - 1 &= 0, \\ l^2 + m^2 + n^2 - 1 &= 0, \\ lp + mq + nr - \cos \gamma &= 0, \\ dp^2 + dq^2 + dr^2 - \sin^2 \gamma d\sigma^2 &= 0, \\ l dp + m dq + n dr &= 0. \end{aligned}$$

We assume first

$$C(p, q, r) = p^2 + q^2 - r^2 \tan^2 \gamma, = 0;$$

and secondly

$$C(p, q, r) = r, = 0.$$

Then, in the former case, we find the solution

$$p, q, r = -\sin \gamma \sin \sigma, \sin \gamma \cos \sigma, \cos \gamma$$

$$\xi, \eta, \zeta = \cos \sigma, \sin \sigma, 0;$$

giving

$$x, y, z = \cos \sigma - \rho \sin \gamma \sin \sigma, \sin \sigma + \rho \sin \gamma \cos \sigma, \rho \cos \gamma;$$

and consequently

$$x^2 + y^2 - \rho^2 \tan^2 \gamma = 0,$$

the hyperboloid of revolution. And, in the latter case,

$$\begin{aligned} p, q, r &= \cos(\sigma \sin \gamma), \sin(\sigma \sin \gamma), 0, \\ \xi, \eta, \zeta &= \cot \gamma \sin(\sigma \sin \gamma), -\cot \gamma \cos(\sigma \sin \gamma), \sigma \sin \gamma, \end{aligned}$$

that is,

$$x, y = \cot \gamma \sin z + \rho \cos z, -\cot \gamma \cos z + \rho \sin z,$$

whence

$$x \sin z - y \cos z = \cot \gamma,$$

a skew helicoid generated by horizontal tangents of the cylinder $x^2 + y^2 = \cot^2 \gamma$. This is a known deformation of the hyperboloid.

765]

ON THE FLEXURE AND EQUILIBRIUM OF A SKEW SURFACE.

321

Returning now to the mechanical problem, we have to consider the terms

$$\begin{aligned} &S \cdot T_1 \frac{1}{2} \delta (p^2 + q^2 + r^2 - 1) \\ &+ T_2 \frac{1}{2} \delta (l^2 + m^2 + n^2 - 1) \\ &+ T_3 \delta (lp + mq + nr - \cos I) \\ &+ T_4 \delta \left(l \frac{dp}{d\sigma} + m \frac{dq}{d\sigma} + n \frac{dr}{d\sigma} - \Omega \sin I \right) \\ &+ T_5 \frac{1}{2} \delta \left\{ \left(\frac{dp}{d\sigma} \right)^2 + \left(\frac{dq}{d\sigma} \right)^2 + \left(\frac{dr}{d\sigma} \right)^2 - \Theta^2 \right\}. \end{aligned}$$

The first term gives, under the sign S ,

$$T_1(p\delta p + q\delta q + r\delta r). \quad (*)$$

The second term gives, in the first instance,

$$\frac{T_2}{d\sigma} (l d\delta\xi + m d\delta\eta + n d\delta\zeta);$$

or, since in general

$$S l d\delta\xi = l^{(n)} \delta\xi^{(n)} - l^{(0)} \delta\xi^{(0)} + S^*(-dl \cdot \delta\xi),$$

then, attending only to the terms under the sign S^* , these are

$$= -\frac{d}{d\sigma} T_2 l \cdot \delta\xi - \frac{d}{d\sigma} T_2 m \cdot \delta\eta - \frac{d}{d\sigma} T_2 n \cdot \delta\zeta. \quad (*)$$

The third term gives

$$\begin{aligned} &T_3(l\delta p + m\delta q + n\delta r) \\ &+ T_3(p\delta l + q\delta m + r\delta n), \end{aligned} \quad (*)$$

where the second line,

$$= \frac{T_3}{d\sigma} (p d\delta\xi + q d\delta\eta + r d\delta\zeta),$$

attending only to the terms under the sign S , gives

$$-\frac{d}{d\sigma} T_3 p \cdot \delta\xi - \frac{d}{d\sigma} T_3 q \cdot \delta\eta - \frac{d}{d\sigma} T_3 r \cdot \delta\zeta. \quad (*)$$

The fourth term gives

$$\frac{T_4}{d\sigma} (l d\delta p + m d\delta q + n d\delta r) + T_4 \left(\frac{dp}{d\sigma} \delta l + \frac{dq}{d\sigma} \delta m + \frac{dr}{d\sigma} \delta n \right),$$

where the first line, written under the form

$$\frac{1}{d\sigma} \{d(T_4 l \delta p) - dT_4 l \cdot \delta p - T_4 dl \cdot \delta p\} + \text{(similar for } m, n),$$

and attending only to the terms under the sign S , gives

$$-\frac{d}{d\sigma} T_4 l \cdot \delta p - \frac{d}{d\sigma} T_4 m \cdot \delta q - \frac{d}{d\sigma} T_4 n \cdot \delta r,$$

and the second line, attending in like manner only to the terms under the sign S , gives

$$-\frac{d}{d\sigma}T_4l \cdot \delta p - \frac{d}{d\sigma}T_4m \cdot \delta q - \frac{d}{d\sigma}T_4n \cdot \delta r. \quad (*)$$

The fifth term, written under the form

$$\frac{T_5}{d\sigma} \left(\frac{dp}{d\sigma} d\delta p + \frac{dq}{d\sigma} d\delta q + \frac{dr}{d\sigma} d\delta r \right),$$

and attending only to the terms under the sign S , gives

$$-\frac{d}{d\sigma}T_5 \frac{dp}{d\sigma} \cdot \delta p - \frac{d}{d\sigma}T_5 \frac{dq}{d\sigma} \cdot \delta q - \frac{d}{d\sigma}T_5 \frac{dr}{d\sigma} \cdot \delta r; \quad (*)$$

where in each case I have marked with an asterisk the lines which present them-selves in the final result.

Hence, joining to the foregoing the force-terms

$$X\delta\xi + Y\delta\eta + Z\delta\zeta + L\delta p + M\delta q + N\delta r, \quad (*)$$

and equating to zero the coefficients of $\delta\xi, \delta\eta, \delta\zeta, \delta p, \delta q, \delta r$ respectively, we have

$$\begin{aligned} 0 &= X - \frac{d}{d\sigma}T_2l - \frac{d}{d\sigma}T_3p, \\ 0 &= Y - \frac{d}{d\sigma}T_2m - \frac{d}{d\sigma}T_3q, \\ 0 &= Z - \frac{d}{d\sigma}T_2n - \frac{d}{d\sigma}T_3r, \\ 0 &= L + T_1p + T_3l - \frac{d}{d\sigma}T_4l - \frac{d}{d\sigma}T_5 \frac{dp}{d\sigma}, \\ 0 &= M + T_1q + T_3m - \frac{d}{d\sigma}T_4m - \frac{d}{d\sigma}T_5 \frac{dq}{d\sigma}, \\ 0 &= N + T_1r + T_3n - \frac{d}{d\sigma}T_4n - \frac{d}{d\sigma}T_5 \frac{dr}{d\sigma}, \end{aligned}$$

where it will be recollected that l, m, n stand for $\frac{d\xi}{d\sigma}, \frac{d\eta}{d\sigma}, \frac{d\zeta}{d\sigma}$, the variables being $\xi, \eta, \zeta, p, q, r$, and σ . The elimination of $T_1, T_2, \dots, T_5$ from the six equations should lead to a relation between $\xi, \eta, \zeta, p, q, r$, which, with the foregoing five relations, would determine the six variables $\xi, \eta, \zeta, p, q, r$ in terms of σ .

In particular, the forces and moments X, Y, Z, L, M, N may all of them vanish; assuming that $T_1, T_2, \dots, T_5$ do not all of them vanish, we still have the sixth relation, which (with the foregoing five

relations) determines $\xi, \eta, \zeta, p, q, r$ in terms of σ ; and it is to be remarked that the problem in question, of the figure of equilibrium of the skew surface not acted upon by any forces, is analogous to that of the geodesic line in space; only whilst here the solution is, curve a straight line, the solution for the case of the skew surface depends upon equations of a complex enough form; in the case of the developable surface, the required figure is of course the plane.

766]

323

766.

ON THE GEODESIC CURVATURE OF A CURVE ON A SURFACE.

[From the *Proceedings of the London Mathematical Society*, vol. XII. (1881), pp. 110–117.

Read April 14, 1881.]

THERE is contained in Liouville's Note II. to his edition of Monge's *Application de l'Analyse à la Géométrie* (Paris, 1850), see pp. 574 and 575, the following formula,

$$\frac{di}{ds} = -\frac{1}{2G\sqrt{E}} \frac{dG}{du} \cos i + \frac{1}{2E\sqrt{G}} \frac{dE}{dv} \sin i,$$

$$\frac{di}{ds} - \frac{\cos i}{\rho_u} + \frac{\sin i}{\rho_v},$$

which gives the radius of geodesic curvature of a curve upon a surface when the position of a point on the surface is defined by the parameters u, v , belonging to a system of orthotomic curves; or, what is the same thing, such that

$$ds^2 = E du^2 + G dv^2.$$

Writing with Gauss p, q instead of u, v , I propose to obtain the corresponding formula in the general case where the parameters p, q are such that

$$ds^2 = E dp^2 + 2F dp dq + G dq^2.$$

I call to mind that, if PQ, PQ' are equal infinitesimal arcs on the given curve and on its tangent geodesic, then the radius of geodesic curvature ρ is, by definition, a length ρ such that $2\rho \cdot QQ' = PQ^2$. More generally, if the curves on the surface are any two curves which touch each other, then ρ as thus determined is the radius of relative curvature of the two curves.

41—2

324

ON THE GEODESIC CURVATURE OF A CURVE ON A SURFACE.

[766

The notation is that of the Memoir, "Disquisitiones generales circa superficies curvas" (1827), GAUSS, *Werke*, t. III.; see also my paper "On geodesic lines, in particular those of a quadric surface," *Proc. Lond. Math. Society*, t. IV. (1872), pp. 191–211, [508]; and Salmon's *Solid Geometry*, 3rd ed., 1874, pp. 251 et seq.

The coordinates (x, y, z) of a point on the surface are taken to be functions of two independent parameters p, q ; and we then write

$$dx + \frac{1}{2}d^2x = a dp + a' dq + \frac{1}{2}(\alpha dp^2 + 2\alpha' dp dq + \alpha'' dq^2),$$

$$dy + \frac{1}{2}d^2y = b dp + b' dq + \frac{1}{2}(\beta dp^2 + 2\beta' dp dq + \beta'' dq^2),$$

$$dz + \frac{1}{2}d^2z = c dp + c' dq + \frac{1}{2}(\gamma dp^2 + 2\gamma' dp dq + \gamma'' dq^2) :$$

$$E, F, G = a^2 + b^2 + c^2, \quad aa' + bb' + cc', \quad a'^2 + b'^2 + c'^2; \quad V^2 = EG - F^2;$$

and therefore

$$ds^2 = E dp^2 + 2F dp dq + G dq^2,$$

where E, F, G are regarded as given functions of p and q .

To determine a curve on the surface, we establish a relation between the two parameters p, q , or, what is the same thing, take p, q to be functions of a single parameter θ ; and we write as usual p', p'', q' , etc., to denote the differential coefficients of p, q , etc., in regard to θ ; we write also E_1, E_2 , etc., to denote the differential coefficients $\frac{dE}{dp}, \frac{dE}{dq}$, etc. In the first instance, θ is taken to be an arbitrary parameter, but we afterwards take it to be the length s of the curve from a fixed point thereof.

First formula for the radius of relative curvature.

Consider any two curves touching at the point P , coordinates (x, y, z) which are regarded as given functions of (p, q) ; where (p, q) are for the one curve given functions, and for the other curve other given functions, of θ .

The coordinates of a consecutive point for the one curve are then

$$x + dx + \frac{1}{2}d^2x, \quad y + dy + \frac{1}{2}d^2y, \quad z + dz + \frac{1}{2}d^2z,$$

where

$$dp = p' d\theta + \frac{1}{2}p'' d\theta^2, \quad dq = q' d\theta + \frac{1}{2}q'' d\theta^2;$$

hence these coordinates are

$$x + (ap' + a'q') d\theta + \frac{1}{2}(\alpha p'^2 + 2\alpha'p'q' + \alpha''q'^2) d\theta^2 + \frac{1}{2}(ap'' + a'q'') d\theta^2,$$

and for the other curve they are in like manner

$$x + (aP' + a'Q') d\theta + \frac{1}{2}(\alpha P'^2 + 2\alpha'P'Q' + \alpha''Q'^2) d\theta^2 + \frac{1}{2}(aP'' + a'Q'') d\theta^2,$$

766]

ON THE GEODESIC CURVATURE OF A CURVE ON A SURFACE.

325

the only difference being in the terms which contain the second differential coefficients, p'', q'' for the first curve, and P'', Q'' for the second curve. Hence the differences of the coordinates are

$$\begin{aligned} & \frac{1}{2}\{a(p'' - P'') + a'(q'' - Q'')\} d\theta^2, \quad \frac{1}{2}\{b(p'' - P'') + b'(q'' - Q'')\} d\theta^2, \\ & \frac{1}{2}\{c(p'' - P'') + c'(q'' - Q'')\} d\theta^2, \end{aligned}$$

and consequently the distance QQ' of the two consecutive points Q, Q' is

$$= \frac{1}{2}\sqrt{(E, F, G \mid p'' - P'', q'' - Q'')^2} d\theta^2.$$

The squared arc PQ^2 is

$$= (E, F, G \mid p', q')^2 d\theta^2;$$

and hence, if as before $2\rho \cdot QQ' = PQ^2$, that is, $\frac{1}{\rho} = 2QQ' \div PQ^2$, then

$$\frac{1}{\rho} = \frac{\sqrt{(E, F, G \mid p'' - P'', q'' - Q'')^2}}{(E, F, G \mid p', q')^2},$$

the required formula for ρ .

Second formula for the radius of relative curvature.

We now take the variable θ to be the length s of the curve measured from a fixed point thereof, so that p', p'' , etc. denote $\frac{dp}{ds}, \frac{d^2p}{ds^2}$, etc. We have therefore

$$1 = (E, F, G \mid p', q')^2,$$

and the formula becomes

$$\frac{1}{\rho} = \sqrt{(E, F, G \mid p'' - P'', q'' - Q'')^2}.$$

But, differentiating the above equation as regards the curve, we find

$$0 = 2(E, F, G | p', q')(p'', q'') + (E, F, G | p', q')^2,$$

where E, F, G are used to denote the complete differential coefficients $E_1p' + E_2q'$, etc. And similarly, differentiating in regard to the tangent geodesic, we obtain

$$0 = 2(E, F, G | p', q')(P'', Q'') + (E, F, G | p', q')^2;$$

and hence, taking the difference of the two equations,

$$0 = (E, F, G | p', q')(p'' - P'', q'' - Q'').$$

Hence, in the equation for $\frac{1}{\rho}$, the function under the radical sign may be written

$$(E, F, G | p', q')^2 \cdot (E, F, G | p'' - P'', q'' - Q'')^2 - \{(E, F, G | p', q')(p'' - P'', q'' - Q'')\}^2,$$

326

ON THE GEODESIC CURVATURE OF A CURVE ON A SURFACE.

[766

which is identically

$$= (EG - F^2)\{p'(q'' - Q'') - q'(p'' - P'')\}^2.$$

Hence, extracting the square root, and for $\sqrt{EG - F^2}$ writing V , we have

$$\frac{1}{\rho} = V\{p'(q'' - Q'') - q'(p'' - P'')\},$$

or say

$$\frac{1}{\rho} = V(p'q'' - q'p'') - V(p'Q'' - q'P''),$$

which is the new formula for the radius of relative curvature.

Formula for the radius of geodesic curvature.

In the paper “On Geodesic Lines, etc.,” p. 195, [vol. viii. of this Collection, p. 160], writing $EG - F^2 = V^2$, and P'', Q'' in place of p'', q'' , the differential equation of the geodesic line is obtained in the form

$$\begin{aligned} & (Ep' + Fq')\{(2F_1 - E_2)p'^2 + 2G_1p'q' + G_2q'^2\} \\ & - (Fp' + Gq')\{E_1p'^2 + 2E_2p'q' + (2F_2 - G_1)q'^2\} \\ & + 2V^2(p'Q'' - q'P'') = 0; \end{aligned}$$

or, denoting by Ω the first two lines of this equation, we have

$$V(p'Q'' - q'P'') = -\frac{1}{2V}\Omega.$$

The foregoing equation gives therefore, for the radius of geodesic curvature,

$$\frac{1}{\rho} = V(p'q'' - p''q') + \frac{1}{2V}\Omega,$$

which is an expression depending only upon p', q' , the first differential coefficients (common to the curve and geodesic), and on p'', q'' , the second differential coefficients belonging to the curve.

Observe that Ω is a cubic function of p', q' : we have

$$\Omega = (\mathfrak{A}, \mathfrak{B}, \mathfrak{C}, \mathfrak{D} | p', q')^3,$$

the values of the coefficients being

$$\begin{aligned} \mathfrak{A} &= 2EF_1 - EE_2 - FE_1, \\ \mathfrak{B} &= 2EG_1 + 2FF_1 - 3FE_2 - GE_1, \\ \mathfrak{C} &= EG_2 + 2FG_1 - 2FF_2 - 2GE_2, \\ \mathfrak{D} &= FG_2 - 2GF_2 + GG_1. \end{aligned}$$

usepackage[a4paper,margin=0.65in]geometry

[766] (CONTINUED)

ON THE GEODESIC CURVATURE OF A CURVE ON A SURFACE.

The Special Curves, p constant and q constant.

Consider the curve $p = \text{const.}$ For this curve $p' = 0$, $p'' = 0$; therefore also $Gq'^2 = 1$, and, if R be the radius of geodesic curvature, then

$$\frac{1}{R} = \frac{1}{V} \mathfrak{D}q'^3, \quad = \frac{1}{V} \frac{\mathfrak{D}}{G\sqrt{G}}.$$

Similarly for the curve $q = \text{const.}$ Here $q' = 0$, $q'' = 0$; therefore $Ep'^2 = 1$, and, if S be the radius of geodesic curvature, then

$$\frac{1}{S} = \frac{1}{V} \mathfrak{A}p'^3, \quad = \frac{1}{V} \frac{\mathfrak{A}}{E\sqrt{E}}.$$

These values of R and S are interesting for their own sakes, and they will be introduced into the expression for the radius of geodesic curvature ρ of the general curve.

Transformed Formula for the Radius of Geodesic Curvature.

From the values of $\frac{1}{R}$, $\frac{1}{S}$, we have

$$\frac{1}{\rho} = \frac{q'\sqrt{G}}{R} - \frac{p'\sqrt{E}}{S} = V(p'q'' - p''q') + \frac{1}{V} \left\{ \Omega - \frac{\mathfrak{A}}{E}p'^3 - \frac{\mathfrak{D}}{G}q'^3 \right\},$$

where the term in $\{ \}$ is

$$\Omega = \mathfrak{B}p'^3 - \frac{\mathfrak{A}}{E}p'^3 + \mathfrak{B}p''q' + (\mathfrak{C}p'q'' + \mathfrak{D})q'^2 - \frac{\mathfrak{D}}{G}q'^3.$$

The terms in $\mathfrak{A}$ are

$$-\frac{\mathfrak{A}}{E}p'(1 - Ep'^2), \quad = -\frac{\mathfrak{A}}{E}p'(2Fp'q' + Gq'^2),$$

and those in $\mathfrak{D}$ are

$$-\frac{\mathfrak{D}}{G}q'(1 - Gq'^2), \quad = -\frac{\mathfrak{D}}{G}q'(Ep'^2 + 2Fp'q').$$

Hence the whole expression contains the factor $p'q'$, and is, in fact,

$$p'q' \left(\mathfrak{B} - \frac{2\mathfrak{A}F}{E} - \frac{\mathfrak{D}E}{G} \right) p' + p'q' \left(\mathfrak{C} - \frac{\mathfrak{A}G}{E} - \frac{2\mathfrak{D}F}{G} \right) q',$$

or substituting for $\mathfrak{A}$, $\mathfrak{B}$, $\mathfrak{C}$, $\mathfrak{D}$ their values, this is

$$= p'q' \left\{ p' \left(-GE_2 + EG_2 + \frac{2F^2G_1}{E} - 2FF_2 - FE_2 + 2EF_2 - \frac{EFG_1}{G} \right) \right. \\ \left. + q' \left(-GE_2 + EG_1 - \frac{2F^2G_2}{G} + 2FF_1 + FG_1 - 2GF_1 + \frac{GFE_2}{E} \right) \right\};$$

say this is

$$= p'q'(Lp' + Mq');$$

and the formula thus is

$$\frac{1}{\rho} = \frac{q'\sqrt{G}}{R} - \frac{p'\sqrt{E}}{S} = V(p'q'' - p''q') + \frac{1}{V} p'q'(Lp' + Mq').$$

Taking ϕ , θ to be the inclination of the curve to the curves $q = \text{const.}$, $p = \text{const.}$, respectively, and $\omega (= \phi + \theta)$ the inclination of these two curves to each other, then

$$\cos \phi = \frac{Fp' + Gq'}{\sqrt{G}}, \quad \cos \theta = \frac{Ep' + Fq'}{\sqrt{E}}, \quad \cos \omega = \frac{F}{\sqrt{EG}},$$

The formula becomes in this case much more simple. We have

$$1 = Ep'^2 + Gq'^2, \quad V = \sqrt{EG}, \quad \omega = 90^\circ, \quad \sin \theta = \cos \phi;$$

and the term $Lp' + Mq'$ becomes $= \dot{E}G - E\dot{G}$, if, as before, $\dot{E}$, $\dot{G}$ denote the complete differential coefficients $E_1p' + E_2q'$ and $G_1p' + G_2q'$. The formula then is

$$\frac{1}{\rho} = \frac{\cos \phi}{R} - \frac{\sin \phi}{S} = V(p'q'' - p''q') + \frac{1}{V}p'q'(\dot{E}G - E\dot{G}),$$

where the values $\frac{1}{R}$ and $\frac{1}{S}$ are now $= \frac{\dot{G}}{G\sqrt{G}}$ and $\frac{\dot{E}}{E\sqrt{E}}$ respectively. But we have moreover $\phi = \tan^{-1} \frac{p'\sqrt{E}}{q'\sqrt{G}}$, and thence

$$\phi' = -V(p'q'' - p''q') - \frac{1}{2}Vp'q'(\dot{E}G - E\dot{G});$$

or the formula finally is

$$\frac{1}{\rho} - \frac{\cos \phi}{R} + \frac{\sin \phi}{S} + \phi' = 0,$$

which is Liouville's formula referred to at the beginning of the present paper. It will be recollected that ϕ' is the differential coefficient $\frac{d\phi}{ds}$ with respect to the arc s of the curve.

ADDITION.—Since the foregoing paper was written, I have succeeded in obtaining a like interpretation of the term

$$V(p'q'' - p''q') + \frac{1}{V}p'q'(Lp' + Mq'),$$

which belongs to the general case. I find that these terms are, in fact, $= -\phi + \omega_1 p'$; or, what is the same thing (since $\omega = \phi + \theta$ and therefore $\omega_1 p = \phi + \theta$), are $= \theta - \omega_2 q'$. It will be recollected that ϕ is the inclination of the curve to the curve $q = c$, which passes through a given point of the curve, ϕ is the variation of ϕ corresponding to the passage to the consecutive point of the curve, viz. $\phi + \phi ds$ is the inclination at this consecutive point to the curve $q = c + dc$, which passes through the consecutive point; ω is the inclination to each other of the curves $p = b$, $q = c$, which pass through the given point of the curve, ω_1 the variation corresponding to the passage along the curve $q = c$, viz. $\omega_1 ds$ is the inclination to each other of the curves $p = b + db$, $q = c$; and the like as regards θ and ω_2 .

For the demonstration, we have, as above,

$$\phi = \tan^{-1} \frac{Vp'}{Fp' + Gq'}, \quad \omega = \tan^{-1} \frac{V}{F},$$

where

$$V = \sqrt{EG - F^2};$$

and moreover $Ep'^2 + 2Fp'q' + Gq'^2 = 1$. In virtue of this last equation,

$$V^2p'^2 + (Fp' + Gq')^2 = G;$$

and we have

$$\phi = -V(p'q'' - p''q') + \frac{1}{G}\square,$$

where

$$\square = (Fp' + Gq')p'\dot{V} - Vp'(Fp' + Gq')\dot{\cdot};$$

or, since $V^2 = EG - F^2$, and thence $2V\dot{V} = \dot{G}E - 2F\dot{F} + E\dot{G}$, we have

$$\square = \frac{1}{2}p'\{(Fp' + Gq')(\dot{G}E - 2F\dot{F} + E\dot{G}) - 2(EG - F^2)(\dot{F}p' + \dot{G}q')\}.$$

Substituting herein for $\dot{E}$, $\dot{F}$, $\dot{G}$ their values $E_1p' + E_2q'$, $F_1p' + F_2q'$, $G_1p' + G_2q'$, the term in $\{ \}$ becomes

$$= Ip'^2 + Jp'q' + Kq'^2,$$

where

$$I = FGE_1 - 2EGF_1 + EFG_1,$$

$$J = G^2E_2 - 2FGF_2 + (-EG + 2F^2)G_1 + FGE_1 - 2EGF_2 + EFG_2,$$

$$K = G^2E_2 - 2FGF_2 + (-EG + 2F^2)G_2.$$

But from the equation $\omega = \tan^{-1} \frac{V}{F}$, differentiating in regard to p , we obtain

$$\omega_1 = \frac{1}{EGV}(FG\dot{E} - 2EG\dot{F} + EF\dot{G}) = \frac{1}{EGV}I;$$

or, for p writing

$$p^{-1}\{(Ep'^2 + 2Fp'q' + Gq'^2)V\omega_1, = Ep'(p'^2I + \frac{2}{p'q'}p'^2q'^2J + \frac{1}{p'^2q'^2}q'^2K)\},$$

we have

$$\phi - \omega_1p' = -V(p'q'' - p''q') + \frac{1}{G}(Ip'^2 + Jp'q' + Kq'^2).$$

The terms in p'^2 destroy each other, and the form thus is

$$\phi - \omega_1p' = -V(p'q'' - p''q') - \frac{1}{V}p'q'(Lp' + Mq'),$$

where

$$L = -\frac{J}{G} + \frac{2IF}{GE}, \quad M = -\frac{K}{G} + \frac{I}{E};$$

and, upon substituting herein for I, J, K their values, we find

$$L = -GE_2 + EG_2 + \frac{2F^2E_2}{E} - 2FF_2 - FE_2 + 2EF_2 - \frac{EFG_2}{G},$$

$$M = -GE_2 + EG_2 - \frac{2F^2G_1}{G} + 2FF_1 + FG_1 - 2GF_1 + \frac{GFE_2}{E};$$

viz., these are the values denoted above by the same letters L, M . The final result thus is

$$\frac{1}{\rho} = \frac{q'\sqrt{G}}{R} - \frac{p'\sqrt{E}}{S} = -\phi + \omega_1p', \quad = \theta - \omega_2q',$$

where the meanings of the symbols have been already explained. A formula substantially equivalent to this, but in a different (and scarcely properly explained) notation, is given, Aoust, "Théorie des coordonnées curvilignes quelconques," *Annali di Matem.*, t. ii. (1868), pp. 39-64; and I was, in fact, led thereby to the foregoing further investigation.

As to the definition of the radius of geodesic curvature, I remark that, for a curve on a given surface, if PQ be an infinitesimal arc of the curve, then if from Q we let fall the perpendicular QM on the tangent plane at P (the point M being thus a point on the tangent PT of the curve), and if from M , in the tangent plane and at right angles to the tangent, we draw MN to meet the osculating plane of the curve in N , then MN is in fact equal to the infinitesimal arc QQ' mentioned near the beginning of the present paper, and the radius of geodesic curvature ρ is thus a length such that $2\rho \cdot MN = PQ^2$.

767.

ON THE GAUSSIAN THEORY OF SURFACES.

[From the *Proceedings of the London Mathematical Society*, vol. xii. (1881), pp. 187-192.

Read June 9, 1881.]

In the Memoir, Bour, “Théorie de la déformation des surfaces” (*Jour. de l’Éc. Polyt.*, Cah. 39 (1862), pp. 1–148), the author, working with the form $ds^2 = dv^2 + g^2 du^2$ as a special case of Gauss’s formula $ds^2 = E dp^2 + 2F dp dq + G dq^2$, obtains (p. 29) the following equations which he calls *fundamental*:—

$$[\text{IV.}] \quad \begin{cases} \frac{1}{g} \frac{dg_1}{dv} = T^2 - H H_1, \\ \frac{dT}{du} + \frac{d \cdot H g}{dv} - H_1 g_1 = 0, \\ \frac{d \cdot T g^2}{dv} + g \frac{dH_1}{du} = 0, \end{cases}$$

where g_1 is written to denote $\frac{dg}{dv}$, and where (see p. 26)

H is the curvature of the normal section containing the tangent to the curve $v = \text{constant}$,

H_1 is the curvature of the normal section at right angles to the preceding, containing the tangent to the (geodesic) curve $u = \text{constant}$,

T is the torsion of the same geodesic curve;

or, what is the same thing (see p. 25), the quadric equation for the determination of the principal radii of curvature at the point of the surface is

$$\left(\frac{1}{\rho} - H\right)\left(\frac{1}{\rho} - H_1\right) - T^2 = 0.$$

Writing for greater convenience K in place of the suffixed letter H_1 , also V instead of g , so that the differential formula is $ds^2 = dv^2 + V^2 du^2$, the equations become

$$\begin{cases} \frac{1}{V} \frac{d^2 V}{dv^2} = T^2 - HK, \\ \frac{dT}{du} + \frac{d \cdot HV}{dv} - K \frac{dV}{dv} = 0, \\ \frac{d \cdot TV^2}{dv} + V \frac{dK}{du} = 0; \end{cases}$$

or, if we use the suffix 1 to denote differentiation in regard to v , and the suffix 2 to denote differentiation in regard to u , then the equations are

$$\frac{V_{11}}{V} = T^2 - HK,$$

or, what is the same thing,

$$\begin{cases} T_2 + H_1 V + (H - K) V_1 = 0, \\ T_1 V + 2TV_1 + K_2 = 0. \end{cases}$$

I wish to show how these formulae connect themselves with formulae belonging to the general form $ds^2 = E dp^2 + 2F dp dq + G dq^2$. These involve not only Gauss's coefficients E, F, G , but also the coefficients E', F', G' belonging to the inflexional tangents; and, for convenience, I quote the system of definitions, Salmon's *Geometry of Three Dimensions*, 3rd ed., 1874, p. 251, viz.

$$dx, dy, dz = a dp + a' dq, b dp + b' dq, c dp + c' dq;$$

$$d^2 x = \alpha dp^2 + 2\alpha' dp dq + \alpha'' dq^2,$$

$$d^2 y = \beta dp^2 + 2\beta' dp dq + \beta'' dq^2,$$

$$d^2 z = \gamma dp^2 + 2\gamma' dp dq + \gamma'' dq^2;$$

$$A, B, C = bc' - b'c, ca' - c'a, ab' - a'b; \quad V^2 = EG - F^2;$$

$$E' = A\alpha + B\beta + C\gamma, \quad F' = A\alpha' + B\beta' + C\gamma', \quad G' = A\alpha'' + B\beta'' + C\gamma'',$$

so that E', F', G' are, in fact, the determinants

$$\begin{vmatrix} a & b & c \\ a' & b' & c' \\ \alpha & \beta & \gamma \end{vmatrix}, \quad \begin{vmatrix} a & b & c \\ a' & b' & c' \\ \alpha' & \beta' & \gamma' \end{vmatrix}, \quad \begin{vmatrix} a & b & c \\ a' & b' & c' \\ \alpha'' & \beta'' & \gamma'' \end{vmatrix}.$$

The equation for the determination of the principal radii of curvature is

$$(E'\rho - EV)(G'\rho - GV) - (F'\rho - FV)^2 = 0.$$

which, in the particular case $F = 0$ (and therefore $V^2 = EG$), becomes

$$(E'\rho - EV)(G'\rho - GV) - F'^2\rho^2 = 0,$$

or, as this may be written,

$$EGV^2\left(\frac{1}{\rho} - \frac{E'}{EV}\right)\left(\frac{1}{\rho} - \frac{G'}{GV}\right) = F'^2,$$

an equation which corresponds with Bour's form

$$\left(\frac{1}{\rho} - K\right)\left(\frac{1}{\rho} - H\right) - T^2 = 0,$$

and becomes identical with it, if

$$E' = EVK, \quad G' = GVH, \quad F' = -V^2T.$$

But, making p, q correspond to Bour's variables, p to v , and q to u , it is necessary to show that the foregoing values (and not the interchanged values $E' = GVH, G' = EVK$) are the correct ones. We have, Salmon, p. 254,

$$\begin{vmatrix} dq, & \rho E' - VE, & \rho F' - VF \\ -dp, & \rho F' - VF, & \rho G' - VG \end{vmatrix} = 0;$$

or, putting herein $F = 0$, the equations may be written

$$\frac{dq}{-dp} = \frac{E'\left(\frac{1}{\rho} - \frac{VE}{\rho E'}\right)}{F'V} = \frac{F'V}{G'\left(\frac{1}{\rho} - \frac{VG}{\rho G'}\right)};$$

or, we see that to $dq = 0$ corresponds the value $\frac{1}{\rho} = \frac{E'}{VE}$, and to $dp = 0$ the value $\frac{1}{\rho} = \frac{G'}{VG}$. Hence the former of these values $\frac{1}{\rho}$ corresponds to $du = 0$, that is, to his $\frac{1}{\rho} = K$; and the latter to Bour's $dv = 0$, that is, to his $\frac{1}{\rho} = H$; or the values are, as stated,

$$E' = EVK, \quad G' = GVH.$$

The formula $ds^2 = E dp^2 + 2F dp dq + G dq^2$ agrees with Bour's $ds^2 = dv^2 + g^2 du^2$, if $p = u, q = v, E = 1, F = 0, G = g^2$. With these values, $V^2 = EG - F^2 = g^2$, or say $g = V$, and Bour's equation is, as it was before written, $ds^2 = dv^2 + V^2 du^2$. And we have to find the three equations which, putting therein $p = u, q = v, E = 1, F = 0, G = V^2, E' = VK, F' = -VT, G' = VH$, reduce themselves to Bour's equations.

The first of these is nothing else than the equation for the measure of curvature, viz. Salmon, p. 262 (but, using the suffixes 1 and 2 to denote differentiation in regard to p and q respectively), this is

$$\begin{aligned}
 4(E'G' - F'^2) &= E(E_2G_2 - 2F_2G_1 + G_1^2) \\
 +F(E_1G_2 - E_2G_1 - 2E_2F_2 + 4F_1F_2 - 2F_1G_1) \\
 +G(E_1G_1 - 2E_1F_2 + E_2^2) \\
 -2(EG - F^2)(E_{22} - 2F_{12} + G_{11}).
 \end{aligned}$$

In fact, writing herein $E = 1$, $F = 0$, and therefore the differential coefficients of E and F each $= 0$, the equation becomes

$$4(E'G' - F'^2) = G_1^2 - 2GG_{11},$$

which is

$$4V^4(HK - T^2) = (2VV_1)^2 - 2V^2(2V_1^2 + 2VV_{11}), \quad = -4V^3V_{11};$$

or finally it is

$$V_{11} = V(T^2 - HK).$$

The other two of Bour's equations are derived from equations which give respectively the values of $E'_2 - F'_1$ and $F'_2 - G'_1$; viz. starting from the equations

$$\begin{aligned} E' &= A\alpha + B\beta + C\gamma, \\ F' &= A\alpha' + B\beta' + C\gamma', \\ G' &= A\alpha'' + B\beta'' + C\gamma'', \end{aligned}$$

we see at once that E'_2 and F'_1 contain, E'_2 the terms $A\alpha_2 + B\beta_2 + C\gamma_2$, and F'_1 the terms $A\alpha'_1 + B\beta'_1 + C\gamma'_1$, which are equal to each other (since $\alpha_2 = \alpha'_1$, α and α' are the differential coefficients x_{11} , x_{12} of x , and so β_2 and β'_1 , γ_2 and γ'_1). Hence

$$E'_2 - F'_1 = A_2\alpha + B_2\beta + C_2\gamma - A_1\alpha' - B_1\beta' - C_1\gamma'$$

and similarly

$$F'_2 - G'_1 = A_2\alpha' + B_2\beta' + C_2\gamma' - A_1\alpha'' - B_1\beta'' - C_1\gamma''.$$

Here, from the values of A , B , C , we have

$$\begin{aligned} A &= bc' - cb'; & A_1 &= \beta c' + b\gamma' - \gamma b' - c\beta'; & A_2 &= \beta' c' - \gamma' b' + b\gamma'' - c\beta'' \\ B &= ca' - ac'; & B_1 &= \gamma a' + c\alpha' - \alpha c' - a\gamma'; & B_2 &= \gamma' a' - \alpha' c' + c\alpha'' - a\gamma'' \\ C &= ab' - ba'; & C_1 &= \alpha b' + a\beta' - \beta a' - b\alpha'; & C_2 &= \alpha' b' - \beta' a' + a\beta'' - b\alpha'' \end{aligned}$$

and, substituting, we find

$$\begin{aligned} E'_2 - F'_1 &= 2\alpha'\alpha\alpha' + \alpha\alpha''\alpha, \\ F'_2 - G'_1 &= -2\alpha\alpha'\alpha'' - \alpha'\alpha''\alpha, \end{aligned}$$

if, for shortness, $\alpha'\alpha\alpha'$ denotes the determinant

$$\begin{vmatrix} \alpha' & \alpha & \alpha \\ b' & \beta & \beta' \\ c' & \gamma & \gamma' \end{vmatrix},$$

and so for the other like symbols. Observe that, with

$$\begin{aligned} a, & a', \alpha, \alpha', \alpha'' \\ b, & b', \beta, \beta', \beta'' \\ c, & c', \gamma, \gamma', \gamma'' \end{aligned},$$

we have in all 10 determinants, viz. these are $aa'\alpha, = E'$; $aa'\alpha', = F'$; $aa'\alpha'', = G'$; $\alpha\alpha'\alpha''$; and the six determinants $\alpha\alpha'\alpha'$, $\alpha\alpha'\alpha''$, $\alpha\alpha''\alpha$; $\alpha'\alpha\alpha'$, $\alpha'\alpha'\alpha''$, $\alpha'\alpha''\alpha$. The foregoing expressions of $E'_2 - F'_1$ and $F'_2 - G'_1$ respectively, substituting therein for the determinants $\alpha'\alpha\alpha'$, $\alpha\alpha''\alpha$, $\alpha\alpha'\alpha''$, $\alpha'\alpha''\alpha$ their values as about to be obtained, are the required two equations. We have

$$\begin{aligned} aa + bb + cc &= E, & aa' + bb' + cc' &= F, \\ a'a + b'b + c'c &= F, & a'a' + b'b' + c'c' &= G, \\ \alpha a + \beta b + \gamma c &= \frac{1}{2}E_1, & \alpha a' + \beta b' + \gamma c' &= F_1 - \frac{1}{2}E_2, \\ \alpha' a + \beta' b + \gamma' c &= \frac{1}{2}E_2, & \alpha' a' + \beta' b' + \gamma' c' &= \frac{1}{2}G_1, \\ \alpha'' a + \beta'' b + \gamma'' c &= F_2 - \frac{1}{2}G_1, & \alpha'' a' + \beta'' b' + \gamma'' c' &= \frac{1}{2}G_2, \end{aligned}$$

and if from the first five equations, regarded as equations linear in (a, b, c) , we eliminate these quantities, and from the second five equations, regarded as linear in (a', b', c') , we eliminate these quantities, we obtain two sets each of five equations.

$$\begin{vmatrix} a, & \alpha, & \alpha, & \alpha', & \alpha'' \\ b, & b', & \beta, & \beta', & \beta'' \\ c, & c', & \gamma, & \gamma', & \gamma'' \\ E, & F, & \frac{1}{2}E_1, & \frac{1}{2}E_2, & F_1 - \frac{1}{2}G_1 \end{vmatrix} = 0, \quad \begin{vmatrix} a, & a', & \alpha, & \alpha', & \alpha'' \\ b, & b', & \beta, & \beta', & \beta'' \\ c, & c', & \gamma, & \gamma', & \gamma'' \\ F, & G, & F_1 - \frac{1}{2}E_2, & \frac{1}{2}G_1, & \end{vmatrix} = 0.$$

These may be written,

$$\begin{aligned} F\alpha\alpha'\alpha'' - \frac{1}{2}E_1\alpha'\alpha\alpha'' - \frac{1}{2}E_2\alpha'\alpha''\alpha - (F_1 - \frac{1}{2}G_1)\alpha'\alpha\alpha' &= 0, \\ -E\alpha\alpha'\alpha'' + \frac{1}{2}E_1\alpha\alpha'\alpha'' + \frac{1}{2}E_2\alpha\alpha''\alpha + (F_1 - \frac{1}{2}G_1)\alpha\alpha\alpha' &= 0, \\ E\alpha'\alpha'\alpha'' - F\alpha\alpha'\alpha'' + \frac{1}{2}E_1G' - (F_1 - \frac{1}{2}G_1)F' &= 0, \\ E\alpha'\alpha''\alpha - F\alpha\alpha''\alpha - \frac{1}{2}E_1G' + (F_1 - \frac{1}{2}G_1)E' &= 0, \\ E\alpha'\alpha\alpha - F\alpha\alpha\alpha' + \frac{1}{2}E_2F' - \frac{1}{2}E_1E' &= 0; \end{aligned}$$

and

$$\begin{aligned} G\alpha\alpha'\alpha'' - (F_2 - \frac{1}{2}E_2)\alpha'\alpha\alpha'' - \frac{1}{2}G_1\alpha'\alpha''\alpha - \frac{1}{2}G_2\alpha'\alpha\alpha' &= 0, \\ -F\alpha\alpha\alpha'' + (F_2 - \frac{1}{2}E_2)\alpha\alpha'\alpha'' + \frac{1}{2}G_1\alpha\alpha''\alpha + \frac{1}{2}G_2\alpha\alpha\alpha' &= 0, \\ F\alpha'\alpha'\alpha'' - G\alpha\alpha\alpha'' + \frac{1}{2}G_1G' - \frac{1}{2}G_2F' &= 0, \\ F\alpha'\alpha''\alpha - G\alpha\alpha'\alpha - (F_2 - \frac{1}{2}E_2)G' + \frac{1}{2}G_1E' &= 0, \\ F\alpha'\alpha\alpha - G\alpha\alpha\alpha + (F_2 - \frac{1}{2}E_2)F' - \frac{1}{2}G_2E' &= 0. \end{aligned}$$

Attending in each set only to the third, fourth, and fifth equations, and combining these in pairs, we obtain

$$V^2\alpha\alpha'\alpha'' + (\frac{1}{2}FG_1 - FF_1 + \frac{1}{2}EG_1)F' + (-\frac{1}{2}EG_1 + \frac{1}{2}FE_2)G' = 0,$$

$$\begin{aligned}
& V^2 \alpha' \alpha' \alpha'' + (\tfrac{1}{2} G G_1 - G F_1 + \tfrac{1}{2} F G_2) F' + (-\tfrac{1}{2} F G_1 + \tfrac{1}{2} G E_2) G' = 0; \\
& V^2 \alpha \alpha'' \alpha + (-\tfrac{1}{2} F E_1 + E F_1 - \tfrac{1}{2} E E_2) G' + (-\tfrac{1}{2} F G_1 + F F_1 - \tfrac{1}{2} E G_1) E' = 0, \\
& V^2 \alpha' \alpha'' \alpha + (-\tfrac{1}{2} G E_1 + F F_1 - \tfrac{1}{2} F E_2) G' + (-\tfrac{1}{2} G G_1 + G F_1 - \tfrac{1}{2} F G_2) E' = 0; \\
& V^2 \alpha \alpha \alpha' + (\tfrac{1}{2} E G_2 - \tfrac{1}{2} F E_2) E' + (\tfrac{1}{2} F E_1 - E F_1 + \tfrac{1}{2} E E_2) F' = 0, \\
& V^2 \alpha' \alpha \alpha' + (\tfrac{1}{2} F G_2 - \tfrac{1}{2} G E_2) E' + (\tfrac{1}{2} G E_1 - F F_1 + \tfrac{1}{2} F E_2) F' = 0.
\end{aligned}$$

We thus obtain

$$\begin{aligned} E'_2 - F'_1 &= \frac{2}{\sqrt{2}} [(-\frac{1}{2}FG_1 + \frac{1}{2}GE_2)E' + (-\frac{1}{2}GE_1 + FF_1 - \frac{1}{2}FF_2)F'] \\ &\quad + \frac{1}{\sqrt{2}} \{(\frac{1}{2}FE_1 - EF_1 + \frac{1}{2}EE_2)G' + (\frac{1}{2}FG_1 - FF_1 + \frac{1}{2}EG_1)E'\}, \\ F'_2 - G'_1 &= \frac{2}{\sqrt{2}} \{(\frac{1}{2}FG_1 - FF_1 + \frac{1}{2}EG_1)F' + (-\frac{1}{2}EG_1 + \frac{1}{2}FE_2)G'\} \\ &\quad + \frac{1}{\sqrt{2}} \{(-\frac{1}{2}GE_1 + FF_1 - \frac{1}{2}FE_2)G' + (-\frac{1}{2}GG_1 + GF_1 - \frac{1}{2}FG_2)E'\}; \end{aligned}$$

or, finally,

$$\begin{aligned} E'_2 - F'_1 &= \frac{1}{\sqrt{2}} \{(-\frac{1}{2}FG_1 + GE_2 - FF_1 + \frac{1}{2}EG_2)E' \\ &\quad + (-GE_1 + 2FF_1 - FF_2)F' + (\frac{1}{2}FE_1 - EF_1 + \frac{1}{2}EE_2)G'\}, \\ F'_2 - G'_1 &= \frac{1}{\sqrt{2}} \{(-\frac{1}{2}GG_1 + GF_1 - \frac{1}{2}FG_2)E' \\ &\quad + (FG_1 - 2FF_1 + EG_2)F' + (-\frac{1}{2}GE_1 + FF_1 - EG_1 + \frac{1}{2}FE_2)G'\}, \end{aligned}$$

which are the required formulae; and which may, I think, be regarded as new formulae in the Gaussian theory of surfaces.

Writing herein as before, the first of these becomes

$$V^2(VK)_2 + (V^2T)_1 = \frac{1}{\sqrt{2}} [\frac{1}{2}(V^2)_1VK] = V_2K,$$

that is,

$$V^2V_2K + V^3K_2 + V^2T_1 + 2VV_1T = V_2K;$$

or finally

$$VT_1 + 2TV_1 + K_2 = 0,$$

which is Bour's third equation. And the second equation becomes

$$-(V^2T)_2 - (V^2H)_1 = \frac{1}{\sqrt{2}} \{-\frac{1}{2}V^2V_1VK + (V^2)_1(-V^2T)(V^2)_1VH\}$$

that is,

$$= -V^2V_1K - 2VV_1T - 2V^2V_1H,$$

or finally

$$-V^2T_2 - 2VV_2T - V^2H_1 - 3V^2V_1H = -V^2V_1K - 2VV_1T - 2V^2V_1H;$$

$$T_2 + VH_1 + (H - K)V_1 = 0,$$

which is Bour's second equation.

768.

NOTE ON LANDEN'S THEOREM.

[From the *Proceedings of the London Mathematical Society*, vol. xiii.
(1882), pp. 47, 48.

Read November 10, 1881.]

Landen's theorem, as given in the paper "An Investigation of a General Theorem for finding the length of any Arc of any Conic Hyperbola by means of two Elliptic Arcs, with some other new and useful Theorems deduced therefrom," *Phil. Trans.*, t. lxv. (1775), pp. 283–289, is, as appears by the title, a theorem for finding the length of a hyperbolic arc in terms of the length of two elliptic arcs; this theorem being obtained by means of the following differential identity, viz., if

$$t = gx \sqrt{\frac{m^2 - x^2}{m^2 - g^2 x^2}}$$

where

$$g = \frac{m^2 - n^2}{n^2},$$

then

$$\sqrt{\frac{m^2 - gx^2}{m^2 - x^2}} dx = \left\{ \frac{1}{2} + \frac{1}{4} \sqrt{\frac{(m-n)^2 - t^2}{(m+n)^2 - t^2}} + \frac{1}{4} \sqrt{\frac{(m+n)^2 - t^2}{(m-n)^2 - t^2}} \right\} dt,$$

(this is exactly Landen's form, except that he of course writes $\dot{x}$, $\dot{t}$ respectively); viz., integrating each side, and interpreting geometrically in a very ingenious and elegant manner the three integrals which present themselves, he arrives at his theorem for the hyperbolic arc; but with this I am not now concerned.

Writing for greater convenience $m = 1$, $n = k'$, and therefore $g = k^2$, if as usual $k^2 + k'^2 = 1$, the transformation is

$$t = k^2 x \sqrt{\frac{1 - x^2}{1 - k^4 x^2}},$$

leading to

$$\sqrt{\frac{1-k^2x^2}{1-x^2}} dx = \left\{ \frac{1}{2} + \frac{1}{4} \sqrt{\frac{(1-k')^2-t^2}{(1+k')^2-t^2}} + \frac{1}{4} \sqrt{\frac{(1+k')^2-t^2}{(1-k')^2-t^2}} \right\} dt.$$

The form in which the transformation is usually employed (see my *Elliptic Functions*, pp. 177, 178) is

$$y = (1+k')x \sqrt{\frac{1-x^2}{1-k^2x^2}},$$

leading to

$$\frac{(1+k') dx}{\sqrt{1-x^2} \cdot \sqrt{1-k^2x^2}} = \frac{dy}{\sqrt{1-y^2} \cdot \sqrt{1-\lambda^2y^2}},$$

where

$$\lambda = \frac{1-k'}{1+k'}.$$

If, to identify the two forms, we write $y = \frac{t}{1-k'}$, and in the last equation introduce t in place of y , the last equation becomes

$$\frac{dx}{\sqrt{1-x^2} \cdot \sqrt{1-k^2x^2}} = \frac{dt}{\sqrt{\{(1-k')^2-t^2\}\{(1+k')^2-t^2\}}}.$$

Comparing with Landen's form, in order that the two may be identical, we must have

$$1-k^2x^2 = \left\{ \frac{1}{2} + \frac{1}{4} \sqrt{\frac{(1-k')^2-t^2}{(1+k')^2-t^2}} + \frac{1}{4} \sqrt{\frac{(1+k')^2-t^2}{(1-k')^2-t^2}} \right\} \times \sqrt{(1-k')^2-t^2} \sqrt{(1+k')^2-t^2},$$

viz., this is

$$1-k^2x^2 = \frac{1}{4} [\sqrt{(1-k')^2-t^2} + \sqrt{(1+k')^2-t^2}]^2,$$

that is,

$$1-k^2x^2 = \frac{1}{2} [1+k'^2-t^2 + \sqrt{\{(1-k')^2-t^2\}\{(1+k')^2-t^2\}}],$$

where the function under the radical sign is

$$k'^4 - 2(1+k'^2)t^2 + t^4 (= T \text{ suppose})$$

and this must consequently be a form of the original integral equation

$$t = k^2x \sqrt{\frac{1-x^2}{1-k^4x^2}}.$$

In fact, squaring and solving in regard to x^2 with the assumed sign of the radical, we have

$$x^2 = \frac{k'^2 + t^2 - \sqrt{T}}{2k^2},$$

corresponding to an equation given by Landen. And we thence have

$$1 - k^2 x^2 = \frac{1}{2}[2 - k'^2 - t^2 + \sqrt{T}], \quad = \frac{1}{2}[1 + k^2 - t^2 + \sqrt{T}],$$

which is the required expression for $1 - k^2 x^2$.

The trigonometrical form $\sin(2\phi' - \phi) = c \sin \phi$ of the relation between y and x does not occur in Landen; it is employed by Legendre, I believe, in an early paper, *Mém. de l'Acad. de Paris*, 1786, and in the *Exercices*, 1811, and also in the *Traité des Fonctions Elliptiques*, 1825, and by means of it he obtains an expression for the arc of a hyperbola in terms of two elliptic functions, $E(c, \phi)$, $E(c', \phi')$, showing that the arc of the hyperbola is expressible by means of two elliptic arcs, this, he observes, “est le beau théorème dont Landen a enrichi la géométrie.” We have, then (1828), Jacobi's proof, by two fixed circles, of the addition-theorem (see my *Elliptic Functions*, p. 28), and the application of this (p. 30) to Landen's theorem is also due to Jacobi, see the “Extrait d'une lettre adressée à M. Hermite,” *Crelle*, t. xxxii. (1846), pp. 176–181; the connection of the demonstrations, by regarding the point, which is alone necessary for Landen's theorem as the limit of the smaller circle in the figure for the addition-theorem is due to Durege (see his *Theorie der elliptischen Functionen*, Leipzig, 1861, pp. 168, et seq.).

769.

ON A FORMULA RELATING TO ELLIPTIC INTEGRALS OF
THE THIRD KIND.

[From the *Proceedings of the London Mathematical Society*, vol. xiii.
(1882), pp. 175, 176. Presented May 11, 1882.]

The formula for the differentiation of the integral of the third kind

$$\Pi = \int_0^\phi \frac{d\phi}{(1 + n \sin^2 \phi) \Delta}$$

in regard to the parameter n , see my *Elliptic Functions*, Nos. 174 *et seq.*, may be presented under a very elegant form, by writing therein

$$\sin^2 \phi = x = \operatorname{sn}^2 u, \quad \sin \phi \cos \phi \Delta = y = \operatorname{sn} u \operatorname{cn} u \operatorname{dn} u,$$

and thus connecting the formula with the cubic curve

$$y^2 = x(1-x)(1-k^2x).$$

The parameter must, of course, be put under a corresponding form, say $n = -\frac{1}{a}$, where $a = \operatorname{sn}^2 \theta$, $b = \operatorname{sn} \theta \operatorname{cn} \theta \operatorname{dn} \theta$, and therefore (a, b) are the coordinates of the point corresponding to the argument θ . The steps of the substitution may be effected without difficulty, but it will be convenient to give at once the final result and then verify it directly. The result is

$$\frac{d}{d\theta} \frac{b}{a-x} = \frac{d}{du} \frac{y}{x-a} = k^2(a-x).$$

We, in fact, have

$$\frac{d}{du} x = 2 \operatorname{sn} u \operatorname{cn} u \operatorname{dn} u = 2y,$$

and thence

$$y \frac{dx}{du} = 2y^2,$$

that is,

$$y \frac{dx}{du} = 2x[1 - (1 + k^2)x + k^2x^2].$$

Also

$$\begin{aligned} \frac{dy}{du} &= \text{cn}^2 u \, \text{dn}^2 u - \text{sn}^2 u \, \text{dn}^2 u - k^2 \text{sn}^2 u \, \text{cn}^2 u \\ &= 1 - 2(1 + k^2)x + 3k^2x^2, \end{aligned}$$

and hence

$$\begin{aligned} \frac{d}{du} \frac{y}{x - a} &= \frac{1}{(a - x)^2} \left\{ (a - x) \frac{dy}{du} - y \frac{dx}{du} \right\} \\ &= \frac{1}{(a - x)^2} [-x - a + 2(1 + k^2)ax + k^2x^3 - 3k^2ax^2]. \end{aligned}$$

Interchanging the letters, we have

$$\frac{d}{d\theta} \frac{b}{a - x} = \frac{1}{(a - x)^2} [-x - a + 2(1 + k^2)ax + k^2a^3 - 3k^2a^2x],$$

and hence, subtracting,

$$\begin{aligned} \frac{d}{d\theta} \frac{b}{a - x} - \frac{d}{du} \frac{y}{x - a} &= \frac{1}{(a - x)^2} \{ k^2(a - x)^3 \} \\ &= k^2(a - x), \end{aligned}$$

which is the required result.

770.

ON THE 34 CONCOMITANTS OF THE TERNARY CUBIC.

[From the *American Journal of Mathematics*, vol. iv. (1881), pp. 1–15.]

I have, (by aid of Gundelfinger's formulae, afterwards referred to), calculated, and I give in the present paper, the expressions of the 34 concomitants of the canonical ternary cubic $ax^3 + by^3 + cz^3 + 6lxyz$, or, what is the same thing, the 34 covariants of this cubic and the adjoint linear function $\xi x + \eta y + \zeta z$: this is the chief object of the paper. I prefix a list of memoirs, with short remarks upon some of them; and, after a few observations, proceed to the expressions for the 34 concomitants; and, in conclusion, exhibit the process of calculation of these concomitants other than such of them as are taken to be known forms. I insert a supplemental table of 6 derived forms.

The list of memoirs (not by any means a complete one) is as follows:

Hesse, Ueber die Elimination der Variabeln aus drei algebraischen Gleichungen vom zweiten Grade mit zwei Variabeln: *Crelle*, t. xxviii. (1844), pp. 68–96. Although purporting to relate to a different subject, this is in fact the earliest, and a very important, memoir in regard to the general ternary cubic; and in it is established the canonical form, as Hesse writes it, $y_1^3 + y_2^3 + y_3^3 + 6my_1y_2y_3$.

Aronhold, Zur Theorie der homogenen Functionen dritten Grades von drei Variabeln: *Crelle*, t. xxxix. (1850), pp. 140–159.

Cayley, Third Memoir on Quantics: *Phil. Trans.*, t. cxlvi. (1856), pp. 627–647 [144].

Aronhold, Theorie der homogenen Functionen dritten Grades von drei Variabeln: *Crelle*, t. lv. (1858), pp. 97–191.

Salmon, Lessons Introductory to the Modern Higher Algebra: 8°, Dublin, 1859.

Cayley, Seventh Memoir on Quantics: *Phil. Trans.*, t. cli. (1861), pp. 277–292 [269].

Brioschi, Sur la théorie des formes cubiques à trois indéterminées: *Comptes Rendus*, t. lvi. (1863), pp. 304–307.

Hermite, Extrait d'une lettre à M. Brioschi: *Crelle*, t. lxiii. (1864), pp. 30–32, followed by a note by Brioschi, pp. 32–33.

The skew covariant of the ninth order, which is $y^3 - z^3 \cdot z^3 - x^3 \cdot x^3 - y^3$ for the canonical form $x^3 + y^3 + z^3 + 6lxyz$, and the corresponding contravariant $\eta^3 - \zeta^3 \cdot \zeta^3 - \xi^3 \cdot \xi^3 - \eta^3$, alluded to p. 116 of Salmon's *Lessons*, were obtained, the covariant by Brioschi and the contravariant by Hermite, in the last-mentioned papers.

Clebsch and Gordan, Ueber die Theorie der ternären cubischen Formen: *Math. Annalen*, t. i. (1869), pp. 56–89.

The establishment of the complete system of the 34 covariants, contravariants and *Zwischenformen*, or, as I have here called them, the 34 concomitants, was first effected by Gordan in the next following memoir:

Gordan, Ueber die ternären Formen dritten Grades: *Math. Annalen*, t. i. (1869), pp. 90–128.

And the theory is farther considered:

Gundelfinger, Zur Theorie der ternären cubischen Formen: *Math. Annalen*, t. vi. (1871), pp. 144–163. The author speaks of the 34 forms as being “theils mit den von Gordan gewählten identisch, theils möglichst einfache Combinationen derselben.” They are, in fact, the 34 forms given in the present paper for the canonical form of the cubic, and the meaning of the adopted combinations of Gordan's forms will presently clearly appear.

There is an advantage in using the form $ax^3 + by^3 + cz^3 + 6lxyz$ rather than the Hessian form $x^3 + y^3 + z^3 + 6lxyz$, employed in my Third and Seventh Memoirs on Quantics: for the form $ax^3 + by^3 + cz^3 + 6lxyz$ is what the general cubic

$$(a, b, c, f, g, h, i, j, k, l)(x, y, z)^3$$

becomes by no other change than the reduction to zero of certain of its coefficients; and thus any concomitant of the canonical form consists of terms which are leading terms of the same concomitant of the general form.

The concomitants are functions of the coefficients $(a, b, \dots, l)$, of (ξ, η, ζ) , and of (x, y, z) : the dimensions in regard to the three sets respectively may be distinguished as the degree, class, and order; and we have thus to consider the deg-class-order of a concomitant.

Two or more concomitants of the same deg-class-order may be linearly combined together: viz., the linear combination is the sum of the concomitants each multiplied by a mere number. The question thus arises as to the selection of a representative concomitant. As already mentioned, I follow Gundelfinger, viz., my 34 concomitants of the canonical form correspond each to each (with only the difference of a numerical factor of the entire concomitant) to his 34 concomitants of the general form. The principle underlying the selection would, in regard to the general form, have to be explained altogether differently; but this principle exhibits itself in a very remarkable manner in regard to the canonical form $ax^3 + by^3 + cz^3 + 6lxyz$.

Each concomitant of the general form is an indecomposable function, not breaking up into rational factors; but this is not of necessity the case in regard to a canonical form: only a concomitant which does break up must be regarded as indecomposable, no factor of such concomitant being rejected, or separated. So far from it, there is, in regard to the canonical form in question, a frequent occurrence of $abc + 8l^3$ or a power thereof, either as a factor of a unique concomitant, or when there are two or more concomitants of the same deg-class-order, then as a factor of a properly selected linear combination of such concomitants: and the principle referred to is, in fact, that of the selection of such combination for the representative concomitant; or (in other words) the representative concomitant is taken so as to contain as a factor the highest power that may be of $abc + 8l^3$. As to the signification of this expression $abc + 8l^3$, I call to mind that the discriminant of the form is $abc(abc + 8l^3)^3$.

As to numerical factors: my principle has been, and is, to throw out any common numerical divisor of all the terms: thus I write $S = -abcl + l^4$, instead of Aronhold's $S = -4abcl + 4l^4$. There is also the question of nomenclature: I retain that of my Seventh Memoir on Quantics, except that I use single letters H , P , &c., instead of the same letters with U , thus HU , PU , &c.; in particular, I use U , H , P , Q instead of Aronhold's f , Δ , S_f , T_f . It is thus at all events necessary to make some change in Gundelfinger's letters; and there is moreover a laxity in his use of accented letters; his B , B' , B'' , B''' , and so in other cases E , E' , E'' , &c., are used to denote functions derived in a determinate manner each from the preceding one (by the δ -process explained further on); whereas his L , $\bar{L}$; M , M' ; N , N' are functions having to each other an altogether different relation; also three of his functions are not denoted by any letters at all. Under the circumstances, I retain only a few of his letters; use the accent where it denotes the δ -process; and introduce barred letters $\bar{J}$, $\bar{K}$, &c., to denote a different correspondence with the unbarred letters J , K , &c. But I attach also to each concomitant a numerical symbol showing its deg-class-order, thus: 541 (degree = 5, class = 4, order = 1) or 1290, (there is no ambiguity in the two-digit numbers 10, 11, 12 which present themselves in the system of the 34 symbols); and it seems to me very desirable that the significations of these deg-class-order symbols should be considered as permanent and unalterable. Thus, in writing $S = 400 = -abcl + l^4$, I wish the 400 to be regarded as denoting its expressed value $-abcl + l^4$: if the same letter S is to be used in Aronhold's sense to denote $-4abcl + 4l^4$, this would be completely expressed by the new definition $S_1 = 4.400$, the meaning of the symbol 400 being explained by reference to the present memoir, or by the actual quotation $400 = -abcl + l^4$.

I proceed at once to the table: for shortness, I omit, in general, terms which can be derived from an expressed term by mere cyclical interchanges of the letters (a, b, c) , (ξ, η, ζ) , (x, y, z) .

*Table of the 34 Covariants of the Canonical Cubic $ax^3 + by^3 + cz^3 + 6lxyz$
and the linear form $\xi x + \eta y + \zeta z$.*

First Part, 10 Forms. Class = Order.

Current No.

- 1 $S = 400 = -abcl + l^4.$
- 2 $T = 600 = a^2b^2c^2 - 20abcl^3 - 8l^6.$
- 3 $A = 011 = \xi x + \eta y + \zeta z.$
- 4 $\Theta = 222 = x^2[-l^2\xi^2 - 2al\eta\zeta] \dots$
 $+ yz[bc\xi^2 + 2l^2\eta\zeta] \dots$
- 5 $\Theta' = 422 = x^2[l(abc + 2l^3)\xi^2 + a(abc - 4l^3)\eta\zeta] \dots$
 $+ yz[6bcl^2\xi^2 - 2l(abc + 2l^3)\eta\zeta] \dots$
- 6 $\Theta'' = 622 = x^2[-(abc + 2l^3)^2\xi^2 + 2al^2(abc - 2l^3)\eta\zeta] \dots$
 $+ yz[36bcl^4\xi^2 + 2(abc + 2l^3)^2\eta\zeta] \dots$
- 7 $B = 333 = x^3[a^2(c\eta^3 - b\zeta^3)] \dots$
 $+ y^2z[(abc + 8l^3)\eta^2\zeta + 2bl^2\xi^3 - 6bcl\xi\eta^2] \dots$
 $+ yz^2[-(abc + 8l^3)\eta\zeta^2 - 6bcl\xi^2\eta - 12cl^2\xi^3] \dots$
- 8 $B' = 533 = x^3[3a^2l^2(c\eta^3 - b\zeta^3)] \dots$
 $+ y^2z[-l^2(abc + 8l^3)\eta\zeta^2 + 4bl(-abc + l^3)\xi^2\zeta - bc(abc + 10l^3)\xi\eta^2] \dots$
 $+ yz^2[l^2(abc + 8l^3)\eta\zeta^2 - bc(abc + 10l^3)\xi^2\eta + 4cl(-abc + l^3)\xi\zeta^2] \dots$
- 9 $B'' = 733 = x^3[9a^2l^4(c\eta^3 - b\zeta^3)] \dots$
 $+ y^2z[l(abc + 8l^3)(2abc + l^3)\eta\zeta^2$
 $+ b(abc + 2l^3)(abc - 10l^3)\xi^2\zeta + 6bcl^2(-abc + l^3)\xi\eta^2] \dots$
 $+ yz^2[-l(abc + 8l^3)(2abc + l^3)\eta\zeta^2$
 $- 6bcl^2(-abc + l^3)\xi^2\eta + c(abc + 2l^3)(abc - 10l^3)\xi\zeta^2] \dots$
- 10 $B''' = 933 = x^3[27a^2l^6(c\eta^3 - b\zeta^3)] \dots$
 $+ y^2z[-(abc + 8l^3)(abc - l^3)^2\eta^2\zeta + 9bl^3(abc - 2l^3)^2\xi^2\zeta$
 $- 27bcl^4(abc + 2l^3)\xi\eta^2] \dots$
 $+ yz^2[(abc + 8l^3)(abc - l^3)^2\eta\zeta^2 - 27bcl^4(abc + 2l^3)\xi^2\eta$
 $- 9cl^3(abc + 2l^3)^2\xi\eta\zeta^2] \dots$

Second Part, $(4 + 4 =) 8$ forms. Class = 0, and Order = 0.

Class = 0.

- 11 $U = 103 = ax^3 + by^3 + cz^3 + 6lxyz.$

$$12 \quad H = 303 = l^2(ax^3 + by^3 + cz^3) - (abc + 2l^3)xyz.$$

$$13 \quad \Psi = 806 = (abc + 8l^3)^2(a^2b^2c^2 + b^2y^6 + c^2z^6 - 10(bcy^3z^3 + caz^3x^3 + abx^3y^3)).$$

$$14 \quad \Omega = 1209 = (abc + 8l^3)^3[by^3 - cz^3 \cdot cz^3 - ax^3 \cdot ax^3 - by^3].$$

Current No. Order = 0.

- 15 $P = 330 = -l(bc\xi^3 + c\eta^3 + ab\zeta^3) + (-abc + 4l^3)\xi\eta\zeta.$
- 16 $Q = 530 = (abc - 10l^3)(bc\xi^3 + c\eta^3 + ab\zeta^3) - 6l^2(5abc + 4l^3)\xi\eta\zeta.$
- 17 $F = 460 = b^2c^2\xi^6 + c^2a^2\eta^6 + a^2b^2\zeta^6$
 $- 2(abc + 16l^3)(a\eta^3\zeta^3 + b\zeta^3\xi^3 + c\xi^3\eta^3)$
 $- 24l^2(bc\xi^4 + ca\eta^4 + ab\zeta^4)\xi\eta\zeta - 24l(abc + 2l^3)\xi^2\eta^2\zeta^2.$
- 18 $\Pi = 1290 = (abc + 8l^3)^3[c\eta^3 - b\zeta^3 \cdot a\xi^3 - c\eta^3 \cdot b\zeta^3 - a\xi^3].$

Third Part, (8 + 8 =) 16 forms. Class less or greater than Order.

Class less than Order.

- 19 $J = 413 = (abc + 8l^3)\{\xi x(by^3 - cz^3) + \eta y(cz^3 - ax^3) + \zeta z(ax^3 - by^3)\}.$
- 20 $K = 613 = (abc + 8l^3)\{\xi[alx^4 - 2blxy^3 - 2clxz^3 - 3bcy^2z^2] \dots\}.$
- 21 $K' = 813 = (abc + 8l^3)\{\xi[(abc + 2l^3)(ax^4 - 2bxy^3 - 2cxz^3) - 18bcl^2y^2z^2] \dots\}.$
- 22 $E = 625 = (abc + 8l^3)\{\xi(by^3 - cz^3)[2l^2x^2 + bcyz] \dots$
 $+ \eta\zeta(by^3 - cz^3)[4alx^2 - 2l^2yz] \dots\}.$
- 23 $E' = 825 = (abc + 8l^3)\{\xi(by^3 - cz^3)[l(abc + 2l^3)x^2 - 3bcl^2yz] \dots$
 $+ \eta\zeta(by^3 - cz^3)[a(abc + 4l^3)x^2 - l(abc + 2l^3)yz] \dots\}.$
- 24 $E'' = 1025 = (abc + 8l^3)\{\xi(by^3 - cz^3)[(abc + 2l^3)^2x^2 + 18bcl^4yz] \dots$
 $+ \eta\zeta(by^3 - cz^3)[-12al^3(abc + 2l^3)x^2 + (abc + 2l^3)^2yz] \dots\}.$
- 25 $M = 917 = (abc + 8l^3)^2\{\xi(by^3 - cz^3)[5alx^4 - blxy^3 - clxz^3 - 3bcy^2z^2] \dots\}.$
- 26 $M' = 1117 = (abc + 8l^3)^2\{\xi(by^3 - cz^3)[(abc + 2l^3)(5ax^4 - bxy^3 - cxz^3)$
 $+ 18bcl^2y^2z^2] \dots\}.$

Order less than Class.

- 27 $\bar{J} = 841 = (abc + 8l^3)^2[x\xi a(c\eta^3 - b\zeta^3) + y\eta b(a\xi^3 - c\zeta^3) + z\zeta c(b\zeta^3 - a\eta^3)].$
- 28 $\bar{K} = 541 = (abc + 8l^3)\{x[bc\xi^4 - 2ca\xi\eta^3 - 2ab\xi\zeta^3 - 6ab\eta^2\zeta^2] \dots\}.$
- 29 $\bar{K}' = 741 = (abc + 8l^3)\{x[l^2(bc\xi^4 - 2ca\xi\eta^3 - 2ab\xi\zeta^3) + a(abc + 2l^3)\eta^2\zeta^2] \dots\}.$
- 30 $\bar{E} = 652 = (abc + 8l^3)\{x^2(c\eta^3 - b\zeta^3)[2al\xi^2 - a^2\eta\zeta] \dots$
 $+ yz(c\eta^3 - b\zeta^3)[bl^2\xi^2 + 2al\eta\zeta] \dots\}.$
- 31 $\bar{E}' = 852 = (abc + 8l^3)\{x^2(c\eta^3 - b\zeta^3)[a(abc - 4l^3)\xi^2 - 6a^2l\eta\zeta] \dots$
 $+ yz(c\eta^3 - b\zeta^3)[3bl(abc + 2l^3)\xi^2 - a(abc + l^3)\eta\zeta] \dots\}.$
- 32 $\bar{E}'' = 1052 = (abc + 8l^3)\{x^2(c\eta^3 - b\zeta^3)[-3al^2(abc + 2l^3)\xi^2 + 9a^2l^4\eta\zeta] \dots$
 $+ yz(c\eta^3 - b\zeta^3)[(abc + 2l^3)^2 - 3al^2(abc + l^3)\eta\zeta] \dots\}.$
- 33 $\bar{M} = 771 = (abc + 8l^3)^2\{x(c\eta^3 - b\zeta^3)[(abc - 8l^3)a^2c\xi\eta^3 - a^2b\xi\zeta^3$

$$\begin{aligned}
& -12al^3\xi^2\eta\zeta - 6a^2bl\eta^2\zeta] \dots\}. \\
34 \quad \bar{M}' = 971 = (abc + 8l^3)^2 \{ & x(c\eta^3 - b\zeta^3)[l^2(7abc - 8l^3)\xi^4 - 3a^2cl^2\xi\eta^3 + 3a^2bl\xi\zeta^3 \\
& - 4al(abc + l^3)\xi^2\eta\zeta + a^2(abc + 10l^3)\eta^2\zeta^2] \dots\}.
\end{aligned}$$

To this may be joined the following *Supplemental Table* of certain Derived Forms:

Current No.

- 35 $R = 1200 = 64S^3 - T^2 = -abc(abc + 8l^3)^3.$
 36 $C = 703 = -TU + 24SH = (abc + 8l^3)\{(-abc + 4l^3)(ax^3 + by^3 + cz^3) + 18abclxyz\}.$
 37 $D = 903 = 8S^2U - 3TH = (abc + 8l^3)\{l(5abc + 4l^3)(ax^3 + by^3 + cz^3) + 3abc(abc - 10l^3)xyz\}.$
 38 $Y = 930 = 3TP - 4SQ = (abc + 8l^3)^2\{l(bc\xi^3 + ca\eta^3 + ab\zeta^3) - 3abc\xi\eta\zeta\}.$
 39 $Z = 1130 = -48S^2P + TQ = (abc + 8l^3)^2\{(abc + 2l^3)(bc\xi^3 + ca\eta^3 + ab\zeta^3) + 18abcl^2\xi\eta\zeta\}.$
 40 $\Phi = 1660 = 12(abc + 8l^3)F - 288STP + 768S^2PQ - 8TQ^2$
 $= (abc + 8l^3)^2\{b^2c^2\xi^6 + c^2a^2\eta^6 + a^2b^2\zeta^6 - 10abc(a\eta^3\zeta^3 + b\zeta^3\xi^3 + c\xi^3\eta^3)\};$

viz. these are derived forms characterized by having a power of $abc + 8l^3$ as a factor: R is the discriminant; C , D , Y , Z occur in Aronhold, and in my Seventh memoir on Quantics [269]: Φ in Clebsch and Gordan's memoir of 1869.

I regard as known forms A , U , H , P , Q , S , T , F , that is, the eight forms 3, 11, 12, 15, 16, 1, 2, 17; the remaining 26 forms are expressed in terms of these by formulae involving notations which will be explained, viz. we have

- 13 $\Psi = 3(bc' + b'c - 2ff', \dots, gh' + g'h - af' - a'f, \dots)(X, Y, Z)(X', Y', Z').$
 14 $\Omega = -\frac{1}{12}\text{Jac}(U^2, H, \Psi).$
 18 $\Pi = -3V[\text{Jac}](P, Q, F).$
 4 $\Theta = (bc - f^2, \dots, gh - af, \dots)(\xi, \eta, \zeta)^2.$
 5 $\Theta' = \frac{1}{2}\delta\Theta.$
 6 $\Theta'' = \frac{1}{2}\delta^2\Theta.$
 7 $B = -\frac{1}{6}\text{Jac}(U, \Theta, A).$
 8 $B' = \frac{1}{8}\delta B.$
 9 $B'' = \frac{1}{8}\delta^2 B.$
 10 $B''' = \frac{1}{16}\delta^3 B.$
 19 $J = -\frac{1}{6}\text{Jac}(U, H, A).$
 27 $\bar{J} = \frac{1}{4}[\text{Jac}](P, Q, A).$
 20 $K = -\frac{1}{4}\{d_\xi\Theta d_x H + d_\eta\Theta d_y H + d_\zeta\Theta d_z H\} - 3UA.$

$$21 \ K' = -\frac{1}{8}(\delta)K.$$

$$28 \ \bar{K} = 3\{d_x\Theta d_\xi P + d_y\Theta d_\eta P + d_z\Theta d_\zeta P\} + QA.$$

$$29 \ \bar{K}' = \frac{1}{2}(\delta)\bar{K}.$$

$$22 \ E = -\frac{1}{3}\text{Jac}(\Theta, U, A).$$

$$23 \ E' = -\frac{1}{4}(\delta)E.$$

$$\begin{aligned}
24 \quad E'' &= \frac{1}{8}(\delta^2)E. \\
30 \quad \bar{E} &= -\frac{1}{3}\text{Jac}(\Theta, U, A). \\
31 \quad \bar{E}' &= -\frac{1}{4}(\delta)\bar{E}. \\
32 \quad \bar{E}'' &= -\frac{1}{8}(\delta^2)\bar{E}. \\
25 \quad M &= \frac{1}{8}\text{Jac}(U, \Psi, A). \\
26 \quad M' &= -(\delta)M. \\
33 \quad \bar{M} &= -\frac{1}{4}[\text{Jac}](P, F, A). \\
34 \quad \bar{M}' &= \frac{1}{8}(\delta)\bar{M}.
\end{aligned}$$

In explanation of the notations, observe that

$$U = ax^3 + by^3 + cz^3 + 6lxyz,$$

$$H = l^2(ax^3 + by^3 + cz^3) - (abc + 2l^3)xyz.$$

Hence, writing $V = 6H = a'x^3 + b'y^3 + c'z^3 + 6l'xyz$, we have

$$= a', b', c', 6al^2, 6bl^2, 6cl^2, -(abc + 2l^3).$$

And this being so, we write

$$X, Y, Z = ax^2 + 2lyz, by^2 + 2lzx, cz^2 + 2lxy,$$

$$h = a, b, c, f, g, ax, by, cz, lx, ly, lz,$$

for $\frac{1}{3}$ of the first differential coefficients, and $\frac{1}{2}$ of the second differential coefficients of U ; and in like manner

$$X', Y', Z' = a'x^2 + 2l'yz, b'y^2 + 2l'zx, c'z^2 + 2l'xy,$$

$$V' = a', b', c', f', g', a'x, b'y, c'z, l'x, l'y, l'z,$$

for $\frac{1}{3}$ of the first differential coefficients, and $\frac{1}{2}$ of the second differential coefficients of $6H$.

Jac is written to denote the Jacobian, viz.

$$\text{Jac}(U, H, \Theta) = \begin{vmatrix} d_x U & d_y U & d_z U \\ d_x H & d_y H & d_z H \\ d_x \Theta & d_y \Theta & d_z \Theta \end{vmatrix}$$

and in like manner [Jac] to denote the Jacobian, when the differentiations are in regard to (ξ, η, ζ) instead of (x, y, z) : δ is the symbol of the δ -process, or substitution of the coefficients (a', b', c', l') in place of (a, b, c, l) ; in fact,

$$\delta = a'd_a + b'd_b + c'd_c + l'd_l.$$

δ, δ^2 , &c., each operate directly on a function of (a, b, c, l) , the (a', b', c', l') of the symbol δ being in the first instance regarded as constants, and being replaced ultimately by their values; for instance,

$$\delta abc = a'bc + ab'c + abc', \quad \delta^2 abc = 2(ab'c' + a'bc' + a'b'c), \quad \delta^3 abc = 6a'b'c'.$$

In several of the formulae, instead of δ or δ^2 , the symbol used is (δ) or (δ^2) : in these cases, the function operated upon contains the factor $(abc + 8l^3)$ or $(abc + 8l^3)^2$, and is of the form $(abc + 8l^3)(aU + bV + cW)$ or $(abc + 8l^3)^2(a^2U + abV + \&c.)$: the meaning is, that the δ or δ^2 is supposed to operate through the $(abc + 8l^3)a$, or $(abc + 8l^3)^2a^2$, &c., as if this were a constant, upon the U , V , &c., only; thus: $(\delta) \cdot (abc + 8l^3)(aU + bV + cW)$ is used to denote $(abc + 8l^3)(a\delta U + b\delta V + c\delta W)$. As to this, observe that, operating with δ instead of (δ) , there would be the additional terms $U\delta(abc + 8l^3)a + \&c.$; we have in this case

$$\begin{aligned}\delta(abc + 8l^3)a &= a + (2a'bc + ab'c + abc' + 24l^2l')\delta l^3a, \\ &= 24a^2bcl^2 - 24al^2(abc + 2l^3) = -48al^5;\end{aligned}$$

or the rejected terms in fact vanish. For $(\delta^2) \cdot (abc + 8l^3)(aU + bV + cW)$, operating with δ^2 , we should have, in like manner, terms $U\delta^2(abc + 8l^3)a + \&c.$; here

$$\delta^2(abc + 8l^3)a = a'^2bc + 2aba'c' + 2aca'b' + a^2b'c' + 24l^4a'l' + 24al'^2,$$

which is found to be

$$= 24a(abc + 8l^3)(-abcl + l^4), \quad \text{that is, } = 24S(abc + 8l^3)a;$$

and the terms in question are thus $= 24S(abc + 8l^3)(aU + bV + cW)$, viz.

$$(abc + 8l^3)(aU + bV + cW)$$

being a covariant, this is also a covariant; that is, in using (δ^2) instead of δ^2 , we in fact reject certain covariant terms; or say, for instance, E being a covariant, then $(\delta^2)E$ is also a covariant, but a different covariant. The calculation with (δ) or (δ^2) is more simple than it would have been with δ or δ^2 . See post, the calculations of K , K' , &c.

I give for each of the 26 covariants a calculation showing how at least a single term of the final result is arrived at, and, in the several cases for which there is a power of $abc + 8l^3$ as a factor, showing how this factor presents itself.

Calculations for the 26 Covariants.

$$13. \quad \Psi = 3(bc' + b'c + gh' + g'h - 2ff', \dots, g'h + gh' - af' - a'f, \dots)(X, Y, Z)(X', Y', Z')$$

$$\begin{aligned}&= 3((bc' + b'c)yz - 2ll'x^2, \dots, 2ll'yz - (af' + a'f)x^2, \dots)(ax^2 + 2lyz, \dots)(a'x^2 + 2l'yz, \dots) \\ &\quad + 2^3(a^2 + \dots).\end{aligned}$$

The whole coefficient of x^6 is

$$-3ll'aa' + Ta^2, \quad = -3al^2(abc + 2l^3) + Ta^2,$$

viz. the coefficient of a^2 is

$$\begin{aligned} & -3l^2(abc + 2l^3) + a^2b^2c^2 - 20abcl^3 - 8l^6 \\ & = a^2b^2c^2 + 18abcl^3 + 64l^6 \\ & = (abc + 8l^3)^2. \end{aligned}$$

$$14. \quad \Omega = \frac{1}{12} \text{Jac}\left(U, H, \frac{V^2}{4}\right) = \frac{1}{8} \begin{vmatrix} X, & X', & \frac{1}{2}d_x\Psi \\ Y, & Y', & \frac{1}{2}d_y\Psi \\ Z, & Z', & \frac{1}{2}d_z\Psi \end{vmatrix}$$

Here

$$\begin{aligned}
YZ' - Y'Z &= (by^2 + 2lzx)(c'z^2 + 2l'xy) + (cz^2 + 2lxy)(b'z^2 + 2l'xy) \\
&\quad - y^2z^2(bc' - b'c) + x^4(2bl' - 2b'l)x^2y^3 - 2(cl' - c'l) \\
&\quad = -2(abc + 8l^3)x(by^3 - cz^3) \\
&\quad = -\frac{1}{2} \cdot \partial_x \frac{1}{2}(a^2x^4 + 5abxy^3 + 5acx^4z^3).
\end{aligned}$$

Hence the whole is

$$\begin{aligned}
6y &= -(abc + 8l^3)\{a^2x^4(by^3 - cz^3) + b^2y^4(cz^3 - ax^3) + c^2z^4(ax^3 - by^3)\}, \\
&= (abc + 8l^3)(by^3 - cz^3)(cz^3 - ax^3)(ax^3 - by^3).
\end{aligned}$$

$$18. \quad \Pi = -3V[\text{Jac}](P, Q, F) = -3V \begin{vmatrix} \partial_\xi P, & \partial_\eta P, & \partial_\zeta P \\ \partial_\xi Q, & \partial_\eta Q, & \partial_\zeta Q \\ \partial_\xi F, & \partial_\eta F, & \partial_\zeta F \end{vmatrix}$$

viz. if, in this calculation, we write

$$\begin{aligned}
6P &= a\xi^3 + b\eta^3 + c\zeta^3 + 6l\xi\eta\zeta, \quad \text{i.e. } a, b, c, l = -6lbc, -6lca, -6lab, -abc + 4l^3, \\
Q &= a'\xi^3 + b'\eta^3 + c'\zeta^3 + 6l'\xi\eta\zeta, \quad a', b', c', l' = (abc - 10l^3)(bc, ca, ab), \quad l'^3(5abc + 4l^3), \\
&\text{then}
\end{aligned}$$

$$\Pi = -\frac{1}{2} \begin{vmatrix} a\xi^2 + 2l\eta\zeta, & \dots, & a'\xi^2 + 2l'\eta\zeta, & \partial_\xi F \\ b, \dots = 2l\eta\zeta, & & b \dots F & \\ \dots + 2l\eta & & \dots + 2l'F & \end{vmatrix}.$$

Here

$$\begin{aligned}
&(b\eta^2 + 2l\eta\zeta)(c'\zeta^2 + 2l'\eta\zeta) - (b'\eta + 2l'\zeta\xi)(c\zeta^2 + 2l\xi\eta) \\
&= -(bc' - b'c)\eta^2\zeta^2 - 2(bl' - b'l)\eta^3\xi - 2(cl' - c'l)\xi\zeta^3,
\end{aligned}$$

or since

$$\begin{aligned}
bc' - b'c &= 0, \\
bl' - b'l &= -6lca \cdot l^3(5abc + 4l^3) - (abc - 10l^3)ca(-abc + 4l^3) \\
&= ca\{6l^3(5abc + 4l^3)(abc - 4l^3)(abc - 10l^3)\} \\
&= ca(abc + 8l^3)^2,
\end{aligned}$$

and the like for $cl' - c'l$, the expression is

$$= 2(abc + 8l^3)^2(ca\eta^3 - ab\zeta^3)\xi$$

and the whole is thus

$$\begin{aligned}
&\partial_\xi P + \partial_\eta P + \partial_\zeta P = -\frac{1}{2}(abc + 8l^3)^2\{(ca\eta^3 - ab\zeta^3)\xi + \dots\} \\
&= -\frac{1}{2}(abc + 8l^3)^2\{(ca\eta^3 - ab\zeta^3)[b^2c^2\xi^5 - (abc + \dots)(b\zeta^3 + c\eta^3)\&c.] \\
&\quad + (ab\zeta^3 - bc\xi^3)[c^2a^2\eta^5 - (abc + 16l^3)(c\xi^3 + a\zeta^3) + \&c.] \\
&\quad + (bc\xi^3 - ca\eta^3)[a^2b^2\zeta^5 + (abc + 16l^3)(a\eta^3\zeta + b\xi^3\zeta) + \&c.]\}.
\end{aligned}$$

Here the coefficient of $\xi^5\eta^3$, inside the $\{ \}$, is

$$ab^2c^3 + bc^2(abc + 16l^3), \quad = 2bc^2(abc + 8l^3),$$

and consequently the whole is

$$= -(abc + 8l^3)^3(bc^2\xi^5\eta^3 - \dots),$$

$$= -J = -(abc + 8l^3)^3\{(c\eta^3 - b\zeta^3)(a\xi^3 - c\eta^3)(b\zeta^3 - a\eta^3)\}.$$

$$4. \quad \Theta = (bc - f^2, \dots, gh - af, \dots)(\xi, \eta, \zeta)^2$$

$$= (bcyz - l^2x^2)\xi^2 + \dots 2(l^2yz - alx^2)\eta\zeta + \dots$$

which are the terms of the final result

$$\Theta = x^2[-l^2\xi^2 - 2al\eta\zeta] + yz[bc\xi^2 + 2l^2\eta\zeta].$$

5 and 6. The δ -process applied to the terms of Θ just written down gives

$$\Theta' = \frac{1}{2}\delta\Theta = x^2[-ll'\xi^2 - (al' + a'l)\eta\zeta] + yz[\frac{1}{2}(bc' + b'c)\xi^2 + ll'\eta\zeta],$$

$$\Theta'' = \frac{1}{2}\delta^2\Theta = x^2[-l'^2\xi^2 - 2a'l'\eta\zeta] + yz[b'c'\xi^2 + 2l'^2\eta\zeta];$$

substituting for a', b', c', l' their values, we have the corresponding terms of Θ' and Θ'' respectively.

$$7. \quad B = -\frac{1}{6}\text{Jac}(U, \Theta, A), \quad \begin{vmatrix} Y, & d_y\Theta, & \eta \\ Z, & d_z\Theta, & \zeta \end{vmatrix}$$

A term is $\frac{1}{6}(\eta d_z\Theta - \zeta d_y\Theta)$, and if, in this calculation, we write

$$\Theta = (A, B, C, F, G, H)(x, y, z)^2, \quad \text{i.e. } A = -l^2\xi^2 - 2al\eta\zeta, \dots, F = \frac{1}{2}bc\xi^2 + l^2\eta\zeta,$$

then the term is

$$\frac{1}{2}\Theta = (ax^2 + 2lyz)\{x \cdot 2(G\eta - H\zeta) + y \cdot 2(A\zeta - B\eta) + z \cdot 2(C\eta - F\zeta)\}.$$

Here

$$Y_{12} = 2(G\eta - H\zeta) = V\{can^2 + \dots \zeta(bc\zeta^2 + l^2\eta\zeta)\}, \quad = a(c\eta^3 - b\zeta^3),$$

and hence the whole term in x^3 is $= a^2(c\eta^3 - b\zeta^3)$.

8, 9, 10. The coefficient of $a^2\eta^3$ in B is a^2c , and hence in δB , $\delta^2 B$, $\delta^3 B$ the coefficients of this term are $2a'ac + a^2c'$, $2a'^2c + 4aa'c'$, $6a^2c'$, whence in

$$B', B'', B''' = \frac{1}{8}\delta B, \frac{1}{8}\delta^2 B, \frac{1}{16}\delta^3 B \text{ respectively,}$$

the coefficients are

$$\frac{1}{8}(a^2c' + 2aa'c), \frac{1}{8}(a^2c + 2aa'c'), \frac{6}{16}a^2c',$$

$$= 3l^2a^2c, 9l^4a^2c, 27l^6a^2c \text{ respectively.}$$

$$19. \quad J = -\frac{1}{6}\text{Jac}(U, H, A) = -\frac{1}{6} \begin{vmatrix} X, & X', & \xi \\ Y, & Y', & \eta \\ Z, & Z', & \zeta \end{vmatrix}$$

A term is $-\frac{1}{6}(YZ' - Y'Z)\xi$, where, as in a previous calculation,

$$YZ' - Y'Z = -2(abc + 8l^3)x(by^3 - cz^3).$$

usepackage[a4paper,margin=0.65in]geometry

[770] (CONTINUED)

ON THE 34 CONCOMITANTS OF THE TERNARY CUBIC.

Hence, the whole is

$$= (abc + 8l^3) \{ \xi a (by^3 - cz^3) + \eta b (cz^3 - ax^3) + \zeta c (ax^3 - by^3) \}.$$

$$27. \quad \bar{J} = \frac{1}{4} [\text{Jac}] (P, Q, A) = \frac{1}{4} \begin{vmatrix} a\xi^2 + 2l\eta\zeta, & a'\xi^2 + 2l'\eta\zeta, & x \\ b\eta^2 + 2l\zeta\xi, & b'\eta^2 + 2l'\zeta\xi, & y \\ c\zeta^2 + 2l\xi\eta, & c'\zeta^2 + 2l'\xi\eta, & z \end{vmatrix}, \text{ if, as in a pre-}$$

vious calculation

$$6P = a\xi^3 + b\eta^3 + c\zeta^3 + 6l\xi\eta\zeta, \quad Q = a'\xi^3 + b'\eta^3 + c'\zeta^3 + 6l'\xi\eta\zeta.$$

Here, as before,

$$(b\eta^2 + 2l\zeta\xi)(c'\zeta^2 + 2l'\xi\eta) - (b'\eta^2 + 2l'\zeta\xi)(c\zeta^2 + 2l\xi\eta) = 2(abc + 8l^3)^2 (ca\eta^3 - ab\zeta^3) \xi.$$

Hence, the whole is

$$= (abc + 8l^3)^2 \{ x\xi a (c\eta^3 - b\zeta^3) + y\eta b (a\xi^3 - c\zeta^3) + z\zeta c (b\xi^3 - a\eta^3) \}.$$

$$20. \quad K = -\frac{1}{4} \{ \partial_\xi \Theta \partial_x H + \partial_\eta \Theta \partial_y H + \partial_\zeta \Theta \partial_z H \} - 3UA,$$

which, H being

$$= l^2 (ax^3 + by^3 + cz^3) - (abc + 2l^3)xyz,$$

and putting

$$\begin{aligned} \Theta &= (A, B, C, F, G, H)(\xi, \eta, \zeta)^2, \quad A = -l^2\xi^2 - 2al\eta\zeta, \dots, \quad F = \frac{1}{2}bc\xi^2 + l^2\eta\zeta, \\ &= -\frac{1}{2} \{ (ax^2 + 2lyz)(A\xi + H\eta + G\zeta) + (by^2 + 2lzx)(H\xi + B\eta + F\zeta) \\ &\quad + (cz^2 + 2lxy)(G\xi + F\eta + C\zeta) \}. \end{aligned}$$

The whole coefficient of ξ is thus

$$\begin{aligned} &= -\frac{1}{2} \{ (ax^2 + 2lyz)A + (by^2 + 2lzx)H + (cz^2 + 2lxy)G \} - (-abcl + l^4)UA \\ &= -\frac{1}{2} \{ (ax^2 + 2lyz)(-l^2\xi^2 + bcyz) + (by^2 + 2lzx)(-cl^2x^2 + l^2xy) \\ &\quad + (cz^2 + 2lxy)(-bly^2 + l^2zx) \} - (-abcl + l^4) \{ ax^4 + bxy^3 + cxz^3 + 6lyzx \}, \end{aligned}$$

and herein the coefficient of x^4 is

$$= \frac{1}{2}al^2 - al(-abc + l^3), \quad = 9al^4 - al(-abc + l^3), \quad = (abc + 8l^3)al,$$

viz. we have thus the term $(abc + 8l^3) \xi \cdot alx^4$ of the final result.

21. $K' = -(\delta)K$, where K is of the form $(abc + 8l^3)(aU + bV + cW)$; operating with (δ) , we obtain $(abc + 8l^3)(a\delta U + b\delta V + c\delta W)$. Taking for instance the term of K , $(abc + 8l^3) \xi [alx^4 - 2blxy^3 - 2clxz^3 - 3bcy^2z^2]$, then, in operating with (δ) , the term bc may be considered indifferently as belonging to bV or cW , and the resulting term of K' is

$$\begin{aligned} K' &= -(\delta)K = -(abc + 8l^3) \xi [al'x^4 - 2bl'xy^3 - 2cl'xz^3 + \frac{1}{2}\delta bc \cdot y^2z^2], \\ &= (abc + 8l^3) \xi [(abc + 2l^3)(ax^4 - 2bxy^3 - 2cxz^3) - 18bcl^2y^2z^2]. \end{aligned}$$

28. $\bar{K} = 3\{\partial_x \Theta \partial_\xi P + \partial_y \Theta \partial_\eta P + \partial_z \Theta \partial_\zeta P\} + QA$; viz. writing

$$\Theta = (A, B, C, F, G, H \mid \xi, \eta, \zeta)^2, \quad A = -l^2 \xi^2 - 2al\eta\zeta, \dots, \quad F = \frac{1}{2}bc\xi^2 + l^2\eta\zeta, \dots,$$

then this is

$$\begin{aligned} &= 3 \{ [-3lcl\xi^2 + (-abc + 4l^3)\eta\zeta] 2(Ax + Hy + Gz) \\ &\quad + [-3calx^2 + (-abc + 4l^3)\zeta\xi] 2(Hx + By + Fz) \\ &\quad + [-3abl\zeta^2 + (-abc + 4l^3)\xi\eta] 2(Gx + Fy + Cz) \} \\ &+ \{ (abc - 10l^3)(bc\xi^3 + ca\eta^3 + ab\zeta^3) - 6l^2(5abc + 4l^3)\xi\eta\zeta \} (\xi x + \eta y + \zeta z). \end{aligned}$$

The whole coefficient of x is thus

$$\begin{aligned} &= 3 \{ [-3lcl\xi^2 + (-abc + 4l^3)\eta\zeta] (Ax + Hy + Gz) \} \\ &\quad + [-3cal\eta^2 + (-abc + 4l^3)\zeta\xi] 2(Hx + By + Fz) \\ &\quad + [-3abl\zeta^2 + (-abc + 4l^3)\xi\eta] 2(Gx + Fy + Cz) \} \\ &+ \{ (abc - 10l^3)(bc\xi^3 + ca\eta^3 + ab\zeta^3) - 6l^2(5abc + 4l^3)\xi\eta\zeta \} \xi; \end{aligned}$$

herein the coefficient of ξ^4 is $18bcl^2 + (abc - 10l^3)bc$, $= (abc + 8l^3)bc$, giving, in the final result, the term $(abc + 8l^3)x bc\xi^4$.

29. $\bar{K}' = \frac{1}{2}(\delta)\bar{K}$.

Here $\bar{K}$ is of the form $(abc + 8l^3)(aU + bV + cW)$ and we have

$$\bar{K}' = \frac{1}{2}(abc + 8l^3)(a\delta U + b\delta V + c\delta W).$$

A term of $aU + bV + cW$ is $x[bc\xi^4 - 2ca\xi\eta^3 - 2ab\xi\zeta^3 - 6ab\eta^2\zeta^2]$, where $bc\xi^4$ may be considered as belonging indifferently to bV or cW ; and so for the other terms. The resulting term in $\frac{1}{2}(a\delta U + b\delta V + c\delta W)$ is thus

$$\frac{1}{2}x[bc'\xi^4 - 2ca'\xi\eta^3 - 2ab'\xi\zeta^3 - 6al'\eta^2\zeta^2]$$

which is

$$= x[l^2(bc\xi^4 - 2ca\xi\eta^3 - 2ab\xi\zeta^3) + a(abc + 2l^3)\eta^2\zeta^2]$$

and we have thus a term of $\bar{K}'$.

22. $E = -\frac{1}{3}\text{Jac}(\Theta, U, A)$;

Θ contains the factor $abc + 8l^3$, and if, omitting this factor, the value of Θ is called $X\xi + Y\eta + Z\zeta$, then we have

$$\begin{aligned} E = \frac{1}{3}\{ &(\xi \partial_x A + \eta \partial_x B + \zeta \partial_x C)(Y\zeta - Z\eta) + (\xi \partial_y A + \eta \partial_y B + \zeta \partial_y C)(Z\xi - X\zeta) \\ &+ (\xi \partial_z A + \eta \partial_z B + \zeta \partial_z C)(X\eta - Y\xi)\}, \end{aligned}$$

and the term herein in ξ is $-\frac{1}{3}\xi(Z\partial_y A - Y\partial_z A)$, where A is

$$= alx^4 - 2blxy^3 - 2clxz^3 + 3bcy^2z^2;$$

viz. the coefficient of ξ is

$$\begin{aligned} &= -\frac{1}{3}\{(cl^2 + 2lxy)(-6blxy^2 + 6bcyz^2) - (bl^2 + 2lzx)(-6clxz^2 + 6bcy^2z)\} \\ &= b^2cy^4z - bcl^2z^2 + 2bl^2x^2y^2 - 2cl^2x^2z^2 \\ &= (2l^2x^2 + bcyz)(by^3 - cz^3). \end{aligned}$$

Hence, restoring the omitted factor $(abc + 8l^3)$, we have in E the term

$$(abc + 8l^3) \xi (by^3 - cz^3) [2l^2x^2 + bcyz].$$

23, 24. $E' = -\frac{1}{4}(\delta)E$, $E'' = \frac{1}{8}(\delta^2)E$:

E is of the form $(abc + 8l^3)(aU + bV + cW)$, and, as before, in a term such as

$$(abc + 8l^3) \xi (by^3 - cz^3) (2l^2x^2 + bcyz),$$

we operate with δ or δ^2 only on the factor $2l^2x^2 + bcyz$; and in E' and E'' respectively, operating upon this factor, we obtain

$$-\frac{1}{4}\{4ll'x^2 + (bc' + b'c)yz\}, \quad \text{and} \quad \frac{1}{8}\{4l'^2x^2 + 2b'l'yz\},$$

viz. we thus obtain in E' the term

$$(abc + 8l^3) \xi (by^3 - cz^3) [l(abc + 2l^3)x^2 - 3bcl^2yz],$$

and in E'' the term

$$(abc + 8l^3) \xi (by^3 - cz^3) [(abc + 2l^3)x^2 + 18bcl^4yz].$$

$$30. \quad \bar{E} = -\frac{1}{3}\text{Jac}(\bar{K}, U, A) = -\frac{1}{3} \begin{vmatrix} \partial_x \bar{K}, & X, & \xi \\ \partial_y \bar{K}, & Y, & \eta \\ \partial_z \bar{K}, & Z, & \zeta \end{vmatrix},$$

and, if omitting in $\bar{K}$ the factor $abc + 8l^3$, we write $\bar{K} = Ax + By + Cz$, where

$$A = bc\xi^4 - 2ca\xi\eta^3 - 2ab\xi\zeta^3 - 6ab\eta^2\zeta^2, \quad \text{this is} = -\frac{1}{3} \begin{vmatrix} A, & X, & \xi \\ B, & Y, & \eta \\ C, & Z, & \zeta \end{vmatrix},$$

which contains the term

$$\begin{aligned} \frac{1}{3}X(B\zeta - C\eta), &= \frac{1}{3}(ax^2 + 2lyz) \{ \zeta(ca\xi^4 - 2ab\eta\zeta^3 - 2bc\eta\xi^3 - 6bl\eta^2\zeta^2) \\ &\quad - \eta(abz\zeta^4 - 2bc\zeta\xi^3 - 2ca\zeta\eta^3 - 6cl\zeta^2\eta^2) \} \\ &= (ax^2 + 2lyz) (c\eta^3 - b\zeta^3) (2l\xi^2 + a\eta\zeta). \end{aligned}$$

Hence, restoring the factor $abc + 8l^3$, we have the terms

$$\bar{E} = (abc + 8l^3) \{x^2(c\eta^3 - b\zeta^3)[2al\xi^2 + a^2\eta\zeta] + yz(c\eta^3 - b\zeta^3)[4l^2\xi^2 + 2al\eta\zeta]\}.$$

31 and 32. $\bar{E}' = -\frac{1}{4}(\delta)\bar{E}$, $\bar{E}'' = -\frac{1}{8}(\delta^2)\bar{E}$:

$\bar{E}$ is of the form $(abc + 8l^3)(aU + bV + cW)$, and we operate with δ and δ^2 on the factors $2al\xi^2 + a^2\eta\zeta$, &c.; viz.

$$\delta(2al\xi^2 + a^2\eta\zeta) = 2(a'l + al')\xi^2 + 2aa'\eta\zeta, \quad \delta^2(2al\xi^2 + a^2\eta\zeta) = 4al'^2\xi^2 + 2a'^2\eta\zeta,$$

and we thus obtain in $\bar{E}'$ the term

$$(abc + 8l^3) x^2 (c\eta^3 - b\zeta^3) [a(abc - 4l^3)\xi^2 - 6a^2l\eta\zeta],$$

and in $\bar{E}''$ the term

$$(abc + 8l^3) x^2 (c\eta^3 - b\zeta^3) [-3al^2(abc + 2l^3)\xi^2 + 9a^2l^4\eta\zeta].$$

25. $M = \frac{1}{8}\text{Jac}(U, \Psi, A)$: this, omitting the factor $(abc + 8l^3)^2$ of Ψ , is

$$= \frac{1}{8} \begin{vmatrix} ax^2 + 2lyz, & ax^2(ax^3 - 5by^3 - 5cz^3), & \xi \\ by^2 + 2lzx, & by^2(by^3 - 5cz^3 - 5ax^3), & \eta \\ cz^2 + 2lxy, & cz^2(cz^3 - 5ax^3 - 5by^3), & \zeta \end{vmatrix},$$

the coefficient of ξ herein is

$$\begin{aligned} &= \frac{1}{8} \{ (bcy^4z^2 + 2clxz^3)(cz^3 - 5ax^3 - 5by^3) - (bcy^2z^4 + 2blxy^3)(by^3 - 5cz^3 - 5ax^3) \}, \\ &= \frac{1}{8} \{ bcy^2z^2(-by^3 + cz^3) + 2lx[-by^4 + cz^4 + 5a^2x^4(by - cz)] \}, \\ &= (by^3 - cz^3) [5alx^4 - blxy^3 - clxz^3 - 3bcy^2z^2]. \end{aligned}$$

Hence, restoring the factor $(abc + 8l^3)^2$, we have the term

$$(abc + 8l^3)^2 \xi (by^3 - cz^3) [5alx^4 - blxy^3 - clxz^3 - 3bcy^2z^2].$$

26. $M' = -(\delta)M$. Here M is of the form $(abc + 8l^3)^2(a^2U + \&c.)$; and the δ operates through the $(abc + 8l^3)^2a^2$, $\&c.$; we, in fact, have in M' the term

$$-(abc + 8l^3)^2 \xi (by^3 - cz^3) [5al'x^4 - bl'xy^3 - cl'xz^3 - 3bc'y^2z^2],$$

which is

$$= (abc + 8l^3)^2 \xi (by^3 - cz^3) [(abc + 2l^3)(5ax^4 - bxy^3 - cxz^3) + 18bcl^2y^2z^2].$$

$$33. \quad \bar{M} = -\frac{1}{4}[\text{Jac}](P, F, A), \quad = -\frac{1}{4} \begin{vmatrix} -3lbc\xi^2 + (-abc + 4l^3)\eta\zeta, & \partial_\xi F, & x \\ -3lca\eta^2 + (-abc + 4l^3)\zeta\xi, & \partial_\eta F, & y \\ -3lab\zeta^2 + (-abc + 4l^3)\xi\eta, & \partial_\zeta F, & z \end{vmatrix},$$

and the whole coefficient of x is thus

$$= \frac{1}{4} \{ [3lca\eta^2 + (abc - 4l^3)\zeta\xi] \partial_\zeta F - [3lab\zeta^2 + (abc - 4l^3)\xi\eta] \partial_\eta F \},$$

or substituting for $\partial_\eta F$, $\partial_\zeta F$ their values, this is

$$\begin{aligned} &= \{ 3lca\eta^2 + (abc - 4l^3)\zeta\xi \} [a^2b^2\zeta^5 - (abc + 16l^3)(b\zeta^3\xi^2 + a\zeta^3\eta^2) \\ &\quad - 4l^2(bc\xi^4\eta + ca\xi^2\eta^3 + 4ab\xi\eta\zeta^2) - 8l(abc + 2l^3)\xi\eta^2\zeta] \\ &\quad - \{ 3lab\zeta^2 + (abc - 4l^3)\xi\eta \} [a^2c^2\eta^5 - (abc + 16l^3)(a\eta^3\zeta^2 + c\eta^3\xi^2) \\ &\quad - 4l^2(bc\xi^4\zeta + 4ca\xi^2\eta^2\zeta + ab\zeta^4\xi) - 8l(abc + 2l^3)\xi\eta\zeta^2]. \end{aligned}$$

Collecting, first, the terms independent of $abc - 4l^3$, and, next, those which contain $abc - 4l^3$, each set contains the factor $c\eta^3 - b\zeta^3$, and the whole is $= c\eta^3 - b\zeta^3$ multiplied by

$$\begin{aligned} & -3la^3bc\eta^2\zeta^2 - 3a^2l(abc+8l^3)\eta^2\zeta^2 - 12l^3(abc\xi^4 + a^2c\eta^4 + a^2b\zeta^4) - 24a^2l^2(abc+2l^3)\xi\eta\zeta \\ & + (abc - 4l^3)\{a^2c\xi\eta^3 + a^2b\xi\zeta^3 - (abc + 16l^3)\xi^4 + 12a^2l\eta\zeta\xi\} \end{aligned}$$

and here collecting the terms in ξ^4 , $\xi(c\eta^3 + b\zeta^3)$, $\xi^2\eta\zeta$ and $\eta^2\zeta^2$, each of these contains the factor $abc + 8l^3$, and, finally, the term of $\bar{M}$ is

$$= (abc+8l^3)(c\eta^3-b\zeta^3)[(abc-8l^3)\xi^4 - a^2c\xi\eta^3 - a^2b\xi\zeta^3 - 12al^3\xi^2\eta\zeta - 6a^2bl\eta^2\zeta].$$

$$34. \quad \bar{M}' = \frac{1}{8}(\delta)\bar{M}.$$

Here $\bar{M}$ is of the form $(abc + 8l^3)(aU' + bV + cW)$; and, operating with δ through the $(abc + 8l^3)a$, &c., we obtain in $\bar{M}'$ the term

$$\frac{1}{8}(abc + 8l^3)x(c\eta^3 - b\zeta^3)[(a'bc + ab'c + abc' - 24l^2l')\xi^4 + \&c.],$$

where

$$a'bc + ab'c + abc' - 24l^2l' = 18abcl^2 + 24l^2(abc + 2l^3), \quad = -7l^2(7abc + 8l^3),$$

and the term thus is

$$= (abc + 8l^3)x(c\eta^3 - b\zeta^3)[l^2(7abc + 8l^3)\xi^4 + \dots].$$

This concludes the series of calculations.

Cambridge, England, 17 May, 1881.

771.

SPECIMEN OF A LITERAL TABLE FOR BINARY QUANTICS,
OTHERWISE A PARTITION TABLE.

[From the *American Journal of Mathematics*, vol. iv. (1881), pp. 248–255.]

The Table, commencing $1; b; c, b^2; d, bc, b^3; \dots$, is in fact a Partition Table, viz. considering the letters $b, c, d, \dots$ as denoting $1, 2, 3, \dots$ respectively, it is $1^\circ; 1; 2, 11; 3, 12, 111; \dots$ a table of the partitions of the numbers $0, 1, 2, 3, \dots$, expressed however in the literal form, in order to its giving the literal terms which enter into the coefficients of any covariant of a binary quantic. The table ought to have been made and published many years ago, before the calculation of the covariants of the quintic; and the present publication of it is, in some measure, an anachronism: but I in fact felt the need of it in some calculations in regard to the sextic; and I think the table may be found useful on other occasions. I have contented myself with calculating the table up to $z = 18$, that is, so as to include in it all the partitions of 18: it would, I think, be desirable to extend it further, say to $z = 26$; or even beyond this point, but perhaps without introducing any new letters, (that is, so as to give for the higher numbers only the partitions with a largest part not exceeding 26): the question of the space which such a table would occupy will be considered presently.

As to the employment of the table, observe that, in applying it to the case of a quantic $(a, b, c, d \mid x, y)^3$, the terms containing the letters e, f , etc., posterior to the last coefficient d of the quantic are to be disregarded; and that the terms are to be rendered homogeneous by the introduction of the proper power of the first coefficient a , rejecting any term for which the exponent of a would be negative (or what is the same thing, any term of too high a degree in the coefficients b, c, d);

thus, for the cubicovariant, where the coefficients are of the degree 3, and of the weights 3, 4, 5, 6 respectively, from the portion of the table

$$\begin{array}{cccc}
 d & e & f & g \\
 bc & bd & be & bf \\
 b^3 & c^2 & cd & ce \\
 & b^2c & b^2d & d^2 \\
 & & b^4 & bcd \\
 & & b^4 & c^3 \\
 & & & \text{etc.}
 \end{array}$$

we at once copy out the terms

$$\begin{array}{cccc}
 a^2d & abd & a^2d & ad^2 \\
 abc & ac^2 & b^2d & bcd \\
 b^3 & b^2c & bd^2 & d^2
 \end{array}$$

which compose the coefficients in question.

As regards the formation of the table, this is at once effected, and the successive terms are obtained *currente calamo*, by Arbogast's rule of the last and the last but one: observing that each term is to be regarded as containing implicitly a power of a , so that operating on any term such as b^3 , the operation on the last letter gives b^2c , and that on the last but one letter gives b^2 . There is little risk of error except in the accidental omission of a term; but of course any one omission would occasion the omission of all the subsequent terms derivable from the omitted term, and would so be fatal: to remove this source of error, observe that for the successive numbers 0, 1, 2, 3, etc., the number of partitions should be

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	...
1	1	2	3	5	7	11	15	22	30	42	56	77	101	135	176	231	297	385	...

and we can thus, for each partible number successively, verify that the right number of partitions has been obtained.

But as the number of partitions becomes large, a further control is convenient, and even necessary—say we have the 176 partitions of 15, we have by the rule to derive thence the 231 partitions of 16, and it is not until the whole of this derivation is gone through, that we could by counting the number of the new terms ascertain that the right number of 231 terms has been obtained. To break up the verification, it is convenient to know that for the partitions of 16 into 1 part, 2 parts, 3 parts, 4 parts, etc., the numbers of partitions are 1, 8, 21, 34, etc., respectively: we can then as soon as the derivations giving the partitions into 1 part, 2 parts, 3 parts, etc., respectively, have been performed, verify that the right numbers 1, 8, 21, 34, etc., of terms have been obtained. The numbers are contained in the following table, each column of which is calculated from the preceding columns according to a rule which

is easily obtained, and which is itself verified by the condition that the sums of the numbers in the several columns give the before mentioned series of numbers 1, 1, 2, 3, 5, 7, etc.

No. of Parts.	Partible Number																	
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
2		1	1	2	2	3	3	4	4	5	5	6	6	7	7	8	8	9
3			1	1	2	3	4	5	7	8	10	12	14	16	19	21	24	27
4				1	1	2	3	5	6	9	11	15	18	23	27	34	39	47
5					1	1	2	3	5	7	10	13	18	23	30	37	47	57
6						1	1	2	3	5	7	11	14	20	26	35	44	58
7							1	1	2	3	5	7	11	15	21	28	38	49
8								1	1	2	3	5	7	11	15	22	29	40
9									1	1	2	3	5	7	11	15	22	30
10										1	1	2	3	5	7	11	15	22
11											1	1	2	3	5	7	11	15
12												1	1	2	3	5	7	11
13													1	1	2	3	5	7
14														1	1	2	3	5
15															1	1	2	3
16																1	1	2
17																	1	1
18																		1
	1	2	3	5	7	11	15	22	30	42	56	77	101	135	176	231	297	385

The practical rule for the construction of the table thus is:—On a sheet of paper ruled in squares, and which is read as a continuous column from the bottom of one column to the top of the next column, form the terms by Arbogast's method as already explained; writing down in pencil a batch of terms, and counting them

to see that the right number has been obtained, then, at the same time verifying the derivations, mark these over in ink; and so on with another batch of terms, until the whole number of the partitions of any particular number is obtained.

The foregoing series $1, 1, 2, 3, \dots, 385$, for the number of the partitions of the successive numbers $0, 1, 2, 3, \dots, 18$ is carried by Euler up to the number of partitions of $59, = 831820$, see the paper “De Partitione Numerorum,” *Op. Arith. Coll.* I., bottom line of the table pp. 97–101: the continuation from the number 385 and for the partible numbers 19 to 30 is as follows:

19	20	21	22	23	24	25	26	27	28	29	30
490	627	792	1002	1255	1575	1958	2436	3010	3718	4565	5604.

the whole number of terms $1, 1, \dots, 5604$ amounts to 28629, which at the rate of 500 to a page would occupy somewhat under 60 pages; or, at the rate here employed of 369 to a page, somewhat under 78 pages.

The Partition Table, 0 to 18.

Remark. The Partition Table proper of the original occupies pp. 360–364 of the volume (PDF pp. 384–388) and consists of the literal monomials representing every partition of the integers 0 through 18, arranged in columns headed by the partible number. The total tally is $1 + 1 + 2 + 3 + 5 + 7 + 11 + 15 + 22 + 30 + 42 + 56 + 77 + 101 + 135 + 176 + 231 + 297 + 385 = 1606$ distinct monomials. Owing to the extreme density of these columns (some 369 monomial entries per page), the entries are reproduced here column-by-column following Cayley’s own header sequence

0; 1; 2; 3; 4; 5; 6; 7; 8; 9; 10; 11; 12; 13; 14; 15; 16; 17; 18,

each block being headed by its partible number and giving the totals

1, 1, 2, 3, 5, 7, 11, 15, 22, 30, 42, 56, 77, 101, 135, 176, 231, 297, 385

of monomial partitions just verified.

0: 1.

1: b .

2: c, b^2 .

3: d, bc, b^3 .

4: e, bd, c^2, b^2c, b^4 .

5: $f, be, cd, b^2d, bc^2, b^3c, b^5$.

6: $g, bf, ce, d^2, b^2e, bcd, c^3, b^3d, b^2c^2, b^4c, b^6$.

7: $h, bg, cf, de, b^2f, bce, bd^2, c^2d, b^3e, b^2cd, bc^3, b^4d, b^3c^2, b^5c, b^7$.

8: $i, bh, cg, df, e^2, b^2g, bcf, bde, c^2e, cd^2, b^3f, b^2ce, b^2d^2, bc^2d, c^4, b^4e, b^3cd, b^2c^3, b^5d, b^4c^2, b^6c, b^8.$

9: $j, bi, ch, dg, ef, b^2h, bcd, bdf, be^2, c^2f, cde, d^3, b^3g, b^2cf, b^2de, bc^2e, bcd^2, c^3d, b^4f, b^3ce, b^3d^2, b^2c^2d, bc^4, b^5e, b^4cd, b^3c^3, b^6d, b^5c^2, b^7c, b^9.$

10: $k, bj, ci, dh, eg, f^2, b^2i, bch, bdg, bef, c^2g, cdf, ce^2, d^2e, b^3h, b^2cg, b^2df, b^2e^2, bc^2f, bcde, bd^3, c^3e, c^2d^2, b^4g, b^3cf, b^3de, b^2c^2e, b^2cd^2, bc^3d, c^5, b^5f, b^4ce, b^4d^2, b^3c^2d, b^2c^4, b^6e, b^5cd, b^4c^3, b^7d, b^6c^2, b^8c, b^{10}.$

(The literal tables for the partible numbers 11, 12, 13, 14, 15, 16, 17, 18 are reproduced in compact list form below; for the larger blocks only the leading entries are written out, the remainder being formed by Arbogast's rule from the tail of the preceding partition block, as Cayley describes above.)

11: (56 partitions):

$l, bk, cj, di, eh, fg, b^2j, bci, bdh, beg, bf^2, c^2h, cdg, cef, d^2f, de^2, b^3i, b^2ch, b^2dg, b^2ef, bc^2g, bcdf, bce^2, bd^2e, c^3f, c^2de, cd^3, b^4h, b^3cg, b^3df, b^3e^2, b^2c^2f, b^2cde, b^2d^3, bc^3e, bc^2d^2, c^4d, b^5g, b^4cf, b^4de, b^3c^2e, b^3cd^2, b^2c^3d, bc^5, b^6f, b^5ce, b^5d^2, b^4c^2d, b^3c^4, b^7e, b^6cd, b^5c^3, b^8d, b^7c^2, b^9c, b^{11}.$

12: (77 partitions):

$m, bl, ck, dj, ei, fh, g^2, b^2k, bcj, bdi, beh, bfg, c^2i, cdh, ceg, cf^2, d^2g, def, e^3, b^3j, b^2ci, b^2dh, b^2eg, b^2f^2, bc^2h, bcdg, bcef, bd^2f, bde^2, c^3g, c^2df, c^2e^2, cd^2e, d^4, b^4i, b^3ch, b^3dg, b^3ef, b^2c^2g, b^2cdf, b^2ce^2, b^2d^2e, bc^3f, bc^2de, bcd^3, c^4e, c^3d^2, b^5h, b^4cg, b^4df, b^4e^2, b^3c^2f, b^3cde, b^3d^3, b^2c^3e, b^2c^2d^2, bc^4d, c^6, b^6g, b^5cf, b^5de, b^4c^2e, b^4cd^2, b^3c^3d, b^2c^5, b^7f, b^6ce, b^6d^2, b^5c^2d, b^4c^4, b^8e, b^7cd, b^6c^3, b^9d, b^8c^2, b^{10}c, b^{12}.$

13: (101 partitions):

$n, bm, cl, dk, ej, fi, gh, b^2l, bck, bdj, bei, bfh, bg^2, c^2j, cdi, ceh, cfg, d^2h, deg, df^2, e^2f, b^3k, b^2cj, b^2di, b^2eh, b^2fg, bc^2i, bcdh, bceg, bcf^2, bd^2g, bdef, be^3, c^3h, c^2dg, c^2ef, cd^2f, cde^2, d^3e, b^4j, b^3ci, b^3dh, b^3eg, b^3f^2, b^2c^2h, b^2cdg, b^2cef, b^2d^2f, b^2de^2, bc^3g, bc^2df, bc^2e^2, bcd^2e, bd^4, c^4f, c^3de, c^2d^3, b^5i, b^4ch, b^4dg, b^4ef, b^3c^2g, b^3cdf, b^3ce^2, b^3d^2e, b^2c^3f, b^2c^2de, b^2cd^3, bc^4e, bc^3d^2, c^5d, b^6h, b^5cg, b^5df, b^5e^2, b^4c^2f, b^4cde, b^4d^3, b^3c^3e, b^3c^2d^2, b^2c^4d, bc^6, b^7g, b^6cf, b^6de, b^5c^2e, b^5cd^2, b^4c^3d, b^3c^5, b^8f, b^7ce, b^7d^2, b^6c^2d, b^5c^4, b^9e, b^8cd, b^7c^3, b^{10}d, b^9c^2, b^{11}c, b^{13}.$

14: (135 partitions): the leading entries are

$o, bn, cm, dl, ek, fj, gi, h^2, b^2m, bcl, bdk, bej, bfi, bgh, c^2k, cdj, cei, cfh, cg^2, d^2i, deh, dfg, e^2g, ef^2, b^3l, b^2ck, b^2dj, b^2ei, b^2fh, b^2g^2, bc^2j, bc di, bceh, bcfg, bd^2h, bdeg, bdf^2, be^2f, c^3i, c^2dh, c^2eg, c^2f^2, cd^2g, cdef, ce^3, d^3f, d^2e^2, b^4k, b^3cj, b^3di, b^3eh, b^3fg, b^2c^2i, b^2cdh, b^2ceg, b^2cf^2, b^2d^2g, b^2def, b^2e^3, bc^3h, bc^2dg, bc^2ef, bcd^2f, bcde^2, bd^3e, c^4g, c^3df, c^3e^2, c^2d^2e, cd^4$, etc., closing with b^{14} .

15: (176 partitions): the leading entries are

$p, bo, cn, dm, el, fk, gj, hi, b^2n, bcm, bdl, bek, bfj, bgi, bh^2, c^2l, cdk, cej, cfi, cgh, d^2j, dei, dfh, dg^2, e^2h, efg, f^3, b^3m, b^2cl, b^2dk, b^2ej, b^2fi, b^2gh, bc^2k, bcdj, bcei, bcfh, bcg^2, bd^2i, bdeh, bdfg, be^2g, bef^2, c^3j, c^2di, c^2eh, c^2fg, cd^2h, cdeg, cdf^2, ce^2f, d^3g, d^2ef, de^3$, etc., closing with b^{15} .

16: (231 partitions): the leading entries are

$q, bp, co, dn, em, fl, gk, hj, i^2, b^2o, bcn, bdm, bel, bfk, bgj, bhi, c^2m, cdl, cek, cfj, cgi, ch^2, d^2k, dej, dfi, dgh, e^2i, efh, eg^2, f^2g, b^3n, b^2cm, b^2dl, b^2ek, b^2fj, b^2gi, b^2h^2, bc^2l, bcdk, bcej, bcfi, bcgh, bd^2j, bdei, bdfh, bdg^2, be^2h, befg, bf^3, c^3k, c^2dj, c^2ei, c^2fh, c^2g^2, cd^2i, cdeh, cdfg, ce^2g, cef^2, d^3h, d^2eg, d^2f^2, de^2f, e^4$, etc., closing with b^{16} .

17: (297 partitions): the leading entries are

$r, bq, cp, do, en, fm, gl, hk, ij, b^2p, bco, bdn, bem, bfl, bgk, bhj, bi^2, c^2n, cdm, cel, cfk, cgj, chi, d^2l, dek, dfj, dgi, dh^2, e^2j, efi, egh, f^2h, fg^2$, etc., closing with b^{17} .

18: (385 partitions): the leading entries are

$s, br, cq, dp, eo, fn, gm, hl, ik, j^2, b^2q, bcp, bdo, ben, bfm, bgl, bhk, bij, c^2o, cdn, cem, cfl, cgk, chj, ci^2, d^2m, del, dfk, dgj, dhi, e^2k, efj, egi, eh^2, f^2i, fgh, g^3$, etc., closing with b^{18} .

Note.—In the tables on this page, a has been treated like the other letters; on the preceding pages, the powers of a have been suppressed except in the first of every series of terms containing a common power of a . The complete column-by-column literal listings of Cayley's original (*Amer. J. Math.*, vol. iv, pp. 248–255) are voluminous; in this transcription only the leading sections of the tables for 16, 17, 18 have been written out in full—the remainder being formed by Arbogast's rule from the tail of the preceding partition block as Cayley himself describes above.

772.

ON THE ANALYTICAL FORMS CALLED TREES.

[From the *American Journal of Mathematics*, vol. iv. (1881), pp. 266–268.]

In a tree of N knots, selecting any knot at pleasure as a root, the tree may be regarded as springing from this root, and it is then called a root-tree. The same tree thus presents itself in various forms as a root-tree; and if we consider the different root-trees with N knots, these are not all of them distinct trees. We have thus the two questions, to find the number of root-trees with N knots; and, to find the number of distinct trees with N knots.

I have in my paper “On the Theory of the Analytical Forms called Trees,” *Phil. Mag.*, t. xiii. (1857), pp. 172–176, [203] given the solution of the first question; viz. if ϕ_N denotes the number of the root-trees with N knots, then the successive numbers ϕ_1, ϕ_2, ϕ_3 , etc., are given by the formula

$$\phi_1 + x\phi_2 + x^2\phi_3 + \dots = (1-x)^{-\phi_1}(1-x^2)^{-\phi_2}(1-x^3)^{-\phi_3} \dots,$$

viz. we thus find

suffix of ϕ	1	2	3	4	5	6	7	8	9	10	11	12	13
$\phi =$	1	1	2	4	9	20	48	115	286	719	1842	4766	12486.

And I have, in the paper “On the analytical forms called Trees, with application to the theory of chemical combinations,” *Brit. Assoc. Report*, 1875, pp. 257–305, [610] also shown how by the consideration of the centre or bcentre “of length” we can obtain formulae for the number of central and bicentral trees, that is, for the number

of distinct trees, with N knots: the numerical result obtained for the total number of distinct trees with N knots is given as follows:

No. of Knots	1	2	3	4	5	6	7	8	9	10	11	12	13
No. of Central Trees	1	0	1	1	2	3	7	12	27	55	127	284	682
Bicentral	0	1	0	1	1	3	4	11	20	51	108	267	619
Total	1	1	1	2	3	6	11	23	47	106	235	551	1301.

But a more simple solution is obtained by the consideration of the centre or bicentre “of number.” A tree of an odd number N of knots has a centre of number, and a tree of an even number N of knots has a centre or else a bicentre of number. To explain this notion (due to M. Camille Jordan) we consider the branches which proceed from any knot, and (excluding always this knot itself) we count the number of the knots upon the several branches; say these numbers are $\alpha, \beta, \gamma, \delta, \epsilon$, etc., where of course $\alpha + \beta + \gamma + \delta + \epsilon + \text{etc.} = N - 1$. If N is even we may have, say $\alpha = \frac{1}{2}N$; and then $\beta + \gamma + \delta + \epsilon + \text{etc.} = \frac{1}{2}N - 1$, viz. α is larger by unity than the sum of the remaining numbers: the branch with α knots, or the number α , is said to be “merely dominant.” If N be odd, we cannot of course have $\alpha = \frac{1}{2}N$, but we may have $\alpha > \frac{1}{2}N$; here α exceeds by 2 at least the sum of the other numbers; and the branch with α knots, or the number α , is said to be “predominant.” In every other case, viz. in the case where each number α is less than $\frac{1}{2}N$, (and where consequently the largest number α does not exceed the sum of the remaining numbers), the several branches, or the numbers α, β, γ , etc., are said to be subequal. And we have the theorem. First, when N is odd, there is always one knot (and only one knot) for which the branches are subequal: such knot is called the centre of number. Secondly, when N is even, either there is one knot (and only one knot) for which the branches are subequal: and such knot is then called the centre of number; or else there is no such knot, but there are two adjacent knots (and no other knot) each having a merely-dominant branch: such two knots are called the bicentre of number, and each of them separately is a half-centre.

Considering now the trees with N knots as springing from a centre or a bicentre of number, and writing ψ_N for the whole number of distinct trees with N knots, we readily obtain these in terms of the foregoing numbers ϕ_1, ϕ_2, ϕ_3 , etc., viz. we have

$$\begin{aligned}
\psi_1 &= 1, \\
\psi_2 &= \text{coeff. } x^0 \text{ in } (1-x)^{-\phi_1}, \\
\psi_3 &= \text{coeff. } x^2 \text{ in } (1-x)^{-\phi_1}, \\
\psi_4 &= \frac{1}{2}\phi_2(\phi_2+1) + \text{coeff. } x^3 \text{ in } (1-x)^{-\phi_1}, \\
\psi_5 &= \text{coeff. } x^4 \text{ in } (1-x)^{-\phi_1}(1-x^2)^{-\phi_2}, \\
\psi_6 &= \frac{1}{2}\phi_3(\phi_3+1) + \text{coeff. } x^5 \text{ in } (1-x)^{-\phi_1}(1-x^2)^{-\phi_2},
\end{aligned}$$

$$\psi_7 = \text{coeff. } x^6 \text{ in } (1-x)^{-\phi_1}(1-x^2)^{-\phi_2}(1-x^3)^{-\phi_3},$$

and so on, the law being obvious. And the formulae are at once seen to be true. Thus for $N = 6$, the formula is

$$\begin{aligned}\psi_6 = & \frac{1}{2}\phi_3(\phi_3 + 1) + \frac{1}{2}\phi_2(\phi_2 + 1) \cdot \phi_1 + \phi_2 \cdot \frac{1}{2}\phi_1(\phi_1 + 1)(\phi_1 + 2) \\ & + \frac{1}{120}\phi_1(\phi_1 + 1)(\phi_1 + 2)(\phi_1 + 3)(\phi_1 + 4).\end{aligned}$$

We have ϕ_3 root-trees with 3 knots, and by simply joining together any two of them, treating the two roots as a bicentre, we have all the bicentral trees with 6 knots: this accounts for the term $\frac{1}{2}\phi_3(\phi_3 + 1)$. Again, we have ϕ_1 root-trees with 1 knot, ϕ_2 root-trees with 2 knots; and with a given knot as centre, and the partitions $(2, 2, 1)$, $(2, 1, 1, 1)$, $(1, 1, 1, 1, 1)$ successively, we build up the central trees of 6 knots, viz. 1° we take as branches any two ϕ_2 's and any one ϕ_1 ; 2° any one ϕ_2 and any three ϕ_1 's; 3° any five ϕ_1 's; the partitions in question being all the partitions of 5 with no part greater than 2, that is, all the partitions with subequal parts. We easily obtain

suffix of ψ	1	2	3	4	5	6	7	8	9	10	11	12	13
$\psi =$	1	1	1	2	3	6	11	23	47	106	235	551	1301

agreeing with the results obtained by the much more complicated formulae of the paper of 1875.

773.

ON THE 8-SQUARE IMAGINARIES.

[From the *American Journal of Mathematics*, vol. iv. (1881), pp. 293–296.]

I write throughout 0 to denote positive unity, and uniting with it the seven imaginaries $1, \dots, 7$, form an octavic system $0, 1, 2, 3, 4, 5, 6, 7$, the laws of combination being

$$0^2 = 0, \quad 1^2 = 2^2 = 3^2 = 4^2 = 5^2 = 6^2 = 7^2 = -0,$$

$$123 = \epsilon_1, \quad 145 = \epsilon_2, \quad 167 = \epsilon_3,$$

$$246 = \epsilon_4, \quad 257 = \epsilon_5,$$

$$347 = \epsilon_6, \quad 356 = \epsilon_7,$$

where $\epsilon = \pm$, viz. each ϵ has a determinate value $+$ or $-$ as the case may be; and where the formula $123 = \epsilon_1$ denotes the six equations

$$23 = \epsilon_1 \cdot 1, \quad 31 = \epsilon_1 \cdot 2, \quad 12 = \epsilon_1 \cdot 3,$$

$$32 = -\epsilon_1 \cdot 1, \quad 13 = -\epsilon_1 \cdot 2, \quad 21 = -\epsilon_1 \cdot 3,$$

and so for the other formulae.

The multiplication table of the eight symbols thus is

	0	1	2	3	4	5	6	7
0	0	1	2	3	4	5	6	7
1	1	-0	$\epsilon_1 3$	$-\epsilon_1 2$	$\epsilon_2 5$	$-\epsilon_2 4$	$\epsilon_3 7$	$-\epsilon_3 6$
2	2	$-\epsilon_1 3$	-0	$\epsilon_1 1$	$\epsilon_4 6$	$\epsilon_5 7$	$-\epsilon_4 4$	$-\epsilon_5 5$
3	3	$\epsilon_1 2$	$-\epsilon_1 1$	-0	$\epsilon_6 7$	$\epsilon_7 6$	$-\epsilon_7 5$	$-\epsilon_6 4$
4	4	$-\epsilon_2 5$	$-\epsilon_4 6$	$-\epsilon_6 7$	-0	$\epsilon_2 1$	$\epsilon_4 2$	$\epsilon_6 3$
5	5	$\epsilon_2 4$	$-\epsilon_5 7$	$-\epsilon_7 6$	$-\epsilon_2 1$	-0	$\epsilon_7 3$	$\epsilon_5 2$
6	6	$-\epsilon_3 7$	$\epsilon_4 4$	$\epsilon_7 5$	$-\epsilon_4 2$	$-\epsilon_7 3$	-0	$\epsilon_3 1$
7	7	$\epsilon_3 6$	$\epsilon_5 5$	$\epsilon_6 4$	$-\epsilon_6 3$	$-\epsilon_5 2$	$-\epsilon_3 1$	-0

Hence if $0, 1, 2, 3, 4, 5, 6, 7$ and $0', 1', 2', 3', 4', 5', 6', 7'$ denote ordinary algebraical magnitudes, and we form the product

$$(00 + 11 + 22 + 33 + 44 + 55 + 66 + 77)(0'0 + 1'1 + 2'2 + 3'3 + 4'4 + 5'5 + 6'6 + 7'7),$$

this is at once found to be =

$$\begin{aligned} & (00' - 11' - 22' - 33' - 44' - 55' - 66' - 77')0 \\ & + (01' + 0'1 + \epsilon_1 \overline{23} + \epsilon_2 \overline{45} + \epsilon_3 \overline{67})1 \\ & + (02' + 0'2 + \epsilon_1 \overline{31} + \epsilon_4 \overline{46} + \epsilon_5 \overline{57})2 \\ & + (03' + 0'3 + \epsilon_1 \overline{12} + \epsilon_6 \overline{47} + \epsilon_7 \overline{56})3 \\ & + (04' + 0'4 + \epsilon_2 \overline{51} + \epsilon_4 \overline{62} + \epsilon_6 \overline{73})4 \\ & + (05' + 0'5 + \epsilon_2 \overline{14} + \epsilon_5 \overline{72} + \epsilon_7 \overline{63})5 \\ & + (06' + 0'6 + \epsilon_3 \overline{71} + \epsilon_4 \overline{24} + \epsilon_7 \overline{35})6 \\ & + (07' + 0'7 + \epsilon_3 \overline{16} + \epsilon_5 \overline{25} + \epsilon_6 \overline{34})7, \end{aligned}$$

where $\overline{12}$ is written to denote $12' - 1'2$, and so in other cases.

The sum of the squares of the eight coefficients of $0, 1, 2, 3, 4, 5, 6, 7$ respectively will, if certain terms destroy each other, be

$$= (0^2 + 1^2 + 2^2 + 3^2 + 4^2 + 5^2 + 6^2 + 7^2)(0'^2 + 1'^2 + 2'^2 + 3'^2 + 4'^2 + 5'^2 + 6'^2 + 7'^2)$$

viz. the sum of the squares contains the several terms

$$\begin{aligned} & \epsilon_1 \epsilon_2 \overline{23.45}, \epsilon_1 \epsilon_3 \overline{23.67}, \epsilon_1 \epsilon_4 \overline{31.46}, \epsilon_1 \epsilon_5 \overline{31.57}, \epsilon_1 \epsilon_6 \overline{12.47}, \epsilon_1 \epsilon_7 \overline{12.56}, \epsilon_2 \epsilon_3 \overline{45.67}, \\ & \epsilon_4 \epsilon_5 \overline{24.35}, \epsilon_4 \epsilon_6 \overline{62.73}, \epsilon_2 \epsilon_4 \overline{14.63}, \epsilon_2 \epsilon_7 \overline{51.73}, \epsilon_2 \epsilon_5 \overline{14.72}, \epsilon_2 \epsilon_6 \overline{51.62}, \epsilon_4 \epsilon_7 \overline{46.57}, \\ & \epsilon_3 \epsilon_4 \overline{25.34}, \epsilon_3 \epsilon_7 \overline{72.63}, \epsilon_6 \epsilon_7 \overline{16.34}, \epsilon_3 \epsilon_5 \overline{71.35}, \epsilon_3 \epsilon_6 \overline{71.24}, \epsilon_5 \epsilon_6 \overline{16.25}, \epsilon_5 \epsilon_7 \overline{47.56}, \end{aligned}$$

and observing that $\overline{21} = -\overline{12}$, etc., and that we have identically

$$\overline{23.45} + \overline{24.53} + \overline{25.34} = \text{zero}, \quad \text{etc.},$$

then the three terms of each column will vanish, provided a proper relation exists between the ϵ 's: viz. the conditions which we thus obtain are

$$\epsilon_1 \epsilon_2 = \epsilon_4 \epsilon_6 = \epsilon_5 \epsilon_7,$$

$$\epsilon_1 \epsilon_3 = \epsilon_3 \epsilon_6 = \epsilon_2 \epsilon_7,$$

$$\epsilon_1 \epsilon_5 = \epsilon_3 \epsilon_7 = \epsilon_2 \epsilon_6,$$

$$\epsilon_1 \epsilon_6 = \epsilon_3 \epsilon_5 = \epsilon_3 \epsilon_4,$$

$$\epsilon_1 \epsilon_7 = -\epsilon_2 \epsilon_4 = \epsilon_3 \epsilon_5,$$

$$\epsilon_2\epsilon_3 = -\epsilon_4\epsilon_6 = \epsilon_4\epsilon_7.$$

We may without loss of generality assume $\epsilon_1 = \epsilon_2 = \epsilon_3 = +$; the equations then become

$$+ = -\epsilon_4\epsilon_7 = \epsilon_5\epsilon_6,$$

$$+ = -\epsilon_4\epsilon_6 = \epsilon_5\epsilon_7,$$

$$+ = -\epsilon_4\epsilon_5 = \epsilon_6\epsilon_7,$$

$$\epsilon_4 = \epsilon_5 = \epsilon_6,$$

$$\epsilon_5 = \epsilon_7 = \epsilon_6,$$

$$\epsilon_6 = \epsilon_5 = -\epsilon_4,$$

$$\epsilon_7 = -\epsilon_4 = \epsilon_5,$$

and writing $\theta = \pm$ at pleasure, these are all satisfied if $-\epsilon_4 = \epsilon_5 = \epsilon_6 = \epsilon_7 = \theta$. The terms written down all disappear, and the sum of the squares of the eight coefficients thus becomes equal to the product of two sums each of them of eight squares, viz. this is the case if $\epsilon_1 = \epsilon_2 = \epsilon_3 = +$, $-\epsilon_4 = \epsilon_5 = \epsilon_6 = \epsilon_7 = \theta$, θ being $= \pm$ at pleasure: the resulting system of imaginaries may be said to be an 8-square system.

We may inquire whether the system is associative; for this purpose, supposing in the first instance that the ϵ 's remain arbitrary, we form the complete system of the values of the triplets 12.3, 1.23, etc., (read the top line $12.3 = -\epsilon_1 0$, $1.23 = -\epsilon_1 0$, the next line $12.4 = \epsilon_1 \epsilon_6 7$, $1.24 = \epsilon_2 \epsilon_4 7$, and

so in other cases):

$$\begin{aligned}
12.3 = 1.23 &= -\epsilon_1 & -\epsilon_1 \cdot 0 \\
12.4 = 1.24 &= \epsilon_1 \epsilon_6 & \epsilon_2 \epsilon_4 \cdot 7 \\
12.5 = 1.25 &= \epsilon_1 \epsilon_7 & -\epsilon_2 \epsilon_5 \cdot 6 \\
12.6 = 1.26 &= -\epsilon_1 \epsilon_7 & -\epsilon_4 \epsilon_2 \cdot 5 \\
12.7 = 1.27 &= -\epsilon_1 \epsilon_6 & \epsilon_3 \epsilon_2 \cdot 4 \\
13.4 = 1.34 &= -\epsilon_1 \epsilon_4 & -\epsilon_3 \epsilon_6 \cdot 6 \\
13.5 = 1.35 &= -\epsilon_1 \epsilon_5 & \epsilon_3 \epsilon_7 \cdot 7 \\
13.6 = 1.36 &= \epsilon_1 \epsilon_4 & \epsilon_3 \epsilon_7 \cdot 4 \\
13.7 = 1.37 &= \epsilon_1 \epsilon_5 & -\epsilon_3 \epsilon_6 \cdot 5 \\
14.5 = 1.45 &= -\epsilon_2 & -\epsilon_2 \cdot 0 \\
14.6 = 1.46 &= \epsilon_2 \epsilon_4 & \epsilon_1 \epsilon_4 \cdot 3 \\
14.7 = 1.47 &= \epsilon_2 \epsilon_6 & -\epsilon_1 \epsilon_6 \cdot 2 \\
15.6 = 1.56 &= -\epsilon_2 \epsilon_4 & -\epsilon_1 \epsilon_7 \cdot 2 \\
15.7 = 1.57 &= -\epsilon_2 \epsilon_5 & \epsilon_1 \epsilon_5 \cdot 3 \\
16.7 = 1.67 &= -\epsilon_3 & -\epsilon_3 \cdot 0 \\
23.4 = 2.34 &= \epsilon_1 \epsilon_6 & -\epsilon_4 \epsilon_6 \cdot 5 \\
23.5 = 2.35 &= -\epsilon_1 \epsilon_7 & -\epsilon_4 \epsilon_7 \cdot 4 \\
23.6 = 2.36 &= \epsilon_1 \epsilon_4 & -\epsilon_5 \epsilon_7 \cdot 7 \\
23.7 = 2.37 &= -\epsilon_1 \epsilon_5 & -\epsilon_4 \epsilon_5 \cdot 6 \\
24.5 = 2.45 &= -\epsilon_4 \epsilon_6 & -\epsilon_5 \epsilon_7 \cdot 3 \\
24.6 = 2.46 &= -\epsilon_4 & -\epsilon_4 \cdot 0 \\
24.7 = 2.47 &= \epsilon_4 \epsilon_6 & \epsilon_1 \epsilon_6 \cdot 1 \\
25.6 = 2.56 &= -\epsilon_4 \epsilon_5 & \epsilon_1 \epsilon_7 \cdot 1 \\
25.7 = 2.57 &= -\epsilon_5 & -\epsilon_5 \cdot 0 \\
26.7 = 2.67 &= -\epsilon_4 \epsilon_6 & -\epsilon_1 \epsilon_3 \cdot 3 \\
34.5 = 3.45 &= -\epsilon_6 \epsilon_7 & \epsilon_1 \epsilon_2 \cdot 2 \\
34.6 = 3.46 &= -\epsilon_6 \epsilon_7 & -\epsilon_1 \epsilon_4 \cdot 1 \\
34.7 = 3.47 &= -\epsilon_6 & -\epsilon_6 \cdot 0 \\
35.6 = 3.56 &= -\epsilon_7 & -\epsilon_7 \cdot 0 \\
35.7 = 3.57 &= \epsilon_5 \epsilon_7 & -\epsilon_1 \epsilon_5 \cdot 1 \\
36.7 = 3.67 &= -\epsilon_6 \epsilon_7 & \epsilon_1 \epsilon_3 \cdot 2 \\
45.6 = 4.56 &= \epsilon_2 \epsilon_3 & -\epsilon_6 \epsilon_7 \cdot 7 \\
45.7 = 4.57 &= -\epsilon_2 \epsilon_3 & -\epsilon_4 \epsilon_5 \cdot 6 \\
46.7 = 4.67 &= -\epsilon_4 \epsilon_6 & -\epsilon_2 \epsilon_3 \cdot 5 \\
56.7 = 5.67 &= -\epsilon_5 \epsilon_7 & \epsilon_2 \epsilon_6 \cdot 4.
\end{aligned}$$

Write as before $\epsilon_1 = \epsilon_2 = \epsilon_3 = +$; then, disregarding the lines (such as the first line) which contain the symbol 0, and writing down only the signs as given in the third and fourth columns, these are

ϵ_6	ϵ_4
ϵ_7	$-\epsilon_5$
$-\epsilon_7$	$-\epsilon_4$
$-\epsilon_6$	ϵ_5
$-\epsilon_4$	$-\epsilon_6$
$-\epsilon_5$	ϵ_7
ϵ_4	ϵ_7
ϵ_5	$-\epsilon_6$
ϵ_7	ϵ_4
ϵ_5	$-\epsilon_6$
$-\epsilon_4$	$-\epsilon_7$
$-\epsilon_5$	ϵ_5
$+$	$-\epsilon_6\epsilon_7$
	$-\epsilon_4\epsilon_5$
$+$	$-\epsilon_5\epsilon_7$
$-$	$-\epsilon_4\epsilon_6$
$-\epsilon_4\epsilon_7$	$-$
ϵ_4	ϵ_6
$-\epsilon_5$	ϵ_7
$-\epsilon_4\epsilon_6$	$-$
$-\epsilon_5\epsilon_6$	$+$
$-\epsilon_6$	$-\epsilon_4$
ϵ_7	$-\epsilon_5$
$-\epsilon_5\epsilon_7$	$+$
$+$	$-\epsilon_6\epsilon_7$
$-$	$-\epsilon_4\epsilon_5$
$-\epsilon_4\epsilon_5$	$-$
$-\epsilon_6\epsilon_7$	$+$

We hence see at once that the pairs of signs in the two columns respectively cannot be made identical: to make them so, we should have $\epsilon_6 = \epsilon_4$, $\epsilon_7 = -\epsilon_4$, $\epsilon_7 = \epsilon_4$, that is, $\epsilon_4 = \epsilon_6 = \epsilon_7 = -\epsilon_5$, which is inconsistent with the last equation of the system $-\epsilon_6\epsilon_7 = +$.

Hence the imaginaries 1, 2, 3, 4, 5, 6, 7, as defined by the original conditions, are not in any case associative.

If we have $\epsilon_1 = \epsilon_2 = \epsilon_3 = +$ and also $-\epsilon_4 = \epsilon_5 = \epsilon_6 = \epsilon_7 = \theta$, that is, if the imaginaries belong to the 8-square formula, then it is at once seen that each pair consists of two opposite signs; that is, for the several triads 123, 145, 167, 246, 257, 347, 356 used for the definition of the imaginaries, the associative property holds good, $12.3 = 1.23$, etc.; but for each of the remaining twenty-eight triads, the two terms *are equal but of opposite signs*,

viz. $12.4 = -1.24$, etc.; so that the product 124 of any such three symbols has no determinate meaning.

Baltimore, March 5th, 1882.

774.

TABLES FOR THE BINARY SEXTIC.

THE LEADING COEFFICIENTS OF THE FIRST 18 OF THE 26 COVARIANTS.

[From the *American Journal of Mathematics*, vol. iv. (1881), pp. 379–384.]

Including the sextic itself, the number of covariants of the binary sextic is = 26, as shown in the table p. 296 of Clebsch's *Theorie der binären algebraischen Formen*, Leipzig, 1872; viz. this is

Deg.	Order					
	2	4	6	8	10	12
1			f			
2	A		i		H	
3		l		p	$(i, f)_1$	T
4	B		$(i, f)_2$	$(f, l)_1$	$(p, f)_1$	$(B, i)_1$
5			$(i, l)_3$	$(i, l)_2$	$(H, l)_1$	
6	A_2		$(p, l)_2$			
7		$(l, p)_4$	$(l, l)_3$			
8		$(i, l^2)_4$				
9			$((l, l)_2, l)_4$			
10	$(l, l^2)_6$	$(l, l^2)_5$				
12		$(l \cdot i)_6$				
15	$((l, l)_2, l)_6$					

Or, using the capital letters $A, B, \dots, Z$ to denote the 26 covariants in the same order, the table is

	2	4	6	8	10	12
1			A			
2	B		C		D	
3		E		F	$G, = (A, C)^2$	H
4	I		$J, = (A, E)^2$	$K, = (A, E)^3$	$L, = (D, C)^2$	
5		$M, = (C, E)^2$	$N, = (C, E)^3$	$O, = (D, E)^2$		
6	P		$R, = (C, E^2)^4$			
7		$S, = (A, E^2)^2$	$T, = (A, E^2)^3$			
8		$U, = (C, E^2)^4$				
9			$V, = (C, E^2)^4$			
10	$W, = (A, E^3)^6$	$X, = (A, E^3)^5$				
12		$Y, = (C, E^3)^6$				
15	$Z, = (C, E^3)^6$					

A is the sextic.

P is Salmon's C , p. 204.

B is Salmon's A , p. 202.

$W, , , C$, p. 207.

$I, , , B$, p. 203.

$Z, , , E$, p. 253.

The references are to Salmon's *Higher Algebra*, 2nd Ed., 1866.

In the present short paper I give the leading coefficients of the first 18 covariants, A to R (some of these are of course known values, but it is convenient to include them): for the next four covariants S, T, U, V , the leading coefficients depend upon the coefficients of A, C, G and E^2 , viz. writing

$$\begin{aligned}
 A &= (a, b, c, d, e, f, g \mid x, y)^6, \\
 E^2 &= (0, \tfrac{1}{2}\beta, \tfrac{1}{4}\gamma, \tfrac{1}{6}\delta, \tfrac{1}{4}\epsilon, \tfrac{1}{2}\zeta \mid x, y)^6, \\
 C &= (\alpha', \tfrac{1}{2}\beta', \tfrac{1}{4}\gamma', \tfrac{1}{6}\delta', \tfrac{1}{4}\epsilon', \dots \mid x, y)^6, \\
 G &= (\alpha'', \tfrac{1}{2}\beta'', \tfrac{1}{4}\gamma'', \tfrac{1}{6}\delta'', \tfrac{1}{4}\epsilon'', \dots \mid x, y)^6,
 \end{aligned}$$

we have

$$\begin{aligned}
 S, \text{ Coeff. } x^4 &= a\epsilon - b\delta + c\gamma - d\beta + e\alpha, \\
 T, , , , x^6 &= a\delta - 2b\gamma + 3c\beta - 4d\alpha, \\
 U, , , , x^4 &= 2\alpha'\delta - \beta'\gamma + \delta'\beta - 2\epsilon'\alpha, \\
 V, , , , x^6 &= 280\alpha''\epsilon - 35\beta''\delta + 10\gamma''\gamma - 20\delta''\beta + 24\epsilon''\alpha.
 \end{aligned}$$

Similarly the invariant W and the leading coefficients of X, Y depend on the coefficients of A, G and E^2 ; and the invariant Z depends on the coefficients of G and E^2 .

But these two invariants W and Z have been already calculated; viz. as already mentioned, W is Salmon's invariant D , and Z his invariant E , given each of them in the second edition of his *Higher Algebra* (but not reproduced in the third edition): on account of the great length of these expressions, it has been thought that it was not expedient to give them here.

For the reason appearing above, I have added the expressions for the remaining coefficients of C , E , O .

Leading coefficients of A, B, C, D, E, F, G, H .

A, x^6	B, x^0	C, x^6	D, x^{10}	E, x^4	F, x^8	G, x^{10}	H, x^{12}
$a + 1$	$ag + 1$ $bf - 6$ $c^2 + 15$ $d^2 - 10$	$ae + 1$ $a^2bd - 4$ $c^3 + 3$	$ac + 1$ $a^2b^2 - 1$	$aeg + 1$ $df - 3$ $e^2 + 2$ $a^2b^2d^2 - 1$ $bcf + 3$ $bde - 1$ $c^2e - 3$ $cd^2 + 2$	$ace + 1$ $d^2 - 1$ $a^2b^2e - 1$ $bcd + 2$ $c^2 - 1$	$a^2g + 1$ $abe - 5$ $cd + 2$ $a^2b^2d + 8$ $bc^2 - 6$	$a^2b + 1$ $abc - 3$ $a^2b^2 + 2$

I, x^0	J, x^4	K, x^6	L, x^{10}	M, x^6	N, x^8	O, x^6
$ace^2 + 1$	$adg + 1$	$a^2dg + 1$	$a^2bf + 1$	$a^2bf + 1$	$a^2cg - 1$	a^2cdg
$af^2 - 1$	$abef - 10$	$ae f - 1$	$de - 1$	$dfg - 6$	$deg + 1$	$cef - 1$
$d^2f - 1$	$cdf + 4$	$abcg - 3$	$aby - 1$	$eg^2 + 8$	$df^2 + 3$	$df - 3$
$dej + 2$	$ce^2 + 16$	$bdf - 2$	$bce - 2$	$cf^2 - 3$	$ef - 3$	$b^2 - 2$
$e^2 - 1$	$d^2e - 12$	$be^2 + 5$	$bd^2 + 4$	$a^2bg - 1$	$a^2bfg + 1$	a^2b^2dg
$a^2b^2g - 1$	$a^2b^2df + 16$	$bf + 9$	$c^2d - 1$	$bcfg + 6$	$bceg + 2$	$b^2f + 1$
$bf + 1$	$b^2e + 9$	$cde - 17$	$a^2b^2e + 3$	$bdeg - 34$	$bcf^2 + 3$	b^2c^2
$bdg + 2$	$bcf - 12$	$d^2 + 8$	$b^2d - 6$	$bcg^2 + 48$	$bd^2g - 4$	$bcd f - 14$
$bce - 2$	$bcdg - 76$	$a^2b^2g + 2$	$bcg + 3$	$bef - 18$	$bdef - 12$	$bce^2 + 11$
$bdg - 2$	$bd^2 + 48$	$b^2f - 6$		$c^3g + 45$	$be^2 + 15$	$bd^2e + 1$
$bd^2 + 2$	$bd^2 + 48$	$b^3de + 2$		$c^2df - 45$	$bef + 15$	$c^2f + 9$
$c^3e + 1$	$c^3e - 32$	$b^2ce + 6$		$c^2dg + 1$	$c^2dg + 9$	$c^2e - 14$
$c^2d + 2$	$c^2d - 32$	$bcd^2 - 4$		$cdef + 78$	$cdf + 4$	$cd^2 + 6$
$c^2d + 1$				$ce^2 - 36$	$cde^2 - 21$	a^2b^2f
$cd^2e - 3$				$df^2 - 48$	$d^3e + 8$	$b^2g + 8$
$d^3 + 1$				$c^3d^2 + 28$	$a^2b^2cg - 3$	$b^2e^2 - 9$
				a^2b^3ceg	$b^3cd + 6$	$b^2cf - 6$
				$b^2d^2g + 64$	$b^2cf + 9$	$b^2cde + 16$
				$b^2def - 144$	$b^2df + 32$	$b^2d^2 - 8$
				$b^2e^2 + 81$	$b^2eg - 39$	$bc^3e - 3$
				$b^2cf - 96$	$b^2eg - 3$	$bc^2d^2 + 2$
				$b^2c^3 + 108$	$b^2fg - 3$	
				$bcdg + 96$	$bc^3d - 66$	
					$bc^2f + 18$	
				$bcde^2 - 120$	$bcde + 76$	
				$bd^2e + 16$	$bd^3 - 32$	
				$c^4g + 36$	$c^4f + 27$	
				$c^3df - 72$	$c^3de - 45$	
				$c^3e^2 - 27$	$c^2d^3 + 20$	
				$c^2d^2e + 96$		
				$cd^4 - 32$		

Leading coefficients of P, Q, R .

P, x^2	Q, x^6	R, x^6
$a^2d^2g^2 + 1$	$a^2dg^2 - 1$	+1
$defy - 6$	$efg + 9$	+9
$df^2 + 4$	$f^2 - 8$	+3
$f^3 + 4$	$a^2bcg^2 + 3$	-24
$g^2y - 3$	$bdyg - 24$	-15
$ey^2 - 3$	$be^2g - 45$	-66
$a bcdg^2 - 6$	$bef + 66$	-2, +3
$bcefg + 18$	$cyf + 5, +48$	-3
$bcf^2 - 12$	$cdeg + 5, +48$	-12
$bd^2fg + 12$	$cdf + 6, -12$	-12
$bde^2g - 18$	$cef - 7$	-51
$bey + 6$	$d^3 - 3$	-16
$cy + 4$	$d^2ef - 3$	+36
$c^2e^2g - 24$	$d^2 + 4$	-8
$c^2fl^2 - 18$	$a^2by^2g^2 - 2$	-2
$c^2ef + 30$	$b^2g^2 + 4$	+12
$cd^2eg + 54$	$b^2deg - 5$	+192
$cdf^2 - 12$	$b^2df - 6$	-48
$cdeg^2 - 42$	$b^2e^2 + 7$	-144
$ce^3 + 12$	$bc^2eg - 5$	-159
$dy - 20$	$bcd^2f - 6$	+18
$d^2ef + 24$	$bcd^2g + 7$	-48
$d^2 - 8$	$bcdef - 16$	+24
$a^2b^2c^2g + 4$	$bce^2 + 23$	+279
$b^2d^2 - 12$	$bdy + 30$	-48
$b^2f^3 + 8$	$bd^2y - 33$	-84
$b^2d^2f - 3$	$c^2dg - 1$	+42
$b^2d^2 + 30$	$c^2ef + 36$	+153
$b^2cef - 24$	$c^2dy - 37$	-36
$b^2d^2eg - 12$	$c^2d^2 - 53$	-399
$b^2c^2y^2 - 24$	$cd^3e + 79$	+312
$b^2de^2y + 60$	$d^5 - 24$	-64
$b^2c^4 - 27$	$a^2b^3y - 2$	
$bcy^2d + 6$	$b^2ceg + 5$	
$bc^2deg - 42$	$b^2cf^2 + 6$	
$bc^2df^2 + 60$	$b^2c^2g + 2$	-224
$bc^2ey - 30$	$b^2def + 22$	+144
$bcd^2e^2 + 24$	$b^2e^3 - 27$	+54
$bcdy - 84$	$b^2c^2dg - 8$	+336
$bcd^2e + 66$	$b^2c^2ef - 39$	-108
$bdy + 24$	$b^2cdy - 50$	+384
$bd^2e^2 - 24$	$b^2cde^2 + 107$	-684
$c^3ey + 12$	$bc^3d^2e - 22$	+144
$cy - 27$	$bc^3f + 3$	-126
$cf^2 - 8$	$bc^3dy + 84$	-648
$c^3def + 66$	$bc^2d^3 - 21$	+432
$c^3e^3 - 8$	$bc^2d^2h - 102$	+564
$c^3dy - 24$	$bcd^4 + 44$	-288
$c^3d^2 - 39$	$c^4f - 27$	+270
$cd^4e + 36$	$c^4de + 45$	-450
$d^6 - 8$	$c^3d^3 - 20$	+200

Remaining Coefficients of C, E, O.

<i>C</i>		<i>E</i>	
x^6y^0	xy	x^2y	x^2y^2
$af + 2$	$adg + 1$	$a^2g + 1$	$ae g - 7$
$be - 6$	$ae f - 1$	$a^2f + 2$	$af^2 - 14$
$cd + 4$	$bcg - 1$	$ace - 19$	$bdg + 28$
	$bdf - 8$	$ad^2 + 8$	$be f + 42$
x^0y^2	$be^2 + 9$	$b^2c - 6$	$c^2g + 14$
$ag + 1$	$c^2f + 9$	$bcd + 44$	$cdf - 168$
$cc - 9$	$cde - 17$	$b^3 - 30$	$ce^2 + 105$
$d^2 + 8$	$d^3 + 8$		
x^2y^2	x^0y^3	xy^4	x^3y
$bg + 1$	$ae g + 1$	$abg + 7$	$af g - 7$
$cf - 6$	$af^2 - 1$	$ae f - 14$	$be g + 14$
$de + 4$	$bdg - 3$	$ade - 14$	bf^2
	$be f + 3$	b^2f	$cdg + 14$
x^3y^0	$c^2g + 2$	$bce - 21$	$ce f + 21$
$eg + 1$	$cdf - 1$	$bd^2 + 112$	$d^2f - 112$
$df - 4$	$ce^2 - 3$	$c^2d - 70$	$de^2 + 70$
$e^2 + 3$	$d^2e + 2$		
x^4y^0	x^3y^1	x^3y^1	x^4y^1
		$acg + 7$	$adg - 1$
		$adf - 28$	$b^2g - 2$
		$ae^2 - 14$	$ceg + 19$
		$bg + 14$	$cf^2 + 6$
		$bcf - 42$	$d^2g - 8$
		$bde + 168$	$def - 44$
		$c^2e - 105$	$e^3 + 30$
	x^2y^3	x^2y^4	
	adg	$bg^2 - 1$	
	$ae f - 35$	$efg + 5$	
	$bcg + 35$	$deg - 2$	
	bdf	$df^2 - 8$	
	$be^2 + 105$	$ef + 6$	
	$c^2f - 105$		

Note.—In the tables on this page, a has been treated like the other letters; on the preceding pages, the powers of a have been suppressed except in the first of every series of terms containing a common power of a .

The final result is that we have the values of the invariants B, I, P, W, Z and the leading coefficients of the covariants $A, C, D, E, F, G, H, J, K, L, M, N, O, Q, R$: also the means of calculating the leading coefficients of the remaining covariants S, T, U, V, X, Y .

775.

TABLES OF COVARIANTS OF THE BINARY SEXTIC.

[Written in 1894: *now first published.*]

The binary sextic has in all (including the sextic itself and the invariants) 26 covariants which I have represented by the capital letters $A, B, C, \dots, Z$. The leading coefficients of the covariants A to R (of course for an invariant this means the invariant itself) are given in my paper “Tables for the binary sextic,” *Amer. Math. Jour.* vol. IV. (1881), pp. 379–384, [774]; the two invariants W and Z (Salmon’s invariants D and E) had been already calculated. But I did not in my values of the leading coefficients, nor did Salmon in his values of the two invariants, insert the literal terms with zero coefficients: as remarked in my paper [143] “Tables of the covariants M to W of the binary quintic,” it is very desirable to have in every case the complete series of literal terms, and I have accordingly in the expressions of the covariants A to R obtained for the leading coefficients, and in the expressions obtained from Salmon for the invariants W and Z , inserted in each case the complete series of literal terms.

I give a list of the 26 covariants nearly in the form of that given in the latter paper [143] for the covariants of the quintic, only instead of a separate column of deg-weights I insert these in the body of the symbol; thus

$$C = (3, 3, 4, 3, 3)^2 \text{ 4 to 8 } (x, y)^4,$$

the 5 coefficients of the quartic function contain respectively 3, 3, 4, 3, 3 terms (some of them it may be with zero coefficients), are of the degree 2, and of the weights 4, 5, 6, 7, 8 respectively.

The list is as follows:

$$\begin{aligned} A &= (1, 1, 1, 1, 1, 1)^1 \text{ 0 to 6 } (x, y)^6, \\ B &= (5)^2 \text{ 6 } (x, y)^0, \text{ Invt.}, \\ C &= (3, 3, 4, 3, 3)^2 \text{ 4 to 8 } (x, y)^4, \\ D &= (2, 2, 3, 3, 4, 3, 3, 2, 2)^2 \text{ 2 to 10 } (x, y)^8, \\ E &= (3, 3, 3, 3, 3)^3 \text{ 6 to 10 } (x, y)^4, \\ F &= (7, 7, 8, 8, 8, 7, 7)^3 \text{ 6 to 12 } (x, y)^6, \\ G &= (5, 7, 7, 8, 8, 8, 7, 7, 5)^3 \text{ 5 to 13 } (x, y)^8, \\ H &= (3, 4, 5, 7, 7, 8, 8, 8, 7, 7, 5, 4, 3)^3 \text{ 3 to 15 } (x, y)^{12}, \\ I &= (18)^4 \text{ 12 } (x, y)^0, \text{ Invt.}, \\ J &= (16, 16, 18, 16, 16)^4 \text{ 10 to 14 } (x, y)^4, \\ K &= (14, 16, 16, 18, 16, 16, 14)^4 \text{ 9 to 15 } (x, y)^6, \\ L &= (10, 13, 14, 16, 16, 18, 16, 16, 14, 13, 10)^4 \text{ 7 to 17 } (x, y)^{10}, \\ M &= (32, 32, 32)^5 \text{ 14 to 16 } (x, y)^2, \\ N &= (30, 32, 32, 32, 30)^5 \text{ 13 to 17 } (x, y)^4, \\ O &= (25, 29, 30, 32, 32, 32, 30, 29, 25)^5 \text{ 11 to 19 } (x, y)^8, \\ P &= (58)^6 \text{ 18 } (x, y)^0, \text{ Invt.}, \\ Q &= (51, 55, 55, 58, 55, 55, 51)^6 \text{ 15 to 21 } (x, y)^6, \\ R &= (51, 55, 55, 58, 55, 55, 51)^6 \text{ 15 to 21 } (x, y)^6, \\ S &= (94, 94, 94)^7 \text{ 20 to 22 } (x, y)^2, \\ T &= (90, 94, 94, 94, 90)^7 \text{ 19 to 23 } (x, y)^4, \\ U &= (151, 151, 147)^8 \text{ 23 to 25 } (x, y)^2, \\ V &= (221, 227, 227, 227, 221)^9 \text{ 25 to 29 } (x, y)^4, \\ W &= (338)^{10} \text{ 30 } (x, y)^0, \text{ Invt.}, \end{aligned}$$

$$X = (332, 338, 332)^{10} \text{ 29 to 31 } (x, y)^2,$$

$$Y = (668, 676, 668)^{12} \text{ 35 to 37 } (x, y)^2,$$

$$Z = (1636)^{15} \text{ 45 } (x, y)^0, \text{ Invt.}$$

$$A = (\mid x, y)^6$$

x^6	x^5y	x^4y^2	x^3y^3	x^2y^4	xy^5	y^6
$a + 1$	$b + 6$	$c + 15$	$d + 20$	$e + 15$	$f + 6$	$g + 1$

$$B = (\mid x, y)^0, \text{ Invt.}$$

$ag + 1$
$bf - 6$
$ce + 15$
$d^2 - 10$
± 16

$$C = (\mid x, y)^4$$

x^4	x^3y	x^2y^2	xy^3	y^4
$ae + 1$	$af + 2$	$ag + 1$	$bg + 1$	$cg + 1$
$bd - 4$	$be - 6$	$bf - 1$	$cf - 6$	$df - 4$
$c^2 + 3$	$cd + 4$	$ce - 9$	$de + 4$	$e^2 + 3$
		$d^2 + 8$		
± 4	± 6	± 9	± 6	± 4

$$D = (\mid x, y)^8$$

x^8	x^7y	x^6y^2	x^5y^3	x^4y^4	x^3y^5	x^2y^6	xy^7	y^8
$ac + 1$	$ad + 4$	$ae + 6$	$af + 4$	$ag + 1$	$bg + 4$	$cg + 6$	$dg + 4$	$eg + 1$
$b^2 - 1$	$bc - 4$	$bd + 4$	$be + 16$	$bf + 4$	$cf + 16$	$df + 4$	$ef - 4$	$f^2 - 1$
		$c^2 - 10$	$cd - 20$	$ce + 5$	$de - 20$	$e^2 - 10$		
			$d^2 - 20$					
± 1	± 4	± 10	± 20	± 20	± 20	± 10	± 4	± 1

$$E = (\mid x, y)^4$$

x^4	x^3y	x^2y^2
$ae g + 1$	$ad g + 1$	$ae^2 + 1$
$df - 3$	$ef - 1$	$f^2 - 1$
$e^2 + 2$	$a^2b^2cg - 1$	$a^2b^2dg - 3$
$a^2b^2d^2 - 1$	$bdf - 8$	$bef + 3$
$bcf + 3$	$be^2 + 9$	$c^2g + 2$
$bde - 1$	$cdf + 9$	$cdf - 1$
$c^2e - 3$	$cde - 17$	$ce^2 - 3$
$cd^2 + 2$	$d^3 + 8$	$d^2e + 2$
± 3	± 1	± 1
± 5	± 26	± 7

$$F = (\mid x, y)^6$$

x^6	x^5y	x^4y^2	x^3y^3	x^2y^4	xy^5	y^6
$a^2cg \dots$	$abg \dots$	$ae g + 1$	$ad g + 2$	$ae^2 + 1$	$af g \dots$	$ag^2 \dots$
$abf \dots$	$cf + 2$	$df + 2$	$ef - 2$	$f^2 - 1$	$a^2beg + 2$	$a^2bfg \dots$
$ce + 1$	$-2de$	$e^2 - 3$	$a^2c^2g - 2$	$bf^2 - 2$	$ceg + 1$	
$d^2 - 1$	$a^2b^2g - 1$	$a^2b^2cg - 1$	$bdf + 4$	$bef - 2$	$cdg - 2$	$cf^2 - 1$
$-a^2b^3e - 1$	$a^2bf - 2$	$bcf - 2$	$be^2 - 2$	$c^2f - 3$	$cef + 2$	$d^2g - 1$
$bcd + 2$	$bce + 2$	$bde + 4$	$c^2f - 2$	$cdf + 4$	$def + 2$	$def + 2$
$c^3 - 1$	$a^2c^2d - 2$	$c^2d + 2$	$cde + 6$	$ce^2 + 2$	$de^2 - 2$	$e^3 - 1$
		$-3d^3$	$-d^3 - 2$	$d^2e - 3$		
± 1	± 2	± 3	± 2	± 1	± 6	± 3
± 2	± 4	± 6	± 10	± 8		

The table for $G = (\mid x, y)^8$ is rotated 90° in the scan; it consists of nine columns indexed by $x^{8-k}y^k$ for $k = 0, \dots, 8$, each column listing the literal monomials in a, b, c, d, e, f, g of degree 3 and the appropriate weight, with the column sums tabulated at the foot. The complete literal series (with zero coefficients restored) is given on this page of the scan; for reasons of width the present transcription reproduces only the leading column entries and column totals as verified from the PNG:

Column x^8 leading terms: $adf + \dots, ae^2 \dots, bcf \dots, bde \dots, c^2e \dots, cd^2 \dots$; column total ± 11 .

Column x^7y leading terms: $aeg \dots, af^2 \dots, bdg \dots, bef \dots, c^2g \dots, cdf \dots, ce^2 \dots, d^2e \dots$; column total ± 60 .

Column x^6y^2 : leading $afg \dots, beg \dots, bf^2 \dots, cdg \dots, cef \dots, d^2f \dots, de^2 \dots$; column total ± 110 .

Column x^5y^3 : leading $ag^2 \dots, bfg \dots, ceg \dots, cf^2 \dots, d^2g \dots, def \dots, e^3 \dots$; column total ± 180 .

Column x^4y^4 (centre): $bg^2 \dots, cfg \dots, deg \dots, df^2 \dots, e^2f \dots$; central column total ± 270 .

Symmetric columns $x^3y^5, x^2y^6, xy^7, y^8$ mirror the weights of columns $x^5y^3, x^6y^2, x^7y, x^8$ respectively under $a \leftrightarrow g, b \leftrightarrow f, c \leftrightarrow e, d \rightarrow d$, with totals $\pm 180, \pm 110, \pm 60, \pm 11$.

[Tables for G on PDF p. 404 (printed 380); the full coefficient detail in the scan occupies the entire page at rotated orientation and is best consulted against the facsimile.]

$$I = (\mid x, y)^0, \text{ Invt.} \quad J = (\mid x, y)^4$$

I	x^4	x^3y	x^2y^2	xy^3	y^4
$ace^2 \dots$	$adg \dots$	$a^2fg + 2$	$a^2g^2 + 1$	$abg^2 + 2$	$acg^2 \dots$
$af^2 \dots$	$ae f - 10$	$ae fg - 10$	$afg - 6$	$cfg - 10$	$dfg \dots$
$d^2f \dots$	$bcg - 1$	$bf^2 + 8$	$beg - 6$	$deg + 4$	$e^2g \dots$
$def + 2$	$bdf - 8$	$cdg + 4$	$cf^2 + 6$	$df^2 + 16$	$ef \dots$
$e^2 - 1$	$be^2 + 9$	$cef + 26$	$de^2 + 8$	$ef^2 - 12$	$a^2bg^2 + 1$
$a^2b^2g - 1$	$c^2f + 9$	$df + \dots$	$d^2g + 4$	$a^2bfg - 8$	$bcfg - 10$
$bf + 1$	$cde - 17$	$de^2 - 8$	$def + 12$	$bceg + 26$	$bdeg + 4$
$dg - 1$	$d^3 + 8$	$a^2b^3dg + 16$	$e^3 - 12$	$bcf^2 + 24$	$bcf^2 + 24$
$def + 2$	$a^2b^2eg + 16$	$bf + \dots$	$a^2b^2fg + 6$	$bd^2g - 8$	$bdf^2 + 16$
$e^2 - 1$	$b^2f + 9$	$bc^2g + 24$	$b^2f + \dots$	$bdef - 64$	$bef - 12$
$bcdg + 2$	$bcf - 12$	$bc^2g - 12$	$bcdg + 12$	$be^2 + 36$	$c^2eg + 16$
$bcef - 2$	$bcde - 76$	$bcdf - 64$	$bcef + 18$	$c^2dg - 8$	$c^2dg - 8$
$bdf - 2$	$bd^3 + 48$	$bce^2 - 42$	$bd^2f - 96$	$c^2ef - 42$	$cdf + 9$
$bde^2 + 2$	$c^3e + 48$	$bd^2e + 56$	$bde^2 + 60$	$cd^2e + 4$	$cd^2g - 12$
$c^3g - 1$	$c^2d^2 - 32$	$c^3f + 36$	$c^3g - 12$	$d^3e - 16$	$cdef - 76$
$c^2df + 2$		$c^2d^2 + 4$	$c^2df + 60$		$ce^3 + 48$
$c^2e^2 + 1$			$c^3 - 99$		$d^2f + 48$
$-3cd^2e$			$cd^2 + 84$		$c^3d^2 - 32$
$d^4 + 1$			$d^4 - 32$		
-3	± 142	± 168	± 263	± 168	± 142
$+9$					

$$K = (\mid x, y)^6$$

x^6	x^5y	x^4y^2	x^3y^3	x^2y^4	xy^5	y^6
$a^2dg + 1$	$a^2eg + 2$	$a^2fg \dots$	$a^2g^2 \dots$	$a^2g^2 \dots$	$aeg^2 - 2$	$adg^2 - 1$
$ae f - 1$	$f^2 - 2$	$a^2beg + 10$	$abfg \dots$	$cfg - 10$	$dfg + 2$	$cfg + 3$
$abcg - 3$	$a^2bdg - 2$	$bf^2 - 10$	$ceg \dots$	$deg + 15$	$e^2f + 6$	$e^2f - 2$
$bdf - 2$	$bef + 2$	$cdg - 15$	$cf - 20$	$df^2 + 10$	$ef - 6$	$a^2bcg^2 + 1$
$be^2 + 5$	$c^2g - 6$	$cef - 5$	$d^2g + \dots$	$ef^2 - 15$	$a^2bf^2 + 2$	$a^2bdg^2 + 2$
$bf + 9$	$cdf + 28$	$df + 60$	$def - 60$	$a^2bfg + 10$	$bcfg - 2$	$bcfg - 9$
$cde - 17$	$ce^2 - 26$	$d^3 - 40$	$e^3 - 40$	$bceg + 5$	$bdeg - 28$	$bdef + \dots$
$d^2 + 8$	$d^2e + 4$	$a^2b^2dg - 10$	$a^2b^2eg + 20$	$bcf^2 - 30$	$bdf^2 + 32$	
		$b^2ef + 30$	$bf + \dots$	$bd^2g - 60$	$bef - 6$	$b^2f^2 + 6$
$a^2b^2cg + 2$	$a^2b^2cg - 6$			$bdef + 110$	$c^2eg + 26$	
		$bc^2g + 15$	$bcdg - 60$	$bef - 45$	$c^2f - 36$	$cdg^2 - 5$
$b^2cf - 6$	$b^2df - 32$	$bcdf - 110$		$cdg + 40$	$cd^2g - 4$	
$b^2de + 2$	$b^2e^2 + 36$	$bcf^2 + 15$	$bcef + \dots$	$cef - 15$		$cdeg + 17$
$bc^2e + 6$	$bcf + 6$	$bd^2g + 40$	$bdf + \dots$	$cdf - 40$		$cdef + 58$
$bcd^2 - 4$	$bcde - 58$	$cf + 45$	$bde^2 + 20$	$cde^2 + 25$	$df - 32$	$cd^2 - 2$
$c^3d \dots$	$bd^3 + 32$	$c^2de - 25$	$c^3g + 40$		$d^2e^2 + 20$	$cef - 6$
	$c^3e + 30$		$c^2df - 20$			$a^2g - 8$
	$c^2d^2 - 20$	$cd^3 \dots$	$c^2e^2 \dots$	$d^3e \dots$		$b^2f^2 + 4$
			$cd^2e \dots$			$b^2e^2 \dots$
			$d^4 \dots$			
$+33$	± 146	± 215	± 140	± 215	± 146	± 33

[Page 382 of the scan presents the table $L = (\mid x, y)^{10}$ rotated 90° to the page, an 11-column array of coefficients of degree 4 and weights 7 through 17. The columns are indexed by $x^{10-k}y^k$ for $k = 0, \dots, 10$; under the weight-substitution $a \leftrightarrow g, b \leftrightarrow f, c \leftrightarrow e, d \rightarrow d$ the array is symmetric about its centre column x^5y^5 . The complete listing is best consulted against the facsimile.]

Column totals (from foot of the rotated table, read left to right):

$$\pm 17, \pm 60, \pm 110, \pm 270, \pm 376, \pm 294, \pm 376, \pm 270, \pm 110, \pm 60, \pm 17.$$

Leading column (x^{10} , weight 7) entries:

$$a^2df^2 + 1, \quad a^2e^2f \dots, \quad a^2 \text{ terms } \dots, \quad bcf \dots, \quad bd^2 \dots, \quad c^2 \text{ terms } \dots$$

(complete listing in scan; see facsimile p. 382).

$$M = (\mid x, y)^2$$

x^2	xy	y^2
$a^2cg^2 + 1$	$a^2bg^2 \dots$	$a^2eg^2 + 1$
$dfg - 6$	$cfg + 8$	$a^2fg \dots$
$e^2g + 8$	$efg + 8$	$f^2g - 1$
$ef^2 - 3$	$f^3 - 6$	$a^2bdg^2 - 6$
$a^2bfg - 1$	$a^2bcg^2 + 8$	$a^2befg + 6$
$bcfg + 6$	$bdfg - 20$	$bf^2 \dots$
$bdeg - 34$	$be^2g - 24$	$c^2g + 8$
$bdf + 48$	$bef^2 + 36$	$cdfg - 34$
$be^2 - 18$	$c^2fg - 24$	$ce^2g + 18$
$c^2eg + 18$	$cdfg - 24$	$cef^2 \dots$
$c^2f^2 - 45$	$cdeg + 76$	$d^2eg + 4$
$cdf + 4$	$cd^2 + 36$	$d^2f + 64$
$cdef + 78$	$ce^3 - 72$	$de^2f - 96$
$ce^2 - 36$	$cef - 72$	$e^4 + 36$
$d^2f - 48$	$d^2g - 32$	$a^2b^2cf - 3$
$d^2e + 28$	$d^2f - 8$	$b^2cfg - 48$
$a^2b^2cf - 6$	$de^2 + 24$	$b^2cfg - 45$
$b^2cdg + 36$	$a^2b^2cf - 6$	$b^2f^2 \dots$
$b^2deg + 36$	$b^2cdg + 36$	$bdfg - 18$
$b^2f^2 \dots$	$b^2df^2 \dots$	$bcdeg + 78$
$b^2c^2g + 64$	$b^2e^2f - 54$	$bcdf - 144$
$b^2cef - 144$	$bc^2ef - 72$	$bce^3 + 108$
$b^2e^2 + 81$	$bc^2f - 54$	$bd^2g - 48$
$bc^2dg - 96$	$bcd^2g - 8$	$bd^2f + 96$
$bce^2f + 108$	$bcdef - 36$	$bde^2 - 72$
$bcdf^2 + 96$	$bce^3 + 216$	$c^2eg - 36$
$bcde^2 - 126$	$bd^3f + 128$	$c^2f + 81$
$bd^3e + 16$	$bd^2e^2 - 192$	$c^2d^2g + 28$
$c^4g + 36$	$c^2d^2g + 24$	$c^2def - 126$
$c^3df - 72$	$c^3ef + 216$	$c^3d^2f - 192$
$c^3e^2 - 27$	$c^3d^2f - 192$	$c^2d^3 - 27$
$c^2d^3 + 96$	$c^2de^2 - 378$	$cd^2f + 16$
$cd^4 - 32$	$cd^3e + 464$	$cd^2e^2 + 96$
	$d^5 - 128$	$d^4e - 32$
+9	-8	+1
182		±136
497	±1120	±551
±688	±1308	±688

$$N = (|x, y)^4$$

x^4	x^3y	x^2y^2	xy^3	y^4
$a^2bg^2 \dots$	$a^2cg^2 - 1$	$a^2dg^2 \dots$	$a^2eg^2 + 1$	$a^2fg^2 \dots$
$cfg - 1$	$dfg + 4$	$efg - 1$	$f^2g - 1$	$abeg^2 + 1$
$deg + 1$	$e^2g \dots$	$f^2 - 3$	$a^2bdg^2 - 4$	$b f^2g - 1$
$df^2 + 3$	$ef^2 - 3$	$a^2bcg^2 - 3$	$b^2fg + 4$	$c dg^2 - 1$
$ef - 3$	$a^2b^2g^2 + 1$	$bdfg \dots$	$bf^3 \dots$	$c efg - 2$
$e^2 \dots$	$b^2fg - 4$	$be^2g - 15$	$c^2f \dots$	$c f^2 + 3$
a^2b^2cg	$bdeg - 16$	$bef + 18$	$cdfg + 16$	$c^2fg + 4$
$bceg + 2$	$bdf^2 \dots$	$c^2fg + 15$	$c^2g - 22$	$de^2g - 1$
$bcf - 3$	$b^2f + 18$	$cdeg \dots$	$cef^2 + 6$	$d^2f^2 - 6$
$bd^2g - 4$	$c^2eg + 22$	$cdf^2 - 36$	$d^2eg + 8$	$ef + 3$
$bdef - 12$	$c^2f^2 + 3$	$cef + 9$	$df - 32$	$a^2b^2df - 3$
$be^3 + 15$	$cd^2g - 8$	$df \dots$	$d^2f + 36$	$b^2efg + 3$
$c^2dg + 1$	$cdef - 48$	$d^2f + 24$	$e^4 - 12$	$b f^3 \dots$
$c^2ef + 9$	$ce^2 + 12$	$d^2f - 12$	$a^2b^2cg - 3$	$b c^3f + 3$
$cdf^2 + 4$	$d^2f + 32$	$a^2b^2fg + 3$	$b^2c^2g \dots$	$b cdfg + 12$
$cd^2 - 21$	$d^2e^2 - 12$	$b^2ef - 18$	$b^2f - 3$	$b c^2g - 9$
$d^4 + 8$	$a^2b^2f^2 \dots$	$b^2deg + 36$	$b^2ef \dots$	$b cef - 9$
$a^2b^3eg - 3$	$a^2b^2cf \dots$	$b^2cf \dots$	$b^2cg - 18$	$b d^2eg - 4$
$b^2f^2 \dots$	$b^2cef - 6$	$b^2cef - 27$	$bcdeg + 48$	$b df^2 - 32$
$b^2cdg + 6$	$b^2d^2g + 32$	$b^2c^2eg - 9$	$bcd f^2 \dots$	$b d^2f + 66$
$b^2cef + 9$	$b^2def \dots$	$b^2cf^2 + 27$	$bce^2 - 18$	$b e^2 - 27$
$b^2df^2 + 32$	$b^2e^2 - 27$	$b^2cd^2g - 24$	$bd^2g - 32$	$c^2efg - 15$
$b^2de^2 - 39$	$b^2c^2dg - 36$	$b^2cdef \dots$	$b^2d^2f + 32$	$c^2deg + 21$
$b c^2g - 3$	$b^2c^2ef + 18$	$b^2cef + 27$	$bde^3 - 12$	$c^2df^2 + 39$
$b c^2df - 66$	$b^2cd^2f - 32$	$b^2cd^2f \dots$	$c^2eg - 12$	$c^2ef - 18$
$b c^2f + 18$	$b^2cde^2 + 84$	$b^2d^2e^2 - 12$	$df + 27$	$c d^2g - 8$
$b cd^2e + 76$	$b^2d^2e - 32$	$b c^2dg^2 + 12$	$c^2d^2g + 12$	$cd^2ef - 76$
$b d^4 - 32$	$c^4g + 12$	$b c^2ef - 27$	$c^2def - 84$	$cd^2e^2 + 45$
$c^4f + 27$	$c^4df + 12$	$b c^2df + 12$	$c^3f + 45$	$d^4f + 32$
$c^3de - 45$	$-45c^3e^2$		$cd^2f + 32$	$d^4e^2 - 20$
$c^2d^3 + 20$	$c^3d^2 + 20$	$c^2de^2 \dots$	$cd^3e^2 - 20$	
	$c^2d^3 \dots$	$c^2d^2e \dots$	$d^4e \dots$	
		$cd^3 \dots$	d^5	

[Page 385 presents the table $O = (\mid x, y)^8$, a 9-column array indexed by $x^{8-k}y^k$ ($k = 0, \dots, 8$) of coefficients of degree 5 and weights 11 through 19. The complete listing occupies the full page; column totals were not separately tabulated in the scan.]

Selected leading entries by column:

x^8 : $a^2fg^2 \dots, a^2beg^2 \dots, cfg^2 - 3, efg - 5, f^2 \dots, \dots$ (full series in scan).

x^7y : $a^2bfg^2 \dots, abf^2g \dots, cef^2 \dots, df^3 + 2, e^2f^2 - 10, \dots$

x^6y^2 : leading $ae fg^2 \dots, bf^2g - 3, cdfg + 10, defg - 5, ef^3 \dots$

x^5y^3 : leading $a^2g^3 \dots, a^2bfg^2 \dots, beg^2 - 5, cf^2g - 7, d^2fg + 2, de^2g - 10$

x^4y^4 (centre): $a^2f^2g \dots, abefg + 5, cef^2 - 12, deg^2 + 10, e^2fg - 5$

Subsequent columns $x^3y^5, x^2y^6, xy^7, y^8$ are obtained from the four leading columns by the duality $a \leftrightarrow g, b \leftrightarrow f, c \leftrightarrow e, d \leftrightarrow d$.

[Full coefficient detail in scan p. 385 (printed); see facsimile.]

$$P = (\mid x, y)^0, \text{ Invt.}$$

$a^2b^2g^4 \dots$
$a^2bfg^3 \dots$
$a^2b^2eg^3 \dots$
$cf^3 \dots$
$d^2g^2 + 1$
$defg - 6$
$df^2 + 4$
$e^2g + 4$
$ef^2 - 3$
$ab^2efg^2 \dots$
$f^3g \dots$
$abcdg^2 - 6$
$cefg + 18$
$cf^2 - 12$
$d^2fg + 12$
$d^2eg - 18$
$de^2f + 6$
$b^2cf + 4$
$a^2b^2dfg - 18$
$c^2e^2g - 24$
$c^2ef^2 + 30$
$cd^2eg + 54$
$cdf^2 - 12$
$cdeg^2 - 42$
$ce^3 + 12$
$d^3g - 20$
$d^2ef + 24$
$d^2 - 8$
$a^2b^2dg^2 + 4$
$efg - 12$
$f^2 + 8$
$b^2fg^2 - 3$
$cdfg \dots$
$ce^2g + 30$
$cef^2 - 24$
$d^2fg - 12$
$df^2 - 24$
$de^2f + 60$
$e^4 - 27$
$bc^3fg + 6$
$c^2deg - 42$
$c^2df^2 + 60$
$c^2e^2f - 30$
$cd^2g + 24$
$cdf^2 - 84$
$cde^2 + 66$
$d^4f + 24$
$d^4e^2 - 24$
$b^2c^2eg + 12$
$cf^2 - 27$
$-c^2d^2g \ 8$
$c^2def + 66$
$c^2e^2 - 8$
$cdf - 24$
$c^2d^2e^2 - 39$
$cd^2e + 36$
$d^4 - 8$

[Page 387 presents the table $Q = (\mid x, y)^6$, a 7-column array indexed by the column headers 51, 55, 55, 58, 55, 55, 51 at the top—these are the term-counts per column—and $x^{6-k}y^k$ for $k = 0, \dots, 6$. Each column lists the literal monomials of degree 6 and weights 15 through 21. The full content spans this page and the next (p. 388, marked “ Q (continued)”), and is best consulted against the facsimile.]

Selected leading entries from the principal columns of Q :

Column 1 (x^6 , header 51, weight 15): $a^2bdg^3 \dots, efg \dots, f^2 - 2, a^2bcf + 5, dfg \dots, e^2 - 8, a^2b^2cf - 2, cdfg + 6, c^2g + 8, cef - 10, -d^2eg + 10, df + 18, de^3 - 22, e^4 + 12, ab^2 + 4, dfg - 6, e^2g - 8, \dots$; column total ± 110 .

Column 2 (x^5y , header 55, weight 16): $a^2beg^3 \dots, f^2g \dots, bef \dots, cdfg + 6, c^2f + 8, cef - 10, d^2eg + 18, def - 22, e^3 + 12, abf^2 + 4, \dots$

Column 3 (x^4y^2 , header 55, weight 17): $a^2bfg^3 \dots, a^2b^2eg^2 \dots, cefg + 15, cf^2 - 10, dg - 15, dfg + 20, de^2g - 25, def - 10, e^2f + 15, a^2bf + 4, bdfg + 5, bef - 90, ce^2g + 40, cef^2 + 40, d^2fg + 50, df^2 + 10, de^2f - 40, e^4 - 15, bdef + 35, c^2deg - 45, \dots$

Column 4 (x^3y^3 , header 58, weight 18; centre): $a^2fg^3 \dots, a^2bfg^3 \dots, cf^2g \dots, df^2 \dots, defg + 20, e^2fg - 20, ef^3 + 20, a^2b^2fg^2 - 6, c^4g \dots, c^2f^2 - 20, c^2deg \dots, c^2df + 60, c^2e^3 - 20, cd^2eg \dots, cd^2f + 62, cd^2 - 60, bcd^2f + 66, bcde - 126, bd^3e + 24, c^4f - 50, c^3e + 60, c^2d^3 - 30, c^2de^2 + 60, c^2d^2 - 15, cd^3 \dots, d^6 \dots$

Columns 5, 6, 7 (x^2y^4 , xy^5 , y^6) are the weight-mirrors of columns 3, 2, 1 under the involution $a \leftrightarrow g$, $b \leftrightarrow f$, $c \leftrightarrow e$, $d \rightarrow d$.

[Full coefficient detail for table Q on facsimile pp. 387–388.]

$Q = (\mid x, y)^6$ (continued).

[Continuation of the table Q from p. 387 above; remaining literal monomials for each of the 7 columns and column totals. See facsimile.]

Selected continued entries:

Column 1 (continued, x^6): $a^2b^2c^2g + 76, cdef + 125, b^2 - 24, d^3 \dots, bcdef + 105, b^2ce^2 - 44, \dots, a^2b^2fg - 2, a^2b^2cg - 4, \dots, bdcef + 50, cdf - 44, -c^2d^2eg - 112, c^2d^3 - 12, c^2df + 28, cd^2e \dots, d^5 + 32, cd^4 \dots$

Column 4 (centre, x^3y^3 , continued): $bc^2d^2f + 74, a^2d^4 + 22, \dots, cd^3 + 10, c^4 - 25, d^3 \dots$

Column 7 (y^6 , continued, mirror of column 1): g -terms mirror a -terms above.

[End of paper 775; column totals at foot of p. 388 of scan.]

776.

ON THE JACOBIAN SEXTIC EQUATION.

[From the *Quarterly Journal of Pure and Applied Mathematics*, vol. XVIII. (1882), pp. 52–65.]

The Jacobian sextic equation has been discussed under the form

$$(z - \alpha)^6 - 4a(z - \alpha)^5 + 10b(z - \alpha)^3 - 4c(z - \alpha) + 5b^2 - 4ac = 0,$$

(see references at end of paper), but the connexion of this form with the general sextic equation has not, so far as I am aware, been considered. And although this is probably known, I do not find it to have been explicitly stated that the group of the equation is the positive half-group, or group of the 60 positive substitutions out of the 120 substitutions, which leave unaltered Serret's 6-valued function of six letters.

Invariantive Property of the Jacobian Sextic.

Taking $z - \alpha$ as the variable, and comparing the equation with the general sextic equation

$$= (a, b, c, d, e, f, g \mid z - \alpha, 1)^6 0,$$

we have

$$a, b, c, d, e, f, g = 1, -\frac{2}{3}a, 0, \frac{1}{2}b, 0, -\frac{2}{3}c, 5b^2 - 4ac;$$

the Jacobian equation is thus an equation

$$(a, b, c, d, e, f, g \mid x, y)^6 = 0,$$

for which $c = 0, e = 0, ag + 9bf - 20d^2 = 0$; but of course any equation, which can be by a linear transformation upon the variables brought into this form, may be regarded as a Jacobian equation.

Hence, using henceforward the small italic in place of the small roman letters, the Jacobian sextic may be regarded as an equation

$$= (a, b, c, d, e, f, g \mid x, y)^6 0,$$

linearly transformable into the form

$$= (a, b, 0, d, 0, f, g \mid x, y)^6 0,$$

where $ag + 9bf - 20d^2 = 0$. It is to be shown, that this implies a single relation between the four invariants A, B, C , and D of the sextic function.

I call to mind that the general sextic has five invariants A, B, C, D, E of the orders 2, 4, 6, 10, 15 respectively; the last of them E is not independent, but its square is equal to a rational and integral function of A, B, C, D ; and instead of D , we consider the discriminant Δ which is an invariant of the same order 10. The values of A, B, C are given. Table Nos. 31, 34, and 35 of my Third Memoir on Quantics, *Phil. Trans.*, vol. CXLVI. (1856), pp. 627–647, [144]; those of D, Δ, E were obtained by Dr Salmon, see his *Higher Algebra*, second ed. 1866, where the values of A, B, C, D, Δ, E are all given; only those of A, B, C , are reproduced in the third edition, 1876.

It may be remarked, that for the general form we have $A = ag - 6bf + 15ce - 10d^2$, and that B is the determinant

$$B = \begin{vmatrix} a, & b, & c, & d \\ b, & c, & d, & e \\ c, & d, & e, & f \\ d, & e, & f, & g \end{vmatrix},$$

containing 24 terms. C and Δ are complicated forms, the latter of them containing 246 terms. But writing $c = 0, e = 0$, there is a great reduction; we have

$A =$	$B =$	$C =$	$D =$
$ag + 1$	$ad^2g - 1$	$a^2d^2f + 1$	$ag^5 + 1$
$bf - 6$	$bf^2 + 1$	$, df^2 + 4$	
$d^2 - 10$	$a^2b^2fg - 2$	$abd^2fg + 12$	$a^2b^2fg^4 - 30$
	$d^4 + 1$	$, d^4g - 20$	$, , d^2g^3 - 300$
		$a^2b^3df + 4$	$, , d^4g^2 - 2500$
		$, , bf^3 + 8$	$, , d^6g + 3125$
		$, , b^3d^3 - 24$	$, , df^3g^3 - 15$
		$, , bd^3f + 24$	$, , f^4 - 4800$
		$, , d^5 - 8$	$, , bdf^3g^2 + 7500$
			$, , bdf^2g \dots$
			$, , bd^3f^2 \dots$
			$\dots$
			$, , d^6g + 30000$
			$, , d^8g + 50000$
			$a^2b^2dg^4 - 2500$
			$, , b^2d^2fg^3 \dots$
			$, , b^2d^3g^2 - 171300$
			$, , b^2d^4fg - 240000$
			$, , bd^5g^2 + 780000$
			$, , b^2d^5 + 1200000$
			$, , d^7 - 1000000$
			$, , d^8 - 1600000$
			$, , d^9 - 7500$
			$\dots - 11520$
			ab^3dfg^3
			$+ 50000$
			$, , b^3f^4g$
			$, , b^3d^4fg^2 + 83200$
			$a^2b^4g^4 - 3125$
			$, , b^4d^2fg^2 - 240000$
			$, , b^4f^4 - 331776$
			$, , b^4d^2f^3 + 1200000$
			$, , b^4d^4f^2 + 1843200$
			$, , b^4d^6fg - 1600000$
			$, , b^4d^8 - 2560000$

It is clear that these are all functions of ag, bf, d^2 and $a^2f^2 + b^2g^2$, say of α, β, δ and ϕ . In fact, A and B are functions of α, β, δ ; C contains two terms, coefficient 4, which are $\pm\sqrt{\delta} \cdot 4 \cdot \phi$; Δ contains two terms

$$-3125(a^4f^4 + b^4g^4),$$

which are $= -3125(\phi^2 - 2\alpha^2\beta^2)$; and also several pairs of terms, each which pair contains the factor ϕ . We thus have

$A =$	$B =$	$C =$	$\Delta =$
$\alpha + 1$	$\alpha\delta - 1$	$\alpha^2\delta + 1$	$\alpha^4 + 1$
$\beta - 6$	$\beta^2 + 1$	$\alpha\beta\delta + 12$	$\alpha^3\beta - 30$
$\delta - 10$	$\beta\delta - 2$	$\alpha\delta^2 - 20$	$\alpha^3\delta - 300$
	$\delta^2 + 1$	$\beta^3 + 8$	$\alpha^2\delta^2 + 5840 (= 6250 - 410)$
		$\beta^2\delta - 24$	$\alpha^2\beta^2 - 15$
		$\beta\delta^2 + 24$	$\alpha^2\beta\delta - 4800$
		$\delta^3 - 8$	$\alpha^2\beta^3 + 3000$
			$\alpha^2\beta^2\delta - 171300$
			$\alpha^2\beta\delta^2 + 780000$
			$\alpha^2\delta^3 - 1000000$
			$\alpha^2\beta^4 - 11520$
			$\alpha^2\beta^3\delta + 83200$
			$\beta^4 - 331776$
			$\beta^3\delta + 1843200$
			$\beta^2\delta^2 - 2560000$
			$\phi\sqrt{\delta} \cdot \alpha^3 - 2500$
			, , $\alpha^2\beta - 7500$
			, , $\alpha\delta + 50000$
			, , $\beta^2 - 240000$
			, , $\beta\delta + 1200000$
			, , $\delta^2 - 1600000$
			$\phi^2 - 3125$

We have *ante*, the relation $\alpha + 9\beta - 20\delta = 0$, and using this to eliminate α , we have A, B, C as functions of β, δ, ϕ (that is, of bf, d^2 and $a^2f^2 + b^2g^2$). Effecting the substitution, we find the values of A, B, C without difficulty. As regards the value of Δ , this is

$$= -3125\phi^2 + 2\phi K\sqrt{\delta} + \text{terms without } \phi,$$

where

$$\begin{aligned} 2K = & -2500(81\beta^2 - 360\beta\delta + 400\delta^2) \\ & + 7500(-9\beta^2 + 20\beta\delta) \\ & + 50000(-9\beta\delta + 20\delta^2) \\ & - 240000(\beta^2) \\ & + 1200000(\beta\delta) \\ & - 1600000(\delta^2), \end{aligned}$$

or, reducing and dividing by 2,

$$K = -3125(60\beta^2 - 240\beta\delta + 256\delta^2).$$

The calculation of the terms without ϕ is much more laborious, but they come out

$$= -3125(60\beta^2 - 240\beta\delta + 256\delta^2)^2 \delta.$$

Hence the value of Δ is

$$\begin{aligned} \Delta = & -3125 \{ \phi^2 \\ & + 2\phi(60\beta^2 - 240\beta\delta + 256\delta^2)\sqrt{\delta} \\ & + (60\beta^2 - 240\beta\delta + 256\delta^2)^2 \delta \}, \end{aligned}$$

say this is

$$\Delta = -3125 h^2,$$

where

$$h = \phi + (60\beta^2 - 240\beta\delta + 256\delta^2)\sqrt{\delta},$$

that is,

$$= a^2 f^2 + b^2 g^2 + (60b^2 d^2 f^2 - 240bd^4 f + 256d^6).$$

The values of A, B, C , and the foregoing value of h then are

$$\begin{array}{llll} A = & B = & C = & h = \\ \beta - 15 & \beta^2 + 1 & \beta^3 + \beta\delta & \beta^2\delta + 60 \\ \delta + 10 & \beta\delta + 7 & \beta^2\delta + 51 & \beta\delta\sqrt{\delta} - 240 \\ & \delta^2 - 19 & \beta\delta^2 + 84 & \delta^2\sqrt{\delta} + 256 \\ & & \delta^3 - 8 & \phi + 1 \\ & & \phi\sqrt{\delta} + 4 & \end{array}$$

We may, if we please, regard β, δ, ϕ as irrational invariants of the sextic, viz. A, B, C being rational and integral functions of β, δ, ϕ , we have conversely β, δ, ϕ irrational functions of A, B, C ; and then the equation for h , say

$$\frac{h^2 (= \Delta)}{25\phi\sqrt{\delta}} = \phi + \sqrt{\delta}(60\beta^2 - 240\beta\delta + 256\delta^2),$$

is the invariantive relation which characterises the Jacobian sextic.

The expression for Δ in terms of A, B, C, D is

$$\Delta = A^5 - 375A^3B - 625A^2C + 3125D,$$

and it was in the foregoing investigation proper to use Δ in place of D . But I annex the value of D for the case in question $b = 0, f = 0$; and also its value in terms of $\alpha, \beta, \delta, \phi$. These are

$D =$	$D =$
$ag^5 - 1$	$\phi^2 - 1$
$, d^2fg - 12$	$\alpha^2f^2 + 2$
$, d^5 + 5$	$\alpha g^5 - 12$
$a^2b^2d^2f - 90$	$\beta - 72$
$, , b^2df^2 - 48$	
$, bdf^2f + 246$	$bc^2f^4 + 168$
$, , bd^3f^2 + 480$	$, , \alpha^2$
$, d^6f - 258$	$, , \beta^4 + 552$
$, d^8 - 432$	$\delta^2\alpha^2 + 5$
$ab^3df^4 - 12$	$, , \alpha\beta + 246$
$, b^3f^5 - 12$	$\delta\alpha\beta^2 + 240$
$, , b^3d^3f^2g + 168$	$, , \beta^3 - 976$
$, , b^3d^5fg + 240$	$\delta^2\alpha^2 - 258$
$, , bd^6fg - 168$	$\delta\alpha\beta - 168$
$, , d^8g - 228$	$, , \beta^2 + 336$
$a^2b^4f^6 - 1$	$\delta^3\alpha - 228$
$, b^4d^2f^4 - 48$	$\delta\beta - 408$
$, , b^4d^4f^2 - 72$	$\delta^4 - 240$
$, , b^4d^4f^2 + 480$	$\phi\sqrt{\delta}\alpha\beta - 12$
$, b^4d^6f + 552$	$, \beta^2 - 48$
$, , b^4d^8 - 432$	$\phi\sqrt{\delta}\beta - 480$
$, , b^4d^8f^2 - 976$	$\phi\sqrt{\delta} - 432$
$, , b^4d^6f^4 + 336$	
$, , b^6d^8 + 408$	
$, , d^{10} - 248$	

The Group of the Jacobian Sextic.

The solution of the Jacobian sextic equation depends upon that of a quintic; in fact, calling the roots $z_1, z_2, z_3, z_4, z_5, z_6$, then there exists a quintic having the roots

$$\begin{aligned} &\sqrt{(z_1 - z_2.z_1 - z_3.z_4 - z_5)}, \\ &\sqrt{(z_2 - z_4.z_2 - z_5.z_3 - z_6)}, \\ &\sqrt{(z_3 - z_5.z_4 - z_6.z_1 - z_2)}, \\ &\sqrt{(z_4 - z_6.z_5 - z_1.z_2 - z_3)}, \\ &\sqrt{(z_5 - z_1.z_6 - z_2.z_3 - z_4)}, \end{aligned}$$

the coefficients of which are rational functions of the coefficients a, b, d, f, g , and of the fourth root of the discriminant, i.e. $\sqrt[4]{h}$. But the meaning of this has not, so far as I am aware, been noticed. Passing to the quintic whose roots are the squares of the foregoing values, i.e. $z_1 - z_2, z_1 - z_3, z_4 - z_5$, &c., the coefficients are here rational functions of a, b, d, f, g and h ; that is, they are rational functions of a, b, d, f, g . The symmetrical functions of these roots $z_1 - z_2, z_1 - z_3, z_4 - z_5$, &c., are thus rational functions of the coefficients of the sextic; each such rational function is a 12-valued function of $z_1, z_2, z_3, z_4, z_5, z_6$, invariable by all the substitutions of a group of 60 substitutions; and therefore also every like 12-valued function of the roots $z_1, z_2, z_3, z_4, z_5, z_6$ is invariable by the substitutions of this group of 60; or, in other words, this group of 60 is the group of the Jacobian sextic equation.

I write for convenience, in this section only,

$$z_1, z_2, z_3, z_4, z_5, z_6 = f, a, b, c, d, e;$$

and writing further ab for shortness instead of $a - b$, &c., (so that of course $ba = -ab$), and putting

$$B = -ab.cd.ef, \quad C = -ac.bf.de, \quad D = ad.bc.ef, \quad E = ae.bd.cf, \quad F = af.be.cd,$$

then the five functions are B, C, D, E, F , and the group of 60 which leaves unaltered every symmetrical function of these functions is made up of the substitutions

1.			1	
$ab \cdot ce,$	$ab \cdot df,$	$ce \cdot df,$	15	
$ac \cdot bf,$	$ac \cdot de,$	$bf \cdot de,$		
$ad \cdot be,$	$ad \cdot ef,$	$be \cdot ef,$		
$ae \cdot bd,$	$ae \cdot cf,$	$bd \cdot cf,$		
$af \cdot bc,$	$af \cdot cd,$	$bc \cdot cd,$		
$abcde,$	$acebd,$	$adbec,$	$aedbc,$	24
$afbce,$				
$abdef,$	$abefc,$	$ae fcb,$	$aecbf,$	
$afced,$				
$afdbc,$	$adfbe,$	$aebfd,$	$afedb,$	
$bdcef,$				
	$acdf e,$	$ae fde,$	$adecf,$	
	$adcfb,$	$ab fed,$	$ac bdf,$	
	$bc fde,$	$bedfe,$	$bfecd,$	
$abc \cdot dfe,$	$acb \cdot def,$		20	
$abd \cdot cfe,$	$adb \cdot cef,$			
$abe \cdot cfd,$	$aeb \cdot cdf,$			
$abf \cdot ced,$	$afb \cdot cde,$			
$acd \cdot bef,$	$adc \cdot bfe,$			
$ace \cdot bfd,$	$aec \cdot bdf,$			
$acf \cdot bed,$	$afc \cdot bde,$			
$ade \cdot bfc,$	$aed \cdot bcf,$			
$adf \cdot bce,$	$afd \cdot bce,$			
$ae f \cdot bcd,$	$afe \cdot bcd,$		60	

where the symbols ab , $abcde$, abc , &c. denote cyclical substitutions. It is easy to verify that each of these substitutions does in fact merely permute B, C, D, E, F ; thus

$$\begin{array}{rccccc}
 & B & C & D & E & F \\
 abcde \text{ on} & -ab.ce.df, & -ac.bf.de, & ad.bc.ef, & ae.bd.cf, & af.be.cd \\
 = & -bc.da.ef, & bd.cf.ea, & be.cd.af, & ba.ce.df, & bf.ca.de \\
 = & +ad.bc.ef, & ae.bd.cf, & -af.be.cd, & -ab.ce.df, & ac.bf.de \\
 = & D & E & F & B & C,
 \end{array}$$

which (expressed as a cyclical substitution) is $= BDFCE$, and so in other cases.

We may to the foregoing 60 substitutions join the 60 other substitutions:

$$\begin{array}{lll}
 cdef, & cfed, & 30 \\
 bdf e, & bef d, & \\
 bcef, & bfce, & \\
 bcdf, & bf dc, & \\
 bced, & bdec, & \\
 aedf, & afde, & \\
 acef, & afec, & \\
 acfd, & adfc, & \\
 adce, & aecd, & \\
 & aefb, & \\
 & afbd, & \\
 abfe, & & \\
 adbf, & & \\
 abcd, & adcb, & \\
 abcf, & afcb, & \\
 aebe, & acbe, & \\
 abdc, & acdb, & \\
 ab \cdot cd \cdot ef, & ab \cdot ef \cdot de, & 10 \\
 ac \cdot bd \cdot ef, & ac \cdot be \cdot df, & \\
 ad \cdot be \cdot cf, & ad \cdot bf \cdot ce, & \\
 ae \cdot bc \cdot df, & ae \cdot bf \cdot cd, & \\
 af \cdot bc \cdot de, & af \cdot bd \cdot ce, & \\
 abcefd, & adfecb, & 20 \\
 abfdec, & acedfb, & \\
 abecdf, & afdceb, & \\
 abdfce, & aecfdb, & \\
 acfbde, & aedbf c, & \\
 acbfed, & adefbc, & \\
 acdebf, & afbedc, & \\
 adbcfe, & aefcbd, & \\
 adebcf, & afebcd, & \\
 ae bdcf, & afcdbe, & 60
 \end{array}$$

each of which changes B, C, D, E, F into a permutation of $-B, -C, -D, -E, -F$.

The 60 and 60 substitutions form together a group of 120 substitutions, which leave unaltered any even symmetrical function of B, C, D, E, F , or say any symmetrical function of B^2, C^2, D^2, E^2, F^2 ; such a function is thus a 6-valued function of a, b, c, d, e, f , viz. it is Serret's 6-valued function of 6 letters.

Transformation of the Jacobian Sextic into the Resolvent Sextic of a special quintic equation.

Starting from the Jacobian Sextic Equation

$$(a, b, 0, d, 0, f, g \mid z, 1)^6 = 0,$$

$ag + 9bf - 20d^2 = 0$, I effect upon it the Tschirnhausen transformation

$$X = -az^2 - 6bz - 10d;$$

which, it may be remarked, is a particular case of the Tschirnhausen-Hermite form

$$\begin{aligned} X(= A + B) &= (az + b)B + (az^2 + 6bz + 5c)C + (az^3 + 6bz^2 + 15cz + 10d)D \\ &+ (az^4 + 6bz^3 + 15cz^2 + 20dz + 10e)E \\ &+ (az^5 + 6bz^4 + 15cz^3 + 20dz^2 + 15ez + 6f)F. \end{aligned}$$

Writing for convenience $Y = X + 10d$, $Z = X - 10d$, this is

$$+Y az^2 + 6bz - \dots = 0,$$

and we thence have

$$\begin{aligned} az^3 + 6bz^2 + \dots + Yz \cdot &= 0, \\ +az^4 + 6bz^3 \cdot + Yz^2 \cdot &= 0, \\ -Zz^2 \cdot -6fz - g &= 0, \\ -Zz^3 \cdot -6fz^2 - gz \cdot &= 0, \\ -Zz^4 \cdot -6fz^3 - gz^2 \cdot \cdot &= 0, \end{aligned}$$

or, eliminating, the resulting equation is

$$\begin{vmatrix} \cdot & a, & 6b, & \cdot & Y, & \cdot \\ \cdot & \cdot & a, & 6b, & \cdot & Y \\ a, & 6b, & \cdot & Y, & \cdot & \cdot \\ \cdot & \cdot & Z, & \cdot, & 6f, & g \\ \cdot & Z, & \cdot & 6f, & g, & \cdot \\ Z, & \cdot & 6f & g, & \cdot & \cdot \end{vmatrix} = 0.$$

The developed form is most easily obtained by expanding the determinant in the form

$$123 \cdot 456 - 456 \cdot 123; \text{ \&c.},$$

where the terms 123, &c., belong to the matrix

$$\begin{bmatrix} \cdot & a, & 6b, & \cdot & Y, & \cdot \\ \cdot & \cdot & a, & 6b, & \cdot & Y \\ a, & 6b, & \cdot & Y, & \cdot & \cdot \end{bmatrix}$$

and those of $\overline{123}$, &c., to the matrix

$$\begin{bmatrix} \cdot & \cdot & Z, & \cdot & 6f, & g \\ \cdot & Z, & \cdot & 6f, & g, & \cdot \\ Z, & \cdot & 6f, & g, & \cdot & \cdot \end{bmatrix}$$

The several terms are

$$\begin{array}{lll} 123 \cdot 456 & + - a^3 & \cdot - g^2 \\ & - 6a^2b & \\ -124 \cdot 356 & \cdot + & - 6fg^2 \\ +125 \cdot 346 & & \cdot \\ -126 \cdot 345 & - a^2Y & - g^2 - 216f^2 \\ & + - 36ab^2 & fZ - 216f \\ +134 \cdot 256 & - a^2Y & \cdot - fZ \\ & + - 6abY & \\ -135 \cdot 246 & + 6abY & - 6fgz \\ & - \cdot & - 6fgz \\ +136 \cdot 245 & & - 36f^2z \\ +145 \cdot 236 & & \\ -146 \cdot 235 & & \\ +156 \cdot 234 & & \\ -234 \cdot 156 & - a^2F - 216b^2 & - gZ^2 \\ +235 \cdot 146 & + 6abY & \\ -236 \cdot 145 & - - 36b^2Y & fZ \\ -245 \cdot 136 & - 36b^2Y & \\ +246 \cdot 135 & & - gZ \\ -256 \cdot 134 & & \\ +345 \cdot 126 & & 36fz \\ -346 \cdot 125 & & \\ +356 \cdot 124 & + aY^2 & gz^2 \\ -456 \cdot 123 & & \\ & - 6bY^2 & \cdot - 6fZ^2 \\ & Y^2 + -rt & - gZ^2 \\ & - 6bY^2 & - 6fZ^2 \\ & + & \\ & - - Y^3 & Z^3 \end{array}$$

Hence, collecting and reducing, the equation is

$$\begin{aligned} 0 &= Y^3Z^3. \\ &+ Y^2Z^2 \cdot (3ag + 72bf) \\ &+ YZ \cdot (3ag^2 + 36agbf + 1296b^2f^2) \\ &+ Y \cdot 216a^2f^3 \\ &+ Z \cdot -216b^3g^2 \\ &+ a^3g^3 - 36ag^2bf, \end{aligned}$$

where Y, Z denote $X + 10d, X - 10d$ respectively, and consequently $YZ = X^2 - 100d^2$. Hence, writing as before $\alpha, \beta, \delta, \phi$ to denote ag, bf, d^2 and $a^2f^2 + b^2g^2$ respectively, the result finally is

$$\left(1, \alpha + 3, \alpha^2 + 3, \alpha^3 + 216, \phi\sqrt{\delta} + 2160 \mid X, 1 \right)^6 = 0,$$

with the developed coefficients

$$\begin{aligned} &+0, \\ &\beta + 72, \\ &\alpha\beta + 36, \quad b^3f - 216, \\ &\delta - 300, \\ &\alpha\delta - 600, \\ &\beta^2 + 1296, \\ &\beta\delta - 14400, \\ &\delta^3 + 30000, \\ &-\alpha^3\beta - 36, \\ &\alpha^2\delta - 30, \\ &\alpha^2\delta - 360, \\ &\alpha\delta^2 + 30000, \\ &\beta^2\delta + 12960, \\ &\beta\delta^2 + 720000, \\ &\delta^3 - 1000000. \end{aligned}$$

where observe that the coefficient of the term in X is $216(a^2f^3 - b^3g^2) = 216\sqrt{\phi^2 - 4\alpha^2\beta^3}$.

We have as before $ag + 9bf - 20d^2 = 0$, that is, $\alpha + 9\beta - 20\delta = 0$; and using this equation to eliminate α , also in the constant term writing its value for ϕ in terms of h ,

$$\phi = h + (-60\beta^2 + 240\beta\delta - 256\delta^2)\sqrt{\delta},$$

the new equation is

$$\left(1, \beta - 9, \beta^2 - 243, -216\sqrt{h}, A\sqrt{\delta} + 432 \mid X, 1 \right)^6 = 0,$$

with developed coefficients

$$\begin{aligned} &\delta + 48, \quad \beta\delta - 1872, \\ &\beta^2 + 3840, \\ &\beta^3 - 729, \\ &\beta^2\delta + 4184, \\ &\beta\delta^2 - 11520, \\ &\delta^3 + 8292, \end{aligned}$$

where

$$\begin{aligned} \Lambda &= (h + (-60\beta^2 + 240\beta\delta - 256\delta^2)\sqrt{\delta})^2 - 4(-9\beta + 20\delta)^2\beta^3 \\ &= h^2 + 2h\sqrt{\delta} \begin{vmatrix} \beta^2 - 60 \\ \beta\delta + 240 \\ \delta^2 - 256 \end{vmatrix} - 4(\beta - 4\delta)^2(9\beta - 16\delta)^2. \end{aligned}$$

It is to be shown that this Tschirnhausen-transformation of the Jacobian sextic is, in fact, the resolvent sextic of the quintic equation

$$(a, 0, c, 0, e, f \mid x, 1)^5 = 0,$$

where

$$a = 1, \quad c = 2d, \quad e = -9bf + 36d^2, \quad f^2 = 216h.$$

I consider the general quintic $f^5 = (a, b, c, d, e, f \mid x, 1)^5 = 0$; taking the roots to be x_1, x_2, x_3, x_4, x_5 , and writing

$$\begin{aligned}\phi_1 &= 12345 - 24135, \\ \phi_2 &= 13425 - 32145, \\ \phi_3 &= 14235 - 43125, \\ \phi_4 &= 21435 - 13245, \\ \phi_5 &= 31245 - 14325, \\ \phi_6 &= 41325 - 12435,\end{aligned}$$

where 12345 is used to denote the function

$$= (x_1x_2 + x_2x_3 + x_3x_4 + x_4x_5 + x_5x_1)\sqrt{(20)},$$

(this numerical factor $\sqrt{(20)}$ being inserted for greater convenience), then the equation whose roots are $\phi_1, \phi_2, \phi_3, \phi_4, \phi_5, \phi_6$, which equation may be regarded as the resolvent sextic of the given quintic equation, is

$$\left(1, -5a^2x, 5a^4x^2, -\sqrt{(\square)} \cdot a^6, +5 \mid \phi, 1\right)^6 = 0,$$

with developed terms (selected):

$$\begin{aligned}&ae + 1 \\ &-a^2df + 1 \\ &+1 \\ &+a^3cf^2 + 1 \\ &a^2e^2 + 1 \\ &-2a^3def \\ &+ \&c. \\ &\&c.\end{aligned}$$

$D = +a^4f^4 \&c.$, the discriminant of the quintic: see p. 274* of my paper “On a new auxiliary equation in the theory of equations of the fifth order,” *Phil. Trans.*, t. CLI. (1861), pp. 263–276, [268].

I now write $b = 0$, $d = 0$, but, to avoid confusion again, write roman instead of italic letters, viz. I consider the resolvent sextic of the quintic equation

$$(a, 0, c, 0, e, f \mid x, 1)^5 = 0.$$

Many of the terms thus vanish, and the equation assumes the form

$$\left(1, -5a^4, 5a^8, -a^{12}\sqrt{\square}, +5 \mid \phi X, 1\right)^6 = 0,$$

with developed coefficients

$$\begin{aligned}&ae + 1, -3a^2cf^2 + 1 \\ &c^2 + 3, ac^2e - 2 \\ &c^2 + 15, ac^2f^2 + 1 \\ &a^3c^2e^2 - 11 \\ &ac^3e + 35 \\ &c^5 - 25\end{aligned}$$

and then if, as before,

$$\begin{aligned}a &= 1, c = 2d, e = -9bf + 36d^2, f^2 = 216h, \\ a &= 1, c = 2\sqrt{\delta}, e = -9\beta + 36\delta, f^2 = 216h,\end{aligned}$$

or say

*[This Collection, vol. iv., p. 321.]

this becomes identical with the foregoing Tschirnhausen-transformation equation; thus

$$\begin{aligned} -9ae + 3c^2 &= -9\beta + 36\delta + 12\delta, \\ &= \delta + 48; \end{aligned}$$

and similarly

$$\begin{aligned} 3a^3e^2 - 2ac^2e + 15c^4 &= \beta^2 + 243, \\ -\beta\delta - 1872, \\ +\delta^2 + 3840. \end{aligned}$$

So for the constant term, $+a^6cf^2$ gives the term $432h\sqrt{\delta}$, and $+a^6e^5$, &c., give the remaining terms $-729\beta^6$, &c. of the value in question.

It only remains to verify the equality of the coefficients of X ,

$$216\sqrt{h} = \sqrt{D} \quad \text{or} \quad 46656h = D.$$

Here D , the discriminant of the quintic $(a, 0, c, 0, e, f \mid x, 1)^5$, from the general form (see my Second Memoir on Quantics, [141], or Salmon's *Higher Algebra*, third edition, p. 209) putting therein $b = 0$, $d = 0$, is

$$\begin{aligned} D &= +a^4f^4 \cdot 1, \\ &+ a^3ce^2f + 160, \\ &a^3e^5 + 256, \\ &a^2c^3ef^2 - 1440, \\ &a^2c^3e^3 - 2560, \\ &ac^5f + 3456, \\ &ac^4e^2 + 6400, \end{aligned}$$

and writing for a, c, e, f their values $1, 2\sqrt{\delta}, 9(-\beta + 4\delta), 216h$, the value becomes

$$\begin{aligned} D &= (216)^2 \cdot h^2. \\ &+ 432h\sqrt{\delta} \cdot 12960 (\beta - 4\delta)^2 \\ &+ 34560 (\beta - 4\delta) \delta \\ &+ 55296 \delta^2 \\ &- 256 \cdot 9^5 \cdot -(\beta - 4\delta)^5 \\ &- 10240 \cdot 9^4 \cdot (\beta - 4\delta)^4 \delta \\ &- 102400 \cdot 9^3 \cdot (\beta - 4\delta)^3 \delta^2. \end{aligned}$$

The whole divides by $(216)^2$, and we thus obtain

$$\begin{aligned} \Lambda &= h^2 + 2h\sqrt{\delta} \cdot 60 (-\beta + 4\delta)^2 \cdot +(-\beta - 4\delta)^5 \cdot 324 (\beta - 4\delta)^4 \\ &\quad -240 (-\beta + 4\delta) \delta \quad + 1440 (\beta - 4\delta) \delta \\ &\quad +256 \delta^2 \quad + 1600 \delta^2, \end{aligned}$$

which is, in fact, equal to the foregoing value of Λ .

The conclusion is that, starting from the Jacobian sextic

$$f = (a, b, 0, d, 0, f, g \mid z, 1)^6 = 0,$$

where $ag + 9bf - 20d^2 = 0$, and effecting upon it the Tschirnhausen-transformation

$$X = -az^2 - 6bz - 10d,$$

so as to obtain from it a sextic equation in X , this sextic equation in X is the resolvent sextic of the quintic equation

$$(1, 0, c, 0, e, f \mid x, 1)^5 = 0,$$

where

$$c = 2d, \quad e = -9bf + 36d^2, \quad f = \sqrt{(216h)},$$

and, Δ being the discriminant of the Jacobian sextic, then

$$h = \frac{1}{5\sqrt{5}}\sqrt{(-\Delta)}, \quad = a^2f^2 + b^2g^2 + 60b^2d^2f^2 - 240bd^4f + 256d^6.$$

As to the subject of the present paper, see in particular Brioschi, “Ueber die Auflösung der Gleichungen vom fünften Grade,” *Math. Annalen*, t. XIII. (1878), pp. 109–160, and the third Appendix to his translation of my *Elliptic Functions*, Milan, 1880, each containing references to the earlier papers.

777.

A SOLVABLE CASE OF THE QUINTIC EQUATION.

[From the *Quarterly Journal of Pure and Applied Mathematics*, vol. XVIII. (1882), pp. 154–157.]

The roots of the general quintic equation

$$(a, b, c, d, e, f \mid x, 1)^5 = 0$$

may be taken to be

$$\begin{aligned} & -\frac{b}{a} + B + C + D + E, \\ & -\frac{b}{a} + \omega B + \omega^2 C + \omega^3 D + \omega^4 E, \\ & -\frac{b}{a} + \omega^2 B + \omega^4 C + \omega D + \omega^3 E, \\ & -\frac{b}{a} + \omega^3 B + \omega C + \omega^4 D + \omega^2 E, \\ & -\frac{b}{a} + \omega^4 B + \omega^3 C + \omega^2 D + \omega E, \end{aligned}$$

where ω is an imaginary fifth root of unity; and if one of the four functions B, C, D, E is = 0, say if $E = 0$ (this implies of course a single relation between the coefficients), then the equation is solvable.

Writing $x = \xi - \frac{b}{a}$, we have

$$(a, b, c, d, e, f \mid \xi - \frac{b}{a}, 1)^5 = (a', 0, c', d', e', f' \mid \xi, 1)^5,$$

where

$$\begin{aligned} a' &= a, \\ ac' &= ac - b^2, \\ a^2 d' &= a^2 d - 3abc + 2b^3, \\ a^3 e' &= a^3 e - 4a^2 bd + 6ab^2 c - 3b^4, \\ a^4 f' &= a^4 f - 5a^3 be + 10a^2 b^2 d - 10ab^3 c + 4b^5, \end{aligned}$$

and the roots of the new equation

$$(a', 0, c', d', e', f' \mid \xi, 1)^5 = 0$$

have the above-mentioned values, omitting therefrom the term $-\frac{b}{a}$: we find without difficulty

$$\begin{aligned} 2\frac{c'}{a} &= -BE - CD, \\ 2\frac{d'}{a} &= -B^2 D - BC^2 - CE^2 - D^2 E, \\ \frac{e'}{a} &= -B^3 C - B^2 E^2 + BCDE + BD^3 + C^3 E + C^2 D^2 - DE^3, \\ \frac{f'}{a} &= -B^5 + 5B^2 DE - 5B^2 C^2 E - 5B^2 CD^2 + 5BC^3 D + 5BCE^3 \\ &\quad - 5BD^3 E^2 - C^5 + 5CD^3 E - 5CD^2 E^2 - D^5 - E^5, \end{aligned}$$

and hence, when $E = 0$, we have

$$\begin{aligned} 2\frac{c'}{a} &= -CD, \\ 2\frac{d'}{a} &= -B^2 D - BC^2, \end{aligned}$$

$$\begin{aligned}\frac{e'}{a} &= -B^3C - BD^3 - C^2D^2, \\ \frac{f'}{a} &= -B^5 - 5B^2CD^2 + 5BC^3D - C^5 - D^5,\end{aligned}$$

or, as these may be written,

$$\begin{aligned}-2\frac{c'}{a} &= CD, \\ -2\frac{d'}{a} &= B^2D + BC^2, \\ -\frac{e'}{a} - \frac{c'^2}{a^2} &= B^3C - BD^3, \\ -\frac{f'}{a} &= B^5 + C^5 + D^5 - 10\frac{c'}{a}(B^2D - BC^2),\end{aligned}$$

equations which imply a single relation between the coefficients a' , c' , d' , e' , f' . Supposing this satisfied, we may attend only to the first three equations; or, writing for convenience,

$$\begin{aligned}\gamma &= -2\frac{c'}{a} = -\frac{2}{a^2}(ac - b^2), \\ \delta &= -2\frac{d'}{a} = -\frac{2}{a^3}(a^2d - 3abc + 2b^3), \\ \theta &= -\frac{e'}{a} - \frac{c'^2}{a^2} = -\frac{1}{a^4}\{a^2(ae - 4bd + 3c^2) + (ac - b^2)^2\},\end{aligned}$$

the equations are

$$\begin{aligned}\gamma &= CD, \\ \delta &= B(BD + C^2), \\ \theta &= B(D^3 - B^2C).\end{aligned}$$

51–2

The first equation gives $C = \frac{\gamma}{D}$, and substituting this value in the other two equations, we have

$$\begin{aligned}B^2D^2 + B\gamma^2 - \delta D &= 0, \\ B^3\gamma^2 + BD^4 + \theta D &= 0.\end{aligned}$$

Eliminating B , the result is obtained in the form $\det. = 0$, where in the last column of the determinant each term is divisible by D ; and omitting this factor, the result is

$$\begin{vmatrix} D^2, & \gamma^2, & 0, & -\delta D \\ 0, & D^2, & \gamma^2, & -\delta D \\ \gamma^2, & 0, & D^4, & \theta \\ 0, & \gamma^2, & 0, & D^4, \quad \theta \end{vmatrix} = 0.$$

If, in order to develop the determinant, we consider it as a sum of products, each first factor being a minor composed out of columns 1 and 2, and the second factor being the complementary minor composed out of columns 3, 4, 5 (the several products being of course taken each with its proper sign), the expansion presents itself in the form

$$\begin{aligned}& D^4(-\gamma^2\theta D^2 + \theta^2 D^3) \\ & - D^2\gamma^2(-\theta\gamma^2 D^2 + \theta D^5 - \delta^2 D^3) \\ & - \gamma^2 D^2(-\delta D^6 - \theta\gamma^4) \\ & + \gamma^4(-\gamma^4\delta D^2 - \delta^3 D^4 - \delta\gamma^2 D^4) \\ & - \gamma^8 D^2.\end{aligned}$$

Hence, collecting, and changing the sign of the whole expression, we obtain

$$BD^6 - (2\gamma^2\delta + \theta^2)D^4 + (-\gamma^4\delta + \delta^2\gamma^2\theta + \delta^2)D^3 + \gamma^8 = 0,$$

viz. a cubic equation for D^2 . We have then as above $C = \frac{\gamma}{D}$, and B is given rationally as the common root of the foregoing quadric and cubic equations satisfied by B .

Substituting for γ, δ, θ their values in terms of the original coefficients, the equation for D^2 becomes

$$\begin{cases} 2(a^2d - 3abc + 2b^3)(aD)^6 \\ + a^2[(ae - 4bd + 3c^2)^2a^2 + a^2(ac - b^2)^2(ae - 4bd + 3c^2) - 16(ac - b^2)^3(a^2d - 3abc + 2b^3)](aD)^4 \\ + [28(ac - b^2)^2(a^2d - 3abc + 2b^3) \\ + 4(ae - b^2)^2\{4a^2(ac - b^2)(a^2d - 3abc + 2b^3)(ae - 4bd + 3c^2) + 8(a^2d - 3abc + 2b^3)^2\}](aD)^3 \\ - 128(ac - b^2)^3\{a^2(ae - 4bd + 3c^2)(ac^2 - b^2)^2\} = 0, \end{cases}$$

and the solution of the given quintic equation thus ultimately depends upon that of this cubic equation.

778.

[ADDITION TO MR HUDSON'S PAPER "ON EQUAL ROOTS OF EQUATIONS."]

[From the *Quarterly Journal of Pure and Applied Mathematics*, vol. XVIII. (1882), pp. 226–229.]

It seems desirable to present in a more developed form some of the results of the foregoing paper.

Thus, if the equation $(a_0, a_1, \dots, a_n \mid x, 1)^n = 0$ of the order n has $n - v$ equal roots, where v is not $> \frac{1}{2}n - 1$, then we have $\psi(r, v + 1, m) = 0$, where m has any one of the values $0, 1, \dots, n - 2v - 2$, and r any one of the values

$$2v + 2, 2v + 3, \dots, n - m.$$

The signification is

$$\begin{aligned} \psi(r, v + 1, m) = & r \cdot 1 \cdot \frac{1}{[r]^{v+1}} a_m a_{r+m} \\ & - (r - 2) \cdot \frac{v + 1}{1} \cdot \frac{1}{[r - 1]^{v+1}} a_{m+1} a_{r+m-1} \\ & + (r - 4) \cdot \frac{v + 1 \cdot v + 2}{1 \cdot 2} \cdot \frac{1}{[r - 2]^{v+1}} a_{m+2} a_{r+m-2} \\ & + (-)^s (r - 2s) \cdot \frac{[v + 1]^s}{[s]^s} \cdot \frac{1}{[r - s]^{v+1}} a_{m+s} a_{r+m-s} \\ & \dots \\ & + (-)^{v+1} (r - 2v - 2) \cdot \frac{1}{[r - v - 1]^{v+1}} a_{m+v+1} a_{r+m-v-1}. \end{aligned}$$

Thus, when $v = 0$, the condition is

$$r \cdot \frac{1}{r \cdot r - 1} a_m a_{r+m} - (r - 2) \cdot \frac{1}{r - 1 \cdot r - 2} a_{m+1} a_{r+m-1} = 0,$$

that is,

$$a_m a_{r+m} - a_{m+1} a_{r+m-1} = 0,$$

satisfied when the equation has all its roots equal.

The values of m are $0, 1, 2, \dots, n - 2$, and those of r are $2, 3, \dots, n - m$; in particular, if $m = 0$, the values of r are $2, 3, \dots, n$, and the corresponding conditions are

$$\begin{aligned} a_0 a_2 - a_1^2 &= 0, \\ a_0 a_3 - a_1 a_2 &= 0, \\ a_0 a_n - a_1 a_{n-1} &= 0, \end{aligned}$$

and so for the different values of m up to the final value $r = 2$, for which $m = n - 2$, and the condition is

$$a_{n-2} a_n - a_{n-1}^2 = 0;$$

we have thus, it is clear, the whole series of conditions included in

$$\begin{vmatrix} a_0, & a_1, & a_2, & \dots, & a_{n-1} \\ a_1, & a_2, & a_3, & \dots, & a_n \end{vmatrix} = 0,$$

which are obviously satisfied in the case in question of the roots being all equal.

Again, when $v = 1$, the condition for $n - 1$ equal roots is

$$r \cdot 1 \cdot \frac{1}{r \cdot r - 1 \cdot r - 2} a_m a_{r+m} - (r - 2) \cdot \frac{2}{1} \cdot \frac{1}{r - 1 \cdot r - 2 \cdot r - 3} a_{m+1} a_{r+m-1} + (r - 4) \cdot 1 \cdot \frac{1}{r - 2 \cdot r - 3 \cdot r - 4} a_{m+2} a_{r+m-2} = 0;$$

that is,

$$r - 1 \cdot r - 2 - r - 1 \cdot r - 3 + r - 2 \cdot r - 3 = 0;$$

or, what is the same thing,

$$(r - 3) a_m a_{r+m} - 2(r - 2) a_{m+1} a_{r+m-1} + (r - 1) a_{m+2} a_{r+m-2} = 0,$$

where $n = 4$ at least, and m, r have the values

$$m = 0, 1, 2, \dots, n - 4$$

$$r = \begin{cases} 4, 4, \dots, 4 \\ 5, 5 \\ \vdots \\ n - 1 \\ n \end{cases}$$

thus, when $n = 4$, the only values are $m = 0, r = 4$, and the condition is

$$a_0 a_4 - 4a_1 a_3 + 3a_2^2 = 0.$$

Similarly, when $v = 2$, the condition for $n - 2$ equal roots is found to be

$$\frac{a_m a_{r+m}}{r - 1 \cdot r - 2 \cdot r - 3} - \frac{3a_{m+1} a_{r+m-1}}{r - 1 \cdot r - 2 \cdot r - 4} + \frac{3a_{m+2} a_{r+m-2}}{r - 2 \cdot r - 3 \cdot r - 5} - \frac{a_{m+3} a_{r+m-3}}{r - 3 \cdot r - 4 \cdot r - 5} = 0;$$

or, what is the same thing,

$$\begin{aligned} & r - 4 \cdot r - 5 \cdot a_m a_{r+m} \\ & - 3 \cdot r - 2 \cdot r - 5 \cdot a_{m+1} a_{r+m-1} \\ & + 3 \cdot r - 1 \cdot r - 4 \cdot a_{m+2} a_{r+m-2} \\ & - r - 1 \cdot r - 2 \cdot a_{m+3} a_{r+m-3} = 0, \end{aligned}$$

where $n = 6$ at least, and m, r have the values

$$m = 0, 1, \dots, n - 6$$

$$r = \begin{cases} 6, 6, \dots, 6 \\ 7, 7 \\ \vdots \\ n - 1 \\ n \end{cases}$$

Observe that the sum of the coefficients is 0, *viz.*

$$(r - 4)(r - 5) - 3(r - 2)(r - 5) + 3(r - 1)(r - 4) - (r - 1)(r - 2) = 0,$$

this should obviously be the case, since the conditions for $n - 2$ equal roots must be satisfied when the roots are all of them equal; and the property serves as a verification.

It is to be remarked that the equation $\psi(r, v + 1, m) = 0$ does not in all cases give all the conditions for the existence of $n - v$ equal roots in an equation of the order n ; thus when $n = 3$ and $v = 1$, we cannot by means of it obtain the condition that a cubic equation may have 2 equal roots. The problem really considered is that of the determination of those quadric functions of the coefficients which vanish in the case of $n - v$ equal roots; and in the case in question ($n = 3, v = 1$) there is no quadric function which vanishes, but the condition depends on a cubic function.

The question of the quadric functions which vanish in the case of $n - v$ equal roots, and to a small extent that of the cubic functions which thus vanish, is considered in Dr Salmon's "Note on the conditions that an equation may have equal roots," *Camb. and Dublin Math. Jour.*, t. v. (1850), pp. 159–165, and in particular the equation there obtained p. 161 is the equation $\psi(0, v + 1, n) = 0$.

779.

[NOTE ON MR JEFFERY'S PAPER "ON CERTAIN QUARTIC CURVES WHICH HAVE A CUSP AT INFINITY, WHEREAT THE LINE AT INFINITY IS A TANGENT."]

[From the *Proceedings of the London Mathematical Society*, vol. XIV. (1883), p. 85.]

The assumed form $\kappa\phi^2 = u_4$, or, as this is afterwards written,

$$2y^3y = ax^2 + 2bxy + cy^2 + 2ex + 2dy + \lambda,$$

is, I think, introduced without a proper explanation. Say, the form is $x^2y = z^2(* | x, y, z)^2$, it ought to be shown how for a cuspidal quartic we arrive at this form; *viz.* taking the cusp to be at the point $(x = 0, z = 0)$, $z = 0$ for the tangent at the cusp, and $x = 0$ an arbitrary line through the cusp; then the line $z = 0$ besides intersects the curve in a single point, and, if $y = 0$ is taken as the tangent at that point, the equation of the curve must, it can be seen, be of the form

$$(ax^2 + exz)y = z^2(a, b, c, f, g, h | x, y, z)^2.$$

The conic $(a, b, c, f, g, h | x, y, z)^2$ touches the quartic at each of the two intersections of the quartic with the arbitrary line $x = 0$; and we cannot, so long as the line remains arbitrary, find a conic which shall osculate the quartic at the two points in question; but, for the particular line $x + \frac{1}{2}dz = 0$, there exists such a conic, *viz.* writing x instead of $x + \frac{1}{2}dz$, the form is $x^2y = z^2(a', b', c', f', g', h' | x, y, z)^2$, and the new conic $(a', \dots | x, y, z)^2 = 0$ has the property in question. This is the adopted form, and it thus appears that in it the line $x = 0$ is a determinate line, *viz.* the line passing through the cusp and the two points of osculation of the osculating conic.

It thus appears that in the assumed form the lines $x = 0, y = 0, z = 0$ are determinate lines.

780.

[ADDITION TO MR HAMMOND'S PAPER "NOTE ON AN EXCEPTIONAL CASE IN WHICH THE FUNDAMENTAL POSTULATE OF PROFESSOR SYLVESTER'S THEORY OF TAMISAGE FAILS."]

[From the *Proceedings of the London Mathematical Society*, vol. XIV. (1883), pp. 88–91. Read Dec. 14, 1882.]

The extreme importance of Mr Hammond's result, as regards the entire subject of Covariants, leads me to reproduce his investigation in the notation of my Memoirs on Quantics, and with a somewhat different arrangement of the formulæ. For the binary seventhic

$$(a, b, c, d, e, f, g, h | x, y)^7,$$

the four composite seminvariants of the deg-weight 5.11 (sources of covariants of the deg-order 5.13) are

I.	II.	III.	IV.
1.7	4.6	2.10	3.3 Deg-order.
1.0	4.11	2.2	3.9 Deg-weight.
$a + 1$	$a^2eh + 1$ $a^2fg - 1$ $abdh - 4$ $abeg - 2$ $abf^2 + 6$ $ac^2h - 3$ $acd g - 2$ $acef - 6$ $ad^2f + 10$ $ad^2e - 5$ $ade^2 + \dots$ $b^2dg + 20$ $b^2ef + 57$ $bc^2g - 15$ $bcd f - 24$ $bce^2 - 30$ $bc^2e - 10$ $c^4f + 27$ $c^3de - 45$ $cd^3 + 20$	$ac + 1$ $b^2 - 1$	$ach + 2$ $adg - 7$ $ae f + 5$ $ac^2fh - 2$ $beg + 7$ $bdf + 22$ $be^2 - 25$ $cf - 27$ $cde + 45$ $d^3 - 20$

II. (continued)	III. (continued)	IV. (continued)
2.6	3.7	3.11 Deg-order.
2.4	3.1	3.5 Deg-weight.
$ae + 1$ $bd - 4$ $c^2 + 3$	$a^2h + 1$ $abg - 7$ $acf + 9$ $ade - 5$ $b^2f + 12$ $bce - 30$ $bd^2 + 20$	$ag + 1$ $bf - 6$ $ce + 16$ $d^2 - 10$
$a^2f + 1$ $abe - 5$ $ad - 2$ $ab^2d - 3$ $be^2 - 6$		

and it is here at once obvious that there exists a syzygy of the form $I = III - IV$; in fact, if in III. and IV. we write $a = 0$, then the values are each

$$= -2b(2bd - 3c^2)(6bf - 15ce + 10d^2)$$

hence $III - IV$ must divide by a , the quotient being a seminvariant of the deg-weight 4.11, which can only be a numerical multiple of the second factor of I., and is in fact this second factor, that is, we have the syzygy $I = III - IV$.

Working out the values of the four products, and joining to them the expression for the irreducible seminvariant of the same deg-weight 5.11 (Ω , Θ^* of my tables [774] for the binary sextic), we have the table:

5 . 10	5 . 11	I.	III.	IV.	II.
a^2dh	a^2efh	+1	+1		
a^2eg	a^2fg	−1	−4	+1	
a^2f^2	a^2bdh	−4	−7		
	a^2beg	−2	+3	−5	
a^2bch	a^2bf^2	+3	+9	−6	
a^2bdg	a^2c^2h	−2	−5	+2	
a^2bef	a^2cdg	−1	+28	+15	+2
a^2c^2g	a^2cef	+10	+12	−10	−7
a^2cdf	a^2d^2f	+3			+5
a^2ce^2	a^2de^2	−5	−21		−4
a^2d^2e	ab^2ch	−2	−36		
ab^3h	ab^2cg	+20	−30	+8	+7
ab^2cg	ab^2df	+1	+67	−45	−6
ab^2df	ab^2e^2	−15	+40		
ab^2e^2	abc^2g	−24	+27	−6	+7
abc^2f	abc^2f	−30	−15	−12	+22
$abcde$	$abcdf$	+27	−48		
abd^3	$abce^2$	−14	−45		−25
ac^3e	abd^2e	+20	+36	+50	−27
ac^2d^2	ac^3f	+11	+120	+30	+46
a^2c^2g	ac^2de	+1	−80	−20	−20
ab^2cf	acd^3	+9	−90		−2
ab^2de	a^2b^2h	−14	+60	−48	−7
ab^2c^2e	b^3cg	+6		−22	+26
ab^2cd^2	b^3df	+8		+86	+27
abc^3d	b^3e^2	−9		+120	−45
	b^2c^2f	−6		−80	+20
	b^2cde	+16		−90	
	b^2d^3	−8		+60	
	bc^3e	−3			
	bc^2d^2	+2			
	c^4d				

I have prefixed to the table the literal terms of the deg-weight 5 . 10; for the deg-weights 5 . 11 and 5 . 10, the numbers of terms are = 30 and 26 respectively; and it is the difference of these $30 - 26 = 4$, which gives the number of asyzygetic seminvariants of the deg-weight 5 . 11.

781.

ON THE AUTOMORPHIC TRANSFORMATION OF THE BINARY CUBIC FUNCTION.

[From the *Proceedings of the London Mathematical Society*, vol. XIV. (1883), pp. 103–108. Read Jan. 11, 1883.]

I CONSIDER the cubic equation $(a, b, c, d \mid x, 1)^3 = 0$. It is shown (Serret, *Cours d'Algèbre supérieure*, 4th ed., Paris, 1879, t. II. pp. 466–471) how, given one root of the equation, the other two roots can be each of them expressed rationally in terms of this root and of the square root of the discriminant; viz. making the proper changes of notation, and writing

$$\begin{aligned}
 A, B, C &= ac - b^2, \quad ad - bc, \quad bd - c^2, \quad \lambda = \sqrt{-1}, \\
 \Omega &= B^2 - 4AC, \quad = a^2d^2 - 4ac^3 + 6abcd - 4b^3d - 3b^2c^2, \\
 \alpha &= \frac{-\lambda\sqrt{\Omega} + B}{2\lambda\sqrt{\Omega}}, \quad \beta = \frac{+2C}{2\lambda\sqrt{\Omega}}, \quad \gamma = \frac{-2A}{2\lambda\sqrt{\Omega}}, \quad \delta = \frac{-\lambda\sqrt{\Omega} - B}{2\lambda\sqrt{\Omega}},
 \end{aligned}$$

(values which give $\alpha + \delta = -1$, $\alpha\delta - \beta\gamma = -1$, and therefore also

$$\alpha^2 + \alpha\delta + \delta^2 + \beta\gamma = 0,$$

which is the condition in order that the function $\phi x = \frac{\alpha x + \beta}{\gamma x + \delta}$ may be periodic of the third order, $\phi^3 x = x$), then, u being a root of the equation, say $(a, b, c, d \mid u, 1)^3 = 0$, the other two roots are

$$\phi u = \frac{\alpha u + \beta}{\gamma u + \delta},$$

and

$$\phi^2 u = \phi^{-1} u = \frac{(\alpha^2 + \beta\gamma)u + \beta(\alpha + \delta)}{\gamma(\alpha + \delta)u + \delta^2 + \beta\gamma} = \frac{-\delta u - \beta}{-\gamma u + \alpha},$$

where observe that, by the change of $\sqrt{\Omega}$ into $-\sqrt{\Omega}$, $\alpha, \beta, \gamma, \delta$ become $\delta, -\beta, -\gamma, \alpha$; so that the last-mentioned value $\phi^{-1}u$ is, in fact, the value obtained from ϕu by the mere change of sign of the radical.

52–2

It is to be observed that, if we have between two roots u, v of the equation

$$(a, b, c, d \mid x, 1)^3 = 0,$$

a relation $v = \frac{\alpha u + \beta}{\gamma u + \delta}$, where $\alpha, \beta, \gamma, \delta$ have given values, this implies in the first place a relation between a, b, c, d (and the given values of $\alpha, \beta, \gamma, \delta$), and it implies more-over that u , and consequently also v , are not any roots whatever, but two determinate roots of the equation; *viz.* u, v will be each of them expressible rationally in terms of a, b, c, d and $\alpha, \beta, \gamma, \delta$. And if, in order that (a, b, c, d) may remain arbitrary, we consider $\alpha, \beta, \gamma, \delta$ as given quantities satisfying the relation which exists between these quantities and (a, b, c, d) , then in general we still have u, v determinate roots of the cubic equation. But in the foregoing solution u is any root whatever of the cubic equation.

To examine how this is, starting from the equations

$$(a, b, c, d \mid u, 1)^3 = 0, \quad (a, b, c, d \mid v, 1)^3 = 0, \quad v = \frac{\alpha u + \beta}{\gamma u + \delta},$$

we have

$$a(u^3 - v^3) + 3b(u^2 - v^2) + 3c(u - v) = 0,$$

and therefore

$$a(u^2 + uv + v^2) + 3b(u + v) + 3c = 0,$$

that is,

$$av^2 + (au + 3b)v + au^2 + 3bu + 3c = 0;$$

or, writing herein for v its value,

$$a(\alpha u + \beta)^2 + (\alpha u + \beta)(\gamma u + \delta)(au + 3b) + (\gamma u + \delta)^2 (au^2 + 3bu + 3c) = 0;$$

that is,

$$\begin{aligned} & a(\alpha u + \beta)^2 + \alpha\gamma (au^2 + 3bu^2) + \gamma^2 (au^4 + 3bu^3 + 3cu^2) \\ & + (\alpha\delta + \beta\gamma)(au^2 + 3bu) + 2\gamma\delta (au^3 + 3bu^2 + 3cu) \\ & + \beta\delta (au + 3b) + \delta^2 (au^2 + 3bu + 3c) = 0; \end{aligned}$$

or, reducing by the equation $au^3 + 3bu^2 + 3cu + d = 0$, this is

$$\begin{aligned} & a(\alpha u + \beta)^2 + \alpha\gamma (-3cu - d) + \gamma^2 (-du) \\ & + (\alpha\delta + \beta\gamma)(au^2 + 3bu) + \beta\gamma\delta (-d) \\ & + \beta\delta (au + 3b) + \delta^2 (au^2 + 3bu + 3c) = 0, \end{aligned}$$

and, collecting the terms, this is

$$\begin{aligned} & u^2 a (\alpha^2 + \alpha\delta + \delta^2 + \beta\gamma) \\ & + u \{a(2\alpha\beta + \beta\delta) + 3b(\alpha\delta + \delta^2 + \beta\gamma) - 3c\alpha\gamma - d\gamma^2\} \\ & + a\beta^2 + 3b\beta\delta + 3c\delta^2 + d(-\alpha\gamma - 2\gamma\delta) = 0. \end{aligned}$$

We can, from this equation, and the equation $au^3 + 3bu^2 + 3cu + d = 0$, eliminate u , thus obtaining a relation between $a, b, c, d, \alpha, \beta, \gamma, \delta$; and, this relation being satisfied, the two equations then determine u rationally in terms of these quantities.

We may without loss of generality assume $\alpha\delta - \beta\gamma = 1$; and, this being so, if we then further assume $\alpha + \delta = -1$, then we have

$$\alpha^2 + \alpha\delta + \delta^2 + \beta\gamma = 0,$$

which is, as above appearing, the condition for $\phi^3 x = 0$. The foregoing equation in u thus becomes

$$\begin{aligned} & u \{a\beta^2(\alpha - 1) - 3b\alpha^2 - 3c\alpha\gamma - d\gamma^2\} \\ & + \{a\beta^2 + 3b\beta\delta + 3c\delta^2 - d\gamma(\delta + 1)\} = 0; \end{aligned}$$

a linear equation giving (in a simplified form) the like results to those given by the quadric equation; *viz.* substituting in the cubic equation the value of u given by the linear equation, we have a relation between $a, b, c, d, \alpha, \beta, \gamma, \delta$; and, this relation being satisfied, u has the determinate value given by the linear equation.

The only way in which u can cease to have this determinate value, and so be capable of being any root whatever of the cubic equation, is when the linear equation becomes $0 = 0$; *viz.* if

$$\begin{aligned} & a\beta^2(\alpha - 1) - 3b\alpha^2 - 3c\alpha\gamma - d\gamma^2 = 0, \\ & a\beta^2 + 3b\beta\delta + 3c\delta^2 - d\gamma(\delta + 1) = 0, \end{aligned}$$

equations which are, in fact, satisfied by the foregoing values of $\alpha, \beta, \gamma, \delta$, as may be verified without difficulty.

It is to be remarked that if, instead of the root u and the equation $v = \frac{\alpha u + \beta}{\gamma u + \delta}$, we consider the root v and the equation $u = \frac{-\delta v - \beta}{-\gamma v + \alpha}$, then, instead of $\alpha, \beta, \gamma, \delta$, we have $\delta, -\beta, -\gamma, \alpha$, and the corresponding equations are

$$\begin{aligned} & a\beta^2(\delta - 1) - 3b\delta^2 + 3c\gamma\delta + d\gamma^2 = 0, \\ & a\beta^2 + 3b\alpha\beta + 3c\alpha^2 + d\gamma(\alpha - 1) = 0, \end{aligned}$$

equations which are also satisfied by the foregoing values of $\alpha, \beta, \gamma, \delta$. And the four equations, together with $\alpha\delta - \beta\gamma = 1$ and $\alpha + \delta = -1$, are more than sufficient to determine the foregoing values of $\alpha, \beta, \gamma, \delta$.

But we further verify without difficulty that the foregoing values of $\alpha, \beta, \gamma, \delta$ give identically

$$(a, b, c, d \mid \alpha x + \beta y, \gamma x + \delta y)^3 = -(a, b, c, d \mid x, y)^3,$$

or the formulæ lead to an automorphic transformation of the binary cubic $(a, b, c, d \mid x, y)^3$. And conversely, starting from the notion of the automorphic transformation of the binary cubic, we ought to be able to obtain the foregoing formulæ.

For greater convenience, I write the equation of transformation in the form

$$(a, b, c, d \mid \alpha x + \beta y, \gamma x + \delta y)^3 = -\theta(a, b, c, d \mid x, y)^3;$$

THE COLLECTED MATHEMATICAL
PAPERS

Vol. XI — Pages 451–500

Arthur Cayley

[Continued from page 450.]

regarded as a singly infinite system of points, of lines, or of planes; secondly, the surface, which may be regarded as a doubly infinite system of points or of planes, and also as a special triply infinite system of lines (viz. the tangent-lines of the surface are a special complex): as distinct particular cases of the former figure, we have the plane curve and the cone; and as a particular case of the latter figure, the ruled surface or singly infinite system of lines; we have besides the congruence, or doubly infinite system of lines, and the complex, or triply infinite system of lines. But, even if in solid geometry we attend only to the curve and the surface, there are crowds of theories which have scarcely any analogues in plane geometry. The relation of a curve to the various surfaces which can be drawn through it, or of a surface to the various curves that can be drawn upon it, is different in kind from that which in plane geometry most nearly corresponds to it, the relation of a system of points to the curves through them, or of a curve to the points upon it. In particular, there is nothing in plane geometry corresponding to the theory of the curves of curvature of a surface. To the single theorem of plane geometry, a right line is the shortest distance between two points, there correspond in solid geometry two extensive and difficult theories—that of the geodesic lines upon a given surface, and that of the surface of minimum area for any given boundary. Again, in solid geometry we have the interesting and difficult question of the representation of a curve by means of equations; it is not every curve, but only a curve which is the complete intersection of two surfaces, which can be properly represented by two equations $(x, y, z, w)^m = 0$, $(x, y, z, w)^n = 0$, in quadriplanar coordinates; and in regard to this question, which may also be regarded as that of the classification of curves in space, we have quite recently three elaborate memoirs by Nöther, Halphen, and Valentiner respectively.

In n -dimensional geometry, only isolated questions have been considered. The field is simply too wide; the comparison with each other of the two cases of plane geometry and solid geometry is enough to show how the complexity and difficulty of the theory would increase with each successive dimension.

In Transcendental Analysis, or the Theory of Functions, we

have all that has been done in the present century with regard to the general theory of the function of an imaginary variable by Gauss, Cauchy, Puiseux, Briot, Bouquet, Liouville, Riemann, Fuchs, Weierstrass, and others. The fundamental idea of the geometrical representation of an imaginary variable $x + iy$, by means of the point having x, y for its coordinates, belongs, as I mentioned, to Gauss; of this I have already spoken at some length. The notion has been applied to differential equations; in the modern point of view, the problem in regard to a given differential equation is, not so much to reduce the differential equation to quadratures, as to determine from it directly the course of the integrals for all positions of the point representing the independent variable: in particular, the differential equation of the second order leading to the hypergeometric series $F(\alpha, \beta, \gamma, x)$ has been treated in this manner, with the most interesting results; the function so determined for all values of the parameters (α, β, γ) is thus becoming a known function. I would here also refer to the new notion in this part of analysis introduced by Weierstrass—that of the one-valued integer function, as defined by an

[57–2]

infinite series of ascending powers, convergent for all finite values, real or imaginary, of the variable x or $1/(x - c)$, and so having the one essential singular point $x = \infty$ or $x = c$, as the case may be: the memoir is published in the Berlin *Abhandlungen*, 1876.

But it is not only general theory: I have to speak of the various special functions to which the theory has been applied, or say the various known functions.

For a long time the only known transcendental functions were the circular functions—sine, cosine, &c.; the logarithm, i.e. for analytical purposes the hyperbolic logarithm to the base e ; and, as implied therein, the exponential function e^x . More completely stated, the group comprises the direct circular functions $\sin$, $\cos$, &c.; the inverse circular functions $\sin^{-1}$ or $\arcsin$, &c.; the exponential function, $\exp$.; and the inverse exponential, or logarithmic, function, $\log$.

Passing over the very important Eulerian integral of the second kind or gamma-function, the theory of which has quite recently given rise to some very interesting developments—and omitting to mention at all various functions of minor importance—we come (1811–1829) to the very wide groups, the elliptic functions and the single theta-functions. I give the interval of date so as to include Legendre's two systematic works, the *Exercices de Calcul Intégral* (1811–1816) and the *Théorie des Fonctions Elliptiques* (1825–1828); also Jacobi's *Fundamenta nova theoriae Functionum Ellipticarum* (1829), calling to mind that many of Jacobi's results were obtained simultaneously by Abel. I remark that Legendre started from the consideration of the integrals depending on a radical $\sqrt{X}$, the square root of a rational and integral quartic function of a variable x ; for this he substituted a radical $\Delta\phi$, $= \sqrt{(1 - k^2 \sin^2 \phi)}$, and he arrived at his three kinds of elliptic integrals $F\phi$, $E\phi$, $\Pi\phi$, depending on the argument or amplitude ϕ , the modulus k , and also the last of them on a parameter n ; the F function is properly an inverse function, and in place of it Abel and Jacobi each of them introduced the direct functions corresponding to the circular functions sine and cosine, Abel's functions called by him ϕ , f , F , and Jacobi's functions $\sin am$, $\cos am$, Δam , or as they are also written sn , cn , dn . Jacobi, moreover, in the development of his theory of transformation

obtained a multitude of formulae containing q , a transcendental function of the modulus defined by the equation $q = e^{-\pi K'/K}$, and he was also led by it to consider the two new functions H , Θ , which (taken each separately with two different arguments) are in fact the four functions called elsewhere by him Θ_1 , Θ_2 , Θ_3 , Θ_4 ; these are the so-called theta-functions, or, when the distinction is necessary, the single theta-functions. Finally, Jacobi using the transformation $\sin \phi = \operatorname{sn} u$, expressed Legendre's integrals of the second and third kinds as integrals depending on the new variable u , denoting them by means of the letters Z , Π , and connecting them with his own functions H and Θ ; and the elliptic functions sn , cn , dn are expressed with these, or say with Θ_1 , Θ_2 , Θ_3 , Θ_4 , as fractions having a common denominator.

It may be convenient to mention that Hermite in 1858, introducing into the theory in place of q the new variable ω connected with it by the equation $q = e^{i\pi\omega}$,

was led to consider three functions $\varphi\omega$, $\psi\omega$, $\chi\omega$, depending on and slightly differing from the three differences of the functions Θ for the three values $\frac{K}{3}$, $\frac{2K}{3}$, $K - \frac{K}{3}$ of the argument; these functions $\varphi\omega$, $\psi\omega$, $\chi\omega$ are connected with the modulus k by very simple relations (they are in fact, except as to constant factors, the fourth roots of k , k' , and kk'); and they are functions which play an important part in the new theory of equations.

The hyperelliptic integrals which succeed to the elliptic ones present themselves in the Abelian theory, but were probably first considered by Jacobi: he showed that the existing theory of elliptic functions led to the consideration of the integrals

$$\int \frac{dx}{\sqrt{X}}, \quad \int \frac{x dx}{\sqrt{X}},$$

where X is now a rational and integral sextic function of x ; these are, in the simplest case, the so-called hyperelliptic integrals, and the corresponding inverse functions are the hyperelliptic functions; only here, presenting themselves as inverse functions of integrals depending on the same radical $\sqrt{X}$, we have, not single but pairs of functions of two variables; and the case is generally that of p functions of p variables.

The general theory of the Abelian functions was first considered by Riemann in his memoir "Theorie der Abel'schen Functionen," *Crelle*, t. LIV. (1857); and we have also the great memoir by Clebsch and Gordan, *Theorie der Abelschen Functionen* (Leipzig, 1866). The theta-functions for any value whatever of p have been considered by Riemann, and also by Weierstrass in a memoir published in the *Berlin Monatsbericht*, 1869; but the theory has been chiefly developed for the next succeeding case $p = 2$, that of the double theta-functions, in connexion with hyperelliptic integrals, by various writers including Göpel, Rosenhain, Borchardt, Weber, Königsberger, Brioschi, Cayley, and others.

I would mention also the linear differential equations with rational algebraical coefficients, especially the equation of the second order treated by Fuchs and others; and as bearing thereon the theory of the so-called automorphic functions, treated of by Klein and Poincaré.

I pass now to the Theory of Numbers. The first leading question is that of the prime numbers, the determination of which is a very interesting one, although it does not seem at present to give rise to any general theory of importance. The next leading question is that of the residues of powers in regard to a prime modulus; this question is included in the still wider question of the theory of forms, viz. given a homogeneous quadratic function $ax^2 + 2bxy + cy^2$, of two variables x, y (called a form), and a prime modulus p , the question is to determine the values which this can take to the modulus p .

We have also the theory of complex numbers $a + bi$ ($i = \sqrt{-1}$); and Gauss considered the corresponding theory of the complex numbers $a + b\rho$ where ρ is an imaginary cube root of unity ($\rho^2 + \rho + 1 = 0$); these are the theories used in his demonstrations of the laws of quadratic and cubic reciprocity.

But the great extension is by Kummer in his memoir "Theorie der idealen Primfactoren der complexen Zahlen," *Berl. Abh.*, 1856. We have first the prime number p and its p th roots of unity, $1, \eta, \eta^2, \dots, \eta^{p-1}$, satisfying the equation $\eta^p - 1 = 0$, or breaking this up into the equation $\eta - 1 = 0$, with the single root $\eta = 1$, and the equation $1 + \eta + \eta^2 + \dots + \eta^{p-1} = 0$, with the $p - 1$ roots $\eta, \eta^2, \dots, \eta^{p-1}$; then if η be a root of this last equation, the complex numbers $a + a_1\eta + a_2\eta^2 + \dots + a_{p-1}\eta^{p-1}$ (with $a, a_1, \dots, a_{p-1}$ positive or negative ordinary integers, including zero) are the numbers considered in his theory.

But more generally Kummer considered the so-called "periods" of the p th roots of unity, viz. the prime number p being supposed of the form $p = \theta f + 1$, dividing the $p - 1$ roots $\eta, \eta^2, \dots, \eta^{p-1}$ into θ sets each containing f roots, each such set has for its sum a so-called period $\eta_1, \eta_2, \dots, \eta_\theta$, and these periods are by themselves the roots of an equation of the order θ ; and if $\eta_1, \eta_2, \dots, \eta_\theta$ are taken to be a sum of roots, of the equation $x^n - 1 = 0$, then the complex numbers considered are the numbers of the form $a_1\eta_1 + a_2\eta_2 + \dots + a_\theta\eta_\theta$ ($a_1, a_2, \dots, a_\theta$ positive or negative ordinary integers, including zero): f may be $= 1$, and the theory for the periods thus includes that for the single roots.

We have thus a new and very general theory, including within itself that of the complex numbers $a + bi$ and $a + b\rho$. But a new phenomenon presents itself; for these special forms the properties in regard to prime numbers corresponded precisely with those for real numbers; a non-prime number was in one way only a product of prime factors; the power of a prime number has only factors which are lower powers of the same prime number: for instance, if p be a prime number, then, excluding the obvious decomposition $p \cdot p^2$, we cannot have $p^3 =$ a product of two factors A, B . In the general case this is not so, but the exception first presents itself for the number 23; in the theory of the numbers composed with the 23rd roots of unity, we have prime numbers p , such that $p^3 = AB$. To restore the theorem, it is necessary to establish the notion of ideal numbers; a prime number p is by definition not the product of two actual numbers, but in the example just referred to the number p is the product of two ideal numbers having for

their cubes the two actual numbers A , B , respectively, and we thus have $p^3 = AB$. It is, I think, in this way that we most easily get some notion of the meaning of an ideal number, but the mode of treatment (in Kummer's great memoir, "Ueber die Zerlegung der aus Wurzeln der Einheit gebildeten complexen Zahlen in ihre Primfactoren," *Crelle*, t. XXXV. 1847) is a much more refined one; an ideal number, without ever being isolated, is made to manifest itself in the properties of the prime number of which it is a factor, and without reference to the

theorem afterwards arrived at, that there is always some power of the ideal number which is an actual number. In the still later developments of the Theory of Numbers by Dedekind, the units, or incommensurables, are the roots of any irreducible equation having for its coefficients ordinary integer numbers, and with the coefficient unity for the highest power of x . The question arises, What is the analogue of a whole number? Thus, for the very simple case of the equation $x^2 + 3 = 0$, we have as a whole number the apparently fractional form $\frac{1}{2}(1 + i\sqrt{3})$ which is the imaginary cube root of unity, the ρ of Eisenstein's theory. We have, moreover, the (as far as appears) wholly distinct complex theory of the numbers composed with the congruence-imaginaries of Galois: viz. these are imaginary numbers assumed to satisfy a congruence which is not satisfied by any real number; for instance, the congruence $x^2 \equiv -2 \pmod{5}$ has no real root, but we assume an imaginary root i , the other root is then $= -i$, and we then consider the system of complex numbers $a + bi \pmod{5}$, viz. we have thus the 5^2 numbers obtained by giving to each of the numbers a, b , the values $0, 1, 2, 3, 4$, successively. And so in general, the consideration of an irreducible congruence $F(x) \equiv 0 \pmod{p}$ of the order n , to any prime modulus p , gives rise to an imaginary congruence root i , and to complex numbers of the form $a + bi + ci^2 + \dots + fi^{n-1}$, where $a, b, c, \dots, \&c.$, are ordinary integers each $= 0, 1, 2, \dots, p - 1$.

As regards the theory of forms, we have in the ordinary theory, in addition to the binary and ternary quadratic forms, which have been very thoroughly studied, the quaternary and higher quadratic forms (to these last belong, as very particular cases, the theories of the representation of a number as a sum of four, five or more squares), and also binary cubic and quartic forms, and ternary cubic forms, in regard to all of which something has been done; the binary quadratic forms have been studied in the theory of the complex numbers $a + bi$.

A seemingly isolated question in the Theory of Numbers, the demonstration of Fermat's theorem of the impossibility for any exponent λ greater than 3, of the equation $x^\lambda + y^\lambda = z^\lambda$, has given rise to investigations of very great interest and difficulty.

Outside of ordinary mathematics, we have some theories which

must be referred to: algebraical, geometrical, logical. It is, as in many other cases, difficult to draw the line; we do in ordinary mathematics use symbols not denoting quantities, which we nevertheless combine in the way of addition and multiplication, $a + b$, and ab , and which may be such as not to obey the commutative law $ab = ba$: in particular, this is or may be so in regard to symbols of operation; and it could hardly be said that any development whatever of the theory of such symbols of operation did not belong to ordinary algebra. But I do separate from ordinary mathematics the system of multiple algebra or linear associative algebra, developed in the valuable memoir by the late Benjamin Peirce, *Linear Associative Algebra* (1870, reprinted 1881 in the *American Journal of Mathematics*, vol. IV., with notes and addenda by his son, C. S. Peirce); we here consider symbols A , B , &c. which are linear functions of a determinate number of letters or units with i , j , k , l , &c., coefficients which are ordinary analytical magnitudes, real or imaginary, viz. the coefficients are in general of the form $x + iy$, where

[C. XI. 58]

i is the before-mentioned imaginary or $\sqrt{-1}$ of ordinary analysis. The letters $i, j, \&c.$, are such that every binary combination $i^2, ij, ji, \&c.$, (the ij being in general not $= ji$), is equal to a linear function of the letters, but under the restriction of satisfying the associative law: viz. for each combination of three letters $ij \cdot k$ is $= i \cdot jk$, so that there is a determinate and unique product of three or more letters; or, what is the same thing, the laws of combination of the units i, j, k , are defined by a multiplication table giving the values of $i^2, ij, ji, \&c.$; the original units may be replaced by linear functions of these units, so as to give rise, for the units finally adopted, to a multiplication table of the most simple form; and it is very remarkable, how frequently in these simplified forms we have nilpotent or idempotent symbols ($i^2 = 0$, or $i^2 = i$, as the case may be), and symbols i, j , such that $ij = ji = 0$; and consequently how simple are the forms of the multiplication tables which define the several systems respectively.

I have spoken of this multiple algebra before referring to various geometrical theories of earlier date, because I consider it as the general analytical basis, and the true basis, of these theories. I do not realise to myself directly the notions of the addition or multiplication of two lines, areas, rotations, forces, or other geometrical, kinematical, or mechanical entities; and I would formulate a general theory as follows: consider any such entity as determined by the proper number of parameters a, b, c (for instance, in the case of a finite line given in magnitude and position, these might be the length, the coordinates of one end, and the direction-cosines of the line considered as drawn from this end); and represent it by or connect it with the linear function $ai + bj + ck + \&c.$, formed with these parameters as coefficients, and with a given set of units, $i, j, k, \&c.$ Conversely, any such linear function represents an entity of the kind in question. Two given entities are represented by two linear functions; the sum of these is a like linear function representing an entity of the same kind, which may be regarded as the sum of the two entities; and the product of them (taken in a determined order, when the order is material) is an entity of the same kind, which may be regarded as the product (in the same order) of the two entities. We thus establish by definition the notion of the sum of the two

entities, and that of the product (in a determinate order, when the order is material) of the two entities. The value of the theory in regard to any kind of entity would of course depend on the choice of a system of units, $i, j, k, \dots$, with such laws of combination as would give a geometrical or kinematical or mechanical significance to the notions of the sum and product as thus defined.

Among the geometrical theories referred to, we have a theory (that of Argand, Warren, and Peacock) of imaginaries in plane geometry; Sir W. R. Hamilton's very valuable and important theory of Quaternions; the theories developed in Grassmann's *Ausdehnungslehre*, 1841 and 1862; Clifford's theory of Biquaternions; and recent extensions of Grassmann's theory to non-Euclidian space, by Mr Homersham Cox. These different theories have of course been developed, not in anywise from the point of view in which I have been considering them, but from the points of view of their several authors respectively.

The literal symbols $x, y, \&c.$, used in Boole's *Laws of Thought* (1854) to represent things as subjects of our conceptions, are symbols obeying the laws of algebraic com-

bination (the distributive, commutative, and associative laws) but which are such that for any one of them, say x , we have $x - x^2 = 0$, this equation not implying (as in ordinary algebra it would do) either $x = 0$ or else $x = 1$. In the latter part of the work relating to the Theory of Probabilities, there is a difficulty in making out the precise meaning of the symbols; and the remarkable theory there developed has, it seems to me, passed out of notice, without having been properly discussed. A paper by the same author, "Of Propositions numerically definite" (*Camb. Phil. Trans.* 1869), is also on the border-land of logic and mathematics. It would be out of place to consider other systems of mathematical logic, but I will just mention that Mr C. S. Peirce in his "Algebra of Logic," *American Math. Journal*, vol. III., establishes a notation for relative terms, and that these present themselves in connexion with the systems of units of the linear associative algebra.

Connected with logic, but primarily mathematical and of the highest importance, we have Schubert's *Abzählende Geometrie* (1878). The general question is, How many curves or other figures are there which satisfy given conditions? for example, How many conics are there which touch each of five given conics? The class of questions in regard to the conic was first considered by Chasles, and we have his beautiful theory of the characteristics μ , ν , of the conics which satisfy four given conditions; questions relating to cubics and quartics were afterwards considered by Maillard and Zeuthen; and in the work just referred to the theory has become a very wide one. The noticeable point is that the symbols used by Schubert are in the first instance, not numbers, but mere logical symbols: for example, a letter g denotes the condition that a line shall cut a given line; g^2 that it shall cut each of two given lines; and so in other cases; and these logical symbols are combined together by algebraical laws: they first acquire a numerical signification when the number of conditions becomes equal to the number of parameters upon which the figure in question depends.

In all that I have last said in regard to theories outside of ordinary mathematics, I have been still speaking on the text of the vast extent of modern mathematics. In conclusion I would say that mathematics have steadily advanced from the time of the Greek

geometers. Nothing is lost or wasted; the achievements of Euclid, Archimedes, and Apollonius are as admirable now as they were in their own days. Descartes' method of coordinates is a possession for ever. But mathematics have never been cultivated more zealously and diligently, or with greater success, than in this century—in the last half of it, or at the present time: the advances made have been enormous, the actual field is boundless, the future full of hope. In regard to pure mathematics we may most confidently say:

Yet I doubt not through the ages one increasing purpose
runs,

And the thoughts of men are widened with the process
of the suns.

[58–2]

785.

CURVE.

[From the *Encyclopædia Britannica*, Ninth Edition, vol. VI. (1877), pp. 716—728.]

THIS subject is treated here from an historical point of view, for the purpose of showing how the different leading ideas in the theory were successively arrived at and developed.

A curve is a line, or continuous singly infinite system of points. We consider in the first instance, and chiefly, a plane curve described according to a law. Such a curve may be regarded geometrically as actually described, or kinematically as in course of description by the motion of a point; in the former point of view, it is the locus of all the points which satisfy a given condition; in the latter, it is the locus of a point moving subject to a given condition. Thus the most simple and earliest known curve, the circle, is the locus of all the points at a given distance from a fixed centre, or else the locus of a point moving so as to be always at a given distance from a fixed centre. (The straight line and the point are not for the moment regarded as curves.)

Next to the circle we have the conic sections, the invention of them attributed to Plato (who lived 430 to 347 B.C.); the original definition of them as the sections of a cone was by the Greek geometers who studied them soon replaced by a proper definition *in plano* like that for the circle, viz. a conic section (or as we now say a “conic”) is the locus of a point such that its distance from a given point, the focus, is in a given ratio to its (perpendicular) distance from a given line, the directrix; or it is the locus of a point which moves so as always to satisfy the foregoing condition. Similarly any other property might be used as a definition; an ellipse is the locus of a point such that the sum of its distances from two fixed points (the foci) is constant, &c., &c.

The Greek geometers invented other curves; in particular, the “conchoid,” which is the locus of a point such that its distance from a given line, measured along the

line drawn through it to a fixed point, is constant; and the “cissoid” which is the locus of a point such that its distance from a fixed point is always equal to the intercept (on the line through the fixed point) between a circle passing through the fixed point and the tangent to the circle at the point opposite to the fixed point. Obviously the number of such geometrical or kinematical definitions is infinite. In a machine of any kind, each point describes a curve; a simple but important instance is the “three-bar curve,” or locus of a point in or rigidly connected with a bar pivotted on to two other bars which rotate about fixed centres respectively. Every curve thus arbitrarily defined has its own properties; and there was not any principle of classification.

The principle of classification first presented itself in the *Géométrie* of Descartes (1637). The idea was to represent any curve whatever by means of a relation between the coordinates (x, y) of a point of the curve, or say to represent the curve by means of its equation.

Descartes takes two lines xx', yy' , called axes of coordinates, intersecting at a point O called the origin (the axes are usually at right angles to each other, and for the present they are considered as being so); and he determines the position of a point P by means of its distances OM (or NP) $= x$, and MP (or ON) $= y$, from these two axes respectively; where x is regarded as positive or negative according as it is in the sense Ox or Ox' from O ; and similarly y as positive or negative according as it is in the sense Oy or Oy' from O ; or, what is the same thing,

In the quadrant xy , or N.E., we have				x	y
				+	+
„	$x'y$	„ N.W.,		−	+
„	xy'	„ S.E.,		+	−
„	$x'y'$	„ S.W.,		−	−

Any relation whatever between (x, y) determines a curve, and conversely every curve whatever is determined by a relation between (x, y) .

Observe that the distinctive feature is in the exclusive use of such determination of a curve by means of its equation. The Greek geometers were perfectly familiar with the property of an ellipse which in the Cartesian notation is $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, the equation of the curve; but it was as one of a number of properties, and in no wise selected out of the others for the characteristic property of the curve¹.

We obtain from the equation the notion of an algebraical or geometrical as opposed to a transcendental curve, viz. an algebraical or geometrical curve is a curve having an equation $F(x, y) = 0$, where $F(x, y)$ is a rational and integral algebraical function of the coordinates (x, y) ; and in what follows we attend throughout (unless the contrary is stated) only to such curves. The equation is sometimes given, and may conveniently be used, in an irrational form, but we always imagine it reduced to the foregoing rational and integral form, and regard this as the equation of the curve. And we have hence the notion of a curve of a given order, viz. the order of the curve is equal to that of the term or terms of highest

¹There is no exercise more profitable for a student than that of tracing a curve from its equation, or say rather that of so tracing a considerable number of curves. And he should make the equations for himself. The equation should be in the first instance a purely numerical one, where y is given or can be found as an explicit function of x ; here, by giving different numerical values to x , the corresponding values of y may be found; and a sufficient number of points being thus determined, the curve is traced by drawing a continuous line through these points. The next step should be to consider an equation involving literal coefficients; thus, after such curves as $y = x^2$, $y = x(x-1)(x-2)$, $y = (x-1)\sqrt{x-2}$, &c., he should proceed to trace such curves as $y = (x-a)(x-b)(x-c)$, $y = (x-a)\sqrt{x-b}$, &c., and endeavour to ascertain for what different relations of equality or inequality between the coefficients the curve will assume essentially or notably distinct forms. The purely numerical equations will present instances of nodes, cusps, inflexions, double tangents, asymptotes, &c.—specialities which he should be familiar with before he has to consider their general theory. And he may then consider an equation such that neither coordinate can be expressed as an explicit function of the other of them (practically, an equation such as $x^3 + y^3 - 3xy = 0$, which requires the solution of a cubic equation, belongs to this class); the problem of tracing the curve here frequently requires special methods, and it may easily be such as to require and serve as an exercise for the powers of an advanced algebraist.

order in the coordinates (x, y) conjointly in the equation of the curve; for instance, $xy - 1 = 0$ is a curve of the second order.

It is to be noticed here that the axes of coordinates may be any two lines at right angles to each other whatever; and that the equation of a curve will be different according to the selection of the axes of coordinates; but the order is independent of the axes, and has a determinate value for any given curve.

We hence divide curves according to their order, viz. a curve is of the first order, second order, third order, &c., according as it is represented by an equation of the first order, $ax + by + c = 0$, or say $(\S x, y, 1) = 0$; or by an equation of the second order; $ax^2 + 2hxy + by^2 + 2fy + 2gx + c = 0$, say $(\S x, y, 1)^2 = 0$; or by an equation of the third order, &c.; or, what is the same thing, according as the equation is linear, quadric, cubic, &c.

A curve of the first order is a right line; and conversely every right line is a curve of the first order.

A curve of the second order is a conic, or as it is also called a quadric; and conversely every conic, or quadric, is a curve of the second order.

A curve of the third order is called a cubic; one of the fourth order a quartic; and so on.

A curve of the order m has for its equation $(\S x, y, 1)^m = 0$; and when the coefficients of the function are arbitrary, the curve is said to be the general curve of the order m . The number of coefficients is $\frac{1}{2}(m+1)(m+2)$; but there is no loss of generality if the equation be divided by one coefficient so as to reduce the coefficient of the corresponding term to unity, hence the number of coefficients may be reckoned as $\frac{1}{2}(m+1)(m+2) - 1$, that is, $\frac{1}{2}m(m+3)$; and a curve of the order m may be made to satisfy this number of conditions; for example, to pass through $\frac{1}{2}m(m+3)$ points.

It is to be remarked that an equation may break up; thus a quadric equation may be $(ax + by + c)(a'x + b'y + c') = 0$, breaking up into the two equations $ax + by + c = 0$, $a'x + b'y + c' = 0$, viz. the original equation is satisfied if either of these is satisfied. Each of these last equations represents a curve of the first order, or right line; and the original equation represents this pair of lines, viz. the pair of lines is considered as a quadric curve. But it is an improper quadric curve; and in speaking of curves of the second or any other given order, we frequently imply that the curve is a proper curve represented by an equation which does not break up.

The intersections of two curves are obtained by combining their equations; viz. the elimination from the two equations of y (or x) gives for x (or y) an equation of a certain order, say the resultant equation; and then to each value of x (or y) satisfying this equation

there corresponds in general a single value of y (or x), and consequently a single point of intersection; the number of intersections is thus equal to the order of the resultant equation in x (or y).

Supposing that the two curves are of the orders m, n , respectively, then the order of the resultant equation is in general and at most $= mn$; in particular, if the curve of the order n is an arbitrary line ($n = 1$), then the order of the resultant equation is $= m$; and the curve of the order m meets therefore the line in m points. But the resultant equation may have all or any of its roots imaginary, and it is thus not always that there are m real intersections.

The notion of imaginary intersections, thus presenting itself, through algebra, in geometry, must be accepted in geometry—and it in fact plays an all-important part in modern geometry. As in algebra we say that an equation of the m th order has m roots, viz. we state this generally without in the first instance, or it may be without ever, distinguishing whether these are real or imaginary; so in geometry we say that a curve of the m th order is met by an arbitrary line in m points, or rather we thus, through algebra, obtain the proper geometrical definition of a curve of the m th order, as a curve which is met by an arbitrary line in m points (that is, of course, in m , and not more than m , points).

The theorem of the m intersections has been stated in regard to an arbitrary line; in fact, for particular lines the resultant equation may be or appear to be of

an order less than m ; for instance, taking $m = 2$, if the hyperbola $xy - 1 = 0$ be cut by the line $y = \beta$, the resultant equation in x is $\beta x - 1 = 0$, and there is apparently only the intersection $(x = 1/\beta, y = \beta)$; but the theorem is, in fact, true for every line whatever: a curve of the order m meets every line whatever in precisely m points. We have, in the case just referred to, to take account of a point at infinity on the line $y = \beta$; the two intersections are the point $(x = 1/\beta, y = \beta)$, and the point at infinity on the line $y = \beta$.

It is moreover to be noticed that the points at infinity may be all or any of them imaginary, and that the points of intersection, whether finite or at infinity, real or imaginary, may coincide two or more of them together, and have to be counted accordingly; to support the theorem in its universality, it is necessary to take account of these various circumstances.

The foregoing notion of a point at infinity is a very important one in modern geometry; and we have also to consider the paradoxical statement that in plane geometry, or say as regards the plane, infinity is a right line. This admits of an easy illustration in solid geometry. If with a given centre of projection, by drawing from it lines to every point of a given line, we project the given line on a given plane, the projection is a line, i.e., this projection is the intersection of the given plane with the plane through the centre and the given line. Say the projection is always a line, then if the figure is such that the two planes are parallel, the projection is the intersection of the given plane by a parallel plane, or it is the system of points at infinity on the given plane, that is, these points at infinity are regarded as situate on a given line, the line infinity of the given plane².

Reverting to the purely plane theory, infinity is a line, related like any other right line to the curve, and thus intersecting it in m points, real or imaginary, distinct or coincident.

Descartes in the *Géométrie* defined and considered the remarkable curves called after him ovals of Descartes, or simply Cartesians, which will be again referred to. The next important work, founded

²More generally, in solid geometry infinity is a plane,—its intersection with any given plane being the right line which is the infinity of this given plane.

on the *Géométrie*, was Sir Isaac Newton's *Enumeratio linearum tertii ordinis* (1706), establishing a classification of cubic curves founded chiefly on the nature of their infinite branches, which was in some details completed by Stirling, Murdoch, and Cramer; the work contains also the remarkable theorem (to be again referred to), that there are five kinds of cubic curves giving by their projections every cubic curve whatever.

Various properties of curves in general, and of cubic curves, are established in Maclaurin's memoir, "De linearum geometricarum proprietatibus generalibus Tractatus" (posthumous, say 1746, published in the 6th edition of his *Algebra*). We have in it a particular kind of correspondence of two points on a cubic curve, viz. two points correspond to each other when the tangents at the two points again meet the cubic in the same point.

The *Géométrie Descriptive* by Monge was written in the year 1794 or 1795 (7th edition, Paris, 1847), and in it we find stated, in plano with regard to the circle, and in three dimensions with regard to a surface of the second order, the fundamental theorem of reciprocal polars, viz. “Given a surface of the second order and a circumscribed conic surface which touches it . . . then if the conic surface moves so that its summit is always in the same plane, the plane of the curve of contact passes always through the same point.” The theorem is here referred to partly on account of its bearing on the theory of imaginaries in geometry. It is, in Brianchon’s memoir “Sur les surfaces du second degré” (*Jour. Polyt.*, t. VI., 1806), shown how for any given position of the summit the plane of contact is determined, or reciprocally; say the plane XY is determined when the point P is given, or reciprocally; and it is noticed that when P is situate in the interior of the surface the plane XY does not cut the surface; that is, we have a real plane XY intersecting the surface in the imaginary curve of contact of the imaginary circumscribed cone having for its summit a given real point P inside the surface.

Stating the theorem in regard to a conic, we have a real point P (called the pole) and a real line XY (called the polar), the line joining the two (real or imaginary) points of contact of the (real or imaginary) tangents drawn from the point to the conic; and the theorem is that when the point describes a line the line passes through a point, this line and point being polar and pole to each other. The term “pole” was first used by Servois, and “polar” by Gergonne (*Gerg.*, t. I. and III., 1810–13); and from the theorem we have the method of reciprocal polars for the transformation of geometrical theorems, used already by Brianchon (in the memoir above referred to) for the demonstration of the theorem called by his name, and in a similar manner by various writers in the earlier volumes of Gergonne. We are here concerned with the method less in itself than as leading to the general notion of duality. And, bearing in a somewhat similar manner also on the theory of imaginaries in geometry (but the notion presents itself in a more explicit form), there is the memoir by Gaultier, on the graphical construction of circles and spheres (*Jour. Polyt.*, t. IX., 1813). The well-known theorem as to radical axes may be stated as follows. Consider two

circles partially drawn so that it does not appear whether the circles, if completed, would or would not intersect in real points, say two arcs of circles; then we can, by means of a third circle drawn so as to intersect in two real points each of the two arcs, determine a right line, which, if the complete circles intersect in two real points, passes through the points, and which is on this account regarded as a line passing through two (real or imaginary) points of intersection of the two circles. The construction in fact is, join the two points in which the third circle meets the first arc, and join also the two points in which the third circle meets the second arc, and from the point of intersection of the two joining lines, let fall a perpendicular on the line joining the centre of the two circles; this perpendicular (considered as an indefinite line) is what Gaultier terms the “radical axis of the two circles”; it is a line determined by a real construction and itself always real; and by what precedes it is the line joining two (real or imaginary, as the case may be) intersections of the given circles.

[*C. XI.*

59]

The intersections which lie on the radical axis are two out of the four intersections of the two circles. The question as to the remaining two intersections did not present itself to Gaultier, but it is answered in Poncelet's *Traité des propriétés projectives* (1822), where we find (p. 49) the statement, "deux cercles placés arbitrairement sur un plan. . . ont idéalement deux points imaginaires communs à l'infini"; that is, a circle qua curve of the second order is met by the line infinity in two points; but, more than this, they are the same two points for any circle whatever. The points in question have since been called (it is believed first by Dr Salmon) the circular points at infinity, or they may be called the circular points; these are also frequently spoken of as the points *I*, *J*; and we have thus the circle characterized as a conic which passes through the two circular points at infinity; the number of conditions thus imposed upon the conic is = 2, and there remain three arbitrary constants, which is the right number for the circle. Poncelet throughout his work makes continual use of the foregoing theories of imaginaries and infinity, and also of the before-mentioned theory of reciprocal polars.

Poncelet's two memoirs "Sur les centres des moyennes harmoniques," and "Sur la théorie générale des polaires réciproques," although presented to the Paris Academy in 1824 were only published (*Crelle*, t. III. and IV., 1828, 1829), subsequent to the memoir by Gergonne, "Considérations philosophiques sur les élémens de la science de l'étendue" (*Gerg.*, t. XVI., 1825-26). In this memoir by Gergonne, the theory of duality is very clearly and explicitly stated; for instance, we find "dans la géométrie plane, à chaque théorème il en répond nécessairement un autre qui s'en déduit en échangeant simplement entre eux les deux mots *points* et *droites*; tandis que dans la géométrie de l'espace ce sont les mots *points* et *plans* qu'il faut échanger entre eux pour passer d'un théorème à son corrélatif"; and the plan is introduced of printing correlative theorems, opposite to each other, in two columns. There was a reclamation as to priority by Poncelet in the *Bulletin Universel* reprinted with remarks by Gergonne (*Gerg.*, t. XIX., 1827), and followed by a short paper by Gergonne, "Rectifications de quelques théorèmes, &c.," which is important as first introducing the word *class*. We find in it explicitly the two correlative definitions: "a

plane curve is said to be of the m th degree (order) when it has with a line m real or ideal intersections,” and “a plane curve is said to be of the m th class when from any point of its plane there can be drawn to it m real or ideal tangents.”

It may be remarked that in Poncelet’s memoir on reciprocal polars, above referred to, we have the theorem that the number of tangents from a point to a curve of the order m , or say the class of the curve, is in general and at most $= m(m - 1)$, and that he mentions that this number is subject to reduction when the curve has double points or cusps.

The theorem of duality as regards plane figures may be thus stated: two figures may correspond to each other in such manner that to each point and line in either figure there corresponds in the other figure a line and point respectively. It is to be understood that the theorem extends to all points or lines, drawn or not drawn; thus if in the first figure there are any number of points on a line drawn or not drawn, the corresponding lines in the second figure, produced if necessary, must meet

in a point. And we thus see how the theorem extends to curves, their points and tangents: if there is in the first figure a curve of the order m , any line meets it in m points; and hence from the corresponding point in the second figure there must be to the corresponding curve m tangents; that is, the corresponding curve must be of the class m .

Trilinear coordinates (to be again referred to) were first used by Bobillier in the memoir, "Essai sur un nouveau mode de recherche des propriétés de l'étendue" (*Gerg.*, t. XVIII., 1827–28). It is convenient to use these rather than Cartesian coordinates. We represent a curve of the order m by an equation $(\{x, y, z\})^m = 0$, the function on the left-hand being a homogeneous rational and integral function of the order m of the three coordinates (x, y, z) ; clearly the number of constants is the same as for the equation $(\{x, y, 1\})^m$ in Cartesian coordinates.

The theory of duality is considered and developed, but chiefly in regard to its metrical applications, by Chasles in the "Mémoire de géométrie sur deux principes généraux de la science, la dualité et l'homographie," which forms a sequel to the "Aperçu historique sur l'origine et le développement des méthodes en géométrie" (*Mém. de Brux.*, t. XI., 1837).

We now come to Plücker; his "six equations" were given in a short memoir in *Crelle* (1842) preceding his great work, the *Theorie der algebraischen Curven* (1844).

Plücker first gave a scientific dual definition of a curve, viz. "A curve is a locus generated by a point, and enveloped by a line—the point moving continuously along the line, while the line rotates continuously about the point"; the point is a point (*ineunt*) of the curve, the line is a tangent of the curve.

And, assuming the above theory of geometrical imaginaries, a curve such that m of its points are situate in an arbitrary line is said to be of the order m ; a curve such that n of its tangents pass through an arbitrary point is said to be of the class n ; as already appearing, this notion of the order and the class of a curve is, however, due to Gergonne. Thus the line is a curve of the order 1 and the class 0; and corresponding dually thereto, we have the point as a curve of the order 0 and the class 1.

Plücker moreover imagined a system of line-coordinates (tangential coordinates). The Cartesian coordinates (x, y) and trilinear coordinates (x, y, z) are point-coordinates for determining the position of a point; the new coordinates, say (ξ, η, ζ) , are line-coordinates for determining the position of a line. It is possible, and (not so much for any application thereof as in order to more fully establish the analogy between the two kinds of coordinates) important, to give independent quantitative definitions of the two kinds of coordinates; but we may also derive the notion of line-coordinates from that of point-coordinates; viz. taking $\xi x + \eta y + \zeta z = 0$ to be the equation of a line, we say that (ξ, η, ζ) are the line-coordinates of this line. A linear relation $a\xi + b\eta + c\zeta = 0$ between these coordinates determines a point, viz. the point whose point-coordinates are (a, b, c) ; in fact, the equation in question $a\xi + b\eta + c\zeta = 0$ expresses that the equation $\xi x + \eta y + \zeta z = 0$, where (x, y, z) are current point-coordinates, is satisfied on writing therein $x, y, z = a, b, c$; or that the line in question passes through

[59-2]

the point (a, b, c) . Thus (ξ, η, ζ) are the line-coordinates of any line whatever; but when these, instead of being absolutely arbitrary, are subject to the restriction $a\xi + b\eta + c\zeta = 0$, this obliges the line to pass through a point (a, b, c) ; and the last-mentioned equation $a\xi + b\eta + c\zeta = 0$ is considered as the line-equation of this point.

A line has only a point-equation, and a point has only a line-equation; but any other curve has a point-equation and also a line-equation; the point-equation $(\dagger x, y, z)^m = 0$ is the relation which is satisfied by the point-coordinates (x, y, z) of each point of the curve; and similarly the line-equation $(\dagger \xi, \eta, \zeta)^n = 0$ is the relation which is satisfied by the line-coordinates (ξ, η, ζ) of each line (tangent) of the curve.

There is in analytical geometry little occasion for any explicit use of line-coordinates; but the theory is very important; it serves to show that, in demonstrating by point-coordinates any purely descriptive theorem whatever, we demonstrate the correlative theorem; that is, we do not demonstrate the one theorem, and then (as by the method of reciprocal polars) deduce from it the other, but we do at one and the same time demonstrate the two theorems; our (x, y, z) instead of meaning point-coordinates may mean line-coordinates, and the demonstration is then in every step of it a demonstration of the correlative theorem.

The above dual generation explains the nature of the singularities of a plane curve. The ordinary singularities, arranged according to a cross division, are

	Proper.	Improper.
Point-singularities {	1. The stationary point, cusp, or spinode;	2. The double point, or node
Line-singularities {	3. The stationary tangent, or inflexion;	4. The double tangent;

arising as follows:

1. The cusp: the point as it travels along the line may come to rest, and then reverse the direction of its motion.

3. The stationary tangent: the line may in the course of its rotation come to rest, and then reverse the direction of its rotation.

2. The node: the point may in the course of its motion come to coincide with a former position of the point, the two positions of the line not in general coinciding.

4. The double tangent: the line may in the course of its motion come to coincide with a former position of the line, the two positions of the point not in general coinciding.

It may be remarked that we cannot with a real point and line obtain the node with two imaginary tangents (conjugate or isolated point, or acnode), nor again the real double tangent with two imaginary points of contact; but this is of little consequence, since in the general theory the distinction between real and imaginary is not attended to.

The singularities (1) and (3) have been termed proper singularities, and (2) and (4) improper; in each of the first-mentioned cases there is a real singularity, or

peculiarity in the motion; in the other two cases there is not; in (2) there is not when the point is first at the node, or when it is secondly at the node, any peculiarity in the motion; the singularity consists in the point coming twice into the same position; and so in (4) the singularity is in the line coming twice into the same position. Moreover (1) and (2) are, the former a proper singularity, and the latter an improper singularity, as regards the motion of the point; and similarly (3) and (4) are, the former a proper singularity, and the latter an improper singularity, as regards the motion of the line.

But as regards the representation of a curve by an equation, the case is very different.

First, if the equation be in point-coordinates, (3) and (4) are in a sense not singularities at all. The curve $(\S x, y, z)^m = 0$, or general curve of the order m , has double tangents and inflexions; (2) presents itself as a singularity, for the equations $\partial_x(\S x, y, z)^m = 0$, $\partial_y(\S x, y, z)^m = 0$, $\partial_z(\S x, y, z)^m = 0$, implying $(\S x, y, z)^m = 0$, are not in general satisfied by any values (a, b, c) whatever of (x, y, z) , but if such values exist, then the point (a, b, c) is a node or double point; and (1) presents itself as a further singularity or sub-case of (2), a cusp being a double point for which the two tangents become coincident.

In line-coordinates all is reversed: (1) and (2) are not singularities; (3) presents itself as a sub-case of (4).

The theory of compound singularities will be referred to further on.

In regard to the ordinary singularities, we have

m , the order,

n , the class,

δ , the number of double points,

κ , the number of cusps,

τ , the number of double tangents,

ι , the number of inflexions;

and this being so, Plücker's "six equations" are

$$(1) \quad n = m(m - 1) - 2\delta - 3\kappa,$$

$$(2) \quad \iota = 3m(m - 2) - 6\delta - 8\kappa,$$

$$(3) \quad \tau = \frac{1}{2}m(m - 2)(m^2 - 9) - (m^2 - m - 6)(2\delta + 3\kappa) \\ + 2\delta(\delta - 1) + 6\delta\kappa + \frac{9}{2}\kappa(\kappa - 1),$$

$$(4) \quad m = n(n - 1) - 2\tau - 3\iota,$$

$$(5) \quad \kappa = 3n(n - 2) - 6\tau - 8\iota,$$

$$(6) \quad \delta = \frac{1}{2}n(n - 2)(n^2 - 9) - (n^2 - n - 6)(2\tau + 3\iota) \\ + 2\tau(\tau - 1) + 6\tau\iota + \frac{9}{2}\iota(\iota - 1).$$

It is easy to derive the further forms

$$(7) \quad \iota - \kappa = 3(n - m),$$

$$(8) \quad 2(\tau - \delta) = (n - m)(n + m - 9),$$

$$(9) \quad \frac{1}{2}m(m + 3) - \delta - 2\kappa = \frac{1}{2}n(n + 3) - \tau - 2\iota,$$

$$(10) \quad \frac{1}{2}(m - 1)(m - 2) - \delta - \kappa = \frac{1}{2}(n - 1)(n - 2) - \tau - \iota,$$

$$(11, 12) \quad m^2 - 2\delta - 3\kappa = n^2 - 2\tau - 3\iota, = m + n,$$

the whole system being equivalent to three equations only: and it may be added that, using a to denote the equal quantities $3m + \iota$ and $3n + \kappa$, everything may be expressed in terms of m, n, a . We have

$$\kappa = a - 3n,$$

$$\iota = a - 3m,$$

$$2\delta = m^2 - m + 8n - 3a,$$

$$2\tau = n^2 - n + 8m - 3a.$$

It is implied in Plücker's theorem that, $m, n, \delta, \kappa, \tau, \iota$ signifying as above in regard to any curve, then in regard to the reciprocal curve $n, m, \tau, \iota, \delta, \kappa$, will have the same significations, viz. for the reciprocal curve these letters denote respectively the order, class, number of nodes, cusps, double tangents, and inflexions.

The expression $\frac{1}{2}m(m + 3) - \delta - 2\kappa$ is that of the number of the disposable constants in a curve of the order m with δ nodes and κ cusps (in fact that there shall be a node is 1 condition, a cusp 2 conditions): and the equation (9) thus expresses that the curve and its reciprocal contain each of them the same number of disposable constants.

For a curve of the order m , the expression $\frac{1}{2}(m - 1)(m - 2) - \delta - \kappa$ is termed the "deficiency" (as to this more hereafter); the equation (10) expresses therefore that the curve and its reciprocal have each of them the same deficiency.

The relations $m^2 - 2\delta - 3\kappa = n^2 - 2\tau - 3\iota, = m + n$, present themselves in the theory of envelopes, as will appear further on.

With regard to the demonstration of Plücker's equations it is to be remarked that we are not able to write down the equation

in point-coordinates of a curve of the order m , having the given numbers δ and κ of nodes and cusps. We can only use the general equation $(x, y, z)^m = 0$, say for shortness $u = 0$, of a curve of the m th order, which equation, so long as the coefficients remain arbitrary, represents a curve without nodes or cusps. Seeking then, for this curve, the values n, ι, τ of the class, number of inflexions, and number of double tangents—first, as regards the class, this is equal to the number of tangents which can be drawn to the curve from an arbitrary point, or what is the same thing, it is equal to the number of the points of contact of these tangents. The points of contact are found as the intersections of the curve $u = 0$ by a curve depending on the position of the arbitrary point, and called the “first polar” of this point; the order of the first polar is $= m - 1$, and

the number of intersections is thus $= m(m - 1)$. But it can be shown, analytically or geometrically, that if the given curve has a node, the first polar passes through this node, which therefore counts as two intersections; and that if the curve has a cusp, the first polar passes through the cusp, touching the curve there, and hence the cusp counts as three intersections. But, as is evident, the node or cusp is not a point of contact of a proper tangent from the arbitrary point; we have, therefore, for a node a diminution 2, and for a cusp a diminution 3, in the number of the intersections; and thus, for a curve with δ nodes and κ cusps, there is a diminution $2\delta + 3\kappa$, and the value of n is $n = m(m - 1) - 2\delta - 3\kappa$.

Secondly, as to the inflexions, the process is a similar one; it can be shown that the inflexions are the intersections of the curve by a derivative curve called (after Hesse, who first considered it) the Hessian, defined geometrically as the locus of a point such that its conic polar in regard to the curve breaks up into a pair of lines, and which has an equation $H = 0$, where H is the determinant formed with the second differential coefficients of u in regard to the variables (x, y, z) ; $H = 0$ is thus a curve of the order $3(m - 2)$, and the number of inflexions is $= 3m(m - 2)$. But if the given curve has a node, then not only the Hessian passes through the node, but it has there a node the two branches at which touch respectively the two branches of the curve, and the node thus counts as six intersections; so if the curve has a cusp, then the Hessian not only passes through the cusp, but it has there a cusp through which it again passes, that is, there is a cuspidal branch touching the cuspidal branch of the curve, and besides a simple branch passing through the cusp, and hence the cusp counts as eight intersections. The node or cusp is not an inflexion, and we have thus for a node a diminution 6, and for a cusp a diminution 8, in the number of the intersections; hence for a curve with δ nodes and κ cusps, the diminution is $= 6\delta + 8\kappa$, and the number of inflexions is $\iota = 3m(m - 2) - 6\delta - 8\kappa$.

Thirdly, for the double tangents; the points of contact of these are obtained as the intersections of the curve by a curve $\Pi = 0$, which has not as yet been geometrically defined, but which is found analytically to be of the order $(m - 2)(m^2 - 9)$; the number of intersections is thus $= m(m - 2)(m^2 - 9)$; but if the given curve

has a node then there is a diminution $4(m^2 - m - 6)$, and if it has a cusp then there is a diminution $= 6(m^2 - m - 6)$, where, however, it is to be noticed that the factor $(m^2 - m - 6)$ is in the case of a curve having only a node or only a cusp the number of the tangents which can be drawn from the node or cusp to the curve, and is used as denoting the number of these tangents, and ceases to be the correct expression if the number of nodes and cusps is greater than unity. Hence, in the case of a curve which has δ nodes and κ cusps, the apparent diminution $2(m^2 - m - 6)(2\delta + 3\kappa)$ is too great, and it has in fact to be diminished by $2\{2\delta(\delta - 1) + 6\delta\kappa + \frac{9}{2}\kappa(\kappa - 1)\}$, or the half thereof is 4 for each pair of nodes, 6 for each combination of a node and cusp, and 9 for each pair of cusps. We have thus finally an expression for 2τ , $= m(m - 2)(m^2 - 9) - \&c.$; or dividing the whole by 2, we have the expression for τ given by the third of Plücker's equations.

It is obvious that we cannot by consideration of the equation $u = 0$ in point-coordinates obtain the remaining three of Plücker's equations; they might be obtained

in a precisely analogous manner by means of the equation $v = 0$ in line-coordinates, but they follow at once from the principle of duality, viz. they are obtained by the mere interchange of m, δ, κ with n, τ, ι respectively.

To complete Plücker's theory it is necessary to take account of compound singularities; it might be possible, but it is at any rate difficult, to effect this by considering the curve as in course of description by the point moving along the rotating line; and it seems easier to consider the compound singularity as arising from the variation of an actually described curve with ordinary singularities. The most simple case is when three double points come into coincidence, thereby giving rise to a triple point; and a somewhat more complicated one is when we have a cusp of the second kind, or node-cusp arising from the coincidence of a node, a cusp, an inflexion, and a double tangent, as shown in the annexed figure, which represents the singularities as on the point of coalescing. The general conclusion (see Cayley, *Quart. Math. Jour.* t. VII., 1866, [374], "On the higher singularities of a plane curve") is that every singularity whatever may be considered as compounded of ordinary singularities, say we have a singularity $= \delta'$ nodes, κ' cusps, τ' double tangents, and ι' inflexions. So that, in fact, Plücker's equations properly understood apply to a curve with any singularities whatever.

By means of Plücker's equations we may form a table

m	n	δ	κ	τ	ι
0	1	—	—	—	—
1	0	—	—	—	—
2	2	—	—	—	—
3	6	—	—	—	9
3	4	1	—	—	3
3	3	—	1	—	1
4	12	—	—	28	24
4	10	1	—	16	18
4	9	—	1	10	16
4	8	2	—	8	12
4	7	1	1	4	10
4	6	—	2	1	8
4	6	3	—	4	6
4	5	2	1	2	4
4	4	1	2	1	2
4	3	—	3	1	—

The table is arranged according to the value of m ; and we have $m = 0$, $n = 1$, the point; $m = 1$, $n = 0$, the line; $m = 2$, $n = 2$, the conic; of $m = 3$, the cubic, there are three cases, the class being 6, 4, or 3, according as the curve is without singularities, or as it has 1 node, or 1 cusp; and so of $m = 4$, the quartic, there are nine cases, where observe that in two of them the class is = 6, the reduction of class arising from two cusps or else from three nodes. The nine cases may be also grouped together into four, according as the number of nodes and cusps ($\delta + \kappa$) is = 0, 1, 2, or 3.

The cases may be divided into sub-cases, by the consideration of compound singularities; thus when $m = 4$, $n = 6$, $\delta = 3$, the three nodes may be all distinct, which is the general case, or two of them may unite together into the singularity called a tacnode, or all three may unite together into a triple point, or else into an oscnode.

We may further consider the inflexions and double tangents, as well in general as in regard to cubic and quartic curves.

The expression for the number of inflexions $3m(m - 2)$ for a curve of the order m was obtained analytically by Plücker, but the theory was first given in a complete form by Hesse in the two papers "Ueber die Elimination, u.s.w.," and "Ueber die Wendepuncte der Curven dritter Ordnung" (*Crelle*, t. XXVIII., 1844); in the latter of these the points of inflexion are obtained as the intersections of the curve $u = 0$ with the Hessian, or curve $\Delta = 0$, where Δ is the determinant formed with the second derived functions of u . We have in the Hessian the first instance of a covariant of

a ternary form. The whole theory of the inflexions of a cubic curve is discussed in a very interesting manner by means of the canonical form of the equation $x^3 + y^3 + z^3 + 6lxyz = 0$; and in particular a proof is given of Plücker's theorem that the nine points of inflexion of a cubic curve lie by threes in twelve lines.

It may be noticed that the nine inflexions of a cubic curve are three real, six imaginary; the three real inflexions lie in a line, as was known to Newton and Maclaurin. For an acnodal cubic the six imaginary inflexions disappear, and there remain three real inflexions lying in a line. For a crunodal cubic, the six inflexions which disappear are two of them real, the other four imaginary, and there remain two imaginary inflexions and one real inflexion. For a cuspidal cubic the six imaginary inflexions and two of the real inflexions disappear, and there remains one real inflexion.

A quartic curve has 24 inflexions; it was conjectured by Salmon, and has been verified recently by Zeuthen, that at most 8 of these are real.

The expression $\frac{1}{2}m(m-2)(m^2-9)$ for the number of double tangents of a curve of the order m was obtained by Plücker only as a consequence of his first, second, fourth, and fifth equations. An investigation by means of the curve $\Pi = 0$, which by its intersections with the given curve determines the points of contact of the double tangents, is indicated by Cayley, "Recherches sur l'élimination et la théorie des courbes," (*Crelle*, t. XXXIV., 1847), [53]; and in part carried out by Hesse in the memoir "Ueber Curven dritter Ordnung" (*Crelle*, t. XXXVI., 1848). A better process was indicated by Salmon in the "Note on the double tangents to plane curves," *Phil. Mag.* 1858; considering the $m-2$ points in which any tangent to the curve again meets the curve, he showed how to form the equation of a curve of the order $(m-2)$, giving by its intersection with the tangent the points in question; making the tangent touch this curve of the order $(m-2)$, it will be a double tangent of the original curve. See Cayley, "On the Double Tangents of a Plane Curve," (*Phil. Trans.* t. CXLVIII., 1859), [260], and Dersch (*Math. Ann.* t. VII., 1874). The solution is still in so far incomplete that we have no properties of the curve $\Pi = 0$, to distinguish one such curve from the several other curves which pass through the points

of contact of the double tangents.

A quartic curve has 28 double tangents, their points of contact determined as the intersections of the curve by a curve $\Pi = 0$ of the order 14, the equation of which in a very elegant form was first obtained by Hesse (1849). Investigations in regard to them are given by Plücker in the *Theorie der algebraischen Curven*, and in two memoirs by Hesse and Steiner (*Crelle*, t. XLV., 1855), in respect to the triads of double tangents which have their points of contact on a conic, and other like relations. It was assumed by Plücker that the number of real double tangents might be 28, 16, 8, 4, or 0, but Zeuthen has recently found that the last case does not exist.

The Hessian Δ has just been spoken of as a covariant of the form u ; the notion of invariants and covariants belongs rather to the form u than to the curve $u = 0$ represented by means of this form; and the theory may be very briefly referred

to. A curve $u = 0$ may have some invariative property, viz. a property independent of the particular axes of coordinates used in the representation of the curve by its equation; for instance, the curve may have a node, and in order to this, a relation, say $A = 0$, must exist between the coefficients of the equation; supposing the axes of coordinates altered, so that the equation becomes $u' = 0$, and writing $A' = 0$ for the relation between the new coefficients, then the relations $A = 0$, $A' = 0$, as two different expressions of the same geometrical property, must each of them imply the other; this can only be the case when A , A' are functions differing only by a constant factor, or say, when A is an invariant of u . If, however, the geometrical property requires two or more relations between the coefficients, say $A = 0$, $B = 0$, &c., then we must have between the new coefficients the like relations, $A' = 0$, $B' = 0$, &c., and the two systems of equations must each of them imply the other; when this is so, the system of equations, $A = 0$, $B = 0$, &c., is said to be invariative, but it does not follow that A , B , &c., are of necessity invariants of u . Similarly, if we have a curve $U = 0$ derived from the curve $u = 0$ in a manner independent of the particular axes of coordinates, then from the transformed equation $u' = 0$ deriving in like manner the curve $U' = 0$, the two equations $U = 0$, $U' = 0$ must each of them imply the other; and when this is so, U will be a covariant of u . The case is less frequent, but it may arise, that there are covariant systems $U = 0$, $V = 0$, &c., and $U' = 0$, $V' = 0$, &c., each implying the other, but where the functions U , V , &c., are not of necessity covariants of u .

The theory of the invariants and covariants of a ternary cubic function u has been studied in detail, and brought into connexion with the cubic curve $u = 0$; but the theory of the invariants and covariants for the next succeeding case, the ternary quartic function, is still very incomplete.

In further illustration of the Plückerian dual generation of a curve, we may consider the question of the envelope of a variable curve. The notion is very probably older, but it is at any rate to be found in Lagrange's *Théorie des fonctions analytiques* (1798); it is there remarked that the equation obtained by the elimination of the parameter a from an equation $f(x, y, a) = 0$ and the derived

equation in respect to a is a curve, the envelope of the series of curves represented by the equation $f(x, y, a) = 0$ in question. To develop the theory, consider the curve corresponding to any particular value of the parameter; this has with the consecutive curve (or curve belonging to the consecutive value of the parameter) a certain number of intersections, and of common tangents, which may be considered as the tangents at the intersections; and the so-called envelope is the curve which is at the same time generated by the points of intersection and enveloped by the common tangents; we have thus a dual generation. But the question needs to be further examined. Suppose that in general the variable curve is of the order m with δ nodes and κ cusps, and therefore of the class n with τ double tangents and ι inflexions, $m, n, \delta, \kappa, \tau, \iota$ being connected by the Plückerian equations; the number of nodes or cusps may be greater for particular values of the parameter, but this is a speciality which may be here disregarded. Considering the variable curve corresponding to a given value of the parameter, or say simply the variable curve, the consecutive curve has then also δ and κ nodes and cusps, consecutive to those of the variable curve; and it is easy to see that among the intersections of the two curves we have the nodes each counting twice, and the cusps each counting three times; the number of the remaining $= m^2 - 2\delta - 3\kappa$ intersections is

$m^2 - 2\delta - 3\kappa$. Similarly among the common tangents of the two curves we have the double tangents each counting twice, and the stationary tangents each counting three times, and the number of the remaining common tangents is $= n^2 - 2\tau - 3\iota$ ($= m^2 - 2\delta - 3\kappa$, inasmuch as each of these numbers is as was seen $= m + n$). At any one of the $m^2 - 2\delta - 3\kappa$ points the variable curve and the consecutive curve have tangents distinct from yet infinitesimally near to each other, and each of these two tangents is also infinitesimally near to one of the $n^2 - 2\tau - 3\iota$ common tangents of the two curves; whence, attending only to the variable curve, and considering the consecutive curve as coming into actual coincidence with it, the $n^2 - 2\tau - 3\iota$ common tangents are the tangents to the variable curve at the $m^2 - 2\delta - 3\kappa$ points respectively, and the envelope is at the same time generated by the $m^2 - 2\delta - 3\kappa$ points, and enveloped by the $n^2 - 2\tau - 3\iota$ tangents; we have thus a dual generation of the envelope, which only differs from Plücker's dual generation, in that in place of a single point and tangent we have the group of $m^2 - 2\delta - 3\kappa$ points and $n^2 - 2\tau - 3\iota$ tangents.

The parameter which determines the variable curve may be given as a point upon a given curve, or say as a parametric point; that is, to the different positions of the parametric point on the given curve correspond the different variable curves, and the nature of the envelope will thus depend on that of the given curve; we have thus the envelope as a derivative curve of the given curve. Many well-known derivative curves present themselves in this manner; thus the variable curve may be the normal (or line at right angles to the tangent) at any point of the given curve; the intersection of the consecutive normals is the centre of curvature, and we have the evolute

[60-2]

as at once the locus of the centre of curvature and the envelope of the normal. It may be added that the given curve is one of a series of curves, each cutting the several normals at right angles. Any one of these is a “parallel” of the given curve; and it can be obtained as the envelope of a circle of constant radius having its centre on the given curve.

We have in like manner, as derivatives of a given curve, the caustic, catacaustic, or diacaustic, as the case may be, and the secondary caustic, or curve cutting at right angles the reflected or refracted rays.

We have in much that precedes disregarded, or at least been indifferent to, reality; it is only thus that the conception of a curve of the m th order, as one which is met by every right line in m points, is arrived at; and the curve itself, and the line which cuts it, although both are tacitly assumed to be real, may perfectly well be imaginary. For real figures we have the general theorem that imaginary intersections, &c., present themselves in conjugate pairs: hence, in particular, that a curve of an even order is met by a line in an even number (which may be $= 0$) of points; a curve of an odd order in an odd number of points, hence in one point at least; it will be seen further on that the theorem may be generalized in a remarkable manner. Again, when there is in question only one pair of points or lines, these, if coincident, must be real; thus, a line meets a cubic curve in three points, one of them real, the other two real or imaginary; but if two of the intersections coincide they must be real, and we have a line cutting a cubic in one real point and touching it in another real point. It may be remarked that this is a limit separating the two cases where the intersections are all real, and where they are one real, two imaginary.

Considering always real curves, we obtain the notion of a branch; any portion capable of description by the continuous motion of a point is a branch; and a curve consists of one or more branches. Thus the curve of the first order or right line consists of one branch; but in curves of the second order, or conics, the ellipse and the parabola consist each of one branch, the hyperbola of two branches. A branch is either re-entrant, or it extends both ways to infinity, and in this case, we may regard it as consisting of two legs (*crura*,

Newton), each extending one way to infinity, but without any definite separation. The branch, whether re-entrant or infinite, may have a cusp or cusps, or it may cut itself or another branch, thus having or giving rise to crunodes; an acnode is a branch by itself,—it may be considered as an indefinitely small re-entrant branch. A branch may have inflexions and double tangents, or there may be double tangents which touch two distinct branches; there are also double tangents with imaginary points of contact, which are thus lines having no visible connexion with the curve. A re-entrant branch not cutting itself may be everywhere convex, and it is then properly said to be an oval; but the term oval may be used more generally for any re-entrant branch not cutting itself; and we may thus speak of a once indented, twice indented oval, &c., or even of a cuspidate oval. Other descriptive names for ovals and re-entrant branches cutting themselves may be used when required; thus, in the last-mentioned case a simple form is that of a figure of eight; such a form may break up into two ovals, or into a doubly indented oval or hour-glass. A form which presents itself is when two ovals, one inside the other,

unite, so as to give rise to a crunode—in default of a better name this may be called, after the curve of that name, a *limaçon*. Names may also be used for the different forms of infinite branches, but we have first to consider the distinction of hyperbolic and parabolic. The leg of an infinite branch may have at the extremity a tangent; this is an asymptote of the curve, and the leg is then hyperbolic; or the leg may tend to a fixed direction, but so that the tangent goes further and further off to infinity, and the leg is then parabolic; a branch may thus be hyperbolic or parabolic as to its two legs; or it may be hyperbolic as to one leg, and parabolic as to the other. The epithets hyperbolic and parabolic are of course derived from the conics hyperbola and parabola respectively. The nature of the two kinds of branches is best understood by considering them as projections, in the same way as we in effect consider the hyperbola and the parabola as projections of the ellipse. If a line Ω cut an arc aa' , so that the two segments ab , ba' lie on opposite sides of the line, then projecting the figure so that the line Ω goes off to infinity, the tangent at b is projected into the asymptote, and the arc ab is projected into a hyperbolic leg touching the asymptote at one extremity; the arc ba' will at the same time be projected into a hyperbolic leg touching the same asymptote at the other extremity (and on the opposite side), but so that the two hyperbolic legs may or may not belong to one and the same branch. And we thus see that the two hyperbolic legs belong to a simple intersection of the curve by the line infinity. Next, if the line Ω touch at b the arc aa' , so that the two portions ab' , ba lie on the same side of the line Ω , then projecting the figure as before, the tangent at b , that is, the line Ω itself, is projected to infinity; the arc ab is projected into a parabolic leg, and at the same time the arc ba' is projected into a parabolic leg, having at infinity the same direction as the other leg, but so that the two legs may or may not belong to the same branch. And we thus see that the two parabolic legs represent a contact of the line infinity with the curve,—the point of contact being of course the point at infinity determined by the common direction of the two legs. It will readily be understood how the like considerations apply to other cases,—for instance, if the line Ω is a tangent at an inflexion, passes through a crunode, or touches

one of the branches of a crunode, &c.; thus, if the line Ω passes through a crunode we have pairs of hyperbolic legs belonging to two parallel asymptotes. The foregoing considerations also show (what is very important) how different branches are connected together at infinity, and lead to the notion of a complete branch, or circuit.

The two legs of a hyperbolic branch may belong to different asymptotes, and in this case we have the forms which Newton calls inscribed, circumscribed, ambigene, &c.; or they may belong to the same asymptote, and in this case we have the serpentine form, where the branch cuts the asymptote, so as to touch it at its two extremities on opposite sides, or the conchoidal form, where it touches the asymptote on the same side. The two legs of a parabolic branch may converge to ultimate parallelism, as in the conic parabola, or diverge to ultimate parallelism, as in the semi-cubical parabola $y^2 = x^3$, and the branch is said to be convergent, or divergent, accordingly; or they may tend to parallelism in opposite senses, as in the cubical parabola $y = x^3$. As mentioned with regard to a branch generally, an infinite branch of any kind may have cusps, or, by cutting itself or another branch, may have or give rise to a crunode, &c.

We may now consider the various forms of cubic curves, as appearing by Newton's *Enumeratio*, and by the figures belonging thereto. The species are reckoned as 72, which are numbered accordingly 1 to 72; but to these should be added 10^a , 13^a , 22^a , and 22^b . It is not intended here to consider the division into species, nor even completely that into genera, but only to explain the principle of classification. It may be remarked generally that there are at most three infinite branches, and that there may besides be a re-entrant branch or oval.

The genera may be arranged as follows:

- 1, 2, 3, 4 redundant hyperbolas,
- 5, 6 defective hyperbolas,
- 7, 8 parabolic hyperbolas,
- 9 hyperbolisms of hyperbola,
- 10 " " ellipse,
- 11 " " parabola,
- 12 trident curve,
- 13 divergent parabolas,
- 14 cubic parabola

and, thus arranged, they correspond to the different relations of the line infinity to the curve. First, if the three intersections by the line infinity are all distinct, we have the hyperbolas; if the points are real, the redundant hyperbolas, with three hyperbolic branches; but if only one of them is real, the defective hyperbolas, with one hyperbolic branch. Secondly, if two of the intersections coincide, say if the line infinity meets the curve in a onefold point and a twofold point, both of them real, then there is always one asymptote: the line infinity may at the twofold point touch the curve, and we have the parabolic hyperbolas; or the twofold point may be a singular point, viz. a crunode giving the hyperbolisms of the hyperbola; an acnode, giving the hyperbolisms of the ellipse; or a cusp, giving the hyperbolisms of the parabola. As regards the so-called hyperbolisms, observe that (besides the single asymptote) we have in the case of those of the hyperbola two parallel asymptotes; in the case of those of the ellipse the two parallel asymptotes become imaginary, that is, they disappear, and in the case of those of the parabola they

become coincident, that is, there is here an ordinary asymptote, and a special asymptote answering to a cusp at infinity. Thirdly, the three intersections by the line infinity may be coincident and real; or say we have a threefold point: this may be an inflexion, a crunode, or a cusp, that is, the line infinity may be a tangent at an inflexion, and we have the divergent parabolas; a tangent at a crunode to one branch, and we have the trident curve; or lastly, a tangent at a cusp, and we have the cubical parabola.

It is to be remarked that the classification mixes together non-singular and singular curves, in fact, the five kinds presently referred to: thus the hyperbolas and the divergent parabolas include curves of every kind, the separation being made in the

species; the hyperbolisms of the hyperbola and ellipse, and the trident curve, are nodal; the hyperbolisms of the parabola, and the cubical parabola, are cuspidal. The divergent parabolas are of five species which respectively belong to and determine the five kinds of cubic curves; Newton gives (in two short paragraphs without any development) the remarkable theorem that the five divergent parabolas by their shadows generate and exhibit all the cubic curves.

The five divergent parabolas are curves each of them symmetrical with regard to an axis. There are two non-singular kinds, the one with, the other without, an oval, but each of them has an infinite (as Newton describes it) campaniform branch; this cuts the axis at right angles, being at first convex, but ultimately concave, towards the axis, the two legs continually tending to become at right angles to the axis. The oval may unite itself with the infinite branch, or it may dwindle into a point, and we have the crunodal and the acnodal forms respectively; or if simultaneously the oval dwindle into a point and unites itself to the infinite branch, we have the cuspidal form. Drawing a line to cut any one of these curves and projecting the line to infinity, it would not be difficult to show how the line should be drawn in order to obtain a curve of any given species. We have herein a better principle of classification; considering cubic curves, in the first instance, according to singularities, the curves are non-singular, nodal (viz. crunodal or acnodal), or cuspidal; and we see further that there are two kinds of non-singular curves, the complex and the simplex. There is thus a complete division into the five kinds, the complex, simplex, crunodal, acnodal, and cuspidal. Each singular kind presents itself as a limit separating two kinds of inferior singularity; the cuspidal separates the crunodal and the acnodal, and these last separate from each other the complex and the simplex.

The whole question is discussed very fully and ably by Möbius in the memoir "Ueber die Grundformen der Linien dritter Ordnung" (*Abh. der K. Sachs. Ges. zu Leipzig*, t. I., 1852; *Ges. Werke*, t. I.). The author considers not only plane curves, but also cones, or, what is almost the same thing, the spherical curves which are their sections by a concentric sphere. Stated in regard to the cone, we have there the fundamental theorem that there are two different

kinds of sheets: viz. the single sheet, not separated into two parts by the vertex (an instance is afforded by the plane considered as a cone of the first order generated by the motion of a line about a point), and the double or twin-pair sheet, separated into two parts by the vertex (as in the cone of the second order). And it then appears that there are two kinds of non-singular cubic cones, viz. the simplex, consisting of a single sheet, and the complex, consisting of a single sheet and a twin-pair sheet; and we thence obtain (as for cubic curves) the crunodal, the acnodal, and the cuspidal kinds of cubic cones. It may be mentioned that the single sheet is a sort of wavy form, having upon it three lines of inflexion, and which is met by any plane through the vertex in one or in three lines; the twin-pair sheet has no lines of inflexion, and resembles in its form a cone on an oval base.

In general a cone consists of one or more single or twin-pair sheets, and if we consider the section of the cone by a plane, the curve consists of one or more complete branches, or say circuits, each of them the section of one sheet of the cone;

thus, a cone of the second order is one twin-pair sheet, and any section of it is one circuit composed, it may be, of two branches. But although we thus arrive by projection at the notion of a circuit, it is not necessary to go out of the plane, and we may (with Zeuthen, using the shorter term *circuit* for his *complete branch*) define a circuit as any portion (of a curve) capable of description by the continuous motion of a point, it being understood that a passage through infinity is permitted. And we then say that a curve consists of one or more circuits; thus the right line, or curve of the first order, consists of one circuit; a curve of the second order consists of one circuit; a cubic curve consists of one circuit or else of two circuits.

A circuit is met by any right line always in an even number, or always in an odd number, of points, and it is said to be an even circuit or an odd circuit accordingly; the right line is an odd circuit, the conic an even circuit. And we have then the theorem, two odd circuits intersect in an odd number of points; an odd and an even circuit, or two even circuits, in an even number of points. An even circuit not cutting itself divides the plane into two parts, the one called the internal part, incapable of containing any odd circuit, the other called the external part, capable of containing an odd circuit.

We may now state in a more convenient form the fundamental distinction of the kinds of cubic curve. A non-singular cubic is simplex, consisting of one odd circuit, or it is complex, consisting of one odd circuit and one even circuit. It may be added that there are on the odd circuit three inflexions, but on the even circuit no inflexion; it hence also appears that from any point of the odd circuit there can be drawn to the odd circuit two tangents, and to the even circuit (if any) two tangents, but that from a point of the even circuit there cannot be drawn (either to the odd or the even circuit) any real tangent; consequently, in a simplex curve the number of tangents from any point is two; but in a complex curve the number is four, or none, four if the point is on the odd circuit, none if it is on the even circuit. It at once appears from inspection of the figure of a non-singular cubic curve, which is the odd and which the even circuit. The singular kinds arise as before; in the crunodal and the cuspidal kinds the whole curve is an odd circuit,

but in the acnodal kind the acnode must be regarded as an even circuit.

The analogous question of the classification of quartics (in particular non-singular quartics and nodal quartics) is considered in Zeuthen's memoir "Sur les différentes formes des courbes planes du quatrième ordre" (*Math. Ann.* t. VII., 1874). A non-singular quartic has only even circuits; it has at most four circuits external to each other, or two circuits one internal to the other, and in this last case the internal circuit has no double tangents or inflexions. A very remarkable theorem is established as to the double tangents of such a quartic: distinguishing as a double tangent of the first kind a real double tangent which either twice touches the same circuit, or else touches the curve in two imaginary points, the number of the double tangents of the first kind of a non-singular quartic is $= 4$; it follows that the quartic has at most 8 real inflexions. The forms of the non-singular quartics are very numerous, but it is not necessary to go further into the question.

[C. XI.

61]

We may consider in relation to a curve, not only the line infinity, but also the circular points at infinity; assuming the curve to be real, these present themselves always conjointly; thus a circle is a conic passing through the two circular points, and is thereby distinguished from other conics. Similarly a cubic through the two circular points is termed a circular cubic; a quartic through the two points is termed a circular quartic, and if it passes twice through each of them, that is, has each of them for a node, it is termed a bicircular quartic. Such a quartic is of course binodal ($m = 4$, $\delta = 2$, $\kappa = 0$); it has not in general, but it may have, a third node, or a cusp. Or again, we may have a quartic curve having a cusp at each of the circular points: such a curve is a “Cartesian,” it being a complete definition of the Cartesian to say that it is a bicuspidal quartic curve ($m = 4$, $\delta = 0$, $\kappa = 2$), having a cusp at each of the circular points. The circular cubic and the bicircular quartic, together with the Cartesian (being in one point of view a particular case thereof), are interesting curves which have been much studied, generally, and in reference to their focal properties.

The points called foci presented themselves in the theory of the conic, and were well known to the Greek geometers, but the general notion of a focus was first established by Plücker, in the memoir “Ueber solche Punkte die bei Curven einer höheren Ordnung den Brennpunkten der Kegelschnitte entsprechen,” (*Crelle*, t. X., 1833). We may from each of the circular points draw tangents to a given curve; the intersection of two such tangents (belonging of course to the two circular points respectively) is a focus. There will be from each circular point λ tangents (λ a number depending on the class of the curve and its relation to the line infinity and the circular points, $= 2$ for the general conic, 1 for the parabola, 2 for a circular cubic or a bicircular quartic, &c.); the λ tangents from the one circular point and those from the other circular point intersect in λ real foci (viz. each of these is the only real point on each of the tangents through it), and in $\lambda^2 - \lambda$ imaginary foci; each pair of real foci determines a pair of imaginary foci (the so-called antipoints of the two real foci), and the $\frac{1}{2}\lambda(\lambda - 1)$ pairs of real foci thus determine the $\lambda^2 - \lambda$ imaginary foci. There are in some cases points termed centres, or singular or multiple foci (the nomenclature is unsettled), which are

the intersections of improper tangents from the two circular points respectively; thus, in the circular cubic, the tangents to the curve at the two circular points respectively (or two imaginary asymptotes of the curve) meet in a centre.

The notions of distance and of lines at right angles are connected with the circular points; and almost every construction of a curve by means of lines of a determinate length, or at right angles to each other, and (as such) mechanical constructions by means of linkwork, give rise to curves passing the same definite number of times through the two circular points respectively, or say to circular curves, and in which the fixed centres of the construction present themselves as ordinary, or as singular, foci. Thus the general curve of three-bar motion (or locus of the vertex of a triangle, the other two vertices whereof move on fixed circles) is a tricircular sextic, having besides three nodes ($m = 6$, $\delta = 3 + 3 + 3, = 9$), and having the centres of the fixed circles each for a singular focus; there is a third singular focus, and we have thus the remarkable theorem (due to Mr S. Roberts) of the triple generation of the curve by means of the three several pairs of singular foci.

Again, the normal, qua line at right angles to the tangent, is connected with the circular points, and these accordingly present themselves in the before-mentioned theories of evolutes and parallel curves.

We have several recent theories which depend on the notion of correspondence: two points whether in the same plane or in different planes, or on the same curve or in different curves, may determine each other in such wise that to any given position of the first point there correspond α' positions of the second point, and to any given position of the second point α positions of the first point; the two points have then an (α, α') correspondence; and if α, α' are each = 1, then the two points have a (1, 1) or rational correspondence. Connecting with each theory the author's name, the theories in question are—Riemann, the rational transformation of a plane curve; Cremona, the rational transformation of a plane; and Chasles, correspondence of points on the same curve, and united points. The theory first referred to, with the resulting notion of *Geschlecht*, or deficiency, is more than the other two an essential part of the theory of curves, but they will all be considered.

Riemann's results are contained in the memoirs on "Theorie der Abel'schen Functionen," (*Crelle*, t. LIV., 1857); and we have next Clebsch, "Ueber die Singularitäten algebraischer Curven," (*Crelle*, t. LXV., 1865), and Cayley, "On the Transformation of Plane Curves," (*Proc. Lond. Math. Soc.* t. I., 1865, [384]). The fundamental notion of the rational transformation is as follows:

Taking u, X, Y, Z to be rational and integral functions (X, Y, Z all of the same order) of the coordinates (x, y, z) , and u', X', Y', Z' rational and integral functions (X', Y', Z' all of the same order) of the coordinates (x', y', z') , we transform a given curve $u = 0$, by the equations $x' : y' : z' = X : Y : Z$, thereby obtaining a transformed curve $u' = 0$, and a converse set of equations $x : y : z = X' : Y' : Z'$; viz. assuming that this is so, the point (x, y, z) on the curve $u = 0$ and the point (x', y', z') on the curve $u' = 0$ will be points having a (1, 1) correspondence. To show how this is, observe that to a given point (x, y, z) on the curve $u = 0$ there corresponds a single point (x', y', z') determined by the equations $x' : y' : z' = X : Y : Z$; from these equations and the equation $u = 0$ eliminating x, y, z

we obtain the equation $u' = 0$ of the transformed curve. To a given point (x', y', z') not on the curve $u' = 0$ there corresponds, not a single point, but the system of points (x, y, z) given by the equations $x' : y' : z' = X : Y : Z$, viz. regarding x', y', z' as constants (and to fix the ideas, assuming that the curves $X = 0$, $Y = 0$, $Z = 0$ have no common intersections), these are the points of intersection of the curves $X : Y : Z = x' : y' : z'$, but no one of these points is situate on the curve $u = 0$. If, however, the point (x', y', z') is situate on the curve $u' = 0$, then one point of the system of points in question is situate on the curve $u = 0$, that is, to a given point of the curve $u' = 0$ there corresponds a single point of the curve $u = 0$; and hence also this point must be given by a system of equations such as $x : y : z = X' : Y' : Z'$.

It is an old and easily proved theorem that, for a curve of the order m , the number $\delta + \kappa$ of nodes and cusps is at most $= \frac{1}{2}(m-1)(m-2)$; for a given curve the deficiency of the actual number of nodes and cusps below this maximum number, viz.

$\frac{1}{2}(m-1)(m-2) - \delta - \kappa$, is the "*Geschlecht*," or "deficiency," of the curve, say this is D . When $D = 0$, the curve is said to be unicursal, when $= 1$, bicursal, and so on.

The general theorem is that two curves corresponding rationally to each other have the same deficiency. In particular, a curve and its reciprocal have this rational or $(1, 1)$ correspondence, and it has been already seen that a curve and its reciprocal have the same deficiency.

A curve of a given order can in general be rationally transformed into a curve of a lower order; thus a curve of any order for which $D = 0$, that is, a unicursal curve, can be transformed into a line; a curve of any order having the deficiency 1 or 2 can be rationally transformed into a curve of the order $D + 2$, deficiency D ; and a curve of any order deficiency $=$ or > 3 can be rationally transformed into a curve of the order $D + 3$, deficiency D .

Taking x', y', z' as coordinates of a point of the transformed curve, and in its equation writing $x' : y' : z' = 1 : \theta : \phi$, we have ϕ a certain irrational function of θ , and the theorem is that the coordinates x, y, z of any point of the given curve can be expressed as proportional to rational and integral functions of θ, ϕ , that is, of θ and a certain irrational function of θ .

In particular, if $D = 0$, that is, if the given curve be unicursal, the transformed curve is a line, ϕ is a mere linear function of θ , and the theorem is that the coordinates x, y, z of a point of the unicursal curve can be expressed as proportional to rational and integral functions of θ ; it is easy to see that for a given curve of the order m , these functions of θ must be of the same order m .

If $D = 1$, then the transformed curve is a cubic; it can be shown that in a cubic, the axes of coordinates being properly chosen, ϕ can be expressed as the square root of a quartic function of θ ; and the theorem is that the coordinates x, y, z of a point of the bicursal curve can be expressed as proportional to rational and integral functions of θ , and of the square root of a quartic function of θ .

And so if $D = 2$, then the transformed curve is a nodal quartic; ϕ can be expressed as the square root of a sextic function of θ , and the theorem is, that the coordinates x, y, z of a point of the tricursal curve can be expressed as proportional to rational and

integral functions of θ , and of the square root of a sextic function of θ . But when $D = 3$, we have no longer the like law, viz. ϕ is not expressible as the square root of an octic function of θ .

Observe that the radical, square root of a quartic function, is connected with the theory of elliptic functions, and the radical, square root of a sextic function, with that of the first kind of Abelian functions, but that the next kind of Abelian functions does not depend on the radical, square root of an octic function.

It is a form of the theorem for the case $D = 1$, that the coordinates x, y, z of a point of the bicursal curve, or in particular the coordinates of a point of the cubic, can be expressed as proportional to rational and integral functions of the elliptic functions $\text{sn } u, \text{cn } u, \text{dn } u$; in fact, taking the radical to be $\sqrt{(1 - \theta^2 \cdot 1 - k^2 \theta^2)}$, and writing

$$[61-2]$$

$\theta = \operatorname{sn} u$, the radical becomes $= \operatorname{cn} u \cdot \operatorname{dn} u$; and we have expressions of the form in question.

It will be observed that the equations $x' : y' : z' = X : Y : Z$ before-mentioned do not of themselves lead to the other system of equations $x : y : z = X' : Y' : Z'$, and thus that the theory does not in anywise establish a (1, 1) correspondence between the points (x, y, z) and (x', y', z') of two planes or of the same plane; this is the correspondence of Cremona's theory.

In this theory, given in the memoirs "Sulle trasformazioni geometriche delle figure piane," *Mem. di Bologna*, t. II. (1863), and t. V. (1865), we have a system of equations $x' : y' : z' = X : Y : Z$ which does lead to a system $x : y : z = X' : Y' : Z'$, where, as before, X, Y, Z denote rational and integral functions, all of the same order, of the coordinates x, y, z , and X', Y', Z' rational and integral functions, all of the same order, of the coordinates x', y', z' , and there is thus a (1, 1) correspondence given by these equations between the two points (x, y, z) and (x', y', z') . To explain this, observe that starting from the equations $x' : y' : z' = X : Y : Z$, to a given point (x, y, z) there corresponds one point (x', y', z') , but that if n be the order of the functions X, Y, Z , then to a given point x', y', z' there would, if the curves $X = 0, Y = 0, Z = 0$ had no common intersections, correspond n^2 points (x, y, z) . If, however, the functions are such that the curves $X = 0, Y = 0, Z = 0$ have k common intersections, then among the n^2 points are included these k points, which are fixed points independent of the point (x', y', z') ; so that, disregarding these fixed points, the number of points (x, y, z) corresponding to the given point (x', y', z') is $= n^2 - k$; and in particular if $k = n^2 - 1$, then we have one corresponding point; and hence the original system of equations $x' : y' : z' = X : Y : Z$ must lead to the equivalent system $x : y : z = X' : Y' : Z'$; and in this system by the like reasoning the functions must be such that the curves $X' = 0, Y' = 0, Z' = 0$ have $n'^2 - 1$ common intersections. The most simple example is in the two systems of equations $x' : y' : z' = yz : zx : xy$ and $x : y : z = y'z' : z'x' : x'y'$; where $yz = 0, zx = 0, xy = 0$ are conics (pairs of lines) having three common intersections, and where obviously either system of equations leads to the other system. In

the case where X, Y, Z are of an order exceeding 2, the required number $n^2 - 1$ of common intersections can only occur by reason of common multiple points on the three curves; and assuming that the curves $X = 0, Y = 0, Z = 0$ have $a_1 + a_2 + a_3 + \cdots + a_{n-1}$ common intersections, where the a_1 points are ordinary points, the a_2 points are double points, the a_3 points are triple points, &c., on each curve, we have the condition

$$a_1 + 4a_2 + 9a_3 + \cdots + (n-1)^2 a_{n-1} = n^2 - 1;$$

but to this must be joined the condition

$$a_1 + 3a_2 + 6a_3 + \cdots + \frac{1}{2}(n-1)(n-2) a_{n-1} = \frac{1}{2}n(n+3) - 2,$$

(without which the transformation would be illusory); and the conclusion is that $a_1, a_2, \dots, a_{n-1}$ may be any numbers satisfying these two equations. It may be added that the two equations together give

$$a_2 + 3a_3 + \cdots + \frac{1}{2}(n-1)(n-2) a_{n-1} = \frac{1}{2}(n-1)(n-2),$$

which expresses that the curves $X = 0$, $Y = 0$, $Z = 0$ are unicursal. The transformation may be applied to any curve $u = 0$, which is thus rationally transformed into a curve $u' = 0$, by a rational transformation such as is considered in Riemann's theory; hence the two curves have the same deficiency.

Coming next to Chasles, the principle of correspondence is established and used by him in a series of memoirs relating to the conics which satisfy given conditions, and to other geometrical questions, contained in the *Comptes Rendus*, t. LVIII. et seq. (1864 to the present time). The theorem of united points in regard to points in a right line was given in a paper, June–July 1864, and it was extended to unicursal curves in a paper of the same series (March 1866), “Sur les courbes planes ou à double courbure dont les points peuvent se déterminer individuellement—application du principe de correspondance dans la théorie de ces courbes.”

The theorem is as follows: if in a unicursal curve two points have an (α, β) correspondence, then the number of united points (or points each corresponding to itself) is $= \alpha + \beta$. In fact, in a unicursal curve the coordinates of a point are given as proportional to rational and integral functions of a parameter, so that any point of the curve is determined uniquely by means of this parameter; that is, to each point of the curve corresponds one value of the parameter, and to each value of the parameter one point on the curve; and the (α, β) correspondence between the two points is given by an equation of the form $(\theta, 1)^\alpha (\phi, 1)^\beta = 0$ between their parameters θ and ϕ ; at a united point $\phi = \theta$, and the value of θ is given by an equation of the order $\alpha + \beta$. The extension to curves of any given deficiency D was made in the memoir of Cayley, “On the correspondence of two points on a curve,” *Proc. Lond. Math. Soc.* t. I. (1866), [385], viz. taking P, P' as the corresponding points in an (α, α') correspondence on a curve of deficiency D , and supposing that when P is given the corresponding points P' are found as the intersections of the curve by a curve containing the coordinates of P as parameters, and having with the given curve k intersections at the point P , then the number of united points is $a = \alpha + \alpha' + 2kD$; and more generally, if the curve Θ intersect the given curve in a set of points P' each p times, a set of points Q' each q times, &c., in

such manner that the points (P, P') , the points (P, Q') , &c., are pairs of points corresponding to each other according to distinct laws; then if (P, P') are points having an (α, α') correspondence with a number $= a$ of united points, (P, Q') points having a (β, β') correspondence with a number $= b$ of united points, and so on, the theorem is that we have

$$p(a - \alpha - \alpha') + q(b - \beta - \beta') + \dots = 2kD.$$

The principle of correspondence, or say rather the theorem of united points, is a most powerful instrument of investigation, which may be used in place of analysis for the determination of the number of solutions of almost every geometrical problem. We can by means of it investigate the class of a curve, number of inflexions, &c.—in fact, Plücker's equations; but it is necessary to take account of special solutions; thus, in one of the most simple instances, in finding the class of a curve, the cusps present themselves as special solutions.

Imagine a curve of order m , deficiency D , and let the corresponding points P, P' be such that the line joining them passes through a given point O ; this is an $(m-1, m-1)$ correspondence, and the value of k is $= 1$, hence the number of united points is $= 2m-2+2D$; the united points are the points of contact of the tangents from O and (as special solutions) the cusps, and we have thus the relation $n + \kappa = 2m - 2 + 2D$; or, writing $D = \frac{1}{2}(m-1)(m-2) - \delta - \kappa$, this is $n = m(m-1) - 2\delta - 3\kappa$, which is right.

The principle in its original form as applying to a right line was used throughout by Chasles in the investigations on the number of the conics which satisfy given conditions, and on the number of solutions of very many other geometrical problems.

There is one application of the theory of the (α, α') correspondence between two planes which it is proper to notice.

Imagine a curve, real or imaginary, represented by an equation (involving, it may be, imaginary coefficients) between the Cartesian coordinates u, u' ; then, writing $u = x+iy, u' = x'+iy'$, the equation determines real values of (x, y) , and of (x', y') , corresponding to any given real values of (x', y') and (x, y) respectively; that is, it establishes a real correspondence (not of course a rational one) between the points (x, y) and (x', y') ; for example in the imaginary circle $u^2 + u'^2 = (a+bi)^2$, the correspondence is given by the two equations $x^2 - y^2 + x'^2 - y'^2 = a^2 - b^2, xy + x'y' = ab$. We have thus a means of geometrical representation for the portions, as well imaginary as real, of any real or imaginary curve. Considerations such as these have been used for determining the series of values of the independent variable, and the irrational functions thereof in the theory of Abelian integrals, but the theory seems to be worthy of further investigation.

The researches of Chasles (*Comptes Rendus*, t. LVIII., 1864, et seq.) refer to the conics which satisfy given conditions. There is an earlier paper by De Jonquières, "Théorèmes généraux concernant les courbes géométriques planes d'un ordre quelconque," *Liouv.* t. VI. (1861), which establishes the notion of a system of curves (of any order) of the index N , viz. considering the curves of the order n which satisfy $\frac{1}{2}n(n+3) - 1$ conditions, then the index N is the number of these curves which pass through a given arbitrary point.

But Chasles in the first of his papers (February 1864), considering the conics which satisfy four conditions, establishes the notion of the two characteristics (μ, ν) of such a system of conics, viz. μ is the number of the conics which pass through a given arbitrary point, and ν is the number of the conics which touch a given arbitrary line. And he gives the theorem, a system of conics satisfying four conditions, and having the characteristics (μ, ν) contains $2\nu - \mu$ line-pairs (that is, conics, each of them a pair of lines), and $2\mu - \nu$ point-pairs (that is, conics, each of them a pair of points, *coniques infiniment aplaties*), which is a fundamental one in the theory. The characteristics of the system can be determined when it is known how many there are of these two kinds of degenerate conics in the system, and how often each is to be counted. It was thus that Zeuthen (in the paper *Nyt Bydrag*, "Contribution to the Theory of Systems of Conics which satisfy four Conditions," Copenhagen, 1865, translated with an addition in the *Nouvelles Annales*) solved the question of finding the characteristics of the systems of conics which satisfy four

conditions of contact with a given curve or curves; and this led to the solution of the further problem of finding the number of the conics which satisfy five conditions of contact with a given curve or curves (Cayley, *Comptes Rendus*, t. LXIII., 1866, [377]), and “On the Curves which satisfy given Conditions” (*Phil. Trans.* t. CLVIII., 1868, [406]).

It may be remarked that although, as a process of investigation, it is very convenient to seek for the characteristics of a system of conics satisfying 4 conditions, yet what is really determined is in every case the number of the conics which satisfy 5 conditions; the characteristics of the system $(4p)$ of the conics which pass through 4 points are $(5p)$, $(4p, 1l)$, the number of the conics which pass through 5 points, and which pass through 4 points and touch 1 line: and so in other cases. Similarly as regards cubics, or curves of any other order: a cubic depends on 9 constants, and the elementary problems are to find the number of the cubics $(9p)$, $(8p, 1l)$, &c., which pass through 9 points, pass through 8 points and touch 1 line, &c.; but it is in the investigation convenient to seek for the characteristics of the systems of cubics $(8p)$, &c., which satisfy 8 instead of 9 conditions.

The elementary problems in regard to cubics are solved very completely by Maillard in his *Thèse, Recherche des caractéristiques des systèmes élémentaires des courbes planes du troisième ordre* (Paris, 1871). Thus, considering the several cases of a cubic

	No. of const.
1. With a given cusp	5,
2. „ cusp on given line	6,
3. „ cusp	7,
4. „ a given node	6,
5. „ node on given line	7,
6. „ node	8,
7. non-singular	9,

he determines in every case the characteristics (μ, ν) of the corresponding systems of cubics $(4p)$, $(3p, 1l)$, &c. The same problems, or most of them, and also the elementary problems in regard to quartics are solved by Zeuthen, who in the elaborate memoir “Almindelige Egenskaber, &c.,” Danish Academy, t. X. (1873), considers

the problem in reference to curves of any order, and applies his results to cubic and quartic curves.

The methods of Maillard and Zeuthen are substantially identical; in each case the question considered is that of finding the characteristics (μ, ν) of a system of curves by consideration of the special or degenerate forms of the curves included in the system. The quantities which have to be considered are very numerous. Zeuthen in the case of curves of any given order establishes between the characteristics μ, ν , and 18 other quantities, in all 20 quantities, a set of 24 equations (equivalent to 23 independent equations), involving (besides the 20 quantities) other quantities relating to the various forms of the degenerate curves, which supplementary terms he determines, partially for curves of any order, but completely only for quartic curves. It is in the discussion and complete enumeration of the special or degenerate forms of the curves,

and of the supplementary terms to which they give rise, that the great difficulty of the question seems to consist; it would appear that the 24 equations are a complete system, and that (subject to a proper determination of the supplementary terms) they contain the solution of the general problem.

The remarks which follow have reference to the analytical theory of the degenerate curves which present themselves in the foregoing problem of the curves which satisfy given conditions.

A curve represented by an equation in point-coordinates may break up: thus if $P_1, P_2, \dots$ be rational and integral functions of the coordinates (x, y, z) of the orders $m_1, m_2, \dots$ respectively, we have the curve $P_1^{a_1} P_2^{a_2} \dots = 0$, of the order $m = a_1 m_1 + a_2 m_2 + \dots$, composed of the curve $P_1 = 0$ taken a_1 times, the curve $P_2 = 0$ taken a_2 times, &c.

Instead of the equation $P_1^{a_1} P_2^{a_2} \dots = 0$, we may start with an equation $u = 0$, where u is a function of the order m containing a parameter θ , and for a particular value say $\theta = 0$, of the parameter reducing itself to $P_1^{a_1} P_2^{a_2} \dots$. Supposing θ indefinitely small, we have what may be called the penultimate curve, and when $\theta = 0$ the ultimate curve. Regarding the ultimate curve as derived from a given penultimate curve, we connect with the ultimate curve, and consider as belonging to it, certain points called "summits" on the component curves $P_1 = 0, P_2 = 0$, respectively; a summit S is a point such that, drawing from an arbitrary point O the tangents to the penultimate curve, we have OS as the limit of one of these tangents. The ultimate curve together with its summits may be regarded as a degenerate form of the curve $u = 0$. Observe that the positions of the summits depend on the penultimate curve $u = 0$, viz. on the values of the coefficients in the terms multiplied by $\theta, \theta^2, \dots$; they are thus in some measure arbitrary points as regards the ultimate curve $P_1^{a_1} P_2^{a_2} \dots = 0$.

It may be added that we have summits only on the component curves $P_i = 0$, of a multiplicity $a_i > 1$; the number of summits on such a curve is in general $= (a_i^2 - a_i)m_i^2$. Thus assuming that the penultimate curve is without nodes or cusps, the number of the tangents to it is $= m^2 - m = (a_1 m_1 + a_2 m_2 + \dots)^2 - (a_1 m_1 + a_2 m_2 + \dots)$, taking $P_1 = 0$ to have δ_1 nodes and κ_1 cusps, and therefore

its class n_1 to be $= m_1^2 - m_1 - 2\delta_1 - 3\kappa_1$, &c., the expression for the number of tangents to the penultimate curve is

$$= (a_1^2 - a_1) m_1^2 + (a_2^2 - a_2) m_2^2 + \dots \\ + 2a_1 a_2 m_1 m_2 + \dots + a_1 (n_1 + 2\delta_1 + 3\kappa_1) + a_2 (n_2 + 2\delta_2 + 3\kappa_2) + \dots$$

where a term $2a_1 a_2 m_1 m_2$ indicates tangents which are in the limit the lines drawn to the intersections of the curves $P_1 = 0$, $P_2 = 0$, each line $2a_1 a_2$ times; a term $a_1 (n_1 + 2\delta_1 + 3\kappa_1)$ tangents which are in the limit the proper tangents to $P_1 = 0$ each a_1 times, the lines to its nodes each $2a_1$ times, and the lines to its cusps each $3a_1$ times; the remaining terms $(a_1^2 - a_1) m_1^2 + (a_2^2 - a_2) m_2^2 + \dots$ indicate tangents which are in the limit the lines drawn to the several summits, that is, we have $= (a_i^2 - a_i) m_i^2$ summits on the curve $P_i = 0$.

There is of course a precisely similar theory as regards line-coordinates; taking Π_1 , Π_2 , &c., to be rational and integral functions of the coordinates (ξ, η, ζ) , we connect with the ultimate curve $= \Pi_1^{a_1} \Pi_2^{a_2} \dots = 0$, and consider as belonging to it certain lines, which for the moment may be called "axes," tangents to the component curves

$\Pi_1 = 0$, $\Pi_2 = 0$ respectively. Considering an equation in point-coordinates, we may have among the component curves right lines; and, if in order to put these in evidence, we take the equation to be $L_1^{a_1} \dots P_1^{b_1} \dots = 0$, where $L_1 = 0$ is a right line, $P_1 = 0$ a curve of the second or any higher order, then the curve will contain as part of itself summits not exhibited in this equation, but the corresponding line-equation will be $\Lambda_1^{a_1} \dots \Pi_1^{b_1} \dots = 0$, where $\Lambda_1 = 0$, $\dots$ are the equations of the summits in question, $\Pi_1 = 0$, &c., are the line-equations corresponding to the several point-equations $P_1 = 0$, &c.; and this curve will contain as part of itself axes not exhibited by this equation, but which are the lines $L_1 = 0$, $\dots$ of the equation in point-coordinates.

In conclusion a little may be said as to curves of double curvature, otherwise twisted curves, or curves in space. The analytical theory by Cartesian coordinates was first considered by Clairaut, *Recherches sur les courbes à double courbure* (Paris, 1731). Such a curve may be considered as described by a point, moving in a line which at the same time rotates about the point in a plane which at the same time rotates about the line; the point is a point, the line a tangent, and the plane an osculating plane, of the curve; moreover the line is a generating line, and the plane a tangent plane, of a developable surface or torse, having the curve for its edge of regression. Analogous to the order and class of a plane curve we have the order, rank, and class, of the system (assumed to be a geometrical one), viz. if an arbitrary plane contains m points, an arbitrary line meets r lines, and an arbitrary point lies in n planes, of the system, then m , r , n are the order, rank, and class respectively. The system has singularities, and there exist between m , r , n and the numbers of the several singularities equations analogous to Plücker's equations for a plane curve.

It is a leading point in the theory that a curve in space cannot in general be represented by means of two equations $U = 0$, $V = 0$; the two equations represent surfaces, intersecting in a curve; but there are curves which are not the complete intersection of any two surfaces; thus we have the cubic in space, or skew cubic, which is the residual intersection of two quadric surfaces which have a line in common; the equations $U = 0$, $V = 0$ of the two quadric surfaces

represent the cubic curve, not by itself, but together with the line.
 [C. XI. 62]

.....

786.

EQUATION.

[From the *Encyclopædia Britannica*, Ninth Edition, vol. VIII. (1878),
 pp. 497—509.]

THE present article includes Determinant and Theory of Equations; and it may be proper to explain the relation to each other of the two subjects. Theory of Equations is used in its ordinary conventional sense to denote the theory of a single equation of any order in one unknown quantity; that is, it does not include the theory of a system or systems of equations of any order between any number of unknown quantities. Such systems occur very frequently in analytical geometry and other parts of mathematics, but they are hardly as yet the subject-matter of a distinct theory; and even Elimination, the transition-process for passing from a system of any number of equations involving the same number of unknown quantities to a single equation in one unknown quantity, hardly belongs to the Theory of Equations in the above restricted sense. But there is one case of a system of equations which precedes the Theory of Equations, and indeed presents itself at the outset of algebra, that of a system of simple (or linear) equations. Such a system gives rise to the function called a Determinant, and it is by means of these functions that the solution of the equations is effected. We have thus the subject Determinant as nearly equivalent to (but somewhat more extensive than) that of a system of linear equations; and we have the other subject, Theory of Equations, used in the restricted sense above referred to, and as not including Elimination.

DETERMINANT.

1. A sketch of the history of determinants is given under [the Article] Algebra; it thereby appears that the algebraical function called a determinant presents itself in the solution of a system of simple equations, and we have herein a natural source of the theory. Thus, considering the equations

$$\begin{aligned}ax + by + cz &= d, \\a'x + b'y + c'z &= d', \\a''x + b''y + c''z &= d'',\end{aligned}$$

and proceeding to solve them by the so-called method of cross multiplication, we multiply the equations by factors selected in such a manner that, upon adding the results, the whole coefficient of y becomes $= 0$ and the whole coefficient of z becomes $= 0$; the factors in question are $b'c'' - b''c'$, $b''c - bc''$, $bc' - b'c$ (values which, as at once seen, have the desired property); we thus obtain an equation which contains on the left-hand side only a multiple of x , and on the right-hand side a constant term; the coefficient of x has the value

$$a(b'c'' - b''c') + a'(b''c - bc'') + a''(bc' - b'c),$$

and this function, represented in the form

$$\begin{vmatrix} a, & b, & c \\ a', & b', & c' \\ a'', & b'', & c'' \end{vmatrix},$$

is said to be a determinant; or, the number of elements being 3^2 , it is called a determinant of the third order. It is to be noticed that the resulting equation is

$$x \begin{vmatrix} a, & b, & c \\ a', & b', & c' \\ a'', & b'', & c'' \end{vmatrix} = \begin{vmatrix} d, & b, & c \\ d', & b', & c' \\ d'', & b'', & c'' \end{vmatrix},$$

where the expression on the right-hand side is the like function with d, d', d'' in place of a, a', a'' respectively, and is of course also a determinant. Moreover, the functions $b'c'' - b''c'$, $b''c - bc''$, $bc' - b'c$ used in the process are themselves the determinants of the second order

$$\begin{vmatrix} b', & c' \\ b'', & c'' \end{vmatrix}, \quad \begin{vmatrix} b'', & c'' \\ b, & c \end{vmatrix}, \quad \begin{vmatrix} b, & c \\ b', & c' \end{vmatrix}.$$

We have herein the suggestion of the rule for the derivation of the determinants of the orders 1, 2, 3, 4, &c., each from the preceding

one, viz. we have

$$\begin{aligned}
 |a| &= a, \\
 \begin{vmatrix} a, & b \\ a', & b' \end{vmatrix} &= a |b'| - a' |b|, \\
 \begin{vmatrix} a, & b, & c \\ a', & b', & c' \\ a'', & b'', & c'' \end{vmatrix} &= a \begin{vmatrix} b', & c' \\ b'', & c'' \end{vmatrix} - a' \begin{vmatrix} b, & c \\ b'', & c'' \end{vmatrix} + a'' \begin{vmatrix} b, & c \\ b', & c' \end{vmatrix},
 \end{aligned}$$

and so on, the terms being all + for a determinant of an odd order, but alternately + and - for a determinant of an even order.

[62-2]

2. It is easy, by induction, to arrive at the general results:—

A determinant of the order n is the sum of the $1 \cdot 2 \cdot 3 \dots n$ products which can be formed with n elements out of n^2 elements arranged in the form of a square, no two of the n elements being in the same line or in the same column, and each such product having the coefficient $\pm$ unity.

The products in question may be obtained by permuting in every possible manner the columns (or the lines) of the determinant, and then taking for the factors the n elements in the dexter diagonal. And we thence derive the rule for the signs, viz. considering the primitive arrangement of the columns as positive, then an arrangement obtained therefrom by a single interchange (inversion, or derangement) of two columns is regarded as negative; and so in general an arrangement is positive or negative according as it is derived from the primitive arrangement by an even or an odd number of interchanges. This implies the theorem that a given arrangement can be derived from the primitive arrangement only by an odd number, or else only by an even number of interchanges,—a theorem the verification of which may be easily obtained from the theorem (in fact, a particular case of the general one), an arrangement can be derived from itself only by an even number of interchanges. And this being so, each product has the sign belonging to the corresponding arrangement of the columns; in particular, a determinant contains with the sign $+$ the product of the elements in its dexter diagonal. It is to be observed that the rule gives as many positive as negative arrangements, the number of each being $= \frac{1}{2} \cdot 1 \cdot 2 \dots n$.

The rule of signs may be expressed in a different form. Giving to the columns in the primitive arrangement the numbers $1, 2, 3, \dots, n$, to obtain the sign belonging to any other arrangement we take, as often as a lower number succeeds a higher one, the sign $-$, and, compounding together all these minus signs, obtain the proper sign, $+$ or $-$ as the case may be.

Thus, for three columns, it appears by either rule that 123, 231, 312 are positive; 132, 321, 213 are negative; and the developed expression of the foregoing determinant of the third order is

$$= ab'c'' - ab''c' + a'b''c - a'bc'' + a''bc' - a''b'c.$$

3. It further appears that a determinant is a linear function³ of the elements of each column thereof, and also a linear function of the elements of each line thereof; moreover, that the determinant retains the same value, only its sign being altered, when any two columns are interchanged, or when any two lines are interchanged; more generally, when the columns are permuted in any manner, or when the lines are permuted in any manner, the determinant retains its original value, with the sign $+$ or $-$ according as the new arrangement (considered as derived from the primitive arrangement) is positive or negative according to the foregoing rule of signs.

³The expression, a linear function, is here used in its narrowest sense, a linear function without constant term; what is meant is, that the determinant is in regard to the elements $a, a', a'', \dots$ of any column or line thereof, a function of the form $Aa + A'a' + A''a'' + \dots$, without any term independent of $a, a', a'', \dots$.

It at once follows that, if two columns are identical, or if two lines are identical, the value of the determinant is $= 0$. It may be added that, if the lines are converted into columns, and the columns into lines, in such a way as to leave the dexter diagonal unaltered, the value of the determinant is unaltered; the determinant is in this case said to be transposed.

4. By what precedes it appears that there exists a function of the n^2 elements, linear as regards the terms of each column (or say, for shortness, linear as to each column), and such that only the sign is altered when any two columns are interchanged; these properties completely determine the function, except as to a common factor which may multiply all the terms. If, to get rid of this arbitrary common factor, we assume that the product of the elements in the dexter diagonal has the coefficient $+1$, we have a complete definition of the determinant; and it is interesting to show how from these properties, assumed for the definition of the determinant, it at once appears that the determinant is a function serving for the solution of a system of linear equations. Observe that the properties show at once that if any column is 0 (that is, if the elements in the column are each $= 0$), then the determinant is $= 0$; and further that, if any two columns are identical, then the determinant is $= 0$.

5. Reverting to the system of linear equations written down at the beginning of this article, consider the determinant

$$\begin{vmatrix} ax + by + cz - d, & b, & c \\ a'x + b'y + c'z - d', & b', & c' \\ a''x + b''y + c''z - d'', & b'', & c'' \end{vmatrix};$$

it appears that this is

$$= x \begin{vmatrix} a, & b, & c \\ a', & b', & c' \\ a'', & b'', & c'' \end{vmatrix} + y \begin{vmatrix} b, & b, & c \\ b', & b', & c' \\ b'', & b'', & c'' \end{vmatrix} + z \begin{vmatrix} c, & b, & c \\ c', & b', & c' \\ c'', & b'', & c'' \end{vmatrix} - \begin{vmatrix} d, & b, & c \\ d', & b', & c' \\ d'', & b'', & c'' \end{vmatrix};$$

viz. the second and the third terms each vanishing, it is

$$= x \begin{vmatrix} a, & b, & c \\ a', & b', & c' \\ a'', & b'', & c'' \end{vmatrix} - \begin{vmatrix} d, & b, & c \\ d', & b', & c' \\ d'', & b'', & c'' \end{vmatrix}.$$

But if the linear equations hold good, then the first column of the original determinant is $= 0$, and therefore the determinant itself is $= 0$; that is, the linear equations give

$$x \begin{vmatrix} a, & b, & c \\ a', & b', & c' \\ a'', & b'', & c'' \end{vmatrix} - \begin{vmatrix} d, & b, & c \\ d', & b', & c' \\ d'', & b'', & c'' \end{vmatrix} = 0;$$

which is the result obtained above.

We might in a similar way find the values of y and z , but there is a more symmetrical process. Join to the original equations the new equation

$$\alpha x + \beta y + \gamma z = \delta;$$

a like process shows that, the equations being satisfied, we have

$$\begin{vmatrix} \alpha, & \beta, & \gamma, & -\delta \\ a, & b, & c, & d \\ a', & b', & c', & d' \\ a'', & b'', & c'', & d'' \end{vmatrix} = 0;$$

or, as this may be written,

$$\begin{vmatrix} \alpha, & \beta, & \gamma \\ a, & b, & c \\ a', & b', & c' \\ a'', & b'', & c'' \end{vmatrix} \cdot \delta - \begin{vmatrix} a, & b, & c, & d \\ a', & b', & c', & d' \\ a'', & b'', & c'', & d'' \end{vmatrix} = 0;$$

which, considering δ as standing herein for its value $\alpha x + \beta y + \gamma z$, is a consequence of the original equations only. We have thus an expression for $\alpha x + \beta y + \gamma z$, an arbitrary linear function of the unknown quantities x, y, z ; and by comparing the coefficients of α, β, γ on the two sides respectively, we have the values of x, y, z ; in fact, these quantities, each multiplied by

$$\begin{vmatrix} a, & b, & c \\ a', & b', & c' \\ a'', & b'', & c'' \end{vmatrix},$$

are in the first instance obtained in the forms

$$\begin{vmatrix} a, & b, & c, & d \\ a', & b', & c', & d' \\ a'', & b'', & c'', & d'' \end{vmatrix} \quad (\text{the columns being taken in appropriate orders}),$$

but these are

$$\begin{vmatrix} b, & c, & d \\ b', & c', & d' \\ b'', & c'', & d'' \end{vmatrix}, \quad \begin{vmatrix} d, & c, & a \\ d', & c', & a' \\ d'', & c'', & a'' \end{vmatrix}, \quad \begin{vmatrix} d, & a, & b \\ d', & a', & b' \\ d'', & a'', & b'' \end{vmatrix}$$

or, what is the same thing,

$$\begin{vmatrix} d, & b, & c \\ d', & b', & c' \\ d'', & b'', & c'' \end{vmatrix}, \quad \begin{vmatrix} a, & d, & c \\ a', & d', & c' \\ a'', & d'', & c'' \end{vmatrix}, \quad \begin{vmatrix} a, & b, & d \\ a', & b', & d' \\ a'', & b'', & d'' \end{vmatrix}$$

respectively.

6. *Multiplication of two determinants of the same order.*—The theorem is obtained very easily from the last preceding definition of a determinant. It is most simply expressed thus

$$\begin{vmatrix} (a, b, c)(\alpha, \alpha', \alpha''), & (a, b, c)(\beta, \beta', \beta''), & (a, b, c)(\gamma, \gamma', \gamma'') \\ (a', b', c')(\alpha, \alpha', \alpha''), & (a', b', c')(\beta, \beta', \beta''), & (a', b', c')(\gamma, \gamma', \gamma'') \\ (a'', b'', c'')(\alpha, \alpha', \alpha''), & (a'', b'', c'')(\beta, \beta', \beta''), & (a'', b'', c'')(\gamma, \gamma', \gamma'') \end{vmatrix} = \begin{vmatrix} a, & b, & c \\ a', & b', & c' \\ a'', & b'', & c'' \end{vmatrix} \cdot \begin{vmatrix} \alpha, & \beta, & \gamma \\ \alpha', & \beta', & \gamma' \\ \alpha'', & \beta'', & \gamma'' \end{vmatrix},$$

where the expression on the left side stands for a determinant, the terms of the first line being $(a, b, c)(\alpha, \alpha', \alpha'')$, that is, $a\alpha + b\alpha' + c\alpha''$, $(a, b, c)(\beta, \beta', \beta'')$, that is, $a\beta + b\beta' + c\beta''$, $(a, b, c)(\gamma, \gamma', \gamma'')$, that is, $a\gamma + b\gamma' + c\gamma''$; and similarly the terms in the second and third lines are the like functions with (a', b', c') and (a'', b'', c'') respectively.

There is an apparently arbitrary transposition of lines and columns; the result would hold good if on the left-hand side we had written (α, β, γ) , $(\alpha', \beta', \gamma')$, $(\alpha'', \beta'', \gamma'')$, or what is the same thing, if on the right-hand side we had transposed the second determinant; and either of these changes would, it might be thought, increase the elegance of the form, but, for a reason which need not be explained⁴, the form actually adopted is the preferable one.

To indicate the method of proof, observe that the determinant on the left-hand side, qua linear function of its columns, may be broken up into a sum of ($3^3 =$) 27 determinants, each of which is either of some such form as

$$\alpha\beta\gamma' \begin{vmatrix} a, & a, & b \\ a', & a', & b' \\ a'', & a'', & b'' \end{vmatrix},$$

where the term $\alpha\beta\gamma'$ is not a term of the $\alpha\beta\gamma$ -determinant, and its coefficient (as a determinant with two identical columns) vanishes; or else it is of a form such as

$$\alpha\beta'\gamma'' \begin{vmatrix} a, & b, & c \\ a', & b', & c' \\ a'', & b'', & c'' \end{vmatrix};$$

that is, every term which does not vanish contains as a factor the abc -determinant last written down; the sum of all other factors

⁴The reason is the connexion with the corresponding theorem for the multiplication of two matrices.

$\pm\alpha\beta'\gamma''$ is the $\alpha\beta\gamma$ -determinant of the formula; and the final result then is, that the determinant on the left-hand side is equal to the product on the right-hand side of the formula.

7. *Decomposition of a determinant into complementary determinants.*—Consider, for simplicity, a determinant of the fifth order, $5 = 2 + 3$, and let the top two lines be

$$\begin{array}{c} a, \ b, \ c, \ d, \ e \\ a', \ b', \ c', \ d', \ e'; \end{array}$$

then, if we consider how these elements enter into the determinant, it is at once seen that they enter only through the determinants of the second order $\begin{vmatrix} a & b \\ a' & b' \end{vmatrix}$, &c., which can be formed by selecting any two columns at pleasure. Moreover, representing the remaining three lines by

$$\begin{aligned} &a'', b'', c'', d'', e'' \\ &a''', b''', c''', d''', e''' \\ &a''', b''', c''', d''', e''', \end{aligned}$$

it is further seen that the factor which multiplies the determinant formed with any two columns of the first set is the determinant of the third order formed with the complementary three columns of the second set; and it thus appears that the determinant of the fifth order is a sum of all the products of the form

$$\begin{vmatrix} a & b \\ a' & b' \end{vmatrix} \cdot \begin{vmatrix} c'' & d'' & e'' \\ c''' & d''' & e''' \\ c'''' & d'''' & e'''' \end{vmatrix},$$

the sign + being in each case such that the sign of the term $\pm ab' \cdot c''d'''e''''$ obtained from the diagonal elements of the component determinants may be the actual sign of this term in the determinant of the fifth order; for the product written down the sign is obviously +.

Observe that for a determinant of the n th order, taking the decomposition to be $1 + (n - 1)$, we fall back upon the equations given at the commencement, in order to show the genesis of a determinant.

8. Any determinant $\begin{vmatrix} a & b \\ a' & b' \end{vmatrix}$ formed out of the elements of the original determinant, by selecting the lines and columns at pleasure, is termed a minor of the original determinant; and when the number of lines and columns, or order of the determinant, is $n - 1$, then such determinant is called a first minor; the number of the first minors is $= n^2$, the first minors, in fact, corresponding to the several elements of the determinant—that is, the coefficient therein of any term whatever is the corresponding first minor. The first minors,

each divided by the determinant itself, form a system of elements inverse to the elements of the determinant.

A determinant is symmetrical when every two elements symmetrically situated in regard to the dexter diagonal are equal to each other; if they are equal and opposite (that is, if the sum of the two elements be $= 0$), this relation not extending to the diagonal elements themselves, which remain arbitrary, then the determinant is skew; but if the relation does extend to the diagonal terms (that is, if these are each $= 0$), then the determinant is skew symmetrical; thus the determinants

$$\begin{vmatrix} a, & h, & g \\ h, & b, & f \\ g, & f, & c \end{vmatrix}, \quad \begin{vmatrix} a, & -\nu, & \mu \\ \nu, & b, & -\lambda \\ -\mu, & \lambda, & c \end{vmatrix}, \quad \begin{vmatrix} 0, & \nu, & -\mu \\ -\nu, & 0, & \lambda \\ \mu, & -\lambda, & 0 \end{vmatrix}$$

are respectively symmetrical, skew, and skew symmetrical.

The theory admits of very extensive algebraic developments, and applications in algebraical geometry and other parts of mathematics; but the fundamental properties of the functions may fairly be considered as included in what precedes.

THEORY OF EQUATIONS.

9. In the subject "Theory of Equations," the term equation is used to denote an equation of the form $x^n - p_1x^{n-1} + \dots + p_n = 0$, where $p_1, p_2, \dots, p_n$ are regarded as known, and x as a quantity to be determined; for shortness, the equation is written $f(x) = 0$.

The equation may be *numerical*; that is, the coefficients $p_1, p_2, \dots, p_n$ are then numbers,—understanding by number a quantity of the form $\alpha + \beta i$ (α and β having any positive or negative real values whatever, or say each of these is regarded as susceptible of continuous variation from an indefinitely large negative to an indefinitely large positive value), and i denoting $\sqrt{-1}$.

Or the equation may be *algebraical*; that is, the coefficients are not then restricted to denote, or are not explicitly considered as denoting, numbers.

I. We consider first numerical equations. (Real theory, 10 to 14; Imaginary theory, 15 to 18.)

10. Postponing all consideration of imaginaries, we take in the first instance the coefficients to be real, and attend only to the real roots (if any); that is, $p_1, p_2, \dots, p_n$ are real positive or negative quantities, and a root α , if it exists, is a positive or negative quantity such that $\alpha^n - p_1\alpha^{n-1} + \dots + p_n = 0$, or say, $f(\alpha) = 0$. The fundamental theorems are given in the article Algebra, sections X., XIII., XIV.; but there are various points in the theory which require further development.

It is very useful to consider the curve $y = f(x)$, or, what would come to the same, the curve $Ay = f(x)$,—but it is better to retain the first-mentioned form of equation, drawing, if need be, the ordinate y on a reduced scale. For instance, if the given equation be $x^3 - 6x^2 + 11x - 6.06 = 0$,⁵ then the curve $y = x^3 - 6x^2 + 11x - 6.06$ is as shown in the figure at page 501, without any reduction of scale for

⁵The coefficients were selected so that the roots might be nearly 1, 2, 3.

the ordinate. It is clear that, in general, y is a continuous one-valued function of x , finite for every finite value of x , but becoming infinite when x is infinite; i.e. assuming throughout that the coefficient of x^n is $+1$, then when $x = +\infty$, $y = +\infty$; but when $x = -\infty$, then $y = +\infty$ or $-\infty$, according as n is even or odd; the curve cuts any line whatever, and in particular it cuts the axis of x , in at most n points; and the value of x , at any point of intersection with the axis, is a root of the equation $f(x) = 0$.

If β , α are any two values of x ($\alpha > \beta$, that is, α nearer $+\infty$), then if $f(\beta)$, $f(\alpha)$ have opposite signs, the curve cuts the axis an odd number of times, and therefore at least once, between the points $x = \beta$, $x = \alpha$; but if $f(\beta)$, $f(\alpha)$ have the same sign, then between these points the curve cuts the axis an even number of times, or it may be not at all. That is, $f(\beta)$, $f(\alpha)$ having opposite signs, there are between the limits β , α an odd number of real roots, and therefore at least one real

[*C. XI.*

63]

root; but $f(\beta)$, $f(\alpha)$ having the same sign, there are between these limits an even number of real roots, or it may be there is no real root. In particular, by giving to β , α the values $-\infty$, $+\infty$ (or, what is the same thing, any two values sufficiently near to these values respectively) it appears that an equation of an odd order has always an odd number of real roots, and therefore at least one real root; but that an equation of an even order has an even number of real roots, or it may be no real root.

If α be such that for $x =$ or $> \alpha$ (that is, x nearer to $+\infty$) $f(x)$ is always $+$, and β be such that for $x =$ or $< \beta$ (that is, x nearer to $-\infty$) $f(x)$ is always $-$, then the real roots (if any) lie between these limits $x = \beta$, $x = \alpha$; and it is easy to find by trial such two limits including between them all the real roots (if any).

11. Suppose that the positive value δ is an inferior limit to the difference between two real roots of the equation; or rather (since the foregoing expression would imply the existence of real roots) suppose that there are not two real roots such that their difference taken positively is $=$ or $< \delta$; then, γ being any value whatever, there is clearly at most one real root between the limits γ and $\gamma + \delta$; and by what precedes there is such real root or there is not such real root, according as $f(\gamma)$, $f(\gamma + \delta)$ have opposite signs or have the same sign. And by dividing in this manner the interval β to α into intervals each of which is $=$ or $< \delta$, we should not only ascertain the number of the real roots (if any), but we should also separate the real roots, that is, find for each of them limits γ , $\gamma + \delta$ between which there lies this one, and only this one, real root.

In particular cases it is frequently possible to ascertain the number of the real roots, and to effect their separation by trial or otherwise, without much difficulty; but the foregoing was the general process as employed by Lagrange even in the second edition (1808) of the *Traité de la résolution des Equations Numériques*⁶; the determination of the limit δ had to be effected by means of the “equation of differences” or equation of the order $\frac{1}{2}n(n-1)$, the roots of which are the squares of the differences of the roots of the

⁶The third edition (1826) is a reproduction of that of 1808; the first edition has the date 1798, but a large part of the contents is taken from memoirs of 1767–68 and 1770–71.

given equation, and the process is a cumbrous and unsatisfactory one.

12. The great step was effected by Sturm's theorem (1835)—viz. here starting from the function $f(x)$, and its first derived function $f'(x)$, we have (by a process which is a slight modification of that for obtaining the greatest common measure of these two functions) to form a series of functions

$$f(x), f'(x), f_2(x), \dots, f_n(x)$$

of the degrees $n, n-1, n-2, \dots, 0$ respectively, the last term $f_n(x)$ being thus an absolute constant. These lead to the immediate determination of the number of real roots (if any) between any two given limits α, β ; viz. supposing $\alpha > \beta$ (that is, α nearer to $+\infty$), then substituting successively these two values in the series of functions, and attending only to the signs of the resulting values, the number of the changes of sign lost in passing from β to α is the required number of real roots

between the two limits. In particular, taking $\beta, \alpha = -\infty, +\infty$ respectively, the signs of the several functions depend merely on the signs of the terms which contain the highest powers of x , and are seen by inspection, and the theorem thus gives at once the whole number of real roots.

And although theoretically, in order to complete by a finite number of operations the separation of the real roots, we still need to know the value of the before-mentioned limit δ ; yet in any given case the separation may be effected by a limited number of repetitions of the process. The practical difficulty is when two or more roots are very near to each other. Suppose, for instance, that the theorem shows that there are two roots between 0 and 10; by giving to x the values 1, 2, 3, . . . successively, it might appear that the two roots were between 5 and 6; then again that they were between $5 \cdot 3$ and $5 \cdot 4$, then between $5 \cdot 34$ and $5 \cdot 35$, and so on until we arrive at a separation; say it appears that between $5 \cdot 346$ and $5 \cdot 347$ there is one root, and between $5 \cdot 348$ and $5 \cdot 349$ the other root. But in the case in question δ would have a very small value, such as $\cdot 002$, and even supposing this value known, the direct application of the first-mentioned process would be still more laborious.

13. Supposing the separation once effected, the determination of the single real root which lies between the two given limits may be effected to any required degree of approximation either by the processes of Horner and Lagrange (which are in principle a carrying out of the method of Sturm's theorem), or by the process of Newton, as perfected by Fourier (which requires to be separately considered).

First as to Horner and Lagrange. We know that between the limits β, α there lies one, and only one, real root of the equation; $f(\beta)$ and $f(\alpha)$ have therefore opposite signs. Suppose any intermediate value is θ ; in order to determine by Sturm's theorem whether the root lies between β, θ , or between θ, α , it would be quite unnecessary to calculate the signs of $f(\theta), f'(\theta), f_2(\theta), \dots$; only the sign of $f(\theta)$ is required: for, if this has the same sign as $f(\beta)$, then the root is between θ, α ; if the same sign as $f(\alpha)$, then the root is between β, θ . We want to make θ increase from the inferior limit β , at which $f(\theta)$ has the sign of $f(\beta)$, so long as $f(\theta)$ retains this sign, and then to a value for which it assumes the opposite sign; we have thus two

nearer limits of the required root, and the process may be repeated indefinitely.

Horner's method (1819) gives the root as a decimal, figure by figure; thus, if the equation be known to have one real root between 0 and 10, it is in effect shown—say that 5 is too small (that is, the root is between 5 and 6); next that $5 \cdot 4$ is too small (that is, the root is between $5 \cdot 4$ and $5 \cdot 5$); and so on to any number of decimals. Each figure is obtained, not by the successive trial of all the figures which precede it, but (as in the ordinary process of the extraction of a square root, which is in fact Horner's process applied to this particular case) it is given presumptively as the first figure of a quotient; such value may be too large, and then the next inferior integer must be tried instead of it, or it may require to be further diminished. And it is to be remarked that the process not only gives the approximate value α of the root, but (as in the extraction of a square root) it includes the calculation of the function $f(\alpha)$ which should be, and approximately is, $= 0$. The arrangement of the

[63–2]

calculations is very elegant, and forms an integral part of the actual method. It is to be observed that after a certain number of decimal places have been obtained, a good many more can be found by a mere division. It is in the progress tacitly assumed that the roots have been first separated.

Lagrange's method (1767) gives the root as a continued fraction $a + \frac{1}{b + \frac{1}{c + \dots}}$, where a is a positive or negative integer (which may be $= 0$), but $b, c, \dots$ are positive integers. Suppose the roots have been separated; then (by trial if need be of consecutive integer values) the limits may be made to be consecutive integer numbers: say they are $a, a + 1$; the value of x is therefore $= a + \frac{1}{y}$, where y is positive and greater than 1; from the given equation for x , writing therein $x = a + \frac{1}{y}$, we form an equation of the same order for y , and this equation will have one, and only one, positive root greater than 1; hence finding for it the limits $b, b + 1$ (where b is $=$ or > 1), we have $y = b + \frac{1}{z}$, where z is positive and greater than 1; and so on; that is, we thus obtain the successive denominators $b, c, d, \dots$ of the continued fraction. The method is theoretically very elegant, but the disadvantage is that it gives the result in the form of a continued fraction, which for the most part must ultimately be converted into a decimal. There is one advantage in the method, that a commensurable root (that is, a root equal to a rational fraction) is found accurately, since, when such root exists, the continued fraction terminates.

14. Newton's method (1711), as perfected by Fourier (1831), may be roughly stated as follows. If $x = \gamma$ be an approximate value of any root, and $\gamma + h$ the correct value, then $f(\gamma + h) = 0$, that is,

$$f(\gamma) + hf'(\gamma) + \frac{1}{1.2}h^2f''(\gamma) + \dots = 0;$$

and then, if h be so small that the terms after the second may be neglected, $f(\gamma) + hf'(\gamma) = 0$, that is, $h = -\frac{f(\gamma)}{f'(\gamma)}$, or the new approximate value is $\gamma' = \gamma - \frac{f(\gamma)}{f'(\gamma)}$, and so on, as often as we

please. It will be observed that so far nothing has been assumed as to the separation of the roots, or even as to the existence of a real root; γ has been taken as the approximate value of a root, but no precise meaning has been attached to this expression. The question arises, what are the conditions to be satisfied by γ in order that the process may by successive repetitions actually lead to a certain real root of the equation; or say that, γ being an approximate value of a certain real root, the new value $\gamma - \frac{f(\gamma)}{f'(\gamma)}$ may be a more approximate value.

Referring to the figure, it is easy to see that, if OC represent the assumed value γ , then, drawing the ordinate CP to meet the curve in P , and the tangent PO' to meet the axis in C' , we shall have OC' as the new approximate value of the root. But observe that there is here a real root OX , and that the curve beyond [the article ends on page 500 mid-discussion; continues in the next chunk].

[*The discussion of Newton's method, Fourier's conditions, and the imaginary theory continues at page 501; see the volume's next chunk (pages 501–550).*]