

Seminar on Algebraic Geometry of Bois Marie

1966–67

Intersection Theory and the Riemann–Roch Theorem

(SGA 6)

a seminar directed by

P. Berthelot, A. Grothendieck, L. Illusie

with the collaboration of

D. Ferrand, J.-P. Jouanolou, O. Jussila, S. Kleiman, M. Raynaud, J.-P. Serre

Preface

The present edition of *Séminaire de Géométrie Algébrique 6* reproduces, up to minor changes, the first edition, distributed by the I.H.E.S. in the form of mimeographed exposés. The corrections made concern essentially printing errors, except in Exposés I and III, where slight modifications were made by L. Illusie to a few statements which were incorrect in the primitive version. Footnotes have also been added, especially in Exposé XIV by A. Grothendieck, in order to point out recent progress on questions that were still open at the time of the seminar.

Finally, note that Exposé XI, by D. Ferrand, on determinant theory for perfect complexes, was not written up and therefore will not be published; to avoid errors in references, we have left unchanged the numbering of the following exposés.

June 1971

Introduction

The purpose of the present seminar is to develop a global theory of intersections, and a Riemann–Roch formula, for arbitrary schemes. The reader will find a description of the programme of the seminar in Exposé 0. The possibility in principle of proving a Riemann–Roch formula for general regular schemes, along the lines of the well-known Borel–Serre report of 1958, was already clear at that time, at least for a projective morphism. The extension to the general case is less evident; the programme in which it belongs, and which is partly carried out in this seminar, goes back to 1960. As was also the case for the duality theory of coherent sheaves, an essential tool for formulating a satisfactory theory is Verdier’s theory of derived categories; familiarity with it is indispensable for understanding the seminar.

Apart from this theory, the study of the seminar requires little more than a general knowledge of the foundations of algebraic geometry as set out in EGA, chapters I, II, III. We shall only occasionally refer to EGA IV, for certain facts concerning flat or smooth morphisms, which for the most part also appear in the first exposés of SGA 1. Finally, to develop some results with the full generality desirable for applications, we sometimes use the notion of a ringed site and of a ringed topos, for which we refer to SGA 4, Exposés I to IV. A reader unfamiliar with the language of sites and topoi may everywhere replace these objects by ordinary topological spaces, the objects of the topos then being replaced by open subsets of these spaces; nevertheless,

we recommend that the reader assimilate the language of topoi, which provides an extremely convenient unifying principle.

The fundamental notion for the theory presented here is that of a perfect complex of modules, together with its various variants, developed in Exposés I, II, III. It seems clear that these notions, also important in other contexts in algebraic geometry, notably for formulas of Lefschetz–Verdier type in cohomology with discrete coefficients (SGA 5) or coherent coefficients, will also have a role to play outside algebraic geometry, especially in the formulation of a complex analytic variant of the Riemann–Roch theorem presented here, or of suitable variants of the Atiyah–Singer index theorem. Some indications in this direction will be given in Exposé II. Unfortunately, at present there is still no statement, even heuristic, which would encompass these two theorems, the first of which remains conjectural for the moment. A new idea seems to be missing, just as one is also missing in algebraic geometry for proving Riemann–Roch beyond projective hypotheses (cf. Exposé XIV, no. 2).

We shall make no use in this seminar of the local theory of intersections, set out in J.-P. Serre, *Algèbre locale. Multiplicités* (Lecture Notes in Mathematics no. 11, Springer), and merely point out here that Serre’s course had an evident influence on the genesis of the theory in 1957. Nor shall we use the theory of the Chow ring developed in the Chevalley Seminar of 1958, Exposés 2 and 3 (École Normale Supérieure). That theory is essentially tied to nonsingularity hypotheses, whereas the aim of our seminar is, on the contrary, to develop a theory of intersections on general schemes, and even on general locally ringed topoi. On this matter see the comments in Exposé XIV, nos. 4 and 8, which give the relations between our theory and that of the Chow ring and raise the question of a generalization of the latter. Thus one may say that the seminar presents a coherent and self-contained theory of intersections, but one which is by no means exhaustive of the different points of view useful, indeed indispensable, in intersection theory; consequently it should be complemented by the cited exposés of Serre and Chevalley.

We have appended to the seminar, as an appendix to Exposé 0, Grothendieck’s 1957 report, “Classes of sheaves and the Riemann–Roch theorem”, which served as the basis for the Borel–Serre seminar at Princeton in the same year and for their already cited article. That report sketches two proofs of the Riemann–Roch theorem for nonsingular quasi-projective varieties. The first, valid for the time being only in characteristic zero but giving, in return, a slightly more precise result in the case of an immersion, has not yet appeared in the literature, apart from the work of Borel–Serre and the present seminar. The report presupposes, of course, none of the other exposés of the seminar, and can even serve, like Exposé 0, as an introduction to the study of the latter, especially for a reader not yet familiar with Borel–Serre. The proof just alluded to applies essentially unchanged to the case of a projective morphism of complex analytic spaces, and will doubtless also apply in arbitrary characteristic once the problem of the operations “exterior powers” in the derived category is solved (cf. Exposé XIV, nos. 1, 2, 3). The absence of a study of this question is no doubt the most serious gap in this seminar, from the point of view we have adopted.

One will notice the absence, in the table of contents of the present seminar, of two of the participants mentioned on the cover page. One of them, J.-P. Jouanolou, took an active part in the technical elaboration of the first part of the seminar, but was prevented from taking part in the oral exposés; in particular, he is responsible for the linear-independence assertion contained in the important statement VI 1.1, giving the structure of the ring K° of classes of locally free modules on a projective bundle, which had previously been proved only under the assumption that the base scheme has an ample invertible module. The other, J.-P. Serre, gave two oral exposés which were logically independent of the rest of the seminar, and therefore preferred to publish them in the form of a separate article (cf. J.-P. Serre, “Grothendieck groups of split reductive group schemes”, to appear in *Publications Mathématiques*, no. 34).

As in the other parts of the *Séminaire de Géométrie Algébrique du Bois Marie*, the sigla SGA 1 to SGA 6 refer to the different parts of that seminar; the Roman numeral following the siglum indicates the number of the exposé, and the Arabic numerals following it correspond to the internal numbering of the exposé in question. For internal references within the present seminar the siglum SGA 6 is omitted, and references of the form I 4.9 are used, meaning: Exposé I, statement 4.9. The siglum EGA refers to the *Éléments de Géométrie Algébrique* of J. Dieudonné and A. Grothendieck.

Table of contents

Exposé	Title and author	Page
0	Outline of a programme for a theory of intersections on general schemes, by A. Grothendieck	1
Appendix	Classes of sheaves and the Riemann–Roch theorem, by A. Grothendieck	20
I	Generalities on finiteness conditions in derived categories, by L. Illusie	56
II	Existence of global resolutions, by L. Illusie	160
III	Relative finiteness conditions, by L. Illusie	190
IV	Grothendieck groups of ringed topoi, by L. Illusie	222
V	Generalities on λ -rings, by P. Berthelot	297
VI	The K° of a projective bundle: computations and consequences, by P. Berthelot	365
VII	Regular immersions and the computation of K° of a blow-up, by P. Berthelot	416
VIII	The Riemann–Roch theorem, by P. Berthelot	466
IX	Some computations of $K_\bullet$ -groups, by P. Berthelot	498
X	Formalism of intersections on proper algebraic schemes, by O. Jussila, with an appendix by A. Grothendieck, “Specialization in intersection theory”	519
XI	Not written up	–
XII	A relative representability theorem for the Picard functor, by M. Raynaud, written by S. Kleiman	595
XIII	Finiteness theorems for the Picard functor, by S. Kleiman	616
XIV	Open problems in intersection theory, by A. Grothendieck	667

Exposé 0

Outline of a programme for a theory of intersections on general schemes

by A. Grothendieck

The present exposé is introductory in nature, and its reading is not logically indispensable for the study of the seminar. It is addressed more particularly to readers familiar with the provisional version of the Riemann–Roch theorem, as it is presented in the Borel–Serre report or in Grothendieck’s report already mentioned in the introduction, cited below as [RRR], and reproduced as an appendix at the end of the present exposé.

1. Reminder on the Riemann–Roch formula

Recall the Riemann–Roch formula for a proper morphism

$$f : X \longrightarrow Y$$

of quasi-projective smooth schemes over a field k , and a coherent sheaf F on X , whose class in the group $K(X)$ of classes of coherent sheaves on X is denoted by $\mathrm{cl}(F)$:

$$\mathrm{Todd}(T_Y) \, \mathrm{ch}_Y(f_!(\mathrm{cl}(F))) = f_*(\mathrm{Todd}(T_X) \, \mathrm{ch}_X(F)). \quad (1.1)$$

This formula is valid in $A(Y) \otimes \mathbb{Q}$, where $A(Y)$ is the Chow ring of Y . The f_* on the right hand side is obtained by tensoring with $\mathbb{Q}$ the homomorphism “direct image of cycles”

$$f_* : A(X) \longrightarrow A(Y),$$

while the term on the left is the Euler–Poincaré characteristic of F relative to f :

$$f_!(\mathrm{cl}(F)) = \sum_i (-1)^i \mathrm{cl}(R^i f_*(F)).$$

The symbols ch_X , ch_Y denote the Chern characters on X and on Y , and T_X , T_Y the tangent bundles of X and Y . As is known, $\mathrm{Todd}(-)$ and $\mathrm{ch}(-)$ are universal polynomials with coefficients in $\mathbb{Q}$ in the Chern classes of the argument. Since the constant term of $\mathrm{Todd}(-)$ is 1, it is an invertible element for every value of the argument. Thus, after multiplying by $\mathrm{Todd}(T_Y)^{-1}$, formula (1.1) takes the equivalent form, more convenient for our purposes,

$$\mathrm{ch}_Y(f_!(\mathrm{cl}(F))) = f_*(\mathrm{Todd}(T_f) \, \mathrm{ch}_X(F)), \quad (1.2)$$

where

$$T_f = T_X - f^* T_Y \in K(X). \quad (1.3)$$

Thus T_f plays the role of a virtual relative tangent bundle of X over Y . When f is smooth, one simply has $T_f = T_{X/Y}$, whereas when $f : X \rightarrow Y$ is an immersion, one obtains

$$T_f = -\check{N}_{X/Y},$$

where $N_{X/Y}$ denotes the normal sheaf of X in Y .

One of the main aims of the seminar is to generalize (1.2) simultaneously in two directions:

- a) to get rid of the hypothesis that a base field k exists;
- b) to replace the regularity hypotheses on Y, X by a hypothesis of “local regularity” on f .

Finally, along the way, we shall also deal with the following problem:

- c) to eliminate the quasi-projectivity hypotheses which, in the absence of a base field, are expressed by the existence of ample invertible modules on X and Y .

2. Removing the base field

Let us first examine generalization a), while retaining the hypotheses of regularity and of the existence of ample invertible modules on X and on Y .

The definitions of $K(X)$, $K(Y)$ and of the homomorphism $f_! : K(X) \rightarrow K(Y)$ then present no new problem, thanks to the fact that X and Y are regular. The most natural way to give

a meaning to (1.2) would seem to consist in defining Chow rings $A(X), A(Y)$ and a group homomorphism

$$f_* : A(X) \longrightarrow A(Y),$$

in establishing a theory of Chern classes, giving maps

$$c_i : K(X) \longrightarrow A(X),$$

and similarly for Y , and finally in describing an element, the virtual relative tangent bundle,

$$T_f \in K(X).$$

2.1

For the latter, a definition presents itself rather evidently. One uses the fact that the morphism f , thanks to the hypothesis that ample invertible modules exist on X and on Y , may be factored as

$$X \xrightarrow{i} X' \xrightarrow{f'} Y,$$

where i is a closed immersion and f' is smooth; for example, one may take $X' = \mathbb{P}_Y^r$ for suitable r . One then sets

$$T_f = i^* T_{X'/Y} - \check{N}_{X/X'} \in K(X), \quad (2.1)$$

where $\Omega_{X'/Y}^1$ is the locally free module of relative 1-differentials of X' over Y , or the relative cotangent module of X' over Y , and $N_{X/X'}$ is the conormal module of X in X' , by definition equal to J/J^2 , where J is the ideal on X' defining X . The regularity hypotheses on X and X' also imply that $N_{X/X'}$ is locally free. Finally, $\check{N}$ denotes the dual of N . One verifies without difficulty, in Exposé VIII, that the element T_f defined by (2.1) is independent of the chosen factorization of f .

2.2

As for a definition, under the conditions considered, of a Chow ring and a corresponding theory of Chern classes, although it is plausible that such a theory should be developable on the model of smooth varieties over a field, it seems that this work has not been done at the present time. We shall therefore adopt another viewpoint, by directly defining a ring which replaces the Chow ring, and whose tensor product with $\mathbb{Q}$ is indeed isomorphic to $A(X) \otimes \mathbb{Q}$ in the classical case.

A first natural filtration of $K(X)$, the one considered in [RRR], is the topological filtration by the codimension of the support of coherent sheaves: $\text{Fil}^i K(X)$ is generated by the $\text{cl}(F)$ with F coherent and $\text{codim}(\text{Supp } F, X) \geq i$. A first question is the compatibility of this filtration with the multiplicative structure of $K(X)$. In [RRR], this compatibility is obtained, for X smooth over a field, by means of Chow's moving lemma. Assuming this assertion, one obtains a graded ring $\text{Gr}_{\text{top}}(X)$ and a canonical surjective homomorphism

$$\varphi : A(X) \longrightarrow \text{Gr}_{\text{top}}(X),$$

sending the class of a prime cycle Z to $\text{cl}(\mathcal{O}_Z)$; it then turns out, Exposé XIV 4.2, that the kernel is a torsion subgroup. Moreover, [RRR] shows how the images in $\text{Gr}_{\text{top}}(X)$ of the Chern classes $c_i(F)$ can be computed directly in terms of $\text{cl}(F) \in K(X)$ and the exterior-power operations λ^i in $K(X)$.

This shows that, for a noetherian regular scheme X endowed with an ample invertible module, the ring $\text{Gr}_{\text{top}}(X)$ is a convenient substitute for the hypothetical Chow ring $A(X)$. It also has

the advantage over the latter that certain computations of [RRR], leading to important formulas in this ring, are performed directly in it—formulas which, even for X projective and smooth over a field, have not at present been established in $A(X)$ itself (Exp. XIV, 4.3 and 6.4). To justify this viewpoint completely, however, one would have to solve the problem mentioned above of the compatibility of the topological filtration of $K(X)$ with its ring structure, a problem which seems essentially connected with Chow's moving lemma, itself the essential technical ingredient of the theory of the Chow ring which we have claimed to be able to avoid.

2.3

To avoid this problem, one endows $K(X)$ with a second filtration, finer than the topological filtration, and describable in purely algebraic terms from the structure of $K(X)$ as an augmented λ -ring. The definition of this filtration is suggested by the expression given in [RRR] for the Chern classes of an element $x \in K(X)$ in $\mathrm{Gr}_{\mathrm{top}}(X)$:

$$c_i(x) = \gamma^i(x - \epsilon(x)) \mod \mathrm{Fil}^{i+1} K(X), \quad (2.3)$$

where γ^i denotes a certain linear combination of the λ^j ($j \leq i$), made explicit in loc. cit., and where ϵ is the augmentation. One shows that $\gamma^i(x - \epsilon(x))$ is indeed of topological filtration at least i , which gives meaning to formula (2.3). One then defines the λ -filtration of $K(X)$ by the ideals $\mathrm{Fil}_\lambda^i K(X)$, where $\mathrm{Fil}_\lambda^i K(X)$ consists of finite sums of expressions of the form

$$\gamma^{i_1}(x_1 - \epsilon(x_1)) \cdots \gamma^{i_r}(x_r - \epsilon(x_r)), \quad i_1 + \cdots + i_r \geq i.$$

The associated graded ring for this filtration will be denoted $\mathrm{Gr}^\bullet(X)$; it is this ring which will replace the Chow ring of X in the seminar. In the case under consideration here, one can moreover show that the canonical homomorphism

$$\mathrm{Gr}^\bullet(X) \longrightarrow \mathrm{Gr}_{\mathrm{top}}(X) \quad (2.4)$$

is an isomorphism modulo torsion, justifying the viewpoint that, after tensoring with $\mathbb{Q}$, this ring can play exactly the same role as the Chow ring tensored with $\mathbb{Q}$. On the other hand, exactly what was needed has been done so that formula (2.3), with $\mathrm{Fil}_{\mathrm{top}}$ replaced by Fil_λ , defines a theory of Chern classes with values in $\mathrm{Gr}^\bullet(X)$; the Chern classes in $\mathrm{Gr}_{\mathrm{top}}(X)$ are then deduced by means of the homomorphism (2.4). It is shown in Exposé V, using the fact proved in Exposé VI that $K(X)$ is a special λ -ring in the sense of [RRR], that is, a λ -ring in the sense of Exposé V, that this notion of Chern class satisfies all the usual formal properties reviewed in [RRR].

2.4

One should note a very appreciable advantage of the definition of $\mathrm{Gr}^\bullet(X)$ over that of $A(X)$ and $\mathrm{Gr}_{\mathrm{top}}(X)$: it remains meaningful for an absolutely arbitrary scheme X , and more generally for an arbitrary locally ringed topos X . One then takes for $K(X)$ the group of classes constructed from locally free modules on X , which is again an augmented λ -ring, hence equipped with a corresponding filtration and an associated graded $\mathrm{Gr}^\bullet(X)$. The price of this generality, however, is that when one is not willing to neglect torsion phenomena, i.e. to tensor with $\mathbb{Q}$, the properties of $\mathrm{Gr}^\bullet(X)$ are in several respects less satisfactory than those of its two competitors (Exp. XIV no. 4). Thus, assuming again that $f : X \rightarrow Y$ is a proper morphism of noetherian regular schemes endowed with ample invertible modules, it is not true in general that the homomorphism

$$f_* : K(X) \longrightarrow K(Y)$$

maps $\mathrm{Fil}_\lambda^i K(X)$ into $\mathrm{Fil}_\lambda^{i-d} K(Y)$, where d is the relative dimension of X over Y , and hence defines, by passage to associated graded, a degree $-d$ homomorphism from $\mathrm{Gr}^\bullet(X)$ to $\mathrm{Gr}^\bullet(Y)$. However, Exposé VII proves that this is true after tensoring with $\mathbb{Q}$, so that one obtains a degree $-d$ homomorphism

$$f_* : \mathrm{Gr}^\bullet(X) \otimes \mathbb{Q} \longrightarrow \mathrm{Gr}^\bullet(Y) \otimes \mathbb{Q}, \quad (2.5)$$

which plays the role of the homomorphism “direct image of cycles” $A(X) \rightarrow A(Y)$.

Having said this, we see that we possess all the structural elements needed to give meaning to formula (1.2) when the existence of a base field is not postulated, with $f : X \rightarrow Y$ a proper morphism of noetherian regular schemes admitting ample invertible modules.

3. The case of general schemes

We now show how one can still give meaning to formula (1.2) when the regularity hypothesis on X, Y is abandoned and replaced by a local hypothesis on the morphism f , to be specified below.

3.1

A first difficulty comes from the fact that the observation which served as the starting point of the theory [RRR], namely that the group $K(X)$ of classes of sheaves on X is the same whether it is defined in terms of coherent sheaves or in terms of locally free sheaves, is essentially tied to the regularity hypothesis on X . For a general scheme X , say noetherian for simplicity, one must distinguish between these two groups; we shall denote them respectively by $K_\bullet(X)$ and $K^\bullet(X)$. The first is covariant in X for proper morphisms, the second contravariant in X for arbitrary morphisms. Notice that $K^\bullet(X)$ is naturally endowed with a ring structure, indeed with an augmented λ -ring structure, whereas $K_\bullet(X)$ no longer has, in general, a ring structure, but only a module structure over $K^\bullet(X)$. The homomorphism $x \mapsto x \cdot \mathrm{cl}(\mathcal{O}_X)$ from $K^\bullet(X)$ to $K_\bullet(X)$ is no longer an isomorphism in general. For our formulation of the generalized Riemann–Roch theorem, we shall work exclusively with $K^\bullet(X)$ and $K^\bullet(Y)$, to the exclusion of $K_\bullet(X)$ and $K_\bullet(Y)$. As noted in 2.3, these augmented λ -rings, equipped with their natural λ -filtration, give rise to graded rings $\mathrm{Gr}^\bullet(X)$, $\mathrm{Gr}^\bullet(Y)$, and to Chern-class homomorphisms

$$c_i : K^\bullet(X) \longrightarrow \mathrm{Gr}^\bullet(X),$$

and similarly for Y . The difficulty, however, lies in defining a direct-image homomorphism

$$f_* : K^\bullet(X) \longrightarrow K^\bullet(Y), \quad (3.1)$$

which can no longer be defined by the naive formula

$$f_*(\mathrm{cl}(F)) = \sum_i (-1)^i \mathrm{cl}(R^i f_*(F)),$$

because even if F is locally free, the modules $R^i f_*(F)$ are in general no longer locally free, nor even of finite Tor-dimension, and the preceding formula loses its meaning. A solution to this difficulty is furnished by the viewpoint of derived categories. Indeed, whereas the $R^i f_*(F)$ are in general “bad” sheaves, say of infinite Tor-dimension, the complex $Rf_*(F)$ on Y , the direct image of F in the sense of derived categories, tends to be excellent. More precisely, when F has finite Tor-dimension over Y , for instance when f has finite Tor-dimension and F is locally free on X , then $Rf_*(F)$ is perfect on Y , that is, has coherent cohomology and globally finite Tor-dimension (Exp. III). It follows easily, Exp. IV, that its class in the Grothendieck group of perfect complexes has a meaning in $K^\bullet(Y)$.

3.2

To give a meaning to (1.2), it remains to impose on f a hypothesis making it possible to give a meaning to the homomorphism (2.5), i.e. ensuring that the homomorphism (3.1) just defined is, up to a shift and modulo torsion, compatible with the filtrations of these rings; and finally one must again define an element, the “virtual relative tangent bundle”,

$$T_f \in K^\bullet(X).$$

For the latter, returning to the definition proposed in 2.1, one sees that a hypothesis is needed ensuring that, with the notation of loc. cit., the conormal module $N_{X/X'}$ is locally free. The most natural hypothesis to this end is that i be a “regular immersion”, i.e. identify X with a subscheme of X' defined by an ideal which locally can be defined by an $\mathcal{O}_{X'}$ -regular sequence of sections of $\mathcal{O}_{X'}$. This hypothesis, purely local in nature, in fact does not depend on the choice of the factorization of f into an immersion followed by a smooth morphism (Exp. VIII), and, as above, the same is true for the class T_f defined by formula (2.1). When the hypothesis just considered is satisfied, the morphism f is called a “locally complete intersection” morphism. Such a morphism is also locally of finite Tor-dimension.

We shall also allow ourselves to omit the word “locally” in this terminology, which cannot cause any confusion.

If f is a locally complete intersection morphism, one introduces a natural notion of “virtual relative dimension” $d(f)$ by the formula

$$d(f)(x) = \dim_x(f^{-1}(f(x))) - \operatorname{codim}_x(X, X').$$

It is an integer-valued function, locally constant at the point $x \in X$, hence constant if X is connected; moreover it does not depend on the choice of the factorization (2.1). When this virtual relative dimension d is constant, when f is proper, and when X and Y admit ample invertible modules, one can prove (Exposé VIII), although this is far from a trivial result, that

$$f_*(\operatorname{Fil}^i K^\bullet(X)_{\mathbb{Q}}) \subset \operatorname{Fil}^{i-d} K^\bullet(Y)_{\mathbb{Q}}, \quad (3.4)$$

whence a homomorphism (3.3) of degree $-d$.

3.3

We have therefore assembled all the structural elements needed to give a meaning to formula (1.2) in the case of a proper locally complete intersection morphism of noetherian schemes X, Y , both admitting ample invertible modules. With a little additional care, this formulation in fact retains a meaning independently of any noetherian hypothesis.

In this general form, this Riemann–Roch formula will be proved in Exposé VIII. The reader will observe that the proof given in this seminar combines elements of the two proofs which appear in [RRR] and which were discussed in the Introduction. The need to resort to the methods of the first of these proofs, involving more K -formalism than the second, followed in Borel–Serre, is due to the need for a proof of the inclusion (3.4), a question which did not arise in the more restricted framework of [RRR] and Borel–Serre [2], where one could work with the Chow ring or the ring $\operatorname{Gr}_{\operatorname{top}}^\bullet$. From the second proof, we retain the introduction of the scheme obtained by blowing up Y along X in the case where f is an immersion. This technical artifice will no doubt disappear, and the proof in the case of an immersion will extend to the case where one no longer postulates the existence of an ample invertible module on Y , once the question raised in Exposé XIV, no. 1, concerning the introduction of exterior-power operations in the derived category $D^-(Y)$ has been resolved.

4. Removing the ample invertible module hypothesis

We must finally give a few indications of how one can still give a meaning to (1.2) in the absence of a hypothesis asserting the existence of ample invertible modules on X and Y .

4.1

First note that for an arbitrary scheme X , or more generally an arbitrary locally ringed topos, it seems beyond doubt that the “right” invariant which should replace the ring $K(X)$ of [RRR] is not the ring $K^\bullet(X)$ considered above, defined in terms of locally free modules on X , but rather the group $K(\text{Parf}(X))$ considered in 3.1, defined in terms of perfect complexes on X . We shall denote this group here by $K^\bullet(X)$, and distinguish it from the “naive” group of 3.1, denoted $K_{\text{naive}}^\bullet(X)$. This point of view is forced by the need to define a direct-image homomorphism (3.1) for a proper morphism

$$f : X \longrightarrow Y$$

of finite Tor-dimension, say between noetherian schemes; the non-noetherian case requires somewhat stronger local finiteness properties on f , cf. Exposé III. With the definition of $K^\bullet(X)$ now adopted, this homomorphism f_* is indeed defined in a trivial way by formula (3.2), where now F is an arbitrary perfect complex on X , which implies that $Rf_*(F)$ is also perfect.

4.2

This new definition of $K^\bullet(X)$ raises, however, a new problem in homological algebra: that of defining a λ -structure on $K^\bullet(X)$ compatible with its natural ring structure coming from tensor product in the derived category. If L and L' are two perfect complexes, then so is their total tensor product $L \otimes^{\mathbf{L}} L'$. More precisely, one would need to define “exterior-power” operations in the derived category $D^-(X)$, transforming perfect complexes into perfect complexes. This question has not yet been clarified at present, and, as was said in the Introduction, it constitutes the most serious gap in the seminar. Once this question is solved, and proceeding as in 2.3, one obtains a natural filtration on $K^\bullet(X)$, allowing one to form the associated graded ring $\text{Gr}^\bullet(X)$, which will play the role of the Chow ring, and to define a theory of Chern classes

$$c_i : K^\bullet(X) \longrightarrow \text{Gr}^i(X).$$

4.3

To give a meaning to formula (1.2) for a proper locally complete intersection morphism

$$f : X \longrightarrow Y$$

of noetherian schemes, one must still include in the statement of this formula the nontrivial inclusions (3.4), where d is the virtual relative dimension, in order to be able to define (3.3); and finally one must define the virtual relative tangent class

$$T_f \in K^\bullet(X).$$

When f admits a factorization (2.1) through an immersion into a relative smooth scheme, the definition of T_f presents no new difficulty. It can moreover be reformulated in that case, in a way more consonant with the spirit of derived categories, by considering the well-known canonical homomorphism

$$N_{X/X'} \longrightarrow \Omega_{X'/Y}^1 \otimes_{\mathcal{O}_{X'}} \mathcal{O}_X \tag{4.1}$$

induced by the exterior differential, and the chain complex $L_{X/Y}$ defined from (4.1) by “extending by zeros”, with the complex equal in degree 1, respectively 0, to the source, respectively the target, of the homomorphism (4.1). Formula (2.2) then takes the more satisfactory form

$$T_f = \text{cl}(L_{X/Y}^\vee), \quad (4.2)$$

a formula which makes sense in $K^\bullet(X)$ when f is a locally complete intersection morphism, since in that case the complex $L_{X/Y}$ is manifestly perfect. The assertion that T_f is invariant relative to the chosen factorization of f can then be made precise as an invariance assertion for the complex $L_{X/Y}$: up to canonical isomorphism in the derived category $D(X)$, this complex is likewise independent of the chosen factorization (Exposé VIII).

4.4

When $f : X \rightarrow Y$ is an arbitrary morphism of schemes, one can still construct a canonical object $L_{X/Y}$ of the derived category $D(X)$, defined up to unique isomorphism, which on each affine open of X lying above an affine open of Y is isomorphic to the complex constructed according to the principle explained in the preceding paragraph. In the general case, however, “smooth” should be replaced by “formally smooth”, so that every algebra B over a ring A occurs as a quotient of a formally smooth algebra, for example a polynomial algebra. For such a definition it is not possible to proceed by simple gluing from the affine constructions, since the objects of the derived category $D(X)$ are, essentially, not glueable.

Here is a general construction of $L_{X/Y}$, valid more generally for every morphism

$$f : X \longrightarrow Y$$

of commutatively ringed topoi. Let $B = \mathcal{O}_X$ and $A = f^{-1}\mathcal{O}_Y$, so that A is a ring on the space X and B is an A -algebra. Let $T \rightarrow B$ be a homomorphism of sheaves of sets which generates B as an A -algebra, i.e. such that the corresponding homomorphism of A -algebras

$$C = A[T] \longrightarrow B$$

is an epimorphism of sheaves of sets; here $A[T]$ denotes the commutative A -algebra freely generated by T , that is, the sheaf associated with the presheaf

$$U \longmapsto A(U)[T(U)].$$

Let J be the kernel of $C \rightarrow B$, and consider, as in 4.3, the chain complex of length 1 defined by the natural homomorphism

$$J/J^2 \longrightarrow \Omega_{C/A}^1 \otimes_C B.$$

It is not difficult to see that, up to canonical isomorphism in the category $D(X)$, the complex so constructed is independent of the choice of T and of $T \rightarrow B$, so that it deserves the designation of the relative cotangent complex $L_{X/Y}$ of X over Y . One also verifies that, on every affine open of X over an affine open of Y , it is given, up to canonical isomorphism, by the procedure of 4.3.

In the case where f is locally complete intersection, one sees moreover that the result is a perfect complex, which makes it possible again to define an element T_f of $K^\bullet(X)$ by formula (4.2). Here again, in general, it would not have been possible to define T_f as an element of $K_{\text{naive}}^\bullet(X)$, and nothing permits one to suppose that T_f is globally isomorphic in $D(X)$ to a bounded complex with locally free components of finite type. This is therefore an additional justification for the point of view advanced in 4.1 in favour of the “sophisticated” $K^\bullet(X)$, in preference to $K_{\text{naive}}^\bullet(X)$.

4.5

Subject to the question raised in 4.2, we have therefore again assembled all the structural elements needed to give a meaning to (1.2) for a proper locally complete intersection morphism of arbitrary schemes; the noetherian hypothesis implicitly made in the preceding sections can in fact be eliminated without any essential difficulty. Unfortunately, this does not mean that, even subject to this reservation, the formula in question is established, even in the case where Y is the spectrum of an algebraically closed field of characteristic $p > 0$ and X is, say, a proper smooth algebraic scheme of dimension 3. See the comments in Exposé XIV, no. 2.

5. Analytic variant

In the form presented in no. 4, the Riemann–Roch formula (1.2) also has a meaning for a proper locally complete intersection morphism of complex analytic spaces, or of rigid analytic spaces in the sense of Tate. But of course the proof given in [RRR] applies to this case only when one assumes in addition that the morphism f is projective; and, as in the algebraic case, it seems that an essentially new idea is needed to treat the general case. The formula we have in view here is a formula with values in an “analytic Chow ring” $\mathrm{Gr}^\bullet(X)$, or rather $\mathrm{Gr}^\bullet(X)_\mathbb{Q}$, constructed according to the principle explained in 4.2, the difficulty encountered there disappearing, moreover, because one is in characteristic zero, as is noted in Exposé XIV, no. 1.

When X and Y are nonsingular, it is possible to give another version of the Riemann–Roch formula by proceeding as follows. First let X be an arbitrary complex analytic space, and let X^{top} be the same space equipped with the sheaf of rings of continuous complex-valued functions. We therefore have a canonical morphism of ringed spaces

$$X^{\mathrm{top}} \longrightarrow X,$$

which, by the evident contravariant nature of the functor $K^\bullet(X)$ in the variable ringed space X , defines a homomorphism of rings

$$K^\bullet(X) \longrightarrow K^\bullet(X^{\mathrm{top}}). \quad (5.1)$$

For simplicity suppose that X^{top} is compact; then, as in the case of a scheme admitting an ample invertible module, one proves (Exposé II) that every perfect complex on X^{top} is globally isomorphic to a bounded complex, locally free in each degree, i.e. to a complex of complex vector bundles on X^{top} , and hence that $K^\bullet(X^{\mathrm{top}})$ is canonically isomorphic to the well-known group $K(X^{\mathrm{top}})$ of topologists [1], defined in terms of complex vector bundles on the compact space X^{top} . Thus one may interpret (5.1) as a homomorphism from the $K^\bullet(X)$ defined in terms of the analytic structure to the well-known ring $K(X^{\mathrm{top}})$ of topologists. If now

$$f : X \longrightarrow Y$$

is a morphism of compact nonsingular analytic spaces, one knows how to associate to it, thanks to Atiyah–Hirzebruch [1], a group homomorphism

$$f_*^{\mathrm{top}} : K^\bullet(X^{\mathrm{top}}) \longrightarrow K^\bullet(Y^{\mathrm{top}}). \quad (5.2)$$

Using likewise the homomorphism f_* of (3.1), defined by formula (3.2), which here implicitly uses Grauert’s finiteness theorem [4] for a proper morphism of analytic spaces, one obtains a square of natural homomorphisms involving only groups of type $K^\bullet$, to the exclusion of rings of Chow-ring or cohomology type:

$$\begin{array}{ccc} K^\bullet(X) & \longrightarrow & K^\bullet(X^{\mathrm{top}}) \\ \downarrow f_* & & \downarrow f_*^{\mathrm{top}} \\ K^\bullet(Y) & \longrightarrow & K^\bullet(Y^{\mathrm{top}}). \end{array} \quad (5.3)$$

The Riemann–Roch type formula to which we are alluding consists in asserting the commutativity of this square. Note the difference in nature between this assertion, which does not neglect torsion phenomena and is situated in a group $K^\bullet(Y^{\text{top}})$ of an essentially discrete nature, and the Riemann–Roch formula of the type considered in no. 4, which is situated in a group of “Chow” type no longer of a discrete nature, but which, on the other hand, must be tensored with $\mathbb{Q}$. Let us note that formula (5.3) is proved at any rate when X and Y are nonsingular projective varieties [7], as is the formula of the type considered in no. 4, proved in this case in [RRR]. What is involved here is a plausible variant of the Atiyah–Singer index theorem, and of its generalizations due to Shih [7] and Atiyah, which obviously calls for a common generalization. Once the commutativity of (5.3) is proved, one would immediately deduce from it, using the “differentiable Riemann–Roch theorem” of Atiyah–Hirzebruch [1], a cohomological variant of the complex analytic Riemann–Roch formula, namely the commutativity of the diagram

$$\begin{array}{ccc} K^\bullet(X) & \xrightarrow{x \mapsto \text{ch}(x) \text{ Todd}(T_X)} & H^{2\bullet}(X, \mathbb{Q}) \\ \downarrow f_* & & \downarrow f_* \\ K^\bullet(Y) & \xrightarrow{y \mapsto \text{ch}(y) \text{ Todd}(T_Y)} & H^{2\bullet}(Y, \mathbb{Q}), \end{array} \quad (5.4)$$

where the horizontal arrows are deduced from cohomological Chern classes, themselves deduced from the homomorphism (5.1) and from the ordinary Chern-class homomorphism

$$K^\bullet(X^{\text{top}}) \longrightarrow H^{2\bullet}(X, \mathbb{Z}),$$

the first vertical arrow is the same as in (5.3), and the second is the Gysin homomorphism in rational cohomology.

6. The covariant invariant $K_\bullet(X)$

In all that precedes, scarcely anything has been said except about the contravariant invariant $K^\bullet(X)$, to the exclusion of the covariant invariant $K_\bullet(X)$ whose existence was indicated in 3.1. If one regards $K^\bullet(X)$ as giving an analogue of the integral cohomology ring of a topological space, one should regard $K_\bullet(X)$ as giving an analogue of integral homology. The module structure of $K_\bullet(X)$ over $K^\bullet(X)$ then corresponds to the cap-product operation. The fact that, when X is regular, the natural homomorphism $K^\bullet(X) \rightarrow K_\bullet(X)$ is an isomorphism should be regarded as the analogue of the Poincaré duality theorem for oriented topological varieties, asserting that cap product with the fundamental homology class defines an isomorphism from cohomology to homology. These analogies moreover suggest the usefulness of also defining, for a finite polyhedron X , say, a covariant invariant $K_\bullet(X)$ whose relations to integral homology – spectral sequence, Chern homomorphism, and so on – would be analogous to those existing between $K^\bullet(X)$ and integral cohomology. Nor does it seem excluded that there might exist, in terms of these groups $K_\bullet(X)$, variants of the differentiable Riemann–Roch theorem valid for spaces more general than compact differentiable manifolds.

Returning to the case where X is an arbitrary noetherian scheme, one should endow $K_\bullet(X)$ with an increasing filtration, $\text{Fil}_i K_\bullet(X)$ being generated by classes $\text{cl}_\bullet(F)$, with F a coherent sheaf on X such that $\dim \text{Supp } F \leq i$. One will prove (Exposé X) the compatibility of this filtration with the module structure and with the filtration of $K^\bullet(X)$, i.e. the formula

$$\text{Fil}_i K^\bullet(X) \cdot \text{Fil}_j K_\bullet(X) \subset \text{Fil}_{j-i} K_\bullet(X), \quad (6.1)$$

which allows the graded group associated with the filtered module $K_\bullet(X)$, denoted $\text{Gr}_\bullet(X)$, to be endowed with a module structure over $\text{Gr}^\bullet(X)$. Its elements may in fact be interpreted as

classes of algebraic cycles on X , for an equivalence relation less fine than rational equivalence when X is of finite type over a base field k . If

$$f : X \longrightarrow Y$$

is a proper morphism of noetherian schemes, then

$$f_* : K_\bullet(X) \longrightarrow K_\bullet(Y) \quad (6.2)$$

is compatible with the filtrations and defines a homomorphism of graded groups

$$f_* : \mathrm{Gr}_\bullet(X) \longrightarrow \mathrm{Gr}_\bullet(Y), \quad (6.3)$$

giving rise to a “projection formula”, i.e. a formula which says that it is $\mathrm{Gr}^\bullet(Y)$ -linear; this formula is obtained by passage to associated gradeds from the analogous projection formula for the homomorphism (6.2), which is $K^\bullet(Y)$ -linear.

From the point of view proposed in the seminar, a manageable theory of intersections on noetherian schemes X , not necessarily regular, should consist in the combined study of the contravariant invariants $K^\bullet(X), \mathrm{Gr}^\bullet(X)$ and the covariant invariants $K_\bullet(X), \mathrm{Gr}_\bullet(X)$. These play roles of essentially different nature, and their properties complement each other. Thus the covariant groups $K_\bullet(X), \mathrm{Gr}_\bullet(X)$ are easier to compute for certain nonproper fibrations, thanks to the “homotopy exact sequence” which one can establish for them ([RRR] and Exposé IX). For example, one proves (Exposé IX) canonical isomorphisms such as

$$K_\bullet(X[t]) \simeq K_\bullet(X), \quad \mathrm{Gr}_\bullet(X[t]) \simeq \mathrm{Gr}_\bullet(X),$$

which fail in general for $K^\bullet$ when X is a nonregular scheme. On the other hand, $K^\bullet(X)$ and $\mathrm{Gr}^\bullet(X)$ lend themselves well to computations, thanks to their ring structures.

When one considers schemes X proper over a field, the projection

$$\pi_X : X \longrightarrow (\text{point})$$

defines a homomorphism, the “Euler–Poincaré characteristic”,

$$\chi_X = \pi_{X*} : K_\bullet(X) \longrightarrow K_\bullet(\text{point}) = \mathbb{Z}, \quad (6.4)$$

which induces on the subgroup $\mathrm{Gr}_0(X) = \mathrm{Fil}_0 K_\bullet(X)$ the homomorphism “degree of zero-cycles”

$$\deg_X : \mathrm{Gr}_0(X) \longrightarrow \mathbb{Z}. \quad (6.5)$$

The multiplication homomorphism

$$\mathrm{Gr}^i(X) \times \mathrm{Gr}_i(X) \longrightarrow \mathrm{Gr}_0(X),$$

composed with the degree homomorphism (6.5), then furnishes a pairing

$$\mathrm{Gr}^i(X) \times \mathrm{Gr}_i(X) \longrightarrow \mathbb{Z}, \quad (6.6)$$

which explains that, in certain respects, $\mathrm{Gr}^\bullet(X)$ and $\mathrm{Gr}_\bullet(X)$ play dual roles, just as cohomology and homology do. Thus, if $f : X \rightarrow Y$ is a morphism of proper algebraic schemes, the two homomorphisms

$$f^* : \mathrm{Gr}^\bullet(Y) \longrightarrow \mathrm{Gr}^\bullet(X), \quad f_* : \mathrm{Gr}_\bullet(X) \longrightarrow \mathrm{Gr}_\bullet(Y)$$

are formally transposes of one another, in the sense of the pairings (6.6).

The homomorphisms (6.4) and (6.5), and the pairing (6.6) deduced from them, may be regarded as the source of numerical relations in intersection theory on proper algebraic schemes over a given field k . This point of view will be developed in some detail in Exposé X, which is essentially independent of Exposés VI, VII, VIII, centred on the proof of the Riemann–Roch theorem, and of Exposé IX. Some notions concerning the behaviour of $K^\bullet(X)$ and $\mathrm{Gr}^\bullet(X)$ under specialization will also be given there.

7. Picard groups and the determinant functor

As in every intersection theory, it is important to specify the place occupied in the theory by the Picard group, or group of classes of divisors in the classical terminology. Subject to a slight restriction, automatically satisfied when X is for instance quasi-compact, one establishes in Exposé X a canonical isomorphism

$$c_1 : \text{Pic}(X) \xrightarrow{\sim} \text{Gr}^1(X), \quad (7.1)$$

which shows the exact role of the Picard group in the theory we are proposing. This role justifies a more detailed study of the Picard group in Exposés XII and XIII, which are logically independent of the text of the seminar. Kleiman, in Exposé XIII, studies numerical equivalence and certain finiteness theorems for the Picard group, announced without proof in [6], using purely numerical techniques, not relying on the theory of the Picard functor and its representability, due to himself and Mumford, and markedly simpler than Grothendieck's initial proofs. The only result refractory to Kleiman's methods, and which he is obliged to use, is [6, C-07 (i)]; its proof and certain refinements, including certain representability theorems for the Picard functor, occupy Exposé XII. The latter is moreover of a spirit and technique entirely foreign to the rest of the seminar, and is connected with [5]; this exposé is logically independent of all the others.

Finally, in Exposé XI, of a nature appreciably more general than XII and XIII and closer to the general spirit of the seminar, one studies, on an arbitrary locally ringed topos, a remarkable functor called the “determinant functor”:

$$\overset{\bullet}{\det} : \text{Parf}_{\text{is}}(X) \longrightarrow \text{Inv}(X) \quad (7.2)$$

from the category of perfect complexes on X , whose morphisms are only the isomorphisms in the sense of the derived category $D(X)$, to the category of invertible modules on X . This functor is described, up to very little, by the requirement that it be of “local nature” and that it be given, for a bounded complex L with locally free components, by the formula

$$\overset{\bullet}{\det}(L) = \bigotimes_i \det(L_i)^{(-1)^i}, \quad (7.3)$$

where $\det(L_i)$ denotes the exterior power of maximal degree of the locally free module L_i . Although this exposé is also logically independent of the rest of the seminar, except for Exposé I, it nevertheless constitutes an important element of the general “yoga” proposed in the seminar and developed essentially in Exposés I to XI.

Bibliography

- [1] M. Atiyah and F. Hirzebruch, *Vector bundles and homogeneous spaces*, Symposia in Pure Mathematics, vol. 3, Differential Geometry, pp. 7–38, 1961.
- [2] A. Borel and J.-P. Serre, *Le théorème de Riemann–Roch*, Bull. Soc. math. France, t. 86, pp. 97–136 (1958).
- [3] H. Grauert, *Ein Theorem der analytischen Garbentheorie*, Pub. Math. no. 5, pp. 5–64 (1960).
- [4] A. Grothendieck, *Classes de faisceaux et théorème de Riemann–Roch*, mimeographed notes, Princeton 1957, cited as [RRR] and reproduced as an appendix to Exposé 0 of the seminar.

- [5] A. Grothendieck, *Technique de descente et théorèmes d'existence en Géométrie Algébrique*, in *Fondements de la Géométrie Algébrique*, extracts from the Bourbaki Seminar 1957–1962, Secrétariat Mathématique, Paris.
- [6] W. Shih, Séminaire de Topologie à l'I.H.E.S., 1966/67.
- [7] M. Atiyah and F. Hirzebruch, *The Riemann–Roch theorem for analytic embeddings*, Topology, vol. 1, pp. 151–166 (1962).