

ELEMENTARY GLOBAL STUDY OF SOME CLASSES OF MORPHISMS (EGA II)

CONTENTS

Summary	2
1. Affine morphisms	3
1.1 S -preschemes and $\mathcal{O}_S$ -algebras	3
1.2 Affine preschemes over a prescheme	4
1.3 Affine preschemes over S associated to an $\mathcal{O}_S$ -algebra	5
1.4 Quasi-coherent sheaves over an affine prescheme over S	6
1.5 Change of base prescheme	8
1.6 Affine morphisms	9
1.7 Vector bundle associated with a sheaf of modules	10
2. Homogeneous prime spectra	13
2.1 Generalities on graded rings and modules	13
2.2 Rings of fractions of a graded ring	16
2.3 Homogeneous prime spectrum of a graded ring	18
2.4 The scheme structure on $\text{Proj}(S)$	20
2.5 The sheaf associated to a graded module	22
2.6 The graded S -module associated to a sheaf on $\text{Proj}(S)$	26
2.7 Finiteness conditions	28
2.8 Functorial behaviour	30
2.9 Closed subschemes of a scheme $\text{Proj}(S)$	35
3. Homogeneous spectrum of a sheaf of graded algebras	36
3.1 Homogeneous spectrum of a quasi-coherent graded $\mathcal{O}_Y$ -algebra	36
3.2 Sheaf on $\text{Proj}(\mathcal{S})$ associated to a graded $\mathcal{S}$ -module	40
3.3 Graded $\mathcal{S}$ -module associated to a sheaf on $\text{Proj}(\mathcal{S})$	42
3.4 Finiteness conditions	44
3.5 Functorial behaviour	46
3.6 Closed subpreschemes of a prescheme $\text{Proj}(\mathcal{S})$	48
3.7 Morphisms from a prescheme to a homogeneous spectrum	49
3.8 Criteria for immersion into a homogeneous spectrum	51
4. Projective bundles; ample sheaves	53
4.1 Definition of projective bundles	53
4.2 Morphisms from a prescheme to a projective bundle	54
4.3 The Segre morphism	57
4.4 Immersions into projective bundles; very ample sheaves	59
4.5 Ample sheaves	62
4.6 Relatively ample sheaves	66
5. Quasi-affine morphisms; quasi-projective morphisms; proper morphisms; projective morphisms	70
5.1 Quasi-affine morphisms	70
5.2 Serre's criterion	72
5.3 Quasi-projective morphisms	74
5.4 Proper morphisms and universally closed morphisms	75
5.5 Projective morphisms	77
5.6 Chow's lemma	80
6. Integral morphisms and finite morphisms	82
6.1 Preschemes integral over another prescheme	82
6.2 Quasi-finite morphisms	86

6.3	Integral closure of a prescheme	88
6.4	Determinant of an endomorphism of $\mathcal{O}_X$ -modules	91
6.5	Norm of an invertible sheaf	94
6.6	Application: criteria for ampleness	98
6.7	Chevalley's theorem	102
7.	Valuative criteria	103
7.1	Reminder on valuation rings	104
7.2	Valuative criterion for separatedness	106
7.3	Valuative criterion for properness	107
7.4	Algebraic curves and function fields of dimension 1	111
8.	Blowup schemes; projecting cones; projective closure	114
8.1	Blowup preschemes	114
8.2	Preliminary results on the localisation of graded rings	117
8.3	Projecting cones	122
8.4	Projective closure of a vector bundle	126
8.5	Functorial behaviour	127
8.6	A canonical isomorphism for punctured cones	129
8.7	Blowing up projecting cones	130
8.8	Ample sheaves and contractions	133
8.9	Grauert's ampleness criterion: statement	137
8.10	Grauert's ampleness criterion: proof	138
8.11	Uniqueness of contractions	142
8.12	Quasi-coherent sheaves on projecting cones	144
8.13	Projective closures of subsheaves and closed subschemes	147
8.14	Supplement on sheaves associated to graded $\mathcal{S}$ -modules	148
	Bibliography	153
	Bibliography	153
	Index of notation	154
	Index of notation	154
	Terminological index	156
	Terminological index	156
	Original table of contents	159
	Original table of contents	159
	Errata and Addenda (List 1)	161
	Errata and Addenda	161

SUMMARY

- §1. Affine morphisms.
- §2. Homogeneous prime spectra.
- §3. Homogeneous spectrum of a sheaf of graded algebras.
- §4. Projective bundles; ample sheaves.
- §5. Quasi-affine morphisms; quasi-projective morphisms; proper morphisms; projective morphisms.
- §6. Integral morphisms and finite morphisms.
- §7. Valuative criteria.
- §8. Blowup schemes; projective cones; projective closure.

[II:5] The various classes of morphisms studied in this chapter are treated without making much use of cohomological methods; a deeper study making use of those methods will be carried out in Chapter III, where we shall use chiefly §§2, 4, and 5 of Chapter II. §8 may be omitted on a first reading; it gives some supplements to the formalism developed in §§1 through 3, reducing to easy

applications of that formalism, and we shall use it less consistently than the other results of this chapter.

§1. AFFINE MORPHISMS

Most of the results in this section are the “global” counterparts of those in Chapter I, §1; they are therefore not essentially new and merely provide convenient language for what follows.

1.1. S -preschemes and $\mathcal{O}_S$ -algebras

(1.1.1). Let S be a prescheme, X an S -prescheme, and $f : X \rightarrow S$ its structure morphism. We know (0, 4.2.4) that the direct image $f_*(\mathcal{O}_X)$ is an $\mathcal{O}_S$ -algebra, which we ^[II-6] denote $\mathcal{A}(X)$ when there is little chance of confusion; if U is an open subset of S , then we have

$$\mathcal{A}(f^{-1}(U)) = \mathcal{A}(X)|_U.$$

Similarly, for every $\mathcal{O}_X$ -module $\mathcal{F}$ (resp. every $\mathcal{O}_X$ -algebra $\mathcal{B}$), we write $\mathcal{A}(\mathcal{F})$ (resp. $\mathcal{A}(\mathcal{B})$) for the direct image $f_*(\mathcal{F})$ (resp. $f_*(\mathcal{B})$) which is an $\mathcal{A}(X)$ -module (resp. an $\mathcal{A}(X)$ -algebra) and not only an $\mathcal{O}_S$ -module (resp. an $\mathcal{O}_S$ -algebra).

(1.1.2). Let Y be a second S -prescheme, $g : Y \rightarrow S$ its structure morphism, and $h : X \rightarrow Y$ an S -morphism; we then have the commutative diagram

$$(1.1.2.1) \quad \begin{array}{ccc} X & \xrightarrow{h} & Y \\ & \searrow f & \swarrow g \\ & S. & \end{array}$$

We have by definition $h = (\psi, \theta)$, where $\theta : \mathcal{O}_Y \rightarrow h_*(\mathcal{O}_X) = \psi_*(\mathcal{O}_X)$ is a homomorphism of sheaves of rings; we thus obtain (0, 4.2.2) a homomorphism of $\mathcal{O}_S$ -algebras $g_*(\theta) : g_*(\mathcal{O}_Y) \rightarrow g_*(h_*(\mathcal{O}_X)) = f_*(\mathcal{O}_X)$, in other words, a homomorphism of $\mathcal{O}_S$ -algebras $\mathcal{A}(Y) \rightarrow \mathcal{A}(X)$, which we denote by $\mathcal{A}(h)$. If $h' : Y \rightarrow Z$ is a second S -morphism, then it is immediate that $\mathcal{A}(h' \circ h) = \mathcal{A}(h) \circ \mathcal{A}(h')$. We have thus defined a *contravariant functor* $X \mapsto \mathcal{A}(X)$ on the category of S -preschemes, with values in the category of $\mathcal{O}_S$ -algebras.

Now let $\mathcal{F}$ be an $\mathcal{O}_X$ -module, $\mathcal{G}$ an $\mathcal{O}_Y$ -module, and $u : \mathcal{G} \rightarrow \mathcal{F}$ an h -morphism, that is (0, 4.4.1) a homomorphism of $\mathcal{O}_Y$ -modules $\mathcal{G} \rightarrow h_*(\mathcal{F})$. Then $g_*(u) : g_*(\mathcal{G}) \rightarrow g_*(h_*(\mathcal{F})) = f_*(\mathcal{F})$ is a homomorphism $\mathcal{A}(\mathcal{G}) \rightarrow \mathcal{A}(\mathcal{F})$ of $\mathcal{O}_S$ -modules, which we denote by $\mathcal{A}(u)$; in addition, the pair $(\mathcal{A}(h), \mathcal{A}(u))$ forms a *di-homomorphism* from the $\mathcal{A}(Y)$ -module $\mathcal{A}(\mathcal{G})$ to the $\mathcal{A}(X)$ -module $\mathcal{A}(\mathcal{F})$.

(1.1.3). If we fix the prescheme S , then we can consider the pairs $(X, \mathcal{F})$, where X is an S -prescheme and $\mathcal{F}$ is an $\mathcal{O}_X$ -module, as forming a *category*, by defining a *morphism* $(X, \mathcal{F}) \rightarrow (Y, \mathcal{G})$ as a pair (h, u) , where $h : X \rightarrow Y$ is an S -morphism and $u : \mathcal{G} \rightarrow \mathcal{F}$ is an h -morphism. We can then say that $(X, \mathcal{F}) \mapsto (\mathcal{A}(X), \mathcal{A}(\mathcal{F}))$ is a *contravariant functor* with values in the category whose objects are pairs consisting of an $\mathcal{O}_S$ -algebra and a module over that algebra, and whose morphisms are the di-homomorphisms.

1.2. Affine preschemes over a prescheme

Definition (1.2.1). — Let X be an S -prescheme, and $f : X \rightarrow S$ its structure morphism. We say that X is *affine over S* if there exists a cover (S_α) of S by affine open sets such that for all α , the prescheme induced by X on the open set $f^{-1}(S_\alpha)$ is affine.

Example (1.2.2). — Every closed subprescheme of S is an affine S -prescheme over S ((I, 4.2.3) and (I, 4.2.4)).

Remark (1.2.3). — An affine prescheme X over S is not necessarily an affine scheme, as the example $X = S$ shows (1.2.2). On the other hand, if an affine scheme X is an S -prescheme, then X is not necessarily affine over S (see Example (1.3.3)). However, recall that if S is a *scheme*, then every S -prescheme which is an affine scheme is affine over S (I, 5.5.10).

Proposition (1.2.4). — *Every S -prescheme which is affine over S is separated over S (in other words, it is an S -scheme).*

Proof. This follows immediately from (I, 5.5.5) and (I, 5.5.8). $\square$

Proposition (1.2.5). — *Let X be an S -scheme affine over S , and $f : X \rightarrow S$ its structure morphism. For every open $U \subset S$, $f^{-1}(U)$ is affine over U .*

Proof. By Definition (1.2.1), we can reduce to the case where $S = \text{Spec}(A)$ and $X = \text{Spec}(B)$ are affine; then $f = ({}^a\phi, \tilde{\phi})$, where $\phi : A \rightarrow B$ is a homomorphism. As the $D(g)$ for $g \in A$ form a basis for S , we reduce to the case where $U = D(g)$; but we then know that $f^{-1}(U) = D(\phi(g))$ (I, 1.2.2.2), hence the proposition. $\square$

Proposition (1.2.6). — *Let X be an S -scheme affine over S , and $f : X \rightarrow S$ its structure morphism. For every quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$, $f_*(\mathcal{F})$ is a quasi-coherent $\mathcal{O}_S$ -module.*

Proof. Taking into account Proposition (1.2.4), this follows from (I, 9.2.2, a). $\square$

In particular, the $\mathcal{O}_S$ -algebra $\mathcal{A}(X) = f_*(\mathcal{O}_X)$ is *quasi-coherent*.

Proposition (1.2.7). — *Let X be an S -scheme affine over S . For every S -prescheme Y , the map $h \mapsto \mathcal{A}(h)$ from the set $\text{Hom}_S(Y, X)$ to the set $\text{Hom}(\mathcal{A}(X), \mathcal{A}(Y))$ (1.1.2) is bijective.*

Proof. Let $f : X \rightarrow S$ and $g : Y \rightarrow S$ be the structure morphisms. First, suppose that $S = \text{Spec}(A)$ and $X = \text{Spec}(B)$ are affine; we must prove that for every homomorphism $\omega : f_*(\mathcal{O}_X) \rightarrow g_*(\mathcal{O}_Y)$ of $\mathcal{O}_S$ -algebras, there exists a unique S -morphism $h : Y \rightarrow X$ such that $\mathcal{A}(h) = \omega$. By definition, for every open $U \subset S$, ω defines a homomorphism $\omega_U = \Gamma(U, \omega) : \Gamma(f^{-1}(U), \mathcal{O}_X) \rightarrow \Gamma(g^{-1}(U), \mathcal{O}_Y)$ of $\Gamma(U, \mathcal{O}_S)$ -algebras. In particular, for $U = S$, this gives a homomorphism $\phi : \Gamma(X, \mathcal{O}_X) \rightarrow \Gamma(Y, \mathcal{O}_Y)$ of $\Gamma(S, \mathcal{O}_S)$ -algebras, to which corresponds a well-defined S -morphism $h : Y \rightarrow X$, since X is affine (I, 2.2.4). It remains to prove that $\mathcal{A}(h) = \omega$, or, in other words, that, for every open set U of a basis for S , ω_U coincides with the homomorphism of algebras ϕ_U corresponding to the S -morphism $g^{-1}(U) \rightarrow f^{-1}(U)$, a restriction of h . We can reduce to the case where $U = D(\lambda)$, with $\lambda \in A$; then, if $f = ({}^a\rho, \tilde{\rho})$, where $\rho : A \rightarrow B$ is a ring homomorphism, we have $f^{-1}(U) = D(\mu)$, where $\mu = \rho(\lambda)$, and $\Gamma(f^{-1}(U), \mathcal{O}_X)$ is the ring of fractions B_μ ; the diagram

$$\begin{array}{ccc} B & \xrightarrow{\phi} & \Gamma(Y, \mathcal{O}_Y) \\ \downarrow & & \downarrow \\ B_\mu & \xrightarrow{\phi_U} & \Gamma(g^{-1}(U), \mathcal{O}_Y) \end{array}$$

is commutative, and so is the analogous diagram where ϕ_U is replaced by ω_U ; the equality $\phi_U = \omega_U$ then follows from the universal property of rings of fractions (0, 1.2.4).

We now pass to the general case; let (S_α) be a cover of S by affine open sets ^[II:8] such that the $f^{-1}(S_\alpha)$ are affine. Then every homomorphism $\omega : \mathcal{A}(X) \rightarrow \mathcal{A}(Y)$ of $\mathcal{O}_S$ -algebras gives by restriction a family of homomorphisms

$$\omega_\alpha : \mathcal{A}(f^{-1}(S_\alpha)) \longrightarrow \mathcal{A}(g^{-1}(S_\alpha))$$

of $\mathcal{O}_{S_\alpha}$ -algebras, hence a family of S_α -morphisms $h_\alpha : g^{-1}(S_\alpha) \rightarrow f^{-1}(S_\alpha)$ by the above. It remains to see that for every affine open set of a basis for $S_\alpha \cap S_\beta$, the restrictions of h_α and h_β to $g^{-1}(U)$ coincide, which is evident since by the above, these restrictions both correspond to the homomorphism $\mathcal{A}(X)|_U \rightarrow \mathcal{A}(Y)|_U$, a restriction of ω . $\square$

Corollary (1.2.8). — *Let X and Y be two S -schemes which are affine over S . For an S -morphism $h : Y \rightarrow X$ to be an isomorphism, it is necessary and sufficient for $\mathcal{A}(h) : \mathcal{A}(X) \rightarrow \mathcal{A}(Y)$ to be an isomorphism.*

Proof. This follows immediately from Proposition (1.2.7) and from the functorial nature of $\mathcal{A}(X)$. $\square$

1.3. Affine preschemes over S associated to an $\mathcal{O}_S$ -algebra

Proposition (1.3.1). — *Let S be a prescheme. For every quasi-coherent $\mathcal{O}_S$ -algebra $\mathcal{B}$, there exists a prescheme X affine over S , defined up to unique S -isomorphism, such that $\mathcal{A}(X) = \mathcal{B}$.*

Proof. The uniqueness follows from Corollary (1.2.8); we prove the existence of X . For every affine open $U \subset S$, let X_U be the prescheme $\text{Spec}(\Gamma(U, \mathcal{B}))$; as $\Gamma(U, \mathcal{B})$ is a $\Gamma(U, \mathcal{O}_S)$ -algebra, X_U is an S -prescheme (I, 1.6.1). In addition, as $\mathcal{B}$ is quasi-coherent, the $\mathcal{O}_S$ -algebra $\mathcal{A}(X_U)$ canonically identifies with $\mathcal{B}|_U$ (I, 1.3.7), (I, 1.3.13), (I, 1.6.3)). Let V be a second affine open subset of S , and let $X_{U,V}$ be the prescheme induced by X_U on $f_U^{-1}(U \cap V)$, where f_U denotes the structure morphism $X_U \rightarrow S$; $X_{U,V}$ and $X_{V,U}$ are affine over $U \cap V$ (1.2.5), and by definition $\mathcal{A}(X_{U,V})$ and $\mathcal{A}(X_{V,U})$ canonically identify with $\mathcal{B}|_{(U \cap V)}$. Hence there is (1.2.8) a canonical S -isomorphism $\theta_{U,V} : X_{V,U} \rightarrow X_{U,V}$; in addition, if W is a third affine open subset of S , and if $\theta'_{U,V}$, $\theta'_{V,W}$, and $\theta'_{U,W}$ are the restrictions of $\theta_{U,V}$, $\theta_{V,W}$, and $\theta_{U,W}$ to the inverse images of $U \cap V \cap W$ in X_V , X_W , and X_W respectively under the structure morphisms, then we have $\theta'_{U,V} \circ \theta'_{V,W} = \theta'_{U,W}$. As a result, there exists a prescheme X , a cover (T_U) of X by affine open sets, and for every U an isomorphism $\phi_U : X_U \rightarrow T_U$, such that ϕ_U maps $f_U^{-1}(U \cap V)$ to $T_U \cap T_V$, and we have $\theta_{U,V} = \phi_U^{-1} \circ \phi_V$ (I, 2.3.1). The morphism $g_U = f_U \circ \phi_U^{-1}$ makes T_U an S -prescheme, and the morphisms g_U and g_V coincide on $T_U \cap T_V$, hence X is an S -prescheme. In addition, it is clear by definition that X is affine over S and that $\mathcal{A}(T_U) = \mathcal{B}|_U$, hence $\mathcal{A}(X) = \mathcal{B}$. $\square$

We say that the S -scheme X defined in this way is *associated to the $\mathcal{O}_S$ -algebra $\mathcal{B}$* , or is the *spectrum of $\mathcal{B}$* , and we denote it by $\text{Spec}(\mathcal{B})$.

Corollary (1.3.2). — *Let X be a prescheme affine over S , $f : X \rightarrow S$ the structure morphism. For every affine open $U \subset S$, the induced prescheme on $f^{-1}(U)$ is the affine scheme with ring $\Gamma(U, \mathcal{A}(X))$.*

Proof. [II-9] As we can suppose that X is associated to an $\mathcal{O}_S$ -algebra by Propositions (1.2.6) and (1.3.1), the corollary follows from the construction of X described in Proposition (1.3.1). $\square$

Example (1.3.3). — Let S be the affine plane over a field K , where the point 0 has been doubled (I, 5.5.11); with the notation of (I, 5.5.11), S is the union of two affine open sets Y_1 and Y_2 ; if f is the open immersion $Y_1 \rightarrow S$, then $f^{-1}(Y_2) = Y_1 \cap Y_2$ is not an affine open set in Y_1 (*loc. cit.*), hence we have an example of an affine scheme which is not affine over S .

Corollary (1.3.4). — *Let S be an affine scheme; for an S -prescheme X to be affine over S , it is necessary and sufficient for X to be an affine scheme.*

Corollary (1.3.5). — *Let X be a prescheme affine over a prescheme S , and let Y be an X -prescheme. For Y to be affine over S , it is necessary and sufficient for Y to be affine over X .*

Proof. We immediately reduce to the case where S is an affine scheme, and then X is likewise an affine scheme (1.3.4); the two conditions in the statement then both express that Y is an affine scheme (1.3.4). $\square$

(1.3.6). Let X be a prescheme affine over S . To define a prescheme Y affine over X , it is equivalent, by Corollary (1.3.5), to give a prescheme Y affine over S , and an S -morphism $g : Y \rightarrow X$; in other words (Proposition (1.3.1) and (1.2.7)), it is equivalent to give a quasi-coherent $\mathcal{O}_S$ -algebra $\mathcal{B}$ and a homomorphism $\mathcal{A}(X) \rightarrow \mathcal{B}$ of $\mathcal{O}_S$ -algebras (which can be considered as defining on $\mathcal{B}$ an $\mathcal{A}(X)$ -algebra structure). If $f : X \rightarrow S$ is the structure morphism, then we have $\mathcal{B} = f_*(g_*(\mathcal{O}_Y))$.

Corollary (1.3.7). — *Let X be a prescheme affine over S ; for X to be of finite type over S , it is necessary and sufficient for the quasi-coherent $\mathcal{O}_S$ -algebra $\mathcal{A}(X)$ to be of finite type (I, 9.6.2).*

Proof. By definition (I, 9.6.2), we can reduce to the case where S is affine; then X is an affine scheme (1.3.4), and if $S = \operatorname{Spec}(A)$, $X = \operatorname{Spec}(B)$, then $\mathcal{A}(X)$ is the $\mathcal{O}_S$ -algebra $\tilde{B}$; as $\Gamma(S, \tilde{B}) = B$, the corollary follows from (I, 9.6.2) and (I, 6.3.3). $\square$

Corollary (1.3.8). — *Let X be a prescheme affine over S ; for X to be reduced, it is necessary and sufficient for the quasi-coherent $\mathcal{O}_S$ -algebra $\mathcal{A}(X)$ to be reduced (0, 4.1.4).*

Proof. The question is local on S ; by Corollary (1.3.2), the corollary follows from (I, 5.1.1) and (I, 5.1.4). $\square$

1.4. Quasi-coherent sheaves over an affine prescheme over S

Proposition (1.4.1). — *Let X be a prescheme affine over S , Y an S -prescheme, and $\mathcal{F}$ (resp. $\mathcal{G}$) a quasi-coherent $\mathcal{O}_X$ -module (resp. an $\mathcal{O}_Y$ -module). Then the map $(h, u) \mapsto (\mathcal{A}(h), \mathcal{A}(u))$ from the set of morphisms $(Y, \mathcal{G}) \rightarrow (X, \mathcal{F})$ to the set of di-homomorphisms $(\mathcal{A}(X), \mathcal{A}(\mathcal{F})) \rightarrow (\mathcal{A}(Y), \mathcal{A}(\mathcal{G}))$ ((1.1.2) and (1.1.3)) is bijective.*

Proof. The proof follows exactly as that of Proposition (1.2.7) by using (I, 2.2.5) and (I, 2.2.4), and the details are left to the reader. $\square$

Corollary (1.4.2). — *If, in addition to the hypotheses of Proposition (1.4.1), we suppose that Y is affine over S , then for (h, u) to be an isomorphism, it is necessary and sufficient for $(\mathcal{A}(h), \mathcal{A}(u))$ to be a di-isomorphism.*

Proposition (1.4.3). — ^[III:10] *For every pair $(\mathcal{B}, \mathcal{M})$ consisting of a quasi-coherent $\mathcal{O}_S$ -algebra $\mathcal{B}$ and a quasi-coherent $\mathcal{B}$ -module $\mathcal{M}$ (considered as an $\mathcal{O}_S$ -module or as a $\mathcal{B}$ -module, which are equivalent (I, 9.6.1)), there exists a pair $(X, \mathcal{F})$ consisting of a prescheme X affine over S and of a quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$, such that $\mathcal{A}(X) = \mathcal{B}$ and $\mathcal{A}(\mathcal{F}) = \mathcal{M}$; in addition, this pair is determined up to unique isomorphism.*

Proof. The uniqueness follows from Proposition (1.4.1) and Corollary (1.4.2); the existence is proved as in Proposition (1.3.1), and we leave the details to the reader. $\square$

We denote by $\tilde{\mathcal{M}}$ the $\mathcal{O}_X$ -module $\mathcal{F}$, and we say that it is *associated* to the quasi-coherent $\mathcal{B}$ -module $\mathcal{M}$; for every affine open $U \subset S$, $\tilde{\mathcal{M}}|_{p^{-1}(U)}$ (where p is the structure morphism $X \rightarrow S$) is canonically isomorphic to $(\Gamma(U, \mathcal{M}))^\sim$.

Corollary (1.4.4). — *On the category of quasi-coherent $\mathcal{B}$ -modules, $\mathcal{M} \mapsto \tilde{\mathcal{M}}$ is an additive covariant exact functor which commutes with inductive limits and direct sums.*

Proof. We immediately reduce to the case where S is affine, and the corollary then follows from (I, 1.3.5), (I, 1.3.9), and (I, 1.3.11). $\square$

Corollary (1.4.5). — *Under the hypotheses of Proposition (1.4.3), for $\tilde{\mathcal{M}}$ to be an $\mathcal{O}_X$ -module of finite type, it is necessary and sufficient for $\mathcal{M}$ to be a $\mathcal{B}$ -module of finite type.*

Proof. We immediately reduce to the case where $S = \operatorname{Spec}(A)$ is an affine scheme. Then $\mathcal{B} = \tilde{B}$, where B is an A -algebra, and $\mathcal{M} = \tilde{M}$, where M is a B -module (I, 1.3.13); over the prescheme X , $\mathcal{O}_X$ is associated to the ring B and $\tilde{\mathcal{M}}$ to the B -module M ; for $\tilde{\mathcal{M}}$ to be of finite type, it is therefore necessary and sufficient for M to be of finite type (I, 1.3.13), hence our assertion. $\square$

Proposition (1.4.6). — *Let Y be a prescheme affine over S , X and X' two preschemes affine over Y (hence also over S (1.3.5)). Let $\mathcal{B} = \mathcal{A}(Y)$, $\mathcal{A} = \mathcal{A}(X)$, and $\mathcal{A}' = \mathcal{A}(X')$. Then $X \times_Y X'$ is affine over Y (thus also over S), and $\mathcal{A}(X \times_Y X')$ canonically identifies with $\mathcal{A} \otimes_{\mathcal{B}} \mathcal{A}'$.*

Proof. By (I, 9.6.1), $\mathcal{A} \otimes_{\mathcal{B}} \mathcal{A}'$ is a quasi-coherent $\mathcal{B}$ -algebra, thus also a quasi-coherent $\mathcal{O}_S$ -algebra (I, 9.6.1); let Z be the spectrum of $\mathcal{A} \otimes_{\mathcal{B}} \mathcal{A}'$ (1.3.1). The canonical $\mathcal{B}$ -homomorphisms $\mathcal{A} \rightarrow \mathcal{A} \otimes_{\mathcal{B}} \mathcal{A}'$ and $\mathcal{A}' \rightarrow \mathcal{A} \otimes_{\mathcal{B}} \mathcal{A}'$ correspond (1.2.7) to Y -morphisms $p : Z \rightarrow X$ and $p' : Z \rightarrow X'$. To see that the triple (Z, p, p') is a product $X \times_Y X'$, we can reduce to the case where S is an affine scheme

with ring C (I, 3.2.6.4). But then Y , X , and X' are affine schemes (1.3.4) whose rings B , A , and A' are C -algebras such that $\mathcal{B} = \tilde{B}$, $\mathcal{A} = \tilde{A}$, and $\mathcal{A}' = \tilde{A}'$. We then know (I, 1.3.13) that $\mathcal{A} \otimes_{\mathcal{B}} \mathcal{A}'$ canonically identifies with the $\mathcal{O}_S$ -algebra $(A \otimes_B A')^\sim$, hence the ring $A(Z)$ identifies with $A \otimes_B A'$ and the morphisms p and p' correspond to the canonical homomorphisms $A \rightarrow A \otimes_B A'$ and $A' \rightarrow A \otimes_B A'$. The proposition then follows from (I, 3.2.2). $\square$

Corollary (1.4.7). — *Let $\mathcal{F}$ (resp. $\mathcal{F}'$) be a quasi-coherent $\mathcal{O}_X$ -module (resp. $\mathcal{O}_{X'}$ -module); then $\mathcal{A}(\mathcal{F} \otimes_Y \mathcal{F}')$ canonically identifies with $\mathcal{A}(\mathcal{F}) \otimes_{\mathcal{A}(Y)} \mathcal{A}(\mathcal{F}')$.*

Proof. We know that $\mathcal{F} \otimes_Y \mathcal{F}'$ is quasi-coherent over $X \times_Y X'$ (I, 9.1.2). Let $g : Y \rightarrow S$, $f : X \rightarrow Y$, and $f' : X' \rightarrow Y$ be the structure morphisms, such that the structure morphism $h : Z \rightarrow S$ is equal to $g \circ f \circ p$ and to $g \circ f' \circ p'$. We define a canonical homomorphism

$$\mathcal{A}(\mathcal{F}) \otimes_{\mathcal{A}(Y)} \mathcal{A}(\mathcal{F}') \longrightarrow \mathcal{A}(\mathcal{F} \otimes_Y \mathcal{F}')$$

in the following way: for every open $U \subset S$, we have canonical homomorphisms $\Gamma(f^{-1}(g^{-1}(U)), \mathcal{F}) \rightarrow \Gamma(h^{-1}(U), p^*(\mathcal{F}))$ and $\Gamma(f'^{-1}(g^{-1}(U)), \mathcal{F}') \rightarrow \Gamma(h^{-1}(U), p'^*(\mathcal{F}'))$ (0, 4.4.3), thus we obtain a canonical homomorphism

$$\Gamma(f^{-1}(g^{-1}(U)), \mathcal{F}) \otimes_{\Gamma(g^{-1}(U), \mathcal{O}_Y)} \Gamma(f'^{-1}(g^{-1}(U)), \mathcal{F}') \longrightarrow \Gamma(h^{-1}(U), p^*(\mathcal{F})) \otimes_{\Gamma(h^{-1}(U), \mathcal{O}_Z)} \Gamma(h^{-1}(U), p'^*(\mathcal{F}')).$$

To see that we have defined an isomorphism of $\mathcal{A}(Z)$ -modules, we can reduce to the case where S (and as a result X , X' , Y , and $X \times_Y X'$) are affine schemes, and (with the notation of Proposition (1.4.6)), $\mathcal{F} = \tilde{M}$, $\mathcal{F}' = \tilde{M}'$, where M (resp. M') is an A -module (resp. an A' -module). Then $\mathcal{F} \otimes_Y \mathcal{F}'$ identifies with the sheaf on $X \times_Y X'$ associated to the $(A \otimes_B A')$ -module $M \otimes_B M'$ (I, 9.1.3), and the corollary follows from the canonical identification of the $\mathcal{O}_S$ -modules $(M \otimes_B M')^\sim$ and $\tilde{M} \otimes_{\tilde{B}} \tilde{M}'$ (where M , M' , and B are considered as C -modules) ((I, 1.3.13) and (I, 1.6.3)). $\square$

If we apply Corollary (1.4.7) in particular to the case where $X = Y$ and $\mathcal{F}' = \mathcal{O}_{X'}$, then we see that the $\mathcal{A}'$ -module $\mathcal{A}(f'^*(\mathcal{F}))$ identifies with $\mathcal{A}(\mathcal{F}) \otimes_{\mathcal{B}} \mathcal{A}'$.

(1.4.8). In particular, when $X = X' = Y$ (X being affine over S), we see that if $\mathcal{F}$ and $\mathcal{G}$ are two quasi-coherent $\mathcal{O}_X$ -modules, then we have

$$(1.4.8.1) \quad \mathcal{A}(\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{G}) = \mathcal{A}(\mathcal{F}) \otimes_{\mathcal{A}(X)} \mathcal{A}(\mathcal{G})$$

up to canonical functorial isomorphism. If in addition $\mathcal{F}$ admits a finite presentation, then it follows from (I, 1.6.3) and (I, 1.3.12) that

$$(1.4.8.2) \quad \mathcal{A}(\mathcal{H}om_X(\mathcal{F}, \mathcal{G})) = \mathcal{H}om_{\mathcal{A}(X)}(\mathcal{A}(\mathcal{F}), \mathcal{A}(\mathcal{G}))$$

up to canonical isomorphism.

Remark (1.4.9). — If X and X' are two preschemes affine over S , then the sum $X \sqcup X'$ is also affine over S , as the sum of two affine schemes is an affine scheme.

Proposition (1.4.10). — *Let S be a prescheme, $\mathcal{B}$ a quasi-coherent $\mathcal{O}_S$ -algebra, and $X = \text{Spec}(\mathcal{B})$. For a quasi-coherent sheaf of ideals $\mathcal{J}$ of $\mathcal{B}$, $\tilde{\mathcal{J}}$ is a quasi-coherent sheaf of ideals of $\mathcal{O}_X$, and the closed subscheme Y of X defined by $\tilde{\mathcal{J}}$ is canonically isomorphic to $\text{Spec}(\mathcal{B}/\mathcal{J})$.*

Proof. It follows immediately from (I, 4.2.3) that Y is affine over S ; by Proposition (1.3.1), we reduce to the case where S is affine, and the proposition then follows immediately from (I, 4.1.2). $\square$

We can also express the result of Proposition (1.4.10) by saying that if $h : \mathcal{B} \rightarrow \mathcal{B}'$ is a surjective homomorphism of quasi-coherent $\mathcal{O}_S$ -algebras, $^ah : \text{Spec}(\mathcal{B}') \rightarrow \text{Spec}(\mathcal{B})$ is a closed immersion.

Proposition (1.4.11). — ^[II:12] *Let S be a prescheme, $\mathcal{B}$ a quasi-coherent $\mathcal{O}_S$ -algebra, and $X = \text{Spec}(\mathcal{B})$. For every quasi-coherent sheaf of ideals $\mathcal{K}$ of $\mathcal{O}_S$, we have (denoting by f the structure morphism $X \rightarrow S$) $f^*(\mathcal{K})\mathcal{O}_X = (\mathcal{K}\mathcal{B})^\sim$ up to canonical isomorphism.*

Proof. Since the question is local on S , we can reduce to the case where $S = \text{Spec}(A)$ is affine, and in this case the proposition is none other than (I, 1.6.9). $\square$

1.5. Change of base prescheme

Proposition (1.5.1). — *Let X be a prescheme affine over S . For every extension $g : S' \rightarrow S$ of the base prescheme, $X' = X_{(S')} = X \times_S S'$ is affine over S' .*

Proof. If f' is the projection $X' \rightarrow S'$, then it suffices to prove that $f'^{-1}(U')$ is an affine open set for every affine open subset U' of S' such that $g(U')$ is contained in an affine open subset U of S (1.2.1); we can thus reduce to the case where S and S' are affine, and it suffices to prove that X' is then an affine scheme (1.3.4). But then X is an affine scheme (1.3.4), and if A , A' , and B are the rings of S , S' , and X respectively, then we know that X' is the affine scheme with ring $A' \otimes_A B$ (I, 3.2.2). $\square$

Corollary (1.5.2). — *Under the hypotheses of Proposition (1.5.1), let $f : X \rightarrow S$ be the structure morphism, $f' : X' \rightarrow S'$ and $g' : X' \rightarrow X$ the projections, such that the diagram*

$$\begin{array}{ccc} X & \xleftarrow{g'} & X' \\ f \downarrow & & \downarrow f' \\ S & \xleftarrow{g} & S' \end{array}$$

is commutative. For every quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$, there exists a canonical isomorphism of $\mathcal{O}_{S'}$ -modules

$$(1.5.2.1) \quad u : g^*(f_*(\mathcal{F})) \simeq f'_*(g'^*(\mathcal{F})).$$

In particular, there exists a canonical isomorphism from $\mathcal{A}(X')$ to $g^(\mathcal{A}(X))$.*

Proof. To define u , it suffices to define a homomorphism

$$v : f_*(\mathcal{F}) \longrightarrow g_*(f'_*(g'^*(\mathcal{F}))) = f_*(g'_*(g'^*(\mathcal{F})))$$

and to set $u = v^\#$ (0, 4.4.3). We take $v = f_*(\rho)$, where ρ is the canonical homomorphism $\mathcal{F} \rightarrow g'_*(g'^*(\mathcal{F}))$ (0, 4.4.3). To prove that u is an isomorphism, we can reduce to the case where S and S' are affine, and hence so are X and X' ; with the notation of Proposition (1.5.1), we then have $\mathcal{F} = \tilde{M}$, where M is a B -module. We then note immediately that $g^*(f_*(\mathcal{F}))$ and $f'_*(g'^*(\mathcal{F}))$ are both equal to the $\mathcal{O}_{S'}$ -module associated to the A' -module $A' \otimes_A M$ (where M is considered as an A -module), and that u is the homomorphism associated to the identity ((I, 1.6.3), (I, 1.6.5), (I, 1.6.7)). $\square$

Remark (1.5.3). — One must not suppose that Corollary (1.5.2) remains true when X is no longer assumed affine over S , even when $S' = \text{Spec}(k(s))$ ($s \in S$) and $S' \rightarrow S$ is the canonical morphism (I, 2.4.5)—in which case X' is none other than the fibre $f^{-1}(s)$ (I, 3.6.2). In other words, when X is not affine over S , the operation ^[II:13] “direct image of quasi-coherent sheaves” does not commute with the operation of “passing to fibres”. However, we will see in Chapter III (III, 4.2.4) a result in this sense, of an “asymptotic” nature, valid for *coherent* sheaves on X when f is proper (5.4) and S is Noetherian.

Corollary (1.5.4). — *For every prescheme X affine over S and every $s \in S$, the fibre $f^{-1}(s)$ (where f denotes the structure morphism $X \rightarrow S$) is an affine scheme.*

Proof. It suffices to apply Proposition (1.5.1) with $S' = \text{Spec}(k(s))$ and to use Corollary (1.3.4). $\square$

Corollary (1.5.5). — *Let X be an S -prescheme and S' a prescheme affine over S ; then $X' = X_{(S')}$ is a prescheme affine over X . In addition, if $f : X \rightarrow S$ is the structure morphism, then there is a canonical isomorphism of $\mathcal{O}_X$ -algebras $\mathcal{A}(X') \simeq f^*(\mathcal{A}(S'))$, and for every quasi-coherent $\mathcal{A}(S')$ -module $\mathcal{M}$, a canonical di-isomorphism $f^*(\mathcal{M}) \simeq \mathcal{A}(f'^*(\tilde{\mathcal{M}}))$, denoting by $f' = f_{(S')}$ the structure morphism $X' \rightarrow S'$.*

Proof. It suffices to swap the roles of X and S' in (1.5.1) and (1.5.2). $\square$

(1.5.6). Now let S, S' be two preschemes, $q : S' \rightarrow S$ a morphism, $\mathcal{B}$ (resp. $\mathcal{B}'$) a quasi-coherent $\mathcal{O}_S$ -algebra (resp. $\mathcal{O}_{S'}$ -algebra), $u : \mathcal{B} \rightarrow \mathcal{B}'$ a q -morphism (that is, a homomorphism $\mathcal{B} \rightarrow q_*(\mathcal{B}')$ of $\mathcal{O}_S$ -algebras). If $X = \text{Spec}(\mathcal{B})$, $X' = \text{Spec}(\mathcal{B}')$, then we canonically obtain a morphism

$$v = \text{Spec}(u) : X' \longrightarrow X$$

such that the diagram

$$(1.5.6.1) \quad \begin{array}{ccc} X' & \xrightarrow{v} & X \\ \downarrow & & \downarrow \\ S' & \xrightarrow{q} & S \end{array}$$

is commutative (the vertical arrows being the structure morphisms). Indeed, the data of u is equivalent to that of a homomorphism of quasi-coherent $\mathcal{O}_{S'}$ -algebras $u^\sharp : q^*(\mathcal{B}) \rightarrow \mathcal{B}'$ (0, 4.4.3); this thus canonically defines an S' -morphism

$$w : \operatorname{Spec}(\mathcal{B}') \longrightarrow \operatorname{Spec}(q^*(\mathcal{B}))$$

such that $\mathcal{A}(w) = u^\sharp$ (1.2.7). On the other hand, it follows from (1.5.2) that $\operatorname{Spec}(q^*(\mathcal{B}))$ canonically identifies with $X \times_S S'$; the morphism v is the composition $X' \xrightarrow{w} X \times_S S' \xrightarrow{p_1} X$ of w with the first projection, and the commutativity of (1.5.6.1) follows from the definitions. Let U (resp. U') be an affine open of S (resp. S') such that $q(U') \subset U$, $A = \Gamma(U, \mathcal{O}_S)$, $A' = \Gamma(U', \mathcal{O}_{S'})$ their rings, $B = \Gamma(U, \mathcal{B})$, $B' = \Gamma(U', \mathcal{B}')$; the restriction of u , as a $(q|U')$ -morphism $\mathcal{B}|U \rightarrow \mathcal{B}'|U'$, corresponds to a di-homomorphism of algebras $B \rightarrow B'$; if V, V' are the inverse images of U, U' in X, X' respectively, under the structure morphisms, then the morphism $V' \rightarrow V$, the restriction of v , corresponds (I, 1.7.3) to the above di-homomorphism.

(1.5.7). Under the same hypotheses as in (1.5.6), let $\mathcal{M}$ be a quasi-coherent $\mathcal{B}$ -module; there is then a canonical isomorphism of $\mathcal{O}_{X'}$ -modules

$$(1.5.7.1) \quad v^*(\widetilde{\mathcal{M}}) \simeq (q^*(\mathcal{M}) \otimes_{q^*(\mathcal{B})} \mathcal{B}')^\sim.$$

[II-14] Indeed, the canonical isomorphism (1.5.2.1) gives a canonical isomorphism from $p_1^*(\widetilde{\mathcal{M}})$ to the sheaf on $\operatorname{Spec}(q^*(\mathcal{B}))$ associated to the $q^*(\mathcal{B})$ -module $q^*(\mathcal{M})$, and it then suffices to apply (1.4.7).

1.6. Affine morphisms

(1.6.1). We say that a morphism $f : X \rightarrow Y$ of preschemes is *affine* if it defines X as a prescheme affine over Y . The properties of preschemes affine over another translate as follows in this language:

Proposition (1.6.2). —

- (i) A closed immersion is affine.
- (ii) The composition of two affine morphisms is affine.
- (iii) If $f : X \rightarrow Y$ is an affine S -morphism, then $f_{(S')} : X_{(S')} \rightarrow Y_{(S')}$ is affine for every base change $S' \rightarrow S$.
- (iv) If $f : X \rightarrow Y$ and $f' : X' \rightarrow Y'$ are two affine S -morphisms, then

$$f \times_S f' : X \times_S X' \longrightarrow Y \times_S Y'$$

is affine.

- (v) If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are two morphisms such that $g \circ f$ is affine and g is separated, then f is affine.
- (vi) If f is affine, then so is f_{red} .

Proof. By (I, 5.5.12), it suffices to prove (i), (ii), and (iii). But (i) is none other than Example (1.2.2), and (ii) is none other than Corollary (1.3.5); finally, (iii) follows from Proposition (1.5.1), since $X_{(S')}$ identifies with the product $X \times_Y Y_{(S')}$ (I, 3.3.11). $\square$

Corollary (1.6.3). — If X is an affine scheme and Y is a scheme, then every morphism $f : X \rightarrow Y$ is affine.

Proposition (1.6.4). — Let Y be a locally Noetherian prescheme, $f : X \rightarrow Y$ a morphism of finite type. For f to be affine, it is necessary and sufficient for f_{red} to be.

Proof. By (1.6.2, vi), we need only prove the sufficiency of the condition. It suffices to prove that if Y is affine and Noetherian, then X is affine; but Y_{red} is then affine, so the same is true for X_{red} by hypothesis. Now X is Noetherian, so the conclusion follows from (I, 6.1.7). $\square$

1.7. Vector bundle associated with a sheaf of modules

(1.7.1). Let A be a ring, E an A -module. Recall that the *symmetric algebra* of E , denoted by $\mathbf{S}(E)$ (or $\mathbf{S}_A(E)$), is the quotient of the tensor algebra $\mathbf{T}(E)$ by the two-sided ideal generated by the elements $x \otimes y - y \otimes x$, where x and y range over E . The algebra $\mathbf{S}(E)$ is characterized by the following universal property: if σ is the canonical map $E \rightarrow \mathbf{S}(E)$ (obtained by composing $E \rightarrow \mathbf{T}(E)$ with the canonical map $\mathbf{T}(E) \rightarrow \mathbf{S}(E)$), then every A -linear map $E \rightarrow B$, where B is a commutative A -algebra, factors uniquely as $E \xrightarrow{\sigma} \mathbf{S}(E) \xrightarrow{g} B$, where g is an A -homomorphism of algebras. We immediately deduce from this characterization that for two A -modules E and F , we have

$$\mathbf{S}(E \oplus F) = \mathbf{S}(E) \otimes \mathbf{S}(F)$$

[II:15] up to canonical isomorphism; in addition, $\mathbf{S}(E)$ is a covariant functor in E , from the category of A -modules to that of commutative A -algebras; finally, the above characterization also shows that if $E = \varinjlim E_\lambda$, then we have $\mathbf{S}(E) = \varinjlim \mathbf{S}(E_\lambda)$ up to canonical isomorphism. By abuse of language, a product $\sigma(x_1)\sigma(x_2)\cdots\sigma(x_n)$, where $x_i \in E$, is often denoted by $x_1x_2\cdots x_n$ if no confusion follows. The algebra $\mathbf{S}(E)$ is *graded*, $\mathbf{S}_n(E)$ being the set of linear combinations of products of n elements of E ($n \geq 0$); the algebra $\mathbf{S}(A)$ is canonically isomorphic to the polynomial algebra $A[T]$ in one indeterminate, and $\mathbf{S}(A^n)$ is canonically isomorphic to the polynomial algebra $A[T_1, \dots, T_n]$ in n indeterminates.

(1.7.2). Let ϕ be a ring homomorphism $A \rightarrow B$. If F is a B -module, then the canonical map $F \rightarrow \mathbf{S}(F)$ gives a canonical map $F_{[\phi]} \rightarrow \mathbf{S}(F)_{[\phi]}$, which thus factors as $F_{[\phi]} \rightarrow \mathbf{S}(F_{[\phi]}) \rightarrow \mathbf{S}(F)_{[\phi]}$; the canonical homomorphism $\mathbf{S}(F_{[\phi]}) \rightarrow \mathbf{S}(F)_{[\phi]}$ is surjective, but not necessarily bijective. If E is an A -module, then every di-homomorphism $E \rightarrow F$ (that is to say, every A -homomorphism $E \rightarrow F_{[\phi]}$) thus canonically gives an A -homomorphism of algebras $\mathbf{S}(E) \rightarrow \mathbf{S}(F_{[\phi]}) \rightarrow \mathbf{S}(F)_{[\phi]}$, that is to say a di-homomorphism of algebras $\mathbf{S}(E) \rightarrow \mathbf{S}(F)$.

With the same notations, for every A -module E , $\mathbf{S}(E \otimes_A B)$ canonically identifies with the algebra $\mathbf{S}(E) \otimes_A B$; this follows immediately from the universal property of $\mathbf{S}(E)$ (1.7.1).

(1.7.3). Let R be a multiplicative subset of the ring A . Applying (1.7.2) to the ring $B = R^{-1}A$ and recalling that $R^{-1}E = E \otimes_A R^{-1}A$, we obtain $\mathbf{S}(R^{-1}E) = R^{-1}\mathbf{S}(E)$ up to canonical isomorphism. In addition, if $R' \supset R$ is a second multiplicative subset of A , then the diagram

$$\begin{array}{ccc} R^{-1}E & \longrightarrow & R'^{-1}E \\ \downarrow & & \downarrow \\ \mathbf{S}(R^{-1}E) & \longrightarrow & \mathbf{S}(R'^{-1}E) \end{array}$$

is commutative.

(1.7.4). Now let $(S, \mathcal{A})$ be a ringed space, and let $\mathcal{E}$ be a $\mathcal{A}$ -module over S . By associating to each open subset $U \subset S$ the $\Gamma(U, \mathcal{A})$ -algebra $\mathbf{S}(\Gamma(U, \mathcal{E}))$, we obtain (by the functoriality of $\mathbf{S}$, (1.7.2)) a presheaf of algebras. Its associated sheaf, denoted by $\mathbf{S}(\mathcal{E})$ or $\mathbf{S}_{\mathcal{A}}(\mathcal{E})$, is called the *symmetric $\mathcal{A}$ -algebra* of the $\mathcal{A}$ -module $\mathcal{E}$. It follows immediately from (1.7.1) that $\mathbf{S}(\mathcal{E})$ is a solution to a universal problem: every homomorphism of $\mathcal{A}$ -modules $\mathcal{E} \rightarrow \mathcal{B}$, where $\mathcal{B}$ is an $\mathcal{A}$ -algebra, factors uniquely as $\mathcal{E} \rightarrow \mathbf{S}(\mathcal{E}) \rightarrow \mathcal{B}$, the second arrow being a homomorphism of $\mathcal{A}$ -algebras. There is thus a bijective correspondence between homomorphisms $\mathcal{E} \rightarrow \mathcal{B}$ of $\mathcal{A}$ -modules and homomorphisms $\mathbf{S}(\mathcal{E}) \rightarrow \mathcal{B}$ of $\mathcal{A}$ -algebras. In particular, every homomorphism $u : \mathcal{E} \rightarrow \mathcal{F}$ of $\mathcal{A}$ -modules defines a homomorphism $\mathbf{S}(u) : \mathbf{S}(\mathcal{E}) \rightarrow \mathbf{S}(\mathcal{F})$ of $\mathcal{A}$ -algebras, and $\mathbf{S}(\mathcal{E})$ is thus a covariant functor in $\mathcal{E}$.

[II:16] By (1.7.2) and the fact that $\mathbf{S}$ commutes with inductive limits, we have $(\mathbf{S}(\mathcal{E}))_x = \mathbf{S}(\mathcal{E}_x)$ for every point $x \in S$. If $\mathcal{E}, \mathcal{F}$ are two $\mathcal{A}$ -modules, then $\mathbf{S}(\mathcal{E} \oplus \mathcal{F})$ canonically identifies with $\mathbf{S}(\mathcal{E}) \otimes_{\mathcal{A}} \mathbf{S}(\mathcal{F})$, as we see for the corresponding presheaves.

We also note that $\mathbf{S}(\mathcal{E})$ is a graded $\mathcal{A}$ -algebra, the infinite direct sum of the $\mathbf{S}_n(\mathcal{E})$, where the $\mathcal{A}$ -module $\mathbf{S}_n(\mathcal{E})$ is the sheaf associated to the presheaf $U \mapsto \mathbf{S}_n(\Gamma(U, \mathcal{E}))$. Taking in particular $\mathcal{E} = \mathcal{A}$, we see that $\mathbf{S}_{\mathcal{A}}(\mathcal{A})$ identifies with $\mathcal{A}[T] = \mathcal{A} \otimes_{\mathbf{Z}} \mathbf{Z}[T]$ (T an indeterminate, and $\mathbf{Z}$ regarded as a constant sheaf).

(1.7.5). Let $(T, \mathcal{B})$ be a second ringed space, f a morphism $(S, \mathcal{A}) \rightarrow (T, \mathcal{B})$. If $\mathcal{F}$ is a $\mathcal{B}$ -module, then $\mathbf{S}(f^*(\mathcal{F}))$ canonically identifies with $f^*(\mathbf{S}(\mathcal{F}))$; indeed, if $f = (\psi, \theta)$, then by definition (0, 4.3.1),

$$\mathbf{S}(f^*(\mathcal{F})) = \mathbf{S}(\psi^*(\mathcal{F}) \otimes_{\psi^*(\mathcal{B})} \mathcal{A}) = \mathbf{S}(\psi^*(\mathcal{F})) \otimes_{\psi^*(\mathcal{B})} \mathcal{A}$$

(1.7.2); for every open U of S and every section h of $\mathbf{S}(\psi^*(\mathcal{F}))$ over U , h coincides, in a neighbourhood V of every point $s \in U$, with an element of $\mathbf{S}(\Gamma(V, \psi^*(\mathcal{F})))$; if we refer to the definition of $\psi^*(\mathcal{F})$ (0, 3.7.1) and take into account that every element of $\mathbf{S}(E)$ for a module E is a linear combination of a finite number of products of elements of E , then we see that there is a neighbourhood W of $\psi(s)$ in T , a section h' of $\mathbf{S}(\mathcal{F})$ over W , and a neighbourhood $V' \subset V \cap \psi^{-1}(W)$ of s such that h coincides with $t \mapsto h'(\psi(t))$ over V' ; hence our assertion.

Proposition (1.7.6). — *Let A be a ring, $S = \text{Spec}(A)$ its prime spectrum, $\mathcal{E} = \tilde{M}$ the $\mathcal{O}_S$ -module associated to an A -module M ; then the $\mathcal{O}_S$ -algebra $\mathbf{S}(\mathcal{E})$ is associated to the A -algebra $\mathbf{S}(M)$.*

Proof. For every $f \in A$, $\mathbf{S}(M_f) = (\mathbf{S}(M))_f$ (1.7.3), and the proposition thus follows from Definition (I, 1.3.4). $\square$

Corollary (1.7.7). — *If S is a prescheme, $\mathcal{E}$ a quasi-coherent $\mathcal{O}_S$ -module, then the $\mathcal{O}_S$ -algebra $\mathbf{S}(\mathcal{E})$ is quasi-coherent. If in addition $\mathcal{E}$ is of finite type, then each of the $\mathcal{O}_S$ -modules $\mathbf{S}_n(\mathcal{E})$ is of finite type.*

Proof. The first assertion is an immediate consequence of (1.7.6) and of (I, 1.4.1); the second follows from the fact that if E is an A -module of finite type, then $\mathbf{S}_n(E)$ is an A -module of finite type; we then apply (I, 1.3.13). $\square$

Definition (1.7.8). — Let $\mathcal{E}$ be a quasi-coherent $\mathcal{O}_S$ -module. We call the *vector bundle* over S defined by $\mathcal{E}$ and denote by $\mathbf{V}(\mathcal{E})$ the spectrum (1.3.1) of the quasi-coherent $\mathcal{O}_S$ -algebra $\mathbf{S}(\mathcal{E})$.

By (1.2.7), for every S -prescheme X , there is a canonical bijective correspondence between the S -morphisms $X \rightarrow \mathbf{V}(\mathcal{E})$ and the homomorphisms of $\mathcal{O}_S$ -algebras $\mathbf{S}(\mathcal{E}) \rightarrow \mathcal{A}(X)$, thus also between these S -morphisms and the homomorphisms of $\mathcal{O}_S$ -modules $\mathcal{E} \rightarrow \mathcal{A}(X) = f_*(\mathcal{O}_X)$ (where f is the structure morphism $X \rightarrow S$). In particular:

(1.7.9). Take X to be the subscheme of S induced on an open subset $U \subset S$. Then the S -morphisms $U \rightarrow \mathbf{V}(\mathcal{E})$ are none other than the U -sections (I, 2.5.5) of the U -prescheme induced by $\mathbf{V}(\mathcal{E})$ on the open $p^{-1}(U)$ (where p is the structure morphism $\mathbf{V}(\mathcal{E}) \rightarrow S$). From what we have just seen, these U -sections correspond bijectively to homomorphisms of $\mathcal{O}_S$ -modules $\mathcal{E} \rightarrow j_*(\mathcal{O}_S|U)$ (where j is the canonical injection $U \rightarrow S$), or ^[II:17] equivalently, by (0, 4.4.3), to the $(\mathcal{O}_S|U)$ -homomorphisms $j^*(\mathcal{E}) = \mathcal{E}|U \rightarrow \mathcal{O}_S|U$. In addition, it is immediate that the restriction to an open $U' \subset U$ of an S -morphism $U \rightarrow \mathbf{V}(\mathcal{E})$ corresponds to the restriction to U' of the corresponding homomorphism $\mathcal{E}|U \rightarrow \mathcal{O}_S|U$. We conclude that the sheaf of germs of S -sections of $\mathbf{V}(\mathcal{E})$ is canonically identified with the dual $\mathcal{E}^\vee$ of $\mathcal{E}$.

In particular, if we set $X = U = S$, then the zero homomorphism $\mathcal{E} \rightarrow \mathcal{O}_S$ corresponds to a canonical S -section of $\mathbf{V}(\mathcal{E})$, called the *zero S -section* (cf. (8.3.3)).

(1.7.10). Now take X to be the spectrum $\{\xi\}$ of a field K ; the structure morphism $f : X \rightarrow S$ then corresponds to a monomorphism $k(s) \rightarrow K$, where $s = f(\xi)$ (I, 2.4.6); the S -morphisms $\{\xi\} \rightarrow \mathbf{V}(\mathcal{E})$ are none other than the *geometric points of $\mathbf{V}(\mathcal{E})$ with values in the extension K of $k(s)$* (I, 3.4.5), lying over the points of $p^{-1}(s)$. The set of these points, which we can call the *rational geometric fibre over K of $\mathbf{V}(\mathcal{E})$ over the point s* , is identified by (1.7.8) with the set of homomorphisms of $\mathcal{O}_S$ -modules $\mathcal{E} \rightarrow f_*(\mathcal{O}_X)$, or, equivalently (0, 4.4.3) with the set of homomorphisms of $\mathcal{O}_X$ -modules $f^*(\mathcal{E}) \rightarrow \mathcal{O}_X = K$. But we have by definition (0, 4.3.1) $f^*(\mathcal{E}) = \mathcal{E}_s \otimes_{\mathcal{O}_s} K = \mathcal{E}^s \otimes_{k(s)} K$, setting $\mathcal{E}^s = \mathcal{E}_s / \mathfrak{m}_s \mathcal{E}_s$; the geometric fibre of $\mathbf{V}(\mathcal{E})$ rational over K over s thus identifies with the dual of the K -vector space $\mathcal{E}^s \otimes_{k(s)} K$; if $\mathcal{E}^s$ or K is of finite dimension over $k(s)$, then this dual also identifies with $(\mathcal{E}^s)^\vee \otimes_{k(s)} K$, denoting by $(\mathcal{E}^s)^\vee$ the dual of the $k(s)$ -vector space $\mathcal{E}^s$.

Proposition (1.7.11). —

- (i) $\mathbf{V}(\mathcal{E})$ is a contravariant functor in $\mathcal{E}$ from the category of quasi-coherent $\mathcal{O}_S$ -modules to the category of affine S -schemes.

- (ii) If $\mathcal{E}$ is an $\mathcal{O}_S$ -module of finite type, then $\mathbf{V}(\mathcal{E})$ is of finite type over S .
- (iii) If $\mathcal{E}$ and $\mathcal{F}$ are two quasi-coherent $\mathcal{O}_S$ -modules, then $\mathbf{V}(\mathcal{E} \oplus \mathcal{F})$ canonically identifies with $\mathbf{V}(\mathcal{E}) \times_S \mathbf{V}(\mathcal{F})$.
- (iv) Let $g : S' \rightarrow S$ be a morphism; for every quasi-coherent $\mathcal{O}_S$ -module $\mathcal{E}$, $\mathbf{V}(g^*(\mathcal{E}))$ canonically identifies with $\mathbf{V}(\mathcal{E})_{(S')} = \mathbf{V}(\mathcal{E}) \times_S S'$.
- (v) A surjective homomorphism $\mathcal{E} \rightarrow \mathcal{F}$ of quasi-coherent $\mathcal{O}_S$ -modules corresponds to a closed immersion $\mathbf{V}(\mathcal{F}) \rightarrow \mathbf{V}(\mathcal{E})$.

Proof. (i) is an immediate consequence of (1.2.7), taking into account that every homomorphism of $\mathcal{O}_S$ -modules $\mathcal{E} \rightarrow \mathcal{F}$ canonically defines a homomorphism of $\mathcal{O}_S$ -algebras $\mathbf{S}(\mathcal{E}) \rightarrow \mathbf{S}(\mathcal{F})$. (ii) follows immediately from the definition (I, 6.3.1) and the fact that if E is an A -module of finite type, then $\mathbf{S}(E)$ is an A -algebra of finite type. To prove (iii), it suffices to start with the canonical isomorphism $\mathbf{S}(\mathcal{E} \oplus \mathcal{F}) \simeq \mathbf{S}(\mathcal{E}) \otimes_{\mathcal{O}_S} \mathbf{S}(\mathcal{F})$ (1.7.4) and to apply (1.4.6). Similarly, to prove (iv), it suffices to start with the canonical isomorphism $\mathbf{S}(g^*(\mathcal{E})) \simeq g^*(\mathbf{S}(\mathcal{E}))$ (1.7.5) and to apply (1.5.2). Finally, to establish (v), it suffices to remark that if the homomorphism $\mathcal{E} \rightarrow \mathcal{F}$ is surjective, then so is the corresponding homomorphism $\mathbf{S}(\mathcal{E}) \rightarrow \mathbf{S}(\mathcal{F})$ of $\mathcal{O}_S$ -algebras, and the conclusion follows from (1.4.10). $\square$

(1.7.12). Take in particular $\mathcal{E} = \mathcal{O}_S$; the prescheme $\mathbf{V}(\mathcal{O}_S)$ is the affine S -scheme that is the spectrum of the $\mathcal{O}_S$ -algebra $\mathbf{S}(\mathcal{O}_S)$, which identifies with the $\mathcal{O}_S$ -algebra $\mathcal{O}_S[T] = \mathcal{O}_S \otimes_{\mathbf{Z}} \mathbf{Z}[T]$ [II:18] (T indeterminate); this is evident when $S = \text{Spec}(\mathbf{Z})$, by virtue of (1.7.6), and we pass from there to the general case by considering the structure morphism $S \rightarrow \text{Spec}(\mathbf{Z})$ and using (1.7.11, iv). Because of this result, we set $\mathbf{V}(\mathcal{O}_S) = S[T]$, and we thus have the formula

$$(1.7.12.1) \quad S[T] = S \otimes_{\mathbf{Z}} \mathbf{Z}[T].$$

The identification of the sheaf of germs of S -sections of $S[T]$ with $\mathcal{O}_S$, already seen in (I, 3.3.15), reappears here in a more general context as a special case of (1.7.9).

(1.7.13). For every S -prescheme X , we have seen (1.7.8) that $\text{Hom}_S(X, S[T])$ canonically identifies with $\text{Hom}_{\mathcal{O}_S}(\mathcal{O}_S, \mathcal{A}(X))$, which is canonically isomorphic to $\Gamma(S, \mathcal{A}(X))$, and is therefore equipped with a ring structure; in addition, every S -morphism $h : X \rightarrow Y$ induces a ring homomorphism $\Gamma(\mathcal{A}(h)) : \Gamma(S, \mathcal{A}(Y)) \rightarrow \Gamma(S, \mathcal{A}(X))$ (1.1.2). With the ring structure just defined on $\text{Hom}_S(X, S[T])$, we may therefore regard $\text{Hom}_S(X, S[T])$ as a *contravariant functor* in X , from the category of S -preschemes to that of rings. On the other hand, $\text{Hom}_S(X, \mathbf{V}(\mathcal{E}))$ is likewise identified with $\text{Hom}_{\mathcal{O}_S}(\mathcal{E}, \mathcal{A}(X))$ (where $\mathcal{A}(X)$ is considered as an $\mathcal{O}_S$ -module); as a result, we can canonically equip it with a *module* structure over the ring $\text{Hom}_S(X, S[T])$, and we see as above that the pair

$$(\text{Hom}_S(X, S[T]), \text{Hom}_S(X, \mathbf{V}(\mathcal{E})))$$

is a contravariant functor in X , with values in the category whose objects are the pairs (A, M) consisting of a ring A and an A -module M , the morphisms being di-homomorphisms.

We will interpret these facts by saying that $S[T]$ is an *S-scheme of rings* and that $\mathbf{V}(\mathcal{E})$ is an *S-scheme of modules* on the S -scheme of rings $S[T]$ (cf. Chapter 0, §8).

(1.7.14). We will see that the structure of an S -scheme of modules defined on the S -scheme $\mathbf{V}(\mathcal{E})$ allows us to reconstruct the $\mathcal{O}_S$ -module $\mathcal{E}$ up to unique isomorphism: for this, we will show that $\mathcal{E}$ is canonically isomorphic to an $\mathcal{O}_S$ -submodule of $\mathbf{S}(\mathcal{E}) = \mathcal{A}(\mathbf{V}(\mathcal{E}))$, defined by means of this structure. Indeed, by (1.7.4), the set $\text{Hom}_{\mathcal{O}_S}(\mathbf{S}(\mathcal{E}), \mathcal{A}(X))$ of homomorphisms of $\mathcal{O}_S$ -algebras is canonically identified with $\text{Hom}_{\mathcal{O}_S}(\mathcal{E}, \mathcal{A}(X))$, the set of homomorphisms of $\mathcal{O}_S$ -modules: if h and h' are two elements of this latter set, s_i ($1 \leq i \leq n$) sections of $\mathcal{E}$ over an open $U \subset S$, t a section of $\mathcal{A}(X)$ over U , then we have by definition

$$(h + h')(s_1 s_2 \cdots s_n) = \prod_{i=1}^n (h(s_i) + h'(s_i))$$

and

$$(t \cdot h)(s_1 s_2 \cdots s_n) = t^n \prod_{i=1}^n h(s_i).$$

This being so, if z is a section of $\mathbf{S}(\mathcal{E})$ over U , then $h \mapsto h(z)$ is a map from $\text{Hom}_S(X, \mathbf{V}(\mathcal{E})) = \text{Hom}_{\mathcal{O}_S}(\mathbf{S}(\mathcal{E}), \mathcal{A}(X))$ to $\Gamma(U, \mathcal{A}(X))$. We will ^[II:19] show that $\mathcal{E}$ is identified with a submodule of $\mathbf{S}(\mathcal{E})$ such that, for every open $U \subset S$, every section z of this $\mathcal{O}_S$ -submodule over U , and every S -prescheme X , the map $h \mapsto h(z)$ from $\text{Hom}_{\mathcal{O}_S}(\mathbf{S}(\mathcal{E})|_U, \mathcal{A}(X)|_U)$ to $\Gamma(U, \mathcal{A}(X))$ is a homomorphism of $\Gamma(U, \mathcal{A}(X))$ -modules.

It is immediate that $\mathcal{E}$ has this property; to show the converse, we can reduce to proving that when $S = \text{Spec}(A)$, $\mathcal{E} = \tilde{M}$, a section z of $\mathbf{S}(\mathcal{E})$ over S that (for $U = S$) has the property stated above is necessarily a section of $\mathcal{E}$; we then have $z = \sum_{n=0}^{\infty} z_n$, where $z_n \in \mathbf{S}_n(M)$, and it is a question of proving that $z_n = 0$ for $n \neq 1$. Set $B = \mathbf{S}(M)$ and take for X the prescheme $\text{Spec}(B[T])$, where T is an indeterminate. The set $\text{Hom}_{\mathcal{O}_S}(\mathbf{S}(\mathcal{E}), \mathcal{A}(X))$ identifies with the set of ring homomorphisms $h : B \rightarrow B[T]$ (I, 1.3.13), and from what we saw above, we have $(T \cdot h)(z) = \sum_{n=0}^{\infty} T^n h(z_n)$: the hypothesis on z implies that we have $\sum_{n=0}^{\infty} T^n h(z_n) = T \cdot \sum_{n=0}^{\infty} h(z_n)$ for every homomorphism h . Taking h in particular to be the canonical injection, we obtain $\sum_{n=0}^{\infty} T^n z_n = T \cdot \sum_{n=0}^{\infty} z_n$, which gives the desired conclusion $z_n = 0$ for $n \neq 1$.

Proposition (1.7.15). — *Let Y be a prescheme whose underlying space is Noetherian, or a quasi-compact scheme. Every affine Y -scheme X of finite type over Y is Y -isomorphic to a closed Y -subscheme of a Y -scheme of the form $\mathbf{V}(\mathcal{E})$, where $\mathcal{E}$ is a quasi-coherent $\mathcal{O}_Y$ -module of finite type.*

Proof. The quasi-coherent $\mathcal{O}_Y$ -algebra $\mathcal{A}(X)$ is of finite type (1.3.7). The hypotheses imply that $\mathcal{A}(X)$ is generated by a quasi-coherent $\mathcal{O}_Y$ -submodule of finite type $\mathcal{E}$ (I, 9.6.5); by definition, this implies that the canonical homomorphism $\mathbf{S}(\mathcal{E}) \rightarrow \mathcal{A}(X)$ canonically extending the injection $\mathcal{E} \rightarrow \mathcal{A}(X)$ is surjective; the conclusion then follows from (1.4.10). $\square$

§2. HOMOGENEOUS PRIME SPECTRA

2.1. Generalities on graded rings and modules

Notation (2.1.1). — Given a positively graded ring S , we denote by S_n the subset of S consisting of homogeneous elements of degree n ($n \geq 0$), by S_+ the (direct) sum of the S_n for $n > 0$; we have $1 \in S_0$, S_0 is a subring of S , S_+ is a graded ideal of S , and S is the direct sum of S_0 and S_+ . If M is a graded module over S (with positive or negative degrees), we similarly denote by M_n the S_0 -module consisting of homogeneous elements of M of degree n (with $n \in \mathbf{Z}$).

For every integer $d > 0$, we denote by $S^{(d)}$ the direct sum of the S_{nd} ; by considering the elements of S_{nd} as homogeneous of degree n , the S_{nd} define on $S^{(d)}$ a graded ring structure.

For every integer k such that $0 \leq k \leq d-1$, we denote by $M^{(d,k)}$ the direct sum ^[II:20] of the M_{nd+k} ($n \in \mathbf{Z}$); this is a graded $S^{(d)}$ -module when we consider the elements of M_{nd+k} as homogeneous of degree n . We write $M^{(d)}$ in place of $M^{(d,0)}$.

With the above notation, for every integer n (positive or negative), we denote by $M(n)$ the graded S -module defined by $(M(n))_k = M_{n+k}$ for every $k \in \mathbf{Z}$. In particular, $S(n)$ will be a graded S -module such that $(S(n))_k = S_{n+k}$, by agreeing to set $S_n = 0$ for $n < 0$. We say that a graded S -module M is *free* if it is isomorphic, as a graded module, to a direct sum of modules of the form $S(n)$; since $S(n)$ is a cyclic S -module generated by the element 1 of S , regarded as an element of degree $-n$, this is equivalent to saying that M admits a *basis* over S consisting of *homogeneous* elements.

We say that a graded S -module M *admits a finite presentation* if there exists an exact sequence $P \rightarrow Q \rightarrow M \rightarrow 0$, where P and Q are finite direct sums of modules of the form $S(n)$ and the homomorphisms are of degree 0 (cf. (2.1.2)).

(2.1.2). Let M and N be two graded S -modules; we define on $M \otimes_S N$ a graded S -module structure in the following way. On the tensor product $M \otimes_{\mathbf{Z}} N$, we can define a graded $\mathbf{Z}$ -module structure (where $\mathbf{Z}$ is graded by $\mathbf{Z}_0 = \mathbf{Z}$, $\mathbf{Z}_n = 0$ for $n \neq 0$) by setting $(M \otimes_{\mathbf{Z}} N)_q = \bigoplus_{m+n=q} M_m \otimes_{\mathbf{Z}} N_n$ (as M and N are respectively direct sums of the M_m and the N_n , we know that we can canonically identify $M \otimes_{\mathbf{Z}} N$ with the direct sum of all the $M_m \otimes_{\mathbf{Z}} N_n$). This being so, we have $M \otimes_S N = (M \otimes_{\mathbf{Z}} N)/P$, where P is the $\mathbf{Z}$ -submodule of $M \otimes_{\mathbf{Z}} N$ generated by the elements $(xs) \otimes y - x \otimes (sy)$ for $x \in M$, $y \in N$, $s \in S$; it is clear that P is a graded $\mathbf{Z}$ -submodule of $M \otimes_{\mathbf{Z}} N$, and we see immediately that we obtain a graded S -module structure on $M \otimes_S N$ by passing to the quotient.

For two graded S -modules M and N , recall that a homomorphism $u : M \rightarrow N$ of S -modules is said to be of degree k if $u(M_j) \subset N_{j+k}$ for all $j \in \mathbf{Z}$. If H_n denotes the set of all the homomorphisms of degree n from M to N , then we denote by $\text{Hom}_S(M, N)$ the (direct) sum of the H_n ($n \in \mathbf{Z}$) in the S -module H of all the homomorphisms (of S -modules) from M to N ; in general, $\text{Hom}_S(M, N)$ is not equal to the latter. However, we have $H = \text{Hom}_S(M, N)$ when M is of finite type; indeed, we can then suppose that M is generated by a finite number of homogeneous elements x_i ($1 \leq i \leq n$), and every homomorphism $u \in H$ can be written in a unique way as $\sum_{k \in \mathbf{Z}} u_k$, where for each k , $u_k(x_i)$ is equal to the homogeneous component of degree $k + \deg(x_i)$ of $u(x_i)$ ($1 \leq i \leq n$), which implies that $u_k = 0$ except for a finite number of indices; we have by definition that $u_k \in H_k$, hence the conclusion.

The elements of degree 0 of $\text{Hom}_S(M, N)$ are called the *homomorphisms of graded S -modules*. It is clear that $S_m H_n \subset H_{m+n}$, so the H_n define on $\text{Hom}_S(M, N)$ a graded S -module structure.

It follows immediately from these definitions that we have

$$(2.1.2.1) \quad M(m) \otimes_S N(n) = (M \otimes_S N)(m+n),$$

$$(2.1.2.2) \quad \text{Hom}_S(M(m), N(n)) = (\text{Hom}_S(M, N))(n-m),$$

for two graded S -modules M and N .

[II:21] Let S and S' be two graded rings; a homomorphism of graded rings $\phi : S \rightarrow S'$ is a homomorphism of rings such that $\phi(S_n) \subset S'_n$ for all $n \in \mathbf{Z}$ (in other words, ϕ must be a homomorphism of degree 0 of graded $\mathbf{Z}$ -modules). Such a homomorphism endows S' with a graded S -module structure; equipped with this structure and with its graded ring structure, S' is called a *graded S -algebra*.

If M is also a graded S -module, then the tensor product $M \otimes_S S'$ of graded S -modules is equipped in a natural way with a graded S' -module structure, the grading being defined as above.

Lemma (2.1.3). — *Let S be a ring graded in positive degrees. For a subset E of S_+ consisting of homogeneous elements to generate S_+ as an S -module, it is necessary and sufficient for E to generate S as an S_0 -algebra.*

Proof. The condition is evidently sufficient; we show that it is necessary. Let E_n (resp. E^n) be the set of elements of E of degree equal to n (resp. of degree $\leq n$); it suffices to show, by induction on $n > 0$, that S_n is the S_0 -module generated by the elements of degree n which are products of elements of E^n . This is evident for $n = 1$ by virtue of the hypothesis; the latter also shows that $S_n = \sum_{p=0}^{n-1} S_p E_{n-p}$, and the induction argument is then immediate. $\square$

Corollary (2.1.4). — *For S_+ to be an ideal of finite type, it is necessary and sufficient for S to be an S_0 -algebra of finite type.*

Proof. We can always assume that a finite system of generators of the S_0 -algebra S (resp. of the S -ideal S_+) consists of homogeneous elements, by replacing each of the generators considered by its homogeneous components. $\square$

Corollary (2.1.5). — *For S to be Noetherian, it is necessary and sufficient for S_0 to be Noetherian and for S to be an S_0 -algebra of finite type.*

Proof. The condition is evidently sufficient; it is necessary, since S_0 is isomorphic to S/S_+ and S_+ must be an ideal of finite type (2.1.4). $\square$

Lemma (2.1.6). — *Let S be a ring graded in positive degrees, which is an S_0 -algebra of finite type. Let M be a graded S -module of finite type. Then:*

- (i) *The M_n are S_0 -modules of finite type, and there exists an integer n_0 such that $M_n = 0$ for $n \leq n_0$.*
- (ii) *There exists an integer n_1 and an integer $h > 0$ such that, for every integer $n \geq n_1$, we have $M_{n+h} = S_h M_n$.*
- (iii) *For every pair of integers (d, k) such that $d > 0$, $0 \leq k \leq d-1$, $M^{(d,k)}$ is an $S^{(d)}$ -module of finite type.*
- (iv) *For every integer $d > 0$, $S^{(d)}$ is an S_0 -algebra of finite type.*
- (v) *There exists an integer $h > 0$ such that $S_{mh} = (S_h)^m$ for all $m > 0$.*
- (vi) *For every integer $n > 0$, there exists an integer m_0 such that $S_m \subset S_+^n$ for all $m \geq m_0$.*

Proof. We can assume that S is generated (as an S_0 -algebra) by homogeneous elements f_i , of degrees h_i ($1 \leq i \leq r$), and M is generated (as an S -module) by homogeneous elements x_j of degrees k_j ($1 \leq j \leq s$). It is clear that M_n is formed by linear combinations, ^[II:22] with coefficients in S_0 , of elements $f_1^{\alpha_1} \cdots f_r^{\alpha_r} x_j$ such that the α_i are integers ≥ 0 satisfying $k_j + \sum_i \alpha_i h_i = n$; for each j , there are only finitely many systems (α_i) satisfying this equation, since the h_i are > 0 , hence the first assertion of (i); the second is evident. On the other hand, let h be the l.c.m. of the h_i and set $g_i = f_i^{h/h_i}$ ($1 \leq i \leq r$), so that all the g_i are of degree h ; let z_μ be the elements of M of the form $f_1^{\alpha_1} \cdots f_r^{\alpha_r} x_j$ with $0 \leq \alpha_i < h/h_i$ for $1 \leq i \leq r$; there are finitely many of these elements, so let n_1 be the largest of their degrees. It is clear that for $n \geq n_1$, every element of M_{n+h} is a linear combination of the z_μ whose coefficients are monomials of positive degree in the g_i , so we have $M_{n+h} = S_h M_n$, which establishes (ii). In a similar way, we see (for all $d > 0$) that an element of $M^{(d,k)}$ is a linear combination, with coefficients in S_0 , of elements of the form $g^d f_1^{\alpha_1} \cdots f_r^{\alpha_r} x_j$ with $0 \leq \alpha_i < d$, where g is a homogeneous element of S ; hence (iii); (iv) then follows from (iii) and from Lemma (2.1.3), by taking $M = S_+$, since $(S_+)^{(d)} = (S^{(d)})_+$. The assertion of (v) is deduced from (ii) by taking $M = S$. Finally, for a given n , there are finitely many systems (α_i) such that $\alpha_i \geq 0$ and $\sum_i \alpha_i h_i < n$; if m_0 is chosen strictly larger than the greatest value of $\sum_i \alpha_i h_i$ among these systems, then $S_m \subset S_+^n$ for $m \geq m_0$, which proves (vi). $\square$

Corollary (2.1.7). — *If S is Noetherian, then so is $S^{(d)}$ for every integer $d > 0$.*

Proof. This follows from (2.1.5) and (2.1.6, iv). $\square$

(2.1.8). Let $\mathfrak{p}$ be a *graded* prime ideal of the graded ring S ; $\mathfrak{p}$ is thus a direct sum of the subgroups $\mathfrak{p}_n = \mathfrak{p} \cap S_n$. Suppose that $\mathfrak{p}$ does not contain S_+ . Then if $f \in S_+$ is not in $\mathfrak{p}$, the relation $f^n x \in \mathfrak{p}$ is equivalent to $x \in \mathfrak{p}$; in particular, if $f \in S_d$ ($d > 0$), for all $x \in S_{m-nd}$, then the relation $f^n x \in \mathfrak{p}_m$ is equivalent to $x \in \mathfrak{p}_{m-nd}$.

Proposition (2.1.9). — *Let n_0 be an integer > 0 ; for all $n \geq n_0$, let $\mathfrak{p}_n$ be a subgroup of S_n . For there to exist a graded prime ideal $\mathfrak{p}$ of S not containing S_+ and such that $\mathfrak{p} \cap S_n = \mathfrak{p}_n$ for all $n \geq n_0$, it is necessary and sufficient for the following conditions to be satisfied:*

- (1st) $S_m \mathfrak{p}_n \subset \mathfrak{p}_{m+n}$ for all $m \geq 0$ and all $n \geq n_0$.
- (2nd) For $m \geq n_0$, $n \geq n_0$, $f \in S_m$, $g \in S_n$, the relation $fg \in \mathfrak{p}_{m+n}$ implies $f \in \mathfrak{p}_m$ or $g \in \mathfrak{p}_n$.
- (3rd) $\mathfrak{p}_n \neq S_n$ for at least one $n \geq n_0$.

In addition, the graded prime ideal $\mathfrak{p}$ is then unique.

Proof. It is evident that the conditions (1st) and (2nd) are necessary. In addition, if $\mathfrak{p} \not\supset S_+$, then there exists at least one $k > 0$ such that $\mathfrak{p} \cap S_k \neq S_k$; if $f \in S_k$ is not in $\mathfrak{p}$, the relation $\mathfrak{p} \cap S_n = S_n$ implies $\mathfrak{p} \cap S_{n-mk} = S_{n-mk}$ according to (2.1.8); therefore, if $\mathfrak{p} \cap S_n = S_n$ for all sufficiently large n , we would have $\mathfrak{p} \supset S_+$, contrary to the hypothesis, which proves that (3rd) is necessary. Conversely, suppose that the conditions (1st), (2nd), and (3rd) are satisfied. Note that if for an integer $d \geq n_0$, $f \in S_d$ is not in $\mathfrak{p}_d$, then, if $\mathfrak{p}$ exists, $\mathfrak{p}_m$, for $m < n_0$, is necessarily equal to the set of the $x \in S_m$ such that $f^r x \in \mathfrak{p}_{m+rd}$, except for a finite number of values of r . This already proves that if $\mathfrak{p}$ exists, then it is unique. It remains to show that if we define the $\mathfrak{p}_m$ for $m < n_0$ by the previous condition, then $\mathfrak{p} = \sum_{n=0}^{\infty} \mathfrak{p}_n$ is a prime ideal. First, note that by virtue of (2nd), for $m \geq n_0$, $\mathfrak{p}_m$ is also defined as the set of the $x \in S_m$ such that $f^r x \in \mathfrak{p}_{m+rd}$ except for a finite number of values of r . This ^[II:23] being so, if $g \in S_m$, $x \in \mathfrak{p}_n$, then we have $f^r g x \in \mathfrak{p}_{m+n+rd}$ except for a finite number of values of r , so $gx \in \mathfrak{p}_{m+n}$, which proves that $\mathfrak{p}$ is an ideal of S . To establish that this ideal is prime, in other words that the ring $S/\mathfrak{p}$, graded by the subgroups $S_n/\mathfrak{p}_n$, is an integral domain, it suffices (by considering the highest-degree components of two elements of $S/\mathfrak{p}$) to prove that if $x \in S_m$ and $y \in S_n$ are such that $x \notin \mathfrak{p}_m$ and $y \notin \mathfrak{p}_n$, then $xy \notin \mathfrak{p}_{m+n}$. If not, for r large enough, we would have $f^{2r} xy \in \mathfrak{p}_{m+n+2rd}$; but we have $f^r y \notin \mathfrak{p}_{n+rd}$ for all $r > 0$; it then follows from (2nd) that, except for a finite number of values of r , we have $f^r x \in \mathfrak{p}_{m+rd}$, and we conclude that $x \in \mathfrak{p}_m$ contrary to the hypothesis. $\square$

(2.1.10). We say that a subset $\mathfrak{J}$ of S_+ is an *ideal* of S_+ if it is an ideal of S , and $\mathfrak{J}$ is a *graded prime ideal* of S_+ if it is the intersection of S_+ and a graded prime ideal of S not containing S_+ (this prime ideal is also unique according to Proposition (2.1.9)). If $\mathfrak{J}$ is an ideal of S_+ , the *radical* of $\mathfrak{J}$ in S_+ is

the set of elements of S_+ which have a power in $\mathfrak{J}$, in other words the set $\tau_+(\mathfrak{J}) = \tau(\mathfrak{J}) \cap S_+$; in particular, the radical of 0 in S_+ is then called the *nilradical* of S_+ and denoted by $\mathfrak{N}_+$: this is the set of nilpotent elements of S_+ . If $\mathfrak{J}$ is an *graded* ideal of S_+ , then its radical $\tau_+(\mathfrak{J})$ is a *graded* ideal: by passing to the quotient ring $S/\mathfrak{J}$, we can reduce to the case $\mathfrak{J} = 0$, and it remains to see that if $x = x_h + x_{h+1} + \cdots + x_k$ is nilpotent, then so are the $x_i \in S_i$ ($1 \leq h \leq i \leq k$); we can assume $x_k \neq 0$ and the component of highest degree of x^n is then x_k^n , hence x_k is nilpotent, and we then argue by induction on k . We say that the graded ring S is *essentially reduced* if $\mathfrak{N}_+ = 0$, in other words, if S_+ does not contain nilpotent elements $\neq 0$.

(2.1.11). We note that if, in the graded ring S , an element x is a zero-divisor, then so is its component of highest degree. We say that a ring S is *essentially integral* if the ring S_+ (without a unit element) contains no zero-divisors and is nonzero; for this, it is enough that no nonzero homogeneous element of S_+ be a zero-divisor in this ring. It is clear that if $\mathfrak{p}$ is a graded prime ideal of S_+ , then $S/\mathfrak{p}$ is essentially integral.

Let S be an essentially integral graded ring, and let $x_0 \in S_0$: if there then exists a homogeneous element $f \neq 0$ of S_+ such that $x_0 f = 0$, then we have $x_0 S_+ = 0$, since we have $(x_0 g)f = (x_0 f)g = 0$ for all $g \in S_+$, and the hypothesis thus implies $x_0 g = 0$. For S to be integral, it is necessary and sufficient for S_0 to be integral and the annihilator of S_+ in S_0 to be 0.

2.2. Rings of fractions of a graded ring

(2.2.1). Let S be a graded ring, in positive degrees, f a *homogeneous* element of S , of degree $d > 0$; then the ring of fractions $S' = S_f$ is graded, taking for S'_n the set of the x/f^k , where $x \in S_{n+kd}$ with $k \geq 0$ (we observe here that n can take arbitrary negative values); we denote the subring $S'_0 = (S_f)_0$ of S' consisting of elements of degree 0 by the notation $S_{(f)}$.

If $f \in S_d$, then the monomials $(f/1)^h$ in S_f (h a positive or negative integer) form a *free system* over the ring $S_{(f)}$, and the set of their linear combinations is none other than ^[II.24] the ring $(S^{(d)})_f$, which is thus *isomorphic* to $S_{(f)}[T, T^{-1}] = S_{(f)} \otimes_{\mathbb{Z}} \mathbb{Z}[T, T^{-1}]$ (where T is an indeterminate). Indeed, if we have a relation $\sum_{h=-a}^b z_h (f/1)^h = 0$ with $z_h = x_h/f^m$, where the x_h are in S_{md} , then this relation is equivalent by definition to the existence of a $k > -a$ such that $\sum_{h=-a}^b f^{h+k} x_h = 0$, and as the degrees of the terms of this sum are distinct, we have $f^{h+k} x_h = 0$ for all h , hence $z_h = 0$ for all h .

If M is a graded S -module, then $M' = M_f$ is a graded S_f -module, M'_n being the set of the z/f^k with $z \in M_{n+kd}$ ($k \geq 0$); we denote by $M_{(f)}$ the set of the homogeneous elements of degree 0 of M' ; it is immediate that $M_{(f)}$ is an $S_{(f)}$ -module and that we have $(M^{(d)})_f = M_{(f)} \otimes_{S_{(f)}} (S^{(d)})_f$.

Lemma (2.2.2). — *Let d and e be integers > 0 , $f \in S_d$, $g \in S_e$. There exists a canonical ring isomorphism*

$$S_{(fg)} \simeq (S_{(f)})_{g^d/f^e};$$

if we canonically identify these two rings, then there exists a canonical module isomorphism

$$M_{(fg)} \simeq (M_{(f)})_{g^d/f^e}.$$

Proof. Indeed, fg divides $f^e g^d$, and this latter element divides $(fg)^{de}$, so the graded rings S_{fg} and $S_{f^e g^d}$ are canonically identified; on the other hand, $S_{f^e g^d}$ also identifies with $(S_{f^e})_{g^d/f^e}$ (0, 1.4.6), and as $f^e/1$ is invertible in S_{f^e} , $S_{f^e g^d}$ also identifies with $(S_{f^e})_{g^d/f^e}$. The element g^d/f^e is of degree 0 in S_{f^e} ; we immediately conclude that the subring of $(S_{f^e})_{g^d/f^e}$ consisting of elements of degree 0 is $(S_{(f^e)})_{g^d/f^e}$, and as we evidently have $S_{(f^e)} = S_{(f)}$, this proves the first part of the lemma; the second is established in a similar way. $\square$

(2.2.3). Under the hypotheses of (2.2.2), it is clear that the canonical homomorphism $S_f \rightarrow S_{fg}$ (0, 1.4.1), which sends x/f^k to $g^k x/(fg)^k$, is of degree 0, thus gives by restriction a *canonical homomorphism* $S_{(f)} \rightarrow S_{(fg)}$, such that the diagram

$$\begin{array}{ccc} & S_{(f)} & \\ \swarrow & & \searrow \\ S_{(fg)} & \xrightarrow{\sim} & (S_{(f)})_{g^d/f^e} \end{array}$$

is commutative. We similarly define a canonical homomorphism $M_{(f)} \rightarrow M_{(fg)}$.

Lemma (2.2.4). — *If f and g are two homogeneous elements of S_+ , then the ring $S_{(fg)}$ is generated by the union of the canonical images of $S_{(f)}$ and $S_{(g)}$.*

Proof. By virtue of Lemma (2.2.2), it suffices to see that $1/(g^d/f^e) = f^{d+e}/(fg)^d$ belongs to the canonical image of $S_{(g)}$ in $S_{(fg)}$, which is evident by definition. $\square$

Proposition (2.2.5). — *Let d be an integer > 0 and let $f \in S_d$. Then there exists a canonical ring isomorphism $S_{(f)} \simeq S^{(d)}/(f-1)S^{(d)}$; if we identify these two rings by this isomorphism, then there exists a canonical module isomorphism $M_{(f)} \simeq M^{(d)}/(f-1)M^{(d)}$.*

Proof. The first of these isomorphisms is defined by sending x/f^n , where $x \in S_{nd}$, to the element $\bar{x}$, the class of x mod. $(f-1)S^{(d)}$; this map is well-defined, because we have the congruence $f^h x \equiv x \pmod{(f-1)S^{(d)}}$ for all $x \in S^{(d)}$, so if $f^h x = 0$ for an $h > 0$, [III:25] then we have $\bar{x} = 0$. On the other hand, if $x \in S_{nd}$ is such that $x = (f-1)y$ with $y = y_{hd} + y_{(h+1)d} + \cdots + y_{kd}$ with $y_{jd} \in S_{jd}$ and $y_{hd} \neq 0$, then we necessarily have $h = n$ and $x = -y_{hd}$, as well as the relations $y_{(j+1)d} = f y_{jd}$ for $h \leq j \leq k-1$, $f y_{kd} = 0$, which ultimately gives $f^{k-n+1}x = 0$; we send every class $\bar{x}$ mod. $(f-1)S^{(d)}$ of an element $x \in S_{nd}$ to the element x/f^n of $S_{(f)}$, since the preceding remark shows that this map is well-defined. It is immediate that these two maps thus defined are ring homomorphisms, each the reciprocal of the other. We proceed exactly the same way for M . $\square$

Corollary (2.2.6). — *If S is Noetherian, then so is $S_{(f)}$ for f homogeneous of degree > 0 .*

Proof. This follows immediately from Corollary (2.1.7) and Proposition (2.2.5). $\square$

(2.2.7). Let T be a multiplicative subset of S_+ consisting of *homogeneous* elements; $T_0 = T \cup \{1\}$ is then a multiplicative subset of S ; as the elements of T_0 are homogeneous, the ring $T_0^{-1}S$ is still graded in the evident way; we denote by $S_{(T)}$ the subring of $T_0^{-1}S$ consisting of elements of degree 0, that is to say, the elements of the form x/h , where $h \in T$ and x is homogeneous of degree equal to that of h . We know from (0, 1.4.5) that $T_0^{-1}S$ is canonically identified with the inductive limit of the rings S_f , where f varies over T (with respect to the canonical homomorphisms $S_f \rightarrow S_{fg}$); as this identification respects the degrees, it identifies $S_{(T)}$ with the *inductive limit* of the $S_{(f)}$ for $f \in T$. For every graded S -module M , we similarly define the module $M_{(T)}$ (over the ring $S_{(T)}$) consisting of elements of degree 0 of $T_0^{-1}M$, and we see that this module is the inductive limit of the $M_{(f)}$ for $f \in T$.

If $\mathfrak{p}$ is a graded prime ideal of S_+ , then we denote by $S_{(\mathfrak{p})}$ and $M_{(\mathfrak{p})}$ the ring $S_{(T)}$ and the module $M_{(T)}$ respectively, where T is the set of *homogeneous* elements of S_+ which do not belong to $\mathfrak{p}$.

2.3. Homogeneous prime spectrum of a graded ring

(2.3.1). Given a graded ring S , in positive degrees, we call the *homogeneous prime spectrum* of S and denote it by $\text{Proj}(S)$ the set of graded prime ideals of S_+ (2.1.10), or equivalently the set of graded prime ideals of S not containing S_+ ; we will define a *scheme* structure having $\text{Proj}(S)$ as the underlying set.

(2.3.2). For every subset E of S , let $V_+(E)$ be the set of graded prime ideals of S containing E and not containing S_+ ; this is thus the subset $V(E) \cap \text{Proj}(S)$ of $\text{Spec}(S)$. From (I, 1.1.2) we deduce:

$$(2.3.2.1) \quad V_+(0) = \text{Proj}(S), \quad V_+(S) = V_+(S_+) = \emptyset,$$

$$(2.3.2.2) \quad V_+(\bigcup_\lambda E_\lambda) = \bigcap_\lambda V_+(E_\lambda),$$

$$(2.3.2.3) \quad V_+(EE') = V_+(E) \cup V_+(E').$$

We do not change $V_+(E)$ by replacing E with the graded ideal generated by E ; in addition, if $\mathfrak{J}$ is a graded ideal of S , then we have

$$(2.3.2.4) \quad V_+(\mathfrak{J}) = V_+(\bigcup_{q \geq n} (\mathfrak{J} \cap S_q))$$

[II:26] for all $n > 0$: indeed, if $\mathfrak{p} \in \text{Proj}(S)$ contains the homogeneous elements of $\mathfrak{J}$ of degree $\geq n$, then as by hypothesis there exists a homogeneous element $f \in S_d$ not contained in $\mathfrak{p}$, for every $m \geq 0$ and every $x \in S_m \cap \mathfrak{J}$, we have $f^r x \in \mathfrak{J} \cap S_{m+rd}$ for all but finitely many values of r , so $f^r x \in \mathfrak{p} \cap S_{m+rd}$, which implies that $x \in \mathfrak{p} \cap S_m$ (2.1.9).

Finally, we have, for every graded ideal $\mathfrak{J}$ of S ,

$$(2.3.2.5) \quad V_+(\mathfrak{J}) = V_+(\mathfrak{r}_+(\mathfrak{J})).$$

(2.3.3). By definition, the $V_+(E)$ are the closed subsets of $X = \text{Proj}(S)$ for the topology induced by the spectral topology of $\text{Spec}(S)$, which we also call the *spectral topology* on X . For all $f \in S$, we set

$$(2.3.3.1) \quad D_+(f) = D(f) \cap \text{Proj}(S) = \text{Proj}(S) - V_+(f)$$

and so, for any two elements f and g of S (I, 1.1.9.1),

$$(2.3.3.2) \quad D_+(fg) = D_+(f) \cap D_+(g).$$

Proposition (2.3.4). — *The $D_+(f)$, as f runs over the set of homogeneous elements of S_+ , form a base for the topology of $X = \text{Proj}(S)$.*

Proof. It follows from (2.3.2.2) and (2.3.2.4) that every closed subset of X is the intersection of sets of the form $V_+(f)$, where f is homogeneous of degree > 0 . $\square$

(2.3.5). Let f be a homogeneous element of S_+ , of degree $d > 0$; for every graded prime ideal $\mathfrak{p}$ of S that does not contain f , we know that the set of the x/f^n , where $x \in \mathfrak{p}$ and $n \geq 0$, is a prime ideal of the ring of fractions S_f (0, 1.2.6); its intersection with $S_{(f)}$ is thus a prime ideal of $S_{(f)}$, which we denote by $\psi_f(\mathfrak{p})$: it is the set of the x/f^n for $n \geq 0$ and $x \in \mathfrak{p} \cap S_{nd}$. We have thus defined a map

$$\psi_f : D_+(f) \longrightarrow \text{Spec}(S_{(f)});$$

furthermore, if $g \in S_e$ is another homogeneous element of S_+ , then we have a commutative diagram

$$(2.3.5.1) \quad \begin{array}{ccc} D_+(f) & \xrightarrow{\psi_f} & \text{Spec}(S_{(f)}) \\ \uparrow & & \uparrow \\ D_+(fg) & \xrightarrow{\psi_{fg}} & \text{Spec}(S_{(fg)}) \end{array}$$

where the vertical arrow on the left is the inclusion, and the vertical arrow on the right is the map ${}^a\omega_{fg,f}$ induced by the canonical homomorphism $\omega = \omega_{fg,f} : S_{(f)} \rightarrow S_{(fg)}$ (I, 1.2.1). Indeed, if $x/f^n \in \omega^{-1}(\psi_{fg}(\mathfrak{p}))$, with $fg \notin \mathfrak{p}$, then, by definition, $g^n x / (fg)^n \in \psi_{fg}(\mathfrak{p})$, so $g^n x \in \mathfrak{p}$, and so $x \in \mathfrak{p}$; the converse is evident.

Proposition (2.3.6). — *The map ψ_f is a homeomorphism from $D_+(f)$ to $\text{Spec}(S_{(f)})$.*

Proof. First, ψ_f is continuous; indeed, if $h \in S_{nd}$, then $h/f^n \in \psi_f(\mathfrak{p})$ if and only if $h \in \mathfrak{p}$, and so $\psi_f^{-1}(D(h/f^n)) = D_+(hf)$; our claim then follows from (2.3.3.2). Furthermore, the $D_+(hf)$, where h runs over the sets S_{nd} , form a basis for the topology of $D_+(f)$, by (2.3.4) and (2.3.3.2); the [II:27] above thus proves, taking into account the (T_0) axiom, which holds in $D_+(f)$ and in $\text{Spec}(S_{(f)})$, that ψ_f is injective and that the inverse map $\psi_f(D_+(f)) \rightarrow D_+(f)$ is continuous. Finally, to see that ψ_f is surjective, we note that, if $\mathfrak{q}_0$ is a prime ideal of $S_{(f)}$, and if, for all $n > 0$, we denote by $\mathfrak{p}_n$ the set of $x \in S_n$ such that $x^d/f^n \in \mathfrak{q}_0$, then the $\mathfrak{p}_n$ satisfy the conditions of (2.1.9): if $x, y \in S_n$ are such that $x^d/f^n, y^d/f^n \in \mathfrak{q}_0$, then $(x+y)^{2d}/f^{2n} \in \mathfrak{q}_0$, whence $(x+y)^d/f^n \in \mathfrak{q}_0$, since $\mathfrak{q}_0$ is prime; this proves that the $\mathfrak{p}_n$ are subgroups of the S_n , and the verification of the other conditions of (2.1.9) is immediate, taking into account the fact that $\mathfrak{q}_0$ is prime. If $\mathfrak{p}$ is the graded prime ideal of S thus defined, then indeed $\psi_f(\mathfrak{p}) = \mathfrak{q}_0$, since, for $x \in S_{nd}$, the conditions $x/f^n \in \mathfrak{q}_0$ and $x^d/f^{nd} \in \mathfrak{q}_0$ are equivalent because $\mathfrak{q}_0$ is prime. $\square$

Corollary (2.3.7). — *To have $D_+(f) = \emptyset$, it is necessary and sufficient for f to be nilpotent.*

Proof. To have $\text{Spec}(S_{(f)}) = \emptyset$, it is necessary and sufficient to have $S_{(f)} = 0$, or equivalently to have $1 = 0$ in S_f , which means, by definition, that f is nilpotent. $\square$

Corollary (2.3.8). — *Let E be a subset of S_+ . Then the following conditions are equivalent:*

- (a) $V_+(E) = X = \text{Proj}(S)$.
- (b) Every element of E is nilpotent.
- (c) The homogeneous components of every element of E are nilpotent.

Proof. It is clear that (c) implies (b), and that (b) implies (a). If $\mathfrak{J}$ is the graded ideal of S generated by E , then condition (a) is equivalent to requiring that $V_+(\mathfrak{J}) = X$; a fortiori, (a) implies that every homogeneous element $f \in \mathfrak{J}$ is such that $V_+(f) = X$, and so f is nilpotent by (2.3.7). $\square$

Corollary (2.3.9). — *If $\mathfrak{J}$ is a graded ideal of S_+ , then $\mathfrak{r}_+(\mathfrak{J})$ is the intersection of the graded prime ideals of S_+ that contain $\mathfrak{J}$.*

Proof. By considering the graded ring $S/\mathfrak{J}$, we can reduce to the case where $\mathfrak{J} = 0$. We need to prove that, if $f \in S_+$ is not nilpotent, then there exists a graded prime ideal of S that does not contain f ; but at least one of the homogeneous components of f is not nilpotent, and we can thus suppose f to be homogeneous; the claim then follows from (2.3.7). $\square$

(2.3.10). For every subset Y of $X = \text{Proj}(S)$, let $j_+(Y)$ be the set of $f \in S_+$ such that $Y \subset V_+(f)$; this is equivalent to saying that $j_+(Y) = j(Y) \cap S_+$; then $j_+(Y)$ is an ideal of S_+ that is equal to its radical in S_+ .

Proposition (2.3.11). — (i) For every subset E of S_+ , $j_+(V_+(E))$ is the radical in S_+ of the graded ideal of S_+ generated by E .
(ii) For every subset Y of X , $V_+(j_+(Y)) = \overline{Y}$, where $\overline{Y}$ is the closure of Y in X .

Proof.

- (i) If $\mathfrak{J}$ is the graded ideal of S_+ generated by E , then $V_+(E) = V_+(\mathfrak{J})$, and the claim then follows from (2.3.9).
- (ii) Since $V_+(\mathfrak{J}) = \bigcap_{f \in \mathfrak{J}} V_+(f)$, having $Y \subset V_+(\mathfrak{J})$ implies that $Y \subset V_+(f)$ for every $f \in \mathfrak{J}$, and thus $j_+(Y) \supset \mathfrak{J}$, whence $V_+(j_+(Y)) \subset V_+(\mathfrak{J})$, which proves (ii) by the definition of the closed subsets. $\square$

Corollary (2.3.12). — *The closed subsets Y of $X = \text{Proj}(S)$ are in bijective correspondence with the graded ideals of S_+ that are equal to their radical in S_+ , via the inclusion-reversing maps $Y \mapsto j_+(Y)$ and $\mathfrak{J} \mapsto V_+(\mathfrak{J})$; the union $Y_1 \cup Y_2$ of two closed subsets of X corresponds [II:28] to $j_+(Y_1) \cap j_+(Y_2)$, and the intersection of an arbitrary family (Y_λ) of closed subsets corresponds to the radical in S_+ of the sum of the $j_+(Y_\lambda)$.*

Corollary (2.3.13). — Let $\mathfrak{J}$ be a graded ideal of S_+ ; to have $V_+(\mathfrak{J}) = \emptyset$, it is necessary and sufficient for every element f of S_+ to have a power f^n in $\mathfrak{J}$.

The preceding corollary can also be expressed in either of the following equivalent forms:

Corollary (2.3.14). — Let (f_α) be a family of homogeneous elements of S_+ . For the $D_+(f_\alpha)$ to form a cover of $X = \text{Proj}(S)$, it is necessary and sufficient for every element of S_+ to have a power in the ideal generated by the f_α .

Corollary (2.3.15). — Let (f_α) be a family of homogeneous elements of S_+ , and f an element of S_+ . Then the following are equivalent:

- (a) $D_+(f) \subset \bigcup_\alpha D_+(f_\alpha)$;
- (b) $V_+(f) \supset \bigcap_\alpha V_+(f_\alpha)$;
- (c) f has a power in the ideal generated by the f_α .

Corollary (2.3.16). — For $X = \text{Proj}(S)$ to be empty, it is necessary and sufficient for every element of S_+ to be nilpotent.

Corollary (2.3.17). — In the bijective correspondence described in (2.3.12), the irreducible closed subsets of X correspond to the graded prime ideals of S_+ .

Proof. If $Y = Y_1 \cup Y_2$, where Y_1 and Y_2 are proper closed subsets of Y , then

$$j_+(Y) = j_+(Y_1) \cap j_+(Y_2)$$

with the ideals $j_+(Y_1)$ and $j_+(Y_2)$ being distinct from $j_+(Y)$, and so $j_+(Y)$ is not prime. Conversely, if $\mathfrak{J}$ is a graded ideal of S_+ that is not prime, then there exist elements $f, g \in S_+$ such that $f \notin \mathfrak{J}$ and $g \notin \mathfrak{J}$, but $fg \in \mathfrak{J}$; then $V_+(f) \not\supset V_+(\mathfrak{J})$ and $V_+(g) \not\supset V_+(\mathfrak{J})$, but $V_+(\mathfrak{J}) \subset V_+(f) \cup V_+(g)$, by (2.3.2.3); we thus conclude that $V_+(\mathfrak{J})$ is the union of the closed subsets $V_+(f) \cap V_+(\mathfrak{J})$ and $V_+(g) \cap V_+(\mathfrak{J})$, which are distinct from $V_+(\mathfrak{J})$. $\square$

2.4. The scheme structure on $\text{Proj}(S)$

(2.4.1). Let f and g be homogeneous elements of S_+ ; consider the affine schemes $Y_f = \text{Spec}(S_{(f)})$, $Y_g = \text{Spec}(S_{(g)})$, and $Y_{fg} = \text{Spec}(S_{(fg)})$. By (2.2.2), the morphism $w_{fg,f} = ({}^a\omega_{fg,f}, \tilde{\omega}_{fg,f})$ from Y_{fg} to Y_f , corresponding to the canonical homomorphism $\omega_{fg,f} : S_{(f)} \rightarrow S_{(fg)}$, is an open immersion (I, 1.3.6). Using the inverse homeomorphism of $\psi_f : D_+(f) \rightarrow Y_f$ (2.3.6), we can transport the affine scheme structure of Y_f to $D_+(f)$; by the commutativity of diagram (2.3.5.1), the affine scheme $D_+(fg)$ can thus be identified with the induced scheme on the open subset $D_+(fg)$ of the underlying space of the affine scheme $D_+(f)$. It is then clear (taking (2.3.4) into account) that $X = \text{Proj}(S)$ is endowed with a unique prescheme structure, whose restriction to each $D_+(f)$ is the affine scheme that we have just defined. Furthermore:

Proposition (2.4.2). — The prescheme $\text{Proj}(S)$ is a scheme.

Proof. It suffices (I, 5.5.6) to show, for any homogeneous f and g in S_+ , that $D_+(f) \cap D_+(g) = D_+(fg)$ is affine, and that its ring is generated by the canonical images of the rings of $D_+(f)$ and $D_+(g)$; the first point is evident by definition, and the second follows from (2.2.4). $\square$

[II:29] Whenever we speak of the homogeneous prime spectrum $\text{Proj}(S)$ as a scheme, we always mean it with the structure just defined.

Example (2.4.3). — Let $S = K[T_1, T_2]$, where K is a field, T_1 and T_2 are indeterminates, and S is graded by total degree. It follows from (2.3.14) that $\text{Proj}(S)$ is the union of $D_+(T_1)$ and $D_+(T_2)$; we immediately see that these affine schemes are canonically isomorphic to $\text{Spec}(K[T])$, and that $\text{Proj}(S)$ is obtained by gluing these two affine schemes as described in (I, 2.3.2) (cf. (7.4.14)).

Proposition (2.4.4). — Let S be a positively graded ring, and let X be the scheme $\text{Proj}(S)$.

- (i) If $\mathfrak{N}_+$ is the nilradical of S_+ (2.1.10), then the scheme X_{red} is canonically isomorphic to $\text{Proj}(S/\mathfrak{N}_+)$; in particular, if S is essentially reduced, then $\text{Proj}(S)$ is reduced.
- (ii) If S is essentially reduced, then, for X to be integral, it is necessary and sufficient for S to be essentially integral.

Proof.

- (i) Let $\bar{S}$ be the graded ring $S/\mathfrak{N}_+$, and denote by $x \mapsto \bar{x}$ the canonical homomorphism $S \rightarrow \bar{S}$ of degree 0. For every $f \in S_d$ ($d > 0$), the canonical homomorphism $S_f \rightarrow \bar{S}_{\bar{f}}$ (0, 1.5.1) is surjective and of degree 0, and thus gives, by restriction, a surjective homomorphism $S_{(f)} \rightarrow \bar{S}_{(\bar{f})}$; if we suppose that $f \notin \mathfrak{N}_+$, then we immediately see that $\bar{S}_{(\bar{f})}$ is reduced, and that the kernel of the above homomorphism is the nilradical of $S_{(f)}$, or, in other words, that $\bar{S}_{(\bar{f})} = (S_{(f)})_{\text{red}}$. So to this homomorphism corresponds a closed immersion $D_+(\bar{f}) \rightarrow D_+(f)$ that identifies $D_+(\bar{f})$ with $(D_+(f))_{\text{red}}$ (I, 5.1.2), and which is, in particular, a homeomorphism of the underlying spaces of these two affine schemes. Furthermore, if $g \notin \mathfrak{N}_+$ is another homogeneous element of S_+ , then the diagram

$$\begin{array}{ccc} S_{(f)} & \longrightarrow & \bar{S}_{(\bar{f})} \\ \downarrow & & \downarrow \\ S_{(fg)} & \longrightarrow & \bar{S}_{(\bar{f}\bar{g})} \end{array}$$

is commutative; since, furthermore, the $D_+(f)$, for homogeneous f of degree > 0 with $f \notin \mathfrak{N}_+$, form a cover of $X = \text{Proj}(S)$ (2.3.7), we see that the morphisms $D_+(\bar{f}) \rightarrow D_+(f)$ are the restrictions of a closed immersion $\text{Proj}(\bar{S}) \rightarrow \text{Proj}(S)$ which is a homeomorphism of the underlying spaces; whence the conclusion (I, 5.1.2).

- (ii) Suppose that S is essentially integral, or, in other words, that (0) is a graded prime ideal of S_+ that is distinct from S_+ ; then X is reduced, by (i), and irreducible, by (2.3.17). Conversely, suppose that S is essentially reduced and that X is integral; then, for homogeneous $f \neq 0$ in S_+ , we have that $D_+(f) \neq \emptyset$ (2.3.7); the hypothesis that X is irreducible implies that $D_+(f) \cap D_+(g) \neq \emptyset$ for homogeneous $f, g \neq 0$ in S_+ ; thus $fg \neq 0$, by (2.3.3.2), and we thus conclude that S_+ has no zero divisors, whence the first claim. $\square$

(2.4.5). Given a commutative ring A , recall that we say that a graded ring S is a *graded A -algebra* if it is endowed with the structure of an A -algebra such that each of its subgroups S_n is an A -module; for this, it suffices for S_0 to be ^[II:30] an A -algebra, or, in other words, we define the structure of a graded A -algebra on S by defining the structure of an A -algebra on S_0 and setting $\alpha \cdot x = (\alpha \cdot 1)x$ for $\alpha \in A$ and $x \in S_n$.

Proposition (2.4.6). — Suppose that S is a graded A -algebra. Then, on $X = \text{Proj}(S)$, the structure sheaf $\mathcal{O}_X$ is a sheaf of A -algebras (where A is regarded as the constant sheaf on X); in other words, X is a scheme over $\text{Spec}(A)$.

Proof. It suffices to note that, for every homogeneous f in S_+ , $S_{(f)}$ is an algebra over A , and that the diagram

$$\begin{array}{ccc} S_{(f)} & \xrightarrow{\quad} & S_{(fg)} \\ & \nwarrow \quad \nearrow & \\ & A & \end{array}$$

is commutative, for homogeneous f, g in S_+ . $\square$

Proposition (2.4.7). — Let S be a positively graded ring.

- (i) For every integer $d > 0$, there exists a canonical isomorphism from the scheme $\text{Proj}(S)$ to the scheme $\text{Proj}(S^{(d)})$.
- (ii) Let S' be the graded ring such that $S'_0 = \mathbf{Z}$ and $S'_n = S_n$ (considered as a $\mathbf{Z}$ -module) for $n > 0$. Then there exists a canonical isomorphism from the scheme $\text{Proj}(S)$ to the scheme $\text{Proj}(S')$.

Proof.

- (i) We first show that the map $\mathfrak{p} \mapsto \mathfrak{p} \cap S^{(d)}$ is a bijection from the set $\text{Proj}(S)$ to the set $\text{Proj}(S^{(d)})$. Indeed, suppose that we have a graded prime ideal $\mathfrak{p}' \in \text{Proj}(S^{(d)})$, and let $\mathfrak{p}_{nd} = \mathfrak{p}' \cap S_{nd}$ ($n \geq 0$). For all $n > 0$ that are not multiples of d , define $\mathfrak{p}_n$ as the set of $x \in S_n$ such that $x^d \in \mathfrak{p}_{nd}$; if $x, y \in \mathfrak{p}_n$, then $(x + y)^{2d} \in \mathfrak{p}_{2nd}$, and so $(x + y)^d \in \mathfrak{p}_{nd}$, since $\mathfrak{p}'$ is prime; it is immediate that the $\mathfrak{p}_n$ thus defined, for $n \geq 0$, satisfy the conditions of (2.1.9), and so there exists a unique prime ideal $\mathfrak{p} \in \text{Proj}(S)$ such that $\mathfrak{p} \cap S^{(d)} = \mathfrak{p}'$. Since, for every homogeneous f in S_+ , we have that $V_+(f) = V_+(f^d)$ (2.3.2.3), we see that the above bijection is a *homeomorphism* of topological spaces. Finally, with the same notation, $S_{(f)}$ and $S_{(f^d)}$ are canonically identified (2.2.2), and so $\text{Proj}(S)$ and $\text{Proj}(S^{(d)})$ are canonically identified as *schemes*.
- (ii) If, to each $\mathfrak{p} \in \text{Proj}(S)$, we associate the unique prime ideal $\mathfrak{p}' \in \text{Proj}(S')$ such that $\mathfrak{p}' \cap S_n = \mathfrak{p} \cap S_n$ for every $n > 0$ (2.1.9), then it is clear that we have defined a canonical homeomorphism $\text{Proj}(S) \xrightarrow{\sim} \text{Proj}(S')$ of the underlying spaces, since $V_+(f)$ is the same set for S and S' when f is a homogeneous element of S_+ . Since, further, $S_{(f)} = S'_{(f)}$, $\text{Proj}(S)$ and $\text{Proj}(S')$ can be identified as *schemes*.

□

Corollary (2.4.8). — *If S is a graded A -algebra, and if S'_A is the graded A -algebra such that $(S'_A)_0 = A$ and $(S'_A)_n = S_n$ for $n > 0$, then there exists a canonical isomorphism from $\text{Proj}(S)$ to $\text{Proj}(S'_A)$.*

Proof. In fact, these two schemes are both canonically isomorphic to $\text{Proj}(S')$, using the notation of (2.4.7, (ii)). □

2.5. The sheaf associated to a graded module

(2.5.1). Let M be a graded S -module. Then, for every homogeneous f in S_+ , $M_{(f)}$ is an $S_{(f)}$ -module, and thus has a corresponding quasi-coherent associated sheaf $(M_{(f)})^\sim$ on the affine scheme $\text{Spec}(S_{(f)})$, identified with $D_+(f)$ (I, 1.3.4).

[II:31]

Proposition (2.5.2). — *There exists on $X = \text{Proj}(S)$ exactly one quasi-coherent $\mathcal{O}_X$ -module $\tilde{M}$ such that, for f homogeneous in S_+ , we have $\Gamma(D_+(f), \tilde{M}) = M_{(f)}$, with the restriction homomorphism $\Gamma(D_+(f), \tilde{M}) \rightarrow \Gamma(D_+(fg), \tilde{M})$, for f and g homogeneous in S_+ , being the canonical homomorphism $M_{(f)} \rightarrow M_{(fg)}$ (2.2.3).*

Proof. Suppose that $f \in S_d$ and $g \in S_e$. Since $D_+(fg)$ can be identified with the prime spectrum of $(S_{(f)})_{g^d/f^e}$ by (2.2.2), the restriction to $D_+(fg)$ of the sheaf $(M_{(f)})^\sim$ on $D_+(f)$ is canonically identified with the sheaf associated to the module $(M_{(f)})_{g^d/f^e}$ (I, 1.3.6), and thus also with $(M_{(fg)})^\sim$ (2.2.2); we thus conclude that there exists a canonical isomorphism

$$\theta_{g,f} : (M_{(f)})^\sim|_{D_+(fg)} \xrightarrow{\sim} (M_{(g)})^\sim|_{D_+(fg)}$$

such that, if h is a third homogeneous element of S_+ , then $\theta_{f,h} = \theta_{f,g} \circ \theta_{g,h}$ in $D_+(fgh)$. Consequently (0, 3.3.1) there exists a quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$ on X , and, for every homogeneous f in S_+ , an isomorphism η_f from $\mathcal{F}|_{D_+(f)}$ to $(M_{(f)})^\sim$ such that $\theta_{g,f} = \eta_g \circ \eta_f^{-1}$. If we then consider the sheaf $\mathcal{G}$ associated to the presheaf (on the base for the topology of X given by the $D_+(f)$) defined by $D_+(f) \mapsto M_{(f)}$, with the canonical homomorphisms $M_{(f)} \rightarrow M_{(fg)}$ as restriction homomorphisms, then the above proves that $\mathcal{F}$ and $\mathcal{G}$ are isomorphic (taking (I, 1.3.7) into account); the sheaf $\mathcal{G}$ is denoted by $\tilde{M}$, and indeed satisfies the conditions of the statement. We have, in particular, $\tilde{S} = \mathcal{O}_X$. □

Definition (2.5.3). — We say that the quasi-coherent $\mathcal{O}_X$ -module $\tilde{M}$ defined in (2.5.2) is *associated* to the graded S -module M .

Recall that the graded S -modules form a category when we restrict from arbitrary homomorphisms of graded modules to homomorphisms of degree 0. With this convention:

Proposition (2.5.4). — *The functor $M \mapsto \tilde{M}$ is an exact additive covariant functor from the category of graded S -modules to the category of quasi-coherent $\mathcal{O}_X$ -modules, and it commutes with inductive limits and direct sums.*

Proof. Indeed, since these properties are local, it suffices to show that they are satisfied for the sheaves of the form $\tilde{M}|D_+(f) = (M_{(f)})^\sim$; but the localization functor $M \mapsto M_{(f)}$, the degree-zero functor $N \mapsto N_0$ on graded S_f -modules, and the associated-sheaf functor $P \mapsto \tilde{P}$ on $S_{(f)}$ -modules all have these three properties: exactness and commutation with inductive limits and direct sums ((I, 1.3.5) and (I, 1.3.9)); whence the proposition. $\square$

We denote by $\tilde{u}$ the homomorphism $\tilde{M} \rightarrow \tilde{N}$ corresponding to a homomorphism $u : M \rightarrow N$ of degree 0. We immediately deduce from (2.5.4) that the results of (I, 1.3.9) and (I, 1.3.10) still hold for graded S -modules and homomorphisms of degree 0 (with the meaning given here to $\tilde{M}$), since the proofs are purely formal.

Proposition (2.5.5). — *For all $\mathfrak{p} \in X = \text{Proj}(S)$, we have $\tilde{M}_{\mathfrak{p}} = M_{(\mathfrak{p})}$.*

Proof. By definition, $\tilde{M}_{\mathfrak{p}} = \varinjlim \Gamma(D_+(f), \tilde{M})$, where f runs over the set of homogeneous elements of S_+ such that $f \notin \mathfrak{p}$; since $\Gamma(D_+(f), \tilde{M}) = M_{(f)}$, the proposition follows from the definition of $M_{(\mathfrak{p})}$ (2.2.7). $\square$

[II:32] In particular, the local ring $(\mathcal{O}_X)_{\mathfrak{p}}$ is exactly the ring $S_{(\mathfrak{p})}$, the set of fractions x/f with f homogeneous in S_+ and not belonging to $\mathfrak{p}$, and with x homogeneous of the same degree as f .

More particularly, if S is *essentially integral*, so that $\text{Proj}(S) = X$ is *integral* (2.4.4), and if $\xi = (0)$ is the generic point of X , then the field of rational functions $R(X) = \mathcal{O}_{\xi}$ is the field consisting of the elements fg^{-1} , where f and g are homogeneous of the same degree in S_+ and $g \neq 0$.

Proposition (2.5.6). — *If, for all $z \in M$ and all homogeneous f in S_+ , there exists a power of f that annihilates z , then $\tilde{M} = 0$. This sufficient condition is also necessary if the S_0 -algebra S is generated by the set S_1 of homogeneous elements of degree 1.*

Proof. The condition $\tilde{M} = 0$ is equivalent to $M_{(f)} = 0$ for all homogeneous f in S_+ . On the other hand, if $f \in S_d$, the equality $M_{(f)} = 0$ means that, for every homogeneous $z \in M$ whose degree is a multiple of d , there exists a power f^n such that $f^n z = 0$. If $M_{(f)} = 0$ for all $f \in S_1$, then the condition of the statement is thus satisfied for all $f \in S_1$; the condition is *a fortiori* satisfied for all homogeneous f in S_+ if S_1 generates S , since every homogeneous element of S_+ is then a linear combination of products of elements of S_1 . $\square$

Proposition (2.5.7). — *Let $d > 0$ be an integer, and let $f \in S_d$. Then, for all $n \in \mathbf{Z}$, the $(\mathcal{O}_X|D_+(f))$ -module $(S(nd))^\sim|D_+(f)$ is canonically isomorphic to $\mathcal{O}_X|D_+(f)$.*

Proof. Indeed, multiplication by the invertible element $(f/1)^n$ of S_f gives a bijection from $S_{(f)} = (S_f)_0$ to $(S_f)_{nd} = (S_f(nd))_0 = (S(nd)_f)_0 = S(nd)_{(f)}$; in other words, the $S_{(f)}$ -modules $S_{(f)}$ and $S(nd)_{(f)}$ are canonically isomorphic, whence the proposition. $\square$

Corollary (2.5.8). — *On the open subset $U = \bigcup_{f \in S_d} D_+(f)$, the restriction of the $\mathcal{O}_X$ -module $(S(nd))^\sim$ is an invertible $(\mathcal{O}_X|U)$ -module (0, 5.4.1).*

Corollary (2.5.9). — *If the ideal S_+ of S is generated by the set S_1 of homogeneous elements of degree 1, then the $\mathcal{O}_X$ -module $(S(n))^\sim$ is invertible for all $n \in \mathbf{Z}$.*

Proof. It suffices to remark that $X = \bigcup_{f \in S_1} D_+(f)$, by the hypothesis (2.3.14) and to apply (2.5.8) with $U = X$. $\square$

(2.5.10). We set, for the rest of this section,

$$(2.5.10.1) \quad \mathcal{O}_X(n) = (S(n))^\sim$$

for all $n \in \mathbf{Z}$, and, for every open subset U of X , and every $(\mathcal{O}_X|U)$ -module $\mathcal{F}$,

$$(2.5.10.2) \quad \mathcal{F}(n) = \mathcal{F} \otimes_{\mathcal{O}_X|U} (\mathcal{O}_X(n)|U)$$

for all $n \in \mathbf{Z}$. If the ideal S_+ is generated by S_1 , then the functor $\mathcal{F} \mapsto \mathcal{F}(n)$ is *exact* for every $n \in \mathbf{Z}$, since $\mathcal{O}_X(n)$ is then an invertible $\mathcal{O}_X$ -module.

(2.5.11). Let M and N be graded S -modules. For all $f \in S_d$ ($d > 0$), we define a canonical functorial homomorphism of $S_{(f)}$ -modules by

$$(2.5.11.1) \quad \lambda_f : M_{(f)} \otimes_{S_{(f)}} N_{(f)} \longrightarrow (M \otimes_S N)_{(f)}$$

[II:33] by composing the homomorphism $M_{(f)} \otimes_{S_{(f)}} N_{(f)} \rightarrow M_f \otimes_{S_f} N_f$ (coming from the injections $M_{(f)} \rightarrow M_f, N_{(f)} \rightarrow N_f$, and $S_{(f)} \rightarrow S_f$) with the canonical isomorphism $M_f \otimes_{S_f} N_f \xrightarrow{\sim} (M \otimes_S N)_f$ (0, 1.3.4), and by noting that, by the definition of the tensor product of two graded modules, this latter isomorphism preserves degrees; for $x \in M_{md}$ and $y \in N_{nd}$ ($m, n \geq 0$), we thus have

$$\lambda_f((x/f^m) \otimes (y/f^n)) = (x \otimes y)/f^{m+n}.$$

It immediately follows from this definition that, if $g \in S_e$ ($e > 0$), then the diagram

$$\begin{array}{ccc} M_{(f)} \otimes_{S_{(f)}} N_{(f)} & \xrightarrow{\lambda_f} & (M \otimes_S N)_{(f)} \\ \downarrow & & \downarrow \\ M_{(fg)} \otimes_{S_{(fg)}} N_{(fg)} & \xrightarrow{\lambda_{fg}} & (M \otimes_S N)_{(fg)} \end{array}$$

(where the vertical arrow on the right is the canonical homomorphism, and the one on the left comes from the canonical homomorphisms) commutes. Thus λ induces a canonical functorial homomorphism of $\mathcal{O}_X$ -modules

$$(2.5.11.2) \quad \lambda : \tilde{M} \otimes_{\mathcal{O}_X} \tilde{N} \longrightarrow (M \otimes_S N)^\sim.$$

Consider, in particular, graded ideals $\mathfrak{J}$ and $\mathfrak{K}$ of S ; since $\tilde{\mathfrak{J}}$ and $\tilde{\mathfrak{K}}$ are sheaves of ideals of $\mathcal{O}_X$, we have a canonical homomorphism $\tilde{\mathfrak{J}} \otimes_{\mathcal{O}_X} \tilde{\mathfrak{K}} \rightarrow \mathcal{O}_X$; the diagram

$$(2.5.11.3) \quad \begin{array}{ccc} \tilde{\mathfrak{J}} \otimes_{\mathcal{O}_X} \tilde{\mathfrak{K}} & \xrightarrow{\lambda} & (\mathfrak{J} \otimes_S \mathfrak{K})^\sim \\ & \searrow & \swarrow \\ & \mathcal{O}_X & \end{array}$$

then commutes. Indeed, we can reduce to verifying this on each open subset $D_+(f)$ (for f homogeneous in S_+), and this follows immediately from the definition (2.5.11.1) of λ_f and from (I, 1.3.13).

Finally, note that, if M, N , and P are graded S -modules, then the diagram

$$(2.5.11.4) \quad \begin{array}{ccc} \tilde{M} \otimes_{\mathcal{O}_X} \tilde{N} \otimes_{\mathcal{O}_X} \tilde{P} & \xrightarrow{\lambda \otimes 1} & (M \otimes_S N)^\sim \otimes_{\mathcal{O}_X} \tilde{P} \\ 1 \otimes \lambda \downarrow & & \downarrow \lambda \\ \tilde{M} \otimes_{\mathcal{O}_X} (N \otimes_S P)^\sim & \xrightarrow{\lambda} & (M \otimes_S N \otimes_S P)^\sim \end{array}$$

[II:34] commutes. It again suffices to verify this on each open subset $D_+(f)$, and this follows immediately from the definitions and from (I, 1.3.13).

(2.5.12). Under the hypotheses of (2.5.11), we define a functorial canonical homomorphism of $S_{(f)}$ -modules

$$(2.5.12.1) \quad \mu_f : (\mathrm{Hom}_S(M, N))_{(f)} \longrightarrow \mathrm{Hom}_{S_{(f)}}(M_{(f)}, N_{(f)})$$

by sending u/f^n , where u is a homomorphism of degree nd , to the homomorphism $M_{(f)} \rightarrow N_{(f)}$ that sends x/f^m ($x \in M_{md}$) to $u(x)/f^{m+n}$. For $g \in S_e$ ($e > 0$), we again have a commutative diagram:

$$\begin{array}{ccc} (\mathrm{Hom}_S(M, N))_{(f)} & \xrightarrow{\mu_f} & \mathrm{Hom}_{S_{(f)}}(M_{(f)}, N_{(f)}) \\ \downarrow & & \downarrow \\ (\mathrm{Hom}_S(M, N))_{(fg)} & \xrightarrow{\mu_{fg}} & \mathrm{Hom}_{S_{(fg)}}(M_{(fg)}, N_{(fg)}) \end{array}$$

(where the vertical arrow on the left is the canonical homomorphism, and the one on the right comes from the canonical homomorphisms). We thus again conclude (taking (I, 1.3.8) into account) that the μ_f define a functorial canonical homomorphism of $\mathcal{O}_X$ -modules

$$(2.5.12.2) \quad \mu : (\mathrm{Hom}_S(M, N))^\sim \longrightarrow \mathcal{H}om_{\mathcal{O}_X}(\tilde{M}, \tilde{N})$$

Proposition (2.5.13). — Suppose that the ideal S_+ is generated by S_1 . Then the homomorphism λ (2.5.11.2) is an isomorphism; so too is the homomorphism μ (2.5.12.2) if the graded S -module M admits a finite presentation (2.1.1).

Proof. Since X is the union of the $D_+(f)$ for $f \in S_1$ (2.3.14), we are led to proving that λ_f and μ_f are isomorphisms, under the given hypotheses, whenever f is homogeneous and of degree 1. But we can then define a $\mathbf{Z}$ -bilinear map $M_m \times N_n \rightarrow M_{(f)} \otimes_{S_{(f)}} N_{(f)}$ by sending (x, y) to the element $(x/f^m) \otimes (y/f^n)$ (if $m < 0$, we write x/f^m to mean $f^{-m}x/1$); these maps define a $\mathbf{Z}$ -linear map $M \otimes_{\mathbf{Z}} N \rightarrow M_{(f)} \otimes_{S_{(f)}} N_{(f)}$, and, if $s \in S_q$, this map sends $(sx) \otimes y$ to $(s/f^q)((x/f^m) \otimes (y/f^n))$ (for $x \in M_m$ and $y \in N_n$). We thus obtain a di-homomorphism of modules $\gamma_f : M \otimes_S N \rightarrow M_{(f)} \otimes_{S_{(f)}} N_{(f)}$, with respect to the canonical homomorphism $S \rightarrow S_{(f)}$ (sending $s \in S_q$ to s/f^q). Suppose furthermore that, for an element $\sum_i (x_i \otimes y_i)$ of $M \otimes_S N$ (with x_i and y_i homogeneous of degree m_i and n_i , respectively), we have that $f^r \sum_i (x_i \otimes y_i) = 0$, or, in other words, that $\sum_i (f^r x_i \otimes y_i) = 0$. We thus deduce, by (0, 1.3.4), that $\sum_i (f^r x_i / f^{m_i+r}) \otimes (y_i / f^{n_i}) = 0$, i.e. $\gamma_f(\sum_i (x_i \otimes y_i)) = 0$. Then γ_f factors as $M \otimes_S N \rightarrow (M \otimes_S N)_f \xrightarrow{\gamma'_f} M_{(f)} \otimes_{S_{(f)}} N_{(f)}$; if λ'_f is the restriction of γ'_f [II:35] to $(M \otimes_S N)_{(f)}$, then we can immediately show that λ_f and λ'_f are inverse $S_{(f)}$ -homomorphisms, whence the first part of the proposition.

To prove the second part, suppose that M is the cokernel of a homomorphism $P \rightarrow Q$ of graded S -modules, with P and Q being direct sums of a finite number of modules of the form $S(n)$; using the left-exactness of the functor $L \mapsto \mathrm{Hom}_S(L, N)$ and the exactness of the functor $M \mapsto M_{(f)}$, we can immediately reduce to proving that μ_f is an isomorphism whenever $M = S(n)$. But, for any homogeneous z in N , let u_z be the homomorphism from $S(n)$ to N such that $u_z(1) = z$; we immediately see that $\eta : z \rightarrow u_z$ is an isomorphism of degree 0 from $N(-n)$ to $\mathrm{Hom}_S(S(n), N)$. There is a corresponding isomorphism

$$\eta_f : (N(-n))_{(f)} \longrightarrow (\mathrm{Hom}_S(S(n), N))_{(f)}.$$

Now let η'_f be the isomorphism $N_{(f)} \rightarrow \mathrm{Hom}_{S_{(f)}}(S(n)_{(f)}, N_{(f)})$ that, to any $z' \in N_{(f)}$, associates the homomorphism $v_{z'}$ that is such that $v_{z'}(s/f^k) = sz'/f^{n+k}$ (for $s \in S_{n+k} = (S(n))_k$). We easily note that the composed map

$$(N(-n))_{(f)} \xrightarrow{\eta_f} (\mathrm{Hom}_S(S(n), N))_{(f)} \xrightarrow{\mu_f} \mathrm{Hom}_{S_{(f)}}(S(n)_{(f)}, N_{(f)}) \xrightarrow{\eta'^{-1}_f} N_{(f)}$$

is the isomorphism $z/f^h \mapsto z/f^{h-n}$ from $(N(-n))_{(f)}$ to $N_{(f)}$, and thus μ_f is an isomorphism. $\square$

If the ideal S_+ is generated by S_1 , then we deduce from (2.5.13) that, for every graded ideal $\mathfrak{J}$ of S , and for every graded S -module M , we have

$$(2.5.13.1) \quad \tilde{\mathfrak{J}} \cdot \tilde{M} = (\mathfrak{J} \cdot M)^\sim$$

up to canonical isomorphism; this follows from the commutativity of the diagram

$$\begin{array}{ccc} \tilde{\mathfrak{J}} \otimes_{\mathcal{O}_X} \tilde{M} & \xrightarrow{\lambda} & (\mathfrak{J} \otimes_S M)^\sim \\ & \searrow & \swarrow \\ & \tilde{M} & \end{array}$$

which we can verify as we did for (2.5.11.3).

Corollary (2.5.14). — Suppose that S is generated by S_1 . For any $m, n \in \mathbf{Z}$, we then have:

$$(2.5.14.1) \quad \mathcal{O}_X(m) \otimes_{\mathcal{O}_X} \mathcal{O}_X(n) = \mathcal{O}_X(m+n)$$

$$(2.5.14.2) \quad \mathcal{O}_X(n) = (\mathcal{O}_X(1))^{\otimes n}$$

up to canonical isomorphism.

Proof. The first equation follows from (2.5.13) and from the existence of the canonical isomorphism $S(m) \otimes_S S(n) \xrightarrow{\sim} S(m+n)$ of degree 0 that sends the element $1 \otimes 1$ (where the first 1 is in $(S(m))_{-m}$ and the second is in $(S(n))_{-n}$) to the element $1 \in (S(m+n))_{-(m+n)}$. It then suffices to prove the second equation for $n = -1$, and, by (2.5.13), this reduces to seeing that $\text{Hom}_S(S(1), S)$ is canonically isomorphic to $S(-1)$, which follows immediately from the definitions (2.1.2) and the fact that $S(1)$ is a cyclic S -module. $\square$

[II:36]

Corollary (2.5.15). — Suppose that S is generated by S_1 . Then, for every graded S -module M , and for every $n \in \mathbf{Z}$, we have

$$(2.5.15.1) \quad (M(n))^\sim = \tilde{M}(n)$$

up to canonical isomorphism.

Proof. This follows from definitions (2.5.10.2) and (2.5.10.1), from Proposition (2.5.13), and from the existence of a canonical isomorphism $M(n) \xrightarrow{\sim} M \otimes_S S(n)$ of degree 0 that, to every $z \in (M(n))_h = M_{n+h}$, associates $z \otimes 1 \in M_{n+h} \otimes (S(n))_{-n} \subset (M \otimes_S S(n))_h$. $\square$

(2.5.16). We denote by S' the graded ring such that $S'_0 = \mathbf{Z}$, and $S'_n = S_n$ for $n > 0$. Then, if $f \in S_d$ ($d > 0$), we have that $(S(n))_{(f)} = (S'(n))_{(f)}$ for all $n \in \mathbf{Z}$, since an element of $(S'(n))_{(f)}$ is of the form x/f^k , with $x \in S'_{n+kd}$ ($k > 0$), and we can always take k to be such that $n + kd \neq 0$. Since $X = \text{Proj}(S)$ and $X' = \text{Proj}(S')$ are canonically identified (2.4.7, (ii)), we see that, for all $n \in \mathbf{Z}$, $\mathcal{O}_X(n)$ and $\mathcal{O}_{X'}(n)$ are the images of one another under the above identification.

Note also that, for all $d > 0$ and all $n \in \mathbf{Z}$, we have

$$(S^{(d)}(n))_h = S_{(n+h)d} = (S(nd))_{hd}$$

for $f \in S_d$, and thus $(S^{(d)}(n))_{(f)} = (S(nd))_{(f)}$. We know that the schemes $X = \text{Proj}(S)$ and $X^{(d)} = \text{Proj}(S^{(d)})$ are canonically identified (2.4.7, (i)); the above shows that, if the S_0 -algebra $S^{(d)}$ is generated by S_d , then $\mathcal{O}_X(nd)$ and $\mathcal{O}_{X^{(d)}}(n)$ are the images of one another under this identification, for all $n \in \mathbf{Z}$.

Proposition (2.5.17). — Let $d > 0$ be an integer, and let $U = \bigcup_{f \in S_d} D_+(f)$. Then the restriction to U of the canonical homomorphism $\mathcal{O}_X(nd) \otimes_{\mathcal{O}_X} \mathcal{O}_X(-nd) \rightarrow \mathcal{O}_X$ is an isomorphism for every integer n .

Proof. By (2.5.16), we can restrict to the case where $d = 1$, and the conclusion then follows from the proof of (2.5.13). $\square$

2.6. The graded S -module associated to a sheaf on $\text{Proj}(S)$ Throughout this section, we assume that the ideal S_+ of S is generated by the set S_1 of homogeneous elements of degree 1.

(2.6.1). The $\mathcal{O}_X$ -module $\mathcal{O}_X(1)$ is invertible (2.5.9); we thus define, for every $\mathcal{O}_X$ -module $\mathcal{F}$ (0, 5.4.6),

$$(2.6.1.1) \quad \Gamma_\bullet(\mathcal{F}) = \Gamma_\bullet(\mathcal{O}_X(1), \mathcal{F}) = \bigoplus_{n \in \mathbf{Z}} \Gamma(X, \mathcal{F}(n))$$

taking (2.5.14.2) into account. Recall (0, 5.4.6) that $\Gamma_\bullet(\mathcal{O}_X)$ is endowed with the structure of a graded ring, and $\Gamma_\bullet(\mathcal{F})$ with the structure of a graded $\Gamma_\bullet(\mathcal{O}_X)$ -module.

Since $\mathcal{O}_X(n)$ is locally free, $\Gamma_\bullet(\mathcal{F})$ is a left exact additive covariant functor in $\mathcal{F}$; in particular, if $\mathcal{I}$ is a sheaf of ideals of $\mathcal{O}_X$, then $\Gamma_\bullet(\mathcal{I})$ is canonically identified with a graded ideal of $\Gamma_\bullet(\mathcal{O}_X)$.

(2.6.2). Let M be a graded S -module; for every $f \in S_d$ ($d > 0$), $x \mapsto x/1$ is a homomorphism of abelian groups $M_0 \rightarrow M_{(f)}$, and, since $M_{(f)}$ is canonically identified [II:37] with $\Gamma(D_+(f), \tilde{M})$, we

thus obtain a homomorphism of abelian groups $\alpha_0^f : M_0 \rightarrow \Gamma(D_+(f), \tilde{M})$. It is clear that, for every $g \in S_e$ ($e > 0$), the diagram

$$\begin{array}{ccc} & & \Gamma(D_+(f), \tilde{M}) \\ & \nearrow \alpha_0^f & \downarrow \\ M_0 & & \\ & \searrow \alpha_0^{fg} & \downarrow \\ & & \Gamma(D_+(fg), \tilde{M}) \end{array}$$

commutes; this implies that, for every $x \in M_0$, the sections $\alpha_0^f(x)$ and $\alpha_0^g(x)$ of $\tilde{M}$ agree on $D_+(f) \cap D_+(g)$, and thus there exists a unique section $\alpha_0(x) \in \Gamma(X, \tilde{M})$ whose restriction to each $D_+(f)$ is $\alpha_0^f(x)$. We have thus defined (without using the hypothesis that S be generated by S_1) a homomorphism of abelian groups

$$(2.6.2.1) \quad \alpha_0 : M_0 \longrightarrow \Gamma(X, \tilde{M}).$$

Applying this result to the graded S -module $M(n)$ (for each $n \in \mathbf{Z}$), we obtain, for each $n \in \mathbf{Z}$, a homomorphism of abelian groups

$$(2.6.2.2) \quad \alpha_n : M_n = (M(n))_0 \longrightarrow \Gamma(X, \tilde{M}(n))$$

(taking (2.5.15) into account); whence we obtain a functorial homomorphism (of degree 0) of graded abelian groups

$$(2.6.2.3) \quad \alpha : M \longrightarrow \Gamma_\bullet(\tilde{M})$$

(also denoted by α_M) which, on each M_n , agrees with α_n .

If we take, in particular, $M = S$, then we immediately see (taking into account the definition (0, 5.4.6) of multiplication in $\Gamma_\bullet(\mathcal{O}_X)$) that $\alpha : S \rightarrow \Gamma_\bullet(\mathcal{O}_X)$ is a homomorphism of graded rings, and that, for every graded S -module M , (2.6.2.3) is a di-homomorphism of graded modules.

Proposition (2.6.3). — *For every $f \in S_d$ ($d > 0$), $D_+(f)$ is identical to the set of $\mathfrak{p} \in X$ on which the section $\alpha_d(f)$ of $\mathcal{O}_X(d)$ does not vanish (0, 5.5.2).*

Proof. Since $X = \bigcup_{g \in S_1} D_+(g)$ by hypothesis, it suffices to show that, for all $g \in S_1$, the set of $\mathfrak{p} \in D_+(g)$ on which $\alpha_d(f)$ does not vanish is identical to $D_+(fg)$. But the restriction of $\alpha_d(f)$ to $D_+(g)$ is, by definition, the section corresponding to the element $f/1$ of $(S(d))_{(g)}$; under the canonical isomorphism $(S(d))_{(g)} \xrightarrow{\sim} S_{(g)}$ (2.5.7), this section of $\mathcal{O}_X(d)$ over $D_+(g)$ is identified with the section of $\mathcal{O}_X$ over $D_+(g)$ that corresponds to the element f/g^d of $S_{(g)}$; to say that this section vanishes at $\mathfrak{p} \in D_+(g)$ means that $f/g^d \in \mathfrak{q}$, where $\mathfrak{q}$ is the prime ideal of $S_{(g)}$ corresponding to $\mathfrak{p}$ (2.3.6); by definition, this is equivalent to $f \in \mathfrak{p}$, whence the proposition. $\square$

(2.6.4). Now let $\mathcal{F}$ be an $\mathcal{O}_X$ -module, and set $M = \Gamma_\bullet(\mathcal{F})$; by the existence of the homomorphism of graded rings $\alpha : S \rightarrow \Gamma_\bullet(\mathcal{O}_X)$, we can consider M as a graded S -module. For every $f \in S_d$ ($d > 0$), it follows from (2.6.3) that the restriction to $D_+(f)$ of the section $\alpha_d(f)$ of $\mathcal{O}_X(d)$ is invertible; thus so too is the restriction to $D_+(f)$ of the section $\alpha_{nd}(f^n)$ of $\mathcal{O}_X(nd)$, for every $n > 0$. So let $z \in M_{nd} = \Gamma(X, \mathcal{F}(nd))$ ($n > 0$); if there exists an integer $k \geq 0$ such that the restriction to $D_+(f)$ of $f^k z$, i.e. the $[[3.38]]$ section $(z|_{D_+(f)})(\alpha_{kd}(f^k)|_{D_+(f)})$ of $\mathcal{F}((n+k)d)$, is zero, then, by the above remark, we also have that $z|_{D_+(f)} = 0$. This shows that we have defined an $S_{(f)}$ -homomorphism $\beta_f : M_{(f)} \rightarrow \Gamma(D_+(f), \mathcal{F})$ by sending the element z/f^n to the section $(z|_{D_+(f)})(\alpha_{nd}(f^n)|_{D_+(f)})^{-1}$ of $\mathcal{F}$ over $D_+(f)$. We immediately verify that the diagram

$$(2.6.4.1) \quad \begin{array}{ccc} M_{(f)} & \xrightarrow{\beta_f} & \Gamma(D_+(f), \mathcal{F}) \\ \downarrow & & \downarrow \\ M_{(fg)} & \xrightarrow{\beta_{fg}} & \Gamma(D_+(fg), \mathcal{F}) \end{array}$$

commutes for $g \in S_e$ ($e > 0$). If we recall that $M_{(f)}$ is canonically identified with $\Gamma(D_+(f), \tilde{M})$, and that the $D_+(f)$ form a base for the topology of X (2.3.4), then we see that the β_f come from a unique canonical homomorphism of $\mathcal{O}_X$ -modules

$$(2.6.4.2) \quad \beta : (\Gamma_\bullet(\mathcal{F}))^\sim \longrightarrow \mathcal{F}$$

(also denoted by $\beta_{\mathcal{F}}$) which is evidently functorial.

Proposition (2.6.5). — *Let M be a graded S -module, and $\mathcal{F}$ an $\mathcal{O}_X$ -module; then the composite homomorphisms*

$$(2.6.5.1) \quad \tilde{M} \xrightarrow{\tilde{\alpha}} (\Gamma_\bullet(\tilde{M}))^\sim \xrightarrow{\beta} \tilde{M}$$

$$(2.6.5.2) \quad \Gamma_\bullet(\mathcal{F}) \xrightarrow{\alpha} \Gamma_\bullet((\Gamma_\bullet(\mathcal{F}))^\sim) \xrightarrow{\Gamma_\bullet(\beta)} \Gamma_\bullet(\mathcal{F})$$

are the identity isomorphisms.

Proof. The proof for (2.6.5.1) is local: in an open subset $D_+(f)$, it follows immediately from the definitions, along with the fact that β , applied to quasi-coherent sheaves, is determined by its action on the sections over $D_+(f)$ (I, 1.3.8). The assertion (2.6.5.2) is verified separately in each degree: if we set $M = \Gamma_\bullet(\mathcal{F})$, then $M_n = \Gamma(X, \mathcal{F}(n))$, and $(\Gamma_\bullet(\tilde{M}))_n = \Gamma(X, \tilde{M}(n)) = \Gamma(X, (M(n))^\sim)$. But if $f \in S_1$ and $z \in M_n$, then $\alpha_n^f(z)$ is the element $z/1$ of $(M(n))_{(f)}$, equal to $(f/1)^n(z/f^n)$; it corresponds, via β_f , to the section

$$\left((\alpha_1(f))^n |_{D_+(f)} \right) \left((z |_{D_+(f)}) ((\alpha_1(f))^n |_{D_+(f)})^{-1} \right)$$

over $D_+(f)$, i.e. the restriction of z to $D_+(f)$, which finishes the proof for (2.6.5.2). $\square$

2.7. Finiteness conditions

Proposition (2.7.1). — (i) *If S is a graded Noetherian ring, then $X = \text{Proj}(S)$ is a Noetherian scheme.*
(ii) *If S is a graded A -algebra of finite type, then $X = \text{Proj}(S)$ is a scheme of finite type over $Y = \text{Spec}(A)$.*

[II:39]

Proof.

- (i) If S is Noetherian, then the ideal S_+ admits a finite system of homogeneous generators f_i ($1 \leq i \leq p$), thus (2.3.14) the underlying space X is the union of the $D_+(f_i) = \text{Spec}(S_{(f_i)})$, and everything then reduces to showing that each of the $S_{(f_i)}$ is Noetherian, which follows from (2.2.6).
- (ii) The hypothesis implies that S_0 is an A -algebra of finite type, and that S is an S_0 -algebra of finite type, and so S_+ is an ideal of finite type (2.1.4). We are thus reduced, as in (i), to showing that $S_{(f)}$ is an A -algebra of finite type for all $f \in S_d$. By (2.2.5), it suffices to show that $S^{(d)}$ is an A -algebra of finite type, which follows from (2.1.6). $\square$

(2.7.2). In what follows, we consider the following finiteness conditions for a graded S -module M :

- (TF) There exists an integer n such that the submodule $\bigoplus_{k \geq n} M_k$ is an S -module of finite type.
- (TN) There exists an integer n such that $M_k = 0$ for $k \geq n$.

If M satisfies (TN), then $M_{(f)} = 0$ for all homogeneous f in S_+ , and thus $\tilde{M} = 0$.

Let M and N be graded S -modules; we say that a homomorphism $u : M \rightarrow N$ of degree 0 is (TN)-injective (resp. (TN)-surjective, (TN)-bijective) if there exists an integer n such that $u_k : M_k \rightarrow N_k$ is injective (resp. surjective, bijective) for $k \geq n$. Saying that u is (TN)-injective (resp. (TN)-surjective) thus reduces to saying that $\text{Ker } u$ (resp. $\text{Coker } u$) satisfies (TN). By (2.5.4), if u is (TN)-injective (resp. (TN)-surjective, (TN)-bijective), then $\tilde{u}$ is injective (resp. surjective, bijective); if u is (TN)-bijective, then we also say that u is a (TN)-isomorphism.

Proposition (2.7.3). — *Let S be a graded ring such that the ideal S_+ is of finite type, and let M be a graded S -module.*

- (i) If M satisfies condition (TF) then the $\mathcal{O}_X$ -module $\widetilde{M}$ is of finite type.
- (ii) Suppose that M satisfies (TF); for $\widetilde{M} = 0$, it is necessary and sufficient for M to satisfy (TN).

Proof. We have just seen that condition (TN) implies that $\widetilde{M} = 0$. If M satisfies (TF), then the graded submodule $M' = \bigoplus_{k \geq n} M_k$, which is, by hypothesis, of finite type, is such that M/M' satisfies (TN); thus $(M/M')^\sim = 0$, and the exactness of the functor $M \mapsto \widetilde{M}$ (2.5.4) implies that $\widetilde{M} = \widetilde{M}'$; to prove that $\widetilde{M}$ is of finite type, we can thus reduce to the case where M is of finite type. Since the question is local, it suffices to prove that $M_{(f)}$ is an $S_{(f)}$ -module of finite type (I, 1.3.9); but $M^{(d)}$ is an $S^{(d)}$ -module of finite type (2.1.6, iii), and our claim then follows from (2.2.5).

Now suppose that M satisfies (TF) and that $\widetilde{M} = 0$; then, with the same notation as above, we have that $\widetilde{M}' = 0$, and condition (TN) for M' is equivalent to condition (TN) for M , so to prove that $\widetilde{M} = 0$ implies that M satisfies (TN), we can again restrict to the case where M is generated by a finite number of homogeneous elements x_i ($1 \leq i \leq p$); let $(f_j)_{1 \leq j \leq q}$ be a system of homogeneous generators of the ideal S_+ . By hypothesis, $M_{(f_j)} = 0$ for all j , and so there exists an integer n such that $f_j^n x_i = 0$ for any i and j . Let n_j be the degree of f_j , and let m be the largest value of $\sum_j r_j n_j$ over the finitely many systems of nonnegative integers (r_j) such that $\sum_j r_j \leq nq$; it is then clear [II:40] that, if $k > m$, then $S_k x_i = 0$ for all i ; if h is the largest of the degrees of the x_i , then we conclude that $M_k = 0$ for $k > h + m$, which finishes the proof. $\square$

Corollary (2.7.4). — Let S be a graded ring such that the ideal S_+ is of finite type; for $X = \text{Proj}(S) = \emptyset$, it is necessary and sufficient for there to exist n such that $S_k = 0$ for $k \geq n$.

Proof. The condition $X = \emptyset$ is equivalent to $\mathcal{O}_X = \widetilde{S} = 0$, and S is a cyclic S -module. $\square$

Theorem (2.7.5). — Suppose that the ideal S_+ is generated by a finite number of homogeneous elements of degree 1; let $X = \text{Proj}(S)$. Then, for every quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$, the canonical homomorphism $\beta : (\Gamma_\bullet(\mathcal{F}))^\sim \rightarrow \mathcal{F}$ (2.6.4) is an isomorphism.

Proof. If S_+ is generated by a finite number of elements $f_i \in S_1$, then X is the union of the subspaces $\text{Spec}(S_{(f_i)})$ (2.3.6), which are quasi-compact, and so X is quasi-compact; furthermore, X is a scheme (2.4.2); by (I, 9.3.2), (2.5.14.2), and (2.6.3), we have, for all $f \in S_d$ ($d > 0$), a canonical isomorphism $(\Gamma_\bullet(\mathcal{F}))_{(\alpha_d(f))} \xrightarrow{\sim} \Gamma(D_+(f), \mathcal{F})$; also, by definition, $(\Gamma_\bullet(\mathcal{F}))_{(\alpha_d(f))}$ (where $\Gamma_\bullet(\mathcal{F})$ is considered as a $\Gamma_\bullet(\mathcal{O}_X)$ -module) is exactly $(\Gamma_\bullet(\mathcal{F}))_{(f)}$ (where $\Gamma_\bullet(\mathcal{F})$ is considered as an S -module); if we refer to the definition (I, 9.3.1) of the above canonical isomorphism, then we see that it agrees with the homomorphism β_f , whence the theorem. $\square$

Remark (2.7.6). — If we suppose that the graded ring S is Noetherian, then the condition of (2.7.5) is satisfied *ipso facto* as soon as we suppose that the ideal S_+ is generated by the set S_1 of homogeneous elements of degree 1.

Corollary (2.7.7). — Under the hypotheses of (2.7.5), every quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$ is isomorphic to an $\mathcal{O}_X$ -module of the form $\widetilde{M}$, where M is a graded S -module.

Corollary (2.7.8). — Under the hypotheses of (2.7.5), every quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$ of finite type is isomorphic to an $\mathcal{O}_X$ -module of the form $\widetilde{N}$, where N is a graded S -module of finite type.

Proof. We can suppose that $\mathcal{F} = \widetilde{M}$, where M is a graded S -module (2.7.7). Let $(f_\lambda)_{\lambda \in L}$ be a system of homogeneous generators of M ; for every finite subset H of L , let M_H be the graded submodule of M generated by the f_λ such that $\lambda \in H$; it is clear that M is the inductive limit of its submodules M_H , and so $\mathcal{F}$ is the inductive limit of its sub- $\mathcal{O}_X$ -modules $\widetilde{M}_H$ (2.5.4). But, since $\mathcal{F}$ is of finite type, and since the underlying space of X is quasi-compact, it follows from (0, 5.2.3) that $\mathcal{F} = \widetilde{M}_H$ for some finite subset H of L . $\square$

Corollary (2.7.9). — Under the hypotheses of (2.7.5), let $\mathcal{F}$ be a quasi-coherent $\mathcal{O}_X$ -module of finite type. Then there exists an integer n_0 such that, for all $n \geq n_0$, $\mathcal{F}(n)$ is isomorphic to a quotient of an $\mathcal{O}_X$ -module of the form $\mathcal{O}_X^k$ (for some $k > 0$ depending on n), and is thus generated by a finite number of its sections over X (0, 5.1.1).

Proof. By (2.7.8), we can suppose that $\mathcal{F} = \tilde{M}$, where M is a quotient of a finite direct sum of S -modules of the form $S(m_i)$; by (2.5.4), we can thus restrict to the case where $M = S(m)$, and so $\mathcal{F}(n) = (S(m+n))^\sim = \mathcal{O}_X(m+n)$ (2.5.15). It thus suffices to prove

[41]

Lemma (2.7.9.1). — *Under the hypotheses of (2.7.5), for all $n \geq 0$, there exists an integer k (depending on n) and a surjective homomorphism $\mathcal{O}_X^k \rightarrow \mathcal{O}_X(n)$.*

It suffices (2.7.2) to show that, for suitable k , there is a (TN)-surjective homomorphism u of degree 0 from the graded product S -module S^k to $S(n)$. But $(S(n))_0 = S_n$, and, by hypothesis, $S_h = S_1^h$ for all $h > 0$, and so $SS_n = S_n + S_{n+1} + \dots$. Since S_n is an S_0 -module of finite type ((2.1.5) and (2.1.6, i)), consider a system of generators $(a_i)_{1 \leq i \leq k}$ of this module; consider the homomorphism u that sends the i -th element of the canonical basis of S^k to a_i ($1 \leq i \leq k$); since $\text{Coker } u$ can then be identified with $(S(n))_{-n} + \dots + (S(n))_{-1}$, u is indeed the desired homomorphism. $\square$

Corollary (2.7.10). — *Under the hypotheses of (2.7.5), let $\mathcal{F}$ be a quasi-coherent $\mathcal{O}_X$ -module of finite type. Then there exists an integer n_0 such that, for all $n \geq n_0$, $\mathcal{F}$ is isomorphic to a quotient of an $\mathcal{O}_X$ -module of the form $(\mathcal{O}_X(-n))^k$ (for some k depending on n).*

Proposition (2.7.11). — *Suppose that the hypotheses of (2.7.5) are satisfied, and let M be a graded S -module. Then:*

- (i) *The canonical homomorphism $\tilde{\alpha} : \tilde{M} \rightarrow (\Gamma_\bullet(\tilde{M}))^\sim$ is an isomorphism.*
- (ii) *Let $\mathcal{G}$ be a quasi-coherent sub- $\mathcal{O}_X$ -module of $\tilde{M}$, and let N be the graded sub- S -module of M given by the inverse image of $\Gamma_\bullet(\mathcal{G})$ under α . Then $\tilde{N} = \mathcal{G}$ (where $\tilde{N}$ is identified, by (2.5.4), with a sub- $\mathcal{O}_X$ -module of $\tilde{M}$).*

Proof. Since $\beta : (\Gamma_\bullet(\tilde{M}))^\sim \rightarrow \tilde{M}$ is an isomorphism (by (2.7.5)), $\tilde{\alpha}$ is the inverse isomorphism (by (2.6.5.1)), whence (i). Let P be the graded submodule $\alpha(M)$ of $\Gamma_\bullet(\tilde{M})$; since the functor $M \mapsto \tilde{M}$ is exact (2.5.4), the image of $\tilde{M}$ under $\tilde{\alpha}$ is equal to $\tilde{P}$, and so, by (i), $\tilde{P} = (\Gamma_\bullet(\tilde{M}))^\sim$. Set $Q = \Gamma_\bullet(\mathcal{G}) \cap P$; by the above, and by (2.5.4), we have that $\tilde{Q} = (\Gamma_\bullet(\mathcal{G}))^\sim$, and so the restriction of β to $\tilde{Q}$ is an isomorphism from this $\mathcal{O}_X$ -module to $\mathcal{G}$ by (2.7.5). But, by the definition of N , and by (2.5.4), the restriction of the isomorphism $\tilde{\alpha}$ to $\tilde{N}$ is an isomorphism from $\tilde{N}$ to $\tilde{Q}$, whence the conclusion, by (2.6.5.1). $\square$

2.8. Functorial behaviour

(2.8.1). Let S and S' be positively graded rings, and $\phi : S' \rightarrow S$ a homomorphism of graded rings. We denote by $G(\phi)$ the open subset of $X = \text{Proj}(S)$ given by the complement of $V_+(\phi(S'_+))$, or, equivalently, the union of the $D_+(\phi(f'))$ where f' runs over the set of homogeneous elements of S'_+ . The restriction to $G(\phi)$ of the continuous map ${}^a\phi$ from $\text{Spec}(S)$ to $\text{Spec}(S')$ (I, 1.2.1) is thus a continuous map from $G(\phi)$ to $\text{Proj}(S')$, which we again denote, with an abuse of language, by ${}^a\phi$. If $f' \in S'_+$ is homogeneous, then

$$(2.8.1.1) \quad {}^a\phi^{-1}(D_+(f')) = D_+(\phi(f'))$$

taking into account the fact that ${}^a\phi$ sends $G(\phi)$ to $\text{Proj}(S')$, as well as (I, 1.2.2.2). The homomorphism ϕ also canonically defines (with the same notation) a homomorphism of graded rings $S'_{f'} \rightarrow S_f$, whence, by restriction to the degree 0 elements, [III:42] a homomorphism $S'_{(f')} \rightarrow S_{(f)}$, which we denote by $\phi_{(f)}$; there is a corresponding (I, 1.6.1) morphism of affine schemes $({}^a\phi_{(f)}, \tilde{\phi}_{(f)}) : \text{Spec}(S_{(f)}) \rightarrow \text{Spec}(S'_{(f')})$. If we canonically identify $\text{Spec}(S_{(f)})$ with the scheme induced by $\text{Proj}(S)$ on $D_+(f)$ (2.3.6), then we have defined a morphism $\Phi_f : D_+(f) \rightarrow D_+(f')$, and ${}^a\phi_{(f)}$ is identified with the restriction of ${}^a\phi$ to $D_+(f)$. It is also immediate that, if g' is another homogeneous element

of S'_+ , and $g = \phi(g')$, then the diagram

$$\begin{array}{ccc} D_+(f) & \xrightarrow{\Phi_f} & D_+(f') \\ \uparrow & & \uparrow \\ D_+(fg) & \xrightarrow{\Phi_{fg}} & D_+(f'g') \end{array}$$

commutes, by the fact that the diagram

$$\begin{array}{ccc} S'_{(f')} & \xrightarrow{\phi_{(f)}} & S_{(f)} \\ \omega_{f'g',f'} \downarrow & & \downarrow \omega_{fg,f} \\ S'_{(f'g')} & \xrightarrow{\phi_{(fg)}} & S_{(fg)} \end{array}$$

commutes. Taking the definition of $G(\phi)$, along with (2.3.3.2), we thus see that:

Proposition (2.8.2). — *Given a homomorphism of graded rings $\phi : S' \rightarrow S$, there exists exactly one morphism $(^a\phi, \tilde{\phi})$ from the induced prescheme $G(\phi)$ to $\text{Proj}(S')$ (said to be associated to ϕ , and denoted by $\text{Proj}(\phi)$) such that, for every homogeneous element $f' \in S'_+$, the restriction of this morphism to $D_+(\phi(f'))$ agrees with the morphism associated to the homomorphism $S'_{(f')} \rightarrow S_{(\phi(f'))}$ corresponding to ϕ .*

Proof. With the above notation, if $f' \in S'_d$, then the diagram

$$(2.8.2.1) \quad \begin{array}{ccc} S'_{(f')} & \xrightarrow{\phi_{(f)}} & S_{(f)} \\ \sim \downarrow & & \downarrow \sim \\ S'^{(d)} / (f' - 1)S'^{(d)} & \longrightarrow & S^{(d)} / (f - 1)S^{(d)} \end{array}$$

commutes (the vertical arrows being the isomorphisms (2.2.5)). □

Corollary (2.8.3). — (i) *The morphism $\text{Proj}(\phi)$ is affine.*

(ii) *If $\text{Ker}(\phi)$ is nilpotent (and, in particular, if ϕ is injective), then the morphism $\text{Proj}(\phi)$ is dominant.*

Proof. Claim (i) is an immediate consequence of (2.8.2) and (2.8.1.1). Claim (ii) follows since, if $\text{Ker}(\phi)$ is nilpotent, then, for every homogeneous $f' \in S'_+$, we immediately see that $\text{Ker}(\phi_f)$ (with $f = \phi(f')$) is nilpotent, and thus so too is $\text{Ker}(\phi_{(f)})$; we then apply (2.8.2) and (I, 1.2.7). □

[II:43] We note that there are, in general, morphisms from $\text{Proj}(S)$ to $\text{Proj}(S')$ that are not affine, and that thus do not come from homomorphisms of graded rings $S' \rightarrow S$; an example is the structure morphism $\text{Proj}(S) \rightarrow \text{Spec}(A)$ when A is a field ($\text{Spec}(A)$ thus being identified with $\text{Proj}(A[T])$, cf. (3.1.7)); indeed, this follows from (I, 2.3.2).

(2.8.4). Let S'' be another positively graded ring, and $\phi' : S'' \rightarrow S'$ a homomorphism of graded rings, and set $\phi'' = \phi \circ \phi'$. By (2.8.1.1) and the formula $^a\phi'' = (^a\phi') \circ (^a\phi)$, we immediately see that $G(\phi'') \subset G(\phi)$, and that, if Φ, Φ' , and Φ'' are the morphisms associated to ϕ, ϕ' , and ϕ'' (respectively), then $\Phi'' = \Phi' \circ (\Phi|_{G(\phi'')})$.

(2.8.5). Suppose that S (resp. S') is a graded A -algebra (resp. a graded A' -algebra), and let $\psi : A' \rightarrow A$ be a ring homomorphism such that the diagram

$$(2.8.5.1) \quad \begin{array}{ccc} A' & \xrightarrow{\psi} & A \\ \downarrow & & \downarrow \\ S' & \xrightarrow{\phi} & S \end{array}$$

commutes. We can then consider $G(\phi)$ (resp. $\text{Proj}(S')$) as a scheme over $\text{Spec}(A)$ (resp. $\text{Spec}(A')$); if Φ (resp. Ψ) is the morphism associated to ϕ (resp. ψ), then the diagram

$$\begin{array}{ccc} G(\phi) & \xrightarrow{\Phi} & \text{Proj}(S') \\ \downarrow & & \downarrow \\ \text{Spec}(A) & \xrightarrow{\Psi} & \text{Spec}(A') \end{array}$$

commutes: it suffices to prove this for the restriction of Φ to $D_+(f)$, where $f = \phi(f')$, with f' homogeneous in S'_+ ; this then follows from the fact that the diagram

$$\begin{array}{ccc} A' & \xrightarrow{\psi} & A \\ \downarrow & & \downarrow \\ S'_{(f')} & \xrightarrow{\phi_{(f)}} & S_{(f)} \end{array}$$

commutes.

(2.8.6). Now let M be a graded S -module, and consider the S' -module $M_{[\phi]}$, which is evidently graded. Let f' be homogeneous in S'_+ , and let $f = \phi(f')$; we know (0, 1.5.2) that there is a canonical isomorphism $(M_{[\phi]})_{f'} \xrightarrow{\sim} (M_f)_{[\phi_f]}$, and it is immediate that this isomorphism preserves degree, whence a canonical isomorphism $(M_{[\phi]})_{(f')} \xrightarrow{\sim} (M_{(f)})_{[\phi_{(f)}]}$. To this isomorphism, there canonically corresponds an isomorphism of sheaves $(M_{[\phi]})^\sim|_{D_+(f')} \xrightarrow{\sim} (\Phi_f)_*(\tilde{M}|_{D_+(f)})$ ((2.5.2) and (I, 1.6.3)). Furthermore, [II:4.4] if g' is another homogeneous element of S'_+ , and $g = \phi(g')$, then the diagram

$$\begin{array}{ccc} (M_{[\phi]})_{(f')} & \xrightarrow{\sim} & (M_{(f)})_{[\phi_{(f)}]} \\ \downarrow & & \downarrow \\ (M_{[\phi]})_{(f'g')} & \xrightarrow{\sim} & (M_{(fg)})_{[\phi_{(fg)}]} \end{array}$$

commutes, whence we immediately conclude that the isomorphism

$$(M_{[\phi]})^\sim|_{D_+(f'g')} \xrightarrow{\sim} (\Phi_{fg})_*(\tilde{M}|_{D_+(fg)})$$

is the restriction to $D_+(f'g')$ of the isomorphism $(M_{[\phi]})^\sim|_{D_+(f')} \xrightarrow{\sim} (\Phi_f)_*(\tilde{M}|_{D_+(f)})$. Since Φ_f is the restriction to $D_+(f)$ of the morphism Φ , we see that, taking (2.8.1.1) into account, and setting $X' = \text{Proj}(S')$:

Proposition (2.8.7). — *There exists a canonical functorial isomorphism from the $\mathcal{O}_{X'}$ -module $(M_{[\phi]})^\sim$ to the $\mathcal{O}_{X'}$ -module $\Phi_*(\tilde{M}|_{G(\phi)})$.*

We thus immediately deduce a canonical functorial map from the set of ϕ -morphisms $M' \rightarrow M$ from a graded S' -module to the graded S -module M , to the set of Φ -morphisms $\tilde{M}' \rightarrow \tilde{M}|_{G(\phi)}$. With the notation of (2.8.4), if M'' is a graded S'' -module, then, to the composition of a ϕ -morphism $M' \rightarrow M$ and a ϕ' -morphism $M'' \rightarrow M'$, canonically corresponds the composition of $\tilde{M}'|_{G(\phi')} \rightarrow \tilde{M}|_{G(\phi'')} and $\tilde{M}'' \rightarrow \tilde{M}'|_{G(\phi')}$.$

Proposition (2.8.8). — *Under the hypotheses of (2.8.1), let M' be a graded S' -module. Then there exists a canonical functorial homomorphism ν from the $(\mathcal{O}_X|_{G(\phi)})$ -module $\Phi^*(\tilde{M}')$ to the $(\mathcal{O}_X|_{G(\phi)})$ -module $(M' \otimes_{S'} S)^\sim|_{G(\phi)}$. If the ideal S'_+ is generated by S'_1 , then ν is an isomorphism.*

Proof. Indeed, for $f' \in S'_d$ ($d > 0$), we define a canonical functorial homomorphism of $S_{(f)}$ -modules (where $f = \phi(f')$)

$$(2.8.8.1) \quad \nu_f : M'_{(f')} \otimes_{S'_{(f')}} S_{(f)} \longrightarrow (M' \otimes_{S'} S)_{(f)}$$

by composing the homomorphism $M'_{(f')} \otimes_{S'_{(f')}} S_{(f)} \rightarrow M'_{f'} \otimes_{S'_{f'}} S_f$ and the canonical isomorphism $M'_{f'} \otimes_{S'_{f'}} S_f \xrightarrow{\sim} (M' \otimes_{S'} S)_f$ (0, 1.5.4), and noting that the latter preserves degrees. We can immediately verify the compatibility of ν_f with the restriction operators from $D_+(f)$ to $D_+(fg)$ (for any $g' \in S'_+$ and $g = \phi(g')$), whence the definition of the homomorphism

$$\nu : \Phi^*(\widetilde{M}') \longrightarrow (M' \otimes_{S'} S)^\sim | G(\phi)$$

taking (I, 1.6.5) into account. To prove the second claim, it suffices to show that ν_f is an isomorphism for all $f' \in S'_1$, since $G(\phi)$ is then a union of the $D_+(\phi(f'))$. We first define a $\mathbf{Z}$ -bilinear $M'_m \times S_n \rightarrow M'_{(f')} \otimes_{S'_{(f')}} S_{(f)}$ by sending (x', s) to the element $(x' / f'^m) \otimes (s / f'^n)$ (with the convention that x' / f'^m is $f'^{-m} x' / 1$ when $m < 0$). [II:45] We claim that, in the proof of (2.5.13), this map gives rise to a di-homomorphism of modules

$$\eta_f : M' \otimes_{S'} S \longrightarrow M'_{(f')} \otimes_{S'_{(f')}} S_{(f)}.$$

Furthermore, if, for $r > 0$, we have $f'^r \sum_i (x'_i \otimes s_i) = 0$, then this can also be written as $\sum_i (f'^r x'_i \otimes s_i) = 0$, whence, by (0, 1.5.4), $\sum_i (f'^r x'_i / f'^{m_i+r}) \otimes (s_i / f'^{n_i}) = 0$, i.e. $\eta_f(\sum_i x'_i \otimes s_i) = 0$, which proves that η_f factors as $M' \otimes_{S'} S \rightarrow (M' \otimes_{S'} S)_f \xrightarrow{\eta'_f} M'_{(f')} \otimes_{S'_{(f')}} S_{(f)}$; we finally can prove that η'_f and ν_f are inverse isomorphisms to one another.

In particular, it follows from (2.1.2.1) that we have a canonical homomorphism

$$(2.8.8.2) \quad \Phi^*(\mathcal{O}_{X'}(n)) \xrightarrow{\sim} \mathcal{O}_X(n) | G(\phi)$$

for all $n \in \mathbf{Z}$. □

(2.8.9). Let A and A' be rings, and $\psi : A' \rightarrow A$ a ring homomorphism, defining a morphism $\Psi : \text{Spec}(A) \rightarrow \text{Spec}(A')$. Let S' be a positively graded A' -algebra, and set $S = S' \otimes_{A'} A$, which is evidently an A -algebra graded by the $S'_n \otimes_{A'} A$; the map $\phi : s' \rightarrow s' \otimes 1$ is then a graded ring homomorphism that makes the diagram (2.8.5.1) commute. Since S_+ is here the A -module generated by $\phi(S'_+)$, we have $G(\phi) = \text{Proj}(S) = X$; whence, setting $X' = \text{Proj}(S')$, we have the commutative diagram

$$(2.8.9.1) \quad \begin{array}{ccc} X & \xrightarrow{\Phi} & X' \\ p \downarrow & & \downarrow \\ Y & \xrightarrow{\Psi} & Y' \end{array}$$

Now let M' be a graded S' -module, and set $M = M' \otimes_{A'} A = M' \otimes_{S'} S$. Under these conditions:

Proposition (2.8.10). — *The diagram (2.8.9.1) identifies the scheme X with the product $X' \times_{Y'} Y$; furthermore, the canonical homomorphism $\nu : \Phi^*(\widetilde{M}') \rightarrow \widetilde{M}$ (2.8.8) is an isomorphism.*

Proof. The first claim will be proven if we show that, for every homogeneous f' in S'_+ , setting $f = \phi(f')$, the restrictions of Φ and p to $D_+(f)$ identify this scheme with the product $D_+(f') \times_{Y'} Y$ (I, 3.2.6.2); in other words, it suffices (I, 3.2.2) to prove that $S_{(f)}$ is canonically identified with $S'_{(f')} \otimes_{A'} A$, which follows immediately from the canonical degree-preserving isomorphism $S_f \xrightarrow{\sim} S'_{f'} \otimes_{A'} A$ (0, 1.5.4). The second claim then follows from the fact that $M'_{(f')} \otimes_{S'_{(f')}} S_{(f)}$ can be identified, by the above, with $M'_{(f')} \otimes_{A'} A$, and this can be identified with $M_{(f)}$, since M_f is canonically identified with $M'_{f'} \otimes_{A'} A$ by an isomorphism that preserves degrees. □

Corollary (2.8.11). — *For every integer $n \in \mathbf{Z}$, $\widetilde{M}(n)$ can be identified with $\Phi^*(\widetilde{M}'(n)) = \widetilde{M}'(n) \otimes_{Y'} \mathcal{O}_Y$; in particular, $\mathcal{O}_X(n) = \Phi^*(\mathcal{O}_{X'}(n)) = \mathcal{O}_{X'}(n) \otimes_{Y'} \mathcal{O}_Y$.*

Proof. This follows from (2.8.10) and (2.5.15). □

(2.8.12). Under the hypotheses of (2.8.9), for $f' \in S'_d$ ($d > 0$) and $f = \phi(f')$, the diagram

$$(2.8.12.1) \quad \begin{array}{ccc} M'_{(f')} & \xrightarrow{\sim} & M'^{(d)} / (f' - 1)M'^{(d)} \\ \downarrow & & \downarrow \\ M_{(f)} & \xrightarrow{\sim} & M^{(d)} / (f - 1)M^{(d)} \end{array}$$

(cf. (2.2.5)) commutes.

(2.8.13). Keep the notation and hypotheses of (2.8.9), and let $\mathcal{F}'$ be an $\mathcal{O}_{X'}$ -module; if we set $\mathcal{F} = \Phi^*(\mathcal{F}')$, then, for all $n \in \mathbf{Z}$, we have $\mathcal{F}(n) = \Phi^*(\mathcal{F}'(n))$, by (2.8.11) and (0, 4.3.3). Then, by (0, 3.7.1), we have a canonical homomorphism

$$\Gamma(\rho) : \Gamma(X', \mathcal{F}'(n)) \longrightarrow \Gamma(X, \mathcal{F}(n))$$

which gives a canonical di-homomorphism of graded modules

$$\Gamma_{\bullet}(\mathcal{F}') \longrightarrow \Gamma_{\bullet}(\mathcal{F}).$$

Suppose that the ideal S_+ is generated by S_1 , and that $\mathcal{F}' = \widetilde{M}'$, thus $\mathcal{F} = \widetilde{M}$ with $M = M' \otimes_{A'} A$. If f' is homogeneous in S'_+ , and $f = \phi(f')$, then we have seen that $M_{(f)} = M'_{(f')} \otimes_{A'} A$, and the diagram

$$\begin{array}{ccc} M'_0 & \longrightarrow & M'_{(f')} & = \Gamma(D_+(f'), \widetilde{M}') \\ \downarrow & & \downarrow & \\ M_0 & \longrightarrow & M_{(f)} & = \Gamma(D_+(f), \widetilde{M}) \end{array}$$

thus commutes; we immediately conclude from this remark, and from the definition of the homomorphism α (2.6.2), that the diagram

$$(2.8.13.1) \quad \begin{array}{ccc} M' & \xrightarrow{\alpha_{M'}} & \Gamma_{\bullet}(\widetilde{M}') \\ \downarrow & & \downarrow \\ M & \xrightarrow{\alpha_M} & \Gamma_{\bullet}(\widetilde{M}) \end{array}$$

commutes. Similarly, the diagram

$$(2.8.13.2) \quad \begin{array}{ccc} (\Gamma_{\bullet}(\mathcal{F}'))^{\sim} & \xrightarrow{\beta_{\mathcal{F}'}} & \mathcal{F}' \\ \downarrow & & \downarrow \\ (\Gamma_{\bullet}(\mathcal{F}))^{\sim} & \xrightarrow{\beta_{\mathcal{F}}} & \mathcal{F} \end{array}$$

commutes (the vertical arrow on the right being the canonical Φ -morphism $\mathcal{F}' \rightarrow \Phi^*(\mathcal{F}') = \mathcal{F}$).

[II:47]

(2.8.14). Still keeping the notation and hypotheses of (2.8.9), let N' be another graded S' -module, and let $N = N' \otimes_{A'} A$. It is immediate that the canonical di-homomorphisms $M' \rightarrow M$ and $N' \rightarrow N$ give a di-homomorphism $M' \otimes_{S'} N' \rightarrow M \otimes_S N$ (with respect to the canonical ring homomorphism $S' \rightarrow S$), and thus also an S -homomorphism $(M' \otimes_{S'} N') \otimes_{A'} A \rightarrow M \otimes_S N$ of degree 0, to which corresponds (taking (2.8.10) into account) a homomorphism of $\mathcal{O}_X$ -modules

$$(2.8.14.1) \quad \Phi^*((M' \otimes_{S'} N')^{\sim}) \longrightarrow (M \otimes_S N)^{\sim}.$$

Furthermore, we can immediately verify that the diagram

$$(2.8.14.2) \quad \begin{array}{ccc} \Phi^*(\widetilde{M}' \otimes_{\mathcal{O}_{X'}} \widetilde{N}') & \xrightarrow{\sim} & \widetilde{M} \otimes_{\mathcal{O}_X} \widetilde{N} & = & \Phi^*(\widetilde{M}') \otimes_{\mathcal{O}_X} \Phi^*(\widetilde{N}') \\ \Phi^*(\lambda) \downarrow & & \downarrow \lambda & & \\ \Phi^*((M' \otimes_S N')^\sim) & \longrightarrow & (M \otimes_S N)^\sim & & \end{array}$$

commutes, with the first line being the canonical isomorphism (0, 4.3.3). If the ideal S'_+ is generated by S'_1 , then it is clear that S_+ is generated by S_1 , and the two vertical arrows of (2.8.14.2) are then isomorphisms (2.5.13); it is thus also the case for (2.8.14.1).

We similarly have a canonical di-homomorphism $\text{Hom}_{S'}(M', N') \rightarrow \text{Hom}_S(M, N)$ by sending a homomorphism u' of degree k to the homomorphism $u' \otimes 1$, which is also of degree k ; from this, we again deduce a S -homomorphism of degree 0

$$(\text{Hom}_{S'}(M', N')) \otimes_{A'} A \longrightarrow \text{Hom}_S(M, N)$$

whence a homomorphism of $\mathcal{O}_X$ -modules

$$(2.8.14.3) \quad \Phi^*((\text{Hom}_{S'}(M', N'))^\sim) \longrightarrow (\text{Hom}_S(M, N))^\sim.$$

Furthermore, the diagram

$$\begin{array}{ccc} \Phi^*((\text{Hom}_{S'}(M', N'))^\sim) & \longrightarrow & (\text{Hom}_S(M, N))^\sim \\ \Phi^*(\mu) \downarrow & & \downarrow \mu \\ \Phi^*(\mathcal{H}om_{\mathcal{O}_{X'}}(\widetilde{M}', \widetilde{N}')) & \longrightarrow & \mathcal{H}om_{\mathcal{O}_X}(\widetilde{M}, \widetilde{N}) \end{array}$$

commutes (the second horizontal line being the canonical homomorphism (0, 4.4.6)).

[II:48]

(2.8.15). With the notation and hypotheses of (2.8.1), it follows from (2.4.7) that we do not change the morphism Φ , up to isomorphism, when we replace S_0 and S'_0 by $\mathbb{Z}$, and ϕ_0 by the identity map, and thus when we replace S and S' by $S^{(d)}$ and $S'^{(d)}$ (respectively) ($d > 0$), and ϕ by its restriction $\phi^{(d)}$ to $S^{(d)}$.

2.9. Closed subschemes of a scheme $\text{Proj}(S)$

(2.9.1). If $\phi : S \rightarrow S'$ is a homomorphism of graded rings, then we say that ϕ is (TN)-surjective (resp. (TN)-injective, (TN)-bijective) if there exists an integer n such that, for $k \geq n$, $\phi_k : S_k \rightarrow S'_k$ is surjective (resp. injective, bijective). It then follows from (2.8.15) that the study of Φ may be reduced to the case where ϕ is surjective (resp. injective, bijective). Instead of saying that ϕ is (TN)-bijective, we sometimes say that it is a (TN)-isomorphism.

Proposition (2.9.2). — *Let S be a positively graded ring, and let $X = \text{Proj}(S)$.*

- (i) *If $\phi : S \rightarrow S'$ is a (TN)-surjective homomorphism of graded rings, then the corresponding morphism Φ (2.8.1) is defined on the whole of $\text{Proj}(S')$, and is a closed immersion of $\text{Proj}(S')$ into X . If $\mathfrak{J}$ is the kernel of ϕ , then the closed subscheme of X associated to Φ is defined by the quasi-coherent sheaf of ideals $\widetilde{\mathfrak{J}}$ of $\mathcal{O}_X$.*
- (ii) *Suppose further that the ideal S_+ is generated by a finite number of homogeneous elements of degree 1. Let X' be a closed subscheme of X defined by a quasi-coherent sheaf of ideals $\mathcal{J}$ of $\mathcal{O}_X$. Let $\mathfrak{J}$ be the graded ideal of S given by the inverse image of $\Gamma_\bullet(\mathcal{J})$ under the canonical homomorphism $\alpha : S \rightarrow \Gamma_\bullet(\mathcal{O}_X)$ (2.6.2), and set $S' = S/\mathfrak{J}$. Then X' is the subscheme associated to the closed immersion $\text{Proj}(S') \rightarrow X$ corresponding to the canonical homomorphism of graded rings $S \rightarrow S'$.*

Proof.

- (i) We can suppose that ϕ is surjective (2.9.1). Since, by hypothesis, $\phi(S_+)$ generates S'_+ , we have $G(\phi) = \text{Proj}(S')$. Now, the second claim can be checked locally on X ; so let f be a homogeneous element of S_+ , and set $f' = \phi(f)$. Since ϕ is a surjective homomorphism of

graded rings, we immediately see that $\phi_{(f')} : S_{(f)} \rightarrow S'_{(f')}$ is surjective, and that its kernel is $\mathfrak{J}_{(f)}$, which proves (i) (I, 4.2.3).

- (ii) By (i), we are led to proving that the homomorphism $\tilde{j} : \tilde{\mathfrak{J}} \rightarrow \mathcal{O}_X$ induced by the canonical injection $j : \mathfrak{J} \rightarrow S$ is an isomorphism from $\tilde{\mathfrak{J}}$ to $\mathcal{J}$, which follows from (2.7.11). $\square$

We note that $\mathfrak{J}$ is the *largest* of the graded ideals $\mathfrak{J}'$ of S such that $\tilde{j}(\mathfrak{J}') = \mathcal{J}$, since we can immediately show, using the definitions (2.6.2), that this equation implies that $\alpha(\mathfrak{J}') \subset \Gamma_\bullet(\mathcal{J})$.

Corollary (2.9.3). — Suppose that the hypotheses of (2.9.2, (i)) are satisfied, and further that the ideal S_+ is generated by S_1 ; then $\Phi^*((S(n))^\sim)$ is canonically isomorphic to $(S'(n))^\sim$ for all $n \in \mathbf{Z}$, and so $\Phi^*(\mathcal{F}(n))$ is canonically isomorphic to $\Phi^*(\mathcal{F})(n)$ for every $\mathcal{O}_X$ -module $\mathcal{F}$.

Proof. This is a particular case of (2.8.8), taking (2.5.10.2) into account. $\square$

Corollary (2.9.4). — Suppose that the hypotheses of (2.9.2, (ii)) are satisfied. For the closed sub-prescheme X' of X to be integral, it is necessary and sufficient for the graded ideal $\mathfrak{J}$ to be prime in S .

[II:49]

Proof. Since X' is isomorphic to $\text{Proj}(S/\mathfrak{J})$, the condition is sufficient by (2.4.4). To see that it is necessary, consider the exact sequence $0 \rightarrow \mathcal{J} \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_X/\mathcal{J} \rightarrow 0$, which gives the exact sequence

$$0 \rightarrow \Gamma_\bullet(\mathcal{J}) \rightarrow \Gamma_\bullet(\mathcal{O}_X) \rightarrow \Gamma_\bullet(\mathcal{O}_X/\mathcal{J}).$$

It suffices to prove that, if $f \in S_m$ and $g \in S_n$ are such that the image in $\Gamma_\bullet(\mathcal{O}_X/\mathcal{J})$ of $\alpha_{n+m}(fg)$ is zero, then the image of either $\alpha_m(f)$ or $\alpha_n(g)$ is zero. But, by definition, these images are sections of invertible $(\mathcal{O}_X/\mathcal{J})$ -modules $\mathcal{L} = (\mathcal{O}_X/\mathcal{J})(m)$ and $\mathcal{L}' = (\mathcal{O}_X/\mathcal{J})(n)$ over the integral scheme X' ; the hypothesis implies that the product of these two sections is zero in $\mathcal{L} \otimes \mathcal{L}'$ ((2.9.3) and (2.5.14.1)), and so one of them is zero by (I, 7.4.4). $\square$

Corollary (2.9.5). — Let A be a ring, M an A -module, S a graded A -algebra generated by the set S_1 of homogeneous elements of degree 1, $u : M \rightarrow S_1$ a surjective homomorphism of A -modules, and $\bar{u} : \mathbf{S}(M) \rightarrow S$ the homomorphism (of A -algebras) from the symmetric algebra $\mathbf{S}(M)$ of M to S that extends u . Then the morphism corresponding to $\bar{u}$ is a closed immersion of $\text{Proj}(S)$ into $\text{Proj}(\mathbf{S}(M))$.

Proof. Indeed, $\bar{u}$ is surjective by hypothesis, and so it suffices to apply (2.9.2). $\square$

§3. HOMOGENEOUS SPECTRUM OF A SHEAF OF GRADED ALGEBRAS

3.1. Homogeneous spectrum of a quasi-coherent graded $\mathcal{O}_Y$ -algebra

(3.1.1). Let Y be a prescheme, $\mathcal{S}$ a graded $\mathcal{O}_Y$ -algebra, and $\mathcal{M}$ a graded $\mathcal{S}$ -module. If $\mathcal{S}$ is *quasi-coherent*, then each of its homogeneous components $\mathcal{S}_n$ is a *quasi-coherent* $\mathcal{O}_Y$ -module, since they are the images of $\mathcal{S}$ under a homomorphism from $\mathcal{S}$ to itself ((I, 1.3.8) and (I, 1.3.9)); similarly, if $\mathcal{M}$ is quasi-coherent as an $\mathcal{O}_Y$ -module, then its homogeneous components $\mathcal{M}_n$ are also quasi-coherent, and the converse is also true. For an integer $d > 0$, we denote by $\mathcal{S}^{(d)}$ the direct sum of the $\mathcal{O}_Y$ -modules $\mathcal{S}_{nd}$ (for $n \in \mathbf{Z}$), which is quasi-coherent if $\mathcal{S}$ is quasi-coherent (I, 1.3.9); for every integer k such that $0 \leq k \leq d-1$, we denote by $\mathcal{M}^{(d,k)}$ (or $\mathcal{M}^{(d)}$, for $k=0$) the direct sum of the $\mathcal{M}_{nd+k}$ (for $n \in \mathbf{Z}$), which is a graded $\mathcal{S}^{(d)}$ -module, and quasi-coherent if both $\mathcal{S}$ and $\mathcal{M}$ are quasi-coherent (I, 9.6.1). We denote by $\mathcal{M}(n)$ the graded $\mathcal{S}$ -module such that $(\mathcal{M}(n))_k = \mathcal{M}_{n+k}$ for all $k \in \mathbf{Z}$; if $\mathcal{S}$ and $\mathcal{M}$ are quasi-coherent, then $\mathcal{M}(n)$ is a quasi-coherent graded $\mathcal{S}$ -module (I, 9.6.1).

We say that $\mathcal{M}$ is a graded $\mathcal{S}$ -module of *finite type* (resp. admitting a *finite presentation*) if, for all $y \in Y$, there exists an open neighbourhood U of y , along with integers n_i (resp. integers m_i and n_j) such that there is a surjective degree 0 homomorphism $\bigoplus_{i=1}^r (\mathcal{S}(n_i)|U) \rightarrow \mathcal{M}|U$ (resp. such that $\mathcal{M}|U$ is isomorphic to the cokernel of a degree 0 homomorphism $\bigoplus_{i=1}^r (\mathcal{S}(m_i)|U) \rightarrow \bigoplus_{j=1}^s (\mathcal{S}(n_j)|U)$).

Let U be an affine open of Y , with ring $A = \Gamma(U, \mathcal{O}_Y)$; by hypothesis, the graded $(\mathcal{O}_Y|U)$ -algebra $\mathcal{S}|U$ is isomorphic to $\tilde{S}$, where $S = \Gamma(U, \mathcal{S})$ is a graded A -algebra (I, 1.4.3); [II:50] set $X_U = \text{Proj}(\Gamma(U, \mathcal{S}))$. Let $U' \subset U$ be another affine open of Y , with ring A' , and let $j : U' \rightarrow U$ be

the canonical injection, which corresponds to the restriction homomorphism $A \rightarrow A'$; we have that $\mathcal{S}|U' = j^*(\mathcal{S}|U)$, and so $S' = \Gamma(U', \mathcal{S})$ is canonically identified with $S \otimes_A A'$ (I, 1.6.5). We conclude (2.8.10) that $X_{U'}$ is canonically identified with $X_U \times_U U'$, and thus also with $f_U^{-1}(U')$, where we denote by f_U the structure morphism $X_U \rightarrow U$ (I, 4.4.1). We denote by $\sigma_{U',U}$ the canonical isomorphism $f_U^{-1}(U') \xrightarrow{\sim} X_{U'}$ thus defined, and by $\rho_{U',U}$ the open immersion $X_{U'} \rightarrow X_U$ obtained by composing $\sigma_{U',U}^{-1}$ with the canonical injection $f_U^{-1}(U') \rightarrow X_U$. It is immediate that, if $U'' \subset U'$ is another affine open of Y , then $\rho_{U'',U} = \rho_{U',U} \circ \rho_{U'',U'}$.

Proposition (3.1.2). — *Let Y be a prescheme. For every quasi-coherent positively graded $\mathcal{O}_Y$ -algebra $\mathcal{S}$, there exists exactly one (up to Y -isomorphism) prescheme X over Y satisfying the following property: if $f : X \rightarrow Y$ is the structure morphism, then, for every affine open U of Y , there exists an isomorphism η_U from the induced prescheme $f^{-1}(U)$ to $X_U = \text{Proj}(\Gamma(U, \mathcal{S}))$ such that, if V is another affine open of Y that is contained in U , then the diagram*

$$(3.1.2.1) \quad \begin{array}{ccc} f^{-1}(V) & \xrightarrow{\eta_V} & X_V \\ \downarrow & & \downarrow \rho_{V,U} \\ f^{-1}(U) & \xrightarrow{\eta_U} & X_U \end{array}$$

commutes.

Proof. Given affine opens U and V of Y , let $X_{U,V}$ be the prescheme induced on $f_U^{-1}(U \cap V)$ by X_U ; we are going to define a Y -isomorphism $\theta_{U,V} : X_{U,V} \xrightarrow{\sim} X_{U,V}$. For this, we consider an affine open $W \subset U \cap V$: by composing the isomorphisms

$$f_U^{-1}(W) \xrightarrow{\sigma_{W,U}} X_W \xrightarrow{\sigma_{W,V}^{-1}} f_V^{-1}(W),$$

we obtain an isomorphism τ_W , and we immediately see that, if $W' \subset W$ is an affine open, then $\tau_{W'}$ is the restriction of τ_W to $f_U^{-1}(W')$; the τ_W are thus indeed the restrictions of a Y -isomorphism $\theta_{U,V}$. Further, if U, V , and W are affine open subsets of Y , and $\theta'_{U,V}, \theta'_{V,W}$, and $\theta'_{U,W}$ the restrictions of $\theta_{U,V}, \theta_{V,W}$, and $\theta_{U,W}$ (respectively) to the inverse images of $U \cap V \cap W$ in X_U, X_V , and X_W (respectively), then it follows from the above definitions that we have $\theta'_{U,V} \circ \theta'_{V,W} = \theta'_{U,W}$. The existence of some X satisfying the properties in the statement thus follows from (I, 2.3.1); its uniqueness up to Y -isomorphism is trivial, taking (3.1.2.1) into account. $\square$

(3.1.3). We say that the prescheme X defined in (3.1.2) is the *homogeneous spectrum* of the quasi-coherent graded $\mathcal{O}_Y$ -algebra $\mathcal{S}$, and we denote it by $\text{Proj}(\mathcal{S})$. It is immediate that $\text{Proj}(\mathcal{S})$ is *separated over Y* ((2.4.2) and (I, 5.5.5)); if $\mathcal{S}$ is an $\mathcal{O}_Y$ -algebra of *finite type* (I, 9.6.2), then $\text{Proj}(\mathcal{S})$ is of *finite type over Y* ((2.7.1, (ii)) and (I, 6.3.1)).

If f is the structure morphism $X \rightarrow Y$, then it is immediate that, for every prescheme induced by Y on an open subset U of Y , $f^{-1}(U)$ can be identified with the homogeneous spectrum $\text{Proj}(\mathcal{S}|U)$.

Proposition (3.1.4). — *Let $f \in \Gamma(Y, \mathcal{S}_d)$ for $d > 0$. Then there exists an open subset X_f of the underlying space of $X = \text{Proj}(\mathcal{S})$ that satisfies the following property: for every affine open U of Y , we have $X_f \cap \phi^{-1}(U) = D_+(f|U)$ in $\phi^{-1}(U)$ identified with $X_U = \text{Proj}(\Gamma(U, \mathcal{S}))$, where ϕ denotes the structure morphism $X \rightarrow Y$. [II:51] Furthermore, the prescheme induced on X_f is affine over Y , and is canonically isomorphic to $\text{Spec}(\mathcal{S}^{(d)} / (f - 1)\mathcal{S}^{(d)})$ (1.3.1).*

Proof. We have $f|U \in \Gamma(U, \mathcal{S}_d) = (\Gamma(U, \mathcal{S}))_d$. If U and U' are affine opens of Y such that $U' \subset U$, then $f|U'$ is the image of $f|U$ under the restriction homomorphism

$$\Gamma(U, \mathcal{S}) \longrightarrow \Gamma(U', \mathcal{S})$$

and so $D_+(f|U')$ is equal (with the notation of (3.1.1)) to the prescheme induced on the inverse image $\rho_{U',U}^{-1}(D_+(f|U))$ in $X_{U'}$ (2.8.1); whence the first claim. Furthermore, the prescheme induced on $D_+(f|U)$ by X_U is canonically identified with $\text{Spec}((\Gamma(U, \mathcal{S}))_{(f|U)})$, with these identifications being compatible with the restriction homomorphisms (2.8.1); the second claim then follows from (2.2.5) and from the commutativity of the diagram (2.8.2.1). $\square$

We also say that X_f (as an open subset of the underlying space X) is the set of $x \in X$ where f does not vanish.

Corollary (3.1.5). — If $f \in \Gamma(Y, \mathcal{S}_d)$ and $g \in \Gamma(Y, \mathcal{S}_e)$, then

$$(3.1.5.1) \quad X_{fg} = X_f \cap X_g.$$

Proof. It suffices to consider the intersection of the two sets with a set $\phi^{-1}(U)$, where U is an affine open in Y , and to then apply formula (2.3.3.2). $\square$

Corollary (3.1.6). — Let (f_α) be a family of sections of $\mathcal{S}$ over Y such that $f_\alpha \in \Gamma(Y, \mathcal{S}_{d_\alpha})$; if the sheaf of ideals of $\mathcal{S}$ generated by this family (0, 5.1.1) contains all the $\mathcal{S}_n$ starting from a certain rank, then the underlying space X is the union of the X_{f_α} .

Proof. For every affine open U of Y , $\phi^{-1}(U)$ is the union of the $X_{f_\alpha} \cap \phi^{-1}(U)$ (2.3.14). $\square$

Corollary (3.1.7). — Let $\mathcal{A}$ be a quasi-coherent $\mathcal{O}_Y$ -algebra; set

$$\mathcal{S} = \mathcal{A}[T] = \mathcal{A} \otimes_{\mathbf{Z}} \mathbf{Z}[T]$$

where T is an indeterminate (and $\mathbf{Z}$ and $\mathbf{Z}[T]$ are considered as simple sheaves over Y). Then $X = \text{Proj}(\mathcal{S})$ is canonically identified with $\text{Spec}(\mathcal{A})$. In particular, $\text{Proj}(\mathcal{O}_Y[T])$ is identified with Y .

Proof. By applying (3.1.6) to the unique section $f \in \Gamma(Y, \mathcal{S})$ that is equal to T at each point of Y we see that $X_f = X$. Further, here we have $d = 1$, and $\mathcal{S}^{(1)} / (f - 1)\mathcal{S}^{(1)} = \mathcal{S} / (f - 1)\mathcal{S}$ is canonically isomorphic to $\mathcal{A}$, whence the corollary (1.2.2). $\square$

Let $g \in \Gamma(Y, \mathcal{O}_Y)$; if we take $\mathcal{S} = \mathcal{O}_Y[T]$, then $g \in \Gamma(Y, \mathcal{S}_0)$; let

$$h = gT \in \Gamma(Y, \mathcal{S}_1).$$

If $X = \text{Proj}(\mathcal{S})$, then the canonical identification defined in (3.1.7) identifies X_h with the open subset Y_g of Y (in the sense of (0, 5.5.2)): indeed, we can restrict to the case where $Y = \text{Spec}(A)$ is affine, and everything then reduces (taking (2.2.5) into account) to the fact that the ring of fractions A_g is canonically identified with $A[T] / (gT - 1)A[T]$ (0, 1.2.3).

Proposition (3.1.8). — Let $\mathcal{S}$ be a quasi-coherent positively graded $\mathcal{O}_Y$ -algebra. Then

- (i) For all $d > 0$, there exists a canonical Y -isomorphism from $\text{Proj}(\mathcal{S})$ to $\text{Proj}(\mathcal{S}^{(d)})$. [III:52]
- (ii) Let $\mathcal{S}'$ be the graded $\mathcal{O}_Y$ -algebra given by the direct sum of $\mathcal{O}_Y$ with the $\mathcal{S}_n$ (for $n \geq 0$); then $\text{Proj}(\mathcal{S}')$ and $\text{Proj}(\mathcal{S})$ are canonically Y -isomorphic.
- (iii) Let $\mathcal{L}$ be an invertible $\mathcal{O}_Y$ -module (0, 5.4.1), and let $\mathcal{S}_{(\mathcal{L})}$ be the graded $\mathcal{O}_Y$ -algebra given by the direct sum of the $\mathcal{S}_d \otimes \mathcal{L}^{\otimes d}$ (for $d \geq 0$); then $\text{Proj}(\mathcal{S})$ and $\text{Proj}(\mathcal{S}_{(\mathcal{L})})$ are canonically Y -isomorphic.

Proof. In each of the three cases, it suffices to define the isomorphism locally on Y , since the verification of compatibility with the restriction operations from one open subset to a smaller one is trivial. We can thus suppose that Y is affine, and then (i) follows from (2.4.7, (i)), and (ii) follows from (2.4.8). As for (iii), if we further suppose that $\mathcal{L}$ is isomorphic to $\mathcal{O}_Y$ (which we are allowed to do, since the question is local on Y), then the isomorphism between $\text{Proj}(\mathcal{S})$ and $\text{Proj}(\mathcal{S}_{(\mathcal{L})})$ is evident; to define a canonical isomorphism, let $Y = \text{Spec}(A)$ and $\mathcal{S} = \tilde{S}$, where S is a graded A -algebra, and let c be a generator of the free A -module L such that $\mathcal{L} = \tilde{L}$; then, for all $n > 0$, $x_n \mapsto x_n \otimes c^{\otimes n}$ is an A -isomorphism from S_n to $S_n \otimes L^{\otimes n}$, and these A -isomorphisms define an A -isomorphism of graded algebras $\phi_c : S \rightarrow S_{(L)} = \bigoplus_{n \geq 0} S_n \otimes L^{\otimes n}$. So let $f \in S_+$ be homogeneous of degree d ; for all $x \in S_{nd}$, we have that

$$\frac{x \otimes c^{\otimes nd}}{(f \otimes c^{\otimes d})^n} = \frac{x \otimes (\varepsilon c)^{\otimes nd}}{(f \otimes (\varepsilon c)^{\otimes d})^n}$$

for every invertible element $\varepsilon \in A$, which shows that the isomorphism $S_{(f)} \rightarrow (S_{(L)})_{(f \otimes c^{\otimes d})}$ induced from ϕ_c is independent of the generator c of L considered, and thus finishes the proof. $\square$

(3.1.9). Recall ((0, 4.1.3) and (I, 1.3.14)) that, for the quasi-coherent graded $\mathcal{O}_Y$ -algebra $\mathcal{S}$ to be generated by the $\mathcal{O}_Y$ -module $\mathcal{S}_1$, it is necessary and sufficient for there to exist an affine open cover (U_α) of Y such that the graded algebra $\Gamma(U_\alpha, \mathcal{S})$ over $\Gamma(U_\alpha, \mathcal{S}_0)$ is generated by the set $\Gamma(U_\alpha, \mathcal{S}_1)$ of its homogeneous elements of degree 1. For every open V of Y , $\mathcal{S}|_V$ is then generated by the $(\mathcal{O}_Y|_V)$ -module $\mathcal{S}_1|_V$.

Proposition (3.1.10). — Suppose that there exists a finite affine open cover (U_i) of Y such that each graded algebra $\Gamma(U_i, \mathcal{S})$ is of finite type over $\Gamma(U_i, \mathcal{O}_Y)$. Then there exists $d > 0$ such that $\mathcal{S}^{(d)}$ is generated by $\mathcal{S}_d$, with $\mathcal{S}_d$ an $\mathcal{O}_Y$ -module of finite type.

Proof. Indeed, it follows from (2.1.6, (v)) that, for each i , there exists an integer m_i such that $\Gamma(U_i, \mathcal{S}_{nm_i}) = (\Gamma(U_i, \mathcal{S}_{m_i}))^n$ for all $n > 0$; it suffices to take d to be a common multiple of all the m_i , taking (2.1.6, (i)) into account. $\square$

Corollary (3.1.11). — Under the hypotheses of (3.1.10), $\text{Proj}(\mathcal{S})$ is Y -isomorphic to a homogeneous spectrum $\text{Proj}(\mathcal{S}')$, where $\mathcal{S}'$ is a graded $\mathcal{O}_Y$ -algebra generated by $\mathcal{S}'_1$, with $\mathcal{S}'_1$ an $\mathcal{O}_Y$ -module of finite type.

Proof. It suffices to take $\mathcal{S}' = \mathcal{S}^{(d)}$, where d satisfies the property of (3.1.10), and to then apply (3.1.8, (i)). $\square$

(3.1.12). If $\mathcal{S}$ is a quasi-coherent positively-graded $\mathcal{O}_Y$ -algebra, we know (I, 5.1.1) that its nilradical $\mathcal{N}$ is a quasi-coherent $\mathcal{O}_Y$ -module; we say that $\mathcal{N}_+ = \mathcal{N} \cap \mathcal{S}_+$ is the nilradical of $\mathcal{S}_+$; this is a quasi-coherent graded $\mathcal{S}_0$ -module, since we can immediately reduce to the case where Y is affine, and the proposition then follows from (2.1.10). For all $y \in Y$, $(\mathcal{N}_+)_y$ is then the nilradical of $(\mathcal{S}_+)_y = (\mathcal{S}_y)_+$ (I, 5.1.1). We say that the graded $\mathcal{O}_Y$ -algebra $\mathcal{S}$ is *essentially reduced* if $\mathcal{N}_+ = 0$, which is equivalent [II:53] to saying that $\mathcal{S}_y$ is an essentially reduced graded $\mathcal{O}_y$ -algebra for all $y \in Y$. For every graded $\mathcal{O}_Y$ -algebra $\mathcal{S}$, $\mathcal{S}/\mathcal{N}_+$ is essentially reduced.

We say that $\mathcal{S}$ is *integral* if, for all $y \in Y$, $\mathcal{S}_y$ is an integral ring and, furthermore, $(\mathcal{S}_y)_+ = (\mathcal{S}_+)_y \neq 0$.

Proposition (3.1.13). — Let $\mathcal{S}$ be a positively-graded $\mathcal{O}_Y$ -algebra. If $X = \text{Proj}(\mathcal{S})$, then the Y -scheme X_{red} is canonically isomorphic to $\text{Proj}(\mathcal{S}/\mathcal{N}_+)$; in particular, if $\mathcal{S}$ is essentially reduced, then X is reduced.

Proof. The fact that $X' = \text{Proj}(\mathcal{S}/\mathcal{N}_+)$ is reduced follows immediately from (2.4.4, (i)), since the property is local; further, for every affine open $U \subset Y$, $\phi'^{-1}(U)$ is equal to $(\phi^{-1}(U))_{\text{red}}$ (where we denote by ϕ and ϕ' the structure morphisms $X \rightarrow Y$ and $X' \rightarrow Y$, respectively); we immediately see that the canonical U -morphisms $\phi'^{-1}(U) \rightarrow \phi^{-1}(U)$ are compatible with the restriction operations, and thus define a closed immersion $X' \rightarrow X$ that is a homeomorphism of the underlying spaces; whence the conclusion (I, 5.1.2). $\square$

Proposition (3.1.14). — Let Y be an integral prescheme, and $\mathcal{S}$ a quasi-coherent graded $\mathcal{O}_Y$ -algebra such that $\mathcal{S}_0 = \mathcal{O}_Y$.

- (i) If $\mathcal{S}$ is integral (3.1.12), then $X = \text{Proj}(\mathcal{S})$ is integral, and the structure morphism $\phi : X \rightarrow Y$ is dominant.
- (ii) Suppose furthermore that $\mathcal{S}$ is essentially reduced. Then, conversely, if X is integral and ϕ is dominant, then $\mathcal{S}$ is integral.

Proof.

- (i) If (U_α) is a base of Y consisting of non-empty affine opens, then it suffices to prove the proposition in the case where Y is replaced by one of the U_α , and $\mathcal{S}$ by $\mathcal{S}|_{U_\alpha}$: indeed, on the one hand it will follow that the underlying spaces $\phi^{-1}(U_\alpha)$ are irreducible open subsets (and thus non-empty) of X such that $\phi^{-1}(U_\alpha) \cap \phi^{-1}(U_\beta) \neq \emptyset$ for any pair of indices (since $U_\alpha \cap U_\beta$ contains some U_γ), and so X is irreducible (0, 2.1.4); on the other hand, X will be reduced, since this is a local property, and so X will indeed be integral, with $\phi(X)$ dense in Y .

So suppose that $Y = \text{Spec}(A)$, where A is integral (I, 5.1.4), and that $\mathcal{S} = \tilde{S}$, where S is a graded A -algebra; the hypothesis is that, for every $y \in Y$, $\tilde{S}_y = S_y$ is an integral graded ring such that $(S_y)_+ \neq 0$. It suffices to prove that S is an *integral* ring, since then we will

have that $S_+ \neq 0$, and we can then apply (2.4.4, (ii)). But let $f, g \neq 0$ be elements of S , and suppose that $fg = 0$; for all $y \in Y$ we then have that $(f/1)(g/1) = 0$ in S_y , and so $f/1 = 0$ or $g/1 = 0$ by hypothesis. Suppose, for example, that $f/1 = 0$ in S_y ; this implies that there exists $a \in A$ such that $a \notin \mathfrak{y}_y$ and $af = 0$; we then have, for all $z \in Y$, that $(a/1)(f/1) = 0$ in the integral ring S_z , and since $a/1 \neq 0$ (since A is integral), $f/1 = 0$, which implies that $f = 0$.

- (ii) Since the question is local on Y , we can again suppose that $Y = \text{Spec}(A)$, with A integral, and that $\mathcal{S} = \tilde{S}$. By hypothesis, for all $y \in Y$, $(S_y)_+$ does not contain any non-zero nilpotent elements, and the same is true of $(S_0)_y = A_y$ by hypothesis; so S_y is a reduced ring for all $y \in Y$, and we thus conclude first of all that S itself is reduced (I, 5.1.1). The hypothesis that X is integral implies that S is essentially integral (2.4.4, (ii)), and everything then reduces to showing that the annihilator $\mathfrak{J}$ of S_+ in $A = S_0$ is just 0 (2.1.11). If this were not the case, we would have that $(S_h)_+ = 0$ for some $h \neq 0$ in $\mathfrak{J}$, and thus (3.1.1) that $\phi^{-1}(D(h)) = \emptyset$, and $\phi(X)$ would not be dense in Y , contradicting the hypothesis (since $D(h) \neq \emptyset$, since h is not nilpotent). [II:54]

□

3.2. Sheaf on $\text{Proj}(\mathcal{S})$ associated to a graded $\mathcal{S}$ -module

(3.2.1). Let Y be a prescheme, $\mathcal{S}$ a quasi-coherent positively graded $\mathcal{O}_Y$ -algebra, and $\mathcal{M}$ a quasi-coherent graded $\mathcal{S}$ -module (on $(Y, \mathcal{O}_Y)$ or, equivalently (I, 9.6.1), on the ringed space $(Y, \mathcal{S})$). With the notation of (3.1.1), we denote by $\tilde{\mathcal{M}}_U$ the quasi-coherent $\mathcal{O}_{X_U}$ -module $(\Gamma(U, \mathcal{M}))^\sim$; for $U' \subset U$, $\Gamma(U', \mathcal{M})$ is canonically identified with $\Gamma(U, \mathcal{M}) \otimes_A A'$ (I, 1.6.4); thus we have $\tilde{\mathcal{M}}_{U'} = \rho_{U',U}^*(\tilde{\mathcal{M}}_U)$ (2.8.11).

Proposition (3.2.2). — *There exists on $\text{Proj}(\mathcal{S}) = X$ a unique quasi-coherent $\mathcal{O}_X$ -module $\tilde{\mathcal{M}}$ such that, for every affine open U of Y , we have $\eta_U^*((\Gamma(U, \mathcal{M}))^\sim) = \tilde{\mathcal{M}}|_{f^{-1}(U)}$ (where η_U denotes the isomorphism $f^{-1}(U) \xrightarrow{\sim} \text{Proj}(\Gamma(U, \mathcal{S}))$), with f the structure morphism $X \rightarrow Y$.*

Proof. Since $\rho_{U',U}$ is identified with the injection morphism $f^{-1}(U') \rightarrow f^{-1}(U)$ (3.1.2.1), the proposition follows immediately from the relation $\tilde{\mathcal{M}}_{U'} = \rho_{U',U}^*(\tilde{\mathcal{M}}_U)$ and from the gluing principle for sheaves (0, 3.3.1). □

We say that $\tilde{\mathcal{M}}$ is the $\mathcal{O}_X$ -module associated to the quasi-coherent graded $\mathcal{S}$ -module $\mathcal{M}$.

Proposition (3.2.3). — *Let $\mathcal{M}$ be a quasi-coherent graded $\mathcal{S}$ -module, and let $f \in \Gamma(Y, \mathcal{S}_d)$ (for $d > 0$). If ξ_f is the canonical isomorphism from X_f to the Y -prescheme $Z_f = \text{Spec}(\mathcal{S}^{(d)} / (f - 1)\mathcal{S}^{(d)})$ (3.1.4), then $(\xi_f)_*(\tilde{\mathcal{M}}|_{X_f})$ is the $\mathcal{O}_{Z_f}$ -module $(\mathcal{M}^{(d)} / (f - 1)\mathcal{M}^{(d)})^\sim$ (1.4.3).*

Proof. Since the question is local on Y , we can immediately reduce to (2.2.5), taking into account the commutativity of the diagram in (2.8.12.1). □

Proposition (3.2.4). — *The assignment $\mathcal{M} \mapsto \tilde{\mathcal{M}}$ is an exact additive covariant functor from the category of quasi-coherent graded $\mathcal{S}$ -modules to the category of quasi-coherent $\mathcal{O}_X$ -modules, and it commutes with inductive limits and direct sums.*

Proof. Since the question is local on Y , we can reduce to (I, 1.3.11), (I, 1.3.9), and (2.5.4). □

In particular, if $\mathcal{N}$ is a quasi-coherent graded sub- $\mathcal{S}$ -module of $\mathcal{M}$, then $\tilde{\mathcal{N}}$ is canonically identified with a quasi-coherent sub- $\mathcal{O}_X$ -module of $\tilde{\mathcal{M}}$; in particular, for every quasi-coherent graded sheaf $\mathcal{I}$ of ideals of $\mathcal{S}$, $\tilde{\mathcal{I}}$ is a quasi-coherent sheaf of ideals of $\mathcal{O}_X$.

If $\mathcal{M}$ is a quasi-coherent graded $\mathcal{S}$ -module, and $\mathcal{I}$ a quasi-coherent sheaf of ideals of $\mathcal{O}_Y$, then $\mathcal{I} \cdot \mathcal{M}$ is a quasi-coherent graded sub- $\mathcal{S}$ -module of $\mathcal{M}$, and we have

$$(3.2.4.1) \quad (\mathcal{I} \cdot \mathcal{M})^\sim = \tilde{\mathcal{I}} \cdot \tilde{\mathcal{M}}$$

(where the right-hand side is defined as in (0, 4.3.5)). It suffices to verify this formula in the case where $Y = \text{Spec}(A)$ is affine, $\mathcal{S} = \tilde{S}$, with S a graded A -algebra, $\mathcal{M} = \tilde{M}$, with M a graded S -module, and $\mathcal{I} = \tilde{\mathfrak{I}}$, with $\mathfrak{I}$ an ideal of A . [II:55] For every homogeneous element f of S_+ , the

restriction to $D_+(f) = \text{Spec}(S_{(f)})$ of the left-hand side of (3.2.4.1) can be associated with $(\mathfrak{I}M)_{(f)} = \mathfrak{I} \cdot M_{(f)}$, and the same is true of the restriction of the right-hand side, given (I, 1.3.13) and (I, 1.6.9).

Proposition (3.2.5). — *Let $f \in \Gamma(Y, \mathcal{S}_d)$ (for $d > 0$). On the open subset X_f , the $(\mathcal{O}_X|_{X_f})$ -module $(\mathcal{S}(nd))^\sim|_{X_f}$ is canonically isomorphic to $\mathcal{O}_X|_{X_f}$ for all $n \in \mathbf{Z}$. In particular, if the $\mathcal{O}_Y$ -algebra $\mathcal{S}$ is generated by $\mathcal{S}_1$ (3.1.9), then the $\mathcal{O}_X$ -modules $(\mathcal{S}(n))^\sim$ are invertible for all $n \in \mathbf{Z}$.*

Proof. Indeed, for every affine open U of Y , we defined in (2.5.7) a canonical isomorphism from $(\mathcal{S}(nd))^\sim|(X_f \cap \phi^{-1}(U))$ to $\mathcal{O}_X|(X_f \cap \phi^{-1}(U))$, taking (3.1.4) into account (where ϕ is the structure morphism $X \rightarrow Y$); it is immediate that these isomorphisms are compatible with the restriction from U to an affine open $U' \subset U$, whence the first claim. To prove the second, it suffices to note that, if $\mathcal{S}$ is generated by $\mathcal{S}_1$, then there is a cover (U_α) of Y by affine opens such that $\Gamma(U_\alpha, \mathcal{S})$ is generated by $\Gamma(U_\alpha, \mathcal{S})_1 = \Gamma(U_\alpha, \mathcal{S}_1)$; we can then apply the result of (2.5.9), since the property of being invertible is local. $\square$

We again set, for all $n \in \mathbf{Z}$,

$$(3.2.5.1) \quad \mathcal{O}_X(n) = (\mathcal{S}(n))^\sim$$

and, for all $\mathcal{O}_X$ -modules $\mathcal{F}$,

$$(3.2.5.2) \quad \mathcal{F}(n) = \mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{O}_X(n).$$

It follows immediately from these definitions that, for every open subset U of Y , we have

$$((\mathcal{S}|_U)(n))^\sim = \mathcal{O}_X(n)|_{f^{-1}(U)}$$

where f is the structure morphism $X \rightarrow Y$.

Proposition (3.2.6). — *Let $\mathcal{M}$ and $\mathcal{N}$ be quasi-coherent graded $\mathcal{S}$ -modules. Then there exists a canonical functorial (in $\mathcal{M}$ and $\mathcal{N}$) homomorphism*

$$(3.2.6.1) \quad \lambda : \widetilde{\mathcal{M}} \otimes_{\mathcal{O}_X} \widetilde{\mathcal{N}} \longrightarrow (\mathcal{M} \otimes_{\mathcal{S}} \mathcal{N})^\sim$$

and a canonical functorial (in $\mathcal{M}$ and $\mathcal{N}$) homomorphism

$$(3.2.6.2) \quad \mu : (\mathcal{H}om_{\mathcal{S}}(\mathcal{M}, \mathcal{N}))^\sim \longrightarrow \mathcal{H}om_{\mathcal{O}_X}(\widetilde{\mathcal{M}}, \widetilde{\mathcal{N}}).$$

Furthermore, if $\mathcal{S}$ is generated by $\mathcal{S}_1$ (3.1.9), then λ is an isomorphism; if, further, $\mathcal{M}$ admits a finite presentation (3.1.1), then μ is an isomorphism.

Proof. The homomorphisms λ and μ were defined in (2.5.11.2) and (2.5.12.2) in the case where Y is affine; since these definitions are local, they transfer immediately to the general case considered here, taking (2.8.14) into account. $\square$

Corollary (3.2.7). — *If $\mathcal{S}$ is generated by $\mathcal{S}_1$, then, for any $m, n \in \mathbf{Z}$,*

$$(3.2.7.1) \quad \mathcal{O}_X(m) \otimes_{\mathcal{O}_X} \mathcal{O}_X(n) = \mathcal{O}_X(m+n)$$

$$(3.2.7.2) \quad \mathcal{O}_X(n) = (\mathcal{O}_X(1))^{\otimes n}$$

up to canonical isomorphism.

[II:56]

Corollary (3.2.8). — *If $\mathcal{S}$ is generated by $\mathcal{S}_1$, then, for any graded $\mathcal{S}$ -module $\mathcal{M}$ and any $n \in \mathbf{Z}$,*

$$(3.2.8.1) \quad (\mathcal{M}(n))^\sim = \widetilde{\mathcal{M}}(n)$$

up to canonical isomorphism.

Proof. This follows from the corresponding properties in the case where Y is affine ((2.5.14) and (2.5.15)), along with (2.8.11). $\square$

Remarks (3.2.9). —

- (1) If $\mathcal{S} = \mathcal{A}[T]$, with $\mathcal{A}$ a quasi-coherent $\mathcal{O}_Y$ -algebra (3.1.7), then we immediately see that all the invertible $\mathcal{O}_X$ -modules $\mathcal{O}_X(n)$ are canonically isomorphic to $\mathcal{O}_X$.
 Furthermore, let $\mathcal{N}$ be a quasi-coherent $\mathcal{A}$ -module, and set $\mathcal{M} = \mathcal{N} \otimes_{\mathcal{A}} \mathcal{A}[T]$. It then follows from (3.2.3) and (3.1.7) that, under the canonical identification of $X = \text{Proj}(\mathcal{A}[T])$ with $X' = \text{Spec}(\mathcal{A})$, the $\mathcal{O}_X$ -module $\mathcal{M}$ is identified with the $\mathcal{O}_{X'}$ -module $\widetilde{\mathcal{N}}$ associated to $\mathcal{N}$ (in the sense of (1.4.3)).
- (2) Let $\mathcal{S}$ be an arbitrary graded $\mathcal{O}_Y$ -algebra, and $\mathcal{S}'$ the graded $\mathcal{O}_Y$ -algebra such that $\mathcal{S}'_0 = \mathcal{O}_Y$ and $\mathcal{S}'_n = \mathcal{S}_n$ for all $n > 0$; then the canonical isomorphism from $X = \text{Proj}(\mathcal{S})$ to $X' = \text{Proj}(\mathcal{S}')$ (3.1.8, (ii)) identifies $\mathcal{O}_X(n)$ with $\mathcal{O}_{X'}(n)$ for all $n \in \mathbb{Z}$. This follows from the same proposition for the affine case (2.5.16) and from the fact that the identifications, for the affine opens of Y , commute with the restriction operations. Similarly, let $X^{(d)} = \text{Proj}(\mathcal{S}^{(d)})$; then the canonical isomorphism from X to $X^{(d)}$ (3.1.8, (i)) identifies $\mathcal{O}_X(nd)$ with $\mathcal{O}_{X^{(d)}}(n)$ for all $n \in \mathbb{Z}$.

Proposition (3.2.10). — *Let $\mathcal{L}$ be an invertible $\mathcal{O}_Y$ -module, and g the canonical isomorphism from $X_{(\mathcal{L})} = \text{Proj}(\mathcal{S}_{(\mathcal{L})})$ to $X = \text{Proj}(\mathcal{S})$ (3.1.8, (iii)). Then, for any $n \in \mathbb{Z}$, $g_*(\mathcal{O}_{X_{(\mathcal{L})}}(n))$ is canonically isomorphic to $\mathcal{O}_X(n) \otimes_Y \mathcal{L}^{\otimes n}$.*

Proof. Suppose first of all that Y is affine, with ring A , and that $\mathcal{L} = \widetilde{L}$, where L is a free A -module of rank one. With the notation from the proof of (3.1.8, (iii)), we define, for $f \in S_d$, an isomorphism from $(S(n))_{(f)} \otimes_A L^{\otimes n}$ to $(S_{(L)}(n))_{(f \otimes c^{\otimes d})}$ by sending $(x/f^k) \otimes c^{\otimes n}$, where $x \in S_{kd+n}$, to the element $(x \otimes c^{\otimes(n+kd)})/(f \otimes c^{\otimes d})^k$; it is immediate that this isomorphism is independent of the chosen generator c of L ; further, the isomorphisms thus defined for each $f \in S_+$ are compatible with the restriction operators $D_+(f) \rightarrow D_+(fg)$. Finally, in the general case, we easily see, from the definitions (3.1.1), that the isomorphisms thus defined for each affine open U of Y are compatible with passing from U to an affine open $U' \subset U$. $\square$

3.3. Graded $\mathcal{S}$ -module associated to a sheaf on $\text{Proj}(\mathcal{S})$ Throughout this entire section we suppose that the graded $\mathcal{O}_Y$ -algebra $\mathcal{S}$ is generated by $\mathcal{S}_1$ (3.1.9). Recall that, by (3.1.8, (i)), this restrictive assumption is not essential, thanks to finiteness conditions (3.1.10).

(3.3.1). Let p be the structure morphism $X = \text{Proj}(\mathcal{S}) \rightarrow Y$. For every $\mathcal{O}_X$ -module $\mathcal{F}$, set

$$(3.3.1.1) \quad \Gamma_*(\mathcal{F}) = \bigoplus_{n \in \mathbb{Z}} p_*(\mathcal{F}(n))$$

[II:57] and, in particular,

$$(3.3.1.2) \quad \Gamma_*(\mathcal{O}_X) = \bigoplus_{n \in \mathbb{Z}} p_*(\mathcal{O}_X(n)).$$

We know (0, 4.2.2) that there exists a canonical homomorphism

$$p_*(\mathcal{F}) \otimes_{\mathcal{O}_Y} p_*(\mathcal{G}) \longrightarrow p_*(\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{G})$$

for $\mathcal{O}_X$ -modules $\mathcal{F}$ and $\mathcal{G}$; we thus deduce from (3.2.7.1) that $\Gamma_*(\mathcal{O}_X)$ is endowed with the structure of a graded $\mathcal{O}_Y$ -algebra, and (3.2.5.2) defines the structure of a graded $\Gamma_*(\mathcal{O}_X)$ -module on $\Gamma_*(\mathcal{F})$.

By (3.2.5), and by left exactness of the functor p_* (0, 4.2.1), the assignment $\mathcal{F} \mapsto \Gamma_*(\mathcal{F})$ is an additive *left-exact* covariant functor from the category of $\mathcal{O}_X$ -modules to the category of graded $\mathcal{O}_Y$ -modules (where the morphisms are the homomorphisms of degree 0). In particular, if $\mathcal{I}$ is a sheaf of ideals of $\mathcal{O}_X$, then $\Gamma_*(\mathcal{I})$ can be identified with a *graded sheaf of ideals* in $\Gamma_*(\mathcal{O}_X)$.

(3.3.2). Let $\mathcal{M}$ be a quasi-coherent graded $\mathcal{S}$ -module. For every affine open U of Y , we defined in (2.6.2) a homomorphism of abelian groups

$$\alpha_{0,U} : \Gamma(U, \mathcal{M}_0) \longrightarrow \Gamma(p^{-1}(U), \widetilde{\mathcal{M}}).$$

It is immediate that these homomorphisms commute with the restriction operations (2.8.13.1) and thus define (without using the hypothesis that $\mathcal{S}$ is generated by $\mathcal{S}_1$) a homomorphism of sheaves of abelian groups

$$(3.3.2.1) \quad \alpha_0 : \mathcal{M}_0 \longrightarrow p_*(\widetilde{\mathcal{M}}).$$

Applying this result to each of the $\mathcal{M}_n = (\mathcal{M}(n))_0$, and taking (3.2.8.1) into account, we can define a homomorphism of sheaves of abelian groups

$$(3.3.2.2) \quad \alpha_n : \mathcal{M}_n \longrightarrow p_*(\widetilde{\mathcal{M}}(n))$$

for all $n \in \mathbf{Z}$, whence a functorial homomorphism (of degree 0) of graded sheaves of abelian groups

$$(3.3.2.3) \quad \alpha : \mathcal{M} \longrightarrow \Gamma_*(\widetilde{\mathcal{M}})$$

(also denoted $\alpha_{\mathcal{M}}$).

By taking $\mathcal{M} = \mathcal{S}$ in particular, we see that $\alpha : \mathcal{S} \rightarrow \Gamma_*(\mathcal{O}_X)$ is a homomorphism of graded $\mathcal{O}_Y$ -algebras, and that (3.3.2.3) is a semilinear homomorphism of graded modules with respect to this homomorphism of graded algebras.

We again note that to each of the α_n there corresponds (0, 4.4.3) a canonical homomorphism of $\mathcal{O}_X$ -modules

$$(3.3.2.4) \quad \alpha_n^\# : p^*(\mathcal{M}_n) \longrightarrow \widetilde{\mathcal{M}}(n).$$

We can easily verify that this homomorphism is exactly the one which corresponds functorially (3.2.4) to the canonical homomorphism (of degree 0) of graded $\mathcal{O}_Y$ -modules

$$(3.3.2.5) \quad \mathcal{M}_n \otimes_{\mathcal{O}_Y} \mathcal{S} \longrightarrow \mathcal{M}(n)$$

[II:58] where the grading of the right-hand side comes naturally from that of $\mathcal{S}$. We can restrict to the case where $Y = \text{Spec}(A)$ is affine, $\mathcal{M} = \widetilde{M}$, and $\mathcal{S} = \widetilde{S}$, with the graded A -algebra S being generated by S_1 , so that, as f runs over S_1 , the $D_+(f)$ form a cover of X . By the definitions (2.6.2), we then see, taking (I, 1.6.7) into account, that the restriction to $D_+(f)$ of the homomorphism (3.3.2.4) corresponds (I, 1.3.8) to the homomorphism of $S_{(f)}$ -modules $M_n \otimes_A S_{(f)} \rightarrow (M(n))_{(f)}$ that sends $x \otimes 1$ (where $x \in M_n$) to $x/1$; this proves the claim.

Proposition (3.3.3). — *For every section $f \in \Gamma(Y, \mathcal{S}_d)$ (where $d > 0$), X_f is precisely the set of points of X where $\alpha_d(f)$ (considered as a section of $\mathcal{O}_X(d)$) does not vanish (0, 5.5.2).*

Proof. (Note that $\alpha_d(f)$ is a section of $p_*(\mathcal{O}_X(d))$ over Y , but by definition such a section is also a section of $\mathcal{O}_X(d)$ over X (0, 4.2.1).) The definition of X_f (3.1.4) lets us reduce to the case where Y is affine, which has already been dealt with in (2.6.3). $\square$

(3.3.4). From now on, we suppose, in addition to the hypothesis at the start of this section, that, for every quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$, the $p_*(\mathcal{F}(n))$ are *quasi-coherent* on Y , so that $\Gamma_*(\mathcal{F}) = \bigoplus_{n \in \mathbf{Z}} p_*(\mathcal{F}(n))$ is also a quasi-coherent $\mathcal{O}_Y$ -module ((I, 1.4.1) and (I, 1.3.9)); this will always be the case if X is of *finite type* over Y (I, 9.2.2). We thus conclude that $(\Gamma_*(\mathcal{F}))^\sim$ is defined, and is a quasi-coherent $\mathcal{O}_X$ -module. For every affine open U of Y , we have ((I, 1.3.9) and (I, 2.5.4))

$$\begin{aligned} \left(\Gamma(U, \bigoplus_{n \in \mathbf{Z}} p_*(\mathcal{F}(n))) \right)^\sim &= \bigoplus_{n \in \mathbf{Z}} \left(\Gamma(U, p_*(\mathcal{F}(n))) \right)^\sim \\ &= \bigoplus_{n \in \mathbf{Z}} \left(\Gamma(p^{-1}(U), \mathcal{F}(n)) \right)^\sim \\ &= \left(\bigoplus_{n \in \mathbf{Z}} \Gamma(p^{-1}(U), \mathcal{F}(n)) \right)^\sim \\ &= (\Gamma_*(\mathcal{F}|_{p^{-1}(U)}))^\sim \end{aligned}$$

and hence, by (2.6.4), we have a canonical homomorphism

$$\beta_U : \left(\Gamma(U, \bigoplus_{n \in \mathbf{Z}} p_*(\mathcal{F}(n))) \right)^\sim \longrightarrow \mathcal{F}|_{p^{-1}(U)}.$$

Furthermore, the commutativity of (2.8.13.2) shows that these homomorphisms commute with the restriction operations on Y ; we thus obtain a canonical functorial homomorphism

$$(3.3.4.1) \quad \beta : (\Gamma_*(\mathcal{F}))^\sim \longrightarrow \mathcal{F}$$

(also denoted $\beta_{\mathcal{F}}$) for quasi-coherent $\mathcal{O}_X$ -modules.

Proposition (3.3.5). — Let $\mathcal{M}$ be a quasi-coherent graded $\mathcal{S}$ -module, and $\mathcal{F}$ a quasi-coherent $\mathcal{O}_X$ -module; then the composite homomorphisms

$$(3.3.5.1) \quad \widetilde{\mathcal{M}} \xrightarrow{\alpha} (\Gamma_*(\widetilde{\mathcal{M}}))^\sim \xrightarrow{\beta} \widetilde{\mathcal{M}}$$

$$(3.3.5.2) \quad \Gamma_*(\mathcal{F}) \xrightarrow{\alpha} \Gamma_*((\Gamma_*(\mathcal{F}))^\sim) \xrightarrow{\Gamma_*(\beta)} \Gamma_*(\mathcal{F})$$

are the identity isomorphisms.

Proof. The question is local on Y , so we can reduce to (2.6.5). $\square$

3.4. Finiteness conditions [II:59]

Proposition (3.4.1). — Let Y be a prescheme, and $\mathcal{S}$ a quasi-coherent $\mathcal{O}_Y$ -algebra generated by $\mathcal{S}_1$ (3.1.9); suppose further that $\mathcal{S}_1$ is of finite type. Then $X = \text{Proj}(\mathcal{S})$ is of finite type over Y .

Proof. Since the question is local on Y , we can suppose that Y is affine of ring A ; then $\mathcal{S} = \widetilde{S}$, where $S = \Gamma(Y, \mathcal{S})$, and by hypothesis S is an A -algebra generated by $S_1 = \Gamma(Y, \mathcal{S}_1)$, where we can further suppose that S_1 is an A -module of finite type ((I, 1.3.9) and (I, 1.3.12)). Then S is a graded A -algebra of finite type, and we can reduce to (2.7.1, (ii)). $\square$

(3.4.2). Let $\mathcal{S}$ be a quasi-coherent graded $\mathcal{O}_Y$ -algebra; for a quasi-coherent graded $\mathcal{S}$ -module $\mathcal{M}$, consider the following finiteness conditions:

(TF) There exists an integer n such that the $\mathcal{S}$ -module $\bigoplus_{k \geq n} \mathcal{M}_k$ is of finite type.

(TN) There exists an integer n such that $\mathcal{M}_k = 0$ for $k \geq n$.

If $\mathcal{M}$ satisfies (TN), then $\widetilde{\mathcal{M}} = 0$, since this is a local property on Y (2.7.2).

Let $\mathcal{M}$ and $\mathcal{N}$ be quasi-coherent graded $\mathcal{S}$ -modules; we say that a homomorphism $u : \mathcal{M} \rightarrow \mathcal{N}$ of degree 0 is (TN)-injective (resp. (TN)-surjective, (TN)-bijective) if there exists an integer n such that $u_k : \mathcal{M}_k \rightarrow \mathcal{N}_k$ is injective (resp. surjective, bijective) for $k \geq n$; then $\widetilde{u} : \widetilde{\mathcal{M}} \rightarrow \widetilde{\mathcal{N}}$ is injective (resp. surjective, bijective) by (2.7.2), since this is a local property on Y , and taking (I, 1.3.9) into account; if u is (TN)-bijective, then we also say that u is a (TN)-isomorphism.

Proposition (3.4.3). — Let Y be a prescheme, and $\mathcal{S}$ a quasi-coherent graded $\mathcal{O}_Y$ -algebra generated by $\mathcal{S}_1$, with $\mathcal{S}_1$ assumed to be of finite type. Let $\mathcal{M}$ be a quasi-coherent graded $\mathcal{S}$ -module.

(i) If $\mathcal{M}$ satisfies (TF), then $\widetilde{\mathcal{M}}$ is of finite type.

(ii) If $\mathcal{M}$ satisfies (TF), then $\widetilde{\mathcal{M}} = 0$ if and only if $\mathcal{M}$ satisfies (TN).

Proof. Since the questions are local on Y , we can reduce to the case where Y is affine of ring A , $\mathcal{S} = \widetilde{S}$, where S is a graded A -algebra such that the ideal S_+ is of finite type, and $\mathcal{M} = \widetilde{M}$, where M is a graded S -module; the proposition then follows from (2.7.3). $\square$

Theorem (3.4.4). — Let Y be a prescheme, and $\mathcal{S}$ a quasi-coherent graded $\mathcal{O}_Y$ -algebra generated by $\mathcal{S}_1$, where $\mathcal{S}_1$ is assumed to be of finite type; let $X = \text{Proj}(\mathcal{S})$. For every quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$, the canonical homomorphism β (3.3.4) is an isomorphism.

Proof. Note first of all that β is defined, by (3.4.1). To see that β is an isomorphism, we can reduce to the case where Y is affine of ring A , $\mathcal{S} = \widetilde{S}$, where S is a graded A -algebra generated by S_1 , and S_1 is an A -module of finite type. It then suffices to apply (2.7.5). $\square$

Corollary (3.4.5). — Under the hypotheses of (3.4.4), every quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$ is isomorphic to an $\mathcal{O}_X$ -module of the form $\widetilde{\mathcal{M}}$, where $\mathcal{M}$ is a quasi-coherent graded $\mathcal{S}$ -module. If, further, $\mathcal{F}$ is of finite type, and if we assume that Y is a quasi-compact scheme, or that the underlying space of Y is Noetherian, then we can take $\mathcal{M}$ to be of finite type.

[II:60]

Proof. The first claim follows immediately from (3.4.4) by taking $\mathcal{M} = \Gamma_*(\mathcal{F})$. To prove the second, it suffices to show that $\mathcal{M}$ is the inductive limit of its graded sub- $\mathcal{S}$ -modules of finite type $\mathcal{N}_\lambda$: indeed, it will follow from this that $\widetilde{\mathcal{M}}$ is the inductive limit of the $\widetilde{\mathcal{N}_\lambda}$ (3.2.4), and so $\mathcal{F}$ is the

inductive limit of the $\beta(\widetilde{\mathcal{N}}_\lambda)$; since X is quasi-compact ((3.4.1) and (I, 6.3.1)) and $\mathcal{F}$ is of finite type, $\mathcal{F}$ will necessarily be equal to one of the $\beta(\widetilde{\mathcal{N}}_\lambda)$ (0, 5.2.3).

To define the $\mathcal{N}_\lambda$ having $\mathcal{M}$ as their inductive limit, it suffices to consider, for each $n \in \mathbf{Z}$, the quasi-coherent $\mathcal{O}_Y$ -module $\mathcal{M}_n$, which is the inductive limit of its sub- $\mathcal{O}_Y$ -modules $\mathcal{M}_n^{(\mu_n)}$ of finite type, by the hypotheses on Y (I, 9.4.9); it is immediate that $\mathcal{P}_{\mu_n} = \mathcal{S} \cdot \mathcal{M}_n^{(\mu_n)}$ is a graded $\mathcal{S}$ -module of finite type, and we immediately see that taking $\mathcal{N}_\lambda$ to be the finite sums of the $\mathcal{S}$ -modules of the form $\mathcal{P}_{\mu_n}$ gives the desired objects. $\square$

Corollary (3.4.6). — Suppose that the hypotheses of (3.4.4) are satisfied, and further that the underlying space of Y is quasi-compact; let $\mathcal{F}$ be a quasi-coherent $\mathcal{O}_X$ -module of finite type. Then there exists n_0 such that, for all $n \geq n_0$, the canonical homomorphism $\sigma : p^*(p_*(\mathcal{F}(n))) \rightarrow \mathcal{F}(n)$ (0, 4.4.3) is surjective.

Proof. For all $y \in Y$, let U be an affine open neighbourhood of y in Y . There exists an integer $n_0(U)$ such that, for all $n \geq n_0(U)$, $\mathcal{F}(n)|_{p^{-1}(U)}$ is generated by a finite number of its sections over $p^{-1}(U)$ (2.7.9); but these sections are the canonical images of sections of $p^*(p_*(\mathcal{F}(n)))$ over $p^{-1}(U)$ ((0, 3.7.1) and (0, 4.4.3)), so $\mathcal{F}(n)|_{p^{-1}(U)}$ is equal to the canonical image of $p^*(p_*(\mathcal{F}(n)))|_{p^{-1}(U)}$. Finally, since Y is quasi-compact, there exists a finite cover of Y by affine opens U_i , and taking n_0 to be the largest of the $n_0(U_i)$ finishes the proof. $\square$

Remarks (3.4.7). — If $p = (\psi, \theta) : X \rightarrow Y$ is a morphism of ringed spaces, and $\mathcal{F}$ an $\mathcal{O}_X$ -module, the fact that the canonical homomorphism $\sigma : p^*(p_*(\mathcal{F})) \rightarrow \mathcal{F}$ is surjective can be explained in the following way (0, 4.4.1): for all $x \in X$, and every section s of $\mathcal{F}$ over an open neighbourhood V of x , there exists an open neighbourhood U of $p(x)$ in Y , a finite number of sections t_i (for $1 \leq i \leq m$) of $\mathcal{F}$ over $p^{-1}(U)$, a neighbourhood $W \subset V \cap p^{-1}(U)$ of x , and sections a_i (for $1 \leq i \leq m$) of $\mathcal{O}_X$ over W such that

$$s|_W = \sum_{i=1}^m a_i \cdot (t_i|_W).$$

If Y is an affine scheme and $p_*(\mathcal{F})$ is quasi-coherent, this condition is equivalent to $\mathcal{F}$ being generated by its sections over X (0, 5.5.1): indeed, if $Y = \text{Spec}(A)$, we can suppose that $U = D(f)$ with $f \in A$; then there exists an integer $n > 0$ and sections s_i of $\mathcal{F}$ over X such that t_i is the restriction to $p^{-1}(U)$ of $s_i g^n$, where $g = \theta(f)$ (by applying (I, 1.4.1) to $p_*(\mathcal{F})$); since g is invertible over $p^{-1}(U)$, we thus have

$$s|_W = \sum_i b_i \cdot (s_i|_W)$$

where $b_i = a_i(g|_W)^{-n}$, whence the claim. If Y is affine, then the corollary (3.4.6) recovers (2.7.9).

[II:61] We thus conclude that, when Y is an arbitrary prescheme, the following three conditions, for a quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$ such that $p_*(\mathcal{F})$ is quasi-coherent, are equivalent:

- a) The canonical homomorphism $\sigma : p^*(p_*(\mathcal{F})) \rightarrow \mathcal{F}$ is surjective.
- b) There exists a quasi-coherent $\mathcal{O}_Y$ -module $\mathcal{G}$ and a surjective homomorphism $p^*(\mathcal{G}) \rightarrow \mathcal{F}$.
- c) For every affine open U of Y , $\mathcal{F}|_{p^{-1}(U)}$ is generated by its sections over $p^{-1}(U)$.

Indeed, we have just proven the equivalence between a) and c). It is also clear that a) implies b), since $p_*(\mathcal{F})$ is quasi-coherent by hypothesis. Conversely, every homomorphism $u : p^*(\mathcal{G}) \rightarrow \mathcal{F}$ factors as $p^*(\mathcal{G}) \rightarrow p^*(p_*(\mathcal{F})) \xrightarrow{\sigma} \mathcal{F}$ (0, 3.5.4.4), so if u is surjective then so too is σ , which proves that b) implies a).

Corollary (3.4.8). — Suppose that the hypotheses of (3.4.4) are satisfied, and suppose further that Y is a quasi-compact scheme, or that the underlying space of Y is Noetherian. Let $\mathcal{F}$ be a quasi-coherent $\mathcal{O}_X$ -module of finite type; then there exists an integer n_0 such that, for all $n \geq n_0$, $\mathcal{F}$ is isomorphic to a quotient of an $\mathcal{O}_X$ -module of the form $(p^*(\mathcal{G}))(-n)$, where $\mathcal{G}$ is a quasi-coherent $\mathcal{O}_Y$ -module of finite type (that depends on n).

Proof. Since the structure morphism $X \rightarrow Y$ is separated and of finite type, $p_*(\mathcal{F}(n))$ is quasi-coherent (I, 9.2.2, b)), and thus the inductive limit of its quasi-coherent sub- $\mathcal{O}_Y$ -modules of finite type, by the hypotheses on Y (I, 9.4.9). We thus deduce, by (3.4.6), (0, 4.3.2), and (0, 5.2.3), that $\mathcal{F}(n)$ is the canonical image of an $\mathcal{O}_X$ -module of the form $p^*(\mathcal{G})$, where $\mathcal{G}$ is a quasi-coherent sub- $\mathcal{O}_Y$ -module of $p_*(\mathcal{F}(n))$ of finite type; the corollary then follows from (3.2.5.2) and (3.2.7.1). $\square$

3.5. Functorial behaviour

(3.5.1). Let Y be a prescheme, and $\mathcal{S}$ and $\mathcal{S}'$ quasi-coherent positively graded $\mathcal{O}_Y$ -algebras; let $X = \text{Proj}(\mathcal{S})$ and $X' = \text{Proj}(\mathcal{S}')$, and let p and p' be the structure morphisms from X and X' to Y . Let $\phi : \mathcal{S}' \rightarrow \mathcal{S}$ be an $\mathcal{O}_Y$ -homomorphism of graded algebras. For every affine open U of Y , let $S_U = \Gamma(U, \mathcal{S})$ and $S'_U = \Gamma(U, \mathcal{S}')$; the homomorphism ϕ defines a homomorphism $\phi_U : S'_U \rightarrow S_U$ of graded A_U -algebras, where $A_U = \Gamma(U, \mathcal{O}_Y)$. Corresponding to it are an open subset $G(\phi_U)$ of $p^{-1}(U)$ and a morphism $\Phi_U : G(\phi_U) \rightarrow p'^{-1}(U)$ (2.8.1). Furthermore, if $V \subset U$ is an affine open, then the diagram

$$(3.5.1.1) \quad \begin{array}{ccc} S'_U & \xrightarrow{\phi_U} & S_U \\ \downarrow & & \downarrow \\ S'_V & \xrightarrow{\phi_V} & S_V \end{array}$$

commutes, and we immediately see, by definition (2.8.1), that we have

$$G(\phi_V) = G(\phi_U) \cap p^{-1}(V)$$

and that Φ_V is the restriction to $G(\phi_V)$ of Φ_U . We have thus defined an open subset $G(\phi)$ of X such that $G(\phi) \cap p^{-1}(U) = G(\phi_U)$ for every affine open $U \subset Y$, and an affine Y -morphism $\Phi : G(\phi) \rightarrow X'$, which we say is *associated* to ϕ , and which we denote by $\text{Proj}(\phi)$. [II-62] If, for all $y \in Y$, there is an affine neighbourhood U of y such that the $\Gamma(U, \mathcal{O}_Y)$ -module $\Gamma(U, \mathcal{S}_+)$ is generated by $\phi(\Gamma(U, \mathcal{S}'_+))$, then $G(\phi_U) = p^{-1}(U)$, and so $G(\phi) = X$.

Proposition (3.5.2). —

- (i) If $\mathcal{M}$ is a quasi-coherent graded $\mathcal{S}$ -module, then there exists a canonical functorial isomorphism from the $\mathcal{O}_{X'}$ -module $(\mathcal{M}|_{G(\phi)})^\sim$ to the $\mathcal{O}_{X'}$ -module $\Phi_*(\widetilde{\mathcal{M}}|G(\phi))$.
- (ii) If $\mathcal{M}'$ is a quasi-coherent graded $\mathcal{S}'$ -module, then there exists a canonical functorial homomorphism v from the $(\mathcal{O}_X|G(\phi))$ -module $\Phi^*(\widetilde{\mathcal{M}'})$ to the $(\mathcal{O}_X|G(\phi))$ -module $(\mathcal{M}' \otimes_{\mathcal{S}'} \mathcal{S})^\sim|G(\phi)$. If $\mathcal{S}'$ is generated by $\mathcal{S}'_+$, then v is an isomorphism.

Proof. The homomorphisms in question are indeed already defined if Y is affine ((2.8.7) and (2.8.8)), and in the general case it suffices to check that they are compatible with the restriction of an affine open of Y to a smaller open, which follows immediately from the commutativity of (3.5.1.1). $\square$

In particular, for all $n \in \mathbb{Z}$, we have a canonical homomorphism

$$(3.5.2.1) \quad \Phi^*(\mathcal{O}_{X'}(n)) \longrightarrow \mathcal{O}_X(n)|G(\phi).$$

Proposition (3.5.3). — Let Y and Y' be preschemes, $\psi : Y' \rightarrow Y$ a morphism, and $\mathcal{S}$ a quasi-coherent graded $\mathcal{O}_Y$ -algebra; set $\mathcal{S}' = \psi^*(\mathcal{S})$. Then the Y' -scheme $X' = \text{Proj}(\mathcal{S}')$ is canonically identified with $\text{Proj}(\mathcal{S}) \times_Y Y'$. Furthermore, if $\mathcal{M}$ is a quasi-coherent graded $\mathcal{S}$ -module, then the $\mathcal{O}_{X'}$ -module $(\psi^*(\mathcal{M}))^\sim$ can be identified with $\widetilde{\mathcal{M}} \otimes_Y \mathcal{O}_{Y'}$.

Proof. Note first of all that $\psi^*(\mathcal{S})$ and $\psi^*(\mathcal{M})$ are quasi-coherent $\mathcal{O}_{Y'}$ -modules, as are their homogeneous components (0, 5.1.4). Let U be an affine open of Y , $U' \subset \psi^{-1}(U)$ an affine open of Y' , and A and A' the rings of U and U' , respectively; then $\mathcal{S}|U = \widetilde{S}$, where S is a graded A -algebra, and $\mathcal{S}'|U'$ can be identified with $(S \otimes_A A')^\sim$ (I, 1.6.5); the first claim then follows from (2.8.10) and (I, 3.2.6.2), since we immediately see that the projection $\text{Proj}(\mathcal{S}'|U') \rightarrow \text{Proj}(\mathcal{S}|U)$ defined by the above identification is compatible with the restriction operations on U and U' , and thus indeed defines a morphism $\text{Proj}(\mathcal{S}') \rightarrow \text{Proj}(\mathcal{S})$. Now let

$$\begin{aligned} q &: \text{Proj}(\mathcal{S}) \longrightarrow Y \\ q' &: \text{Proj}(\mathcal{S}') \longrightarrow Y' \end{aligned}$$

be the structure morphisms; $q'^{-1}(U')$ can then be identified with $q^{-1}(U) \times_U U'$, and the two sheaves $(\psi^*(\mathcal{M}))^\sim|q'^{-1}(U')$ and $(\widetilde{\mathcal{M}} \otimes_Y \mathcal{O}_{Y'})|q'^{-1}(U')$ are then both canonically identified with $(M \otimes_A A')^\sim$, where we set $M = \Gamma(U, \mathcal{M})$, by (2.8.10) and (I, 1.6.5); whence the second claim, since

we can again immediately see the compatibility of the above identifications with the restriction operations. $\square$

Corollary (3.5.4). — *With the notation of (3.5.3), $\mathcal{O}_{X'}(n)$ is canonically identified with $\mathcal{O}_X(n) \otimes_Y \mathcal{O}_{Y'}$ for all $n \in \mathbf{Z}$ (where $X = \text{Proj}(\mathcal{S})$).*

Proof. Indeed, with the notation of (3.5.3), it is clear that $\psi^*(\mathcal{S}(n)) = \mathcal{S}'(n)$ for all $n \in \mathbf{Z}$. $\square$

(3.5.5). Keeping the above notation, denote by Ψ the canonical projection $X' \rightarrow X$, and set $\mathcal{M}' = \psi^*(\mathcal{M})$; we further suppose that $\mathcal{S}$ is generated by $\mathcal{S}_1$, and that X is of finite type over Y ; ^[II:63] it then follows that $\mathcal{S}'$ is generated by $\mathcal{S}'_1$ (as can be seen by reducing to the case where Y and Y' are affine), and that X' is of finite type over Y' (I, 6.3.4). Let $\mathcal{F}$ be an $\mathcal{O}_X$ -module, and set $\mathcal{F}' = \Psi^*(\mathcal{F})$; it then follows from (3.5.4) and (0, 4.3.3) that $\mathcal{F}'(n) = \Psi^*(\mathcal{F}(n))$ for all $n \in \mathbf{Z}$. We further define a canonical ψ -homomorphism $\theta_n : q_*(\mathcal{F}(n)) \rightarrow q'_*(\mathcal{F}'(n))$ in the following way: given the commutativity of the diagram

$$\begin{array}{ccc} X & \xleftarrow{\Psi} & X' \\ q \downarrow & & \downarrow q' \\ Y & \xleftarrow{\psi} & Y' \end{array}$$

it is enough to define a homomorphism $q_*(\mathcal{F}(n)) \rightarrow \psi_*(q'_*(\Psi^*(\mathcal{F}(n)))) = q_*(\Psi_*(\Psi^*(\mathcal{F}(n))))$, and it suffices to take the homomorphism $\theta_n = q_*(\rho_n)$, where ρ_n is the canonical homomorphism $\mathcal{F}(n) \rightarrow \Psi_*(\Psi^*(\mathcal{F}(n)))$ (0, 4.4.3). It is immediate that, for every affine open U of Y and every affine open U' of Y' such that $U' \subset \psi^{-1}(U)$, the homomorphism θ_n gives, on sections, the canonical homomorphism (0, 3.7.2) $\Gamma(q^{-1}(U), \mathcal{F}(n)) \rightarrow \Gamma(q'^{-1}(U'), \mathcal{F}'(n))$. The commutativity of (2.8.13.2) then shows that, if $\mathcal{F}$ is quasi-coherent, the diagram

$$\begin{array}{ccc} \mathcal{F} & \xrightarrow{\rho} & \mathcal{F}' \\ \beta_{\mathcal{F}} \uparrow & & \uparrow \beta_{\mathcal{F}'} \\ (\Gamma_*(\mathcal{F}))^\sim & \xrightarrow{\theta} & (\Gamma_*(\mathcal{F}'))^\sim \end{array}$$

commutes (the top horizontal arrow being the canonical Ψ -morphism $\mathcal{F} \rightarrow \Psi^*(\mathcal{F})$).

Similarly, the commutativity of (2.8.13.1) shows that the diagram

$$\begin{array}{ccc} \Gamma_*(\widetilde{\mathcal{M}}) & \xrightarrow{\theta} & \Gamma_*(\widetilde{\mathcal{M}'}) \\ \alpha_{\mathcal{M}} \uparrow & & \uparrow \alpha_{\mathcal{M}'} \\ \mathcal{M} & \xrightarrow{\psi} & \mathcal{M}' \end{array}$$

commutes (the bottom horizontal arrow being the canonical ψ -morphism $\mathcal{M} \rightarrow \psi^*(\mathcal{M})$).

(3.5.6). Now consider preschemes Y and Y' , a morphism $g : Y' \rightarrow Y$, a quasi-coherent graded $\mathcal{O}_Y$ -algebra $\mathcal{S}$, a quasi-coherent graded $\mathcal{O}_{Y'}$ -algebra $\mathcal{S}'$, and a g -morphism of graded algebras $u : \mathcal{S} \rightarrow \mathcal{S}'$, i.e. a $\mathcal{O}_Y$ -homomorphism of graded algebras $\mathcal{S} \rightarrow g_*(\mathcal{S}')$; we already know that this is equivalent to giving a $\mathcal{O}_{Y'}$ -homomorphism of graded algebras $u^\sharp : g^*(\mathcal{S}) \rightarrow \mathcal{S}'$. We thus canonically obtain from $u^\sharp$ a Y' -morphism $W = \text{Proj}(u^\sharp) : G(u^\sharp) \rightarrow \text{Proj}(g^*(\mathcal{S}))$, where $G(u^\sharp)$ is an open subset of $X' = \text{Proj}(\mathcal{S}')$ (3.5.1). We also know that $X'' = \text{Proj}(g^*(\mathcal{S}))$ is canonically identified with $X \times_Y Y'$, where $X = \text{Proj}(\mathcal{S})$ (3.5.3); ^[III:64] composing the first projection $p : X \times_Y Y' \rightarrow X$ with $\text{Proj}(u^\sharp)$, we thus obtain a morphism $v : G(u^\sharp) \rightarrow X$, which we denote by $\text{Proj}(u)$, and which is such that the diagram

$$\begin{array}{ccc} G(u^\sharp) & \xrightarrow{v} & X \\ \downarrow & & \downarrow \\ Y' & \xrightarrow{g} & Y \end{array}$$

commutes.

Furthermore, for every quasi-coherent graded $\mathcal{S}$ -module $\mathcal{M}$, we have a canonical v -morphism

$$(3.5.6.1) \quad v : \widetilde{\mathcal{M}} \longrightarrow (g^*(\mathcal{M}) \otimes_{g^*(\mathcal{S})} \mathcal{S}')^\sim | G(u^\sharp).$$

Indeed, $v^\sharp$ is given by composing the homomorphisms

$$v^*(\widetilde{\mathcal{M}}) = W^*(p^*(\widetilde{\mathcal{M}})) \longrightarrow W^*((g^*(\mathcal{M}))^\sim) \longrightarrow (g^*(\mathcal{M}) \otimes_{g^*(\mathcal{S})} \mathcal{S}')^\sim | G(u^\sharp)$$

where the first arrow comes from the isomorphism (3.5.3) and the second is the homomorphism (3.5.2, (ii)); if $\mathcal{S}$ is generated by $\mathcal{S}_1$, then it follows from (3.5.2) that $v^\sharp$ is an isomorphism.

As a particular case of (3.5.6.1), we have, for all $n \in \mathbf{Z}$, a canonical v -morphism

$$(3.5.6.2) \quad v : \mathcal{O}_X(n) \longrightarrow \mathcal{O}_{X'}(n) | G(u^\sharp).$$

3.6. Closed subschemes of a prescheme $\text{Proj}(\mathcal{S})$

(3.6.1). Let Y be a prescheme, and $\phi : \mathcal{S} \rightarrow \mathcal{S}'$ a degree 0 homomorphism of quasi-coherent graded $\mathcal{O}_Y$ -algebras. We say that ϕ is (TN)-surjective (resp. (TN)-injective, (TN)-bijective) if there exists n such that, for all $k \geq n$, $\phi_k : \mathcal{S}_k \rightarrow \mathcal{S}'_k$ is surjective (resp. injective, bijective). If this is the case, then we can reduce the study of the corresponding morphism $\Phi : \text{Proj}(\mathcal{S}') \rightarrow \text{Proj}(\mathcal{S})$ to the case where ϕ is surjective (resp. injective, bijective). We prove this as in (2.9.1) (which is the particular case where Y is affine) by using (3.1.8). Instead of saying that ϕ is (TN)-bijective, we also say that it is a (TN)-isomorphism.

Proposition (3.6.2). — Let Y be a prescheme, and $\mathcal{S}$ a quasi-coherent graded $\mathcal{O}_Y$ -algebra; set $X = \text{Proj}(\mathcal{S})$.

- (i) If $\phi : \mathcal{S} \rightarrow \mathcal{S}'$ is a (TN)-surjective homomorphism of graded $\mathcal{O}_Y$ -algebras, then the corresponding morphism $\Phi = \text{Proj}(\phi)$ (3.5.1) is defined on all of $\text{Proj}(\mathcal{S}')$ and is a closed immersion of $\text{Proj}(\mathcal{S}')$ into X . If $\mathcal{J}$ is the kernel of ϕ , then the closed subscheme of X associated to Φ is defined by the quasi-coherent sheaf of ideals $\widetilde{\mathcal{J}}$ in $\mathcal{O}_X$.
- (ii) Suppose further that $\mathcal{S}_0 = \mathcal{O}_Y$, that $\mathcal{S}$ is generated by $\mathcal{S}_1$, and that $\mathcal{S}_1$ is of finite type. Let X' be a closed subscheme of $X = \text{Proj}(\mathcal{S})$, defined by a quasi-coherent sheaf of ideals $\mathcal{I}$ in $\mathcal{O}_X$. [III:65] Let $\widetilde{\mathcal{J}}$ be the quasi-coherent graded sheaf of ideals of $\mathcal{S}$, given by the inverse image of $\Gamma_*(\mathcal{I})$ by the canonical homomorphism $\alpha : \mathcal{S} \rightarrow \Gamma_*(\mathcal{O}_X)$ (3.3.2), and set $\mathcal{S}' = \mathcal{S} | \widetilde{\mathcal{J}}$. Then X' is the subscheme associated (I, 4.2.1) to the closed immersion $\text{Proj}(\mathcal{S}') \rightarrow X$ corresponding to the canonical homomorphism $\mathcal{S} \rightarrow \mathcal{S}'$ of graded $\mathcal{O}_Y$ -algebras.

Proof.

- (i) We can assume that ϕ is surjective (3.6.1). Then, for every affine open U of Y , $\Gamma(U, \mathcal{S}) \rightarrow \Gamma(U, \mathcal{S}')$ is surjective (I, 1.3.9), so $G(\phi) = \text{Proj}(\mathcal{S}')$ by (3.5.1). We can then immediately reduce to proving the proposition in the case where Y is affine, and this follows from (2.9.2, (i)).
- (ii) We can reduce to proving that the homomorphism $\widetilde{\mathcal{J}} \rightarrow \mathcal{O}_X$ induced by the canonical injection $\mathcal{J} \rightarrow \mathcal{S}$ is an isomorphism from $\widetilde{\mathcal{J}}$ to $\mathcal{I}$; since the question is local on Y , we can take Y to be affine of ring A , which implies that $\mathcal{S} = \widetilde{S}$, where S is a graded A -algebra generated by S_1 , with S_1 of finite type over A . It then suffices to apply (2.9.2, (ii)).

□

Corollary (3.6.3). — Under the conditions of (3.6.2, (i)), suppose further that $\mathcal{S}$ is generated by $\mathcal{S}_1$. Then $\Phi^*(\mathcal{O}_{X'}(n))$ is canonically identified with $\mathcal{O}_{X'}(n)$ for all $n \in \mathbf{Z}$.

Proof. We have defined such a canonical isomorphism when Y is affine (2.9.3); in the general case, it suffices to show that the isomorphisms thus defined for each affine open U of Y are compatible with the passage from U to an affine open $U' \subset U$, which is immediate. □

Corollary (3.6.4). — Let Y be a prescheme, $\mathcal{S}$ a quasi-coherent graded $\mathcal{O}_Y$ -algebra generated by $\mathcal{S}_1$, $\mathcal{M}$ a quasi-coherent $\mathcal{O}_Y$ -module, u a surjective $\mathcal{O}_Y$ -homomorphism $\mathcal{M} \rightarrow \mathcal{S}_1$, and $\bar{u} : \mathbf{S}_{\mathcal{O}_Y}(\mathcal{M}) \rightarrow \mathcal{S}$ the homomorphism of graded $\mathcal{O}_Y$ -algebras that extends u (1.7.4). Then the morphism corresponding to $\bar{u}$ is a closed immersion of $\text{Proj}(\mathcal{S})$ into $\text{Proj}(\mathbf{S}_{\mathcal{O}_Y}(\mathcal{M}))$.

Proof. Indeed, $\bar{u}$ is surjective by hypothesis, and we apply (3.6.2, (i)). $\square$

3.7. Morphisms from a prescheme to a homogeneous spectrum

(3.7.1). Let $q : X \rightarrow Y$ be a morphism of preschemes, $\mathcal{L}$ an invertible $\mathcal{O}_X$ -module, and $\mathcal{S}$ a quasi-coherent positively graded $\mathcal{O}_Y$ -algebra; then $q^*(\mathcal{S})$ is a quasi-coherent positively graded $\mathcal{O}_X$ -algebra. Consider the quasi-coherent graded $\mathcal{O}_X$ -algebra $\mathcal{S}' = \bigoplus_{n \geq 0} \mathcal{L}^{\otimes n}$; suppose that we have an $\mathcal{O}_X$ -homomorphism of graded algebras

$$\psi : q^*(\mathcal{S}) \longrightarrow \mathcal{S}' = \bigoplus_{n \geq 0} \mathcal{L}^{\otimes n}$$

which is equivalent to having a q -morphism of graded algebras

$$\psi^b : \mathcal{S} \longrightarrow q_*(\mathcal{S}').$$

We know that $\text{Proj}(\mathcal{S}')$ is canonically identified with X ((3.1.7) and (3.1.8, (iii))); we thus canonically obtain from ψ an open subset $G(\psi)$ of X , and a Y -morphism

$$(3.7.1.1) \quad r_{\mathcal{L}, \psi} : G(\psi) \longrightarrow \text{Proj}(\mathcal{S}) = P$$

[II:66] which we call the morphism *associated to $\mathcal{L}$ and ψ* ; recall (3.5.6) that this morphism is obtained by composing the Y -morphism

$$\tau = \text{Proj}(\psi) : G(\psi) \longrightarrow \text{Proj}(q^*(\mathcal{S}'))$$

with the first projection $\pi : \text{Proj}(q^*(\mathcal{S}')) = P \times_Y X \rightarrow P$.

(3.7.2). We will explicitly describe $r = r_{\mathcal{L}, \psi}$ in the case where $Y = \text{Spec}(A)$ is affine, so that $\mathcal{S} = \tilde{S}$, where S is a positively graded A -algebra. Suppose first of all that $X = \text{Spec}(B)$ is also affine, and that we have $\mathcal{L} = \tilde{L}$, where L is a free B -module of rank 1. Then $q^*(\mathcal{S}) = (S \otimes_A B)^\sim$ (I, 1.6.5); if c is a generator of L , then $\psi_n : q^*(\mathcal{S}_n) \rightarrow \mathcal{L}^{\otimes n}$ corresponds to a homomorphism $w_n : s \otimes b \mapsto bv_n(s)c^{\otimes n}$ from $S_n \otimes_A B$ to $L^{\otimes n}$, where $v_n : S_n \rightarrow B$ is a homomorphism of A -modules, the v_n together constituting a homomorphism of algebras $S \rightarrow B$. Let $f \in S_d$ (where $d > 0$), and set $g = v_d(f)$; we have $\pi^{-1}(D_+(f)) = D_+(f \otimes 1)$ by (2.8.10) and the identification of $D_+(f)$ with $\text{Spec}(S_{(f)})$ (2.3.6); we also know, from (2.8.1.1) (taking into account the canonical identification of X with $\text{Proj}(\mathcal{S}')$), that

$$\tau^{-1}(D_+(f \otimes 1)) = D(g)$$

whence

$$(3.7.2.1) \quad r^{-1}(D_+(f)) = D(g).$$

Furthermore, the morphism $\tau = \text{Proj}(\psi)$, restricted to $D(g)$, corresponds to the homomorphism that sends $(s \otimes 1)/(f \otimes 1)^n$ (where $s \in S_{nd}$) to $v_{nd}(s)/g^n$ (2.8.1), and the projection π , restricted to $D_+(f \otimes 1)$, corresponds to the homomorphism $s/f^n \mapsto (s \otimes 1)/(f \otimes 1)^n$; we thus conclude that r , restricted to $D(g)$, corresponds to the homomorphism of A -algebras $\omega : S_{(f)} \rightarrow B_g$ such that $\omega(s/f^n) = v_{nd}(s)/g^n$ (where $s \in S_{nd}$ and $n > 0$). Passing to the case where X is arbitrary (but Y still affine), we thus obtain, taking (2.8.1) into account, the following:

Proposition (3.7.3). — *If $Y = \text{Spec}(A)$ is affine and $\mathcal{S} = \tilde{S}$, where S is a graded A -algebra, then, for all $f \in S_d = \Gamma(Y, \mathcal{S}_d)$, we have*

$$(3.7.3.1) \quad r_{\mathcal{L}, \psi}^{-1}(D_+(f)) = X_{\psi^b(f)}$$

(where $\psi^b(f) \in \Gamma(X, \mathcal{L}^{\otimes d})$) and the restriction $X_{\psi^b(f)} \rightarrow D_+(f) = \text{Spec}(S_{(f)})$ of $r_{\mathcal{L}, \psi}$ corresponds (I, 2.2.4) to the homomorphism of algebras

$$(3.7.3.2) \quad \psi_{(f)}^b : S_{(f)} \longrightarrow \Gamma(X_{\psi^b(f)}, \mathcal{O}_X)$$

such that, for $s \in S_{nd} = \Gamma(Y, \mathcal{S}_{nd})$,

$$(3.7.3.3) \quad \psi_{(f)}^b(s/f^n) = (\psi^b(s)|_{X_{\psi^b(f)}})(\psi^b(f)|_{X_{\psi^b(f)}})^{-n}.$$

We say that $r_{\mathcal{L},\psi}$ is *everywhere defined* if $G(\psi) = X$. For this to be the case, it is evidently necessary and sufficient that $G(\psi) \cap q^{-1}(U) = q^{-1}(U)$ for every affine open $U \subset Y$; in other words, the question is *local* on Y . If Y is affine, then $G(\psi)$ is the union of the $r^{-1}(D_+(f))$ over f homogeneous in S_+ (2.8.1); by (3.7.3.1), the $X_{\psi^b(f)}$ must then form a *cover* of X ; in other words:

Corollary (3.7.4). — *Under the hypotheses of (3.7.3), for $r_{\mathcal{L},\psi}$ to be everywhere defined, it is necessary and sufficient that, for every $x \in X$, there exist an integer $n > 0$ and a section s of $\mathcal{S}_n$ over Y such that, if we set $t = \psi^b(s) \in \Gamma(X, \mathcal{L}^{\otimes n})$, then $t(x) \neq 0$.*

[II:67] Note that this condition is always satisfied if ψ is (TN)-surjective.

Similarly, the question of whether $r_{\mathcal{L},\psi}$ is *dominant* is local on Y , and we have:

Corollary (3.7.5). — *Under the hypotheses of (3.7.3), for $r_{\mathcal{L},\psi}$ to be dominant, it is necessary and sufficient that, for every integer $n > 0$, every section $s \in S_n$ such that $\psi^b(s) \in \Gamma(X, \mathcal{L}^{\otimes n})$ is locally nilpotent is itself nilpotent.*

Proof. We have to show that $r_{\mathcal{L},\psi}^{-1}(D_+(s))$ is nonempty whenever $D_+(s)$ is nonempty, and the corollary thus follows from (3.7.3.1) and (2.3.7). $\square$

Proposition (3.7.6). — *Let $q : X \rightarrow Y$ be a morphism, $\mathcal{L}$ an invertible $\mathcal{O}_X$ -module, $\mathcal{S}$ and $\mathcal{S}'$ quasi-coherent graded $\mathcal{O}_Y$ -algebras, $u : \mathcal{S}' \rightarrow \mathcal{S}$ a homomorphism of graded algebras, and $\psi : q^*(\mathcal{S}) \rightarrow \bigoplus_{n \geq 0} \mathcal{L}^{\otimes n}$ a homomorphism of graded algebras; let $\psi' = \psi \circ q^*(u)$ be the composite homomorphism. If $r_{\mathcal{L},\psi'}$ is everywhere defined, then so too is $r_{\mathcal{L},\psi}$; if u is (TN)-surjective, then if $r_{\mathcal{L},\psi'}$ is dominant, so too is $r_{\mathcal{L},\psi}$; conversely, if u is (TN)-injective, then if $r_{\mathcal{L},\psi}$ is dominant, so too is $r_{\mathcal{L},\psi'}$.*

Proof. We know that $G(\psi') \subset G(\psi)$ (2.8.4), whence the first claim; if u is (TN)-surjective, then $\text{Proj}(u) : \text{Proj}(\mathcal{S}') \rightarrow \text{Proj}(\mathcal{S})$ is everywhere defined and a closed immersion; since $r_{\mathcal{L},\psi'}$ is the composition of $\text{Proj}(u)$ and the restriction to $G(\psi')$ of $r_{\mathcal{L},\psi}$, we thus conclude that, if $r_{\mathcal{L},\psi'}$ is dominant, then so too is $r_{\mathcal{L},\psi}$. Finally, if u is (TN)-injective, then we know that $\text{Proj}(u)$ is a dominant morphism from $G(u)$ to $\text{Proj}(\mathcal{S}')$ (2.8.3); since $G(\psi')$ is the inverse image of $G(u)$ under $r_{\mathcal{L},\psi}$, we see that, if $r_{\mathcal{L},\psi}$ is dominant, then so too is $r_{\mathcal{L},\psi'}$. $\square$

Proposition (3.7.7). — *Let Y be a quasi-compact prescheme, $q : X \rightarrow Y$ a quasi-compact morphism, $\mathcal{L}$ an invertible $\mathcal{O}_X$ -module, and $\mathcal{S}$ a quasi-coherent graded $\mathcal{O}_Y$ -algebra given by the filtered inductive limit of an inductive system $(\mathcal{S}^\lambda)$ of quasi-coherent graded $\mathcal{O}_Y$ -algebras. Let $\phi_\lambda : \mathcal{S}^\lambda \rightarrow \mathcal{S}$ be the canonical homomorphism, $\psi : q^*(\mathcal{S}) \rightarrow \bigoplus_{n \geq 0} \mathcal{L}^{\otimes n}$ a homomorphism of graded algebras, and set $\psi_\lambda = \psi \circ q^*(\phi_\lambda)$. For $r_{\mathcal{L},\psi}$ to be everywhere defined, it is necessary and sufficient that there exist some λ such that $r_{\mathcal{L},\psi_\lambda}$ is everywhere defined; $r_{\mathcal{L},\psi_\mu}$ is then everywhere defined for $\mu \geq \lambda$.*

Proof. The condition is sufficient by (3.7.6). Conversely, suppose that $r_{\mathcal{L},\psi}$ is everywhere defined; we can reduce to the case where Y is affine, since, if, for every affine open $U \subset Y$ there exists $\lambda(U)$ such that the restriction of $r_{\mathcal{L},\psi_{\lambda(U)}}$ to $q^{-1}(U)$ is everywhere defined, then it will suffice (Y being quasi-compact) to cover Y by a finite number of affine opens U_i , and to take $\lambda \geq \lambda(U_i)$ for all indices i , by (3.7.6). If Y is affine, then the hypothesis implies that, for all $x \in X$, there exists a section $s^{(x)}$ of S_n such that, if we set $t^{(x)} = \psi^b(s^{(x)})$, then $t^{(x)}(x) \neq 0$ (where $t^{(x)}$ is considered as a section of $\mathcal{L}^{\otimes n}$ over X), which implies that $t^{(x)}(z) \neq 0$ for any z in a neighbourhood $V(x)$ of x . Cover X by a finite number of $V(x_i)$, and let $s^{(i)}$ be the corresponding sections of S ; then there exists some λ such that the $s^{(i)}$ are all of the form $\phi_\lambda(s'_\lambda{}^{(i)})$, with $s'_\lambda{}^{(i)} \in S^\lambda$ for all i ; it then follows from (3.7.4) that $r_{\mathcal{L},\psi_\lambda}$ is everywhere defined. The final claim is a trivial consequence of (3.7.6). $\square$

Corollary (3.7.8). — *Under the hypotheses of (3.7.7), if the $r_{\mathcal{L},\psi_\lambda}$ are dominant, then so too is $r_{\mathcal{L},\psi}$; the converse is true if the ϕ_λ are all injective.*

[II:68]

Proof. The second claim is a particular case of (3.7.6); to show that $r_{\mathcal{L},\psi}$ is dominant, we can restrict to the case where Y is affine; if $s \in S$ is such that $\psi^b(s)$ is locally nilpotent, since we can write $s = \phi_\lambda(s_\lambda)$ for at least one λ , we conclude from the hypotheses and from (3.7.5) that s_λ is nilpotent, and thus so too is s , and the criteria of (3.7.5) then apply. $\square$

Remarks (3.7.9). —

- (i) With the notation of (3.7.1), and taking (3.2.10) into account, we have, for all $n \in \mathbb{Z}$, a canonical homomorphism

$$(3.7.9.1) \quad \theta : r_{\mathcal{L}, \psi}^*(\mathcal{O}_P(n)) \longrightarrow \mathcal{L}^{\otimes n}$$

defined in a general way in (3.5.6.2). We immediately see that, under the conditions of (3.7.3), the restriction of this homomorphism to $X_{\psi^b(f)}$ sends s/f^k (where $s \in S_{n+kd}$) to the element $(\psi^b(s)|X_{\psi^b(f)})(\psi^b(f)|X_{\psi^b(f)})^{-k}$.

- (ii) Let $\mathcal{F}$ be a quasi-coherent $\mathcal{O}_X$ -module, and suppose that q is quasi-compact and separated, so that, for all $n \geq 0$, $q_*(\mathcal{F} \otimes \mathcal{L}^{\otimes n})$ is a quasi-coherent $\mathcal{O}_Y$ -module (I, 9.2.2). Let $\mathcal{M}' = \bigoplus_{n \geq 0} \mathcal{F} \otimes \mathcal{L}^{\otimes n}$, which is a quasi-coherent graded $\mathcal{S}'$ -module, and consider its image $\mathcal{M} = q_*(\mathcal{M}') = \bigoplus_{n \geq 0} q_*(\mathcal{F} \otimes \mathcal{L}^{\otimes n})$ (which is a quasi-coherent $\mathcal{S}$ -module, via the homomorphism ψ^b). We are going to show the existence of a canonical homomorphism of $\mathcal{O}_X$ -modules

$$(3.7.9.2) \quad \zeta : r_{\mathcal{L}, \psi}^*(\widetilde{\mathcal{M}}) \longrightarrow \mathcal{F}|G(\psi).$$

Indeed, we have already defined (3.5.6.1) a canonical homomorphism

$$(3.7.9.3) \quad r_{\mathcal{L}, \psi}^*(\widetilde{\mathcal{M}}) \longrightarrow (q^*(\mathcal{M}) \otimes_{q^*(\mathcal{S})} \mathcal{S}')^\sim |G(\psi)$$

where the right-hand side is considered as a quasi-coherent sheaf on $\text{Proj}(\mathcal{S}')$. We also have a canonical homomorphism

$$(3.7.9.4) \quad q^*(q_*(\mathcal{M}')) \otimes_{q^*(\mathcal{S})} \mathcal{S}' \longrightarrow \mathcal{M}'$$

for any quasi-coherent graded $\mathcal{S}'$ -module $\mathcal{M}'$: for any open U of X , any section t' of $q^*(q_*(\mathcal{M}'_h))$ over U , and any section b' of $\mathcal{S}'_k$ over U , we send $t' \otimes b'$ to the section $b'\sigma(t')$ of $\mathcal{M}'_{h+k}$, where $\sigma(t')$ is the section of $\mathcal{M}'_h$ over U that corresponds canonically (0, 4.4.3) to t' . We thus obtain a canonical homomorphism

$$(3.7.9.5) \quad (q^*(q_*(\mathcal{M}')) \otimes_{q^*(\mathcal{S})} \mathcal{S}')^\sim |G(\psi) \longrightarrow \widetilde{\mathcal{M}}|G(\psi)$$

and, finally, since $\widetilde{\mathcal{M}}$ is canonically identified with $\mathcal{F}$ (3.2.9, (i)), we indeed obtain the desired canonical homomorphism.

Under the conditions of (3.7.3), the restriction of this homomorphism to $X_{\psi^b(f)}$ acts as follows: given a section t_{nd} of $\mathcal{F} \otimes \mathcal{L}^{\otimes nd}$ over X , if t'_{nd} is the section t_{nd} considered as a section of $q_*(\mathcal{F} \otimes \mathcal{L}^{\otimes nd})$ over Y , then it sends the element t'_{nd}/f^n to the section $(t_{nd}|X_{\psi^b(f)})(\psi^b(f)|X_{\psi^b(f)})^{-n}$ of $\mathcal{F}$ over $X_{\psi^b(f)}$.

3.8. Criteria for immersion into a homogeneous spectrum [II:69]

(3.8.1). With the notation of (3.7.1), the question of whether $r_{\mathcal{L}, \psi}$ is an immersion (resp. an open immersion, a closed immersion) is clearly *local on Y* .

Proposition (3.8.2). — *Under the hypotheses of (3.7.3), for $r_{\mathcal{L}, \psi}$ to be everywhere defined and an immersion, it is necessary and sufficient that there exists a family of sections $s_\alpha \in S_{n_\alpha}$ (where $n_\alpha > 0$) such that, if we set $f_\alpha = \psi^b(s_\alpha)$, the following conditions are satisfied:*

- (i) *The X_{f_α} form a cover of X .*
- (ii) *The X_{f_α} are affine opens.*
- (iii) *For every α and every $t \in \Gamma(X_{f_\alpha}, \mathcal{O}_X)$, there exists an integer $m > 0$ and some $s \in S_{mn_\alpha}$ such that $t = (\psi^b(s)|X_{f_\alpha})(f_\alpha|X_{f_\alpha})^{-m}$.*

For $r_{\mathcal{L}, \psi}$ to be everywhere defined and an open immersion, it is necessary and sufficient that there exists a family (s_α) satisfying conditions (i), (ii), and (iii) above, as well as:

- (iv) *For all $n > 0$ and every $s \in S_{nn_\alpha}$ such that $\psi^b(s)|X_{f_\alpha} = 0$, there exists an integer $k > 0$ such that $s_\alpha^k = 0$.*

For $r_{\mathcal{L}, \psi}$ to be everywhere defined and a closed immersion, it is necessary and sufficient that there exists a family (s_α) satisfying conditions (i), (ii), and (iii) above, as well as:

(v) The $D_+(s_\alpha)$ form a cover of $P = \text{Proj}(S)$.

Proof. For r to be an immersion (resp. a closed immersion), it is necessary and sufficient that there exists a cover of $r(G(\psi))$ (resp. of P) by the $D_+(s_\alpha)$, such that, if we set $V_\alpha = r^{-1}(D_+(s_\alpha))$, then the restriction of r to V_α is a closed immersion of V_α into $D_+(s_\alpha)$ (I, 4.2.4). Condition (i) says that r is everywhere defined and also that the $D_+(s_\alpha)$ cover $r(X)$, by (3.7.3.1); since $D_+(s_\alpha)$ is affine, conditions (ii) and (iii) say that the restriction of r to X_{f_α} is a closed immersion into $D_+(s_\alpha)$, by (I, 4.2.3); finally, since conditions (iii) and (iv) say that $\psi^b_{(s_\alpha)}$ is bijective (using the notation of (3.7.3.2)), conditions (ii), (iii), and (iv) say that the restriction of r to X_{f_α} is an isomorphism to $D_+(s_\alpha)$ for all α , and so conditions (i), (ii), (iii), and (iv) together say that r is an open immersion. $\square$

Corollary (3.8.3). — *Under the hypotheses of (3.7.6), if $r_{\mathcal{L},\psi'}$ is everywhere defined and an immersion, then so too is $r_{\mathcal{L},\psi}$. If we further suppose that u is (TN)-surjective, then if $r_{\mathcal{L},\psi'}$ is an open (resp. closed) immersion, so too is $r_{\mathcal{L},\psi}$.*

Proof. By (3.8.2), there exists a family $s'_\alpha \in S'_{n_\alpha}$ such that, if we set $f_\alpha = \psi^b(s'_\alpha)$, conditions (i), (ii), and (iii) are satisfied. But if we set $s_\alpha = u(s'_\alpha)$, then we also have that $f_\alpha = \psi^b(s_\alpha)$, and if $t = (\psi^b(s')|X_{f_\alpha})(f_\alpha|X_{f_\alpha})^{-m}$, then also $t = (\psi^b(s)|X_{f_\alpha})(f_\alpha|X_{f_\alpha})^{-m}$ by setting $s = u(s')$, whence the first claim. The second claim follows immediately from the fact that $\text{Proj}(u)$ is a closed immersion. $\square$

Proposition (3.8.4). — *Suppose that the hypotheses of (3.7.7) are satisfied, and further that $q : X \rightarrow Y$ is a morphism of finite type. Then, for $r_{\mathcal{L},\psi}$ to be everywhere defined and an immersion, it is necessary and sufficient that there exist some λ such that $r_{\mathcal{L},\psi_\lambda}$ is everywhere defined and an immersion; in this case, $r_{\mathcal{L},\psi_\mu}$ is also everywhere defined and an immersion for all $\mu \geq \lambda$.*

[II:70]

Proof. Taking (3.8.3) into account, it suffices to show that, if $r_{\mathcal{L},\psi}$ is everywhere defined and an immersion, then so too is $r_{\mathcal{L},\psi_\lambda}$ for at least one λ . By the same argument as in (3.7.7), using the quasi-compactness of Y , we first reduce to the case where $Y = \text{Spec}(A)$ is affine. Since X is quasi-compact, (3.8.2) then shows the existence of a finite family $(s_i \in S_{n_i})$ of elements of S satisfying conditions (i), (ii), and (iii). The morphism $X_{f_i} \rightarrow Y$ (where $f_i = \psi^b(s_i)$) is of finite type: it is a morphism of affine schemes, and is thus quasi-compact (I, 6.6.1), and also locally of finite type, since q is of finite type (I, 6.3.2), and the conclusion then follows from (I, 6.6.3). The ring B_i of X_{f_i} is thus an A -algebra of finite type (I, 6.3.3); let (t_{ij}) be a family of generators of this algebra. By hypothesis, there exist elements $s'_{ij} \in S_{m_{ij}n_i}$ such that

$$t_{ij} = (\psi^b(s'_{ij})|X_{f_i})(\psi^b(s_i)|X_{f_i})^{-m_{ij}}.$$

Also by hypothesis, there exists some λ and elements $s_{i\lambda} \in S_{n_i}^\lambda$ and $s'_{ij\lambda} \in S_{m_{ij}n_i}^\lambda$ whose images under ϕ_λ are s_i and s'_{ij} , respectively; it is clear that $r_{\mathcal{L},\psi_\lambda}$ satisfies conditions (i), (ii), and (iii) of (3.8.2). $\square$

Proposition (3.8.5). — *Let Y be a quasi-compact scheme, or a prescheme whose underlying space is Noetherian, $q : X \rightarrow Y$ a morphism of finite type, $\mathcal{L}$ an invertible $\mathcal{O}_X$ -module, $\mathcal{S}$ a quasi-coherent graded $\mathcal{O}_Y$ -algebra, and $\psi : q^*(\mathcal{S}) \rightarrow \bigoplus_{n \geq 0} \mathcal{L}^{\otimes n}$ a homomorphism of graded algebras. For $r_{\mathcal{L},\psi}$ to be everywhere defined and an immersion, it is necessary and sufficient that there exist an integer $n > 0$ and a quasi-coherent sub- $\mathcal{O}_Y$ -module of finite type $\mathcal{E}$ of $\mathcal{S}_n$ such that*

- a) the homomorphism $\psi_n \circ q^*(j_n) : q^*(\mathcal{E}) \rightarrow \mathcal{L}^{\otimes n}$ (where $j_n : \mathcal{E} \rightarrow \mathcal{S}$ is the canonical injection) is surjective; and
- b) if we denote by $\mathcal{S}'$ the (graded) sub- $\mathcal{O}_Y$ -algebra of $\mathcal{S}$ generated by $\mathcal{E}$, and by ψ' the homomorphism $\psi \circ q^*(j')$ (where j' is the injection of $\mathcal{S}'$ into $\mathcal{S}$), $r_{\mathcal{L},\psi'}$ is everywhere defined and an immersion.

If this is the case, then every quasi-coherent sub- $\mathcal{O}_Y$ -module $\mathcal{E}'$ of $\mathcal{S}_n$ that contains $\mathcal{E}$ also possesses the same property, as does the image of $\otimes^k \mathcal{E}$ in $\mathcal{S}_{kn}$ for all $k > 0$.

Proof. The fact that the condition is sufficient, and the last two claims, are particular cases of (3.8.3), taking into account the canonical isomorphism between $\text{Proj}(\mathcal{S})$ and $\text{Proj}(\mathcal{S}^{(k)})$ (3.1.8).

Let (U_i) be a finite cover of Y by affine opens, and set $A_i = A(U_i)$. Since $q^{-1}(U_i)$ is quasi-compact, the hypothesis that $r_{\mathcal{L},\psi}$ be everywhere defined and an immersion implies, along with (3.8.2), the existence of a finite family $(s_{ij} \in S_{n_{ij}}^{(i)})$ of elements of $S^{(i)} = \Gamma(U_i, \mathcal{S})$ satisfying conditions (i), (ii), and (iii). We see, as in the proof of (3.8.4), that the morphism $X_{f_{ij}} \rightarrow U_i$ (where $f_{ij} = \psi^b(s_{ij})$) is of finite type, and that the ring B_{ij} of $X_{f_{ij}}$ is thus an A_i -algebra of finite type, having a system of generators of the form $(\psi^b(t_{ijk})|X_{f_{ij}})(f_{ij}|X_{f_{ij}})^{-m_{ijk}}$, where $t_{ijk} \in S_{m_{ijk}n_{ij}}^{(i)}$. Let n be a common multiple of the $m_{ijk}n_{ij}$; replacing (for each (i, j, k)) s_{ij} by some power s_{ij}^ρ such that $\rho m_{ijk}n_{ij} = n$, and multiplying t_{ijk} by $s_{ij}^{\rho-m_{ijk}}$, we can assume that, for each i , the s_{ij} and t_{ijk} belong to $S_n^{(i)}$ and that $m_{ijk} = 1$. Let E_i be the sub- A_i -module of $S^{(i)}$ generated by these elements (for fixed i). Then there exists a coherent sub- $\mathcal{O}_Y$ -module $\mathcal{E}_i$ of $\mathcal{S}_n$ of finite type such that $\mathcal{E}_i|_{U_i} = (E_i)^\sim$ (I, 9.4.7). [II:71] It is clear that the sub- $\mathcal{O}_Y$ -module $\mathcal{E}$ of $\mathcal{S}_n$ given by the sum of the $\mathcal{E}_i$ is the desired object (since each section f_{ij} is such that, for all $x \in X_{f_{ij}}$, there exists an affine neighbourhood $V \subset X_{f_{ij}}$ of x such that $f_{ij}|_V$ is a basis for $\Gamma(V, \mathcal{L}^{\otimes n})$). $\square$

§4. PROJECTIVE BUNDLES; AMPLE SHEAVES

4.1. Definition of projective bundles

Definition (4.1.1). — Let Y be a prescheme, $\mathcal{E}$ a quasi-coherent $\mathcal{O}_Y$ -module, and $\mathbf{S}_{\mathcal{O}_Y}(\mathcal{E})$ the symmetric $\mathcal{O}_Y$ -algebra of $\mathcal{E}$ (1.7.4), which is quasi-coherent (1.7.7). We define the *projective bundle on Y defined by $\mathcal{E}$* , denoted $\mathbf{P}(\mathcal{E})$, to be the Y -scheme $P = \text{Proj}(\mathbf{S}_{\mathcal{O}_Y}(\mathcal{E}))$. The $\mathcal{O}_P$ -module $\mathcal{O}_P(1)$ is called the *fundamental sheaf on P* .

When Y is affine of ring A , then we have $\mathcal{E} = \tilde{E}$ for some A -module E , and we then write $\mathbf{P}(E)$ instead of $\mathbf{P}(\tilde{E})$.

When we take $\mathcal{E} = \mathcal{O}_Y^n$, we write $\mathbf{P}_Y^{n-1}$ instead of $\mathbf{P}(\mathcal{E})$; if, further, Y is affine of ring A , then we also write $\mathbf{P}_A^{n-1}$ instead of $\mathbf{P}_Y^{n-1}$. Since $\mathbf{S}_{\mathcal{O}_Y}(\mathcal{O}_Y)$ is canonically identified with $\mathcal{O}_Y[T]$ (1.7.4), $\mathbf{P}_Y^0$ is canonically identified with Y (3.1.7); Example (2.4.3) is then exactly $\mathbf{P}_K^1$.

(4.1.2). Let $\mathcal{E}$ and $\mathcal{F}$ be quasi-coherent $\mathcal{O}_Y$ -modules; let $u : \mathcal{E} \rightarrow \mathcal{F}$ be an $\mathcal{O}_Y$ -homomorphism; there is a corresponding canonical homomorphism $\mathbf{S}(u) : \mathbf{S}_{\mathcal{O}_Y}(\mathcal{E}) \rightarrow \mathbf{S}_{\mathcal{O}_Y}(\mathcal{F})$ of graded $\mathcal{O}_Y$ -algebras (1.7.4). If u is *surjective*, then so too is $\mathbf{S}(u)$, and thus (3.6.2, (i)) $\text{Proj}(\mathbf{S}(u))$ is a *closed immersion* $\mathbf{P}(\mathcal{F}) \rightarrow \mathbf{P}(\mathcal{E})$, which we denote by $\mathbf{P}(u)$. We can thus say that $\mathbf{P}(\mathcal{E})$ is a *contravariant functor* in $\mathcal{E}$, with the condition that we only consider *surjective* morphisms of quasi-coherent $\mathcal{O}_Y$ -modules.

Still supposing that u is surjective, and letting $P = \mathbf{P}(\mathcal{E})$, $Q = \mathbf{P}(\mathcal{F})$, and $j = \mathbf{P}(u)$, we have, up to isomorphism, that

$$(4.1.2.1) \quad j^*(\mathcal{O}_P(n)) = \mathcal{O}_Q(n) \quad \text{for all } n \in \mathbf{Z}$$

by (3.6.3).

(4.1.3). Now let $\psi : Y' \rightarrow Y$ be a morphism, and let $\mathcal{E}' = \psi^*(\mathcal{E})$; then $\mathbf{S}_{\mathcal{O}_{Y'}}(\mathcal{E}') = \psi^*(\mathbf{S}_{\mathcal{O}_Y}(\mathcal{E}))$ (1.7.5); thus (3.5.3)

$$(4.1.3.1) \quad \mathbf{P}(\psi^*(\mathcal{E})) = \mathbf{P}(\mathcal{E}) \times_Y Y'$$

up to canonical isomorphism; furthermore, we clearly have that

$$\psi^*((\mathbf{S}_{\mathcal{O}_Y}(\mathcal{E}))(n)) = (\mathbf{S}_{\mathcal{O}_{Y'}}(\mathcal{E}'))(n)$$

for all $n \in \mathbf{Z}$, whence, letting $P = \mathbf{P}(\mathcal{E})$ and $P' = \mathbf{P}(\mathcal{E}')$, we have (3.5.4), up to isomorphism, that

$$(4.1.3.2) \quad \mathcal{O}_{P'}(n) = \mathcal{O}_P(n) \otimes_Y \mathcal{O}_{Y'} \quad \text{for all } n \in \mathbf{Z}.$$

[II:72]

Proposition (4.1.4). — Let $\mathcal{L}$ be an invertible $\mathcal{O}_Y$ -module. For every quasi-coherent $\mathcal{O}_Y$ -module $\mathcal{E}$, there exists a canonical Y -isomorphism $i_{\mathcal{L}} : \mathbf{P}(\mathcal{E}) \xrightarrow{\sim} \mathbf{P}(\mathcal{E} \otimes \mathcal{L})$; furthermore, if we let $P = \mathbf{P}(\mathcal{E})$ and $Q = \mathbf{P}(\mathcal{E} \otimes \mathcal{L})$, then $i_{\mathcal{L}}^*(\mathcal{O}_Q(n))$ is canonically isomorphic to $\mathcal{O}_P(n) \otimes_Y \mathcal{L}^{\otimes n}$ for all $n \in \mathbf{Z}$.

Proof. First note that, if A is a ring, E an A -module, and L a free rank-one A -module, then we can canonically define a homomorphism of A -modules

$$\mathbf{S}_n(E \otimes L) \longrightarrow \mathbf{S}_n(E) \otimes L^{\otimes n}$$

by sending $(x_1 \otimes y_1) \dots (x_n \otimes y_n)$ to the element

$$(x_1 x_2 \dots x_n) \otimes (y_1 \otimes y_2 \otimes \dots \otimes y_n) \quad (x_i \in E, y_i \in L, \text{ for } 1 \leq i \leq n);$$

we can immediately see (by restricting to the case where $L = A$) that this homomorphism is in fact an isomorphism. We thus obtain a canonical isomorphism of graded A -algebras $\mathbf{S}_A(E \otimes L) \xrightarrow{\sim} \bigoplus_{n \geq 0} \mathbf{S}_n(E) \otimes L^{\otimes n}$. Returning to the setting of (4.1.4), the preceding remarks allow us to define a canonical isomorphism of graded $\mathcal{O}_Y$ -algebras

$$(4.1.4.1) \quad \mathbf{S}_{\mathcal{O}_Y}(\mathcal{E} \otimes_{\mathcal{O}_Y} \mathcal{L}) \xrightarrow{\sim} \bigoplus_{n \geq 0} \mathbf{S}_n(\mathcal{E}) \otimes_{\mathcal{O}_Y} \mathcal{L}^{\otimes n}$$

by defining this isomorphism as an isomorphism of presheaves, and taking into account (1.7.4), (I, 1.3.9), and (I, 1.3.12). The proposition then follows from (3.1.8, (iii)) and (3.2.10). $\square$

(4.1.5). With the hypotheses of (4.1.1), let $P = \mathbf{P}(\mathcal{E})$, and denote by p the structure morphism $P \rightarrow Y$. Since, by definition, $\mathcal{E} = (\mathbf{S}_{\mathcal{O}_Y}(\mathcal{E}))_1$, we have a canonical homomorphism $\alpha_1 : \mathcal{E} \rightarrow p_*(\mathcal{O}_P(1))$ (3.3.2.2), and thus (0, 4.4.3) also a canonical homomorphism

$$(4.1.5.1) \quad \alpha_1^\sharp : p^*(\mathcal{E}) \longrightarrow \mathcal{O}_P(1).$$

Proposition (4.1.6). — *The canonical homomorphism (4.1.5.1) is surjective.*

Proof. We have seen, in (3.3.2), that $\alpha_1^\sharp$ corresponds functorially to the canonical homomorphism $\mathcal{E} \otimes_{\mathcal{O}_Y} \mathbf{S}_{\mathcal{O}_Y}(\mathcal{E}) \rightarrow (\mathbf{S}_{\mathcal{O}_Y}(\mathcal{E}))(1)$; since, by definition, $\mathcal{E}$ generates $\mathbf{S}_{\mathcal{O}_Y}(\mathcal{E})$, this homomorphism is surjective, whence the conclusion, by (3.2.4). $\square$

4.2. Morphisms from a prescheme to a projective bundle

(4.2.1). Keeping the notation of (4.1.5), let X be a Y -prescheme, $q : X \rightarrow Y$ the structure morphism, and let $r : X \rightarrow P$ be a Y -morphism such that the following diagram commutes:

$$\begin{array}{ccc} P & \xleftarrow{r} & X \\ p \downarrow & & \searrow q \\ Y & & \end{array}$$

[II:73]

Since the functor r^* is right exact (0, 4.3.1), we obtain, from the surjective homomorphism in (4.1.5.1), a surjective homomorphism

$$r^*(\alpha_1^\sharp) : r^*(p^*(\mathcal{E})) \longrightarrow r^*(\mathcal{O}_P(1)).$$

But $r^*(p^*(\mathcal{E})) = q^*(\mathcal{E})$, and $r^*(\mathcal{O}_P(1))$ is locally isomorphic to $r^*(\mathcal{O}_P) = \mathcal{O}_X$; in other words, it is an invertible $\mathcal{O}_X$ -module $\mathcal{L}_r$. Thus, from r , we have defined a canonical surjective $\mathcal{O}_X$ -homomorphism

$$(4.2.1.1) \quad \phi_r : q^*(\mathcal{E}) \longrightarrow \mathcal{L}_r.$$

When $Y = \text{Spec}(A)$ is affine and $\mathcal{E} = \tilde{E}$, we can describe this homomorphism more explicitly as follows. given $f \in E$, it follows from (2.6.3) that

$$(4.2.1.2) \quad r^{-1}(D_+(f)) = X_{\phi_r^\flat(f)}.$$

Now let V be an affine open subset of X contained in $r^{-1}(D_+(f))$, and let B be its ring, which is an A -algebra; let $S = \mathbf{S}_A(E)$; the restriction of r to V corresponds to an A -homomorphism $\omega : S_{(f)} \rightarrow B$, and we have $q^*(\mathcal{E})|_V = (E \otimes_A B)^\sim$ and $\mathcal{L}_r|_V = \tilde{L}_r$, whence $L_r = (S(1))_{(f)} \otimes_{S_{(f)}} B_{[\omega]}$ (I, 1.6.5). The restriction of ϕ_r to $q^*(\mathcal{E})|_V$ corresponds to the B -homomorphism $u : E \otimes_A B \rightarrow L_r$, which sends $x \otimes 1$ to $(x/1) \otimes f = (f/1) \otimes \omega(x/f)$. The canonical extension of ϕ_r to a homomorphism of $\mathcal{O}_X$ -algebras

$$\psi_r : q^*(\mathbf{S}(\mathcal{E})) = \mathbf{S}(q^*(\mathcal{E})) \longrightarrow \mathbf{S}(\mathcal{L}_r) = \bigoplus_{n \geq 0} \mathcal{L}_r^{\otimes n}$$

is therefore such that its restriction to $q^*(\mathbf{S}_n(\mathcal{E}))|V$ corresponds to the homomorphism $\mathbf{S}_n(E \otimes_A B) = \mathbf{S}_n(E) \otimes_A B \rightarrow L_r^{\otimes n}$ that sends $s \otimes 1$ to $(f/1)^{\otimes n} \otimes \omega(s/f^n)$.

(4.2.2). Conversely, suppose that we are given a morphism $q : X \rightarrow Y$, an invertible $\mathcal{O}_X$ -module $\mathcal{L}$, and a quasi-coherent $\mathcal{O}_Y$ -module $\mathcal{E}$; every homomorphism $\phi : q^*(\mathcal{E}) \rightarrow \mathcal{L}$ canonically determines a homomorphism of quasi-coherent $\mathcal{O}_X$ -algebras

$$\psi : \mathbf{S}(q^*(\mathcal{E})) = q^*(\mathbf{S}(\mathcal{E})) \longrightarrow \bigoplus_{n \geq 0} \mathcal{L}^{\otimes n}$$

and thus (3.7.1) a Y -morphism $r_{\mathcal{L}, \psi} : G(\psi) \rightarrow \text{Proj}(\mathbf{S}(\mathcal{E})) = \mathbf{P}(\mathcal{E})$, which we denote by $r_{\mathcal{L}, \phi}$. If ϕ is *surjective*, then so too is ψ , and thus (3.7.4) $r_{\mathcal{L}, \phi}$ is *everywhere defined*. Furthermore, with the notation of (4.2.1) and (4.2.2):

Proposition (4.2.3). — *Given a morphism $q : X \rightarrow Y$ and a quasi-coherent $\mathcal{O}_Y$ -module $\mathcal{E}$, the assignments $r \mapsto (\mathcal{L}_r, \phi_r)$ and $(\mathcal{L}, \phi) \mapsto r_{\mathcal{L}, \phi}$ give a bijective correspondence between the set of Y -morphisms $r : X \rightarrow \mathbf{P}(\mathcal{E})$ and the set of equivalence classes of pairs $(\mathcal{L}, \phi)$ consisting of an invertible $\mathcal{O}_X$ -module $\mathcal{L}$ and a surjective homomorphism $\phi : q^*(\mathcal{E}) \rightarrow \mathcal{L}$, where two pairs $(\mathcal{L}, \phi)$ and $(\mathcal{L}', \phi')$ are equivalent if there exists an $\mathcal{O}_X$ -isomorphism $\tau : \mathcal{L} \xrightarrow{\sim} \mathcal{L}'$ such that $\phi' = \tau \circ \phi$.*

Proof. Start first with a Y -morphism $r : X \rightarrow \mathbf{P}(\mathcal{E})$, and construct $\mathcal{L}_r$ and ϕ_r (4.2.1), and let $r' = r_{\mathcal{L}_r, \phi_r}$; it follows immediately from (4.2.1) and (3.7.2) that the morphisms r and r' are identical (by taking the generator of $\mathcal{L}_r$ to be the element $(f/1) \otimes 1$ to define the homomorphisms v_n of (3.7.2)). Conversely, take a pair $(\mathcal{L}, \phi)$ and construct $r'' = r_{\mathcal{L}, \phi}$, and then $\mathcal{L}_{r''}$ and $\phi_{r''}$; we will show that there exists a canonical isomorphism $\tau : \mathcal{L}_{r''} \xrightarrow{\sim} \mathcal{L}$ such that $\phi = \tau \circ \phi_{r''}$; to define it, we can restrict to the case where $Y = \text{Spec}(A)$ and $X = \text{Spec}(B)$ are affine, and (with the notation of (4.2.1) and (3.7.2)) associate to each element $(x/1) \otimes 1$ of $L_{r''}$ (where $x \in E$) the element $v_1(x)c$ of L . We immediately see that τ does not depend on the chosen generator c of L ; since v_1 is surjective by hypothesis, to prove that τ is an isomorphism it suffices to show that, if $x/1 = 0$ in $(S(1))_{(f)}$, then $v_1(x)/1 = 0$ in B_g ; but the first equality implies that $f^n x = 0$ in $\mathbf{S}_{n+1}(E)$ for some n , and this implies that $v_{n+1}(f^n x) = g^n v_1(x) = 0$ in B , whence the conclusion. Finally, it is immediate that, for any two equivalent pairs $(\mathcal{L}, \phi)$ and $(\mathcal{L}', \phi')$, we have $r_{\mathcal{L}, \phi} = r_{\mathcal{L}', \phi'}$. $\square$

In particular, for $X = Y$:

Theorem (4.2.4). — *The set of Y -sections of $\mathbf{P}(\mathcal{E})$ is in canonical bijective correspondence with the set of quasi-coherent sub- $\mathcal{O}_Y$ -modules $\mathcal{F}$ of $\mathcal{E}$ such that $\mathcal{E}/\mathcal{F}$ is invertible.*

We note that this property corresponds to the classical definition of “projective space” as the set of hyperplanes of a vector space (the classical case corresponding to $Y = \text{Spec}(K)$, where K is a field, and $\mathcal{E} = \tilde{E}$, where E is a finite-dimensional K -vector space; the $\mathcal{F}$ having the property described in (4.2.4) then correspond to the hyperplanes of E , and we already know that the Y -sections of $\mathbf{P}(\mathcal{E})$ are then the K -rational points of $\mathbf{P}(\mathcal{E})$ (I, 3.4.5)).

Remark (4.2.5). — Since there is a canonical bijective correspondence between Y -morphisms from X to P and their graph morphisms, X -sections of $P \times_Y X$ (I, 3.3.14), we see that, conversely, (4.2.3) can be deduced from (4.2.4). Denote by $\text{Hyp}_Y(X, \mathcal{E})$ the set of quasi-coherent sub- $\mathcal{O}_X$ -modules $\mathcal{F}$ of $\mathcal{E} \otimes_Y \mathcal{O}_X = q^*(\mathcal{E})$ such that $q^*(\mathcal{E})/\mathcal{F}$ is an invertible $\mathcal{O}_X$ -module. If $g : X' \rightarrow X$ is a Y -morphism, then the right exactness of g^* gives

$$g^*(q^*(\mathcal{E})/\mathcal{F}) = g^*q^*(\mathcal{E})/g^*(\mathcal{F}),$$

so the latter sheaf is invertible. Thus $X \mapsto \text{Hyp}_Y(X, \mathcal{E})$ is a *contravariant functor* on the category of Y -preschemes. We can therefore interpret theorem (4.2.4) as defining a *canonical isomorphism* between the functors $X \mapsto \text{Hom}_Y(X, \mathbf{P}(\mathcal{E}))$ and $X \mapsto \text{Hyp}_Y(X, \mathcal{E})$, both contravariant in X on the category of Y -preschemes. This also gives a characterisation of the projective bundle $P = \mathbf{P}(\mathcal{E})$ by the following *universal property*, which is much closer to the geometric intuition than the constructions from §2-3: for every morphism $q : X \rightarrow Y$ and every invertible $\mathcal{O}_X$ -module $\mathcal{L}$ that is a quotient of $\mathcal{E} \otimes_Y \mathcal{O}_X$, there exists a unique Y -morphism $r : X \rightarrow P$ such that $\mathcal{L} = r^*(\mathcal{O}_P(1))$.

We will see later that we can similarly define, amongst other things, “Grassmannian” schemes.

Corollary (4.2.6). — Suppose that every invertible $\mathcal{O}_Y$ -module is trivial (I, 2.4.8). Let V be the group $\text{Hom}_{\mathcal{O}_Y}(\mathcal{E}, \mathcal{O}_Y)$, considered as a module over the ring $A = \Gamma(Y, \mathcal{O}_Y)$, and let V^* be the subset of V consisting of surjective homomorphisms. Then the set of Y -sections of $\mathbf{P}(\mathcal{E})$ is canonically identified with V^*/A^* , where A^* is the group of units of A .

[II:75] In particular:

- (1) The corollary (4.2.6) applies whenever Y is a *local scheme* (I, 2.4.8). Let Y be an arbitrary prescheme, y a point of Y , and $Y' = \text{Spec}(k(y))$; then the fibre $p^{-1}(y)$ of $\mathbf{P}(\mathcal{E})$ can, by (4.1.3.1), be identified with $\mathbf{P}(\mathcal{E}^y)$, where $\mathcal{E}^y = \mathcal{E}_y \otimes_{\mathcal{O}_y} k(y) = \mathcal{E}_y / \mathfrak{m}_y \mathcal{E}_y$ is considered as a vector space over $k(y)$. More generally, if K is an extension of $k(y)$, then $p^{-1}(y) \otimes_{k(y)} K$ can be identified with $\mathbf{P}(\mathcal{E}^y \otimes_{k(y)} K)$. The corollary (4.2.6) then shows that the set of *geometric points of $\mathbf{P}(\mathcal{E})$ with values in the extension K of $k(y)$* (I, 3.4.5), which we can also call the *rational geometric fibre over K of $\mathbf{P}(\mathcal{E})$ over the point y* , can be identified with the *projective space associated to the dual of the K -vector space $\mathcal{E}^y \otimes_{k(y)} K$* .
- (2) Suppose that Y is affine of ring A , and, further, that every invertible $\mathcal{O}_Y$ -module is trivial; further, take $\mathcal{E} = \mathcal{O}_Y^n$; then, in (4.2.6), V can be identified with A^n (I, 1.3.8), and V^* with the set of systems $(f_i)_{1 \leq i \leq n}$ of elements of A that generate the ideal A ; any two such systems define the same Y -section of $\mathbf{P}_Y^{n-1} = \mathbf{P}_A^{n-1}$, or, in other words, *the same point of $\mathbf{P}_A^{n-1}$ with values in A* , if and only if one of them can be obtained from the other by multiplication by an invertible element of A .

These properties justify the terminology “projective bundle” for $\mathbf{P}(\mathcal{E})$. We note that the definition of “projective space” obtained in this way is in fact *dual* to the classical definition; this is imposed upon us by the necessity of being able to define $\mathbf{P}(\mathcal{E})$ for *arbitrary* quasi-coherent $\mathcal{O}_Y$ -modules $\mathcal{E}$, and not just locally free ones.

Remark (4.2.7). — We will see, in Chapter V, that, if Y is connected and locally Noetherian, and if $\mathcal{E}$ is locally free, then, letting $P = \mathbf{P}(\mathcal{E})$, every invertible $\mathcal{O}_P$ -module is isomorphic to an $\mathcal{O}_P$ -module of the form $\mathcal{L}' \otimes_{\mathcal{O}_Y} \mathcal{O}_P(m)$, with $\mathcal{L}'$ some invertible $\mathcal{O}_Y$ -module, well defined up to isomorphism, and m some well defined integer. In other words, $H^1(P, \mathcal{O}_P^*)$ is isomorphic to $\mathbf{Z} \times H^1(Y, \mathcal{O}_Y^*)$ (0, 5.4.7). We will also see ((III, 2.1.14), taking (0, 5.4.10) into account) that $p_*(\mathcal{L}^{\otimes m}) = 0$ if $m < 0$, and $p_*(\mathcal{L}^{\otimes m})$ is isomorphic to $\mathcal{L}' \otimes_{\mathcal{O}_Y} (\mathbf{S}_{\mathcal{O}_Y}(\mathcal{E}))_m$ if $m \geq 0$. If $\mathcal{F}$ is a quasi-coherent $\mathcal{O}_Y$ -module, then every Y -morphism $\mathbf{P}(\mathcal{E}) \rightarrow \mathbf{P}(\mathcal{F})$ is determined by the data of an invertible $\mathcal{O}_Y$ -module, an integer $m \geq 0$, and an $\mathcal{O}_Y$ -homomorphism $\psi : \mathcal{F} \rightarrow \mathcal{L}' \otimes_{\mathcal{O}_Y} (\mathbf{S}_{\mathcal{O}_Y}(\mathcal{E}))_m$ such that the corresponding homomorphism $\psi^\sharp$ of $\mathcal{O}_{\mathbf{P}(\mathcal{F})}$ -modules is surjective. We will also see that, if the Y -morphism in question is an isomorphism, then $m = 1$ and $\mathcal{F}$ is isomorphic to $\mathcal{E} \otimes_{\mathcal{O}_Y} \mathcal{L}'$ (the converse of (4.1.4)). This will allow us to determine the sheaf of germs of automorphisms of $\mathbf{P}(\mathcal{E})$ as the quotient of the sheaf of groups $\mathcal{A}ut(\mathcal{E})$ (which is locally isomorphic to $\mathbf{GL}(n, \mathcal{O}_Y)$ if $\mathcal{E}$ is of rank n) by $\mathcal{O}_Y^*$.

(4.2.8). Keeping the notation of (4.2.1), let $u : X' \rightarrow X$ be a morphism; if the Y -morphism $r : X \rightarrow P$ corresponds to the homomorphism $\phi : q^*(\mathcal{E}) \rightarrow \mathcal{L}$, then the Y -morphism $r \circ u$ corresponds to $u^*(\phi) : u^*(q^*(\mathcal{E})) \rightarrow u^*(\mathcal{L})$, as follows immediately from the definitions.

(4.2.9). Let $\mathcal{E}$ and $\mathcal{F}$ be quasi-coherent $\mathcal{O}_Y$ -modules, $v : \mathcal{E} \rightarrow \mathcal{F}$ a surjective homomorphism, and $j = \mathbf{P}(v)$ the corresponding closed immersion $\mathbf{P}(\mathcal{F}) \rightarrow \mathbf{P}(\mathcal{E})$ (4.1.2). If the Y -morphism $r : X \rightarrow \mathbf{P}(\mathcal{F})$ corresponds to the homomorphism $\phi : q^*(\mathcal{F}) \rightarrow \mathcal{L}$, then the [II:76] Y -morphism $j \circ r$ corresponds to $q^*(\mathcal{E}) \xrightarrow{q^*(v)} q^*(\mathcal{F}) \xrightarrow{\phi} \mathcal{L}$; this again follows from the definition given in (4.2.1).

(4.2.10). Let $\psi : Y' \rightarrow Y$ be a morphism, and let $\mathcal{E}' = \psi^*(\mathcal{E})$. If the Y -morphism $r : X \rightarrow P$ corresponds to the homomorphism $\phi : q^*(\mathcal{E}) \rightarrow \mathcal{L}$, then the Y' -morphism

$$r_{(Y')} : X_{(Y')} \longrightarrow P' = \mathbf{P}(\mathcal{E}')$$

corresponds to $\phi_{(Y')} : q_{(Y')}^*(\mathcal{E}') = q^*(\mathcal{E}) \otimes_Y \mathcal{O}_{Y'} \rightarrow \mathcal{L} \otimes_Y \mathcal{O}_{Y'}$. Indeed, by (4.1.3.1), we have the commutative diagram

$$\begin{array}{ccccc} Y' & \xleftarrow{p_{(Y')}} & P' = P_{(Y')} & \xleftarrow{r_{(Y')}} & X_{(Y')} \\ \downarrow & & \downarrow u & & \downarrow v \\ Y & \xleftarrow{p} & P & \xleftarrow{r} & X \end{array}$$

From (4.1.3.2), we have

$$(r_{(Y')})^*(\mathcal{O}_{P'}(1)) = (r_{(Y')})^*(u^*(\mathcal{O}_P(1))) = v^*(r^*(\mathcal{O}_P(1))) = v^*(\mathcal{L}) = \mathcal{L} \otimes_Y \mathcal{O}_{Y'};$$

we also know that $u^*(\alpha_1^\sharp)$ is exactly the canonical homomorphism $\alpha_1^\sharp : (p_{(Y')})^*(\mathcal{E}') \rightarrow \mathcal{O}_{P'}(1)$; we can see this by explicitly calculating the canonical homomorphisms $\alpha_1^\sharp$ to P and P' as in (4.1.6). Whence our claim.

4.3. The Segre morphism

(4.3.1). Let Y be a prescheme, and $\mathcal{E}$ and $\mathcal{F}$ quasi-coherent $\mathcal{O}_Y$ -modules; let $P_1 = \mathbf{P}(\mathcal{E})$ and $P_2 = \mathbf{P}(\mathcal{F})$, and denote the structure morphisms by $p_1 : P_1 \rightarrow Y$ and $p_2 : P_2 \rightarrow Y$. Let $Q = P_1 \times_Y P_2$, and let $q_1 : Q \rightarrow P_1$ and $q_2 : Q \rightarrow P_2$ be the canonical projections; then the $\mathcal{O}_Q$ -module $\mathcal{L} = \mathcal{O}_{P_1}(1) \otimes_Y \mathcal{O}_{P_2}(1) = q_1^*(\mathcal{O}_{P_1}(1)) \otimes_{\mathcal{O}_Q} q_2^*(\mathcal{O}_{P_2}(1))$ is invertible, since it is the tensor product of two invertible $\mathcal{O}_Q$ -modules (0, 5.4.4). Also, if $r = p_1 \circ q_1 = p_2 \circ q_2$ is the structure morphism $Q \rightarrow Y$, then $r^*(\mathcal{E} \otimes_{\mathcal{O}_Y} \mathcal{F}) = q_1^*(p_1^*(\mathcal{E})) \otimes_{\mathcal{O}_Q} q_2^*(p_2^*(\mathcal{F}))$ (0, 4.3.3); the canonical surjective homomorphisms (4.1.5.1) $p_1^*(\mathcal{E}) \rightarrow \mathcal{O}_{P_1}(1)$ and $p_2^*(\mathcal{F}) \rightarrow \mathcal{O}_{P_2}(1)$ thus give, by taking the tensor product, a canonical homomorphism

$$(4.3.1.1) \quad s : r^*(\mathcal{E} \otimes_{\mathcal{O}_Y} \mathcal{F}) \longrightarrow \mathcal{L}$$

which is evidently surjective; from this we obtain (4.2.2) a canonical morphism, called the *Segre morphism*:

$$(4.3.1.2) \quad \varsigma : \mathbf{P}(\mathcal{E}) \times_Y \mathbf{P}(\mathcal{F}) \longrightarrow \mathbf{P}(\mathcal{E} \otimes_{\mathcal{O}_Y} \mathcal{F}).$$

We can study the morphism ς more explicitly in the case where $Y = \text{Spec}(A)$ is affine, and $\mathcal{E} = \tilde{E}$ and $\mathcal{F} = \tilde{F}$, where E and F are A -modules, whence $\mathcal{E} \otimes_{\mathcal{O}_Y} \mathcal{F} = (E \otimes_A F)^\sim$ (I, 1.3.12); let $R = \mathbf{S}_A(E)$, $S = \mathbf{S}_A(F)$, and $T = \mathbf{S}_A(E \otimes_A F)$; let $f \in E$ and $g \in F$, and consider the affine open

$$D_+(f) \times_Y D_+(g) = \text{Spec}(B)$$

[II:77] of Q , where $B = R_{(f)} \otimes_A S_{(g)}$; the restriction of $\mathcal{L}$ to this affine open is $\tilde{L}$, where

$$L = (R(1))_{(f)} \otimes_A (S(1))_{(g)}$$

and the element $c = (f/1) \otimes (g/1)$ is a generator of L considered as a free B -module (2.5.7). The homomorphism (4.3.1.1) corresponds to the homomorphism

$$(x \otimes y) \otimes b \longmapsto b((x/1) \otimes (y/1))$$

from $(E \otimes_A F) \otimes_A B$ to L . With the notation of (3.7.2), we thus have that $v_1(x \otimes y) = (x/f) \otimes (y/g)$; the restriction of ς to $D_+(f) \times_Y D_+(g)$ is a morphism from this affine scheme to $D_+(f \otimes g)$, corresponding to the ring homomorphism $\omega : T_{(f \otimes g)} \rightarrow R_{(f)} \otimes_A S_{(g)}$ defined by

$$(4.3.1.3) \quad \omega((x \otimes y)/(f \otimes g)) = (x/f) \otimes (y/g)$$

for $x \in E$ and $y \in F$.

(4.3.2). It follows from (4.2.3) that we have a canonical isomorphism

$$(4.3.2.1) \quad \tau : \varsigma^*(\mathcal{O}_P(1)) \xrightarrow{\sim} \mathcal{O}_{P_1}(1) \otimes_Y \mathcal{O}_{P_2}(1)$$

where we let $P = \mathbf{P}(\mathcal{E} \otimes_{\mathcal{O}_Y} \mathcal{F})$. We will show that, for $x \in \Gamma(Y, \mathcal{E})$ and $y \in \Gamma(Y, \mathcal{F})$, we have

$$(4.3.2.2) \quad \tau(\alpha_1(x \otimes y)) = \alpha_1(x) \otimes \alpha_1(y).$$

Indeed, we can restrict to the case where Y is affine, and we then have, with the notation of (4.3.1) and (2.6.2), that $\alpha_1^{f \otimes g}(x \otimes y) = (x \otimes y)/1$ in $(T(1))_{(f \otimes g)}$, that $\alpha_1^f(x) = x/1$ in $(R(1))_{(f)}$, and

that $\alpha_1^g(y) = y/1$ in $(S(1))_{(g)}$. The definition of τ given in (4.2.3) and the calculation of v_1 done in (4.3.1) then immediately prove the claim (4.3.2.2). From this we obtain the equation

$$(4.3.2.3) \quad \varsigma^{-1}(P_{x \otimes y}) = (P_1)_x \times_Y (P_2)_y$$

with the notation of (3.1.4). Indeed, taking (3.3.3) into account, the equation (4.3.2.2) (by restricting to the affine case, with the help of (I, 3.2.7) and (I, 3.2.3)) leaves us only to prove the following lemma:

Lemma (4.3.2.4). — *Let B and B' be A -algebras, and let $Y = \text{Spec}(A)$, $Z = \text{Spec}(B)$, and $Z' = \text{Spec}(B')$; then $D(t \otimes t') = D(t) \times_Y D(t')$ for any $t \in B$, $t' \in B'$.*

Proof. Indeed, if $p : Z \times_Y Z' \rightarrow Z$ and $p' : Z \times_Y Z' \rightarrow Z'$ are the canonical projections, then it follows from (I, 1.2.2.2) that $p^{-1}(D(t)) = D(t \otimes 1)$ and $p'^{-1}(D(t')) = D(1 \otimes t')$; the conclusion follows from (I, 3.2.7) and (I, 1.1.9.1), since $(t \otimes 1)(1 \otimes t') = t \otimes t'$. $\square$

Proposition (4.3.3). — *The Segre morphism is a closed immersion.*

Proof. Since the question is local on Y , we can restrict to the case where Y is affine. With the notation of (4.3.1) and (4.3.2), the $D_+(f \otimes g)$ then form a basis for the topology of P , since the $f \otimes g$ generate T when f runs over E and g runs over F . By (4.3.2.3), we also know that $\varsigma^{-1}(D_+(f \otimes g)) = D_+(f) \times_Y D_+(g)$. It thus suffices (I, 4.2.4) to prove that the restriction of ς to $D_+(f) \times_Y D_+(g)$ is a closed immersion into $D_+(f \otimes g)$. But, with the same notation, the equation (4.3.1.3) shows that ω is surjective, which completes the proof. $\square$

(4.3.4). The Segre morphism is functorial in $\mathcal{E}$ and $\mathcal{F}$, if we consider only [II:78] surjective homomorphisms of quasi-coherent $\mathcal{O}_Y$ -modules. Indeed, we must then show that, if $\mathcal{E} \rightarrow \mathcal{E}'$ is a surjective $\mathcal{O}_Y$ -homomorphism, then the diagram

$$\begin{array}{ccc} \mathbf{P}(\mathcal{E}') \times_Y \mathbf{P}(\mathcal{F}) & \xrightarrow{j \times 1} & \mathbf{P}(\mathcal{E}) \times_Y \mathbf{P}(\mathcal{F}) \\ \varsigma \downarrow & & \downarrow \varsigma \\ \mathbf{P}(\mathcal{E}' \otimes_{\mathcal{O}_Y} \mathcal{F}) & \longrightarrow & \mathbf{P}(\mathcal{E} \otimes_{\mathcal{O}_Y} \mathcal{F}) \end{array}$$

commutes, where j denotes the canonical closed immersion $\mathbf{P}(\mathcal{E}') \rightarrow \mathbf{P}(\mathcal{E})$. Let $P'_1 = \mathbf{P}(\mathcal{E}')$ and keep the notation from (4.3.1); then $j \times 1$ is a closed immersion (I, 4.3.1) and, up to isomorphism,

$$(j \times 1)^*(\mathcal{O}_{P_1}(1) \otimes \mathcal{O}_{P_2}(1)) = j^*(\mathcal{O}_{P_1}(1)) \otimes \mathcal{O}_{P_2}(1) = \mathcal{O}_{P'_1}(1) \otimes \mathcal{O}_{P_2}(1)$$

by (4.1.2.1) and (I, 9.1.5); our claim then follows from (4.2.8) and (4.2.9).

(4.3.5). With the notation of (4.3.1), let $\psi : Y' \rightarrow Y$ be a morphism, and let $\mathcal{E}' = \psi^*(\mathcal{E})$ and $\mathcal{F}' = \psi^*(\mathcal{F})$; then the Segre morphism $\mathbf{P}(\mathcal{E}') \times \mathbf{P}(\mathcal{F}') \rightarrow \mathbf{P}(\mathcal{E}' \otimes \mathcal{F}')$ can be identified with $\varsigma_{(Y')}$. Indeed, keeping the notation of (4.3.1), let $P'_1 = \mathbf{P}(\mathcal{E}')$ and $P'_2 = \mathbf{P}(\mathcal{F}')$; we know (4.1.3.1) that P'_i can be identified with $(P_i)_{(Y')}$ ($i = 1, 2$), and so the structure morphism $P'_1 \times P'_2 \rightarrow Y'$ can be identified with $r_{(Y')}$. Also $\mathcal{E}' \otimes \mathcal{F}'$ can be identified with $\psi^*(\mathcal{E} \otimes \mathcal{F})$, and so $\mathbf{P}(\mathcal{E}' \otimes \mathcal{F}')$ can be identified with $(\mathbf{P}(\mathcal{E} \otimes \mathcal{F}))_{(Y')}$ (4.1.3.1). Finally, $\mathcal{O}_{P'_1}(1) \otimes_{Y'} \mathcal{O}_{P'_2}(1) = \mathcal{L}'$ can be identified with $\mathcal{L} \otimes_Y \mathcal{O}_{Y'}$, by (4.1.3.1) and (I, 9.1.11). The canonical homomorphism $(r_{(Y')})^*(\mathcal{E}' \otimes \mathcal{F}') \rightarrow \mathcal{L}'$ can then be identified with $s_{(Y')}$, and our claim follows from (4.2.10).

Remark (4.3.6). — The prescheme given by the sum of $\mathbf{P}(\mathcal{E})$ and $\mathbf{P}(\mathcal{F})$ is likewise canonically isomorphic to a closed subprescheme of $\mathbf{P}(\mathcal{E} \oplus \mathcal{F})$. Indeed, the surjective homomorphisms $\mathcal{E} \oplus \mathcal{F} \rightarrow \mathcal{E}$ and $\mathcal{E} \oplus \mathcal{F} \rightarrow \mathcal{F}$ correspond to closed immersions $\mathbf{P}(\mathcal{E}) \rightarrow \mathbf{P}(\mathcal{E} \oplus \mathcal{F})$ and $\mathbf{P}(\mathcal{F}) \rightarrow \mathbf{P}(\mathcal{E} \oplus \mathcal{F})$; everything then reduces to showing that the underlying spaces of the closed subpreschemes of $\mathbf{P}(\mathcal{E} \oplus \mathcal{F})$ obtained in this way have empty intersection. Since the question is local on Y , we can adopt the notation of (4.3.1); but $\mathbf{S}_n(E)$ and $\mathbf{S}_n(F)$ can be identified with submodules of $\mathbf{S}_n(E \oplus F)$ with intersection consisting only of 0; if $\mathfrak{p}$ is a graded prime ideal of $\mathbf{S}(E)$ such that $\mathfrak{p} \cap \mathbf{S}_n(E) \neq \mathbf{S}_n(E)$ for any $n \geq 0$, then there exists a corresponding graded prime ideal of $\mathbf{S}(E \oplus F)$ whose intersection with $\mathbf{S}_n(E)$ is $\mathfrak{p} \cap \mathbf{S}_n(E)$, but which also contains $\mathbf{S}_+(F)$, as we immediately see; thus no point in $\text{Proj}(\mathbf{S}(E))$ can have the same image in $\text{Proj}(\mathbf{S}(E \oplus F))$ as any point in $\text{Proj}(\mathbf{S}(F))$.

4.4. Immersions into projective bundles; very ample sheaves

Proposition (4.4.1). — Let Y be a quasi-compact scheme, or a prescheme whose underlying space is Noetherian, $q : X \rightarrow Y$ a morphism of finite type, and $\mathcal{L}$ an invertible $\mathcal{O}_X$ -module.

- (i) Let $\mathcal{S}$ be a quasi-coherent positively-graded $\mathcal{O}_Y$ -algebra, and $\psi : q^*(\mathcal{S}) \rightarrow \bigoplus_{n \geq 0} \mathcal{L}^{\otimes n}$ a homomorphism of graded algebras. For $r_{\mathcal{L}, \psi}$ to be everywhere defined and an immersion, it is necessary and [II:79] sufficient for there to exist an integer $n \geq 0$ and a quasi-coherent sub- $\mathcal{O}_Y$ -module of finite type $\mathcal{E}$ of $\mathcal{S}_n$ such that the homomorphism $\psi' = \psi_n \circ q^*(j) : q^*(\mathcal{E}) \rightarrow \mathcal{L}^{\otimes n} = \mathcal{L}'$ (where j is the injection $\mathcal{E} \rightarrow \mathcal{S}_n$) is surjective and such that the morphism $r_{\mathcal{L}', \psi'} : X \rightarrow \mathbf{P}(\mathcal{E})$ is an immersion.
- (ii) Let $\mathcal{F}$ be a quasi-coherent $\mathcal{O}_Y$ -module, and $\phi : q^*(\mathcal{F}) \rightarrow \mathcal{L}$ a surjective homomorphism. For the morphism $r_{\mathcal{L}, \phi}$ to be an immersion $X \rightarrow \mathbf{P}(\mathcal{F})$, it is necessary and sufficient for there to exist a quasi-coherent sub- $\mathcal{O}_Y$ -module of finite type $\mathcal{E}$ of $\mathcal{F}$ such that the homomorphism $\phi' = \phi \circ q^*(j) : q^*(\mathcal{E}) \rightarrow \mathcal{L}$ (where j is the canonical injection $\mathcal{E} \rightarrow \mathcal{F}$) is surjective and such that the morphism $r_{\mathcal{L}, \phi'} : X \rightarrow \mathbf{P}(\mathcal{E})$ is an immersion.

Proof.

- (i) The fact that $r_{\mathcal{L}, \psi}$ is everywhere defined and is an immersion is equivalent, by (3.8.5), to the existence of some $n \geq 0$ and $\mathcal{E}$ such that, if $\mathcal{S}'$ is the subalgebra of $\mathcal{S}$ generated by $\mathcal{E}$, the homomorphism $q^*(\mathcal{E}) \rightarrow \mathcal{L}^{\otimes n}$ is surjective and the morphism $r_{\mathcal{L}', \psi'} : X \rightarrow \text{Proj}(\mathcal{S}')$ is everywhere defined and is an immersion. We already have a canonical surjective homomorphism $\mathbf{S}(\mathcal{E}) \rightarrow \mathcal{S}'$ to which there exists a corresponding closed immersion $\text{Proj}(\mathcal{S}') \rightarrow \mathbf{P}(\mathcal{E})$ (3.6.2); whence the conclusion.
- (ii) Since $\mathcal{F}$ is the inductive limit of its quasi-coherent submodules of finite type $\mathcal{E}_\lambda$ (I, 9.4.9), $\mathbf{S}(\mathcal{F})$ is the inductive limit of the $\mathbf{S}(\mathcal{E}_\lambda)$; the conclusion then follows from (3.8.4), by observing that we can take all the n_i in the proof of (3.8.4) to be equal to 1: indeed, supposing that Y is affine, if $r = r_{\mathcal{L}, \phi}$ is an immersion, then $r(X)$ is a quasi-compact subspace of $\mathbf{P}(\mathcal{F})$ that we can cover by finitely many open subsets of $\mathbf{P}(\mathcal{F})$ of the form $D_+(f)$, with $f \in F$, such that $D_+(f) \cap r(X)$ is closed.

□

Definition (4.4.2). — Let Y be a prescheme, and $q : X \rightarrow Y$ a morphism. We say that an invertible $\mathcal{O}_X$ -module $\mathcal{L}$ is *very ample relative to q* (or *relative to Y* , or simply *very ample*, if q is clear from the context) if there exists a quasi-coherent $\mathcal{O}_Y$ -module $\mathcal{E}$ and a Y -immersion $i : X \rightarrow P = \mathbf{P}(\mathcal{E})$ such that $\mathcal{L}$ is isomorphic to $i^*(\mathcal{O}_P(1))$.

It is equivalent (4.2.3) to say that there exists a quasi-coherent $\mathcal{O}_Y$ -module $\mathcal{E}$ and a surjective homomorphism $\phi : q^*(\mathcal{E}) \rightarrow \mathcal{L}$ such that $r_{\mathcal{L}, \phi} : X \rightarrow \mathbf{P}(\mathcal{E})$ is an immersion.

We note that the existence of a very ample (for Y) $\mathcal{O}_X$ -module implies that q is separated ((3.1.3) and (I, 5.5.1, (i) and (ii))).

Corollary (4.4.3). — Suppose that there exists a quasi-coherent graded $\mathcal{O}_Y$ -algebra $\mathcal{S}$, generated by $\mathcal{S}_1$, and a Y -immersion $i : X \rightarrow P = \text{Proj}(\mathcal{S})$ such that $\mathcal{L}$ is isomorphic to $i^*(\mathcal{O}_P(1))$; then $\mathcal{L}$ is very ample relative to q .

Proof. If $\mathcal{F} = \mathcal{S}_1$, then the canonical homomorphism $\mathbf{S}(\mathcal{F}) \rightarrow \mathcal{S}$ is surjective, and so, by composing the corresponding closed immersion $\text{Proj}(\mathcal{S}) \rightarrow \mathbf{P}(\mathcal{F})$ (3.6.2) with the immersion i , we obtain an immersion $j : X \rightarrow \mathbf{P}(\mathcal{F}) = P'$ such that $\mathcal{L}$ is isomorphic to $j^*(\mathcal{O}_{P'}(1))$ (3.6.3). □

Proposition (4.4.4). — Let $q : X \rightarrow Y$ be a quasi-compact morphism, and $\mathcal{L}$ an invertible $\mathcal{O}_X$ -module. Then the following properties are equivalent:

- (a) $\mathcal{L}$ is very ample relative to q .
- (b) $q_*(\mathcal{L})$ is quasi-coherent, the canonical homomorphism $\sigma : q^*(q_*(\mathcal{L})) \rightarrow \mathcal{L}$ is surjective, and the morphism $r_{\mathcal{L}, \sigma} : X \rightarrow \mathbf{P}(q_*(\mathcal{L}))$ is an immersion.

Proof. Since q is quasi-compact, we know that $q_*(\mathcal{L})$ is quasi-coherent if q is separated (I, 9.2.2).

[II:80] We know (3.4.7) that the existence of a surjective homomorphism $\phi : q^*(\mathcal{E}) \rightarrow \mathcal{L}$ (with $\mathcal{E}$ a quasi-coherent $\mathcal{O}_Y$ -module) implies that σ is surjective; furthermore, given the factorisation

$q^*(\mathcal{E}) \rightarrow q^*(q_*(\mathcal{L})) \xrightarrow{\sigma} \mathcal{L}$ of ϕ , there is a canonically corresponding factorisation

$$q^*(\mathbf{S}(\mathcal{E})) \longrightarrow q^*(\mathbf{S}(q_*(\mathcal{L}))) \longrightarrow \bigoplus_{n \geq 0} \mathcal{L}^{\otimes n}$$

and therefore, by (3.8.3), the hypothesis that $r_{\mathcal{L},\phi}$ is an immersion implies that so too is $j = r_{\mathcal{L},\sigma}$; furthermore, by (4.2.4), $\mathcal{L}$ is isomorphic to $j^*(\mathcal{O}_{P'}(1))$, where $P' = \mathbf{P}(q_*(\mathcal{L}))$. We thus see that (a) and (b) are equivalent. $\square$

Corollary (4.4.5). — Suppose that q is quasi-compact. For $\mathcal{L}$ to be very ample relative to Y , it is necessary and sufficient for there to exist an open cover (U_α) of Y such that $\mathcal{L}|_{q^{-1}(U_\alpha)}$ is very ample relative to U_α for every α .

Proof. Indeed, condition (b) of (4.4.4) is local on Y . $\square$

Proposition (4.4.6). — Let Y be a quasi-compact scheme, or a prescheme whose underlying space is Noetherian, $q : X \rightarrow Y$ a morphism of finite type, and $\mathcal{L}$ an invertible $\mathcal{O}_X$ -module. Then conditions (a) and (b) of (4.4.4) are equivalent to the following:

- (a') There exists a quasi-coherent $\mathcal{O}_Y$ -module $\mathcal{E}$ of finite type and a surjective homomorphism $\phi : q^*(\mathcal{E}) \rightarrow \mathcal{L}$ such that $r_{\mathcal{L},\phi}$ is an immersion.
- (b') There exists a coherent sub- $\mathcal{O}_Y$ -module $\mathcal{E}$ of $q_*(\mathcal{L})$ of finite type that has the properties stated in condition (a').

Proof. It is clear that (a') or (b') imply (a); also (a) implies (a'), by (4.4.1), and similarly (b) implies (b'). $\square$

Corollary (4.4.7). — Suppose that Y is a quasi-compact scheme, or a Noetherian prescheme. If $\mathcal{L}$ is very ample for q , then there exists a quasi-coherent graded $\mathcal{O}_Y$ -algebra $\mathcal{S}$ such that $\mathcal{S}_1$ is of finite type and generates $\mathcal{S}$, and also a dominant open Y -immersion $i : X \rightarrow P = \text{Proj}(\mathcal{S})$ such that $\mathcal{L}$ is isomorphic to $i^*(\mathcal{O}_P(1))$.

Proof. Indeed, condition (b') of (4.4.6) is satisfied; the structure morphism $p : \mathbf{P}(\mathcal{E}) = P' \rightarrow Y$ is then separated and of finite type (3.1.3), and so P' is a quasi-compact scheme (resp. a Noetherian prescheme) if Y is a quasi-compact scheme (resp. a Noetherian prescheme). Let Z be the closure (I, 9.5.11) of the subprescheme X' of P' associated to the immersion $j = r_{\mathcal{L},\phi}$ from X into P' ; it is clear that j factors as a dominant open immersion $i : X \rightarrow Z$ followed by the canonical injection $Z \rightarrow P'$. But Z can be identified with a prescheme $\text{Proj}(\mathcal{S})$, where $\mathcal{S}$ is a graded $\mathcal{O}_Y$ -algebra equal to the quotient of $\mathbf{S}(\mathcal{E})$ by a quasi-coherent graded sheaf of ideals (3.6.2), and it is clear that $\mathcal{S}_1$ is of finite type and generates $\mathcal{S}$; furthermore, $\mathcal{O}_Z(1)$ is the inverse image of $\mathcal{O}_{P'}(1)$ by the canonical injection (3.6.3), and so $\mathcal{L} = i^*(\mathcal{O}_Z(1))$. $\square$

Proposition (4.4.8). — Let $q : X \rightarrow Y$ be a morphism, $\mathcal{L}$ a very ample (relative to q) $\mathcal{O}_X$ -module, and $\mathcal{L}'$ an invertible $\mathcal{O}_X$ -module, such that there exists a quasi-coherent $\mathcal{O}_Y$ -module $\mathcal{E}'$ and a surjective homomorphism $q^*(\mathcal{E}') \rightarrow \mathcal{L}'$. Then $\mathcal{L} \otimes_{\mathcal{O}_X} \mathcal{L}'$ is very ample relative to q .

Proof. The hypothesis implies the existence of a Y -morphism $r' : X \rightarrow P' = \mathbf{P}(\mathcal{E}')$ such that $\mathcal{L}' = r'^*(\mathcal{O}_{P'}(1))$ (4.2.1). There is, by hypothesis, a quasi-coherent $\mathcal{O}_Y$ -module $\mathcal{E}$ and a ^[II:81] Y -immersion $r : X \rightarrow P = \mathbf{P}(\mathcal{E})$ such that $\mathcal{L} = r^*(\mathcal{O}_P(1))$. Let $Q = \mathbf{P}(\mathcal{E} \otimes \mathcal{E}')$, and consider the Segre morphism $\zeta : P \times_Y P' \rightarrow Q$ (4.3.1). Since r is an immersion, so too is $(r, r')_Y : X \rightarrow P \times_Y P'$ (I, 5.3.14); but since ζ is an immersion (4.3.3), so too is $r'' : X \xrightarrow{(r, r')} P \times_Y P' \xrightarrow{\zeta} Q$. But also (4.3.2.1) $\zeta(\mathcal{O}_Q(1))$ is isomorphic to $\mathcal{O}_P(1) \otimes_Y \mathcal{O}_{P'}(1)$, and so (I, 9.1.4) $r''^*(\mathcal{O}_Q(1))$ is isomorphic to $\mathcal{L} \otimes \mathcal{L}'$, which proves the proposition. $\square$

Corollary (4.4.9). — Let $q : X \rightarrow Y$ be a morphism.

- (1) Let $\mathcal{L}$ be an invertible $\mathcal{O}_X$ -module, and $\mathcal{K}$ an invertible $\mathcal{O}_Y$ -module. For $\mathcal{L}$ to be very ample relative to q , it is necessary and sufficient for $\mathcal{L} \otimes q^*(\mathcal{K})$ to be so.
- (2) If $\mathcal{L}$ and $\mathcal{L}'$ are very ample (relative to q) $\mathcal{O}_X$ -modules, then so too is $\mathcal{L} \otimes \mathcal{L}'$; in particular, $\mathcal{L}^{\otimes n}$ is very ample relative to q for all $n > 0$.

Proof. Claim (ii) is an immediate consequence of (4.4.8), as well as the necessity of condition (i); conversely, if $\mathcal{L} \otimes q^*(\mathcal{K})$ is very ample, then so too is $(\mathcal{L} \otimes q^*(\mathcal{K})) \otimes q^*(\mathcal{K}^{-1})$, by the above, and the latter $\mathcal{O}_X$ -module is isomorphic to $\mathcal{L}$ ((0, 5.4.3) and (0, 5.4.5)). $\square$

Proposition (4.4.10). —

- (i) For every prescheme Y , every invertible $\mathcal{O}_Y$ -module $\mathcal{L}$ is very ample relative to the identity morphism 1_Y .
- (i bis) Let $f : X \rightarrow Y$ be a morphism, and $j : X' \rightarrow X$ an immersion. If $\mathcal{L}$ is a very ample (relative to f) $\mathcal{O}_X$ -module, then $j^*(\mathcal{L})$ is very ample relative to $f \circ j$.
- (ii) Let Z be a quasi-compact prescheme, $f : X \rightarrow Y$ a morphism of finite type, $g : Y \rightarrow Z$ a quasi-compact morphism, $\mathcal{L}$ a very ample (relative to f) $\mathcal{O}_X$ -module, and $\mathcal{K}$ a very ample (relative to g) $\mathcal{O}_Y$ -module. Then there exists some integer $n_0 > 0$ such that $\mathcal{L} \otimes f^*(\mathcal{K}^{\otimes n})$ is very ample relative to $g \circ f$ for all $n \geq n_0$.
- (iii) Let $f : X \rightarrow Y$ and $g : Y' \rightarrow Y$ be morphisms, and let $X' = X_{(Y')}$. If $\mathcal{L}$ is a very ample (relative to f) $\mathcal{O}_X$ -module, then $\mathcal{L}' = \mathcal{L} \otimes_Y \mathcal{O}_{Y'}$ is a very ample (relative to $f_{(Y')}$) $\mathcal{O}_{X'}$ -module.
- (iv) Let $f_i : X_i \rightarrow Y_i$ ($i = 1, 2$) be S -morphisms. If $\mathcal{L}_i$ is a very ample (relative to f_i) $\mathcal{O}_{X_i}$ -module ($i = 1, 2$), then $\mathcal{L}_1 \otimes_S \mathcal{L}_2$ is very ample relative to $f_1 \times_S f_2$.
- (v) Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be morphisms. If an $\mathcal{O}_X$ -module $\mathcal{L}$ is very ample relative to $g \circ f$, then it is also very ample relative to f .
- (vi) Let $f : X \rightarrow Y$ be a morphism, and j the canonical injection $X_{\text{red}} \rightarrow X$. If an $\mathcal{O}_X$ -module $\mathcal{L}$ is very ample relative to f , then $j^*(\mathcal{L})$ is very ample relative to f_{red} .

Proof. Property (i bis) follows immediately from the definition (4.4.2), and it is immediate that (vi) follows formally from (i bis) and (v), by an argument copied from the proof of (I, 5.5.12), which we leave to the reader. To prove (v), we consider, as in (I, 5.5.12), the factorisation $X \xrightarrow{\Gamma_f} X \times_Z Y \xrightarrow{p_2} Y$, where $p_2 = (g \circ f) \times 1_Y$. It follows from the hypothesis and from (i) and (iv) that $\mathcal{L} \otimes_{\mathcal{O}_Z} \mathcal{O}_Y$ is very ample for p_2 ; but also $\mathcal{L} = \Gamma_f^*(\mathcal{L} \otimes_{\mathcal{O}_Z} \mathcal{O}_Y)$ (I, 9.1.4), and Γ_f is an immersion (I, 5.3.11); we can thus apply (i bis).

[II:82] To prove (i), we apply the definition (4.4.2) with $\mathcal{E} = \mathcal{L}$, and note that then $\mathbf{P}(\mathcal{E})$ can be identified with Y (4.1.4).

Now we prove (iii). There exists a quasi-coherent $\mathcal{O}_Y$ -module $\mathcal{E}$ and a Y -immersion $i : X \rightarrow \mathbf{P}(\mathcal{E}) = P$ such that $\mathcal{L} = i^*(\mathcal{O}_P(1))$; if we let $\mathcal{E}' = g^*(\mathcal{E})$, then $\mathcal{E}'$ is a quasi-coherent $\mathcal{O}_{Y'}$ -module, and we have that $P' = \mathbf{P}(\mathcal{E}') = P_{(Y')}$ (4.1.3.1), that $i_{(Y')}$ is an immersion from $X_{(Y')}$ into P' (I, 4.3.2), and that $\mathcal{L}'$ is isomorphic to $(i_{(Y')})^*(\mathcal{O}_{P'}(1))$ (4.2.10).

To prove (iv), note that there is, by hypothesis, a Y_i -immersion $r_i : X_i \rightarrow P_i = \mathbf{P}(\mathcal{E}_i)$, where $\mathcal{E}_i$ is a quasi-coherent $\mathcal{O}_{Y_i}$ -module, and $\mathcal{L}_i = r_i^*(\mathcal{O}_{P_i}(1))$ ($i = 1, 2$); $r_1 \times_S r_2$ is an S -immersion of $X_1 \times_S X_2$ into $P_1 \times_S P_2$ (I, 4.3.1), and the inverse image of $\mathcal{O}_{P_1}(1) \otimes_S \mathcal{O}_{P_2}(1)$ under this immersion is $\mathcal{L}_1 \otimes_S \mathcal{L}_2$. Now let $T = Y_1 \times_S Y_2$, and let p_1 and p_2 be the projections from T to Y_1 and Y_2 , respectively. If we let $P'_i = \mathbf{P}(p_i^*(\mathcal{E}_i))$ ($i = 1, 2$), then $P'_i = P_i \times_{Y_i} T$, by (4.1.3.1), and so

$$P'_1 \times_T P'_2 = (P_1 \times_{Y_1} T) \times_T (P_2 \times_{Y_2} T) = P_1 \times_{Y_1} (T \times_{Y_2} P_2) = P_1 \times_{Y_1} (Y_1 \times_S P_2) = P_1 \times_S P_2$$

up to canonical isomorphism. Similarly, $\mathcal{O}_{P'_i}(1) = \mathcal{O}_{P_i}(1) \otimes_{Y_i} \mathcal{O}_T$ (4.1.3.2), and an analogous calculation (based in particular on (I, 9.1.9.1) and (I, 9.1.2.2)) shows that, in the above identification, $\mathcal{O}_{P'_1}(1) \otimes_T \mathcal{O}_{P'_2}(1)$ can be identified with $\mathcal{O}_{P_1}(1) \otimes_S \mathcal{O}_{P_2}(1)$. We can thus consider $r_1 \times_S r_2$ as a T -immersion from $X_1 \times_S X_2$ into $P'_1 \times_T P'_2$, with the inverse image of $\mathcal{O}_{P'_1}(1) \otimes_T \mathcal{O}_{P'_2}(1)$ under this immersion being $\mathcal{L}_1 \otimes_S \mathcal{L}_2$. We then finish the argument as in (4.4.8) by using the Segre morphism.

It remains only to prove (ii). We can first of all restrict to the case where Z is an affine scheme, since, in general, there exists a finite cover (U_i) of Z by affine opens; if the proposition were proven for $\mathcal{K}|_{g^{-1}(U_i)}$, $\mathcal{L}|_{f^{-1}(g^{-1}(U_i))}$, and an integer n_i , then it would suffice to take n_0 to be the largest of the n_i to prove the proposition for $\mathcal{K}$ and $\mathcal{L}$ (4.4.5). The hypothesis implies that f and g are separated morphisms, and so X and Y are quasi-compact schemes.

There is an immersion $r : X \rightarrow P = \mathbf{P}(\mathcal{E})$, where $\mathcal{E}$ is a quasi-coherent $\mathcal{O}_Y$ -module of finite type, and $\mathcal{L} = r^*(\mathcal{O}_P(1))$, by (4.4.6). We will see that there exists a very ample (relative to the composed morphism $P \xrightarrow{h} Y \xrightarrow{g} Z$) $\mathcal{O}_P$ -module $\mathcal{M}$ such that $\mathcal{O}_P(1)$ is isomorphic to $\mathcal{M} \otimes_Y \mathcal{K}^{\otimes(-m)}$ for some

integer m . For $n \geq m + 1$, $\mathcal{O}_P(1) \otimes_Y \mathcal{K}^{\otimes n}$ will then be very ample for Z , by hypothesis and by (iv) applied to the morphisms $h : P \rightarrow Y$ and 1_Y ; since r is an immersion and $\mathcal{L} \otimes f^*(\mathcal{K}^{\otimes n}) = r^*(\mathcal{O}_P(1) \otimes_Y \mathcal{K}^{\otimes n})$, the conclusion will then follow from (i bis). To prove our claim concerning $\mathcal{O}_P(1)$, we will use the following lemma:

Lemma (4.4.10.1). — *Let Z be a quasi-compact scheme, or a prescheme whose underlying space is Noetherian, and let $g : Y \rightarrow Z$ be a quasi-compact morphism, $\mathcal{K}$ a very ample (with respect to g) invertible $\mathcal{O}_Y$ -module, and $\mathcal{E}$ a quasi-coherent $\mathcal{O}_Y$ -module of finite type. Then there exists an integer m_0 such that, for all $m \geq m_0$, $\mathcal{E}$ is isomorphic to a quotient of an $\mathcal{O}_Y$ -module of the form $g^*(\mathcal{F}) \otimes \mathcal{K}^{\otimes(-m)}$, where $\mathcal{F}$ is a quasi-coherent $\mathcal{O}_Z$ -module of finite type (depending on m).*

This lemma will be proven in (4.5.10.1); the reader can verify that (4.4.10) is not used anywhere in (4.5).

[II:83] Assuming this lemma, there exists a closed immersion j_1 from P to

$$P_1 = \mathbf{P}(g^*(\mathcal{F}) \otimes \mathcal{K}^{\otimes(-m)})$$

such that $\mathcal{O}_P(1)$ is isomorphic to $j_1^*(\mathcal{O}_{P_1}(1))$ (4.1.2). Now, there exists an isomorphism from P_1 to $P_2 = \mathbf{P}(g^*(\mathcal{F}))$, sending $\mathcal{O}_{P_2}(1) \otimes_Y \mathcal{K}^{\otimes(-m)}$ to $\mathcal{O}_{P_1}(1)$ (4.1.4); we thus have a closed immersion $j_2 : P \rightarrow P_2$ such that $\mathcal{O}_P(1)$ is isomorphic to $j_2^*(\mathcal{O}_{P_2}(1)) \otimes_Y \mathcal{K}^{\otimes(-m)}$. Finally, P_2 can be identified with $P_3 \times_Z Y$, where $P_3 = \mathbf{P}(\mathcal{F})$, and $\mathcal{O}_{P_2}(1)$ with $\mathcal{O}_{P_3}(1) \otimes_Z \mathcal{O}_Y$ (4.1.3). By definition, $\mathcal{O}_{P_3}(1)$ is very ample for Z ; since so too is $\mathcal{K}$, we conclude, from (iv), that $\mathcal{O}_{P_2}(1) \otimes_Y \mathcal{K}$ is very ample for Z ; so too is $\mathcal{M} = j_2^*(\mathcal{O}_{P_2}(1) \otimes_Y \mathcal{K})$ by (i bis), and $\mathcal{O}_P(1)$ is isomorphic to $\mathcal{M} \otimes_Y \mathcal{K}^{\otimes(-m-1)}$, which finishes the proof. $\square$

Proposition (4.4.11). — *Let $f : X \rightarrow Y$ and $f' : X' \rightarrow Y$ be morphisms, X'' the sum prescheme $X \sqcup X'$, and f'' the morphism $X'' \rightarrow Y$ that agrees with f (resp. f') on X (resp. X'). Let $\mathcal{L}$ (resp. $\mathcal{L}'$) be an invertible $\mathcal{O}_X$ -module (resp. invertible $\mathcal{O}_{X'}$ -module), and let $\mathcal{L}''$ be the invertible $\mathcal{O}_{X''}$ -module that agrees with $\mathcal{L}$ (resp. $\mathcal{L}'$) on X (resp. X'). For $\mathcal{L}''$ to be very ample relative to f'' , it is necessary and sufficient for $\mathcal{L}$ to be very ample relative to f and for $\mathcal{L}'$ to be very ample relative to f' .*

Proof. We can immediately restrict to the case where Y is affine. If $\mathcal{L}''$ is very ample then so too are $\mathcal{L}$ and $\mathcal{L}'$, by (4.4.10, (i bis)). Conversely, if $\mathcal{L}$ and $\mathcal{L}'$ are very ample, then it follows immediately from the definition (4.4.2) and from (4.3.6) that $\mathcal{L}''$ is very ample. $\square$

4.5. Ample sheaves

(4.5.1). Given a prescheme X and an invertible $\mathcal{O}_X$ -module $\mathcal{L}$, we define, for every $\mathcal{O}_X$ -module $\mathcal{F}$ (when there will be no confusion possible over $\mathcal{L}$) $\mathcal{F}(n) = \mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{L}^{\otimes n}$ ($n \in \mathbf{Z}$); we also define $S = \bigoplus_{n \geq 0} \Gamma(X, \mathcal{L}^{\otimes n})$ (a graded subring of the ring $\Gamma_{\bullet}(\mathcal{L})$ defined in (0, 5.4.6)). If we consider X as a $\mathbf{Z}$ -prescheme, and we denote by p the structure morphism $X \rightarrow \text{Spec}(\mathbf{Z})$, then there is a bijective correspondence between homomorphisms $p^*(\tilde{S}) \rightarrow \bigoplus_{n \geq 0} \mathcal{L}^{\otimes n}$ of graded $\mathcal{O}_X$ -algebras and endomorphisms of the graded ring S (I, 2.2.5); the homomorphism $\varepsilon : p^*(\tilde{S}) \rightarrow \bigoplus_{n \geq 0} \mathcal{L}^{\otimes n}$ that then corresponds to the identity automorphism of S is said to be *canonical*. There is a corresponding (3.7.1) morphism $G(\varepsilon) \rightarrow \text{Proj}(S)$ that is also said to be *canonical*.

Theorem (4.5.2). — *Let X be a quasi-compact scheme or a prescheme whose underlying space is Noetherian, $\mathcal{L}$ an invertible $\mathcal{O}_X$ -module, and S the graded ring $\bigoplus_{n \geq 0} \Gamma(X, \mathcal{L}^{\otimes n})$. Then the following conditions are equivalent:*

- (a) When f runs over the set of homogeneous elements of S_+ , the X_f form a base for the topology of X .
- (a') When f runs over the set of homogeneous elements of S_+ , the X_f that are affine form a cover of X .
- (b) The canonical morphism $G(\varepsilon) \rightarrow \text{Proj}(S)$ (4.5.1) is everywhere defined and is a dominant open immersion. [II:84]
- (b') The canonical morphism $G(\varepsilon) \rightarrow \text{Proj}(S)$ is everywhere defined and is a homeomorphism from the underlying space of X to a subspace of $\text{Proj}(S)$.
- (c) For every quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$, if we denote by $\mathcal{F}_n$ the sub- $\mathcal{O}_X$ -module of $\mathcal{F}(n)$ generated by the sections of $\mathcal{F}(n)$ over X , then $\mathcal{F}$ is the sum of the sub- $\mathcal{O}_X$ -modules $\mathcal{F}_n(-n)$ over the integers $n > 0$.
- (c') Property (c) holds for every quasi-coherent sheaf of ideals of $\mathcal{O}_X$.

Furthermore, if (f_α) is a family of homogeneous elements of S_+ such that the X_{f_α} are affine, then the restriction to $\bigcup_\alpha X_{f_\alpha}$ of the canonical morphism $X \rightarrow \text{Proj}(S)$ is an isomorphism from $\bigcup_\alpha X_{f_\alpha}$ to $\bigcup_\alpha (\text{Proj}(S))_{f_\alpha}$.

Proof. It is clear that (b) implies (b'), and (b') implies (a) by (3.7.3.1) (taking into account the fact that ε^b is the identity). Condition (a) implies (a'), since every $x \in X$ has an affine neighbourhood U such that $\mathcal{L}|_U$ is isomorphic to $\mathcal{O}_X|_U$; if $f \in \Gamma(X, \mathcal{L}^{\otimes n})$ is such that $x \in X_f \subset U$, then X_f is also the set of $x' \in U$ such that $(f|_U)(x') \neq 0$, and it is thus an affine open subset (I, 1.3.6). To prove that (a') implies (b), it suffices to prove the last claim of the theorem, and to further prove that, if $X = \bigcup_\alpha X_{f_\alpha}$, then condition (iv) of (3.8.2) is satisfied. This latter point follows immediately from (I, 9.3.1, (i)). As for the last claim of (4.5.2), since X_{f_α} is the inverse image of $(\text{Proj}(S))_{f_\alpha}$ under $G(\varepsilon) \rightarrow \text{Proj}(S)$, it suffices to apply (I, 9.3.2). Thus (a), (a'), (b), and (b') are all equivalent.

To show that (a') implies (c), note that, if X_f is affine (with $f \in S_k$), then $\mathcal{F}|_{X_f}$ is generated by its sections over X_f (I, 1.3.9); on the other hand (I, 9.3.1, (ii)), such a section s is of the form $(t|_{X_f}) \otimes (f|_{X_f})^{-m}$, where $t \in \Gamma(X, \mathcal{F}(km))$; by definition, t is also a section of $\mathcal{F}_{km}$, so s is indeed a section of $\mathcal{F}_{km}(-km)$ over X_f , which proves (c). It is clear that (c) implies (c'), so it remains only to show that (c') implies (a). But let U be an open neighbourhood of $x \in X$, and let $\mathcal{J}$ be a quasi-coherent sheaf of ideals of $\mathcal{O}_X$ defining a closed subscheme of X that has $X - U$ as its underlying space (I, 5.2.1). Hypothesis (c') implies that there exists an integer $n > 0$ and a section f of $\mathcal{J}(n)$ over X such that $f(x) \neq 0$. But we clearly have $f \in S_n$, and $x \in X_f \subset U$, which proves (a). $\square$

When X is a prescheme whose underlying space is Noetherian, the equivalent conditions of (4.5.2) imply that X is a *scheme*, since it is isomorphic to a subscheme of the scheme $\text{Proj}(S)$, by (4.5.2, (b)).

Definition (4.5.3). — We say that an invertible $\mathcal{O}_X$ -module $\mathcal{L}$ is *ample* if X is a quasi-compact scheme and if the equivalent conditions of (4.5.2) are satisfied.

It evidently follows from criterion (a) of (4.5.2) that, if $\mathcal{L}$ is an ample $\mathcal{O}_X$ -module, then, for every open subset U of X , $\mathcal{L}|_U$ is an ample $(\mathcal{O}_X|_U)$ -module.

It follows from the proof of (4.5.2) that the *affine* X_f form a base for the topology of X . Furthermore:

Corollary (4.5.4). — Let $\mathcal{L}$ be an ample $\mathcal{O}_X$ -module. For every finite subspace Z of X and every neighbourhood U of Z , there exists an integer n and some $f \in \Gamma(X, \mathcal{L}^{\otimes n})$ such that X_f is an affine neighbourhood of Z contained in U .

Proof. [II:85] By (4.5.2, (b)), it suffices to prove that, for every finite subset Z' of $\text{Proj}(S)$ and every open neighbourhood U of Z' , there exists a homogeneous element $f \in S_+$ such that $Z' \subset (\text{Proj}(S))_f \subset U$ (2.4.1). But, by definition, the closed set Y , complement of U in $\text{Proj}(S)$, is of the form $V_+(\mathfrak{J})$, where $\mathfrak{J}$ is a graded ideal of S that does not contain S_+ (2.3.2); also, the points of Z' are, by definition, graded prime ideals $\mathfrak{p}_i$ of S_+ that do not contain $\mathfrak{J}$ (2.3.1). There thus exists an element $f \in \mathfrak{J}$ that does not belong to any of the $\mathfrak{p}_i$ (Bourbaki, *Alg. comm.*, chap. II, §1, no. 1, prop. 2), and, since the $\mathfrak{p}_i$ are graded, the argument made *loc. cit.* shows that we can even take f to be homogeneous; this element then satisfies the claim. $\square$

Proposition (4.5.5). — Suppose that X is a quasi-compact scheme or a prescheme whose underlying space is Noetherian. Then conditions (a) to (c') of (4.5.2) are equivalent to the following:

- (d) For every quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$ of finite type, there exists an integer n_0 such that, for all $n \geq n_0$, $\mathcal{F}(n)$ is generated by its sections over X .
- (d') For every quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$ of finite type, there exist integers $n > 0$ and $k > 0$ such that $\mathcal{F}$ is isomorphic to a quotient of the $\mathcal{O}_X$ -module $\mathcal{L}^{\otimes(-n)} \otimes \mathcal{O}_X^k$.
- (d'') Property (d') holds for every quasi-coherent sheaf of ideals of $\mathcal{O}_X$ of finite type.

Proof. Since X is quasi-compact, if a quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$ of finite type is such that $\mathcal{F}(n)$ (which is of finite type) is generated by its sections over X , then $\mathcal{F}(n)$ is generated by a finite number of these sections (0, 5.2.3), and so (d) implies (d'), and it is clear that (d') implies (d''). Since every quasi-coherent $\mathcal{O}_X$ -module $\mathcal{G}$ is the inductive limit of its sub- $\mathcal{O}_X$ -modules of finite type (I, 9.4.9), to satisfy condition (c') of (4.5.2), it suffices to do so for a quasi-coherent sheaf of ideals of

$\mathcal{O}_X$ that is of finite type, and (d'') thus implies (c'). It remains only to show that, if $\mathcal{L}$ is ample, then property (d) is satisfied. Consider a finite cover of X by X_{f_i} ($f_i \in S_{n_i}$), that we can assume to be affine; by replacing the f_i with suitable powers (which does not alter the X_{f_i}), we can assume that all the n_i are equal to one single integer m . The sheaf $\mathcal{F}|_{X_{f_i}}$, being of finite type, by hypothesis, is generated by a finite number of its sections h_{ij} over X_{f_i} (I, 1.3.13); so there exists an integer k_0 such that the section $h_{ij} \otimes f_i^{\otimes k_0}$ extends to a section of $\mathcal{F}(k_0 m)$ over X for every pair (i, j) (I, 9.3.1). A fortiori, the $h_{ij} \otimes f_i^{\otimes k_0}$ extend to sections of $\mathcal{F}(km)$ over X for every $k \geq k_0$, and, for these values of k , $\mathcal{F}(km)$ is thus generated by its sections over X . For every p such that $0 < p < m$, $\mathcal{F}(p)$ is also of finite type, and so there exists an integer k_p such that $\mathcal{F}(p)(km) = \mathcal{F}(p + km)$ is generated by its sections over X for all $k \geq k_p$. Taking n_0 to be the largest of the $k_p m$, we thus conclude that $\mathcal{F}(n)$ is generated by its sections over X for all $n \geq n_0$, since such an n is of the form $n = km + p$, with $k \geq k_p$ and $0 \leq p < m$. $\square$

Proposition (4.5.6). — *Let X be a quasi-compact scheme, and $\mathcal{L}$ an invertible $\mathcal{O}_X$ -module.*

- (i) *Let $n > 0$ be an integer. For $\mathcal{L}$ to be ample, it is necessary and sufficient for $\mathcal{L}^{\otimes n}$ to be ample.*
- (ii) *Let $\mathcal{L}'$ be an invertible $\mathcal{O}_X$ -module such that, for all $x \in X$, there exists an integer $n > 0$ [II:86] and a section s' of $\mathcal{L}'^{\otimes n}$ over X such that $s'(x) \neq 0$. Then, if $\mathcal{L}$ is ample, so too is $\mathcal{L} \otimes \mathcal{L}'$.*

Proof. Property (i) is an evident consequence of criterion (a) of (4.5.2), since $X_{f^{\otimes n}} = X_f$. On the other hand, if $\mathcal{L}$ is ample, then, for every $x \in X$ and every neighbourhood U of x , there exists some $m > 0$ and $f \in \Gamma(X, \mathcal{L}^{\otimes m})$ such that $x \in X_f \subset U$ (4.5.2, (a)); if $f' \in \Gamma(X, \mathcal{L}'^{\otimes n})$ is such that $f'(x) \neq 0$, then $s(x) \neq 0$ for $s = f^{\otimes n} \otimes f'^{\otimes m} \in \Gamma(X, (\mathcal{L} \otimes \mathcal{L}')^{\otimes mn})$, and so $x \in X_s \subset X_f \subset U$, which proves that $\mathcal{L} \otimes \mathcal{L}'$ is ample (4.5.2, (a)). $\square$

Corollary (4.5.7). — *The tensor product of two ample $\mathcal{O}_X$ -modules is ample.*

Corollary (4.5.8). — *Let $\mathcal{L}$ be an ample $\mathcal{O}_X$ -module, and $\mathcal{L}'$ an invertible $\mathcal{O}_X$ -module; then there exists an integer $n_0 > 0$ such that $\mathcal{L}^{\otimes n} \otimes \mathcal{L}'$ is ample and generated by its sections over X for $n \geq n_0$.*

Proof. It follows from (4.5.5) that there exists an integer m_0 such that $\mathcal{L}^{\otimes m} \otimes \mathcal{L}'$ is generated by its sections over X for all $m \geq m_0$; by (4.5.6), we can then take $n_0 = m_0 + 1$. $\square$

Remark (4.5.9). — Let $P = H^1(X, \mathcal{O}_X^\times)$ be the group of classes of invertible $\mathcal{O}_X$ -modules (0, 5.4.7), and let P^+ be the subset of P consisting of classes of ample sheaves. Suppose that P^+ is non-empty. Then it follows from (4.5.7) and (4.5.8) that

$$P^+ + P^+ \subset P^+ \quad \text{and} \quad P^+ - P^+ = P$$

or, in other words, $P^+ \cup \{0\}$ is the set of *positive* elements in P for a *preorder* structure on P that is compatible with its group structure, and is even *archimedean*, by (4.5.8). This is why we sometimes say “positive sheaf” instead of ample sheaf, and “negative sheaf” for the inverse of an ample sheaf (but we will not use this terminology).

Proposition (4.5.10). — *Let Y be an affine scheme, $q : X \rightarrow Y$ a quasi-compact separated morphism, and $\mathcal{L}$ an invertible $\mathcal{O}_X$ -module.*

- (i) *If $\mathcal{L}$ is very ample for q , then $\mathcal{L}$ is ample.*
- (ii) *Suppose further that the morphism q is of finite type. Then, for $\mathcal{L}$ to be ample, it is necessary and sufficient for it to possess one of the following properties:*
 - (e) *There exists $n_0 > 0$ such that, for every integer $n \geq n_0$, $\mathcal{L}^{\otimes n}$ is very ample for q .*
 - (e') *There exists $n > 0$ such that $\mathcal{L}^{\otimes n}$ is very ample for q .*

Proof. The first claim follows from the definition (4.4.2) of a very ample $\mathcal{O}_X$ -module: if A is the ring of Y , then there exists an A -module E and a surjective homomorphism

$$\psi : q^*((\mathbf{S}(E))^\sim) \longrightarrow \bigoplus_{n \geq 0} \mathcal{L}^{\otimes n}$$

such that $i = r_{\mathcal{L}, \psi}$ is an everywhere-defined immersion $X \rightarrow P = \mathbf{P}(\tilde{E})$ and such that $\mathcal{L} = i^*(\mathcal{O}_P(1))$; since the $D_+(f)$ for f homogeneous in $(\mathbf{S}(E))_+$ form a base for the topology of P , and

since $i^{-1}(D_+(f)) = X_{\psi^b(f)}$, by (3.7.3.1), we see that condition (a) of (4.5.2) is satisfied, and so $\mathcal{L}$ is ample.

Now to prove that, if q is of finite type and $\mathcal{L}$ is ample, then condition (e) is satisfied. Firstly, it follows from criterion (b) of (4.5.2) and from (4.4.1, (i)) that there exists ^[II:87] an integer k_0 such that $\mathcal{L}^{\otimes k_0}$ is very ample relative to q . Also, by (4.5.5), there exists an integer m_0 such that, for all $m \geq m_0$, $\mathcal{L}^{\otimes m}$ is generated by its sections over X . Let $n_0 = k_0 + m_0$; if $n \geq n_0$, then we can write $n = k_0 + m$ with $m \geq m_0$, whence $\mathcal{L}^{\otimes n} = \mathcal{L}^{\otimes k_0} \otimes \mathcal{L}^{\otimes m}$. Since $\mathcal{L}^{\otimes m}$ is generated by its sections over X , it follows from (4.4.8) and (3.4.7) that $\mathcal{L}^{\otimes n}$ is very ample relative to q . Finally, it is clear that (e) implies (e'), and (e') implies that $\mathcal{L}$ is ample by (i) and by (4.5.6, (i)).

(4.5.10.1). [Proof of Lemma (4.4.10.1)]. Let $\mathcal{E}(n) = \mathcal{E} \otimes \mathcal{K}^{\otimes n}$; since g is separated (4.4.2), the argument of (3.4.8) applies and reduces the assertion to proving that the canonical homomorphism $g^*(g_*(\mathcal{E}(n))) \rightarrow \mathcal{E}(n)$ is surjective for all sufficiently large n . Furthermore, since Z is quasi-compact, the argument of (3.4.6) reduces us to proving the claim when Z is affine. But $\mathcal{K}$ is then ample, by (4.5.10, (ii)), and the conclusion follows from (4.5.5, (d')).

□

Corollary (4.5.11). — Let Y be an affine scheme, $q : X \rightarrow Y$ a separated morphism of finite type, $\mathcal{L}$ an ample $\mathcal{O}_X$ -module, and $\mathcal{L}'$ an invertible $\mathcal{O}_X$ -module. Then there exists an integer n_0 such that, for all $n \geq n_0$, $\mathcal{L}^{\otimes n} \otimes \mathcal{L}'$ is very ample relative to q .

Proof. There exists m_0 such that, for $m \geq m_0$, $\mathcal{L}^{\otimes m} \otimes \mathcal{L}'$ is generated by its sections over X (4.5.8); there also exists k_0 such that $\mathcal{L}^{\otimes k}$ is very ample relative to q for $k \geq k_0$. Then $\mathcal{L}^{\otimes(k+m_0)} \otimes \mathcal{L}'$ is very ample if $k \geq k_0$ ((4.4.8) and (3.4.7)).

□

Remark (4.5.12). — We do not know if the hypothesis that an $\mathcal{O}_X$ -module $\mathcal{L}$ is such that $\mathcal{L}^{\otimes n}$ is very ample (relative to q) implies the same conclusion for $\mathcal{L}^{\otimes(n+1)}$.

Proposition (4.5.13). — Let X be a quasi-compact prescheme, Z a closed prescheme of X defined by a nilpotent quasi-coherent sheaf $\mathcal{J}$ of ideals of $\mathcal{O}_X$, and j the canonical injection $Z \rightarrow X$. For an invertible $\mathcal{O}_X$ -module $\mathcal{L}$ to be ample, it is necessary and sufficient for $\mathcal{L}' = j^*(\mathcal{L})$ to be an ample $\mathcal{O}_Z$ -module.

Proof. The condition is *necessary*. Indeed, for every section f of $\mathcal{L}^{\otimes n}$ over X , let f' be its canonical image $f \otimes 1$, which is a section of $\mathcal{L}'^{\otimes n} = \mathcal{L}^{\otimes n} \otimes_{\mathcal{O}_X} (\mathcal{O}_X / \mathcal{J})$ over the space Z (which is identical to X); it is clear that $X_f = Z_{f'}$, and so the criterion (a) of (4.5.2) shows that $\mathcal{L}'$ is ample.

To see that the condition is *sufficient*, note first of all that we can restrict to the case where $\mathcal{J}^2 = 0$, by considering the (finite) sequence of preschemes $X_k = (X, \mathcal{O}_X / \mathcal{J}^{k+1})$ with each prescheme being a closed subprescheme of the next, defined by a square-zero sheaf of ideals. But X is a scheme, since X_{red} is a scheme by hypothesis ((4.5.3) and (I, 5.5.1)). Criterion (a) of (4.5.2) shows that it suffices to prove

Lemma (4.5.13.1). — Under the hypotheses of (4.5.13), suppose further that $\mathcal{J}$ is square-zero; with $\mathcal{L}$ being an invertible $\mathcal{O}_X$ -module, let g be a section of $\mathcal{L}'^{\otimes n}$ over Z such that Z_g is affine. Then there exists an integer $m > 0$ such that $g^{\otimes m}$ is the canonical image of a section f of $\mathcal{L}^{\otimes nm}$ over X .

We have the exact sequence of $\mathcal{O}_X$ -modules

$$0 \rightarrow \mathcal{J}(n) \rightarrow \mathcal{O}_X(n) = \mathcal{L}^{\otimes n} \rightarrow \mathcal{O}_Z(n) = \mathcal{L}'^{\otimes n} \rightarrow 0$$

^[II:88] since $\mathcal{F}(n)$ is an exact functor in $\mathcal{F}$; from this, we have the exact sequence of cohomology

$$0 \rightarrow \Gamma(X, \mathcal{J}(n)) \rightarrow \Gamma(X, \mathcal{L}^{\otimes n}) \rightarrow \Gamma(X, \mathcal{L}'^{\otimes n}) \xrightarrow{\partial} H^1(X, \mathcal{J}(n))$$

that sends, in particular, g to an element $\partial g \in H^1(X, \mathcal{J}(n))$. Note that, since $\mathcal{J}^2 = 0$, $\mathcal{J}$ can be considered as a quasi-coherent $\mathcal{O}_Z$ -module, and, for all k , $\mathcal{L}'^{\otimes k} \otimes_{\mathcal{O}_Z} \mathcal{J}(n) = \mathcal{J}(n+k)$; for every section $s \in \Gamma(X, \mathcal{L}'^{\otimes k})$, tensor multiplication with s is thus a homomorphism $\mathcal{J}(n) \rightarrow \mathcal{J}(n+k)$ of $\mathcal{O}_Z$ -modules, which then gives a homomorphism $H^i(X, \mathcal{J}(n)) \xrightarrow{s} H^i(X, \mathcal{J}(n+k))$ of cohomology groups.

With this, we will see that

$$(4.5.13.2) \quad g^{\otimes m} \otimes \partial g = 0$$

for $m > 0$ large enough. In fact, Z_g is an *affine open* subset of Z , and so $H^1(Z_g, \mathcal{J}(n)) = 0$ when $\mathcal{J}(n)$ is considered as an $\mathcal{O}_Z$ -module (I, 5.1.9.2). In particular, if we set $g' = g|_{Z_g}$, and if we consider its image under the map $\partial : \Gamma(Z_g, \mathcal{L}'^{\otimes n}) \rightarrow H^1(Z_g, \mathcal{J}(n))$, then $\partial g' = 0$. To make this equation explicit, note that, in degree 1, the cohomology of a sheaf of abelian groups agrees with its Čech cohomology (G, II, 5.9); to calculate ∂g , we must thus consider a fine-enough open cover (U_α) of X , that we can suppose to be *finite* and consisting of affine opens, and take, for each α , a section $g_\alpha \in \Gamma(U_\alpha, \mathcal{L}^{\otimes n})$ whose canonical image in $\Gamma(U_\alpha, \mathcal{L}'^{\otimes n})$ is $g|_{U_\alpha}$, and to consider the cocycle class $(g_{\alpha|\beta} - g_{\beta|\alpha})$, with $g_{\alpha|\beta}$ being the restriction of g_α to $U_\alpha \cap U_\beta$ (with this cocycle taking values in $\mathcal{J}(n)$). We can further suppose that $\partial g'$ is calculated in the same way, by means of a cover given by the $U_\alpha \cap Z_g$, and restrictions $g_\alpha|_{(U_\alpha \cap Z_g)}$ (by replacing, if necessary, (U_α) by a finer cover); the equation $\partial g' = 0$ then implies that there exists, for each α , a section $h_\alpha \in \Gamma(U_\alpha \cap Z_g, \mathcal{J}(n))$ such that $(g_{\alpha|\beta} - g_{\beta|\alpha})|_{(U_\alpha \cap U_\beta \cap Z_g)} = h_{\alpha|\beta} - h_{\beta|\alpha}$, where $h_{\alpha|\beta}$ denotes the restriction of h_α to $U_\alpha \cap U_\beta \cap Z_g$ (G, II, 5.11). Then there exists an integer $m > 0$ such that $g^{\otimes m} \otimes h_\alpha$ is the restriction to $U_\alpha \cap Z_g$ of a section $t_\alpha \in \Gamma(U_\alpha, \mathcal{J}(n + nm))$ for all α (I, 9.3.1); thus $g^{\otimes m} \otimes (g_{\alpha|\beta} - g_{\beta|\alpha}) = t_{\alpha|\beta} - t_{\beta|\alpha}$ for every pair of indices, which proves (4.5.13.2).

Now note that, if $s \in \Gamma(X, \mathcal{O}_Z(p))$ and $t \in \Gamma(X, \mathcal{O}_Z(q))$, then, in the group $H^1(X, \mathcal{J}(p+q))$,

$$(4.5.13.3) \quad \partial(s \otimes t) = (\partial s) \otimes t + s \otimes (\partial t).$$

Indeed, we can again calculate the two members by considering an open cover (U_α) of X , and, for each α , a section $s_\alpha \in \Gamma(U_\alpha, \mathcal{O}_X(p))$ (resp. $t_\alpha \in \Gamma(U_\alpha, \mathcal{O}_X(q))$) whose canonical image in $\Gamma(U_\alpha, \mathcal{O}_Z(p))$ (resp. $\Gamma(U_\alpha, \mathcal{O}_Z(q))$) is $s|_{U_\alpha}$ (resp. $t|_{U_\alpha}$); equation (4.5.13.3) then follows from the equations

$$(s_{\alpha|\beta} \otimes t_{\alpha|\beta}) - (s_{\beta|\alpha} \otimes t_{\beta|\alpha}) = (s_{\alpha|\beta} - s_{\beta|\alpha}) \otimes t_{\alpha|\beta} + s_{\beta|\alpha} \otimes (t_{\alpha|\beta} - t_{\beta|\alpha})$$

with the same notation as above. By induction on k , we thus have

$$(4.5.13.4) \quad \partial(g^{\otimes k}) = (kg^{\otimes(k-1)}) \otimes (\partial g)$$

[II:89] and we thus conclude from (4.5.13.2) and (4.5.13.4) that $\partial(g^{\otimes(m+1)}) = 0$; thus $g^{\otimes(m+1)}$ is the canonical image of a section f of $\mathcal{L}^{\otimes n(m+1)}$ over X , which proves (4.5.13). $\square$

Corollary (4.5.14). — *Let X be a Noetherian scheme, and j the canonical injection $X_{\text{red}} \rightarrow X$. For an invertible $\mathcal{O}_X$ -module $\mathcal{L}$ to be ample, it is necessary and sufficient for $j^*(\mathcal{L})$ to be an ample $\mathcal{O}_{X_{\text{red}}}$ -module.*

Proof. This follows from (I, 6.1.6) $\square$

4.6. Relatively ample sheaves

Definition (4.6.1). — Let $f : X \rightarrow Y$ be a quasi-compact morphism, and $\mathcal{L}$ an invertible $\mathcal{O}_X$ -module. We say that $\mathcal{L}$ is *ample relative to f* , or *relative to Y* , or *f -ample*, or *Y -ample* (or even simply *ample* if no confusion may arise with the notion defined in (4.5.3)) if there exists an affine open cover (U_α) of Y such that, if we set $X_\alpha = f^{-1}(U_\alpha)$, then $\mathcal{L}|_{X_\alpha}$ is an ample $\mathcal{O}_{X_\alpha}$ -module for all α .

The existence of an f -ample $\mathcal{O}_X$ -module implies that f is necessarily *separated* ((4.5.3) and (I, 5.5.5)).

Proposition (4.6.2). — *Let $f : X \rightarrow Y$ be a quasi-compact morphism, and $\mathcal{L}$ an invertible $\mathcal{O}_X$ -module. If $\mathcal{L}$ is very ample relative to f , then it is ample relative to f .*

Proof. This follows from the local (on Y) character of the notion of a very ample sheaf (4.4.5), from the definition (4.6.1), and from criterion (4.5.10, (i)). $\square$

Proposition (4.6.3). — *Let $f : X \rightarrow Y$ be a quasi-compact morphism, and $\mathcal{L}$ an invertible $\mathcal{O}_X$ -module, and let $\mathcal{S}$ be the graded $\mathcal{O}_Y$ -algebra $\bigoplus_{n \geq 0} f_*(\mathcal{L}^{\otimes n})$. Then the following conditions are equivalent:*

- (a) $\mathcal{L}$ is f -ample.
- (b) $\mathcal{S}$ is quasi-coherent, and the canonical homomorphism $\sigma : f^*(\mathcal{S}) \rightarrow \bigoplus_{n \geq 0} \mathcal{L}^{\otimes n}$ (0, 4.4.3) is such that the Y -morphism $r_{\mathcal{L}, \sigma} : G(\sigma) \rightarrow \text{Proj}(\mathcal{S}) = P$ is everywhere defined and is a dominant open immersion.
- (b') The morphism f is separated, and the Y -morphism $r_{\mathcal{L}, \sigma}$ is everywhere defined and is a homeomorphism from the underlying space of X to a subspace of $\text{Proj}(\mathcal{S})$.

Furthermore, if any of the above are satisfied, then, for all $n \in \mathbf{Z}$, the canonical homomorphism

$$(4.6.3.1) \quad r_{\mathcal{L},\sigma}^*(\mathcal{O}_P(n)) \longrightarrow \mathcal{L}^{\otimes n}$$

defined in (3.7.9.1) is an isomorphism.

Finally, for every quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$, if we set $\mathcal{M} = \bigoplus_{n \geq 0} f_*(\mathcal{F} \otimes \mathcal{L}^{\otimes n})$, then the canonical homomorphism

$$(4.6.3.2) \quad r_{\mathcal{L},\sigma}^*(\widetilde{\mathcal{M}}) \longrightarrow \mathcal{F}$$

defined in (3.7.9.2) is an isomorphism.

Proof. We note that (a) implies that f is separated, and thus that $\mathcal{S}$ is quasi-coherent (I, 9.2.2, (a)). Since $r_{\mathcal{L},\sigma}$ being an everywhere defined immersion is of a local (on Y) character, to prove that (a) implies (b) we can suppose that Y is affine and $\mathcal{L}$ ample; [II:90] the claim then follows from (4.5.2, (b)). It is clear that (b) implies (b'); finally, to prove that (b') implies (a), it suffices to consider an affine open cover (U_α) of Y and to apply the criterion (4.5.2, (b')) to each sheaf $\mathcal{L}|_{f^{-1}(U_\alpha)}$.

For the final two claims, we use the fact that σ^b is here the identity, and the clarification of the homomorphisms (3.7.9.1) and (3.7.9.2); from this, it immediately follows that (4.6.3.1) is an isomorphism. As for (4.6.3.2), we can restrict to the case where Y is affine, and thus $\mathcal{L}$ ample; it is clear that the homomorphism (4.6.3.2) is injective, and criterion (4.5.2, (c)) shows that it is surjective, whence the conclusion. $\square$

Corollary (4.6.4). — *Let (U_α) be an open cover of Y . For $\mathcal{L}$ to be ample relative to Y , it is necessary and sufficient for $\mathcal{L}|_{f^{-1}(U_\alpha)}$ to be ample relative to U_α for all α .*

Proof. Condition (b) is in fact local on Y . $\square$

Corollary (4.6.5). — *Let $\mathcal{K}$ be an invertible $\mathcal{O}_Y$ -module. For $\mathcal{L}$ to be Y -ample, it is necessary and sufficient for $\mathcal{L} \otimes f^*(\mathcal{K})$ to be Y -ample.*

Proof. This is an evident consequence of (4.6.4), by taking the U_α to be such that $\mathcal{K}|_{U_\alpha}$ is isomorphic to $\mathcal{O}_Y|_{U_\alpha}$ for all α . $\square$

Corollary (4.6.6). — *Suppose that Y is affine; for $\mathcal{L}$ to be Y -ample, it is necessary and sufficient for $\mathcal{L}$ to be ample.*

Proof. This is an immediate consequence of the definition (4.6.1) and the criteria (4.6.3, (b)) and (4.5.2, (b)), since here $\text{Proj}(\mathcal{S}) = \text{Proj}(\Gamma(Y, \mathcal{S}))$ by definition. $\square$

Corollary (4.6.7). — *Let $f : X \rightarrow Y$ be a quasi-compact morphism. Suppose that there exists a quasi-coherent $\mathcal{O}_Y$ -module $\mathcal{E}$, and a Y -morphism $g : X \rightarrow P = \mathbf{P}(\mathcal{E})$ that is a homeomorphism from the underlying space of X to a subspace of P ; then $\mathcal{L} = g^*(\mathcal{O}_P(1))$ is Y -ample.*

Proof. We can assume Y to be affine; the corollary then follows from the criterion (4.5.2, (a)), from equation (3.7.3.1), and from (4.2.3). $\square$

Proposition (4.6.8). — *Let X be a quasi-compact scheme or a prescheme whose underlying space is Noetherian, and let $f : X \rightarrow Y$ be a quasi-compact separated morphism. For an invertible $\mathcal{O}_X$ -module $\mathcal{L}$ to be f -ample, it is necessary and sufficient for one of the following equivalent conditions to be satisfied:*

- (c) *For every quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$ of finite type, there exists an integer $n_0 > 0$ such that, for all $n \geq n_0$, the canonical homomorphism $\sigma : f^*(f_*(\mathcal{F} \otimes \mathcal{L}^{\otimes n})) \rightarrow \mathcal{F} \otimes \mathcal{L}^{\otimes n}$ is surjective.*
- (c') *For every quasi-coherent sheaf $\mathcal{J}$ of ideals of $\mathcal{O}_X$ of finite type, there exists an integer $n > 0$ such that the canonical homomorphism $\sigma : f^*(f_*(\mathcal{J} \otimes \mathcal{L}^{\otimes n})) \rightarrow \mathcal{J} \otimes \mathcal{L}^{\otimes n}$ is surjective.*

Proof. Since X is quasi-compact, so too is $f(X)$, and so there exists a finite cover (U_i) of $f(X)$ consisting of affine open subsets U_i of Y . To prove condition (c) when $\mathcal{L}$ is f -ample, we can replace Y by the U_i , and X by the $f^{-1}(U_i)$, since, if we obtain, for each i , an integer n_i such that (c) holds true (for U_i , $f^{-1}(U_i)$, and $\mathcal{L}|_{f^{-1}(U_i)}$) for all $n \geq n_i$, then it suffices to take n_0 to be the largest of the n_i in order to obtain (c) for Y , X , and $\mathcal{L}$. But if Y is affine, condition (c) follows from (4.5.5, (d)), taking (4.6.6) into account. It is trivial that (c) implies (c'). Finally, to prove that (c') implies that $\mathcal{L}$ is f -ample, we can again restrict to the case where Y is affine: in fact, every quasi-coherent sheaf $\mathcal{J}_i$

of ideals of $\mathcal{O}_X|f^{-1}(U_i)$ of finite type is the restriction of a coherent sheaf of ideals of $\mathcal{O}_X$ of finite type (I, 9.4.7), and hypothesis (c') implies that $\mathcal{J}_i \otimes (\mathcal{L}^{\otimes n}|f^{-1}(U_i))$ is generated by its sections (taking (I, 9.2.2) and (3.4.7) into account); it thus suffices to apply criterion (4.5.5, (d')). $\square$

Proposition (4.6.9). — *Let $f : X \rightarrow Y$ be a quasi-compact morphism, and $\mathcal{L}$ an invertible $\mathcal{O}_X$ -module.*

- (i) *Let $n > 0$ be an integer. For $\mathcal{L}$ to be f -ample, it is necessary and sufficient for $\mathcal{L}^{\otimes n}$ to be f -ample.*
- (ii) *Let $\mathcal{L}'$ be an invertible $\mathcal{O}_X$ -module, and suppose that there exists an integer $n > 0$ such that the canonical homomorphism $\sigma : f^*(f_*(\mathcal{L}'^{\otimes n})) \rightarrow \mathcal{L}'^{\otimes n}$ is surjective. Then, if $\mathcal{L}$ is f -ample, so too is $\mathcal{L} \otimes \mathcal{L}'$.*

Proof. We can in fact immediately restrict to the case where Y is affine, and the proposition is then an immediate consequence of (4.5.6). $\square$

Corollary (4.6.10). — *The tensor product of two f -ample $\mathcal{O}_X$ -modules is f -ample.*

Proposition (4.6.11). — *Let Y be a quasi-compact prescheme, $f : X \rightarrow Y$ a morphism of finite type, and $\mathcal{L}$ an invertible $\mathcal{O}_X$ -module. For $\mathcal{L}$ to be f -ample, it is necessary and sufficient for it to possess one of the following equivalent properties:*

- (d) *There exists some $n_0 > 0$ such that, for every integer $n \geq n_0$, $\mathcal{L}^{\otimes n}$ is very ample relative to f .*
- (d') *There exists some $n > 0$ such that $\mathcal{L}^{\otimes n}$ is very ample relative to f .*

Proof. If $\mathcal{L}$ is ample relative to f , then there exists a finite cover (U_i) of Y by affine open subsets such that the $\mathcal{L}|f^{-1}(U_i)$ are ample. We thus conclude (4.5.10) that there exists an integer n_0 such that $\mathcal{L}^{\otimes n}|f^{-1}(U_i)$ is very ample relative to $f^{-1}(U_i) \rightarrow U_i$ for all $n \geq n_0$ and every i , and so $\mathcal{L}^{\otimes n}$ is very ample relative to f (4.4.5). Conversely, (d') already implies that $\mathcal{L}^{\otimes n}$ is f -ample (4.6.2), and thus so too is $\mathcal{L}$ (4.6.9, (i)). $\square$

Corollary (4.6.12). — *Let Y be a quasi-compact prescheme, $f : X \rightarrow Y$ a morphism of finite type, and $\mathcal{L}$ and $\mathcal{L}'$ invertible $\mathcal{O}_X$ -modules. If $\mathcal{L}$ is f -ample, then there exists some n_0 such that $\mathcal{L}^{\otimes n} \otimes \mathcal{L}'$ is very ample relative to f for all $n \geq n_0$.*

Proof. We argue as in (4.6.11), by using a finite affine open cover of Y and (4.5.11). $\square$

Proposition (4.6.13). — (i) *For every prescheme Y , every invertible $\mathcal{O}_Y$ -module $\mathcal{L}$ is ample relative to the identity morphism 1_Y .*

- (i bis) *Let $f : X \rightarrow Y$ be a quasi-compact morphism, and $j : X' \rightarrow X$ a quasi-compact morphism that is a homeomorphism from the underlying space of X' to a subspace of X . If $\mathcal{L}$ is an $\mathcal{O}_X$ -module that is ample relative to f , then $j^*(\mathcal{L})$ is ample relative to $f \circ j$.*
- (ii) *Let Z be a quasi-compact prescheme, $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ quasi-compact morphisms, $\mathcal{L}$ an $\mathcal{O}_X$ -module that is ample relative to f , and $\mathcal{K}$ an $\mathcal{O}_Y$ -module that is ample relative to g . Then there exists an integer $n_0 > 0$ such that $\mathcal{L} \otimes f^*(\mathcal{K}^{\otimes n})$ is ample relative to $g \circ f$ for all $n \geq n_0$.*
- (iii) *Let $f : X \rightarrow Y$ be a quasi-compact morphism, and $g : Y' \rightarrow Y$ a morphism, and let $X' = X_{(Y')}$. If $\mathcal{L}$ is an $\mathcal{O}_X$ -module that is ample relative to f , then $\mathcal{L}' = \mathcal{L} \otimes_Y \mathcal{O}_{Y'}$ is an $\mathcal{O}_{X'}$ -module that is ample relative to $f_{(Y')}$.*
- (iv) *Let $f_i : X_i \rightarrow Y_i$ ($i = 1, 2$) be quasi-compact S -morphisms. If $\mathcal{L}_i$ is an $\mathcal{O}_{X_i}$ -module that is ample relative to f_i ($i = 1, 2$), then $\mathcal{L}_1 \otimes_S \mathcal{L}_2$ is ample relative to $f_1 \times_S f_2$.*
- (v) *Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be morphisms such that $g \circ f$ is quasi-compact. If an $\mathcal{O}_X$ -module $\mathcal{L}$ is ample relative to $g \circ f$, and if g is separated or the underlying space of X is locally Noetherian, then $\mathcal{L}$ is ample relative to f .*
- (vi) *Let $f : X \rightarrow Y$ be a quasi-compact morphism, and j the canonical injection $X_{\text{red}} \rightarrow X$. If $\mathcal{L}$ is an $\mathcal{O}_X$ -module that is ample relative to f , then $j^*(\mathcal{L})$ is ample relative to f_{red} .*

Proof. Note first of all that (v) and (vi) follow from (i), (i bis), and (iv) by the same argument as in (4.4.10), by using (4.6.4) instead of (4.4.5); we leave the details of the argument to the reader. Claim (i) is trivially a consequence of (4.4.10, (i)) and (4.6.2). To prove (i bis), (iii), and (iv), we will use the following lemma:

Lemma (4.6.13.1). — (i) *Let $u : Z \rightarrow S$ be a morphism, $\mathcal{L}$ an invertible $\mathcal{O}_S$ -module, and s a section of $\mathcal{L}$ over S , and let s' be the canonically corresponding section $u^*(\mathcal{L}) = \mathcal{L}'$ over Z . Then $Z_{s'} = u^{-1}(S_s)$.*

- (ii) Let Z and Z' be S -preschemes, p and p' the projections of $T = Z \times_S Z'$, $\mathcal{L}$ (resp. $\mathcal{L}'$) an invertible $\mathcal{O}_Z$ -module (resp. invertible $\mathcal{O}_{Z'}$ -module), and let t (resp. t') be a section of $\mathcal{L}$ (resp. $\mathcal{L}'$) over Z (resp. Z'), and s (resp. s') the corresponding section of $p^*(\mathcal{L})$ (resp. $p'^*(\mathcal{L}')$) over $Z \times_S Z'$. Then $T_{s \otimes s'} = Z_t \times_S Z'_{t'}$.

Proof. It follows from the definitions that we can restrict to the case where all the preschemes in question are affine. Furthermore, in (i), we can suppose that $\mathcal{L} = \mathcal{O}_B$; claim (i) then follows immediately from (I, 1.2.2.2). Similarly, in (ii), we can restrict to the case where $\mathcal{L} = \mathcal{O}_Z$ and $\mathcal{L}' = \mathcal{O}_{Z'}$, and then the claim reduces to Lemma (4.3.2.4). $\square$

We now prove (i bis). We can assume that Y is affine (4.6.4), and thus that $\mathcal{L}$ is ample (4.6.6); if s runs over the set of the union of the $\Gamma(X, \mathcal{L}^{\otimes n})$ ($n > 0$), then the X_s form a base for the topology of X (4.5.2, (a)), and so, by hypothesis, the $j^{-1}(X_s)$ form a base for the topology of X' ; we thus conclude, by Lemma (4.6.13.1, (i)) and (4.5.2, (a)), that $j^*(\mathcal{L})$ is ample.

Next we prove (iii). We can again suppose that Y and Y' are affine (4.6.4), whence it follows that the projection $h : X' \rightarrow X$ is affine (1.5.5). Since $\mathcal{L}$ is ample (4.6.6), if s runs over the set of sections of the $\mathcal{L}^{\otimes n}$ ($n > 0$) over X such that X_s is affine, then the X_s cover X (4.5.2, (a')), and so the $h^{-1}(X_s)$ are affine (1.2.5) and cover X' ; it thus follows again from Lemma (4.6.13.1, (i)) and from (4.5.2, (a')) that $\mathcal{L}'$ is ample, since the morphism $f_{(Y')}$ is quasi-compact (I, 6.6.4, (iii)).

To prove (iv), note first of all that $f_1 \times_S f_2$ is quasi-compact (I, 6.6.4, (iv)). We can further suppose that S , Y_1 , and Y_2 are affine ((4.6.4) and (I, 3.2.7)), and thus that $\mathcal{L}_i$ is ample ($i = 1, 2$) (4.6.6). The open subsets $(X_1)_{s_1} \times_S (X_2)_{s_2}$ form a cover of $X_1 \times_S X_2$, where s_i runs over the sections of $\mathcal{L}_i^{\otimes n_i}$ such that $(X_i)_{s_i}$ is affine (4.5.2, (a')). Then, replacing s_1 and s_2 with suitable powers, which does not change the $(X_i)_{s_i}$, we can assume that $n_1 = n_2$. We thus deduce, from (4.6.13.1, (ii)) and (4.5.2, (a')), that $\mathcal{L}_1 \otimes_S \mathcal{L}_2$ is ample, whence the claim, since $Y_1 \times_S Y_2$ is affine (4.6.6).

It remains only to prove (ii). By the same argument as in (4.4.10), but here using (4.6.4), we can restrict to the case where Z is affine. Since $\mathcal{K}$ is then ample, and Y quasi-compact, there exists a finite number of sections $s_i \in \Gamma(Y, \mathcal{K}^{\otimes k_i})$ such that the Y_{s_i} ^[II-93] are affine and cover Y (4.5.2, (a')); replacing the s_i with suitable powers, we can further suppose that all the k_i are equal to one single integer k . Let s'_i be the sections of $f^*(\mathcal{K}^{\otimes k})$ over X that canonically correspond to the s_i , so that the $X_{s'_i} = f^{-1}(Y_{s_i})$ (4.6.13.1, (i)) cover X . Since $\mathcal{L}|_{X_{s'_i}}$ is ample ((4.6.4) and (4.6.6)), there exists, for each i , a finite number of sections $t_{ij} \in \Gamma(X, \mathcal{L}^{\otimes n_{ij}})$ such that the X_{ij} are affine, contained in the $X_{s'_i}$, and cover $X_{s'_i}$ (4.5.2, (a')); we can also suppose that all the n_{ij} are equal to one single integer n . With this in mind, X is separated and quasi-compact, and so there exists an integer $m > 0$, and, for every (i, j) , a section

$$u_{ij} \in \Gamma(X, \mathcal{L}^{\otimes n} \otimes_X f^*(\mathcal{K}^{\otimes mk}))$$

such that $t_{ij} \otimes s_i'^{\otimes m}$ is the restriction to $X_{s'_i}$ of u_{ij} (I, 9.3.1); furthermore, $X_{u_{ij}} = X_{t_{ij}}$, and so the $X_{u_{ij}}$ are affine and cover X . We can also suppose that m is of the form nr ; if we set $n_0 = rk$, then we see (4.5.2, (a')) that $\mathcal{L} \otimes_{\mathcal{O}_X} f^*(\mathcal{K}^{\otimes n_0})$ is ample. Furthermore, there exists $h_0 > 0$ such that $\mathcal{K}^{\otimes h}$ is generated by its sections over Y for all $h \geq h_0$ (4.5.5); a fortiori, $f^*(\mathcal{K}^{\otimes h})$ is generated by its sections over X for all $h \geq h_0$, by definition of the inverse images ((0, 3.7.1) and (4.4.1)). We thus conclude that $\mathcal{L} \otimes f^*(\mathcal{K}^{\otimes n_0+h})$ is ample for all $h \geq h_0$ (4.5.6), which finishes the proof. $\square$

Remark (4.6.14). — Under the conditions of (ii), one should not conclude that $\mathcal{L} \otimes f^*(\mathcal{K})$ is ample for $g \circ f$; in fact, since $\mathcal{L} \otimes f^*(\mathcal{K}^{-1})$ is also ample for f (4.6.5), we would conclude that $\mathcal{L}$ is ample for $g \circ f$; taking, in particular, g to be the identity morphism, every invertible $\mathcal{O}_X$ -module would be ample for f , which is not the case in general (see (5.1.6), (5.3.4, (i)), and (5.3.1)).

Proposition (4.6.15). — Let $f : X \rightarrow Y$ be a quasi-compact morphism, $\mathcal{J}$ a quasi-coherent locally-nilpotent sheaf of ideals of $\mathcal{O}_X$, Z the closed subscheme of X defined by $\mathcal{J}$, and $j : Z \rightarrow X$ the canonical injection. For an invertible $\mathcal{O}_X$ -module $\mathcal{L}$ to be ample for f , it is necessary and sufficient for $j^*(\mathcal{L})$ to be ample for $f \circ j$.

Proof. Since the question is local on Y (4.6.4), we can suppose Y to be affine; since X is then quasi-compact, we can suppose $\mathcal{J}$ to be nilpotent. Taking (4.6.6) into account, the proposition is then exactly the same as (4.5.13). $\square$

Corollary (4.6.16). — *Let X be a locally Noetherian prescheme, and $f : X \rightarrow Y$ a quasi-compact morphism. For an invertible $\mathcal{O}_X$ -module $\mathcal{L}$ to be ample for f , it is necessary and sufficient for its inverse image $\mathcal{L}'$ under the canonical injection $X_{\text{red}} \rightarrow X$ to be ample for f_{red} .*

Proof. We have already seen that the condition is necessary (4.6.13, (vi)); conversely, if it is satisfied, then we can restrict, to prove that $\mathcal{L}$ is ample for f , to the case where Y is affine (4.6.4); then Y_{red} is also affine, and so $\mathcal{L}'$ is ample (4.6.6), and so too is $\mathcal{L}$ by (4.5.13), since X is then Noetherian and X_{red} a closed subscheme of X defined by a quasi-coherent nilpotent sheaf of ideals (I, 6.1.6). $\square$

Proposition (4.6.17). — *With the notation and hypotheses of (4.4.11), for $\mathcal{L}''$ to be ample relative to f'' , it is necessary and sufficient for $\mathcal{L}$ to be ample relative to f and $\mathcal{L}'$ ample relative to f' .*

[II:94]

Proof. The necessity of the condition follows from (4.6.13, (i bis)). To see that the condition is sufficient, we can restrict to the case where Y is affine, and then the fact that $\mathcal{L}''$ is ample follows from criterion (4.5.2, (a)) applied to $\mathcal{L}$, $\mathcal{L}'$, and $\mathcal{L}''$, noting that a section of $\mathcal{L}$ over X can be extended (by 0) to a section of $\mathcal{L}''$ over X'' . $\square$

Proposition (4.6.18). — *Let Y be a quasi-compact prescheme, $\mathcal{S}$ a quasi-coherent graded $\mathcal{O}_Y$ -algebra of finite type, and $X = \text{Proj}(\mathcal{S})$, and let $f : X \rightarrow Y$ be the structure morphism. Then f is of finite type, and there exists an integer $d > 0$ such that $\mathcal{O}_X(d)$ is invertible and f -ample.*

Proof. By (3.1.10), there exists an integer $d > 0$ such that $\mathcal{S}^{(d)}$ is generated by $\mathcal{S}_d$. We know that, under the canonical isomorphism between X and $X^{(d)} = \text{Proj}(\mathcal{S}^{(d)})$, $\mathcal{O}_X(d)$ is identified with $\mathcal{O}_{X^{(d)}}(1)$ (3.2.9, (ii)). We thus see that we can restrict to the case where $\mathcal{S}$ is generated by $\mathcal{S}_1$; the proposition then follows from (4.4.3) and (4.6.2) (taking into account the fact that f is a morphism of finite type (3.4.1)). $\square$

§5. QUASI-AFFINE MORPHISMS; QUASI-PROJECTIVE MORPHISMS; PROPER MORPHISMS; PROJECTIVE MORPHISMS

5.1. Quasi-affine morphisms

Definition (5.1.1). — We define a quasi-affine scheme to be a scheme isomorphic to some subscheme induced on some quasi-compact open subset of an affine scheme. We say that a morphism $f : X \rightarrow Y$ is quasi-affine, or that X (considered as a Y -prescheme via f) is a quasi-affine Y -scheme, if there exists a cover (U_α) of Y by affine open subsets such that the $f^{-1}(U_\alpha)$ are quasi-affine schemes.

It is clear that a quasi-affine morphism is *separated* (II, 5.5.5) and (I, 5.5.8)) and *quasi-compact* (I, 6.6.1); every affine morphism is evidently quasi-affine.

Recall that, for any prescheme X , setting $A = \Gamma(X, \mathcal{O}_X)$, the identity homomorphism $A \rightarrow A = \Gamma(X, \mathcal{O}_X)$ defines a morphism $X \rightarrow \text{Spec}(A)$, said to be *canonical* (I, 2.2.4); this is nothing but the canonical morphism defined in (4.5.1) for the specific case where $\mathcal{L} = \mathcal{O}_X$, if we remember that $\text{Proj}(A[T])$ is canonically identified with $\text{Spec}(A)$ (3.1.7).

Proposition (5.1.2). — *Let X be a quasi-compact scheme or a prescheme whose underlying space is Noetherian, and A the ring $\Gamma(X, \mathcal{O}_X)$. The following conditions are equivalent.*

- (a) X is a quasi-affine scheme.
- (b) The canonical morphism $u : X \rightarrow \text{Spec}(A)$ is an open immersion.
- (b') The canonical morphism $u : X \rightarrow \text{Spec}(A)$ is a homeomorphism from X to some subspace of the underlying space of $\text{Spec}(A)$.
- (c) The $\mathcal{O}_X$ -module $\mathcal{O}_X$ is very ample relative to u (4.4.2).
- (c') The $\mathcal{O}_X$ -module $\mathcal{O}_X$ is ample (4.5.1).
- (d) When f ranges over A , the X_f form a basis for the topology of X .
- (d') When f varies over A , the X_f that are affine form a cover of X . [II:95]
- (e) Every quasi-coherent $\mathcal{O}_X$ -module is generated by its sections over X .
- (e') Every quasi-coherent sheaf of ideals of $\mathcal{O}_X$ of finite type is generated by its sections over X .

Proof. It is clear that (b) implies (a), and (a) implies (c) by (4.4.4, b) applied to the identity morphism (taking into account the remark preceding this proposition); Furthermore, (c) implies (c') (4.5.10, i), and (c'), (b), and (b') are all equivalent by (4.5.2, b) and (4.5.2, b'). Finally, (c') is the same as each of (d), (d'), (e), and (e') by (4.5.2, a), (4.5.2, a'), (4.5.2, c), and (4.5.5, d''). $\square$

We further observe that, with the previous notation, the X_f that are affine form a *basis* for the topology of X , and that the canonical morphism u is *dominant* (4.5.2).

Corollary (5.1.3). — *Let X be a quasi-compact prescheme. If there exists a morphism $v : X \rightarrow Y$ from X to some affine scheme Y that is a homeomorphism from X to some open subspace of Y , then X is quasi-affine.*

Proof. There exists a family (g_α) of sections of $\mathcal{O}_Y$ over Y such that the $D(g_\alpha)$ form a basis for the topology of $v(X)$; if $v = (\psi, \theta)$ and we set $f_\alpha = \theta(g_\alpha)$, then we have $X_{f_\alpha} = \psi^{-1}(D(g_\alpha))$ (I, 2.2.4.1), so the X_{f_α} form a basis for the topology of X , and the criterion (5.1.2, d) is satisfied. $\square$

Corollary (5.1.4). — *If X is a quasi-affine scheme, then every invertible $\mathcal{O}_X$ -module is very ample (relative to the canonical morphism), and a fortiori ample.*

Proof. Such a module $\mathcal{L}$ is generated by its sections over X (5.1.2, e), so $\mathcal{L} \otimes \mathcal{O}_X = \mathcal{L}$ is very ample (4.4.8). $\square$

Corollary (5.1.5). — *Let X be a quasi-compact prescheme. If there exists an invertible $\mathcal{O}_X$ -module $\mathcal{L}$ such that $\mathcal{L}$ and $\mathcal{L}^{-1}$ are ample, then X is a quasi-affine scheme.*

Proof. Indeed, $\mathcal{O}_X = \mathcal{L} \otimes \mathcal{L}^{-1}$ is then ample (4.5.7). $\square$

Proposition (5.1.6). — *Let $f : X \rightarrow Y$ be a quasi-compact morphism. Then the following conditions are equivalent.*

- (a) *The morphism f is quasi-affine.*
- (b) *The $\mathcal{O}_Y$ -algebra $f_*(\mathcal{O}_X) = \mathcal{A}(X)$ is quasi-coherent, and the canonical morphism $X \rightarrow \text{Spec}(\mathcal{A}(X))$ corresponding to the identity morphism $\mathcal{A}(X) \rightarrow \mathcal{A}(X)$ (1.2.7) is an open immersion.*
- (b') *The $\mathcal{O}_Y$ -algebra $\mathcal{A}(X)$ is quasi-coherent, and the canonical morphism $X \rightarrow \text{Spec}(\mathcal{A}(X))$ is a homeomorphism from X to some subspace of $\text{Spec}(\mathcal{A}(X))$.*
- (c) *The $\mathcal{O}_X$ -module $\mathcal{O}_X$ is very ample for f .*
- (c') *The $\mathcal{O}_X$ -module $\mathcal{O}_X$ is ample for f .*
- (d) *The morphism f is separated, and, for every quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$, the canonical homomorphism $\sigma : f^*(f_*(\mathcal{F})) \rightarrow \mathcal{F}$ (0, 4.4.3) is surjective.*

Furthermore, whenever f is quasi-affine, every invertible $\mathcal{O}_X$ -module $\mathcal{L}$ is very ample relative to f .

Proof. The equivalence between (a) and (c') follows from the local (on Y) character of the f -ampleness (4.6.4), Definition (5.1.1), and the criterion (5.1.2, c'). The other properties are local on Y [II:96] and thus follow immediately from (5.1.2) and (5.1.4), taking into account the fact that $f_*(\mathcal{F})$ is quasi-coherent whenever f is separated (I, 9.2.2, a). $\square$

Corollary (5.1.7). — *Let $f : X \rightarrow Y$ be a quasi-affine morphism. For every open subset U of Y , the restriction $f^{-1}(U) \rightarrow U$ of f is quasi-affine.*

Corollary (5.1.8). — *Let Y be an affine scheme, and $f : X \rightarrow Y$ a quasi-compact morphism. For f to be quasi-affine, it is necessary and sufficient for X to be a quasi-affine scheme.*

Proof. This is an immediate consequence of (5.1.6) and (4.6.6). $\square$

Corollary (5.1.9). — *Let Y be a quasi-compact scheme or a prescheme whose underlying space is Noetherian, and $f : X \rightarrow Y$ a morphism of finite type. If f is quasi-affine, then there exists a quasi-coherent $\mathcal{O}_Y$ -subalgebra $\mathcal{B}$ of $\mathcal{A}(X) = f_*(\mathcal{O}_X)$ of finite type (I, 9.6.2) such that the morphism $X \rightarrow \text{Spec}(\mathcal{B})$ corresponding to the canonical injection $\mathcal{B} \rightarrow \mathcal{A}(X)$ is an immersion. Further, every quasi-coherent $\mathcal{O}_Y$ -subalgebra $\mathcal{B}'$ of finite type of $\mathcal{A}(X)$ containing $\mathcal{B}$ has the same property.*

Proof. Indeed, $\mathcal{A}(X)$ is the inductive limit of its quasi-coherent $\mathcal{O}_Y$ -subalgebras of finite type (I, 9.6.5); the result is then a particular case of (3.8.4), taking into account the identification of $\text{Spec}(\mathcal{A}(X))$ with $\text{Proj}(\mathcal{A}(X)[T])$ (3.1.7). $\square$

Proposition (5.1.10). —

- (i) A quasi-compact morphism $X \rightarrow Y$ that is a homeomorphism from the underlying space of X to some subspace of the underlying space of Y (so, in particular, any closed immersion) is quasi-affine.
- (ii) The composition of any two quasi-affine morphisms is quasi-affine.
- (iii) If $f : X \rightarrow Y$ is a quasi-affine S -morphism, then $f_{(S')} : X_{(S')} \rightarrow Y_{(S')}$ is a quasi-affine morphism for any extension $S' \rightarrow S$ of the base prescheme.
- (iv) If $f : X \rightarrow Y$ and $g : X' \rightarrow Y'$ are quasi-affine S -morphisms, then $f \times_S g$ is quasi-affine.
- (v) If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are morphisms such that $g \circ f$ is quasi-affine, and if g is separated or the underlying space of X is locally Noetherian, then f is quasi-affine.
- (vi) If f is a quasi-affine morphism, then so is f_{red} .

Proof. Taking into account the criterion (5.1.6, c'), all of (i), (iii), (iv), (v), and (vi) follow immediately from (4.6.13, i bis), (4.6.13, iii), (4.6.13, iv), (4.6.13, v), and (4.6.13, vi) (respectively). To prove (ii), we can restrict to the case where Z is affine, and then the claim follows directly from applying (4.6.13, ii) to $\mathcal{L} = \mathcal{O}_X$ and $\mathcal{K} = \mathcal{O}_Y$. $\square$

Remark (5.1.11). — Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be morphisms such that $X \times_Z Y$ is locally Noetherian. Then the graph immersion $\Gamma_f : X \rightarrow X \times_Z Y$ is quasi-affine, since it is quasi-compact (I, 6.3.5), and since (I, 5.5.12) shows that, in (v), the conclusion still holds true if we remove the hypothesis that g is separated.

Proposition (5.1.12). — Let $f : X \rightarrow Y$ be a quasi-compact morphism, and $g : X' \rightarrow X$ a quasi-affine morphism. If $\mathcal{L}$ is an ample (for f) $\mathcal{O}_X$ -module, then $g^*(\mathcal{L})$ is an ample (for $f \circ g$) $\mathcal{O}_{X'}$ -module.

Proof. Since $\mathcal{O}_{X'}$ is very ample for g , and the question is local on Y (4.6.4), it follows from (4.6.13, ii) that there exists (for Y affine) an integer n such that

$$g^*(\mathcal{L}^{\otimes n}) = (g^*(\mathcal{L}))^{\otimes n}$$

is ample for $f \circ g$, and so $g^*(\mathcal{L})$ is ample for $f \circ g$ (4.6.9). $\square$

5.2. Serre's criterion

Theorem (5.2.1). — (Serre's criterion). Let X be a quasi-compact scheme or a prescheme whose underlying space is Noetherian. The following conditions are equivalent.

- (a) X is an affine scheme.
- (b) There exists a family of elements $f_\alpha \in A = \Gamma(X, \mathcal{O}_X)$ such that the X_{f_α} are affine, and such that the ideal generated by the f_α in A is equal to A itself.
- (c) The functor $\Gamma(X, \mathcal{F})$ is exact in $\mathcal{F}$ on the category of quasi-coherent $\mathcal{O}_X$ -modules, or, in other words, if

$$(*) \quad 0 \longrightarrow \mathcal{F}' \longrightarrow \mathcal{F} \longrightarrow \mathcal{F}'' \longrightarrow 0$$

is an exact sequence of quasi-coherent $\mathcal{O}_X$ -modules, then the sequence

$$0 \longrightarrow \Gamma(X, \mathcal{F}') \longrightarrow \Gamma(X, \mathcal{F}) \longrightarrow \Gamma(X, \mathcal{F}'') \longrightarrow 0$$

is also exact.

- (c') Condition (c) holds for every exact sequence (*) of quasi-coherent $\mathcal{O}_X$ -modules such that $\mathcal{F}$ is isomorphic to an $\mathcal{O}_X$ -submodule of $\mathcal{O}_X^n$ for some finite n .
- (d) $H^1(X, \mathcal{F}) = 0$ for every quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$.
- (d') $H^1(X, \mathcal{I}) = 0$ for every quasi-coherent sheaf of ideals $\mathcal{I}$ of $\mathcal{O}_X$.

Proof. It is evident that (a) implies (b); furthermore, (b) implies that the X_{f_α} cover X , because, by hypothesis, the section 1 is a linear combination of the f_α , and the $D(f_\alpha)$ thus cover $\text{Spec}(A)$. The final claim of (4.5.2) thus implies that $X \rightarrow \text{Spec}(A)$ is an isomorphism.

We know that (a) implies (c) (I, 1.3.11), and (c) trivially implies (c'). We now prove that (c') implies (b). First of all, (c') implies that, for every closed point $x \in X$ and every open neighbourhood U of x , there exists some $f \in A$ such that $x \in X_f \subset U$. Let $\mathcal{I}$ (resp. $\mathcal{I}'$) be the quasi-coherent sheaf of ideals of $\mathcal{O}_X$ defining the closed reduced subprescheme of X that has $X - U$ (resp. $(X - U) \cup \{x\}$) as its underlying space (I, 5.2.1); it is clear that we have $\mathcal{I}' \subset \mathcal{I}$, and that $\mathcal{I}'' = \mathcal{I} / \mathcal{I}'$ is a quasi-coherent $\mathcal{O}_X$ module that has support equal to $\{x\}$, and such that $\mathcal{I}''_x = k(x)$. Hypothesis (c')

applied to the exact sequence $0 \rightarrow \mathcal{J}' \rightarrow \mathcal{J} \rightarrow \mathcal{J}'' \rightarrow 0$ shows that $\Gamma(X, \mathcal{J}) \rightarrow \Gamma(X, \mathcal{J}'')$ is surjective. The section of $\mathcal{J}''$ whose germ at x is 1_x is thus the image of some section $f \in \Gamma(X, \mathcal{J}) \subset \Gamma(X, \mathcal{O}_X)$, and we have, by definition, that $f(x) = 1_x$ and $f(y) = 0$ in $X - U$, which establishes our claim. Now, if U is affine, then so is X_f (I, 1.3.6), so the union of the X_f that are affine ($f \in A$) is an open set Z that contains *all the closed points* of X ; since X is a quasi-compact Kolmogoroff space, we necessarily have $Z = X$ (0, 2.1.3). Because X is quasi-compact, there are a *finite* number of elements $f_i \in A$ ($1 \leq i \leq n$) such that the X_{f_i} are affine and cover X . So consider the homomorphism $\mathcal{O}_X^n \rightarrow \mathcal{O}_X$ defined by the sections f_i (0, 5.1.1); since, for all $x \in X$, at least one of the $(f_i)_x$ is invertible, this homomorphism is *surjective*, and we thus have an exact sequence $0 \rightarrow \mathcal{R} \rightarrow \mathcal{O}_X^n \rightarrow \mathcal{O}_X \rightarrow 0$, where $\mathcal{R}$ is a quasi-coherent $\mathcal{O}_X$ -submodule of $\mathcal{O}_X^n$. It then follows [II:98] from (c') that the corresponding homomorphism $\Gamma(X, \mathcal{O}_X^n) \rightarrow \Gamma(X, \mathcal{O}_X)$ is surjective, which proves (b).

Finally, (a) implies (d) (I, 5.1.9.2), and (d) trivially implies (d'). It remains to show that (d') implies (c'). But if $\mathcal{F}'$ is a quasi-coherent $\mathcal{O}_X$ -submodule of $\mathcal{O}_X^n$, then the filtration $0 \subset \mathcal{O}_X \subset \mathcal{O}_X^2 \subset \dots \subset \mathcal{O}_X^n$ defines a filtration of $\mathcal{F}'$ given by the $\mathcal{F}'_k = \mathcal{F}' \cap \mathcal{O}_X^k$ ($0 \leq k \leq n$), which are quasi-coherent $\mathcal{O}_X$ -modules (I, 4.1.1), and $\mathcal{F}'_{k+1}/\mathcal{F}'_k$ is isomorphic to a quasi-coherent $\mathcal{O}_X$ -submodule of $\mathcal{O}_X^{k+1}/\mathcal{O}_X^k = \mathcal{O}_X$, which is to say, a quasi-coherent sheaf of ideals of $\mathcal{O}_X$. Hypothesis (d') thus implies that $H^1(X, \mathcal{F}'_{k+1}/\mathcal{F}'_k) = 0$; the exact cohomology sequence $H^1(X, \mathcal{F}'_k) \rightarrow H^1(X, \mathcal{F}'_{k+1}) \rightarrow H^1(X, \mathcal{F}'_{k+1}/\mathcal{F}'_k) = 0$ then lets us prove by induction on k that $H^1(X, \mathcal{F}'_k) = 0$ for all k . $\square$

Remark (5.2.1.1). — When X is a *Noetherian* prescheme, we can replace “quasi-coherent” by “coherent” in the statements of (c') and (d'). Indeed, in the proof of the fact that (c') implies (b), $\mathcal{J}$ and $\mathcal{J}'$ are then *coherent* sheaves of ideals, and, furthermore, every quasi-coherent submodule of a coherent module is coherent (I, 6.1.1); whence the conclusion.

Corollary (5.2.2). — *Let $f : X \rightarrow Y$ be a separated quasi-compact morphism. The following conditions are equivalent.*

- (a) *The morphism f is an affine morphism.*
- (b) *The functor f_* is exact on the category of quasi-coherent $\mathcal{O}_X$ -modules.*
- (c) *For every quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$, we have $R^1 f_*(\mathcal{F}) = 0$.*
- (c') *for every quasi-coherent sheaf of ideals $\mathcal{J}$ of $\mathcal{O}_X$, we have $R^1 f_*(\mathcal{J}) = 0$.*

Proof. All these conditions are local on Y , by definition of the functor $R^1 f_*$ (T, 3.7.3), and so we can assume that Y is affine. If f is affine, then X is affine, and property (b) is nothing more than (I, 1.6.4). Conversely, we now show that (b) implies (a): for every quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$, we have that $f_*(\mathcal{F})$ is a quasi-coherent $\mathcal{O}_Y$ -module (I, 9.2.2, a). By hypothesis, the functor $f_*(\mathcal{F})$ is exact in $\mathcal{F}$, and the functor $\Gamma(Y, \mathcal{G})$ is exact in $\mathcal{G}$ (in the category of quasi-coherent $\mathcal{O}_Y$ -modules) because Y is affine (I, 1.3.11); so $\Gamma(Y, f_*(\mathcal{F})) = \Gamma(X, \mathcal{F})$ is exact in $\mathcal{F}$, which proves our claim, by (5.2.1, c).

If f is affine, then $f^{-1}(U)$ is affine for every affine open subset U of Y (1.3.2), and so $H^1(f^{-1}(U), \mathcal{F}) = 0$ (5.2.1, d), which, by definition, implies that $R^1 f_*(\mathcal{F}) = 0$. Finally, suppose that condition (c') is satisfied; the exact sequence of terms of low degree in the Leray spectral sequence (G, II, 4.17.1 and I, 4.5.1) gives, in particular, the exact sequence

$$0 \longrightarrow H^1(Y, f_*(\mathcal{J})) \longrightarrow H^1(X, \mathcal{J}) \longrightarrow H^0(Y, R^1 f_*(\mathcal{J})).$$

Since Y is affine, and $f_*(\mathcal{J})$ quasi-coherent (I, 9.2.2, a), we have that $H^1(Y, f_*(\mathcal{J})) = 0$ (5.2.1); hypothesis (c') thus implies that $H^1(X, \mathcal{J}) = 0$, and we conclude, by (5.2.1), that X is an affine scheme. $\square$

Corollary (5.2.3). — *If $f : X \rightarrow Y$ is an affine morphism, then, for every quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$, the canonical homomorphism $H^1(Y, f_*(\mathcal{F})) \rightarrow H^1(X, \mathcal{F})$ is bijective.*

[II:99]

Proof. We have the exact sequence

$$0 \longrightarrow H^1(Y, f_*(\mathcal{F})) \longrightarrow H^1(X, \mathcal{F}) \longrightarrow H^0(Y, R^1 f_*(\mathcal{F}))$$

of terms of low degree in the Leray spectral sequence, and the conclusion follows from (5.2.2). $\square$

Remark (5.2.4). — In Chapter III, §1, we prove that, if X is affine, then we have $H^i(X, \mathcal{F}) = 0$ for all $i > 0$ and all quasi-coherent $\mathcal{O}_X$ -modules $\mathcal{F}$.

5.3. Quasi-projective morphisms

Definition (5.3.1). — We say that a morphism $f : X \rightarrow Y$ is *quasi-projective*, or that X (considered as a Y -prescheme via f) is *quasi-projective over Y* , or that X is a *quasi-projective Y -scheme*, if f is of finite type and there exists an invertible f -ample $\mathcal{O}_X$ -module.

We note that this notion is *not local on Y* : the counterexamples of Nagata [26] and Hironaka show that, even if X and Y are non-singular algebraic schemes over an algebraically closed field, every point of Y can have an affine neighbourhood U such that $f^{-1}(U)$ is quasi-projective over U , without f being quasi-projective.

We note that a quasi-projective morphism is necessarily *separated* (4.6.1). When Y is quasi-compact, it is equivalent to say either that f is quasi-projective, or that f is of finite type and there exists a *very ample* (relative to f) $\mathcal{O}_X$ -module ((4.6.2) and (4.6.11)). Further:

Proposition (5.3.2). — Let Y be a quasi-compact scheme or a prescheme whose underlying space is Noetherian, and let X be a Y -prescheme. The following conditions are equivalent.

- (a) X is a quasi-projective Y -scheme.
- (b) X is of finite type over Y , and there exists some quasi-coherent $\mathcal{O}_Y$ -module $\mathcal{E}$ of finite type such that X is Y -isomorphic to a subprescheme of $\mathbf{P}(\mathcal{E})$.
- (c) X is of finite type over Y , and there exists some quasi-coherent graded $\mathcal{O}_Y$ -algebra $\mathcal{S}$ such that $\mathcal{S}_1$ is of finite type and generates $\mathcal{S}$, and such that X is Y -isomorphic to a subprescheme induced on some everywhere-dense open subset of $\text{Proj}(\mathcal{S})$.

Proof. This follows immediately from the previous remark and from (4.4.3), (4.4.6), and (4.4.7). $\square$

We note that, whenever Y is a Noetherian prescheme, we can, in conditions (b) and (c) of (5.3.2), remove the hypothesis that X is of finite type over Y , since this is then automatically satisfied (I, 6.3.5).

Corollary (5.3.3). — Let Y be a quasi-compact scheme such that there exists an ample $\mathcal{O}_Y$ -module $\mathcal{L}$ (4.5.3). For a Y -scheme X to be quasi-projective, it is necessary and sufficient for it to be of finite type over Y and also isomorphic to a Y -subscheme of a projective bundle of the form $\mathbf{P}_Y^r$.

Proof. If $\mathcal{E}$ is a quasi-coherent $\mathcal{O}_Y$ -module of finite type, then $\mathcal{E}$ is isomorphic to a quotient of an $\mathcal{O}_Y$ -module $\mathcal{L}^{\otimes(-n)} \otimes_{\mathcal{O}_Y} \mathcal{O}_Y^k$ (4.5.5), and so $\mathbf{P}(\mathcal{E})$ is isomorphic to a closed subscheme of $\mathbf{P}_Y^{k-1}$ ((4.1.2) and (4.1.4)). $\square$

Proposition (5.3.4). —

- (i) A quasi-affine morphism of finite type (and, in particular, a quasi-compact immersion, or an affine morphism of finite type) is quasi-projective.
- (ii) If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are quasi-projective, and if Z is quasi-compact, then $g \circ f$ is quasi-projective. [III:100]
- (iii) If $f : X \rightarrow Y$ is a quasi-projective S -morphism, then $f_{(S')} : X_{(S')} \rightarrow Y_{(S')}$ is quasi-projective for every extension $S' \rightarrow S$ of the base prescheme.
- (iv) If $f : X \rightarrow Y$ and $g : X' \rightarrow Y'$ are quasi-projective S -morphisms, then $f \times_S g$ is quasi-projective.
- (v) If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are morphisms such that $g \circ f$ is quasi-projective, and if g is separated or X locally Noetherian, then f is quasi-projective.
- (vi) If f is a quasi-projective morphism, then so is f_{red} .

Proof. (i) follows from (5.1.6) and (5.1.10, i). The other claims are immediate consequences of Definition (5.3.1), of the properties of morphisms of finite type (I, 6.3.4), and of (4.6.13). $\square$

Remark (5.3.5). — We note that it can happen that f_{red} is quasi-projective without f being quasi-projective, even if we assume that Y is the spectrum of an algebra of finite rank over $\mathbb{C}$ and that f is proper.

Corollary (5.3.6). — *If X and X' are quasi-projective Y -schemes, then $X \sqcup X'$ is a quasi-projective Y -scheme.*

Proof. This follows from (4.6.18). □

5.4. Proper morphisms and universally closed morphisms

Definition (5.4.1). — We say that a morphism of preschemes $f : X \rightarrow Y$ is *proper* if it satisfies the following two conditions:

- (a) f is separated and of finite type; and
- (b) for every prescheme Y' and every morphism $Y' \rightarrow Y$, the projection $f_{(Y')} : X \times_Y Y' \rightarrow Y'$ is a closed morphism (I, 2.2.6).

When this is the case, we also say that X (considered as a Y -prescheme with structure morphism f) is *proper over Y* .

It is immediate that conditions (a) and (b) are *local* on Y . To show that the image of a closed subset Z of $X \times_Y Y'$ under the projection $q : X \times_Y Y' \rightarrow Y'$ is closed in Y' , it suffices to see that $q(Z) \cap U'$ is closed in U' for every affine open subset U' of Y' ; since $q(Z) \cap U' = q(Z \cap q^{-1}(U'))$, and since $q^{-1}(U')$ can be identified with $X \times_Y U'$ (I, 4.4.1), we see that to satisfy condition (b) of Definition (5.4.1), we can restrict to the case where Y is an *affine* scheme. We will see below (5.6.3) that, if Y is locally Noetherian, then we can even restrict to proving (b) in the case where Y' is of finite type over Y .

It is clear that every proper morphism is *closed*.

Proposition (5.4.2). —

- (i) *A closed immersion is a proper morphism.*
- (ii) *The composition of two proper morphisms is proper.*
- (iii) *If X and Y are S -preschemes, and $f : X \rightarrow Y$ a proper S -morphism, then $f_{(S')} : X_{(S')} \rightarrow Y_{(S')}$ is proper for every extension $S' \rightarrow S$ of the base prescheme.*
- (iv) *If $f : X \rightarrow Y$ and $g : X' \rightarrow Y'$ are proper S -morphisms, then $f \times_S g : X \times_S X' \rightarrow Y \times_S Y'$ is a proper S -morphism.*

[II:101]

Proof. It suffices to prove (i), (ii), and (iii) (I, 3.5.1). In each of the three cases, verifying condition (a) of (5.4.1) follows from previous results ((I, 5.5.1) and (I, 6.3.4)); it remains to verify condition (b). It is immediate in case (i), because if $X \rightarrow Y$ is a closed immersion, then so is $X \times_Y Y' \rightarrow Y \times_Y Y' = Y'$ ((I, 4.3.2) and (3.3.3)). To prove (ii), consider two proper morphisms $X \rightarrow Y$ and $Y \rightarrow Z$, and a morphism $Z' \rightarrow Z$. We can write $X \times_Z Z' = X \times_Y (Y \times_Z Z')$ (I, 3.3.9.1), and so the projection $X \times_Z Z' \rightarrow Z'$ factors as $X \times_Y (Y \times_Z Z') \rightarrow Y \times_Z Z' \rightarrow Z'$. Taking the initial remark into account, (ii) follows from the fact that the composition of two closed morphisms is closed. Finally, for every morphism $S' \rightarrow S$, we can identify $X_{(S')}$ with $X \times_Y Y_{(S')}$ (I, 3.3.11); for every morphism $Z \rightarrow Y_{(S')}$, we can write

$$X_{(S')} \times_{Y_{(S')}} Z = (X \times_Y Y_{(S')}) \times_{Y_{(S')}} Z = X \times_Y Z;$$

since by hypothesis $X \times_Y Z \rightarrow Z$ is closed, this proves (iii). □

Corollary (5.4.3). — *Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be morphisms such that $g \circ f$ is proper.*

- (i) *If g is separated, then f is proper.*
- (ii) *If g is separated and of finite type, and if f is surjective, then g is proper.*

Proof. (i) follows from (5.4.2) by the general procedure (I, 5.5.12). To prove (ii), we need only verify that condition (b) of Definition (5.4.1) is satisfied. For every morphism $Z' \rightarrow Z$, the diagram

$$\begin{array}{ccc} X \times_Z Z' & \xrightarrow{f \times 1_{Z'}} & Y \times_Z Z' \\ & \searrow p & \downarrow p' \\ & & Z' \end{array}$$

(where p and p' are the projections) commutes (I, 3.2.1); furthermore, $f \times 1_{Z'}$ is surjective because f is surjective (I, 3.5.2), and p is a closed morphism by hypothesis. Every closed subset F of $Y \times_Z Z'$ is thus the image under $f \times 1_{Z'}$ of some closed subset E of $X \times_Z Z'$, so $p'(F) = p(E)$ is closed in Z' by hypothesis, whence the corollary. $\square$

Corollary (5.4.4). — *If X is a proper prescheme over Y , and $\mathcal{S}$ a quasi-coherent $\mathcal{O}_Y$ -algebra, then every Y -morphism $f : X \rightarrow \text{Proj}(\mathcal{S})$ is proper (and a fortiori closed).*

Proof. The structure morphism $p : \text{Proj}(\mathcal{S}) \rightarrow Y$ is separated, and $p \circ f$ is proper by hypothesis. $\square$

Corollary (5.4.5). — *Let $f : X \rightarrow Y$ be a separated morphism of finite type. Let $(X_i)_{1 \leq i \leq n}$ (resp. $(Y_i)_{1 \leq i \leq n}$) be a finite family of closed subpreschemes of X (resp. Y), and j_i (resp. h_i) the canonical injection $X_i \rightarrow X$ (resp. $Y_i \rightarrow Y$). Suppose that the underlying space of X is the union of the X_i , and that, for all i , there is a morphism $f_i : X_i \rightarrow Y_i$, such that the diagram*

$$\begin{array}{ccc} X_i & \xrightarrow{f_i} & Y_i \\ j_i \downarrow & & \downarrow h_i \\ X & \xrightarrow{f} & Y \end{array}$$

commutes. Then, for f to be proper, it is necessary and sufficient for all of the f_i to be proper.

[II:102]

Proof. If f is proper, then so is $f \circ j_i$, because j_i is a closed immersion (5.4.2); since h_i is a closed immersion, and thus a separated morphism, f_i is proper, by (5.4.3). Conversely, suppose that all of the f_i are proper, and consider the prescheme Z given by the sum of the X_i ; let u be the morphism $Z \rightarrow X$ which reduces to j_i on each X_i . The restriction of $f \circ u$ to each X_i is equal to $f \circ j_i = h_i \circ f_i$, and is thus proper, because both the h_i and the f_i are (5.4.2); it then follows immediately from Definition (5.4.1) that $f \circ u$ is proper. But since by hypothesis u is surjective, we conclude that f is proper by (5.4.3). $\square$

Corollary (5.4.6). — *Let $f : X \rightarrow Y$ be a separated morphism of finite type; for f to be proper, it is necessary and sufficient for $f_{\text{red}} : X_{\text{red}} \rightarrow Y_{\text{red}}$ to be proper.*

Proof. This is a particular case of (5.4.5), with $n = 1$, $X_1 = X_{\text{red}}$, and $Y_1 = Y_{\text{red}}$ (I, 5.1.5). $\square$

(5.4.7). If X and Y are Noetherian preschemes, and $f : X \rightarrow Y$ a separated morphism of finite type, then we can, to show that f is proper, restrict to the case of *dominant* morphisms and *integral* preschemes. Indeed, let X_i ($1 \leq i \leq n$) be the (finitely many) irreducible components of X , and consider, for each i , the unique closed reduced subprescheme of X that has X_i as its underlying space, which we again denote by X_i (I, 5.2.1). Let Y_i be the unique closed reduced subprescheme of Y that has $f(X_i)$ as its underlying space. If g_i (resp. h_i) is the injection morphism $X_i \rightarrow X$ (resp. $Y_i \rightarrow Y$), then we conclude that $f \circ g_i = h_i \circ f_i$, where f_i is a dominant morphism $X_i \rightarrow Y_i$ (I, 5.2.2); we are then under the right conditions to apply (5.4.5), and for f to be proper, it is necessary and sufficient for all the f_i to be proper.

Corollary (5.4.8). — *Let X and Y be separated S -preschemes of finite type over S , and $f : X \rightarrow Y$ an S -morphism. For f to be proper, it is necessary and sufficient that, for every S -prescheme S' , the morphism $f \times_S 1_{S'} : X \times_S S' \rightarrow Y \times_S S'$ be closed.*

Proof. First note that, if $g : X \rightarrow S$ and $h : Y \rightarrow S$ are the structure morphisms, then we have, by definition, $g = h \circ f$, and so f is separated and of finite type ((I, 5.5.1) and (I, 6.3.4)). If f is proper, then so is $f \times_S 1_{S'}$ (5.4.2); a fortiori, $f \times_S 1_{S'}$ is closed. Conversely, suppose that the conditions of the statement are satisfied, and let Y' be a Y -prescheme; Y' can also be considered as an S -prescheme, and since $Y \rightarrow S$ is separated, $X \times_Y Y'$ can be identified with a closed subprescheme of $X \times_S Y'$

(I, 5.4.2). In the commutative diagram

$$\begin{array}{ccc} X \times_Y Y' & \xrightarrow{f \times 1_{Y'}} & Y \times_Y Y' = Y' \\ \downarrow & & \downarrow \\ X \times_S Y' & \xrightarrow{f \times 1_{Y'}} & Y \times_S Y', \end{array}$$

the vertical arrows are closed immersions; it thus immediately follows that if $f \times 1_{Y'}$ over S is a closed morphism, then so is the base change $f \times 1_{Y'}$ over Y . $\square$

Remark (5.4.9). — We say that a morphism $f : X \rightarrow Y$ is *universally closed* if it satisfies condition (b) of Definition (5.4.1). The reader will observe that, ^[II-103] in (5.4.2) to (5.4.8), we can replace every occurrence of “proper” with “universally closed” without changing the validity of the results (and in the hypotheses of (5.4.3), (5.4.5), (5.4.6), and (5.4.8), we can omit the finiteness conditions).

(5.4.10). Let $f : X \rightarrow Y$ be a morphism of finite type. We say that a closed subset Z of X is *proper on Y* (or *Y -proper*, or *proper for f*) if the restriction of f to a closed subprescheme of X , with underlying space Z (I, 5.2.1), is *proper*. Since this restriction is then separated, it follows from (5.4.6) and (I, 5.5.1, vi) that the preceding property *does not depend* on the closed subprescheme of X that has Z as its underlying space. If $g : X' \rightarrow X$ is a *proper* morphism, then $g^{-1}(Z)$ is a *proper* subset of X' : if T is a subprescheme of X that has Z as its underlying space, it suffices to note that the restriction of g to the closed subprescheme $g^{-1}(T)$ of X' is a proper morphism $g^{-1}(T) \rightarrow T$, by (5.4.2, iii), and to then apply (5.4.2, ii). Further, if X'' is a Y -scheme of finite type, and $u : X \rightarrow X''$ a Y -morphism, then $u(Z)$ is a *proper* subset of X'' ; indeed, let us take T to be the closed reduced subprescheme of X having Z as its underlying space; then the restriction of f to T is proper, and thus so is the restriction of u to T (5.4.3, i), thus $u(Z)$ is closed in X'' ; let T'' be a closed subprescheme of X'' having $u(Z)$ as its underlying space (I, 5.2.1), such that $u|_T$ factors as $T \xrightarrow{v} T'' \xrightarrow{j} X''$, where j is the canonical injection (I, 5.2.2), and v is thus proper and surjective (5.4.5); if g is the restriction to T'' of the structure morphism $X'' \rightarrow Y$, then g is separated and of finite type, and we have that $f|_T = g \circ v$; it thus follows from (5.4.3, ii) that g is proper, whence our assertion.

It follows, in particular, from these remarks that, if Z is a Y -proper subset of X , then

- (1) for every closed subprescheme X' of X , $Z \cap X'$ is a Y -proper subset of X' ; and
- (2) if X is a subprescheme of a Y -scheme of finite type X'' , then Z is also a Y -proper subset of X'' (and so, in particular, is *closed in X''*).

5.5. Projective morphisms

Proposition (5.5.1). — *Let X be a Y -prescheme. The following conditions are equivalent.*

- (a) *X is Y -isomorphic to a closed subprescheme of a projective bundle $\mathbf{P}(\mathcal{E})$, where $\mathcal{E}$ is a quasi-coherent $\mathcal{O}_Y$ -module of finite type.*
- (b) *There exists a quasi-coherent graded $\mathcal{O}_Y$ -algebra $\mathcal{S}$ such that $\mathcal{S}_1$ is of finite type and generates $\mathcal{S}$, and such that X is Y -isomorphic to $\text{Proj}(\mathcal{S})$.*

Proof. Condition (a) implies (b), by (3.6.2, ii): if $\mathcal{I}$ is a quasi-coherent graded sheaf of ideals of $\mathbf{S}(\mathcal{E})$, then the quasi-coherent graded $\mathcal{O}_Y$ -algebra $\mathcal{S} = \mathbf{S}(\mathcal{E})/\mathcal{I}$ is generated by $\mathcal{S}_1$, and $\mathcal{S}_1$, the canonical image of $\mathcal{E}$, is an $\mathcal{O}_Y$ -module of finite type. Condition (b) implies (a) by (3.6.2) applied to the case where $\mathcal{M} \rightarrow \mathcal{S}_1$ is the identity map. $\square$

[II:104]

Definition (5.5.2). — We say that a Y -prescheme X is *projective* over Y , or is a *projective Y -scheme*, if it satisfies either of the (equivalent) conditions (a) and (b) of (5.5.1). We say that a morphism $f : X \rightarrow Y$ is *projective* if it makes X a projective Y -scheme.

It is clear that if $f : X \rightarrow Y$ is projective, then there exists a *very ample* (relative to f) $\mathcal{O}_X$ -module (4.4.2).

Theorem (5.5.3). —

- (i) Every projective morphism is quasi-projective and proper.
- (ii) Conversely, let Y be a quasi-compact scheme or a prescheme whose underlying space is Noetherian; then every morphism $f : X \rightarrow Y$ that is quasi-projective and proper is projective.

Proof.

- (i) It is clear that if $f : X \rightarrow Y$ is projective, then it is of finite type and quasi-projective (thus, in particular, separated); furthermore, it follows immediately from (5.5.1, b) and (3.5.3) that if f is projective, then so is $f \times_Y 1_{Y'} : X \times_Y Y' \rightarrow Y'$ for every morphism $Y' \rightarrow Y$. To show that f is universally closed, it is thus enough to show that a projective morphism f is closed. Since the question is local on Y , we can suppose that $Y = \text{Spec}(A)$, thus (5.5.1) $X = \text{Proj}(S)$, where S is a graded A -algebra generated by a finite number of elements of S_1 . For all $y \in Y$, the fibre $f^{-1}(y)$ can be identified with $\text{Proj}(S \otimes_A k(y))$ (I, 3.6.1), and so also with $\text{Proj}(S \otimes_A k(y))$ (2.8.10); so $f^{-1}(y)$ is empty if and only if $S \otimes_A k(y)$ satisfies condition (TN) (2.7.4), or, in other words, if $S_n \otimes_A k(y) = 0$ for sufficiently large n . But since $(S_n)_y$ is an $\mathcal{O}_y$ -module of finite type, the preceding condition implies that $(S_n)_y = 0$ for sufficiently large n , by Nakayama's lemma. If $\mathfrak{a}_n$ is the annihilator in A of the A -module S_n , then the preceding condition also implies that $\mathfrak{a}_n \not\subset \mathfrak{j}_y$ for sufficiently large n (0, 1.7.4). But since $S_n S_1 = S_{n+1}$, by hypothesis, we have that $\mathfrak{a}_n \subset \mathfrak{a}_{n+1}$, and if $\mathfrak{a}$ is the union of the $\mathfrak{a}_n$, then we see that $f(X) = V(\mathfrak{a})$, which proves that $f(X)$ is closed in Y . If now X' is an arbitrary closed subset of X , then there exists a closed subprescheme of X that has X' as its underlying space (I, 5.2.1), and it is clear (5.5.1, a) that the morphism $X' \rightarrow X \xrightarrow{f} Y$ is projective, and so $f(X')$ is closed in Y .
- (ii) The hypothesis on Y and the fact that f is quasi-projective implies the existence of a quasi-coherent $\mathcal{O}_Y$ -module $\mathcal{E}$ of finite type, as well as a Y -immersion $j : X \rightarrow \mathbf{P}(\mathcal{E})$ (5.3.2). But since f is proper, j is closed, by (5.4.4), and so f is projective.

□

Remark (5.5.4). —

- (i) Let $f : X \rightarrow Y$ be a morphism such that f is proper, such that there exists a *very ample* (relative to f) $\mathcal{O}_X$ -module $\mathcal{L}$, and such that the quasi-coherent $\mathcal{O}_Y$ -module $\mathcal{E} = f_*(\mathcal{L})$ is of finite type. Then f is a *projective* morphism: indeed (4.4.4), there is then a Y -immersion $r : X \rightarrow \mathbf{P}(\mathcal{E})$, and, since f is proper, r is a *closed* immersion (5.4.4). We will see in Chapter III, §3, that when Y is *locally Noetherian*, the third condition above ($\mathcal{E}$ being of finite type) is a consequence of the first two, and so the first two conditions *characterise*, in this case, the projective morphisms, and if Y is quasi-compact, then we can replace the second condition (the existence of a very ample (relative to f) $\mathcal{O}_X$ -module $\mathcal{L}$) by the hypothesis that there exists an *ample* (relative to f) $\mathcal{O}_X$ -module (4.6.11).
- (ii) Let Y be a quasi-compact scheme such that there exists an ample $\mathcal{O}_Y$ -module. For a Y -scheme X to be *projective*, it is necessary and sufficient for it to be Y -isomorphic to a *closed* Y -subscheme of a projective bundle of the form $\mathbf{P}_Y^r$. The condition is clearly sufficient. [III:105] Conversely, if X is projective over Y , then it is quasi-projective, and so there exists a Y -immersion j of X into some $\mathbf{P}_Y^r$ (5.3.3) that is *closed*, by (5.4.4) and (5.5.3).
- (iii) The argument of (5.5.3) shows that, for every prescheme Y and every integer $r \geq 0$, the structure morphism $\mathbf{P}_Y^r \rightarrow Y$ is *surjective*, because if we set $\mathcal{S} = \mathbf{S}_{\mathcal{O}_Y}(\mathcal{O}_Y^{r+1})$, then we evidently have $\mathcal{S}_y = \mathbf{S}_{k(y)}(k(y)^{r+1})$ (1.7.3), and so $(\mathcal{S}_n)_y \neq 0$ for every $y \in Y$ and every $n \geq 0$.
- (iv) It follows from the examples of Nagata [26] that there exist proper morphisms that are not quasi-projective.

Proposition (5.5.5). —

- (i) A closed immersion is a projective morphism.
- (ii) If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are projective morphisms, and if Z is a quasi-compact scheme or a prescheme whose underlying space is Noetherian, then $g \circ f$ is projective.

- (iii) If $f : X \rightarrow Y$ is a projective S -morphism, then $f_{(S')} : X_{(S')} \rightarrow Y_{(S')}$ is projective for every extension $S' \rightarrow S$ of the base prescheme.
- (iv) If $f : X \rightarrow Y$ and $g : X' \rightarrow Y'$ are projective S -morphisms, then so is $f \times_S g$.
- (v) If $g \circ f$ is a projective morphism, and if g is separated, then f is projective.
- (vi) If f is projective, then so is f_{red} .

Proof. (i) follows immediately from (3.1.7). We have to show (iii) and (iv) separately, because of the restriction introduced on Z in (ii) (cf. (I, 3.5.1)). To show (iii), we restrict to the case where $S = Y$ (I, 3.3.11), and the claim then immediately follows from (5.5.1, b) and (3.5.3). To show (iv), we are immediately led to the case where $X = \mathbf{P}(\mathcal{E})$ and $X' = \mathbf{P}(\mathcal{E}')$, where $\mathcal{E}$ (resp. $\mathcal{E}'$) is a quasi-coherent $\mathcal{O}_Y$ -module (resp. quasi-coherent $\mathcal{O}_{Y'}$ -module) of finite type. Let p and p' be the canonical projections of $T = Y \times_S Y'$ to Y and Y' (respectively); by (4.1.3.1), we have $\mathbf{P}(p^*(\mathcal{E})) = \mathbf{P}(\mathcal{E}) \times_Y T$ and $\mathbf{P}(p'^*(\mathcal{E}')) = \mathbf{P}(\mathcal{E}') \times_{Y'} T$; whence

$$\begin{aligned} \mathbf{P}(p^*(\mathcal{E})) \times_T \mathbf{P}(p'^*(\mathcal{E}')) &= (\mathbf{P}(\mathcal{E}) \times_Y T) \times_T (T \times_{Y'} \mathbf{P}(\mathcal{E}')) \\ &= \mathbf{P}(\mathcal{E}) \times_Y (T \times_{Y'} \mathbf{P}(\mathcal{E}')) = \mathbf{P}(\mathcal{E}) \times_S \mathbf{P}(\mathcal{E}') \end{aligned}$$

by replacing T with $Y \times_S Y'$, and using (I, 3.3.9.1). But $p^*(\mathcal{E})$ and $p'^*(\mathcal{E}')$ are of finite type over T (0, 5.2.4), and thus so is $p^*(\mathcal{E}) \otimes_{\mathcal{O}_T} p'^*(\mathcal{E}')$; since $\mathbf{P}(p^*(\mathcal{E})) \times_T \mathbf{P}(p'^*(\mathcal{E}'))$ can be identified with a closed subprescheme of $\mathbf{P}(p^*(\mathcal{E}) \otimes_{\mathcal{O}_T} p'^*(\mathcal{E}'))$ (4.3.3), this proves (iv). To show (v) and (vi), we can apply (I, 5.5.13), because every closed subprescheme of a projective Y -scheme is a projective Y -scheme, by (5.5.1, a).

It remains to prove (ii); by the hypothesis on Z , this follows from (5.5.3), (5.3.4, ii), and (5.4.2, ii). $\square$

Proposition (5.5.6). — *If X and X' are projective Y -schemes, then $X \sqcup X'$ is a projective Y -scheme.*

Proof. This is an evident consequence of (5.5.2) and (4.3.6). $\square$

Proposition (5.5.7). — *Let X be a projective Y -scheme, and $\mathcal{L}$ a Y -ample $\mathcal{O}_X$ -module; then, for every section f of $\mathcal{L}$ over X , X_f is affine over Y .*

[II:106]

Proof. Since the question is local on Y , we can assume that $Y = \text{Spec}(A)$; furthermore, $X_{f^{\otimes n}} = X_f$, so by replacing $\mathcal{L}$ with some suitable $\mathcal{L}^{\otimes n}$, we can assume that $\mathcal{L}$ is very ample relative to the structure morphism $q : X \rightarrow Y$ (4.6.11). The canonical homomorphism $\sigma : q^*(q_*(\mathcal{L})) \rightarrow \mathcal{L}$ is thus surjective, and the corresponding morphism

$$r = r_{\mathcal{L}, \sigma} : X \longrightarrow P = \mathbf{P}(q_*(\mathcal{L}))$$

is an immersion such that $\mathcal{L} = r^*(\mathcal{O}_P(1))$ (4.4.4); furthermore, since X is proper over Y , the immersion r is closed (5.4.4). But by definition, $f \in \Gamma(Y, q_*(\mathcal{L}))$, and σ^b is the identity of $q_*(\mathcal{L})$; it then follows from Equation (3.7.3.1) that we have $X_f = r^{-1}(D_+(f))$; so X_f is a closed subprescheme of the affine scheme $D_+(f)$, and is thus also an affine scheme. $\square$

In the particular case where $Y = X$, we obtain (taking (4.6.13, i) into account) the following corollary, whose direct proof is immediate anyway:

Corollary (5.5.8). — *Let X be a prescheme, and $\mathcal{L}$ an invertible $\mathcal{O}_X$ -module. For every section f of $\mathcal{L}$ over X , X_f is affine over X (and thus also an affine scheme whenever X is an affine scheme).*

5.6. Chow's lemma

Theorem (5.6.1). — (Chow's lemma). *Let S be a prescheme, and X an S -scheme of finite type. Suppose that one of the following conditions is satisfied:*

- (a) S is Noetherian;
- (b) S is a quasi-compact scheme, and X has a finite number of irreducible components.

Under these hypotheses,

- (i) *there exists a quasi-projective S -scheme X' , and an S -morphism $f : X' \rightarrow X$ that is both projective and surjective;*
- (ii) *we can take X' and f to be such that there exists an open subset $U \subset X$ for which $U' = f^{-1}(U)$ is dense in X' , and for which the restriction of f to U' is an isomorphism $U' \simeq U$; and*
- (iii) *if X is reduced (resp. irreducible, integral), then we can assume that X' is reduced (resp. irreducible, integral).*

Proof. The proof proceeds in multiple steps.

- (A) We can first restrict to the case where X is *irreducible*. Indeed, in hypothesis (a), X is Noetherian, and so, under either hypothesis, the irreducible components X_i of X are finite in number. If the theorem is shown to be true for each of the closed reduced preschemes of X having the X_i as their underlying spaces, and if X'_i and $f_i : X'_i \rightarrow X_i$ are the prescheme and the morphism corresponding to X_i (respectively), then the prescheme X' given by the sum of the X'_i , and the morphism $f : X' \rightarrow X$ whose restriction to each X'_i is $j_i \circ f_i$ (where j_i is the canonical injection $X_i \rightarrow X$) satisfy the conclusion of the theorem. It is immediate that X' is reduced if all of the X'_i are; furthermore, we can satisfy (ii) by taking U to be the union of the sets $U_i \cap \mathbb{C} \left(\bigcup_{j \neq i} X_j \right)$. Finally, since the X'_i are quasi-projective over S , so is X' ^[III:107] (5.3.6); similarly, the morphisms $X'_i \rightarrow X$ are projective by (5.5.5, i) and (5.5.5, ii), and so f is projective (5.5.6), and is clearly surjective, by definition.
- (B) Now suppose that X is *irreducible*. Since the structure morphism $r : X \rightarrow S$ is of finite type, there exists a finite cover (S_i) of S by affine open subsets, and for each i there is a finite cover (T_{ij}) of $r^{-1}(S_i)$ by affine open subsets, and the morphisms $T_{ij} \rightarrow S_i$ are of finite type, and so quasi-projective (5.3.4, i); since in both hypotheses (a) and (b) the immersion $S_i \rightarrow S$ is quasi-compact, it is also quasi-projective (5.3.4, i), and so the restriction of r to T_{ij} is a quasi-projective morphism (5.3.4, ii). Denote the T_{ij} by U_k ($1 \leq k \leq n$). There exists, for each index k , an open immersion $\phi_k : U_k \rightarrow P_k$, where P_k is projective over S ((5.3.2) and (5.5.2)). Let $U = \bigcap_k U_k$; since X is irreducible, and the U_k nonempty, U is nonempty, and thus dense in X ; the restrictions of the ϕ_k to U define a morphism

$$\phi : U \longrightarrow P = P_1 \times_S P_2 \times_S \cdots \times_S P_n$$

such that the diagrams

$$(5.6.1.1) \quad \begin{array}{ccc} U & \xrightarrow{\phi} & P \\ j_k \downarrow & & \downarrow p_k \\ U_k & \xrightarrow{\phi_k} & P_k \end{array}$$

commute, where j_k is the canonical injection $U \rightarrow U_k$, and p_k the canonical projection $P \rightarrow P_k$. If j is the canonical injection $U \rightarrow X$, then the morphism $\psi = (j, \phi)_S : U \rightarrow X \times_S P$ is an *immersion* (I, 5.3.14). In hypothesis (a), $X \times_S P$ is locally Noetherian ((3.4.1), (I, 6.3.7), and (I, 6.3.8)); in hypothesis (b), $X \times_S P$ is a quasi-compact scheme ((I, 5.5.1) and (I, 6.6.4)); in both cases, the *closure* X' in $X \times_S P$ of the subscheme Z associated to ψ (and so with underlying space $\psi(U)$) exists, and ψ factors as

$$(5.6.1.2) \quad \psi : U \xrightarrow{\psi'} X' \xrightarrow{h} X \times_S P$$

where ψ' is an *open immersion* and h a *closed immersion* (I, 9.5.10). Let $q_1 : X \times_S P \rightarrow X$ and $q_2 : X \times_S P \rightarrow P$ be the canonical projections; we set

$$(5.6.1.3) \quad f : X' \xrightarrow{h} X \times_S P \xrightarrow{q_1} X,$$

$$(5.6.1.4) \quad g : X' \xrightarrow{h} X \times_S P \xrightarrow{q_2} P.$$

We will see that X' and f satisfy the conclusion of the theorem.

- (C) First we show that f is *projective* and *surjective*, and that the restriction of f to $U' = f^{-1}(U)$ is an *isomorphism* from U' to U . Since the P_k are projective over S , so is P (5.5.5, iv), and so $X \times_S P$ is projective over X (5.5.5, iii), and thus so is X' , which is a closed subscheme of $X \times_S P$. Furthermore, we have $f \circ \psi' = q_1 \circ (h \circ \psi') = q_1 \circ \psi = j$, so $f(X')$ contains the open everywhere-dense subset U of X ; but f is a *closed* morphism (5.5.3), so $f(X') = X$. Now note that $q_1^{-1}(U) = U \times_S P$ is induced on an open subset of $X \times_S P$, and, by definition, the prescheme $U' = h^{-1}(U \times_S P)$ is induced by X' on the open subset U' ; it is thus the closure relative to [II:108] $U \times_S P$ of the prescheme Z (I, 9.5.8). But the immersion ψ factors as $U \xrightarrow{\Gamma_\phi} U \times_S P \xrightarrow{j \times 1} X \times_S P$, and since P is separated over S , the graph morphism Γ_ϕ is a closed immersion (I, 5.4.3), and so Z is a *closed* subscheme of $U \times_S P$, whence $U' = Z$. Since ψ is an immersion, the restriction of f to U' is an isomorphism onto U , and the inverse of ψ' ; finally, by the definition of X' , U' is dense in X' .
- (D) We now show that g is an *immersion*, which will imply that X' is *quasi-projective* over S , because P is projective over S . Set

$$V_k = \phi_k(U_k) \quad (\text{open subset of } P_k)$$

$$W_k = p_k^{-1}(V_k) \quad (\text{open subset of } P)$$

$$U'_k = f^{-1}(U_k) \quad (\text{open subset of } X')$$

$$U''_k = g^{-1}(W_k) \quad (\text{open subset of } X').$$

It is clear that the U'_k form an open cover of X' ; we will first see that the U''_k also form an open cover of X' , by showing that $U'_k \subset U''_k$. For this, it will suffice to show that the diagram

$$(5.6.1.5) \quad \begin{array}{ccc} U'_k & \xrightarrow{g|_{U'_k}} & P \\ f|_{U'_k} \downarrow & & \downarrow p_k \\ U_k & \xrightarrow{\phi_k} & P_k \end{array}$$

commutes. But the prescheme $U'_k = h^{-1}(U_k \times_S P)$ is induced by X' on the open subset U'_k , and is thus the closure of $Z = U' \subset U'_k$ relative to U'_k (I, 9.5.8). To show the commutativity of (5.6.1.5), it thus suffices (since P_k is an S -scheme) to show that composing the diagram with the canonical injection $U' \rightarrow U'_k$ (or, equivalently, thanks to the isomorphism from U' to U , with ψ) gives us a commutative diagram (I, 9.5.6). But, by definition, the diagram thus obtained is exactly (5.6.1.1), whence our claim.

The W_k thus form an open cover of $g(X')$; to show that g is an immersion, it suffices to show that each of the restrictions $g|_{U''_k}$ is an immersion into W_k (I, 4.2.4). For this, consider

the morphism $u_k : W_k \xrightarrow{p_k} V_k \xrightarrow{\phi_k^{-1}} U_k \rightarrow X$; since X is separated over S , the graph morphism $\Gamma_{u_k} : W_k \rightarrow X \times_S W_k$ is a closed immersion (I, 5.4.3), and so the graph $T_k = \Gamma_{u_k}(W_k)$ is a closed subscheme of $X \times_S W_k$; if we show that $U' \rightarrow X \times_S W_k$ factors through this subscheme, then the map from the subscheme induced by X' on the open subset U''_k of X' to $X \times_S W_k$ will also factor through this graph, by (I, 9.5.8). Since the restriction of q_2 to T_k is an isomorphism onto W_k , the restriction of g to U''_k will be an immersion into W_k , and our claim will be proven. Let v_k be the canonical injection $U' \rightarrow X \times_S W_k$; we have to show that there exists a morphism $w_k : U' \rightarrow W_k$ such that $v_k = \Gamma_{u_k} \circ w_k$. By the definition of the product, it suffices to prove that $q_1 \circ v_k = u_k \circ q_2 \circ v_k$ (I, 3.2.1), or, by composing on the right [II:109] with the isomorphism $\psi' : U \rightarrow U'$, that $q_1 \circ \psi = u_k \circ q_2 \circ \psi$. But since $q_1 \circ \psi = j$ and $q_2 \circ \psi = \phi$, our claim follows from the commutativity of (5.6.1.1), taking into account the definition of u_k .

- (E) It is clear that since U , and thus U' , is irreducible, so is the X' from the preceding construction, and the morphism f is thus *birational* (I, 2.2.9). If in addition X is reduced, then so is U' , and hence X' is also reduced (I, 9.5.9). This finishes the proof.

□

Corollary (5.6.2). — Suppose that one of the hypotheses, (a) and (b), of (5.6.1) is satisfied. For X to be proper over S , it is necessary and sufficient for there to exist a projective scheme X' over S , and a surjective S -morphism $f : X' \rightarrow X$ (which is thus projective, by (5.5.5, v)). Whenever this is the case, we can further choose f to be such that there exists a dense open subset U of X for which the restriction of f to $f^{-1}(U)$ is an isomorphism $f^{-1}(U) \simeq U$, and for which $f^{-1}(U)$ is dense in X' . If in addition X is irreducible (resp. reduced), then we can assume that X' is also irreducible (resp. reduced); when X and X' are irreducible, f is a birational morphism.

Proof. The condition is sufficient, by (5.5.3) and (5.4.3, ii). It is necessary because, with the notation of (5.6.1), if X is proper over S , then X' is proper over S , because it is projective over X , and thus proper over X (5.5.3), and our claim follows from (5.4.2, ii); furthermore, since X' is quasi-projective over S , it is projective over S , by (5.5.3). □

Corollary (5.6.3). — Let S be a locally Noetherian prescheme, and X an S -scheme of finite type over S , with structure morphism $f_0 : X \rightarrow S$. For X to be proper over S , it is necessary and sufficient that, for every morphism of finite type $S' \rightarrow S$, $(f_0)_{(S')} : X_{(S')} \rightarrow S'$ be a closed morphism. It even suffices for this condition to be verified only for every S -prescheme of the form $S' = S \otimes_{\mathbf{Z}} \mathbf{Z}[T_1, \dots, T_n]$ (where the T_i are indeterminates).

Proof. The condition being clearly necessary, we now show that it is sufficient. Since the question is local on S and S' (5.4.1), we can suppose that S and S' are affine and Noetherian. By Chow's lemma, there exists a projective S -scheme P , an immersion $j : X' \rightarrow P$, and a surjective projective morphism $f : X' \rightarrow X$, such that the diagram

$$\begin{array}{ccc} X & \xleftarrow{f} & X' \\ f_0 \downarrow & & \downarrow j \\ S & \xleftarrow{r} & P \end{array}$$

commutes. Since P is of finite type over S , the first hypothesis implies that the projection $q_2 : X \times_S P \rightarrow P$ is a closed morphism. The morphism $(f, j) : X' \rightarrow X \times_S P$ factors as the graph immersion $\Gamma_j : X' \rightarrow X' \times_S P$, which is closed because P is separated over S , followed by the base change $f \times 1_P : X' \times_S P \rightarrow X \times_S P$. But f , being projective, is proper (5.5.3), and so $f \times 1_P$ is closed; hence (f, j) is closed. Since $j = q_2 \circ (f, j)$ and j is already an immersion, we conclude that j is a closed immersion, and thus proper (5.4.2, i). Furthermore, the structure morphism $r : P \rightarrow S$ is projective, and thus proper (5.5.3), so $f_0 \circ f = r \circ j$ is proper (5.4.2, ii); finally, since f is surjective, f_0 is proper, by (5.4.3).

To prove the proposition using only the second, weaker hypothesis (where S' is of the form $S \otimes_{\mathbf{Z}} \mathbf{Z}[T_1, \dots, T_n]$), it suffices to show that it implies the first. But, if S' is affine and of finite type over $S = \text{Spec}(A)$, [III-110] then we have $S' = \text{Spec}(A[c_1, \dots, c_n])$ (I, 6.3.3), and S' is thus isomorphic to a closed subscheme of $S'' = \text{Spec}(A[T_1, \dots, T_n])$ (where the T_i are indeterminates). In the commutative diagram

$$\begin{array}{ccc} X \times_S S' & \xrightarrow{1_X \times j} & X \times_S S'' \\ (f_0)_{(S')} \downarrow & & \downarrow (f_0)_{(S'')} \\ S' & \xrightarrow{j} & S'' \end{array}$$

both j and $1_X \times j$ are closed immersions (I, 4.3.1), and $(f_0)_{(S'')}$ is closed by hypothesis; thus $(f_0)_{(S')}$ is also closed. □

§6. INTEGRAL MORPHISMS AND FINITE MORPHISMS

6.1. Preschemes integral over another prescheme

Definition (6.1.1). — Let X be an S -prescheme, with structure morphism $f : X \rightarrow S$. We say that X is *integral over S* , or that f is an *integral morphism*, if there exists a cover (S_α) of S by affine opens

such that, for all α , the induced prescheme $f^{-1}(S_\alpha)$ is an affine scheme whose ring B_α is an integral algebra over the ring A_α of S_α . We say that X is *finite over S* , or that f is a *finite morphism* if X is integral and of finite type over S .

If S is affine of ring A , then we also say “integral (resp. finite) over A ” instead of “integral (resp. finite) over S ”.

(6.1.2). It is clear that, if X is integral over S , then it is *affine* over S . For an affine prescheme X over S to be integral (resp. finite) over S it is necessary and sufficient that the associated quasi-coherent $\mathcal{O}_S$ -algebra $\mathcal{A}(X)$ be such that there exist a cover (S_α) of S by affine opens having the property that, for all α , $\Gamma(S_\alpha, \mathcal{A}(X))$ is an integral (resp. integral and of finite type) algebra over $\Gamma(S_\alpha, \mathcal{O}_S)$. A quasi-coherent $\mathcal{O}_S$ -algebra with this property is said to be *integral* (resp. *finite*) over $\mathcal{O}_S$. Giving an integral (resp. finite) prescheme over S is thus (1.3.1) the same as giving a quasi-coherent $\mathcal{O}_S$ -algebra that is integral (resp. finite) over $\mathcal{O}_S$. Note that a quasi-coherent $\mathcal{O}_S$ -algebra $\mathcal{B}$ is finite if and only if it is an $\mathcal{O}_S$ -module of finite type (I, 1.3.9); it is equivalent to say that $\mathcal{B}$ is an *integral $\mathcal{O}_S$ -algebra of finite type*, since an algebra that is integral and of finite type over a ring A is an A -module of finite type.

Proposition (6.1.3). — *Let S be a locally Noetherian prescheme. For an affine prescheme X over S to be finite over S , it is necessary and sufficient that the $\mathcal{O}_S$ -algebra $\mathcal{A}(X)$ be coherent.*

Proof. Taking the preceding remark into account, this reduces to noting that, if S is locally Noetherian, then the quasi-coherent $\mathcal{O}_S$ -modules of finite type are exactly the coherent $\mathcal{O}_S$ -modules (I, 1.5.1). $\square$

Proposition (6.1.4). — *Let X be an integral (resp. finite) prescheme over S , with structure morphism $f : X \rightarrow S$. Then, for every affine open $U \subset S$ of ring A , $f^{-1}(U)$ is an affine scheme whose ring B is an integral (resp. finite) algebra over A .*

[II:111]

Proof. We first prove the following lemma:

Lemma (6.1.4.1). — *Let A be a ring, M an A -module, and $(g_i)_{1 \leq i \leq m}$ a finite system of elements of A such that the $D(g_i)$ (for $1 \leq i \leq m$) cover $\text{Spec}(A)$. If, for all i , M_{g_i} is an A_{g_i} -module of finite type, then M is an A -module of finite type.*

Proof. We can assume that M_{g_i} admits a finite system of generators (m_{ij}/g_i^n) with $m_{ij} \in M$, with n the same for all indices i . We will show that the m_{ij} form a system of generators of M . By hypothesis, for each i , there exist $a_{ij} \in A$ and some integer p (independent of i) such that, in M_{g_i} , $m/1 = (\sum_j a_{ij} m_{ij})/g_i^p$; this implies that there exists an integer $r \geq p$ such that, for all i , we have $g_i^r m \in M'$. But, since the $D(g_i^r) = D(g_i)$ cover $\text{Spec}(A)$, the ideal of A generated by the g_i^r is equal to A , or, in other words, there exist elements $a_i \in A$ such that $\sum_i a_i g_i^r = 1$; then $m = (\sum_i a_i g_i^r) m \in M'$, whence the lemma. $\square$

Now we already know (1.3.2) that $f^{-1}(U)$ is affine. If ϕ is the homomorphism $A \rightarrow B$ corresponding to f , then there exists a finite cover of U by opens $D(g_i)$ (where $g_i \in A$) such that, if $h_i = \phi(g_i)$, B_{h_i} is an integral (resp. integral and finite) algebra over A_{g_i} . Indeed, there exists a cover of U by affine opens $V_\alpha \subset U$ such that, if $A_\alpha = A(V_\alpha)$, $B_\alpha = A(f^{-1}(V_\alpha))$ is an integral (resp. finite) algebra over A_α . Every $x \in U$ belongs to some V_α , so there exists $g \in A$ such that $x \in D(g) \subset V_\alpha$; if g_α is the image of g in A_α , then $A(D(g)) = A_g = (A_\alpha)_{g_\alpha}$; let $h = \phi(g)$, and let h_α be the image of g_α in B_α ; we have

$$A(D(h)) = B_h = (B_\alpha)_{h_\alpha}$$

and, since B_α is integral (resp. finite) over A_α , $(B_\alpha)_{h_\alpha}$ is integral (resp. finite) over $(A_\alpha)_{g_\alpha}$. It now suffices to use the fact that U is quasi-compact to obtain the desired cover.

If we suppose first of all that the B_{h_i} are integral and finite over the A_{g_i} , then since B_{h_i} can also be written as B_{g_i} as an A_{g_i} -module, Lemma (6.1.4.1) shows that, in this case, B is an A -module of finite type.

Now suppose only that each B_{h_i} is integral over A_{g_i} ; let $b \in B$, and let C be the sub- A -algebra of B generated by b . For all i , C_{h_i} is the algebra over A_{g_i} generated by $b/1$ in B_{h_i} ; it follows from the

hypothesis that each C_{h_i} is an A_{g_i} -module of finite type, and so (6.1.4.1) C is an A -module of finite type, which proves that B is integral over A . $\square$

Proposition (6.1.5). —

- (i) *A closed immersion is finite (and a fortiori integral).*
- (ii) *The composition of two finite (resp. integral) morphisms is finite (resp. integral).*
- (iii) *If $f : X \rightarrow Y$ is a finite (resp. integral) S -morphism, then $f_{(S')} : X_{(S')} \rightarrow Y_{(S')}$ is finite (resp. integral) for any base extension $S' \rightarrow S$.*
- (iv) *If $f : X \rightarrow Y$ and $g : X' \rightarrow Y'$ are finite (resp. integral) S -morphisms, then $f \times_S g : X \times_S X' \rightarrow Y \times_S Y'$ is finite (resp. integral).*
- (v) *If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are morphisms such that $g \circ f$ is finite (resp. integral), if g is separated, then f is finite (resp. integral).*
- (vi) *If $f : X \rightarrow Y$ is a finite (resp. integral) morphism, then f_{red} is finite (resp. integral).*

[II:112]

Proof. By (I, 5.5.12), it suffices to prove (i), (ii), and (iii). To prove that a closed immersion $X \rightarrow S$ is finite, we can restrict to the case where $S = \text{Spec}(A)$, and everything then follows from noting that a quotient ring $A/\mathfrak{J}$ is a monogeneous A -module. To prove that the composition of two finite (resp. integral) morphisms $X \rightarrow Y$, $Y \rightarrow Z$ is finite (resp. integral), we can again assume that Z (and thus X and Y (1.3.4)) is affine, and then the claim is equivalent to saying that, if B is a finite (resp. integral) A -algebra and C a finite (resp. integral) B -algebra, then C is a finite (resp. integral) A -algebra, which is immediate. Finally, to prove (iii), we can restrict to the case where $S = Y$, since $X_{(S')}$ can be identified with $X \times_Y Y_{(S')}$ (I, 3.3.11); we can further suppose that $S = \text{Spec}(A)$ and $S' = \text{Spec}(A')$; then X is affine of ring B (1.3.4), and $X_{(S')}$ affine of ring $A' \otimes_A B$, and it suffices to note that, if B is a finite (resp. integral) A -algebra, then $A' \otimes_A B$ is a finite (resp. integral) A' -algebra. $\square$

We also note that, if X and Y are S -preschemes that are finite (resp. integral) over S , then their sum $X \sqcup Y$ is a finite (resp. integral) prescheme over S , since this reduces to showing that, if B and C are finite (resp. integral) A -algebras over A , then so too is $B \times C$.

Corollary (6.1.6). — *If X is an integral (resp. finite) prescheme over S , then, for every open $U \subset S$, $f^{-1}(U)$ is integral (resp. finite) over U .*

Proof. This is a particular case of (6.1.5, (iii)). $\square$

Corollary (6.1.7). — *Let $f : X \rightarrow Y$ be a finite morphism. Then, for all $y \in Y$, the fibre $f^{-1}(y)$ is a finite algebraic scheme over $k(y)$, and a fortiori its underlying space is discrete and finite.*

Proof. Indeed, as a $k(y)$ -prescheme, $f^{-1}(y)$ can be identified with $X \times_Y \text{Spec}(k(y))$ (I, 3.6.1), which is finite over $\text{Spec}(k(y))$ (6.1.5, (iii)); it is thus an affine scheme whose ring is an algebra of finite rank over $k(y)$ (6.1.4). The corollary then follows from (I, 6.4.4). $\square$

Corollary (6.1.8). — *Let X and S be integral preschemes, and $f : X \rightarrow S$ a dominant morphism. If f is integral (resp. finite) then the field $R(X)$ of rational functions on X is algebraic (resp. algebraic of finite degree) over the field $R(S)$ of rational functions on S .*

Proof. Let s be the generic point of S ; the $k(s)$ -prescheme $f^{-1}(s)$ is integral (resp. finite) over $\text{Spec}(k(s))$ (6.1.5, (iii)) and contains, by hypothesis, the generic point x of X ; the local ring of x in $f^{-1}(s)$ is equal to $k(x)$ (I, 3.6.5), and is thus a local ring of an integral (resp. finite) algebra over $k(s)$ (6.1.4), whence the corollary. $\square$

Remark (6.1.9). — The hypothesis that g be separated is essential for the validity of (6.1.5, (v)): if Y is not separated over Z , then the identity 1_Y is the composite morphism $Y \xrightarrow{\Delta_Y} Y \times_Z Y \xrightarrow{p_1} Y$, but Δ_Y is not an integral morphism, as follows from (6.1.10):

Proposition (6.1.10). — *Every integral morphism is universally closed.*

Proof. Let $f : X \rightarrow Y$ be an integral morphism; by (6.1.5, (iii)), it suffices to show that f is *closed*. Let Z be a closed subset of X ; ^[II:113] then there exists a subprescheme of X whose underlying space is Z (I, 5.2.1), and it thus follows from (6.1.5, (i) and (ii)) that it suffices to prove that $f(X)$ is *closed* in Y . By (6.1.5, (vi)), we can suppose that X and Y are *reduced*; further, if T is the closed reduced subprescheme of Y whose underlying space is $\overline{f(X)}$ (I, 5.2.1), then we know that f factors as $X \xrightarrow{g} T \xrightarrow{j} Y$, where j is the injection morphism (I, 5.2.2), and since j is separated (I, 5.5.1, (i)), it follows from (6.1.5, (v)) that g is an integral morphism. We can thus suppose that $f(X)$ is *dense* in Y . Finally, since the question is local on Y , we can restrict to the case where $Y = \text{Spec}(A)$. Then $X = \text{Spec}(B)$, where B is an A -algebra that is integral over A (6.1.4); furthermore, A is reduced (I, 5.1.4), and the hypothesis that $f(X)$ be dense in Y implies that the homomorphism $\phi : A \rightarrow B$ corresponding to f is *injective* (I, 1.2.7). Under these conditions, to say that $f(X) = Y$ is to say that every prime ideal of A is the intersection with A of a prime ideal of B , which is exactly the first theorem of Cohen–Seidenberg ([13, t. I, p. 257, th. 3]). $\square$

Corollary (6.1.11). — *Every finite morphism $f : X \rightarrow Y$ is projective.*

Proof. Since f is affine, $\mathcal{O}_X$ is a *very ample* $\mathcal{O}_X$ -module with respect to f (5.1.2); furthermore, $f_*(\mathcal{O}_X)$ is a quasi-coherent $\mathcal{O}_Y$ -module of *finite type* (6.1.2); finally, f is separated, of finite type, and universally closed (6.1.10), and thus satisfies the conditions of criterion (5.5.4, (i)). $\square$

Proposition (6.1.12). — *Let $f : X' \rightarrow X$ be a finite morphism, and let $\mathcal{B} = f_*(\mathcal{O}_{X'})$ (which is a quasi-coherent $\mathcal{O}_X$ -algebra, and a $\mathcal{O}_X$ -module of finite type). Let $\mathcal{F}'$ be a quasi-coherent $\mathcal{O}_{X'}$ -module; for $\mathcal{F}'$ to be locally free of rank r , it is necessary and sufficient that $f_*(\mathcal{F}')$ be a locally free $\mathcal{B}$ -module of rank r .*

Proof. It is clear that, if $f_*(\mathcal{F}')|U$ is isomorphic to $\mathcal{B}^r|U$ (where $U \subset X$ is open), then $\mathcal{F}'|f^{-1}(U)$ is isomorphic to $\mathcal{O}_{X'}^r|f^{-1}(U)$ (1.4.2). Conversely, suppose that $\mathcal{F}'$ is locally free of rank r ; we will show that $f_*(\mathcal{F}')$ is locally isomorphic to $\mathcal{B}^r$ as a $\mathcal{B}$ -module. Let x be a point of X ; as U runs over a fundamental system of affine neighbourhoods of x , $f^{-1}(U)$ runs over a fundamental system of affine neighbourhoods (1.2.5) of the finite set $f^{-1}(x)$, since f is closed (6.1.10). The proposition then follows from the following lemma: $\square$

Lemma (6.1.12.1). — *Let Y be a prescheme, $\mathcal{E}$ a locally free $\mathcal{O}_Y$ -module of rank r , and Z a finite subset of Y contained inside some affine open V . Then there exists a neighbourhood $U \subset V$ of Z such that $\mathcal{E}|U$ is isomorphic to $\mathcal{O}_Y^r|U$.*

Proof. We can evidently assume that Y is affine; for all $z_i \in Z$, there exists in the closure $\overline{\{z_i\}}$ at least one closed point z'_i (0, 2.1.3); if Z' is the set of the z'_i then every neighbourhood of Z' is a neighbourhood of Z , and we can thus assume that Z is discrete and closed in Y . Consider the closed reduced subprescheme of Y that has Z as its underlying space (I, 5.2.1) and let $j : Z \rightarrow Y$ be the canonical injection; $j^*(\mathcal{E}) = \mathcal{E} \otimes_Y \mathcal{O}_Z$ is locally free of rank r on the discrete scheme Z , and is thus isomorphic to $\mathcal{O}_Z^r$; in other words, there exist r sections s_i (for $1 \leq i \leq r$) of $\mathcal{E} \otimes_Y \mathcal{O}_Z$ over Z such that the homomorphism $\mathcal{O}_Z^r \rightarrow \mathcal{E} \otimes_Y \mathcal{O}_Z$ defined by these sections is bijective. But $Y = \text{Spec}(A)$ is affine, Z is defined by an ideal $\mathfrak{J}$ of A , and we have $\mathcal{E} = \tilde{M}$, where M is an A -module; the s_i are elements of $M \otimes_A (A/\mathfrak{J})$ and are thus images of r elements $t_i \in M = \Gamma(Y, \mathcal{E})$. ^[II:114] For all $z_j \in Z$ there is thus a neighbourhood V_j of z_j such that the restrictions of the t_i to V_j define an isomorphism $\mathcal{O}_Y^r|V_j \rightarrow \mathcal{E}|V_j$ (0, 5.5.4); the neighbourhood U given by the union of the V_j is the desired neighbourhood. $\square$

Proposition (6.1.13). — *Let $g : X' \rightarrow X$ be an integral morphism of preschemes, Y a normal locally integral prescheme, and f a rational map from Y to X' such that $g \circ f$ is an everywhere defined rational map (I, 7.2.1). Then f is everywhere defined.*

Proof. If f_1 and f_2 are morphisms (from dense open subsets of Y to X') in the class f , then it is clear that $g \circ f_1$ and $g \circ f_2$ are equivalent morphisms, which justifies the notation $g \circ f$ for their equivalence class. We recall also that, if we further suppose Y to be *locally Noetherian*, then the hypothesis that Y is normal already implies that Y is locally integral (I, 6.1.13).

To prove the proposition, note first of all that the question is local on Y , and so we can suppose that there exists in the class $g \circ f$ a morphism $h : Y \rightarrow X$. Consider the inverse image

$Y' = X'_{(h)} = X'_{(Y)}$, and note that the morphism $g' = g_{(Y)} : Y' \rightarrow Y$ is integral (6.1.5, (iii)). Given the correspondence between rational maps from Y to X' and rational Y -sections of Y' (I, 7.1.2), we see that it suffices to prove the specific case of (6.1.13) where $X = Y$; in other words, the following: $\square$

Corollary (6.1.14). — *Let X be a normal locally integral prescheme, $g : X' \rightarrow X$ an integral morphism, and f a rational X -section of X' . Then f is everywhere defined.*

Proof. Since the question is local on X , we can assume that X is integral, and then f is identified with a morphism from an open U of X to X' (I, 7.2.2) that is a U -section of $g^{-1}(U)$. Since g is separated, f is a closed immersion from U into $g^{-1}(U)$ (I, 5.4.6); let Z be the closed subscheme of $g^{-1}(U)$ associated to f (I, 4.2.1), which is isomorphic to U , and thus integral; let X_1 be the reduced subscheme of X' whose underlying space is the closure $\bar{Z}$ of Z in X' (I, 5.2.1); then Z is an induced subscheme on an open of X_1 (I, 5.2.3), and, since it is irreducible, so too is X_1 , which is thus integral. The morphism f can then be considered as a rational X -section of X_1 ; since the restriction of g to X_1 is an integral morphism (6.1.5, (i) and (ii)), we can finally reduce to proving (6.1.14) in the specific case where $X' = X_1$; in other words, the following: $\square$

Corollary (6.1.15). — *Let X be a normal integral prescheme, X' an integral prescheme, and $g : X' \rightarrow X$ an integral morphism. If there exists a rational X -section f of X' , then g is an isomorphism.*

Proof. Since the question is local on X , we can assume that X is affine of integral ring A , and then X' is affine of ring A' with A' integral over A (6.1.4) and integral; furthermore, the argument of (6.1.14) shows that there exists a dense open of X that is isomorphic to a dense open of X' , and so A and A' have the same field of fractions. Also, by (I, 8.2.1.1), and the hypothesis that the $\mathcal{O}_x$ are integrally closed, the ring A is integrally closed, and so $A' = A$, which finishes the proof of (6.1.13). $\square$

6.2. Quasi-finite morphisms

Proposition (6.2.1). — *Let $f : X \rightarrow Y$ be a morphism locally of finite type, and x a point of X . Then the following conditions are equivalent:* [II:115]

- a) *The point x is isolated in its fibre $f^{-1}(f(x))$.*
- b) *The ring $\mathcal{O}_x$ is a quasi-finite $\mathcal{O}_{f(x)}$ -module (0, 7.4.1).*

Proof. Since the question is clearly local on X and on Y , we can assume that $X = \text{Spec}(A)$ and $Y = \text{Spec}(B)$ are affine, with A a B -algebra of finite type (I, 6.3.3). Furthermore, we can replace X by $X \times_Y \text{Spec}(\mathcal{O}_{f(x)})$ without changing either the fibre $f^{-1}(f(x))$ or the local ring $\mathcal{O}_x$ (I, 3.6.5); we can thus assume that B is a local ring (equal to $\mathcal{O}_{f(x)}$); if $\mathfrak{n}$ is the maximal ideal of B , then $f^{-1}(f(x))$ is an affine scheme of ring $A/\mathfrak{n}A$, of finite type over $k(f(x)) = B/\mathfrak{n}$ (I, 6.4.11). With this, if (a) is satisfied then we can further suppose that $f^{-1}(f(x))$ consists of the single point x ; thus $A/\mathfrak{n}A$ is of finite rank over $B/\mathfrak{n}$ (I, 6.4.4), or, in other words, A is a quasi-finite B -module. Conversely, if A is a quasi-finite B -module, then $f^{-1}(f(x))$ is an Artinian affine scheme, and thus discrete (I, 6.4.4); so x is isolated in its fibre, and this shows that (b) implies (a). $\square$

Corollary (6.2.2). — *Let $f : X \rightarrow Y$ be a morphism of finite type. Then the following conditions are equivalent:*

- a) *Every point $x \in X$ is isolated in its fibre $f^{-1}(f(x))$ (or, in other words, the subspace $f^{-1}(f(x))$ is discrete).*
- b) *For all $x \in X$, the prescheme $f^{-1}(f(x))$ is a finite $k(f(x))$ -prescheme.*
- c) *For all $x \in X$, the ring $\mathcal{O}_x$ is a quasi-finite $\mathcal{O}_{f(x)}$ -module.*

Proof. The equivalence of (a) and (c) follows from (6.2.1). Since $f^{-1}(f(x))$ is an algebraic $k(f(x))$ -prescheme (I, 6.4.11), the equivalence of (a) and (b) follows from (I, 6.4.4). $\square$

Definition (6.2.3). — *If $f : X \rightarrow Y$ is a morphism of finite type satisfying the equivalent conditions of (6.2.2), we say that f is quasi-finite, or that X is quasi-finite over Y .*

It is clear that every finite morphism is quasi-finite (6.1.7).

Proposition (6.2.4). —

- (i) An immersion $X \rightarrow Y$ that is closed, or such that X is Noetherian, is a quasi-finite morphism.
- (ii) If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are quasi-finite morphisms, then $g \circ f$ is quasi-finite.
- (iii) If X and Y are S -preschemes, and $f : X \rightarrow Y$ a quasi-finite S -morphism, then $f_{(S')} : X_{(S')} \rightarrow Y_{(S')}$ is quasi-finite for any base extension $g : S' \rightarrow S$.
- (iv) If $f : X \rightarrow Y$ and $g : X' \rightarrow Y'$ are quasi-finite S -morphisms, then

$$f \times_S g : X \times_S X' \longrightarrow Y \times_S Y'$$

is quasi-finite.

- (v) Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be morphisms such that $g \circ f$ is quasi-finite; if, further, g is separated, or X is Noetherian, or $X \times_Z Y$ is locally Noetherian, then f is quasi-finite.
- (vi) If f is quasi-finite, then f_{red} is quasi-finite.

Proof. If $f : X \rightarrow Y$ is an immersion, then every fibre has at most one point, and claim (i) then follows from (I, 6.3.4, (i)) and (I, 6.3.5). To prove (ii), note first of all that $h = g \circ f$ is of finite type (I, 6.3.4, (ii)); furthermore, if $z = h(x)$ and $y = f(x)$, then y is isolated in $g^{-1}(z)$, and so there exists an open neighbourhood V of y in Y that does not meet any point of $g^{-1}(z)$ apart from y ; thus $f^{-1}(V)$ is an open neighbourhood of x that does not meet any of the $f^{-1}(y')$, where $y' \neq y$ is in $g^{-1}(z)$; ^[II.116] since x is isolated in $f^{-1}(y)$, it is isolated in $h^{-1}(z) = f^{-1}(g^{-1}(z))$. To prove (iii), we can reduce to the case where $Y = S$ (I, 3.3.11); we again note first of all that $f' = f_{(S')}$ is of finite type (I, 6.3.4, (iii)); also, if $x' \in X' = X_{(S')}$, and if we set $y' = f'(x')$ and $y = g(y')$, then $f'^{-1}(y')$ can be identified with $f^{-1}(y) \otimes_{k(y)} k(y')$ (I, 3.6.5); since $f^{-1}(y)$ is of finite rank over $k(y)$ by hypothesis, $f'^{-1}(y')$ is of finite rank over $k(y')$, and thus discrete. Claims (iv), (v), and (vi) follow from (i), (ii), and (iii) by the general method (I, 5.5.12), except when the hypothesis in (v) is not “ g is separated”; in these cases, we remark first of all that, if x is isolated in $f^{-1}(g^{-1}(g(f(x))))$, then it is *a fortiori* isolated in $f^{-1}(f(x))$; the fact that f is of finite type then follows from (I, 6.3.6). $\square$

Proposition (6.2.5). — Let A be a complete local Noetherian ring, X an A -scheme locally of finite type, and x a point of X over the closed point y of $Y = \text{Spec}(A)$. Suppose that x is isolated in its fibre $f^{-1}(y)$ (where f is the structure morphism $X \rightarrow Y$). Then $\mathcal{O}_x$ is an A -module of finite type, and X is Y -isomorphic to the sum (I, 3.1) of $X' = \text{Spec}(\mathcal{O}_x)$ (which is a finite Y -scheme) and an A -scheme X'' .

Proof. It follows from (6.2.1) that $\mathcal{O}_x$ is a quasi-finite A -module. Since $\mathcal{O}_x$ is Noetherian (I, 6.3.7), and the homomorphism $A \rightarrow \mathcal{O}_x$ is local, the hypothesis that A is complete implies that $\mathcal{O}_x$ is an A -module of finite type (0, 7.4.3). Let $X' = \text{Spec}(\mathcal{O}_x)$ be the local scheme of X at the point x (I, 2.4.1), and let $g : X' \rightarrow X$ be the canonical morphism. Since the composite $X' \xrightarrow{g} X \xrightarrow{f} Y$ is finite (6.1.1), and since f is separated, g is finite (6.1.5, (v)), and so $g(X')$ is closed in X (6.1.10); also, since g is of finite type, g is a local isomorphism at the closed point x' of X' by the definition of g and (I, 6.5.4); but since X' is the only open neighbourhood of x' , this implies that g is an open immersion, and so $g(X')$ is also open in X , which finishes the proof. $\square$

Corollary (6.2.6). — Let A be a complete local Noetherian ring, $Y = \text{Spec}(A)$, and $f : X \rightarrow Y$ a separated quasi-finite morphism. Then X is Y -isomorphic to a sum $X' \sqcup X''$, where X' is a finite Y -scheme, and X'' is a quasi-finite Y -scheme such that, if y is the closed point of Y , then $X'' \cap f^{-1}(y) = \emptyset$.

Proof. Indeed, the fibre $f^{-1}(y)$ is finite and discrete by hypothesis, and the corollary then follows, by induction on the number of points in this fibre, from (6.2.5). $\square$

Remark (6.2.7). — In Chapter V, we will see that, if Y is locally Noetherian, then a separated quasi-finite morphism $X \rightarrow Y$ is necessarily quasi-affine.

6.3. Integral closure of a prescheme

Proposition (6.3.1). — *Let $(X, \mathcal{A})$ be a ringed space, $\mathcal{B}$ a (commutative) $\mathcal{A}$ -algebra, and f a section of $\mathcal{B}$ over X . Then the following properties are equivalent:*

- a) *The sub- $\mathcal{A}$ -module of $\mathcal{B}$ generated by the f^n for $n \geq 0$ (0, 5.1.1) is of finite type.*
- b) *There exists a sub- $\mathcal{A}$ -algebra $\mathcal{C}$ of $\mathcal{B}$ that is an $\mathcal{A}$ -module of finite type and such that $f \in \Gamma(X, \mathcal{C})$.*
- c) *For all $x \in X$, f_x is integral over the stalk $\mathcal{A}_x$.*

[II:117]

Proof. Since the sub- $\mathcal{A}$ -module of $\mathcal{B}$ generated by the f^n is an $\mathcal{A}$ -algebra, it is clear that (a) implies (b). On the other hand, (b) implies that, for all $x \in X$, the $\mathcal{A}_x$ -module $\mathcal{C}_x$ is of finite type, which implies that every element of the algebra $\mathcal{C}_x$, and, in particular, f_x , is integral over $\mathcal{A}_x$. Finally, if we have, for some $x \in X$, a relation of the form

$$f_x^n + (a_1)_x f_x^{n-1} + \dots + (a_n)_x = 0$$

where the a_i (for $1 \leq i \leq n$) are sections of $\mathcal{A}$ over an open neighbourhood U of x , then the section $f^n|U + a_1 \cdot f^{n-1}|U + \dots + a_n$ is zero over a neighbourhood $V \subset U$ of x , from which it immediately follows that the $f^k|V$ (for $k \geq 0$) are linear combinations of the $f^j|V$ (for $0 \leq j \leq n-1$) with coefficients in $\Gamma(V, \mathcal{A})$; we thus conclude that (c) implies (a). $\square$

If the equivalent conditions of (6.3.1) are satisfied then we say that the section f is *integral over $\mathcal{A}$* .

Corollary (6.3.2). — *Under the hypotheses of (6.3.1), there exists a (unique) sub- $\mathcal{A}$ -module $\mathcal{A}'$ of $\mathcal{B}$ such that, for all $x \in X$, $\mathcal{A}'_x$ is the set of germs $f_x \in \mathcal{B}_x$ that are integral over $\mathcal{A}_x$. For every open $U \subset X$, the sections of $\mathcal{A}'$ over U are the sections $f \in \Gamma(U, \mathcal{B})$ that are integral over $\mathcal{A}|U$.*

Proof. The existence of $\mathcal{A}'$ is immediate, by taking $\Gamma(U, \mathcal{A}')$ to be the set of $f \in \Gamma(U, \mathcal{B})$ such that f_x is integral over $\mathcal{A}_x$ for all $x \in U$. The second claim follows immediately from (6.3.1). $\square$

It is clear that $\mathcal{A}'$ is a sub- $\mathcal{A}$ -algebra of $\mathcal{B}$; we say that it is the *integral closure of $\mathcal{A}$ in $\mathcal{B}$* .

(6.3.3). Let $(X, \mathcal{A})$ and $(Y, \mathcal{B})$ be ringed spaces, and let

$$g = (\psi, \theta) : X \longrightarrow Y$$

be a morphism. Let $\mathcal{C}$ (resp. $\mathcal{D}$) be an $\mathcal{A}$ -algebra (resp. $\mathcal{B}$ -algebra), and let

$$u : \mathcal{D} \longrightarrow \mathcal{C}$$

be a g -morphism (0, 4.4.1). Then, if $\mathcal{A}'$ (resp. $\mathcal{B}'$) is the integral closure of $\mathcal{A}$ (resp. $\mathcal{B}$) in $\mathcal{C}$ (resp. $\mathcal{D}$), the restriction of u to $\mathcal{B}'$ is a g -morphism

$$u' : \mathcal{B}' \longrightarrow \mathcal{A}'.$$

Indeed, if j is the canonical injection $\mathcal{B}' \rightarrow \mathcal{D}$, then it suffices to show that

$$v = u^\# \circ g^*(j) : g^*(\mathcal{B}') \longrightarrow \mathcal{C}$$

sends $g^*(\mathcal{B}')$ to $\mathcal{A}'$. But an element of $g^*(\mathcal{B}')_x = \mathcal{B}'_{\psi(x)} \otimes_{\mathcal{B}_{\psi(x)}} \mathcal{A}_x$ is integral over $\mathcal{A}_x$ by the definition of $\mathcal{B}'$, and thus so too is its image under v_x , which proves the claim.

Proposition (6.3.4). — *Let X be a prescheme, and $\mathcal{A}$ a quasi-coherent $\mathcal{O}_X$ -algebra. Then the integral closure $\mathcal{O}'_X$ of $\mathcal{O}_X$ in $\mathcal{A}$ is a quasi-coherent $\mathcal{O}_X$ -algebra, and, for every affine open U of X , $\Gamma(U, \mathcal{O}'_X)$ is the integral closure of $\Gamma(U, \mathcal{O}_X)$ in $\Gamma(U, \mathcal{A})$.*

Proof. We can restrict to the case where $X = \text{Spec}(B)$ is affine, and $\mathcal{A} = \tilde{A}$, where A is a B -algebra; [II:118] let B' be the integral closure of B in A . Everything then reduces to proving that, for all $x \in X$, an element of A_x which is integral over B_x necessarily belongs to B'_x , which follows from the fact that, for a commutative ring C , the operations of taking the integral closure in a C -algebra and passing to a ring of fractions (with respect to a multiplicative subset of C) commute [13, t. I, pp. 261 and 257]. $\square$

The X -scheme $X' = \operatorname{Spec}(\mathcal{O}'_X)$ is then called the *integral closure of X with respect to $\mathcal{A}$* (or with respect to $\operatorname{Spec}(\mathcal{A})$); it is clear that X' is *integral* over X (6.1.2).

We immediately deduce from (6.3.4) that, if $f : X' \rightarrow X$ is the structure morphism, then, for every open U of X , $f^{-1}(U)$ is the *integral closure of the prescheme induced by X on U , with respect to $\mathcal{A}|_U$* .

(6.3.5). Let X and Y be preschemes, $f : X \rightarrow Y$ a morphism, $\mathcal{A}$ (resp. $\mathcal{B}$) a quasi-coherent $\mathcal{O}_X$ -algebra (resp. quasi-coherent $\mathcal{O}_Y$ -algebra), and let $u : \mathcal{B} \rightarrow \mathcal{A}$ be an f -morphism. We have seen (6.3.3) that we obtain an f -morphism $u' : \mathcal{O}'_Y \rightarrow \mathcal{O}'_X$, where $\mathcal{O}'_X$ (resp. $\mathcal{O}'_Y$) is the integral closure of $\mathcal{O}_X$ (resp. $\mathcal{O}_Y$) in $\mathcal{A}$ (resp. $\mathcal{B}$). Then, if X' (resp. Y') is the integral closure of X (resp. Y) with respect to $\mathcal{A}$ (resp. $\mathcal{B}$), we canonically obtain from u a morphism $f' = \operatorname{Spec}(u') : X' \rightarrow Y'$ (1.5.6) such that the diagram

$$(6.3.5.1) \quad \begin{array}{ccc} X' & \xrightarrow{f'} & Y' \\ \downarrow & & \downarrow \\ X & \xrightarrow{f} & Y \end{array}$$

commutes.

(6.3.6). Suppose that X has only a *finite* number of irreducible components X_i (for $1 \leq i \leq r$), with generic points ξ_i , and consider, in particular, the integral closure of X with respect to some quasi-coherent $\mathcal{R}(X)$ -algebra $\mathcal{A}$ (quasi-coherent as either an $\mathcal{O}_X$ -algebra or as an $\mathcal{R}(X)$ -algebra, since the two are equivalent). We know (I, 7.3.5) that $\mathcal{A}$ is a direct sum of r quasi-coherent $\mathcal{O}_X$ -algebras $\mathcal{A}_i$, where the support of $\mathcal{A}_i$ is contained inside X_i , and the sheaf induced by $\mathcal{A}_i$ on X_i is a simple sheaf whose fibre A_i is an algebra over $\mathcal{O}_{\xi_i}$. It is then clear (6.3.4) that the integral closure $\mathcal{O}'_X$ of $\mathcal{O}_X$ in $\mathcal{A}$ is the *direct sum* of the integral closures $\mathcal{O}_X^{(i)}$ of $\mathcal{O}_X$ in each of the $\mathcal{A}_i$, and thus the integral closure $X' = \operatorname{Spec}(\mathcal{O}'_X)$ of X with respect to $\mathcal{A}$ is the X -scheme given by the *sum* of the $\operatorname{Spec}(\mathcal{O}_X^{(i)}) = X'_i$ (for $1 \leq i \leq r$).

Suppose further that the $\mathcal{O}_X$ -algebra $\mathcal{A}$ is *reduced*, or, equivalently, that each of the algebras A_i is reduced, and can thus be considered as an algebra over the field $k(\xi_i)$ (equal to the field of rational functions of the reduced subprescheme of X that has X_i as its underlying space); then (1.3.8) each of the X'_i is a *reduced* X -scheme, and X' is also the integral closure of X_{red} . Suppose further that each of the algebras A_i is a *direct sum of a finite number of fields* K_{ij} (for $1 \leq j \leq s_i$); if $\mathcal{K}_{ij}$ is the sub-algebra of $\mathcal{A}_i$ corresponding to K_{ij} , then it is clear that $\mathcal{O}_X^{(i)}$ is the *direct sum* of the integral closures $\mathcal{O}_X^{(ij)}$ of $\mathcal{O}_X$ in each of the $\mathcal{K}_{ij}$. Then X'_i is the *sum* of the X -schemes $X'_{ij} = \operatorname{Spec}(\mathcal{O}_X^{(ij)})$ (for $1 \leq j \leq s_i$). Furthermore, under these hypotheses, and with this notation, we have:

[II:119]

Proposition (6.3.7). — *Each of the X'_{ij} is an integral normal X -prescheme, and its field of rational functions $R(X'_{ij})$ is canonically identified with the algebraic closure K'_{ij} of $k(\xi_i)$ in K_{ij} .*

Proof. By the above, we can suppose that X is integral, and so $r = 1$ and $s_1 = 1$, so that the unique algebra A_1 is a field K ; let ξ be the generic point of X , and let $f : X' \rightarrow X$ be the structure morphism. For every non-empty affine open U of X , $f^{-1}(U)$ can be identified with the spectrum of the integral closure B'_U in the field K of the integral ring $B_U = \Gamma(U, \mathcal{O}_X)$ (6.3.4); since the ring B'_U is integral and integrally closed, so too are the local rings of the points of its spectrum, and thus $f^{-1}(U)$ is by definition an *integral* and *normal* scheme ((0, 4.1.4) and (I, 5.1.4)). Furthermore, since (0) is the only prime ideal of B'_U over the prime ideal (0) of B_U [13, t. I, p. 259], $f^{-1}(\xi)$ consists of a single point ξ' , and $k(\xi')$ is the field of fractions K' of B'_U , which is exactly the algebraic closure of $k(\xi)$ in K . Finally, X' is irreducible, since, if U runs over the set of non-empty affine opens of X , then the $f^{-1}(U)$ give an open cover of X' by irreducible opens; furthermore, the intersection $f^{-1}(U \cap V)$ of any two of these opens contains ξ' , and is thus non-empty, and we conclude by (0, 2.1.4). $\square$

Corollary (6.3.8). — Let X be a reduced prescheme that has only a finite number of irreducible components X_i (where $1 \leq i \leq r$), and let ξ_i be the generic point of X_i . Then the integral closure X' of X with respect to $\mathcal{R}(X)$ is the sum of the r X -schemes X'_i , which are integral and normal. If $f : X' \rightarrow X$ is the structure morphism, then $f^{-1}(\xi_i)$ consists only of the generic point ξ'_i of X'_i , and we have $k(\xi'_i) = k(\xi_i)$, or, in other words, that f is birational.

In this case, we say that X' is the *normalisation* of the reduced prescheme X ; we note that f , since it is birational and integral, is *surjective* (6.1.10). Then in order to have $X' = X$, it is necessary and sufficient that X be *normal*. If X is an integral prescheme, then it follows from (6.3.8) that its normalisation X' is integral.

(6.3.9). Let X and Y be integral preschemes, $f : X \rightarrow Y$ a dominant morphism, and $L = R(X)$ (resp. $K = R(Y)$) the field of rational functions of X (resp. Y); there is a canonical injection $K \rightarrow L$ corresponding to f , and if we identify K (resp. L) with the simple sheaf $\mathcal{R}(Y)$ (resp. $\mathcal{R}(X)$), this injection is an f -morphism. Let K_1 (resp. L_1) be an extension of K (resp. L), and suppose we have a morphism $K_1 \rightarrow L_1$ such that the diagram

$$\begin{array}{ccc} K_1 & \longrightarrow & L_1 \\ \uparrow & & \uparrow \\ K & \longrightarrow & L \end{array}$$

commutes; if K_1 (resp. L_1) is considered as a simple sheaf on Y (resp. X), and thus as an $\mathcal{R}(Y)$ -algebra (resp. $\mathcal{R}(X)$ -algebra), this implies that $K_1 \rightarrow L_1$ is an f -morphism. With this, if X' (resp. Y') is the integral closure of X (resp. Y) with respect to L_1 (resp. K_1), then X' (resp. Y') is a normal integral prescheme (6.3.6) whose field of rational functions is canonically identified with the algebraic closure L' (resp. K') of L (resp. K) in L_1 (resp. K_1), and there exists a (necessarily dominant) canonical morphism $f' : X' \rightarrow Y'$ that makes the diagram (6.3.5.1) commute. [II:120]

The most important case is when we take $L_1 = L$, with K_1 then being an extension of K contained in L , and when we suppose X to be integral and *normal*, so that $X' = X$. The above then shows that, since X is normal, if Y' is the integral closure of Y with respect to a field $K_1 \subset L = R(X)$, then every dominant morphism $f : X \rightarrow Y$ factors as

$$f : X \xrightarrow{f'} Y' \longrightarrow Y$$

where f' is dominant; also, since the monomorphism $K_1 \rightarrow L$ is given, f' is necessarily unique, as we see by reducing to the case where X and Y are affine. We thus see that, given Y , L , and a K -monomorphism $K_1 \rightarrow L$, the integral closure Y' of Y with respect to K_1 is the solution of a *universal problem*.

Remark (6.3.10). — Consider the hypotheses of (6.3.6), and suppose further that each of the algebras A_i is of *finite rank* over $k(\xi_i)$ (which implies that A_i is a direct sum of a finite number of fields); we can then show, in certain cases, that the structure morphism $X' \rightarrow X$ is not just integral, but even *finite*. Restrict to the case where X is *reduced*; since the question is local on X , we can further suppose that X is affine of ring C , and that C has only a finite number of minimal prime ideals $\mathfrak{p}_i$ (for $1 \leq i \leq r$) such that the $C_i = C/\mathfrak{p}_i$ are integral; then X' is finite over X if the integral closure of each C_i in every finite-degree extension of its field of fractions is a C -module of *finite type* (6.3.4). We know that this condition is always satisfied if C is an *algebra of finite type over a field* [13, t. I, p. 267, th. 9], or *over $\mathbf{Z}$* [9, I, p. 93, th. 3], or *over a complete Noetherian local ring* [25, p. 298]. We thus conclude that $X' \rightarrow X$ will be a finite morphism whenever X is a scheme of *finite type* over a field, or over $\mathbf{Z}$, or over a complete Noetherian local ring.

6.4. Determinant of an endomorphism of $\mathcal{O}_X$ -modules

(6.4.1). Let A be a (commutative) ring, E a free A -module of rank n , and u an endomorphism of E ; recall that, to define the *characteristic polynomial* of u , we consider the endomorphism $u \otimes 1$ of the free $A[T]$ -module of rank n , $E \otimes_A A[T]$ (where T is an indeterminate), and we set

$$(6.4.1.1) \quad P(u, T) = \det(T \cdot I - (u \otimes 1))$$

(where I is the identity automorphism of $E \otimes_A A[T]$). We have

$$(6.4.1.2) \quad P(u, T) = T^n - \sigma_1(u)T^{n-1} + \dots + (-1)^n \sigma_n(u)$$

where $\sigma_i(u)$ is an element of A , equal to a homogeneous polynomial of degree i (with integer coefficients) with respect to the entries of the matrix of u in some arbitrary basis of E ; we say that the $\sigma_i(u)$ are the *elementary symmetric functions* of u ; in particular, we have that $\sigma_1(u) = \text{Tr } u$ and $\sigma_n(u) = \det u$. Recall that, by the Hamilton–Cayley theorem, we have

$$(6.4.1.3) \quad P(u, u) = u^n - \sigma_1(u)u^{n-1} + \dots + (-1)^n \sigma_n(u) = 0$$

[II:121] which can also be written as

$$(6.4.1.4) \quad (\det u) \cdot I_E = uQ(u)$$

(where I_E is the identity automorphism of E), where

$$(6.4.1.5) \quad Q(u) = (-1)^{n+1}(u^{n-1} - \sigma_1(u)u^{n-2} + \dots + (-1)^{n-1}\sigma_{n-1}(u)).$$

Let $\phi : A \rightarrow B$ be a ring homomorphism, which makes B into an A -algebra; consider the B -module $E_{(B)} = E \otimes_A B$, which is free of rank n , and the extension $u \otimes 1$ of u to an endomorphism of $E_{(B)}$; it is immediate that we have $\sigma_i(u \otimes 1) = \phi(\sigma_i(u))$ for all indices i .

(6.4.2). Now suppose that A is an *integral* ring, with field of fractions K , and let E be an A -module of *finite type* (but not necessarily free). Let n be the *rank* of E , i.e. the rank of the free K -module $E \otimes_A K$; to every endomorphism u of E there canonically corresponds the endomorphism $u \otimes 1$ of $E \otimes_A K$; by an abuse of language, we also define the *characteristic polynomial* of u , denoted by $P(u, T)$, to be the polynomial $P(u \otimes 1, T)$, whose coefficients $\sigma_i(u \otimes 1)$ (which belong to K) are also denoted by $\sigma_i(u)$ and called the *elementary symmetric functions* of u ; in particular, $\det u = \det(u \otimes 1)$ by definition. With this notation, Equations (6.4.1.3) to (6.4.1.5) make sense and still hold true, if we interpret u^j as the composite

$$E \xrightarrow{x \mapsto x \otimes 1} E \otimes_A K \xrightarrow{(u \otimes 1)^j} E \otimes_A K.$$

If F is the torsion submodule of E , and if $E_0 = E/F$, then $u(F) \subset F$, and so, by passing to quotients, u gives an endomorphism u_0 of E_0 ; furthermore, $E \otimes_A K$ can be identified with $E_0 \otimes_A K$, and $u \otimes 1$ with $u_0 \otimes 1$, and so $\sigma_i(u) = \sigma_i(u_0)$ for $1 \leq i \leq n$.

If E is *torsion free*, then E is canonically identified with a sub- A -module of $E \otimes_A K$, and the relation $u \otimes 1 = 0$ is equivalent to $u = 0$. If E is a free A -module, then the two definitions of $\sigma_i(u)$ given in (6.4.1) and (6.4.2) coincide, by the above, which justifies the notation used.

We also note that, if E is a *torsion module*, or, in other words, if $E_0 = \{0\}$, then the exterior algebra of E_0 consists of K , and the determinant of the unique endomorphism u_0 of E_0 is equal to 1.

Proposition (6.4.3). — *Let A be an integral ring, E an A -module of finite type, and u an endomorphism of E ; then the elementary symmetric functions $\sigma_i(u)$ of u (and, in particular, $\det u$) are elements of K that are integral over A .*

Proof. Let A' be the integral closure of A ; since $A'[T]$ is an integrally closed ring [24, p. 99], it is the integral closure of $A[T]$ in its field of fractions $K(T)$. Replacing u by $T \cdot I - u \otimes 1$, and A by $A[T]$, we see that we can reduce to proving that $\det u$ is integral over A . If n is the rank of E , then $\det(u) = \det(\wedge^n u)$ and $(\wedge^n u) \otimes 1 = \wedge^n(u \otimes 1)$, so we can suppose that $n = 1$. But then the map $u \mapsto \det u$ is a homomorphism from the A -module $\text{Hom}_A(E, E)$ to K ; since E is of finite type, $\text{Hom}_A(E, E)$ is isomorphic to a sub- A -module of the A -module of finite type E^m if E admits a system of m generators, and so the elements $\det u$ belong to a sub- A -module of K of finite type, and are thus integral over A . $\square$

[II:122]

Corollary (6.4.4). — Under the hypotheses of (6.4.3), if we further suppose A to be normal, then the $\sigma_i(u)$ (and, in particular, $\text{Tr } u$ and $\det u$) belong to A .

Proposition (6.4.5). — Let A be an integral ring, E an A -module of finite type of rank n , and u an endomorphism of E such that the $\sigma_i(u)$ belong to A . For u to be an automorphism of E , it is necessary that $\det u$ be invertible in A ; this condition is sufficient if E is torsion free.

Proof. The condition is sufficient when E is torsion free, since the hypothesis and equations (6.4.1.4) and (6.4.1.5) (which hold in E , and not just in $E \otimes_A K$, since E is torsion free) prove that $(\det u)^{-1}Q(u)$ is the inverse of u .

The condition is necessary, since, if u is invertible, then it follows from (6.4.3) that $\det(u^{-1})$ belongs to the integral closure A' of A in its field of fractions K , and is evidently the inverse of $\det u$ in A' . Our claim then follows from: $\square$

Lemma (6.4.5.1). — Let A be a subring of a ring A' such that A' is integral over A . If an element $x \in A$ is invertible in A' , then it is also invertible in A .

Proof. In the contrary case, x would belong to a maximal ideal $\mathfrak{m}$ of A , and it follows from the first theorem of Cohen–Seidenberg [13, t. I, p. 257, th. 3] that there would be a maximal ideal $\mathfrak{m}'$ of A' such that $\mathfrak{m} = \mathfrak{m}' \cap A$; we would then have that $x \in \mathfrak{m}'$, which is a contradiction. $\square$

Corollary (6.4.6). — Let A be an integral and integrally closed ring, E a torsion-free A -module of finite type, and u an endomorphism of E . For u to be an automorphism of E , it is necessary and sufficient that $\det u$ be invertible in A .

Proof. This follows from (6.4.4) and (6.4.5). $\square$

Remark (6.4.7). — We will later need a generalisation of the above results. Consider a reduced Noetherian ring A ; let $\mathfrak{p}_\alpha$ (for $1 \leq \alpha \leq r$) be its minimal prime ideals, K_α the field of fractions of the integral ring $A/\mathfrak{p}_\alpha$, and K the total ring of fractions of A , which is the direct sum of the fields K_α . Let E be an A -module of finite type, and suppose that $E \otimes_A K$ is a free K -module of rank n (which here is no longer a consequence of the other hypotheses); equivalently, we can ask that all the K_α -vector spaces $E \otimes_A K_\alpha = E_\alpha$ have the same dimension n ; if then u is an endomorphism of E , we again set $P(u, T) = P(u \otimes 1, T)$ and $\sigma_j(u) = \sigma_j(u \otimes 1)$, and, in particular, $\det u = \det(u \otimes 1)$; the $\sigma_j(u)$ are thus elements of K . It is immediate that $E \otimes_A K$ is the direct sum of the E_α , and that the latter are stable under $u \otimes 1$; the restriction of $u \otimes 1$ to E_α is exactly the extension u_α of u to this K_α -vector space; we thus conclude that $\sigma_j(u)$ is the element of K whose components in the K_α are the $\sigma_j(u_\alpha)$. Since the integral closure of A in K is the direct sum of the integral closures of A in the K_α , the $\sigma_j(u)$ are integral over A .

Lemma (6.4.7.1). — The sub- A -algebra of K generated by all the elements $\sigma_j(u)$ (for $1 \leq j \leq n$), where u runs over $\text{Hom}_A(E, E)$, is an A -module of finite type.

Proof. It suffices to prove that the sub- $A[T]$ -algebra of $K[T]$ generated by the $P(u, T)$ is an $A[T]$ -module of finite type, since if the $F_i(T)$ (for $1 \leq i \leq m$) form a system of generators for this $A[T]$ -module, then the coefficients of the $F_i(T)$ are integral over A , and thus generate an A -algebra that is an A -module of finite type [13, t. I, p. 255, th. 1]. [II:123] We can thus replace A by $A[T]$ (which is Noetherian), and E by $E \otimes_A A[T] = E'$, which is such that $E' \otimes_{A[T]} K[T] = E \otimes_A K[T]$ is a free $K[T]$ -module of rank n . Using the initial notation, we thus see that it suffices to prove that the A -module generated by the elements $\det u$, where u runs over $\text{Hom}_A(E, E)$, is of finite type; a fortiori (since every submodule of an A -module of finite type is of finite type) it suffices to prove that, as v runs over the set of endomorphisms of $\wedge^n E$, the A -module generated by the $\det v$ is of finite type; in other words, we can again reduce to the case where $n = 1$. But then the proposition follows from the fact that $\text{Hom}_A(E, E)$ is an A -module of finite type, and that $v \mapsto \det v$ is a homomorphism from this A -module into K . $\square$

Let F be the kernel of the canonical homomorphism $E \rightarrow E \otimes_A K$, and let $E_0 = E/F$; we see, as above, that $E \otimes_A K$ can be identified with $E_0 \otimes_A K$, that $u(F) \subset F$, and that, if u_0 is the endomorphism of E_0 induced from u by passing to the quotient, then $u \otimes 1$ can be identified with $u_0 \otimes 1$, and

$\sigma_j(u) = \sigma_j(u_0)$ for all j . If we have $F = 0$, then equations (6.4.1.3) and (6.4.1.5) still make sense and hold true whenever E is identified with a submodule of $E \otimes_A K$, and the u^j with homomorphisms $E \rightarrow E \otimes_A K$; then Proposition (6.4.5) extends also to this case, with the same proof.

(6.4.8). Let $(X, \mathcal{A})$ be a ringed space, $\mathcal{E}$ a locally free $\mathcal{A}$ -module (of finite rank), and u an endomorphism of $\mathcal{E}$. By hypothesis, there exists a base $\mathfrak{B}$ for the topology of X such that, for all $V \in \mathfrak{B}$, $\mathcal{E}|_V$ is isomorphic to some $\mathcal{A}^n|_V$ (where n may vary with V). For all $V \in \mathfrak{B}$, u_V is thus an endomorphism of the $\Gamma(V, \mathcal{A})$ -module $\Gamma(V, \mathcal{E})$, which is free by hypothesis; the determinant $\det u_V$ is thus defined and belongs to $\Gamma(V, \mathcal{A})$. Furthermore, if $e_1, \dots, e_n$ form a basis of $\Gamma(V, \mathcal{E})$, then their restrictions to every open $W \subset V$ form a basis of $\Gamma(W, \mathcal{E})$ over $\Gamma(W, \mathcal{A})$, and so $\det u_W$ is the restriction of $\det u_V$ to W . There thus exists exactly one section of $\mathcal{A}$ over X , which we denote by $\det u$, and call the *determinant* of u , such that the restriction of $\det u$ to each $V \in \mathfrak{B}$ is $\det u_V$. It is clear that, for all $x \in X$, we have $(\det u)_x = \det u_x$; for endomorphisms u and v of $\mathcal{E}$, we have

$$(6.4.8.1) \quad \det(u \circ v) = (\det u)(\det v)$$

as well as

$$(6.4.8.2) \quad \det(1_{\mathcal{E}}) = 1_{\mathcal{A}}$$

and, if the rank of $\mathcal{E}$ is constant (which will be the case (0, 5.4.1) if X is connected) and equal to n ,

$$(6.4.8.3) \quad \det(s \cdot u) = s^n \det u$$

for all $s \in \Gamma(X, \mathcal{A})$ (note that $\det(0) = 0_{\mathcal{A}}$ if $n \geq 1$, but $\det(0) = 1_{\mathcal{A}}$ for $n = 0$). Furthermore, for u to be an *automorphism* of $\mathcal{E}$, it is necessary and sufficient that $\det u$ be *invertible* in $\Gamma(X, \mathcal{A})$.

If the rank of $\mathcal{E}$ is constant, then we can similarly define the elementary symmetric functions $\sigma_i(u)$, which are elements of $\Gamma(X, \mathcal{A})$; we again have the relations in (6.4.1.3) and (6.4.1.5).

[II:124] We have thus defined a homomorphism $\det : \text{Hom}_{\mathcal{A}}(\mathcal{E}, \mathcal{E}) \rightarrow \Gamma(X, \mathcal{A})$ of multiplicative monoids; if we note that $\text{Hom}_{\mathcal{A}}(\mathcal{E}, \mathcal{E}) = \Gamma(X, \mathcal{H}\text{om}_{\mathcal{A}}(\mathcal{E}, \mathcal{E}))$ by definition, then we see that we can replace X in this definition by an arbitrary open U of X , which immediately defines a homomorphism $\det : \mathcal{H}\text{om}_{\mathcal{A}}(\mathcal{E}, \mathcal{E}) \rightarrow \mathcal{A}$ of sheaves of multiplicative monoids. If $\mathcal{E}$ has constant rank, then we similarly define homomorphisms $\sigma_i : \mathcal{H}\text{om}_{\mathcal{A}}(\mathcal{E}, \mathcal{E}) \rightarrow \mathcal{A}$ of sheaves of sets; for $i = 1$, the homomorphism $\sigma_1 = \text{Tr}$ is a homomorphism of $\mathcal{A}$ -modules.

Let $(Y, \mathcal{B})$ be another ringed space, and let $f : (X, \mathcal{A}) \rightarrow (Y, \mathcal{B})$ be a morphism of ringed spaces; if $\mathcal{F}$ is a locally free $\mathcal{B}$ -module, then $f^*(\mathcal{F})$ is a locally free $\mathcal{A}$ -module (which is of the same rank as $\mathcal{F}$ if the latter is of constant rank) (0, 5.4.5). For every endomorphism v of $\mathcal{F}$, $f^*(v)$ is an endomorphism of $f^*(\mathcal{F})$, and it follows immediately from the definitions that $\det f^*(v)$ is the section of $\mathcal{A} = f^*(\mathcal{B})$ over X that canonically corresponds to $\det v \in \Gamma(Y, \mathcal{B})$. We can further say that the homomorphism $f^*(\det) : f^*(\mathcal{H}\text{om}_{\mathcal{B}}(\mathcal{F}, \mathcal{F})) \rightarrow f^*(\mathcal{B}) = \mathcal{A}$ is the composition

$$f^*(\mathcal{H}\text{om}_{\mathcal{B}}(\mathcal{F}, \mathcal{F})) \xrightarrow{\gamma^{\sharp}} \mathcal{H}\text{om}_{\mathcal{A}}(f^*(\mathcal{F}), f^*(\mathcal{F})) \xrightarrow{\det} \mathcal{A}$$

(0, 4.4.6). We have analogous results for the σ_i .

(6.4.9). Now suppose that X is a *locally integral prescheme*, so that the sheaf $\mathcal{R}(X)$ of rational functions on X is a locally simple sheaf of fields (I, 7.3.4), and quasi-coherent as an $\mathcal{O}_X$ -module. If $\mathcal{E}$ is a quasi-coherent $\mathcal{O}_X$ -module of *finite type*, then $\mathcal{E}' = \mathcal{E} \otimes_{\mathcal{O}_X} \mathcal{R}(X)$ is a locally free $\mathcal{R}(X)$ -module (I, 7.3.6); for every endomorphism u of $\mathcal{E}$, $u \otimes 1_{\mathcal{R}(X)}$ is then an endomorphism of $\mathcal{E}'$, and $\det(u \otimes 1)$ is a section of $\mathcal{R}(X)$ over X , which we also call the *determinant* of u , and also denote by $\det u$. It follows from (6.4.3) that $\det u$ is a section of the *integral closure* of $\mathcal{O}_X$ in $\mathcal{R}(X)$ (6.3.2); if, further, X is *normal*, then $\det u$ is a section of $\mathcal{O}_X$ over X , and if we further suppose that $\mathcal{E}$ is *torsion free*, then for u to be an automorphism of $\mathcal{E}$ it is necessary and sufficient that $\det u$ be *invertible*, by (6.4.6). Equations (6.4.8.1) to (6.4.8.3) still hold true; from the homomorphism $u \mapsto \det u$, applied to the modules of sections of $\mathcal{H}\text{om}_{\mathcal{O}_X}(\mathcal{E}, \mathcal{E})$, we obtain a homomorphism of sheaves $\det : \mathcal{H}\text{om}_{\mathcal{O}_X}(\mathcal{E}, \mathcal{E}) \rightarrow \mathcal{R}(X)$, which takes values in $\mathcal{O}_X$ if X is normal. We have analogous definitions and results for the other elementary symmetric functions $\sigma_j(u)$, if $\mathcal{E}'$ is of *constant rank*; if, further, X is *normal*, then the $\sigma_j(u)$ are sections of $\mathcal{O}_X$ over X .

Finally, let X and Y be integral preschemes, and $f : X \rightarrow Y$ a *dominant* morphism. We know that there exists a canonical homomorphism $f^*(\mathcal{R}(Y)) \rightarrow \mathcal{R}(X)$ (I, 7.3.8), whence we obtain, for every

quasi-coherent $\mathcal{O}_Y$ -module $\mathcal{F}$ of finite type, a canonical homomorphism $\theta : f^*(\mathcal{F} \otimes_{\mathcal{O}_Y} \mathcal{R}(Y)) \rightarrow f^*(\mathcal{F}) \otimes_{\mathcal{O}_X} \mathcal{R}(X)$. [II:125] If v is an endomorphism of $\mathcal{F}$, then $f^*(v \otimes 1_{\mathcal{R}(Y)})$ is an endomorphism of $f^*(\mathcal{F} \otimes_{\mathcal{O}_Y} \mathcal{R}(Y))$, and we have a commutative diagram

$$\begin{array}{ccc} f^*(\mathcal{F} \otimes_{\mathcal{O}_Y} \mathcal{R}(Y)) & \xrightarrow{f^*(v \otimes 1)} & f^*(\mathcal{F} \otimes_{\mathcal{O}_Y} \mathcal{R}(Y)) \\ \theta \downarrow & & \downarrow \theta \\ f^*(\mathcal{F}) \otimes_{\mathcal{O}_X} \mathcal{R}(X) & \xrightarrow{f^*(v) \otimes 1} & f^*(\mathcal{F}) \otimes_{\mathcal{O}_X} \mathcal{R}(X) \end{array}$$

We thus easily conclude that $\det f^*(v)$ is the canonical image, under the homomorphism $f^*(\mathcal{R}(Y)) \rightarrow \mathcal{R}(X)$, of the section $\det v$ of $\mathcal{R}(Y)$; indeed, we can immediately reduce to the case where $X = \text{Spec}(A)$ and $Y = \text{Spec}(B)$ are affine, with A and B integral rings having respective fields of fractions K and L , and with the homomorphism $B \rightarrow A$ injective and therefore extending to a monomorphism $L \rightarrow K$; if $\mathcal{F} = \tilde{M}$, where M is a B -module of finite type, then the rank of $M \otimes_B L$ over L is equal to that of $(M \otimes_B A) \otimes_A K$ over K , and $\det((u \otimes 1) \otimes 1)$ is the image in K of $\det(u \otimes 1)$ for any endomorphism u of M , whence the conclusion.

(6.4.10). Finally, suppose that X is a *locally Noetherian reduced prescheme*, so that the sheaf $\mathcal{R}(X)$ of rational functions on X is again a quasi-coherent $\mathcal{O}_X$ -module (I, 7.3.4); let $\mathcal{E}$ be a *coherent* $\mathcal{O}_X$ -module such that $\mathcal{E}' = \mathcal{E} \otimes_{\mathcal{O}_X} \mathcal{R}(X)$ is *locally free* (of finite rank). By (6.4.7), if $\mathcal{E}'$ is of constant rank, then we can, for any endomorphism u of $\mathcal{E}$, define the $\sigma_j(u)$, which are sections of $\mathcal{R}(X)$ over X . If we do not suppose $\mathcal{E}'$ to be of constant rank, we can still define the homomorphism $\det : \mathcal{H}om_{\mathcal{O}_X}(\mathcal{E}, \mathcal{E}) \rightarrow \mathcal{R}(X)$.

6.5. Norm of an invertible sheaf

(6.5.1). Let $(X, \mathcal{A})$ be a ringed space and $\mathcal{B}$ a (commutative) $\mathcal{A}$ -algebra. The $\mathcal{A}$ -module $\mathcal{B}$ is canonically identified with a sub- $\mathcal{A}$ -module of $\mathcal{H}om_{\mathcal{A}}(\mathcal{B}, \mathcal{B})$, and a section f of $\mathcal{B}$ over an open U of X is identified with multiplication by this section. If $(X, \mathcal{A})$ and $\mathcal{B}$ satisfy one of the conditions given in (6.4.8), (6.4.9), or (6.4.10), then we can define $\det(f)$ (and, in certain cases, the $\sigma_j(f)$) as sections of $\mathcal{A}$ or $\mathcal{R}(X)$ over U ; we call these the *norm* of f (resp. the *elementary symmetric functions* of f), and denote the former by $N_{\mathcal{B}/\mathcal{A}}(f)$. We will suppose that *one* of the following conditions is satisfied:

- (I) $\mathcal{B}$ is a *locally free* $\mathcal{A}$ -module (of finite rank).
- (II) $(X, \mathcal{A})$ is a *locally Noetherian reduced prescheme*, $\mathcal{B}$ is a *coherent* $\mathcal{A}$ -module such that $\mathcal{B} \otimes_{\mathcal{A}} \mathcal{R}(X)$ is a *locally free* $\mathcal{R}(X)$ -module, and, for every section $f \in \Gamma(U, \mathcal{B})$ over an open U of X , the norm $N_{\mathcal{B}/\mathcal{A}}(f)$ is a section of $\mathcal{A}$ over U .

[II:126] Note that the latter condition is automatically satisfied whenever the locally Noetherian prescheme X is *normal* (6.4.9).

The hypothesis that $\mathcal{B} \otimes_{\mathcal{A}} \mathcal{R}(X)$ is locally free can also be expressed in the following way: denote by X_α the closed reduced subpreschemes of X whose underlying spaces are the irreducible components of X (I, 5.2.1), which are thus locally Noetherian integral preschemes. Every $x \in X$ belongs to a finite number of the subspaces X_α ; on the other hand, $\mathcal{B} \otimes_{\mathcal{A}} \mathcal{R}(X_\alpha)$ is a locally free $\mathcal{R}(X_\alpha)$ -module of constant rank k_α (I, 7.3.6); to say that $\mathcal{B} \otimes_{\mathcal{A}} \mathcal{R}(X)$ is a locally free $\mathcal{R}(X)$ -module means that, for all $x \in X$, the ranks k_α that correspond to the indices such that $x \in X_\alpha$ are equal. This is a local statement, and we can reduce to the case $X = \text{Spec}(C)$, where C is a reduced Noetherian ring, and $\mathcal{B} = \tilde{D}$, where D is a C -algebra that is a C -module of finite type; if $\mathfrak{p}_i$ (for $1 \leq i \leq m$) are the minimal prime ideals of C , then the total ring of fractions L of C is the direct sum of the fields of fractions K_i of the integral rings $C/\mathfrak{p}_i$, and $D \otimes_C L$ is the direct sum of the $D \otimes_C K_i$, whence the conclusion.

It is clear that, under hypothesis (I) or (II), we have thus defined a *homomorphism of sheaves of multiplicative monoids* $N_{\mathcal{B}/\mathcal{A}} : \mathcal{B} \rightarrow \mathcal{A}$, also denoted by N if no confusion may arise, and we call this homomorphism the *norm*. For sections f and g of $\mathcal{B}$ over the same open U , we thus have

$$(6.5.1.1) \quad N_{\mathcal{B}/\mathcal{A}}(fg) = N_{\mathcal{B}/\mathcal{A}}(f)N_{\mathcal{B}/\mathcal{A}}(g)$$

for the corresponding sections of $\mathcal{A}$ over U ;

$$(6.5.1.2) \quad N_{\mathcal{B}/\mathcal{A}}(1_{\mathcal{B}}) = 1_{\mathcal{A}};$$

finally, for any section s of $\mathcal{A}$ over U ,

$$(6.5.1.3) \quad N_{\mathcal{B}/\mathcal{A}}(s \cdot 1_{\mathcal{B}}) = s^n$$

if the rank of $\mathcal{B}$ is constant and equal to n (for $s = 0_{\mathcal{A}}$, this formula gives $N(0_{\mathcal{B}}) = 0_{\mathcal{A}}$ if $n \geq 1$, and $N(0_{\mathcal{B}}) = N(1_{\mathcal{B}}) = 1_{\mathcal{A}}$ if $n = 0$).

In hypothesis (I), for $f \in \Gamma(U, \mathcal{B})$ to be invertible it is necessary and sufficient that $N(f) \in \Gamma(U, \mathcal{A})$ be invertible. In hypothesis (II), this condition is necessary; it is sufficient (by (6.4.7)) if we suppose that $\mathcal{B} \rightarrow \mathcal{B} \otimes_{\mathcal{A}} \mathcal{R}(X)$ is *injective* and that the following more restrictive hypothesis is satisfied:

(II bis) $(X, \mathcal{A})$ is a locally Noetherian reduced prescheme, $\mathcal{B}$ is a coherent $\mathcal{A}$ -module such that $\mathcal{B} \otimes_{\mathcal{A}} \mathcal{R}(X)$ is a locally free $\mathcal{R}(X)$ -module, and, for any section $f \in \Gamma(U, \mathcal{B})$ over an open U such that $\mathcal{B} \otimes_{\mathcal{A}} \mathcal{R}(X)|_U$ is of constant rank n on $\mathcal{R}(X)|_U$, the $\sigma_j(f)$ (for $1 \leq j \leq n$) are sections of $\mathcal{A}$ over U .

(We again note that this condition is satisfied if X is *normal*.)

(6.5.2). Suppose that one of the hypotheses (I) or (II) of (6.5.1) is satisfied, and let $\mathcal{L}'$ be an invertible $\mathcal{B}$ -module. We will canonically associate to it (up to unique isomorphism) an invertible $\mathcal{A}$ -module $\mathcal{L}$ in the following way. Denote by $\mathcal{A}^*$ (resp. $\mathcal{B}^*$) the subsheaf of $\mathcal{A}$ (resp. of $\mathcal{B}$) such that $\Gamma(U, \mathcal{A}^*)$ (resp. $\Gamma(U, \mathcal{B}^*)$) is the set of invertible elements of $\Gamma(U, \mathcal{A})$ (resp. of $\Gamma(U, \mathcal{B})$) for every open $U \subset X$; [II:127] these are sheaves of multiplicative groups, and $N_{\mathcal{B}/\mathcal{A}}$ restricted to $\mathcal{B}^*$ is a homomorphism $\mathcal{B}^* \rightarrow \mathcal{A}^*$ of sheaves of groups (6.5.1). Let $\mathfrak{L}$ be the set of pairs (U_λ, ν_λ) that have the following property: U_λ is an open of X , and ν_λ is an isomorphism $\mathcal{L}'|_{U_\lambda} \xrightarrow{\sim} \mathcal{B}|_{U_\lambda}$ of $(\mathcal{B}|_{U_\lambda})$ -modules. By hypothesis, the U_λ form a cover of X ; for arbitrary indices λ and μ , set $\omega_{\lambda\mu} = (\nu_\lambda|_{U_\lambda \cap U_\mu}) \circ (\nu_\mu|_{U_\lambda \cap U_\mu})^{-1}$, which is an automorphism of $\mathcal{B}|_{U_\lambda \cap U_\mu}$ canonically identified with a section of $\mathcal{B}^*$ over $U_\lambda \cap U_\mu$, and $(\omega_{\lambda\mu})$ is a 1-cocycle for the cover $\mathfrak{U} = (U_\lambda)$ with values in $\mathcal{B}^*$ (0, 5.4.7). The fact that $N_{\mathcal{B}/\mathcal{A}} : \mathcal{B}^* \rightarrow \mathcal{A}^*$ is a homomorphism implies that $(N_{\mathcal{B}/\mathcal{A}}\omega_{\lambda\mu})$ is a 1-cocycle of $\mathfrak{U}$ with values in $\mathcal{A}^*$, and thus corresponds (up to unique isomorphism) to an invertible $\mathcal{A}$ -module $\mathcal{L}$ (0, 5.4.7) which we denote by $N_{\mathcal{B}/\mathcal{A}}(\mathcal{L}')$, and call the *norm* of the invertible $\mathcal{B}$ -module $\mathcal{L}'$.

Let $\mathfrak{M}$ be a subset of $\mathfrak{L}$ such that the corresponding U_λ still form a cover of X , and let $\mathfrak{B}$ be the resulting cover; the restriction of the cocycle $(\omega_{\lambda\mu})$ to $\mathfrak{B}$ defines a 1-cocycle $(N_{\mathcal{B}/\mathcal{A}}\omega_{\lambda\mu})$ of $\mathfrak{B}$ with values in $\mathcal{A}^*$, the restriction of the 1-cocycle $(N_{\mathcal{B}/\mathcal{A}}\omega_{\lambda\mu})$ of $\mathfrak{U}$; it is clear that there is a canonical isomorphism from the invertible $\mathcal{A}$ -module defined by this 1-cocycle of $\mathfrak{B}$ to $N_{\mathcal{B}/\mathcal{A}}(\mathcal{L}')$, which allows us to define the invertible $\mathcal{A}$ -module by an arbitrary sub-cover of $\mathfrak{U}$. This possibility immediately shows that, if $\mathcal{L}'_1$ and $\mathcal{L}'_2$ are invertible $\mathcal{B}$ -modules, then, by (6.5.1.1) and (6.5.1.2),

$$(6.5.2.1) \quad N(\mathcal{L}'_1 \otimes_{\mathcal{B}} \mathcal{L}'_2) = N(\mathcal{L}'_1) \otimes_{\mathcal{A}} N(\mathcal{L}'_2)$$

and

$$(6.5.2.2) \quad N_{\mathcal{B}/\mathcal{A}}(\mathcal{B}) = \mathcal{A}$$

as well as

$$(6.5.2.3) \quad N(\mathcal{L}'^{-1}) = (N(\mathcal{L}'))^{-1}$$

up to canonical isomorphisms. Furthermore, it follows from (6.5.1.3) that, if $\mathcal{L}$ is an invertible $\mathcal{A}$ -module and if $\mathcal{B}$ is of constant rank n on $\mathcal{A}$ in case (I) (resp. $\mathcal{B} \otimes_{\mathcal{A}} \mathcal{R}(X)$ is of constant rank n on $\mathcal{R}(X)$ in case (II)), we have, up to canonical isomorphism,

$$(6.5.2.4) \quad N_{\mathcal{B}/\mathcal{A}}(\mathcal{L} \otimes_{\mathcal{A}} \mathcal{B}) = \mathcal{L}^{\otimes n}.$$

(6.5.3). We now show that $N_{\mathcal{B}/\mathcal{A}}$ is a covariant functor in the category of invertible $\mathcal{B}$ -modules. Let $h' : \mathcal{L}'_1 \rightarrow \mathcal{L}'_2$ be a homomorphism of invertible $\mathcal{B}$ -modules, and let $\mathfrak{B} = (U_\lambda)$ be an open cover of X such that, for all λ , we have $(\mathcal{B}|_{U_\lambda})$ -isomorphisms $\eta_\lambda^{(1)} : \mathcal{L}'_1|_{U_\lambda} \xrightarrow{\sim} \mathcal{B}|_{U_\lambda}$ and $\eta_\lambda^{(2)} : \mathcal{L}'_2|_{U_\lambda} \xrightarrow{\sim} \mathcal{B}|_{U_\lambda}$; there is thus, for each λ , an endomorphism h'_λ of $\mathcal{B}|_{U_\lambda}$ such that $h'_\lambda \circ \eta_\lambda^{(1)} = \eta_\lambda^{(2)} \circ (h'|_{U_\lambda})$, and we can evidently identify h'_λ with a section of $\mathcal{B}$ over U_λ (0, 5.1.1). So, for every pair (λ, μ)

of indices, the restrictions to $U_\lambda \cap U_\mu$ of $(\eta_\lambda^{(2)})^{-1} \circ h'_\lambda \circ \eta_\lambda^{(1)}$ and $(\eta_\mu^{(2)})^{-1} \circ h'_\mu \circ \eta_\mu^{(1)}$ agree. We thus obtain, for 1-cocycles $(\omega_{\lambda\mu}^{(1)})$ and $(\omega_{\lambda\mu}^{(2)})$ with values in $\mathcal{B}^*$ corresponding to $\mathcal{L}'_1$ and $\mathcal{L}'_2$, the relation

$$\omega_{\lambda\mu}^{(2)} h'_\mu = h'_\lambda \omega_{\lambda\mu}^{(1)}.$$

[II:128] If we set $h_\lambda = N(h'_\lambda)$, we thus have the analogous relations

$$N(\omega_{\lambda\mu}^{(2)}) h_\mu = h_\lambda N(\omega_{\lambda\mu}^{(1)})$$

and so the h_λ define a homomorphism $N(\mathcal{L}'_1) \rightarrow N(\mathcal{L}'_2)$ which we denote by $N_{\mathcal{B}/\mathcal{A}}(h')$, or $N(h')$. In hypothesis (I), for h' to be an *isomorphism*, it is necessary and sufficient (since it is a local question) for $N_{\mathcal{B}/\mathcal{A}}(h')$ to be an isomorphism. In hypothesis (II), this condition is again necessary; it is sufficient if hypothesis (II bis) is satisfied and $\mathcal{B} \rightarrow \mathcal{B} \otimes_{\mathcal{A}} \mathcal{R}(X)$ is injective.

Take, in particular, $\mathcal{L}'_1 = \mathcal{B}$; the homomorphisms $\mathcal{B} \rightarrow \mathcal{L}'$ can then be identified (0, 5.1.1) with the sections of $\mathcal{L}'$ over X , whence a canonical map

$$N_{\mathcal{B}/\mathcal{A}} : \Gamma(X, \mathcal{L}') \longrightarrow \Gamma(X, N_{\mathcal{B}/\mathcal{A}}(\mathcal{L}')).$$

It again follows from (6.5.1.1) that, if $f'_1 \in \Gamma(X, \mathcal{L}'_1)$ and $f'_2 \in \Gamma(X, \mathcal{L}'_2)$, then

$$(6.5.3.1) \quad N(f'_1 \otimes f'_2) = N(f'_1) \otimes N(f'_2).$$

For every invertible $\mathcal{A}$ -module $\mathcal{L}$ and every section $f \in \Gamma(X, \mathcal{L})$, we have, taking (6.5.2.4) into account, that

$$(6.5.3.2) \quad N_{\mathcal{B}/\mathcal{A}}(f \otimes 1_{\mathcal{B}}) = f^{\otimes n}$$

whenever $\mathcal{B}$ is of constant rank n in hypothesis (I) (resp. whenever $\mathcal{B} \otimes_{\mathcal{A}} \mathcal{R}(X)$ is of constant rank n in hypothesis (II)). Finally, for the homomorphism $\mathcal{B} \rightarrow \mathcal{L}'$ corresponding to a section f' of $\mathcal{L}'$ over X to be an isomorphism, it is necessary and sufficient for f'_x to be a basis for $\mathcal{L}'_x$ for all $x \in X$; in hypothesis (I), this condition is thus equivalent to saying that $(N(f'))_x$ is a basis for $(N(\mathcal{L}'))_x$ for all x ; in hypothesis (II), this condition is again necessary, and it is sufficient whenever $\mathcal{B}$ satisfies hypothesis (II bis) and $\mathcal{B} \rightarrow \mathcal{B} \otimes_{\mathcal{A}} \mathcal{R}(X)$ is injective.

(6.5.4). Let $(X, \mathcal{A})$ and $(X', \mathcal{A}')$ be ringed spaces, $f : X' \rightarrow X$ a morphism, $\mathcal{B}$ an $\mathcal{A}$ -algebra, and $\mathcal{B}' = f^*(\mathcal{B})$ the inverse image $\mathcal{A}'$ -algebra. Suppose that one of the following hypotheses is satisfied:

- (1) $\mathcal{B}$ satisfies hypothesis (I) of (6.5.1).
- (2) $(X, \mathcal{A})$ and $\mathcal{B}$ satisfy hypothesis (II) of (6.5.1), $(X', \mathcal{A}')$ is a locally Noetherian reduced prescheme, and, if we denote by X_α and X'_β the closed reduced subpreschemes of X and X' (respectively) that have the irreducible components of these spaces as their underlying spaces, then the restriction of f to each X'_β is a *dominant* morphism from X'_β to X_α .

Under these conditions, $\mathcal{B}'$ satisfies hypothesis (I) or hypothesis (II) (respectively) of (6.5.1). The first claim is immediate; to establish the second, it suffices to show that, for all $x' \in X'$, the ranks of the $\mathcal{B}' \otimes_{\mathcal{O}_{X'}} \mathcal{R}(X'_\beta)$ are the same for the β such that $x' \in X'_\beta$. But if the restriction of f to X'_β is a dominant morphism to X_α , then the rank of $\mathcal{B}' \otimes_{\mathcal{O}_{X'}} \mathcal{R}(X'_\beta)$ is equal to that of $\mathcal{B} \otimes_{\mathcal{O}_X} \mathcal{R}(X_\alpha)$ (as we immediately see by reducing to the affine case, as in (6.4.9)), whence our claim, by hypothesis (II) and (6.5.1).

[II:129] With this in mind, it follows from (6.4.8), (6.4.9), and (6.4.10) that, if s is a section of $\mathcal{B}$ over an open $U \subset X$, and s' the corresponding section of $\mathcal{B}'$ over $f^{-1}(U)$, then $N_{\mathcal{B}'/\mathcal{A}'}(s')$ is the section of $\mathcal{A}'$ over $f^{-1}(U)$ that corresponds to the section $N_{\mathcal{B}/\mathcal{A}}(s)$ of $\mathcal{A}$ over U .

If $\mathcal{M}$ is an invertible $\mathcal{B}$ -module, then we deduce from the above that, if $\mathcal{M}' = f^*(\mathcal{M})$ (which is an invertible $\mathcal{B}'$ -module), we have $N_{\mathcal{B}'/\mathcal{A}'}(\mathcal{M}') = f^*(N_{\mathcal{B}/\mathcal{A}}(\mathcal{M}))$ up to canonical isomorphism.

(6.5.5). Suppose from now on that $(X, \mathcal{A})$ is a *prescheme*. The data of a quasi-coherent $\mathcal{A}$ -algebra $\mathcal{B}$, which is an $\mathcal{A}$ -module of finite type, is then equivalent, as we know, to that of a *finite* morphism $g : X' \rightarrow X$ such that $g_*(\mathcal{O}_{X'}) = \mathcal{B}$, defined up to X -isomorphism ((6.1.2) and (1.3.1)). Furthermore, the data of a quasi-coherent $\mathcal{O}_{X'}$ -module $\mathcal{F}'$ is equivalent to that of a quasi-coherent $\mathcal{B}$ -module $\mathcal{F}$ such that $g_*(\mathcal{F}') = \mathcal{F}$ (1.4.3), and for $\mathcal{F}'$ to be invertible it is necessary and sufficient that $\mathcal{F}$ be

invertible (6.1.12). To translate the above results in terms of finite morphisms g , it will be necessary to suppose either that $g_*(\mathcal{O}_{X'})$ is a *locally free* $\mathcal{O}_X$ -module (of finite type) or that $(X, \mathcal{O}_X)$ and $g_*(\mathcal{O}_{X'})$ satisfy hypothesis (II). For every invertible $\mathcal{O}_{X'}$ -module $\mathcal{L}'$, we thus set

$$(6.5.5.1) \quad N_{X'/X}(\mathcal{L}') = N_{g_*(\mathcal{O}_{X'})/\mathcal{O}_X}(g_*(\mathcal{L}'))$$

and we call this the *norm* (with respect to g) of $\mathcal{L}'$. Similarly, if $h' : \mathcal{L}'_1 \rightarrow \mathcal{L}'_2$ is a homomorphism of invertible $\mathcal{O}_{X'}$ -modules, then we set

$$(6.5.5.2) \quad N_{X'/X}(h') = N_{g_*(\mathcal{O}_{X'})/\mathcal{O}_X}(g_*(h')) : N_{X'/X}(\mathcal{L}'_1) \rightarrow N_{X'/X}(\mathcal{L}'_2).$$

In particular, for $\mathcal{L}'_1 = \mathcal{O}_{X'}$, we thus obtain a canonical map

$$(6.5.5.3) \quad N_{X'/X} : \Gamma(X', \mathcal{L}') \rightarrow \Gamma(X, N_{X'/X}(\mathcal{L}')).$$

We leave to the reader the majority of these translations, and we restrict ourselves to spelling out the details of the following:

Proposition (6.5.6). — *Let $g : X' \rightarrow X$ be a finite morphism, and suppose that either $g_*(\mathcal{O}_{X'})$ is a locally free $\mathcal{O}_X$ -module or that $(X, \mathcal{O}_X)$ and $g_*(\mathcal{O}_{X'})$ satisfy (II bis) (which will be the case, in particular, if X is locally Noetherian and normal). For a homomorphism $h' : \mathcal{L}'_1 \rightarrow \mathcal{L}'_2$ of invertible $\mathcal{O}_{X'}$ -modules to be an isomorphism, it is necessary and sufficient, in the first hypothesis, that $N_{X'/X}(h')$ be an isomorphism; in the second hypothesis, this condition is again necessary, and is sufficient if the homomorphism $g_*(\mathcal{O}_{X'}) \rightarrow g_*(\mathcal{O}_{X'}) \otimes_{\mathcal{O}_X} \mathcal{R}(X)$ is injective.*

Proof. Note that we use here the fact that, for $g_*(h')$ to be an isomorphism, it is necessary and sufficient for h' to be an isomorphism (1.4.2). $\square$

Corollary (6.5.7). — *Let $g : X' \rightarrow X$ be a finite morphism, and suppose that either $g_*(\mathcal{O}_{X'})$ is a locally free $\mathcal{O}_X$ -module or that $(X, \mathcal{O}_X)$ and $g_*(\mathcal{O}_{X'})$ satisfy (II bis) and $g_*(\mathcal{O}_{X'}) \rightarrow g_*(\mathcal{O}_{X'}) \otimes_{\mathcal{O}_X} \mathcal{R}(X)$ is injective. Let $\mathcal{L}'$ be an invertible $\mathcal{O}_{X'}$ -module, f' a section of $\mathcal{L}'$ over X' , and $f = N_{X'/X}(f')$ the corresponding section of $\mathcal{L} = N_{X'/X}(\mathcal{L}')$ over X (6.5.5.3). [II:130] Then $g(X' \setminus X'_{f'}) = X \setminus X_f$, and X_f is the largest open subset U of X such that $g^{-1}(U) \subset X'_{f'}$.*

Proof. Indeed, $g(X' \setminus X'_{f'})$ is closed in X (6.1.10); it thus suffices to prove the last claim. But the relation $U \subset X_f$ is equivalent to the fact that the homomorphism $\mathcal{O}_X|_U \rightarrow \mathcal{L}|_U$ defined by $f|_U$ is an isomorphism. By (6.5.6), this is equivalent to saying that the homomorphism $\mathcal{O}_{X'}|_{g^{-1}(U)} \rightarrow \mathcal{L}'|_{g^{-1}(U)}$ defined by $f'|_{g^{-1}(U)}$ is an isomorphism, which is equivalent to the relation $g^{-1}(U) \subset X'_{f'}$. $\square$

Proposition (6.5.8). — *Let $g : X' \rightarrow X$ be a finite morphism, and $f : Y \rightarrow X$ a morphism; let $Y' = X'_{(Y)}$, $g' = g_{(Y)}$, and $f' = f_{(X')}$, so that the diagram*

$$\begin{array}{ccc} X' & \xleftarrow{f'} & Y' \\ g \downarrow & & \downarrow g' \\ X & \xleftarrow{f} & Y \end{array}$$

commutes.

Suppose that either $g_(\mathcal{O}_{X'})$ is a locally free $\mathcal{O}_X$ -module or that $(X, \mathcal{O}_X)$ and $g_*(\mathcal{O}_{X'})$ satisfy (II). Suppose further that Y is a locally Noetherian reduced prescheme, and that the restriction of f to any irreducible component of Y is a dominant morphism to an irreducible component of X . Then, for every invertible $\mathcal{O}_{X'}$ -module $\mathcal{L}'$, we have*

$$N_{Y'/Y}(f'^*(\mathcal{L}')) = f^*(N_{X'/X}(\mathcal{L}'))$$

up to canonical isomorphism.

Proof. Note that we have $f^*(g_*(\mathcal{L}')) = g'_*(f'^*(\mathcal{L}'))$, by (1.5.2), and, in particular, $g'_*(\mathcal{O}_{Y'}) = f^*(g_*(\mathcal{O}_{X'}))$; if $g_*(\mathcal{O}_{X'})$ is locally free, then so too is $g'_*(\mathcal{O}_{Y'})$. The conclusion then follows from the definitions and (6.5.4). $\square$

Remark (6.5.9). — We later generalise the notion of norm developed above, by placing it in relation to the notion of direct image of a divisor.

6.6. Application: criteria for ampleness

Proposition (6.6.1). — Let Y be a prescheme, $f : X \rightarrow Y$ a quasi-compact morphism, and $g : X' \rightarrow X$ a finite and surjective morphism. Suppose that either $g_*(\mathcal{O}_{X'})$ is a locally free $\mathcal{O}_X$ -module or that $(X, \mathcal{O}_X)$ and $g_*(\mathcal{O}_{X'})$ satisfy (II bis). Then, for every invertible $\mathcal{O}_{X'}$ -module $\mathcal{L}'$ that is ample for $f \circ g$, $\mathcal{L} = N_{X'/X}(\mathcal{L}')$ is ample for f .

Proof. We can suppose Y to be affine (4.6.4), and then, by (4.6.6), the statement is equivalent to: $\square$

Corollary (6.6.2). — Let X be a quasi-compact prescheme, $g : X' \rightarrow X$ a finite surjective morphism, such that either $g_*(\mathcal{O}_{X'})$ is a locally free $\mathcal{O}_X$ -module or $(X, \mathcal{O}_X)$ and $g_*(\mathcal{O}_{X'})$ satisfy (II bis). Then, for every ample $\mathcal{O}_{X'}$ -module $\mathcal{L}'$, $\mathcal{L} = N_{X'/X}(\mathcal{L}')$ is ample.

Proof. In the second hypothesis, we can further suppose that the canonical homomorphism $g_*(\mathcal{O}_{X'}) \rightarrow g_*(\mathcal{O}_{X'}) \otimes_{\mathcal{O}_X} \mathcal{R}(X)$ is injective. ^[II:131] Indeed, if not, let $\mathcal{T}$ be the kernel of this homomorphism, which is a coherent ideal of $\mathcal{B} = g_*(\mathcal{O}_{X'})$ (I, 6.1.1), and set $X'' = \text{Spec}(\mathcal{B}/\mathcal{T})$; we thus have a commutative diagram

$$\begin{array}{ccc} X'' & \xrightarrow{h} & X' \\ & \searrow g' & \swarrow g \\ & X & \end{array}$$

where h is a closed immersion (1.4.10). Furthermore, we know that the support of $\mathcal{T}$ is a closed set (0, 5.2.2) that is nowhere dense in X (I, 7.4.6), whence we conclude that, for the generic point x of an irreducible component of X , there is an affine open neighbourhood U of x such that $\mathcal{B}|_U = (\mathcal{B}/\mathcal{T})|_U$. Since g is, by hypothesis, surjective, we thus conclude that $x \in g'(X'')$; g' is thus dominant, and, since it is a finite morphism, it is surjective (6.1.10); by definition,

$$g'_*(\mathcal{O}_{X''}) \otimes_{\mathcal{O}_X} \mathcal{R}(X) = (\mathcal{B}/\mathcal{T}) \otimes_{\mathcal{O}_X} \mathcal{R}(X) = g_*(\mathcal{O}_{X'}) \otimes_{\mathcal{O}_X} \mathcal{R}(X)$$

thus $(X, \mathcal{O}_X)$ and $g'_*(\mathcal{O}_{X''})$ satisfy (II bis), and furthermore $g'_*(\mathcal{O}_{X''}) \rightarrow g'_*(\mathcal{O}_{X''}) \otimes_{\mathcal{O}_X} \mathcal{R}(X)$ is injective. Finally, $h^*(\mathcal{L}') = \mathcal{L}''$ is an ample $\mathcal{O}_{X''}$ -module (4.6.13, (i bis)), and we have that $N_{X''/X}(\mathcal{L}'') = N_{X'/X}(\mathcal{L}')$. Indeed, to define these two invertible $\mathcal{O}_X$ -modules, we can use the same affine open cover (U_λ) of X such that the restrictions of $g_*(\mathcal{L}')$ and $g'_*(\mathcal{L}'')$ to U_λ are isomorphic to $\mathcal{B}|_{U_\lambda}$ and $(\mathcal{B}/\mathcal{T})|_{U_\lambda}$ (respectively). We immediately see that, for every isomorphism $\eta_\lambda : g_*(\mathcal{L}')|_{U_\lambda} \rightarrow \mathcal{B}|_{U_\lambda}$, there is a canonically corresponding isomorphism

$$\eta'_\lambda : g'_*(\mathcal{L}'')|_{U_\lambda} \rightarrow (\mathcal{B}/\mathcal{T})|_{U_\lambda}$$

so that, if $(\omega_{\lambda\mu})$ and $(\omega'_{\lambda\mu})$ are the 1-cocycles corresponding to the systems of isomorphisms (η_λ) and (η'_λ) (6.5.2), $\omega'_{\lambda\mu}$ is the canonical image in $\Gamma(U_\lambda \cap U_\mu, \mathcal{B}/\mathcal{T})$ of $\omega_{\lambda\mu} \in \Gamma(U_\lambda \cap U_\mu, \mathcal{B})$. By the definition of $\mathcal{T}$, we thus conclude that

$$N_{\mathcal{B}/\mathcal{A}}(\omega_{\lambda\mu}) = N_{(\mathcal{B}/\mathcal{T})/\mathcal{A}}(\omega'_{\lambda\mu})$$

(where $\mathcal{A} = \mathcal{O}_X$), whence the claimed equality.

So suppose that the homomorphism $g_*(\mathcal{O}_{X'}) \rightarrow g_*(\mathcal{O}_{X'}) \otimes_{\mathcal{O}_X} \mathcal{R}(X)$ is injective when we are in hypothesis (II bis). It suffices to prove that, as f runs over the sections of $\mathcal{L}^{\otimes n}$ (for $n > 0$) over X , the X_f form a base for the topology of X (4.5.2). But let $x \in X$, and let U be an arbitrary neighbourhood of x ; since $g^{-1}(x)$ is finite (6.1.7) and $\mathcal{L}'$ is ample, there exists an integer $n > 0$ and a section f' of $\mathcal{L}'^{\otimes n}$ over X' such that $X'_{f'}$ is a neighbourhood of $g^{-1}(x)$ contained inside $g^{-1}(U)$ (4.5.4). Since

$$\mathcal{L}^{\otimes n} = N_{X'/X}(\mathcal{L}'^{\otimes n})$$

it suffices to take $f = N_{X'/X}(f')$; indeed, then $X \setminus X_f = g(X' \setminus X'_{f'})$ (6.5.7), and so $x \in X_f \subset U$. $\square$

Corollary (6.6.3). — Under the hypotheses of (6.6.1), for an invertible $\mathcal{O}_X$ -module $\mathcal{L}$ to be ample for f , it is necessary and sufficient that $\mathcal{L}' = g^*(\mathcal{L})$ be ample for $f \circ g$.

Proof. The condition is necessary, since g is affine (5.1.12). To prove that the condition is sufficient, we can suppose that Y is affine (4.6.4), and so X and X' are quasi-compact and $\mathcal{L}'$ is ample (4.6.6), and we need to show that $\mathcal{L}$ is ample. [II.132] But the set of points $x \in X$ that admit a neighbourhood where $g_*(\mathcal{O}_{X'})$ (resp. $g_*(\mathcal{O}_{X'}) \otimes_{\mathcal{O}_X} \mathcal{R}(X)$) has a given rank n in the first (resp. second) hypothesis is simultaneously open and closed in X , and so X is the prescheme given by the sum of a finite number of these opens, and we can thus suppose that it is equal to one of them (4.6.17). But then $N_{X'/X}(\mathcal{L}') = \mathcal{L}^{\otimes n}$, and so $\mathcal{L}^{\otimes n}$ is ample by (6.6.2), and thus so too is $\mathcal{L}$ (4.5.6). $\square$

Corollary (6.6.4). — Suppose that the hypotheses of (6.6.1) are satisfied, and suppose further that $f : X \rightarrow Y$ is of finite type. Then, for f to be quasi-projective, it is necessary and sufficient that $f \circ g$ be quasi-projective. If we further suppose that Y is a quasi-compact scheme, or a prescheme whose underlying space is Noetherian, then for f to be projective, it is necessary and sufficient that $f \circ g$ be projective.

Proof. The hypothesis implies that $f \circ g$ is of finite type. Taking into account the definition of quasi-projective morphisms (5.3.1), the first claim follows from (6.6.1) and (6.6.3). Taking into account this result, along with (5.5.3, (ii)), it remains to show that, if f is quasi-projective, then for f to be proper, it is necessary and sufficient that $f \circ g$ be proper. But f is then separated (5.3.1) and of finite type; since g is surjective, our claim follows from (5.4.2, (ii)) and (5.4.3, (ii)). $\square$

In particular:

Corollary (6.6.5). — Let X be a prescheme of finite type over a field K , and K' a finite-degree extension of K . For X to be projective (resp. quasi-projective) over K , it is necessary and sufficient that $X' = X \otimes_K K'$ be projective (resp. quasi-projective) over K' .

Proof. The condition is necessary ((5.3.4, (iii)) and (5.5.5, (iii))). Conversely, suppose that it is satisfied, and let $g : X' \rightarrow X$ be the canonical projection. It is clear that g is a finite morphism (6.1.5, (iii)) and surjective (I, 3.5.2, (ii)). Furthermore, $g_*(\mathcal{O}_{X'})$ is a locally free $\mathcal{O}_X$ -module, since it is isomorphic to $\mathcal{O}_X \otimes_K K'$ (1.5.2). It then follows from the hypothesis and from (6.1.11) and (5.5.5, (ii)) that X' is projective (resp. quasi-projective) over K ; we then deduce from (6.6.4) that X is projective (resp. quasi-projective) over K . $\square$

In Chapter V, we will show that the statement of (6.6.5) remains true when K' is an arbitrary extension of K .

The rest of this section is dedicated to the proof of the criterion in (6.6.11), which is a rather technical refinement of (6.6.1); it can be omitted on a first reading.

Lemma (6.6.6). — Let X be a reduced Noetherian prescheme, and $\mathcal{E}$ a coherent $\mathcal{O}_X$ -module such that $\mathcal{E} \otimes_{\mathcal{O}_X} \mathcal{R}(X)$ is a locally free $\mathcal{R}(X)$ -module of constant rank n . Then there exists a reduced Noetherian prescheme Z and a birational finite morphism $h : Z \rightarrow X$ that has the following property: the morphisms of sheaves of sets $\sigma_i : \mathcal{H}om_{\mathcal{O}_X}(\mathcal{E}, \mathcal{E}) \rightarrow \mathcal{R}(X)$ (for $1 \leq i \leq n$) (cf. (6.4.10)) send $\mathcal{H}om_{\mathcal{O}_X}(\mathcal{E}, \mathcal{E})$ to the coherent $\mathcal{O}_X$ -algebra $h_*(\mathcal{O}_Z)$.

Proof. Consider an affine open U of X , of ring $A(U) = A$; let $E = \Gamma(U, \mathcal{E})$, and let C_U be the subalgebra of $R(U)$ generated by the $\sigma_i(u)$ where u runs over $\text{Hom}_A(E, E)$; we have seen (6.4.7.1) that this A -algebra is an A -module of finite type. Furthermore, it is clear that forming the algebras C_U commutes with the restriction operations of an affine open U to an affine open $U' \subset U$. We have thus defined a finite sub- $\mathcal{O}_X$ -algebra $\mathcal{C}$ of $\mathcal{R}(X)$ such that $\Gamma(U, \mathcal{C}) = C_U$ for every affine open U of X . We take $Z = \text{Spec}(\mathcal{C})$, and take h to be the structure morphism, which is thus finite (6.1.2); since $\mathcal{C}$ is reduced, Z is a reduced Noetherian prescheme (1.3.8). Finally, the total ring of fractions of C_U is $R(U)$, by definition, and since C_U is contained inside the integral closure of $A(U)$ in $R(U)$, there is a bijective correspondence between minimal prime ideals of $A(U)$ and minimal prime ideals of C_U [13, t. I, p. 259], which proves that h is birational and finishes the proof. $\square$

Corollary (6.6.7). — Under the hypotheses of (6.6.6), let W be an open of X such that, for all $x \in W$, either X is normal at the point x , or $\mathcal{E}_x$ is a free $\mathcal{O}_x$ -module. Then we can suppose h to be defined such that the restriction of h to $h^{-1}(W)$ is an isomorphism from $h^{-1}(W)$ to W .

Proof. Either hypothesis implies that, if $U \subset W$ is an affine open, then, with the notation of (6.6.6), $(\sigma_i(u))_x \in A_x$ for all $x \in U$ (6.4.3), and so $\sigma_i(u) \in A$, and the conclusion follows from the definition of h given in (6.6.6). $\square$

(6.6.8). Let X be a reduced Noetherian prescheme, and $g : X' \rightarrow X$ a finite surjective morphism, so that $\mathcal{B} = g_*(\mathcal{O}_{X'})$ is a coherent $\mathcal{O}_X$ -algebra; suppose further that $\mathcal{B} \otimes_{\mathcal{O}_X} \mathcal{R}(X)$ is a locally free $\mathcal{R}(X)$ -module of constant rank n . We can then apply Lemma (6.6.6), taking $\mathcal{E} = \mathcal{B}$, whence, with the notation of (6.6.6), we obtain a homomorphism of sheaves of multiplicative monoids $\sigma_n : \mathcal{H}om_{\mathcal{O}_X}(\mathcal{B}, \mathcal{B}) \rightarrow h_*(\mathcal{O}_Z)$, and by composing this homomorphism with the canonical homomorphism $\mathcal{B} \rightarrow \mathcal{H}om_{\mathcal{O}_X}(\mathcal{B}, \mathcal{B})$ (6.5.1), we thus obtain a homomorphism of sheaves of multiplicative monoids:

$$(6.6.8.1) \quad N' : \mathcal{B} = g_*(\mathcal{O}_{X'}) \longrightarrow h_*(\mathcal{O}_Z) = \mathcal{C}.$$

With this in mind, for every invertible $\mathcal{O}_{X'}$ -module $\mathcal{L}'$, $g_*(\mathcal{L}')$ is an invertible $\mathcal{B}$ -module (6.1.12), and the method of (6.5.2) allows us to functorially associate to $\mathcal{L}'$ an invertible $\mathcal{C}$ -module, which we denote by $N'(g_*(\mathcal{L}'))$.

Lemma (6.6.9). — *Let X be a reduced Noetherian prescheme, and $g : X' \rightarrow X$ a finite surjective morphism such that $g_*(\mathcal{O}_{X'}) \otimes_{\mathcal{O}_X} \mathcal{R}(X)$ is a locally free $\mathcal{R}(X)$ -module of constant rank. Then there exists a reduced Noetherian prescheme Z and a finite birational morphism $h : Z \rightarrow X$ that has the following property: for every ample $\mathcal{O}_{X'}$ -module $\mathcal{L}'$, the invertible $\mathcal{O}_Z$ -module $\mathcal{M}$ such that $h_*(\mathcal{M}) = N'(g_*(\mathcal{L}'))$ (using the notation of (6.6.8)) is ample.*

Proof. Suppose first of all that the homomorphism $\mathcal{B} \rightarrow \mathcal{B} \otimes_{\mathcal{O}_X} \mathcal{R}(X)$ is injective. Define Z and h as in (6.6.6) (with $\mathcal{E} = g_*(\mathcal{O}_{X'})$). Let $z \in Z$; we have to show that there exists an integer $m > 0$ and a section t of $\mathcal{M}^{\otimes m}$ over Z such that Z_t is an affine open that contains z (4.5.2). Let $x = h(z)$, and let U be an affine open of X that contains x ; then $h^{-1}(U)$ is an affine open neighbourhood of z , and it suffices to find t such that $z \in Z_t \subset h^{-1}(U)$, since Z_t will then necessarily be affine (5.5.8). There exists, by hypothesis, an integer $n > 0$ and a section s' of $\mathcal{L}'^{\otimes n}$ over X' such that

$$(6.6.9.1) \quad g^{-1}(x) \subset X'_{s'} \subset g^{-1}(U)$$

by (4.5.4). By definition, s' is also a section of $g_*(\mathcal{L}'^{\otimes n})$ over X , and it corresponds, as in (6.5.2), to a section

$$s = N'(s') \quad \text{of} \quad N'(g_*(\mathcal{L}'^{\otimes n})) = N'(g_*(\mathcal{L}'))^{\otimes n}$$

over X . [II-134] We will show that, if t is the section s considered as a section of $\mathcal{M}^{\otimes n}$ over Z , then t is the desired section. Set

$$(6.6.9.2) \quad V = X \setminus g(X' \setminus X'_{s'})$$

which is an open of X that contains x and is contained in U , by (6.6.9.1) and (6.1.10). We will show that

$$(6.6.9.3) \quad h^{-1}(V) \subset Z_t \subset h^{-1}(U)$$

which will finish the proof. It is equivalent to say that the set T of $y \in X$ such that s_y is invertible contains V and is contained in U . For this, consider first of all an affine open W contained in V ; then $g^{-1}(W)$ is an affine open in X' , and by (6.6.9.2) $s'_{y'}$ is invertible for all $y' \in g^{-1}(W)$; by the hypotheses on X and $\mathcal{B}$, we can apply the results of (6.4.7), and we see that, if $y = g(y')$, then s_y is invertible; in other words, $V \subset T$. On the other hand, it also follows from (6.4.7) that, conversely, if s_y is invertible, then so too is $s'_{y'}$, which implies that $y' \in g^{-1}(U)$ by (6.6.9.1), and so $y \in U$, whence $T \subset U$ in this case.

We pass from this to the general case by the same argument as in (6.6.2), replacing X' by X'' such that $g'_*(\mathcal{O}_{X''}) \rightarrow g'_*(\mathcal{O}_{X''}) \otimes_{\mathcal{O}_X} \mathcal{R}(X)$ is injective, and $\mathcal{L}'$ by an ample $\mathcal{O}_{X''}$ -module $\mathcal{L}''$ such that $N'(g_*(\mathcal{L}')) = N'(g'_*(\mathcal{L}''))$. Lemma (6.6.9) is then proven in all cases (with a suitable choice of h). $\square$

Corollary (6.6.10). — *Suppose that the hypotheses of (6.6.9) are satisfied; for every invertible $\mathcal{O}_X$ -module $\mathcal{L}$ such that $g^*(\mathcal{L})$ is ample, $h^*(\mathcal{L})$ is ample.*

Proof. If we set $\mathcal{L}' = g^*(\mathcal{L})$, then $g_*(\mathcal{L}') = \mathcal{L} \otimes_{\mathcal{O}_X} \mathcal{B}$ (0, 5.4.10), so

$$N'(g_*(\mathcal{L}')) = (\mathcal{L} \otimes_{\mathcal{O}_X} \mathcal{C})^{\otimes n}$$

(by the same argument as for (6.5.2.4)). We thus conclude that $\mathcal{M} = (h^*(\mathcal{L}))^{\otimes n}$, and since $\mathcal{M}$ is ample, so too is $h^*(\mathcal{L})$ (4.5.6). $\square$

Proposition (6.6.11). — *Let Y be an affine scheme, X a reduced Noetherian prescheme, $f : X \rightarrow Y$ a quasi-compact morphism, and $g : X' \rightarrow X$ a finite surjective morphism. Let W be an open subset of X such that, for all $x \in W$, either X is normal at the point x or there exists an open neighbourhood $T \subset W$ of x such that $(g_*(\mathcal{O}_{X'}))|_T$ is a locally free $(\mathcal{O}_X|_T)$ -module. Then there exists a reduced Y -prescheme Z and a finite birational Y -morphism $h : Z \rightarrow X$ such that the restriction of h to $h^{-1}(W)$ is an isomorphism from $h^{-1}(W)$ to W and has the following property: for every invertible $\mathcal{O}_X$ -module $\mathcal{L}$ such that $g^*(\mathcal{L})$ is ample for $f \circ g$, $h^*(\mathcal{L})$ is ample for $f \circ h$.*

Proof. Since Y is affine, $g^*(\mathcal{L})$ is ample, and the problem thus reduces to proving that, for a suitable choice of h , $h^*(\mathcal{L})$ is ample (4.6.6). We will show that we can replace g by a finite surjective morphism $g' : X'' \rightarrow X$ such that $g'^*(\mathcal{L})$ is ample and $g'_*(\mathcal{O}_{X''}) \otimes_{\mathcal{O}_X} \mathcal{R}(X)$ is a locally free $\mathcal{R}(X)$ -module of constant rank; we will then have reduced to the conditions of (6.6.10), and the proposition will thus be proven.

For this, let $\mathcal{B} = g_*(\mathcal{O}_{X'})$; denote by X_i (for $1 \leq i \leq n$) the closed reduced subpreschemes of X that have the irreducible components of X as their underlying space (I, 5.2.1); they are integral by hypothesis. ^[II-135] Let X'_i be the closed subprescheme $g^{-1}(X_i)$ of X' , and $g_i : X'_i \rightarrow X_i$ the morphism g restricted to X'_i , which is finite (6.1.5, (iii)) and surjective; let k_i be the rank of the $\mathcal{O}_{X_i}$ -algebra $\mathcal{B}_i = (g_i)_*(\mathcal{O}_{X'_i})$. Since $\mathcal{B}_i \otimes_{\mathcal{O}_{X_i}} \mathcal{R}(X_i)$ is a constant presheaf (I, 7.3.5), the rank k_i is also the rank of the $(\mathcal{O}_X|_U)$ -algebra $\mathcal{B}|_U$ for every open U of X that meets *only* the irreducible component X_i . If T is an open subset of X such that $\mathcal{B}|_T$ is isomorphic to $\mathcal{O}_X^m|_T$, then it follows from the above remark that the numbers k_i are equal to m for all the indices i such that $T \cap X_i \neq \emptyset$. So let U be the open subset of X given by the points that admit a neighbourhood on which $\mathcal{B}$ is a locally free $\mathcal{O}_X$ -module, and let U_j (for $1 \leq j \leq s$) be its connected components, which are open in X and finitely many (since U is Noetherian); denote by V_j the closed subprescheme of X' given by the closure of the subprescheme induced on the open $g^{-1}(U_j)$ (I, 9.5.11). By the above, for all indices i such that $X_i \cap U_j \neq \emptyset$, the ranks k_i are all equal to a single integer m_j ; note also that a single X_i cannot meet two U_j with different indices j . Let i_λ be the indices i such that $X_i \cap U = \emptyset$. Consider the product k of all the k_i , set $n_i = k/k_i$, and let X'' be the prescheme defined as follows. For each j (for $1 \leq j \leq s$), consider k/m_j preschemes isomorphic to V_j , and for each λ , k/k_{i_λ} preschemes isomorphic to X'_{i_λ} ; then X'' is the sum of all these preschemes. We define a morphism $g'' : X'' \rightarrow X'$ that reduces to the canonical injection on each of the summands of X'' ; it is clear that g'' is a finite dominant morphism, and thus surjective (since a finite morphism is closed (6.1.10)); set $g' = g \circ g''$, which is a finite surjective morphism $X'' \rightarrow X$; we have $g'^*(\mathcal{L}) = g''^*(g^*(\mathcal{L}))$, and so $g'^*(\mathcal{L})$ is an ample $\mathcal{O}_{X''}$ -module (5.1.12). It is then clear that, for this new prescheme X'' , the ranks defined as the k_i for X' are all equal to k ; taking (I, 7.3.3) into account, we immediately conclude that, for every affine open T of X , $(g'_*(\mathcal{O}_{X''}) \otimes_{\mathcal{O}_X} \mathcal{R}(X))|_T$ is an $(\mathcal{R}(X)|_T)$ -module isomorphic to $(\mathcal{R}(X)|_T)^k$. $\square$

Corollary (6.6.12). — *If, in the statement of (6.6.11), we have $W = X$, then for an invertible $\mathcal{O}_X$ -module $\mathcal{L}$ to be ample for f , it is necessary and sufficient that $g^*(\mathcal{L})$ be ample for $f \circ g$.*

Remark (6.6.13). — In Chapter III, we will see that, if Y is Noetherian, if f is of finite type, and if the restriction of f to the closed reduced subprescheme of X that has $X \setminus W$ as its underlying space is *proper*, then the conclusion of (6.6.12) still holds true. But we will give, in Chapter V, examples of algebraic schemes X over a field K (with the structure morphism $X \rightarrow \text{Spec}(K)$ not proper) whose normalisation X' is quasi-affine, but which are not quasi-affine (in that $\mathcal{O}_X$ is not ample, even though $\mathcal{O}_{X'}$ is (5.1.12) and the morphism $X' \rightarrow X$ is finite and surjective (6.3.10)). We will see in the next section that this circumstance cannot happen when we replace “quasi-affine” by “affine”.

6.7. Chevalley's theorem We are going to prove (with the help of Serre's criterion (5.2.1)) the following theorem, which was proven by C. Chevalley by other methods, in the case of *algebraic schemes*.

[II:136]

Theorem (6.7.1). — *Let X be an affine scheme, Y a Noetherian prescheme, and $f : X \rightarrow Y$ a finite surjective morphism. Then Y is an affine scheme.*

Proof. It is clear that $f_{\text{red}} : X_{\text{red}} \rightarrow Y_{\text{red}}$ is finite (6.1.5, (vi)); since X_{red} is an affine scheme, and since saying that Y is affine is equivalent to saying that Y_{red} is affine (since Y is Noetherian (I, 6.1.7)), we see that we can suppose Y to be *reduced*. For every closed subset Y' of Y , there exists exactly one reduced subprescheme of Y that has Y' as its underlying space (I, 5.1.2); its inverse image $f^{-1}(Y')$, which is canonically isomorphic to $X \times_Y Y'$ (I, 4.4.1), is affine as a closed subprescheme of X , and the restriction of f to $f^{-1}(Y')$, which can be identified with $f \times_Y 1_{Y'}$, is a finite surjective morphism (6.1.5, (iii)). By the principle of Noetherian induction (0, 2.2.2), we can thus (taking (I, 6.1.7) into account) reduce to proving the theorem under the hypothesis that for *every* closed subset $Y' \neq Y$, every closed subprescheme of Y that has Y' as its underlying space is affine. We thus conclude that, for every coherent $\mathcal{O}_Y$ -module $\mathcal{F}$ whose (closed) support Z is distinct from Y , we have $H^1(Y, \mathcal{F}) = 0$. Indeed, there exists a closed subprescheme Y' of Y that has Z as its underlying space and is such that, if $j : Y' \rightarrow Y$ is the canonical injection, then $\mathcal{F} = j_*(j^*(\mathcal{F}))$ (I, 9.3.5); then (5.2.3) $H^1(Y, \mathcal{F}) = H^1(Y', j^*(\mathcal{F})) = 0$, by (I, 5.1.9.2).

Suppose first of all that Y is not irreducible, and let Y' be an irreducible component of Y , and $Y'' = Y \setminus Y'$; we again denote by Y' the closed reduced subprescheme of Y that has Y' as its underlying space, and by j the canonical injection $Y' \rightarrow Y$. Let $\mathcal{F}$ be a coherent $\mathcal{O}_Y$ -module, and consider the canonical homomorphism

$$\rho : \mathcal{F} \longrightarrow \mathcal{F}' = j_*(j^*(\mathcal{F}))$$

(0, 4.4.3); $\mathcal{F}'$ is a coherent $\mathcal{O}_Y$ -module by (0, 5.3.10) and (0, 5.3.12), since $j_*(\mathcal{O}_{Y'}) = \mathcal{O}_Y / \mathcal{J}$, where we denote by $\mathcal{J}$ the sheaf of ideals of $\mathcal{O}_Y$ that defines the subprescheme Y' . Then $\mathcal{G} = \text{Ker } \rho$ and $\mathcal{K} = \text{Im } \rho$ are also coherent $\mathcal{O}_Y$ -modules (0, 5.3.4); but, by definition, the fibre $\mathcal{F}'_y$ of $\mathcal{F}'$ at the generic point y of Y' is equal to $\mathcal{F}_y$, since y is interior to Y' and thus $\mathcal{J}_y = 0$, since Y is reduced. We thus conclude that y is not contained in the (closed) support of $\mathcal{G}$; also, the support of $\mathcal{F}'$ (and *a fortiori* that of $\mathcal{K}$) is contained in Y' ; in other words, the supports of $\mathcal{G}$ and $\mathcal{K}$ are *distinct* from Y . We thus deduce that $H^1(Y, \mathcal{G}) = H^1(Y, \mathcal{K}) = 0$, and the exact sequence of cohomology applied to the exact sequence $0 \rightarrow \mathcal{G} \rightarrow \mathcal{F} \rightarrow \mathcal{K} \rightarrow 0$ implies that $H^1(Y, \mathcal{F}) = 0$. We thus conclude by Serre's criterion (5.2.1).

So suppose that Y is irreducible, and thus *integral*. We can also suppose that X is *integral*: if we denote by X_i the closed reduced subpreschemes of X that have the irreducible components of X as their underlying space (I, 5.2.1), and by f_i the restriction of f to X_i , then at least one of the f_i is dominant, and since it is a finite morphism (6.1.5), it is surjective (6.1.10); since X_i is also an affine scheme, we see that we can replace X by X_i in the statement.

Lemma (6.7.1.1). — *Let X and Y be integral Noetherian preschemes, x (resp. y) the generic point of X (resp. Y), and $f : X \rightarrow Y$ a finite surjective morphism. Let $\mathcal{L}$ be an invertible $\mathcal{O}_X$ -module such that there exists an affine open neighbourhood U of y and a section $g \in \Gamma(X, \mathcal{L})$ such that $x \in X_g \subset f^{-1}(U)$. [II:137] Then there exist integers $m, n > 0$, a homomorphism $u : \mathcal{O}_Y^m \rightarrow f_*(\mathcal{L}^{\otimes n})$, and an open neighbourhood V of y such that the restriction $u|_V$ is an isomorphism from $\mathcal{O}_Y^m|_V$ to $f_*(\mathcal{L}^{\otimes n})|_V$.*

Proof. Let C be the (integral) ring of U , and $k = \mathcal{O}_y$ its field of fractions: since f is finite, $U' = f^{-1}(U)$ is affine (1.3.2); let D be its (integral) ring with field of fractions $K = \mathcal{O}_x$; by hypothesis, D is a C -module of finite type (6.1.4), and so K is an extension of finite rank of k . The fibre $f^{-1}(y) = X \times_Y \text{Spec}(k(y)) = X \times_Y \text{Spec}(\mathcal{O}_y)$ can be identified with $\text{Spec}(K)$ (I, 3.6.6); let s_i (for $1 \leq i \leq m$) be elements of D that form a basis of K over k . There exists $n > 0$ such that the sections $(s_i|_{X_g})g^{\otimes n}$ of $\mathcal{L}^{\otimes n}$ over X_g extend to sections b_i (for $1 \leq i \leq m$) of $\mathcal{L}^{\otimes n}$ over X (I, 9.3.1). The b_i are also, by definition, sections of $f_*(\mathcal{L}^{\otimes n})$ over Y , and thus define a homomorphism $u : \mathcal{O}_Y^m \rightarrow f_*(\mathcal{L}^{\otimes n})$ (0, 5.1.1); we will show that u is the desired homomorphism. We have that $\mathcal{L}^{\otimes n}|_{U'} = \tilde{M}$, where M is a D -module of finite type, so if ϕ is the injection $C \rightarrow D$ corresponding to the morphism

$f^{-1}(U) \rightarrow U$ given by the restriction of f , then $M_{[\phi]}$ is a C -module of finite type; then

$$f_*(\mathcal{L}^{\otimes n})|U = (M_{[\phi]})^\sim$$

(I, 1.6.3) is coherent, and since U is an arbitrary affine open of Y , $f_*(\mathcal{L}^{\otimes n})$ is coherent; furthermore, $u|U = \tilde{\theta}$, where θ is a C -homomorphism $C^m \rightarrow M_{[\phi]}$, and $u_y = \theta_y$ is the homomorphism $\theta \otimes 1 : k^m = C^m \otimes_C k \rightarrow M_{[\phi]} \otimes_C k$; but the latter is, by definition, an *isomorphism*, since the $(b_i)_x$ form a basis of $M_{[\phi]} \otimes k$ over k , with g_x being, by hypothesis, a generator of $(\mathcal{L}^{\otimes n})_x$. We thus conclude that the supports of the $\mathcal{O}_Y$ -modules $\text{Ker } u$ and $\text{Coker } u$ do not contain y ; since these $\mathcal{O}_Y$ -modules are coherent (0, 5.3.3), their supports are closed (0, 5.2.2), whence the lemma. $\square$

With this, the hypotheses of Lemma (6.7.1.1) are satisfied in the case that we are considering, by taking $\mathcal{L} = \mathcal{O}_X$, since X is affine (I, 1.1.10); set $\mathcal{A} = \mathcal{O}_Y$ and $\mathcal{B} = f_*(\mathcal{O}_X)$. By Serre's criterion (5.2.1.1), it suffices to prove that, for every coherent $\mathcal{O}_Y$ -module $\mathcal{F}$, we have that $H^1(Y, \mathcal{F}) = 0$; it even suffices to prove this in the case where $\mathcal{F} \subset \mathcal{O}_Y$, which implies that $\mathcal{F}$ is torsion free, since Y is integral; in fact, we will show that $H^1(Y, \mathcal{F}) = 0$ for every coherent *torsion-free* $\mathcal{O}_Y$ -module $\mathcal{F}$. But the homomorphism $u : \mathcal{A}^m \rightarrow \mathcal{B}$ defines a homomorphism

$$v : \mathcal{G} = \mathcal{H}om_{\mathcal{A}}(\mathcal{B}, \mathcal{F}) \longrightarrow \mathcal{H}om_{\mathcal{A}}(\mathcal{A}^m, \mathcal{F}) = \mathcal{F}^m.$$

We will first show that v is *injective*: by hypothesis, $\mathcal{T} = \text{Coker } u$ has a support that does not meet V , and is thus a *torsion* $\mathcal{O}_Y$ -module (I, 7.4.6); the exact sequence

$$\mathcal{A}^m \longrightarrow \mathcal{B} \longrightarrow \mathcal{T} \longrightarrow 0$$

gives, by left exactness of the functor $\mathcal{H}om_{\mathcal{A}}$, the exact sequence

$$0 \longrightarrow \mathcal{H}om_{\mathcal{A}}(\mathcal{T}, \mathcal{F}) \longrightarrow \mathcal{G} \xrightarrow{v} \mathcal{F}^m.$$

But since $\mathcal{F}$ is torsion free, we have that $\mathcal{H}om_{\mathcal{A}}(\mathcal{T}, \mathcal{F}) = 0$ (0, 5.2.6), whence our claim. We thus have the exact sequence

$$0 \longrightarrow \mathcal{G} \longrightarrow \mathcal{F}^m \longrightarrow \text{Coker } v \longrightarrow 0$$

[II:138] where $\mathcal{G}$ and $\text{Coker } v$ are coherent $\mathcal{O}_Y$ -modules ((0, 5.3.4) and (0, 5.3.5)); by the exact sequence of cohomology, it would suffice to show that $H^1(Y, \mathcal{G}) = H^1(Y, \text{Coker } v) = 0$ in order to deduce that $H^1(Y, \mathcal{F}^m) = (H^1(Y, \mathcal{F}))^m = 0$, and thus that $H^1(Y, \mathcal{F}) = 0$. But the restriction $v|V$ is an isomorphism, and so the support of $\text{Coker } v$ is distinct from Y , whence $H^1(Y, \text{Coker } v) = 0$ by the hypothesis at the start. Now, $\mathcal{G}$ is a coherent $\mathcal{B}$ -module (I, 9.6.4); since X is affine over Y , there exists a quasi-coherent $\mathcal{O}_X$ -module $\mathcal{K}$ such that $\mathcal{G}$ is isomorphic to $f_*(\mathcal{K})$ (1.4.3); since X is affine, we have that $H^1(X, \mathcal{K}) = 0$ (I, 5.1.9.2), and so $H^1(Y, \mathcal{G}) = 0$ by (5.2.3), which finishes the proof of Theorem (6.7.1). $\square$

Corollary (6.7.2). — *Let X be a Noetherian prescheme, $(X_i)_{1 \leq i \leq n}$ a finite cover of the space X consisting of closed subsets. For X to be affine, it is necessary and sufficient that, for each i , there exist a closed subprescheme of X that has X_i as its underlying space and is affine.*

Proof. Indeed, if this is the case, then let X' be the scheme given by the *sum* of the X_i ; it is clear that X' is affine, and we define a surjective morphism $f : X' \rightarrow X$ by taking the restriction of f to X_i to be the canonical injection. Everything reduces to showing that f is *finite*, by (6.7.1), and this we have already seen in (6.1.5). $\square$

Corollary (6.7.3). — *For a Noetherian prescheme X to be affine, it is necessary and sufficient that the closed reduced subpreschemes whose underlying spaces are the irreducible components of X be affine.*

§7. VALUATIVE CRITERIA

In this section, we give valuative criteria for separation and properness for a given morphism, that is, criteria which introduce a variable auxiliary scheme of the form $\text{Spec}(A)$, where A is a valuation ring. Under certain suitable “Noetherian” hypotheses, we can refine our criteria and restrict to the case where A is a *discrete* valuation ring. This will be the only case that we need to concern ourselves with in all that follows, and we introduce arbitrary valuation rings, in the general case, only to discuss the links with the classical study of such objects.

7.1. Reminder on valuation rings

(7.1.1). Amongst the many diverse equivalent properties that characterise valuation rings, we will use the following: a ring A is said to be a *valuation ring* if it is an integral ring which is not a field, and A is *maximal* in the set of local rings strictly contained in the field of fractions K of A under the domination relation (I, 8.1.1). Recall that a valuation ring is *integrally closed*. If A is a valuation ring, then so too is $A_{\mathfrak{p}}$ for any prime ideal $\mathfrak{p} \neq 0$ of A .

(7.1.2). Let K be a field, and A a local subring of K that is not a field; [II:139] then there exists a valuation ring that both dominates A and has K as its field of fractions ([1, p. 1-07, lemma 2]).

Now let B be a valuation ring, k its residue field, K its field of fractions, and L an extension of k . Then there exists a *complete* valuation ring C that dominates B and whose residue field is L . Indeed, L is an algebraic extension of a pure transcendental extension $L' = k(T_{\mu})_{\mu \in M}$; we know that we can extend the valuation of B corresponding to B to a valuation of $K' = K(T_{\mu})_{\mu \in M}$ in such a way that L' is the residue field of this valuation ([24, p. 98]); replacing B by the completion of the ring of this extended valuation, we see that we can restrict to the case where B is complete and L is an algebraic closure of k . If $\bar{K}$ is an algebraic closure of K , we can then extend the valuation that defines B to $\bar{K}$, and the corresponding residue field is an algebraic closure of k , as we can see by lifting to $\bar{K}$ the coefficients of a monic polynomial of $k[T]$. We are thus finally led to the case where $L = k$, and it then suffices to take C to be the completion of B in order to satisfy our claim.

(7.1.3). Let K be a field, and A a subring of K ; the integral closure A' of A in K is the intersection of the valuation rings that contain A and have K as their field of fractions ([11, p. 51, th. 2]). Proposition (7.1.2) can then be expressed geometrically in an equivalent form:

Proposition (7.1.4). — *Let Y be a prescheme, $p : X \rightarrow Y$ a morphism, x a point of X , $y = p(x)$, and $y' \neq y$ a specialisation (0, 2.1.2) of y . Then there exists a local scheme Y' which is the spectrum of some valuation ring, and a separated morphism $f : Y' \rightarrow Y$ such that, denoting the unique closed point of Y' by a and the generic point of Y' by b , we have $f(a) = y'$ and $f(b) = y$. We can furthermore suppose that one of the following two additional properties is satisfied:*

- (i) *Y' is the spectrum of a complete valuation ring whose residue field is algebraically closed, and there exists a $k(y)$ -homomorphism $k(x) \rightarrow k(b)$.*
- (ii) *There exists a $k(y)$ -isomorphism $k(x) \xrightarrow{\sim} k(b)$.*

Proof. Let Y_1 be the closed reduced subprescheme of Y that has $\overline{\{y\}}$ as its underlying space (I, 5.2.1), and let X_1 be the closed subprescheme given by the inverse image $p^{-1}(Y_1)$; since $y' \in \overline{\{y\}}$ by hypothesis, and since $k(x)$ is the same in X and in X_1 , we can assume that Y is *integral*, with generic point y ; $\mathcal{O}_{y'}$ is then an integral local ring that is not a field, and whose field of fractions is $\mathcal{O}_y = k(y)$, and $k(x)$ is then an extension of $k(y)$. To satisfy the conditions $f(a) = y'$ and $f(b) = y$ as well as the additional condition (i) (resp. (ii)), we take $Y' = \text{Spec}(A')$, where A' is a complete valuation ring that dominates $\mathcal{O}_{y'}$ and whose algebraically closed residue field extends $k(x)$ (resp. a valuation ring that dominates $\mathcal{O}_{y'}$ and whose field of fractions is $k(x)$); the existence of such an A' is guaranteed by (7.1.2). $\square$

(7.1.5). Recall that a local ring A is said to be of *dimension 1* if there exists a prime ideal distinct from the maximal ideal $\mathfrak{m}$, and if every prime ideal of A distinct from $\mathfrak{m}$ is a *minimal* prime ideal; when A is *integral*, it is equivalent to ask that $\mathfrak{m}$ and (0) be the only prime ideals, with $\mathfrak{m} \neq (0)$; in other words, $Y = \text{Spec}(A)$ consists of two [II:140] points a and b : a is the unique *closed* point, we have $\mathfrak{j}_a = \mathfrak{m}$, and $k(a) = k$ is the *residue field* $k = A/\mathfrak{m}$; b is the *generic point* of Y , $\mathfrak{j}_b = (0)$, with the set $\{b\}$ being the unique open subset of Y distinct from both $\emptyset$ and Y (an open subset which is thus *everywhere dense*), and $k(b) = K$ is the *field of fractions* of A .

(7.1.6). For a Noetherian local ring A of dimension 1, we know ([1, pp. 2-08 and 17-01]) that the following conditions are equivalent:

- (a) A is normal;
- (b) A is regular;
- (c) A is a valuation ring;

furthermore, A is then a *discrete valuation ring*. Propositions (7.1.2) and (7.1.3) then have the following analogues for discrete valuation rings:

Proposition (7.1.7). — *Let A be an integral local Noetherian ring that is not a field, K its field of fractions, and L an extension of finite type of K ; then there exists a discrete valuation ring that dominates A and has L as its field of fractions.*

Proof. Suppose first of all that $L = K$. Let $\mathfrak{m}$ be the maximal ideal of A , $(x_1, \dots, x_n)$ a system of non-null generators of $\mathfrak{m}$, and B the subring $A[x_2/x_1, \dots, x_n/x_1]$ of K , which is Noetherian. It is immediate that the ideal $\mathfrak{m}B$ of B is identical to the principal ideal x_1B ; if $\mathfrak{p}$ is a minimal prime ideal over x_1B , then $\mathfrak{p}$ has height 1 ([13, t. I, p. 277]); in other words, $B_{\mathfrak{p}}$ is a local Noetherian ring of dimension 1; it is clear that $\mathfrak{p}B_{\mathfrak{p}} \cap A$ is an ideal of A that contains $\mathfrak{m}$ and that does not contain 1, and is thus equal to $\mathfrak{m}$, and so $B_{\mathfrak{p}}$ dominates A (I, 8.1.1). It follows from the Krull-Akizuki Theorem ([25, p. 293]) that the integral closure C of $B_{\mathfrak{p}}$ is a Noetherian ring (even though C is not necessarily a $B_{\mathfrak{p}}$ -module of finite type); if $\mathfrak{n}$ is a maximal ideal of C , then $C_{\mathfrak{n}}$ is a normal local Noetherian ring of dimension 1 ([25, p. 295]), and thus a discrete valuation ring that dominates $B_{\mathfrak{p}}$ and *a fortiori* A .

Now, if L is an extension of finite type of K , we can, by the above, restrict to the case where A is already a discrete valuation ring. Let w be a valuation of K associated to A ; there exists a discrete valuation w' of L that extends w : we can restrict, by induction on the number of generators of L , to the case where $L = K(\alpha)$, and then the proposition is classical ([24, p. 106]). $\square$

Corollary (7.1.8). — *Let A be a Noetherian integral ring, K its field of fractions, and L an extension of finite type of K . Then the integral closure of A in L is the intersection of the discrete valuation rings that have L as their field of fractions and that contain A .*

Proof. Indeed, such a discrete valuation ring, being normal, contains *a fortiori* every element of L that is integral over A . It thus suffices to prove that, if $x \in L$ is not integral over A , then there exists a discrete valuation ring C that has L as its field of fractions, contains A , and does not contain x . The hypothesis on x implies that $x \notin B = A[1/x]$, or, in other words, that $1/x$ is not invertible in the Noetherian ring B . There is thus a prime ideal $\mathfrak{p}$ of B that contains $1/x$. The integral local ring $B_{\mathfrak{p}}$ is Noetherian and contained in L , which is an extension of finite type of the field of fractions of $B_{\mathfrak{p}}$ (with the latter containing K). By (7.1.7), there thus exists a discrete valuation ring C that dominates $B_{\mathfrak{p}}$ and has L as its field of fractions; since $1/x \in \mathfrak{p}B_{\mathfrak{p}}$ belongs to the maximal ideal of C , we have that $x \notin C$, which concludes the proof. $\square$

The geometric form of (7.1.7) is the following:

[II:141]

Proposition (7.1.9). — *Let Y be a locally Noetherian prescheme, $p : X \rightarrow Y$ a morphism locally of finite type, x a point of X , $y = p(x)$, and $y' \neq y$ a specialisation of y . Then there exists a local scheme Y' , spectrum of a discrete valuation ring, a separated morphism $f : Y' \rightarrow Y$, and a rational Y -map g from Y' to X , such that, denoting the closed point of Y' by a , and the generic point of Y' by b , we have $f(a) = y'$, $f(b) = y$, $g(b) = x$, and such that, in the commutative diagram*

(7.1.9.1)

$$\begin{array}{ccc} & k(x) & \\ \gamma \swarrow & \uparrow \pi & \\ k(b) & \xleftarrow{\phi} & k(y) \end{array}$$

(where π , ϕ , and γ are the homomorphisms corresponding to p , f , and g , respectively) the homomorphism γ is bijective.

Proof. As in (7.1.4), we can restrict to the case where Y is integral with generic point y (taking (I, 6.3.4, iv) into account), and, since the question is local on X and Y , we can assume that p is of finite type; we are then in the situation of (7.1.4), with the additional property that $k(x)$ is an extension of finite type of $k(y)$ (I, 6.4.11) and that $\mathcal{O}_{y'}$ is Noetherian; this lets us apply (7.1.7) and take $Y' = \text{Spec}(A')$, where A' is a discrete valuation ring that dominates $\mathcal{O}_{y'}$ and whose field of fractions is $k(x)$. We have thus defined a commutative diagram (7.1.9.1) where γ is a bijection, with π and

ϕ corresponding to the morphisms p and f . Furthermore, since X and Y are locally Noetherian (I, 6.6.2) and since Y' is integral, there exists exactly one rational Y -map g from Y' to X to which corresponds the isomorphism γ (I, 7.1.15), which finishes the proof. $\square$

7.2. Valuitive criterion for separatedness

Proposition (7.2.1). — *Let X and Y be preschemes, and $f : X \rightarrow Y$ a quasi-compact morphism. The following two conditions are equivalent:*

- (a) *The morphism f is closed.*
- (b) *For all $x \in X$, and every specialisation $y' \neq y$ of $y = f(x)$, there exists a specialisation x' of x such that $f(x') = y'$.*

Proof. Condition (b) says that $f(\overline{\{x\}}) = \overline{\{y\}}$, and is thus a consequence of (a). To show that (b) implies (a), consider a closed subset X' of the underlying space X ; let $Y' = \overline{f(X')}$, and show that $Y' = f(X')$ as follows. Consider the closed reduced subpreschemes of X and Y whose underlying spaces are X' and Y' (respectively) (I, 5.2.1); there then exists a morphism $f' : X' \rightarrow Y'$ such that the diagram

$$\begin{array}{ccc} X' & \xrightarrow{f'} & Y' \\ \downarrow & & \downarrow \\ X & \xrightarrow{f} & Y \end{array}$$

commutes (I, 5.2.2), and, since f is quasi-compact, so too is f' . We are thus led to proving that, if f is a quasi-compact and *dominant* morphism, then ^[II:142] condition (b) implies that $f(X) = Y$. But let y' be a point of Y , and let y be the generic point of an irreducible component of Y that contains y' ; by (b), it suffices to show that $f^{-1}(y)$ is not empty. But we know that this property is a consequence of the fact that f is quasi-compact and dominant (I, 6.6.5). $\square$

Corollary (7.2.2). — *Let $f : X \rightarrow Y$ be a quasi-compact immersion. For the underlying space X to be closed in Y , it is necessary and sufficient for it to contain every specialisation (in Y) of all of its points.*

Proposition (7.2.3). — *Let Y be a prescheme (resp. a locally Noetherian prescheme), and $f : X \rightarrow Y$ a morphism (resp. a morphism locally of finite type). The following conditions are equivalent:*

- (a) *f is separated.*
- (b) *The diagonal morphism $X \rightarrow X \times_Y X$ is quasi-compact, and, for every Y -prescheme of the form $Y' = \text{Spec}(A)$, with A some valuation ring (resp. some discrete valuation ring), any two Y -morphisms from Y' to X that agree on the generic point of Y' are equal.*
- (c) *The diagonal morphism $X \rightarrow X \times_Y X$ is quasi-compact, and, for every Y -prescheme of the form $Y' = \text{Spec}(A)$, with A some valuation ring (resp. some discrete valuation ring), any two Y' -sections of $X' = X_{(Y')}$ that agree on the generic point of Y' are equal.*

Proof. The equivalence of (b) and (c) follows from the bijective correspondence between Y -morphisms from Y' to X and Y' -sections of X' (I, 3.3.14). If X is separated over Y , condition (b) is satisfied, by (I, 7.2.2.1), since Y' is integral. It remains to show that (b) implies that the diagonal morphism $\Delta : X \rightarrow X \times_Y X$ is closed, and it is equivalent to show that it satisfies the criterion of (7.2.2). But let z be a point of the diagonal $\Delta(X)$, and $z' \neq z$ a specialisation of z in $X \times_Y X$. There then exists (7.1.4) a valuation ring A and a morphism f from $Y' = \text{Spec}(A)$ to $X \times_Y X$ such that f sends the closed point a of Y' to z' , and the generic point b of Y' to z ; this morphism makes Y' an $(X \times_Y X)$ -prescheme, and *a fortiori* a Y -prescheme. If we compose the two projections of $X \times_Y X$ with f , then we obtain two Y -morphisms, g_1 and g_2 , from Y' to X , which, by hypothesis, agree on the point b ; they are thus equal to one single morphism g , which implies (I, 5.3.1) that f factors as $f = \Delta \circ g$, and thus $z' \in \Delta(X)$. If we suppose that Y is locally Noetherian and f is locally of finite type, then $X \times_Y X$ is locally Noetherian (I, 6.6.7); we can thus follow the same argument as before by supposing that A is a discrete valuation ring, by (7.1.9). $\square$

Remark (7.2.4). — (i) The hypothesis that the morphism Δ is quasi-compact is always satisfied whenever Y is locally Noetherian and f is locally of finite type, because $X \times_Y X$ is then

locally Noetherian (I, 6.6.4, i). In the general case, this also means that, for every cover (U_α) of X by affine opens, the sets $U_\alpha \cap U_\beta$ are *quasi-compact*.

- (ii) For f to be separated, it is *sufficient* to require condition (b) or (c) only for *complete* valuation rings A whose residue fields are *algebraically closed*; this follows from the proof of (7.2.3) and from (7.1.4).

[II:143]

7.3. Valuative criterion for properness

Proposition (7.3.1). — *Let A be a valuation ring, $Y = \text{Spec}(A)$, b the generic point of Y , X an integral scheme, and $f : X \rightarrow Y$ a closed morphism such that $f^{-1}(b)$ consists of a single point x and such that the corresponding homomorphism $k(b) \rightarrow k(x)$ is bijective. Then f is an isomorphism.*

Proof. Since f is closed and dominant, we have that $f(X) = Y$; it suffices (I, 4.2.2) to prove that, for all $y' \neq b$ in Y , there exists *exactly one* point x' such that $f(x') = y'$, and that the corresponding homomorphism $\mathcal{O}_{y'} \rightarrow \mathcal{O}_{x'}$ is bijective, since then f will be a homeomorphism. But if $f(x') = y'$ then $\mathcal{O}_{x'}$ is a local ring contained in $K = k(x) = k(b)$ and dominating $\mathcal{O}_{y'}$; the latter is the local ring $A_{y'}$, and is thus a valuation ring (7.1.1) that has K as its field of fractions. Also, $\mathcal{O}_{x'} \neq K$, since x' is not the generic point of X (0, 2.1.3); we thus conclude that $\mathcal{O}_{x'} = \mathcal{O}_{y'}$. Since X is an integral scheme, the fact that $\mathcal{O}_{x'} = \mathcal{O}_{x''}$ implies that $x' = x''$ (I, 8.2.2), which finishes the proof. $\square$

(7.3.2). Let A be a valuation ring, K its field of fractions, $Y = \text{Spec}(A)$, and b the generic point of Y , such that $\mathcal{O}_b = k(b)$ is equal to K ; let $f : X \rightarrow Y$ be a morphism. We know (I, 7.1.4) that the *rational Y -sections* of X are in bijective correspondence with the *germs* of Y -sections (defined in a neighbourhood of b) at the point b , whence we have a canonical map

$$(7.3.2.1) \quad \Gamma_{\text{rat}}(X/Y) \longrightarrow \Gamma(f^{-1}(b)/\text{Spec}(K))$$

with the elements of $\Gamma(f^{-1}(b)/\text{Spec}(K))$ being identified, by definition (I, 3.4.5), with the *points* of $f^{-1}(b) = X \otimes_A K$ that are *rational over K* . When f is *separated*, it follows from (I, 5.4.7) that the map (7.3.2.1) is *injective*, since Y is an integral scheme.

Composing (7.3.2.1) with the canonical map $\Gamma(X/Y) \rightarrow \Gamma_{\text{rat}}(X/Y)$ (I, 7.1.2), we obtain a canonical map

$$(7.3.2.2) \quad \Gamma(X/Y) \longrightarrow \Gamma(f^{-1}(b)/\text{Spec}(K)).$$

When f is *separated*, this map is again *injective* (I, 5.4.7).

Proposition (7.3.3). — *Let A be a valuation ring with field of fractions K , $Y = \text{Spec}(A)$, b the generic point of Y , and $f : X \rightarrow Y$ a separated and closed morphism. Then the canonical map (7.3.2.2) is bijective (which is equivalent to saying that it is surjective, and implies that the rational Y -sections of X are everywhere defined).*

Proof. So let x be a point of $f^{-1}(b)$ that is *rational over K* . Since f is separated, so too is the morphism $f^{-1}(b) \rightarrow \text{Spec}(K)$ corresponding to f (I, 5.5.1, iv), and, since every section of $f^{-1}(b)$ is a closed immersion (I, 5.4.6), $\{x\}$ is closed in $f^{-1}(b)$. Consider the closed reduced subprescheme X' of X that has the closure $\overline{\{x\}}$ of $\{x\}$ in X as its underlying space. It is clear that the restriction of f to X' satisfies the hypotheses of (7.3.1), and is thus an *isomorphism* from X' to Y , whose inverse isomorphism is the desired Y -section of X . $\square$

(7.3.4). To state the two following results, we use a terminology that will be justified and discussed in chapter IV: if F is a subset [II:144] of a prescheme Y , we define the *codimension* of F in Y , denoted $\text{codim}_Y F$, to be the lower bound of the integers $\dim(\mathcal{O}_z)$ over all z in F .

Corollary (7.3.5). — *Let Y be a locally Noetherian reduced prescheme, and N the set of points $y \in Y$ where Y is not regular (0, 4.1.4); suppose that $\text{codim}_Y N \geq 2$. Let $f : X \rightarrow Y$ be a morphism of finite type, both separated and closed, and let g be a rational Y -section of X ; if Y' is the set of points of Y where g is not defined (a set which is closed (I, 7.2.1)), then $\text{codim}_Y Y' \geq 2$.*

Proof. It suffices to prove that g is defined at every point $z \in Y$ such that $\dim \mathcal{O}_z \leq 1$. If $\dim \mathcal{O}_z = 0$, then z is the generic point of an irreducible component of Y (I, 1.1.14), and so belongs to every everywhere-dense open subset of Y , and, in particular, to the domain of definition of g . So suppose that $\dim \mathcal{O}_z = 1$; by hypothesis, $\mathcal{O}_z$ is then a regular Noetherian local ring, and thus, by (7.1.6), a discrete valuation ring. Let $Z = \operatorname{Spec}(\mathcal{O}_z)$; since $U = Y - Y'$ is everywhere dense, $U \cap Z$ is nonempty (I, 2.4.2); let g' be the rational map from Z to X induced by g (I, 7.2.8); it suffices to show that g' is a morphism (I, 7.2.9). But g' can be considered as a rational Z -section of the Z -prescheme $f^{-1}(Z) = X \times_Y Z$; it is clear that the morphism $f^{-1}(Z) \rightarrow Z$ corresponding to f is closed, and it follows from (I, 5.5.1, i) that it is separated; we thus conclude from (7.3.3) that g' is everywhere defined; since Z is reduced, and X is separated over Y , g' is a morphism (I, 7.2.2). $\square$

Corollary (7.3.6). — *Let S be a locally Noetherian prescheme, and X and Y both S -preschemes; suppose that Y is reduced, and further that the set N of points $y \in Y$ where Y is not regular is such that $\operatorname{codim}_Y N \geq 2$; suppose finally that the structure morphism $X \rightarrow S$ is proper. Let f be a rational S -map from Y to X , and let Y' be the points of Y where f is not defined; then $\operatorname{codim}_Y Y' \geq 2$.*

Proof. We know (I, 7.1.2) that we can identify the rational S -maps from Y to X with the rational Y -sections of $X \times_S Y$; since the structure morphism $X \times_S Y \rightarrow Y$ is closed (5.4.1), we can apply (7.3.5), whence the corollary. $\square$

Remark (7.3.7). — The hypotheses on Y in (7.3.5) and (7.3.6) will be satisfied in particular when Y is normal (0, 4.1.4), by (7.1.6).

We can characterise the universally closed morphisms (resp. proper morphisms) by a converse of (7.3.3):

Theorem (7.3.8). — *Let Y be a prescheme (resp. a locally Noetherian prescheme), and $f : X \rightarrow Y$ a quasi-compact separated morphism (resp. a morphism of finite type). The following conditions are equivalent:*

- (a) *f is universally closed (resp. proper).*
- (b) *For every Y -scheme of the form $Y' = \operatorname{Spec}(A)$, where A is a valuation ring (resp. a discrete valuation ring) with field of fractions K , the canonical map*

$$\operatorname{Hom}_Y(Y', X) \longrightarrow \operatorname{Hom}_Y(\operatorname{Spec}(K), X)$$

corresponding to the canonical injection $A \rightarrow K$ is surjective (resp. bijective). [II:145]

- (c) *For every Y -scheme of the form $Y' = \operatorname{Spec}(A)$, where A is a valuation ring (resp. a discrete valuation ring), the canonical map (7.3.2.2) with respect to the Y' -prescheme $X_{(Y')}$ is surjective (resp. bijective).*

Proof. The equivalence of (b) and (c) follows immediately from (I, 3.3.14); (a) implies (c), since (a) implies, in either case, that $f_{(Y')}$ is separated (I, 5.5.1, iv) and closed, and it suffices to apply (7.3.3). It remains to prove that (b) implies (a). We first consider the case where Y is arbitrary, and f is separated and quasi-compact. If condition (b) is satisfied by f , then it is also satisfied by $f_{(Y'')} : X_{(Y'')} \rightarrow Y''$, where Y'' is an arbitrary Y -prescheme, thanks to the equivalence between (b) and (c), and the fact that $X_{(Y'')} \times_{Y''} Y' = X \times_Y Y'$ for every morphism $Y' \rightarrow Y''$ (I, 3.3.9.1); since, further, $f_{(Y'')}$ is separated and quasi-compact whenever f is ((I, 5.5.1, iv) and (I, 6.6.4, iii)), we are led to proving that (b) implies that f is closed. For this, it suffices to verify condition (b) of (7.2.1). So let $x \in X$, and y' be a specialisation of $y = f(x)$, distinct from y ; by (7.1.4), there exists a scheme Y' , the spectrum of some valuation ring, and a separated morphism $g : Y' \rightarrow Y$ such that, letting a denote the closed point and b the generic point of Y' , we have that $g(a) = y'$, $g(b) = y$, and that there exists a $k(y)$ -homomorphism $k(x) \rightarrow k(b)$. The latter corresponds canonically to a Y -morphism $\operatorname{Spec}(k(b)) \rightarrow X$ (I, 2.4.6), and it thus follows from (b) that there exists a Y -morphism $h : Y' \rightarrow X$ to which the previous morphism corresponds. We then have that $h(b) = x$; if we set $h(a) = x'$, then x' is a specialisation of x , and we have that $f(x') = f(h(a)) = g(a) = y'$.

If now Y is locally Noetherian and f of finite type, then hypothesis (b) implies, first of all, that f is separated, by (7.2.3), with the diagonal morphism $X \rightarrow X \times_Y X$ being quasi-compact (7.2.4). Further, to show that f is proper, it suffices to show that $f_{(Y'')} : X_{(Y'')} \rightarrow Y''$ is closed for every Y -prescheme Y'' of finite type, taking (5.6.3) into account. Since Y'' is then locally Noetherian, we can follow the

same reasoning as in the first case by taking Y' to be the spectrum of a discrete valuation ring, and applying (7.1.9) instead of (7.1.4). $\square$

- Remarks (7.3.9).** — (i) Whenever Y is an arbitrary prescheme and f a separated morphism, for f to be universally closed, it suffices that condition (b) or (c) be satisfied for the *complete* valuation rings A whose residue field is *algebraically closed*; this follows from the above proof and from (7.1.4).
- (ii) From criterion (c) of (7.3.8) we obtain a new proof of the fact that a projective morphism $X \rightarrow Y$ is closed (5.5.3), and it is closer to the classical approach. We can indeed assume that Y is affine, and thus that X can be identified with a closed subscheme of a projective bundle $\mathbf{P}_Y^n$ (5.3.3); to prove that $X \rightarrow Y$ is closed, it suffices to verify that the structure morphism $\mathbf{P}_Y^n \rightarrow Y$ is closed, and criterion (c) of (7.3.8), combined with (4.1.3.1), tells us that we can reduce to proving the following fact: *if Y is the spectrum of a valuation ring A , with field of fractions K , then every point of $\mathbf{P}_Y^n$ with values in K comes from (by restriction to the generic point of Y) a point of $\mathbf{P}_Y^n$ with values in A .* But every invertible $\mathcal{O}_Y$ -module is trivial (I, 2.4.8); so it follows from (4.2.6) that a point of $\mathbf{P}_Y^n$ with values in K can be identified with a class of tuples $(\zeta c_0, \zeta c_1, \dots, \zeta c_n)$, where $\zeta \in K^\times$ and the $c_i \in K$ are not all zero. However, by multiplying the c_i by an element of A of ^[II:146] suitable valuation, we can suppose that the c_i all belong to A , and that at least one of them is invertible. But then, by (4.2.6), the system $(c_0, \dots, c_n)$ also defines a point of $\mathbf{P}_Y^n$ with values in A , which proves our claim.
- (iii) Criteria (7.2.3) and (7.3.8) are particularly simple when we consider the data of a Y -prescheme X as being equivalent to the data of the functor

$$X(Y') = \text{Hom}_Y(Y', X)$$

for Y -preschemes Y' ; these criteria allow us, for example, to prove that, under certain conditions, the “Picard schemes” are proper.

Corollary (7.3.10). — *Let Y be an integral scheme (resp. a locally Noetherian integral scheme), X an integral scheme, and $f : X \rightarrow Y$ a dominant morphism.*

- (i) *If f is quasi-compact and universally closed, then every valuation ring whose field of fractions is the field $R(X)$ of rational functions on X , and which dominates a local ring of Y , also dominates a local ring of X .*
- (ii) *Conversely, suppose that f is of finite type, and that the property described in (i) is verified by every valuation ring (resp. every discrete valuation ring) that has $R(X)$ as its field of fractions. Then f is proper.*

Proof. Note first of all that the hypotheses imply, in any case, that f is separated (I, 5.5.9).

- (i) Let $K = R(Y)$, $L = R(X)$, y a point of Y , and A a valuation ring that dominates $\mathcal{O}_y$ and has L as its field of fractions; the injection $\mathcal{O}_y \rightarrow A$ then defines a morphism h from $Y' = \text{Spec}(A)$ to Y (I, 2.4.4) such that $h(a) = y$, where we write a to denote the closed point of Y' ; furthermore, if η is the generic point of Y , which is also the generic point of $\text{Spec}(\mathcal{O}_y)$, then we have $h(b) = \eta$, writing b to denote the generic point of Y' (since $K \subset L$ by hypothesis). If ξ is the generic point of X , then $k(\xi) = k(b) = L$ by hypothesis, whence we have a Y -morphism $g : \text{Spec}(L) \rightarrow X$ such that $g(b) = \xi$; by (7.3.8), g comes from a Y -morphism $g' : Y' \rightarrow X$. If $x = g'(a)$, it is clear that A dominates $\mathcal{O}_x$.
- (ii) Since the question is local on Y , we can always suppose that Y is affine (resp. affine and Noetherian). Since f is of finite type, we can apply, in either case, Chow’s lemma (5.6.1). There is thus a projective morphism $p : P \rightarrow Y$, an immersion $j : X' \rightarrow P$, and a projective morphism $g : X' \rightarrow X$ that is both surjective and birational, with X' integral, such that the diagram

$$\begin{array}{ccc} P & \xleftarrow{j} & X' \\ p \downarrow & & \downarrow g \\ Y & \xleftarrow{f} & X \end{array}$$

commutes. It suffices to prove that j is a *closed* immersion, since then $f \circ g = p \circ j$ will be a projective morphism, and thus proper, and, since g is surjective, f will also be proper (5.4.3). Let Z be the closed reduced subscheme of P that has $\overline{j(X')}$ as its underlying space (I, 5.2.1); since X' is integral, j factors as $i \circ h$, where $i : Z \rightarrow P$ is the canonical injection, $h : X' \rightarrow Z$ a dominant open immersion (I, 5.2.3), and Z is integral; [II:147] furthermore, Z is projective over Y , and we see that we can restrict to the case where P is *integral* and j is *dominant* and *birational*, and everything then reduces to showing that j is *surjective*. But let $z \in P$; then $\mathcal{O}_z$ is an integral (resp. integral and Noetherian) local ring whose field of fractions is

$$L = R(P) = R(X') = R(X).$$

We can restrict to the case where z is not the generic point of P . There then exists ((7.1.2) and (7.1.7)) a valuation ring (resp. a discrete valuation ring) A which dominates $\mathcal{O}_z$ and has L as its field of fractions. *A fortiori*, A dominates $\mathcal{O}_y$, where $y = p(z)$, and, by hypothesis, there thus exists some $x \in X$ such that A dominates $\mathcal{O}_x$. Since g is proper, the first part of the proof shows that A also dominates $\mathcal{O}_{x'}$ for some $x' \in X'$; it then follows that $\mathcal{O}_z$ and $\mathcal{O}_{j(x')} = \mathcal{O}_{x'}$ are allied (I, 8.1.4), and, since P is a scheme, this implies that $z = j(x')$ (I, 8.2.2) and finishes the proof. □

Corollary (7.3.11). — *Let X and Y be integral schemes, and $f : X \rightarrow Y$ a dominant, quasi-compact, and universally closed morphism. Suppose further that Y is affine of (integral) ring B . Then $\Gamma(X, \mathcal{O}_X)$ is canonically isomorphic to a subring of the integral closure of B in $R(X)$.*

Proof. Indeed (I, 8.2.1.1), $\Gamma(X, \mathcal{O}_X)$ can be identified with the intersection of the $\mathcal{O}_x$ over $x \in X$; by (7.3.10), (7.1.2), and (7.1.3), $\Gamma(X, \mathcal{O}_X)$ is then contained in the intersection of the valuation rings that contain B and that have $R(X)$ as their field of fractions; the conclusion then follows from (7.1.3). □

Remarks (7.3.12). — Under the hypotheses of (7.3.11), and when we suppose that $R(X)$ is an extension of finite type of $R(Y)$, we can, in many cases, conclude that $\Gamma(X, \mathcal{O}_X)$ is a *module of finite type* over the ring $B = \Gamma(Y, \mathcal{O}_Y)$. For example, this will be the case whenever B is an *algebra of finite type over a field*, since we then know that the integral closure of B in an extension of finite type of its field of fractions is a B -module of finite type ([13, t. I, p. 267, th. 9]); the conclusion then follows from (7.3.11) and the fact that B is Noetherian.

In particular, a *proper affine scheme X over a field K is finite*. Indeed, by (1.6.4), (5.4.6), and (I, 6.4.4, (c)), we can restrict to the case where X is *reduced*. Furthermore, it suffices to prove that each of the closed subschemes of X that have an irreducible component of X as their underlying space (of which there are finitely many) is finite over K , which means (taking (5.4.5) into account) that we are finally reduced to the case where X is *integral*. But then the result follows from the above remarks.

In chapter III, we will again prove the preceding proposition by other methods, and as a consequence of more general results, by showing that, if $f : X \rightarrow Y$ is proper and Y is locally Noetherian, $f_*(\mathcal{F})$ is coherent for any coherent $\mathcal{O}_X$ -module $\mathcal{F}$ (III, 4.4.2).

Finally, note that criterion (7.3.10) is taken as the *definition* of proper morphisms in classical algebraic geometry. We only mention this here as a remark, since criterion (7.3.8) seems more manageable in all the applications with which we are familiar.

7.4. Algebraic curves and function fields of dimension 1 The aim of this section is to show how to formulate the classical notion of algebraic curves (as presented, for example, in the book of C. Chevalley [23]) in the language of schemes. Throughout this section, *we write k to mean a field, all the schemes in question are k -schemes of finite type, and all the morphisms are k -morphisms.*

Proposition (7.4.1). — *Let X be a prescheme of finite type over k (and thus Noetherian); let x_i ($1 \leq i \leq n$) be the generic points of the irreducible components X_i of X , and let $K_i = k(x_i)$ ($1 \leq i \leq n$). Then the following conditions are equivalent:*

- (a) *Each of the K_i is an extension of k with transcendence degree equal to 1.*
- (b) *For every closed point x of X , the local ring $\mathcal{O}_x$ is of dimension 1 (7.1.5).*
- (c) *The closed irreducible subsets of X that are distinct from the X_i are exactly the closed points of X .*

Proof. Since X is quasi-compact, every closed irreducible subset F of X contains a closed point (0, 2.1.3). By (I, 2.4.2), there is a bijective correspondence between the prime ideals of $\mathcal{O}_x$ and the closed irreducible subsets of X that contain x (I, 1.1.14); the equivalence between (b) and (c) follows immediately from this. Now, if $\mathfrak{p}_\alpha$ ($1 \leq \alpha \leq r$) are the minimal prime ideals of the local Noetherian ring $\mathcal{O}_x$, then the local rings $\mathcal{O}_x/\mathfrak{p}_\alpha$ are integral, and have the K_i such that $x \in X_i$ as their fields of fractions. Furthermore, we know ([1, p. 4-06, th. 2]) that the dimension of a local integral k -algebra of finite type is equal to the transcendence degree over k of its field of fractions. Finally, the dimension of $\mathcal{O}_x$ is the supremum of the dimensions of the $\mathcal{O}_x/\mathfrak{p}_\alpha$; but condition (a) implies that these dimensions are equal to 1, and so (a) implies (b); conversely, if $\mathcal{O}_x$ is of dimension 1, then none of the $\mathfrak{p}_\alpha$ can be equal to the maximal ideal of $\mathcal{O}_x$, otherwise $\mathcal{O}_x$ would be of dimension 0; thus each of the $\mathcal{O}_x/\mathfrak{p}_\alpha$ is of dimension 1, which shows that (b) implies (a). $\square$

We note that, under the conditions of (7.4.1), the set X is either *empty* or *infinite*, as an immediate result of (I, 6.4.4).

Definition (7.4.2). — We define an *algebraic curve over k* to be a non-empty algebraic scheme over k that satisfies the conditions of (7.4.1).

In the language of dimensions, which will be introduced in Chapter IV, this can be expressed by saying that an algebraic curve over k is a non-empty algebraic k -scheme *whose irreducible components are all of dimension 1*.

We note that, if X is an algebraic curve over k , then the closed reduced subpreschemes X_i ($1 \leq i \leq n$) of X whose underlying spaces are the irreducible components of X are also algebraic curves over k .

Corollary (7.4.3). — *Let X be an irreducible algebraic curve. The only non-closed point of X is its generic point. The closed subsets of X that are distinct from X are the finite sets of closed points; these are also the only subsets of X that are not everywhere dense.*

Proof. If a point $x \in X$ is not closed, then its closure in X is an irreducible closed subset of X , and thus necessarily the whole of X , by (7.4.1), and thus x is the generic point of X . A closed subset F of X that is distinct from X cannot contain [III:149] the generic point of X , and so all its points are closed (in X , and *a fortiori* in F); by considering the closed reduced subpreschemes of X that have F as their underlying space (I, 5.2.1), it thus follows from (I, 6.2.2) that F is finite and discrete. The closure in X of any infinite subset of X is thus necessarily equal to X itself. $\square$

If X is an arbitrary algebraic curve, by applying (7.4.3) to the irreducible components of X , we see that the only non-closed points of X are the generic points of these components.

Corollary (7.4.4). — *Let X and Y be irreducible algebraic curves over k , and $f : X \rightarrow Y$ a k -morphism. For f to be dominant, it is necessary and sufficient for $f^{-1}(y)$ to be finite for all $y \in Y$.*

Proof. Indeed, if f is not dominant, then $f(X)$ is necessarily a *finite* subset of Y , by (7.4.3), and so it is not possible for $f^{-1}(y)$ to be finite for every point of Y , since otherwise X would be finite, which is a contradiction (7.4.1). Conversely, if f is dominant, then for any $y \in Y$ distinct from the generic point η of Y , we have that $f^{-1}(y)$ is closed in X , since $\{y\}$ is closed in Y (7.4.3); also, by hypothesis, $f^{-1}(y)$ does not contain the generic point ζ of X , and is thus finite, by (7.4.3). Finally, to see that, when f is dominant, $f^{-1}(\eta)$ is finite, we note that the fibre $f^{-1}(\eta)$ is an irreducible scheme of finite

type over $k(\eta)$, and with generic point ξ ((I, 6.3.9) and (I, 6.4.11)). Since $k(\xi)$ and $k(\eta)$ are extensions of finite type of k , both of transcendence degree 1, we have that $k(\xi)$ is necessarily an extension of finite degree of $k(\eta)$, and so ξ is closed in $f^{-1}(\eta)$ (I, 6.4.2), and $f^{-1}(\eta)$ thus consists of a single point ξ . $\square$

We will see, in Chapter III, that, if $f : X \rightarrow Y$ is a *proper* morphism of Noetherian preschemes such that $f^{-1}(y)$ is finite for all $y \in Y$, then f is necessarily *finite*; it will thus follow from (7.4.4) that a proper dominant morphism from an irreducible algebraic curve to an algebraic curve is *finite*.

Corollary (7.4.5). — *Let X be an algebraic curve over k . For X to be regular, it is necessary and sufficient for X to be normal, or for the local rings of its closed points to be discrete valuation rings.*

Proof. This follows immediately from (7.4.1, (b)) and (7.1.6). $\square$

Corollary (7.4.6). — *Let X be a reduced algebraic curve, and $\mathcal{A}$ a reduced coherent $\mathcal{R}(X)$ -algebra; then the integral closure X' of X with respect to $\mathcal{A}$ (6.3.4) is a normal algebraic curve, and the canonical morphism $X' \rightarrow X$ is finite.*

Proof. The fact that $X' \rightarrow X$ is finite follows from (6.3.10); X' is thus an algebraic k -scheme; furthermore, if x_i ($1 \leq i \leq n$) are the generic points of the irreducible components of X , and x'_j ($1 \leq j \leq m$) the generic points of the irreducible components of X' , then each of the $k(x'_j)$ is a finite algebraic extension of one of the $k(x_i)$ (6.3.6), and thus of transcendence degree 1 over k . So X' is indeed an algebraic curve over k , and, furthermore, we know that X' is a finite sum of normal integral schemes ((6.3.6) and (6.3.7)). $\square$

(7.4.7). We say that an algebraic curve X over k is *complete* if it is *proper* over k .

Corollary (7.4.8). — *For a reduced algebraic curve X over k to be complete, it is necessary and sufficient for its normalisation X' to be complete.*

[II:150]

Proof. The canonical morphism $f : X' \rightarrow X$ is finite (7.4.6), and thus proper (6.1.11) and surjective (6.3.8); if $g : X \rightarrow \text{Spec}(k)$ is the structure morphism, then g is proper if and only if $g \circ f$ is proper, by (5.4.2, (ii)) and (5.4.3, (ii)), since g is separated by hypothesis. $\square$

Proposition (7.4.9). — *Let X be a normal algebraic curve over k , and Y a proper algebraic k -scheme. Then every rational k -map from X to Y is everywhere defined, or, in other words, is a morphism.*

Proof. It follows from (7.3.7) that, at the points $x \in X$ where such a map is not defined, the dimension of $\mathcal{O}_x$ must be ≥ 2 , and so the set of such points is empty; the final claim follows from (I, 7.2.3). $\square$

Corollary (7.4.10). — *A normal algebraic curve over k is quasi-projective over k .*

Proof. Since X is a finite sum of normal integral algebraic curves (6.3.8), we can restrict to the case where X is integral (5.3.6). Since X is quasi-compact, it is covered by a finite number of affine open subsets U_i ($1 \leq i \leq n$), and, since each of these U_i is of finite type over k , for each i there exists some integer n_i along with a k -immersion $f_i : U_i \rightarrow \mathbf{P}_k^{n_i}$ ((5.3.3) and (5.3.4, (i))). Since U_i is dense in X , it follows from (7.4.9) that f_i can be extended to a k -morphism $g_i : X \rightarrow \mathbf{P}_k^{n_i}$, whence we obtain a k -morphism $g = (g_1, \dots, g_n)_k$ from X to the product P of the $\mathbf{P}_k^{n_i}$ over k . Furthermore, for each i , since the restriction of g_i to U_i is an immersion, so too is the restriction of g to U_i (I, 5.3.14). Since the U_i cover X , and since g is separated (I, 5.5.1, (v)), g is an immersion from X into P (I, 8.2.8). Since the Segre morphism (4.3.3) gives an immersion of P into $\mathbf{P}_k^N$, this proves that X is quasi-projective. $\square$

Corollary (7.4.11). — *Any normal algebraic curve X is isomorphic to an everywhere-dense open subscheme of a complete normal algebraic curve $\hat{X}$, and this $\hat{X}$ is unique up to unique isomorphism.*

Proof. If X_1 and X_2 are complete normal curves, then it follows from (7.4.9) that every isomorphism from any dense open U_1 in X_1 to any dense open U_2 in X_2 can be uniquely extended to an isomorphism from X_1 to X_2 ; whence the uniqueness claim. To prove the existence of $\hat{X}$, it suffices to note that we can consider X as a subscheme of a projective bundle $\mathbf{P}_k^n$ (7.4.10). Let $\bar{X}$ be the closure of X

in $\mathbf{P}_k^n$ (I, 9.5.11); since X is induced by $\bar{X}$ on a dense open subset of $\bar{X}$ (I, 9.5.10), the generic points x_i of the irreducible components of X are also the generic points of the irreducible components of $\bar{X}$, and the $k(x_i)$ are the same for both of these schemes, and so (7.4.1) $\bar{X}$ is an algebraic curve over k that is reduced (I, 9.5.9) and projective over k (5.5.1), whence complete (5.5.3). So we take for $\hat{X}$ the normalisation of $\bar{X}$, which is again complete (7.4.8); furthermore, if $h : \hat{X} \rightarrow \bar{X}$ is the canonical morphism, then the restriction of h to $h^{-1}(X)$ is an isomorphism to X , since X is normal (6.3.4), and since $h^{-1}(X)$ contains the generic points of the irreducible components of $\hat{X}$ (6.3.8), it is dense in $\hat{X}$, which finishes the proof. $\square$

[II:151]

Remark (7.4.12). — We will show, in Chapter V, that the conclusion of (7.4.10) still holds true without the assumption that the curve is normal (or even reduced); we will also show that, for an algebraic curve (reduced or not) to be affine, it is necessary and sufficient for its (reduced) irreducible components to *not* be complete.

Corollary (7.4.13). — *Let X be a normal irreducible curve over k , with function field $K = R(X)$, and let Y be a complete integral curve over k , with function field $L = R(Y)$. Then there is a canonical bijective correspondence between dominant k -morphisms $X \rightarrow Y$ and k -monomorphisms $L \rightarrow K$.*

Proof. By (7.4.9), rational k -maps from X to Y can be identified with k -morphisms $u : X \rightarrow Y$. Since the dominant morphisms $u : X \rightarrow Y$ are characterised by being those such that $u(x) = y$ (writing x and y to denote the generic points of X and Y , respectively), the corollary follows from these remarks and from (I, 7.1.13). $\square$

(7.4.14). We can refine the result of (7.4.13) in the case where Y is the projective line $\mathbf{P}_k^1 = \text{Proj}(k[T_0, T])$, where T_0 and T are indeterminates. Then Y is an integral scheme (2.4.4), and the scheme induced on the open subset $D_+(T_0)$ of Y is isomorphic to $\text{Spec}(k[T])$ (2.3.6), and so the generic point of Y is the ideal (0) of $k[T]$, and the field of rational functions of Y is $k(T)$, which proves that Y is a complete algebraic curve over k . Furthermore, the only graded prime ideal of $S = k[T_0, T]$ that contains T_0 and is distinct from S_+ is the principal ideal (T_0) , and so the complement of $D_+(T_0)$ in $Y = \mathbf{P}_k^1$ consists of one closed point, called the “point at infinity”, which we denote by ∞ (for a general study of the links between vector bundles and projective bundles, see (8.4)). With these notations:

Corollary (7.4.15). — *Let X be a normal irreducible curve over k , with function field $K = R(X)$. Then there exists a canonical bijective correspondence between the set K and the set of morphisms u from X to $\mathbf{P}_k^1$ that are distinct from the constant morphism with value ∞ . For such a morphism to be dominant, it is necessary and sufficient for the corresponding element of K to be transcendental over k .*

Proof. This claim follows immediately from (7.4.9) and the following:

Lemma. — *Let X be an integral prescheme over k , and let $K = R(X)$ be its field of rational functions. Then there exists a canonical bijective correspondence between the set K and the set of rational maps u from X to $\mathbf{P}_k^1$ that are distinct from the constant morphism with value ∞ . For such a rational map to be dominant, it is necessary and sufficient for the corresponding element of K to be transcendental over k .*

First of all, rational maps from X to $\mathbf{P}_k^1$ correspond bijectively to points of $\mathbf{P}_k^1$ with values in the extension K of k (I, 7.1.12). If such a point is located (I, 3.4.5) at the generic point of $\mathbf{P}_k^1$, then the corresponding rational map is clearly dominant. In the converse case, since every point of $\mathbf{P}_k^1$ that is distinct from the generic point is closed (7.4.3), the image of the domain of definition U of u by the unique morphism $U \rightarrow \mathbf{P}_k^1$ of the class u (I, 7.2.2) consists of one closed point y of $\mathbf{P}_k^1$, and this morphism (which is not necessarily everywhere defined on X) is thus not dominant; as an abuse of language, we thus say that the rational map u is “constant, of value y ”. It remains to place in bijective correspondence the points of $\mathbf{P}_k^1$ with value in K that are located (I, 3.4.5) *not* at ∞ , and the elements of K , and then to verify that the location of such a point is the generic point of $\mathbf{P}_k^1$ if and only if it corresponds to an [II:152] element that is transcendental over k . But this is immediate (4.2.6, example 1). $\square$

Corollary (7.4.16). — *Let X and Y be algebraic curves over k that are normal, complete, and irreducible; let $K = R(X)$ and $L = R(Y)$ be their fields. Then there exists a canonical bijective correspondence between the set of k -isomorphisms $X \xrightarrow{\sim} Y$ and the set of k -isomorphisms $L \xrightarrow{\sim} K$.*

Proof. This is an evident consequence of (7.4.13). $\square$

(7.4.17). This corollary (7.4.16) shows that an algebraic curve over k that is normal, complete, and irreducible, is determined by its field of rational functions K up to unique isomorphism; by definition, K is an extension of finite type of k , of transcendence degree 1 (we classically call this a *field of algebraic functions of one variable*). Furthermore:

Proposition (7.4.18). — *For every extension K of k of finite type and of transcendence degree 1, there exists an algebraic curve X (determined up to unique isomorphism) that is normal, complete, and irreducible, and such that $R(X) = K$. The set of local rings of X can be identified (I, 8.2.1) with the set consisting of the field K and the valuation rings that contain k and have K as their field of fractions.*

Proof. Indeed, K is an extension of finite degree of a pure transcendental extension $k(T)$ of k , which can be identified, as we have seen, with the field of rational functions of the projective line $Y = \mathbf{P}_k^1$. Let X be the integral closure of Y with respect to K (6.3.4); then X is a normal algebraic curve over k , with function field K (6.3.7), and it is complete, since the morphism $X \rightarrow Y$ is finite (7.4.6). The local rings $\mathcal{O}_x$ of X are either the field K , when x is the generic point, or discrete valuation rings that contain k and have K as their field of fractions, when x is distinct from the generic point (7.4.5). Conversely, let A be such a ring; since the morphism $X \rightarrow \text{Spec}(k)$ is proper, the fact that A dominates k implies that A also dominates a local ring $\mathcal{O}_x$ of X (7.3.10); since the latter is a valuation ring that has K as a field of fractions, it is necessarily equal to A . $\square$

Remarks (7.4.19). — It follows from (7.4.16) and (7.4.18) that the data of an algebraic curve over k that is normal, complete, and irreducible, is essentially equivalent to the data of an extension K of k that is of finite type and of transcendence degree 1. We note that, if k' is an extension of the base field k , then $X \otimes_k k'$ will again be a complete algebraic curve over k' (5.4.2, (iii)), but, in general, it will be neither reduced nor irreducible. It will, however, be both reduced and irreducible if K is a separable extension of k , and k is algebraically closed in K (this can be expressed, in classical terminology, which we will not use, by saying that K is a “regular extension” of k). But even in this case, it is possible for $X \otimes_k k'$ to not be normal. The reader will find details on these questions in Chapter IV.

§8. BLOWUP SCHEMES; PROJECTING CONES; PROJECTIVE CLOSURE

8.1. Blowup preschemes

(8.1.1). Let Y be a prescheme, and, for every integer $n \geq 0$, let $\mathcal{I}_n$ be a quasi-coherent sheaf of ideals of $\mathcal{O}_Y$; suppose that the following conditions are satisfied:

$$(8.1.1.1) \quad \mathcal{I}_0 = \mathcal{O}_Y, \quad \mathcal{I}_n \subset \mathcal{I}_m \text{ for } m \leq n,$$

$$(8.1.1.2) \quad \mathcal{I}_m \mathcal{I}_n \subset \mathcal{I}_{m+n} \text{ for any } m, n.$$

[II:153] We note that these hypotheses imply

$$(8.1.1.3) \quad \mathcal{I}_1^n \subset \mathcal{I}_n.$$

Set

$$(8.1.1.4) \quad \mathcal{S} = \bigoplus_{n \geq 0} \mathcal{I}_n.$$

It follows from (8.1.1.1) and (8.1.1.2) that $\mathcal{S}$ is a quasi-coherent graded $\mathcal{O}_Y$ -algebra, and thus defines a Y -scheme $X = \text{Proj}(\mathcal{S})$. If $\mathcal{J}$ is an invertible sheaf of ideals of $\mathcal{O}_Y$, then $\mathcal{I}_n \otimes_{\mathcal{O}_Y} \mathcal{J}^{\otimes n}$ is canonically identified with $\mathcal{I}_n \mathcal{J}^n$. If we then replace the $\mathcal{I}_n$ by the $\mathcal{I}_n \mathcal{J}^n$, and, in doing so, replace $\mathcal{S}$ by a quasi-coherent $\mathcal{O}_Y$ -algebra $\mathcal{S}(\mathcal{J})$, then $X(\mathcal{J}) = \text{Proj}(\mathcal{S}(\mathcal{J}))$ is canonically isomorphic to X (3.1.8).

(8.1.2). Suppose that Y is *locally integral*, so that the sheaf $\mathcal{R}(Y)$ of rational functions is a quasi-coherent $\mathcal{O}_Y$ -algebra (I, 7.3.7). We say that an $\mathcal{O}_Y$ -submodule $\mathcal{I}$ of $\mathcal{R}(Y)$ is a *fractional ideal* of $\mathcal{R}(Y)$ if it is of *finite type* (0, 5.2.1). Suppose we have, for all $n \geq 0$, a quasi-coherent fractional ideal $\mathcal{I}_n$ of $\mathcal{R}(Y)$, such that $\mathcal{I}_0 = \mathcal{O}_Y$, and such that condition (8.1.1.2) (but not necessarily the second condition (8.1.1.1)) is satisfied; we can then again define a quasi-coherent graded $\mathcal{O}_Y$ -algebra by Equation (8.1.1.4), and the corresponding Y -scheme $X = \text{Proj}(\mathcal{S})$; we will again have a canonical isomorphism from X to $X_{\mathcal{I}}$ for every *invertible* fractional ideal $\mathcal{I}$ of $\mathcal{R}(Y)$.

Definition (8.1.3). — Let Y be a prescheme (resp. a locally integral prescheme), and $\mathcal{I}$ a quasi-coherent ideal of $\mathcal{O}_Y$ (resp. a quasi-coherent fractional ideal of $\mathcal{R}(Y)$). We say that the Y -scheme $X = \text{Proj}(\bigoplus_{n \geq 0} \mathcal{I}^n)$ is obtained by blowing up the ideal $\mathcal{I}$, or is the blow-up prescheme of Y relative to $\mathcal{I}$. When $\mathcal{I}$ is a quasi-coherent ideal of $\mathcal{O}_Y$, and Y' is the closed subscheme of Y defined by $\mathcal{I}$, we also say that X is the Y -scheme obtained by blowing up Y' .

By definition, $\mathcal{S} = \bigoplus_{n \geq 0} \mathcal{I}^n$ is then generated by $\mathcal{S}_1 = \mathcal{I}$; if $\mathcal{I}$ is an $\mathcal{O}_Y$ -module of *finite type*, then X is *projective* over Y (5.5.2). Without any hypotheses on $\mathcal{I}$, the $\mathcal{O}_X$ -module $\mathcal{O}_X(1)$ is *invertible* (3.2.5) and *very ample*, by (4.4.3) applied to the structure morphism $X \rightarrow Y$.

We note that, if $f : X \rightarrow Y$ is the structure morphism, then the restriction of f to $f^{-1}(Y - Y')$ is an *isomorphism* onto $Y - Y'$ whenever $\mathcal{I}$ is an *ideal* of $\mathcal{O}_Y$ and Y' is the closed subscheme that it defines: indeed, since the question is local on Y , it suffices to assume that $\mathcal{I} = \mathcal{O}_Y$, and our claim then follows from (3.1.7).

If we replace $\mathcal{I}$ by $\mathcal{I}^d$ ($d > 0$), then the blow-up Y -scheme X is replaced by a canonically isomorphic Y -scheme X' (8.1.1); similarly, for every *invertible* ideal (resp. *invertible* fractional ideal) $\mathcal{J}$, the blow-up prescheme $X_{(\mathcal{I})}$ relative to the ideal $\mathcal{I}\mathcal{J}$ is canonically isomorphic to X (8.1.1).

In particular, whenever $\mathcal{I}$ is an *invertible* ideal (resp. *invertible* fractional ideal), the Y -scheme obtained by blowing up $\mathcal{I}$ is *isomorphic* to Y (3.1.7).

Proposition (8.1.4). — Let Y be an *integral* prescheme.

- (i) For every sequence $(\mathcal{I}_n)$ of quasi-coherent fractional ideals of $\mathcal{R}(Y)$ that satisfies (8.1.1.2) [II:154] and such that $\mathcal{I}_0 = \mathcal{O}_Y$, the Y -scheme $X = \text{Proj}(\bigoplus_{n \geq 0} \mathcal{I}_n)$ is *integral*, and the structure morphism $f : X \rightarrow Y$ is *dominant*.
- (ii) Let $\mathcal{I}$ be a quasi-coherent fractional ideal of $\mathcal{R}(Y)$, and let X be the Y -scheme given by the blow up of Y relative to $\mathcal{I}$. If $\mathcal{I} \neq 0$, then the structure morphism $f : X \rightarrow Y$ is then *birational* and *surjective*.

Proof.

- (i) This follows from the fact that $\mathcal{S} = \bigoplus_{n \geq 0} \mathcal{I}_n$ is an *integral* $\mathcal{O}_Y$ -algebra ((3.1.12) and (3.1.14)), since, for all $y \in Y$, $\mathcal{O}_y$ is an integral ring (I, 5.1.4).
- (ii) By (i), X is *integral*; if, furthermore, x and y are the generic points of X and Y (respectively), then we have $f(x) = y$, and it remains to show that $k(x)$ has dimension one over $k(y)$. But x is also the generic point of the fibre $f^{-1}(y)$; if ψ is the canonical morphism $Z \rightarrow Y$, where $Z = \text{Spec}(k(y))$, then the prescheme $f^{-1}(y)$ can be identified with $\text{Proj}(\mathcal{S}')$, where $\mathcal{S}' = \psi^*(\mathcal{S})$ (3.5.3). But it is clear that $\mathcal{S}' = \bigoplus_{n \geq 0} (\mathcal{I}_y)^n$, and, since $\mathcal{I}$ is a quasi-coherent fractional ideal of $\mathcal{R}(Y)$ that is not zero, $\mathcal{I}_y \neq 0$ (I, 7.3.6), whence $\mathcal{I}_y = k(y)$; then $\text{Proj}(\mathcal{S}')$ can be identified with $\text{Spec}(k(y))$ (3.1.7), whence the conclusion. □

We show a *converse* of (8.1.4) in (III, 2.3.8).

(8.1.5). We return to the setting and notation of (8.1.1). By definition, the injection homomorphisms $\mathcal{S}_{n+1} \rightarrow \mathcal{S}_n$ (8.1.1.1) define, for every $k \in \mathbf{Z}$, an injective homomorphism of degree zero of graded $\mathcal{S}$ -modules

$$(8.1.5.1) \quad u_k : \mathcal{S}_+(k+1) \longrightarrow \mathcal{S}(k);$$

since $\mathcal{S}_+(k+1)$ and $\mathcal{S}(k+1)$ are canonically (TN)-isomorphic, they give a canonical correspondence between u_k and an injective homomorphism of $\mathcal{O}_X$ -modules (3.4.2):

$$(8.1.5.2) \quad \tilde{u}_k : \mathcal{O}_X(k+1) \longrightarrow \mathcal{O}_X(k).$$

Recall as well (3.2.6) that we have defined canonical homomorphisms

$$(8.1.5.3) \quad \lambda : \mathcal{O}_X(h) \otimes_{\mathcal{O}_X} \mathcal{O}_X(k) \longrightarrow \mathcal{O}_X(h+k)$$

and, since the diagram

$$\begin{array}{ccc} \mathcal{S}(h) \otimes_{\mathcal{S}} \mathcal{S}(k) \otimes_{\mathcal{S}} \mathcal{S}(l) & \longrightarrow & \mathcal{S}(h+k) \otimes_{\mathcal{S}} \mathcal{S}(l) \\ \downarrow & & \downarrow \\ \mathcal{S}(h) \otimes_{\mathcal{S}} \mathcal{S}(k+l) & \longrightarrow & \mathcal{S}(h+k+l) \end{array}$$

commutes, it follows from the functoriality of the λ (3.2.6) that the homomorphisms (8.1.5.3) define the structure of a *quasi-coherent graded $\mathcal{O}_X$ -algebra* on

$$(8.1.5.4) \quad \mathcal{S}_X = \bigoplus_{n \in \mathbb{Z}} \mathcal{O}_X(n).$$

Furthermore, the diagram

$$\begin{array}{ccc} \mathcal{S}(h) \otimes_{\mathcal{S}} \mathcal{S}(k+1) & \longrightarrow & \mathcal{S}(h+k+1) \\ \downarrow 1 \otimes u_k & & \downarrow u_{k+h} \\ \mathcal{S}(h) \otimes_{\mathcal{S}} \mathcal{S}(k) & \longrightarrow & \mathcal{S}(h+k) \end{array}$$

commutes; the functoriality of the λ then implies that we have a commutative diagram

$$(8.1.5.5) \quad \begin{array}{ccc} \mathcal{O}_X(h) \otimes_{\mathcal{O}_X} \mathcal{O}_X(k+1) & \xrightarrow{\lambda} & \mathcal{O}_X(h+k+1) \\ \downarrow 1 \otimes \tilde{u}_k & & \downarrow \tilde{u}_{k+h} \\ \mathcal{O}_X(h) \otimes_{\mathcal{O}_X} \mathcal{O}_X(k) & \xrightarrow{\lambda} & \mathcal{O}_X(h+k) \end{array}$$

where the horizontal arrows are the canonical homomorphisms. We can thus say that the $\tilde{u}_k$ define an *injective homomorphism* (of degree zero) of graded $\mathcal{S}_X$ -modules

$$(8.1.5.6) \quad \tilde{u} : \mathcal{S}_X(1) \longrightarrow \mathcal{S}_X.$$

(8.1.6). Keeping the notation from (8.1.5), we now note that, for $n \geq 0$, the composite homomorphism $\tilde{v}_n = \tilde{u}_{n-1} \circ \tilde{u}_{n-2} \circ \dots \circ \tilde{u}_0$ is an *injective homomorphism* $\mathcal{O}_X(n) \rightarrow \mathcal{O}_X$; we denote by $\mathcal{I}_{n,X}$ its image, which is thus a quasi-coherent ideal of $\mathcal{O}_X$, *isomorphic* to $\mathcal{O}_X(n)$. Furthermore, the diagram

$$\begin{array}{ccc} \mathcal{O}_X(m) \otimes_{\mathcal{O}_X} \mathcal{O}_X(n) & \xrightarrow{\lambda} & \mathcal{O}_X(m+n) \\ \downarrow \tilde{v}_m \otimes \tilde{v}_n & & \downarrow \tilde{v}_{m+n} \\ \mathcal{O}_X & \xrightarrow{\text{id}} & \mathcal{O}_X \end{array}$$

commutes for $m \geq 0, n \geq 0$. We thus deduce the following inclusions:

$$(8.1.6.1) \quad \mathcal{I}_{0,X} = \mathcal{O}_X, \quad \mathcal{I}_{n,X} \subset \mathcal{I}_{m,X} \quad \text{for } 0 \leq m \leq n;$$

$$(8.1.6.2) \quad \mathcal{I}_{m,X} \mathcal{I}_{n,X} \subset \mathcal{I}_{m+n,X} \quad \text{for } m \geq 0, n \geq 0.$$

[II:156]

Proposition (8.1.7). — *Let Y be a prescheme, $\mathcal{I}$ a quasi-coherent ideal of $\mathcal{O}_Y$, and $X = \text{Proj}(\bigoplus_{n \geq 0} \mathcal{I}^n)$ the Y -scheme given by blowing up $\mathcal{I}$. We then have, for all $n > 0$, a canonical isomorphism*

$$(8.1.7.1) \quad \mathcal{O}_X(n) \xrightarrow{\sim} \mathcal{I}^n \mathcal{O}_X = \mathcal{I}_{n,X}$$

(cf. (0, 4.3.5)), and thus that $\mathcal{I}^n \mathcal{O}_X$ is a very-ample invertible $\mathcal{O}_X$ -module if $n > 0$.

Proof. The last claim is immediate, since $\mathcal{O}_X(1)$ is invertible (3.2.5) and very ample for Y by definition ((4.4.3) and (4.4.9)). Also by definition, the image of v_n is exactly $\mathcal{I}^n \mathcal{S}$, and (8.1.7.1) then follows from the exactness of the functor $\mathcal{M}$ (3.2.4) and from Equation (3.2.4.1). $\square$

Corollary (8.1.8). — Under the hypotheses of (8.1.7), if $f : X \rightarrow Y$ is the structure morphism, and Y' the closed subprescheme of Y defined by $\mathcal{I}$, then the closed subprescheme $X' = f^{-1}(Y')$ of X is defined by $\mathcal{I}\mathcal{O}_X$ (which is canonically isomorphic to $\mathcal{O}_X(1)$), from which we obtain a canonical short exact sequence

$$(8.1.8.1) \quad 0 \longrightarrow \mathcal{O}_X(1) \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{O}_{X'} \longrightarrow 0.$$

Proof. This follows from (8.1.7.1) and from (I, 4.4.5). $\square$

(8.1.9). Under the hypotheses of (8.1.7), we can be more precise about the structure of the $\mathcal{I}_{n,X}$. Note that the homomorphism

$$\tilde{u}_{-1} : \mathcal{O}_X \longrightarrow \mathcal{O}_X(-1)$$

canonically corresponds to a section s of $\mathcal{O}_X(-1)$ over X , which we call the *canonical section* (relative to $\mathcal{I}$) (0, 5.1.1). In the diagram in (8.1.5.5), the horizontal arrows are isomorphisms (3.2.7); by replacing h with k , and k with -1 in this diagram, we obtain that $\tilde{u}_k = 1_k \otimes \tilde{u}_{-1}$ (where 1_k denotes the identity on $\mathcal{O}_X(k)$), or, equivalently, that the homomorphism $\tilde{u}_k$ is given exactly by *tensoring with the canonical section s* (for all $k \in \mathbf{Z}$). The homomorphism $\tilde{u}$ (8.1.5.6) can then be understood in the same way.

Thus, for all $n \geq 0$, the homomorphism $\tilde{v}_n : \mathcal{O}_X(n) \rightarrow \mathcal{O}_X$ is given exactly by tensoring with $s^{\otimes n}$; we thus deduce:

Corollary (8.1.10). — With the notation of (8.1.8), the underlying space of X' is the set of $x \in X$ such that $s(x) = 0$, where s denotes the canonical section of $\mathcal{O}_X(-1)$.

Proof. Indeed, if c_x is a generator of the fibre $(\mathcal{O}_X(1))_x$ at a point x , then $s_x \otimes c_x$ is canonically identified with a generator of the fibre of $\mathcal{I}_{1,X}$ at the point x , and is thus invertible if and only if $s_x \notin \mathfrak{m}_x(\mathcal{O}_X(-1))_x$, or, equivalently, if and only if $s(x) \neq 0$. $\square$

Proposition. — Let Y be an integral prescheme, $\mathcal{I}$ a quasi-coherent fractional ideal of $\mathcal{B}(Y)$, and X the Y -scheme given by blowing up $\mathcal{I}$. Then $\mathcal{I}\mathcal{O}_X$ is an invertible $\mathcal{O}_X$ -module that is very ample for Y .

Proof. Since the question is local on Y (4.4.5), we can reduce to the case where $Y = \text{Spec}(A)$, with A an integral domain with field of fractions K , and $\mathcal{I} = \tilde{\mathcal{I}}$, with $\mathcal{I}$ some fractional ideal of K ; there then exists an element $a \neq 0$ of A such that $a\mathcal{I} \subset A$. Let $S = \bigoplus_{n \geq 0} \mathcal{I}^n$; the map $x \mapsto ax$ is an A -isomorphism from $\mathcal{I}^{n+1} = (S(1))_n$ to $a\mathcal{I}^{n+1} = a\mathcal{I}S_n \subset \mathcal{I}^n = S_n$, [II:157] and thus defines a (TN)-isomorphism of degree zero of graded S -modules $S_+(1) \rightarrow a\mathcal{I}S$. On the other hand, $x \mapsto a^{-1}x$ is an isomorphism of degree zero of graded S -modules $a\mathcal{I}S \xrightarrow{\sim} \mathcal{I}S$. We thus obtain, by composition (3.2.4), an isomorphism of $\mathcal{O}_X$ -modules $\mathcal{O}_X(1) \xrightarrow{\sim} \mathcal{I}\mathcal{O}_X$, and, since S is generated by $S_1 = \mathcal{I}$, $\mathcal{O}_X(1)$ is invertible (3.2.5) and very ample ((4.4.3) and (4.4.9)), whence our claim. $\square$

8.2. Preliminary results on the localisation of graded rings

(8.2.1). Let S be a graded ring, which for the moment we do not assume to be concentrated in nonnegative degrees. We define

$$(8.2.1.1) \quad S^{\geq} = \bigoplus_{n \geq 0} S_n, \quad S^{\leq} = \bigoplus_{n \leq 0} S_n$$

which are both graded subrings of S , in only positive and negative degrees (respectively). If f is a homogeneous element of degree d (positive or negative) in S , then the localization $S_f = S'$ is again endowed with the structure of a graded ring, by taking S'_n ($n \in \mathbf{Z}$) to be the set of the x/f^k for $x \in S_{n+kd}$ ($k \geq 0$); we define $S_{(f)} = S'_0$, and will write $S_f^{\geq}$ and $S_f^{\leq}$ for $S'^{\geq}$ and $S'^{\leq}$ (respectively). If $d > 0$, then

$$(8.2.1.2) \quad (S^{\geq})_f = S_f$$

since, if $x \in S_{n+kd}$ with $n + kd < 0$, then we can write $x/f^k = x f^h / f^{h+k}$, and we also have that $n + (h+k)d > 0$ for h sufficiently large and > 0 . We thus conclude, by definition, that

$$(8.2.1.3) \quad (S^{\geq})_{(f)} = (S_f^{\geq})_0 = S_{(f)}.$$

If M is a graded S -module, then we similarly define

$$(8.2.1.4) \quad M^{\geq} = \bigoplus_{n \geq 0} M_n, \quad M^{\leq} = \bigoplus_{n \leq 0} M_n$$

which are (respectively) a graded $S^{\geq}$ -module and a graded $S^{\leq}$ -module, and their intersection is the S_0 module M_0 . If $f \in S_d$, then we define M_f to be the graded S_f -module whose elements of degree n are the z/f^k for $z \in M_{n+kd}$ ($k \geq 0$); we denote by $M_{(f)}$ the set of elements of degree zero of M_f , and this is an $S_{(f)}$ -module, and we will write $M_f^{\geq}$ and $M_f^{\leq}$ to mean $(M_f)^{\geq}$ and $(M_f)^{\leq}$ (respectively). If $d > 0$, then we see, as above, that

$$(8.2.1.5) \quad (M^{\geq})_f = M_f$$

and

$$(8.2.1.6) \quad (M^{\geq})_{(f)} = (M_f^{\geq})_0 = M_{(f)}.$$

(8.2.2). Let z be an indeterminate, which we will call the *homogenisation variable*. If S is a graded ring (with arbitrary integer degrees), then the polynomial algebra¹

$$(8.2.2.1) \quad \hat{S} = S[z]$$

[II:158] is a graded S -algebra, where we define the degree of fz^n ($n \geq 0$), with f homogeneous, as

$$(8.2.2.2) \quad \deg(fz^n) = n + \deg f.$$

Lemma (8.2.3). — (i) *There are canonical isomorphisms of (non-graded) rings*

$$(8.2.3.1) \quad \hat{S}_{(z)} \xrightarrow{\sim} \hat{S}/(z-1)\hat{S} \xrightarrow{\sim} S.$$

(ii) *There is a canonical isomorphism of (non-graded) rings*

$$(8.2.3.2) \quad \hat{S}_{(f)} \xrightarrow{\sim} S_f^{\leq}$$

for all $f \in S_d$ with $d > 0$.

Proof. The first of the isomorphisms in (8.2.3.1) was defined in (2.2.5), and the second is trivial; the isomorphism $\hat{S}_{(z)} \xrightarrow{\sim} S$ thus defined thus gives a correspondence between xz^n/z^{n+k} (where $\deg(x) = k$ for $k \geq -n$) and the element x . The homomorphism (8.2.3.2) gives a correspondence between xz^n/f^k (where $\deg(x) = kd - n$) and the element x/f^k of degree $-n$ in $S_f^{\leq}$, and it is again clear that this does indeed give an isomorphism. $\square$

(8.2.4). Let M be a graded S -module. It is clear that the S -module

$$(8.2.4.1) \quad \hat{M} = M \otimes_S \hat{S} = M \otimes_S S[z]$$

is the direct sum of the S -modules $M \otimes_S Sz^n$, and thus of the abelian groups $M_k \otimes_S Sz^n$ ($k \in \mathbf{Z}$, $n \geq 0$); we define on $\hat{M}$ the structure of a graded $\hat{S}$ -module by setting

$$(8.2.4.2) \quad \deg(x \otimes z^n) = n + \deg x$$

for all homogeneous x in M . We leave it to the reader to prove the analogue of (8.2.3):

Lemma (8.2.5). — (i) *There is a canonical di-isomorphism of (non-graded) modules*

$$(8.2.5.1) \quad \hat{M}_{(z)} \xrightarrow{\sim} M.$$

(ii) *For all $f \in S_d$ ($d > 0$), there is a di-isomorphism of (non-graded) modules*

$$(8.2.5.2) \quad \hat{M}_{(f)} \xrightarrow{\sim} M_f^{\leq}.$$

¹This should not be confused with the use of the notation $\hat{S}$ for the separated completion of a ring.

(8.2.6). Let S be a *positively-graded* ring, and consider the decreasing sequence of graded ideals of S

$$(8.2.6.1) \quad S_{[n]} = \bigoplus_{m \geq n} S_m \quad (n \geq 0)$$

(so, in particular, we have $S_{[0]} = S$ and $S_{[1]} = S_+$). Since it is evident that $S_{[m]}S_{[n]} \subset S_{[m+n]}$, we can define a *graded ring* $S^\natural$ by setting

$$(8.2.6.2) \quad S^\natural = \bigoplus_{n \geq 0} S_n^\natural \quad \text{with} \quad S_n^\natural = S_{[n]}.$$

$S_0^\natural$ is then the ring S considered as a *non-graded* ring, and $S^\natural$ is thus an $S_0^\natural$ -algebra. For every homogeneous element $f \in S_d$ ($d > 0$), we denote by $f^\natural$ the element f considered as belonging to $S_{[d]} = S_d^\natural$. With this notation:

[II:159]

Lemma (8.2.7). — *Let S be a positively-graded ring, and f a homogeneous element of S_d ($d > 0$). There are canonical ring isomorphisms*

$$(8.2.7.1) \quad S_f \xrightarrow{\sim} \bigoplus_{n \in \mathbf{Z}} S(n)_{(f)}$$

$$(8.2.7.2) \quad (S_f^\geq)_{f/1} \xrightarrow{\sim} S_f$$

$$(8.2.7.3) \quad S_{(f^\natural)}^\natural \xrightarrow{\sim} S_f^\geq$$

where the first two are isomorphisms of graded rings.

Proof. It is immediate, by definition, that we have $(S_f)_n = (S(n)_f)_0$, whence the isomorphism in (8.2.7.1), which is exactly the identity. Next, since $f/1$ is invertible in S_f , there is a canonical isomorphism $S_f \xrightarrow{\sim} (S_f^\geq)_{f/1} = (S_f)_{f/1}$, by (8.2.1.2) applied to S_f ; the inverse isomorphism is, by definition, the isomorphism in (8.2.7.2). Finally, if $x = \sum_{m \geq n} y_m$ is an element of $S_{[n]}$ with $n = kd$, then the element $x/(f^\natural)^k$ corresponds to the element $\sum y_m/f^k$ of $S_f^\geq$, and we can quickly verify that this defines an isomorphism (8.2.7.3). $\square$

(8.2.8). If M is a graded S -module, then we similarly define, for all $n \in \mathbf{Z}$,

$$(8.2.8.1) \quad M_{[n]} = \bigoplus_{m \geq n} M_m$$

and, since $S_{[m]}M_{[n]} \subset M_{[m+n]}$ ($m \geq 0$), we can define a graded $S^\natural$ -module $M^\natural$ by setting

$$(8.2.8.2) \quad M^\natural = \bigoplus_{n \in \mathbf{Z}} M_n^\natural \quad \text{with} \quad M_n^\natural = M_{[n]}.$$

We leave to the reader the proof of:

Lemma (8.2.9). — *With the notation of (8.2.7) and (8.2.8), there are canonical di-isomorphisms of modules*

$$(8.2.9.1) \quad M_f \xrightarrow{\sim} \bigoplus_{n \in \mathbf{Z}} M(n)_{(f)}$$

$$(8.2.9.2) \quad (M_f^\geq)_{f/1} \xrightarrow{\sim} M_f$$

$$(8.2.9.3) \quad M_{(f^\natural)}^\natural \xrightarrow{\sim} M_f^\geq$$

where the first two are di-isomorphisms of graded modules.

Lemma (8.2.10). — *Let S be a positively-graded ring.*

- (i) *For $S^\natural$ to be an $S_0^\natural$ -algebra of finite type (resp. a Noetherian ring), it is necessary and sufficient for S to be an S_0 -algebra of finite type (resp. a Noetherian ring).*
- (ii) *For $S_{n+1}^\natural = S_1^\natural S_n^\natural$ ($n \geq n_0$), it is necessary and sufficient for $S_{n+1} = S_1 S_n$ ($n \geq n_0$).*
- (iii) *For $S_n^\natural = (S_1^\natural)^n$ ($n \geq n_0$), it is necessary and sufficient for $S_n = S_1^n$ ($n \geq n_0$).*

- (iv) If (f_α) is a set of homogeneous elements of S_+ such that S_+ is the radical in S of the ideal of S generated by the f_α , then $S_+^\natural$ is the radical in $S^\natural$ of the ideal of $S^\natural$ generated by the $f_\alpha^\natural$.

Proof.

- (i) If $S^\natural$ is an $S_0^\natural$ -algebra of finite type, then $S_+ = S_1^\natural$ is a module of finite type over $S = S_0^\natural$, by (2.1.6, i), and so S is an S_0 -algebra of finite type (2.1.4); if $S^\natural$ is a Noetherian ring, then so too is $S_0^\natural = S$ (2.1.5). Conversely, if S is an S_0 -algebra [II:160] of finite type, then we know (2.1.6, ii) that there exist $h > 0$ and $m_0 > 0$ such that $S_{n+h} = S_h S_n$ for $n \geq m_0$; we can clearly assume that $m_0 \geq h$. Furthermore, the S_m are S_0 -modules of finite type (2.1.6, i). So, if $n \geq m_0 + h$, then $S_n^\natural = S_h S_{n-h}^\natural = S_h^\natural S_{n-h}^\natural$; and if $m < m_0 + h$ then, letting $E = S_{m_0} + \dots + S_{m_0+h-1}$, we have that

$$S_m^\natural = S_m + \dots + S_{m_0+h-1} + S_h E + S_h^2 E + \dots$$

For $1 \leq m \leq m_0$, let G_m be the union of the finite systems of generators of the S_0 -modules S_i for $m \leq i \leq m_0 + h - 1$, considered as a subset of $S_{[m]}$. For $m_0 + 1 \leq m \leq m_0 + h - 1$, let G_m be the union of the finite system of generators of the S_0 -modules S_i for $m \leq i \leq m_0 + h - 1$ and of $S_h E$, considered as a subset of $S_{[m]}$. It is clear that $S_m^\natural = S_0^\natural G_m$ for $1 \leq m \leq m_0 + h - 1$, and thus the union G of the G_m for $1 \leq m \leq m_0 + h - 1$ is a system of generators of the $S_0^\natural$ -algebra $S^\natural$. We thus conclude that, if $S = S_0^\natural$ is a Noetherian ring, then so too is $S^\natural$.

- (ii) It is clear that, if $S_{n+1} = S_1 S_n$ for $n \geq n_0$, then $S_{n+1}^\natural = S_1 S_n^\natural$, and *a fortiori* $S_{n+1}^\natural = S_1^\natural S_n^\natural$ for $n \geq n_0$. Conversely, this last equality can be written as

$$S_{n+1} + S_{n+2} + \dots = (S_1 + S_2 + \dots)(S_n + S_{n+1} + \dots)$$

and comparing terms of degree $n + 1$ (in S) on both sides gives that $S_{n+1} = S_1 S_n$.

- (iii) If $S_n = S_1^n$ for $n \geq n_0$, then $S_n^\natural = S_1^n + S_1^{n+1} + \dots$; since $S_1^\natural$ contains $S_1 + S_1^2 + \dots$, we have that $S_n^\natural \subset S_1^{\natural n}$, and thus $S_n^\natural = S_1^{\natural n}$ for $n \geq n_0$. Conversely, the only terms of $S_1^{\natural n} = (S_1 + S_2 + \dots)^n$ that are of degree n in S are those of S_1^n ; the equality $S_n^\natural = S_1^{\natural n}$ thus implies that $S_n = S_1^n$.
- (iv) It suffices to show that, if an element $g \in S_{k+h}$ is considered as an element of $S_k^\natural$ ($k > 0$, $h \geq 0$), then there exists an integer $n > 0$ such that g^n is a linear combination (in $S_{kn}^\natural$) of the $f_\alpha^\natural$ with coefficients in $S^\natural$. By hypothesis, there exists an integer m_0 such that, for $m \geq m_0$, we have, in S , that $g^m = \sum_\alpha c_{\alpha m} f_\alpha$, where the indices α here are independent of m ; furthermore, we can clearly assume that the $c_{\alpha m}$ are homogeneous, with

$$\deg(c_{\alpha m}) = m(k + h) - \deg f_\alpha$$

in S . So take m_0 sufficiently large to ensure that $km_0 > \deg f_\alpha$ for all the f_α that appear in g^{m_0} ; for all α , let $c'_{\alpha m}$ be the element $c_{\alpha m}$ considered as having degree $km - \deg f_\alpha$ in $S^\natural$; we then have, in $S^\natural$, that $g^m = \sum_\alpha c'_{\alpha m} f_\alpha^\natural$, which finishes the proof. $\square$

(8.2.11). Consider the graded S_0 -algebra

$$(8.2.11.1) \quad S^\natural \otimes_S S_0 = S^\natural / S_+ S^\natural = \bigoplus_{n \geq 0} S_{[n]} / S_+ S_{[n]}.$$

Since S_n is a quotient S_0 -module of $S_{[n]} / S_+ S_{[n]}$, there is a canonical homomorphism of graded S_0 -algebras

$$(8.2.11.2) \quad S^\natural \otimes_S S_0 \longrightarrow S$$

which is clearly *surjective*, and thus corresponds (2.9.2) to a canonical *closed immersion*

$$(8.2.11.3) \quad \text{Proj}(S) \longrightarrow \text{Proj}(S^\natural \otimes_S S_0).$$

Proposition (8.2.12). — *The canonical morphism (8.2.11.3) is bijective. For the homomorphism (8.2.11.2) to be (TN)-bijective, it is necessary and sufficient for there to exist some n_0 such that $S_{n+1} = S_1 S_n$ for $n \geq n_0$. If this latter condition is satisfied, then (8.2.11.3) is an isomorphism; the converse is true whenever S is Noetherian.*

Proof. To prove the first claim, it suffices (2.8.3) to show that the kernel $\mathfrak{I}$ of the homomorphism (8.2.11.2) consists of nilpotent elements. But if $f \in S_{[n]}$ is an element whose class modulo $S_+ S_{[n]}$ belongs to this kernel, then this implies that $f \in S_{[n+1]}$; then f^{n+1} , considered as an element of $S_{[n(n+1)]}$, is also an element of $S_+ S_{[n(n+1)]}$, since it can be written as $f \cdot f^n$; so the class of f^{n+1} modulo $S_+ S_{[n(n+1)]}$ is zero, which proves our claim. Since the hypothesis that $S_{n+1} = S_1 S_n$ for $n \geq n_0$ is equivalent to $S_{n+1}^\natural = S_1^\natural S_n^\natural$ for $n \geq n_0$ (8.2.10, ii), this hypothesis is equivalent, by definition, to the fact that (8.2.11.2) is (TN)-injective, and thus (TN)-bijective, and so (8.2.11.3) is an isomorphism, by (2.9.1). Conversely, if (8.2.11.3) is an isomorphism, then the sheaf $\tilde{\mathfrak{I}}$ on $\text{Proj}(S^\natural \otimes_S S_0)$ is zero (2.9.2, i); since $S^\natural \otimes_S S_0$ is Noetherian, as a quotient of $S^\natural$ (8.2.10, i), we conclude from (2.7.3) that $\mathfrak{I}$ satisfies condition (TN), and so $S_{n+1}^\natural = S_1^\natural S_n^\natural$ for $n \geq n_0$, and this finishes the proof, by (8.2.10, ii). $\square$

(8.2.13). Consider now the canonical injections $(S_+)^n \rightarrow S_{[n]}$, which define an injective homomorphism of degree zero of graded rings

$$(8.2.13.1) \quad \bigoplus_{n \geq 0} (S_+)^n \longrightarrow S^\natural.$$

Proposition (8.2.14). — *For the homomorphism (8.2.13.1) to be a (TN)-isomorphism, it is necessary and sufficient for there to exist some n_0 such that $S_n = S_1^n$ for all $n \geq n_0$. Whenever this is the case, the morphism corresponding to (8.2.13.1) is everywhere defined and is an isomorphism*

$$\text{Proj}(S^\natural) \xrightarrow{\sim} \text{Proj}\left(\bigoplus_{n \geq 0} (S_+)^n\right);$$

the converse is true whenever S is Noetherian.

Proof. The first two claims are evident, given (8.2.10, iii) and (2.9.1). The third will follow from (8.2.10, i and iii) and the following lemma:

Lemma (8.2.14.1). — *Let T be a positively-graded ring that is also a T_0 -algebra of finite type. If the morphism corresponding to the injective homomorphism $\bigoplus_{n \geq 0} T_1^n \rightarrow T$ is everywhere defined and also an isomorphism $\text{Proj}(T) \rightarrow \text{Proj}(\bigoplus_{n \geq 0} T_1^n)$, then there exists some n_0 such that $T_n = T_1^n$ for $n \geq n_0$.*

Let g_i ($1 \leq i \leq r$) be generators of the T_0 -module T_1 . The hypothesis implies first of all that the $D_+(g_i)$ cover $\text{Proj}(T)$ (2.8.1). Let $(h_j)_{1 \leq j \leq s}$ be a system of homogeneous elements of T_+ , with $\deg(h_j) = n_j$, that form, with the g_i , a system of generators of the ideal T_+ , or, equivalently (2.1.3), a system of generators of T as a T_0 -algebra; if we set $T' = \bigoplus_{n \geq 0} T_1^n$, then the element $h_j / g_i^{n_j}$ of the ring $T_{(g_i)}$ must, by hypothesis, belong to the subring $T'_{(g_i)}$, and so there exists some integer k such that $T_1^k h_j \subset T_1^{k+n_j}$ for all j . We thus conclude, by induction on the exponent r , that $T_1^k h_j' \subset T'$ for all $r \geq 1$, and, by definition of the h_j , we thus have that $T_1^k T \subset T'$. Also, there exists, for all j , an integer m_j such that $h_j^{m_j}$ belongs to the ideal of T generated by the g_i (2.3.14), so $h_j^{m_j} \in T_1 T$, and $h_j^{m_j k} \in T_1^k T \subset T'$. There is thus an integer $m_0 \geq k$ such that $h_j^m \in T_1^{m m_j}$ for $m \geq m_0$. So, if q is the largest of the integers n_j , then $n_0 = q m_0 + k$ is the required number. Indeed, an element of T_n , for $n \geq n_0$, is the sum of monomials belonging to $T_1^\alpha u$, where u is a product of powers of the h_j ; if $\alpha \geq k$, then it follows from the above that $T_1^\alpha u \subset T_1^n$; in the other case, one of the exponents of the h_j is $\geq m_0$, so $u \in T_1^\beta v$, where $\beta \geq k$ and v is again a product of powers of the h_j ; we can then reduce to the previous case, and so we conclude that $T_1^\alpha u \subset T_1^n$ in all cases. $\square$

Remark (8.2.15). — The condition $S_n = S_1^n$ for $n \geq n_0$ clearly implies that $S_{n+1} = S_1 S_n$ for $n \geq n_0$, but the converse is not necessarily true, even if we assume that S is Noetherian. For example, let K be a field, $A = K[x]$, and $B = K[y]/y^2 K[y]$, where x and y are indeterminates, with x taken to have degree 1 and y to have degree 2, and let $S = A \otimes_K B$, so that S is a graded algebra over K that has

a basis given by the elements $1, \mathbf{x}^n$ ($n \geq 1$), and $\mathbf{x}^n \mathbf{y}$ ($n \geq 0$). It is immediate that $S_{n+1} = S_1 S_n$ for $n \geq 2$, but $S_1^n = K\mathbf{x}^n$ while $S_n = K\mathbf{x}^n + K\mathbf{x}^{n-2}\mathbf{y}$ for $n \geq 2$.

8.3. Projecting cones

(8.3.1). Let Y be a prescheme; in all of this section, we will consider only Y -preschemes and Y -morphisms. Let $\mathcal{S}$ be a quasi-coherent positively-graded $\mathcal{O}_Y$ -algebra; we further assume that $\mathcal{S}_0 = \mathcal{O}_Y$. Following the notation introduced in (8.2.2), we let

$$(8.3.1.1) \quad \widehat{\mathcal{S}} = \mathcal{S}[\mathbf{z}] = \mathcal{S} \otimes_{\mathcal{O}_Y} \mathcal{O}_Y[\mathbf{z}]$$

which we consider as a positively-graded $\mathcal{O}_Y$ -algebra by defining the degrees as in (8.2.2.2), so that, for every affine open subset U of Y , we have

$$\Gamma(U, \widehat{\mathcal{S}}) = (\Gamma(U, \mathcal{S}))[\mathbf{z}].$$

In what follows, we write

$$(8.3.1.2) \quad X = \text{Proj}(\mathcal{S}), \quad C = \text{Spec}(\mathcal{S}), \quad \widehat{C} = \text{Proj}(\widehat{\mathcal{S}})$$

(where, in the definition of C , we consider $\mathcal{S}$ as a non-graded $\mathcal{O}_Y$ -algebra), and we say that C (resp. $\widehat{C}$) is the *affine cone* (resp. *projective cone*) defined by $\mathcal{S}$; we will sometimes say “cone” instead of “affine cone”. By an abuse of language, we also say that C (resp. $\widehat{C}$) is the *affine projecting cone* of X (resp. the *projective projecting cone* of X), with the implicit understanding that the prescheme X is given in the form $\text{Proj}(\mathcal{S})$; finally, we say that $\widehat{C}$ is the *projective closure* of C (with the data of $\mathcal{S}$ being implicit in the structure of C).

Proposition (8.3.2). — *There exist canonical Y -morphisms*

$$(8.3.2.1) \quad Y \xrightarrow{\varepsilon} C \xrightarrow{i} \widehat{C}$$

$$(8.3.2.2) \quad X \xrightarrow{j} \widehat{C}$$

such that ε and j are closed immersions, and i is an affine morphism, which is a dominant open immersion, for which

$$(8.3.2.3) \quad i(C) = \widehat{C} - j(X);$$

furthermore, $\widehat{C}$ is the smallest closed subprescheme of $\widehat{C}$ containing $i(C)$.

Proof. To define i , consider the open subset of $\widehat{C}$ given by

$$(8.3.2.4) \quad \widehat{C}_{\mathbf{z}} = \text{Spec}(\widehat{\mathcal{S}}/(\mathbf{z} - 1)\widehat{\mathcal{S}})$$

(3.1.4), where $\mathbf{z}$ is canonically identified with a section of $\widehat{\mathcal{S}}$ over Y . The isomorphism $i : C \xrightarrow{\sim} \widehat{C}_{\mathbf{z}}$ then corresponds to the canonical isomorphism (8.2.3.1)

$$\widehat{\mathcal{S}}/(\mathbf{z} - 1)\widehat{\mathcal{S}} \xrightarrow{\sim} \mathcal{S}.$$

The morphism ε corresponds to the augmentation homomorphism $\mathcal{S} \rightarrow \mathcal{S}_0 = \mathcal{O}_Y$, which has kernel $\mathcal{S}_+$ (1.2.7), and, since the latter is surjective, ε is a closed immersion (1.4.10). Finally, j corresponds (3.5.1) to the surjective homomorphism of degree zero $\widehat{\mathcal{S}} \rightarrow \mathcal{S}$, which restricts to the identity on $\mathcal{S}$ and is zero on $\mathbf{z}\widehat{\mathcal{S}}$, which is its kernel; j is everywhere defined, and is a closed immersion, by (3.6.2).

To prove the other claims of (8.3.2), we can clearly restrict to the case where $Y = \text{Spec}(A)$ is affine, and $\mathcal{S} = \widehat{S}$, with S a graded A -algebra, whence $\widehat{\mathcal{S}} = (\widehat{S})^\sim$; the homogeneous elements f of S_+ can then be identified with sections of $\widehat{\mathcal{S}}$ over Y , and the open subset of $\widehat{C}$, denoted $D_+(f)$ in (2.3.3), can then be written as $\widehat{C}_f$ (3.1.4); similarly, the open subset of C denoted $D(f)$ in (I, 1.1.1) can be written as C_f (0, 5.5.2). With this in mind, it follows from (2.3.14) and from the definition of $\widehat{S}$ that, in this case, the open subsets $\widehat{C}_{\mathbf{z}} = i(C)$ and $\widehat{C}_f$ (with f homogeneous in S_+) form a cover of $\widehat{C}$. Furthermore, with this notation,

$$(8.3.2.5) \quad i^{-1}(\widehat{C}_f) = C_f;$$

indeed, $\widehat{C}_f \cap i(C) = \widehat{C}_f \cap \widehat{C}_z = \widehat{C}_{fz} = \text{Spec}(\widehat{S}_{(fz)})$. But, if $d = \deg(f)$, then $\widehat{S}_{(fz)}$ is canonically isomorphic to $(\widehat{S}_{(z)})_{f/z^d}$ (2.2.2), and it follows from the definition of the isomorphism in (8.2.3.1) that the image of $(\widehat{S}_{(z)})_{f/z^d}$ under the corresponding isomorphism of rings of fractions is exactly S_f . Since $C_f = \text{Spec}(S_f)$, this proves (8.3.2.5) and shows, at the same time, that the morphism i is affine; furthermore, the restriction of i to C_f , considered as a morphism to $\widehat{C}_f$, corresponds (I, 1.7.3) to the canonical homomorphism $\widehat{S}_{(f)} \rightarrow \widehat{S}_{(fz)}$, and, by the above and (8.2.3.2), we can claim the following result:

(8.3.2.6). If $Y = \text{Spec}(A)$ is affine, and $\mathcal{S} = \widetilde{S}$, then, for every homogeneous f in S_+ , $\widehat{C}_f$ is canonically identified with $\text{Spec}(S_f^{\leq})$, and the morphism $C_f \rightarrow \widehat{C}_f$ given by restricting i then corresponds to the canonical injection $S_f^{\leq} \rightarrow S_f$.

Now note that (for Y affine) the complement of $\widehat{C}_z$ in $\widehat{C} = \text{Proj}(\widehat{S})$ [II:164] is, by definition, the set of graded prime ideals of $\widehat{S}$ containing z , which is exactly $j(X)$, by definition of j , which proves (8.3.2.3).

Finally, to prove the last claim of (8.3.2), we can assume that Y is affine. With the above notation, note that, in the ring $\widehat{S}$, z is not a zero divisor; since the closure of $i(C)$ is $\widehat{C}$, it suffices to prove the following lemma:

Lemma (8.3.2.7). — Let T be a positively-graded ring, $Z = \text{Proj}(T)$, and g a homogeneous element of T of degree $d > 0$. If g is not a zero divisor in T , then Z is the smallest closed subprescheme of Z that contains $Z_g = D_+(g)$.

By (I, 4.1.9), the question is local on Z ; for every homogeneous element $h \in T_e$ ($e > 0$), it thus suffices to prove that Z_h is the smallest closed subprescheme of Z_h that contains Z_{gh} ; it follows from the definitions and from (I, 4.3.2) that this condition is equivalent to asking for the canonical homomorphism $T_{(h)} \rightarrow T_{(gh)}$ to be *injective*. But this homomorphism can be identified with the canonical homomorphism $T_{(h)} \rightarrow (T_{(h)})_{g^e/h^d}$ (2.2.3). But since g^e is not a zero divisor in T , g^e/h^d is not a zero divisor in T_h (nor *a fortiori* in $T_{(h)}$), since the fact that $(g^e/h^d)(t/h^m) = 0$ (for $t \in T$ and $m > 0$) implies the existence of some $n > 0$ such that $h^n g^e t = 0$, whence $h^n t = 0$, and thus $t/h^m = 0$ in T_h . This thus finishes the proof (0, 1.2.2). $\square$

(8.3.3). We will often identify the affine cone C with the subprescheme induced by the projective cone $\widehat{C}$ on the open subset $i(C)$ by means of the open immersion i . The closed subprescheme of C associated to the closed immersion ε is called the *vertex prescheme* of C ; we also say that ε , which is a Y -section of C , is the *vertex section*, or the *zero section*, of C ; we can identify Y with the vertex prescheme of C by means of ε . Also, $i \circ \varepsilon$ is a Y -section of $\widehat{C}$, and thus also a closed immersion (I, 5.4.6), corresponding to the canonical surjective homomorphism of degree zero $\widehat{\mathcal{S}} = \mathcal{S}[\mathbf{z}] \rightarrow \mathcal{O}_Y[\mathbf{z}]$ (3.1.7), whose kernel is $\mathcal{S}_+[\mathbf{z}] = \mathcal{S}_+ \widehat{\mathcal{S}}$; the subprescheme of $\widehat{C}$ associated to this closed immersion is also called the *vertex prescheme* of $\widehat{C}$, and $i \circ \varepsilon$ the *vertex section* of $\widehat{C}$; it can be identified with Y by means of $i \circ \varepsilon$. Finally, the closed subprescheme of $\widehat{C}$ associated to j is called the *part at infinity* of $\widehat{C}$, and can be identified with X by means of j .

(8.3.4). The subpreschemes of C (resp. $\widehat{C}$) induced on the *open* subsets

$$(8.3.4.1) \quad E = C - \varepsilon(Y), \quad \widehat{E} = \widehat{C} - i(\varepsilon(Y))$$

are called (by an abuse of language) the *punctured affine cone* and the *punctured projective cone* (respectively) defined by $\mathcal{S}$; we note that, despite this nomenclature, E is *not necessarily affine* over Y , nor $\widehat{E}$ projective over Y (8.4.3). When we identify C with $i(C)$, we thus have the underlying spaces

$$(8.3.4.2) \quad C \cup \widehat{E} = \widehat{C}, \quad C \cap \widehat{E} = E$$

so that $\widehat{C}$ can be considered as being obtained by *gluing* the open subpreschemes C and $\widehat{E}$; furthermore, by (8.3.2.3),

$$(8.3.4.3) \quad E = \widehat{E} - j(X).$$

If $Y = \operatorname{Spec}(A)$ is affine, then, with the notation of (8.3.2),

$$(8.3.4.4) \quad E = \bigcup C_f, \quad \widehat{E} = \bigcup \widehat{C}_f, \quad C_f = C \cap \widehat{C}_f$$

where f runs over the set of homogeneous elements of S_+ (or only a subset M of this set, with M generating an ideal of S_+ whose radical in S_+ is S_+ itself, or, equivalently, such that the X_f for $f \in M$ cover X (2.3.14)). The gluing of C and $\widehat{C}_f$ along C_f is thus determined by the injection morphisms $C_f \rightarrow C$ and $C_f \rightarrow \widehat{C}_f$, which, as we have seen (8.3.2.6), correspond (respectively) to the canonical homomorphisms $S \rightarrow S_f$ and $S_f^{\leq} \rightarrow S_f$.

Proposition (8.3.5). — *With the notation of (8.3.1) and (8.3.4), the morphism associated (3.5.1) to the canonical injection $\phi : \mathcal{S} \rightarrow \widehat{\mathcal{S}} = \mathcal{S}[\mathbf{z}]$ is a surjective affine morphism (called the canonical retraction)*

$$(8.3.5.1) \quad p : \widehat{E} \longrightarrow X$$

such that

$$(8.3.5.2) \quad p \circ j = 1_X.$$

Proof. To prove the proposition, we can restrict to the case where Y is affine. Taking into account the expression in (8.3.4.4) for $\widehat{E}$, the fact that the domain of definition $G(\phi)$ of p is equal to $\widehat{E}$ will follow from the first of the following claims:

(8.3.5.3). If $Y = \operatorname{Spec}(A)$ is affine, and $\mathcal{S} = \widetilde{S}$, then, for all homogeneous $f \in S_+$,

$$(8.3.5.4) \quad p^{-1}(X_f) = \widehat{C}_f$$

and the restriction of p to $\widehat{C}_f = \operatorname{Spec}(S_f^{\leq})$, considered as a morphism from $\widehat{C}_f$ to X_f , corresponds to the canonical injection $S_{(f)} \rightarrow S_f^{\leq}$. If, further, $f \in S_1$, then $\widehat{C}_f$ is isomorphic to $X_f \otimes_{\mathbf{Z}} \mathbf{Z}[T]$ (where T is an indeterminate).

Indeed, Equation (8.3.5.4) is exactly a particular case of (2.8.1.1), and the second claim is exactly the definition of $\operatorname{Proj}(\phi)$ whenever Y is affine (2.8.1). Equation (8.3.5.2) and the surjectivity of p then follow from the fact that the composition $\mathcal{S} \rightarrow \widehat{\mathcal{S}} \rightarrow \mathcal{S}$ of the canonical homomorphisms is the identity on $\mathcal{S}$. Finally, the last claim of (8.3.5.3) follows from the fact that $S_f^{\leq}$ is isomorphic to $S_{(f)}[T]$ whenever $f \in S_1$ (2.2.1). $\square$

Corollary (8.3.6). — *The restriction*

$$(8.3.6.1) \quad \pi : E \longrightarrow X$$

of p to E is a surjective affine morphism. If Y is affine and f homogeneous in S_+ , then

$$(8.3.6.2) \quad \pi^{-1}(X_f) = C_f$$

and the restriction of π to C_f corresponds to the canonical injection $S_{(f)} \rightarrow S_f$. If, further, $f \in S_1$, then C_f is isomorphic to $X_f \otimes_{\mathbf{Z}} \mathbf{Z}[T, T^{-1}]$ (where T is an indeterminate).

Proof. Equation (8.3.6.2) follows immediately from (8.3.5.3) and (8.3.2.5), and shows the surjectivity of π ; we have already seen that the immersion i , restricted to C_f , corresponds [II:166] to the injection $S_f^{\leq} \rightarrow S_f$ (8.3.2). Finally, the last claim is a consequence of the fact that, for $f \in S_1$, S_f is isomorphic to $S_{(f)}[T, T^{-1}]$ (2.2.1). $\square$

Remark (8.3.7). — Whenever Y is affine, the elements of the underlying space of E are the (not-necessarily-graded) prime ideals $\mathfrak{p}$ of S not containing S_+ , by definition of the immersion ε (8.3.2). For such an ideal $\mathfrak{p}$, the $\mathfrak{p} \cap S_n$ clearly satisfy the conditions of (2.1.9), and so there exists exactly one graded prime ideal $\mathfrak{q}$ of S such that $\mathfrak{q} \cap S_n = \mathfrak{p} \cap S_n$ for all n ; the map $\pi : E \rightarrow X$ of underlying spaces can then be understood via the equation

$$(8.3.7.1) \quad \pi(\mathfrak{p}) = \mathfrak{q}.$$

Indeed, to prove this equation, it suffices to consider some homogeneous f in S_+ such that $\mathfrak{p} \in D(f)$, and to note that $\mathfrak{q}_{(f)}$ is the inverse image of $\mathfrak{p}_f$ under the injection $S_{(f)} \rightarrow S_f$.

Corollary (8.3.8). — If $\mathcal{S}$ is generated by $\mathcal{S}_1$, then the morphisms p and π are of finite type; for all $x \in X$, the fibre $p^{-1}(x)$ is isomorphic to $\text{Spec}(k(x)[T])$, and the fibre $\pi^{-1}(x)$ is isomorphic to $\text{Spec}(k(x)[T, T^{-1}])$.

Proof. This follows immediately from (8.3.5) and (8.3.6) by noting that, whenever Y is affine and S is generated by S_1 , the X_f , for $f \in S_1$, form a cover of X (2.3.14). $\square$

Remark (8.3.9). — The punctured affine cone corresponding to the graded $\mathcal{O}_Y$ -algebra $\mathcal{O}_Y[T]$ (where T is an indeterminate) can be identified with $G_m = \text{Spec}(\mathcal{O}_Y[T, T^{-1}])$, since it is exactly C_T , as we have seen in (8.3.2) (see (8.4.4) for a more general result). This prescheme is canonically endowed with the structure of a “ Y -scheme in commutative groups”. This idea will be explained in detail later on, but, for now, can be quickly summarised as follows. A Y -scheme in groups is a Y -scheme G endowed with two Y -morphisms, $p : G \times_Y G \rightarrow G$ and $s : G \rightarrow G$, that satisfy conditions formally analogous to the axioms of the composition law and the symmetry law of a group: the diagram

$$\begin{array}{ccc} G \times G \times G & \xrightarrow{p \times 1} & G \times G \\ 1 \times p \downarrow & & \downarrow p \\ G \times G & \xrightarrow{p} & G \end{array}$$

should commute (“associativity”), and there should be a condition which corresponds to the fact that, for groups, the maps

$$(x, y) \mapsto (x, x^{-1}, y) \mapsto (x, x^{-1}y) \mapsto x(x^{-1}y)$$

and

$$(x, y) \mapsto (x, x^{-1}, y) \mapsto (x, yx^{-1}) \mapsto (yx^{-1})x$$

should both reduce to $(x, y) \mapsto y$; the sequence of morphisms corresponding, for example, to the first composite map is

$$G \times G \xrightarrow{(1, s) \times 1} G \times G \times G \xrightarrow{1 \times p} G \times G \xrightarrow{p} G$$

and the reader should write down the second sequence.

[II:167] It is immediate (I, 3.4.3) that the data of a structure of a Y -scheme in groups on a Y -scheme G is equivalent to the data, for every Y -prescheme Z , of a group structure on the set $\text{Hom}_Y(Z, G)$, where these structures should be such that, for every Y -morphism $Z \rightarrow Z'$, the corresponding map $\text{Hom}_Y(Z', G) \rightarrow \text{Hom}_Y(Z, G)$ is a group homomorphism. In the particular case of G_m that we consider here, $\text{Hom}_Y(Z, G_m)$ can be identified with the set of Z -sections of $Z \times_Y G_m$ (I, 3.3.14), and thus with the set of Z -sections of $\text{Spec}(\mathcal{O}_Z[T, T^{-1}])$; finally, the same reasoning as in (I, 3.3.15) shows that this set is canonically identified with the set of invertible elements of the ring $\Gamma(Z, \mathcal{O}_Z)$, and the group structure on this set is the structure coming from the multiplication in the ring $\Gamma(Z, \mathcal{O}_Z)$. The reader can verify that the morphisms p and s from above are obtained in the following way: they correspond, by (1.2.7) and (1.4.6), to the homomorphisms of $\mathcal{O}_Y$ -algebras

$$\begin{aligned} \pi : \mathcal{O}_Y[T, T^{-1}] &\longrightarrow \mathcal{O}_Y[T, T^{-1}, T', T'^{-1}] \\ \sigma : \mathcal{O}_Y[T, T^{-1}] &\longrightarrow \mathcal{O}_Y[T, T^{-1}] \end{aligned}$$

and are entirely defined by the data of $\pi(T) = TT'$ and $\sigma(T) = T^{-1}$.

With this in mind, G_m can be considered as a “universal domain of operators” for every affine cone $C = \text{Spec}(\mathcal{S})$, where $\mathcal{S}$ is a quasi-coherent positively-graded $\mathcal{O}_Y$ -algebra. This means that we can canonically define a Y -morphism $G_m \times_Y C \rightarrow C$ which has the formal properties of an external law of a set endowed with a group of operators; or, again, as above for schemes in groups, we can give, for every Y -prescheme Z , an external law on $\text{Hom}_Y(Z, C)$, having the group $\text{Hom}_Y(Z, G_m)$ as its set of operators, with the usual axioms of sets endowed with a group of operators, and a compatibility condition with respect to the Y -morphisms $Z \rightarrow Z'$. In the current case, the morphism $G_m \times_Y C \rightarrow C$ is defined by the data of a homomorphism of $\mathcal{O}_Y$ -algebras $\mathcal{S} \rightarrow \mathcal{S} \otimes_{\mathcal{O}_Y} \mathcal{O}_Y[T, T^{-1}] = \mathcal{S}[T, T^{-1}]$, which associates, to each section $s_n \in \Gamma(U, \mathcal{S}_n)$ (where U is an open subset of Y), the section $s_n T^n \in \Gamma(U, \mathcal{S} \otimes_{\mathcal{O}_Y} \mathcal{O}_Y[T, T^{-1}])$.

Conversely, suppose that we are given a quasi-coherent, *a priori non-graded*, $\mathcal{O}_Y$ -algebra $\mathcal{S}$, and, on $C = \operatorname{Spec}(\mathcal{S})$, a structure of a “ Y -scheme in sets endowed with a group of operators” that has the Y -scheme in groups G_m as its domain of operators; then we canonically obtain a grading of $\mathcal{O}_Y$ -algebras on $\mathcal{S}$. Indeed, the data of a Y -morphism $G_m \times_Y C \rightarrow C$ is equivalent to that of a homomorphism of $\mathcal{O}_Y$ -algebras $\psi : \mathcal{S} \rightarrow \mathcal{S}[T, T^{-1}]$, which can be written as $\psi = \sum_{n \in \mathbb{Z}} \psi_n T^n$, where the $\psi_n : \mathcal{S} \rightarrow \mathcal{S}$ are homomorphisms of $\mathcal{O}_Y$ -modules (with $\psi_n(s) = 0$ except for finitely many n for every section $s \in \Gamma(U, \mathcal{S})$, for any open subset U of Y). We can then prove that the axioms of sets endowed with a group of operators imply that the $\psi_n(\mathcal{S}) = \mathcal{S}_n$ define a grading (with arbitrary integer degrees) of $\mathcal{O}_Y$ -algebras on $\mathcal{S}$, with the ψ_n being the corresponding projectors. We thus obtain the notion of a structure of an “affine cone” on every affine Y -scheme, defined in a “geometric” way without any reference to any prior grading. ^[II:168] We will not further develop this point of view here, and we leave the work of precisely formulating the definitions and results corresponding to the information given above to the reader.

8.4. Projective closure of a vector bundle

(8.4.1). Let Y be a prescheme, and $\mathcal{E}$ a quasi-coherent $\mathcal{O}_Y$ -module. If we take $\mathcal{S}$ to be the graded $\mathcal{O}_Y$ -algebra $\mathbf{S}_{\mathcal{O}_Y}(\mathcal{E})$, then Definition (8.3.1.1) shows that $\widehat{\mathcal{S}}$ can be identified with $\mathbf{S}_{\mathcal{O}_Y}(\mathcal{E} \oplus \mathcal{O}_Y)$. With the affine cone $\operatorname{Spec}(\mathcal{S})$ defined by $\mathcal{S}$ being, by definition, $\mathbf{V}(\mathcal{E})$, and $\operatorname{Proj}(\mathcal{S})$ being, by definition, $\mathbf{P}(\mathcal{E})$, we see that:

Proposition (8.4.2). — *The projective closure of a vector bundle $\mathbf{V}(\mathcal{E})$ on Y is canonically isomorphic to $\mathbf{P}(\mathcal{E} \oplus \mathcal{O}_Y)$, and the part at infinity of the latter is canonically isomorphic to $\mathbf{P}(\mathcal{E})$.*

Remark (8.4.3). — Take, for example, $\mathcal{E} = \mathcal{O}_Y^r$ with $r \geq 2$; then the punctured cones E and $\widehat{E}$ defined by $\mathcal{S}$ are neither affine nor projective over Y if $Y \neq \emptyset$. The second claim is immediate, because $\widehat{C} = \mathbf{P}(\mathcal{O}_Y^{r+1})$ is projective over Y , and the underlying spaces of E and $\widehat{E}$ are non-closed open subsets of $\widehat{C}$, and so the canonical immersions $E \rightarrow \widehat{E}$ and $\widehat{E} \rightarrow \widehat{C}$ are not projective (5.5.3); we then conclude by (5.5.5, v). Now, supposing, for example, that $Y = \operatorname{Spec}(A)$ is affine, and $r = 2$, then $C = \operatorname{Spec}(A[T_1, T_2])$, and E is then the prescheme induced by C on the open subset $D(T_1) \cup D(T_2)$; but we have already seen that the latter is not affine (I, 5.5.11); *a fortiori* $\widehat{E}$ cannot be affine, since E is the open subset where the section $\mathbf{z}$ over $\widehat{E}$ does not vanish (8.3.2).

However:

Proposition (8.4.4). — *If $\mathcal{L}$ is an invertible $\mathcal{O}_Y$ -module, then there are canonical isomorphisms for both the punctured cones E and $\widehat{E}$ corresponding to $C = \mathbf{V}(\mathcal{L})$:*

$$(8.4.4.1) \quad \operatorname{Spec} \left(\bigoplus_{n \in \mathbb{Z}} \mathcal{L}^{\otimes n} \right) \xrightarrow{\sim} E$$

$$(8.4.4.2) \quad \mathbf{V}(\mathcal{L}^{-1}) \xrightarrow{\sim} \widehat{E}.$$

Furthermore, there exists a canonical isomorphism from the projective closure of $\mathbf{V}(\mathcal{L})$ to the projective closure of $\mathbf{V}(\mathcal{L}^{-1})$ that sends the zero section (resp. the part at infinity) of the former to the part at infinity (resp. the zero section) of the latter.

Proof. We have here that $\mathcal{S} = \bigoplus_{n \geq 0} \mathcal{L}^{\otimes n}$; the canonical injection

$$\mathcal{S} \longrightarrow \bigoplus_{n \in \mathbb{Z}} \mathcal{L}^{\otimes n}$$

defines a canonical dominant morphism

$$(8.4.4.3) \quad \operatorname{Spec} \left(\bigoplus_{n \in \mathbb{Z}} \mathcal{L}^{\otimes n} \right) \longrightarrow \mathbf{V}(\mathcal{L}) = \operatorname{Spec} \left(\bigoplus_{n \geq 0} \mathcal{L}^{\otimes n} \right)$$

and it suffices to prove that this morphism is an isomorphism from the scheme $\operatorname{Spec}(\bigoplus_{n \in \mathbb{Z}} \mathcal{L}^{\otimes n})$ to E . Since the question is local on Y , we can assume that $Y = \operatorname{Spec}(A)$ is affine ^[II:169] and that $\mathcal{L} = \mathcal{O}_Y$, and so $\mathcal{S} = (A[T])^\sim$ and $\bigoplus_{n \in \mathbb{Z}} \mathcal{L}^{\otimes n} = (A[T, T^{-1}])^\sim$. But $A[T, T^{-1}]$ is the localization $A[T]_T$ of $A[T]$, and thus (8.4.4.3) identifies $\operatorname{Spec}(\bigoplus_{n \in \mathbb{Z}} \mathcal{L}^{\otimes n})$ with the prescheme induced by $C = \mathbf{V}(\mathcal{L})$ on

the open subset $D(T)$; the complement $V(T)$ of this open subset in C is the underlying space of the closed subscheme of C defined by the ideal $TA[T]$, which is exactly the zero section of C , and so $E = D(T)$.

The isomorphism in (8.4.4.2) will be a consequence of the last claim, since $\mathbf{V}(\mathcal{L}^{-1})$ is the complement of the part at infinity of its projective closure, and $\hat{E}$ is the complement of the zero section of the projective closure of $C = \mathbf{V}(\mathcal{L})$. But these projective closures are $\mathbf{P}(\mathcal{L}^{-1} \oplus \mathcal{O}_Y)$ and $\mathbf{P}(\mathcal{L} \oplus \mathcal{O}_Y)$ (respectively); but we can write $\mathcal{L} \oplus \mathcal{O}_Y = \mathcal{L} \otimes (\mathcal{L}^{-1} \oplus \mathcal{O}_Y)$. The existence of the desired canonical isomorphism then follows from (4.1.4), and everything reduces to showing that this isomorphism swaps the zero sections and the parts at infinity. For this, we can reduce to the case where $Y = \operatorname{Spec}(A)$ is affine, $L = Ac$, and $L^{-1} = Ac'$, with the canonical isomorphism $L \otimes L^{-1} \rightarrow A$ sending $c \otimes c'$ to the element 1 of A . Then $\mathbf{S}(L \oplus A)$ is the tensor product of $A[\mathbf{z}]$ with $\bigoplus_{n \geq 0} Ac^{\otimes n}$, and $\mathbf{S}(L^{-1} \oplus A)$ is the tensor product of $A[\mathbf{z}]$ with $\bigoplus_{n \geq 0} Ac'^{\otimes n}$, and the isomorphism defined in (4.1.4) sends $\mathbf{z}^h \otimes c'^{\otimes(n-h)}$ to the element $\mathbf{z}^{n-h} \otimes c^{\otimes h}$. But, in $\mathbf{P}(\mathcal{L}^{-1} \oplus \mathcal{O}_Y)$, the part at infinity is the set of points where the section $\mathbf{z}$ vanishes, and the zero section is the set of points where the section c' vanishes; since we have analogous definitions for $\mathbf{P}(\mathcal{L} \oplus \mathcal{O}_Y)$, the conclusion follows immediately from the above explanation. $\square$

8.5. Functorial behaviour

(8.5.1). Let Y and Y' be preschemes, $q : Y' \rightarrow Y$ a morphism, and $\mathcal{S}$ (resp. $\mathcal{S}'$) a quasi-coherent positively-graded $\mathcal{O}_Y$ -algebra (resp. quasi-coherent positively-graded $\mathcal{O}_{Y'}$ -algebra). Consider a q -morphism of graded algebras

$$(8.5.1.1) \quad \phi : \mathcal{S} \longrightarrow \mathcal{S}'.$$

We know (1.5.6) that this corresponds, canonically, to a morphism

$$\Phi = \operatorname{Spec}(\phi) : \operatorname{Spec}(\mathcal{S}') \longrightarrow \operatorname{Spec}(\mathcal{S})$$

such that the diagram

$$(8.5.1.2) \quad \begin{array}{ccc} C' & \xrightarrow{\Phi} & C \\ \downarrow & & \downarrow \\ Y' & \xrightarrow{q} & Y \end{array}$$

commutes, where we write $C = \operatorname{Spec}(\mathcal{S})$ and $C' = \operatorname{Spec}(\mathcal{S}')$. Suppose, further, that $\mathcal{S}_0 = \mathcal{O}_Y$ and $\mathcal{S}'_0 = \mathcal{O}_{Y'}$; let $\varepsilon : Y \rightarrow C$ and $\varepsilon' : Y' \rightarrow C'$ be the canonical immersions (8.3.2); we then have a commutative diagram

$$(8.5.1.3) \quad \begin{array}{ccc} Y' & \xrightarrow{q} & Y \\ \varepsilon' \downarrow & & \downarrow \varepsilon \\ C' & \xrightarrow{\Phi} & C \end{array}$$

[II:170] which corresponds to the diagram

$$\begin{array}{ccc} \mathcal{S} & \xrightarrow{\phi} & \mathcal{S}' \\ \downarrow & & \downarrow \\ \mathcal{O}_Y & \longrightarrow & \mathcal{O}_{Y'} \end{array}$$

where the vertical arrows are the augmentation homomorphisms, and so the commutativity follows from the hypothesis that ϕ is assumed to be a homomorphism of graded algebras.

Proposition (8.5.2). — *If E (resp. E') is the punctured affine cone defined by $\mathcal{S}$ (resp. $\mathcal{S}'$), then $\Phi^{-1}(E) \subset E'$; if, further, $\operatorname{Proj}(\phi) : G(\phi) \rightarrow \operatorname{Proj}(\mathcal{S})$ is everywhere defined (or, equivalently, if $G(\phi) = \operatorname{Proj}(\mathcal{S}')$), then $\Phi^{-1}(E) = E'$, and conversely.*

Proof. The first claim follows from the commutativity of (8.5.1.3). To prove the second, we can restrict to the case where $Y = \text{Spec}(A)$ and $Y' = \text{Spec}(A')$ are affine, and $\mathcal{S} = \tilde{S}$ and $\mathcal{S}' = \tilde{S}'$. For every homogeneous f in S_+ , writing $f' = \phi(f)$, we have that $\Phi^{-1}(C_f) = C'_{f'}$ (I, 2.2.4.1); saying that $G(\phi) = \text{Proj}(S')$ implies that the radical (in S'_+) of the ideal generated by the $f' = \phi(f)$ is S'_+ itself ((2.8.1) and (2.3.14)), and this is equivalent to saying that the $C'_{f'}$ cover E' (8.3.4.4). $\square$

(8.5.3). The q -morphism ϕ canonically extends to a q -morphism of graded algebras

$$(8.5.3.1) \quad \hat{\phi} : \hat{\mathcal{S}} \longrightarrow \hat{\mathcal{S}'}$$

by letting $\hat{\phi}(\mathbf{z}) = \mathbf{z}$. This induces a morphism

$$\hat{\Phi} = \text{Proj}(\hat{\phi}) : G(\hat{\phi}) \longrightarrow \hat{C} = \text{Proj}(\hat{\mathcal{S}'})$$

such that the diagram

$$\begin{array}{ccc} G(\hat{\phi}) & \xrightarrow{\hat{\Phi}} & \hat{C} \\ \downarrow & & \downarrow \\ Y' & \xrightarrow{q} & Y \end{array}$$

commutes (3.5.6). It follows immediately from the definitions that, if we write $i : C \rightarrow \hat{C}$ and $i' : C' \rightarrow \hat{C}'$ to mean the canonical open immersions (8.3.2), then $i'(C') \subset G(\hat{\phi})$, and the diagram

$$(8.5.3.2) \quad \begin{array}{ccc} C' & \xrightarrow{\Phi} & C \\ i' \downarrow & & \downarrow i \\ G(\hat{\phi}) & \xrightarrow{\hat{\Phi}} & \hat{C} \end{array}$$

commutes. Finally, if we let $X = \text{Proj}(\mathcal{S})$ and $X' = \text{Proj}(\mathcal{S}')$, and if $j : X \rightarrow \hat{C}$ and $j' : X' \rightarrow \hat{C}'$ are the canonical closed immersions (8.3.2), then it follows from the definition of these immersions that $j'(G(\phi)) \subset G(\hat{\phi})$, and that the diagram [III:171]

$$(8.5.3.3) \quad \begin{array}{ccc} G(\phi) & \xrightarrow{\text{Proj}(\phi)} & X \\ j' \downarrow & & \downarrow j \\ G(\hat{\phi}) & \xrightarrow{\hat{\Phi}} & \hat{C} \end{array}$$

commutes.

Proposition (8.5.4). — *If $\hat{E}$ (resp. $\hat{E}'$) is the punctured projective cone defined by $\mathcal{S}$ (resp. by $\mathcal{S}'$), then $\hat{\Phi}^{-1}(\hat{E}) \subset \hat{E}'$; furthermore, if $p : \hat{E} \rightarrow X$ and $p' : \hat{E}' \rightarrow X'$ are the canonical retractions, then $p'(\hat{\Phi}^{-1}(\hat{E})) \subset G(\phi)$, and the diagram*

$$(8.5.4.1) \quad \begin{array}{ccc} \hat{\Phi}^{-1}(\hat{E}) & \xrightarrow{\hat{\Phi}} & \hat{E} \\ p' \downarrow & & \downarrow p \\ G(\phi) & \xrightarrow{\text{Proj}(\phi)} & X \end{array}$$

commutes. If $\text{Proj}(\phi)$ is everywhere defined, then so too is $\hat{\Phi}$, and we have that $\hat{\Phi}^{-1}(\hat{E}) = \hat{E}'$.

Proof. The first claim follows from the commutativity of Diagrams (8.5.1.3) and (8.5.3.2), and the two following claims from the definition of the canonical retractions (8.3.5) and the definition of $\hat{\phi}$. To see that $\hat{\Phi}$ is everywhere defined whenever $\text{Proj}(\phi)$ is, we can restrict to the case where $Y = \text{Spec}(A)$ and $Y' = \text{Spec}(A')$ are affine, and where $\mathcal{S} = \tilde{S}$ and $\mathcal{S}' = \tilde{S}'$; the hypothesis is that, when f runs over the set of homogeneous elements of S_+ , the radical in S'_+ of the ideal generated in S'_+ by the $\phi(f)$ is S'_+ itself; we thus immediately conclude that the radical in $(S'[\mathbf{z}])_+$ of the ideal

generated by $\mathbf{z}$ and the $\phi(f)$ is $(S'[\mathbf{z}])_+$ itself, whence our claim; this also shows that $\widehat{E}'$ is the union of the $\widehat{C}'_{\phi(f)}$, and hence equal to $\widehat{\Phi}^{-1}(\widehat{E})$. $\square$

Corollary (8.5.5). — *Whenever $\text{Proj}(\phi)$ is everywhere defined, the inverse image under $\widehat{\Phi}$ of the underlying space of the part at infinity (resp. of the vertex prescheme) of $\widehat{C}$ is the underlying space of the part at infinity (resp. of the vertex prescheme) of $\widehat{C}'$.*

Proof. This follows immediately from (8.5.4) and (8.5.2), taking into account the equalities (8.3.4.1) and (8.3.4.2). $\square$

8.6. A canonical isomorphism for punctured cones

(8.6.1). Let Y be a prescheme, $\mathcal{S}$ a quasi-coherent positively-graded $\mathcal{O}_Y$ -algebra such that $\mathcal{S}_0 = \mathcal{O}_Y$, and let X be the Y -scheme $\text{Proj}(\mathcal{S})$. We are going to apply the results of (8.5) to the case where $Y' = X$, and $q : X \rightarrow Y$ is the structure morphism; let

$$(8.6.1.1) \quad \mathcal{S}_X = \bigoplus_{n \in \mathbb{Z}} \mathcal{O}_X(n)$$

[II:172] which is a quasi-coherent graded $\mathcal{O}_X$ -algebra, with multiplication defined by means of the canonical homomorphisms (3.2.6.1)

$$\mathcal{O}_X(m) \otimes_{\mathcal{O}_X} \mathcal{O}_X(n) \longrightarrow \mathcal{O}_X(m+n)$$

whose associativity is ensured by the commutative diagram in (2.5.11.4). Let $\mathcal{S}'$ be the quasi-coherent positively-graded $\mathcal{O}_X$ -subalgebra $\mathcal{S}'_X = \bigoplus_{n \geq 0} \mathcal{O}_X(n)$ of $\mathcal{S}_X$.

Finally, consider the canonical q -morphism

$$(8.6.1.2) \quad \alpha : \mathcal{S} \longrightarrow \mathcal{S}'_X$$

defined in (3.3.2.3) as a homomorphism $\mathcal{S} \rightarrow q_*(\mathcal{S}_X)$, but which clearly sends $\mathcal{S}$ to $q_*(\mathcal{S}'_X)$. Write

$$(8.6.1.3) \quad C_X = \text{Spec}(\mathcal{S}'_X), \quad \widehat{C}_X = \text{Proj}(\mathcal{S}'_X[\mathbf{z}]), \quad X' = \text{Proj}(\mathcal{S}'_X)$$

and denote by E_X and $\widehat{E}_X$ the corresponding punctured affine and punctured projective cones (respectively); denote the canonical morphisms defined in (8.3) by $\varepsilon_X : X \rightarrow C_X$, $i_X : C_X \rightarrow \widehat{C}_X$, $j_X : X' \rightarrow \widehat{C}_X$, $p_X : \widehat{E}_X \rightarrow X'$, and $\pi_X : E_X \rightarrow X'$.

Proposition (8.6.2). — *The structure morphism $u : X' \rightarrow X$ is an isomorphism, and the morphism $\text{Proj}(\alpha)$ is everywhere defined and identical to u . The morphism $\text{Proj}(\widehat{\alpha}) : \widehat{C}_X \rightarrow \widehat{C}$ is everywhere defined, and its restrictions to $\widehat{E}_X$ and E_X are isomorphisms to $\widehat{E}$ and E (respectively). Finally, if we identify X' with X via u , then the morphisms p_X and π_X are identified with the structure morphisms of the X -preschemes $\widehat{E}_X$ and E_X .*

Proof. We can clearly restrict to the case where $Y = \text{Spec}(A)$ is affine, and $\mathcal{S} = \widetilde{S}$; then X is the union of affine open subsets X_f , where f runs over the set of homogeneous elements of S_+ , with the ring of each X_f being $S_{(f)}$. It follows from (8.2.7.1) that

$$(8.6.2.1) \quad \Gamma(X_f, \mathcal{S}'_X) = S_f.$$

So $u^{-1}(X_f) = \text{Proj}(S_f^{\geq})$. But if $f \in S_d$ ($d > 0$), then $\text{Proj}(S_f^{\geq})$ is canonically isomorphic to $\text{Proj}((S_f^{\geq})^{(d)})$ (2.4.7), and we also know that $(S_f^{\geq})^{(d)} = (S^{(d)})_f^{\geq}$ can be identified with $S_{(f)}[T]$ (2.2.1) by the map $T \mapsto f/1$; we thus conclude (3.1.7) that the structure morphism $u^{-1}(X_f) \rightarrow X_f$ is an isomorphism, whence the first claim. To prove the second, note that the restriction $u^{-1}(X_f) \cap G(\alpha) \rightarrow X = \text{Proj}(S)$ of $\text{Proj}(\alpha)$ corresponds to the canonical map $x \mapsto x/1$ from S to $S_f^{\geq}$ (2.6.2); we thus deduce, first of all, that $G(\alpha) = X'$, and then, taking into account the fact that $u^{-1}(X_f) = (u^{-1}(X_f))_{f/1}$, that it follows from (2.8.1.1) that the image of $u^{-1}(X_f)$ under $\text{Proj}(\alpha)$ is contained in X_f , and the restriction of $\text{Proj}(\alpha)$ to $u^{-1}(X_f)$, considered as a morphism to $X_f = \text{Spec}(S_{(f)})$, is indeed identical to that of u . Finally, applying (8.3.5.4) to p_X instead of p , we see that $p_X^{-1}(u^{-1}(X_f)) = \text{Spec}((S_f^{\geq})_{f/1}^{\leq})$, and this open subset is, by (8.5.4.1), the inverse image under $\text{Proj}(\widehat{\alpha})$ of $p^{-1}(X_f) =$

$\text{Spec}(S_f^{\leq})$ (8.3.5.3). Taking (8.2.3.2) into account, the restriction of $\text{Proj}(\hat{\alpha})$ to $p_X^{-1}(u^{-1}(X_f))$ corresponds to the isomorphism inverse to (8.2.7.2), restricted to $S_f^{\leq}$, whence the third claim; the last claim is evident by definition.

[II:173] We note also that it follows from the commutative diagram in (8.5.3.2) that the restriction to C_X of $\text{Proj}(\hat{\alpha})$ is exactly the morphism $\text{Spec}(\alpha)$. $\square$

Corollary (8.6.3). — Considered as X -schemes, $\hat{E}_X$ is canonically isomorphic to $\text{Spec}(\mathcal{S}_X^{\leq})$, and E_X to $\text{Spec}(\mathcal{S}_X)$.

Proof. Since we know that the morphisms p_X and π_X are affine ((8.3.5) and (8.3.6)), it suffices (given (1.3.1)) to prove the corollary in the case where $Y = \text{Spec}(A)$ is affine and $\mathcal{S} = \tilde{\mathcal{S}}$. The first claim follows from the existence of the canonical isomorphisms (8.2.7.2) $(S_f^{\geq})_{f/1}^{\leq} \xrightarrow{\sim} S_f^{\leq}$ and from the fact that these isomorphisms are compatible with the map sending f to fg (where f and g are homogeneous in S_+). Similarly, applying (8.3.6.2) to π_X instead of π , we see that $\pi_X^{-1}(u^{-1}(X_f)) = \text{Spec}((S_f^{\geq})_{f/1})$ for f homogeneous in S_+ , and the second claim then follows from the existence of the canonical isomorphisms (8.2.7.2) $(S_f^{\geq})_{f/1} \xrightarrow{\sim} S_f$.

We can then say that $\hat{C}_X$, considered as an X -scheme, is given by *gluing* the affine X -schemes $C_X = \text{Spec}(\mathcal{S}_X^{\geq})$ and $\hat{E}_X = \text{Spec}(\mathcal{S}_X^{\leq})$ over X , where the intersection of the two affine X -schemes is the open subset $E_X = \text{Spec}(\mathcal{S}_X)$. $\square$

Corollary (8.6.4). — Assume that $\mathcal{O}_X(1)$ is an invertible $\mathcal{O}_X$ -module, and that $\mathcal{S}_X$ is isomorphic to $\bigoplus_{n \in \mathbf{Z}} (\mathcal{O}_X(1))^{\otimes n}$ (which will be the case, in particular, whenever $\mathcal{S}$ is generated by $\mathcal{S}_1$ ((3.2.5) and (3.2.7))). Then the punctured projective cone $\hat{E}$ can be identified with the rank-1 vector bundle $\mathbf{V}(\mathcal{O}_X(-1))$ on X , and the punctured affine cone E with the subprescheme of this vector bundle induced on the complement of the zero section. With this identification, the canonical retraction $\hat{E} \rightarrow X$ is identified with the structure morphism of the X -scheme $\mathbf{V}(\mathcal{O}_X(-1))$. Finally, there exists a canonical Y -morphism $\mathbf{V}(\mathcal{O}_X(1)) \rightarrow C$, whose restriction to the complement of the zero section of $\mathbf{V}(\mathcal{O}_X(1))$ is an isomorphism from this complement to the punctured affine cone E .

Proof. If we write $\mathcal{L} = \mathcal{O}_X(1)$, then $\mathcal{S}_X^{\geq}$ is identical to $\mathbf{S}_{\mathcal{O}_X}(\mathcal{L})$, and so $\hat{E}_X$ is canonically identified with $\mathbf{V}(\mathcal{L}^{-1})$, by (8.6.3), and C_X with $\mathbf{V}(\mathcal{L})$. The morphism $\mathbf{V}(\mathcal{L}) \rightarrow C$ is the restriction of $\text{Proj}(\hat{\alpha})$, and the claims of the corollary are then particular cases of (8.6.2). $\square$

We note that the inverse image under the morphism $\mathbf{V}(\mathcal{O}_X(1)) \rightarrow C$ of the underlying space of the vertex prescheme of C is the underlying space of the zero section of $\mathbf{V}(\mathcal{O}_X(1))$ (8.5.5); but, in general, the corresponding subpreschemes of C and of $\mathbf{V}(\mathcal{O}_X(1))$ are not isomorphic. This problem will be studied below.

8.7. Blowing up projecting cones

(8.7.1). Under the conditions of (8.6.1), we have, writing $r = \text{Proj}(\hat{\alpha})$, a commutative diagram

$$(8.7.1.1) \quad \begin{array}{ccc} X & \xrightarrow{i_X \circ \varepsilon_X} & \hat{C}_X \\ q \downarrow & & \downarrow r \\ Y & \xrightarrow{i_Y \circ \varepsilon} & \hat{C} \end{array}$$

[II:174] by (8.5.1.3) and (8.5.3.2); furthermore, the restriction of r to the complement $\hat{C}_X - i_X(\varepsilon_X(X))$ of the zero section is an isomorphism to the complement $\hat{C} - i(\varepsilon(Y))$ of the zero section, by (8.6.2). If we suppose, to simplify things, that Y is affine and that $\mathcal{S}$ is of finite type and generated by $\mathcal{S}_1$, then X is projective over Y and $\hat{C}_X$ is projective over X (5.5.1); hence $\hat{C}_X$ is projective over Y (5.5.5, ii), and *a fortiori* over $\hat{C}$ (5.5.5, v). We then have a projective Y -morphism $r : \hat{C}_X \rightarrow \hat{C}$ (whose restriction to C_X is a projective Y -morphism $C_X \rightarrow C$) that *contracts* X to Y and induces an isomorphism when restricted to the complements of X and Y . We thus have a connection between C_X and C , analogous to that which exists between a blow-up prescheme and the original prescheme (8.1.3). We will effectively show that C_X can be identified with the homogeneous spectrum of a graded $\mathcal{O}_C$ -algebra.

(8.7.2). Keeping the notation of (8.6.1), consider, for all $n \geq 0$, the quasi-coherent ideal

$$(8.7.2.1) \quad \mathcal{S}_{[n]} = \bigoplus_{m \geq n} \mathcal{S}_m$$

of the graded $\mathcal{O}_Y$ -algebra $\mathcal{S}$. It is clear that

$$(8.7.2.2) \quad \mathcal{S}_{[0]} = \mathcal{S}, \quad \mathcal{S}_{[n]} \subset \mathcal{S}_{[m]} \quad \text{for } m \leq n$$

$$(8.7.2.3) \quad \mathcal{S}_{[m]} \mathcal{S}_{[n]} \subset \mathcal{S}_{[m+n]}.$$

Consider the $\mathcal{O}_C$ -module associated to $\mathcal{S}_{[n]}$, which is a quasi-coherent ideal of $\mathcal{O}_C = \widetilde{\mathcal{S}}$ (1.4.4)

$$(8.7.2.4) \quad \mathcal{I}_n = (\mathcal{S}_{[n]})^\sim.$$

We thus deduce, from (8.7.2.2) and (8.7.2.3), using (1.4.4) and (1.4.8.1), the analogous formulas

$$(8.7.2.5) \quad \mathcal{I}_0 = \mathcal{O}_C, \quad \mathcal{I}_n \subset \mathcal{I}_m \quad \text{for } m \leq n$$

$$(8.7.2.6) \quad \mathcal{I}_m \mathcal{I}_n \subset \mathcal{I}_{m+n}.$$

We are thus in the setting of (8.1.1), which leads us to introduce the quasi-coherent graded $\mathcal{O}_C$ -algebra

$$(8.7.2.7) \quad \mathcal{S}^\natural = \bigoplus_{n \geq 0} \mathcal{I}_n = \left(\bigoplus_{n \geq 0} \mathcal{S}_{[n]} \right)^\sim.$$

Proposition (8.7.3). — *There is a canonical C -isomorphism*

$$(8.7.3.1) \quad h : C_X \xrightarrow{\sim} \text{Proj}(\mathcal{S}^\natural).$$

Proof. Suppose first of all that $Y = \text{Spec}(A)$ is affine, so that $\mathcal{S} = \widetilde{S}$, with S a positively-graded A -algebra, and $C = \text{Spec}(S)$. Definition (8.7.2.4) then shows, with the notation of (8.2.6), that $\mathcal{S}^\natural = (S^\natural)^\sim$. To define (8.7.3.1), consider a homogeneous element $f \in S_d$ ($d > 0$) and the corresponding element $f^\natural \in S^\natural$ (8.2.6); the S -isomorphism in (8.2.7.3) then defines a C -isomorphism

$$(8.7.3.2) \quad \text{Spec}(S_f^\geq) \xrightarrow{\sim} \text{Spec}(S_{(f^\natural)}^\natural).$$

[II:175] But with the notation of (8.6.2), if $v : C_X \rightarrow X$ is the structure morphism, then it follows from (8.6.2.1) that $v^{-1}(X_f) = \text{Spec}(S_f^\geq)$. We also have that $\text{Spec}(S_{(f^\natural)}^\natural) = D_+(f^\natural)$, which means that (8.7.3.2) defines an isomorphism $v^{-1}(X_f) \rightarrow D_+(f^\natural)$. Furthermore, if $g \in S_e$ ($e > 0$), then the diagram

$$\begin{array}{ccc} v^{-1}(X_{fg}) & \xrightarrow{\sim} & D_+(f^\natural g^\natural) \\ \downarrow & & \downarrow \\ v^{-1}(X_f) & \xrightarrow{\sim} & D_+(f^\natural) \end{array}$$

commutes, by definition of the isomorphism in (8.2.7.3). Finally, by definition, S_+ is generated by the homogeneous f , and so it follows from (8.2.10, iv) and from (2.3.14) that the $D_+(f^\natural)$ form a cover of $\text{Proj}(S^\natural)$, and that the $v^{-1}(X_f)$ form a cover of C_X , since the X_f form a cover of X ; in this case, we have thus defined the isomorphism (8.7.3.1).

To prove (8.7.3) in the general case, it suffices to show that, if U and U' are affine open subsets of Y , given by rings A and A' (respectively), and such that $U' \subset U$, then, setting $\mathcal{S}|_U = \widetilde{S}$ and $\mathcal{S}|_{U'} = \widetilde{S}'$, the diagram

$$(8.7.3.3) \quad \begin{array}{ccc} C_{U'} & \longrightarrow & \text{Proj}(\mathcal{S}'^\natural) \\ \downarrow & & \downarrow \\ C_U & \longrightarrow & \text{Proj}(\mathcal{S}^\natural) \end{array}$$

commutes. But S' is canonically identified with $S \otimes_A A'$, and so $S'^\natural$ is canonically identified with

$$S^\natural \otimes_S S' = S^\natural \otimes_A A';$$

thus $\text{Proj}(S^{\natural}) = \text{Proj}(S^{\natural}) \times_U U'$ (2.8.10); similarly, if $X = \text{Proj}(S)$ and $X' = \text{Proj}(S')$, then $X' = X \times_U U'$ and $\mathcal{S}_{X'} = \mathcal{S}_X \otimes_{\mathcal{O}_U} \mathcal{O}_{U'}$ (3.5.4), or, equivalently, $\mathcal{S}_{X'} = j^*(\mathcal{S}_X)$, where j is the projection $X' \rightarrow X$. We then have, by (1.5.2), that $C_{U'} = C_U \times_X X' = C_U \times_U U'$, and the commutativity of (8.7.3.3) is immediate. $\square$

Remark (8.7.4). — (i) The end of the proof of (8.7.3) can be immediately generalised in the following way. Let $g : Y' \rightarrow Y$ be a morphism, $\mathcal{S}' = g^*(\mathcal{S})$, and $X' = \text{Proj}(\mathcal{S}')$; then we have a commutative diagram

$$(8.7.4.1) \quad \begin{array}{ccc} C_{X'} & \longrightarrow & \text{Proj}(\mathcal{S}'^{\natural}) \\ \downarrow & & \downarrow \\ C_X & \longrightarrow & \text{Proj}(\mathcal{S}^{\natural}) \end{array}$$

Now let $\phi : \mathcal{S}'' \rightarrow \mathcal{S}$ be a homomorphism of graded $\mathcal{O}_Y$ -algebras such that, if we write $X'' = \text{Proj}(\mathcal{S}'')$, then $u = \text{Proj}(\phi) : X \rightarrow X''$ is everywhere defined; we also have [II:176] a Y -morphism $v : C \rightarrow C''$ (with $C'' = \text{Spec}(\mathcal{S}'')$) such that $\mathcal{A}(v) = \phi$, and, since ϕ is a homomorphism of graded algebras, ϕ induces a v -morphism of graded algebras $\psi : \mathcal{S}''^{\natural} \rightarrow \mathcal{S}^{\natural}$ (1.4.1). Furthermore, it follows from (8.2.10, iv) and from the hypothesis on ϕ that $\text{Proj}(\psi)$ is everywhere defined. Finally, taking (3.5.6.1) into account, there is a canonical u -morphism $\mathcal{S}_{X''} \rightarrow \mathcal{S}_X$, whence (1.5.6) a morphism $w : C_{X''} \rightarrow C_X$. With this in mind, the diagram

$$(8.7.4.2) \quad \begin{array}{ccc} C_{X''} & \xrightarrow{\sim} & \text{Proj}(\mathcal{S}''^{\natural}) \\ w \downarrow & & \downarrow \text{Proj}(\psi) \\ C_X & \xrightarrow{\sim} & \text{Proj}(\mathcal{S}^{\natural}) \end{array}$$

is commutative, as we can immediately verify by restricting to the case where Y is affine.

- (ii) Note that, by (8.7.2.5) and (8.7.2.6), we have $\mathcal{I}_1^m \subset \mathcal{I}_m \subset \mathcal{I}_1$ for all $m > 0$. But, by definition, $\mathcal{I}_1 = (\mathcal{S}_+)^{\sim}$, and so $\mathcal{I}_1$ defines the closed subprescheme $\varepsilon(Y)$ in C ((1.4.10) and (8.3.2)); we thus conclude that, for all $m > 0$, the support of $\mathcal{O}_C/\mathcal{I}_m$ is contained in the underlying space of the vertex prescheme $\varepsilon(Y)$; on the inverse image of the punctured affine cone E , the structure morphism $\text{Proj}(\mathcal{S}^{\natural}) \rightarrow C$ thus restricts to an isomorphism (by (8.7.3) and (8.7.1)). Furthermore, by canonically identifying C with an open subset of $\widehat{C}$ (8.3.3), we can clearly extend the ideals $\mathcal{I}_m$ of $\mathcal{O}_C$ to ideals $\mathcal{I}_m$ of $\mathcal{O}_{\widehat{C}}$, by asking them to agree with $\mathcal{O}_{\widehat{C}}$ on the open subset $\widehat{E}$ of $\widehat{C}$. If we define $\mathcal{T} = \bigoplus_{n \geq 0} \mathcal{I}_n$, which is a quasi-coherent graded $\mathcal{O}_{\widehat{C}}$ -algebra, we can extend the isomorphism (8.7.3.1) to a $\widehat{C}$ -isomorphism

$$(8.7.4.3) \quad \widehat{C}_X \xrightarrow{\sim} \text{Proj}(\mathcal{T}).$$

Indeed, over $\widehat{E}$, it follows from the above that $\text{Proj}(\mathcal{T})$ is canonically identified with $\widehat{E}$, and we thus define the isomorphism (8.7.4.3) over $\widehat{E}$ by asking for it to agree with the canonical isomorphism $\widehat{E}_X \rightarrow \widehat{E}$ (8.6.2); it is clear that this isomorphism and (8.7.3.1) then agree over E .

Corollary (8.7.5). — Suppose that there exists some $n_0 > 0$ such that

$$(8.7.5.1) \quad \mathcal{S}_{n+1} = \mathcal{S}_1 \mathcal{S}_n \quad \text{for } n \geq n_0.$$

Then the vertex subprescheme of C_X (isomorphic to X) is the inverse image under the canonical morphism $r : C_X \rightarrow C$ of the vertex subprescheme of C (isomorphic to Y). Conversely, if this property is true, and if we further assume that Y is Noetherian and that $\mathcal{S}$ is of finite type, then there exists some $n_0 > 0$ such that (8.7.5.1) holds true.

Proof. Since the first claim is local on Y , we can assume that $Y = \text{Spec}(A)$ is affine, so that $\mathcal{S} = \widetilde{S}$, with S a positively-graded A -algebra. The claim then follows from (8.2.12), since $\text{Proj}(S^{\natural} \otimes_S S_0) = C_X \times_C \varepsilon(Y)$ (by the identification in (8.7.3.1)), or, in other words, since this prescheme is the inverse image of $\varepsilon(Y)$ in C_X (I, 4.4.1). The converse also follows from (8.2.12) whenever Y is Noetherian

affine and S is of finite type. ^[II:177] If Y is Noetherian (but not necessarily affine) and $\mathcal{S}$ is of finite type, then there exists a finite cover of Y by Noetherian affine open subsets U_i , and we then deduce from the above that, for all i , there exists an integer n_i such that $\mathcal{S}_{n+1}|U_i = (\mathcal{S}_1|U_i)(\mathcal{S}_n|U_i)$ for $n \geq n_i$; the largest of the n_i then ensures that (8.7.5.1) holds true. $\square$

(8.7.6). Now consider the C -prescheme Z given by *blowing up* the *vertex subprescheme* $\varepsilon(Y)$ in the affine cone C ; by Definition (8.1.3), it is exactly the prescheme $\text{Proj}(\bigoplus_{n \geq 0} \mathcal{S}_+^n)$; the canonical injection

$$(8.7.6.1) \quad \iota : \bigoplus_{n \geq 0} \mathcal{S}_+^n \longrightarrow \mathcal{S}_+^\natural$$

defines (by the identification in (8.7.3)) a canonical dominant C -morphism

$$(8.7.6.2) \quad G(\iota) \longrightarrow Z$$

where $G(\iota)$ is an open subset of C_X (3.5.1); note that it could be the case that $G(\iota) \neq C_X$, as shown by the example where $Y = \text{Spec}(K)$, with K a field, and $\mathcal{S} = \tilde{S}$, with $S = K[\mathbf{y}]$, where $\mathbf{y}$ is an indeterminate of degree 2; if R_n denotes the set $(S_+)^n$, considered as a subset of $S_{[n]} = S_n^\natural$, then $S_+^\natural$ is not the radical in $S^\natural$ of the ideal generated by the union of the R_n (cf. (2.3.14)).

Corollary (8.7.7). — Assume that there exists some $n_0 > 0$ such that

$$(8.7.7.1) \quad \mathcal{S}_n = \mathcal{S}_1^n \quad \text{for } n \geq n_0.$$

Then the canonical morphism (8.7.6.2) is everywhere defined, and is an isomorphism $C_X \xrightarrow{\sim} Z$. Conversely, if this property is true, and if we further assume that Y is Noetherian and that $\mathcal{S}$ is of finite type, then there exists some n_0 such that (8.7.7.1) holds true.

Proof. The first claim is local on Y , and thus follows from (8.2.14); the converse follows similarly, arguing as in (8.7.5). $\square$

Remark (8.7.8). — Since condition (8.7.7.1) implies (8.7.5.1), we see that, whenever it holds true, not only can C_X be identified with the prescheme given by blowing up the vertex (identified with Y) of the affine cone C , but also the vertex (identified with X) of C_X can be identified with the closed subprescheme given by the inverse image of the vertex Y of C . Furthermore, hypothesis (8.7.7.1) implies that, on $X = \text{Proj}(\mathcal{S})$, the $\mathcal{O}_X$ -modules $\mathcal{O}_X(n)$ are invertible ((3.2.5) and (3.2.9)), and that $\mathcal{O}_X(n) = \mathcal{L}^{\otimes n}$ with $\mathcal{L} = \mathcal{O}_X(1)$ ((3.2.7) and (3.2.9)); by Definition (8.6.1.1), C_X is thus the *vector bundle* $\mathbf{V}(\mathcal{L})$ on X , and its vertex is the *zero section* of this vector bundle.

8.8. Ample sheaves and contractions

(8.8.1). Let Y be a prescheme, $f : X \rightarrow Y$ a *separated* and *quasi-compact* morphism, and $\mathcal{L}$ an invertible $\mathcal{O}_X$ -module that is *ample relative to* f . Consider the positively-graded $\mathcal{O}_Y$ -algebra

$$(8.8.1.1) \quad \mathcal{S} = \mathcal{O}_Y \oplus \bigoplus_{n \geq 1} f_*(\mathcal{L}^{\otimes n})$$

^[II:178] which is quasi-coherent (I, 9.2.2, a). There is a canonical homomorphism of graded $\mathcal{O}_X$ -algebras

$$(8.8.1.2) \quad \tau : f^*(\mathcal{S}) \longrightarrow \bigoplus_{n \geq 0} \mathcal{L}^{\otimes n}$$

which, in degrees ≥ 1 , agrees with the canonical homomorphism $\sigma : f^*(f_*(\mathcal{L}^{\otimes n})) \rightarrow \mathcal{L}^{\otimes n}$ (0, 4.4.3), and is the identity in degree 0. The hypothesis that $\mathcal{L}$ is *f-ample* then implies ((4.6.3) and (3.6.1)) that the corresponding Y -morphism

$$(8.8.1.3) \quad r = r_{\mathcal{L}, \tau} : X \longrightarrow P = \text{Proj}(\mathcal{S})$$

is everywhere defined and is a *dominant open immersion*, and that

$$(8.8.1.4) \quad r^*(\mathcal{O}_P(n)) = \mathcal{L}^{\otimes n} \quad \text{for all } n \in \mathbf{Z}.$$

Proposition (8.8.2). — Let $C = \operatorname{Spec}(\mathcal{S})$ be the affine cone defined by $\mathcal{S}$; if $\mathcal{L}$ is f -ample, then there exists a canonical Y -morphism

$$(8.8.2.1) \quad g : V = \mathbf{V}(\mathcal{L}) \longrightarrow C$$

such that the diagram

$$(8.8.2.2) \quad \begin{array}{ccccc} X & \xrightarrow{j} & \mathbf{V}(\mathcal{L}) & \xrightarrow{\pi} & X \\ f \downarrow & & \downarrow g & & \downarrow f \\ Y & \xrightarrow{\varepsilon} & C & \xrightarrow{\psi} & Y \end{array}$$

commutes, where ψ and π are the structure morphisms, and j and ε the canonical immersions sending X and Y (respectively) to the zero section of $\mathbf{V}(\mathcal{L})$ and the vertex prescheme of C (respectively). Furthermore, the restriction of g to $\mathbf{V}(\mathcal{L}) - j(X)$ is an open immersion

$$(8.8.2.3) \quad \mathbf{V}(\mathcal{L}) - j(X) \longrightarrow E = C - \varepsilon(Y)$$

into the punctured affine cone E corresponding to $\mathcal{S}$.

Proof. With the notation of (8.8.1), let $\mathcal{S}_P^{\geq} = \bigoplus_{n \geq 0} \mathcal{O}_P(n)$ and $C_P = \operatorname{Spec}(\mathcal{S}_P^{\geq})$. We know (8.6.2) that there is a canonical morphism $h = \operatorname{Spec}(\alpha) : C_P \rightarrow C$ such that the diagram

$$(8.8.2.4) \quad \begin{array}{ccc} C_P & \longrightarrow & P \\ h \downarrow & & \downarrow p \\ C & \xrightarrow{\psi} & Y \end{array}$$

commutes; furthermore, if $\varepsilon_P : P \rightarrow C_P$ is the canonical immersion, then the diagram

$$(8.8.2.5) \quad \begin{array}{ccc} P & \xrightarrow{p} & C_P \\ \varepsilon_P \downarrow & & \downarrow h \\ Y & \xrightarrow{\varepsilon} & C \end{array}$$

commutes (8.7.1.1), and, finally, the restriction of h to the punctured affine cone E_P is an isomorphism $E_P \xrightarrow{\sim} E$ (8.6.2). It follows from (8.8.1.4) that

$$r^*(\mathcal{S}_P^{\geq}) = \mathbf{S}_{\mathcal{O}_X}(\mathcal{L})$$

[II:179] and so we have a canonical P -morphism $q : \mathbf{V}(\mathcal{L}) \rightarrow C_P$, with the commutative diagram

$$(8.8.2.6) \quad \begin{array}{ccc} \mathbf{V}(\mathcal{L}) & \xrightarrow{\pi} & X \\ q \downarrow & & \downarrow r \\ C_P & \longrightarrow & P \end{array}$$

identifying $\mathbf{V}(\mathcal{L})$ with the product $C_P \times_P X$ (1.5.2); since r is an open immersion, so too is q (I, 4.3.2). Furthermore, the restriction of q to $\mathbf{V}(\mathcal{L}) - j(X)$ sends this prescheme to E_P , by (8.5.2), and the diagram

$$(8.8.2.7) \quad \begin{array}{ccc} X & \xrightarrow{j} & \mathbf{V}(\mathcal{L}) \\ r \downarrow & & \downarrow q \\ P & \xrightarrow{\varepsilon_P} & C_P \end{array}$$

is commutative (since it is a particular case of (8.5.1.3)). The claims of (8.8.2) immediately follow from these facts, by taking g to be the composite morphism $h \circ q$. $\square$

Remark (8.8.3). — Assume further that Y is a Noetherian prescheme, and that f is a proper morphism. Since r is then proper (5.4.4), and thus closed, and since it is also a dominant open immersion, r is necessarily an isomorphism $X \xrightarrow{\sim} P$. Furthermore, we will see, in Chapter III (III, 2.3.5.1), that $\mathcal{S}$ is then necessarily an $\mathcal{O}_Y$ -algebra of finite type. It then follows that $\mathcal{S}^{\natural}$ is an $\mathcal{S}_0^{\natural}$ -algebra of finite

type ((8.2.10, i) and (8.7.2.7)); since C_P is C -isomorphic to $\text{Proj}(\mathcal{S}^{\natural})$ (8.7.3), we see that the morphism $h : C_P \rightarrow C$ is *projective*; since the morphism r is an isomorphism, so too is $q : \mathbf{V}(\mathcal{L}) \rightarrow C_P$, and we thus conclude that the morphism $g : \mathbf{V}(\mathcal{L}) \rightarrow C$ is *projective*. Furthermore, since the restriction of h to E_P is an isomorphism to E , and since q is an isomorphism, the restriction (8.8.2.3) of g is an isomorphism $\mathbf{V}(\mathcal{L}) - j(X) \xrightarrow{\sim} E$.

If we further assume that $\mathcal{L}$ is *very ample* for f , then, as we will also see in Chapter III (III, 2.3.5.1), there exists some integer $n_0 > 0$ such that $\mathcal{S}_n = \mathcal{S}_1^n$ for $n \geq n_0$. We then conclude, by (8.7.7), that $\mathbf{V}(\mathcal{L})$ can be identified with the prescheme Z given by *blowing up the vertex prescheme* (identified with Y) in the affine cone C , and that the zero section of $\mathbf{V}(\mathcal{L})$ (identified with Y) is the *inverse image* of the vertex subprescheme Y of C .

Some of the above results can in fact be proven even without the Noetherian hypothesis:

Corollary (8.8.4). — *Let Y be a prescheme (resp. a quasi-compact scheme), $f : X \rightarrow Y$ a proper morphism, and $\mathcal{L}$ an invertible $\mathcal{O}_X$ -module that is ample relative to f . Then the morphism in (8.8.2.1) is proper (resp. projective), and its restriction (8.8.2.3) is an isomorphism.*

Proof. To prove that g is proper, we can restrict to the case where Y is affine, and it then suffices to consider the case where Y is a quasi-compact scheme. The same arguments as in (8.8.3) first of all show that r is an isomorphism $X \xrightarrow{\sim} P$; then q is also an isomorphism, and, since the restriction of h to E_P is an isomorphism $E_P \xrightarrow{\sim} E$, we have already seen that (8.8.2.3) is an isomorphism. It remains only to prove that g is *projective*.

Since f is of finite type, by hypothesis, we can apply (3.8.5) to the homomorphism [II:180] τ from (8.8.1.2): there is an integer $d > 0$ and a quasi-coherent $\mathcal{O}_Y$ -submodule $\mathcal{E}$ of finite type of $\mathcal{S}_d$ such that, if $\mathcal{S}'$ is the $\mathcal{O}_Y$ -subalgebra of $\mathcal{S}$ generated by $\mathcal{E}$, and $\tau' = \tau \circ f^*(\phi)$ (where ϕ is the canonical injection $\mathcal{S}' \rightarrow \mathcal{S}$), then $r' = r_{\mathcal{L}, \tau'}$ is an immersion

$$X \longrightarrow P' = \text{Proj}(\mathcal{S}').$$

Furthermore, since ϕ is injective, r' is also a *dominant immersion* (3.7.6); the same argument as for r then shows that r' is a *surjective closed immersion*; since r' factors as $X \xrightarrow{r} \text{Proj}(\mathcal{S}) \xrightarrow{\Phi} \text{Proj}(\mathcal{S}')$, where $\Phi = \text{Proj}(\phi)$, we thus conclude that Φ is also a *surjective closed immersion*. But this implies that Φ is an *isomorphism*; we can restrict to the case where $Y = \text{Spec}(A)$ is affine, and $\mathcal{S} = \tilde{S}$ and $\mathcal{S}' = \tilde{S}'$, with S a graded A -algebra and S' a graded subalgebra of S . For every homogeneous element $t \in S'$, we have that $S'_{(t)}$ is a subring of $S_{(t)}$; if we return to the definition of $\text{Proj}(\phi)$ (2.8.1), we see that it suffices to prove that, if B' is a subring of a ring B , and if the morphism $\text{Spec}(B) \rightarrow \text{Spec}(B')$ corresponding to the canonical injection $B' \rightarrow B$ is a closed immersion, then this morphism is necessarily an *isomorphism*; but this follows from (I, 4.2.3). Furthermore, $\Phi^*(\mathcal{O}_{P'}(n)) = \mathcal{O}_P(n)$ ((3.5.2, ii) and (3.5.4)), and so $r'^*(\mathcal{O}_{P'}(n))$ is isomorphic to $\mathcal{L}^{\otimes n}$ (4.6.3). Let $\mathcal{S}'' = \mathcal{S}'^{(d)}$, so that (3.1.8, i) X is canonically identified with $P'' = \text{Proj}(\mathcal{S}'')$, and $\mathcal{L}'' = \mathcal{L}^{\otimes d}$ with $\mathcal{O}_{P''}(1)$ (3.2.9, ii).

Now, if $C'' = \text{Spec}(\mathcal{S}'')$, then $\mathcal{S}_{P''}^{\geq} = \bigoplus_{n \geq 0} \mathcal{O}_{P''}(n)$ can be identified with $\bigoplus_{n \geq 0} \mathcal{L}''^{\otimes n}$, and thus $C_{P''} = \text{Spec}(\mathcal{S}_{P''}^{\geq})$ with $\mathbf{V}(\mathcal{L}'')$; we also know (8.7.3) that $C_{P''}$ is C'' -isomorphic to $\text{Proj}(\mathcal{S}''^{\natural})$; by the definition of $\mathcal{S}''$, we know that $\mathcal{S}''^{\natural}$ is generated by $\mathcal{S}_1''^{\natural}$, and that $\mathcal{S}_1''^{\natural}$ is of finite type over $\mathcal{S}_0''^{\natural} = \mathcal{S}''$ ((8.2.10, i and iii)), and so $\text{Proj}(\mathcal{S}''^{\natural})$ is *projective* over C'' (5.5.1). Consider the diagram

$$(8.8.4.1) \quad \begin{array}{ccc} \mathbf{V}(\mathcal{L}) & \xrightarrow{g} & \text{Spec}(\mathcal{S}) = C \\ u \downarrow & & \downarrow v \\ \mathbf{V}(\mathcal{L}'') & \xrightarrow{g''} & \text{Spec}(\mathcal{S}'') = C'' \end{array}$$

where g and g'' correspond, by (1.5.6), to the canonical f -morphisms

$$\mathcal{S} \longrightarrow \bigoplus_{n \geq 0} \mathcal{L}^{\otimes n} \quad \text{and} \quad \mathcal{S}'' \longrightarrow \bigoplus_{n \geq 0} \mathcal{L}''^{\otimes n}$$

(3.3.2.3) (see (8.8.5) below), and v and u to the inclusion morphisms $\mathcal{S}'' \rightarrow \mathcal{S}$ and $\bigoplus_{n \geq 0} \mathcal{L}^{\otimes nd} \rightarrow \bigoplus_{n \geq 0} \mathcal{L}^{\otimes n}$ (respectively); it is immediate (3.3.2) that this diagram is commutative. We have just seen that g'' is a *projective* morphism; we also know that u is a *finite* morphism. Since the question

is local on X , we can assume that X is affine of ring A , and that $\mathcal{L} = \mathcal{O}_X$; everything then reduces to noting that the ring $A[T]$ is a module of finite type over its subring $A[T^d]$ (with T an indeterminate). Since Y is a quasi-compact scheme, and since C'' is affine over Y , we know that C'' is also a quasi-compact scheme, [III:181] and so $g'' \circ u$ is a projective morphism (5.5.5, ii); by commutativity of (8.8.4.1), $v \circ g$ is also projective, and, since v is affine, thus separated, we finally conclude that g is projective (5.5.5, v). $\square$

(8.8.5). Consider again the situation in (8.8.1). We will see that the morphism $g : V(\mathcal{L}) \rightarrow C$ can also be defined in a way that works for any invertible (but not necessarily ample) $\mathcal{O}_X$ -module $\mathcal{L}$. For this, consider the f -morphism

$$(8.8.5.1) \quad \tau^\flat : \mathcal{S} \longrightarrow \bigoplus_{n \geq 0} \mathcal{L}^{\otimes n}$$

corresponding to the morphism τ from (8.8.1.2). This induces (1.5.6) a morphism $g' : V \rightarrow C$ such that, if $\pi : V \rightarrow X$ and $\psi : C \rightarrow Y$ are the structure morphisms, the diagrams

$$(8.8.5.2) \quad \begin{array}{ccc} X & \xleftarrow{\pi} & V \\ f \downarrow & & \downarrow g' \\ Y & \xleftarrow{\psi} & C \end{array} \quad \begin{array}{ccc} X & \xrightarrow{j} & V \\ f \downarrow & & \downarrow g' \\ Y & \xrightarrow{\varepsilon} & C \end{array}$$

commute ((8.5.1.2) and (8.5.1.3)). We will show that (if we assume that $\mathcal{L}$ is f -ample) *the morphisms g and g' are identical*.

Since the question is local on Y , we can assume that $Y = \text{Spec}(A)$ is affine, and (by (8.8.1.3)) identify X with an open subset of $P = \text{Proj}(S)$, where $S = A \oplus \bigoplus_{n \geq 1} \Gamma(X, \mathcal{L}^{\otimes n})$; we then deduce, by (8.8.1.4), that $\Gamma(X, \mathcal{O}_P(n)) = \Gamma(X, \mathcal{L}^{\otimes n})$ for all $n \in \mathbb{Z}$. Taking into account the definition of $h = \text{Spec}(\alpha)$, where α is the canonical p -morphism $\tilde{S} \rightarrow \mathcal{S}_P^{\geq}$ (8.6.1.2), we have to show that the restriction to X of $\alpha^\sharp : p^*(\tilde{S}) \rightarrow \mathcal{S}_P^{\geq}$ is identical to τ . Taking (0, 4.4.3) into account, it suffices to show that, if we compose the canonical homomorphism $\alpha_n : S_n \rightarrow \Gamma(P, \mathcal{O}_P(n))$ with the restriction homomorphism $\Gamma(P, \mathcal{O}_P(n)) \rightarrow \Gamma(X, \mathcal{O}_P(n)) = \Gamma(X, \mathcal{L}^{\otimes n})$, then we obtain the identity, for all $n > 0$; but this follows immediately from the definition of the algebra S and of α_n (2.6.2).

Proposition (8.8.6). — Assume (with the notation of (8.8.5)) that, if we write $f = (f_0, \lambda)$, then the homomorphism $\lambda : \mathcal{O}_Y \rightarrow f_*(\mathcal{O}_X)$ is bijective; then:

- (i) if we write $g = (g_0, \mu)$, then $\mu : \mathcal{O}_C \rightarrow g_*(\mathcal{O}_V)$ is an isomorphism; and
- (ii) if X is integral (resp. locally integral and normal), then C is integral (resp. normal).

Proof. Indeed, the f -morphism $\tau^\flat$ is then an isomorphism

$$\tau^\flat : \mathcal{S} = \psi_*(\mathcal{O}_C) \longrightarrow f_*(\pi_*(\mathcal{O}_V)) = \psi_*(g_*(\mathcal{O}_V))$$

and the Y -morphism g can be considered as that for which the homomorphism $\mathcal{A}(g)$ (1.1.2) is equal to $\tau^\flat$. To see that μ is an isomorphism of $\mathcal{O}_C$ -modules, it suffices (1.4.2) to see that $\mathcal{A}(\mu) : \psi_*(\mathcal{O}_C) \rightarrow \psi_*(g_*(\mathcal{O}_V))$ is an isomorphism. But, by Definition (1.1.2), we have that $\mathcal{A}(\mu) = \mathcal{A}(g)$, whence the conclusion of (i).

To prove (ii), we can restrict to the case where Y is affine, and so $\mathcal{S} = \tilde{S}$, with [III:182] $S = \bigoplus_{n \geq 0} \Gamma(X, \mathcal{L}^{\otimes n})$; the hypothesis that X is integral implies that the ring S is integral (I, 7.4.4), and thus so too is C (I, 5.1.4). To show that C is normal, we will use the following lemma:

Lemma (8.8.6.1). — Let Z be a normal integral prescheme. Then the ring $\Gamma(Z, \mathcal{O}_Z)$ is integral and integrally closed.

Proof. It follows from (I, 8.2.1.1) that $\Gamma(Z, \mathcal{O}_Z)$ is the intersection, in the field of rational functions $R(Z)$, of the integrally closed rings $\mathcal{O}_z$ over all $z \in Z$. $\square$

With this in mind, we first show that V is *locally integral* and *normal*; for this, we can restrict to the case where $X = \text{Spec}(A)$ is affine, with ring A integral and integrally closed (6.3.8), and where $\mathcal{L} = \mathcal{O}_X$. Since then $V = \text{Spec}(A[T])$, and $A[T]$ is integral and integrally closed [24, p. 99], this proves our claim. For every affine open subset U of C , $g^{-1}(U)$ is quasi-compact, since the

morphism g is quasi-compact; since V is locally integral, the connected components of $g^{-1}(U)$ are open integral preschemes in $g^{-1}(U)$, and thus finite in number, and, since V is normal, these preschemes are also normal (6.3.8). Then $\Gamma(U, \mathcal{O}_C)$, which is equal to $\Gamma(g^{-1}(U), \mathcal{O}_V)$, by (i), is the finite direct product of integral and integrally closed rings (8.8.6.1), which proves that C is normal (6.3.4). $\square$

8.9. Grauert's ampleness criterion: statement We intend to show that the properties proven in (8.8.2) characterise f -ample $\mathcal{O}_X$ -modules, and, more precisely, to prove the following criterion:

Theorem (8.9.1). — (Grauert's criterion). *Let Y be a prescheme, $p : X \rightarrow Y$ a separated and quasi-compact morphism, and $\mathcal{L}$ an invertible $\mathcal{O}_X$ -module. For $\mathcal{L}$ to be ample relative to p , it is necessary and sufficient for there to exist a Y -prescheme C , a Y -section $\varepsilon : Y \rightarrow C$ of C , and a Y -morphism $q : \mathbf{V}(\mathcal{L}) \rightarrow C$, satisfying the following properties:*

(i) *the diagram*

$$(8.9.1.1) \quad \begin{array}{ccc} X & \xrightarrow{j} & \mathbf{V}(\mathcal{L}) \\ p \downarrow & & \downarrow q \\ Y & \xrightarrow{\varepsilon} & C \end{array}$$

commutes, where j is the zero section of the vector bundle $\mathbf{V}(\mathcal{L})$; and

(ii) *the restriction of q to $\mathbf{V}(\mathcal{L}) - j(X)$ is a quasi-compact open immersion*

$$\mathbf{V}(\mathcal{L}) - j(X) \longrightarrow C$$

whose image does not intersect $\varepsilon(Y)$.

Note that, if C is separated over Y , we can, in condition (ii), remove the hypothesis that the open immersion is quasi-compact; to see that this property (of quasi-compactness) is in fact a consequence of the other conditions, we can restrict to the case where Y is affine, and the claim then follows from (I, 5.5.1, i) and (I, 5.5.10). We can also remove [II:183] the same hypothesis if we assume that X is Noetherian, since then V is also Noetherian, and the claim follows from (I, 6.3.5).

Corollary (8.9.2). — *If the morphism $p : X \rightarrow Y$ is proper, then we can, in the statement of Theorem (8.9.1), assume that q is proper, and replace “open immersion” by “isomorphism”.*

In a more suggestive manner, we can say (whenever $p : X \rightarrow Y$ is proper) that $\mathcal{L}$ is ample relative to p if and only if we can “contract” the zero section of the vector bundle $\mathbf{V}(\mathcal{L})$ to the base prescheme Y . An important particular case is that where Y is the spectrum of a field, and where the operation of “contraction” consists in contracting the zero section of $\mathbf{V}(\mathcal{L})$ to a single point.

(8.9.3). The necessity of the conditions in Theorem (8.9.1) and Corollary (8.9.2) follow immediately from (8.8.2) and (8.8.4).

To show that the conditions of (8.9.1) suffice, consider a slightly more general situation. For this, let (with the notation of (8.8.2))

$$\mathcal{S}' = \bigoplus_{n \geq 0} \mathcal{L}^{\otimes n}$$

and

$$V = \mathbf{V}(\mathcal{L}) = \text{Spec}(\mathcal{S}').$$

The closed subprescheme $j(X)$, the zero section of $\mathbf{V}(\mathcal{L})$, is defined by the quasi-coherent sheaf of ideals $\mathcal{J} = (\mathcal{S}'_+)^{\sim}$ of $\mathcal{O}_V$ (1.4.10). This $\mathcal{O}_V$ -module is invertible, since this property is local on X , and this reduces to remarking that the ideal $TA[T]$ in a ring of polynomials $A[T]$ is a free cyclic $A[T]$ -module. Furthermore, it is immediate (again, because the question is local on X) that

$$\mathcal{L} = j^*(\mathcal{J})$$

and

$$j_*(\mathcal{L}) = \mathcal{J} / \mathcal{J}^2.$$

Now, if

$$\pi : \mathbf{V}(\mathcal{L}) \longrightarrow X$$

is the structure morphism, then $\pi_*(\mathcal{J}) = \mathcal{J}'_+$ and $\pi_*(\mathcal{J} / \mathcal{J}^2) = \mathcal{L}$; there are thus canonical homomorphisms $\mathcal{L} \rightarrow \pi_*(\mathcal{J}) \rightarrow \mathcal{L}$, the first being the canonical injection $\mathcal{L} \rightarrow \mathcal{J}'_+$, and the second the canonical projection from $\mathcal{J}'_+$ to $\mathcal{J}'_1 = \mathcal{L}$, and their composition being the identity. We can also canonically embed $\pi_*(\mathcal{J}) = \mathcal{J}'_+ = \bigoplus_{n \geq 1} \mathcal{L}^{\otimes n}$ into the product $\prod_{n \geq 1} \mathcal{L}^{\otimes n} = \varprojlim_n \pi_*(\mathcal{J} / \mathcal{J}^{n+1})$ (since $\pi_*(\mathcal{J} / \mathcal{J}^{n+1}) = \mathcal{L} \oplus \mathcal{L}^{\otimes 2} \oplus \dots \oplus \mathcal{L}^{\otimes n}$), and we thus have canonical homomorphisms

$$(8.9.3.1) \quad \mathcal{L} \longrightarrow \varprojlim \pi_*(\mathcal{J} / \mathcal{J}^{n+1}) \longrightarrow \mathcal{L}$$

whose composition is the identity.

With this in mind, the generalisation of (8.9.1) that we are going to prove is the following:

Proposition (8.9.4). — *Let Y be a prescheme, V a Y -prescheme, and X a closed subprescheme of V defined by an ideal $\mathcal{J}$ of $\mathcal{O}_V$, which is an invertible $\mathcal{O}_V$ -module; if $j : X \rightarrow V$ is ^[III:184] the canonical injection, then let $\mathcal{L} = j^*(\mathcal{J}) = \mathcal{J} \otimes_{\mathcal{O}_V} \mathcal{O}_X$, so that $j_*(\mathcal{L}) = \mathcal{J} / \mathcal{J}^2$. Assume that the structure morphism $p : X \rightarrow Y$ is separated and quasi-compact, and that the following conditions are satisfied:*

- (i) *there exists a Y -morphism $\pi : V \rightarrow X$ of finite type such that $\pi \circ j = 1_X$, and so $\pi_*(\mathcal{J} / \mathcal{J}^2) = \mathcal{L}$;*
- (ii) *there exists a homomorphism of $\mathcal{O}_X$ -modules $\phi : \mathcal{L} \rightarrow \varprojlim \pi_*(\mathcal{J} / \mathcal{J}^{n+1})$ such that the composition*

$$\mathcal{L} \xrightarrow{\phi} \varprojlim \pi_*(\mathcal{J} / \mathcal{J}^{n+1}) \xrightarrow{\alpha} \pi_*(\mathcal{J} / \mathcal{J}^2) = \mathcal{L}$$

(where α is the canonical homomorphism) is the identity;

- (iii) *there exists a Y -prescheme C , a Y -section ε of C , and a Y -morphism $q : V \rightarrow C$ such that the diagram*

$$(8.9.4.1) \quad \begin{array}{ccc} X & \xrightarrow{j} & V \\ p \downarrow & & \downarrow q \\ Y & \xrightarrow{\varepsilon} & C \end{array}$$

commutes; and

- (iv) *the restriction of q to $W = V - j(X)$ is a quasi-compact open immersion into C , whose image does not intersect $\varepsilon(Y)$.*

Then $\mathcal{L}$ is ample relative to p .

8.10. Grauert's ampleness criterion: proof

Lemma (8.10.1). — *Let $\pi : V \rightarrow X$ be a morphism, $j : X \rightarrow V$ an X -section of V that is also a closed immersion, and $\mathcal{J}$ a quasi-coherent ideal of $\mathcal{O}_V$ that defines the closed subprescheme of V associated to j . Then the following all hold true.*

- (i) *For all $n \geq 0$, $\pi_*(\mathcal{O}_V / \mathcal{J}^{n+1})$ and $\pi_*(\mathcal{J} / \mathcal{J}^{n+1})$ are quasi-coherent $\mathcal{O}_X$ -modules, and $\pi_*(\mathcal{O}_V / \mathcal{J}) = \mathcal{O}_X$ and $\pi_*(\mathcal{J} / \mathcal{J}^2) = j^*(\mathcal{J})$.*
- (ii) *If $X = \{\xi\} = \text{Spec}(k)$, where k is a field, then $\varprojlim \pi_*(\mathcal{O}_V / \mathcal{J}^{n+1})$ is isomorphic to the separated completion of the local ring $\mathcal{O}_{j(\xi)}$ for the $\mathfrak{m}_{j(\xi)}$ -adic topology.*
- (iii) *Assume that $\mathcal{J}$ is an invertible $\mathcal{O}_V$ -module (which implies that*

$$\mathcal{L} = j^*(\mathcal{J}) = \pi_*(\mathcal{J} / \mathcal{J}^2)$$

is an invertible $\mathcal{O}_X$ -module), and that there exists a homomorphism $\phi : \mathcal{L} \rightarrow \varprojlim \pi_(\mathcal{J} / \mathcal{J}^{n+1})$ such that the composition $\mathcal{L} \xrightarrow{\phi} \varprojlim \pi_*(\mathcal{J} / \mathcal{J}^{n+1}) \xrightarrow{\alpha} \pi_*(\mathcal{J} / \mathcal{J}^2)$ (where α is the canonical homomorphism) is the identity. If we write $\mathcal{S} = \bigoplus_{n \geq 0} \mathcal{L}^{\otimes n}$, then ϕ canonically induces an isomorphism of $\mathcal{O}_X$ -algebras from the completion $\widehat{\mathcal{S}}$ of $\mathcal{S}$ relative to its canonical filtration (the completion being isomorphic to the product $\prod_{n \geq 0} \mathcal{L}^{\otimes n}$) to $\varprojlim \pi_*(\mathcal{O}_V / \mathcal{J}^{n+1})$.*

Proof. Note first of all that the support of the $\mathcal{O}_V$ -module $\mathcal{O}_V / \mathcal{J}^{n+1}$ is $j(X)$, and the support of $\mathcal{J} / \mathcal{J}^{n+1}$ is contained in $j(X)$. In the case of (ii), $j(X)$ is a closed point $j(\xi)$ of V , ^[III:185] and, by definition, $\pi_*(\mathcal{O}_V / \mathcal{J}^{n+1})$ is the fibre of $\mathcal{O}_V / \mathcal{J}^{n+1}$ at the point $j(\xi)$, or, equivalently, setting $C = \mathcal{O}_{j(\xi)}$, and denoting by $\mathfrak{m}$ the maximal ideal of C , the C -module $C / \mathfrak{m}^{n+1}$; claim (ii) is then evident.

To prove (i), note that the question is local on X ; we can thus restrict to the case where X is affine. Let U be an affine open subset of V ; then $j(X) \cap U$ is an affine open subset of $j(X)$, so $U_0 = \pi(j(X) \cap U)$, which is isomorphic to it, is an affine open subset of X ; for every affine open subset $W_0 \subset U_0$ in X , $W = \pi^{-1}(W_0) \cap U$ is an affine open subset of V , since X is a scheme (I, 5.5.10); in particular, $U' = U \cap \pi^{-1}(U_0)$ is an affine open subset of V , and clearly $\pi(U') = U_0$ and $j(U_0) = j(X) \cap U$. Then, by definition, $\Gamma(W_0, \pi_*(\mathcal{O}_V / \mathcal{I}^{n+1})) = \Gamma(\pi^{-1}(W_0), \mathcal{O}_V / \mathcal{I}^{n+1})$; but since every point of $\pi^{-1}(W_0)$ not belonging to $j(W_0)$ has an open neighbourhood in $\pi^{-1}(W_0)$ not intersecting $j(X)$, and in which $\mathcal{O}_V / \mathcal{I}^{n+1}$ is thus zero, it is clear that the sections of $\mathcal{O}_V / \mathcal{I}^{n+1}$ over $\pi^{-1}(W_0)$ and over W are in bijective correspondence. In other words, if π' is the restriction of π to U' , then the $(\mathcal{O}_X|_{U_0})$ -modules $\pi_*(\mathcal{O}_V / \mathcal{I}^{n+1})|_{U_0}$ and $\pi'_*((\mathcal{O}_V / \mathcal{I}^{n+1})|_{U'})$ are identical. Since U' and U_0 are affine, and since the U_0 cover X , we thus conclude (I, 1.6.3) that $\pi_*(\mathcal{O}_V / \mathcal{I}^{n+1})$ is quasi-coherent, and the proof is identical for $\pi_*(\mathcal{I} / \mathcal{I}^{n+1})$.

Finally, to prove (iii), note that $\mathcal{S}$ is exactly $\mathbf{S}_{\mathcal{O}_X}(\mathcal{L})$; so ϕ canonically induces a homomorphism of $\mathcal{O}_X$ -algebras $\psi : \mathcal{S} \rightarrow \varprojlim \pi_*(\mathcal{O}_V / \mathcal{I}^{n+1})$ (1.7.4); furthermore, this homomorphism sends $\mathcal{L}^{\otimes n}$ to $\varprojlim_m \pi_*(\mathcal{I}^n / \mathcal{I}^{m+1})$, and is thus continuous for the topologies considered, and indeed then extends to a homomorphism $\hat{\psi} : \hat{\mathcal{S}} \rightarrow \varprojlim \pi_*(\mathcal{O}_V / \mathcal{I}^{n+1})$. To see that this is indeed an isomorphism, we can, as in the proof of (i), restrict to the case where $X = \text{Spec}(A)$ and $V = \text{Spec}(B)$ are affine, with $\mathcal{I} = \tilde{\mathfrak{J}}$, where $\mathfrak{J}$ is an ideal of B ; there is an injection $A \rightarrow B$ corresponding to π that identifies A with a subring of B complementary to $\mathfrak{J}$, and $\mathcal{L}$ (resp. $\pi_*(\mathcal{O}_V / \mathcal{I}^{n+1})$) is the quasi-coherent $\mathcal{O}_X$ -module associated to the A -module $L = \mathfrak{J} / \mathfrak{J}^2$ (resp. $B / \mathfrak{J}^{n+1}$). Since $\mathcal{I}$ is an invertible $\mathcal{O}_V$ -module, we can further assume that $\mathfrak{J} = Bt$, where t is not a zero divisor in B . From the fact that $B = A \oplus Bt$, we deduce that, for all $n > 0$,

$$B = A \oplus At \oplus At^2 \oplus \dots \oplus At^n \oplus Bt^{n+1}$$

and so there exists a canonical A -isomorphism from the ring of formal series $A[[T]]$ to $C = \varprojlim B / \mathfrak{J}^{n+1}$ that sends T to t . We also have that $L = A\bar{t}$, where $\bar{t}$ is the class of t modulo Bt^2 , and the homomorphism ϕ sends, by hypothesis, $\bar{t}$ to an element $t' \in C$ that is congruent to t modulo Ct^2 . We thus deduce, by induction on n , that

$$A \oplus At' \oplus \dots \oplus A(t')^n \oplus Ct^{n+1} = A \oplus At \oplus \dots \oplus At^n \oplus Ct^{n+1}$$

which proves that the homomorphism $\hat{\psi}$ does indeed correspond to an isomorphism from $\prod_{n \geq 0} L^{\otimes n}$ to C . $\square$

Lemma (8.10.2). — *Under the hypotheses of Lemma (8.10.1), let $g : X' \rightarrow X$ be a morphism, ^[II:186] write $V' = V \times_X X'$, and let $\pi' : V' \rightarrow X'$ and $g' : V' \rightarrow V$ be the canonical projections, so that we have the commutative diagram*

$$\begin{array}{ccc} V & \xleftarrow{g'} & V' \\ \pi \downarrow & & \downarrow \pi' \\ X & \xleftarrow{g} & X' \end{array}$$

Then $j' = j \times 1_{X'}$ is an X' -section of V' that is also a closed immersion, and $\mathcal{I}' = g'^(\mathcal{I})\mathcal{O}_{V'}$ is the quasi-coherent ideal of $\mathcal{O}_{V'}$ that defines the closed subscheme of V' associated to j' . Furthermore, $\pi'_*(\mathcal{O}_{V'} / \mathcal{I}'^{n+1}) = g^*(\pi_*(\mathcal{O}_V / \mathcal{I}^{n+1}))$. Finally, $\mathcal{I}'$ is an $\mathcal{O}_{V'}$ -module that is canonically isomorphic to $g'^*(\mathcal{I})$, and is, in particular, invertible if $\mathcal{I}$ is an invertible $\mathcal{O}_V$ -module.*

Proof. The fact that j' is a closed immersion follows from (I, 4.3.1), and it is an X' -section of V' by functoriality of extension of the base prescheme. Furthermore, if Z (resp. Z') is the closed subscheme of V (resp. V') associated to j (resp. j'), then $Z' = g'^{-1}(Z)$ (I, 4.3.1), and the second claim then follows from (I, 4.4.5). To prove the other claims, we see, as in (8.10.1), that we can restrict to the case where X , V , and X' (and thus also V') are affine; we keep the notation from the proof of (8.10.1), and let $X' = \text{Spec}(A')$. Then $V' = \text{Spec}(B')$, where $B' = B \otimes_A A'$, and $\mathcal{I}' = \tilde{\mathfrak{J}}'$, where $\mathfrak{J}' = \text{Im}(\mathfrak{J} \otimes_A A')$. Then $B' / \mathfrak{J}'^{n+1} = (B / \mathfrak{J}^{n+1}) \otimes_A A'$; furthermore, since $\mathfrak{J}$ is a direct factor

(as an A -module) of B , $\mathfrak{J} \otimes_A A'$ is a direct factor (as an A' -module) of B' , and is thus canonically identified with $\mathfrak{J}'$. $\square$

Corollary (8.10.3). — Assume that the hypotheses of Lemma (8.10.1) are satisfied, and assume further that π is of finite type, and that $\mathcal{J}$ is an invertible $\mathcal{O}_V$ -module. Then, for all $x \in X$, the local ring at the point $j(x)$ of the fibre $\pi^{-1}(x)$ is a regular (thus integral) ring of dimension 1, whose completion is isomorphic to the formal series ring $k(x)[[T]]$ (where T is an indeterminate); furthermore, there exists exactly one irreducible component of $\pi^{-1}(x)$ that contains $j(x)$.

Proof. Since $\pi^{-1}(x) = V \times_X \text{Spec}(k(x))$, we are led, by (8.10.2), to the case where X is the spectrum of a field K . Since π is of finite type (I, 6.3.4, iv), $\mathcal{O}_{j(x)}$ is a Noetherian local ring, and thus separated for the $\mathfrak{m}_{j(x)}$ -adic topology (0, 7.3.5); it follows from (8.10.1, ii and iii) that the completion of this ring is isomorphic to $K[[T]]$, and so $\mathcal{O}_{j(x)}$ is regular and of dimension 1 ([1, p. 17-01, th. 1]); finally, since $\mathcal{O}_{j(x)}$ is integral, $j(x)$ belongs to exactly one of the (finitely many) irreducible components of $\pi^{-1}(x)$ (I, 5.1.4). $\square$

Corollary (8.10.4). — Suppose that the hypotheses of Lemma (8.10.1) are satisfied, and assume further that $\mathcal{J}$ is an invertible $\mathcal{O}_V$ -module. Let $W = V - j(X)$; for every quasi-coherent ideal $\mathcal{K}$ of $\mathcal{O}_X$, let $\mathcal{K}_V = \pi^*(\mathcal{K})\mathcal{O}_V$ and $\mathcal{K}_W = \mathcal{K}_V|_W$. Then $\mathcal{K}_V$ is the largest quasi-coherent ideal of $\mathcal{O}_V$ whose restriction to W is $\mathcal{K}_W$.

Proof. Indeed, we see as in (8.10.1) that the question is local on X and V ; we can thus reuse the notation from the proof of (8.10.1), with $\mathfrak{J} = Bt$, where t is not a zero divisor in B . Furthermore, we have $W = \text{Spec}(B_t)$ and $\mathcal{K} = \tilde{\mathfrak{K}}$, where $\mathfrak{K}$ is an ideal of A ; whence $\pi^*(\mathcal{K})\mathcal{O}_V = (\mathfrak{K}.B)^\sim$ (I, 1.6.9), $\mathcal{K}_W = (\mathfrak{K}.B_t)^\sim$, and the largest ideal ^[III:187] of B whose canonical image in B_t is $\mathfrak{K}.B_t$ is the inverse image of $\mathfrak{K}.B_t$, that is, the set of $s \in B$ such that, for some integer $n > 0$, we have $t^n s \in \mathfrak{K}.B$. We have to show that this last relation implies that $s \in \mathfrak{K}.B$, or again that the canonical image of t is not a zero divisor in $B/\mathfrak{K}B = (A/\mathfrak{K}) \otimes_A B$, which follows from (8.10.2) applied to $X' = \text{Spec}(A/\mathfrak{K})$. $\square$

Corollary (8.10.5). — Suppose that the hypotheses of (8.10.3) are satisfied; let $W = V - j(X)$, x be a point of X , $\mathcal{K}$ a quasi-coherent ideal of $\mathcal{O}_X$, and z the generic point of the irreducible component of $\pi^{-1}(x)$ that contains $j(x)$ (8.10.3).

- (i) Let g be a section of $\mathcal{O}_V$ over V such that $g|_W$ is a section of $\mathcal{K}_W$ over W (using the notation from (8.10.4)). Then g is a section of $\mathcal{K}_V$; if further $g(z) \neq 0$, and if, for every integer $m > 0$, we denote by g_m^x the germ at the point x of the canonical image g_m of g in $\Gamma(X, \pi_*(\mathcal{O}_V/\mathcal{J}^{m+1}))$, then there exists an integer $m > 0$ such that the image of g_m^x in

$$(\pi_*(\mathcal{O}_V/\mathcal{J}^{m+1}))_x \otimes_{\mathcal{O}_x} k(x)$$

is $\neq 0$.

- (ii) Suppose further that the conditions of (8.10.1, iii) are fulfilled. Then, if there exists a section g of $\mathcal{K}_V$ over V such that $g(z) \neq 0$, then there exists an integer $n \geq 0$ and a section f of $\mathcal{K}.\mathcal{L}^{\otimes n} = \mathcal{K} \otimes \mathcal{L}^{\otimes n} \subset \mathcal{L}^{\otimes n}$ such that $f(x) \neq 0$. If g is a section of $\mathcal{J}$, we can take $n > 0$.

Proof.

- (i) Since the ideal of $\mathcal{O}_W$ generated by $g|_W$ is contained in $\mathcal{K}_W$ by hypothesis, the ideal of $\mathcal{O}_V$ generated by g is contained in $\mathcal{K}_V$ by (8.10.4), or, in other words, g is a section of $\mathcal{K}_V$. To prove the second claim of (i), we can again assume that X and V are affine, and reuse the notation from (8.10.1); the fibre $\pi^{-1}(x)$ is then affine of ring $B' = B \otimes_A k(x)$, and there exists in B' an element t' which is not a zero divisor and is such that $B' = k(x) \oplus B't'$. Since $j(x)$ is a specialisation of z and since $g(z) \neq 0$, we necessarily have that $g_{j(x)} \neq 0$. But $\mathcal{O}_{j(x)}$ is a separated local ring (8.10.3), and thus embeds into its completion, and the image of g in this completion is thus not null. But this completion is isomorphic to $\varprojlim_n (B'/B't'^{n+1})$ (8.10.3); if $g' = g \otimes 1 \in B'$, there then exists an integer m such that $g' \notin B't'^{m+1}$, or, again, the image g'_m of g' in $B'/B't'^{m+1}$ is nonzero. But since g'_m is exactly the image of g_m^x , our claim is proved.

- (ii) By (8.10.1, iii), $\pi_*(\mathcal{O}_V / \mathcal{I}^{m+1})$ is isomorphic to the direct sum of the $\mathcal{L}^{\otimes k}$ for $0 \leq k \leq m$; we denote by f_k the section of $\mathcal{L}^{\otimes k}$ over X that is the component of the element of $\bigoplus_{k=0}^m \Gamma(X, \mathcal{L}^{\otimes k})$ which corresponds to g_m by this isomorphism. Choosing m as in (i), there is thus an index k such that $f_k(x) \neq 0$, by (i). To see that f_k is a section of $\mathcal{K} \mathcal{L}^{\otimes k}$, it suffices to consider, as above, the case where X and V are affine, and this follows immediately from the fact that $g \in \mathfrak{K}.B$ (with the notation from (8.10.4)). The final claim follows from the fact that the hypothesis $g \in \Gamma(V, \mathcal{I})$ implies that $f_0 = 0$.

□

(8.10.6). *Proof of (8.9.4).* The question is local on Y (4.6.4); since ε is a Y -section, we can thus replace C by an affine open neighbourhood U of a point of $\varepsilon(Y)$ such that $\varepsilon(Y) \cap U$ is closed in U . In other words, we can assume that C is affine, and that Y is a closed subscheme of C (and thus also affine) defined by a quasi-coherent sheaf $\mathcal{I}$ of ideals of $\mathcal{O}_C$. Since p is separated and quasi-compact, X is thus a quasi-compact scheme, and we are reduced to proving that $\mathcal{L}$ is ample (4.6.4). By criterion (4.5.2, a), we must thus prove the following: for every quasi-coherent ideal $\mathcal{K}$ of $\mathcal{O}_X$ and every point $x \in X$ not belonging to the support of $\mathcal{O}_X / \mathcal{K}$, there exists an integer $n > 0$ and a section f of $\mathcal{K} \otimes \mathcal{L}^{\otimes n}$ over X such that $f(x) \neq 0$.

For this, set

$$\begin{aligned}\mathcal{K}_V &= \pi^*(\mathcal{K})\mathcal{O}_V \\ \mathcal{K}_W &= \mathcal{K}_V|_W\end{aligned}$$

where $W = V - j(X)$; since the restriction of q to W is a quasi-compact open immersion into C , it follows from (I, 9.4.2) that $\mathcal{K}_W$ is the restriction to W of a quasi-coherent ideal $\mathcal{K}'_V$ of $\mathcal{O}_V$ of the form

$$\mathcal{K}'_V = q^*(\mathcal{K}_C)\mathcal{O}_V$$

where $\mathcal{K}_C$ is a quasi-coherent ideal of $\mathcal{O}_C$. Furthermore, since, by hypotheses, $q^{-1}(Y) \subset j(X)$, and since Y is defined by the ideal $\mathcal{I}$, the restriction to W of $q^*(\mathcal{I})\mathcal{O}_V$ is identical to that of $\mathcal{O}_V$, and so $\mathcal{K}_W$ is also the restriction to W of $q^*(\mathcal{I}\mathcal{K}_C)\mathcal{O}_V$, and we can thus suppose that $\mathcal{K}_C \subset \mathcal{I}$, whence

$$(8.10.6.1) \quad \mathcal{K}'_V \subset q^*(\mathcal{I})\mathcal{O}_V \subset \mathcal{I}$$

taking into account (I, 4.4.6) and the commutativity of (8.9.4.1). Furthermore, we deduce from (8.10.4) that

$$(8.10.6.2) \quad \mathcal{K}'_V \subset \mathcal{K}_V.$$

With this in mind, it follows from (8.10.3) that $j(x)$ belongs to exactly one irreducible component of $\pi^{-1}(x)$; let z be the generic point of this component, and let $z' = q(z)$. By (8.10.5), the proof will be finished (taking (8.10.6.1) and (8.10.6.2) into account) if we show the existence of a section g of $\mathcal{K}'_V$ over V such that $g(z) \neq 0$. But, by hypothesis, $\mathcal{K}$ has a restriction equal to that of $\mathcal{O}_X$ in an open neighbourhood of x ; also, it follows from (8.10.3) that $z \neq j(x)$, and so $z \in W$, and thus $(\mathcal{K}_W)_z = \mathcal{O}_{V,z}$, whence, by definition, $(\mathcal{K}_C)_{z'} = \mathcal{O}_{C,z'}$. Since C is affine, there is thus a section g' of $\mathcal{K}_C$ over C such that $g'(z') \neq 0$, and by taking g to be the section of $\mathcal{K}'_V$ corresponding canonically to g' , we indeed have $g(z) \neq 0$, which finishes the proof.

Remark (8.10.7). — We ignore the question of whether or not condition (ii) in (8.9.4) is superfluous or not. In any case, the conclusion does not hold if we do not assume the existence of a Y -morphism $\pi : V \rightarrow X$ such that $\pi \circ j = 1_X$; we briefly point out how we can indeed construct a counterexample, whose details will not be developed until later on. We take $Y = \text{Spec}(k)$, where k is a field, and $C = \text{Spec}(A)$, where $A = k[T_1, T_2]$, and the Y -section ε corresponding to the augmentation homomorphism $A \rightarrow k$. We denote by C' the scheme induced by C by blowing up the closed point $a = \varepsilon(Y)$ of C ; if D is the inverse image of a in C' , we consider in D a closed point b , ^[II:189] and we denote by V the scheme induced by C' by blowing up b ; X is the closed subscheme of V given by the inverse image of a by the structure morphism $q : V \rightarrow C$. We now show that X is the union of two irreducible components, X_1 and X_2 , where X_1 is the inverse image of b in V . It is immediate that the ideal $\mathcal{I}$ of $\mathcal{O}_V$ that defines X is again invertible, but we can show that $j^*(\mathcal{I}) = \mathcal{L}$ (where j is the canonical injection $X \rightarrow V$) is not ample, by considering the “degree” of the inverse image

of $\mathcal{L}$ in X_1 , which would be > 0 if $\mathcal{L}$ were ample, but which can be shown (by an elementary intersection calculation) to be equal to 0.

8.11. Uniqueness of contractions

Lemma (8.11.1). — *Let U and V be preschemes, and $h = (h_0, \lambda) : U \rightarrow V$ a surjective morphism. Suppose that*

- (1) $\lambda : \mathcal{O}_V \rightarrow h_*(\mathcal{O}_U) = (h_0)_*(\mathcal{O}_U)$ is an isomorphism;
- (2) *the underlying space of V can be identified with the quotient of the underlying space of U by the relation $h_0(x) = h_0(y)$ (a condition which always holds whenever the morphism h is open or closed, or, a fortiori when h is proper.)*

Then, for every prescheme W , the map

$$(8.11.1.1) \quad \text{Hom}(V, W) \longrightarrow \text{Hom}(U, W)$$

that, to each morphism $v = (v_0, \nu)$ from V to W , associates the morphism $u = v \circ h = (u_0, \mu)$, is a bijection from $\text{Hom}(V, W)$ to the set of u such that u_0 is constant on every fibre $h_0^{-1}(x)$.

Proof. It is clear that, if $u = v \circ h$, so that $u_0 = v_0 \circ h_0$, then u_0 is constant on every set $h_0^{-1}(x)$. Conversely, if u has this property, we will show that there exists exactly one $v \in \text{Hom}(V, W)$ such that $u = v \circ h$. The existence and uniqueness of the continuous map $v_0 : V \rightarrow W$ such that $u_0 = v_0 \circ h_0$ follows from the hypotheses, since h_0 can be identified with the canonical map from U to U/R . We can also, replacing V by some isomorphic prescheme if necessary, suppose that λ is the identity; by hypothesis, μ is then a homomorphism $\mu : \mathcal{O}_W \rightarrow (u_0)_*(\mathcal{O}_U) = (v_0)_*((h_0)_*(\mathcal{O}_U))$ such that the corresponding homomorphism $\mu^\# : u_0^*(\mathcal{O}_W) \rightarrow \mathcal{O}_U$ is local on every fibre. Since $(v_0)_*((h_0)_*(\mathcal{O}_U)) = (v_0)_*(\mathcal{O}_V)$, we necessarily have that $\nu = \mu$, and everything then reduces to showing that the corresponding homomorphism $\nu^\# : v_0^*(\mathcal{O}_W) \rightarrow \mathcal{O}_V$ is local on every fibre. But every $y \in V$ is of the form $h_0(x)$ for some $x \in U$; let $z = v_0(y) = u_0(x)$. Then (0, 3.5.5) the homomorphism $\mu_x^\#$ factors as

$$\mu_x^\# : \mathcal{O}_z \xrightarrow{\nu_y^\#} \mathcal{O}_y \xrightarrow{\lambda_x^\#} \mathcal{O}_x.$$

By hypothesis, $\lambda_x^\#$ and $\mu_x^\#$ are local homomorphisms; thus $\lambda_x^\#$ sends every invertible element of $\mathcal{O}_y$ to an invertible element of $\mathcal{O}_x$; if $\nu_y^\#$ sent a non-invertible element of $\mathcal{O}_z$ to an invertible element of $\mathcal{O}_y$, then $\mu_x^\#$ would send this element of $\mathcal{O}_z$ to an invertible element of $\mathcal{O}_x$, contradicting the hypothesis, whence the lemma. $\square$

Corollary (8.11.2). — *Let U be an integral prescheme, and V a normal prescheme; then every morphism $h : U \rightarrow V$ that is universally closed, birational, and radicial, is also an isomorphism.*

Proof. If $h = (h_0, \lambda)$, then it follows from the hypotheses that h_0 is injective and closed, and that $h_0(U)$ ^[II:190] is dense in V , and so h_0 is a homeomorphism from U to V . To prove the corollary, it will suffice to show that $\lambda : \mathcal{O}_V \rightarrow (h_0)_*(\mathcal{O}_U)$ is an isomorphism: we can then apply (8.11.1), which proves that the map (8.11.1.1) is bijective (the fibres $h_0^{-1}(x)$ each consisting of a single point); thus h will be an isomorphism. The question clearly being local on V , we can suppose that $V = \text{Spec}(A)$ is affine, of an integral and integrally closed ring (8.8.6.1); h then corresponds (I, 2.2.4) to a homomorphism $\phi : A \rightarrow \Gamma(U, \mathcal{O}_U)$, and everything reduces to showing that ϕ is an isomorphism. But, if K is the field of fractions of A , then $\Gamma(U, \mathcal{O}_U)$ has, by hypothesis, K as its field of fractions, and A is a subring of $\Gamma(U, \mathcal{O}_U)$, with ϕ being the canonical injection (I, 8.2.7). Since the morphism h satisfies the hypotheses of (7.3.11), $\Gamma(U, \mathcal{O}_U)$ is a subring of the integral closure of A in K , and is thus identical to A by hypothesis. $\square$

Remark (8.11.3). — We will see in chapter III (III, 4.4.11) that, whenever V is a *locally Noetherian* prescheme, every morphism $h : U \rightarrow V$ that is proper and quasi-finite (in particular, every morphism satisfying the hypotheses of (8.11.2)) is necessarily *finite*. The conclusion of (8.11.2) then follows in this case from (6.1.15).

(8.11.4). We will now see that, in Grauert's criterion, we can often prove that the prescheme C and the "contraction" q are determined in an *essentially unique* manner.

Lemma (8.11.5). — *Let Y be a prescheme, $p : X \rightarrow Y$ a proper morphism, $\mathcal{L}$ a p -ample invertible $\mathcal{O}_X$ -module, C a Y -prescheme, $\varepsilon : Y \rightarrow C$ a Y -section, and $q : V = \mathbf{V}(\mathcal{L}) \rightarrow C$ a Y -morphism, all such that the diagram in (8.9.1.1) commutes. Suppose further that, if $p = (p_0, \theta)$, then $\theta : \mathcal{O}_Y \rightarrow p_*(\mathcal{O}_X)$ is an isomorphism. Let $\mathcal{S}' = \bigoplus_{n \geq 0} p_*(\mathcal{L}^{\otimes n})$ and $C' = \text{Spec}(\mathcal{S}')$, and let $q' : \mathbf{V}(\mathcal{L}) \rightarrow C'$ be the canonical Y -morphism (8.8.5). Then there exists exactly one Y -morphism $u : C' \rightarrow C$ such that $q = u \circ q'$.*

Proof. The hypothesis on θ implies, in particular, that p is surjective; since, by (8.8.4), the restriction of q' to $\mathbf{V}(\mathcal{L}) - j(X)$ is an isomorphism to $C' - \varepsilon'(Y)$ (where ε' is the vertex section of C'), it follows from (8.8.4) that q' is proper and surjective; furthermore, by (8.8.6), if we let $q' = (q'_0, \tau)$, then $\tau : \mathcal{O}_{C'} \rightarrow q'_*(\mathcal{O}_V)$ is an isomorphism. We are thus in a situation where we can apply (8.11.1), and we will have proven the lemma if we show that q is constant on every fibre $q'^{-1}(z')$, where $z' \in C'$. But this condition is trivially satisfied for $z' \notin \varepsilon'(Y)$. If $z' \in \varepsilon'(Y)$, then there exists exactly one $y \in Y$ such that $z' = \varepsilon'(y)$, and, by commutativity of (8.8.5.2) and the fact that q' sends $\mathbf{V}(\mathcal{L}) - j(X)$ to $C' - \varepsilon'(Y)$, $q'^{-1}(z') = j(p^{-1}(y))$; the commutativity of the diagram in (8.9.1.1) then proves our claim. $\square$

Corollary (8.11.6). — *Under the hypotheses of (8.11.5), suppose further that q is proper, and that the restriction of q to $\mathbf{V}(\mathcal{L}) - j(X)$ is an isomorphism to $C - \varepsilon(Y)$. Then the morphism u is universally closed, surjective, and radicial, and its restriction to $C' - \varepsilon'(Y)$ is an isomorphism to $C - \varepsilon(Y)$.*

Proof. Since q' is an isomorphism from $\mathbf{V}(\mathcal{L}) - j(X)$ to $C' - \varepsilon'(Y)$ (8.8.4), the last claim follows immediately from the fact that $q = u \circ q'$. Furthermore, the commutativity of the diagrams ^[II:191] in (8.8.5.2) and (8.9.1.1) shows that the restriction of u to the closed subprescheme $\varepsilon'(Y)$ of C' is an isomorphism to the closed subprescheme $\varepsilon(Y)$ of C , from which we immediately deduce that, for all $z' \in \varepsilon'(Y)$, if $z = u(z')$, then u defines an isomorphism from $k(z)$ to $k(z')$. These remarks prove that u is bijective and radicial; furthermore, if $\psi : C \rightarrow Y$ and $\psi' : C' \rightarrow Y$ are the structure morphisms, then $\psi' = \psi \circ u$, and, since ψ' is separated (1.2.4), so too is u (I, 5.5.1, v). We have already seen, in the proof of (8.11.5), that q' is surjective; since $q = u \circ q'$ is proper, we finally conclude, from (5.4.3) and (5.4.9), that u is universally closed. $\square$

Proposition (8.11.7). — *Let Y be a prescheme, X an integral prescheme, $p : X \rightarrow Y$ a proper morphism, $\mathcal{L}$ a p -ample invertible $\mathcal{O}_X$ -module, C a normal Y -prescheme, $\varepsilon : Y \rightarrow C$ a Y -section, and $q : V = \mathbf{V}(\mathcal{L}) \rightarrow C$ a Y -morphism, all such that the diagram in (8.9.1.1) commutes. Suppose further that, if $p = (p_0, \theta)$, then $\theta : \mathcal{O}_Y \rightarrow p_*(\mathcal{O}_X)$ is an isomorphism. Let $\mathcal{S}' = \bigoplus_{n \geq 0} p_*(\mathcal{L}^{\otimes n})$ and $C' = \text{Spec}(\mathcal{S}')$, and let $q' : \mathbf{V}(\mathcal{L}) \rightarrow C'$ be the canonical Y -morphism (8.8.5). Then the unique Y -morphism $u : C' \rightarrow C$ such that $q = u \circ q'$ is an isomorphism.*

Proof. It follows from (8.8.6) that C' is integral; since u is a homeomorphism of the underlying subspaces $C' \rightarrow C$ (u being bijective and closed, by (8.11.6)), C is irreducible, thus integral, and, since the restriction of u to a non-empty open subset of C' is an isomorphism to an open subset of C , u is birational. Since C is assumed to be normal, it suffices to apply (8.11.2) to obtain the conclusion. $\square$

Remark (8.11.8). — (i) The hypothesis that C is normal implies that X is also normal. Indeed, $C' = \text{Spec}(\mathcal{S}')$ is then normal, being isomorphic to C , and integral, by (8.8.6); we thus conclude that $\text{Proj}(\mathcal{S}')$ is normal. Indeed, the question is local on Y ; if Y is affine, with $\mathcal{S}' = \tilde{S}'$, then the ring $S' = \Gamma(C', \mathcal{O}_{C'})$ is integral and integrally closed (8.8.6.1), and so, for every homogeneous element $f \in S'_+$, the graded ring S'_f is integral and integrally closed [13, t. I, p. 257 and 261], and thus so too is the ring $S'_{(f)}$ of its degree-zero terms, because the intersection of S'_f with the field of fractions of $S'_{(f)}$ is equal to $S'_{(f)}$; this proves our claim (6.3.4). Finally, since X is isomorphic to an open subprescheme of $\text{Proj}(\mathcal{S}')$ (8.8.1), X is indeed normal. We can thus express (8.11.7) in the following form: *If X is integral and normal, and $p = (p_0, \theta) : X \rightarrow Y$ is a proper morphism such that $\theta : \mathcal{O}_Y \rightarrow p_*(\mathcal{O}_X)$ is an isomorphism, then, for every p -ample $\mathcal{O}_X$ -module $\mathcal{L}$, there exists exactly one way of contracting the zero section of $V = \mathbf{V}(\mathcal{L})$ to obtain a normal Y -scheme C and a proper Y -morphism $q : V \rightarrow C$.*

- (ii) When p is proper, the hypothesis $p_*(\mathcal{O}_X) = \mathcal{O}_Y$ can be considered as an auxiliary hypothesis, not really restricting the generality of the result. Indeed, if it is not satisfied, then it suffices to replace Y with the Y -scheme $Y' = \text{Spec}(p_*(\mathcal{O}_X))$, and to consider X as a Y' -scheme. We will return to this general method in chapter III, § 4.

8.12. Quasi-coherent sheaves on projecting cones

(8.12.1). Let us use the hypotheses and notation of (8.3.1). Let $\mathcal{M}$ be a quasi-coherent graded $\mathcal{S}$ -module; to avoid any confusion, we denote by $\widetilde{\mathcal{M}}$ the quasi-coherent $\mathcal{O}_C$ -module ^[II:192] associated to $\mathcal{M}$ (1.4.3) when $\mathcal{M}$ is considered as a non-graded $\mathcal{S}$ -module, and by $\mathcal{P}roj_0(\mathcal{M})$ the quasi-coherent $\mathcal{O}_X$ -module associated to $\mathcal{M}$, $\mathcal{M}$ being considered this time as a graded $\mathcal{S}$ -module (in other words, the $\mathcal{O}_X$ -module denoted by $\widetilde{\mathcal{M}}$ in (3.2.2)). In addition, we set

$$(8.12.1.1) \quad \mathcal{M}_X = \mathcal{P}roj(\mathcal{M}) = \bigoplus_{n \in \mathbb{Z}} \mathcal{P}roj_0(\mathcal{M}(n));$$

the quasi-coherent graded $\mathcal{O}_X$ -algebra $\mathcal{S}_X$ being defined by (8.6.1.1), $\mathcal{P}roj(\mathcal{M})$ is equipped with a structure of a (quasi-coherent) graded $\mathcal{S}_X$ -module, by means of the canonical homomorphisms (3.2.6.1)

$$(8.12.1.2) \quad \mathcal{O}_X(m) \otimes_{\mathcal{O}_X} \mathcal{P}roj_0(\mathcal{M}(n)) \longrightarrow \mathcal{P}roj_0(\mathcal{S}(m) \otimes_{\mathcal{S}} \mathcal{M}(n)) \longrightarrow \mathcal{P}roj_0(\mathcal{M}(m+n)),$$

the verification of the axioms of sheaves of modules being done using the commutative diagram in (2.5.11.4).

If $Y = \text{Spec}(A)$ is affine, $\mathcal{S} = \widetilde{S}$, and $\mathcal{M} = \widetilde{M}$, where S is a graded A -algebra and M is a graded S -module, then, for every homogeneous element $f \in S_+$, we have

$$(8.12.1.3) \quad \Gamma(X_f, \mathcal{P}roj(\widetilde{M})) = M_f$$

by the definitions and (8.2.9.1).

Now consider the quasi-coherent graded $\widehat{\mathcal{S}}$ -module

$$(8.12.1.4) \quad \widehat{\mathcal{M}} = \mathcal{M} \otimes_{\mathcal{S}} \widehat{\mathcal{S}}$$

($\widehat{\mathcal{S}}$ being defined by (8.3.1.1)); this induces a quasi-coherent graded $\mathcal{O}_{\widehat{C}}$ -module $\mathcal{P}roj_0(\widehat{\mathcal{M}})$, which we will also denote by

$$(8.12.1.5) \quad \mathcal{M}^\square = \mathcal{P}roj_0(\widehat{\mathcal{M}}).$$

It is clear (3.2.4) that $\mathcal{M}^\square$ is an additive functor which is exact in $\mathcal{M}$, commuting with direct sums and with inductive limits.

Proposition (8.12.2). — *With the notation of (8.3.2), we have canonical functorial isomorphisms*

$$(8.12.2.1) \quad i^*(\mathcal{M}^\square) \xrightarrow{\sim} \widetilde{\mathcal{M}}, \quad j^*(\mathcal{M}^\square) \xrightarrow{\sim} \mathcal{P}roj_0(\mathcal{M}).$$

Indeed, $i^*(\mathcal{M}^\square)$ is canonically identified with $(\widehat{\mathcal{M}}/(\mathbf{z}-1)\widehat{\mathcal{M}})^\sim$ on $\text{Spec}(\widehat{\mathcal{S}}/(\mathbf{z}-1)\widehat{\mathcal{S}})$ by (3.2.3); the first of the canonical isomorphisms (8.12.2.1) is then immediately induced (1.4.1) by the canonical isomorphism $\widehat{\mathcal{M}}/(\mathbf{z}-1)\widehat{\mathcal{M}} \xrightarrow{\sim} \mathcal{M}$. The canonical immersion $j : X \rightarrow \widehat{C}$ corresponds to the canonical homomorphism $\widehat{\mathcal{S}} \rightarrow \mathcal{S}$ with kernel $\mathbf{z}\widehat{\mathcal{S}}$ (8.3.2); the second homomorphism (8.12.2.1) is the particular case of the canonical homomorphism (3.5.2, ii), since here we have $\widehat{\mathcal{M}} \otimes_{\widehat{\mathcal{S}}} \mathcal{S} = \mathcal{M}$; to verify that this is an isomorphism, we can restrict to the case where $Y = \text{Spec}(A)$ is affine, $\mathcal{S} = \widetilde{S}$, and $\mathcal{M} = \widetilde{M}$; by appealing to (2.8.8), the proof that, for all homogeneous f in S_+ , the preceding homomorphism, restricted to X_f , restricts to an isomorphism, is then immediate.

^[II:193]

By an abuse of language, we again say, thanks to the existence of the first isomorphism (8.12.2.1), that $\mathcal{M}^\square$ is the projective closure of the $\mathcal{O}_X$ -module $\widetilde{\mathcal{M}}$ (it being implicit that the data of the $\mathcal{O}_C$ -module $\widehat{\mathcal{M}}$ includes the grading of the $\mathcal{S}$ -module $\mathcal{M}$).

(8.12.3). With the notation of (8.3.5), we have a canonical functorial homomorphism

$$(8.12.3.1) \quad p^*(\mathcal{P}roj(\mathcal{M})) \longrightarrow \mathcal{M}^\square|_{\widehat{E}}.$$

Indeed, this is a particular case of the homomorphism $\nu^\sharp$ defined more generally in (3.5.6). If $Y = \operatorname{Spec}(A)$ is affine, $\mathcal{S} = \tilde{S}$, and $\mathcal{M} = \tilde{M}$, then, by appealing to (2.8.8), the restriction of (8.12.3.1) to $p^{-1}(X_f) = \hat{C}_f$ (for some homogeneous f in S_+) corresponds to the canonical homomorphism

$$(8.12.3.2) \quad M_{(f)} \otimes_{S_{(f)}} S_f^{\leq} \longrightarrow M_f^{\leq}$$

taking into account (8.2.3.2) and (8.2.5.2).

(8.12.4). Let us place ourselves in the settings of (8.5.1), and assume its hypotheses and keep its notation. It follows from (1.5.6) that, for every quasi-coherent graded $\mathcal{S}$ -module $\mathcal{M}$, we have, on one hand, a canonical isomorphism

$$(8.12.4.1) \quad \Phi^*(\tilde{\mathcal{M}}) \xrightarrow{\sim} (q^*(\mathcal{M}) \otimes_{q^*(\mathcal{S})} \mathcal{S}')^\sim$$

of $\mathcal{O}_C$ -modules; on the other hand, (3.5.6) implies the existence of a canonical $\operatorname{Proj}(\phi)$ -morphism

$$(8.12.4.2) \quad \mathcal{P}\mathcal{r}\mathcal{o}\mathcal{j}_0(\mathcal{M}) \longrightarrow \mathcal{P}\mathcal{r}\mathcal{o}\mathcal{j}_0(q^*(\mathcal{M}) \otimes_{q^*(\mathcal{S})} \mathcal{S}')|G(\phi)$$

and also of a canonical $\hat{\Phi}$ -morphism

$$(8.12.4.3) \quad \mathcal{P}\mathcal{r}\mathcal{o}\mathcal{j}_0(\widehat{\mathcal{M}}) \longrightarrow \mathcal{P}\mathcal{r}\mathcal{o}\mathcal{j}_0(q^*(\widehat{\mathcal{M}}) \otimes_{q^*(\widehat{\mathcal{S}})} \widehat{\mathcal{S}}')|G(\hat{\phi}).$$

(8.12.5). Consider now the setting of (8.6.1), with the same notation; we thus take $Y' = X$, the morphism $q : X \rightarrow Y$ being the structure morphism, and ϕ the canonical q -morphism (8.6.1.2). We then have a canonical isomorphism

$$(8.12.5.1) \quad q^*(\mathcal{M}) \otimes_{q^*(\mathcal{S})} \mathcal{S}_X^{\geq} \xrightarrow{\sim} \mathcal{M}_X^{\geq}$$

by setting $\mathcal{M}_X^{\geq} = \bigoplus_{n \geq 0} \mathcal{P}\mathcal{r}\mathcal{o}\mathcal{j}_0(\mathcal{M}(n))$. We can indeed restrict to the case where $Y = \operatorname{Spec}(A)$ is affine, $\mathcal{S} = \tilde{S}$, and $\mathcal{M} = \tilde{M}$, and define the isomorphism (8.12.5.1) on each of the affine open subsets X_f (where f is homogeneous in S_+), by verifying the compatibility with taking a homogeneous multiple of f . But the restriction to X_f of the left-hand side of (8.12.5.1) is $\tilde{M}' = ((M \otimes_A S_{(f)}) \otimes_{S \otimes_A S_{(f)}} S_f^{\geq})^\sim$ by (8.6.2.1); since we have a canonical isomorphism from $M \otimes_A S_{(f)}$ to $M \otimes_S (S \otimes_A S_{(f)})$, we have an induced isomorphism from $\tilde{M}'$ to $(M \otimes_S S_f^{\geq})^\sim$, and the latter is canonically isomorphic, by (8.2.9.1), to the restriction to X_f of the right-hand side of (8.12.5.1), and satisfies the required compatibility conditions. [II:194]

Replacing $\mathcal{M}$ by $\widehat{\mathcal{M}}$, $\mathcal{S}$ by $\widehat{\mathcal{S}}$, and $\mathcal{S}_X$ by $(\mathcal{S}_X^{\geq})^\wedge$ in the previous argument, we similarly have a canonical isomorphism

$$(8.12.5.2) \quad q^*(\widehat{\mathcal{M}}) \otimes_{q^*(\widehat{\mathcal{S}})} (\mathcal{S}_X^{\geq})^\wedge \xrightarrow{\sim} (\mathcal{M}_X^{\geq})^\wedge.$$

If we recall (8.6.2) that the structure morphism $u : \operatorname{Proj}(\mathcal{S}_X^{\geq}) \rightarrow X$ is an isomorphism, then we deduce, first of all, from the above, that we have a canonical u -isomorphism

$$(8.12.5.3) \quad \mathcal{P}\mathcal{r}\mathcal{o}\mathcal{j}_0 \mathcal{M} \xrightarrow{\sim} \mathcal{P}\mathcal{r}\mathcal{o}\mathcal{j}_0(\mathcal{M}_X^{\geq})$$

as a particular case of (8.12.4.2). We note that, with the notation from the proof of (8.6.2), this reduces to seeing that the canonical homomorphism $M_{(f)} \otimes_{S_{(f)}} (S_f^{\geq})^{(d)} \rightarrow (M_f^{\geq})^{(d)}$ is an isomorphism whenever $f \in S_d$, which is immediate.

Secondly, the isomorphism (8.12.5.2) gives us, by this time applying (8.12.4.3) to the canonical morphism $r = \operatorname{Proj}(\hat{\alpha}) : \hat{C}_X \rightarrow \hat{C}$, a canonical r -morphism

$$(8.12.5.4) \quad \mathcal{M}^\square \longrightarrow (\mathcal{M}_X^{\geq})^\square.$$

Recall now (8.6.2) that the restrictions of r to the punctured cones $\hat{E}_X$ and E_X are isomorphisms to $\hat{E}$ and E (respectively). Furthermore:

Proposition (8.12.6). — *The restrictions to $\hat{E}_X$ and E_X of the canonical r -morphism (8.12.5.4) are isomorphisms*

$$(8.12.6.1) \quad \mathcal{M}^\square|_{\hat{E}} \xrightarrow{\sim} (\mathcal{M}_X^{\geq})^\square|_{\hat{E}_X}$$

$$(8.12.6.2) \quad \widetilde{\mathcal{M}}|_E \xrightarrow{\sim} (\mathcal{M}_X^{\geq})^\sim|_{E_X}.$$

Proof. We restrict to the case where Y is affine, as in the proof of (8.6.2) (whose notation we adopt); by reducing to definitions (2.8.8), we have to show that the canonical homomorphism

$$\widehat{M}_{(f)} \otimes_{\widehat{S}_{(f)}} (S_f^{\geq})_{(f/1)}^{\wedge} \longrightarrow (M \otimes_S S_f^{\geq})_{(f/1)}^{\wedge}$$

is an isomorphism; but, by (8.2.3.2) and (8.2.5.2), the left-hand side is canonically identified with $M_f^{\leq} \otimes_{S_f^{\leq}} (S_f^{\geq})_{f/1}^{\leq}$, and thus with $M_f^{\leq}$, by (8.2.7.2), and the right-hand side with $(M_f^{\geq})_{f/1}^{\leq}$, and thus also with $M_f^{\leq}$, by (8.2.9.2), whence the conclusion concerning (8.12.6.1); (8.12.6.2) then follows from (8.12.6.1) and (8.12.2.1). $\square$

Corollary (8.12.7). — *With the identifications of (8.6.3), the restriction of $(\mathcal{M}_X^{\geq})^{\square}$ to $\widehat{E}_X$ can be identified with $(\mathcal{M}_X^{\leq})^{\sim}$, and the restriction of $(\mathcal{M}_X^{\geq})^{\square}$ to E_X with $\widetilde{\mathcal{M}}_X$.*

Proof. We can restrict to the affine case, and this follows from the identification of $(M_f^{\geq})_{f/1}^{\leq}$ with $M_f^{\leq}$, and of $(M_f^{\geq})_{f/1}$ with M_f (8.2.9.2). $\square$

Proposition (8.12.8). — *Under the hypotheses of (8.6.4), the canonical homomorphism (8.12.3.1) is an isomorphism.*

Proof. Taking into account the fact that $\text{Proj}(\mathcal{S}_X^{\geq}) \rightarrow X$ is an isomorphism (8.6.2), and the [II:195] isomorphisms (8.12.5.4) and (8.12.6.1), we are led to proving the corresponding proposition for the canonical homomorphism $p_X^*(\text{Proj}_0(\mathcal{M}_X^{\geq})) \rightarrow (\mathcal{M}_X^{\geq})^{\square}|_{\widehat{E}_X}$, or, in other words, we can restrict to the case where $\mathcal{S}_1$ is an invertible $\mathcal{O}_Y$ -module, and where $\mathcal{S}$ is generated by $\mathcal{S}_1$. With the notation of (8.12.3), we then have, for some $f \in S_1$, that $S_f^{\leq} = S_{(f)}[1/f]$, and the canonical homomorphism $M_{(f)} \otimes_{S_{(f)}} S_f^{\leq} \rightarrow M_f^{\leq}$ is an isomorphism, by the definition of $M_f^{\leq}$. $\square$

(8.12.9). Consider now the quasi-coherent $\mathcal{S}$ -modules

$$\mathcal{M}_{[n]} = \bigoplus_{m \geq n} \mathcal{M}_m$$

and (with the notation of (8.7.2)) the quasi-coherent graded $\mathcal{S}^{\natural}$ -module

$$(8.12.9.1) \quad \mathcal{M}^{\natural} = \left(\bigoplus_{n \geq 0} \mathcal{M}_{[n]} \right)^{\sim}.$$

We have seen (8.7.3) that there exists a canonical C-isomorphism $h : C_X \xrightarrow{\sim} \text{Proj}(\mathcal{S}^{\natural})$. Furthermore:

Proposition (8.12.10). — *There exists a canonical h-isomorphism*

$$(8.12.10.1) \quad \text{Proj}_0(\mathcal{M}^{\natural}) \xrightarrow{\sim} \widetilde{\mathcal{M}}_X.$$

Proof. We argue as in (8.7.3), this time using the existence of the di-isomorphism (8.2.9.3) instead of (8.2.7.3). We leave the details to the reader. $\square$

8.13. Projective closures of subsheaves and closed subschemes

(8.13.1). With hypotheses and notation as in (8.12.1), consider a *not-necessarily graded* quasi-coherent sub- $\mathcal{S}$ -module $\mathcal{N}$ of $\mathcal{M}$. We can then consider the quasi-coherent $\mathcal{O}_C$ -module $\widetilde{\mathcal{N}}$ associated to $\mathcal{N}$, which is a sub- $\mathcal{O}_C$ -module of $\widetilde{\mathcal{M}}$. We have seen elsewhere (8.12.2.1) that $\widetilde{\mathcal{M}}$ can be identified with the restriction of $\mathcal{M}^\square$ to C . Since the canonical injection $i : C \rightarrow \widehat{C}$ is an affine morphism (8.3.2), and *a fortiori* quasi-compact, the *canonical extension* $(\widetilde{\mathcal{N}})^-$, the largest sub- $\mathcal{O}_{\widehat{C}}$ -module contained in $\mathcal{M}^\square$ and inducing $\widetilde{\mathcal{N}}$ on C , is a *quasi-coherent* $\mathcal{O}_{\widehat{C}}$ -module (I, 9.4.2). We will give a more explicit description by using a graded $\widehat{\mathcal{S}}$ -module.

(8.13.2). For this, consider, for every integer $n \geq 0$, the homomorphism $\bigoplus_{i \leq n} \mathcal{M}_i \rightarrow \mathcal{M}$ which, for every open U of Y , sends the family

$$(s_i) \in \bigoplus_{i \leq n} \Gamma(U, \mathcal{M}_i)$$

to the section $\sum_i s_i \in \Gamma(U, \mathcal{M})$. Denote by $\mathcal{N}'_n$ the inverse image of $\mathcal{N}$ by this homomorphism, which is a quasi-coherent sub- $\mathcal{O}_Y$ -module of $\bigoplus_{i \leq n} \mathcal{M}_i$. Now consider the homomorphism $\bigoplus_{i \leq n} \mathcal{M}_i \rightarrow \widehat{\mathcal{M}} = \mathcal{M}[\mathbf{z}]$ which sends (s_i) to the section $\sum_{i \leq n} s_i \mathbf{z}^{n-i} \in \Gamma(U, \widehat{\mathcal{M}}_n)$, and let $\mathcal{N}_n$ be the image of $\mathcal{N}'_n$ under this homomorphism; we immediately have that $\overline{\mathcal{N}} = \bigoplus_{n \geq 0} \mathcal{N}_n$ is a (quasi-coherent) sub- $\widehat{\mathcal{S}}$ -module of $\widehat{\mathcal{M}}$; we say that $\overline{\mathcal{N}}$ is induced from $\mathcal{N}$ by *homogenisation*, via the “homogenising variable” $\mathbf{z}$. We note [II:196] that, if $\mathcal{N}$ is already a *graded* sub- $\mathcal{S}$ -module of $\mathcal{M}$, then $\overline{\mathcal{N}}$ can be identified with the direct sum of the components $\mathcal{N}_n$ of degree $n \geq 0$ in $\widehat{\mathcal{N}} = \mathcal{N}[\mathbf{z}]$.

Proposition (8.13.3). — *The $\mathcal{O}_{\widehat{C}}$ -module $\mathcal{P}\mathcal{r}\mathcal{O}_0(\overline{\mathcal{N}})$ is the canonical extension $(\widetilde{\mathcal{N}})^-$ of $\widetilde{\mathcal{N}}$ to $\widehat{C}$.*

Proof. The question is local on Y and $\widehat{C}$ by the definition of the canonical extension (I, 9.4.1). We can thus already suppose that $Y = \text{Spec}(A)$ is affine, with $\mathcal{S} = \widetilde{S}$, $\mathcal{M} = \widetilde{M}$, and $\mathcal{N} = \widetilde{N}$, where N is a non-necessarily-graded sub- S -module of M . Furthermore (8.3.2.6), $\widehat{C}$ is a union of affine opens $\widehat{C}_z = C$ and $\widehat{C}_f = \text{Spec}(S_f^\leq)$ (with f homogeneous in S_+). It thus suffices to show that: (1) the restriction of $\mathcal{P}\mathcal{r}\mathcal{O}_0(\overline{\mathcal{N}})$ to C is $\widetilde{\mathcal{N}}$; (2) the restriction of $\mathcal{P}\mathcal{r}\mathcal{O}_0(\overline{\mathcal{N}})$ to each $\widehat{C}_f$ is the canonical extension of the restriction of $\mathcal{N}$ to $C \cap \widehat{C}_f = \text{Spec}(S_f)$ (8.3.2.6). For the first point, note that $\mathcal{P}\mathcal{r}\mathcal{O}_0(\overline{\mathcal{N}})|_C$ can be identified with $(\overline{N}_{(\mathbf{z})})^\sim$ (8.3.2.4); but $\overline{N}_{(\mathbf{z})}$ is canonically identified (2.2.5) with the image of $\overline{N}$ in $\widehat{M}/(\mathbf{z}-1)\widehat{M}$, and by the canonical isomorphism of the latter with M (8.2.5), this image can be identified with N , by the definition of $\overline{N}$ given in (8.13.2).

To prove the second point, note that the injection $i : C \cap \widehat{C}_f \rightarrow \widehat{C}_f$ corresponds to the canonical injection $S_f^\leq \rightarrow S_f$ (8.3.2.6); we also have that $\Gamma(\widehat{C}_f, \mathcal{M}^\square) = M_f^\leq$, that $\Gamma(\widehat{C}_f, i_*(\widetilde{\mathcal{N}})) = N_f$, and, by (8.12.2.1), that $\Gamma(\widehat{C}_f, i_*(i^*(\mathcal{M}^\square))) = M_f$. Taking (I, 9.4.2) into account, we are thus led to showing that $\overline{N}_{(f)} \subset \widehat{M}_{(f)} = M_f^\leq$ is canonically identified with the inverse image of N_f under the canonical injection $M_f^\leq \rightarrow M_f$. Indeed, let $d = \deg(f) > 0$, and suppose that an element $(\sum_{k \leq md} x_k)/f^m$ of M_f (with $x_k \in M_k$) is of the form y/f^m with $y \in N$. By multiplying y and the x_k by one single suitable f^h , we can already assume that $\sum_{k \leq md} x_k = y$. But in the identification of (8.2.5.2), $(\sum_{k \leq md} x_k)/f^m$ corresponds to $\sum_{k \leq md} x_k \mathbf{z}^{md-k}/f^m$, and this is indeed an element of $\overline{N}_{(f)}$, since $\sum_{k \leq md} x_k \in N$; the converse is evident. $\square$

Remark (8.13.4). — (i) The most important case of application of (8.13.3) is that where $\mathcal{M} = \mathcal{S}$, with $\widetilde{\mathcal{N}}$ then being an *arbitrary* quasi-coherent sheaf of ideals $\mathcal{I}$ of $\mathcal{O}_C$ (1.4.3), corresponding bijectively to a *closed subprescheme* Z of C . Then the canonical extension $(\mathcal{I})^-$ of $\mathcal{I}$ is the quasi-coherent sheaf of ideals of $\mathcal{O}_{\widehat{C}}$ that defines the *closure* $\overline{Z}$ of Z in $\widehat{C}$ (I, 9.5.10); Proposition (8.13.3) gives a canonical way of defining $\overline{Z}$ by using a graded ideal in $\widehat{\mathcal{S}} = \mathcal{S}[\mathbf{z}]$.

(ii) Suppose, to simplify things, that Y is affine, and adopt the notation from the proof of (8.13.3). For every non-zero $x \in N$, let $d(x)$ be the largest degree of the homogeneous components x_i of x in M ; by definition, $\overline{N}$ is the submodule of $\widehat{M}$ consisting of 0 and elements of the

form $h(x, k) = \mathbf{z}^k \sum_{i \leq d(x)} x_i \mathbf{z}^{d(x)-i}$ (for integral $k \geq 0$); it is thus generated, as a module over $\widehat{S} = S[\mathbf{z}]$, by the elements of the form

$$h(x, 0) = \sum_{i \leq d(x)} x_i \mathbf{z}^{d(x)-i}.$$

[II:197] We say that $h(x, 0)$ is induced from x by *homogenisation* via the “homogenising variable” $\mathbf{z}$. But since $h(x, 0)$ does not depend additively on x (nor *a fortiori* S -linearly), *we will refrain from believing* (even when $M = S$) that the $h(x, 0)$ form a *system of generators* of the graded $\widehat{S}$ -module $\overline{N}$ when we let x run over a *system of generators* of the S -module N . This is, however, the case when N is a *free cyclic* S -module (the only case considered in elementary algebraic geometry), since, if t is a basis of N , then $h(t, 0)$ generates the $\widehat{S}$ -module $\overline{N}$.

8.14. Supplement on sheaves associated to graded $\mathcal{S}$ -modules

(8.14.1). Let Y be a prescheme, $\mathcal{S}$ a positively-graded quasi-coherent $\mathcal{O}_Y$ -algebra, $X = \text{Proj}(\mathcal{S})$, and $q : X \rightarrow Y$ the structure morphism (which is separated, by (3.1.3)). Using the notation of (8.12.1), we have defined a functor $\mathcal{M}_X = \text{Proj}(\mathcal{M})$ in $\mathcal{M}$, from the category of quasi-coherent graded $\mathcal{S}$ -modules to the category of quasi-coherent graded $\mathcal{S}_X$ -modules; it is further clear (3.2.4) that this is an *additive* and *exact* functor, commuting with inductive limits.

Note, furthermore, that it follows immediately from the definition (8.12.1.1) that we have

$$(8.14.1.1) \quad \text{Proj}(\mathcal{M}(n)) = (\text{Proj}(\mathcal{M}))(n) \quad \text{for all } n \in \mathbf{Z}.$$

(8.14.2). We will first extend the canonical homomorphisms λ and μ , defined in (3.2.6), to $\mathcal{S}_X$ -modules of the form $\text{Proj}(\mathcal{M})$. For this, note that, for any $m \in \mathbf{Z}$ and $n \in \mathbf{Z}$, we have, by (2.1.2.1), a canonical homomorphism of $\mathcal{O}_X$ -modules

$$(8.14.2.1) \quad \lambda_{mn} : \text{Proj}_0(\mathcal{M}(m)) \otimes_{\mathcal{O}_X} \text{Proj}_0(\mathcal{N}(n)) \longrightarrow \text{Proj}_0((\mathcal{M} \otimes_{\mathcal{S}} \mathcal{N})(m+n)),$$

and likewise, by (2.1.2.2), a canonical homomorphism of $\mathcal{O}_X$ -modules

$$(8.14.2.2) \quad \mu_{mn} : \text{Proj}_0((\text{Hom}_{\mathcal{S}}(\mathcal{M}, \mathcal{N}))(n-m)) \longrightarrow \text{Hom}_{\mathcal{O}_X}(\text{Proj}_0(\mathcal{M}(m)), \text{Proj}_0(\mathcal{N}(n)))$$

for any quasi-coherent graded $\mathcal{S}$ -modules $\mathcal{M}$ and $\mathcal{N}$. This induces a homomorphism

$$\mu_k : \text{Proj}_0((\text{Hom}_{\mathcal{S}}(\mathcal{M}, \mathcal{N}))(k)) \longrightarrow (\text{Hom}_{\mathcal{S}_X}(\text{Proj}(\mathcal{M}), \text{Proj}(\mathcal{N})))_k$$

given by sending every $u \in \Gamma(U, \text{Proj}_0((\text{Hom}_{\mathcal{S}}(\mathcal{M}, \mathcal{N}))(k)))$ to the homomorphism $\mu_k(u)$, of degree k , of graded $\mathbf{Z}$ -modules $\Gamma(U, \text{Proj}(\mathcal{M})) \rightarrow \Gamma(U, \text{Proj}(\mathcal{N}))$ (where U is open in X) which, in each $\Gamma(U, \text{Proj}_0(\mathcal{M}(m)))$, agrees with $\mu_{m,m+k}(u)$; furthermore, by returning to the definition of the μ_{mn} (2.5.12.1), we immediately see that $\mu_k(u)$ is in fact a homomorphism of degree k of graded $\Gamma(U, \mathcal{S}_X)$ -modules, and, furthermore, that the μ_k define a homomorphism of *graded* $\mathcal{S}_X$ -modules

$$(8.14.2.3) \quad \text{Proj}(\text{Hom}_{\mathcal{S}}(\mathcal{M}, \mathcal{N})) \longrightarrow \text{Hom}_{\mathcal{S}_X}(\text{Proj}(\mathcal{M}), \text{Proj}(\mathcal{N})).$$

Similarly, taking the associativity diagram (2.5.11.4) into account, the homomorphisms (8.14.2.1) give a homomorphism of *graded* $\mathcal{S}_X$ -modules

$$(8.14.2.4) \quad \lambda : \text{Proj}(\mathcal{M}) \otimes_{\mathcal{S}_X} \text{Proj}(\mathcal{N}) \longrightarrow \text{Proj}(\mathcal{M} \otimes_{\mathcal{S}} \mathcal{N}).$$

[II:198]

Proposition (8.14.3). — *The homomorphism (8.14.2.4) is bijective; so too is (8.14.2.3) whenever the graded $\mathcal{S}$ -module $\mathcal{M}$ admits a finite presentation (3.1.1).*

Proof. The question is clearly local on X and Y ; we can thus suppose that $Y = \text{Spec}(A)$ is affine, with $\mathcal{S} = \widetilde{S}$, $\mathcal{M} = \widetilde{M}$, and $\mathcal{N} = \widetilde{N}$, where S is a positively-graded A -algebra, and M and N are graded S -modules. If f is a homogeneous element of S_+ , then the homomorphisms (8.14.2.1) and (8.14.2.2), restricted to the affine open $D_+(f)$, correspond to the canonical homomorphisms (2.5.11.1) and (2.5.12.1):

$$\begin{aligned} M(m)_{(f)} \otimes_{S_{(f)}} N(n)_{(f)} &\longrightarrow (M \otimes_S N)_{(f)}(m+n) \\ (\text{Hom}_S(M, N))_{(f)}(n-m) &\longrightarrow \text{Hom}_{S_{(f)}}(M(m)_{(f)}, N(n)_{(f)}). \end{aligned}$$

If we refer to the definitions of these homomorphisms, we thus see (taking (8.2.9.1) into account) that the restriction of (8.14.2.4) to $D_+(f)$ corresponds to the canonical homomorphism

$$M_f \otimes_{S_f} N_f \longrightarrow (M \otimes_S N)_f$$

defined in (0, 1.3.4), and we know that this latter homomorphism is an isomorphism. Similarly, the restriction of (8.14.2.3) to $D_+(f)$ corresponds to the canonical homomorphism (0, 1.3.5)

$$(\mathrm{Hom}_S(M, N))_f \longrightarrow \mathrm{Hom}_{S_f}(M_f, N_f)$$

taking into account the fact that, since M is of finite type, the module $\mathrm{Hom}_S(M, N)$, the direct sum of the subgroups consisting of *homogeneous* homomorphisms of S -modules (2.1.2), agrees with the set of *all* homomorphisms $M \rightarrow N$ of S -modules. The hypothesis that M admits a finite presentation then implies (0, 1.3.5) that the canonical homomorphism in question is indeed an isomorphism. $\square$

Proposition (8.14.4). — *If U is a quasi-compact open of X , then there exists an integer d such that, for every integer n that is a multiple of d , $\mathcal{O}_X(n)|_U$ is invertible, with its inverse being $\mathcal{O}_X(-n)|_U$.*

Proof. Since $q(U)$ is quasi-compact, it is covered by a finite number of affine opens V_i , and so every $x \in U$ is contained in some affine open of the form $D_+(f)$, where f is a homogeneous element of degree > 0 of one of the rings $\Gamma(V_i, \mathcal{S})$. Since U is quasi-compact, we can cover it by a finite number of such opens $D_+(f_j)$; let d be a common multiple of the degrees of the f_j . This d satisfies the desired property, by (2.5.17). $\square$

(8.14.5). With the hypotheses and notation of (8.14.1), we defined, in (3.3.2), canonical homomorphisms of $\mathcal{O}_Y$ -modules

$$(8.14.5.1) \quad \alpha_n : \mathcal{M}_n \longrightarrow q_*(\mathcal{P}roj_0(\mathcal{M}(n))) \quad (n \in \mathbf{Z}).$$

Generalising the notation of (3.3.1), we set, for every graded $\mathcal{S}_X$ -module $\mathcal{F}$,

$$(8.14.5.2) \quad \Gamma_*(\mathcal{F}) = \bigoplus_{n \in \mathbf{Z}} q_*(\mathcal{F}_n).$$

In particular, $\Gamma_*(\mathcal{S}_X) = \bigoplus_{n \in \mathbf{Z}} q_*(\mathcal{O}_X(n))$ is the graded $\mathcal{O}_Y$ -algebra denoted by $\Gamma_*(\mathcal{O}_X)$ in (3.3.1.2); it is clear that $\Gamma_*(\mathcal{F})$ is a graded $\Gamma_*(\mathcal{S}_X)$ -module (0, 4.2.2). When [III:199] we take $\mathcal{M} = \mathcal{S}$ in the homomorphisms (8.14.5.1), we obtain the homomorphism of graded $\mathcal{O}_Y$ -algebras

$$(8.14.5.3) \quad \alpha : \mathcal{S} \longrightarrow \Gamma_*(\mathcal{S}_X)$$

previously defined in (3.3.2), and which makes $\Gamma_*(\mathcal{F})$ a graded $\mathcal{S}$ -module; the homomorphisms (8.14.5.1) then define a homomorphism (of degree 0) of graded $\mathcal{S}$ -modules

$$(8.14.5.4) \quad \alpha : \mathcal{M} \longrightarrow \Gamma_*(\mathcal{P}roj(\mathcal{M})).$$

(8.14.6). In general, for a quasi-coherent graded $\mathcal{S}_X$ -module $\mathcal{F}$, it is not certain that the graded $\mathcal{S}$ -module $\Gamma_*(\mathcal{F})$ will necessarily be quasi-coherent. Consider an open X' of X such that the restriction $q' : X' \rightarrow Y$ of q to X' is a *quasi-compact* morphism. Since q' is further separated, $q'_*(\mathcal{F}')$ is then a quasi-coherent $\mathcal{O}_Y$ -module for every quasi-coherent $\mathcal{O}_{X'}$ -module $\mathcal{F}'$ (I, 9.2.2, b). We set

$$(8.14.6.1) \quad \mathcal{S}_{X'} = \mathcal{S}_X|_{X'} = \bigoplus_{n \in \mathbf{Z}} \mathcal{O}_X(n)|_{X'}$$

and, for every graded $\mathcal{S}_{X'}$ -module $\mathcal{F}'$,

$$(8.14.6.2) \quad \Gamma'_*(\mathcal{F}') = \bigoplus_{n \in \mathbf{Z}} q'_*(\mathcal{F}'_n).$$

The previous remark then shows that, if $\mathcal{F}'$ is a quasi-coherent $\mathcal{S}_{X'}$ -module, then $\Gamma'_*(\mathcal{F}')$ is a graded *quasi-coherent* $\mathcal{S}$ -module (I, 9.6.1).

We note also that the canonical injection $j : X' \rightarrow X$ is *quasi-compact*, because $q' = q \circ j$ is quasi-compact and q is separated (I, 6.6.4, v). Then $\mathcal{F} = j_*(\mathcal{F}')$ is a quasi-coherent graded $\mathcal{S}_X$ -module for every quasi-coherent graded $\mathcal{S}_{X'}$ -module $\mathcal{F}'$, and it follows from the previous definitions that

$$(8.14.6.3) \quad \Gamma'_*(\mathcal{F}') = \Gamma_*(\mathcal{F}).$$

With the same hypotheses on X' , for every quasi-coherent graded $\mathcal{S}$ -module $\mathcal{M}$, we set

$$(8.14.6.4) \quad \mathcal{P}roj'(\mathcal{M}) = \mathcal{P}roj(\mathcal{M})|X'$$

which is a quasi-coherent graded $\mathcal{S}_{X'}$ -module. The canonical homomorphism

$$\mathcal{P}roj(\mathcal{M}) \longrightarrow j_*(\mathcal{P}roj'(\mathcal{M}))$$

(0, 4.4.3) thus gives a canonical homomorphism $\Gamma_*(\mathcal{P}roj(\mathcal{M})) \rightarrow \Gamma'_*(\mathcal{P}roj'(\mathcal{M}))$ of graded $\mathcal{S}$ -modules, and, by composition with (8.14.5.4), we obtain a functorial canonical homomorphism (of degree 0) of quasi-coherent graded $\mathcal{S}$ -modules

$$(8.14.6.5) \quad \alpha' : \mathcal{M} \longrightarrow \Gamma'_*(\mathcal{P}roj'(\mathcal{M})).$$

(8.14.7). Keeping the hypotheses on X' from (8.14.6), let $\mathcal{F}'$ be a quasi-coherent graded $\mathcal{S}_{X'}$ -module, so that $\mathcal{P}roj'(\Gamma'_*(\mathcal{F}'))$ is also a graded quasi-coherent $\mathcal{S}_{X'}$ -module. ^[II:200] We will define a functorial canonical homomorphism (of degree 0) of graded $\mathcal{S}_{X'}$ -modules

$$(8.14.7.1) \quad \beta' : \mathcal{P}roj'(\Gamma'_*(\mathcal{F}')) \longrightarrow \mathcal{F}'.$$

Suppose first of all that $Y = \text{Spec}(A)$ is affine, and that $\mathcal{S} = \tilde{S}$, where S is a positively-graded A -algebra; then $\Gamma'_*(\mathcal{F}') = \tilde{M}$, where $M = \bigoplus_{n \in \mathbb{Z}} \Gamma(X', \mathcal{F}'_n)$ is a graded S -module. Let $f \in S_d$ be such that $D_+(f) \subset X'$; by definition (2.6.2), $\alpha_d(f)$ restricted to $D_+(f)$ is the section of $\mathcal{O}_X(d)$ over $D_+(f)$ corresponding to the element $f/1$ of $(S(d))_{(f)}$, and is thus invertible; thus so too is $\alpha_{nd}(f^n)$ for every $n > 0$. From this, we immediately conclude that we have defined an S_f -homomorphism (of degree 0) of graded modules $\beta_f : M_f \rightarrow \Gamma(D_+(f), \mathcal{F}')$ by sending each element $z/f^n \in M_f$ (where $z \in M$) to the section $(z|D_+(f))(\alpha_d(f^n)|D_+(f))^{-1}$ of $\mathcal{F}'$ over $D_+(f)$. Furthermore, we have a commutative diagram corresponding to (2.6.4.1), whence the definition of β' in this case. To pass to the general case, we must consider an A -algebra A' , the graded A' -algebra $S' = S \otimes_A A'$, and use the commutative diagram analogous to (2.8.13.2); we leave the details to the reader.

Proposition (8.14.8). — *If X' is an open of $X = \text{Proj}(\mathcal{S})$ such that $q' : X' \rightarrow Y$ is quasi-compact, then the homomorphism β' defined in (8.14.7) is bijective.*

Proof. We can clearly restrict to the case where Y is affine, and everything then reduces to proving (with the notation of (8.14.7)) that the homomorphism $\beta_f : M_f \rightarrow \Gamma(D_+(f), \mathcal{F}')$ is an isomorphism. But replacing f by one of its powers changes neither $D_+(f)$ nor β_f ; since X' is quasi-compact by hypothesis, we can always assume, by (8.14.4), that the sheaf $\mathcal{O}_X(d)$ is invertible. Since X' is a scheme (because q' is separated), the proposition is then exactly (I, 9.3.1). $\square$

Corollary (8.14.9). — *Under the hypotheses of (8.14.8), every quasi-coherent graded $\mathcal{S}_{X'}$ -module is isomorphic to a graded $\mathcal{S}_{X'}$ -module of the form $\mathcal{P}roj'(\mathcal{M})$, where $\mathcal{M}$ is a quasi-coherent graded $\mathcal{S}$ -module. Further, if $\mathcal{F}'$ is of finite type, and if we assume that Y is a quasi-compact scheme, or a prescheme whose underlying space is Noetherian, then we can assume that $\mathcal{M}$ is of finite type.*

Proof. The proof starting from (8.14.8) follows exactly the same route as the proof of (3.4.5) starting from (3.4.4), and we leave the details to the reader. $\square$

Proposition (8.14.10). — *Under the hypotheses of (8.14.7), let $\mathcal{M}$ be a quasi-coherent graded $\mathcal{S}$ -module, and $\mathcal{F}'$ a quasi-coherent graded $\mathcal{S}_{X'}$ -module; the composite homomorphisms*

$$(8.14.10.1) \quad \mathcal{P}roj'(\mathcal{M}) \xrightarrow{\mathcal{P}roj'(\alpha')} \mathcal{P}roj'(\Gamma'_*(\mathcal{P}roj'(\mathcal{M}))) \xrightarrow{\beta'} \mathcal{P}roj'(\mathcal{M})$$

$$(8.14.10.2) \quad \Gamma'_*(\mathcal{F}') \xrightarrow{\alpha'} \Gamma'_*(\mathcal{P}roj'(\Gamma'_*(\mathcal{F}'))) \xrightarrow{\Gamma'_*(\beta')} \Gamma'_*(\mathcal{F}')$$

are the identity isomorphisms.

Proof. The question is local on Y , and the proof follows as in (2.6.5); we leave the details to the reader. $\square$

Remark (8.14.11). — In chapter III (III, 2.3.1), we will see that, when Y is locally Noetherian, and $\mathcal{S}$ is a quasi-coherent graded $\mathcal{O}_Y$ -algebra of finite type (in which case ^[II:201] we can take $X' = X$), then the homomorphism α (8.14.5.4) is (TN)-bijective for every quasi-coherent graded $\mathcal{S}$ -module $\mathcal{M}$ satisfying condition (TF).

Remark (8.14.12). — The situation described in (8.14.4) is a particular case of the following. Let X be a ringed space, and $\mathcal{S}$ a (positively- and negatively-) graded $\mathcal{O}_X$ -algebra; suppose that there exists an integer $d > 0$ such that $\mathcal{S}_d$ and $\mathcal{S}_{-d}$ are *invertible*, with the canonical homomorphism

$$(8.14.12.1) \quad \mathcal{S}_d \otimes_{\mathcal{O}_X} \mathcal{S}_{-d} \longrightarrow \mathcal{O}_X$$

being an *isomorphism* (such that $\mathcal{S}_{-d}$ is identified with $\mathcal{S}_d^{-1}$). We then say that the graded $\mathcal{O}_X$ -algebra $\mathcal{S}$ is *periodic*, of *period* d . This nomenclature stems from the following property: *under the preceding hypotheses, for every graded $\mathcal{S}$ -module $\mathcal{F}$, the canonical homomorphism*

$$(8.14.12.2) \quad \mathcal{S}_d \otimes_{\mathcal{O}_X} \mathcal{F}_n \longrightarrow \mathcal{F}_{n+d}$$

is an isomorphism for all $n \in \mathbb{Z}$. Indeed, the question is local on X , and we can assume that $\mathcal{S}_d$ has an *invertible* section s over X , with its inverse s' being a section of $\mathcal{S}_{-d}$. The homomorphism $\mathcal{F}_{n+d} \rightarrow \mathcal{S}_d \otimes \mathcal{F}_n$, which sends each section $z \in \Gamma(U, \mathcal{F}_{n+d})$ to the section $(s|U) \otimes (s'|U)z$ of $\mathcal{S}_d \otimes \mathcal{F}_n$ over U , is then the inverse of (8.14.12.2), whence our claim. This induces, for all $k \in \mathbb{Z}$, a canonical isomorphism

$$(\mathcal{S}_d)^{\otimes k} \otimes \mathcal{F}_n \xrightarrow{\sim} \mathcal{F}_{n+kd}.$$

Then the data of a graded $\mathcal{S}$ -module $\mathcal{F}$ is equivalent to the data of $\mathcal{S}_0$ -modules $\mathcal{F}_i$ ($0 \leq i \leq d-1$) and canonical homomorphisms

$$\mathcal{S}_i \otimes \mathcal{F}_j \longrightarrow \mathcal{F}_{i+j} \quad \text{for } 0 \leq i, j \leq d-1$$

(setting $\mathcal{F}_{i+j} = \mathcal{S}_d \otimes \mathcal{S}_0 \mathcal{F}_{i+j-d}$ whenever $i+j \geq d$). Of course, for these homomorphisms to give a well-defined $\mathcal{S}$ -module structure on the direct sum of the $(\mathcal{S}_d)^{\otimes k} \otimes \mathcal{F}_i$ ($k \in \mathbb{Z}$, $0 \leq i \leq d-1$), they should satisfy some associativity conditions that we will not explain.

In the case where $d = 1$ (which is the one considered in (3.3)), we can thus say that the category of graded $\mathcal{S}$ -modules (resp. quasi-coherent $\mathcal{S}$ -modules if X is a prescheme and $\mathcal{S}$ is quasi-coherent) is *equivalent* to the category of arbitrary $\mathcal{S}_0$ -modules (resp. quasi-coherent $\mathcal{S}_0$ -modules); it is in this way that we can think of the results of this paragraph as generalising those of §3. Furthermore, we see that, under suitable finiteness conditions, the results of this paragraph (along with (8.14.11)) reduce, in some sense, the study of quasi-coherent graded algebras on a prescheme, and graded modules “modulo (TN)” on such algebras, to the study of the particular case where the algebras in question are *periodic* (and where condition (TN) for $\mathcal{M}$ (3.4.2) thus implies that $\mathcal{M} = 0$).

Remark (8.14.13). — Under the hypotheses of (8.14.1), let d be an integer > 0 ; we have defined a canonical Y -isomorphism h from X to $X^{(d)} = \text{Proj}(\mathcal{S}^{(d)})$ (3.1.8). For every ^[II-202] quasi-coherent graded $\mathcal{S}$ -module $\mathcal{M}$ and every integer k such that $0 \leq k \leq d-1$, we also have (with the notation of (3.1.1)) a canonical h -isomorphism

$$(8.14.13.1) \quad (\mathcal{P}roj(\mathcal{M}))^{(d,k)} \xleftarrow{\sim} \mathcal{P}roj(\mathcal{M}^{(d,k)}).$$

Suppose, first of all, that $Y = \text{Spec}(A)$ is affine, $\mathcal{S} = \tilde{S}$, and $\mathcal{M} = \tilde{M}$, where S is a positively-graded A -algebra, and M a graded S -module. We know, for every $f \in S_e$ ($e > 0$), that h sends $D_+(f)$ to $D_+(f^d)$, and corresponds to the canonical isomorphism $S_{(f^d)} \rightarrow S_{(f)}$ (2.2.2). The restriction of (8.14.13.1) to $D_+(f^d)$ then corresponds to the canonical di-isomorphism $M_{f^d} \rightarrow M_f$ restricted to the elements of M_{f^d} whose degree is congruent to k (modulo d). We leave to the reader the task of showing that these isomorphisms are compatible with passing from f to some homogeneous multiple fg , and then that there is an analogous compatibility with passing from S to a graded A' -algebra $S' = S \otimes_A A'$, where A' is some A -algebra. In particular, this gives us an h -isomorphism

$$(8.14.13.2) \quad (\mathcal{S}^{(d)})_{X^{(d)}} \xrightarrow{\sim} (\mathcal{S}_X)^{(d)}$$

that respects the multiplicative structures of both the source and the target, and that, thanks to (8.14.13.1), becomes an h -di-isomorphism from a graded $(\mathcal{S}^{(d)})_{X^{(d)}}$ -module to a graded $(\mathcal{S}_X)^{(d)}$ -module. Similarly, we have an h -isomorphism

$$(8.14.13.3) \quad \mathcal{P}roj_0(\mathcal{S}^{(d,k)}(n)) \xrightarrow{\sim} \mathcal{O}_X(nd+k),$$

which completes the result of (3.2.9, ii).

The isomorphism in (8.14.13.1) immediately induces an isomorphism of graded $\mathcal{S}^{(d)}$ -modules

$$(8.14.13.4) \quad \Gamma_*^{(d)}(\mathcal{P}roj(\mathcal{M}^{(d,k)})) \xrightarrow{\sim} \Gamma_*((\mathcal{P}roj(\mathcal{M}))^{(d,k)})$$

where $\Gamma_*^{(d)}$ corresponds to the structure morphism $q^{(d)} : X^{(d)} \rightarrow Y$; it can be immediately verified that the canonical homomorphism α (8.14.5.4), and the analogous homomorphism $\alpha^{(d)}$ for $X^{(d)}$, make the following diagram commute:

$$(8.14.13.5) \quad \begin{array}{ccc} & \mathcal{M}^{(d,k)} & \\ \alpha^{(d)} \swarrow & & \searrow \alpha \\ \Gamma_*^{(d)}(\mathcal{P}roj(\mathcal{M}^{(d,k)})) & \xrightarrow{\sim} & \Gamma_*((\mathcal{P}roj(\mathcal{M}))^{(d,k)}) \end{array}$$

the verification being carried out by supposing that Y is affine and then calculating the restrictions of the images under $\alpha^{(d)}$ and α of a single element of $M^{(d,k)}$ to the open subsets $D_+(f^d)$ and $D_+(f)$ (using the same notation as above).

Proposition (8.14.14). — *Let Y be a quasi-compact prescheme, $\mathcal{S}$ a quasi-coherent graded $\mathcal{O}_Y$ -algebra of finite type, and $\mathcal{M}$ a quasi-coherent graded $\mathcal{S}$ -module satisfying condition (TF); let $X = \text{Proj}(\mathcal{S})$. Then $\mathcal{S}_X$ is a periodic graded $\mathcal{O}_X$ -algebra (8.14.12), and there exists ^[II:203] some period d of $\mathcal{S}_X$ such that the $(\mathcal{P}roj(\mathcal{M}))^{(d,k)}$ ($0 \leq k \leq d-1$) are $(\mathcal{S}_X)^{(d)}$ -modules of finite type.*

Proof. Indeed, (3.1.10) proves that there exists some d such that $\mathcal{S}^{(d)}$ is generated by $\mathcal{S}_d = (\mathcal{S}^{(d)})_1$, with the latter being an $\mathcal{S}_0$ -module of finite type. To prove the first claim, we can thus, by (8.14.13.2), restrict to the case where $d = 1$, and the proposition then follows from (3.2.7). Furthermore, taking (8.14.13.1) into account, the second claim is a consequence of (2.1.6, iii) and (3.4.3). $\square$

Editorial note. For standalone use, entries [1], [9], [11], and [13], which are cited in this volume, are reproduced here from the preceding series bibliography. The EGA II authority itself prints only the continuation [23]–[26] on printed page 205.

BIBLIOGRAPHY

- [1] H. Cartan and C. Chevalley, *Séminaire de l'École Normale Supérieure, 8th year (1955–56): Géométrie algébrique*.
- [9] M. Nagata, *A general theory of algebraic geometry over Dedekind domains*, Amer. Math. J. **78** (1956), 78–116, and **80** (1958), 382–420.
- [11] P. Samuel, *Commutative algebra* (notes by D. Herzig), Cornell University, 1953.
- [13] P. Samuel and O. Zariski, *Commutative algebra*, 2 vols., Van Nostrand, New York, 1958–60.
[II:205]
- [23] C. Chevalley, *Introduction to the theory of algebraic functions of one variable*, Mathematical Surveys, no. 6, American Mathematical Society, New York, 1951.
- [24] P. Jaffard, *Les systèmes d'idéaux*, Dunod, Paris, 1960.
- [25] M. Nagata, *On the derived normal rings of Noetherian integral domains*, Mem. Coll. Sci. Kyoto, Ser. A, **29** (1955), 293–303.
- [26] M. Nagata, *Existence theorems for non-projective complete algebraic varieties*, Illinois J. Math. **2** (1958), 490–498.

[II:207]

INDEX OF NOTATION

- $\mathcal{A}(X), \mathcal{A}(\mathcal{F}), \mathcal{A}(\mathcal{B})$ (X an S -prescheme, $\mathcal{F}$ an $\mathcal{O}_X$ -module, $\mathcal{B}$ an $\mathcal{O}_X$ -algebra): 1.1.1.
 $\mathcal{A}(h)$ (h an S -morphism): 1.1.2.
 $\text{Spec}(\mathcal{B})$ ($\mathcal{B}$ an $\mathcal{O}_S$ -algebra): 1.3.1.
 $\widetilde{\mathcal{M}}$ ($\mathcal{M}$ a $\mathcal{B}$ -module, $\mathcal{B}$ an $\mathcal{O}_S$ -algebra): 1.4.3.
 $\mathbf{T}(E), \mathbf{S}(E), \mathbf{S}_A(E)$ (E an A -module): 1.7.1.
 $\mathbf{S}(\mathcal{E}), \mathbf{S}_{\mathcal{A}}(\mathcal{E})$ ($\mathcal{E}$ an $\mathcal{A}$ -module): 1.7.4.
 $\mathbf{V}(\mathcal{E})$ ($\mathcal{E}$ a quasi-coherent $\mathcal{O}_S$ -module): 1.7.8.
 $S_n, S_+, M_n, S^{(d)}, M^{(d,k)}, M^{(d)}$ (S a graded ring, M a graded S -module): 2.1.1.
 $M(n), S(n)$ (S a graded ring, M a graded S -module): 2.1.1.
 $\text{Hom}_S(M, N)$ (S a graded ring, M, N graded S -modules): 2.1.2.
 $\mathfrak{N}_+, \mathfrak{r}_+(\mathfrak{J})$ ($\mathfrak{J}$ an ideal of S_+): 2.1.10.
 $S_{(f)}, M_{(f)}$ (f a homogeneous element of S_+): 2.2.1.
 $S_{(T)}, M_{(T)}, S_{(\mathfrak{p})}, M_{(\mathfrak{p})}$ (T a multiplicative subset of S_+ consisting of homogeneous elements, $\mathfrak{p}$ a graded prime ideal of S_+): 2.2.7.
 $\text{Proj}(S)$ (S a graded ring): 2.3.1.
 $V_+(E)$ (E a subset of S_+): 2.3.2.
 $D_+(f)$ (f an element of S): 2.3.3.
 $j_+(Y)$ (Y a subset of $\text{Proj}(S)$): 2.3.10.
 $\widetilde{M}$ (M a graded S -module): 2.5.2.
 $\widetilde{u}$ (u a homomorphism of degree zero): 2.5.4.
 $\mathcal{O}_X(n), \mathcal{F}(n)$ ($X = \text{Proj}(S)$, $\mathcal{F}$ an $\mathcal{O}_X$ -module): 2.5.10.
 λ (canonical homomorphism): 2.5.11.
 μ (canonical homomorphism): 2.5.12.
 $X^{(d)}$ ($X = \text{Proj}(S)$, d an integer > 0): 2.5.16.
 $\Gamma_*(\mathcal{F})$ ($\mathcal{F}$ an $\mathcal{O}_X$ -module, $X = \text{Proj}(S)$): 2.6.1.
 $\alpha_n, \alpha, \alpha_{\mathcal{M}}$ (canonical homomorphisms): 2.6.2.
 $\beta, \beta_{\mathcal{F}}$ (canonical homomorphisms): 2.6.4.
 $G(\phi), {}^a\phi$ (by abuse of language; ϕ a homomorphism of graded rings): 2.8.1.
 $\text{Proj}(\phi)$ (ϕ a homomorphism of graded rings): 2.8.2.
 $\mathcal{S}^{(d)}, \mathcal{M}^{(d,k)}, \mathcal{M}^{(d)}, \mathcal{M}(n)$ ($\mathcal{S}$ a graded $\mathcal{O}_Y$ -algebra, $\mathcal{M}$ a graded $\mathcal{S}$ -module): 3.1.1.
 $\text{Proj}(\mathcal{S})$ ($\mathcal{S}$ a quasi-coherent graded $\mathcal{O}_Y$ -algebra): 3.1.3.
 X_f ($X = \text{Proj}(\mathcal{S})$, f a section of $\mathcal{S}_d$ over Y): 3.1.4.
 $\widetilde{\mathcal{M}}$ ($\mathcal{M}$ a graded $\mathcal{S}$ -module): 3.2.2.
 $\mathcal{O}_X(n), \mathcal{F}(n)$ ($X = \text{Proj}(\mathcal{S})$, $\mathcal{F}$ an $\mathcal{O}_X$ -module): 3.2.5.
 λ, μ (canonical homomorphisms): 3.2.6.
 $\Gamma_*(\mathcal{F})$ ($\mathcal{F}$ an $\mathcal{O}_X$ -module, $X = \text{Proj}(\mathcal{S})$): 3.3.1.
 $\alpha_n, \alpha, \alpha_{\mathcal{M}}$ (canonical homomorphisms): 3.3.2.
 $\beta, \beta_{\mathcal{F}}$ (canonical homomorphisms): 3.3.4.
 $G(\phi), \text{Proj}(\phi)$ (ϕ a homomorphism of graded $\mathcal{O}_Y$ -algebras): 3.5.1.
 $\text{Proj}(u)$ (u a g -morphism from a graded $\mathcal{O}_Y$ -algebra to a graded $\mathcal{O}_{Y'}$ -algebra): 3.5.6.
 $r_{\mathcal{L}, \psi}$ ($\mathcal{L}$ an invertible $\mathcal{O}_X$ -module): 3.7.1.
 $\mathbf{P}(\mathcal{E}), \mathbf{P}(E), \mathbf{P}_Y^n, \mathbf{P}_A^n$: 4.1.1.
 $\mathbf{P}(u)$ (u a surjective homomorphism of $\mathcal{O}_Y$ -modules): 4.1.2.
 $\text{Hyp}_Y(X, \mathcal{E})$ ($\mathcal{E}$ a quasi-coherent $\mathcal{O}_Y$ -module): 4.2.5.
 ς (Segre morphism): 4.3.1.
 $\mathcal{F}(n)$ ($\mathcal{F}$ an $\mathcal{O}_X$ -module and X equipped with an invertible $\mathcal{O}_X$ -module): 4.5.1.
 ε (identity automorphism): 4.5.1.

$P(u, T), \sigma_i(u)$ (u an endomorphism of a free module with a finite basis): 6.4.1.

$P(u, T), \sigma_i(u), \det u$ (u an endomorphism of a finite-type module over an integral ring or a reduced Noetherian ring): 6.4.2 and 6.4.7.

$\sigma_i(u), \det u$ (u an endomorphism of a locally free $\mathcal{A}$ -module): 6.4.8.

$\sigma_i, \det$ (homomorphisms of sheaves): 6.4.8, 6.4.9, and 6.4.10.

$\sigma_i(u), \det u$ (u an endomorphism of a finite-type $\mathcal{O}_X$ -module on a locally integral prescheme or on a locally Noetherian reduced prescheme): 6.4.9 and 6.4.10.

$N_{\mathcal{B}/\mathcal{A}}(f)$ (f a section of the $\mathcal{A}$ -algebra $\mathcal{B}$): 6.5.1.

$N_{\mathcal{B}/\mathcal{A}}(\mathcal{L}')$ ($\mathcal{L}'$ an invertible $\mathcal{B}$ -module): 6.5.2.

$N_{\mathcal{B}/\mathcal{A}}(h')$ (h' a homomorphism of invertible $\mathcal{B}$ -modules): 6.5.3.

$N_{X'/X}(\mathcal{L}')$ (X' finite over X , $\mathcal{L}'$ an invertible $\mathcal{O}_{X'}$ -module): 6.5.5.

$N_{X'/X}(h')$ (h' a homomorphism of invertible $\mathcal{O}_{X'}$ -modules): 6.5.5.

$\mathcal{S}_X$ ($X = \text{Proj}(\mathcal{S})$, $\mathcal{S}$ a sum of ideals of $\mathcal{O}_Y$): 8.1.5.

$S^{\geq}, S^{\leq}, S_f^{\geq}, S_f^{\leq}, M^{\geq}, M^{\leq}, M_f^{\geq}, M_f^{\leq}$ (S a graded ring, M a graded S -module, f a homogeneous element of S_+): 8.2.1.

$\widehat{S}$ (S a graded ring): 8.2.2.

$\widehat{M}$ (M a graded S -module): 8.2.4.

$S_{[n]}, S^{\natural}, f^{\natural}$ (S a graded ring, f homogeneous in S): 8.2.6.

$M_{[n]}, M^{\natural}$ (M a graded S -module): 8.2.8.

$\mathcal{S}$ ($\mathcal{S}$ a graded $\mathcal{O}_Y$ -algebra): 8.3.1.

$\mathbf{G}_m$ (group scheme): 8.3.9.

$\mathcal{S}_X$ ($X = \text{Proj}(\mathcal{S})$, $\mathcal{S}$ a graded $\mathcal{O}_Y$ -algebra): 8.6.1.

$\mathcal{S}_X, C_X, \widehat{C}_X, E_X, \widehat{E}_X$ ($X = \text{Proj}(\mathcal{S})$, $\mathcal{S}$ a graded $\mathcal{O}_Y$ -algebra with $\mathcal{S}_0 = \mathcal{O}_Y$): 8.6.1.

$\mathcal{S}_{[n]}, \mathcal{S}^{\natural}$ ($\mathcal{S}$ a graded $\mathcal{O}_Y$ -algebra): 8.7.2.

$\mathcal{P}\text{roj}_0(\mathcal{M}), \mathcal{P}\text{roj}(\mathcal{M}), \mathcal{M}_X, \widehat{\mathcal{M}}, \mathcal{M}^{\square}$ ($\mathcal{M}$ a graded $\mathcal{S}$ -module): 8.12.1.

$\mathcal{M}_X^{\geq}$ ($X = \text{Proj}(\mathcal{S})$, $\mathcal{M}$ a graded $\mathcal{S}$ -module): 8.12.5.

$\mathcal{M}_{[n]}, \mathcal{M}^{\natural}$ ($\mathcal{M}$ a graded $\mathcal{S}$ -module): 8.12.9.

$(\widetilde{\mathcal{N}})^-$ ($\mathcal{N}$ a sub- $\mathcal{S}$ -module of a graded $\mathcal{S}$ -module): 8.13.1.

$\overline{\mathcal{N}}$ ($\mathcal{N}$ a sub- $\mathcal{S}$ -module of a graded $\mathcal{S}$ -module): 8.13.2.

$\Gamma_*(\mathcal{F})$ ($X = \text{Proj}(\mathcal{S})$, $\mathcal{F}$ a graded $\mathcal{S}_X$ -module): 8.14.5.

$\mathcal{S}_{X'}, \Gamma'_*(\mathcal{F}'), \mathcal{P}\text{roj}'(\mathcal{M}), \alpha', \beta'$ (X' open in $X = \text{Proj}(\mathcal{S})$, $\mathcal{F}'$ a graded $\mathcal{S}_{X'}$ -module, $\mathcal{M}$ a graded $\mathcal{S}$ -module): 8.14.6.

[II:209]

TERMINOLOGICAL INDEX

- Affine morphism: 1.6.1.
 Affine prescheme over a prescheme: 1.2.1.
 Graded algebra over a graded ring: 2.1.2.
 Symmetric algebra of an A -module: 1.7.1.
 Symmetric $\mathcal{A}$ -algebra of an $\mathcal{A}$ -module: 1.7.4.
 Essentially reduced graded $\mathcal{O}_Y$ -algebra: 3.1.12.
 Integral graded $\mathcal{O}_Y$ -algebra: 3.1.12.
 Ample invertible $\mathcal{O}$ -module: 4.5.3.
 Ample relative to f ; f -ample; Y -ample invertible $\mathcal{O}$ -module: 4.6.1.
 Essentially integral graded ring: 2.1.11.
 Essentially reduced graded ring: 2.1.10.
 Sheaf associated to an $\mathcal{O}$ -module: 1.4.3.
 Sheaf associated to a graded S -module: 2.5.3.
 Sheaf associated to a graded $\mathcal{S}$ -module: 3.2.2.
 S -scheme associated to an $\mathcal{O}_S$ -algebra: 1.3.1.
 Condition (TF); condition (TN): 2.7.2.
 Condition (TF); condition (TN): 3.4.2.
 Affine cone and projective cone defined by a graded $\mathcal{O}_Y$ -algebra: 8.3.1.
 Punctured affine cone and punctured projective cone defined by a graded $\mathcal{O}_Y$ -algebra: 8.3.4.
 Affine projecting cone and projective projecting cone of a prescheme $\text{Proj}(\mathcal{S})$: 8.3.1.
 Algebraic curve over a field: 7.4.2.
 Complete algebraic curve: 7.4.7.
 Serre's criterion: 5.2.1.
 Determinant of an endomorphism of a finite-type module over an integral ring: 6.4.2.
 Determinant of an endomorphism of a finite-type module over a reduced Noetherian ring: 6.4.7.
 Determinant of an endomorphism of a locally free $\mathcal{A}$ -module: 6.4.8.
 Determinant of an endomorphism of a finite-type $\mathcal{O}_X$ -module on a locally integral prescheme X : 6.4.9.
 Determinant of an endomorphism of a finite-type $\mathcal{O}_X$ -module on a locally Noetherian reduced prescheme X : 6.4.10.
 Blow-up prescheme: 8.1.3.
 Integral morphism: 6.1.1.
 Integral prescheme over a prescheme: 6.1.1.
 Integral quasi-coherent $\mathcal{O}_S$ -algebra: 6.1.2.
 Section of an $\mathcal{A}$ -algebra integral over $\mathcal{A}$: 6.3.1.
 Tautological sheaf of a projective bundle: 4.1.1.
 Integral closure of a sheaf of rings $\mathcal{A}$ in an $\mathcal{A}$ -algebra: 6.3.2.
 Integral closure of a prescheme X relative to an $\mathcal{O}_X$ -algebra: 6.3.4.
 Projective closure of an affine cone: 8.3.1.
 [II:210] Rational geometric fibre of a projective bundle over an extension K of $k(y)$: 4.2.6.
 Projective bundle defined by an $\mathcal{O}_Y$ -module: 4.1.1.
 Vector bundle defined by an $\mathcal{O}_S$ -module: 1.7.8.
 Finite morphism: 6.1.1.
 Finite prescheme over a prescheme: 6.1.1.
 Finite quasi-coherent $\mathcal{O}_S$ -algebra: 6.1.2.
 Elementary symmetric functions of an endomorphism of a locally free module: 6.4.1.
 Elementary symmetric functions of an endomorphism of a finite-type module over an integral ring: 6.4.2.
 Elementary symmetric functions of an endomorphism of a finite-type module over a reduced Noetherian ring: 6.4.7.

- Elementary symmetric functions of an endomorphism of a locally free $\mathcal{A}$ -module of constant rank: 6.4.8.
- Elementary symmetric functions of an endomorphism of a finite-type $\mathcal{O}_X$ -module on a locally integral prescheme X : 6.4.9.
- Elementary symmetric functions of an endomorphism of a finite-type $\mathcal{O}_X$ -module on a locally Noetherian reduced prescheme X : 6.4.10.
- Elementary symmetric functions of a section of an $\mathcal{A}$ -algebra: 6.5.1.
- Homogenization of a sub- $\mathcal{S}$ -module of a graded $\mathcal{S}$ -module: 8.13.2.
- Homomorphism of graded rings: 2.1.2.
- Homomorphism of degree k of graded modules: 2.1.2.
- Ideal of S_+ ; graded prime ideal of S_+ (S a graded ring): 2.1.10.
- Chow's lemma: 5.6.1.
- Locus at infinity of a projective cone: 8.3.3.
- Canonical morphism $G(\mathcal{E}) \rightarrow \text{Proj}(S)$, $S = \bigoplus_{n \geq 0} \Gamma(X, \mathcal{L}^{\otimes n})$: 4.5.1.
- Canonical morphism $X \rightarrow \text{Spec}(\Gamma(X, \mathcal{O}_X))$: 5.1.1.
- Segre morphism: 4.3.1.
- Nilradical of S_+ (S a graded ring): 2.1.10.
- Nilradical of $\mathcal{S}_+$ ($\mathcal{S}$ a quasi-coherent graded $\mathcal{O}_Y$ -algebra): 3.1.12.
- Norm of a section of an $\mathcal{A}$ -algebra: 6.5.1.
- Norm of an invertible $\mathcal{A}$ -module: 6.5.2.
- Norm of a section of an invertible $\mathcal{A}$ -module: 6.5.3.
- Norm of an invertible $\mathcal{O}_{X'}$ -module: 6.5.5.
- Periodic graded algebra: 8.14.12.
- Characteristic polynomial of an endomorphism of a finite-type module over an integral ring: 6.4.2.
- Characteristic polynomial of an endomorphism of a finite-type module over a reduced Noetherian ring: 6.4.7.
- Apex prescheme of an affine or projective cone: 8.3.3.
- Graded module having a finite presentation: 2.1.1.
- Graded $\mathcal{S}$ -module having a finite presentation: 3.1.1.
- Graded tensor product of two graded modules: 2.1.2.
- Projective morphism: 5.5.2.
- Projective prescheme over a prescheme: 5.5.2.
- Proper morphism: 5.4.1.
- Proper subset: 5.4.10.
- Proper prescheme over a prescheme: 5.4.1.
- Quasi-affine morphism: 5.1.1.
- Quasi-affine prescheme over a prescheme: 5.1.1.
- Quasi-affine scheme: 5.1.1.
- [II:211] Quasi-finite morphism: 6.2.3.
- Quasi-finite prescheme over a prescheme: 6.2.3.
- Quasi-projective morphism: 5.3.1.
- Quasi-projective prescheme over a prescheme: 5.3.1.
- Radical of an ideal of S_+ (S a graded ring): 2.1.10.
- Canonical retraction of a punctured projective cone onto its locus at infinity: 8.3.5.
- Canonical section of $\mathcal{O}_X(-1)$ (X a blow-up prescheme): 8.1.9.
- Zero section of an affine cone: 8.3.3.
- Zero section of a vector bundle: 1.7.9.
- Apex section of an affine or projective cone: 8.3.3.
- Spectrum of an $\mathcal{O}_S$ -algebra: 1.3.1.
- Homogeneous prime spectrum of a graded ring: 2.3.1.
- Homogeneous spectrum of a quasi-coherent graded $\mathcal{O}_S$ -algebra: 3.1.3.
- (TN)-injective, (TN)-surjective, or (TN)-bijective homomorphism of graded modules: 2.7.2.

(TN)-isomorphism of graded modules: 2.7.2.

(TN)-injective, (TN)-surjective, or (TN)-bijective homomorphism of graded rings: 2.9.1.

(TN)-isomorphism of graded rings: 2.9.1.

(TN)-injective, (TN)-surjective, or (TN)-bijective homomorphism of graded $\mathcal{S}$ -modules: 3.4.2.

(TN)-isomorphism of graded $\mathcal{S}$ -modules: 3.4.2.

(TN)-injective, (TN)-surjective, or (TN)-bijective homomorphism of graded $\mathcal{O}_Y$ -algebras: 3.6.1.

(TN)-isomorphism of graded $\mathcal{O}_Y$ -algebras: 3.6.1.

Spectral topology on $\text{Proj}(S)$: 2.3.3.

Very ample for q ; very ample for Y (invertible $\mathcal{O}$ -module): 4.4.2.

Graded $\mathcal{S}$ -module of finite type: 3.1.1.

Universally closed morphism: 5.4.9.

[II:213]

ORIGINAL TABLE OF CONTENTS

Unit	Title	Original page
II	Elementary global study of some classes of morphisms	5
§1	Affine morphisms	5
1.1	S -preschemes and $\mathcal{O}_S$ -algebras	5
1.2	Affine preschemes over a prescheme	6
1.3	Affine prescheme over S associated to an $\mathcal{O}_S$ -algebra	8
1.4	Quasi-coherent sheaves on a prescheme affine over S	9
1.5	Change of the base prescheme	12
1.6	Affine morphisms	14
1.7	Vector bundle associated to a sheaf of modules	14
§2	Homogeneous prime spectra	19
2.1	Generalities on graded rings and modules	19
2.2	Rings of fractions of a graded ring	23
2.3	Homogeneous prime spectrum of a graded ring	25
2.4	The scheme structure on $\text{Proj}(S)$	28
2.5	Sheaf associated to a graded module	30
2.6	Graded S -module associated to a sheaf on $\text{Proj}(S)$	36
2.7	Finiteness conditions	38
2.8	Functorial behaviour	41
2.9	Closed subpreschemes of a scheme $\text{Proj}(S)$	48
§3	Homogeneous spectrum of a sheaf of graded algebras	49
3.1	Homogeneous spectrum of a quasi-coherent graded $\mathcal{O}_Y$ -algebra	49
3.2	Sheaf on $\text{Proj}(\mathcal{S})$ associated to a graded $\mathcal{S}$ -module	54
3.3	Graded $\mathcal{S}$ -module associated to a sheaf on $\text{Proj}(\mathcal{S})$	56
3.4	Finiteness conditions	59
3.5	Functorial behaviour	61
3.6	Closed subpreschemes of a prescheme $\text{Proj}(\mathcal{S})$	64
3.7	Morphisms from a prescheme into a homogeneous spectrum	65
3.8	Criteria for immersion into a homogeneous spectrum	69
[II:214] §4	Projective bundles; ample sheaves	71
4.1	Definition of projective bundles	71
4.2	Morphisms from a prescheme into a projective bundle	72
4.3	The Segre morphism	76
4.4	Immersions into projective bundles; very ample sheaves	78
4.5	Ample sheaves	83
4.6	Relatively ample sheaves	89
§5	Quasi-affine morphisms; quasi-projective morphisms; proper morphisms; projective morphisms	94
5.1	Quasi-affine morphisms	94
5.2	Serre's criterion	97
5.3	Quasi-projective morphisms	99
5.4	Proper morphisms and universally closed morphisms	100
5.5	Projective morphisms	103
5.6	Chow's lemma	106
§6	Integral morphisms and finite morphisms	110
6.1	Preschemes integral over another	110
6.2	Quasi-finite morphisms	114
6.3	Integral closure of a prescheme	116
6.4	Determinant of an endomorphism of an $\mathcal{A}$ -module	120
6.5	Norm of an invertible sheaf	125
6.6	Application: criteria for ampleness	130
6.7	Chevalley's theorem	135
§7	Valuative criteria	138

7.1	Review of valuation rings	138
7.2	Valuative criterion for separatedness	141
7.3	Valuative criterion for properness	143
7.4	Algebraic curves and function fields of dimension one	148
§8	Blow-up schemes; projecting cones; projective closure	152
8.1	Blow-up preschemes	152
8.2	Preliminary results on localization in graded rings	157
8.3	Projecting cones	162
8.4	Projective closure of a vector bundle	168
8.5	Functorial behaviour	169
8.6	A canonical isomorphism for punctured cones	171
[II:215] 8.7	Blow-up of projecting cones	173
8.8	Ample sheaves and contractions	177
8.9	Grauert's ampleness criterion: statement	182
8.10	Grauert's ampleness criterion: proof	184
8.11	Uniqueness of contractions	189
8.12	Quasi-coherent sheaves on projecting cones	191
8.13	Projective closure of subsheaves and closed subschemes	195
8.14	Complements on sheaves associated to graded $\mathcal{S}$ -modules	197
	Bibliography (continued)	205
	Index of notation	207
	Terminological index	209
	Errata and Addenda (List 1)	217

Received July 22, 1960.

[II:217]

ERRATA AND ADDENDA

(List 1)

Chapter 0

(0, 2.1.3). On line 15 from the bottom of p. 21, replace “inter-section” by “intersection”.

(0, 2.1.6). On line 13 of p. 22, replace

$$Z_i \cap C \left(\bigcup_{j \neq i} Z_j \right) \quad \text{by} \quad \bigcup_{j \neq i} Z_j.$$

(0, 3.1.6). On line 17 from the bottom of p. 25, replace $\Gamma(U, \mathcal{F})$ by $\Gamma(V, \mathcal{F})$.(0, 4.1.1). On line 12 of p. 36, replace $\mathcal{G} \rightarrow \mathcal{F}$ by $\mathcal{B} \rightarrow \mathcal{A}$.

(0, 4.2.1). On line 11 from the bottom of p. 39, replace “sheaves” by “sheaf”.

(0, 5.1.2). After line 14 of p. 45, add:

If $\mathcal{G}$ is an $\mathcal{O}_Y$ -module generated by its sections over Y , then $f^*(\mathcal{G})$ is generated by its sections over X , since f^* is a right-exact functor.

(0, 5.2.4). On line 18 from the bottom of p. 46, replace f_U by f_U^* .

(0, 5.3.4). Before line 16 from the bottom of p. 47, add:

If

$$\mathcal{A} \longrightarrow \mathcal{B} \longrightarrow \mathcal{C} \longrightarrow \mathcal{D} \longrightarrow \mathcal{E}$$

is an exact sequence of $\mathcal{O}_X$ -modules, and if $\mathcal{A}, \mathcal{B}, \mathcal{D}, \mathcal{E}$ are coherent, then $\mathcal{C}$ is coherent.

(0, 5.4.1). After line 2 of p. 49, add:

Suppose that $\mathcal{O}_X$ is coherent, and let $\mathcal{F}$ be a coherent $\mathcal{O}_X$ -module. If, at a point $x \in X$, $\mathcal{F}_x$ is a free $\mathcal{O}_x$ -module of rank n , then there exists a neighbourhood U of x such that $\mathcal{F}|_U$ is locally free of rank n ; indeed, $\mathcal{F}_x$ is then isomorphic to $\mathcal{O}_x^n$, and the assertion follows from (0, 5.2.7).

(0, 6.6.2). Replace lines 10 and 11 of p. 59 by:

Indeed, this follows from (0, 6.4.1, d).

(0, 6.7.1). Before line 10 from the bottom of p. 59, add:

If f is a flat morphism, we also say that X is *flat over* Y , or *Y-flat*.

(0, 7.2.9). On line 11 from the bottom of p. 65, replace M_n by M_{n-1} .**Chapter I**(I, 1.2.1). On line 18 of p. 83, replace K by k (twice).(I, 1.2.7). On line 16 from the bottom of p. 84, replace A by A' (twice) and $\mathfrak{r}$ by $\mathfrak{r}'$; on line 17 from the bottom, replace X by X' .

(I, 1.7.3) and (I, 2.2.4). The statements given under these numbers can, in accordance with a remark of J. Tate, be generalized as follows.

1.8. Morphisms from locally ringed spaces to affine schemes

Proposition (1.8.1). — *Let $(S, \mathcal{O}_S)$ be an affine scheme and $(X, \mathcal{O}_X)$ a locally ringed space. There is a canonical bijection from the set of ring homomorphisms*

$$\Gamma(S, \mathcal{O}_S) \longrightarrow \Gamma(X, \mathcal{O}_X)$$

to the set of morphisms of ringed spaces $(\psi, \theta) : (X, \mathcal{O}_X) \rightarrow (S, \mathcal{O}_S)$ such that, for every $x \in X$, $\theta_x^\# : \mathcal{O}_{\psi(x)} \rightarrow \mathcal{O}_x$ is a local homomorphism.

[II:218]

Proof. First note that, if $(X, \mathcal{O}_X)$ and $(S, \mathcal{O}_S)$ are any two ringed spaces, a morphism (ψ, θ) from $(X, \mathcal{O}_X)$ to $(S, \mathcal{O}_S)$ canonically defines a ring homomorphism

$$\Gamma(\theta) : \Gamma(S, \mathcal{O}_S) \longrightarrow \Gamma(X, \mathcal{O}_X),$$

and hence a first map

$$(1.8.1.1) \quad \rho : \text{Hom}((X, \mathcal{O}_X), (S, \mathcal{O}_S)) \longrightarrow \text{Hom}(\Gamma(S, \mathcal{O}_S), \Gamma(X, \mathcal{O}_X)).$$

Conversely, under the hypotheses of the proposition, put $A = \Gamma(S, \mathcal{O}_S)$ and consider a ring homomorphism $\phi : A \rightarrow \Gamma(X, \mathcal{O}_X)$. For every $x \in X$, the set of $f \in A$ such that $\phi(f)(x) = 0$ is a prime ideal of A , since $\mathcal{O}_x/\mathfrak{m}_x = k(x)$ is a field. It is therefore an element of $S = \text{Spec}(A)$, which we again denote by ${}^a\phi(x)$. Moreover, for every $f \in A$, by definition (0, 5.5.2),

$${}^a\phi^{-1}(D(f)) = X_f,$$

which proves that ${}^a\phi$ is a continuous map $X \rightarrow S$.

We next define a homomorphism of $\mathcal{O}_S$ -modules

$$\tilde{\phi} : \mathcal{O}_S \rightarrow {}^a\phi_*(\mathcal{O}_X).$$

For every $f \in A$, one has $\Gamma(D(f), \mathcal{O}_S) = A_f$ (I, 1.3.6); to $s/f \in A_f$ associate

$$(\phi(s)|X_f)(\phi(f)|X_f)^{-1} \in \Gamma(X_f, \mathcal{O}_X) = \Gamma(D(f), {}^a\phi_*(\mathcal{O}_X)).$$

Passing from $D(f)$ to $D(fg)$ shows at once that this defines a homomorphism of sheaves and hence a morphism of ringed spaces $({}^a\phi, \tilde{\phi})$. With the same notation, and writing $y = {}^a\phi(x)$, one has (0, 3.7.1)

$$\tilde{\phi}_x^\#(s_y/f_y) = (\phi(s)_x)(\phi(f)_x)^{-1}.$$

Since $s_y \in \mathfrak{m}_y$ is, by definition, equivalent to $\phi(s)_x \in \mathfrak{m}_x$, the homomorphism $\tilde{\phi}_x^\# : \mathcal{O}_y \rightarrow \mathcal{O}_x$ is local. We have therefore defined a second map

$$\sigma : \text{Hom}(\Gamma(S, \mathcal{O}_S), \Gamma(X, \mathcal{O}_X)) \rightarrow \mathfrak{L},$$

where $\mathfrak{L}$ denotes the set of morphisms $(\psi, \theta) : (X, \mathcal{O}_X) \rightarrow (S, \mathcal{O}_S)$ for which $\theta_x^\#$ is local for every $x \in X$.

It remains to prove that σ and the restriction of ρ to $\mathfrak{L}$ are inverses. The definition of $\tilde{\phi}$ immediately gives $\Gamma(\tilde{\phi}) = \phi$, and hence $\rho \circ \sigma$ is the identity. To see that $\sigma \circ \rho$ is the identity, begin with $(\psi, \theta) \in \mathfrak{L}$ and put $\phi = \Gamma(\theta)$. The hypothesis on $\theta_x^\#$ gives, after passage to residue fields, a monomorphism

$$\theta^x : k(\psi(x)) \rightarrow k(x)$$

such that, for every $f \in A = \Gamma(S, \mathcal{O}_S)$,

$$\theta^x(f(\psi(x))) = \phi(f)(x).$$

Thus $f(\psi(x)) = 0$ is equivalent to $\phi(f)(x) = 0$, which proves that ${}^a\phi = \psi$. On the other hand, the definitions make the diagram

$$\begin{array}{ccc} A & \xrightarrow{\phi} & \Gamma(X, \mathcal{O}_X) \\ \downarrow & & \downarrow \\ A_{\psi(x)} & \xrightarrow{\theta_x^\#} & \mathcal{O}_x \end{array}$$

commutative; the analogous diagram with $\theta_x^\#$ replaced by $\tilde{\phi}_x^\#$ is also commutative. It follows from (0, 1.2.4) that $\tilde{\phi}_x^\# = \theta_x^\#$, and consequently that $\tilde{\phi} = \theta$. $\square$

(1.8.2). When $(X, \mathcal{O}_X)$ and $(Y, \mathcal{O}_Y)$ are locally ringed spaces, we shall consider morphisms $(\psi, \theta) : (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ such that, for every $x \in X$, $\theta_x^\# : \mathcal{O}_{\psi(x)} \rightarrow \mathcal{O}_x$ is a local homomorphism.

[II:219] Whenever we speak henceforth of a *morphism of locally ringed spaces*, we shall always mean a morphism of this kind. With this definition of morphisms, locally ringed spaces form a category. For two objects X, Y of this category, $\text{Hom}(X, Y)$ will therefore denote the set of morphisms of locally ringed spaces from X to Y (the set denoted $\mathfrak{L}$ in (1.8.1)). When we wish to consider the set of morphisms of ringed spaces from X to Y , we shall write $\text{Hom}_{\text{rs}}(X, Y)$ to avoid ambiguity. The map (1.8.1.1) is thus written

$$(1.8.2.1) \quad \rho : \text{Hom}_{\text{rs}}(X, Y) \rightarrow \text{Hom}(\Gamma(Y, \mathcal{O}_Y), \Gamma(X, \mathcal{O}_X)),$$

and its restriction

$$(1.8.2.2) \quad \rho' : \text{Hom}(X, Y) \rightarrow \text{Hom}(\Gamma(Y, \mathcal{O}_Y), \Gamma(X, \mathcal{O}_X))$$

is functorial in X and Y in the category of locally ringed spaces.

Corollary (1.8.3). — *Let $(Y, \mathcal{O}_Y)$ be a locally ringed space. For Y to be an affine scheme, it is necessary and sufficient that, for every locally ringed space $(X, \mathcal{O}_X)$, the map (1.8.2.2) be bijective.*

Proof. Proposition (1.8.1) shows that the condition is necessary. Conversely, suppose it holds and put $A = \Gamma(Y, \mathcal{O}_Y)$. The hypothesis and (1.8.1) show that the functors

$$X \longmapsto \text{Hom}(X, Y) \quad \text{and} \quad X \longmapsto \text{Hom}(X, \text{Spec}(A))$$

from locally ringed spaces to sets are isomorphic. This implies the existence of a canonical isomorphism $Y \rightarrow \text{Spec}(A)$ (cf. 0, 8). $\square$

(1.8.4). Let $S = \text{Spec}(A)$ be an affine scheme. Denote by (S', A') the ringed space whose underlying space consists of one point and whose structure sheaf A' is the (necessarily simple) sheaf on S' defined by the ring A . Let $\pi : S \rightarrow S'$ be the unique map. For every open subset U of S , the canonical map

$$\Gamma(S', A') = \Gamma(S, \mathcal{O}_S) \longrightarrow \Gamma(U, \mathcal{O}_S)$$

defines a π -morphism $\iota : A' \rightarrow \mathcal{O}_S$ of sheaves of rings. Thus there is a canonical morphism of ringed spaces

$$i = (\pi, \iota) : (S, \mathcal{O}_S) \longrightarrow (S', A').$$

For every A -module M , denote by M' the simple sheaf on S' defined by M , which is evidently an A' -module. It is clear that $i_*(\tilde{M}) = M'$ (I, 1.3.7).

Lemma (1.8.5). — *With the notation of (1.8.4), for every A -module M , the canonical functorial $\mathcal{O}_S$ -homomorphism (0, 4.4.3.3)*

$$(1.8.5.1) \quad i^*(i_*(\tilde{M})) \longrightarrow \tilde{M}$$

is an isomorphism.

Proof. Both sides of (1.8.5.1) are right exact in M (the functor $M \mapsto i_*(\tilde{M})$ being evidently exact) and commute with direct sums. By viewing M as the cokernel of a homomorphism $A^{(I)} \rightarrow A^{(J)}$, one is reduced to the case $M = A$, where the assertion is immediate. $\square$

Corollary (1.8.6). — *Let $(X, \mathcal{O}_X)$ be a ringed space and let $u : X \rightarrow S$ be a morphism of ringed spaces.*

[II:220] *For every A -module M , there is, with the notation of (1.8.4), a canonical functorial isomorphism of $\mathcal{O}_X$ -modules*

$$(1.8.6.1) \quad u^*(\tilde{M}) \simeq u^*(i^*(M')).$$

Corollary (1.8.7). — *Under the hypotheses of (1.8.6), for every A -module M and every $\mathcal{O}_X$ -module $\mathcal{F}$, there is a canonical isomorphism, functorial in M and $\mathcal{F}$,*

$$(1.8.7.1) \quad \text{Hom}_{\mathcal{O}_S}(\tilde{M}, u_*(\mathcal{F})) \simeq \text{Hom}_A(M, \Gamma(X, \mathcal{F})).$$

Proof. By (0, 4.4.3) and Lemma (1.8.5), there is a canonical isomorphism of bifunctors

$$\text{Hom}_{\mathcal{O}_S}(\tilde{M}, u_*(\mathcal{F})) \simeq \text{Hom}_{A'}(M', i_*(u_*(\mathcal{F}))).$$

The right-hand side is plainly $\text{Hom}_A(M, \Gamma(X, \mathcal{F}))$. Under the canonical homomorphism (1.8.7.1), an $\mathcal{O}_S$ -homomorphism $h : \tilde{M} \rightarrow u_*(\mathcal{F})$ (equivalently, a u -morphism $\tilde{M} \rightarrow \mathcal{F}$) corresponds to

$$\Gamma(h) : M \longrightarrow \Gamma(S, u_*(\mathcal{F})) = \Gamma(X, \mathcal{F}).$$

$\square$

(1.8.8). With the notation of (1.8.4), it is clear from (0, 4.1.1) that every morphism of ringed spaces $(\psi, \theta) : X \rightarrow S'$ is equivalent to the data of a ring homomorphism $A \rightarrow \Gamma(X, \mathcal{O}_X)$. Proposition (1.8.1) may therefore be interpreted as giving a canonical bijection

$$\text{Hom}(X, S) \simeq \text{Hom}(X, S'),$$

where the right-hand side means morphisms of ringed spaces, since A is not in general a local ring.

More generally, let X, Y be locally ringed spaces and let (Y', A') be the ringed space whose underlying space is one point and whose sheaf of rings is the simple sheaf defined by $\Gamma(Y, \mathcal{O}_Y)$. Then (1.8.2.1) may be interpreted as a map

$$(1.8.8.1) \quad \rho : \text{Hom}_{\text{rs}}(X, Y) \longrightarrow \text{Hom}(X, Y').$$

Corollary (1.8.3) says that affine schemes are characterized among locally ringed spaces as those for which the restriction of ρ to $\text{Hom}(X, Y)$,

$$(1.8.8.2) \quad \rho' : \text{Hom}(X, Y) \longrightarrow \text{Hom}(X, Y'),$$

is bijective for every locally ringed space X . In a later chapter we shall generalize this definition, thereby associating to an arbitrary ringed space Z (and not only to one whose underlying space is a point) a locally ringed space which we shall again call $\text{Spec}(Z)$. This will be the starting point of a “relative” theory of preschemes over arbitrary ringed spaces, extending the results of Chapter I.

(1.8.9). The pairs $(X, \mathcal{F})$, where X is a locally ringed space and $\mathcal{F}$ an $\mathcal{O}_X$ -module, form a category: a morphism $(X, \mathcal{F}) \rightarrow (Y, \mathcal{G})$ is a pair (u, h) consisting of a morphism of locally ringed spaces

[II:221] $u : X \rightarrow Y$ and a u -morphism $h : \mathcal{G} \rightarrow \mathcal{F}$ of modules. For fixed $(X, \mathcal{F})$ and $(Y, \mathcal{G})$, these morphisms form a set denoted $\text{Hom}((X, \mathcal{F}), (Y, \mathcal{G}))$. The assignment $(u, h) \mapsto (\rho'(u), \Gamma(h))$ is a canonical map

$$(1.8.9.1) \quad \text{Hom}((X, \mathcal{F}), (Y, \mathcal{G})) \longrightarrow \text{Hom}((\Gamma(Y, \mathcal{O}_Y), \Gamma(Y, \mathcal{G})), (\Gamma(X, \mathcal{O}_X), \Gamma(X, \mathcal{F}))),$$

functorial in $(X, \mathcal{F})$ and $(Y, \mathcal{G})$; the right-hand side is the set of di-homomorphisms corresponding to the rings and modules in question (0, 1.0.2).

Corollary (1.8.10). — *Let Y be a locally ringed space and $\mathcal{G}$ an $\mathcal{O}_Y$ -module. For Y to be an affine scheme and $\mathcal{G}$ a quasi-coherent $\mathcal{O}_Y$ -module, it is necessary and sufficient that, for every pair $(X, \mathcal{F})$ consisting of a locally ringed space X and an $\mathcal{O}_X$ -module $\mathcal{F}$, the canonical map (1.8.9.1) be bijective.*

We leave to the reader the task of spelling out the proof, which is modelled on that of (1.8.3) and uses (1.8.1) and (1.8.7).

Remark (1.8.11). — The statements (I, 1.7.3), (I, 1.7.4), and (I, 2.2.4), as well as the definition (I, 1.6.1), are special cases of (1.8.1); similarly, (I, 2.2.5) follows from (1.8.7). Corollary (1.8.7) also gives (I, 1.6.3), and hence (I, 1.6.4), as a special case. Indeed, if X is affine and $\Gamma(X, \mathcal{F}) = N$, the functors

$$M \mapsto \text{Hom}_{\mathcal{O}_S}(\tilde{M}, u_*(\tilde{N})) \quad \text{and} \quad M \mapsto \text{Hom}_{\mathcal{O}_S}(\tilde{M}, (N_{[\phi]})^\sim)$$

(where $\phi : A \rightarrow \Gamma(X, \mathcal{O}_X)$ corresponds to u) are isomorphic by (1.8.7) and (I, 1.3.8). Finally, (I, 1.6.5), and hence (I, 1.6.6), follows from (1.8.6) and from the fact that, for every $f \in A$, the A_f -modules

$$N' \otimes_{A'} A_f \quad \text{and} \quad (N' \otimes_{A'} A)_f$$

(with the notation of (I, 1.6.5)) are canonically isomorphic.

Insert after (I, 3.2.8):

(3.2.9). It follows from (1.8.1) that (I, 3.2.2) may be sharpened as follows. The prescheme

$$Z = \text{Spec}(B \otimes_A C)$$

is not only a product of $X = \text{Spec}(B)$ and $Y = \text{Spec}(C)$ in the category of S -preschemes, but also in the category of locally ringed spaces over S (with the definition of S -morphisms modelled on (I, 2.5.2)). The proof of (I, 3.2.6) also shows that, for arbitrary S -preschemes X, Y , the prescheme $X \times_S Y$ is not only their product in the category of S -preschemes, but also in the category of locally ringed spaces over the prescheme S .

(I, 4.4.5). On line 14 from the bottom of p. 126, replace B by A ; on line 13 from the bottom, replace “ A -algebra” by “ B -algebra”.

(I, 5.3.5). On line 2 from the bottom of p. 132, replace $Y \rightarrow S$ by $g : Y \rightarrow S$.

(I, 6.3.2.1). On line 6 from the bottom of p. 144, read

$$D(g_i) \subset W.$$

(I, 7.3.8). Replace the present text, which is erroneous, by the following:

Let X, Y be integral preschemes; then $\mathcal{R}(X)$ and $\mathcal{R}(Y)$ are respectively a quasi-coherent $\mathcal{O}_X$ -module and a quasi-coherent $\mathcal{O}_Y$ -module (I, 7.3.3). If $f : X \rightarrow Y$ is a dominant morphism, there is a canonical homomorphism of $\mathcal{O}_X$ -modules

$$(7.3.8.1) \quad \tau : f^*(\mathcal{R}(Y)) \longrightarrow \mathcal{R}(X).$$

[II:222] First suppose that $X = \text{Spec}(A)$ and $Y = \text{Spec}(B)$ are affine, with A, B integral domains. Then f corresponds to an injective homomorphism $B \rightarrow A$, which extends to a monomorphism $L \rightarrow K$ from the fraction field L of B into the fraction field K of A . The homomorphism (7.3.8.1) then corresponds to the canonical homomorphism

$$L \otimes_B A \longrightarrow K$$

of (I, 1.6.5).

In the general case, for every pair of nonempty affine open subsets $U \subset X$, $V \subset Y$ such that $f(U) \subset V$, the preceding construction defines a homomorphism $\tau_{U,V}$. If $U' \subset U$, $V' \subset V$, and $f(U') \subset V'$, then $\tau_{U,V}$ extends $\tau_{U',V'}$; the assertion follows by gluing. If x, y are the generic points of X, Y , respectively, then $f(x) = y$ and

$$(f^*(\mathcal{R}(Y)))_x = \mathcal{O}_y \otimes_{\mathcal{O}_y} \mathcal{O}_x = \mathcal{O}_x$$

by (0, 4.3.1), so τ_x is an isomorphism.

(I, 9.5.2). In the last three lines of p. 176 and the first five lines of p. 177, replace X' everywhere by Y' and X'' by Y'' .

(I, 10.5.4). On line 11 from the bottom of p. 187, replace (10.3.4) by (10.3.5).

(I, 10.6.3). On line 3 from the bottom of p. 189, replace X by $\mathfrak{X}$.

(I, 10.14.2). On line 5 of p. 210, replace “coherent $\mathcal{O}_X$ -module” by “coherent ideal of $\mathcal{O}_X$ ”; on line 7, replace “coherent $\mathcal{O}_X$ -modules” by “coherent ideals of $\mathcal{O}_X$ ”.