

The Weil conjecture for K3 surfaces

Pierre Deligne* (Bures-sur-Yvette)

1. Statement of the theorem

Let $\mathbb{F}_q$ be a field with q elements, let $\overline{\mathbb{F}}_q$ be an algebraic closure of $\mathbb{F}_q$, let $\varphi \in \text{Gal}(\overline{\mathbb{F}}_q/\mathbb{F}_q)$ be the Frobenius substitution $x \mapsto x^q$, and let $F = \varphi^{-1}$ be the “geometric Frobenius”.

Let X be a scheme, separated and of finite type, over $\mathbb{F}_q$, and let $\overline{X}$ be the scheme over $\overline{\mathbb{F}}_q$ obtained from it by extension of scalars. For every closed point x of X , let $\deg(x) = [k(x) : \mathbb{F}_q]$ be the degree of the residue-field extension over $\mathbb{F}_q$. The zeta function $Z(X, t) \in \mathbb{Z}[[t]]$ is defined by

$$Z(X, t) = \prod_{x \in X} (1 - t^{\deg(x)})^{-1} \quad (x \text{ a closed point of } X),$$

$$\log Z(X, t) = \sum_{n>0} \#X(\mathbb{F}_{q^n}) \frac{t^n}{n}.$$

For every prime number ℓ prime to q , the ℓ -adic cohomology with proper supports $H_c^i(\overline{X}, \mathbb{Q}_\ell)$ of $\overline{X}$ is a finite-dimensional vector space over $\mathbb{Q}_\ell$, on which $\text{Gal}(\overline{\mathbb{F}}_q/\mathbb{F}_q)$ acts by transport of structure. By Grothendieck, one has

$$Z(X, t) = \prod_i \det(1 - Ft, H_c^i(\overline{X}, \mathbb{Q}_\ell))^{(-1)^{i+1}}.$$

Take X to be a nonsingular projective surface. In this case it is known that the factors

$$P_i(t) = \det(1 - Ft, H^i(\overline{X}, \mathbb{Q}_\ell))$$

of $Z(X, t)$ are polynomials with rational integral coefficients independent of ℓ . The Weil conjecture asserts here that the roots of $P_i(t)$ have complex absolute value $q^{-i/2}$. This conjecture is invariant under finite extension of the finite base field. For X as above, and $i \neq 2$, it follows from Weil [9]. If, for simplicity, X is assumed geometrically connected, one indeed has

$$P_0(t) = 1 - t, \quad P_4(t) = 1 - q^2 t,$$

$$P_1(t) = \det(1 - \varphi t; T_\ell(\text{Pic}^0(X)_{\text{red}})), \quad P_3(t) = P_1(qt).$$

Let k be a field of characteristic 0 and let X be a nonsingular projective geometrically connected surface over k . We say that X is a K3 surface if X is regular and has trivial canonical divisor, that is, if $H^1(X, \mathcal{O}) = 0$ and Ω_X^2 is isomorphic to $\mathcal{O}_X$. For the miraculous properties of these surfaces, we refer to [8]. We shall mainly use the following facts.

1.1. Lemma. (i) One has $h^{2,0} = h^{0,2} = 1$.

(ii) One has $H^2(X, T_X^1) = 0$; infinitesimal deformations are unobstructed. The map induced by interior product

$$H^1(X, T_X^1) \otimes H^0(X, \Omega_X^2) \xrightarrow{\sim} H^1(X, \Omega_X^1) \tag{1.1.1}$$

is bijective, and $H^0(X, T_X^1) = 0$; X has no infinitesimal automorphisms.

One has

$$h^{2,0} =_{\text{dfn}} \dim H^0(X, \Omega_X^2) = \dim H^0(X, \mathcal{O}_X) = 1,$$

*This article was written during a stay at the University of Warwick, which I thank for its hospitality.

and (i) follows by Hodge symmetry. Since $H^1(X, \mathcal{O}_X) = 0$, Hodge symmetry and Serre duality give

$$h^{0,1} = h^{1,0} = h^{2,1} = h^{1,2} = 0.$$

Since $\Omega_X^2 = \mathcal{O}_X$, the interior-product isomorphism

$$T_X^1 \otimes \Omega_X^2 \xrightarrow{\sim} \Omega_X^1$$

induces isomorphisms $T_X^1 \simeq \Omega_X^1$. The vanishing of $H^2(X, T_X^1)$ therefore follows from that of $h^{1,2}$, and it is clear that (1.1.1) is bijective. The same argument gives $H^0(X, T_X^1) = 0$.

1.2. Lemma. *Let Y_0 be a nonsingular projective surface over a finite field $\mathbb{F}_q$. The following conditions are equivalent.*

- (i) *There exist a complete discrete valuation ring V of unequal characteristic, with finite residue field k , a proper smooth morphism $f_V : Y \rightarrow \text{Spec}(V)$ whose generic fiber is a K3 surface, and $i : \mathbb{F}_q \rightarrow k$ such that $Y_0 \otimes_{\mathbb{F}_q} k \simeq Y \otimes_V k$.*
- (ii) *There exist an integral scheme S whose function field has characteristic 0, a point $s \in S$, a proper smooth morphism $f : Y \rightarrow S$ whose generic fiber is a K3 surface, and $i : \mathbb{F}_q \rightarrow k(s)$ such that $Y_0 \otimes_{\mathbb{F}_q} k(s) \simeq Y_s$.*

The proof uses a standard passage to the limit. If the equivalent conditions of 1.2 are satisfied, we say that Y_0 lifts to a K3 surface.

1.3. Theorem. *A nonsingular projective surface over a finite field $\mathbb{F}_q$ which lifts to a K3 surface satisfies the Weil conjecture.*

This theorem was obtained independently by Pjateckii-Shapiro and Safarevitch. It applies, for example, to surfaces of degree 4 in $\mathbb{P}^3$. The Betti numbers of a K3 surface are $(1, 0, 22, 0, 1)$; the theorem therefore implies that

$$|\#X(\mathbb{F}_q) - \#\mathbb{P}^2(\mathbb{F}_q)| \leq 21q.$$

Grothendieck's conjectural theory of motives, in particular his theory of the “motivic Galois group”, was very useful to me in constructing the proof. In that language, which will not be used below, the idea is roughly to construct, by reduction from $\mathbb{C}$ to $\mathbb{F}_q$, an abelian variety A and an injective “morphism” of motives

$$H^2(X) \hookrightarrow H^1(A) \otimes H^1(A),$$

in order to deduce the Weil conjecture for X from Weil's theorem for A . However, we do not define this “morphism” by an algebraic cycle; it is therefore not a morphism in the sense of [5]. In Hodge theory, its mode of definition has the following analogue, which will not be used below.

1.4. Lemma. *Let H be a variation of Hodge structures of weight n on a scheme S . If the local system $H_{\mathbb{Z}}$ has no global sections except the section s and its multiples, then at every point of S the section s is of type $(n/2, n/2)$.*

1.5. Notation. 1.5.1. Scalar extensions will be denoted by subscripts: if M_A is a module over a commutative ring with unit A , or a scheme over A , for example an algebraic group over A , then M_B will denote the module over the A -algebra B , or the scheme over B , obtained by extension of scalars.

1.5.2. If a group G acts on a set X , $g * x$ denotes the transform of x by g .

1.5.3. When an equality is the definition of one of its sides, it will be denoted $=_{\text{dfn}}$.

2. Polarized Hodge structures

To handle Hodge structures, we shall use the formalism described in [4] and briefly recalled below.

2.1. Let $\underline{S}$ be the real algebraic group of invertible elements of the $\mathbb{R}$ -algebra $\mathbb{C}$, let $w : \mathbb{G}_{m,\mathbb{R}} \rightarrow \underline{S}$ be the inclusion of $\mathbb{R}^*$ in $\mathbb{C}^*$, and let $t : \underline{S} \rightarrow \mathbb{G}_{m,\mathbb{R}}$ be the inverse of the norm; then $tw(x) = x^{-2}$. Let V be a finite-dimensional real vector space. It is equivalent to give:

- a) $\rho : \underline{S} \rightarrow \mathrm{GL}(V)$ of weight n : $\rho w(x) * v = x^n v$;
- b) a Hodge bigrading of weight n of $V_{\mathbb{C}} = V \otimes_{\mathbb{R}} \mathbb{C}$:

$$V_{\mathbb{C}} = \bigoplus_{p+q=n} V^{p,q}, \quad \overline{V^{p,q}} = V^{q,p};$$

- c) a Hodge filtration of weight n of $V_{\mathbb{C}} = V \otimes_{\mathbb{R}} \mathbb{C}$: this is a decreasing filtration such that

$$F^p(V_{\mathbb{C}}) \oplus \overline{F^{n-p+1}(V_{\mathbb{C}})} \xrightarrow{\sim} V_{\mathbb{C}}.$$

One has

$$F^p(V_{\mathbb{C}}) = \bigoplus_{p' \geq p} V^{p',q'};$$

for $z \in \underline{S}(\mathbb{R}) = \mathbb{C}^*$ and $v^{p,q} \in V^{p,q}$, one has

$$z * v^{p,q} = z^p \bar{z}^q v^{p,q}.$$

Put $C = \rho(i)$: this is Weil's operator C . One has $Cv^{p,q} = i^{p-q}v^{p,q}$. For $\mathcal{S} \subset \mathbb{Z} \times \mathbb{Z}$, a Hodge bigrading is said to be of type $\mathcal{S}$ if the Hodge numbers $h^{p,q} = \dim V^{p,q}$ vanish for $(p,q) \notin \mathcal{S}$.

2.2. Here a Hodge structure H of weight n will mean a free finite-type $\mathbb{Z}$ -module $H_{\mathbb{Z}}$ together with a Hodge bigrading of weight n of $H_{\mathbb{C}} = H_{\mathbb{Z}} \otimes \mathbb{C}$. The Tate Hodge structure $\mathbb{Z}(n)$ is the Hodge structure of type $\{(-n, -n)\}$ given by

$$\mathbb{Z}(n)_{\mathbb{C}} = \mathbb{C}, \quad \mathbb{Z}(n)_{\mathbb{Z}} = (2\pi i)^n \mathbb{Z} \subset \mathbb{C}.$$

A polarization of a Hodge structure H of weight n is a morphism of Hodge structures

$$\psi : H \otimes H \longrightarrow \mathbb{Z}(-n)$$

such that, on $H_{\mathbb{R}} = H_{\mathbb{Z}} \otimes \mathbb{R}$, the real bilinear form

$$(2\pi i)^n \psi(x, Cy)$$

is symmetric and positive definite. The form ψ is then $(-1)^n$ -symmetric.

A rational Hodge structure of weight n consists of a $\mathbb{Q}$ -vector space $H_{\mathbb{Q}}$ and a Hodge bigrading of weight n of $H_{\mathbb{C}} = H_{\mathbb{Q}} \otimes \mathbb{C}$. Let $\mathbb{Q}(n)$ denote the rational Hodge structure underlying $\mathbb{Z}(n)$. The definition of polarizations is left to the reader.

The following statement reformulates a theorem of Riemann.

2.3. Scholium. *The functor $A \mapsto H^1(A, \mathbb{Z})$ identifies polarized abelian varieties with polarized Hodge structures of type $\{(0, 1), (1, 0)\}$.*

From the end of this section, we shall use only 2.11(iii') $\Rightarrow$ (ii). To prove this implication, one may restrict to connected groups, which makes 2.5 useless or evident.

2.4. Let G be a real linear algebraic group. We say that G is compact if $G(\mathbb{R})$ is compact and every connected component of $G(\mathbb{C})$ has a real point. The evident functor is then an equivalence from the category of algebraic linear representations of G to the category of linear representations of the compact group $G(\mathbb{R})$.

2.5. Lemma. *Every real algebraic subgroup H of a compact real algebraic group G is compact.*

It is known that the map

$$(g, \ell) \mapsto g \cdot \exp(i\ell) : G(\mathbb{R}) \times \text{Lie}(G) \longrightarrow G(\mathbb{C})$$

is bijective. It is enough to prove that, for every real subgroup H of G and every $h = g \cdot \exp(i\ell) \in H(\mathbb{C})$, one has $\ell \in \text{Lie}(H)$. Since $\bar{h} \in H(\mathbb{C})$, one has

$$\exp(2i\ell) = \bar{h}^{-1}h \in H(\mathbb{C}).$$

Regard $G(\mathbb{C})$ as the set of real points of a real algebraic group. Let $\bar{J} \subset H(\mathbb{C})$ be the Zariski closure, in this group, of $J = \{\exp(ni\ell) \mid n \in 2\mathbb{Z}\}$. The elements $g \in J$, and hence the elements $g \in \bar{J}$, satisfy $g^{-1} = \bar{g}$, whence $\text{Lie}(\bar{J}) \subset i\text{Lie}(G)$. The Lie algebra $\text{Lie}(\bar{J})$ is abelian, and $\exp(\text{Lie}(\bar{J}))$ is a group, necessarily of finite index in $\bar{J}$. There are therefore n and $\ell' \in \text{Lie}(\bar{J})$ such that

$$\exp(ni\ell) = \exp(\ell').$$

Thus $i\ell \in \text{Lie}(\bar{J}) \subset \text{Lie}(H)$ and $\ell \in \text{Lie}(H)$, completing the proof.

2.6. Lemma. *Let G be a real algebraic group. The following conditions are equivalent:*

- (i) G is compact;
- (ii) for every real, respectively complex, linear representation of G , there exists a positive-definite G -invariant symmetric bilinear form, respectively hermitian sesquilinear form;
- (iii) for one faithful representation of G , there exists a form as in (ii).

For G connected, these conditions are also equivalent to

- (iii') for one representation of G with finite kernel, there exists a form as in (ii).

For G compact, G -invariance is equivalent to $G(\mathbb{R})$ -invariance, and (ii) is classical. If (iii) holds, G is a subgroup of an orthogonal or unitary group, and one applies 2.5. Condition (iii') implies that $G(\mathbb{R})$ is compact, as a finite cover of a compact group; if G is connected, G is therefore compact.

2.7. Let G be a real linear algebraic group. For every involutive automorphism σ of G , let $G^{(\sigma)}$ be the real form of $G_{\mathbb{C}}$ defined by the antilinear involution $g \mapsto \sigma(\bar{g})$. We say that σ is a Cartan involution if $G^{(\sigma)}$ is compact.

Let C be an element of $G(\mathbb{R})$ whose square is central. A real, respectively complex, representation of G will be called C -polarizable if there exists on V a G -invariant bilinear, respectively sesquilinear, form ψ such that $\psi(x, Cy)$ is symmetric, respectively hermitian, and positive definite. We then say that ψ is a C -polarization of V . A special case of the following lemma is stated without proof in [4].

2.8. Lemma. *a) Whether ψ is a C -polarization of V depends only on the $G(\mathbb{R})$ -conjugacy class of C .*

b) The following conditions are equivalent:

- (i) $\text{ad } C$ is a Cartan involution of G ;
- (ii) every real, respectively complex, representation of G is C -polarizable;
- (iii) G has a faithful real, respectively complex, C -polarizable representation.

If G is connected, these conditions are also equivalent to

- (iii') G has a real, respectively complex, C -polarizable representation with finite kernel.

To prove a), note that $\psi(x, gCg^{-1}y) = \psi(g^{-1}x, Cg^{-1}y)$. A sesquilinear form ψ on a complex representation of G is G -invariant if and only if

$$\psi(gx, \bar{g}y) = \psi(x, y) \quad \text{for } g \in G(\mathbb{C}).$$

This condition is equivalent to

$$\psi(gx, CC^{-1}\bar{g}Cy) = \psi(x, Cy) \quad \text{for } g \in G(\mathbb{C})$$

(replace y by Cy), that is, to the $G^{(\text{ad } C)}$ -invariance of $\psi(x, Cy)$. The hermitian form in 2.8 therefore follows from 2.6. The non-hermitian form follows from it:

- a) the complexification, respectively realification, of a real, respectively complex, C -polarizable representation is C -polarizable;

- b) a polarization of V induces a polarization of every subrepresentation of V ;
- c) for V real, respectively complex, one has $V \hookrightarrow V \otimes_{\mathbb{R}} \mathbb{C}$.

2.9. Let G be a reductive group over $\mathbb{Q}$, and let t and w be morphisms defined over $\mathbb{Q}$

$$\mathbb{G}_m \xrightarrow{w} G \xrightarrow{t} \mathbb{G}_m,$$

with $tw(x) = x^{-2}$ and w central. Suppose also given a commutative diagram

$$\begin{array}{ccccc} \mathbb{G}_{m,\mathbb{R}} & \xrightarrow{w} & \underline{S} & \xrightarrow{t} & \mathbb{G}_{m,\mathbb{R}} \\ \parallel & & \downarrow h & & \parallel \\ \mathbb{G}_{m,\mathbb{R}} & \xrightarrow{w} & G_{\mathbb{R}} & \xrightarrow{t} & \mathbb{G}_{m,\mathbb{R}}. \end{array}$$

A representation $V_{\mathbb{Q}}$ of G , defined over $\mathbb{Q}$, is called homogeneous of weight n if $w(x) * v = x^n v$. Every representation is a direct sum of homogeneous representations. Let $V_{\mathbb{Q}}$ be a representation of weight n . Then h endows $V_{\mathbb{Q}}$ with a rational Hodge structure of weight n .

We shall regard $\mathbb{Q}(n)_{\mathbb{Q}}$ as a representation of G by the rule

$$g * x = t(g)^n x.$$

This convention is reasonable, since $\underline{S}$ acts on $\mathbb{Q}(n)_{\mathbb{R}}$ by the same formula. A polarization of a representation of weight n $V_{\mathbb{Q}}$ of G is a G -invariant form

$$\psi : V_{\mathbb{Q}} \otimes V_{\mathbb{Q}} \longrightarrow \mathbb{Q}(n)$$

such that $\psi(x, h(i)y)$ is a symmetric positive-definite form on $V_{\mathbb{R}}$. A polarization ψ , being G -invariant, is also $\underline{S}$ -invariant, hence is a polarization of the rational Hodge structure $V_{\mathbb{Q}}$.

2.10. Lemma. *For the representation $V_{\mathbb{Q}}$ to be polarizable, it is necessary and sufficient that the representation $V_{\mathbb{R}}$ be $h(i)$ -polarizable.*

Let $P_{\mathbb{Q}}$, respectively $P_{\mathbb{R}}$, be the space of G -invariant $(-1)^n$ -symmetric bilinear forms on $V_{\mathbb{Q}}$, respectively on $V_{\mathbb{R}}$. One has $P_{\mathbb{R}} = \mathbb{R} \otimes P_{\mathbb{Q}}$. The $h(i)$ -polarizations form an open subset of $P_{\mathbb{R}}$, and the polarizations of $V_{\mathbb{Q}}$ form the trace of this open subset on $P_{\mathbb{Q}}$. The lemma follows.

2.11. Proposition. *a) Whether ψ is a polarization of V depends only on the $G(\mathbb{R})$ -conjugacy class of h .*

b) The following conditions are equivalent:

- (i) $\text{ad } h(i)$ is a Cartan involution of $\text{Ker}(t)_{\mathbb{R}}$;
- (ii) every homogeneous representation of G is polarizable;
- (iii) G has a faithful family of polarizable homogeneous representations.

If G is connected, these conditions are equivalent to

- (iii') G has a polarizable representation ρ such that $\text{Ker}(\rho) \cap \text{Ker}(t)$ is finite.

Assertion a) follows from 2.8.a). If G is connected, then $\text{Ker}(t)$ is connected: otherwise t would be of the form t_0^n , with $n > 1$. Since $twx = x^{-2}$, one would have $n = 2$ and $w(-1) \notin \text{Ker}(t)^0$. This is absurd, since

$$w(-1) \in h(\text{Ker}(t : \underline{S} \rightarrow \mathbb{G}_{m,\mathbb{R}})),$$

which is connected. With this said, b) follows from 2.10 and 2.8.b).

3. Reminders on the spinor group

For the reminders contained in this section, one may consult [3].

3.1. Let V be a free module over a ring A , endowed with a quadratic form Q . The module V is identified with a submodule of the Clifford algebra $C(V)$; in $C(V)$ one has

$$v \cdot v = Q(v).$$

Let $C^+(V)$ denote the even part of $C(V)$.

If V is a free module over A , endowed with a symmetric bilinear form ψ , let $C(V)$, or $C(V, \psi)$, denote the Clifford algebra of V endowed with the quadratic form $\psi(x, x)$.

3.2. Suppose that A is a field of characteristic 0, and that Q is nondegenerate. Write $\text{CSpin}(V)$ for the Clifford group. It is the algebraic group of invertible elements g of $C^+(V)$ such that $gVg^{-1} = V$. Define a morphism from $\text{CSpin}(V)$ to $\text{SO}(V)$ by

$$g \mapsto (v \mapsto gv g^{-1}).$$

The kernel of this morphism is reduced to the scalars: one has an exact sequence of algebraic groups

$$0 \longrightarrow \mathbb{G}_m \xrightarrow{w} \text{CSpin}(V) \longrightarrow \text{SO}(V) \longrightarrow 0.$$

The spinor group $\text{Spin}(V)$ is the algebraic subgroup of $\text{CSpin}(V)$ which is the kernel of the spinor norm

$$N : \text{CSpin}(V) \longrightarrow \mathbb{G}_m.$$

Let t denote the inverse of N ; one has a commutative diagram

$$\begin{array}{ccccccc} & & 0 & & & & \\ & & \downarrow & & & & \\ & & \text{Spin}(V) & & & & \\ & & \downarrow & \searrow & & & \\ 0 & \longrightarrow & \mathbb{G}_m & \xrightarrow{w} & \text{CSpin}(V) & \longrightarrow & \text{SO}(V) \longrightarrow 0 \\ & & \searrow \scriptstyle x \mapsto x^{-2} & & \downarrow \scriptstyle t & & \\ & & & & \mathbb{G}_m & & \\ & & & & \downarrow & & \\ & & & & 0 & & \end{array} \quad (3.2.1)$$

By definition, one also has

$$\alpha : \text{CSpin}(V) \longrightarrow C^+(V)^*. \quad (3.2.2)$$

3.3. Let $(C^+(V))_s$ denote the representation of $\text{CSpin}(V)$ on $C^+(V)$ given by

$$g * x = \alpha(g) \cdot x. \quad (3.3.1)$$

Let $(C^+(V))_{\text{ad}}$ denote the representation of $\text{CSpin}(V)$ on $C^+(V)$ deduced from the action of $\text{SO}(V)$ on $C^+(V)$ by transport of structure. One has

$$g *_{\text{ad}} x = \alpha(g) \cdot x \cdot \alpha(g)^{-1}. \quad (3.3.2)$$

The action of $\text{CSpin}(V)$ on $(C^+(V))_s$ is compatible with the right $C^+(V)$ -module structure of $(C^+(V))_s$. One has an isomorphism of representations

$$\text{End}_{C^+(V)}((C^+(V))_s) = (C^+(V))_{\text{ad}}. \quad (3.3.3)$$

One has an isomorphism of representations

$$(C^+(V))_{\text{ad}} = \bigoplus_i \bigwedge^{2i} V. \quad (3.3.4)$$

3.4. Suppose A algebraically closed. There are two cases.

1) $\dim(V)$ odd: $C^+(V)$ is here a matrix algebra. Let W be a simple $C^+(V)$ -module; W is a spin representation of $\text{CSpin}(V)$:

$$g * w = \alpha(g) \cdot w.$$

One has isomorphisms of representations

$$(C^+(V))_s = \text{a sum of copies of } W, \quad (3.4.1)$$

$$(C^+(V))_{\text{ad}} = \text{End}_A(W). \quad (3.4.2)$$

2) $\dim(V)$ even: $C^+(V)$ is here a product of two matrix algebras. Let W_1 and W_2 be two nonisomorphic simple $C^+(V)$ -modules, and put $W = W_1 + W_2$; W_1 and W_2 are the half-spin representations. One has isomorphisms of representations

$$(C^+(V))_s = \text{a sum of copies of } W, \quad (3.4.3)$$

$$(C^+(V))_{\text{ad}} = \text{End}_A(W_1) \times \text{End}_A(W_2). \quad (3.4.4)$$

In both cases, $C^+(V)$ is the bicommutant of the representation W .

3.5. Proposition. *Let Γ be a Zariski-dense subgroup of $\text{Spin}(V)$ and let a be an automorphism of the A -algebra $C^+(V)$ which commutes with the action of Γ given by $\gamma \mapsto \alpha(\gamma) \cdot x \cdot \alpha(\gamma)^{-1}$. Then a is the identity.*

It is enough to treat the case where A is an algebraically closed field and Γ is the whole group Spin ; indeed, commuting with a is a Zariski-closed condition. With the notation of 3.4, identify $C^+(V)$ with a subalgebra of $\text{End}_A(W)$.

There is an automorphism $\bar{a}$ of W such that

$$a(x) = \bar{a}x\bar{a}^{-1} \quad (x \in C^+(V)).$$

For $\gamma \in \text{Spin}(V)$, by hypothesis one has

$$\alpha(\gamma)\bar{a}\alpha(\gamma)^{-1}\bar{a}^{-1} \cdot x \cdot \bar{a}\alpha(\gamma)\bar{a}^{-1}\alpha(\gamma)^{-1} = x \quad (x \in C^+(V));$$

the commutator $(\alpha(\gamma), \bar{a}) = \alpha(\gamma)\bar{a}\alpha(\gamma)^{-1}\bar{a}^{-1}$ is therefore in the center of $C^+(V)$.

1) $\dim(V)$ odd: here $C^+(V) = \text{End}_A(W)$, and $\det_W((\alpha(\gamma), \bar{a})) = 1$; hence $\alpha(\gamma)\bar{a}\alpha(\gamma)^{-1}\bar{a}^{-1}$ is a root of unity ν . Since $\text{Spin}(V)$ is connected, and since for $\gamma = e$ one has $\nu = 1$, one even has

$$(\alpha(\gamma), \bar{a}) = 1, \quad \text{i.e.} \quad \alpha(\gamma)\bar{a} = \bar{a}\alpha(\gamma).$$

2) $\dim(V)$ even: here $C^+(V) = \text{End}_A(W_1) \times \text{End}_A(W_2)$, and $\bar{a}$ either permutes the W_i or preserves them. In both cases, from

$$\det_{W_i}(\alpha(\gamma)) = 1 \quad (i = 1, 2)$$

one deduces that

$$\det_{W_i}((\alpha(\gamma), \bar{a})) = 1 \quad (i = 1, 2),$$

and one concludes as above that

$$\alpha(\gamma)\bar{a} = \bar{a}\alpha(\gamma).$$

In both cases, $\bar{a}$ lies in the commutant of the representation, and therefore commutes with the bicommutant $C^+(V)$: a is the identity.

4. Construction of abelian varieties

In this paragraph and in part of the next, we present results of [7].

4.1. Let $V_{\mathbb{Z}}$ be a free $\mathbb{Z}$ -module of rank $n + 2$ endowed with a bilinear form of nonzero discriminant

$$B : V_{\mathbb{Z}} \otimes V_{\mathbb{Z}} \longrightarrow \mathbb{Z}.$$

Assume that B has index $(n+, 2-)$. We are interested in morphisms of weight 0, $h : \underline{S} \rightarrow \mathrm{SO}(V_{\mathbb{R}})$, such that

- 1) B is a polarization of the representation $V_{\mathbb{Q}}$ of $\mathrm{SO}(V_{\mathbb{Q}})$;
- 2) the Hodge numbers of $V_{\mathbb{C}}$ are

$$h^{-1,1} = h^{1,-1} = 1 \quad \text{and} \quad h^{0,0} = n.$$

4.2. A morphism h as above lifts uniquely to $\tilde{h} : \underline{S} \rightarrow \mathrm{CSpin}(V_{\mathbb{R}})$ making the diagram

$$\begin{array}{ccccc} \mathbb{G}_{m,\mathbb{R}} & \xrightarrow{w} & \underline{S} & \xrightarrow{t} & \mathbb{G}_{m,\mathbb{R}} \\ \parallel & & \downarrow \tilde{h} & & \parallel \\ \mathbb{G}_{m,\mathbb{R}} & \xrightarrow{w} & \mathrm{CSpin}(V_{\mathbb{R}}) & \xrightarrow{t} & \mathbb{G}_{m,\mathbb{R}} \end{array}$$

commutative.

4.3. Lemma. Relative to $\tilde{h}$, every representation of $\mathrm{CSpin}(V_{\mathbb{Q}})$ is polarizable (2.9).

Apply criterion 2.11 b)(iii') to the representation $V_{\mathbb{Q}}$ of $\mathrm{CSpin}(V_{\mathbb{Q}})$.

4.4. Lemma. The Hodge structure on $C^+(V_{\mathbb{Q}})_{\mathrm{ad}}$ defined by $\tilde{h}$ is of type $\{(-1, 1), (0, 0), (1, -1)\}$.

One has

$$C^+(V_{\mathbb{Q}})_{\mathrm{ad}} \simeq \bigwedge^2 V_{\mathbb{Q}},$$

and one concludes by using $h^{-1,1}(V_{\mathbb{Q}}) = 1$.

4.5. Proposition. The Hodge structure on $C^+(V_{\mathbb{Q}})_s$ defined by $\tilde{h}$ is of type $\{(0, 1), (1, 0)\}$. The Hodge structure $C^+(V_{\mathbb{Q}})_s$, endowed with the lattice $C^+(V_{\mathbb{Z}})$, therefore defines an abelian variety.

First suppose that n is odd. After extension of scalars to $\mathbb{C}$, $C^+(V_{\mathbb{Q}})$ becomes a matrix algebra

$$C^+(V_{\mathbb{C}}) \simeq \mathrm{End}(W).$$

On W , $\mathrm{CSpin}(V)_{\mathbb{C}}$ acts by $g \cdot w = \alpha(g)w$: this is the spin representation. The representation $C^+(V_{\mathbb{Q}})_s$ is a sum of $\dim(W)$ copies of W , so it is enough to prove that the weight-one representation W is purely of type $\{(0, 1), (1, 0)\}$. Otherwise, the representation $\mathrm{End}(W) \simeq C^+(V_{\mathbb{C}})_{\mathrm{ad}}$ could not be purely of type $\{(-1, 1), (0, 0), (1, -1)\}$.

For n even, similarly

$$C^+(V_{\mathbb{C}}) \simeq \mathrm{End}(W') \oplus \mathrm{End}(W''),$$

where W' and W'' are the semi-spin representations. As above, one proves that W' and W'' are of type $\{(0, 1), (1, 0)\}$, and hence so is $C^+(V_{\mathbb{Q}})_s$.

The second assertion follows from 2.3 and 3.3.

4.6. The construction given in this section hides the following fact, which we shall not use, but which is read from the Dynkin diagram and is the real reason everything works.


Let $V_{\mathbb{Z}}$ be a polarized Hodge structure of weight 0 and type $\{(1, -1), (0, 0), (-1, 1)\}$, with $h^{1,-1} = 0$. Let

$$r : \mathbb{G}_m \longrightarrow \mathrm{SO}(V_{\mathbb{C}})$$

be the homomorphism such that $r(x)v^{pq} = x^p v^{pq}$ for v^{pq} of type (p, q) . Let T be a maximal torus of $\mathrm{SO}(V_{\mathbb{C}})$ with a system of simple roots Φ , let $\tilde{T}$ be the inverse image of T in $\mathrm{Spin}(V_{\mathbb{C}})$, and let $(w_{\alpha})_{\alpha \in \Phi}$ be the fundamental weights of $\tilde{T}$. The homomorphism r has a unique conjugate

$$r_0 : \mathbb{G}_m \longrightarrow T$$

such that the rational numbers $\langle w_\alpha, r_0 \rangle \in \frac{1}{2}\mathbb{Z}$ are all positive. One checks that r_0 is the homomorphism $H_{\alpha_1} : \mathbb{G}_m \rightarrow T$ relative to the root denoted below by α_1 . The numbers $\langle w_\alpha, r_0 \rangle$ are as follows:



Let w be a dominant weight of a spin or semi-spin representation, and let σ be the opposition involution. The point is that

$$\langle w, r_0 \rangle + \langle \sigma w, r_0 \rangle = 1.$$

5. Construction of families of abelian varieties

We shall use the following theorem of A. Borel, to appear.

5.1. Theorem (A. Borel). Let X/Γ be the quotient of a Hermitian symmetric domain by a torsion-free arithmetic group. It is known [2] that X/Γ is naturally a quasi-projective algebraic variety. Let S be a reduced scheme over $\mathbb{C}$ and let $f : S^{an} \rightarrow X/\Gamma$ be a morphism of analytic spaces. Then f is algebraic.

Let S be smooth over $\mathbb{C}$. For the notion of a variation of Hodge structures $\underline{H}$ on S , and of a polarization of $\underline{H}$, we refer to Griffiths [6]. We shall denote by $H_{\mathbb{Z}}$ the local system of free $\mathbb{Z}$ -modules on S defined by $\underline{H}$, and by H_s the Hodge structure which is the fiber of $\underline{H}$ at $s \in S$.

For X the Siegel half-space, Theorem 5.1 has the following corollary, which generalizes 2.3.

5.2. Scholium. The functor $(a : A \rightarrow S) \mapsto R^1 a_* \mathbb{Z}$ identifies polarized abelian schemes A over S with polarized variations of Hodge structures of type $\{(0, 1), (1, 0)\}$ on S .

5.3. In the category of variations of Hodge structures on S , all usual tensor operations make sense. Thus:

- a) if $\underline{H}$ and $\underline{H}'$ are variations of Hodge structures, there are variations of Hodge structures

$$\underline{H} \otimes \underline{H}', \quad \text{Hom}(\underline{H}, \underline{H}'), \quad \bigwedge^n \underline{H}, \quad \underline{H}^\vee, \quad \text{End}(\underline{H}), \dots;$$

- b) if $\underline{H}$ is a variation of Hodge structures of weight 0, and ψ is a polarization of $\underline{H}$, or any symmetric morphism of Hodge structures $\underline{H} \otimes \underline{H} \rightarrow \mathbb{Z}$, then $C(\underline{H}, \psi)$ and $C^+(\underline{H}, \psi)$ are defined. At every point $t \in S$, with the notation of 3.4, one has

$$C^+(\underline{H}, \psi)_t = C^+(H_t, \psi_t).$$

5.4. Let $(V_{\mathbb{Z}}, B)$ be as in 4.1, and let X be the set of $h : \underline{S} \rightarrow \text{SO}(V_{\mathbb{R}})$ of the type considered in 4.1. Giving such an h is equivalent to giving a two-dimensional subspace $V_{\mathbb{R}}^-$ of $V_{\mathbb{R}}$, on which B is negative definite, together with an orientation of $V_{\mathbb{R}}^-$: for $z = re^{i\theta} \in \mathbb{C}^*$, $h(z)$ induces the identity on the orthogonal complement of $V_{\mathbb{R}}^-$ and the rotation through angle 2θ on $V_{\mathbb{R}}^-$. The group $\text{O}(V_{\mathbb{R}})$ acts on X by transport of structure, $(g * h)(z) = gh(z)g^{-1}$, and the description above makes it geometrically evident that X is a homogeneous space under $\text{SO}(V_{\mathbb{R}})$. The stabilizer of $h \in X$ is a subgroup $\text{SO}(2) \times \text{SO}(n)$, the identity component of a maximal compact subgroup.

5.5. For $h \in X$, let F_h be the corresponding Hodge filtration of $V_{\mathbb{C}}$. The function $h \mapsto F_h^1$ identifies X with an open subset of the space of isotropic lines in $V_{\mathbb{C}}$. This construction endows X with a complex structure for which F_h is a holomorphic function of h . It is known that, for this structure, the two connected components of X are Hermitian symmetric domains.

5.6. Let $W_{\mathbb{Q}}$ be a weight- n representation of the algebraic group $\mathrm{CSpin}(V_{\mathbb{Q}})$: $w(x)y = x^n y$. Let $W_{\mathbb{Q}}$ also denote the constant local system on X with fiber $W_{\mathbb{Q}}$. The group $\mathrm{CSpin}(V_{\mathbb{Q}})$ acts on $(X, W_{\mathbb{Q}})$ by

$$\gamma * (w \text{ at } h \in X) = (\gamma w \text{ at } \gamma h \gamma^{-1} \in X).$$

Each $h \in X$ defines $h : \underline{S} \rightarrow \mathrm{CSpin}(V_{\mathbb{R}})$, hence a $\mathbb{Q}$ -Hodge structure on the fiber of $W_{\mathbb{Q}}$ at $h \in X$. Thus one obtains a variation of $\mathbb{Q}$ -Hodge structures of weight n on X , again denoted $W_{\mathbb{Q}}$:

- a) the Hodge filtration is a holomorphic function of h ;
- b) the transversality axiom is satisfied; we shall use it only in a trivial case.

This construction is compatible with tensor products. In particular, a polarization $\psi : W_{\mathbb{Q}} \otimes W_{\mathbb{Q}} \rightarrow \mathbb{Q}(-n)$ of the representation $W_{\mathbb{Q}}$ defines a polarization of $W_{\mathbb{Q}}$.

Let $W_{\mathbb{Z}}$ be an integral lattice in $W_{\mathbb{Q}}$ and let Γ be a subgroup of $\mathrm{CSpin}(V_{\mathbb{Q}})$, discrete in $\mathrm{CSpin}(V_{\mathbb{R}})$, acting freely on X and satisfying $\Gamma W_{\mathbb{Z}} = W_{\mathbb{Z}}$. Then the constant local system $W_{\mathbb{Z}}$ on X , with fiber $W_{\mathbb{Z}}$, underlies a Γ -equivariant variation of Hodge structure W on X as above. Passing to the quotient, it defines a variation of Hodge structures W on X/Γ .

If $W_{\mathbb{Q}}$ is a representation of $\mathrm{SO}(V_{\mathbb{Q}})$, the same construction applies to discrete subgroups of $\mathrm{SO}(V_{\mathbb{Q}})$.

5.7. Proposition. Let $(\underline{H}, \psi)$ be a polarized variation of Hodge structures of type $\{(-1, 1), (0, 0), (1, -1)\}$, with $h^{1, -1} = 1$, on a smooth connected scheme S . There exist:

- a) a finite surjective étale cover $u : S_1 \rightarrow S$;
- b) an abelian scheme A over S_1 ;
- c) a $\mathbb{Z}$ -algebra C , and $\mu : C \rightarrow \mathrm{End}_{S_1}(A)$;
- d) an isomorphism of variations of Hodge structures

$$u : C^+(\underline{H}, \psi) \xrightarrow{\sim} \mathrm{End}_C(R^1 a_* \mathbb{Z})$$

inducing an isomorphism of local systems of algebras

$$u_{\mathbb{Z}} : C^+(H_{\mathbb{Z}}, \psi) \xrightarrow{\sim} \mathrm{End}_C(R^1 a_* \mathbb{Z}).$$

Let $(V_{\mathbb{Z}}, B)$ be a $\mathbb{Z}$ -module endowed with a symmetric bilinear form, isomorphic to the $(H_{s\mathbb{Z}}, \psi)$, and resume the notation of 5.4. Let n be an integer,

$$\Gamma = \{\gamma \in \mathrm{SO}(V_{\mathbb{Z}}) \mid \gamma \equiv 1 \pmod{n}\}$$

and

$$\tilde{\Gamma} = \{\gamma \in \mathrm{CSpin}(V_{\mathbb{Q}}) \mid \alpha(\gamma) \equiv 1 \pmod{n} \text{ in } C^+(V_{\mathbb{Z}})\}.$$

Assume n large enough that Γ and $\tilde{\Gamma}$ are torsion-free.

There exists a finite étale cover $v : S'_1 \rightarrow S$ such that the local system $v^* H_{\mathbb{Z}/n\mathbb{Z}}$ is trivial. Choose an isomorphism σ_n between this local system and the constant local system $V_{\mathbb{Z}/n\mathbb{Z}}$. Let $s \in S'_1$ and let $\sigma : V_{\mathbb{Z}} \xrightarrow{\sim} H_{s\mathbb{Z}}$ be an isometry such that $\sigma = \sigma_n \pmod{n}$. The Hodge structure h_s of H_s defines $\sigma^{-1}(h_s) \in X$. The image of $\sigma^{-1}(h_s)$ in X/Γ depends only on s . Thus one defines

$$m : S'_1 \longrightarrow X/\Gamma$$

and an isomorphism compatible with polarizations

$$\underline{H} \simeq m^* \underline{V}.$$

Put

$$S_1 = S'_1 \times_{X/\Gamma} X/\tilde{\Gamma} :$$

$$\begin{array}{ccccc}
& & S_1 & & \\
& \swarrow p_1 & & \searrow p_2 & \\
S & \xrightarrow{v} & S'_1 & & X/\tilde{\Gamma} \\
& & \downarrow m & \swarrow & \\
& & X/\Gamma & &
\end{array}$$

On S_1 one has:

- a) an isomorphism $(vp_1)^* \underline{H} \simeq (mp_1)^* \underline{V}$;
- b) a variation of Hodge structure, the inverse image under p_2 of the one on $X/\tilde{\Gamma}$ defined by the representation $(C^+(V_{\mathbb{Q}}))_s$ and the lattice $C^+(V_{\mathbb{Z}})$.

By 5.2, 5.6, and 4.5, this variation of Hodge structures corresponds to an abelian scheme $a : A \rightarrow S_1$.

Let C be the algebra $C^+(V_{\mathbb{Z}})$. The abelian scheme constructed in b) has complex multiplication by C , and the variation of Hodge structures $\text{End}_C(R^1 a_* \mathbb{Z})$ is

$$p_2^*((C^+(V_{\mathbb{Z}}))_{\text{ad}}).$$

Combining this with a), one obtains an isomorphism of local systems of algebras and of variations of Hodge structures

$$C^+((vp_1)^* \underline{H}) \simeq \text{End}_C(R^1 a_* \mathbb{Z}).$$

This proves 5.7.

6. K3 surfaces

6.1. A polarized $K3$ surface over a field K of characteristic 0 is a $K3$ surface over K , say Y , endowed with an element $\eta \in \text{NS}(Y) = \text{Pic}(Y)$ which is the class of an ample invertible sheaf. For $K = \mathbb{C}$, we shall identify $\text{NS}(Y)$ with a subset of $H^2(Y, \mathbb{Z})$.

6.2. A family of $K3$ surfaces parametrized by a characteristic-zero scheme S is a proper flat morphism $f : Y \rightarrow S$ whose fibers are $K3$ surfaces. A polarization of $f : Y \rightarrow S$ is a section η of $\text{Pic}_S(Y)$ which, for every $s \in S$, is a polarization of $Y_s = f^{-1}(s)$.

For every prime number ℓ , identify η with a section of $R^2 f_* \mathbb{Z}_{\ell}$; let $P^2 f_* \mathbb{Z}_{\ell}$ denote the orthogonal complement of η for the cup product; it is a sub- $\mathbb{Z}_{\ell}$ -sheaf of $R^2 f_* \mathbb{Z}_{\ell}$. The cup product

$$R^2 f_* \mathbb{Z}_{\ell} \otimes R^2 f_* \mathbb{Z}_{\ell} \longrightarrow R^4 f_* \mathbb{Z}_{\ell}$$

and the trace morphism

$$R^4 f_* \mathbb{Z}_{\ell} \xrightarrow{\sim} \mathbb{Z}_{\ell}(-2)$$

induce

$$\psi_{\ell} : P^2 f_* \mathbb{Z}_{\ell}(1) \otimes P^2 f_* \mathbb{Z}_{\ell}(1) \longrightarrow \mathbb{Z}_{\ell}. \quad (6.2.1)$$

6.3. Take S to be a scheme of finite type over $\mathbb{C}$. Then $R^2 f_* \mathbb{Z}$ is a variation of Hodge structures on S ; η is everywhere of type $(1, 1)$ and its orthogonal complement $P^2 f_* \mathbb{Z}$ is again a variation of Hodge structures. As above, the cup product defines

$$\psi : P^2 f_* \mathbb{Z}(1) \otimes P^2 f_* \mathbb{Z}(1) \longrightarrow \mathbb{Z}. \quad (6.3.1)$$

This time, the opposite of ψ is a polarization of the variation of Hodge structures $P^2 f_* \mathbb{Z}(1)$. Moreover, ψ_{ℓ} is deduced from ψ via the isomorphism

$$(P^2 f_* \mathbb{Z}(1)_{\mathbb{Z}}) \otimes \mathbb{Z}_{\ell} \simeq P^2 f_* \mathbb{Z}_{\ell}(1).$$

For every $s \in S$, the fiber $(P^2 f_* \mathbb{Z}(1))_s$ is the primitive part of the cohomology $H^2(Y_s, \mathbb{Z})$. The Hodge structure on $(P^2 f_* \mathbb{Z}(1))_s$ is of type $\{(-1, 1), (0, 0), (1, -1)\}$, with $h^{-1, 1} = 1$ ((1.1)(i)). The action of $\pi_1(S, s)$ on $H^2(Y_s, \mathbb{Z})$ induces

$$\pi : \pi_1(S, s) \longrightarrow \text{O}(P^2(Y_s, \mathbb{Z})(1)_{\mathbb{Z}}, \psi). \quad (6.3.2)$$

6.4. Proposition. For every complex polarized $K3$ surface Y_0 , there exists a family of polarized $K3$ surfaces $f : Y \rightarrow S$ as above, with $Y_s \simeq Y_0$ for some $s \in S$, and such that the image of π (6.3.2) is of finite index.

Let $f^\wedge : Y^\wedge \rightarrow T^\wedge$ be the versal formal deformation of Y_0 . Since $H^0(Y_0, T^1) = 0$, $T^\wedge$ is even a formal moduli scheme for Y_0 , but this is unimportant. Since $H^2(Y_0, T^1) = 0$ ((1.1)(i)), $T^\wedge$ is the spectrum of a power-series ring over $\mathbb{C}$. From $f^\wedge$ one obtains a non-polarized variation of Hodge structures on $T^\wedge$. By (1.1.i), it is the universal deformation of the Hodge structure $H^2(Y_0, \mathbb{Z})$ endowed with the cup product.

Let $\tilde{\eta}$ be the image of η under the composite map

$$H^2(Y_0, \mathbb{Z}) \longrightarrow H_{DR}^2(Y^\wedge/T^\wedge) \longrightarrow R^2 f_*^\wedge \mathcal{O}.$$

Let $S^\wedge$ be the formal subscheme of $T^\wedge$ with equation $\tilde{\eta} = 0$, and let $f^\wedge : Y^\wedge \rightarrow S^\wedge$ be induced by $f^\wedge$. Since $h^{0,2} = 1$, $S^\wedge \subset T^\wedge$ is defined by one equation; this equation belongs to a regular system of parameters, so $S^\wedge$ is a power-series ring. On $S^\wedge$, η comes from an ample invertible sheaf $\mathcal{O}(1)$ on $Y^\wedge$; hence $f^\wedge$ is algebraizable. Moreover, on $S^\wedge$, the polarized variation of Hodge structure

$$(P^2(Y^\wedge/S^\wedge, \mathbb{Z})(1), \psi)$$

is a universal deformation of $(P^2(Y_0, \mathbb{Z})(1), \psi)$.

Put $S^\wedge = \text{Spec}(A^\wedge)$, and express $A^\wedge$ as an inductive limit of finite-type algebras over $\mathbb{C}$. Since $Y^\wedge$ is algebraizable, there exists a Cartesian diagram

$$\begin{array}{ccc} Y^\wedge & \longrightarrow & Y \\ f^\wedge \downarrow & & \downarrow f \\ S^\wedge & \longrightarrow & S \end{array}$$

with S of finite type over $\mathbb{C}$ and Y a polarized family of $K3$ surfaces over S ; one does not suppose that $S^\wedge$ is a completion of S . One may take S normal and connected; let $s \in S$ be the image of the closed point of $S^\wedge$.

In fact, using Artin's approximation theorem [1], one can find S such that $S^\wedge$ is a completion of S . Then S is smooth in a neighborhood of s . The simplifications resulting from such a choice are mostly psychological.

We show that $f : Y \rightarrow S$ satisfies 6.2. After replacing S by a finite étale cover, one defines, as in 5.7, a holomorphic map

$$m : S \longrightarrow X/\Gamma$$

from S to a moduli space of polarized Hodge structures such that the inverse image of the universal family of Hodge structures on X/Γ is $(P^2(Y/S, \mathbb{Z})(1), \psi)$.

By Borel (5.1), m is a morphism of schemes. The morphism m is dominant, because the composite $m^\wedge : S^\wedge \rightarrow X/\Gamma$ is dominant. It follows, for example from P. Deligne, Théorie de Hodge II, Publ. Math. IHES 40, Lemma 4.4.17, that $m(\pi_1(S, s))$ is of finite index in

$$\pi_1(X/\Gamma, m(s)) = \Gamma,$$

itself of finite index in the orthogonal group.

6.5. Proposition. Let Y_0 be a polarized $K3$ surface over a field K of characteristic 0. There exist:

- a) a scheme S of finite type over $\mathbb{Q}$, and a family $f : Y \rightarrow S$ of polarized $K3$ surfaces, parametrized by S ;
- b) a finite extension K' of K and $v : \text{Spec}(K') \rightarrow S$ such that v^*Y is isomorphic to $Y_0 \otimes_K K'$;
- c) an abelian scheme $a : A \rightarrow S$ over S ;
- d) a $\mathbb{Z}$ -algebra C and $\mu : C \rightarrow \text{End}_S(A)$;
- e) an isomorphism of sheaves of $\mathbb{Z}_\ell$ -algebras on S

$$u : C^+(P^2 f_* \mathbb{Z}_\ell(1), \psi_\ell) \xrightarrow{\sim} \text{End}_C(R^1 a_* \mathbb{Z}_\ell).$$

One is reduced to the case where K is of finite type over $\mathbb{Q}$. Choose a complex embedding $\sigma : K \rightarrow \mathbb{C}$ and apply 6.4 to $Y_0 \otimes_{K, \sigma} \mathbb{C}$ to obtain a family of polarized $K3$ surfaces $f_1 : Y_1 \rightarrow S_1$. Apply 5.7 to f_1 ; one obtains

a family $f_2 : Y_2 \rightarrow S_2$ still satisfying the conclusion of 6.4, and, over S_2 , an abelian scheme $a_2 : A_2 \rightarrow S_2$ with complex multiplication $\mu_2 : C \rightarrow \text{End}_{S_2}(A_2)$ and an isomorphism

$$u_2 : C^+(P^2 f_{2*} \mathbb{Z}(1), \psi) \xrightarrow{\sim} \text{End}_C(R^1 a_{2*} \mathbb{Z})$$

of local systems of algebras. The system of complex schemes and morphisms

$$\Xi_2 = (S_2, f_2, Y_2, a_2, A_2, \mu_2)$$

is defined by finitely many equations. Hence there exists a finite-type extension K_3 of K , with $K \subset K_3 \subset \mathbb{C}$, and, over K_3 , a system

$$\Xi_3 = (S_3; f_3, Y_3, a_3, A_3, \mu_3)$$

which, after extension of scalars from K_3 to $\mathbb{C}$, gives Ξ_2 .

6.5.1. Lemma. There exists a unique isomorphism of sheaves of $\mathbb{Z}_\ell$ -algebras

$$u_3 : C^+(P^2 f_{3*} \mathbb{Z}_\ell(1), \psi_\ell) \xrightarrow{\sim} \text{End}_C(R^1 a_{3*} \mathbb{Z}_\ell).$$

It is enough to verify that such an isomorphism exists and is unique after extension of scalars to an algebraic closure of K_3 , for instance $\mathbb{C}$. Existence holds by construction, and uniqueness follows from 3.5. Indeed, for $s \in S_2$, the image of $\pi_1(S_2, s)$ in $O(P^2(Y_{2s}, \mathbb{Z}), \psi)$ has finite index and hence contains a Zariski-dense subgroup of the group SO .

Let T be an integral scheme of finite type over $\mathbb{Q}$, with function field K_3 . For suitable T , S_3 is the generic fiber of a system

$$\Xi = (S, f, Y, a, A, \mu)$$

over T , and u_3 extends to

$$u : C^+(P^2 f_* \mathbb{Z}_\ell(1), \psi_\ell) \xrightarrow{\sim} \text{End}_C(R^1 a_* \mathbb{Z}_\ell).$$

One easily verifies the existence of K' as in b), and this completes the proof.

6.6. We prove Theorem 1.3. Let $f_V : Y_V \rightarrow \text{Spec}(V)$ be as in 1.2(i). Let K be the fraction field of V , and apply 6.5 to the surface

$$Y_K = Y \otimes_V K$$

over K , endowed with any polarization. After replacing V by its normalization in a finite extension of K , one may suppose that there exists

$$\Xi = (S, f, Y, a, A, \mu, u)$$

as in 6.5, such that Y_K is the inverse image of Y under $v : \text{Spec}(K) \rightarrow S$. By inverse image, one deduces an abelian variety with complex multiplication (A_K, μ) over K .

Let $\bar{K}$ be an algebraic closure of K , and let $\bar{k}$ be the corresponding algebraic closure of k , the residue field of the normalization of V in $\bar{K}$. From u , one obtains an isomorphism of $\text{Gal}(\bar{K}/K)$ -modules

$$C^+(P^2(Y_{\bar{K}}, \mathbb{Z}_\ell)(1), \psi_\ell) \xrightarrow{\sim} \text{End}_C(H^1(A_{\bar{K}}, \mathbb{Z}_\ell)). \quad (6.6.1)$$

Since Y_V is smooth over V , the inertia group I acts trivially on

$$H^2(Y_{\bar{K}}, \mathbb{Z}_\ell) \simeq H^2(Y_{\bar{k}}, \mathbb{Z}_\ell).$$

The polarization defines an invariant η in $H^2(Y_{\bar{k}}, \mathbb{Z}_\ell)(1)$, with orthogonal complement $P^2(Y_{\bar{k}}, \mathbb{Z}_\ell)(1)$, and I acts trivially on

$$C^+(P^2(Y_{\bar{K}}, \mathbb{Z}_\ell)(1), \psi_\ell) \simeq C^+(P^2(Y_{\bar{k}}, \mathbb{Z}_\ell)(1), \psi_\ell).$$

Since I acts trivially on $\text{End}_C(H^1(A_{\bar{K}}, \mathbb{Z}_\ell))$, and commutes with the complex multiplication, it acts through the center of $C \otimes \mathbb{Q}_\ell$. On the other hand, it is known that a subgroup of finite index in I acts on $H^1(A_{\bar{K}}, \mathbb{Z}_\ell)$ unipotently; it follows that I acts through one of its finite quotients. After again replacing K by a finite extension, one may therefore suppose that A_K has good reduction over V .

For $A_{\bar{k}}$ the special fiber of its reduction, (6.6.1) reduces to an isomorphism of $\text{Gal}(\bar{k}/k)$ -modules

$$C^+(P^2(Y_{\bar{k}}, \mathbb{Z}_\ell)(1), \psi_\ell) \xrightarrow{\sim} \text{End}_C(H^1(A_{\bar{k}}, \mathbb{Z}_\ell)). \quad (6.6.2)$$

By Weil's theory for A_k , the eigenvalues of Frobenius acting on

$$C^+(P^2(Y_{\bar{k}}, \mathbb{Z}_\ell)(1), \psi_\ell)$$

are algebraic numbers all of whose complex conjugates have absolute value 1. One has

$$\bigwedge^2 (P^2(Y_{\bar{k}}, \mathbb{Z}_\ell)(1)) \subset C^+(P^2(Y_{\bar{k}}, \mathbb{Z}_\ell)(1), \psi_\ell),$$

and, since $\dim P^2(Y_{\bar{k}}, \mathbb{Z}_\ell) > 2$ in the present case, it follows that the eigenvalues of Frobenius acting on $P^2(Y_{\bar{k}}, \mathbb{Z}_\ell)(1)$ also have complex absolute value 1. The same assertion holds for $H^2(Y_{\bar{k}}, \mathbb{Z}_\ell)(1)$, and the theorem follows.

7. Remarks

7.1. Let H be a rational Hodge structure. The Mumford–Tate group of H is the smallest algebraic subgroup G of $\text{GL}(H_{\mathbb{Q}})$, defined over $\mathbb{Q}$, such that $G(\mathbb{R})$ contains the image of $h : \underline{S} \rightarrow \text{GL}(H_{\mathbb{R}})$. If H is polarizable, G is reductive. This definition differs from Mumford's: if H has weight $n \neq 0$, our G contains the homotheties. By 2.9, every homogeneous representation of G is endowed with a natural Hodge structure.

One of the key points in the proof of 1.3 was that the Hodge structure $H^2(X, \mathbb{Q})$, for X a $K3$, can be expressed in terms of abelian varieties, in the following sense.

7.2. Definition. A rational Hodge structure H is expressible in terms of abelian varieties if it belongs to the smallest category of rational Hodge structures which is stable under direct factors, direct sums, and tensor products, and which contains the $H^1(A, \mathbb{Q})$ for A an abelian variety, as well as the $\mathbb{Q}(n)$.

In this subsection, we show that this property of $K3$ surfaces is exceptional.

Consider the following property of a rational Hodge structure, with Mumford–Tate group G :

- (*) the Hodge structure of the adjoint representation $\text{Lie}(G)$ of G is purely of type $\{(-1, 1), (0, 0), (1, -1)\}$.

7.3. Proposition. If a rational Hodge structure H is expressible in terms of abelian varieties, then condition (*) is satisfied.

Indeed:

- (a) the Hodge structures $H^1(A, \mathbb{Q})$ and $\mathbb{Q}(n)$ satisfy (*);
- (b) condition (*) is stable under direct factors, direct sums, and tensor products.

7.4. Analogous arguments show that if H is expressible in terms of abelian varieties, then the reductive group $G_{\mathbb{C}}$ has no factors of exceptional type.

7.5. Proposition. Let $\underline{H}$ be a polarized variation of Hodge structures on an irreducible scheme S . For $s \in S$ outside a meagre subset of S , there exists a finite-index subgroup of $\pi_1(S, s)$ whose image in $\text{GL}(H_{s\mathbb{Z}})$ is contained in the Mumford–Tate group of H_s .

At every point of S , the Mumford–Tate group G_s of H_s may be described as follows. It is the algebraic subgroup of $\text{GL}(H_{s\mathbb{Q}})$ which fixes all tensors $t \in H_{s\mathbb{Q}}^{\otimes n} \otimes H_{s\mathbb{Q}}^{*\otimes n}$, $n > 0$, that are rational of type $(0, 0)$. From this description one easily deduces that, outside a meagre subset M of S , G_s is locally constant: for $s \notin M$, the largest subspace of type $(0, 0)$

$$V(n)_s \subset H_{s\mathbb{Q}}^{\otimes n} \otimes H_{s\mathbb{Q}}^{*\otimes n}$$

is locally constant and defines a local system $V(n)$ on S . This local system underlies a polarized variation of Hodge structures of type $(0, 0)$; hence on $V(n)_s$ there exists a positive-definite quadratic form invariant under $\pi_1(S, s)$, and $\pi_1(S, s)$ acts on $V(n)_s$ through a finite group. One concludes by observing that there exists a finite sequence of integers n_i such that G_s is the algebraic subgroup of $\text{GL}(H_{s\mathbb{Q}})$ acting trivially on the $V(n_i)_s$.

7.6. Consider a Lefschetz pencil of hypersurfaces of degree d and odd dimension $2r - 1$:

$$\begin{array}{ccc} X & \longrightarrow & \mathbb{P}^{2r}(\mathbb{C}) \times \mathbb{P}^1(\mathbb{C}) \\ f \downarrow & \swarrow \text{pr}_2 & \\ \mathbb{P}^1(\mathbb{C}) & & \end{array}$$

Let $U = \mathbb{P}^1(\mathbb{C})$ be the complement of the finite set of $t \in \mathbb{P}^1(\mathbb{C})$ such that the hypersurface $X_t = f^{-1}(t)$ is singular, and let $s \in U$. The cup product is an alternating form ψ on $H^{2r-1}(X_s, \mathbb{Z})$, respected by the monodromy, whence

$$m : \pi_1(U, s) \longrightarrow \text{Sp}(H^{2r-1}(X_s, \mathbb{Z}), \psi).$$

By a theorem of Kazhdan and Margulis, the image of m is Zariski-dense in the symplectic group. By 7.5, for s outside a meagre, in fact countable, subset of $\mathbb{P}^1(\mathbb{C})$, the Mumford–Tate group of the rational Hodge structure $H^{2r-1}(X_s, \mathbb{Q})$ is therefore the whole group of symplectic similitudes. If this Hodge structure is not of Hodge level at most one, i.e. if some h^{pq} with $|q - p| \neq 1$ is nonzero, it follows from 7.3 that it is not expressible in terms of abelian varieties.

Bibliography

1. Artin, M.: Algebraisation of formal moduli I. Global analysis. Papers in honor of K. Kodaira. Princeton University Press.
2. Baily, W. L., Borel, A.: Compactification of arithmetic quotients of bounded symmetric domains. *Annals of Math.* (2) **84**, 442–528 (1966). (MR 35, 6870).
3. Bourbaki, N.: *Algèbre – Chapitre 9*. Paris: Hermann 1959. Act. Sci. et Ind. 1272.
4. Deligne, P.: *Travaux de Griffiths – Séminaire Bourbaki 376 – mai 1970*. Lecture Notes in mathematics 180. Berlin-Heidelberg-New York: Springer 1971.
5. Demazure, M.: *Motifs des variétés algébriques – Séminaire Bourbaki 365 – mai 1970*. Lecture Notes in mathematics 180. Berlin-Heidelberg-New York: Springer 1971.
6. Griffiths, P. A.: Periods of integrals on algebraic manifolds. Summary of main results and discussions of open problems and conjectures. *Bull. Am. Math. Soc.* **75**, 228–296 (1970).
7. Kuga, M., Satake, I.: Abelian varieties attached to polarized K3-surfaces. *Math. Ann.* **169**, 239–242 (1967). (MR 35 1602).
8. Shafarevich, I. R., Averbuch, B. C., Vainberg, Yu. R., Yuzhchenko, A. B., Manin, Yu. I., Moishezon, B. C., Tyurina, C. I., Tyurin, A. I.: *Algebraic surfaces*. Trudi Mat. Inst. Steklov LXXV.
9. Weil, A.: *Variétés abéliennes et courbes algébriques*. Paris: Hermann 1948.

Pierre Deligne
Institut des Hautes Études Scientifiques
35, Route de Chartres
F-91 Bures-sur-Yvette
France

(Received 6 August 1971 / 1 October 1971)