

# Seminar on Algebraic Geometry 3

## Group Schemes

Current-Progress Cumulative English Translation and T<sub>E</sub>X Reconstruction  
Exposés I–XI except XII, plus XIII

SGA 3 English reconstruction project

28 July 2026

# Reconstruction and attribution status

This is a cumulative current-progress English reconstruction containing Exposés I–XI except XII, plus XIII. Exposé XII is represented by an explicit gap page; Exposés XIV–XXVI are outside the currently available translation set. The published Polo–Gille PDFs control the French text, mathematics, numbering, notes, and diagram appearance. The pre-existing user-supplied OCR was used only as a read-only locator and drafting witness. Exposés IV–IX, XI, and XIII use native  $\text{\LaTeX}$  diagrams. Source-page image witnesses still present in Exposés I–III and the four raster diagram placeholders in the newly integrated Exposé X remain explicitly provisional until their own native-diagram pass is complete.

Jacob Reinhold’s CC BY 4.0 English Markdown is credited and was used as a comparison and drafting lineage. It is not the source authority; the Polo–Gille PDFs control every admitted passage, formula, and diagram. The CC BY 4.0 statement belongs to that comparison lineage and does not by itself license the underlying French edition or this reconstruction.

No editable Polo–Gille source was publicly available when this reconstruction began. Formulas and document structure were therefore reconstructed in  $\text{\LaTeX}$ . This checkpoint does not claim completion of SGA 3 as a whole. It binds the latest archive-admitted English scopes into one reading surface while preserving the omissions visibly.

# Contents

|                                                                                                   |            |
|---------------------------------------------------------------------------------------------------|------------|
| <b>Reconstruction status</b>                                                                      | <b>i</b>   |
| <b>Editorial Notice</b>                                                                           | <b>1</b>   |
| <b>Introduction</b>                                                                               | <b>3</b>   |
| <b>Exposé I. Algebraic Structures. Group Cohomology</b>                                           | <b>7</b>   |
| 1. Generalities . . . . .                                                                         | 7          |
| 2. Algebraic structures . . . . .                                                                 | 18         |
| 3. The category of $\mathcal{O}$ -modules, the category of $G$ - $\mathcal{O}$ -modules . . . . . | 24         |
| 4. Algebraic structures in the category of schemes . . . . .                                      | 27         |
| 5. Group cohomology . . . . .                                                                     | 39         |
| 6. $G$ -equivariant objects and modules . . . . .                                                 | 47         |
| Bibliography . . . . .                                                                            | 57         |
| <b>Exposé II. Tangent Bundles – Lie Algebras</b>                                                  | <b>59</b>  |
| 1. The functors $\mathrm{Hom} \, Z/S \, (X, Y)$ . . . . .                                         | 59         |
| 2. The schemes $I \, S \, (M)$ . . . . .                                                          | 62         |
| 3. The tangent bundle, condition (E) . . . . .                                                    | 65         |
| 4. Tangent Space to a Group — Lie Algebras . . . . .                                              | 81         |
| 5. Computation of some Lie algebras . . . . .                                                     | 105        |
| 6. Miscellaneous remarks . . . . .                                                                | 112        |
| Bibliography . . . . .                                                                            | 115        |
| <b>Exposé III. Infinitesimal Extensions</b>                                                       | <b>116</b> |
| 0. Reminders from SGA 1, Exposé III . . . . .                                                     | 117        |
| 1. Extensions and Cohomology . . . . .                                                            | 142        |
| 2. Infinitesimal extensions of a morphism of group schemes . . . . .                              | 151        |
| 3. Infinitesimal extensions of a group scheme . . . . .                                           | 157        |
| 4. Infinitesimal extensions of closed subgroups . . . . .                                         | 161        |
| Bibliography . . . . .                                                                            | 199        |
| <b>Exposé IV. Topologies and Sheaves</b>                                                          | <b>200</b> |
| 1. Universal Effective Epimorphisms . . . . .                                                     | 200        |
| 2. Descent Morphisms . . . . .                                                                    | 204        |
| 3. Universal Effective Equivalence Relations . . . . .                                            | 207        |
| 3.1. Equivalence Relations: Definitions . . . . .                                                 | 207        |
| 3.2. Equivalence Relation Defined by a Group Acting Freely . . . . .                              | 210        |
| 3.3. Universally Effective Equivalence Relations . . . . .                                        | 211        |
| 3.4. (M)-Effectivity . . . . .                                                                    | 213        |

|                                                                                        |            |
|----------------------------------------------------------------------------------------|------------|
| 3.5. Construction of Quotients by Descent . . . . .                                    | 215        |
| 4. Topologies and Sheaves . . . . .                                                    | 216        |
| 4.1. Sieves . . . . .                                                                  | 216        |
| 4.2. Topologies: Definitions . . . . .                                                 | 219        |
| 4.3. Presheaves, Sheaves, and the Sheaf Associated to a Presheaf . . . . .             | 221        |
| 4.4. Exactness Properties of the Category of Sheaves . . . . .                         | 228        |
| 4.5. A Topology Coarser than the Canonical Topology . . . . .                          | 236        |
| 4.6. Quotient of a Sheaf by an Equivalence Relation . . . . .                          | 240        |
| 4.7. Use of Effectivity Criteria: Isomorphism Theorem . . . . .                        | 245        |
| 5. Passage to the Quotient and Algebraic Structures . . . . .                          | 246        |
| 5.1. Principal Homogeneous Bundles . . . . .                                           | 246        |
| 5.2. Group Structures and Passage to the Quotient . . . . .                            | 249        |
| 5.3. Use of Effectivity Criteria: Noether's Theorem . . . . .                          | 251        |
| 6. Topologies in the Category of Schemes . . . . .                                     | 252        |
| 6.1. The Zariski Topology . . . . .                                                    | 252        |
| 6.2. A Procedure for Constructing Topologies . . . . .                                 | 252        |
| 6.3. Application to the Category of Schemes . . . . .                                  | 256        |
| 6.4. Effectivity Conditions . . . . .                                                  | 258        |
| 6.5. Principal Homogeneous Bundles . . . . .                                           | 259        |
| 6.6. Other Topologies . . . . .                                                        | 259        |
| 6.7. Homogeneous Spaces . . . . .                                                      | 259        |
| Bibliography . . . . .                                                                 | 262        |
| <b>Exposé V. Construction of Quotient Schemes</b>                                      | <b>263</b> |
| 1. $\mathcal{C}$ -groupoids . . . . .                                                  | 264        |
| 2. Examples of $\mathcal{C}$ -groupoids . . . . .                                      | 267        |
| 3. Some sorites concerning $\mathcal{C}$ -groupoids . . . . .                          | 269        |
| 4. Passage to the quotient by a finite flat groupoid . . . . .                         | 274        |
| 5. Passage to the quotient by a finite flat groupoid (the general case) . . . . .      | 282        |
| 6. Passage to the quotient when a quasi-section exists . . . . .                       | 285        |
| 7. Quotient by a proper and flat groupoid . . . . .                                    | 289        |
| 8. Passage to the quotient by a flat groupoid that is not necessarily proper . . . . . | 296        |
| 9. Elimination of the Noetherian hypotheses in Theorem 7.1 . . . . .                   | 298        |
| 10. Supplement: quotients by a group scheme . . . . .                                  | 302        |
| 10.2. Stabilizer of the diagonal . . . . .                                             | 305        |
| 10.3. The case in which $F$ is quasi-finite over $X$ . . . . .                         | 306        |
| 10.4. The case in which $F$ is flat over $X$ . . . . .                                 | 307        |
| Bibliography . . . . .                                                                 | 310        |
| <b>Exposé VI-A. Generalities on Algebraic Groups</b>                                   | <b>311</b> |
| 0. Preliminary Remarks . . . . .                                                       | 311        |
| 1. Local properties of an $A$ -group locally of finite type . . . . .                  | 314        |
| 2. Connected components of an $A$ -group locally of finite type . . . . .              | 318        |
| 3. Construction of quotients $F \backslash G$ (for $G, F$ of finite type) . . . . .    | 332        |
| 4. Construction of quotients $F \backslash G$ (general case) . . . . .                 | 339        |
| 5. Connections with Exposé IV and consequences . . . . .                               | 343        |
| 6. Complements on $k$ -groups not necessarily of finite type . . . . .                 | 348        |
| Bibliography . . . . .                                                                 | 351        |

|                                                                                                      |            |
|------------------------------------------------------------------------------------------------------|------------|
| <b>Exposé VI-B. Generalities on Group Schemes</b>                                                    | <b>353</b> |
| 1. Morphisms of groups locally of finite type over a field . . . . .                                 | 353        |
| 2. “Open properties” of groups and group morphisms locally of finite presentation . . . . .          | 361        |
| 3. Identity component of a group locally of finite presentation . . . . .                            | 371        |
| 4. Dimension of the fibers of groups locally of finite presentation . . . . .                        | 378        |
| 5. Separation of groups and homogeneous spaces . . . . .                                             | 382        |
| 5.8. Complements . . . . .                                                                           | 395        |
| 6. Subfunctors and group subschemes . . . . .                                                        | 399        |
| 7. Generated subgroups; commutator group . . . . .                                                   | 413        |
| 8. Solvable or nilpotent group schemes . . . . .                                                     | 429        |
| 9. Quotient sheaves . . . . .                                                                        | 435        |
| 10. Passage to the projective limit in group schemes and schemes with an operator<br>group . . . . . | 438        |
| 11. Affine group schemes . . . . .                                                                   | 447        |
| 12. Complements on $\text{Gaf}$ and the “anti-affine” groups . . . . .                               | 477        |
| 13. Flat affine groups over a regular base of dimension at most 2 . . . . .                          | 483        |
| Bibliography . . . . .                                                                               | 490        |
| <b>Working-state notice for Exposé VII</b>                                                           | <b>493</b> |
| <b>Exposé VII-A. Infinitesimal Study of Group Schemes</b>                                            | <b>494</b> |
| A) Differential Operators and Restricted $p$ -Lie Algebras . . . . .                                 | 494        |
| 1. Differential Operators . . . . .                                                                  | 494        |
| 2. Invariant Differential Operators on Group Schemes . . . . .                                       | 502        |
| 3. Coalgebras and Cartier Duality . . . . .                                                          | 507        |
| 4. “Frobeniusseries” . . . . .                                                                       | 514        |
| 5. Restricted $p$ -Lie Algebras . . . . .                                                            | 525        |
| 6. The $p$ -Lie Algebra of an $S$ -Group Scheme . . . . .                                            | 537        |
| 7. Radicial Groups of Height 1 . . . . .                                                             | 542        |
| 8. The Case of a Base Field . . . . .                                                                | 551        |
| <b>Bibliography</b>                                                                                  | <b>560</b> |
| <b>Exposé VII-B. Infinitesimal Study of Group Schemes</b>                                            | <b>561</b> |
| B) Formal Groups . . . . .                                                                           | 561        |
| 0. Review of Pseudocompact Rings and Modules . . . . .                                               | 561        |
| 0.4. Profinite Algebras . . . . .                                                                    | 576        |
| 0.5. Base Change . . . . .                                                                           | 578        |
| 1. Formal Varieties over a Pseudocompact Ring . . . . .                                              | 580        |
| 1.3. Monomorphisms and topological flatness . . . . .                                                | 589        |
| 1.4. Descent and exactness . . . . .                                                                 | 600        |
| 1.5. The flat topology . . . . .                                                                     | 603        |
| 1.6. Étale formal varieties . . . . .                                                                | 604        |
| 2. Generalities on Formal Groups . . . . .                                                           | 612        |
| 2.1. Formal groups and cogroups . . . . .                                                            | 612        |
| 2.2. Bialgebras over $\mathbf{O}_k$ . . . . .                                                        | 614        |
| 2.3. The covariant algebra . . . . .                                                                 | 617        |
| 2.3.2 . . . . .                                                                                      | 620        |
| 2.4. Quotients of formal groups . . . . .                                                            | 623        |
| 2.5. Étale formal groups . . . . .                                                                   | 629        |

|                                                                                                                                                  |            |
|--------------------------------------------------------------------------------------------------------------------------------------------------|------------|
| 2.6. Lie algebras of formal groups . . . . .                                                                                                     | 634        |
| 2.7. Infinitesimal formal monoids . . . . .                                                                                                      | 640        |
| 2.8. Unipotent group schemes over a field . . . . .                                                                                              | 642        |
| 2.9. Pointed cocommutative Hopf algebras over a field . . . . .                                                                                  | 643        |
| 3. Phenomena specific to characteristic 0 . . . . .                                                                                              | 646        |
| 4. Phenomena specific to characteristic $p > 0$ . . . . .                                                                                        | 656        |
| 5. Homogeneous spaces of infinitesimal formal groups over a field . . . . .                                                                      | 667        |
| 5.3. Remarks . . . . .                                                                                                                           | 692        |
| 5.4. Corollary . . . . .                                                                                                                         | 693        |
| 5.5. Complements . . . . .                                                                                                                       | 693        |
| <b>Bibliography</b>                                                                                                                              | <b>696</b> |
| <b>Exposé VIII. Diagonalizable Groups</b>                                                                                                        | <b>698</b> |
| 1. Biduality . . . . .                                                                                                                           | 698        |
| 2. Scheme-theoretic Properties of Diagonalizable Groups . . . . .                                                                                | 704        |
| 3. Exactness Properties of the Functor $D_S$ . . . . .                                                                                           | 705        |
| 4. Torsors under a Diagonalizable Group . . . . .                                                                                                | 707        |
| 5. Quotient of an Affine Scheme by a Diagonalizable Group Acting Freely . . . . .                                                                | 711        |
| 6. Essentially Free Morphisms and Representability of Certain Functors . . . . .                                                                 | 714        |
| 7. Appendix: On Monomorphisms of Group Preschemes . . . . .                                                                                      | 720        |
| Bibliography . . . . .                                                                                                                           | 726        |
| <b>Exposé IX. Groups of Multiplicative Type: Homomorphisms into a Group Scheme</b>                                                               | <b>727</b> |
| 1. Definitions . . . . .                                                                                                                         | 727        |
| 2. Extension of Certain Properties of Diagonalizable Groups to Groups of Multiplicative Type . . . . .                                           | 729        |
| 3. Infinitesimal Properties: Lifting and Conjugation Theorems . . . . .                                                                          | 733        |
| 4. The Density Theorem . . . . .                                                                                                                 | 735        |
| 5. Central Homomorphisms of Groups of Multiplicative Type . . . . .                                                                              | 744        |
| 6. Monomorphisms of Groups of Multiplicative Type, and Canonical Factorization . . . . .                                                         | 749        |
| 7. Algebraicity of Formal Homomorphisms into an Affine Group . . . . .                                                                           | 754        |
| 8. Subgroups, Quotient Groups, and Extensions of Groups of Multiplicative Type over a Field . . . . .                                            | 758        |
| Bibliography . . . . .                                                                                                                           | 760        |
| <b>Working-state notice for Exposé X</b>                                                                                                         | <b>761</b> |
| <b>Exposé X. Characterization and Classification of Groups of Multiplicative Type</b>                                                            | <b>762</b> |
| 1. Classification of Isotrivial Groups. Case of a Base Field . . . . .                                                                           | 762        |
| 2. Infinitesimal Variations of Structure . . . . .                                                                                               | 764        |
| 3. Finite Variations of Structure: Complete Base Ring . . . . .                                                                                  | 768        |
| 4. Case of an arbitrary base. Quasi-isotriviality theorem . . . . .                                                                              | 772        |
| 5. Scheme of homomorphisms from one group of multiplicative type to another. Twisted constant groups and groups of multiplicative type . . . . . | 778        |
| 6. Infinite Galois principal bundles and the enlarged fundamental group . . . . .                                                                | 784        |
| 7. Classification in terms of the enlarged fundamental group . . . . .                                                                           | 788        |
| 8. Appendix: Elimination of certain affineness hypotheses . . . . .                                                                              | 790        |
| 9. Addenda . . . . .                                                                                                                             | 797        |

|                                                                               |                                                                                                                               |            |
|-------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------|------------|
| 0.9.1                                                                         | Simplicial sets, topoi, groupoids, and topological spaces . . . . .                                                           | 797        |
| 0.9.2                                                                         | Locally constant sheaves; descent data . . . . .                                                                              | 797        |
| 0.9.3                                                                         | Some equivalences of categories . . . . .                                                                                     | 798        |
| 0.9.4                                                                         | Fundamental groups and groupoids . . . . .                                                                                    | 798        |
| 0.9.5                                                                         | Cones . . . . .                                                                                                               | 799        |
| 0.9.6                                                                         | Representability of the functor $\bar{\pi}^1(S'/S, \xi; -)$ . . . . .                                                         | 801        |
| 0.9.7                                                                         | Contractibility of $k$ ; counterexample to the injectivity of $k_{\bullet} \rightarrow K_{\bullet}$ . . .                     | 801        |
|                                                                               | Bibliography . . . . .                                                                                                        | 802        |
| <b>Exposé XI. Representability Criteria and Multiplicative-Type Subgroups</b> |                                                                                                                               | <b>803</b> |
| 0.                                                                            | Introduction . . . . .                                                                                                        | 803        |
| 1.                                                                            | Review of Smooth, Étale, and Unramified Morphisms . . . . .                                                                   | 804        |
| 2.                                                                            | Examples of Formally Smooth Functors from the Theory of Groups of Multi-<br>plicative Type . . . . .                          | 810        |
| 3.                                                                            | Auxiliary Representability Results . . . . .                                                                                  | 813        |
| 4.                                                                            | The Scheme of Multiplicative-Type Subgroups of a Smooth Affine Group . . .                                                    | 824        |
| 5.                                                                            | First Corollaries of the Representability Theorem . . . . .                                                                   | 829        |
| 6.                                                                            | A Rigidity Property for Homomorphisms of Certain Group Schemes, and the<br>Representability of Certain Transporters . . . . . | 833        |
| <b>Exposé XII (not included)</b>                                              |                                                                                                                               | <b>839</b> |
| <b>Exposé XIII. Regular Elements of Algebraic Groups and Lie Algebras</b>     |                                                                                                                               | <b>840</b> |
| 0.10                                                                          | An auxiliary lemma on varieties with operators . . . . .                                                                      | 840        |
| 0.11                                                                          | Density theorem and theory of the regular points of $G$ . . . . .                                                             | 843        |
| 0.12                                                                          | The case of an arbitrary base prescheme . . . . .                                                                             | 853        |
| 0.13                                                                          | Lie algebras over a field: rank, regular elements, Cartan subalgebras . . . .                                                 | 857        |
| 0.14                                                                          | The case of the Lie algebra of a smooth algebraic group: the density theorem                                                  | 862        |
| 0.15                                                                          | Cartan subalgebras and subgroups of type $(C)$ associated with a smooth<br>algebraic group . . . . .                          | 866        |
| <b>Current translation boundary</b>                                           |                                                                                                                               | <b>870</b> |

# Editorial Notice

We present here a slightly revised edition of the original Seminar, whose purpose and contents are described in the Introduction. The revision consisted mainly of correcting typographical errors, adding a few supplementary remarks or references in footnotes, dividing the work in its present form into three volumes, each with a detailed table of contents and an index of notation, and adding a terminological index at the end of Volume 3. In addition, Exposé VI<sub>B</sub>, by J.-E. Bertin, was partially rewritten by the author, especially Sections 5 and 10, so some references to that Exposé differ from the references to the original version. A list of the Exposés in the Seminar appears at the beginning of this volume.

Since the first edition of the present Seminar appeared, all of Chapter IV of the *Éléments de Géométrie Algébrique* has been published, making certain passages of the Seminar unnecessary. We have occasionally indicated in footnotes the relevant references to EGA IV that allow those passages to be bypassed.

For another treatment of algebraic groups that systematically uses the language of schemes, we refer to M. Demazure and P. Gabriel, *Groupes Algébriques* (North-Holland & Masson et Cie). Unlike the present Seminar, that book assumes no knowledge of algebraic geometry, but includes all the required preliminaries from scheme theory; it can therefore serve as an introduction to the study of this Seminar. (It also treats subjects not covered in the Seminar, such as the Dieudonné-style structure theory of commutative affine algebraic groups, in *loc. cit.*, Chapter V.)

Bures-sur-Yvette, March 1970





# Introduction

## 1. The purpose of this Seminar is twofold.

On the one hand, we aim to provide convenient foundations for the theory of group schemes in general. Exposés I–IV will supply the indispensable preliminary exercises in schematic and categorical syntax. To obtain a language that fits geometric intuition without effort, and to avoid circumlocutions that become intolerable in the long run, we always identify a prescheme  $X$  over another prescheme  $S$  with the functor

$$(\mathbf{Sch})_{/S}^{\circ} \longrightarrow \mathbf{Set}$$

that it represents\*. It is therefore necessary to formulate many definitions so that they apply to arbitrary functors, whether representable or not. Moreover, nearly all the functors we shall use will be “sheaves” (for the “faithfully flat quasi-compact topology”). Exposé IV, which treats groups only incidentally, sketches the language of “localization” and sheaves; this language also proves very convenient in questions concerning the representability of functors. Above all, that Exposé will provide the most convenient framework for the questions of passage to a quotient that arise later.

Exposé V gives some general results on the existence of quotients. They are taken up again in Exposé VI<sub>A</sub> for the quotient of an algebraic group over a field (or, more generally, over an Artinian ring) by a subgroup\*\*. That Exposé and the following Exposé VI<sub>B</sub> also contain various elementary results specific to algebraic groups over a field that are used routinely later.

Exposé VII studies certain phenomena related to the characteristic of the base field and, in particular, develops with the appropriate generality the correspondence between radicial group schemes of height 1 and restricted  $p$ -Lie algebras.

Finally, Exposé XVIII contains the scheme-theoretic generalization of Weil’s theorem on the “birational” definition of algebraic groups.

On the other hand, we intend to generalize the Borel–Chevalley structure theory of affine algebraic groups to groups over an arbitrary base prescheme. In fact, while the Seminar notes were being written, it became apparent that the affine hypothesis was unnecessary for many results of the theory. The most complete results are of course obtained for “semisimple

---

\*Such a viewpoint seems to have been considered for the first time eight or nine years ago, in connection with the theory of formal groups, by P. Cartier, who unfortunately did not take the trouble to state it precisely and systematize it as it deserved.

\*\*For a more thorough study of passage to the quotient, especially by reductive groups, see D. Mumford’s important work *Geometric Invariant Theory*, *Ergebnisse der Mathematik*, vol. 34, Springer, 1965. We note that on one important point the terminology of that book does not agree with ours: over a field of characteristic  $p > 0$ , the groups Mumford calls “reductive”<sup>(1)</sup> are the smooth groups of multiplicative type in the sense of the Seminar (cf. Exposé IX). One may reasonably suppose that Mumford adopted this meaning of “reductive,” which loses its significance over a base that is not a field, provisionally and as a kind of stopgap (and this is approximately what Mumford explains, for other reasons, as early as the second paragraph of his preface). <sup>(1)</sup>*Editor’s note:* See the remarks added at the end of this Introduction.

group schemes,” or more generally “reductive group schemes,” with which we shall be exclusively concerned beginning with Exposé XIX. Chevalley himself had already constructed the “Tôhoku groups” over the ring of integers, and that construction will be taken up in the present Seminar. The principal uniqueness theorem<sup>(2)</sup> gives a simple characterization of the “twisted” variants of these Tôhoku groups over a base prescheme  $S$ : they are the affine, smooth group schemes over  $S$  whose geometric fibers are connected semisimple groups in the usual sense<sup>\*\*\*</sup>(2).

## 2.

As in the familiar theory over an algebraically closed field, the subtori of the group schemes under consideration play a crucial role. Thus the preliminary study of tori and, more generally, of “group schemes of multiplicative type”—both intrinsically and as multiplicative-type subgroups of a given group—occupies a fairly large part of this Seminar (Exposés VIII–XII). Their remarkable rigidity (in some respects even greater than that of abelian schemes or semisimple group schemes) makes them highly effective tools for studying certain more general groups.

## 3.

Beginning with Exposé XII (apart from Exposé XVIII, already mentioned), we shall routinely use the theory of affine algebraic groups over an algebraically closed field, which the reader will find in the 1956 Chevalley Seminar, especially in Exposés IV–IX of that Seminar. We shall also use, though to a lesser extent, the later Exposés of the Chevalley Seminar devoted to the structure of semisimple algebraic groups. Indeed, we shall redevelop Chevalley’s theory directly within the framework of schemes; it will be seen that this approach makes the exposition simpler and more precise, even over a base field.

## 4.

The principal purpose of this Seminar is, of course, to develop techniques that apply to the study of group schemes over an arbitrary base, that is, essentially, to the study of families of algebraic groups. The infinitesimal properties of such families, and in particular the case of an Artinian base scheme, therefore play an important role. These properties arise even in the study of algebraic groups over a field  $k$  when the field is not perfect, especially when applying descent in the non-Galois case. Among the new results obtained in this setting, let us mention the existence of maximal tori and Cartan subgroups in an arbitrary smooth algebraic group, the rationality of the variety of maximal tori, and various related results (Exposé XIV<sup>(3)</sup>); another example is the correspondence between the “forms” of a semisimple group and homogeneous principal bundles under a suitable semisimple algebraic group (generally not connected) (Exposé XXIV, 1.16–1.20). In general, one may say that the methods required for work over a nonperfect base field are essentially those used over arbitrary base preschemes and thus lie outside the framework of classical algebraic geometry.

---

<sup>(2)</sup> *Editor’s note:* This refers to Corollary XXIII.5.6, which follows easily from the uniqueness theorem for split reductive groups (cf. XXIII, Theorem 4.1 and Corollary 5.2), since every semisimple  $S$ -group is a “twisted form” (for the étale topology) of a Tôhoku group (cf. XXII, Corollary 2.3).

<sup>\*\*\*</sup> This is the essential result of M. Demazure’s thesis, *Schémas en groupes réductifs*, Bull. Soc. Math. France **93** (1965), 369–413.

<sup>(3)</sup> *Editor’s note:* In particular, Theorems 1.1 and 6.1.

## 5.

It did not seem useful to indicate, at the beginning of the written Exposés, the date or dates of the corresponding oral presentations in the Seminar. We shall merely say that the order of the mimeographed Exposés (I–XXVI) does indeed agree with the order of the oral presentations. Moreover, the final written text was sometimes prepared considerably later than the oral presentation and often differs from it quite substantially: the written text is generally more detailed and more complete (as in Exposés IV and VII<sub>B</sub>), or even appreciably more general (as in Exposés XII and VII<sub>B</sub>), than the oral presentation. Other written Exposés correspond to no oral presentation (VI<sub>B</sub>, VII<sub>A</sub>, XV, XVI, XVII, and most of XXVI); they were written and inserted into the mimeographed Seminar either to provide convenient references for various other Exposés (notably in the case of VI<sub>B</sub>) or because they form a natural continuation of the concepts and techniques already developed. Consequently, Exposés VII<sub>A</sub>, VII<sub>B</sub>, XV, XVI, and XVII need not be read in order to study the rest of the Seminar, particularly the part devoted to reductive group schemes.

## 6.

From scheme theory we shall use primarily the general language of schemes set out in EGA I; the notions of flat, étale, and smooth morphisms set out in SGA 1, I–V; and the theory of faithfully flat descent in SGA 1, VIII. As far as possible, we have avoided imposing unnecessary Noetherian hypotheses. In return, this has required us to replace the usual hypothesis “of finite type” with the hypothesis “of finite presentation.” For the notion of a morphism of finite presentation, the reader should consult EGA IV, 1.4 and 1.6. The results of SGA 1, I–IV, which are most often stated in the Noetherian setting, will be developed in general in EGA IV\*\*\*\*; standard methods for reducing certain kinds of statements involving finite-presentation hypotheses to the Noetherian case will also be developed there in detail (EGA IV, Sections 8, 9, and 11). A reader who does not wish to assume these results from EGA IV may simplify certain statements or their proofs by assuming the base prescheme to be locally Noetherian. This does, however, lead to difficulties when a proof proceeds by descent from  $\hat{A}$  to  $A$ , where  $\hat{A}$  is the completion of a Noetherian local ring  $A$ , because that method introduces the generally non-Noetherian ring

$$\hat{A} \otimes_A \hat{A}.$$

## 7.

References follow the usual decimal system: the reference 5.7.11 denotes the proposition (or lemma, definition, and so forth) with that number in the same Exposé; in a reference such as XVII 7.8, the Roman numeral denotes the Exposé. We use the following abbreviations for our standard references:

**Bible** Chevalley Seminar, *Groupes de Lie algébriques*, 1956/58.

**EGA**  $X$ ,  $x.y.z$  J. Dieudonné and A. Grothendieck, *Éléments de Géométrie Algébrique*, Chapter  $X$ , statement  $x.y.z$  (or subsection  $x.y$ , and so forth).

**SGA**  $n$ ,  $X$   $y.z$  *Séminaire de Géométrie Algébrique du Bois-Marie*, year  $n$ , statement  $y.z$  of Exposé  $X$ .

---

\*\*\*\*Since this Introduction was written, all four parts (Sections 1–21) of EGA IV have appeared.

**TDTE** A. Grothendieck, *Techniques de descente et Théorèmes d'existence en Géométrie Algébrique*, Exposés delivered in the Bourbaki Seminar between 1959 and 1962.

<sup>(1)</sup>*Editor's note.* Concerning quotients by a reductive group, the situation has evolved considerably since A. Grothendieck wrote Note (\*\*). For an affine algebraic group over an arbitrary field  $k$ , the appropriate notion, introduced by Mumford, is that of a “geometrically reductive” group. (On the other hand,  $G$  is said to be “linearly reductive” if every rational representation of  $G$  is completely reducible; as noted in Note (\*\*), however, that condition is too restrictive when  $\text{char}(k) > 0$ .) By results of M. Nagata and W. J. Haboush ([Na64], [NM64], [Ha75]), the geometrically reductive  $k$ -groups are precisely the reductive  $k$ -groups in the sense of the present Seminar. For all of this, see the later editions of Mumford’s book ([MF82]). The notion of geometric reductivity over an arbitrary base, together with its consequences for passage to a quotient, was extended to this setting by C. S. Seshadri ([Se77]); additions due to M. Raynaud appear in the article [CTS79] by J.-L. Colliot-Thélène and J.-J. Sansuc. Finally, for more recent developments on passage to the quotient by an algebraic group or, more generally, by a groupoid (cf. Exposé V of the present Seminar), we refer to the articles [Ko97] and [KM97] by J. Kollár and by S. Keel and S. Mori.

## Bibliography

- [CTS79] J.-L. Colliot-Thélène and J.-J. Sansuc, *Fibrés quadratiques et composantes connexes réelles*, Math. Ann. **244** (1979), 105–134.
- [Ha75] W. J. Haboush, *Reductive groups are geometrically reductive*, Ann. of Math. **102** (1975), no. 1, 67–83.
- [KM97] S. Keel and S. Mori, *Quotient by groupoids*, Ann. of Math. **145** (1997), no. 1, 193–213.
- [Ko97] J. Kollár, *Quotient spaces modulo algebraic groups*, Ann. of Math. **145** (1997), no. 1, 33–79.
- [MF82] D. Mumford and J. Fogarty, *Geometric invariant theory*, 2nd ed., Springer-Verlag, 1982; 3rd ed., with F. Kirwan, 1994.
- [Na64] M. Nagata, *Invariants of a group in an affine ring*, J. Math. Kyoto Univ. **3** (1964), no. 3, 369–377.
- [NM64] M. Nagata and T. Miyata, *Note on semi-reductive groups*, J. Math. Kyoto Univ. **3** (1964), no. 3, 379–382.
- [Se77] C. S. Seshadri, *Geometric reductivity over an arbitrary base*, Adv. Math. **26** (1977), no. 3, 225–274.

# Exposé I

## Algebraic Structures. Group Cohomology

by *M. Demazure*

This exposé consists of two parts: the first collects a number of general definitions and establishes notation that will often be used in the sequel; the second treats group cohomology and culminates in Theorem 5.3.3, on the vanishing of the cohomology of diagonalizable groups.<sup>1</sup>

We choose once and for all a Universe.<sup>2</sup> All definitions and constructions will be relative to this Universe. We shall systematically allow ourselves the following abuse of language. To define a functor  $f : \mathcal{C} \rightarrow \mathcal{C}'$ , we shall be content to define the object  $f(S)$  of  $\mathcal{C}'$  for every object  $S$  of  $\mathcal{C}$ , whenever there is no ambiguity about how to define  $f(h)$  for an arrow  $h$  of  $\mathcal{C}$ . In practice, we shall say: let  $f : \mathcal{C} \rightarrow \mathcal{C}'$  be the functor defined by  $f(S) = \dots$ .

### 1. Generalities

#### 1.1.

Let  $\mathcal{C}$  be a category. We denote by  $\widehat{\mathcal{C}}$  the category

$$\mathrm{Hom}(\mathcal{C}^\circ, \mathbf{Set})$$

of contravariant functors from  $\mathcal{C}$  to the category  $\mathbf{Set}$  of sets.<sup>3</sup> There is a canonical functor

$$\mathbf{h} : \mathcal{C} \longrightarrow \widehat{\mathcal{C}}$$

which associates with every  $X \in \mathrm{Ob}(\mathcal{C})$  the functor  $\mathbf{h}_X$  such that

$$\mathbf{h}_X(S) = \mathrm{Hom}(S, X).$$

For every functor  $\mathbf{F} \in \mathrm{Ob}(\widehat{\mathcal{C}})$ , one has (see, for example, EGA 0<sub>III</sub>, 8.1.4) a bijection

$$\mathrm{Hom}(\mathbf{h}_X, \mathbf{F}) \xrightarrow{\sim} \mathbf{F}(X).$$

4

---

<sup>1</sup>Editor's note: version of October 13, 2024.

<sup>2</sup>Editor's note: see SGA 4, Exposé I, §0 and the Appendix; see also the discussion in [DG70], p. xxvi.

<sup>3</sup>Editor's note: this is called the category of presheaves on  $\mathcal{C}$ ; see IV, 4.3.1.

<sup>4</sup>Editor's note: this result is often called the Yoneda lemma; we use that terminology in other editor's notes.

In particular, for every pair  $X, Y$  of objects of  $\mathcal{C}$ , the canonical map

$$\mathrm{Hom}_{\mathcal{C}}(X, Y) \xrightarrow{\sim} \mathrm{Hom}_{\widehat{\mathcal{C}}}(\mathbf{h}_X, \mathbf{h}_Y)$$

is bijective; that is, the functor  $\mathbf{h}$  is fully faithful. It therefore defines an isomorphism of  $\mathcal{C}$  onto a full subcategory of  $\widehat{\mathcal{C}}$ , and an equivalence of  $\mathcal{C}$  with the full subcategory of  $\widehat{\mathcal{C}}$  formed by the representable functors (that is, the functors isomorphic to one of the form  $\mathbf{h}_X$ ). In the sequel we shall often identify  $X$  with  $\mathbf{h}_X$ . The following paragraphs are intended to show that this identification can be made safely.

**Remark 1.1.1.**<sup>5</sup> We sometimes need the following variant. Let  $\mathcal{D}$  be a full subcategory of  $\mathcal{C}$ , and let  $X, Y \in \mathrm{Ob}(\mathcal{D})$ . Denote by  $\mathbf{h}'_X$  and  $\mathbf{h}'_Y$  the restrictions of  $\mathbf{h}_X$  and  $\mathbf{h}_Y$  to  $\mathcal{D}$ . Then

$$\mathrm{Hom}_{\mathcal{C}}(X, Y) = \mathrm{Hom}_{\mathcal{D}}(X, Y) = \mathrm{Hom}_{\widehat{\mathcal{D}}}(\mathbf{h}'_X, \mathbf{h}'_Y).$$

Thus, to give a morphism  $X \rightarrow Y$  “is the same thing” as to give, for every  $T \in \mathrm{Ob}(\mathcal{D})$ , a map

$$\phi(T) : \mathrm{Hom}(T, X) \longrightarrow \mathrm{Hom}(T, Y)$$

functorially in  $T$ .

## 1.2.

We shall say that  $\mathbf{F}$  is a *subobject* (or a subfunctor) of  $\mathbf{G}$  if  $\mathbf{F}(S)$  is a subset of  $\mathbf{G}(S)$  for every  $S$ .<sup>6</sup>

In  $\widehat{\mathcal{C}}$ , arbitrary inverse limits exist and are computed pointwise:

$$\left( \varprojlim_i \mathbf{F}_i \right) (S) = \varprojlim_i \mathbf{F}_i(S).$$

<sup>7</sup> In particular, fiber products are defined by

$$(\mathbf{F} \times_{\mathbf{G}} \mathbf{F}')(S) = \mathbf{F}(S) \times_{\mathbf{G}(S)} \mathbf{F}'(S).$$

8

As a final object of  $\widehat{\mathcal{C}}$ , we choose the functor  $\mathbf{e}$  such that  $\mathbf{e}(S) = \{\emptyset\}$ .<sup>9</sup> Every  $\mathbf{F} \in \mathrm{Ob}(\widehat{\mathcal{C}})$  has a unique morphism to  $\mathbf{e}$ , and we set

$$\mathbf{F} \times \mathbf{F}' = \mathbf{F} \times_{\mathbf{e}} \mathbf{F}'.$$

The functor  $\mathbf{h}$  commutes with inverse limits. In particular, for  $X \times X'$  to exist, where  $X, X' \in \mathrm{Ob}(\mathcal{C})$ , respectively for  $\mathcal{C}$  to admit a final object  $e$ , it is necessary and sufficient that  $\mathbf{h}_X \times \mathbf{h}_{X'}$ , respectively  $\mathbf{e}$ , be representable; in that case

$$\mathbf{h}_X \times \mathbf{h}_{X'} \simeq \mathbf{h}_{X \times X'}, \quad \mathbf{h}_e \simeq \mathbf{e}.$$

<sup>5</sup>Editor’s note: this remark has been added.

<sup>6</sup>Editor’s note: the order has been changed so that fiber products are introduced before monomorphisms; see editor’s note (9).

<sup>7</sup>Editor’s note: similarly, arbitrary direct limits exist and are computed pointwise:  $(\varinjlim_i \mathbf{F}_i)(S) = \varinjlim_i \mathbf{F}_i(S)$ . In general, however, the functor  $\mathbf{h}$  does not commute with direct limits.

<sup>8</sup>Editor’s note: in particular, the equalizer of a pair of morphisms  $u, v : \mathbf{F} \rightrightarrows \mathbf{G}$  is the subfunctor  $\mathrm{Ker}(u, v)$  of  $\mathbf{F}$  defined by  $\mathrm{Ker}(u, v)(S) = \{x \in \mathbf{F}(S) \mid u(x) = v(x)\}$ .

<sup>9</sup>Editor’s note:  $\{\emptyset\}$ , the power set of the empty set, is a one-element set.

A monomorphism in  $\widehat{\mathcal{C}}$  is precisely a morphism  $\mathbf{F} \rightarrow \mathbf{G}$  such that, for every  $S \in \text{Ob}(\mathcal{C})$ , the corresponding map of sets  $\mathbf{F}(S) \rightarrow \mathbf{G}(S)$  is injective.<sup>10</sup>

The functor  $\Gamma$ . For every  $\mathbf{F} \in \text{Ob}(\widehat{\mathcal{C}})$ , set

$$\Gamma(\mathbf{F}) = \text{Hom}(\mathbf{e}, \mathbf{F}).$$

An element of  $\Gamma(\mathbf{F})$  is therefore a family  $(\gamma_S)_{S \in \text{Ob}(\mathcal{C})}$ , with  $\gamma_S \in \mathbf{F}(S)$ , such that for every arrow  $f : S' \rightarrow S''$  of  $\mathcal{C}$ ,

$$\mathbf{F}(f)(\gamma_{S''}) = \gamma_{S'}.$$

For  $X \in \text{Ob}(\mathcal{C})$ , set  $\Gamma(X) = \Gamma(\mathbf{h}_X)$ . If  $\mathcal{C}$  has a final object  $e$ , then

$$\Gamma(X) \simeq \text{Hom}(e, X).$$

### 1.3.

Let  $S \in \text{Ob}(\mathcal{C})$ . We denote by  $\mathcal{C}_{/S}$  the category of objects of  $\mathcal{C}$  over  $S$ : its objects are the arrows  $f : T \rightarrow S$  of  $\mathcal{C}$ , and  $\text{Hom}(f, f')$  is the subset of  $\text{Hom}(T, T')$  formed by the arrows  $u$  such that  $f = f' \circ u$ . If  $\mathcal{C}$  has a final object  $e$ , then  $\mathcal{C}_{/e}$  is isomorphic to  $\mathcal{C}$ . The category  $\mathcal{C}_{/S}$  has a final object, namely the identity arrow  $S \rightarrow S$ .

If  $f : T \rightarrow S$  is an object of  $\mathcal{C}_{/S}$ , one may form the category  $(\mathcal{C}_{/S})_{/f}$ , which by abuse of language we denote by  $(\mathcal{C}_{/S})_{/T}$ , and there is a canonical isomorphism

$$\mathcal{C}_{/T} \simeq (\mathcal{C}_{/S})_{/T}.$$

This construction also applies to  $\widehat{\mathcal{C}}$ ; in particular, one obtains the category  $\widehat{\mathcal{C}}_{/\mathbf{h}_S}$ . On the other hand, one may form the category  $\widehat{\mathcal{C}}_{/S}$ .

If  $f : T \rightarrow S$  is an object of  $\mathcal{C}_{/S}$ , then  $\Gamma(f)$  is identified with the set  $\Gamma(T/S)$  of sections of  $T$  over  $S$ , that is, arrows  $S \rightarrow T$  that are right inverses to  $f$ . Moreover,  $\mathbf{h}_f : \mathbf{h}_T \rightarrow \mathbf{h}_S$  is then an object of  $\widehat{\mathcal{C}}_{/\mathbf{h}_S}$ , and

$$\Gamma(\mathbf{h}_f) \simeq \Gamma(\mathbf{h}_T/\mathbf{h}_S) \simeq \Gamma(T/S) \simeq \Gamma(f).$$

### 1.4.

We now define an equivalence between the categories  $\widehat{\mathcal{C}}_{/S}$  and  $\widehat{\mathcal{C}}_{/\mathbf{h}_S}$ . In other words, we shall prove that to give a functor on the category of objects of  $\mathcal{C}$  over  $S$  is “the same thing” as to give a functor on  $\mathcal{C}$  equipped with a morphism to  $\mathbf{h}_S$ .

(i) **Construction of  $\alpha_S : \widehat{\mathcal{C}}_{/\mathbf{h}_S} \rightarrow \widehat{\mathcal{C}}_{/S}$ .**

First let  $H : \mathbf{F} \rightarrow \mathbf{h}_S$  be an object of  $\widehat{\mathcal{C}}_{/\mathbf{h}_S}$ . We must define a functor  $\alpha_S(H)$  on  $\mathcal{C}_{/S}$ . For an object  $f : T \rightarrow S$  of  $\mathcal{C}_{/S}$ , define  $\alpha_S(H)(f)$  to be the inverse image of  $f \in \mathbf{h}_S(T)$  under the map

$$H(T) : \mathbf{F}(T) \longrightarrow \mathbf{h}_S(T).$$

11

<sup>10</sup>Editor’s note: if  $\mathbf{F}(S) \rightarrow \mathbf{G}(S)$  is injective for every  $S$ , it is clear that  $\mathbf{F} \rightarrow \mathbf{G}$  is a monomorphism. The converse follows by considering  $\mathbf{F} \times_{\mathbf{G}} \mathbf{F} \rightrightarrows \mathbf{F} \rightarrow \mathbf{G}$ . Thus  $\mathbf{F} \rightarrow \mathbf{G}$  is a monomorphism if and only if the diagonal morphism  $\mathbf{F} \rightarrow \mathbf{F} \times_{\mathbf{G}} \mathbf{F}$  is an isomorphism (see EGA I, 5.3.8). Similarly, a pointwise surjective morphism is an epimorphism, and the converse follows by considering the pushout  $\mathbf{G} \amalg_{\mathbf{F}} \mathbf{G}$ ; see the proof of Lemma 4.4.4 in Exposé IV.

<sup>11</sup>Editor’s note: for example, if  $\mathbf{F} = \mathbf{h}_X$ , then  $H$  corresponds to a morphism  $h : X \rightarrow S$ , and  $H(T) : \text{Hom}_{\mathcal{C}}(T, X) \rightarrow \text{Hom}_{\mathcal{C}}(T, S)$  is the map  $g \mapsto h \circ g$ . Thus  $\alpha_S(\mathbf{h}_X) = \text{Hom}_{\mathcal{C}_{/S}}(-, X)$ .



Next, let  $u : f \rightarrow f'$  be an arrow of  $\mathcal{C}_{/S}$ . Then  $\mathbf{F}(u) : \mathbf{F}(T') \rightarrow \mathbf{F}(T)$  induces a map from  $\alpha_S(H)(f')$  to  $\alpha_S(H)(f)$ , denoted  $\alpha_S(H)(u)$ . It is immediate that the assignments

$$f \mapsto \alpha_S(H)(f), \quad u \mapsto \alpha_S(H)(u)$$

define a functor on  $\mathcal{C}_{/S}$ , hence an object  $\alpha_S(H)$  of  $\widehat{\mathcal{C}_{/S}}$ .

Finally, let  $H : \mathbf{F} \rightarrow \mathbf{h}_S$  and  $H' : \mathbf{F}' \rightarrow \mathbf{h}_S$  be two objects of  $\widehat{\mathcal{C}_{/\mathbf{h}_S}}$ , and let  $U : H \rightarrow H'$  be a morphism in that category:

$$\begin{array}{ccc} \mathbf{F} & \xrightarrow{U} & \mathbf{F}' \\ & \searrow H & \swarrow H' \\ & \mathbf{h}_S & \end{array} .$$

For every  $f : T \rightarrow S$ , the map  $U(T) : \mathbf{F}(T) \rightarrow \mathbf{F}'(T)$  induces a map

$$\alpha_S(U)(f) : \alpha_S(H)(f) \rightarrow \alpha_S(H')(f),$$

which defines a morphism of functors

$$\alpha_S(U) : \alpha_S(H) \rightarrow \alpha_S(H').$$

It is easy to verify that the assignments

$$H \mapsto \alpha_S(H), \quad U \mapsto \alpha_S(U)$$

define a functor

$$\alpha_S : \widehat{\mathcal{C}_{/\mathbf{h}_S}} \rightarrow \widehat{\mathcal{C}_{/S}}.$$

**(ii) Proposition 1.4.1.** *The functor  $\alpha_S : \widehat{\mathcal{C}_{/\mathbf{h}_S}} \rightarrow \widehat{\mathcal{C}_{/S}}$  is an equivalence of categories.*

We indicate only the principle of the construction of a quasi-inverse functor

$$\beta_S : \widehat{\mathcal{C}_{/S}} \rightarrow \widehat{\mathcal{C}_{/\mathbf{h}_S}}.$$

Let  $\mathbf{G}$  be a functor on  $\mathcal{C}_{/S}$ . For every object  $T$  of  $\mathcal{C}$ , set

$$\beta_S(\mathbf{G})(T) = \coprod_{f \in \text{Hom}(T, S)} \mathbf{G}(f), \quad \text{Hom}(T, S) = \mathbf{h}_S(T).$$

This defines a functor  $\beta_S(\mathbf{G})$  on  $\mathcal{C}$ , equipped with the evident projection to  $\mathbf{h}_S$ .

### 1.5.

The equivalence  $\alpha_S$  commutes with the functors  $\Gamma$ . In other words, if  $H : \mathbf{F} \rightarrow \mathbf{h}_S$  is an object of  $\widehat{\mathcal{C}_{/\mathbf{h}_S}}$  and  $\alpha_S(H)$  is the corresponding object of  $\widehat{\mathcal{C}_{/S}}$ , then

$$\Gamma(\alpha_S(H)) \simeq \Gamma(H) \simeq \Gamma(\mathbf{F}/\mathbf{h}_S).$$

The equivalence  $\alpha_S$  also commutes with the functors  $\mathbf{h}$ . That is, if  $f : T \rightarrow S$  is an object of  $\mathcal{C}_{/S}$ , then  $\mathbf{h}_f : \mathbf{h}_T \rightarrow \mathbf{h}_S$  is an object of  $\widehat{\mathcal{C}_{/\mathbf{h}_S}}$ , and its image under  $\alpha_S$  is precisely  $\mathbf{h}_{\mathcal{C}_{/S}}(f)$ , where

$$\mathbf{h}_{\mathcal{C}_{/S}} : \mathcal{C}_{/S} \rightarrow \widehat{\mathcal{C}_{/S}}$$

is the canonical functor.<sup>12</sup> Consequently:

**Proposition 1.5.1.** *Let  $H : \mathbf{F} \rightarrow \mathbf{h}_S$  be an object of  $\widehat{\mathcal{C}}_{\mathbf{h}_S}$ . The functor*

$$\alpha_S(H) : (\mathcal{C}/S)^\circ \longrightarrow \mathbf{Set}$$

*is representable if and only if  $\mathbf{F} : \mathcal{C}^\circ \rightarrow \mathbf{Set}$  is representable. If  $\mathbf{F} \simeq \mathbf{h}_T$ , then  $\alpha_S(H)$  is represented by the object  $T \rightarrow S$  of  $\mathcal{C}/S$ .*

The equivalence  $\alpha_S$  is transitive in  $S$ . If  $f : T \rightarrow S$  is an object of  $\mathcal{C}/S$ , there is a commutative diagram of equivalences:

$$\begin{array}{ccccc} (\widehat{\mathcal{C}}_{\mathbf{h}_S})_{/\mathbf{h}_T} & \xrightarrow{\alpha_S/\mathbf{h}_T} & (\widehat{\mathcal{C}/S})_{/\mathbf{h}_T} & \xrightarrow{\alpha_f} & (\widehat{\mathcal{C}/S})_{/T} \\ & \searrow \cong & & & \swarrow \cong \\ & & \widehat{\mathcal{C}}_{/\mathbf{h}_T} & \xrightarrow{\alpha_T} & \widehat{\mathcal{C}}_{/T} \end{array},$$

Here  $\alpha_S/\mathbf{h}_T$  denotes, provisionally, the restriction (see 1.6) of the functor  $\alpha_S$  to the objects over  $\mathbf{h}_T$ .

## 1.6. Base change in a functor

For every  $S \in \text{Ob}(\mathcal{C})$ , there is a canonical functor

$$i_S : \mathcal{C}/S \longrightarrow \mathcal{C}$$

defined by  $i_S(f) = T$  when  $f$  is the arrow  $T \rightarrow S$ . If  $f : T \rightarrow S$  is an object of  $\mathcal{C}/S$ , we denote by  $i_{T/S} = i_f$  the functor

$$i_{T/S} : (\mathcal{C}/S)_{/T} \longrightarrow \mathcal{C}/S,$$

and there is a commutative diagram:

$$\begin{array}{ccccc} (\mathcal{C}/S)_{/T} & \xrightarrow{i_{T/S}} & \mathcal{C}/S & \xrightarrow{i_S} & \mathcal{C} \\ & \searrow \cong & & & \swarrow i_T \\ & & \mathcal{C}_{/T} & & \end{array},$$

Thus, identifying  $(\mathcal{C}/S)_{/T}$  with  $\mathcal{C}_{/T}$ , as we shall henceforth do,

$$i_S \circ i_{T/S} = i_T.$$

Likewise, if  $\mathcal{C}$  is identified with  $\mathcal{C}_{/e}$  when  $\mathcal{C}$  has a final object  $e$ , then  $i_{S/e} : \mathcal{C}/S \rightarrow \mathcal{C}_{/e}$  is identified with  $i_S$ .

For  $X \in \text{Ob}(\mathcal{C})$  (respectively,  $Y \in \text{Ob}(\mathcal{C}/S)$ ), let  $p_S(X)$  (respectively,  $p_{T/S}(Y)$ ) be the object of  $\mathcal{C}/S$  (respectively, of  $\mathcal{C}_{/T}$ ), when it exists, defined by  $X \times_S$  (respectively,  $Y \times_S T$ ) equipped with its second projection:

<sup>12</sup>Editor's note: see editor's note (10).

$$\begin{array}{ccc}
X \times S & & Y \times_S T \\
p_S(X) \downarrow & \text{resp.} & \downarrow p_{T/S}(Y) \\
S & & T
\end{array}
.$$

The partially defined functor  $p_S$  (respectively,  $p_{T/S}$ ) is called the *base-change functor*. By the definition of a product (respectively, a fiber product), it is right adjoint to  $i_S$  (respectively, to  $i_{T/S}$ ).<sup>13</sup> We also write

$$p_S(X) = X_S, \quad p_{T/S}(Y) = Y_T.$$

The functor  $i_S$  defines a restriction functor

$$i_S^* : \widehat{\mathcal{C}} \longrightarrow \widehat{\mathcal{C}_{/S}}.$$

We write

$$\mathbf{F}_S = i_S^*(\mathbf{F}) = \mathbf{F} \circ i_S.$$

Plainly,

$$i_{T/S}^* \circ i_S^* = i_T^*,$$

that is, for every functor  $\mathbf{F} \in \text{Ob}(\widehat{\mathcal{C}})$ ,

$$(\mathbf{F}_S)_T = \mathbf{F}_T.$$

The notation is justified by the following result.

**Proposition 1.6.1.** *The functor*

$$(\mathbf{h}_X)_S : (\mathcal{C}_{/S})^\circ \longrightarrow \mathbf{Set}$$

*is representable if and only if the product  $X \times S$  exists. In that case*

$$(\mathbf{h}_X)_S \simeq \mathbf{h}_{X_S}.$$

This shows that  $\mathbf{F}_S$  has two interpretations: it is the restriction of  $\mathbf{F}$  to  $\mathcal{C}_{/S}$ , and it is the functor obtained by the base change  $e \leftarrow S$ . This leads to the following notation:

$$\begin{array}{ccccc}
\mathbf{F} & \xrightarrow{\quad} & \mathbf{F}_S & \xrightarrow{\quad} & \mathbf{F}_T \\
\downarrow & & \downarrow & & \downarrow \\
e & \xleftarrow{\quad} & S & \xleftarrow{\quad} & T
\end{array}$$

<sup>13</sup>Editor's note: that is, if  $g : U \rightarrow S$  (respectively,  $h : V \rightarrow T$ ) is an object of  $\mathcal{C}_{/S}$  (respectively, of  $\mathcal{C}_{/T}$ ), then  $i_S(g) = U$ , while  $i_{T/S}(h)$  is the object  $f \circ h : V \rightarrow S$  of  $\mathcal{C}_{/S}$ , and

$$\text{Hom}_{\mathcal{C}_{/S}}(U, X \times S) \simeq \text{Hom}_{\mathcal{C}}(U, X),$$

respectively,

$$\text{Hom}_{\mathcal{C}_{/T}}(V, X \times_S T) \simeq \text{Hom}_{\mathcal{C}_{/S}}(V, X).$$

This diagram reflects both of the preceding interpretations. Notice that

$$\Gamma(\mathbf{F}_S) \simeq \text{Hom}(\mathbf{h}_S, \mathbf{F}) \simeq \mathbf{F}(S),$$

and, in particular,

$$\Gamma(X_S) \simeq \text{Hom}(S, X).$$

### 1.7.0.

Let  $\mathbf{E}$  be an object of  $\widehat{\mathcal{C}}$ .<sup>14</sup> Consider the category  $\mathcal{C}/_{\mathbf{E}}$  of *objects of  $\mathcal{C}$  over  $\mathbf{E}$* . Its objects are pairs  $(V, \rho)$  formed by an object  $V$  of  $\mathcal{C}$  and a morphism  $\rho : \mathbf{h}_V \rightarrow \mathbf{E}$  in  $\widehat{\mathcal{C}}$ , that is,  $\rho \in \mathbf{E}(V)$ . A morphism from  $(V, \rho)$  to  $(V', \rho')$  is a morphism  $f : V \rightarrow V'$  in  $\mathcal{C}$  such that

$$\rho' \circ f = \rho, \quad \text{that is,} \quad \mathbf{E}(f)(\rho') = \rho.$$

Let  $\mathbf{L}$  be the functor

$$\mathbf{L} = \varinjlim_{(V, \rho) \in \mathcal{C}/_{\mathbf{E}}} \mathbf{h}_V.$$

Thus, for every  $S \in \text{Ob}(\mathcal{C})$ ,

$$\mathbf{L}(S) = \varinjlim_{(V, \rho) \in \mathcal{C}/_{\mathbf{E}}} \mathbf{h}_V(S)$$

is the set of equivalence classes of triples  $(V, \rho, v)$ , where  $v : S \rightarrow V$  is a morphism in  $\mathcal{C}$ , and where  $(V, \rho, v)$  is identified with  $(V', \rho', f \circ v)$  for every morphism  $f : V \rightarrow V'$  in  $\mathcal{C}$  such that  $\rho' \circ f = \rho$ .

The map  $\phi_{\mathbf{E}}(S)$  which associates with the class of  $(V, \rho, v)$  the element  $\rho \circ v$  of  $\mathbf{E}(S)$  is well defined and gives a morphism of functors

$$\phi_{\mathbf{E}} : \varinjlim_{(V, \rho) \in \mathcal{C}/_{\mathbf{E}}} \mathbf{h}_V \longrightarrow \mathbf{E}.$$

**Lemma.** *The morphism  $\phi_{\mathbf{E}}$  is an isomorphism.*

Indeed, let  $S \in \text{Ob}(\mathcal{C})$ . Every  $x \in \mathbf{E}(S)$  is the image under  $\phi_{\mathbf{E}}(S)$  of the triple  $(S, x, \text{id}_S)$ ; hence  $\phi_{\mathbf{E}}(S)$  is surjective. Conversely, let  $\ell_1 = (V_1, \rho_1, v_1)$  and  $\ell_2 = (V_2, \rho_2, v_2)$  be two elements of  $\mathbf{L}(S)$  having the same image in  $\mathbf{E}(S)$ , and set

$$\gamma = \rho_1 \circ v_1 = \rho_2 \circ v_2.$$

Then  $\ell_1$  and  $\ell_2$  are both equal in  $\mathbf{L}(S)$  to the class of  $(S, \gamma, \text{id}_S)$ . Thus  $\phi_{\mathbf{E}}(S)$  is injective.

**Corollary.** *For every object  $\mathbf{F}$  of  $\widehat{\mathcal{C}}$ ,*

$$\text{Hom}(\mathbf{E}, \mathbf{F}) = \varprojlim_{(V, \rho) \in \mathcal{C}/_{\mathbf{E}}} \mathbf{F}(V).$$

---

<sup>14</sup>Editor's note: the lemma below has been added (see SGA 4, I.3.4); it is used in the proof of 1.7.1 and will be useful repeatedly in the sequel.

### 1.7. The objects $\underline{\text{Hom}}$ , $\underline{\text{Isom}}$ , etc.

Let  $\mathbf{F}$  and  $\mathbf{G}$  be two objects of  $\widehat{\mathcal{C}}$ . We define another object of  $\widehat{\mathcal{C}}$  by

$$\begin{aligned}\underline{\text{Hom}}(\mathbf{F}, \mathbf{G})(S) &= \text{Hom}_{\widehat{\mathcal{C}}/S}(\mathbf{F}_S, \mathbf{G}_S) \\ &\simeq \text{Hom}_{\widehat{\mathcal{C}}/\mathbf{h}_S}(\mathbf{F} \times \mathbf{h}_S, \mathbf{G} \times \mathbf{h}_S) \\ &\simeq \text{Hom}_{\widehat{\mathcal{C}}}(\mathbf{F} \times \mathbf{h}_S, \mathbf{G}).\end{aligned}$$

The object  $\underline{\text{Hom}}(\mathbf{F}, \mathbf{G})$  has the following properties:

(i)  $\underline{\text{Hom}}(\mathbf{e}, \mathbf{G}) \simeq \mathbf{G}$ .<sup>15</sup>

(ii) The formation of  $\underline{\text{Hom}}$  commutes with base extension:

$$\underline{\text{Hom}}(\mathbf{F}_S, \mathbf{G}_S) \simeq \underline{\text{Hom}}(\mathbf{F}, \mathbf{G})_S.$$

(iii)  $(\mathbf{F}, \mathbf{G}) \mapsto \underline{\text{Hom}}(\mathbf{F}, \mathbf{G})$  is a bifunctor, contravariant in  $\mathbf{F}$  and covariant in  $\mathbf{G}$ .

These three properties follow immediately from the definitions.

We shall prove that, for every object  $\mathbf{E}$  of  $\widehat{\mathcal{C}}$ ,

$$\text{Hom}(\mathbf{E}, \underline{\text{Hom}}(\mathbf{F}, \mathbf{G})) \simeq \text{Hom}(\mathbf{E} \times \mathbf{F}, \mathbf{G}).$$

Let  $\phi : \mathbf{E} \times \mathbf{F} \rightarrow \mathbf{G}$ . We must associate with it a morphism from  $\mathbf{E}$  to  $\underline{\text{Hom}}(\mathbf{F}, \mathbf{G})$ . For an arrow  $S' \rightarrow S$  of  $\mathcal{C}$ , there are maps

$$\mathbf{E}(S) \times \mathbf{F}(S') \longrightarrow \mathbf{E}(S') \times \mathbf{F}(S') \xrightarrow{\phi(S')} \mathbf{G}(S').$$

Every element  $e$  of  $\mathbf{E}(S)$  therefore defines, for each  $S' \rightarrow S$ , a map  $\mathbf{F}(S') \rightarrow \mathbf{G}(S')$  functorial in  $S'$ ; this is an element  $\theta_\phi(e)$  of  $\underline{\text{Hom}}(\mathbf{F}, \mathbf{G})(S)$ . We have thus obtained a map

$$\phi \longmapsto \theta_\phi, \quad \text{Hom}(\mathbf{E} \times \mathbf{F}, \mathbf{G}) \longrightarrow \text{Hom}(\mathbf{E}, \underline{\text{Hom}}(\mathbf{F}, \mathbf{G})),$$

which is functorial in  $\mathbf{E}$ .

**Proposition 1.7.1.**<sup>16</sup> *Let  $\mathbf{E}, \mathbf{F}, \mathbf{G} \in \text{Ob}(\widehat{\mathcal{C}})$ .*

(a) *The map  $\phi \mapsto \theta_\phi$  is a bijection:*

$$\text{Hom}(\mathbf{E} \times \mathbf{F}, \mathbf{G}) \xrightarrow{\sim} \text{Hom}(\mathbf{E}, \underline{\text{Hom}}(\mathbf{F}, \mathbf{G})).$$

(b) *Moreover, there is an isomorphism of functors*

$$\underline{\text{Hom}}(\mathbf{E}, \underline{\text{Hom}}(\mathbf{F}, \mathbf{G})) \simeq \underline{\text{Hom}}(\mathbf{E} \times \mathbf{F}, \mathbf{G}).$$

---

<sup>15</sup>Editor's note: if  $\mathbf{E}$  is a third object of  $\widehat{\mathcal{C}}$ , then

$$\underline{\text{Hom}}(\mathbf{E}, \mathbf{F} \times \mathbf{G}) \simeq \underline{\text{Hom}}(\mathbf{E}, \mathbf{F}) \times \underline{\text{Hom}}(\mathbf{E}, \mathbf{G}).$$

<sup>16</sup>Editor's note: part (b), which will be useful in II.1 and II.3.11, has been added. A second, more direct proof of part (a) has also been sketched.

For (a), regard both sides as functors of  $\mathbf{E}$ . The asserted result is true when  $\mathbf{E} = \mathbf{h}_X$ , since it is then precisely the definition of the functor  $\underline{\text{Hom}}(\mathbf{F}, \mathbf{G})$ . Moreover, both sides, as functors of  $\mathbf{E}$ , take direct limits to inverse limits. Finally, by Lemma 1.7.0, every object  $\mathbf{E}$  of  $\mathcal{C}$  is isomorphic to the direct limit of the  $\mathbf{h}_X$ , as  $X$  ranges over  $\mathcal{C}/\mathbf{E}$ . This proves (a).

Let us also sketch a direct proof of (a). To every

$$\theta \in \text{Hom}(\mathbf{E}, \underline{\text{Hom}}(\mathbf{F}, \mathbf{G}))$$

associate  $\phi_\theta \in \text{Hom}(\mathbf{E} \times \mathbf{F}, \mathbf{G})$  as follows. For every  $S \in \text{Ob}(\mathcal{C})$ , there is a map

$$\theta(S) : \mathbf{E}(S) \longrightarrow \text{Hom}(\mathbf{F} \times S, \mathbf{G}),$$

functorial in  $S$ . If  $(e, f) \in \mathbf{E}(S) \times \mathbf{F}(S)$ , then  $f$  is a morphism  $S \rightarrow \mathbf{F}$ , so  $f \times \text{id}_S$  is a morphism  $S \rightarrow \mathbf{F} \times S$ . On the other hand,  $\theta(S)(e)$  is a morphism  $\mathbf{F} \times S \rightarrow \mathbf{G}$ . Composition therefore gives a morphism

$$\theta(S)(e) \circ (f \times \text{id}_S) : S \longrightarrow \mathbf{G},$$

that is, an element  $\phi_\theta(S)(e, f)$  of  $\mathbf{G}(S)$ . It is easy to check that  $S \mapsto \phi_\theta(S)$  is functorial in  $S$ , and therefore defines a morphism  $\phi_\theta$  from  $\mathbf{E} \times \mathbf{F}$  to  $\mathbf{G}$ . We leave to the reader the verification that  $\theta \mapsto \phi_\theta$  and  $\phi \mapsto \theta_\phi$  are mutually inverse bijections.

For (b), if  $S \in \text{Ob}(\mathcal{C})$ , then by 1.7(ii) and part (a), applied to  $\mathcal{C}/_S$ ,

$$\begin{aligned} & \underline{\text{Hom}}(\mathbf{E}, \underline{\text{Hom}}(\mathbf{F}, \mathbf{G}))(S) \\ & \simeq \text{Hom}_S(\mathbf{E}_S, \underline{\text{Hom}}_S(\mathbf{F}_S, \mathbf{G}_S)) \\ & \simeq \text{Hom}_S(\mathbf{E}_S \times_S \mathbf{F}_S, \mathbf{G}_S) \\ & \simeq \text{Hom}(\mathbf{E} \times \mathbf{F} \times S, \mathbf{G}) \\ & \simeq \underline{\text{Hom}}(\mathbf{E} \times \mathbf{F}, \mathbf{G})(S), \end{aligned}$$

and these isomorphisms are functorial in  $S$ .

**Corollary 1.7.2.** *There are isomorphisms*

$$\text{Hom}(\mathbf{E}, \underline{\text{Hom}}(\mathbf{F}, \mathbf{G})) \simeq \text{Hom}(\mathbf{F}, \underline{\text{Hom}}(\mathbf{E}, \mathbf{G})),$$

and

$$\underline{\text{Hom}}(\mathbf{E}, \underline{\text{Hom}}(\mathbf{F}, \mathbf{G})) \simeq \underline{\text{Hom}}(\mathbf{F}, \underline{\text{Hom}}(\mathbf{E}, \mathbf{G})).$$

In particular, taking  $\mathbf{E} = \underline{\mathbf{e}}$  and using  $\underline{\text{Hom}}(\underline{\mathbf{e}}, \mathbf{G}) \simeq \mathbf{G}$ , we obtain

$$\Gamma(\underline{\text{Hom}}(\mathbf{F}, \mathbf{G})) \simeq \text{Hom}(\mathbf{F}, \mathbf{G}).$$

Composition supplies functorial morphisms

$$\underline{\text{Hom}}(\mathbf{F}, \mathbf{G}) \times \underline{\text{Hom}}(\mathbf{G}, \mathbf{H}) \longrightarrow \underline{\text{Hom}}(\mathbf{F}, \mathbf{H}).$$

If  $\mathbf{F}$  and  $\mathbf{G}$  are two objects of  $\widehat{\mathcal{C}}$ , let  $\text{Isom}(\mathbf{F}, \mathbf{G})$  denote the subset of  $\text{Hom}(\mathbf{F}, \mathbf{G})$  formed by the isomorphisms from  $\mathbf{F}$  onto  $\mathbf{G}$ . Define a subobject  $\underline{\text{Isom}}(\mathbf{F}, \mathbf{G})$  of  $\underline{\text{Hom}}(\mathbf{F}, \mathbf{G})$  by

$$\underline{\text{Isom}}(\mathbf{F}, \mathbf{G})(S) = \text{Isom}(\mathbf{F}_S, \mathbf{G}_S).$$

Then

$$\Gamma(\underline{\text{Isom}}(\mathbf{F}, \mathbf{G})) \simeq \text{Isom}(\mathbf{F}, \mathbf{G}), \quad \underline{\text{Isom}}(\mathbf{F}, \mathbf{G}) \simeq \underline{\text{Isom}}(\mathbf{G}, \mathbf{F}).$$

In the special case  $\mathbf{F} = \mathbf{G}$ , set

$$\begin{aligned}\underline{\text{End}}(\mathbf{F}) &= \underline{\text{Hom}}(\mathbf{F}, \mathbf{F}), & \text{End}(\mathbf{F}) &= \text{Hom}(\mathbf{F}, \mathbf{F}) \simeq \Gamma(\underline{\text{End}}(\mathbf{F})), \\ \underline{\text{Aut}}(\mathbf{F}) &= \underline{\text{Isom}}(\mathbf{F}, \mathbf{F}), & \text{Aut}(\mathbf{F}) &= \text{Isom}(\mathbf{F}, \mathbf{F}) \simeq \Gamma(\underline{\text{Aut}}(\mathbf{F})).\end{aligned}$$

Formation of the objects  $\underline{\text{Hom}}$ ,  $\underline{\text{Isom}}$ ,  $\underline{\text{Aut}}$ , and  $\underline{\text{End}}$  commutes with base change.

**Remark 1.7.3.** <sup>17</sup> An object isomorphic to  $\underline{\text{Isom}}(\mathbf{F}, \mathbf{G})$  may also be constructed as follows. There is a morphism

$$\underline{\text{Hom}}(\mathbf{F}, \mathbf{G}) \times \underline{\text{Hom}}(\mathbf{G}, \mathbf{F}) \longrightarrow \underline{\text{End}}(\mathbf{F}).$$

Interchanging  $\mathbf{F}$  and  $\mathbf{G}$  gives a morphism

$$\underline{\text{Hom}}(\mathbf{F}, \mathbf{G}) \times \underline{\text{Hom}}(\mathbf{G}, \mathbf{F}) \longrightarrow \underline{\text{End}}(\mathbf{F}) \times \underline{\text{End}}(\mathbf{G}).$$

On the other hand, the identity morphism of  $\mathbf{F}$  is an element of  $\text{End}(\mathbf{F})$ , and hence defines a morphism  $\underline{\mathbf{e}} \rightarrow \underline{\text{End}}(\mathbf{F})$ . Doing the same for  $\mathbf{G}$  and taking the product gives a morphism

$$\underline{\mathbf{e}} \longrightarrow \underline{\text{End}}(\mathbf{F}) \times \underline{\text{End}}(\mathbf{G}).$$

It is then immediate that the fiber product of  $\underline{\mathbf{e}}$  and

$$\underline{\text{Hom}}(\mathbf{F}, \mathbf{G}) \times \underline{\text{Hom}}(\mathbf{G}, \mathbf{F})$$

over  $\underline{\text{End}}(\mathbf{F}) \times \underline{\text{End}}(\mathbf{G})$  is isomorphic to  $\underline{\text{Isom}}(\mathbf{F}, \mathbf{G})$ .

All these definitions apply, in particular, when  $\mathbf{F} = \mathbf{h}_X$  and  $\mathbf{G} = \mathbf{h}_Y$ . If  $\underline{\text{Hom}}(\mathbf{h}_X, \mathbf{h}_Y)$  is representable by an object of  $\mathcal{C}$ , we denote that object by  $\underline{\text{Hom}}(X, Y)$ . It has the following property: if  $Z \times X$  exists, then

$$\text{Hom}(Z, \underline{\text{Hom}}(X, Y)) \simeq \text{Hom}(Z \times X, Y).$$

This property characterizes it when products exist in  $\mathcal{C}$ .

Similarly, one defines, when they exist, the objects

$$\underline{\text{Isom}}(X, Y), \quad \underline{\text{End}}(X), \quad \underline{\text{Aut}}(X).$$

We merely note that the construction above shows that  $\underline{\text{Isom}}(X, Y)$  exists whenever fiber products exist in  $\mathcal{C}$  and the objects

$$\underline{\text{Hom}}(X, Y), \quad \underline{\text{Hom}}(Y, X), \quad \underline{\text{End}}(X), \quad \text{and} \quad \underline{\text{End}}(Y)$$

exist.

Everything above also applies to categories of the form  $\mathcal{C}_{/S}$ . The corresponding objects will be denoted as explicitly as possible by appropriate symbols. For example, if  $T$  and  $T'$  are two objects of  $\mathcal{C}$  over  $S$ , we denote by  $\underline{\text{Hom}}_S(T, T')$  the object

$$\underline{\text{Hom}}_{\mathcal{C}_{/S}}(T/S, T'/S).$$

---

<sup>17</sup>Editor's note: the numbering 1.7.3 has been added for later references.

### 1.8. Constant objects

Let  $\mathcal{C}$  be a category in which direct sums and fiber products exist and in which direct sums commute with base change (for example, the category of schemes).<sup>18</sup> For every set  $E$  and every object  $S$  of  $\mathcal{C}$ , set

$$E_S = \{\text{the direct sum of a family } (S_i)_{i \in E} \text{ of objects of } \mathcal{C}, \text{ all isomorphic to } S. \quad (1.8.1)$$

This object is characterized by the formula

$$\mathrm{Hom}_{\mathcal{C}}(E_S, T) = \mathrm{Hom}_{\mathbf{Set}}(E, \mathrm{Hom}_{\mathcal{C}}(S, T)), \quad (1.8.2)$$

for every  $T \in \mathrm{Ob}(\mathcal{C})$ .

The object  $E_S$  has a canonical projection to  $S$ , so that  $E \mapsto E_S$  is in fact a functor from  $\mathbf{Set}$  to  $\mathcal{C}/_S$ .<sup>19</sup> If  $S' \rightarrow S$  is an arrow of  $\mathcal{C}$ , then, because direct sums commute with base change,

$$E_{S'} = (E_S)_{S'}.$$

In particular, if  $\mathcal{C}$  has a final object  $e$ , then

$$E_S = (E_e)_S.$$

The functor  $E \mapsto E_S/S$ , from  $\mathbf{Set}$  to  $\mathcal{C}/_S$ , commutes with finite products. It is enough to prove that

$$E_S \times_S F_S = (E \times F)_S. \quad (\times)$$

Indeed, by the results of 1.7 applied to  $\mathcal{C}/_S$ , for every  $T \in \mathrm{Ob}(\mathcal{C}/_S)$  there are natural isomorphisms (all unspecified  $\mathrm{Hom}$ 's being taken in  $\mathcal{C}/_S$ )

$$\begin{aligned} \mathrm{Hom}((E \times F)_S, T) &\simeq \mathrm{Hom}_{\mathbf{Set}}(E \times F, \mathrm{Hom}(S, T)) \\ &\simeq \mathrm{Hom}_{\mathbf{Set}}(E, \mathrm{Hom}_{\mathbf{Set}}(F, \mathrm{Hom}(S, T))) \\ &\simeq \mathrm{Hom}_{\mathbf{Set}}(E, \mathrm{Hom}(F_S, T)), \end{aligned}$$

and

$$\begin{aligned} \mathrm{Hom}(E_S \times_S F_S, T) &\simeq \mathrm{Hom}(E_S, \underline{\mathrm{Hom}}(F_S, T)) \\ &\simeq \mathrm{Hom}_{\mathbf{Set}}(E, \mathrm{Hom}(S, \underline{\mathrm{Hom}}(F_S, T))). \end{aligned}$$

But

$$\mathrm{Hom}(S, \underline{\mathrm{Hom}}(F_S, T)) \simeq \mathrm{Hom}(F_S, T),$$

which proves  $(\times)$ .

Suppose that, with  $\emptyset$  denoting an initial object of  $\mathcal{C}$ , the following diagram is cartesian:

$$(*) \quad \begin{array}{ccc} \emptyset & \xrightarrow{\quad} & S \\ \downarrow & & \downarrow \\ S & \xrightarrow{\quad} & S \amalg S \end{array}$$

<sup>18</sup>Editor's note: the former terminology "preschemes/schemes" has been replaced by the current terminology "schemes/separated schemes."

<sup>19</sup>Editor's note: in the next three paragraphs, the order of the sentences has been changed and a few details have been added concerning the role of condition  $(*)$  below.



This is the case for the category of schemes.<sup>20</sup> Then the functor  $E \mapsto E_S$  commutes with finite inverse limits.

Indeed, in view of  $(\times)$ , it is enough to show that  $E \mapsto E_S$  commutes with fiber products. Let  $u : E \rightarrow G$  and  $v : F \rightarrow G$  be two maps of sets. Since direct sums in  $\mathcal{C}$  commute with base change, one has

$$F_S \times_{G_S} E_S \simeq \coprod_{f \in F} (S_f \times_{G_S} E_S) \simeq \coprod_{\substack{f \in F \\ x \in E}} (S_f \times_{G_S} S_x). \quad (1)$$

If  $v(f) \neq u(x)$ , there is a morphism in  $\mathcal{C}/_S$

$$S_f \times_{G_S} S_x \longrightarrow S_f \times_{S_{v(f)} \amalg S_{u(x)}} S_x.$$

By condition  $(*)$ , the term on the right is  $\emptyset$ . Hence

$$S_f \times_{G_S} S_x \simeq \emptyset \quad \text{if } v(f) \neq u(x). \quad (2)$$

On the other hand, if  $v(f) = u(x)$ , there is a morphism in  $\mathcal{C}/_S$

$$S \longrightarrow S_f \times_{G_S} S_x.$$

Since  $S \xrightarrow{\text{id}} S$  is the final object of  $\mathcal{C}/_S$ , it follows that

$$S_f \times_{G_S} S_x \simeq S \quad \text{if } v(f) = u(x). \quad (3)$$

Combining (1), (2), and (3), we obtain an isomorphism functorial in  $S$ :

$$F_S \times_{G_S} E_S \simeq \coprod_{F \times_G E} S = (F \times_G E)_S.$$

An object of the form  $E_S$  is called a *constant object*. There is a morphism, functorial in  $E$ ,

$$E \longrightarrow \Gamma(E_S/S),$$

which associates with each  $i \in E$  the section of  $E_S$  over  $S$  defined by the isomorphism from  $S$  onto  $S_i$ . Suppose condition  $(*)$  is satisfied for every object  $S$  of  $\mathcal{C}$ . Then  $E \rightarrow \Gamma(E_S/S)$  is a monomorphism whenever  $S \not\simeq \emptyset$ .

If  $\mathcal{C}$  is the category of schemes, then  $\Gamma(E_S/S)$  is identified with the locally constant maps from the topological space  $S$  to the set  $E$ , the preceding map associating with each element of  $E$  the corresponding constant map. It follows that  $E_S$  may equivalently be defined as representing the functor that assigns to every  $S'$  over  $S$  the set of locally constant functions from the topological space  $S'$  to  $E$ .<sup>21</sup>

## 2. Algebraic structures

Given a species of algebraic structure in the category of sets, we shall extend it to the category  $\mathcal{C}$ . We first treat the example of groups.

<sup>20</sup>Editor's note: Condition  $(*)$  is not satisfied if  $\mathcal{C}$  is the category whose arrows are  $A \rightarrow B$ ,  $\text{id}_A$ , and  $\text{id}_B$ . In that case  $B \amalg B = B$  and  $B \times_B B = B \not\simeq A$ .

<sup>21</sup>Editor's note: the diagonal morphism  $E_S \rightarrow E_S \times_S E_S = (E \times E)_S$  is also a closed immersion; that is,  $E_S$  is separated over  $S$ .

## 2.1. Group structures

We retain the notation of the preceding paragraph.

**Definition 2.1.1.** Let  $\mathbf{G} \in \text{Ob}(\widehat{\mathcal{C}})$ . A  $\widehat{\mathcal{C}}$ -group structure on  $\mathbf{G}$  is the datum, for every  $S \in \text{Ob}(\mathcal{C})$ , of a group structure on the set  $\mathbf{G}(S)$ , such that for every arrow  $f : S' \rightarrow S''$  of  $\mathcal{C}$ , the map

$$\mathbf{G}(f) : \mathbf{G}(S'') \longrightarrow \mathbf{G}(S')$$

is a group homomorphism. If  $\mathbf{G}$  and  $\mathbf{H}$  are two  $\widehat{\mathcal{C}}$ -groups, a *morphism of  $\widehat{\mathcal{C}}$ -groups* from  $\mathbf{G}$  to  $\mathbf{H}$  is a morphism  $u \in \text{Hom}(\mathbf{G}, \mathbf{H})$  such that  $u(S) : \mathbf{G}(S) \rightarrow \mathbf{H}(S)$  is a group homomorphism for every  $S \in \text{Ob}(\mathcal{C})$ .

We denote by  $\text{Hom}_{\widehat{\mathcal{C}}\text{-Gr}}(\mathbf{G}, \mathbf{H})$  the set of these morphisms and by  $(\widehat{\mathcal{C}}\text{-Gr})$  their category.

**Examples.** For  $\mathbf{E} \in \text{Ob}(\widehat{\mathcal{C}})$ , the object  $\underline{\text{Aut}}(\mathbf{E})$  has an evident  $\widehat{\mathcal{C}}$ -group structure. The final object  $\mathbf{e}$  has a unique  $\widehat{\mathcal{C}}$ -group structure, making it a final object of  $(\widehat{\mathcal{C}}\text{-Gr})$ .

For every  $S \in \text{Ob}(\mathcal{C})$ , let  $e_{\mathbf{G}}(S)$  be the identity element of  $\mathbf{G}(S)$ . The family of elements  $e_{\mathbf{G}}(S)$  defines

$$e_{\mathbf{G}} \in \Gamma(\mathbf{G}) = \text{Hom}(\mathbf{e}, \mathbf{G}).$$

It is a morphism of  $\widehat{\mathcal{C}}$ -groups  $\mathbf{e} \rightarrow \mathbf{G}$ , called the *unit section* of  $\mathbf{G}$ .

To give a  $\widehat{\mathcal{C}}$ -group structure on  $\mathbf{G}$  is equivalently to give a composition law on  $\mathbf{G}$ , that is, a  $\widehat{\mathcal{C}}$ -morphism

$$\pi_{\mathbf{G}} : \mathbf{G} \times \mathbf{G} \longrightarrow \mathbf{G}$$

such that  $\pi_{\mathbf{G}}(S)$  equips  $\mathbf{G}(S)$  with a group structure for every  $S \in \text{Ob}(\mathcal{C})$ .

Similarly,  $f : \mathbf{G} \rightarrow \mathbf{H}$  is a morphism of  $\widehat{\mathcal{C}}$ -groups if and only if the following diagram is commutative:

$$\begin{array}{ccc} \mathbf{G} \times \mathbf{G} & \xrightarrow{\pi_{\mathbf{G}}} & \mathbf{G} \\ (f, f) \downarrow & & \downarrow f \\ \mathbf{H} \times \mathbf{H} & \xrightarrow{\pi_{\mathbf{H}}} & \mathbf{H} \end{array}$$

A subobject  $\mathbf{H}$  of  $\mathbf{G}$  such that  $\mathbf{H}(S)$  is a subgroup of  $\mathbf{G}(S)$  for every  $S$  has the evident  $\widehat{\mathcal{C}}$ -group structure induced by that of  $\mathbf{G}$ . It is the unique structure for which the monomorphism  $\mathbf{H} \rightarrow \mathbf{G}$  is a morphism of  $\widehat{\mathcal{C}}$ -groups. With this structure,  $\mathbf{H}$  is called a *sub- $\widehat{\mathcal{C}}$ -group* of  $\mathbf{G}$ .

If  $\mathbf{G}$  and  $\mathbf{H}$  are two  $\widehat{\mathcal{C}}$ -groups, then  $\mathbf{G} \times \mathbf{H}$  has the evident product structure: for every  $S$ , the set  $\mathbf{G}(S) \times \mathbf{H}(S)$  is given the product of the prescribed group structures. The resulting object is called the *product  $\widehat{\mathcal{C}}$ -group* of  $\mathbf{G}$  and  $\mathbf{H}$ ; it is their product in the category of  $\widehat{\mathcal{C}}$ -groups.

If  $\mathbf{G}$  is a  $\widehat{\mathcal{C}}$ -group, then  $\mathbf{G}_S$  is a  $\widehat{\mathcal{C}_{/S}}$ -group for every  $S \in \text{Ob}(\mathcal{C})$ . For two  $\widehat{\mathcal{C}}$ -groups  $\mathbf{G}$  and  $\mathbf{H}$ , define the object  $\underline{\text{Hom}}_{\widehat{\mathcal{C}}\text{-Gr}}(\mathbf{G}, \mathbf{H})$  of  $\widehat{\mathcal{C}}$  by

$$\underline{\text{Hom}}_{\widehat{\mathcal{C}}\text{-Gr}}(\mathbf{G}, \mathbf{H})(S) = \text{Hom}_{\widehat{\mathcal{C}_{/S}}\text{-Gr}}(\mathbf{G}_S, \mathbf{H}_S).$$

Notice that  $\underline{\text{Hom}}_{\widehat{\mathcal{C}}\text{-Gr}}$  is not in general a  $\widehat{\mathcal{C}}$ -group, still less the internal  $\underline{\text{Hom}}$  object in the category  $(\widehat{\mathcal{C}}\text{-Gr})$ .

One similarly defines

$$\underline{\text{Isom}}_{\widehat{\mathcal{C}}\text{-Gr}}(\mathbf{G}, \mathbf{H}), \quad \underline{\text{End}}_{\widehat{\mathcal{C}}\text{-Gr}}(\mathbf{G}), \quad \underline{\text{Aut}}_{\widehat{\mathcal{C}}\text{-Gr}}(\mathbf{G}).$$

**Definition 2.1.2.** Let  $G \in \text{Ob}(\mathcal{C})$ . A  $\mathcal{C}$ -group structure on  $G$  is a  $\widehat{\mathcal{C}}$ -group structure on  $\mathbf{h}_G \in \text{Ob}(\widehat{\mathcal{C}})$ . A morphism from the  $\mathcal{C}$ -group  $G$  to the  $\mathcal{C}$ -group  $H$  is an element

$$u \in \text{Hom}(G, H) \simeq \text{Hom}(\mathbf{h}_G, \mathbf{h}_H)$$

that defines a morphism of  $\widehat{\mathcal{C}}$ -groups from  $\mathbf{h}_G$  to  $\mathbf{h}_H$ .

We denote the category of  $\mathcal{C}$ -groups by  $(\mathcal{C}\text{-Gr})$ . There is a cartesian square in **Cat**:

$$\begin{array}{ccc} (\mathcal{C}\text{-Gr}) & \longrightarrow & (\widehat{\mathcal{C}}\text{-Gr}) \\ \downarrow & & \downarrow \\ \mathcal{C} & \xrightarrow{\mathbf{h}} & \widehat{\mathcal{C}} \end{array} .$$

All the preceding definitions and constructions therefore carry over at once to  $(\mathcal{C}\text{-Gr})$  whenever the functors they use (products, internal  $\underline{\text{Hom}}$  objects, and so on) are representable. They also apply to the categories  $\mathcal{C}/_S$ . In that case we write  $\underline{\text{Hom}}_{S\text{-Gr}}$  for  $\underline{\text{Hom}}_{\mathcal{C}/_S\text{-Gr}}$ , and similarly for the other objects.

## 2.2.

More generally, suppose  $(T)$  is a species of structure on  $n$  base sets defined by finite inverse limits—for example, by commutative diagrams constructed from cartesian products, as in the structures of a monoid, a group, a set with operators, a module over a ring, or a Lie algebra over a ring. The preceding construction defines the notion of a “structure of species  $(T)$ ” on  $n$  objects  $\mathbf{F}_1, \dots, \mathbf{F}_n$  of  $\widehat{\mathcal{C}}$ . Such a structure consists, for every  $S \in \text{Ob}(\mathcal{C})$ , of a structure of species  $(T)$  on the sets  $\mathbf{F}_1(S), \dots, \mathbf{F}_n(S)$ , in such a way that for every arrow  $S' \rightarrow S''$  of  $\mathcal{C}$ , the family of maps

$$(\mathbf{F}_i(S''))_i \longrightarrow (\mathbf{F}_i(S'))_i$$

is a polyhomomorphism for the species  $(T)$ .

Morphisms of structures of species  $(T)$  are defined similarly, giving a category

$$(\widehat{\mathcal{C}} \times \dots \times \widehat{\mathcal{C}})^{(T)}.$$

The fully faithful functor  $\mathbf{h} \times \dots \times \mathbf{h}$  then defines, by inverse image, the category

$$(\mathcal{C} \times \dots \times \mathcal{C})^{(T)}.$$

Since it commutes with inverse limits, all the functorial properties, notions, and notation introduced in  $\widehat{\mathcal{C}}$  carry over to this category.

Now suppose fiber products exist in  $\mathcal{C}$ , and let  $(T)$  be a species of algebraic structure specified by certain morphisms between cartesian products, subject to axioms consisting of commutative diagrams formed with those morphisms. A structure of species  $(T)$  on a family of objects of  $\mathcal{C}$  is then defined by corresponding morphisms between cartesian products satisfying the corresponding commutativity conditions. Consequently, if  $\mathcal{C}$  and  $\mathcal{C}'$

are categories with products and  $f : \mathcal{C} \rightarrow \mathcal{C}'$  is a functor commuting with products, then a structure of species  $(T)$  on a family  $(F_i)$  of objects of  $\mathcal{C}$  induces one on the family  $(f(F_i))$  in  $\mathcal{C}'$ . Thus every  $\mathcal{C}$ -group becomes a  $\mathcal{C}'$ -group, every pair consisting of a  $\mathcal{C}$ -ring and a module over it becomes the analogous pair in  $\mathcal{C}'$ , and so on.

In particular, let  $\mathcal{C}$  satisfy the conditions of 1.8.<sup>22</sup> The functor  $E \mapsto E_S$  defined there commutes with finite inverse limits. It therefore carries groups to  $S$ -groups (that is,  $\mathcal{C}/S$ -groups), rings to  $S$ -rings, and so forth.

**Remark.** It is worth observing that the preceding construction, when applied to  $\widehat{\mathcal{C}}$ , gives back precisely the notions already defined there. In other words, to give an object of  $\widehat{\mathcal{C}}$  a structure of species  $(T)$  when it is viewed as a functor on  $\mathcal{C}$  is the same as to give the representable functor on  $\widehat{\mathcal{C}}$  defined by that object a structure of species  $(T)$ .<sup>23</sup>

We shall treat two further special cases of this construction: structures with operator groups and modules.

### 2.3. Structures with operator groups

**Definition 2.3.1.** Let  $\mathbf{E} \in \text{Ob}(\widehat{\mathcal{C}})$  and  $\mathbf{G} \in \text{Ob}(\widehat{\mathcal{C}}\text{-Gr})$ . A structure on  $\mathbf{E}$  of an object with  $\widehat{\mathcal{C}}$ -operator group  $\mathbf{G}$  (or of a  $\mathbf{G}$ -object) consists in giving  $\mathbf{E}(S)$ , for every  $S \in \text{Ob}(\mathcal{C})$ , the structure of a set with operator group  $\mathbf{G}(S)$ , in such a way that, for every arrow  $S' \rightarrow S''$  in  $\mathcal{C}$ , the map of sets  $\mathbf{E}(S'') \rightarrow \mathbf{E}(S')$  is compatible with the homomorphism of operator groups  $\mathbf{G}(S'') \rightarrow \mathbf{G}(S')$ .

As usual, this is equivalent to giving a morphism

$$\mu : \mathbf{G} \times \mathbf{E} \longrightarrow \mathbf{E}$$

which, for every  $S$ , equips  $\mathbf{E}(S)$  with the structure of a set on which  $\mathbf{G}(S)$  acts. But

$$\text{Hom}(\mathbf{G} \times \mathbf{E}, \mathbf{E}) \simeq \text{Hom}(\mathbf{G}, \underline{\text{End}}(\mathbf{E})),$$

so  $\mu$  defines a morphism  $\mathbf{G} \rightarrow \underline{\text{End}}(\mathbf{E})$ . It is immediate that this morphism maps  $\mathbf{G}$  into  $\underline{\text{Aut}}(\mathbf{E})$  and is a morphism of  $\widehat{\mathcal{C}}$ -groups. Consequently, giving  $\mathbf{E}$  the structure of an object with  $\widehat{\mathcal{C}}$ -operator group  $\mathbf{G}$  is equivalent to giving a morphism of  $\widehat{\mathcal{C}}$ -groups

$$\rho : \mathbf{G} \longrightarrow \underline{\text{Aut}}(\mathbf{E}).$$

In particular, every element  $g \in \mathbf{G}(S)$  defines an automorphism  $\rho(g)$  of the functor  $\mathbf{E}_S$ , that is, an automorphism of  $\mathbf{E} \times \mathbf{h}_S$  commuting with the projection  $\mathbf{E} \times \mathbf{h}_S \rightarrow \mathbf{h}_S$ , and hence, in particular, an automorphism of the set  $\mathbf{E}(S')$  for every  $S' \rightarrow S$ .

**Definition 2.3.2.** We denote by  $\mathbf{E}^{\mathbf{G}}$  the subobject of  $\mathbf{E}$  defined as follows:

$$\mathbf{E}^{\mathbf{G}}(S) = \{x \in \mathbf{E}(S) \mid x_{S'} \text{ is invariant under } \mathbf{G}(S') \text{ for every } S' \rightarrow S\},$$

where  $x_{S'}$  denotes the image of  $x$  under  $\mathbf{E}(S) \rightarrow \mathbf{E}(S')$ .

<sup>22</sup>Editor's note: including condition (\*).

<sup>23</sup>Editor's note: for group structures, for example, let  $\mathbf{G} \in \text{Ob}(\widehat{\mathcal{C}})$ . If the functor

$$\widehat{\mathcal{C}} \longrightarrow \mathbf{Set}, \quad \mathbf{F} \longmapsto \text{Hom}_{\widehat{\mathcal{C}}}(\mathbf{F}, \mathbf{G})$$

has a group structure, then so does its restriction to  $\mathcal{C}$ , namely  $X \mapsto \mathbf{G}(X)$ . Conversely, if  $\mathbf{G}$  is a  $\widehat{\mathcal{C}}$ -group, its multiplication morphism  $\pi_{\mathbf{G}} : \mathbf{G} \times \mathbf{G} \rightarrow \mathbf{G}$  induces, for every  $\mathbf{F} \in \text{Ob}(\widehat{\mathcal{C}})$ , a group structure on  $\text{Hom}_{\widehat{\mathcal{C}}}(\mathbf{F}, \mathbf{G})$ , functorially in  $\mathbf{F}$ .

Thus  $\mathbf{E}^{\mathbf{G}}$ , the “subobject of invariants of  $\mathbf{G}$ ,” is the largest subobject of  $\mathbf{E}$  on which  $\mathbf{G}$  acts trivially.

**Definition 2.3.3.** Let  $\mathbf{F}$  be a subobject of  $\mathbf{E}$ . We denote by  $\underline{\text{Norm}}_{\mathbf{G}}\mathbf{F}$  and  $\underline{\text{Centr}}_{\mathbf{G}}\mathbf{F}$  the sub- $\mathcal{C}$ -groups of  $\mathbf{G}$  defined by

$$\begin{aligned} (\underline{\text{Norm}}_{\mathbf{G}}\mathbf{F})(S) &= \{g \in \mathbf{G}(S) \mid \rho(g)\mathbf{F}_S = \mathbf{F}_S\} \\ &= \{g \in \mathbf{G}(S) \mid \rho(g)\mathbf{F}(S') = \mathbf{F}(S') \text{ for every } S' \rightarrow S\}, \\ (\underline{\text{Centr}}_{\mathbf{G}}\mathbf{F})(S) &= \{g \in \mathbf{G}(S) \mid \rho(g)|_{\mathbf{F}_S} = \text{id}\} \\ &= \{g \in \mathbf{G}(S) \mid \rho(g)|_{\mathbf{F}(S')} = \text{id} \text{ for every } S' \rightarrow S\}. \end{aligned}$$

Here the vertical bar after  $\rho(g)$  denotes restriction.<sup>24</sup>

**Scholium 2.3.3.1.**<sup>25</sup> In particular, let  $x \in \Gamma(\mathbf{E})$ , that is (cf. 1.2), a collection of elements  $x_S \in \mathbf{E}(S)$ ,  $S \in \text{Ob}(\mathcal{C})$ , such that for every arrow  $f : S' \rightarrow S$  one has  $\mathbf{E}(f)(x_S) = x_{S'}$ . If  $\mathcal{C}$  has a final object  $S_0$ , then  $\Gamma(\mathbf{E}) = \mathbf{E}(S_0)$ . The element  $x$  then defines a subfunctor of  $\mathbf{E}$ , denoted by  $\mathbf{x}$ , and

$$\underline{\text{Norm}}_{\mathbf{G}}\mathbf{x} = \underline{\text{Centr}}_{\mathbf{G}}\mathbf{x}.$$

We denote this functor by  $\underline{\text{Stab}}_{\mathbf{G}}(x)$  and call it the *stabilizer* of  $x$ ; thus, for every  $S \in \text{Ob}(\mathcal{C})$ ,

$$\underline{\text{Stab}}_{\mathbf{G}}(x)(S) = \{g \in \mathbf{G}(S) \mid \rho(g)x_S = x_S\}.$$

Suppose that fiber products exist in  $\mathcal{C}$ . If  $\mathbf{G} = \mathbf{h}_G$  (respectively,  $\mathbf{E} = \mathbf{h}_E$ ), where  $G$  is a  $\mathcal{C}$ -group (respectively,  $E \in \text{Ob}(\mathcal{C})$ ), and if  $\mathcal{C}$  has a final object  $S_0$ , so that  $x$  is a morphism  $S_0 \rightarrow E$ , then  $\underline{\text{Stab}}_{\mathbf{G}}(x)$  is represented by the fiber product  $G \times_E S_0$ , where the morphism  $G \rightarrow E$  is the composite of

$$\text{id}_G \times x : G = G \times S_0 \longrightarrow G \times E$$

and

$$\mu : G \times E \longrightarrow E.$$

**Remark 2.3.3.2.**<sup>25</sup> The formation of  $\mathbf{E}^{\mathbf{G}}$ ,  $\underline{\text{Norm}}_{\mathbf{G}}\mathbf{F}$ , and  $\underline{\text{Centr}}_{\mathbf{G}}\mathbf{F}$  commutes with base change; that is, for every  $S \in \text{Ob}(\mathcal{C})$ ,

$$(\mathbf{E}^{\mathbf{G}})_S = (\mathbf{E}_S)^{\mathbf{G}_S}, \quad (\underline{\text{Norm}}_{\mathbf{G}}\mathbf{F})_S \simeq \underline{\text{Norm}}_{\mathbf{G}_S}\mathbf{F}_S, \quad (\underline{\text{Centr}}_{\mathbf{G}}\mathbf{F})_S \simeq \underline{\text{Centr}}_{\mathbf{G}_S}\mathbf{F}_S.$$

**Definition 2.3.4.** If  $G$  is a  $\mathcal{C}$ -group and  $\mathbf{E}$  is an object of  $\widehat{\mathcal{C}}$  (respectively,  $E$  is an object of  $\mathcal{C}$ ), a  $G$ -object structure on  $\mathbf{E}$  (respectively, on  $E$ ) is an  $\mathbf{h}_G$ -object structure on  $\mathbf{E}$  (respectively, on  $\mathbf{h}_E$ ).

In view of this definition, all the notions and notation defined above carry over to  $\mathcal{C}$  whenever they involve only representable functors. For example, if  $\underline{\text{Norm}}_{\mathbf{h}_G}(\mathbf{h}_F)$  is representable, then there is a unique subobject of  $G$  representing it; it is then a sub- $\mathcal{C}$ -group of  $G$ , and is denoted by  $\text{Norm}_G(F)$ , and so forth.

**Definition 2.3.5.**

<sup>24</sup>Editor’s note: the original has been corrected by replacing the inclusion  $\rho(g)\mathbf{F}_S \subset \mathbf{F}_S$  with an equality, so as to ensure that  $\underline{\text{Norm}}_{\mathbf{G}}(\mathbf{F})$  is indeed a group. See VI<sub>B</sub>, 6.4 for conditions under which the “transporter” coincides with the “strict transporter.”

<sup>25</sup>Editor’s note: Scholium 2.3.3.1 and Remark 2.3.3.2 have been added.

- (a) We say that the  $\widehat{\mathcal{C}}$ -group  $\mathbf{G}$  acts on the  $\widehat{\mathcal{C}}$ -group  $\mathbf{H}$  if  $\mathbf{H}$  is equipped with a  $\mathbf{G}$ -object structure such that, for every  $g \in \mathbf{G}(S)$ , the automorphism of  $\mathbf{H}(S)$  defined by  $g$  is a group automorphism.

Equivalently, for every  $g \in \mathbf{G}(S)$ , the automorphism  $\rho(g)$  of  $\mathbf{H}_S$  is an automorphism of  $\widehat{\mathcal{C}}_S$ -groups; or equivalently, the morphism of  $\widehat{\mathcal{C}}$ -groups  $\mathbf{G} \rightarrow \underline{\text{Aut}}(\mathbf{H})$  maps  $\mathbf{G}$  into

$$\underline{\text{Aut}}_{\widehat{\mathcal{C}}\text{-Gr}}(\mathbf{H}).$$

- (b) In the situation above, there is a unique  $\widehat{\mathcal{C}}$ -group structure on  $\mathbf{H} \times \mathbf{G}$  such that, for every  $S$ ,  $(\mathbf{H} \times \mathbf{G})(S)$  is the semidirect product of the groups  $\mathbf{H}(S)$  and  $\mathbf{G}(S)$  for the given action of  $\mathbf{G}(S)$  on  $\mathbf{H}(S)$ . We denote this  $\widehat{\mathcal{C}}$ -group by  $\mathbf{H} \cdot \mathbf{G}$  and call it the *semidirect product of  $\mathbf{H}$  by  $\mathbf{G}$* . Thus, by definition,

$$(\mathbf{H} \cdot \mathbf{G})(S) = \mathbf{H}(S) \cdot \mathbf{G}(S).$$

Let  $\mathbf{G}$  be a  $\widehat{\mathcal{C}}$ -group. For every arrow  $S' \rightarrow S$  in  $\mathcal{C}$  and every  $g \in \mathbf{G}(S)$ , let  $\text{Int}(g)$  be the automorphism of  $\mathbf{G}(S')$  defined by

$$\text{Int}(g)h = ghg^{-1}.$$

This definition extends to a morphism of  $\widehat{\mathcal{C}}$ -groups

$$\text{Int} : \mathbf{G} \longrightarrow \underline{\text{Aut}}_{\widehat{\mathcal{C}}\text{-Gr}}(\mathbf{G}) \subset \underline{\text{Aut}}(\mathbf{G}).$$

Definition 2.3.3 therefore applies, and for every subobject  $\mathbf{E}$  of  $\mathbf{G}$  one has the sub- $\widehat{\mathcal{C}}$ -groups of  $\mathbf{G}$

$$\underline{\text{Norm}}_{\mathbf{G}}(\mathbf{E}) \quad \text{and} \quad \underline{\text{Centr}}_{\mathbf{G}}(\mathbf{E}).$$

**Definition 2.3.6.** The *center* of  $\mathbf{G}$ , denoted  $\underline{\text{Centr}}(\mathbf{G})$ , is the sub- $\widehat{\mathcal{C}}$ -group  $\underline{\text{Centr}}_{\mathbf{G}}(\mathbf{G})$  of  $\mathbf{G}$ . We say that  $\mathbf{G}$  is commutative if  $\underline{\text{Centr}}(\mathbf{G}) = \mathbf{G}$ , or, equivalently, if  $\mathbf{G}(S)$  is commutative for every  $S$ .

We say that the sub- $\widehat{\mathcal{C}}$ -group  $\mathbf{H}$  of  $\mathbf{G}$  is invariant in  $\mathbf{G}$  if  $\underline{\text{Norm}}_{\mathbf{G}}(\mathbf{H}) = \mathbf{G}$ , or, equivalently, if  $\mathbf{H}(S)$  is invariant in  $\mathbf{G}(S)$  for every  $S$ .<sup>26</sup>

**Definition 2.3.6.1.**<sup>27</sup> Let  $f : \mathbf{G} \rightarrow \mathbf{G}'$  be a morphism of  $\widehat{\mathcal{C}}$ -groups. The *kernel* of  $f$ , denoted  $\text{Ker } f$ , is the sub- $\widehat{\mathcal{C}}$ -group of  $\mathbf{G}$  defined by

$$(\text{Ker } f)(S) = \{x \in \mathbf{G}(S) \mid f(S)(x) = 1\} = \text{Ker } f(S)$$

for every  $S \in \text{Ob}(\mathcal{C})$ ; it is an invariant sub- $\widehat{\mathcal{C}}$ -group.

Notice that if  $\mathbf{G} = \mathbf{h}_G$  and  $\mathbf{G}' = \mathbf{h}_{G'}$ , if  $\mathcal{C}$  has a final object  $S_0$ , and if fiber products exist in  $\mathcal{C}$ , then  $\text{Ker}(f)$  is represented by  $S_0 \times_{G'} G$ .

**Definition 2.3.6.2.**<sup>27</sup> Let  $\mathbf{E} \in \widehat{\mathcal{C}}$ , and let  $\mathbf{G}$  be a  $\widehat{\mathcal{C}}$ -group acting on  $\mathbf{E}$ . We say that the action of  $\mathbf{G}$  on  $\mathbf{E}$  is *faithful* if the kernel of the morphism  $\mathbf{G} \rightarrow \underline{\text{Aut}}(\mathbf{E})$  is trivial; that is, if, for every  $S \in \text{Ob}(\mathcal{C})$  and  $g \in \mathbf{G}(S)$ , the condition

$$g_{S'} \cdot x = x \quad \text{for every } S' \rightarrow S \text{ and every } x \in \mathbf{E}(S')$$

implies  $g = 1$ .

<sup>26</sup>Editor's note: in addition,  $\mathbf{H}$  is said to be *central* in  $\mathbf{G}$  if  $\underline{\text{Centr}}_{\mathbf{G}}(\mathbf{H}) = \mathbf{G}$ , or, equivalently, if  $\mathbf{H}(S)$  is central in  $\mathbf{G}(S)$  for every  $S$ .

<sup>27</sup>Editor's note: Definitions 2.3.6.1 and 2.3.6.2 have been added.

Many definitions and propositions from elementary group theory carry over readily. We record only the following one, which will be useful to us.

**Proposition 2.3.7.** *Let  $f : \mathbf{W} \rightarrow \mathbf{G}$  be a morphism of  $\widehat{\mathcal{C}}$ -groups. Put  $\mathbf{H}(S) = \text{Ker } f(S)$ . Let  $u : \mathbf{G} \rightarrow \mathbf{W}$  be a morphism of  $\widehat{\mathcal{C}}$ -groups that is a section of  $f$  (and is therefore necessarily a monomorphism). Then  $\mathbf{W}$  is identified with the semidirect product of  $\mathbf{H}$  by  $\mathbf{G}$  for the action of  $\mathbf{G}$  on  $\mathbf{H}$  defined by*

$$(g, h) \mapsto \text{Int}(u(g))h$$

for  $g \in \mathbf{G}(S)$ ,  $h \in \mathbf{H}(S)$ , and  $S \in \text{Ob}(\mathcal{C})$ .

All these definitions and propositions carry over, as usual, to  $\mathcal{C}$ . In particular, the semidirect product of two  $\mathcal{C}$ -groups  $H$  and  $G$ , with  $G$  acting on  $H$ , is defined whenever the Cartesian product  $H \times G$  exists, and the analog of Proposition 2.3.7 takes the following form.

**Proposition 2.3.8.** *Let*

$$H \xrightarrow{i} W \xrightarrow{f} G$$

*be a sequence of morphisms of  $\mathcal{C}$ -groups such that, for every  $S \in \text{Ob}(\mathcal{C})$ , the pair  $(H(S), i(S))$  is a kernel of  $f(S) : W(S) \rightarrow G(S)$ . Let  $u : G \rightarrow W$  be a morphism of  $\mathcal{C}$ -groups that is a section of  $f$ . Then  $W$  is identified with the semidirect product of  $H$  by  $G$  for the action of  $G$  on  $H$  such that, if  $S \in \text{Ob}(\mathcal{C})$ ,  $g \in G(S)$ , and  $h \in H(S)$ , then*

$$\text{Int}(u(g))i(h) = i({}^gh).$$

### 3. The category of $\mathbf{O}$ -modules, the category of $\mathbf{G}$ - $\mathbf{O}$ -modules

**Definition 3.1.** Let  $\mathbf{O}$  and  $\mathbf{F}$  be two objects of  $\widehat{\mathcal{C}}$ . We say that  $\mathbf{F}$  is a  $\widehat{\mathcal{C}}$ -module over the  $\widehat{\mathcal{C}}$ -ring  $\mathbf{O}$ , or, for short, an  $\mathbf{O}$ -module, if, for every  $S \in \text{Ob}(\mathcal{C})$ , the set  $\mathbf{O}(S)$  has been endowed with a ring structure and  $\mathbf{F}(S)$  with a module structure over that ring, in such a way that for every arrow  $S' \rightarrow S''$  in  $\mathcal{C}$ , the map  $\mathbf{O}(S'') \rightarrow \mathbf{O}(S')$  is a ring homomorphism and  $\mathbf{F}(S'') \rightarrow \mathbf{F}(S')$  is a homomorphism of abelian groups compatible with that ring homomorphism.

Once  $\mathbf{O}$  is fixed, one defines in the usual way a morphism between two  $\mathbf{O}$ -modules  $\mathbf{F}$  and  $\mathbf{F}'$ , hence the abelian group  $\text{Hom}_{\mathbf{O}}(\mathbf{F}, \mathbf{F}')$ , and the category of  $\mathbf{O}$ -modules, denoted  $(\mathbf{O}\text{-Mod.})$ .

**Lemma 3.1.1.**<sup>28</sup> *The category  $(\mathbf{O}\text{-Mod.})$  has the structure of an abelian category, defined objectwise. Moreover,  $(\mathbf{O}\text{-Mod.})$  satisfies axiom (AB 5) (see [Gr57, 1.5]); that is, arbitrary direct sums exist, and if  $\mathbf{M}$  is an  $\mathbf{O}$ -module,  $\mathbf{N}$  a submodule, and  $(\mathbf{F}_i)_{i \in I}$  an increasing filtered family of submodules of  $\mathbf{M}$ , then*

$$\bigcup_{i \in I} (\mathbf{F}_i \cap \mathbf{N}) = \left( \bigcup_{i \in I} \mathbf{F}_i \right) \cap \mathbf{N}.$$

Indeed, let  $f : \mathbf{F} \rightarrow \mathbf{F}'$  be a morphism of  $\mathbf{O}$ -modules. Define the  $\mathbf{O}$ -modules  $\text{Ker } f$  (respectively,  $\text{Im } f$  and  $\text{Coker } f$ ) by setting, for every  $S \in \text{Ob}(\mathcal{C})$ ,

$$(\text{Ker } f)(S) = \text{Ker } f(S)$$

(respectively, the analogous formulas). Then  $\text{Ker } f$  (respectively,  $\text{Coker } f$ ) is a kernel (respectively, a cokernel) of  $f$ , and there is an isomorphism of  $\mathbf{O}$ -modules

$$\mathbf{F}/\text{Ker } f \xrightarrow{\sim} \text{Im } f.$$

This proves that  $(\mathbf{O}\text{-Mod.})$  is an abelian category.

Arbitrary direct sums exist and are defined objectwise. Finally, if  $\mathbf{M}$  is an  $\mathbf{O}$ -module,  $\mathbf{N}$  a submodule, and  $(\mathbf{F}_i)_{i \in I}$  an increasing filtered family of submodules of  $\mathbf{M}$ , then the inclusion

$$\bigcup_{i \in I} (\mathbf{F}_i \cap \mathbf{N}) \subset \left( \bigcup_{i \in I} \mathbf{F}_i \right) \cap \mathbf{N}$$

is an equality. Indeed, if  $S \in \text{Ob}(\mathcal{C})$  and

$$x \in \mathbf{N}(S) \cap \bigcup_i \mathbf{F}_i(S),$$

there is an  $i \in I$  such that  $x \in \mathbf{N}(S) \cap \mathbf{F}_i(S)$ .

**Proposition 3.1.2.**<sup>28</sup> *Suppose that the category  $\mathcal{C}$  is small, that is, that  $\text{Ob}(\mathcal{C})$  is a set. Then  $(\mathbf{O}\text{-Mod.})$  has a generator, and consequently has enough injective objects.*

*Proof.* For every object  $S$  of  $\mathcal{C}$ , define the  $\mathbf{O}$ -module  $\mathbf{O}_S$  as follows. For every object  $S'$ ,  $\mathbf{O}_S(S')$  is a direct sum of copies of  $\mathbf{O}(S')$  indexed by  $\text{Hom}(S', S)$ , that is,

$$\mathbf{O}_S(S') = \bigoplus_{g: S' \rightarrow S} \mathbf{O}(S')g.$$

For every morphism  $f: S'' \rightarrow S'$ , writing  $f^*$  for the ring homomorphism  $\mathbf{O}(S') \rightarrow \mathbf{O}(S'')$ , the morphism

$$\mathbf{O}_S(f): \mathbf{O}_S(S') \longrightarrow \mathbf{O}_S(S'')$$

sends each element  $ag$  of  $\mathbf{O}_S(S')$ , where  $a \in \mathbf{O}(S')$  and  $g: S' \rightarrow S$ , to the element  $f^*(a)(g \circ f)$  of  $\mathbf{O}_S(S'')$ . It is readily checked that this defines a functor in  $\mathbf{O}$ -modules.

Since  $\text{Ob}(\mathcal{C})$  is a set by hypothesis, we may form the direct sum

$$\mathbf{G} = \bigoplus_{S \in \text{Ob}(\mathcal{C})} \mathbf{O}_S.$$

**Lemma 3.1.2.1.** *Let  $\mathbf{F}$  be an  $\mathbf{O}$ -module. For every  $S \in \text{Ob}(\mathcal{C})$ , every element  $x \in \mathbf{F}(S)$  defines a unique morphism of  $\mathbf{O}$ -modules*

$$\mathbf{x}: \mathbf{O}_S \longrightarrow \mathbf{F}$$

such that  $\mathbf{x}(1 \text{ id}_S) = x$ .

*Proof.* Let  $\phi: \mathbf{O}_S \rightarrow \mathbf{F}$  be a morphism of  $\mathbf{O}$ -modules such that  $\phi(1 \text{ id}_S) = x$ . Let  $g: S' \rightarrow S$  and  $a \in \mathbf{O}(S')$ . The  $\mathbf{O}(S')$ -linearity and the commutativity of the diagram

$$\begin{array}{ccc} \mathbf{O}_S(S) & \xrightarrow{\psi} & \mathbf{F}(S) \\ \mathbf{O}_S(g) \downarrow & & \downarrow \mathbf{F}(g) \\ \mathbf{O}_S(S') & \xrightarrow{\phi} & \mathbf{F}(S') \end{array}$$

<sup>28</sup>Editor's note: Statements 3.1.1 and 3.1.2 have been added to show that the category  $(\mathbf{O}\text{-Mod.})$  is abelian and satisfies axiom (AB 5), and moreover has enough injective objects when the category  $\mathcal{C}$  is small. An error in 3.1.2, pointed out in 2016 by Linyuan Liu, is corrected here.



imply that

$$\phi(ag) = a\phi(1g) = a\mathbf{F}(g)(\phi(1 \text{id}_S)) = a\mathbf{F}(g)(x).$$

Thus  $\phi$ , if it exists, is determined by the preceding equality. Conversely, if  $\phi$  is defined in this way, then for every  $f : S'' \rightarrow S'$  one has

$$\begin{aligned} \phi \circ \mathbf{O}_S(f)(ag) &= \phi(f^*(a)(g \circ f)) \\ &= f^*(a)\mathbf{F}(g \circ f)(x) \\ &= \mathbf{F}(f)(a\mathbf{F}(g)(x)) \\ &= \mathbf{F}(f) \circ \phi(ag), \end{aligned}$$

that is, the diagram

$$\begin{array}{ccc} \mathbf{O}_S(S') & \xrightarrow{\phi} & \mathbf{F}(S') \\ \mathbf{O}_S(f) \downarrow & & \downarrow \mathbf{F}(f) \\ \mathbf{O}_S(S'') & \xrightarrow{\phi} & \mathbf{F}(S'') \end{array}$$

is commutative. This proves the lemma.

Now let  $\mathbf{F}$  be an  $\mathbf{O}$ -module. For each  $S \in \text{Ob}(\mathcal{C})$ , let  $F_0(S)$  be a set of generators of the  $\mathbf{O}(S)$ -module  $\mathbf{F}(S)$ , and let  $\mathbf{L}_S$  be the direct sum, indexed by  $F_0(S)$ , of copies of  $\mathbf{O}_S$ . By the lemma, we obtain a morphism of  $\mathbf{O}$ -modules

$$\mathbf{L}_S \longrightarrow \mathbf{F}$$

such that the morphism  $\mathbf{L}_S(S) \rightarrow \mathbf{F}(S)$  is surjective, and hence is an epimorphism in the category of  $\mathbf{O}(S)$ -modules.

Set

$$I = \bigsqcup_{S \in \text{Ob}(\mathcal{C})} F_0(S).$$

We then obtain a morphism of  $\mathbf{O}$ -modules

$$\mathbf{G}^{\oplus I} \longrightarrow \bigoplus_{S \in \text{Ob}(\mathcal{C})} \mathbf{L}_S \longrightarrow \mathbf{F}$$

which is an epimorphism objectwise, and hence an epimorphism of  $(\mathbf{O}\text{-Mod.})$ . This shows that  $\mathbf{G}$  is a generator of  $(\mathbf{O}\text{-Mod.})$  (see [Gr57, 1.9.1]). Since  $(\mathbf{O}\text{-Mod.})$  satisfies (AB 5), it follows from [Gr57, 1.10.1] that  $(\mathbf{O}\text{-Mod.})$  has enough injective objects.

**Remark 3.1.3.** If  $\mathbf{O}_0$  is the  $\widehat{\mathcal{C}}$ -ring defined by

$$\mathbf{O}_0(S) = \mathbb{Z}$$

(which must not be confused with the functor associated with the constant object  $\mathbb{Z}$ ), then the category of  $\mathbf{O}_0$ -modules is isomorphic to the category of commutative  $\widehat{\mathcal{C}}$ -groups.

**Definition 3.1.4.** Notice that if  $\mathbf{F}$  is an  $\mathbf{O}$ -module, then for every  $S \in \text{Ob}(\mathcal{C})$ ,  $\mathbf{F}_S$  is an  $\mathbf{O}_S$ -module. We may therefore define an abelian  $\widehat{\mathcal{C}}$ -group  $\underline{\text{Hom}}_{\mathbf{O}}(\mathbf{F}, \mathbf{F}')$  by

$$\underline{\text{Hom}}_{\mathbf{O}}(\mathbf{F}, \mathbf{F}')(S) = \text{Hom}_{\mathbf{O}_S}(\mathbf{F}_S, \mathbf{F}'_S).$$

In the same way one defines the objects

$$\underline{\text{Isom}}_{\mathbf{O}}(\mathbf{F}, \mathbf{F}'), \quad \underline{\text{End}}_{\mathbf{O}}(\mathbf{F}), \quad \text{and} \quad \underline{\text{Aut}}_{\mathbf{O}}(\mathbf{F}),$$

the last of which is a  $\widehat{\mathcal{C}}$ -group under the structure induced by composition of automorphisms.

**Definition 3.2.** Let  $\mathbf{O}$  be a  $\widehat{\mathcal{C}}$ -ring,  $\mathbf{F}$  an  $\mathbf{O}$ -module, and  $\mathbf{G}$  a  $\widehat{\mathcal{C}}$ -group. A  $\mathbf{G}$ - $\mathbf{O}$ -module structure on  $\mathbf{F}$  is a  $\mathbf{G}$ -object structure such that, for every  $S \in \text{Ob}(\mathcal{C})$  and every  $g \in \mathbf{G}(S)$ , the automorphism of  $\mathbf{F}(S)$  defined by  $g$  is an automorphism of its  $\mathbf{O}(S)$ -module structure.

Equivalently, the morphism of  $\widehat{\mathcal{C}}$ -groups

$$\rho : \mathbf{G} \longrightarrow \underline{\text{Aut}}(\mathbf{F})$$

defined in 2.3 maps  $\mathbf{G}$  into the  $\widehat{\mathcal{C}}$ -subgroup  $\underline{\text{Aut}}_{\mathbf{O}}(\mathbf{F})$  of  $\underline{\text{Aut}}(\mathbf{F})$ . Thus, giving a  $\mathbf{G}$ - $\mathbf{O}$ -module structure on the  $\mathbf{O}$ -module  $\mathbf{F}$  is the same as giving a morphism of  $\widehat{\mathcal{C}}$ -groups

$$\rho : \mathbf{G} \longrightarrow \underline{\text{Aut}}_{\mathbf{O}}(\mathbf{F}).$$

One defines naturally the abelian group  $\text{Hom}_{\mathbf{G-O}}(\mathbf{F}, \mathbf{F}')$ , and hence the additive category of  $\mathbf{G}$ - $\mathbf{O}$ -modules, denoted  $(\mathbf{G-O-Mod.})$ .

**Remark 3.2.1.** The reader will notice that  $(\mathbf{G-O-Mod.})$  may also be defined as the category of  $\mathbf{O}[\mathbf{G}]$ -modules, where  $\mathbf{O}[\mathbf{G}]$  is the algebra of the  $\widehat{\mathcal{C}}$ -group  $\mathbf{G}$  over the  $\widehat{\mathcal{C}}$ -ring  $\mathbf{O}$ , whose definition is clear.<sup>29</sup> Consequently, by 3.1.1 and 3.1.2,  $(\mathbf{G-O-Mod.})$  is an abelian category satisfying axiom (AB 5); moreover, if  $\mathcal{C}$  is small,  $(\mathbf{G-O-Mod.})$  has enough injective objects.

All the constructions of this section specialize immediately to the case where  $\mathbf{G}$  (or  $\mathbf{O}$ , or both) is representable by an object of  $\mathcal{C}$ , which is thereby endowed with the corresponding algebraic structure.

We have treated briefly the principal algebraic structures that occur later in this seminar. For the others (the structure of an  $\mathbf{O}$ -Lie algebra, for example), we believe that the reader has now seen enough examples in this section to make the general mechanism sketched in 2.2 work in each particular case.

We shall now apply what we have just done to the category of schemes, denoted  $(\mathbf{Sch})$ , and more generally to the categories  $(\mathbf{Sch})_{/S}$ , also denoted  $(\mathbf{Sch}_{/S})$ .

## 4. Algebraic structures in the category of schemes

Whenever there is no ambiguity, we shall permit ourselves the following abuses of language. We shall denote by  $T$  the object  $T \xrightarrow{f} S$  of  $\mathcal{C}_{/S}$ , with the datum of  $f$  (the structural morphism of  $T$ ) understood, and we shall identify  $\mathcal{C}$  with a subcategory of  $\widehat{\mathcal{C}}$ . In view of the compatibilities stated in the preceding sections, these identifications may be made without danger.

We shall also simplify terminology according to the following pattern: a  $(\mathbf{Sch})$ -group will also be called a *group scheme*; a  $(\mathbf{Sch})_{/S}$ -group will be called a *group scheme over  $S$* , an  *$S$ -group scheme*, or an  *$S$ -group*; and it will be called an  *$A$ -group* when  $S$  is the spectrum of the ring  $A$ .

<sup>29</sup>Editor's note: the sentence which follows has been added; it will be used in Section 5.

#### 4.1. Constant schemes

The category of schemes satisfies the conditions required in 1.8. Constant objects may therefore be defined in it. For every set  $E$ , there is a scheme  $E_{\mathbb{Z}}$ , and for every scheme  $S$ , an  $S$ -scheme

$$E_S = (E_{\mathbb{Z}})_S.$$

Recall (cf. *loc. cit.*) that for every  $S$ -scheme  $T$ ,

$$\mathrm{Hom}_S(T, E_S)$$

is the set of locally constant maps from the topological space underlying  $T$  to  $E$ .

The functor  $E \mapsto E_S$  commutes with finite inverse limits. In particular, if  $G$  is a group, then  $G_S$  is an  $S$ -group scheme; if  $O$  is a ring, then  $O_S$  is an  $S$ -ring scheme; and similarly for other algebraic structures.

#### 4.2. Affine $S$ -groups

Let us recall some facts about affine  $S$ -schemes (EGA II, Section 1). An  $S$ -scheme  $T$  is said to be *affine over  $S$*  if the inverse image of every affine open subset of  $S$  is affine. The  $\mathcal{O}_S$ -algebra  $f_*\mathcal{O}_T$ , denoted by  $\mathcal{A}(T)$ , is then quasi-coherent, where  $f$  denotes the structural morphism of  $T$ . Conversely, to every quasi-coherent  $\mathcal{O}_S$ -algebra  $\mathcal{A}$ , one can associate an  $S$ -scheme affine over  $S$ , denoted  $\mathrm{Spec}(\mathcal{A})$ . The functors

$$T \longmapsto \mathcal{A}(T), \quad \mathcal{A} \longmapsto \mathrm{Spec}(\mathcal{A})$$

are quasi-inverse equivalences between the category of  $S$ -schemes affine over  $S$  and the category opposite to that of quasi-coherent  $\mathcal{O}_S$ -algebras.

It follows that giving an algebraic structure on an  $S$ -scheme  $T$  affine over  $S$  is equivalent to giving the corresponding structure on  $\mathcal{A}(T)$  in the category opposite to that of quasi-coherent  $\mathcal{O}_S$ -algebras. In particular, if  $G$  is an  $S$ -group affine over  $S$ , then  $\mathcal{A}(G)$  is endowed with the structure of an augmented  $\mathcal{O}_S$ -bialgebra: there are two morphisms of  $\mathcal{O}_S$ -algebras

$$\Delta : \mathcal{A}(G) \longrightarrow \mathcal{A}(G) \otimes_{\mathcal{O}_S} \mathcal{A}(G), \quad \varepsilon : \mathcal{A}(G) \longrightarrow \mathcal{O}_S,$$

corresponding to the morphisms of  $S$ -schemes

$$\pi : G \times G \longrightarrow G, \quad e_G : S \longrightarrow G.$$

The maps  $\Delta$  and  $\varepsilon$  satisfy the following conditions, which express that  $G$  is an  $S$ -monoid.

<sup>30</sup>

**(HA 1)**  $\Delta$  is coassociative: the following diagram is commutative.

---

<sup>30</sup>Editor's note: of course, the inversion morphism  $G \rightarrow G$  induces a morphism of  $\mathcal{O}_S$ -algebras  $\tau : \mathcal{A}(G) \rightarrow \mathcal{A}(G)$  which, together with  $\Delta$  and  $\varepsilon$ , makes  $\mathcal{A}(G)$  into a Hopf  $\mathcal{O}_S$ -algebra.

$$\begin{array}{ccccc}
& & \mathcal{A}(G) \otimes_{\mathcal{O}_S} \mathcal{A}(G) & & \\
& \nearrow \Delta & & \searrow \text{Id} \otimes \Delta & \\
\mathcal{A}(G) & & & & \mathcal{A}(G) \otimes_{\mathcal{O}_S} \mathcal{A}(G) \otimes_{\mathcal{O}_S} \mathcal{A}(G) \\
& \searrow \Delta & & \nearrow \Delta \otimes \text{Id} & \\
& & \mathcal{A}(G) \otimes_{\mathcal{O}_S} \mathcal{A}(G) & &
\end{array}$$

Figure 1: Coassociativity of the diagonal map  $\Delta$ .

Temporary Loop-1 image; source locator: *Polo-Gille Exp1-13oct24.pdf*, component-local p. 22;  
combined-reader p. 32.

(HA 2)  $\Delta$  is compatible with  $\varepsilon$ : the following two composites are the identity:

$$\begin{aligned}
\mathcal{A}(G) &\xrightarrow{\Delta} \mathcal{A}(G) \otimes_{\mathcal{O}_S} \mathcal{A}(G) \xrightarrow{\text{Id} \otimes \varepsilon} \mathcal{A}(G) \otimes_{\mathcal{O}_S} \mathcal{O}_S \xrightarrow{\sim} \mathcal{A}(G), \\
\mathcal{A}(G) &\xrightarrow{\Delta} \mathcal{A}(G) \otimes_{\mathcal{O}_S} \mathcal{A}(G) \xrightarrow{\varepsilon \otimes \text{Id}} \mathcal{O}_S \otimes_{\mathcal{O}_S} \mathcal{A}(G) \xrightarrow{\sim} \mathcal{A}(G).
\end{aligned}$$

Let us take this opportunity to note once again that it follows from the definition of an  $S$ -group structure that, in order to give such a structure on an  $S$ -scheme  $G$  affine over  $S$ , it is not necessary to verify anything on  $\mathcal{A}(G)$ ; one need only endow every  $G(S')$ , for  $S'$  over  $S$ , with a group structure functorial in  $S'$ . This observation applies *mutatis mutandis* to morphisms.

### 4.3. The groups $\mathbb{G}_a$ and $\mathbb{G}_m$ . The ring $\mathcal{O}$

#### 4.3.1.

Let  $\mathbb{G}_a$  be the functor from  $(\mathbf{Sch})^\circ$  to  $\mathbf{Set}$  defined by

$$\mathbb{G}_a(S) = \Gamma(S, \mathcal{O}_S),$$

endowed with the  $(\widehat{\mathbf{Sch}})$ -group structure defined by the additive-group structure of the ring  $\Gamma(S, \mathcal{O}_S)$ . It is representable by an affine scheme, which we denote by  $\mathbb{G}_a$ , and which is therefore a group scheme:

$$\mathbb{G}_a = \text{Spec } \mathbb{Z}[T].$$

Indeed,

$$\mathbb{G}_a(S) = \text{Hom}(S, \mathbb{G}_a) = \text{Hom}_{\text{Alg.}}(\mathbb{Z}[T], \Gamma(S, \mathcal{O}_S)) \simeq \Gamma(S, \mathcal{O}_S).$$

For every scheme  $S$ , one therefore has an  $S$ -group affine over  $S$ , denoted  $\mathbb{G}_{a,S}$ , corresponding to the  $\mathcal{O}_S$ -bialgebra  $\mathcal{O}_S[T]$ , whose diagonal map and augmentation are defined by

$$\Delta(T) = T \otimes 1 + 1 \otimes T, \quad \varepsilon(T) = 0.$$

#### 4.3.2.

Let  $\mathbb{G}_m$  be the functor from  $(\mathbf{Sch})^\circ$  to  $\mathbf{Set}$  defined by

$$\mathbb{G}_m(S) = \Gamma(S, \mathcal{O}_S)^\times,$$

where  $\Gamma(S, \mathcal{O}_S)^\times$  denotes the multiplicative group of invertible elements of the ring  $\Gamma(S, \mathcal{O}_S)$ , endowed with its natural  $\widehat{(\mathbf{Sch})}$ -group structure. It is representable by an affine scheme denoted by  $\mathbb{G}_m$ :

$$\mathbb{G}_m = \mathrm{Spec} \mathbb{Z}[T, T^{-1}] = \mathrm{Spec} \mathbb{Z}[\mathbb{Z}],$$

where  $\mathbb{Z}[\mathbb{Z}]$  denotes the group algebra of  $\mathbb{Z}$  over the ring  $\mathbb{Z}$ . Indeed,

$$\mathbb{G}_m(S) = \mathrm{Hom}_{\mathrm{Alg}}(\mathbb{Z}[T, T^{-1}], \Gamma(S, \mathcal{O}_S)) \simeq \Gamma(S, \mathcal{O}_S)^\times.$$

For every scheme  $S$ , one therefore has an  $S$ -group affine over  $S$ , denoted  $\mathbb{G}_{m,S}$ , corresponding to the  $\mathcal{O}_S$ -algebra  $\mathcal{O}_S[\mathbb{Z}]$ , whose diagonal map and augmentation are defined by

$$\Delta(x) = x \otimes x, \quad \varepsilon(x) = 1, \quad x \in \mathbb{Z}.$$

#### 4.3.3.

Let  $\mathbf{O}$  be the functor  $\mathbb{G}_a$  endowed with its  $\widehat{(\mathbf{Sch})}$ -ring structure. It is represented by the scheme  $\mathrm{Spec} \mathbb{Z}[T]$ , which we denote by  $\mathbf{O}$  when it is regarded as endowed with its ring-scheme structure.

For every scheme  $S$ ,

$$\mathbf{O}_S = S \times_{\mathrm{Spec} \mathbb{Z}} \mathrm{Spec} \mathbb{Z}[T] = \mathrm{Spec}(\mathcal{O}_S[T])$$

is therefore an  $S$ -ring scheme affine over  $S$ . (Note: in EGA II, 1.7.13,  $\mathbf{O}_S$  is denoted by  $S[T]$ .)

**4.3.3.1.** <sup>31</sup> For every object  $X$  of  $\widehat{(\mathbf{Sch})}$ ,

$$\mathbf{O}(X) = \mathrm{Hom}(X, \mathbf{O})$$

has a ring structure, functorial in  $X$ . In particular, if  $X'$  is a scheme and morphisms  $x : X' \rightarrow X$  and  $f : X \rightarrow \mathbf{O}$  are given (that is,  $f \in \mathbf{O}(X)$ ), then

$$f(x) = f \circ x \in \mathbf{O}(X') = \Gamma(X', \mathcal{O}_{X'}).$$

**Definition.** Let  $\pi : M \rightarrow X$  be a morphism in  $\widehat{(\mathbf{Sch})}$ , and set

$$\mathbf{O}_X = \mathbf{O} \times X.$$

We say that  $M$  is an  $\mathbf{O}_X$ -module if, for every  $X$ -scheme  $X'$ , the set

$$\mathrm{Hom}_X(X', M)$$

has been endowed with an  $\mathbf{O}(X')$ -module structure, functorial in  $X'$ .

This is equivalent to giving an  $X$ -abelian-group law

$$\mu : M \times_X M \longrightarrow M$$

and an *external law*

$$\mathbf{O} \times M = \mathbf{O}_X \times_X M \longrightarrow M, \quad (f, m) \longmapsto f \cdot m,$$

which is an  $X$ -morphism, that is,  $\pi(f \cdot m) = \pi(m)$ , and which, for every  $x \in X(S)$ , endows

$$M(x) = \{m \in M(S) \mid \pi(m) = x\}$$

with an  $\mathbf{O}(S)$ -module structure.

In that case, for every  $Y \in \mathrm{Ob}(\widehat{(\mathbf{Sch})}_{/X})$ , not necessarily representable,

$$\mathrm{Hom}_X(Y, M) = \Gamma(M_Y/Y)$$

is an  $\mathbf{O}(Y)$ -module, functorially in  $Y$ .

---

<sup>31</sup>Editor's note: this paragraph has been added; it will be useful later (cf. II, 1.3).

#### 4.4. Diagonalizable groups

**4.4.1.** The construction of  $\mathbb{G}_m$  generalizes as follows. Let  $M$  be a commutative group, and let  $M_{\mathbb{Z}}$  be the group scheme associated with it by the procedure of 4.1. Consider the functor defined by

$$\mathbf{D}(M)(S) = \mathrm{Hom}_{\mathrm{groups}}(M, \mathbb{G}_m(S)) \simeq \underline{\mathrm{Hom}}_{S\text{-}\mathbf{Gr}}(M_S, \mathbb{G}_{m,S}).$$

This is a commutative  $\widehat{(\mathbf{Sch})}$ -group represented by a group scheme denoted  $D(M)$ ; thus, by definition,

$$D(M) \simeq \underline{\mathrm{Hom}}_{(\mathbf{Sch})\text{-}\mathbf{Gr}}(M_{\mathbb{Z}}, \mathbb{G}_m).$$

Indeed, set

$$D(M) = \mathrm{Spec} \mathbb{Z}[M],$$

where  $\mathbb{Z}[M]$  is the group algebra of  $M$  over  $\mathbb{Z}$ . Then

$$D(M)(S) = \mathrm{Hom}_{\mathrm{Alg.}}(\mathbb{Z}[M], \Gamma(S, \mathcal{O}_S)) \simeq \mathrm{Hom}_{\mathrm{groups}}(M, \Gamma(S, \mathcal{O}_S)^{\times})$$

by the very definition of the algebra  $\mathbb{Z}[M]$ .

**4.4.2.** For every scheme  $S$ , one therefore has an  $S$ -group scheme affine over  $S$ ,

$$D_S(M) = D(M)_S = (\underline{\mathrm{Hom}}_{(\mathbf{Sch})\text{-}\mathbf{Gr}}(M_{\mathbb{Z}}, \mathbb{G}_m))_S = \underline{\mathrm{Hom}}_{S\text{-}\mathbf{Gr}}(M_S, \mathbb{G}_{m,S}).$$

It is associated with the  $\mathcal{O}_S$ -bialgebra  $\mathcal{O}_S[M]$ , endowed with the diagonal map and augmentation defined by

$$\Delta(x) = x \otimes x, \quad \varepsilon(x) = 1, \quad x \in M.$$

**4.4.3.** If  $f : M \rightarrow N$  is a homomorphism of commutative groups, one defines in the evident way a morphism of  $S$ -groups

$$D_S(f) : D_S(N) \longrightarrow D_S(M),$$

and hence a functor

$$D_S : M \longmapsto D_S(M)$$

from the category of abelian groups to the category of  $S$ -groups affine over  $S$ . It may also be defined as the composite of the functor  $M \mapsto M_S$  and the functor

$$M_S \longmapsto \underline{\mathrm{Hom}}_{S\text{-}\mathbf{Gr}}(M_S, \mathbb{G}_{m,S}).$$

This functor commutes with base extension.

An  $S$ -group isomorphic to a group of the form  $D_S(M)$  is said to be *diagonalizable*. Notice that the elements of  $M$  are interpreted as certain characters of  $D_S(M)$ , that is, as certain elements of

$$\mathrm{Hom}_{S\text{-}\mathbf{Gr}}(D_S(M), \mathbb{G}_{m,S}).$$

(Compare VIII, 1.)

**4.4.4.** Let us give some examples of diagonalizable groups. First,

$$D(\mathbb{Z}) = \mathbb{G}_m, \quad D(\mathbb{Z}^r) = (\mathbb{G}_m)^r.$$

Set

$$\mu_n = D(\mathbb{Z}/n\mathbb{Z}),$$

and call it the *group of  $n$ -th roots of unity*. Indeed,

$$\mu_n(S) = \text{Hom}_{\text{groups}}(\mathbb{Z}/n\mathbb{Z}, \Gamma(S, \mathcal{O}_S)^\times) = \{f \in \Gamma(S, \mathcal{O}_S) \mid f^n = 1\}.$$

The  $S$ -group  $\mu_{n,S}$  corresponds to the  $\mathcal{O}_S$ -algebra  $\mathcal{O}_S[T]/(T^n - 1)$ . Suppose, in particular, that  $S$  is the spectrum of a field  $k$  of characteristic  $p = n$ . On setting  $T - 1 = s$ , one obtains

$$k[T]/(T^n - 1) = k[s]/(s^p),$$

which shows that the underlying topological space of  $\mu_{p,k}$  consists of one point, whose local ring is the Artinian  $k$ -algebra  $k[s]/(s^p)$ . In the same vein, note that the  $S$ -schemes  $\mathbb{G}_{a,S}$ ,  $\mathbb{G}_{m,S}$ , and  $\mathbf{O}_S$  are smooth over  $S$ , and that  $D_S(M)$  is

flat over  $S$ . Moreover, it is formally smooth (respectively, smooth) over  $S$  if and only if no residue characteristic of  $S$  divides the torsion of  $M$  (respectively, and if in addition  $M$  is of finite type); see Exposé VIII, 2.1.

#### 4.5. Other examples of groups

The preceding procedure applies to the “classical groups” (linear groups  $\text{GL}_n$ , symplectic groups  $\text{Sp}_n$ , orthogonal groups  $\text{O}_n$ , and so forth). For example, one defines  $\text{GL}_n$  as representing the functor  $\mathbf{GL}_n$  such that

$$\mathbf{GL}_n(S) = \text{GL}(n, \Gamma(S, \mathcal{O}_S)) = \text{Aut}_{\mathcal{O}_S}(\mathcal{O}_S^n).$$

It may be constructed, for example, as the open subscheme of  $\text{Spec } \mathbb{Z}[T_{ij}]$ ,  $1 \leq i, j \leq n$ , defined by the function

$$f = \det((T_{ij})_{i,j=1}^n),$$

that is, as  $\text{Spec } \mathbb{Z}[T_{ij}, f^{-1}]$ .

#### 4.6. Functor-modules in the category of schemes

We shall associate with every  $\mathcal{O}_S$ -module on a scheme  $S$  a  $\mathbf{O}_S$ -module, where  $\mathbf{O}_S$  denotes the functor-ring introduced in 4.3.3. This can be done in two different ways, as follows.

**Definition 4.6.1.** Let  $S$  be a scheme. For every  $\mathcal{O}_S$ -module  $\mathcal{F}$ , denote by  $\mathbf{V}(\mathcal{F})$  and  $\mathbf{W}(\mathcal{F})$  the contravariant functors on  $(\mathbf{Sch})_S$  defined by

$$\begin{aligned} \mathbf{V}(\mathcal{F})(S') &= \text{Hom}_{\mathcal{O}_{S'}}(\mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{O}_{S'}, \mathcal{O}_{S'}), \\ \mathbf{W}(\mathcal{F})(S') &= \Gamma(S', \mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{O}_{S'}), \end{aligned}$$

where  $\mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{O}_{S'}$  denotes the inverse image on  $S'$  of the  $\mathcal{O}_S$ -module  $\mathcal{F}$ .

Then  $\mathbf{V}(\mathcal{F})$  and  $\mathbf{W}(\mathcal{F})$  are endowed in the evident way with  $\mathbf{O}_S$ -module structures (recall that  $\mathbf{O}_S(S') = \Gamma(S', \mathcal{O}_{S'}) = \mathbf{W}(\mathcal{O}_S)(S')$ ). Thus one has in fact defined two functors  $\mathbf{V}$  and  $\mathbf{W}$  from the category of  $\mathcal{O}_S$ -modules to the category of  $\mathbf{O}_S$ -modules, with  $\mathbf{V}$  contravariant and  $\mathbf{W}$  covariant.

For the remainder of this paragraph, we restrict to the case in which the  $\mathcal{O}_S$ -modules in question are quasi-coherent. Thus we regard  $\mathbf{V}$  and  $\mathbf{W}$  as functors

$$\mathbf{V} : (\mathcal{O}_S\text{-Mod.q.c.})^\circ \longrightarrow (\mathbf{O}_S\text{-Mod.}), \quad \mathbf{W} : (\mathcal{O}_S\text{-Mod.q.c.}) \longrightarrow (\mathbf{O}_S\text{-Mod.}).$$

**Remark 4.6.1.1.** <sup>32</sup> The reader will notice that, in what follows, every proposition involving only the functor  $\mathbf{W}$  remains valid for arbitrary  $\mathcal{O}_S$ -modules, not necessarily quasi-coherent.

#### Proposition 4.6.2.

<sup>32</sup>Editor’s note: the numbering 4.6.1.1 has been added for later references.

- (i) The functors  $\mathbf{V}$  and  $\mathbf{W}$  commute with base extension: if  $S'$  lies over  $S$  and  $\mathcal{F}$  is a quasi-coherent  $\mathcal{O}_S$ -module, then

$$\mathbf{V}(\mathcal{F} \otimes \mathcal{O}_{S'}) \simeq \mathbf{V}(\mathcal{F})_{S'}, \quad \mathbf{W}(\mathcal{F} \otimes \mathcal{O}_{S'}) \simeq \mathbf{W}(\mathcal{F})_{S'}.$$

- (ii) The functors  $\mathbf{V}$  and  $\mathbf{W}$  are fully faithful: the canonical maps

$$\mathrm{Hom}_{\mathcal{O}_S}(\mathcal{F}, \mathcal{F}') \longrightarrow \mathrm{Hom}_{\mathbf{O}_S}(\mathbf{V}(\mathcal{F}'), \mathbf{V}(\mathcal{F})),$$

$$\mathrm{Hom}_{\mathcal{O}_S}(\mathcal{F}, \mathcal{F}') \longrightarrow \mathrm{Hom}_{\mathbf{O}_S}(\mathbf{W}(\mathcal{F}), \mathbf{W}(\mathcal{F}'))$$

are bijective.

- (iii) The functors  $\mathbf{V}$  and  $\mathbf{W}$  are additive:

$$\mathbf{V}(\mathcal{F} \oplus \mathcal{F}') \simeq \mathbf{V}(\mathcal{F}) \times_S \mathbf{V}(\mathcal{F}'), \quad \mathbf{W}(\mathcal{F} \oplus \mathcal{F}') \simeq \mathbf{W}(\mathcal{F}) \times_S \mathbf{W}(\mathcal{F}').$$

Parts (i) and (iii) are immediate from the definitions. For (ii), take the  $S'$  to be open subschemes of  $S$ . We leave the proof to the reader (for  $\mathbf{V}$ , use EGA II, 1.7.14).

Recall (cf. 3.1.4) that if  $\mathbf{F}$  and  $\mathbf{F}'$  are  $\mathbf{O}_S$ -modules, then  $\underline{\mathrm{Hom}}_{\mathbf{O}_S}(\mathbf{F}, \mathbf{F}')$  denotes the  $S$ -functor in abelian groups that assigns to every  $S' \rightarrow S$  the group

$$\mathrm{Hom}_{\mathbf{O}_{S'}}(\mathbf{F}_{S'}, \mathbf{F}'_{S'}).$$

**Proposition 4.6.3.** *There are canonical morphisms in  $(\mathbf{O}_S\text{-Mod.})$ :* <sup>33</sup>

$$\begin{array}{ccc} \underline{\mathrm{Hom}}_{\mathbf{O}_S}(\mathbf{W}(\mathcal{F}), \mathbf{W}(\mathcal{F}')) & \xlongequal{\quad} & \underline{\mathrm{Hom}}_{\mathbf{O}_S}(\mathbf{V}(\mathcal{F}'), \mathbf{V}(\mathcal{F})) \\ & \nwarrow \quad \nearrow & \\ & \mathbf{W}(\mathrm{Hom}_{\mathcal{O}_S}(\mathcal{F}, \mathcal{F}')) & \end{array} .$$

Figure 2: Canonical morphisms of Proposition 4.6.3.

*Temporary Loop-1 image; source locator: Polo-Gille Exp1-13oct24.pdf, component-local p. 26;  
combined-reader p. 36.*

This follows immediately from 4.6.2(i) and (ii).

**Notation 4.6.3.1.** Let  $\mathcal{F}$  be a quasi-coherent  $\mathcal{O}_S$ -module. By EGA II, 1.7.8, the  $S$ -functor  $\mathbf{V}(\mathcal{F})$  is represented by an  $S$ -scheme affine over  $S$ , denoted  $\mathbb{V}(\mathcal{F})$ , and called the *vector fibration* <sup>34</sup> defined by  $\mathcal{F}$ :

$$\mathbb{V}(\mathcal{F}) = \mathrm{Spec}(\mathcal{S}(\mathcal{F})),$$

where  $\mathcal{S}(\mathcal{F})$  denotes the symmetric algebra of the  $\mathcal{O}_S$ -module  $\mathcal{F}$ . <sup>35</sup>

**Proposition 4.6.4.** *Let  $\mathcal{F}$  and  $\mathcal{F}'$  be two quasi-coherent  $\mathcal{O}_S$ -modules, and let  $\mathcal{A}$  be a quasi-coherent  $\mathcal{O}_S$ -algebra. There is a functorial isomorphism*

$$\mathrm{Hom}_S(\mathrm{Spec}(\mathcal{A}), \underline{\mathrm{Hom}}_{\mathbf{O}_S}(\mathbf{W}(\mathcal{F}'), \mathbf{W}(\mathcal{F}))) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{O}_S}(\mathcal{F}', \mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{A}).$$

<sup>33</sup>Editor's note: the isomorphism  $\underline{\mathrm{Hom}}_{\mathbf{O}_S}(\mathbf{W}(\mathcal{F}), \mathbf{W}(\mathcal{F}')) \simeq \underline{\mathrm{Hom}}_{\mathbf{O}_S}(\mathbf{V}(\mathcal{F}'), \mathbf{V}(\mathcal{F}))$  has been added.

<sup>34</sup>Editor's note: "vector bundle" has been replaced by "vector fibration." Current usage calls a vector fibration that is locally trivial of rank  $r$  a "vector bundle of rank  $r$ ," that is, one whose sheaf of sections is locally isomorphic to  $\mathcal{O}_S^{\oplus r}$ .

<sup>35</sup>Editor's note: the articles [Ni04] and [Ni02], respectively, show that if  $S$  is Noetherian and  $\mathcal{F}$  is a coherent  $\mathcal{O}_S$ -module, then  $\mathbf{W}(\mathcal{F})$ , respectively the  $S$ -group assigning  $\mathrm{Aut}_{\mathcal{O}_T}(\mathcal{F} \otimes \mathcal{O}_T)$  to every  $T \rightarrow S$ , is representable if and only if  $\mathcal{F}$  is locally free.



Indeed, put  $X = \operatorname{Spec}(\mathcal{A})$ . The left-hand side is canonically isomorphic to

$$\underline{\operatorname{Hom}}_{\mathbf{O}_S}(\mathbf{W}(\mathcal{F}'), \mathbf{W}(\mathcal{F}))(X),$$

that is, by definition, to

$$\begin{aligned} & \operatorname{Hom}_{\mathbf{O}_X}(\mathbf{W}(\mathcal{F}')_X, \mathbf{W}(\mathcal{F})_X) \\ & \simeq \operatorname{Hom}_{\mathbf{O}_X}(\mathbf{W}(\mathcal{F}' \otimes \mathcal{O}_X), \mathbf{W}(\mathcal{F} \otimes \mathcal{O}_X)) \end{aligned}$$

(cf. 4.6.2(i)). By 4.6.2(ii), this may also be written as

$$\operatorname{Hom}_{\mathcal{O}_X}(\mathcal{F}' \otimes \mathcal{O}_X, \mathcal{F} \otimes \mathcal{O}_X) = \operatorname{Hom}_{\mathcal{O}_S}(\mathcal{F}', \pi_* \pi^*(\mathcal{F})),$$

where  $\pi : X \rightarrow S$  is the structural morphism. But EGA II, 1.4.7 gives

$$\pi_* \pi^*(\mathcal{F}) \simeq \mathcal{F} \otimes \mathcal{A},$$

which completes the proof.

**Corollary 4.6.4.1.** *There is a canonical isomorphism*

$$\mathbf{W}(\mathcal{F} \otimes \mathcal{A}) \simeq \underline{\operatorname{Hom}}_S(\operatorname{Spec}(\mathcal{A}), \mathbf{W}(\mathcal{F})).$$

Indeed, <sup>36</sup> let  $f : S' \rightarrow S$  be an  $S$ -scheme and put  $X' = X \times_S S'$ . There is a Cartesian square

$$\begin{array}{ccc} X' & \xrightarrow{f'} & X \\ \pi' \downarrow & & \downarrow \pi \\ S' & \xrightarrow{f} & S \end{array}$$

Figure 3: The Cartesian base-change square used in Corollary 4.6.4.1.

*Temporary Loop-1 image; source locator: Polo-Gille Exp1-13oct24.pdf, component-local p. 27;  
combined-reader p. 37.*

By EGA II, 1.5.2,  $X'$  is affine over  $S'$  and  $\pi'_* \mathcal{O}_{X'} = f^*(\mathcal{A})$ . Hence

$$\begin{aligned} & \underline{\operatorname{Hom}}_S(\operatorname{Spec}(\mathcal{A}), \mathbf{W}(\mathcal{F}))(S') \\ & = \operatorname{Hom}_{S'}(\operatorname{Spec}(f^* \mathcal{A}), \mathbf{W}(f^* \mathcal{F})). \end{aligned}$$

By 4.6.4, applied to  $f^* \mathcal{F}$ ,  $\mathcal{F}' = \mathcal{O}_{S'}$ , and  $f^* \mathcal{A}$ , this is equal to

$$\begin{aligned} \Gamma(S', f^* \mathcal{F} \otimes f^* \mathcal{A}) &= \Gamma(S', f^*(\mathcal{F} \otimes \mathcal{A})) \\ &= \mathbf{W}(\mathcal{F} \otimes \mathcal{A})(S'). \end{aligned}$$

**Proposition 4.6.5.** *If  $\mathcal{F}$  and  $\mathcal{F}'$  are two  $\mathcal{O}_S$ -modules locally free of finite type, the morphisms of 4.6.3 are isomorphisms.*

Indeed, for every  $S' \rightarrow S$ ,

$$\begin{aligned} & \mathbf{W}(\mathcal{H}om_{\mathcal{O}_S}(\mathcal{F}, \mathcal{F}'))(S') \\ & = \Gamma(S', \mathcal{H}om_{\mathcal{O}_S}(\mathcal{F}, \mathcal{F}') \otimes \mathcal{O}_{S'}) \\ & = \operatorname{Hom}_{\mathcal{O}_{S'}}(\mathcal{F} \otimes \mathcal{O}_{S'}, \mathcal{F}' \otimes \mathcal{O}_{S'}). \end{aligned}$$

<sup>36</sup>Editor's note: the original has been expanded in what follows.

The last member is isomorphic both to

$$\underline{\mathrm{Hom}}_{\mathbf{O}_S}(\mathbf{W}(\mathcal{F}), \mathbf{W}(\mathcal{F}'))(S')$$

and to

$$\underline{\mathrm{Hom}}_{\mathbf{O}_S}(\mathbf{V}(\mathcal{F}'), \mathbf{V}(\mathcal{F}))(S')$$

by 4.6.2(i) and (ii).

**Corollary 4.6.5.1.** *Let  $\mathcal{F}$  be an  $\mathcal{O}_S$ -module locally free of finite type. Put  $\mathcal{F}^\vee = \mathcal{H}om_{\mathcal{O}_S}(\mathcal{F}, \mathcal{O}_S)$ . There are canonical isomorphisms*

$$\mathbf{W}(\mathcal{F}^\vee) \simeq \underline{\mathrm{Hom}}_{\mathbf{O}_S}(\mathbf{W}(\mathcal{F}), \mathbf{O}_S) \simeq \mathbf{V}(\mathcal{F}),$$

$$\mathbf{V}(\mathcal{F}^\vee) \simeq \underline{\mathrm{Hom}}_{\mathbf{O}_S}(\mathbf{V}(\mathcal{F}), \mathbf{O}_S) \simeq \mathbf{W}(\mathcal{F}).$$

Finally, we have the following proposition.

**Proposition 4.6.6.** *Let  $f : \mathcal{F} \rightarrow \mathcal{F}'$  be a morphism of  $\mathcal{O}_S$ -modules locally free of finite rank. The morphism*

$$\mathbf{W}(f) : \mathbf{W}(\mathcal{F}) \longrightarrow \mathbf{W}(\mathcal{F}')$$

*is a monomorphism if and only if  $f$  identifies  $\mathcal{F}$  locally with a direct summand of  $\mathcal{F}'$ .*

The direct assertion is essentially contained in EGA 0<sub>I</sub>, 5.5.5.<sup>37</sup> Conversely, if  $\mathcal{F}$  is locally a direct summand of  $\mathcal{F}'$ , then for every  $\pi : S' \rightarrow S$ , the module  $\pi^*\mathcal{F}$  is a submodule of  $\pi^*\mathcal{F}'$ , since this holds locally as a direct summand. Thus

$$\mathbf{W}(\mathcal{F})(S') = \Gamma(S', \pi^*\mathcal{F})$$

is a submodule of

$$\mathbf{W}(\mathcal{F}')(S') = \Gamma(S', \pi^*\mathcal{F}').$$

#### 4.7. The category of $G$ - $\mathcal{O}_S$ -modules

Let  $G$  be an  $S$ -group and  $\mathcal{F}$  an  $\mathcal{O}_S$ -module;  $\mathbf{W}(\mathcal{F})$  has a  $\mathbf{O}_S$ -module structure.

**Definition 4.7.1.** A  $G$ - $\mathcal{O}_S$ -module structure on  $\mathcal{F}$  is an  $\mathbf{h}_G$ - $\mathbf{O}_S$ -module structure on  $\mathbf{W}(\mathcal{F})$  (cf. 3.2). A morphism of  $G$ - $\mathcal{O}_S$ -modules is, by definition, a morphism of the associated  $\mathbf{h}_G$ - $\mathbf{O}_S$ -modules. This gives the category  $(G\text{-}\mathcal{O}_S\text{-}\mathbf{Mod.})$ ; we denote by  $(G\text{-}\mathcal{O}_S\text{-}\mathbf{Mod.q.c.})$  its full subcategory consisting of those  $G$ - $\mathcal{O}_S$ -modules that are quasi-coherent as  $\mathcal{O}_S$ -modules.

By 3.2, giving a  $G$ - $\mathcal{O}_S$ -module structure on  $\mathcal{F}$  is therefore equivalent to giving a morphism of  $(\widehat{\mathbf{Sch}})_{/S}$ -groups

$$\rho : \mathbf{h}_G \longrightarrow \underline{\mathrm{Aut}}_{\mathbf{O}_S}(\mathbf{W}(\mathcal{F})).$$

**Remark 4.7.1.0.**<sup>38</sup> Since 4.6.2 gives an anti-isomorphism of  $S$ -functors in groups

$$(\dagger) \quad \underline{\mathrm{Aut}}_{\mathbf{O}_S}(\mathbf{W}(\mathcal{F})) \simeq \underline{\mathrm{Aut}}_{\mathbf{O}_S}(\mathbf{V}(\mathcal{F})),$$

it is equivalent to give an  $\mathbf{h}_G$ - $\mathbf{O}_S$ -module structure on  $\mathbf{W}(\mathcal{F})$  or on  $\mathbf{V}(\mathcal{F})$ . Indeed, let

$$\rho : \mathbf{h}_G \longrightarrow \underline{\mathrm{Aut}}_{\mathbf{O}_S}(\mathbf{W}(\mathcal{F}))$$

be as above. For every  $T \rightarrow S$  and  $g \in G(T)$ , denote by  $\rho^*(g)$  the image of  $\rho(g)$  under the anti-isomorphism  $(\dagger)$ . Then

$$\rho^*(gh) = \rho^*(h) \circ \rho^*(g),$$

<sup>37</sup>Editor's note: the following sentence has been expanded.

<sup>38</sup>Editor's note: this remark has been added from [DG], II, Section 2, 1.1.

so  $\rho^*$  defines a *right*  $\mathbf{h}_G\text{-}\mathbf{O}_S$ -module structure on  $\mathbf{V}(\mathcal{F})$ . Setting

$$\rho^\vee(g) = \rho^*(g^{-1}),$$

one obtains an  $\mathbf{h}_G\text{-}\mathbf{O}_S$ -module structure on  $\mathbf{V}(\mathcal{F})$ , whose datum is equivalent to that of  $\rho$ .

**Remark 4.7.1.1.** One may say that the categories just defined have been constructed by the following Cartesian squares.

$$\begin{array}{ccccc} (G\text{-}\mathcal{O}_S\text{-Mod.q.c.}) & \hookrightarrow & (G\text{-}\mathcal{O}_S\text{-Mod.}) & \longrightarrow & (\mathbf{h}_G\text{-}\mathbf{O}_S\text{-Mod.}) \\ \downarrow & & \downarrow & & \downarrow \text{oubli} \\ (\mathcal{O}_S\text{-Mod.q.c.}) & \hookrightarrow & (\mathcal{O}_S\text{-Mod.}) & \xrightarrow{\mathbf{W}} & (\mathbf{O}_S\text{-Mod.}). \end{array}$$

Figure 4: Cartesian squares defining the categories of  $G\text{-}\mathcal{O}_S$ -modules.

*Temporary Loop-1 image; source locator: Polo-Gille Exp1-13oct24.pdf, component-local p. 28; combined-reader p. 38.*

The categories  $(\mathcal{O}_S\text{-Mod.})$  and  $(\mathbf{O}_S\text{-Mod.})$  are abelian, but one should remember that in general the functor  $\mathbf{W}$  is neither left nor right exact.<sup>39</sup>

**Remark 4.7.1.2.**<sup>40</sup> Let  $\mathcal{F}$  be a  $G\text{-}\mathcal{O}_S$ -module. The subsheaf of invariants  $\mathcal{F}^G$  is defined as follows: for every open subset  $U$  of  $S$ ,

$$\mathcal{F}^G(U) = \mathbf{W}(\mathcal{F})^G(U) = \{x \in \mathcal{F}(U) \mid g \cdot x_{S'} = x_{S'} \text{ for every } S' \xrightarrow{f} U, g \in G(S')\},$$

where  $x_{S'}$  denotes the image of  $x$  in

$$\Gamma(S', f^*\mathcal{F}) = \Gamma(U, f_*f^*\mathcal{F}).$$

One should note that the natural morphism

$$\mathbf{W}(\mathcal{F}^G) \longrightarrow \mathbf{W}(\mathcal{F})^G$$

is not an isomorphism in general. For example, let  $S = \text{Spec}(\mathbb{Z})$ , and let  $G$  be the constant group  $\mathbb{Z}/2\mathbb{Z} = \{1, \tau\}$  acting on  $\mathcal{F} = \mathcal{O}_S$  by  $\tau \cdot 1 = -1$ . Then  $\mathcal{F}^G = 0$ , whereas, if  $R$  is an  $\mathbb{F}_2$ -algebra,

$$\mathbf{W}(\mathcal{F})^G(\text{Spec}(R)) = R.$$

**4.7.2.** From now until the end of 4.7, suppose that  $G$  is affine over  $S$ .<sup>41</sup> By 4.6.4, giving a morphism of  $S$ -functors

$$\rho : \mathbf{h}_G \longrightarrow \underline{\text{End}}_{\mathbf{O}_S}(\mathbf{W}(\mathcal{F}))$$

is equivalent to giving a morphism of  $\mathcal{O}_S$ -modules

$$\mu : \mathcal{F} \longrightarrow \mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{A}(G).$$

<sup>39</sup>Editor's note: the original has been corrected by deleting the assertion that the category  $(G\text{-}\mathcal{O}_S\text{-Mod.})$  is abelian; see 4.7.2.1 below.

<sup>40</sup>Editor's note: this remark has been added.

<sup>41</sup>Editor's note: see VI<sub>B</sub>, Sections 11.1–11.6 for the extension of the results of 4.7.2 to the case where  $G$  is not necessarily affine but  $G$  and  $\mathcal{F}$  are assumed flat over  $S$ .

Saying that  $\rho$  is a morphism of  $(\widehat{\mathbf{Sch}})_S$ -groups is then equivalent to saying that  $\mu$  satisfies the following axioms.

(CM 1) The following diagram is commutative.

$$\begin{array}{ccc}
 \mathcal{F} & \xrightarrow{\mu} & \mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{A}(G) \\
 \mu \downarrow & & \downarrow \text{id} \otimes \Delta \\
 \mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{A}(G) & \xrightarrow{\mu \otimes \text{id}} & \mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{A}(G) \otimes_{\mathcal{O}_S} \mathcal{A}(G) .
 \end{array}$$

Figure 5: The coassociativity axiom for the coaction  $\mu$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp1-13oct24.pdf, component-local p. 29;  
combined-reader p. 39.*

(CM 2) The following composite is the identity.

$$\mathcal{F} \xrightarrow{\mu} \mathcal{F} \otimes \mathcal{A}(G) \xrightarrow{\text{id} \otimes \varepsilon} \mathcal{F} \otimes \mathcal{O}_S \xrightarrow{\sim} \mathcal{F} .$$

Figure 6: The counit axiom for the coaction  $\mu$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp1-13oct24.pdf, component-local p. 29;  
combined-reader p. 39.*

These axioms (CM 1) and (CM 2) are those of a right *comodule* structure over the bialgebra  $\mathcal{A}(G)$ .<sup>42</sup>

Put  $\mathcal{A} = \mathcal{A}(G)$ . If  $\mathcal{F}$  and  $\mathcal{F}'$  are  $\mathcal{A}$ -comodules, a morphism of comodules  $f : \mathcal{F} \rightarrow \mathcal{F}'$  is a morphism of  $\mathcal{O}_S$ -modules such that the following diagram is commutative.

$$\begin{array}{ccc}
 \mathcal{F} & \xrightarrow{f} & \mathcal{F}' \\
 \mu_{\mathcal{F}} \downarrow & & \downarrow \mu_{\mathcal{F}'} \\
 \mathcal{F} \otimes \mathcal{A} & \xrightarrow{f \otimes \text{id}} & \mathcal{F}' \otimes \mathcal{A} .
 \end{array}$$

Figure 7: Compatibility required of a morphism of comodules.

*Temporary Loop-1 image; source locator: Polo-Gille Exp1-13oct24.pdf, component-local p. 30;  
combined-reader p. 40.*

This gives the category  $(\mathcal{A}\text{-}\mathbf{Comod})$ ; we denote by  $(\mathcal{A}\text{-}\mathbf{Comod.q.c.})$  its full subcategory consisting of those  $\mathcal{A}$ -comodules that are quasi-coherent as  $\mathcal{O}_S$ -modules. We have therefore proved the following statement.

**Proposition 4.7.2.** *Let  $G$  be an  $S$ -group affine over  $S$ . There are equivalences of categories*

$$\begin{aligned}
 (G\text{-}\mathcal{O}_S\text{-}\mathbf{Mod.}) &\simeq (\mathcal{A}(G)\text{-}\mathbf{Comod.}), \\
 (G\text{-}\mathcal{O}_S\text{-}\mathbf{Mod.q.c.}) &\simeq (\mathcal{A}(G)\text{-}\mathbf{Comod.q.c.}).
 \end{aligned}$$

<sup>42</sup>Editor's note: left  $G\text{-}\mathcal{O}_S$ -modules correspond naturally to right  $\mathcal{A}(G)$ -comodules.

If, in addition,  $S = \operatorname{Spec}(\Lambda)$  is affine and one sets  $\Lambda[G] = \Gamma(S, \mathcal{A}(G))$ , there is an equivalence of categories

$$(\mathcal{A}(G)\text{-}\mathbf{Comod.q.c.}) \simeq (\Lambda[G]\text{-}\mathbf{Comod.}).$$

43

Suppose, moreover, that  $\mathcal{A} = \mathcal{A}(G)$  is a flat  $\mathcal{O}_S$ -module.<sup>44</sup> Let  $\mathcal{E}$  be an  $\mathcal{A}$ -comodule and  $\mathcal{F}$  an  $\mathcal{O}_S$ -submodule of  $\mathcal{E}$ . Since  $\mathcal{A}$  is flat over  $\mathcal{O}_S$ , one may identify  $\mathcal{F} \otimes \mathcal{A}$  (respectively,  $\mathcal{F} \otimes \mathcal{A} \otimes \mathcal{A}$ ) with an  $\mathcal{O}_S$ -submodule of  $\mathcal{E} \otimes \mathcal{A}$  (respectively, of  $\mathcal{E} \otimes \mathcal{A} \otimes \mathcal{A}$ ). Suppose that  $\mu_{\mathcal{E}}$  maps  $\mathcal{F}$  into  $\mathcal{F} \otimes \mathcal{A}$ . Then its restriction

$$\mu_{\mathcal{F}} : \mathcal{F} \longrightarrow \mathcal{F} \otimes \mathcal{A}$$

endows  $\mathcal{F}$  with an  $\mathcal{A}$ -comodule structure; one says that  $\mathcal{F}$  is a *subcomodule* of  $\mathcal{E}$ . Passing to the quotient,  $\mu_{\mathcal{E}}$  defines a morphism of  $\mathcal{O}_S$ -modules

$$\mathcal{E}/\mathcal{F} \longrightarrow (\mathcal{E}/\mathcal{F}) \otimes \mathcal{A},$$

which endows  $\mathcal{E}/\mathcal{F}$  with an  $\mathcal{A}$ -comodule structure. If  $f : \mathcal{E} \rightarrow \mathcal{E}'$  is a morphism of  $\mathcal{A}$ -comodules, then  $\operatorname{Ker} f$  (respectively,  $\operatorname{Im} f$ ) is an  $\mathcal{A}$ -subcomodule of  $\mathcal{E}$  (respectively, of  $\mathcal{E}'$ ), and  $f$  induces an isomorphism of  $\mathcal{A}$ -comodules

$$\mathcal{E}/\operatorname{Ker} f \xrightarrow{\sim} \operatorname{Im} f.$$

Moreover, if  $\mathcal{E}$  and  $\mathcal{E}'$  are quasi-coherent  $\mathcal{O}_S$ -modules, then so are  $\operatorname{Ker} f$  and  $\operatorname{Im} f$ . Consequently,  $(\mathcal{A}\text{-}\mathbf{Comod.})$  and  $(\mathcal{A}\text{-}\mathbf{Comod.q.c.})$  are abelian categories.

**Corollary 4.7.2.1.** *Suppose that  $G$  is affine and flat over  $S$ . Then the category  $(G\text{-}\mathcal{O}_S\text{-}\mathbf{Mod.q.c.})$  (respectively,  $(G\text{-}\mathcal{O}_S\text{-}\mathbf{Mod.})$ ), equivalent to the category of  $\mathcal{A}(G)$ -comodules that are quasi-coherent over  $\mathcal{O}_S$  (respectively, of all  $\mathcal{A}(G)$ -comodules), is abelian.*

**4.7.3.** Suppose now that  $G$  is a *diagonalizable group*, that is, that  $\mathcal{A}(G)$  is the algebra of a commutative group  $M$  over the sheaf of rings  $\mathcal{O}_S$ . If  $\mathcal{F}$  is an  $\mathcal{O}_S$ -module, then

$$\mathcal{F} \otimes \mathcal{A}(G) = \coprod_{m \in M} \mathcal{F} \otimes m\mathcal{O}_S.$$

Giving a morphism of  $\mathcal{O}_S$ -modules

$$\mu : \mathcal{F} \longrightarrow \mathcal{F} \otimes \mathcal{A}(G)$$

is therefore equivalent to giving  $\mathcal{O}_S$ -endomorphisms  $(\mu_m)_{m \in M}$  of  $\mathcal{F}$  such that, for every section  $x$  of  $\mathcal{F}$  on an open subset of  $S$ , the family  $(\mu_m(x))$  is a section of the direct sum  $\coprod_{m \in M} \mathcal{F}$ . This means that on every sufficiently small open subset, only finitely many restrictions of the  $\mu_m(x)$  are nonzero.

For  $\mu$  defined by

$$\mu(x) = \sum_{m \in M} \mu_m(x) \otimes m,$$

condition (CM 1), respectively (CM 2), holds if and only if

$$\mu_m \circ \mu_{m'} = \delta_{mm'} \mu_m, \quad \left( \text{respectively, } \sum_{m \in M} \mu_m = \operatorname{Id}_{\mathcal{F}} \right).$$

<sup>43</sup>Editor's note: the preceding sentence has been added.

<sup>44</sup>Editor's note: the following paragraph has been added from [Se68, Section 1.3].

Thus the  $\mu_m$  are pairwise orthogonal projectors whose sum is the identity. We have proved the following result.

**Proposition 4.7.3.** *If  $G = D_S(M)$  is a diagonalizable  $S$ -group, the category of quasi-coherent  $G$ - $\mathcal{O}_S$ -modules (respectively, of all  $G$ - $\mathcal{O}_S$ -modules) is equivalent to the category of quasi-coherent  $\mathcal{O}_S$ -modules (respectively, of all  $\mathcal{O}_S$ -modules) graded by  $M$ .*

**Remark.** If  $\mathcal{F}$  has an  $\mathcal{O}_S$ -algebra structure preserved by the operations of  $G$ , then the grading of  $\mathcal{F}$  is an algebra grading. More precisely:

**Corollary 4.7.3.1.** *The functor  $\mathcal{A} \mapsto \text{Spec}(\mathcal{A})$  induces an equivalence between the category of quasi-coherent  $\mathcal{O}_S$ -algebras graded by  $M$  and the category opposite to that of  $S$ -schemes affine over  $S$  with  $S$ -operator group  $G = D_S(M)$ .*

**Proposition 4.7.4.** *Let  $G$  be a diagonalizable  $S$ -group. If*

$$0 \longrightarrow \mathcal{F}_1 \longrightarrow \mathcal{F}_2 \longrightarrow \mathcal{F}_3 \longrightarrow 0$$

*is an exact sequence of quasi-coherent  $G$ - $\mathcal{O}_S$ -modules that splits as a sequence of  $\mathcal{O}_S$ -modules, then it also splits as a sequence of  $G$ - $\mathcal{O}_S$ -modules.*

Indeed, if  $G = D_S(M)$ , each  $\mathcal{F}_i$  is graded by the  $(\mathcal{F}_i)_m$ , and for every  $m \in M$  the sequence

$$0 \longrightarrow (\mathcal{F}_1)_m \longrightarrow (\mathcal{F}_2)_m \longrightarrow (\mathcal{F}_3)_m \longrightarrow 0$$

of  $\mathcal{O}_S$ -modules is split. The preceding proposition then gives the result.

## 5. Group cohomology

### 5.1. The standard complex

Let  $\mathcal{C}$  be a category,  $\mathbf{G}$  a  $\widehat{\mathcal{C}}$ -group,  $\mathbf{O}$  a  $\widehat{\mathcal{C}}$ -ring, and  $\mathbf{F}$  a  $\mathbf{G}$ - $\mathbf{O}$ -module.<sup>45</sup> For  $n \geq 0$ , set

$$C^n(\mathbf{G}, \mathbf{F}) = \text{Hom}(\mathbf{G}^n, \mathbf{F}), \quad \underline{C}^n(\mathbf{G}, \mathbf{F}) = \underline{\text{Hom}}(\mathbf{G}^n, \mathbf{F}),$$

where  $\mathbf{G}^0$  is the final object  $\mathbf{e}$ . Then  $\underline{C}^n(\mathbf{G}, \mathbf{F})$  (respectively,  $C^n(\mathbf{G}, \mathbf{F})$ ) is endowed in the evident way with an  $\mathbf{O}$ -module structure (respectively, a  $\Gamma(\mathbf{O})$ -module structure), and

$$C^n(\mathbf{G}, \mathbf{F}) \simeq \Gamma(\underline{C}^n(\mathbf{G}, \mathbf{F})), \quad \underline{C}^n(\mathbf{G}, \mathbf{F})(S) = C^n(\mathbf{G}_S, \mathbf{F}_S).$$

To give an element of  $C^n(\mathbf{G}, \mathbf{F})$  is to give, for every  $S \in \text{Ob}(\mathcal{C})$ , an  $n$ -cochain of  $\mathbf{G}(S)$  with values in  $\mathbf{F}(S)$ , functorially in  $S$ . Recall that the boundary operator

$$\partial : C^n(\mathbf{G}(S), \mathbf{F}(S)) \longrightarrow C^{n+1}(\mathbf{G}(S), \mathbf{F}(S))$$

is given by

$$\begin{aligned} \partial f(g_1, \dots, g_{n+1}) &= g_1 f(g_2, \dots, g_{n+1}) \\ &\quad + \sum_{i=1}^n (-1)^i f(g_1, \dots, g_i g_{i+1}, \dots, g_{n+1}) \\ &\quad + (-1)^{n+1} f(g_1, \dots, g_n). \end{aligned}$$

<sup>45</sup>Editor's note: This complex is often called the Hochschild complex; see, for example, Section II.3 of [DG70].

It is functorial in  $S$ , and hence defines a homomorphism

$$\partial : C^n(\mathbf{G}, \mathbf{F}) \longrightarrow C^{n+1}(\mathbf{G}, \mathbf{F})$$

such that  $\partial \circ \partial = 0$ . We have thus defined a complex of abelian groups (indeed, of  $\Gamma(\mathbf{O})$ -modules), denoted  $C^*(\mathbf{G}, \mathbf{F})$ . In the same way one defines the complex of  $\mathbf{O}$ -modules  $\underline{C}^*(\mathbf{G}, \mathbf{F})$ , and

$$C^*(\mathbf{G}, \mathbf{F}) = \Gamma(\underline{C}^*(\mathbf{G}, \mathbf{F})).$$

We denote by  $H^n(\mathbf{G}, \mathbf{F})$  (respectively,  $\underline{H}^n(\mathbf{G}, \mathbf{F})$ ) the groups (respectively, the  $\widehat{\mathcal{C}}$ -groups) of cohomology of  $C^*(\mathbf{G}, \mathbf{F})$  (respectively,  $\underline{C}^*(\mathbf{G}, \mathbf{F})$ ). In particular,

$$\underline{H}^0(\mathbf{G}, \mathbf{F}) = \mathbf{F}^{\mathbf{G}}, \quad H^0(\mathbf{G}, \mathbf{F}) = \Gamma(\mathbf{F}^{\mathbf{G}}).$$

**Remark 5.1.1.** The “set-theoretic” description of  $\partial$  given above is convenient for checking that  $\partial \circ \partial = 0$ . One may also define  $\partial$  in terms of the multiplication  $m : \mathbf{G} \times \mathbf{G} \rightarrow \mathbf{G}$  and the action  $\mu : \mathbf{G} \times \mathbf{F} \rightarrow \mathbf{F}$  as follows: for every  $f \in C^n(\mathbf{G}, \mathbf{F})$ ,

$$\partial f = \mu \circ (\text{id}_{\mathbf{G}} \times f) + \sum_{i=1}^n (-1)^i f \circ (\text{id}_{\mathbf{G}^{i-1}} \times m \times \text{id}_{\mathbf{G}^{n-i}}) + (-1)^{n+1} f \circ \text{pr}_{[1,n]},$$

where  $\text{pr}_{[1,n]}$  denotes the projection of  $\mathbf{G}^{n+1} = \mathbf{G}^n \times \mathbf{G}$  onto  $\mathbf{G}^n$ . Similarly, for every  $S \in \text{Ob}(\mathcal{C})$  and  $f \in \underline{C}^n(\mathbf{G}, \mathbf{F})(S) = C^n(\mathbf{G}_S, \mathbf{F}_S)$ ,

$$\partial f = \mu_S \circ (\text{id}_{\mathbf{G}_S} \times f) + \sum_{i=1}^n (-1)^i f \circ (\text{id}_{\mathbf{G}_S^{i-1}} \times m_S \times \text{id}_{\mathbf{G}_S^{n-i}}) + (-1)^{n+1} f \circ \text{pr}_{[1,n]},$$

where  $m_S$  and  $\mu_S$  are obtained from  $m$  and  $\mu$  by base change.<sup>46</sup>

## 5.2.

Recall (see Section 3) that  $(\mathbf{G}\text{-}\mathbf{O}\text{-Mod.})$  is an abelian category, with its structure defined objectwise. Thus

$$0 \longrightarrow \mathbf{F}' \longrightarrow \mathbf{F} \longrightarrow \mathbf{F}'' \longrightarrow 0$$

is an exact sequence of  $\mathbf{G}\text{-}\mathbf{O}$ -modules if and only if the sequence of abelian groups

$$0 \longrightarrow \mathbf{F}'(S) \longrightarrow \mathbf{F}(S) \longrightarrow \mathbf{F}''(S) \longrightarrow 0$$

is exact for every  $S \in \text{Ob}(\mathcal{C})$ .<sup>47</sup>

Suppose that  $\mathcal{C}$  is small. Then, by 3.2.1,  $(\mathbf{G}\text{-}\mathbf{O}\text{-Mod.})$  has enough injective objects, so the derived functors of the left exact functors  $\underline{H}^0$  and  $H^0$  are defined. We shall now show that the functors  $\underline{H}^n$  (respectively,  $H^n$ ) are indeed the derived functors of  $\underline{H}^0$  (respectively,  $H^0$ ).<sup>48</sup>

**Definition 5.2.0.** For every  $\mathbf{O}$ -module  $\mathbf{P}$ , let  $E(\mathbf{P})$  denote the object  $\underline{\text{Hom}}(\mathbf{G}, \mathbf{P})$  of  $\widehat{\mathcal{C}}$ , endowed with the following  $\mathbf{G}\text{-}\mathbf{O}$ -module structure. For every  $S \in \text{Ob}(\mathcal{C})$ ,

$$\underline{\text{Hom}}(\mathbf{G}, \mathbf{P})(S) = \text{Hom}_S(\mathbf{G}_S, \mathbf{P}_S),$$

<sup>46</sup>Editor’s note: This remark has been added.

<sup>47</sup>Editor’s note: The preceding reminder has been added.

<sup>48</sup>Editor’s note: The preceding sentence has been added.

and  $g \in \mathbf{G}(S)$  and  $a \in \mathbf{O}(S)$  act on  $\phi \in \text{Hom}_S(\mathbf{G}_S, \mathbf{P}_S)$  by

$$(g \cdot \phi)(h) = \phi(hg), \quad (a \cdot \phi)(h) = a\phi(h),$$

for every  $h \in \mathbf{G}(S')$ ,  $S' \rightarrow S$ . Moreover, for every  $\phi \in \text{Hom}_S(\mathbf{G}_S, \mathbf{P}_S)$ , set

$$\varepsilon(\phi) = \phi(1) \in \mathbf{P}(S),$$

where 1 denotes the unit element of  $\mathbf{G}(S)$ . This defines a functor

$$E : (\mathbf{O}\text{-Mod.}) \longrightarrow (\mathbf{G}\text{-}\mathbf{O}\text{-Mod.})$$

and a natural transformation  $\varepsilon : E \rightarrow \text{Id}$ , where  $\text{Id}$  denotes the identity functor of  $(\mathbf{O}\text{-Mod.})$ .  
49

**Remark 5.2.0.1.** In what follows, let  $\mathbf{G}_1$  and  $\mathbf{G}_2$  denote two copies of  $\mathbf{G}$ . The morphism

$$\mathbf{G}_1 \times E(\mathbf{P}) \longrightarrow E(\mathbf{P}), \quad (g_1, \phi) \longmapsto (g_2 \longmapsto \phi(g_2 g_1))$$

corresponds, via the isomorphisms

$$\begin{aligned} \text{Hom}(\mathbf{G}_1 \times E(\mathbf{P}), E(\mathbf{P})) &\simeq \text{Hom}\left(E(\mathbf{P}), \underline{\text{Hom}}(\mathbf{G}_1, \underline{\text{Hom}}(\mathbf{G}_2, \mathbf{P}))\right) \\ &\simeq \text{Hom}\left(E(\mathbf{P}), \underline{\text{Hom}}(\mathbf{G}_2 \times \mathbf{G}_1, \mathbf{P})\right), \end{aligned}$$

to the morphism

$$\phi \longmapsto ((g_2, g_1) \longmapsto \phi(g_2 g_1)),$$

that is, to the morphism

$$\underline{\text{Hom}}(\mathbf{G}, \mathbf{P}) \longrightarrow \underline{\text{Hom}}(\mathbf{G}_2 \times \mathbf{G}_1, \mathbf{P})$$

induced by the multiplication

$$\mu_{\mathbf{G}} : \mathbf{G} \times \mathbf{G} \longrightarrow \mathbf{G}, \quad (g_2, g_1) \longmapsto g_2 g_1.$$

**Lemma 5.2.0.2.** (i) The functor  $E$  is right adjoint to the forgetful functor

$$(\mathbf{G}\text{-}\mathbf{O}\text{-Mod.}) \longrightarrow (\mathbf{O}\text{-Mod.}).$$

More precisely,  $\varepsilon : E \rightarrow \text{Id}$  induces, for every  $\mathbf{M} \in (\mathbf{G}\text{-}\mathbf{O}\text{-Mod.})$  and  $\mathbf{P} \in \text{Ob}(\mathbf{O}\text{-Mod.})$ , a bijection

$$\text{Hom}_{\mathbf{G}\text{-}\mathbf{O}\text{-Mod.}}(\mathbf{M}, E(\mathbf{P})) \xrightarrow{\sim} \text{Hom}_{\mathbf{O}\text{-Mod.}}(\mathbf{M}, \mathbf{P})$$

functorial in  $\mathbf{M}$  and  $\mathbf{P}$ .

(ii) Consequently, if  $\mathbf{I}$  is an injective object of  $(\mathbf{O}\text{-Mod.})$ , then  $E(\mathbf{I})$  is an injective object of  $(\mathbf{G}\text{-}\mathbf{O}\text{-Mod.})$ .

*Proof.* To every  $\mathbf{O}$ -morphism  $f : \mathbf{M} \rightarrow \mathbf{P}$ , associate the element  $\phi_f$  of  $\text{Hom}_{\mathbf{O}}(\mathbf{M}, E(\mathbf{P}))$  defined as follows. For every  $S \in \text{Ob}(\mathcal{C})$  and  $m \in \mathbf{M}(S)$ , the element  $\phi_f(m)$  of  $\text{Hom}_S(\mathbf{G}_S, \mathbf{P}_S)$  is defined, for every  $g \in \mathbf{G}(S')$ ,  $S' \rightarrow S$ , by

$$\phi_f(m)(g) = f(gm) \in \mathbf{P}(S').$$

---

<sup>49</sup>Editor's note: The original has been modified so as to introduce 5.2.0.1 and 5.2.0.2, which will be useful in the proof of Theorem 5.3.1.



Then, for every  $h \in \mathbf{G}(S)$ , one has  $\phi_f(hm) = h \cdot f(m)$ ; that is,  $\phi_f$  is an element of

$$\mathrm{Hom}_{\mathbf{G}\text{-}\mathbf{O}\text{-Mod.}}(\mathbf{M}, E(\mathbf{P})).$$

If  $\phi \in \mathrm{Hom}_{\mathbf{G}\text{-}\mathbf{O}\text{-Mod.}}(\mathbf{M}, E(\mathbf{P}))$ , and if for every  $m \in \mathbf{M}(S)$  we write  $f(m) = \phi(m)(1)$ , then

$$\phi_f(m)(g) = f(gm) = \phi(gm)(1) = (g \cdot \phi(m))(1) = \phi(m)(g);$$

thus  $\phi_f = \phi$ . Conversely, it is clear that  $\phi_f(m)(1) = f(m)$ . This proves (i), and (ii) follows at once.

**Definition 5.2.0.3.** Let  $\mathbf{M}$  be a  $\mathbf{G}\text{-}\mathbf{O}$ -module. The identity map of  $\mathbf{M}$ , considered as an  $\mathbf{O}$ -module, corresponds under the adjunction to the morphism of  $\mathbf{G}\text{-}\mathbf{O}$ -modules

$$\mu_{\mathbf{M}} : \mathbf{M} \longrightarrow E(\mathbf{M})$$

such that, for every  $S \in \mathrm{Ob}(\mathcal{C})$  and  $m \in \mathbf{M}(S)$ ,  $\mu_{\mathbf{M}}(m)$  is the morphism  $\mathbf{G}_S \rightarrow \mathbf{M}_S$  defined as follows: for every  $S' \rightarrow S$  and  $g \in \mathbf{G}(S')$ ,

$$\mu_{\mathbf{M}}(m)(g) = g \cdot m_{S'} \in \mathbf{M}(S').$$

Notice that  $\mu_{\mathbf{M}}$  is a monomorphism. Indeed,  $\varepsilon_{\mathbf{M}} : E(\mathbf{M}) \rightarrow \mathbf{M}$  is a morphism of  $\mathbf{O}$ -modules such that  $\varepsilon_{\mathbf{M}} \circ \mu_{\mathbf{M}} = \mathrm{id}_{\mathbf{M}}$ . Consequently, as an  $\mathbf{O}$ -module,  $\mathbf{M}$  is a direct summand of  $E(\mathbf{M})$ .

**Proposition 5.2.1.** *Suppose that  $\mathcal{C}$  is small, that finite products exist in  $\mathcal{C}$ , and that  $\mathbf{G}$  is representable. Then the functors  $\underline{H}^n(\mathbf{G}, -)$  (respectively,  $H^n(\mathbf{G}, -)$ ) are the derived functors of the left exact functor  $\underline{H}^0(\mathbf{G}, -)$  (respectively,  $H^0(\mathbf{G}, -)$ ) on the category of  $\mathbf{G}\text{-}\mathbf{O}$ -modules.*

By the well-known general results,<sup>50</sup> it suffices to verify that the  $\underline{H}^n(\mathbf{G}, -)$  (respectively,  $H^n(\mathbf{G}, -)$ ) form an effaceable cohomological functor in degrees  $> 0$ . Let

$$0 \longrightarrow \mathbf{F}' \longrightarrow \mathbf{F} \longrightarrow \mathbf{F}'' \longrightarrow 0$$

be an exact sequence of  $\mathbf{G}\text{-}\mathbf{O}$ -modules, and let  $S \in \mathrm{Ob}(\mathcal{C})$ . By hypothesis,  $\mathbf{G}$  is represented by an object  $G \in \mathrm{Ob}(\mathcal{C})$ , and finite products exist in  $\mathcal{C}$ ; in particular,  $\mathcal{C}$  has a final object  $e$ . Hence every  $\mathbf{G}^n \times \mathbf{h}_S$  is represented by  $G^n \times S$  (with  $G^0 = e$ ), and the sequence

$$0 \longrightarrow \mathbf{F}'(G^n \times S) \longrightarrow \mathbf{F}(G^n \times S) \longrightarrow \mathbf{F}''(G^n \times S) \longrightarrow 0$$

is exact. This shows that the sequence of  $\mathbf{O}$ -modules

$$0 \longrightarrow \underline{C}^n(\mathbf{h}_G, \mathbf{F}') \longrightarrow \underline{C}^n(\mathbf{h}_G, \mathbf{F}) \longrightarrow \underline{C}^n(\mathbf{h}_G, \mathbf{F}'') \longrightarrow 0$$

is exact. It follows that  $\underline{C}^*(\mathbf{G}, -)$ , regarded as a functor on  $(\mathbf{G}\text{-}\mathbf{O}\text{-Mod.})$  with values in the category of complexes of  $(\mathbf{O}\text{-Mod.})$ , is exact. Thus the functors  $\underline{H}^n(\mathbf{G}, -)$  are cohomological.

Since  $\Gamma$  is exact, the same is true of the functors  $H^n(\mathbf{G}, -)$ . It remains only to prove the following.

**Lemma 5.2.2.** *For every  $\mathbf{P} \in \mathrm{Ob}(\mathbf{O}\text{-Mod.})$ , one has*

$$\underline{H}^n(\mathbf{G}, \underline{\mathrm{Hom}}(\mathbf{G}, \mathbf{P})) = 0, \quad H^n(\mathbf{G}, \underline{\mathrm{Hom}}(\mathbf{G}, \mathbf{P})) = 0, \quad n > 0.$$

It is enough to show that  $\underline{C}^*(\mathbf{G}, \underline{\mathrm{Hom}}(\mathbf{G}, \mathbf{P}))$  and  $C^*(\mathbf{G}, \underline{\mathrm{Hom}}(\mathbf{G}, \mathbf{P}))$  are homotopically trivial in degrees  $> 0$ . It is enough, in fact, to do this for the second complex, since the

<sup>50</sup>Editor's note: See [Gr57, 2.2.1] and [Gr57, 2.3]. The original has also been expanded in what follows.

corresponding result for the first follows by base change.<sup>51</sup> For every  $n \geq 0$ , define a morphism

$$\sigma : C^{n+1}(\mathbf{G}, \underline{\mathrm{Hom}}(\mathbf{G}, \mathbf{P})) \longrightarrow C^n(\mathbf{G}, \underline{\mathrm{Hom}}(\mathbf{G}, \mathbf{P}))$$

as follows. Let  $f \in C^{n+1}(\mathbf{G}, \underline{\mathrm{Hom}}(\mathbf{G}, \mathbf{P}))$ . For every  $S \in \mathrm{Ob}(\mathcal{C})$  and  $g_1, \dots, g_n \in \mathbf{G}(S)$ , the element  $\sigma(f)(g_1, \dots, g_n)$  of  $\underline{\mathrm{Hom}}_S(\mathbf{G}_S, \mathbf{P}_S)$  is defined, for every  $S' \rightarrow S$  and  $x \in \mathbf{G}(S')$ , by

$$\sigma(f)(g_1, \dots, g_n)(x) = f(x, g_1, \dots, g_n)(e) \in \mathbf{P}(S'),$$

where  $e$  denotes the unit element of  $\mathbf{G}(S')$ . Then  $\sigma$  is a homotopy operator in degrees  $> 0$ . Indeed, for every  $g_1, \dots, g_{n+1} \in \mathbf{G}(S)$  and  $x \in \mathbf{G}(S')$ , on the one hand,

$$\begin{aligned} \partial\sigma(f)(g_1, \dots, g_{n+1})(x) &= f(xg_1, g_2, \dots, g_{n+1})(e) \\ &\quad + \sum_{i=1}^n (-1)^i f(x, g_1, \dots, g_i g_{i+1}, \dots, g_{n+1})(e) \\ &\quad + (-1)^{n+1} f(x, g_1, \dots, g_n)(e), \end{aligned}$$

and, on the other hand,

$$\begin{aligned} \sigma(\partial f)(g_1, \dots, g_{n+1})(x) &= (xf(g_1, g_2, \dots, g_{n+1}))(e) - f(xg_1, g_2, \dots, g_{n+1})(e) \\ &\quad + \sum_{i=1}^n (-1)^{i+1} f(x, g_1, \dots, g_i g_{i+1}, \dots, g_{n+1}) \\ &\quad + (-1)^{n+2} f(x, g_1, \dots, g_n)(e). \end{aligned}$$

Consequently,

$$(\partial\sigma(f) + \sigma(\partial f))(g_1, \dots, g_{n+1})(x) = f(g_1, \dots, g_{n+1})(x);$$

that is,  $\partial\sigma + \sigma\partial$  is the identity map of  $C^{n+1}(\mathbf{G}, \underline{\mathrm{Hom}}(\mathbf{G}, \mathbf{P}))$  for every  $n \geq 0$ .

**Remark 5.2.3.** The hypothesis that  $\mathcal{C}$  be small is used only to ensure the existence of the derived functors  $R^n \underline{H}^0$  and  $R^n H^0$ . In every case, what precedes shows that the functors  $\underline{H}^n(\mathbf{G}, -)$  (respectively,  $H^n(\mathbf{G}, -)$ ) form an effaceable cohomological functor in degrees  $> 0$ , and hence are the (right) satellite functors of the left exact functor  $\underline{H}^0(\mathbf{G}, -)$  (respectively,  $H^0(\mathbf{G}, -)$ ), in the sense of [Gr57, 2.2]. If  $(\mathbf{G}\text{-}\mathbf{O}\text{-Mod.})$  has enough injective objects (as it does when  $\mathcal{C}$  is small), they coincide with the derived functors (*loc. cit.*, 2.3).<sup>52</sup>

### 5.3. Cohomology of $G\text{-}\mathcal{O}_S$ -modules

Let  $S$  be a scheme,  $G$  an  $S$ -group, and  $\mathcal{F}$  a quasi-coherent  $G\text{-}\mathcal{O}_S$ -module. The cohomology groups of  $G$  with values in  $\mathcal{F}$  are defined by

$$H^n(G, \mathcal{F}) = H^n(\mathbf{h}_G, \mathbf{W}(\mathcal{F}))$$

(for the notation, see 4.6).

Suppose that  $G$  is affine over  $S$ . In view of Proposition 4.6.4, this cohomology is computed as follows:  $H^n(G, \mathcal{F})$  is the  $n$ -th cohomology group of the complex  $C^*(G, \mathcal{F})$ , whose  $n$ -th term is

$$C^n(G, \mathcal{F}) = \Gamma\left(S, \mathcal{F} \otimes \underbrace{\mathcal{A}(G) \otimes \dots \otimes \mathcal{A}(G)}_{n \text{ times}}\right).$$

<sup>51</sup>Editor's note: The original has been expanded in what follows.

<sup>52</sup>Editor's note: This remark has been added.

If  $f$  (respectively,  $a_i$ ) is a section of  $\mathcal{F}$  (respectively, of  $\mathcal{A}(G)$ ) over an open subset of  $S$ , then

$$\begin{aligned} \partial(f \otimes a_1 \otimes \cdots \otimes a_n) &= \mu_{\mathcal{F}}(f) \otimes a_1 \otimes a_2 \otimes \cdots \otimes a_n \\ &\quad + \sum_{i=1}^n (-1)^i f \otimes a_1 \otimes \cdots \otimes \Delta a_i \otimes \cdots \otimes a_n \\ &\quad + (-1)^{n+1} f \otimes a_1 \otimes a_2 \otimes \cdots \otimes a_n \otimes 1, \end{aligned}$$

where

$$\Delta : \mathcal{A}(G) \longrightarrow \mathcal{A}(G) \otimes \mathcal{A}(G), \quad \mu_{\mathcal{F}} : \mathcal{F} \longrightarrow \mathcal{F} \otimes \mathcal{A}(G)$$

describe the coalgebra structure of  $\mathcal{A}(G)$  and the comodule structure of  $\mathcal{F}$ . Notice in passing that the cohomology of  $G$  with values in  $\mathcal{F}$  therefore depends only on the comodule structure of  $\mathcal{F}$ , and in particular only on the  $S$ -monoid structure of  $G$ .

In particular,

$$H^0(G, \mathcal{F}) = \Gamma(S, \mathcal{F}^G),$$

where  $\mathcal{F}^G$ , the *sheaf of invariants of  $\mathcal{F}$* , is the sheaf whose sections over an open subset  $U$  of  $S$  are those sections of  $\mathcal{F}$  over  $U$  whose inverse image in every  $S'$  over  $U$  is invariant under  $G(S')$  (see 4.7.1.2).

**Theorem 5.3.1.** *Let  $S$  be an affine scheme and  $G$  an  $S$ -group that is affine and flat over  $S$ . The functors  $H^n(G, -)$  are the derived functors of  $H^0(G, -)$  on the category of quasi-coherent  $G$ - $\mathcal{O}_S$ -modules.*

*Proof.* Since  $G$  is affine and flat over  $S$ , 4.7.2.1 shows that the category  $(G\text{-}\mathcal{O}_S\text{-Mod.q.c.})$  is equivalent to the category  $(\mathcal{A}(G)\text{-Comod.q.c.})$  of quasi-coherent  $\mathcal{A}(G)$ -comodules over  $\mathcal{O}_S$ , and is therefore abelian. Moreover, since  $\mathcal{A}(G)$  is a flat  $\mathcal{O}_S$ -module, each functor

$$\mathcal{F} \longmapsto \mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{A}(G)^{\otimes n}$$

is exact; since  $S$  is affine, it follows that  $C^*(G, -)$  is an exact functor on  $(G\text{-}\mathcal{O}_S\text{-Mod.q.c.})$ .  
53

Let  $\Delta$  (respectively,  $\eta$ ) denote the comultiplication (respectively, the augmentation) of  $\mathcal{A}(G)$ . For every quasi-coherent  $\mathcal{O}_S$ -module  $\mathcal{P}$ , let

$$\text{Ind}(\mathcal{P}) = \mathcal{P} \otimes_{\mathcal{O}_S} \mathcal{A}(G),$$

endowed with the  $\mathcal{A}(G)$ -comodule structure defined by

$$\text{id}_{\mathcal{P}} \otimes \Delta : \mathcal{P} \otimes_{\mathcal{O}_S} \mathcal{A}(G) \longrightarrow \mathcal{P} \otimes_{\mathcal{O}_S} \mathcal{A}(G) \otimes_{\mathcal{O}_S} \mathcal{A}(G).$$

This defines a functor

$$\text{Ind} : (\mathcal{O}_S\text{-Mod.q.c.}) \longrightarrow (G\text{-}\mathcal{O}_S\text{-Mod.q.c.}).$$

It follows from 4.6.4.1, 5.2.0, and 5.2.0.1 that there is an isomorphism of  $\mathbf{G}\text{-}\mathbf{O}_S$ -modules

$$\mathbf{W}(\text{Ind}(\mathcal{P})) \simeq E(\mathbf{W}(\mathcal{P})) = \underline{\text{Hom}}(\mathbf{G}, \mathbf{W}(\mathcal{P})). \quad (*)$$

---

<sup>53</sup>Editor's note: The original has been modified to show that the category  $(G\text{-}\mathcal{O}_S\text{-Mod.q.c.})$  is abelian and has enough injective objects. Compare [Ja03], Part I, 3.3–3.4, 3.9, 4.2, and 4.14–4.16 (bearing in mind that “ $k$ -group scheme” there means “affine  $k$ -group scheme”; see *loc. cit.*, 2.1).

Under this identification, the morphism  $\varepsilon : E(\mathbf{W}(\mathcal{P})) \rightarrow \mathbf{W}(\mathcal{P})$  corresponds to the morphism of  $\mathcal{O}_S$ -modules

$$\mathrm{id}_{\mathcal{P}} \otimes \eta : \mathrm{Ind}(\mathcal{P}) \longrightarrow \mathcal{P}.$$

We have already used the fact that the functor

$$\mathbf{W} : (\mathcal{O}_S\text{-Mod.}) \longrightarrow (\mathbf{O}_S\text{-Mod.})$$

is fully faithful. By Definition 4.7.1, the same is true of its restriction to  $(G\text{-}\mathcal{O}_S\text{-Mod.})$ : if  $\mathcal{M}$  and  $\mathcal{M}'$  are  $G\text{-}\mathcal{O}_S$ -modules, there is a functorial isomorphism

$$\mathrm{Hom}_{G\text{-}\mathcal{O}_S\text{-Mod.}}(\mathcal{M}, \mathcal{M}') \simeq \mathrm{Hom}_{\mathbf{G}\text{-}\mathbf{O}_S\text{-Mod.}}(\mathbf{W}(\mathcal{M}), \mathbf{W}(\mathcal{M}')).$$

Consequently, Lemma 5.2.0.2 gives the following result.

**Corollary 5.3.1.1.** *(i) The functor  $\mathrm{Ind}$  is right adjoint to the forgetful functor*

$$(G\text{-}\mathcal{O}_S\text{-Mod.q.c.}) \longrightarrow (\mathcal{O}_S\text{-Mod.q.c.}).$$

*More precisely, the map  $\mathrm{id}_{\mathcal{P}} \otimes \eta : \mathrm{Ind}(\mathcal{P}) \rightarrow \mathcal{P}$  induces, for every object  $\mathcal{M}$  of  $(G\text{-}\mathcal{O}_S\text{-Mod.q.c.})$ , a bijection*

$$\mathrm{Hom}_{G\text{-}\mathcal{O}_S\text{-Mod.}}(\mathcal{M}, \mathrm{Ind}(\mathcal{P})) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{O}_S}(\mathcal{M}, \mathcal{P}).$$

*(ii) Consequently, if  $\mathcal{I}$  is an injective object of  $(\mathcal{O}_S\text{-Mod.q.c.})$ , then  $\mathrm{Ind}(\mathcal{I})$  is an injective object of  $(G\text{-}\mathcal{O}_S\text{-Mod.q.c.})$ .*

Let  $\mathcal{F}$  be a  $G\text{-}\mathcal{O}_S$ -module, and let  $\mu_{\mathcal{F}} : \mathcal{F} \rightarrow \mathrm{Ind}(\mathcal{F})$  be the map defining its  $\mathcal{A}(G)$ -comodule structure. It follows from 5.2.0.3 (or directly from axioms (CM 1) and (CM 2) of 4.7.2) that  $\mu_{\mathcal{F}}$  is a morphism of  $G\text{-}\mathcal{O}_S$ -modules, and that

$$(\mathrm{id}_{\mathcal{F}} \otimes \eta) \circ \mu_{\mathcal{F}} = \mathrm{id}_{\mathcal{F}}.$$

Thus  $\mathcal{F}$  is a direct summand of  $\mathrm{Ind}(\mathcal{F})$  as an  $\mathcal{O}_S$ -module; in particular,  $\mu_{\mathcal{F}}$  is a monomorphism. By (\*) and 5.2.2,

$$H^n(\mathbf{G}, \mathbf{W}(\mathrm{Ind}(\mathcal{F}))) \simeq H^n\left(\mathbf{G}, \underline{\mathrm{Hom}}_S(\mathbf{G}, \mathbf{W}(\mathcal{F}))\right) = 0 \quad (n > 0).$$

Therefore  $H^n(G, -)$  is effaceable for  $n > 0$ .

Finally, since  $S$  is affine,  $(\mathcal{O}_S\text{-Mod.q.c.})$  has enough injective objects. Let  $\mathcal{F} \hookrightarrow \mathcal{I}$  be a monomorphism of  $\mathcal{O}_S$ -modules, where  $\mathcal{I}$  is injective in  $(\mathcal{O}_S\text{-Mod.q.c.})$ . Since  $\mathcal{A}(G)$  is flat over  $\mathcal{O}_S$ ,  $\mathrm{Ind}(\mathcal{F})$  is a  $G\text{-}\mathcal{O}_S$ -submodule of  $\mathrm{Ind}(\mathcal{I})$ . Hence:

**Corollary 5.3.1.2.** *The abelian category  $(G\text{-}\mathcal{O}_S\text{-Mod.q.c.})$  has enough injective objects.*

Taking account of [Gr57, 2.2.1 and 2.3], already used in the proof of 5.2.1, completes the proof of Theorem 5.3.1.

**Remark 5.3.1.3.** Corollary 5.3.1.1 may also be proved by the following calculation. To every morphism of  $G\text{-}\mathcal{O}_S$ -modules

$$\phi : \mathcal{M} \longrightarrow \mathcal{P} \otimes_{\mathcal{O}_S} \mathcal{A}(G)$$

associate the  $\mathcal{O}_S$ -morphism

$$(\mathrm{id}_{\mathcal{P}} \otimes \eta) \circ \phi : \mathcal{M} \longrightarrow \mathcal{P}.$$

Conversely, to every  $\mathcal{O}_S$ -morphism  $f : \mathcal{M} \rightarrow \mathcal{P}$ , associate the morphism of  $G\text{-}\mathcal{O}_S$ -modules

$$(f \otimes \mathrm{id}_{\mathcal{A}(G)}) \circ \mu_{\mathcal{M}} : \mathcal{M} \longrightarrow \mathcal{P} \otimes_{\mathcal{O}_S} \mathcal{A}(G).$$

One sees at once that

$$(\mathrm{id}_{\mathcal{P}} \otimes \eta) \circ (f \otimes \mathrm{id}_{\mathcal{A}(G)}) \circ \mu_{\mathcal{M}} = (f \otimes \mathrm{id}_{\mathcal{O}_S}) \circ (\mathrm{id}_{\mathcal{P}} \otimes \eta) \circ \mu_{\mathcal{M}} = f.$$

On the other hand, for every  $\phi$ , the following diagram is commutative.

$$\begin{array}{ccc} \mathcal{M} & \xrightarrow{\psi} & \mathcal{P} \otimes_{\mathcal{O}_S} \mathcal{A}(G) \\ \mu_{\mathcal{M}} \downarrow & & \downarrow \mathrm{id}_{\mathcal{P}} \otimes \Delta \\ \mathcal{M} \otimes_{\mathcal{O}_S} \mathcal{A}(G) & \xrightarrow{\phi \otimes \mathrm{id}_{\mathcal{A}(G)}} & \mathcal{P} \otimes_{\mathcal{O}_S} \mathcal{A}(G) \otimes_{\mathcal{O}_S} \mathcal{A}(G). \end{array}$$

Figure 8: The adjunction square for  $\phi$ , the comodule map  $\mu_{\mathcal{M}}$ , and the comultiplication  $\Delta$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp1-13oct24.pdf, component-local/printed p. 38;  
combined-reader p. 48.*

It follows that

$$\begin{aligned} & \left( ((\mathrm{id}_{\mathcal{P}} \otimes \eta) \circ \phi) \otimes \mathrm{id}_{\mathcal{A}(G)} \right) \circ \mu_{\mathcal{M}} \\ &= (\mathrm{id}_{\mathcal{P}} \otimes \eta \otimes \mathrm{id}_{\mathcal{A}(G)}) \circ (\phi \otimes \mathrm{id}_{\mathcal{A}(G)}) \circ \mu_{\mathcal{M}} \\ &= (\mathrm{id}_{\mathcal{P}} \otimes \eta \otimes \mathrm{id}_{\mathcal{A}(G)}) \circ (\mathrm{id}_{\mathcal{P}} \otimes \Delta) \circ \phi = \phi. \end{aligned}$$

This proves Corollary 5.3.1.1(i), and (ii) follows.

Let  $\mathcal{F}$  be a  $G$ - $\mathcal{O}_S$ -module. We saw above that axiom (CM 2) of 4.7.2 shows that, as an  $\mathcal{O}_S$ -module,  $\mathcal{F}$  is a direct summand of  $E(\mathcal{F})$ . This gives:

**Proposition 5.3.2.** *Let  $S$  be an affine scheme and  $G$  an  $S$ -group that is affine and flat. Suppose that every exact sequence*

$$0 \longrightarrow \mathcal{F}_1 \longrightarrow \mathcal{F}_2 \longrightarrow \mathcal{F}_3 \longrightarrow 0$$

*of quasi-coherent  $G$ - $\mathcal{O}_S$ -modules which splits as a sequence of  $\mathcal{O}_S$ -modules also splits as a sequence of  $G$ - $\mathcal{O}_S$ -modules. Then the functors  $H^n(G, -)$  vanish for  $n > 0$  (or, equivalently, the functor  $H^0(G, -)$  is exact).*

Indeed, by hypothesis, the sequence of  $G$ - $\mathcal{O}_S$ -modules

$$0 \longrightarrow \mathcal{F} \longrightarrow E(\mathcal{F}) \longrightarrow E(\mathcal{F})/\mathcal{F} \longrightarrow 0$$

is split. Thus  $\mathcal{F}$  is a direct summand of  $E(\mathcal{F})$  as a  $G$ - $\mathcal{O}_S$ -module, while the cohomology of the latter vanishes.

Combining this with Proposition 4.7.4 immediately gives:

**Theorem 5.3.3.** *Let  $S$  be an affine scheme and  $G$  a diagonalizable  $S$ -group. For every quasi-coherent  $G$ - $\mathcal{O}_S$ -module  $\mathcal{F}$ , one has*

$$H^n(G, \mathcal{F}) = 0 \quad (n > 0).$$

**Remark.** Proposition 5.3.2 remains valid when  $G$  is not necessarily flat over  $S$ ; the proof then uses relative cohomology. <sup>54</sup>

<sup>54</sup>Editor's note: The editors have not attempted to develop this remark.

## 6. $G$ -equivariant objects and modules

Let  $\mathcal{C}$  be a category having a final object and in which fiber products exist.<sup>55</sup> Let  $G$  be a  $\widehat{\mathcal{C}}$ -group,  $\pi : M \rightarrow X$  a morphism in  $\widehat{\mathcal{C}}$ , and  $\lambda = \lambda_X : G \times X \rightarrow X$  an action of  $G$  on  $X$ . In what follows,  $Y \times_f M$  denotes the fiber product of  $\pi : M \rightarrow X$  and an  $X$ -functor  $f : Y \rightarrow X$ .

For every  $U \in \text{Ob}(\mathcal{C})$  and  $x \in X(U)$ , write  $M_x = U \times_x M$ ; thus, for every  $\phi : U' \rightarrow U$ ,

$$M_x(U') = \{m \in M(U') \mid \pi(m) = x_{U'} = \phi^*(x)\}.$$

Finally, if  $g \in G(U)$ , we also write  $gx$  for the element  $\lambda(g, x)$  of  $X(U)$ .

**Definition 6.1.** (a) We say that  $M$  is a  $G$ -equivariant  $X$ -object if an action  $\Lambda : G \times M \rightarrow M$  of  $G$  on  $M$ , lifting  $\lambda$ , has been given; that is, if the following square commutes.

$$\begin{array}{ccc} G \times M & \xrightarrow{\Lambda} & M \\ \text{id}_G \times \pi \downarrow & & \downarrow \pi \\ G \times X & \xrightarrow{\lambda} & X \end{array}$$

Figure 9: The action of  $G$  on  $M$  lifts its action on  $X$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp1-13oct24.pdf, component-local p. 39; combined-reader p. 49.*

Equivalently, for every morphism  $(g, x) : U \rightarrow G \times X$ , one is given maps

$$\Lambda_x^U(g) : M_x(U) \longrightarrow M_{gx}(U), \quad m \longmapsto g \cdot m,$$

which satisfy  $1 \cdot m = m$  and  $g \cdot (h \cdot m) = (gh) \cdot m$ , and which are functorial in the  $(G \times X)$ -object  $U$ . Equivalently again, one is given morphisms of  $U$ -objects

$$\Lambda_x(g) : M_x \longrightarrow M_{gx}$$

satisfying  $\Lambda_x(1) = \text{id}$  and  $\Lambda_{hx}(g) \circ \Lambda_x(h) = \Lambda_x(gh)$ .

(b) Now let  $\mathbf{O}$  be a  $\widehat{\mathcal{C}}$ -ring and put  $\mathbf{O}_X = \mathbf{O} \times X$ . Under the hypotheses of (a), we say that  $M$  is a  $G$ -equivariant  $\mathbf{O}_X$ -module if it is an  $\mathbf{O}_X$ -module (cf. Definition 4.3.3.1, which is valid for every ring functor on a category  $\mathcal{C}$ ) and if the action  $\Lambda$  is compatible with the  $\mathbf{O}_X$ -module structure on  $M$ . Thus, for every  $(g, x) \in G(U) \times X(U)$ ,  $U \in \text{Ob}(\mathcal{C})$ , the map

$$\Lambda_x(g) : M_x \longrightarrow M_{gx}, \quad m \longmapsto g \cdot m,$$

is a morphism of  $\mathbf{O}_U$ -modules.

**Remark 6.2.** (i) In (a), the conditions  $\Lambda_x(1) = \text{id}$  and  $\Lambda_{hx}(g) \circ \Lambda_x(h) = \Lambda_x(gh)$  plainly imply that each  $\Lambda_x(g)$  is an isomorphism, with inverse  $\Lambda_{gx}(g^{-1})$ . Conversely, if each  $\Lambda_x(g)$  is an isomorphism, applying the second condition with  $h = 1$  gives  $\Lambda_x(1) = \text{id}$ .

<sup>55</sup>Editor's note: This section has been added.

(ii) Let  $M$  be an  $\mathbf{O}_X$ -module. Giving a morphism  $\Lambda : G \times M \rightarrow M$  which makes the diagram of Definition 6.1 commute and for which every morphism

$$\Lambda_x(g) : M_x \longrightarrow M_{gx}, \quad m \longmapsto g \cdot m,$$

is an isomorphism of  $\mathbf{O}_U$ -modules is equivalent to giving an isomorphism of  $\mathbf{O}_{G \times X}$ -modules

$$\theta : G \times M = (G \times X) \times_{\text{pr}_X} M \xrightarrow{\sim} (G \times X) \times_{\lambda} M, \quad (g, x, m) \longmapsto (g, x, g \cdot m).$$

Since each  $\Lambda_x(g)$  is assumed to be an isomorphism, the equality  $\Lambda_x(1) = \text{id}$  follows from  $\Lambda_{hx}(g) \circ \Lambda_x(h) = \Lambda_x(gh)$  with  $h = 1$ . Consequently,  $\Lambda$  is an action of  $G$  on  $M$ , that is,  $g \cdot (h \cdot m) = (gh) \cdot m$ , if and only if the following diagram of  $(G \times G \times X)$ -isomorphisms commutes. Here  $\mu$  denotes multiplication in  $G$ , and  $f^*(\theta)$  is obtained from  $\theta$  by base change along  $f : G \times G \times X \rightarrow G \times X$ .

$$\begin{array}{ccc}
 (G \times G \times X) \times_{\text{pr}_X \circ \text{pr}_{2,3}} M & \xrightarrow[\sim]{\text{pr}_{2,3}^*(\theta)} & (G \times G \times X) \times_{\lambda \circ \text{pr}_{2,3}} M \\
 \parallel & & \parallel \\
 (G \times G \times X) \times_{\text{pr}_X \circ (\mu \times \text{id}_X)} M & & (G \times G \times X) \times_{\text{pr}_X \circ (\text{id}_G \times \lambda)} M \\
 \downarrow (\mu \times \text{id}_X)^*(\theta) \wr & & \downarrow \wr (\text{id}_G \times \lambda)^*(\theta) \\
 (G \times G \times X) \times_{\lambda \circ (\mu \times \text{id}_X)} M & \xlongequal{\quad} & (G \times G \times X) \times_{\lambda \circ (\text{id}_G \times \lambda)} M
 \end{array}$$

Figure 10: The cocycle condition for the isomorphism  $\theta$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp1-13oct24.pdf, component-local p. 40;  
combined-reader p. 50.*

Thus, giving a  $G$ -equivariant  $\mathbf{O}_X$ -module structure on  $M$  is equivalent to giving an isomorphism  $\theta$  of  $\mathbf{O}_{G \times X}$ -modules as above for which this diagram commutes.

(iii) Everything above extends to the case in which  $G$  is merely a  $\widehat{\mathcal{C}}$ -monoid. In that case, giving an action  $\Lambda : G \times M \rightarrow M$  lifting  $\lambda$ , such that every  $\Lambda_x(g) : M_x \rightarrow M_{gx}$  is a morphism of  $\mathbf{O}_U$ -modules, is equivalent to giving a morphism  $\theta$  of  $\mathbf{O}_{G \times X}$ -modules as in (ii), for which the diagram above commutes without the isomorphism signs under the arrows and which satisfies

$$p_M \circ \theta \circ (\varepsilon_G \times \text{id}_M) = \text{id}_M,$$

where  $\varepsilon_G$  is the unit section of  $G$  and  $p_M$  is the projection onto  $M$ .

### 6.3. $G$ -equivariant morphisms

Let  $Y$  be a second object of  $\widehat{\mathcal{C}}$ , endowed with an action  $\lambda_Y : G \times Y \rightarrow Y$ , and let  $N$  be a second  $G$ -equivariant  $\mathbf{O}_X$ -module. A morphism  $f : Y \rightarrow X$  in  $\widehat{\mathcal{C}}$  (respectively, a morphism

of  $\mathbf{O}_X$ -modules  $\phi : M \rightarrow N$ ) is called *G-equivariant* if it commutes with the action of  $G$ . In set-theoretic notation, this says

$$f(g \cdot y) = g \cdot f(y) \quad (\text{respectively, } \phi(g \cdot m) = g \cdot \phi(m)),$$

or, equivalently,

$$f \circ \lambda_Y = \lambda_X \circ (\text{id}_G \times f), \quad \phi \circ \Lambda_M = \Lambda_N \circ (\text{id}_G \times \phi).$$

The following lemma is immediate.

**Lemma 6.3.1.** *Let  $f : Y \rightarrow X$  be a  $G$ -equivariant morphism and let  $M$  be a  $G$ -equivariant  $\mathbf{O}_X$ -module. Then the inverse image  $f^*(M) = Y \times_f M$  is a  $G$ -equivariant  $\mathbf{O}_Y$ -module.*

On the other hand,  $G$  acts on  $\text{Hom}_{\widehat{\mathcal{C}}}(Y, X)$ . Indeed, let  $T \in \text{Ob}(\mathcal{C})$ ,  $g \in G(T)$ , and let  $\phi$  be a  $T$ -morphism  $Y_T \rightarrow X_T$ . Then  $g$  defines automorphisms  $\lambda_Y(g)$  and  $\lambda_X(g)$  of  $Y_T$  and  $X_T$ , and we write  $g \cdot \phi$ , or  $g\phi g^{-1}$ , for the morphism

$$\lambda_X(g) \circ \phi \circ \lambda_Y(g^{-1}).$$

This defines an action of  $G(T)$  on  $\text{Hom}_T(Y_T, X_T)$ , functorially in  $T$ .

**Definition 6.3.2.** *If  $\phi : Y \rightarrow X$  is an arbitrary morphism in  $\widehat{\mathcal{C}}$ , we may therefore consider its stabilizer  $\text{Stab}_G(\phi)$  (cf. 2.3.3.1). For every  $T \in \text{Ob}(\mathcal{C})$ ,  $\text{Stab}_G(\phi)(T)$  is the subgroup of  $G(T)$  formed by those  $g$  for which  $g \circ \phi_T = \phi_T \circ g$ , that is, for which the following diagram commutes.*

$$\begin{array}{ccc} Y_T & \xrightarrow{\phi_T} & X_T \\ g \downarrow & & \downarrow g \\ Y_T & \xrightarrow{\phi_T} & X_T \end{array}$$

Figure 11: The defining square for  $\text{Stab}_G(\phi)$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp1-13oct24.pdf, component-local p. 41;  
combined-reader p. 51.*

*Then  $\phi : Y \rightarrow X$  is equivariant under  $\text{Stab}_G(\phi)$ .*

## 6.4. Global sections

Let  $M$  be a  $G$ -equivariant  $\mathbf{O}_X$ -module. Let  $S_0$  denote the final object of  $\mathcal{C}$ , and denote by  $\prod_{X/S_0} M$  the *functor of sections of  $M$  over  $X$*  (cf. Exposé II, 1.1): to every  $T \in \text{Ob}(\mathcal{C})$  it assigns

$$\text{Hom}_X(X_T, M) = \text{Hom}_{X_T}(X_T, M_T) = \Gamma(M_T/X_T).$$

Recall also that every morphism  $g : Z \rightarrow Y$  of  $\widehat{\mathcal{C}}$ -objects over  $X$  induces a morphism of abelian groups

$$M(g) : M(Y) \longrightarrow M(Z),$$



compatible with the ring homomorphism  $g^* : \mathbf{O}(Y) \rightarrow \mathbf{O}(Z)$ . In particular, when  $Z = Y$ , so that  $g$  is an  $X$ -endomorphism of  $Y$ , one obtains a morphism of abelian groups

$$M(g) : M(Y) \longrightarrow M(Y)$$

which is not generally a morphism of  $\mathbf{O}(Y)$ -modules, but which satisfies

$$M(g)(\alpha \cdot m) = g^*(\alpha) \cdot M(g)(m)$$

for every  $m \in M(Y)$  and  $\alpha \in \mathbf{O}(Y)$ . In what follows, we simply write  $g^*$  instead of  $M(g)$ .

Let  $T \in \text{Ob}(\mathcal{C})$ , put  $X_T = X \times T$ , and let  $\text{pr}_X : X_T \rightarrow X$  be the projection. For  $\alpha \in \mathbf{O}(X_T)$  and  $g \in G(T)$ , set

$$g(\alpha) = (g^{-1})^*(\alpha).$$

Set-theoretically, this element of  $\mathbf{O}(X_T)$  is defined, for every  $S \in \text{Ob}(\mathcal{C})$ ,  $x \in X(S)$ , and  $t \in T(S)$ , by

$$g(\alpha)(x, t) = \alpha(g^{-1}x, t).$$

This gives a left action of  $G(T)$  by ring automorphisms on  $\mathbf{O}(X_T)$ , functorially in  $T$ , and  $g(\alpha) = \alpha$  when  $\alpha$  is the image in  $\mathbf{O}(X_T)$  of an element of  $\mathbf{O}(T)$ .

Let  $\phi$  now be the identity morphism of  $X_T$  (cf. 6.3 and the generalization below to a morphism  $\phi : Y \rightarrow X$ ). Denote  $\text{Hom}_X(X_T, M)$  by  $M(\phi g^{-1})$  or  $M(\phi)$  according as  $X_T$  is regarded as an  $X$ -object through  $\text{pr}_X \circ \lambda(g^{-1})_T$  or through  $\text{pr}_X$ , respectively.

Then  $\lambda(g^{-1})_T$  is an  $X$ -morphism between these two  $X$ -objects. Hence there is a morphism of abelian groups

$$(g^{-1})^* : M(\phi) \longrightarrow M(\phi g^{-1}), \quad m \longmapsto m \circ \lambda(g^{-1})_T,$$

and

$$(g^{-1})^*(\alpha \cdot m) = (g\alpha) \cdot (g^{-1})^*(m)$$

for  $m \in M(\phi)$  and  $\alpha \in \mathbf{O}(X_T)$ . If  $m$  is a section of  $M_T$  over  $X_T$ , then  $(g^{-1})^*(m)$  is the section defined set-theoretically by  $(x, t) \mapsto m(g^{-1}x, t)$ . In particular,  $(g^{-1})^*$  is a morphism of  $\mathbf{O}(T)$ -modules.

The functoriality of the  $\mathbf{O}(X_T)$ -module morphisms  $\Lambda_x(g)$  gives the following commutative diagram.

$$\begin{array}{ccc} \mathbf{M}(\phi) & \xrightarrow{(g^{-1})^*} & \mathbf{M}(\phi g^{-1}) \\ \Lambda_\phi(g) \downarrow & & \downarrow \Lambda_{\phi g^{-1}}(g) \\ \mathbf{M}(g\phi) & \xrightarrow{(g^{-1})^*} & \mathbf{M}(g\phi g^{-1}) \end{array}$$

Figure 12: Naturality of the maps  $\Lambda_x(g)$  on sections.

*Temporary Loop-1 image; source locator: Polo-Gille Exp1-13oct24.pdf, component-local p. 42;  
combined-reader p. 52.*

Moreover,  $g\phi g^{-1} = \phi$ , because  $\phi$  is the identity. Writing

$$A(g) = (g^{-1})^* \circ \Lambda_\phi(g),$$

we obtain a morphism of abelian groups

$$A(g) : M(\phi) \longrightarrow M(\phi)$$

which is compatible with the action of  $G(T)$  on  $\mathbf{O}(X_T)$ , namely

$$A(g)(\alpha \cdot m) = (g\alpha) \cdot A(g)(m).$$

Finally, if  $h$  is a second element of  $G(T)$ , functoriality of  $M$  and of the morphisms  $\Lambda_x(g)$  gives the following commutative diagram.

$$\begin{array}{ccccc}
 \mathbf{M}(\phi) & & & & \\
 \Lambda_\phi(g) \downarrow & \searrow \Lambda_\phi(hg) & & & \\
 \mathbf{M}(g\phi) & \xrightarrow{\Lambda_\phi(h)} & \mathbf{M}(hg\phi) & & \\
 (g^{-1})^* \downarrow & & (g^{-1})^* \downarrow & \searrow ((hg)^{-1})^* & \\
 \mathbf{M}(\phi) & \xrightarrow{\Lambda_\phi(h)} & \mathbf{M}(h\phi) & \xrightarrow{(h^{-1})^*} & \mathbf{M}(\phi)
 \end{array}$$

Figure 13: Composition of the induced action on global sections.

*Temporary Loop-1 image; source locator: Polo-Gille Exp1-13oct24.pdf, component-local p. 42;  
combined-reader p. 52.*

It follows that  $A(h) \circ A(g) = A(hg)$ , and hence:

**Proposition 6.4.1.** *For every  $T$ , the  $\mathbf{O}(X_T)$ -module*

$$\left( \prod_{X/S_0} M \right) (T) = \Gamma(M_T/X_T)$$

*is endowed, functorially in  $T$ , with an action of  $G(T)$  compatible with the action of  $G(T)$  on  $\mathbf{O}(X_T)$ ; that is,*

$$A(g)(\alpha \cdot m) = g(\alpha) \cdot A(g)(m).$$

Since  $g(\alpha) = \alpha$  for  $\alpha \in \mathbf{O}(T)$ , it follows in particular that  $\prod_{X/S_0} M$  is a  $G$ - $\mathbf{O}_{S_0}$ -module.

More generally, let  $Y$  be a second  $G$ -equivariant  $\widehat{\mathcal{C}}$ -object, let  $\phi : Y \rightarrow X$  be a morphism in  $\widehat{\mathcal{C}}$ , and put  $H = \text{Stab}_G(\phi)$ , as in 6.3. Then

$$M_\phi = Y \times_\phi M$$

is an  $H$ -equivariant  $\mathbf{O}_Y$ -module. We obtain:

**Corollary 6.4.2.** *The functor  $\prod_{Y/S_0} M_\phi$  is a  $\text{Stab}_G(\phi)$ - $\mathbf{O}_{S_0}$ -module.*

### 6.5. $G$ -equivariant $\mathcal{O}_X$ -modules

Apply the preceding discussion in the following situation. Let  $S$  be a scheme, let  $G$  be an  $S$ -group scheme acting on an  $S$ -scheme  $X$ , and let  $\mathcal{F}$  be an  $\mathcal{O}_X$ -module, not necessarily quasi-coherent.

**Definition 6.5.1.** We say that  $\mathcal{F}$  is a  $G$ -equivariant  $\mathcal{O}_X$ -module if the  $\mathbf{O}_X$ -module  $M = W(\mathcal{F})$  is  $G$ -equivariant. Thus, for every morphism  $(g, x) : U \rightarrow G \times_S X$ , one is given isomorphisms of  $\mathcal{O}_U$ -modules

$$\Lambda_x(g) : W(x^*(\mathcal{F})) \xrightarrow{\sim} W((gx)^*(\mathcal{F})),$$

functorial in  $U$  and satisfying  $\Lambda_{hx}(g) \circ \Lambda_x(h) = \Lambda_x(gh)$ . Because  $W$  is fully faithful (cf. 4.6.1.1 and 4.6.2), these yield isomorphisms of  $\mathcal{O}_U$ -modules

$$\Lambda_x(g) : x^*(\mathcal{F}) \xrightarrow{\sim} (gx)^*(\mathcal{F}),$$

where  $gx$  denotes  $\lambda \circ (g, x) : U \rightarrow X$ . Applying this to the identity morphism of  $G \times_S X$  gives an isomorphism of  $\mathcal{O}_{G \times_S X}$ -modules

$$(\star) \quad \theta : \mathrm{pr}_X^*(\mathcal{F}) \xrightarrow{\sim} \lambda^*(\mathcal{F})$$

such that the following diagram  $(\star\star)$  of morphisms of sheaves on  $G \times_S G \times_S X$  commutes.

$$\begin{array}{ccc}
 \mathrm{pr}_{2,3}^* \circ \mathrm{pr}_X^*(\mathcal{F}) & \xrightarrow[\sim]{\mathrm{pr}_{2,3}^*(\theta)} & \mathrm{pr}_{2,3}^* \circ \lambda^*(\mathcal{F}) \xlongequal{\quad} (\mathrm{id}_G \times \lambda)^* \circ \mathrm{pr}_X^*(\mathcal{F}) \\
 \parallel & & \downarrow \scriptstyle (\mathrm{id}_G \times \lambda)^*(\theta) \\
 (\mu \times \mathrm{id}_X)^* \circ \mathrm{pr}_X^*(\mathcal{F}) & \xrightarrow[\sim]{(\mu \times \mathrm{id}_X)^*(\theta)} & (\mu \times \mathrm{id}_X)^* \circ \lambda^*(\mathcal{F}) \xlongequal{\quad} (\mathrm{id}_G \times \lambda)^* \circ \lambda^*(\mathcal{F}) .
 \end{array}
 \quad (\star\star)$$

Figure 14: The cocycle condition  $(\star\star)$  for a  $G$ -equivariant  $\mathcal{O}_X$ -module.

Temporary Loop-1 image; source locator: Polo-Gille Exp1-13oct24.pdf, component-local p. 43;  
combined-reader p. 53.

If  $\mathcal{E}$  is a second  $G$ -equivariant  $\mathcal{O}_X$ -module, a morphism  $\phi : \mathcal{E} \rightarrow \mathcal{F}$  of  $\mathcal{O}_X$ -modules is called  $G$ -equivariant if  $W(\phi) : W(\mathcal{E}) \rightarrow W(\mathcal{F})$  is  $G$ -equivariant. With the notation above, this is equivalent to

$$\theta_{\mathcal{F}} \circ \mathrm{pr}_X^*(\phi) = \lambda^*(\phi) \circ \theta_{\mathcal{E}}.$$

**Remark 6.5.2.** If  $f : Y \rightarrow X$  is  $G$ -equivariant, Lemma 6.3.1 shows that  $f^*(\mathcal{F})$  is a  $G$ -equivariant  $\mathcal{O}_Y$ -module.

**Remark 6.5.3.** Suppose in addition that  $\mathcal{F}$  is quasi-coherent. Giving a  $G$ -equivariant  $\mathcal{O}_X$ -module structure on  $M = W(\mathcal{F})$  is then equivalent to giving such a structure on  $V(\mathcal{F})$ , and this in turn is equivalent to giving an action of  $G$  on the vector fibration  $V(\mathcal{F})$ , compatible with the action on  $X$  and linear on the fibers.

Indeed, let

$$\varphi : \lambda^*(\mathcal{F}) \xrightarrow{\sim} \mathrm{pr}_X^*(\mathcal{F})$$

be the inverse of  $\theta$ . For every morphism  $(g, x) : U \rightarrow G \times_S X$ , the isomorphism of  $\mathcal{O}_U$ -modules

$$\varphi_x(g) = \Lambda_x(g)^{-1} = \Lambda_{gx}(g^{-1})$$

from  $(gx)^*(\mathcal{F})$  to  $x^*(\mathcal{F})$  induces an isomorphism of  $\mathcal{O}_U$ -modules

$$V((gx)^*(\mathcal{F})) \xrightarrow{\sim} V(x^*(\mathcal{F})).$$

Denote the latter by  ${}^t\varphi_x(g)$ . Then

$${}^t\varphi_{gx}(h) \circ {}^t\varphi_x(g) = {}^t(\Lambda_{gx}(h) \circ \varphi_x(g))^{-1} = \Lambda_x(hg)^{-1} = {}^t\varphi_x(hg).$$

For every  $X$ -scheme  $f : Y \rightarrow X$ , one has

$$V(\mathcal{F}) \times_f Y \simeq V(f^*(\mathcal{F})) \quad (\text{cf. 4.6.2}).$$

Therefore the isomorphism

$$\begin{aligned} {}^t\varphi : G \times_S V(\mathcal{F}) &= (G \times_S X) \times_{\text{pr}_X} V(\mathcal{F}) = V(\text{pr}_X^*(\mathcal{F})) \\ &\xrightarrow{\sim} V(\lambda^*(\mathcal{F})) = (G \times_S X) \times_\lambda V(\mathcal{F}) \end{aligned}$$

endows  $V(\mathcal{F})$  with a  $G$ -equivariant  $\mathcal{O}_X$ -module structure. Identifying every functor  $V(-)$  with the vector fibration which represents it, and composing  ${}^t\varphi$  with the projection onto  $V(\mathcal{F})$ , gives an action of  $G$  on  $V(\mathcal{F})$ , compatible with the action on  $X$  and linear on the fibers.

This recovers the definition in [GIT, Chapter 1, Section 3], except that Mumford there considers a locally free  $\mathcal{O}_X$ -module  $\mathcal{E}$  of finite rank and defines an action of  $G$  on  $V(\mathcal{E}) = W(\mathcal{E}^\vee)$ . Indeed, the preceding diagram  $(\star\star)$ , with  $\theta$  replaced by  $\varphi$  and the directions of the arrows reversed, is exactly the diagram in *loc. cit.*, Definition 1.6, and the isomorphism  ${}^t\varphi$  above coincides with the isomorphism  $\Phi$  of *loc. cit.*, p. 31.

**Remark 6.5.4.** Consider in particular the case  $X = S$ , with the trivial action of  $G$ . A  $G$ -equivariant  $\mathcal{O}_S$ -module  $\mathcal{F}$  is then the same thing as a  $G$ - $\mathcal{O}_S$ -module (cf. 4.7.1). If  $f : G \rightarrow S$  denotes the morphism which here equals both  $\text{pr}_X$  and  $\lambda$ , the isomorphism

$$(\star) \quad \theta : f^*(\mathcal{F}) \xrightarrow{\sim} f^*(\mathcal{F})$$

is an element of

$$\begin{aligned} \text{Hom}_{\mathcal{O}_G}(f^*(\mathcal{F}), f^*(\mathcal{F})) &= \text{Hom}_{\mathcal{O}_G}(W(f^*(\mathcal{F})), W(f^*(\mathcal{F}))) \\ &= \text{End}_{\mathcal{O}_S}(W(\mathcal{F}))(G). \end{aligned}$$

It is precisely the morphism

$$\rho : G \longrightarrow \text{End}_{\mathcal{O}_S}(W(\mathcal{F}))$$

which defines the action of  $G$  on  $W(\mathcal{F})$ . Moreover,  $\theta$  corresponds by adjunction to the morphism of  $\mathcal{O}_S$ -modules

$$f_*(\theta) \circ \tau : \mathcal{F} \longrightarrow f_*f^*(\mathcal{F}),$$

where  $\tau : \mathcal{F} \rightarrow f_*f^*(\mathcal{F})$  is the unit morphism of the adjunction. This will be used in Exposé VI<sub>B</sub>, 11.10 bis.

## 6.6. The functors $\prod_{X/S} W(\mathcal{F})$ and $W(p_*(\mathcal{F}))$

Let  $S$  be a scheme, let  $G$  be an  $S$ -group scheme acting on  $S$ -schemes  $X$  and  $Y$  through

$$\lambda : G \times_S X \rightarrow X, \quad \mu : G \times_S Y \rightarrow Y,$$

and let  $p : X \rightarrow Y$  be a  $G$ -equivariant morphism. Suppose that  $p : X \rightarrow Y$  and  $\nu_Y : Y \rightarrow S$  are quasi-compact and quasi-separated and that  $\pi : G \rightarrow S$  is flat.

The projection  $\mathrm{pr}_Y : G \times_S Y \rightarrow Y$  is then flat, as is  $\mu$ , since  $\mu$  is the composite of  $\mathrm{pr}_Y$  and the automorphism  $(g, y) \mapsto (g, gy)$ . For a variable  $S$ -scheme  $f : T \rightarrow S$ , denote by  $p_T : X_T \rightarrow T$  and  $f_X : X_T \rightarrow X$  the morphisms obtained from  $p$  and  $f$  by base change.

Let  $\mathcal{F}$  be a quasi-coherent  $G$ -equivariant  $\mathcal{O}_X$ -module, and let

$$\theta : \mathrm{pr}_X^*(\mathcal{F}) \xrightarrow{\sim} \lambda^*(\mathcal{F})$$

be the isomorphism of 6.5.1 (\*). Since  $p$  is quasi-compact and quasi-separated,  $p_*(\mathcal{F})$  is quasi-coherent (cf. EGA I, 9.2.1).

**Lemma 6.6.1.** *The quasi-coherent  $\mathcal{O}_Y$ -module  $p_*(\mathcal{F})$  is  $G$ -equivariant.*

Indeed, one has the following two cartesian squares.

$$\begin{array}{ccccc} G \times_S X & \xrightarrow{\mathrm{pr}_X} & X & \xleftarrow{\lambda} & G \times_S X \\ \downarrow q & & \downarrow p & & \downarrow q \\ G \times_S Y & \xrightarrow{\mathrm{pr}_Y} & Y & \xleftarrow{\mu} & G \times_S Y. \end{array}$$

Figure 15: The two cartesian base-change squares used in Lemma 6.6.1.

*Temporary Loop-1 image; source locator: Polo-Gille Exp1-13oct24.pdf, component-local p. 45;  
combined-reader p. 55.*

Since  $p$  is quasi-compact and quasi-separated and  $\mathrm{pr}_Y$  and  $\mu$  are flat, EGA III, 1.4.15, completed by EGA IV<sub>1</sub>, 1.7.21, gives

$$q_* \mathrm{pr}_X^*(\mathcal{F}) = \mathrm{pr}_Y^* p_*(\mathcal{F}), \quad q_* \lambda^*(\mathcal{F}) = \mu^* p_*(\mathcal{F}).$$

Consequently,  $\theta$  induces an isomorphism

$$\theta_Y : \mathrm{pr}_Y^* p_*(\mathcal{F}) \xrightarrow{\sim} \mu^* p_*(\mathcal{F}).$$

The same argument shows that  $\theta_Y$  satisfies the cocycle condition 6.5.1 (\*\*), because the morphisms  $G \times_S G \times_S Y \rightarrow G \times_S Y$  involved are flat. This proves the lemma.

In particular, take  $Y = S$  with the trivial action of  $G$ . By Remark 6.5.4,  $p_*(\mathcal{F})$  is then a  $G$ - $\mathcal{O}_S$ -module. If, in addition,  $G$  is affine over  $S$ , and if

$$\mathcal{A}(G) = \pi_*(\mathcal{O}_G),$$

then  $p_*(\mathcal{F})$  is an  $\mathcal{A}(G)$ -comodule, by 4.7.2.

On the other hand, by Proposition 6.4.1, the functor  $\prod_{X/S} W(\mathcal{F})$ , which assigns to every  $f : T \rightarrow S$

$$W(f_X^*(\mathcal{F}))(X_T) = \Gamma(X_T, f_X^*(\mathcal{F})) = \Gamma(T, p_{T*} f_X^*(\mathcal{F})),$$

is a  $G$ - $\mathcal{O}_S$ -module. There is moreover a canonical morphism

$$\tau : W(p_*(\mathcal{F})) \longrightarrow \prod_{X/S} W(\mathcal{F}),$$

which, for each  $f : T \rightarrow S$ , is given by the canonical morphism

$$\Gamma(T, f^* p_*(\mathcal{F})) \longrightarrow \Gamma(T, p_{T*} f_X^*(\mathcal{F})).$$

It is an isomorphism when restricted to the full subcategory of schemes  $T$  flat over  $S$ , and it is immediate that  $\tau$  is a morphism of  $G\text{-}\mathcal{O}_S$ -modules. We have therefore proved the following proposition; for part (ii), compare [GIT], p. 32.

**Proposition 6.6.2.** *Let  $S$  be a scheme, let  $G$  be an  $S$ -group scheme acting on an  $S$ -scheme  $X$ , and let  $\mathcal{F}$  be a quasi-coherent  $G$ -equivariant  $\mathcal{O}_X$ -module. Suppose that  $\pi : G \rightarrow S$  is flat and  $p : X \rightarrow S$  is quasi-compact and quasi-separated.*

(i) *Then  $p_*(\mathcal{F})$  is a quasi-coherent  $G\text{-}\mathcal{O}_S$ -module. The canonical morphism*

$$W(p_*(\mathcal{F})) \longrightarrow \prod_{X/S} W(\mathcal{F})$$

*is a morphism of  $G\text{-}\mathcal{O}_S$ -modules, and the two functors coincide on the category of flat  $S$ -schemes.*

(ii) *If, in addition,  $G$  is affine over  $S$ , and if  $\mathcal{A}(G) = \pi_*(\mathcal{O}_G)$ , then  $p_*(\mathcal{F})$  has the structure of an  $\mathcal{A}(G)$ -comodule.<sup>56</sup>*

## 6.7. Stabilizers

Let  $S$  be a scheme, let  $G$  be an  $S$ -group scheme acting on an  $S$ -scheme  $X$ , and let  $\mathcal{F}$  be a quasi-coherent  $G$ -equivariant  $\mathcal{O}_X$ -module. Let  $Y$  be a second  $S$ -scheme with an action of  $G$ , possibly the trivial action, let  $\phi : Y \rightarrow X$  be an  $S$ -morphism, not necessarily  $G$ -equivariant, and let  $\text{Stab}_G(\phi)$  be the stabilizer of  $\phi$  in  $G$  (cf. 6.3.2).

Let us point out at once (see Exposé VI<sub>B</sub>, Section 6) that  $\text{Stab}_G(\phi)$  is representable by a closed subgroup scheme  $H$  of  $G$  if  $X$  is separated over  $S$  and  $Y$  is essentially free over  $S$  (cf. *loc. cit.*, Definition 6.2.1). Indeed, consider

$$r : G \times_S Y \longrightarrow X \times_S X, \quad r(g, y) = (\phi(y), g\phi(g^{-1}y)),$$

put  $P = G \times_S Y$ , and let  $P'$  be the inverse image under  $r$  of the diagonal  $\Delta_{X/S}$ . Then (cf. *loc. cit.*, 6.2.4(a))

$$\text{Stab}_G(\phi) = \prod_{P/G} P'.$$

It follows that  $\text{Stab}_G(\phi)$  is representable by a closed subgroup scheme  $H$  of  $G$  if  $X$  is separated over  $S$  and  $Y$  is essentially free over  $S$ . The latter condition is automatic if  $S$  is the spectrum of a field or if  $Y = S$ . Under these hypotheses,  $\phi^*(\mathcal{F})$  is a quasi-coherent  $H$ -equivariant  $\mathcal{O}_Y$ -module (cf. 6.5.2).

If, in addition,  $\pi : G \rightarrow S$  and  $p : Y \rightarrow S$  are quasi-compact and quasi-separated over  $S$ , and  $\pi$  is flat, then  $p_*\phi^*(\mathcal{F})$  is an  $H\text{-}\mathcal{O}_S$ -module by Proposition 6.6.2. In particular:

**Corollary 6.7.1.** *Let  $S$  be a scheme, let  $G$  be an  $S$ -group scheme acting on an  $S$ -scheme  $X$ , and let  $\mathcal{F}$  be a quasi-coherent  $G$ -equivariant  $\mathcal{O}_X$ -module. Suppose that  $\pi : G \rightarrow S$  is flat and  $X \rightarrow S$  is quasi-compact and separated. Let  $\tau : S \rightarrow X$  be a section of  $X$  over  $S$ .*

(i) *The stabilizer  $H = \text{Stab}_G(\tau)$  is a closed subgroup scheme of  $G$ , and  $\tau^*(\mathcal{F})$  is a quasi-coherent  $H\text{-}\mathcal{O}_S$ -module.*

(ii) *If  $H$  is moreover affine over  $S$ , for example if  $G$  is, and if  $\mathcal{A}(H) = \pi_*(\mathcal{O}_H)$ , then  $\tau^*(\mathcal{F})$  is an  $\mathcal{A}(H)$ -comodule.*

<sup>56</sup>Editor's note: The same holds if  $G$  is flat, quasi-compact, and quasi-separated over  $S$ , and if  $p_*(\mathcal{F})$  is a flat  $\mathcal{O}_S$ -module; cf. Exposé VI<sub>B</sub>, 11.6.1(ii).

### 6.8. $G$ -equivariant sheaves on $G$

To conclude, we give two results, 6.8.1 and 6.8.6, which will be used in Exposés II and III, in particular in III, 4.25.

**Proposition 6.8.1.** *Let  $S$  be a scheme, let  $\pi : G \rightarrow S$  be an  $S$ -group scheme, and let  $\varepsilon : S \rightarrow G$  be the unit section. Consider the action of  $G$  on itself by left translations. The functors*

$$\mathcal{E} \mapsto \pi^*(\mathcal{E}), \quad \mathcal{F} \mapsto \varepsilon^*(\mathcal{F})$$

*induce mutually quasi-inverse equivalences between the category of quasi-coherent  $\mathcal{O}_S$ -modules and the category of quasi-coherent  $G$ -equivariant  $\mathcal{O}_G$ -modules.*

*Proof.* Let  $\mu$  denote multiplication in  $G$ , and let  $\text{pr}_2 : G \times_S G \rightarrow G$  be the second projection. Since  $\pi \circ \mu = \pi \circ \text{pr}_2$ , every quasi-coherent  $\mathcal{O}_S$ -module  $\mathcal{E}$  has a canonical isomorphism

$$\mu^* \pi^*(\mathcal{E}) = \text{pr}_2^* \pi^*(\mathcal{E}).$$

It is immediate that this isomorphism satisfies the cocycle condition 6.5.1 ( $\star\star$ ), so  $\pi^*(\mathcal{E})$  is a  $G$ -equivariant  $\mathcal{O}_G$ -module. Since  $\varepsilon^* \pi^*(\mathcal{E}) = \mathcal{E}$ , the functor  $\mathcal{E} \mapsto \pi^*(\mathcal{E})$  is

fully faithful. It remains to show that, for every  $G$ -equivariant  $\mathcal{O}_G$ -module  $\mathcal{F}$ ,

$$\mathcal{F} \simeq \pi^* \varepsilon^*(\mathcal{F}).$$

By hypothesis, there is an isomorphism

$$\theta : \text{pr}_2^*(\mathcal{F}) \xrightarrow{\sim} \mu^*(\mathcal{F}).$$

Pulling  $\theta$  back along the morphism

$$\tau : G \longrightarrow G \times_S G, \quad \tau = (\text{id}_G, \varepsilon \circ \pi),$$

gives an isomorphism  $\mathcal{F} \xrightarrow{\sim} \pi^* \varepsilon^*(\mathcal{F})$ .

**Remarks 6.8.2.** (a) Consider the action of  $G \times_S G$  on  $G$  defined by

$$(g_1, g_2) \cdot g = g_1 g g_2^{-1}.$$

The stabilizer of the unit section  $\varepsilon : S \rightarrow G$  is the diagonal subgroup  $H$  of  $G \times_S G$ . Consequently, if  $\mathcal{F}$  is a quasi-coherent  $(G \times_S G)$ -equivariant  $\mathcal{O}_G$ -module, then  $\mathcal{E} = \varepsilon^*(\mathcal{F})$  has the structure of an  $H$ - $\mathcal{O}_S$ -module, by 6.5.2.

(b) One can show that  $\mathcal{F} \mapsto \varepsilon^*(\mathcal{F})$  is an equivalence between the category of quasi-coherent  $(G \times_S G)$ -equivariant  $\mathcal{O}_G$ -modules and the category of quasi-coherent  $H$ - $\mathcal{O}_S$ -modules. This is a special case of more general descent results (cf. Exposé IV, Section 2, and SGA 1, VIII); see, for example, [Th87], 1.2–1.3.

**Remark 6.8.3.** Retain the notation of 6.8.1. For every quasi-coherent  $\mathcal{O}_S$ -module  $\mathcal{E}$ , let  $\pi_*^G \pi^*(\mathcal{E})$  be the  $\mathcal{O}_S$ -submodule of  $\pi_* \pi^*(\mathcal{E})$  whose sections over an open subset  $V$  of  $S$  are the elements

$$\gamma \in \Gamma(\pi^{-1}(V), \pi^*(\mathcal{E}))$$

such that  $g \cdot \gamma_{S'} = \gamma_{S'}$  for every  $S' \rightarrow V$  and  $g \in G(S')$ . Then the natural morphism

$$\mathcal{E} \longrightarrow \pi_*^G \pi^*(\mathcal{E})$$

is an isomorphism. This is immediate if

$$\pi_* \pi^*(\mathcal{E}) = \mathcal{E} \otimes_{\mathcal{O}_S} \pi_*(\mathcal{O}_G),$$

for example if  $G \rightarrow S$  is affine, or if  $G \rightarrow S$  is quasi-compact and quasi-separated and  $\mathcal{E}$  is flat; the general case is verified without difficulty.

**Remark 6.8.4.** Let  $S$  be a scheme, let  $H$  be an  $S$ -group scheme acting on an  $S$ -scheme  $X$ , and let  $\mathcal{F}$  be an  $\mathcal{O}_X$ -module. Suppose  $H$  is flat over  $S$ , and let  $W_P(\mathcal{F})$  denote the restriction of  $W(\mathcal{F})$  to the full subcategory of flat  $S$ -schemes. By 6.5.1, endowing  $\mathcal{F}$  with an  $H$ -equivariant module structure is equivalent to giving an isomorphism

$$\theta : \mathrm{pr}_X^*(\mathcal{F}) \longrightarrow \lambda^*(\mathcal{F})$$

of sheaves on  $H \times_S X$  which satisfies the cocycle condition  $(\star\star)$ . Hence, to endow  $\mathcal{F}$  with an  $H$ -equivariant module structure, it suffices to endow  $W_P(\mathcal{F})$  with such a structure.

**Recollection 6.8.5.** Let  $X$  be an  $S$ -scheme and  $Y$  a closed  $S$ -subscheme of  $X$ . Let  $\mathcal{N}_{Y/X}$  denote the conormal sheaf of the immersion  $i : Y \hookrightarrow X$  (cf. EGA IV<sub>4</sub>, 16.1.2). If  $S' \rightarrow S$  is flat and  $i' : Y' \hookrightarrow X'$  is obtained from  $i$  by base change, then *loc. cit.*, 16.2.2(iii), gives

$$\mathcal{N}_{Y'/X'} \otimes_{\mathcal{O}_{Y'}} \mathcal{O}_{Y'} = \mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y'}.$$

**Proposition 6.8.6.** *Let  $S$  be a scheme, let  $X$  be an  $S$ -group scheme, and let  $Y$  be an  $S$ -subgroup scheme of  $X$ . Suppose that  $Y$  is flat over  $S$ . Then the conormal sheaf  $\mathcal{N}_{Y/X}$  is a  $(Y \times_S Y)$ -equivariant  $\mathcal{O}_Y$ -module.*

Indeed,  $H = Y \times_S Y$  is flat over  $S$ . By Remark 6.8.4, it is therefore enough to endow  $W_P(\mathcal{N}_{Y/X})$  with an  $H$ -equivariant module structure. Let  $S'$  be a flat  $S$ -scheme, let  $Y' \hookrightarrow X'$  be the immersion obtained by base change, and put

$$\mathcal{N}' = \mathcal{N}_{Y'/X'} \otimes_{\mathcal{O}_{Y'}} \mathcal{O}_{Y'}.$$

By 6.8.5,  $\mathcal{N}' = \mathcal{N}_{Y'/X'}$ . Every  $h \in H(S')$  induces an automorphism of  $Y'$ , and hence, for every  $y \in Y'(S')$ , isomorphisms

$$\Gamma(S', y^*(\mathcal{N}')) \xrightarrow{\sim} \Gamma(S', y^* h^*(\mathcal{N}'))$$

which endow  $W_P(\mathcal{N}_{Y/X})$  with an  $H$ -equivariant module structure (cf. 6.1).

## Bibliography

57

- [DG70] M. Demazure and P. Gabriel, *Groupes Algébriques*, Masson & North-Holland, 1970.
- [Gr57] A. Grothendieck, "Sur quelques points d'algèbre homologique," *Tôhoku Math. J.* **9** (1957), 119–221.
- [Ja03] J. C. Jantzen, *Representations of Algebraic Groups*, Academic Press, 1987; second edition, American Mathematical Society, 2003.
- [GIT] D. Mumford, *Geometric Invariant Theory*, Springer-Verlag, 1965; second edition with J. Fogarty, 1982; third edition with J. Fogarty and F. Kirwan, 1994.
- [Ni02] N. Nitsure, "Representability of  $\mathrm{GL}_{\mathcal{E}}$ ," *Proc. Indian Acad. Sci.* **112** (2002), no. 4, 539–542.

---

<sup>57</sup>Editor's note: These are additional references cited in this Exposé.



- [Ni04] N. Nitsure, "Representability of Hom implies flatness," *Proc. Indian Acad. Sci.* **114** (2004), no. 1, 7–14.
- [Se68] J.-P. Serre, "Groupes de Grothendieck des schémas en groupes réductifs déployés," *Publ. Math. I.H.É.S.* **34** (1968), 37–52.
- [Th87] R. W. Thomason, "Equivariant resolution, linearization, and Hilbert's fourteenth problem over arbitrary base schemes," *Adv. Math.* **65** (1987), 16–34.

# Exposé II

## Tangent Bundles – Lie Algebras

by *M. Demazure*

We propose in this exposé to construct, in the theory of schemes, the analogues of the tangent bundles and Lie algebras of classical theory. It will, however, be useful not to restrict ourselves to schemes proper, but also to consider certain functors on the category of schemes that are not necessarily representable (for example, the functors Hom, Norm, and so forth). As announced in the preceding exposé (cf. I 1.1), we shall identify a scheme with its associated functor.<sup>0</sup>

On the other hand, the constructions presented below go beyond the setting of scheme theory. They are equally valid, for example, in the theory of analytic spaces with nilpotent elements, apart from a few changes of detail.

Before beginning this construction, we must give a few general definitions that supplement those of I 1.7.

### 1. The functors $\underline{\mathrm{Hom}}_{Z/S}(X, Y)$

We resume the notation of I 1.1. We identify the category  $\mathcal{C}$  with a full subcategory of

$$\widehat{\mathcal{C}} = \mathrm{Hom}(\mathcal{C}^\circ, \mathbf{Set})$$


(in particular, we suppress the underlinings that allowed us to distinguish graphically an object of  $\widehat{\mathcal{C}}$  from an object of  $\mathcal{C}$ ).<sup>1</sup>

Consider the following situation: four objects of  $\widehat{\mathcal{C}}$ , denoted by  $S, X, Y, Z$ , the first of which is in fact an object of  $\mathcal{C}$ , with  $X$  and  $Y$  over  $Z$ , and  $Z$  over  $S$ .

---

<sup>0</sup>Editor's note: version of October 14, 2024.

<sup>1</sup>Editor's note: in this edition these are rendered by bold letters **F, G, V, W**; cf. Exposé I. For functors such as Norm and Centr (cf. 5.2), however, the underlinings of the original have been retained, and one has been added to the functor Lie, in order to distinguish the functor Lie( $G/S$ ) from the  $\Gamma(S, \mathcal{O}_S)$ -Lie algebra  $\mathrm{Lie}(G/S) = \underline{\mathrm{Lie}}(G/S)(S)$ , used, for example, in Exposé VII<sub>A</sub>.

Figure 16: The objects  $X$  and  $Y$  over  $Z$ , and  $Z$  over  $S$ .

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 2;  
combined-reader p. 60.*

**Definition 1.1.**

We define an object of  $\widehat{\mathcal{C}}_S$ , denoted  $\underline{\mathrm{Hom}}_{Z/S}(X, Y)$ , by

$$\underline{\mathrm{Hom}}_{Z/S}(X, Y)(S') = \mathrm{Hom}_{Z_{S'}}(X_{S'}, Y_{S'}) = \mathrm{Hom}_Z(X \times_S S', Y), \quad (1)$$

for every object  $S'$  of  $\mathcal{C}_S$ . One sees at once that  $\underline{\mathrm{Hom}}_{Z/S}(X, Y)$  is simply the subobject of  $\underline{\mathrm{Hom}}_S(X, Y)$  formed by the morphisms compatible with  $p_X$  and  $p_Y$ ; that is, the kernel of the pair of morphisms

$$\underline{\mathrm{Hom}}_S(X, Y) \rightrightarrows \underline{\mathrm{Hom}}_S(X, Z),$$

the first defined by composition with  $p_Y$ , and the second being the constant morphism whose “image” is  $p_X$ .

As in I 1.7, for every object  $T$  of  $\widehat{\mathcal{C}}$  over  $S$ , there is a natural bijection

$$\mathrm{Hom}_S(T, \underline{\mathrm{Hom}}_{Z/S}(X, Y)) \simeq \mathrm{Hom}_Z(X \times_S T, Y). \quad (2)$$

Moreover, by I 1.7.1, if  $E, F$  are objects of  $\widehat{\mathcal{C}}$  over  $Z$ , then

$$\mathrm{Hom}_Z(E, \underline{\mathrm{Hom}}_Z(F, Y)) \simeq \mathrm{Hom}_Z(E \times_Z F, Y) \simeq \mathrm{Hom}_Z(F, \underline{\mathrm{Hom}}_Z(E, Y)).$$

Applying this with  $E = X$  and  $F = Z \times_S T$ , one obtains natural bijections, for every object  $T$  of  $\widehat{\mathcal{C}}_S$ ,

$$\begin{aligned}
 \mathrm{Hom}_S(T, \underline{\mathrm{Hom}}_{Z/S}(X, Y)) &\simeq \mathrm{Hom}_Z(X \times_S T, Y) \\
 &\simeq \begin{cases} \mathrm{Hom}_Z(Z \times_S T, \underline{\mathrm{Hom}}_Z(X, Y)), \\ \mathrm{Hom}_Z(X, \underline{\mathrm{Hom}}_Z(Z \times_S T, Y)). \end{cases}
 \end{aligned} \quad (3)$$

These bijections are functorial in  $T$ , and hence give isomorphisms of  $S$ -functors.<sup>2</sup>

$$\begin{array}{ccc}
 \underline{\mathrm{Hom}}_S(T, \underline{\mathrm{Hom}}_{Z/S}(X, Y)) & \xrightarrow{\sim} & \underline{\mathrm{Hom}}_{Z/S}(X, \underline{\mathrm{Hom}}_Z(Z \times_S T, Y)) \\
 \searrow \simeq & & \nearrow \simeq \\
 & \underline{\mathrm{Hom}}_{Z/S}(X \times_S T, Y) &
 \end{array} \quad (4)$$

Figure 17: The functorial isomorphisms (4).

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 2;  
combined-reader p. 60.*

<sup>2</sup>Editor’s note: points (2), (3), and (4) have been expanded; in particular, (4) will be used in 3.11.

Let us point out two special cases of the definition. If  $Z = S$ , then

$$\underline{\mathrm{Hom}}_{S/S}(X, Y) = \underline{\mathrm{Hom}}_S(X, Y).$$

On the other hand, when  $X = Z$ , we set

$$\prod_{Z/S} Y = \underline{\mathrm{Hom}}_{Z/S}(Z, Y), \quad (5)$$

so that, by definition,

$$\left( \prod_{Z/S} Y \right) (S') = \mathrm{Hom}_Z(Z \times_S S', Y) \simeq \Gamma(Y_{S'}/Z_{S'}).$$

The functor  $\prod_{Z/S} : \widehat{\mathcal{C}}_{/Z} \rightarrow \widehat{\mathcal{C}}_{/S}$  is right adjoint to the base-change functor from  $S$  to  $Z$ : for every  $S$ -functor  $U$ ,

$$\mathrm{Hom}_S \left( U, \prod_{Z/S} Y \right) = \mathrm{Hom}_Z(U \times_S Z, Y).$$

If  $\mathcal{C} = \mathbf{Sch}$  and  $Z$  is an  $S$ -scheme, the functor  $\prod_{Z/S}$  is called the “Weil restriction of scalars.”

<sup>3</sup> One also has an isomorphism

$$\underline{\mathrm{Hom}}_{Z/S}(X, Y) \simeq \underline{\mathrm{Hom}}_{X/S}(X, Y \times_Z X) = \prod_{X/S} (Y \times_Z X), \quad (6)$$

which gives in particular, when  $Z = S$ , an isomorphism

$$\underline{\mathrm{Hom}}_S(X, Y) \simeq \prod_{X/S} Y_X. \quad (7)$$

### Remark 1.2.

The functor  $Y \mapsto \underline{\mathrm{Hom}}_{Z/S}(X, Y)$  commutes with products in the following sense: there is a functorial isomorphism

$$\underline{\mathrm{Hom}}_{Z/S}(X, Y \times_Z Y') \simeq \underline{\mathrm{Hom}}_{Z/S}(X, Y) \times_S \underline{\mathrm{Hom}}_{Z/S}(X, Y'). \quad (*)$$

It follows that if  $Y$  is a  $Z$ -group, respectively a  $Z$ -ring, and so on, then  $\underline{\mathrm{Hom}}_{Z/S}(X, Y)$  is an  $S$ -group, respectively an  $S$ -ring, and so on.

### Remark 1.3.

Let  $\pi : M \rightarrow Y$  be a  $Y$ -functor in  $\mathcal{O}_Y$ -modules (cf. I 4.3.3.1), and set  $H = \underline{\mathrm{Hom}}_{Z/S}(X, Y)$ . Then  $\underline{\mathrm{Hom}}_{Z/S}(X, M)$  has a natural  $\mathcal{O}_H$ -module structure; more precisely, for every  $H'$  over  $H$ , the set

$$\mathrm{Hom}_H(H', \underline{\mathrm{Hom}}_{Z/S}(X, M))$$

has a natural  $\mathcal{O}(X \times_S H')$ -module structure. <sup>4</sup>

Indeed, let  $m : M \times_Y M \rightarrow M$  and  $\lambda : \mathcal{O}_Y \times_Y M \rightarrow M$  denote the morphisms defining the structures of a  $Y$ -abelian group and an  $\mathcal{O}_Y$ -module. Let  $H'$  be an  $S$ -scheme over

<sup>3</sup>Editor’s note: the preceding two sentences have been added.

<sup>4</sup>Editor’s note: this remark has been added; it is useful for Corollary 3.11.1.

$H = \underline{\mathrm{Hom}}_{Z/S}(X, Y)$ ; equivalently, one has been given a  $Z$ -morphism  $f : X \times_S H' \rightarrow Y$ , which makes  $X \times_S H'$  a  $Y$ -object. Then

$$\mathrm{Hom}_H(H', \underline{\mathrm{Hom}}_{Z/S}(X, M))$$

is the set of  $Z$ -morphisms  $\phi : X \times_S H' \rightarrow M$  such that  $\pi \circ \phi = f$ , that is, the set of  $Y$ -morphisms  $X \times_S H' \rightarrow M$ .

If  $\phi, \psi$  are two such morphisms, define  $\phi + \psi$  to be the composite  $Y$ -morphism

$$X \times_S H' \xrightarrow{\phi \times \psi} M \times_Y M \xrightarrow{m} M.$$

This makes  $\underline{\mathrm{Hom}}_{Z/S}(X, M)$  an abelian group over  $H = \underline{\mathrm{Hom}}_{Z/S}(X, Y)$ .<sup>5</sup>

Similarly, if  $a \in \mathcal{O}(X \times_S H')$ , that is, if  $a : X \times_S H' \rightarrow \mathcal{O}_S$  is an  $S$ -morphism, define  $a\phi$  to be the composite  $\lambda \circ (a \times \phi)$ , where  $a \times \phi$  is the  $Y$ -morphism from  $X \times_S H'$  to  $\mathcal{O}_Y \times_Y M \simeq \mathcal{O}_S \times_S M$  with components  $a$  and  $\phi$ . This gives  $\mathrm{Hom}_H(H', \underline{\mathrm{Hom}}_{Z/S}(X, M))$  an  $\mathcal{O}(X \times_S H')$ -module structure, functorial in the  $H$ -object  $H'$ .

## 2. The schemes $I_S(\mathcal{M})$

### Definition 2.1.

Let  $S$  be a scheme and  $\mathcal{M}$  a quasi-coherent  $\mathcal{O}_S$ -module. Denote by  $\mathcal{D}_{\mathcal{O}_S}(\mathcal{M})$  the quasi-coherent  $\mathcal{O}_S$ -algebra  $\mathcal{O}_S \oplus \mathcal{M}$ , where  $\mathcal{M}$  is regarded as an ideal of square zero. Denote by

$$I_S(\mathcal{M}) = \mathrm{Spec} \mathcal{D}_{\mathcal{O}_S}(\mathcal{M})$$

the corresponding  $S$ -scheme.<sup>6</sup> In particular, put

$$\mathcal{D}_{\mathcal{O}_S} = \mathcal{D}_{\mathcal{O}_S}(\mathcal{O}_S), \quad I_S = I_S(\mathcal{O}_S),$$

and call these, respectively, the algebra of dual numbers over  $S$  and the scheme of dual numbers over  $S$ .

The assignment  $\mathcal{M} \mapsto I_S(\mathcal{M})$  is a contravariant functor from quasi-coherent  $\mathcal{O}_S$ -modules to  $S$ -schemes. In particular, the morphisms  $0 \rightarrow \mathcal{M}$  and  $\mathcal{M} \rightarrow 0$  define, respectively, the structural morphism

$$\rho : I_S(\mathcal{M}) \longrightarrow I_S(0) = S$$

and a section  $\varepsilon_{\mathcal{M}}$  of it, called the *zero section*.<sup>7</sup>

#### 2.1.1.

Since  $\mathcal{M} \mapsto I_S(\mathcal{M})$  is contravariant, every  $a \in \mathrm{End}_{\mathcal{O}_S}(\mathcal{M})$  defines an  $S$ -endomorphism  $a^*$  of  $I_S(\mathcal{M})$ , and

$$1^* = \mathrm{id}, \quad (ab)^* = b^* \circ a^*, \quad 0^* = \varepsilon_{\mathcal{M}} \circ \rho, \quad a^* \circ \varepsilon_{\mathcal{M}} = \varepsilon_{\mathcal{M}}.$$

Consequently,  $I_S(\mathcal{M})$  carries a right action of the multiplicative monoid of  $\mathrm{End}_{\mathcal{O}_S}(\mathcal{M})$ , commuting with the  $S$ -morphisms  $I_S(\mathcal{M}) \rightarrow I_S(\mathcal{M}')$  arising from morphisms  $\mathcal{M}' \rightarrow \mathcal{M}$ ; in particular, the operators  $a^*$  preserve the zero section.<sup>8</sup>

<sup>5</sup>Editor's note: if  $M$  is a  $Y$ -group, not necessarily abelian, and one sets  $\phi \cdot \psi = m \circ (\phi \times \psi)$ , this makes  $\underline{\mathrm{Hom}}_{Z/S}(X, M)$  a group over  $\underline{\mathrm{Hom}}_{Z/S}(X, Y)$ .

<sup>6</sup>Editor's note: the underlying space of  $I_S(\mathcal{M})$  is the same as that of  $S$ .

<sup>7</sup>Editor's note: compare 3.1.1.

<sup>8</sup>Editor's note: what follows expands the original; the numbering 2.1.1–2.1.3 has also been added.

For every  $a \in \text{End}_{\mathcal{O}_S}(\mathcal{M})$ , every  $f : S' \rightarrow S$ , and every  $m \in I_S(\mathcal{M})(S')$ , write  $m \cdot a = a^*(m)$ . Then

$$m \cdot 1 = m, \quad (m \cdot a) \cdot b = m \cdot (ab), \quad m \cdot 0 = \varepsilon_{\mathcal{M}}(\rho(m)),$$

and if  $m = \varepsilon_{\mathcal{M}}(f)$ , then  $m \cdot a = m$ .<sup>9</sup>

### Remark 2.1.2.

Formation of  $I_S(\mathcal{M})$  commutes with base extension: there are canonical isomorphisms

$$I_S(\mathcal{M})_{S'} \simeq I_{S'}(\mathcal{M} \otimes_{\mathcal{O}_S} \mathcal{O}_{S'}).$$

For simplicity write  $I_{S'}(\mathcal{M}) = I_S(\mathcal{M})_{S'}$ . More generally, if  $X$  is an  $S$ -functor, not necessarily representable, write  $I_X(\mathcal{M}) = I_S(\mathcal{M}) \times_S X$ .

### 2.1.3.

By the preceding discussion, the multiplicative monoid of  $\mathcal{O}(S')$  acts on the  $S'$ -scheme  $I_{S'}(\mathcal{M})$ , functorially in  $\mathcal{M}$ . In other words, the  $S$ -scheme  $I_S(\mathcal{M})$  is an object with an operator-monoid structure  $\mathcal{O}_S$ , functorial in  $\mathcal{M}$ . Thus there is a morphism of  $S$ -schemes

$$\lambda : I_S(\mathcal{M}) \times_S \mathcal{O}_S \longrightarrow I_S(\mathcal{M}),$$

satisfying the evident conditions. For every  $S$ -functor  $X$ , base change gives a morphism of  $X$ -functors

$$\lambda_X : I_X(\mathcal{M}) \times_S \mathcal{O}_S \longrightarrow I_X(\mathcal{M}),$$

making  $I_X(\mathcal{M})$  an object with operator monoid  $\mathcal{O}(X)$ . Every  $a \in \mathcal{O}(X) = \text{Hom}_S(X, \mathcal{O}_S)$  defines the  $X$ -endomorphism

$$a^* = \lambda_X \circ (\text{id}_{I_X(\mathcal{M})} \times (a \circ \text{pr}_X)).$$

Explicitly, if  $x \in X(S')$  and  $m \in I_S(\mathcal{M})(S') = I_{S'}(\mathcal{M})(S')$ , then  $a(x) = a \circ x \in \mathcal{O}(S')$  and

$$(m, x) \cdot a = (m \cdot a(x), x).$$

This operation is functorial in  $\mathcal{M}$  and preserves the zero section  $\varepsilon_{\mathcal{M}} : X \rightarrow I_X(\mathcal{M})$ ; thus  $a^* \circ \varepsilon_{\mathcal{M}} = \varepsilon_{\mathcal{M}}$ .<sup>10</sup>

The operation is also “functorial in  $X$ ” in the following sense. If  $\pi : Y \rightarrow X$  is a morphism of  $S$ -functors and  $u : \mathcal{O}(X) \rightarrow \mathcal{O}(Y)$  the corresponding ring morphism, so that  $u(a) = a \circ \pi$ , then the following diagram is commutative.

$$\begin{array}{ccc} I_X(\mathcal{M}) & \xrightarrow{a^*} & I_X(\mathcal{M}) \\ \pi \uparrow & & \uparrow \pi \\ I_Y(\mathcal{M}) & \xrightarrow{u(a)^*} & I_Y(\mathcal{M}) \end{array} \quad .$$

Figure 18: Functoriality of the operator action under  $Y \rightarrow X$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp2-14oct24.pdf, component-local p. 5;  
combined-reader p. 63.*

<sup>9</sup>Editor’s note: later we shall be chiefly interested in  $a \in \mathcal{O}(S)$ , acting by homotheties on  $\mathcal{M}$ . For example, if  $\mathcal{M}$  is free of rank  $r$ , then for every  $S' \rightarrow S$ ,  $\text{Hom}_S(S', I_S(\mathcal{M}))$  identifies with the set of  $r$ -tuples  $(\epsilon_1, \dots, \epsilon_r) \in \mathcal{O}(S')^r$  satisfying  $\epsilon_i \epsilon_j = 0$  for all  $i, j$ , and  $(\epsilon_1, \dots, \epsilon_r) \cdot a = (a\epsilon_1, \dots, a\epsilon_r)$  for every  $a \in \mathcal{O}(S)$ .

<sup>10</sup>Editor’s note: this paragraph has been added; it will be useful in 3.4.2 and 4.6.2.

**2.2.**

Let  $\mathcal{M}$  and  $\mathcal{N}$  be two quasi-coherent  $\mathcal{O}_S$ -modules. The commutative diagram below induces the corresponding diagram of  $S$ -schemes on the following page.




Figure 19: The direct-sum diagram of  $\mathcal{O}_S$ -modules.

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 5;  
combined-reader p. 63.*




Figure 20: The induced commutative diagram of  $S$ -schemes.

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 6;  
combined-reader p. 64.*

**Proposition 2.2.**

For every  $S$ -scheme  $X$ , the diagram of functors over  $S$  obtained by applying  $\underline{\mathrm{Hom}}_S(-, X)$  to the preceding diagram is cartesian.





Figure 21: The cartesian diagram of  $\underline{\mathrm{Hom}}_S$ -functors.

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 6;  
combined-reader p. 64.*

Indeed, for every  $S' \rightarrow S$ , one must verify that the diagram of sets obtained by evaluating the functors on  $S'$  is cartesian. Since formation of  $I_S(\mathcal{P})$  commutes with base extension, it is enough to take  $S' = S$ , and hence to verify that the following diagram of sets is cartesian.

Figure 22: The cartesian diagram after evaluation on  $S$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp2-14oct24.pdf, component-local p. 6;  
combined-reader p. 64.*

If  $x \in X(S)$ , it follows from SGA 1, III 5.1 that  $X(\varepsilon_{\mathcal{M}})^{-1}(x)$  is isomorphic, functorially in  $\mathcal{M}$ , to

$$\mathrm{Hom}_{\mathcal{O}_S}(x^*(\Omega_{X/S}^1), \mathcal{M}),$$

where  $\Omega_{X/S}^1$  denotes the sheaf of relative differentials of  $X$  over  $S$ . This latter functor in  $\mathcal{M}$  plainly carries a direct sum of  $\mathcal{O}_S$ -modules to the product of the corresponding sets, which proves the proposition.<sup>11</sup>

### Corollary 2.2.1.

Let  $X$  be an  $S$ -scheme and  $\mathcal{M}$  a finite free  $\mathcal{O}_S$ -module. The functor  $\underline{\mathrm{Hom}}_S(I_S(\mathcal{M}), X)$ , as a functor over  $X$ , is isomorphic to a finite product over  $X$  of copies of  $\underline{\mathrm{Hom}}_S(I_S, X)$ .

### Nota 2.2.2.

It follows from the proof of the proposition that  $\underline{\mathrm{Hom}}_S(I_S, X)$ , as an  $X$ -functor, is isomorphic to  $\mathbf{V}(\Omega_{X/S}^1)$  (I 4.6.1), and is therefore represented by the vector fibration<sup>12</sup>

$$V(\Omega_{X/S}^1).$$

## 3. The tangent bundle, condition (E)

Throughout this section, unless otherwise stated, the letters  $\mathcal{M}, \mathcal{M}', \mathcal{N}, \dots$  will denote finite free  $\mathcal{O}_S$ -modules (that is, modules isomorphic to finite direct sums of copies of  $\mathcal{O}_S$ ).

We shall systematically use the identifications established in Exposé I. Thus, by a “functor over  $S$ ” we shall mean indifferently either a functor equipped with a morphism to  $S = h_S$ , or a functor on the category of objects over  $S$ . We shall use “group functor over  $S$ ,” and similar expressions, in the same way.

### Definition 3.1.

Let  $S$  be a scheme, let  $\mathcal{M}$  be a finite free  $\mathcal{O}_S$ -module, and let  $X$  be a functor over  $S$ . The *tangent bundle of  $X$  over  $S$ , relative to the  $\mathcal{O}_S$ -module  $\mathcal{M}$* , is the  $S$ -functor

$$T_{X/S}(\mathcal{M}) = \underline{\mathrm{Hom}}_S(I_S(\mathcal{M}), X).$$

<sup>11</sup>Editor’s note: see also the addition 0.1.8 in Exposé III.

<sup>12</sup>Editor’s note: cf. editor’s note (33) of Exposé I.



In particular, the *tangent bundle of  $X$  over  $S$*  is <sup>13</sup>

$$T_{X/S} = T_{X/S}(\mathcal{O}_S) = \underline{\mathrm{Hom}}_S(I_S, X).$$

The assignment  $\mathcal{M} \mapsto T_{X/S}(\mathcal{M})$  is a covariant functor from finite free  $\mathcal{O}_S$ -modules to  $S$ -functors. In particular, the morphisms  $\mathcal{M} \rightarrow 0$  and  $0 \rightarrow \mathcal{M}$  define, respectively, an  $S$ -morphism

$$\pi_{\mathcal{M}} : T_{X/S}(\mathcal{M}) \longrightarrow T_{X/S}(0) \simeq X$$

and a section  $\tau_0$  of it, called the *zero section*. <sup>14</sup>

### Remark 3.1.1.

<sup>15</sup> The projection  $\pi_{\mathcal{M}} : T_{X/S}(\mathcal{M}) \rightarrow X$  is induced by the zero section  $\varepsilon_{\mathcal{M}} : S \rightarrow I_S(\mathcal{M})$ , whereas the zero section  $\tau_0 : X \rightarrow T_{X/S}(\mathcal{M})$  is induced by the structural morphism  $\rho : I_S(\mathcal{M}) \rightarrow S$ . More explicitly, let  $t \in T_{X/S}(\mathcal{M})(S')$  correspond to the  $S$ -morphism  $f : S' \times_S I_S(\mathcal{M}) \rightarrow X$ , and let  $x \in X(S')$  correspond to  $g : S' \rightarrow X$ . Then

$$\pi(t) = f \circ (\mathrm{id}_{S'} \times \varepsilon_{\mathcal{M}}), \quad \tau_0(x) = g \circ (\mathrm{id}_{S'} \times \rho).$$

It follows from 3.1 that  $\mathcal{M} \mapsto T_{X/S}(\mathcal{M})$  is a covariant functor from finite free  $\mathcal{O}_S$ -modules to functors over  $X$ . In particular,  $\mathcal{O}(S)$  is an operator monoid of the  $X$ -functor  $T_{X/S}(\mathcal{M})$ , compatible with functoriality in  $\mathcal{M}$ .

### Scholium 3.1.2.

<sup>15</sup> The preceding discussion means, in particular, the following. For every  $S$ -morphism  $\varphi : X' \rightarrow X$ , put

$$\Sigma(X', \mathcal{M}) = \mathrm{Hom}_X(X', T_{X/S}(\mathcal{M})).$$

The multiplicative monoid  $\mathrm{End}_{\mathcal{O}_S}(\mathcal{M})$  acts on  $\Sigma(X', \mathcal{M})$ ; write the action as  $(\lambda, x) \mapsto \lambda * x$ . Then

$$\lambda * (\mu * x) = (\lambda\mu) * x, \quad 1 * x = x, \quad 0 * x = \tau_0 \circ \varphi.$$

There is likewise an action of  $\mathrm{End}_{\mathcal{O}_S}(\mathcal{M} \oplus \mathcal{M})$  on  $\Sigma(X', \mathcal{M} \oplus \mathcal{M})$ . <sup>16</sup>

Let  $m : \mathcal{M} \oplus \mathcal{M} \rightarrow \mathcal{M}$  be addition, and let  $\delta : \mathcal{M} \rightarrow \mathcal{M} \oplus \mathcal{M}$  be the diagonal. Write

$$m_{X'} : \Sigma(X', \mathcal{M} \oplus \mathcal{M}) \rightarrow \Sigma(X', \mathcal{M}), \quad \delta_{X'} : \Sigma(X', \mathcal{M}) \rightarrow \Sigma(X', \mathcal{M} \oplus \mathcal{M})$$

for the induced maps. For  $\lambda, \mu \in \mathcal{O}(S)$ , let  $\ell_{\lambda}$  denote multiplication by  $\lambda$  on  $\mathcal{M}$ , and let  $\ell_{\lambda, \mu}$  denote multiplication by  $(\lambda, \mu)$  on  $\mathcal{M} \oplus \mathcal{M}$ . Since

$$m \circ \ell_{\lambda, \lambda} = \ell_{\lambda} \circ m, \quad m \circ \ell_{\lambda, \mu} \circ \delta = \ell_{\lambda + \mu},$$

we have, for every  $z \in \Sigma(X', \mathcal{M} \oplus \mathcal{M})$  and  $x \in \Sigma(X', \mathcal{M})$ ,

$$\lambda * m(z) = m((\lambda, \lambda) * z), \quad m((\lambda, \mu) * \delta(x)) = (\lambda + \mu) * x. \quad (\dagger)$$

<sup>13</sup>Editor's note: when  $S = \mathrm{Spec}(k)$  and  $X$  is a  $k$ -scheme, one has  $T_{X/S}(S') = \mathrm{Hom}_S(S' \otimes_k k[\epsilon], X) = X(S' \otimes_k k[\epsilon])$ ; this recovers one of the usual definitions of the tangent bundle.

<sup>14</sup>Editor's note: the original order " $0 \rightarrow \mathcal{M}$  and  $\mathcal{M} \rightarrow 0$ " has been corrected to " $\mathcal{M} \rightarrow 0$  and  $0 \rightarrow \mathcal{M}$ ."

<sup>15</sup>Editor's note: paragraphs 3.1.1 and 3.1.2 have been added.

<sup>16</sup>Editor's note: in fact, only the actions of  $\mathcal{O}(S)$  (respectively  $\mathcal{O}(S) \times \mathcal{O}(S)$ ) on  $\Sigma(X', \mathcal{M})$  (respectively  $\Sigma(X', \mathcal{M} \oplus \mathcal{M})$ ) will be used; see below and the proof of 3.6.

**Definition 3.2.**

Let  $u \in X(S) = \text{Hom}_S(S, X) = \Gamma(X/S)$ . The *tangent space of  $X$  over  $S$  at  $u$ , relative to  $\mathcal{M}$* , denoted  $L_{X/S}^u(\mathcal{M})$ , is the  $S$ -functor obtained from the  $X$ -functor  $T_{X/S}(\mathcal{M})$  by pullback along  $u : S \rightarrow X$ .

$$\begin{array}{ccc} L_{X/S}^u(\mathcal{M}) & \longrightarrow & T_{X/S}(\mathcal{M}) \\ \downarrow & & \downarrow \pi \\ S & \xrightarrow{u} & X \end{array}$$

Figure 23: The cartesian square defining  $L_{X/S}^u(\mathcal{M})$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp2-14oct24.pdf, component-local p. 8;  
combined-reader p. 66.*

In particular,  $L_{X/S}^u(\mathcal{O}_S)$  is denoted  $L_{X/S}^u$  and is called the *tangent space of  $X$  over  $S$  at  $u$* .

**Remark 3.2.1.**

<sup>17</sup> It follows from 3.1.1 that, for every  $t : S' \rightarrow S$ ,  $L_{X/S}^u(\mathcal{M})(S')$  is the set of  $S$ -morphisms  $f : I_{S'}(\mathcal{M}) \rightarrow X$  such that  $f \circ \varepsilon_{\mathcal{M}} = u \circ t$ , where  $\varepsilon_{\mathcal{M}} : S' \rightarrow I_{S'}(\mathcal{M})$  is the zero section.

**Proposition 3.3.**

If  $X$  is represented by an  $S$ -scheme  $\tilde{X}$ , then  $T_{X/S}(\mathcal{M})$  and  $L_{X/S}^u(\mathcal{M})$  are representable. In particular,  $T_{X/S}$  and  $L_{X/S}^u$  are represented by the vector fibrations

$$V(\Omega_{\tilde{X}/S}^1) \quad \text{and} \quad V(u^* \Omega_{\tilde{X}/S}^1)$$

over  $\tilde{X}$  and  $S$ , respectively.

It plainly suffices to prove the assertion for  $T_{X/S}(\mathcal{M})$ , since the corresponding assertion for  $L_{X/S}^u(\mathcal{M})$  follows by pullback. By Corollary 2.2.1, it even suffices to treat  $T_{X/S}$ , in which case the result is exactly Nota 2.2.2.

**Remark 3.3.1.**

The proposition gives a particularly simple description of the vector fibration representing  $L_{X/S}^u$ . The image of the section  $u$  of  $X$  over  $S$  is locally closed,<sup>18</sup> and hence is defined by a quasi-coherent ideal  $\mathfrak{m}$  in the scheme induced on an open subscheme of  $X$ . The quotient  $\mathfrak{m}/\mathfrak{m}^2$  may be regarded as a quasi-coherent module on  $S$ ; it is this module that defines the vector fibration in question.

For example, let  $X$  be an algebraic scheme over a field  $k$ , and let  $u$  be a  $k$ -rational point of  $X$ . Let  $\mathfrak{m}$  be the maximal ideal of the local ring  $\mathcal{O}_{X,u}$ , and let  $\mathfrak{t}$  be the  $k$ -vector space dual to  $\mathfrak{m}/\mathfrak{m}^2$ ; this is the Zariski tangent space of  $\mathcal{O}_{X,u}$  at  $u$ . In the notation of I 4.6.5.1,

$$L_{X/k}^u = V(\mathfrak{m}/\mathfrak{m}^2) = W(\mathfrak{t}).$$

<sup>17</sup>Editor's note: this remark has been added.

<sup>18</sup>Editor's note: cf. EGA I, 5.3.11.

Returning to the general situation, observe first that  $L_{X/S}^u(\mathcal{M})$  is covariant from finite free  $\mathcal{O}_S$ -modules to functors over  $S$ . In particular,  $\mathcal{O}(S)$  is a set of operators of the  $S$ -functor  $L_{X/S}^u(\mathcal{M})$ , compatible with functoriality in  $\mathcal{M}$ .<sup>19</sup>

**Proposition 3.4.**

Formation of  $T_{X/S}(\mathcal{M})$  and  $L_{X/S}^u(\mathcal{M})$  commutes with base extension. For every  $S$ -scheme  $S'$ , there are isomorphisms, functorial in  $\mathcal{M}$ ,

$$T_{X_{S'}/S'}(\mathcal{M} \otimes_{\mathcal{O}_S} \mathcal{O}_{S'}) \xrightarrow{\sim} T_{X/S}(\mathcal{M})_{S'},$$

$$L_{X_{S'}/S'}^{u'}(\mathcal{M} \otimes_{\mathcal{O}_S} \mathcal{O}_{S'}) \xrightarrow{\sim} L_{X/S}^u(\mathcal{M})_{S'}, \quad u' = u_{S'}.$$

This follows immediately from the compatibility of Hom with base extension.

**Corollary 3.4.1.**

The  $X$ -functor  $T_{X/S}(\mathcal{M})$ , respectively the  $S$ -functor  $L_{X/S}^u(\mathcal{M})$ , is naturally an object with operators  $\mathcal{O}_X$ , respectively  $\mathcal{O}_S$ , and these structures are functorial in  $\mathcal{M}$ .

For  $L_{X/S}^u(\mathcal{M})$ , this is immediate. For each  $S'$  over  $S$ , the ring  $\mathcal{O}(S')$  acts on  $\mathcal{M} \otimes \mathcal{O}_{S'}$ , hence on

$$L_{X_{S'}/S'}^{u'}(\mathcal{M} \otimes \mathcal{O}_{S'}) = L_{X/S}^u(\mathcal{M})_{S'};$$

this action is functorial in  $S'$ .

The case of  $T_{X/S}(\mathcal{M})$  requires a little more detail. For an  $X$ -object  $f : X' \rightarrow X$ , put  $T_{X/S}(\mathcal{M})_{X'} = T_{X/S}(\mathcal{M}) \times_X X'$ . We must equip

$$T_{X/S}(\mathcal{M})_{X'}(X') = \mathrm{Hom}_X(X', T_{X/S}(\mathcal{M}))$$

with an action of  $\mathcal{O}(X')$ , functorially in  $X'$ . Let  $X_{X'} = X \times_S X'$ , and let  $f'$  be the section of  $X_{X'}$  over  $X'$  defined by  $f$ . The following diagram provides the required identification.




Figure 24: The base-change diagram used to define the  $\mathcal{O}_X$ -operator structure.

*Temporary Loop-1 image; source locator: Polo-Gille Exp2-14oct24.pdf, component-local p. 10;  
combined-reader p. 68.*

<sup>19</sup>Editor's note: cf. 3.1.2.

Together with 3.2.1, the diagram identifies  $T_{X/S}(\mathcal{M})_{X'}(X')$  with

$$L_{X_{X'}/X'}^{f'}(\mathcal{M})(X') = \{ X'\text{-morphisms } \psi : I_{X'}(\mathcal{M}) \rightarrow X_{X'} \mid \psi \circ \varepsilon_{\mathcal{M}} = f' \}.$$

Every  $a \in \mathcal{O}(X')$  acts through its action on  $I_{X'}(\mathcal{M})$ : in the notation of 2.1.1,

$$a\psi = \psi \circ a^*, \quad (a\psi)(x) = \psi(x \cdot a)$$

for every  $X'' \rightarrow X'$  and every  $x \in I_{X'}(\mathcal{M})(X'')$ . This construction is plainly functorial in  $X'$ .<sup>20</sup> The isomorphisms of Proposition 3.4 are therefore, by construction, isomorphisms for the corresponding  $\mathcal{O}_{X_{S'}}$ - and  $\mathcal{O}_{S'}$ -operator structures.

**Remark 3.4.2.**

<sup>21</sup> The action of  $\mathcal{O}_X$  on  $T_{X/S}(\mathcal{M})$  can be seen more directly as follows. For every  $f : X' \rightarrow X$ ,

$$\mathrm{Hom}_X(X', T_{X/S}(\mathcal{M})) = \{ \varphi \in \mathrm{Hom}_S(I_{X'}(\mathcal{M}), X) \mid \varphi \circ \varepsilon_{\mathcal{M}} = f \}.$$

By 2.1.3, the  $S$ -functor  $I_{X'}(\mathcal{M})$  carries an action of the monoid  $\mathcal{O}(X')$  preserving the zero section. Thus, if  $a^*$  denotes the  $X'$ -endomorphism defined by  $a \in \mathcal{O}(X')$ , then

$$a\varphi = \varphi \circ a^*, \quad (a\varphi)(m, x') = \varphi(m \cdot a(x'), x')$$

for every  $S' \rightarrow S$  and every  $(m, x') \in \mathrm{Hom}_S(S', I_S(\mathcal{M}) \times_S X')$ . Notice that  $a^* \circ \varepsilon_{\mathcal{M}} = \varepsilon_{\mathcal{M}}$ , so that  $(a\varphi) \circ \varepsilon_{\mathcal{M}} = f$ .

Likewise, for  $t : S' \rightarrow S$ , the set  $L_{X/S}^u(\mathcal{M})(S')$  consists of the  $S$ -morphisms  $\varphi : I_{S'}(\mathcal{M}) \rightarrow X$  satisfying  $\varphi \circ \varepsilon_{\mathcal{M}} = u \circ t$ , and  $a\varphi = \varphi \circ a^*$  for  $a \in \mathcal{O}(S')$ .

**Remark 3.4.3.**

<sup>21</sup> The observations of 3.1.2 apply equally to the action of  $\mathcal{O}_S$  on  $L_{X/S}^u(\mathcal{M})$  and to that of  $\mathcal{O}_X$  on  $T_{X/S}(\mathcal{M})$ .

**Definition 3.5.**

Let  $S$  be a scheme and  $X$  an  $S$ -functor. We say that  $X$  *satisfies condition (E) relative to  $S$*  if, for every  $S'$  over  $S$  and every pair of finite free  $\mathcal{O}_{S'}$ -modules  $\mathcal{M}, \mathcal{N}$ , the diagram of sets obtained by applying  $X$  to diagram (\*) of 2.1 is cartesian.<sup>22</sup>

$$\begin{array}{ccc} & X(I_{S'}(\mathcal{M} \oplus \mathcal{N})) & \\ \swarrow & & \searrow \\ X(I_{S'}(\mathcal{M})) & & X(I_{S'}(\mathcal{N})) \\ \searrow & & \swarrow \\ & X(S') & \end{array},$$

Figure 25: The cartesian diagram in condition (E).

*Temporary Loop-1 image; source locator: Polo-Gille Exp2-14oct24.pdf, component-local p. 11;  
combined-reader p. 69.*

<sup>20</sup>Editor's note: the original has been expanded in what follows.

<sup>21</sup>Editor's note: Remarks 3.4.2 and 3.4.3 have been added.

<sup>22</sup>Editor's note: for examples of  $S$ -functors and  $S$ -groups that do not satisfy (E), see 6.2 and 6.3.

**3.5.1.**

<sup>23</sup> Equivalently, the functor  $\mathcal{M} \mapsto T_{X/S}(\mathcal{M})$  carries direct sums of finite free  $\mathcal{O}_S$ -modules to products of  $X$ -functors. In that case, the same is true of

$$\mathcal{M} \mapsto L_{X/S}^u(\mathcal{M}) = S \times_X T_{X/S}(\mathcal{M})$$

for every  $u \in \Gamma(X/S)$ .

Proposition 2.2 shows that every representable functor satisfies condition (E). We shall sometimes abbreviate “ $X$  satisfies condition (E) relative to  $S$ ” to “ $X/S$  satisfies condition (E).”

If  $X/S$  satisfies (E), then  $\mathcal{M} \mapsto T_{X/S}(\mathcal{M})$  commutes with products and hence carries groups to groups. In particular,  $T_{X/S}(\mathcal{M})$  is a commutative  $X$ -group; for the same reason,  $L_{X/S}^u(\mathcal{M})$  is a commutative  $S$ -group.

**Proposition 3.6.**

If  $X/S$  satisfies (E), then the abelian-group structure on  $T_{X/S}(\mathcal{M})$ , respectively on  $L_{X/S}^u(\mathcal{M})$ , together with the action of  $\mathcal{O}_X$ , respectively of  $\mathcal{O}_S$ , makes it an  $\mathcal{O}_X$ -module, respectively an  $\mathcal{O}_S$ -module.

The operator action is functorial in  $\mathcal{M}$ , and therefore respects the abelian-group structure obtained by functoriality from that of  $\mathcal{M}$ . <sup>24</sup> Indeed, retaining the notation of 3.1.2, the abelian  $X$ -group law on  $T_{X/S}(\mathcal{M})$  is the composite

$$T_{X/S}(\mathcal{M}) \times_X T_{X/S}(\mathcal{M}) \simeq T_{X/S}(\mathcal{M} \oplus \mathcal{M}) \xrightarrow{m} T_{X/S}(\mathcal{M}),$$

whereas

$$T_{X/S}(\mathcal{M}) \xrightarrow{\delta} T_{X/S}(\mathcal{M} \oplus \mathcal{M}) \simeq T_{X/S}(\mathcal{M}) \times_X T_{X/S}(\mathcal{M})$$

is the diagonal morphism.

By Remark 3.4.3 and the identities (†) of 3.1.2,

$$\lambda(x + y) = \lambda x + \lambda y, \quad (\lambda + \mu)x = \lambda x + \mu x$$

for every  $f : X' \rightarrow X$ , every  $x, y \in \text{Hom}_X(X', T_{X/S}(\mathcal{M}))$ , and every  $\lambda, \mu \in \mathcal{O}(X')$ .

**Remark 3.6.1.**

If  $X$  is representable, then it satisfies (E), and  $T_{X/S}$  and  $L_{X/S}^u$  are represented by vector fibrations. The preceding laws are then exactly those induced by the vector-fibration structures (cf. I 4.6). <sup>25</sup>

**Proposition 3.4 bis.**

If  $X/S$  satisfies (E), then  $X_{S'}/S'$  satisfies (E), and the isomorphisms of 3.4 respect the  $\mathcal{O}_{X_{S'}}$ -module and  $\mathcal{O}_{S'}$ -module structures, respectively.

No further comment is needed.

<sup>23</sup>Editor’s note: what follows has been added.

<sup>24</sup>Editor’s note: what follows has been expanded.

<sup>25</sup>Editor’s note: explicitly, for every  $S$ -morphism  $f : X' \rightarrow X$ , the action of  $\mathcal{O}(X')$  on  $\text{Hom}_X(X', T_{X/S}(\mathcal{M}))$  corresponds, under the identification

$$\text{Hom}_X(X', T_{X/S}(\mathcal{M})) \simeq \text{Hom}_{\mathcal{O}_{X'}}(f^* \Omega_{X/S}^1, \mathcal{M} \otimes_{\mathcal{O}_S} \mathcal{O}_{X'}),$$

to the natural action on the right. This follows from the proof of SGA 1, III 5.1; see also the addition 0.1.8 in Exposé III.

**Proposition 3.7.**

The functors  $T_{X/S}(\mathcal{M})$  and  $L_{X/S}^u(\mathcal{M})$  are functorial in  $X$ . Thus, if  $f : X \rightarrow X'$  is an  $S$ -morphism, the following diagrams commute.

$$\begin{array}{ccc} T_{X/S}(\mathcal{M}) & \xrightarrow{T(f)} & T_{X'/S}(\mathcal{M}) \\ \downarrow & & \downarrow \\ X & \xrightarrow{\quad} & X' \end{array}, \quad \begin{array}{ccc} L_{X/S}^u(\mathcal{M}) & \xrightarrow{L(f)} & L_{X'/S}^{f \circ u} \\ & \searrow & \swarrow \\ & S & \end{array}$$

Figure 26: Functoriality of tangent bundles and tangent spaces in  $X$ .

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 12; combined-reader p. 70.*

Moreover, if  $f$  is a monomorphism, then so are  $T(f)$  and  $L(f)$ .<sup>26</sup> The existence of  $T(f)$  and  $L(f)$ , and the last assertion, follow immediately from the definitions. Commutativity follows from functoriality in  $\mathcal{M}$  and from  $T_{X/S}(0) = X$ .

**Remark 3.7.1.**

<sup>27</sup> Suppose that  $X$  and  $X'$  are representable, and let  $r$  be the rank of  $\mathcal{M}$ . By 2.2.2,  $T_{X/S}(\mathcal{M})$  is isomorphic to the product over  $X$  of  $r$  copies of  $V(\Omega_{X/S}^1)$ , and similarly for  $T_{X'/S}(\mathcal{M})$ . Hence the square in 3.7 is cartesian when  $f$  is an open immersion, or more generally when  $f^*\Omega_{X'/S}^1 = \Omega_{X/S}^1$ , for example when  $f$  is étale. Under these conditions there is an isomorphism of  $S$ -functors

$$L_{X/S}^u(\mathcal{M}) \xrightarrow{\sim} L_{X'/S}^{f \circ u}(\mathcal{M}).$$

In general, the square in 3.7 defines the following morphism of  $X$ -functors.

$$\begin{array}{ccc} T_{X/S}(\mathcal{M}) & \longrightarrow & T_{X'/S}(\mathcal{M}) \times_{X'} X \\ & \searrow & \swarrow \\ & X & \end{array}.$$

Figure 27: The induced morphism over  $X$  in the general case.

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 13; combined-reader p. 71.*

**Proposition 3.7 bis.**

Let  $f : X \rightarrow X'$  be an  $S$ -morphism. If  $X$  and  $X'$  satisfy (E) relative to  $S$ , then

$$T_{X/S}(\mathcal{M}) \xrightarrow{T(f)} T_{X'/S}(\mathcal{M})_X, \quad \text{respectively} \quad L_{X/S}^u(\mathcal{M}) \xrightarrow{L(f)} L_{X'/S}^{f \circ u}(\mathcal{M}),$$

is a morphism of  $\mathcal{O}_X$ -modules, respectively of  $\mathcal{O}_S$ -modules.

This follows from Proposition 3.7 by functoriality in  $\mathcal{M}$ .

<sup>26</sup>Editor's note: the preceding sentence has been added.

<sup>27</sup>Editor's note: the hypothesis that  $X$  and  $X'$  be representable has been added, and what follows has been expanded.

**Proposition 3.8.**

For two functors  $X, Y$  over  $S$ , there are isomorphisms, functorial in  $\mathcal{M}$ ,

$$T_{X/S}(\mathcal{M}) \times_S T_{Y/S}(\mathcal{M}) \xrightarrow{\sim} T_{(X \times_S Y)/S}(\mathcal{M}), \quad (3.8.1)$$

$$L_{X/S}^u(\mathcal{M}) \times_S L_{Y/S}^v(\mathcal{M}) \xrightarrow{\sim} L_{(X \times_S Y)/S}^{(u,v)}(\mathcal{M}). \quad (3.8.2)$$

The first isomorphism follows from 1.2 (\*);<sup>28</sup> the second follows from it by base change along  $(u, v) : S \rightarrow X \times_S Y$ .

**Remark 3.8.0.**

The isomorphism (3.8.1) may also be read as an isomorphism of  $X \times_S Y$ -functors

$$(T_{X/S}(\mathcal{M}) \times_X (X \times_S Y)) \times_{X \times_S Y} (T_{Y/S}(\mathcal{M}) \times_Y (X \times_S Y)) \xrightarrow{\sim} T_{(X \times_S Y)/S}(\mathcal{M}).$$

**Corollary 3.8.1.**

If  $X/S$  carries an algebraic structure defined by finite cartesian products, then  $T_{X/S}(\mathcal{M})$  carries a structure of the same kind, and the projection  $T_{X/S}(\mathcal{M}) \rightarrow X$  is a morphism for that structure.

**Proposition 3.8 bis.**

If  $X/S$  and  $Y/S$  satisfy (E), then  $(X \times_S Y)/S$  satisfies (E), and (3.8.1), respectively (3.8.2), is an isomorphism of  $\mathcal{O}_{X \times_S Y}$ -modules, respectively of  $\mathcal{O}_S$ -modules.

*Proof.*<sup>29</sup> By 3.5.1 and (3.8.1), the product  $(X \times_S Y)/S$  satisfies (E). Let us prove the assertion about (3.8.1). Given an  $S$ -morphism  $(x, y) : Z \rightarrow X \times_S Y$ , Remark 3.4.2 reduces the claim to the  $\mathcal{O}(Z)$ -linearity of the map

$$\begin{aligned} & \{\varphi \in \text{Hom}_S(I_Z(\mathcal{M}), X) \mid \varphi \circ \varepsilon_{\mathcal{M}} = x\} \times \{\psi \in \text{Hom}_S(I_Z(\mathcal{M}), Y) \mid \psi \circ \varepsilon_{\mathcal{M}} = y\} \\ & \longrightarrow \{\theta \in \text{Hom}_S(I_Z(\mathcal{M}), X \times_S Y) \mid \theta \circ \varepsilon_{\mathcal{M}} = (x, y)\}, \end{aligned}$$

which sends  $(\varphi, \psi)$  to  $\varphi \times \psi$ . This is clear, since for  $a \in \mathcal{O}(Z)$ ,

$$(\varphi \circ a^*) \times (\psi \circ a^*) = (\varphi \times \psi) \circ a^* = a \cdot (\varphi \times \psi).$$

The assertion about (3.8.2) follows similarly from 3.2.1.

**3.9.0.**

<sup>30</sup> If  $X$  is an  $S$ -group with unit section  $e : S \rightarrow X$ , write

$$\underline{\text{Lie}}(X/S, \mathcal{M}) = L_{X/S}^e(\mathcal{M}).$$

Thus  $\underline{\text{Lie}}(X/S, \mathcal{M})$  is defined by the following cartesian square.

<sup>28</sup>Editor's note: what follows has been expanded.

<sup>29</sup>Editor's note: this proof has been added.

<sup>30</sup>Editor's note: this paragraph has been added to explain the introduction of the notion of an  $S$ - $H$ -functor.

$$\begin{array}{ccc}
\underline{\mathrm{Lie}}(X/S, \mathcal{M}) & \xrightarrow{i} & T_{X/S}(\mathcal{M}) \\
\downarrow & & \downarrow p \\
S & \xrightarrow{e} & X.
\end{array}$$

Figure 28: The cartesian square defining  $\underline{\mathrm{Lie}}(X/S, \mathcal{M})$ .

Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 14;  
combined-reader p. 72.

By Corollary 3.8.1, the projection  $p : T_{X/S}(\mathcal{M}) \rightarrow X$  is a morphism of  $S$ -groups. Hence  $\underline{\mathrm{Lie}}(X/S, \mathcal{M})$  is an  $S$ -group, identified by  $i$  with the kernel of  $p$ .

If  $X/S$  also satisfies (E), Proposition 3.9 will show that this group structure, induced by that of  $X$ , agrees with the abelian-group structure induced by functoriality in  $\mathcal{M}$  (cf. 3.5.1). In fact, the result holds under the weaker assumption that  $X$  be an  $S$ -functor in monoids, or more generally an  $S$ -functor in  $H$ -sets, as defined next.

### Definitions 3.9.0.1.

- (a) An  $H$ -set is a set  $X$  equipped with a composition law having a two-sided identity, denoted  $e_X$  or simply  $e$ .<sup>31</sup> If  $f : X \rightarrow Y$  is a morphism of  $H$ -sets, its kernel is  $\ker f = f^{-1}(e_Y)$ , a sub- $H$ -set of  $X$ .
- (b) An  $H$ -object in a category  $\mathcal{C}$  is an object  $X$  equipped with a morphism  $X \times X \rightarrow X$  and a section of  $X$  over the final object that is a two-sided identity. Every  $\mathcal{C}$ -monoid, and in particular every  $\mathcal{C}$ -group, is a  $\mathcal{C}$ - $H$ -object. An  $H$ -object in the category of functors over  $S$  will be called an  $S$ - $H$ -functor.
- (c) If  $X$  is an  $S$ - $H$ -functor (for example, an  $S$ -group), and  $e : S \rightarrow X$  is its identity section, write

$$\underline{\mathrm{Lie}}(X/S, \mathcal{M}) = L_{X/S}^e(\mathcal{M}), \quad \underline{\mathrm{Lie}}(X/S) = \underline{\mathrm{Lie}}(X/S, \mathcal{O}_S).$$

We now spell out the following special case of 3.8.1.<sup>32</sup>

### Corollary 3.9.0.2.

If  $X$  is an  $S$ - $H$ -functor, respectively an  $S$ -group, then  $T_{X/S}(\mathcal{M})$  and  $\underline{\mathrm{Lie}}(X/S, \mathcal{M})$  are likewise  $S$ - $H$ -functors, respectively  $S$ -groups, and there are morphisms of  $S$ - $H$ -functors, respectively of  $S$ -groups, as follows.

$$\underline{\mathrm{Lie}}(X/S, \mathcal{M}) \xrightarrow{i} T_{X/S}(\mathcal{M}) \begin{array}{c} \xrightarrow{p} \\ \xleftarrow{s} \end{array} X,$$

Figure 29: The inclusion, projection, and zero section for the tangent functor.

Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 15;  
combined-reader p. 73.

Here  $i$  identifies  $\underline{\mathrm{Lie}}(X/S, \mathcal{M})$  with  $\ker p$ , and  $s$  is a section of  $p$ .

<sup>31</sup>Editor's note: this terminology is inspired by the notion of an  $H$ -space in topology.

<sup>32</sup>Editor's note: Corollary 3.9.0.2 has been added; it will be used in 4.1.



**Proposition 3.9.**

Let  $X$  be an  $S$ - $H$ -functor satisfying (E) relative to  $S$ . The  $S$ - $H$ -functor structure on  $\underline{\text{Lie}}(X/S, \mathcal{M})$  induced by that of  $X$  agrees with the  $S$ -group structure defined in 3.5.1.

The preceding discussion shows that  $\underline{\text{Lie}}(X/S, \mathcal{M})$  is an  $H$ -object in the category of  $\mathcal{O}_S$ -modules. The proposition follows from the next lemma. <sup>33</sup>

**Lemma 3.10.**

Let  $\mathcal{C}$  be a category. Let  $G$  be an  $H$ -object in the category of  $\mathcal{C}$ - $H$ -objects. Thus  $G$  is a  $\mathcal{C}$ - $H$ -object, whose composition law we denote  $f : G \times G \rightarrow G$ , together with a morphism of  $\mathcal{C}$ - $H$ -objects  $h : G \times G \rightarrow G$ . <sup>34</sup> Then  $f = h$ , and  $f$  is commutative.

Evaluating the functors on a variable argument reduces the proof, as usual, to the case in which  $\mathcal{C}$  is the category of sets. We then have a set  $G$  and two maps  $f, h : G \times G \rightarrow G$  satisfying

$$h(f(x, y), f(z, t)) = f(h(x, z), h(y, t)).$$

Let  $e, u \in G$  be the respective identities for  $f$  and  $h$ , so that

$$f(e, x) = f(x, e) = x, \quad h(u, x) = h(x, u) = x.$$

First,

$$h(f(u, y), f(x, u)) = f(x, y) = h(f(x, u), f(u, y)).$$

Putting  $y = e$ , respectively  $x = e$ , gives

$$\begin{aligned} x = f(x, e) &= h(f(u, e), f(x, u)) = h(u, f(x, u)) = f(x, u), \\ y = f(e, y) &= h(f(e, u), f(u, y)) = h(u, f(u, y)) = f(u, y). \end{aligned}$$

Substituting these identities into the original relation yields

$$h(y, x) = f(x, y) = h(x, y).$$

This proves the lemma and Proposition 3.9. The following corollaries now follow from 3.9.

**Corollary 3.9.1.**

If  $X$  is an  $S$ - $H$ -functor satisfying (E) relative to  $S$ , every element of  $X(I_S(\mathcal{M}))$  that maps to the identity element of  $X(S)$  is invertible.

**Corollary 3.9.2.**

If  $X$  is an  $S$ -monoid satisfying (E) relative to  $S$ , an element of  $X(I_S(\mathcal{M}))$  is invertible if and only if its image in  $X(S)$  is invertible.

**Corollary 3.9.3.**

If  $X$  is an  $S$ -group satisfying (E) relative to  $S$ , the two  $S$ -group laws on  $\underline{\text{Lie}}(X/S, \mathcal{M})$  coincide.

<sup>33</sup>Editor's note: this is inspired by the standard proof that the fundamental group of an  $H$ -space is abelian.

<sup>34</sup>Editor's note:  $G \times G$  is equipped with the law  $(G \times G) \times (G \times G) \rightarrow G \times G, ((x, z), (y, t)) \mapsto (f(x, y), f(z, t))$ .

**Corollary 3.9.4.**

<sup>35</sup> Let  $G$  be an  $S$ -group satisfying (E) relative to  $S$ . For  $n \in \mathbb{Z}$ , let  $n_G : G \rightarrow G$  be the morphism of  $S$ -functors defined by  $g \mapsto g^n$ . Then the derived morphism

$$L(n_G) : \underline{\mathrm{Lie}}(G/S) \longrightarrow \underline{\mathrm{Lie}}(G/S)$$

is multiplication by  $n$ ; that is, it sends  $x \in \underline{\mathrm{Lie}}(G/S)(S')$  to  $nx$ .

The map  $n_G$  is not generally a group homomorphism, but it preserves the identity section  $e : S \rightarrow G$ , so its derivative indeed carries  $\underline{\mathrm{Lie}}(G/S) = L_{G/S}^e$  to itself. If  $i$  denotes the inclusion  $\underline{\mathrm{Lie}}(G/S) \hookrightarrow T_{G/S}$ , then  $L(n_G)$  is characterized by

$$i(L(n_G)(x)) = i(x)^n.$$

By 3.9, the group laws on  $\underline{\mathrm{Lie}}(G/S)$  coming from (E) and from the law on  $G$  agree. Hence  $i(x)^n = i(nx)$ , and therefore  $L(n_G)(x) = nx$ .

Before drawing further consequences from Proposition 3.9, we prove one more functoriality result.

**Proposition 3.11.**

In the situation of Section 1, there is an isomorphism, functorial in  $\mathcal{M}$ ,

$$T_{\underline{\mathrm{Hom}}_{Z/S}(X,Y)/S}(\mathcal{M}) \xrightarrow{\sim} \underline{\mathrm{Hom}}_{Z/S}(X, T_{Y/Z}(\mathcal{M})).$$

Indeed, by definition,

$$T_{\underline{\mathrm{Hom}}_{Z/S}(X,Y)/S}(\mathcal{M}) = \underline{\mathrm{Hom}}_S(I_S(\mathcal{M}), \underline{\mathrm{Hom}}_{Z/S}(X, Y)).$$

Applying the isomorphism (4) of 1.1 with  $T = I_S(\mathcal{M})$  gives

$$\begin{aligned} \underline{\mathrm{Hom}}_S(I_S(\mathcal{M}), \underline{\mathrm{Hom}}_{Z/S}(X, Y)) \\ \simeq \underline{\mathrm{Hom}}_{Z/S}(X, \underline{\mathrm{Hom}}_Z(Z \times_S I_S(\mathcal{M}), Y)). \end{aligned}$$

Since  $Z \times_S I_S(\mathcal{M}) \simeq I_Z(\mathcal{M})$ , this is

$$\underline{\mathrm{Hom}}_{Z/S}(X, \underline{\mathrm{Hom}}_Z(I_Z(\mathcal{M}), Y)) = \underline{\mathrm{Hom}}_{Z/S}(X, T_{Y/Z}(\mathcal{M})).$$

**Corollary 3.11.1.**

If  $Y/Z$  satisfies (E), then  $\underline{\mathrm{Hom}}_{Z/S}(X, Y)/S$  satisfies (E), and the isomorphism of 3.11 respects the module structures over  $\underline{\mathrm{Hom}}_{Z/S}(X, Y)$ . <sup>36</sup>

*Proof.* Let  $\mathcal{M}, \mathcal{N}$  be finite free  $\mathcal{O}_S$ -modules. If  $Y/Z$  satisfies (E), then

$$T_{Y/Z}(\mathcal{M} \oplus \mathcal{N}) \simeq T_{Y/Z}(\mathcal{M}) \times_Y T_{Y/Z}(\mathcal{N}).$$

The right-hand side is a subfunctor of  $T_{Y/Z}(\mathcal{M}) \times_S T_{Y/Z}(\mathcal{N})$ , and 1.2 (\*) therefore gives

$$\begin{aligned} \underline{\mathrm{Hom}}_{Z/S}(X, T_{Y/Z}(\mathcal{M} \oplus \mathcal{N})) \\ \simeq \underline{\mathrm{Hom}}_{Z/S}(X, T_{Y/Z}(\mathcal{M})) \times_{\underline{\mathrm{Hom}}_{Z/S}(X, Y)} \underline{\mathrm{Hom}}_{Z/S}(X, T_{Y/Z}(\mathcal{N})). \end{aligned}$$

<sup>35</sup>Editor's note: this corollary has been added; it amplifies Remark 4.1.1.2 below.

<sup>36</sup>Editor's note: writing  $H = \underline{\mathrm{Hom}}_{Z/S}(X, Y)$ , this implies in particular that  $T_{H/S}(\mathcal{M})$  is naturally a module over  $\prod_{X \times_S H/H} \mathcal{O}_H$ . This result appeared somewhat surprising to the editors, so the proof has been given in detail.

Together with 3.11, this says that  $\underline{\text{Hom}}_{Z/S}(X, Y)$  satisfies (E), proving the first assertion.

For the second, put  $H = \underline{\text{Hom}}_{Z/S}(X, Y)$ , and let  $\Delta : H' \rightarrow H$  be an  $S$ -morphism. Equivalently, let  $\delta : H' \times_S X \rightarrow Y$  be the corresponding  $Z$ -morphism; it makes  $H' \times_S X$  a  $Y$ -object. On the one hand, we have the following commutative diagram.

$$\begin{array}{ccc}
 \text{Hom}_H(H', \underline{\text{Hom}}_{Z/S}(X, T_{Y/Z}(\mathcal{M}))) & \hookrightarrow & \text{Hom}_S(H', \underline{\text{Hom}}_{Z/S}(X, T_{Y/Z}(\mathcal{M}))) \\
 \parallel & & \parallel \\
 \text{Hom}_Y(H' \times_S X, T_{Y/Z}(\mathcal{M})) & \hookrightarrow & \text{Hom}_Z(H' \times_S X, T_{Y/Z}(\mathcal{M})) \\
 \parallel & & \parallel \\
 \{\psi \in \text{Hom}_Z(I_{H' \times_S X}(\mathcal{M}), Y) \mid \psi \circ \varepsilon_{\mathcal{M}} = \delta\} & \hookrightarrow & \text{Hom}_Z(I_{H' \times_S X}(\mathcal{M}), Y).
 \end{array}$$

Figure 30: The first comparison diagram in the proof of Corollary 3.11.1.

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 17;  
combined-reader p. 75.*

By 1.3, the action of  $\alpha \in \mathcal{O}(H' \times_S X)$  on  $\Psi \in \text{Hom}_Y(H' \times_S X, T_{Y/Z}(\mathcal{M}))$  is given, for every  $U \rightarrow S$  and every  $(h, x) \in \text{Hom}_S(U, H' \times_S X)$ , by

$$(\alpha\Psi)(h, x) = \alpha(h, x)\Psi(h, x),$$

where  $U$  is viewed over  $Y$  through  $\delta \circ (h, x)$ , and  $\alpha(h, x) \in \mathcal{O}(U)$  acts through the  $\mathcal{O}_Y$ -module structure on  $T_{Y/Z}(\mathcal{M})$ . By 3.4.2, under the identification

$$\text{Hom}_Y(H' \times_S X, T_{Y/Z}(\mathcal{M})) = \{\psi \in \text{Hom}_Z(I_{H' \times_S X}(\mathcal{M}), Y) \mid \psi \circ \varepsilon_{\mathcal{M}} = \delta\},$$

this action is

$$(\alpha\psi)(m, h, x) = \psi(m \cdot \alpha(h, x), h, x) \tag{1}$$

for every  $(m, h, x) \in \text{Hom}_S(U, I_S(\mathcal{M}) \times_S H' \times_S X)$ .

On the other hand, consider  $T_{H/S}(\mathcal{M}) = \underline{\text{Hom}}_S(I_S(\mathcal{M}), H)$ . We have a second commutative diagram.

$$\begin{array}{ccc}
 \text{Hom}_H(H', T_{H/S}(\mathcal{M})) & \hookrightarrow & \text{Hom}_S(H', T_{H/S}(\mathcal{M})) \\
 \parallel & & \parallel \\
 \{\Phi \in \text{Hom}_S(I_{H'}(\mathcal{M}), H) \mid \Phi \circ \varepsilon_{\mathcal{M}} = \Delta\} & \hookrightarrow & \text{Hom}_S(I_{H'}(\mathcal{M}), H) \\
 (*) \parallel & & \parallel \\
 \{\phi \in \text{Hom}_Z(I_{H' \times_S X}(\mathcal{M}), Y) \mid \phi \circ \varepsilon_{\mathcal{M}} = \delta\} & \hookrightarrow & \text{Hom}_Z(I_{H' \times_S X}(\mathcal{M}), Y),
 \end{array}$$

Figure 31: The second comparison diagram in the proof of Corollary 3.11.1.

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 18;  
combined-reader p. 76.*

The bijection  $(*)$  in the diagram is described as follows (cf. 1.1(2) and I 1.7.1). For every  $U \rightarrow S$  and  $(m, h, x) \in \text{Hom}_S(U, I_S(\mathcal{M}) \times_S H' \times_S X)$ , where  $U$  lies over  $Z$  through  $U \xrightarrow{x} X \rightarrow Z$ , one has  $\Phi(m, h) \in \text{Hom}_Z(X \times_S U, Y)$  and

$$\varphi(m, h, x) = \Phi(m, h) \circ (x \times \text{id}_U) \in \text{Hom}_Z(U, Y). \tag{\dagger}$$

By 3.4.2, with  $X$  replaced by  $H$  and  $X'$  by  $H'$ , the action of  $a \in \mathcal{O}(H')$  on  $\Phi \in \text{Hom}_S(I_{H'}(\mathcal{M}), H)$  is

$$(a\Phi)(m, h) = \Phi(m \cdot a(h), h).$$

Consequently, if  $\varphi$ , respectively  $a\varphi$ , corresponds to  $\Phi$ , respectively  $a\Phi$ , then  $(\dagger)$  gives

$$(a\varphi)(m, h, x) = \Phi(m \cdot a(h), h) \circ (x \times \text{id}_U) = \varphi(m \cdot a(h), h, x). \quad (2)$$

Together with (1), this proves that the isomorphism

$$T_{H/S}(\mathcal{M}) \xrightarrow{\sim} \underline{\text{Hom}}_{Z/S}(X, T_{Y/Z}(\mathcal{M}))$$

of 3.11.1 is an isomorphism of  $\mathcal{O}(H)$ -modules. Moreover, for every  $H' \rightarrow H$ , the  $\mathcal{O}(H')$ -module structure on  $\text{Hom}_H(H', T_{H/S}(\mathcal{M}))$  extends, functorially in  $H'$ , to an  $\mathcal{O}(H' \times_S X)$ -module structure.

Taking  $Z = S$  yields the next corollary.

### Corollary 3.11.2.

There is an isomorphism, functorial in  $\mathcal{M}$ ,

$$T_{\underline{\text{Hom}}_S(X, Y)/S}(\mathcal{M}) \xrightarrow{\sim} \underline{\text{Hom}}_S(X, T_{Y/S}(\mathcal{M})).$$

If  $Y/S$  satisfies (E), then  $\underline{\text{Hom}}_S(X, Y)/S$  satisfies (E), and this isomorphism respects the module structures over  $\underline{\text{Hom}}_S(X, Y)$ .

Let  $u : X \rightarrow Y$  be an  $S$ -morphism, identified with the constant morphism  $u : S \rightarrow \underline{\text{Hom}}_S(X, Y)$  such that  $u(f) = u$  for every  $f : S' \rightarrow S$ . The fiber product of  $u$  with

$$\underline{\text{Hom}}_S(X, T_{Y/S}(\mathcal{M})) \longrightarrow \underline{\text{Hom}}_S(X, Y)$$

identifies with  $\underline{\text{Hom}}_{Y/S}(X, T_{Y/S}(\mathcal{M}))$ , where  $X$  lies over  $Y$  through  $u$ . Hence the definition of the tangent space and Corollary 3.11.2 give the following result. <sup>37</sup>

### Corollary 3.11.3.

For an  $S$ -morphism  $u : X \rightarrow Y$ , there is an isomorphism, functorial in  $\mathcal{M}$ ,

$$L_{\underline{\text{Hom}}_S(X, Y)/S}^u(\mathcal{M}) \xrightarrow{\sim} \underline{\text{Hom}}_{Y/S}(X, T_{Y/S}(\mathcal{M})),$$

where on the right  $X$  lies over  $Y$  through  $u$ . If  $Y/S$  satisfies (E), this is an isomorphism of  $\mathcal{O}_S$ -modules. <sup>38</sup>

In particular, for  $Y = X$ , the functor  $\underline{\text{End}}_S(X)$  is an  $S$ -functor in monoids, hence an  $S$ - $H$ -functor. Recall that  $\underline{\text{Lie}}(\underline{\text{End}}_S(X)/S, \mathcal{M})$  denotes  $L_{\underline{\text{End}}_S(X)/S}^e(\mathcal{M})$ , with  $e$  the identity section.

### Corollary 3.11.4.

There is an isomorphism, functorial in  $\mathcal{M}$ ,

$$\underline{\text{Lie}}(\underline{\text{End}}_S(X)/S, \mathcal{M}) \xrightarrow{\sim} \prod_{X/S} T_{X/S}(\mathcal{M}).$$

If  $X/S$  satisfies (E), this is an isomorphism of  $\mathcal{O}_S$ -modules. <sup>38</sup>

<sup>37</sup>Editor's note: the preceding sentences have been added.

<sup>38</sup>Editor's note: the preceding sentence has been added.

**Remark 3.11.5.**

<sup>39</sup> Suppose that  $X/S$  satisfies (E). Then

$$\prod_{X/S} T_{X/S}(\mathcal{M}) = \underline{\mathrm{Hom}}_{X/S}(X, T_{X/S}(\mathcal{M}))$$

is a module over  $\prod_{X/S} \mathcal{O}_X$ . Explicitly, for every  $S' \rightarrow S$ ,

$$\begin{aligned} \underline{\mathrm{Hom}}_{X/S}(X, T_{X/S}(\mathcal{M}))(S') \\ = \{ \psi \in \mathrm{Hom}_X(I_{S'}(\mathcal{M}) \times_S X, X) \mid \psi \circ (\varepsilon_{\mathcal{M}} \times \mathrm{id}_X) = \mathrm{pr}_X \} \end{aligned}$$

is functorially an  $\mathcal{O}(X \times_S S')$ -module. This follows either from 3.6 and the properties of  $\prod_{X/S}$  (cf. 1.2), or from the proof of 3.11.1.

We now give a geometric interpretation of the tangent bundle. Let  $U$  be an  $S$ -functor.

<sup>40</sup> By I 1.7.2, there are isomorphisms, functorial in  $\mathcal{M}$ ,

$$\begin{aligned} T_{X/S}(\mathcal{M})(U) &= \mathrm{Hom}_S(U, \underline{\mathrm{Hom}}_S(I_S(\mathcal{M}), X)) \\ &\simeq \mathrm{Hom}_S(I_S(\mathcal{M}), \underline{\mathrm{Hom}}_S(U, X)) \\ &\simeq \mathrm{Hom}_{I_S(\mathcal{M})}(U_{I_S(\mathcal{M})}, X_{I_S(\mathcal{M})}). \end{aligned}$$

In particular, the morphism  $\mathcal{M} \rightarrow 0$  gives the following commutative diagram.

$$\begin{array}{ccc} \mathrm{Hom}_S(U, T_{X/S}(\mathcal{M})) & \xrightarrow{\sim} & \mathrm{Hom}_{I_S(\mathcal{M})}(U_{I_S(\mathcal{M})}, X_{I_S(\mathcal{M})}) \\ \pi_{\mathcal{M}} \downarrow & & \downarrow \\ \mathrm{Hom}_S(U, X) & \xrightarrow{\text{identité}} & \mathrm{Hom}_S(U, X) \end{array} ,$$

Figure 32: The geometric interpretation of the projection  $\pi_{\mathcal{M}}$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp2-14oct24.pdf, component-local p. 19; combined-reader p. 77.*

The second vertical arrow is obtained by base change along  $\varepsilon_{\mathcal{M}} : S \rightarrow I_S(\mathcal{M})$ . <sup>41</sup>

**Proposition 3.12.**

Let  $h_0 : U \rightarrow X$  be an  $S$ -morphism. Then  $\mathrm{Hom}_X(U, T_{X/S}(\mathcal{M}))$  identifies with the set of  $I_S(\mathcal{M})$ -morphisms

$$U_{I_S(\mathcal{M})} \longrightarrow X_{I_S(\mathcal{M})}$$

whose restriction to  $U$  is  $h_0$ , where  $U$  is regarded as a subobject of  $U \times_S I_S(\mathcal{M})$  through  $\mathrm{id}_U \times_S \varepsilon_{\mathcal{M}}$ .

Taking  $U = X$  and  $h_0 = \mathrm{id}_X$  gives the following. <sup>42</sup>

<sup>39</sup>Editor's note: this remark has been added.

<sup>40</sup>Editor's note: the original has been simplified in what follows.

<sup>41</sup>Editor's note: via

$$\mathrm{Hom}_{I_S(\mathcal{M})}(U_{I_S(\mathcal{M})}, X_{I_S(\mathcal{M})}) \simeq \mathrm{Hom}_S(U \times_S I_S(\mathcal{M}), X),$$

this is equivalent to Remark 3.1.1. Likewise, 3.12 is equivalent to the first part of Remark 3.4.2.

<sup>42</sup>Editor's note: the original has been expanded in what follows.

**Corollary 3.12.1.**

The set  $\Gamma(T_{X/S}(\mathcal{M})/X)$  identifies with the set of  $I_S(\mathcal{M})$ -endomorphisms  $\phi$  of  $X_{I_S(\mathcal{M})}$  inducing the identity on  $X$ ; equivalently, the following diagram commutes.<sup>43</sup>

$$\begin{array}{ccc} I_X(\mathcal{M}) & \xrightarrow{\phi} & I_X(\mathcal{M}) \\ \varepsilon_{\mathcal{M}} \uparrow & & \uparrow \varepsilon_{\mathcal{M}} \\ X & \xrightarrow{\text{id}} & X. \end{array}$$

Figure 33: An infinitesimal endomorphism inducing the identity on  $X$ .

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 20; combined-reader p. 78.*

On the other hand, by 3.11.2,

$$\Gamma(T_{X/S}(\mathcal{M})/X) \simeq \underline{\text{Lie}}(\underline{\text{End}}_S(X)/S, \mathcal{M})(S).$$

If  $X/S$  satisfies (E), then  $\underline{\text{End}}_S(X)/S$  satisfies (E), so the latter Lie functor is an  $\mathcal{O}_S$ -module by 3.6 (indeed, by 3.11.5, it is a  $\prod_{X/S} \mathcal{O}_X$ -module). Applying 3.9 gives the next proposition.<sup>44</sup>

**Proposition 3.13.**

If  $X/S$  satisfies (E), the abelian group  $\Gamma(T_{X/S}(\mathcal{M})/X)$  identifies with the set of  $I_S(\mathcal{M})$ -endomorphisms of  $X_{I_S(\mathcal{M})}$  that induce the identity on  $X$ . Consequently, every such endomorphism is an automorphism.

**Corollary 3.13.1.**

Let  $u : X \rightarrow Y$  be an  $S$ -isomorphism, with  $Y/S$  satisfying (E). Every  $I_S(\mathcal{M})$ -morphism  $X_{I_S(\mathcal{M})} \rightarrow Y_{I_S(\mathcal{M})}$  extending  $u$  is an isomorphism.

**Corollary 3.13.2.**

If  $Y/S$  satisfies (E), the monomorphism

$$\underline{\text{Isom}}_S(X, Y) \longrightarrow \underline{\text{Hom}}_S(X, Y)$$

induces, for every  $u \in \underline{\text{Isom}}_S(X, Y)$ , an isomorphism

$$L_{\underline{\text{Isom}}_S(X, Y)/S}^u(\mathcal{M}) \xrightarrow{\sim} L_{\underline{\text{Hom}}_S(X, Y)/S}^u(\mathcal{M}).$$

*Proof.*<sup>45</sup> For every  $S' \rightarrow S$ , we must show that this map is a bijection on  $S'$ -points. By base change (cf. 3.4), it suffices to take  $S' = S$ . The right-hand side then parametrizes the  $I_S(\mathcal{M})$ -morphisms  $X_{I_S(\mathcal{M})} \rightarrow Y_{I_S(\mathcal{M})}$  extending  $u$ , whereas the left-hand side parametrizes those that are isomorphisms. The assertion is therefore Corollary 3.13.1.

<sup>43</sup>Editor's note: when  $\mathcal{M} = \mathcal{O}_X$ , these are called *infinitesimal endomorphisms* of  $X$ ; see 6.2 at the end of this exposé.

<sup>44</sup>Editor's note: the original has been expanded in what follows.

<sup>45</sup>Editor's note: the following proof has been added.

**Corollary 3.13.3.**

If  $X/S$  satisfies (E), the monomorphism  $\underline{\mathrm{Aut}}_S(X) \rightarrow \underline{\mathrm{End}}_S(X)$  induces, for every  $u \in \underline{\mathrm{Aut}}_S(X)$ , an isomorphism

$$L_{\underline{\mathrm{Aut}}_S(X)/S}^u(\mathcal{M}) \xrightarrow{\sim} L_{\underline{\mathrm{End}}_S(X)/S}^u(\mathcal{M}).$$

In particular,

$$\underline{\mathrm{Lie}}(\underline{\mathrm{Aut}}_S(X)/S, \mathcal{M}) \xrightarrow{\sim} \underline{\mathrm{Lie}}(\underline{\mathrm{End}}_S(X)/S, \mathcal{M}) \xrightarrow{\sim} \prod_{X/S} T_{X/S}(\mathcal{M}),$$

so that  $\underline{\mathrm{Lie}}(\underline{\mathrm{Aut}}_S(X)/S, \mathcal{M})$  is a  $\prod_{X/S} \mathcal{O}_X$ -module.<sup>46</sup>

**3.14.**

Finally, suppose that  $X$  is representable.<sup>47</sup> By 2.2.2, the  $X$ -functor  $T_{X/S}$  is represented by  $V(\Omega_{X/S}^1)$ , whence bijections

$$\Gamma(T_{X/S}/X) \simeq \mathrm{Hom}_X(\Omega_{X/S}^1, \mathcal{O}_X) \simeq \mathrm{Der}_{\mathcal{O}_S}(\mathcal{O}_X).$$

This also follows from the preceding results. By 3.13,  $\Gamma(T_{X/S}/X)$  identifies with the set of infinitesimal endomorphisms of  $X$ , that is, the  $I_S$ -endomorphisms of  $X_{I_S}$  that induce the identity on  $X$ . The schemes  $X$  and  $X_{I_S}$  have the same underlying topological space, and their structure sheaves are  $\mathcal{O}_X$  and

$$\mathcal{D}_{\mathcal{O}_X} = \mathcal{O}_X \oplus \mathcal{M}, \quad \mathcal{M} = \mathcal{O}_X,$$

where  $\mathcal{M}$  is a square-zero ideal. Let  $\pi : \mathcal{D}_{\mathcal{O}_X} \rightarrow \mathcal{O}_X$  be the morphism of  $\mathcal{O}_X$ -algebras vanishing on  $\mathcal{M}$ . Giving an infinitesimal endomorphism of  $X$  is equivalent to giving a morphism of  $\mathcal{O}_S$ -algebras  $\phi : \mathcal{O}_X \rightarrow \mathcal{D}_{\mathcal{O}_X}$  satisfying  $\pi \circ \phi = \mathrm{id}_{\mathcal{O}_X}$ , hence to giving an  $\mathcal{O}_S$ -derivation of  $\mathcal{O}_X$ .

If  $D, D' \in \mathrm{Der}_{\mathcal{O}_S}(\mathcal{O}_X)$ , and  $\phi_D$  denotes the infinitesimal endomorphism corresponding to  $D$ , then

$$\phi_{D+D'} = \phi_D \circ \phi_{D'}.$$

Thus the identification

$$\{\text{infinitesimal endomorphisms of } X\} \simeq \mathrm{Der}_{\mathcal{O}_S}(\mathcal{O}_X)$$

is an isomorphism of abelian groups. In view of 3.13 and 3.11.5, we have therefore constructed an isomorphism of abelian groups, indeed of  $\mathcal{O}(X)$ -modules,

$$\Gamma(T_{X/S}/X) \xrightarrow{\sim} \mathrm{Der}_{\mathcal{O}_S}(\mathcal{O}_X),$$

recovering the classical interpretation of tangent vector fields as derivations of the structure sheaf.<sup>48</sup> Finally,

$$\Gamma(T_{X/S}/X) = H^0(X, \mathfrak{g}_{X/S}),$$

where  $\mathfrak{g}_{X/S}$  is the dual of  $\Omega_{X/S}^1$ .

<sup>46</sup>Editor's note: the preceding assertion has been added.

<sup>47</sup>Editor's note: the reminder of 2.2.2 has been added, and the sequel has been expanded.

<sup>48</sup>Editor's note: this isomorphism is also functorial in  $S$ . If  $S' \rightarrow S$  and  $X' = X_{S'}$ , then

$$\underline{\mathrm{Lie}}(\underline{\mathrm{Aut}}_S(X)/S)(S') \simeq \Gamma(T_{X'/S'}/X') \simeq \mathrm{Der}_{\mathcal{O}_{S'}}(\mathcal{O}_{X'}).$$

## 4. Tangent Space to a Group — Lie Algebras

### 4.1.

Let  $G$  be a group functor over  $S$ . By 3.9.0.2,  $T_{G/S}(\mathcal{M})$  and  $\underline{\mathrm{Lie}}(G/S, \mathcal{M})$  are endowed with group structures over  $S$ , and there are group morphisms

$$\underline{\mathrm{Lie}}(G/S, \mathcal{M}) \xrightarrow{i} T_{G/S}(\mathcal{M}) \xrightleftharpoons[s]{p} G \quad ;$$

Figure 34: The split tangent sequence of the group functor  $G$ .

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 22;  
combined-reader p. 80; printed p. 70.*

By definition,  $i$  identifies  $\underline{\mathrm{Lie}}(G/S, \mathcal{M})$  with the kernel of  $p$ , and  $s$  is a section of  $p$ . It follows from I 2.3.7 that this sequence identifies  $T_{G/S}(\mathcal{M})$  with the semidirect product of  $G$  by  $\underline{\mathrm{Lie}}(G/S, \mathcal{M})$ .

#### Definition 4.1.A.

The corresponding action of  $G$  on  $\underline{\mathrm{Lie}}(G/S, \mathcal{M})$  is denoted

$$\mathrm{Ad} : G \longrightarrow \underline{\mathrm{Aut}}_{S\text{-gr.}}(\underline{\mathrm{Lie}}(G/S, \mathcal{M}))$$

and is called the *adjoint representation* of  $G$ , relative to  $\mathcal{M}$ . Thus, by definition, for  $x \in G(S')$  and  $X \in \underline{\mathrm{Lie}}(G/S, \mathcal{M})(S')$ ,

$$\mathrm{Ad}(x)X = i^{-1}(s(x)i(X)s(x)^{-1}).$$

If  $G$  is commutative, then  $T_{G/S}(\mathcal{M})$  is commutative as well and  $\mathrm{Ad}(x)X = X$ .<sup>49</sup>

#### Definition 4.1.B.

Let  $G, H$  be group functors over  $S$ , and let  $f : G \rightarrow H$  be a group morphism. Functoriality gives a morphism of exact sequences compatible with the canonical sections.

$$\begin{array}{ccccccc} 1 & \longrightarrow & \underline{\mathrm{Lie}}(G/S, \mathcal{M}) & \longrightarrow & T_{G/S}(\mathcal{M}) & \longrightarrow & G \longrightarrow 1 \\ & & \downarrow L(f) & & \downarrow T(f) & & \downarrow f \\ 1 & \longrightarrow & \underline{\mathrm{Lie}}(H/S, \mathcal{M}) & \longrightarrow & T_{H/S}(\mathcal{M}) & \longrightarrow & H \longrightarrow 1 \end{array} \quad ;$$

Figure 35: Functoriality of the split tangent sequence.

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 22;  
combined-reader p. 80; printed p. 70.*

The morphism  $L(f)$ , also denoted  $\underline{\mathrm{Lie}}(f)$ , is called the *derived morphism* of  $f$ .<sup>50</sup>

<sup>49</sup>Editor’s note: the numbering 4.1.A and 4.1.B has been added in order to bring these definitions into relief.

<sup>50</sup>Editor’s note: if  $f$  is a monomorphism, then so are  $L(f)$  and  $T(f)$ ; cf. 3.7.



**Remark 4.1.C.**

If  $G/S$  and  $H/S$  satisfy condition  $E$ , then  $L(f)$  respects the  $\mathcal{O}_S$ -module structures obtained from “functoriality in  $\mathcal{M}$ ” (cf. 3.6).<sup>51</sup>

**Proposition 4.1.1.**

Let  $g \in G(S)$ . Then

$$\mathrm{Ad}(g) : \underline{\mathrm{Lie}}(G/S, \mathcal{M}) \longrightarrow \underline{\mathrm{Lie}}(G/S, \mathcal{M})$$

is the derived morphism of  $\mathrm{Int}(g) : G \rightarrow G$ .

Indeed,  $\mathrm{Ad}(g)X = i^{-1}(\mathrm{Int}(g)i(X))$ , which is precisely  $L(\mathrm{Int}(g))(X)$  by the definition of the derived morphism.

Suppose that  $G/S$  satisfies  $E$ . By Proposition 3.9, the group structure just defined on  $\underline{\mathrm{Lie}}(G/S, \mathcal{M})$  is exactly the one induced by its  $\mathcal{O}_S$ -module structure (which is defined by (E)). Proposition 4.1.1 and the functoriality of the  $\mathcal{O}_S$ -action (cf. 3.6) therefore yield the following corollary.

**Corollary 4.1.1.1.**

Suppose that  $G/S$  satisfies  $E$ . Then  $\mathrm{Ad}$  maps  $G$  into the subgroup

$$\underline{\mathrm{Aut}}_{\mathcal{O}_S\text{-mod.}}(\underline{\mathrm{Lie}}(G/S, \mathcal{M}))$$

of  $\underline{\mathrm{Aut}}_{S\text{-gr.}}(\underline{\mathrm{Lie}}(G/S, \mathcal{M}))$ . In other words, for every  $g \in G(S')$ , the automorphism  $\mathrm{Ad}(g)$  respects the  $\mathcal{O}_{S'}$ -module structure on  $\underline{\mathrm{Lie}}(G'/S', \mathcal{M})$ , where  $G' = G \times_S S'$ . Thus  $\mathrm{Ad}$  is a *linear representation* (cf. I 3.2) of  $G$  on the  $\mathcal{O}_S$ -module  $\underline{\mathrm{Lie}}(G/S, \mathcal{M})$ .

**Remarks 4.1.1.2.**

a) The functor  $G/S$  satisfies  $E$  if and only if, for every pair  $(\mathcal{M}, \mathcal{N})$  of finite free  $\mathcal{O}_S$ -modules, the following diagram, obtained by applying  $\underline{\mathrm{Lie}}(G/S, -)$  to diagram (\*) of 2.1, is cartesian.

$$\begin{array}{ccc} & \underline{\mathrm{Lie}}(G/S, \mathcal{M} \oplus \mathcal{N}) & \\ \swarrow & & \searrow \\ \underline{\mathrm{Lie}}(G/S, \mathcal{M}) & & \underline{\mathrm{Lie}}(G/S, \mathcal{N}) \\ \searrow & & \swarrow \\ & \underline{\mathrm{Lie}}(G/S, 0) = S & \end{array},$$

Figure 36: The cartesian criterion for condition  $E$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp2-14oct24.pdf, component-local p. 23;  
combined-reader p. 81; printed p. 71.*

b) Suppose that  $G/S$  satisfies  $E$ . The derived morphism of the group law  $\pi : G \times_S G \rightarrow G$  is the addition law on  $\underline{\mathrm{Lie}}(G/S, \mathcal{M})$ . Notice that  $\pi$  is not a group morphism, but  $\pi(e, e) = e$ ; hence the derived morphism  $L(\pi)$  maps

$$T_{(G \times_S G)/S}^{(e,e)}(\mathcal{M}) = \underline{\mathrm{Lie}}(G/S, \mathcal{M}) \times_S \underline{\mathrm{Lie}}(G/S, \mathcal{M})$$

to  $\underline{\mathrm{Lie}}(G/S, \mathcal{M})$ ; cf. 3.7 and 3.8. Likewise, for  $n \in \mathbb{Z}$ , let  $n_G : G \rightarrow G$  denote the  $S$ -functor morphism  $g \mapsto g^n$ . Then  $L(n_G)$  is multiplication by  $n$  on  $\underline{\mathrm{Lie}}(G/S)$ ; cf. 3.9.4.

<sup>51</sup>Editor’s note: this remark has been added.

**4.1.2.0.**

Consider the  $S$ -functor  $\underline{\mathrm{Hom}}_{G/S}(G, T_{G/S}(\mathcal{M}))$ .<sup>52</sup> For every  $S' \rightarrow S$ , one has  $T_{G/S}(\mathcal{M})_{S'} \simeq T_{G_{S'}/S'}(\mathcal{M})$  (cf. 3.4), and therefore

$$\begin{aligned} \underline{\mathrm{Hom}}_{G/S}(G, T_{G/S}(\mathcal{M}))(S') \\ \simeq \mathrm{Hom}_{G_{S'}}(G_{S'}, T_{G_{S'}/S'}(\mathcal{M})) = \Gamma(T_{G_{S'}/S'}(\mathcal{M})/G_{S'}). \end{aligned}$$

First note the isomorphism, functorial in  $S'$ ,

$$\mathrm{Hom}_{S'}(G_{S'}, \underline{\mathrm{Lie}}(G_{S'}/S', \mathcal{M})) \xrightarrow{\sim} \Gamma(T_{G_{S'}/S'}(\mathcal{M})/G_{S'}), \quad (*)$$

which sends  $f : G_{S'} \rightarrow \underline{\mathrm{Lie}}(G_{S'}/S', \mathcal{M})$  to the section  $s_f : G_{S'} \rightarrow T_{G_{S'}/S'}(\mathcal{M})$  defined, for every  $S'' \rightarrow S'$  and  $g \in G(S'')$ , by

$$s_f(g) = i(f(g))s(g).$$

Let  $h$  be an automorphism of the functor  $G_{S'}$  over  $S'$ , not necessarily preserving the group structure. To every section  $\tau$  of  $T_{G_{S'}/S'}(\mathcal{M})$  one associates, by transport of structure, a section  $h(\tau)$ : it is the unique section that makes the following diagram commute.

$$\begin{array}{ccc} G_{S'} & \xrightarrow{\tau} & T_{G_{S'}/S'}(\mathcal{M}) \\ h \downarrow & & \downarrow \mathrm{T}(h) \\ G_{S'} & \xrightarrow{h(\tau)} & T_{G_{S'}/S'}(\mathcal{M}) \end{array} \quad .$$

Figure 37: Transport of a tangent section by an automorphism  $h$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp2-14oct24.pdf, component-local p. 24;  
combined-reader p. 82; printed p. 72.*

In particular, take  $h = t_x$ , right translation by an element  $x \in G(S')$ ; thus  $h(g) = t_x(g) = g \cdot x$  for all  $g \in G(S'')$ ,  $S'' \rightarrow S'$ . One then has

$$t_x(s_f) = s_{t_x(f)},$$

where  $t_x(f) : G_{S'} \rightarrow \underline{\mathrm{Lie}}(G_{S'}/S', \mathcal{M})$  is given by

$$t_x(f)(g) = f(g \cdot x^{-1}).$$

Let  $G$  act by right translations on

$$\underline{\mathrm{Hom}}_{G/S}(G, T_{G/S}(\mathcal{M})) \quad \text{and} \quad \underline{\mathrm{Hom}}_S(G, \underline{\mathrm{Lie}}(G/S, \mathcal{M}))$$

as follows. For  $S' \rightarrow S$  and  $x \in G(S')$ , let

$$\begin{aligned} \sigma &\in \Gamma(T_{G_{S'}/S'}(\mathcal{M})/G_{S'}), \\ f &\in \mathrm{Hom}_{S'}(G_{S'}, \underline{\mathrm{Lie}}(G_{S'}/S', \mathcal{M})). \end{aligned}$$

Set

$$(\sigma \cdot x)(g) = \sigma(g \cdot x^{-1}) \cdot s(x), \quad (f \cdot x)(g) = f(g \cdot x^{-1})$$

<sup>52</sup>Editor's note: the original has been expanded in what follows to display the action of  $G$  on the functors  $\underline{\mathrm{Hom}}_{G/S}(G, T_{G/S}(\mathcal{M}))$  and  $\underline{\mathrm{Hom}}_S(G, \underline{\mathrm{Lie}}(G/S, \mathcal{M}))$ .

for every  $g \in G(S'')$ ,  $S'' \rightarrow S'$ . The isomorphism  $(*)$  respects these  $G$ -actions.

Consequently, under this isomorphism, the elements of

$$\underline{\mathrm{Hom}}_{G/S}(G, T_{G/S}(\mathcal{M}))^G(S')$$

correspond to the constant morphisms from  $G_{S'}$  to  $\underline{\mathrm{Lie}}(G_{S'}/S', \mathcal{M})$ , that is, to the morphisms factoring through  $G_{S'} \rightarrow S'$ ; equivalently, they correspond to the elements of  $\underline{\mathrm{Lie}}(G/S, \mathcal{M})(S')$ .

**Terminology.** The elements of  $\underline{\mathrm{Hom}}_{G/S}(G, T_{G/S}(\mathcal{M}))^G(S')$  are called the sections of  $T_{G_{S'}/S'}(\mathcal{M})$  invariant under right translation.

### Proposition 4.1.2.

The map

$$\underline{\mathrm{Lie}}(G/S, \mathcal{M})(S) \longrightarrow \Gamma(T_{G/S}(\mathcal{M})/G), \quad X \longmapsto (x \mapsto X \cdot x),$$

is a bijection onto the set of sections invariant under right translation.<sup>53</sup>

Likewise, let  $G$  act on the right on  $\underline{\mathrm{End}}_{I_S(\mathcal{M})/S}(G_{I_S(\mathcal{M})})$  by

$$(u \cdot x)(g) = u(g \cdot x^{-1}) \cdot x,$$

for  $S' \rightarrow S$ ,  $x \in G(S')$ ,  $u \in \underline{\mathrm{End}}_{I_{S'}(\mathcal{M})}(G_{I_{S'}(\mathcal{M})})$ , and  $g \in G(S'')$ ,  $S'' \rightarrow I_{S'}(\mathcal{M})$ . The morphism of 3.12.1

$$\underline{\mathrm{Hom}}_{G/S}(G, T_{G/S}(\mathcal{M})) \longrightarrow \underline{\mathrm{End}}_{I_S(\mathcal{M})/S}(G_{I_S(\mathcal{M})})$$

respects the  $G$ -actions. For each  $S' \rightarrow S$ , it therefore induces a bijection between  $\Gamma(T_{G_{S'}/S'}(\mathcal{M})/G_{S'})$  and the set of  $I_{S'}(\mathcal{M})$ -endomorphisms  $u$  of  $G_{I_{S'}(\mathcal{M})}$  that induce the identity on  $G$  and “commute with right translations,” meaning that  $u_{S''} \cdot x = u_{S''}$  for every  $S'' \rightarrow S'$  and  $x \in G(S'')$ .

### Proposition 4.1.3.

There is a bijection, functorial in  $G$ , between  $\underline{\mathrm{Lie}}(G/S, \mathcal{M})(S)$  and the set of  $I_S(\mathcal{M})$ -endomorphisms of  $G_{I_S(\mathcal{M})}$  that induce the identity on  $G$  and commute with the right translations of  $G$ , in the sense just specified.

Taking 3.13 into account gives the following theorem.

### Theorem 4.1.4.

Let  $G$  be an  $S$ -functor in groups, and suppose that  $G/S$  satisfies  $E$ . Then  $\underline{\mathrm{Lie}}(G/S, \mathcal{M})(S)$  is identified, functorially in  $G$ , with the group of  $I_S(\mathcal{M})$ -automorphisms of  $G_{I_S(\mathcal{M})}$  that induce the identity on  $G$  and commute with the right translations of  $G$ .

For  $\mathcal{M} = \mathcal{O}_S$ , this recovers one of the classical definitions of the Lie algebra of a group.

<sup>53</sup>Statements 4.1.2, 4.1.3, and 4.1.4 can be obtained more simply by observing that the automorphisms of  $G$  invariant under right translations are the left translations. Editor’s note: see, for example, the proofs of Propositions 4.6 and 2.5 of [DG70], §II.4.

**4.2.0.**

Before proceeding, let us establish further consequences of 3.11.<sup>54</sup> Let  $X, Y$  be over  $Z$ , with  $Z$  over  $S$ , as in Section 1. As was seen in 3.11, the isomorphisms 1.1(4)

$$\begin{array}{ccc} \underline{\mathrm{Hom}}_S(\mathrm{I}_S(\mathcal{M}), \underline{\mathrm{Hom}}_{Z/S}(X, Y)) & \xrightarrow{\sim} & \underline{\mathrm{Hom}}_{Z/S}(X, \underline{\mathrm{Hom}}_Z(\mathrm{I}_Z(\mathcal{M}), Y)) \\ & \searrow \simeq & \nearrow \simeq \\ & \underline{\mathrm{Hom}}_{Z/S}(X \times_S \mathrm{I}_S(\mathcal{M}), Y) & \end{array}$$

Figure 38: The adjunction isomorphisms of 1.1(4).

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 25;  
combined-reader p. 83; printed p. 73.*

induce the following isomorphism  $\theta$ .

$$\begin{array}{ccc} T_{\underline{\mathrm{Hom}}_{Z/S}(X, Y)}(\mathcal{M}) & \xrightarrow[\sim]{\theta} & \underline{\mathrm{Hom}}_{Z/S}(X, T_{Y/Z}(\mathcal{M})) \\ & \searrow \simeq & \nearrow \simeq \\ & \underline{\mathrm{Hom}}_{Z/S}(X \times_S \mathrm{I}_S(\mathcal{M}), Y) & \end{array} .$$

Figure 39: The induced tangent-space isomorphism  $\theta$ .

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 25;  
combined-reader p. 83; printed p. 73.*

By 1.3, if  $Y$  is a  $Z$ -group, then  $\underline{\mathrm{Hom}}_Z(V, Y)$  is a  $Z$ -group for every  $V \rightarrow Z$ , in particular for  $V = \mathrm{I}_Z(\mathcal{M})$ . Explicitly, if  $Z'' \rightarrow Z' \rightarrow Z$  and  $\varphi, \psi \in \underline{\mathrm{Hom}}_Z(V_{Z'}, Y)(Z')$ , their product is defined by

$$(\varphi \cdot \psi)(v) = \varphi(v)\psi(v) \quad (v \in V_{Z'}(Z'')).$$

Suppose now that  $X$  and  $Y$  are  $Z$ -groups.

**Definition 4.2.0.1.**

We denote by  $\underline{\mathrm{Hom}}_{(Z/S)\text{-gr.}}(X, Y)$  the subfunctor of  $\underline{\mathrm{Hom}}_{Z/S}(X, Y)$  defined, for every  $S' \rightarrow S$ , by

$$\underline{\mathrm{Hom}}_{(Z/S)\text{-gr.}}(X, Y)(S') = \underline{\mathrm{Hom}}_{Z_{S'}\text{-gr.}}(X_{S'}, Y_{S'}). \quad (3)$$

The same definition applies with  $Y$  replaced by the  $Z$ -group  $T_{Y/Z}(\mathcal{M})$ .

Under the isomorphisms (2) above,  $T_{\underline{\mathrm{Hom}}_{(Z/S)\text{-gr.}}(X, Y)/S}(\mathcal{M})(S')$  corresponds to the  $Z_{S'}$ -morphisms

$$\varphi : X_{S'} \times_{S'} \mathrm{I}_{S'}(\mathcal{M}) \longrightarrow Y_{S'}$$

that are multiplicative in  $X$ , that is,

$$\varphi(x_1 x_2, m) = \varphi(x_1, m) \varphi(x_2, m).$$

These in turn correspond to the morphisms of  $Z_{S'}$ -groups  $X_{S'} \rightarrow T_{Y/Z}(\mathcal{M})_{S'}$ . Hence:

<sup>54</sup>Editor's note: paragraph 4.2.0 has been added; its results are used implicitly in 4.2 and 4.7 of the original.

**Proposition 4.2.0.2.**

Let  $X, Y$  be  $Z$ -groups, where  $Z$  is over  $S$ . There are isomorphisms of  $S$ -functors, functorial in  $\mathcal{M}$ ,

$$T_{\underline{\text{Hom}}_{(Z/S)\text{-gr.}}(X,Y)/S}(\mathcal{M}) \xrightarrow{\sim} \underline{\text{Hom}}_{(Z/S)\text{-gr.}}(X, T_{Y/Z}(\mathcal{M})).$$

Taking  $Z = S$  gives the following consequence. If  $Y$  is a commutative  $S$ -group, then so is  $T_{Y/S}(\mathcal{M})$ , and hence so are  $H = \underline{\text{Hom}}_{S\text{-gr.}}(X, Y)$ ,  $\underline{\text{Hom}}_{S\text{-gr.}}(X, T_{Y/S}(\mathcal{M}))$ , and  $T_{H/S}(\mathcal{M})$ .

**Corollary 4.2.0.3.**

Let  $X, Y$  be  $S$ -groups. There are isomorphisms of  $S$ -functors, functorial in  $\mathcal{M}$ ,

$$T_{\underline{\text{Hom}}_{S\text{-gr.}}(X,Y)/S}(\mathcal{M}) \xrightarrow{\sim} \underline{\text{Hom}}_{S\text{-gr.}}(X, T_{Y/S}(\mathcal{M})).$$

If  $Y$  is commutative, these are moreover isomorphisms of abelian  $S$ -groups.

**Definition 4.2.0.4.**

If  $Y$  is an  $\mathcal{O}_S$ -module, the functor  $T_{Y/S}(\mathcal{M})$ , respectively  $\underline{\text{Lie}}(Y/S, \mathcal{M})$ , carries an  $\mathcal{O}_S$ -module structure induced by that of  $Y$ . Equipped with this structure, it will be denoted  $T'_{Y/S}(\mathcal{M})$ , respectively  $\underline{\text{Lie}}'(Y/S, \mathcal{M})$ .<sup>55</sup>

Consequently, if  $X, Y$  are  $\mathcal{O}_S$ -modules, then  $T'_{Y/S}(\mathcal{M}) = \underline{\text{Hom}}_S(I_S(\mathcal{M}), Y)$ . If  $H = \underline{\text{Hom}}_{\mathcal{O}_S\text{-mod.}}(X, Y)$ , then  $\underline{\text{Hom}}_{\mathcal{O}_S\text{-mod.}}(X, T'_{Y/S}(\mathcal{M}))$  and  $T'_{H/S}(\mathcal{M})$  also carry  $\mathcal{O}_S$ -module structures.

**Corollary 4.2.0.5.**

If  $X, Y$  are  $\mathcal{O}_S$ -modules, there are isomorphisms of  $\mathcal{O}_S$ -modules, functorial in  $\mathcal{M}$ ,

$$T'_{\underline{\text{Hom}}_{\mathcal{O}_S\text{-mod.}}(X,Y)/S}(\mathcal{M}) \xrightarrow{\sim} \underline{\text{Hom}}_{\mathcal{O}_S\text{-mod.}}(X, T'_{Y/S}(\mathcal{M})).$$

**Definition 4.2.A.**

Let  $X, L$  be  $S$ -groups, with  $X$  acting on  $L$  by group automorphisms (cf. I 2.3.5). We define the subfunctor  $\underline{Z}_S^1(X, L)$  of  $\underline{\text{Hom}}_S(X, L)$  by

$$\underline{Z}_S^1(X, L)(S') = \{ \varphi \in \text{Hom}_{S'}(X_{S'}, L_{S'}) \mid \varphi(x_1 x_2) = \varphi(x_1)(x_1 \cdot \varphi(x_2)), \\ x_1, x_2 \in X(S''), S'' \rightarrow S' \}.$$

It is called the *functor of crossed homomorphisms* from  $X$  to  $L$ .<sup>56</sup>

**Remark 4.2.B.**

a) If  $L'$  is another  $S$ -group on which  $X$  acts by group automorphisms, then

$$\underline{Z}_S^1(X, L \times_S L') \simeq \underline{Z}_S^1(X, L) \times_S \underline{Z}_S^1(X, L').$$

b) If  $L$  is an  $X$ - $\mathcal{O}_S$ -module, then  $\underline{Z}_S^1(X, L)$  is the kernel of the differential

$$\partial : \underline{\text{Hom}}_S(X, L) \longrightarrow \underline{\text{Hom}}_S(X^2, L)$$

defined in I 5.1. In particular,  $\underline{Z}_S^1(X, L)$  is then an  $\mathcal{O}_S$ -module.

<sup>55</sup>Editor's note: the prime is used to distinguish the  $\mathcal{O}_S$ -module structure induced by that of  $Y$  from the one that will be obtained from condition E.

<sup>56</sup>Editor's note: the original has been expanded below; in particular, Definition 4.2.A and Remark 4.2.B have been added.

## 4.2.

Let  $u : X \rightarrow Y$  be a morphism of  $S$ -groups. We describe  $L_{\underline{\text{Hom}}_{S\text{-gr.}}(X,Y)/S}^u(\mathcal{M})$ . First, 3.11.3 gives an isomorphism of  $S$ -functors, functorial in  $\mathcal{M}$ ,

$$L_{\underline{\text{Hom}}_{S(X,Y)/S}}^u(\mathcal{M}) \xrightarrow{\sim} \underline{\text{Hom}}_{Y/S}(X, T_{Y/S}(\mathcal{M})). \quad (\dagger)$$

Since  $Y$  is an  $S$ -group,

$$T_{Y/S}(\mathcal{M}) = \underline{\text{Lie}}(Y/S, \mathcal{M}) \cdot Y = \underline{\text{Lie}}(Y/S, \mathcal{M})_Y,$$

and one obtains an isomorphism, functorial in  $\mathcal{M}$ ,

$$L_{\underline{\text{Hom}}_{S(X,Y)/S}}^u(\mathcal{M}) \xrightarrow{\sim} \underline{\text{Hom}}_S(X, \underline{\text{Lie}}(Y/S, \mathcal{M})).$$

For  $S' \rightarrow S$ , let  $u' : X' \rightarrow Y'$  be the base change of  $u$ . The  $S$ -functor<sup>57</sup>

$$\underline{\text{Hom}}_{(Y/S)\text{-gr.}}(X, \underline{\text{Lie}}(Y/S, \mathcal{M}) \cdot Y)$$

is defined by

$$\begin{aligned} & \underline{\text{Hom}}_{(Y/S)\text{-gr.}}(X, \underline{\text{Lie}}(Y/S, \mathcal{M}) \cdot Y)(S') \\ &= \text{Hom}_{Y'\text{-gr.}}(X', (\underline{\text{Lie}}(Y/S, \mathcal{M}) \cdot Y)_{S'}) \\ &= \text{Hom}_{Y'\text{-gr.}}(X', \underline{\text{Lie}}(Y'/S', \mathcal{M}) \cdot Y'). \end{aligned}$$

Thus  $(\dagger)$  induces an isomorphism

$$L_{\underline{\text{Hom}}_{S\text{-gr.}}(X,Y)/S}^u(\mathcal{M}) \xrightarrow{\sim} \underline{\text{Hom}}_{(Y/S)\text{-gr.}}(X, \underline{\text{Lie}}(Y/S, \mathcal{M}) \cdot Y). \quad (\dagger')$$

On the other hand, the composite

$$X \xrightarrow{u} Y \xrightarrow{\text{Ad}} \underline{\text{Aut}}_{S\text{-gr.}}(\underline{\text{Lie}}(Y/S, \mathcal{M}))$$

Figure 40: The action of  $X$  on  $\underline{\text{Lie}}(Y/S, \mathcal{M})$ .

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 27;  
combined-reader p. 85; printed p. 75.*

defines an action of  $X$  on  $L = \underline{\text{Lie}}(Y/S, \mathcal{M})$  by group automorphisms.

Let  $\Phi \in \underline{\text{Hom}}_{Y/S}(X, \underline{\text{Lie}}(Y/S, \mathcal{M}) \cdot Y)$ . For  $S'' \rightarrow S' \rightarrow S$  and  $x \in X(S'')$ , there is a unique expression

$$\Phi(S')(x) = \varphi(S')(x) \cdot u'(x), \quad \varphi(S')(x) \in \underline{\text{Lie}}(Y'/S', \mathcal{M})(S'').$$

This determines an element  $\varphi \in \underline{\text{Hom}}_S(X, \underline{\text{Lie}}(Y/S, \mathcal{M}))$ . The map  $\Phi(S')$  is a group morphism if and only if, for all  $x_1, x_2 \in X(S'')$ ,

$$\begin{aligned} \varphi(S')(x_1 x_2) &= \varphi(S')(x_1) (u(x_1) \varphi(S')(x_2) u(x_1)^{-1}) \\ &= \varphi(S')(x_1) (x_1 \cdot \varphi(S')(x_2)), \end{aligned}$$

that is, precisely when  $\varphi \in \underline{Z}_S^1(X, \underline{\text{Lie}}(Y/S, \mathcal{M}))$ .

<sup>57</sup>Editor's note: this is Definition 4.2.0.1 applied with  $Z = Y$ , to the  $Y$ -groups  $u : X \rightarrow Y$  and  $T_{Y/S}(\mathcal{M}) \rightarrow Y$ .

**Proposition 4.2.**

Let  $u : X \rightarrow Y$  be a morphism of  $S$ -groups. There is an isomorphism of  $S$ -functors, functorial in  $\mathcal{M}$ ,

$$L_{\underline{\text{Hom}}_{S\text{-gr.}}(X,Y)/S}^u(\mathcal{M}) \xrightarrow{\sim} \underline{Z}_S^1(X, \underline{\text{Lie}}(Y/S, \mathcal{M})).$$

Suppose in addition that  $Y/S$  satisfies  $E$ . It follows from 4.2.0.3, just as in the proof of 3.11.1, that  $\underline{\text{Hom}}_{S\text{-gr.}}(X, Y)/S$  satisfies  $E$ . Thus (cf. 3.5.1)

$$\begin{aligned} L_{\underline{\text{Hom}}_{S\text{-gr.}}(X,Y)/S}^u(\mathcal{M} \oplus \mathcal{N}) &\simeq L_{\underline{\text{Hom}}_{S\text{-gr.}}(X,Y)/S}^u(\mathcal{M}) \\ &\quad \times_S L_{\underline{\text{Hom}}_{S\text{-gr.}}(X,Y)/S}^u(\mathcal{N}). \end{aligned}$$

This also follows directly from Proposition 4.2 and Remark 4.2.Ba. Consequently, both sides of the isomorphism in Proposition 4.2 carry  $\mathcal{O}_S$ -module structures obtained from functoriality in  $\mathcal{M}$ , and that isomorphism respects them.

**Proposition 4.2 bis.**

Let  $u : X \rightarrow Y$  be a morphism of  $S$ -groups, and suppose that  $Y/S$  satisfies  $E$ . There is an isomorphism of  $\mathcal{O}_S$ -modules, functorial in  $\mathcal{M}$ ,

$$L_{\underline{\text{Hom}}_{S\text{-gr.}}(X,Y)/S}^u(\mathcal{M}) \xrightarrow{\sim} \underline{Z}_S^1(X, \underline{\text{Lie}}(Y/S, \mathcal{M})).$$

58

When  $Y/S$  satisfies  $E$ , 3.13.1 also gives, just as in the proof of 3.13.2, an isomorphism functorial in  $\mathcal{M}$ , for every  $u \in \underline{\text{Isom}}_{S\text{-gr.}}(X, Y)$ ,

$$L_{\underline{\text{Isom}}_{S\text{-gr.}}(X,Y)/S}^u(\mathcal{M}) \xrightarrow{\sim} L_{\underline{\text{Hom}}_{S\text{-gr.}}(X,Y)/S}^u(\mathcal{M}). \quad (*)$$

This yields the following two corollaries.

**Corollary 4.2.1.**

Let  $u : X \rightarrow Y$  be a morphism of  $S$ -groups. If  $Y/S$  satisfies  $E$ , there is an isomorphism of  $\mathcal{O}_S$ -modules, functorial in  $\mathcal{M}$ ,

$$L_{\underline{\text{Isom}}_{S\text{-gr.}}(X,Y)/S}^u(\mathcal{M}) \xrightarrow{\sim} \underline{Z}_S^1(X, \underline{\text{Lie}}(Y/S, \mathcal{M})).$$

**Corollary 4.2.2.**

Let  $X$  be an  $S$ -group. If  $X/S$  satisfies  $E$ , there is an isomorphism of  $\mathcal{O}_S$ -modules, functorial in  $\mathcal{M}$ ,

$$\underline{\text{Lie}}(\underline{\text{Aut}}_{S\text{-gr.}}(X)/S, \mathcal{M}) \xrightarrow{\sim} \underline{Z}_S^1(X, \underline{\text{Lie}}(X/S, \mathcal{M})).$$

If  $Y$  is commutative, its adjoint action on  $L = \underline{\text{Lie}}(Y/S, \mathcal{M})$  is trivial, so  $\underline{Z}_S^1(X, L) = \underline{\text{Hom}}_{S\text{-gr.}}(X, L)$ .<sup>59</sup>

**Corollary 4.2.3.**

Let  $Y$  be a commutative  $S$ -group. There is an isomorphism of  $S$ -functors, functorial in  $\mathcal{M}$ ,

$$L_{\underline{\text{Hom}}_{S\text{-gr.}}(X,Y)/S}^u(\mathcal{M}) \xrightarrow{\sim} \underline{\text{Hom}}_{S\text{-gr.}}(X, \underline{\text{Lie}}(Y/S, \mathcal{M})).$$

<sup>58</sup>Editor's note: the original has been expanded from this point; in particular, Proposition 4.2 bis has been added.

<sup>59</sup>Editor's note: the following sentence has been added.

## 4.3.

We now consider the case in which  $X$  and  $Y$  are  $\mathcal{O}_S$ -modules. Recall (cf. 4.2.0.4) that  $T'_{Y/S}(\mathcal{M})$ , respectively  $\underline{\text{Lie}}'(Y/S, \mathcal{M})$ , denotes the functor  $T_{Y/S}(\mathcal{M})$ , respectively  $\underline{\text{Lie}}(Y/S, \mathcal{M})$ , equipped with the  $\mathcal{O}_S$ -module structure induced by that of  $Y$ .

When  $Y/S$  satisfies  $E$ , we shall always write  $\underline{\text{Lie}}(Y/S, \mathcal{M})$  for this functor equipped instead with the  $\mathcal{O}_S$ -module structure defined for every functor satisfying  $E$ . By 3.9, the underlying abelian-group structures of  $\underline{\text{Lie}}(Y/S, \mathcal{M})$  and  $\underline{\text{Lie}}'(Y/S, \mathcal{M})$  coincide, but their module structures need not coincide (see the counterexample in 6.3). For  $S' \rightarrow S$ ,  $a \in \mathcal{O}(S')$ , and  $m \in \underline{\text{Lie}}(Y/S, \mathcal{M})(S')$ , we write  $a \cdot' m$ , respectively  $a \cdot m$ , for the action coming from  $Y$ , respectively the action obtained from condition  $E$ . We use the same convention for the two actions on  $T_{Y/S}(\mathcal{M})$ .

There is an isomorphism of  $\mathcal{O}_S$ -modules

$$T'_{Y/S}(\mathcal{M}) \simeq \underline{\text{Lie}}'(Y/S, \mathcal{M}) \oplus Y.$$

Exactly as for Propositions 4.2 and 4.2 bis, one therefore obtains:

**Proposition 4.3.**

Let  $u : X \rightarrow Y$  be a morphism of  $\mathcal{O}_S$ -modules. There is an isomorphism of  $S$ -functors, functorial in  $\mathcal{M}$ ,

$$L^u_{\underline{\text{Hom}}_{\mathcal{O}_S\text{-mod.}}(X, Y)/S}(\mathcal{M}) \xrightarrow{\sim} \underline{\text{Hom}}_{\mathcal{O}_S\text{-mod.}}(X, \underline{\text{Lie}}'(Y/S, \mathcal{M})). \quad (*)$$

If  $Y/S$  satisfies  $E$ , then  $\underline{\text{Hom}}_{\mathcal{O}_S\text{-mod.}}(X, Y)/S$  satisfies  $E$ , and  $(*)$  is an isomorphism of  $\mathcal{O}_S$ -modules when both sides carry the structures obtained from functoriality in  $\mathcal{M}$ .<sup>60 61</sup>

**Remark 4.3 bis.**

Let  $u : X \rightarrow Y$  be a morphism of  $\mathcal{O}_S$ -modules.<sup>62</sup> Denote by  $\tau_u$  the map that sends each  $\mathcal{O}_S$ -linear morphism  $\varphi : X \rightarrow \underline{\text{Lie}}'(Y/S, \mathcal{M})$  to

$$u \oplus \varphi : X \longrightarrow T'_{Y/S}(\mathcal{M}) = Y \oplus \underline{\text{Lie}}'(Y/S, \mathcal{M}).$$

The isomorphism of 4.3 belongs to the following commutative diagram, functorial in  $\mathcal{M}$ .

$$\begin{array}{ccc} L^u_{\underline{\text{Hom}}_{\mathcal{O}_S\text{-mod.}}(X, Y)/S}(\mathcal{M}) & \xrightarrow{\sim} & \underline{\text{Hom}}_{\mathcal{O}_S\text{-mod.}}(X, \underline{\text{Lie}}'(Y/S, \mathcal{M})) \\ \downarrow & & \downarrow \tau_u \\ T_{\underline{\text{Hom}}_{\mathcal{O}_S\text{-mod.}}(X, Y)/S}(\mathcal{M}) & \xrightarrow{\sim} & \underline{\text{Hom}}_{\mathcal{O}_S\text{-mod.}}(X, T'_{Y/S}(\mathcal{M})). \end{array}$$

Figure 41: Compatibility of the isomorphism of 4.3 with  $\tau_u$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp2-14oct24.pdf, component-local p. 30;  
combined-reader p. 88; printed p. 78.*

Moreover, when  $Y/S$  satisfies  $E$ , 3.13.1 gives, just as in the proof of 3.13.2, for every  $u \in \underline{\text{Isom}}_{\mathcal{O}_S\text{-mod.}}(X, Y)$ ,

$$L^u_{\underline{\text{Isom}}_{\mathcal{O}_S\text{-mod.}}(X, Y)/S}(\mathcal{M}) = L^u_{\underline{\text{Hom}}_{\mathcal{O}_S\text{-mod.}}(X, Y)/S}(\mathcal{M}). \quad (*)$$

<sup>60</sup>Editor's note: the following sentence has been added.

<sup>61</sup>Editor's note: on the right-hand side, the scalar action is  $(a\varphi)(x) = a\varphi(x)$ , where  $a\varphi(x)$  denotes the action of  $a$  on  $\varphi(x) \in \underline{\text{Lie}}(Y/S, \mathcal{M})$ . This differs from the action  $(a \cdot' \varphi)(x) = a \cdot' \varphi(x) = \varphi(ax)$ , but the two agree when  $\underline{\text{Lie}}(Y/S, \mathcal{M}) = \underline{\text{Lie}}'(Y/S, \mathcal{M})$ .

<sup>62</sup>Editor's note: this remark has been added; it will be useful in 4.5.1.



**Corollary 4.3.1.**

Let  $X$  be an  $\mathcal{O}_S$ -module satisfying  $E$  over  $S$ . There is an isomorphism, functorial in  $\mathcal{M}$ ,

$$\underline{\mathrm{Lie}}(\underline{\mathrm{Aut}}_{\mathcal{O}_S\text{-mod.}}(X)/S, \mathcal{M}) \xrightarrow{\sim} \underline{\mathrm{Hom}}_{\mathcal{O}_S\text{-mod.}}(X, \underline{\mathrm{Lie}}'(X/S, \mathcal{M})),$$

and this isomorphism respects the  $\mathcal{O}_S$ -module structures obtained from functoriality in  $\mathcal{M}$ . In particular,  $\underline{\mathrm{Aut}}_{\mathcal{O}_S\text{-mod.}}(X)/S$  satisfies  $E$ .<sup>63</sup>

**Proof.** The first assertion follows from  $(*)$  and 4.3. For the second, since  $X/S$  satisfies  $E$ , there are isomorphisms of  $\mathcal{O}_S$ -modules

$$\underline{\mathrm{Lie}}'(X/S, \mathcal{M} \oplus \mathcal{N}) \simeq \underline{\mathrm{Lie}}'(X/S, \mathcal{M}) \times_S \underline{\mathrm{Lie}}'(X/S, \mathcal{N}).$$

Consequently,

$$\begin{aligned} & \underline{\mathrm{Lie}}(\underline{\mathrm{Aut}}_{\mathcal{O}_S\text{-mod.}}(X)/S, \mathcal{M} \oplus \mathcal{N}) \\ & \simeq \underline{\mathrm{Lie}}(\underline{\mathrm{Aut}}_{\mathcal{O}_S\text{-mod.}}(X)/S, \mathcal{M}) \times_S \underline{\mathrm{Lie}}(\underline{\mathrm{Aut}}_{\mathcal{O}_S\text{-mod.}}(X)/S, \mathcal{N}). \end{aligned}$$

Remark 4.1.1.2a now proves the assertion.

**4.3.2.**

Before continuing in this direction, let us examine more closely the relations among  $Y$ ,  $\underline{\mathrm{Lie}}(Y/S)$ , and  $\underline{\mathrm{Lie}}'(Y/S)$ . First,

$$\underline{\mathrm{Lie}}(\mathcal{O}_S/S, \mathcal{M}) = \underline{\mathrm{Lie}}'(\mathcal{O}_S/S, \mathcal{M}) = \mathbf{W}(\mathcal{M}), \quad (1)$$

where  $\mathbf{W}(\mathcal{M})$  is defined in I 4.6. There is therefore a canonical isomorphism

$$d : \mathcal{O}_S \xrightarrow{\sim} \underline{\mathrm{Lie}}(\mathcal{O}_S/S). \quad (2)$$

Let  $F$  be an  $\mathcal{O}_S$ -module. For every  $S_2 \rightarrow S_1 \rightarrow S$ , there is a dihomomorphism

$$\begin{cases} F(S_1) \longrightarrow F(S_2), \\ \mathcal{O}(S_1) \longrightarrow \mathcal{O}(S_2), \end{cases} \quad (3)$$

and hence a morphism of  $\mathcal{O}(S_2)$ -modules

$$F(S_1) \otimes_{\mathcal{O}(S_1)} \mathcal{O}(S_2) \longrightarrow F(S_2).$$

<sup>64</sup>

In particular, take  $S_1 = S'$  and  $S_2 = I_{S'}(\mathcal{M})$ . One obtains morphisms of  $\mathcal{O}(S')$ -modules, functorial in  $\mathcal{M}$ ,

$$F(S') \otimes_{\mathcal{O}(S')} T_{\mathcal{O}_S/S}(\mathcal{M})(S') \longrightarrow T'_{F/S}(\mathcal{M})(S').$$

As  $S'$  varies, these give morphisms of  $\mathcal{O}_S$ -modules

$$F \otimes_{\mathcal{O}_S} T_{\mathcal{O}_S/S}(\mathcal{M}) \longrightarrow T'_{F/S}(\mathcal{M}), \quad (4)$$

functorial in  $\mathcal{M}$ . Compatibility with the tangent-bundle projections onto their bases then yields morphisms

$$F \otimes_{\mathcal{O}_S} \underline{\mathrm{Lie}}(\mathcal{O}_S/S, \mathcal{M}) \longrightarrow \underline{\mathrm{Lie}}'(F/S, \mathcal{M}), \quad (5)$$

<sup>63</sup>Editor's note: compare editor's note (61). The final sentence and its proof have also been added.

<sup>64</sup>Editor's note:  $S' \rightarrow S$  has been replaced by  $S_2 \rightarrow S_1 \rightarrow S$ , since both  $S_2$  and  $S_1$  must vary below. The original has also been expanded in what follows.

and the following diagram commutes.

$$\begin{array}{ccccccc}
 0 & \longrightarrow & F \otimes_{\mathcal{O}_S} \underline{\mathrm{Lie}}(\mathcal{O}_S/S, \mathcal{M}) & \longrightarrow & F \otimes_{\mathcal{O}_S} T_{\mathcal{O}_S/S}(\mathcal{M}) & \longrightarrow & F \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \parallel \\
 0 & \longrightarrow & \underline{\mathrm{Lie}}'(F/S, \mathcal{M}) & \longrightarrow & T'_{F/S}(\mathcal{M}) & \longrightarrow & F \longrightarrow 0.
 \end{array}$$

Figure 42: The morphisms (4) and (5) in the split tangent sequences.

*Temporary Loop-1 image; source locator: Polo-Gille Exp2-14oct24.pdf, component-local p. 31;  
combined-reader p. 89; printed p. 79.*

The morphisms (5) may be regarded as morphisms of abelian  $S$ -groups

$$F \otimes_{\mathcal{O}_S} \underline{\mathrm{Lie}}(\mathcal{O}_S/S, \mathcal{M}) \longrightarrow \underline{\mathrm{Lie}}(F/S, \mathcal{M}). \quad (6)$$

Tensoring  $F$  with the isomorphism  $d : \mathcal{O}_S \xrightarrow{\sim} \underline{\mathrm{Lie}}(\mathcal{O}_S/S)$  gives, when  $\mathcal{M} = \mathcal{O}_S$ , a morphism of abelian  $S$ -groups

$$F \xrightarrow{\sim} F \otimes_{\mathcal{O}_S} \underline{\mathrm{Lie}}(\mathcal{O}_S/S) \longrightarrow \underline{\mathrm{Lie}}(F/S), \quad (7)$$

also denoted  $d : F \rightarrow \underline{\mathrm{Lie}}(F/S)$ .

### Remark 4.3.3.

When  $F/S$  satisfies  $E$ , the morphisms (6) and (7) need not be  $\mathcal{O}_S$ -linear if both sides are equipped with the module structures obtained from  $\mathcal{M}$  by condition  $E$ .

For example, let  $k$  be a field of characteristic  $p > 0$ , let  $S = \mathrm{Spec}(k)$ , and let  $F$  be the  $\mathcal{O}_S$ -module that assigns to every  $S$ -scheme  $T$  the set  $F(T) = \Gamma(T, \mathcal{O}_T)$ , with scalars acting through the  $p$ -th power:

$$r \cdot f = r^p f \quad (r \in \mathcal{O}(T), f \in F(T)).$$

As an  $S$ -functor in groups,  $F \simeq \mathbf{G}_{a,S}$ . Hence  $F$  satisfies  $E$  and  $\underline{\mathrm{Lie}}(F/S) \simeq \underline{\mathrm{Lie}}(\mathbf{G}_{a,S}/S) \simeq \mathcal{O}_S$ . For every  $T \rightarrow S$ , the canonical morphism  $d : F(T) \rightarrow \mathcal{O}(T)$  is the identity: it respects the abelian-group structure but not the  $\mathcal{O}_S$ -module structure. <sup>65</sup>

### Remark 4.3.4.

The morphisms (4) and (5) can be made explicit as follows. The morphism

$$\Theta : F \otimes_{\mathcal{O}_S} T_{\mathcal{O}_S/S}(\mathcal{M}) \longrightarrow T'_{F/S}(\mathcal{M}) = \underline{\mathrm{Hom}}_S(I_S(\mathcal{M}), F)$$

is defined, for  $S' \rightarrow S$ ,  $\alpha \in \mathcal{O}(I_{S'}(\mathcal{M}))$ , and  $f : S' \rightarrow F$ , by

$$\Theta(f \otimes \alpha) = \alpha \cdot (\tau_0 \circ f) = \alpha \cdot (f \circ \rho),$$

where  $\tau_0 : F \rightarrow T'_{F/S}(\mathcal{M})$  is the zero section and  $\rho : I_{S'}(\mathcal{M}) \rightarrow S'$  is the structural morphism.

The morphism  $\Theta$  induces

$$\theta : F \otimes_{\mathcal{O}_S} \underline{\mathrm{Lie}}(\mathcal{O}_S/S, \mathcal{M}) \longrightarrow \underline{\mathrm{Lie}}'(F/S, \mathcal{M}).$$

<sup>65</sup>Editor's note: the original asserted that, when  $F/S$  satisfies  $E$ , (6), and hence (7), are morphisms of  $\mathcal{O}_S$ -modules. The counterexample in 6.3 contradicts this. That assertion has been removed, the preceding example inserted, and Definition 4.4 modified accordingly. This does not affect what follows.

This follows from the functoriality in  $\mathcal{M}$  noted after (4), and can also be seen directly. Indeed,

$$\underline{\mathrm{Lie}}'(F/S, \mathcal{M})(S') = \{ \varphi \in \mathrm{Hom}_S(I_{S'}(\mathcal{M}), F) \mid \varphi \circ \varepsilon_{\mathcal{M}} = e \},$$

where  $e : S' \rightarrow F$  is the unit section. On the other hand,  $\underline{\mathrm{Lie}}(\mathcal{O}_S/S, \mathcal{M})(S')$  equals  $\Gamma(S', \mathcal{M})$ , the kernel of the augmentation

$$\eta : \mathcal{O}(I_{S'}(\mathcal{M})) \longrightarrow \mathcal{O}(S').$$

It remains to see that if  $f \in F(S')$  and  $\alpha \in \Gamma(S', \mathcal{M})$ , then  $\alpha \cdot (f \circ \rho) \circ \varepsilon_{\mathcal{M}} = e$ .

Apply the dihomomorphism (3) to the  $S$ -morphism  $\varepsilon_{\mathcal{M}} : S' \rightarrow I_{S'}(\mathcal{M})$ . The following diagram commutes.

$$\begin{array}{ccc} F(I_{S'}(\mathcal{M})) & \xrightarrow{F(\varepsilon_{\mathcal{M}})} & F(S') \\ \alpha \downarrow & & \downarrow \eta(\alpha) \\ F(I_{S'}(\mathcal{M})) & \xrightarrow{F(\varepsilon_{\mathcal{M}})} & F(S') \end{array} .$$

Figure 43: Compatibility of scalar action with  $F(\varepsilon_{\mathcal{M}})$ .

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 32;  
combined-reader p. 90; printed p. 80.*

Thus, for every  $\varphi : I_{S'}(\mathcal{M}) \rightarrow F$ ,

$$(\alpha \cdot \varphi) \circ \varepsilon_{\mathcal{M}} = \eta(\alpha) \cdot (\varphi \circ \varepsilon_{\mathcal{M}}),$$

and therefore  $(\alpha \cdot \varphi) \circ \varepsilon_{\mathcal{M}} = e$  when  $\eta(\alpha) = 0$ .

In particular, taking  $\mathcal{M} = t\mathcal{O}_S$ , one has  $\underline{\mathrm{Lie}}(\mathcal{O}_S/S) = t\mathcal{O}_S$ , and the morphism  $F \rightarrow \underline{\mathrm{Lie}}'(F/S)$  is  $f \mapsto t \cdot (f \circ \rho)$ .

<sup>66</sup>

### Remark 4.3.5.

Let  $F$  be an  $\mathcal{O}_S$ -module, set

$$E = \underline{\mathrm{End}}_{\mathcal{O}_S\text{-mod.}}(F),$$

and denote by  $d_F$  and  $d_E$  the  $\mathcal{O}_S$ -linear morphisms given by 4.3.2(5):

$$d_F : F \otimes_{\mathcal{O}_S} \underline{\mathrm{Lie}}(\mathcal{O}_S/S, \mathcal{M}) \longrightarrow \underline{\mathrm{Lie}}'(F/S, \mathcal{M}),$$

$$d_E : E \otimes_{\mathcal{O}_S} \underline{\mathrm{Lie}}(\mathcal{O}_S/S, \mathcal{M}) \longrightarrow \underline{\mathrm{Lie}}'(E/S, \mathcal{M}).$$

Remark 4.3.4 gives the following commutative diagram of  $\mathcal{O}_S$ -linear morphisms.

$$\begin{array}{ccc} E \otimes_{\mathcal{O}_S} \underline{\mathrm{Lie}}(\mathcal{O}_S/S, \mathcal{M}) & \xrightarrow{d_E} & \underline{\mathrm{Lie}}'(E/S, \mathcal{M}) \\ \parallel & & \uparrow \simeq (*) \\ \underline{\mathrm{Hom}}_{\mathcal{O}_S}(F, F \otimes_{\mathcal{O}_S} \underline{\mathrm{Lie}}(\mathcal{O}_S/S, \mathcal{M})) & \xrightarrow{d_F} & \underline{\mathrm{Hom}}_{\mathcal{O}_S}(F, \underline{\mathrm{Lie}}'(F/S, \mathcal{M})) \end{array}$$

Figure 44: Compatibility of  $d_E$  and  $d_F$ .

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 32;  
combined-reader p. 90; printed p. 80.*

<sup>66</sup>Editor's note: this remark has been added; it will be useful in 4.6.2.

The right vertical arrow is the isomorphism  $(*)$  of 4.3. Thus, by that result, if  $F/S$  satisfies  $E$ , then  $E/S$  satisfies  $E$ , and  $(*)$  is also an isomorphism of  $\mathcal{O}_S$ -modules when the right-hand terms are equipped with the module structures obtained from  $E$ .<sup>67</sup>

**Remark 4.4.0.**

In 4.3.2, the morphisms (4) are isomorphisms if and only if the morphisms (5) are. If these equivalent conditions hold, then  $F/S$  satisfies  $E$ . Indeed, it is enough to verify that

$$\underline{\mathrm{Lie}}'(F/S, \mathcal{M} \oplus \mathcal{N}) \simeq \underline{\mathrm{Lie}}'(F/S, \mathcal{M}) \times_S \underline{\mathrm{Lie}}'(F/S, \mathcal{N}).$$

Under the hypothesis, the horizontal arrows in the following commutative diagram are isomorphisms.

$$\begin{array}{ccc} F \otimes_{\mathcal{O}_S} \underline{\mathrm{Lie}}(\mathcal{O}_S/S, \mathcal{M} \oplus \mathcal{N}) & \xrightarrow{\sim} & \underline{\mathrm{Lie}}'(F/S, \mathcal{M} \oplus \mathcal{N}) \\ \downarrow \wr & & \downarrow \\ F \otimes_{\mathcal{O}_S} (\underline{\mathrm{Lie}}(\mathcal{O}_S/S, \mathcal{M}) \times_S \underline{\mathrm{Lie}}(\mathcal{O}_S/S, \mathcal{N})) & \xrightarrow{\sim} & \underline{\mathrm{Lie}}'(F/S, \mathcal{M}) \times_S \underline{\mathrm{Lie}}'(F/S, \mathcal{N}); \end{array}$$

Figure 45: The direct-sum comparison used to verify condition  $E$ .

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 33;  
combined-reader p. 91; printed p. 81.*

The second vertical arrow is therefore an isomorphism as well, which is exactly condition  $E$  for  $F/S$ .<sup>68</sup>

**Definition 4.4.**

An  $\mathcal{O}_S$ -module  $F$  is called *good* if the morphisms

$$F \otimes_{\mathcal{O}_S} T_{\mathcal{O}_S/S}(\mathcal{M}) \longrightarrow T_{F/S}(\mathcal{M}),$$

or, equivalently,

$$F \otimes_{\mathcal{O}_S} \underline{\mathrm{Lie}}(\mathcal{O}_S/S, \mathcal{M}) \longrightarrow \underline{\mathrm{Lie}}(F/S, \mathcal{M}),$$

are isomorphisms of abelian  $S$ -groups (so that  $F/S$  satisfies (E)), and if moreover they respect the  $\mathcal{O}_S$ -module structures obtained from condition  $E$ .

**Corollary 4.4.1.**

Let  $F$  be an  $\mathcal{O}_S$ -module, and consider the following conditions:<sup>69</sup>

- (i)  $F$  is a good  $\mathcal{O}_S$ -module;
- (ii)  $F/S$  satisfies  $E$ , and  $d : F \rightarrow \underline{\mathrm{Lie}}(F/S)$  is an isomorphism of  $\mathcal{O}_S$ -modules;
- (iii)  $\underline{\mathrm{Lie}}(F/S, \mathcal{M}) = \underline{\mathrm{Lie}}'(F/S, \mathcal{M})$ .

Then

$$(i) \iff (ii) \implies (iii).$$

<sup>67</sup>Editor's note: this remark has been added; it will be useful in 4.5 and 4.7.

<sup>68</sup>Editor's note: in view of the changes made in Definition 4.4 (cf. editor's note (65)), this remark has been inserted here; in the original it occurred in the proof of 4.4.1.

<sup>69</sup>Editor's note: the original proved that (i) implies (ii) and (iii); the implication (ii) implies (i) has been added.

**Proof.** The implication (i)  $\Rightarrow$  (ii) follows from the definition. To prove (ii)  $\Rightarrow$  (i), it remains to show that the morphisms of abelian  $S$ -groups, functorial in  $\mathcal{M}$ ,

$$F \otimes_{\mathcal{O}_S} \underline{\mathrm{Lie}}(\mathcal{O}_S/S, \mathcal{M}) \xrightarrow{\sim} \underline{\mathrm{Lie}}(F/S, \mathcal{M})$$

are isomorphisms of  $\mathcal{O}_S$ -modules. Since  $F/S$  satisfies  $E$ , both sides carry finite direct sums of copies of  $\mathcal{O}_S$  to finite products of abelian  $S$ -groups. We are therefore reduced to the case  $\mathcal{M} = \mathcal{O}_S$ , which is precisely the hypothesis.

Finally, (i)  $\Rightarrow$  (iii) follows from the definition and from the fact that the isomorphisms

$$F \otimes_{\mathcal{O}_S} \underline{\mathrm{Lie}}(\mathcal{O}_S/S, \mathcal{M}) \xrightarrow{\sim} \underline{\mathrm{Lie}}'(F/S, \mathcal{M})$$

of 4.3.2(5) are  $\mathcal{O}_S$ -linear.

### Examples 4.4.2.

For every quasi-coherent  $\mathcal{O}_S$ -module  $\mathcal{E}$ , the  $\mathcal{O}_S$ -modules  $\mathbf{V}(\mathcal{E})$  and  $\mathbf{W}(\mathcal{E})$  defined in I 4.6 are good.

Indeed, for every  $f : S' \rightarrow S$ , the morphisms

$$\begin{aligned} \mathbf{V}(\mathcal{E})(S') \otimes_{\mathcal{O}(S')} \mathcal{O}(I_{S'}(\mathcal{M})) &\longrightarrow T_{\mathbf{V}(\mathcal{E})/S}(\mathcal{M})(S'), \\ \mathbf{W}(\mathcal{E})(S') \otimes_{\mathcal{O}(S')} \mathcal{O}(I_{S'}(\mathcal{M})) &\longrightarrow T_{\mathbf{W}(\mathcal{E})/S}(\mathcal{M})(S') \end{aligned}$$

correspond respectively to

$$\begin{aligned} \mathrm{Hom}_{\mathcal{O}_{S'}\text{-mod.}}(f^*(\mathcal{E}), \mathcal{O}_{S'}) \otimes_{\mathcal{O}(S')} \Gamma(S', D_{\mathcal{O}_{S'}}(\mathcal{M})) \\ \longrightarrow \mathrm{Hom}_{\mathcal{O}_{S'}\text{-mod.}}(f^*(\mathcal{E}), D_{\mathcal{O}_{S'}}(\mathcal{M})) \quad (\mathrm{V}) \\ \Gamma(S', f^*(\mathcal{E})) \otimes_{\mathcal{O}(S')} \Gamma(S', D_{\mathcal{O}_{S'}}(\mathcal{M})) \\ \longrightarrow \Gamma(S', f^*(\mathcal{E}) \otimes_{\mathcal{O}_{S'}} D_{\mathcal{O}_{S'}}(\mathcal{M})) \quad (\mathrm{W}). \end{aligned}$$

These are isomorphisms because, as an  $\mathcal{O}_{S'}$ -module,  $D_{\mathcal{O}_{S'}}(\mathcal{M})$  is isomorphic to a finite direct sum of copies of  $\mathcal{O}_{S'}$ .<sup>70</sup>

**Proposition 4.5.** *Let  $F$  be a good  $\mathcal{O}_S$ -module. Then:*

(i) *The  $S$ -group  $\underline{\mathrm{Aut}}_{\mathcal{O}_S\text{-mod.}}(F)$  satisfies condition (E), and there is a functorial isomorphism*

$$\underline{\mathrm{Lie}}(\underline{\mathrm{Aut}}_{\mathcal{O}_S\text{-mod.}}(F)/S, \mathcal{M}) \xrightarrow{\sim} \underline{\mathrm{Hom}}_{\mathcal{O}_S\text{-mod.}}(F, \underline{\mathrm{Lie}}(F/S, \mathcal{M})),$$

*which respects the  $\mathcal{O}_S$ -module structures obtained from condition (E). In particular, there is an isomorphism of  $\mathcal{O}_S$ -modules*

$$\underline{\mathrm{Lie}}(\underline{\mathrm{Aut}}_{\mathcal{O}_S\text{-mod.}}(F)/S) \xrightarrow{\sim} \underline{\mathrm{End}}_{\mathcal{O}_S\text{-mod.}}(F).$$

(ii)<sup>71</sup> *Moreover,  $\underline{\mathrm{End}}_{\mathcal{O}_S\text{-mod.}}(F)$  is a good  $\mathcal{O}_S$ -module.*

Indeed, by 4.4.1,  $F/S$  satisfies (E), and

$$\underline{\mathrm{Lie}}(F/S, \mathcal{M}) = \underline{\mathrm{Lie}}'(F/S, \mathcal{M}) \simeq F \otimes_{\mathcal{O}_S} \underline{\mathrm{Lie}}(\mathcal{O}_S/S, \mathcal{M}). \quad (1)$$

Assertion (i) now follows from 4.3.1. Put  $E = \underline{\mathrm{End}}_{\mathcal{O}_S\text{-mod.}}(F)$ . By (1) and Remark 4.3.5, one has the following commutative diagram of morphisms of abelian  $S$ -groups.

<sup>70</sup>Editor's note: the preceding proof has been added.

<sup>71</sup>Editor's note: part (ii), an immediate consequence of what precedes, has been added.

$$\begin{array}{ccc}
\underline{\text{End}}_{\mathcal{O}_S\text{-mod.}}(F) \otimes_{\mathcal{O}_S} \underline{\text{Lie}}(\mathcal{O}_S/S, \mathcal{M}) & \xrightarrow{d_E} & \underline{\text{Lie}}(\underline{\text{End}}_{\mathcal{O}_S\text{-mod.}}(F)/S, \mathcal{M}) \\
\parallel & & \simeq \uparrow (*) \\
\underline{\text{Hom}}_{\mathcal{O}_S}(F, F \otimes_{\mathcal{O}_S} \underline{\text{Lie}}(\mathcal{O}_S/S, \mathcal{M})) & \xrightarrow{d_F} & \underline{\text{Hom}}_{\mathcal{O}_S}(F, \underline{\text{Lie}}(F/S, \mathcal{M}))
\end{array}$$

Figure 46: The comparison diagram for the differential  $d_E$ .

Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 34;  
combined-reader p. 92; printed p. 82.

Here  $d_F$  and the vertical isomorphism  $(*)$  are isomorphisms of  $\mathcal{O}_S$ -modules; consequently, so is  $d_E$ . This proves (ii).

**Scholium 4.5.1.** <sup>72</sup> Put (cf. 2.1)  $\mathcal{O}_{I_S} = \mathcal{O}_S \oplus t\mathcal{O}_S$ , with  $t^2 = 0$ , and let  $F$  be a good  $\mathcal{O}_S$ -module. For every  $S' \rightarrow S$ , the morphism

$$F(S') \oplus tF(S') = F(S') \otimes_{\mathcal{O}(S')} \mathcal{O}(I_{S'}) \longrightarrow F(I_{S'}) = F(S') \oplus \underline{\text{Lie}}(F/S)(S'),$$

which is the identity on  $F(S')$ , induces an isomorphism of  $\mathcal{O}(S')$ -modules

$$tF(S') \xrightarrow{\sim} \underline{\text{Lie}}(F/S)(S').$$

As  $S'$  varies, this gives an isomorphism that we may write

$$\underline{\text{Lie}}(F/S) \simeq tF.$$

Thus, for every  $S' \rightarrow S$ , Proposition 4.5 gives a commutative diagram.

$$\begin{array}{ccccc}
\underline{\text{End}}_{\mathcal{O}_{S'}\text{-mod.}}(F_{S'}) & \xrightarrow{\sim} & \underline{\text{Hom}}_{\mathcal{O}_{S'}\text{-mod.}}(F_{S'}, tF_{S'}) & \xrightarrow{\sim} & \underline{\text{Lie}}(\underline{\text{Aut}}_{\mathcal{O}_S\text{-mod.}}(F)/S)(S') \\
& & \downarrow & & \downarrow \\
& & \underline{\text{Aut}}_{\mathcal{O}_{I_{S'}}\text{-mod.}}(F_{I_{S'}}) & \xlongequal{\quad} & T_{\underline{\text{Aut}}_{\mathcal{O}_S\text{-mod.}}(F)/S}(S')
\end{array}$$

Figure 47: Endomorphisms and automorphisms over the dual numbers.

Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 35;  
combined-reader p. 93; printed p. 83.

It follows from 4.3 bis that every

$$X \in \underline{\text{End}}_{\mathcal{O}_{S'}\text{-mod.}}(F_{S'})$$

corresponds to  $\text{id} + tX \in \underline{\text{Aut}}_{\mathcal{O}_{I_{S'}}\text{-mod.}}(F_{I_{S'}})$ .

**Definition 4.6.** An  $S$ -functor in groups  $G$  is called *good* if  $G/S$  satisfies condition (E) and  $\underline{\text{Lie}}(G/S)$  is a good  $\mathcal{O}_S$ -module.

<sup>73</sup> Notice that if  $F$  is a good  $\mathcal{O}_S$ -module, then it is a good  $S$ -group: indeed,  $F/S$  satisfies (E), and  $\underline{\text{Lie}}(F/S) \simeq F$  is a good  $\mathcal{O}_S$ -module.

<sup>72</sup>Editor's note: this scholium, implicit in the original, has been added for use in 4.7.1. Here and below,  $t, t'$ , and so forth denote square-zero variables; the letter  $\varepsilon$  is reserved for the unit section of a group.

<sup>73</sup>Editor's note: the following sentence has been added.

**Example 4.6.1.** *If  $G$  is representable, it is good. Indeed,  $G/S$  satisfies (E), and  $\underline{\mathrm{Lie}}(G/S)$  has the form  $V(E)$ , hence is good by 4.4.2.*

**Lemma 4.6.2.** <sup>74</sup> *Let  $G$  be an  $S$ -functor in groups such that  $G/S$  satisfies (E), and put  $F = \underline{\mathrm{Lie}}(G/S)$ . Then  $F$  satisfies (E), and the morphism of abelian groups*

$$d : F \longrightarrow \underline{\mathrm{Lie}}(F/S)$$

*respects the  $\mathcal{O}_S$ -module structures. Consequently,  $G$  is good if and only if  $F \rightarrow \underline{\mathrm{Lie}}(F/S)$  is bijective.*

*Proof.* Suppose that  $G/S$  satisfies (E). Let  $\mathcal{M}, \mathcal{N}$  be finite free  $\mathcal{O}_S$ -modules, write

$$F(\mathcal{N}) = \underline{\mathrm{Lie}}(G/S, \mathcal{N}),$$

and let  $e$  denote the unit section of  $G$ . For every  $S' \rightarrow S$ ,

$$F(\mathcal{N})(S') = \{g \in \mathrm{Hom}_S(I_{S'}(\mathcal{N}), G) \mid g \circ \varepsilon_{\mathcal{N}} = e\}.$$

Moreover,  $\underline{\mathrm{Lie}}(F(\mathcal{N})/S, \mathcal{M})(S')$  identifies with the set of morphisms

$$\left\{ \Phi \in \mathrm{Hom}_S(I_{S'}(\mathcal{N}) \times_{S'} I_{S'}(\mathcal{M}), G) \left| \begin{array}{l} \Phi \circ (\varepsilon_{\mathcal{N}} \times \mathrm{id}_{I_{S'}(\mathcal{M})}) = e \in G(I_{S'}(\mathcal{M})), \\ \Phi \circ (\mathrm{id}_{I_{S'}(\mathcal{N})} \times \varepsilon_{\mathcal{M}}) = e \in G(I_{S'}(\mathcal{N})). \end{array} \right. \right\}.$$

This proves

$$\underline{\mathrm{Lie}}(F(\mathcal{N})/S, \mathcal{M}) \simeq \underline{\mathrm{Lie}}(F(\mathcal{M})/S, \mathcal{N}). \quad (1)$$

Since  $G/S$  satisfies (E), it follows that

$$\begin{aligned} \underline{\mathrm{Lie}}(F(\mathcal{N})/S, \mathcal{M}_1 \oplus \mathcal{M}_2) &\simeq \underline{\mathrm{Lie}}(F(\mathcal{M}_1 \oplus \mathcal{M}_2)/S, \mathcal{N}) \\ &\simeq \underline{\mathrm{Lie}}((F(\mathcal{M}_1) \times_S F(\mathcal{M}_2))/S, \mathcal{N}). \end{aligned}$$

By 3.8, the last term is isomorphic to

$$\begin{aligned} &\underline{\mathrm{Lie}}(F(\mathcal{M}_1)/S, \mathcal{N}) \times_S \underline{\mathrm{Lie}}(F(\mathcal{M}_2)/S, \mathcal{N}) \\ &\simeq \underline{\mathrm{Lie}}(F(\mathcal{N})/S, \mathcal{M}_1) \times_S \underline{\mathrm{Lie}}(F(\mathcal{N})/S, \mathcal{M}_2). \end{aligned}$$

Hence

$$\underline{\mathrm{Lie}}(F(\mathcal{N})/S, \mathcal{M}_1 \oplus \mathcal{M}_2) \simeq \underline{\mathrm{Lie}}(F(\mathcal{N})/S, \mathcal{M}_1) \times_S \underline{\mathrm{Lie}}(F(\mathcal{N})/S, \mathcal{M}_2), \quad (2)$$

so  $F(\mathcal{N})/S$  satisfies (E), by Remark 4.1.1.2(a).

It remains to show that the morphism of abelian groups

$$d : F(\mathcal{N}) \longrightarrow \underline{\mathrm{Lie}}(F(\mathcal{N})/S)$$

respects the  $\mathcal{O}_S$ -module structures. Consider the free  $\mathcal{O}_S$ -module  $\mathcal{M} = t\mathcal{O}_S$ . Then

$$\mathcal{O}(I_{S'}(\mathcal{M})) = \mathcal{O}(S')[t]/(t^2) = \mathcal{O}(S') \oplus t\mathcal{O}(S'), \quad \underline{\mathrm{Lie}}(\mathbf{O}_S/S) \simeq t\mathcal{O}_S.$$

Let  $\rho_t : I_S(\mathcal{M}) \rightarrow S$  be the structural morphism, corresponding to the injection

$$u_t : \mathcal{O}_S \hookrightarrow D_{\mathcal{O}_S}(\mathcal{M}) = \mathcal{O}_S \oplus t\mathcal{O}_S.$$

---

<sup>74</sup>Editor's note: this lemma has been added.

Recall (cf. 3.4.2) that for every  $S$ -functor  $X$ ,  $F(\mathcal{N})(X)$  is the set of  $S$ -morphisms  $\phi : I_X(\mathcal{N}) \rightarrow G$  such that  $\phi \circ \varepsilon_{\mathcal{N}} = e$ , and the action of  $a \in \mathcal{O}(X)$  is

$$a \cdot \phi = \phi \circ a^*,$$

where  $a^*$  is the endomorphism of  $I_X(\mathcal{N})$  associated with  $a$ ; see 2.1.3.

Consequently, by 4.3.4, the composite

$$d : F(\mathcal{N}) \xrightarrow{\sim} F(\mathcal{N}) \otimes_{\mathcal{O}_S} t\mathcal{O}_S \longrightarrow \underline{\text{Lie}}(F(\mathcal{N})/S)$$

is given, for  $S' \rightarrow S$  and  $f \in \text{Hom}_S(S', F(\mathcal{N}))$ , by

$$f \longmapsto f \otimes t \longmapsto t \cdot (f \circ \rho_t) = f \circ \rho_t \circ t^*.$$

Here  $f \circ \rho_t \in F(\mathcal{N})(I_{S'}(\mathcal{M}))$  is viewed as an  $S$ -morphism

$$\phi : I_{I_{S'}(\mathcal{M})}(\mathcal{N}) \longrightarrow G$$

such that  $\phi \circ \varepsilon_{\mathcal{N}} = e$ . We must prove that  $d$  is  $\mathcal{O}_S$ -linear, or equivalently that

$$d(a \cdot f) = u_t(a) \cdot d(f) \quad (a \in \mathcal{O}(S')).$$

Viewing  $f \in F(\mathcal{N})(S')$  as a morphism  $I_{S'}(\mathcal{N}) \rightarrow G$ , one likewise has  $a \cdot f = f \circ a^*$ . Functoriality in  $X$  of the action of  $\mathcal{O}(X)$  on  $I_X(\mathcal{N})$  gives the following commutative diagram.

$$\begin{array}{ccc} I_{S'}(\mathcal{N}) & \xrightarrow{a^*} & I_{S'}(\mathcal{N}) \\ \rho_t \uparrow & & \uparrow \rho_t \\ I_{I_{S'}(\mathcal{M})}(\mathcal{N}) & \xrightarrow{u_t(a)^*} & I_{I_{S'}(\mathcal{M})}(\mathcal{N}) \end{array} .$$

Figure 48: Compatibility of scalar action with the structural morphism  $\rho_t$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp2-14oct24.pdf, component-local p. 36;  
combined-reader p. 94; printed p. 84.*

Therefore

$$\begin{aligned} d(a \cdot f) &= f \circ a^* \circ \rho_t \circ t^* \\ &= f \circ \rho_t \circ u_t(a)^* \circ t^* \\ &= f \circ \rho_t \circ t^* \circ u_t(a)^* = u_t(a) \cdot d(f). \end{aligned}$$

The penultimate equality follows from the commutativity of  $\mathcal{O}$ . This completes the proof of Lemma 4.6.2.

**Theorem 4.7.** *If  $F$  is a good  $\mathcal{O}_S$ -module, then the  $S$ -group  $\underline{\text{Aut}}_{\mathcal{O}_S\text{-mod.}}(F)$  is good.*

<sup>75</sup> Indeed, by 4.5,  $\underline{\text{Aut}}_{\mathcal{O}_S\text{-mod.}}(F)/S$  satisfies (E), while

$$\underline{\text{Lie}}(\underline{\text{Aut}}_{\mathcal{O}_S\text{-mod.}}(F)/S) \simeq \underline{\text{End}}_{\mathcal{O}_S\text{-mod.}}(F)$$

is a good  $\mathcal{O}_S$ -module.

<sup>75</sup>Editor's note: the original has been simplified here by using the additions made in 4.3.1 and 4.5.



**4.7.1.**

<sup>76</sup> Let  $G$  be an  $S$ -group and  $F$  a good  $\mathcal{O}_S$ -module. Suppose given a linear representation of  $G$  in  $F$ , that is (I 4.7.1), a morphism of  $S$ -groups

$$\rho : G \longrightarrow \underline{\mathrm{Aut}}_{\mathcal{O}_S\text{-mod.}}(F).$$

If  $G/S$  satisfies (E), then 4.1.C and 4.5 give a morphism of  $\mathcal{O}_S$ -modules, denoted  $\rho'$  or  $d\rho$ ,

$$\underline{\mathrm{Lie}}(G/S) \longrightarrow \underline{\mathrm{Lie}}(\underline{\mathrm{Aut}}_{\mathcal{O}_S\text{-mod.}}(F)/S) \simeq \underline{\mathrm{End}}_{\mathcal{O}_S\text{-mod.}}(F).$$

<sup>77</sup> Furthermore, put  $\mathcal{O}_{I_S} = \mathcal{O}_S \oplus t\mathcal{O}_S$ , with  $t^2 = 0$ . It follows from 4.5.1 that, if  $S' \rightarrow S$  and

$$X \in \underline{\mathrm{Lie}}(G/S)(S') \subset G(I_{S'}),$$

then in  $\underline{\mathrm{Aut}}_{\mathcal{O}_{I_{S'}}\text{-mod.}}(F_{I_{S'}})$  one has

$$\rho(X) = \mathrm{id} + t\rho'(X). \quad (*)$$

In other words, for every  $S'' \rightarrow I_{S'}$  and  $f \in F(S'')$ ,

$$\rho(X)(f) = f + t\rho'(X)(f) \quad \text{in } F(S'').$$

**Definition 4.7.2.** Let  $G$  be a good  $S$ -group. Then  $\underline{\mathrm{Lie}}(G/S)$  is a good  $\mathcal{O}_S$ -module, and there is a morphism of  $S$ -groups

$$\mathrm{Ad} : G \longrightarrow \underline{\mathrm{Aut}}_{\mathcal{O}_S\text{-mod.}}(\underline{\mathrm{Lie}}(G/S)).$$

By 4.7.1, it induces a morphism of  $\mathcal{O}_S$ -modules

$$\mathrm{ad} : \underline{\mathrm{Lie}}(G/S) \longrightarrow \underline{\mathrm{End}}_{\mathcal{O}_S\text{-mod.}}(\underline{\mathrm{Lie}}(G/S)),$$

or, equivalently, an  $\mathcal{O}_S$ -bilinear morphism

$$\underline{\mathrm{Lie}}(G/S) \times_S \underline{\mathrm{Lie}}(G/S) \longrightarrow \underline{\mathrm{Lie}}(G/S), \quad (x, y) \longmapsto [x, y] = \mathrm{ad}(x) \cdot y.$$

Here  $x, y$  are arbitrary elements of

$$\underline{\mathrm{Lie}}(G/S)(S') = \underline{\mathrm{Lie}}(G_{S'}/S')(S').$$

If  $G$  is commutative, then  $[x, y] = 0$ .

**4.7.3.**

The bracket admits the following equivalent description. It is enough to consider sections over  $S$ , that is,

$$x, y \in \underline{\mathrm{Lie}}(G/S)(S).$$

There is a canonical isomorphism  $I_S \times_S I_S \simeq I_{I_S}$ . To avoid confusion, let  $I$  and  $I'$  denote two copies of  $I_S$ , and put

$$\mathcal{O}_I = \mathcal{O}_S[t], \quad \mathcal{O}_{I'} = \mathcal{O}_S[t'], \quad t^2 = 0 = (t')^2.$$

<sup>76</sup>Editor's note: the numbering 4.7.1, 4.7.2, and 4.7.3 has been added.

<sup>77</sup>Editor's note: the following sentence has been added.

There is then a commutative square

$$\begin{array}{ccc} I \times I' & \longrightarrow & I' \\ \downarrow & & \downarrow \\ I & \longrightarrow & S \end{array},$$

Figure 49: The two dual-number structures on  $I \times_S I'$ .

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 37; combined-reader p. 95; printed p. 85.*

in which the two arrows from  $I \times I'$  identify it with the scheme of dual numbers over  $I$  or over  $I'$ , respectively. Writing  $L = \underline{\mathrm{Lie}}(G/S)$ , one obtains the following commutative diagram of groups.

$$(1) \quad \begin{array}{ccccccc} & & 1 & & 1 & & \\ & & \downarrow & & \downarrow & & \\ & & L(I) & \longrightarrow & L(S) & \longrightarrow & 1 \\ & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & L(I') & \longrightarrow & G(I \times I') & \longrightarrow & G(I') \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & L(S) & \longrightarrow & G(I) & \longrightarrow & G(S) \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & 1 & & 1 & & 1 \end{array}.$$

Figure 50: The first exact-sequence diagram used to construct the bracket.

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 38; combined-reader p. 96; printed p. 86.*

The ninth object in this diagram is  $\underline{\mathrm{Lie}}(L/S)(S)$ . If  $G$  is good, this is  $L(S)$ , and hence one obtains the next commutative diagram. Its rows and columns are exact sequences of groups; the five groups denoted  $L(-)$  are commutative.<sup>78</sup> Under the identifications

$$L(I) = L(S) \oplus tL(S), \quad L(I') = L(S) \oplus t'L(S),$$

the injections  $L(S) \hookrightarrow L(I)$  and  $L(S) \hookrightarrow L(I')$  are  $u \mapsto tu$  and  $u \mapsto t'u$ , respectively.

<sup>78</sup>Editor's note: the original has been expanded in what follows.

$$(2) \quad \begin{array}{ccccccc} z \in & L(S) & \xrightarrow{t} & L(I) & \longrightarrow & L(S) & \ni x \\ & \downarrow t' & & \downarrow & & \downarrow & \\ & L(I') & \longrightarrow & G(I \times I') & \longrightarrow & G(I') & \\ & \downarrow & & \downarrow & & \downarrow & \\ y \in & L(S) & \longrightarrow & G(I) & \longrightarrow & G(S) & . \end{array}$$

Figure 51: The exact-sequence diagram defining  $[x, y]$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp2-14oct24.pdf, component-local p. 38;  
combined-reader p. 96; printed p. 86.*

In such a diagram, take  $x, y$  as indicated and choose arbitrary lifts  $\tilde{x} \in L(I)$  of  $x$  and  $y' \in L(I')$  of  $y$ . The commutator

$$\tilde{x} y' \tilde{x}^{-1} (y')^{-1} \quad \text{in } G(I \times I')$$

is independent of the chosen lifts and is the image of a uniquely determined element  $z$  as shown. One has  $z = [x, y]$ .<sup>79</sup>

Indeed, use the same letters  $x$  and  $y$  for their images under the canonical sections  $L(S) \rightarrow L(I)$  and  $L(S) \rightarrow L(I')$ . Then

$$\tilde{x} = xu, \quad y' = yv, \quad u, v \in L(S) = L(I) \cap L(I').$$

Since  $L(I)$  and  $L(I')$  are commutative,

$$\begin{aligned} \tilde{x} y' \tilde{x}^{-1} (y')^{-1} &= xuyvu^{-1}x^{-1}v^{-1}y^{-1} \\ &= xuyu^{-1}vx^{-1}v^{-1}y^{-1} \\ &= xyx^{-1}y^{-1}. \end{aligned}$$

This element maps to the unit in both  $G(I)$  and  $G(I')$ , and therefore comes from the unique  $z$  indicated above. Finally, viewing  $y$  as an element of  $L_{I'}(I')$ , and  $x$  as an element of  $L(S) \subset G(I')$ , formula 4.7.1 (\*) gives

$$xyx^{-1} = \text{Ad}(x)(y) = (\text{id} + t' \text{ad}(x))(y) = y + t'[x, y].$$

Thus  $xyx^{-1}y^{-1} \in L(I')$  is the image of  $z = [x, y] \in L(S)$ .

Two properties are now apparent from this construction:

(i) the bracket is functorial in  $G$ . More precisely,

$$G \longmapsto \underline{\text{Lie}}(G/S)$$

is a functor from good  $S$ -groups to good  $\mathcal{O}_S$ -modules equipped with an  $\mathcal{O}_S$ -bilinear composition law;

(ii)  $[x, y] + [y, x] = 0$ , since the diagram is symmetric about its main diagonal.<sup>80</sup>

<sup>79</sup>Editor's note: the original said in a footnote that the author, at Gabriel's request, acknowledged that the exercise was not immediate, which was precisely why it had not been placed in the text. The details have been supplied here, taking account of the addition in 4.7.1; see also [DG70], §II.4, 4.2.

<sup>80</sup>Editor's note: that is,  $x$  and  $y$  play symmetric roles. The image of  $[y, x]$  in  $G(I \times I')$  is the commutator  $y'\tilde{x}(y')^{-1}\tilde{x}^{-1}$ , the inverse of  $\tilde{x}y'\tilde{x}^{-1}(y')^{-1} = [x, y]$ . On the other hand (cf. 4.10 below), it is not known at this point whether  $[x, x] = 0$  necessarily holds: if  $x$  is represented by  $\tilde{x} \in L(I)$  and by  $x' \in L(I')$ , it is not a priori clear that  $\tilde{x}$  and  $x'$  commute.

**Proposition 4.8.** *Let  $F$  be a good  $\mathcal{O}_S$ -module. Under the identification*

$$\underline{\mathrm{Lie}}(\underline{\mathrm{Aut}}_{\mathcal{O}_S\text{-mod.}}(F)/S) = \underline{\mathrm{End}}_{\mathcal{O}_S\text{-mod.}}(F),$$

*one has*

$$\mathrm{Ad}(g) \cdot Y = g \circ Y \circ g^{-1}, \quad [X, Y] = X \circ Y - Y \circ X,$$

*for every  $S' \rightarrow S$ ,  $g \in \underline{\mathrm{Aut}}_{\mathcal{O}_{S'}\text{-mod.}}(F_{S'})$ , and*

$$X, Y \in \underline{\mathrm{Lie}}(\underline{\mathrm{Aut}}_{\mathcal{O}_S\text{-mod.}}(F)/S)(S') = \underline{\mathrm{End}}_{\mathcal{O}_{S'}\text{-mod.}}(F_{S'}).$$

*Proof.*<sup>81</sup> After base change, we may assume  $S' = S$ . Put  $I = I_S$  and  $\mathcal{O}_I = \mathcal{O}_S[t]$ , with  $t^2 = 0$ . Recall from 4.5.1 that the inclusion

$$i : \underline{\mathrm{End}}_{\mathcal{O}_S\text{-mod.}}(F) \hookrightarrow \underline{\mathrm{Aut}}_{\mathcal{O}_I\text{-mod.}}(F_I)$$

sends  $Y$  to  $\mathrm{id} + tY$ . By the definition of  $\mathrm{Ad}(g)$  in 4.1.A,

$$\mathrm{id} + t \mathrm{Ad}(g)(Y) = g \circ (\mathrm{id} + tY) \circ g^{-1} = \mathrm{id} + t(g \circ Y \circ g^{-1}),$$

and therefore  $\mathrm{Ad}(g)(Y) = g \circ Y \circ g^{-1}$ .

Let  $I'$  be a second copy of  $I_S$ , and put  $\mathcal{O}_{I'} = \mathcal{O}_S[t']$ , with  $(t')^2 = 0$ . Apply 4.7.3 to

$$G = \underline{\mathrm{Aut}}_{\mathcal{O}_S\text{-mod.}}(F), \quad L = \underline{\mathrm{Lie}}(G/S) = \underline{\mathrm{End}}_{\mathcal{O}_S\text{-mod.}}(F).$$

Here  $L = \underline{\mathrm{End}}(F)$  is the differential identification of Proposition 4.5. The re-edition prints  $\underline{\mathrm{Aut}}(F)$  at this point, but  $\underline{\mathrm{End}}(F)$  is the type-correct object used by the ensuing endomorphism calculation. Identify  $X$  with its image under the canonical section  $L(S) \hookrightarrow L(I)$ . Its image in  $G(I \times I')$  is  $\mathrm{id} + t'X$ , whose inverse is  $\mathrm{id} - t'X$ . Similarly,  $Y$  lifts to  $\mathrm{id} + tY$ , with inverse  $\mathrm{id} - tY$ . Their commutator is

$$\begin{aligned} & (\mathrm{id} + t'X) \circ (\mathrm{id} + tY) \circ (\mathrm{id} - t'X) \circ (\mathrm{id} - tY) \\ &= \mathrm{id} + tt'(X \circ Y - Y \circ X). \end{aligned}$$

This is the image in  $G(I \times I')$  of  $Z = X \circ Y - Y \circ X \in L(S)$ : first  $Z$  maps to  $tZ \in L(I)$ , and then to  $\mathrm{id} + t'tZ \in G(I \times I')$ . By 4.7.3,  $[X, Y] = X \circ Y - Y \circ X$ .

**Corollary 4.8.1.** *Let  $G$  be a good  $S$ -group and let  $x, y, z \in \underline{\mathrm{Lie}}(G/S)(S')$ . Then*

$$[x, [y, z]] + [y, [z, x]] + [z, [x, y]] = 0.$$

<sup>82</sup> Indeed, since  $G$  is good,  $\underline{\mathrm{Lie}}(G/S)$  is a good  $\mathcal{O}_S$ -module; hence Theorem 4.7 shows that

$$\underline{\mathrm{Aut}}_{\mathcal{O}_S\text{-mod.}}(\underline{\mathrm{Lie}}(G/S))$$

is a good  $S$ -group. The morphism

$$\mathrm{Ad} : G \longrightarrow \underline{\mathrm{Aut}}_{\mathcal{O}_S\text{-mod.}}(\underline{\mathrm{Lie}}(G/S))$$

and the functoriality in 4.7.3(i) yield

$$\mathrm{ad}[x, y] = [\mathrm{ad} x, \mathrm{ad} y].$$

<sup>81</sup>Editor's note: the original has been expanded and simplified in what follows.

<sup>82</sup>Editor's note: the original has been expanded in what follows.

Together with Proposition 4.8, this gives

$$\mathrm{ad}[x, y] = \mathrm{ad} x \circ \mathrm{ad} y - \mathrm{ad} y \circ \mathrm{ad} x.$$

Applying both sides to  $z$  gives the Jacobi identity.

**Corollary 4.8.2.** *Let  $G$  be a good  $S$ -group acting linearly on a good  $\mathcal{O}_S$ -module  $F$ ; equivalently, let  $F$  be a  $G$ - $\mathcal{O}_S$ -module, with  $G$  and  $F$  good. Then the linear map*

$$\rho' : \underline{\mathrm{Lie}}(G/S) \longrightarrow \underline{\mathrm{End}}_{\mathcal{O}_S\text{-mod.}}(F)$$

*is a representation; that is,*

$$\rho'([x, y]) = \rho'(x) \circ \rho'(y) - \rho'(y) \circ \rho'(x).$$

**Scholium 4.9.** To every good  $S$ -group  $G$  (for example, every representable one) we have functorially associated a good  $\mathcal{O}_S$ -module  $\underline{\mathrm{Lie}}(G/S)$  equipped with a bilinear map satisfying

$$[x, y] + [y, x] = 0, \quad [x, [y, z]] + [y, [z, x]] + [z, [x, y]] = 0.$$

We shall call  $\underline{\mathrm{Lie}}(G/S)$ , with this structure, the *Lie algebra* of  $G$  over  $S$ . The qualification is that it is not yet known whether it is an  $\mathcal{O}_S$ -Lie algebra in the strict sense.<sup>83</sup> Every linear representation of  $G$  in a good  $\mathcal{O}_S$ -module  $F$  induces a representation of its Lie algebra. In particular, the adjoint representation of  $G$  induces the adjoint representation of its Lie algebra.

**Definition 4.10.** An  $S$ -functor in groups  $G$  is called *very good* if it is good and  $\underline{\mathrm{Lie}}(G/S)$  is an  $\mathcal{O}_S$ -Lie algebra, that is, if  $[x, x] = 0$  identically.

**Examples 4.10.1.** The following  $S$ -groups are very good:  $\underline{\mathrm{Aut}}_{\mathcal{O}_S\text{-mod.}}(F)$  for every good  $\mathcal{O}_S$ -module  $F$  (by 4.7 and 4.8); every representable group (as shown below); every good  $S$ -group admitting a monomorphism into a very good  $S$ -group, in particular every good subgroup functor of a representable group; and every good  $S$ -group admitting a faithful linear representation in a good  $\mathcal{O}_S$ -module, for example every good  $S$ -group for which  $\mathrm{Ad}$  is a monomorphism.

#### 4.11.

Suppose now that  $G$  is a group scheme over  $S$ . By 4.1.4,  $\underline{\mathrm{Lie}}(G/S)(S)$  identifies with the group of infinitesimal automorphisms of  $G/S$  invariant under right translation; by 3.14, this is the group of derivations of  $\mathcal{O}_G$  over  $\mathcal{O}_S$  invariant under right translation. This identification respects the module structure and is an anti-isomorphism of Lie algebras.<sup>84</sup>

To see this, argue as in 4.7.3. Put

$$\mathcal{O}_I = \mathcal{O}_S[t], \quad \mathcal{O}_{I'} = \mathcal{O}_S[t'], \quad t^2 = 0 = (t')^2,$$

and let  $x \in L(I)$ ,  $y \in L(I')$ , where  $L = \underline{\mathrm{Lie}}(G/S)$ . Left translation  $\lambda_x$ , respectively  $\lambda_y$ , is an  $S$ -automorphism of  $G_{I \times I'}$  inducing the identity on  $G_{I'}$ , respectively on  $G_I$ . It corresponds to the  $\mathcal{O}_S$ -automorphism

$$u = \mathrm{id} + t d_x, \quad \text{respectively} \quad v = \mathrm{id} + t' d_y$$

<sup>83</sup>Editor's note: the unresolved point is whether  $[x, x] = 0$ ; see 4.10.

<sup>84</sup>Editor's note: a sign error in the original has been corrected here; cf. [DG70], §II.4, 4.4 and 4.6.

of

$$\mathcal{O}_{G_{I \times I'}} = \mathcal{O}_G[t, t']/(t^2, (t')^2),$$

where  $d_x, d_y$  are  $\mathcal{O}_S$ -derivations of  $\mathcal{O}_G$  invariant under right translation. The correspondence between  $S$ -automorphisms of  $G_{I \times I'}$  and  $\mathcal{O}_S$ -automorphisms of  $\mathcal{O}_{G_{I \times I'}}$  is contravariant. Therefore

$$\lambda_x \lambda_y \lambda_x^{-1} \lambda_y^{-1}$$

corresponds to

$$v^{-1}u^{-1}vu = \text{id} + tt'(d_y d_x - d_x d_y).$$

It follows from 4.7.3 that

$$x \mapsto -d_x$$

is an isomorphism of Lie algebras; for further details, see [DG70], §II.4, 4.4 and 4.6. The same argument applies to

$$\underline{\text{Lie}}(G/S)(S') = \underline{\text{Lie}}(G_{S'}/S')(S')$$

for every  $S' \rightarrow S$ . Thus one recovers the classical definition.<sup>85</sup>

**Scholium 4.11.1.** *Via the isomorphism  $x \mapsto -d_x$ ,  $\underline{\text{Lie}}(G/S)$  identifies with the functor that assigns to each  $S' \rightarrow S$  the  $\mathcal{O}(S')$ -Lie algebra of derivations of  $G_{S'}$  relative to  $S'$  invariant under right translation.*

Since every representable group is already known to be good by 4.6.1, the preceding discussion gives:

**Corollary 4.11.2.** *Every representable group is very good.*

Let  $\varepsilon : S \rightarrow G$  be the unit section of  $G$ , and put

$$\omega_{G/S}^1 = \varepsilon^*(\Omega_{G/S}^1).$$

Recall from 3.3 that  $\underline{\text{Lie}}(G/S)$  is represented by the vector fibration  $V(\omega_{G/S}^1)$ .

**Scholium 4.11.3.** *Thus to every  $S$ -group scheme  $G$  we have functorially associated a vector fibration*

$$\text{Lie}(G/S) = V(\omega_{G/S}^1)$$

*over  $S$ , representing the functor  $\underline{\text{Lie}}(G/S)$ , and hence endowed with the structure of an  $S$ -scheme in  $\mathcal{O}_S$ -Lie algebras. Moreover, this construction commutes with base extension and finite products (cf. 3.4 and 3.8).*

**Remarks 4.11.4.** <sup>86</sup> Let  $\pi : G \rightarrow S$  be the structural morphism.

(a) The  $\mathcal{O}_G$ -module  $\Omega_{G/S}^1$  is naturally  $(G \times_S G)$ -equivariant (cf. I, §6). Hence, by I 6.8.1,

$$\Omega_{G/S}^1 \simeq \pi^*(\omega_{G/S}^1).$$

For example,  $\Omega_{G/S}^1$  is locally free, respectively locally free of finite type, whenever  $\omega_{G/S}^1$  is. This holds in particular when  $S$  is the spectrum of a field, respectively when  $S$  is the spectrum of a field and  $G$  is locally of finite type over  $S$ .

(b) By I 6.8.2,  $\omega_{G/S}^1$  has a canonical  $G$ - $\mathcal{O}_S$ -module structure, which induces the adjoint action on

$$V(\omega_{G/S}^1) = \text{Lie}(G/S).$$

<sup>85</sup>Editor's note: the original has been expanded in what follows. In particular, the numbering 4.11.1–4.11.8 has been added for later reference.

<sup>86</sup>Editor's note: what follows has been expanded, taking account of the additions made in Exposé I, §6.8.

87

(c) By EGA I, 5.3.11 and 5.4.6,  $\varepsilon : S \rightarrow G$  is an immersion; it is a closed immersion if  $G$  is separated over  $S$ . Thus  $\omega_{G/S}^1$  identifies with  $\mathcal{I}/\mathcal{I}^2$ , where  $\mathcal{I}$  is the quasi-coherent ideal defining  $\varepsilon(S)$  in an open subscheme  $U \subset G$  in which  $\varepsilon(S)$  is closed. If  $G \rightarrow S$  is separated, one may take  $U = G$ . If  $G = \operatorname{Spec} A(G)$  is affine over  $S$ , then  $\mathcal{I}$  is the augmentation ideal of  $A(G)$ , namely the kernel of

$$\varepsilon^\sharp : A(G) \longrightarrow \mathcal{O}_S$$

(cf. I 4.2).<sup>88</sup>

**Remark 4.11.5.** The isomorphism  $\Omega_{G/S}^1 \simeq \pi^*(\omega_{G/S}^1)$  identifies the  $\mathcal{O}_S$ -module  $\omega_{G/S}^1$  with the sheaf

$$\pi_*(\Omega_{G/S}^1)^G$$

of differentials of  $G$  relative to  $S$  invariant under right translation: its sections over an open  $U \subset S$  are the sections of  $\Omega_{G/S}^1$  over  $\pi^{-1}(U)$  invariant under right translation (cf. I 6.8.3; compare also VII<sub>A</sub>, 2.4).<sup>89</sup>

**Notation 4.11.6.** Write  $\mathcal{L}ie(G/S)$  for the sheaf of sections of the vector fibration  $\operatorname{Lie}(G/S) \rightarrow S$ . It is the  $\mathcal{O}_S$ -module

$$\mathcal{L}ie(G/S) = (\omega_{G/S}^1)^\vee = \mathcal{H}om_{\mathcal{O}_S}(\omega_{G/S}^1, \mathcal{O}_S),$$

dual to  $\omega_{G/S}^1$  (cf. EGA II, 1.7.9), and it carries a structure of  $\mathcal{O}_S$ -Lie algebra.

In general this construction does not commute with base extension. Consequently, the Lie-algebra structure on this module does not determine the structure of an  $S$ -scheme in  $\mathcal{O}_S$ -Lie algebras on  $\operatorname{Lie}(G/S)$ .<sup>90</sup>

Nevertheless:

**Lemma 4.11.7.** *Suppose that  $\omega_{G/S}^1$  is locally free of finite type. This is the case, in particular, if  $G$  is smooth over  $S$  (cf. EGA IV<sub>4</sub>, 17.2.4), or if  $S$  is the spectrum of a field and  $G$  is locally of finite type over  $S$ . Then*

$$\mathcal{L}ie(G/S)^\vee \simeq (\omega_{G/S}^1)^{\vee\vee} \simeq \omega_{G/S}^1,$$

and therefore

$$\operatorname{Lie}(G/S) = V(\omega_{G/S}^1) = V(\mathcal{L}ie(G/S)^\vee) = W(\mathcal{L}ie(G/S)),$$

where the final equality follows from I 4.6.5.1.

Finally, let  $G \rightarrow H$  be a monomorphism of functors in groups. Then

$$\operatorname{Lie}(G/S) \longrightarrow \operatorname{Lie}(H/S)$$

<sup>87</sup>Editor's note: for this reason, the linear action of  $G$  on  $\omega_{G/S}^1$  could be called the pre-adjoint action; by an abuse of terminology it will still be called the adjoint action. A slightly more general construction, used in Exposé III (especially III 0.8), is as follows. Suppose that to every  $Y \rightarrow S$  one assigns an  $\mathcal{O}_Y$ -module  $\mathcal{N}(Y)$ , and to every  $S$ -morphism  $\phi : Z \rightarrow Y$  a functorial morphism of  $\mathcal{O}_Z$ -modules  $\mathcal{N}(\phi) : \phi^*\mathcal{N}(Y) \rightarrow \mathcal{N}(Z)$ . For example, if  $\mathcal{J}$  is a quasi-coherent ideal of  $\mathcal{O}_S$ , one may take  $\mathcal{N}(Y) = \mathcal{J}\mathcal{O}_Y$ . The action of  $G$  on  $\omega_{G/S}^1$  then induces a generalized adjoint action of  $G$  on the  $S$ -functor in abelian groups that assigns to  $f : Y \rightarrow S$  the group  $\operatorname{Hom}_{\mathcal{O}_Y}(f^*\omega_{G/S}^1, \mathcal{N}(Y))$ .

<sup>88</sup>Editor's note: the preceding description of  $\omega_{G/S}^1$ , which is useful later (for example in VII<sub>A</sub>, 5.5), has been added.

<sup>89</sup>Editor's note: this description of  $\omega_{G/S}^1$  in terms of invariant differentials is not used below.

<sup>90</sup>Editor's note: the reason is that  $\mathcal{L}ie(G/S) \simeq (\omega_{G/S}^1)^\vee$  need not determine  $\omega_{G/S}^1$ . For example, let  $S = \mathbb{A}_k^1$ , where  $k$  is a field, and let  $G$  be the  $S$ -group whose fiber at  $s = 0$  is  $\mathbb{G}_{a,k}$ , while all other fibers are the unit group. Then  $\mathcal{L}ie(G/S) = 0$ , but  $\operatorname{Lie}(G_s/k)(k) = k$ .

is also a monomorphism (cf. 3.7). Every vector monomorphism of vector fibrations is a closed immersion.<sup>91</sup> It follows that:

**Corollary 4.11.8.** *Let  $G \hookrightarrow H$  be a monomorphism of  $S$ -group schemes.*

(i) *The morphism*

$$\mathrm{Lie}(G/S) \longrightarrow \mathrm{Lie}(H/S)$$

*is a closed immersion; consequently*

$$\omega_{H/S}^1 \longrightarrow \omega_{G/S}^1$$

*is an epimorphism.*

(ii) *If  $\omega_{G/S}^1$  is locally free of finite type, then the corresponding morphism*

$$\mathcal{L}ie(G/S) \longrightarrow \mathcal{L}ie(H/S)$$

*identifies  $\mathcal{L}ie(G/S)$  with an  $\mathcal{O}_S$ -submodule of  $\mathcal{L}ie(H/S)$  that is locally a direct summand.*

## 5. Computation of some Lie algebras

### 5.1. Examples of Lie algebras: diagonalizable groups

Let  $G = D_S(M)$  be a diagonalizable group over  $S$  (I 4.4). Since the formation of  $\underline{\mathrm{Lie}}(G/S)$  commutes with base extension, it is enough to carry out the construction for  $G = D(M)$ . We then have

$$G(I_S) = \mathrm{Hom}_{\mathrm{gr}}(M, \Gamma(I_S, \mathcal{O}_{I_S})^\times) = \mathrm{Hom}_{\mathrm{gr}}(M, \Gamma(S, \mathcal{D}_{\mathcal{O}_S})^\times).$$

There is a split exact sequence

$$1 \longrightarrow \Gamma(S, \mathcal{O}_S) \longrightarrow \Gamma(S, \mathcal{D}_{\mathcal{O}_S})^\times \longrightarrow \Gamma(S, \mathcal{O}_S)^\times \longrightarrow 1,$$

which shows that  $\underline{\mathrm{Lie}}(G)(S)$  identifies with  $\mathrm{Hom}_{\mathrm{gr}}(M, \mathcal{O}(S))$ , endowed with its evident  $\mathcal{O}(S)$ -module structure. After base change we therefore obtain:

**Proposition 5.1.** *There are isomorphisms*

$$\underline{\mathrm{Hom}}_{S\text{-gr}}(M_S, \mathbf{O}_S) \xrightarrow{\sim} \underline{\mathrm{Lie}}(D_S(M)/S)$$

and

$$\mathcal{H}om_{\mathrm{gr}}(\widetilde{M}_S, \mathcal{O}_S) \xrightarrow{\sim} \mathcal{L}ie(D_S(M)/S),$$

where, in the second isomorphism,  $\widetilde{M}_S$  denotes the constant sheaf of groups on  $S$  defined by  $M$ , and  $\mathcal{H}om_{\mathrm{gr}}$  denotes the sheaf of homomorphisms of sheaves of groups.

**Corollary 5.1.1.** *If  $M$  is free of finite type—or, as we shall say later, if  $D_S(M)$  is a split torus—then, with  $W$  as in I 4.6.5,*

$$W(\mathcal{L}ie(D_S(M)/S)) \xrightarrow{\sim} \underline{\mathrm{Lie}}(D_S(M)/S), \quad M^\vee \otimes_{\mathbb{Z}} \mathcal{O}_S \xrightarrow{\sim} \mathcal{L}ie(D_S(M)/S).$$

Here  $M^\vee$  denotes the dual of the abelian group  $M$ . In particular,

$$\mathbf{O}_S \xrightarrow{\sim} \underline{\mathrm{Lie}}(\mathbb{G}_{m,S}/S), \quad \mathcal{O}_S \xrightarrow{\sim} \mathcal{L}ie(\mathbb{G}_{m,S}/S).$$

<sup>91</sup>Editor's note: let  $f : M \rightarrow N$  be a morphism of  $\mathcal{O}_S$ -modules and let  $P = \mathrm{Coker}(f)$ . If  $V(N) \rightarrow V(M)$  is a monomorphism, the surjective morphism  $\mathrm{Sym}(N) \rightarrow \mathrm{Sym}(P)$  factors through  $\mathcal{O}_S$ ; hence  $P = 0$ .



## 5.2. Normalizers and centralizers

We first prove a few lemmas. Recall (I 3.1.1) that a sequence

$$0 \longrightarrow F' \longrightarrow F \longrightarrow F'' \longrightarrow 0$$

of  $\mathbf{O}_S$ -modules is called *exact* if, for every  $S' \rightarrow S$ , the sequence

$$0 \longrightarrow F'(S') \longrightarrow F(S') \longrightarrow F''(S') \longrightarrow 0$$

of  $\mathbf{O}(S')$ -modules is exact. Likewise, a sequence  $1 \rightarrow G' \rightarrow G \rightarrow G'' \rightarrow 1$  of  $S$ -groups is called *exact* if the corresponding sequence of groups on  $S'$ -points is exact for every  $S' \rightarrow S$ .

**Lemma 5.2.1.** *Let  $1 \rightarrow G' \rightarrow G \rightarrow G'' \rightarrow 1$  be an exact sequence of  $S$ -groups.* <sup>92</sup>

(i) *The sequences*

$$1 \longrightarrow T_{G'/S}(\mathcal{M}) \longrightarrow T_{G/S}(\mathcal{M}) \longrightarrow T_{G''/S}(\mathcal{M}) \longrightarrow 1$$

*and*

$$1 \longrightarrow \underline{\mathrm{Lie}}(G'/S, \mathcal{M}) \longrightarrow \underline{\mathrm{Lie}}(G/S, \mathcal{M}) \longrightarrow \underline{\mathrm{Lie}}(G''/S, \mathcal{M}) \longrightarrow 1$$

*are exact.*

(ii) *Let  $1 \rightarrow H' \rightarrow H \rightarrow H'' \rightarrow 1$  be a second sequence of groups. It is exact if and only if*

$$1 \longrightarrow G' \times_S H' \longrightarrow G \times_S H \longrightarrow G'' \times_S H'' \longrightarrow 1$$

*is exact.*

(iii) *If two of  $G', G, G''$  satisfy condition (E), then so does the third.*

(iv) *If  $0 \rightarrow F' \rightarrow F \rightarrow F'' \rightarrow 0$  is exact and two of the  $\mathbf{O}_S$ -modules  $F', F, F''$  are good, then so is the third.*

(v) *If two of the  $S$ -groups  $G', G, G''$  are good, then so is the third.*

The first part of (i) is immediate, and the second follows from it. Likewise, (ii) is immediate. We prove (iii). For brevity, write  $L(\mathcal{M}) = \underline{\mathrm{Lie}}(G, \mathcal{M})$ ,  $L'(\mathcal{M}) = \underline{\mathrm{Lie}}(G', \mathcal{M})$ , and similarly for  $L''$ . We have the following commutative diagram. <sup>93</sup>

$$\begin{array}{ccccccc} 1 & \longrightarrow & L'(\mathcal{M} \oplus \mathcal{N}) & \longrightarrow & L(\mathcal{M} \oplus \mathcal{N}) & \longrightarrow & L''(\mathcal{M} \oplus \mathcal{N}) \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & L'(\mathcal{M}) \times L'(\mathcal{N}) & \longrightarrow & L(\mathcal{M}) \times L(\mathcal{N}) & \longrightarrow & L''(\mathcal{M}) \times L''(\mathcal{N}) \longrightarrow 1 \end{array}$$

Figure 52: The exact-sequence diagram used to prove Lemma 5.2.1(iii).

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 44;  
combined-reader p. 102.*

The upper row is exact by (i), and the lower row is exact by (i) and (ii). Assertion (iii) follows.

<sup>92</sup>Editor's note: The statement and proof of this lemma have been expanded; this will be useful in 5.3.1.

<sup>93</sup>Editor's note: The proofs of (iii) and (v) have been added.

We prove (iv). There is a commutative diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & F' \otimes_{\mathbf{O}_S} T_{\mathbf{O}_S/S}(\mathcal{M}) & \longrightarrow & F \otimes_{\mathbf{O}_S} T_{\mathbf{O}_S/S}(\mathcal{M}) & \longrightarrow & F'' \otimes_{\mathbf{O}_S} T_{\mathbf{O}_S/S}(\mathcal{M}) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & T_{F'/S}(\mathcal{M}) & \longrightarrow & T_{F/S}(\mathcal{M}) & \longrightarrow & T_{F''/S}(\mathcal{M}) \longrightarrow 0
 \end{array}$$

Figure 53: The exact-sequence diagram used to prove Lemma 5.2.1(iv).

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 44;  
combined-reader p. 102.*

with exact rows: the first because  $T_{\mathbf{O}_S/S}(\mathcal{M})$  is a free, hence flat,  $\mathbf{O}_S$ -module, and the second by (i). Thus, if two of  $F', F, F''$  are good, so is the third. Finally, (v) follows from (iii) and (iv).

**Lemma 5.2.2.** *Let  $G$  be an  $S$ -group, let  $E, H$  be two  $G$ -objects, and let  $F$  be a  $G$ - $\mathbf{O}_S$ -module.*

- (i) *The canonical homomorphism  $E^G \times_S H^G \rightarrow (E \times_S H)^G$  is an isomorphism.*
- (ii) *If  $F$  is a good  $\mathbf{O}_S$ -module, then so is  $F^G$ .*

The first assertion is immediate; we prove the second. For every  $S' \rightarrow S$ , there is a commutative diagram

$$\begin{array}{ccc}
 F^G(I_{S'}(\mathcal{M})) & \xhookrightarrow{\quad} & F(I_{S'}(\mathcal{M})) \\
 \uparrow \phi & & \uparrow \simeq \phi_1 \\
 F^G(S') \otimes_{\mathbf{O}(S')} \mathbf{O}(I_{S'}(\mathcal{M})) & \xhookrightarrow{\quad} & F(S') \otimes_{\mathbf{O}(S')} \mathbf{O}(I_{S'}(\mathcal{M}))
 \end{array}$$

Figure 54: The invariants-and-base-change square in the proof of Lemma 5.2.2.

*Temporary Loop-1 image; source locator: Polo–Gille Exp2-14oct24.pdf, component-local p. 45;  
combined-reader p. 103.*

and we must prove that  $\phi$  is bijective. It is plainly injective. To prove surjectivity, let  $(t_0, \dots, t_n)$  be a basis of the free  $\mathbf{O}(S')$ -module  $\mathbf{O}(I_{S'}(\mathcal{M}))$ , and let

$$u = \sum_{i=0}^n f_i \otimes t_i$$

be an element of  $F(S') \otimes_{\mathbf{O}(S')} \mathbf{O}(I_{S'}(\mathcal{M}))$  such that  $\phi_1(u)$  belongs to  $F^G(I_{S'}(\mathcal{M}))$ . We show that each  $f_i$  belongs to  $F^G(S')$ .

Let  $S'' \rightarrow S'$ . Via the zero section  $\varepsilon_{\mathcal{M}}$ , we may regard  $S''$  as lying over  $I_{S'}(\mathcal{M})$ . Then, for every  $g \in G(S'')$ ,

$$u_{S''} = g \cdot u_{S''} = \sum_{i=0}^n g \cdot (f_i)_{S''} \otimes (t_i)_{S''}.$$

Since the  $(t_i)_{S''}$  form a basis of

$$\mathbf{O}(I_{S''}(\mathcal{M})) \simeq \mathbf{O}(I_{S'}(\mathcal{M})) \otimes_{\mathbf{O}(S')} \mathbf{O}(S'')$$

over  $\mathbf{O}(S'')$ , it follows that  $g \cdot (f_i)_{S''} = (f_i)_{S''}$ , hence  $f_i \in F^G(S')$  for all  $i$ .

**Notation.** If  $E$  is an  $S$ -group and  $F$  a sub- $S$ -group, write  $E/F$  for the  $S$ -functor sending  $S' \rightarrow S$  to the set  $E(S')/F(S')$  of classes  $\bar{x} = xF(S')$ . If  $E$  is commutative, then  $E/F$  is a commutative  $S$ -group.<sup>94</sup>

Now let  $G$  be an  $S$ -group and  $K$  a sub- $S$ -group, and set  $E = \underline{\mathrm{Lie}}(G/S, \mathcal{M})$  and  $F = \underline{\mathrm{Lie}}(K/S, \mathcal{M})$ . The adjoint action of  $K$  on  $E$  stabilizes  $F$ , and therefore induces an action on  $E/F$ . For every  $S' \rightarrow S$ ,

$$(E/F)^K(S') = \{\bar{x} \in E(S')/F(S') \mid \text{for every } f : S'' \rightarrow S' \text{ and } k \in K(S''), \\ f^*(\bar{x}^{-1}) \mathrm{Ad}(k)(f^*(\bar{x})) \in F(S'')\},$$

where  $f^*(\bar{x})$  denotes the image of  $\bar{x}$  in  $E(S'')$ .

**Theorem 5.2.3.** *Let  $G$  be an  $S$ -group and  $K$  a sub- $S$ -group. Put (I 2.3.3)*

$$N = \underline{\mathrm{Norm}}_G(K), \quad Z = \underline{\mathrm{Centr}}_G(K),$$

and let  $K$  act on  $\underline{\mathrm{Lie}}(G/S, \mathcal{M})$  through the adjoint representation of  $G$ .

(i) *If the group law of  $\underline{\mathrm{Lie}}(G/S, \mathcal{M})$  is commutative,<sup>9596</sup> then*

$$\frac{\underline{\mathrm{Lie}}(N/S, \mathcal{M})}{\underline{\mathrm{Lie}}(K/S, \mathcal{M})} = \left( \frac{\underline{\mathrm{Lie}}(G/S, \mathcal{M})}{\underline{\mathrm{Lie}}(K/S, \mathcal{M})} \right)^K.$$

(ii) *If the same group law is commutative, then*

$$\underline{\mathrm{Lie}}(Z/S, \mathcal{M}) = \underline{\mathrm{Lie}}(G/S, \mathcal{M})^K.$$

(iii) *If  $G$  satisfies (E), then  $Z$  satisfies (E); if  $G$  and  $K$  satisfy (E), then  $N$  satisfies (E).*

(iv) *If  $G$  is good, then  $Z$  is good; if, moreover,  $G$  is very good, then  $Z$  is very good.*

(v) *If  $G$  and  $K$  are good, then  $N$  is good; if, moreover,  $G$  is very good, then  $N$  is very good.*

To prove (i) and (ii),<sup>97</sup> we use the following lemma, which follows from the diagram of exact sequences considered in 4.1.B, with  $G$  and  $H$  interchanged.

**Lemma 5.2.3.0.** *Let  $G$  be an  $S$ -group,  $H$  a sub- $S$ -group, and  $\mathcal{M}$  a free  $\mathcal{O}_S$ -module of finite type. Then  $T_{H/S}(\mathcal{M})$  and  $\underline{\mathrm{Lie}}(G/S, \mathcal{M})$  are sub- $S$ -groups of  $T_{G/S}(\mathcal{M})$ , and*

$$T_{H/S}(\mathcal{M}) \cap \underline{\mathrm{Lie}}(G/S, \mathcal{M}) = \underline{\mathrm{Lie}}(H/S, \mathcal{M}),$$

where

$$T_{H/S}(\mathcal{M}) \cap \underline{\mathrm{Lie}}(G/S, \mathcal{M}) := T_{H/S}(\mathcal{M}) \times_{T_{G/S}(\mathcal{M})} \underline{\mathrm{Lie}}(G/S, \mathcal{M}).$$

<sup>94</sup>Editor's note: These notations have been expanded.

<sup>95</sup>This condition is automatic when  $G$  satisfies (E), for example when  $G$  is representable.

<sup>96</sup>Editor's note: For an example in which the  $S$ -group  $\underline{\mathrm{Lie}}(G/S)$  is noncommutative, see 6.3.

<sup>97</sup>Editor's note: The proof has been expanded, and in particular Lemma 5.2.3.0, implicit in the original, has been added.

Since the functors in (i) and (ii) commute with base extension, it is enough to prove the equalities on  $S$ -points. Put  $H = N$  (respectively,  $H = Z$ ) and let  $\alpha = \pm 1$ . By the preceding lemma and the definitions of  $N$  and  $Z$  (I 2.3.3),

$$\begin{aligned} \underline{\text{Lie}}(H/S, \mathcal{M})(S) = \{ X \in \underline{\text{Lie}}(G/S, \mathcal{M})(S) \subset G(I_S(\mathcal{M})) \mid \\ \text{for every } f : S' \rightarrow I_S(\mathcal{M}) \text{ and } u \in K(S'), \\ f^*(X^\alpha) u f^*(X^{-\alpha}) u^{-1} \in K(S'), \\ \text{respectively } f^*(X) u f^*(X^{-1}) u^{-1} = 1 \}. \end{aligned} \quad (*)$$

Here  $f^*(X)$  denotes the image of  $X$  in  $G(S')$ .

For brevity, write

$$\mathfrak{g} = \underline{\text{Lie}}(G/S, \mathcal{M}), \quad \mathfrak{k} = \underline{\text{Lie}}(K/S, \mathcal{M}), \quad \mathfrak{n} = \underline{\text{Lie}}(N/S, \mathcal{M}), \quad \mathfrak{z} = \underline{\text{Lie}}(Z/S, \mathcal{M}).$$

If  $X \in \underline{\text{Lie}}(H/S, \mathcal{M})(S)$ , the equalities (\*) hold for every  $f : S' \rightarrow S$ , because  $f$  factors through  $I_S(\mathcal{M})$  as

$$f = \rho \circ \varepsilon_{\mathcal{M}} \circ f,$$

where  $\rho : I_S(\mathcal{M}) \rightarrow S$  is the structural morphism and  $\varepsilon_{\mathcal{M}} : S \rightarrow I_S(\mathcal{M})$  is the zero section. Consequently,

$$\mathfrak{n}(S)/\mathfrak{k}(S) \subset (\mathfrak{g}/\mathfrak{k})^K(S), \quad \mathfrak{z}(S) \subset \mathfrak{g}^K(S).$$

For the reverse inclusions, assume henceforth that  $\mathfrak{g}$  is commutative. Then  $\mathfrak{g}^K$  and  $(\mathfrak{g}/\mathfrak{k})^K$  are commutative  $S$ -groups. Let  $X \in \mathfrak{g}(S)$ , and write  $\bar{X}$  for its image in  $(\mathfrak{g}/\mathfrak{k})(S)$ . Suppose that  $\bar{X} \in (\mathfrak{g}/\mathfrak{k})^K(S)$  (respectively,  $X \in \mathfrak{g}^K(S)$ ); we show that  $X \in \mathfrak{n}(S)$  (respectively,  $X \in \mathfrak{z}(S)$ ).

Let  $f : S' \rightarrow I_S(\mathcal{M})$ . Consider the cartesian square

$$\begin{array}{ccc} I_{S'}(\mathcal{M}) & \xrightarrow{p} & I_S(\mathcal{M}) \\ \rho' \downarrow & & \downarrow \rho \\ S' & \xrightarrow{\rho \circ f} & S \end{array}$$

Figure 55: The cartesian square used in the reverse-inclusion argument.

*Temporary Loop-1 image; source locator: Polo-Gille Exp2-14oct24.pdf, component-local p. 47; combined-reader p. 105.*

and let  $h$  be the section of  $\rho'$  defined by  $f$ . It is enough to show that, for every  $v \in K(I_{S'}(\mathcal{M}))$ ,

$$\left. \begin{aligned} & p^*(X^\alpha) v p^*(X^{-\alpha}) v^{-1} \in K(I_{S'}(\mathcal{M})), \\ \text{respectively } & p^*(X) v p^*(X^{-1}) v^{-1} = 1. \end{aligned} \right\} \quad (**)$$

Indeed, take  $v = \rho'^*(u)$  and apply  $h^*$  to (\*\*). This gives (\*), because  $\rho' \circ h = \text{id}_{S'}$  and  $p \circ h = f$ .

We now prove (\*\*), writing  $X$  for  $p^*(X)$ . Every  $v \in K(I_{S'}(\mathcal{M}))$  is uniquely of the form  $Yk$ , with  $Y \in \underline{\text{Lie}}(K/S, \mathcal{M})(S')$  and  $k \in K(S')$ . The expression  $X^\alpha v X^{-\alpha} v^{-1}$  becomes

$$X^\alpha Y (k X^{-\alpha} k^{-1}) Y^{-1} = X^\alpha \text{Ad}(k)(X^{-\alpha}),$$

because  $\underline{\text{Lie}}(G/S, \mathcal{M})$  is commutative. This element lies in  $\underline{\text{Lie}}(G/S, \mathcal{M})(S')$ . By Lemma 5.2.3.0, condition (\*\*) is therefore equivalent, for every  $k \in K(S')$ , to

$$\begin{cases} \text{Ad}(k)(X) = X, & H = Z, \\ X^\alpha \text{Ad}(k)(X^{-\alpha}) \in \underline{\text{Lie}}(K/S, \mathcal{M})(S'), & H = N. \end{cases}$$

For  $H = Z$ , this follows from  $X \in \mathfrak{g}^K(S)$ . For  $H = N$ , the condition may equally be written

$$\text{Ad}(k)(\bar{X}) = \bar{X}, \quad \text{Ad}(k)(\bar{X}^{-1}) = \bar{X}^{-1}. \quad (*')$$

The second equality follows from the first because the action of  $K$  on  $\mathfrak{g}/\mathfrak{k}$  respects its group structure. Thus (\*\*) also follows from  $\bar{X} \in (\mathfrak{g}/\mathfrak{k})^K(S)$ , proving (i) and (ii).

For (iii)–(v), put

$$\mathfrak{g}(\mathcal{M}) = \underline{\text{Lie}}(G/S, \mathcal{M})$$

and define  $\mathfrak{k}(\mathcal{M})$ ,  $\mathfrak{z}(\mathcal{M})$ , and  $\mathfrak{n}(\mathcal{M})$  similarly. If  $G/S$  satisfies (E), then

$$\mathfrak{g}(\mathcal{M} \oplus \mathcal{N}) \simeq \mathfrak{g}(\mathcal{M}) \times_S \mathfrak{g}(\mathcal{N}),$$

and hence, by Lemma 5.2.2(i),

$$\mathfrak{g}(\mathcal{M} \oplus \mathcal{N})^K \simeq \mathfrak{g}(\mathcal{M})^K \times_S \mathfrak{g}(\mathcal{N})^K.$$

It follows that  $\mathfrak{z}(\mathcal{M} \oplus \mathcal{N}) \simeq \mathfrak{z}(\mathcal{M}) \times_S \mathfrak{z}(\mathcal{N})$ , so  $Z$  satisfies (E). If  $K/S$  also satisfies (E), then

$$\begin{aligned} (\mathfrak{g}/\mathfrak{k})(\mathcal{M} \oplus \mathcal{N}) &\simeq (\mathfrak{g}/\mathfrak{k})(\mathcal{M}) \times_S (\mathfrak{g}/\mathfrak{k})(\mathcal{N}), \\ (\mathfrak{g}/\mathfrak{k})^K(\mathcal{M} \oplus \mathcal{N}) &\simeq (\mathfrak{g}/\mathfrak{k})^K(\mathcal{M}) \times_S (\mathfrak{g}/\mathfrak{k})^K(\mathcal{N}), \end{aligned}$$

and we have the following commutative diagram.

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathfrak{k}(\mathcal{M} \oplus \mathcal{N}) & \longrightarrow & \mathfrak{n}(\mathcal{M} \oplus \mathcal{N}) & \longrightarrow & (\mathfrak{n}/\mathfrak{k})(\mathcal{M} \oplus \mathcal{N}) \longrightarrow 0 \\ & & \downarrow \wr & & \downarrow & & \downarrow \wr \\ 0 & \longrightarrow & \mathfrak{k}(\mathcal{M}) \times_S \mathfrak{k}(\mathcal{N}) & \longrightarrow & \mathfrak{n}(\mathcal{M}) \times_S \mathfrak{n}(\mathcal{N}) & \longrightarrow & (\mathfrak{n}/\mathfrak{k})(\mathcal{M}) \times_S (\mathfrak{n}/\mathfrak{k})(\mathcal{N}) \longrightarrow 0 \end{array}$$

Figure 56: The exact-sequence comparison proving condition (E) for  $N$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp2-14oct24.pdf, component-local p. 48;  
combined-reader p. 106.*

It follows that

$$\mathfrak{n}(\mathcal{M} \oplus \mathcal{N}) \simeq \mathfrak{n}(\mathcal{M}) \times_S \mathfrak{n}(\mathcal{N}),$$

so  $N$  satisfies (E).

Now write  $\mathfrak{g} = \mathfrak{g}(\mathcal{O}_S) = \underline{\text{Lie}}(G/S)$ , and similarly  $\mathfrak{k}, \mathfrak{z}, \mathfrak{n}$ . If  $G$  is good, then  $\mathfrak{g}$  is a good  $\mathbf{O}_S$ -module, and Lemma 5.2.2(ii) shows that  $\mathfrak{z} \simeq \mathfrak{g}^K$  is good; since  $Z$  satisfies (E),  $Z$  is

good. If  $K$  is also good, then  $\mathfrak{k}$  is good, and Lemmas 5.2.1(iv) and 5.2.2(ii) show that  $\mathfrak{g}/\mathfrak{k}$  and  $(\mathfrak{g}/\mathfrak{k})^K$  are good. From the exact sequence

$$0 \longrightarrow \mathfrak{k} \longrightarrow \mathfrak{n} \longrightarrow (\mathfrak{g}/\mathfrak{k})^K \longrightarrow 0$$

and Lemma 5.2.1(iv),  $\mathfrak{n}$  is good. Finally, if  $G$  is very good, so that  $[x, x] = 0$  identically for  $x \in \mathfrak{g}$ , then  $Z$  and  $N$  are plainly very good. This proves (iii)–(v).

**Corollary 5.2.3.1.** *If the group law of  $\underline{\text{Lie}}(G/S)$  is commutative, then*

$$\underline{\text{Lie}}(\underline{\text{Centr}}(G)/S) = \underline{\text{Lie}}(G/S)^G.$$

**Corollary 5.2.3.2.** *If the group law of  $\underline{\text{Lie}}(G/S)$  is commutative and  $K$  is a normal subgroup of  $G$ , then*

$$(\underline{\text{Lie}}(G/S)/\underline{\text{Lie}}(K/S))^K = \underline{\text{Lie}}(G/S)/\underline{\text{Lie}}(K/S).$$

### 5.3. Linear representations

Let  $G$  be a good  $S$ -group acting linearly on a good  $\mathbf{O}_S$ -module  $F$  through

$$\rho : G \longrightarrow \underline{\text{Aut}}_{\mathbf{O}_S\text{-mod}}(F).$$

In 4.7.1 and 4.8.2 we defined the corresponding linear representation

$$\rho' : \underline{\text{Lie}}(G/S) \longrightarrow \underline{\text{End}}_{\mathbf{O}_S\text{-mod}}(F).$$

The sub- $S$ -groups  $\underline{\text{Norm}}_G(E)$  and  $\underline{\text{Centr}}_G(E)$  are defined for every subset  $E$  of  $F$ , and in particular for every  $\mathbf{O}_S$ -submodule  $E$ .

**Definition 5.3.0.** For every  $S' \rightarrow S$ , put

$$\underline{\text{Norm}}_{\underline{\text{Lie}}(G/S)}(E)(S') = \{X \in \underline{\text{Lie}}(G/S)(S') \mid \rho'(X)E_{S'} \subset E_{S'}\},$$

$$\underline{\text{Centr}}_{\underline{\text{Lie}}(G/S)}(E)(S') = \{X \in \underline{\text{Lie}}(G/S)(S') \mid \rho'(X)E_{S'} = 0\}.$$

This construction works for every linear representation of an  $\mathbf{O}_S$ -Lie algebra in the sense of 4.9, and the two subobjects obtained are  $\mathbf{O}_S$ -submodules stable under the bracket.

**Theorem 5.3.1.** *Let  $G$  be a good  $S$ -group acting linearly on a good  $\mathbf{O}_S$ -module  $F$ , and let  $E$  be an  $\mathbf{O}_S$ -submodule of  $F$ .*

(i) *One has*

$$\underline{\text{Lie}}(\underline{\text{Centr}}_G(E)/S) = \underline{\text{Centr}}_{\underline{\text{Lie}}(G/S)}(E),$$

*and  $\underline{\text{Centr}}_G(E)$  is a good  $S$ -group; it is very good if  $G$  is very good.*

(ii) *Suppose that  $E$  is a good  $\mathbf{O}_S$ -module. Then<sup>98</sup>*

$$\underline{\text{Lie}}(\underline{\text{Norm}}_G(E)/S) = \underline{\text{Norm}}_{\underline{\text{Lie}}(G/S)}(E),$$

*and  $\underline{\text{Norm}}_G(E)$  is a good  $S$ -group; it is very good if  $G$  is very good.*

---

<sup>98</sup>Editor's note: The original stated the following equality without a hypothesis on  $E$ ; this has been corrected. See 6.5.1 for counterexamples.

The proof is left to the reader.

**5.3.2.** Let  $G$  be a good  $S$ -group. The preceding results apply in particular to the adjoint representation. Let  $E$  be a good  $\mathbf{O}_S$ -submodule of  $\underline{\mathrm{Lie}}(G/S)$ .<sup>99</sup> It has a centralizer and a normalizer in  $G$ , and by 5.3.1 their Lie algebras are the usual centralizer and normalizer of  $E$ , computed using the bracket:

$$\begin{aligned}\underline{\mathrm{Centr}}_{\underline{\mathrm{Lie}}(G/S)}(E)(S') &= \{X \in \underline{\mathrm{Lie}}(G/S)(S') \mid [X, E_{S'}] = 0\}, \\ \underline{\mathrm{Norm}}_{\underline{\mathrm{Lie}}(G/S)}(E)(S') &= \{X \in \underline{\mathrm{Lie}}(G/S)(S') \mid [X, E_{S'}] \subset E_{S'}\}.\end{aligned}$$

**5.3.3.** Let  $K$  be a sub- $S$ -group of  $G$ . Then  $\underline{\mathrm{Lie}}(K/S)$  is an  $\mathbf{O}_S$ -submodule of  $\underline{\mathrm{Lie}}(G/S)$ . Suppose that it is a good  $\mathbf{O}_S$ -module,<sup>100</sup> as happens when  $K$  is a good  $S$ -group. Plainly,

$$\begin{aligned}\underline{\mathrm{Centr}}_G(K) &\subset \underline{\mathrm{Centr}}_G(\underline{\mathrm{Lie}}(K/S)), \\ \underline{\mathrm{Norm}}_G(K) &\subset \underline{\mathrm{Norm}}_G(\underline{\mathrm{Lie}}(K/S)).\end{aligned}$$

Hence, by 5.3.1,

$$\begin{aligned}\underline{\mathrm{Lie}}(\underline{\mathrm{Centr}}_G(K)/S) &\subset \underline{\mathrm{Centr}}_{\underline{\mathrm{Lie}}(G/S)}(\underline{\mathrm{Lie}}(K/S)), \\ \underline{\mathrm{Lie}}(\underline{\mathrm{Norm}}_G(K)/S) &\subset \underline{\mathrm{Norm}}_{\underline{\mathrm{Lie}}(G/S)}(\underline{\mathrm{Lie}}(K/S)).\end{aligned}$$

None of these four inclusions is an equality in general; many examples will appear later. In particular, if  $K$  is a normal subgroup of  $G$ , then  $\underline{\mathrm{Lie}}(K/S)$  is an ideal of  $\underline{\mathrm{Lie}}(G/S)$ .

## 6. Miscellaneous remarks

### 6.1.

The bracket of two infinitesimal automorphisms can be defined for an  $S$ -functor  $X$  which is not necessarily a group: apply the results of this exposé to  $\underline{\mathrm{Aut}}_S(X)$ . To obtain a convenient formalism, one is led to assume that  $X$  is good, meaning that the  $\mathbf{O}_X$ -module  $T_{X/S}$  is good. If  $X$  is an  $S$ -group, this agrees with Definition 4.6.

### 6.2.

There exist functors with infinitesimal endomorphisms that are not automorphisms and hence, a fortiori, do not satisfy condition (E).<sup>101</sup> For a pointed set  $(E, x_0)$ , let  $M_A(E)$  be the free abelian monoid generated by  $E$ , and let  $M_{A,P}(E, x_0)$  be the quotient of  $M_A(E)$  by the equivalence relation generated by  $m \sim x_0 + m$ . The assignment

$$(E, x_0) \longmapsto M_{A,P}(E, x_0)$$

is left adjoint to the forgetful functor from abelian monoids to pointed sets;  $M_{A,P}(E, x_0)$  is the free abelian monoid on the pointed set  $(E, x_0)$ .

Now let  $X$  be the functor assigning to every scheme  $S$  the free abelian monoid on  $\mathbf{O}(S)$ , pointed by zero. Every morphism

$$f : S \longrightarrow I_{\mathbb{Z}} = \mathrm{Spec}(\mathbb{Z}[\varepsilon])$$

<sup>99</sup>Editor's note: This hypothesis has been added; cf. 6.5.1.

<sup>100</sup>Editor's note: This hypothesis has been added; cf. 6.5.1.

<sup>101</sup>Editor's note: The original has been expanded here. In particular, O. Gabber pointed out the correct definition of the free abelian monoid on a pointed set.

corresponds to a square-zero element  $u_f \in \mathbf{O}(S)$ , and hence defines an endomorphism of  $X(S)$  by  $x \mapsto x + u_f$ , where the sum is taken in  $M_{A,P}(\mathbf{O}(S), 0)$ . Thus we obtain an endomorphism  $\phi$  of

$$X_{I_{\mathbb{Z}}} = X \times_{\mathbb{Z}} I_{\mathbb{Z}}$$

defined, for  $f \in I_{\mathbb{Z}}(S)$  and  $x \in X(S)$ , by

$$\phi((x, f)) = (x + u_f, f).$$

If  $f_0 : S \rightarrow I_{\mathbb{Z}}$  is the composite of the structural morphism  $S \rightarrow \mathrm{Spec}(\mathbb{Z})$  and the zero section, then  $u_{f_0} = 0$ , so  $\phi((x, f_0)) = (x, f_0)$ . Thus  $\phi$  induces the identity on  $X$ : it is an infinitesimal endomorphism of  $X$  which is plainly not an automorphism.

### 6.3.

There exist modules which are not good. First, the preceding counterexample can be modified slightly.<sup>102</sup> Let  $F(S)$  be the free  $\mathbf{O}(S)$ -module with basis the elements of  $\mathbf{O}(S)$ . Then  $F$  does not satisfy (E) relative to  $\mathrm{Spec}(\mathbb{Z})$ . Moreover, let  $G$  be the  $\mathbb{Z}$ -group defined by

$$G(S) = \mathrm{GL}_n(R(S)), \quad n \geq 2,$$

where  $R(S)$  is the  $\mathbf{O}(S)$ -algebra of the abelian group  $\mathbf{O}(S)$ . Then the  $\mathbb{Z}$ -group  $\underline{\mathrm{Lie}}(G/\mathbb{Z})$  is not commutative.

There are also the following counterexamples.<sup>103</sup> Let  $S = \mathrm{Spec}(k)$ , where  $k$  is a field of characteristic  $p > 0$ .

(a) Let  $F$  be the  $\mathbf{O}_S$ -module assigning to every  $S$ -scheme  $T$  the group

$$F(T) = \Gamma(T, \mathcal{O}_T),$$

with its  $\mathbf{O}(T)$ -module structure defined by  $r \cdot f = r^p f$ . As an  $S$ -functor of groups,  $F$  is isomorphic to  $\mathbb{G}_{a,S}$ . Thus  $F/S$  satisfies (E), and

$$\underline{\mathrm{Lie}}(F/S) \simeq \underline{\mathrm{Lie}}(\mathbb{G}_{a,S}/S) \simeq \mathbf{O}_S.$$

For each  $T \rightarrow S$ , the canonical morphism  $F \rightarrow \underline{\mathrm{Lie}}(F/S)$  is the identity map  $F(T) \rightarrow \mathbf{O}(T)$ . It respects the abelian-group structure but not the module structure, so  $F$  is not good (4.4.1).

(b) Let  $\alpha_{p,k}$  be the  $k$ -functor of groups defined by

$$\alpha_{p,k}(T) = \{x \in \mathbf{O}(T) \mid x^p = 0\}.$$

It is represented by  $\mathrm{Spec}(k[X]/(X^p))$ , hence is a very good  $S$ -group. It also carries an  $\mathbf{O}_S$ -module structure, but is not a good  $\mathbf{O}_S$ -module, because the canonical morphism

$$\alpha_{p,k} \longrightarrow \underline{\mathrm{Lie}}(\alpha_{p,k}/k) = \mathbb{G}_{a,k}$$

is not bijective.

<sup>102</sup>Editor's note: The following observation has been added.

<sup>103</sup>Editor's note: Example (a) has been expanded and example (b) has been added.



**6.4.**

For a group-valued functor  $G$  over  $S$ , the definitions give

$$(G/S \text{ satisfies (E)}) \iff (G \text{ is good}) \iff (G \text{ is very good}).$$

It would be interesting to prove or disprove the reverse implications.<sup>104</sup>

**6.5.**

Let  $\text{Nil}$  be the  $\mathbb{Z}$ -functor of groups defined by

$$\text{Nil}(S) = \{x \in \mathbf{O}(S) \mid \text{there exists } n \in \mathbb{N} \text{ with } x^n = 0\}.$$

<sup>105</sup> The functor  $\text{Nil}$  is very good but not representable. Let  $\text{Nil}^2$ ,  $\mathbf{O}_{\text{red}}$ , and  $F$  be the  $\mathbb{Z}$ -functors of groups assigning to  $S$ , respectively, the ideal  $\text{Nil}(S)^2$  and the groups

$$\mathbf{O}_{\text{red}}(S) = \mathbf{O}(S)/\text{Nil}(S), \quad F(S) = \mathbf{O}(S)/\text{Nil}(S)^2.$$

It is easy to see that

$$\underline{\text{Lie}}(\mathbf{O}_{\text{red}}/\mathbb{Z}) = 0,$$

so the  $\mathbf{O}_{\mathbb{Z}}$ -module  $\mathbf{O}_{\text{red}}$  is not good, although it is a good  $\mathbb{Z}$ -group. If  $\mathcal{M}, \mathcal{N}$  are free  $\mathbb{Z}$ -modules of finite rank, then

$$\begin{aligned} \text{Nil}^2(I_S(\mathcal{M} \oplus \mathcal{N})) &\simeq \text{Nil}^2(S) \oplus \text{Nil}(S) \otimes_{\mathbb{Z}} \mathcal{M} \\ &\quad \oplus \text{Nil}(S) \otimes_{\mathbb{Z}} \mathcal{N}, \end{aligned}$$

and hence

$$\begin{aligned} F(I_S(\mathcal{M} \oplus \mathcal{N})) &\simeq F(S) \oplus \mathbf{O}_{\text{red}}(S) \otimes_{\mathbb{Z}} \mathcal{M} \\ &\quad \oplus \mathbf{O}_{\text{red}}(S) \otimes_{\mathbb{Z}} \mathcal{N}. \end{aligned}$$

Thus  $F$  satisfies condition (E), while

$$\underline{\text{Lie}}(F/\mathbb{Z}) = \mathbf{O}_{\text{red}}.$$

Since  $\mathbf{O}_{\text{red}}$  is not a good  $\mathbf{O}_{\mathbb{Z}}$ -module,  $F$  is a  $\mathbb{Z}$ -group which satisfies (E) but is not good.

**6.5.1.** We now give the counterexamples announced in 5.3.1–5.3.3. Let  $S$  be a scheme, let  $F = \mathbf{O}_S^{\oplus 2}$  with the natural action of the good  $S$ -group  $G = \text{GL}_{2,S}$ , and let  $E$  be the  $\mathbf{O}_S$ -submodule of  $F$  formed by the pairs  $(x_1, x_2)$  for which  $x_2$  is nilpotent. Put

$$N = \underline{\text{Norm}}_G(E).$$

Then

$$\underline{\text{Lie}}(N/S) = \underline{\text{Lie}}(G/S),$$

whereas, for every  $S' \rightarrow S$ ,

$$\underline{\text{Norm}}_{\underline{\text{Lie}}(G/S)}(E)(S') = \left\{ \begin{pmatrix} a & b \\ x & c \end{pmatrix} \mid a, b, c, x \in \mathbf{O}(S'), \ x \text{ nilpotent} \right\}.$$

<sup>104</sup>Editor's note: A representable  $G$  is very good (4.11). For representability criteria see, for example, [MO67]. For automorphisms of an affine algebraic group over an algebraically closed field of characteristic zero, see [HM69].

<sup>105</sup>Editor's note: This paragraph has been added.

Consequently,

$$\underline{\mathrm{Lie}}(\underline{\mathrm{Norm}}_G(E)/S) \neq \underline{\mathrm{Norm}}_{\underline{\mathrm{Lie}}(G/S)}(E).$$

Taking the semidirect product  $G_1 = F \rtimes G$  gives an analogous counterexample in which  $E$  is an  $\mathbf{O}_S$ -submodule of  $\underline{\mathrm{Lie}}(G_1/S)$ . Moreover, with the notation above,

$$E = \underline{\mathrm{Lie}}(K/S),$$

where  $K$  is the subgroup  $\mathbf{O}_S \oplus \mathrm{Nil}^2$  of  $F$ ; that is,  $K(S')$  consists of the pairs  $(x_1, x_2)$  with  $x_2 \in \mathrm{Nil}(S')^2$ .

## Bibliography

106

- [DG70] M. Demazure and P. Gabriel, *Groupes Algébriques*, Masson & North-Holland, 1970.
- [HM69] G. Hochschild and G. D. Mostow, *Automorphisms of affine algebraic groups*, *J. Algebra* **13** (1969), 535–543.
- [MO67] H. Matsumura and F. Oort, *Representability of group functors and automorphisms of algebraic schemes*, *Invent. Math.* **4** (1967), 1–25.

---

<sup>106</sup>Editor’s note: Additional references cited in this exposé.

# Exposé III

## Infinitesimal Extensions

by *M. Demazure*

In this exposé we work in the following general situation. We have a scheme  $S$  and a coherent nilpotent ideal  $\mathcal{I}$  on  $S$ . We denote by  $S_n$  the closed subscheme of  $S$  defined by the ideal  $\mathcal{I}^{n+1}$ , for  $n \geq 0$ . In particular,  $S_0$  is defined by  $\mathcal{I}$ . Since  $\mathcal{I}$  is nilpotent,  $S_n = S$  for all sufficiently large  $n$ , and the  $S_i$  have the same underlying topological space. A typical example is the following:  $S$  is the spectrum of a local Artinian ring  $A$ , and  $\mathcal{I}$  is the ideal defined by the radical of  $A$ , so that  $S_0$  is the spectrum of the residue field of  $A$ .<sup>0</sup>

In this situation one is given certain data over  $S_0$  and seeks data over  $S$  that lift them, that is, data whose base change from  $S$  to  $S_0$  recovers the original data. This is done step by step, through the schemes  $S_n$ . At each step we shall define the obstructions that arise and classify the resulting solutions when they exist.

The passage from  $S_n$  to  $S_{n+1}$  may be generalized as follows. We have a scheme  $S$  and two ideals  $\mathcal{I} \supset \mathcal{J}$  such that  $\mathcal{I}\mathcal{J} = 0$ . In the preceding case  $S$ ,  $\mathcal{I}$ , and  $\mathcal{J}$  are respectively

$$S_{n+1}, \quad \mathcal{I}/\mathcal{I}^{n+2}, \quad \mathcal{I}^{n+1}/\mathcal{I}^{n+2}.$$

We denote by  $S_0$ , respectively  $S_{\mathcal{J}}$ , the closed subscheme of  $S$  defined by  $\mathcal{I}$ , respectively  $\mathcal{J}$ , and pose a problem of extending data from  $S_{\mathcal{J}}$  to  $S$ .

SGA 1, Exposé III, treated the problems of extending morphisms of schemes and of extending schemes. Here we consider the problems of extending group morphisms, group structures, and subgroups.

In §0 we have collected those results of SGA 1, Exposé III, that will be useful to us, putting them into the form best suited to our purpose and sparing the reader constant reference to that exposé.<sup>1</sup> Section 1 collects computations in group cohomology that will be used later and have nothing to do with scheme theory. Sections 2 and 3 treat, respectively, the extension of group morphisms and the extension of group structures. In §4 we briefly recall the proof of a result stated in TDTE IV concerning the extension of subschemes, and apply it to the problem of extending subgroups. For the remainder of the Seminar, only the result of §2 concerning the extension of group morphisms is indispensable.<sup>2</sup>

The idea of reducing problems of infinitesimal extension to the usual cohomological computations in group extensions was suggested by J. Giraud during the oral exposé, whose computations were appreciably more complicated and less transparent. Unfortunately, this

<sup>0</sup>Editor's note: version of October 14, 2024.

<sup>1</sup>Editor's note: the reader may prefer to begin with §1, which is easier and may serve as motivation and a guide to the results of §0.

<sup>2</sup>Editor's note: nevertheless, 3.10 is used in XXIV 1.13. Moreover, 4.37 is used in the proof of IX 3.2 bis and 3.6 bis, although those results also follow from results of Exposé X that do not use 4.37.

method seems to work well only for the first two problems studied, and we have been unable to avoid rather laborious computations for extensions of subgroups.

To simplify the language, we shall call a  $Y$ -functor, respectively a  $Y$ -scheme, and so on, a functor, respectively a scheme, and so on, equipped with a morphism to the functor  $Y$ . This extends the definitions of Exposé I, which concerned only representable  $Y$ .

## 0. Reminders from SGA 1, Exposé III, and miscellaneous remarks

We begin with a general definition.

**Definition 0.1.** *Let  $\mathcal{C}$  be a category,  $X$  an object of  $\widehat{\mathcal{C}}$ , and  $G$  a  $\widehat{\mathcal{C}}$ -group acting on  $X$ . We say that  $X$  is formally principal homogeneous<sup>3</sup> under  $G$  if the following equivalent conditions hold:*

- (i) *for every object  $T$  of  $\mathcal{C}$ , the set  $X(T)$  is either empty or principal homogeneous under  $G(T)$ ;*
- (ii) *the morphism of functors*

$$G \times X \longrightarrow X \times X, \quad (g, x) \longmapsto (gx, x),$$

*is an isomorphism.*

We shall now put the results of SGA 1, III §5<sup>4</sup> into the form most useful here. Throughout this section we use the following notation. On a scheme  $S$  we have two quasi-coherent ideals  $\mathcal{I}$  and  $\mathcal{J}$  such that

$$\mathcal{I} \supset \mathcal{J}, \quad \mathcal{I}\mathcal{J} = 0.$$

In particular,  $\mathcal{J}^2 = 0$ . We denote by  $S_0$ , respectively  $S_{\mathcal{J}}$ , the closed subscheme of  $S$  defined by  $\mathcal{I}$ , respectively  $\mathcal{J}$ . For every  $S$ -functor  $X$ , we systematically denote by  $X_0$  and  $X_{\mathcal{J}}$  the functors obtained by base change from  $S$  to  $S_0$  and  $S_{\mathcal{J}}$ . We use the same notation for a morphism.

**Definition 0.1.1.**<sup>5</sup> *Let  $X$  be an  $S$ -functor. Define a functor  $X^+$  over  $S$  by*

$$\mathrm{Hom}_S(Y, X^+) = \mathrm{Hom}_{S_{\mathcal{J}}}(Y_{\mathcal{J}}, X_{\mathcal{J}}) = \mathrm{Hom}_S(Y_{\mathcal{J}}, X)$$

*for a variable  $S$ -scheme  $Y$ . In the notation of Exposé II, §1,*

$$X^+ \simeq \prod_{S_{\mathcal{J}}/S} X_{\mathcal{J}}.$$

The identity morphism of  $X_{\mathcal{J}}$  defines, by construction, an  $S$ -morphism

$$p_X : X \longrightarrow X^+.$$

Explicitly,<sup>6</sup> for every  $S$ -scheme  $Y$ , the map

$$p_X(Y) : \mathrm{Hom}_S(Y, X) \longrightarrow \mathrm{Hom}_S(Y, X^+) = \mathrm{Hom}_S(Y_{\mathcal{J}}, X)$$

<sup>3</sup>Editor's note: one also calls such an object a *pseudo-torsor*; see EGA IV<sub>4</sub>, 16.5.15. The more general notion of a formally homogeneous object, not necessarily principal homogeneous, is defined in Exposé IV, 6.7.1.

<sup>4</sup>Editor's note: see also EGA IV<sub>4</sub>, 16.5.14–18.

<sup>5</sup>Editor's note: the numbering 0.1.1 through 0.1.13 has been added to make the definitions and results contained here more visible.

<sup>6</sup>Editor's note: the following sentence has been added, and the next two remarks have been expanded.

is induced by the morphism  $Y_{\mathcal{J}} \rightarrow Y$ .

**Remark 0.1.2.** If  $X$  is an  $S$ -group functor, then  $X_{\mathcal{J}}$  is an  $S_{\mathcal{J}}$ -group functor. The defining formula for  $X^+$  therefore equips  $X^+$  with an  $S$ -group-functor structure, and the description above shows that

$$p_X : X \longrightarrow X^+$$

is a morphism of  $S$ -group functors.

**Remark 0.1.3.** For every  $S$ -group functor  $Y$ , one has

$$\mathrm{Hom}_{S\text{-gr}}(Y, X^+) = \mathrm{Hom}_{S_{\mathcal{J}}\text{-gr}}(Y_{\mathcal{J}}, X_{\mathcal{J}}).$$

Indeed, let  $f \in \mathrm{Hom}_S(Y, X^+)$  correspond to  $f_{\mathcal{J}} \in \mathrm{Hom}_{S_{\mathcal{J}}}(Y_{\mathcal{J}}, X_{\mathcal{J}})$ . The condition that  $f$  be an  $S$ -group morphism is that, for every  $T \rightarrow S$  and every  $y, y' \in Y(T)$ ,

$$f(y \cdot y') = f(y) \cdot f(y'),$$

which is equivalent to

$$f_{\mathcal{J}}(y_{\mathcal{J}}) \cdot f_{\mathcal{J}}(y'_{\mathcal{J}}) = f_{\mathcal{J}}((y \cdot y')_{\mathcal{J}}).$$

Since  $(y \cdot y')_{\mathcal{J}} = y_{\mathcal{J}} \cdot y'_{\mathcal{J}}$ , this is precisely the condition that  $f_{\mathcal{J}}$  be a group morphism. Applying this to  $Y = X$ , we again see that  $p_X$ , which corresponds to  $\mathrm{id}_{X_{\mathcal{J}}}$ , is a morphism of  $S$ -group functors.

Return now to the general case and suppose that  $X$  is an  $S$ -scheme. An  $S$ -morphism from a variable  $S$ -scheme  $Y$  to  $X^+$  is, by definition, an  $S_{\mathcal{J}}$ -morphism  $g_{\mathcal{J}} : Y_{\mathcal{J}} \rightarrow X_{\mathcal{J}}$ . We define an  $X^+$ -functor in abelian groups  $L_X$  as follows.

**Scholium 0.1.4.**<sup>7</sup> If  $\pi : Y \rightarrow S$  is a morphism of schemes and  $M$  is an  $\mathcal{O}_S$ -module, we denote the inverse image  $\pi^*M$  by

$$M \otimes_{\mathcal{O}_S} \mathcal{O}_Y.$$

If  $\mathcal{J}$  is an ideal of  $\mathcal{O}_S$ , we denote by  $\mathcal{J}\mathcal{O}_Y$  the ideal of  $\mathcal{O}_Y$  that is the image of

$$\pi^*(\mathcal{J}) = \mathcal{J} \otimes_{\mathcal{O}_S} \mathcal{O}_Y \longrightarrow \mathcal{O}_Y.$$

For every morphism of  $S$ -schemes  $f : Z \rightarrow Y$ , there is an epimorphism of  $\mathcal{O}_Z$ -modules

$$f^*(\mathcal{J}\mathcal{O}_Y) \longrightarrow \mathcal{J}\mathcal{O}_Z. \quad (0.1.4)$$

It follows from the commutative diagram below.

$$\begin{array}{ccc} \mathcal{J} \otimes_{\mathcal{O}_S} \mathcal{O}_Y \otimes_{\mathcal{O}_Y} \mathcal{O}_Z & \xlongequal{\quad} & \mathcal{J} \otimes_{\mathcal{O}_S} \mathcal{O}_Z \\ \downarrow & & \downarrow \\ f^*(\mathcal{J}\mathcal{O}_Y) = (\mathcal{J}\mathcal{O}_Y) \otimes_{\mathcal{O}_Y} \mathcal{O}_Z & \longrightarrow & \mathcal{J}\mathcal{O}_Z. \end{array}$$

Figure 57: The tensor-product diagram yielding the epimorphism (0.1.4).

*Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 4;  
combined-reader p. 114.*

<sup>7</sup>Editor's note: this scholium has been added.

**Definition 0.1.5.**<sup>8</sup> Let  $X$  be an  $S$ -scheme. For every  $X^+$ -scheme  $Y$ , given by a morphism

$$g_{\mathcal{J}} : Y_{\mathcal{J}} \longrightarrow X_{\mathcal{J}},$$

set

$$\mathrm{Hom}_{X^+}(Y, L_X) = \mathrm{Hom}_{\mathcal{O}_{Y_0}}(g_0^*(\Omega_{X_0/S_0}^1), \mathcal{J} \mathcal{O}_Y),$$

where  $\Omega_{X_0/S_0}^1$  is the module of relative differentials of  $X_0$  over  $S_0$ , and  $\mathcal{J} \mathcal{O}_Y$  is regarded as an  $\mathcal{O}_{Y_0}$ -module because it is annihilated by  $\mathcal{J}$ .

Then  $L_X$  is an  $X^+$ -functor in abelian groups.<sup>9</sup> Indeed, for every  $X^+$ -morphism  $f : Z \rightarrow Y$ , the functor  $f_0^*$  and the morphism

$$f_0^*(\mathcal{J} \mathcal{O}_Y) \longrightarrow \mathcal{J} \mathcal{O}_Z$$

of (0.1.4) induce a natural homomorphism  $L_X(f)$ :

$$\begin{aligned} & \mathrm{Hom}_{\mathcal{O}_{Y_0}}(g_0^*(\Omega_{X_0/S_0}^1), \mathcal{J} \mathcal{O}_Y) \\ & \longrightarrow \mathrm{Hom}_{\mathcal{O}_{Z_0}}(f_0^* g_0^*(\Omega_{X_0/S_0}^1), f_0^*(\mathcal{J} \mathcal{O}_Y)) \\ & \longrightarrow \mathrm{Hom}_{\mathcal{O}_{Z_0}}(f_0^* g_0^*(\Omega_{X_0/S_0}^1), \mathcal{J} \mathcal{O}_Z). \end{aligned}$$

Finally,  $L_X(f)$  has the following simple local description. The schemes  $Y$  and  $Y_{\mathcal{J}}$  have the same underlying topological space, as do  $V$  and  $V_{\mathcal{J}}$  for every open subset  $V$  of  $Y$ . Let

$$U = \mathrm{Spec}(A), \quad V = \mathrm{Spec}(B), \quad W = \mathrm{Spec}(C)$$

be affine opens such that  $U$  lies over  $\mathrm{Spec}(\Lambda) \subset S$ ,  $g_{\mathcal{J}}(V_{\mathcal{J}}) \subset U$ , and  $W \subset f^{-1}(V)$ . Write  $I, J$  for the ideals of  $\Lambda$  corresponding to  $\mathcal{J}, \mathcal{J}$ . Then  $f$ , respectively  $g_{\mathcal{J}}$ , induces a morphism of  $\Lambda$ -algebras

$$\theta : B \longrightarrow C, \quad \text{respectively} \quad \varphi : A \longrightarrow B/JB,$$

and clearly  $\theta(JB) \subset JC$ . On the other hand,

$$m \in \mathrm{Hom}_{\mathcal{O}_{Y_0}}(g_0^*(\Omega_{X/S}^1), \mathcal{J} \mathcal{O}_Y)$$

induces an element  $D$  of

$$\begin{aligned} \mathrm{Hom}_{\mathcal{O}_{V_0}}(g_0^*(\Omega_{U/S}^1), \mathcal{J} \mathcal{O}_V) &= \mathrm{Hom}_{B/IB}(\Omega_{A/\Lambda}^1 \otimes_A B/IB, JB) \\ &= \mathrm{Der}_{\Lambda}(A, JB), \end{aligned}$$

and the image of  $L_X(f)(m)$  in

$$\begin{aligned} \mathrm{Hom}_{\mathcal{O}_{W_0}}(f_0^* g_0^*(\Omega_{U/S}^1), \mathcal{J} \mathcal{O}_Z) &= \mathrm{Hom}_{C/IC}(\Omega_{A/\Lambda}^1 \otimes_A C/IC, JC) \\ &= \mathrm{Der}_{\Lambda}(A, JC) \end{aligned}$$

is simply  $\theta \circ D$ .

<sup>8</sup>Editor's note: introducing the functor  $L_X$  gives an extension, functorial in  $Y$  and in  $X$ , of SGA 1, III, Proposition 5.1; see 0.1.8 and 0.1.9 below, as well as 0.1.10 and 0.2. Moreover, when  $X$  is an  $S$ -group functor,  $L_X$  gives rise, under certain hypotheses, to an  $S$ -group functor  $L'_X$  and to an exact sequence  $1 \rightarrow L'_X \rightarrow X \rightarrow X^+$ ; see 0.4, 0.9, and 0.11. These play an essential role in this exposé; see Theorems 2.1, 3.5, and 4.21.

<sup>9</sup>Editor's note: what follows has been added.

**Remark 0.1.6.**<sup>10</sup> Let  $f : X \rightarrow W$  be an  $S$ -morphism. It induces an  $S$ -morphism

$$f^+ : X^+ \longrightarrow W^+$$

defined as follows. If  $g \in \text{Hom}_S(Y, X^+)$  corresponds to an  $S$ -morphism  $g_{\mathcal{J}} : Y_{\mathcal{J}} \rightarrow X$ , then  $f^+(g)$  is the element  $f \circ g_{\mathcal{J}}$  of  $\text{Hom}_S(Y, W^+)$ . The following diagram is commutative.

$$\begin{array}{ccc} X & \xrightarrow{f} & W \\ p_X \downarrow & & \downarrow p_W \\ X^+ & \xrightarrow{f^+} & W^+ \end{array} .$$

Figure 58: Naturality of the morphisms  $p_X$  and  $p_W$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 5;  
combined-reader p. 115.*

**Reminders 0.1.7.**<sup>11</sup> In this paragraph, given an  $S$ -scheme  $X$ , we “recall” certain functorial properties of the module of differentials  $\Omega_{X/S}^1$  and of the first infinitesimal neighborhood of the diagonal,  $\Delta_{X/S}^{(1)}$ . These properties follow readily from EGA IV<sub>4</sub>, §§16.1–16.4, but do not appear there explicitly.

(a) Let us begin by recalling the following facts (cf. EGA II, §§1.2–1.5). Let  $g : Y \rightarrow X$  be a morphism of schemes, let  $\pi : X' \rightarrow X$  be an affine  $X$ -scheme, and let  $\mathcal{B} = \pi_*(\mathcal{O}_{X'})$  be the corresponding quasi-coherent  $\mathcal{O}_X$ -algebra. Then the  $Y$ -scheme  $Y \times_X X'$  is affine and corresponds to the quasi-coherent  $\mathcal{O}_Y$ -algebra  $g^*(\mathcal{B})$ , and there is a commutative diagram of bijections:

$$\begin{array}{ccc} \text{Hom}_X(Y, X') & \xrightarrow{\sim} & \text{Hom}_Y(Y, Y \times_X X') \\ \wr \downarrow & & \downarrow \wr \\ \text{Hom}_{\mathcal{O}_X\text{-alg.}}(\mathcal{B}, g_*(\mathcal{O}_Y)) & \xrightarrow{\sim} & \text{Hom}_{\mathcal{O}_Y\text{-alg.}}(g^*(\mathcal{B}), \mathcal{O}_Y). \end{array}$$

Figure 59: The affine-scheme/algebra bijections after base change by  $g$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 5;  
combined-reader p. 115.*

Moreover, these bijections are functorial in the pair  $(X, X')$ . More precisely, let  $W'$  be an affine  $W$ -scheme corresponding to the quasi-coherent  $\mathcal{O}_W$ -algebra  $\mathcal{A}$ , and suppose that one has a commutative diagram of morphisms of schemes as follows.

<sup>10</sup>Editor’s note: this remark has been expanded.

<sup>11</sup>Editor’s note: This paragraph has been added (see also [DG70], §I.4, nos. 1–2).

$$\begin{array}{ccccc}
 & & X' & \xrightarrow{f'} & W' \\
 & \nearrow g' & \downarrow & & \downarrow \\
 Y & \xrightarrow{g} & X & \xrightarrow{f} & W
 \end{array}$$

Figure 60: The scheme morphisms used to express functoriality in  $(X, X')$ .

Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 5;  
combined-reader p. 115.

Denote by

$$\varphi : \mathcal{A} \longrightarrow f_*(\mathcal{B}), \quad \varphi^\sharp : f^*(\mathcal{A}) \longrightarrow \mathcal{B}$$

the algebra morphisms associated with  $f'$ , and by

$$\theta : \mathcal{B} \longrightarrow g_*(\mathcal{O}_Y), \quad \theta^\sharp : g^*(\mathcal{B}) \longrightarrow \mathcal{O}_Y$$

those associated with a variable  $X$ -morphism  $g' : Y \rightarrow X'$ . Then the following diagram is commutative, where  $Y$  is viewed as a  $W$ -scheme through  $f \circ g$ .

$$\begin{array}{ccc}
 \mathrm{Hom}_X(Y, X') & \xrightarrow{g' \mapsto f' \circ g'} & \mathrm{Hom}_W(Y, W') \\
 \downarrow \wr & \searrow \simeq & \\
 & & \mathrm{Hom}_{\mathcal{O}_W\text{-alg.}}(\mathcal{A}, f_*g_*(\mathcal{O}_Y)) \\
 & \nearrow \theta \mapsto f_*(\theta) \circ \phi & \downarrow \wr \\
 \mathrm{Hom}_{\mathcal{O}_X\text{-alg.}}(\mathcal{B}, g_*(\mathcal{O}_Y)) & \xrightarrow{\theta \mapsto \theta \circ \phi^\sharp} & \mathrm{Hom}_{\mathcal{O}_X\text{-alg.}}(f^*(\mathcal{A}), g_*(\mathcal{O}_Y)) \\
 \downarrow \wr & & \downarrow \wr \\
 \mathrm{Hom}_{\mathcal{O}_Y\text{-alg.}}(g^*(\mathcal{B}), \mathcal{O}_Y) & \xrightarrow{\theta^\sharp \mapsto \theta^\sharp \circ g^*(\phi^\sharp)} & \mathrm{Hom}_{\mathcal{O}_Y\text{-alg.}}(g^*f^*(\mathcal{A}), \mathcal{O}_Y).
 \end{array}$$

Figure 61: Functoriality of the affine-scheme/algebra correspondences.

Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 6;  
combined-reader p. 116.

(b) Let now  $X$  be an  $S$ -scheme. Let  $\Omega_{X/S}^1$  be the module of differentials of  $X$  over  $S$ , and let  $\Delta_{X/S}^{(1)}$  be the first infinitesimal neighborhood of the diagonal immersion  $\delta_X : X \rightarrow X \times_S X$ . It is a subscheme of  $X \times_S X$ , in which the diagonal  $\Delta_{X/S}$  is a closed subscheme. Write  $\mathrm{pr}_X^i$ ,  $i = 1, 2$ , for the two projections  $X \times_S X \rightarrow X$ , and write  $\pi_X$  for the restriction of  $\mathrm{pr}_X^1$  to  $\Delta_{X/S}^{(1)}$ .

On the one hand, every morphism  $f : X \rightarrow W$  of  $S$ -schemes induces an  $S$ -morphism

$$\Delta^{(1)}f : \Delta_{X/S}^{(1)} \longrightarrow \Delta_{W/S}^{(1)}$$



for which the following diagram is commutative.

$$\begin{array}{ccccccc}
 X & \xrightarrow{\delta_X} & \Delta_{X/S}^{(1)} & \longrightarrow & X \times_S X & \xrightarrow{\text{pr}_X^i} & X \\
 \downarrow f & & \downarrow \Delta^{(1)} f & & \downarrow f \times f & & \downarrow f \\
 W & \xrightarrow{\delta_W} & \Delta_{W/S}^{(1)} & \longrightarrow & W \times_S W & \xrightarrow{\text{pr}_W^i} & W.
 \end{array}$$

Figure 62: Functoriality of the first infinitesimal neighborhood of the diagonal.

*Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 6; combined-reader p. 116.*

On the other hand, through the projection  $\pi_X$ , the scheme  $\Delta_{X/S}^{(1)}$  is an affine  $X$ -scheme: it is the spectrum of the augmented quasi-coherent  $\mathcal{O}_X$ -algebra

$$\mathcal{P}_{X/S}^1 = \mathcal{O}_X \oplus \Omega_{X/S}^1,$$

where  $\Omega_{X/S}^1$  is a square-zero ideal. The augmentation is the  $\mathcal{O}_X$ -algebra morphism

$$\varepsilon_X : \mathcal{P}_{X/S}^1 \longrightarrow \mathcal{O}_X$$

that vanishes on  $\Omega_{X/S}^1$  and corresponds to the closed immersion  $\delta_X : X \hookrightarrow \Delta_{X/S}^{(1)}$ . Hence every morphism of  $S$ -schemes  $f : X \rightarrow W$  induces a morphism of augmented  $\mathcal{O}_X$ -algebras

$$f^*(\mathcal{P}_{W/S}^1) = \mathcal{O}_X \oplus f^*(\Omega_{W/S}^1) \longrightarrow \mathcal{P}_{X/S}^1 = \mathcal{O}_X \oplus \Omega_{X/S}^1,$$

or, equivalently, a morphism of  $\mathcal{O}_X$ -modules

$$f_{X/W/S} : f^*(\Omega_{W/S}^1) \longrightarrow \Omega_{X/S}^1;$$

see EGA IV<sub>4</sub>, (16.4.3.6), and (16.4.18.2) for the notation  $f_{X/W/S}$ .

Since  $\pi_X : \Delta_{X/S}^{(1)} \rightarrow X$  is affine, part (a) shows that  $\Delta^{(1)}f$  is completely determined by  $f_{X/W/S}$ . For every  $X$ -scheme  $g : Y \rightarrow X$ , the set

$$\text{Hom}_X(Y, \Delta_{X/S}^{(1)}) \simeq \text{Hom}_{\mathcal{O}_Y\text{-alg}}(\mathcal{O}_Y \oplus g^*(\Omega_{X/S}^1), \mathcal{O}_Y)$$

identifies with a subset of  $\text{Hom}_{\mathcal{O}_Y}(g^*(\Omega_{X/S}^1), \mathcal{O}_Y)$ , namely

$$\widetilde{\text{Hom}}_{\mathcal{O}_Y}(g^*(\Omega_{X/S}^1), \mathcal{O}_Y),$$

the subset formed by the  $\mathcal{O}_Y$ -morphisms  $\psi : g^*(\Omega_{X/S}^1) \rightarrow \mathcal{O}_Y$  for which  $\text{Im}(\psi)$  is a square-zero ideal of  $\mathcal{O}_Y$ .<sup>12</sup>

Consequently, apply part (a) to the following diagram.

<sup>12</sup>Editor's note: In the proof of SGA 1, III 5.1, the sentence "But the algebra homomorphisms  $\mathcal{A} \rightarrow \mathcal{O}_Y$  correspond ..." must be corrected accordingly; the remainder of the cited passage is unchanged.

$$\begin{array}{ccccc}
& & \Delta_{X/S}^{(1)} & \xrightarrow{\Delta^{(1)}f} & \Delta_{W/S}^{(1)} \\
& \nearrow g' & \downarrow & & \downarrow \\
Y & \xrightarrow{g} & X & \xrightarrow{f} & W
\end{array}$$

Figure 63: The diagram to which the affine functoriality statement is applied.

*Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 7; combined-reader p. 117.*

Taking into account that  $\Delta^{(1)}f$  is the restriction of  $f \times f$  to  $\Delta_{X/S}^{(1)}$ , one obtains the following commutative diagram, functorial in the  $X$ -scheme  $g : Y \rightarrow X$ .

$$\begin{array}{ccc}
\mathrm{Hom}_X(Y, X \times_S X) & \xrightarrow{(g, g') \mapsto (f \circ g, f \circ g')} & \mathrm{Hom}_W(Y, \Delta_{W/S}^{(1)}) \\
\uparrow & & \uparrow \\
(0.1.7 (*)) \quad \mathrm{Hom}_X(Y, \Delta_{X/S}^{(1)}) & \xrightarrow{g' \mapsto (\Delta^{(1)}f) \circ g'} & \mathrm{Hom}_W(Y, \Delta_{W/S}^{(1)}) \\
\downarrow \simeq & & \downarrow \simeq \\
\mathrm{Hom}_{\mathcal{O}_Y}^\square(g^*(\Omega_{X/S}^1), \mathcal{O}_Y) & \xrightarrow{\psi \mapsto \psi \circ g^*(f_{X/W/S})} & \mathrm{Hom}_{\mathcal{O}_Y}^\square(g^*f^*(\Omega_{W/S}^1), \mathcal{O}_Y).
\end{array}$$

Figure 64: The functorial comparison of first-neighborhood morphisms and differential maps, equation 0.1.7 (\*).

*Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 7; combined-reader p. 117.*

**Remark 0.1.7.1.** Let us conclude this paragraph with the following remark, which will be useful later (cf. 0.1.10). Denote by  $\tilde{L}_X$  the  $X$ -functor that associates to every  $X$ -scheme  $g : Y \rightarrow X$  the set

$$\mathrm{Hom}_{\mathcal{O}_Y}(g^*(\Omega_{X/S}^1), \mathcal{O}_Y),$$

and denote by  $\tilde{L}_f : \tilde{L}_X \rightarrow \tilde{L}_W$  the morphism of functors defined above: it sends  $\psi \in \tilde{L}_X(Y)$  to  $\psi \circ g^*(f_{X/W/S})$ . The preceding discussion gives a commutative diagram of functors.

$$\begin{array}{ccccc}
X \times_X L_X^\square & \xleftarrow{\sim} & \Delta_{X/S}^{(1)} & \hookrightarrow & X \times_S X \\
\downarrow f \times L_f^\square & & \downarrow \Delta^{(1)} f & & \downarrow f \times f \\
W \times_W L_W^\square & \xleftarrow{\sim} & \Delta_{W/S}^{(1)} & \hookrightarrow & W \times_S W.
\end{array}$$

Figure 65: Functoriality of  $\tilde{L}_X$  and the first infinitesimal neighborhood.

Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 7;  
combined-reader p. 117.

**Theorem 0.1.8.** (SGA 1, III 5.1)<sup>13</sup> Let  $Y$  and  $X$  be two  $S$ -schemes, let  $\mathcal{J}$  be a quasi-coherent square-zero ideal of  $\mathcal{O}_Y$ , let  $Y_{\mathcal{J}}$  be the closed subscheme of  $Y$  defined by  $\mathcal{J}$ , and let  $g_{\mathcal{J}} : Y_{\mathcal{J}} \rightarrow X$  be an  $S$ -morphism.

- (a) The set  $P(g_{\mathcal{J}})$  of  $S$ -morphisms  $g : Y \rightarrow X$  extending  $g_{\mathcal{J}}$  is either empty or principal homogeneous under the abelian group

$$\mathrm{Hom}_{\mathcal{O}_{Y_{\mathcal{J}}}}(g_{\mathcal{J}}^*(\Omega_{X/S}^1), \mathcal{J}).$$

- (b) Suppose that  $i : Y_0 \hookrightarrow Y_{\mathcal{J}}$  is the closed immersion defined by a quasi-coherent ideal  $\mathcal{I} \supset \mathcal{J}$  such that  $\mathcal{I}\mathcal{J} = 0$ , and set  $g_0 = g_{\mathcal{J}} \circ i$ . Then the preceding abelian group is isomorphic to

$$\mathrm{Hom}_{\mathcal{O}_{Y_0}}(g_0^*(\Omega_{X/S}^1), \mathcal{I}).$$

*Proof.* Part (b) follows at once from part (a). Indeed, since  $\mathcal{J}$  is annihilated by  $\mathcal{I}$ , it may be regarded as an  $\mathcal{O}_{Y_0}$ -module, and adjunction gives

$$\mathrm{Hom}_{\mathcal{O}_{Y_{\mathcal{J}}}}(g_{\mathcal{J}}^*(\Omega_{X/S}^1), \mathcal{J}) = \mathrm{Hom}_{\mathcal{O}_{Y_0}}(i^*g_{\mathcal{J}}^*(\Omega_{X/S}^1), \mathcal{J}).$$

To prove part (a), we may suppose that  $P(g_{\mathcal{J}}) \neq \emptyset$ , that is, that an  $S$ -morphism  $g : Y \rightarrow X$  extending  $g_{\mathcal{J}}$  exists. Write  $j : Y_{\mathcal{J}} \hookrightarrow Y$  for the immersion. Then  $P(g_{\mathcal{J}})$  is the set of  $S$ -morphisms  $g' : Y \rightarrow X$  such that  $g' \circ j = g_{\mathcal{J}}$ . Giving such a  $g'$  is equivalent to giving an  $S$ -morphism

$$h : Y \longrightarrow X \times_S X$$

such that  $\mathrm{pr}_1 \circ h = g$  and  $h_{\mathcal{J}} = \delta \circ g_{\mathcal{J}}$ , where  $h_{\mathcal{J}} = h \circ j$  and  $\delta : X \hookrightarrow X \times_S X$  is the diagonal immersion.

<sup>13</sup>Editor's note: The statement and proof of SGA 1, III 5.1 are reproduced here, since they are needed below to prove Proposition 0.2; see already the following Corollary 0.1.9. Moreover,  $\mathcal{J}$  (respectively  $\mathcal{I}$ ) is another calligraphic form of the letter  $J$  (respectively  $I$ ). Returning to the earlier notation, with  $\mathcal{J} \subset \mathcal{I}$  ideals of  $\mathcal{O}_S$  such that  $\mathcal{I}\mathcal{J} = 0$ , this result will be applied with  $\mathcal{J} = \mathcal{J}\mathcal{O}_Y$  (respectively  $\mathcal{I} = \mathcal{I}\mathcal{O}_Y$ ).

$$\begin{array}{ccc}
X \times_S X & \xleftarrow{h_{\mathcal{J}} = \delta \circ g_{\mathcal{J}}} & Y_{\mathcal{J}} \\
\text{pr}_1 \downarrow & \nwarrow h & \downarrow j \\
X & \xleftarrow{g} & Y.
\end{array}$$

Figure 66: The morphism  $h$  determined by two extensions of  $g_{\mathcal{J}}$ .

Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 8;  
combined-reader p. 118.

Since  $h_{\mathcal{J}}$  factors through  $\delta$ , and since  $Y$  lies in the first infinitesimal neighborhood of the immersion  $j : Y_{\mathcal{J}} \rightarrow Y$  because  $\mathcal{J}^2 = 0$ , functoriality (EGA IV<sub>4</sub>, 16.2.2(i)) shows that the desired morphisms  $h$  factor uniquely through  $\Delta_{X/S}^{(1)}$ ; see 0.1.7. Put

$$Y' = \Delta_{X/S}^{(1)} \times_X Y, \quad Y'_{\mathcal{J}} = Y' \times_Y Y_{\mathcal{J}} = \Delta_{X/S}^{(1)} \times_X Y_{\mathcal{J}}.$$

The desired morphisms  $h$  are then in bijection with the sections  $u$  of  $Y' \rightarrow Y$  extending the section  $u_{\mathcal{J}} = (\delta \circ g_{\mathcal{J}}, \text{id})$  of  $Y'_{\mathcal{J}} \rightarrow Y_{\mathcal{J}}$ . On the other hand,  $Y'$  (respectively  $Y'_{\mathcal{J}}$ ) is affine over  $Y$  (respectively  $Y_{\mathcal{J}}$ ) and corresponds to the quasi-coherent algebra

$$\mathcal{A} = \mathcal{O}_Y \oplus g^*(\Omega_{X/S}^1), \quad \text{respectively} \quad \mathcal{A}_{\mathcal{J}} = \mathcal{A} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y_{\mathcal{J}}} = \mathcal{O}_{Y_{\mathcal{J}}} \oplus g_{\mathcal{J}}^*(\Omega_{X/S}^1).$$

Let  $\varepsilon : \mathcal{A} \rightarrow \mathcal{O}_Y$  be the canonical augmentation, that is, the  $\mathcal{O}_Y$ -algebra morphism that vanishes on  $g^*(\Omega_{X/S}^1)$ , and define  $\varepsilon_{\mathcal{J}} : \mathcal{A}_{\mathcal{J}} \rightarrow \mathcal{O}_{Y_{\mathcal{J}}}$  similarly. Then

$$\Gamma(Y'/Y) \simeq \text{Hom}_{\mathcal{O}_Y\text{-alg}}(\mathcal{A}, \mathcal{O}_Y), \quad \Gamma(Y'_{\mathcal{J}}/Y_{\mathcal{J}}) \simeq \text{Hom}_{\mathcal{O}_{Y_{\mathcal{J}}}\text{-alg}}(\mathcal{A}_{\mathcal{J}}, \mathcal{O}_{Y_{\mathcal{J}}}).$$

Under these isomorphisms, the section  $u = (\delta \circ g, \text{id})$  (respectively  $u_{\mathcal{J}}$ ) corresponds to  $\varepsilon$  (respectively  $\varepsilon_{\mathcal{J}}$ ). Consequently,  $P(g_{\mathcal{J}})$  is in bijection with the algebra morphisms  $\mathcal{A} \rightarrow \mathcal{O}_Y$  that reduce to  $\varepsilon_{\mathcal{J}}$ ; under this bijection,  $g$  corresponds to  $\varepsilon$ .

Put  $M = g^*(\Omega_{X/S}^1)$ . Then  $\text{Hom}_{\mathcal{O}_Y\text{-alg}}(\mathcal{A}, \mathcal{O}_Y)$  identifies with the set of  $\mathcal{O}_Y$ -morphisms  $\psi : M \rightarrow \mathcal{O}_Y$  whose image is a square-zero ideal. We are interested in those morphisms that induce the zero morphism  $M \rightarrow \mathcal{O}_{Y_{\mathcal{J}}} = \mathcal{O}_Y/\mathcal{J}$ , that is, those that carry  $M$  into  $\mathcal{J}$ . Conversely, since  $\mathcal{J}^2 = 0$ , every  $\mathcal{O}_Y$ -morphism  $\phi : M \rightarrow \mathcal{J}$  comes from a unique algebra morphism  $\mathcal{A} \rightarrow \mathcal{O}_Y$  reducing to  $\varepsilon_{\mathcal{J}}$ . Finally,

$$\text{Hom}_{\mathcal{O}_Y}(g^*(\Omega_{X/S}^1), \mathcal{J}) = \text{Hom}_{\mathcal{O}_{Y_{\mathcal{J}}}}(g_{\mathcal{J}}^*(\Omega_{X/S}^1), \mathcal{J})$$

because  $\mathcal{J}^2 = 0$ , as in the proof of part (b). Thus one obtains a bijection

$$P(g_{\mathcal{J}}) \simeq \text{Hom}_{\mathcal{O}_{Y_{\mathcal{J}}}}(g_{\mathcal{J}}^*(\Omega_{X/S}^1), \mathcal{J})$$

under which  $g$  corresponds to the zero morphism.

For every

$$m \in \text{Hom}_{\mathcal{O}_{Y_{\mathcal{J}}}}(g_{\mathcal{J}}^*(\Omega_{X/S}^1), \mathcal{J}),$$

write  $m \cdot g$  for the element of  $P(g_{\mathcal{J}})$  associated with  $g$  and  $m$  by the preceding bijection. We have already seen that  $0 \cdot g = g$ ; it remains to prove

$$m' \cdot (m \cdot g) = (m + m') \cdot g. \quad (0.1.8(*))$$

This may be checked locally.<sup>14</sup> Indeed, the two preceding morphisms  $Y \rightarrow X$  induce the same continuous map as  $g$  on the underlying topological spaces. It is therefore enough to check that, for every affine open  $U = \text{Spec}(A)$  of  $X$  lying over an affine open  $\text{Spec}(\Lambda)$  of  $S$ , and every affine open  $V = \text{Spec}(B)$  of  $g^{-1}(U)$ , they induce the same morphism of  $\Lambda$ -algebras  $A \rightarrow B$ .

Set  $J = \Gamma(V, \mathcal{J})$ , and let  $\varphi, \psi, \eta : A \rightarrow B$  be the morphisms induced respectively by  $g$ ,  $m \cdot g$ , and  $m' \cdot (m \cdot g)$ . They agree modulo  $J$ . There are unique expressions

$$\psi = \varphi + D, \quad \eta = \psi + D',$$

where  $D$  belongs to

$$\text{Der}_{\varphi}(A, J) = \{\delta \in \text{Hom}_{\Lambda}(A, J) \mid \delta(ab) = \varphi(a)\delta(b) + \varphi(b)\delta(a)\},$$

and  $D'$  belongs to  $\text{Der}_{\psi}(A, J)$ . But  $\text{Der}_{\varphi}(A, J) = \text{Der}_{\psi}(A, J)$ , since  $J^2 = 0$ , and both identify with

$$\text{Hom}_{B/J}(\Omega_{A/\Lambda}^1 \otimes_A B/J, J).$$

Under this identification,  $D$  corresponds to  $m$  and  $D'$  to  $m'$ . Hence  $\eta = \varphi + D + D'$ , while  $D + D'$  corresponds to  $m + m'$ , proving (0.1.8(\*)).

**Corollary 0.1.9.**<sup>15</sup> *Let  $X$  be an  $S$ -scheme, and retain the notation of 0.1.5. Then  $X$  carries a left action of the abelian  $X^+$ -group  $L_X$ , making  $X$  formally principal homogeneous under  $L_X$  over  $X^+$ . In other words, there is an isomorphism of  $X^+$ -functors*

$$L_X \times_{X^+} X \xrightarrow{\sim} X \times_{X^+} X, \quad (m, x) \mapsto (x, m \cdot x).$$

*Proof.* Let  $i_0 : X_0 \hookrightarrow X$  be the immersion. Since  $X_0 = X \times_S S_0$ , one first has

$$i_0^*(\Omega_{X/S}^1) \simeq \Omega_{X_0/S_0}^1$$

(EGA IV, 16.4.5).

Let  $Y$  be an  $X^+$ -scheme, given by an  $S$ -morphism  $g_{\mathcal{J}} : Y_{\mathcal{J}} \rightarrow X$ , and let  $g_0 : Y_0 \rightarrow X_0$  be obtained by base change. By 0.1.8, if  $\text{Hom}_{X^+}(Y, X)$  is nonempty, it is principal homogeneous under

$$\text{Hom}_{\mathcal{O}_{Y_0}}(g_0^* i_0^*(\Omega_{X/S}^1), \mathcal{J} \mathcal{O}_Y),$$

which identifies with

$$\text{Hom}_{\mathcal{O}_{Y_0}}(g_0^*(\Omega_{X_0/S_0}^1), \mathcal{J} \mathcal{O}_Y) = L_X(Y).$$

Thus there is a bijection

$$L_X(Y) \times \text{Hom}_{X^+}(Y, X) \xrightarrow{\sim} \text{Hom}_{X^+}(Y, X \times_{X^+} X), \quad (m, g) \mapsto (g, m \cdot g).$$

We show that this is functorial in  $Y$ . Let  $f : Z \rightarrow Y$  be a morphism of  $S$ -schemes. It is enough to prove that the following diagram is commutative.

<sup>14</sup>Editor's note: The preceding argument is reproduced almost word for word from SGA 1, III 5.1; what follows has been expanded.

<sup>15</sup>Editor's note: This corollary has been added to SGA 1, III 5.1; it proves part (i) of Proposition 0.2 below.

$$\begin{array}{ccc}
L_X(Y) \times \mathrm{Hom}_{X+}(Y, X) & \xrightarrow{\sim} & \mathrm{Hom}_{X+}(Y, X \times_{X+} X) \\
\downarrow L_X(f) \times f & & \downarrow f \times f \\
L_X(Z) \times \mathrm{Hom}_{X+}(Z, X) & \xrightarrow{\sim} & \mathrm{Hom}_{X+}(Z, X \times_{X+} X).
\end{array}$$

Figure 67: Functoriality in  $Y$  of the principal-homogeneous bijection.

*Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 10; combined-reader p. 120.*

If  $\mathrm{Hom}_{X+}(Y, X) = \emptyset$ , there is nothing to prove. It is therefore enough to show that, for every  $S$ -morphism  $g : Y \rightarrow X$  extending  $g_{\mathcal{J}}$  and every  $m \in L_X(Y)$ , one has

$$(m \cdot g) \circ f = L_X(f)(m) \cdot (g \circ f). \quad (0.1.9(*))$$

These two  $S$ -morphisms  $Z \rightarrow X$  agree on  $Z_{\mathcal{J}}$  with  $g_{\mathcal{J}} \circ f_{\mathcal{J}}$ ; in particular, they induce the same continuous map as  $g \circ f$  on the underlying topological spaces. It therefore suffices to check the equality on local rings. Let  $z \in Z$ , put  $y = f(z)$ ,  $x = g(y)$ , and let  $A, B, C$  denote respectively the local rings  $\mathcal{O}_{X,x}, \mathcal{O}_{Y,y}, \mathcal{O}_{Z,z}$ . Let  $s$  be the image of  $x$  in  $S$ , put  $\Lambda = \mathcal{O}_{S,s}$ , let  $J$  and  $I$  be the ideals of  $\Lambda$  corresponding to  $\mathcal{J}$  and  $\mathcal{I}$ , and let  $\varphi, \psi : A \rightarrow B$  and  $\theta : B \rightarrow C$  be the  $\Lambda$ -algebra morphisms induced by  $g, m \cdot g$ , and  $f$ , respectively. Then  $m$  induces an element  $D$  of

$$\mathrm{Hom}_{B/IB}(\Omega_{A/\Lambda}^1 \otimes_A B/IB, JB) = \mathrm{Der}_{\Lambda}(A, JB),$$

and  $\psi = \varphi + D$ . Thus  $(m \cdot g) \circ f$  corresponds to

$$\theta \circ \psi = \theta \circ \varphi + \theta \circ D.$$

By 0.1.5,  $\theta \circ D$  is the image of  $L_X(f)(m)$  in

$$\mathrm{Hom}_{C/IC}(\Omega_{A/\Lambda}^1 \otimes_A C/IC, JC) = \mathrm{Der}_{\Lambda}(A, JC).$$

Consequently,  $\theta \circ \varphi + \theta \circ D$  is the image of  $L_X(f)(m) \cdot (g \circ f)$ . This proves (0.1.9(\*)).

**Corollary 0.1.10.**<sup>16</sup>

- (a) *The functor  $L_X$  depends functorially on  $X$ . For every  $S$ -morphism  $f : X \rightarrow W$ , there is an  $S$ -morphism  $L_f : L_X \rightarrow L_W$  that is a morphism of abelian groups over  $f^+$ ; that is, the following diagram is commutative,*

<sup>16</sup>Editor's note: The functoriality of  $L_X$  in  $X$  has been made explicit, in particular in part (b) below.

$$\begin{array}{ccc}
L_X & \xrightarrow{L_f} & L_W \\
\downarrow & & \downarrow \\
X^+ & \xrightarrow{f^+} & W^+
\end{array}$$

Figure 68: The morphism  $L_f$  over  $f^+$ .

*Temporary Loop-1 image; source locator: Polo–Gille Exp3-14oct24.pdf, component-local p. 11;  
combined-reader p. 121.*

and, for every  $Y \rightarrow X^+$ , the map

$$L_f(Y) : \text{Hom}_{X^+}(Y, L_X) \longrightarrow \text{Hom}_{W^+}(Y, L_W),$$

where  $Y$  lies over  $W^+$  through  $f^+$ , is a homomorphism of abelian groups.

(b) Moreover, the following diagram is commutative.

$$\begin{array}{ccc}
L_X \times_{X^+} X & \xrightarrow{\sim} & X \times_{X^+} X \\
L_f \times f \downarrow & & \downarrow f \times f \\
L_W \times_{W^+} W & \xrightarrow{\sim} & W \times_{W^+} W
\end{array} .$$

Figure 69: Compatibility of  $L_f$  with the formally principal homogeneous actions.

*Temporary Loop-1 image; source locator: Polo–Gille Exp3-14oct24.pdf, component-local p. 11;  
combined-reader p. 121.*

*Proof.* For part (a),  $L_f$  is induced by the morphism of  $\mathcal{O}_{X_0}$ -modules

$$f_{X_0/W_0/S_0} : f_0^*(\Omega_{W_0/S_0}^1) \longrightarrow \Omega_{X_0/S_0}^1$$

(cf. 0.1.7(b)). For every  $X^+$ -scheme  $Y$ , given by an  $S$ -morphism  $g_{\mathcal{J}} : Y_{\mathcal{J}} \rightarrow X$ , one has the following commutative diagram, functorial in  $Y$ .

$$\begin{array}{ccc}
\text{Hom}_{\mathcal{O}_{Y_0}}(g_0^*(\Omega_{X_0/S_0}^1), \mathcal{J}\mathcal{O}_Y) & \xrightarrow{L_f(Y)} & \text{Hom}_{\mathcal{O}_{Y_0}}(g_0^*f_0^*(\Omega_{W_0/S_0}^1), \mathcal{J}\mathcal{O}_Y) \\
\downarrow & & \downarrow \\
\{g_{\mathcal{J}}\} & \xrightarrow{\quad\quad\quad} & \{f_{\mathcal{J}} \circ g_{\mathcal{J}}\}
\end{array}$$

Figure 70: The homomorphism  $L_f(Y)$  on the differential-module descriptions.

*Temporary Loop-1 image; source locator: Polo–Gille Exp3-14oct24.pdf, component-local p. 11;  
combined-reader p. 121.*

Here  $L_f(Y)$  is the map

$$\psi \longmapsto \psi \circ g_0^*(f_{X_0/W_0/S_0}),$$

which is indeed a homomorphism of abelian groups.<sup>17</sup>

We now prove part (b). Let  $Y$  be an  $X^+$ -scheme. If  $\text{Hom}_{X^+}(Y, X) = \emptyset$ , there is nothing to prove. Thus let  $g \in \text{Hom}_{X^+}(Y, X)$ . We must show that, for every  $m \in L_X(Y)$ ,

$$f \circ (m \cdot g) = L_f(Y)(m) \cdot (f \circ g). \quad (0.1.10 (*))$$

With  $g$  fixed,  $\text{Hom}_X(Y, X \times_{X^+} X)$  is a subset of  $\text{Hom}_X(Y, \Delta_{X/S}^{(1)})$ , and

$$\begin{aligned} L_X(Y) &= \text{Hom}_{\mathcal{O}_{Y_0}}(g_0^*(\Omega_{X_0/S_0}^1), \mathcal{I}_{\mathcal{O}_Y}) \\ &= \text{Hom}_{\mathcal{O}_Y}(g^*(\Omega_{X/S}^1), \mathcal{I}_{\mathcal{O}_Y}), \end{aligned}$$

which is a subset of  $\tilde{L}_X(Y)$ ; see 0.1.7. Moreover,  $L_f(Y)$  is the restriction of  $\tilde{L}_f(Y)$  to  $L_X(Y)$ . Finally, the bijection

$$L_X(Y) \xrightarrow{\sim} \text{Hom}_X(Y, X \times_{X^+} X), \quad m \longmapsto (g, m \cdot g),$$

is the inverse of the restriction to  $L_X(Y) \subset \tilde{L}_X(Y)$  of the bijection

$$\text{Hom}_X(Y, \Delta_{X/S}^{(1)}) \xrightarrow{\sim} \{g\} \times \tilde{L}_X(Y)$$

considered in 0.1.7.1. Consequently, equality (0.1.10 (\*)) follows from (0.1.7 (\*)). Indeed, let  $g'$  be the  $X$ -morphism  $Y \rightarrow \Delta_{X/S}^{(1)}$  defined by  $(g, m \cdot g)$ . The element of  $\tilde{L}_W(Y)$  corresponding to  $(f \circ g, f \circ g')$  is  $\tilde{L}_f(Y)(m) = L_f(Y)(m)$ . Thus

$$L_f(m) \cdot (f \circ g) = f \circ (m \cdot g).$$

**Lemma 0.1.11.** *Let  $X, X'$  be two  $S$ -schemes. There is a commutative diagram*

$$\begin{array}{ccc} L_X \times_S L_{X'} & \xrightarrow{\sim} & L_{X \times_S X'} \\ \downarrow & & \downarrow \\ X^+ \times_S X'^+ & \xrightarrow{\sim} & (X \times_S X')^+ \end{array} .$$

Figure 71: Compatibility of  $L_X$  and  $X^+$  with products.

*Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 12; combined-reader p. 122.*

*Proof.*<sup>18</sup> For every  $S$ -scheme  $Y$ ,

$$\begin{aligned} \text{Hom}_S(Y, X^+ \times_S X'^+) &= \text{Hom}_S(Y, X^+) \times \text{Hom}_S(Y, X'^+) \\ &\simeq \text{Hom}_S(Y_{\mathcal{I}}, X) \times \text{Hom}_S(Y_{\mathcal{I}}, X') \\ &= \text{Hom}_S(Y, (X \times_S X')^+). \end{aligned}$$

<sup>17</sup>Editor's note: Observe that  $L_f$  depends only on  $f_0 : X_0 \rightarrow W_0$ .

<sup>18</sup>Editor's note: the proof has been expanded.



Thus

$$X^+ \times_S X'^+ \simeq (X \times_S X')^+.$$

Now let  $Y$  be a scheme over  $X^+ \times_S X'^+$ , represented by a morphism

$$h : Y_{\mathcal{J}} \longrightarrow X \times_S X'.$$

Let  $p, q$  be the projections from  $X \times_S X'$  to  $X, X'$ , and put  $f = p \circ h$ ,  $g = q \circ h$ . Since

$$\Omega_{(X_0 \times_{S_0} X'_0)/S_0}^1 \simeq p_0^*(\Omega_{X_0/S_0}^1) \oplus q_0^*(\Omega_{X'_0/S_0}^1)$$

(EGA IV<sub>4</sub>, 16.4.23), there is a natural isomorphism

$$\begin{aligned} & \mathrm{Hom}_{\mathcal{O}_{Y_0}}(f_0^*(\Omega_{X_0/S_0}^1), \mathcal{J} \mathcal{O}_Y) \times \mathrm{Hom}_{\mathcal{O}_{Y_0}}(g_0^*(\Omega_{X'_0/S_0}^1), \mathcal{J} \mathcal{O}_Y) \\ & \simeq \mathrm{Hom}_{\mathcal{O}_{Y_0}}(h_0^*(\Omega_{(X_0 \times_{S_0} X'_0)/S_0}^1), \mathcal{J} \mathcal{O}_Y). \end{aligned}$$

In other words,

$$L_X(Y) \times L_{X'}(Y) \simeq L_{X \times_S X'}(Y).$$

**Remark 0.1.12.**<sup>19</sup> Let  $\mathcal{C}$  be a category stable under fiber products, let  $S$  be an object of  $\mathcal{C}$ , and let  $T_1, T_2$  be two objects over  $S$ . For  $i = 1, 2$ , let  $L_i$  and  $X_i$  be two objects over  $T_i$ , as in the diagram below.




Figure 72: The objects  $L_i, X_i$  over  $T_i$ , and  $T_i$  over  $S$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 12;  
combined-reader p. 122.*

There is then a natural isomorphism

$$(L_1 \times_{T_1} X_1) \times_S (L_2 \times_{T_2} X_2) \simeq (L_1 \times_S L_2) \times_{T_1 \times_S T_2} (X_1 \times_S X_2).$$

Consequently, the preceding lemma gives:

**Corollary 0.1.13.** *Let  $X_1, X_2$  be two  $S$ -schemes. There is a commutative diagram of isomorphisms:*

$$\begin{array}{ccc} L_{X_1 \times_S X_2} \times_{(X_1 \times_S X_2)^+} (X_1 \times_S X_2) & & \\ \downarrow (0.1.11) \simeq & \searrow \simeq & \\ (L_{X_1} \times_S L_{X_2}) \times_{(X_1^+ \times_S X_2^+)} (X_1 \times_S X_2) & \xrightarrow[(0.1.12)]{\simeq} & (L_{X_1} \times_{X_1^+} X_1) \times_S (L_{X_2} \times_{X_2^+} X_2). \end{array}$$

Figure 73: The product isomorphisms of Corollary 0.1.13.

*Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 13;  
combined-reader p. 123.*

<sup>19</sup>Editor's note: this remark and the corollary that follows have been added.

We can now state:

**Proposition 0.2.** *For every  $S$ -scheme  $X$ , one can define a left action of the  $X^+$ -abelian group  $L_X$  on the  $X^+$ -object  $X$ , with the following properties:*

- (i) *this action makes  $X$  formally principal homogeneous under  $L_X$  over  $X^+$ ; that is, the morphism*

$$L_X \times_{X^+} X \longrightarrow X \times_{X^+} X$$

*is an isomorphism of  $X^+$ -functors;*

- (ii) *the action is functorial in the  $S$ -scheme  $X$ : for every  $S$ -morphism  $f : X \rightarrow W$ , the following diagram is commutative;*

$$\begin{array}{ccc} L_X \times_{X^+} X & \longrightarrow & X \\ L_f \times f \downarrow & & \downarrow f \\ L_W \times_{W^+} W & \longrightarrow & W \end{array} \quad ;$$

Figure 74: Functoriality of the action in Proposition 0.2(ii).

*Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 13;  
combined-reader p. 123.*

- (iii) *the action commutes with fiber products: for all  $S$ -schemes  $X_1, X_2$ , the following diagram is commutative.*

$$\begin{array}{ccc} L_{X_1 \times_S X_2} \times_{(X_1 \times_S X_2)^+} (X_1 \times_S X_2) & \longrightarrow & X_1 \times_S X_2 \\ \simeq \downarrow & \searrow \simeq & \uparrow \\ (L_{X_1} \times_S L_{X_2}) \times_{(X_1 \times_S X_2)^+} (X_1 \times_S X_2) & \xrightarrow{\sim} & (L_{X_1} \times_{X_1^+} X_1) \times_S (L_{X_2} \times_{X_2^+} X_2) \end{array}$$

Figure 75: Compatibility of the action with fiber products.

*Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 13;  
combined-reader p. 123.*

*Proof.*<sup>20</sup> Parts (i) and (ii) follow respectively from Corollaries 0.1.9 and 0.1.10. To prove (iii), write

$$P(X) = L_X \times_{X^+} X$$

for every  $S$ -scheme  $X$ . Applying (ii) to the projections

$$p_i : X_1 \times_S X_2 \longrightarrow X_i$$

gives the following commutative squares.

<sup>20</sup>Editor's note: the original has been modified and expanded to account for the preceding additions; in particular, those additions incorporate, with details, the content of original page 89.

$$\begin{array}{ccc}
P(X_1 \times_S X_2) & \longrightarrow & X_1 \times_S X_2 \\
L_{p_i} \times p_i \downarrow & & \downarrow p_i \\
P(X_i) & \longrightarrow & X_i
\end{array}$$

Figure 76: The square induced by each projection  $p_i$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 13;  
combined-reader p. 123.*

For  $i = 1, 2$ , these yield the commutative square

$$\begin{array}{ccc}
P(X_1 \times_S X_2) & \longrightarrow & X_1 \times_S X_2 \\
\downarrow & & \parallel \\
P(X_1) \times_S P(X_2) & \longrightarrow & X_1 \times_S X_2.
\end{array}$$

Figure 77: The product square in the proof of Proposition 0.2(iii).

*Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 14;  
combined-reader p. 124.*

Combining this square with Corollary 0.1.13 shows that its vertical arrow is an isomorphism and gives precisely the commutative diagram asserted in (iii).

**Remark 0.3.** Suppose that the  $X^+$ -scheme  $Y$  is flat over  $S$  (cf. SGA 1, IV). We may then write

$$\mathrm{Hom}_{X^+}(Y, L_X) = \mathrm{Hom}_{\mathcal{O}_{Y_0}}(g_0^*(\Omega_{X_0/S_0}^1), \mathcal{I} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Y_0}).$$

**Remark 0.4.** Let  $\pi_0 : X_0 \rightarrow S_0$  be the structure morphism, and suppose that there is an  $\mathcal{O}_{S_0}$ -module  $\omega_{X_0/S_0}^1$  such that

$$\Omega_{X_0/S_0}^1 \simeq \pi_0^*(\omega_{X_0/S_0}^1).$$

(This occurs, in particular, when  $X_0$  is an  $S_0$ -group; cf. II 4.11.) Define a functor  $L'_X$  over  $S$  by

$$\mathrm{Hom}_S(Y, L'_X) = \mathrm{Hom}_{\mathcal{O}_{Y_0}}(\omega_{X_0/S_0}^1 \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Y_0}, \mathcal{I} \mathcal{O}_Y). \quad (0.4.1)$$

Then

$$\mathrm{Hom}_{X^+}(Y, L_X) = \mathrm{Hom}_S(Y, L'_X)$$

for every  $X^+$ -scheme  $Y$ ; in other words,

$$L_X = L'_X \times_S X^+.$$

Since  $L_X \times_{X^+} X = L'_X \times_S X$ , the action of  $L_X$  on  $X$  induces an action of  $L'_X$  on  $X$ , and this action respects  $p_X : X \rightarrow X^+$ .<sup>21</sup> Indeed, if  $Y$  is an  $S$ -scheme,  $h : Y \rightarrow X$  is an

<sup>21</sup>Editor's note: what follows has been expanded.

$S$ -morphism, and  $m \in L'_X(Y)$ , then  $h$  and  $m \cdot h$  have the same restriction to  $Y_{\mathcal{J}}$ ; that is,

$$p_X(m \cdot h) = p_X(h).$$

**Remark 0.5.** Keep the hypotheses and notation of 0.4, and suppose in addition that  $Y$  is an  $S$ -scheme flat over  $S$ . Then

$$\mathrm{Hom}_{X^+}(Y, L_X) = \mathrm{Hom}_S(Y, L'_X) = \mathrm{Hom}_{S_0}(Y_0, L_{0X}),$$

where the  $S_0$ -functor in abelian groups  $L_{0X}$  is defined by the following identity (with  $T$  the variable  $S_0$ -scheme):

$$\mathrm{Hom}_{S_0}(T, L_{0X}) = \mathrm{Hom}_{\mathcal{O}_T}(\omega_{X_0/S_0}^1 \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_T, \mathcal{J} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_T). \quad (0.5.1)$$

Thus, in the notation of II 1, we have shown that  $L'_X$  and  $\prod_{S_0/S} L_{0X}$  have the same restriction to the full subcategory of  $(\mathbf{Sch})/S$  whose objects are the  $S$ -schemes flat over  $S$ .

**Remark 0.6.** Keep the hypotheses and notation of 0.5,<sup>22</sup> and suppose in addition that  $\pi_0 : X_0 \rightarrow S_0$  has a section  $\varepsilon_0$ . Then

$$\omega_{X_0/S_0}^1 \simeq \varepsilon_0^*(\Omega_{X_0/S_0}^1).$$

First, independently of this last hypothesis, one has

$$\mathrm{Hom}_{S_0}(T, L_{0X}) = \Gamma\left(T, \mathcal{H}om_{\mathcal{O}_T}(\omega_{X_0/S_0}^1 \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_T, \mathcal{J} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_T)\right).$$

Suppose now that  $\omega_{X_0/S_0}^1$  admits a finite presentation (cf. EGA 0<sub>I</sub>, 5.2.5), as is in particular the case when  $X_0$  is locally of finite presentation over  $S_0$  (cf. EGA IV<sub>4</sub>, 16.4.22). If  $T$  is flat over  $S_0$ , it then follows from EGA 0<sub>I</sub>, 6.7.6 that

$$\mathcal{H}om_{\mathcal{O}_T}(\omega_{X_0/S_0}^1 \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_T, \mathcal{J} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_T) \simeq \mathcal{H}om_{\mathcal{O}_{S_0}}(\omega_{X_0/S_0}^1, \mathcal{J}) \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_T,$$

whence

$$\mathrm{Hom}_{S_0}(T, L_{0X}) = \Gamma\left(T, \mathcal{H}om_{\mathcal{O}_{S_0}}(\omega_{X_0/S_0}^1, \mathcal{J}) \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_T\right).$$

Using the notation  $\mathbf{W}(\cdot)$  of I 4.6.1, we have therefore proved that, for every  $S_0$ -scheme  $T$  flat over  $S_0$ ,

$$\mathrm{Hom}_{S_0}(T, L_{0X}) = \mathrm{Hom}_{S_0}\left(T, \mathbf{W}\left(\mathcal{H}om_{\mathcal{O}_{S_0}}(\omega_{X_0/S_0}^1, \mathcal{J})\right)\right).$$

In summary, if  $\omega_{X_0/S_0}^1$  admits a finite presentation and one restricts to the category of  $S$ -schemes flat over  $S$ , then

$$L'_X = \prod_{S_0/S} \mathbf{W}\left(\mathcal{H}om_{\mathcal{O}_{S_0}}(\omega_{X_0/S_0}^1, \mathcal{J})\right), \quad (0.6.1)$$

and  $\mathcal{H}om_{\mathcal{O}_{S_0}}(\omega_{X_0/S_0}^1, \mathcal{J})$  is a quasi-coherent  $\mathcal{O}_{S_0}$ -module, by EGA I, 9.1.1.

Finally, suppose moreover that  $\omega_{X_0/S_0}^1$  is locally free of finite rank, for example that  $X_0$  is smooth over  $S_0$  (in which case it is automatically locally of finite presentation over  $S_0$ ). Then

$$\mathcal{H}om_{\mathcal{O}_{S_0}}(\omega_{X_0/S_0}^1, \mathcal{J}) \simeq \underline{\mathrm{Lie}}(X_0/S_0) \otimes_{\mathcal{O}_{S_0}} \mathcal{J}, \quad (0.6.2)$$

<sup>22</sup>Editor's note: the following hypothesis, which was implicit in the original, has been added.

where, by an abuse of language (since  $X_0$  need not be an  $S_0$ -group),  $\underline{\text{Lie}}(X_0/S_0)$  denotes the dual of the  $\mathcal{O}_{S_0}$ -module  $\omega_{X_0/S_0}^1$ .<sup>23</sup>

Proposition 0.2 (and its proof) has two important corollaries.<sup>24</sup>

**Corollary 0.7.** *Let  $X$  be an  $S$ -scheme.*

- (a) *Every  $S$ -endomorphism of  $X$  inducing the identity on  $X_{\mathcal{J}}$  is an automorphism.*
- (b) *There is an exact sequence of groups*

$$0 \longrightarrow \text{Hom}_{\mathcal{O}_{X_0}}(\Omega_{X_0/S_0}^1, \mathcal{J} \mathcal{O}_X) \xrightarrow{i} \text{Aut}_S(X) \longrightarrow \text{Aut}_{S_{\mathcal{J}}}(X_{\mathcal{J}}).$$

- (c) *Let  $\text{Aut}_S(X)$  act on the first group by transport of structure. Then, for every  $u \in \text{Aut}_S(X)$  and  $m \in \text{Hom}_{\mathcal{O}_{X_0}}(\Omega_{X_0/S_0}^1, \mathcal{J} \mathcal{O}_X)$ ,*

$$i(u \cdot m) = u i(m) u^{-1}.$$

*Proof.* By 0.2(i),  $\text{Hom}_{X^+}(X, X)$  is a principal homogeneous set under  $\text{Hom}_{X^+}(X, L_X)$ , since it is certainly nonempty: it contains the marked point given by the identity automorphism of  $X$ .<sup>25</sup> Consequently, the map  $m \mapsto m \cdot \text{id}_X$  induces a bijection

$$\text{Hom}_{\mathcal{O}_{X_0}}(\Omega_{X_0/S_0}^1, \mathcal{J} \mathcal{O}_X) = L_X(X) \xrightarrow{\sim} \text{Hom}_{X^+}(X, X).$$

Let  $m \in L_X(X)$ , and let  $f = m' \cdot \text{id}_X$  be an element of  $\text{Hom}_{X^+}(X, X)$ . Applying 0.2(ii) to  $f$  gives

$$f \circ (m \cdot \text{id}_X) = L_f(X)(m) \cdot f = L_f(X)(m) \cdot (m' \cdot \text{id}_X).$$

On the other hand, because  $f$  is an  $X^+$ -endomorphism of  $X$ , one has  $f_{\mathcal{J}} = \text{id}_{X_{\mathcal{J}}}$ , hence  $f_0 = \text{id}_{X_0}$ . Since  $L_f$  depends only on  $f_0$  (cf. the editor's note (17) in 0.1.10), it follows that  $L_f(X)(m) = m$ . Thus the preceding equality becomes

$$(m' \cdot \text{id}_X) \circ (m \cdot \text{id}_X) = m \cdot (m' \cdot \text{id}_X) = (m + m') \cdot \text{id}_X.$$

This shows that the bijection  $m \mapsto m \cdot \text{id}_X$  carries the group law of  $\text{Hom}_{X^+}(X, L_X)$  to composition of  $X^+$ -endomorphisms of  $X$ .

It follows first that every element of  $\text{Hom}_{X^+}(X, X)$  is invertible, proving part (a), and then that there is an exact sequence

$$0 \longrightarrow \text{Hom}_{X^+}(X, L_X) \xrightarrow{i} \text{Aut}_S(X) \longrightarrow \text{Aut}_{S_{\mathcal{J}}}(X_{\mathcal{J}}),$$

which proves part (b).

The morphism  $i$  just defined is functorial in  $X$  with respect to isomorphisms, because it is defined structurally from the action of  $L_X$  on  $X$  over  $X^+$ , an action that is itself

<sup>23</sup>Editor's note: this is justified by II 3.3 and 4.11. Indeed, the functor  $L_{X_0/S_0}^{\varepsilon_0}$ , the “tangent space to  $X_0$  over  $S_0$  at  $\varepsilon_0$ ,” is represented over  $S_0$  by the vector bundle  $\mathbf{V}(\varepsilon_0^*(\Omega_{X_0/S_0}^1)) = \mathbf{V}(\omega_{X_0/S_0}^1)$ . Its sheaf of sections is the dual module  $\mathcal{H}om_{\mathcal{O}_{S_0}}(\omega_{X_0/S_0}^1, \mathcal{O}_{S_0})$ ; this is  $\underline{\text{Lie}}(X_0/S_0)$  when  $X_0$  is an  $S_0$ -group and  $\varepsilon_0$  is the identity section.

<sup>24</sup>Editor's note: Corollary 0.7 bis, which was used implicitly in the proof of 0.8, has been added to 0.7. We point out that 0.8–0.12, as well as 0.17, which play an important technical role later in this exposé, are consequences of 0.7 and 0.7 bis.

<sup>25</sup>Editor's note: what follows has been expanded.

functorial in  $X$  by Proposition 0.2(ii).<sup>26</sup> Hence every automorphism  $u$  of  $X$  over  $S$  induces, by transport of structure, isomorphisms

$$h : \mathrm{Hom}_{X^+}(X, L_X) \xrightarrow{\sim} \mathrm{Hom}_{X^+}(X, L_X) \quad \text{and} \quad f : \mathrm{Aut}_S(X) \xrightarrow{\sim} \mathrm{Aut}_S(X)$$

such that the following diagram commutes.

$$\begin{array}{ccc} \mathrm{Hom}_{X^+}(X, L_X) & \xrightarrow{i} & \mathrm{Aut}_S(X) \\ h \downarrow & & \downarrow f \\ \mathrm{Hom}_{X^+}(X, L_X) & \xrightarrow{i} & \mathrm{Aut}_S(X) \end{array}$$

Figure 78: Functoriality of  $i$  under transport of structure.

*Temporary Loop-1 image; source locator: Polo–Gille Exp3-14oct24.pdf, component-local p. 16;  
combined-reader p. 126; printed p. 116.*

Thus  $f \circ i = i \circ h$ . On the other hand,  $f$  is described by the following commutative diagram.

$$\begin{array}{ccc} X & \xrightarrow{a} & X \\ u \downarrow & & \downarrow u \\ X & \xrightarrow{f(a)} & X \end{array}$$

Figure 79: The automorphism  $f$  acts by conjugation with  $u$ .

*Temporary Loop-1 image; source locator: Polo–Gille Exp3-14oct24.pdf, component-local p. 16;  
combined-reader p. 126; printed p. 116.*

In other words,

$$f(a) = u \circ a \circ u^{-1} \quad \text{for every } a \in \mathrm{Aut}_S(X).$$

Writing  $i(h(m)) = f(i(m))$  now gives the asserted formula.

**Corollary 0.7 bis.** *Let  $X$  be an  $S$ -scheme such that  $X_{\mathcal{J}}$  is an  $S_{\mathcal{J}}$ -monoid. Then  $L_X$  has the structure of an  $S$ -monoid, and there is a split exact sequence of  $S$ -monoids:*

$$1 \longrightarrow L'_X \xrightarrow{i} L_X \begin{array}{c} \xrightarrow{p} \\ \xleftarrow{s} \end{array} X^+ \longrightarrow 1$$

Figure 80: The split exact sequence of  $S$ -monoids in Corollary 0.7 bis.

*Temporary Loop-1 image; source locator: Polo–Gille Exp3-14oct24.pdf, component-local p. 17;  
combined-reader p. 127; printed p. 117.*

*The monoid law induced on  $L'_X$  coincides with its abelian-group structure. In particular, if  $X_{\mathcal{J}}$  is an  $S_{\mathcal{J}}$ -group, then  $L_X$  is an  $S$ -group and is the semidirect product of  $X^+$  and  $L'_X$ .*

<sup>26</sup>Editor's note: what follows has been expanded.

*Proof.* Since  $X_{\mathcal{J}}$  is an  $S_{\mathcal{J}}$ -monoid,

$$X^+ = \prod_{S_{\mathcal{J}}/S} X_{\mathcal{J}}$$

is an  $S$ -monoid; indeed,  $X^+(Y) = X_{\mathcal{J}}(Y_{\mathcal{J}})$  for every  $Y \rightarrow S$ . For every  $S$ -scheme  $Y$ , let  $\tilde{Y}_{\mathcal{J}}$  denote the affine  $Y_{\mathcal{J}}$ -scheme corresponding to the quasi-coherent  $\mathcal{O}_{Y_{\mathcal{J}}}$ -algebra

$$\mathcal{O}_{Y_{\mathcal{J}}} \oplus \mathcal{J} \mathcal{O}_Y,$$

that is, the graded algebra associated with the filtration  $\mathcal{O}_Y \supset \mathcal{J} \mathcal{O}_Y$ . Then  $L_X(Y)$  identifies with  $X_{\mathcal{J}}(\tilde{Y}_{\mathcal{J}})$ , and  $L'_X(Y)$  identifies with the kernel of the morphism

$$p : X_{\mathcal{J}}(\tilde{Y}_{\mathcal{J}}) \longrightarrow X_{\mathcal{J}}(Y_{\mathcal{J}})$$

induced by the zero section  $Y_{\mathcal{J}} \rightarrow \tilde{Y}_{\mathcal{J}}$ . Equivalently, that section is induced by the morphism of  $\mathcal{O}_{Y_{\mathcal{J}}}$ -algebras

$$\mathcal{O}_{\tilde{Y}_{\mathcal{J}}} \longrightarrow \mathcal{O}_{Y_{\mathcal{J}}}$$

that vanishes on the ideal  $\mathcal{J} \mathcal{O}_Y$ . Thus, for every  $Y \rightarrow S$ , there is a split exact sequence of monoids, functorial in  $Y$ :

$$1 \longrightarrow L'_X(Y) \xrightarrow{i} L_X(Y) \xrightleftharpoons[s]{p} X^+(Y) \longrightarrow 1.$$

Figure 81: The pointwise split exact sequence for  $L_X(Y)$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 17;  
combined-reader p. 127; printed p. 117.*

It remains to show that the monoid law induced on  $L'_X$  coincides with its abelian-group structure. Let  $\mu$  be the monoid law of  $L_X$ , and let  $e$  be its unit section. We must prove that, for every  $m, m' \in L'_X(Y)$ ,

$$\mu(m \cdot e, m' \cdot e) = (m + m') \cdot e.$$

This may be seen in either of two ways. First, repeat the proof of (0.1.10 (\*)), replacing the morphism  $f : X \rightarrow W$  there by

$$\mu : L_X \times_S L_X \longrightarrow L_X.$$

Identify  $X^+(Y) = X_{\mathcal{J}}(Y_{\mathcal{J}})$  with its image under  $s$  in  $L_X(Y) = X_{\mathcal{J}}(\tilde{Y}_{\mathcal{J}})$ . For all  $g, g' \in X_{\mathcal{J}}(Y_{\mathcal{J}})$  and  $m, m' \in L'_X(Y)$ , one then obtains

$$\mu(m \cdot g, m' \cdot g') = L_{\mu}^{(g, g')}(m, m') \cdot \mu(g, g'). \quad (\star)$$

Here  $L_{\mu}^{(g, g')}$  denotes the morphism derived from  $\mu$  at the point  $(g, g')$ ; in other words,  $\tilde{Y}_{\mathcal{J}}$  lies over  $L_X \times_S L_X$  through  $(g, g')$ . In particular,

$$\mu(m \cdot e, m' \cdot e) = L_{\mu}^{(e, e)}(m, m') \cdot e.$$

But

$$L_{\mu}^{(e, e)}(m, m') = L_{\ell_e}(m') + L_{r_e}(m),$$

where  $\ell_e$ , respectively  $r_e$ , denotes left, respectively right, translation by  $e$ . Each is the identity map of  $X_{\mathcal{J}}$ , and therefore  $L_{\mu}^{(e,e)}(m, m') = m + m'$ .

Alternatively, one may argue as follows (cf. the proof of [DG70], §II.4, Theorem 3.5). By Lemma 0.1.11, formation of  $X^+$  and of  $L_X$  commutes with products, and hence so does formation of  $L'_X$ . It follows that the morphism

$$\mu' : L'_X \times L'_X \longrightarrow L'_X$$

induced by  $\mu$  is a homomorphism for the abelian-group structure. Lemma 3.10 of Exposé II then shows that  $\mu'$  coincides with the abelian-group law.

**0.8.**<sup>27</sup> Now let  $X$  be an  $S$ -scheme such that  $X_{\mathcal{J}}$  is an  $S_{\mathcal{J}}$ -group. Suppose that there is an  $S$ -morphism

$$P : X \times_S X \longrightarrow X$$

whose base change

$$P_{\mathcal{J}} : X_{\mathcal{J}} \times_{S_{\mathcal{J}}} X_{\mathcal{J}} \longrightarrow X_{\mathcal{J}}$$

is the group law of  $X_{\mathcal{J}}$ . An important special case is that in which  $X$  is an  $S$ -group and  $P$  is its group law. We obtain a morphism

$$L_P : L_X \times_S L_X \simeq L_{X \times_S X} \longrightarrow L_X.$$

In fact, this morphism does not depend on  $P$ , since it can be computed from the group law  $P_{\mathcal{J}}$  of  $X_{\mathcal{J}}$ , as we now show.<sup>28</sup>

By Proposition 0.2(ii) and (iii), for every  $Y \rightarrow S$ , every  $x, x' \in X(Y)$ , and every  $m, m' \in L'_X(Y)$ ,

$$\begin{aligned} P(m \cdot x, m' \cdot x') &= P((m, m') \cdot (x, x')) \\ &= L_P^{(x, x')}(m, m') \cdot \mu(g, g'), \end{aligned}$$

where  $g$ , respectively  $g'$ , is the image of  $x$ , respectively  $x'$ , in  $X^+(Y)$ . Moreover (cf. the proof of 0.10),  $L_P^{(x, x')} = L_{\mu}^{(g, g')}$ , and by 0.7 bis  $(\star)$  this is the element of  $L'_X(Y)$  defined by the equality

$$L_{\mu}^{(g, g')}(m, m') \cdot \mu(g, g') = \mu(m \cdot g, m' \cdot g')$$

in  $L_X(Y)$ .

Write  $\times$ , rather than  $\mu$ , for the group law of  $L_X$ , and write  $\text{Ad}$  for the *adjoint action* of  $X^+$  on  $L'_X$ . This action factors through  $X_0$  and is induced by the adjoint action of  $X_0$  on  $\omega_{X_0/S_0}^1$ . Then

$$\begin{aligned} L_{\mu}^{(g, g')}(m, m') \times g \times g' &= m \times g \times m' \times g' \\ &= (m \times \text{Ad}(g)(m')) \times g \times g', \end{aligned}$$

and hence

$$L_P^{(x, x')}(m, m') = m \times \text{Ad}(g)(m').$$

We have therefore proved:

**Proposition 0.8.** *Let  $P : X \times_S X \rightarrow X$  be an  $S$ -morphism such that  $P_{\mathcal{J}}$  equips  $X_{\mathcal{J}}$  with an  $S_{\mathcal{J}}$ -group structure. Let  $\times$  denote the group law of  $L'_X$ , let  $(m, x) \mapsto m \cdot x$  denote the morphism  $L'_X \times_S X \rightarrow X$  defining the action of  $L'_X$  on  $X$ , and let*

$$\text{Ad} : X^+ \longrightarrow \text{Aut}_{S\text{-gr}}(L'_X)$$

<sup>27</sup>Editor's note: the number 0.8 has been inserted here to mark the return to the original text.

<sup>28</sup>Editor's note: the original has been expanded in what follows.



be the adjoint action of  $X^+$  on  $L'_X$ , induced by the adjoint action of  $X_0$  on  $\omega_{X_0/S_0}^1$ . Then, for every  $S' \rightarrow S$ , every  $x, x' \in X(S')$ , and every  $m, m' \in L'_X(S')$ ,

$$P(m \cdot x, m' \cdot x') = (m \times \text{Ad}(p_X(x))(m')) \cdot P(x, x'). \quad (0.8.1)$$

If  $X$  is an  $S$ -group, we shall write  $*$  for its law,  $e$  for its unit section, and  $i$  for the  $S$ -morphism defined by

$$i(m) = m \cdot e$$

for every  $S' \rightarrow S$  and every  $m \in L'_X(S')$ .

**Corollary 0.9.** *Let  $X$  be an  $S$ -group. Then  $X^+$  is naturally an  $S$ -group, and  $p_X$  is a morphism of  $S$ -groups. Moreover, the  $S$ -morphism*

$$i : L'_X \longrightarrow X, \quad m \longmapsto m \cdot e,$$

*is an isomorphism of  $S$ -groups from  $L'_X$  onto  $\text{Ker}(p_X)$ , and, for every  $S' \rightarrow S$ , every  $x' \in X(S')$ , and every  $m \in L'_X(S')$ ,*

$$m \cdot x' = (m \cdot e) * x' = i(m) * x'. \quad (0.9.1)$$

The first two assertions were already proved in 0.1.2. Since  $X$  is formally principal homogeneous over  $X^+$  under  $L_X = L'_X \times_S X^+$ , the morphism  $i$  is indeed an isomorphism of  $S$ -functors from  $L'_X$  onto the kernel of  $p_X$ . The fact that  $i$  is a group morphism and formula (0.9.1) follow from (0.8.1), applied respectively with  $x = x' = e$ , and with  $x = e$ ,  $m' = 1$ .

**Corollary 0.10.** *Let  $X$  be an  $S$ -group. With the preceding notation, for every  $S' \rightarrow S$ , every  $x \in X(S')$ , and every  $m' \in L'_X(S')$ ,*

$$x * i(m') * x^{-1} = i(\text{Ad}(p_X(x))(m')). \quad (0.10.1)$$

This follows from the equality  $i(m') * x^{-1} = m' \cdot x^{-1}$  and from (0.8.1), applied with  $m = 1$  and  $x' = x^{-1}$ .

Thus, when  $X$  is an  $S$ -group, we have explicitly determined the kernel of  $X \rightarrow X^+$  and the action of the inner automorphisms of  $X$  on this kernel. We shall now see that the same can be done for certain, not necessarily representable,  $S$ -group functors. One case will be useful to us, namely the functors  $\underline{\text{Aut}}$  of Exposé I, 1.7. We state it at once in the next proposition.

**Proposition 0.11.** *Let  $E$  be an  $S$ -scheme, and put  $X = \underline{\text{Aut}}_S(E)$ . The kernel of the morphism of  $S$ -group functors*

$$p_X : X \longrightarrow X^+$$

*is canonically identified with the commutative  $S$ -group functor  $L'_X$  defined by*

$$\text{Hom}_S(Y, L'_X) = \text{Hom}_{\mathcal{O}_{E_0 \times_{S_0} Y_0}} \left( \Omega_{E_0/S_0}^1 \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Y_0}, \mathcal{I} \mathcal{O}_{E \times_S Y} \right),$$

where  $Y$  denotes a variable  $S$ -scheme.

Indeed, for a variable  $S$ -scheme  $Y$ , we have

$$\text{Hom}_S(Y, X) = \text{Aut}_Y(E \times_S Y)$$

and

$$\begin{aligned} \text{Hom}_S(Y, X^+) &= \text{Hom}_S(Y_{\mathcal{I}}, X) \\ &= \text{Aut}_{Y_{\mathcal{I}}}(E \times_S Y_{\mathcal{I}}) \\ &= \text{Aut}_{Y_{\mathcal{I}}}((E \times_S Y) \times_Y Y_{\mathcal{I}}). \end{aligned}$$

Applying 0.7 (b) to the  $Y$ -scheme  $E \times_S Y$ , we obtain an isomorphism of groups

$$\mathrm{Hom}_S(Y, L'_X) \simeq \mathrm{Ker}(\mathrm{Hom}_S(Y, X) \longrightarrow \mathrm{Hom}_S(Y, X^+)).$$

This isomorphism is readily checked to be functorial in the  $S$ -scheme  $Y$ . Hence it yields an isomorphism of  $S$ -groups

$$L'_X \simeq \mathrm{Ker}(X \longrightarrow X^+),$$

which completes the proof of Proposition 0.11.

**Corollary 0.12.**<sup>29</sup> *Retain the notation of 0.11:  $E$  is an  $S$ -scheme and  $X = \underline{\mathrm{Aut}}_S(E)$ . There is a natural action  $f$  of  $X$  on  $L'_X$ , defined as follows. For every  $S$ -scheme  $Y$ ,*

$$\mathrm{Hom}_S(Y, X) = \mathrm{Aut}_Y(E \times_S Y)$$

and

$$\mathrm{Hom}_S(Y, L'_X) = \mathrm{Hom}_{\mathcal{O}_{E_0 \times_{S_0} Y_0}} \left( \Omega_{E_0 \times_{S_0} Y_0 / Y_0}^1, \mathcal{I} \mathcal{O}_{E \times_S Y} \right).$$

Here

$$\Omega_{E_0/S_0}^1 \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Y_0} \simeq \Omega_{E_0 \times_{S_0} Y_0 / Y_0}^1$$

(see EGA IV<sub>4</sub>, 16.4.5). The first group acts on the second by transport of structure, and this action is functorial in  $Y$ . One then has

$$x i(m) x^{-1} = i(f(x)m), \quad (0.12.1)$$

for every  $Y \rightarrow S$ ,  $x \in \mathrm{Hom}_S(Y, X)$ , and  $m \in \mathrm{Hom}_S(Y, L'_X)$ .

Indeed, this follows from 0.7 (c), applied to the  $Y$ -scheme  $E \times_S Y$ .

**Reminder 0.13.** The direct image of a quasi-coherent module under a morphism of finite presentation is quasi-coherent. Under the same hypotheses, formation of the direct image commutes with flat base change. In the situation

$$\begin{array}{ccc} T & \xleftarrow{g'} & T' = T \times_S S' \\ f \downarrow & & \downarrow f' \\ S & \xleftarrow{g} & S' \end{array}$$

Figure 82: The flat base-change square in Reminder 0.13.

*Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 20;  
combined-reader p. 130; printed p. 120.*

if  $f$  (and hence  $f'$ ) is of finite presentation and  $g$  (and hence  $g'$ ) is flat, then for every quasi-coherent  $\mathcal{O}_T$ -module  $\mathcal{F}$ ,

$$f_*(\mathcal{F}) \otimes_{\mathcal{O}_S} \mathcal{O}_{S'} = f'_*(\mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{O}_{S'}),$$

or, more elegantly,

$$g^*(f_*\mathcal{F}) = f'_*(g'^*\mathcal{F}).$$

<sup>29</sup>Editor's note: "Remark" has been changed to "Corollary".

These two facts hold more generally for a quasi-compact and quasi-separated morphism  $f$ ; see EGA I, 9.2.1, and EGA III<sub>1</sub>, 1.4.15 in the quasi-compact separated case (taking EGA III<sub>2</sub>, Err<sub>III</sub> 25 into account), as well as EGA IV<sub>1</sub>, 1.7.4 and 1.7.21.

**Remark 0.14.**<sup>30</sup> Return to the notation of 0.11: let  $E$  be an  $S$ -scheme, put  $X = \underline{\text{Aut}}_S(E)$ , and let  $L'_X$  be the commutative  $S$ -group functor defined by

$$\begin{aligned} L'_X(Y) &= \text{Hom}_{\mathcal{O}_{E_0 \times_{S_0} Y_0}} \left( \Omega_{E_0/S_0}^1 \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Y_0}, \mathcal{I} \mathcal{O}_{E \times_S Y} \right) \\ &= \text{Hom}_{\mathcal{O}_{E \times_S Y}} \left( \Omega_{E/S}^1 \otimes_{\mathcal{O}_S} \mathcal{O}_Y, \mathcal{I} \mathcal{O}_{E \times_S Y} \right) \\ &= \Gamma \left( E \times_S Y, \widetilde{\text{Hom}}_{\mathcal{O}_{E \times_S Y}} \left( \Omega_{E/S}^1 \otimes_{\mathcal{O}_S} \mathcal{O}_Y, \mathcal{I} \mathcal{O}_{E \times_S Y} \right) \right). \end{aligned}$$

Suppose that  $Y$  is flat over  $S$ . There are then isomorphisms

$$\mathcal{I} \mathcal{O}_{E \times_S Y} \xleftarrow{\sim} (\mathcal{I} \mathcal{O}_E) \otimes_{\mathcal{O}_S} \mathcal{O}_Y \xrightarrow{\sim} (\mathcal{I} \mathcal{O}_E) \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Y_0}.$$

Suppose in addition that  $E$  is of finite presentation over  $S$ . Then  $\Omega_{E/S}^1$  is an  $\mathcal{O}_E$ -module of finite presentation (EGA IV<sub>4</sub>, 16.4.22), and therefore, by EGA 0<sub>I</sub>, 6.7.6,

$$\begin{aligned} \widetilde{\text{Hom}}_{\mathcal{O}_{E \times_S Y}} \left( \Omega_{E/S}^1 \otimes_{\mathcal{O}_S} \mathcal{O}_Y, (\mathcal{I} \mathcal{O}_E) \otimes_{\mathcal{O}_S} \mathcal{O}_Y \right) \\ \simeq \widetilde{\text{Hom}}_{\mathcal{O}_E} \left( \Omega_{E/S}^1, \mathcal{I} \mathcal{O}_E \right) \otimes_{\mathcal{O}_S} \mathcal{O}_Y. \end{aligned}$$

Let  $\pi : E \rightarrow S$  and  $g : Y \rightarrow S$  be the structural morphisms. Apply 0.13 to the diagram

$$\begin{array}{ccc} E & \xleftarrow{g'} & E \times_S Y \\ \pi \downarrow & & \downarrow \pi' \\ S & \xleftarrow{g} & Y \end{array}$$

Figure 83: The structural base-change square used in Remark 0.14.

*Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 21;  
combined-reader p. 131; printed p. 121.*

and to the  $\mathcal{O}_E$ -module

$$\mathcal{F} = \widetilde{\text{Hom}}_{\mathcal{O}_E} \left( \Omega_{E/S}^1, \mathcal{I} \mathcal{O}_E \right).$$

We obtain

$$\begin{aligned} \Gamma(E \times_S Y, g'^* \mathcal{F}) &= \Gamma(Y, \pi'_* g'^* \mathcal{F}) \\ &= \Gamma(Y, g^* \pi_* \mathcal{F}) \\ &= W(\pi_* \mathcal{F})(Y). \end{aligned}$$

Thus, if  $E$  is of finite presentation over  $S$ , we have

$$L'_X = W \left( \pi_* \widetilde{\text{Hom}}_{\mathcal{O}_E} (\Omega_{E/S}^1, \mathcal{I} \mathcal{O}_E) \right) \quad (0.14.1)$$

<sup>30</sup>Editor's note: The first part of this remark has been expanded, and a second part has been added.

on the category of  $S$ -schemes flat over  $S$ . Notice also that the module to which  $W$  is applied is quasi-coherent, by EGA I, 9.1.1 and 9.2.1.

Let  $L_0$  be the  $S_0$ -functor<sup>31</sup>

$$L_0 = W\left(\pi_{0*}\widetilde{\mathrm{Hom}}_{\mathcal{O}_{E_0}}(\Omega_{E_0/S_0}^1, \mathcal{I}\mathcal{O}_E)\right).$$

Returning to the definition of  $L'_X(Y)$ , and using the isomorphism

$$\mathcal{I}\mathcal{O}_{E \times_S Y} \simeq (\mathcal{I}\mathcal{O}_E) \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Y_0},$$

the same reasoning gives

$$L'_X(Y) = L_0(Y_0) = L_0(Y \times_S S_0) = \left(\prod_{S_0/S} L_0\right)(Y).$$

Consequently, on the category of  $S$ -schemes flat over  $S$ ,

$$L'_X = \prod_{S_0/S} W\left(\pi_{0*}\widetilde{\mathrm{Hom}}_{\mathcal{O}_{E_0}}(\Omega_{E_0/S_0}^1, \mathcal{I}\mathcal{O}_E)\right).$$

It is not obvious that the action of  $X$  on  $L'_X$  defined in 0.12 comes from an action of  $X_0 = \underline{\mathrm{Aut}}_{S_0}(E_0)$  on  $L_0$ . This is nevertheless the case when  $E$  is also flat over  $S$ .

Indeed, in that case there are canonical isomorphisms

$$\mathcal{I}\mathcal{O}_E \simeq \mathcal{I} \otimes_{\mathcal{O}_S} \mathcal{O}_E \simeq \mathcal{I} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{E_0},$$

and hence

$$L_0 \simeq W\left(\pi_{0*}\widetilde{\mathrm{Hom}}_{\mathcal{O}_{E_0}}\left(\Omega_{E_0/S_0}^1, \mathcal{I} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{E_0}\right)\right),$$

and

$$L'_X = \prod_{S_0/S} W\left(\pi_{0*}\widetilde{\mathrm{Hom}}_{\mathcal{O}_{E_0}}\left(\Omega_{E_0/S_0}^1, \mathcal{I} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{E_0}\right)\right). \quad (0.14.2)$$

For every  $S_0$ -scheme  $T$ , we then have

$$L_0(T) \simeq \mathrm{Hom}_{\mathcal{O}_{E_0 \times_{S_0} T}}\left(\Omega_{E_0 \times_{S_0} T/T}^1, \mathcal{I} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{E_0 \times_{S_0} T}\right).$$

Moreover,

$$\mathrm{Hom}_{S_0}(T, X_0) = \mathrm{Aut}_T(E_0 \times_{S_0} T)$$

acts on  $L_0(T)$  by transport of structure, functorially in  $T$ . Finally, for every  $S$ -scheme  $Y$  flat over  $S$ , the action by transport of structure of

$$\mathrm{Hom}_S(Y, X) = \mathrm{Aut}_Y(E \times_S Y)$$

on  $L'_X(Y) = L_0(Y_0)$  factors through  $\mathrm{Aut}_{Y_0}(E_0 \times_{S_0} Y_0)$ .

Finally, let us extract the following two propositions from SGA 1, Exposé III.

**Proposition 0.15.** (SGA 1, Exposé III, 6.8)<sup>32</sup> *For every affine  $S_{\mathcal{J}}$ -scheme  $Y$  smooth over  $S_{\mathcal{J}}$ , there exists an  $S$ -scheme  $X$ , smooth over  $S$ , such that  $X \times_S S_{\mathcal{J}} \simeq Y$ ; such an  $X$  is unique up to a non-unique isomorphism.*

<sup>31</sup>Editor's note: What follows has been added.

<sup>32</sup>Editor's note: This "global" lifting result uses the "local" lifting result of loc. cit., 4.1, which is stated for  $S$  locally Noetherian; see EGA IV<sub>4</sub>, 18.1.1 for the general case.

**Proposition 0.16.** (SGA 1, Exposé III, 5.5)<sup>33</sup> Let  $X$  be an  $S$ -scheme smooth over  $S$ . For every affine  $S$ -scheme  $Y$ , the canonical map

$$p_X(Y) : \mathrm{Hom}_S(Y, X) \longrightarrow \mathrm{Hom}_S(Y, X^+) = \mathrm{Hom}_{S_{\mathcal{J}}}(Y_{\mathcal{J}}, X_{\mathcal{J}})$$

is surjective.

**Corollary 0.17.** Let  $E$  be an affine  $S$ -scheme smooth over  $S$ , and put  $X = \underline{\mathrm{Aut}}_S(E)$ . For every affine  $S$ -scheme  $Y$ , the canonical map

$$\begin{aligned} \mathrm{Aut}_Y(E \times_S Y) &= \mathrm{Hom}_S(Y, X) \longrightarrow \mathrm{Hom}_S(Y, X^+) \\ &= \mathrm{Aut}_{Y_{\mathcal{J}}}(E_{\mathcal{J}} \times_{S_{\mathcal{J}}} Y_{\mathcal{J}}) \end{aligned}$$

is surjective.

Indeed,  $Y \times_S E$  is affine over the affine scheme  $Y$ , hence is affine. Applying 0.16, we see that every  $S_{\mathcal{J}}$ -morphism

$$Y_{\mathcal{J}} \times_{S_{\mathcal{J}}} E_{\mathcal{J}} \longrightarrow E_{\mathcal{J}}$$

extends to an  $S$ -morphism

$$Y \times_S E \longrightarrow E.$$

In other words,<sup>34</sup> every  $Y_{\mathcal{J}}$ -endomorphism of  $Y_{\mathcal{J}} \times_{S_{\mathcal{J}}} E_{\mathcal{J}}$  lifts to a  $Y$ -endomorphism of  $Y \times_S E$ . Then 0.7 (a) shows that every  $Y_{\mathcal{J}}$ -automorphism of  $Y_{\mathcal{J}} \times_{S_{\mathcal{J}}} E_{\mathcal{J}}$  lifts to a  $Y$ -automorphism of  $Y \times_S E$ , which is the asserted property.

## 1. Extensions and Cohomology

### 1.1.

Let  $\mathcal{C}$  be a category stable under fiber products.<sup>35</sup> Let  $S$  be an object of  $\mathcal{C}$ , let  $G$  be a representable  $S$ -group, and let  $F$  be an  $S$ -functor in commutative groups on which  $G$  acts. The cohomology groups  $H^n(G, F)$  were defined in I 5.1. Recall that they are the cohomology groups of a complex denoted  $C^*(G, F)$ . Writing

$$(G/S)^n = G \times_S \cdots \times_S G \quad (n \text{ factors}),$$

one has

$$C^n(G, F) = \mathrm{Hom}_S((G/S)^n, F).$$

Since  $G$ , and hence each  $(G/S)^n$ , is representable, one also has

$$C^n(G, F) = F((G/S)^n).$$

It follows from this and from the definition of the coboundary operator that the complex  $C^*(G, F)$  depends only on the restriction of  $F$  to the full subcategory of  $\mathcal{C}/S$  whose objects are the Cartesian powers of  $G$  over  $S$ . Consequently:

**Lemma 1.1.1.** Let  $\mathcal{C}$  be a category stable under fiber products, let  $S$  be an object of  $\mathcal{C}$ , and let  $G$  be a representable  $S$ -group. Denote by  $\mathcal{C}(G)$  the full subcategory of  $\mathcal{C}/S$  whose objects

<sup>33</sup>Editor's note: This is an immediate consequence of the definition of “(formally) smooth” adopted in EGA IV<sub>4</sub>, 17.3.1 (and 17.1.1).

<sup>34</sup>Editor's note: The original has been modified slightly so that the hypotheses of 0.7 apply exactly.

<sup>35</sup>Editor's note: the hypothesis that  $\mathcal{C}$  is stable under fiber products has been added. It holds in the applications in which  $\mathcal{C}$  is the category of schemes (**Sch**), or the category of functors (**Sch**)<sup>o</sup>  $\rightarrow$  (**Ens**). If this hypothesis is omitted, one must instead assume below that the fiber products  $G \times_S \cdots \times_S G$  are representable.

are the Cartesian powers of  $G$  over  $S$ . Let  $F$  and  $F'$  be two  $S$ -functors in commutative groups on which  $G$  acts. If  $F$  and  $F'$  have the same restriction to  $\mathcal{C}(G)$ , there is a canonical isomorphism

$$H^*(G, F) \xrightarrow{\sim} H^*(G, F').$$

### 1.1.2. Cohomology and restriction of scalars.<sup>36</sup>

We record another comparison result. Let  $T \rightarrow S$  be a morphism in  $\mathcal{C}$ . If  $F$  is a  $T$ -functor in commutative groups, then the functor obtained by restriction of scalars (cf. Exposé II, 1),

$$F_1 = \prod_{T/S} F,$$

is an  $S$ -functor in commutative groups, and there is a morphism of  $S$ -functors in groups<sup>37</sup>

$$u : \prod_{T/S} \underline{\text{Aut}}_{T\text{-gr.}}(F) \longrightarrow \underline{\text{Aut}}_{S\text{-gr.}}(F_1).$$

Now let  $G$  be an  $S$ -functor in groups, and suppose an action of  $G_T$  on  $F$  is given by

$$G_T \longrightarrow \underline{\text{Aut}}_{T\text{-gr.}}(F).$$

By the definition of  $\prod_{T/S}$ , this gives a morphism of  $S$ -functors in groups

$$G \longrightarrow \prod_{T/S} \underline{\text{Aut}}_{T\text{-gr.}}(F).$$

Composing with  $u$  therefore gives an action of  $G$  on  $F_1 = \prod_{T/S} F$ .<sup>38</sup>

**Lemma 1.1.2.** *Under the preceding assumptions, there is a canonical isomorphism*

$$H^*\left(G, \prod_{T/S} F\right) \simeq H^*(G_T, F).$$

Indeed, by the definition of cohomology, the standard complexes are canonically isomorphic.

<sup>36</sup>Editor's note: the heading of this paragraph has been added.

<sup>37</sup>Editor's note: explicitly, let  $S' \rightarrow S$  and  $\alpha \in \underline{\text{Aut}}_{(T \times_S S')\text{-gr.}}(F_{T \times_S S'})$ . For every  $U \rightarrow T \times_S S'$ , let  $\alpha_U \in \text{Aut}_{\text{gr.}}(F_{T \times_S S'}(U))$  be the element defined by  $\alpha$ . Then  $u(S')(\alpha)$  is the element  $\beta$  of

$$\underline{\text{Aut}}_{S\text{-gr.}}(F_1)(S') = \underline{\text{Aut}}_{S'\text{-gr.}}\left(\prod_{T \times_S S'/S'} F_{T \times_S S'}\right)$$

such that  $\beta_{S''} = \alpha_{T \times_S S''}$  for every  $S'' \rightarrow S'$ .

<sup>38</sup>Editor's note: explicitly, for every  $S' \rightarrow S$ , the action of  $G(S')$  on  $F_1(S') = F(T \times_S S')$  is given by the action of  $G_T(T \times_S S') = G(T \times_S S')$  on  $F(T \times_S S')$ , together with the group morphism  $G(S') \rightarrow G(T \times_S S')$  corresponding to the projection  $T \times_S S' \rightarrow S'$ . Equivalently, the action of  $G_T$  on  $F$  is a morphism  $G_T \times_T F \rightarrow F$ . Applying  $\prod_{T/S}$ , and writing  $G_1 = \prod_{T/S} G_T$ , gives, by II 1.2, a morphism  $G_1 \times_S F_1 \rightarrow F_1$ , hence an action of  $G_1$  on  $F_1$ . The action of  $G$  is obtained through the natural morphism  $G \rightarrow G_1$ , which corresponds by adjunction to  $\text{id}_G$ .

**1.2. Lifting group morphisms.**<sup>39</sup>

Following the general principles, we make the following definition.

**Definition 1.2.1.** *Let*

$$1 \longrightarrow M \xrightarrow{u} E \xrightarrow{v} G$$

*be a sequence of morphisms of  $\widehat{\mathcal{C}}$ -groups. It is called exact if the following equivalent conditions hold:*

(i) *for every  $S \in \text{Ob } \mathcal{C}$ , the following sequence of ordinary groups is exact:*

$$1 \longrightarrow M(S) \xrightarrow{u(S)} E(S) \xrightarrow{v(S)} G(S);$$

(ii) *for every object  $H$  of  $\widehat{\mathcal{C}}$ , the following sequence of ordinary groups is exact:*

$$1 \longrightarrow \text{Hom}(H, M) \xrightarrow{u(H)} \text{Hom}(H, E) \xrightarrow{v(H)} \text{Hom}(H, G).$$

Taking  $H = G$  in (ii), one sees that the set of sections of  $v$  (which need not preserve the group structures) is either empty or principal homogeneous under  $\text{Hom}(G, M)$ . Suppose that it is nonempty, and let

$$s : G \longrightarrow E$$

be a section of  $v$ . For every  $S \in \text{Ob } \mathcal{C}$  and every  $x \in G(S)$ , the element  $s(x) \in E(S)$  defines an inner automorphism of  $E_S$  that normalizes  $M_S$  (more precisely, the image of  $M_S$  under  $u_S$ ), and hence an automorphism of  $M_S$ .

**Scholium 1.2.1.1.**<sup>40</sup> *If  $M$  is commutative, one sees set-theoretically that this automorphism is independent of the chosen section and depends only on  $x$ , and that it depends multiplicatively on  $x$ . In summary, every exact sequence*

$$(E) \quad 1 \longrightarrow M \xrightarrow{u} E \xrightarrow{v} G$$

*in which  $M$  is commutative and  $v$  admits a section determines a morphism of  $\widehat{\mathcal{C}}$ -groups*

$$G \longrightarrow \underline{\text{Aut}}_{\widehat{\mathcal{C}}\text{-gr.}}(M),$$

*called the action of  $G$  on  $M$  defined by the extension  $(E)$ .*

**Definition 1.2.1.2.** We saw in I 2.3.7 that  $v$  admits a section that is a morphism of  $\widehat{\mathcal{C}}$ -groups if and only if the extension  $(E)$  is isomorphic, as an extension, to the semidirect product of  $M$  by  $G$  for the preceding action. Such a section of  $v$  is called a *section of the extension  $(E)$* , or simply a *section of  $(E)$* .

If  $s$  is a section of  $(E)$  and

$$m \in \Gamma(M) \simeq \text{Ker}(\Gamma(E) \rightarrow \Gamma(G))$$

(for the definition of  $\Gamma$ , see I 1.2), then the morphism  $G \rightarrow E$  defined by

$$x \longmapsto u(m)s(x)u(m)^{-1}$$

<sup>39</sup>Editor's note: this heading has been added.

<sup>40</sup>Editor's note: the numbering 1.2.1.1 and 1.2.1.2 has been added to highlight the notions introduced here.

is also a section of  $(E)$ .<sup>41</sup> It is called the section deduced from  $s$  by the inner automorphism defined by  $m$ , or by  $u(m)$ .

**Lemma 1.2.2.** *Let*

$$(E) \quad 1 \longrightarrow M \xrightarrow{u} E \xrightarrow{v} G$$

*be an exact sequence of  $\widehat{\mathcal{C}}$ -groups in which  $M$  is commutative and  $v$  admits a section. Let  $G$  act on  $M$  in the manner defined by  $(E)$ .*

- (i) *The extension  $(E)$  canonically defines a class  $c(E) \in H^2(G, M)$ , whose vanishing is necessary and sufficient for the existence of a section of  $(E)$ .*
- (ii) *If  $c(E) = 0$ , the set of sections of  $(E)$  is principal homogeneous under  $Z^1(G, M)$ , and the set of sections of  $(E)$ , modulo the action of the inner automorphisms defined by the elements of  $\Gamma(M)$ , is principal homogeneous under  $H^1(G, M)$ .*
- (iii) <sup>42</sup> *Let  $s$  be a section of  $(E)$ . The set of conjugates of  $s$  by the inner automorphisms defined by  $\Gamma(M)$  is in bijection with  $\Gamma(M)/H^0(G, M)$ .*

The proof is exactly the same as for ordinary groups; starting from a section of  $v$  ensures the functoriality of the set-theoretic computations. We briefly indicate the principal steps.

(a) To every section  $s$  of  $v$ , associate the morphism

$$D_s : G \times G \longrightarrow M$$

defined set-theoretically by

$$u(D_s(x, y)) = s(xy)s(y)^{-1}s(x)^{-1}.$$

The following computation shows that  $D_s$  is a 2-cocycle.<sup>43</sup> By the definition of the differential in the standard complex (I 5.1),

$$(\partial^2 D_s)(x, y, z) = (s(x)D_s(y, z)s(x)^{-1}) \cdot D_s(x, y)^{-1} \cdot D_s(xy, z)^{-1} \cdot D_s(x, yz).$$

Substituting the definition of  $D_s$  into this formula gives, without using commutativity,

$$D_s \in Z^2(G, M).$$

(b) If  $s$  and  $s'$  are two sections of  $v$ , there is a morphism  $f : G \rightarrow M$  such that  $s(x) = f(x)s'(x)$ . Then

$$\begin{aligned} D_{s'}(x, y) &= f^{-1}(xy)D_s(x, y)s(x)f(y)s(x)^{-1}f(x) \\ &= D_s(x, y) \cdot f^{-1}(xy) \cdot (s(x)f(y)s(x)^{-1}) \cdot f(x), \end{aligned}$$

and therefore

$$D_{s'} = D_s \cdot \partial^1 f.$$

This shows that the class of  $D_s$  in  $H^2(G, M)$  is independent of the chosen section  $s$  of  $v$ ; it is the class  $c(E)$  of the extension  $(E)$ .<sup>44</sup>

<sup>41</sup>Editor's note: in the right-hand side,  $x$  has been corrected to  $s(x)$ .

<sup>42</sup>Editor's note: this item has been added as the counterpart of 1.2.4(iii). The assertion is valid for every section of  $v$ ; see the proof.

<sup>43</sup>Editor's note: the original has been corrected to make it compatible with I 5.1.

<sup>44</sup>Editor's note: the preceding sentence has been added.



(c) Let  $s$  and  $s'$  be two sections of  $v$ , and let  $m \in \Gamma(M)$ . The equality

$$s(x) = m^{-1}s'(x)m \quad (x \in G(S), S \in \text{Ob } \mathcal{C})$$

is equivalent to

$$s(x) = m^{-1}s'(x)m s'(x)^{-1}s'(x), \quad \text{that is,} \quad s = \partial^0 m \cdot s'.$$

In particular, the stabilizer of  $s$  in  $\Gamma(M)$  is the subgroup of elements  $m$  such that  $\partial^0 m = e_M$ , namely  $H^0(G, M)$ . This already proves (iii).

(d) The argument is now familiar.<sup>45</sup> Let  $s_0$  be an arbitrary section of  $v$ . There exists a section  $s$ , necessarily of the form  $s = f \cdot s_0$ , that is a group morphism, that is, satisfies  $D_s = 0$ , if and only if

$$(D_{s_0})^{-1} = \partial^1 f,$$

for some  $f : G \rightarrow M$ , or equivalently if and only if  $c(E) = 0$ . This proves (i).

In that case, the sections of  $(E)$  are precisely the sections  $s' = h \cdot s$ , where  $h : G \rightarrow M$  satisfies  $\partial^1 h = 0$ , that is,  $h \in Z^1(G, M)$ . Moreover, by (c), two such sections  $h_1 \cdot s$  and  $h_2 \cdot s$  are conjugate under  $\Gamma(M)$  if and only if  $h_1$  and  $h_2$  have the same image in  $H^1(G, M)$ . This proves (ii).

Still let

$$(E) \quad 1 \longrightarrow M \xrightarrow{u} E \xrightarrow{v} G$$

be an exact sequence of  $\widehat{\mathcal{C}}$ -groups with  $M$  commutative, and let

$$f : H \longrightarrow G$$

be a morphism of  $\widehat{\mathcal{C}}$ -groups.

Consider  $E_f = H \times_G E$ . It is a  $\widehat{\mathcal{C}}$ -group, and the projection  $v_f : E_f \rightarrow H$  is a morphism of  $\widehat{\mathcal{C}}$ -groups; the same holds for  $e_f : E_f \rightarrow E$ . Sending  $M$  into  $E$  by  $u$  and into  $H$  by the unit morphism also defines a morphism of  $\widehat{\mathcal{C}}$ -groups  $u_f : M \rightarrow E_f$ . We have therefore constructed the following commutative diagram of  $\widehat{\mathcal{C}}$ -groups.

$$\begin{array}{ccccccc} (E) & 1 & \longrightarrow & M & \xrightarrow{u} & E & \xrightarrow{v} & G \\ & & & \uparrow \text{id} & & \uparrow e_f & & \uparrow f \\ (E_f) & 1 & \longrightarrow & M & \xrightarrow{u_f} & E_f & \xrightarrow{v_f} & H \end{array}$$

Figure 84: The pullback extension  $(E_f)$  and its morphism to  $(E)$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 27;  
combined-reader p. 137; printed p. 127.*

It follows immediately that:

**Lemma 1.2.3.**

(i) *The sequence  $(E_f)$  is exact.*

---

<sup>45</sup>Editor's note: what follows has been expanded.

- (ii) The map  $s \mapsto e_f \circ s = f'$  gives a bijective correspondence between the morphisms

$$s : H \longrightarrow E_f$$

such that  $v_f \circ s = \text{id}$ , that is, the sections of  $v_f$ , and the morphisms

$$f' : H \longrightarrow E$$

such that  $v \circ f' = f$ , that is, the morphisms  $f'$  lifting  $f$ .

- (iii) Under this correspondence, the sections of  $(E_f)$  correspond to the group morphisms  $f'$  lifting  $f$ .

Applying Lemma 1.2.2 to the extension  $(E_f)$ , and taking 1.2.3 into account, gives the following proposition, which formally contains Lemma 1.2.2.

**Proposition 1.2.4.** *Let*

$$(E) \quad 1 \longrightarrow M \longrightarrow E \xrightarrow{v} G$$

be an exact sequence of  $\widehat{\mathcal{C}}$ -groups with  $M$  commutative. Let  $f : H \rightarrow G$  be a morphism of  $\widehat{\mathcal{C}}$ -groups, and suppose that it lifts to a morphism, not necessarily a group morphism,  $f' : H \rightarrow E$ . Let  $H$  act on  $M$  by the composite morphism, which is multiplicative and independent of the choice of  $f'$ ,

$$H \xrightarrow{f'} E \xrightarrow{\text{int}} \underline{\text{Aut}}_{\widehat{\mathcal{C}}\text{-gr.}}(M).$$

- (i) The morphism  $f$  canonically defines a class  $c(f) \in H^2(H, M)$ , whose vanishing is necessary and sufficient for the existence of a morphism of  $\widehat{\mathcal{C}}$ -groups

$$f' : H \longrightarrow E$$

lifting  $f$ .

- (ii) If  $c(f) = 0$ , the set of morphisms of  $\widehat{\mathcal{C}}$ -groups  $f'$  lifting  $f$ , modulo the action of the inner automorphisms defined by elements of  $\Gamma(M)$  (that is, by elements  $m \in \Gamma(E)$  such that  $v(m) = e$ ), is principal homogeneous under  $H^1(H, M)$ .
- (iii) If  $f' : H \rightarrow E$  is a group morphism lifting  $f$ , the set of transforms of  $f'$  by the inner automorphisms defined by the elements of  $\Gamma(M)$  is isomorphic to

$$\Gamma(M)/\Gamma(M^H) = \Gamma(M)/H^0(H, M).$$

### 1.3. Extensions of group laws.

Consider the following situation: one has a morphism in  $\widehat{\mathcal{C}}$ ,

$$(\dagger) \quad p : X \longrightarrow Y,$$

and a commutative  $\widehat{\mathcal{C}}$ -group  $M$  acting on  $X$ , such that  $X$  is formally principal homogeneous over  $Y$  under  $M_Y$ .

If  $g : Y \rightarrow Z$  is an arbitrary morphism in  $\widehat{\mathcal{C}}$ , then  $g \circ p : X \rightarrow Z$  is invariant under  $M$ : for every  $S \in \text{Ob } \widehat{\mathcal{C}}$ , the map  $(g \circ p)(S)$  is invariant under the action of  $M(S)$  on  $X(S)$ . Conversely, we shall assume that the following condition holds for  $n = 1, 2, 3, 4$ :

$(+)_n$ : Every morphism from  $X^n$  to  $M$  that is invariant under the action of  $M^n$  on  $X^n$  factors uniquely through  $p^n : X^n \rightarrow Y^n$ , where the superscript  $n$  denotes a Cartesian power.

**Lemma 1.3.1.**

- (i) If  $h$  is a morphism from  $Y$  to  $M$ , the automorphism  $u_h$  of  $X$ , defined set-theoretically by  $x \mapsto h(p(x)) \cdot x$ , preserves the fibers of  $p$  and commutes with the operations of  $M$  on  $X$ .<sup>46</sup> Thus, for every  $S \in \text{Ob } \mathcal{C}$ ,  $x \in X(S)$ , and  $m \in M(S)$ , one has

$$p(h(p(x)) \cdot x) = p(x), \quad m \cdot h(p(x)) \cdot x = h(p(m \cdot x)) \cdot m \cdot x.$$

- (ii) This construction gives a bijective correspondence between morphisms from  $Y$  to  $M$  and automorphisms of  $X$  that preserve the fibers of  $p$  and commute with the operations of  $M$ .

The first assertion is clear, since  $p(m \cdot x) = p(x)$  and  $M$  is commutative. Conversely, an automorphism  $u$  of  $X$  preserving the fibers of  $p$  can be written set-theoretically as  $x \mapsto g(x)x$ , where  $g$  is a certain morphism from  $X$  to  $M$ . If  $u$  commutes with the operations of  $M$ , then  $g$  is invariant under  $M$ ,<sup>47</sup> and the conclusion follows from condition  $(+)_1$ .

Suppose now, in addition, that a group law on  $Y$  and an action of  $Y$  on  $M$  are given, that is, a morphism of  $\widehat{\mathcal{C}}$ -groups

$$(\ddagger) \quad f : Y \longrightarrow \underline{\text{Aut}}_{\widehat{\mathcal{C}}\text{-gr.}}(M).$$

**Definition 1.3.2.** A composition law on  $X$ ,

$$P : X \times X \longrightarrow X,$$

is called *admissible* if it satisfies the following two conditions:

- (i)  $P$  lifts the group law of  $Y$ ; that is, the following diagram is commutative.

$$\begin{array}{ccc} X \times X & \xrightarrow{P} & X \\ (p,p) \downarrow & & \downarrow p \\ Y \times Y & \xrightarrow{\quad} & Y \end{array}$$

Figure 85: The compatibility of an admissible composition law with the group law on  $Y$ .

*Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 29;  
combined-reader p. 139; printed p. 129.*

- (ii) For every  $S \in \text{Ob } \mathcal{C}$  and all  $x, y \in X(S)$ ,  $m, n \in M(S)$ , one has

$$(++) \quad P(m \cdot x, n \cdot y) = m \cdot f(p(x))(n) \cdot P(x, y).$$

**Proposition 1.3.3.** For a group law  $*$  on  $X$  to be admissible, it is necessary and sufficient that the following four conditions hold:

- (i)  $p : X \rightarrow Y$  is a group morphism.  
(ii) The morphism  $i : M \rightarrow X$ , defined by  $i(m) = m \cdot e_X$ , is a group isomorphism from  $M$  onto  $\ker(p)$ ; in other words, set-theoretically,

$$(m \cdot e_X) * (n \cdot e_X) = (mn) \cdot e_X.$$

<sup>46</sup>Editor's note: here and below, "commutes with  $p$  and the operations of  $M$ " has been replaced by "preserves the fibers of  $p$  and commutes with the operations of  $M$ ." The equalities that follow have also been added.

<sup>47</sup>Editor's note: indeed,  $mg(x)x = mu(x) = u(mx) = g(mx)mx$ , and hence  $g(mx) = g(x)$ .

- (iii) One has  $m \cdot x = (m \cdot e_X) * x = i(m) * x$  for every  $m \in M(S)$  and  $x \in X(S)$ .
- (iv) The inner automorphisms of  $X$  act on  $\ker(p)$  according to the set-theoretic formula

$$x * i(m) * x^{-1} = i(f(p(x))m).$$

The proof is immediate.

**Lemma 1.3.4.**<sup>48</sup> Let  $h : Y \rightarrow M$  be a morphism and let  $u_h$  be the automorphism  $x \mapsto h(p(x)) \cdot x$  of  $X$  (cf. 1.3.1). Let  $P$  be an admissible composition law (respectively, an admissible group law) on  $X$ , and let  $P'$  be the composition law on  $X$  obtained from  $P$  by means of  $u_h$ , that is,

$$P'(x, y) = u_h^{-1}(P(u_h(x), u_h(y))).$$

Then:

- (i)  $P'$  is an admissible composition law (respectively, an admissible group law).
- (ii) For every  $x, y \in X(S)$ , with  $S \in \text{Ob } \mathcal{C}$ , set  $v = p(x)$  and  $w = p(y)$ . Then

$$P'(x, y) = h(vw)^{-1} \cdot h(v) \cdot f(v)(h(w)) \cdot P(x, y) = (\partial^1 h)(p(x), p(y)) \cdot P(x, y).$$

*Proof.* One has  $u_h^{-1} = u_{h^\vee}$ , where  $h^\vee : Y \rightarrow M$  is defined by  $h^\vee(y) = h(y)^{-1}$ . By 1.3.2(i) and (ii),

$$P(h(v) \cdot x, h(w) \cdot y) = h(v) \cdot f(v)(h(w)) \cdot P(x, y), \quad p(P(x, y)) = vw,$$

and hence

$$P'(x, y) = h(vw)^{-1} \cdot h(v) \cdot f(v)(h(w)) \cdot P(x, y) = (\partial^1 h)(p(x), p(y)) \cdot P(x, y).$$

It follows immediately that  $P'$  satisfies conditions (i) and (ii) of 1.3.2.

**Definition 1.3.5.** Two admissible composition laws obtained from one another by the procedure of 1.3.4 are called equivalent.<sup>49</sup>

**Proposition 1.3.6.** Suppose that an admissible composition law on  $X$  exists. Then:

- (i) There exists a canonically determined class  $c \in H^3(Y, M)$ , whose vanishing is necessary and sufficient for the existence of an admissible associative composition law on  $X$ .
- (ii) If  $c = 0$ , the set of admissible associative composition laws on  $X$  (respectively, the set of their equivalence classes) is principal homogeneous under  $Z^2(Y, M)$  (respectively,  $H^2(Y, M)$ ).

The proof proceeds in several steps.

(a) Let  $P$  be an admissible composition law on  $X$ . Since  $P$  lifts the associative composition law of  $Y$ , there is a unique morphism  $a : X^3 \rightarrow M$  such that

$$(*) \quad P(x, P(y, z)) = a(x, y, z) \cdot P(P(x, y), z).$$

<sup>48</sup>Editor's note: the statement and its proof have been expanded with a view to step (e) in the proof of 1.3.6.

<sup>49</sup>Editor's note: this is indeed an equivalence relation, since  $u_h^{-1} = u_{h^\vee}$ .

Applying conditions 1.3.2(i) and (ii), one sees at once that  $a$  is invariant under the action of  $M^3$  on  $X^3$ .<sup>50</sup> Condition  $(+)_3$  therefore gives:

(1) *There exists a unique morphism  $DP : Y^3 \rightarrow M$  such that*

$$P(x, P(y, z)) = DP(p(x), p(y), p(z)) \cdot P(P(x, y), z),$$

and  $P$  is associative if and only if  $DP = 0$ .

(b) Let us compute  $P(P(P(x, y), z), t)$  step by step using the preceding formula. Put

$$p(x) = u, \quad p(y) = v, \quad p(z) = w, \quad p(t) = h.$$

One obtains<sup>51</sup> the following pentagonal diagram, in which an arrow  $a \xrightarrow{m} b$  means that  $b = m \cdot a$ .




Figure 86: The pentagonal comparison used to compute the associativity obstruction.

*Temporary Loop-1 image; source locator: Polo-Gille Exp3-14oct24.pdf, component-local p. 30;  
combined-reader p. 140; printed p. 130.*

Thus one finds

$$\begin{aligned}
 & DP(u, v, w) \cdot DP(u, vw, h) \cdot f(u)(DP(v, w, h)) \\
 & \cdot DP(u, v, wh)^{-1} \cdot DP(uv, w, h)^{-1} = e_M,
 \end{aligned}$$

that is,  $\partial^3 DP(u, v, w, h) = e_M$ . Moreover, the left-hand side of the preceding formula can be expressed, by means of  $P$  and  $a$ , as the value at  $(x, y, z, t)$  of a certain morphism  $X^4 \rightarrow M$ .

It follows from the uniqueness assertion in  $(+)_4$  that  $\partial^3 DP$  and  $e_M$ , which factor the same morphism, are equal. Hence:

(2)  *$DP$  is a cocycle; that is,  $DP \in Z^3(Y, M)$ .*

(c) If  $P$  and  $P'$  are two admissible composition laws on  $X$ , there is a unique morphism

$$b : X^2 \longrightarrow M$$

<sup>50</sup>Editor's note: for  $x, y, z \in X(S)$ , set  $u = p(x)$ ,  $v = p(y)$ , and  $w = p(z)$ . Then

$$\begin{aligned}
 P(mx, P(m'y, m''z)) &= P(mx, m'f(v)(m'')P(y, z)) \\
 &= mf(u)(m')f(uv)(m'')P(x, P(y, z)).
 \end{aligned}$$

On the other hand, since  $p(P(x, y)) = uv$ , one also has

$$\begin{aligned}
 P(P(mx, m'y), m''z) &= P(mf(u)(m')P(x, y), m''z) \\
 &= mf(u)(m')f(uv)(m'')P(P(x, y), z).
 \end{aligned}$$

Comparison of these equalities with  $(*)$  gives  $a(mx, m'y, m''z) = a(x, y, z)$ .

<sup>51</sup>Editor's note: the diagram that follows has been added.

such that  $P'(x, y) = b(x, y)P(x, y)$ . Applying 1.3.2(ii) to  $P$  and  $P'$ , one sees that  $b$  is invariant under  $M^2$ , and hence, by  $(+)_2$ :

(3) For every pair  $(P, P')$  of admissible composition laws there exists a unique morphism  $d(P, P') : Y^2 \rightarrow M$  such that

$$P'(x, y) = d(P, P')(p(x), p(y)) \cdot P(x, y),$$

and the set of admissible composition laws is thereby made principal homogeneous under

$$\mathrm{Hom}(Y^2, M) = C^2(Y, M).$$

(d) Under the preceding conditions one has

$$(4) \quad DP' - DP = \partial^2 d(P, P').$$

(e) The laws  $P$  and  $P'$  are equivalent if and only if there is a morphism

$$h \in C^1(Y, M) = \mathrm{Hom}(Y, M)$$

such that  $d(P, P') = \partial^1 h$ . This follows from the definition of equivalence and from 1.3.4(ii).

(f) It remains only to conclude. We seek an associative  $P'$ , that is, one for which  $DP' = e_M$ . Now  $DP$  is a cocycle whose class in  $H^3(Y, M)$  does not depend on the chosen admissible composition law  $P$ , by (3) and (4). This class is the required obstruction  $c$ . An appropriate  $P'$  can be chosen if and only if  $c = 0$ . Indeed, after choosing an arbitrary  $P$ , one must solve, by (1),

$$0 = DP' = DP + \partial^2 d(P, P'),$$

which is possible, by (3) and (4), if and only if  $c = 0$ . Again by (3) and (4), the set of associative laws  $P'$  is principal homogeneous under  $Z^2(Y, M)$ . By (e), the set of associative laws  $P'$  modulo equivalence is principal homogeneous under  $H^2(Y, M)$ .

## 2. Infinitesimal extensions of a morphism of group schemes

Retain the notation of §0. Let  $Y$  and  $X$  be two  $S$ -group functors, and let  $M$  be the kernel of the group homomorphism  $p_X : X \rightarrow X^+$ . Thus there is an exact sequence of  $S$ -group functors

$$1 \longrightarrow M \longrightarrow X \xrightarrow{p_X} X^+.$$

By the definition of  $X^+$ , there are isomorphisms

$$\begin{aligned} \mathrm{Hom}_S(Y, X^+) &\xrightarrow{\sim} \mathrm{Hom}_{S_{\mathcal{J}}}(Y_{\mathcal{J}}, X_{\mathcal{J}}), \\ \mathrm{Hom}_{S\text{-gr}}(Y, X^+) &\xrightarrow{\sim} \mathrm{Hom}_{S_{\mathcal{J}}\text{-gr}}(Y_{\mathcal{J}}, X_{\mathcal{J}}). \end{aligned}$$

The map

$$\mathrm{Hom}_S(Y, p_X) : \mathrm{Hom}_S(Y, X) \longrightarrow \mathrm{Hom}_S(Y, X^+)$$

sends an  $S$ -morphism  $f : Y \rightarrow X$  to the  $S$ -morphism  $f^+ : Y \rightarrow X^+$  corresponding under these isomorphisms to the  $S_{\mathcal{J}}$ -morphism  $f_{\mathcal{J}} : Y_{\mathcal{J}} \rightarrow X_{\mathcal{J}}$  obtained from  $f$  by base change. If  $M$  is commutative, Proposition 1.2.4 applies to this situation.

**2.0.**<sup>52</sup> In what follows we shall consider the case in which  $Y$  is flat over  $S$ , and  $X$  is an  $S$ -group functor of one of the following two types:

<sup>52</sup>Editor's note: The numbering 2.0 has been added for later references.

*a*  $X$  is an  $S$ -group scheme;

*b*  $X = \underline{\text{Aut}}_S(E)$ , where  $E$  is an  $S$ -scheme of finite presentation over  $S$ .

Write  $(\text{Flat})/S$  for the category of  $S$ -schemes flat over  $S$ . In case *a*, respectively *b*, the  $S$ -group functor  $M = \text{Ker}(X \rightarrow X^+)$ , its restriction  $L$  to  $(\text{Flat})/S$ , and the actions on  $M$  of the inner automorphisms of  $X$  were calculated in 0.9, 0.5, and 0.10, respectively in 0.11, 0.14, and 0.12.

More explicitly, in case *a*, let  $L_0$  be the  $S_0$ -functor in commutative groups defined, for every  $S_0$ -scheme  $T_0$ , by

$$\text{Hom}_{S_0}(T_0, L_0) = \text{Hom}_{\mathcal{O}_{T_0}} \left( \omega_{X_0/S_0}^1 \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{T_0}, \mathcal{J} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{T_0} \right).$$

The group  $X_0$  acts on this functor through its adjoint representation on  $\omega_{X_0/S_0}^1$ , and

$$L = \prod_{S_0/S} L_0; \quad L(T) = L_0(T \times_S S_0)$$

for every  $S$ -scheme  $T$ .

In case *b*, let  $\pi : E \rightarrow S$  be the structural morphism. Then  $L$  is the abelian-group functor on  $(\text{Flat})/S$  defined by

$$\text{Hom}_S(T, L) = \Gamma \left( T, \pi_* \text{Hom}_{\mathcal{O}_E} (\Omega_{E/S}^1, \mathcal{J} \mathcal{O}_E) \otimes_{\mathcal{O}_S} \mathcal{O}_T \right),$$

on which  $X$ , regarded as a functor on  $(\text{Flat})/S$ , acts as described in 0.12. In either case there is an exact sequence of group functors on  $(\text{Flat})/S$ :

$$1 \longrightarrow L \longrightarrow X \longrightarrow X^+. \quad (\text{E})$$

Since  $Y$  is flat over  $S$ , Lemma 1.1.1 shows that the groups  $H^i(Y, M)$  depend only on the restriction  $L$  of  $M$  to  $(\text{Flat})/S$ . In case *a*, where  $L = \prod_{S_0/S} L_0$ , Lemma 1.1.2 gives isomorphisms

$$H^i(Y, L) \simeq H^i(Y_0, L_0).$$

Taking all this into account, Proposition 1.2.4 yields the following result.<sup>53</sup>

**Theorem 2.1.** *Let  $S$  be a scheme, and let  $\mathcal{I} \supset \mathcal{J}$  be two quasi-coherent ideals satisfying  $\mathcal{I} \mathcal{J} = 0$ , defining the closed subschemes  $S_0$  and  $S_{\mathcal{J}}$ . Let*

*$X$  be an  $S$ -group functor of type *a* or *b*, with  $L_0$  and  $L$  as above;*

*$Y$  be an  $S$ -group scheme flat over  $S$ , and  $f_{\mathcal{J}} : Y_{\mathcal{J}} \rightarrow X_{\mathcal{J}}$  a morphism of  $S_{\mathcal{J}}$ -groups.*

*Then:*

*i* *The morphism  $f_{\mathcal{J}}$  lifts to a morphism of  $S$ -groups  $Y \rightarrow X$  if and only if the following two conditions hold:*

*$i_1$   $f_{\mathcal{J}}$  lifts to a morphism of  $S$ -functors  $Y \rightarrow X$ . By Proposition 1.2.4 this defines an action of  $Y$  on  $L$ , independent of the chosen lift. In case *a*, the resulting action of  $Y_0$  on  $L_0$  comes from  $f_0 : Y_0 \rightarrow X_0$  and the adjoint action of  $X_0$  on  $L_0$ .*

---

<sup>53</sup>Editor's note: The definitions that appeared in the original statement of Theorem 2.1 and occupied most of original page 113 have been moved above.

- i*<sub>2</sub> The canonically defined obstruction  $c(f_{\mathcal{J}}) \in H^2(Y, L)$  vanishes. In case *a*, this group is isomorphic to  $H^2(Y_0, L_0)$ .
- (ii) If the conditions in *i* hold, the set  $\mathcal{E}$  of morphisms of  $S$ -group functors  $Y \rightarrow X$  extending  $f_{\mathcal{J}}$  is principal homogeneous under  $Z^1(Y, L)$ , isomorphic in case *a* to  $Z^1(Y_0, L_0)$ . Moreover, the quotient of  $\mathcal{E}$  by the action of inner automorphisms of  $X$  defined by sections over  $S$  that induce the unit section of  $X_{\mathcal{J}}$  over  $S_{\mathcal{J}}$  is principal homogeneous under  $H^1(Y, L)$ , isomorphic in case *a* to  $H^1(Y_0, L_0)$ .
- (iii) If  $f : Y \rightarrow X$  is a morphism of  $S$ -group functors extending  $f_{\mathcal{J}}$ , the set of morphisms  $Y \rightarrow X$  obtained from  $f$  by those inner automorphisms is isomorphic to

$$\Gamma(L)/H^0(Y, L),$$

and in case *a* to  $\Gamma(L_0)/H^0(Y_0, L_0)$ .

**Remark 2.1.1.**<sup>54</sup> If  $f, f' : Y \rightarrow X$  are morphisms of  $S$ -group functors extending  $f_{\mathcal{J}}$ , one obtains a cocycle

$$d(f, f') \in Z^1(Y, L) \quad (\simeq Z^1(Y_0, L_0) \text{ in case (a)})$$

such that

$$f' = d(f, f') \cdot f. \quad (*)$$

<sup>55</sup> Write  $\bar{d}(f, f')$  for the image of  $d(f, f')$  in  $H^1(Y, L)$ , or equivalently in  $H^1(Y_0, L_0)$  in case *a*.

**Remark 2.2.** Keep the preceding notation; in particular,  $Y$  is flat over  $S$ . In case *b*, formula (0.14.1) identifies  $L$  with the restriction to  $(\text{Flat})/S$  of

$$\mathbf{W}\left(\pi_* \text{Hom}_{\mathcal{O}_E}(\Omega_{E/S}^1, \mathcal{J} \mathcal{O}_E)\right).$$

In case *a*, suppose in addition that  $X$  is locally of finite presentation over  $S$ . Formula (0.6.1) then identifies  $L$  with the restriction to  $(\text{Flat})/S$  of

$$\prod_{S_0/S} \mathbf{W}\left(\text{Hom}_{\mathcal{O}_{S_0}}(\omega_{X_0/S_0}^1, \mathcal{J})\right).$$

In both cases the module to which  $\mathbf{W}$  is applied is quasi-coherent, by EGA I, 9.1.1. Suppose further that  $Y$  is affine over  $S$ .<sup>56</sup> By I, 5.3, one obtains

$$\begin{aligned} H^i(Y, L) &= H^i(Y_0, L_0) = H^i\left(Y_0, \text{Hom}_{\mathcal{O}_{S_0}}(\omega_{X_0/S_0}^1, \mathcal{J})\right) && \text{in case (a),} \\ H^i(Y, L) &= H^i\left(Y, \pi_* \text{Hom}_{\mathcal{O}_E}(\Omega_{E/S}^1, \mathcal{J} \mathcal{O}_E)\right) && \text{in case (b).} \end{aligned}$$

**Remark 2.3.**

<sup>54</sup>Editor's note: This remark, analogous to 4.5.1, has been added to introduce the notation  $d(f, f')$  and  $\bar{d}(f, f')$ , used in 4.38. Consequently, the part of 2.1(ii) concerning  $\mathcal{E}$  itself was also added.

<sup>55</sup>Editor's note: We follow the sign conventions of the original. One could instead write  $f' = d(f', f) \cdot f$ , but then, in order to retain the equality  $d_1(d(i, i')) = d(Y, i'(Y))$  in 4.27 for two immersions  $i, i' : Y \hookrightarrow X$ , the sign of the class  $d(Y, Y')$  introduced in 4.5.1 would have to be changed. This would in turn alter the signs in 4.8, 4.14, and 4.17. We have preferred to retain all the signs of the original, which are correct.

<sup>56</sup>Editor's note: This hypothesis and the reference to I, 5.3 have been added.



(1) By 0.16 and 0.17, condition  $i_1$  is automatic when  $Y$  is affine and

$$\begin{cases} X \text{ is smooth over } S, & \text{in case (a),} \\ E \text{ is smooth and affine over } S, & \text{in case (b).} \end{cases}$$

(2) Under these hypotheses, with  $Y$  still flat over  $S$  as in 2.0, formulas 2.2(a) and (0.6.2) give in case  $a$

$$H^i(Y, L) = H^i(Y_0, L_0) = H^i\left(Y_0, \text{Lie}(X_0/S_0) \otimes_{\mathcal{O}_{S_0}} \mathcal{J}\right),$$

and in case  $b$ , by (0.14.2), 1.1.2, and I, 5.3,

$$H^i(Y, L) = H^i\left(Y_0, \pi_{0*} \text{Hom}_{\mathcal{O}_{E_0}}(\Omega_{E_0/S_0}^1, \mathcal{J} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{E_0})\right).$$

57

We now state several corollaries for a diagonalizable  $S$ -group  $Y$  (I, 4.4). If  $S$  is affine, then, by I, 5.3.3,  $H^n(Y, F) = 0$  for every  $n > 0$  and every quasi-coherent  $\mathcal{O}_S$ -module  $F$ .

**Corollary 2.4.** *Let  $S_0 \hookrightarrow S$  be a closed subscheme defined by a nilpotent ideal, and let  $Y$  be a diagonalizable  $S$ -group. Let*

*a*  $X$  be an  $S$ -group locally of finite presentation over  $S$ , or

*b*  $X = \underline{\text{Aut}}_S(E)$ , where  $E$  is an  $S$ -scheme locally of finite presentation.

*If  $f : Y \rightarrow X$  is a morphism of  $S$ -groups whose base change  $f_0 : Y_0 \rightarrow X_0$  is the unit morphism, then  $f$  is the unit morphism.*

Indeed, the question is local on  $S$ , and in case  $b$  also local on  $E$ . We may therefore suppose that  $S$  is affine and, in case  $b$ , that  $E$  is of finite presentation over  $S$ . Introduce the closed subschemes  $S_n \subset S$  defined by the powers of the ideal defining  $S_0$ . This reduces the assertion to the case in which  $S_0$  is defined by a square-zero ideal, where it follows from Theorem 2.1 and Remark 2.2.

If  $f_0$  is not required to be the unit morphism, one has the following statement.

**Corollary 2.5.** *Let  $S$  and  $S_0$  be as in 2.4, with  $S$  affine. Let  $Y$  be a diagonalizable  $S$ -group,  $X$  an  $S$ -group functor, and  $f_0 : Y_0 \rightarrow X_0$  a morphism of  $S_0$ -group functors.*

*i* 58

(ii) *Suppose either that  $X$  is a smooth  $S$ -group, or that  $X = \underline{\text{Aut}}_S(E)$  with  $E$  smooth and affine over  $S$ . Then  $f_0$  extends to a morphism of  $S$ -groups  $Y \rightarrow X$ , and any two extensions are conjugate by an inner automorphism of  $X$  defined by a section over  $S$  inducing the unit section of  $X_0$  over  $S_0$ .*

<sup>57</sup>Editor's note: The preceding material has been added; see Editor's Note 31 in 0.14.

<sup>58</sup>Editor's note: The original stated: "Suppose one of the following two properties holds: *a*  $X$  is an  $S$ -group locally of finite presentation; *b*  $X = \underline{\text{Aut}}_S(E)$ , where  $E$  is of finite presentation over  $S$ . Then  $f_0$  extends to a morphism of  $S$ -groups  $Y \rightarrow X$  if and only if it extends to a morphism of  $S$ -functors  $Y \rightarrow X$ ." It added that this followed step by step from part (ii) of the theorem. That argument appears insufficient. For example, if  $\mathcal{J}^3 = 0$  and  $\mathcal{J} = \mathcal{J}^2$ , and if  $f : Y \rightarrow X$  is an  $S$ -functor morphism lifting  $f_0$ , then  $f_0$  lifts to an  $S_{\mathcal{J}}$ -group morphism  $g_{\mathcal{J}} : Y_{\mathcal{J}} \rightarrow X_{\mathcal{J}}$ ; but must  $g_{\mathcal{J}}$  then lift to an  $S$ -functor morphism  $g : Y \rightarrow X$ ? In any event, assertion *i* is not used later, where  $X$  is always assumed smooth over  $S$ .

Introduce  $S_n$  as above.<sup>59</sup> For (ii), a scheme smooth over  $S$  is necessarily locally of finite presentation. Thus, in case  $b$ , a smooth affine  $E/S$  is of finite presentation, so the hypothesis in 2.0(b) is indeed satisfied.

Condition  $i_1$  of Theorem 2.1 is automatic by 0.16 and 0.17. Moreover, every section of  $X_{S_n}$  over  $S_n$  lifts to a section of  $X_{S_{n+1}}$  over  $S_{n+1}$ : this follows from smoothness in case  $a$ , and from 0.17 in case  $b$ . Consequently, if  $f$  and  $f'$  are two lifts of  $f_0$ , one may arrange step by step that  $f_n = f'_n$  by lifting the inner automorphism supplied by Theorem 2.1(ii). This proves the assertion.

The same argument, together with Remark 2.3, gives:

**Corollary 2.6.** *Let  $\mathcal{I}$  be a nilpotent ideal of  $S$  defining  $S_0$ , let  $Y$  be an affine  $S$ -group flat over  $S$ , and let  $X$  be a smooth  $S$ -group.*

*i If, for every  $n \geq 0$ ,*

$$H^2\left(Y_0, \mathrm{Lie}(X_0/S_0) \otimes_{\mathcal{O}_{S_0}} \mathcal{I}^{n+1}/\mathcal{I}^{n+2}\right) = 0,$$

*then every  $S_0$ -group morphism  $f_0 : Y_0 \rightarrow X_0$  lifts to an  $S$ -group morphism  $f : Y \rightarrow X$ .*

*(ii) If, for every  $n \geq 0$ ,*

$$H^1\left(Y_0, \mathrm{Lie}(X_0/S_0) \otimes_{\mathcal{O}_{S_0}} \mathcal{I}^{n+1}/\mathcal{I}^{n+2}\right) = 0,$$

*then any two such lifts are conjugate by an inner automorphism of  $X$  defined by a section over  $S$  inducing the unit section of  $X_0$  over  $S_0$ .*

**Lemma 2.7.** *Let  $S$  be affine,  $G$  an affine  $S$ -group,  $\mathcal{F}$  a quasi-coherent  $\mathcal{O}_S$ -module, and  $\mathcal{L}$  a locally free  $\mathcal{O}_S$ -module. Suppose  $G$  acts on  $\mathcal{F}$  in the sense of Exposé I, hence acts on  $\mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{L}$ . Let  $\Lambda$  be the ring of  $S$ , and let  $L$  be the projective  $\Lambda$ -module defining  $\mathcal{L}$ . Then there is a canonical isomorphism*

$$H^*(G, \mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{L}) \simeq H^*(G, \mathcal{F}) \otimes_{\Lambda} L.$$

60

Let  $\mathcal{A}$  be the  $\mathcal{O}_S$ -algebra  $\mathcal{A}(G)$ , and consider the complex  $\mathcal{C}$  of quasi-coherent  $\mathcal{O}_S$ -modules

$$0 \longrightarrow \mathcal{F} \longrightarrow \mathcal{F} \otimes \mathcal{A} \longrightarrow \mathcal{F} \otimes \mathcal{A} \otimes \mathcal{A} \longrightarrow \cdots.$$

61

By I, 5.3,  $H^*(G, \mathcal{F})$ , respectively  $H^*(G, \mathcal{F} \otimes \mathcal{L})$ , is the cohomology of  $\Gamma(S, \mathcal{C})$ , respectively  $\Gamma(S, \mathcal{C} \otimes \mathcal{L})$ . Since  $S$  is affine, EGA I, 1.3.12 gives

$$\Gamma(S, \mathcal{C} \otimes \mathcal{L}) \simeq \Gamma(S, \mathcal{C}) \otimes_{\Lambda} L.$$

Because  $L$  is projective, hence flat,

$$H^*(\Gamma(S, \mathcal{C}) \otimes_{\Lambda} L) \simeq H^*(\Gamma(S, \mathcal{C})) \otimes_{\Lambda} L,$$

which proves the lemma.

<sup>59</sup>Editor's note: What follows has been slightly modified and expanded from the original.

<sup>60</sup>Editor's note: Here  $\mathcal{L}$  is given the trivial action of  $G$ .

<sup>61</sup>Editor's note: The original proof has been expanded and simplified. It also invoked the isomorphisms  $H^n(\mathcal{C} \otimes \mathcal{L}) \simeq H^n(\mathcal{C}) \otimes \mathcal{L}$  and  $H_n(\Gamma(-)) \simeq \Gamma(H_n(-))$ , where  $H^n(\mathcal{C})$  denotes the cohomology sheaf of the complex  $\mathcal{C}$ .

Using the lemma, Corollary 2.6 becomes:

**Corollary 2.8.** *Let  $S$  be affine and let  $\mathcal{I}$  be a nilpotent ideal defining  $S_0$ . Suppose every  $\mathcal{I}^{n+1}/\mathcal{I}^{n+2}$  is locally free on  $S_0$ . Let  $Y$  be an affine  $S$ -group flat over  $S$ , let  $X$  be a smooth  $S$ -group, and let  $f_0 : Y_0 \rightarrow X_0$  be an  $S_0$ -group morphism.*

- i If  $H^2(Y_0, \text{Lie}(X_0/S_0)) = 0$ , then  $f_0$  lifts to an  $S$ -group morphism  $Y \rightarrow X$ .*
- (ii) If  $H^1(Y_0, \text{Lie}(X_0/S_0)) = 0$ , then any two such lifts are conjugate by an inner automorphism of  $X$  defined by a section over  $S$  inducing the unit section of  $X_0$  over  $S_0$ .*

Taking  $Y = X$  gives:

**Corollary 2.9.** *Let  $S$  and  $S_0$  be as above, and let  $X$  be a smooth affine  $S$ -group.*

- i If  $H^1(X_0, \text{Lie}(X_0/S_0)) = 0$ , every endomorphism of  $X$  over  $S$  that induces the identity on  $X_0$  is the inner automorphism defined by a section over  $S$  inducing the unit section of  $X_0$  over  $S_0$ .*
- (ii) If  $H^2(X_0, \text{Lie}(X_0/S_0)) = 0$ , every  $S_0$ -automorphism of  $X_0$  extends to an  $S$ -automorphism of  $X$ .<sup>62</sup>*

**Remark 2.10.** The assertions involving  $H^1$  have converses by Theorem 2.1. For example, let  $S = I_{S_0}$  be the dual-number scheme over  $S_0$  (II, 2.1), and let  $X$  be a flat  $S$ -group such that every automorphism over  $S$  inducing the identity over  $S_0$  is an inner automorphism defined by a section over  $S$  inducing the unit section of  $X_0$ . Then

$$H^1(X_0, \text{Lie}(X_0/S_0)) = 0.$$

63

**Corollary 2.11.** *Let  $S, \mathcal{I}, \mathcal{J}$  be as in 2.1. Let  $Y$  be an  $S$ -group scheme flat over  $S$ , let  $X$  be an  $S$ -group scheme, and let  $f : Y \rightarrow X$  be an  $S$ -group morphism. The set of morphisms from  $Y$  to  $X$  obtained from  $f$  by conjugation with elements  $x \in X(S)$  inducing the unit of  $X(S_{\mathcal{I}})$  is isomorphic to the quotient*

$$\mathcal{E} = \text{Hom}_{\mathcal{O}_{S_0}}(\omega_{X_0/S_0}^1, \mathcal{J}) / \text{Hom}_{\mathcal{O}_{S_0}}(\omega_{X_0/S_0}^1, \mathcal{J})^{\text{ad}(Y_0)}.$$

Here the second group consists of those  $\mathcal{O}_{S_0}$ -morphisms  $\omega_{X_0/S_0}^1 \rightarrow \mathcal{J}$  which, after every base change  $S' \rightarrow S_0$ , yield morphisms

$$\omega_{X_0/S_0}^1 \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{S'} \longrightarrow \mathcal{J} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{S'}$$

invariant under the action of  $Y_0(S')$  on the first factor.

By Theorem 2.1(iii), this set is  $\Gamma(L_0)/H^0(Y_0, L_0)$ . Now

$$\Gamma(L_0) = \text{Hom}_{\mathcal{O}_{S_0}}(\omega_{X_0/S_0}^1, \mathcal{J}),$$

and  $H^0(Y_0, L_0)$  is precisely  $\Gamma(L_0)^{\text{ad}(Y_0)}$  in the sense just stated.

<sup>62</sup>Editor's note: Let  $f_0$  be an  $S_0$ -automorphism of  $X_0$ , with inverse  $g_0$ . By 2.8(i), lift  $f_0$  and  $g_0$  to  $S$ -endomorphisms  $f$  and  $g$  of  $X$ . Then  $g \circ f$  and  $f \circ g$  induce the identity on  $X_0$ , so by 0.7 they are  $S$ -automorphisms. Hence  $f$  and  $g$  are  $S$ -automorphisms as well.

<sup>63</sup>Editor's note: This is used in Exposé XXIV, 1.13.

**Corollary 2.12.** *Under the hypotheses of 2.11, suppose in addition that  $\omega_{X_0/S_0}^1$  is locally free of finite rank. Then*

$$\mathcal{E} \simeq \Gamma\left(S_0, \operatorname{Lie}(X_0/S_0) \otimes_{\mathcal{O}_{S_0}} \mathcal{J}\right) / H^0\left(Y_0, \operatorname{Lie}(X_0/S_0) \otimes_{\mathcal{O}_{S_0}} \mathcal{J}\right).$$

Indeed,<sup>64</sup> local freeness of finite rank gives

$$\operatorname{Hom}_{\mathcal{O}_{S_0}}(\omega_{X_0/S_0}^1, \mathcal{J}) \simeq \operatorname{Lie}(X_0/S_0) \otimes_{\mathcal{O}_{S_0}} \mathcal{J}.$$

**Corollary 2.13.** *Suppose in addition that  $Y_0$  is diagonalizable. Then*

$$\mathcal{E} \simeq \Gamma\left(S_0, \operatorname{Lie}(X_0/S_0) \otimes_{\mathcal{O}_{S_0}} \mathcal{J}\right) / \Gamma\left(S_0, \operatorname{Lie}(X_0/S_0)^{\operatorname{ad}(Y_0)} \otimes_{\mathcal{O}_{S_0}} \mathcal{J}\right),$$

where  $\operatorname{Lie}(X_0/S_0)^{\operatorname{ad}(Y_0)}$  is the factor of the decomposition in I, 4.7.3 corresponding to the zero character of  $Y_0$ .

Indeed, if  $Y_0 \simeq D_{S_0}(M)$ , the cited result gives a direct-sum decomposition

$$\operatorname{Lie}(X_0/S_0) = \operatorname{Lie}(X_0/S_0)_0 \oplus \bigoplus_{\substack{m \in M \\ m \neq 0}} \operatorname{Lie}(X_0/S_0)_m.$$

Tensoring with  $\mathcal{J}$  gives the analogous decomposition for  $\operatorname{Lie}(X_0/S_0) \otimes \mathcal{J}$ , whence

$$H^0(Y_0, \operatorname{Lie}(X_0/S_0) \otimes \mathcal{J}) \simeq \Gamma\left(S_0, \operatorname{Lie}(X_0/S_0)_0 \otimes_{\mathcal{O}_{S_0}} \mathcal{J}\right).$$

**Corollary 2.14.** *Suppose in addition that  $S_0$  is affine. Then*

$$\mathcal{E} \simeq \Gamma\left(S_0, \left[\operatorname{Lie}(X_0/S_0) / \operatorname{Lie}(X_0/S_0)^{\operatorname{ad}(Y_0)}\right] \otimes_{\mathcal{O}_{S_0}} \mathcal{J}\right).$$

### 3. Infinitesimal extensions of a group scheme

Still using the notation of §0 ( $S$ ,  $\mathcal{J}$ ,  $\mathcal{J}$ , and so on), let  $X$  be an  $S$ -scheme, and suppose that  $X_{\mathcal{J}}$  is endowed with a group structure. We propose to determine the  $S$ -group structures on  $X$  that induce the given structure on  $X_{\mathcal{J}}$ .

From now on, assume that  $X$  is flat over  $S$ . Let  $\mathcal{C}$  be the category of  $S$ -schemes flat over  $S$ ; thus  $X \in \operatorname{Ob} \mathcal{C}$ . Denote by  $Y$ , respectively  $M$ , the functor on  $\mathcal{C}$  defined by  $X^+$ , respectively  $L'_X$ . The canonical morphism  $p_X : X \rightarrow X^+$  defines a morphism in  $\widehat{\mathcal{C}}$

$$p : X \longrightarrow Y,$$

and the action of  $L'_X$  on  $X$  in  $(\widehat{\operatorname{Sch}})/S$  defines an action of  $M$  on  $X$  in  $\widehat{\mathcal{C}}$ . One checks at once that  $X$  is thereby formally principal homogeneous under  $M_Y$  over  $Y$  (cf. 0.2(i) and 0.4).

The action of  $X^+$  on  $L'_X$  defined in 0.8 (denoted  $\operatorname{Ad}$  there) defines an action, denoted  $\tilde{f}$ , of  $Y$  on  $M$ . On the other hand, by 0.5,

$$\operatorname{Hom}_{\widehat{\mathcal{C}}}(Z, M) \simeq \operatorname{Hom}_{S_0}(Z_0, L_0), \quad Z \in \operatorname{Ob} \mathcal{C},$$

where  $L_0$  is the functor defined in 0.5.

**Lemma 3.1.**

<sup>64</sup>Editor's note: The following sentence has been added.

- (i) Condition  $(+)_n$  of 1.3 holds for every positive integer  $n$ .
- (ii) Let the  $S_0$ -group  $X_0$  act on the  $S_0$ -functor  $L_0$  through its adjoint representation. Then there is a canonical isomorphism

$$H^*(X_0, L_0) \simeq H^*(Y, M),$$

where the first cohomology is computed in  $(\text{Sch})/S_0$ , and the second in  $\mathcal{C}$ .

Both parts of the lemma follow from the relation

$$\begin{aligned} \text{Hom}_{\widehat{\mathcal{C}}}(Y, M) &\simeq \text{Hom}_{\widehat{(\text{Sch})/S_0}}(X^+ \times_S S_0, L_0) \\ &\simeq \text{Hom}_{S_0}(X_0, L_0) \\ &\simeq \text{Hom}_{\widehat{\mathcal{C}}}(X, M), \end{aligned}$$

which follows at once from the definition of  $M$  as  $\prod_{S_0/S} L_0$ . More generally, the same relation holds with  $X, Y$  replaced by  $X^n, Y^n$ . Consequently, every morphism  $X^n \rightarrow M$  factors uniquely through  $Y^n$ , which gives  $(+)_n$ . It also gives the relation  $C^*(Y, M) = C^*(X_0, L_0)$ , and hence (ii).

We may therefore apply the constructions of 1.3. In particular:

**Lemma 3.2.** *Let  $P : X \times_S X \rightarrow X$  be a morphism. Then  $P$  induces the group law of  $X_{\mathcal{J}}$  if and only if  $P$  is an admissible composition law on  $X$  (cf. 1.3.2).*

Indeed,  $P$  induces the group law of  $X_{\mathcal{J}}$  if and only if it lifts the group law of  $X^+$ , or equivalently that of  $Y$ . Thus it remains only to show that every morphism  $P$  lifting the group law of  $X_{\mathcal{J}}$  satisfies identity  $((+))$  of 1.3.2(ii), which is exactly what was proved in 0.8.

**Proposition 3.3.** *Let  $S$  be a scheme and  $S_0$  a closed subscheme defined by a nilpotent ideal. Let  $X$  be an  $S$ -scheme that is flat and either quasi-compact or locally of finite presentation over  $S$ . Let  $P : X \times_S X \rightarrow X$  be a composition law on  $X$ . Then  $P$  is a group law if and only if the following two conditions hold:*

- (i)  $P$  is associative;
- (ii)  $P$  induces a group law on  $X_0 = X \times_S S_0$ .

These conditions are clearly necessary. We prove that they are sufficient. Suppose first that  $X \rightarrow S$  has a section. Since  $X(S')$  is then nonempty for every  $S' \rightarrow S$ , it suffices<sup>65</sup> to show that, for every  $x \in X(S')$ , left and right translation by  $x$  are isomorphisms of  $X_{S'}$ .<sup>66</sup>

<sup>65</sup>Editor's note: The original has been corrected by deleting the inappropriate reference to an exercise of Bourbaki on semigroups (cf. [BAI], §I.2, Exercises 9–13) and by indicating the role of the left and right translations; see the following editor's note.

<sup>66</sup>Editor's note: Let  $E$  be a nonempty set endowed with an associative composition law such that every left translation  $\ell_x$  is bijective, and fix  $x_0 \in E$ . There is a unique  $e \in E$  such that  $x_0 e = x_0$ ; then  $x_0 e x = x_0 x$  implies  $e x = x$  for every  $x \in E$ . Moreover, for every  $x$  there is a unique  $x'$  such that  $x x' = e$ . Suppose in addition that there is a  $b \in E$  for which right translation  $r_b$  is injective. Then  $x e b = x b$  gives  $x e = x$ , so  $e$  is a unit, and  $x x' x = x = x e$  gives  $x' x = e$ ; hence  $x'$  is both a left and a right inverse of  $x$ , and  $E$  is a group.

The hypothesis that  $r_b$  be injective is necessary: on any set  $E$ , define  $x y = y$  for all  $x, y \in E$ . Every left translation is then the identity (and the law is associative), whereas  $r_y(E) = \{y\}$  for every  $y$ ; thus  $E$  is not a group when  $|E| > 1$ .

We may plainly assume  $S' = S$ . The translation in question,  $t$ , induces on  $X_0$  a translation  $t_0$ , which is an automorphism since  $X_0$  is a group. The conclusion follows by flatness (SGA 1 III, 4.2).<sup>67</sup>

Now drop the assumption that  $X$  has a section over  $S$ , and suppose that there exists  $S' \rightarrow S$  such that  $X_{S'}$  has a section over  $S'$ . By what has just been proved,  $X_{S'}$  is then an  $S'$ -group; let  $e'$  be its unit section. The inverse image of  $e'$  under  $\text{pr}_i : S' \times_S S' \rightarrow S'$ ,  $i = 1, 2$ , is the unit section of  $X_{S''}$  for the group law obtained by pulling  $P_{S'}$  back along  $\text{pr}_i$ . But since  $P$  is “defined over  $S$ ,” these two group laws coincide, and so do their unit sections. Hence

$$\text{pr}_1^*(e') = \text{pr}_2^*(e').$$

If  $S' \rightarrow S$  is a descent morphism (cf. Exposé IV, no. 2), there is a section of  $X$  whose base extension is  $e'$ , and the proof is complete. Since  $X_X$  has a section over  $X$ , namely the diagonal section, it is now enough to prove that  $X \rightarrow S$  is a descent morphism. It is flat and surjective, and is either quasi-compact or locally of finite presentation; it is therefore an (fpqc) covering, hence a descent morphism (Exposé IV, no. 6).

**Remark.** In fact, the hypothesis that  $X \rightarrow S$  be quasi-compact or locally of finite presentation is superfluous, by the following result, which the reader may prove as an exercise on Exposé IV:

Under the stated hypotheses on  $S$  and  $S_0$ , if  $X \rightarrow S$  is flat and  $X_0 \rightarrow S_0$  is an (fpqc) covering, then  $X \rightarrow S$  is a descent morphism.

**Lemma 3.4.** *Two admissible composition laws on  $X$  are equivalent (cf. 1.3.5) if and only if one is obtained from the other by an automorphism of  $X$  over  $S$  that induces the identity on  $X_{\mathcal{J}}$ .*

Indeed, the morphisms constructed in 1.3.1 are precisely those in the preceding statement, by 0.7.<sup>68</sup>

Taking account of all the preceding results, Proposition 1.3.6 gives:

**Theorem 3.5.** *Let  $S$  be a scheme, and let  $\mathcal{I}$  and  $\mathcal{J}$  be two ideals on  $S$  such that  $\mathcal{I} \supset \mathcal{J}$  and  $\mathcal{I}\mathcal{J} = 0$ . Let  $S_0$  and  $S_{\mathcal{J}}$  be the closed subschemes of  $S$  that they define. Let  $X$  be an  $S$ -scheme flat over  $S$  (and either locally of finite presentation or quasi-compact over  $S$ ), and let  $X_0$  and  $X_{\mathcal{J}}$  be the schemes obtained by base change. Suppose  $X_{\mathcal{J}}$  is endowed with an  $S_{\mathcal{J}}$ -group structure, and let  $L_0$  denote the  $S_0$ -functor in commutative groups defined by*

$$\text{Hom}_{S_0}(T, L_0) = \text{Hom}_{\mathcal{O}_T} \left( \omega_{X_0/S_0}^1 \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_T, \mathcal{J} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_T \right),$$

on which  $X_0$  acts through its adjoint representation.

- (i) *An  $S$ -group structure exists on  $X$  that induces the given structure on  $X_{\mathcal{J}}$  if and only if the following conditions hold:*

- (i<sub>1</sub>) *there is a morphism of  $S$ -schemes  $X \times_S X \rightarrow X$  inducing the group law of  $X_{\mathcal{J}}$ ;*

<sup>67</sup>Editor’s note: Since  $X$  and  $X_0$  have the same underlying topological space and  $t_0$  is an automorphism,  $t$  is a homeomorphism, hence an affine morphism; cf. Exposé VI<sub>B</sub>, 2.9.1, or EGA IV<sub>4</sub>, 18.12.7.1. It therefore suffices to show the following. Let  $\mathcal{J}$  be a nilpotent ideal of a ring  $\Lambda$ , and let  $\varphi : A \rightarrow B$  be a morphism of  $\Lambda$ -algebras, with  $B$  flat over  $\Lambda$ , such that  $\varphi \otimes_{\Lambda} (\Lambda/\mathcal{J})$  is bijective. Then  $\varphi$  is bijective. By the “nilpotent Nakayama lemma,”  $\varphi$  is surjective; moreover, since  $B$  is flat over  $\Lambda$ , one also has  $\text{Ker}(\varphi) \otimes_{\Lambda} (\Lambda/\mathcal{J}) = 0$ , whence  $\text{Ker}(\varphi) = 0$ . Thus  $\varphi$  is bijective.

<sup>68</sup>Editor’s note: Indeed, by the proof of 0.7, the  $S$ -endomorphisms of  $X$  inducing the identity on  $X_{\mathcal{J}}$  are the automorphisms  $m \cdot \text{id}_X$ , where  $m$  ranges over  $M(X) = \text{Hom}_S(X, M)$ . For every  $S' \rightarrow S$  and every  $x \in X(S')$ , one has  $(m \cdot \text{id}_X)(x) = m(x) \cdot x$ . By the proof of 3.1, each  $m : X \rightarrow M$  factors uniquely through a morphism  $h : Y = X^+ \rightarrow M$ ; hence  $m \cdot \text{id}_X$  is the automorphism  $u_h$  introduced in 1.3.1. The lemma now follows from the definition of equivalence; cf. 1.3.4 and 1.3.5.

- (i<sub>2</sub>) a certain obstruction class in  $H^3(X_0, L_0)$ , canonically defined by  $X$  and the group law of  $X_{\mathcal{J}}$ , is zero.
- (ii) If the conditions in (i) hold, the set  $\mathcal{E}$  of group laws on  $X$  inducing the given law on  $X_{\mathcal{J}}$  is principal homogeneous under  $Z^2(X_0, L_0)$ . Moreover, the quotient of  $\mathcal{E}$  by the  $S$ -automorphisms of  $X$  inducing the identity on  $X_{\mathcal{J}}$  is principal homogeneous under  $H^2(X_0, L_0)$ .

<sup>69</sup> Indeed, by 3.2 every morphism of  $S$ -schemes  $f : X \times_S X \rightarrow X$  inducing the group law of  $X_{\mathcal{J}}$  is an admissible composition law on  $X$ . By 1.3.6(i), the existence of an associative admissible composition law  $P : X \times_S X \rightarrow X$  is equivalent to the vanishing of a certain class  $c(f) \in H^3(X_0, L_0)$ ; in that case, 3.3 shows that  $P$  is a group law. This proves (i), and (ii) then follows from 3.3 and 1.3.6(ii).

**Remark 3.5.1.** <sup>70</sup> If  $\mu$  and  $\mu'$  are group laws on  $X$  inducing the given law on  $X_{\mathcal{J}}$ , one obtains a cocycle  $\delta(\mu, \mu') \in Z^2(X_0, L_0)$ . The chosen sign convention is  $\mu' = \delta(\mu, \mu') \cdot \mu$ ; that is, for every  $S' \rightarrow S$  and  $x, y \in X(S')$ ,

$$\mu'(x, y) = \delta(\mu, \mu')(x_0, y_0) \cdot \mu(x, y).$$

71

Write  $\bar{\delta}(\mu, \mu')$  for the image of  $\delta(\mu, \mu')$  in  $H^2(X_0, L_0)$ . Finally, if  $X$  endowed with  $\mu$ , respectively  $\mu'$ , is denoted simply by  $X$ , respectively  $X'$ , write  $\delta(X, X')$  in place of  $\delta(\mu, \mu')$ , and similarly for  $\bar{\delta}(X, X')$ .

**Remark 3.6.** Let  $X_{\mathcal{J}}$  be a smooth affine  $S_{\mathcal{J}}$ -scheme. By 0.15, up to isomorphism there is a unique  $S$ -scheme  $X$ , smooth over  $S$ , whose reduction modulo  $\mathcal{J}$  is  $X_{\mathcal{J}}$ . If  $X_{\mathcal{J}}$  is endowed with an  $S_{\mathcal{J}}$ -group structure, condition (i<sub>1</sub>) is automatic by 0.16. Moreover, 0.6 simplifies the definition of  $L_0$  and gives:

**Corollary 3.7.** Let  $S, \mathcal{I}, \mathcal{J}$  be as in 3.1, and let  $X_{\mathcal{J}}$  be a smooth affine  $S_{\mathcal{J}}$ -group.

- (i) The set of smooth  $S$ -groups whose reduction modulo  $\mathcal{J}$  is  $X_{\mathcal{J}}$ , up to isomorphism inducing the identity on  $X_{\mathcal{J}}$ , is either empty or principal homogeneous under

$$H^2\left(X_0, \text{Lie}(X_0/S_0) \otimes_{\mathcal{O}_{S_0}} \mathcal{J}\right).$$

- (ii) A smooth  $S$ -group whose reduction modulo  $\mathcal{J}$  is  $X_{\mathcal{J}}$  exists if and only if a certain obstruction in

$$H^3\left(X_0, \text{Lie}(X_0/S_0) \otimes_{\mathcal{O}_{S_0}} \mathcal{J}\right)$$

is zero.

As usual, one obtains the following corollaries.

**Corollary 3.8.** Let  $S$  be a scheme and  $S_0$  a closed subscheme defined by a nilpotent ideal  $\mathcal{J}$ . Let  $X_0$  be a smooth affine  $S_0$ -group.

<sup>69</sup>Editor's note: The following argument has been added.

<sup>70</sup>Editor's note: This remark, analogous to 4.5.1, has been added to introduce the notation  $\delta(\mu, \mu')$ , or  $\delta(X, X')$ , used in 4.38. Consequently, the part of 3.5(ii) concerning  $\mathcal{E}$  itself has also been added.

<sup>71</sup>Editor's note: We follow the sign conventions of the original in order to have, in 4.38(5), the equality  $\partial^1 \bar{d}(X, X') = \bar{\delta}(X, X')$ ; see also Editor's Note 54.

(i) If, for every  $n \geq 0$ ,

$$H^2\left(X_0, \text{Lie}(X_0/S_0) \otimes_{\mathcal{O}_{S_0}} \mathcal{I}^{n+1}/\mathcal{I}^{n+2}\right) = 0,$$

then any two smooth  $S$ -groups whose reduction is  $X_0$  are isomorphic by an isomorphism inducing the identity on  $X_0$ .

(ii) If, for every  $n \geq 0$ ,

$$H^3\left(X_0, \text{Lie}(X_0/S_0) \otimes_{\mathcal{O}_{S_0}} \mathcal{I}^{n+1}/\mathcal{I}^{n+2}\right) = 0,$$

then there exists a smooth  $S$ -group whose reduction is  $X_0$ .

**Corollary 3.9.** *Let  $S$  be affine and  $S_0$  a closed subscheme defined by a nilpotent ideal  $\mathcal{I}$ . Suppose that every  $\mathcal{I}^{n+1}/\mathcal{I}^{n+2}$  is locally free on  $S_0$ . Let  $X_0$  be a smooth affine  $S_0$ -group.*

(i) *If  $H^2(X_0, \text{Lie}(X_0/S_0)) = 0$ , then any two smooth  $S$ -groups whose reduction is  $X_0$  are isomorphic.*

(ii) *If  $H^3(X_0, \text{Lie}(X_0/S_0)) = 0$ , then there exists a smooth  $S$ -group whose reduction is  $X_0$ .*

**Corollary 3.10.** *Let  $S_0$  be a scheme and let  $S = I_{S_0}$  be the dual-number scheme over  $S_0$ . Let  $X_0$  be a smooth  $S_0$ -group. Then every smooth  $S$ -group  $Y$  for which  $Y_0$  is  $S_0$ -isomorphic to  $X_0$  is  $S$ -isomorphic to  $X = X_0 \times_{S_0} S$  if and only if*

$$H^2(X_0, \text{Lie}(X_0/S_0)) = 0.$$

72

Indeed, by 3.5 the set of classes of such groups  $Y$ , up to an  $S$ -group isomorphism inducing the identity on  $X_0$ , is in bijection with  $H^2(X_0, \text{Lie}(X_0/S_0))$ . Hence their classes up to an arbitrary  $S$ -group isomorphism are in bijection with

$$H^2(X_0, \text{Lie}(X_0/S_0))/\Gamma_0,$$

where

$$\Gamma_0 = \text{Aut}_{S_0\text{-gr.}}(X_0),$$

which acts in the evident way on  $H^2$ . The conclusion follows at once. <sup>73</sup>

## 4. Infinitesimal extensions of closed subgroups

We first state a result valid in an arbitrary abelian category.

**Lemma 4.1.** *Let*

$$0 \longrightarrow A' \xrightarrow{i} A \xrightarrow{p} A'' \longrightarrow 0$$

*be an exact sequence, let  $\phi : A' \rightarrow Q$  be a morphism, and let  $\pi : A'' \rightarrow P$  be an epimorphism with kernel  $C$ . Let  $E$  be the set, up to isomorphism, of quadruples  $(B, f, g, h)$  such that the sequence*

$$0 \longrightarrow Q \xrightarrow{f} B \xrightarrow{g} P \longrightarrow 0$$

*is exact and the following diagram commutes:*

<sup>72</sup>Editor's note: This result is used in XXIV, 1.13.

<sup>73</sup>Editor's note: Indeed,  $\text{Aut}_{S_0\text{-gr.}}(X_0)$  acts by group automorphisms on the abelian group  $H^2(X_0, \text{Lie}(X_0/S_0))$ , so the orbit of 0 is the singleton  $\{0\}$ . Consequently, the quotient set is a singleton if and only if  $H^2(X_0, \text{Lie}(X_0/S_0)) = \{0\}$ .



$$\begin{array}{ccccccccc}
0 & \longrightarrow & A' & \xrightarrow{i} & A & \xrightarrow{p} & A'' & \longrightarrow & 0 \\
& & \downarrow \phi & & \downarrow h & & \downarrow \pi & & \\
0 & \longrightarrow & Q & \xrightarrow{f} & B & \xrightarrow{g} & P & \longrightarrow & 0.
\end{array}$$

- (i) The set  $E$  is nonempty if and only if the image in  $\text{Ext}^1(C, Q)$  of the element  $A$  of  $\text{Ext}^1(A'', A')$  is zero.
- (ii) Under these conditions,  $E$  is a principal homogeneous set under the abelian group  $\text{Hom}(C, Q)$ .

Introduce the amalgamated sum

$$B' = A \sqcup_{A'} Q.$$

There is then a commutative diagram with exact rows: <sup>74</sup>

$$\begin{array}{ccccccccc}
0 & \longrightarrow & A' & \xrightarrow{i} & A & \xrightarrow{p} & A'' & \longrightarrow & 0 \\
& & \downarrow \phi & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & Q & \xrightarrow{j} & B' & \longrightarrow & A'' & \longrightarrow & 0,
\end{array}$$

It is clear that the category of solutions to the problem is canonically isomorphic to the category of solutions to the corresponding problem for the sequence

$$0 \longrightarrow Q \longrightarrow B' \longrightarrow A'' \longrightarrow 0$$

and the morphisms  $\text{id}_Q$  and  $\pi : A'' \rightarrow P$ . <sup>75</sup> In this case,  $E$  is in bijection with the set of subobjects  $N$  of  $B'$  for which  $B' \rightarrow A''$  induces an isomorphism of  $N$  with the kernel  $C$  of  $A'' \rightarrow P$ ; equivalently, it is the set of morphisms  $e : C \rightarrow B'$  lifting the canonical morphism  $C \rightarrow A''$ . The abelian group  $G = \text{Hom}(C, Q)$  acts on  $E$  by  $g \cdot e = g + e$ , with addition in  $\text{Hom}(C, B')$ ; if  $E \neq \emptyset$ , this makes  $E$  a principal homogeneous set under  $G$ .

We deduce the following.

**Proposition 4.2.** <sup>76</sup> Let  $S$  be a scheme, let  $S_{\mathcal{J}}$  be the closed subscheme defined by a quasi-coherent square-zero ideal  $\mathcal{J}$ , let  $X$  be an  $S$ -scheme, and let  $\mathcal{F}$  be an  $\mathcal{O}_X$ -module. Put

$$X_{\mathcal{J}} = X \times_S S_{\mathcal{J}}, \quad \mathcal{F}_{\mathcal{J}} = \mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{O}_{S_{\mathcal{J}}},$$

and let  $\mathcal{G}_{\mathcal{J}} = \mathcal{F}_{\mathcal{J}} / \mathcal{H}_{\mathcal{J}}$  be a quotient module of  $\mathcal{F}_{\mathcal{J}}$ . Suppose given a morphism of  $\mathcal{O}_{X_{\mathcal{J}}}$ -modules

$$f : \mathcal{J} \otimes_{\mathcal{O}_S} \mathcal{G}_{\mathcal{J}} \longrightarrow \mathcal{Q}.$$

<sup>74</sup>Editor's note: One has  $\text{Coker}(j) = B' \sqcup_Q 0 = A \sqcup_{A'} 0 = A''$ , and  $\text{Ker}(j) \simeq \text{Ker}(i) = 0$ , as one sees by reasoning "as if the category were a category of modules." For a proof entirely in terms of arrows, see for example [Fr64], Theorem 2.5.4 (\*).

<sup>75</sup>Editor's note: In what follows,  $A$  has been replaced by  $B'$ , and the end of the argument has been expanded.

<sup>76</sup>Editor's note: The statement has been rewritten so as to fit exactly the application made in 4.3; the proof has also been expanded, following indications supplied by M. Demazure.

Let  $\mathcal{E}$  be the sheaf of sets on  $X$  defined as follows: for each open subset  $U$  of  $X$ ,  $\mathcal{E}(U)$  is the set of quotient modules  $\mathcal{G}$  of  $\mathcal{F}|_U$  such that  $\mathcal{G}/\mathcal{I}\mathcal{G} = \mathcal{G}_{\mathcal{I}}|_U$  and for which there exists an isomorphism

$$h : \mathcal{I}\mathcal{G} \xrightarrow{\sim} \mathcal{Q}|_U$$

making the following diagram commute:

$$\begin{array}{ccc} \mathcal{I} \otimes_{\mathcal{O}_{S_{\mathcal{I}}}} (\mathcal{G}_{\mathcal{I}}|_U) & \xrightarrow{f|_U} & \mathcal{Q}|_U \\ \text{can.} \downarrow & \nearrow h \simeq & \\ \mathcal{I}\mathcal{G} & & \end{array} .$$

The isomorphism  $h$  is then unique, since  $\mathcal{I} \otimes_{\mathcal{O}_{S_{\mathcal{I}}}} (\mathcal{G}_{\mathcal{I}}|_U) \rightarrow \mathcal{I}\mathcal{G}$  is an epimorphism. Then  $\mathcal{E}$  is a sheaf formally principal homogeneous under the sheaf of commutative groups

$$\mathcal{A} = \text{Hom}_{\mathcal{O}_X}(\mathcal{H}_{\mathcal{I}}, \mathcal{Q}) = \text{Hom}_{\mathcal{O}_{X_{\mathcal{I}}}}(\mathcal{H}_{\mathcal{I}}, \mathcal{Q}).$$

*Proof.* If  $\mathcal{E}(U) = \emptyset$ , there is nothing to prove; we may therefore suppose that  $\mathcal{E}(U)$  contains an element  $\mathcal{G}$ . In the following diagram,  $h$  is an isomorphism and all the arrows are epimorphisms:

$$\begin{array}{ccccc} \mathcal{I} \otimes_{\mathcal{O}_{S_{\mathcal{I}}}} (\mathcal{F}_{\mathcal{I}}|_U) & \longrightarrow & \mathcal{I} \otimes_{\mathcal{O}_{S_{\mathcal{I}}}} (\mathcal{G}_{\mathcal{I}}|_U) & \xrightarrow{f|_U} & \mathcal{Q}|_U \\ \text{can.} \downarrow & & \text{can.} \downarrow & \nearrow h \simeq & \\ \mathcal{I}\mathcal{F}|_U & \longrightarrow & \mathcal{I}\mathcal{G} & & \end{array} .$$

Thus the morphism

$$\mathcal{I} \otimes_{\mathcal{O}_{S_{\mathcal{I}}}} (\mathcal{F}_{\mathcal{I}}|_U) \longrightarrow \mathcal{Q}|_U$$

induces a necessarily unique epimorphism  $\phi : \mathcal{I}\mathcal{F}|_U \rightarrow \mathcal{Q}|_U$ . If  $\mathcal{G}$  is an  $\mathcal{O}_U$ -module such that  $\mathcal{G}/\mathcal{I}\mathcal{G} = \mathcal{G}_{\mathcal{I}}|_U$  and there is a commutative diagram with exact rows

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{I}\mathcal{F}|_U & \longrightarrow & \mathcal{F}|_U & \longrightarrow & \mathcal{F}_{\mathcal{I}}|_U & \longrightarrow & 0 \\ & & \phi \downarrow & & \downarrow & & \downarrow \pi & & \\ 0 & \longrightarrow & \mathcal{Q}|_U & \longrightarrow & \mathcal{G} & \xrightarrow{p_{\mathcal{I}}} & \mathcal{G}_{\mathcal{I}}|_U & \longrightarrow & 0 \end{array}$$

(where  $p_{\mathcal{I}}$  is the projection  $\mathcal{G} \rightarrow \mathcal{G}/\mathcal{I}\mathcal{G} = \mathcal{G}_{\mathcal{I}}|_U$ , so that  $\mathcal{Q}|_U = \text{Ker}(p_{\mathcal{I}}) = \mathcal{I}\mathcal{G}$ ), then  $\mathcal{G}$  may be identified with a quotient module of  $\mathcal{F}|_U$ . Consequently, by 4.1(ii), the set  $\mathcal{E}(U)$  is principal homogeneous under the abelian group

$$\text{Hom}_{\mathcal{O}_X}(\mathcal{H}_{\mathcal{I}}, \mathcal{Q})(U) = \text{Hom}_{\mathcal{O}_{X_{\mathcal{I}}}}(\mathcal{H}_{\mathcal{I}}, \mathcal{Q})(U).$$

**Proposition 4.3.** (TDTE IV 5.1) *Let  $S$  be a scheme, let  $S_{\mathcal{J}}$  be the closed subscheme defined by a quasi-coherent square-zero ideal  $\mathcal{J}$ , let  $X$  be an  $S$ -scheme, and let  $\mathcal{F}$  be a quasi-coherent  $\mathcal{O}_X$ -module. Put*

$$X_{\mathcal{J}} = X \times_S S_{\mathcal{J}}, \quad \mathcal{F}_{\mathcal{J}} = \mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{O}_{S_{\mathcal{J}}}.$$

*Let  $\mathcal{G}_{\mathcal{J}} = \mathcal{F}_{\mathcal{J}} / \mathcal{H}_{\mathcal{J}}$  be a quasi-coherent quotient module of  $\mathcal{F}_{\mathcal{J}}$ , flat over  $S_{\mathcal{J}}$ . For each open subset  $U$  of  $X$ , let  $\mathcal{E}(U)$  be the set of quasi-coherent quotient modules  $\mathcal{G}^{77}$  of  $\mathcal{F}|_U$ , flat over  $S$ , such that  $\mathcal{G} / \mathcal{J}\mathcal{G} \simeq \mathcal{G}_{\mathcal{J}}|_U$ . Then the  $\mathcal{E}(U)$  form a sheaf of sets  $\mathcal{E}$  on  $X$ , formally principal homogeneous under the sheaf of commutative groups*

$$\mathcal{A} = \mathrm{Hom}_{\mathcal{O}_{X_{\mathcal{J}}}} \left( \mathcal{H}_{\mathcal{J}}, \mathcal{J} \otimes_{\mathcal{O}_{S_{\mathcal{J}}}} \mathcal{G}_{\mathcal{J}} \right).$$

*Proof.* Let  $\pi : X \rightarrow S$  be the structural morphism. Let  $U$  be an open subset of  $X$ , and let  $\mathcal{G}$  be an  $\mathcal{O}_U$ -module flat over  $S$ , such that  $\mathcal{G} / \mathcal{J}\mathcal{G} \simeq \mathcal{G}_{\mathcal{J}}|_U$ . For every  $x \in U$ , the stalk  $\mathcal{G}_x$  is a flat module over the local ring  $\mathcal{O}_{S,s}$ , where  $s = \pi(x)$ . Hence the morphism

$$\mathcal{J}_s \otimes_{\mathcal{O}_{S,s}} (\mathcal{G} / \mathcal{J}\mathcal{G})_x = \mathcal{J}_s \otimes_{\mathcal{O}_{S,s}} \mathcal{G}_x \longrightarrow (\mathcal{J}\mathcal{G})_x$$

is bijective. We therefore have an exact sequence

$$0 \longrightarrow \mathcal{J} \otimes_{\mathcal{O}_S} (\mathcal{G}_{\mathcal{J}}|_U) \longrightarrow \mathcal{G} \longrightarrow \mathcal{G}_{\mathcal{J}}|_U \longrightarrow 0.$$

Since  $\mathcal{J} \otimes_{\mathcal{O}_S} (\mathcal{G}_{\mathcal{J}}|_U)$  and  $\mathcal{G}_{\mathcal{J}}|_U$  are quasi-coherent  $\mathcal{O}_U$ -modules,  $\mathcal{G}$  is quasi-coherent as well (EGA III, 1.4.17).

Conversely, since  $\mathcal{G}_{\mathcal{J}}$  is flat over  $S_{\mathcal{J}}$ , if  $\mathcal{G}$  is a quasi-coherent  $\mathcal{O}_U$ -module such that  $\mathcal{G} / \mathcal{J}\mathcal{G} \simeq \mathcal{G}_{\mathcal{J}}$  and the morphism

$$\mathcal{J} \otimes_{\mathcal{O}_{S_{\mathcal{J}}}} \mathcal{G}_{\mathcal{J}} \longrightarrow \mathcal{J}\mathcal{G}$$

is bijective, then  $\mathcal{G}$  is flat over  $S$ , by the “fundamental criterion of flatness” (SGA 1, IV, 5.5).<sup>78</sup>

Consequently, the set  $\mathcal{E}(U)$  considered here is the set considered in 4.2 when  $f$  is the identity morphism of  $\mathcal{J} \otimes_{\mathcal{O}_{S_{\mathcal{J}}}} \mathcal{G}_{\mathcal{J}}$ ; the conclusion therefore follows from 4.2.  $\square$

Retain the preceding notation.<sup>79</sup> Let  $Y_{\mathcal{J}}$  be a closed subscheme of  $X_{\mathcal{J}}$ , defined by a quasi-coherent ideal  $\mathcal{I}_{Y_{\mathcal{J}}}$ , and suppose that  $Y_{\mathcal{J}}$  is flat over  $S_{\mathcal{J}}$ . Applying 4.3 to  $\mathcal{F} = \mathcal{O}_X$  and

$$\mathcal{G}_{\mathcal{J}} = \mathcal{O}_{Y_{\mathcal{J}}} = \mathcal{O}_{X_{\mathcal{J}}} / \mathcal{I}_{Y_{\mathcal{J}}},$$

we obtain the following corollary.

**Corollary 4.3.1.** *Let  $S, S_{\mathcal{J}}, \mathcal{J}, X, X_{\mathcal{J}}, Y_{\mathcal{J}}$ , and  $\mathcal{I}_{Y_{\mathcal{J}}}$  be as above, and suppose that  $Y_{\mathcal{J}}$  is flat over  $S_{\mathcal{J}}$ . Let  $\mathcal{A}_{\mathcal{J}}$  be the sheaf of commutative groups*

$$\mathrm{Hom}_{\mathcal{O}_{X_{\mathcal{J}}}} \left( \mathcal{I}_{Y_{\mathcal{J}}}, \mathcal{J} \otimes_{\mathcal{O}_{S_{\mathcal{J}}}} \mathcal{O}_{Y_{\mathcal{J}}} \right)$$

<sup>77</sup>Editor’s note: To lighten the statement, the hypothesis that  $\mathcal{G}$  be quasi-coherent has been added here; the remark that this hypothesis is automatically satisfied has been moved to the proof, which has been expanded accordingly.

<sup>78</sup>Editor’s note: See also [BAC], §III.5, Theorem 1.

<sup>79</sup>Editor’s note: The discussion that follows has been expanded and Corollary 4.3.1 has been added. Recall also that “pseudo-torsor” is synonymous with “formally principal homogeneous.”

on  $X_{\mathcal{J}}$ , and put  $\mathcal{A} = i_*(\mathcal{A}_{\mathcal{J}})$ , where  $i : X_{\mathcal{J}} \hookrightarrow X$  is the immersion. For each open subset  $U$  of  $X$ , let  $\mathcal{E}(U)$  be the set of closed subschemes  $Y$  of  $U$ , flat over  $S$ , such that

$$Y \times_S S_{\mathcal{J}} = Y_{\mathcal{J}} \cap U.$$

Then  $\mathcal{E}$  is an  $\mathcal{A}$ -pseudo-torsor.

If, moreover, such a  $Y$  exists locally—that is, if every  $x \in X$  has an open neighborhood  $U$  with  $\mathcal{E}(U) \neq \emptyset$ —then  $\mathcal{E}$  is an  $\mathcal{A}$ -torsor. The  $\mathcal{A}$ -torsors on  $X$  are parametrized by

$$H^1(X, \mathcal{A}) = H^1(X_{\mathcal{J}}, \mathcal{A}_{\mathcal{J}})$$

(see, for example, EGA IV<sub>4</sub>, 16.5.15), and  $\mathcal{E}$  has a global section—that is,  $\mathcal{E}(X) \neq \emptyset$ —if and only if its corresponding cohomology class is zero. Thus we obtain:

**Corollary 4.4.** *Let  $S, S_{\mathcal{J}}, \mathcal{J}, X, X_{\mathcal{J}}, Y_{\mathcal{J}}$ , and  $\mathcal{I}_{Y_{\mathcal{J}}}$  be as above, and suppose that  $Y_{\mathcal{J}}$  is flat over  $S_{\mathcal{J}}$ . Let  $E$  be the set of closed subschemes  $Y$  of  $X$ , flat over  $S$ , such that  $Y \times_S S_{\mathcal{J}} = Y_{\mathcal{J}}$ .*

- (i) *The set  $E$  is either empty or principal homogeneous under the abelian group*

$$H^0(X, \mathcal{A}) = H^0(X_{\mathcal{J}}, \mathcal{A}_{\mathcal{J}}) = \operatorname{Hom}_{\mathcal{O}_{X_{\mathcal{J}}}} \left( \mathcal{I}_{Y_{\mathcal{J}}}, \mathcal{J} \otimes_{\mathcal{O}_{S_{\mathcal{J}}}} \mathcal{O}_{Y_{\mathcal{J}}} \right).$$

- (ii) *The set  $E$  is nonempty if and only if both of the following conditions hold:*

- (a) *The problem has a solution locally on  $X$ .*  
 (b) *A certain obstruction vanishes; it lies in*

$$H^1 \left( X_{\mathcal{J}}, \operatorname{Hom}_{\mathcal{O}_{X_{\mathcal{J}}}} \left( \mathcal{I}_{Y_{\mathcal{J}}}, \mathcal{J} \otimes_{\mathcal{O}_{S_{\mathcal{J}}}} \mathcal{O}_{Y_{\mathcal{J}}} \right) \right).$$

**Complement 4.4.1.** <sup>80</sup> Retain the notation of 4.4, and suppose that  $E$  contains an element  $Y$ . Let  $\mathcal{I}_Y$  be the ideal of  $\mathcal{O}_X$  defining  $Y$ , and let  $\mathcal{I}_{Y_{\mathcal{J}}}$  be its image in  $\mathcal{O}_{X_{\mathcal{J}}}$ . As we saw in the proof of 4.2, there is a commutative diagram

$$\begin{array}{ccc} \mathcal{J} \otimes_{\mathcal{O}_{S_{\mathcal{J}}}} \mathcal{O}_{X_{\mathcal{J}}} & \twoheadrightarrow & \mathcal{J} \mathcal{O}_X \\ \downarrow & \swarrow \text{dashed} & \downarrow \\ \mathcal{J} \otimes_{\mathcal{O}_{S_{\mathcal{J}}}} \mathcal{O}_{Y_{\mathcal{J}}} & \xrightarrow{\sim} & \mathcal{J} \mathcal{O}_Y \end{array}$$

and hence an epimorphism of  $\mathcal{O}_X$ -modules

$$\phi : \mathcal{J} \mathcal{O}_X \longrightarrow \mathcal{J} \otimes_{\mathcal{O}_{S_{\mathcal{J}}}} \mathcal{O}_{Y_{\mathcal{J}}}.$$

Let  $\mathcal{K}$  be its kernel. For every element  $Y'$  of  $E$ , the morphism  $\mathcal{O}_X \rightarrow \mathcal{O}_{Y'}$  then factors through  $\mathcal{O}_X / \mathcal{K}$ , which is the amalgamated sum  $B'$  in the proof of Lemma 4.1. If  $\mathcal{I}_{Y'}$  denotes the ideal of  $Y'$  in  $\mathcal{O}_X$ , there is a commutative diagram

<sup>80</sup>Editor's note: This complement has been added; it is useful in proving Proposition 4.8(ii).

$$\begin{array}{ccccccc}
& & & 0 & & 0 & \\
& & & \downarrow & & \downarrow & \\
& & & \mathcal{I}_{Y'}/\mathcal{K} & \xrightarrow{\sim} & \mathcal{I}_{Y_{\mathcal{J}}} & \\
& & & \downarrow & & \downarrow & \\
0 & \longrightarrow & (\mathcal{I}\mathcal{O}_X)/\mathcal{K} & \longrightarrow & \mathcal{O}_X/\mathcal{K} & \longrightarrow & \mathcal{O}_{X_{\mathcal{J}}} \longrightarrow 0 \\
& & \downarrow \wr & & \downarrow & & \downarrow \\
0 & \longrightarrow & \mathcal{I} \otimes_{\mathcal{O}_{S_{\mathcal{J}}}} \mathcal{O}_{Y_{\mathcal{J}}} & \longrightarrow & \mathcal{O}_{Y'} & \longrightarrow & \mathcal{O}_{Y_{\mathcal{J}}} \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
& & 0 & & 0 & & 0
\end{array}$$

Replacing  $X$  by the closed subscheme defined by  $\mathcal{K}$ , we are thus reduced to the case  $\mathcal{K} = 0$ . The datum of  $Y'$  is then equivalent to that of the  $\mathcal{O}_X$ -submodule  $\mathcal{I}_{Y'}$  of  $\mathcal{O}_X$  which maps bijectively onto  $\mathcal{I}_{Y_{\mathcal{J}}}$  under the projection  $p : \mathcal{O}_X \rightarrow \mathcal{O}_{X_{\mathcal{J}}}$ . Let

$$f' : \mathcal{I}_{Y_{\mathcal{J}}} \xrightarrow{\sim} \mathcal{I}_{Y'}, \quad f : \mathcal{I}_{Y_{\mathcal{J}}} \xrightarrow{\sim} \mathcal{I}_Y$$

denote the respective inverse isomorphisms. Then  $f' - f$  is an element of

$$\mathrm{Hom}_{\mathcal{O}_{X_{\mathcal{J}}}}(\mathcal{I}_{Y_{\mathcal{J}}}, \mathcal{I} \otimes_{\mathcal{O}_{S_{\mathcal{J}}}} \mathcal{O}_{Y_{\mathcal{J}}}) = \mathrm{Hom}_{\mathcal{O}_X}(\mathcal{I}_{Y_{\mathcal{J}}}, \mathcal{I} \mathcal{O}_Y),$$

which we denote by  $d(Y', Y)$ . Notice that  $d(Y, Y') = -d(Y', Y)$ .

For our fixed  $Y$  and variable  $Y'$ , consider the morphism

$$\mathcal{I}_{Y'} \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{O}_Y = \mathcal{O}_X / \mathcal{I}_Y.$$

Since its composite with  $\mathcal{O}_Y \rightarrow \mathcal{O}_{Y_{\mathcal{J}}}$  is zero, it takes values in

$$\mathcal{I} \mathcal{O}_Y = \mathcal{I} \otimes_{\mathcal{O}_{S_{\mathcal{J}}}} \mathcal{O}_{Y_{\mathcal{J}}}.$$

More precisely, let  $V$  be an open subset of  $X$ , let  $x'$  be a section of  $\mathcal{I}_{Y'}$  on  $V$ , and let  $x_{\mathcal{J}}$  be its image in  $\Gamma(V, \mathcal{I}_{Y_{\mathcal{J}}})$ . Then

$$x' = f'(x_{\mathcal{J}}) = f(x_{\mathcal{J}}) + (f' - f)(x_{\mathcal{J}}) = f(x_{\mathcal{J}}) + d(Y', Y)(x_{\mathcal{J}}).$$

Consequently, the morphism  $\mathcal{I}_{Y'} \rightarrow \mathcal{I} \mathcal{O}_Y$  is given by  $d(Y', Y)$ .

**4.5.0.**<sup>81</sup> Retain the notation of 4.3.1 and 4.4, and make the following transformations. The quasi-coherent  $\mathcal{O}_{X_{\mathcal{J}}}$ -module

$$\mathcal{I}_{Y_{\mathcal{J}}} / \mathcal{I}_{Y_{\mathcal{J}}}^2$$

is annihilated by  $\mathcal{I}_{Y_{\mathcal{J}}}$ , and is therefore the direct image of a quasi-coherent  $\mathcal{O}_{Y_{\mathcal{J}}}$ -module, denoted by  $\mathcal{N}_{Y_{\mathcal{J}}/X_{\mathcal{J}}}$ , and called the *conormal sheaf* of  $Y_{\mathcal{J}}$  in  $X_{\mathcal{J}}$ .<sup>82</sup> Since  $\mathcal{I} \otimes_{\mathcal{O}_{S_{\mathcal{J}}}} \mathcal{O}_{Y_{\mathcal{J}}}$

<sup>81</sup>Editor's note: The numbering 4.5.0 has been added to mark the return to the original.

<sup>82</sup>Editor's note: The following sentence has been corrected.

is annihilated by  $\mathcal{I}_{Y_{\mathcal{J}}}$ , the sheaf of commutative groups  $\mathcal{A}_{\mathcal{J}}$  of 4.3.1 identifies with

$$\begin{aligned} & \mathrm{Hom}_{\mathcal{O}_{Y_{\mathcal{J}}}} \left( \mathcal{I}_{Y_{\mathcal{J}}} / \mathcal{I}_{Y_{\mathcal{J}}}^2, \mathcal{I} \otimes_{\mathcal{O}_{S_{\mathcal{J}}}} \mathcal{O}_{Y_{\mathcal{J}}} \right) \\ &= \mathrm{Hom}_{\mathcal{O}_{Y_{\mathcal{J}}}} \left( \mathcal{N}_{Y_{\mathcal{J}}/X_{\mathcal{J}}}, \mathcal{I} \otimes_{\mathcal{O}_{S_{\mathcal{J}}}} \mathcal{O}_{Y_{\mathcal{J}}} \right), \end{aligned}$$

and hence, for every  $i \geq 0$ ,

$$H^i(X_{\mathcal{J}}, \mathcal{A}_{\mathcal{J}}) = H^i \left( Y_{\mathcal{J}}, \mathrm{Hom}_{\mathcal{O}_{Y_{\mathcal{J}}}} \left( \mathcal{N}_{Y_{\mathcal{J}}/X_{\mathcal{J}}}, \mathcal{I} \otimes_{\mathcal{O}_{S_{\mathcal{J}}}} \mathcal{O}_{Y_{\mathcal{J}}} \right) \right).$$

<sup>83</sup> One may now drop the hypothesis that  $Y$  be closed, as follows. First observe that every open subset  $U_{\mathcal{J}}$  of  $X_{\mathcal{J}}$  is obtained by base change from the open subscheme  $U$  of  $X$  having the same underlying topological space as  $U_{\mathcal{J}}$ . Let  $Y_{\mathcal{J}}$  now be a closed subscheme of  $U_{\mathcal{J}}$ , flat over  $S_{\mathcal{J}}$ , and let  $\mathcal{I}_{Y_{\mathcal{J}}}$  be the quasi-coherent ideal of  $\mathcal{O}_{U_{\mathcal{J}}}$  defining  $Y_{\mathcal{J}}$ . If  $Y_{\mathcal{J}}$  lifts to a subscheme  $Y$  of  $X$ , then  $Y$ , having the same underlying topological space as  $Y_{\mathcal{J}}$ , is a closed subscheme of  $U$ . Consequently, the obstruction to lifting  $Y_{\mathcal{J}}$  to a subscheme, flat over  $S$ , of  $X$  or of  $U$  is the same; it lies in

$$H^1 \left( Y_{\mathcal{J}}, \mathrm{Hom}_{\mathcal{O}_{Y_{\mathcal{J}}}} \left( \mathcal{N}_{Y_{\mathcal{J}}/X_{\mathcal{J}}}, \mathcal{I} \otimes_{\mathcal{O}_{S_{\mathcal{J}}}} \mathcal{O}_{Y_{\mathcal{J}}} \right) \right).$$

Finally, return to the notation of §0. Let  $\mathcal{I}$  be a quasi-coherent ideal of  $\mathcal{O}_S$  such that  $\mathcal{J} \subset \mathcal{I}$  and  $\mathcal{I}\mathcal{J} = 0$ , and let  $S_0$  be the closed subscheme of  $S_{\mathcal{J}}$  defined by  $\mathcal{I}$ . For every  $S$ -scheme  $Z$ , write

$$Z_{\mathcal{J}} = Z \times_S S_{\mathcal{J}}, \quad Z_0 = Z \times_S S_0.$$

Since  $\mathcal{J}$  is annihilated by  $\mathcal{I}$ , the notation of 4.4 gives

$$\mathcal{I} \otimes_{\mathcal{O}_{S_{\mathcal{J}}}} \mathcal{O}_{Y_{\mathcal{J}}} = \mathcal{I} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Y_0},$$

and

$$\begin{aligned} & \mathrm{Hom}_{\mathcal{O}_{Y_{\mathcal{J}}}} \left( \mathcal{N}_{Y_{\mathcal{J}}/X_{\mathcal{J}}}, \mathcal{I} \otimes_{\mathcal{O}_{S_{\mathcal{J}}}} \mathcal{O}_{Y_{\mathcal{J}}} \right) \\ &= \mathrm{Hom}_{\mathcal{O}_{Y_0}} \left( \mathcal{N}_{Y_{\mathcal{J}}/X_{\mathcal{J}}} \otimes_{\mathcal{O}_{Y_{\mathcal{J}}}} \mathcal{O}_{Y_0}, \mathcal{I} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Y_0} \right), \end{aligned}$$

and similarly for the other terms. Thus one obtains:

**Proposition 4.5.** *Let  $S$  be a scheme, and let  $S_0$  and  $S_{\mathcal{J}}$  be the closed subschemes defined by quasi-coherent ideals  $\mathcal{I}$  and  $\mathcal{J}$ , where  $\mathcal{I} \supset \mathcal{J}$  and  $\mathcal{I}\mathcal{J} = 0$ . Let  $X$  be an  $S$ -scheme and let  $Y_{\mathcal{J}}$  be a subscheme of  $X_{\mathcal{J}}$ , flat over  $S_{\mathcal{J}}$ . Define the  $\mathcal{O}_{Y_0}$ -module*

$$\mathcal{A}_0 = \mathrm{Hom}_{\mathcal{O}_{Y_0}} \left( \mathcal{N}_{Y_{\mathcal{J}}/X_{\mathcal{J}}} \otimes_{\mathcal{O}_{Y_{\mathcal{J}}}} \mathcal{O}_{Y_0}, \mathcal{I} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Y_0} \right).$$

Then:

- (i) *A subscheme  $Y$  of  $X$ , reducing to  $Y_{\mathcal{J}}$  and flat over  $S$ , exists if and only if the following two conditions hold:*
  - (a) *Such a  $Y$  exists locally on  $X$ .*
  - (b) *A certain obstruction in  $R^1\Gamma(Y_0, \mathcal{A}_0)$  vanishes.*<sup>84</sup>

<sup>83</sup>Editor's note: The discussion that follows has been expanded.

<sup>84</sup>Editor's note: Here  $R^1\Gamma(Y_0, \mathcal{A}_0)$  denotes the "coherent" cohomology group  $H^1(Y_0, \mathcal{A}_0)$  of the  $\mathcal{O}_{Y_0}$ -module  $\mathcal{A}_0$ , in order to distinguish it from the "Hochschild" cohomology groups  $H^i(Y_0, M_0)$ , where  $Y_0$  is an  $S_0$ -group and  $M_0$  an  $\mathcal{O}_{S_0}$ -module, which will be considered beginning in 4.16.

- (ii) Under these conditions, the set of subschemes  $Y$  satisfying the stated requirements is principal homogeneous under the abelian group  $\Gamma(Y_0, \mathcal{A}_0)$ .

**Remark 4.5.1.**<sup>85</sup> It follows from 4.5(ii) that for every pair  $(Y, Y')$  of subschemes<sup>86</sup> of  $X$ , flat over  $S$  and reducing to  $Y_{\mathcal{J}}$ , there is a “deviation”

$$\begin{aligned} d(Y', Y) &\in \Gamma(Y_0, \mathcal{A}_0) \\ &= \operatorname{Hom}_{\mathcal{O}_{Y_0}} \left( \mathcal{N}_{Y_{\mathcal{J}}/X_{\mathcal{J}}} \otimes_{\mathcal{O}_{Y_{\mathcal{J}}}} \mathcal{O}_{Y_0}, \mathcal{J} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Y_0} \right). \end{aligned}$$

With the sign convention adopted in 4.4.1,  $d(Y', Y)$  corresponds to the morphism of  $\mathcal{O}_X$ -modules

$$\mathcal{I}_{Y'} \hookrightarrow \mathcal{O}_X \longrightarrow \mathcal{O}_Y,$$

which takes values in  $\mathcal{J} \mathcal{O}_Y \simeq \mathcal{J} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Y_0}$ , and factors through  $\mathcal{I}_{Y'} = \mathcal{I}_{Y_{\mathcal{J}}}$ , then through  $\mathcal{N}_{Y_{\mathcal{J}}/X_{\mathcal{J}}}$ .

**Remark 4.6.**<sup>87</sup> If  $X$  is flat over  $S$  and  $Y_{\mathcal{J}}$  is locally complete intersection in  $X_{\mathcal{J}}$ , then condition (a) is always satisfied, and every  $Y$  flat over  $S$  lifting  $Y_{\mathcal{J}}$  is locally complete intersection in  $X$ . If, moreover,  $Y_0$  is affine, condition (b) is also satisfied.

**Definition 4.6.1.** (cf. SGA 6, VII 1.1) Let  $B$  be a commutative ring, let  $f : E \rightarrow B$  be a  $B$ -linear morphism, where  $E$  is a free  $B$ -module of finite rank  $d$ , and let  $I = f(E)$ . If a basis of  $E$  is chosen,  $f$  is given by a  $d$ -tuple  $(f_1, \dots, f_d)$  of elements of  $B$ , and  $I$  is the ideal generated by the  $f_i$ . The Koszul complex  $K_{\bullet}(f)$  is the graded  $B$ -module  $\bigwedge_{\bullet}^{\bullet} E$ , equipped with the differential of degree  $-1$

$$x_1 \wedge \cdots \wedge x_i \mapsto \sum_{j=1}^i (-1)^{j-1} f(x_j) x_1 \wedge \cdots \wedge \widehat{x_j} \wedge \cdots \wedge x_i.$$

Thus there is an augmented chain complex, with  $B/I$  in degree  $-1$ ,

$$\cdots \longrightarrow \bigwedge^2 E \longrightarrow E \xrightarrow{f} B \longrightarrow B/I \longrightarrow 0,$$

which is exact in degree 0 by definition, since  $I = f(E)$ . The morphism  $f$  is said to be *regular* if  $K_{\bullet}(f)$  is acyclic in degrees  $> 0$ , that is, if the augmented complex above is a resolution of  $C = B/I$ .

In that case, the proof of SGA 6, VII 1.2(b) shows that the  $C$ -modules  $I^n/I^{n+1}$ , for  $n \in \mathbb{N}$ , are free, with  $I/I^2$  of rank  $d$ .

**Definition 4.6.2.** (cf. SGA 6, VII 1.4) Let  $X$  be a scheme,  $Y$  a subscheme, and  $U$  an open subset of  $X$  such that  $Y$  is a closed subscheme of  $U$ , defined by the quasi-coherent ideal  $\mathcal{I}_Y$ .

The subscheme  $Y$  is said to be *locally complete intersection* in  $X$  if  $Y \hookrightarrow X$  is a regular immersion in the sense of SGA 6, VII 1.4; that is, if for every  $y \in Y$  there exist an affine open neighborhood  $V$  of  $y$  in  $U$ , a finite free  $\mathcal{O}_V$ -module  $\mathcal{E}$ , and a regular morphism  $f : \mathcal{E} \rightarrow \mathcal{O}_V$  with image  $\mathcal{I}_Y|_V$ , such that  $K_{\bullet}(f)$  is a resolution of  $\mathcal{O}_{Y \cap V}$ .

<sup>85</sup>Editor’s note: This remark has been placed here; it replaces Remark 4.7 of the original.

<sup>86</sup>Editor’s note: “closed subschemes” has been corrected to “subschemes.”

<sup>87</sup>Editor’s note: We have retained, for reference, Remark 4.6 of the original, which did not include the definition of “locally complete intersection.” It is followed by the “correct” definition, taken from SGA 6, VII 1.4 (and replacing that of EGA IV<sub>4</sub>, 16.9.2), and by proofs of the three assertions made in the remark.

This implies that the immersion  $Y \hookrightarrow X$  is locally of finite presentation and, by 4.6.1, that the conormal sheaf

$$\mathcal{N}_{Y/X} = \mathcal{I}_Y / \mathcal{I}_Y^2$$

is a finite locally free  $\mathcal{O}_Y$ -module.

**Lemma 4.6.3.**<sup>88</sup> *Let  $A$  be a ring, let  $J$  be a square-zero ideal of  $A$ , put  $\bar{A} = A/J$ , let  $B$  be a flat  $A$ -algebra, let  $E$  be a free  $B$ -module of finite rank, and let  $f : E \rightarrow B$  be a morphism of  $B$ -modules. Suppose that the morphism*

$$g : \bar{E} = E \otimes_A \bar{A} \longrightarrow \bar{B} = B \otimes_A \bar{A}$$

*induced by  $f$  is regular and that  $\bar{B}/g(\bar{E})$  is flat over  $\bar{A}$ . Then  $f$  is regular and  $B/f(E)$  is flat over  $A$ .*

*Proof.* Put

$$C = B/f(E), \quad \bar{C} = C \otimes_A \bar{A} = \bar{B}/g(\bar{E}).$$

First, the  $\bigwedge_B^i E$  are free  $B$ -modules, and hence flat  $A$ -modules, since  $B$  is flat over  $A$ . Since

$$\bigwedge_B^\bullet E \otimes_A \bar{A} \simeq \bigwedge_{\bar{B}}^\bullet \bar{E},$$

there is an exact sequence of complexes

$$0 \longrightarrow J \otimes_A \bigwedge_B^\bullet E \longrightarrow \bigwedge_B^\bullet E \longrightarrow \bigwedge_{\bar{B}}^\bullet \bar{E} \longrightarrow 0.$$

Moreover, since  $J^2 = 0$ , for every  $A$ -module  $M$  one has

$$J \otimes_A M \simeq J \otimes_{\bar{A}} (\bar{A} \otimes_A M).$$

Writing the augmentation arrows as dashed arrows, and denoting the rank of  $E$  by  $d$ , one obtains the following bicomplex, whose rows are exact.

---

<sup>88</sup>Editor's note: To prove the results stated in Remark 4.6, Lemmas 4.6.3 and 4.6.4, Proposition 4.6.5, and Remark 4.6.6 have been added.



$$\begin{array}{ccccccc}
& & 0 & & 0 & & 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & J \otimes_{\bar{A}} \bigwedge_{\bar{B}}^d \bar{E} & \longrightarrow & \bigwedge_{\bar{B}}^d E & \longrightarrow & \bigwedge_{\bar{B}}^d \bar{E} \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
& & \vdots & & \vdots & & \vdots \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & J \otimes_{\bar{A}} \bar{E} & \longrightarrow & E & \longrightarrow & \bar{E} \longrightarrow 0 \\
& & \downarrow \text{id} \otimes g & & \downarrow f & & \downarrow g \\
0 & \longrightarrow & J \otimes_{\bar{A}} \bar{B} & \longrightarrow & B & \longrightarrow & \bar{B} \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
& & J \otimes_{\bar{A}} \bar{C} & \longrightarrow & C & \longrightarrow & \bar{C} \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
& & 0 & & 0 & & 0
\end{array}$$

The right and left columns are also exact, since  $K_{\bullet}(g)$  is a resolution of  $\bar{C}$ , and  $\bar{C}$  is flat over  $\bar{A}$ . The long exact homology sequence associated with the exact sequence of unaugmented complexes therefore shows that  $K_{\bullet}(f)$  is acyclic in degrees  $> 0$ , and in degree 0 gives an exact sequence

$$0 \longrightarrow J \otimes_{\bar{A}} \bar{C} \longrightarrow C \longrightarrow \bar{C} \longrightarrow 0.$$

It follows from the “fundamental criterion of flatness” that  $C$  is flat over  $A$  (cf. [BAC], §III.5, Theorem 1).  $\square$

**Lemma 4.6.4.**<sup>88</sup> *Let  $A$  be a commutative ring, let  $J$  be a nilpotent ideal, and let  $N \subset M$  be  $A$ -modules such that  $M/N$  is flat over  $A$ . If  $x_1, \dots, x_n$  are elements of  $N$  whose images generate the image  $\bar{N}$  of  $N$  in  $M/JM$ , then they generate  $N$ .*

Indeed, let  $N'$  be the submodule of  $N$  generated by the  $x_i$ , and put  $Q = N/N'$ . The morphism  $N' \otimes_A (A/J) \rightarrow \bar{N}$  is surjective. On the other hand, since  $M/N$  is flat over  $A$ , the morphism  $N \otimes_A (A/J) \rightarrow \bar{N}$  is bijective. Thus  $Q \otimes_A (A/J) = 0$ , whence  $Q = 0$  by the “nilpotent Nakayama lemma”:

$$Q = JQ = J^2Q = \dots = 0.$$

We can now prove the following proposition.

**Proposition 4.6.5.**<sup>88</sup> *Let  $S, \mathcal{I}, \mathcal{J}$ , and  $X, Y_{\mathcal{J}}$  be as in 4.5. Suppose in addition that  $X$  is flat over  $S$  and that  $Y_{\mathcal{J}}$  is locally complete intersection in  $X_{\mathcal{J}}$ .*

- (a) Condition (a) of 4.5(i) is then satisfied; moreover, every  $Y$  flat over  $S$  lifting  $Y_{\mathcal{J}}$  is locally complete intersection in  $X$ .
- (b) If, in addition,  $Y_0$  is affine, condition (b) of the cited result is also satisfied.

*Proof.* The first assertion of (a) follows from Lemma 4.6.3, and the second then follows from Lemma 4.6.4. Moreover, the hypothesis implies (cf. 4.6.2) that  $\mathcal{N}_{Y_{\mathcal{J}}/X_{\mathcal{J}}}$  is a finite locally free  $\mathcal{O}_{Y_{\mathcal{J}}}$ -module. Hence the  $\mathcal{O}_{Y_0}$ -module

$$\mathcal{A}_0 = \mathrm{Hom}_{\mathcal{O}_{Y_0}} \left( \mathcal{N}_{Y_{\mathcal{J}}/X_{\mathcal{J}}} \otimes_{\mathcal{O}_{Y_{\mathcal{J}}}} \mathcal{O}_{Y_0}, \mathcal{I} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Y_0} \right)$$

is quasi-coherent (cf. EGA I, 1.3.12), and therefore  $R^1\Gamma(Y_0, \mathcal{A}_0) = 0$  when  $Y_0$  is affine.  $\square$

**Remark 4.6.6.**<sup>88</sup> Let us conclude this paragraph with an example showing that, under the hypotheses of Lemma 4.6.3, a regular sequence  $(\bar{g}_1, \bar{g}_2)$  generating the ideal  $\bar{I} = g(\bar{E})$  need not lift to a regular sequence in  $B$ .

Let  $k$  be a field, put  $\bar{A} = k[X, Y]$ , let  $k\varepsilon$  denote the  $\bar{A}$ -module  $\bar{A}/(X, Y)$ , so that

$$P\varepsilon = P(0, 0)\varepsilon \quad (P \in \bar{A}),$$

and let  $A = \bar{A} \oplus k\varepsilon$ , where  $J = k\varepsilon$  is a square-zero ideal. Thus  $A/J = \bar{A}$ .

The algebra  $B = A \otimes_k k[Z, T]$  is free, and hence flat, over  $A$ , and  $\bar{B} = k[X, Y, Z, T]$ . Put

$$\bar{g}_1 = XZ - YT, \quad \bar{g}_2 = XZ - 1.$$

Since  $\bar{g}_1$  is irreducible,  $\bar{B}/(\bar{g}_1)$  is an integral domain; hence  $(\bar{g}_1, \bar{g}_2)$  is a regular sequence in  $\bar{B}$ , generating the ideal

$$\bar{I} = (XZ - 1, YT - 1).$$

Therefore

$$\bar{C} = \bar{B}/\bar{I} = k[X, Y, X^{-1}, Y^{-1}] = \bar{A}[X^{-1}, Y^{-1}]$$

is a flat  $\bar{A}$ -algebra, and also a flat  $A$ -algebra. But every lift of  $\bar{g}_1$  to  $B$  has the form

$$XZ - YT + \lambda\varepsilon, \quad \lambda \in k[Z, T],$$

and therefore annihilates  $\varepsilon$ .

*Source note.* The Polo-Gille PDF prints  $XY - ZT + \lambda\varepsilon$  for this lift. Since the element being lifted is  $\bar{g}_1 = XZ - YT$ , the displayed lift above restores that same polynomial before adding  $\lambda\varepsilon$ .

**4.7.** Remark 4.7 has been suppressed here and placed in 4.5.1.

**Remark 4.8.0.**<sup>89</sup> Let  $S$  be a scheme,  $S'$  a closed subscheme of  $S$ ,  $X$  an  $S$ -scheme,  $Y$  an  $S$ -subscheme of  $X$ , and put  $X' = X \times_S S'$  and  $Y' = Y \times_S S'$ . Then there is a surjective morphism of  $\mathcal{O}_{Y'}$ -modules

$$\mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y'} \twoheadrightarrow \mathcal{N}_{Y'/X'}.$$

Indeed, after replacing  $X$  by a suitable open subscheme, one may suppose that  $Y$  is closed and is defined by an ideal  $\mathcal{I}_Y$  of  $\mathcal{O}_X$ . The image of  $\mathcal{I}_Y$  in  $\mathcal{O}_{X'}$  is then the ideal  $\mathcal{I}_{Y'}$  defining  $Y'$ , and there is a surjective morphism of  $\mathcal{O}_{Y'}$ -modules

$$\pi : (\mathcal{I}_Y/\mathcal{I}_Y^2) \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y'} \twoheadrightarrow \mathcal{I}_{Y'}/\mathcal{I}_{Y'}^2.$$

<sup>89</sup>Editor's note: This remark, used in the proposition that follows, has been inserted here; it appeared in 4.10 of the original.

Suppose in addition that  $\mathcal{O}_Y = \mathcal{O}_X/\mathcal{I}_Y$  is flat over  $\mathcal{O}_S$ . The natural morphism

$$\mathcal{I}_Y \otimes_{\mathcal{O}_X} \mathcal{O}_{X'} \longrightarrow \mathcal{I}_{Y'}$$

is then bijective (cf. EGA IV<sub>2</sub>, 2.1.8).

There is therefore the following commutative diagram with exact rows:

$$\begin{array}{ccccccc} \mathcal{I}_Y^2 \otimes_{\mathcal{O}_X} \mathcal{O}_{X'} & \longrightarrow & \mathcal{I}_Y \otimes_{\mathcal{O}_X} \mathcal{O}_{X'} & \longrightarrow & (\mathcal{I}_Y/\mathcal{I}_Y^2) \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y'} & \longrightarrow & 0 \\ \text{surj.} \downarrow & & \downarrow \wr & & \pi \downarrow \text{surj.} & & \\ 0 \longrightarrow & \mathcal{I}_{Y'}^2 & \longrightarrow & \mathcal{I}_{Y'} & \longrightarrow & \mathcal{I}_{Y'}/\mathcal{I}_{Y'}^2 & \longrightarrow 0 \end{array}$$

It follows from the snake lemma that<sup>90</sup>

$$\mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y'} \xrightarrow{\sim} \mathcal{N}_{Y'/X'} \quad \text{if } Y \text{ is flat over } S. \quad (4.8.0)$$

**Proposition 4.8.** *Let  $S, S_0, S_{\mathcal{J}}$ , and  $\mathcal{I}, \mathcal{J}$  be as in 4.5.<sup>91</sup> Let  $X$  be an  $S$ -scheme, let  $Y$  be a subscheme of  $X$ , and let  $i : Y \hookrightarrow X$  be the immersion.*

- (i) *For every  $S$ -morphism  $f : T \rightarrow X$  such that  $f_{\mathcal{J}} : T_{\mathcal{J}} \rightarrow X_{\mathcal{J}}$  factors through  $Y_{\mathcal{J}}$ , one can define an obstruction*

$$c(X, Y, f) \in \text{Hom}_{\mathcal{O}_{T_0}}(f_0^*(\mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y_0}), \mathcal{J} \mathcal{O}_T) \quad (*)$$

*whose vanishing is equivalent to the existence of a factorization of  $f$  through  $Y$ .*

- (ii) *Let  $Y'$  be a second subscheme of  $X$ . Suppose that  $Y'_{\mathcal{J}} = Y_{\mathcal{J}}$  and that  $Y$  and  $Y'$  are flat over  $S$ . Then there are isomorphisms (cf. 4.8.0)*

$$\mathcal{J} \mathcal{O}_Y \simeq \mathcal{J} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Y_0} \simeq \mathcal{J} \mathcal{O}_{Y'}, \quad \mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y_{\mathcal{J}}} \xrightarrow{\sim} \mathcal{N}_{Y_{\mathcal{J}}/X_{\mathcal{J}}},$$

*and hence an isomorphism*

$$\begin{aligned} u : \text{Hom}_{\mathcal{O}_{Y_0}}(\mathcal{N}_{Y_{\mathcal{J}}/X_{\mathcal{J}}} \otimes_{\mathcal{O}_{Y_{\mathcal{J}}}} \mathcal{O}_{Y_0}, \mathcal{J} \mathcal{O}_Y) \\ \xrightarrow{\sim} \text{Hom}_{\mathcal{O}_{Y_0}}(\mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y_0}, \mathcal{J} \mathcal{O}_{Y'}). \end{aligned}$$

*Writing  $i' : Y' \hookrightarrow X$  for the canonical immersion and  $d(Y, Y')$  for the deviation of 4.5.1, one has<sup>92</sup>*

$$c(X, Y, i') = u(d(Y, Y')). \quad (**)$$

- (iii) *The canonical morphism*

$$\mathcal{N}_{Y/X} \xrightarrow{D} i^*(\Omega_{X/S}^1)$$

<sup>90</sup>Editor's note: In the original, this was stated in Remark 4.10 under the additional hypothesis that  $Y'$  be locally complete intersection in  $X'$ . Consequently, that hypothesis also appeared in statements 4.12–4.14; it seems in fact superfluous, and has been deleted from those statements.

<sup>91</sup>Editor's note: The hypothesis that  $\mathcal{J}$  be nilpotent has been deleted, since it appears to be superfluous (cf. the proof).

<sup>92</sup>Editor's note: See also 4.27 below.

(cf. SGA 1, II, formula 4.3)<sup>93</sup> induces a morphism

$$D_0 : \mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y_0} \longrightarrow \Omega_{X_0/S_0}^1 \otimes_{\mathcal{O}_{X_0}} \mathcal{O}_{Y_0},$$

and hence, for every  $S$ -morphism  $f : T \rightarrow X$  as in (i), a morphism

$$\begin{aligned} v_{f_0} : \operatorname{Hom}_{\mathcal{O}_{T_0}} \left( f_0^*(\Omega_{X_0/S_0}^1), \mathcal{I} \mathcal{O}_T \right) \\ \longrightarrow \operatorname{Hom}_{\mathcal{O}_{T_0}} \left( f_0^*(\mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y_0}), \mathcal{I} \mathcal{O}_T \right), \\ a \longmapsto a \circ f_0^*(D_0), \end{aligned}$$

where the first group above is  $\operatorname{Hom}_{X^+}(T, L_X)$ , cf. 0.1.5. For  $a \in \operatorname{Hom}_{X^+}(T, L_X)$ , one has

$$c(X, Y, a \cdot f) - c(X, Y, f) = v_{f_0}(a), \quad (***)$$

where  $a \cdot f$  denotes the composite morphism

$$T \xrightarrow{a \times f} L_X \times_{X^+} X \longrightarrow X.$$

We shall prove part (i) of the proposition, leaving it to the reader to (not) verify assertions (ii) and (iii); this verification proceeds by reduction to the affine case and then by comparison of the explicit definitions.<sup>94</sup>

Let us therefore prove (i). The morphism  $f : T \rightarrow X$  defines a morphism of sheaves of rings<sup>95</sup>

$$\phi : \mathcal{O}_X \longrightarrow f_*(\mathcal{O}_T).$$

Let  $U$  be an open subscheme of  $X$  in which  $Y$  is closed. Since  $T$ , respectively  $Y_{\mathcal{I}}$ , has the same underlying space as  $T_{\mathcal{I}}$ , respectively  $Y$ , the continuous map underlying  $f$  sends  $T$  into  $U$ . Since  $U$  is open in  $X$ , the morphism  $\phi$  induces a morphism of sheaves of rings

$$\mathcal{O}_U = \mathcal{O}_X|_U \longrightarrow f_*(\mathcal{O}_T),$$

that is,  $f$  factors through  $U$ .

We may therefore restrict to the case in which  $Y$  is closed, hence defined by a sheaf of ideals  $\mathcal{I}_Y$ . The morphism  $f$  factors through  $Y$  if and only if the composite

$$\mathcal{I}_Y \longrightarrow \mathcal{O}_X \longrightarrow f_*(\mathcal{O}_T)$$

is zero. Since  $f_{\mathcal{I}}$  factors through  $Y_{\mathcal{I}}$ , the composite

$$\mathcal{I}_{Y_{\mathcal{I}}} \longrightarrow \mathcal{O}_{X_{\mathcal{I}}} \longrightarrow f_*(\mathcal{O}_{T_{\mathcal{I}}})$$

is zero. Consider the following commutative diagram, whose first row is exact:

<sup>93</sup>Editor's note: See also EGA IV<sub>4</sub>, 16.4.21. Recall that if  $U$  is an affine open subset of  $X$  such that  $Y \cap U$  is defined by the ideal  $I$  of  $A = \mathcal{O}_X(U)$ , if  $d$  denotes the differential  $A \rightarrow \Gamma(U, \Omega_{X/S}^1)$ , and if  $x \in I$ , then  $D(x + I^2)$  is the element  $d(x) \otimes 1$  of  $\Gamma(U, \Omega_{X/S}^1) \otimes_A (A/I)$ .

<sup>94</sup>Editor's note: These verifications have been carried out below.

<sup>95</sup>Editor's note: On the one hand, the hypothesis that  $\mathcal{I}$  be nilpotent—that is, that  $X_0$  have the same underlying topological space as  $X$ —has been deleted; on the other hand, the sentence that follows has been expanded.

$$\begin{array}{ccccccc}
0 & \longrightarrow & f_*(\mathcal{I}\mathcal{O}_T) & \longrightarrow & f_*(\mathcal{O}_T) & \longrightarrow & f_*(\mathcal{O}_{T_{\mathcal{J}}}) \\
& & \nearrow \phi & & \uparrow \phi & & \uparrow \phi_{\mathcal{J}} \\
& & & & \mathcal{O}_X & \longrightarrow & \mathcal{O}_{X_{\mathcal{J}}} \\
& & & & \uparrow & & \uparrow \\
& & & & \mathcal{I}_Y & \longrightarrow & \mathcal{I}_{Y_{\mathcal{J}}}
\end{array}$$

It follows that  $\phi$  maps  $\mathcal{I}_Y$  into  $f_*(\mathcal{I}\mathcal{O}_T)$ .<sup>96</sup> Since  $\mathcal{J}^2 = 0$ , the  $\mathcal{O}_X$ -module  $f_*(\mathcal{I}\mathcal{O}_T)$ , with its  $\mathcal{O}_X$ -module structure induced by  $\phi$ , is annihilated by  $\mathcal{I}_Y$ . Consequently,  $\phi$  induces a morphism of  $\mathcal{O}_X$ -modules

$$h : i_*(\mathcal{N}_{Y/X}) = \mathcal{I}_Y/\mathcal{I}_Y^2 \longrightarrow f_*(\mathcal{I}\mathcal{O}_T).$$

On the other hand, one has the following cartesian squares:

$$\begin{array}{ccccc}
T_0 & \xrightarrow{f_0} & X_0 & \xleftarrow{i_0} & Y_0 \\
\downarrow \tau_{T_0} & & \downarrow \tau_{X_0} & & \downarrow \tau_{Y_0} \\
T & \xrightarrow{f} & X & \xleftarrow{i} & Y.
\end{array}$$

Here  $\tau_{T_0}, \tau_{X_0}, \tau_{Y_0}$  are the closed immersions obtained by base change from  $S_0 \hookrightarrow S$ . Since  $\mathcal{I}\mathcal{O}_T$  is a quasi-coherent  $\mathcal{O}_T$ -module annihilated by  $\mathcal{I}$ , there is an isomorphism

$$\mathcal{I}\mathcal{O}_T \simeq (\tau_{T_0})_* \tau_{T_0}^*(\mathcal{I}\mathcal{O}_T),$$

and hence

$$f_*(\mathcal{I}\mathcal{O}_T) \simeq (\tau_{X_0})_*(f_0)_* \tau_{T_0}^*(\mathcal{I}\mathcal{O}_T).$$

Thus, by adjunction,  $h$  corresponds to a morphism of  $\mathcal{O}_{T_0}$ -modules

$$h_0 : f_0^* \tau_{X_0}^* i_*(\mathcal{N}_{Y/X}) \longrightarrow \tau_{T_0}^*(\mathcal{I}\mathcal{O}_T).$$

*Source notation note.* The Polo–Gille prose and this  $h_0$ -display print  $i_{T_0}$  for the closed immersion labeled  $\tau_{T_0}$  in the cartesian square and the preceding base-change formulas. We use  $\tau_{T_0}$  consistently.

Now

$$\tau_{X_0}^* i_*(\mathcal{N}_{Y/X}) \simeq (i_0)_* \tau_{Y_0}^*(\mathcal{N}_{Y/X}) = \mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y_0}.$$

Returning to the customary abuse of notation  $\tau_{T_0}^*(\mathcal{I}\mathcal{O}_T) = \mathcal{I}\mathcal{O}_T$ , the morphism  $h_0$  identifies with a morphism of  $\mathcal{O}_{T_0}$ -modules

$$h_0 : f_0^*(\mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y_0}) \longrightarrow \mathcal{I}\mathcal{O}_T,$$

which is the required obstruction  $c(X, Y, f)$ . This proves (i).

<sup>96</sup>Editor's note: The argument that follows has been expanded.

When  $f$  is the immersion  $i' : Y' \hookrightarrow X$ , one sees that  $c(X, Y, i')$  comes from the morphism

$$\mathcal{I}_Y \hookrightarrow \mathcal{O}_X \longrightarrow \mathcal{O}_{Y'},$$

and hence corresponds, by 4.4.1 and 4.5.1, to the class  $d(Y, Y')$ . This proves (ii).

*Source notation note.* At this point the Polo–Gille proof prints  $i' : Y \hookrightarrow X$ , while part (ii) defines  $i'$  as the immersion of  $Y'$ . The latter domain is used here.

Let us prove (iii). First,  $D : \mathcal{N}_{Y/X} \rightarrow i^*(\Omega_{X/S}^1)$  induces a morphism

$$\begin{aligned} D_0 : \tau_{Y_0}^*(\mathcal{N}_{Y/X}) &\longrightarrow \tau_{Y_0}^* i^*(\Omega_{X/S}^1) \\ &= i_0^* \tau_{X_0}^*(\Omega_{X/S}^1). \end{aligned}$$

Since  $X_0 = X \times_S S_0$ , one has

$$\tau_{X_0}^*(\Omega_{X/S}^1) \simeq \Omega_{X_0/S_0}^1 \quad (\text{cf. EGA IV}_4, 16.4.5).$$

This gives the asserted morphism

$$D_0 : \mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y_0} \longrightarrow \Omega_{X_0/S_0}^1 \otimes_{\mathcal{O}_{X_0}} \mathcal{O}_{Y_0}.$$

We shall verify equality  $(***)$  after the following remark.

**Remark 4.9.** The obstruction  $c(X, Y, f)$  is computed locally on  $T$ . Let  $U = \text{Spec}(C)$  be an affine open subset of  $T$  lying over an affine open subset  $\text{Spec}(A)$  of  $X$ , itself lying over an affine open subset  $\text{Spec}(\Lambda)$  of  $S$ . Let  $J \subset I \subset \Lambda$ , respectively  $I_Y \subset A$ , be the ideals corresponding to  $\mathcal{J} \subset \mathcal{I}$ , respectively  $\mathcal{I}_Y$ , put  $B = A/I_Y$ , and let  $\phi : A \rightarrow C$  be the morphism of  $\Lambda$ -algebras corresponding to  $f : T \rightarrow X$ . Since  $f(T_{\mathcal{J}}) \subset Y_{\mathcal{J}}$ , one has  $\phi(I_Y) \subset JC$ , and  $\phi$  therefore induces a morphism of  $\Lambda$ -algebras

$$B \longrightarrow C/JC \longrightarrow C_0 = C/IC.$$

The obstruction  $c = c(X, Y, f)$  is then computed by the following commutative diagram:

$$\begin{array}{ccccc} I_Y & \longrightarrow & A & \xrightarrow{\phi} & C \\ \downarrow & & & & \uparrow \\ I_Y/I_Y^2 & \longrightarrow & I_Y/I_Y^2 \otimes_B C_0 & \xrightarrow{c} & JC \end{array} \quad ,$$

That is, over  $U$ , it is the unique element of

$$\text{Hom}_{C_0}(I_Y/I_Y^2 \otimes_B C_0, JC) = \text{Hom}_{B_0}(I_Y/I_Y^2 \otimes_B B_0, JC)$$

such that, with the evident notation,

$$c(\bar{x} \otimes_B 1) = \phi(x) \quad \text{for every } x \in I_Y.$$

<sup>97</sup> We can now complete the proof of 4.8(iii). Equality  $(***)$  may be checked locally on  $T$ , so we are reduced to the affine situation described above. Let  $d_{A/\Lambda} : A \rightarrow \Omega_{A/\Lambda}^1$  denote the differential. Over  $U$ , the element  $a$  corresponds to an element  $a_U$  of

$$\begin{aligned} &\text{Hom}_{C_0}(\Omega_{A_0/\Lambda_0}^1 \otimes_{A_0} C_0, JC) \\ &\simeq \text{Hom}_{B_0}(\Omega_{A/\Lambda}^1 \otimes_A B_0, JC) \simeq \text{Hom}_A(\Omega_{A/\Lambda}^1, JC). \end{aligned}$$

<sup>97</sup>Editor's note: What follows has been added.

On the one hand,  $v_{f_0}(a)$  corresponds over  $U$  to  $a_U \circ D_0$ , where  $D_0$  is the morphism of  $B$ -modules<sup>98</sup>

$$\begin{aligned} I_Y/I_Y^2 &\longrightarrow \Omega_{A/\Lambda}^1 \otimes_A B_0, \\ x + I_Y^2 &\longmapsto d_{A/\Lambda}(x) \otimes 1. \end{aligned}$$

On the other hand (cf. the proofs of 0.1.8 and 0.1.9), the morphism of  $\Lambda$ -algebras  $\phi' : A \rightarrow C$  corresponding to  $a \cdot f$  differs from  $\phi$  by the  $\Lambda$ -derivation  $A \rightarrow JC$  associated with  $a_U$ ; that is,

$$\phi' = \phi + a_U \circ d_{A/\Lambda} = \phi + a_U \circ (d_{A/\Lambda} \otimes 1).$$

Consequently, writing  $c' = c(X, Y, a \cdot f)$ , for every  $x \in I_Y$ , and writing  $\bar{x}$  for its image in  $I_Y/I_Y^2$ , one has

$$\begin{aligned} (c' - c)(\bar{x} \otimes 1) &= a_U(d_{A/\Lambda}(x) \otimes 1) \\ &= (a_U \circ D_0)(\bar{x}) = v_{f_0}(a)(\bar{x}). \end{aligned}$$

This proves that  $c' - c = v_{f_0}(a)$ .

**4.10.** Remark 4.10 of the original has been suppressed, since it was made obsolete by the addition of Remark 4.8.0.

**4.11.** We now propose to study the following situation. Let  $S, S_{\mathcal{J}}$ , and  $S_0$  be as in 4.8. We have three  $S$ -schemes  $X, X', T$ , a subscheme  $Y$  of  $X$  (respectively a subscheme  $Y'$  of  $X'$ ), and morphisms  $f : T \rightarrow X'$  and  $g : X' \rightarrow X$ .

$$\begin{array}{ccccc} & & Y' & & Y \\ & & \downarrow i' & & \downarrow i \\ T & \xrightarrow{f} & X' & \xrightarrow{g} & X \end{array} .$$

We suppose that, after reduction modulo  $\mathcal{J}$ , this diagram can be completed to a commutative diagram

$$\begin{array}{ccccc} & & Y'_{\mathcal{J}} & \dashrightarrow & Y_{\mathcal{J}} \\ & \nearrow & \downarrow i'_{\mathcal{J}} & & \downarrow i_{\mathcal{J}} \\ T_{\mathcal{J}} & \xrightarrow{f_{\mathcal{J}}} & X'_{\mathcal{J}} & \xrightarrow{g_{\mathcal{J}}} & X_{\mathcal{J}} \end{array} .$$

By 4.8 we therefore have the obstructions

$$\begin{aligned} c(X, Y, g \circ i') &\in \text{Hom}_{\mathcal{O}_{Y'_0}}(i'^*_0 g^*_0(\mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y_0}), \mathcal{J} \mathcal{O}_{Y'}), \\ c(X', Y', f) &\in \text{Hom}_{\mathcal{O}_{T_0}}(f^*_0(\mathcal{N}_{Y'/X'} \otimes_{\mathcal{O}_{Y'}} \mathcal{O}_{Y'_0}), \mathcal{J} \mathcal{O}_T), \\ c(X, Y, g \circ f) &\in \text{Hom}_{\mathcal{O}_{T_0}}(f^*_0 g^*_0(\mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y_0}), \mathcal{J} \mathcal{O}_T), \end{aligned}$$

<sup>98</sup>Editor's note: Cf. Editor's Note 93.

whose relations we seek to compute.<sup>99</sup>

**Lemma 4.12.** *Suppose that  $Y'$  is flat over  $S$ , so that*

$$\mathcal{I} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Y'_0} = \mathcal{I} \mathcal{O}_{Y'}.$$

*Then:*

(i) *There is a natural morphism*

$$\begin{aligned} b_{f_0} : \operatorname{Hom}_{\mathcal{O}_{Y'_0}}(i_0'^* g_0^*(\mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y_0}), \mathcal{I} \mathcal{O}_{Y'}) \\ \longrightarrow \operatorname{Hom}_{\mathcal{O}_{T_0}}(f_0^* g_0^*(\mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y_0}), \mathcal{I} \mathcal{O}_T). \end{aligned}$$

(ii) *There is also a natural morphism, functorial in  $T$ ,*<sup>100</sup>

$$\begin{aligned} a_{g_0}(f_0) : \operatorname{Hom}_{\mathcal{O}_{T_0}}(f_0^*(\mathcal{N}_{Y'/X'} \otimes_{\mathcal{O}_{Y'}} \mathcal{O}_{Y'_0}), \mathcal{I} \mathcal{O}_T) \\ \longrightarrow \operatorname{Hom}_{\mathcal{O}_{T_0}}(f_0^* g_0^*(\mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y_0}), \mathcal{I} \mathcal{O}_T). \end{aligned}$$

*Proof.*<sup>101</sup> First note that, with  $X, X', Y, Y'$  fixed, giving a  $T$  as above is equivalent to giving a morphism  $(f, f_{\mathcal{I}}) : T \rightarrow X' \times_{X'+} Y'^+$ . Put  $Z = X' \times_{X'+} Y'^+$ , and let  $M$  and  $M'$  be the  $Z$ -functors defined, for every  $f : T \rightarrow Z$ , by

$$\begin{aligned} M(T) &= \operatorname{Hom}_{\mathcal{O}_{T_0}}(f_0^* g_0^*(\mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y_0}), \mathcal{I} \mathcal{O}_T), \\ M'(T) &= \operatorname{Hom}_{\mathcal{O}_{T_0}}(f_0^*(\mathcal{N}_{Y'/X'} \otimes_{\mathcal{O}_{Y'}} \mathcal{O}_{Y'_0}), \mathcal{I} \mathcal{O}_T). \end{aligned}$$

<sup>102</sup>

In all cases there is a commutative diagram

$$\begin{array}{ccc} f_0^*(\mathcal{I} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Y'_0}) & \xlongequal{\quad} & \mathcal{I} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{T_0} \\ \downarrow & & \downarrow \\ f_0^*(\mathcal{I} \mathcal{O}_{Y'}) & \dashrightarrow & \mathcal{I} \mathcal{O}_T \end{array}$$

Since  $Y'$  is flat over  $S$ , the left vertical arrow is an isomorphism, and hence we obtain a morphism of  $\mathcal{O}_{T_0}$ -modules

$$f_0^*(\mathcal{I} \mathcal{O}_{Y'}) \longrightarrow \mathcal{I} \mathcal{O}_T.$$

<sup>99</sup>Editor's note: Starting with 4.17, this will be applied when  $X$  is an  $S$ -group,  $g : X \times_S X \rightarrow X$  is multiplication,  $Y$  is a subscheme of  $X$  such that  $Y_{\mathcal{I}}$  is a subgroup of  $X_{\mathcal{I}}$ ,  $Y' = Y \times_S Y$ , and the two morphisms  $Y^3 \rightarrow X^2$  send  $(y_1, y_2, y_3)$  respectively to  $(y_1 y_2, y_3)$  and  $(y_1, y_2 y_3)$ . In that case, comparison of the obstructions above will show that the obstruction to  $Y$  being a subgroup of  $X$  lies in a certain Hochschild cohomology group  $H^2(Y_0, N_0)$ .

<sup>100</sup>Editor's note: We have removed the hypothesis that  $Y'_0$  be locally complete intersection in  $X'_0$ , which is superfluous by 4.8.0. We have also added that  $a_{g_0}(f_0)$  is "functorial in  $T$ ", a fact that plays a crucial role in the proof of 4.17.

<sup>101</sup>Editor's note: The proof has been expanded to make the "functoriality in  $T$ " of  $a_{g_0}$  explicit.

<sup>102</sup>Editor's note: The situation simplifies from 4.16 onward: we shall restrict to schemes flat over  $S$ ,  $Y$  will be a flat  $S$ -group, and  $Y' = Y \times_S Y$ ; this will give  $S_0$ -functors  $N_0$  and  $N'_0$ .



It induces a morphism of abelian groups

$$\mathrm{Hom}_{\mathcal{O}_{T_0}}(f_0^* g_0^*(\mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y_0}), f_0^*(\mathcal{I} \mathcal{O}_{Y'})) \longrightarrow M(T),$$

and, on composing with the morphism

$$M(Y') \longrightarrow \mathrm{Hom}_{\mathcal{O}_{T_0}}(f_0^* g_0^*(\mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y_0}), f_0^*(\mathcal{I} \mathcal{O}_{Y'}))$$

induced by  $f_0^*$ , we obtain the morphism  $b_{f_0} : M(Y') \rightarrow M(T)$ .

Likewise, in all cases there is a diagram

$$\begin{array}{ccc} g_0^*(\mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y_0}) & \xrightarrow{\quad} & \mathcal{N}_{Y'/X'} \otimes_{\mathcal{O}_{Y'}} \mathcal{O}_{Y'_0} \\ \downarrow & & \downarrow \\ g_0^*(\mathcal{N}_{Y_0/X_0}) & \longrightarrow & \mathcal{N}_{Y'_0/X'_0} \end{array}$$

Since  $Y'$  is flat over  $S$ , the second vertical arrow is an isomorphism by 4.8.0. We therefore obtain an  $\mathcal{O}_{Y'_0}$ -morphism

$$i_0'^* g_0^*(\mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y_0}) \longrightarrow \mathcal{N}_{Y'/X'} \otimes_{\mathcal{O}_{Y'}} \mathcal{O}_{Y'_0}.$$

This induces a morphism  $a_{g_0}(\mathrm{id}_{Y'}) : M'(Y') \rightarrow M(Y')$  and, for every  $f : T \rightarrow Z$ , a morphism  $a_{g_0}(f) : M'(T) \rightarrow M(T)$ , such that the following diagram commutes:

$$\begin{array}{ccc} M'(Y') & \xrightarrow{a_{g_0}(\mathrm{id}_{Y'})} & M(Y') \\ \downarrow b'_{f_0} & & \downarrow b_{f_0} \\ M'(T) & \xrightarrow{a_{g_0}(f_0)} & M(T) \end{array}$$

Here  $b'_{f_0}$  is defined in the same way as  $b_{f_0}$ .

Q.E.D.

**Remark 4.12.1.** <sup>103</sup> Let  $M_0$  and  $M'_0$  be the  $Y'_0$ -functors defined, for every  $f : T_0 \rightarrow Y'_0$ , by

$$\begin{aligned} M_0(T) &= \mathrm{Hom}_{\mathcal{O}_{T_0}}(f_0^* g_0^*(\mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y_0}), \mathcal{I} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{T_0}), \\ M'_0(T) &= \mathrm{Hom}_{\mathcal{O}_{T_0}}(f_0^*(\mathcal{N}_{Y'/X'} \otimes_{\mathcal{O}_{Y'}} \mathcal{O}_{Y'_0}), \mathcal{I} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{T_0}). \end{aligned}$$

Note at once that  $Z_0 = Y'_0$ , and that on the category of  $Z$ -schemes  $T$  flat over  $S$ , the functors  $M$  and  $M'$  coincide respectively with  $\prod_{S_0/S} M_0$  and  $\prod_{S_0/S} M'_0$ . In that case,  $b_{f_0}$  is simply the morphism

$$f_0^* : M_0(Y'_0) \longrightarrow M(T_0)$$

induced by  $f_0$ . Moreover, for every morphism  $u : U \rightarrow T$ , if  $h = f \circ u$ , the following diagram commutes:

<sup>103</sup>Editor's note: This remark has been added for use in the proof of 4.17.

$$\begin{array}{ccc}
M'_0(T_0) & \xrightarrow{a_{g_0}(f_0)} & M_0(T_0) \\
u_0^* \downarrow & & \downarrow u_0^* \\
M'_0(U_0) & \xrightarrow{a_{g_0}(h_0)} & M_0(U_0)
\end{array}$$

Thus  $a_{g_0}$  becomes a morphism of functors

$$\prod_{S_0/S} M'_0 \longrightarrow \prod_{S_0/S} M_0.$$

**Proposition 4.13.** *Suppose that  $Y'$  is flat over  $S$ . Then*

$$c(X, Y, g \circ f) = a_{g_0}(c(X', Y', f)) + b_{f_0}(c(X, Y, g \circ i')).$$

Since the definitions of the various obstructions and of the morphisms  $a_{g_0}$  and  $b_{f_0}$  are local, it plainly suffices to verify the formula when all the schemes in question are affine. Write

$$\begin{aligned}
S &= \operatorname{Spec}(\Lambda), & S_{\mathcal{J}} &= \operatorname{Spec}(\Lambda/\mathcal{J}), & S_0 &= \operatorname{Spec}(\Lambda/\mathcal{J}), \\
T &= \operatorname{Spec}(C), & X &= \operatorname{Spec}(A), & Y &= \operatorname{Spec}(A/I_Y) = \operatorname{Spec}(B), \\
X' &= \operatorname{Spec}(A'), & Y' &= \operatorname{Spec}(A'/I_{Y'}) = \operatorname{Spec}(B').
\end{aligned}$$

There is therefore a diagram of rings and ideals <sup>104</sup>

$$\begin{array}{ccccc}
& & B' & & B \\
& & \uparrow \pi' & & \uparrow \pi \\
C & \xleftarrow{f} & A' & \xleftarrow{g} & A \\
& & \uparrow & & \uparrow \\
& & I_{Y'} & & I_Y
\end{array}$$

Let us study the terms of the formula to be proved. In what follows, for  $x \in I_Y$  (respectively  $u \in I_{Y'}$ ), write  $\bar{x}$  (respectively  $\bar{u}$ ) for its image in  $I_Y/I_Y^2$  (respectively  $I_{Y'}/I_{Y'}^2$ ). If  $m$  belongs to a  $\Lambda$ -module  $M$ , write  $\bar{m}_0$  for its image in  $M_0 = M/\mathcal{J}M$ .

We saw that  $c = c(X, Y, g \circ f)$  is the unique  $C_0$ -morphism

$$I_Y/I_Y^2 \otimes_B C_0 \longrightarrow \mathcal{J}C$$

<sup>104</sup>Editor's note: We retain the notation of the original, writing  $f : A' \rightarrow C$  and  $g : A \rightarrow A'$  for the ring homomorphisms corresponding to  $f : T \rightarrow X'$  and  $g : X' \rightarrow X$ . This explains the formula  $c(X, Y, g \circ f)(\bar{x} \otimes 1) = f(g(x))$ , for  $x \in I_Y$ .

such that  $c(\bar{x} \otimes 1) = f(g(x))$  for every  $x \in I_Y$ .

Fix  $x \in I_Y$ . Since  $g_{\mathcal{J}}(Y'_{\mathcal{J}}) \subset Y_{\mathcal{J}}$ , one has  $g(x) \in I_{Y'} + \mathcal{J}A'$ . Write

$$g(x) = x' + \sum_i \lambda_i a'_i, \quad x' \in I_{Y'}, \quad \lambda_i \in \mathcal{J}, \quad a'_i \in A'.$$

Then

$$c(X, Y, g \circ f)(\bar{x} \otimes 1) = f(g(x)) = f(x') + \sum_i \lambda_i f(a'_i). \quad (1)$$

Consider  $a_{g_0}(c(X', Y', f))$ . By the definitions just given, it is defined by the following diagram:

$$\begin{array}{ccccc} & & I_{Y'} & \xrightarrow{f} & C \\ & & \downarrow & & \uparrow \\ I_{Y'_0}/I_{Y'_0}^2 \otimes_{B'_0} C_0 & \xleftarrow{\sim} & I_{Y'}/I_{Y'}^2 \otimes_{B'} C_0 & \xrightarrow{c(X', Y', f)} & JC \\ \uparrow \bar{g}_0 & & \uparrow & \nearrow a_{g_0}(c(X', Y', f)) & \\ I_{Y_0}/I_{Y_0}^2 \otimes_{B_0} C_0 & \xleftarrow{\quad} & I_Y/I_Y^2 \otimes_B C_0 & & \end{array} .$$

Thus  $a_{g_0}(c(X', Y', f))(\bar{x} \otimes 1) = f(u)$ , where  $u \in I_{Y'}$  has image  $\bar{u}$  in  $I_{Y'_0}/I_{Y'_0}^2$  satisfying

$$\bar{u}_0 \otimes 1 = \bar{g}_0(\bar{x}_0) \otimes 1 = g_0(\bar{x}_0) \otimes 1.$$

We may therefore take  $u = x'$ , which gives

$$a_{g_0}(c(X', Y', f))(\bar{x} \otimes 1) = f(x'). \quad (2)$$

Finally consider  $b_{f_0}(c(X, Y, g \circ i'))$ . By hypothesis, the morphism of  $\Lambda_0$ -algebras  $f_0 : A'_0 \rightarrow C_0$  factors through  $B'_0$ . Since  $Y'$  is flat, and hence

$$\mathcal{J} \otimes_{\Lambda_0} B'_0 \xrightarrow{\sim} \mathcal{J}B',$$

we obtain a morphism of  $B'_0$ -modules  $\psi : \mathcal{J}B' \rightarrow \mathcal{J}C$  making the following diagram commute:

$$\begin{array}{ccccc} J \otimes_{\Lambda_0} A'_0 & \xrightarrow{\text{id} \otimes \pi'} & J \otimes_{\Lambda_0} B'_0 & \xrightarrow{\text{id} \otimes f_0} & J \otimes_{\Lambda_0} C_0 \\ \downarrow & & \downarrow \wr & & \downarrow \\ JA' & \xrightarrow{\pi'} & JB' & \xrightarrow{\psi} & JC. \end{array}$$

Let

$$\varphi : \mathcal{J} B' \otimes_{B'_0} C_0 \longrightarrow \mathcal{J} C$$

be the morphism of  $C_0$ -modules induced by  $\psi$ . Then, for every  $a' \in A'$  and  $\lambda \in \mathcal{J}$ ,

$$\varphi(\lambda \pi'(a') \otimes 1) = \lambda f(a'). \quad (\dagger)$$

The element  $b_{f_0}(c(X, Y, g \circ i'))$  is then defined by the commutative diagram

$$\begin{array}{ccc}
 I_Y & \xrightarrow{\pi' \circ g} & B' \\
 \downarrow & & \uparrow \\
 I_Y/I_Y^2 \otimes_B B'_0 & \xrightarrow{c(X, Y, g \circ i')} & JB' \\
 \downarrow & & \downarrow \\
 I_Y/I_Y^2 \otimes_B C_0 & \xrightarrow{\quad} & JB' \otimes_{B'_0} C_0 \\
 & \searrow b_{f_0}(c(X, Y, g \circ i')) & \downarrow \phi \\
 & & JC.
 \end{array}$$

Consequently,

$$b_{f_0}(c(X, Y, g \circ i'))(\bar{x} \otimes 1) = \varphi \left( \sum_i \lambda_i \pi'(a'_i) \otimes 1 \right) = \sum_i \lambda_i f(a'_i), \quad (3)$$

the last equality following from  $(\dagger)$ . Comparison of the three explicit formulas (1), (2), and (3) gives the required formula.

**Corollary 4.14.** *Let  $Y, Y'$  be two flat subschemes of  $X$ , both reducing to  $Y_{\mathcal{J}}$ , and suppose that  $Y_0$  is locally complete intersection in  $X_0$ . If  $f : T \rightarrow X$  is an  $S$ -morphism such that  $f_{\mathcal{J}}$  factors through  $Y_{\mathcal{J}} \rightarrow X_{\mathcal{J}}$ , then*

$$c(X, Y, f) - c(X, Y', f) = b_{f_0}(d(Y, Y')).$$

Indeed, applying the preceding formula to the diagram

$$\begin{array}{ccccc}
 & & Y' & & Y \\
 & & \downarrow i' & & \downarrow i \\
 T & \xrightarrow{f} & X & \xrightarrow{\text{id}} & X
 \end{array}$$

we find  $c(X, Y, f) - c(X, Y', f) = b_{f_0}(c(X, Y, i'))$ . Moreover, by 4.8(ii),  $c(X, Y, i') = d(Y, Y')$ .

**Proposition 4.15.** *Let  $X$  be an  $S$ -group smooth over  $S$ , and let  $Y$  be a sub- $S$ -group flat and locally of finite presentation over  $S$ . Then  $Y$  is locally complete intersection in  $X$  (cf. 4.6.2).*

*Proof.*<sup>105</sup> We show that the immersion  $Y \rightarrow X$  is regular in the sense of EGA IV<sub>4</sub>, 16.9.2. By EGA IV<sub>4</sub>, 19.5.1, this implies that it is regular also in the sense of 4.6.2 (indeed, by loc. cit., the two definitions are equivalent when  $S$  is locally noetherian). Thus “regular immersion” will here mean regular in the sense of EGA IV<sub>4</sub>, 16.9.2. Since  $X$  and  $Y$  are flat and locally of finite presentation over  $S$ , EGA IV<sub>4</sub>, 19.2.4 reduces the assertion to proving that  $Y_s \rightarrow X_s$  is a regular immersion for every  $s \in S$ . By EGA IV<sub>4</sub>, 19.1.5(ii), it is enough to check this on the geometric fibers of  $S$ , and hence when  $S$  is the spectrum of an algebraically closed field  $k$ .

By Exposé VI<sub>A</sub>, 3.2, the quotient  $X/Y$  exists and is smooth, the morphism  $\pi : X \rightarrow X/Y$  is flat, and the following square is cartesian:

$$\begin{array}{ccc} Y & \xrightarrow{f} & X \\ \downarrow & & \downarrow \pi \\ \bar{e} & \xrightarrow{i} & X/Y \end{array}$$

Here  $\bar{e}$  is the image in  $X/Y$  of the unit point of  $X$ . By flat base change (cf. EGA IV<sub>4</sub>, 19.1.5(iii)), it therefore suffices to show that  $i$  is a regular immersion. This is immediate: the noetherian local ring  $\mathcal{O}_{X/Y, \bar{e}}$  is smooth (hence regular), so its maximal ideal is generated by a regular sequence.

**4.16.**<sup>106</sup> Let  $X$  be an  $S$ -group smooth over  $S$ , and denote its group law by  $\mu : X \times_S X \rightarrow X$ . Let  $Y_{\mathcal{J}}$  be a sub- $S_{\mathcal{J}}$ -group of  $X_{\mathcal{J}}$ , flat and locally of finite presentation over  $S_{\mathcal{J}}$ . By 4.15,  $Y_{\mathcal{J}}$  is locally complete intersection in  $X_{\mathcal{J}}$ . Hence, by 4.6.5, every flat<sup>107</sup>  $S$ -scheme  $Y$  lifting

<sup>105</sup>Editor’s note: We have added to the statement the hypothesis that  $Y$  be locally of finite presentation over  $S$ , and supplied the proof that follows, which is more direct than the one sketched in the original. For completeness, we also spell out the latter. As in the proof above, first reduce to the case  $S = \text{Spec}(k)$ , where  $k$  is algebraically closed. By EGA IV<sub>4</sub>, 16.9.10 and 19.3.2, it suffices to see that, for every  $y \in Y$ , the completion of the local ring  $\mathcal{O}_{Y, y}$  is a quotient of a complete noetherian local ring by a regular sequence. By loc. cit., 19.3.3, the set of  $y \in Y$  with this property is an open subset  $U \subset Y$ . Since  $Y$  is of finite type over  $k$ , it suffices to show that  $U$  contains every closed point. Because  $Y$  is a  $k$ -group, a translation argument reduces this to the completion of  $\mathcal{O}_{Y, e}$ , that is, to the formal group  $\hat{Y}$  corresponding to  $Y$  (cf. Exposé VII<sub>B</sub>). Since  $X$  is smooth, the affine algebra  $\mathcal{A}(\hat{X})$  is a power series algebra  $k[[X_1, \dots, X_n]]$ . The Dieudonné structure theorem then shows that  $\mathcal{A}(\hat{Y})$  is isomorphic to a quotient

$$k[[X_1, \dots, X_{r+s}]]/(X_1^{p^{n_1}}, \dots, X_r^{p^{n_r}}),$$

where  $p$  is the characteristic exponent of  $k$  and  $r + s \leq n$ ; cf. Exposé VII<sub>B</sub>, Remark 5.5.2(b).

<sup>106</sup>Editor’s note: We have reorganized 4.16 by collecting here, on the one hand, the hypotheses stated at the end of 4.15 and, on the other, the definition of the obstruction  $DY$ .

<sup>107</sup>Editor’s note: The original has been corrected by adding the word “flat”.

$Y_{\mathcal{J}}$  is locally complete intersection in  $X$ . For such a  $Y$ , 4.8.0 gives

$$\mathcal{N}_{Y/X} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y_0} = \mathcal{N}_{Y_0/X_0} = \mathcal{N}_{Y_{\mathcal{J}}/X_{\mathcal{J}}} \otimes_{\mathcal{O}_{Y_{\mathcal{J}}}} \mathcal{O}_{Y_0}. \quad (4.16.1)$$

On the other hand, let  $\varepsilon_0 : S_0 \rightarrow Y_0$  be the unit section of  $Y_0$ , and let  $\mathbf{n}_{Y_0/X_0}$  be the quasi-coherent  $\mathcal{O}_{S_0}$ -module

$$\mathbf{n}_{Y_0/X_0} = \varepsilon_0^*(\mathcal{N}_{Y_0/X_0}).$$

Since  $Y_0$  and  $X_0$  are  $S_0$ -groups, it is immediate that  $\mathcal{N}_{Y_0/X_0}$  is invariant under, say, left translations by  $Y_0$ . Hence <sup>108</sup> it is the inverse image of  $\mathbf{n}_{Y_0/X_0}$  under  $Y_0 \rightarrow S_0$ ; that is,

$$\mathcal{N}_{Y_0/X_0} = \mathbf{n}_{Y_0/X_0} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Y_0}. \quad (4.16.2)$$

Taking (4.16.1) and (4.16.2) into account, we deduce first from 4.5 that the set of subschemes  $Y$  of  $X$ , flat over  $S$  and lifting  $Y_{\mathcal{J}}$ , is either empty or principal homogeneous under

$$\mathrm{Hom}_{\mathcal{O}_{Y_0}}(\mathbf{n}_{Y_0/X_0} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Y_0}, \mathcal{J} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Y_0}). \quad (4.16.3)$$

On the other hand, 4.8(i) shows that, for every such  $Y$  and every  $S$ -morphism  $f : T \rightarrow X$  such that  $f_{\mathcal{J}} : T_{\mathcal{J}} \rightarrow X_{\mathcal{J}}$  factors through  $Y_{\mathcal{J}}$ , the obstruction  $c(X, Y, f)$  to  $f$  factoring through  $Y$  belongs to

$$\mathrm{Hom}_{\mathcal{O}_{T_0}}(\mathbf{n}_{Y_0/X_0} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{T_0}, \mathcal{J} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{T_0}).$$

If  $T$  is also flat over  $S$ , this last group is equal to

$$\mathrm{Hom}_{\mathcal{O}_{T_0}}(\mathbf{n}_{Y_0/X_0} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{T_0}, \mathcal{J} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{T_0}).$$

This leads us to introduce the following group functor  $N_0$ .

**Definition 4.16.1.** *Let  $N_0$  be the  $S_0$ -functor in commutative groups defined, for every  $Z \in \mathrm{Ob}(\mathbf{Sch})_{/S_0}$ , by*

$$\mathrm{Hom}_{S_0}(Z, N_0) = \mathrm{Hom}_{\mathcal{O}_Z}(\mathbf{n}_{Y_0/X_0} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_Z, \mathcal{J} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_Z). \quad (*)$$

*Then the set of subschemes  $Y$  of  $X$ , flat over  $S$  and lifting  $Y_{\mathcal{J}}$ , is either empty or principal homogeneous under*

$$\mathrm{Hom}_{S_0}(Y_0, N_0) = C^1(Y_0, N_0).$$

For every such  $Y$ , consider the diagram

$$\begin{array}{ccc} Y \times_S Y & & Y \\ \downarrow (i, i) & & \downarrow i \\ X \times_S X & \xrightarrow{\mu} & X \end{array}$$

---

<sup>108</sup>Editor's note: See 4.25 below.

and let

$$DY = c(X, Y, \mu \circ (i, i))$$

be the obstruction to  $\mu \circ (i, i)$  factoring through  $Y$ , that is, to  $Y$  being stable under the group law of  $X$ . By what precedes,  $DY$  is an element of

$$N_0(Y_0 \times_{S_0} Y_0) = C^2(Y_0, N_0).$$

**Lemma 4.17.**<sup>109</sup> *Let  $X$  be an  $S$ -group smooth over  $S$ , and let  $Y_{\mathcal{J}}$  be a sub- $S_{\mathcal{J}}$ -group of  $X_{\mathcal{J}}$ , flat and locally of finite presentation over  $S_{\mathcal{J}}$ . For every subscheme  $Y$  of  $X$ , flat over  $S$  and lifting  $Y_{\mathcal{J}}$ , consider the obstruction defined in 4.16.1:*

$$DY \in \text{Hom}_{S_0}(Y_0 \times_{S_0} Y_0, N_0) = C^2(Y_0, N_0).$$

- (i) *For  $Y$  to be a sub- $S$ -group of  $X$ , it is necessary and sufficient that  $DY = 0$ .*
- (ii) *If  $Y_0$  is made to act on  $N_0$ , by functoriality, through the inner automorphisms of  $Y_0$ , then  $DY \in Z^2(Y_0, N_0)$ .*
- (iii) *If  $Y$  and  $Y'$  are two subschemes of  $X$ , flat over  $S$ , lifting  $Y_{\mathcal{J}}$  (so that the deviation  $d(Y, Y') \in C^1(Y_0, N_0)$  is defined; cf. 4.5.1), then*

$$DY' - DY = \partial^1 d(Y, Y').$$

110

We prove these assertions in succession.

**4.18. — Proof of 4.17(i).** By definition,  $DY = 0$  if and only if  $Y$  is stable under the group law of  $X$ . Thus  $DY = 0$  if  $Y$  is a subgroup of  $X$ . Conversely, if  $DY = 0$ , then  $Y$  is endowed with the induced law  $\mu_Y$ , which is associative and reduces modulo  $\mathcal{J}$  to the group law on  $X_{\mathcal{J}}$ . Since  $Y$  is flat and locally of finite presentation over  $S$ , it follows from 3.3 that  $\mu_Y$  is a group law.

**4.19. — Proof of 4.17(ii).** We compare the two values of  $u = c(X, Y, \mu^2 \circ (i, i, i))$  computed in the following two diagrams  $(D_j)$ ,  $j = 1, 2$ :

$$(D_j) \quad \begin{array}{ccccc} Y \times_S Y \times_S Y & & Y \times_S Y & & Y \\ \downarrow (i, i, i) & & \downarrow (i, i) & & \downarrow i \\ X \times_S X \times_S X & \xrightarrow{f_j} & X \times_S X & \xrightarrow{\mu} & X \end{array}$$

Here  $f_1 = (1, \pi)$ ,  $f_2 = (\pi, 1)$ , and  $\mu^2$  denotes the morphism

$$\mu \circ f_1 = \mu \circ f_2 : X \times_S X \times_S X \longrightarrow X.$$

<sup>111</sup> Put

$$\mu_Y = \mu \circ (i, i), \quad f_j^Y = f_j \circ (i, i, i), \quad \mu^{2,Y} = \mu^2 \circ (i, i, i).$$

<sup>109</sup>Editor's note: Sections 4.17 and 4.18 have been modified to take account of the additions made in 4.16.

<sup>110</sup>Editor's note: The original has  $DY' - DY = -\partial d(Y, Y')$ , but the operator  $\partial$  there is the negative of the differential  $\partial^1$  defined in I, 5.1.

<sup>111</sup>Editor's note: The notation has been slightly modified, and the beginning of the argument has been expanded.

For  $j = 1, 2$ , let  $a_j$  and  $b_j$  be the morphisms

$$a_j = a_{\mu_0}((f_j^Y)_0), \quad b_j = b_{(f_j^Y)_0},$$

associated by Lemma 4.12 with the pair of morphisms  $(f_j^Y, \mu)$ . Thus

$$\begin{cases} a_j : \text{Hom}_{\mathcal{O}_{Y_0^3}} \left( (f_j^Y)_0^* \mathcal{N}_{Y_0 \times Y_0 / X_0 \times X_0}, \mathcal{I} \mathcal{O}_{Y_0^3} \right) \longrightarrow \\ \quad \text{Hom}_{\mathcal{O}_{Y_0^3}} \left( (\mu^{2,Y})_0^* \mathcal{N}_{Y_0 / X_0}, \mathcal{I} \mathcal{O}_{Y_0^3} \right), \\ b_j : \text{Hom}_{\mathcal{O}_{Y_0^2}} \left( (\mu_Y)_0^* \mathcal{N}_{Y_0 / X_0}, \mathcal{I} \mathcal{O}_{Y_0^2} \right) \longrightarrow \\ \quad \text{Hom}_{\mathcal{O}_{Y_0^3}} \left( (\mu^{2,Y})_0^* \mathcal{N}_{Y_0 / X_0}, \mathcal{I} \mathcal{O}_{Y_0^3} \right). \end{cases} \quad (\dagger)$$

Since

$$\mathcal{N}_{Y_0 \times Y_0 / X_0 \times X_0} \simeq \text{pr}_1^* \mathcal{N}_{Y_0 / X_0} \oplus \text{pr}_2^* \mathcal{N}_{Y_0 / X_0}$$

(because  $X_0$  and  $Y_0$  are flat over  $S_0$ ), and

$$\mathcal{N}_{Y_0 / X_0} \simeq \mathbf{n}_{Y_0 / X_0} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Y_0},$$

we have

$$(f_j^Y)_0^* \mathcal{N}_{Y_0 \times Y_0 / X_0 \times X_0} \simeq (\mathbf{n}_{Y_0 / X_0} \oplus \mathbf{n}_{Y_0 / X_0}) \otimes \mathcal{O}_{Y_0^3},$$

and likewise

$$(\mu^{2,Y})_0^* \mathcal{N}_{Y_0 / X_0} \simeq \mathbf{n}_{Y_0 / X_0} \otimes \mathcal{O}_{Y_0^3},$$

$$(\mu_Y)_0^* \mathcal{N}_{Y_0 / X_0} \simeq \mathbf{n}_{Y_0 / X_0} \otimes \mathcal{O}_{Y_0^2}.$$

Moreover, since  $Y_0^2$  and  $Y_0^3$  are flat over  $S_0$ ,  $(\dagger)$  may be rewritten as

$$\begin{cases} a_j : \text{Hom}_{S_0}(Y_0^3, N_0 \oplus N_0) \longrightarrow \text{Hom}_{S_0}(Y_0^3, N_0), \\ b_j : \text{Hom}_{S_0}(Y_0^2, N_0) \longrightarrow \text{Hom}_{S_0}(Y_0^3, N_0). \end{cases} \quad (\ddagger)$$

Applying 4.13 twice to  $c(X, Y, \mu^{2,Y}) = u$ , we obtain

$$\begin{aligned} a_1(c(X^2, Y^2, f_1)) + b_1(c(X, Y, \mu_Y)) &= u \\ &= a_2(c(X^2, Y^2, f_2)) + b_2(c(X, Y, \mu_Y)). \end{aligned}$$

Now  $c(X, Y, \mu_Y) = DY$ , and, since  $f_1 = (1, \mu)$  and  $f_2 = (\mu, 1)$ , we have, with the evident notation,

$$c(X^2, Y^2, f_1) = (0, DY), \quad c(X^2, Y^2, f_2) = (DY, 0).$$

Consequently,

$$u = a_1((0, DY)) + b_1(DY) = a_2((DY, 0)) + b_2(DY).$$

The first point to note is that  $b_j$  is simply  $\text{Hom}_{S_0}((f_j^Y)_0, N_0)$ , that is, the morphism deduced from  $(f_j^Y)_0$  by functoriality. The preceding identity therefore becomes

$$\begin{aligned} a_1((0, DY)) - \text{Hom}((\mu, 1), N_0)(DY) + \text{Hom}((1, \mu), N_0)(DY) \\ - a_2((DY, 0)) = 0. \end{aligned}$$

The two middle terms are the  $DY(xy, z)$  and  $DY(x, yz)$  parts of the formula for the 2-coboundary. It remains only to identify the other two terms.



We first compute the map  $a_j$ . It is obtained, by pullback along  $(f_j^Y)_0$ , from the morphism of  $\mathcal{O}_{Y_0^2}$ -modules

$$P : \mathbf{n}_{Y_0/X_0} \otimes \mathcal{O}_{Y_0^2} \longrightarrow (\mathbf{n}_{Y_0/X_0} \oplus \mathbf{n}_{Y_0/X_0}) \otimes \mathcal{O}_{Y_0^2}$$

induced by multiplication in  $Y_0$ . Consider the vector bundle  $V = \mathbf{V}(\mathbf{n}_{Y_0/X_0})$ . By duality,  $P$  gives

$$\mathbf{V}(P) : V \times_{S_0} V \times_{S_0} Y_0 \times_{S_0} Y_0 \longrightarrow V \times_{S_0} Y_0 \times_{S_0} Y_0,$$

which is expressed set-theoretically by

$$\mathbf{V}(P)(u, v, a, b) = (u + \text{Ad}(a)v, ab, b).$$

<sup>112</sup> This is proved exactly as is the corresponding fact for Lie algebras, that is, for the module  $\omega_{Y_0/S_0}^1$ . First observe that, by functoriality in  $Y_0$ ,  $V$  is endowed with a group structure in the category of vector bundles over  $S_0$ . By the lemma already used for Lie algebras (Exposé II, 3.10), this structure coincides with the group structure underlying its  $\mathcal{O}_S$ -module structure. Next,

$$\mathbf{V}(\mathbf{n}_{Y_0/X_0} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Y_0}) = \mathbf{V}(\mathcal{N}_{Y_0/X_0})$$

is itself endowed with an  $S_0$ -group structure, namely the semidirect product of the structure on  $V$  by that on  $Y_0$ . It remains only to identify the operations of  $Y_0$  on  $V$  to obtain the desired formula.

We now compute the two remaining terms. Consider first  $a_1((0, DY))$ . It is computed by the following diagram, in which  $\mathbf{n}$  denotes  $\mathbf{n}_{Y_0/X_0}$ :

$$\begin{array}{ccc} \mathbf{n} \otimes \mathcal{O}_{Y_0^2} & \xrightarrow{P} & (\mathbf{n} + \mathbf{n}) \otimes \mathcal{O}_{Y_0^2} \\ \downarrow (f_1^Y)_0^* & & \downarrow (f_1^Y)_0^* \\ \mathbf{n} \otimes \mathcal{O}_{Y_0^3} & & (\mathbf{n} + \mathbf{n}) \otimes \mathcal{O}_{Y_0^3} \\ & \searrow a_1((0, DY)) & \downarrow (0, DY) \\ & & \mathcal{J} \otimes \mathcal{O}_{Y_0^3} \end{array} .$$

Regard the vector bundles defined by these modules as schemes over  $S_0$ , and take points with values in an arbitrary scheme. If  $(u, x, y, z)$  denotes a point of  $\mathbf{V}(\mathcal{J}) \times Y_0^3$ , we obtain the following diagram:

<sup>112</sup>Editor's note: The entries  $a, b$  have been replaced by  $ab, b$  to show that  $\mathbf{V}(P)$  is obtained, by pullback to  $Y_0^2$ , from the multiplication morphism  $V_{Y_0} \times_{S_0} V_{Y_0} \rightarrow V_{Y_0}$ .

$$\begin{array}{ccc}
(\mathrm{Ad}(x)\mathrm{DY}_{y,z}(u), x, yz) & \longleftarrow & (0 + \mathrm{DY}_{y,z}(u), x, yz) \\
\uparrow & & \uparrow \\
& & (0 + \mathrm{DY}_{y,z}(u), x, y, z) \\
& & \uparrow \\
(\mathrm{Ad}(x)\mathrm{DY}_{y,z}(u), x, y, z) & \longleftarrow & (u, x, y, z)
\end{array} .$$

Thus

$$a_1((0, DY))(x, y, z) = \mathrm{Ad}(x)DY(y, z),$$

which is indeed the first term of the coboundary. Likewise,  $a_2((DY, 0))(x, y, z) = DY(x, y)$ , and therefore

$$\begin{aligned}
0 &= \mathrm{Ad}(x)DY(y, z) - DY(xy, z) + DY(x, yz) - DY(x, y) \\
&= (\partial^2 DY)(x, y, z).
\end{aligned}$$

113

**4.20. — Proof of 4.17(iii).** <sup>114</sup> We compare the two values of  $v = c(X, Y, \mu \circ (i', i'))$  computed in the following two diagrams:

$$\begin{array}{ccc}
(*) & & \begin{array}{ccc} & Y' & Y \\ & \downarrow i' & \downarrow i \\ Y' \times_S Y' & \xrightarrow{\mu \circ (i', i')} & X = X \end{array} \\
& & \\
(\dagger) & & \begin{array}{ccc} & Y \times_S Y & Y \\ & \downarrow (i, i) & \downarrow i \\ Y' \times_S Y' & \xrightarrow{(i', i')} X \times_S X \xrightarrow{\mu} & X. \end{array}
\end{array}$$

Let  $f = \mu \circ (i', i')$ . Diagram  $(*)$  gives

$$v = DY' + f_0^*(c(X, Y, i')). \quad (1)$$

Now  $Y'_0 = Y_0$ , and  $f_0$  is the multiplication  $Y_0^2 \rightarrow Y_0$ ; hence

$$f_0^*(c(X, Y, i'))(x_0, y_0) = c(X, Y, i')(x_0 y_0). \quad (2)$$

Put  $c = c(X, Y, i')$ . Under the identification  $N'_0 \simeq N_0 \oplus N_0$ , the element

$$c(X \times_S X, Y \times_S Y, (i', i'))$$

<sup>113</sup>Editor's note: The signs have been changed to make them compatible with I, 5.1.

<sup>114</sup>Editor's note: The argument that follows has been expanded from the original.

identifies with the pair  $(c, c)$ . Thus, putting  $h = (i', i')$ , diagram  $(\dagger)$  gives

$$v = h_0^*(DY) + a_{\mu_0}(c, c). \quad (3)$$

Since  $h_0$  is the identity map on  $Y_0^2$ , we have  $h_0^*(DY) = DY$ . Finally, by the preceding calculation of  $a_{\mu_0}$ , for every  $S' \rightarrow S$  and every  $x_0, y_0 \in Y_0(S'_0)$ ,

$$a_{\mu_0}(c, c)(x_0, y_0) = c(x_0) + \text{Ad}(x_0)(c(y_0)). \quad (4)$$

It follows that

$$\begin{aligned} (DY' - DY)(x_0, y_0) &= \text{Ad}(x_0)(c(X, Y, i')(y_0)) - c(X, Y, i')(x_0 y_0) + c(X, Y, i')(x_0) \\ &= (\partial^1 c(X, Y, i'))(x_0, y_0). \end{aligned} \quad (5)$$

Since  $c(X, Y, i') = d(Y, Y')$  (cf. 4.8(iii)), this proves that

$$DY' - DY = \partial^1 d(Y, Y').$$

**Theorem 4.21.** *Let  $S$  be a scheme and let  $\mathcal{I}$  and  $\mathcal{J}$  be two ideals<sup>115</sup> on  $S$ , such that  $\mathcal{I} \supset \mathcal{J}$  and  $\mathcal{I}\mathcal{J} = 0$ . Let  $X$  be an  $S$ -group smooth over  $S$ , and let  $Y_{\mathcal{J}}$  be a sub- $S_{\mathcal{J}}$ -group of  $X_{\mathcal{J}}$ , flat and locally of finite presentation over  $S_{\mathcal{J}}$ . Consider the  $S_0$ -functor of commutative groups  $N_0$  defined by*

$$\begin{aligned} \text{Hom}_{S_0}(T, N_0) &= \text{Hom}_{\mathcal{O}_T} \left( \mathbf{n}_{Y_0/X_0} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_T, \mathcal{J} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_T \right), \\ &\quad T \in \text{Ob}(\mathbf{Sch}/S_0), \end{aligned}$$

on which  $Y_0$  acts through the inner automorphisms of  $X_0$ .

(i) *For there to exist a sub- $S$ -group of  $X$ , flat over  $S$ , which reduces to  $Y_{\mathcal{J}}$ , it is necessary and sufficient that the following two conditions hold:*

(i<sub>1</sub>) *There exists a subscheme  $Y$  of  $X$ , flat over  $S$ , lifting  $Y_{\mathcal{J}}$ . This condition is automatically satisfied when  $Y_0$  is affine (cf. 4.6.5).*

(i<sub>2</sub>) *A certain canonical obstruction, an element of  $H^2(Y_0, N_0)$ , vanishes.*

(ii) *If the conditions in (i) hold, the set of sub- $S$ -groups  $Y$  of  $X$ , flat over  $S$  and reducing to  $Y_{\mathcal{J}}$ , is a principal homogeneous set under  $Z^1(Y_0, N_0)$ .<sup>116</sup>*

Indeed, condition (i<sub>1</sub>) is necessary. Suppose that it holds, and let  $Y$ , flat over  $S$ , lift  $Y_{\mathcal{J}}$ . We must find a  $Y'$ , also flat over  $S$  and lifting  $Y_{\mathcal{J}}$ , such that  $DY' = 0$ ,<sup>117</sup> cf. 4.17(i). By 4.17(iii), this amounts to finding an element  $d(Y', Y) \in C^1(Y_0, N_0)$  such that

$$DY = \partial^1 d(Y', Y).$$

<sup>118</sup> Let  $c \in H^2(Y_0, N_0)$  be the class represented by  $DY$ , which is a cocycle by 4.17(ii). By 4.17(iii), this class does not depend on the choice of  $Y$ , and its vanishing is necessary and sufficient for the existence of  $d(Y', Y)$  satisfying the preceding equation. This proves (i).

If  $Y$  has now been chosen so that  $DY = 0$ , the equation to be solved becomes  $\partial^1 d(Y', Y) = 0$ , which proves (ii).

<sup>115</sup>Editor's note: The hypothesis that  $\mathcal{J}$  be nilpotent, which appears superfluous, has been removed.

<sup>116</sup>Editor's note: The question whether this set, modulo conjugation by the elements  $x \in X(S)$  inducing the identity element of  $X(S_{\mathcal{J}})$ , is a principal homogeneous set under  $H^1(Y_0, N_0)$ , occupies 4.23–4.36.

<sup>117</sup>Editor's note: The expression  $\partial DY'$  in the original has been corrected to  $DY'$ .

<sup>118</sup>Editor's note: See the editor's note to 4.17(iii).

**Remark 4.22.** Retain the notation of 4.21. By 4.15,  $Y_0$  is locally a complete intersection in  $X_0$ ; hence  $\mathcal{N}_{Y_0/X_0}$  is a finite locally free  $\mathcal{O}_{Y_0}$ -module, and therefore

$$\mathbf{n}_{Y_0/X_0} = \varepsilon_0^*(\mathcal{N}_{Y_0/X_0})$$

is a finite locally free  $\mathcal{O}_{S_0}$ -module. Thus, writing

$$\mathbf{n}_{Y_0/X_0}^\vee = \mathrm{Hom}_{\mathcal{O}_{Y_0}}(\mathbf{n}_{Y_0/X_0}, \mathcal{O}_{Y_0}),$$

we have

$$\begin{aligned} \mathrm{Hom}_{\mathcal{O}_T} \left( \mathbf{n}_{Y_0/X_0} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_T, \mathcal{J} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_T \right) \\ \simeq \mathbf{n}_{Y_0/X_0}^\vee \otimes_{\mathcal{O}_{S_0}} \mathcal{J} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_T \end{aligned}$$

for every  $T \rightarrow S_0$ .<sup>119</sup> Consequently, the  $S_0$ -functor  $N_0$  is isomorphic to

$$\mathbf{W} \left( \mathbf{n}_{Y_0/X_0}^\vee \otimes_{\mathcal{O}_{S_0}} \mathcal{J} \right) \simeq \mathbf{W} \left( \mathrm{Hom}_{\mathcal{O}_{S_0}}(\mathbf{n}_{Y_0/X_0}, \mathcal{J}) \right).$$

It follows that there are isomorphisms<sup>120</sup>

$$\begin{aligned} H^2(Y_0, N_0) &\simeq H^2 \left( Y_0, \mathrm{Hom}_{\mathcal{O}_{S_0}}(\mathbf{n}_{Y_0/X_0}, \mathcal{J}) \right) \\ &\simeq H^2 \left( Y_0, \mathbf{n}_{Y_0/X_0}^\vee \otimes_{\mathcal{O}_{S_0}} \mathcal{J} \right), \\ Z^1(Y_0, N_0) &\simeq Z^1 \left( Y_0, \mathrm{Hom}_{\mathcal{O}_{S_0}}(\mathbf{n}_{Y_0/X_0}, \mathcal{J}) \right) \\ &\simeq Z^1 \left( Y_0, \mathbf{n}_{Y_0/X_0}^\vee \otimes_{\mathcal{O}_{S_0}} \mathcal{J} \right). \end{aligned}$$

**4.23.** — Still under the hypotheses of 4.21, we shall now study how the set of subgroups  $Y$  lifting  $Y_{\mathcal{J}}$  behaves under conjugation by sections of  $X$ . If  $x$  is a section of  $X$  over  $S$  inducing the unit section of  $X_{\mathcal{J}}$ , the inner automorphism  $\mathrm{Int}(x)$  defined by  $x$  sends flat subgroups of  $X$  lifting  $Y_{\mathcal{J}}$  to flat subgroups of  $X$  lifting  $Y_{\mathcal{J}}$ . Under the conditions of 4.21(ii), the set of these subgroups is a principal homogeneous space under  $Z^1(Y_0, N_0)$ . We shall see that there is then a subgroup  $\Gamma$  of  $B^1(Y_0, N_0)$ <sup>121</sup> such that two subgroups of  $X$ , flat over  $S$  and lifting  $Y_{\mathcal{J}}$ , are conjugate by an  $x \in X(S)$  inducing the unit of  $X(S_{\mathcal{J}})$  if and only if their *difference* in  $Z^1(Y_0, N_0)$  belongs to  $\Gamma$ . In the best cases we shall show that  $\Gamma = B^1(Y_0, N_0)$ . Thus the set of flat subgroups of  $X$  lifting  $Y_{\mathcal{J}}$ , modulo conjugation by the elements  $x \in X(S)$  inducing the unit of  $X(S_{\mathcal{J}})$ , is either empty or a principal homogeneous space under  $H^1(Y_0, N_0)$  (cf. 4.29 and 4.36).

**4.24.** — We retain the notation of 4.21. Let  $Y$  be a flat subgroup of  $X$  whose reduction is  $Y_{\mathcal{J}}$ . Recall that in 0.5 we introduced the functor  $L_0^X$  (respectively  $L_0^Y$ ) defined, for a variable  $S_0$ -scheme  $T$ , by the identity

$$\mathrm{Hom}_{S_0}(T, L_0^X) = \mathrm{Hom}_{\mathcal{O}_T} \left( \omega_{X_0/S_0}^1 \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_T, \mathcal{J} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_T \right),$$

<sup>119</sup>Editor's note: The preceding argument has been expanded. It shows that the following isomorphism is valid without a flatness hypothesis. On the other hand, since 4.16 the discussion has been restricted to  $S$ -schemes  $f: T \rightarrow S$  flat over  $S$ , in order to ensure that the group

$$\mathrm{Hom}_{\mathcal{O}_{T_0}} \left( \mathbf{n}_{Y_0/X_0} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{T_0}, \mathcal{J} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{T_0} \right),$$

in which the obstruction  $c(X, Y, f)$  lies, coincides with  $N_0(T_0)$ ; see the end of 4.16.

<sup>120</sup>Editor's note: With the notation of I, 5.3, and assuming that  $Y_0$  is affine over  $S_0$ .

<sup>121</sup>Editor's note:  $Z^1(Y_0, N_0)$  has been replaced by  $B^1(Y_0, N_0)$ , since the proof shows that  $\Gamma$  is a subgroup of  $B^1(Y_0, N_0)$ ; cf. 4.27–4.29.

and similarly with  $Y$  in place of  $X$ , as well as the functor

$$L'_X = \prod_{S_0/S} L_0^X.$$

We then have:

**Lemma 4.25.** *There is a canonical exact sequence of  $Y_0$ - $\mathcal{O}_{S_0}$ -modules*

$$\mathbf{n}_{Y_0/X_0} \xrightarrow{d} \omega_{X_0/S_0}^1 \longrightarrow \omega_{Y_0/S_0}^1 \longrightarrow 0 \quad (+)$$

having the following properties:

- (i) *After pullback to  $Y_0$ ,  $d$  gives the morphism  $\overline{D}_0$  of 4.8(iii).*
- (ii) *If  $X_0$  and  $Y_0$  are smooth over  $S_0$ , then  $d$  is injective. Since the two modules of differentials are then locally free of finite type, the same is true of  $\mathbf{n}_{Y_0/X_0}$ , and the sequence is locally split.*

*Proof.*<sup>122</sup> Let  $\pi_0 : Y_0 \rightarrow S_0$  denote the structure morphism. By SGA 1 II, formula (4.3) (see also EGA IV<sub>4</sub>, 16.4.21), there is a canonical exact sequence of  $\mathcal{O}_{Y_0}$ -modules

$$\mathcal{N}_{Y_0/X_0} \xrightarrow{\overline{D}_0} \Omega_{X_0/S_0}^1 \otimes_{\mathcal{O}_{X_0}} \mathcal{O}_{Y_0} \longrightarrow \Omega_{Y_0/S_0}^1 \longrightarrow 0. \quad (\dagger)$$

Since this sequence consists of  $(Y_0 \times_S Y_0)$ -equivariant modules and morphisms, its pullback (+) by  $\varepsilon_0^*$  is an exact sequence of  $Y_0$ - $\mathcal{O}_{S_0}$ -modules, and  $(\dagger)$  is the pullback of (+) by  $\pi_0^*$  (cf. Exposé I, Section 6.8). This proves (i).

Suppose in addition that  $X_0$  and  $Y_0$  are smooth over  $S_0$ . Then, by SGA 1 II, 4.10 (see also EGA IV<sub>4</sub>, 17.2.3(i) and 17.2.5),  $\overline{D}_0$  is injective and the sequence  $(\dagger)$  consists of  $\mathcal{O}_{Y_0}$ -modules locally free of finite type (and is therefore locally split). By the equivalence of categories I, 6.8.1,  $d$  is also injective, and consequently the sequence (+) has the asserted properties.

**4.26.** —<sup>123</sup> For every  $S_0$ -scheme  $f : T \rightarrow S_0$ , the sequence (+) gives an exact sequence of  $Y_0(T)$ - $\mathcal{O}(T)$ -modules

$$\begin{aligned} 0 \longrightarrow \mathrm{Hom}_{\mathcal{O}_T} \left( f^* \omega_{Y_0/S_0}^1, f^* \mathcal{J} \right) &\longrightarrow \mathrm{Hom}_{\mathcal{O}_T} \left( f^* \omega_{X_0/S_0}^1, f^* \mathcal{J} \right) \\ &\longrightarrow \mathrm{Hom}_{\mathcal{O}_T} \left( f^* \mathbf{n}_{Y_0/X_0}, f^* \mathcal{J} \right). \end{aligned}$$

Thus there is an exact sequence of  $Y_0$ - $\mathcal{O}_{S_0}$ -modules

$$0 \longrightarrow L_0^Y \longrightarrow L_0^X \xrightarrow{d} N_0. \quad (4.26.1)$$

It gives an exact sequence of complexes of abelian groups

$$0 \longrightarrow C^*(Y_0, L_0^Y) \longrightarrow C^*(Y_0, L_0^X) \xrightarrow{d^*} C^*(Y_0, N_0),$$

and, in particular, the following commutative diagram with exact rows:

<sup>122</sup>Editor's note: The proof has been expanded to take account of the additions made in Exposé I, Section 6.8.

<sup>123</sup>Editor's note: The original has been expanded to make clear that this is an exact sequence of  $Y_0$ - $\mathcal{O}_{S_0}$ -modules.

$$\begin{array}{ccccccc}
0 & \longrightarrow & C^0(Y_0, L_{0Y}) & \longrightarrow & C^0(Y_0, L_{0X}) & \xrightarrow{d^0} & C^0(Y_0, N_0) \\
& & \downarrow \partial & & \downarrow \partial & & \downarrow \partial \\
0 & \longrightarrow & C^1(Y_0, L_{0Y}) & \longrightarrow & C^1(Y_0, L_{0X}) & \xrightarrow{d^1} & C^1(Y_0, N_0) \quad .
\end{array}$$

Observe that  $C^0(Y_0, L_0^Y)$  (respectively  $C^0(Y_0, L_0^X)$ ) is just

$$\mathrm{Hom}_{S_0}(S_0, L_0^Y) = \mathrm{Hom}_S(S, L_Y')$$

(respectively the analogous group with  $X$  in place of  $Y$ ); by 0.9, it is the group of sections of  $Y$  (respectively  $X$ ) over  $S$  inducing the unit section of  $X_{\mathcal{J}}$ . Also,  $d^1$  is precisely the morphism  $v_{i_{Y_0}}$  of 4.8(iii), where  $i_{Y_0} : Y_0 \rightarrow X_0$  is the canonical immersion.<sup>124</sup>

**Lemma 4.27.** *Under the conditions of 4.21 for  $S$ ,  $\mathcal{J}$ ,  $\mathcal{J}'$ , and  $X$ , let  $Y$  be a subgroup of  $X$ , flat over  $S$  and lifting  $Y_{\mathcal{J}}$ . Let  $i : Y \hookrightarrow X$  denote the canonical immersion.<sup>125</sup>*

- (i) *Let  $i' : Y \rightarrow X$  be a morphism of  $S$ -schemes lifting  $i_0$  (so that  $i'$  is also an immersion), let  $Y' = i'(Y)$ , and let  $d(i, i')$  be the element of  $C^1(Y_0, L_0^X)$  such that  $i' = d(i, i') \cdot i$  (cf. 1.2.4). Then the deviation  $d(Y, Y') \in C^1(Y_0, N_0)$  (cf. 4.5.1) is given by*

$$d(Y, Y') = d^1(d(i, i')).$$

- (ii) *Let  $x \in C^0(Y_0, L_0^X)$  be a section of  $X$  over  $S$  inducing the unit section of  $X_{\mathcal{J}}$  over  $S_{\mathcal{J}}$ . Then the deviation  $d(Y, \mathrm{Int}(x)Y) \in C^1(Y_0, N_0)$  (cf. 4.5.1) is given by*

$$-d(Y, \mathrm{Int}(x)Y) = d^1 \partial x = \partial d^0 x.$$

Indeed,  $Y'$  is the image of  $Y$  under the composite morphism<sup>126</sup>

$$Y \xrightarrow{(d(i, i'), i)} L_X' \times_S X \longrightarrow X,$$

which is denoted  $d(i, i') \cdot i$  in 4.8(iii). By loc. cit. and the equality  $v_{i_0} = d^1$ , we therefore have

$$\begin{aligned}
c(X, Y', d(i, i') \cdot i) - c(X, Y', i) \\
= v_{i_0}(d(i, i')) = d^1(d(i, i')).
\end{aligned}$$

But  $d(i, i') \cdot i = i'$  factors through  $Y'$  by definition, so the first term is zero. Moreover, by 4.8(ii),

$$c(X, Y', i) = d(Y', Y) = -d(Y, Y').$$

Hence  $d(Y, Y') = d^1(d(i, i'))$ , which proves (i).

Now let  $x$  be as in (ii). From the formula

$$xyx^{-1} = xyx^{-1}y^{-1}y = (x - \mathrm{Ad}(y)x)y = (-\partial x)(y) \cdot y,$$

we see that  $Y'$  is the image of  $Y$  under the immersion  $i' = (-\partial x) \cdot i_Y$ . Part (i) therefore gives

$$-d(Y, \mathrm{Int}(x)Y) = d^1 \partial x = \partial d^0 x.$$

<sup>124</sup>Editor's note: This follows from the definition of  $d : \mathbf{n}_{Y_0/X_0} \rightarrow \Omega_{X_0/S_0}^1$  (cf. 4.25) and from the definition of  $v_{i_{Y_0}}$  (cf. 4.8).

<sup>125</sup>Editor's note: Part (i) below has been added; it will be useful in 4.35.1 and then in 4.38(4) and (5).

<sup>126</sup>Editor's note: Recall that  $L_X' = \prod_{S_0/S} L_0^X$ .

**Corollary 4.28.** *Two subgroups  $Y$  and  $Y'$  of  $X$ , flat over  $S$  and lifting  $Y_{\mathcal{J}}$ , are conjugate by a section of  $X$  over  $S$  inducing the unit section of  $X_{\mathcal{J}}$  if and only if*

$$d(Y, Y') \in \partial d^0 C^0(Y_0, L_0^X) \subset \partial C^0(Y_0, N_0) = B^1(Y_0, N_0).$$

*Source note.* The French re-edition prints  $G$  here; the ambient group throughout this section is  $X$ .

**Corollary 4.29.** *If  $d^0$  is surjective, then  $Y$  and  $Y'$  as above are conjugate by a section of  $X$  over  $S$  inducing the unit section of  $X_{\mathcal{J}}$  if and only if  $d(Y, Y') \in B^1(Y_0, N_0)$ .*

**Corollary 4.30.** *Let  $Y$  be as in 4.27. The set of conjugates of  $Y$  by sections of  $X$  over  $S$  inducing the unit section of  $X_{\mathcal{J}}$  is isomorphic to*

$$d^1 \partial C^0(Y_0, L_0^X) = C^0(Y_0, L_0^X) / \ker(d^1 \partial).$$

Observe now that  $C^0(Y_0, L_0^X) / \ker(d^1 \partial)$  is computed using only the left-hand square of the commutative diagram in 4.26. In particular, it can also be computed in any diagram of the same type having the same left-hand square. Consider the functor  $L_0^X / L_0^Y$  over  $S_0$ , defined by

$$\mathrm{Hom}_{S_0}(T, L_0^X / L_0^Y) = \mathrm{Hom}_{S_0}(T, L_0^X) / \mathrm{Hom}_{S_0}(T, L_0^Y).$$

There is a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & C^0(Y_0, L_{0Y}) & \longrightarrow & C^0(Y_0, L_{0X}) & \longrightarrow & C^0(Y_0, L_{0X}/L_{0Y}) \longrightarrow 0 \\ & & \downarrow \partial & & \downarrow \partial & & \downarrow \partial \\ 0 & \longrightarrow & C^1(Y_0, L_{0Y}) & \longrightarrow & C^1(Y_0, L_{0X}) & \longrightarrow & C^1(Y_0, L_{0X}/L_{0Y}) \longrightarrow 0 \end{array} ,$$

whence, by the preceding observation:

**Corollary 4.31.** *Let  $Y$  be as in 4.27. The set of conjugates of  $Y$  by sections of  $X$  over  $S$  inducing the unit section of  $X_{\mathcal{J}}$  is isomorphic to*

$$E = \partial C^0(Y_0, L_0^X / L_0^Y) = C^0(Y_0, L_0^X / L_0^Y) / H^0(Y_0, L_0^X / L_0^Y).$$

**Corollary 4.32.** *Suppose in addition that  $S_0$  is affine and that  $\omega_{Y_0/S_0}^1$  is finite locally free.<sup>127</sup> If*

$$\mathcal{F}_0 = (\mathrm{Lie}(X_0/S_0) / \mathrm{Lie}(Y_0/S_0)) \otimes_{\mathcal{O}_{S_0}} \mathcal{J},$$

*then  $E = \Gamma(S_0, \mathcal{F}_0) / H^0(Y_0, \mathcal{F}_0)$ .*

Indeed,<sup>128</sup> since  $\omega_{Y_0/S_0}^1$  is finite locally free, as is  $\omega_{X_0/S_0}^1$  (because  $X$  is assumed smooth over  $S$ ), 0.6 gives

$$L_0^Y = \mathbf{W}(\mathrm{Lie}(Y_0/S_0) \otimes_{\mathcal{O}_{S_0}} \mathcal{J}), \quad L_0^X = \mathbf{W}(\mathrm{Lie}(X_0/S_0) \otimes_{\mathcal{O}_{S_0}} \mathcal{J}).$$

On the other hand, by 4.25 there is an exact sequence of  $Y_0$ - $\mathcal{O}_{S_0}$ -modules

$$0 \longrightarrow \mathcal{K} \longrightarrow \omega_{X_0/S_0}^1 \xrightarrow{\varphi} \omega_{Y_0/S_0}^1 \longrightarrow 0, \quad \mathcal{K} = \ker(\varphi).$$

Since  $\omega_{Y_0/S_0}^1$  and  $\omega_{X_0/S_0}^1$  are finite locally free, there is a locally split exact sequence

$$0 \longrightarrow \mathrm{Lie}(Y_0/S_0) \otimes_{\mathcal{O}_{S_0}} \mathcal{J} \longrightarrow \mathrm{Lie}(X_0/S_0) \otimes_{\mathcal{O}_{S_0}} \mathcal{J} \longrightarrow \mathcal{F}_0 \longrightarrow 0.$$

<sup>127</sup>Editor's note:  $\mathrm{Lie}(Y_0/S_0)$  has been corrected to  $\omega_{Y_0/S_0}^1$ .

<sup>128</sup>Editor's note: The original has been expanded in what follows.

It follows that there is an exact sequence of  $Y_0\text{-}\mathcal{O}_{S_0}$ -modules

$$0 \longrightarrow L_0^Y \longrightarrow L_0^X \longrightarrow \mathbf{W}(\mathcal{F}_0).$$

By the argument used to prove 4.31, we may compute  $E$  as the image of the composite map

$$C^0(Y_0, L_0^X) \xrightarrow{\pi} C^0(Y_0, \mathbf{W}(\mathcal{F}_0)) \xrightarrow{\partial} C^1(Y_0, \mathbf{W}(\mathcal{F}_0)).$$

Now  $\pi$  is the map

$$\Gamma(S_0, \text{Lie}(X_0/S_0) \otimes_{\mathcal{O}_{S_0}} \mathcal{J}) \longrightarrow \Gamma(S_0, \mathcal{F}_0).$$

Since  $S_0$  is affine,  $\pi$  is surjective, and this proves the assertion.

**Corollary 4.33.** *Let  $S, \mathcal{J}, \mathcal{J}$ , and  $X$  be as in 4.21, and let  $Y$  be a diagonalizable subgroup of  $X$ . Suppose that  $\omega_{Y_0/S_0}^1$  is finite locally free and that  $S_0$  is affine.<sup>129</sup> The set of subgroups of  $X$  conjugate to  $Y$  by a section of  $X$  over  $S$  inducing the unit section of  $X_{\mathcal{J}}$  is isomorphic to*

$$E = \Gamma\left(S_0, \frac{\text{Lie}(X_0/S_0)}{\text{Lie}(X_0/S_0)^{\text{ad}(Y_0)}} \otimes_{\Gamma(S_0, \mathcal{O}_{S_0})} \Gamma(S_0, \mathcal{J})\right),$$

that is, to  $L_0^X(Y_0)/H^0(Y_0, L_0^X)$ .<sup>130</sup>

Indeed, by I, 4.7.3 (cf. 2.13), write

$$\text{Lie}(X_0/S_0) = \text{Lie}(X_0/S_0)^{\text{ad}(Y_0)} \oplus \mathcal{R}.$$

Since  $Y_0$  is commutative,

$$\text{Lie}(Y_0/S_0) \subset \text{Lie}(X_0/S_0)^{\text{ad}(Y_0)},$$

and hence

$$\begin{aligned} \mathcal{F}_0 &= \left( \frac{\text{Lie}(X_0/S_0)^{\text{ad}(Y_0)}}{\text{Lie}(Y_0/S_0)} \right) \otimes_{\mathcal{O}_{S_0}} \mathcal{J} \oplus \mathcal{R} \otimes_{\mathcal{O}_{S_0}} \mathcal{J}, \\ \mathcal{F}_0^{\text{ad}(Y_0)} &= \left( \frac{\text{Lie}(X_0/S_0)^{\text{ad}(Y_0)}}{\text{Lie}(Y_0/S_0)} \right) \otimes_{\mathcal{O}_{S_0}} \mathcal{J}. \end{aligned}$$

By 4.32, therefore,  $E \simeq \Gamma(S_0, \mathcal{R} \otimes_{\mathcal{O}_{S_0}} \mathcal{J})$ . Returning to the definition of  $\mathcal{R}$  completes the proof.

**Corollary 4.34.** *Let  $S, \mathcal{J}, \mathcal{J}$ , and  $X$  be as in 4.21, and let  $Y$  be a diagonalizable subgroup of  $X$ . Suppose that  $\omega_{Y_0/S_0}^1$  is finite locally free and that  $S_0$  is affine.<sup>131</sup> If  $x \in X(S)$  induces the unit section of  $X_{\mathcal{J}}$  and normalizes  $Y$ , then it centralizes  $Y$ .*

This follows immediately by comparing the preceding corollary with 2.14. Indeed, 4.33 shows that the elements of  $C^0(Y_0, L_0^X)$  which globally preserve  $Y$  are the elements of  $H^0(Y_0, L_0^X)$ , and 2.14 shows that these are precisely the elements which act trivially on the canonical immersion  $Y \rightarrow X$ .

**4.35.** — Return to the general situation of 4.21 and suppose that  $Y_{\mathcal{J}}$  is smooth over  $S_{\mathcal{J}}$ . By 4.25(ii), there is then an exact sequence of  $Y_0\text{-}\mathcal{O}_{S_0}$ -modules

$$0 \longrightarrow \text{Lie}(Y_0/S_0) \longrightarrow \text{Lie}(X_0/S_0) \longrightarrow \mathbf{n}_{Y_0/X_0}^{\vee} \longrightarrow 0, \quad (*)$$

and these are finite locally free  $\mathcal{O}_{S_0}$ -modules.

<sup>129</sup>Editor's note: The hypothesis on  $\omega_{Y_0/S_0}^1$  has been added, and the hypothesis " $S$  affine" has been replaced by " $S_0$  affine."

<sup>130</sup>Editor's note: What follows has been added; cf. 4.34.

<sup>131</sup>Editor's note: Cf. Editor's Note 129.



On the other hand, by SGA 1, II, 4.10, every subscheme  $Y$  of  $X$  lifting  $Y_{\mathcal{J}}$  and flat over  $S$  is smooth over  $S$ .<sup>132</sup> Suppose in addition that  $S_0$  and  $Y_{\mathcal{J}}$  are affine. Since  $\mathbf{n}_{Y_0/X_0}^\vee$  is a locally free  $\mathcal{O}_{S_0}$ -module, the sequence  $(*)$  remains exact after tensoring with  $\mathcal{J}$  over  $\mathcal{O}_{S_0}$ , and then pulling back to  $Y_0^n$ . Since the  $Y_0^n$  are affine, this gives an exact sequence of complexes of abelian groups

$$0 \longrightarrow C^*(Y_0, L_0^Y) \longrightarrow C^*(Y_0, L_0^X) \xrightarrow{d^*} C^*(Y_0, N_0) \longrightarrow 0,$$

and, in particular, a commutative diagram with exact rows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & C^1(Y_0, L_{0Y}) & \longrightarrow & C^1(Y_0, L_{0X}) & \xrightarrow{d^1} & C^1(Y_0, N_0) \longrightarrow 0 \\ & & \downarrow \partial & & \downarrow \partial & & \downarrow \partial \\ 0 & \longrightarrow & C^2(Y_0, L_{0Y}) & \longrightarrow & C^2(Y_0, L_{0X}) & \xrightarrow{d^2} & C^2(Y_0, N_0) \longrightarrow 0 \end{array} \quad .$$

Let now  $Y$  and  $Y'$  be two subgroups of  $X$  lifting  $Y_{\mathcal{J}}$  and flat, hence smooth, over  $S$ . Since  $Y_{\mathcal{J}}$  is affine, 0.15 shows that  $Y$  and  $Y'$  are isomorphic as schemes lifting  $Y_{\mathcal{J}}$ ; that is, there is an isomorphism of  $S$ -schemes

$$f : Y \xrightarrow{\sim} Y'$$

inducing the identity on  $Y_{\mathcal{J}}$ . On the one hand, by 1.2.4,  $f$  defines an element  $a \in C^1(Y_0, L_0^X)$  such that

$$f(y) = a(y_0)y$$

for every  $y \in Y(S')$ ,  $S' \rightarrow S$ , and by 4.27(i),

$$d^1(a) = d(Y, Y').$$

Moreover, because  $Y$  and  $Y'$  are subgroups of  $X$ , the last element belongs to  $Z^1(Y_0, N_0)$  (cf. 4.21). Thus  $\partial a$  is an element of  $Z^2(Y_0, L_0^Y)$ , whose image  $\overline{\partial a}$  in  $H^2(Y_0, L_0^Y)$  depends only on the class  $\overline{d(Y, Y')} \in H^1(Y_0, N_0)$ . This is precisely the definition of the connecting map

$$\partial^1 : H^1(Y_0, N_0) \longrightarrow H^2(Y_0, L_0^Y),$$

so that

$$\partial^1(\overline{d(Y, Y')}) = \overline{\partial a}.$$

On the other hand, transport the group structure of  $Y'$  by  $f$ , and let  $Y_1$  be the resulting group, whose underlying scheme is therefore  $Y$ . Thus its group law  $\mu_1$  is defined, for every  $S' \rightarrow S$  and  $x, y \in Y(S')$ , by

$$\mu_1(x, y) = f^{-1}(f(x)f(y)).$$

By 3.5.1,  $Y_1$  defines a cocycle

$$\delta(Y, Y_1) \in Z^2(Y_0, \text{Lie}(Y_0/S_0))$$

<sup>132</sup>Editor's note: What follows, together with Proposition 4.35.1, has been added; it is implicit in the original. Cf. 4.38(5).

such that, for every  $S' \rightarrow S$  and  $x, y \in Y(S')$ ,

$$\delta(Y, Y_1)(x_0, y_0)xy = \mu_1(x, y) = f^{-1}(f(x)f(y)).$$

Put  $b = \delta(Y, Y_1)$ . For every  $S' \rightarrow S$  and  $x, y \in Y(S')$ , we have  $(b(x_0, y_0)xy)_0 = x_0y_0$ . Hence  $f(b(x_0, y_0)xy)$  is equal, on the one hand, to

$$a(x_0y_0)b(x_0, y_0)xy,$$

and, on the other hand, to

$$f(x)f(y) = a(x_0)x a(y_0)y = a(x_0) \operatorname{Ad}(x_0)(a(y_0))xy.$$

Comparing these two expressions, we find that  $b(x_0, y_0)$  equals

$$\begin{aligned} & a(x_0y_0)^{-1}a(x_0) \operatorname{Ad}(x_0)(a(y_0)) \\ &= \operatorname{Ad}(x_0)(a(y_0)) - a(x_0y_0) + a(x_0) = (\partial a)(x_0, y_0). \end{aligned}$$

Thus  $\delta(Y, Y_1) = \partial a$ , and we have proved:

**Proposition 4.35.1.**<sup>132</sup> *Under the hypotheses of 4.21, suppose in addition that  $S_0$  is affine and that  $Y_{\mathcal{J}}$  is smooth over  $S_{\mathcal{J}}$  and affine. Let  $Y, Y'$  be two subgroups of  $X$  lifting  $Y_{\mathcal{J}}$  and flat (hence smooth) over  $S$ . Let  $f : Y \xrightarrow{\sim} Y'$  be an isomorphism of  $S$ -schemes inducing the identity on  $Y_{\mathcal{J}}$ , and let  $Y_1$  denote the group obtained by transporting the group structure of  $Y'$  by  $f$ . Then*

$$\partial^1(\overline{d(Y, Y')}) = \overline{\delta(Y, Y_1)}.$$

**Proposition 4.36.** *Under the hypotheses of 4.21, suppose in addition that  $Y_{\mathcal{J}}$  is smooth over  $S_{\mathcal{J}}$  and that  $S_0$  is affine. The set of sub- $S$ -groups  $Y$  of  $X$ , flat (or smooth) over  $S$ , reducing to  $Y_{\mathcal{J}}$ , modulo conjugation by sections of  $X$  over  $S$  inducing the unit section of  $X_{\mathcal{J}}$ , is either empty or a principal homogeneous set under the group*

$$H^1\left(Y_0, [\operatorname{Lie}(X_0/S_0)/\operatorname{Lie}(Y_0/S_0)] \otimes_{\mathcal{O}_{S_0}} \mathcal{J}\right).$$

It is enough to verify that Corollary 4.29 applies, that is, that

$$d^0 : \operatorname{Hom}_{\mathcal{O}_{S_0}}(\omega_{X_0/S_0}^1, \mathcal{J}) \longrightarrow \operatorname{Hom}_{\mathcal{O}_{S_0}}(\mathbf{n}_{Y_0/X_0}, \mathcal{J})$$

is surjective. This follows from the fact that the sequence (+) of 4.25(ii) is split, since  $S_0$  is affine.<sup>133</sup>

Let us finally state a corollary common to 4.21 and 4.36. In fact, this will be the only form in which we shall subsequently use the general results of this section.<sup>134</sup>

**Corollary 4.37.** *Let  $S$  be a scheme and let  $S_0$  be the closed subscheme defined by a nilpotent ideal  $\mathcal{J}$ . Let  $X$  be an  $S$ -group smooth over  $S$ , and let  $Y_0$  be a sub- $S_0$ -group of  $X_0$ , flat over  $S_0$ .*

(i) *If  $S_0$  is affine,  $Y_0$  is smooth over  $S_0$ , and if*

$$H^1\left(Y_0, [\operatorname{Lie}(X_0/S_0)/\operatorname{Lie}(Y_0/S_0)] \otimes_{\mathcal{O}_{S_0}} \mathcal{J}^{n+1}/\mathcal{J}^{n+2}\right) = 0$$

*for every  $n \geq 0$ , then any two sub- $S$ -groups of  $X$ , flat (or smooth) over  $S$  and reducing to  $Y_0$ , are conjugate by a section of  $X$  over  $S$  inducing the unit section of  $X_0$ .*

<sup>133</sup>Editor's note: This also follows from the proof of 4.32.

<sup>134</sup>Editor's note: For example, 4.37 is used in Exposé IX to prove statements 3.2 bis and 3.6 bis.

(ii) If  $Y_0$  is affine and of finite presentation, and if<sup>135</sup>

$$H^2\left(Y_0, \mathbf{n}_{Y_0/X_0}^\vee \otimes_{\mathcal{O}_{S_0}} \mathcal{I}^{n+1}/\mathcal{I}^{n+2}\right) = 0$$

for every  $n \geq 0$ , then there exists a sub- $S$ -group of  $X$ , flat over  $S$ , reducing to  $Y_0$ .

**4.38.** It remains to relate the three constructions made in this Exposé. To avoid inessential complications, we shall place ourselves in the following situation:  $S_0$  is the spectrum of a field  $k$ ,  $S$  is the spectrum of the dual numbers  $D(k)$ ,  $G$  is an  $S$ -group smooth over  $S$ , and  $K$  is a sub- $S$ -group, smooth over  $S$  and affine.<sup>136</sup>

Put

$$\mathfrak{g}_0 = \text{Lie } G_0 = \Gamma(S_0, \text{Lie } G_0) = \text{Lie}(G_0/S_0)(S_0), \quad \mathfrak{k}_0 = \text{Lie } K_0.$$

We have an exact sequence of  $k$ -vector spaces<sup>137</sup>

$$0 \longrightarrow \mathfrak{k}_0 \xrightarrow{i} \mathfrak{g}_0 \xrightarrow{d} \mathbf{n}_{K_0/G_0}^\vee \longrightarrow 0,$$

which gives rise to an exact cohomology sequence

$$\begin{aligned} 0 \longrightarrow H^0(K_0, \mathfrak{k}_0) &\xrightarrow{i^0} H^0(K_0, \mathfrak{g}_0) \xrightarrow{d^0} H^0(K_0, \mathfrak{g}_0/\mathfrak{k}_0) \\ &\xrightarrow{\partial^0} H^1(K_0, \mathfrak{k}_0) \xrightarrow{i^1} H^1(K_0, \mathfrak{g}_0) \xrightarrow{d^1} H^1(K_0, \mathbf{n}_{K_0/G_0}^\vee) \xrightarrow{\partial^1} H^2(K_0, \mathfrak{k}_0). \end{aligned}$$

These various groups all have a geometric meaning.

a)  $H^0(K_0, \mathfrak{k}_0) = \text{Lie } \underline{\text{Centr}}(K_0)$ <sup>138</sup> by II, 5.2.3.

b)  $H^0(K_0, \mathfrak{g}_0) = \text{Lie } \underline{\text{Centr}}_{G_0}(K_0)$ <sup>138</sup> (idem).

c)  $H^0(K_0, \mathfrak{g}_0/\mathfrak{k}_0) = \text{Lie } \underline{\text{Norm}}_{G_0}(K_0)/\mathfrak{k}_0$ <sup>138</sup> (idem).

d)

$$H^1(K_0, \mathfrak{k}_0) = \text{Lie}(\underline{\text{Aut}}_{S_0\text{-gr.}}(K_0)) / \text{Im}(\mathfrak{k}_0),$$

where  $\text{Im}(\mathfrak{k}_0)$  denotes the image of  $\mathfrak{k}_0$  under the morphism  $\text{Lie}(\text{Int}_0)$  deduced from

$$\text{Int}_0 : K_0 \longrightarrow \underline{\text{Aut}}_{S_0\text{-gr.}}(K_0).$$

Indeed, it follows from 2.1(ii), applied with  $Y = X = K$  and  $f_0 = \text{id}_{K_0}$ , that  $Z^1(K_0, \mathfrak{k}_0)$  is the group of infinitesimal automorphisms of the  $S_0$ -group  $K_0$ , and that  $H^1(K_0, \mathfrak{k}_0)$  is obtained by quotienting by inner infinitesimal automorphisms, that is, by the image of  $\mathfrak{k}_0$ . Moreover, by II, 4.2.2, we also have

$$Z^1(K_0, \mathfrak{k}_0) = \text{Lie}(\underline{\text{Aut}}_{S_0\text{-gr.}}(K_0)).$$

139

<sup>135</sup>Editor's note: We have replaced  $\text{Hom}_{\mathcal{O}_{S_0}}(\mathbf{n}_{Y_0/X_0}, \mathcal{I}^{n+1}/\mathcal{I}^{n+2})$  by  $\mathbf{n}_{Y_0/X_0}^\vee \otimes_{\mathcal{O}_{S_0}} \mathcal{I}^{n+1}/\mathcal{I}^{n+2}$ , in accordance with Remark 4.22.

<sup>136</sup>Editor's note: We have slightly modified the original in what follows. In particular, we have replaced  $X$  by  $G$  and  $Y$  by  $K$ , and denoted their Lie algebras by  $\mathfrak{g}_0$  and  $\mathfrak{k}_0$ . We have also written  $H^i(K_0, \cdot)$  explicitly in place of the abbreviation  $H^i(\cdot)$  used in the original.

<sup>137</sup>Editor's note: Equipped with the adjoint action of  $K_0$ .

<sup>138</sup>Editor's note: Since the formation of centralizers and normalizers commutes with base change (cf. I, 2.3.3.1), we have written  $\underline{\text{Centr}}(K_0)$  in place of  $\underline{\text{Centr}}(K)_0$  in the original, and likewise  $\underline{\text{Centr}}_{G_0}(K_0)$  and  $\underline{\text{Norm}}_{G_0}(K_0)$  in place of  $\underline{\text{Centr}}_G(K)_0$  and  $\underline{\text{Norm}}_G(K)_0$ .

<sup>139</sup>Editor's note: This is the Lie algebra  $\text{Der}_k(\mathfrak{k}_0)$  of derivations of  $\mathfrak{k}_0$ ; hence  $H^1(K_0, \mathfrak{k}_0)$  is the quotient of  $\text{Der}_k(\mathfrak{k}_0)$  by the inner derivations (that is, by the image of  $\text{ad} : \mathfrak{k}_0 \rightarrow \text{Der}_k(\mathfrak{k}_0)$ ).

- e) By 2.1(ii),  $H^1(K_0, \mathfrak{g}_0)$  is the group of deviations between homomorphisms  $K \rightarrow G$  extending the canonical immersion  $i_0 : K_0 \rightarrow G_0$ , modulo the deviations obtained through the action of the inner automorphisms of  $G$  defined by elements of  $G(S)$  which give the unit of  $G(S_0)$  (that is, elements of  $\mathfrak{g}_0$ ).
- f) By 4.36,  $H^1(K_0, \mathbf{n}_{K_0/G_0}^\vee)$  is the group of deviations between subgroups  $K'$  of  $G$  extending  $K_0$  and flat over  $S$  (hence smooth over  $S$ , cf. SGA 1, II, 4.10), modulo the deviations obtained through the action of the inner automorphisms of  $G$  constructed as above.
- g) By 3.5(ii),  $H^2(K_0, \mathfrak{k}_0)$  is the group of deviations between group structures on  $K$  extending that of  $K_0$ , modulo the  $S$ -automorphisms of  $K$  inducing the identity on  $K_0$ .

We now propose to make explicit the six morphisms of the preceding exact sequence in terms of the geometric interpretation just given.

- 1) The morphisms  $i^0$  and  $d^0$  are simply those obtained by passing to the Lie algebra (and then, for  $d^0$ , to the quotient) in the canonical monomorphisms

$$\underline{\text{Centr}}(K_0) \longrightarrow \underline{\text{Centr}}_{G_0}(K_0) \longrightarrow \underline{\text{Norm}}_{G_0}(K_0).$$

This follows immediately from the definition of the identifications a), b), and c).

- 2) The morphism  $\partial^0$  is constructed as follows. Let

$$\bar{x} \in \text{Lie } \underline{\text{Norm}}_{G_0}(K_0)/\mathfrak{k}_0.$$

Lift it to an element

$$x \in \text{Lie } \underline{\text{Norm}}_{G_0}(K_0) \subset \underline{\text{Norm}}_G(K)(S).$$

Then  $\text{Int}(x)$  defines an automorphism of  $K$  inducing the identity on  $K_0$ , hence an element of  $\text{Lie } \underline{\text{Aut}}_{S_0\text{-gr.}}(K_0)$ . Denote by  $\overline{\text{Int}(x)}$  the image of this element in

$$\text{Lie } \underline{\text{Aut}}_{S_0\text{-gr.}}(K_0)/\text{Im}(\mathfrak{k}_0).$$

Then

$$\partial^0(\bar{x}) = -\overline{\text{Int}(x)} = \overline{\text{Int}(x^{-1})}. \quad (*)$$

Indeed, let us compute the element of  $\text{Lie } \underline{\text{Aut}}_{S_0\text{-gr.}}(K_0)$  defined by  $\text{Int}(x)$ . By definition it corresponds to an element  $a \in Z^1(K_0, \mathfrak{k}_0)$  such that

$$xyx^{-1} = a(y_0)y, \quad y \in K(S'), \quad S' \longrightarrow S.$$

But this can also be written

$$a(y_0) = xyx^{-1}y^{-1} = x - \text{Ad}(y)x = -\partial(x)(y_0),$$

whence  $a = -\partial(x)$ .

On the other hand,<sup>140</sup> the image under  $\partial$  of

$$x \in \text{Lie } \underline{\text{Norm}}_{G_0}(K_0) \subset \mathfrak{g}_0$$

is an element of  $Z^1(K_0, \mathfrak{k}_0)$ , whose image  $\overline{\partial(x)}$  in  $H^1(K_0, \mathfrak{k}_0)$  depends only on  $\bar{x}$ . By definition of the connecting map  $\partial^0$ ,

$$\partial^0(\bar{x}) = \overline{\partial(x)}.$$

Combined with  $a = -\partial(x)$ , this proves (\*).

- 3) <sup>141</sup> Let  $i : K \rightarrow G$  denote the canonical immersion. Let  $\bar{u} \in H^1(K_0, \mathfrak{k}_0)$  be the image

<sup>140</sup>Editor's note: We have added the following sentence.

<sup>141</sup>Editor's note: We have expanded the original in what follows, and in (\*\*) we have corrected  $u \circ i$  to  $i \circ u$ .

of an element

$$u \in \text{Lie } \underline{\text{Aut}}_{S_0\text{-gr.}}(K_0) \subset \underline{\text{Aut}}_{S\text{-gr.}}(K).$$

Then

$$i^1(\bar{u}) = \bar{d}(i, i \circ u), \quad (**)$$

where  $\bar{d}(i, i \circ u)$  is the class defined in 2.1.1.

Indeed,  $\bar{u}$  is the image of an element  $v \in Z^1(K_0, \mathfrak{k}_0)$  such that  $u(y) = v(y_0)y$ , and  $i^1(\bar{u})$  is the image in  $H^1(K_0, \mathfrak{g}_0)$  of the cocycle  $i \circ v \in Z^1(K_0, \mathfrak{g}_0)$ . Since  $i$  is a group morphism, the equality  $u(y) = v(y_0)y$  implies

$$i(u(y)) = i(v(y_0))i(y).$$

It follows that  $i \circ v = d(i, i \circ u)$ , proving (\*\*).

- 4) Let  $i' : K \rightarrow G$  be a group morphism lifting  $i_0$ , let  $\bar{d}(i, i')$  be the class defined in 2.1.1, and let

$$d(K, i'(K)) \in C^1(K_0, \mathbf{n}_{K_0/G_0}^\vee)$$

be the deviation defined in 4.5.1. By 4.21,  $d(K, i'(K))$  belongs to  $Z^1(K_0, \mathbf{n}_{K_0/G_0}^\vee)$ . Denote its image in  $H^1(K_0, \mathbf{n}_{K_0/G_0}^\vee)$  by  $\bar{d}(K, i'(K))$ . Then, by 4.27(i),

$$d^1(\bar{d}(i, i')) = \bar{d}(K, i'(K)). \quad (\dagger)$$

*Source note.* In these three adjacent occurrences the French re-edition prints  $\mathbf{n}_{K_0/X_0}$ , without the dual. The English uses  $\mathbf{n}_{K_0/G_0}^\vee$ , as required by the exact sequence and cohomology sequence established at the start of 4.38.

- 5) Finally, let  $K'$  be a subgroup of  $G$  lifting  $K_0$  and flat, hence smooth, over  $S$ . We have assumed that  $K_0$  is affine. We therefore know that  $K$  and  $K'$  are isomorphic as schemes extending  $K_0$  (cf. 0.15), so there exists an isomorphism of  $S$ -schemes

$$f : K \xrightarrow{\sim} K'$$

inducing the identity on  $K_0$ . Transport the group structure of  $K'$  by  $f$ , and let  $K_1$  be the group thereby obtained (whose underlying scheme is thus  $K$ ); that is, the group law  $\mu_1$  of  $K_1$  is defined, for every  $S' \rightarrow S$  and every  $x, y \in K(S')$ , by

$$\mu_1(x, y) = f^{-1}(f(x)f(y)).$$

By 3.5.1,<sup>142</sup>  $K_1$  defines a cocycle  $\delta(K, K_1) \in Z^2(K_0, \mathfrak{k}_0)$  such that, for every  $S' \rightarrow S$  and every  $x, y \in K(S')$ ,

$$\delta(K, K_1)(x_0, y_0)xy = \mu_1(x, y) = f^{-1}(f(x)f(y)).$$

Then, by 4.35.1,

$$\partial^1 \bar{d}(K, K') = \bar{\delta}(K, K_1). \quad (\ddagger)$$

---

<sup>142</sup>Editor's note: We have modified the original in what follows, taking account of the additions made in 3.5.1 and 4.35.1.

## Bibliography

<sup>143</sup>

- [BA1g] N. Bourbaki, *Algèbre*, Chap. I–III, Hermann, 1970.
- [BAC] N. Bourbaki, *Algèbre commutative*, Chap. I–IV, Masson, 1985.
- [DG70] M. Demazure and P. Gabriel, *Groupes algébriques*, Masson & North-Holland, 1970.
- [Fr64] P. Freyd, *Abelian categories*, Harper and Row, 1964.

---

<sup>143</sup>Editor’s note: Additional references cited in this Exposé.

# Exposé IV

## Topologies and Sheaves

by M. Demazure\*

This exposé is intended to acquaint the reader with the essentials of the language of topologies and sheaves (without cohomology), a language particularly convenient for questions concerning the formation of quotients, among others.<sup>0</sup>

The first three sections develop the language of forming quotients. The fourth, which is the central part, presents the theory of sheaves, oriented mainly toward its application to questions about quotients; the fifth applies this theory to the formation of quotients in groups and to principal homogeneous bundles. The final section concerns the category of schemes more specifically and defines several useful topologies on that category.

The reader will profitably consult [AS], [MA], [D], and SGA 4; in particular, [D] for the applications of topologies to descent theory, and SGA 4 for questions concerning universes (which are treated especially poorly in this exposé).

### 1. Universal Effective Epimorphisms

Throughout the remainder of this exposé, a category  $\mathcal{C}$  is fixed.

**Definition 1.1.** A morphism  $u : T \rightarrow S$  is called an epimorphism if, for every object  $X$ , the corresponding map

$$X(S) = \text{Hom}(S, X) \longrightarrow X(T) = \text{Hom}(T, X)$$

is injective.<sup>1</sup> It is called a universal epimorphism if, for every morphism  $S' \rightarrow S$ , the fiber product  $T' = T \times_S S'$  exists and  $u' : T' \rightarrow S'$  is an epimorphism.

**Definition 1.2.** A diagram of maps of sets

$$A \xrightarrow{u} B \begin{matrix} \xrightarrow{v_1} \\ \xrightarrow{v_2} \end{matrix} C$$

is said to be exact if  $u$  is injective and its image consists of those elements  $b \in B$  for which  $v_1(b) = v_2(b)$ . A diagram of the same form in  $\mathcal{C}$  is said to be exact if, for every object  $X$  of  $\mathcal{C}$ , the corresponding diagram of sets

$$A(X) \longrightarrow B(X) \rightrightarrows C(X)$$

---

\*This text develops the substance of two lectures given by A. Grothendieck, supplementing them at several important points that had been passed over in silence or only briefly touched upon.

<sup>0</sup>Editor's note: version of October 13, 2024.

<sup>1</sup>Editor's note: that is, if  $u$  is right-cancellable.

is exact; one then also says that  $u$  makes  $A$  a kernel of the pair of arrows  $(v_1, v_2)$ . Dually, a diagram in  $\mathcal{C}$

$$C \begin{array}{c} \xrightarrow{v_1} \\ \xrightarrow{v_2} \end{array} B \xrightarrow{u} A$$

is said to be exact if it is exact as a diagram in the opposite category  $\mathcal{C}^\circ$ ; that is, if for every object  $X$  of  $\mathcal{C}$ , the corresponding diagram of sets

$$X(A) \longrightarrow X(B) \rightrightarrows X(C)$$

is exact.<sup>2</sup> One also says that  $u$  makes  $A$  a cokernel of the pair of arrows  $(v_1, v_2)$ .

**Definition 1.3.** A morphism  $u : T \rightarrow S$  is called an effective epimorphism if the fiber square  $T \times_S T$  exists and the diagram

$$T \times_S T \begin{array}{c} \xrightarrow{\text{pr}_1} \\ \xrightarrow{\text{pr}_2} \end{array} T \xrightarrow{u} S$$

is exact; that is, if  $u$  makes  $S$  a cokernel of  $(\text{pr}_1, \text{pr}_2)$ . It is called a universal effective epimorphism if, for every morphism  $S' \rightarrow S$ , the fiber product  $T' = T \times_S S'$  exists and the morphism  $u' : T' \rightarrow S'$  is an effective epimorphism.

There are evidently implications

$$\begin{array}{ccc} \text{universal effective epimorphism} & \implies & \text{effective epimorphism} \\ \Downarrow & & \Downarrow \\ \text{universal epimorphism} & \implies & \text{epimorphism,} \end{array}$$

but in general no other implication holds.<sup>3</sup>

**Definition 1.4.0.**<sup>4</sup> We “recall” that a morphism  $u : T \rightarrow S$  is said to be *squarable* if, for every morphism  $S' \rightarrow S$ , the fiber product  $T \times_S S'$  exists.

**Lemma 1.4.** Consider morphisms  $U \xrightarrow{v} T \xrightarrow{u} S$ . Then:

- (a) if  $u$  and  $v$  are epimorphisms, then  $uv$  is an epimorphism; and if  $uv$  is an epimorphism, then  $u$  is an epimorphism;
- (b) if  $u$  and  $v$  are universal epimorphisms, then  $uv$  is a universal epimorphism; and if  $uv$  is a universal epimorphism and  $u$  is squarable, then  $u$  is a universal epimorphism.

Lemma 1.4 follows immediately from the definitions. We deduce:

**Corollary 1.5.** Let  $u : X \rightarrow Y$  and  $u' : X' \rightarrow Y'$  be universal epimorphisms, and suppose that  $Y \times Y'$  exists. Then  $X \times X'$  exists and

$$u \times u' : X \times X' \longrightarrow Y \times Y'$$

<sup>2</sup>Editor’s note: this implies, in particular, that  $u$  is an epimorphism.

<sup>3</sup>Editor’s note: for example, if  $\mathcal{C} = \mathbf{Sch}$  is the category of schemes, it is easy to see that every universal epimorphism is surjective. Let

$$T = \coprod_{p \text{ prime}} \text{Spec}(\mathbf{F}_p), \quad S = \text{Spec}(\mathbf{Z}).$$

Then the morphism  $u : T \rightarrow S$  is an epimorphism that is not universal. On the other hand,  $T \times_S T$  identifies with  $T$ , so the two projections  $T \times_S T \rightarrow T$  coincide. Since  $\text{id}_T$  does not descend to a morphism  $S \rightarrow T$ , this shows that  $u$  is not an effective epimorphism.

<sup>4</sup>Editor’s note: the numbering 1.4.0 has been added for later reference.



is a universal epimorphism.

We also record the following.

**Definition 1.6.0.**<sup>5</sup> An object  $S$  of  $\mathcal{C}$  is called *squarable* if its product with every object of  $\mathcal{C}$  exists. (If  $\mathcal{C}$  has a final object  $e$ , this is equivalent to saying that the morphism  $S \rightarrow e$  is squarable; cf. 1.4.0.)

**Lemma 1.6.** Let  $u : X \rightarrow Y$  be a morphism in  $\mathcal{C}/_S$ . For it to be an epimorphism (respectively, a universal epimorphism, an effective epimorphism, or a universal effective epimorphism), it is sufficient that the corresponding morphism in  $\mathcal{C}$  have the same property; it is also necessary if  $S$  is assumed to be a squarable object of  $\mathcal{C}$ .

The proof is immediate and is left to the reader. The hypothesis that  $S$  is squarable is used to interpret the  $\mathcal{C}$ -morphisms from an object  $Y$  of  $\mathcal{C}/_S$  to an object  $Z$  of  $\mathcal{C}$  as the  $\mathcal{C}/_S$ -morphisms from  $Y$  to  $Z \times S$ .

**Lemma 1.7.** With the notation of 1.4, if  $u$  and  $v$  are effective epimorphisms and  $v$  is a universal epimorphism, then  $uv$  is an effective epimorphism.

To see this, consider the diagram

$$\begin{array}{ccccc}
 S & \xleftarrow{u} & T & \rightleftharpoons & T \times_S T \\
 & & \uparrow v & & \uparrow v \times_S v \\
 & & U & \rightleftharpoons & U \times_S U \\
 & & \uparrow & \nearrow & \\
 & & U \times_T U & & 
 \end{array} \tag{1.7.1}$$

By hypothesis, the first row and the first column are exact; moreover, by 1.5 and 1.6,  $v \times_S v$  is an epimorphism, since  $v$  is a universal epimorphism. The conclusion follows by an evident diagram chase. Indeed, if an element of  $X(U)$  has the same two images in  $X(U \times_S U)$ , then it has, *a fortiori*, the same two images in  $X(U \times_T U)$ , and hence comes from an element of  $X(T)$ , since the first column is exact.

Since the first row is exact, it remains only to verify that the element in question has the same two images in  $X(T \times_S T)$ . Because  $v \times_S v$  is an epimorphism, it is enough to verify that their images in  $X(U \times_S U)$  are the same, which is indeed the case.

**Proposition 1.8.** Consider morphisms  $U \xrightarrow{v} T \xrightarrow{u} S$ . If  $u$  and  $v$  are universal effective epimorphisms, then  $uv$  is a universal effective epimorphism. If  $uv$  is a universal effective epimorphism and  $u$  is squarable, then  $u$  is a universal effective epimorphism.

The first implication follows at once from 1.7. For the second, consider the following diagram (of “bisimplicial” type):

$$\begin{array}{ccccc}
 S & \xleftarrow{u} & T & \rightleftharpoons & T \times_S T \\
 \uparrow uv & & \uparrow & & \uparrow \\
 U & \longleftarrow & U \times_S T & \rightleftharpoons & U \times_S T \times_S T \\
 \uparrow & & \uparrow & & \uparrow \\
 U \times_S U & \longleftarrow & U \times_S U \times_S T & \rightleftharpoons & U \times_S U \times_S T \times_S T.
 \end{array} \tag{1.8.1}$$

The first, second, and third columns are exact by the hypothesis that  $uv$  is a universal effective epimorphism. The second row is exact because  $U \times_S T \rightarrow U$  is an effective epimorphism (it has a section over  $U$ ), and the same is true of the third row, for the same reason. An evident diagram chase then shows that the first row is exact, that is, that  $u$  is an effective epimorphism. Since the hypotheses are invariant under an arbitrary base change  $S' \rightarrow S$ , it follows that  $u$  is in fact a universal effective epimorphism.

<sup>5</sup>Editor’s note: the numbering 1.6.0 has been added for later reference.

**Corollary 1.9.** *Let  $u : X \rightarrow Y$  and  $u' : X' \rightarrow Y'$  be universal effective epimorphisms, and suppose that  $Y \times Y'$  exists. Then  $X \times X'$  exists and*

$$u \times u' : X \times X' \longrightarrow Y \times Y'$$

*is a universal effective epimorphism.*

The proof is the same as for 1.5, using the diagram

$$\begin{array}{ccccc}
 & & Y' & \xleftarrow{u'} & X' \\
 & & \uparrow & & \uparrow \\
 X & \xleftarrow{\quad} & X \times Y' & \xleftarrow{\quad} & X \times X' \\
 u \downarrow & & \downarrow & \swarrow u \times u' & \\
 Y & \xleftarrow{\quad} & Y \times Y' & & .
 \end{array} \tag{1.9.1}$$

**Corollary 1.10.** *Let  $u : T \rightarrow S$  be a squarable morphism, and let  $S' \rightarrow S$  be a base-change morphism that is a universal effective epimorphism. Then  $u$  is a universal effective epimorphism if and only if*

$$u' : T' = T \times_S S' \longrightarrow S'$$

*is one:*

$$\begin{array}{ccc}
 T & \xleftarrow{v'} & T' \\
 u \downarrow & & \downarrow u' \\
 S & \xleftarrow{v} & S'.
 \end{array} \tag{1.10.1}$$

Only sufficiency requires proof. If  $u'$  is a universal effective epimorphism, then so is  $vu'$ , by 1.8. Since  $vu' = uv'$ , another application of 1.8 shows that  $u$  is a universal effective epimorphism.

**Remark 1.11.** *The same reasoning shows that in 1.10 one may replace “universal effective epimorphism” by “universal epimorphism,” by “universal and effective epimorphism,” or simply by “epimorphism” (and in the last case the hypothesis that  $u$  be squarable is plainly unnecessary).*

In the proof of 1.8 we used the following result, which deserves to be stated explicitly.

**Proposition 1.12.** *Let  $u : T \rightarrow S$  be a morphism admitting a section. Then  $u$  is an epimorphism; if  $T \times_S T$  exists, it is an effective epimorphism; and if, in addition,  $u$  is squarable, it is a universal effective epimorphism.*

The first assertion is contained in 1.4(a), and the third follows at once from the second, so it is enough to establish the second. In fact, there is a stronger conclusion: for every functor  $F : \mathcal{C}^\circ \rightarrow \mathbf{Set}$ , not necessarily representable, the diagram of sets

$$F(S) \longrightarrow F(T) \rightrightarrows F(T \times_S T)$$

is exact. This may be regarded as a special case of the formalism of Čech cohomology (in dimension 0!), which we merely recall here. Assume only that  $T \times_S T$  exists, and set

$$\check{H}^0(T/S, F) = \text{Ker}(F(T) \rightrightarrows F(T \times_S T)).$$

One may plainly regard  $\check{H}^0(T/S, F)$  as a contravariant functor of the variable  $T$  in  $\mathcal{C}/S$ : every  $S$ -morphism  $T' \rightarrow T$  defines a map

$$\check{H}^0(T/S, F) \longrightarrow \check{H}^0(T'/S, F). \tag{+}$$

Fix  $T, T'$  in  $\mathcal{C}/S$ . A familiar calculation shows that, if there exists an  $S$ -morphism  $T' \rightarrow T$ , the corresponding map  $(+)$  is independent of the choice of that morphism.<sup>6</sup>

Thus  $\check{H}^0(T/S, F)$  may be regarded as a functor on the category associated with the set  $\text{Ob}(\mathcal{C}/S)$ , preordered by the relation of domination:  $T'$  dominates  $T$  if there exists an  $S$ -morphism  $T' \rightarrow T$ . In particular, if  $T$  and  $T'$  are isomorphic in this latter category, that is, if each dominates the other, then  $(+)$  is an isomorphism of sets. This applies in particular when  $T'$  is the final object of  $\mathcal{C}/S$ , essentially  $S$  itself. The object  $T$  always dominates  $T' = S$ , while the converse holds precisely when  $T/S$  has a section. This proves 1.12 in the stronger form announced.

**Remark 1.13.** *For various applications, the notions introduced and the results stated in this exposé must be developed more generally for a family of morphisms  $u_i : T_i \rightarrow S$  with the same target, rather than for a single morphism  $u : T \rightarrow S$ . Such a family is called epimorphic if, for every object  $X$  of  $\mathcal{C}$ , the corresponding map*

$$X(S) \longrightarrow \prod_i X(T_i)$$

*is injective. One similarly introduces the notion of an effective epimorphic family and the universal variants of these notions. Whenever needed below, we shall take for granted that the results of this exposé extend to this more general situation.*

**Remark 1.14.** *Every morphism that is both a monomorphism and an effective epimorphism is an isomorphism.*

Indeed, in the notation of 1.3, the fact that  $T \rightarrow S$  is a monomorphism implies that the two morphisms

$$\text{pr}_1, \text{pr}_2 : T \times_S T \rightrightarrows T$$

are equal (and are isomorphisms). But a diagram

$$C \begin{array}{c} \xrightarrow{v} \\ \rightrightarrows \\ \xrightarrow{v} \end{array} B \xrightarrow{u} A$$

is exact only if  $u$  is an isomorphism, as follows immediately from the definition.<sup>7</sup>

## 2. Descent Morphisms

Recall the following definitions.

**Definition 2.1.** *Let  $f : S' \rightarrow S$  be a morphism such that  $S'' = S' \times_S S'$  exists, and let  $u' : X' \rightarrow S'$  be an object over  $S'$ . A gluing datum on  $X'/S'$ , relative to  $f$ , is an  $S''$ -isomorphism*

$$c : X''_1 \xrightarrow{\sim} X''_2,$$

---

<sup>6</sup>Editor's note: this is the following argument, communicated by M. Demazure. Let  $f, g : T' \rightarrow T$ , and let  $\phi : T' \rightarrow T \times_S T$  be the morphism with components  $f$  and  $g$ , so that  $p_1 \circ \phi = f$  and  $p_2 \circ \phi = g$ . Then  $F(\phi) : F(T \times_S T) \rightarrow F(T')$  satisfies  $F(\phi) \circ F(p_1) = F(f)$  and  $F(\phi) \circ F(p_2) = F(g)$ . For every  $x \in \check{H}^0(T/S, F)$ , however, one has  $F(p_1)(x) = F(p_2)(x)$ . Applying  $F(\phi)$  to both sides gives  $F(f)(x) = F(g)(x)$ , which shows that  $f$  and  $g$  induce the same map.

<sup>7</sup>Editor's note: a monomorphism that is an epimorphism need not be an isomorphism. For example, in  $\mathcal{C} = \mathbf{Sch}$ , the morphism

$$\text{Spec}(\mathbf{F}_p) \amalg \text{Spec}(\mathbf{Z}[1/p]) \longrightarrow \text{Spec}(\mathbf{Z})$$

is a monomorphism and a surjective epimorphism, but is not an isomorphism.

where  $X_i''$ , for  $i = 1, 2$ , denotes the inverse image, assumed to exist, of  $X'/S'$  under the projection  $\text{pr}_i : S'' \rightarrow S'$ . The gluing datum  $c$  is called a descent datum if it satisfies the “cocycle condition”

$$\text{pr}_{3,1}^*(c) = \text{pr}_{3,2}^*(c) \text{pr}_{2,1}^*(c),$$

where  $\text{pr}_{i,j}$ , for  $1 \leq j < i \leq 3$ , are the canonical projections from

$$S''' = S' \times_S S' \times_S S'$$

to  $S''$  (and we now assume that  $S'''$  also exists), where  $\text{pr}_{i,j}^*(c)$  is the inverse image of  $c$ , viewed as an  $S'''$ -morphism from  $X_j'''$  to  $X_i'''$ , and where, for each integer  $k$  from 1 to 3,  $X_k'''$  denotes the inverse image, assumed to exist, of  $X'/S'$  under the  $k$ -th projection  $q_k : S''' \rightarrow S'$ .

In the second part of the definition we have therefore used customary identifications and abuses of notation,<sup>8</sup> which experience shows to be harmless, but which must plainly be avoided in a rigorous exposition of descent theory—a theory that must itself justify, to some extent, these customary abuses of language. Such a formal exposition ([D]) was written by Giraud, with a view to justifying and making precise SGA 1, Exposé VII, which was never written. For a detailed account of the faithfully flat descent results used constantly in this Seminar, one may consult SGA 1, Exposé VIII.

Continue to let  $f : S' \rightarrow S$  be a morphism for which  $S'' = S' \times_S S'$  exists, and let  $X$  be an object over  $S$  such that  $X' = X \times_S S'$  and  $X'' = X \times_S S''$  exist. The inverse images of  $X'$  under  $\text{pr}_i$ , for  $i = 1, 2$ , then exist and are canonically isomorphic; consequently,  $X'/S'$  carries a canonical gluing datum relative to  $f$ . When  $S'''$  and  $X''' = X \times_S S'''$  exist, this is in fact a descent datum. If  $Y$  is another object over  $S$  satisfying the same conditions as  $X/S$ , then for every  $S$ -morphism  $X \rightarrow Y$ , the corresponding  $S'$ -morphism  $X' \rightarrow Y'$  is compatible with the canonical gluing data on  $X'$  and  $Y'$ . In particular, if  $S' \rightarrow S$  is squarable, then

$$X \mapsto X' = X \times_S S'$$

is a functor from  $\mathcal{C}_S$  to the “category of objects over  $S'$  equipped with a descent datum relative to  $f$ ”—a category whose definition is left to the reader and which is a full subcategory of the “category of objects over  $S'$  equipped with a gluing datum relative to  $f$ .”

With this understood, we make the following definition.

**Definition 2.2.** A morphism  $f : S' \rightarrow S$  is called a descent morphism (respectively, an effective descent morphism) if  $f$  is squarable (that is,  $X \times_S S'$  exists for every  $X \rightarrow S$ ) and the preceding functor

$$X \mapsto X' = X \times_S S'$$

from the category  $\mathcal{C}_S$  of objects over  $S$  to the category of objects over  $S'$  equipped with a descent datum relative to  $f$  is fully faithful (respectively, an equivalence of categories).

The first of these notions can be expressed using gluing data alone, and therefore without invoking the triple fiber product  $S'''$ :  $f$  is a descent morphism precisely when it is squarable and  $X \mapsto X'$  is a fully faithful functor from  $\mathcal{C}_S$  to the category of objects over  $S'$  equipped with a gluing datum relative to  $f$ . Written out, this means that for any two objects  $X, Y$  over  $S$ , the following diagram of sets is exact:

$$\text{Hom}_S(X, Y) \longrightarrow \text{Hom}_{S'}(X', Y') \begin{matrix} \xrightarrow{p_1} \\ \xrightarrow{p_2} \end{matrix} \text{Hom}_{S''}(X'', Y''). \quad (\text{x})$$

<sup>8</sup>Editor’s note: for example, we have identified, on the one hand,  $\text{pr}_{2,1}^*(X_1'') = \text{pr}_{3,1}^*(X_1'')$ , and, on the other hand,  $\text{pr}_{2,1}^*(X_2'') = \text{pr}_{3,2}^*(X_2'')$ .

Here  $p_i(u')$  denotes the inverse image of  $u' \in \text{Hom}_{S'}(X', Y')$  under the projection  $\text{pr}_i : S'' \rightarrow S'$ , for  $i = 1, 2$ . Indeed, by definition, the kernel of the pair  $(p_1, p_2)$  is precisely the set of  $S'$ -morphisms  $X' \rightarrow Y'$  compatible with the canonical gluing data.

By the definition of the inverse images  $Y'$  and  $Y''$ , there are canonical bijections

$$\text{Hom}_{S'}(X', Y') \simeq \text{Hom}_S(X', Y), \quad \text{Hom}_{S''}(X'', Y'') \simeq \text{Hom}_S(X'', Y).$$

Thus exactness of (x) is equivalent to exactness of

$$\text{Hom}_S(X, Y) \longrightarrow \text{Hom}_S(X', Y) \rightrightarrows \text{Hom}_S(X'', Y), \quad (\text{xx})$$

which is obtained by applying  $\text{Hom}_S(-, Y)$  to the diagram in  $\mathcal{C}/_S$

$$X'' \rightrightarrows X' \longrightarrow X, \quad (\text{xxx})$$

itself obtained from

$$S'' \rightrightarrows S' \longrightarrow S$$

by the base change  $X \rightarrow S$ . In view of 1.6, this proves the first part of the following proposition.

**Proposition 2.3.** *Let  $f : S' \rightarrow S$  be a morphism. If  $f$  is a universal effective epimorphism, then it is a descent morphism. Conversely, every descent morphism is a universal effective epimorphism if  $S$  is a squarable object of  $\mathcal{C}$ , that is, if its product with every object of  $\mathcal{C}$  exists.*

It remains to prove the converse. If  $f$  is a descent morphism, we must show that it is a universal effective epimorphism: for every morphism  $X \rightarrow S$ , diagram (xxx) must be exact, or equivalently, its image under  $\text{Hom}(-, Z)$  must be an exact diagram of sets for every object  $Z$  of  $\mathcal{C}$ . By hypothesis,  $Z \times S$  exists. Let  $Y$  be the object of  $\mathcal{C}/_S$  that it defines. The image of (xxx) under  $\text{Hom}(-, Z)$  is then isomorphic to its image under  $\text{Hom}_S(-, Y)$ , and the latter is exact by the hypothesis on  $f$ .

The results concerning descent morphisms may therefore be applied to universal effective epimorphisms. For example:

**Proposition 2.4.** *Let  $f : S' \rightarrow S$  be a descent morphism (for example, a universal effective epimorphism). Then:*

- (a) *For every  $S$ -morphism  $u : X \rightarrow Y$ , the morphism  $u$  is an isomorphism (respectively, a monomorphism) if and only if  $u' : X' \rightarrow Y'$  is one.*
- (b) *Let  $X, Y$  be two subobjects of  $S$ , and let  $X', Y'$  be their inverse images in  $S'$ . Then  $X$  is contained in  $Y$  (respectively, is equal to  $Y$ ) if and only if the same is true of  $X'$  and  $Y'$ .*

For (a), the definition shows that if  $u'$  is an isomorphism in the category of objects with gluing datum, then  $u$  is an isomorphism. Every isomorphism of objects over  $S'$  that is compatible with gluing data is, however, an isomorphism of objects with gluing datum: its inverse is also compatible with the gluing data. For (b), it remains to show that if  $X'$  is contained in  $Y'$ , that is, if there exists an  $S'$ -morphism  $X' \rightarrow Y'$ , then the same is true of  $X, Y$  over  $S$ . Since  $Y' \rightarrow S'$ , and hence  $Y'' \rightarrow S''$ , is a monomorphism, the morphism  $X' \rightarrow Y'$  is automatically compatible with the gluing data and therefore comes from an  $S$ -morphism  $X \rightarrow Y$ .

The proof applies more generally to two objects  $X, Y$  over  $S$ , with  $Y \rightarrow S$  a monomorphism, when one asks whether  $X \rightarrow S$  factors through  $Y$ : it is enough that  $X' \rightarrow S'$  factor through  $Y'$ .

**Corollary 2.5.** *Let  $f : S' \rightarrow S$  be a universal effective epimorphism and  $g : S \rightarrow T$  a morphism such that  $S \times_T S$  exists. Suppose that  $S'' = S' \times_S S'$  is also a fiber product of  $S'$  with itself over  $T$ , that is, that the canonical morphism*

$$S' \times_S S' \xrightarrow{\sim} S' \times_T S'$$

*is an isomorphism. Then  $g : S \rightarrow T$  is a monomorphism (and the converse is of course true).*

Indeed, consider the cartesian diagram

$$\begin{array}{ccc} S' \times_S S' & \xrightarrow{\sim} & S' \times_T S' \\ \downarrow & & \downarrow f \\ S & \longrightarrow & S \times_T S, \end{array} \quad (2.5.1)$$

where the second horizontal arrow is the diagonal morphism. By 1.9, the second vertical arrow  $f$  is a universal effective epimorphism; by hypothesis, the first horizontal arrow is an isomorphism. It follows from 2.4(a), or equally from 2.4(b),<sup>9</sup> that  $S \rightarrow S \times_T S$  is also an isomorphism, which says precisely that  $g : S \rightarrow T$  is a monomorphism.

**Remark 2.6.** *The notions introduced in 2.1 in terms of morphisms between certain inverse limits have an evident expression in terms of the contravariant functors defined by the objects  $S, S', X'$  under consideration. Subject to the existence of the fiber products considered in 2.1, there is a one-to-one correspondence between gluing data (respectively, descent data) on  $X'/S'$  relative to  $S' \rightarrow S$ , and gluing data (respectively, descent data) for the corresponding objects of*

$$\widehat{\mathcal{C}} = \text{Hom}(\mathcal{C}^\circ, \mathbf{Set}).$$

*This makes it possible, if desired, to extend these notions to the case in which no existence hypothesis is imposed on fiber products in  $\mathcal{C}$ .*

**Remark 2.7.** *The notions introduced in this section generalize to families of morphisms. They can also be presented more intrinsically by means of the notion of a sieve (4.1). For these matters the reader is referred to [D].*

### 3. Universal Effective Equivalence Relations

#### 3.1. Equivalence Relations: Definitions

**Definition 3.1.1.** *A  $\mathcal{C}$ -equivalence relation in an object  $X \in \text{Ob } \mathcal{C}$  is a representable subfunctor  $R$  of the functor  $X \times X$  such that, for every  $S \in \text{Ob } \mathcal{C}$ ,  $R(S)$  is the graph of an equivalence relation on  $X(S)$ .*

This definition applies in particular to the category  $\widehat{\mathcal{C}}$ . If  $X$  is regarded as an object of  $\widehat{\mathcal{C}}$ , then a  $\widehat{\mathcal{C}}$ -equivalence relation in  $X$  is simply a subfunctor  $R$  of  $X \times X$  such that  $R(S)$  is the graph of an equivalence relation on  $X(S)$ , for every object  $S$  of  $\mathcal{C}$ .<sup>10</sup>

If  $R$  is a  $\mathcal{C}$ -equivalence relation in  $X$ , let  $p_i$  denote the morphism from  $R$  to  $X$  induced by the projection  $\text{pr}_i : X \times X \rightarrow X$ . Thus there is a diagram

$$p_1, p_2 : R \rightrightarrows X.$$

<sup>9</sup>Editor's note: applied to  $f : S' \times_T S' \rightarrow S \times_T S = Y$ .

<sup>10</sup>Editor's note: the condition is plainly necessary. Conversely, if  $R(S)$  is the graph of an equivalence relation for every  $S \in \text{Ob } \mathcal{C}$ , this equivalence relation extends to  $R(F)$  for every  $F \in \text{Ob } \widehat{\mathcal{C}}$ : two morphisms  $\varphi, \psi : F \rightarrow R$  are declared equivalent if, for every  $S \in \text{Ob } \mathcal{C}$  and every  $x \in F(S)$ , the elements  $\varphi(x)$  and  $\psi(x)$  are equivalent in  $X(S)$ .

**Definition 3.1.2.** A morphism  $u : X \rightarrow Z$  is said to be compatible with  $R$  if  $up_1 = up_2$ . An object of  $\mathcal{C}$  that is a cokernel of the pair  $(p_1, p_2)$  is also called a quotient object of  $X$  by  $R$ , and is denoted  $X/R$ .

Thus one has an exact diagram

$$\begin{array}{ccc} R & \xrightarrow[p_2]{p_1} & X \xrightarrow{p} X/R, \end{array}$$

and  $X/R$  represents the covariant functor

$$\mathrm{Hom}_{\mathcal{C}}(X/R, Z) = \{\text{morphisms from } X \text{ to } Z \text{ compatible with } R\}.$$

Since quotient objects are taken in  $\mathcal{C}$ , the quotient  $X/R$  is unique whenever it exists.

These definitions extend at once to a  $\widehat{\mathcal{C}}$ -equivalence relation in  $X$ . One should note, however, that the identification of the objects of  $\mathcal{C}$  with their images in  $\widehat{\mathcal{C}}$  does not commute with the formation of quotients: the quotient  $X/R$  of  $X$  by  $R$  in  $\mathcal{C}$  is not *a priori* a quotient of  $X$  by  $R$  in  $\widehat{\mathcal{C}}$ . One must therefore avoid identifying  $\mathcal{C}$  uncritically with its image in  $\widehat{\mathcal{C}}$  in questions involving passage to a quotient.<sup>11</sup>

In what follows, we shall simply say *equivalence relation* to mean a  $\widehat{\mathcal{C}}$ -equivalence relation; whenever necessary, we shall specify that a  $\mathcal{C}$ -equivalence relation is intended.<sup>12</sup>

**Definition 3.1.3.** If  $X$  is an object of  $\mathcal{C}$  over  $S$ , an equivalence relation in  $X$  over  $S$  is an equivalence relation  $R$  in  $X$  for which the structure morphism  $X \rightarrow S$  is compatible with  $R$ .

The canonical monomorphism  $R \rightarrow X \times X$  then factors through the monomorphism

$$X \times_S X \longrightarrow X \times X$$

and defines an equivalence relation in the object  $X \rightarrow S$  of  $\mathcal{C}/_S$ . Whenever the quotient  $X/R$  exists, it is equipped with a canonical morphism to  $S$ , and the corresponding object of  $\mathcal{C}/_S$  is a quotient of  $X \in \mathrm{Ob} \mathcal{C}/_S$  by the preceding equivalence relation. Conversely, if  $S$  is a squarable object of  $\mathcal{C}$  and  $Y \rightarrow S$  is a quotient of  $X$  by this equivalence relation in  $\mathcal{C}/_S$ , then  $Y$  is a quotient of  $X$  by  $R$  in  $\mathcal{C}$ . In any event, we shall never have to consider quotients in  $\mathcal{C}/_S$  that are not already quotients in  $\mathcal{C}$ .

**Definition 3.1.4.** Let  $X$  (respectively  $X'$ ) be an object of  $\mathcal{C}$  equipped with an equivalence relation  $R$  (respectively  $R'$ ). A morphism

$$u : X \longrightarrow X'$$

is said to be compatible with  $R$  and  $R'$  if the following equivalent conditions hold:

- (i) for every  $S \in \mathrm{Ob} \mathcal{C}$ , two points of  $X(S)$  congruent modulo  $R(S)$  are carried by  $u$  to two points of  $X'(S)$  congruent modulo  $R'(S)$ ;

<sup>11</sup>Editor's note: here is an illustration that also previews the sequel of this exposé. Let  $G$  be a  $\mathcal{C}$ -group, let  $H$  be a  $\mathcal{C}$ -subgroup, and let  $R$  be the equivalence relation in  $G$  defined by  $G \times H \rightarrow G \times G$ ,  $(g, h) \mapsto (g, gh)$  (cf. 3.2). The functor  $Q$  defined by  $Q(S) = G(S)/H(S)$  is a quotient in  $\widehat{\mathcal{C}}$  (by 4.4.9 applied to the coarsest topology; cf. 4.4.2), but it is not in general the quotient one wants. For example, when  $\mathcal{C} = \mathbf{Sch}$ , there is an exact sequence of affine group schemes

$$1 \longrightarrow \mu_2 \longrightarrow \mathbf{G}_m \xrightarrow{p} \mathbf{G}_m \longrightarrow 1$$

that identifies  $\mathbf{G}_m$  with the quotient  $\mathbf{G}_m/\mu_2$ . Moreover, because  $p$  is finite and locally free,  $\mathbf{G}_m$  is the sheaf quotient of  $\mathbf{G}_m$  by  $\mu_2$  in the larger category of sheaves for the fppf topology; cf. 4.6.6(ii) and 6.3.2. By contrast, the quotient  $Q$  in  $\widehat{\mathcal{C}}$  is not isomorphic to  $\mathbf{G}_m$ , since, for example,  $Q(\mathbf{Z}) = \{1\}$ , whereas  $\mathbf{G}_m(\mathbf{Z}) = \{\pm 1\}$ . Thus  $Q$  is not an fppf sheaf and, *a fortiori*, is not representable.

<sup>12</sup>Editor's note: even for a  $\widehat{\mathcal{C}}$ -equivalence relation in  $X$ , the question of interest is the existence of a quotient in  $\mathcal{C}$ .

(ii) *there exists a necessarily unique morphism  $R \rightarrow R'$  making the diagram*

$$\begin{array}{ccc} R & \longrightarrow & R' \\ \downarrow & & \downarrow \\ X \times X & \xrightarrow{u \times u} & X' \times X' \end{array}$$

*commute.*

By the universal property of  $X/R$ , whenever  $X/R$  and  $X'/R'$  exist there is a unique morphism  $v$  making the diagram

$$\begin{array}{ccc} X & \xrightarrow{p} & X/R \\ u \downarrow & & \downarrow v \\ X' & \xrightarrow{p'} & X'/R' \end{array}$$

commute.

**Definition 3.1.5.** *A subobject (that is, a representable subfunctor)  $Y$  of  $X$  is said to be stable under the equivalence relation  $R$  if the following equivalent conditions hold:*

- (i) *for every  $S \in \text{Ob } \mathcal{C}$ , the subset  $Y(S)$  of  $X(S)$  is stable under  $R(S)$ ;*
- (ii) *the two subobjects of  $R$  that are the inverse images of  $Y$  under  $p_1$  and  $p_2$  are identical.*

An important special case of a stable subobject of  $X$  is the following: the quotient  $X/R$  exists and  $Y$  is the inverse image in  $X$  of a subobject of  $X/R$ .

**Definition 3.1.6.** *Let  $R$  be an equivalence relation in  $X$ , and let  $X' \rightarrow X$  be a morphism. The equivalence relation  $R'$  in  $X'$  obtained from the cartesian diagram*

$$\begin{array}{ccc} R' & \longrightarrow & R \\ \downarrow & & \downarrow \\ X' \times X' & \longrightarrow & X \times X \end{array}$$

*is called the equivalence relation in  $X'$  that is the inverse image of the equivalence relation  $R$  in  $X$ . In particular, if  $X'$  is a subobject of  $X$ , it is called the equivalence relation induced on  $X'$  by  $R$ , and is denoted  $R_{X'}$ .*

The morphism  $X' \rightarrow X$  is compatible with  $R'$  and  $R$ ; hence, when the quotients exist, it induces a morphism  $X'/R' \rightarrow X/R$  (3.1.4). If  $X'$  is a subobject of  $X$ , we shall see later that in certain cases one can prove that  $X'/R' \rightarrow X/R$  is a monomorphism and therefore identifies  $X'/R'$  with a subobject of  $X/R$ . In that situation, the inverse image of this subobject in  $X$  is a subobject containing  $X'$  and stable under  $R$ : the *saturation* of  $X'$  with respect to the equivalence relation  $R$ .

**Proposition 3.1.7.** *If the subobject  $Y$  of  $X$  is stable under  $R$ , then for  $i = 1, 2$  the following two squares are cartesian:*

$$\begin{array}{ccc} R_Y & \longrightarrow & R \\ p_i \downarrow & & \downarrow p_i \\ Y & \longrightarrow & X. \end{array}$$

The proof is immediate.



### 3.2. Equivalence Relation Defined by a Group Acting Freely

**Definition 3.2.1.** Let  $X$  be an object of  $\mathcal{C}$ , and let  $H$  be a  $\mathcal{C}$ -group acting on  $X$ , that is, equipped with a morphism of  $\mathcal{C}$ -groups

$$H \longrightarrow \text{Aut}(X).$$

The group  $H$  is said to act freely on  $X$  if the following equivalent conditions hold:

- (i) for every  $S \in \text{Ob } \mathcal{C}$ , the group  $H(S)$  acts freely on  $X(S)$ ;
- (ii) the morphism of functors

$$H \times X \longrightarrow X \times X$$

defined on sets by  $(h, x) \mapsto (hx, x)$  is a monomorphism.

In the preceding definition the group  $H$  was assumed to act on  $X$  on the left. There is of course an analogous notion when the group acts on the right, that is, when one is given a group morphism from the opposite group  $H^\circ$  to  $\text{Aut}(X)$ .

If  $H$  acts freely on  $X$ , the image of  $H \times X$  under the morphism in (ii) is an equivalence relation in  $X$ , called the *equivalence relation defined by the action of  $H$  on  $X$* . When the quotient of  $X$  by this equivalence relation exists, it is denoted  $H \backslash X$  (or  $X/H$  when  $H$  acts on the right). It represents the following covariant functor: for every object  $Z$  of  $\mathcal{C}$ ,

$$\text{Hom}(H \backslash X, Z) = \{\text{morphisms from } X \text{ to } Z \text{ invariant under } H\},$$

where a morphism  $f : X \rightarrow Z$  is said to be invariant under  $H$  if, for every  $S \in \text{Ob } \mathcal{C}$ , the corresponding morphism  $X(S) \rightarrow Z(S)$  is invariant under the group  $H(S)$ .

**Lemma 3.2.2.** Under the assumptions of 3.2.1, let  $Y$  be a subobject of  $X$ . The following conditions are equivalent:

- (i)  $Y$  is stable under the equivalence relation defined by  $H$  (3.1.5);
- (ii) for every  $S \in \text{Ob } \mathcal{C}$ , the subset  $Y(S)$  of  $X(S)$  is stable under  $H(S)$ ;
- (iii) there exists a necessarily unique morphism  $f$  making the diagram

$$\begin{array}{ccc} H \times Y & \xrightarrow{f} & Y \\ \downarrow & & \downarrow \\ H \times X & \longrightarrow & X \end{array}$$

commute.

Under these conditions,  $f$  defines a morphism of  $\widehat{\mathcal{C}}$ -groups

$$H \longrightarrow \text{Aut}(Y),$$

and the equivalence relation in  $Y$  defined by this action of  $H$  is precisely the equivalence relation induced on  $Y$  by the equivalence relation in  $X$  defined by the action of  $H$ .

The proof is immediate. There is of course an analogous statement for a right action. The action of  $H$  on  $Y$  defined above will be called the action induced on  $Y$  by the given action of  $H$  on  $X$ .

Consider now two  $\mathcal{C}$ -groups  $H$  and  $G$ , together with a group morphism

$$u : H \longrightarrow G.$$

Then  $H$  acts on  $G$  by translations (set-theoretically,  $hg = u(h)g$ ), and it acts freely if and only if  $u$  is a monomorphism. When it exists, the quotient of  $G$  by this action of  $H$  is denoted  $H \backslash G$ . Similarly, one defines a right action of  $H$  on  $G$  and a quotient  $G/H$ . These quotients are functorial with respect to the groups involved. More precisely, one has the following lemma, stated for right quotients.

**Lemma 3.2.3.** *Let  $u : H \rightarrow G$  and  $u' : H' \rightarrow G'$  be monomorphisms of  $\mathcal{C}$ -groups, and suppose that a morphism of  $\mathcal{C}$ -groups*

$$f : G \longrightarrow G'$$

*is given. The following conditions are equivalent:*

- (i)  *$f$  is compatible with the equivalence relations in  $G$  and  $G'$  defined by  $H$  and  $H'$ ;*
- (ii) *for every  $S \in \text{Ob } \mathcal{C}$ , one has  $fu(H(S)) \subseteq u'(H'(S))$ ;*
- (iii) *there exists a necessarily unique multiplicative morphism  $g : H \rightarrow H'$  such that the diagram*

$$\begin{array}{ccc} H & \xrightarrow{g} & H' \\ u \downarrow & & \downarrow u' \\ G & \xrightarrow{f} & G' \end{array}$$

*commutes.*

*Under these conditions, if the quotients  $G/H$  and  $G'/H'$  exist, there is a unique morphism  $\bar{f}$  making the diagram*

$$\begin{array}{ccc} G & \xrightarrow{f} & G' \\ p \downarrow & & \downarrow p' \\ G/H & \xrightarrow{\bar{f}} & G'/H' \end{array}$$

*commute.*

The first assertion is proved by reduction to the set-theoretic case. The second follows immediately from (i).

The notions introduced above for general equivalence relations could all be translated into the present setting. We record only the following lemma, whose proof is immediate by reduction to the set-theoretic case.

**Lemma 3.2.4.** *Let  $u : H \rightarrow G$  be a monomorphism of  $\mathcal{C}$ -groups, and let  $G'$  be a  $\mathcal{C}$ -subgroup of  $G$ . The subobject  $G'$  of  $G$  is stable under the equivalence relation defined by  $H$  if and only if  $u$  factors through the canonical monomorphism  $G' \rightarrow G$ . Under this condition, the action of  $H$  on  $G'$  induced by the action on  $G$  defined by  $u$  is precisely the action arising from the monomorphism  $H \rightarrow G'$  through which  $u$  factors.*

### 3.3. Universally Effective Equivalence Relations

**Definition 3.3.1.** *Let  $f : X \rightarrow Y$  be a morphism. The equivalence relation defined by  $f$  in  $X$ , denoted  $\mathcal{R}(f)$ , is the  $\mathcal{C}$ -equivalence relation in  $X$  that is the image of the canonical monomorphism*

$$X \times_Y X \longrightarrow X \times X.$$

**Definition 3.3.2.** *An equivalence relation  $R$  in  $X$  is said to be effective if:*

- (i)  *$R$  is representable, that is, it is a  $\mathcal{C}$ -equivalence relation;*

(ii) the quotient  $Y = X/R$  exists in  $\mathcal{C}$ ;<sup>13</sup>

(iii) the diagram

$$\begin{array}{ccc} & p_1 & \\ R & \rightrightarrows & X \xrightarrow{p} Y \\ & p_2 & \end{array}$$

makes  $R$  the fiber square of  $X$  over  $Y$ , that is,  $R$  is the equivalence relation defined by  $p$ .

**Scholium 3.3.2.1.**<sup>14</sup> If  $R$  is an effective equivalence relation in  $X$ , then  $p$  is an effective epimorphism (1.3). If  $f : X \rightarrow Y$  is an effective epimorphism, then  $\mathcal{R}(f)$  is an effective equivalence relation in  $X$ , one quotient of which is  $Y$ . Thus there is a one-to-one “Galois” correspondence between effective equivalence relations in  $X$  and effective quotients of  $X$ , that is, equivalence classes of effective epimorphisms with source  $X$ .

**Definition 3.3.3.** An equivalence relation  $R$  in  $X$  is said to be *universally effective* if the quotient  $Y = X/R$  exists and, for every  $Y' \rightarrow Y$ , the fiber products  $X' = X \times_Y Y'$  and  $R' = R \times_Y Y'$  exist and  $R'$  is the fiber square of  $X'$  over  $Y'$ . Equivalently,  $R$  is effective and  $p : X \rightarrow X/R$  is a universal effective epimorphism.

**Scholium 3.3.3.1.**<sup>14</sup> As above, there is a one-to-one correspondence between universally effective equivalence relations in  $X$  and universal effective quotients of  $X$ .

**Remark 3.3.3.2.**<sup>14</sup> Suppose that  $\mathcal{C}$  is the category of  $S$ -schemes, and write  $\mathbf{A}_S^1$  for affine one-space over  $S$ . Let  $R \subset X \times_S X$  be a universally effective equivalence relation, and let  $p : X \rightarrow Y$  be its quotient. Then, for every open subset  $U$  of  $Y$ ,

$$\mathcal{O}(U) = \text{Hom}_S(U, \mathbf{A}_S^1)$$

is the set of elements  $\varphi$  of

$$\mathcal{O}(p^{-1}(U)) = \text{Hom}_S(p^{-1}(U), \mathbf{A}_S^1)$$

such that  $\varphi \circ \text{pr}_1 = \varphi \circ \text{pr}_2$ . In particular, if  $R$  is defined by the action of a group  $H$  acting freely on the right on  $X$  (cf. 3.2.1), then  $\mathcal{O}(U)$  is the set of all  $\varphi \in \mathcal{O}(p^{-1}(U))$  such that

$$\varphi(xh) = \varphi(x)$$

for every  $S' \rightarrow S$ ,  $x \in X(S')$ , and  $h \in H(S')$ .

**Proposition 3.3.4.** Let  $R$  be a universally effective equivalence relation in  $X$ . Let  $f : X \rightarrow Z$  be a morphism compatible with  $R$ , hence factoring through a morphism  $g : X/R \rightarrow Z$ . The following conditions are equivalent:

- (i)  $g$  is a monomorphism;
- (ii)  $R$  is the equivalence relation defined by  $f$ .

Indeed, (i) plainly implies (ii), while the converse is precisely 2.5.

**Definition 3.3.5.** Let  $H$  be a  $\mathcal{C}$ -group acting freely on  $X$ . One says that  $H$  acts *effectively*, or that the action of  $H$  on  $X$  is *effective* (respectively *universally effective*), if the equivalence relation in  $X$  defined by the action of  $H$  is effective (respectively *universally effective*).

<sup>13</sup>Editor’s note: the words “in  $\mathcal{C}$ ” have been added.

<sup>14</sup>Editor’s note: the numbering 3.3.2.1 (respectively 3.3.3.1) has been added for later reference; Remark 3.3.3.2 has also been added.

### 3.4. $(M)$ -Effectivity

In practice, universal effective epimorphisms are most often difficult to characterize. Nevertheless, one frequently has at hand a number of morphisms of this kind—in scheme theory, for example, faithfully flat quasi-compact morphisms. This leads to the developments below.

**3.4.1.** We first state several conditions on a family  $(M)$  of morphisms of  $\mathcal{C}$ :

- (a)  $(M)$  is stable under base change: every  $u : T \rightarrow S$  belonging to  $(M)$  is squarable (cf. 1.4.0), and for every  $S' \rightarrow S$ , the base change

$$u' : T \times_S S' \longrightarrow S'$$

belongs to  $(M)$ ;

- (b) the composite of two elements of  $(M)$  belongs to  $(M)$ ;  
 (c) every isomorphism belongs to  $(M)$ ;  
 (d) every element of  $(M)$  is an effective epimorphism.

Conditions (a) and (b) imply:

- (a') the cartesian product of two elements of  $(M)$  belongs to  $(M)$ . More precisely, let  $u : X \rightarrow Y$  and  $u' : X' \rightarrow Y'$  be two  $S$ -morphisms belonging to  $(M)$ . If  $Y \times_S Y'$  exists, then  $X \times_S X'$  exists and  $u \times_S u'$  belongs to  $(M)$ .

This follows from the diagram

$$\begin{array}{ccccc}
 & & Y' & \xleftarrow{u'} & X' \\
 & & \uparrow & & \uparrow \\
 X & \longleftarrow & X \times_S Y' & \longleftarrow & X \times_S X' \\
 u \downarrow & & \downarrow & & \swarrow \\
 Y & \longleftarrow & Y \times_S Y' & & u \times_S u'
 \end{array}$$

Likewise, (a) and (d) imply:

- (d') every element of  $(M)$  is a universal effective epimorphism.

**3.4.2.** The family  $(M_0)$  of universal effective epimorphisms satisfies conditions (a)–(d) of 3.4.1. Indeed, (a), (c), and (d) hold by definition, and (b) follows from 1.8. Henceforth we assume given a family  $(M)$  of morphisms of  $\mathcal{C}$  satisfying these conditions; our results apply in particular to the family  $(M_0)$ .

**Definition 3.4.3.** An equivalence relation  $R$  in  $X$  is said to be *of type  $(M)$*  if it is representable and  $p_1 \in (M)$ ; conditions (b) and (c) then imply that  $p_2 \in (M)$ .

The relation  $R$  is said to be  *$(M)$ -effective* if it is effective and the canonical morphism  $X \rightarrow X/R$  belongs to  $(M)$ .

A quotient  $Y$  of  $X$  is said to be  *$(M)$ -effective* if the canonical morphism  $X \rightarrow Y$  belongs to  $(M)$ .

The following consequences are immediate from the definition.<sup>15</sup>

#### Proposition 3.4.3.1.

<sup>15</sup>Editor's note: the numbering 3.4.3.1 has been added for later reference, and the proof of part (i) has been expanded.

- (i) An  $(M)$ -effective equivalence relation is of type  $(M)$  and is universally effective.
- (ii) An  $(M)$ -effective quotient is universally effective (cf. 3.3.3).
- (iii) The maps  $R \mapsto X/R$  and  $p \mapsto \mathcal{R}(p)$  establish a one-to-one correspondence between  $(M)$ -effective equivalence relations in  $X$  and  $(M)$ -effective quotients of  $X$ .
- (iv) To be  $(M_0)$ -effective is equivalent to being universally effective.

We prove (i). Since  $R$  is  $(M)$ -effective, there is a cartesian square

$$\begin{array}{ccc} R & \xrightarrow{p_2} & X \\ p_1 \downarrow & & \downarrow p \\ X & \xrightarrow{p} & X/R, \end{array}$$

and  $p \in (M)$ . By 3.4.1(a),  $p_1$  and  $p_2$  belong to  $(M)$ , so  $R$  is of type  $(M)$ .

Set  $Y = X/R$ , and let  $Y' \rightarrow Y$  be any morphism. By 3.4.1(a), the fiber products  $X' = X \times_Y Y'$  and  $R' = R \times_Y Y'$  exist, and the morphisms  $X' \rightarrow Y'$  and  $p'_i : R' \rightarrow X'$  belong to  $(M)$ . Finally, since  $R = X \times_Y X$ , associativity of the fiber product gives

$$R' = X \times_Y X \times_Y Y' = X' \times_{Y'} X'.$$

This shows that  $R'$  is  $(M)$ -effective; in particular,  $R$  is universally effective. This proves (i), as well as (iv). Parts (ii) and (iii) follow, taking Definition 3.3.2 into account.

**3.4.4.** Let  $H$  be an  $S$ -group whose structure morphism belongs to  $(M)$ . If  $H$  acts freely on the  $S$ -object  $X$ , then it defines an equivalence relation of type  $(M)$  in  $X$ . Indeed, by (a), the fiber product  $H \times_S X$  exists and  $\text{pr}_2 : H \times_S X \rightarrow X$  belongs to  $(M)$ . The action of  $H$  is said to be  $(M)$ -effective if the equivalence relation in  $X$  defined by this action is  $(M)$ -effective.

**Proposition 3.4.5 (( $M$ )-effectivity and base change).** *Let  $R$  be an  $(M)$ -effective equivalence relation in  $X$  over  $S$ , and set  $Y = X/R$ . Let  $S' \rightarrow S$  be a base change for which  $Y' = Y \times_S S'$  exists. Then  $X' = X \times_S S'$  exists,  $R' = R \times_S S'$  is an  $(M)$ -effective equivalence relation in  $X'$  over  $S'$ , and*

$$X'/R' \simeq (X/R)'.$$

Indeed, the canonical morphisms  $X \rightarrow Y$  and  $R \rightarrow Y$  belong to  $(M)$ ; hence, by (a'),  $X'$  and  $R'$  are representable. By associativity of the fiber product,  $R'$  is the equivalence relation in  $X'$  defined by the canonical morphism  $X' \rightarrow Y'$ , which belongs to  $(M)$ . This proves the assertion.

**Proposition 3.4.6 (( $M$ )-effectivity and cartesian products).** *Let  $R$  (respectively  $R'$ ) be an  $(M)$ -effective equivalence relation in  $X$  (respectively  $X'$ ) over  $S$ . If  $(X/R) \times_S (X'/R')$  exists, then  $X \times_S X'$  exists,  $R \times_S R'$  is an  $(M)$ -effective equivalence relation in  $X \times_S X'$  over  $S$ , and*

$$(X \times_S X')/(R \times_S R') \simeq (X/R) \times_S (X'/R').$$

Set  $Y = X/R$  and  $Y' = X'/R'$ . By (a'), the fiber product  $X \times_S X'$  exists and the canonical morphism

$$q : X \times_S X' \longrightarrow Y \times_S Y'$$

belongs to  $(M)$ . The formula

$$(X \times_S X') \times_{(Y \times_S Y')} (X \times_S X') \simeq (X \times_Y X) \times_S (X' \times_{Y'} X')$$

(all unlabelled fiber products being taken over  $S$ ) shows that  $R \times_S R'$  is the equivalence relation in  $X \times_S X'$  defined by  $q$ , which completes the proof.

**3.4.7.**<sup>16</sup> Suppose that  $\mathcal{C}$  has a final object  $e$ , and let  $f : G \rightarrow G'$  be a morphism of  $\mathcal{C}$ -groups belonging to  $(M)$ . By 3.4.1(a), the kernel  $\text{Ker}(f)$  is represented by  $e \times_{G'} G$ , and the morphism  $\text{Ker}(f) \rightarrow e$  belongs to  $(M)$ .

On the other hand, the equivalence relation defined by  $f$  is the same as that defined by the action of  $\text{Ker}(f)$ , say on the right, on  $G$ : it is the image of the morphism

$$G \times \text{Ker}(f) \longrightarrow G \times G$$

defined on sets by  $(g, h) \mapsto (g, gh)$ . Consequently, Scholium 3.3.2.1 gives the following corollary.

**Corollary 3.4.7.1.** *Suppose that  $\mathcal{C}$  has a final object  $e$ , and let  $f : G \rightarrow G'$  be a morphism of  $\mathcal{C}$ -groups belonging to  $(M)$ . Then the action of  $\text{Ker}(f)$  on  $G$  is  $(M)$ -effective, and  $G'$  is the quotient  $G/\text{Ker}(f)$ .*

### 3.5. Construction of Quotients by Descent

It frequently happens that one cannot construct a quotient directly, but can do so after a suitable base change. This subsection gives a useful criterion for that situation.

In 2.1 we defined a descent datum on an object  $X'$  over  $S'$ , relative to a morphism  $S' \rightarrow S$ .

**Definition 3.5.1.** *A descent datum on  $X'$ , relative to  $S' \rightarrow S$ , is said to be effective if  $X'$ , equipped with this descent datum, is isomorphic to the inverse image over  $S'$  of an object  $X$  over  $S$ , equipped with its canonical descent datum.*

If  $S' \rightarrow S$  is a descent morphism (2.2), then the object  $X$  in the definition is unique up to unique isomorphism. To say that  $S' \rightarrow S$  is an effective descent morphism (2.2) means that it is a descent morphism and that every descent datum relative to it is effective.

Now consider an equivalence relation  $R$  in an object  $X$  over  $S$ . Let  $X'$ ,  $X''$ , and  $X'''$  be the inverse images of  $X$  over

$$S', \quad S'' = S' \times_S S', \quad S''' = S' \times_S S' \times_S S',$$

respectively, and let  $R'$ ,  $R''$ , and  $R'''$  be the equivalence relations obtained from  $R$  by inverse image. Suppose that  $R'$  in  $X'$  is  $(M)$ -effective, and consider the quotient  $Y' = X'/R'$ , an object over  $S'$ . By 3.4.5, its two inverse images over  $S''$  are isomorphic to  $X''/R''$ . Thus the  $S'$ -object  $Y'$  has a canonical gluing datum. Applying in the same way the uniqueness of  $X'''/R'''$ , one sees that this is in fact a descent datum. (In this argument it has been tacitly assumed that all the displayed fiber products exist. This holds, in particular, if  $S' \rightarrow S$  is squarable, for example if it is a descent morphism.)

**Proposition 3.5.2.** *Let  $R$  be an equivalence relation in the object  $X$  over  $S$ , and let  $S' \rightarrow S$  be a universal effective epimorphism. Assume that every  $S$ -morphism whose inverse image over  $S'$  belongs to  $(M)$  itself belongs to  $(M)$ . The following conditions are equivalent:*

- (i)  $R$  is  $(M)$ -effective in  $X$ ;
- (ii)  $R'$  is  $(M)$ -effective in  $X'$ , and the canonical descent datum on  $X'/R'$  is effective.

---

<sup>16</sup>Editor's note: this paragraph has been added.

When these conditions hold, the object descended from  $X'/R'$  is canonically isomorphic to  $X/R$ .

The implication (i)  $\Rightarrow$  (ii) is simply 3.4.4 translated into the language of descent. Once the converse is proved, the last assertion of the proposition follows from the fact that a universal effective epimorphism is a descent morphism (2.3), and hence the descended object is unique up to unique isomorphism.

We prove (ii)  $\Rightarrow$  (i). Let  $Y' = X'/R'$ , and let  $Y$  be the descended object. The canonical morphism  $p' : X' \rightarrow X'/R' = Y'$  is compatible by construction with the descent data: its two inverse images over  $S''$  coincide with the canonical morphism  $X'' \rightarrow X''/R''$ . It is therefore the inverse image over  $S'$  of an  $S$ -morphism  $p : X \rightarrow Y$ . Since  $p'$  belongs to  $(M)$ , the assumption on  $S' \rightarrow S$  implies that  $p$  also belongs to  $(M)$ . Moreover, because  $p'$  is compatible with the equivalence relation  $R'$ , the morphism  $p$  is compatible with  $R$ , again because a universal effective epimorphism is a descent morphism. Hence there is a morphism

$$R \longrightarrow X \times_Y X.$$

To prove that  $R$  is  $(M)$ -effective and that  $Y$  is isomorphic to  $X/R$ , it suffices to prove that this morphism is an isomorphism. After base change from  $S$  to  $S'$ , it becomes an isomorphism because  $R'$  is effective; it is therefore an isomorphism for the same reason as in 2.4.

Notice that the hypothesis in the proposition is satisfied if  $(M)$  is the family  $(M_0)$  of universal effective epimorphisms and if  $\mathcal{C}$  has fiber products (1.10). We obtain the following.

**Corollary 3.5.3.** *Suppose that  $\mathcal{C}$  has fiber products (fiber products over  $S$  would suffice). Let  $R$  be an equivalence relation in  $X$  over  $S$ , and let  $S' \rightarrow S$  be a universal effective epimorphism. The following conditions are equivalent:*

- (i)  $R$  is universally effective in  $X$ ;
- (ii)  $R'$  is universally effective in  $X'$ , and the canonical descent datum on  $X'/R'$  is effective.

When these conditions hold, the object descended from  $X'/R'$  is canonically isomorphic to  $X/R$ .

## 4. Topologies and Sheaves

The notion of a sieve, and the presentation of the notion of a topology adopted here (4.2.1), which is more intrinsic and in many respects more convenient than the presentation by covering families in [MA], are due to J. Giraud [AS].

### 4.1. Sieves

**Definition 4.1.1.** *A sieve of the category  $\mathcal{C}$  is a subfunctor  $C$  of the final functor*

$$e : \mathcal{C}^\circ \longrightarrow \mathbf{Set}.$$

To every sieve  $C$  of  $\mathcal{C}$  one associates the set  $E(C)$  of objects  $X$  of  $\mathcal{C}$  such that  $C(X) \neq \emptyset$ , that is, such that the structural morphism  $X \rightarrow e$  factors through  $C$ . Thus

$$\begin{cases} X \in E(C) & \Longleftrightarrow & C(X) = e(X) = \{\emptyset\}, \\ X \notin E(C) & \Longleftrightarrow & C(X) = \emptyset. \end{cases} \quad (+)$$

The set  $E = E(C)$  has the following property:

$$X \in E \text{ and } \text{Hom}(Y, X) \neq \emptyset \implies Y \in E. \quad (++)$$

(If the set  $\text{Ob } \mathcal{C}$  is endowed with its natural preorder, with  $Y$  dominating  $X$  when there is an arrow from  $Y$  to  $X$ , then the sets  $E$  satisfying  $(++)$  are precisely the subsets of  $\text{Ob } \mathcal{C}$  that contain every object dominating one of their elements.)<sup>17</sup>

Conversely, if  $E$  is a subset of  $\text{Ob } \mathcal{C}$  having property  $(++)$ , then  $E$  can be written uniquely in the form  $E(C)$ , with  $C$  defined by the formulas  $(+)$ . Thus sieves of  $\mathcal{C}$  correspond bijectively to subsets of  $\text{Ob } \mathcal{C}$  satisfying  $(++)$ . By abuse of language, we shall sometimes call the set  $E(C)$  itself a sieve of  $\mathcal{C}$ .

Let  $C$  and  $C'$  be two sieves of  $\mathcal{C}$ .<sup>18</sup> Since both are subfunctors of the final functor  $e$ , it is equivalent to say that  $C$  dominates  $C'$  (for the domination relation in  $\text{Ob } \widehat{\mathcal{C}}$ ), that  $C$  is a subfunctor of  $C'$ , or that  $E(C) \subset E(C')$ . In this case we say that  $C$  is *finer* than  $C'$ . This is an order relation on the set of sieves of  $\mathcal{C}$ . Moreover,

$$E(C) \cap E(C') = E(C \times C'),$$

so the set of sieves of  $\mathcal{C}$  is filtered for the order relation “is finer than.”

Every subset  $E$  of  $\text{Ob } \widehat{\mathcal{C}}$ , and in particular every subset of  $\text{Ob } \mathcal{C}$ , defines a sieve  $C(E)$ : the set of  $X \in \text{Ob } \mathcal{C}$  such that  $F(X) \neq \emptyset$  for at least one  $F \in E$  satisfies  $(++)$ , and hence defines the desired sieve.

This sieve may also be defined as the image of the family of morphisms  $\{F \rightarrow e \mid F \in E\}$ , in the sense of the following definition.

**Definition 4.1.2.** Let  $\{F_i \rightarrow F\}$  be a family of morphisms of  $\widehat{\mathcal{C}}$  with common target  $F$ . The image of this family is the subfunctor of  $F$  defined by

$$S \longmapsto \bigcup_i \text{Im } F_i(S) \subset F(S).$$

**Proposition 4.1.3.** Formation of the image commutes with base change. In the preceding notation let  $I$  be the image of the family  $\{F_i \rightarrow F\}$ . For every morphism  $G \rightarrow F$  in  $\widehat{\mathcal{C}}$ , the image of the family

$$\{F_i \times_F G \longrightarrow G\}$$

is the subfunctor  $I \times_F G$  of  $G$ .

**Definition 4.1.4.0.**<sup>19</sup> Let  $C$  be a sieve of  $\mathcal{C}$ . If  $E$  is a subset of  $\text{Ob } \mathcal{C}$  such that  $C(E) = C$ , then  $E$  is called a *base* of  $C$ . Every sieve  $C$  has a base, for example  $E(C)$ .

We now describe  $\text{Hom}(C, F)$ , where  $C$  is a sieve of  $\mathcal{C}$  and  $F \in \text{Ob } \widehat{\mathcal{C}}$ , by means of a base  $\{S_i\}$  of  $C$ . For every pair  $(i, j)$  there is a diagram in  $\widehat{\mathcal{C}}$ :

$$\begin{array}{ccc} S_i \times S_j & \longrightarrow & S_i \\ \downarrow & & \downarrow \\ S_j & \longrightarrow & e. \end{array}$$

<sup>17</sup>Editor's note: Here “dominating” is meant in the sense of the preorder just mentioned:  $Y$  dominates  $X$  if there is an arrow  $Y \rightarrow X$ . On the other hand, if  $X, Y$  are two subobjects of an object  $Z$ , one says (cf. 2.4) that  $Y$  contains  $X$  if  $X \subset Y$ . To avoid any ambiguity between these usages, the sequel uses “dominating” in the first case and “containing” in the second.

<sup>18</sup>Editor's note: The following sentence has been expanded.

<sup>19</sup>Editor's note: The numbering 4.1.4.0 has been added to highlight this definition.



It gives a diagram of sets

$$\Gamma(F) = \text{Hom}(e, F) \xrightarrow{\sigma} \prod_i \text{Hom}(S_i, F) \xrightarrow[\tau_2]{\tau_1} \prod_{i,j} \text{Hom}(S_i \times S_j, F),$$

with  $\tau_1\sigma = \tau_2\sigma$ . Hence there is a morphism

$$\text{Hom}(e, F) \longrightarrow \text{Ker} \left( \prod_i \text{Hom}(S_i, F) \rightrightarrows \prod_{i,j} \text{Hom}(S_i \times S_j, F) \right).$$

One checks immediately:

**Proposition 4.1.4.** *There is an isomorphism, functorial in  $F$ ,*

$$\text{Hom}(C, F) \xrightarrow{\sim} \text{Ker} \left( \prod_i \text{Hom}(S_i, F) \rightrightarrows \prod_{i,j} \text{Hom}(S_i \times S_j, F) \right),$$

for which the diagram

$$\begin{array}{ccc} \text{Hom}(e, F) & \longrightarrow & \text{Ker} \left( \prod_i \text{Hom}(S_i, F) \rightrightarrows \prod_{i,j} \text{Hom}(S_i \times S_j, F) \right) \\ \parallel & & \uparrow \sim \\ \text{Hom}(e, F) & \longrightarrow & \text{Hom}(C, F) \end{array}$$

is commutative; the bottom arrow is induced by the canonical morphism  $C \rightarrow e$ .

**Corollary 4.1.5.** *Suppose that the fiber products  $S_i \times S_j$  are representable, for example that the  $S_i$  are squarable. Then, for every  $F \in \text{Ob } \mathcal{C}$ , there is an isomorphism*

$$\text{Hom}(C, F) \xrightarrow{\sim} \text{Ker} \left( \prod_i F(S_i) \rightrightarrows \prod_{i,j} F(S_i \times S_j) \right).$$

**Remark 4.1.6.** Let  $R$  be a sieve of  $\mathcal{C}$ , let  $\overline{R}$  denote the full subcategory of  $\mathcal{C}$  whose object set is  $E(R)$ , and let

$$i_R : \overline{R} \longrightarrow \mathcal{C}$$

be the inclusion functor. There is an isomorphism, functorial in  $F \in \text{Ob } \widehat{\mathcal{C}}$ ,

$$\text{Hom}(R, F) \xrightarrow{\sim} \Gamma(F \circ i_R),$$

for which the diagram

$$\begin{array}{ccc} \text{Hom}(e, F) & \longrightarrow & \text{Hom}(R, F) \\ \downarrow \sim & & \downarrow \sim \\ \Gamma(F) & \longrightarrow & \Gamma(F \circ i_R) \end{array}$$

is commutative; the bottom arrow is induced by  $i_R$ .

**Definition 4.1.7.** *Let  $\mathcal{C}$  be a category. A sieve of the object  $S$  of  $\mathcal{C}$  is a sieve of the category  $\mathcal{C}_{/S}$ .*

A sieve of  $S$  is therefore a subobject of  $S$  in  $\widehat{\mathcal{C}}$ . It corresponds canonically to a subset of  $\text{Ob } \mathcal{C}_{/S}$  that contains the source of every arrow whose target it contains. By abuse of language, such a subset will also be called a sieve of  $S$ .

## 4.2. Topologies: Definitions

**Definition 4.2.1.** A topology on a category  $\mathcal{C}$  consists, for every object  $S$  of  $\mathcal{C}$ , of a set  $J(S)$  of sieves of  $S$ , called covering sieves or refinements of  $S$ , satisfying the following axioms:

- (T 1) For every refinement  $R$  of  $S$  and every morphism  $T \rightarrow S$ , the sieve  $R \times_S T$  of  $T$  is covering (“stability under base change”).
- (T 2) If  $R, C$  are two sieves of  $S$ , if  $R$  is covering, and if for every  $T \in \text{Ob } \mathcal{C}$  and every morphism  $T \rightarrow R$ , the sieve  $C \times_S T$  of  $T$  is covering, then  $C$  is a refinement of  $S$ .<sup>20</sup>
- (T 3) If  $C \supset R$  are two sieves of  $S$  and  $R$  is covering, then  $C$  is covering.
- (T 4) For every  $S$ ,  $S$  is a refinement of  $S$ .

These axioms may be reformulated as follows. Suppose that a topology  $S \mapsto J(S)$  on  $\mathcal{C}$  is given. For every  $F \in \text{Ob } \widehat{\mathcal{C}}$ , let  $J(F)$  be the set of subfunctors  $R$  of  $F$  such that, for every morphism  $T \rightarrow F$  in  $\widehat{\mathcal{C}}$  with  $T$  representable, the sieve  $R \times_F T$  of  $T$  is covering. By (T 1), this notation agrees with the preceding one. An  $R \in J(F)$  is again called a *refinement* of  $F$ . The preceding axioms immediately imply:

- (T'0) If  $F \supset G$  are objects of  $\widehat{\mathcal{C}}$ , and if  $G \times_F S \in J(S)$  for every  $S \in \text{Ob } \mathcal{C}$  and every morphism  $S \rightarrow F$ , then  $G \in J(F)$ .
- (T'1) If  $G \in J(F)$  and  $H \rightarrow F$  is a morphism in  $\widehat{\mathcal{C}}$ , then  $G \times_F H \in J(H)$ .
- (T'2) If  $F \supset G \supset H$ , if  $G \in J(F)$ , and if  $H \in J(G)$ , then  $H \in J(F)$ .
- (T'3) If  $F \supset G \supset H$  and  $H \in J(F)$ , then  $G \in J(F)$ .
- (T'4) For every  $F \in \text{Ob } \widehat{\mathcal{C}}$ , one has  $F \in J(F)$ .

Conversely, if for every  $F \in \text{Ob } \widehat{\mathcal{C}}$  one is given a set  $J(F)$  of subobjects of  $F$  satisfying T'0–T'4, then  $S \mapsto J(S)$  defines a topology on  $\mathcal{C}$ , and these two constructions are inverse to each other.

Properties T'1, T'2, and T'3 imply the further property<sup>21</sup>:

- (T'5) If  $G$  and  $H$  are two subobjects of  $F$  and  $G, H \in J(F)$ , then  $G \cap H \in J(F)$ .

Thus  $J(F)$ , ordered by  $\supset$ , is filtered. This observation will be useful below.

**4.2.2.** The topology defined by  $J$  is said to be *finer* than the topology defined by  $J'$  if  $J(S) \supset J'(S)$  for every  $S \in \text{Ob } \mathcal{C}$ . Equivalently,  $J(F) \supset J'(F)$  for every  $F \in \text{Ob } \widehat{\mathcal{C}}$ .

Every set of topologies on  $\mathcal{C}$  has a greatest lower bound. More precisely, let  $I$  be an index set and, for each  $i \in I$ , let  $S \mapsto J_i(S)$  be a topology on  $\mathcal{C}$ . Then

$$J(S) = \bigcap_{i \in I} J_i(S)$$

defines a topology, and it is the greatest lower bound of the given set.

In particular, suppose that for every  $S \in \text{Ob } \mathcal{C}$  a set  $E(S)$  of sieves of  $S$  is given. The *topology generated by these sets* is the least fine topology for which every element of  $E(S)$  is a refinement of  $S$ .

<sup>20</sup>Editor's note: In other words, if  $C$  is covering “locally with respect to the covering sieve  $R$ ,” then  $C$  is covering.

<sup>21</sup>Editor's note: T'1 and T'2 suffice:  $G \cap H = G \times_F H$  belongs to  $J(H)$  by T'1, and hence to  $J(F)$  by T'2.

**Definition 4.2.3.** Let  $\{F_i \rightarrow F\}$  be a family of morphisms in  $\widehat{\mathcal{C}}$ , and let  $G \subset F$  be its image (4.1.2). The family is *covering* if  $G \in J(F)$ . A morphism is *covering* if the one-member family consisting of that morphism is covering.

This definition applies in particular to an inclusion: a sieve  $C$  of  $S$  is covering if and only if the canonical morphism  $C \rightarrow S$  is covering.

Axioms  $T'0$ – $T'5$  give the following properties of covering families:

- (C 0) If  $\{F_i \rightarrow F\}$  is a family in  $\widehat{\mathcal{C}}$  such that every representable base change  $\{F_i \times_F S \rightarrow S\}$  is covering, then the original family is covering.
- (C 1) If  $\{F_i \rightarrow F\}$  is covering and  $G \rightarrow F$  is any morphism, then  $\{F_i \times_F G \rightarrow G\}$  is covering (“stability under base change”).
- (C 2) If  $\{F_i \rightarrow F\}$  is covering and, for every  $i$ ,  $\{F_{ij} \rightarrow F_i\}$  is covering, then the composite family  $\{F_{ij} \rightarrow F\}$  is covering (“stability under composition”).
- (C 3) If  $\{G_j \rightarrow F\}$  is covering and  $\{F_i \rightarrow F\}$  is a family such that for every  $j$  there is an  $i$  for which  $G_j \rightarrow F$  factors through  $F_i \rightarrow F$ , then  $\{F_i \rightarrow F\}$  is covering (“saturation”).
- (C 4) Every one-member family consisting of an isomorphism is covering.

Notice that (C 2) and (C 3) also imply:

- (C 5) If  $\{F_i \rightarrow F\}$  is a family for which there exists a covering family  $\{G_j \rightarrow F\}$  such that  $\{F_i \times_F G_j \rightarrow G_j\}$  is covering for every  $j$ , then  $\{F_i \rightarrow F\}$  is covering (“a locally covering family is covering”).

**4.2.4.** Conversely, let  $\mathcal{C}$  be a category having fiber products, and suppose that for every  $S \in \text{Ob } \mathcal{C}$  one is given a set of families of morphisms in  $\mathcal{C}$  with target  $S$ , called covering families, satisfying (C 1)–(C 4), and hence also (C 5). Let  $J(S)$  be the set of sieves of  $S$  having a covering base (or, equivalently by (C 3), all of whose bases are covering). Then  $S \mapsto J(S)$  defines a topology on  $\mathcal{C}$ , and the two preceding constructions are inverse to one another.

In applications it is inconvenient to consider all covering families, since one often has a fairly simple description of a “sufficient” number of them. This leads to the following definitions.

**Definition 4.2.5.0.**<sup>22</sup> Let  $\mathcal{C}$  be a category, and suppose given, for every  $S \in \text{Ob } \mathcal{C}$ , a set  $P(S)$  of families of morphisms in  $\mathcal{C}$  with target  $S$ . The topology generated by  $P$  is the least fine topology for which the given families are covering.

**Definition 4.2.5.** A pretopology on a category  $\mathcal{C}$  consists, for every  $S \in \text{Ob } \mathcal{C}$ , of a set  $R(S)$  of families  $\{S_i \rightarrow S\}$ , called covering for the pretopology, satisfying:

- (P 1) For every  $\{S_i \rightarrow S\} \in R(S)$  and every morphism  $T \rightarrow S$ , the fiber products  $S_i \times_S T$  exist and  $\{S_i \times_S T \rightarrow T\} \in R(T)$ .
- (P 2) If  $\{S_i \rightarrow S\} \in R(S)$  and, for each  $i$ ,  $\{T_{ij} \rightarrow S_i\} \in R(S_i)$ , then the composite family  $\{T_{ij} \rightarrow S\}$  belongs to  $R(S)$ .
- (P 3) Every one-member family consisting of an isomorphism is covering.

<sup>22</sup>Editor’s note: This definition, which appeared after 4.2.5 in the original, has been placed here because it will be used in 6.2.1 in a slightly more general setting than that of 4.2.5.

**Proposition 4.2.6.** *For each  $S$ , let  $J(S)$  be the set of sieves of  $S$  that are covering for the topology generated by the pretopology  $R$ . Let  $J_R(S)$  be the subset of  $J(S)$  consisting of the sieves defined by the families in  $R(S)$ . Then  $J_R(S)$  is cofinal in  $J(S)$ : every refinement of  $S$  contains a sieve defined by a family in  $R(S)$ .*

For every  $S$ , let  $J'(S)$  be the set of sieves of  $S$  containing a sieve in  $J_R(S)$ . Clearly  $J'(S) \subset J(S)$ . To prove  $J(S) = J'(S)$ , it suffices to prove that the  $J'(S)$  define a topology, that is, satisfy (T 1)–(T 4). Axioms (T 1), (T 3), and (T 4) are immediate; only (T 2) remains.<sup>23</sup>

Let  $U \in J'(S)$ , let  $C$  be a sieve of  $S$ , and suppose that  $C \times_S T \in J'(T)$  for every  $T \rightarrow U$ . We must prove  $C \in J'(S)$ . By definition,  $U$  contains a refinement  $U'$  defined by a family  $\{S_i \rightarrow S\} \in R(S)$ . Since (T 3) has already been verified, it is enough to prove  $U' \cap C \in J'(S)$ ; hence we may suppose  $U = U'$ . By hypothesis,  $C \times_S S_i \in J'(S_i)$  for every  $i$ . Thus for each  $i$  there is a covering family  $\{T_{ij} \rightarrow S_i\} \in R(S_i)$  such that  $T_{ij} \rightarrow S_i$  factors through  $C \times_S S_i \rightarrow S_i$ . Consequently  $T_{ij} \rightarrow S$  factors through  $C \rightarrow S$ . Hence  $C$  contains the sieve defined by the composite family  $\{T_{ij} \rightarrow S\}$ , and (P 2) completes the proof.

Axioms (P 1)–(P 3) are those of [MA]. Because pretopologies are so useful in practice, we shall interpret every important result in terms of a pretopology defining the topology under consideration.

**Remark 4.2.7.** One may introduce a slightly more general notion: for every  $S$ , one specifies a set of covering families satisfying (P 1), (P 3), and the conclusion of Proposition 4.2.6. This occurs in particular when the given families satisfy (P 1), (P 3), and (C 5). See [D].

**Definition 4.2.8.** *Let  $\mathcal{C}$  be equipped with a topology, let  $S$  be an object of  $\mathcal{C}$ , and let  $P(S')$  be a condition on an argument  $S' \in \text{Ob } \mathcal{C}/_S$ . Suppose that  $\text{Hom}(S'', S') \neq \emptyset$  implies  $P(S') \Rightarrow P(S'')$ . The condition  $P$  is said to hold locally on  $S$  for the topology under consideration if the following equivalent conditions hold:*

- (i) *The set of  $S' \rightarrow S$  for which  $P(S')$  holds is a refinement of  $S$ .*
- (ii) *There is a refinement of  $S$  such that  $P(S')$  holds for every  $S'$  in that refinement.*
- (iii) *If the topology is defined by a pretopology, there is a covering family for that pretopology such that  $P(S')$  holds for every  $S'$  in the family.*

**Example 4.2.9.** *Let  $f : X \rightarrow Y$  be an  $S$ -morphism. We say that  $f$  is locally an isomorphism if there is a covering family  $\{S_i \rightarrow S\}$  such that  $f \times_S S_i$  is an isomorphism for every  $i$ . Equivalently, there is a refinement  $R$  of  $S$  such that  $X(T) \rightarrow Y(T)$  is an isomorphism for every  $T \rightarrow R$ .*

Many other examples of this “local” language will appear below.

### 4.3. Presheaves, Sheaves, and the Sheaf Associated to a Presheaf

**Definition 4.3.1.** *Let  $\mathcal{C}$  be a category. A presheaf of sets on  $\mathcal{C}$  is a contravariant functor from  $\mathcal{C}$  to the category of sets. The category*

$$\widehat{\mathcal{C}} = \text{Hom}(\mathcal{C}^\circ, \mathbf{Set})$$

*is called the category of presheaves on  $\mathcal{C}$ . If  $\mathcal{C}$  is equipped with a topology, a presheaf  $P$  is separated (respectively, is a sheaf) if, for every  $S \in \text{Ob } \mathcal{C}$  and every  $R \in J(S)$ , the canonical map*

$$P(S) = \text{Hom}(S, P) \longrightarrow \text{Hom}(R, P) \quad (+)$$

<sup>23</sup>Editor’s note: Typographical errors in the original have been corrected in what follows.

is injective (respectively, bijective). The full subcategory of  $\widehat{\mathcal{C}}$  whose objects are the sheaves is called the category of sheaves and is denoted by  $\widetilde{\mathcal{C}}$ .<sup>24</sup>

**Proposition 4.3.2.** *Let  $P$  be a separated presheaf (respectively, a sheaf). For every  $H \in \text{Ob } \mathcal{C}$  and every  $R \in J(H)$ , the canonical map*

$$\text{Hom}(H, P) \longrightarrow \text{Hom}(R, P) \quad (+)$$

*is injective (respectively, bijective).*

Let  $P$  be separated, let  $H$  be a presheaf, let  $R \in J(H)$ , and let  $u, v : H \rightarrow P$  satisfy  $uj = vj$ . For every  $f : S \rightarrow H$ , with  $S \in \text{Ob } \mathcal{C}$ , the sieve  $R \times_H S$  is a refinement of  $S$ , and  $ufj_S = v f j_S$ . This is expressed by the diagram

$$\begin{array}{ccccc} R \times_H S & \xrightarrow{j_S} & S & & \\ f_R \downarrow & & f \downarrow & & \\ R & \xrightarrow{j} & H & \xrightarrow[u]{v} & P. \end{array}$$

Since  $P$  is separated, it follows that  $uf = vf$ . This holds for every representable  $S$ , so  $u = v$ .

Now suppose that  $P$  is a sheaf. Let  $g : R \rightarrow P$ ; we show that it factors through  $H$ . For every  $f : S \rightarrow H$ , with  $S \in \text{Ob } \mathcal{C}$ , the morphism  $g \circ f_R : R \times_H S \rightarrow P$  factors uniquely through  $S$ , thereby defining a morphism  $h : S \rightarrow P$ . By uniqueness,  $h$  is plainly functorial in  $f$ :

$$\begin{array}{ccccc} R \times_H S & \xrightarrow{f_R} & R & \xrightarrow{g} & P \\ j_S \downarrow & & j \downarrow & \nearrow & \\ S & \xrightarrow{f} & H & \xrightarrow{h} & P \end{array}$$

Thus, for every  $S$ , we have defined a map  $H(S) \rightarrow P(S)$  functorial in  $S$ , and hence a morphism  $H \rightarrow P$  with the required properties.

**Corollary 4.3.2.1.**<sup>25</sup> *Let  $R$  and  $F$  be two sheaves. If  $R$  is a refinement of  $F$ , then  $R = F$ .*

Indeed, suppose that  $R$  is a refinement of  $F$ , and write  $j : R \hookrightarrow F$  for the inclusion. By 4.3.2,  $\text{Hom}(F, R) = \text{End}(R)$ ; hence there is a  $\pi : F \rightarrow R$  such that  $\pi j = \text{id}_R$ . Likewise,  $\text{End}(F) = \text{Hom}(R, F)$ , and  $j\pi j = j$  implies  $j\pi = \text{id}_F$ . Thus  $j$  is an isomorphism.

**Proposition 4.3.3 ([AS], 1.3).** *Let  $\mathcal{C}$  be a category and  $P$  a presheaf on  $\mathcal{C}$ . For every  $S \in \text{Ob } \mathcal{C}$ , let  $J(S)$  be the set of sieves  $R$  of  $S$  such that, for every  $T \rightarrow S$ , the map*

$$\text{Hom}(T, P) \longrightarrow \text{Hom}(R \times_S T, P) \quad (+)$$

*is injective (respectively, bijective). Then the  $J(S)$  define a topology on  $\mathcal{C}$ ; that is, they satisfy (T 1)–(T 4).*

**Corollary 4.3.4.** *For every  $S \in \text{Ob } \mathcal{C}$ , let  $K(S)$  be a family of sieves satisfying (T 1), and let  $P$  be a presheaf on  $\mathcal{C}$ . Then  $P$  is separated (respectively, a sheaf) for the topology generated by the  $K(S)$  if and only if, for every  $S \in \text{Ob } \mathcal{C}$  and every  $R \in K(S)$ , the canonical map*

$$\text{Hom}(S, P) \longrightarrow \text{Hom}(R, P) \quad (+)$$

<sup>24</sup>Editor's note: If  $Q$  is a subpresheaf of a separated presheaf  $P$ , then  $Q$  is separated. Indeed, for every sieve  $R$  of  $S$ , the composite  $Q(S) \hookrightarrow P(S) \hookrightarrow P(R)$  is injective and factors through  $Q(S) \rightarrow Q(R)$ .

<sup>25</sup>Editor's note: This corollary has been added.

is injective (respectively, bijective).

**Corollary 4.3.5.** *For each  $S \in \text{Ob } \mathcal{C}$ , let  $R(S)$  be a set of families of morphisms in  $\mathcal{C}$  with target  $S$ , satisfying (P 1), for example a pretopology. Let  $P$  be a presheaf on  $\mathcal{C}$ . Then  $P$  is separated (respectively, a sheaf) for the topology generated by  $R$  if and only if, for every  $S \in \text{Ob } \mathcal{C}$  and every family  $\{S_i \rightarrow S\} \in R(S)$ , the map*

$$P(S) \longrightarrow \prod_i P(S_i)$$

is injective (respectively, the diagram

$$P(S) \longrightarrow \prod_i P(S_i) \rightrightarrows \prod_{i,j} P(S_i \times_S S_j)$$

is exact).

**Definition 4.3.6.** *The canonical topology on a category  $\mathcal{C}$  is the finest topology for which every representable functor is a sheaf.*

**Corollary 4.3.7.** *A sieve  $R$  of  $S$  is a refinement for the canonical topology if and only if, for every morphism  $T \rightarrow S$  in  $\mathcal{C}$  and every  $X \in \text{Ob } \mathcal{C}$ , the canonical map*

$$\text{Hom}(T, X) \longrightarrow \text{Hom}(R \times_S T, X)$$

is bijective.

**Definition 4.3.8.** *A sieve covering for the canonical topology is called a universal effective epimorphic sieve.*

**Corollary 4.3.9.** *A universal effective epimorphic family defines a universal effective epimorphic sieve. Conversely, every squarable family defining a universal effective epimorphic sieve is universal effective epimorphic.*

Return now to a category  $\mathcal{C}$  equipped with an arbitrary topology and to the construction of the sheaf associated to a presheaf  $P$ . Let  $S$  be an object of  $\mathcal{C}$ . If  $R \supset R'$  are two refinements of  $S$ , there is a diagram

$$\begin{array}{ccc} \text{Hom}(S, P) & \longrightarrow & \text{Hom}(R, P) \\ & \searrow & \downarrow \\ & & \text{Hom}(R', P). \end{array}$$

As already observed, the ordered set  $J(S)$  is filtered. Since  $S \in J(S)$ , there is an evident morphism

$$\text{Hom}(S, P) \longrightarrow \varinjlim_{R \in J(S)} \text{Hom}(R, P).$$

**Definition 4.3.10.0.**<sup>26</sup> Set

$$\check{H}^0(S, P) = \varinjlim_{R \in J(S)} \text{Hom}(R, P).$$

---

<sup>26</sup>Editor's note: The numbering 4.3.10.0 has been added for later reference. Moreover, it follows from the definition that if  $Q \rightarrow P$  is a monomorphism, then the same is true of  $LP \rightarrow LQ$ ; thus  $L$  “preserves monomorphisms” (see also 4.3.16 for the more general statement that  $L$  “commutes with finite inverse limits”).

One checks that  $\check{H}^0(S, P)$  depends functorially on  $S$ , and hence defines a functor  $LP$  by

$$\mathrm{Hom}(S, LP) = \check{H}^0(S, P) = \varinjlim_{R \in J(S)} \mathrm{Hom}(R, P). \quad (++)$$

By construction there are morphisms

$$\ell_P : P \longrightarrow LP, \quad z_R : \mathrm{Hom}(R, P) \longrightarrow \mathrm{Hom}(S, LP).$$

**Lemma 4.3.10.**

(i) For every refinement  $R$  of  $S$  and every  $u : R \rightarrow P$ , the diagram

$$\begin{array}{ccc} P & \xrightarrow{\ell_P} & LP \\ u \uparrow & & \uparrow z_R(u) \\ R & \xrightarrow{i_R} & S \end{array}$$

is commutative.

- (ii) For every morphism  $v : S \rightarrow LP$ , there exist a refinement  $R$  of  $S$  and a morphism  $u : R \rightarrow P$  such that  $v = z_R(u)$ .
- (iii) Let  $Q$  be a functor and let  $u, v : Q \rightarrow P$  satisfy  $\ell_P u = \ell_P v$ . Then the kernel of the pair  $(u, v)$  is a refinement of  $Q$ .
- (iv) Let  $u : R \rightarrow P$  and  $u' : R' \rightarrow P$ . The equality  $z_R(u) = z_{R'}(u')$  holds if and only if there is a refinement  $R'' \subset R \cap R'$  of  $S$  on which  $u$  and  $u'$  coincide.

*Proof.* For (i), we must prove  $z_R(u)i_R = \ell_P u$ . It is enough to compare the composites of these two morphisms with every morphism  $g : T \rightarrow R$ , where  $T$  is representable. Put  $f = i_R g$  and  $R' = R \times_S T$ .<sup>27</sup> We have

$$\begin{array}{ccc} P & \xrightarrow{\ell_P} & LP \\ u \uparrow & & \uparrow z_R(u) \\ R & \xrightarrow{i_R} & S \\ p \uparrow & \swarrow g & \uparrow f \\ R' & \xrightarrow[\sim]{i_{R'}} & T. \end{array}$$

By the definition of  $\ell_P$ ,  $\ell_P u g = z_R(u) f$ ; this is the special case of the assertion in which  $i_R$  is an isomorphism. But  $z_R(u) f = z_R(u) i_R g$ , proving (i).

Assertions (ii) and (iv) merely express the definition of  $\mathrm{Hom}(S, LP)$  as a direct limit.

For (iii), let  $K$  be the kernel of  $(u, v)$ . For every morphism  $f : S \rightarrow Q$ , with  $S$  representable,  $K \times_Q S$  is a subfunctor of the kernel of  $uf, vf : S \rightarrow P$ . By  $T'0$ , it is therefore enough to prove the assertion when  $Q = S$  is representable. In that case, (ii) and (iv) show that  $K$  contains a refinement of  $S$ , and hence is itself a refinement of  $S$ .

<sup>27</sup>Editor's note: Let  $p : R' \rightarrow R$  be the projection. Since  $i_R$  is a monomorphism,  $p = gi_{R'}$ . Let  $g'$  be the section of  $i_{R'}$  defined by  $g$ . Then  $pg' = g$ , hence  $pg'i_{R'} = gi_{R'} = p$ . This implies  $g'i_{R'} = \mathrm{id}_{R'}$ , so  $i_{R'} : R' \rightarrow T$  is an isomorphism with inverse  $g'$ .

One finally checks that  $P \mapsto LP$  defines a functor

$$L : \widehat{\mathcal{C}} \longrightarrow \widehat{\mathcal{C}},$$

and that  $P \mapsto \ell_P$  defines a morphism of functors

$$\ell : \text{Id}_{\widehat{\mathcal{C}}} \longrightarrow L.$$

We now state the essential result.

**Proposition 4.3.11.**

- (i) For every presheaf  $P$ , the presheaf  $LP$  is separated and  $\ell_P : P \rightarrow LP$  is covering (4.2.3).
- (ii) If  $P$  is a sheaf, then  $P \rightarrow LP$  is an isomorphism.
- (iii) For every presheaf  $P$  and every separated presheaf (respectively, sheaf)  $F$ , the map

$$\text{Hom}(\ell_P, F) : \text{Hom}(LP, F) \longrightarrow \text{Hom}(P, F)$$

is injective (respectively, bijective).

- (iv) If  $P$  is separated, then  $\ell_P : P \rightarrow LP$  is a covering monomorphism (so  $P$  is a refinement of  $LP$ ), and  $LP$  is a sheaf.

*Proof.* (i) First,  $P \rightarrow LP$  is covering. Indeed, for every  $v : S \rightarrow LP$ , Lemma 4.3.10(i),(ii) supplies a refinement  $R$  of  $S$  and a commutative diagram (the map from  $R$  to the fiber product is induced by  $v' : R \rightarrow P$  and  $i_R : R \rightarrow S$ )

$$\begin{array}{ccccc} & & P & \xrightarrow{\ell_P} & LP \\ & \nearrow v' & \uparrow & & \uparrow v \\ & & P \times_{LP} S & \xrightarrow{\quad} & S \\ & \nearrow & \uparrow & & \uparrow \\ R & \xrightarrow{i_R} & & & \end{array}$$

so  $P \times_{LP} S \rightarrow S$  is covering. By (C 0),  $P \rightarrow LP$  is covering.

Now suppose that two morphisms  $v_1, v_2 : S \rightarrow LP$  induce the same morphism on a refinement  $R$  of  $S$ . There are refinements  $R_i$ ,  $i = 1, 2$ , and morphisms  $u_i : R_i \rightarrow P$  with  $z_{R_i}(u_i) = v_i$ . Replacing  $R$  by a smaller refinement, we may suppose  $R_1 = R_2 = R$ . The commutative diagram of Lemma 4.3.10(i) then gives  $\ell_P u_1 = \ell_P u_2$ . By 4.3.10(iii),  $u_1$  and  $u_2$  coincide on a refinement of  $R$ , hence on a refinement of  $S$ ; by 4.3.10(iv),  $z_R(u_1) = z_R(u_2)$ . Thus  $LP$  is separated.

(ii) This is clear: if  $P$  is a sheaf, then  $\text{Hom}(S, P) \rightarrow \text{Hom}(R, P)$  is already an isomorphism for every refinement  $R$  of  $S$ .

(iii) Let  $u, v : LP \rightarrow F$  satisfy  $u\ell_P = v\ell_P$ . To show  $u = v$ , it is enough to prove  $uf = vf$  for every  $f : S \rightarrow LP$  with  $S$  representable. There are a refinement  $R$  of  $S$  and a morphism  $g : R \rightarrow P$  with  $f = z_R(g)$ . The morphisms  $uf$  and  $vf$  agree on  $R$ , where both equal  $u\ell_P g = v\ell_P g$ . If  $F$  is separated, they therefore agree on  $S$ , proving injectivity. If  $F$  is a sheaf, the commutative diagram

$$\begin{array}{ccc} P & \xrightarrow{\ell_P} & LP \\ \downarrow & & \downarrow \\ F & \xrightarrow{\sim} & LF \end{array}$$



shows that  $\text{Hom}(\ell_P, F)$  is also surjective.

(iv) Suppose that  $P$  is separated. To show that  $P \rightarrow LP$  is a monomorphism, it is enough to consider  $u, v : S \rightarrow P$ , with  $S$  representable, satisfying  $\ell_P u = \ell_P v$ . By 4.3.10(iii),  $u$  and  $v$  agree on a refinement of  $S$ , and therefore agree because  $P$  is separated. Thus  $P \rightarrow LP$  is a monomorphism; since it is covering by (i),  $P$  is a refinement of  $LP$ .

It remains to show that  $LP$  is a sheaf. We already know from (i) that it is separated. Let  $S \in \text{Ob } \mathcal{C}$ , let  $R$  be a refinement of  $S$ , and let  $h : R \rightarrow LP$ . We must find  $u : S \rightarrow LP$  with  $ui_R = h$ . Since  $P$  is a refinement of  $LP$ ,  $R' = P \times_{LP} R$  is a refinement of  $R$ , and hence of  $S$ . Put  $u = z_{R'}(h')$ ; the situation is

$$\begin{array}{ccccc} P & \xrightarrow{\ell_P} & LP & & \\ h' \uparrow & & \uparrow h & \swarrow u & \\ R' & \xrightarrow{j} & R & \xrightarrow{i_R} & S. \end{array}$$

Here the diagonal arrow is  $u : S \rightarrow LP$ . We have  $ui_{R'} = \ell_P h' = hj$ , hence  $ui_R j = hj$ . Since  $R'$  is a refinement of  $R$  and  $LP$  is separated, Proposition 4.3.2 gives  $ui_R = h$ .

**Corollary 4.3.12.**<sup>28</sup> *Let  $F$  be a sheaf and let  $R$  be a subobject of  $F$  in  $\widehat{\mathcal{C}}$ . Then  $R$  is a separated presheaf,  $\ell_R : R \rightarrow LR$  is a covering monomorphism, and there is a commutative diagram*

$$\begin{array}{ccc} R & \xrightarrow{i} & F \\ \ell_R \searrow & & \uparrow j \\ & & LR. \end{array}$$

Consequently,  $R$  is a refinement of  $F$  if and only if  $j$  is an isomorphism.

We already observed that  $R$  is separated and that  $j$  is a monomorphism; see editor notes (24) and (26). By Proposition 4.3.11(iv),  $\ell_R$  is a covering monomorphism. Thus, if  $j$  is an isomorphism,  $R$  is a refinement of  $F$ . Conversely, if  $i$  is covering, then so is  $j$ , and Proposition 4.3.2.1 shows that  $j$  is an isomorphism.

**Remark 4.3.13.** If  $J'(S)$  is a cofinal subset of  $J(S)$ , then

$$\text{Hom}(S, LP) = \varinjlim_{R \in J'(S)} \text{Hom}(R, P).$$

In particular, let  $S \mapsto R(S)$  be a pretopology generating the given topology. The functor  $L$  can be described using the covering families belonging to  $R(S)$ . Writing out the preceding formula recovers the construction of [MA].

Let  $i : \widehat{\mathcal{C}} \rightarrow \widetilde{\mathcal{C}}$  be the inclusion functor. Proposition 4.3.11 gives the following theorem.

**Theorem 4.3.14.** *There is a unique functor  $a : \widehat{\mathcal{C}} \rightarrow \widetilde{\mathcal{C}}$  for which the diagram*

$$\begin{array}{ccc} \widehat{\mathcal{C}} & \xrightarrow{L} & \widehat{\mathcal{C}} \\ a \downarrow & & \downarrow L \\ \widetilde{\mathcal{C}} & \xrightarrow{i} & \widehat{\mathcal{C}} \end{array}$$

is commutative; that is,  $L(L(P))$  is a sheaf for every presheaf  $P$ . The functors  $i$  and  $a$  are adjoint: for every presheaf  $P$  and every sheaf  $F$ , there is an isomorphism, functorial in  $P$  and  $F$ ,

$$\text{Hom}_{\widetilde{\mathcal{C}}}(P, i(F)) \simeq \text{Hom}_{\widehat{\mathcal{C}}}(a(P), F),$$

<sup>28</sup>Editor's note: The statement and proof of this corollary have been expanded.

or simply

$$\mathrm{Hom}(P, F) \simeq \mathrm{Hom}(a(P), F).$$

**Definition 4.3.15.** The sheaf  $a(P)$  is called the sheaf associated to the presheaf  $P$ .

**Remark 4.3.16.** Since  $L$  is constructed using inverse limits and filtered direct limits, it commutes with finite inverse limits.<sup>29</sup>

Moreover, under the identification  $L(P \times P) = LP \times LP$ , the morphism  $\ell_{P \times P}$  is identified with  $\ell_P \times \ell_P$ . Thus, for example, if  $P$  is a presheaf of groups, then  $LP$  is canonically a presheaf of groups and the canonical morphism  $P \rightarrow LP$  is a morphism of groups. The same holds for the functor  $a$ . Consequently, if  $P$  is a presheaf of groups and  $F$  a sheaf of groups, there is an isomorphism

$$\mathrm{Hom}_{\widehat{\mathcal{C}\text{-gr.}}}(P, i(F)) \simeq \mathrm{Hom}_{\widehat{\mathcal{C}\text{-gr.}}}(a(P), F).$$

See [D] for further details.

**4.3.17.** Let  $\mathcal{V}$  be any category. A *presheaf on  $\mathcal{C}$  with values in  $\mathcal{V}$*  is a contravariant functor from  $\mathcal{C}$  to  $\mathcal{V}$ . Before defining sheaves with values in  $\mathcal{V}$ , we recall the definition of the inverse limit of a functor. Let  $\mathcal{R}$  and  $\mathcal{V}$  be categories, and let

$$F : \mathcal{R}^\circ \longrightarrow \mathcal{V}$$

be a contravariant functor. Write  $\varprojlim F$  for the object of  $\widehat{\mathcal{V}}$  defined by

$$\begin{aligned} (\varprojlim F)(X) &= \mathrm{Hom}_{\widehat{\mathcal{V}}}(X, \varprojlim F) \\ &= \varprojlim_{U \in \mathrm{Ob} \mathcal{R}} \mathrm{Hom}_{\mathcal{V}}(F(U), X) \\ &= \mathrm{Hom}(c_X, F), \end{aligned}$$

where  $X$  varies over the objects of  $\mathcal{V}$ , where  $c_X$  denotes the contravariant functor from  $\mathcal{R}$  to  $\mathcal{V}$  that sends every object to  $X$  and every arrow to  $\mathrm{id}_X$ , and where the last  $\mathrm{Hom}$  is taken in  $\mathrm{Hom}(\mathcal{R}^\circ, \mathcal{V})$ . If  $\mathcal{R}$  has a final object  $e_{\mathcal{R}}$ , then  $\varprojlim F = F(e_{\mathcal{R}})$ . If  $\mathcal{V} = \mathbf{Set}$ , the functor  $\varprojlim$  is identified with  $\Gamma$ .

Let  $S$  be an object of  $\mathcal{C}$  and  $R$  a sieve of  $S$ . Let  $\overline{R}$  be the full subcategory of  $\mathcal{C}_{/S}$  whose object set is  $E(R)$ , and let  $i_R : \overline{R} \rightarrow \mathcal{C}_{/S}$  be the canonical functor. A presheaf  $P$  on  $\mathcal{C}$  with values in  $\mathcal{V}$  restricts to a functor

$$P_S : (\mathcal{C}_{/S})^\circ \longrightarrow \mathcal{V}.$$

The functor  $i_R$  induces a morphism in  $\widehat{\mathcal{V}}$ :

$$P(S) = P_S(S) = \varprojlim P_S \longrightarrow \varprojlim (P_S \circ i_R).$$

We write

$$\varprojlim_R P = \varprojlim (P_S \circ i_R)$$

for the latter object of  $\widehat{\mathcal{V}}$ . In view of 4.1.6, Definition 4.3.1 generalizes as follows.

<sup>29</sup>Editor's note: In particular, if  $K$  is the kernel of a pair of morphisms of presheaves  $u, v : Q \rightrightarrows P$ , then  $LK$  is the kernel of  $Lu, Lv : LQ \rightrightarrows LP$ . This will be used in 4.4.5.

**Definition 4.3.18.** A presheaf  $P$  on  $\mathcal{C}$  with values in  $\mathcal{V}$  is separated (respectively, a sheaf) if, for every  $S \in \text{Ob } \mathcal{C}$  and every  $R \in J(S)$ , the canonical morphism in  $\mathcal{V}$

$$P(S) \longrightarrow \varprojlim_R P$$

is a monomorphism (respectively, an isomorphism).

When  $\mathcal{V}$  is the category **Gr** of groups (or any other category of sets endowed with algebraic structures defined by finite inverse limits), the following notions are equivalent (cf. [D]): a presheaf on  $\mathcal{C}$  with values in **Gr** whose underlying presheaf of sets is a sheaf, and a group object in the category of sheaves of sets. With these identifications understood, we shall always regard sheaves with values in a category of sets endowed with algebraic structures defined by finite inverse limits as sheaves of sets equipped, in  $\tilde{\mathcal{C}}$ , with the corresponding algebraic structure.

#### 4.4. Exactness Properties of the Category of Sheaves

##### Theorem 4.4.1.

- (i) Arbitrary inverse limits exist in  $\tilde{\mathcal{C}}$ , and “are computed in  $\hat{\mathcal{C}}$ ”; that is, the inclusion functor

$$i : \tilde{\mathcal{C}} \longrightarrow \hat{\mathcal{C}}$$

commutes with inverse limits. More explicitly, if  $(X_\alpha)$  is an inverse system of sheaves, the presheaf

$$\varprojlim_\alpha i(X_\alpha) : S \longmapsto \varprojlim_\alpha X_\alpha(S)$$

is a sheaf, and

$$i\left(\varprojlim_\alpha X_\alpha\right) = \varprojlim_\alpha i(X_\alpha).$$

- (ii) Arbitrary direct limits exist in  $\tilde{\mathcal{C}}$ . If  $(X_\alpha)$  is a direct system of sheaves, then

$$\varinjlim_\alpha X_\alpha = a\left(\varinjlim_\alpha i(X_\alpha)\right),$$

where  $\varinjlim_\alpha i(X_\alpha)$  is the presheaf direct limit of the  $i(X_\alpha)$ :

$$\varinjlim_\alpha i(X_\alpha) : S \longmapsto \varinjlim_\alpha X_\alpha(S).$$

- (iii) The functor

$$a : \hat{\mathcal{C}} \longrightarrow \tilde{\mathcal{C}}$$

commutes with arbitrary direct limits and finite inverse limits.

Assertions (i) and (ii) follow formally from the adjunction formula (4.3.14), and<sup>30</sup> the first assertion of (iii) follows from (ii). The second assertion of (iii) was already noted in 4.3.16.

**Scholium 4.4.2.** This theorem makes it possible to use the following method for proving in  $\tilde{\mathcal{C}}$  a statement involving both arbitrary direct limits and finite inverse limits (for example, “every epimorphism is universally effective”; cf. below). One begins by proving the

<sup>30</sup>Editor’s note: the original has been modified here.

corresponding statement in the category of sets, then extends it pointwise to the category of presheaves. One then uses the preceding theorem to pass from the category of presheaves to the category of sheaves. Many examples of this method will occur below (4.4.3, 4.4.6, 4.4.9, and so forth).

Finally, observe that statements concerning the category of presheaves are formally corollaries of the corresponding statements concerning the category of sheaves. It is enough to take the coarsest (“chaotic”) topology, namely the topology defined by

$$J(S) = \{S\} \quad \text{for every } S \in \text{Ob } \mathcal{C};$$

every functor is a sheaf for this topology.

**Proposition 4.4.3.** *Let  $\mathcal{F} = \{F_i \rightarrow F\}$  be a family of morphisms of sheaves. The following conditions are equivalent:*

- (i)  $\mathcal{F}$  is an epimorphic family;
- (ii)  $\mathcal{F}$  is a universal effective epimorphic family (1.13);
- (iii)  $\mathcal{F}$  is covering (4.2.3);
- (iv) the sheaf image of  $\mathcal{F}$ , that is, the sheaf associated with the presheaf image of  $\mathcal{F}$  (4.1.2), is  $F$ .

The equivalence of (iii) and (iv) follows from 4.3.12. The remaining equivalences will follow from the next lemmas.

**Lemma 4.4.4.** *Let  $f : X \rightarrow Y$  be a monomorphism of sheaves that is also an epimorphism. Then  $f$  is an isomorphism.*

The lemma is clear first of all in the category of sets. Let us next prove it in the category of presheaves. Consider the presheaf

$$V : S \mapsto Y(S) \amalg_{X(S)} Y(S);$$

this is the amalgamated sum of  $Y$  and  $Y$  over  $X$  in the category of presheaves.<sup>31</sup> Let  $i_1, i_2 : Y \rightrightarrows V$  be the two coordinate morphisms. If  $X \rightarrow Y$  is an epimorphism in the category of presheaves, then  $i_1 = i_2$ . In that case, for every  $S$ , the map  $X(S) \rightarrow Y(S)$  is surjective; since it is also injective, it is bijective, and hence  $X \rightarrow Y$  is an isomorphism.

Finally, work in the category  $\mathcal{C}$  of sheaves. By 4.4.1(ii), the amalgamated sum in  $\mathcal{C}$  of the sheaves  $Y$  and  $Y$  over  $X$  is the sheaf  $Z$  associated with the presheaf  $V$ . Consider the diagram

$$X \xrightarrow{f} Y \xrightleftharpoons[i_2]{i_1} V \xrightarrow{\tau} Z = a(V).$$

We have  $i_1 f = i_2 f$ , hence  $\tau i_1 f = \tau i_2 f$ , and therefore  $\tau i_1 = \tau i_2$ , since  $f$  is an epimorphism in  $\mathcal{C}$ . By part (iii) of the following lemma, the presheaf  $V$  is separated, so  $\tau$  is a monomorphism (4.3.11(iv)). Thus  $i_1 = i_2$ , and we saw above that this implies that  $f : X \rightarrow Y$  is an isomorphism. This proves Lemma 4.4.4, once it has been verified that  $V$  is separated.

Let  $R_\emptyset$  denote the presheaf that assigns the empty set to every  $S \in \text{Ob } \mathcal{C}$ ; it is an initial object of  $\mathcal{C}$ .<sup>32</sup>

**Lemma 4.4.5.**

<sup>31</sup>Editor’s note: the remainder of the proof has been slightly modified.

<sup>32</sup>Editor’s note: part (i) of Lemma 4.4.5 has been corrected, and the proofs of all three parts have been given in detail.

- (i) Suppose that  $R_\emptyset \notin J(S)$  for every  $S \in \text{Ob } \mathcal{C}$ . If  $(X_i)$  is a family of separated presheaves, then the coproduct presheaf  $\coprod_i X_i$  is separated.<sup>33</sup>
- (ii) Consider an equivalence relation in the category of presheaves

$$X \begin{smallmatrix} \xrightarrow{u} \\ \xrightarrow{v} \end{smallmatrix} Y,$$

and let  $w : Y \rightarrow Z$  be the quotient. If  $X$  is a sheaf and  $Y$  is separated, then  $Z$  is separated.

- (iii) Consider an amalgamated sum in the category of presheaves, in which  $u$  and  $u'$  are monomorphisms:

$$\begin{array}{ccc} X & \xrightarrow{u} & Y \\ u' \downarrow & & \downarrow \\ Y' & \longrightarrow & V. \end{array}$$

If  $Y$  and  $Y'$  are separated and  $X$  is a sheaf, then  $V$  is separated.

*Proof.* For (i), set  $X = \coprod_i X_i$ . Let  $S \in \text{Ob } \mathcal{C}$ , let  $\tau : R \hookrightarrow S$  be a refinement of  $S$ , and let  $x_1, x_2 \in X(S)$  satisfy  $x_1 \circ \tau = x_2 \circ \tau$ . There are indices  $i, j$  such that  $x_1 \in X_i(S)$  and  $x_2 \in X_j(S)$ . Since  $R \neq R_\emptyset$ , there is a morphism  $\phi : T \rightarrow R$  with  $T \in \text{Ob } \mathcal{C}$ . Then

$$x_1 \circ \tau \circ \phi = x_2 \circ \tau \circ \phi.$$

Since  $X(T)$  is the disjoint union of the  $X_k(T)$ , it follows that  $i = j$ . Because  $X_i$  is separated and is a subobject of  $X$ , the map

$$X_i(S) \longrightarrow X_i(R) \longrightarrow X(R)$$

is injective. Hence  $x_1 = x_2$ , which proves that  $X$  is separated.

We next prove (iii). Let  $i : Y \rightarrow V$  and  $j : Y' \rightarrow V$  be the canonical morphisms, and let  $K$  be the kernel of the pair

$$Y \times Y' \begin{smallmatrix} \xrightarrow{p} \\ \xrightarrow{q} \end{smallmatrix} V, \quad p = i \circ \text{pr}_1, \quad q = j \circ \text{pr}_2.$$

For  $S \in \text{Ob } \mathcal{C}$ , the monomorphisms  $u, u'$  allow us to identify  $X(S)$  with its image in  $Y(S)$ , respectively in  $Y'(S)$ . Write  $Z(S)$ , respectively  $Z'(S)$ , for the complement. Then  $V(S)$  identifies with the disjoint union of  $Z(S)$ ,  $Z'(S)$ , and  $X(S)$ . It follows at once that

$$i(S) : Y(S) \longrightarrow V(S), \quad j(S) : Y'(S) \longrightarrow V(S)$$

are injective and that

$$(u \times u')(S) : X(S) \longrightarrow K(S)$$

is bijective. Consequently  $i$  and  $j$  are monomorphisms, and  $u \times u' : X \rightarrow K$  is an isomorphism.

<sup>33</sup>Editor's note: in general, the coproduct of two sheaves  $F, G$  is not a sheaf. Indeed, let  $S_1, S_2 \in \text{Ob } \mathcal{C}$ . Suppose that the coproduct  $S = S_1 \coprod S_2$  exists in  $\mathcal{C}$ , and that the fiber product  $S_1 \times_S S_2$  is an initial object  $\emptyset$  of  $\mathcal{C}$  (cf. I, 1.8). Let  $R$  be the sieve of  $S$  with basis  $\{S_1, S_2\}$ . Then  $(F \coprod G)(R)$  is the disjoint union of  $F(S) \coprod G(S)$  and the sets  $F(S_i) \times G(S_j)$ , for  $i \neq j$ ; thus  $F \coprod G$  is not a sheaf in general. On the other hand, let  $\mathcal{C}$  be the category with a single object  $S$  and  $\text{id}_S$  as its only morphism, equipped with the topology defined by  $J(S) = \{R_\emptyset, S\}$ . Then the only separated presheaves are  $R_\emptyset$  and  $X = \mathbf{h}_S$  (which is a sheaf), and  $X \coprod X$  is not separated.

Set  $U = Y \amalg Y'$ . For a refinement  $\tau : R \hookrightarrow S$ , there is a commutative diagram

$$\begin{array}{ccc} U(S) & \xrightarrow{U(\tau)} & U(R) \\ \downarrow & & \downarrow \\ V(S) & \xrightarrow{V(\tau)} & V(R). \end{array}$$

Let  $v_1, v_2 \in V(S)$  have the same image in  $V(R)$ . By the definition of  $V$ , they lift to elements  $y_1, y_2 \in U(S)$ ; let  $z_1, z_2$  denote their images in  $U(R)$ . Then  $z_1, z_2$  have the same image in  $V(R)$ .

Since  $i : Y \rightarrow V$  and  $j : Y' \rightarrow V$  are monomorphisms, the maps  $Y(R) \rightarrow V(R)$  and  $Y'(R) \rightarrow V(R)$  are injective. Thus, because  $Y$  and  $Y'$  are separated, if  $y_1, y_2$  both belong to  $Y(S)$ , or both belong to  $Y'(S)$ , then  $y_1 = y_2$ . Otherwise we may suppose that  $y_1 \in Y(S)$  and  $y_2 \in Y'(S)$ , so that  $z_1 \in Y(R)$  and  $z_2 \in Y'(R)$ . Since  $i \circ z_1 = j \circ z_2$ , the morphism

$$z_1 \boxtimes z_2 : R \longrightarrow Y \times Y'$$

factors through  $K = X$ . Moreover, since  $X$  is a sheaf, the restriction map identifies  $X(S)$  with  $X(R)$ . Hence there is an  $x \in X(S)$  such that

$$u(x) \circ \tau = z_1 = y_1 \circ \tau, \quad u'(x) \circ \tau = z_2 = y_2 \circ \tau.$$

Since  $Y$  and  $Y'$  are separated,  $u(x) = y_1$  and  $u'(x) = y_2$ , and therefore  $v_1 = v_2$ . This proves that  $V$  is separated.

Finally, we prove (ii). First observe that the morphism of presheaves

$$X \xrightarrow{u \boxtimes v} K,$$

where  $K$  is the kernel of the pair

$$w \circ \text{pr}_i : Y \times Y \rightrightarrows Z,$$

is an isomorphism. Indeed, for every  $T \in \text{Ob } \mathcal{C}$ ,  $X(T)$  is an equivalence relation on  $Y(T)$ , and hence the diagram

$$X(T) \xrightarrow{u \boxtimes v} Y(T) \times Y(T) \begin{array}{c} \xrightarrow{w \circ \text{pr}_1} \\ \xleftarrow{w \circ \text{pr}_2} \end{array} Z(T)$$

is exact in the category of sets.

Now let  $S \in \text{Ob } \mathcal{C}$ , and let  $g_1, g_2 : S \rightarrow Z$  be two morphisms that agree on a refinement  $\tau : R \hookrightarrow S$ . By the construction of  $Z$ ,

$$Z(S) = Y(S)/X(S),$$

so there are morphisms  $f_1, f_2 : S \rightarrow Y$  such that  $wf_i = g_i$ . We have  $wf_1\tau = wf_2\tau$ ; by the preceding observation, there is a morphism  $\phi : R \rightarrow X$  such that

$$u\phi = f_1\tau, \quad v\phi = f_2\tau.$$

Since  $X$  is a sheaf, there is a morphism  $\psi : S \rightarrow X$  with  $\phi = \psi\tau$ . Thus, in  $Y(R)$ ,

$$u\psi\tau = f_1\tau, \quad v\psi\tau = f_2\tau.$$

Because  $Y(S) \rightarrow Y(R)$  is injective,  $Y$  being separated, it follows that  $u\psi = f_1$  and  $v\psi = f_2$ , hence  $g_1 = g_2$ . Thus  $Z$  is separated.

**Lemma 4.4.6.**<sup>34</sup>

<sup>34</sup>Editor's note: the statement of the lemma and its proof have been given in detail.

- (i) Let  $\{F_i \xrightarrow{f_i} F\}$  be a family of morphisms of presheaves, and let  $G \subset F$  be its presheaf image. Then the following diagram in  $\widehat{\mathcal{C}}$  is exact:

$$\left(\coprod_i F_i\right) \times_G \left(\coprod_j F_j\right) \xrightleftharpoons[\text{pr}_2]{\text{pr}_1} \coprod_i F_i \longrightarrow G.$$

Equivalently, for every presheaf  $H$ , the following diagram of sets is exact:

$$\text{Hom}(G, H) \longrightarrow \prod_i \text{Hom}(F_i, H) \xrightleftharpoons[q]{p} \prod_{i,j} \text{Hom}(F_i \times_G F_j, H). \quad (*)$$

- (ii) Every covering family of morphisms of sheaves is a universal effective epimorphic family.

*Proof.* For (i), let  $H$  be a presheaf. The map

$$f^* : \text{Hom}(G, H) \longrightarrow \prod_i \text{Hom}(F_i, H), \quad \phi \longmapsto (\phi \circ f_i)_i,$$

is injective. Indeed, for each  $S \in \text{Ob } \mathcal{C}$ , the map  $\phi(S)$  is determined by the maps  $(\phi \circ f_i)(S)$ , since the family

$$f_i(S) : F_i(S) \longrightarrow G(S)$$

is surjective. It is clear that the image of  $f^*$  is contained in  $\ker(p, q)$ .

Conversely, let  $\phi_i : F_i \rightarrow H$  be a family of morphisms such that, for every  $i, j$ , the diagram

$$\begin{array}{ccc} F_i \times_G F_j & \longrightarrow & F_j \\ \downarrow & & \downarrow \phi_j \\ F_i & \xrightarrow{\phi_i} & H \end{array}$$

commutes. For every  $S$ , the map

$$\prod_i F_i(S) \longrightarrow H(S)$$

then factors uniquely through a map  $\phi(S) : G(S) \rightarrow H(S)$ . These maps define a morphism  $\phi : G \rightarrow H$  satisfying  $f^*(\phi) = (\phi_i)$ . This proves the exactness of  $(*)$ , and hence (i).

We prove (ii). Since covering families are stable under base change, it is enough to show that every covering family is effective epimorphic. Let  $\{F_i \rightarrow F\}$  be a covering family of morphisms of sheaves, and let  $G$  be its presheaf image. Because the family is covering, the monomorphism  $G \hookrightarrow F$  is covering as well; therefore, by 4.3.12,

$$a(G) = F.$$

Moreover, since  $G \hookrightarrow F$  is a monomorphism, the fiber products  $F_i \times_G F_j$  and  $F_i \times_F F_j$  coincide. Part (i) therefore shows that, for every presheaf  $H$ , the diagram

$$\text{Hom}(G, H) \longrightarrow \prod_i \text{Hom}(F_i, H) \xrightleftharpoons[q]{p} \prod_{i,j} \text{Hom}(F_i \times_F F_j, H) \quad (**)$$

is exact. If, in addition,  $H$  is a sheaf, then

$$\text{Hom}(G, H) = \text{Hom}(a(G), H) = \text{Hom}(F, H).$$

Thus (\*\*) shows that  $\{F_i \rightarrow F\}$  is an effective epimorphic family in the category  $\tilde{\mathcal{C}}$  of sheaves.

**Lemma 4.4.7.** *Every family of morphisms of sheaves  $\{F_i \rightarrow F\}$  factors as a covering family  $\{F_i \rightarrow G\}$  followed by a monomorphism  $G \rightarrow F$ .*

Indeed, it is enough to take  $G$  to be the sheaf image of the given family.

*Proof of Proposition 4.4.3.* We saw in 4.4.6 that (iii) implies (ii), while (ii) evidently implies (i). Finally, let  $\{F_i \rightarrow F\}$  be an epimorphic family. By Lemma 4.4.7 it factors as a covering family followed by a monomorphism. The latter monomorphism, being dominated<sup>35</sup> by an epimorphic family, is itself an epimorphism; hence it is an isomorphism by 4.4.4.

**Remark 4.4.8.**<sup>36</sup> Because the presheaf image of the family  $\mathcal{F} = \{F_i \rightarrow F\}$  is separated, the construction of the associated sheaf shows that the conditions of Proposition 4.4.3 are also equivalent to the following:

- (v) for every  $S \in \text{Ob } \mathcal{C}$ , each  $f \in F(S)$  lies locally in the image of  $\mathcal{F}$ ; that is,
- (vi) for every  $S \in \text{Ob } \mathcal{C}$  and every  $f \in F(S)$ , the set of morphisms  $S' \rightarrow S$  such that the image of  $f$  in  $F(S')$  belongs to the image of one of the  $F_i(S')$  is a refinement of  $S$ ;
- (vii) if the topology is defined by a pretopology  $S \mapsto \mathcal{R}(S)$ , then, for every  $S \in \text{Ob } \mathcal{C}$  and every  $f \in F(S)$ , there is a family  $\{S_j \rightarrow S\} \in \mathcal{R}(S)$  such that, for each  $j$ , the image  $f_j$  of  $f$  in  $F(S_j)$  belongs to the image of one of the  $F_i(S_j)$ .

**Remark 4.4.8 bis.** If the sheaf  $F$  is representable, the preceding conditions are also equivalent to:

- (viii) the set of morphisms  $T \rightarrow F$ , with  $T \in \text{Ob } \mathcal{C}$ , for which there exist an  $i$  and a commutative diagram

$$\begin{array}{ccc} & F_i & \\ & \downarrow & \\ T & \longrightarrow & F \end{array}$$

is a refinement of  $F$ .

Indeed, if (viii) holds, the presheaf image of the  $F_i$  is dominated by<sup>37</sup> a refinement of  $F$ , which implies that the family is covering. Conversely, apply (vi) to  $\text{id}_F \in F(F)$ .

In pictorial language, this condition says: *locally on  $F$ , there is an  $i$  for which  $F_i \rightarrow F$  has a section*. In particular, a morphism

$$G \longrightarrow F,$$

where  $G$  is a sheaf and  $F$  is a representable sheaf, is covering if and only if it has a section locally on  $F$ .

The following lemmas will be useful in VI<sub>B</sub>, 8.1 and 8.2.<sup>38</sup>

**Lemma 4.4.8.1.** *Let  $Q \subset P$  be presheaves of groups on  $\mathcal{C}$ , with  $Q$  normal in  $P$ . Suppose that  $P$  is separated and that  $Q$  is a sheaf. Then the presheaf of groups  $P/Q$  is separated. Consequently,*

$$a(P/Q) = L(P/Q) = \varinjlim_{R \in J(S)} (P/Q)(R).$$

<sup>35</sup>Editor's note: recall (cf. editor's note 17) that a morphism  $g : G \rightarrow H$  is said to be *dominated* by a family of morphisms  $F_i \rightarrow H$  if that family factors through  $g$ .

<sup>36</sup>Editor's note: the numbering of the original, which had two paragraphs numbered 4.4.7, has been corrected to agree with later references.

<sup>37</sup>Editor's note: "contains" has been replaced by "is dominated by"; cf. editor's note 17.

<sup>38</sup>Editor's note: Lemmas 4.4.8.1 and 4.4.8.2 have been added.



It is enough to prove the first assertion, since the second follows from it by 4.3.11(iv) and (ii). Let  $S \in \text{Ob } \mathcal{C}$ , let  $R$  be a sieve of  $S$ , and let  $y \in P(S)/Q(S)$  have identity image in  $(P/Q)(R)$ . We must show that  $y = 1$ .

By hypothesis, the morphism  $P(S) \rightarrow P(R)$  is injective, while  $Q(S) \rightarrow Q(R)$  is an isomorphism, and the upper row in the following commutative diagram is exact:

$$\begin{array}{ccccccccc} 1 & \longrightarrow & Q(S) & \longrightarrow & P(S) & \longrightarrow & P(S)/Q(S) & \longrightarrow & 1 \\ & & \downarrow \sim & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & Q(R) & \longrightarrow & P(R) & \longrightarrow & (P/Q)(R) & & \end{array}$$

The result follows if the lower row is exact. Let  $f : R \rightarrow P$  have identity image in  $(P/Q)(R)$ , and let  $\phi : T \rightarrow R$  with  $T \in \text{Ob } \mathcal{C}$ . Then  $f \circ \phi$  is the identity in

$$(P/Q)(T) = P(T)/Q(T),$$

that is,  $f \circ \phi \in Q(T)$ . Hence  $\phi \mapsto f \circ \phi$  is a functorial map  $R(T) \rightarrow Q(T)$ , and so defines a morphism of functors  $\pi : R \rightarrow Q$  such that  $\pi(\text{id}_R) = f$ . Thus  $f \in Q(R)$ . This proves the exactness of the lower row and establishes the lemma.

**Lemma 4.4.8.2.** *Let  $H \subset G$  be presheaves of groups on  $\mathcal{C}$ .*

- (i) *If  $H$  is normal in  $G$ , then  $L(H)$  is normal in  $L(G)$ .*
- (ii) *If  $H$  is central in  $G$ , then  $L(H)$  is central in  $L(G)$ .*

Let  $S \in \text{Ob } \mathcal{C}$ . We must show (cf. I 2.3.6) that  $L(H)(S)$  is normal, respectively central, in  $L(G)(S)$ . Let

$$h \in L(H)(S), \quad g \in L(G)(S).$$

There are a sieve  $R$  of  $S$  and elements  $h' \in H(R)$ ,  $g' \in G(R)$  such that

$$h = z_R(h'), \quad g = z_R(g')$$

in the notation of 4.3.10. Since  $z_R$  is a group morphism,

$$ghg^{-1}h^{-1} = z_R(g'h'g'^{-1}h'^{-1}).$$

In case (i),

$$g'h'g'^{-1}h'^{-1} \in H(R),$$

and therefore  $ghg^{-1}h^{-1} \in L(H)(S)$ . In case (ii),  $g'h'g'^{-1}h'^{-1} = 1$ , and hence  $ghg^{-1}h^{-1} = 1$ .

**Proposition 4.4.9.** *Every equivalence relation in  $\tilde{\mathcal{C}}$  is universally effective (3.3.3). More precisely, let  $R$  be a  $\tilde{\mathcal{C}}$ -equivalence relation on the sheaf  $X$ . Then the sheaf associated with the separated presheaf*

$$i(X)/i(R) : S \longmapsto X(S)/R(S)$$

*is a universal effective quotient of  $X$  by  $R$ .*

Let  $X/R$  denote the sheaf quotient of  $X$  by  $R$ , which exists by 4.4.1(ii):

$$X/R = a(i(X)/i(R)).$$

We must show that  $X \rightarrow X/R$  is a universal effective epimorphism and that the morphism

$$f : R \longrightarrow X \times_{X/R} X$$

is an isomorphism. The first assertion has already been proved in 4.4.3. As for  $f$ , it is obtained by applying the functor  $a$  to the morphism

$$i(R) \longrightarrow i(X) \times_{i(X/R)} i(X),$$

or, since  $i(X)/i(R)$  is separated by 4.4.5(ii), so that

$$i(X)/i(R) \longrightarrow i(X/R)$$

is a monomorphism, to the canonical morphism

$$i(R) \longrightarrow i(X) \times_{i(X)/i(R)} i(X).$$

Thus we are reduced to proving the same statement in the category of presheaves. But  $i(X)/i(R)$  is the presheaf

$$S \longmapsto X(S)/R(S),$$

so the assertion reduces to its set-theoretic analogue, where it is immediate.

**Proposition 4.4.10.** *Under the hypotheses of 4.4.9, let  $Y$  be a subsheaf of  $X$ , and write  $R_Y$  for the equivalence relation induced on  $Y$  by  $R$ . Then the canonical morphism (3.1.6)*

$$Y/R_Y \longrightarrow X/R$$

*is a monomorphism. It identifies  $Y/R_Y$  with a subsheaf of  $X/R$ , namely the sheaf image of the composite*

$$Y \longrightarrow X \longrightarrow X/R.$$

The morphism of presheaves

$$i(Y)/i(R_Y) = i(Y)/i(R)_{i(Y)} \longrightarrow i(X)/i(R)$$

is a monomorphism. Since the functor  $a$  is left exact (4.3.16), it carries monomorphisms to monomorphisms; hence

$$Y/R_Y \longrightarrow X/R$$

is a monomorphism. The final assertion follows from the commutative diagram

$$\begin{array}{ccc} Y & \longrightarrow & X \\ \downarrow & & \downarrow \\ Y/R_Y & \longrightarrow & X/R \end{array}$$

and the fact that  $Y \rightarrow Y/R_Y$  is covering.

In view of this proposition, we shall always identify  $Y/R_Y$  with a subsheaf of  $X/R$ .

**Proposition 4.4.11.** *Let  $R$  be a  $\mathcal{C}$ -equivalence relation on the sheaf  $X$ . For every subsheaf  $Y$  of  $X$  stable under  $R$ , let  $Y' = Y/R_Y$ , regarded as a subsheaf of  $X' = X/R$ . Then*

$$Y = Y' \times_{X'} X,$$

*and the maps*

$$Y \longmapsto Y/R_Y, \quad Y' \longmapsto Y' \times_{X'} X$$

*define a bijective correspondence between the subsheaves  $Y$  of  $X$  stable under  $R$  and the subsheaves  $Y'$  of  $X'$ .*

If  $Y'$  is a subsheaf of  $X'$ , then  $Y' \times_{X'} X$  is a subsheaf of  $X$  stable under  $R$ .<sup>39</sup> If  $Y'$  is obtained by passage to the quotient from a subsheaf  $Y$  of  $X$ , then  $Y$  is a subobject of  $Y' \times_{X'} X$ . It is therefore enough to prove the following: if  $Y$  and  $Y_1$  are two subsheaves of  $X$ , both stable under  $R$ , with  $Y_1$  containing  $Y$ , and if their quotients  $Y/R_Y$  and  $Y_1/R_{Y_1}$  are identical, then  $Y = Y_1$ .

We may plainly reduce to the case  $Y_1 = X$ . Write

$$P = i(X)/i(R), \quad Q = i(Y)/i(R_Y)$$

for the corresponding presheaves. The diagram

$$\begin{array}{ccc} Y & \longrightarrow & Q \\ \downarrow & & \downarrow \\ X & \longrightarrow & P \end{array}$$

is Cartesian. On the other hand, there is a commutative diagram

$$\begin{array}{ccc} Q & \longrightarrow & a(Q) \\ \downarrow & & \downarrow \sim \\ P & \longrightarrow & a(P). \end{array}$$

Since  $Q \hookrightarrow a(Q)$  is covering by 4.3.11, the monomorphism  $Q \hookrightarrow P$  is covering, so  $Q$  is a refinement of  $P$ . After base change,  $Y$  is a refinement of  $X$ . Because  $X$  and  $Y$  are sheaves, 4.3.12 then gives  $Y = X$ .

**4.4.12.** In particular, if  $Y$  is a subsheaf of  $X$  and  $Y' = Y/R_Y$ , the preceding correspondence defines a subsheaf  $\bar{Y}$  of  $X$ , stable under  $R$ , containing  $Y$ , and minimal with these properties. It is called the *saturation* of  $Y$  for the equivalence relation  $R$ .

## 4.5. The Case of a Topology Coarser than the Canonical Topology

By 4.3.6 and 4.3.8, the following conditions on a topology  $\mathcal{T}$  on  $\mathcal{C}$  are equivalent:

- (i)  $\mathcal{T}$  is coarser than the canonical topology of  $\mathcal{C}$ ;
- (ii) every representable presheaf is a sheaf for  $\mathcal{T}$ ;
- (iii) every covering sieve for  $\mathcal{T}$  is universal effective epimorphic.

If  $\mathcal{T}$  is defined by a pretopology  $S \mapsto \mathcal{R}(S)$ , these conditions are also equivalent to:

- (iv) every family belonging to  $\mathcal{R}(S)$  is universal effective epimorphic.

When these conditions hold, the canonical functor  $\mathcal{C} \rightarrow \hat{\mathcal{C}}$  factors through a functor

$$j_{\mathcal{C}} = j : \mathcal{C} \longrightarrow \tilde{\mathcal{C}};$$

we also write  $j(S) = \tilde{S}$ .<sup>40</sup>

**Proposition 4.5.1.** *The functor  $j : \mathcal{C} \rightarrow \tilde{\mathcal{C}}$  is fully faithful and commutes with arbitrary inverse limits. In particular, it is left exact and therefore preserves algebraic structures defined by finite inverse limits.*

<sup>39</sup>Editor's note: moreover,  $(Y' \times_{X'} X)/R = Y'$ .

<sup>40</sup>Editor's note: we shall also write  $\mathbf{h}_S = \tilde{S}$ ; cf. the first commutative diagram in 4.5.4.

This follows immediately from the commutative factorization

$$\begin{array}{ccc} & \mathcal{C} & \\ j \swarrow & & \searrow \text{canonical} \\ \mathcal{C} & \xrightarrow{i} & \widehat{\mathcal{C}} \end{array}$$

and from 4.4.1(i).

Before giving further properties of  $j$ , we must define the topology induced on a category  $\mathcal{C}/S$ . We no longer suppose here that the given topology is necessarily coarser than the canonical topology. A sieve of an object  $T$  in  $\mathcal{C}$ , together with a morphism  $T \rightarrow S$ , naturally defines a sieve of  $T$  in  $\mathcal{C}/S$ , because the definition of a sieve of  $T$  depends only on the category

$$\mathcal{C}/T = (\mathcal{C}/S)/T.$$

For example, if the sieve is defined by a family  $\{T_i \rightarrow T\}$ , its image in  $\mathcal{C}/S$  is defined by the same family, now regarded as a family of morphisms in  $\mathcal{C}/S$ .

Declaring these images of covering sieves to be covering defines a topology on  $\mathcal{C}/S$ , called the *topology induced* by the given topology. In the terminology of [AS], 2.3, it is the coarsest topology on  $\mathcal{C}/S$  for which the canonical functor

$$i_S : \mathcal{C}/S \longrightarrow \mathcal{C}$$

is a comorphism.<sup>41</sup> By definition, the identifications

$$(\mathcal{C}/S)/T = \mathcal{C}/T$$

respect these topologies.

**Proposition 4.5.2.** *Let  $S$  be a representable sheaf on  $\mathcal{C}$ , and let  $F \rightarrow S$  be a morphism in  $\widehat{\mathcal{C}}$ . The presheaf*

$$S' \longmapsto \text{Hom}_S(S', F)$$

*on  $\mathcal{C}/S$  is separated (respectively, is a sheaf) if and only if  $F$  is a separated presheaf (respectively, a sheaf) on  $\mathcal{C}$ .*

For every functor  $P$ , one has (I, 1.4.1)

$$\text{Hom}(P, F) = \coprod_{f \in \text{Hom}(P, S)} \text{Hom}_f(P, F).$$

<sup>42</sup> Let  $S' \in \text{Ob}(\mathcal{C})$ , and let  $\eta : C' \rightarrow S'$  be a covering sieve. Since  $S$  is a sheaf, the map  $f \mapsto f \circ \eta$  is a bijection

$$\text{Hom}(S', S) \xrightarrow{\sim} \text{Hom}(C', S).$$

Consequently, the map

$$\text{Hom}_{\widehat{\mathcal{C}}}(S', F) \longrightarrow \text{Hom}_{\widehat{\mathcal{C}}}(C', F)$$

decomposes as the “disjoint union” of the maps

$$\text{Hom}_f(S', F) \longrightarrow \text{Hom}_{f \circ \eta}(C', F).$$

The proposition follows.

<sup>41</sup>Editor’s note: that is, for every object  $T \rightarrow S$  of  $\mathcal{C}/S$ , every covering sieve  $R'$  of  $i_S(T \rightarrow S) = T$ , when regarded as a sieve of the object  $(T \rightarrow S) \in \text{Ob}(\mathcal{C}/S)$ , is covering.

<sup>42</sup>Editor’s note: the argument implicit in the original has been expanded below.

**Corollary 4.5.3.** *If a topology on  $\mathcal{C}$  is coarser than the canonical topology, then the topology it induces on  $\mathcal{C}/S$  is coarser than the canonical topology of  $\mathcal{C}/S$ .*

**Corollary 4.5.4.** *Suppose that the given topology on  $\mathcal{C}$  is coarser than the canonical topology. For every  $S \in \text{Ob}(\mathcal{C})$ , there is an equivalence of categories*

$$\widetilde{\mathcal{C}}_{/\widetilde{S}} \xrightarrow{\sim} \widetilde{\mathcal{C}/S}.$$

The following diagrams commute up to isomorphism; every unlabelled arrow in them is an equivalence:

$$\begin{array}{ccccccc} \mathcal{C}_{/S} & \xrightarrow{(j_{\mathcal{C}})_{/S}} & \widetilde{\mathcal{C}}_{/\widetilde{S}} & \xrightarrow{(i_{\mathcal{C}})_{/\widetilde{S}}} & \widehat{\mathcal{C}}_{/\widehat{S}} & \xrightarrow{(a_{\mathcal{C}})_{/\widehat{S}}} & \widetilde{\mathcal{C}}_{/\widetilde{S}} \\ \parallel & & \downarrow \wr & & \downarrow \wr & & \downarrow \wr \\ \mathcal{C}_{/S} & \xrightarrow{j_{\mathcal{C}/S}} & \widetilde{\mathcal{C}/S} & \xrightarrow{i_{\mathcal{C}/S}} & \widehat{\mathcal{C}/S} & \xrightarrow{a_{\mathcal{C}/S}} & \widetilde{\mathcal{C}/S}. \end{array}$$

If  $T \rightarrow S$  is an object of  $\mathcal{C}/S$ , one also has the commutative diagram

$$\begin{array}{ccc} (\widetilde{\mathcal{C}}_{/\widetilde{S}})_{/\widetilde{T}} & \longrightarrow & (\widetilde{\mathcal{C}/S})_{/\widetilde{T}} \\ \downarrow \wr & & \searrow \\ \widetilde{\mathcal{C}}_{/\widetilde{T}} & \longrightarrow & \widetilde{\mathcal{C}/T} \end{array} \quad \begin{array}{c} \nearrow \\ \widetilde{\mathcal{C}/S}_{/T} \end{array}$$

The commutativity of the first two squares follows from the definition of the equivalence  $\widetilde{\mathcal{C}}_{/\widetilde{S}} \rightarrow \widetilde{\mathcal{C}/S}$ . To prove the commutativity of the last square, it is enough to observe that the following square commutes:

$$\begin{array}{ccc} \widehat{\mathcal{C}}_{/\widehat{S}} & \xrightarrow{L'} & \widehat{\mathcal{C}}_{/\widehat{S}} \\ \downarrow & & \downarrow \\ \widehat{\mathcal{C}/S} & \xrightarrow{L''} & \widehat{\mathcal{C}/S}. \end{array}$$

Here  $L'$  is the restriction of the functor  $L_{\mathcal{C}}$  to  $\widehat{\mathcal{C}}_{/\widehat{S}}$ , and  $L''$  is the functor  $L_{\mathcal{C}/S}$ . This is seen immediately by returning to the definition of the functors  $L$ , given after 4.3.9. The second diagram is simply the restriction to the categories of sheaves of the corresponding diagram for the categories of presheaves (Exposé I, no. 1), which is commutative.

**Scholium 4.5.5.** The preceding assertions show that, when the given topology is coarser than the canonical topology, one may identify  $\mathcal{C}$  with a full subcategory of  $\widetilde{\mathcal{C}}$ , itself a full subcategory of  $\widehat{\mathcal{C}}$ . Under these identifications one may make the customary abuses of language concerning  $\mathcal{C} \rightarrow \widehat{\mathcal{C}}$ , justified by the preceding commutative diagrams. In particular, the first diagram of 4.5.4 shows that the functor  $a$  may be used without any special precaution.

In the next subsection we shall see that the identification of  $\mathcal{C}$  with a full subcategory of  $\widetilde{\mathcal{C}}$ , unlike the corresponding identification with  $\widehat{\mathcal{C}}$ , commutes with the formation of certain direct limits; we shall then explain how this fact is to be used.

From now on, unless expressly stated otherwise, we assume that the given topology is coarser than the canonical topology, and we systematically make the identifications described above.

**Proposition 4.5.6.** *Let  $F$  and  $G$  be two sheaves over  $S$ , and let  $f : F \rightarrow G$  be an  $S$ -morphism. The following conditions on  $f$  are equivalent:*

- (i)  $f$  is a monomorphism (respectively, an epimorphism, respectively, an isomorphism) in  $\mathcal{C}$ ;

- (ii)  $f$  is a monomorphism (respectively, an epimorphism, respectively, an isomorphism) in  $\widetilde{\mathcal{C}}/S = \widetilde{\mathcal{C}}/S$ .

For monomorphisms and isomorphisms this is immediate, since it is a question about presheaves. For epimorphisms it follows from the description of epimorphisms as covering morphisms (4.4.3) and from the fact that, by the definition of the induced topology, these are the same in  $\widetilde{\mathcal{C}}$  and in  $\widetilde{\mathcal{C}}/S$ .

**Proposition 4.5.7.** *Let  $f : F \rightarrow G$  be a morphism of sheaves. The following conditions are equivalent:*

- (i)  $f$  is a monomorphism (respectively, an epimorphism, respectively, an isomorphism);
- (ii) for every  $S \in \text{Ob}(\mathcal{C})$ , the morphism  $f_S : F_S \rightarrow G_S$  is locally a monomorphism (respectively, an epimorphism, respectively, an isomorphism), that is:
- (iii) for every  $S \in \text{Ob}(\mathcal{C})$ , the set of morphisms  $T \rightarrow S$  for which  $F_T \rightarrow G_T$  is a monomorphism (respectively, an epimorphism, respectively, an isomorphism) is a refinement of  $S$ .

If the given topology is defined by a pretopology  $\mathcal{R}$ , these conditions are also equivalent to:

- (iv) for every  $S \in \text{Ob}(\mathcal{C})$ , there is a covering family  $\{S_i \rightarrow S\} \in \mathcal{R}(S)$  such that, for every  $i$ ,  $F_{S_i} \rightarrow G_{S_i}$  is a monomorphism (respectively, an epimorphism, respectively, an isomorphism).

If  $\mathcal{C}$  has a final object  $e$ , it is enough in conditions (ii), (iii), and (iv) to take  $S = e$ .

Clearly (ii), (iii), and (iv) are equivalent. To prove the equivalence of (i) and (ii), as well as the assertion concerning a final object, it is enough to show that (ii) implies (i) and that the properties in question are stable under base change. We prove the latter point first. For monomorphisms and isomorphisms it is immediate, since it is a question about presheaves. For epimorphisms it follows from the fact that every epimorphism of sheaves is universal (4.4.3).

It remains to show that (ii) implies (i). Suppose first that  $f_S : F_S \rightarrow G_S$  is locally a monomorphism (respectively, an isomorphism). There is then a covering sieve  $C$  of  $S$  such that  $f_T$  is a monomorphism (respectively, an isomorphism) for every  $T \rightarrow S$  belonging to  $C$ . Since an inverse limit of monomorphisms (respectively, isomorphisms) is again one,  $\text{Hom}(C, f)$  is a monomorphism (respectively, an isomorphism); see 4.1.4. The commutative diagram

$$\begin{array}{ccc} \text{Hom}(C, F) & \xrightarrow{\text{Hom}(C, f)} & \text{Hom}(C, G) \\ \uparrow \wr & & \uparrow \wr \\ F(S) & \xrightarrow{f(S)} & G(S) \end{array}$$

therefore shows that  $f(S)$  is injective (respectively, bijective).

Finally, suppose that  $f : F \rightarrow G$  is locally an epimorphism, and let  $H \subset G$  be its image. For every  $S \rightarrow G$ , the sheaf  $H \times_G S$  is the image of

$$f \times_G S : F \times_G S \longrightarrow S.$$

To prove that  $f$  is an epimorphism, it is enough to prove that  $H$  is a refinement of  $G$ , or equivalently that  $H \times_G S$  is a refinement of  $S$  for every  $S$ . But this holds after each base change  $T \rightarrow S$  belonging to a refinement of  $S$ , since  $f \times_G S$  is locally covering. Hence  $H \times_G S$  is itself a refinement of  $S$ , by Axiom (T 2).

**Corollary 4.5.8.** *Let  $F$  and  $G$  be two sheaves over  $S$ , and let  $f : F \rightarrow G$  be an  $S$ -morphism. Then  $f$  is a monomorphism, respectively an epimorphism, respectively an isomorphism, if and only if it is so locally on  $S$ .*

**Remark 4.5.9.** The proof of the proposition shows that its assertion about monomorphisms (respectively, isomorphisms) remains valid if one assumes only that  $F$  is a separated presheaf (respectively, a sheaf) and  $G$  is an arbitrary presheaf (respectively, a separated presheaf).

We return temporarily to the case of an arbitrary topology and make the following definition.

**Definition 4.5.10.** *Let  $G \rightarrow F$  be a morphism in  $\widehat{\mathcal{C}}$ . We say that  $G$  is a relative sheaf over  $F$  if, for every  $F$ -functor  $H$  and every refinement  $R$  of  $H$ , the canonical map*

$$\mathrm{Hom}_F(H, G) \xrightarrow{\sim} \mathrm{Hom}_F(R, G) \quad (+)$$

*is bijective.*

Proposition 4.5.2 immediately gives the following generalization.

**Proposition 4.5.11.** *If  $F$  is a sheaf, then  $G$  is a relative sheaf over  $F$  if and only if  $G$  is a sheaf.*

**Lemma 4.5.12.** *In a sequence*

$$X \longrightarrow T \longrightarrow S,$$

*where  $X, T, S$  are three objects of  $\widehat{\mathcal{C}}$ , suppose that  $X$  is a relative sheaf over  $T$ . Then*

$$U = \prod_{T/S} X$$

*is a relative sheaf over  $S$ .*

Indeed, for every  $S$ -functor  $Y$ , one has

$$\mathrm{Hom}_S(Y, U) = \mathrm{Hom}_T(T \times_S Y, X).$$

If  $C$  is a sieve of  $Y$ , then  $T \times_S C$  is a sieve of  $T \times_S Y$ , and the conclusion is immediate.

**Corollary 4.5.13.** *The presheaves  $\mathrm{Hom}_{T/S}(X, Y)$ ,  $\mathrm{Isom}_S(X, Y)$ , and the analogous presheaves are sheaves whenever all the arguments occurring in them are sheaves.*

Indeed, all these presheaves are constructed using fiber products and presheaves  $\prod_{T/S} X$  (I, 1.7 and II, 1). It is therefore enough to verify the assertion for a presheaf  $\prod_{T/S} X$ ; in that case it follows from 4.5.11 and 4.5.12.

## 4.6. Description of the Quotient of a Sheaf by an Equivalence Relation

Recall that the given topology  $\mathcal{T}$  is assumed to be coarser than the canonical topology.

**Proposition 4.6.1.** *Let  $R \xrightarrow[p_2]{p_1} X$  be a  $\widetilde{\mathcal{C}}$ -equivalence relation in the sheaf  $X$ . Define  $F \in \mathrm{Ob}(\widehat{\mathcal{C}})$  as follows: for each  $S \in \mathrm{Ob}(\mathcal{C})$ ,<sup>43</sup>*

$$F(S) = \left\{ \begin{array}{l} \text{sub-}S\text{-sheaves } Z \text{ of } X_S, \text{ stable under } R \times S, \text{ whose quotient} \\ \text{by } R_Z \text{ is } S, \text{ that is, for which } R_Z \rightrightarrows Z \rightarrow S \text{ is exact} \end{array} \right\}.$$

<sup>43</sup>Editor's note:  $R \times S$  denotes the equivalence relation in  $X \times S$  defined by  $R \times S_{\mathrm{diag}} \subset X \times X \times S \times S$ , and  $R_Z$  denotes the equivalence relation that it induces on  $Z$  (cf. 3.1.6).

For every sheaf  $Y$ ,  $\text{Hom}(Y, F)$  is naturally identified with the set

$$\left\{ \begin{array}{l} \text{sub-}Y\text{-sheaves of } X \times Y, \text{ stable under } R \times Y, \\ \text{and having quotient } Y \end{array} \right\}.$$

In particular, the subsheaf  $R \subset X \times X$  corresponds to an element  $p \in \text{Hom}(X, F)$ , and the exact diagram

$$R \xrightarrow[p_2]{p_1} X \xrightarrow{p} F$$

identifies  $F$  with the sheaf quotient  $X/R$ .

Indeed, put  $Q = X/R$ . For every sheaf  $Y$  and every morphism  $f \in \text{Hom}(Y, Q)$ , corresponding to a section  $s : Y \hookrightarrow Q \times Y$ , consider the diagram

$$\begin{array}{ccccc} R \times Y & \rightrightarrows & X \times Y & \longrightarrow & Q \times Y \\ & & \uparrow & & \uparrow_s \\ & & Z & \longrightarrow & Y, \end{array} \quad (*)$$

in which the right-hand square is cartesian. By 4.4.11,  $Z$  is immediately seen to be a sub- $Y$ -sheaf of  $X \times Y$ , stable under  $R \times Y$ , with quotient  $Y$ . Conversely, every  $Z$  of this kind arises from a unique section of  $Q \times Y$  over  $Y$ . Taking  $Y$  first to be representable gives an isomorphism  $Q \simeq F$ ; taking arbitrary  $Y$  then gives the asserted description of  $\text{Hom}(Y, F)$ . Finally, the canonical morphism  $X \rightarrow Q$  corresponds to the sub- $X$ -sheaf  $R \subset X \times X$ , which completes the proof.

**Corollary 4.6.2.** *Let  $G$  be any subfunctor of  $F$  such that  $\text{Hom}(X, G) \subset \text{Hom}(X, F)$  contains  $R$ . Then the canonical morphism  $p : X \rightarrow F$  factors through  $G$ .*

Since  $p$  is covering (4.4.9 and 4.4.3), it follows that  $G$  is a refinement of  $F$ . In particular, every subsheaf  $G$  of  $F$  satisfying the preceding condition equals  $F$  (4.3.12).

**4.6.3.** We now consider the case in which  $X$  and  $R$  are representable. We begin with some terminology. Besides the conditions (a)–(d) introduced in 3.4.1, we shall use the following conditions on a family  $(M)$  of morphisms of  $\mathcal{C}$ ; for completeness, we repeat conditions (a)–(c):

- (a)  $(M)$  is stable under base change.
- (b) The composite of two elements of  $(M)$  belongs to  $(M)$ .
- (c) Every isomorphism belongs to  $(M)$ .
- ( $d_{\mathcal{T}}$ ) Every element of  $(M)$  is covering. <sup>44</sup>
- ( $e_{\mathcal{T}}$ ) Let  $f : X \rightarrow Y$  be a morphism of  $\mathcal{C}$ . If there is a refinement  $R$  of  $Y$  such that, for every  $Y' \rightarrow R$ , the morphism  $X \times_Y Y' \rightarrow Y'$  belongs to  $(M)$ , then  $f$  belongs to  $(M)$ .

Conditions (a) and (b) imply

- ( $a'$ ) the fiber product of two elements of  $(M)$  belongs to  $(M)$ .

On the other hand, by 4.3.9, conditions (a) and ( $d_{\mathcal{T}}$ ) imply

- ( $d'$ ) every element of  $(M)$  is a universal effective epimorphism.

<sup>44</sup>Editor's note:  $\mathcal{T}$  denotes the given topology on  $\mathcal{C}$ , which is coarser than the canonical topology.



**4.6.4.** When  $\mathcal{C}$  has fiber products, the family of covering morphisms, denoted  $(M_{\mathcal{T}})$ , satisfies the preceding conditions. Indeed (cf. 4.2.3), (a) follows from (C 1), (b) from (C 2), (c) from (C 4),  $(d_{\mathcal{T}})$  from the definition, and  $(e_{\mathcal{T}})$  from (C 5). Thus the results established below for a family satisfying these conditions apply in particular to  $(M_{\mathcal{T}})$ . In particular,  $\mathcal{T}$  may be taken to be the canonical topology and  $(M)$  to be the family of universal effective epimorphisms.

**Lemma 4.6.5.** *Let  $(M)$  be a family of morphisms satisfying conditions (a)– $(e_{\mathcal{T}})$  above. Let  $R$  be a  $\mathcal{C}$ -equivalence relation of type  $(M)$  in  $X \in \text{Ob}(\mathcal{C})$ . Let  $\tilde{X}$  be the sheaf defined by  $X$ , let  $\tilde{R}$  be the  $\mathcal{C}$ -equivalence relation in  $\tilde{X}$  defined by  $R$ , and let  $\tilde{X}/\tilde{R}$  be the sheaf quotient. Then  $R$  is  $(M)$ -effective if and only if  $\tilde{X}/\tilde{R}$  is representable. When these conditions hold,  $\tilde{X}/\tilde{R}$  is represented by the quotient  $X/R$ .*

Suppose first that  $R$  is  $(M)$ -effective, and put  $Y = X/R$ . The canonical morphism  $p : X \rightarrow Y$  belongs to  $(M)$ , hence is covering by  $(d_{\mathcal{T}})$ . The corresponding morphism

$$\tilde{p} : \tilde{X} \longrightarrow \tilde{X}/\tilde{R}$$

is therefore a universal effective epimorphism in  $\mathcal{C}$  (4.4.3). Consequently it identifies  $\tilde{X}/\tilde{R}$  with the quotient of  $\tilde{X}$  by the equivalence relation  $R'$  defined by  $\tilde{p}$ . Since the canonical functor  $\mathcal{C} \rightarrow \tilde{\mathcal{C}}$  commutes with fiber products,  $R'$  is precisely  $\tilde{R}$ , because  $R$  is the equivalence relation defined by  $p$ .

Conversely, suppose that  $\tilde{X}/\tilde{R}$  is represented by an object  $Y$  of  $\mathcal{C}$ . Let  $p : X \rightarrow Y$  be the morphism deduced from the canonical morphism  $\tilde{X} \rightarrow \tilde{X}/\tilde{R}$ ; by 4.4.3 it is covering. As before,  $R$  is plainly the equivalence relation defined by  $p$ . It remains only to show that  $p \in (M)$ . The cartesian diagram

$$\begin{array}{ccccc} R & \xrightarrow{\sim} & X \times_Y X & \longrightarrow & X \\ p_2 \searrow & & \downarrow \text{pr}_2 & & \downarrow p \\ & & X & \xrightarrow{p} & Y \end{array}$$

shows that, after the covering base change  $p$ , the morphism  $p$  becomes  $p_2$ , which belongs to  $(M)$ . The assertion follows from  $(e_{\mathcal{T}})$ .

**Corollary 4.6.5.1.** <sup>45</sup> *Let  $(M)$  satisfy conditions (a)– $(e_{\mathcal{T}})$  above, and let  $f : G \rightarrow G'$  be a morphism of  $\mathcal{C}$ -groups such that  $f \in (M)$ . Suppose that  $\text{Ker}(f)$  is representable (as is the case when  $\mathcal{C}$  has a final object  $e$ ). Then the equivalence relation in  $G$  defined by  $H = \text{Ker}(f)$  is  $(M)$ -effective, and  $G'$  represents the sheaf quotient  $\tilde{G}/\tilde{H}$  for the topology  $\mathcal{T}$ .*

This follows from 4.6.5 and 3.4.7.1.

We can now state the main result of this subsection.

**Theorem 4.6.6.** *Let  $(M)$  be a family of morphisms satisfying Axioms (a)– $(e_{\mathcal{T}})$  of 4.6.3. Let  $R$  be a  $\mathcal{C}$ -equivalence relation of type  $(M)$  (cf. 3.4.3) in an object  $X$  of  $\mathcal{C}$ . Consider the functor  $F \in \text{Ob}(\widehat{\mathcal{C}})$  defined by*

$$F(S) = \left\{ \begin{array}{l} \text{sub-}S\text{-sheaves } Z \text{ of } X_S, \text{ stable under } R \times S, \\ \text{whose quotient by } R_Z \text{ is } S \end{array} \right\}.$$

Let  $F_0$  be the subfunctor of  $F$  defined as follows:  $F_0(S)$  consists of the representable  $Z \in F(S)$ , that is,

$$F_0(S) = \left\{ \begin{array}{l} \text{sub-}\mathcal{C}/S\text{-objects } Z \text{ of } X_S, \text{ stable under } R \times S, \text{ such that} \\ R_Z \text{ is } (M)\text{-effective with quotient } S, \text{ i.e. such that} \\ Z \rightarrow S \text{ belongs to } (M) \text{ and } R_Z \simeq Z \times_S Z \end{array} \right\}.$$

Then:

<sup>45</sup>Editor's note: This corollary has been added.

- (i) The morphism  $p \in \text{Hom}(X, F) = F(X)$  defined by the subobject  $R \subset X \times X$  identifies  $F$  with the sheaf quotient of  $X$  by  $R$ .
- (ii) The following conditions are equivalent:
  - (a)  $F$  is representable;
  - (b)  $F_0$  is representable;
  - (c)  $R$  is  $(M)$ -effective.

Under these conditions,  $F = F_0 = X/R$ .

- (iii) Let  $(N)$  be a family of morphisms stable under base change, such that for every covering family  $\{S_i \rightarrow S\}$  and every family  $\{T_i \rightarrow S_i\}$  of morphisms in  $(N)$ , every descent datum on the  $T_i$  relative to  $\{S_i \rightarrow S\}$  is effective. Suppose that  $X$  is squarable (cf. 1.6.0) and that  $R \rightarrow X \times X$  belongs to  $(N)$ . Then  $F_0 = F$ .

*Proof.* Assertion (i) was proved in 4.6.1.

For (ii), we have already proved the equivalence of (a) and (c), as well as the equality  $F = X/R$ . It remains to prove that either (b) or (c) implies  $F_0 = F$ . First note, as stated in the theorem, that  $F_0$  is indeed a subfunctor of  $F$ : if  $S \in \text{Ob}(\mathcal{C})$  and  $Z \in F_0(S)$ , then  $Z \rightarrow S$  is squarable, and hence  $Z \times_S S' \in F_0(S')$  for every  $S' \rightarrow S$ . Since  $R \in F(X)$  belongs to  $F_0(X)$ , Corollary 4.6.2 shows that (b) implies  $F_0 = F$ .

Now suppose that (c) holds, and let  $Q$  be an object of  $\mathcal{C}$  representing  $X/R$ . Then  $X \rightarrow Q$  belongs to  $(M)$ . For every  $S \in \text{Ob}(\mathcal{C})$  and every  $Z \in F(S)$ , diagram (\*) of 4.6.1 gives

$$Z = S \times_{Q \times S} (X \times S).$$

Thus  $Z$  is representable and  $Z \rightarrow S$  belongs to  $(M)$ , so  $Z \in F_0(S)$ .

For (iii), let  $f \in \text{Hom}(S, F)$  correspond to  $Z \in F(S)$ . We must show that  $f$  factors through  $F_0$ , or equivalently that  $Z$  is representable. This is immediate when  $f$  factors through  $X$ , by the following lemma.

**Lemma 4.6.7.** *Let  $x_0 \in X(S)$ . The image of  $x_0$  in  $F(S)$  corresponds to the subsheaf  $Z \subset X_S$  defined by the two cartesian squares*

$$\begin{array}{ccccc} X_S & \xrightarrow{\text{id}_{X_S} \times x_0} & X_S \times_S X_S & \longrightarrow & X \times X \\ \uparrow & & \uparrow & & \uparrow \\ Z & \longrightarrow & R_S & \longrightarrow & R. \end{array}$$

This follows at once from the description of the morphism  $X \rightarrow F$ .

Returning to the proof of the theorem, suppose first that  $f$  factors through  $X$ . Then  $Z$  is representable; moreover, since  $R \rightarrow X \times X$  belongs to  $(N)$ , so does  $Z \rightarrow X_S$ .

In general,  $f$  need not factor through  $X$ . Since  $X \rightarrow F$  is covering (4.4.3), however, 4.4.8(vii) gives a covering family  $\{S_i \rightarrow S\}$  and, for each  $i$ , a morphism  $S_i \rightarrow X$  making the diagram

$$\begin{array}{ccc} X & \longrightarrow & F \\ \uparrow & & \uparrow f \\ S_i & \longrightarrow & S \end{array}$$

commute. By the preceding argument, the morphism  $f_i : S_i \rightarrow F$  defined by this diagram belongs to  $\text{Hom}(S_i, F_0)$  and corresponds to the subsheaf  $Z \times_S S_i \subset X_{S_i}$ . The morphism  $Z \times_S S_i \rightarrow X_{S_i}$  belongs to  $(N)$ , and the family  $\{X_{S_i} \rightarrow X_S\}$  is covering. It therefore remains only to prove the following proposition.

**Proposition 4.6.8.** *Let  $\{S_i \rightarrow S\}$  be a covering family and let  $Z$  be a sheaf over  $S$ . Suppose that, for every  $i$ , the  $S_i$ -functor  $Z \times_S S_i$  is represented by an object  $T_i$ . Then the family  $(T_i)$  carries a canonical descent datum relative to  $\{S_i \rightarrow S\}$ . The sheaf  $Z$  is representable if and only if this datum is effective; in that case the descended object represents  $Z$ .*

First observe that, by 4.4.3,  $\{S_i \rightarrow S\}$  is a universal effective epimorphic family in  $\widetilde{\mathcal{C}}$ , and hence a descent family in  $\mathcal{C}$  (2.3). If  $Z$  is represented by an object  $T$ , then  $T \times_S S_i$ , regarded as a sheaf, is isomorphic to  $Z \times_S S_i$ . Thus the descent datum on the  $T_i$  is effective, and its uniquely determined descended object is isomorphic to  $T$ , hence represents  $Z$ .

Conversely, suppose that the canonical descent datum on the  $T_i$  is effective, and let  $T$  be the descended object. Because  $\{S_i \rightarrow S\}$  is a descent family in  $\mathcal{C}$ , there is an  $S$ -morphism  $T \rightarrow Z$  whose base change to each  $S_i$  is the canonical morphism  $T_i \rightarrow Z \times_S S_i$ . This morphism is locally an isomorphism; since both  $T$  and  $Z$  are sheaves, 4.5.8 implies that it is an isomorphism.

**Corollary 4.6.9.** *Let  $R$  be an  $(M)$ -effective equivalence relation in  $X$ . For every sheaf  $F$ , the map*

$$\mathrm{Hom}(X/R, F) \longrightarrow \mathrm{Hom}(X, F)$$

*identifies the first set with the subset of the second consisting of morphisms compatible with  $R$ .*

**Corollary 4.6.10.** *Let  $\mathcal{T}'$  be a topology coarser than  $\mathcal{T}$ , for which the morphisms in  $(M)$  are covering. Under the hypotheses of 4.6.6(iii),  $X/R$  is also the sheaf quotient of  $X$  by  $R$  for every topology lying between  $\mathcal{T}'$  and the canonical topology.*

**Remark 4.6.11.** If, in the statement of 4.6.6(iii), one assumes in addition that the descended morphism  $T \rightarrow S$  belongs to  $(N)$  whenever the hypotheses there are satisfied, then the inclusion morphisms  $Z \hookrightarrow X_S$  also belong to  $(N)$ . This follows immediately from the construction of  $Z$  by descent.

**Remark 4.6.12.** The implications

$$(c) \implies (b) \implies (a) \quad \text{and} \quad (c) \implies [F_0 = F = X/R]$$

were proved without using the sufficiency direction of Lemma 4.6.5, the only place where condition  $(e\mathcal{T})$  enters. They therefore remain valid when  $(M)$  satisfies only conditions (a)– $(d\mathcal{T})$ . The family of squarable covering morphisms is an example of such a family  $(M)$  (compare 4.6.4). For the canonical topology these are precisely the universal effective epimorphisms. Hence:

**Corollary 4.6.13.** *Let  $R$  be a universal effective equivalence relation in  $X$ . Then the object  $X/R$  of  $\mathcal{C}$  is the sheaf quotient of  $X$  by  $R$  for the canonical topology. It represents the following functor:  $(X/R)(S)$  is the set of sub- $\mathcal{C}/S$ -objects  $Z$  of  $X_S$ , stable under  $R \times S$ , for which the induced equivalence relation is universal effective and has quotient  $S$ .*

Similarly, for an arbitrary topology:

**Corollary 4.6.14.** *Let  $(M)$  be the family of squarable covering morphisms. If  $R$  is an  $(M)$ -effective equivalence relation in  $X$ , then the object  $X/R$  of  $\mathcal{C}$  is the sheaf quotient of  $X$  by  $R$ , and it represents the functor  $F_0$  of 4.6.6.*

**Scholium 4.6.15.** We can now make the following clarifications to 4.5.5. In questions involving inverse limits alone—fiber products, algebraic structures, and so forth—the results of Exposé I and 4.5.5 allow us to identify  $\mathcal{C}$ , without distinction, with a full subcategory of either  $\mathcal{C}$  or  $\widetilde{\mathcal{C}}$ . The same is not true for questions that mix inverse and direct limits.

Whenever both inverse and direct limits occur, and especially when passing to quotients (for example, when putting a group structure on the quotient of a group by an invariant subgroup), we shall regard the given category as embedded in the category of sheaves. Thus, if  $R$  is a  $\mathcal{C}$ -equivalence relation in an object  $X$  of  $\mathcal{C}$ , the notation  $X/R$  will mean the sheaf quotient of  $X$  by  $R$ , formerly denoted  $j(X)/j(R)$ , and, when this sheaf is representable, also the object that represents it. The preceding results show that, in the most important cases, a quotient in  $\mathcal{C}$  is also a quotient in the category of sheaves. In any event, we shall not use the notation  $X/R$  for a quotient in  $\mathcal{C}$  that does not coincide with the quotient in  $\widetilde{\mathcal{C}}$ —for instance, one that is not universal. This modifies the definitions of no. 3.

To study a problem of this kind, one therefore works first in the category of sheaves, where all the usual results hold (cf. no. 4.4), and then specializes the resulting statements to the original category, using the results of the present subsection and, when available, criteria for the effectivity of descent. Examples of this method will appear in the following subsections.

#### 4.7. Use of Effectivity Criteria: Isomorphism Theorem

In this subsection we give an example of the use of effectivity criteria. The initial data are a topology  $\mathcal{T}$  on  $\mathcal{C}$ , still assumed coarser than the canonical topology; a family  $(M)$  of morphisms of  $\mathcal{C}$  satisfying Axioms (a)– $(e_{\mathcal{T}})$  of 4.6.3; and a family  $(N)$  of morphisms of  $\mathcal{C}$  which may satisfy the following axioms:

- (a)  $(N)$  is stable under base change;
- $(f_{\mathcal{T}})$  morphisms in  $(N)$  descend for the given topology. More precisely, for every  $S \in \text{Ob}(\mathcal{C})$ , every covering family  $\{S_i \rightarrow S\}$ , and every family  $\{T_i \rightarrow S_i\}$  of morphisms in  $(N)$ , every descent datum on the  $T_i$  relative to  $\{S_i \rightarrow S\}$  is effective; if  $T$  denotes the descended object, then  $T \rightarrow S$  belongs to  $(N)$ .

Since every element of  $(M)$  is covering (condition 4.6.3  $(d_{\mathcal{T}})$ ), Axiom  $(f_{\mathcal{T}})$  implies the following axiom:<sup>46</sup>

- $(f_M)$  if  $Y \rightarrow X$  belongs to  $(N)$  and  $X \rightarrow X_1$  belongs to  $(M)$ , every descent datum on  $Y$  relative to  $X \rightarrow X_1$  is effective; if  $Y_1$  denotes the descended object, then  $Y_1 \rightarrow X_1$  belongs to  $(N)$ .

Here is an example, to be treated later:  $\mathcal{C}$  is the category of schemes,  $\mathcal{T}$  is the faithfully flat quasi-compact topology,  $(M)$  is the family of faithfully flat quasi-compact morphisms, and  $(N)$  is the family of closed immersions, or the family of quasi-compact immersions.<sup>47</sup>

Taking 4.6.11 into account, the principal result of 4.6.6 may be recalled as follows.

**Proposition 4.7.1.** *Let  $X$  be a squarable object of  $\mathcal{C}$ , and let  $R$  be an equivalence relation of type  $(M)$  in  $X$ . Suppose that  $R \rightarrow X \times X$  belongs to  $(N)$ , where  $(N)$  satisfies (a) and  $(f_{\mathcal{T}})$ . Then the sheaf quotient  $X/R$  is given by*

$$(X/R)(S) = \{Z \subseteq X_S \mid Z \text{ is an } S\text{-subobject stable under } R \times S, \\ Z \rightarrow X_S \text{ belongs to } (N), \quad Z \rightarrow S \text{ is covering or belongs to } (M), \\ R_Z \simeq Z \times_S Z\}.$$

We also have the following result.

<sup>46</sup>Editor's note: Axiom  $(f_M)$ , which originally appeared before Proposition 4.7.2, has been moved here.

<sup>47</sup>Editor's note: cf. §6.4; see also VI<sub>A</sub>, 5.3.1.

**Proposition 4.7.2.** *Let  $X \in \text{Ob}(\mathcal{C})$ , let  $R$  be an  $(M)$ -effective equivalence relation in  $X$ , and let  $(N)$  be a family of morphisms satisfying (a) and  $(f_M)$ . For every subobject  $Y$  of  $X$  which is stable under  $R$  and for which  $Y \rightarrow X$  belongs to  $(N)$ , the equivalence relation  $R_Y$  induced in  $Y$  by  $R$  is  $(M)$ -effective. Moreover, the quotient*

$$Y/R_Y = Y'$$

*is a subobject of  $X' = X/R$ , and  $Y' \rightarrow X'$  belongs to  $(N)$ .*

*The map  $Y \mapsto Y'$  is a bijection from the set of subobjects  $Y$  of  $X$  which are stable under  $R$  and for which  $Y \rightarrow X$  belongs to  $(N)$ , onto the set of subobjects  $Y'$  of  $X'$  for which  $Y' \rightarrow X'$  belongs to  $(N)$ . Its inverse is*

$$Y' \mapsto Y' \times_{X'} X.$$

Indeed, since  $R$  is  $(M)$ -effective, the morphism  $X \rightarrow X'$  belongs to  $(M)$ . Let  $Y'$  be a subobject of  $X'$  such that the canonical morphism  $Y' \rightarrow X'$  belongs to  $(N)$ . Then  $Y = Y' \times_{X'} X$  is a subobject of  $X$  stable under  $R$ , and the morphisms  $Y \rightarrow X$  and  $Y \rightarrow Y'$  belong respectively to  $(N)$  and  $(M)$ , since both families are stable under base change. Let  $R_Y$  denote the equivalence relation induced in  $Y$  by  $R$ . By 4.4.11, the sheaf quotient  $Y/R_Y$  is represented by  $Y'$ ; hence 4.6.5 shows that  $R_Y$  is  $(M)$ -effective.

Conversely, let  $Y$  be a subobject of  $X$ , stable under  $R$ , such that  $Y \rightarrow X$  belongs to  $(N)$ . Since  $Y$  is stable under  $R$ , its two inverse images in

$$R = X \times_{X'} X$$

are identical, and  $Y$  is equipped with a descent datum relative to  $X \rightarrow X'$ . The required conclusion follows from Axiom  $(f_M)$  for the family  $(N)$ .

**Corollary 4.7.3.** *Let  $X \in \text{Ob}(\mathcal{C})$ , and let  $R$  be an  $(M)$ -effective equivalence relation in  $X$ . Suppose in addition that  $R \rightarrow X \times X$  belongs to  $(N)$ , where  $(N)$  satisfies (a) and  $(f_T)$ . Then, for every  $Y$  as in 4.7.2,  $R_Y \rightarrow Y \times Y$  also belongs to  $(N)$ , and consequently 4.7.1 gives*

$$\begin{aligned} (Y/R_Y)(S) = \{ & Z \subseteq Y_S \mid Z \text{ is an } S\text{-subobject stable under } R_Y \times S, \\ & Z \rightarrow Y_S \text{ belongs to } (N) \text{ (hence so does } Z \rightarrow X_S), \\ & Z \rightarrow S \text{ is covering, } R_Z \simeq Z \times_S Z \}. \end{aligned}$$

## 5. Passage to the Quotient and Algebraic Structures

### 5.1. Principal Homogeneous Bundles

**Definition 5.1.0.**<sup>48</sup> We recall (III 0.1) that an object  $X$  equipped with a right action of  $H$  is said to be *formally principal homogeneous*<sup>49</sup> under  $H$  if the canonical morphism of functors

$$X \times H \longrightarrow X \times X, \quad (x, h) \longmapsto (x, xh),$$

is an isomorphism. Equivalently (cf. *loc. cit.*), for every  $S \in \text{Ob } \mathcal{C}$ , the set  $X(S)$  is formally principal homogeneous under  $H(S)$ : it is either empty or principal homogeneous under

<sup>48</sup>Editor's note: the numbering 5.1.0 has been introduced for later reference.

<sup>49</sup>Editor's note: one also says "pseudo-torsor"; cf. EGA IV<sub>4</sub>, 16.5.15. The more general notion of a formally homogeneous object (not necessarily principal homogeneous) is defined in Addendum 6.7.1 at the end of this exposé.

$H(S)$ . In particular, when  $H$  acts on itself by right translations, it is formally principal homogeneous under itself.

**Definition 5.1.1.** An object  $X$  with an action of  $H$  is said to be *trivial* if it is isomorphic, as an  $H$ -object, to  $H$  with its translation action.

**Proposition 5.1.2.** *If  $X$  is formally principal homogeneous under  $H$ , there is an isomorphism*

$$\Gamma(X) \xrightarrow{\sim} \text{Isom}_{H\text{-obj.}}(H, X)$$

*of principal homogeneous sets under  $\Gamma(H)$ .*

To a section  $x$  of  $X$ , associate the morphism  $H \rightarrow X$  defined on sets by  $h \mapsto xh$ . The assertion follows immediately by reduction to the set-theoretic case.

**Corollary 5.1.3.** *There is an isomorphism of  $H$ -objects*

$$X \xrightarrow{\sim} \text{Isom}_{H\text{-obj.}}(H, X).$$

**Corollary 5.1.4.** *An object with a group action is trivial if and only if it is formally principal homogeneous and has a section.*

**Definition 5.1.5.** Let  $\mathcal{C}$  be a category equipped with a topology. An  $S$ -object  $X$  with an action of the  $S$ -group  $H$  is called a *principal homogeneous bundle under  $H$*  if it is locally trivial, that is, if the following equivalent conditions hold:<sup>50</sup>

- (i) the collection of morphisms  $T \rightarrow S$  for which the functor  $X \times_S T$  is trivial under  $H \times_S T$  is a refinement of  $S$ ;
- (ii) for a pretopology defining the given topology, there is a covering family  $\{S_i \rightarrow S\}$  such that, for every  $i$ , the  $S_i$ -functor  $X \times_S S_i$  is trivial under the  $S_i$ -group functor  $H \times_S S_i$ , equivalently, it has a section over  $S_i$ .

**Proposition 5.1.6.** *Let  $\mathcal{C}$  be a category equipped with a topology  $\mathcal{T}$ , and let  $(M)$  be a family of morphisms of  $\mathcal{C}$  satisfying axioms (a)–(e $\mathcal{T}$ ) of 4.6.3. Let  $H$  be an  $S$ -group whose structure morphism  $H \rightarrow S$  belongs to  $(M)$ , and let  $P$  be an  $S$ -object with an action of  $H$ . The following conditions are equivalent:*

- (i)  $P$  is a principal homogeneous bundle under  $H$  (Definition 5.1.5);
- (ii)  $P$  is formally principal homogeneous under  $H$ , and its structure morphism  $P \rightarrow S$  belongs to  $(M)$ ;
- (iii) there exists a morphism  $S' \rightarrow S$  belonging to  $(M)$  such that, after base change from  $S$  to  $S'$ ,  $P$  becomes trivial; that is,  $P \times_S S'$  is trivial under  $H \times_S S'$ ;
- (iv)  $H$  acts freely and  $(M)$ -effectively on  $P$ , and the quotient  $P/H$  is isomorphic to  $S$ .

First, (ii) and (iv) are equivalent. In either case  $P \rightarrow S$  belongs to  $(M)$ , hence is squarable, which ensures that the fiber products  $H \times_S P$  and  $P \times_S P$  are representable. Condition (ii) plainly implies (iii): one may take  $S' = P$ , since formal principal homogeneity implies that  $P \times_S P$  is trivial under  $H \times_S P$  by 5.1.4, its diagonal section providing a section. Condition (iii) plainly implies (i), because  $\{S' \rightarrow S\}$  is a covering family by axiom (d $\mathcal{T}$ ).

It remains to prove (i) $\Rightarrow$ (ii). The morphism of sheaves

$$P \times_S H \longrightarrow P \times_S P$$

---

<sup>50</sup>Editor's note: in this case one also says that  $X$  is an  $H$ -torsor.

is locally an isomorphism and hence is an isomorphism (4.5.8); thus  $P$  is formally principal homogeneous. The structure morphism  $P \rightarrow S$  is locally isomorphic to the structure morphism  $H \rightarrow S$ , which belongs to  $(M)$ . It therefore belongs to  $(M)$  by axiom  $(e_{\mathcal{T}})$ .

The equivalence between (i) and (iv) admits the following generalization.

**Proposition 5.1.7.** *Under the same assumptions on  $\mathcal{C}$  and  $(M)$ , let  $H$  be an  $S$ -group and let  $X$  be an  $S$ -object on which  $H$  acts on the right. Suppose that  $H \rightarrow S$  belongs to  $(M)$ . The following conditions are equivalent:*

- (i)  $H$  acts freely and  $(M)$ -effectively on  $X$ ;
- (ii) there exists an  $S$ -morphism  $p : X \rightarrow Y$ , compatible with the equivalence relation on  $X$  defined by the action of  $H$ , such that the induced action of  $H \times_S Y$  on  $X$  over  $Y$  makes  $X$  a principal homogeneous bundle under  $H_Y$  over  $Y$ .

Under these conditions,  $p$  identifies  $Y$  with the quotient  $X/H$ .

If  $p : X \rightarrow Y$  is compatible with the action of  $H$ , then the induced action of  $H \times_S Y$  on  $X$  over  $Y$  defines the same equivalence relation on  $X$  as the original action of  $H$ , by the isomorphism

$$H_Y \times_Y X \xrightarrow{\sim} H \times_S X.$$

The proposition follows from this observation and the preceding equivalence (iv) $\Leftrightarrow$ (i).

**Corollary 5.1.7.1.**<sup>51</sup> *Let  $\mathcal{C}$  be a category having a final object and fiber products, and let it be equipped with a topology  $\mathcal{T}$  coarser than the canonical topology. Let  $f : G \rightarrow H$  be a morphism of  $\mathcal{C}$ -groups, put  $K = \text{Ker}(f)$ , and suppose that  $f$  is covering for  $\mathcal{T}$ . Then  $H$  represents the sheaf quotient  $G/K$ , and  $f$  is a  $K_H$ -torsor. Equivalently, one may say that  $G$  is a  $K$ -torsor over  $H$ .*

Indeed, because  $f$  is covering, it is a universal effective epimorphism (4.4.3). Hence, by 3.3.3.1,  $H$  is the quotient of  $G$  by the equivalence relation

$$\mathcal{R}(f) = G \times_H G.$$

Moreover, the morphism

$$G \times K \longrightarrow G \times_H G, \quad (g, k) \longmapsto (g, gk),$$

is an isomorphism of  $K_G = G \times_H K_H$ -objects; its inverse is  $(g, g') \mapsto (g, g^{-1}g')$ . Thus  $\mathcal{R}(f)$  is the equivalence relation defined by  $K$ . Since  $f : G \rightarrow H$  is covering, it is a  $K_H$ -torsor by 5.1.6(ii), or directly by Definition 5.1.5(ii).

We can now sharpen Theorem 4.6.6 in the case of a quotient by a group action.

**Proposition 5.1.8.** *Under the assumptions of 5.1.7, let  $F_0$  denote the functor over  $S$  defined as follows: for each  $S' \rightarrow S$ ,  $F_0(S')$  is the set of representable  $S'$ -subfunctors  $Z$  of  $X \times_S S'$  that are stable under  $H \times_S S'$  and are principal homogeneous bundles under this  $S'$ -group for the induced action (3.2.2).*

(i) *The following conditions are equivalent:*

- (a) *the action of  $H$  on  $X$  is  $(M)$ -effective and free;*<sup>52</sup>
- (b)  *$F_0$  is representable.*

<sup>51</sup>Editor's note: this special case has been added as a corollary; it will be used several times in the following exposés.

<sup>52</sup>Editor's note: the words "and free" have been added.

Under these conditions,  $F_0 = X/H$ .

- (ii) Let  $(N)$  be a family of morphisms stable under base change, such that for every covering family  $\{S'_i \rightarrow S'\}$  and every family of morphisms  $\{T_i \rightarrow S'_i\}$  in  $(N)$ , every descent datum on the  $T_i$  relative to  $\{S'_i \rightarrow S'\}$  is effective. Suppose that  $X \times_S H \rightarrow X \times_S X$  belongs to  $(N)$ , and that  $X$  is squarable. Then the element  $p \in \text{Hom}(X, F_0)$  corresponding to the subobject  $X \times_S H$  of  $X \times_S X$  identifies  $F_0$  with the sheaf quotient  $X/H$ .

## 5.2. Group Structures and Passage to the Quotient

In this subsection we are concerned with the algebraic structures that can be placed on the quotient  $(G/H)$  of a group by a subgroup. We first work in the category of sheaves on  $\mathcal{C}$  for an arbitrary topology. Taking the canonical topology and using 4.5.12 will then give results concerning universal effective passage to the quotient in  $\mathcal{C}$ .

**Proposition 5.2.1.** *Let  $u : H \rightarrow G$  be a monomorphism of sheaves of groups. There is a unique  $G$ -object structure on the quotient sheaf  $G/H$  for which the canonical morphism*

$$p : G \longrightarrow G/H$$

*is a morphism of  $G$ -objects. This structure is functorial in the pair  $(G, H)$ : given a commutative diagram*

$$\begin{array}{ccc} H & \longrightarrow & G \\ \downarrow & & \downarrow f \\ H' & \longrightarrow & G', \end{array}$$

*the morphism  $f : G/H \rightarrow G'/H'$  of 3.2.3 is compatible with the morphism  $f$  between the acting groups.*

Indeed,  $G/H$  is the sheaf associated with the presheaf

$$i(G)/i(H) : S \longmapsto G(S)/H(S).$$

Since the functor  $a$  is left exact, it carries objects with a group action to objects with a group action. The presheaf  $i(G)/i(H)$  is an  $i(G)$ -object, and therefore

$$G/H = a(i(G)/i(H))$$

is an  $a(i(G)) = G$ -object. This construction plainly has all the stated properties.

**Corollary 5.2.2.** *Let  $u : H \rightarrow G$  be a monomorphism of  $\mathcal{C}$ -groups, and suppose that the action of  $H$  on  $G$  is universally effective. There is a unique  $G$ -object structure on the quotient object  $G/H \in \text{Ob } \mathcal{C}$  for which  $p : G \rightarrow G/H$  is a morphism of objects with an action. In the sense of the preceding proposition, this structure is functorial in the pair  $(H, G)$ , with  $H$  acting universally effectively on  $G$ .*

**Proposition 5.2.3.** *Let  $u : H \rightarrow G$  be a monomorphism of sheaves of groups that identifies  $H$  with an invariant subgroup sheaf of  $G$ . There is a unique sheaf-of-groups structure on the quotient sheaf  $G/H$  for which the canonical morphism  $p : G \rightarrow G/H$  is a group morphism. This structure is functorial in the pair  $(H, G)$ , with  $H$  invariant.*

The proof is similar to that of 5.2.1.

**Corollary 5.2.4.** *Let  $u : H \rightarrow G$  be a monomorphism of  $\mathcal{C}$ -groups that identifies  $H$  with an invariant subgroup of  $G$ . Suppose that the action of  $H$  on  $G$  is universally effective. There*



is a unique group structure on the quotient object  $G/H \in \text{Ob } \mathcal{C}$  for which the canonical morphism  $G \rightarrow G/H$  is a group morphism. This structure is functorial in the pair  $(H, G)$ , with  $H$  invariant and acting universally effectively.

The group structure of  $G/H$  can be characterized more explicitly.

**Proposition 5.2.5.** *Under the assumptions of 5.2.4, let  $K$  be a  $\mathcal{C}$ -group and let  $f : G \rightarrow K$  be a morphism. The following conditions are equivalent:*

- (i)  *$f$  is a group morphism compatible with the equivalence relation defined by  $H$ ;*
- (ii)  *$f$  is a group morphism whose restriction  $H \rightarrow K$  is the trivial morphism;*
- (iii)  *$f$  factors through a group morphism  $G/H \rightarrow K$ .*

In particular, there is an isomorphism, functorial in the group  $K$ ,

$$\text{Hom}_{\mathcal{C}\text{-gr.}}(G/H, K) \xrightarrow{\sim} \{f \in \text{Hom}_{\mathcal{C}\text{-gr.}}(G, K) \mid f \circ u = e\}.$$

The equivalence of (i) and (ii) is proved set-theoretically, and (iii) $\Rightarrow$ (ii) is immediate. The equivalence of (iii) and (ii) follows from the formula

$$\text{Hom}(G/H, K) \simeq \text{Hom}(i(G)/i(H), K)$$

and the definition of the group structure on  $G/H$ .

**Remark 5.2.6.** In the preceding situation, if the kernel of  $f$  is exactly  $H$ , then the morphism  $G/H \rightarrow K$  through which  $f$  factors is a monomorphism. This follows at once from 3.3.4.

For sheaves of groups, 4.4.11 can be sharpened as follows.

**Proposition 5.2.7.** *Let  $G$  be a sheaf of groups and  $H$  an invariant subgroup sheaf. For every subgroup sheaf  $K$  of  $G$  containing  $H$ , let  $K' = K/H$ , regarded as a subgroup of  $G' = G/H$ . Then*

$$K = K' \times_{G'} G,$$

and the maps

$$K \longmapsto K/H, \quad K' \longmapsto K' \times_{G'} G$$

give a bijection between the subgroup sheaves of  $G$  containing  $H$  and the subgroup sheaves of  $G'$ . Under this correspondence, the invariant subgroup sheaves of  $G$  and of  $G'$  correspond.

The first assertion follows readily from 4.4.11 and 3.2.4. It remains to show that  $K$  is invariant in  $G$  if and only if  $K'$  is invariant in  $G'$ . If  $K$  is invariant in  $G$ , then the presheaf  $i(K)/i(H)$  is invariant in  $i(G)/i(H)$ ; by the usual argument, the same is true of the associated sheaves. Conversely, if  $K'$  is invariant in  $G'$ , then the fiber product  $K' \times_{G'} G$  is invariant in  $G$ , as is immediate.

Now let  $L$  be an arbitrary subgroup sheaf of  $G$ . Denote by  $\bar{L}$  the saturation of  $L$  for the equivalence relation defined by  $H$ ; we also write  $\bar{L} = L \cdot H$ .

**Proposition 5.2.8.** *Under the preceding assumptions,  $L \cdot H$  is a subgroup sheaf of  $G$  containing  $H$ , and the image of  $L$  in  $G/H$  is identified with*

$$(L \cdot H)/H \simeq L/(H \cap L).$$

Indeed, let  $L'$  denote the image sheaf of  $L$  in  $G/H$ . Under the correspondence of the preceding proposition, it is the subgroup sheaf of  $G/H$  corresponding to  $L \cdot H$ . Since  $L \rightarrow L'$  is covering, and hence a universal effective epimorphism of sheaves, it follows from 4.4.9 that  $L'$  is identified with the quotient of  $L$  by the kernel of  $L \rightarrow L'$ , which is plainly  $H \cap L$ .

Finally, consider a sheaf of groups  $G$ , a subgroup sheaf  $K$ , and an invariant subgroup sheaf  $H$  of  $K$ . We first define a right action of the sheaf of groups  $H \backslash K (= K/H)$  on  $G/H$ . The group  $K$  acts on  $G$  by right translations. Since  $H$  is invariant in  $K$ , this action is compatible with the equivalence relation defined by the action of  $H$ , and therefore induces an action of  $K$  on  $G/H$ , that is, a morphism

$$K^\circ \longrightarrow \text{Aut}(G/H)$$

from the group opposite to  $K$ . Since the target is a sheaf (4.5.13) and this morphism is trivial on  $H$ , it factors through  $K/H$  and defines the desired action. The right and left actions of  $G$  on itself commute; hence the actions of  $G$  and  $K/H$  on  $G/H$  commute.

**Proposition 5.2.9.** *Under the preceding assumptions,  $K/H$  acts freely on the right on  $G/H$ , and there is a canonical isomorphism of  $G$ -objects*

$$(G/H)/(K/H) \simeq G/K.$$

*If  $K$  is invariant in  $G$ , in which case  $K/H$  is invariant in  $G/H$  by 5.2.7, this isomorphism respects the group structures on both sides.*

There is an isomorphism of presheaves

$$(i(G)/i(H))/(i(K)/i(H)) \xleftarrow{\sim} i(G)/i(K),$$

which respects the  $i(G)$ -object structures. Applying the functor  $a$  to this relation gives the asserted result.

**Corollary 5.2.10.** *Let  $G$  be a  $\mathcal{C}$ -group,  $K$  a  $\mathcal{C}$ -subgroup of  $G$ , and  $H$  an invariant  $\mathcal{C}$ -subgroup of  $K$ . Let  $(M)$  be a family of morphisms of  $\mathcal{C}$  satisfying axioms (a)–(e $\mathcal{T}$ ). Suppose that the right action of  $H$  on  $G$  (respectively, on  $K$ ) is  $(M)$ -effective. Then  $K/H$  acts naturally and freely on the right on  $G/H$ , and this action commutes with that of  $G$ . The following conditions are equivalent:*

- (i) *the action of  $K$  on  $G$  is  $(M)$ -effective;*
- (ii) *the action of  $K/H$  on  $G/H$  is  $(M)$ -effective.*

*Under these conditions, there is an isomorphism of objects<sup>53</sup> with a  $G$ -action*

$$(G/H)/(K/H) \simeq G/K.$$

### 5.3. Use of Effectivity Criteria: Noether's Theorem

<sup>54</sup> Let  $\mathcal{C}$ ,  $\mathcal{T}$ , and  $(M)$  be as usual. Let  $(N)$  be a family of morphisms satisfying axioms (a) and  $(f_M)$  of 4.7. Combining 5.2.7 and 4.7.2 gives the following result.

**Proposition 5.3.1.** *Let  $G$  be a  $\mathcal{C}$ -group, and let  $H$  be an invariant  $\mathcal{C}$ -subgroup of  $G$  that acts  $(M)$ -effectively on  $G$ . For every  $\mathcal{C}$ -subgroup  $K$  of  $G$  containing  $H$  such that  $K \rightarrow G$  belongs to  $(N)$ , the action of  $H$  on  $K$  is  $(M)$ -effective, and the quotient  $K/H = K'$  is a  $\mathcal{C}$ -subgroup of  $G/H = G'$  such that  $K' \rightarrow G'$  belongs to  $(N)$ . The map  $K \mapsto K'$  is a bijection*

<sup>53</sup>Editor's note: of  $\mathcal{C}$ .

<sup>54</sup>Editor's note: in fact, the isomorphisms established in 5.2.8–5.2.10 deserve to be called “Noether isomorphism theorems.”

between the set of  $\mathcal{C}$ -subgroups  $K$  of  $G$  that contain  $H$  and for which  $K \rightarrow G$  belongs to  $(N)$ , and the set of  $\mathcal{C}$ -subgroups  $K'$  of  $G'$  for which  $K' \rightarrow G'$  belongs to  $(N)$ . Its inverse is

$$K' \mapsto K \times_{G'} G.$$

Under this correspondence, the invariant subgroups of  $G$  and of  $G'$  correspond.

**Corollary 5.3.2.** *If  $H \rightarrow G$  belongs to  $(N)$ , then  $\mathcal{C}$  has a final object  $e$ , and the unit section  $e \rightarrow G/H$  belongs to  $(N)$ .<sup>55</sup>*

## 6. Topologies in the Category of Schemes

### 6.1. The Zariski Topology

This is the topology generated by the following pretopology: a family of morphisms  $\{S_i \rightarrow S\}$  is covering if each morphism is an open immersion and the union of the images of the  $S_i$  is all of  $S$ . It is denoted by  $(\text{Zar})$ .

**Definition 6.1.1.** *A sheaf for the Zariski topology is also called a functor of local nature: it is a contravariant functor from  $\mathbf{Sch}$  to  $\mathbf{Set}$  such that, for every scheme  $S$  and every covering of  $S$  by open subschemes  $S_i$ , the diagram*

$$F(S) \longrightarrow \prod_i F(S_i) \rightrightarrows \prod_{i,j} F(S_i \cap S_j)$$

*is exact.*

In particular, a functor of local nature carries direct sums to products. Since every representable functor is a sheaf, the Zariski topology is coarser than the canonical topology.

As a matter of terminology, whenever we say "local" or "locally" without further qualification, the reference is to the Zariski topology and hence to the usual meaning of these words.

### 6.2. A Procedure for Constructing Topologies

**Proposition 6.2.1.** *Let  $\mathcal{C}$  be a category, let  $\mathcal{C}'$  be a full subcategory, and let  $P$  be a set of families of morphisms in  $\mathcal{C}$  with common target, stable under base change and composition (that is, satisfying axioms (P 1) and (P 2) of 4.2.5). Let  $P'$  be a set of families of morphisms in  $\mathcal{C}'$  that contains every singleton family consisting of an identity isomorphism. Equip  $\mathcal{C}$  with the topology generated by  $P$  and  $P'$  (cf. 4.2.5.0), and suppose that the following three conditions hold:*

- (a) *If  $\{S_i \rightarrow S\} \in P'$ , so that  $S_i, S \in \text{Ob } \mathcal{C}'$ , and  $T \rightarrow S$  is a morphism in  $\mathcal{C}'$ , then the fiber products  $S_i \times_S T$ , formed in  $\mathcal{C}$ , exist and the family  $\{S_i \times_S T \rightarrow T\}$  belongs to  $P'$ ; in particular,  $S_i \times_S T \in \text{Ob } \mathcal{C}'$ .*

*This condition implies that  $P'$  is stable under base change in  $\mathcal{C}'$ , but is not equivalent to that assertion: it also requires the inclusion functor  $\mathcal{C}' \hookrightarrow \mathcal{C}$  to commute with certain fiber products.*

- (b) *For every  $S \in \text{Ob } \mathcal{C}$ , there exists a family  $\{S_i \rightarrow S\} \in P$  with  $S_i \in \text{Ob } \mathcal{C}'$  for every  $i$ .*

---

<sup>55</sup>Editor's note: this follows by applying the proposition with  $K = H$ .

(c) In a situation

$$\begin{array}{ccc}
 & & S_{ijk} \\
 & & \downarrow (P') \\
 S_i & \xleftarrow{(P)} & S_{ij} \\
 \downarrow (P') & & \\
 S & & 
 \end{array} \quad (6.2.1.c)$$

where  $S, S_i, S_{ij}, S_{ijk}$  are objects of  $\mathcal{C}'$ , where  $\{S_i \rightarrow S\} \in P'$ ,  $\{S_{ij} \rightarrow S_i\} \in P$  for each  $i$ , and  $\{S_{ijk} \rightarrow S_{ij}\} \in P'$  for each pair  $ij$ , there exists a family  $\{T_n \rightarrow S\} \in P'$ , and for each  $n$  a multi-index  $(ijk)$ , fitting into a commutative diagram

$$\begin{array}{ccc}
 S_{ijk} & \xleftarrow{\quad} & T_n \\
 & \searrow & \downarrow \\
 & & S.
 \end{array} \quad (6.2.1.d)$$

Under these hypotheses, a sieve  $R$  of an object  $S \in \text{Ob } \mathcal{C}$  is covering if and only if there is a composite family

$$\begin{array}{ccc}
 R & \xleftarrow{\quad} & S_{ij} \\
 \downarrow & & \downarrow (P') \\
 S & \xleftarrow{(P)} & S_i
 \end{array} \quad (6.2.1.e)$$

where  $S_i, S_{ij} \in \text{Ob } \mathcal{C}'$ ,  $\{S_i \rightarrow S\} \in P$ , and  $\{S_{ij} \rightarrow S_i\} \in P'$  for every  $i$ , such that the resulting morphisms  $S_{ij} \rightarrow S$  factor through  $R$ . Equivalently, the sieve generated by this composite family is contained in  $R$ .

*Proof.* Families belonging to  $P$  and  $P'$  are covering, and therefore so is a composite of such families, by (C 2). A sieve of the displayed form is thus covering because it contains a covering sieve.

Conversely, it is enough to show that the sieves of this form constitute a topology, that is, to verify axioms (T 1)–(T 4) of 4.2.1.

*Axiom (T 4).* Let  $S \in \text{Ob } \mathcal{C}$ . By (b), there exists a family  $\{S_i \rightarrow S\} \in P$  with  $S_i \in \text{Ob } \mathcal{C}'$ .

The singleton families  $\{S_i \xrightarrow{\text{id}_{S_i}} S_i\}$  belong to  $P'$  by hypothesis. Hence the maximal sieve  $S$  of  $S$  has the required form:

$$\begin{array}{ccc}
 S_i & \longrightarrow & S \\
 \downarrow (P') \text{ id} & & \downarrow \text{id}_S \\
 S_i & \xrightarrow{(P)} & S.
 \end{array} \quad (6.2.1.T4)$$

*Axiom (T 3).* This is evident.

*Axiom (T 2).* Let  $R$  be a sieve of  $S$  of the required form, and let  $C$  be a sieve<sup>56</sup> of  $S$  such that, for every  $T \in \text{Ob } \mathcal{C}$  and every morphism  $T \rightarrow S$  factoring through  $R$ , the sieve  $C \times_S T$  of  $T$  has the required form. Since  $S_{ij} \rightarrow S$  factors through  $R$ , the sieve

$$C_{ij} = C \times_S S_{ij}$$

<sup>56</sup>Editor's note: the original has been corrected here.

of  $S_{ij}$  has the required form; it is represented schematically by

$$\begin{array}{ccccc}
 & S_{ij} & \longleftarrow & C_{ij} & \\
 \swarrow & (P') \downarrow & & \downarrow & \\
 & S_i & & & \\
 & (P) \downarrow & & \downarrow & \\
 R \longrightarrow & S & \longleftarrow & C. & 
 \end{array} \tag{6.2.1.T2a}$$

Consequently, for every pair  $ij$ , there is a diagram of the form

$$\begin{array}{ccc}
 S_{ijkl} & & \\
 (P') \downarrow & \searrow & \\
 S_{ijk} & & \\
 (P) \downarrow & & \\
 S_{ij} & \longleftarrow & C_{ij}.
 \end{array} \tag{6.2.1.T2b}$$

We have therefore produced a composite family

$$S_{ijkl} \xrightarrow{(P')} S_{ijk} \xrightarrow{(P)} S_{ij} \xrightarrow{(P')} S_i \xrightarrow{(P)} S \tag{6.2.1.T2c}$$

belonging to  $P \circ P' \circ P \circ P'$ , factoring through  $C$ , and having every object other than  $S$  in  $\mathcal{C}'$ .

Apply condition (c) to each family  $\{S_{ijkl} \rightarrow S_i\}$ . For every  $i$  we obtain a family  $\{T_{in} \rightarrow S_i\} \in P'$  such that  $T_{in} \rightarrow S$  factors through one of the  $S_{ijkl}$ , and hence through  $C$ :

$$\begin{array}{ccc}
 T_{in} & \longrightarrow & S_{ijkl} \\
 (P') \downarrow & & \downarrow \\
 S_i & & C \\
 (P) \downarrow & \swarrow & \\
 S & & 
 \end{array} \tag{6.2.1.T2d}$$

Thus the sieve  $C$  of  $S$  has the required form, completing the verification of (T 2).

*Axiom (T 1).* Let  $R$  be a sieve of  $S$  of the prescribed form and let  $T \rightarrow S$  be a morphism in  $\mathcal{C}$ . We show that the sieve  $R \times_S T$  of  $T$  again has the prescribed form. Consider the commutative configuration

$$\begin{array}{ccccccc}
 & S_{ij} & \longleftarrow & & U_{ikj} & & \\
 \swarrow & (P') \downarrow & & & (P') \downarrow & & \\
 & S_i & \longleftarrow & T_i & \xleftarrow{(P)} & U_{ik} & \\
 & (P) \downarrow & & (P) \downarrow & \swarrow (P) & & \\
 R \longrightarrow & S & \longleftarrow & T & & & 
 \end{array} \tag{6.2.1.T1a}$$

Put  $T_i = S_i \times_S T$ . The family  $\{T_i \rightarrow T\}$  belongs to  $P$ , by (P 1). By (b), choose families  $\{U_{ik} \rightarrow T_i\} \in P$  with  $U_{ik} \in \text{Ob } \mathcal{C}'$ . By (P 2),  $\{U_{ik} \rightarrow T\} \in P$ . Condition (a) shows that

$$U_{ik} \times_{S_i} S_{ij} = U_{ikj}$$

is an object of  $\mathcal{C}'$  and that, for each  $ik$ ,  $\{U_{ikj} \rightarrow U_{ik}\} \in P'$ . The commutative diagram

$$\begin{array}{ccccc}
 R & \longleftarrow & U_{ikj} & & \\
 \downarrow & & (P') \downarrow & & \\
 S & \longleftarrow & T & \xleftarrow{(P)} & U_{ik}
 \end{array} \tag{6.2.1.T1b}$$

shows that the morphisms  $U_{ikj} \rightarrow T$  factor through the sieve  $T \times_S R$  of  $T$ . This sieve therefore has the required form, and the proof is complete.

**Corollary 6.2.2.** *Let  $S \in \text{Ob } \mathcal{C}'$ , and let  $R$  be a sieve of  $S$ . Then  $R$  is covering if and only if there exists a family  $\{T_i \rightarrow S\} \in P'$  that factors through  $R$ .*

Such a sieve is plainly covering. Conversely, apply condition (c) with the family  $\{S_i \rightarrow S\}$  reduced to the identity isomorphism of  $S$ ; the proposition then shows that every covering sieve has the stated form.

**Corollary 6.2.3.** *A presheaf  $F$  on  $\mathcal{C}$  is separated (respectively, is a sheaf) if and only if the map*

$$F(S) \longrightarrow \prod_i F(S_i)$$

*is injective (respectively, the diagram*

$$F(S) \longrightarrow \prod_i F(S_i) \rightrightarrows \prod_{i,j} F(S_i \times_S S_j)$$

*is exact) in each of the following two cases:*

- (i)  $\{S_i \rightarrow S\} \in P$ ;
- (ii)  $S, S_i \in \text{Ob } \mathcal{C}'$  and  $\{S_i \rightarrow S\} \in P'$ .

The conditions are necessary because the families in question are covering. Let  $C$  be the sieve of  $S$  that is the image of a composite family

$$\{S_{ij} \xrightarrow{(P')} S_i \xrightarrow{(P)} S\}.$$

A straightforward diagram chase shows that the conditions of the corollary imply that

$$\text{Hom}(S, F) \xrightarrow{g} \text{Hom}(C, F)$$

is injective, respectively bijective. Every covering sieve  $R$  of  $S$  contains a sieve  $C$  of this form, and there is a commutative diagram

$$\begin{array}{ccc} \text{Hom}(S, F) & \xrightarrow{f} & \text{Hom}(R, F) \\ & \searrow g & \downarrow h \\ & & \text{Hom}(C, F). \end{array} \quad (6.2.3.1)$$

Since  $g$  is injective, so is  $f$ , and hence  $F$  is separated. Moreover,  $R$  refines  $C$ , so  $h$  is also injective. If  $g$  is bijective, then  $f$  is bijective as well, and  $F$  is a sheaf.

**Remark 6.2.4.** *The preceding corollary does not follow from 4.3.5, since  $P'$  is not stable under base extension.*

**Remark 6.2.5.** *Condition (c) holds in particular when:*

- (i)  $P'$  is stable under composition;
- (ii) whenever  $\{S_j \rightarrow S\}$  is a family of morphisms in  $\mathcal{C}'$  belonging to  $P$ , it contains a subfamily belonging to  $P'$ .

### 6.3. Application to the Category of Schemes

Take  $\mathcal{C}$  to be the category of schemes,  $\mathcal{C}'$  the full subcategory of affine schemes, and  $P$  the set of surjective families of open immersions. We shall consider the following sets  $P'_i$ :

- $P'_1$  : finite surjective families of flat morphisms;
- $P'_2$  : finite surjective families of flat, finitely presented, quasi-finite morphisms;
- $P'_3$  : finite surjective families of étale morphisms;
- $P'_4$  : finite surjective families of finite étale morphisms.

The description of  $P'_2$  is accompanied by the following clarification.<sup>57</sup>

For each  $P'_i$ , except  $P'_4$ , the conditions of Proposition 6.2.1 hold. Condition (c) follows from 6.2.5: since an affine scheme is quasi-compact, every family of morphisms in  $\mathcal{C}'$  belonging to  $P$  contains a finite subfamily that still belongs to  $P$ , and hence belongs to  $P'_i$  for  $i = 1, 2, 3$ . The topology  $T_i$  generated by  $P$  and  $P'_i$  is denoted and named as follows:<sup>57</sup>

- $T_1$  = (fpqc) the faithfully flat quasi-compact topology,
- $T_2$  = (fppf) the faithfully flat topology (locally of finite presentation),
- $T_3$  = (ét) the étale topology,
- $T_4$  = (étf) the finite étale topology.

Since  $P'_1 \supset P'_2 \supset P'_3 \supset P'_4$ , one has

$$(\text{fpqc}) \geq (\text{fppf}) \geq (\text{ét}) \geq (\text{étf}) \geq (\text{Zar}).$$

#### Proposition 6.3.1.

- (i) A sieve  $R$  of  $S$  is covering for  $T_i$ ,  $1 \leq i \leq 3$ , if and only if there exist a covering  $(S_p)$  of  $S$  by affine open subschemes and, for every  $p$ , a family  $\{S_{pq} \rightarrow S_p\} \in P'_i$ , with all  $S_{pq}$  affine, such that every morphism  $S_{pq} \rightarrow S$  factors through  $R$ .<sup>58</sup>
- (ii) A presheaf  $F$  on **Sch** is a sheaf for (fpqc) (respectively, (fppf), (ét), or (étf)) if and only if:
  - (a)  $F$  is a sheaf for (Zar), that is, a functor of local nature;
  - (b) for every faithfully flat (respectively, faithfully flat, finitely presented and quasi-finite; surjective étale; or surjective finite étale) morphism  $T \rightarrow S$ , with  $T$  and  $S$  affine, the diagram

$$F(S) \longrightarrow F(T) \rightrightarrows F(T \times_S T)$$

is exact.

- (iii) The topologies  $T_i$ , for  $i = 1, 2, 3, 4$ , are coarser than the canonical topology.
- (iv) Every surjective family of flat open morphisms (respectively, flat morphisms locally of finite presentation; étale morphisms; or finite étale morphisms) is covering for (fpqc) (respectively, (fppf), (ét), or (étf)).

<sup>57</sup>Editor's note: let  $P'_2''$  be the set of finite surjective families of morphisms in  $\mathcal{C}'$  that are flat and of finite presentation. By Proposition 6.3.1 below, the topology  $T_2$  generated by  $P$  and  $P'_2$  coincides with the topology generated by  $P$  and  $P'_2''$ . This follows from the results of EGA IV<sub>4</sub>, §17.16 on quasi-sections; see the proof of 6.3.1. We recall the following special case of EGA IV<sub>4</sub>, 17.16.2. Let  $S$  be affine and let  $f : X \rightarrow S$  be a surjective flat morphism locally of finite presentation. Then there exist a faithfully flat, finitely presented, quasi-finite morphism  $S' \rightarrow S$ , with  $S'$  affine, and an  $S$ -morphism  $S' \rightarrow X$ .

<sup>58</sup>Editor's note: every family  $\{S_{pq} \rightarrow S_p\} \in P'_i$  is finite. Hence  $S'_p = \coprod_q S_{pq}$  is affine, and the family may be replaced by the morphism  $S'_p \rightarrow S_p$ , which still belongs to  $P'_i$ .

- (v) *Every finite surjective family of flat quasi-compact morphisms is covering for (fpqc). In particular, every faithfully flat quasi-compact morphism is covering for (fpqc).*

*Proof.* Part (i) follows from 6.2.1 and part (ii) from 6.2.3, taking into account that a sheaf for the Zariski topology carries direct sums to products. Every representable functor is a sheaf for (Zar) and satisfies condition (ii)(b), by SGA 1, VIII, 5.3. Thus  $T_1$  is coarser than the canonical topology, which proves (iii).

We prove (iv). Let  $\{S_i \rightarrow S\}$  be a family as in the statement. By covering  $S$  with affine open subschemes, we reduce immediately to the case in which  $S$  is affine.<sup>59</sup>

First suppose that the morphisms  $S_i \rightarrow S$  are flat and open (respectively, étale). Cover each  $S_i$  by affine open subschemes  $S_{ij}$ . Since the morphisms are open, the images  $T_{ij}$  of the  $S_{ij}$  form an open covering of  $S$ . The affine scheme  $S$  is quasi-compact, so finitely many  $T_{ij}$ , indexed by a finite set  $E$ , cover it. Then

$$S' = \coprod_{(i,j) \in E} S_{ij}$$

is affine, and  $S' \rightarrow S$  belongs to  $P'_1$ , respectively  $P'_3$ , and is therefore covering. Since it factors through the given family, that family is covering.

In the (étf) case, every  $S_i$  is finite over  $S$  and hence affine. In the preceding argument one may take the singleton covering  $\{S_i\}$  of  $S_i$ , which gives  $S' \rightarrow S \in P'_4$ .

Now suppose that the morphisms  $f_i : S_i \rightarrow S$  are flat and locally of finite presentation. For every  $s \in S$ , the proof of EGA IV<sub>4</sub>, 17.16.2 supplies an affine subscheme  $X(s)$  of some  $S_i$ , with  $s \in f_i(X(s))$ , such that the restriction

$$g_i : X(s) \longrightarrow S$$

is flat, finitely presented, and quasi-finite. Its image  $g_i(X(s))$  is an open neighborhood  $U(s)$  of  $s$ , by EGA IV<sub>2</sub>, 2.4.6. Since  $S$  is affine, finitely many such opens  $U(s_j)$ ,  $j = 1, \dots, n$ , cover  $S$ . Consequently,

$$X' = \coprod_{j=1}^n X(s_j)$$

is affine and  $X' \rightarrow S$  is surjective, flat, finitely presented, and quasi-finite, so it belongs to  $P'_2$ .<sup>60</sup> This completes the proof of (iv).

We prove (v). Let  $\{S_i \rightarrow S\}$  be a finite surjective family of flat quasi-compact morphisms, and cover  $S$  by affine open subschemes  $T_j$ . The schemes

$$S_{ij} = T_j \times_S S_i$$

are quasi-compact, and therefore admit finite affine open coverings  $T_{ijk}$ . Each  $T_{ijk} \rightarrow T_j$  is flat, and the family  $\{T_{ijk} \rightarrow T_j\}$  is finite and surjective, hence covering for  $T_j$ . By composition,  $\{T_{ijk} \rightarrow S\}$  is covering as well. It factors through the given family, which is therefore covering:

$$\begin{array}{ccccc} S_i & \longleftarrow & S_{ij} & \longleftarrow & T_{ijk} \\ \downarrow & & \downarrow & \swarrow & \\ S & \longleftarrow & T_j & & \end{array} \quad (6.3.1.v)$$

<sup>59</sup>Editor's note: the argument that follows has been simplified by using the fact that  $S$  is now assumed affine.

<sup>60</sup>Editor's note: this shows that if  $P'_2$  denotes the set of finite surjective families of flat, finitely presented morphisms in  $\mathcal{C}'$ , then the topology generated by  $P$  and  $P'_2$  equals  $T_2$ . Moreover, in the notation used at the beginning of the proof of (iv), if each  $S_i$  is covered by affine opens of finite presentation over  $S$ , then the resulting morphism  $S' \rightarrow S$  belongs to  $P'_2$ .



**Corollary 6.3.2.** *Let  $(M_i)$  be the following families of morphisms:*

- $(M_1)$  : faithfully flat quasi-compact morphisms;
- $(M_2)$  : faithfully flat morphisms locally of finite presentation;
- $(M_3)$  : surjective étale morphisms;
- $(M_4)$  : surjective finite étale morphisms.

*The family  $(M_i)$  satisfies axioms (a), (b), (c),  $(d_{T_i})$ , and  $(e_{T_i})$  of 4.6.3.*<sup>61</sup>

Conditions (a), (b), and (c) are classical (see EGA and SGA 1, *passim*).<sup>62</sup> By 6.3.1(iv),(v),  $(M_i)$  satisfies  $(d_{T_i})$ . It remains to verify  $(e_{T_i})$ . It is enough to verify  $(e_{T_1})$ , which implies all the others; this follows from SGA 1, VIII, §§4–5.

**Corollary 6.3.3.** *Let  $X$  be a scheme and let  $R$  be an equivalence relation in  $X$  of type  $(M_i)$ . Then  $R$  is  $(M_i)$ -effective if and only if the sheaf quotient of  $X$  by  $R$  for  $T_i$  is representable; in that case it is represented by the quotient  $X/R$ .*

This is 4.6.5.

## 6.4. Effectivity Conditions

We now seek families  $(N)$  of morphisms satisfying axiom  $(f_T)$  of 4.7. Observe first that  $(f_{T_1})$  implies  $(f_{T_i})$ , so it is enough to consider the (fpqc) topology.

**Lemma 6.4.1.** *The following families of morphisms satisfy axiom  $(f_{T_1})$  of 4.7, that is, they "descend by (fpqc)":*

- $(N)$  : open immersions;
- $(N')$  : closed immersions;
- $(N'')$  : quasi-compact immersions.

By 6.3.1(ii), it is enough to verify that these families descend for the Zariski topology and along a faithfully flat quasi-compact morphism. The first assertion is clear. For  $(N)$ , the second is SGA 1, VIII, 4.4; for  $(N')$ , it is *loc. cit.*, 1.9; for  $(N'')$ , argue as in *loc. cit.*, 5.5, using the two preceding results.

**Corollary 6.4.2.** *The same result holds for quasi-compact open immersions.*

These results make the general results of 4.7.1, 4.7.2, 5.1.8, 5.3.1, and so on applicable to the present situation. We state the first as an example.

**Corollary 6.4.3.**  $(= 4.7.1 + 4.6.10)$ . *Let  $X$  be a scheme and  $R$  an equivalence relation in  $X$ . Suppose that  $R \rightarrow X$  is faithfully flat and quasi-compact, and that  $R \rightarrow X \times X$  is a closed immersion (respectively, open, quasi-compact, or quasi-compact open). Then the sheaf quotient  $X/R$  is the same for the (fpqc) topology as for the canonical topology. For every scheme  $S$ ,*

$$(X/R)(S) = \left\{ \begin{array}{l} \text{closed (respectively, open, retrocompact, or open retrocompact)} \\ \text{subschemes } Z \text{ of } X_S, \text{ stable under } R \times S, \text{ such that} \\ Z \rightarrow S \text{ is faithfully flat and quasi-compact and} \\ R_Z \rightrightarrows Z \rightarrow S \text{ is exact} \end{array} \right\}. \quad (6.4.3.1)$$

63

<sup>61</sup>Editor's note: the original has been corrected by adding surjectivity to  $(M_3)$  and  $(M_4)$ ; it is automatically satisfied in the other two cases.

<sup>62</sup>Editor's note: see EGA I, 6.6.4 for "quasi-compact"; EGA II, 6.1.5 for "finite"; EGA IV<sub>1</sub>, 1.6.2 for "locally of finite presentation"; EGA IV<sub>2</sub>, 2.2.13 for "faithfully flat"; and EGA IV<sub>4</sub>, 17.3.3 for "étale".

<sup>63</sup>Editor's note: recall that a subscheme  $Z$  of a scheme  $T$  is called *retrocompact* if the immersion  $Z \hookrightarrow T$  is quasi-compact; see EGA 0<sub>III</sub>, 9.1.1.

## 6.5. Principal Homogeneous Bundles


We merely record the terminology:

| topology | principal homogeneous bundles                     |         |
|----------|---------------------------------------------------|---------|
| (fpqc)   | principal homogeneous bundles (without qualifier) | (6.5.1) |
| (ét)     | quasi-isotrivial principal homogeneous bundles    |         |
| (étf)    | locally isotrivial principal homogeneous bundles  |         |
| (Zar)    | locally trivial principal homogeneous bundles.    |         |

## 6.6. Other Topologies

Other topologies on the category of schemes are sometimes useful. One is the *global finite étale topology* (étfg), generated by the pretopology whose covering families are the surjective families of finite étale morphisms. It is not finer than the Zariski topology. The corresponding principal homogeneous bundles are called *isotrivial*.

The position of this topology, and of the chaotic topology, in the lattice is as follows:



64

## 6.7. Homogeneous Spaces

Let<sup>65</sup>  $G$  be an  $S$ -group scheme, let  $X$  be an  $S$ -scheme equipped with a left action of  $G$ , and let

$$\Phi : G \times_S X \longrightarrow X \times_S X$$

be the morphism of  $S$ -schemes defined on points by

$$(g, x) \longmapsto (gx, x).$$

Recall (cf. 5.1.0 and III.0.1) that  $X$  is called a *formally principal homogeneous space* under  $G$  if the following equivalent conditions hold:

- (i) for every  $T \rightarrow S$ , the set  $X(T)$  is empty or is a principal homogeneous space under  $G(T)$ ;

<sup>64</sup>Editor's note: recall (cf. 4.4.2) that the chaotic topology is the coarsest topology, defined by  $J(S) = \{S\}$  for every  $S \in \text{Ob } \mathcal{C}$ .

<sup>65</sup>Editor's note: the numbering that follows has been added.

- (ii)  $\Phi$  is an isomorphism of  $S$ -functors;
- (iii)  $\Phi$  is an isomorphism of  $S$ -schemes.

The equivalence (i)  $\iff$  (ii) is clear, and (ii)  $\iff$  (iii) holds because  $\mathcal{C} = (\mathbf{Sch}/S)$  is a full subcategory of  $\widehat{\mathcal{C}}$ .

The notion of a *formally homogeneous space*, not necessarily principal homogeneous, is obtained by requiring  $\Phi$  to be an epimorphism in the category of sheaves for an appropriate topology  $\mathcal{T}$ . Indeed, requiring  $\Phi$  to be an epimorphism of  $S$ -functors is equivalent to requiring that, for every  $T \rightarrow S$ , the set  $X(T)$  be empty or homogeneous (not necessarily principal homogeneous) under  $G(T)$ . This is too restrictive, as the following simple example shows.

Let

$$S = \operatorname{Spec} R, \quad G = \mathbf{G}_{m,R}, \quad X = \mathbf{G}_{m,R},$$

and let  $G$  act on  $X$  by  $t \cdot x = t^2 x$ . Then  $\Phi$  is surjective, finite, and étale, and hence is an epimorphism in the category of sheaves for the (étf) topology (and, *a fortiori*, an epimorphism of  $S$ -schemes). On the other hand, the points 1 and  $-1$  of  $X(R)$  are not conjugate under any element of  $G(R)$ . Thus

$$G(R) \times X(R) \longrightarrow X(R) \times X(R)$$

is not surjective.<sup>66</sup> We are therefore led to the following definition.

**Definition 6.7.1.** *Let  $G$  be an  $S$ -group, let  $X$  be an  $S$ -scheme equipped with an action of  $G$ , and let  $\mathcal{T}$  be a topology on  $(\mathbf{Sch}/S)$  coarser than the canonical topology. The scheme  $X$  is called a *formally homogeneous space* under  $G$ , relative to  $\mathcal{T}$ , if the following equivalent conditions hold:*

- (i) *the morphism  $\Phi : G \times_S X \rightarrow X \times_S X$  is an epimorphism in the category of sheaves for  $\mathcal{T}$ ;*
- (ii) *for every  $T \rightarrow S$  and every  $x, y \in X(T)$ , there exist a  $\mathcal{T}$ -covering morphism  $T' \rightarrow T$  and an element  $g \in G(T')$  such that  $y_{T'} = g \cdot x_{T'}$ .*

**Remark 6.7.2.** *Condition (i) implies in particular that  $\Phi$  is a universal effective epimorphism in  $(\mathbf{Sch}/S)$ , by 4.4.3. As is readily seen, it follows that  $\Phi$  is surjective; cf. 1.3, Editor's note (3).*

**Proposition and Definition 6.7.3.** <sup>67</sup> *Let  $G$  be an  $S$ -group, let  $X$  be an  $S$ -scheme equipped with an action of  $G$ , and let  $\mathcal{T}$  be a topology on  $(\mathbf{Sch}/S)$  coarser than the canonical topology. The following conditions are equivalent:*

- (i)  *$X$  satisfies both of the following conditions:*
  - (1) *the morphism  $\Phi : G \times_S X \rightarrow X \times_S X$  is covering, that is,  $X$  is a formally homogeneous  $G$ -space;*

<sup>66</sup>Editor's note: this difficulty plainly comes from the following fact. Let  $\mathcal{C}'$  be a full subcategory of  $\widehat{\mathcal{C}}$  containing  $\mathcal{C}$ , for example the category  $\widehat{\mathcal{C}}_{\mathcal{T}}$  of sheaves on  $\mathcal{C}$  for a topology  $\mathcal{T}$  coarser than the canonical topology. If  $f : X \rightarrow Y$  is a morphism in  $\mathcal{C}$ , then the implications

$$f \text{ epimorphic in } \widehat{\mathcal{C}} \implies f \text{ epimorphic in } \mathcal{C}' \implies f \text{ epimorphic in } \mathcal{C}$$

are generally strict.

<sup>67</sup>Editor's note: cf. [Ray70], Definition VI.1.1.1.

- (2) the structural morphism  $X \rightarrow S$  is also covering, that is, locally for  $\mathcal{T}$  it has a section (cf. 4.4.8 bis, (b)).
- (ii) Locally on  $S$  for the topology  $\mathcal{T}$ , the scheme  $X$ , with its  $G$ -action, is isomorphic to the quotient sheaf of  $G$  by a subgroup scheme  $H$ , for  $\mathcal{T}$ . In other words, there is a covering family  $\{S_i \rightarrow S\}$  such that every  $X \times_S S_i$  represents the sheaf quotient of  $G \times_S S_i$  by some subgroup scheme  $H_i$ .

Under these conditions,  $X$  is called a  $G$ -homogeneous space, relative to  $\mathcal{T}$ .

*Proof.* Assume (ii). Put

$$G_i = G \times_S S_i, \quad X_i = X \times_S S_i.$$

The scheme  $X_i$  has a section over  $S_i$ , namely the composite of the unit section  $\varepsilon_i : S_i \rightarrow G_i$  with the projection  $\pi_i : G_i \rightarrow X_i = G_i/H_i$ . It follows that  $X \rightarrow S$  is covering.

Moreover,  $\pi_i$  is covering, so  $\pi_i \times \pi_i$  is covering as well, by 4.2.3 (C 1), (C 2). There is a commutative diagram<sup>68</sup>

$$\begin{array}{ccc} G_i \times_{S_i} X_i & \xrightarrow{\Phi_i} & X_i \times_{S_i} X_i \\ \text{id} \times \pi_i \uparrow & & \pi_i \times \pi_i \uparrow \\ G_i \times_{S_i} G_i & \xrightarrow[\sim]{\Psi_i} & G_i \times_{S_i} G_i, \end{array} \quad (6.7.3.1)$$

where  $\Phi_i$  is obtained from  $\Phi$  by the base change  $S_i \rightarrow S$ , and  $\Psi_i$  is the isomorphism defined on points by

$$(g, g') \mapsto (gg', g).$$

The composite  $(\pi_i \times \pi_i) \circ \Psi_i$  is covering; hence  $\Phi_i$  is covering by 4.2.3 (C 3). Thus  $\Phi$  is locally covering and is therefore covering, by 4.2.3 (C 5). This proves (ii)  $\implies$  (i).

Conversely, assume (i), and first suppose in addition that the structural morphism  $X \rightarrow S$  has a section  $\sigma$ . By EGA I, 5.3.13,  $\sigma$  is an immersion. Define  $H = G \times_X S$  by the following diagram, whose two squares are cartesian:

$$\begin{array}{ccccc} H & \longrightarrow & G & \xrightarrow{\text{id}_G \boxtimes \sigma} & G \times_S X \\ \downarrow & & \pi \downarrow & & \Phi \downarrow \\ S & \xrightarrow{\sigma} & X & \xrightarrow{\text{id}_X \boxtimes \sigma} & X \times_S X. \end{array} \quad (6.7.3.2)$$

Here  $\pi$ ,  $\text{id}_G \boxtimes \sigma$ , and  $\text{id}_X \boxtimes \sigma$  are the morphisms defined on points, for  $T \rightarrow S$ ,  $g \in G(T)$ , and  $x \in X(T)$ , by

$$\pi(g) = g \cdot \sigma_T, \quad (\text{id}_G \boxtimes \sigma)(g) = (g, \sigma_T), \quad (\text{id}_X \boxtimes \sigma)(x) = (x, \sigma_T).$$

Then  $\pi$  is covering, and  $H$  is a subgroup scheme of  $G$  representing the stabilizer  $\text{Stab}_G(\sigma)$  of  $\sigma$  (cf. I, 2.3.3). Thus, for every  $T \rightarrow S$ ,

$$H(T) = \{g \in G(T) \mid g \cdot \sigma_T = \sigma_T\}.$$

Let  $G/H$  denote the presheaf

$$T \mapsto G(T)/H(T),$$

<sup>68</sup>Editor's note: the Polo–Gille re-edition prints  $G_i \times_{S_i} X_i$  at the upper right. We write  $X_i \times_{S_i} X_i$ , since  $\Phi_i$ , as the base change of  $\Phi : G \times_S X \rightarrow X \times_S X$ , and  $\pi_i \times \pi_i$  both have this codomain.

and let  $a(G/H)$  be its associated sheaf for  $\mathcal{T}$ . The preceding construction gives a commutative diagram of presheaves equipped with a  $G$ -action:

$$\begin{array}{ccc} G & \xrightarrow{\pi} & X \\ \downarrow & \nearrow \bar{\pi} & \\ G/H & & \end{array} \quad (6.7.3.3)$$

where  $\bar{\pi}$  is a monomorphism. Since  $\pi$  is covering,  $\bar{\pi}$  is covering as well. By 4.3.12,  $\bar{\pi}$  therefore induces an isomorphism

$$a(G/H) \xrightarrow{\sim} X.$$

We have proved: if  $X$  is a  $G$ -homogeneous space such that  $X \rightarrow S$  has a section  $\sigma$ , then  $X$  represents the sheaf quotient  $G/H$ , where  $H = G \times_X S$  is the stabilizer of  $\sigma$ .

In general, by hypothesis there is a covering family  $\{S_i \rightarrow S\}$  such that every morphism

$$X_i = X \times_S S_i \longrightarrow S_i$$

has a section  $\sigma_i$ . Put  $G_i = G \times_S S_i$ . The morphism

$$\Phi_i : G_i \times_{S_i} X_i \longrightarrow X_i \times_{S_i} X_i$$

obtained from  $\Phi$  by base change along  $S_i \rightarrow S$  is still covering.

The preceding argument then gives

$$X_i \simeq G_i/H_i,$$

where  $H_i$  is the stabilizer of  $\sigma_i$  in  $G_i$ . This completes the proof of (i)  $\implies$  (ii).

## Bibliography

- [AS] J. Giraud, *Analysis Situs*, Séminaire Bourbaki, Exposé 256, May 1963.
- [D] J. Giraud, *Méthode de la descente*, Mémoires de la Société Mathématique de France, vol. 2 (1964), pp. iii–viii and 1–150.
- [MA] M. Artin, *Grothendieck Topologies*, mimeographed notes, Harvard, 1962.
- [SGA 1] *Séminaire de Géométrie Algébrique du Bois-Marie 1960–61: Revêtements étales et groupe fondamental*, Lecture Notes in Mathematics 224 (1971); recomposed and annotated edition, Documents Mathématiques 3, Société Mathématique de France, 2003.
- [SGA 4] *Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964: Théorie des topos et cohomologie étale des schémas*, vols. I–III, Lecture Notes in Mathematics 269, 270 (1972) and 305 (1973).
- [TDTE I] A. Grothendieck, *Techniques de descente et théorèmes d’existence en géométrie algébrique I. Généralités. Descente par morphismes fidèlement plats*, Séminaire Bourbaki, Exposé 190, December 1959.
- [Ray70] M. Raynaud, *Faisceaux amples sur les schémas en groupes et les espaces homogènes*, Lecture Notes in Mathematics 119, Springer-Verlag, 1970.<sup>69</sup>

---

<sup>69</sup>Editor’s note: this reference has been added.

# Exposé V

## Construction of Quotient Schemes

by P. Gabriel

The purpose of this exposé is to prove the theorems stated in TDTE III.<sup>1</sup> If  $X$  and  $T$  are two objects of a category  $\mathcal{C}$ , we write  $X(T)$  instead of  $\mathrm{Hom}_{\mathcal{C}}(T, X)$ . Similarly, if  $\varphi : Y \rightarrow X$  is an arrow (respectively, if  $T$  is an object) of  $\mathcal{C}$ , then  $\varphi(T)$  denotes the map  $g \mapsto \varphi \circ g$  from  $Y(T)$  to  $X(T)$ :

$$\begin{array}{ccc} T & & \\ \downarrow g & \searrow \varphi \circ g & \\ Y & \xrightarrow{\varphi} & X \end{array}$$

Figure 1: The map  $\varphi(T) : Y(T) \rightarrow X(T)$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille Exp5-13oct24.pdf, component-local p. 1; combined-reader p. 263; printed p. 251.*

Likewise,  $T(\varphi)$  denotes the map  $g \mapsto g \circ \varphi$  from  $T(X)$  to  $T(Y)$ :

$$\begin{array}{ccc} Y & \xrightarrow{\varphi} & X \\ & \searrow g \circ \varphi & \downarrow g \\ & & T \end{array}$$

Figure 2: The map  $T(\varphi) : T(X) \rightarrow T(Y)$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille Exp5-13oct24.pdf, component-local p. 1; combined-reader p. 263; printed p. 251.*

Finally, if  $P$  is a scheme, we denote by  $\underline{P}$  the underlying set of  $P$ .

As an exception, in the present exposé we do not follow the convention on quotient notation stated in IV 4.6.15 (*loc. cit.*, top of p. 227 of the original), because we wish to give here a construction of quotients that also applies to “pre-equivalence relations”<sup>2</sup> that are not equivalence relations.

<sup>1</sup>Editor’s note: namely, Theorems 5.1, 5.3, 6.1, 6.2, and 7.2 of TDTE III. The first two (respectively, the next two) correspond to Theorem 4.1 (respectively, Theorems 7.1 and 8.1) of this exposé. Theorem 7.2 of TDTE III is proved in Exposé VI<sub>A</sub>, 3.2 and 3.3.

<sup>2</sup>Editor’s note: that is, groupoids with base  $X$ ; see the terminology at the end of Section 1. When  $\mathcal{C}$  is the

## 1. $\mathcal{C}$ -groupoids

a) Let  $\mathcal{C}$  be a category in which products and fiber products exist. Recall first that a diagram

$$X_1 \begin{array}{c} \xrightarrow{d_1} \\ \xrightarrow{d_0} \end{array} X_0 \xrightarrow{p} Y$$

Figure 3: A parallel pair followed by its proposed cokernel.

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille Exp5-13oct24.pdf, component-local p. 2; combined-reader p. 264; printed p. 252.*

in  $\mathcal{C}$  is said to be *exact* if  $pd_0 = pd_1$  and if, for every  $T \in \mathcal{C}$ , the map  $T(p)$  is a bijection from  $T(Y)$  onto the subset of  $T(X_0)$  consisting of the arrows  $f : X_0 \rightarrow T$  such that  $fd_0 = fd_1$ . One also says that  $(Y, p)$  is the *cokernel* of  $(d_0, d_1)$ , and writes

$$(Y, p) = \text{Coker}(d_0, d_1).$$

b) For example, let  $\mathcal{C}$  be the category **(Esp.An)** of ringed spaces. In this case a cokernel  $(Y, p)$  always exists, and it may be described as follows. The underlying topological space of  $Y$  is obtained from  $X_0$  by identifying the points  $d_0(x)$  and  $d_1(x)$ , and by giving  $Y$  the quotient topology. The canonical map  $\pi : X_0 \rightarrow Y$ , together with  $d_0, d_1$ , then induces a pair of arrows of sheaves of rings on  $Y$ :

$$\pi_*(\mathcal{O}_0) \begin{array}{c} \xrightarrow{\delta_0} \\ \xrightarrow{\delta_1} \end{array} \pi_*(d_{0*}\mathcal{O}_1) = \pi_*(d_{1*}\mathcal{O}_1)$$

Figure 4: The pair of induced arrows between direct-image sheaves.

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille Exp5-13oct24.pdf, component-local p. 2; combined-reader p. 264; printed p. 252.*

Here  $\mathcal{O}_i$  is the structure sheaf of  $X_i$ . As the sheaf of rings on  $Y$ , one chooses the subsheaf of  $\pi_*(\mathcal{O}_0)$  whose sections  $s$  satisfy  $\delta_0(s) = \delta_1(s)$ . The arrow  $p$  is defined in the evident way. Consider<sup>3</sup> the following diagram in **(Esp.An)**:

$$X_1 \begin{array}{c} \xrightarrow{d_1} \\ \xrightarrow{d_0} \end{array} X_0$$

Figure 5: The pair  $(d_0, d_1) : X_1 \rightrightarrows X_0$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille Exp5-13oct24.pdf, component-local p. 2; combined-reader p. 264; printed p. 252.*

Let  $(Y, p)$  be its cokernel. An open subset  $U$  of  $X_0$  is said to be *saturated* if

$$d_0^{-1}(U) = d_1^{-1}(U),$$

which is equivalent to saying  $U = p^{-1}(p(U))$ . In this case, since  $Y$  has the quotient topology,  $p(U)$  is an open subset of  $Y$ .

---

category of schemes, the quotient  $p : X \rightarrow Y$  of a groupoid  $X_*$  with base  $X$  exists under certain hypotheses (see 4.1, 6.1, and 7.1). If, in addition,  $X_*$  is an equivalence relation, then under the same hypotheses  $p$  is faithfully flat and quasi-compact, hence a universal epimorphism; see *loc. cit.*.

<sup>3</sup>Editor's note: Lemmas 1.1 and 1.2 have been added; they are used several times in Sections 5–9.

**Lemma 1.1.** *Let  $U$  be a saturated open subset of  $X$ , and set  $V = p(U)$ . Denote by  $U_1 = d_0^{-1}(U) = d_1^{-1}(U)$  the corresponding open subset of  $X_1$ , and by  $\tilde{d}_0, \tilde{d}_1$ , and  $\tilde{p}$  the restrictions of  $d_0, d_1$  to  $U_1$ , and of  $p$  to  $U$ , respectively. Then  $(V, \tilde{p})$  is a cokernel in **(Esp.An)** of:<sup>4</sup>*

$$U_1 \begin{array}{c} \xrightarrow{\tilde{d}_1} \\ \xrightarrow{\tilde{d}_0} \end{array} U \xrightarrow{\tilde{p}} V$$

Figure 6: The restricted pair and its cokernel in Lemma 1.1.

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille Exp5-13oct24.pdf, component-local p. 2; combined-reader p. 264; printed p. 252.*

The verification is easy.

**Lemma 1.2.**— *Let  $d_0, d_1 : X_1 \rightrightarrows X_0$  be a diagram in **(Sch)**, and let  $(Y, p)$  be its cokernel in **(Esp.An)**.*

- (i) *If  $Y$  is a scheme and  $p$  is a morphism of schemes, then  $(Y, p)$  is a cokernel of  $(d_0, d_1)$  in **(Sch)**.*
- (ii) *Suppose that every point of  $X_0$  has a saturated open neighborhood  $U$  with the following property. Let  $\tilde{d}_0$  and  $\tilde{d}_1$  be the restrictions of  $d_0$  and  $d_1$  to  $d_0^{-1}(U) = d_1^{-1}(U)$ , and let  $(Q, q)$  be the cokernel of  $(\tilde{d}_0, \tilde{d}_1)$  in **(Esp.An)**. If  $Q$  is a scheme and  $q$  is a morphism of schemes, then  $(Y, p)$  is a cokernel of  $(d_0, d_1)$  in **(Sch)**.*

Part (i) is proved in §4(c); since the proof is short, let us repeat it here. Let  $f : X_0 \rightarrow Z$  be a morphism of schemes such that  $fd_0 = fd_1$ . By hypothesis there is a unique morphism of ringed spaces  $r : Y \rightarrow Z$  such that  $f = rp$ . It remains to show that, for every  $y \in Y$ , the homomorphism  $\mathcal{O}_{r(y)} \rightarrow \mathcal{O}_y$  induced by  $r$  is local. This follows because  $p$  is surjective, so  $y = p(x)$  for some  $x$ , and because the homomorphism  $\mathcal{O}_{f(x)} \rightarrow \mathcal{O}_x$  induced by  $f$  is local.

Part (ii) follows from part (i) and the preceding lemma.

**c)** In this exposé we study the existence of  $\text{Coker}(d_0, d_1)$  when the double arrow  $(d_0, d_1)$  is placed in a richer context. More precisely, let

$$X_2 = X_1 \times_{d_1, d_0} X_1$$

be the fiber product of the diagram

$$\begin{array}{ccc} & X_1 & \\ & \downarrow d_1 & \\ X_1 & \xrightarrow{d_0} & X_0 \end{array} \quad (*)$$

<sup>4</sup>Editor's note: this is not the case in the category of schemes. For example, let  $S = \text{Spec}(\mathbf{C})$ , let  $X_0 = \mathbf{A}_S^2 = \text{Spec}(\mathbf{C}[x_1, x_2])$ , let  $d_1 : \mathbf{G}_{m,S} \times_S \mathbf{A}_S^2 \rightarrow \mathbf{A}_S^2$  be the action of  $\mathbf{G}_{m,S}$  on  $\mathbf{A}_S^2$  by homotheties, let  $d_0$  be the projection onto the second factor, and let  $U = \mathbf{A}_S^2 \setminus \{m\}$ , where  $m = (0, 0)$ . Then the projective line  $\mathbf{P}_S^1$  is the cokernel of  $(\tilde{d}_0, \tilde{d}_1)$  in both **(Esp.An)** and **(Sch)**, whereas the cokernel  $Y$  of  $(d_0, d_1)$  in **(Esp.An)** is the union of  $\mathbf{P}_S^1$  and the point  $y_0 = \{p(m)\}$ . The only open subset containing  $y_0$  is  $Y$ , and  $\Gamma(Y, \mathcal{O}_Y) = \mathbf{C}$ . If  $f : \mathbf{A}_S^2 \rightarrow T$  is a morphism of  $S$ -schemes such that  $fd_0 = fd_1$ , and if  $V = \text{Spec}(A)$  is an affine open subset of  $T$  containing  $t_0 = f(y_0)$ , then  $f^{-1}(V) = \mathbf{A}^2$ , and the ring homomorphism  $A \rightarrow \mathbf{C}[x_1, x_2]$  factors through  $\mathbf{C}$ . This shows that  $S = \text{Spec}(\mathbf{C})$  is the cokernel of  $(d_0, d_1)$  in the category **(Sch/S)**.



and let  $d'_0, d'_2 : X_2 \rightarrow X_1$  be the two canonical projections. Thus, by definition, the square

$$\begin{array}{ccc} X_2 & \xrightarrow{d'_0} & X_1 \\ d'_2 \downarrow & & \downarrow d_1 \\ X_1 & \xrightarrow{d_0} & X_0 \end{array} \quad (0)$$

is Cartesian.

In addition, suppose that a third arrow  $d'_1 : X_2 \rightarrow X_1$  is given. We say that

$$(d_0, d_1 : X_1 \rightrightarrows X_0, d'_1)$$

is a  $\mathcal{C}$ -groupoid if, for every object  $T$  of  $\mathcal{C}$ ,  $X_1(T)$  is the set of arrows of a groupoid  $X_*(T)$  whose set of objects is  $X_0(T)$ , whose source map is  $d_1(T)$ , whose target map is  $d_0(T)$ , and whose composition map is  $d'_1(T)$ . As usual, we identify

$$(X_1 \times_{d_1, d_0} X_1)(T) \simeq X_1(T) \times_{d_1(T), d_0(T)} X_1(T),$$

and recall that a groupoid is a category in which every arrow is invertible.<sup>5</sup>

If  $\varphi$  is an arrow of the groupoid  $X_*(T)$ , the map  $f \mapsto \varphi \circ f$  is a bijection from the set of arrows  $f$  whose target equals the source of  $\varphi$  onto the set of arrows having the same target as  $\varphi$ . This is equivalently expressed by saying that the square

$$\begin{array}{ccc} X_2 & \xrightarrow{d'_1} & X_1 \\ d'_0 \downarrow & & \downarrow d_0 \\ X_1 & \xrightarrow{d_0} & X_0 \end{array} \quad (1)$$

is Cartesian.

Similarly,  $g \mapsto g \circ \varphi$  is a bijection from the set of arrows  $g$  of  $X_*(T)$  whose source equals the target of  $\varphi$  onto the set of arrows having the same source as  $\varphi$ . Equivalently, the square

$$\begin{array}{ccc} X_2 & \xrightarrow{d'_1} & X_1 \\ d'_2 \downarrow & & \downarrow d_1 \\ X_1 & \xrightarrow{d_1} & X_0 \end{array} \quad (2)$$

is Cartesian.

On the other hand, let  $s : X_0 \rightarrow X_1$  be the unique arrow of  $\mathcal{C}$  such that, for every  $T$ , the map  $s(T) : X_0(T) \rightarrow X_1(T)$  sends each object of  $X_*(T)$  to its identity arrow.<sup>6</sup> The arrow  $s$  satisfies

$$d_1 s = \text{id}_{X_0}, \quad (3)$$

$$d_0 s = \text{id}_{X_0}. \quad (3 \text{ bis})$$

<sup>5</sup>Editor's note: in this case  $X_2(T)$  is the set of pairs  $(f_2, f_1)$  of composable arrows, that is, pairs such that  $d_0(f_1) = d_1(f_2)$ ; the maps  $d'_0, d'_1, d'_2$  send  $(f_2, f_1)$ , respectively, to  $f_2, f_2 \circ f_1, f_1$ .

<sup>6</sup>Editor's note:  $T \mapsto s(T)$  defines an element of  $\text{Hom}(h_{X_0}, h_{X_1})$ , and the latter is equal to  $\text{Hom}(X_0, X_1)$  by the Yoneda lemma.

Finally, associativity of the composition maps  $d'_1(T)$  is equivalent to commutativity of the diagram

$$\begin{array}{ccc}
 X_1 \times_{d_1, d_0} X_1 \times_{d_1, d_0} X_1 & \xrightarrow{d'_1 \times X_1} & X_1 \times_{d_1, d_0} X_1 \\
 \downarrow X_1 \times d'_1 & & \downarrow d'_1 \\
 X_1 \times_{d_1, d_0} X_1 & \xrightarrow{d'_1} & X_1
 \end{array} \quad (4)$$

Conversely, conditions (1), (2), and (4), together with the existence of an arrow  $s$  satisfying (3), imply that  $(X_1 \rightrightarrows X_0, d'_1)$  is a  $\mathcal{C}$ -groupoid. Condition (3) is harmless: it merely ensures that  $d_1(T) : X_1(T) \rightarrow X_0(T)$  is surjective for every  $T \in \mathcal{C}$ . In the sequel we shall mainly use the Cartesian squares (0), (1), and (2), summarized in the diagram

$$\begin{array}{ccccc}
 X_2 & \xrightarrow{d'_1} & X_1 & \xrightarrow{d_0} & X_0 \\
 d'_2 \downarrow & & d'_0 & & \downarrow d_1 \\
 X_1 & \xrightarrow{d_1} & X_0 & & \\
 & \xrightarrow{d_0} & & & 
 \end{array} \quad (0,1,2)$$

The two left-hand squares in this diagram (that is, squares (0) and (2)) are Cartesian; the first row is exact; and  $X_2$  is identified with the fiber product  $X_1 \times_{d_0, d_0} X_1$ .

We use associativity only indirectly, for example to ensure the existence of an arrow  $s$  satisfying (3) and (3 bis), or to ensure the existence of an arrow

$$\sigma : X_1 \longrightarrow X_1, \quad d_0 \sigma = d_1, \quad d_1 \sigma = d_0. \quad (\dagger)$$

Here  $\sigma$  is chosen so that  $\sigma(T) : X_1(T) \rightarrow X_1(T)$  sends every arrow of  $X_*(T)$  to its inverse.<sup>7</sup>

By an abuse of language, we shall sometimes call a diagram

$$X_2 \xrightarrow{d'_0, d'_1, d'_2} X_1 \xrightarrow{d_0, d_1} X_0$$

a  $\mathcal{C}$ -groupoid when (0), (1), and (2) are Cartesian, (4) is commutative, and there exists an  $s$  satisfying (3). Thus  $X_2$  may be “a” fiber product of  $(*)$  without being “the” fiber product of  $(*)$ .<sup>8</sup>

*Terminology.* In place of the  $\mathcal{C}$ -groupoid  $X_*$ , we shall also speak of the *groupoid*  $X_*$  with base  $X_0$ , or of the *pre-equivalence relation*  $X_*$  in  $X_0$ .

## 2. Examples of $\mathcal{C}$ -groupoids

**a)** Let  $X$  be an object of  $\mathcal{C}$ , and let  $G$  be a  $\mathcal{C}$ -group acting on the left on  $X$ . We denote by  $d_0 : G \times X \rightarrow X$  the arrow defining the action of  $G$  on  $X$ , by  $d_1 : G \times X \rightarrow X$  the projection onto the second factor, and by  $\mu : G \times G \rightarrow G$  the arrow defining

the  $\mathcal{C}$ -group structure on  $G$ , and finally by  $\text{pr}_{2,3}$  the projection of  $G \times G \times X = G \times (G \times X)$  onto the second factor. Then

<sup>7</sup>Editor's note: the Yoneda lemma shows that  $\sigma$  is an involutive automorphism of  $X_1$ ; this will be used, for example, in 3(e) and in Theorem 4.1.

<sup>8</sup>Editor's note: see Example 2(a) below.

$$G \times G \times X \begin{array}{c} \xrightarrow{\text{pr}_{2,3}} \\ \xrightarrow[\mu \times X]{G \times d_0} \end{array} G \times X \xrightarrow[\begin{array}{c} d_1 \\ d_0 \end{array}]{\quad} X$$

Figure 7: The groupoid attached to the action of the  $\mathcal{C}$ -group  $G$  on  $X$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille Exp5-13oct24.pdf, component-local p. 6; combined-reader p. 268; page header p. 256.*

is a  $\mathcal{C}$ -groupoid.

**b)** Let  $d_0, d_1 : X_1 \rightarrow X_0$  be an *equivalence pair*. In other words, if  $d_0 \boxtimes d_1 : X_1 \rightarrow X_0 \times X_0$  is the arrow with components  $d_0$  and  $d_1$ , we suppose that, for every object  $T$  of  $\mathcal{C}$ , the map  $(d_0 \boxtimes d_1)(T)$  is a bijection from  $X_1(T)$  onto the graph of an equivalence relation on  $X_0(T)$ . Consequently,  $X_1(T)$  is identified with the set of pairs  $(x, y)$  of elements of  $X_0(T)$  such that  $x \sim y$ . Likewise,

$$X_2(T) = (X_1 \times_{d_1, d_0} X_1)(T)$$

is identified with the set of triples  $(x, y, z)$  of elements of  $X_0(T)$  such that  $x \sim y$  and  $y \sim z$ . There is therefore a unique arrow  $d'_1 : X_2 \rightarrow X_1$  making squares (1) and (2) commute:  $d'_1(T)$  must send  $(x, y, z) \in X_2(T)$  to  $(x, z) \in X_1(T)$ . With this choice of  $d'_1$ ,

$$(d_0, d_1 : X_1 \rightrightarrows X_0, d'_1)$$

is a  $\mathcal{C}$ -groupoid.

Conversely, consider a  $\mathcal{C}$ -groupoid  $X_*$  for which  $d_0 \boxtimes d_1 : X_1 \rightarrow X_0 \times X_0$  is a monomorphism. Then  $(d_0, d_1)$  is an equivalence pair, and  $X_*$  can be reconstructed from  $(d_0, d_1)$  as explained a few lines above.<sup>9</sup>

**c)** If  $p : X \rightarrow Y$  is any arrow of  $\mathcal{C}$ , and if  $\text{pr}_1$  and  $\text{pr}_2$  are the two projections of  $X \times_{p,p} X$  onto  $X$ , then

$$(\text{pr}_1, \text{pr}_2) : X \times_{p,p} X \rightrightarrows X$$

is an equivalence pair. The arrow  $p$  is called an *effective epimorphism* if the diagram

$$X \times_{p,p} X \begin{array}{c} \xrightarrow{\text{pr}_1} \\ \xrightarrow[\text{pr}_2]{\quad} \end{array} X \xrightarrow{p} Y$$

Figure 8: The defining diagram for an effective epimorphism.

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille Exp5-13oct24.pdf, component-local p. 6; combined-reader p. 268; page header p. 256.*

is exact, that is, if

$$(Y, p) = \text{Coker}(\text{pr}_1, \text{pr}_2).$$

For example, let  $S$  be a Noetherian scheme and let  $\mathcal{C}$  be the category of schemes finite over  $S$ . We show that an epimorphism in  $\mathcal{C}$  need not be effective. Take

$$S = \text{Spec } k[T^3, T^5], \quad Y = S, \quad X = \text{Spec } k[T],$$

where  $k$  is a commutative field. If  $i$  is the inclusion of  $B = k[T^3, T^5]$  into  $A = k[T]$ , take  $p = \text{Spec } i$ . Then  $X \times_{p,p} X$  is identified with  $\text{Spec}(A \otimes_B A)$ , and  $\text{Coker}(\text{pr}_1, \text{pr}_2)$  with  $\text{Spec } B'$ , where  $B'$  is the subring of  $A$  consisting of the elements  $a$  such that  $a \otimes_B 1 = 1 \otimes_B a$ . But

$$T^7 \otimes_B 1 = (T^2 T^5) \otimes_B 1 = T^2 \otimes_B T^5 = T^2 \otimes_B (T^3 T^2) = T^5 \otimes_B T^2 = 1 \otimes_B T^7.$$

<sup>9</sup>Editor's note: in particular, if  $G$  is a  $\mathcal{C}$ -group acting on the left on an object  $X$  of  $\mathcal{C}$ , and if  $X_*$  is the  $\mathcal{C}$ -groupoid defined in 2(a), then  $(d_0, d_1)$  is an equivalence pair if and only if  $G$  acts freely on  $X$ ; cf. Exposé III, 3.2.1.

Thus  $T^7$  belongs to  $B'$ , does not belong to  $B$ , and  $\text{Spec } B'$  is different from  $\text{Spec } B$ , which gives the counterexample.<sup>10</sup>

### 3. Some sorites concerning $\mathcal{C}$ -groupoids

Here are, in no particular order, some remarks used below.

a) Let

$$X_2 \begin{array}{c} \xrightarrow{d'_0, d'_1, d'_2} \\ \xrightarrow{\quad\quad\quad} \\ \xrightarrow{\quad\quad\quad} \end{array} X_1 \xrightarrow{d_0, d_1} X_0$$

Figure 9: The  $\mathcal{C}$ -groupoid  $X_*$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille  
Exp5-13oct24.pdf, component-local p. 7; combined-reader p. 269; page header p. 257.*

be a  $\mathcal{C}$ -groupoid, and let  $f_0 : Y_0 \rightarrow X_0$  be an arrow of  $\mathcal{C}$ . We shall define a  $\mathcal{C}$ -groupoid with base  $Y_0$ ,

$$Y_2 \begin{array}{c} \xrightarrow{e'_0, e'_1, e'_2} \\ \xrightarrow{\quad\quad\quad} \\ \xrightarrow{\quad\quad\quad} \end{array} Y_1 \xrightarrow{e_0, e_1} Y_0$$

Figure 10: The induced  $\mathcal{C}$ -groupoid  $Y_*$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille  
Exp5-13oct24.pdf, component-local p. 7; combined-reader p. 269; page header p. 257.*

which will be said to be *induced by  $X_*$  and  $f_0$* . We also say that  $Y_*$  is the *inverse image of  $X_*$  under the base change  $f_0$* .

We choose  $Y_1$  to be the fiber product of the diagram

$$\begin{array}{ccc} Y_1 & \xrightarrow{\quad f_1 \quad} & X_1 \\ \downarrow & & \downarrow d_0 \boxtimes d_1 \\ Y_0 \times Y_0 & \xrightarrow{\quad f_0 \times f_0 \quad} & X_0 \times X_0 \end{array}$$

Figure 11: The fiber-product definition of  $Y_1$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille  
Exp5-13oct24.pdf, component-local p. 7; combined-reader p. 269; page header p. 257.*

and take  $e_0$  and  $e_1$  to be the composites of the canonical arrow  $Y_1 \rightarrow Y_0 \times Y_0$  with the first and second projections of  $Y_0 \times Y_0$ , respectively. The morphism  $Y_1 \rightarrow Y_0 \times Y_0$  is then  $e_0 \boxtimes e_1$ , and

$$f_0 \circ e_i = d_i \circ f_1 \quad (i = 0, 1),$$

where  $f_1$  denotes the projection of  $Y_1$  onto  $X_1$ .

Set  $Y_2 = Y_1 \times_{e_0, e_1} Y_1$ , cf. 1(c). One may say that the pair  $(e_0, e_1)$  is defined so that, for every  $T \in \mathcal{C}$  and every pair  $(y, x)$  of elements of  $Y_0(T)$ , there is a certain bijective correspondence

$$\psi \longmapsto {}_y\psi_x$$

<sup>10</sup>Editor's note: the same argument applies to  $B = k[T^3, T^4]$  and  $T^5 \otimes_B 1$ ; more generally, it applies to  $B = k[T^n, T^{n+r}]$  and the element  $T^{n+2r} \otimes_B 1$ , provided that  $n$  does not divide  $2r$ .

between the arrows  $\psi$  of  $X_*(T)$  with source  $f_0(x)$  and target  $f_0(y)$ , and the arrows  ${}_y\psi_x$  of  $Y_*(T)$  with source  $x$  and target  $y$ . Thus  $e'_1 : Y_2 \rightarrow Y_1$  is determined by defining, for every  $T \in \mathcal{C}$ , the composition of arrows in  $Y_*(T)$  by the formula

$${}_z\psi_y \circ {}_y\varphi_x = {}_z(\psi \circ \varphi)_x.$$

It is clear that this definition makes each  $Y_*(T)$  a groupoid.

**b)** Given the  $\mathcal{C}$ -groupoid  $X_*$  and the base change  $f_0 : Y_0 \rightarrow X_0$ , one can reconstruct the pair  $(e_0, e_1) : Y_1 \rightrightarrows Y_0$  in another way.<sup>11</sup> Construct  $Y_0 \times_{X_0} X_1$ ,  $\text{pr}_1$ , and  $\text{pr}_2$  so that the square

$$\begin{array}{ccc} Y_0 \times_{X_0} X_1 & \xrightarrow{\text{pr}_2} & X_1 \\ \text{pr}_1 \downarrow & & \downarrow d_0 \\ Y_0 & \xrightarrow{f_0} & X_0 \end{array}$$

Figure 12: The Cartesian square defining  $Y_0 \times_{X_0} X_1$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille Exp5-13oct24.pdf, component-local p. 7; combined-reader p. 269; page header p. 257.*

is Cartesian. A straightforward reduction to the set-theoretic case then shows that the following square is Cartesian:

$$\begin{array}{ccc} Y_1 & \xrightarrow{e_0 \boxtimes f_1} & Y_0 \times_{X_0} X_1 \\ e_1 \downarrow & & \downarrow d_1 \circ \text{pr}_2 \\ Y_0 & \xrightarrow{f_0} & X_0 \end{array}$$

Figure 13: The second Cartesian square in the reconstruction of  $(e_0, e_1)$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille Exp5-13oct24.pdf, component-local p. 8; combined-reader p. 270; page header p. 258.*

Here  $f_1$  denotes the canonical projection of

$$Y_1 = (Y_0 \times Y_0) \times_{(X_0 \times X_0)} X_1$$

onto  $X_1$ .

**c)** We give two examples of inverse images of a  $\mathcal{C}$ -groupoid. Take  $Y_0 = X_1$  and  $f_0 = d_0$ . For every object  $T$  of  $\mathcal{C}$ , the set  $Y_1(T)$  is then identified with the set of diagrams of the form

$$\begin{array}{ccc} b & \xrightarrow{\varphi} & d \\ f \uparrow & & \uparrow g \\ a & & c \end{array}$$

Figure 14: An arrow of the first inverse-image groupoid.

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille Exp5-13oct24.pdf, component-local p. 8; combined-reader p. 270; page header p. 258.*

<sup>11</sup>Editor’s note: this second viewpoint is used in 3(f) and in the proof of 6.1.

in  $X_*(T)$ . The source of such a diagram is the arrow  $f$ , and its target is the arrow  $g$ . These diagrams compose in the evident way.

Now set  $Y'_0 = X_1$  and  $f'_0 = d_1$  (we add the primes<sup>12</sup> to avoid any confusion with the preceding example). In this case, for every  $T \in \mathcal{C}$ , the set  $Y'_1(T)$  is identified with the set of diagrams of the form

$$\begin{array}{ccc} b & & d \\ \uparrow f & & \uparrow g \\ a & \xrightarrow{\psi} & c \end{array}$$

Figure 15: An arrow of the second inverse-image groupoid.

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille Exp5-13oct24.pdf, component-local p. 8; combined-reader p. 270; page header p. 258.*

in the groupoid  $X_*(T)$ . The source of such a diagram is  $f$ , its target is  $g$ , and these diagrams again compose in the evident way.

It is now clear that the identity map of  $Y_0(T)$ , together with the map

$$\begin{array}{ccc} b & \xrightarrow{\varphi} & d \\ \uparrow f & & \uparrow g \\ a & & c \end{array} \quad \mapsto \quad \begin{array}{ccc} b & & d \\ \uparrow f & & \uparrow g \\ a & \xrightarrow{g^{-1} \circ \varphi \circ f} & c \end{array}$$

Figure 16: The map from  $Y_1(T)$  to  $Y'_1(T)$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille Exp5-13oct24.pdf, component-local p. 8; combined-reader p. 270; page header p. 258.*

defines an isomorphism from the groupoid  $Y_*(T)$  onto  $Y'_*(T)$ . Moreover, this isomorphism depends functorially on  $T$ , so the  $\mathcal{C}$ -groupoids  $Y_*$  and  $Y'_*$  are isomorphic.<sup>13</sup>

d)

**Proposition 3.1.**— *Keep the notation of 3(a), and suppose that  $f_0$  is an effective and universal epimorphism. Then  $\text{Coker}(d_0, d_1)$  exists if and only if  $\text{Coker}(e_0, e_1)$  exists.<sup>14</sup> Moreover, in this case  $f_0$  induces an isomorphism*

$$\text{Coker}(d_0, d_1) \xrightarrow{\sim} \text{Coker}(e_0, e_1).$$

Recall first that an epimorphism  $f_0 : Y_0 \rightarrow X_0$  is called *universal* if, in every Cartesian square

$$\begin{array}{ccc} Y' & \longrightarrow & Y_0 \\ f' \downarrow & & \downarrow f_0 \\ X' & \longrightarrow & X_0 \end{array}$$

Figure 17: A base change of the universal epimorphism  $f_0$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille Exp5-13oct24.pdf, component-local p. 9; combined-reader p. 271; page header p. 259.*

<sup>12</sup>Editor’s note: “accents” in the original.

<sup>13</sup>Editor’s note: this plays a crucial role in the proof of Lemma 6.1.

<sup>14</sup>Editor’s note: the original has been modified to display explicitly the isomorphism below.

the arrow  $f'$  is an epimorphism. Now let  $C(d_0, d_1)$  denote the covariant functor from  $\mathcal{C}$  to sets which assigns to every  $T \in \mathcal{C}$  the kernel of the pair

$$T(d_0), T(d_1) : T(X_0) \rightrightarrows T(X_1),$$

and define  $C(e_0, e_1)$  in the same way. For every  $T \in \mathcal{C}$ , there is therefore a commutative diagram

$$\begin{array}{ccccc} C(d_0, d_1)(T) & \longrightarrow & T(X_0) & \xrightleftharpoons[T(d_0)]{T(d_1)} & T(X_1) \\ T(f) \downarrow & & T(f_0) \downarrow & & T(f_1) \downarrow \\ C(e_0, e_1)(T) & \longrightarrow & T(Y_0) & \xrightleftharpoons[T(e_0)]{T(e_1)} & T(Y_1) \end{array}$$

Figure 18: The comparison of the functors  $C(d_0, d_1)$  and  $C(e_0, e_1)$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille Exp5-13oct24.pdf, component-local p. 9; combined-reader p. 271; page header p. 259.*

Here  $T(f)$  is the injection induced by the injection  $T(f_0)$ . If we show that  $T(f)$  is surjective for every  $T$ , we obtain a functorial isomorphism

$$f : C(d_0, d_1) \xrightarrow{\sim} C(e_0, e_1).$$

Thus one of these functors is representable if and only if the other is, which proves the proposition.

To prove that  $T(f)$  is surjective, consider the diagram

$$\begin{array}{ccccc} & & Y_1 & \xrightarrow{f_1} & X_1 \\ & \nearrow \Delta & \downarrow e_0 & \downarrow e_1 & \downarrow d_0 \downarrow d_1 \\ Y_0 \times_{X_0} Y_0 & \xrightleftharpoons[\text{pr}_1]{\text{pr}_2} & Y_0 & \xrightarrow{f_0} & X_0 \end{array}$$

Figure 19: The diagram used to prove the surjectivity of  $T(f)$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille Exp5-13oct24.pdf, component-local p. 9; combined-reader p. 271; page header p. 259.*

Here  $\Delta$  is the section of  $Y_1 \rightarrow Y_0 \times Y_0$  defined by the morphism

$$s \circ f_0 \circ \text{pr}_1 : Y_0 \times Y_0 \longrightarrow X_1,$$

where  $s : X_0 \rightarrow X_1$  satisfies (3) and (3 bis) of Section 1.

If  $g : Y_0 \rightarrow T$  satisfies  $g \circ e_0 = g \circ e_1$ , then

$$g \circ e_0 \circ \Delta = g \circ e_1 \circ \Delta,$$

so  $g \circ \text{pr}_1 = g \circ \text{pr}_2$ . Since  $f_0$  is an effective epimorphism,  $g$  factors as  $h \circ f_0$  for an arrow  $h : X_0 \rightarrow T$ ; in other words,  $g = T(f_0)(h)$ . It remains to prove that  $h \in C(d_0, d_1)(T)$ , or equivalently that  $hd_0 = hd_1$ . But

$$hd_0f_1 = hf_0e_0 = ge_0 = ge_1 = hf_0e_1 = hd_1f_1,$$

and the desired equality follows because  $f_1$  is an epimorphism (since  $f_0$  is a universal epimorphism).

e) Now let  $S$  be a scheme and take  $\mathcal{C} = (\mathbf{Sch}/S)$ . The data of a  $\mathcal{C}$ -groupoid

$$X_2 \begin{array}{c} \xrightarrow{d'_2} \\ \xrightarrow{d'_1} \\ \xrightarrow{d'_0} \end{array} X_1 \begin{array}{c} \xrightarrow{d_1} \\ \xrightarrow{d_0} \end{array} X_0$$

Figure 20: A  $(\mathbf{Sch}/S)$ -groupoid.

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille Exp5-13oct24.pdf, component-local p. 10; combined-reader p. 272; page header p. 260.*

defines an equivalence relation on the set  $\underline{X_0}$  underlying the scheme  $X_0$ : for  $x, y \in \underline{X_0}$ , write  $x \sim y$  if there exists  $z \in \underline{X_1}$  such that  $x = d_1 z$  and  $y = d_0 z$ . Reflexivity and symmetry are clear.<sup>15</sup> Let us prove transitivity. If  $x \sim y$  and  $y \sim z$ , there exist  $u, v \in \underline{X_1}$  such that

$$x = d_1 u, \quad y = d_0 u, \quad y = d_1 v, \quad z = d_0 v.$$

It follows that  $(v, u)$  belongs to the set-theoretic fiber product  $\underline{X_1} \times_{\underline{d_1}, \underline{d_0}} \underline{X_1}$ . Since the canonical map

$$\underline{X_1} \times_{\underline{d_1}, \underline{d_0}} \underline{X_1} \longrightarrow \underline{X_1} \times_{\underline{d_1}, \underline{d_0}} \underline{X_1}$$

Figure 21: The canonical map from the underlying set of the fiber product to the fiber product of the underlying sets.

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille Exp5-13oct24.pdf, component-local p. 10; combined-reader p. 272; page header p. 260.*

is surjective,  $(v, u)$  is the image of some  $w \in \underline{X_2}$ . Hence

$$x = d_1 d'_1 w, \quad z = d_0 d'_1 w,$$

and therefore  $x \sim z$ .

f) Keep the notation of 3(a) and 3(b), still taking  $\mathcal{C} = (\mathbf{Sch}/S)$ . If  $x, y$  are points of  $Y_0$ , then

$$x \sim y \iff f_0(x) \sim f_0(y).$$

Thus the inverse image of the equivalence relation defined by a groupoid is the equivalence relation defined by the inverse image of that groupoid.

Indeed, suppose that  $x \sim y$ . There is a  $z \in \underline{Y_1}$  such that  $x = e_1(z)$  and  $y = e_0(z)$ . Since  $f_0 \circ e_i = d_i \circ f_1$  for  $i = 0, 1$ , it follows that

$$f_0(x) = d_1 f_1(z), \quad f_0(y) = d_0 f_1(z),$$

and hence  $f_0(x) \sim f_0(y)$ .

Conversely, suppose that  $f_0(x) \sim f_0(y)$ , and let  $z \in \underline{X_1}$  satisfy

$$f_0(y) = d_1(z), \quad f_0(x) = d_0(z).$$

<sup>15</sup>Editor's note: reflexivity follows from the existence of  $s : X_0 \rightarrow X_1$ , which is a section of both  $d_0$  and  $d_1$ ; symmetry follows from the existence of the involution  $\sigma$  of  $X_1$  which “exchanges  $d_0$  and  $d_1$ ”, that is, which satisfies  $d_0 \sigma = d_1$  and  $d_1 \sigma = d_0$ ; cf. Section 1, (3), (3 bis), and (†).



Using the construction and notation of 3(b), there is a point  $t \in \overline{Y_0 \times_{X_0} X_1}$  such that  $\mathrm{pr}_1(t) = x$  and  $\mathrm{pr}_2(t) = z$ . Since  $f_0(y) = d_1 \mathrm{pr}_2(t)$ , there is likewise an  $s \in \underline{Y_1}$  such that

$$y = e_1(s), \quad (e_0 \boxtimes f_1)(s) = t.$$

It follows that

$$e_0(s) = \mathrm{pr}_1(e_0 \boxtimes f_1)(s) = \mathrm{pr}_1(t) = x,$$

so  $x \sim y$ .

#### 4. Passage to the quotient by a finite flat groupoid (proof of a special case)

**Theorem 4.1.**— *Consider a  $(\mathbf{Sch}/S)$ -groupoid*

$$X_2 \begin{array}{c} \xrightarrow{d'_2} \\ \xrightarrow[d'_0]{d'_1} \end{array} X_1 \xrightarrow[d_0]{d_1} X_0$$

*and suppose that the following conditions hold:*<sup>16</sup>

- a)  $d_1$  is finite locally free;
- b) for every  $x \in X_0$ , the set  $d_0 d_1^{-1}(x)$  is contained in an affine open subset of  $X_0$ .<sup>17</sup>

*Then:*

- (i) *there exists a cokernel  $(Y, p)$  of  $(d_0, d_1)$  in  $(\mathbf{Sch}/S)$ . Moreover, such a pair  $(Y, p)$  is a cokernel of  $(d_0, d_1)$  in the category of all ringed spaces;*
- (ii)  *$p$  is integral and open, and  $Y$  is affine when  $X_0$  is affine;*<sup>18</sup>
- (iii) *the morphism*

$$X_1 \longrightarrow X_0 \times_Y X_0$$

*with components  $d_0$  and  $d_1$  is surjective;*

- (iv) *if  $(d_0, d_1)$  is an equivalence pair, then  $X_1 \rightarrow X_0 \times_Y X_0$  is an isomorphism<sup>19</sup> and  $p : X_0 \rightarrow Y$  is finite locally free.<sup>20</sup> Moreover,  $(Y, p)$  is a cokernel of  $(d_0, d_1)$  in the category of sheaves for the fppf topology and, for every base change  $Y' \rightarrow Y$ , the scheme  $Y'$  is the cokernel of the groupoid  $X_* \times_Y Y'$  obtained from  $X_*$  by the base change  $X_0 \times_Y Y' \rightarrow X_0$ .*

<sup>16</sup>Editor's note: since  $d_0 = d_1 \sigma$ , where  $\sigma$  is an involutive automorphism of  $X_1$ , these two conditions are symmetric in  $d_1$  and  $d_0$ . Moreover,  $d_0 d_1^{-1}(x) = d_1 d_0^{-1}(x)$ .

<sup>17</sup>Editor's note: condition b) cannot be omitted. H. Hironaka gave an example of an action of the finite group  $G = \mathbb{Z}/2\mathbb{Z}$  on a proper complex variety  $X_0$  for which the quotient  $X_0/G$  is an algebraic space that is not a scheme; see [Hi62], and also [Mum65], Chapter 4, §3.

<sup>18</sup>Editor's note: the assertion that  $p$  is open has been added by repeating the analogous argument given in 6.1.

<sup>19</sup>Editor's note: in this case  $X_1 \rightarrow X_0 \times_S X_0$  is therefore an immersion (EGA I, 5.3.10); see also Exposé VI<sub>B</sub>, 9.2.1. For existence of the quotient—in the category of schemes or in that of algebraic spaces—under the weaker hypothesis that  $d_0$  and  $d_1$  are finite but not necessarily flat, see [An73], §1.1, [Fe03], [Ko08], and the references there.

<sup>20</sup>Editor's note: the consequences that follow have been made explicit; see [Ray67a], Theorem 1(iii), and the proof given below at the end of Section 5.

In particular, for every base change  $S' \rightarrow S$ , the scheme  $Y' = Y \times_S S'$  is the cokernel of the  $S'$ -groupoid  $X'_* = X_* \times_S S'$ . Thus, in this case, formation of the quotient commutes with base change.

It follows immediately from (i) that the underlying topological space of  $Y$  is the quotient of the underlying topological space of  $X_0$  by the equivalence relation defined by the  $(\mathbf{Sch}/S)$ -groupoid  $X_*$ .

We shall first prove the theorem when  $X_0$  is affine and  $d_1$  is locally free of constant rank  $n$ . We shall then see how to reduce the general situation to this special case.

In the case now under consideration,  $X_0, X_1, X_2$  are affine. We may therefore suppose that

$$X_i = \operatorname{Spec} A_i, \quad d_j = \operatorname{Spec} \delta_j, \quad d'_k = \operatorname{Spec} \delta'_k,$$

where the  $A_i$  are commutative rings and the  $\delta_j, \delta'_k$  are ring homomorphisms. The diagram  $(0,1,2)$  may then be replaced by the following dual diagram:

$$(0,1,2)^* \quad \begin{array}{ccccc} & & \delta'_1 & & \\ & & \longleftarrow & & \\ A_2 & \xleftarrow{\delta'_0} & A_1 & \xleftarrow{\delta_0} & A_0 \\ \delta'_2 \uparrow & & \delta_1 \uparrow & & \\ A_1 & \xleftarrow{\delta_1} & A_0 & & \\ & & \delta_0 & & \end{array}$$

The two left-hand squares in this diagram are cocartesian.

Let  $B$  be the subring of  $A_0$  consisting of the elements  $a \in A_0$  for which  $\delta_0(a) = \delta_1(a)$ .

a) We first show that  $A_0$  is integral over  $B$ . For  $a \in A_0$ , let

$$P_{\delta_1}(T, \delta_0(a)) = T^n - \sigma_1 T^{n-1} + \cdots + (-1)^n \sigma_n \quad (1)$$

be the characteristic polynomial of  $\delta_0(a)$ , where  $A_1$  is regarded as an  $A_0$ -algebra through  $\delta_1$ ; see Bourbaki, *Algèbre*, VIII, §12, and *Algèbre commutative*, II, §5, Exercice 9. Since the two left-hand squares of the diagram  $(0,1,2)^*$  are cocartesian, one has

$$\delta_0(P_{\delta_1}(T, \delta_0(a))) = P_{\delta'_2}(T, \delta'_0 \delta_0(a)), \quad (2)$$

$$\delta_1(P_{\delta_1}(T, \delta_0(a))) = P_{\delta'_2}(T, \delta'_1 \delta_0(a)). \quad (3)$$

Because  $\delta'_0 \delta_0 = \delta'_1 \delta_0$ , it follows that

$$\delta_0(P_{\delta_1}(T, \delta_0(a))) = \delta_1(P_{\delta_1}(T, \delta_0(a)));$$

equivalently,

$$\delta_0(\sigma_i) = \delta_1(\sigma_i) \quad \text{for every } i.$$

On the other hand, the Cayley–Hamilton theorem gives

$$\delta_0(a)^n - \delta_1(\sigma_1) \delta_0(a)^{n-1} + \cdots + (-1)^n \delta_1(\sigma_n) = 0.$$

Since  $\delta_1(\sigma_i) = \delta_0(\sigma_i)$ , one also has

$$\delta_0(a)^n - \delta_0(\sigma_1) \delta_0(a)^{n-1} + \cdots + (-1)^n \delta_0(\sigma_n) = 0.$$

Hence

$$a^n - \sigma_1 a^{n-1} + \cdots + (-1)^n \sigma_n = 0,$$

because there is a homomorphism  $\tau : A_1 \rightarrow A_0$  such that  $\tau\delta_0 = \text{id}_{A_0}$ , so  $\delta_0$  is injective. It follows that  $A_0$  is integral over  $B$ .

**b)** Consider two prime ideals  $x$  and  $y$  of  $A_0$ . We shall show that the equality

$$x \cap B = y \cap B$$

implies the existence of a prime ideal  $z$  of  $A_1$  such that

$$x = d_0(z), \quad y = d_1(z).$$

Indeed, if this assertion were false,  $x$  would be distinct from  $\delta_0^{-1}(t)$  for every prime ideal  $t$  of  $A_1$  such that  $\delta_1^{-1}(t) = y$ . For every such  $t$ , one would have

$$\delta_0^{-1}(t) \cap B = \delta_1^{-1}(t) \cap B = y \cap B = x \cap B.$$

It would follow from Cohen–Seidenberg (see Bourbaki, *Algèbre commutative*, V, §2, Corollary 1 to Proposition 1) that  $x$  is contained in none of the ideals  $\delta_0^{-1}(t)$ .<sup>21</sup> There are at most  $n$  prime ideals  $t$  of  $A_1$  satisfying  $\delta_1^{-1}(t) = y$  (see *loc. cit.*, Proposition 3). Hence, by the Prime Avoidance Lemma (*loc. cit.*, II, §1, Proposition 3), there would be an element  $a \in x$  belonging to none of the ideals  $\delta_0^{-1}(t)$ .

Consequently,  $\delta_0(a)$  would belong to none of these prime ideals  $t$ , and the lemma below would imply that the norm  $N_{\delta_1}(\delta_0(a))$  does not belong to  $y$ . Here the norm is computed by regarding  $A_1$  as an  $A_0$ -algebra through  $\delta_1$ ; with the notation of part a),

$$N_{\delta_1}(\delta_0(a)) = \sigma_n.$$

On the other hand,

$$(-1)^{n-1}\sigma_n = a^n + \sum_{i=1}^{n-1} (-1)^i \sigma_i a^{n-i},$$

so this norm belongs to  $B \cap x = B \cap y$ , a contradiction.

**Lemma 4.1.1.**— *Let  $A \rightarrow A'$  be a homomorphism of commutative rings which makes  $A'$  a projective  $A$ -module of rank  $n$ . Let  $p \in \text{Spec}(A)$ , let  $q_1, \dots, q_r$  be the points of  $\text{Spec}(A')$  lying above  $p$ , and let  $a \in A'$ . Then  $a \in q_1 \cup \dots \cup q_r$  if and only if its norm  $N(a)$  belongs to  $p$ .*

Indeed, after replacing  $A$  and  $A'$  by the localizations  $A_p$  and  $A'_p$ , we reduce to the case in which  $(A, p)$  is local and  $A'$  is semilocal, with

$$\text{Spec}(A') = \{q_1, \dots, q_r\}.$$

In this case  $A'$  is a free  $A$ -module of rank  $n$  (see Bourbaki, *Algèbre commutative*, II, §3.2, Corollary 2 to Proposition 5), and  $N(a)$  is the determinant of the endomorphism

$$\ell_a : A' \longrightarrow A', \quad a' \longmapsto aa'.$$

Thus one has the equivalences

$$N(a) \notin p \iff N(a) \text{ is invertible} \iff \ell_a \text{ is invertible} \iff a \notin q_1 \cup \dots \cup q_r.$$

<sup>21</sup>Editor's note: the original has been expanded in what follows; in particular, Lemma 4.1.1, taken from [DG70], III, §2.4, Lemma 4.3, has been added.

**c) Proof of (i).** Set  $Y = \operatorname{Spec} B$  and  $p = \operatorname{Spec} i$ , where  $i : B \hookrightarrow A_0$  is the inclusion. By part a), the morphism  $p : X_0 \rightarrow Y$  is surjective. We first show that  $(Y, p)$  is a cokernel of  $(d_0, d_1)$  in the category of all ringed spaces. Indeed, part b) shows that the set underlying  $\operatorname{Spec} B$  is obtained from the set underlying  $X_0$  by identifying the points  $x$  and  $y$  for which there exists  $z \in X_1$  such that

$$d_1 z = y, \quad d_0 z = x.$$

Moreover, since  $i$  is integral,  $p = \operatorname{Spec} i$  is closed, and therefore  $Y$  carries the quotient topology induced by  $X_0$ .

It follows that  $p$  is open. Indeed, let  $U'$  be any open subset of  $X_0$ . Since  $d_1$  is surjective and finite locally free, it is faithfully flat and of finite presentation, hence open. Thus

$$U = d_1(d_0^{-1}(U')),$$

the saturation of  $U'$  for the equivalence relation defined by  $X_*$ , is open. Since  $Y$  has the quotient topology,  $p(U') = p(U)$  is open.

Finally, the choice of  $B$ , together with the fact that  $p, d_0$ , and  $d_1$  are affine, shows that the canonical sequence of sheaves of rings

$$\mathcal{O}_Y \longrightarrow p_* \mathcal{O}_{X_0} \xrightarrow[p_*(\delta_0)]{p_*(\delta_1)} p_* d_{0*} \mathcal{O}_{X_1} = p_* d_{1*} \mathcal{O}_{X_1}$$

is exact.

It remains to show that  $(Y, p)$  is also the cokernel of  $(d_0, d_1)$  in the category of schemes—more generally, in the category of locally ringed spaces. Let  $q : X_0 \rightarrow Z$  be a morphism of schemes such that  $qd_0 = qd_1$ . By what precedes, there is a unique morphism of ringed spaces  $r : Y \rightarrow Z$  such that  $q = rp$ . We must show that, for every  $y \in Y$ , the homomorphism

$$\mathcal{O}_{Z, r(y)} \longrightarrow \mathcal{O}_{Y, y}$$

induced by  $r$  is local. Since  $p$  is surjective,  $y = p(x)$  for some  $x \in X_0$ , and the assertion follows from the fact that the homomorphism

$$\mathcal{O}_{Z, q(x)} \longrightarrow \mathcal{O}_{X_0, x}$$

induced by  $q$  is local.

**d) Proof of (ii).** This follows from parts a) and c).

**e) Proof of (iii).** Recall that  $\underline{P}$  denotes the set underlying a scheme  $P$ , and that  $\underline{d} : \underline{P} \rightarrow \underline{Q}$  denotes the map induced by a morphism  $d : P \rightarrow Q$ .

**Lemma 4.1.2.**<sup>22</sup> *Let  $(A, \mathfrak{m})$  be a local ring, let  $k$  be its residue field, and let  $K/k$  be a field extension. There exists a local, flat  $A$ -algebra  $B$  such that  $B/\mathfrak{m}B$  is  $k$ -isomorphic to  $K$ . If  $K/k$  is finite,  $B$  may moreover be chosen finite and free over  $A$ .*

This is proved in EGA 0<sub>III</sub>, 10.3.1, where it is also shown that  $B$  may be chosen Noetherian when  $A$  is Noetherian. For the reader's convenience, we recall the proof.

Put  $A' = A[T]$ , where  $T$  is an indeterminate. If  $K = k(T)$ , let  $\mathfrak{p} = \mathfrak{m}A'$  and  $B = A'_{\mathfrak{p}}$ . Then

$$B/\mathfrak{m}B \simeq k[T]_{(0)} = k(T).$$

The ring  $B$  is flat over  $A'$ , which is a free  $A$ -module, so  $B$  is flat over  $A$ .

<sup>22</sup>Editor's note: this lemma, which is used repeatedly in this Exposé and in later Exposés (VI<sub>A</sub>, VI<sub>B</sub>), has been inserted here. It appeared as Lemma VI<sub>B</sub>, 4.5.1, in the original 1965 edition of SGA 3.

Suppose next that  $K = k(t) = k[t]$ , where  $t$  is algebraic over  $k$ . Set  $B = A'/(F)$ , where  $F \in A'$  is a monic polynomial whose image in  $k[T]$  is the minimal polynomial  $f$  of  $t$  over  $k$ . Then  $B$  is a free  $A$ -module of finite rank

$$\deg(F) = \deg(f).$$

In particular,  $B$  is integral over  $A$ , and hence every maximal ideal of  $B$  contains  $\mathfrak{m}$ . Since

$$B/\mathfrak{m}B \simeq k[T]/(f) \simeq K,$$

$B$  is local, with maximal ideal  $\mathfrak{m}B$ . This already proves that  $B$  can be chosen finite and free over  $A$  when  $[K : k] < \infty$ .

In the general case, choose a family  $(t_i)_{i \in I}$  of generators of  $K/k$ , and well-order  $I$ ; that is, give  $I$  a total order  $\leq$  in which every nonempty subset has a least element. For  $i \in I$ , let  $k_i$  (respectively  $k_{<i}$ ) be the subfield of  $K$  generated by the  $t_j$  with  $j \leq i$  (respectively  $j < i$ ). By adding one element if necessary, we may suppose that  $I$  has a greatest element  $\xi$ , so that  $K = k_\xi$ .

Let  $J \subseteq I$  be the set of indices  $i$  for which there exists a direct system  $(A_j)_{j \leq i}$  of local, flat  $A$ -algebras such that

$$A_j/\mathfrak{m}A_j \simeq k_j$$

and  $A_j$  is flat over  $A_\ell$  whenever  $\ell < j$ . Suppose that  $I \setminus J$  is nonempty, let  $i$  be its least element, and put

$$A' = \varinjlim_{j < i} A_j.$$

Since tensor products commute with direct limits,  $A'$  is flat over  $A$  and over every  $A_j$ ,  $j < i$ , and

$$A'/\mathfrak{m}A' \simeq A' \otimes_A (A/\mathfrak{m}) \simeq k_{<i}.$$

Moreover,  $A'$  is local, with maximal ideal  $\mathfrak{m}A'$ . Indeed, if  $x = f_j(x_j)$  is not invertible, then  $x_j$  is not invertible and hence belongs to the maximal ideal  $\mathfrak{m}A_j$  of  $A_j$ ; therefore  $x \in \mathfrak{m}A'$ .

The monogenic case treated above now gives a local, flat  $A'$ -algebra  $A_i$  such that

$$A_i/\mathfrak{m}A_i \simeq k_{<i}(t_i) = k_i.$$

The algebra  $A_i$  is flat over every  $A_j$ ,  $j < i$ , so  $i \in J$ , contrary to the choice of  $i$ . This contradiction proves that  $J = I$ , and  $A_\xi$  has the required properties. This proves Lemma 4.1.2.

Let us now prove Theorem 4.1(iii). Recall once more that  $\underline{P}$  denotes the set underlying a scheme  $P$ , and that  $\underline{d} : \underline{P} \rightarrow \underline{Q}$  is the map induced by a morphism  $d : P \rightarrow Q$ . Part b) can then be restated by saying that the map

$$\underline{d}_0 \boxtimes \underline{d}_1 : \underline{X}_1 \longrightarrow \underline{X}_0 \times_Y \underline{X}_0$$

with components  $\underline{d}_0$  and  $\underline{d}_1$  is surjective. This map factors as

$$\underline{X}_1 \xrightarrow{\underline{d}_0 \boxtimes \underline{d}_1} \underline{X}_0 \times_Y \underline{X}_0 \xrightarrow{q} \underline{X}_0 \times_Y \underline{X}_0,$$

where  $q$  is the canonical map. The image of  $\underline{d}_0 \boxtimes \underline{d}_1$  therefore contains every point  $v \in \underline{X}_0 \times_Y \underline{X}_0$  such that

$$\{v\} = q^{-1}(q(v)).$$

This condition is satisfied in particular when  $v$  is *rational over*  $Y$ , that is, when the residue field  $\kappa(v)$  is identified with the residue field  $\kappa(w)$  of the image  $w$  of  $v$  in  $Y$ .<sup>23</sup>

<sup>23</sup>Editor's note: the pages of Lecture Notes in Mathematics 151 are permuted here; their actual order is 265–266–268–269–267–270–271.

If  $v \in \underline{X_0 \times_Y X_0}$  is not rational over  $Y$ , again let  $w$  be its image in  $Y$ . By Lemma 4.1.2, there exist a local ring  $C$  with radical  $\mathfrak{m}$  and a local, flat homomorphism

$$f : \mathcal{O}_{Y,w} \longrightarrow C$$

such that  $C/\mathfrak{m}$  is isomorphic to  $\kappa(v)$  as a  $\kappa(w)$ -algebra. Put  $Y' = \operatorname{Spec} C$ , and let  $\pi : Y' \rightarrow Y$  be the morphism induced by  $f$ . The canonical projection

$$(X_0 \times_Y X_0) \times_Y Y' \longrightarrow X_0 \times_Y X_0$$

sends some point  $v'$ , rational over  $Y'$ , to  $v$ . Since

$$(X_0 \times_Y X_0) \times_Y Y' \simeq (X_0 \times_Y Y') \times_{Y'} (X_0 \times_Y Y'),$$

and since the hypotheses of Theorem 4.1 and the preceding results, in particular part b), remain valid after the base change  $\pi : Y' \rightarrow Y$ , the point  $v'$  is the image of an element

$$u' \in \underline{X_1 \times_Y Y'}$$

under the base change of  $d_0 \boxtimes d_1$ . If  $u$  is the image of  $u'$  in  $X_1$ , then

$$v = (d_0 \boxtimes d_1)(u),$$

as required.

**f) Proof of (iv).**<sup>24</sup>

**Lemma 4.1.3.**— *If a monomorphism of schemes  $f : T \rightarrow Z$  is finite, it is a closed immersion.*

Covering  $Z$  by affine open subsets  $Z_i$ , and replacing  $f$  by the induced morphisms  $f^{-1}(Z_i) \rightarrow Z_i$ , reduces us—since a finite morphism is affine—to the case

$$Z = \operatorname{Spec} B, \quad T = \operatorname{Spec} A.$$

Because  $f$  is a monomorphism, the diagonal morphism

$$T \longrightarrow T \times_Z T$$

is an isomorphism (EGA I, 5.3.8); equivalently, multiplication gives an isomorphism

$$A \otimes_B A \xrightarrow{\sim} A.$$

Consequently, for every maximal ideal  $\mathfrak{m}$  of  $B$ , there is an isomorphism

$$(A/\mathfrak{m}A) \otimes_k (A/\mathfrak{m}A) \xrightarrow{\sim} A/\mathfrak{m}A, \quad k = B/\mathfrak{m}.$$

Since  $A$  is finite over  $B$ , the  $k$ -vector space  $A/\mathfrak{m}A$  has finite dimension  $d$ , and this isomorphism implies  $d = 0$  or  $d = 1$ . Hence the map

$$k = B/\mathfrak{m} \longrightarrow A/\mathfrak{m}A$$

is surjective. Nakayama's lemma, applied after localization at  $\mathfrak{m}$ , shows that  $B_{\mathfrak{m}} \rightarrow A_{\mathfrak{m}}$  is surjective. It follows that  $B \rightarrow A$  is surjective: its cokernel  $C$  satisfies  $C_{\mathfrak{m}} = 0$  for every maximal ideal  $\mathfrak{m}$ , and therefore vanishes. This proves the lemma.

<sup>24</sup>Editor's note: the following lemma has been added from [DG70], I, §5, Proposition 1.5; see also the proof of EGA IV<sub>3</sub>, 8.11.5. It was used implicitly in the original and explicitly in [DG70], III, §2, 4.6. It is also useful in Theorem 7.1 below.

Let us now prove (iv). By hypothesis,

$$X_0 = \operatorname{Spec} A_0, \quad X_1 = \operatorname{Spec} A_1,$$

and, for  $i = 0, 1$ , the homomorphism  $\delta_i : A_0 \rightarrow A_1$  makes  $A_1$  a finite  $A_0$ -module. Therefore, a fortiori, the homomorphism

$$A_0 \otimes_B A_0 \longrightarrow A_1$$

is finite.

We assume in addition that

$$d = d_0 \boxtimes d_1 : X_1 \longrightarrow X_0 \times_Y X_0$$

is a monomorphism. By Lemma 4.1.3, the corresponding homomorphism  $A_0 \otimes_B A_0 \rightarrow A_1$  is surjective. We shall prove that it is an isomorphism; in the course of doing so, we shall also prove that  $p : X_0 \rightarrow Y$  is finite locally free.

It is enough to prove that, for every prime ideal  $\mathfrak{p}$  of  $B$ , the homomorphism

$$(A_0)_{\mathfrak{p}} \otimes_{B_{\mathfrak{p}}} (A_0)_{\mathfrak{p}} \longrightarrow (A_1)_{\mathfrak{p}},$$

with components  $\delta_{0\mathfrak{p}}$  and  $\delta_{1\mathfrak{p}}$ , is bijective. Thus we may suppose that  $B$  is local. Part b) then shows that  $(A_0)_{\mathfrak{p}}$  is semilocal. Indeed, if  $\mathfrak{m}$  is one of its maximal ideals, the others have the form  $\delta_0^{-1}(\mathfrak{n})$ , where  $\mathfrak{n}$  ranges over the prime ideals of  $A_1$  satisfying  $\delta_1^{-1}(\mathfrak{n}) = \mathfrak{m}$ . There are at most

$$n = [A_1 : A_0]$$

such prime ideals  $\mathfrak{n}$ .

After a faithfully flat base change, we may also suppose that the residue field of  $B$  is infinite.<sup>25</sup> We can then use the following lemma.

**Lemma 4.2.**— *Let  $B$  be a local ring with infinite residue field, let  $A$  be a semilocal ring, and let  $i : B \rightarrow A$  be a homomorphism which sends the maximal ideal  $\mathfrak{n}$  of  $B$  into the radical  $\mathfrak{r}$  of  $A$ . Let  $M$  be a free  $A$ -module of rank  $n$ , and let  $N$  be a  $B$ -submodule of  $M$  which generates  $M$  as an  $A$ -module. Then  $N$  contains an  $A$ -basis of  $M$ .*

Recall that a sequence  $m_1, \dots, m_n$  of elements of  $M$  is an  $A$ -basis if and only if its canonical images in  $M/\mathfrak{r}M$  form a basis over  $A/\mathfrak{r}$ . We may therefore replace

$$M \text{ by } M/\mathfrak{r}M, \quad N \text{ by } N/(N \cap \mathfrak{r}M), \quad A \text{ by } A/\mathfrak{r}, \quad B \text{ by } B/\mathfrak{n}.$$

In this case the lemma is elementary. Namely, if

$$A = K_1 \times \cdots \times K_r, \quad M = K_1^n \times \cdots \times K_r^n,$$

choose, for each  $j$ , an element  $x_j \in N$  whose  $j$ -th component is nonzero. Because the field  $B$  is infinite, a suitable  $B$ -linear combination  $x$  of the  $x_j$  has all its components nonzero. Replace  $M$  by  $M/Ax$ , and proceed by induction on  $n$ .

Apply the lemma with  $A = A_0$ , with  $i : B \hookrightarrow A_0$ , with  $M = A_1$  regarded as an  $A_0$ -module through  $\delta_1$ , and with  $N = \delta_0(A_0)$ . Indeed, since

$$d_0 \boxtimes d_1 : X_1 \longrightarrow X_0 \times_Y X_0$$

is a closed immersion, the homomorphism

$$A_0 \otimes_B A_0 \longrightarrow A_1$$

---

<sup>25</sup>Editor's note: see Lemma 4.1.2.

with components  $\delta_0, \delta_1$  is surjective. This says exactly that  $\delta_0(A_0)$  generates the  $A_0$ -module  $A_1$ .

Choose  $a_1, \dots, a_n \in A_0$  so that  $\delta_0(a_1), \dots, \delta_0(a_n)$  form an  $A_0$ -basis of  $A_1$ . If  $a_1, \dots, a_n$  form a  $B$ -basis of  $A_0$ , then  $A_0 \otimes_B A_0 \rightarrow A_1$  sends the basis

$$(1 \otimes a_i)_{1 \leq i \leq n}$$

to the basis

$$(\delta_0(a_i))_{1 \leq i \leq n},$$

and is therefore bijective. Thus, if

$$\varepsilon : \mathbb{Z}^n \longrightarrow A_0$$

is the homomorphism of abelian groups which sends the standard basis of  $\mathbb{Z}^n$  to  $a_1, \dots, a_n$ , it remains only to prove that the map

$$B \otimes_{\mathbb{Z}} \mathbb{Z}^n \longrightarrow A_0$$

with components  $i$  and  $\varepsilon$  is bijective.

The dual diagram  $(0, 1, 2)^*$  considered at the beginning of this proof induces the following commutative diagram:

$$\begin{array}{ccccc}
 A_2 & \xleftarrow[\delta'_0]{\delta'_1} & A_1 & \xleftarrow{\delta_0} & A_0 \\
 u_2 \uparrow & & u_1 \uparrow \cong & & u_0 \uparrow \\
 A_1 \otimes_{\mathbb{Z}} \mathbb{Z}^n & \xleftarrow[\delta_0 \otimes \mathbb{Z}^n]{\delta_1 \otimes \mathbb{Z}^n} & A_0 \otimes_{\mathbb{Z}} \mathbb{Z}^n & \xleftarrow{i \otimes \mathbb{Z}^n} & B \otimes_{\mathbb{Z}} \mathbb{Z}^n
 \end{array}$$

Figure 22: The commutative ring-and-module diagram concluding the affine proof of Theorem 4.1(iv).

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille Exp5-13oct24.pdf, component-local p. 17; combined-reader p. 279; re-edited page header p. 267.*

Here  $u_0, u_1, u_2$  have, respectively, the component pairs

$$(i, \varepsilon), \quad (\delta_1, \delta_0 \varepsilon), \quad (\delta'_2, \delta'_0 \delta_0 \varepsilon).$$

We know that  $u_1$  is an isomorphism. Since the two left-hand squares of  $(0, 1, 2)^*$  are co-cartesian,  $u_2$  is bijective. The two horizontal rows of the diagram are exact, and therefore  $u_0$  is bijective.<sup>26</sup> Thus  $A_0$  is a locally free  $B$ -module of rank  $n$ . By the preceding reductions, it follows that

$$\delta_0 \otimes \delta_1 : A_0 \otimes_B A_0 \xrightarrow{\sim} A_1$$

is an isomorphism. This completes the proof of Theorem 4.1 in the special case under consideration:  $X_0$  is affine and  $d_1$  is locally free of constant rank  $n$ .

<sup>26</sup>Editor's note: the argument which follows has been added.



## 5. Passage to the quotient by a finite flat groupoid (the general case)

a) Let  $U^{(n)}$  be the largest open subset of  $X_0$  over which  $d_1$  is finite locally free of rank  $n$ . The scheme  $X_0$  is the disjoint union of the  $U^{(n)}$ . The two Cartesian squares

$$\begin{array}{ccc} X_2 & \xrightarrow{d'_0} & X_1 \\ d'_2 \downarrow & & \downarrow d_1 \\ X_1 & \xrightarrow{d_0} & X_0 \end{array} \quad \text{and} \quad \begin{array}{ccc} X_2 & \xrightarrow{d'_1} & X_1 \\ d'_2 \downarrow & & \downarrow d_1 \\ X_1 & \xrightarrow{d_1} & X_0 \end{array}$$

Figure 23: The two Cartesian squares controlling the rank strata  $U^{(n)}$ .

*Native Loop-2 reconstructions; source locator: Polo–Gille Exp5-13oct24.pdf, component-local p. 17; combined-reader p. 279; re-edited page header p. 267.*

show that the inverse images of  $U^{(n)}$  under  $d_0$  and  $d_1$  both coincide with the largest open subset of  $X_1$  over which  $d'_2$  is locally free of rank  $n$ .<sup>27</sup> Thus

$$d_0^{-1}(U^{(n)}) = d_1^{-1}(U^{(n)}),$$

and the groupoid  $X_*$  is the disjoint union of the groupoids  $X_*^{(n)}$  induced by  $X_*$  on the open-and-closed subschemes  $U^{(n)}$ . It is therefore enough to prove Theorem 4.1 for each  $X_*^{(n)}$ . We are reduced to the case in which  $d_1$  is finite locally free of rank  $n$ .

b) *We can now prove the theorem in the general case.*

By part a), we may suppose that  $d_1$  is locally free of rank  $n$ . Let  $(Y, p)$  be a cokernel of  $(d_0, d_1)$  in the category of all ringed spaces. The argument at the end of 4(c) shows that, in order to prove Theorem 4.1(i), it is enough to show that  $Y$  is a scheme and that  $p : X_0 \rightarrow Y$  is a morphism of schemes. By Lemma 1.2, this question is local on  $Y$ . Let  $y \in Y$  and choose  $x \in X_0$  with  $p(x) = y$ . If  $x$  has a saturated affine open neighborhood  $U$ , then Section 4 shows that  $p(U)$  is an affine open subset of  $Y$ , and  $p|_U : U \rightarrow p(U)$  is a morphism of schemes. Thus it remains to prove that every  $x \in X_0$  has a saturated affine open neighborhood. We proceed as follows; the argument is taken from SGA 1, VIII, Corollary 7.6.

$$\begin{array}{ccccccc} d_1(d_0^{-1}(x)) & \mathcal{U} = (V_f)' & \subset & V_f & \subset & V' & \subset & V & \subset & X_0 \\ & \uparrow & & \uparrow & & \uparrow & & \nwarrow & & \\ & \text{largest saturated} & & \text{special affine} & & \text{largest saturated} & & \text{affine open} & & \\ & \text{open in } V_f & & \text{open in } V & & \text{open in } V & & & & \end{array}$$

Figure 24: The nested open subsets used to construct a saturated affine neighborhood of  $x$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Polo–Gille Exp5-13oct24.pdf, component-local p. 18; combined-reader p. 280; re-edited page header p. 268.*

<sup>27</sup>Editor's note: since  $d_0$  (respectively  $d_1$ ) is surjective, flat, and finite, hence faithfully flat and affine,  $d'_2$  has rank  $n$  over a neighborhood of a point  $x \in X_1$  if and only if  $d_1$  has rank  $n$  over a neighborhood of  $d_0(x)$  (respectively  $d_1(x)$ ).

By condition 4.1(b), there is an affine open subset  $V \subseteq X_0$  which contains  $d_1(d_0^{-1}(x))$ .<sup>28</sup> Put  $F = X_0 \setminus V$ . Since  $d_1$  is integral,  $d_1(d_0^{-1}(F))$  is closed, and

$$V' = X_0 \setminus d_1(d_0^{-1}(F))$$

is the largest saturated open subset contained in  $V$ . The set  $V'$  is a neighborhood of the finite set  $d_1(d_0^{-1}(x))$ . Hence there is a section  $f$  of the structure sheaf of  $V$ , vanishing on  $V \setminus V'$ , such that  $d_1(d_0^{-1}(x))$  is contained in the principal open subset  $V_f \subseteq V$ . We shall prove that the largest saturated open subset  $(V_f)'$  contained in  $V_f$  is affine; it will then be the required neighborhood.

Put  $Z(f) = V' \setminus V_f$ . Then  $d_0^{-1}(Z(f))$  is the set of points of

$$d_0^{-1}(V') = d_1^{-1}(V')$$

at which  $d_0^*(f)$  vanishes. The morphism induced by  $d_1$  from this common inverse image to  $V'$  is locally free of rank  $n$ .<sup>29</sup> Lemma 4.1.1 therefore shows that  $d_1(d_0^{-1}(Z(f)))$  is precisely the zero locus of the norm  $N$  of  $d_0^*(f)$  relative to  $d_1$ . It follows that

$$(V_f)' = V' \setminus d_1(d_0^{-1}(Z(f)))$$

is the set of points of  $V_f$  at which  $N$  does not vanish. Consequently,  $(V_f)'$  is affine.

This proves Theorem 4.1(i); assertions (ii), (iii), and the first part of (iv) are then clear. It remains to prove the consequences stated at the end of (iv); see [Ray67a], Theorem 1(iii).

By hypothesis, the groupoid  $X_*$  comes from an equivalence relation

$$i : R \longrightarrow X_0 \times X_0,$$

where  $i$  is an immersion (see editor's note 19). We have shown that  $R$  is effective (see Exposé IV, 3.3.2), and that

$$p : X_0 \longrightarrow Y = X_0/R$$

is surjective and finite locally free; in particular, it is faithfully flat and of finite presentation.

Consequently, if  $(M)$  denotes the family of faithfully flat morphisms locally of finite presentation, then  $R$  is  $(M)$ -effective. By Exposé IV, 6.3.3,  $(Y, p)$  therefore represents the quotient sheaf of  $X_0$  by  $R$  for the fppf topology, and the base-change assertions follow from Exposé IV, 3.4.3.1.

**Remark 5.1.**—<sup>30</sup> Keep the hypotheses and notation of Theorem 4.1, and suppose in addition that  $S$  is locally Noetherian and that  $\pi_0 : X_0 \rightarrow S$  is quasi-projective. We shall prove that  $\pi : Y \rightarrow S$  is quasi-projective.

The preceding hypotheses imply that  $Y \rightarrow S$  is of finite type; see the proof of 6.1(ii). Let  $\mathcal{A}$  be an invertible  $\mathcal{O}_{X_0}$ -module which is ample for  $\pi_0$ . By EGA II, 6.1.12,  $p_*\mathcal{A}$  is an invertible  $p_*\mathcal{O}_{X_0}$ -module. Hence  $Y$  has an affine open cover  $(V_i)_{i \in I}$  such that  $\mathcal{A}$  is trivial over each of the saturated affine open subsets

$$U_i = p^{-1}(V_i).$$

For each  $i$ , put

$$A_{i,0} = \mathcal{O}_{X_0}(U_i),$$

<sup>28</sup>Editor's note:  $d_1(d_0^{-1}(x)) = d_0(d_1^{-1}(x))$ ; see editor's note 16 in Theorem 4.1.

<sup>29</sup>Editor's note: the reference to Lemma 4.1.1 has been added; see [DG70], III, §5.2, p. 313.

<sup>30</sup>Editor's note: this paragraph has been added.

and let  $A_{i,1}$  be the ring of the affine open subset  $d_0^{-1}(U_i) = d_1^{-1}(U_i)$  of  $X_1$ . Let  $\delta_{i,0}$  (respectively  $\delta_{i,1}$ ) be the homomorphism  $A_{i,0} \rightarrow A_{i,1}$  induced by  $d_0$  (respectively  $d_1$ ), and put

$$B_i = \mathcal{O}_Y(V_i) = \{b \in A_{i,0} \mid \delta_{i,0}(b) = \delta_{i,1}(b)\}.$$

Following EGA II, §6.5, consider the invertible  $\mathcal{O}_{X_0}$ -module

$$N_{d_1}(d_0^* \mathcal{A}),$$

the norm, relative to the finite locally free morphism  $d_1 : X_1 \rightarrow X_0$ , of the invertible  $\mathcal{O}_{X_1}$ -module  $d_0^* \mathcal{A}$ . Suppose that  $\mathcal{A}$ , relative to the cover  $(U_i)_{i \in I}$ , is given by transition functions

$$c_{ij} \in \mathcal{O}_{X_0}(U_i \cap U_j)^\times.$$

Then  $N_{d_1}(d_0^* \mathcal{A})$  is given by the transition functions

$$N_{\delta_1}(\delta_0(c_{ij})) \in \mathcal{O}_{X_0}(U_i \cap U_j)^\times.$$

By 4(a), these elements belong to  $\mathcal{O}_Y(V_i \cap V_j)^\times$ , and hence define an invertible  $\mathcal{O}_Y$ -module  $\mathcal{L}$  such that

$$p^* \mathcal{L} = N_{d_1}(d_0^* \mathcal{A}).$$

Moreover, for every  $n \in \mathbb{N}^*$ ,

$$p^*(\mathcal{L}^n) = N_{d_1}(d_0^*(\mathcal{A}^n));$$

see *loc. cit.*, (6.5.2.1).

We now show that  $\mathcal{L}$  is ample for  $\pi : Y \rightarrow S$ . Replacing  $S$  by an affine open subset, we may suppose that  $S$  is affine. Let  $y \in Y$ , choose  $x \in X_0$  with  $p(x) = y$ , let  $V$  be an affine open neighborhood of  $y$  in  $Y$ , and put  $U = p^{-1}(V)$ . Since  $\mathcal{A}$  is  $\pi_0$ -ample, there exist  $n \in \mathbb{N}^*$  and a section

$$s \in \Gamma(X_0, \mathcal{A}^n)$$

such that

$$x \in (X_0)_s \subseteq U.$$

With the notation above,  $s$  is represented by sections

$$a_i \in A_{i,0} = \mathcal{O}_{X_0}(U_i), \quad a_i = c_{ij} a_j \quad \text{on } U_i \cap U_j,$$

and  $(X_0)_s$  is the union of the open subsets

$$U'_i = \{\mathfrak{p} \in \text{Spec}(A_{i,0}) \mid a_i \notin \mathfrak{p}\}.$$

For every  $i$ , put

$$N(a_i) = N_{\delta_1}(\delta_0(a_i)) \in B_i.$$

Theorem 4.1(i) and Lemma 4.1.1 give

$$p(U'_i) = p d_1 d_0^{-1}(U'_i) = p d_1(\{\mathfrak{q} \in \text{Spec}(A_{i,1}) \mid \delta_{i,0}(a_i) \notin \mathfrak{q}\}),$$

while

$$d_1(\{\mathfrak{q} \in \text{Spec}(A_{i,1}) \mid \delta_{i,0}(a_i) \notin \mathfrak{q}\}) = \{\mathfrak{p} \in \text{Spec}(A_{i,0}) \mid N_{\delta_1}(\delta_{i,0}(a_i)) \notin \mathfrak{p}\}.$$

It follows that

$$p(U'_i) = \{\mathfrak{p} \in \text{Spec}(B_i) \mid N(a_i) \notin \mathfrak{p}\}.$$

Consequently,

$$p((X_0)_s) = Y_{N(s)},$$

where  $N(s)$  is the section of  $\mathcal{L}^n$  on  $Y$  defined by the sections  $N(a_i) \in \mathcal{O}_Y(V_i)$ . Thus

$$y \in p((X_0)_s) = Y_{N(s)} \subseteq p(U) = V. \quad (*)$$

This proves that  $\mathcal{L}$  is ample for  $\pi : Y \rightarrow S$ , and hence completes the proof that  $\pi$  is quasi-projective.

## 6. Passage to the quotient when a quasi-section exists

We shall now prove a technical lemma that will be used in the proof of the two theorems we have in view. Let  $S$  be a scheme and let

$$X_2 \begin{array}{c} \xrightarrow{d'_0, d'_1, d'_2} \\ \xrightarrow{\quad\quad\quad} \\ \xrightarrow{\quad\quad\quad} \end{array} X_1 \xrightarrow{d_0, d_1} X_0$$

Figure 25: The groupoid  $X_*$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 20 / combined-reader p. 282 / re-edited header p. 270.*

be a  $(\mathbf{Sch}/S)$ -groupoid. We call a *quasi-section* of the groupoid  $X_*$  any subscheme  $U$  of  $X_0$  for which the following two conditions hold:

- (1) The restriction  $v$  of  $d_1$  to  $d_0^{-1}(U)$  is a finite, locally free, surjective morphism from  $d_0^{-1}(U)$  onto  $X_0$ .
- (2) Every subset  $E$  of  $U$  consisting of points that are pairwise equivalent for the equivalence relation defined by  $X_*$  (§3.e)) is contained in an affine open subset of  $U$ .<sup>31</sup>

If  $U$  is a quasi-section of  $X_*$ , the  $(\mathbf{Sch}/S)$ -groupoid

$$U_2 \begin{array}{c} \xrightarrow{u'_0, u'_1, u'_2} \\ \xrightarrow{\quad\quad\quad} \\ \xrightarrow{\quad\quad\quad} \end{array} U_1 \xrightarrow{u_0, u_1} U$$

Figure 26: The groupoid  $U_*$  induced by  $X_*$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 20 / combined-reader p. 282 / re-edited header p. 270.*

induced by  $X_*$  and the inclusion  $U \hookrightarrow X_0$  (see §3.a)) satisfies the hypotheses of Theorem 4.1. Indeed, put

$$V = d_0^{-1}(U),$$

and let  $u$  and  $v$  be the morphisms with source  $V$  induced by  $d_0$  and  $d_1$ , respectively:

$$X_0 \xleftarrow{v} V \xrightarrow{u} U$$

Figure 27: The morphisms  $v : V \rightarrow X_0$  and  $u : V \rightarrow U$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 20 / combined-reader p. 282 / re-edited header p. 270.*

By §3.b), there is a Cartesian square

---

<sup>31</sup>Editor's note: If  $x, y \in E$ , there is a  $z \in X_1$  such that  $d_1(z) = x$  and  $d_0(z) = y$ ; thus  $z$  belongs to  $v^{-1}(x)$ , which is finite by (1). Hence  $E$  is contained in the finite set  $d_0(v^{-1}(x)) = d_0 d_1^{-1}(x) \cap U$ . The source PDF prints  $d_1(x) = x$  in the first equality; the type-correct reading  $d_1(z) = x$  is used here.

$$\begin{array}{ccc}
 U_1 & \longrightarrow & V \\
 u_1 \downarrow & & \downarrow v \\
 U & \xrightarrow{\text{inclusion}} & X_0
 \end{array}$$

Figure 28: The Cartesian square defining  $U_1$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 20 / combined-reader p. 282 / re-edited header p. 270.*

so  $u_1$  is surjective and finite locally free by condition (1). Together with condition (2), condition (1) therefore ensures that the groupoid  $U_*$  satisfies the hypotheses of Theorem 4.1. In particular,  $\text{Coker}(u_0, u_1)$  exists in  $(\mathbf{Sch}/S)$ . Moreover,  $d_0$  has a section, so  $u$  is a universal effective epimorphism (see Exposé III, 1.12). Proposition 3.1 then shows that  $\text{Coker}(u_0, u_1)$  coincides with the cokernel  $\text{Coker}(v_0, v_1)$  of the groupoid  $V_*$ :

$$\begin{array}{ccccc}
 V_2 & \xrightleftharpoons[v'_0]{v'_1} & V_1 & \xrightleftharpoons[v_0]{v_1} & V
 \end{array}$$

Figure 29: The groupoid  $V_*$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 20 / combined-reader p. 282 / re-edited header p. 270.*

This is the inverse image of  $U_*$  under the base change  $u : V \rightarrow U$ , and hence also the inverse image of  $X_*$  under the base change

$$V \xrightarrow{\text{inclusion}} X_1 \xrightarrow{d_0} X_0$$

Figure 30: The base change defining  $V_*$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 21 / combined-reader p. 283 / re-edited header p. 271.*

By §3.c),  $V_*$  is isomorphic to the groupoid  $V'_*$ , the inverse image of  $X_*$  under the base change

$$v : \quad V \xrightarrow{\text{inclusion}} X_1 \xrightarrow{d_1} X_0$$

Figure 31: The base change defining  $V'_*$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 21 / combined-reader p. 283 / re-edited header p. 271.*

Thus  $V'_*$  admits a cokernel in  $(\mathbf{Sch}/S)$ . Since  $v : V \rightarrow X_0$  is flat, surjective, and finite, it is faithfully flat and quasi-compact, and therefore a universal effective epimorphism by Exposé III, 6.3.2. Proposition 3.1 consequently shows that  $X_*$  admits a cokernel  $\text{Coker}(d_0, d_1)$  in  $(\mathbf{Sch}/S)$ . We have therefore proved the first assertion of part (i) of the following lemma.<sup>32</sup>

<sup>32</sup>Editor's note: The argument that follows has been slightly modified. In particular, the additional

**Lemma 6.1.**—*Suppose that the  $(\mathbf{Sch}/S)$ -groupoid  $X_*$  possesses a quasi-section. Then:*

- (i) *There exists a cokernel  $(Y, p)$  of  $(d_0, d_1)$  in  $(\mathbf{Sch}/S)$ . Moreover, every such  $(Y, p)$  is a cokernel of  $(d_0, d_1)$  in the category of all ringed spaces.*
- (i') *The morphism  $p$  is surjective; it is open (respectively, universally closed) if  $d_0$  is.*
- (ii) *Suppose that  $S$  is locally Noetherian and that  $X_0$  is locally of finite type (respectively, of finite type) over  $S$ . Then  $p$  and  $Y \rightarrow S$  are locally of finite presentation (respectively, of finite presentation).*
- (iii) *The morphism  $X_1 \rightarrow X_0 \times_Y X_0$  with components  $d_0$  and  $d_1$  is surjective.*
- (iv) *If  $(d_0, d_1)$  is an equivalence pair, then  $X_1 \rightarrow X_0 \times_Y X_0$  is an isomorphism. Moreover, if  $d_0 : X_1 \rightarrow X_0$  is flat, then  $p$  is faithfully flat.*

Before proving the second assertion of part (i), we shall prove parts (i'), (ii), and (iii).

**a) Proof of (i') and (ii).**—We have just seen that  $(Y, p)$  identifies both with  $\text{Coker}(v_0, v_1)$  and with  $\text{Coker}(u_0, u_1)$ . Let  $q$  and  $r$  be the canonical epimorphisms from  $U$  and  $V$ , respectively, to  $Y$ :

$$\begin{array}{ccccc} X_0 & \xleftarrow{v} & V & \xrightarrow{u} & U \\ & \searrow p & \downarrow r & \swarrow q & \\ & & Y & & \end{array}$$

Figure 32: The canonical epimorphisms to  $Y$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 21 / combined-reader p. 283 / re-edited header p. 271.*

By hypothesis,  $v$  is surjective and finite locally free, hence open.

On the other hand, if  $d_0 : X_1 \rightarrow X_0$  is open (respectively, universally closed), then  $u$ , being obtained from  $d_0$  by base change, has the same property. By Theorem 4.1,  $q$  is surjective, integral, and open. It follows that  $r$  is surjective and is open (respectively, universally closed) if  $d_0$  is. Since  $v$  is surjective, the same is true of  $p$ . This proves part (i').

Now suppose that  $S$  is locally Noetherian and that  $X_0$  is locally of finite type over  $S$ , so that  $X_0$  is locally Noetherian.

We show that  $Y$  is locally of finite presentation over  $S$ . Let

$$S' = \text{Spec } R$$

be an affine open subset of  $S$ , let

$$Y' = \text{Spec } B$$

be an affine open subset of  $Y$  mapping into  $S'$ , and let

$$U' = \text{Spec } A$$

be the inverse image of  $Y'$  in  $U$ . Since  $R$  is Noetherian, it is enough to show that  $B$  is a finitely generated  $R$ -algebra. By §§4 and 5,  $B$  is contained in  $A$ , which is a finitely generated  $R$ -algebra; the assertion therefore follows because  $R$  is Noetherian and  $A$  is integral over  $B$ .

---

hypothesis that  $d_0$  be flat has been moved to part (iv); the former part (ii) has been split into parts (i') and (ii); and the second assertion of part (i') has been added.

Finally, since  $X_0 \rightarrow S$  is locally of finite type, so is  $p$  (EGA I, 6.6.6). Hence  $p$  is locally of finite presentation because  $Y$  is locally Noetherian.

It remains to prove the final assertion of part (ii). Suppose in addition that  $X_0$  is of finite type over  $S$ . Since  $p$  is surjective,  $Y$  is then quasi-compact over  $S$ , and hence of finite type over  $S$ . Because  $S$  is locally Noetherian, the morphisms  $X_0 \rightarrow S$  and  $Y \rightarrow S$  are of finite presentation, and therefore so is  $p : X_0 \rightarrow Y$  (EGA IV<sub>1</sub>, 1.6.2(v)).

**b) Proof of (iii).**—The groupoid  $V_*$  with base  $V$  is isomorphic both to the inverse image of  $U_*$  under the base change  $u$  and to the inverse image of  $X_*$  under the base change  $v$ . We therefore have a double Cartesian square:

$$\begin{array}{ccccc} X_1 & \xleftarrow{\quad} & V_1 & \xrightarrow{\quad} & U_1 \\ d_0 \boxtimes d_1 \downarrow & & v_0 \boxtimes v_1 \downarrow & & u_0 \boxtimes u_1 \downarrow \\ X_0 \times_Y X_0 & \xleftarrow{v \times v} & V \times_Y V & \xrightarrow{u \times u} & U \times_Y U \end{array}$$

Figure 33: The double Cartesian square used in the proof of Lemma 6.1(iii).

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 22 / combined-reader p. 284 / re-edited header p. 272.*

Since  $u_0 \boxtimes u_1$  is surjective, so is  $v_0 \boxtimes v_1$ . Since  $v \times v$  is surjective, the composite morphism

$$V_1 \longrightarrow X_0 \times_Y X_0$$

is surjective, and hence so is  $d_0 \boxtimes d_1$ .

**c) Proof of (i).**—It remains to prove that  $(Y, p)$  is a cokernel of  $(d_0, d_1)$  in the category of all ringed spaces. We first show that  $Y$  is obtained from  $X_0$  by identifying those points  $x$  and  $y$  for which there exists  $z \in X_1$  with

$$d_0(z) = x, \quad d_1(z) = y.$$

Indeed,  $p$  is surjective and  $pd_0 = pd_1$ . Conversely, if  $p(x) = p(y)$ , there is a point  $z' \in X_0 \times_Y X_0$  whose first projection is  $x$  and whose second projection is  $y$ . If  $z \in X_1$  satisfies

$$(d_0 \boxtimes d_1)(z) = z',$$

then  $d_0(z) = x$  and  $d_1(z) = y$ , as required.

On the other hand, if  $W$  is a saturated open subset of  $X_0$ , then  $W \cap U$  is a saturated open subset of  $U$ . By Theorem 4.1,  $q(W \cap U)$  is open in  $Y$ . Since  $q(W \cap U) = p(W)$ , the topology on  $Y$  is the quotient topology induced by  $X_0$ .

It remains to show that the canonical sequence of sheaves of rings

$$\mathcal{O}_Y \longrightarrow p_* \mathcal{O}_{X_0} \rightrightarrows p_* d_{0*} \mathcal{O}_{X_1} = p_* d_{1*} \mathcal{O}_{X_1} \quad (4)$$

is exact.

Let  $Y'$  be an open subset of  $Y$ , and put

$$U' = q^{-1}(Y'), \quad X'_0 = p^{-1}(Y'),$$

and so forth.<sup>33</sup> Then  $U'$  is an open subset of  $U$  saturated for the equivalence relation defined by the groupoid  $U_*$ . It follows from Lemmas 1.1 and 1.2 that  $Y'$  is the cokernel,

<sup>33</sup>Editor's note: The argument that follows has been expanded, in particular the verification that  $U'$  is a quasi-section of the groupoid induced on  $X'_0$ .

both in  $(\mathbf{Sch}/S)$  and in the category of ringed spaces, of the groupoid induced by  $U_*$  on  $U'$ . Likewise,  $X'_0$  is an open subset of  $X_0$  saturated for the equivalence relation defined by  $X_*$ , and we have the following commutative diagram, in which both squares are Cartesian:

$$\begin{array}{ccccc} X'_0 & \xleftarrow{\tilde{d}_1} & V' = d_0^{-1}(U') & \xrightarrow{\tilde{d}_0} & U' \\ & & \downarrow & & \downarrow \\ X_0 & \xleftarrow{d_1} & V = d_0^{-1}(U) & \xrightarrow{d_0} & U \end{array}$$

Figure 34: The restricted groupoids over  $X'_0$  and  $X_0$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 23 / combined-reader p. 285 / re-edited header p. 273; lower-right U corrected from source-PDF U'.*

Here  $\tilde{d}_1$  is surjective and finite locally free. On the other hand, let  $x \in U'$ . Since  $U$  is a quasi-section, the set

$$E = d_0 d_1^{-1}(x) \cap U$$

is finite and is contained in an affine open subset  $W$  of  $U$ . Then  $E' = E \cap U'$  is a finite set contained in the quasi-affine open subset  $W \cap U'$ . Consequently there is an affine open subset  $W'$  of  $W \cap U'$  that contains  $E'$ . This proves that  $U'$  is a quasi-section of the groupoid  $X'_*$  induced by  $X_*$  on  $X'_0$ . The first assertion of part (i), applied to  $X'_*$  and  $U'$ , now shows that  $Y'$  is the cokernel of  $X'_*$  in  $(\mathbf{Sch}/S)$ .

In particular, for every  $S$ -scheme  $T$ , the sequence

$$T(Y') \xrightarrow{T(p|_{X'_0})} T(X'_0) \xrightleftharpoons[T(d_0|_{X'_1})]{T(d_1|_{X'_1})} T(X'_1) \quad (5)$$

is exact. If  $T$  is the affine line  $\mathbf{G}_{a,S}$  (Exposé I, 4.3), this sequence identifies with

$$\begin{aligned} \Gamma(Y', \mathcal{O}_Y) &\longrightarrow \Gamma(p^{-1}(Y'), \mathcal{O}_{X_0}) \xrightleftharpoons[\delta_0]{\delta_1} \Gamma(d_0^{-1}p^{-1}(Y'), \mathcal{O}_{X_1}) \\ &= \Gamma(d_1^{-1}p^{-1}(Y'), \mathcal{O}_{X_1}), \end{aligned} \quad (6)$$

which is therefore exact for every open subset  $Y'$ . This completes the proof of Lemma 6.1(i).

**d) Proof of (iv).**—If  $(d_0, d_1)$  is an equivalence pair, then so is  $(u_0, u_1)$ . It follows from Theorem 4.1 that

$$u_0 \boxtimes u_1 : U_1 \longrightarrow U \times_Y U$$

is an isomorphism, and hence so is  $v_0 \boxtimes v_1$ ; see the Cartesian squares in part (b). Since  $v \times v$  is faithfully flat and quasi-compact,  $d_0 \boxtimes d_1$  is an isomorphism (SGA 1, VIII, 5.4).

Moreover, if  $d_0$  is flat, then  $u$  is flat. The morphism  $q$  is flat by Theorem 4.1, so  $r$  is flat as well. Since  $v$  is faithfully flat,  $p$  is flat; being also surjective, it is faithfully flat.

## 7. Quotient by a proper and flat groupoid

**Theorem 7.1.**—<sup>34</sup>Let  $S$  be a locally Noetherian scheme and let

<sup>34</sup>Editor's note: We mention here the article of S. Keel and S. Mori [KM97], which establishes the following theorem. Let  $X$  be an algebraic space of finite type over a locally Noetherian base  $S$ , and let  $j : R \rightarrow X \times_S X$



$$\begin{array}{ccccc}
 & d'_2 & & d_1 & \\
 X_2 & \xrightarrow{\quad d'_1 \quad} & X_1 & \xrightarrow{\quad d_1 \quad} & X_0 \\
 & d'_0 & & d_0 & 
 \end{array}$$

Figure 35: The groupoid occurring in Theorem 7.1.

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 24 / combined-reader p. 286 / re-edited header p. 274.*

be a **(Sch/S)**-groupoid such that  $d_1$  is proper and flat,  $X_0$  is quasi-projective over  $S$ ,<sup>35</sup> and the morphism

$$d : X_1 \longrightarrow X_0 \times_S X_0$$

with components  $d_0$  and  $d_1$  is quasi-finite. Then:

- (i) There exists a cokernel  $(Y, p)$  of  $(d_0, d_1)$  in **(Sch/S)**. Moreover, every such  $(Y, p)$  is a cokernel of  $(d_0, d_1)$  in the category of all ringed spaces.
- (ii) The morphism  $p$  is surjective, open, proper, and of finite presentation, and  $Y \rightarrow S$  is separated and of finite presentation.<sup>36</sup>
- (iii) The morphism

$$X_1 \longrightarrow X_0 \times_Y X_0$$

with components  $d_0$  and  $d_1$  is surjective.

- (iv) If  $(d_0, d_1)$  is an equivalence pair, then  $X_1 \rightarrow X_0 \times_Y X_0$  is an isomorphism and  $p$  is faithfully flat.<sup>37</sup> Moreover,  $(Y, p)$  is a cokernel of  $(d_0, d_1)$  in the category of sheaves for the fppf topology, and, for every base change  $Y' \rightarrow Y$ , the scheme  $Y'$  is the cokernel of the groupoid  $X_* \times_Y Y'$  obtained from  $X_*$  by the base change  $X_0 \times_Y Y' \rightarrow X_0$ .

In particular, for every base change  $S' \rightarrow S$ , the scheme

$$Y' = Y \times_S S'$$

is the cokernel of the  $S'$ -groupoid

$$X'_* = X_* \times_S S'.$$

Thus, in this case, “formation of the quotient commutes with base change”.

be a flat groupoid whose stabilizer  $j^{-1}(\Delta_X)$  is finite over  $X$ . There then exists an algebraic space that is a geometric quotient of  $X$  by  $R$  and a uniform categorical quotient; moreover, if  $j$  is separated, this quotient is separated. In particular, if a flat  $S$ -group scheme  $G$  acts properly on  $X$ , with finite stabilizer (that is, if

$$G \times_S X \longrightarrow X \times_S X, \quad (g, x) \longmapsto (x, g \cdot x)$$

is proper and the stabilizer of the diagonal is finite over  $X$ ), then there exists a geometric quotient  $X \rightarrow X/G$ . For a reductive  $S$ -group scheme  $G$ , this is a result of J. Kollár [Ko97].

<sup>35</sup>Editor’s note: This hypothesis on  $X_0$  is necessary; see editor’s note (17) to Theorem 4.1.

<sup>36</sup>Editor’s note: TDTE III, Theorem 6.1, states that  $Y \rightarrow S$  is quasi-projective if  $S$  is Noetherian. The editors did not see how to deduce this from the local existence of quasi-sections.

<sup>37</sup>Editor’s note: The consequences that follow have been made explicit; see [Ray67a], Theorem 1(iii), and the end of the proof of the theorem. We also mention that another proof of Theorem 7.1 in the case of an equivalence relation, based on the existence of Hilbert schemes, is given in [AK80], Theorem 2.9; see also [BLR90], §8.2, Theorem 12. In that case it is furthermore shown that  $Y \rightarrow S$  is quasi-projective.

Let  $(Y, p)$  be the cokernel of  $(d_0, d_1)$  in the category of all ringed spaces. Lemma 1.2 shows that, to prove part (i), it is enough to show that every point  $z \in X_0$  has a saturated open neighborhood  $U_z$  such that, if  $\tilde{d}_0$  and  $\tilde{d}_1$  denote the restrictions of  $d_0$  and  $d_1$  to

$$d_0^{-1}(U_z) = d_1^{-1}(U_z),$$

and if  $(Q, q)$  denotes the cokernel of  $(\tilde{d}_0, \tilde{d}_1)$  in the category of ringed spaces, then  $Q$  is a scheme and  $q$  is a morphism of schemes.

By Lemma 6.1(i), it is therefore enough to show that every point  $z \in X_0$  has a saturated open neighborhood  $U_z$  such that the groupoid induced by  $X_*$  on  $U_z$  has a quasi-section. We may even assume that  $z$  is closed in its fiber over  $S$  (we shall say that  $z$  is *closed relative to*  $S$ ).<sup>38</sup> The existence of  $U_z$  follows from the lemmas below.

**Lemma 7.2.**—*Let  $T$  be an affine Noetherian scheme, let  $X, Y, Z$  be  $T$ -schemes of finite type, with  $X$  quasi-projective over  $T$ , and let*

$$\begin{array}{ccc} Y & \xrightarrow{u} & X \\ v \downarrow & & \downarrow \\ Z & \dashrightarrow & T \end{array}$$

Figure 36: The diagram in Lemma 7.2.

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 25 / combined-reader p. 287 / re-edited header p. 275.*

*be a diagram in  $(\mathbf{Sch}/T)$ . Let  $z$  be a point of  $v(Y)$  that is closed relative to  $T$ , and suppose that  $v$  is flat at the points of  $v^{-1}(z)$ . Then there exists a closed subscheme  $F$  of  $X$  such that*

$$u(u^{-1}(F) \cap v^{-1}(z))$$

*is finite and nonempty, and such that the restriction of  $v$  to  $u^{-1}(F)$  is flat at the points of  $v^{-1}(z)$ .*

Put  $T = \operatorname{Spec} A$ . We may suppose that

$$X = \operatorname{Proj} \mathcal{S},$$

where  $\mathcal{S}$  is the symmetric algebra of a finitely generated  $A$ -module  $E$ .

If  $u(v^{-1}(z))$  is finite, we may take  $F = X$ . Otherwise, let

$$y_1, \dots, y_n$$

be the points of the fiber  $v^{-1}(z)$  associated with its structure sheaf  $\mathcal{O}_{v^{-1}(z)}$ . Thus, if  $\mathcal{O}_i$  denotes the local ring of  $v^{-1}(z)$  at  $y_i$ , the maximal ideal of  $\mathcal{O}_i$  consists of zero divisors. Let  $t$  be the image of  $z$  in  $T$ . The set  $u(v^{-1}(z))$  is an infinite constructible subset of the fiber of  $X$  over  $t$ . Consequently, that fiber contains a closed point  $x \in u(v^{-1}(z))$  distinct from

<sup>38</sup>Editor's note: Indeed, if such an open neighborhood  $U_z$  has been constructed for every point  $z$  that is closed relative to  $S$ , then the union of these  $U_z$  covers  $X_0$ : every fiber over  $S$  of the closed complement is a Noetherian scheme with no closed points, and is therefore empty.

$u(y_1), \dots, u(y_n)$ . The open subset  $X - \{x\}$  is a neighborhood of  $u(y_1), \dots, u(y_n)$ , and hence contains an open neighborhood of the form  $D_+(f)$ , where  $f \in \mathcal{S}$  is homogeneous of degree  $d$  (we use the notation of EGA II, §2.3).

The closed subscheme

$$X_1 = V_+(f)$$

defined by  $f$  therefore contains  $x$  and avoids the points  $u(y_1), \dots, u(y_n)$ . Its inverse image

$$Y_1 = u^{-1}(V_+(f))$$

is consequently distinct from  $Y$  and meets  $v^{-1}(z)$ . We shall in fact prove that the restriction  $v_1$  of  $v$  to  $Y_1$  is flat at the points of  $v^{-1}(z)$ . If  $u(v_1^{-1}(z))$  is finite, we may then take  $F = X_1$ . Otherwise, we repeat the argument with  $Y_1, v_1$ , and the morphism  $u_1$  induced by  $u$  on  $Y_1$ , in place of  $Y, v, u$ . In this way we obtain a descending sequence

$$X \supset X_1 \supset \dots$$

of closed subschemes of  $X$ . Since such a sequence terminates,

$$u(u^{-1}(X_n) \cap v^{-1}(z))$$

is finite and nonempty for some  $n$ , and we then take  $F = X_n$ .

It remains to show that  $v_1$  is flat at the points of  $v^{-1}(z)$ . Let  $y \in Y_1$  lie over  $z$ , let  $\mathcal{O}_y$  be the local ring of  $Y$  at  $y$ , let  $\overline{\mathcal{O}}_y$  be the local ring of  $v^{-1}(z)$  at  $y$ , and let  $\mathcal{O}_{v(y)}$  be the local ring of  $Z$  at  $v(y)$ . Choose  $g \in \mathcal{S}_1$  such that  $D_+(g)$  is a neighborhood of  $u(y)$  in  $X$ , and let  $\varphi$  and  $\overline{\varphi}$  be the respective images of  $f/g^d$  in  $\mathcal{O}_y$  and  $\overline{\mathcal{O}}_y$ . By the construction of  $f$ , the element  $\overline{\varphi}$  is not a zero divisor in  $\overline{\mathcal{O}}_y$ . Since  $\mathcal{O}_y$  is flat over  $\mathcal{O}_z = \mathcal{O}_{v(y)}$ , the element  $\varphi$  is not a zero divisor in  $\mathcal{O}_y$ , and

$$\mathcal{O}_y / \mathcal{O}_y \varphi$$

is flat over  $\mathcal{O}_z$  (SGA 1, IV, 5.7). This quotient is precisely the local ring of  $Y_1$  at  $y$ .

**Lemma 7.3.**—*Retain the notation and hypotheses of Theorem 7.1. Every point  $z \in X_0$  that is closed relative to  $S$  has a saturated open neighborhood  $U_z$  such that the groupoid induced by  $X_*$  on  $U_z$  has a quasi-section.*

The assertion is local on  $S$ , so we may suppose that  $S$  is affine and Noetherian and apply the preceding lemma to the diagram

$$\begin{array}{ccc} X_1 & \xrightarrow{d_0} & X_0 \\ d_1 \downarrow & & \downarrow \\ X_0 & \dashrightarrow & S \end{array}$$

Figure 37: The diagram to which Lemma 7.2 is applied.

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 26 / combined-reader p. 288 / re-edited header p. 276.*

in  $(\mathbf{Sch}/S)$ . Thus let  $F$  be a closed subscheme of  $X_0$  such that

$$d_0(d_0^{-1}(F) \cap d_1^{-1}(z))$$

is finite and nonempty, and such that the restriction of  $d_1$  to  $d_0^{-1}(F)$  is flat at the points of  $d_1^{-1}(z)$ .

Let  $F_1$  and  $F_2$  be the inverse images of  $F$  under  $d_0$  and  $d_0 d'_0 = d_0 d'_1$ , respectively, and let  $\tilde{d}_0, \tilde{d}_1, \dots$  denote the morphisms induced by  $d_0, d_1, \dots$ . We then have a commutative diagram

$$\begin{array}{ccccc}
 F_2 & \xrightarrow{\tilde{d}'_1} & F_1 & \xrightarrow{\tilde{d}_0} & F \\
 \downarrow \tilde{d}'_2 & & \downarrow \tilde{d}_1 & & \downarrow \tilde{q} \\
 X_1 & \xrightarrow[d_0]{d_1} & X_0 & \xrightarrow{q} & S
 \end{array}$$

Figure 38: The induced diagram on  $F_2, F_1, F$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 26 / combined-reader p. 288 / re-edited header p. 276.*

in which the two left-hand squares are Cartesian, the first row is exact (cf. the diagram (0, 1, 2) in §1), and  $q, \tilde{q}$  are the structure morphisms.

We first show that only finitely many points of  $F_1$  lie over  $z$ .<sup>39</sup> Let  $s$  be the image of  $z$  in  $S$ . Since  $F$  is of finite type over  $S$ , the fiber  $\tilde{q}^{-1}(s)$  is a Noetherian scheme. Since  $\tilde{d}_0$  is proper,

$$\tilde{d}_0(\tilde{d}_1^{-1}(z))$$

is a closed subscheme of  $\tilde{q}^{-1}(s)$  consisting of finitely many points. Consequently (cf. EGA I, 6.2.2), those points are closed in  $\tilde{q}^{-1}(s)$ , and also in the fiber  $q^{-1}(s)$  of  $X_0$ , because  $F$  is closed in  $X_0$ .

Let  $y$  be one of these points. Since  $q^{-1}(s)$  is of finite type over  $\kappa(s)$ , it has affine open neighborhoods

$$\operatorname{Spec} B \quad \text{and} \quad \operatorname{Spec} C$$

of  $y$  and  $z$ , respectively, with  $B$  and  $C$  finitely generated  $\kappa(s)$ -algebras. The points  $y$  and  $z$  correspond to maximal ideals

$$\mathfrak{p} \subset B, \quad \mathfrak{q} \subset C.$$

The fields  $B/\mathfrak{p}$  and  $C/\mathfrak{q}$  are finite over  $\kappa(s)$ , so

$$(B/\mathfrak{p}) \otimes_{\kappa(s)} (C/\mathfrak{q})$$

is a finite-dimensional  $\kappa(s)$ -algebra. Its maximal ideals correspond precisely to the points of  $X_0 \times_S X_0$  whose second projection is  $z$  and whose first projection is  $y$ . Thus there are only

<sup>39</sup>Editor's note: Details have been added, and the role of the properness hypothesis on  $d_0$  and  $d_1$  in Theorem 7.1 has been made explicit. Compare the statement and proof of Theorem 8.1, where this properness hypothesis is omitted.

finitely many points  $u \in X_0 \times_S X_0$  whose second projection is  $z$  and whose first projection belongs to

$$\tilde{d}_0(\tilde{d}_1^{-1}(z)).$$

Finally, since  $X_1 \rightarrow X_0 \times_S X_0$  has finite fibers, each such point  $u$  is the image of only finitely many points of  $X_1$ . This proves the assertion.

The morphism  $\tilde{d}_1$  is therefore quasi-finite and flat at the points of  $F_1$  lying over  $z$ . Since  $\tilde{d}_1$  is of finite type, SGA 1, IV, 6.10 and EGA III, 4.4.10 imply that the set  $\Phi$  of points of  $F_1$  at which  $\tilde{d}_1$  is not both flat and quasi-finite is closed in  $F_1$ , and hence in  $X_1$  because  $F_1$  is closed in  $X_1$ .<sup>40</sup> Since  $d_1$  is proper,  $\tilde{d}_1(\Phi)$  is closed and, by what precedes, does not contain  $z$ . Put

$$W = \tilde{d}_1(F_1) - \tilde{d}_1(\Phi).$$

The restriction of  $\tilde{d}_1$  to  $\tilde{d}_1^{-1}(W)$  is of finite presentation (by the Noetherian hypotheses), flat, proper, and quasi-finite.<sup>41</sup> It is therefore finite, locally free, and open, by EGA III, 4.4.2 and EGA IV<sub>2</sub>, 2.1.12 and 2.4.6. Consequently,  $\tilde{d}_1(F_1)$  is a neighborhood of  $z$ , and  $W$  is the largest open subset of  $X_0$  contained in  $\tilde{d}_1(F_1)$  over which  $\tilde{d}_1$  is both flat and quasi-finite.

We shall see in Lemma 7.4 that the inverse images of  $\Phi$  under  $\tilde{d}'_1$  and  $\tilde{d}'_0$  both identify with the set of points of  $F_2$  at which  $\tilde{d}'_2$  is not both flat and quasi-finite. It follows that

$$\begin{aligned} d_0^{-1}(W) &= \tilde{d}'_2(F_2) - \tilde{d}'_2((\tilde{d}'_0)^{-1}(\Phi)), \\ d_1^{-1}(W) &= \tilde{d}'_2(F_2) - \tilde{d}'_2((\tilde{d}'_1)^{-1}(\Phi)) \end{aligned}$$

coincide. Thus  $W$  is saturated.

Set

$$W_1 = \tilde{d}_1^{-1}(W).$$

The equality  $d_0^{-1}(W) = d_1^{-1}(W)$  implies

$$(\tilde{d}'_2)^{-1}(d_0^{-1}(W)) = (\tilde{d}'_2)^{-1}(d_1^{-1}(W)),$$

that is,

$$(\tilde{d}'_0)^{-1}(W_1) = (\tilde{d}'_1)^{-1}(W_1).$$

The morphism  $\tilde{d}_0$  is faithfully flat and quasi-compact, because  $d_0$ , like  $d_1$ , is surjective, proper, and flat. Moreover, the square

$$\begin{array}{ccc} F_2 & \xrightarrow{\tilde{d}'_1} & F_1 \\ \tilde{d}'_0 \downarrow & & \downarrow \tilde{d}_0 \\ F_1 & \xrightarrow{\tilde{d}_0} & F \end{array}$$

Figure 39: The Cartesian square used to descend  $W_1$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 27 / combined-reader p. 289 / re-edited header p. 277.*

<sup>40</sup>Editor's note: If  $f : X \rightarrow Y$  is locally of finite type, the set of points  $x \in X$  that are isolated in their fiber  $f^{-1}(f(x))$  is open in  $X$ . In EGA III, 4.4.10, this is deduced from Zariski's Main Theorem when  $Y$  is locally Noetherian. For arbitrary  $Y$ , it follows from Chevalley's semicontinuity theorem (EGA IV<sub>3</sub>, 13.1.3 and 13.1.4). Consequently,  $f$  is quasi-finite at  $x$  if and only if it is of finite type at  $x$  and  $x$  is isolated in  $f^{-1}(f(x))$ . This will be used below; see editor's note (42).

<sup>41</sup>Editor's note: The original has been expanded in what follows.

is Cartesian. It follows that

$$W_1 = \tilde{d}_0^{-1}(U)$$

for an open subset  $U$  of  $F$  (SGA 1, VIII, 4.4). This open subset  $U$  is a quasi-section of the groupoid with base  $W$  induced by  $X_*$ . We may therefore take  $U_z = W$ .

It remains to state the following lemma, whose proof is classical.

**Lemma 7.4.**—*Consider a Cartesian square of schemes*

$$\begin{array}{ccc} F_2 & \xrightarrow{v} & F_1 \\ d' \downarrow & & \downarrow d \\ X_1 & \xrightarrow{u} & X_0 \end{array}$$

Figure 40: The Cartesian square in Lemma 7.4.

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 27 / combined-reader p. 289 / re-edited header p. 277.*

and let  $x$  be a point of  $F_2$ .

- (i) If  $u$  is flat,  $d'$  is flat at  $x$  if and only if  $d$  is flat at  $v(x)$ .
- (ii) If  $d$  is locally of finite type,  $d'$  is quasi-finite at  $x$  if and only if  $d$  is quasi-finite at  $v(x)$ .<sup>42</sup>


We have thus proved that  $X_0$  has a covering by saturated open subsets  $W$  such that the groupoid  $W_*$  induced by  $X_*$  on  $W$  possesses a quasi-section.<sup>43</sup>

By Lemma 6.1 and the reductions stated after Theorem 7.1, this proves assertions (i) and (iii) of that theorem, as well as the facts that  $p$  is surjective and open and that  $p$  and  $Y \rightarrow S$  are locally of finite presentation. Moreover, because  $X_0 \rightarrow S$  is quasi-projective, and hence separated and of finite type,  $p$  is separated; the proof of Lemma 6.1(ii) shows that  $p$  and  $Y \rightarrow S$  are of finite presentation.

To prove that  $p$  is proper, it remains to prove that it is universally closed. Since this assertion is local on  $Y$ , we may work over a saturated open subset  $W$  such that the groupoid  $W_*$  induced by  $X_*$  on  $W$  possesses a quasi-section  $U$  (because  $X_0$  is covered by such open subsets). With the notation of 6(a), there is a commutative diagram

<sup>42</sup>Editor's note: The conditions are sufficient by base change (EGA II, 6.2.4(iii), and EGA IV<sub>2</sub>, 2.1.4). Conversely, put  $y = d'(x)$  and  $z = u(y) = d(v(x))$ , and suppose that  $d'$  is flat at  $x$  and that  $u$  (hence also  $v$ ) is flat. Then  $\mathcal{O}_{v(x)} \rightarrow \mathcal{O}_x$  is faithfully flat, as is  $\mathcal{O}_z \rightarrow \mathcal{O}_y \rightarrow \mathcal{O}_x$ . Consequently,  $\mathcal{O}_z \rightarrow \mathcal{O}_{v(x)}$  is faithfully flat (EGA IV<sub>2</sub>, 2.2.11(iv)). Finally, suppose that  $d$  is locally of finite type and that  $d'$  is quasi-finite at  $x$ . Then  $v(x)$  is isolated in its fiber  $d^{-1}(z)$ , since  $x$  is isolated in its fiber  $d'^{-1}(y) = d^{-1}(z) \otimes_{\kappa(z)} \kappa(y)$ . By Chevalley's semicontinuity theorem there is therefore an open neighborhood of  $v(x)$  every point of which is isolated in its fiber (EGA IV<sub>3</sub>, 13.1.3 and 13.1.4), and hence  $d$  is quasi-finite at  $v(x)$ .

<sup>43</sup>Editor's note: The sequel has been modified to take advantage of the additions made to Lemma 6.1.

Figure 41: The properness diagram over the open subset  $Z \subseteq Y$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 28 / combined-reader p. 290 / re-edited header p. 278.*

where  $Z$  is an open subset of  $Y$ , all the arrows are surjective, and  $q$  is integral. By hypothesis  $d_0 : X_1 \rightarrow X_0$  is proper, so  $u$ , which is obtained from  $d_0$  by base change, is proper as well. Consequently  $r$  is universally closed, and hence so is  $p$ , since  $v$  is surjective.

Finally, because  $p$  is surjective and universally closed and  $X_0$  is quasi-projective, hence separated, the diagonal  $\Delta_{Y/S}(Y)$  is closed in  $Y \times_S Y$ : it is the image under  $p \times p$  of the diagonal  $\Delta_{X_0/S}(X_0)$ . Thus  $Y$  is separated over  $S$ . This completes the proof of Theorem 7.1(ii).

The assertions in Theorem 7.1(iv) are local on  $Y$ . Since the saturated open subsets  $U_z$  cover  $X_0$ , it is enough to verify them after replacing  $X$  and  $Y$  by  $U_z$  and  $V_z = p(U_z)$ , respectively. As observed at the beginning of the proof of Theorem 7.1, Lemmas 1.1, 1.2, and 6.1(i) show that  $(V_z, p|_{U_z})$  is the cokernel, both in **(Sch)** and in the category of ringed spaces, of the groupoid induced by  $X_*$  on  $U_z$ . The hypothesis that  $d = (d_0, d_1)$  is a monomorphism is preserved under the base change  $U_z \rightarrow X_0$ . The first two assertions of Theorem 7.1(iv) therefore follow from Lemma 6.1(iv).

Let us finally prove the consequences stated at the end of part (iv) (see [Ray67a], Theorem 1(iii)). By hypothesis, the groupoid  $X_*$  comes from an equivalence relation  $R \rightarrow X_0 \times X_0$ . We have proved that  $R$  is effective (Exposé IV, 3.3.2) and that  $p : X_0 \rightarrow Y = X_0/R$  is faithfully flat and of finite presentation. Consequently, if  $(M)$  denotes the family of faithfully flat morphisms locally of finite presentation, then  $R$  is  $(M)$ -effective. By Exposé IV, 6.3.3, the pair  $(Y, p)$  therefore represents the quotient sheaf of  $X_0$  by  $R$  for the fppf topology, and the assertions concerning base change follow from Exposé IV, 3.4.3.1.

## 8. Passage to the quotient by a flat groupoid that is not necessarily proper

**Theorem 8.1.**—<sup>44</sup>Let  $S$  be a Noetherian scheme and

<sup>44</sup>Editor's note: There is a largest open subset  $W$  of  $X_0$  satisfying the conclusions of the theorem. Indeed, let  $W$  be an open subset as in the theorem and let  $W^\sharp$  be a dense saturated open subset of  $W$ . Since  $p$  is open,  $V^\sharp = p(W^\sharp)$  is open in  $V$ , and  $W^\sharp = p^{-1}(V^\sharp)$ , because  $W^\sharp$  is saturated. By Lemma 1.1,  $V^\sharp$  is a cokernel for the groupoid induced on  $W^\sharp$ . Thus two open subsets  $W$  and  $W'$  satisfying the conclusions of the theorem may be glued along their intersection  $W^\sharp$ , and conditions (i)–(iv), together with the property that  $p$  and  $V \rightarrow S$  are locally of finite presentation, are preserved. Conclusion (ii') follows, as in the proof of Lemma 6.1(ii), from the hypothesis that  $X_0$  is of finite type over the Noetherian scheme  $S$ .

Lemmas 1.1 and 1.2 also show that the union of all saturated open subsets  $U \subseteq X_0$  such that the open subset  $p(U) \subseteq Y$  is a scheme and  $p|_U : U \rightarrow p(U)$  is a morphism of schemes is the largest saturated open subset  $\Omega \subseteq X_0$  satisfying condition (i) of Theorem 8.1. The theorem shows that  $\Omega$  contains a dense open subset  $W$ , but it is not immediate that  $\Omega$  satisfies properties (ii)–(iv). For more precise results, and for the representability of the fppf quotient sheaf  $\widetilde{X/R}$  (where  $R$  denotes the groupoid  $X_*$  with base  $X = X_0$ ), see [Ray67a], [Ray67b], and Appendix I of [An73]. These works use weaker hypotheses:  $S$  may be arbitrary,  $X$  is locally of finite type over  $S$ , and  $R$  is an  $S$ -groupoid with base  $X$  such that  $d_0$  (and hence  $d_1$ ) is flat and

$$\begin{array}{ccccc}
& & d'_2 & & \\
& & \xrightarrow{\quad} & & \\
X_2 & \xrightarrow{\quad d'_1 \quad} & X_1 & \xrightarrow{\quad d_1 \quad} & X_0 \\
& \xrightarrow{\quad} & & \xrightarrow{\quad d_0 \quad} & \\
& & d'_0 & & 
\end{array}$$

Figure 42: The  $(\mathbf{Sch}/S)$ -groupoid of Theorem 8.1.

Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 29 / combined-reader p. 291 / re-edited header p. 279.

a  $(\mathbf{Sch}/S)$ -groupoid such that  $d_1$  is flat and of finite type,  $X_0$  is of finite type over  $S$ , and the morphism  $X_1 \rightarrow X_0 \times_S X_0$  with components  $d_0$  and  $d_1$  is quasi-finite.

There exists a dense saturated open subset  $W \subseteq X_0$  having the following properties:

- (i) if  $W_2 \xrightarrow{w'_i} W_1 \xrightarrow{w_j} W$  is the groupoid induced by  $X_*$  on  $W$ , then  $(w_0, w_1)$  has a cokernel  $(V, p)$  in  $(\mathbf{Sch}/S)$ ; moreover,  $(V, p)$  is a cokernel of  $(w_0, w_1)$  in the category of all ringed spaces;
- (ii)  $p$  is surjective and open;
- (ii')  $p$  and  $V \rightarrow S$  are of finite presentation;
- (iii) the morphism  $W_1 \rightarrow W \times_V W$  with components  $w_0$  and  $w_1$  is surjective;
- (iv) if  $(d_0, d_1)$  is an equivalence pair, then  $W_1 \rightarrow W \times_V W$  is an isomorphism and  $p$  is faithfully flat.

We shall prove that  $W$  may be chosen so that the  $(\mathbf{Sch}/S)$ -groupoid  $W_*$  induced by  $X_*$  possesses a quasi-section (see §7). Theorem 8.1 will then follow from Lemma 6.1.

Suppose provisionally that for every point  $z \in X_0$  closed relative to  $S$  (see §7) there is a saturated open subset  $W_z$  that possesses a quasi-section and meets every irreducible component of  $X_0$  passing through  $z$ . Then the complement  $X_0 - W_z$  is saturated: its saturation  $d_1(d_0^{-1}(X_0 - W_z))$  is open and does not meet  $W_z$ . If this complement is nonempty, choose in it a point  $z'$  closed relative to  $S$ , and associate to  $z'$  an open subset  $W_{z'}$  as above. We may further suppose that  $W_{z'} \subseteq X_0 - W_z$ . Then  $W_z$  and  $W_{z'}$  are disjoint, and the groupoid induced by  $X_*$  on  $W_z \cup W_{z'}$  possesses a quasi-section. The process must stop, because  $X_0$  has only finitely many irreducible components. It therefore remains to construct  $W_z$ .

For this purpose we may suppose that  $S$  is affine. Let  $y \in X_1$  satisfy  $d_1(y) = z$ , let  $X$  be an affine open subset of  $X_0$  containing  $d_0(y)$ , let  $Y$  be the inverse image of  $X$  under  $d_0 : X_1 \rightarrow X_0$ , and let  $u : Y \rightarrow X$  and  $v : Y \rightarrow X_0$  be the morphisms induced by  $d_0$  and  $d_1$ . Since  $X$  is affine, hence quasi-projective, Lemma 7.2 provides a subscheme  $F \subseteq X_0$  such that  $d_0^{-1}(F) \cap d_1^{-1}(z)$  is nonempty,  $d_0(d_0^{-1}(F) \cap d_1^{-1}(z))$  is finite, and the restriction of  $d_1$  to  $d_0^{-1}(F)$  is flat at the points of  $v^{-1}(z)$ . We can therefore resume the notation of Lemma 7.3, writing  $F_1$  and  $F_2$  for the inverse images of  $F$  in  $X_1$  and  $X_2$ , and so on:

---

of finite presentation. In particular, if  $\widetilde{X/R}$  is represented by an  $S$ -scheme  $Y$ , then  $Y$  is also the cokernel in the category of ringed spaces. The converse is false in general (see [Mum65], Chapter 0, §3, Example 0.4, cited in [Ray67a], Remark 1), but it is true when  $d = (d_0, d_1)$  is an immersion. Under this hypothesis,  $p : \Omega \rightarrow Z := \Omega/R_\Omega$  is faithfully flat and of finite presentation. If, in addition,  $S$  is locally Noetherian, then a codimension-one point  $x$  of  $X$  belongs to  $\Omega$  if and only if the graph of the groupoid induced on  $\text{Spec}(\mathcal{O}_{X,x})$  is closed. See [Ray67a], Proposition 1; [Ray67b], Proposition 1 and Theorems 2, 1, and 4; and [An73], Theorems 5 and 6 on pages 66–67 and Proposition 3.3.1 on page 49. For an action of an algebraic group over an algebraically closed field  $k$ , see also [DR81].



$$\begin{array}{ccccc}
F_2 & \xrightarrow{\tilde{d}'_1} & F_1 & \xrightarrow{\tilde{d}_0} & F \\
& \searrow \tilde{d}'_0 & \downarrow \tilde{d}_1 & & \\
& & X_0 & & \\
\tilde{d}_2 \downarrow & & \uparrow d_1 & & \\
X_1 & \xrightarrow{\quad} & X_0 & \xrightarrow{\quad} & \\
& \searrow d_0 & & & 
\end{array}$$

Figure 43: The induced diagram on  $F_2, F_1, F$  used in the proof of Theorem 8.1.

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 30 / combined-reader p. 292 / re-edited header p. 280.*

As in 7.3, one proves that  $\tilde{d}_1$  is quasi-finite at the points of  $\tilde{d}_1^{-1}(z)$ . It is therefore natural to consider the open subset  $F'_1 \subseteq F_1$  consisting of the points at which  $\tilde{d}_1$  is both flat and quasi-finite. By Lemma 7.4, the inverse images of  $F'_1$  under  $\tilde{d}'_1$  and  $\tilde{d}'_0$  both consist of the points of  $F_2$  at which  $\tilde{d}'_2$  is flat and quasi-finite. Thus these inverse images coincide, and  $F'_1$  is of the form  $\tilde{d}_0^{-1}(F')$  for an open subset  $F' \subseteq F$  (SGA 1, VIII, 4.4). After replacing  $F$  by  $F'$ , we may therefore suppose that  $\tilde{d}_1$  is quasi-finite and flat. Let  $W_z$  be the largest open subset of  $\tilde{d}_1(F_1)$  over which  $\tilde{d}_1$  is finite and flat.

The open subset  $W_z$  need not contain  $z$ , but it contains the generic points of the irreducible components of  $X_0$  passing through  $z$ .<sup>45</sup> Since  $d_0$  (and likewise  $d_1$ ) is faithfully flat and of finite presentation, hence open, SGA 1, VIII, 5.7 shows that  $d_0^{-1}(W_z)$  and  $d_1^{-1}(W_z)$  both coincide with the largest open subset of  $\tilde{d}'_2(F_2)$  over which  $\tilde{d}'_2$  is finite and flat. As in 7.3, the inverse images of  $\tilde{d}_1^{-1}(W_z)$  under  $\tilde{d}'_0$  and  $\tilde{d}'_1$  therefore coincide. Hence  $\tilde{d}_1^{-1}(W_z) = \tilde{d}_0^{-1}(U)$  for an open subset  $U \subseteq F$ , and  $U$  is a quasi-section for the groupoid induced by  $X_*$  on  $W_z$ .

## 9. Elimination of the Noetherian hypotheses in Theorem 7.1

a) Resume the notation and hypotheses of Lemma 6.1, and let  $\pi : S' \rightarrow S$  be an arbitrary base change. If  $f : X \rightarrow Y$  is a morphism of  $S$ -schemes, write  $f' : X' \rightarrow Y'$  for the morphism of  $S'$ -schemes obtained from  $f$  by extension of the base along  $\pi$ . With this convention,  $p' : X'_0 \rightarrow Y'$  is surjective, as is the morphism

$$X'_1 \longrightarrow X'_0 \times_{Y'} X'_0$$

with components  $d'_0$  and  $d'_1$ . Thus the set underlying  $Y'$  is the quotient of the set underlying  $X'_0$  by the equivalence relation on  $X'_0$  defined by the  $S'$ -groupoid  $X'_*$ . Moreover,  $q' : U' \rightarrow Y'$  is integral and surjective. The topology of  $Y'$  is consequently the quotient topology of that of  $U'$ , and hence also of that of  $X'_0$ ; see the proof in §6(c).

On the other hand,  $U'$  is plainly a quasi-section of the  $S'$ -groupoid  $X'_*$ , so Lemma 6.1 applies. In particular,  $X'_*$  has a cokernel  $(Y_1, p_1)$ , and the topological space underlying  $Y_1$  is obtained from that underlying  $X'_0$  by identifying the points equivalent under the relation

<sup>45</sup>Editor's note: Indeed, let  $\eta$  be such a generic point. The hypotheses imply that  $\mathcal{O}_{X_0, \eta}$  is an Artinian local ring and that  $\mathcal{O}_{F_1, \eta}$  is a finitely generated  $\mathcal{O}_{X_0, \eta}$ -module. By SGA 1, VIII, 6.5, there is therefore an open neighborhood of  $\eta$  over which  $\tilde{d}_1$  is finite.

defined by  $X'_*$ . It follows that the canonical morphism  $Y_1 \rightarrow Y'$  is a homeomorphism. In fact, it is a universal homeomorphism. Indeed, if  $S''$  is an  $S'$ -scheme, let  $Y_2$  be the cokernel of  $(d_0 \times_S S'', d_1 \times_S S'')$ . The preceding argument, applied to the base changes  $S'' \rightarrow S'$  and  $S'' \rightarrow S$ , shows that

$$Y_2 \longrightarrow Y_1 \times_{S'} S'' \quad \text{and} \quad Y_2 \longrightarrow Y \times_S S'' \simeq Y' \times_{S'} S''$$

are homeomorphisms. Hence

$$Y_1 \times_{S'} S'' \longrightarrow Y' \times_{S'} S''$$

is a homeomorphism as well.

**b)** The same remarks plainly apply when the groupoid  $X_*$  *locally* possesses quasi-sections (see the proof of Theorem 7.1).<sup>46</sup> For example, we have the following theorem:

**Theorem 9.0.**—*Let  $S$  be an arbitrary scheme and*

$$X_2 \xrightarrow{d'_j} X_1 \xrightarrow{d_i} X_0$$

*a **(Sch/S)**-groupoid such that  $X_0$  is of finite presentation and quasi-projective over  $S$ ,  $d_1$  is of finite presentation, proper, and flat, and the morphism*

$$d_0 \boxtimes d_1 : X_1 \longrightarrow X_0 \times_S X_0$$

*is quasi-finite. Then:*

- (1) *Every point  $x$  of  $X_0$  has a saturated open neighborhood  $W$  such that the groupoid induced by  $X_*$  on  $W$  has a quasi-section.*
- (2) *Let  $(Y, p)$  be the cokernel of  $(d_0, d_1)$  in the category of all ringed spaces. Then  $Y$  is a scheme,  $p$  is a morphism of schemes, and  $(Y, p)$  is a cokernel of  $(d_0, d_1)$  in **(Sch/S)**.*
- (3) *The morphism  $p$  is surjective, open, and universally closed.*
- (4) *The morphism*

$$d : X_1 \longrightarrow X_0 \times_Y X_0$$

*with components  $d_0$  and  $d_1$  is surjective.*

- (5) *If  $(d_0, d_1)$  is an equivalence pair, then:*

- (a) *the morphism*

$$d : X_1 \longrightarrow X_0 \times_Y X_0$$

*is an isomorphism and  $p$  is faithfully flat;*

- (b) *the morphisms  $p$  and  $Y \rightarrow S$  are of finite presentation, and  $(Y, p)$  is a cokernel of  $(d_0, d_1)$  in the category of sheaves for the fppf topology.*

*Proof.* For (1), the question is local on  $S$ , so we may suppose that  $S = \text{Spec } B$  is affine. There then exist a ring  $A$  of finite type over  $\mathbb{Z}$ , a morphism

$$S \longrightarrow T = \text{Spec } A,$$

---

<sup>46</sup>Editor's note: What follows has been expanded in order to bring out Theorem 9.0 below.

and a  $(\mathbf{Sch}/T)$ -groupoid  $Z_*$  such that

$$X_* \simeq Z_* \times_T S.$$

Indeed, apply EGA IV<sub>3</sub>, 8.8.3 to  $S_0 = \operatorname{Spec} \mathbb{Z}$  and  $S_i = \operatorname{Spec} A_i$ , where the  $A_i$  range over the finitely generated  $\mathbb{Z}$ -subalgebras of  $B$ . Moreover, by EGA IV<sub>3</sub>, 8.10.5, we may suppose that  $Z_*$  satisfies the hypotheses of Theorem 7.1. Consequently,  $Z_*$  has quasi-sections locally.

By part (a), the same is therefore true of  $X_*$ , and assertions (2), (3), (4), and (5)(a) follow from Lemma 6.1, just as in the proof of Theorem 7.1.

**c) Proof of (5)(b).**—Let us show that  $Y \rightarrow S$  is of finite presentation.<sup>47</sup> By hypothesis,  $(d_0^{X_*}, d_1^{X_*})$  is an equivalence pair; that is,

$$d^{X_*} : X_1 \longrightarrow X_0 \times_S X_0$$

is a monomorphism. By EGA IV<sub>3</sub>, 8.10.5, after enlarging  $A$  if necessary, we may suppose that

$$d^{Z_*} : Z_1 \longrightarrow Z_0 \times_T Z_0$$

is a monomorphism. Since  $T = \operatorname{Spec} A$ , with  $A$  Noetherian, Theorem 7.1 then shows that the groupoid  $Z_*$  has a cokernel  $(Q, q)$  in  $(\mathbf{Sch}/T)$ , that  $q$  and  $Q \rightarrow T$  are of finite presentation, and, moreover, that

$$q : Z_0 \longrightarrow Q$$

is faithfully flat and  $d^{Z_*}$  induces an isomorphism

$$Z_1 \xrightarrow{\sim} Z_0 \times_Q Z_0.$$

Put  $Q_S = Q \times_T S$ . Since  $X_i \simeq Z_i \times_T S$ , we obtain an isomorphism

$$d^{Z_*} \times_T S : X_1 \xrightarrow{\sim} X_0 \times_{Q_S} X_0.$$

Denote its inverse by  $\phi$ , and let  $\pi$  be the canonical morphism

$$X_0 \times_Y X_0 \longrightarrow X_0 \times_{Q_S} X_0.$$

Then  $\phi \circ \pi$  is the inverse of

$$d_0 \boxtimes d_1 : X_1 \xrightarrow{\sim} X_0 \times_Y X_0.$$

It follows that the equivalence relation defined by  $X_*$ , that is, the monomorphism shown in Figure 44,

$$X_1 \xrightarrow[\sim]{d_0 \boxtimes d_1} X_0 \times_Y X_0 \hookrightarrow X_0 \times_S X_0$$

Figure 44: The equivalence-relation monomorphism defined by  $X_*$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 33 / combined-reader p. 295 / re-edited header p. 283.*

identifies with the equivalence relation  $\mathcal{R}(q_S)$  defined by  $q_S : X_0 \rightarrow Q_S$ . Since  $q_S$  is faithfully flat and of finite presentation, hence a universal effective epimorphism,  $\mathcal{R}(q_S)$  has quotient  $Q_S$ ; see Exposé IV, 3.3.2. Consequently,

$$Y \simeq Q \times_T S,$$

<sup>47</sup>Editor's note: The original states that this follows from Proposition 9.1 below. We were unable to reconstruct that argument. The proof that follows was communicated to us by O. Gabber.

so  $Y \rightarrow S$  and  $p$  are of finite presentation. Moreover, by Exposé IV, 6.3.3,  $(Y, p)$  is also a cokernel of  $(d_0, d_1)$  in the category of sheaves for the fppf topology.

**Proposition 9.1.**—*Consider morphisms of schemes*

$$X_0 \xrightarrow{p} Y \xrightarrow{q} S$$

*such that  $qp$  is of finite type (respectively, of finite presentation) and  $p$  is faithfully flat and of finite presentation. Then  $q$  is of finite type (respectively, of finite presentation).\**

Since  $p$  is surjective and  $qp$  is quasi-compact,  $q$  is quasi-compact. We may therefore suppose that  $S$ ,  $Y$ , and  $X_0$  are affine, with rings  $A$ ,  $B$ , and  $C$ , respectively. We have

$$B = \varinjlim_i B_i,$$

where the  $B_i$  range over the finitely generated  $A$ -subalgebras of  $B$ . Since  $C$  is of finite presentation over  $B$ , there exist an index  $i_0$ , a  $B_{i_0}$ -algebra  $C_{i_0}$  of finite presentation, and an isomorphism

$$C \simeq C_{i_0} \otimes_{B_{i_0}} B.$$

For  $i \geq i_0$ , put

$$C_i = C_{i_0} \otimes_{B_{i_0}} B_i;$$

then

$$C \simeq C_i \otimes_{B_i} B.$$

$$\begin{array}{ccc} B & \longrightarrow & C \\ \uparrow & & \uparrow \\ B_i & \longrightarrow & C_i \\ \uparrow & & \\ A & & \end{array}$$

Figure 45: The compatible  $A$ -algebras used in the limit argument.

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 33 / combined-reader p. 295 / re-edited header p. 283.*

Since  $C$  is faithfully flat over  $B$ , EGA IV<sub>3</sub>, 11.2.6 and 8.10.5(vi) give an index  $i_1 \geq i_0$  such that  $C_{i_1}$  is faithfully flat over  $B_{i_1}$ . Consequently,  $C_i$  is faithfully flat over  $B_i$  for every  $i \geq i_1$ . For  $i \geq i_1$ , the canonical map  $C_i \rightarrow C$  is then injective, because it is obtained from  $B_i \rightarrow B$  by faithfully flat base extension.

If  $C$  is of finite type over  $A$ , it follows that there is an index  $j \geq i_1$  such that  $C_j = C$ , whence  $B_j = B$ , because  $C_j$  is faithfully flat over  $B_j$ . Thus  $B$  is of finite type over  $A$ .

Now suppose that  $C$  is of finite presentation over  $A$ . By what precedes,  $B$  is of finite type over  $A$ , and hence has the form

$$B = \overline{B}/I,$$

---

\*See EGA IV<sub>4</sub>, 17.7.5 for a more general result.

where  $\overline{B}$  is a polynomial algebra over  $A$  in finitely many indeterminates and  $I$  is an ideal of  $\overline{B}$ . The ideal  $I$  is the union of its finitely generated subideals  $I_\alpha$ ; consequently,

$$B = \varinjlim_{\alpha} B_{\alpha}, \quad B_{\alpha} = \overline{B}/I_{\alpha}.$$

Proceeding as above, there exist an index  $\alpha_0$ , a  $B_{\alpha_0}$ -algebra  $C_{\alpha_0}$  of finite presentation, and an isomorphism

$$C \simeq C_{\alpha_0} \otimes_{B_{\alpha_0}} B.$$

For  $\alpha \geq \alpha_0$ , put

$$C_{\alpha} = C_{\alpha_0} \otimes_{B_{\alpha_0}} B_{\alpha},$$

so that

$$C \simeq C_{\alpha} \otimes_{B_{\alpha}} B \quad (\alpha \geq \alpha_0).$$

Again by EGA IV<sub>3</sub>, 11.2.6 and 8.10.5(vi),  $C_{\alpha}$  is faithfully flat over  $B_{\alpha}$  for all sufficiently large  $\alpha$ . In that case, the kernels of the maps  $C_{\alpha} \rightarrow C$  and  $C_{\alpha} \rightarrow C_{\beta}$ , respectively, identify with

$$C_{\alpha} \otimes_{B_{\alpha}} (I/I_{\alpha}) \quad \text{and} \quad C_{\alpha} \otimes_{B_{\alpha}} (I_{\beta}/I_{\alpha}) \quad (\beta \geq \alpha).$$

Since  $C_{\alpha}$  and  $C$  are of finite presentation over  $A$  and  $C_{\alpha} \rightarrow C$  is surjective,

$$C_{\alpha} \otimes_{B_{\alpha}} (I/I_{\alpha})$$

is a finitely generated ideal,<sup>48</sup> and it is the union of the ideals

$$C_{\alpha} \otimes_{B_{\alpha}} (I_{\beta}/I_{\alpha}).$$

Thus, for all sufficiently large  $\beta$ ,

$$C_{\alpha} \otimes_{B_{\alpha}} (I_{\beta}/I_{\alpha}) = C_{\alpha} \otimes_{B_{\alpha}} (I/I_{\alpha}).$$

Since  $C_{\alpha}$  is faithfully flat over  $B_{\alpha}$ , this also gives  $I_{\beta} = I$ . Therefore  $B$  is of finite presentation over  $A$ .

## 10. Supplement: quotients by a group scheme

Sections 10.2–10.4 below, written following suggestions of M. Raynaud, apply the preceding theorems to the case of an action of a group scheme. For the reader's convenience, Section 10.1 first reproduces statements 2.1–2.3 of Exposé XVI.

**10.1. Representability theorems for quotients.**—We first recall the following result.

**Theorem 10.1.1.**—*Let  $S$  be a scheme, let  $X$  and  $Y$  be two  $S$ -schemes, and let  $f : X \rightarrow Y$  be an  $S$ -morphism. Suppose that one of the following two conditions holds:*

- $\alpha$ ) the morphism  $f$  is locally of finite presentation;*
- $\beta$ ) the scheme  $S$  is locally Noetherian and  $X$  is locally of finite type over  $S$ .*

*Then the following conditions are equivalent:*

---

<sup>48</sup>Editor's note: see EGA IV<sub>1</sub>, 1.4.4.

(i) *there exist an  $S$ -scheme  $X'$  and a factorization*

$$f : X \xrightarrow{f'} X' \xrightarrow{f''} Y,$$

*where  $f'$  is a faithfully flat  $S$ -morphism locally of finite presentation and  $f''$  is a monomorphism;*

(ii) *the first projection*

$$p_1 : X \times_Y X \longrightarrow X$$

*is flat.*

Moreover, if these conditions hold, then  $(X', f')$  is a quotient of  $X$  by the equivalence relation defined by  $f$  for the fppf topology. Thus the factorization  $f = f'' \circ f'$  in (i) is unique up to isomorphism.

The case in which  $Y$  is locally Noetherian and  $X$  is of finite type over  $Y$  is treated in [Mur65], Corollary 2 to Theorem 2. We shall see that the general case reduces to that one.

We first make a few remarks.

a) The implication (i) $\Rightarrow$ (ii) is immediate. Indeed, the first projection

$$p'_1 : X \times_{X'} X \longrightarrow X$$

factors through  $X \times_Y X$ :

$$p'_1 : X \times_{X'} X \xrightarrow{u} X \times_Y X \xrightarrow{p_1} X.$$

The morphism  $u$  is an isomorphism because  $f''$  is a monomorphism, and  $p'_1$  is flat because  $f'$  is flat; hence  $p_1$  is flat.

b) The assertions of Theorem 10.1.1 are local on  $Y$ , and hence local on  $S$ . They are also local on  $X$ , as follows readily from the fact that a flat morphism locally of finite presentation is open (EGA IV<sub>3</sub>, 11.3.1).

c) Under hypothesis  $\alpha$  of Theorem 10.1.1, the preceding remarks reduce us to the case where  $X$  and  $Y$  are affine and  $f$  is of finite presentation. Replacing  $S$  by  $Y$ , if necessary, we may suppose that  $X$  and  $Y$  are of finite presentation over  $S$ . EGA IV<sub>3</sub>, 11.2.6 then reduces us to the case where  $S$  is Noetherian.

d) Under hypothesis  $\beta$  of Theorem 10.1.1, we may suppose that  $S$ ,  $X$ , and  $Y$  are affine, that  $S$  is Noetherian, and that  $X$  is of finite type over  $S$ . Write  $Y$  as a filtered inverse limit of affine schemes  $Y_i$  of finite type over  $S$ . The schemes

$$X \times_{Y_i} X$$

form a filtered decreasing family of closed subschemes of  $X \times_S X$ , whose inverse limit is  $X \times_Y X$ . Since  $X \times_S X$  is Noetherian, for all sufficiently large  $i$  we have

$$X \times_{Y_i} X = X \times_Y X.$$

It follows that

$$f_i : X \xrightarrow{f} Y \longrightarrow Y_i$$

satisfies condition (ii) of Theorem 10.1.1 whenever  $f$  does. The equivalence relation on  $X$  defined by  $f$  is the same as the one defined by  $f_i$ . Thus it suffices to prove (ii) $\Rightarrow$ (i) for  $f_i$ , reducing us to the case where  $Y$  is of finite type over  $S$ .

*Application to group schemes.* Let  $S$  be a scheme, let  $G$  be an  $S$ -group scheme locally of finite presentation over  $S$ , and suppose that  $G$  acts on the left on an  $S$ -scheme  $X$ . If  $X \rightarrow S$  has a section  $\xi$ , recall that the stabilizer  $\text{Stab}_G(\xi)$  is representable by a subgroup scheme of  $G$ ; see Exposé I, 2.3.3.

**Theorem 10.1.2.**—*Let  $S$  be a scheme and let  $G$  be an  $S$ -group scheme locally of finite presentation over  $S$ , acting on an  $S$ -scheme  $X$ . Suppose that  $X \rightarrow S$  has a section  $\xi$  whose stabilizer  $H$  in  $G$  is flat over  $S$ . If one of the following hypotheses holds:*

- a)  $X$  is locally of finite type over  $S$ ;
- b)  $S$  is locally Noetherian,

*then the fppf quotient sheaf  $G/H$  is representable by an  $S$ -scheme, locally of finite presentation over  $S$ , and the  $S$ -morphism*

$$f : G \longrightarrow X, \quad g \longmapsto g \cdot \xi$$

*factors as*

$$\begin{array}{ccc} G & & \\ \downarrow p & \searrow f & \\ G/H & \xhookrightarrow{i} & X \end{array}$$

Figure 46: The factorization of the orbit morphism in Theorem 10.1.2.

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 36 / combined-reader p. 298 / re-edited header p. 286.*

*where  $p$  is the canonical projection, which is faithfully flat and locally of finite presentation, and  $i$  is a monomorphism.*

*Proof.*— The morphism  $f$  makes  $G$  into an  $X$ -scheme. By definition of the stabilizer of  $\xi$ , the morphism

$$G \times_S H \longrightarrow G \times_X G, \quad (g, h) \longmapsto (g, gh)$$

is an isomorphism. Since  $H$  is flat over  $S$ , the scheme  $G \times_S H$  is flat over  $G$ ; hence the first projection

$$p_1 : G \times_X G \longrightarrow G$$

is flat.<sup>49</sup> Moreover, if  $X$  is locally of finite type over  $S$ , then  $f$  is locally of finite presentation (EGA IV<sub>1</sub>, 1.4.3(v)); otherwise,  $S$  is assumed locally Noetherian. It therefore suffices to apply Theorem 10.1.1 to  $f$ . It remains only to see that  $G/H$  is locally of finite presentation over  $S$ , and this follows immediately from Proposition 9.1.

**Corollary 10.1.3.**—*Let  $S$  be a scheme and  $u : G \rightarrow H$  a morphism of  $S$ -group schemes. Suppose that  $G$  is locally of finite presentation over  $S$ , and that either  $H$  is locally of finite type over  $S$  or  $S$  is locally Noetherian.*

<sup>49</sup>The source PDF prints  $G \times_S G$  in this projection. The preceding displayed isomorphism and the application of Theorem 10.1.1 require the fiber product  $G \times_X G$ , which is the reading adopted here.

If  $K = \ker(u)$  is flat over  $S$ , then the quotient group  $G/K$  is representable by an  $S$ -group scheme locally of finite presentation over  $S$ , and  $u$  factors as

$$\begin{array}{ccc} G & \xrightarrow{u} & H \\ & \searrow p \quad \nearrow i & \\ & G/K & \end{array}$$

Figure 47: The factorization of  $u$  through  $G/K$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 36 / combined-reader p. 298 / re-edited header p. 286.*

where  $p$  is the canonical projection and  $i$  is a monomorphism.

*Proof.*— Apply Theorem 10.1.2 with  $X = H$  and with the unit section of  $H$  as  $\xi$ .

## 10.2. Stabilizer of the diagonal

Let  $S$  be a Noetherian scheme,  $X$  an  $S$ -scheme of finite type, and  $G$  a flat  $S$ -group scheme of finite type acting on the left on  $X$ . Thus we have an  $S$ -action  $d_0 : G \times_S X \rightarrow X$ . Let  $d_1 : G \times_S X \rightarrow X$  be projection onto the second factor. As in §2(a), we have the groupoid

$$\begin{array}{ccccc} G \times_S G \times_S X & \xrightarrow[\mu \times X]{\text{pr}_{2,3}} & G \times_S X & \xrightarrow[d_0]{d_1} & X \\ & \xrightarrow{G \times d_0} & & & \end{array}$$

Figure 48: The action groupoid attached to  $(X, G)$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 36 / combined-reader p. 298 / re-edited header p. 286.*

whose cokernel, when it exists, is denoted  $G \backslash X$ .

**Definition 10.2.1.**— We denote by  $F \subset G \times_S X$  the stabilizer of the diagonal section; that is,  $F$  is the  $X$ -scheme defined by the Cartesian square

$$\begin{array}{ccc} F & \xrightarrow{\quad} & X \\ \downarrow & & \downarrow \Delta \\ G \times_S X & \xrightarrow{(d_0, d_1)} & X \times_S X \end{array}$$

Figure 49: The stabilizer  $F$  of the diagonal section.

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 37 / combined-reader p. 299 / re-edited header p. 287.*

Then  $F$  is an  $X$ -subgroup scheme of  $G \times_S X$ . Since  $G \times_S X$  is of finite type over the Noetherian scheme  $S$ , it is Noetherian; consequently  $F$  is of finite type both over  $S$  and over



$X$  (EGA I, 6.3.5 and 6.3.6). If  $X \rightarrow S$  is separated, then  $F$  is a closed  $X$ -subgroup scheme of  $G \times_S X$ .

Recall that  $G$  is said to *act freely* on  $X$  when

$$(d_0, d_1) : G \times_S X \longrightarrow X \times_S X$$

is a monomorphism (Exposé III, 3.2.1). Equivalently,  $F$  is the trivial group scheme over  $X$ .

### 10.3. The case in which $F$ is quasi-finite over $X$

Since  $F$  is of finite type over  $X$ , it is quasi-finite over  $X$  if and only if the stabilizers of the geometric points of  $X$  are finite.

**Theorem 10.3.1.**—<sup>50</sup> *Under the hypotheses of 10.2, suppose that  $F$  is quasi-finite over  $X$ . Then there is a dense,  $G$ -saturated open subset  $U \subset X$  having the following properties:*

- (i) *in  $(\mathbf{Sch}/S)$ , the cokernel  $V = G \backslash U$  exists; moreover,  $V$  is a quotient in the category of ringed spaces;*
- (ii)  *$p : U \rightarrow V$  is surjective, open, and of finite presentation;*
- (iii)  *$V$  is of finite presentation over  $S$ ;*
- (iv) *the morphism*

$$G \times_S U \longrightarrow U \times_V U, \quad (g, x) \longmapsto (gx, x),$$

*is surjective;*

- (v) *if, in addition,  $G$  acts freely on  $X$ , then  $U \rightarrow V$  is a left  $G$ -torsor, locally trivial for the fppf topology. In particular,  $U \rightarrow V$  is faithfully flat.<sup>51</sup>*

*Proof.*— By hypothesis, the morphism

$$G \times_S X \longrightarrow X \times_S X, \quad (g, x) \longmapsto (gx, x),$$

is quasi-finite. Theorem 8.1 therefore applies to the groupoid defined by  $(X, G)$ . Hence there is a dense saturated open subset  $U \subset X$  for which the quotient  $G \backslash U$  exists, and it has properties (i), (ii), and (iii); the same theorem also gives (iv).

To prove (v), recall that  $G$  acts freely on  $X$  if and only if  $(d_0, d_1)$  is an equivalence pair (Exposé III, 3.2.1).<sup>52</sup> In that case Theorem 8.1(iv) shows that

$$G \times_S U \longrightarrow U \times_V U$$

is an isomorphism and that  $p$  is faithfully flat and of finite presentation. Thus  $U$  is a  $G$ -torsor over  $V$ , locally trivial for the fppf topology.

<sup>50</sup>Editor's note: here too there is a largest open subset  $U \subset X$  satisfying the conclusions of the theorem; see editor's note 44.

<sup>51</sup>Editor's note: if one assumes in addition that  $G$  is a reductive  $S$ -group scheme and that its free action on  $X$  is linearizable, then  $G \backslash X$  is known to be representable and  $X \rightarrow G \backslash X$  is a left  $G$ -torsor. This follows from results of Raynaud and Seshadri and appears in [CTS79], Proposition 6.11.

<sup>52</sup>At this point the source PDF says *to prove (iv)*. The cited Theorem 8.1(iv) and the conclusion about a torsor establish part (v) in the present numbering; part (iv) follows from Theorem 8.1(iii).

## 10.4. The case in which $F$ is flat over $X$

Write

$$d = (d_0, d_1) : G \times_S X \longrightarrow X \times_S X, \quad d(g, x) = (gx, x).$$

Recall that the sheaf-theoretic graph  $\tilde{\Gamma}$  of the equivalence relation associated with  $(X, G)$  is the fppf  $S$ -subsheaf of  $X \times_S X$  given by the image of  $(d_0, d_1)$ . It is the fppf sheaf associated with the graph functor

$$T \longmapsto \Gamma(T) = \{(x_0, x_1) \in X(T) \times X(T) \mid x_0 \in G(T)x_1\}.$$

Put  $G_X = G \times_S X$ . For every  $S$ -scheme  $T$  there is a surjective map

$$G_X(T) \longrightarrow \Gamma(T), \quad (g, x) \longmapsto (gx, x),$$

which induces a bijection

$$\phi(T) : G_X(T)/F(T) \xrightarrow{\sim} \Gamma(T).$$

Indeed, if  $(g, x), (g', x') \in G_X(T)$  have the same image, then  $x' = x$  and  $g^{-1}g'x = x$ , so that  $(g^{-1}g', x) \in F(T)$ , and  $(g, x)$  and  $(g', x)$  have the same image in  $G_X(T)/F(T)$ .<sup>53</sup>

By definition (see Exposé IV, 4.4.1(ii), or the proof of 5.2.1), the quotient sheaf  $G_X/F$  is the fppf sheaf associated with the presheaf

$$T \longmapsto G_X(T)/F(T) \simeq \Gamma(T).$$

We therefore obtain an isomorphism of sheaves

$$\phi : G_X/F \xrightarrow{\sim} \tilde{\Gamma}.$$

**Theorem 10.4.1.**—<sup>54</sup> *Under the hypotheses of 10.2, the following assertions hold:*

- a) *the sheaf  $\tilde{\Gamma}$  is representable if and only if  $F$  is flat over  $X$ ;*
- b) *suppose that  $F$  is flat over  $X$ . Then the morphisms induced by  $d_1$  and  $d_0$*

$$G_X/F \begin{array}{c} \xrightarrow{\bar{d}_1} \\ \xrightarrow{\bar{d}_0} \end{array} X$$

Figure 50: The two maps induced by  $d_1$  and  $d_0$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 38 / combined-reader p. 300 / re-edited header p. 288.*

*are faithfully flat and of finite presentation.*

<sup>53</sup>The source PDF prints  $(g^{-1}g'x, x) \in F(T)$ . Since  $F(T) \subseteq G(T) \times X(T)$ , the first coordinate must be the group element  $g^{-1}g'$ ; the English adopts this type-correct reading. In the same sentence the source writes  $F_X(T)$  once in the denominator; the displayed bijection immediately above and the presheaf immediately below both use the defined stabilizer  $F(T)$ , which is retained consistently here.

<sup>54</sup>Editor's note: this is part (2) of Theorem 3 of [Ray67b]. That note also sketches another proof of Theorem 10.1.1.

*Proof of a).*— Suppose that the fppf sheaf  $G_X/F$  is represented by an  $X$ -scheme  $Y$ . Then, by Exposé IV, 6.3.3, the morphism  $p : G_X \rightarrow Y$  is faithfully flat and locally of finite presentation, and the second square in the following diagram is Cartesian:

$$\begin{array}{ccccc}
 F & \longrightarrow & F \times_X G_X & \longrightarrow & G_X \\
 \downarrow & & \downarrow & & \downarrow p \\
 X & \xrightarrow{e_X} & G_X & \longrightarrow & Y
 \end{array}$$

Figure 51: The Cartesian square used in the proof of Theorem 10.4.1(a).

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 39 / combined-reader p. 301 / re-edited header p. 289.*

The first square is obtained by base change along the unit section  $e_X : X \rightarrow G_X$ . Since  $p$  is faithfully flat and locally of finite presentation, the same is true of  $F \rightarrow X$ .

Conversely, suppose  $F$  is flat over  $X$ , and put  $X^2 = X \times_S X$ . The morphism  $d : G_X \rightarrow X^2$  gives the fiber square

$$\begin{array}{ccc}
 G_X \times_{X^2} G_X & \longrightarrow & G_X \\
 \downarrow & & \downarrow \\
 G_X & \longrightarrow & X^2
 \end{array}$$

Figure 52: The fiber square associated with  $d : G_X \rightarrow X^2$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 39 / combined-reader p. 301 / re-edited header p. 289.*

The first projection

$$\mathrm{pr}_1 : G_X \times_{X^2} G_X \longrightarrow G_X$$

is then an  $F \times_X G_X$ -torsor over  $G_X$ , and is therefore flat and of finite type, because  $F$  is.<sup>55</sup> By Theorem 10.1.1,  $d$  factors uniquely as

$$G_X \xrightarrow{\psi} Y \xrightarrow{\tau} X \times_S X$$

Figure 53: The unique factorization of  $d$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 39 / combined-reader p. 301 / re-edited header p. 289.*

where  $\psi$  is faithfully flat and of finite type, and  $\tau$  is a monomorphism of schemes.

<sup>55</sup>The source PDF instead prints the morphism  $G_X \times_{X^2} G_X \rightarrow X^2$  and calls it an  $F \times_X X^2$ -torsor over  $X^2$ . That morphism is not generally surjective. The first projection to  $G_X$  is the stabilizer torsor and is the flatness input required by Theorem 10.1.1; the English uses this type-correct reading.

It follows that the morphism of sheaves  $\psi : G_X \rightarrow Y$  is  $F$ -invariant, and hence induces a morphism

$$\bar{\psi} : G_X/F \rightarrow Y.$$

Moreover, because  $\psi$  is faithfully flat and of finite type, the monomorphism of sheaves  $\tau$  factors through the sheaf image of  $d$ , namely  $\tilde{\Gamma}$ . Thus the isomorphism  $G_X/F \simeq \tilde{\Gamma}$  factors through the monomorphism  $Y \rightarrow \tilde{\Gamma}$ . We conclude that  $Y$  represents  $G_X/F$ .

*Proof of b).*— Suppose  $F$  is flat over  $X$ . By part (a) and its proof,  $G_X/F$  is representable, and

$$p : G_X \rightarrow G_X/F$$

is faithfully flat and of finite presentation. On the other hand, the morphisms  $d_i : G_X \rightarrow X$ , for  $i = 0, 1$ , are faithfully flat and of finite presentation by hypothesis. Since  $d_i = \bar{d}_i \circ p$ , it follows from EGA IV<sub>2</sub>, 2.2.13(iii), and EGA IV<sub>3</sub>, 11.3.16, that each  $\bar{d}_i$  is faithfully flat and of finite presentation.

**Theorem 10.4.2.**—<sup>56</sup> *Under the hypotheses of 10.2, suppose that  $F$  is flat over  $X$ . Then there is a dense saturated open subset  $U \subset X$  such that the fppf quotient  $V = G \backslash U$  is an  $S$ -scheme of finite type and  $U \rightarrow V$  is faithfully flat and of finite presentation.*

*Proof.*— Theorem 10.4.1 shows that  $G_X/F \simeq \tilde{\Gamma}$  is representable. The fppf sheaf  $G \backslash X$  is therefore the quotient sheaf of

$$G_X/F \begin{array}{c} \xrightarrow{\bar{d}_1} \\ \xrightarrow{\bar{d}_0} \end{array} X$$

Figure 54: The equivalence pair whose quotient is  $G \backslash X$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 40 / combined-reader p. 302 / re-edited header p. 290.*

By what precedes,

$$\bar{d}_i : G_X/F \rightarrow X$$

is faithfully flat and of finite presentation for  $i = 0, 1$ , and the morphism

$$G_X/F \xrightarrow{\sim} \tilde{\Gamma} \hookrightarrow X \times_S X$$

Figure 55: The monomorphism from the quotient  $G_X/F$  to  $X \times_S X$ .

*Native L<sup>A</sup>T<sub>E</sub>X reconstruction from the corrected Polo–Gille diagram; source locator: Exposé V, component-local p. 40 / combined-reader p. 302 / re-edited header p. 290.*

is a monomorphism. In other words,  $(\bar{d}_0, \bar{d}_1)$  is an equivalence pair. Consequently, Theorem 8.1 applies. There is therefore a dense saturated open subset  $U \subset X$  such that the fppf quotient  $V = G \backslash U$  is an  $S$ -scheme of finite type and  $U \rightarrow V$  is faithfully flat and of finite presentation.

<sup>56</sup>Editor's note: here too there is a largest open subset  $U \subset X$  satisfying the conclusions of the theorem. Moreover, a codimension-one point  $x \in X$  belongs to  $U$  if and only if

$$(G_X/F) \times_X \operatorname{Spec}(\mathcal{O}_{X,x}) \rightarrow \operatorname{Spec}(\mathcal{O}_{X,x}) \times_S \operatorname{Spec}(\mathcal{O}_{X,x})$$

is a closed immersion; see editor's note 44.

Taking the generic-flatness theorem into account (EGA IV<sub>2</sub>, 6.9.3), we obtain the following result.

**Corollary 10.4.3.**— *Under the hypotheses of 10.2, suppose that  $X$  is reduced. Then there is a dense saturated open subset  $U \subset X$  such that the fppf quotient  $G \backslash U$  is an  $S$ -scheme of finite type and  $U \rightarrow G \backslash U$  is faithfully flat and of finite presentation.*

## Bibliography<sup>57</sup>

- [AK80] A. B. Altman and S. L. Kleiman, *Compactifying the Picard Scheme*, Adv. Math. **35** (1980), 50–112.
- [An73] S. Anantharaman, *Schémas en groupes, espaces homogènes et espaces algébriques sur une base de dimension 1*, Mém. Soc. Math. France **33** (1973), 5–79.
- [BLR90] S. Bosch, W. Lütkebohmert, and M. Raynaud, *Néron models*, Springer-Verlag, 1990.
- [CTS79] J.-L. Colliot-Thélène and J.-J. Sansuc, *Fibrés quadratiques et composantes connexes réelles*, Math. Ann. **244** (1979), 105–134.
- [DG70] M. Demazure and P. Gabriel, *Groupes algébriques*, Masson & North-Holland, 1970.
- [DR81] J. Dixmier and M. Raynaud, *Sur le quotient d’une variété algébrique par un groupe algébrique*, pp. 327–344 in *Mathematical Analysis and Applications* (L. Schwartz 65th birthday, ed. L. Nachbin), Adv. Math. Suppl. Stud., Vol. 7A, 1981.
- [Fe03] D. Ferrand, *Conducteur, descente et pincement*, Bull. Soc. Math. France **131** (2003), no. 4, 553–585.
- [Hi62] H. Hironaka, *An example of a non-Kählerian complex analytic deformation of Kählerian complex structures*, Ann. of Math. **75** (1962), no. 1, 190–208.
- [KM97] S. Keel and S. Mori, *Quotient by groupoids*, Ann. of Math. **145** (1997), no. 1, 193–213.
- [Ko97] J. Kollár, *Quotient spaces modulo algebraic groups*, Ann. of Math. **145** (1997), no. 1, 33–79.
- [Ko08] J. Kollár, *Quotients by finite equivalence relations*, arXiv:0812.3608.
- [Mum65] D. Mumford, *Geometric invariant theory*, Springer-Verlag, 1965; 2nd ed. with J. Fogarty, 1982; 3rd ed. with J. Fogarty and F. Kirwan, 1994.
- [Mur65] J. P. Murre, *Representation of unramified functors. Applications* (according to unpublished results of A. Grothendieck), Sémin. Bourbaki, Vol. 9, Exp. 294 (1965), Soc. Math. France, 1995.
- [Ray67a] M. Raynaud, *Passage au quotient par une relation d’équivalence plate*, pp. 78–85 in *Proc. Conf. Local Fields (Driebergen)* (ed. T. A. Springer), Springer-Verlag, 1967.
- [Ray67b] M. Raynaud, *Sur le passage au quotient par un groupoïde plat*, C. R. Acad. Sci. Paris (Sér. A) **265** (1967), 384–387.

---

<sup>57</sup>Editor’s note: additional references cited in this exposé.

# Exposé VI<sub>A</sub>

## Generalities on Algebraic Groups

by P. Gabriel

<sup>0</sup>Throughout this chapter,  $A$  will denote an Artinian local ring with residue field  $k$ . A group scheme  $G$  over  $\operatorname{Spec} A$  will be called simply an  $A$ -group. This  $A$ -group is said to be *locally of finite type* if its underlying scheme is locally of finite type over  $A$ ; it is said to be *algebraic* if its underlying scheme is of finite type over  $A$ .

### 0. Preliminary Remarks

**0.1.** Let us first consider a group scheme  $G$  over an arbitrary scheme  $S$ . We call the structural morphism

$$\mu : G \times_S G \longrightarrow G$$

*multiplication*, and we call the morphism  $c : G \rightarrow G$ , defined by  $c(T)(x) = x^{-1}$  for every scheme  $T$  over  $S$  and every  $x \in G(T)$ , *inversion*. If  $U$  and  $V$  are subsets of the underlying set of  $G$ , we denote by  $U \cdot V$  the image under multiplication of the subset of  $G \times_S G$  consisting of those points whose first projection belongs to  $U$  and whose second projection belongs to  $V$ . Likewise, the notations  $U^{-1}$  and  $c(U)$  are equivalent.

Let  $\operatorname{pr}_1$  be the projection of  $G \times_S G$  onto the first factor, and let

$$\sigma : G \times_S G \longrightarrow G \times_S G$$

be the morphism with components  $\operatorname{pr}_1$  and  $\mu$ . For every  $S$ -scheme  $T$ , the map  $\sigma(T)$  is

$$(x, y) \longmapsto (x, xy);$$

it follows that  $\sigma$  is an automorphism. The composite of this automorphism with the projection  $\operatorname{pr}_2$  of  $G \times_S G$  onto the second factor is the multiplication morphism. When  $G$  is flat over  $S$ ,  $\operatorname{pr}_2$ , and hence  $\mu$ , are flat morphisms; when  $G$  is smooth over  $S$ ,  $\operatorname{pr}_2$ , and hence  $\mu$ , are smooth morphisms, and so on.

**0.2.** From now on we assume that  $S$  is the spectrum of an Artinian local ring  $A$  with residue field  $k$ . We denote by  $(\mathbf{Sch}/k)_{\operatorname{red}}$  the category of reduced schemes over  $k$ . For every scheme  $X$  over  $A$ , the reduced scheme  $X_{\operatorname{red}}$  is an object of  $(\mathbf{Sch}/k)_{\operatorname{red}}$ , and the functor

$$X \longmapsto X_{\operatorname{red}}$$

is right adjoint to the inclusion of  $(\mathbf{Sch}/k)_{\operatorname{red}}$  into  $\mathbf{Sch}/A$ . It follows that, for every  $A$ -group  $G$ ,  $G_{\operatorname{red}}$  is a group in the category  $(\mathbf{Sch}/k)_{\operatorname{red}}$ .<sup>1</sup> In other words, for every reduced  $k$ -scheme

<sup>0</sup>Editor's note: version of October 13, 2024.

<sup>1</sup>Editor's note: the sentences that follow have been added.

$T$ ,  $G_{\text{red}}(T)$  is endowed, functorially in  $T$ , with a group structure. One must bear in mind that  $G_{\text{red}}$  is not necessarily a  $k$ -group, because its “multiplication” is only a morphism

$$(G_{\text{red}} \times_k G_{\text{red}})_{\text{red}} \longrightarrow G_{\text{red}}^2.$$

<sup>3</sup> However, if  $k$  is a perfect field, the inclusion of  $(\mathbf{Sch}/k)_{\text{red}}$  into  $\mathbf{Sch}/k$  commutes with products. Groups in  $(\mathbf{Sch}/k)_{\text{red}}$  may therefore be identified with group schemes over  $k$  whose underlying schemes are reduced. In this case, if  $G$  is a  $k$ -group,  $G_{\text{red}}$  is a group subscheme of  $G$ ; in general, this subgroup is not normal in  $G$ .

For example, if  $k$  is a field of characteristic 3, the constant group  $(\mathbb{Z}/2\mathbb{Z})_k$  acts nontrivially on the diagonalizable group  $D_k(\mathbb{Z}/3\mathbb{Z})$  (cf. Exposé I, 4.1 and 4.4). If  $G$  denotes the semidirect product of  $D_k(\mathbb{Z}/3\mathbb{Z})$  by  $(\mathbb{Z}/2\mathbb{Z})_k$  defined by this action, then  $G_{\text{red}}$  is identified with  $(\mathbb{Z}/2\mathbb{Z})_k$  and is not normal in  $G$ .<sup>4</sup>

Let  $k$  be an arbitrary field, let  $k^{p^{-\infty}}$  be its perfect closure, and let  $H$  be a group in  $(\mathbf{Sch}/k)_{\text{red}}$ . Then

$$(H \otimes_k k^{p^{-\infty}})_{\text{red}}$$

is a group scheme over the perfect field  $k^{p^{-\infty}}$ . Since  $H \otimes_k k^{p^{-\infty}}$  and  $(H \otimes_k k^{p^{-\infty}})_{\text{red}}$  have the same underlying topological space, groups in  $(\mathbf{Sch}/k)_{\text{red}}$  share with  $k$ -groups certain topological properties that are invariant under extension of the base field. For example, it will follow from 0.3 and the preceding remarks that every group in  $(\mathbf{Sch}/k)_{\text{red}}$  is separated.

In what follows, we shall encounter groups in  $(\mathbf{Sch}/k)_{\text{red}}$  whenever we deal with a nonempty locally closed subset  $U$  of an  $A$ -group  $G$  such that

$$U \cdot U = U \quad \text{and} \quad U^{-1} = U.$$

Indeed, the reduced subscheme of  $G$  defined by  $U$  is then a group in  $(\mathbf{Sch}/k)_{\text{red}}$ .

**0.3.** *An  $A$ -group  $G$  is always separated, because the unit section  $e : \text{Spec } A \rightarrow G$  is a closed immersion.* Indeed, let  $x$  be the unique point of  $\text{Spec } A$ , and let  $\eta : G \rightarrow \text{Spec } A$  be the structural morphism. Since  $\eta \circ e = \text{id}_{\text{Spec } A}$ , for every affine open  $U = \text{Spec } B$  of  $G$  containing  $e(x)$ , the homomorphism  $B \rightarrow A$  has a section and is therefore surjective. It follows that  $e$  is a closed immersion.<sup>5</sup>

The diagonal in  $G \times_A G$  is identified with the functor from  $(\mathbf{Sch}/A)^\circ$  to **Sets** which associates to every  $A$ -scheme  $S$  the inverse image of the identity element of  $G(S)$  under the map

$$\varphi(S) : G(S) \times G(S) \longrightarrow G(S), \quad (x, y) \longmapsto x \cdot y^{-1}.$$

We therefore have the following cartesian square. Since the diagonal morphism is obtained from a closed immersion by base change, it is itself a closed immersion.

$$\begin{array}{ccc} G \times_S G & \xrightarrow{\varphi} & G \\ \text{diagonal morphism} \uparrow & & \uparrow \text{unit section} \\ G & \longrightarrow & \text{Spec } A \end{array}$$

<sup>2</sup>Editor's note: for examples of group schemes  $G$  over a nonperfect field  $k$  for which  $G_{\text{red}}$  is not a group scheme over  $k$ , see 1.3.2 below.

<sup>3</sup>Editor's note: the remainder of 0.2, as well as 0.3, has been slightly modified.

<sup>4</sup>Editor's note: for an analogous example with  $G$  connected, one may take, when  $\text{char}(k) = p > 0$ , the semidirect product of  $\alpha_{p,k}$  and  $\mathbb{G}_{m,k}$ .

<sup>5</sup>Editor's note: this argument is valid for every local ring  $A$  of dimension zero, and shows that, if  $S$  is a discrete scheme, every  $S$ -group is separated; cf. VI<sub>B</sub>, 5.2. On the other hand (cf. VI<sub>B</sub>, 5.6.4), if  $S$  contains a nonisolated closed point  $s$ , the  $S$ -scheme  $G$  obtained by gluing two copies of  $S$  along the open subset  $S - \{s\}$  is not separated over  $S$ , although it carries an  $S$ -group structure.

**0.4.**<sup>6</sup> Let  $G$  be an  $A$ -scheme. We shall say that a point  $g$  of  $G$  is *strictly rational over  $A$*  if there exists an  $A$ -morphism

$$s : \operatorname{Spec} A \longrightarrow G$$

which sends the unique point of  $\operatorname{Spec} A$  to  $g$ ; that is, if the homomorphism  $A \rightarrow \mathcal{O}_{G,g}$  admits a retraction. One then has  $\kappa(g) = k$ , so  $g$  is a closed point of  $G$ . Indeed, if  $B$  is the ring of an affine open neighborhood of  $g$ , and if  $\mathfrak{p}$  is the prime ideal of  $B$  corresponding to  $g$ , then  $B/\mathfrak{p} \subset k$  is a finite  $A$ -algebra, hence an integral Artinian ring, and therefore a field.

Suppose henceforth that  $G$  is an  $A$ -group. Such a morphism  $s : \operatorname{Spec} A \rightarrow G$  defines an automorphism  $r_s$  of the scheme  $G$  over  $A$ , called *right translation by  $s$* . For every morphism  $\pi : S \rightarrow \operatorname{Spec} A$ , the automorphism  $r_s(\pi)$  of  $G(S)$  is defined by

$$x \longmapsto x \cdot G(\pi)(s), \quad x \in G(S).$$

Similarly,  $\ell_s$  denotes *left translation by  $s$* , namely the automorphism of  $G$  defined by

$$\ell_s(\pi)(x) = G(\pi)(s) \cdot x, \quad x \in G(S).$$

The schemes  $G \otimes_A k$  and  $G$  have the same underlying topological space  $\underline{G}$ ,  $G \otimes_A k$  is a  $k$ -group, and  $s \otimes_A k$  depends only on  $g$ , not on  $s$ . Consequently, the automorphisms of  $\underline{G}$  induced by  $r_s$  and  $\ell_s$  (or by  $r_{s \otimes k}$  and  $\ell_{s \otimes k}$ ) depend only on  $g$ , not on  $s$ . Thus, if  $P$  is a subset of  $\underline{G}$ , we may write

$$r_g(P) \text{ or } P \cdot g \quad (\text{respectively, } \ell_g(P) \text{ or } g \cdot P)$$

instead of  $r_s(P)$  (respectively,  $\ell_s(P)$ ), in agreement with 0.1.

**Remark 0.4.1.** If  $g$  is a strictly rational point of  $G$ , and if  $A \rightarrow A'$  is a morphism of Artinian local rings, then  $G' = G \otimes_A A'$  has a unique point  $g'$  above  $g$ , and  $g'$  is strictly rational over  $A'$ . Moreover, if  $P'$  denotes the inverse image of  $P$  in  $G'$ , then  $P' \cdot g'$  is the inverse image of  $P \cdot g$  (cf. EGA I, 3.4.8).

**Proposition 0.5.**<sup>7</sup> Let  $G$  be an  $A$ -group, and let  $U, V$  be two dense open subsets of  $G$ . Then  $U \cdot V$ , that is, the image of  $U \times_A V$  under multiplication, is the entire underlying space of  $G$ .

*Proof.* Since  $G$  and  $G \otimes_A k$  have the same underlying space, after replacing  $A$  by  $k$  and  $G$  by  $G \otimes_A k$ , if necessary, we may assume that  $A = k$ . Let  $g \in G$ , and set  $K = \kappa(g)$ . Then left translation  $\ell_g$  is an automorphism of  $G_K$ . Since the projection  $G_K \rightarrow G$  is open,  $U_K$  and  $V_K$  are dense open subsets of  $G_K$ , as is the image of  $V_K$  under  $\ell_g$ . Consequently there exists  $v \in V_K$  such that  $u = \ell_g(v)$  belongs to  $U_K$ .

Let  $L$  be an extension of  $K$  containing  $\kappa(v)$ , and hence  $\kappa(u)$ , and let  $g_L$  and  $v_L$  be the  $L$ -points of  $G_L$  induced by  $g$  and  $v$ . Then

$$g_L \cdot v_L = u'$$

is a point of  $G_L$  above  $u$ , and therefore

$$g_L = u' \cdot v_L^{-1}$$

lies above  $U \cdot V$ . Thus  $g \in U \cdot V$ , proving the proposition.  $\square$

<sup>6</sup>Editor's note: the original has been expanded in what follows; in particular, Remark 0.4.1 has been added.

<sup>7</sup>Editor's note: the original stated this result under the hypothesis that  $G$  be locally of finite type over  $A$ . Since the result is useful in the general case, and since the proof is essentially the same, it has been stated and proved here in full generality. It will be used several times below.



**Corollary 0.5.1.**<sup>8</sup> *If  $G$  is an irreducible  $A$ -group, then  $G$  is quasi-compact.*

*Proof.* Let  $U$  be a nonempty affine open subset of  $G$ . Then  $U$  is dense in  $G$ , so by 0.5 the morphism

$$\mu : U \times_A U \longrightarrow G$$

is surjective. Hence  $G$  is quasi-compact, since  $U \times_A U$  is.  $\square$

**Corollary 0.5.2.**<sup>9</sup> *Let  $G$  be an  $A$ -group scheme, and let  $H$  be a group subscheme of  $G$  over  $A$ . Then  $H$  is closed.*

*Proof.* Let  $k'$  be the perfect closure of the residue field  $k$  of  $A$ . Since the underlying topological spaces of  $G$  and  $H$  are unchanged under the base change  $A \rightarrow k \rightarrow k'$ , we may assume that  $A = k$  is a perfect field. We may then assume that  $G$  and  $H$  are reduced, hence geometrically reduced.

Let  $\overline{H}$  be the closure of  $H$ . Then  $\mu^{-1}(\overline{H})$  is a closed subset of  $G \times G$  containing  $H \times H$ . The morphisms

$$H \longrightarrow \operatorname{Spec} k \quad \text{and} \quad \overline{H} \longrightarrow \operatorname{Spec} k$$

are universally open. Since  $H$  is dense in  $\overline{H}$ ,  $H \times H$  is dense in  $\overline{H} \times H$ , and  $\overline{H} \times H$  is dense in  $\overline{H} \times \overline{H}$ . Thus  $H \times H$  is dense in  $\overline{H} \times \overline{H}$ . Consequently

$$\mu(\overline{H} \times \overline{H}) \subset \overline{H}.$$

Since  $\overline{H} \times \overline{H}$  is reduced,  $\mu$  induces a morphism

$$\mu' : \overline{H} \times \overline{H} \longrightarrow \overline{H}.$$

Now let  $g \in \overline{H}$ , and put  $K = \kappa(g)$ . The projection  $\overline{H}_K \rightarrow \overline{H}$  is open, so  $H_K$  and  $\ell_g(H_K)$  are two dense open subsets of  $\overline{H}_K$ . Hence there exist  $u, v \in H_K$  such that  $\ell_g(v) = u$ . As in the proof of 0.5, it follows that  $g$  belongs to  $H \cdot H = H$ . Therefore  $\overline{H} = H$ .  $\square$

## 1. Local properties of an $A$ -group locally of finite type

We shall first see that, if  $G$  is locally of finite type and flat over  $A$ , one can “make every closed point of  $G$  strictly rational” by means of a finite flat extension of the base.

**1.1.** Unless expressly stated otherwise, from now on we assume that  $G$  is an  $A$ -group locally of finite type. When  $A$  is a field  $k$ , we shall obtain in Exposé VII<sub>B</sub> very precise results on the local rings of  $G$ .<sup>10</sup> Here we content ourselves with a few elementary results.

**Proposition 1.1.1.** *Let  $x$  be a point of an  $A$ -group  $G$  locally of finite type and flat over  $A$ . Then the local ring  $\mathcal{O}_{G,x}$  is Cohen–Macaulay, and there is a system of parameters  $a_1, \dots, a_n$  of  $\mathcal{O}_{G,x}$  such that  $\mathcal{O}_{G,x}/(a_1, \dots, a_n)$  is a finite flat  $A$ -module (hence finite free).*

We first suppose that  $A$  is equal to its residue field  $k$ . It then suffices to prove that  $\mathcal{O}_{G,x}$  is Cohen–Macaulay, and we may restrict to the case in which  $x$  is a closed point (cf. EGA 0<sub>IV</sub>, 16.5.13). By Lemma 1.1.2 below,  $G$  contains a closed point  $y$  such that  $\mathcal{O}_{G,y}$  is Cohen–Macaulay. By SGA 1, I, § 9, this is equivalent to saying that, for every finite extension  $K$  of the base field  $k$  and every point  $\overline{y}$  of  $\overline{G} = G \otimes_k K$  lying above  $y$ , the local

<sup>8</sup>Editor’s note: this corollary has been inserted here; cf. 2.4 and 2.6.2 below.

<sup>9</sup>Editor’s note: this corollary, used implicitly in VIII, 6.7, has been added; see also VI<sub>B</sub>, 6.2.5.

<sup>10</sup>Editor’s note: for example, they are always complete intersections; see Exposé VII<sub>B</sub>, 5.5.1. Moreover, if  $\operatorname{char}(k) = 0$ , then  $G$  is smooth (Exposé VI<sub>B</sub>, 1.6.1; see also Exposé VII<sub>B</sub>, 3.3.1).

ring  $\mathcal{O}_{\bar{G}, \bar{y}}$  is Cohen–Macaulay. If  $K$  has been chosen sufficiently large, that is, if  $K$  contains a normal extension of  $k$  containing the residue fields  $\kappa(y)$  and  $\kappa(x)$ , then  $\bar{y}$  is (strictly) rational over  $K$ , as is every point  $\bar{x}$  of  $\bar{G}$  lying above  $x$ .<sup>11</sup> Since the automorphism

$$r_{\bar{x}} \circ (r_{\bar{y}})^{-1}$$

sends  $\bar{y}$  to  $\bar{x}$ , it follows that  $\mathcal{O}_{\bar{G}, \bar{x}}$ , and hence  $\mathcal{O}_{G, x}$  (SGA 1, I, § 9), are Cohen–Macaulay.

Now let  $A$  again be arbitrary. The preceding argument applies to  $k \otimes_A G$ , so that  $k \otimes_A \mathcal{O}_{G, x}$  is Cohen–Macaulay. If  $a_1, \dots, a_n$  is a sequence of elements of  $\mathcal{O}_{G, x}$  whose image in  $k \otimes_A \mathcal{O}_{G, x}$  is a system of parameters, it follows from SGA 1, IV, 5.7, or EGA 0<sub>IV</sub>, 15.1.16, that  $a_1, \dots, a_n$  is an  $\mathcal{O}_{G, x}$ -regular sequence and that  $\mathcal{O}_{G, x}/(a_1, \dots, a_n)$  is finite and flat (hence finite and free) over  $A$ .

**Lemma 1.1.2.** *Every nonempty scheme  $X$ , locally of finite type over an Artinian ring  $A$ , contains a closed point  $x$  whose local ring is Cohen–Macaulay.*

We may plainly suppose that  $X$  is affine, with algebra  $B$ , and argue by induction on  $\dim X$  (the assertion is clear if  $X$  is discrete, since all its local rings are then Artinian). Since  $B$  is of finite type over  $A$ , if  $\dim B > 0$ , then  $B$  contains an element  $a$  that is not invertible and is not a zero divisor.<sup>12</sup> The closed subscheme

$$X' = \operatorname{Spec} B/(a)$$

of  $X$  then has dimension strictly less than  $\dim X$ , and by the induction hypothesis it contains a closed point  $x$  such that  $\mathcal{O}_{X', x}$  is Cohen–Macaulay. Since  $\mathcal{O}_{X', x} = \mathcal{O}_{X, x}/(a)$ , and  $a$  is neither invertible nor a zero divisor in  $\mathcal{O}_{X, x}$ , the ring  $\mathcal{O}_{X, x}$  is Cohen–Macaulay (see also EGA IV<sub>2</sub>, 6.11.3).

**Proposition 1.2.** *Let  $A$  be an Artinian local ring, let  $G$  be an  $A$ -group locally of finite type and flat over  $A$ , and let  $x$  be a closed point of  $G$ . Then there exists a local  $A$ -algebra  $A'$ , finite and free over  $A$ , such that every point of  $G \otimes_A A'$  lying above  $x$  is strictly rational over  $A'$ .*

<sup>13</sup> Indeed, let  $k_1$  be a finite normal extension of  $k$  containing the residue field  $\kappa(x)$  of  $x$ . By Lemma V.4.1.2, there exists a local  $A$ -algebra  $A_1$ , finite and free over  $A$ , with residue field  $k_1$ . In this case (cf. Editor’s note 11), all the points  $g_1, \dots, g_n$  of  $G \otimes_A A_1$  lying above  $x \in G$  have  $k_1$  as residue field; that is,  $g_1, \dots, g_n$  are rational over  $A_1$  in the sense of Exposé V, § 4.e).

Let  $B_1, \dots, B_n$  be the local rings of  $g_1, \dots, g_n$ . By 1.1.1, the rings  $B_1, \dots, B_n$  have quotients  $B'_1, \dots, B'_n$  that are Artinian and finite and free over  $A_1$ . Set

$$A' = B'_1 \otimes_{A_1} \cdots \otimes_{A_1} B'_n.$$

Then  $A'$  is local, finite, and free over  $A_1$ , and, for each  $i = 1, \dots, n$ , there are surjective homomorphisms

$$B_i \otimes_{A_1} A' \twoheadrightarrow B'_i \otimes_{A_1} A' \twoheadrightarrow A',$$

<sup>11</sup>Editor’s note: indeed, the hypothesis on  $K$  implies that, for every extension  $L$  of  $K$ , every  $k$ -morphism  $\kappa(x) \rightarrow L$  (respectively,  $\kappa(y) \rightarrow L$ ) factors through  $K$ . Consequently, all the points of  $G \otimes_k K$  lying above  $x$  or  $y$  have  $K$  as residue field; that is, they are (strictly) rational over  $K$ .

<sup>12</sup>Editor’s note: indeed,  $B$  is a Noetherian Jacobson ring (cf. [BAC], V, § 3.4). If every noninvertible element is a zero divisor, then every prime ideal is an associated prime ideal of  $B$ ; in particular,  $B$  has only finitely many maximal ideals  $\mathfrak{m}_1, \dots, \mathfrak{m}_n$ . Since  $B$  is a Jacobson ring, the intersection of the  $\mathfrak{m}_i$  is the nilradical of  $B$ . It follows that each  $\mathfrak{m}_i$  is a minimal prime ideal of  $B$ , and hence that  $\dim B = 0$ .

<sup>13</sup>Editor’s note: the original has been expanded in what follows.

the second induced by the multiplication map

$$B'_i \otimes_{A_1} B'_i \rightarrow B'_i.$$

Consequently,  $A'$  has the required property.

**1.3.** Let  $e$  be the unit element (or origin) of  $G$ , that is, the image of the unique point of  $\operatorname{Spec} A$  under the unit section  $\operatorname{Spec} A \rightarrow G$ . By definition,  $e$  is strictly rational over  $A$ .

**Proposition 1.3.1.**<sup>14</sup> *Let  $G$  be a group locally of finite type and flat over an Artinian ring  $A$ , and let  $K$  (respectively,  $\bar{k}$ ) be the perfect closure (respectively, an algebraic closure) of the residue field  $k$  of  $A$ .*

- (1) *For every closed point  $x$  of  $\bar{G} = G \otimes_k \bar{k}$ , the local rings  $\mathcal{O}_{\bar{G},e}$  and  $\mathcal{O}_{\bar{G},x}$  are isomorphic. In particular, the tangent spaces  $T_e \bar{G}$  and  $T_x \bar{G}$  have the same dimension.*
- (2) *The following assertions are equivalent:*
  - (i)  *$G \otimes_A K$  is reduced.*
  - (i bis)  *$\mathcal{O}_{G,e} \otimes_A K$  is reduced.*
  - (ii)  *$G$  is smooth over  $A$ .*
  - (ii bis)  *$G$  is smooth over  $A$  at the origin.*

*Proof.* (1) Let  $x$  be a closed point of  $\bar{G}$ . There is exactly one  $\bar{k}$ -morphism

$$s : \operatorname{Spec} \bar{k} \rightarrow \bar{G}$$

whose image is  $x$ . Right translation  $r_s$  then induces an isomorphism from

$$\mathcal{O}_{\bar{G},e} = \mathcal{O}_{G,e} \otimes_k \bar{k}$$

onto  $\mathcal{O}_{\bar{G},x}$ , which proves (1).

We now prove (2). By SGA 1, II.2.1, we reduce at once to the case in which  $A$  is a field ( $A = k$ ). The implications

$$(i) \Rightarrow (i \text{ bis}), \quad (ii) \Rightarrow (ii \text{ bis}), \quad (ii) \Rightarrow (i), \quad (ii \text{ bis}) \Rightarrow (i \text{ bis})$$

are evident, so it remains only to prove that (i bis) implies (ii).

Now (i bis) implies that  $\mathcal{O}_{G,e} \otimes_k \bar{k}$  is reduced. Hence, by (1),  $\mathcal{O}_{\bar{G},x}$  is reduced for every closed point  $x$  of  $\bar{G}$ , and therefore  $\bar{G}$  is reduced. Since  $\bar{G}$  is locally of finite type over  $\bar{k}$ , there is at least one closed point  $y$  such that  $\mathcal{O}_{\bar{G},y}$  is regular. By (1), the local rings at all closed points of  $\bar{G}$  are isomorphic to  $\mathcal{O}_{\bar{G},e}$ ; thus all these local rings are regular. Consequently,  $\bar{G}$  is smooth over  $\bar{k}$ , and hence  $G$  is smooth over  $k$ .

<sup>15</sup> We may now give the examples below, pointed out by M. Raynaud, of group schemes  $G$  over a nonperfect field  $k$  for which  $G_{\text{red}}$  is not a  $k$ -group scheme.

**Examples 1.3.2.** Let  $k$  be a nonperfect field of characteristic  $p > 0$ , let  $t \in k - k^p$ , let  $\bar{k}$  be an algebraic closure of  $k$ , and let  $\alpha \in \bar{k}$  satisfy  $\alpha^p = t$ .

<sup>14</sup>Editor's note: part (1), which is useful in Examples 1.3.2 below, has been added.

<sup>15</sup>Editor's note: the following sentence and Examples 1.3.2 have been added.

- (1) Consider the additive group  $\mathbb{G}_{a,k} = \operatorname{Spec} k[X]$ , and let  $G$  be the group subscheme, finite over  $k$ , defined by the additive polynomial  $X^{p^2} - tX^p$ . Then

$$G_{\text{red}} = \operatorname{Spec} k[X]/X(X^{p(p-1)} - t)$$

is étale at the origin. If it were a group scheme over  $k$ , it would be smooth (by 1.3.1); but  $G_{\text{red}}$  is not geometrically reduced, so it is not a group scheme over  $k$ .

- (2) Consider

$$\mathbb{G}_{a,k}^4 = \operatorname{Spec} k[X, Y, U, V],$$

and let  $G$  be the group subscheme defined by the ideal  $I$  generated by the additive polynomials

$$P = X^p - tY^p \quad \text{and} \quad Q = U^p - tV^p.$$

Then  $G$  is two-dimensional and irreducible, since

$$(G_{\bar{k}})_{\text{red}} \simeq \operatorname{Spec} \bar{k}[Y, V]$$

is irreducible.

Let  $A = k[X, Y, U, V]$ , and let  $\mathfrak{m}$  be its augmentation ideal. Denote by  $x, y, u, v$  the images of  $dX, dY, dU, dV$  in

$$\Omega_{A/k}^1 \otimes_A (A/\mathfrak{m}),$$

viewed as linear forms on the tangent space

$$k^4 = T_0 \mathbb{G}_{a,k}^4.$$

We shall show that the subspace  $E = T_0 G_{\text{red}}$  equals  $k^4$ . Otherwise, there would be a linear form

$$f = ax + by + a'u + b'v, \quad a, b, a', b' \in k,$$

not all of whose coefficients are zero, that vanishes on  $E$ . Recall that the formation of  $\Omega_{A/k}^1$ , and hence that of tangent spaces, commutes with base change (cf. EGA IV<sub>4</sub>, 16.4.5), and identify  $f$  with its image in  $(\bar{k}^4)^*$ . Since

$$(G_{\bar{k}})_{\text{red}} \subset (G_{\text{red}})_{\bar{k}},$$

the form  $f$  vanishes on the subspace  $T_0(G_{\bar{k}})_{\text{red}}$  of  $\bar{k}^4$ , which is defined by the equations

$$g_1 = x - \alpha y \quad \text{and} \quad g_2 = u - \alpha v.$$

Thus

$$f = \lambda g_1 + \mu g_2, \quad \lambda, \mu \in \bar{k}.$$

But  $\lambda g_1 + \mu g_2$  belongs to  $k^4$  only if  $\lambda = \mu = 0$ . This contradiction shows that  $E = k^4$ , and hence

$$T_0(G_{\text{red}})_{\bar{k}} = \bar{k}^4.$$

On the other hand,  $R = XV - YU$  belongs to  $\sqrt{I}$ , since

$$R^p = (X^p - tY^p)V^p - Y^p(U^p - tV^p).$$

Consequently, the tangent space  $F$  at the point  $(\alpha, 1, \alpha, 1)$  of  $(G_{\text{red}})_{\bar{k}}$  is contained in the hyperplane  $H$  of  $\bar{k}^4$  with equation

$$\alpha dV + dX - dU - \alpha dY = 0,$$

and therefore has dimension at most 3.<sup>16</sup> It follows from part (1) of 1.3.1 that  $G_{\text{red}}$  is not a group scheme over  $k$ .

---

<sup>16</sup>Editor's note: in fact, it is easy to see that  $\sqrt{I}$  is generated by  $P, Q, R$  at every point other than 0, so  $F = H$ .

## 2. Connected components of an $A$ -group locally of finite type

**2.1.** Let us first consider an arbitrary  $A$ -group  $G$ , and let  $G'$  be the connected component of the origin  $e$  in  $G$ . This connected component is plainly closed, so we may identify it with the reduced closed subscheme of  $G$  having  $G'$  as its underlying space.<sup>17</sup>

**Proposition 2.1.1.** *For every extension  $K$  of the residue field  $k$  of  $A$ , the underlying space of  $G' \otimes_A K$  is the connected component of the origin in the  $K$ -group  $G \otimes_A K$ ; in other words,  $G'$  is geometrically connected.*

Indeed, let  $(G \otimes_A K)'$  be the connected component of the origin in  $G \otimes_A K$ . Since the image of  $(G \otimes_A K)'$  in  $G$  is connected and contains the unit element of  $G$ , that image is contained in  $G'$ . Thus  $(G \otimes_A K)'$  is contained in the inverse image  $G' \otimes_A K$  of  $G'$  in  $G \otimes_A K$ . The proposition therefore follows from the connectedness of  $G' \otimes_A K$ , which is proved in the following lemma.

**Lemma 2.1.2.** *Let  $X$  and  $Y$  be two connected schemes over a field  $k$ . If  $X$  contains a rational point, then  $X \times_k Y$  is connected.*

We give a direct proof below of this result from EGA IV<sub>2</sub> (4.5.8 and 4.5.14).

Suppose first that  $Y$  is nonempty, connected, and affine, with algebra  $B$ . Then  $X \times_k Y$  is the spectrum of the quasi-coherent  $\mathcal{O}_X$ -algebra

$$\mathcal{B} = \mathcal{O}_X \otimes_k B.$$

We wish to show that every nonempty subset  $U$  of  $X \times_k Y$  that is both open and closed equals  $X \times_k Y$ . Now  $U$  is affine over  $X$ , and its affine  $\mathcal{O}_X$ -algebra is a direct factor of  $\mathcal{B}$ . It follows from Lemma 2.1.3 below that the image of  $U$  in  $X$  is open and closed, and hence is all of  $X$ . In particular, this image contains a rational point  $x$  of  $X$ , so  $U$  meets the inverse image of  $x$  in  $X \times_k Y$ . Since this inverse image is isomorphic to  $Y$ , and hence connected,  $U$  contains the whole inverse image. The same conclusion would hold for the complement of  $U$  in  $X \times_k Y$  if  $U$  were different from  $X \times_k Y$ , which is absurd.

If  $Y$  is now an arbitrary  $k$ -scheme, the preceding argument shows that the fibers of the canonical projection

$$X \times_k Y \longrightarrow Y$$

are connected. If  $x$  is a rational point of  $X$ , all these fibers meet the subscheme  $\{x\} \times_k Y$ , which is itself connected. This proves the assertion.

**Lemma 2.1.3.** *Let  $X$  be a scheme, and let  $\mathcal{A}$  be a quasi-coherent  $\mathcal{O}_X$ -algebra that is locally<sup>18</sup> a direct factor of a free  $\mathcal{O}_X$ -module. Then the image of  $\text{Spec } \mathcal{A}$  in  $X$  is open and closed.*

Let  $V$  be this image. It is clear that  $V$  is contained in the support of  $\mathcal{A}$ . Conversely, if  $x$  belongs to the support of  $\mathcal{A}$ , then  $\mathcal{A}_x$  is nonzero and is a free  $\mathcal{O}_{X,x}$ -module, since, by Kaplansky, every projective module over a local ring is free. Consequently, the fiber of  $\text{Spec } \mathcal{A}$  at  $x$ , which is affine with algebra

$$\mathcal{A}_x \otimes_{\mathcal{O}_{X,x}} \kappa(x),$$

is nonempty. Thus the image of  $\text{Spec } \mathcal{A}$  coincides with the support of  $\mathcal{A}$ .

If  $s$  is the unit section of  $\mathcal{A}$ , the equality  $s_x = 0$  implies that  $s$ , and hence  $\mathcal{A}$ , vanish in a neighborhood of  $x$ . Therefore the support of  $\mathcal{A}$  is closed. On the other hand, we may

<sup>17</sup>Editor's note: we shall see below (2.2) that  $G'$  is a group in the category  $(\mathbf{Sch}/k)_{\text{red}}$ .

<sup>18</sup>Editor's note: the word "locally" has been added. The reference to Kaplansky is *Projective modules*, Ann. of Math. **68** (1958), 372–377; see also [BAC], § II.3, Ex. 3.

assume that  $\mathcal{A}$  is a direct factor of the direct sum  $\mathcal{L}$  of a family  $(\mathcal{O}_X^\alpha)$  of copies of  $\mathcal{O}_X$ .<sup>19</sup> For each  $\beta$ , let  $\varphi^\beta$  denote the restriction to  $\mathcal{A}$  of the canonical projection

$$\mathcal{L} = \bigoplus_{\alpha} \mathcal{O}_X^\alpha \longrightarrow \mathcal{O}_X^\beta.$$

For  $x \in X$ , let  $\mathcal{A}_x^*$  denote the dual module

$$\mathrm{Hom}_{\mathcal{O}_{X,x}}(\mathcal{A}_x, \mathcal{O}_{X,x}).$$

Since  $\mathcal{A}_x$  is a direct factor of  $\mathcal{L}_x$ , the restriction map

$$\mathrm{Hom}_{\mathcal{O}_{X,x}}(\mathcal{L}_x, \mathcal{O}_{X,x}) \longrightarrow \mathrm{Hom}_{\mathcal{O}_{X,x}}(\mathcal{A}_x, \mathcal{O}_{X,x}) = \mathcal{A}_x^*$$

is surjective. If  $\mathcal{A}_x \neq 0$ , then, since  $\mathcal{A}_x$  is free and nonzero, there exist  $a \in \mathcal{A}_x$  and a linear form  $\varphi \in \mathcal{A}_x^*$  such that  $\varphi(a) = 1$ . There is therefore a family  $(\xi^\alpha)$  of elements of  $\mathcal{O}_{X,x}$  such that, for every  $u \in \mathcal{A}_{X,x}$ ,

$$\varphi(u) = \sum_{\alpha} \xi^\alpha \varphi^\alpha(u);$$

this sum is finite because  $\varphi^\alpha(u) = 0$  except for finitely many  $\alpha$ . Applying this to  $u = a$ , we find  $\alpha_1, \dots, \alpha_n$  such that

$$1 = \varphi(a) = \xi^{\alpha_1} \varphi^{\alpha_1}(a) + \dots + \xi^{\alpha_n} \varphi^{\alpha_n}(a).$$

There is then an open neighborhood  $U$  of  $x$  on which  $a$  and the  $\xi^{\alpha_i}$  come from sections

$$\tilde{a} \in \mathcal{A}(U) \quad \text{and} \quad \tilde{\xi}^{\alpha_i} \in \mathcal{O}_X(U).$$

The equality

$$\sum_i \tilde{\xi}^{\alpha_i} \varphi^{\alpha_i}(\tilde{a}) = 1 \quad \text{on } U$$

shows that  $\tilde{a}_y \neq 0$  for every  $y \in U$ . (“The support of a projective module is open.”)

**2.2.** The notation remains that of 2.1. It is clear that  $G'$  is a reduced  $k$ -scheme. Lemma 2.1.2 shows that  $G' \times_k G'$  is connected, so  $(G' \times_k G')_{\mathrm{red}}$  is the reduced subscheme of  $G \times_A G$  whose underlying space is the connected component of the origin. In particular, the multiplication morphism

$$\mu : G \times_A G \longrightarrow G$$

induces a morphism

$$\mu' : (G' \times_k G')_{\mathrm{red}} \longrightarrow G'$$

that makes  $G'$  a group in  $(\mathbf{Sch}/k)_{\mathrm{red}}$ .

**2.2.bis.**<sup>20</sup> Recall (cf. Exposé V) that, if  $P$  is a scheme,  $\underline{P}$  denotes the underlying topological space of  $P$ . We define an  $A$ -subfunctor  $G^0$  of  $G$  by setting, for every  $A$ -scheme  $S$ ,

$$G^0(S) = \{u \in G(S) \mid u(\underline{S}) \subset \underline{G'}\}.$$

Let  $c : G \rightarrow G$  be the inversion morphism. Since  $c(\underline{G'}) = \underline{G'}$ , we have  $c \circ u \in G^0(S)$  for every  $u \in G^0(S)$ . On the other hand, if  $u, v \in G^0(S)$ , then  $u \boxtimes v$  sends  $\underline{S}$  into the subspace of  $\underline{G \times_A G}$  consisting of the points whose two projections belong to  $\underline{G'}$ . This subspace is identified with the underlying space of  $G' \times_A G'$ , which is connected by Lemma 2.1.2.

<sup>19</sup>Editor's note: the original has been expanded in what follows.

<sup>20</sup>Editor's note: this paragraph has been added in order to make the connection with Exposé VI<sub>B</sub>, § 3.

Consequently,  $\mu \circ (u \boxtimes v)$  sends  $\underline{S}$  into  $\underline{G}'$ . This proves that  $G^0$  is a group subfunctor of  $G$  over  $A$ .

If the connected component of  $e$  is open in  $G$ , then the subfunctor  $G^0$  is represented by the subscheme induced by  $G$  on this open subset, which is therefore a group subscheme of  $G$ ; this subscheme will also be denoted by  $G^0$ . In this case, with the notation of 2.1,

$$G' = (G^0)_{\text{red}},$$

and the topological spaces  $\underline{G}'$  and  $\underline{G}^0$  coincide.<sup>21</sup>

**2.3.** In accordance with our convention in 1.1, we again assume from now on that  $G$  is locally of finite type over  $A$ . Then  $G$  is locally Noetherian, hence locally connected;<sup>22</sup> thus every connected component of  $G$  is open.

We shall write  $G^0$  for the subscheme induced by  $G$  on the connected component  $G'$  of the unit element. By 2.2.bis,  $G^0$  is a group subscheme of  $G$ , which we shall call the *neutral component* of  $G$ . For every  $A$ -scheme  $S$ , therefore,

$$G^0(S) = \{u \in G(S) \mid u(\underline{S}) \subset \underline{G}^0 = \underline{G}'\}.$$

Let  $G^\alpha$  be an arbitrary connected component of  $G$ , and let

$$\nu^\alpha : G^\alpha \times_A G^0 \longrightarrow G$$

be the morphism defined by

$$\nu^\alpha(S)(g, \gamma) = g\gamma g^{-1}$$

for every  $S \in (\mathbf{Sch}/_A)$ ,  $g \in G^\alpha(S)$ , and  $\gamma \in G^0(S)$ . If  $e$  is the origin of  $G$ , the restriction of  $\nu^\alpha$  to  $G^\alpha \times_A \{e\}$  is the trivial morphism. Since  $G^\alpha \times_A G^0$  is connected by 2.1.2, it follows that  $\nu^\alpha$  factors through  $G^0$ . Thus, for every  $A$ -scheme  $S$ ,  $G^0(S)$  is a normal subgroup of  $G(S)$ . We have proved the following proposition.<sup>23</sup>

**Proposition 2.3.1.** *Let  $G$  be an  $A$ -group locally of finite type. Then the neutral component  $G^0$  is an open normal group subscheme of  $G$ .*

**Proposition 2.4.** *Let  $G$  be an  $A$ -group locally of finite type.*

- (i)  $G^0$  is irreducible, and  $G^0 \otimes_A k$  is geometrically irreducible over  $k$ .
- (ii)  $G^0$  is quasi-compact, and hence is of finite type over  $A$ .

<sup>24</sup> *Proof.* (i) Since  $G^0$  and  $G^0 \otimes_A k$  have the same underlying topological space, it suffices to prove the second assertion. Let  $\bar{k}$  be an algebraic closure of  $k$ . By 2.2,  $(G \otimes_A \bar{k})_{\text{red}}$  is a reduced  $\bar{k}$ -group locally of finite type, and hence is smooth over  $\bar{k}$  by 1.3.1. A fortiori, the local rings of  $(G \otimes_A \bar{k})_{\text{red}}$  are integral domains. Therefore,<sup>25</sup> since  $G \otimes_A \bar{k}$  is locally Noetherian, the connected components of  $G \otimes_A \bar{k}$  are irreducible (cf. EGA I, 6.1.10). In particular, the connected component  $G^0 \otimes_A \bar{k}$  (cf. 2.1.1) is irreducible.

<sup>21</sup>Editor's note: in this exposé, the notation  $G^0$  is reserved for the case in which the connected component of  $e$  is open; in Exposé VI<sub>B</sub>, § 3, this connected component will be denoted by  $\underline{G}^0$  in every case. This is a slight abuse of notation, but it is compatible with the preceding discussion when the connected component is the underlying topological space of an open group subscheme  $G^0$  of  $G$ .

<sup>22</sup>Editor's note: indeed, in a Noetherian space there are only finitely many connected components, and each is therefore open; see also EGA I, 6.1.9.

<sup>23</sup>Editor's note: the number 2.3.1 has been added in order to emphasize this statement. Moreover,  $G^0$  is even a characteristic subgroup of  $G$ ; see 2.6.5(ii).

<sup>24</sup>Editor's note: in the statement, " $G^0$  is geometrically irreducible" has been replaced by " $G^0 \otimes_A k$  is geometrically irreducible over  $k$ ," and the proof has been expanded.

<sup>25</sup>Editor's note: the original has been expanded here, and the reference to EGA I, 6.1.10, has been added.

(ii) We now show that  $G^0$  is of finite type over  $A$ . Since  $G^0$  is locally of finite type over  $A$ , it suffices to prove that  $G^0$  is quasi-compact. Since  $G^0$  is irreducible, this follows from 0.5.1.

**Corollary 2.4.1.** *Every connected component of  $G$  is irreducible,<sup>26</sup> of finite type over  $A$ ,<sup>27</sup> and of the same dimension as  $G^0$ .*

Indeed, we may assume that  $A$  equals its residue field  $k$ . Let  $C$  be a connected component of  $G$ , let  $x$  be a closed point of  $C$ , let  $\kappa(x)$  be the residue field of  $x$ , and let  $k'$  be a finite normal extension of  $k$  containing  $\kappa(x)$ . The canonical projection

$$\pi : C \otimes_k k' \longrightarrow C$$

is open and closed.<sup>28</sup> Consequently, if  $C'$  is a connected component of  $C \otimes_k k'$ , the projection  $C' \rightarrow C$  is surjective. Thus  $C'$  contains a point  $y \in \pi^{-1}(x)$ , and such a point is rational over  $k'$  (cf. the proof of 1.2); it follows that  $C'$  is the image of  $G^0 \otimes_k k'$  under right translation  $r_y$ . Now  $G^0 \otimes_k k'$  is of finite type over  $k'$ , by 2.4, and  $\pi^{-1}(x)$  is finite, of cardinality at most  $[k' : k]$ . Consequently,  $C \otimes_k k'$  is of finite type over  $k'$ , and hence  $C$  is of finite type over  $k$ .

Moreover, since  $G^0 \otimes_k k'$  is irreducible by 2.4, the same is true of  $C'$ ,<sup>29</sup> and hence also of  $C$ , because the projection  $C' \rightarrow C$  is surjective.

Finally, we saw above that  $C \otimes_k k'$  is the disjoint union of finitely many translates of  $G^0 \otimes_k k'$ . Since dimension is invariant under extension of the base field (cf. EGA IV<sub>2</sub>, 4.1.4), it follows that  $C$  has the same dimension as  $G^0$ . (Moreover, by EGA IV<sub>2</sub>, 5.2.1, one has  $\dim_g G = \dim G^0$  for every point  $g \in G$ .)

**2.5.** This paragraph has been added. The results that follow appear in Exposé VI<sub>B</sub>, but could (or should) have appeared in VI<sub>A</sub>; it is useful to have them available already, in order to make Theorem 3.2 below more precise.

**Lemma 2.5.1.** *Let  $(A, \mathfrak{m})$  be an Artinian local ring, and let  $k = A/\mathfrak{m}$  be its residue field.*

- (i) *If  $X$  is an  $A$ -scheme such that  $X \otimes_A k$  is locally of finite type (respectively, of finite type) over  $k$ , then  $X$  has the same property over  $A$ .*
- (ii) *Let  $u : X \rightarrow Y$  be a morphism of  $A$ -schemes. If  $u \otimes_A k$  is an immersion (respectively, a closed immersion), then so is  $u$ .*

*Proof.* (i) Suppose that  $X \otimes_A k$  is locally of finite type over  $k$ . Let  $U = \operatorname{Spec} B$  be an affine open subset of  $X$ . By hypothesis, there are elements  $x_1, \dots, x_n$  of  $B$  whose images generate  $B/\mathfrak{m}B$  as a  $k$ -algebra, and the “nilpotent Nakayama lemma” then shows that the  $x_i$  generate  $B$  as an  $A$ -algebra. This proves that  $X$  is locally of finite type over  $A$ . If, in addition,  $X \otimes_A k$  is quasi-compact, then so is  $X$ , which has the same underlying topological space; hence  $X$  is of finite type over  $A$ . This proves (i).

We prove (ii). Suppose that  $u \otimes_A k$  is an immersion (respectively, a closed immersion). Then  $u$  is a homeomorphism of  $X$  onto a locally closed (respectively, closed) subset of  $Y$ , and, for every  $x \in X$ , the ring homomorphism

$$\varphi_x : \mathcal{O}_{Y, u(x)} \longrightarrow \mathcal{O}_{X, x}$$

<sup>26</sup>Editor’s note: one should bear in mind that a nonneutral connected component is not geometrically connected in general. For example, if  $k = \mathbb{R}$ , the group  $\mu_{3, \mathbb{R}}$ , represented by  $\mathbb{R}[X]/(X^3 - 1)$ , has two connected components:  $\{e\} = \operatorname{Spec} \mathbb{R}$  and  $C = \operatorname{Spec} \mathbb{R}[X]/(X^2 + X + 1)$ ; and  $C \otimes_{\mathbb{R}} \mathbb{C}$  has two components.

<sup>27</sup>Editor’s note: the assertion that follows has been added; cf. Exposé VI<sub>B</sub>, 1.5.

<sup>28</sup>Editor’s note: the original has been expanded in what follows.

<sup>29</sup>Editor’s note: the original has been simplified here.



is such that  $\varphi_x \otimes_A k$  is surjective. The nilpotent Nakayama lemma therefore shows that  $\varphi_x$  is surjective, so  $u$  is an immersion (respectively, a closed immersion).

**Proposition 2.5.2.** *Let  $A$  be an Artinian local ring with residue field  $k$ , and let  $u : G \rightarrow H$  be a quasi-compact morphism between  $A$ -groups locally of finite type.*

(a) *The set  $u(G)$  is closed in  $H$ , and its connected components are irreducible and all have the same dimension.*

(b) *One has*

$$\dim G = \dim u(G) + \dim \operatorname{Ker}(u).$$

(c) *If  $u$  is a monomorphism, then it is a closed immersion.*

*Proof.* By the preceding lemma, it suffices to prove the proposition when  $A = k$ . Moreover, the properties under consideration are stable under fpqc descent, and dimension is invariant under extension of the base field, so we may assume that  $k$  is algebraically closed.

We prove (a). Let  $C$  be the reduced subscheme of  $H$  whose underlying topological space is  $u(G)$ . Since  $u(G)$  is stable under the inversion morphism of  $H$ , so is  $C$ . Furthermore,  $u : G \rightarrow C$  is quasi-compact and dominant. Thus, by EGA IV<sub>2</sub>, 2.3.7, the same is true of

$$u \times_k \operatorname{id}_G \quad \text{and} \quad \operatorname{id}_H \times_k u,$$

and hence of their composite

$$u \times_k u : G \times_k G \longrightarrow C \times_k C.$$

Consequently, multiplication on  $H$  sends  $C \times_k C$  into  $C$ , and therefore  $C$  is a group subscheme of  $H$ .

Replacing  $H$  by  $C$ , we are thus reduced to the case in which  $u$  is dominant. Then  $G(k)$  is dense in  $H$ , and hence meets every connected component of  $H$ ; it therefore acts transitively on the set of those connected components. It is consequently enough to show that  $u(G)$  contains  $H^0$ . Replacing  $G$  by  $u^{-1}(H^0)$ , we may therefore suppose that  $H = H^0$ . In that case, 2.4 shows that  $H$  is irreducible and of finite type over  $k$ , and hence Noetherian. Moreover,  $u$  is locally of finite type (cf. EGA I, 6.6.6) and quasi-compact, and is therefore of finite type. By Chevalley's constructibility theorem (cf. EGA IV<sub>1</sub>, 1.8.5),  $u(G)$  is consequently a constructible (and dense) subset of

$$H = \overline{u(G)},$$

and thus contains a dense open subset  $U$  of  $H$  (cf. EGA 0<sub>III</sub>, 9.2.2). By 0.5, we then have

$$H = U \cdot U \subset u(G),$$

whence  $u(G) = H$ . Together with 2.4.1, this proves (a).

We prove (b). Recall first that the functor  $\operatorname{Ker}(u)$  (cf. I, 2.3.6.1) is represented by  $u^{-1}(e)$ , where  $e$  denotes the identity element of  $H$ . Since  $u$  is of finite type,  $\operatorname{Ker}(u)$  is of finite type over  $k$ . On the other hand, replacing  $H$  by the reduced closed subscheme  $u(G)$ , we may assume that  $u$  is surjective. Let  $u^0$  denote the restriction of  $u$  to  $G^0$ . Since  $G$  and  $\operatorname{Ker}(u)$  are equidimensional, and since

$$\operatorname{Ker}(u)^0 \subset \operatorname{Ker}(u^0),$$

we are reduced to the case in which  $G$ , and hence also  $H$ , is irreducible.

By EGA IV<sub>3</sub>, 9.2.6.2 and 10.6.1(ii), the set of  $y \in H$  such that

$$\dim u^{-1}(y) = \dim G - \dim H$$

contains a nonempty open subset  $V$ . Since  $u$  is surjective,  $U = u^{-1}(V)$  is then a nonempty open subset of  $G$ , and therefore contains a closed point  $x$  of  $G$ , because  $G$  is a Jacobson scheme (cf. EGA IV<sub>3</sub>, 10.4.8). Right translation  $r_x$  is an isomorphism from  $\text{Ker}(u)$  onto  $u^{-1}(u(x))$ , whence

$$\dim \text{Ker}(u) = \dim u^{-1}(u(x)) = \dim G - \dim H.$$

We prove (c), following [DG70], I, § 3.4. (Another proof is given in Exposé VI<sub>B</sub>, 1.4.2.) Suppose that  $u$  is a monomorphism. If  $C$  is a connected component of  $G$ , there is a closed point  $x \in G$  such that  $C = r_x(G^0)$ . If  $u_C$  (respectively,  $u^0$ ) denotes the restriction of  $u$  to  $C$  (respectively, to  $G^0$ ), then

$$u_C = r_{u(x)} \circ u^0 \circ r_x^{-1}.$$

It therefore suffices to show that  $u^0$  is a closed immersion. We may thus suppose that  $G = G^0$ , so that  $G$  is irreducible and of finite type over  $k$ .

Let  $\xi$  be the generic point of  $G$ . Then  $\mathcal{O}_{G,\xi}$  is an Artinian local ring; denote its maximal ideal by  $\mathfrak{m}$ . Let  $h = u(\xi)$ , let  $\mathfrak{n}$  be the maximal ideal of  $\mathcal{O}_{H,h}$ , and set

$$A = \mathcal{O}_{G,\xi} / \mathfrak{n} \mathcal{O}_{G,\xi}.$$

Since  $u$  is a monomorphism, so is the morphism

$$u_h : \text{Spec}(A) \longrightarrow \text{Spec}(\kappa(h))$$

obtained by base change. The multiplication morphism

$$A \otimes_{\kappa(h)} A \longrightarrow A$$

is therefore an isomorphism (cf. EGA I, 5.3.8), whence  $A = \kappa(h)$ . Nakayama's lemma then shows that

$$\mathcal{O}_{H,h} \longrightarrow \mathcal{O}_{G,\xi}$$

is surjective, since  $\mathfrak{n} \mathcal{O}_{G,\xi}$  is contained in  $\mathfrak{m}$  and is therefore nilpotent.

Let  $V$  be an affine open subset of  $H$  containing  $h$ , let  $U \neq \emptyset$  be an affine open subset of  $G$  contained in  $u^{-1}(V)$ , and let

$$\varphi : \mathcal{O}_H(V) \longrightarrow \mathcal{O}_G(U)$$

be the homomorphism of  $k$ -algebras induced by  $u$ . Let  $\mathfrak{p}$  be the prime ideal of  $\mathcal{O}_G(U)$  corresponding to  $\xi$ ,\* and set  $\mathfrak{q} = \varphi^{-1}(\mathfrak{p})$ . Since  $G$  is of finite type over  $k$ , the  $k$ -algebra  $\mathcal{O}_G(U)$  is generated by finitely many elements  $a_1, \dots, a_n$ . By the preceding argument, there are elements  $b_1, \dots, b_n, s \in \mathcal{O}_H(V)$ , with  $s \notin \mathfrak{q}$ , such that in  $\mathcal{O}_G(U)_{\mathfrak{p}}$  one has

$$a_i/1 = \varphi(b_i)/\varphi(s).$$

There are therefore elements

$$t_1, \dots, t_n \in \mathcal{O}_G(U) - \mathfrak{p}$$

such that

$$t_i(a_i\varphi(s) - \varphi(b_i)) = 0.$$

Set

$$t = t_1 \cdots t_n \varphi(s) \in \mathcal{O}_G(U) - \mathfrak{p}.$$

---

\*Translator's note: the French source prints  $\eta$  here; the generic point introduced immediately above is  $\xi$ .

The equalities  $a_i/1 = \varphi(b_i)/\varphi(s)$  then already hold in  $\mathcal{O}_G(U)_t$ . Since

$$t \in \mathcal{O}_G(U) = k[a_1, \dots, a_n],$$

there is an element  $b \in \mathcal{O}_H(V)$  such that, for some  $r \in \mathbb{N}$ ,

$$t/1 = \varphi(b)/\varphi(s)^r.$$

Consequently,  $\varphi$  induces a surjection

$$\mathcal{O}_H(V)_{sb} \longrightarrow \mathcal{O}_G(U)_t,$$

and hence  $u$  is a local immersion at  $\xi$ .

The open subset  $W$  of  $G$  consisting of the points at which  $u$  is a local immersion is therefore nonempty. Since  $G$  is a Jacobson scheme,  $W$  contains a closed point  $y$ . To prove that  $W = G$ , it is enough to show that every closed point of  $G$  belongs to  $W$ . But every closed point  $x$  is the image of  $y$  under the translation  $r_x \circ r_y^{-1}$ , and hence belongs to  $W$ . Thus  $W = G$ , which proves that  $u$  is a local immersion.

Since  $G$  is irreducible, it follows that  $u$  is an immersion. Indeed, for every  $x \in G$ , let  $U_x$  and  $V_x$  be open subsets of  $G$  and  $H$ , respectively, such that  $x \in U_x$  and  $u$  induces a closed immersion of  $U_x$  into  $V_x$ . Since  $U_x$  is dense in  $G$ , the subset  $u(U_x)$  is dense in  $u(G) \cap V_x$ . But  $u(U_x)$  is closed in  $V_x$ , so

$$u(U_x) = V_x.$$

Moreover, since  $u$  is injective,  $U_x = u^{-1}(V_x)$ . Thus  $u$  induces a closed immersion of  $G$  into the open subscheme of  $H$  covered by the  $V_x$ . Hence  $u : G \rightarrow H$  is an immersion. We have already seen that  $u(G)$  is closed in  $H$ , so  $u$  is a closed immersion.

**Lemma 2.5.3.**<sup>30</sup> *Let  $A$  be an Artinian local ring, let  $k$  be its residue field, let  $G$  be a flat  $A$ -group, and let  $X$  be an  $A$ -scheme equipped with a left action*

$$\mu : G \times_A X \longrightarrow X$$

*of  $G$  and with a section  $s_0 : \operatorname{Spec} A \rightarrow X$ . (This is the case, for example, if  $G'$  is a second  $A$ -group and one is given a morphism of  $A$ -groups  $G \rightarrow G'$ .)*

*Let  $\varphi$  be the morphism*

$$\mu \circ (\operatorname{id}_G \times s_0) : G = G \times_A A \longrightarrow X.$$

*If  $\varphi$  is flat at a point  $g$  of  $G$ , then  $\varphi$  is flat.*

*Proof.* Since  $G$  is flat over  $A$ , the fiberwise criterion for flatness (EGA IV<sub>3</sub>, 11.3.10.2) shows that it is enough to prove that  $\varphi \otimes_A k$  is flat. We may therefore suppose that  $A = k$ . In that case, giving  $s_0$  is equivalent to giving a  $k$ -point  $x_0 \in X(k)$ , and  $\varphi$  is the morphism  $h \mapsto hx_0$ .

Let  $h \in G$ ; we show that  $\varphi$  is flat at  $h$ . Let  $K$  be an extension of  $k$  containing copies of both  $\kappa(g)$  and  $\kappa(h)$ . We have a cartesian square

$$\begin{array}{ccc} G_K & \xrightarrow{\varphi_K} & X_K \\ \downarrow & & \downarrow \\ G & \xrightarrow{\varphi} & X \end{array}$$

<sup>30</sup>Editor's note: no finiteness hypotheses are made in this statement; this will be useful later (cf. 6.2 and Exposé VI<sub>B</sub>, § 12).

in which the two vertical arrows are faithfully flat. Thus, by Exposé V, 7.4(i),  $\varphi_K$  is flat at every point  $g' \in G_K$  lying over  $g$ ; and, in order to show that  $\varphi$  is flat at  $h$ , it is enough to show that  $\varphi_K$  is flat at one point  $h'$  lying over  $h$ . We are therefore reduced to the case in which  $g$  and  $h$  are rational.

Set  $u = hg^{-1}$ , and let  $\ell_u$  (respectively,  $\mu_u$ ) be left translation on  $G$  (respectively, on  $X$ ) defined by  $u$ . Since  $\varphi \circ \ell_u = \mu_u \circ \varphi$ , we obtain a commutative square

$$\begin{array}{ccc} \mathcal{O}_{G,h} & \xrightarrow{\sim} & \mathcal{O}_{G,g} \\ \uparrow & & \uparrow \\ \mathcal{O}_{X,hx_0} & \xrightarrow{\sim} & \mathcal{O}_{X,gx_0} \end{array}$$

whose horizontal arrows are isomorphisms. Since the morphism

$$\mathcal{O}_{X,gx_0} \longrightarrow \mathcal{O}_{G,g}$$

is flat, the morphism

$$\mathcal{O}_{X,hx_0} \longrightarrow \mathcal{O}_{G,h}$$

is flat as well.

**Proposition 2.5.4.** *Let  $A$  be an Artinian local ring, let  $k$  be its residue field, let  $G$  be an  $A$ -group locally of finite type, and let  $X \neq \emptyset$  be an  $A$ -scheme locally of finite type equipped with a left action of  $G$ . Suppose that the morphism*

$$\varphi : G \times_A X \longrightarrow X \times_A X, \quad (g, x) \longmapsto (gx, x),$$

*defined on underlying sets by the displayed rule, is surjective. Then:*

- (i) *The connected components of  $X$  are of finite type, irreducible, and all of the same dimension.*
- (ii) *More precisely, let  $\bar{k}$  be an algebraic closure of  $k$ , and let  $x$  be a closed point of  $X \otimes_A \bar{k}$ . Its stabilizer  $F$  is a closed group subscheme of  $G \otimes_A \bar{k}$ , and the dimension of the irreducible components of  $X$  is*

$$\dim G - \dim F.$$

In view of Lemma 2.5.1, we may suppose that  $A = k$ . Suppose first that  $k$  is algebraically closed. Then  $G_{\text{red}}$  is a  $k$ -group locally of finite type; hence, replacing  $G$  by  $G_{\text{red}}$  and  $X$  by  $X_{\text{red}}$ , we may suppose that  $G$  and  $X$  are reduced.

Since  $G \times_k X$  is locally of finite type over  $k$ , the morphism  $\varphi$  is locally of finite type (cf. EGA I, 6.6.6(v)), and is therefore locally of finite presentation because  $X \times_k X$  is locally Noetherian. Let  $x$  be a rational point of  $X$ . The morphism

$$\varphi_x : G \longrightarrow X$$

obtained from  $\varphi$  by base change is surjective and locally of finite presentation. If  $\eta$  is a maximal point of  $X$ , then  $\mathcal{O}_{X,\eta}$  is a field, since  $X$  is reduced; consequently,  $\varphi_x$  is flat at every point of  $G$  lying over  $\eta$ . By Lemma 2.5.3,  $\varphi_x$  is therefore flat. Thus  $\varphi_x : G \rightarrow X$  is faithfully flat and locally of finite presentation, and hence open (cf. EGA IV<sub>2</sub>, 2.4.6).

Since  $G^0$  is open in  $G$ , irreducible, and quasi-compact (by 2.4), every orbit

$$G^0 x = \varphi_x(G^0),$$

as  $x$  ranges over the rational points of  $X$ , is an open, irreducible, and quasi-compact subset of  $X$ , and is therefore of finite type over  $k$ , since  $X$  is locally of finite type over  $k$ .

Every nonempty open subset of  $X$  contains a rational point, so these open subsets cover  $X$ . Furthermore, any two of them are either disjoint or equal. Indeed, if

$$\varphi_x(G^0) \cap \varphi_y(G^0)$$

is nonempty, it contains a rational point  $z$ . There are then rational points  $g, h \in G^0$  such that

$$gx = z = hy,$$

whence  $x = g^{-1}z$  and  $y = h^{-1}z$ , and consequently

$$\varphi_x(G^0) = \varphi_z(G^0) = \varphi_y(G^0).$$

It follows that the orbits  $\varphi_x(G^0)$  are also closed, and are therefore at once the connected components and the irreducible components of  $X$ .

Finally, let  $x, y$  be two rational points of  $X$ . Since  $\varphi_x$  is surjective, there is a rational point  $g \in G$  such that  $y = gx$ . Since  $G^0$  is a normal subgroup of  $G$ , the orbit  $G^0y$  is the image of  $G^0x^*$  under translation  $\ell_g$  on  $X$ ; hence  $G^0y$  and  $G^0x$  have the same dimension.

Moreover, by Exposé I, 2.3.3.1, the stabilizer of  $x$  is represented by the closed subscheme  $F$  of  $G$  defined by the following cartesian square:

$$\begin{array}{ccc} F & \longrightarrow & G \\ \downarrow & & \downarrow \varphi_x \\ \text{Spec } k & \longrightarrow & X \end{array}$$

Thus  $F$  is a  $k$ -group locally of finite type, while  $F \cap G^0$  is a  $k$ -group of finite type containing  $F^0$ . By 2.4.1, the schemes  $F$  and  $F \cap G^0$  are equidimensional, of the same dimension as  $F^0$ . Let

$$C = \varphi_x(G^0)$$

be the irreducible component of  $X$  containing  $x$ . Arguing as in the proof of part (b) of 2.5.2, we obtain

$$\dim C = \dim G^0 - \dim F^0 = \dim G - \dim F.$$

In the general case, that is, when  $k$  is an arbitrary field, let  $\bar{k}$  be an algebraic closure of  $k$ . Let  $C$  be a connected component of  $X$ , and let  $C'$  be a connected component of  $C \otimes_k \bar{k}$ . Then  $C'$  is a connected component of

$$X' = X \otimes_k \bar{k}.$$

The morphism  $\pi : X' \rightarrow X$  is open (cf. EGA IV<sub>2</sub>, 2.4.10), and, since it is integral, it is also closed; therefore  $\pi(C') = C$ . Since  $C'$  is irreducible and quasi-compact,  $C$  is irreducible and quasi-compact, and hence is of finite type over  $k$ , because  $X$  is locally of finite type over  $k$ .

Finally, dimension is invariant under extension of the base field (cf. EGA IV<sub>2</sub>, 4.1.4), so  $\dim C = \dim C'$ . Since all the irreducible components of  $X'$  have the same dimension, the same is true of those of  $X$ .

---

\*Translator's note: the French source prints  $G^0$  here; as the object being translated is an orbit in  $X$ , the intended source expression is  $G^0x$ .

**2.6. Complements.**— This paragraph has been added; it is taken from [Per75], II, §§ 1–2, with complements due to O. Gabber.<sup>31</sup> It shows that the preceding results are valid for every group scheme  $G$  over a field  $k$ . (This will be used in Sections 5, 6, and 7 of Exposé VI<sub>B</sub>.)

Fix a field  $k$ . We begin with the following lemma (loc. cit., II, 2.1.1), which does not appear explicitly in EGA IV<sub>2</sub>, § 4.4 (although it can probably be read between the lines at the beginning of loc. cit., § 4.4.1).

**Lemma 2.6.0.** *Let  $X$  be an irreducible  $k$ -scheme, let  $K$  be an extension of  $k$ , and let  $X'$  be an irreducible component of  $X_K$ . The projection  $X' \rightarrow X$  is surjective.*

*Proof.* Let  $B$  be a transcendence basis of  $K$  over  $k$ , and set  $L = k(B) \subset K$ . By EGA IV<sub>2</sub>, 4.3.2 and 4.4.1,  $X_L$  is irreducible and  $X'$  dominates  $X_L$ . Hence  $X' \rightarrow X_L$  is integral and dominant, and therefore surjective. Since  $X_L \rightarrow X$  is surjective (loc. cit., 4.4.1), so is  $X' \rightarrow X$ .

For the rest of 2.6, fix a  $k$ -group scheme  $G$  and an action  $\mu : G \times X \rightarrow X$  of  $G$  on a  $k$ -scheme  $X$ , satisfying the following condition:

$$\Phi : G \times X \longrightarrow X \times X, \quad (g, x) \longmapsto (gx, x) \quad \text{is surjective.} \quad (\star)$$

(This holds in particular when  $G$  acts on itself by left translations.) For short, we shall say: “Let  $X$  be a  $G$ -scheme satisfying  $(\star)$ .” Finally, write  $k'$  for the perfect closure of  $k$ .

**Proposition 2.6.1.** *Let  $X$  be a  $G$ -scheme satisfying  $(\star)$ .*

- (i)  *$X$  is geometrically pointwise irreducible over  $k$ : for every extension  $K$  of  $k$ , each point  $x \in X_K$  belongs to a unique irreducible component of  $X_K$ .*
- (ii) *Every local ring of  $(X_{k'})_{\text{red}}$  is normal.*
- (iii) *Let  $\eta$  be a maximal point of  $(X_{k'})_{\text{red}}$ , put  $C = \overline{\{\eta\}}$ , and let  $L$  be the algebraic closure of  $k$  in  $\kappa(\eta)$ . Then  $C$  is an  $L$ -scheme and is geometrically irreducible over  $L$ ; that is,  $C \otimes_L K$  is irreducible for every extension  $K$  of  $L$ .*
- (iv) *In particular, if  $x$  is a rational point of  $X$ , then the irreducible component of  $X$  containing  $x$  is geometrically irreducible over  $k$ .*

*Proof.* (i) Since hypothesis  $(\star)$  is preserved by every base change  $k \rightarrow K$ , it is enough to show that every  $x \in X$  belongs to a unique irreducible component of  $X$ . Since  $\text{Spec}(k') \rightarrow \text{Spec}(k)$  is a universal homeomorphism, we may further suppose that  $k$  is perfect. We may then suppose that  $G$  and  $X$  are reduced.

Let  $\eta$  be a maximal point of  $X$ , and let  $z$  be any point of  $Z = \overline{\{\eta\}}$ . Since  $X$  is reduced,  $\mathcal{O}_{X,\eta} = \kappa(\eta)$ ; and, since  $k$  is perfect,  $\kappa(\eta) \otimes_k K$  is normal for every extension  $K$  of  $k$  (cf. EGA IV<sub>2</sub>, 6.14.2). Thus every point of  $X_K$  lying above  $\eta$  is normal.

Since  $\Phi$  is surjective, there is a point  $\gamma$  of  $G \times X$  such that the two projections of  $\Phi(\gamma)$  are  $z$  and  $\eta$ . Set  $K = \kappa(\gamma)$ . There are then rational points  $g \in G_K$  and  $\eta' \in X_K$ , with  $\eta'$  lying above  $\eta$ , such that  $z' = g\eta'$  lies above  $z$ . By the preceding paragraph,  $\eta'$  is a normal point of  $X_K$ , and therefore so is  $z'$ . Since  $\pi : X_K \rightarrow X$  is flat, it follows that  $z = \pi(z')$  is a normal point of  $X$  (cf. EGA IV<sub>2</sub>, 2.1.13). This proves (ii) and (i).

The first assertion of (iii) now follows from (ii). Since  $L$  is algebraically closed in  $\kappa(\eta)$ ,  $C$  is geometrically irreducible over  $L$ , by EGA IV<sub>2</sub>, 4.5.9.

<sup>31</sup>Editor’s note: these results were communicated to us by O. Gabber, in particular 2.6.6, which plays an important role in Section 5 of Exposé VI<sub>B</sub>.

Finally, if  $X$  has a rational point  $x$ , it follows from (iii) that  $L = k$ , and hence the irreducible component  $C$  containing  $x$  is geometrically irreducible over  $k$ .<sup>\*</sup> This can also be seen directly as follows (cf. [Per75], II, 2.1). Let  $C$  be the irreducible component of  $X$  containing  $x$ , and let  $K$  be an extension of  $k$ . The scheme  $X_K$  has a unique point  $x_K$  above  $x$ , and by (i),  $x_K$  belongs to a unique irreducible component  $C'$  of  $X_K$ . On the other hand, by 2.6.0, every irreducible component of  $C_K$  contains  $x_K$ , and therefore equals  $C'$ .

**Notation.**— Write provisionally  $C^0$  for the reduced closed subscheme of  $G$  whose underlying space is the unique irreducible component of  $G$  containing the identity element  $e$ .

**Corollary 2.6.2.**  $C^0$  is geometrically irreducible over  $k$  and is set-theoretically stable under the group law; that is,  $(C_{k'}^0)_{\text{red}}$  is a subgroup of  $(G_{k'})_{\text{red}}$ . Consequently,  $C^0$  is quasi-compact.

*Proof.* By 2.6.1,  $C^0$  is geometrically irreducible over  $k$ , so  $C^0 \times C^0$  is irreducible and

$$\nu(C^0 \times C^0) \subset C^0,$$

where  $\nu$  denotes the morphism  $(g, h) \mapsto gh^{-1}$ . Since  $\text{Spec } k' \rightarrow \text{Spec } k$  is a universal homeomorphism, the same conclusion holds for  $C_{k'}^0$ , and then for

$$H = (C_{k'}^0)_{\text{red}}.$$

Since  $H \times_{k'} H$  is reduced,  $\nu$  induces a morphism

$$H \times_{k'} H \longrightarrow H;$$

thus  $H$  is a subgroup of  $(G_{k'})_{\text{red}}$ . Consequently, by 0.5.1,  $H$ , and hence  $C^0$ , is quasi-compact.

**Recall 2.6.3.**— Let  $Y$  be a scheme. Recall (EGA 0<sub>III</sub>, 9.1.1) that a subset  $E$  of  $Y$  is called *retrocompact* if the inclusion  $E \hookrightarrow Y$  is quasi-compact. Moreover, by EGA IV<sub>1</sub>, 1.9.5(v) and 1.10.1, if  $U$  is a retrocompact open subset of  $Y$ , its closure  $\overline{U}$  is the union of the closures  $\overline{\{y\}}$  of the points  $y \in U$ . Of course, it is enough to let  $y$  run through the maximal points of  $U$ , that is, the maximal points of  $Y$  contained in  $U$ .

Consequently, if  $Y$  is quasi-separated and  $\eta$  is a maximal point of  $Y$ , then the intersection of the closed neighborhoods of  $\eta$  equals  $\overline{\{\eta\}}$ . Indeed, if  $y \in Y - \overline{\{\eta\}}$ , then  $y$  belongs to an affine open  $V$  not containing  $\eta$ . Since  $Y$  is quasi-separated,  $V$  is retrocompact, and therefore  $\overline{V}$  is the union of the  $\overline{\{\xi\}}$ , for  $\xi$  running through the maximal points of  $Y$  belonging to  $V$ . Hence  $\eta \notin \overline{V}$ , so  $Y - \overline{V}$  is a closed neighborhood of  $\eta$  not containing  $y$ .

**Proposition 2.6.4.** Let  $X$  be a  $G$ -scheme satisfying  $(\star)$ , and let  $U$  be an open subset of  $X$ .

- (i)  $C^0U$  is an open subset of  $X$ , equal to the union of the irreducible components of  $X$  whose generic point belongs to  $U$ .
- (i') Consequently, if  $X$  is irreducible, it is quasi-compact.
- (ii) If, moreover,  $U$  is retrocompact in  $X$ , then  $C^0U = \overline{U}$ , and this subset is both open and closed in  $X$ .

*Proof.* (i) First,  $C^0U$  is open, since it is the union, for  $g \in C^0$ , of the projections of the open subsets

$$g \cdot U_{\kappa(g)} \subset X_{\kappa(g)},$$

and every projection  $X_{\kappa(g)} \rightarrow X$  is open.

---

<sup>\*</sup>Translator's note: the French source prints  $X$  in this conclusion and then switches from the point  $x$  to undefined symbols  $e$  and  $e_K$ . The statement and argument require the component  $C$  and the points  $x$  and  $x_K$ , respectively.

To prove the second assertion of (i), we may replace  $k$  by  $k'$  (since  $\operatorname{Spec} k' \rightarrow \operatorname{Spec} k$  is a universal homeomorphism), and therefore suppose that  $k$  is perfect. We may then suppose that  $G$  and  $X$  are reduced, and hence geometrically reduced.

Let  $\eta$  be a maximal point of  $X$  contained in  $U$ , and put  $Z = \overline{\{\eta\}}$ . Consider

$$\mu' : C^0 \times Z \longrightarrow X.$$

Since  $C^0$  is geometrically irreducible,  $C^0 \times Z$  is irreducible; write  $\gamma$  for its generic point. The morphism  $\mu'$  sends the point  $\varepsilon(\eta)$ , where  $\varepsilon$  denotes the identity section of  $C^0$ , to  $\eta$ . Hence  $\gamma$  is sent to a generization of  $\eta$ , and therefore to  $\eta$ . Thus  $\mu'$  sends the underlying space of  $C^0 \times Z$  into  $Z$ , and, since  $C^0 \times Z$  is reduced,  $\mu'$  factors through  $Z$ .

Now let  $z \in Z$ , and put  $K = \kappa(z)$ . The morphism

$$\mu_z : G_K \longrightarrow X_K, \quad h \longmapsto h \cdot z,$$

is surjective. Let  $\alpha$  be a maximal point of  $X_K$ . The local ring  $\mathcal{O}_{X_K, \alpha}$  is a field, since  $X_K$  is reduced, so  $\mu_z$  is flat at every point of  $G_K$  above  $\alpha$ . It follows from Lemma 2.5.3 that  $\mu_z$  is flat.

On the other hand,  $\mu_z$  sends the generic point  $\omega$  of  $C_K^0$  to a point  $t \in Z_K$ . Let  $\beta$  be a maximal point of  $Z_K$  such that  $t \in \overline{\{\beta\}}$ . Since  $\mu_z$  is flat, there is a generization  $\xi$  of  $\omega$  such that  $\mu_z(\xi) = \beta$ . But  $\omega$  is a maximal point of  $G_K$ , so necessarily  $\xi = \omega$ . Thus  $\mu_z(\omega) = \beta$ , and  $\beta$  lies above  $\eta$ , because  $Z_K \rightarrow Z$  is flat.

Put  $L = \kappa(\omega)$ , and let  $\omega_L$  and  $z_L$  be the  $L$ -points obtained from  $\omega$  and  $z$ . Then

$$\omega_L \cdot z_L = \beta'$$

is a point of  $Z_L$  lying above  $\beta$ , and hence

$$z_L = \omega_L^{-1} \cdot \beta' \in C_L^0 \cdot U_L.$$

Therefore  $z \in C^0 \cdot U$ , which proves (i).

Since  $C^0$  is quasi-compact by 2.6.2, assertion (i') follows: if  $X$  is irreducible and  $U$  is a nonempty affine open subset, then  $X = C^0 U$ , so  $X$  is the image of the morphism  $C^0 \times U \rightarrow X$ , and is therefore quasi-compact. Finally, if  $U$  is retrocompact in  $X$ , then by 2.6.3,  $\overline{U}$  is the union of the  $\overline{\{\eta\}}$ , where  $\eta$  runs through the maximal points of  $X$  contained in  $U$ . Hence  $\overline{U} = C^0 U$ . This proves (ii).

One then obtains the following result ([Per75], II, Th. 2.4; see also [Per76], Prop. 4.1.1).

**Theorem 2.6.5.** *Let  $k$  be a field, and let  $G$  be a  $k$ -group scheme.*

- (i) *There is a unique group subscheme  $G^0$  of  $G$ , called the identity component of  $G$ , such that:*
  - (a) *the underlying space of  $G^0$  is the irreducible component of the identity element;*
  - (b)  *$G^0 \rightarrow G$  is a flat closed immersion; that is,*

$$\mathcal{O}_{G, g} = \mathcal{O}_{G^0, g} \quad \text{for every } g \in G^0.$$

- (ii) *Moreover,  $G^0$  is quasi-compact, geometrically irreducible, and is a characteristic subgroup of  $G$ .*
- (iii) *If  $G$  is connected, then  $G = G^0$ .*



*Proof.* (i) Recall first that  $G$  is separated (0.3), and hence, a fortiori, quasi-separated. Let  $U$  be an affine open subset of  $G$  containing the generic point  $\omega$  of  $C^0$ . By 2.6.3 and 2.6.4,

$$\overline{U} = C^0 U$$

is both open and closed, and, set-theoretically,

$$C^0 = \overline{\{\omega\}}$$

is the intersection of these open-and-closed subsets.

As  $U$  runs through the affine open subsets containing  $\omega$ , the  $G$ -schemes  $\overline{U}$  form a projective system whose transition morphisms are affine, since they are closed immersions. We can therefore form its projective limit  $G^0$  (cf. EGA IV<sub>3</sub>, 8.2.2). Thus, for every affine open  $V$  of  $G$ ,  $G^0 \cap V$  is the spectrum of the algebra

$$\varinjlim_U \mathcal{O}_G(V \cap \overline{U}) = \mathcal{O}_G(V) / \sum_U I_{\overline{U}}(V),$$

where  $I_{\overline{U}}(V)$  denotes the kernel of

$$\mathcal{O}_G(V) \longrightarrow \mathcal{O}_G(V \cap \overline{U}).$$

It follows that the underlying space of  $G^0$  is  $C^0$ , and that  $G^0 \rightarrow G$  is a closed immersion.

Moreover, for every  $g \in G^0$ ,  $\mathcal{O}_{G^0, g}$  is the inductive limit, as  $V$  runs through the affine open subsets of  $G$  containing  $g$ , of the  $k$ -algebras

$$\mathcal{O}_{G^0}(V \cap G^0) = \varinjlim_U \mathcal{O}_G(V \cap \overline{U}).$$

This double inductive limit is identified with

$$\varinjlim_U \varinjlim_{g \in V \subset \overline{U}} \mathcal{O}_G(V) = \mathcal{O}_{G, g}.$$

Thus

$$\mathcal{O}_{G^0, g} = \mathcal{O}_{G, g}$$

(see also EGA IV<sub>2</sub>, 5.13.3(ii)), and consequently  $i : G^0 \rightarrow G$  is a flat closed immersion. Conversely, this condition implies that  $i^*(\mathcal{O}_G) = \mathcal{O}_{G^0}$ , so  $G^0$  is uniquely determined by conditions (a) and (b). This proves (i).

The first two assertions of (ii) follow from 2.6.2. Finally, let  $S$  be a  $k$ -scheme and let  $\phi$  be an automorphism of the  $S$ -group  $G_S$ . For every  $s \in S$ ,  $\phi_s$  sends  $G_{\kappa(s)}^0$  into itself, and therefore  $\phi(G_S^0) \subset G_S^0$ . Moreover, the closed immersion

$$i_S : G_S^0 \hookrightarrow G_S$$

obtained from  $i$  by base change is flat, so

$$\mathcal{O}_{G_S, \phi(z)} = \mathcal{O}_{G_S^0, \phi(z)} \quad \text{for every } z \in G_S^0.$$

Hence  $\phi \circ i_S$  factors through  $G_S^0$ . This proves that  $G^0$  is a characteristic subgroup of  $G$ , and proves (ii). Finally, (iii) is a special case of part (i) of the following proposition.<sup>32</sup>

**Proposition 2.6.6.** *Let  $k$  be a field, and let  $G$  be a  $k$ -group acting on a  $k$ -scheme  $X$  in such a way that*

$$G \times X \longrightarrow X \times X, \quad (g, x) \longmapsto (gx, x)$$

*is surjective. Suppose that  $X$  is quasi-separated. Then:*

<sup>32</sup>Editor's note: this proposition, as well as the preceding results, was communicated to us by O. Gabber. It will be used to correct the proof of Theorem 5.3 in Exposé VI<sub>B</sub>.

- (i) Every connected component  $C$  of  $X$  is irreducible.
- (ii) Let  $\eta$  be the generic point of  $C$ , and let  $L$  be the algebraic closure of  $k$  in  $\kappa(\eta)$ . Then  $C$  is an  $L$ -scheme, is geometrically irreducible over  $L$ , and the morphism

$$G_L^0 \times_L C \longrightarrow C \times_L C$$

is surjective.

- (iii) In particular, if  $C$  contains a rational point  $x$ , then  $C$  is geometrically irreducible over  $k$ , and

$$\phi : G^0 \longrightarrow C, \quad g \longmapsto gx,$$

is surjective.

*Proof.* (i) Let  $C$  be a connected component of  $X$ , let  $Y$  be an irreducible component of  $X$  contained in  $C$ , let  $\eta$  be the generic point of  $Y$ , and let  $U$  be an affine open subset of  $X$  containing  $\eta$ . Since  $X$  is quasi-separated,  $U$  is retrocompact in  $X$ . Hence, by 2.6.4,

$$\overline{U} = G^0 U$$

is an open-and-closed subset of  $X$  which meets  $C$ , and therefore contains  $C$ . By 2.6.3, the intersection of the  $\overline{U}$ , as  $U$  runs through a fundamental system of affine open neighborhoods of  $\eta$ , is  $\{\eta\}$ . It follows that

$$C = \overline{\{\eta\}},$$

which proves (i).

The first assertion of (ii), and also that of (iii), follows from 2.6.1. We first prove the second assertion of (iii). Let  $\omega$ , respectively  $\eta$ , denote the generic point of  $G^0$ , respectively  $C$ . Let  $z \in C$ , and set  $K = \kappa(z)$ . Since  $G^0$ , respectively  $C$ , is geometrically irreducible over  $k$ , the scheme  $G_K^0$ , respectively  $C_K$ , is irreducible; denote its generic point by  $\xi$ , respectively  $\beta$ . We saw in the proof of 2.6.4 that

$$\mu_z : G_K \longrightarrow C_K, \quad h \longmapsto h \cdot z,$$

sends  $\xi$  to  $\beta$ . Likewise,  $\phi(\omega) = \eta$ .

Put  $L = \kappa(\xi)$ , and let  $\xi_L$  and  $z_L$  be the  $L$ -points obtained from  $\xi$  and  $z$ . Then

$$\xi_L \cdot z_L = \beta'$$

is a point of  $C_L$  lying above  $\beta \in C_K$ , and hence also above  $\eta \in C$ . Consider the cartesian square

$$\begin{array}{ccc} G_L^0 & \xrightarrow{\varphi_L} & C_L \\ \pi_{G^0} \downarrow & & \downarrow \pi_C \\ G^0 & \xrightarrow{\varphi} & C \end{array}$$

Since

$$\phi_L(\pi_{G^0}^{-1}(\omega)) = \pi_C^{-1}(\eta)$$

(cf. EGA I, 3.4.8), there is a  $g \in G_L^0$  such that  $\phi_L(g) = \beta'$ . Therefore

$$z_L = \xi_L^{-1} \cdot \phi_L(g) = \phi_L(\xi_L^{-1}g),$$

whence

$$\phi(\pi_{G^0}(\xi_L^{-1}g)) = \pi_C(z_L) = z.$$

This proves that  $\phi$  is surjective.

We now prove the second assertion of (ii). It is enough to show that, for every  $z \in C$ , the morphism

$$\mu_z : G_L^0 \otimes_L \kappa(z) \longrightarrow C \otimes_L \kappa(z)$$

is surjective. But this follows from (iii), since  $z$  is a rational point of  $C \otimes_L \kappa(z)$ .

**Remark 2.6.7.**— *Under the hypotheses of 2.6.6, if  $C(k) = \emptyset$ , the morphism*

$$G^0 \times_k C \longrightarrow C \times_k C$$

*need not be surjective. For example, let  $k = \mathbb{R}$  and  $G = \{\pm 1\}_{\mathbb{R}}$ . The  $G$ -torsor*

$$X = \operatorname{Spec} \mathbb{R}[X]/(X^2 + 1)$$

*is connected, but*

$$G^0 \times_{\mathbb{R}} X \longrightarrow X \times_{\mathbb{R}} X$$

*is not surjective. (Here  $L = \mathbb{C}$ , and*

$$G^0 \times_{\mathbb{R}} X \longrightarrow X \times_{\mathbb{C}} X$$

*is an isomorphism.)*

### 3. Construction of quotients $F \backslash G$ (for $G, F$ of finite type)

**3.1.**— Let  $A$  be an Artinian local ring, and let  $u : F \rightarrow G$  be a homomorphism of  $A$ -groups. If

$$\mu : F \times_A F \longrightarrow F \quad \text{and} \quad \nu : G \times_A G \longrightarrow G$$

denote the multiplication morphisms, and if  $\lambda$  denotes the composite morphism

$$F \times_A G \xrightarrow{u \times G} G \times_A G \xrightarrow{\nu} G$$

recall that the *left quotient*  $F \backslash G$  of  $G$  by  $F$  is the cokernel of the (Sch/ $A$ )-groupoid  $G_*$  described below:

$$\begin{array}{ccccc} F \times_A F \times_A G & \xrightarrow{F \times \lambda} & & \xrightarrow{\lambda} & G \\ & \xrightarrow{\mu \times G} & F \times_A G & \xrightarrow{\text{pr}_2} & \\ & \xrightarrow{\text{pr}_{2,3}} & & & \end{array}$$

Here  $\text{pr}_2$  and  $\text{pr}_{2,3}$  are the projections of  $F \times_A G$  and  $F \times_A (F \times_A G)$ , respectively, onto the second factors. We shall say that  $G_*$  is the *groupoid with base  $G$  defined by  $u$*  (cf. Exposé V, § 2.a; as in Exposé V, we do not follow in this exposé the convention of IV, 4.6.15).

Since the unique  $A$ -morphism  $F \rightarrow \operatorname{Spec} A$  is universally open (EGA IV<sub>2</sub>, 2.4.9),  $\text{pr}_2$  is an open morphism. The same therefore holds for  $\lambda$ , which is the composite of  $\text{pr}_2$  with the automorphism  $\sigma$  of  $F \times_A G$  defined by

$$\sigma(S)(x, y) = (x, u(S)(x) \cdot y),$$

where  $S$  is a variable  $A$ -scheme and  $x \in F(S)$ ,  $y \in G(S)$ . One sees in the same way that  $\text{pr}_2$  and  $\lambda$  are flat when  $F$  is flat over  $A$ .

Let us also note, to conclude these preliminaries, that every  $A$ -morphism  $s : \operatorname{Spec} A \rightarrow G$  defines an automorphism of the groupoid  $G_*$  which induces on  $G$ ,  $F \times_A G$ , and  $F \times_A F \times_A G$ , respectively, the automorphisms

$$r_s, \quad \operatorname{id}_F \times_A r_s, \quad \operatorname{id}_F \times_A \operatorname{id}_F \times_A r_s.$$

We shall again write  $r_s$  for this automorphism of  $G_*$ , and shall say that  $r_s$  is the *right translation defined by  $s$*  (cf. 0.4).

**Theorem 3.2.** *Let  $F$  and  $G$  be flat groups locally of finite type over an Artinian local ring  $A$ . Let  $u : F \rightarrow G$  be a quasi-compact homomorphism of  $A$ -groups whose kernel is finite over  $A$ . Then:*<sup>33</sup>

- (i) *The left quotient  $F \backslash G$  of  $G$  by  $F$  exists in  $(\operatorname{Sch}/A)$ , and the sequence*

$$F \times_A G \begin{array}{c} \xrightarrow{\lambda} \\ \xrightarrow{\operatorname{pr}_2} \end{array} G \xrightarrow{p} F \backslash G$$

*is exact in the category of all ringed spaces.*

- (ii) *The canonical morphism  $p : G \rightarrow F \backslash G$  is surjective and open, and  $F \backslash G$  is a direct sum of schemes of finite type over  $A$ .*
- (ii') *More precisely,  $X = F \backslash G$  is equipped with a right action of  $G$  such that  $p(e) \cdot g = p(g)$  for every  $g \in G$ . Consequently, the connected components of  $X$  are of finite type over  $A$ , irreducible, and all of dimension  $\dim G - \dim F$ .*

- (iii) *The canonical morphism*

$$F \times_A G \longrightarrow G \times_{(F \backslash G)} G$$

*is surjective.*

- (iv) *If  $u$  is a monomorphism,*<sup>34</sup> *then:*

- (a) *The morphism*

$$F \times_A G \xrightarrow{(\lambda, \operatorname{pr}_2)} G \times_{(F \backslash G)} G$$

*is an isomorphism, and  $G \rightarrow F \backslash G$  is faithfully flat and locally of finite presentation.*

- (a')  *$F \backslash G$  represents the (fppf) quotient sheaf  $\widetilde{F \backslash G}$ , and  $G \rightarrow F \backslash G$  is an  $F$ -torsor locally trivial for the (fppf) topology.*
- (b)  *$F \backslash G$  is flat over  $A$ , and is smooth over  $A$  if  $G$  is so.*
- (c)  *$u : F \rightarrow G$  is a closed immersion, and  $F \backslash G$  is separated.*
- (d) *If, moreover,  $F$  is a normal subgroup of  $G$ , there is a unique  $A$ -group structure on  $F \backslash G$  for which  $p : G \rightarrow F \backslash G$  is a morphism of  $A$ -groups.*

In the proof of this theorem,  $A'$  will denote an  $A$ -algebra which is local, finite, and free over  $A$ . If  $R$  is a relation involving  $A'$ , we shall say that “ $R(A')$  is true when  $A'$  is sufficiently large” if there is an  $A$ -algebra  $A_1$ , local, finite, and free over  $A$ , such that the relation  $R(A')$  holds for every  $A$ -algebra  $A'$  which is local, finite, and free over  $A_1$ .

<sup>33</sup>Editor's note: we have added (ii') and expanded point (iv), taking account of the additions made in 2.5.2, 2.5.4, and Exposé V, 6.1.

<sup>34</sup>Editor's note: in this case, the hypothesis that  $G$  be flat can be omitted; cf. subsection 3.3.

We shall first prove the theorem when  $F$  and  $G$  are of finite type over  $A$ .

**3.2.1.**— Suppose for a moment that every point of  $G$  has an open saturated neighborhood  $W$  such that the groupoid induced by  $G_*$  on  $W$  has a quasi-section (cf. Exposé V, § 6). Then, by Exposé V, 6.1, assertions (i), (ii), (iii), and (iv)(a) hold, and  $F \backslash G$  is of finite type over  $A$ .<sup>\*</sup> Moreover, under the hypothesis of (iv), since

$$G \longrightarrow F \backslash G$$

is faithfully flat and locally of finite presentation, assertion (b) follows from EGA IV, 2.2.14 and 17.7.7. Assertion (iv)(a') follows from (iv)(a), by Exposé IV, 3.4.3.1, 5.2.2, and 5.1.6. Finally, assertion (iv)(c) will be proved in 3.2.5, while (ii') and (iv)(d) will be proved in Section 5.

We now prove the following assertion:

every finite subset of  $F \backslash G$  is contained in an affine open subset. (†)

35

If  $U$  is a quasi-section of the groupoid induced by  $G_*$  on a saturated open subset  $W$  of  $G$ , then  $U \otimes_A A'$  is a quasi-section of the  $(\text{Sch}/A')$ -groupoid induced by  $G_* \otimes_A A'$  on  $W \otimes_A A'$ . Moreover, if  $U_*$  is the  $(\text{Sch}/A)$ -groupoid induced by  $G_*$  on  $U$ , then  $U_* \otimes_A A'$  is identified with the  $(\text{Sch}/A')$ -groupoid induced by  $G_* \otimes_A A'$  on  $U \otimes_A A'$ .<sup>36</sup>

It follows from the proofs in Exposé V that the construction of the quotient

$$X = F \backslash G$$

commutes with a base extension  $A \rightarrow A'$  of the kind considered here.<sup>37</sup>

Let  $x_1, \dots, x_n$  be points of  $X = F \backslash G$ , which we may suppose to be closed,<sup>38</sup> and let  $g_1, \dots, g_n$  be closed points of  $G$  mapping to  $x_1, \dots, x_n$ . Let  $V$  be an everywhere dense affine open subset of  $X$ ,<sup>39</sup> and let  $U$  be its inverse image in  $G$ . By 1.2, there is an  $A$ -algebra  $A'$ , local, finite, and free over  $A$ , such that the points  $g'_1, \dots, g'_p$  of

$$G' = G \otimes_A A'$$

---

<sup>\*</sup>Translator's note: the French source prints  $k$  here and again in editor's note 39. In this part of the proof the base is the Artinian local ring  $A$ , and "finite type over  $A$ " is the required statement.

<sup>35</sup>Editor's note: let us point out that if  $A = k$  is a field, every quasi-compact open subset of  $F \backslash G$  is quasi-projective (a result due to Chow for smooth algebraic groups); cf. [Ray70], VI, 2.6. By contrast, over the Artinian local ring  $A = \mathbb{C}[\varepsilon]/(\varepsilon^2)$ , there exist abelian  $A$ -schemes  $G$  which are not projective (loc. cit., XII, 4.2).

<sup>36</sup>Editor's note: the preceding assertion is valid for every base change  $A \rightarrow A'$ .

<sup>37</sup>Editor's note: this is explained in detail in 4.6 below. One must show that formation of direct images by the morphisms  $p$ ,  $\lambda$ , and  $\text{pr}_2$  commutes with flat base changes  $A \rightarrow A'$ . Since  $F$  and  $G$  are of finite type over the Artinian ring  $A$ , the morphisms  $f$  in question are all quasi-compact and quasi-separated, and

$$f_*(\mathcal{O}_X) \otimes_A A' = f'_*(\mathcal{O}_{X'})$$

(with the evident notation) follows from EGA IV<sub>1</sub>, 1.7.21.

<sup>38</sup>Editor's note: indeed, let  $y_1, \dots, y_n$  be arbitrary points of  $X$ . Since  $X$  is of finite type over  $A$ , the closure of each  $y_i$  contains a closed point  $x_i$ , and every open subset containing  $x_i$  contains  $y_i$ .

<sup>39</sup>Editor's note: such an open subset exists because  $X$  is of finite type over  $A$ :  $X$  has only finitely many irreducible components  $C_1, \dots, C_p$ , and it is enough, for each  $i$ , to choose a nonempty affine open subset contained in

$$C_i - \bigcup_{j \neq i} C_j.$$

(Here, moreover, one knows from (ii') that the  $C_i$  are disjoint.)

lying above  $g_1, \dots, g_n$  are strictly rational over  $A'$ .<sup>40</sup> Since the morphisms  $G' \rightarrow G$  and  $G \rightarrow X$  are open,

$$U' = U \otimes_A A'$$

is dense in  $G'$ . Hence the open subset

$$\bigcap_{i=1}^p (U')^{-1} \cdot g'_i$$

is nonempty and therefore contains a closed point  $x$ . By 1.2 (and 0.4.1), after enlarging  $A'$  if necessary, we may suppose that  $x$  is strictly rational over  $A'$ . Since  $x \in (U')^{-1} \cdot g'_i$ , we have

$$g'_i \in U' \cdot x \quad (1 \leq i \leq p).$$

Write  $V'$  for the inverse image of  $V$  in

$$X' = X \otimes_A A'.$$

It is an affine open subset of  $X'$ , and is also the image of  $U'$  under the projection  $G' \rightarrow X'$ . Since right translation  $r_x$  is an automorphism of the groupoid  $G_* \otimes_A A'$ , it induces an automorphism, again denoted  $r_x$ , of the quotient  $X'$ . Consequently,

$$V' \cdot x = r_x(V'),$$

the image of  $U' \cdot x$  in  $X'$ , is an affine open subset of  $X'$  containing the images  $x'_1, \dots, x'_p$  of  $g'_1, \dots, g'_p$ .

Consider the equivalence relation on  $X' = X \otimes_A A'$  defined by the projection  $X \otimes_A A' \rightarrow X$ :

$$X \otimes_A A' \otimes_A A' \begin{array}{c} \xrightarrow{d_1} \\ \xrightarrow{d_0} \end{array} X \otimes_A A' \longrightarrow X$$

Here  $d_0$  and  $d_1$  are induced by the two canonical injections of  $A'$  into  $A' \otimes_A A'$ . Since  $A'$  is a finite free  $A$ -algebra, say of rank  $n$ ,  $d_0$  and  $d_1$  are finite and locally free of rank  $n$ . We may therefore apply the argument of Exposé V, 5.b, taken from the proof of SGA 1, VIII.7.6. It follows that  $x'_1, \dots, x'_p$  are contained in a saturated affine open subset  $W'$  contained in the affine open subset  $V' \cdot x$ . The image of  $W'$  in  $X$  then contains  $x_1, \dots, x_n$ , and is an affine open subset of  $X$ , by Exposé V, 4.1(ii).

**3.2.2.**— For every  $A$ -algebra  $A'$ , local, finite, and free over  $A$ , let  $U(A')$  denote the set of points of  $G \otimes_A A'$  having an open saturated neighborhood  $W$  such that the groupoid induced by  $G_* \otimes_A A'$  on  $W$  has a quasi-section. It is clear that  $U(A')$  is saturated under the operations of  $G(\text{Spec } A')$  on  $G \otimes_A A'$ . We shall show that, when  $A'$  is sufficiently large,

$$U(A') = G \otimes_A A'.$$

By Theorem V.8.1,  $U(A)$  is nonempty, and therefore contains a closed point  $y$ . We argue by induction on

$$\dim(G - U(A)).$$

Let  $g_1, \dots, g_n$  be closed points belonging to the various irreducible components of  $G - U(A)$ . By 1.2, there is an  $A$ -algebra  $A'$ , local, finite, and free over  $A$ , such that the points

$$g'_1, \dots, g'_p \quad (\text{respectively, } x = x_1, \dots, x_r)$$

<sup>40</sup>Editor's note: the original has been expanded in what follows.

of  $G' = G \otimes_A A'$ , lying above  $g_1, \dots, g_n$  (respectively, above  $y$ ), are strictly rational over  $A'$ . Then  $U(A')$  contains

$$(U(A) \otimes_A A') \cdot x^{-1} g'_i \quad \text{for every } i.$$

Consequently,  $U(A')$  contains  $g'_1, \dots, g'_p$ , and

$$\dim(G' - U(A')) < \dim(G - U(A)).$$

The induction hypothesis therefore gives an algebra  $A''$ , local, finite, and free over  $A'$ , such that

$$U(A'') = G' \otimes_{A'} A'' = G \otimes_A A''.$$

**3.2.3.**— We are now in a position to prove the existence of  $F \backslash G$  when  $F$  and  $G$  are of finite type over  $A$ . Let  $A'$  be sufficiently large over  $A$  that  $U(A')$  coincides with  $G \otimes_A A'$  (cf. 3.2.2). Set

$$A'' = A' \otimes_A A',$$

and, for every  $A$ -scheme  $X$ , write  $X'$  and  $X''$  for the fiber products  $X \otimes_A A'$  and  $X \otimes_A A''$ . By 3.2.1 and 3.2.2, the quotients  $F' \backslash G'$  and  $F'' \backslash G''$  exist, and we have the following commutative diagram, in which the first two rows and columns are exact:

$$(*) \quad \begin{array}{ccccc} F'' \times_{A''} G'' & \xrightarrow{\text{pr}_2''} & G'' & \xrightarrow{p''} & F'' \backslash G'' \\ w_1 \downarrow & \lambda'' \downarrow & v_1 \downarrow & v_2 \downarrow & u_1 \downarrow \quad u_2 \downarrow \\ F' \times_{A'} G' & \xrightarrow{\text{pr}_2'} & G' & \xrightarrow{p'} & F' \backslash G' \\ h \downarrow & \lambda' \downarrow & g \downarrow & & \\ F \times_A G & \xrightarrow{\text{pr}_2} & G & \xrightarrow{\lambda} & \end{array}$$

In this diagram,  $\text{pr}_2'$  and  $\lambda'$  (respectively,  $\text{pr}_2''$  and  $\lambda''$ ) are obtained from  $\text{pr}_2$  and  $\lambda$  by the evident base changes; the morphisms  $g$  and  $h$  are induced by the canonical injection  $A \rightarrow A'$ . We write  $p'$  and  $p''$  for the canonical morphisms, while  $v_1, v_2$  and  $w_1, w_2$  are induced by the two canonical injections of  $A'$  into  $A''$ .

Finally, formation of the quotient  $F' \backslash G'$  commutes with the two base changes

$$f_1, f_2 : \text{Spec } A'' \rightrightarrows \text{Spec } A'.$$

Thus, if

$$\pi' : F' \backslash G' \longrightarrow \text{Spec } A'$$

denotes the structure morphism, there are canonical isomorphisms, for  $i = 1, 2$ ,

$$\tau_i : F'' \backslash G'' \xrightarrow{\sim} (F' \backslash G') \times_{\pi', f_i} \text{Spec } A'',$$

and  $u_i$  is the composite of  $\tau_i$  with the projection

$$(F' \backslash G') \times_{\pi', f_i} \text{Spec } A'' \longrightarrow F' \backslash G'.$$

Now, for a diagram of type  $(*)$  in which the first two rows and columns are exact, one checks readily that  $\text{Coker}(\text{pr}_2, \lambda)$  exists if and only if  $\text{Coker}(u_1, u_2)$  exists, and that these two cokernels are then identified. The existence of  $F \backslash G$  therefore follows from that of  $\text{Coker}(u_1, u_2)$ .

The compatibility of the formation of  $F \backslash G$  with the base extensions considered here (cf. editor's note 37 in 3.2.1 and 4.6 below) shows that the composite

$$(F' \setminus G') \times_{\pi', f_1} \operatorname{Spec} A'' \xrightarrow{\tau_1^{-1}} F'' \setminus G'' \xrightarrow{\tau_2} (F' \setminus G') \times_{\pi', f_2} \operatorname{Spec} A''$$

is a descent datum on  $F' \setminus G'$  relative to

$$f : \operatorname{Spec} A' \longrightarrow \operatorname{Spec} A.$$

By 3.2.1 (†) and SGA 1, VIII.7.6, this descent datum is effective; that is,  $\operatorname{Coker}(u_1, u_2)$  exists. (One could instead apply Exposé V, Theorem 4.1 directly.)

**3.2.4.**— To complete the proof of assertions (i), (ii), (iii), and (iv)(a) of 3.2 in the case where  $F$  and  $G$  are of finite type over  $A$ , it remains to study the quotient  $F \setminus G$ . By Exposé V, 6.1, assertions (ii), (iii), and (iv)(a) “become true” after the base change

$$f : \operatorname{Spec} A' \longrightarrow \operatorname{Spec} A.$$

By EGA IV<sub>2</sub>, 2.6.1, 2.6.2, and 2.7.1, these assertions were therefore true before the base change. Finally, to prove the second assertion of (i), namely that  $F \setminus G$  is the cokernel of  $(\operatorname{pr}_2, \lambda)$  in the category of all ringed spaces, it suffices to refer to Exposé V, §6(c).

**3.2.5.**—<sup>41</sup> We now prove assertion (iv)(c) of 3.2, following the proof of Exposé VI<sub>B</sub>, 9.2.1. Write

$$X = F \setminus G$$

and let  $d$  be the morphism

$$F \times_A G \longrightarrow G \times_A G$$

whose components are  $\lambda$  and  $\operatorname{pr}_2$ .

Since  $u$  is a closed immersion by 2.5.2, and since

$$d = \sigma \circ (u \times \operatorname{id}_G), \quad \sigma(x, y) = (xy, y),$$

where  $\sigma$  is an automorphism of  $G \times_A G$ , the morphism  $d$  is a closed immersion. On the other hand, by (iv)(a), the following square is Cartesian:

$$\begin{array}{ccc} F \times_A G & \xrightarrow{d} & G \times_A G \\ \downarrow & & \downarrow p \times p \\ X & \xrightarrow{\Delta_X} & X \times_A X \end{array}$$

Moreover,  $p$ , and hence also  $p \times p$ , is faithfully flat and locally of finite presentation. Therefore, by fppf descent, since  $d$  is a closed immersion, so is  $\Delta_X$ ; that is,  $X$  is separated.

**3.3.**—<sup>42</sup> In Theorem 3.2, the hypothesis that  $G$  be flat can be omitted when  $u : F \rightarrow G$  is a monomorphism. This generalization is mentioned in Exposé VI<sub>B</sub>, Remark 9.3(b), and also in [Ray67a], Example a)(i), p. 82. The proof, given in [An73], Theorem 4, follows from Theorem 3.2 and from the following theorem of Grothendieck (mentioned in [Ray67a], Theorem 1(ii), and proved in [DG70], § III.2, 7.1). If  $X$  is a scheme and  $R$  an equivalence relation on  $X$ , we write  $\widetilde{X/R}$  for the (fppf) quotient sheaf of  $X$  by  $R$  (cf. Exposé IV, 4.4.9).

**Theorem 3.3.1 (Grothendieck).** *Let  $A$  be a ring, let  $X$  be an  $A$ -scheme, and let*

$$d_0, d_1 : R \rightrightarrows X$$

<sup>41</sup>Editor’s note: we have added this paragraph.

<sup>42</sup>Editor’s note: we have added this subsection.



be an  $A$ -equivalence relation on  $X$ , with each  $d_i$  faithfully flat and of finite presentation. Let  $X_0$  be a saturated closed subscheme of  $X$ , defined by a nilpotent ideal, and let  $R_0$  be the equivalence relation induced by  $R$  on  $X_0$ . If the (fppf) quotient sheaf  $\widetilde{X_0/R_0}$  is representable by an  $A$ -scheme, then so is  $\widetilde{X/R}$ .

For the proof, see [DG70], § III.2, 7.1. Let us now return to the case where  $A$  is an Artinian local ring. Let  $u : F \rightarrow G$  be a quasi-compact morphism between  $A$ -groups locally of finite type, and suppose in addition that  $u$  is a monomorphism. Then  $u$  is a closed immersion by 2.5.2.

We may now state the following variant of Theorem 3.2.

**Theorem 3.3.2.** *Let  $A$  be an Artinian local ring, let  $G$  be an  $A$ -group locally of finite type, and let  $F$  be a closed subgroup of  $G$ , flat over  $A$ .<sup>43</sup> Then:*

- (i) *The (fppf) quotient sheaf  $\widetilde{F \backslash G}$  is representable by a separated  $A$ -scheme  $F \backslash G$ , locally of finite type; moreover, the sequence*

$$F \times_A G \begin{array}{c} \longrightarrow \\ \longrightarrow \end{array} G \xrightarrow{p} F \backslash G$$

*is exact in the category of all ringed spaces.*

- (ii) *The morphism*

$$F \times_A G \xrightarrow{(\lambda, \text{pr}_2)} G \times_{(F \backslash G)} G$$

*is an isomorphism, and  $p : G \rightarrow F \backslash G$  is faithfully flat and locally of finite presentation; hence  $p$  is an  $F$ -torsor locally trivial for the (fppf) topology.*

- (iii) *If  $G$  is flat (respectively, of finite type; respectively, smooth) over  $A$ , then so is  $F \backslash G$ .*
- (iv)  *$X = F \backslash G$  is equipped with a right action of  $G$  such that  $p(e) \cdot g = p(g)$  for every  $g \in G$ . Consequently, the connected components of  $X$  are of finite type over  $A$ , irreducible, and all of dimension  $\dim G - \dim F$ .*
- (v) *If, in addition,  $F$  is a normal subgroup of  $G$ , there is a unique  $A$ -group structure on  $F \backslash G$  for which  $p : G \rightarrow F \backslash G$  is a morphism of  $A$ -groups.*

Assertions (i) and (ii) follow from 3.2 and 3.3.1. Since  $G \rightarrow F \backslash G$  is faithfully flat and locally of finite presentation, assertion (iii) follows from EGA IV, 2.2.14, 2.7.1, and 17.7.7. Assertions (iv) and (v) will be proved in Section 5. Let us record at once the following corollary.

**Corollary 3.3.3.** *Let  $A$  be an Artinian local ring, let  $G$  be an  $A$ -group locally of finite type, and let  $H$  be a closed subgroup of  $G$ , flat over  $A$ . Write  $p$  for the morphism  $G \rightarrow G/H$ , and write  $\lambda$  (respectively,  $\text{pr}_1$ ) for the morphism  $G \times H \rightarrow G$  defined by  $\lambda(g, h) = gh$  (respectively, the projection  $G \times H \rightarrow G$ ). Then, for every open subset  $U$  of  $G/H$ ,*

$$\mathcal{O}(U) = \{\varphi \in \mathcal{O}(p^{-1}(U)) \mid \varphi \circ \lambda = \varphi \circ \text{pr}_1\}.$$

*In other words,  $\mathcal{O}(U)$  is the set of  $\varphi \in \mathcal{O}(p^{-1}(U))$  such that  $\varphi(gh) = \varphi(g)$  for every  $A$ -scheme  $S$  and every  $g \in G(S)$ ,  $h \in H(S)$ .*

Indeed,  $p : G \rightarrow G/H$  is faithfully flat and locally of finite presentation, hence a covering for the (fppf) topology; the result follows from Exposé IV, 3.3.3.2.

<sup>43</sup>Editor's note: for an example in which  $F$  is not flat and  $\widetilde{F \backslash G}$  is not representable, see [DG70], § III.3, no. 3.3.

#### 4. Construction of quotients $F \backslash G$ (general case)

We now assume the hypotheses of Theorem 3.2, without requiring  $F$  and  $G$  to be of finite type over  $A$ .

**4.1.**— Let  $G^\alpha$  be a connected component of  $G$ . We first show that the saturation

$$\mathcal{S}(G^\alpha)$$

<sup>44</sup> of  $G^\alpha$  under the equivalence relation defined by the groupoid  $G_*$  is an open-and-closed subset of  $G$ ; in other words, it is a union of connected components of  $G$ .

This saturation is the image of  $F \times_A G^\alpha$  under  $\lambda$ , and is therefore open in  $G$  (cf. § 3.1). Let  $k$  be the residue field of  $A$ , and let  $\bar{k}$  be an algebraic closure of  $k$ . It remains to show that the image of

$$(F \times_A G^\alpha) \otimes_A k$$

under  $\lambda \otimes_A k$  is closed in  $G \otimes_A k$ , or, equivalently by SGA 1, VIII.4.4, that the image of

$$(F \times_A G^\alpha) \otimes_A \bar{k}$$

under  $\lambda \otimes_A \bar{k}$  is closed in  $G \otimes_A \bar{k}$ . Since  $G^\alpha \otimes_A \bar{k}$  is a union of finitely many connected components of  $G \otimes_A \bar{k}$ , we are reduced to the case where  $A$  is an algebraically closed field, which we now assume. In this case,  $\mathcal{S}(G^\alpha)$  is the union of the images of  $G^\alpha$  under the left translations  $\ell_{u(x)}$ , where  $x$  runs through the closed points of  $F$ . The assertion follows because these images are connected components of  $G$ .

**4.2.**— Take  $G^\alpha$  in particular to be the connected component  $G^0$  of the identity in  $G$ . Then  $\mathcal{S}(G^0)$  plainly contains the image of  $F$  under  $u$ , which is precisely the equivalence class of the identity. On the other hand, if  $F^\beta$  is a connected component of  $F$ , then  $F^\beta \times_A G^0$  is connected by 2.1.2, so its image under  $\lambda$  is contained in the connected component of  $u(F^\beta)$  in  $G$ . In other words,  $\mathcal{S}(G^0)$  is the union of the connected components that meet the image of  $F$ .

Let us also observe that the open subscheme of  $G$  whose underlying space is  $\mathcal{S}(G^0)$  is a subgroup of  $G$ ; we continue to denote it by  $\mathcal{S}(G^0)$ . Indeed, inversion on  $G$  preserves the image of  $F$  and permutes the connected components of  $G$  that meet this image. It therefore suffices to prove that

$$\nu : G \times_A G \longrightarrow G$$

maps  $\mathcal{S}(G^0) \times_A \mathcal{S}(G^0)$  into  $\mathcal{S}(G^0)$ . For this we may suppose that  $A$  is an algebraically closed field: with the notation of 4.1,  $\mathcal{S}(G^0) \otimes_A \bar{k}$  is the saturation of  $(G \otimes_A \bar{k})^0$  for the equivalence relation defined by the homomorphism  $u \otimes_A \bar{k}$ .

If  $G^\gamma$  and  $G^\delta$  are connected components of  $\mathcal{S}(G^0)$ , then  $G^\gamma \times_A G^\delta$  is connected and its image under  $\nu$  meets the image of  $F$ . Consequently,

$$\nu(G^\gamma \times_A G^\delta)$$

is contained in a connected component of  $G$  that meets  $u(F)$ .<sup>\*</sup>

**4.3.**— It follows from the preceding discussion that the groupoid  $G_*$ , with base  $G$ , defined by  $u$ , is the direct sum of the groupoids  $\mathcal{S}(G_*^\alpha)$  induced by  $G_*$  on the various open-and-closed parts of  $G$  of the form  $\mathcal{S}(G^\alpha)$ . The cokernel of  $G_*$  is therefore the direct sum of the cokernels of the groupoids  $\mathcal{S}(G_*^\alpha)$ , which we are thus led to study separately.

<sup>44</sup>Editor's note: we have written  $\mathcal{S}(G^\alpha)$ , rather than  $\overline{G^\alpha}$ , for the saturation of  $G^\alpha$ .

<sup>\*</sup>Translator's note: the French source prints  $u(G^\gamma \times_A G^\delta)$  in this final sentence. The map defined and used throughout the paragraph is the multiplication  $\nu : G \times_A G \rightarrow G$ ; the target therefore uses  $\nu$ .

First consider the groupoid  $\mathcal{S}(G_*^0)$  induced by  $G_*$  on  $\mathcal{S}(G^0)$ . It is clear that  $\mathcal{S}(G_*^0)$  is the groupoid with base  $\mathcal{S}(G^0)$  defined by the homomorphism

$$F \longrightarrow \mathcal{S}(G^0)$$

induced by  $u$  (§ 3.1). The cokernel whose existence we wish to prove is thus identified with  $F \backslash \mathcal{S}(G^0)$ . Consider, on the other hand, the groupoid

$$G_2^0 \begin{array}{c} \xrightarrow{\ell_2'} \\ \xrightarrow{\ell_1'} \\ \xrightarrow{\ell_0'} \end{array} G_1^0 \xrightarrow[\ell_0]{\ell_1} G_0^0 = G^0$$

induced by  $\mathcal{S}(G_*^0)$  on  $G^0$ . Referring to the explicit construction in Exposé V, § 3.b, the object denoted there by  $Y_0 \times_{X_0} X_1$  is precisely  $F \times_A G^0$ . Hence  $G_1^0$  is the inverse image of  $G^0$  under the morphism

$$F \times_A G^0 \longrightarrow \mathcal{S}(G^0)$$

induced by  $\lambda$ .

I claim that this inverse image is  $F_0 \times_A G^0$ , where  $F_0$  denotes the inverse image of  $G^0$  under  $u$ . Indeed, if  $F^\beta$  is a connected component of  $F_0$ , then  $F^\beta \times_A G^0$  is connected by 2.1.2 and  $\lambda(F^\beta \times_A G^0)$  is contained in  $G^0$ .\* Conversely, if  $F^\beta$  is a connected component of  $F$  not contained in  $F_0$ , then the image of  $F^\beta \times_A G^0$  is again connected and contains  $u(F^\beta)$ ; if  $u(F^\beta)$  is not contained in  $G^0$ , then  $\lambda(F^\beta \times_A G^0)$  does not meet  $G^0$ .

It follows that the groupoid  $G_*^0$  induced by  $G_*$  on  $G^0$  is the groupoid with base  $G^0$  defined by the homomorphism

$$F_0 \longrightarrow G^0$$

induced by  $u$ . Since  $G^0$ , and hence  $F_0$ , are of finite type over  $A$ , Section 3 shows that  $G_*^0$  has a cokernel, namely  $F_0 \backslash G^0$ .†

I now claim that  $F_0 \backslash G^0$  is identified with  $F \backslash \mathcal{S}(G^0)$ . The proof is analogous to that of the first part of assertion (i) of Exposé V, Lemma 6.1. Consider the diagram

$$\mathcal{S}(G^0) \xleftarrow[v]{} F \times_A G^0 \xrightarrow{\text{pr}_2} G^0$$

where  $v$  is the morphism induced by  $\lambda$ . Since  $\text{pr}_2$  has a section, it is a universal effective epimorphism. Consequently,  $F_0 \backslash G^0$  coincides with  $\text{Coker}(v_0, v_1)$ , where

$$V_2 \begin{array}{c} \xrightarrow{v_2'} \\ \xrightarrow{v_1'} \\ \xrightarrow{v_0'} \end{array} V_1 \xrightarrow[v_0]{v_1} V = F \times_A G^0$$

is the inverse image under  $\text{pr}_2$  of the groupoid  $G_*^0$  (cf. Exposé V, § 3.a), and hence also the inverse image of  $\mathcal{S}(G_*^0)$  under the composite morphism

$$F \times_A G^0 \xrightarrow{\text{inclusion}} F \times_A \mathcal{S}(G^0) \xrightarrow{\text{pr}_2} \mathcal{S}(G^0)$$

Similarly, since  $v$  is faithfully flat and quasi-compact,  $F \backslash \mathcal{S}(G^0)$  coincides with the cokernel of the inverse image of  $\mathcal{S}(G_*^0)$  under the base change  $v$ . This inverse image is isomorphic

---

\*Translator's note: the French source prints  $\lambda(F \times_A G^0)$  at this point. The statement concerns the connected component  $F^\beta$  just introduced, and the converse immediately below confirms the intended restriction to  $F^\beta \times_A G^0$ .

†Translator's note: the French source says "by paragraph 4." The finite-type quotient invoked here is the result of Section 3; Section 4 is the general-case construction currently in progress.

to  $V_*$  by Exposé V, § 3.c. It follows that the canonical inclusion of  $G_*^0$  in  $\mathcal{S}(G_*^0)$  induces an isomorphism

$$F_0 \backslash G^0 \xrightarrow{\sim} F \backslash \mathcal{S}(G^0).$$

Finally, formation of  $F \backslash \mathcal{S}(G^0)$  commutes with finite locally free base changes, because the same holds for  $F_0 \backslash G^0$  (cf. editor's note 37 and 4.6 below).

**4.4.**— It remains to construct the cokernel of the groupoid  $\mathcal{S}(G_*^\alpha)$  when  $G^\alpha$  is an arbitrary connected component of  $G$ . Let  $A'$  be a sufficiently large local  $A$ -algebra, finite and free over  $A$  (cf. 3.2). Then  $G^\alpha \otimes_A A'$  is the union of finitely many connected components

$$C^1, \dots, C^n$$

of  $G \otimes_A A'$ , each of which has a strictly rational point. For every  $i$ , there is therefore a right translation  $r_i$  of  $G \otimes_A A'$  which maps  $G^0 \otimes_A A'$  onto  $C^i$ . This translation induces an isomorphism

$$\mathcal{S}(G_*^0) \otimes_A A' \xrightarrow{\sim} \mathcal{S}(C_*^i).$$

Consequently, the groupoid induced by  $G_* \otimes_A A'$  on the saturation of  $C^i$  has a cokernel.

Since  $\mathcal{S}(G_*^\alpha) \otimes_A A'$  is the direct sum of some of the groupoids  $\mathcal{S}(C_*^i)$ , it too has a cokernel. This cokernel is a direct sum of some number of copies of

$$(F_0 \otimes_A A') \backslash (G^0 \otimes_A A').$$

Thus every finite subset of it is contained in an affine open subset. Moreover, formation of this cokernel commutes with finite locally free extensions of the base (cf. editor's note 37 and 4.6 below). As in 3.2.3, it follows that this cokernel has the form  $Y \otimes_A A'$ , where  $Y$  is a cokernel of  $\mathcal{S}(G_*^\alpha)$ .

**4.5.**— We have now constructed  $F \backslash G$  and shown that it is a direct sum of schemes of finite type over  $A$ . The remaining assertions of Theorem 3.2 reduce directly to assertions about the groupoids  $\mathcal{S}(G_*^\alpha)$ . As in Exposé V, § 6, the second assertion in (i) follows from the first together with (ii) and (iii); it is therefore enough to prove (ii), (iii), and (iv)(a).

Since  $A'$  is a local  $A$ -algebra, finite and free over  $A$ , the morphism  $A \rightarrow A'$  is faithfully flat and of finite presentation. By SGA 1, VIII (3.1, 4.6, 5.4), it is therefore enough to verify the corresponding assertions for the groupoid

$$\mathcal{S}(G_*^\alpha) \otimes_A A'.$$

This groupoid is isomorphic to a direct sum of finitely many copies of  $\mathcal{S}(G_*^0) \otimes_A A'$  (cf. 4.4), so we are reduced to the groupoid  $\mathcal{S}(G_*^0)$ . For this last groupoid we continue to follow the proof given in Exposé V, § 6, as we began to do in 4.3.

**4.6.**— Let us conclude this section with some remarks concerning Lemma 6.1 and § 9(a) of Exposé V. Under the hypotheses and with the notation of Exposé V, § 9(a), we seek a condition under which formation of the cokernel of the  $(\text{Sch}/S)$ -groupoid  $X_*$  commutes with a base extension

$$\pi : S' \longrightarrow S.$$

The cokernels of  $X_*$  and  $X'_*$  are identified with the cokernels of the groupoids  $U_*$  and  $U'_*$  induced by  $X_*$  and  $X'_*$  on the quasi-sections  $U$  and  $U'$ , respectively. We are therefore reduced to the case of a  $(\text{Sch}/S)$ -groupoid satisfying the hypotheses of Exposé V, Theorem 4.1.

<sup>45</sup>Let  $Y$  be the cokernel of  $U_*$ , put

$$Y' = Y \times_S S',$$

---

<sup>45</sup>Editor's note: we have added the following sentence.

and let  $Y_1$  be the cokernel of  $U'_*$ ; we saw in Exposé V, § 9(a), that the canonical morphism

$$Y_1 \longrightarrow Y'$$

is a homeomorphism (indeed, a universal homeomorphism); we may thus identify  $Y_1$  and  $Y'$  as topological spaces. If  $p : U \rightarrow Y$  is the canonical morphism and  $p' : U' \rightarrow Y'$  is obtained from it by base change, we want the following sequence of  $\mathcal{O}_{Y'}$ -modules to be exact:

$$\mathcal{O}_{Y'} \longrightarrow p'_*(\mathcal{O}_{U'}) \rightrightarrows p'_*u'_{1*}(\mathcal{O}_{U'_1}) = p'_*u'_{0*}(\mathcal{O}_{U'_1}). \quad (*)$$

<sup>46</sup>Since we are under the hypotheses of Exposé V, Theorem 4.1,  $u_0$  and  $u_1$  are finite and locally free; moreover,  $p$  is integral by Exposé V, 4.1(ii). Thus  $p$  and  $p \circ u_i$  are affine, hence separated and quasi-compact.

Consequently, if  $S'$  is flat over  $S$ , EGA III<sub>1</sub>, 1.4.15 (taking account of correction Err<sub>III</sub> 25 in EGA III<sub>2</sub>) identifies the sequence (\*) with the inverse image of the sequence

$$\mathcal{O}_Y \longrightarrow p_*(\mathcal{O}_U) \rightrightarrows p_*u_{1*}(\mathcal{O}_{U_1}) = p_*u_{0*}(\mathcal{O}_{U_1}), \quad (**)$$

which is exact.<sup>47</sup> An analogous argument applies when the groupoid  $X_*$  has quasi-sections “locally” (cf. the proof of Exposé V, Theorem 7.1). We therefore obtain:

**Proposition 4.6.1.** *Formation of the cokernel of  $X_*$  commutes with flat base extensions when  $X_*$  has quasi-sections locally.*

**4.7.**— Consider the groupoid  $G_*$  of Theorem 3.2, supposing temporarily that  $F$  and  $G$  are of finite type over  $A$ . By 3.2.2 there is a local  $A$ -algebra  $A'$ , finite and free over  $A$ , such that  $G_* \otimes_A A'$  has quasi-sections “locally.” For every base extension  $T \rightarrow \operatorname{Spec} A$ , the sequence

$$(F'' \setminus G'') \times_{\operatorname{Spec} A} T \rightrightarrows (F' \setminus G') \times_{\operatorname{Spec} A} T \longrightarrow (F \setminus G) \times_{\operatorname{Spec} A} T$$

obtained from diagram (\*) of 3.2.3 is exact.

If, in addition,  $T$  is flat over  $\operatorname{Spec} A$ , then 4.6 identifies

$$(F'' \setminus G'') \times_{\operatorname{Spec} A} T \quad \text{and} \quad (F' \setminus G') \times_{\operatorname{Spec} A} T$$

respectively with the cokernels of the groupoids

$$(G_* \otimes_A A'') \times_{\operatorname{Spec} A} T \quad \text{and} \quad (G_* \otimes_A A') \times_{\operatorname{Spec} A} T.$$

The diagram obtained from 3.2.3(\*) by the base change  $T \rightarrow \operatorname{Spec} A$  then shows that

$$(F \setminus G) \times_{\operatorname{Spec} A} T$$

is the cokernel of  $G_* \times_{\operatorname{Spec} A} T$ . The same argument applies in the general case, when  $F$  and  $G$  are merely locally of finite type over  $A$ . Thus:

**Proposition 4.7.1.** *Under the hypotheses of Theorem 3.2, for every flat  $A$ -scheme  $T$ , the scheme*

$$(F \setminus G) \times_{\operatorname{Spec} A} T$$

*is identified with the left quotient of  $G \times_{\operatorname{Spec} A} T$  by  $F \times_{\operatorname{Spec} A} T$ .<sup>48</sup>*

<sup>46</sup>Editor’s note: in what follows we have modified the original; the additional hypotheses imposed there on  $X_*$  are unnecessary.

<sup>47</sup>Editor’s note: we have added both the following sentence and the numbering 4.6.1 below, in order to highlight the stated result.

<sup>48</sup>Editor’s note: we have added the numbering 4.7.1 in order to highlight this result.

## 5. Connections with Exposé IV and consequences

**5.1.**—<sup>49</sup> We retain the notation of § 3 and the hypotheses of Theorem 3.2. We then have the following commutative diagram:

$$\begin{array}{ccc}
 F \times_A G \times_A G & \xrightarrow{F \times \nu} & F \times_A G \\
 \text{pr}_2 \times G \downarrow & \lambda \times G \downarrow & \text{pr}_2 \downarrow \lambda \\
 G \times_A G & \xrightarrow{\nu} & G \\
 p \times G \downarrow & & p \downarrow \\
 (F \backslash G) \times_A G & \xrightarrow{\rho} & F \backslash G
 \end{array}$$

It satisfies

$$\text{pr}_2 \circ (F \times \nu) = \nu \circ (\text{pr}_2 \times G), \quad \lambda \circ (F \times \nu) = \nu \circ (\lambda \times G).$$

Moreover, since  $G$  is flat over  $A$ , the left-hand vertical sequence is exact by 4.7. Hence  $\nu$  induces a morphism of  $A$ -schemes

$$\rho : (F \backslash G) \times_A G \longrightarrow F \backslash G.$$

The morphism  $\rho$  makes  $G$  act on the right on  $F \backslash G$ , as is immediately checked; moreover, the canonical morphism  $G \rightarrow F \backslash G$  commutes with the right actions of  $G$  on  $G$  and on  $F \backslash G$ .<sup>50</sup> This proves the first assertion of Theorem 3.2(ii'). By 2.5.4, the connected components of

$$X = F \backslash G$$

are therefore of finite type, irreducible, and all of the same dimension.

To compute this dimension, we may suppose that  $A = k$  and that  $k$  is algebraically closed. By Exposé I, 2.3.3.1, the stabilizer of the  $k$ -point  $p(e)$  is represented by the fiber

$$H = p^{-1}(p(e)).$$

Since  $F \backslash G$  is the quotient of  $G$  by  $F$  in the category of ringed spaces, the underlying space of this fiber is  $u(F)$ . As  $\ker(u)$  is finite, we obtain

$$\dim H = \dim u(F) = \dim F.$$

It follows from 2.5.4(ii) that

$$\dim X = \dim G - \dim F.$$

This proves Theorem 3.2(ii'), and hence also Theorem 3.3.2(iv).

**5.2.**— When the homomorphism of  $A$ -groups  $u : F \rightarrow G$  is a monomorphism, the result of 5.1 may also be recovered from the results of Exposé IV. Indeed, the canonical morphism

$$p : G \longrightarrow F \backslash G$$

is faithfully flat and open by 3.2; it is therefore a covering for the (fpqc) topology by Exposé IV, 6.3.1, and we may apply Exposé IV, Corollaries 5.2.2 and 5.2.4.

<sup>49</sup>Editor's note: we have changed the title of this section, which was called "Complements" in the original.

<sup>50</sup>Editor's note: we have added what follows.

In particular, suppose, in addition to the hypotheses of 3.2, that  $u$  is the inclusion in  $G$  of a normal subgroup  $F$ . Then there is a unique  $A$ -group structure on  $F \backslash G$  for which the canonical morphism

$$p : G \longrightarrow F \backslash G$$

is a homomorphism of  $A$ -groups.<sup>51</sup> This proves Theorem 3.3.2(v).

**5.3.**— We now review several statements from Exposé IV.

**5.3.1.**— The statements of Exposé IV, 5.2.7 and 5.3.1 translate as follows. Let  $F$  and  $G$  be groups locally of finite type and flat over  $A$ , with  $F$  a closed normal subgroup of  $G$ . The maps

$$H \longmapsto F \backslash H, \quad H' \longmapsto H' \times_{(F \backslash G)} G$$

define a bijection between the flat  $A$ -subgroups of  $G$  containing  $F$  and the flat  $A$ -subgroups of  $F \backslash G$ . Under this bijection, the closed (respectively, normal) subgroups of  $G$  containing  $F$  correspond to the closed (respectively, normal) subgroups of  $F \backslash G$ .<sup>52</sup>

**5.3.2.**— Exposé IV, Proposition 5.2.9, gives the following result. Let  $F, H, G$  be groups locally of finite type and flat over  $A$ , and suppose

$$F \subset H \subset G,$$

with  $F$  closed in  $G$  and normal in  $H$ . Then  $F \backslash H$  acts freely on the left on  $F \backslash G$ , the quotient scheme

$$(F \backslash H) \backslash (F \backslash G)$$

exists, and there is a canonical isomorphism of  $G$ -schemes

$$(F \backslash H) \backslash (F \backslash G) \xrightarrow{\sim} H \backslash G.$$

**5.3.3.**— Finally, Exposé IV, 5.2.8, gives the following assertion. Let  $F, H, G$  be groups locally of finite type and flat over  $A$ . Suppose that  $F$  is contained in  $G$ , closed and normal in  $G$ , that  $H$  is contained in  $G$ , and that  $F \cap H$  is flat over  $A$ . Let

$$F \times_A^\tau H$$

be the  $A$ -group whose underlying scheme is  $F \times_A H$ , with multiplication defined by

$$((x, h), (y, h')) \longmapsto (xhyh^{-1}, hh').$$

Likewise, let

$$u : H \cap F \longrightarrow F \times_A^\tau H$$

be the monomorphism  $x \mapsto (x^{-1}, x)$ , and let  $F \cdot H$  be the quotient

$$(F \cap H) \backslash (F \times_A^\tau H).$$

Then there is a canonical isomorphism

$$F \backslash (F \cdot H) \xrightarrow{\sim} (F \cap H) \backslash H.$$

<sup>51</sup>Editor's note: we have added the following sentence.

<sup>52</sup>Editor's note: in addition to the cited statements of Exposé IV, we use the fact that, since  $G \rightarrow F \backslash G$  is faithfully flat, an  $A$ -subgroup  $H$  of  $G$  is flat over  $A$  if and only if  $F \backslash H$  is.

**5.4.**—<sup>53</sup> Let  $u : G \rightarrow H$  be a quasi-compact morphism between  $A$ -groups locally of finite type, such that the kernel  $N$  of  $u$  is flat over  $A$ . In this case, by 3.3.2 and 5.2, the quotient  $A$ -group

$$C = N \backslash G$$

exists, and the morphism  $p : G \rightarrow C$  is faithfully flat and locally of finite presentation. On the other hand, by Exposé IV, 5.2.6,  $u$  induces a monomorphism

$$v : C \longrightarrow H,$$

which is quasi-compact (because  $u$  is quasi-compact and  $G \rightarrow C$  is surjective; cf. EGA IV<sub>1</sub>, 1.1.3), and is therefore a closed immersion by 2.5.2. We have thus obtained the following proposition.

**Proposition 5.4.1.**— *Let  $u : G \rightarrow H$  be a quasi-compact morphism between  $A$ -groups locally of finite type, such that  $N = \ker u$  is flat over  $A$ . Then  $u$  admits the factorization*

$$\begin{array}{ccc} G & \xrightarrow{u} & H \\ p \downarrow & \nearrow i & \\ N \backslash G & & \end{array}$$

where  $p$  is faithfully flat and locally of finite presentation, and  $i$  is a closed immersion.

Suppose in addition that  $G$  is flat over  $A$ . Then, by 3.3.2,  $C = N \backslash G$  is flat over  $A$ , and hence the quotient

$$X = C \backslash H$$

exists in  $(\mathbf{Sch}/_A)$  and represents the quotient (fppf)-sheaf  $\tilde{C} \backslash H$ , while

$$q : H \longrightarrow X$$

is a  $C$ -torsor. Consequently, if

$$e : \operatorname{Spec} A \longrightarrow H$$

denotes the unit section of  $H$ , then  $v$  induces an isomorphism of (fppf)-sheaves between  $\tilde{C}$  and the fiber product of  $q$  with

$$q \circ e : \operatorname{Spec} A \longrightarrow X,$$

which is represented by a closed subscheme of  $H$ . Consequently,  $v$  is an isomorphism of  $C$  onto a closed group subscheme  $K$  of  $H$ , equal to the stabilizer of the  $A$ -point  $q \circ e$  of  $X$ .<sup>\*</sup> (This also gives another proof that every quasi-compact monomorphism  $v : C \rightarrow H$  between  $A$ -groups locally of finite type is a closed immersion; cf. 2.5.2 and Exposé VI<sub>B</sub>, 1.4.2.)

Suppose further that  $C$  is a normal subgroup of  $H$ . In this case, the  $A$ -group

$$\overline{H} = C \backslash H$$

is the cokernel, in the category of  $A$ -groups, of the morphism  $u : G \rightarrow H$ , and  $K$  is the kernel of the morphism

$$H \longrightarrow \overline{H}.$$

<sup>53</sup>Editor's note: we have expanded the original in what follows; in particular, we have added Proposition 5.4.1.

<sup>\*</sup>Translator's note: the French source declares  $e : \operatorname{Spec} A \rightarrow G$ , although the displayed composition  $q \circ e$  requires the unit section of  $H$ , and then prints "a closed group subscheme  $K$  of  $G$ ." Here  $v : C \rightarrow H$ , the displayed fiber product is represented inside  $H$ , and the next paragraph identifies  $K$  with  $\ker(H \rightarrow \overline{H})$ ; accordingly,  $e$  is taken to be the unit section of  $H$ , and the intended ambient group of  $K$  is  $H$ .



When  $G$  and  $H$  are abelian  $A$ -groups,  $K$  is the image of  $u$  in the category of abelian  $A$ -groups, whereas

$$C = (\ker u) \backslash G$$

is the coimage of  $u$ . Taking into account the isomorphism

$$C \xrightarrow{\sim} K$$

just established, we obtain the following.<sup>53</sup>

**Theorem 5.4.2.**— *Let  $k$  be a field. The category of commutative algebraic  $k$ -groups is abelian.*

Indeed, when  $k$  is a field,  $\ker(u)$  is flat over  $k$ , whatever  $u$  may be.

<sup>54</sup> Note that the full subcategory of commutative affine algebraic  $k$ -groups is thick. Indeed, consider an exact sequence of commutative algebraic  $k$ -groups

$$1 \longrightarrow N \longrightarrow G \longrightarrow G/N \longrightarrow 1.$$

If  $G$  is affine, then  $N$  is clearly affine, and  $G/N$  is also affine by a theorem of Chevalley; cf. Exposé VI<sub>B</sub>, 11.17. Conversely, if  $N$  and  $G/N$  are affine, then  $G$  is also affine by Exposé VI<sub>B</sub>, 9.2(viii). We therefore obtain the following corollary.

**Corollary 5.4.3.**— *Let  $k$  be a field. The category of commutative affine algebraic  $k$ -groups is abelian.*

Let us also point out that the category of all commutative affine  $k$ -groups (not necessarily of finite type) is abelian. This follows from Exposé VI<sub>B</sub>, 11.17 and 11.18.2 (cf. [DG70], § III.3, 7.4); see also Exposé VII<sub>B</sub>, 2.4.2 for a proof using formal groups.

**5.5.**— Let  $G$  be a group locally of finite type and flat over an Artinian local ring  $A$ . We know from 2.3 that the connected component  $G^0$  of the identity is a normal open group subscheme of  $G$ , hence is also flat over  $A$ . It then follows from 3.2 and 5.2 that

$$G^0 \backslash G$$

is an  $A$ -group scheme, flat over  $A$ . Moreover, since every connected component  $G^\alpha$  of  $G$  is saturated for the equivalence relation defined by  $G^0$ , the quotient  $G^0 \backslash G$  is the disjoint union of the  $G^0 \backslash G^\alpha$  (cf. 4.3). In particular, the connected component of the identity in  $G^0 \backslash G$  is precisely

$$G^0 \backslash G^0 \simeq \operatorname{Spec} A,$$

and hence the morphism

$$G^0 \backslash G \longrightarrow \operatorname{Spec} A$$

is a local isomorphism at the identity. Consequently,  $G^0 \backslash G$  is étale over  $\operatorname{Spec} A$ , by Exposé VI<sub>B</sub>, 1.3.<sup>55</sup> We therefore obtain the following proposition (for assertion (ii), see [DG70], § II.5, 1.7–1.10).

**Proposition 5.5.1.**— *Let  $A$  be an Artinian local ring and  $G$  an  $A$ -group locally of finite type and flat.*

(i)  $G^0 \backslash G$  is an étale  $A$ -group.

<sup>53</sup>Editor's note: we have added the number 5.4.2 to this theorem.

<sup>54</sup>Editor's note: we have added what follows.

<sup>55</sup>Editor's note: we have expanded the preceding discussion from the original and added Proposition 5.5.1 in order to make this result explicit.

- (ii) Consequently, if  $A = k$  is an algebraically closed field,  $G^0 \backslash G$  is a constant  $k$ -group acting simply transitively on the set of connected components of  $G$ . Thus, if  $G$  is algebraic,  $G^0 \backslash G$  is finite.

**5.6.**— Now let  $k$  be a perfect field and  $G$  a  $k$ -group locally of finite type. We saw in 0.2 that  $G_{\text{red}}$  is then a group subscheme of  $G$ . Moreover, the equivalence class of the identity of  $G$  under the left action of  $G_{\text{red}}$  on  $G$  is the whole underlying space of  $G$ . Theorem 3.2 therefore gives the following result.

**Proposition 5.6.1.**— *Let  $k$  be a perfect field and  $G$  a  $k$ -group locally of finite type. Then the  $k$ -scheme*

$$G_{\text{red}} \backslash G$$

*is the spectrum of a finite local  $k$ -algebra with residue field  $k$ .*

<sup>56</sup> Indeed, by 3.2, the quotient  $G_{\text{red}} \backslash G$  has a single point, with residue field  $k$ , and is a  $k$ -scheme of finite type. It is therefore the spectrum of a finite-dimensional local  $k$ -algebra (cf. EGA I, 6.4.4).

**Proposition 5.6.2.**— *Let  $u : F \rightarrow G$  be a morphism between groups locally of finite type over a perfect field  $k$ . The following assertions are equivalent:*

- (i)  *$u$  is flat.*  
(ii) *The morphism*

$$u^0 : F^0 \longrightarrow G^0$$

*is dominant, and the morphism*

$$v : F_{\text{red}} \backslash F \longrightarrow G_{\text{red}} \backslash G$$

*induced by  $u$  is flat.*

<sup>57</sup> Indeed, consider the following commutative diagram:

$$\begin{array}{ccc} F & \xrightarrow{p} & F_{\text{red}} \backslash F \\ \downarrow u & & \downarrow v \\ G & \xrightarrow{q} & G_{\text{red}} \backslash G \end{array}$$

where  $p$  and  $q$  denote the canonical projections. By 3.2(iv),  $p$  and  $q$  are faithfully flat. Consequently, if  $u$  is flat, then

$$q \circ u = v \circ p$$

is flat, and hence  $v$  is flat as well.

Conversely, suppose that  $v$  is flat and  $u^0$  is dominant. Since  $u^0$  is quasi-compact (the scheme  $F^0$  is of finite type over  $k$  by 2.4, and hence Noetherian), it maps the generic point  $\xi$  of  $F^0$  to the generic point  $\eta$  of  $G^0$ . Let  $R$  be the finite local  $k$ -algebra such that

$$G_{\text{red}} \backslash G = \text{Spec } R,$$

and let  $\mathfrak{m}$  be its maximal ideal. We have local homomorphisms of local rings

$$R \longrightarrow \mathcal{O}_{G, \eta} \longrightarrow \mathcal{O}_{F, \xi}.$$

Observe that the following square is Cartesian:

<sup>56</sup>Editor's note: we have added the number 5.6.1, as well as the proof that follows.

<sup>57</sup>Editor's note: we have expanded the proof of (ii)  $\Rightarrow$  (i) and simplified the diagram below.

$$\begin{array}{ccc}
G_{\text{red}} & \longrightarrow & G \\
\downarrow & & \downarrow q \\
\text{Spec}(R/\mathfrak{m}) & \longrightarrow & \text{Spec}(R)
\end{array}$$

It follows that

$$\mathcal{O}_{G,\eta}/\mathfrak{m}\mathcal{O}_{G,\eta} \simeq \mathcal{O}_{G_{\text{red}},\eta} = \kappa(\eta),$$

so that

$$\mathcal{O}_{F,\xi}/\mathfrak{m}\mathcal{O}_{F,\xi}$$

is flat over  $\mathcal{O}_{G,\eta}/\mathfrak{m}\mathcal{O}_{G,\eta}$ .

On the other hand, since  $q$  and  $v \circ p$  are flat,  $G$  and  $F$  are flat over  $R$ . The local criterion for flatness (cf. EGA IV<sub>3</sub>, 11.3.10.2) therefore shows that  $\mathcal{O}_{F,\xi}$  is flat over  $\mathcal{O}_{G,\eta}$ ; in other words,  $u$  is flat at  $\xi$ . It follows from 2.5.3 that  $u$  is flat.

## 6. Complements on $k$ -groups not necessarily of finite type

<sup>58</sup> Let us also record the following results, which will be useful in the addendum, Exposé VI<sub>B</sub>, § 12. We fix a base field  $k$ .

**Lemma 6.1.**— *Let  $G$  be a  $k$ -group. For every  $x \in G$ , there exists a point  $u \in G \times G$  such that  $\mu(u) = x$  and both projections  $p_1(u)$  and  $p_2(u)$  are maximal points of  $G$ .*

*Proof.* Set  $K = \kappa(x)$ . Since the projection  $G_K \rightarrow G$  sends maximal points to maximal points, we are reduced to the case in which  $x$  is rational. Left translation  $\lambda_x$  (respectively, right translation  $\rho_x$ ) then gives a morphism

$$G \longrightarrow G \times G, \quad g \longmapsto (\lambda_x(g^{-1}), g)$$

(respectively,  $g \longmapsto (g, \rho_x(g^{-1}))$ ), which induces an isomorphism from  $G$  onto  $\mu^{-1}(x)$ , inverse to  $p_2$  (respectively,  $p_1$ ). Thus, if  $u$  is a maximal point of  $\mu^{-1}(x)$ , then  $p_1(u)$  and  $p_2(u)$  are maximal points of  $G$ , and  $u$  has the required properties.

**Corollary 6.2.**— *Let  $f : G \rightarrow H$  be a quasi-compact dominant morphism of  $k$ -groups.*

- (i)  *$f$  is surjective.*
- (ii) *If  $H$  is reduced,  $f$  is faithfully flat.*

*Proof.* Write  $\mu_H$  (respectively,  $\mu_G$ ) for the multiplication of  $H$  (respectively,  $G$ ). Let  $h \in H$ . By 6.1, there exists  $u \in H \times H$  such that  $\mu_H(u) = h$  and  $\alpha = p_1(u)$ ,  $\beta = p_2(u)$  are maximal points of  $H$ . Since  $f$  is quasi-compact and dominant,  $f^{-1}(\alpha)$  and  $f^{-1}(\beta)$  are nonempty (cf. EGA IV<sub>1</sub>, 1.1.5). Hence there exists  $v \in G \times G$  such that

$$(f \times f)(v) = u$$

(cf. EGA I, 3.5.2). Then  $g = \mu_G(v)$  satisfies  $f(g) = h$ . This proves that  $f$  is surjective.

Suppose in addition that  $H$  is reduced. Then  $\mathcal{O}_{H,\alpha}$  is a field, and we have seen above that  $f^{-1}(\alpha) \neq \emptyset$ . Thus  $f$  is flat at every point  $\xi \in f^{-1}(\alpha)$ , and consequently  $f$  is flat by Lemma 2.5.3.

---

<sup>58</sup>Editor's note: we have added the results that follow, taken from [Per75]. Note that Lemma 6.1 can be expressed, in Weil's language, by saying that "every point of  $G$  is a product of two generic points."

**Reminder 6.3.**— Recall (cf. EGA IV<sub>3</sub>, 11.10.1) that a morphism of schemes  $f : X \rightarrow Y$  is said to be *schematically dominant* if it satisfies the following condition: for every open subset  $U$  of  $Y$ , if  $Z$  is a closed subscheme of  $U$  such that the morphism  $f^{-1}(U) \rightarrow U$  factors through  $Z$ , then  $Z = U$ . When  $f$  is quasi-compact and quasi-separated, this is equivalent to saying that the closed image of  $X$  under  $f$  is  $Y$  (cf. loc. cit., 11.10.3(iv), and EGA I, 9.5.8).

**Proposition 6.4.**— *Let  $f : H \rightarrow G$  be a quasi-compact morphism of  $k$ -groups. Then the closed image of  $f$  is a group subscheme  $H'$  of  $G$ , and  $f$  factors as*

$$\begin{array}{ccc} H & \xrightarrow{f} & G \\ f' \downarrow & \nearrow i & \\ H' & & \end{array}$$

where  $f'$  is schematically dominant, quasi-compact, and surjective.

*Proof.* Since  $H$  is separated (0.3),  $f$  is quasi-compact and separated. Thus  $f_*(\mathcal{O}_H)$  is a quasi-coherent  $\mathcal{O}_G$ -module, and the closed image  $H'$  of  $f$  exists and is the closed subscheme of  $G$  defined by the quasi-coherent ideal

$$\mathcal{I} = \ker(\mathcal{O}_G \rightarrow f_*(\mathcal{O}_H))$$

(cf. EGA I, § 9.5).

Write  $c_G, \mu_G$  (respectively,  $c_H, \mu_H$ ) for the inversion and multiplication morphisms of  $G$  (respectively,  $H$ ). Then  $c_G \circ f = f \circ c_H$  factors through the closed subscheme  $c_G(H')$ , whence

$$H' \subset c_G(H') \quad \text{and therefore} \quad H' = c_G(H'),$$

since  $c_G^2 = \text{id}_G$ . Similarly,

$$f \circ \mu_H = \mu_G \circ (f \times f)$$

factors through  $H'$ , so  $f \times f$  factors through the closed subscheme  $\mu_G^{-1}(H')$  of  $G \times G$ .<sup>\*</sup> On the other hand, since formation of the closed image commutes with flat base change (EGA III, 1.4.15, and EGA IV<sub>1</sub>, 1.7.21), the closed image of  $f \times \text{id}_H$  (respectively,  $\text{id}_{H'} \times f$ ) is  $H' \times H$  (respectively,  $H' \times H'$ ). Hence, by “transitivity of closed images” (EGA I, 9.5.5), the closed image of  $f \times f$  is  $H' \times H'$ . It is therefore contained in  $\mu_G^{-1}(H')$ ; that is, the restriction of  $\mu_G$  to  $H' \times H'$  factors through  $H'$ . This proves that  $H'$  is a closed group subscheme of  $G$ . Write  $i : H' \hookrightarrow G$  for the inclusion.

Now  $f = i \circ f'$ , where  $f' : H \rightarrow H'$  is schematically dominant and quasi-compact (because  $f$  is quasi-compact and  $i$  is separated). By 6.2,  $f'$  is therefore surjective. This proves 6.4.

We can now state the following theorem ([Per75], V, 3.1 and 3.2; see also [Per76], 0.0 and 0.1).

**Theorem 6.5 (D. Perrin).**— *Let  $G$  be a quasi-compact  $k$ -group.*

- (i)  *$G$  is the projective limit of a filtered system  $(G_i)$  of  $k$ -groups of finite type (whose transition morphisms  $u_{ij} : G_j \rightarrow G_i$  are affine for  $i$  sufficiently large), and the morphisms  $G \rightarrow G_i$  are faithfully flat (and affine for  $i$  sufficiently large).*
- (ii) *Let  $H$  be a closed  $k$ -subgroup of  $G$ . Then the fpqc quotient sheaf  $\widetilde{G/H}$  is a  $k$ -scheme in either of the following two cases:*

<sup>\*</sup>Translator’s note: in these two occurrences the French source prints  $\pi_H$  and  $\pi_G$ , although the immediately preceding sentence defines the multiplication morphisms as  $\mu_H$  and  $\mu_G$ . The multiplication symbols required by the identity are used here.

- (1) *the immersion  $H \rightarrow G$  is of finite presentation; in this case  $G/H$  is of finite type over  $k$ ;*
- (2)  *$H$  is normal in  $G$ .*

For the proof of this theorem, which rests on several intermediate theorems, we refer to [Per75]. For the reader's convenience, however, we prove the two corollaries below; cf. [Per75], V, 3.3–3.4, or [Per76], 4.2.3–4.2.5.

**Corollary 6.6.**— *Let  $f : G \rightarrow H$  be a quasi-compact morphism of  $k$ -groups.*

- (i) *If  $f$  is schematically dominant, then it is faithfully flat.*
- (ii) *This is so, in particular, if  $H$  is affine and the morphism*

$$f^\natural : \mathcal{O}(H) \longrightarrow \mathcal{O}(G)$$

*is injective.*

*Proof.* Suppose that  $f$  is schematically dominant. By 6.2(i),  $f$  is surjective, so by 2.5.3 it is enough to prove that  $f$  is flat at the identity element of  $G$ . Since

$$\mathcal{O}_{H,e} = \mathcal{O}_{H^0,e}$$

(cf. 2.6.5), we may replace  $H$  by  $H^0$ , and hence suppose that  $H$  is irreducible. Then  $H$  is quasi-compact (loc. cit.); since  $f$  is quasi-compact,  $G$  is also quasi-compact. By 6.5(i),

$$H = \varprojlim_i H_i,$$

where each  $H_i$  is an algebraic  $k$ -group. Let  $N_i$  be the kernel of  $G \rightarrow H_i$ ; it is a closed normal  $k$ -subgroup of  $G$ . Moreover, since the identity section  $\{e\} \rightarrow H_i$  is of finite presentation, the immersion  $N_i \rightarrow G$  is of finite presentation. Hence, by 6.5(ii), the fpqc quotient  $G/N_i$  is an algebraic  $k$ -group. We then have a commutative diagram

$$\begin{array}{ccc} G & \xrightarrow{f} & H \\ q_i \downarrow & & \downarrow p_i \\ G/N_i & \xrightarrow{f_i} & H_i \end{array}$$

in which  $f$  is schematically dominant and  $p_i, q_i$  are faithfully flat, hence schematically dominant. Thus

$$f_i \circ q_i = p_i \circ f$$

is schematically dominant, and therefore so is  $f_i$ . On the other hand,  $f_i$  is a monomorphism, hence a closed immersion because  $G/N_i$  and  $H_i$  are algebraic (2.5.2). It follows that  $f_i$  is an isomorphism, and consequently  $G \rightarrow H_i$  is faithfully flat. By [BAC], I, § 2.7, Proposition 9, the morphism  $G \rightarrow H$  is then flat; it is also surjective by 6.2(i), and is therefore faithfully flat. This proves (i). Assertion (ii) follows, because if  $H$  is affine and  $f^\natural$  is injective, then the closed image of  $f$  is  $H$ , so  $f$  is schematically dominant (cf. 6.4).

**Corollary 6.7.**— *Let  $u : G \rightarrow H$  be a morphism of  $k$ -groups and set  $N = \ker(u)$ . Suppose that  $u$  is quasi-compact.*

- (i) *The fpqc quotient sheaf  $\widetilde{G/N}$  is represented by a  $k$ -group scheme  $G/N$ , and  $u$  factors as*

$$\begin{array}{ccc}
 G & \xrightarrow{u} & H \\
 p \downarrow & \nearrow i & \\
 G/N & & 
 \end{array}$$

where  $p$  is faithfully flat and  $i$  is a closed immersion.

- (ii) In particular, if  $u$  is a monomorphism, then  $u$  is a closed immersion; and if  $u$  is schematically dominant, then  $u$  is faithfully flat.

*Proof.* (i) By 6.4 and 6.6, the closed image  $G'$  of  $u$  is a closed group subscheme of  $H$ , and

$$p : G \longrightarrow G'$$

is faithfully flat and quasi-compact.\* Clearly  $\ker(p) = \ker(u) = N$ . Thus, by Exposé IV, 3.3.2.1 and 5.1.7,  $G'$  represents the fpqc quotient sheaf  $\widehat{G/N}$ .

(ii) The second assertion is contained in 6.6; let us prove the first. If  $u$  is a monomorphism, then so is  $p$ . Thus  $p$  is both a monomorphism and an effective epimorphism, and is therefore an isomorphism (cf. Exposé IV, 1.14). This proves 6.7.

Let us finally record the following corollaries (cf. [Per76], 4.2.6–4.2.8).

**Corollary 6.8.**— *The category of commutative quasi-compact  $k$ -group schemes is abelian.*

Taking 6.7 into account, the proof is analogous to that of 5.4.2.

**Corollary 6.9.**— *If  $\text{char}(k) = 0$ , every  $k$ -group scheme  $G$  is geometrically reduced.*

Indeed, if  $g \in G$  and  $K$  is an algebraic closure of  $\kappa(g)$ , then

$$\mathcal{O}_{G,e} \otimes_k K \simeq \mathcal{O}_{G_K, g_K}.$$

It is therefore enough to prove that  $\mathcal{O}_{G,e} = \mathcal{O}_{G^0,e}$  is geometrically reduced. We are thus reduced to the case in which  $G$  is connected, hence quasi-compact by 2.6.5. The result then follows from 6.5 and Cartier's theorem for algebraic groups (cf. Exposé VI<sub>B</sub>, 1.6.1, or [DG70], § II.6, Theorem 1.1).

**Corollary 6.10.**— *Let  $G$  be a quasi-compact  $k$ -group, and suppose that  $k$  is algebraically closed.*

- (i) If  $f : G \rightarrow H$  is a faithfully flat morphism of  $k$ -groups, then the induced map  $G(k) \rightarrow H(k)$  is surjective.
- (ii) The set of rational points is dense in  $G$ .

For the proof, we refer to [Per75], V, 3.7 and 3.9.

## Bibliography

59

[An73] S. Anantharaman, *Schémas en groupes, espaces homogènes et espaces algébriques sur une base de dimension 1*, Mém. Soc. Math. France **33** (1973), 5–79.

\*Translator's note: the French source prints “a closed group subscheme of  $G$ ” at this point. Since  $u : G \rightarrow H$ , its closed image lies in  $H$ , as the displayed factorization also shows.

<sup>59</sup>Editor's note: additional references cited in this Exposé.

- [BAC] N. Bourbaki, *Algèbre commutative*, Chapters I–IV and V–VII, Masson, 1985.
- [DG70] M. Demazure and P. Gabriel, *Groupes algébriques*, Masson & North-Holland, 1970.
- [Per75] D. Perrin, *Schémas en groupes quasi-compacts sur un corps*, Publ. Math. Orsay no. 165–75.46 (part 1), <http://portail.mathdoc.fr/PM0/>.
- [Per76] D. Perrin, *Approximation des schémas en groupes, quasi-compacts sur un corps*, Bull. Soc. Math. France **104** (1976), 323–335.
- [Ray67a] M. Raynaud, *Passage au quotient par une relation d'équivalence plate*, pp. 78–85 in *Proc. Conf. Local Fields (Driebergen)*, ed. T. A. Springer, Springer-Verlag, 1967.
- [Ray67b] M. Raynaud, *Sur le passage au quotient par un groupoïde plat*, C. R. Acad. Sci. Paris (Sér. A) **265** (1967), 384–387.
- [Ray70] M. Raynaud, *Faisceaux amples sur les schémas en groupes et les espaces homogènes*, Lecture Notes in Mathematics **119**, Springer-Verlag, 1970.

# Exposé VI<sub>B</sub>

## Generalities on Group Schemes

by J.-E. Bertin

<sup>0</sup> This Exposé, which does not correspond to any oral lecture in the seminar, is intended to bring together a number of commonly used technical results concerning group schemes.\*

### 1. Morphisms of groups locally of finite type over a field

**1.1.**— Let  $A$  be an Artinian local ring, let  $G$  and  $H$  be two  $A$ -groups,<sup>2</sup> and let  $u : G \rightarrow H$  be a morphism of  $A$ -groups. Then  $u$  induces a group morphism

$$u(A) : G(A) \longrightarrow H(A).$$

<sup>3</sup> Since  $H(A)$  acts on  $H$  by right translation, restriction along  $u(A)$  defines an action of  $G(A)$  on  $H$ . This action is compatible with  $u$  and with the action of  $G(A)$  on  $G$  by right translation. Since  $G(A)$  acts transitively on the strictly rational points of  $G$  (see Exposé VI<sub>A</sub>, 0.4 for the definition), these points “all behave in the same way with respect to  $u$ .” The following properties spring from this observation.

**Proposition 1.2.**— *Let  $u : G \rightarrow H$  be a quasi-compact morphism between  $A$ -groups locally of finite type over  $A$ . Then the set  $u(G)$  is closed in  $H$ , and*

$$\dim G = \dim u(G) + \dim \ker u.$$

4

Since  $u$  commutes with the inversion morphisms of  $G$  and  $H$ , the image  $u(G)$  is invariant under inversion on  $H$ ; the same is therefore true of its closure  $\overline{u(G)}$  in  $H$ . On the other hand, let  $L$  be the set of points of  $H \times_A H$  whose two projections belong to  $u(G)$ . It is clear that  $L$  is the image of

$$u \times_A u : G \times_A G \longrightarrow H \times_A H.$$

The multiplication morphism of  $H$  consequently sends  $L$  into  $u(G)$ ; in other words,

$$u(G) \cdot u(G) = u(G).$$

---

<sup>0</sup>Editor’s note: version of October 13, 2024.

\*This Exposé was quite substantially revised after its mimeographed edition; in particular, Sections 10 and 11 were completely rewritten.<sup>(1)</sup>

<sup>1</sup>Editor’s note: and also Section 5; cf. Editor’s notes 34 and 46.

<sup>2</sup>Editor’s note: recall the convention adopted at the beginning of Exposé VI<sub>A</sub>: an  $A$ -group is a group scheme over  $A$  (and every such scheme is separated; cf. Exposé VI<sub>A</sub>, 0.3).

<sup>3</sup>Editor’s note: we have replaced “homomorphism” by “morphism.”

<sup>4</sup>Editor’s note: this statement also appears in Exposé VI<sub>A</sub>, 2.5.2.



Lemma 1.2.1 below shows that the closure  $\overline{L}$  of  $L$  in  $H \times_A H$  is the set of points whose two projections belong to  $\overline{u(G)}$ . Hence

$$\overline{u(G)} \cdot \overline{u(G)} = \overline{u(G)}.$$

Thus the reduced subscheme of  $H$  whose underlying space is  $\overline{u(G)}$  is naturally equipped with a group structure in the category  $(\mathbf{Sch}/k)_{\text{red}}$ , where  $k$  is the residue field of  $A$  (cf. Exposé VI<sub>A</sub>, 0.2).

Let us prove the first assertion of 1.2. After replacing  $A$  by an algebraic closure of its residue field  $k$ , we may suppose that  $A$  is an algebraically closed field  $k$  (cf. EGA IV<sub>2</sub>, 2.3.12). Replacing  $u$  by

$$u_{\text{red}} : G_{\text{red}} \longrightarrow H_{\text{red}},$$

we may suppose that  $G$  and  $H$  are reduced. In this case, as we have just seen,  $u(G)$  is the underlying space of a reduced group subscheme of  $H$ ;\* we may therefore suppose that  $u$  is dominant. Then  $G(k)$  acts transitively on the set of connected components of  $H$ , and it is enough to prove that  $u(G) \cap H^0$  is closed. We are thus reduced to the case in which  $H$  is connected, hence irreducible and of finite type (Exposé VI<sub>A</sub>, 2.4). The morphism  $u$  is then of finite type, since it is quasi-compact and locally of finite type. As  $H$  is Noetherian,  $u(G)$  is constructible (EGA IV<sub>1</sub>, 1.8.5), and therefore contains an open subset  $V$  of  $H$  (EGA 0<sub>III</sub>, 9.2.2). By Exposé VI<sub>A</sub>, 0.5, we then have

$$H = V \cdot V \subset u(G) \cdot u(G) = u(G).$$

Let us prove the second assertion. Recall first that the functor  $\ker u$  (cf. Exposé I, 2.3.6.1) is represented by  $u^{-1}(e)$ , where  $e$  denotes the identity element of  $H$ . When  $u$  is locally of finite type,  $\ker u$  is therefore locally of finite type over  $A$ . As above, this time using EGA IV<sub>2</sub>, 4.1.4, we reduce to the case in which  $A$  is an algebraically closed field  $k$ .

We may further suppose that  $G$  and  $H$  are irreducible and of finite type and that  $u$  is dominant. Indeed, since  $k$  is algebraically closed, the connected components of  $G$ , on whose set  $G(k)$  acts transitively, clearly all have the same dimension; and if  $u^0$  denotes the restriction of  $u$  to  $G^0$ , then

$$(\ker u)^0 \subset \ker u^0, \quad \dim \ker u^0 = \dim \ker u.$$

Likewise,  $u$  is then of finite type over  $k$ . If  $\eta$  is the generic point of  $H$ , then

$$\dim u^{-1}(\eta) = \dim G - \dim H$$

(EGA IV<sub>3</sub>, 10.6.1(ii)). By EGA IV<sub>3</sub>, 9.2.3 and 9.2.6, the set of  $y \in H$  such that

$$\dim u^{-1}(y) = \dim u^{-1}(\eta)$$

contains a nonempty open subset  $V$ . Since  $u$  is dominant,  $U = u^{-1}(V)$  is a nonempty open subset of  $G$ ; it contains a closed point  $x$ , because  $G$  is a Jacobson scheme (EGA IV<sub>3</sub>, 10.4.7). Right translation  $r_x$  is an isomorphism from  $\ker u$  onto  $u^{-1}(u(x))$ , and hence

$$\dim \ker u = \dim u^{-1}(u(x)) = \dim u^{-1}(\eta) = \dim G - \dim H.$$

---

\*Translator's note: the French source prints "a reduced group subscheme of  $G$ " at this point. Since  $u(G) \subset H$ , and the preceding paragraph constructs the reduced subscheme inside  $H$ , the ambient group is  $H$ .

**Lemma 1.2.1.**— *Let  $f : X' \rightarrow X$  and  $g : Y' \rightarrow Y$  be two quasi-compact dominant morphisms of schemes over an Artinian local ring  $A$ . Then*

$$f \times_A g : X' \times_A Y' \longrightarrow X \times_A Y$$

*is dominant (and quasi-compact).*

Indeed,

$$f \times_A g = (\mathrm{id}_X \times_A g) \circ (f \times_A \mathrm{id}_{Y'}).$$

It is therefore enough to prove that  $f \times_A \mathrm{id}_{Y'}$  and  $\mathrm{id}_X \times_A g$  are dominant. For this purpose we may replace  $A$  by its residue field  $k$ . Then  $X$  and  $Y'$  are flat over  $A = k$ , and  $f \times_A \mathrm{id}_{Y'}$  (respectively,  $\mathrm{id}_X \times_A g$ ) is obtained from  $f$  (respectively,  $g$ ) by the flat base change  $Y' \rightarrow A$  (respectively,  $X \rightarrow A$ ). It is therefore dominant (and quasi-compact) by EGA IV<sub>2</sub>, 2.3.7.

**Counterexample 1.2.2.**— *Let  $k$  be a field of characteristic zero, let  $G$  be the constant  $k$ -group  $\mathbb{Z}$ , and let  $H$  be the additive  $k$ -group  $\mathbb{G}_{a,k}$ . Let  $u : G \rightarrow H$  be a morphism of  $k$ -groups. If  $u \neq 0$ , then  $u(G)$  is not closed in  $H$ .*

**Proposition 1.3.**—<sup>5</sup> *Let  $A$  be an Artinian local ring with residue field  $k$ , let  $G$  be an  $A$ -group locally of finite type and flat, and let  $u : G \rightarrow H$  be a morphism of  $A$ -groups. The following assertions are equivalent:*

- (i)  *$u$  is flat (respectively, quasi-finite, unramified, smooth, or étale) at some point of  $G$ ;*<sup>6</sup>
- (ii)  *$u$  is flat (respectively, quasi-finite, unramified, smooth, or étale).*

*Proof.* It is enough to prove that (i) implies (ii). We first establish the following lemma.

**Lemma 1.3.0.**— *Let  $A \rightarrow B \rightarrow C$  be morphisms of commutative rings, and let  $\mathfrak{n}$  be a nilpotent ideal of  $A$ . Suppose that  $C/\mathfrak{n}C$  is a  $(B/\mathfrak{n}B)$ -algebra of finite type.*

- (i) *Then  $C$  is a  $B$ -algebra of finite type.*
- (ii) *Moreover, if  $C$  is flat over  $A$  and  $C/\mathfrak{n}C$  is a  $(B/\mathfrak{n}B)$ -algebra of finite presentation, then  $C$  is a  $B$ -algebra of finite presentation.*

Indeed, let  $x_1, \dots, x_n$  be elements of  $C$  whose images generate  $C/\mathfrak{n}C$  as a  $(B/\mathfrak{n}B)$ -algebra. By the nilpotent Nakayama lemma, the  $x_i$  generate  $C$  as a  $B$ -algebra. This proves (i). Let

$$\varphi : B[X_1, \dots, X_n] \longrightarrow C$$

be the resulting surjective morphism, and put  $I = \ker \varphi$ .

Suppose now that  $C$  is flat over  $A$  and that  $C/\mathfrak{n}C$  is of finite presentation over  $B/\mathfrak{n}B$ . On the one hand,  $I/\mathfrak{n}I$  identifies with the kernel of

$$\overline{\varphi} = \varphi \otimes_A (A/\mathfrak{n}).$$

<sup>5</sup>Editor's note: in the original, 1.3, 1.3.1, and 1.3.1.1 were stated with  $A$  a field. For completeness, we treat the case of an Artinian local ring, and for this purpose have added Lemma 1.3.0.

<sup>6</sup>Editor's note: recall that a morphism of schemes  $f : X \rightarrow Y$  is said to be of finite type (respectively, of finite presentation or quasi-finite) at  $x$  if there are affine open subsets  $V$ , containing  $f(x)$ , and  $U$ , containing  $x$ , such that  $f(U) \subset V$  and  $\mathcal{O}_X(U)$  is an  $\mathcal{O}_Y(V)$ -algebra of finite type (respectively, of finite presentation, or with the additional condition that  $f^{-1}(f(x'))$  is finite for every  $x' \in U$ ); see also Editor's note 40 of Exposé V. The morphism is locally quasi-finite (respectively, locally of finite type or locally of finite presentation) if it has the property at every point  $x$ . It is smooth (respectively, unramified or étale) at  $x$  if there is an open neighborhood  $U$  of  $x$  such that  $f|_U : U \rightarrow Y$  is smooth (respectively, unramified or étale). These definitions show that the loci where  $f$  is of finite presentation, of finite type, quasi-finite, smooth, unramified, or étale are open in  $X$ . Moreover, because the flatness locus of a morphism locally of finite presentation is open (EGA IV<sub>3</sub>, 11.3.1), the locus where  $f$  is both of finite presentation and flat is open as well. These facts will be used repeatedly below.

On the other hand, by EGA IV<sub>1</sub>, 1.4.4,  $\ker \bar{\varphi}$  is an ideal of finite type. Choose  $P_1, \dots, P_s \in I$  whose images generate  $I/\mathfrak{n}I$  as an ideal. The nilpotent Nakayama lemma shows that the  $P_i$  generate  $I$ . This proves (ii).

We return to the proof of 1.3. Let  $x$  be an arbitrary point of  $G$ . Since  $G$  is flat over  $A$ , the fiberwise flatness criterion, in the form EGA IV<sub>3</sub>, 11.3.10.2, shows that  $u$  is flat at  $x$  if  $u \otimes_A k$  is. Likewise, the preceding lemma shows that  $u$  is of finite type (respectively, of finite presentation) at  $x$  if  $u \otimes_A k$  is. The remaining properties can then be checked on the fibers (cf. EGA IV<sub>4</sub>, 17.4.1, 17.5.1, and 17.6.1 for the unramified, smooth, and étale cases). We are therefore reduced to the case  $A = k$ .

Now let  $x$  be a point of  $G$  at which one of the conditions in 1.3(i) holds. The properties under consideration are preserved by fpqc descent (cf. EGA IV, 2.5.1, 2.7.1, and 17.7.1). Thus, after replacing  $k$  by an algebraic closure of  $\kappa(x)$ , we reduce to the case in which  $k$  is algebraically closed and  $x \in G(k)$ .

Since  $G$  is a Jacobson scheme (cf. EGA IV<sub>3</sub>, 10.4.7), and the set  $W$  of points of  $G$  at which  $u$  is flat (respectively, quasi-finite, unramified, smooth, or étale) is stable under generization<sup>7</sup> (respectively, open), it is enough to show that every  $y \in G(k)$  belongs to  $W$ . For such a point  $y$ , the translation

$$r_y \circ r_x^{-1}$$

sends  $x$  to  $y$ , so  $u$  has the required property at  $y$ . Thus  $y \in W$ , and the proof is complete.

**Corollary 1.3.1.**— *Let  $A$  be an Artinian local ring with residue field  $k$ , and let  $G$  be a flat  $A$ -group. The following assertions are equivalent:*<sup>8</sup>

- (i)  *$G$  is locally quasi-finite (respectively, unramified, smooth, or étale) over  $A$  at some point;*
- (ii)  *$G$  is locally quasi-finite (respectively, unramified, smooth, or étale) over  $A$ .*

<sup>9</sup> *Proof.* If  $G$  satisfies one of the conditions in (i) at a point  $x$ , there is an open neighborhood  $U$  of  $x$  that is of finite type over  $A$ . Consequently, it is enough to apply 1.3 in the case where  $H$  is the unit  $A$ -group, taking the following lemma into account.

**Lemma 1.3.1.1.**— *Let  $A$  be an Artinian local ring and  $G$  an  $A$ -group. If  $G$  contains a nonempty open subset of finite type over  $A$ , then  $G$  is locally of finite type over  $A$ .*

*Proof.* By Lemma 1.3.0, we may suppose that  $A$  equals its residue field  $k$ . Moreover, by fpqc descent, we may suppose that  $k$  is algebraically closed (cf. EGA IV<sub>2</sub>, 2.7.1). Let  $V$  be the open subset of  $G$  consisting of the points at which  $G$  is of finite type over  $k$ ; by hypothesis,  $V \neq \emptyset$ . Since  $G$  is a Jacobson scheme,  $V$  contains a closed point  $x$ , and to prove that  $V = G$  it is enough to show that every closed point  $y$  of  $G$  belongs to  $V$ . For such a point  $y$ , the translation

$$r_y \circ r_x^{-1}$$

sends  $x$  to  $y$ , and hence  $y \in V$ .

**Corollary 1.3.2.**— *Let  $A$  be an Artinian local ring, and let  $u : G \rightarrow H$  be a morphism between  $A$ -groups locally of finite type. The following assertions are equivalent:*

<sup>7</sup>Editor's note: the original stated that the flat locus  $W$  is open, which does not seem evident a priori.

<sup>8</sup>Editor's note: it is not enough to suppose that  $G$  is flat over  $A$  at one point. For example, suppose that  $A \neq k$ , let  $H$  be the constant  $A$ -group  $(\mathbb{Z}/2\mathbb{Z})_A$ , and let  $G$  be the closed  $A$ -subgroup of  $H$  whose nonidentity component is reduced. Then the structural morphism  $G \rightarrow \operatorname{Spec} A$  is a local isomorphism at the identity point  $e$ , but is not flat at the point  $g \neq e$ .

<sup>9</sup>Editor's note: we have added the sentence that follows and simplified the proof of Lemma 1.3.1.1 by invoking EGA IV, 2.7.1 instead of loc. cit., 17.7.5.

- (i)  $u$  is universally open;
- (ii)  $u$  is open;
- (iii)  $u$  is open at some point of  $G$ ;
- (iv) the morphism  $u^0 : G^0 \rightarrow H^0$  induced by  $u$  is dominant;
- (iv')  $u^0$  is surjective;
- (v) there is a connected component  $C$  of  $G$  such that, if  $D$  denotes the connected component of  $H$  containing  $u(C)$ , the induced morphism  $u' : C \rightarrow D$  is dominant.

*Proof.* The implications

$$(i) \implies (ii) \implies (iii) \quad \text{and} \quad (iv') \implies (iv) \implies (v)$$

are clear. Since  $G^0$  is of finite type over  $A$  (Exposé VI<sub>A</sub>, 2.4), and hence Noetherian,  $u^0$  is quasi-compact. Thus  $u^0(G^0)$  is closed in  $H^0$  by 1.2, and therefore (iv) implies (iv'). On the other hand,  $G^0$  (respectively,  $C$ ) is open in  $G$  (Exposé VI<sub>A</sub>, 2.3), while  $H^0$  (respectively,  $D$ ) is irreducible (Exposé VI<sub>A</sub>, 2.4.1). Hence (ii) implies (iv) (respectively, (iii) implies (v)). It remains to prove that (v) implies (i).

We endow the open subset  $C$  (respectively,  $D$ ) of  $G$  (respectively,  $H$ ) with its induced scheme structure, and let  $u'$  denote the induced morphism  $C \rightarrow D$ . Let  $k$  be the residue field of  $A$ .<sup>10</sup> Since the base change

$$\mathrm{Spec} k \longrightarrow \mathrm{Spec} A$$

is a universal homeomorphism, we may suppose that  $A = k$ . By hypothesis  $u'$  is dominant; and since  $C$  is of finite type over  $k$  (Exposé VI<sub>A</sub>, 2.4.1), and hence Noetherian,  $u'$  is quasi-compact. By EGA IV<sub>2</sub>, 2.3.7,

$$u' \otimes_k \bar{k}$$

is again quasi-compact and dominant, where  $\bar{k}$  denotes an algebraic closure of  $k$ . Since  $C \otimes_k \bar{k}$  is a union of connected components of  $G \otimes_k \bar{k}$ , the morphism

$$u \otimes_k \bar{k} : G \otimes_k \bar{k} \longrightarrow H \otimes_k \bar{k}$$

satisfies assertion (v). In view of EGA IV<sub>2</sub>, 2.6.4, we are thus reduced to the case where  $A = k$  is an algebraically closed field.

In this case we may further replace  $u$  by  $u_{\mathrm{red}}$ , and are thereby reduced to the case where  $H$  is reduced. Let  $\xi$  (respectively,  $\eta$ ) be the generic point of  $C$  (respectively,  $D$ ). Since  $u'$  is quasi-compact and dominant,

$$u'(\xi) = \eta$$

(cf. EGA IV<sub>1</sub>, 1.1.5). Since  $H$  is reduced, the local ring  $\mathcal{O}_{H,\eta}$  is a field, and hence  $u'$  is flat at  $\xi$ .<sup>11</sup> It follows from 1.3 that  $u$  is flat. Moreover, since  $u$  is locally of finite type and  $H$  is locally Noetherian,  $u$  is locally of finite presentation. Therefore  $u$  is universally open by EGA IV<sub>2</sub>, 2.4.6.

**Proposition 1.4.**— *Let  $A$  be an Artinian local ring, and let  $u : G \rightarrow H$  be a quasi-compact morphism between  $A$ -groups locally of finite type. The following assertions are equivalent:*

<sup>10</sup>Editor's note: we have expanded the original in what follows.

<sup>11</sup>Editor's note: the original invoked the generic flatness theorem (EGA IV<sub>2</sub>, 6.9.1), which is not necessary here.

- (i)  $u$  is proper;
- (ii) there exists  $h \in H$  such that the fiber  $u^{-1}(h)$  is nonempty and proper over  $\kappa(h)$ ;
- (iii)  $\ker u$  is proper over  $A$ .

*Proof.* It is clear that (i) implies (iii), and that (iii) implies (ii). The hypotheses imply that  $u$  is of finite type; and since  $G$  is separated (Exposé VI<sub>A</sub>, 0.3),  $u$  is separated as well (EGA I, 5.5.1). It remains to prove that (ii) implies that  $u$  is universally closed. We may therefore suppose that  $A$  equals its residue field  $k$ .<sup>12</sup> Let  $k'$  be an algebraic closure of  $\kappa(h)$ , let

$$u' : G' \longrightarrow H'$$

be obtained from  $u$  by base change, and let  $h' \in H'$  be a point above  $h$ . Then

$$u'^{-1}(h') = u^{-1}(h) \times_{\kappa(h)} k'$$

is nonempty and proper, and it is enough to prove that  $u'$  is proper (EGA IV<sub>2</sub>, 2.6.4). Thus we may suppose that  $k$  is algebraically closed and  $h \in H(k)$ .

By 1.2,  $u(G)$  is then the underlying set of a closed reduced group subscheme of  $H$ . Since every closed immersion is proper (EGA II, 5.4.2), we may suppose that  $u$  is surjective and that  $H$  is reduced. The group  $G(k)$  then acts transitively on the set of closed points of  $H$ . Consequently, for every closed point  $y \in H$ , the fiber  $u^{-1}(y)$  is proper over  $\kappa(y)$ . By EGA IV<sub>3</sub>, 9.6.1, the set of  $y \in H$  for which  $u^{-1}(y)$  is not proper over  $\kappa(y)$  is locally constructible. It contains no closed point, and is therefore empty (cf. EGA IV<sub>3</sub>, 10.3.1 and 10.4.7).

<sup>13</sup> Now consider the generic point  $\eta$  of  $H^0$ . By what precedes, the fiber

$$u^{-1}(\eta) = G \times_H \operatorname{Spec} \kappa(\eta)$$

is proper over  $\kappa(\eta)$ . Since  $H$  is reduced,  $\kappa(\eta) = \mathcal{O}_{H,\eta}$ . The latter ring is the direct limit of the rings  $\mathcal{O}_H(V)$ , as  $V$  ranges over the nonempty open subsets of  $H^0$ . It follows from EGA IV<sub>3</sub>, 8.1.2(a) and 8.10.5(xii), that there exists a nonempty open subset  $V \subset H^0$  such that the restriction of  $u$  over  $V$  is proper. The translates  $g \cdot V$ , for  $g \in G(k)$ , form an open covering of  $H$ ; and for every  $g \in G(k)$ , the restriction of  $u$  over  $g \cdot V$  is proper. Hence  $u$  is proper (cf. EGA II, 5.4.1).

**Corollary 1.4.1.**— *Let  $A$  be an Artinian local ring, and let  $u : G \rightarrow H$  be a morphism between  $A$ -groups locally of finite type. The following assertions are equivalent:*

- (i)  $u$  is locally quasi-finite;
- (ii)  $u$  is quasi-finite at some point;
- (iii)  $\ker u$  is discrete;
- (iv) the restriction of  $u$  to each connected component of  $G$  is finite.

*If  $u$  is quasi-compact, these assertions are also equivalent to*

- (v)  $u$  is finite.

<sup>12</sup>Editor's note: we have added the following sentence.

<sup>13</sup>Editor's note: we have expanded the original in what follows.

It is clear that (iv) implies (iii), that (iii) implies (ii) (EGA I, 6.4.4), and that, when  $u$  is quasi-compact, (iv) and (v) are equivalent. We have already seen in 1.3 that (i) and (ii) are equivalent.

It remains to prove that (i) implies (iv). Let  $C$  be a connected component of  $G$ . Since  $C$  is of finite type over  $A$  (Exposé VI<sub>A</sub>, 2.4.1) and  $G, H$  are separated (Exposé VI<sub>A</sub>, 0.3),\* EGA I, 5.5.1 and 6.3.4 show that the restriction  $u'$  of  $u$  to  $C$  is separated and of finite type.<sup>14</sup> Its fibers are discrete, so  $u'$  is quasi-finite (cf. EGA II, 6.2.2). Since every quasi-finite proper morphism is finite (cf. EGA III<sub>1</sub>, 4.4.2), it is enough to prove that  $u'$  is universally closed.

For this, we may suppose that  $A$  equals its residue field  $k$ . By fpqc descent (cf. EGA IV<sub>2</sub>, 2.6.4), it is enough to prove that

$$u' \otimes_k \bar{k}$$

is universally closed, where  $\bar{k}$  is an algebraic closure of  $k$ . Since  $C$  is of finite type over  $k$ , the scheme  $C \otimes_k \bar{k}$  is the disjoint union of finitely many connected components  $C'_1, \dots, C'_n$  of  $G \otimes_k \bar{k}$ , and it is enough to prove the assertion for each  $C'_i$ . We are thus reduced to the case in which  $k$  is algebraically closed.

Let  $g$  be a closed point of  $C$ , and let

$$u^0 : G^0 \longrightarrow H$$

be the restriction of  $u$  to  $G^0$ . Then

$$u' = r_{u(g)} \circ u^0 \circ r_{g^{-1}},$$

where  $r_g$  denotes right translation by  $g$ . Thus, to prove that  $u'$  is proper, it is enough to prove that  $u^0$  is proper. By hypothesis  $u$  is locally quasi-finite, so the nonempty fiber  $\ker u$  is discrete. We have seen that  $u^0$  is of finite type, hence its fiber  $\ker u^0$  is finite (cf. EGA II, 6.2.2), and is therefore nonempty and proper. Proposition 1.4 now shows that  $u^0$  is proper.

**Corollary 1.4.2.**— *Let  $A$  be an Artinian local ring, and let  $u : G \rightarrow H$  be a quasi-compact morphism between  $A$ -groups locally of finite type. The following assertions are equivalent:*<sup>15</sup>

- (i)  $u$  is a closed immersion;
- (ii)  $u$  is a monomorphism;
- (iii)  $\ker u$  is trivial, that is, isomorphic to the unit  $A$ -group.

*In particular, every group subscheme<sup>16</sup> of  $H$  is closed.*

\* It is clear that (i) implies (ii). Considering the functors represented by  $G$  and  $H$ , one sees immediately that (ii) and (iii) are equivalent. Finally, if  $\ker u$  is the unit  $A$ -group, it is a proper nonempty fiber. Proposition 1.4 therefore shows that  $u$  is a proper monomorphism;

\*Translator's note: the French source cites Exposé VI<sub>A</sub>, 0.2 here; the separation statement is 0.3, as also recalled in Editor's note 2 above.

<sup>14</sup>Editor's note: we have expanded the original in what follows.

<sup>15</sup>Editor's note: the hypothesis that  $G$  and  $H$  be locally of finite type may be removed, because [Per76], 4.2.4, states that every quasi-compact monomorphism  $u : G \rightarrow H$  between group schemes over a field  $k$  is a closed immersion. See also [DG70], III.3.7.2(b), for the case in which  $G$  and  $H$  are affine; in that case every morphism  $G \rightarrow H$  is affine (EGA II, 1.6.2(v)), hence quasi-compact.

<sup>16</sup>Editor's note: in this particular case, see also Exposé VI<sub>A</sub>, 0.5.2, which holds without any finiteness hypothesis.

\*Translator's note: the French source calls the trivial kernel the “unit  $k$ -group” both in the statement and in the proof. Here the base is the Artinian local ring  $A$ , and  $\ker u$  is an  $A$ -group; the unit object required by the monomorphism criterion is therefore  $\text{Spec } A$ , the unit  $A$ -group.

it is of finite presentation because  $H$  is locally Noetherian (EGA IV<sub>1</sub>, 1.6.1), and hence is a closed immersion (EGA IV<sub>3</sub>, 8.11.5).

The final assertion follows because  $H$  is locally Noetherian, so every immersion  $G \rightarrow H$  is quasi-compact (EGA I, 6.6.4).

**Counterexample 1.4.3.**— *Let  $k$  be a field of characteristic zero, let  $G$  be the constant  $k$ -group  $\mathbb{Z}$ , and let  $H$  be the additive  $k$ -group  $\mathbb{G}_{a,k}$ . Let  $u : G \rightarrow H$  be a morphism of  $k$ -groups. If  $u \neq 0$ , then  $\ker u = 0$ , but  $u$  is not a closed immersion (cf. 1.2.2).*

We shall use below the following two results, which should have appeared in Exposé VI<sub>A</sub>.

**Lemma 1.5.**— *Let  $k$  be a field and let  $G$  be a  $k$ -group locally of finite type. Then:*

- (i) *all the irreducible components of  $G$  have the same dimension;*
- (ii) *for every  $g \in G$ ,*

$$\dim_g G = \dim G.$$

<sup>17</sup> Assertion (i) has already been proved in Exposé VI<sub>A</sub>, 2.4.1, and (ii) follows. Indeed, let  $g \in G$ , and let  $C$  be the connected component of  $G$  containing  $g$ . By definition (EGA 0<sub>IV</sub>, 14.1.2),  $\dim_g G$  is the infimum of the integers  $\dim U$ , where  $U$  ranges over the open neighborhoods of  $g$ . Hence

$$\dim_g G = \dim U_0$$

for some  $U_0$ , which we may suppose is contained in  $C$  (because  $\dim V \leq \dim U$  when  $V \subset U$ ). Since  $C$  is irreducible and of finite type over  $k$  (Exposé VI<sub>A</sub>, 2.4.1),

$$\dim U_0 = \dim C = \operatorname{trdeg}_k \kappa(\xi),$$

where  $\xi$  is the generic point of  $C$  (EGA IV<sub>2</sub>, 5.2.1). Therefore

$$\dim_g G = \dim C = \dim G.$$

**Proposition 1.6.**—<sup>18</sup> *Let  $S$  be a scheme of characteristic zero, and let  $G$  be an  $S$ -group scheme locally of finite presentation over  $S$  at every point of the identity section  $\epsilon(S)$ . The scheme  $G$  is smooth over  $S$  at every point of the identity section if and only if the  $\mathcal{O}_S$ -module*

$$\omega_{G/S} = \epsilon^*(\Omega_{G/S}^1),$$

*called the conormal module of the identity section of  $G$ , is locally free.*

Recall that a scheme  $S$  is said to be of characteristic zero if the field  $\kappa(x)$  has characteristic zero for every closed point  $x \in S$ .

Recall also (Exposé II, 4.11) that, if  $\pi : G \rightarrow S$  is the structural morphism, then

$$\Omega_{G/S}^1 = \pi^*(\omega_{G/S}).$$

Thus it is equivalent to say that the  $\mathcal{O}_S$ -module  $\omega_{G/S}$  is locally free or that the  $\mathcal{O}_G$ -module  $\Omega_{G/S}^1$  is locally free.

<sup>17</sup>Editor's note: the implication (ii)  $\Rightarrow$  (i) is a general fact (cf. EGA 0<sub>IV</sub>, 14.1.6), while (i)  $\Rightarrow$  (ii) follows from the fact that if  $X$  is an irreducible scheme of finite type over a field, then  $\dim X = \dim U$  for every nonempty open subset  $U$  of  $X$ . The essential point here is therefore assertion (i), which was already established in an addition to Exposé VI<sub>A</sub>, 2.4.1. We have modified the statement and proof accordingly.

<sup>18</sup>Editor's note: in the statement, we have replaced "along the identity section" by "at every point of the identity section." At the end of the proof, we have also made explicit the cited results from EGA IV<sub>4</sub>.

If there is an open neighborhood  $U$  of  $\epsilon(S)$  that is smooth over  $S$ , then by EGA IV<sub>4</sub>, 17.2.3,  $\Omega_{U/S}^1$  is locally free of finite type, and so is

$$\omega_{G/S} = \epsilon^*(\Omega_{U/S}^1).$$

Conversely, if  $\omega_{G/S}$  is locally free, then so is

$$\Omega_{G/S}^1 = \pi^*(\omega_{G/S}).$$

Since  $S$  has characteristic zero, the Jacobian criterion (EGA IV<sub>4</sub>, 16.12.2) shows that  $G$  is differentially smooth over  $S$ . It follows from EGA IV<sub>4</sub>, 17.12.5, that  $G$  is smooth over  $S$  at every point of the identity section.

**Corollary 1.6.1 (Cartier).**— *Let  $k$  be a field of characteristic zero. Every  $k$ -group locally of finite type over  $k$  is smooth over  $k$ .*

Indeed, the  $k$ -module  $\omega_{G/k}$  is then plainly free. By 1.6,  $G$  is smooth over  $k$  at the identity point  $e$ , and hence smooth over  $k$  by 1.3.1.

## 2. “Open properties” of groups and group morphisms locally of finite presentation

**2.0.**— In all that follows,  $S$  will denote an arbitrary scheme; an  $S$ -group scheme will be called an  $S$ -group. Given an  $S$ -group  $G$ , we shall denote by  $\epsilon$  the identity section, by

$$c : G \longrightarrow G$$

the inversion morphism, and by

$$\mu : G \times_S G \longrightarrow G$$

the multiplication morphism. For every  $S$ -scheme  $X$ , we shall denote by  $\pi$ , or by  $\pi_X$ , the structural morphism

$$X \longrightarrow S.$$

Given a property  $\mathcal{P}(u)$  of a morphism of  $S$ -schemes

$$u : X \longrightarrow Y,$$

we shall say that  $\mathcal{P}(u)$  is *stable under base change* if, whenever  $u$  satisfies  $\mathcal{P}(u)$ , the same is true of the morphism  $u_{(Y')}$ , for every  $S$ -morphism  $Y' \rightarrow Y$ . We say that  $\mathcal{P}(u)$  is *local in nature for the topology  $\mathcal{T}$*  (cf. Exposé IV, §§ 4 and 6) if  $\mathcal{P}(u)$  satisfies the following two conditions:

- a)  $\mathcal{P}(u)$  is stable under base change;
- b) whenever there is a family of  $S$ -morphisms  $Y_i \rightarrow Y$ , covering for the topology  $\mathcal{T}$ , such that each morphism  $u_{(Y_i)}$  satisfies  $\mathcal{P}(u)$ , then  $u$  satisfies  $\mathcal{P}(u)$ .

**Proposition 2.1.**— *Let  $\mathcal{P}(u)$  be a property of a morphism of  $S$ -schemes, local in nature for the fpqc topology. Let  $u : G \rightarrow H$  be a morphism of  $S$ -groups. Suppose that  $G$  is flat and universally open over  $S$ .*

*Let  $W$  be the largest open subset of  $H$  over which  $u$  satisfies the property  $\mathcal{P}(u)$ , and let  $V = u^{-1}(W)$ . Then*

$$U = \pi_G(V)$$



is open in  $S$ , and  $V$  is an open group subscheme of  $G|_U$ .

The existence of a largest open subset  $W$  of  $H$  over which  $u$  satisfies  $\mathcal{P}(u)$  follows from the fact that  $\mathcal{P}(u)$  is local in nature for the Zariski topology. Since  $\pi_G$  is universally open,  $\pi_G(V)$  is an open subset  $U$  of  $S$ . It is enough to show that  $V$  is a group subscheme of  $G|_U$ . We may therefore suppose that  $U = S$ .

Set

$$G' = G \times_S V, \quad H' = H \times_S V, \quad V' = V \times_S V, \quad W' = W \times_S V,$$

and  $u' = u|_{V'}$ . Let  $W'_1$  be the largest open subset of  $H'$  over which  $u'$  satisfies  $\mathcal{P}(u)$ . Since  $V$  is flat and universally open over  $S$ , the same is true of  $H'$  over  $H$ , and Lemma 2.1.1 below shows that

$$W'_1 = W'.$$

Now consider the automorphism  $a$  (respectively,  $b$ ) of the  $V$ -scheme  $G'$  (respectively,  $H'$ ), namely right translation by the inverse of the diagonal section  $\delta$  (respectively, by the inverse of  $u(\delta)$ ), defined by

$$a(g, v) = (g \cdot v^{-1}, v), \quad (\text{respectively, } b(h, v) = (h \cdot u(v^{-1}), v)),$$

for every morphism  $T \rightarrow S$ , every  $g \in G(T)$ ,  $v \in V(T)$ , and  $h \in H(T)$ .<sup>\*</sup> It is clear that

$$u' \circ a = b \circ u',$$

which shows that  $W'$  is stable under  $b$ , hence that  $V'$  is stable under  $a$ , so that  $V$  is a group subscheme of  $G$ .

**Lemma 2.1.1.**— *Let  $\mathcal{P}(u)$  be a property of an  $S$ -morphism  $u$ , local in nature for the fpqc topology. Consider a cartesian square of morphisms of  $S$ -schemes:*

$$\begin{array}{ccc} X' & \xrightarrow{f'} & Y' \\ \downarrow & & \downarrow g \\ X & \xrightarrow{f} & Y \end{array}$$

where  $g$  is flat and open. Let  $W$  (respectively,  $W'_1$ ) be the largest open subset of  $Y$  (respectively,  $Y'$ ) over which  $f$  (respectively,  $f' = f|_{Y'}$ ) satisfies  $\mathcal{P}(u)$ . Then

$$W'_1 = W \times_Y Y'.$$

Set  $W' = W \times_Y Y'$ . Since  $\mathcal{P}(u)$  is stable under base change, we have

$$W' \subset W'_1.$$

Since  $g$  is open,

$$W_1 = g(W'_1)$$

is an open subset of  $Y$ . Set

$$V_1 = f^{-1}(W_1), \quad V'_1 = V_1 \times_{W_1} W'_1;$$

---

<sup>\*</sup>Translator's note: the French source omits  $b$  before  $(h, v)$  in the second displayed definition. The preceding sentence defines the automorphism  $b$ , and the identity immediately below requires the displayed map to be  $b(h, v)$ ; the symbol is supplied here.

it is clear that  $V'_1 = f'^{-1}(W'_1)$ . Since  $g$  is flat and open, the morphism

$$W'_1 \longrightarrow W_1 = g(W'_1)$$

induced by  $g$  is faithfully flat and open, hence is a covering for the fpqc topology (cf. Exposé IV, 6.3.1(iv)). Since the morphism

$$V'_1 \longrightarrow W'_1$$

induced by  $f'$  satisfies  $\mathcal{P}(u)$ , the same is true of the morphism  $V_1 \rightarrow W_1$  induced by  $f$ . Thus

$$W_1 \subset W, \quad W'_1 \subset g^{-1}(W_1) \subset g^{-1}(W) = W',$$

and therefore  $W' = W'_1$ .

**Remark 2.1.2.**— *A great many properties of a morphism are local in nature for the fpqc topology; among them are the properties of being flat, (universally) open, (locally) of finite type, of finite presentation, quasi-finite (cf. EGA IV<sub>2</sub>, 2.5.1, 2.6.1, and 2.7.1), smooth, étale, and unramified (EGA IV<sub>4</sub>, 17.7.3).*

*The proof of 2.1 in fact uses only base changes by flat morphisms. The proposition therefore applies to a property satisfying condition b) of 2.0 relative to the fpqc topology and stable under base change by flat morphisms (for example, the property of being quasi-compact and dominant).*

*Of course, one can state an analogous proposition concerning properties local in nature for a topology  $\mathcal{T}$  finer than the Zariski topology, the condition to be verified by  $G$  then being that  $\pi_G$  be universally open and covering for the topology  $\mathcal{T}$ .*

*In particular, if  $G$  is flat and locally of finite presentation over  $S$ , there is an analogous statement for properties stable under base change by flat morphisms locally of finite presentation and satisfying condition b) of 2.0 relative to the fpqc topology (for example, the properties of being regular, reduced, Cohen–Macaulay, and so on; cf. EGA IV<sub>2</sub>, 6.8).*

**Proposition 2.2.**— *Let  $G$  and  $H$  be two  $S$ -groups, and let  $u : G \rightarrow H$  be a morphism of  $S$ -groups. Then.*<sup>19</sup>

- (i) *Suppose that  $G$  or  $H$  is flat over  $S$ , and that  $G$  or  $H$  is locally of finite presentation over  $S$ . Let  $V$  be the largest open subset of  $G$  such that the restriction of  $u$  to  $V$  is flat and locally of finite presentation (respectively, smooth, or étale). Then  $U = \pi_V(V)$  is open in  $S$ , and  $V$  is an open group subscheme of  $G|_U$ .*
- (ii) *Suppose that  $G$  or  $H$  is universally open over  $S$ , and let  $V$  be the largest open subset of  $G$  such that the restriction of  $u$  to  $V$  is universally open. Then  $U = \pi_V(V)$  is open in  $S$ , and  $V$  is an open group subscheme of  $G|_U$ .*

We first prove (i). We show that the restriction  $\pi_V$  of  $\pi_G$  to  $V$  is flat and locally of finite presentation:

- a) if  $\pi_G$  is flat (respectively, locally of finite presentation), then so is  $\pi_V$ ;
- b) if  $\pi_H$  is flat (respectively, locally of finite presentation), then so is  $\pi_V$ , as the composite of the restriction of  $u$  to  $V$  with  $\pi_H$ .

<sup>19</sup>Editor's note: in case (i), the open subset  $V$  consists of all the points of  $G$  at which  $u$  is smooth, respectively étale, respectively of finite presentation and flat; cf. editor's note 6. By contrast, in case (ii),  $V$  is the largest open subset contained in the set  $E$  of points of  $G$  at which  $u$  is universally open, but  $E$  is not necessarily open, as the following example shows (EGA IV<sub>3</sub>, 14.1.3(i)). Let  $k$  be a field, let  $H = S = \operatorname{Spec} A$ , where  $A = k[x]$ , and let  $G$  be the  $S$ -group  $\operatorname{Spec} A[y]/(xy)$ . Then  $E$  is the identity section of  $G$ , which is not open.

Thus, in the four cases contemplated in the statement,  $\pi_V$  is flat and locally of finite presentation, hence universally open (EGA IV<sub>2</sub>, 2.4.6). Set  $U = \pi_V(V)$ ; then  $U$  is open in  $S$ . It is therefore enough to show that  $V$  is an open group subscheme of  $G|_U$ , and we may suppose that  $U = S$ .

Set

$$G' = G \times_S V, \quad H' = H \times_S V, \quad V' = V \times_S V, \quad \text{and} \quad u' = u_{(V)}.$$

Since  $V$  is flat and locally of finite presentation over  $S$ , the same is true of  $H'$  over  $H$ . By EGA IV<sub>4</sub>, 17.7.4,  $V'$  is then the largest open subset of  $G'$  such that the restriction of  $u'$  to  $V'$  is flat and locally of finite presentation (respectively, smooth, or étale). With the automorphisms  $a$  and  $b$  defined as in the proof of 2.1, it is clear that  $V'$  is stable under  $a$ , hence that  $V$  is a group subscheme of  $G$ .

We now prove (ii). The restriction  $\pi_V$  of  $\pi_G$  to  $V$  is universally open, either because  $\pi_G$  is universally open, or because it is the composite of the restriction of  $u$  to  $V$  with  $\pi_H$ , when  $\pi_H$  is universally open. Set  $U = \pi_V(V)$ ; then  $U$  is open in  $S$ . It is enough to show that  $V$  is an open group subscheme of  $G|_U$ , and we may suppose that  $U = S$ .

Set, as before,

$$G' = G \times_S V, \quad H' = H \times_S V, \quad V' = V \times_S V, \quad \text{and} \quad u' = u_{(V)}.$$

Then  $\pi_V : V \rightarrow S$  is surjective and universally open; the same is therefore true of  $G' \rightarrow G$ . Consequently,  $V'$  is the largest open subset of  $G'$  such that the restriction of  $u'$  to  $V'$  is universally open, by EGA IV<sub>3</sub>, 14.3.4(i),(ii). With the automorphisms  $a$  and  $b$  defined as before, it is clear that  $V'$  is stable under  $a$ , hence that  $V$  is a group subscheme of  $G$ .

**Corollary 2.3.**— *Let  $G$  be an  $S$ -group, and let  $V$  be the largest open subset of  $G$  that is flat and locally of finite presentation (respectively, smooth, étale, or universally open) over  $S$ . Then  $U = \pi_V(V)$  is open in  $S$ , and  $V$  is an open group subscheme of  $G|_U$ .*

It is enough to apply 2.2 when  $H$  is the identity  $S$ -group and  $u$  is the unique morphism of  $S$ -groups  $G \rightarrow H$ , since  $\pi_H$  is then an isomorphism and

$$\pi_G = \pi_H \circ u.$$

**Corollary 2.4.**— *Let  $G$  be an  $S$ -group. If there is a neighborhood  $X$  of the identity section having one (or more) of the following properties:*

*$X$  is flat and locally of finite presentation (respectively, smooth, étale, or universally open) over  $S$ ,*

*then there is an open group subscheme of  $G$  having the same properties.*

It is enough to apply 2.3, observing that here, with the notation of 2.2, we have

$$\epsilon(S) \subset V,$$

and hence  $U = S$ .

**Proposition 2.5.**<sup>20</sup> *Let  $u : G \rightarrow H$  be a morphism of  $S$ -groups.*

<sup>20</sup>Editor's note: we have expanded the statement and proof, and in (i) we have weakened the hypothesis on  $H$ , replacing "of finite presentation" by "of finite type."

- (i) Suppose that  $G$  (respectively,  $H$ ) is of finite presentation and flat (respectively, of finite type) over  $S$  at the points of its identity section. Then the sets

$$U_{\text{plat}} \supset U_{\text{lis}} \supset U_{\text{ét.}},$$

formed by the points  $s \in S$  such that  $u_s$  is flat (respectively, smooth, or étale), are open in  $S$ .

If, moreover,  $G$  (respectively,  $H$ ) is locally of finite presentation and flat (respectively, locally of finite type) over  $S$ , then the set  $V_{\text{plat}}$  (respectively,  $V_{\text{lis}}$  or  $V_{\text{ét.}}$ ) of points of  $G$  at which  $u$  is flat (respectively, smooth, or étale) is the inverse image under  $\pi_G$  of  $U_{\text{plat}}$  (respectively,  $U_{\text{lis}}$  or  $U_{\text{ét.}}$ ).

- (ii) Suppose that, for every  $s \in S$ ,  $G_s$  is locally of finite type over  $\kappa(s)$  (a condition satisfied if  $G$  is of finite type over  $S$  at the points of the identity section (1.3.1.1)), and that  $u$  is locally of finite type (respectively, locally of finite presentation) at the points of the identity section of  $G$ . Then the sets

$$U_{\text{l.q.f.}} \supset U_{\text{n.r.}},$$

formed by the  $s \in S$  such that  $u_s$  is locally quasi-finite (respectively, unramified), are open in  $S$ .

If, moreover,  $u$  is locally of finite type (respectively, locally of finite presentation), then the set  $V_{\text{l.q.f.}}$  (respectively,  $V_{\text{n.r.}}$ ) of points of  $G$  at which  $u$  is quasi-finite (respectively, unramified) is the inverse image under  $\pi_G$  of  $U_{\text{l.q.f.}}$  (respectively,  $U_{\text{n.r.}}$ ).

We prove (i). First note (1.3.1.1) that, for every  $s \in S$ ,  $G_s$  is locally of finite type over  $\kappa(s)$ . Let  $Y$  be the open subset of  $H$  consisting of the points at which  $\pi_H$  is of finite type, and let  $X_1$  be the open subset of  $u^{-1}(Y)$  consisting of the points at which  $\pi_G$  is of finite presentation. By EGA IV<sub>3</sub>, 11.3.1, the set  $X$  of points of  $X_1$  at which  $\pi_G$  is of finite presentation and flat is open in  $X_1$ , hence in  $G$ .

Let

$$u_X : X \longrightarrow Y$$

be the morphism induced by  $u$ . Since  $Y$  (respectively,  $X$ ) contains the identity section of  $H$  (respectively,  $G$ ), the restriction  $\pi_X$  of  $\pi_G$  to  $X$  is surjective, and the morphism

$$S \longrightarrow X$$

induced by  $\epsilon_G$  is an  $S$ -section of  $X$ .

Consider the sets

$$W_{\text{plat}} \supset W_{\text{lis}} \supset W_{\text{ét.}},$$

consisting of the points of  $X$  at which  $u_X$  is flat (respectively, smooth, or étale). It is clear that  $W_{\text{lis}}$  and  $W_{\text{ét.}}$  are open in  $X$ ; cf. Editor's note 6. On the other hand, since  $\pi_Y$  is locally of finite type and

$$\pi_X = \pi_Y \circ u_X$$

is locally of finite presentation, EGA IV<sub>1</sub>, 1.4.3(v), shows that  $u_X$  is locally of finite presentation. Consequently, the flatness locus  $W_X$  of  $u_X$  is open in  $X$  (EGA IV<sub>3</sub>, 11.3.1).

Let  $x \in X$  and set  $s = \pi_X(x)$ . Then, by EGA IV<sub>2</sub>, 11.3.10,\* combined with EGA IV<sub>4</sub>, 17.5.1 (respectively, 17.6.1),  $x$  belongs to  $W_{\text{plat}}$  (respectively,  $W_{\text{lis}}$  or  $W_{\text{ét.}}$ ) if and only if

---

\*Translator's note: the source citation is retained. Section 11 occurs in EGA IV<sub>3</sub>, not EGA IV<sub>2</sub>; the cited result is EGA IV<sub>3</sub>, 11.3.10.

$u_s$  is flat (respectively, smooth, or étale) at the point  $x$ , or, equivalently by 1.3, if and only if  $u_s$  is flat (respectively, smooth, or étale).

Consequently, for

$$“b = \text{flat, smooth, or étale},”$$

we have

$$U_b = \epsilon_G^{-1}(W_b), \quad W_b = \pi_X^{-1}(U_b).$$

Since  $W_b$  is open in  $X$ , hence in  $G$ , the first equality shows that  $U_b$  is open in  $S$ .

The second assertion of (i) follows from what we have just seen, for then

$$Y = H, \quad X = G, \quad V_b = W_b = \pi_G^{-1}(U_b).$$

We prove assertion (ii). Let

$$V_{\text{l.t.f.}} \supset V_{\text{l.q.f.}}$$

be the open subsets of  $G$  consisting of the points at which  $u$  is of finite type (respectively, quasi-finite). Set

$$X = V_{\text{l.t.f.}},$$

and let  $u_X$  denote the restriction of  $u$  to  $X$ . By hypothesis,  $X$  contains  $\epsilon(S)$ . Let  $x \in X$  and set  $s = \pi_X(x)$ .

If  $u$  is quasi-finite at  $x$ , then, by base change,  $u_s$  is quasi-finite at  $x$ ; and since  $G_s$  is assumed to be locally of finite type,  $u_s$  is locally quasi-finite by 1.3.1.

Conversely, if  $u_s$  is locally quasi-finite, then

$$u_X^{-1}(u_X(x)) \subset u_s^{-1}(u_s(x))$$

is a finite set. Since

$$u_X : X \longrightarrow H$$

is locally of finite type, Chevalley's semicontinuity theorem (EGA IV<sub>3</sub>, 13.1.3) shows that there is an open neighborhood  $W$  of  $x$  in  $X$  such that the fiber

$$u_X^{-1}(u_X(x'))$$

is discrete for every  $x' \in W$ . Hence  $u_X$  is quasi-finite at  $x$ , by EGA II, 6.2.2 (and EGA III<sub>2</sub>, Err<sub>III</sub> 20).

Consequently,

$$U_{\text{l.q.f.}} = \epsilon_G^{-1}(V_{\text{l.q.f.}}), \quad V_{\text{l.q.f.}} = \pi_X^{-1}(U_{\text{l.q.f.}}),$$

and the first equality shows that  $U_{\text{l.q.f.}}$  is open in  $S$ .

If, moreover,  $u$  is locally of finite type over  $S$ , then  $X = G$ , and hence  $V_{\text{l.q.f.}}$  is the inverse image under  $\pi_G$  of  $U_{\text{l.q.f.}}$ .<sup>\*</sup> This proves the first half of (ii).

Now consider the open subsets

$$V_{\text{l.p.f.}} \supset V_{\text{n.r.}},$$

consisting of the points at which  $u$  is of finite presentation (respectively, unramified), and suppose that

$$Z = V_{\text{l.p.f.}}$$

contains the identity section of  $G$ . Let  $z \in Z$ ; set

$$s = \pi_Z(z), \quad h = u_Z(z).$$

---

<sup>\*</sup>Translator's note: in this sentence and in the analogous sentence for  $V_{\text{n.r.}}$  below, the French source prints  $G$  in place of  $u$ . The statement assumes that  $u$  is locally of finite type (respectively, locally of finite presentation), and  $X$  and  $Z$  are precisely the corresponding loci for  $u$ ; the required  $u$  is used here.

If  $u$  is unramified at  $z$ , then, by base change,  $u_s$  is unramified at  $z$ ; and since  $G_s$  is assumed to be locally of finite type,  $u_s$  is unramified by 1.3.1.

Conversely, if  $u_s$  is unramified at the point  $z$ , then the fiber

$$u_s^{-1}(h) = u^{-1}(h)$$

is unramified over  $\kappa(h)$  at the point  $z$ . Since  $Z$  is an open subset of  $G$ , the fiber

$$u_Z^{-1}(h) = u_s^{-1}(h) \cap Z$$

is unramified over  $\kappa(h)$  at the point  $z$ . Since  $u_Z$  is locally of finite presentation, EGA IV<sub>4</sub>, 17.4.1, implies that  $u_Z$  is unramified at the point  $z$ .

Thus

$$U_{\text{n.r.}} = \epsilon_G^{-1}(V_{\text{n.r.}}), \quad V_{\text{n.r.}} = \pi_Z^{-1}(U_{\text{n.r.}}),$$

and the first equality shows that  $U_{\text{n.r.}}$  is open in  $S$ .

If, moreover,  $u$  is locally of finite presentation over  $S$ , then  $Z = G$ , and hence  $V_{\text{n.r.}}$  is the inverse image under  $\pi_G$  of  $U_{\text{n.r.}}$ . This completes the proof of assertion (ii).

**Corollary 2.6.**— *Let  $u : G \rightarrow H$  be a morphism of  $S$ -groups that is radicial (as is the case if  $u$  is a monomorphism; EGA I, 3.5.4). Suppose that  $G$  (respectively,  $H$ ) is locally of finite presentation and flat (respectively, locally of finite type) over  $S$ . Then the set  $U$  of  $s \in S$  such that  $u_s$  is an open immersion is open in  $S$ , and the restriction of  $u$  over  $U$  is an open immersion.*

By 2.5(i), the set  $U'$  of points  $s \in S$  such that  $u_s$  is étale is open in  $S$ . Since  $u$  is radicial, so is  $u_s$ , for every  $s \in S$ ; hence EGA IV<sub>4</sub>, 17.9.1, gives

$$U = U',$$

which shows that  $U$  is open. Finally, by 2.5(i), the restriction of  $u$  over  $U$  is étale; since  $u$  is radicial, this restriction is an open immersion (EGA IV<sub>4</sub>, 17.9.1).

**Proposition 2.7.**— *Let  $G$  be an  $S$ -group. The following conditions are equivalent:*

- (i)  $G$  is unramified over  $S$  at the points of the identity section;
- (ii) the identity section is an open immersion;
- (iii)  $G$  is of finite presentation over  $S$  at the points of the identity section, and, for every  $s \in S$ ,  $G_s$  is unramified over  $\kappa(s)$ .

*If, moreover,  $G$  is locally of finite presentation over  $S$ , then each of the three preceding conditions is equivalent to the following:*

- (iv)  $G$  is unramified over  $S$ .

The equivalence of assertions (i) and (ii) follows from the more general Lemma 2.7.1 below. Note (1.3.1.1) that either condition (i) or condition (iii) implies that, for every  $s \in S$ ,  $G_s$  is locally of finite type over  $\kappa(s)$ . Thus, by EGA IV<sub>4</sub>, 17.4.1, assertion (i) is equivalent to the assertion that, for every  $s \in S$ ,  $G_s$  is unramified over  $\kappa(s)$  at the point  $e_s$ , the identity of  $G_s$ , or, equivalently by 1.3.1, to the assertion that  $G_s$  is unramified over  $\kappa(s)$ . Consequently, assertions (i) and (iii) are equivalent. Finally, if  $G$  is locally of finite presentation over  $S$ , assertions (iii) and (iv) are equivalent (EGA IV<sub>4</sub>, 17.4.1).

**Lemma 2.7.1.**— *Let  $G$  be an  $S$ -scheme equipped with a section  $\epsilon$ . In order that  $G$  be unramified over  $S$  at the points of this section, it is necessary and sufficient that  $\epsilon$  be an open immersion.*

The condition is necessary by EGA IV<sub>4</sub>, 17.4.1, a)⇒b''). Conversely, if  $\epsilon$  is an open immersion, then the restriction to  $\epsilon(S)$  of the structural morphism

$$G \longrightarrow S$$

is an isomorphism; hence  $G$  is unramified over  $S$  at the points of  $\epsilon(S)$ .

**Corollary 2.8.**—<sup>21</sup> *Let  $u : G \rightarrow H$  be a morphism of  $S$ -groups. Suppose that, for every  $s \in S$ ,  $G_s$  is locally of finite type over  $\kappa(s)$ .*

a) *If  $u$  is locally of finite type, the following conditions are equivalent:*

- (i)  *$u$  is locally quasi-finite;*
- (ii) *for every  $s \in S$ , the morphism  $u_s : G_s \rightarrow H_s$  is locally quasi-finite;*
- (iii)  *$\text{Ker } u$  is locally quasi-finite over  $S$ ;*
- (iv) *the fibers of  $\text{Ker } u$  are discrete.*

b) *If  $u$  is locally of finite presentation, the following conditions are equivalent:*

- (v)  *$u$  is unramified;*
- (vi) *for every  $s \in S$ , the morphism  $u_s : G_s \rightarrow H_s$  is unramified;*
- (vii)  *$\text{Ker } u$  is unramified;*
- (viii) *the identity section  $S \rightarrow \text{Ker } u$  is an open immersion.*

The equivalences

$$(i) \iff (ii) \quad \text{and} \quad (v) \iff (vi)$$

follow from 2.5(ii), and Reminder 2.8.1 below shows that

$$(i) \implies (iii) \quad \text{and} \quad (v) \implies (vii).$$

For every  $s \in S$ , denote by  $e_s$  the identity element of  $H_s$ . Then (iii) (respectively, (vii)) implies that, for every  $s \in S$ ,

$$(\text{Ker } u)_s = \text{Ker } u_s = u_s^{-1}(e_s)$$

is locally quasi-finite (respectively, unramified) over  $\kappa(s) = \kappa(e_s)$ . It follows that  $u_s$  is quasi-finite (respectively, unramified) at the identity point of  $G_s$ , and hence, by 1.3, that  $u_s$  is locally quasi-finite (respectively, unramified). Thus

$$(iii) \implies (ii) \quad \text{and} \quad (vii) \implies (vi).$$

Finally,

$$(ii) \iff (iv) \quad \text{by 1.4.1,} \quad (vii) \iff (viii) \quad \text{by 2.7.}$$

**Reminder 2.8.1.**— Recall (cf. Exposé I, 2.3.6.1) that, given a morphism  $u : G \rightarrow H$  of  $S$ -groups, the *kernel of  $u$* , denoted  $\text{Ker } u$ , is the group subfunctor of  $G$  defined, for every morphism  $f : T \rightarrow S$ , by

$$(\text{Ker } u)(T) = \{a \in G(T) \mid u \circ a = \epsilon_H \circ f\}.$$

<sup>21</sup>Editor's note: we have changed the presentation by separating the assertions concerning the "quasi-finite" case from those concerning the "unramified" case.

By loc. cit. (or EGA I, 4.4.1), this functor is represented by the  $S$ -group

$$G \times_H S = u^{-1}(\epsilon_H(S)),$$

which is denoted simply by  $\text{Ker } u$ . In particular, the structural morphism

$$\text{Ker } u \longrightarrow S$$

is obtained from  $u$  by base change.

**Lemma 2.9.**— *Let  $\pi : X \rightarrow S$  be a morphism admitting an  $S$ -section  $\epsilon$ .*

- (i) *If  $\pi$  is injective, then it is integral.<sup>(\*)</sup>*
- (ii) *If  $\pi$  is locally of finite type and, for every  $s \in S$ ,  $\pi_s$  is an isomorphism, then  $\pi$  is an isomorphism.<sup>(\*\*)</sup>*

First observe that, by Lemma 2.9.1 below,  $\pi$ , being a homeomorphism, is an affine morphism.

If  $\pi$  is injective, then  $\epsilon$  is surjective. Since  $\epsilon$  is a surjective immersion,  $\epsilon(S)$  is isomorphic to the closed subscheme of  $X$  defined by a nil ideal  $\mathcal{I}$  of  $\mathcal{O}_X$ . Since  $\epsilon$  is an  $S$ -section of  $\pi$ , there is a direct-sum decomposition

$$\mathcal{O}_X = \mathcal{O}_S \oplus \mathcal{I}$$

of  $\mathcal{O}_S$ -modules. Since  $\mathcal{I}$  is a nil ideal of  $\mathcal{O}_X$ , it is plainly integral over  $\mathcal{O}_S$ . Thus  $\mathcal{O}_X$  is integral over  $\mathcal{O}_S$ , and  $\pi$  is integral.

Now suppose that  $\pi$  is locally of finite type. Since  $\pi \circ \epsilon = \text{id}_S$ , the section  $\epsilon$  is locally of finite presentation (cf. EGA IV<sub>1</sub>, 1.4.3(v)); hence  $\mathcal{I}$  is an ideal of finite type in  $\mathcal{O}_X$  (EGA IV<sub>1</sub>, 1.4.1). For every  $s \in S$ , one has

$$\mathcal{O}_{X_s} = \kappa(s) \oplus \mathcal{I} \otimes_{\mathcal{O}_S} \kappa(s).$$

By hypothesis,  $\epsilon_s$  is an isomorphism, so

$$\mathcal{I} \otimes_{\mathcal{O}_S} \kappa(s) = 0$$

for every  $s \in S$ , and therefore, a fortiori,

$$\mathcal{I} \otimes_{\mathcal{O}_X} \kappa(x) = 0$$

for every  $x \in X$ . Nakayama's lemma now gives  $\mathcal{I} = 0$ , and consequently  $\pi$  is an isomorphism.

**Lemma 2.9.1.**— *Let  $f : X \rightarrow Y$  be a morphism of schemes that is a homeomorphism. Then  $f$  is an affine morphism.<sup>(\*\*\*)</sup>*

It is enough to show that every point  $y \in Y$  has an open neighborhood  $W$  such that the restriction of  $f$  over  $W$  is affine. Let  $y \in Y$ , and let  $V$  be an affine open neighborhood of  $y$  in  $Y$ . Set

$$V' = f^{-1}(V).$$

Then  $V'$  is an open neighborhood of  $x = f^{-1}(y)$  in  $X$ . There is an affine open neighborhood  $W'$  of  $x$  in  $X$  contained in  $V'$ . Set

$$W = f(W').$$

<sup>(\*)</sup>This is also a special case of EGA IV<sub>4</sub>, 18.12.11, since  $\pi$  is plainly a universal homeomorphism.

<sup>(\*\*)</sup>This is also an immediate consequence of EGA IV<sub>4</sub>, 18.12.6.

<sup>(\*\*\*)</sup>Cf. EGA IV<sub>4</sub>, 18.12.7.1 for a slightly more general result, proved by the same argument.



Then  $W$  is an open neighborhood of  $y$  in  $Y$ , contained in the affine scheme  $V$ , and therefore  $W$  is separated. Since  $W'$  is affine, the restriction of  $f$  over  $W$  is an affine morphism (EGA II, 1.2.3).

**Corollary 2.10.**— *Let  $G$  be an  $S$ -group locally of finite type. Suppose that, for every  $s \in S$ ,  $G_s$  is the identity  $\kappa(s)$ -group. Then  $G$  is the identity  $S$ -group.*

More generally:

**Corollary 2.11.**— *Let  $f : X \rightarrow Y$  be an  $S$ -morphism locally of finite type. In order that  $f$  be a monomorphism, it is necessary and sufficient that  $f_s$  be a monomorphism for every  $s \in S$ .*

The condition is plainly necessary; we prove that it is sufficient. If  $f_s$  is a monomorphism for every  $s \in S$ , then, a fortiori,  $f_y$  is a monomorphism for every  $y \in Y$ . We may therefore assume that  $Y = S$ .

By EGA I, 5.3.8, to prove that  $f$  is a monomorphism it is enough to show that

$$\Delta_f : X \longrightarrow X \times_S X$$

is an isomorphism, or, equivalently, that the first projection

$$p : X \times_S X \longrightarrow X$$

is an isomorphism. If  $f_s$  is a monomorphism, EGA I, 5.3.8 likewise shows that the first projection

$$X_s \times_{\kappa(s)} X_s \longrightarrow X_s,$$

which is identified with  $p_s$ , is an isomorphism. But  $p$  has the  $S$ -section  $\Delta_f$ ; hence Lemma 2.9 shows that, if  $p_s$  is an isomorphism for every  $s \in S$ , then  $p$  itself is an isomorphism.

**Corollary 2.12.**— *Let  $G$  be an  $S$ -scheme admitting an  $S$ -section  $\epsilon$ , and suppose that the structural morphism  $\pi : G \rightarrow S$  is closed. Let  $s \in S$  be such that  $\pi$  is of finite presentation at  $\epsilon(s)$ , and such that*

$$\epsilon_s : \text{Spec}(\kappa(s)) \longrightarrow G_s$$

*is an isomorphism (or, equivalently, such that*

$$\pi_s : G_s \longrightarrow \text{Spec} \kappa(s)$$

*is an isomorphism). Then there exists an open neighborhood  $U$  of  $s$  in  $S$  such that*

$$\epsilon|_U : U \longrightarrow G \times_S U$$

*is an isomorphism.*

Let  $V$  be the set of points of  $G$  at which  $\pi$  is unramified. This set is known to be open (cf. Editor's note 6), and it contains  $\epsilon(s)$ . Hence

$$W = \epsilon^{-1}(V)$$

is an open subset of  $S$  containing  $s$ , and  $\pi$  is unramified at  $\epsilon(t)$  for every  $t \in W$ . Since  $\pi$  is closed, the same is true of  $\pi|_W$ , and we may therefore suppose that  $W = S$ .

By 2.7.1,

$$X = G - \epsilon(S)$$

is then a closed subset of  $G$  that does not meet  $G_s$ . Since  $\pi$  is closed,

$$F = \pi(X)$$

is a closed subset of  $S$  that does not contain  $s$ . Set

$$U = S - F.$$

Then  $U$  is an open subset of  $S$ , and  $\epsilon|_U$  is an isomorphism from  $U$  onto  $G|_U$ .

### 3. Identity component of a group locally of finite presentation

**3.0.**— Given a subset  $A$  (respectively,  $B$ ) of an  $S$ -scheme  $X$  (respectively,  $Y$ ), by abuse of notation

$$A \times_S B$$

will denote the subset of  $X \times_S Y$  consisting of the points whose first projection belongs to  $A$  and whose second projection belongs to  $B$ .

Given a subset  $A$  of an  $S$ -group  $G$ , we shall say that  $A$  is *stable under the group law of  $G$*  if

$$c(A) \subset A \quad \text{and} \quad \mu(A \times_S A) \subset A.$$

**Definition 3.1.**— Let  $G$  be an  $S$ -functor in groups satisfying the following condition:

$$\text{for every } s \in S, \text{ the functor } G_s = G \otimes_S \kappa(s) \text{ is representable.} \quad (+)$$

We then denote by  $\underline{G}_s^0$  the connected component of the identity element of the  $\kappa(s)$ -group  $G_s$ .<sup>22</sup> We define a sub- $S$ -functor in groups of  $G$ , called the *identity component of  $G$*  and denoted by  $G^0$ , by setting, for every morphism  $T \rightarrow S$ ,

$$G^0(T) = \{u \in G(T) \mid \text{for every } s \in S, u_s(T_s) \subset \underline{G}_s^0\}.$$

Thus we have defined the functor

$$G \longmapsto G^0$$

from

$$(\widehat{\mathbf{Sch}}/S)\text{-gr.} \quad \text{to} \quad (\widehat{\mathbf{Sch}}/S)\text{-gr.}$$

**Remark 3.2.**— (i) Let  $G$  be an  $S$ -functor in groups satisfying condition  $(+)$ . Then

$$\mathrm{Lie}(G^0/S) = \mathrm{Lie}(G/S)$$

by Exposé II.<sup>23</sup> Indeed, for every  $S$ -scheme  $T$ , denote by  $I_T$  the  $T$ -scheme of dual numbers over  $T$ , and by

$$\tau : T \longrightarrow I_T$$

the zero section (cf. Exposé II, 2.1). By definition,

$$\mathrm{Lie}(G/S)(T) = \{u \in G(I_T) \mid u \circ \tau = e\},$$

where  $e : T \rightarrow G$  denotes the composite of  $T \rightarrow S$  with the identity section  $S \rightarrow G$ . Likewise,

$$\begin{aligned} \mathrm{Lie}(G^0/S)(T) &= \{u \in G^0(I_T) \mid u \circ \tau = e\} \\ &= \{u \in G(I_T) \mid u \circ \tau = e \text{ and } u_s((I_T)_s) \subset \underline{G}_s^0 \text{ for every } s \in S\}. \end{aligned}$$

For every  $s \in S$ , the schemes  $T_s$  and

$$(I_T)_s = I_{T_s}$$

have the same underlying set. Hence, if  $u \in \mathrm{Lie}(G/S)(T)$ , then

$$u_s(I_{T_s}) = e_s,$$

<sup>22</sup>Editor's note: this is a slightly abusive notation, but it is compatible with the notation of Exposé VI<sub>A</sub>, 2.3 when this connected component is the underlying topological space of an open group subscheme  $G^0$  of  $G$ ; cf. Exposé VI<sub>A</sub>, 2.2.bis.

<sup>23</sup>Editor's note: we have added what follows.

where  $e_s$  denotes the identity point of  $G_s$ , and therefore  $u \in \text{Lie}(G^0/S)(T)$ . Thus the inclusion

$$\text{Lie}(G^0/S)(T) \subset \text{Lie}(G/S)(T)$$

is an equality for every  $T$ , and consequently

$$\text{Lie}(G^0/S) = \text{Lie}(G/S).$$

(ii) Let  $G$  and  $G'$  be two  $S$ -functors in groups satisfying (+). Then:

- a) if  $G \subset G'$ , then  $G^0 \subset G'^0$ ;
- b) if  $G \subset G'$  and  $G'^0 \subset G$ , then  $G^0 = G'^0$ ;
- c) if, for every  $s \in S$ ,  $G_s$  is locally of finite type over  $\kappa(s)$ , then  $G^0$  satisfies condition (+), by Exposé VI<sub>A</sub>, 2.3, and

$$(G^0)^0 = G^0.$$

**Proposition 3.3.**— *Let  $G$  be an  $S$ -functor in groups satisfying condition (+), and let  $S'$  be an  $S$ -scheme. Then*

$$(G \times_S S')^0 = G^0 \times_S S'.$$

*In other words, the functor  $G \mapsto G^0$  commutes with base change; that is, the following diagram is commutative:*

$$\begin{array}{ccc} (\widehat{\text{Sch}}_S)\text{-gr.} & \xrightarrow{G \mapsto G^0} & (\widehat{\text{Sch}}_S)\text{-gr.} \\ \downarrow & & \downarrow \\ (\widehat{\text{Sch}}_{S'})\text{-gr.} & \longrightarrow & (\widehat{\text{Sch}}_{S'})\text{-gr.} \end{array}$$

It is enough to check that, for every  $s' \in S'$ , whose image in  $S$  we denote by  $s$ ,

$$\begin{aligned} ((G \times_S S') \otimes_{S'} \kappa(s'))^0 &= (G_s \otimes_{\kappa(s)} \kappa(s'))^0 \\ &= G_s^0 \otimes_{\kappa(s)} \kappa(s'). \end{aligned}$$

This follows from Exposé VI<sub>A</sub>, 2.1.2. Note, for later use in 4.2, that we have not used the group structure of  $G_s$ , but only the fact that  $G_s^0$  has a rational point, namely  $\epsilon(s)$ , and is therefore geometrically connected (see also EGA IV<sub>2</sub>, 4.5.14).

**Special Case 3.4.**— Let  $G$  be an  $S$ -group scheme, and denote by  $\underline{G}^0$  the subset of  $G$  equal to the union of the  $\underline{G}_s^0$  as  $s$  runs through  $S$ . Then  $\underline{G}^0$  is a subset of  $G$  stable under the group law of  $G$  (cf. 3.0), and, for every morphism  $S' \rightarrow S$ ,

$$G^0(S') = \{u \in G(S') \mid u(S') \subset \underline{G}^0\}.$$

When  $\underline{G}^0$  is an open subset of  $G$ , it is clear that  $G^0$  is represented by the subscheme of  $G$  induced on the open subset  $\underline{G}^0$ .

**Proposition 3.5.**— *Let  $S$  be a quasi-compact and quasi-separated scheme, and let  $G$  be an  $S$ -group with fibers locally of finite type. Then there is a quasi-compact open subset  $U$  of  $G$  containing  $\underline{G}^0$ .*

Since the identity section  $\epsilon$  is an immersion,  $\epsilon(S)$  is a quasi-compact subspace of  $G$ . Thus there is a quasi-compact open subset  $V$  of  $G$  containing  $\epsilon(S)$ . Since  $S$  is quasi-separated

and  $V$  is quasi-compact,  $V$  is quasi-compact over  $S$  (EGA IV<sub>1</sub>, 1.2.4). Hence  $V \times_S V$  is quasi-compact over  $S$ , and therefore quasi-compact. It follows that

$$V \cdot V = \mu(V \times_S V)$$

is quasi-compact. Set

$$V_s = V \cap G_s, \quad V_s^0 = V \cap \underline{G}_s^0.$$

Then  $V_s^0$  is open in  $\underline{G}_s^0$ , and is dense in  $\underline{G}_s^0$  because  $G_s^0$  is irreducible (Exposé VI<sub>A</sub>, 2.4). Therefore

$$V_s^0 \cdot V_s^0 = \underline{G}_s^0 \quad (\text{Exposé VI}_A, 0.5),$$

which shows that

$$V_s \cdot V_s \supset \underline{G}_s^0, \quad \text{and hence} \quad V \cdot V \supset \underline{G}^0.$$

Finally, since  $V \cdot V$  is quasi-compact, there is a quasi-compact open subset  $U$  of  $G$  containing  $V \cdot V$ , and therefore containing  $\underline{G}^0$ .

**Corollary 3.6.**— *Let  $G$  be an  $S$ -group whose fibers are locally of finite type and connected. Then  $G$  is quasi-compact over  $S$ .*

The assertion is local on  $S$  (EGA I, 6.6.1), so we are reduced to the case where  $S$  is affine. By 3.5, there is then a quasi-compact open subset  $U$  of  $G$  containing

$$\underline{G}^0 = G.$$

Thus  $G$  is quasi-compact, and hence quasi-compact over the affine scheme  $S$  (EGA I, 6.6.4(v)).

**Proposition 3.7.**—<sup>24</sup> *Let  $G$  be an  $S$ -group locally of finite presentation. Then:*

- (i)  $\underline{G}^0$  is ind-constructible in  $G$ .
- (ii) If, moreover,  $G$  is quasi-separated over  $S$ , and  $S$  is quasi-compact and quasi-separated, then  $\underline{G}^0$  is constructible.
- (iii) Consequently, if  $G$  is quasi-separated over  $S$ , then  $\underline{G}^0$  is locally constructible.

We first prove the first assertion. Since

$$\pi : G \longrightarrow S$$

is locally of finite presentation, given  $s \in S$ , there are an open subset  $U$  of  $G$  containing  $\epsilon(s)$  and an open subset  $V$  of  $S$  containing  $s$  such that

$$\pi(U) \subset V$$

<sup>24</sup>Editor's note: we recall the following definitions and results (cf. EGA 0<sub>III</sub>, §9.1, and EGA IV<sub>1</sub>, §§1.8 and 1.9). Let  $X$  be a topological space. (i) An open subset  $U$  of  $X$  is called *retrocompact* if the inclusion  $U \hookrightarrow X$  is quasi-compact, i.e. if  $U \cap V$  is quasi-compact for every quasi-compact open subset  $V \subset X$ . (ii) A subset  $C$  of  $X$  is called *constructible* if it is a finite union of subsets of the form  $U - V$ , where  $U$  and  $V$  are retrocompact open subsets of  $X$ . (iii) A subset  $L$  of  $X$  is called *locally constructible* if, for every  $x \in X$ , there is an open neighborhood  $U$  of  $x$  in  $X$  such that  $L \cap U$  is constructible in  $U$ . (If  $X$  is quasi-compact and quasi-separated, then  $L$  is constructible.) (iv) A subset  $I$  of  $X$  is called *ind-constructible* if, for every  $x \in X$ , there is an open neighborhood  $U$  of  $x$  in  $X$  such that  $I \cap U$  is a union of locally constructible subsets of  $U$ . Now let  $f : X \rightarrow Y$  be a morphism of schemes. By Chevalley's constructibility theorem (cf. EGA IV<sub>1</sub>, 1.8.4 and 1.9.5(viii)), if  $f$  is of finite presentation (respectively, locally of finite presentation), then the image under  $f$  of every locally constructible subset is locally constructible (respectively, ind-constructible).

and the induced morphism

$$\pi' : U \longrightarrow V$$

is of finite presentation. Then

$$T = \epsilon^{-1}(U)$$

is an open subset of  $S$ . Set

$$W = (\pi')^{-1}(T), \quad \pi'' = \pi'|_W.$$

Thus  $\pi'' : W \rightarrow T$  is of finite presentation and admits as a section the morphism

$$\epsilon'' : T \longrightarrow W$$

induced by  $\epsilon$ .

For every  $t \in T$ , since  $\underline{G}_t^0$  is irreducible (VI<sub>A</sub>, 2.4),  $W \cap \underline{G}_t^0$  is dense in  $\underline{G}_t^0$ , hence irreducible and therefore connected. It is thus the connected component of  $(\pi'')^{-1}(t)$  containing  $\epsilon''(t)$ . It follows from EGA IV<sub>3</sub>, 9.7.12, that the union  $W^0$  of the  $W \cap \underline{G}_t^0$ , for  $t \in T$ , is locally constructible in  $W$ . On the other hand, VI<sub>A</sub>, 0.5, gives

$$\underline{G}^0 \cap \pi^{-1}(T) = W^0 \cdot W^0;$$

that is,  $\underline{G}^0 \cap \pi^{-1}(T)$  is the image of

$$W^0 \times_T W^0$$

under the morphism

$$\mu'' : W \times_T W \longrightarrow \pi^{-1}(T)$$

induced by  $\mu$ .<sup>25</sup> Since  $W \times_T W$  (respectively,  $\pi^{-1}(T)$ ) is of finite presentation (respectively, locally of finite presentation) over  $T$ , the morphism  $\mu''$  is locally of finite presentation and quasi-separated, by EGA IV<sub>1</sub>, 1.4.3 and 1.2.2. If, moreover,  $\pi^{-1}(T)$  is quasi-separated over  $T$ , then  $\mu''$  is quasi-compact (loc. cit., 1.2.4), hence of finite presentation. Since  $W^0 \times_T W^0$  is locally constructible in  $W \times_T W$  (because  $W^0$  is locally constructible in  $W$ ), Chevalley's constructibility theorem (loc. cit., 1.8.4 and 1.9.5(viii)) shows that  $\underline{G}^0 \cap \pi^{-1}(T)$  is ind-constructible in  $\pi^{-1}(T)$ , and is locally constructible in  $\pi^{-1}(T)$  if  $G$ , and hence  $\pi^{-1}(T)$ , is quasi-separated over  $T$ . This proves assertions (i) and (iii).

Now suppose that  $G$  is quasi-separated over  $S$ , and that  $S$  is quasi-compact and quasi-separated. By 3.5, there is a quasi-compact open subset  $U$  of  $G$  containing  $\underline{G}^0$ . Since  $G$  is quasi-separated over  $S$ ,  $G$  is quasi-separated; hence the open subset  $U$  is retrocompact (EGA IV<sub>1</sub>, 1.2.7), and it suffices to show that  $\underline{G}^0$  is constructible in  $U$  (EGA 0<sub>III</sub>, 9.1.8).

Moreover,  $U$  is quasi-compact, hence quasi-compact over  $S$  (EGA IV<sub>1</sub>, 1.2.4), and is quasi-separated over  $S$ . Therefore the restriction of  $\pi$  to  $U$  is of finite presentation. By EGA IV<sub>3</sub>, 9.7.12,  $\underline{G}^0$  is locally constructible in  $U$ , hence constructible in  $U$ , since  $U$  is quasi-compact and quasi-separated (EGA IV<sub>1</sub>, 1.8.1). This proves (ii), and it follows that, for every quasi-compact and quasi-separated open subset  $T$  of  $S$  (for example, every affine open subset of  $S$ ),

$$\underline{G}^0 \cap \pi^{-1}(T)$$

is constructible.

<sup>25</sup>Editor's note: we have expanded the original in what follows.

**Corollary 3.8.**— *Let  $S_0$  be a quasi-compact and quasi-separated scheme,  $I$  an increasing filtered preordered set,  $(\mathcal{A}_i)_{i \in I}$  a direct system of commutative quasi-coherent  $\mathcal{O}_{S_0}$ -algebras,*

$$\mathcal{A} = \varinjlim_i \mathcal{A}_i, \quad S_i = \operatorname{Spec} \mathcal{A}_i \quad (i \in I), \quad S = \operatorname{Spec} \mathcal{A}$$

(cf. EGA II, 1.3.1). *Let  $G$  be an  $S_0$ -group scheme locally of finite presentation. Then the canonical map*

$$\varinjlim_i G^0(S_i) \longrightarrow G^0(S)$$

*is bijective.*

Since  $G$  is locally of finite presentation over  $S_0$ ,\* the canonical map

$$\varinjlim_i G(S_i) \longrightarrow G(S)$$

is bijective, by EGA IV<sub>2</sub>, 8.14.2(c). It follows immediately that the canonical map

$$\varinjlim_i G^0(S_i) \longrightarrow G^0(S)$$

is injective. We prove that it is surjective. Let

$$g \in G^0(S) \subset G(S).$$

There is an  $i \in I$  such that  $g$  factors, by means of an element  $g_i \in G(S_i)$ , through  $S \rightarrow S_i$ . By hypothesis,

$$g^{-1}(\underline{G}^0) = S.$$

But  $\underline{G}^0$  is ind-constructible in  $G$ , by 3.7, so

$$g_i^{-1}(\underline{G}^0)$$

is ind-constructible in  $S_i$ . It follows from EGA IV<sub>2</sub>, 8.3.4, that there is an index  $j \geq i$  such that

$$g_j^{-1}(\underline{G}^0) = S_j,$$

where  $g_j$  is the map induced by  $g_i$  under the base change  $S_j \rightarrow S_i$ . Thus  $g_j \in G^0(S_j)$ , and hence  $g$  comes from an element of

$$\varinjlim_i G^0(S_i).$$

**Proposition 3.9.**— *Let  $G$  be an  $S$ -group locally of finite presentation. Suppose that  $G^0$  is representable. Then the canonical morphism*

$$i : G^0 \longrightarrow G$$

*is an open immersion; moreover,  $G^0$  is quasi-compact over  $S$ .*

Since  $G^0$  is a subfunctor of the functor  $G$ , the morphism  $i$  is a monomorphism, hence is radicial. From the definition of the functor  $G^0$ , one checks immediately from the definition (EGA IV<sub>4</sub>, 17.1.1) that  $i$  is formally étale (noting that  $G^0$  is the image of  $i$  in  $G$ ). Finally, by the characterization (EGA IV<sub>3</sub>, 8.14.2(c)) of  $S$ -schemes locally of finite presentation in terms of the functors they represent, and by 3.8,  $G^0$  is locally of finite presentation over  $S$ , since  $G$  is. Hence  $i$  is locally of finite presentation (cf. EGA IV<sub>1</sub>, 1.4.3). It is therefore a radicial étale morphism, and hence an open immersion (EGA IV<sub>4</sub>, 17.9.1).

The last assertion follows from 3.6.

**Theorem 3.10.**— *Let  $G$  be an  $S$ -group. The following conditions are equivalent:*

---

\*Translator's note: the French source prints "over  $S$ " here. The scheme  $G$  was defined over  $S_0$ , while  $S$  and the  $S_i$  are the test schemes of the direct system; the required base is therefore  $S_0$ .

- (i)  $G$  is smooth over  $S$  at the points of the identity section.
- (ii)  $G$  is flat and locally of finite presentation over  $S$  at the points of the identity section, and, for every  $s \in S$ ,  $G_s$  is smooth over  $\kappa(s)$ .
- (iii) There is an open group subscheme  $G'$  of  $G$  that is smooth over  $S$ .
- (iv)  $G^0$  is representable by an open subscheme of  $G$  that is smooth over  $S$ .

It is clear that

$$(iv) \implies (iii) \implies (i),$$

and, by 1.3.1 and 2.4, (i) implies both (ii) and (iii). Moreover, (ii) implies (i) by EGA IV<sub>4</sub>, 17.5.1.

It remains to prove that (iii) implies (iv). Lemma 3.10.1 below shows that  $G'$  contains  $\underline{G}^0$  and that

$$G'^0 = G^0.$$

It therefore suffices to show that  $\underline{G}^0$  is open in  $G$ , for we have already seen in 3.4 that  $G^0$  will then be represented by the smooth group subscheme induced by  $G'$  on the open subset

$$\underline{G}'^0 = \underline{G}^0.$$

We may thus suppose that  $G' = G$ .

To show that  $\underline{G}^0$  is open, it suffices to show that every  $s \in S$  has a neighborhood  $T$  in  $S$  such that

$$\underline{G}^0 \cap \pi^{-1}(T)$$

is open in  $\pi^{-1}(T)$ . Let  $s \in S$ . Since  $G = G'$ , the morphism  $\pi$  is locally of finite presentation. As in the proof of 3.7, we may therefore construct an open subset  $T$  of  $S$  containing  $s$ , and an open subset  $W$  of  $G$  containing  $\epsilon(s)$ , such that the induced morphism

$$\pi'' : W \longrightarrow T$$

is of finite presentation and admits as a section the morphism

$$\epsilon'' : T \longrightarrow W$$

induced by  $\epsilon$ . For every  $t \in T$ ,

$$W \cap \underline{G}_t^0$$

is then the connected component of  $(\pi'')^{-1}(t)$  containing  $\epsilon''(t)$ . Since  $\pi$  is smooth, so is  $\pi''$ , which is therefore smooth and of finite presentation. By EGA IV<sub>3</sub>, 15.6.5, the union  $W^0$  of the  $W \cap \underline{G}_t^0$ , for  $t \in T$ , is open in  $W$ .

On the other hand, VI<sub>A</sub>, 0.5, gives

$$W^0 \cdot W^0 = \underline{G}^0 \cap \pi^{-1}(T),$$

and we must show that this is open in  $\pi^{-1}(T)$ . We may now suppose that  $T = S$ ; it remains to show that  $W^0 \cdot W^0$  is open in  $G$ . Since  $\pi$  is universally open, so is  $\mu$  (VI<sub>A</sub>, 0.1). Hence, since  $W^0$  is open in  $G$ , so is

$$W^0 \cdot W^0 = \mu(W^0 \times_S W^0).$$

This result will be strengthened in 4.4.

**Lemma 3.10.1.**— *Let  $G$  be an  $S$ -group whose fibers are locally of finite type. Then every open group subscheme  $H$  of  $G$  contains  $\underline{G}^0$  and satisfies*

$$G^0 = H^0.$$

Let  $s \in S$ , and set

$$G'_s = H_s \cap \underline{G}_s^0.$$

Then  $G'_s$  is an open subset of  $\underline{G}_s^0$ , and is dense in  $\underline{G}_s^0$  since  $\underline{G}_s^0$  is irreducible (VI<sub>A</sub>, 2.4). Hence

$$G'_s \cdot G'_s = \underline{G}_s^0$$

by VI<sub>A</sub>, 0.5, which shows that

$$\underline{G}_s^0 = G'_s \cdot G'_s \subset H_s \cdot H_s = H_s.$$

Thus  $\underline{G}_s^0 = \underline{H}_s^0$  for every  $s$ , whence

$$\underline{G}^0 \subset H, \quad G^0 = H^0.$$

**Proposition 3.11.**—<sup>26</sup> *Let  $u : G \rightarrow H$  be a morphism between  $S$ -groups locally of finite presentation. If  $u$  is flat, the induced map*

$$u^0 : G^0 \longrightarrow H^0$$

*is surjective. The converse holds if  $G$  is flat over  $S$  and  $H$  has reduced fibers.*

Suppose that  $u$  is flat. Then, for every  $s \in S$ ,  $u_s$  is flat and locally of finite presentation, hence open (EGA IV<sub>2</sub>, 2.4.6). Therefore

$$u_s^0 : \underline{G}_s^0 \longrightarrow \underline{H}_s^0$$

is surjective, by 1.3.2.<sup>27</sup> Thus the map

$$u^0 : G^0 \longrightarrow H^0$$

is surjective.

Conversely, suppose that  $u^0 : G^0 \rightarrow H^0$  is surjective, that  $G$  is flat over  $S$ , and that  $H$  has reduced fibers. Then, for every  $s \in S$ , the morphism

$$u_s^0 : \underline{G}_s^0 \longrightarrow \underline{H}_s^0$$

is surjective, and hence is flat at every point lying above the generic point  $\eta$  of  $\underline{H}_s^0$ , since

$$\mathcal{O}_{\underline{H}_s^0, \eta}$$

is a field. It follows from 1.3 that  $u_s$  is flat. Therefore  $u$  is flat, by the fiberwise flatness criterion (EGA IV<sub>3</sub>, 11.3.11).

<sup>26</sup>Editor's note: compare VI<sub>A</sub>, 5.6.

<sup>27</sup>Editor's note: we have shortened the original by referring here to 1.3.2. We have also simplified the argument below by omitting a reference to the generic flatness theorem (EGA IV<sub>2</sub>, 6.9.1), which is not needed here.



## 4. Dimension of the fibers of groups locally of finite presentation

**Proposition 4.1.**— *Let  $G$  be an  $S$ -scheme locally of finite type, equipped with an  $S$ -section  $\epsilon$ , and suppose that, for every  $s \in S$ ,*

$$\dim G_s = \dim_{\epsilon(s)} G_s$$

(as is the case when  $G$  is an  $S$ -group; 1.5).

(i) *The function*

$$s \longmapsto \dim G_s$$

*is upper semicontinuous on  $S$ .*

(ii) *If, moreover,  $G$  is locally of finite presentation over  $S$ , this function is locally constructible.*

Let

$$\pi : G \longrightarrow S$$

be the structural morphism. Chevalley's semicontinuity theorem (EGA IV<sub>3</sub>, 13.1.3) asserts that the function

$$x \longmapsto \dim_x \pi^{-1}(\pi(x))$$

is upper semicontinuous on  $G$ . For every  $s \in S$ , however,

$$\dim G_s = \dim \pi^{-1}(s) = \dim_{\epsilon(s)} \pi^{-1}(\pi(\epsilon(s))).$$

Since the map  $s \mapsto \epsilon(s)$  is continuous on  $S$ , the composite function

$$s \longmapsto \dim G_s$$

is upper semicontinuous on  $S$ .

Now suppose that  $G$  is locally of finite presentation over  $S$ . Arguing as above, to show that  $s \mapsto \dim G_s$  is locally constructible, it is enough to show that the function

$$x \longmapsto \dim_x \pi^{-1}(\pi(x))$$

is locally constructible on  $G$ , which follows from EGA IV<sub>3</sub>, 9.9.1.

**Proposition 4.2.**— *Let  $\pi : G \rightarrow S$  be an  $S$ -scheme locally of finite presentation, equipped with an  $S$ -section  $\epsilon$ , and satisfying the following two conditions (which hold when  $G$  is an  $S$ -group, by 1.5 and VI<sub>A</sub>, 2.4.1):*

(a) *For every  $s \in S$  and every  $x \in G_s$ , one has*

$$\dim G_s = \dim_x G_s$$

(or equivalently (1.5), for every  $s \in S$ , all the irreducible components of  $G_s$  have the same dimension).

(b) *For every  $s \in S$ , if  $\underline{G}_s^0$  denotes the connected component of  $G_s$  containing  $\epsilon(s)$ , then  $\underline{G}_s^0$  is geometrically irreducible.*

*Let  $s \in S$ . The following conditions are equivalent:*

- (i)  $G$  is universally open over  $S$  at the points of  $\underline{G}_s^0$ .
- (i bis)  $G$  is universally open over  $S$  at every point of a neighborhood of  $\epsilon(s)$  in  $\underline{G}_s^0$ .<sup>28</sup>
- (ii) The function
- $$t \longmapsto \dim G_t$$
- is constant in a neighborhood of  $s$  in  $S$ .
- (iii)  $\underline{G}^0$  is “universally open over  $S$  at the points of  $\underline{G}_s^0$ ,” that is, given  $S' \rightarrow S$ ,  $s' \in S'_s$ ,  $g \in \underline{G}_{s'}^0$ , and an open neighborhood  $V$  of  $g$  in  $G' = G_{S'}$ , then

$$\pi(V \cap \underline{G}'^0)$$

is an open neighborhood of  $s'$  in  $S'$ .<sup>29</sup>

Clearly (i) implies (i bis). By EGA IV<sub>3</sub>, 14.3.3.1(ii), the set of points of  $\underline{G}_s^0$  at which  $\pi_G$  is universally open is closed in  $\underline{G}_s^0$ . Since  $\underline{G}_s^0$  is irreducible, it follows that (i bis) implies (i).

We show that (i) implies (ii). Let  $T$  be the set of  $t \in S$  such that

$$\dim G_t = \dim G_s.$$

By 4.1,  $T$  is locally constructible; hence, by EGA IV<sub>1</sub>, 1.10.1, to show that  $T$  is a neighborhood of  $s$ , it is enough to show that every generalization  $s'$  of  $s$  belongs to  $T$ .

Let  $x$  be the generic point of  $\underline{G}_s^0$ , and let  $U$  be an open subset of  $G$  containing  $x$ . Since  $\pi_G$  is universally open at  $x$ , EGA IV<sub>3</sub>, 14.3.13 gives, for every generalization  $s'$  of  $s$ ,

$$\dim(U \cap G_{s'}) \geq \dim_x(U \cap G_s).$$

In view of hypothesis (a), this implies

$$\dim G_{s'} \geq \dim G_s.$$

On the other hand, by 4.1 the function  $s \mapsto \dim G_s$  is upper semicontinuous, so that

$$\dim G_{s'} \leq \dim G_s.$$

Thus  $s' \in T$ , proving that (i) implies (ii).

It is clear that (iii) implies (i); we prove that (ii) implies (iii). The dimension of the fibers is unchanged by extension of the base field, and the formation of  $\underline{G}^0$  commutes with base change (see the proof of 3.3). We may therefore assume that  $S' = S$  and  $s' = s$ . Moreover, every open subset  $V$  of  $G$  meeting  $\underline{G}_s^0$  contains the generic point  $\eta$  of  $\underline{G}_s^0$ , so we may assume that  $g = \eta$ .

We may further suppose that  $S$  is affine. Let  $U$  be an affine open neighborhood of  $\epsilon(s)$ ; it is then of finite presentation over  $S$ . Replacing  $S$  by  $\epsilon^{-1}(U)$ , and then  $U$  by  $U \cap \pi^{-1}(S)$ , we reduce to the case in which

$$\pi : U \longrightarrow S$$

is of finite presentation and admits  $\epsilon : S \rightarrow U$  as a section. By EGA IV<sub>3</sub>, 9.7.12,  $\underline{G}^0 \cap U$  is constructible in  $U$ . Replacing  $V$  by an affine open subset contained in  $V \cap U$ , we find

<sup>28</sup>Editor's note: the original reads “at the points of  $\epsilon(s)$ .” It is not enough to suppose that  $G$  is universally open over  $S$  at  $\epsilon(s)$ , as the example given in editor's note 19 shows. We have corrected the proof accordingly.

<sup>29</sup>Editor's note: we have added condition (iii), pointed out by O. Gabber; it will be useful later (5.7).

that  $\underline{G}^0 \cap V$  is constructible in  $V$ . Since  $\pi : V \rightarrow S$  is of finite presentation, Chevalley's constructibility theorem (EGA IV<sub>1</sub>, 1.8.4) shows that

$$\pi(\underline{G}^0 \cap V)$$

is locally constructible in  $S$ .

Consequently, by *loc. cit.*, 1.10.1, to show that  $\pi(\underline{G}^0 \cap V)$  is an open neighborhood of  $s$ , it is enough to show that, for every generalization  $t$  of  $s$ , there is a generalization  $\xi$  of  $\eta$  belonging to  $\underline{G}^0$ , and hence to  $\underline{G}^0 \cap V$ . The generic point  $\xi$  of  $\underline{G}_t^0$  is such a generalization of  $\eta$ . Indeed,  $\epsilon(s)$  belongs to the closure  $X$  of  $\{\xi\}$  in  $G$ ; hence, by Chevalley's semicontinuity theorem (EGA IV<sub>3</sub>, 13.1.3),

$$\dim_{\epsilon(s)} X_s \geq \dim_{\xi} X_t.$$

On the other hand, hypothesis (ii) implies

$$\dim_{\xi} \underline{G}_t^0 = \dim_{\epsilon(s)} G_s.$$

It follows that one of the irreducible components of  $X_s$  containing  $\epsilon(s)$  is equal to  $\underline{G}_s^0$ , whence

$$\eta \in \overline{\{\xi\}}.$$

This proves that (ii) implies (iii), and hence proves the proposition.

<sup>30</sup>One can also prove that (ii) implies (i) as follows. Since  $\pi$  is locally of finite presentation, there exist an open subset  $U$  of  $G$  containing  $\epsilon(s)$  and an open subset  $V$  of  $S$  containing  $s$ , such that  $\pi(U) \subset V$  and the morphism

$$\pi' : U \longrightarrow V$$

induced by  $\pi$  is of finite presentation. Set

$$T = \epsilon^{-1}(U), \quad W = \pi'^{-1}(V) = U \cap \pi^{-1}(V).$$

The morphism

$$\pi'' : W \longrightarrow T$$

induced by  $\pi'$  is of finite presentation and admits as a section the morphism

$$\epsilon'' : T \longrightarrow W$$

induced by  $\epsilon$ . Moreover, for every  $t \in T$ , since  $\underline{G}_t^0$  is irreducible,  $W \cap \underline{G}_t^0$  is dense in  $\underline{G}_t^0$ , hence irreducible and therefore connected. It is consequently the connected component of  $\pi''^{-1}(t)$  containing  $\epsilon''(t)$ .

Since  $W \cap \underline{G}_t^0$  is a dense open subset of  $\underline{G}_t^0$ , 1.5 and EGA IV<sub>2</sub>, 5.2.1 give

$$\dim(W \cap \underline{G}_t^0) = \dim \underline{G}_t^0 = \dim G_t.$$

Thus the function

$$t \longmapsto \dim(W \cap \underline{G}_t^0)$$

is constant in a neighborhood of  $s$  in  $T$ . Finally, we show that  $W \cap \underline{G}_t^0$  is geometrically irreducible for every  $t \in T$ . Let  $K$  be an extension of  $\kappa(t)$ . Then

$$(W \cap \underline{G}_t^0) \otimes_{\kappa(t)} K = (W \otimes_{\kappa(t)} K) \cap (\underline{G}_t^0 \otimes_{\kappa(t)} K)$$

---

<sup>30</sup>Editor's note: the preceding argument was communicated to us by O. Gabber. We have also retained the proof of the implication (ii)  $\Rightarrow$  (i) given in the original.

is a nonempty open subset of  $\underline{G}_t^0 \otimes_{\kappa(t)} K$ , and is therefore irreducible because  $\underline{G}_t^0 \otimes_{\kappa(t)} K$  is irreducible.

We are now in the situation of EGA IV<sub>3</sub>, 15.6.6(ii), which asserts that  $\pi''$ , and hence  $\pi$ , is universally open at the points of  $W \cap \underline{G}_s^0$ . But, by EGA IV<sub>3</sub>, 14.3.3.1(ii), the subset of  $G_s$  consisting of the points at which  $\pi$  is universally open is closed in  $G_s$ . Since it contains  $W \cap \underline{G}_s^0$ , it therefore contains its closure  $\underline{G}_s^0$ .

**Corollary 4.3.**— *Let  $G$  be a flat  $S$ -group locally of finite presentation. Then the function*

$$s \longmapsto \dim G_s$$

*is locally constant on  $S$ .*

This follows immediately from 4.2, since every flat morphism locally of finite presentation is universally open (EGA IV<sub>2</sub>, 2.4.6).

**Corollary 4.4.**— *Let  $G$  be an  $S$ -group locally of finite presentation over  $S$  at the points of the identity section. Consider the following conditions:*

(i)  *$G$  is smooth over  $S$  at the points of the identity section (cf. 3.10).*

(ii) *For every  $s \in S$ ,  $G_s$  is smooth over  $\kappa(s)$ , and the function*

$$s \longmapsto \dim G_s$$

*is locally constant on  $S$ .*

(iii) *For every  $s \in S$ ,  $G_s$  is smooth over  $\kappa(s)$ , and there exists a neighborhood  $V$  of the identity section such that*

$$\pi_V : V \longrightarrow S$$

*is universally open.*

(iv) *For every  $s \in S$ ,  $G_s^0$  is smooth over  $\kappa(s)$ , and  $G^0$  is representable by an open group subscheme of  $G$ , universally open over  $S$ .*

*Then the following implications hold:*

$$(i) \implies (ii) \iff (iii) \iff (iv).$$

*If, moreover,  $S$  is reduced, then conditions (i)–(iv) are equivalent and imply that  $G^0$  is smooth over  $S$ .<sup>31</sup>*

Let us show that (i) implies (ii). For every  $x \in G$ , by 1.5,

$$\dim_x \pi^{-1}(\pi(x)) = \dim \pi^{-1}(\pi(x)).$$

Consequently, by EGA IV<sub>4</sub>, 17.10.2, the function

$$x \longmapsto \dim_x \pi^{-1}(\pi(x)) = \dim \pi^{-1}(\pi(x))$$

<sup>31</sup>Editor's note: the hypothesis that  $S$  be reduced cannot be omitted here. Indeed, let  $k$  be a field, let  $S = \operatorname{Spec} A$ , where  $A = k[\delta]$  with  $\delta^2 = 0$ , and let  $G$  be the closed subgroup

$$\operatorname{Spec} A[X]/(\delta X)$$

of  $\mathbb{G}_{a,S}$  (which assigns to every  $A$ -algebra  $R$  the subgroup of those  $r \in R$  such that  $\delta r = 0$ ). Then  $G$  is an  $S$ -group satisfying (ii)–(iv), but it is not flat, and hence not smooth, over  $S$ .

is continuous in a neighborhood of the identity section. Therefore the function  $s \mapsto \dim G_s$  is continuous on  $S$ , and hence locally constant on  $S$ . On the other hand, for every  $s \in S$ ,  $G_s$  is smooth over  $\kappa(s)$ , by 1.3.1.

Let us show that (ii) implies (iv). It is enough to show that  $\underline{G}^0$  is open in  $G$ , for then, by 3.4,  $G^0$  will be representable by the group subscheme induced on  $\underline{G}^0$ , and the properties of  $G^0$  stated above will follow from 2.4 and 4.2. Given  $s \in S$ , construct, as in the proof of 3.10,

$$W, \quad T, \quad \pi'', \quad \epsilon'', \quad \text{and} \quad W^0.$$

Then, by EGA IV<sub>3</sub>, 15.6.7,  $W^0$  is open in  $W$ . On the other hand, under hypothesis 4.4(ii), it follows from 4.2 that  $\pi$  is universally open at every point of  $W^0$ . Therefore, by VI<sub>A</sub>, 0.1,  $\mu$  is universally open at every point of

$$W^0 \times_S W^0,$$

which shows that  $W^0 \cdot W^0$  is open; the proof is completed as in the proof of Theorem 3.10.

It is clear that

$$(iv) \implies (iii),$$

and (iii)  $\implies$  (ii) follows by applying 4.2 to  $V$ .

Finally, suppose that (ii)–(iv) hold, and let us show that  $G^0$  is smooth over  $S$  when  $S$  is reduced.<sup>32</sup> For this, we may suppose that  $G = G^0$ . Then  $G$  is of finite presentation over  $S$ , by 5.5 below. Thus  $\pi_G$  is of finite presentation, has geometrically integral fibers, and has locally constant relative dimension on  $S$ . It follows from EGA IV<sub>3</sub>, 15.6.7 that the morphism

$$G \times_S S_{\text{red}} \longrightarrow S_{\text{red}}$$

induced by  $\pi_G$  is flat. Hence  $\pi_G$  is flat when  $S$  is reduced. In that case,  $G = G^0$  is smooth over  $S$ , by Theorem 3.10.

## 5. Separation of groups and homogeneous spaces

**Proposition 5.1.**— *For an  $S$ -group  $G$  to be separated, it is necessary and sufficient that the identity section of  $G$  be a closed immersion.*

The condition is necessary (EGA I, 5.4.6); it is sufficient by virtue of the following cartesian diagram (cf. VI<sub>A</sub>, 0.3):

$$\begin{array}{ccc} G \times_S G & \xrightarrow{\mu \circ (\text{id}_G \times c)} & G \\ \Delta_{G/S} \uparrow & & \uparrow \epsilon \\ G & \xrightarrow{\pi} & S \end{array}.$$

**Proposition 5.2.**— *If  $S$  is discrete, every  $S$ -group is separated.*

Indeed,

$$S = \coprod_{s \in S} \text{Spec } \mathcal{O}_{S,s},$$

and, by EGA I, 5.5.5, it is enough to show that, for every  $s \in S$ ,

$$G \times_S \text{Spec } \mathcal{O}_{S,s}$$

<sup>32</sup>Editor's note: we have expanded the original in the argument that follows.

is separated. This follows from VI<sub>A</sub>, 0.3, since  $\mathcal{O}_{S,s}$  is a local ring of dimension 0.<sup>33</sup>

**Theorem 5.3.**— *Let  $S$  be a scheme, let  $G$  be an  $S$ -group scheme locally of finite presentation over  $S$  and universally open over  $S$  in a neighborhood of the identity section, and let  $X$  be an  $S$ -scheme on which  $G$  acts in such a way that the morphism*

$$\Phi : G \times_S X \longrightarrow X \times_S X, \quad (g, x) \longmapsto (gx, x),$$

*is surjective. Suppose moreover that, for every  $s \in S$ :*

- (i) *there exists an open subscheme  $U$  of  $X$ , separated over  $S$ , such that  $U_s$  is dense in  $X_s$ ;*
- (ii) *the fiber  $X_s$  is locally of finite type over  $\kappa(s)$ .<sup>34</sup>*

*Then  $X$  is separated over  $S$ .*

**Corollary 5.4.**— *Let  $S, G, X$  be as in the preliminary hypotheses of 5.3. Suppose moreover that  $X$  has fibers locally of finite type and connected. Then:*

- (i)  *$X$  is separated over  $S$ .*
- (ii) *If there exists an open subset  $V$  of  $X$ , quasi-compact over  $S$ , which meets every nonempty fiber  $X_s$ ,<sup>35</sup> then  $X$  is quasi-compact over  $S$ .*

*Proof.* (i) Let  $s \in S$  be such that  $X_s \neq \emptyset$ . The morphism

$$\Phi_s : G_s \times_{\kappa(s)} X_s \longrightarrow X_s \times_{\kappa(s)} X_s$$

obtained from  $\Phi$  by base change is surjective. Since  $X_s$  is connected, it is irreducible by VI<sub>A</sub>, 2.5.4. Thus, if  $U$  is an affine open subset of  $X$  such that  $U_s$  is nonempty, then  $U_s$  is dense in  $X_s$ , and the theorem applies.

To prove (ii), we may suppose that  $S$  is affine. Then  $V$  is quasi-compact and, by 3.5, there exists a quasi-compact open subset  $U$  of  $G$  containing  $\underline{G}^0$ . Let  $s \in S$  be such that  $X_s \neq \emptyset$ . Then  $X_s$  is irreducible (VI<sub>A</sub>, 2.6.6), and, since  $U_s$  contains  $\underline{G}_s^0$ , the morphism

$$U_s \times_{\kappa(s)} V_s \longrightarrow X_s, \quad (g, x) \longmapsto gx,$$

is surjective (VI<sub>A</sub>, 2.6.4). Consequently the morphism

$$U \times_S V \longrightarrow X$$

is surjective, and  $X$  is therefore quasi-compact, since  $U$  and  $V$  are. As  $S$  was assumed affine, hence separated, it follows that  $X$  is quasi-compact over  $S$  (cf. EGA I, 6.6.4(v)).

**Remark 5.4.1.**—<sup>36</sup> We shall see in the course of the proof that the conclusion of Theorem 5.3 remains true if one assumes only:

<sup>33</sup>Editor's note: we have replaced "Artinian" by "of dimension 0" and mentioned this generalization in VI<sub>A</sub>, 0.3.

<sup>34</sup>Editor's note: we have corrected the original by assuming that  $G$  is universally open over  $S$  in a neighborhood of the identity section and by adding hypothesis (ii); see below for examples, due to O. Gabber, showing that the statements 5.3 and 5.4 of the original are false without additional hypotheses. We also point out that Theorem 5.3 is a reworked version of Theorem 5.3A below, which appears in the 1965 edition of SGAD and is due to M. Raynaud; compare the notes (\*) in Exposé X, 8.5 and 8.8.

**Theorem 5.3A (Raynaud).**— *Let  $G$  be an  $S$ -group locally of finite presentation, universally open over  $S$ , and with connected fibers. Then  $G$  is separated over  $S$ . More generally, every  $S$ -scheme  $X$  locally of finite presentation over  $S$ , equipped with an action of  $G$  such that the morphism  $\Phi : G \times_S X \rightarrow X \times_S X$ ,  $(g, x) \mapsto (gx, x)$ , is surjective, is separated over  $S$ .*

<sup>35</sup>Editor's note: without this hypothesis there is the following counterexample, pointed out by O. Gabber. Let  $G = S$  be a local scheme with closed point  $s$ , such that  $S^* = S - \{s\}$  is not quasi-compact, and let  $X$  be the disjoint union of  $\{s\}$  and  $S^*$ .

<sup>36</sup>Editor's note: we have added this remark; compare editor's note 34.

(i') there exists an open subscheme  $U$  of  $X$ , separated over  $S$ , such that  $U_s$  is dense in every irreducible component of  $X_s$ .

This follows from (i) if  $X_s$  locally has a finite number of irreducible components, as is the case under hypothesis (ii). We shall also see later that 5.4 remains true under the hypothesis that every fiber  $X_s$  is quasi-separated and connected.

**Corollary 5.5.**— *Let  $S$  be a scheme, and let  $G$  be an  $S$ -group scheme, locally of finite presentation, with connected fibers, and universally open over  $S$ . Then  $G$  is separated and of finite presentation over  $S$ .<sup>37</sup>*

Indeed, by 3.6 and 5.4,  $G$  is quasi-compact and separated over  $S$ . Since  $G$  is locally of finite presentation over  $S$ , it is therefore of finite presentation over  $S$ .

**5.6.**— *Proof of Theorem 5.3.* Before proving 5.3, let us establish a few lemmas.

**Lemma 5.6.0.**—<sup>38</sup>

(i) *Let  $A \subset B$  be integral domains, with  $B$  integral over  $A$ , and let  $\mathfrak{q} \in \operatorname{Spec}(B)$  be such that  $A$  is unibranch at  $\mathfrak{p} = \mathfrak{q} \cap A$ . Then the morphism*

$$\pi : \operatorname{Spec}(B) \longrightarrow \operatorname{Spec}(A)$$

*is open at  $\mathfrak{q}$ .*

(ii) *Let  $X, Y$  be irreducible preschemes, let  $f : X \rightarrow Y$  be a dominant morphism, and let  $x \in X$  be a point at which  $f$  is quasi-finite, such that  $y = f(x)$  is a unibranch point of  $Y$ . Then  $f$  is open at  $x$ . In particular, if*

$$Y = \operatorname{Spec}(\mathcal{O}_{Y,y})$$

*is a local prescheme with closed point  $y$ , then  $f(U) = Y$  for every neighborhood  $U$  of  $x$ .*

(i) Let  $K$  (respectively,  $L$ ) be the fraction field of  $A$  (respectively,  $B$ ), let  $A'$  be the normalization of  $A$ , and let  $B'$  be the subring of  $L$  generated by  $A'$  and  $B$ . Then  $B'$  is integral over  $A'$ . Set

$$Y = \operatorname{Spec}(A), \quad X = \operatorname{Spec}(B), \quad Y' = \operatorname{Spec}(A'), \quad X' = \operatorname{Spec}(B').$$

We then have the commutative diagram

$$\begin{array}{ccc} X' & \xrightarrow{\pi'} & Y' \\ \downarrow & & \downarrow \phi \\ X & \xrightarrow{\pi} & Y \end{array}$$

<sup>37</sup>Editor's note: we point out here the following result ([Ray70a], VI, 2.5). If  $S$  is normal,  $G$  is smooth with connected fibers, and  $X$  is an (fppf) homogeneous  $G$ -space—that is, the morphisms  $X \rightarrow S$  and  $G \times_S X \rightarrow X \times_S X$  are coverings for the (fppf) topology—locally of finite type over  $S$ , then  $X$  is locally quasi-projective over  $S$ . In particular,  $G$  is quasi-projective over  $S$ . See also editor's note 35 in VI<sub>A</sub>.

<sup>38</sup>Editor's note: we have added this lemma, communicated by O. Gabber. It improves EGA IV<sub>3</sub>, 14.4.1.2 and corrects the proof of *loc. cit.*, 14.4.1.3 without changing its hypotheses (compare the erratum (ErrIV, 38) in EGA IV<sub>4</sub>).

in which all morphisms are integral and surjective.

Since  $A$  is unibranch at  $\mathfrak{p}$ ,  $Y'$  has a unique point  $\mathfrak{p}'$  above  $\mathfrak{p}$ . Consequently, if  $U$  is an open neighborhood of  $\mathfrak{p}'$  in  $Y'$ , then

$$\phi(Y' - U)$$

is a closed subset not containing  $\mathfrak{p}$ , so that its open complement is contained in  $\phi(U)$ . This shows that  $\phi : Y' \rightarrow Y$  is open at  $\mathfrak{p}'$ . It is therefore enough to show that  $\pi'$  is open, and we are reduced to the case where  $A = A'$  is normal.

Let  $N$  be a quasi-Galois extension of  $K$  containing  $L$ , let

$$\tilde{X} = \operatorname{Spec}(\tilde{B}),$$

where  $\tilde{B}$  is the integral closure of  $B$  in  $N$ , and let  $G = \operatorname{Aut}(N/K)$ . Since  $\tilde{X} \rightarrow X$  is surjective, it is enough to show that

$$\tilde{\pi} : \tilde{X} \rightarrow Y$$

is open. Let  $U$  be an open subset of  $\tilde{X}$ , and set

$$U' = \bigcup_{g \in G} gU.$$

Because  $G$  acts transitively on the fibers of  $\tilde{X} \rightarrow Y$  (cf. [BAC], Chap. V, §2.3, Prop. 6), one has

$$\tilde{\pi}(U) = \tilde{\pi}(U'),$$

and the latter is the open complement of the closed subset

$$\tilde{\pi}(\tilde{X} - U').$$

This proves (i).

(ii) We may suppose that  $Y$  and  $X$  are reduced. By Zariski's Main Theorem (cf. EGA IV<sub>3</sub>, 8.12.9), there exist affine open neighborhoods  $U$  of  $x$  and  $V = \operatorname{Spec}(A)$  of  $y$ , such that  $f(U) \subset V$ , and a factorization

$$\begin{array}{ccc} U & \xhookrightarrow{j} & V' \\ & \searrow f & \downarrow u \\ & & V \end{array},$$

where  $j$  is an open immersion and  $u$  is finite. Replacing  $V'$  by the closure of  $j(U)$ , we may suppose that  $V'$  is irreducible, hence that  $V' = \operatorname{Spec}(B)$ , where  $B$  is an integral  $A$ -algebra, finite over  $A$ . Moreover, since  $f$  is dominant, so is  $u$ , and the morphism  $A \rightarrow B$  is therefore injective.

Since  $A$  is unibranch at  $y$ , it follows from (i) that  $u$  is open at  $j(x)$ , and hence that

$$f = u \circ j$$

is open at  $x$ . This proves the first assertion of (ii). The second follows because, if  $Y$  is a local prescheme with closed point  $y$ , every open subset containing  $y$  equals  $Y$ . In the case  $Y = \operatorname{Spec}(\mathcal{O}_{Y,y})$ , one may also use, instead of EGA IV<sub>3</sub>, 8.12.9, the local form of Zariski's Main Theorem found, for example, in [Pes66] or [Ray70b], Chap. IV, Th. 1.



**Lemma 5.6.1.0.**—<sup>39</sup> Let  $f : X \rightarrow S$  be a morphism locally of finite presentation, let  $s \in S$ , and let  $x \in X_s$ . Set  $n = \dim_x(X_s)$ , and let  $q : \mathbb{A}_S^n \rightarrow S$  be the structure morphism.

Suppose that a quasi-finite morphism  $u : X \rightarrow \mathbb{A}_S^n$  is given such that  $f = q \circ u$ , and suppose that  $f$  is universally open at the generic point  $z$  of an irreducible component  $\mathfrak{z}$  of  $X_s$ , containing  $x$  and of dimension  $n$ . Then  $u$  is universally open at the point  $x \in X$ .

First observe that

(†) the hypotheses are preserved under every base change  $\pi : S' \rightarrow S$  covering  $s$

(that is, such that  $\pi^{-1}(s) \neq \emptyset$ ). Indeed, let  $S' \rightarrow S$  be such a morphism, let

$$\begin{array}{ccc} X' & \xrightarrow{u'} & \mathbb{A}_{S'}^n \\ & \searrow f' & \downarrow q' \\ & & S' \end{array}$$

be the diagram obtained by base change, and let  $s'$  be a point of  $S'$  above  $s$  and  $x'$  a point of  $X'_{s'}$  above  $x$ . Since

$$X'_{s'} = X_s \otimes_{\kappa(s)} \kappa(s'),$$

the lifting of generizations and the invariance of dimension under field extension (EGA IV<sub>2</sub>, 2.3.4(i) and 4.1.4) show that  $x'$  is contained in an irreducible component  $\mathfrak{z}'$  of  $X'_{s'}$  whose generic point  $z'$  lies above  $z$ , and

$$n \leq \dim \mathfrak{z}' \leq \dim_{x'} X'_{s'} \leq n.$$

Thus

$$\dim \mathfrak{z}' = n = \dim_{x'} X'_{s'}.$$

Since  $f$  is universally open at  $z$ ,  $f'$  is universally open at  $z'$ .<sup>1</sup>

Set  $Y = \mathbb{A}_S^n$ . By EGA IV<sub>3</sub>, 14.3.3.1(i), to prove that  $u$  is universally open at  $x$ , it is enough to prove that, for every integer  $r \geq 0$  and every point  $x'$  of

$$X' = X[T_1, \dots, T_r]$$

above  $x$ , the morphism

$$u' : X' \longrightarrow Y' = \mathbb{A}_S^n[T_1, \dots, T_r]$$

is open at  $x'$ . Now set

$$S' = S[T_1, \dots, T_r],$$

and let  $q'$  be the projection

$$Y' \simeq \mathbb{A}_{S'}^n \longrightarrow S'.$$

This is the situation obtained by the base change  $S' \rightarrow S$ . Thus, by the preceding observation, we are reduced to showing that  $u$  is open at  $x$ .

Set  $y = u(x)$ . Since  $u$  is locally of finite presentation (because  $f$  and  $q$  are; cf. EGA IV<sub>1</sub>, 1.4.3), EGA IV<sub>1</sub>, 1.10.3 shows that it is enough to prove

$$u(\operatorname{Spec} \mathcal{O}_{X,x}) = \operatorname{Spec} \mathcal{O}_{Y,y}.$$

<sup>39</sup>Editor's note: we have made explicit this lemma, which is used in the proof of Lemma 5.6.1.

<sup>1</sup>Translator's note: the French source prints  $x$  and  $x'$ , respectively, in this sentence. The stated hypothesis and the immediately preceding choice of generic points require  $z$  and  $z'$ .

For this purpose we may assume that  $S$  is affine and integral. Let  $\pi : S' \rightarrow S$  be its normalization, and denote by  $u' : X' \rightarrow Y'$  the morphism obtained from  $u$  by base change. Since

$$\pi_Y : Y' \longrightarrow Y$$

is integral and surjective, one has

$$\mathrm{Spec} \mathcal{O}_{Y,y} = \bigcup_{y'} \pi_Y(\mathrm{Spec} \mathcal{O}_{Y',y'}),$$

where the union ranges over all points  $y'$  of  $Y'$  above  $y$ . It is therefore enough to show that, for every such  $y'$  and every  $x' \in X'_{y'}$  above  $x$ ,

$$u'(\mathrm{Spec} \mathcal{O}_{X',x'}) = \mathrm{Spec} \mathcal{O}_{Y',y'}.$$

Since the hypotheses are preserved under the base change  $\pi : S' \rightarrow S$ , we are thereby reduced to the case of  $S'$ ; that is, we may assume that  $S$  is integral and normal.

The hypotheses on  $f$  now imply, by EGA IV<sub>3</sub>, 14.3.13, that there is an irreducible component  $Z$  of  $X$ , containing  $\mathfrak{z}$  (and hence  $x$ ), dominating  $S$ , and such that

$$\dim_z(Z_s) = n = \dim(Z_\eta),$$

where  $\eta$  is the generic point of  $S$ . Let  $\xi$  be the generic point of  $Z$ . Since  $u$ , and hence also

$$u_\eta : Z_\eta \longrightarrow \mathbb{A}_\eta^n,$$

is quasi-finite, the closure in  $\mathbb{A}_\eta^n$  of the point

$$u_\eta(\xi) = u(\xi)$$

has dimension  $n$ . Thus  $u(\xi)$  is the generic point of  $\mathbb{A}_\eta^n$ , which is also the generic point of  $Y = \mathbb{A}_S^n$ . Consequently, if  $g$  denotes the restriction of  $u$  to  $Z$ , then

$$g : Z \longrightarrow Y$$

is quasi-finite and dominant. Since  $Y$  is normal, Lemma 5.6.0 shows that  $g$  is open. Hence  $u$  is open at every point of  $Z$ , and in particular at  $x$ . Consequently,

$$u(\mathrm{Spec} \mathcal{O}_{X,x}) = \mathrm{Spec} \mathcal{O}_{Y,y},$$

which completes the proof of Lemma 5.6.1.0.

The following lemma advantageously replaces EGA IV<sub>3</sub>, 14.5.10,<sup>40</sup> in that it is independent of noetherian hypotheses.

**Lemma 5.6.1.**— *Let  $S$  be a scheme, let  $f : X \rightarrow S$  be an  $S$ -scheme locally of finite presentation, let  $s$  be a point of  $S$ , and let  $x$  be a closed point of  $X_s$ . Suppose that  $f$  is universally open at the generic point  $z$  of an irreducible component  $Z$  of  $X_s$  containing  $x$ , and that*

$$\dim_x(X_s) = \dim Z.$$

*Then there exists a commutative diagram*

<sup>40</sup>Editor's note: we have corrected the original, which indicated 19.5.10.

$$\begin{array}{ccc}
X & \xrightarrow{f} & S \\
\uparrow h & \nearrow \phi & \uparrow w \\
S'' & \xrightarrow{\pi} & S'
\end{array} ,$$

where  $S'$  is affine,  $w$  is étale,  $\pi$  is finite, surjective, and of finite presentation, and  $\phi^{-1}(s)$  consists of a single point  $s''$ , such that  $h(s'') = x$  and  $\kappa(s'') = \kappa(x)$ .<sup>41</sup>

*Proof.* First, we may suppose that  $S = \operatorname{Spec}(A)$  and  $X = \operatorname{Spec}(B)$ , where  $B$  is an  $A$ -algebra of finite presentation. Let  $\mathfrak{p}$  and  $\mathfrak{q}$  be the prime ideals of  $A$  and  $B$  corresponding respectively to  $s$  and  $x$ , so that

$$\mathfrak{p} = A \cap \mathfrak{q}.$$

Set  $n = \dim_x(X_s)$ , and choose  $t_1, \dots, t_n \in B$  whose images in

$$\mathcal{O}_{X_s, x} = B_{\mathfrak{q}} / \mathfrak{p}B_{\mathfrak{q}}$$

form a system of parameters. Then

$$\mathcal{O}_{X_s, x} / (t_1, \dots, t_n)$$

is finite-dimensional over  $\kappa(x)$ , and hence also over  $\kappa(s)$ , because  $x$  is a closed point of the algebraic  $\kappa(s)$ -scheme  $X_s$ . Consequently, the  $S$ -morphism

$$u : X \longrightarrow \mathbb{A}_S^n = \operatorname{Spec}(A[T_1, \dots, T_n]), \quad T_i \longmapsto t_i,$$

is of finite presentation,  $x$  is isolated in the fiber  $u^{-1}(u(x))$ , and

$$u(x) = \tau_0(s),$$

where  $\tau_0 : S \rightarrow \mathbb{A}_S^n$  denotes the zero section of  $\mathbb{A}_S^n \rightarrow S$ , corresponding to the  $A$ -algebra homomorphism

$$A[T_1, \dots, T_n] \longrightarrow A, \quad T_i \longmapsto 0.$$

The set of points of  $X$  that are isolated in their fibers over  $\mathbb{A}_S^n$  is open (EGA IV<sub>3</sub>, 13.1.4). Thus, after shrinking  $X$ , we may suppose that  $u$  is quasi-finite and that  $u^{-1}(u(x)) = \{x\}$ . By Lemma 5.6.1.0,  $u$  is universally open at  $x$ .

Let

$$B_0 = B / (t_1, \dots, t_n), \quad X_0 = X \times_{\mathbb{A}_S^n} \tau_0(S) = \operatorname{Spec}(B_0),$$

and let  $u_0 : X_0 \rightarrow S$  be obtained from  $u$  by the base change  $\tau_0 : S \hookrightarrow \mathbb{A}_S^n$ . Then  $u_0$  is quasi-finite and of finite presentation, is universally open at  $x$ , and  $x$  is the unique point of  $X_0$  above  $s$ .

Let  $A'$  be the henselization of the local ring

$$A_{\mathfrak{p}} = \mathcal{O}_{S, s},$$

and set

$$S' = \operatorname{Spec}(A'), \quad B'_0 = B_0 \otimes_A A', \quad X'_0 = X_0 \times_S S' = \operatorname{Spec}(B'_0).$$

The closed point  $s'$  of  $S'$  is the unique point of  $S'$  above  $s$ , and  $\kappa(s') = \kappa(s)$ . The scheme  $X'_0$  has a unique point  $x'$  above  $s'$ ; moreover  $\kappa(x') = \kappa(x)$ , and  $x'$  is also the unique point of  $X'_0$

<sup>41</sup>Editor's note: consequently,  $h$  induces a section  $\sigma$  of  $X \times_S S'' \rightarrow S''$  such that  $\sigma(s'')$  lies above  $x$ ; compare EGA IV<sub>3</sub>, 14.5.10.

above both  $s$  and  $x$ . Since  $A'$  is henselian, EGA IV<sub>4</sub>, 18.5.11 gives a disjoint decomposition into open and closed parts

$$X'_0 = V \amalg W, \quad V = \operatorname{Spec}(\mathcal{O}_{X'_0, x'}). \quad (*)$$

The local ring  $\mathcal{O}_{X'_0, x'}$  is finite and of finite presentation over  $A'$ . Hence the restriction  $\pi$  of  $u'_0$  to  $V$  is finite and of finite presentation. Moreover, because  $u'_0 : X'_0 \rightarrow S'$  is open at  $x'$ ,

$$u'_0(V) = \operatorname{Spec}(\mathcal{O}_{S', s'}) = S',$$

so  $\pi$  is surjective.

This proves the assertion when  $S = S'$ . In general,  $A'$  is the filtered direct limit of  $A$ -subalgebras  $A_i$  that are étale over  $A$  and for which

$$S_i = \operatorname{Spec}(A_i)$$

has a unique point  $s_i$  above  $s$ , with  $\kappa(s_i) = \kappa(s)$ . We have

$$B'_0 = \varinjlim_i B_i, \quad B_i = B_0 \otimes_A A_i.$$

Set

$$X_i = \operatorname{Spec}(B_i) = X_0 \times_S S_i, \quad C = \mathcal{O}_{X'_0, x'}.$$

By  $(*)$ , one has

$$C \simeq B'_0 / f B'_0$$

for some idempotent  $f \in B'_0$ . There are an index  $i$  and an element  $f_i \in B_i$  whose image in  $B'_0$  is  $f$ . Put

$$C_i = B_i / f_i B_i, \quad V_i = \operatorname{Spec}(C_i).$$

Then

$$C = C_i \otimes_{A_i} A', \quad V = V_i \times_{S_i} S',$$

and  $C_i$  is an  $A_i$ -algebra of finite presentation, since the same is true of  $B_i$ . Consequently, the morphisms

$$\begin{array}{ccc} X'_0 & & \\ \uparrow h' & \searrow u'_0 & \\ V & \xrightarrow{\pi} & S' \end{array}$$

arise, by the base change  $S' \rightarrow S_i$ , from morphisms of finite presentation

$$\begin{array}{ccc} X_i & & \\ \uparrow h_i & \searrow u_i & \\ V_i & \xrightarrow{\pi_i} & S_i \end{array}$$

For every  $j \geq i$ , let

$$V_j = V_i \times_{S_i} S_j$$

and let  $\pi_j : V_j \rightarrow S_j$  be the morphism of finite presentation obtained from  $\pi_i$  by base change. Since  $\pi : V \rightarrow S'$  is finite and surjective, EGA IV<sub>2</sub>, 8.10.5 shows that, for some  $j$ , the morphism  $\pi_j : V_j \rightarrow S_j$  is finite and surjective. Then  $w : S_j \rightarrow S$  is affine and étale,  $S_j$  has a unique point  $s_j$  above  $s$ , and  $V_j$  has a unique point  $x_j$  above  $s_j$ , since  $x'$  is the unique point of

$$V = V_j \times_{S_j} S'$$

above  $s'$ :

$$\begin{array}{ccccccc}
X_j & \longrightarrow & X_0 & \longrightarrow & X & & \\
\uparrow h_j & & \searrow u_j & & \searrow u_0 & \searrow u & \\
V_j & \xrightarrow{\pi_j} & S_j & \xrightarrow{w} & S & \xrightarrow{\tau_0} & \mathbb{A}_S^n
\end{array}$$

Thus  $x_j$  is the unique point of  $V_j$  above  $s$ , its image under  $V_j \rightarrow X_j \rightarrow X$  is  $x$ , and

$$\kappa(x_j) = \kappa(x).$$

**Lemma 5.6.2.0.**—<sup>42</sup> *Let  $k$  be a field and let  $G$  be a nonempty  $k$ -scheme of finite type.*

- (i) *Let  $K$  be an extension of  $k$ , and let  $W$  be a dense open subset of  $G_K$ . Denote by  $\pi$  the projection  $G_K \rightarrow G$ . Then*

$$U = \{g \in G \mid W \text{ contains every maximal point of } \pi^{-1}(g)\}$$

*is a dense open subset of  $G$ . (N.B. If  $g$  is a closed point of  $G$  belonging to  $U$ , then  $\pi^{-1}(g) \subset W$ .)*

- (ii) *Suppose in addition that  $G$  is geometrically irreducible. Let*

$$\mu : G \times X \longrightarrow Y$$

*be a morphism of  $k$ -schemes, let  $x$  be a point of  $X$ , and let  $\Omega$  be an open subset of  $Y$  such that*

$$\mu_x^{-1}(\Omega_{\kappa(x)}) \neq \emptyset,$$

*where  $\mu_x$  denotes the morphism*

$$G_{\kappa(x)} \longrightarrow Y_{\kappa(x)}, \quad g \longmapsto \mu(g, x).$$

*Then*

$$U = \{g \in G \mid \mu \text{ sends every maximal point of } g \times x \text{ into } \Omega\}$$

*is a dense open subset of  $G$ . For every closed point  $g$  of  $G$  belonging to  $U$ , one has*

$$\mu(g \times x) \subset \Omega,$$

*and hence*

$$\mu(g', x) \in \Omega_{\kappa(x)} \quad (\text{respectively, } \mu(g, x') \in \Omega_{\kappa(g)}),$$

*for every point  $g' \in G_{\kappa(x)}$  above  $g$  (respectively,  $x' \in X_{\kappa(g)}$  above  $x$ ).*

- (iii) *Suppose that  $G = H^0$ , where  $H$  is a  $k$ -group scheme locally of finite type, acting on a nonempty  $k$ -scheme  $X$  in such a way that the morphism*

$$H \times_S X \longrightarrow X \times_S X, \quad (h, x) \longmapsto (hx, x)$$

*is surjective. Let  $U$  be an open subset of  $X$ . Then:*

- (a)  *$G \cdot U$  is an open subset of  $X$ , equal to the union of the irreducible components of  $X$  whose generic point belongs to  $U$ .*

<sup>42</sup>Editor's note: we have added this lemma, communicated by O. Gabber. It makes it possible to simplify the proof of 5.6.2 and to prove Theorem 5.3, as well as 5.4, in a more general form; see 5.7 and 5.8 below.

- (b) Suppose that  $U$  is dense in every irreducible component of  $X$ . Then, for every finite subset  $A$  of  $X$ , there exists a closed point  $g \in G$  such that

$$g \cdot A' \subset U_{\kappa(g)},$$

where  $A'$  is the inverse image of  $A$  in  $X_{\kappa(g)}$ . In particular, the morphism

$$G \times U \longrightarrow X$$

is surjective.

*Proof.* (i) Let  $\bar{k}$  be a separable closure of  $k$ , and let  $L$  be an extension of  $k$  containing a copy of  $\bar{k}$  and of  $K$ . Denote by  $\pi_L$  (respectively,  $\phi$ ) the projection

$$G_L \longrightarrow G \quad (\text{respectively, } G_L \longrightarrow G_K).$$

For every  $g \in G$ , the map  $\phi$  sends the maximal points of  $\pi_L^{-1}(g)$  surjectively onto those of  $\pi^{-1}(g)$ . Consequently,

$$U = \{g \in G \mid W_L \text{ contains every maximal point of } \pi_L^{-1}(g)\}.$$

Thus, replacing  $K$  by  $L$ , we are reduced to the case in which  $K$  contains  $\bar{k}$ .

Since the projection

$$p : G_K \longrightarrow G_{\bar{k}}$$

is surjective and open,  $V = p(W)$  is a dense open subset of  $G_{\bar{k}}$ . Moreover,

$$p^{-1}(g) = \text{Spec}(\kappa(g) \otimes_{\bar{k}} K)$$

is irreducible for every  $g \in G_{\bar{k}}$  (cf. EGA IV<sub>2</sub>, 4.3.3). Hence, for every  $g \in V$ , the generic point of  $p^{-1}(g)$  belongs to  $W$ .

On the other hand, set

$$\mathcal{G} = \text{Gal}(\bar{k}/k).$$

The projection  $q : G_{\bar{k}} \rightarrow G$  is surjective, and  $\mathcal{G}$  acts transitively on the fibers of  $q$ . Therefore

$$U = q(V'),$$

where  $V'$  is the intersection of the  $\mathcal{G}$ -conjugates of  $V$ .

Let  $Z$  be the closed subset  $G_{\bar{k}} - V$ , endowed with its reduced closed subscheme structure. Since  $G_{\bar{k}}$ , and hence  $Z$ , is of finite presentation over  $\bar{k}$ , the scheme  $Z$  is obtained by base change from a reduced closed subscheme

$$Z_1 \subset G \otimes_k k_1$$

for some finite Galois extension  $k_1$  of  $k$ . Thus the  $\mathcal{G}$ -conjugates of  $Z$  are finite in number, and their union is again a nowhere-dense closed subset  $F$  of  $G_{\bar{k}}$ . It follows that

$$V' = G_{\bar{k}} - F$$

is a dense open subset of  $G_{\bar{k}}$ . Since  $q : G_{\bar{k}} \rightarrow G$  is surjective and open,  $U = q(V')$  is therefore a dense open subset of  $G$ .

Moreover, for every closed point  $g$  of  $G$ , the fiber  $\pi^{-1}(g)$  consists of finitely many closed points of  $G_K$ . Thus, if  $g \in U$ , then

$$\pi^{-1}(g) \subset W.$$

This proves (i), and (ii) follows from it.

Assertion (iii)(a) was proved in VI<sub>A</sub>, 2.6.4. Finally, if  $U$  is dense in every irreducible component of  $X$ , then

$$X = G \cdot U.$$

Hence, for every  $x \in X$ ,

$$\mu_x^{-1}(U_{\kappa(x)})$$

is a nonempty open subset of  $G_{\kappa(x)}$ , and (iii)(b) follows from (ii).

**Corollary 5.6.2.**— *Let  $S$ ,  $G$ , and  $X$  be as in the preliminary hypotheses of 5.3, and let  $U$  be an open subset of  $X$ , let  $s \in S$ , and let  $A$  be a finite subset of  $X_s$ . Suppose that  $U_s$  is dense in  $X_s$  and that  $X_s$  is locally of finite type over  $\kappa(s)$ .<sup>43</sup>*

*Then there are morphisms  $f : S'' \rightarrow S$  and  $h : S'' \rightarrow G$ , where  $f$  factors as a finite surjective morphism  $S'' \rightarrow S'$  followed by an étale morphism  $S' \rightarrow S$ , such that the inverse image  $A''$  of  $A$  in  $X \times_S S''$  (i.e., in  $X_s \times_{\text{Spec } \kappa(s)} S''_s$ ) is contained in*

$$\ell_h^{-1}(U_{S''}),$$

*where  $\ell_h$  denotes the translation of  $X_{S''} = X \times_S S''$  defined by the element  $h \in G(S'')$ .*

*Proof.* Since  $X_s$  is locally of finite type over  $\kappa(s)$ , the connected components of  $X_s$  are open and irreducible (cf. VI<sub>A</sub>, 2.5.4). Hence  $U_s$  is dense in every irreducible component of  $X_s$ .

By Lemma 5.6.2.0, there therefore exists a closed point

$$g \in G_s^0$$

such that

$$g \cdot a' \in U_s \otimes_{\kappa(s)} \kappa(g)$$

for every  $a' \in X_{\kappa(g)}$  lying above a point of  $A$ .

By Lemma 5.6.1, there are morphisms  $f : S'' \rightarrow S$  and  $h : S'' \rightarrow G$ , where  $f$  factors as a finite surjective morphism  $S'' \rightarrow S'$  followed by an étale morphism  $S' \rightarrow S$ , such that  $f^{-1}(s)$  consists of a single point  $s_0$ , and such that

$$h(s_0) = g, \quad \kappa(s_0) = \kappa(g).$$

Denote by  $A''$  the inverse image of  $A$  in

$$X_s \times_{\text{Spec } \kappa(s)} S''_s = X_s \otimes_{\kappa(s)} \kappa(s_0) = X_s \otimes_{\kappa(s)} \kappa(g).$$

Then the translation  $\ell_h$  of  $X_{S''}$  sends  $A''$  into

$$U_s \otimes_{\kappa(s)} \kappa(s_0).$$

**Lemma 5.6.3.**— *Let  $X$  be an  $S$ -scheme. The following conditions are equivalent:*

- (i)  *$X$  is separated over  $S$ .*
- (ii) *For every  $S$ -scheme  $T$ , every section  $\sigma : T \rightarrow X_T$  is a closed immersion.*
- (iii) *For every reduced  $S$ -scheme  $T$ , two  $S$ -morphisms  $f_1, f_2 : T \rightarrow X$  that agree on a dense open subset  $U$  of  $T$  are equal.*

<sup>43</sup>Editor's note: we have added the hypothesis on  $X_s$  and simplified (and corrected) the proof, taking account of the addition of 5.6.2.0. Moreover, the proof shows that the conclusion remains valid if one assumes only that  $U_s$  is dense in every irreducible component of  $X_s$ .

(iv) For every  $s \in S$  and every pair of points  $x_1, x_2$  of  $X_s$ , there exist a morphism

$$\phi : S'' \longrightarrow S' \longrightarrow S$$

and an open subscheme  $V$  of  $X_{S''}$ , separated over  $S''$ , such that:

- (a)  $S' \rightarrow S$  is open,  $S'' \rightarrow S'$  is closed and surjective, and  $\phi^{-1}(s) \neq \emptyset$ ;
- (b) the inverse image of  $\{x_1, x_2\}$  in  $X_{S''}$  is contained in  $V$ .

(iv') For every  $s \in S$ , every pair of points  $x_1, x_2 \in X_s$  is contained in an open subset  $V$  of  $X$  that is separated over  $S$ .<sup>44</sup>

*Proof.* The implication (ii) $\Rightarrow$ (i) is clear (take  $T = X$  and let  $\sigma$  be the diagonal section), as are

$$(i) \Rightarrow (iv') \Rightarrow (iv).$$

On the other hand, (i) $\Rightarrow$ (ii) by EGA I, 5.4.6.

(iii) $\Rightarrow$ (ii). Let  $\sigma : T \rightarrow X_T$  be a section of  $p : X_T \rightarrow T$ . By EGA I, 5.3.13,  $\sigma$  is an immersion, that is, an isomorphism of  $T$  onto a locally closed subscheme  $E$  of  $X_T$ . To show that  $E$  is closed, we may suppose that  $T$  and  $E$  are reduced. Let  $\bar{E}$  be the reduced closed subscheme of  $X_T$  whose underlying space is the closure of  $E$ , so that  $E$  is a dense open subscheme of  $\bar{E}$ . The immersion

$$i : \bar{E} \hookrightarrow X_T$$

and  $\sigma \circ p \circ i$  agree on  $E$ , hence on  $\bar{E}$  by hypothesis (iii). Thus every point of  $\bar{E}$  belongs to  $\sigma(T) = E$ , and consequently  $\bar{E} = E$ .

(i) $\Rightarrow$ (iii). Suppose that  $X$  is separated over  $S$ , and let  $T$  be a reduced  $S$ -scheme and  $f_1, f_2 : T \rightarrow X$  two  $S$ -morphisms that agree on a dense open subset  $U$  of  $T$ . Let

$$g : T \longrightarrow X \times_S X$$

be the morphism with components  $f_1$  and  $f_2$ . Since

$$D = \Delta_{X/S}(X)$$

is closed, its inverse image under  $g$  is a closed subset of  $T$  containing  $U$ , and is therefore equal to  $T$ . Since  $T$  is reduced,  $g$  factors through  $D$  (cf. EGA I, 5.2.2); consequently

$$f_1 = p_1 \circ g = p_2 \circ g = f_2.$$

(iv) $\Rightarrow$ (iii). Let  $T$  be a reduced  $S$ -scheme, and let  $f_1, f_2 : T \rightarrow X$  be two  $S$ -morphisms that agree on a dense open subset  $U$ . Since  $T$  is reduced, in order to prove that  $f_1 = f_2$ , it is enough to prove that  $f_1 = f_2$  set-theoretically. Indeed, suppose this has been established, let  $t \in T$ , let  $V$  be an affine open subset of  $X$  containing  $f_1(t) = f_2(t)$ , and let  $W$  be the

<sup>44</sup>Editor's note: we have simplified the formulation of condition (iv) and added condition (iv'). We have also added the proof of the implication (i) $\Rightarrow$ (iii), which is used in the proof of (iv) $\Rightarrow$ (iii). Observe moreover that, if

$$T = \operatorname{Spec} B, \quad B = k[\varepsilon, x]/(\varepsilon^2, \varepsilon x)$$

( $k$  a field) and

$$X = T_{\text{red}} = \operatorname{Spec} k[x],$$

then the morphisms

$$\phi_\lambda : T \longrightarrow X, \quad x \longmapsto x + \lambda \varepsilon \quad (\lambda \in k),$$

agree on the dense open subset  $\operatorname{Spec} B_x$ , but are not equal.



open neighborhood of  $t$  that is the inverse image of  $V$  under the continuous map underlying both  $f_1$  and  $f_2$ . Then the morphisms

$$f_i|_W : W \longrightarrow V$$

agree on the dense open subset  $U \cap W$  of  $W$ . Since  $V$  is separated and  $W$  is reduced, this implies that  $f_1|_W = f_2|_W$ , and hence that  $f_1 = f_2$ .

Now let  $t \in T$ , let  $s$  be its image in  $S$ , and let us show that the points

$$x_1 = f_1(t), \quad x_2 = f_2(t)$$

of  $X_s$  are equal. Let

$$\phi : S'' \longrightarrow S' \longrightarrow S$$

and an open subset  $V$  of  $X \times_S S''$  be as in (iv). Set

$$T' = T \times_S S', \quad T'' = T \times_S S'',$$

and denote by

$$g : T'' \longrightarrow T, \quad f_i'' : T'' \longrightarrow X_{S''} \quad (i = 1, 2)$$

the morphisms obtained by base change.

Since  $U$  is dense in  $T$  and  $T' \rightarrow T$  is open, the inverse image  $U'$  of  $U$  in  $T'$  is dense in  $T'$ . Let  $U''$  be the inverse image of  $U'$  in  $T''$ , and let  $F$  be the reduced subscheme of  $T''$  whose underlying space is  $\overline{U''}$ . Since  $T'' \rightarrow T'$  is surjective and closed, the image of  $F$  contains  $U'$  and is closed, and is therefore equal to  $T'$ . Consequently,

$$F \cap g^{-1}(t)$$

contains a point  $u$ .

For  $i = 1, 2$ , let  $h_i$  be the restriction of  $f_i''$  to  $F$ . Then  $h_i(u)$  is a point of  $X_{S''}$  above  $f_i(t) = x_i$ , and hence belongs to  $V$ , since  $V$  contains the inverse image of  $\{x_1, x_2\}$  in  $X_{S''}$ .

Thus

$$W = h_1^{-1}(V) \cap h_2^{-1}(V)$$

is an open subset of  $F$  containing  $u$ , and the  $S''$ -morphisms

$$h_i|_W : W \longrightarrow V$$

agree on the dense open subset  $U'' \cap W$  of  $W$ . Since  $V$  is separated over  $S''$ , we have

$$h_1|_W = h_2|_W.$$

It follows that  $h_1(u) = h_2(u)$ , and hence that  $f_1(t) = f_2(t)$ . This proves (iv)  $\Rightarrow$  (iii).

Theorem 5.3 now follows from 5.6.2 and the implication (iv)  $\Rightarrow$  (i) of 5.6.3.

**Counterexample 5.6.4.**— *Not every  $S$ -group  $G$  is separated. Let  $S$  be a scheme having a non-isolated closed point  $s$ , and let  $G$  be the scheme obtained by gluing two copies of  $S$  along the open subset  $S - \{s\}$ . It is easily seen that  $G$  is not separated over  $S$ , and that it carries a natural  $S$ -group structure whose fibers are all trivial, except for the fiber  $G_s$ , which is isomorphic to the two-element group  $\mathbb{Z}/2\mathbb{Z}$ .<sup>45</sup>*

<sup>45</sup>Editor's note: we reproduced this example in VI<sub>A</sub>, 0.3, editor's note 5.

**Theorem 5.7.**—<sup>46</sup> *Let  $S$  be a scheme, let  $G$  be an  $S$ -group locally of finite presentation over  $S$ , such that the function*

$$s \longmapsto \dim G_s$$

*is locally constant on  $S$ , let  $X$  be an  $S$ -scheme on which  $G$  acts, and let  $U$  be an open subset of  $X$ , separated over  $S$ . Then  $\underline{G}^0 \cdot U$  is an open subset of  $X$ , separated over  $S$ .*

*Proof.* Let

$$\mu : G \times_S X \longrightarrow X$$

denote the action of  $G$  on  $X$ . It is the composite of the automorphism

$$(g, x) \longmapsto (g, gx)$$

of  $G \times_S X$  and the projection onto  $X$ . Since  $G \times_S U$  is an open subset of  $G \times_S X$ , it follows from 4.2(iii) that

$$V = \underline{G}^0 \cdot U$$

is open in  $X$ .

Then, arguing as in the proof of 5.6.2, we deduce from the implication (iv) $\Rightarrow$ (i) of 5.6.3 that  $V$  is separated over  $S$ .

**Corollary 5.7.1.**— *Let  $S$  and  $G$  be as in 5.7, and let  $\sigma, \tau$  be two  $S$ -sections of  $G^0$ , that is,  $\sigma, \tau \in G^0(S)$ . Then the coincidence subscheme  $S(\sigma, \tau) \subset S$  of  $\sigma$  and  $\tau$  (that is, the inverse image of the diagonal of  $G \times_S G$  under the morphism  $(\sigma, \tau)$ ) is closed.*

Indeed, for every  $s \in S$ , let  $U$  be an affine open subset of  $G$  containing  $\epsilon(s)$ . Then

$$V = \epsilon^{-1}(U)$$

is an open neighborhood of  $s$  in  $S$ , and, since  $G^0 U$  is separated,  $S(\sigma, \tau) \cap V$  is closed in  $V$ .

**Remark 5.7.2.**— Gabber points out that one can show the following: if  $S$  is henselian local, with closed point  $s$ , then the intersection of the open subsets  $G^0 U$ , where  $U$  ranges over the affine open neighborhoods of  $\epsilon(s)$ , is an open group subscheme of  $G$ , separated over  $S$ .

## 5.8. Complements

<sup>47</sup>We begin by recalling Proposition VI<sub>A</sub>, 2.6.6:

**Proposition 5.8.1.**— *Let  $k$  be a field, let  $G$  be a  $k$ -group locally of finite type acting on a  $k$ -scheme  $X$  in such a way that the morphism*

$$G \times X \longrightarrow X \times X, \quad (g, x) \longmapsto (gx, x),$$

*is surjective. Suppose that  $X$  is quasi-separated. Then the connected components of  $X$  are irreducible.*

---

<sup>46</sup>Editor's note: on the one hand, we omitted Corollary 5.6.5, which repeated 5.5. On the other hand, the original stated Corollary 5.7.1 below as Remark 5.7, referring for its proof to 4.7, a number that does not exist in *Lecture Notes in Mathematics* 151 (but did appear in the 1965 edition of SGAD, whose numbers 4.5 and 4.6 became 5.6.1 and 5.6.2); we have added Theorem 5.7, communicated by O. Gabber, which makes the aforementioned statement of SGAD more precise.

<sup>47</sup>Editor's note: we have added this paragraph of complements and counterexamples, all communicated by O. Gabber.

**Corollary 5.8.2.**— *Let  $S, G, X$  be as in the preliminary hypotheses of 5.3. Suppose moreover that every fiber  $X_s$  is quasi-separated and connected. Then  $X$  is separated and quasi-compact over  $S$ .*

Indeed, by the preceding proposition, every fiber  $X_s$  is irreducible, and the rest of the proof is identical to that of 5.4.

**Example 5.8.3.**— Fix an algebraically closed field  $k$ . Recall first that every “locally Boolean” topological space  $X$ , that is, every space having a basis of compact open subsets, becomes a  $k$ -scheme when equipped with the sheaf of locally constant  $k$ -valued functions (cf. [DG70], I, §1, 2.12). Such an  $X$  will be called a *locally Boolean  $k$ -scheme*.

We also note that every topological space  $E$  having a basis of separated open subsets—and likewise every scheme  $X$ —has a dense separated open subset. Indeed, every increasing union of separated open subsets is separated (for a scheme  $X$ , this follows from 5.6.3), so there is a maximal such open subset  $U$ . It must be dense: otherwise there would be a nonempty open subset  $V$  with  $U \cap V = \emptyset$ ; the open set  $V$  would contain a nonempty separated open subset  $W$ , and

$$U \cup W$$

would still be separated, contradicting the maximality of  $U$ .

Now let  $C$  be the Cantor space, viewed as the underlying space of the group

$$(\mathbb{Z}/2\mathbb{Z})^{\mathbb{N}}$$

with its product topology. For every  $p \in C$ , let  $C(p)$  be another copy of  $C$ , and let  $X$  be obtained by gluing each  $C(p)$  to  $C$  along  $C - \{p\}$ . Then  $X$  is a nonseparated locally Boolean  $k$ -scheme, and  $C$  is a dense open subset of  $X$ .

Let  $G$  be the group of automorphisms of the  $k$ -scheme  $X$ , that is, of homeomorphisms of  $X$ . Then  $G$  acts transitively on  $X$ . Indeed,  $X$  is the union of  $C$  and, for each point  $p \in C$ , a second point  $p'$ . This second point is characterized as the unique  $x \in X - \{p\}$  for which

$$(C - \{p\}) \cup \{x\}$$

is separated. Consequently, every automorphism  $\phi$  of  $C$  extends to an automorphism  $\phi_X$  of  $X$  satisfying

$$\phi_X(p') = \phi_X(p)'$$

for every  $p$ . Moreover, the map

$$\theta_p : X \longrightarrow X$$

that exchanges  $p$  and  $p'$  and fixes every other point is an automorphism of  $X$ . Finally, the automorphism group of  $C$  acts transitively on  $C$ : using the group structure of  $C$ , it is enough to consider translations.

Thus the discrete  $k$ -group  $G$ , which is locally of finite type, acts on the  $k$ -scheme  $X$  in such a way that

$$G \times X \longrightarrow X \times X, \quad (g, x) \longmapsto (gx, x),$$

is surjective, but  $X$  is not separated, even though  $C$  is a dense separated open subset.

**Example 5.8.4.**— Retain the notation of the preceding example. Using the description of  $C$  as the set of sequences

$$(u_n)_{n \in \mathbb{N}}, \quad u_n \in \mathbb{Z}/2\mathbb{Z},$$

one sees that  $C$  with one point  $p$  removed is homeomorphic to a countable disjoint union of copies of  $C$ . Indeed, using the group structure of  $C$ , for example, translation reduces us to

the case in which  $p = 0$ , the zero sequence. Then

$$C - \{0\} = \coprod_{i \in \mathbb{N}} C_i,$$

where

$$C_i = \{(u_n)_{n \in \mathbb{N}} \mid u_i = 1 \text{ and } u_j = 0 \text{ for } j < i\},$$

and every  $C_i$  is homeomorphic to  $C$ . It follows by induction on  $|F|$  that, for every nonempty finite subset  $F$  of  $C$ , the space  $C - F$  is homeomorphic to a countable disjoint union of copies of  $C$ , and hence to  $C$  with one point removed.

For each  $F \subset C$  of cardinality 2, let  $C(F)$  be another copy of  $C$ , denote by  $q_F$  the point 0 of  $C(F)$ , and choose a homeomorphism

$$\phi_F : C(F) - \{q_F\} \xrightarrow{\sim} C - F.$$

Let  $X'$  be obtained by gluing every  $C(F)$  to  $C$  by means of  $\phi_F$ . Then  $X'$  is a nonseparated locally Boolean  $k$ -scheme. Moreover, the construction shows that every locally constant function

$$f : X' \longrightarrow k$$

is constant. Indeed, if  $x, y \in C$  and  $F = \{x, y\}$ , then every neighborhood of  $q_F$  meets every neighborhood of  $x$  and of  $y$ . Thus

$$f(x) = f(q_F) = f(y).$$

If  $F' = \{z, t\}$ , with  $z, t \in C$ , then likewise

$$f(q_{F'}) = f(z) = f(q_{\{z, x\}}) = f(x).$$

Consequently,  $X'$  is connected.

Every point  $x \in X'$  also has a pointed neighborhood homeomorphic to  $(C, 0)$ . More precisely, for every  $F$ , fix a homeomorphism of pointed topological spaces

$$h_F : (C, 0) \xrightarrow{\sim} (C(F), q_F),$$

and let  $T$  be the group of translations of  $C$ . If  $x = q_F$ , use  $h_F$ ; if  $x \in C$ , the translation  $t_x \in T$  is a homeomorphism

$$t_x : (C, 0) \xrightarrow{\sim} (C, x),$$

and it is the unique element of  $T$  with this property.<sup>1</sup>

Let  $L$  be the free group generated by the  $h_F$ , and set

$$G = T * L,$$

the free product (coproduct) of  $T$  and  $L$ . For

$$h \in \{h_F\}_{|F|=2} \cup T,$$

let  $\sigma(h)$  denote the corresponding generator of  $G$ , and let  $S(h)$  and  $B(h)$  be respectively the source and the target of  $h$ . It is convenient also to set

$$\sigma(h_F^{-1}) = \sigma(h_F)^{-1}, \quad S(h_F^{-1}) = B(h_F), \quad B(h_F^{-1}) = S(h_F),$$

---

<sup>1</sup>Translator's note: in these two passages the French source prints  $X$  in place of the  $X'$  just constructed, and  $q_F$  in place of the homeomorphism  $h_F$  just defined. The English uses  $X'$  and  $h_F$ , respectively.

and to let  $E$  be the set consisting of the elements of  $T$  and all  $h_F, h_F^{-1}$ .

On the product space

$$P = G \times X',$$

where  $G$  carries the discrete topology, consider the equivalence relation generated by

$$(g\sigma(h), x) \sim (g, h(x)) \quad \text{when } x \in S(h),$$

for every  $h \in E$ , and let  $Z$  be the quotient space. Thus  $Z$  is obtained by gluing open subsets of the disjoint union

$$\coprod_{g \in G} \{g\} \times X'.$$

Consequently, the saturation  $\tilde{\Omega}$  of every open subset  $\Omega$  of  $P$  is open (cf. [BTop], I, §5.1, Example 2). Explicitly, since every open subset of  $P$  is the union of its intersections with the slices  $\{g\} \times X'$ , it is enough to consider  $\{1\} \times W$ , with  $W$  open in  $X'$ . Its saturation is the union of  $\{1\} \times W$  and the sets

$$\begin{aligned} & \{\sigma(h_1)\} \times h_1^{-1}(W \cap B(h_1)), \\ & \{\sigma(h_1)\sigma(h_2)\} \times h_2^{-1}(h_1^{-1}(W \cap B(h_1)) \cap B(h_2)), \end{aligned}$$

and so on, for all finite sequences  $h_1, \dots, h_n$  of elements of  $E$ . It is therefore open, and the projection

$$\pi : P \longrightarrow Z$$

is open.

Moreover, the word

$$\sigma(h_1) \cdots \sigma(h_n)$$

is reduced in  $G$ , unless one of the  $h_i$  is the identity of  $T$ , two consecutive  $h_i$ 's belong to  $T$ , or one of the pairs  $(h_i, h_{i+1})$  is

$$(h_F, h_F^{-1}) \quad \text{or} \quad (h_F^{-1}, h_F).$$

Suppose, therefore, that  $x \in X'$  and that

$$\beta = (\sigma(h_1) \cdots \sigma(h_n), h_n^{-1} \cdots h_1^{-1}(x))$$

belongs to  $\{1\} \times X'$ . We may then suppose that every  $h_i$  is a translation  $t_i$ . In that case,

$$\sigma(t_1 \cdots t_n) = 1$$

implies  $t_1 \cdots t_n = 1$ , and hence  $\beta = (1, x)$ . The equivalence relation is compatible with the action of  $G$  on  $P = G \times X'$  by left translation on the first factor. It follows that, for every  $g \in G$ , the restriction of  $\pi$  to  $\{g\} \times X'$  is injective.

Now let  $z \in Z$  and choose a representative  $(g, x) \in P$ . By the preceding discussion,

$$U = \pi(\{g\} \times X')$$

is an open neighborhood of  $z$ , and the continuous map

$$\{g\} \times X' \longrightarrow U$$

induced by  $\pi$  is open and bijective, hence a homeomorphism. Thus  $Z$  is locally isomorphic to  $X'$ , and hence also to  $C$ ; it is therefore again a locally Boolean  $k$ -scheme.

Finally,  $G$  acts transitively on  $Z$ . Indeed, every  $z \in Z$  is  $G$ -conjugate to an element of the form  $\pi(1, x)$ , so it is enough to show that every  $(1, x) \in P$  is equivalent to some  $(\sigma(h), 0)$ . If  $x \in C$ , take  $h = t_x$ ; if  $x = q_F$ , take  $h = h_F$ . Thus  $Z$  is a locally Boolean  $k$ -scheme endowed with a transitive action of the discrete group  $G$ , but  $Z$  is not separated.

## 6. Subfunctors and group subschemes<sup>\*</sup>

**Definition 6.1.**— (i) Let  $X$  be an  $S$ -functor (that is, a functor from  $(\mathbf{Sch}/S)^\circ$  to  $(\mathbf{Ens})$ ), let  $G$  be an  $S$ -functor in groups, and let  $u$  and  $v$  be two  $S$ -morphisms from  $X$  to  $G$ . The *transporter of  $u$  into  $v$* , denoted  $\underline{\text{Transp}}(u, v)$ , is the sub- $S$ -functor of  $G$  defined by

$$\begin{aligned} \underline{\text{Transp}}(u, v)(S') &= \{g \in G(S') \mid (\text{int } g) \circ u_{S'} = v_{S'}\} \\ &= \left\{ g \in G(S') \mid \begin{array}{l} g_{S''} u_{S''}(x) g_{S''}^{-1} = v_{S''}(x), \\ \text{for every } x \in X(S'') \text{ and every } S'' \rightarrow S' \end{array} \right\}. \end{aligned}$$

In particular,  $\underline{\text{Transp}}(u, u)$  is a group sub- $S$ -functor of  $G$ ; it is called the *centralizer of  $u$*  and is denoted  $\underline{\text{Centr}}(u)$ .

(ii) Let  $G$  be an  $S$ -functor in groups, and let  $X$  and  $Y$  be two sub- $S$ -functors of  $G$ . The *transporter of  $X$  into  $Y$*  (respectively, the *strict transporter of  $X$  into  $Y$* ), denoted  $\underline{\text{Transp}}_G(X, Y)$  (respectively,  $\underline{\text{Transp}}_G^{\text{str}}(X, Y)$ ), is the sub- $S$ -functor of  $G$  defined by

$$\begin{aligned} \underline{\text{Transp}}_G(X, Y)(S') &= \{g \in G(S') \mid (\text{int } g)(X_{S'}) \subset Y_{S'}\} \\ &= \{g \in G(S') \mid g_{S''} X(S'') g_{S''}^{-1} \subset Y(S'') \text{ for every } S'' \rightarrow S'\}, \end{aligned}$$

and, respectively,

$$\begin{aligned} \underline{\text{Transp}}_G^{\text{str}}(X, Y)(S') &= \{g \in G(S') \mid (\text{int } g)(X_{S'}) = Y_{S'}\} \\ &= \{g \in G(S') \mid g_{S''} X(S'') g_{S''}^{-1} = Y(S'') \text{ for every } S'' \rightarrow S'\}. \end{aligned}$$

Note that

$$\underline{\text{Transp}}_G^{\text{str}}(X, Y) = \underline{\text{Transp}}_G(X, Y) \cap c(\underline{\text{Transp}}_G(Y, X)),$$

where  $c$  denotes the inversion morphism of  $G$ .<sup>49</sup>

(iii) Let  $G$  be an  $S$ -functor in groups, let  $H$  be a sub- $S$ -functor of  $G$ , and let  $i : H \rightarrow G$  be the canonical  $S$ -morphism. The *centralizer* and *normalizer of  $H$  in  $G$*  are the following group sub- $S$ -functors of  $G$ :

$$\underline{\text{Centr}}_G H = \underline{\text{Centr}}(i) = \underline{\text{Transp}}(i, i), \quad \underline{\text{Norm}}_G H = \underline{\text{Transp}}_G^{\text{str}}(H, H).$$

Finally, the *center of  $G$*  is the  $S$ -functor in groups

$$\underline{\text{Centr}}(\text{id}_G) = \underline{\text{Centr}}_G G;$$

it is denoted  $\underline{\text{Centr}}G$ .

---

<sup>\*</sup>On the same theme, see also the results of Exposé XI, §6, whose natural place would be in the present Exposé VI<sub>B</sub>. There one finds, in particular, representability criteria for certain group subfunctors of a given group scheme. (48) Editor's note: we have inserted these results below (6.5.2–6.5.5).

<sup>49</sup>Editor's note: if  $u : X \rightarrow G$  and  $v : Y \rightarrow G$  are two arbitrary morphisms of  $S$ -functors, let  $u(X)$  be the image functor of  $u$ , defined by

$$u(X)(S') = u(S')(X(S')) \subset G(S');$$

it is a subfunctor of  $G$ , as is the image functor  $v(Y)$ . One may then consider the transporter of the image of  $u$  into the image of  $v$ ,  $\underline{\text{Transp}}_G(u(X), v(Y))$ . Thus, in part (ii) of the definition, it is not necessary to restrict to subfunctors, that is, to the case in which  $u$  and  $v$  are monomorphisms. In the original, this restriction sometimes imposed repetitions in the hypotheses, such as: “Let  $u, v : X \rightarrow G$  be morphisms of  $S$ -functors and let  $w : H \rightarrow G$  and  $w' : K \rightarrow G$  be monomorphisms; then  $\underline{\text{Transp}}(u, v)$ ,  $\underline{\text{Centr}}_G(w) = \underline{\text{Centr}}_G H$ , and  $\underline{\text{Transp}}_G(H, K)$  satisfy ...” Such repetitions can be avoided by considering  $\underline{\text{Transp}}_G(u(X), v(Y))$ . We have made such modifications in 6.2 and, later, in 10.11.

**Remark 6.1.1.**—<sup>50</sup> It follows from the definitions that the functors  $\underline{\text{Transp}}(u, v)$  and  $\underline{\text{Transp}}_G(X, Y)$  (and hence also  $\underline{\text{Transp}}_G^{\text{str}}(X, Y)$ ) commute with base change: for every  $S' \rightarrow S$ , if  $G', X', Y', u', v'$  are obtained from  $G, X, Y, u, v$  by base change, then

$$\underline{\text{Transp}}(u, v)_{S'} = \underline{\text{Transp}}(u', v'), \quad \underline{\text{Transp}}(X, Y)_{S'} = \underline{\text{Transp}}(X', Y').$$

**Proposition 6.2.**—<sup>51</sup> Let  $G$  be an  $S$ -group. For a subfunctor  $T$  of the functor  $G$ , consider the following property:

$$(+_f) \quad \text{for every } s \in S, T_s \text{ is representable by a closed subscheme of } G_s.$$

Let  $u, w : X \rightarrow G$  and  $v : Y \rightarrow G$  be morphisms of  $S$ -schemes. Then:

- (i)  $\underline{\text{Transp}}(u, w)$  and  $\underline{\text{Centr}}(u) = \underline{\text{Transp}}(u, u)$  satisfy condition  $(+_f)$ .
- (ii)  $\underline{\text{Transp}}_G(u(X), v(Y))$  and  $\underline{\text{Norm}}_G v(Y)$  satisfy condition  $(+_f)$  if, for every  $s \in S$ ,  $v_s$  is a closed immersion.
- (iii)  $\underline{\text{Transp}}_G^{\text{str}}(X, Y)$  satisfies condition  $(+_f)$  if, for every  $s \in S$ ,  $u_s$  and  $v_s$  are closed immersions.

This follows from Remark 6.1.1 and Corollary 6.2.5 below.<sup>52</sup>

**Definition 6.2.1.**— Let  $f : X \rightarrow S$  be a morphism of schemes. One says that  $f$  is *essentially free*, or that  $X$  is *essentially free over  $S$* , if one can find a covering of  $S$  by affine open subsets  $S_i$ , for every  $i$  an affine  $S_i$ -scheme  $S'_i$  faithfully flat over  $S_i$ , and a covering  $(X'_{ij})_j$  of

$$X'_i = X \times_S S'_i$$

by affine open subsets  $X'_{ij}$ , such that, for every  $(i, j)$ , the ring of  $X'_{ij}$  is a projective<sup>53</sup> module over the ring of  $S'_i$ .

**Proposition 6.2.2.**—

- a) If  $X$  is essentially free over  $S$ , it is flat over  $S$ ; the converse holds if  $S$  is Artinian.
- b) If  $S$  is the spectrum of a field, every  $S$ -scheme is essentially free over  $S$ .
- c) If  $X$  is essentially free over  $S$ , then  $X' = X \times_S S'$  is essentially free over  $S'$ , for every  $S' \rightarrow S$ . The converse holds if  $S' \rightarrow S$  is faithfully flat and quasi-compact.

The proof is immediate, using for the converse in a) the fact that a flat module over an Artinian local ring is free.<sup>54</sup>

<sup>50</sup>Editor's note: we have added this remark, which is used implicitly in Proposition 6.2.

<sup>51</sup>Editor's note: we have modified the statement as indicated in editor's note 49.

<sup>52</sup>Editor's note: the original referred to the results of Exposé VIII, §6. For the reader's convenience, we have reproduced these results (with the exception of VIII, 6.3 and 6.8) in 6.2.1–6.2.5 below. Moreover, this had been suggested by A. Grothendieck in a note at the beginning of VIII.6: "The present section is independent of the theory of diagonalizable groups; its natural place would be in VI<sub>B</sub>."

<sup>53</sup>Editor's note: on the one hand, we have replaced the word "free" by "projective", as indicated in VIII, Remark 6.8. On the other hand, the notion of an essentially free  $S$ -scheme is closely related to the geometric notion of an  $S$ -scheme that is flat and pure, introduced and studied by M. Raynaud and L. Gruson ([RG71]); see the addition 6.9 below.

<sup>54</sup>Editor's note: indeed, let  $(A, \mathfrak{m})$  be an Artinian local ring,  $k$  its residue field,  $M$  an arbitrary  $A$ -module, and let  $(x_i)_{i \in I}$  be elements of  $M$  whose images form a basis of  $M/\mathfrak{m}M$  over  $k$ . Let  $F$  be the free  $A$ -module with basis  $(e_i)_{i \in I}$ , and let  $\varphi : F \rightarrow M$  be the  $A$ -morphism defined by  $\varphi(e_i) = x_i$ . Then  $Q = \text{Coker } \varphi$  satisfies  $Q = \mathfrak{m}Q$ , whence, since  $\mathfrak{m}$  is nilpotent,  $Q = 0$ . Suppose moreover that  $M$  is flat over  $A$ ; then  $K = \text{Ker } \varphi$  satisfies  $K \otimes_A k = 0$ , that is,  $K = \mathfrak{m}K$ , whence  $K = 0$ .

The introduction of Definition 6.2.1 is justified by the following theorem.

**Theorem 6.2.3.**— *Let  $S$  be a scheme, let  $Z$  be an  $S$ -scheme that is essentially free, and let  $Y$  be a closed subscheme of  $Z$ . Consider the functor*

$$F = \prod_{Z/S} Y/Z : (\mathbf{Sch}/S)^\circ \longrightarrow \mathbf{Sets}$$

*defined by the following condition:  $F(S') = \emptyset$  when  $Y_{S'} \neq Z_{S'}$ , and  $F(S')$  is a singleton otherwise.*<sup>55</sup>

- (i) *This functor is representable by a closed subscheme  $T$  of  $S$ .*
- (ii) *If, moreover,  $Y \rightarrow Z$  is of finite presentation, then so is  $T \rightarrow S$ .*

*Proof.* First observe that the functor in question is a sheaf for the fpqc topology. Since  $F(S')$  is either  $\emptyset$  or  $\{\text{pt}\}$  for every  $S'$ , this amounts to checking the following assertions. If  $(S_i)$  is an open covering of  $S$  and every

$$Y_{S_i} \longrightarrow Z_{S_i}$$

is an isomorphism, then  $Y \rightarrow Z$  is an isomorphism; and if  $S' \rightarrow S$  is faithfully flat and quasi-compact and

$$Y_{S'} \longrightarrow Z_{S'}$$

is an isomorphism, then again  $Y \rightarrow Z$  is an isomorphism. The first assertion is clear, and the second follows from SGA 1, VIII, 5.4, or EGA IV<sub>2</sub>, 2.7.1.

Moreover, by SGA 1, VIII, 1.9, faithfully flat and quasi-compact morphisms are effective descent morphisms for the fibered category of closed-immersion arrows. With the notation of 6.2.1, this allows us to restrict to the case  $S = S'_i$ .

Let  $(Z_j)$  be a covering of  $Z$  by affine open subsets such that

$$\mathcal{O}(Z_j)$$

is a projective module over  $A = \mathcal{O}(S)$ . Put

$$Y_j = Y \cap Z_j,$$

and let

$$F_j : (\mathbf{Sch}/S)^\circ \longrightarrow \mathbf{Sets}$$

be the functor defined in terms of  $(Z_j, Y_j)$  as  $F$  was defined in terms of  $(Z, Y)$ . It is a subfunctor of the final functor, and clearly

$$F = \bigcap_j F_j.$$

It is therefore enough to prove that every  $F_j$  is representable by a closed subscheme  $T_j$  of  $S$ , for then  $F$  is represented by the closed subscheme

$$T = \bigcap_j T_j$$

---

<sup>55</sup>Editor's note: cf. Exposé II, §1, where this functor is denoted  $\prod_{Z/S} Y$ . For every  $S' \rightarrow S$ ,

$$F(S') = \Gamma(Y_{S'}/Z_{S'}),$$

which here is equal to  $\{\text{id}_{Z_{S'}}\}$  if  $Y_{S'} = Z_{S'}$ , and is empty otherwise. We have also added assertion (ii), which is used in the proof of 6.5.3.



of  $S$ .

We may consequently suppose that  $Z$  is itself affine, say

$$Z = \operatorname{Spec} B,$$

where  $B$  is a projective  $A$ -module. Let  $L$  be a free  $A$ -module with basis  $(e_\lambda)_{\lambda \in \Lambda}$  of which  $B$  is a direct summand as an  $A$ -module, and let

$$\phi_\lambda : L \longrightarrow A$$

be the coordinate forms with respect to this basis. Let  $E$  be a set of generators for the ideal  $J$  of  $B$  defining the closed subscheme  $Y$  of  $Z$ , and let  $I$  be the ideal of  $A$  generated by the coordinates

$$\phi_\lambda(x), \quad x \in E, \quad \lambda \in \Lambda.$$

For every  $A$ -algebra  $C$ , the morphism

$$B \otimes_A C \longrightarrow (B/J) \otimes_A C$$

is an isomorphism if and only if the image of  $x \otimes 1$  in  $L \otimes_A C$  is zero for every  $x \in E$ . This, in turn, is equivalent to saying that the kernel of  $A \rightarrow C$  contains the ideal  $I$ . Hence

$$T = V(I) = \operatorname{Spec}(A/I)$$

has the required property, proving (i).

Finally, if  $Y \rightarrow Z$  is of finite presentation, the set  $E$  may be chosen finite. Then  $I$  is a finitely generated ideal of  $A$ , that is, the closed immersion  $T \rightarrow S$  is of finite presentation. This proves (ii).

**Examples 6.2.4.**— Let us give important examples of functors which reduce to functors

$$\prod_{Z/S} Y/Z$$

of the type considered in 6.2.3, and for which it will be useful to have representability criteria later on. We denote by  $S$  a scheme, and by  $X, Y, Z, \dots$  schemes over  $S$ .

**a)** Suppose given an  $S$ -morphism

$$q : X \longrightarrow \underline{\operatorname{Hom}}_S(Y, Z), \quad (\text{x})$$

(“ $X$  acts on  $Y$ , with values in  $Z$ ”), that is, a morphism

$$r : X \times_S Y \longrightarrow Z. \quad (\text{xx})$$

Consider a subscheme  $Z'$  of  $Z$ , and hence a monomorphism

$$\underline{\operatorname{Hom}}_S(Y, Z') \longrightarrow \underline{\operatorname{Hom}}_S(Y, Z),$$

which makes the first functor a subfunctor of the second. Let  $X'$  be the inverse image of this subfunctor under  $q$ . Thus  $X'$  is the subfunctor of  $X$  for which  $X'(T)$  is the set of  $x \in X(T)$  such that

$$q(x) : Y_T \longrightarrow Z_T$$

factors through  $Z'_T$ .

This functor  $X'$  may be described as follows. Put

$$P = X \times_S Y,$$

and let  $P'$  be the inverse image of  $Z'$  under  $r : P \rightarrow Z$ . Then there is an evident isomorphism

$$X' \simeq \prod_{P/X} P'/P. \quad (\text{xxx})$$

It follows that, if  $Y$  is essentially free over  $S$  and  $Z'$  is closed in  $Z$ , the subfunctor  $X'$  of  $X$  is representable by a closed subscheme of  $X$ .<sup>56</sup> If, moreover,  $Z' \rightarrow Z$  is of finite presentation, then so is  $X' \rightarrow X$ .

**b)** Suppose given two ways of letting  $X$  act on  $Y$ , with values in  $Z$ , that is, two morphisms

$$q_1, q_2 : X \rightrightarrows \underline{\text{Hom}}_S(Y, Z),$$

and put

$$X' = \text{Ker}(q_1, q_2).$$

Thus  $X'$  is the subfunctor of  $X$  for which  $X'(T)$  is the set of  $x \in X(T)$  such that the two morphisms

$$q_1(x), q_2(x) : Y_T \rightrightarrows Z_T$$

are equal. The data of  $q_1, q_2$  are equivalent to the data of a morphism

$$q : X \longrightarrow \underline{\text{Hom}}_S(Y, Z \times_S Z),$$

or, equivalently, of a morphism

$$r : X \times_S Y \longrightarrow Z \times_S Z.$$

Put  $U = Z \times_S Z$ , and let  $U'$  be the diagonal subscheme of  $Z \times_S Z$ . Then  $X'$  is precisely the inverse image under  $q$  of the subfunctor

$$\underline{\text{Hom}}_S(Y, U') \subset \underline{\text{Hom}}_S(Y, U).$$

It can therefore be put in the form (xxx), with  $P = X \times_S Y$  and  $P'$  the inverse image of the diagonal under  $r$ , that is, the kernel of

$$r_1, r_2 : X \times_S Y \rightrightarrows Z.$$

We are thus under the hypotheses of (a).

Consequently, if  $Y$  is essentially free over  $S$  and  $Z$  is separated over  $S$ , the subfunctor  $X'$  of  $X$  is representable by a closed subscheme of  $X$ .<sup>56</sup> If, moreover,  $Z \rightarrow S$  is locally of finite type, then  $X' \rightarrow X$  is of finite presentation.

**c)** Suppose given a morphism

$$q : X \longrightarrow \underline{\text{Hom}}_S(Y, Y),$$

that is, “ $X$  acts on  $Y$ .” Let  $X'$  be the “kernel” of this morphism:  $X'$  is the subfunctor of  $X$  for which  $X'(T)$  is the set of  $x \in X(T)$  such that

$$q(x) : Y_T \longrightarrow Y_T$$

---

<sup>56</sup>Editor’s note: we have added the sentence that follows.

is the identity. This functor falls under (b), as one sees by introducing a second morphism

$$q' : X \longrightarrow \underline{\mathrm{Hom}}_S(Y, Y)$$

by letting  $X$  act trivially on  $Y$ . Hence, if  $Y$  is essentially free and separated over  $S$ , the subfunctor  $X'$ , the kernel of  $q$ , is representable by a closed subscheme of  $X$ .<sup>56</sup> If, moreover,  $Y \rightarrow S$  is locally of finite type, then  $X' \rightarrow X$  is of finite presentation.

**d)** Under the hypotheses of (c), consider the subfunctor  $Y'$  of  $Y$  “of the invariants under  $X$ .” Thus  $Y'(T)$  is the set of  $y \in Y(T)$  such that the corresponding morphism

$$\bar{q}(y) : X_T \longrightarrow Y_T$$

is the constant  $T$ -morphism with value  $y$ . Introducing  $\bar{q}'$  as in (c), and the morphisms corresponding to  $\bar{q}$  and  $\bar{q}'$ ,

$$\bar{q}, \bar{q}' : Y \rightrightarrows \underline{\mathrm{Hom}}_S(X, Y),$$

one sees that

$$Y' = \mathrm{Ker}(\bar{q}, \bar{q}').$$

It therefore falls again under (b), with the roles of  $X$  and  $Y$  reversed and with  $Z = Y$ .

Consequently, if  $X$  is essentially free over  $S$  and  $Y$  is separated over  $S$ , the subfunctor  $Y'$  of  $Y$  of the invariants under  $X$  is representable by a closed subscheme of  $Y$ .<sup>56</sup> If, moreover,  $Y \rightarrow S$  is locally of finite type, then  $Y' \rightarrow Y$  is of finite presentation.

**e)** Constructions of the kind described in the preceding examples occur especially often in group theory. Thus, when  $G$  is an  $S$ -group scheme acting on the  $S$ -scheme  $X$ ,

$$q : G \longrightarrow \underline{\mathrm{Aut}}_S(X),$$

the kernel of  $q$ , the subgroup of  $G$  acting trivially, is a closed subscheme of  $G$ , provided  $X$  is essentially free and separated over  $S$  (example (c)). The subobject  $X^G$  of invariants is a closed subscheme of  $X$ , provided  $G$  is essentially free over  $S$  and  $X$  is separated over  $S$  (example (d)).

Let  $Y, Z$  be subschemes of  $X$ , and consider the subfunctor  $\underline{\mathrm{Transp}}_G(Y, Z)$  of  $G$ , the transporter of  $Y$  into  $Z$ . Its  $T$ -valued points, for  $T$  over  $S$ , are the elements  $g \in G(T)$  such that the corresponding automorphism of  $X_T$  satisfies

$$g(Y_T) \subset Z_T,$$

that is, induces a morphism  $Y_T \rightarrow X_T$  which factors through  $Y_T \rightarrow Z_T$ . Hence, if  $Y$  is essentially free over  $S$  and  $Z$  is closed in  $X$ , then  $\underline{\mathrm{Transp}}_G(Y, Z)$  is a closed subscheme of  $G$  (example (a)).

One may also consider the strict transporter of  $Y$  into  $Z$ ,<sup>57</sup> whose  $T$ -valued points, for  $T$  over  $S$ , are the elements  $g \in G(T)$  such that  $g(Y_T) = Z_T$ . This functor is precisely

$$\underline{\mathrm{Transp}}_G(Y, Z) \cap \sigma(\underline{\mathrm{Transp}}_G(Z, Y)),$$

where  $\sigma$  is the inversion morphism of  $G$ . Consequently, if  $Y$  and  $Z$  are essentially free over  $S$  and closed in  $X$ , the strict transporter of  $Y$  into  $Z$  is a closed subscheme of  $G$ .

An important case is  $X = G$ , with  $G$  acting on itself by inner automorphisms. If  $H$  is a subscheme of  $G$ , the strict transporter of  $H$  into  $H$  is also called the *normalizer of  $H$  in*

<sup>57</sup>Editor's note: denoted  $\underline{\mathrm{Transp}}_G^{\mathrm{str}}(Y, Z)$ .

$G$ , and is denoted  $\underline{\text{Norm}}_G H$ . Hence, if  $H$  is a closed group subscheme of  $G$ , essentially free over  $S$ , then  $\underline{\text{Norm}}_G H$  is representable by a closed group subscheme of  $G$ .

Finally, let  $Z$  be a subscheme of  $G$ . Its centralizer  $\underline{\text{Centr}}_G(Z)$  in  $G$  is the group subfunctor of  $G$  defined by the procedure of (d), when one regards  $Z$  as acting on  $G$  by the operations induced by those of  $G$ . Thus, if  $Z$  is essentially free over  $S$  and  $G$  is separated over  $S$ , then  $\underline{\text{Centr}}_G(Z)$  is a closed group subscheme of  $G$ . In particular, if  $G$  is essentially free and separated over  $S$ , then the center  $C$  of  $G$ , which is none other than  $\underline{\text{Centr}}_G(G)$ , is a closed group subscheme of  $G$ .

When  $S$  is the spectrum of a field, 6.2.2(b) shows that the “essentially free” conditions in examples (a)–(e) above are automatically satisfied, leaving only the separation conditions. Recalling that a group scheme over a field is necessarily separated (VI<sub>A</sub>, 0.3), we obtain, for example:

**Corollary 6.2.5.**— *Let  $G$  be a group scheme over a field  $k$ , and let  $Y, Y'$  be two subschemes of  $G$ . Then:*

- (i) *The centralizer of  $Y$  in  $G$  is a closed group subscheme of  $G$ ; in particular, this holds for the center  $\underline{\text{Centr}}_G(G)$  of  $G$ .*
- (i') *More generally, if  $u, v : X \rightarrow G$  are morphisms of schemes, then  $\underline{\text{Transp}}_G(u, v)$  is representable by a closed subscheme of  $G$ .*
- (ii) *If  $Y$  is closed, the transporter  $\underline{\text{Transp}}_G(Y', Y)$  is a closed subscheme of  $G$ . If  $Y'$  is also closed, the same conclusion holds for  $\underline{\text{Transp}}_G^{\text{str}}(Y', Y)$ .*
- (iii) *For every group subscheme<sup>58</sup>  $H$  of  $G$ ,  $\underline{\text{Norm}}_G(H)$  is a closed group subscheme of  $G$ .*

**Corollary 6.2.6.**—<sup>59</sup> *Let  $k$  be a field and let  $G$  be a connected algebraic  $k$ -group. Then  $\underline{\text{Centr}}_G(G)$  is representable by a closed group subscheme  $Z$  of  $G$ , and  $G/Z$  is an affine algebraic  $k$ -group.*

*Proof.* The first assertion is, of course, contained in 6.2.5(i), but we shall see that it also follows from the proof of the second assertion. Indeed,  $G$  acts by the adjoint representation on the finite-dimensional  $k$ -vector spaces

$$V_n = \mathcal{O}_{G,e} / \mathfrak{m}_e^{n+1},$$

where  $\mathfrak{m}_e$  is the maximal ideal of  $\mathcal{O}_{G,e}$ . Let  $K_n$  be the kernel of the morphism

$$\rho_n : G \longrightarrow \text{GL}(V_n).$$

By VI<sub>A</sub>, 5.4.1,  $\rho_n$  induces a closed immersion

$$G/K_n \hookrightarrow \text{GL}(V_n),$$

so every  $G/K_n$  is affine. Since  $G$  is Noetherian, the intersection  $K$  of the  $K_n$  is equal to one of the  $K_n$ ; hence  $G/K$  is affine.

On the other hand, if  $Z$  denotes the center of  $G$ , it is clear that

$$Z \subset K.$$

<sup>58</sup>Editor's note: indeed, over a field  $k$ , every group subscheme of  $G$  is closed; see VI<sub>A</sub>, 0.5.2. Moreover, 6.2.5 completes the insertion of the results taken from VIII, §6.

<sup>59</sup>Editor's note: we have inserted this corollary here (cf. Exposé XII, 6.1), as it will be useful later. We return in 6.3 to the original text of Exposé VI<sub>B</sub>.

We shall prove the reverse inclusion. Let  $\widehat{\mathcal{O}}_{G,e}$  be the completion of  $\mathcal{O}_{G,e}$  for the  $\mathfrak{m}_e$ -adic topology, and put

$$\widehat{S} = \operatorname{Spec} \widehat{\mathcal{O}}_{G,e}, \quad S = \operatorname{Spec} \mathcal{O}_{G,e}.$$

Since  $\widehat{S} \rightarrow S$  is faithfully flat, and since the two morphisms

$$K \times_k S \longrightarrow S, \quad (g, x) \longmapsto gxg^{-1} \quad \text{and} \quad (g, x) \longmapsto x$$

coincide after the base change  $\widehat{S} \rightarrow S$ , they coincide. Thus  $K$  acts trivially on  $\mathcal{O}_{G,e}$ .

By 6.2.4(e), the subobject  $G^K$  of points of  $G$  fixed by  $K$ , which is precisely  $\underline{\operatorname{Centr}}_G(K)$ , is a closed subscheme of  $G$ . It is therefore defined by a quasi-coherent ideal  $\mathcal{I}$  of  $\mathcal{O}_G$ . Since  $G^K$  contains

$$S = \operatorname{Spec} \mathcal{O}_{G,e}$$

and  $\mathcal{I}$  is of finite type (because  $G$  is Noetherian), there is an open neighborhood  $U$  of  $e$  such that

$$\mathcal{I}|_U = 0.$$

The subgroup

$$G^K = \underline{\operatorname{Centr}}_G(K)$$

therefore contains  $U$ , and hence also  $U \cdot U$ , which is equal to  $G$  because  $G$  is irreducible (VI<sub>A</sub>, 0.5). Consequently,

$$\underline{\operatorname{Centr}}_G(K) = G.$$

Thus  $K \subset Z$ , and therefore  $Z = K$ , as required.

**Remark 6.3.**— *Let  $k$  be an algebraically closed field, let  $G$  be a  $k$ -group, and let  $H$  be a group subscheme of  $G$ . Suppose that  $G$  and  $H$  are of finite type over  $k$  and are reduced. Then  $\underline{\operatorname{Norm}}_G H$  (respectively,  $\underline{\operatorname{Centr}}_G H$ ) is representable by a group subscheme of  $G$ , whose associated reduced subscheme is precisely the normalizer (respectively, centralizer) of  $H$  in  $G$  in the sense of the Bible.*

**Proposition 6.4.**— *Let  $G$  be an  $S$ -group, and let  $u : X \rightarrow G$  be a monomorphism of  $S$ -schemes. Set*

$$T = \underline{\operatorname{Transp}}_G(X, X).$$

*The following conditions are equivalent:*

- (i)  *$T$  is a group sub- $S$ -functor of  $G$ .*
- (ii)

$$T = \underline{\operatorname{Transp}}_G^{\operatorname{str}}(X, X) = \underline{\operatorname{Norm}}_G X.$$

*These conditions hold in each of the following two cases:*

- a)  *$X$  is of finite presentation over  $S$ .*
- b)  *$T$  is representable by a scheme of finite presentation over  $S$ .*

The equivalence of (i) and (ii) follows from the fact that, for every morphism  $S' \rightarrow S$  and all  $t, t' \in T(S')$ , one has

$$tt' \in T(S'),$$

and from the identity

$$\underline{\operatorname{Transp}}_G^{\operatorname{str}}(X, X) = T \cap c(T)$$

(cf. 6.1(ii)).

Suppose first that a) holds. If  $t \in T(S)$ , then  $\text{int}(t)$  is a monomorphism from  $X$  to itself, and therefore an  $S$ -automorphism of  $X$  by EGA IV<sub>4</sub>, 17.9.6. Thus

$$t \in \underline{\text{Transp}}_G^{\text{str}}(X, X),$$

which proves the assertion in case a).

In case b), it is clear that

$$\mu(T \times_S T) \subset T,$$

and the assertion follows from the next lemma.

**Lemma 6.4.2.**—<sup>60</sup> *Let  $G$  be an  $S$ -scheme of finite presentation, equipped with an associative law (in the sense of EGA 0<sub>III</sub>, 8.2.5). Suppose that, for every  $S$ -scheme  $S'$  and every  $g \in G(S')$ , right and left translation by  $g$  in the set  $G(S')$  are injective, and suppose that  $G(S) \neq \emptyset$ . Then  $G$  is an  $S$ -group.*

*Proof.* It is enough to prove that  $G(S')$  is a group for every  $S$ -scheme  $S'$ . The hypothesis immediately implies that right and left translation by every  $g \in G(S')$  on  $G_{S'}$  are  $S'$ -monomorphisms

$$G_{S'} \longrightarrow G_{S'}.$$

They are therefore  $S'$ -automorphisms, since  $G$  is of finite presentation over  $S$  (EGA IV<sub>4</sub>, 17.9.6). Consequently, right and left translation by  $g$  in the set  $G(S')$  are bijective, and the assertion reduces to the following lemma.

**Lemma 6.4.3.**— *Let  $G$  be a nonempty set with an associative law such that right and left translations are bijective. Then  $G$  is a group.*

The proof is left to the reader.

**Definition 6.5.**— *Let  $G$  be an  $S$ -group, let  $H$  be an  $S$ -functor, and let  $u : H \rightarrow G$  be a monomorphism.*

- (i) The *connected centralizer of  $H$  in  $G$* , denoted

$$\underline{\text{Centr}}_G^0 H,$$

is the identity component of the functor  $\underline{\text{Centr}}_G H$  (cf. 3.1 and 6.1(iii)).

- (i') For every morphism  $u : X \rightarrow G$ , one similarly defines the functor

$$\underline{\text{Centr}}^0(u)$$

(cf. 6.2(iv)).

- (ii) When  $u_s$  is a closed immersion for every  $s \in S$ , the *connected normalizer of  $H$  in  $G$* , denoted

$$\underline{\text{Norm}}_G^0 H,$$

is the identity component of the functor  $\underline{\text{Norm}}_G H$ .

*N.B.* By 1.4.2, the hypothesis in (ii) is satisfied when  $H$  is an  $S$ -group scheme,  $G$  and  $H$  are locally of finite type over  $S$ , and  $u$  is a quasi-compact morphism of  $S$ -groups.

**Proposition 6.5.1.**— *Let  $G$  be an  $S$ -group locally of finite presentation and quasi-separated over  $S$ , let  $H$  be a smooth  $S$ -group with connected fibers, and let  $u : H \rightarrow G$  be a monomorphism. Let  $N$  be the normalizer of  $H$  in  $G$  (cf. 6.1). By 6.5.5 below,  $N$  is representable by a closed group subscheme of  $G$ , of finite presentation over  $G$ . Under these circumstances, the following conditions are equivalent:*

<sup>60</sup>Editor's note: we have retained the numbering of the original; there is no number 6.4.1.

- (i) *The canonical morphism  $H \rightarrow N$  is an open immersion.*
- (ii)  $N^0 = H$  (cf. 3.10).
- (iii) *For every  $s \in S$ , one has  $H_s = (N_s)^0$ .*

Condition (i) implies (ii) by Lemma 3.10.1, since  $H^0 = H$ . It is also clear that (ii) implies (iii), for

$$H_s = (N^0)_s = (N_s)^0.$$

Finally, let us prove that (iii) implies (i). Since  $H_s = (N^0)_s$  for every  $s \in S$ , the morphism

$$H_s \longrightarrow N_s$$

is an open immersion. Moreover,  $H$  and  $N$  are locally of finite presentation over  $S$ , and  $H$  is flat over  $S$ . Consequently,  $H \rightarrow N$  is an open immersion (EGA IV<sub>4</sub>, 17.9.5).

For the reader's convenience, the results 6.8–6.11 of Exposé XI have been included below.<sup>61</sup>

**Theorem 6.5.2.**— *Let  $X$  be a smooth scheme over  $S$ , with geometrically irreducible fibers.*<sup>62</sup>

- (i) *For every closed subscheme  $Y$  of  $X$ , the functor*

$$\prod_{X/S} Y/X$$

*is representable by a closed subscheme  $T$  of  $S$ .*

- (ii) *Moreover, if  $Y \rightarrow X$  is of finite presentation, then so is  $T \rightarrow S$ .*

*Proof.* Let  $f : X \rightarrow S$  denote the structure morphism. Since  $f$  is faithfully flat and locally of finite presentation, it is a covering for the fpqc topology. Moreover,

$$T = \prod_{X/S} Y/X$$

is evidently a subsheaf of  $S$  for the fpqc topology. It follows that the representability of  $T$  by a closed subscheme of  $S$  is a question local on  $S$  for the fpqc topology,<sup>63</sup> and the same is true of the question whether  $T$  is of finite presentation over  $S$  (EGA IV<sub>2</sub>, 2.7.1).

After the base change  $S' \rightarrow S$  with  $S' = X$ , we are reduced to the case in which  $X$  admits a section

$$\epsilon : S \longrightarrow X.$$

We may also suppose that  $S$  is affine, and hence quasi-compact. We then have:

**Corollary 6.5.3.**— *Under the hypotheses of 6.5.2, suppose that  $S$  is quasi-compact, that  $X \rightarrow S$  admits a section  $\epsilon$ , and that  $Y \rightarrow X$  is of finite presentation. Then there exists an integer  $n \geq 0$  such that*

$$\prod_{X/S} Y/X = \prod_{X_n/S} Y_n/X_n,$$

<sup>61</sup>Editor's note: Sections 6.5.2–6.5.5, taken from Exposé XI, 6.8–6.11, have been inserted here. This was in fact suggested by A. Grothendieck in a note at the beginning of XI 6: “The present section does not use the results of Sections 3, 4, and 5 [of XI]; its natural place would be in VI<sub>B</sub>.”

<sup>62</sup>Editor's note: on the one hand, we have corrected the original by replacing “connected” with “irreducible” (see the proof). On the other hand, according to [RG71], I, 3.3.4(iii) and 4.1.1, it is enough to assume that  $X$  is flat over  $S$ , with geometrically irreducible fibers and without embedded components.

<sup>63</sup>Editor's note: see, for example, the addition 1.7 in Exposé VIII.

where  $X_n$  is the  $n$ -th infinitesimal neighborhood of the immersion  $\epsilon : S \rightarrow X$ , and  $Y_n = Y \cap X_n$ .

When  $Y$  is of finite presentation over  $X$ , this corollary does indeed imply 6.5.2. Since  $X$  is smooth over  $S$ ,  $X_n$  is finite and locally free over  $S$ , and is therefore, a fortiori, essentially free over  $S$  (see 6.2.1). Hence

$$\prod_{X_n/S} Y_n/X_n$$

is representable by a closed subscheme  $T$  of  $S$ , of finite presentation over  $S$ , by 6.2.3.

Let us first prove 6.5.3, and hence 6.5.2, when  $S$  is Noetherian. Put

$$T_n = \prod_{X_n/S} Y_n/X_n.$$

The  $T_n$  form a decreasing sequence of closed subschemes of  $S$ , and since  $S$  is Noetherian this sequence is stationary. Let

$$R = \bigcap_{n \geq 0} \prod_{X_n/S} Y_n/X_n$$

be their common value for all sufficiently large  $n$ . Clearly  $T \subset R$ , and it is enough to prove  $R \subset T$ .

After the base change  $R \rightarrow S$ , we are reduced to the case

$$R = S,$$

that is,

$$Y_n = X_n \quad \text{for every } n, \quad \text{or equivalently} \quad Y \supset X_n \quad \text{for every } n.$$

We must then prove  $T = S$ , that is,  $Y = X$ .

Now  $Y \supset X_n$  for every  $n$ , together with the fact that  $X$  is locally Noetherian, implies that in a neighborhood of each point of  $\epsilon(S)$ ,  $Y$  is an induced open subscheme of  $X$ .<sup>64</sup> Consequently, there exists an induced open subscheme  $U$  of  $X$ , containing  $\epsilon(S)$ , such that  $U \subset Y$ . By EGA IV<sub>3</sub>, 11.10.10, since the fibers of  $X/S$  are integral,  $U$  is schematically dense in  $X$ . Since  $Y$  is a closed subscheme containing  $U$ , it follows that  $Y = X$ . This proves 6.5.3, and hence 6.5.2, when  $S$  is Noetherian.

The general case is reduced to the preceding one. For every  $s \in S$ , there exist an affine open neighborhood  $U$  of  $s$  and an affine open neighborhood  $V$  of  $\epsilon(s)$  such that  $f(V) \subset U$ . Then  $f(V)$  is an open neighborhood of  $s$  contained in  $U$ . Let  $S'$  be an affine open neighborhood of  $s$  contained in

$$\epsilon^{-1}(V) \cap f(V),$$

and put

$$X' = V \cap f^{-1}(S').$$

Then  $X'$  and  $S'$  are affine open subsets of  $X$  and  $S$ , respectively, and  $X'/S'$  admits a section. By the local nature of 6.5.2 and 6.5.3, we may suppose that  $S = S'$ .

Thus  $X'$  is an affine open subset of  $X$  containing  $\epsilon(S)$ . Since every fiber  $X_s$  is irreducible,  $X'_s$  is schematically dense in  $X_s$ . Consequently, by EGA IV<sub>3</sub>, 11.10.10,  $X'$  is schematically dense in  $X$ ; likewise, after every base change  $S_1 \rightarrow S$ ,

$$X' \times_S S_1$$

---

<sup>64</sup>Editor's note: see, for example, the proof of 6.2.6.



is schematically dense in  $X \times_S S_1$ .

It follows that

$$\prod_{X/S} Y/X = \prod_{X'/S} Y'/X', \quad Y' = Y \cap X'.$$

This reduces us to the case  $X = X'$ , so we may suppose that both  $S$  and  $X$  are affine. Moreover, if

$$X = \operatorname{Spec} B$$

and  $J$  is the ideal of  $B$  defining  $Y$ , then  $J$  is the filtered colimit of its finitely generated subideals. Thus  $Y$  is the intersection of closed subschemes  $Y_i$  of  $X$  that are of finite presentation over  $X$ , and consequently

$$\prod_{X/S} Y/X = \bigcap_i \prod_{X/S} Y_i/X.$$

To prove 6.5.2, we are therefore reduced to the case in which  $Y$  is of finite presentation over  $X$ , with  $S$  and  $X$  affine.

In that case  $X$  and  $Y$  over  $S$  are obtained by base change  $S \rightarrow S_0$  from an analogous situation  $X_0, Y_0$  over a Noetherian scheme  $S_0$ . This reduces us to the Noetherian case already treated, and completes the proof of 6.5.2 and 6.5.3.

**Corollary 6.5.4.**— *Let  $X$  be a smooth  $S$ -group scheme of finite presentation with connected fibers, let  $Y$  be a group scheme of finite presentation over  $S$ , and let  $i : Y \rightarrow X$  be a monomorphism of  $S$ -group schemes.*

(i) *The functor*

$$\prod_{X/S} Y/X$$

*is representable by a closed subscheme of  $S$  of finite presentation.*

(ii) *If, moreover,  $S$  is quasi-compact, then for all sufficiently large  $n$ ,*

$$\prod_{X/S} Y/X = \prod_{X_n/S} Y_n/X_n,$$

*where  $X_n$  denotes the  $n$ -th infinitesimal neighborhood of the unit section  $\epsilon : S \rightarrow X$ , and  $Y_n = X_n \cap Y$ .*

The proof is essentially that of 6.5.3.<sup>65</sup> On the one hand,  $i$  is locally of finite presentation (EGA IV<sub>1</sub>, 1.4.3). On the other hand, the unit sections of  $Y$  and  $X$  induce bijective immersions

$$S \longrightarrow Y_n \quad \text{and} \quad S \longrightarrow X_n,$$

and hence isomorphisms of  $S_{\text{red}}$  with  $(Y_n)_{\text{red}}$  and  $(X_n)_{\text{red}}$ . Consequently,  $i_n$  is quasi-compact, hence of finite type, and one has the commutative diagram

$$\begin{array}{ccc} (Y_n)_{\text{red}} & \xrightarrow{\tau} & Y_n \\ & \searrow \sigma & \downarrow i_n \\ & & X_n \end{array}$$

<sup>65</sup>Editor's note: the original has been expanded in what follows, and references to EGA IV have been added.

where  $\sigma$  and  $\tau$  are closed immersions and  $\tau$  is surjective. Since  $i_n$  is separated (being a monomorphism), it follows that  $i_n$  is proper (EGA II, 5.4.3). Thus  $i_n$  is a proper monomorphism of finite presentation, and therefore a closed immersion (EGA IV<sub>3</sub>, 8.11.5). It follows from 6.2.3 that

$$\prod_{X_n/S} Y_n/X_n$$

is representable by a closed subscheme  $T_n$  of  $S$ , of finite presentation over  $S$ . It remains to prove the last assertion of 6.5.4 when  $S$  is assumed affine as well.

We again reduce immediately to the case where  $S$  is Noetherian. With the notation of the proof of 6.5.3, it is enough to prove  $R = T$ , or equivalently that

$$Y \supset X_n \quad \text{for every } n \quad \implies \quad Y = X.$$

The hypothesis implies that  $i : Y \rightarrow X$  is étale at the points of the unit section of  $Y$  over  $S$ . Thus  $Y$  is smooth over  $S$  at those points, and the open subset  $U$  of points of  $Y$  at which  $Y$  is smooth over  $S$  is an induced open subgroup of  $Y$  (cf. 2.3). The morphism

$$\tau : U \longrightarrow X$$

is then an étale monomorphism by 2.5, and hence an open immersion. Since the fibers of  $X$  are connected and every open subgroup of an algebraic group is also closed,  $\tau$  is a surjective open immersion, that is, an isomorphism. Therefore  $U = X$ , and a fortiori  $Y = X$ . This completes the proof of 6.5.4.

Proceeding as in 6.2.4(e), one deduces Corollary 6.5.5 from 6.5.4.

**Corollary 6.5.5.**— *Let  $G$  be an  $S$ -group scheme locally of finite type and quasi-separated over  $S$ , let  $H$  be a smooth group scheme of finite presentation over  $S$  with connected fibers, and let  $i : H \rightarrow G$  be a monomorphism of  $S$ -groups. Then:*

- a)  $\underline{\text{Centr}}_G(H)$  and  $\underline{\text{Norm}}_G(H)$  are representable by closed subschemes of  $G$ , of finite presentation over  $G$ .
- a') Likewise, for every monomorphism  $j : K \rightarrow G$  of  $S$ -group schemes that is of finite presentation,  $\underline{\text{Transp}}_G(H, K)$  is representable by a closed subscheme of  $G$ , of finite presentation over  $G$ .
- b) If  $G$  is quasi-compact, there exists an integer  $n \geq 0$  such that, if  $H_n$  denotes the  $n$ -th infinitesimal neighborhood of the unit section of  $H$ , then

$$\begin{aligned} \underline{\text{Centr}}_G(H) &= \underline{\text{Centr}}_G(H_n), & \underline{\text{Norm}}_G(H) &= \underline{\text{Norm}}_G(H_n), \\ \underline{\text{Transp}}_G(H, K) &= \underline{\text{Transp}}_G(H_n, K) = \underline{\text{Transp}}_G(H_n, K_n). \end{aligned}$$

*Proof.*<sup>66</sup> First observe that the hypothesis on  $G$  implies that the monomorphism  $H \rightarrow G$  is of finite presentation (EGA IV<sub>1</sub>, 1.2.4 and 1.4.3), as is the diagonal immersion

$$\Delta_{G/S} : G \longrightarrow G \times_S G$$

(loc. cit., 1.4.3.1). The case of  $\underline{\text{Norm}}_G(H)$  therefore reduces to that of the transporter, by taking  $H = K$ .

<sup>66</sup>Editor's note: the argument that follows has been expanded. This also completes the insertion of XI, 6.8–6.11; at 6.6 below we return to the original text of VI<sub>B</sub>.

In view of 6.2.4(e), we apply 6.5.4 to the group scheme

$$X = H_G = H \times_S G$$

over the base scheme  $G$ , and to the following group subscheme  $Y$ .

In the case of  $\underline{\text{Transp}}_G(H, K)$ , take  $Y$  to be the inverse image of  $K_G$  under the morphism of  $G$ -groups

$$H \times_S G \longrightarrow G \times_S G, \quad (h, g) \longmapsto (ghg^{-1}, g).$$

In the case of  $\underline{\text{Centr}}_G(H)$ , take  $Y$  to be the inverse image of the diagonal subgroup

$$\Delta_{G/S}(G)_G$$

of  $(G \times_S G)_G$  under the morphism of  $G$ -groups

$$H \times_S G \longrightarrow G \times_S G \times_S G, \quad (h, g) \longmapsto (h, ghg^{-1}, g).$$

**Definition 6.6.**— Let  $G$  be an  $S$ -functor in groups and let  $H$  be a group sub- $S$ -functor of  $G$ . One says that  $H$  is *invariant* (respectively, *central*, respectively, *characteristic*) in  $G$  if

$$\underline{\text{Norm}}_G H = G$$

(respectively, if  $\underline{\text{Centr}}_G H = G$ , respectively, if for every  $S$ -scheme  $T$  and every automorphism

$$a \in \text{Aut}_{T\text{-gr}}(G_T)$$

one has  $a(H_T) \subset H_T$ ). In other words, for every  $S$ -scheme  $T$ , the subgroup  $H(T)$  of  $G(T)$  is normal in  $G(T)$  (respectively, central in  $G(T)$ , respectively, invariant under every automorphism of  $G_T$ ).

*N.B.* If  $H$  is central (respectively, characteristic), then it is invariant.

**Remark 6.7.**— Let  $G$  and  $H$  be two  $S$ -groups and let  $u : H \rightarrow G$  be a monomorphism. The group  $H$  is invariant (respectively, central) in  $G$  if and only if the morphism

$$\mu \circ (c \circ \text{pr}_2, \mu \circ (u \times \text{id}_G)) : H \times_S G \longrightarrow G$$

(defined by

$$(h, g) \longmapsto g^{-1}hg$$

for  $g \in G(S')$  and  $h \in H(S')$ ) factors through  $u$  (respectively, is equal to  $u \circ \text{pr}_1$ ). The group  $H$  is characteristic in  $G$  if and only if, for every  $S$ -scheme  $T$  and every  $T$ -group automorphism  $a$  of  $G_T$ , the morphism  $a \circ u(T)$  factors through  $u(T)$ .

**Example 6.8.**— Let  $G$  be an  $S$ -functor in groups. Then  $\underline{\text{Centr}}G$  is characteristic and central. If  $G_s$  is representable for every  $s \in S$ , then  $G^0$  is characteristic. This follows from the definitions and 3.3.

**6.9.**—<sup>67</sup> In [RG71], I, 3.3.3, the authors introduce the geometric notion of a *pure  $S$ -scheme*, which is local on  $S$  for the étale topology; we refer to *loc. cit.* for the precise definition. We merely record the following points:

- a) (*loc. cit.*, Theorem 3.3.5) If  $B$  is a flat, finitely presented  $A$ -algebra, then  $B$  is a projective  $A$ -module if and only if  $\text{Spec}(B)$  is pure over  $\text{Spec}(A)$ .

---

<sup>67</sup>Editor's note: this subsection has been added.

- b) Consequently, if  $X \rightarrow S$  is locally of finite presentation, flat, and pure, then  $X$  is essentially free over  $S$ .
- c) (*loc. cit.*, 3.3.4(iii)) If  $X \rightarrow S$  is locally of finite type, flat, has geometrically irreducible fibers, and has no embedded components, then  $X$  is pure over  $S$ .

Every group scheme locally of finite type over a field is Cohen–Macaulay (cf. VI<sub>A</sub>, 1.1.1), and therefore has no embedded components. We obtain in particular:

- d) Every  $S$ -group scheme  $G$  that is locally of finite presentation, flat, and has connected fibers is pure over  $S$ , and hence essentially free over  $S$ .

All the statements of 6.2.4(e) may therefore be used with the results of (b) and (d) above taken into account.

## 7. Generated subgroups; commutator group

Throughout this section,  $k$  denotes a fixed field.

**Proposition 7.1.**— *Let  $G$  be a  $k$ -group, let  $(X_i)_{i \in I}$  be a family of geometrically reduced  $k$ -schemes,<sup>68</sup> and, for every  $i \in I$ , let  $f_i : X_i \rightarrow G$  be a  $k$ -morphism.*

- (i) *There exists a smallest closed  $k$ -group subscheme of  $G$  through which every  $f_i$  factors. It is denoted*

$$\Gamma_G((f_i)_{i \in I}).$$

*It is a geometrically reduced  $k$ -scheme, and hence smooth when  $G$  is locally of finite type over  $k$  (1.3.1).*

- (ii) *Put*

$$X = \coprod_{i \in I} X_i,$$

*and let  $f : X \rightarrow G$  be the morphism whose restriction to  $X_i$  is  $f_i$ , for every  $i \in I$ . Put*

$$X^1 = X \amalg X,$$

*and let  $f^1 : X^1 \rightarrow G$  be the morphism whose restrictions to the two copies of  $X$  are  $f$  and  $c \circ f$ , respectively. For every  $n > 1$ , put*

$$X^n = X^1 \times_k X^{n-1}, \quad f^n = \mu \circ (f^1 \times_k f^{n-1}) : X^n \rightarrow G.$$

*Then  $\Gamma_G((f_i)_{i \in I})$  is the reduced subscheme of  $G$  whose underlying space is the closure of*

$$\bigcup_{n \geq 1} f^n(X^n).$$

- (iii) *For every  $k$ -scheme  $S$ ,  $\Gamma_G((f_i)_{i \in I})_S$  is the smallest closed group subscheme of  $G_S$  through which every*

$$f_{i,S} : (X_i)_S \rightarrow G_S$$

*factors.*

---

<sup>68</sup>Editor's note: here and below, the little-used term “separable” has been replaced by the customary term “geometrically reduced”; see EGA IV<sub>2</sub>, 4.6.2.

(iii') Moreover,  $\Gamma_G((f_i)_{i \in I})_S$  is the smallest group subscheme of  $G_S$ , not necessarily closed, through which every  $f_{i,S}$  factors.<sup>69</sup>

First note that, to prove (i), (iii), and (iii'), after defining  $X$  and  $f$  as in (ii), we are reduced to the case in which  $I$  consists of a single element.

Let  $H$  be the reduced subscheme of  $G$  whose underlying set is

$$\bigcup_{n \geq 1} \overline{f^n(X^n)}.$$

The family of morphisms

$$f^n : X^n \longrightarrow H$$

is schematically dominant (cf. EGA IV<sub>3</sub>, 11.10.4), and therefore every closed subscheme of  $G$  through which all the  $f^n$  factor also contains  $H$ . Moreover, by *loc. cit.*, 11.10.7,  $H$  is geometrically reduced. To prove (i) and (ii), it is therefore enough to show that  $H$  is a group subscheme of  $G$ , that is, that the restriction of  $c$  to  $H$  and the restriction of  $\mu$  to  $H \times_k H$  factor through the inclusion  $H \rightarrow G$ .

Since  $H$  is geometrically reduced,  $H \times_k H$  is reduced, and it is enough to verify, set-theoretically, that

$$c(H) \subset H, \quad \mu(H \times_k H) \subset H.$$

By EGA IV<sub>3</sub>, 11.10.6, the union of the

$$f^n_{(H)}(X^n \times_k H)$$

is schematically dense in  $H \times_k H$ . Similarly, for every  $n \geq 1$ , the union of the

$$f^m_{(X^n)}(X^n \times_k X^m), \quad m \geq 1,$$

is schematically dense in  $X^n \times_k H$ . It is therefore enough to prove

$$\mu\left(f^n_{(H)}\left(f^m_{(X^n)}(X^n \times_k X^m)\right)\right) \subset H$$

and  $c(f^n(X^n)) \subset H$ . But

$$\begin{aligned} \mu\left(f^n_{(H)}\left(f^m_{(X^n)}(X^n \times_k X^m)\right)\right) &= \mu((f^n \times_k f^m)(X^n \times_k X^m)) \\ &= f^{n+m}(X^{n+m}) \subset H. \end{aligned}$$

Also, since  $c(f^1(X^1)) \subset f^1(X^1)$ , for every  $n$  one has

$$c(f^n(X^n)) \subset f^n(X^n) \subset H.$$

This proves (i) and (ii).

We now prove (iii). Let  $G'$  be a closed group subscheme of  $G_S$  through which  $f_S$  factors. We must show that  $G'$  contains  $H_S$ , or equivalently that

$$H_S = H_S \times_{G_S} G'.$$

Put

$$H'_S = H_S \times_{G_S} G' = G' \cap H_S.$$

<sup>69</sup>Editor's note: the original stated (iii') under the additional hypothesis that  $G$  be locally of finite type over  $k$ ; this hypothesis may be omitted in view of VI<sub>A</sub>, 0.5.2.

Every  $f_S^n$  factors through both  $G'$  and  $H_S$ , and hence through  $H'_S$ . Since the family

$$f^n : X^n \longrightarrow H$$

is schematically dominant, EGA IV<sub>3</sub>, 11.10.6 implies that the base-changed family

$$f_S^n : X_S^n \longrightarrow H_S$$

is schematically dominant as well. Because every  $f_S^n$  factors through  $H'_S$ , EGA IV<sub>3</sub>, 11.10.1(c) gives

$$H'_S = H_S.$$

This proves (iii).

Finally, we show that  $H_S$  is the smallest group subscheme, not necessarily closed, of  $G_S$  through which  $f_S$  factors. Let  $G'$  be such a group subscheme and again put

$$H'_S = H_S \times_{G_S} G'.$$

We must show  $H'_S = H_S$ . By (iii), it is enough to show that  $H'_S$  is closed in  $H_S$ . It is enough in turn to show that  $H_S$  and  $H'_S$  have the same underlying set, and for this it suffices to show, for every  $s \in S$ , that

$$H'_s := H'_S \times_S \kappa(s) = H_s \times_{G_s} G'_s = H_s.$$

By VI<sub>A</sub>, 0.5.2, the  $\kappa(s)$ -group subscheme  $G'_s$  is closed in  $G_s$ . Hence  $H'_s$  is closed in  $H_s$ , and the preceding argument, applied to  $H'_s$ ,  $H_s$ , and the  $f_s^n$ , shows that

$$H'_s = H_s.$$

This proves (iii') and completes the proof.

**Corollary 7.1.1.**—<sup>70</sup> *Let  $G$  be a  $k$ -group locally of finite type, let  $A$  and  $B$  be two  $k$ -subgroups of  $G$  that are smooth and of finite type, and let  $i_A$  (respectively,  $i_B$ ) be the inclusion of  $A$  (respectively,  $B$ ) into  $G$ . Suppose that  $B$  normalizes  $A$ . Then*

$$A \cdot B = \Gamma_G(i_A, i_B).$$

*Proof.* Let

$$H = A \rtimes B$$

be the semidirect product of  $A$  and  $B$  (cf. I, 2.3.5); it is a smooth  $k$ -group of finite type. The group morphism

$$u : H \longrightarrow G, \quad (a, b) \longmapsto ab,$$

is therefore quasi-compact. Hence, by 1.2,

$$u(H) = A \cdot B$$

is a closed reduced subscheme of  $G$ , and is a group in the category  $(\text{Sch}/k)_{\text{red}}$ . By EGA IV<sub>3</sub>, 11.10.7,  $A \cdot B = u(H)$  is geometrically reduced, since  $H$  is; it is therefore a closed subgroup of  $G$ . The evident inclusion

$$A \cdot B \subset \Gamma_G(i_A, i_B)$$

now proves the corollary.

---

<sup>70</sup>Editor's note: this corollary has been added to point out this special case of 7.1.

**Definitions and Remarks 7.2.**—<sup>71</sup>

- (i) Let  $G$  be a  $k$ -group, let  $(X_i)_{i \in I}$  be a family of geometrically reduced  $k$ -schemes, and, for every  $i \in I$ , let  $f_i : X_i \rightarrow G$  be a  $k$ -morphism. The *closed group subscheme of  $G$  generated by the family  $(f_i)_{i \in I}$*  is the smallest closed group subscheme of  $G$  through which every  $f_i$  factors; in this subsection it is denoted

$$\Gamma_G((f_i)_{i \in I}).$$

If  $X$  is a subscheme of  $G$ , geometrically reduced over  $k$ , and  $f : X \hookrightarrow G$  is the immersion, we write  $\Gamma_G(X)$  instead of  $\Gamma_G(f)$ .

- (i') With the notation of 7.1(ii), we shall sometimes put

$$\Gamma'_G(f) = \bigcup_{n \geq 1} f^n(X^n).$$

Notice that  $\Gamma'_G(f)$  is a subset of  $G$  stable under the group law, in the sense of 3.0.

- (ii) If  $X_1$  and  $X_2$  are geometrically reduced  $k$ -schemes and  $f_1 : X_1 \rightarrow G$  and  $f_2 : X_2 \rightarrow G$  are  $k$ -morphisms such that

$$f_1(X_1) = f_2(X_2)$$

as sets, then it is clear that

$$\Gamma_G(f_1) = \Gamma_G(f_2).$$

- (iii) Let  $E$  be a subset of  $G$  such that the reduced subscheme  $\overline{E}$  of  $G$  is geometrically reduced. The *closed group subscheme of  $G$  generated by  $E$*  is the group subscheme

$$\Gamma_G(E) := \Gamma_G(i),$$

where  $i : \overline{E} \hookrightarrow G$  is the inclusion of that reduced subscheme.

- (iv) Since every group subscheme of  $G$  is closed, by VI<sub>A</sub>, 0.5.2,<sup>72</sup> we shall speak of the “generated group subscheme” rather than the “generated closed group subscheme.”
- (v) Let  $X$  be a geometrically reduced  $k$ -scheme and let  $f : X \rightarrow G$  be a  $k$ -morphism. Suppose that  $f(X)$  contains the unit element  $e$  of  $G$ . Put

$$X'^1 = X \times_k X, \quad f'^1 = \mu \circ (f \times_k (c \circ f)),$$

and, for  $n > 1$ , put

$$X'^n = X'^1 \times_k X'^{n-1}, \quad f'^n = \mu \circ (f'^1 \times_k f'^{n-1}).$$

Then  $\Gamma_G(f)$  is the reduced subscheme of  $G$  whose underlying space is the closure of

$$\bigcup_{n \geq 1} f'^n(X'^n).$$

Indeed, recall the notation of 7.1(ii):

$$X^1 = X \amalg X, \quad X^n = X^1 \times_k X^{n-1} \quad (n \geq 2).$$

<sup>71</sup>Editor's note: item (viii) of the original has been placed in (i'), and items (vi) and (vii) have been singled out as Corollary 7.2.1 and Definition 7.2.2 below.

<sup>72</sup>Editor's note: the hypothesis that  $G$  be locally of finite type over  $k$  has been removed here.

The hypothesis  $e \in f(X)$  gives, for every  $n \geq 1$ ,

$$f^n(X^n) \subset f'^n(X'^n) \subset f^{2n}(X^{2n}).$$

Consequently, there exists an integer  $n$  such that  $f^n(X^n) = \Gamma_G(f)$  if and only if there exists an integer  $m$  such that  $f'^m(X'^m) = \Gamma_G(f)$ .

From Remark 7.2(v) we deduce the following result.

**Corollary 7.2.1.**— *Let  $X$  be a geometrically reduced and geometrically connected  $k$ -scheme, and let  $f : X \rightarrow G$  be a  $k$ -morphism such that  $f(X)$  contains the unit element of  $G$ . Then the  $k$ -group  $\Gamma_G(f)$  is connected, and hence irreducible.*

*Proof.* Each  $X'^n$  is connected, so the union of the

$$f'^n(X'^n),$$

all of which contain the unit element, is connected. Its closure  $\Gamma_G(f)$  is therefore connected as well. It follows that  $\Gamma_G(f)$  is irreducible, by VI<sub>A</sub>, 2.4 when  $G$ , and hence also  $\Gamma_G(f)$ , is locally of finite type over  $k$ , and by VI<sub>A</sub>, 2.6.5(iii) in the general case.

**Definition 7.2.2.**— Let  $G$  be a  $k$ -group.

- a) Let  $A$  and  $B$  be two geometrically reduced  $k$ -group subschemes of  $G$ . The *commutator group subscheme of  $A$  and  $B$  in  $G$* , denoted

$$(A, B)_G \quad \text{or simply} \quad (A, B),$$

is the closed group subscheme of  $G$  generated by the morphism

$$\nu : A \times_k B \longrightarrow G$$

defined, for every  $k$ -scheme  $S$ , by

$$(a, b) \longmapsto aba^{-1}b^{-1}, \quad a \in A(S), \quad b \in B(S).$$

- b) Suppose that  $G$  is geometrically reduced over  $k$ . The *derived group of  $G$* , denoted  $\mathfrak{D}(G)$ ,<sup>73</sup> is the group

$$\mathfrak{D}(G) = (G, G).$$

*N.B.* The group  $G$  is commutative if and only if  $\mathfrak{D}(G)$  is the unit  $k$ -group.

**Reminders 7.3.0.**—<sup>74</sup> Let  $\varphi : Y \rightarrow Z$  be a morphism of  $S$ -schemes. Recall that the image presheaf  $\varphi(Y)$  is the  $S$ -functor which assigns to every  $S'$  over  $S$  the subset

$$\varphi(Y(S')) \subset Z(S').$$

Suppose that  $T$  is a subscheme of  $Z$ . If the inclusion of presheaves

$$\varphi(Y) \hookrightarrow Z$$

factors through  $T$ , that is, if

$$\varphi \circ h \in T(S')$$

<sup>73</sup>Editor's note: the notation  $\text{Der}(G)$  of the original has been changed to avoid any risk of confusion with a space of derivations.

<sup>74</sup>Editor's note: these reminders have been added, and the statement of 7.3 has consequently been modified and expanded.



for every  $S$ -morphism  $h : S' \rightarrow Y$ , then, set-theoretically,  $\varphi(Y) \subset T$ : take  $h = \text{id}_Y$ . Conversely, if  $Y$  is reduced and  $\varphi(Y) \subset T$  set-theoretically, then  $\varphi$  factors through  $T$ .

Recall also that, by 6.2, if  $H$  is a closed  $k$ -group subscheme of  $G$ , then

$$\underline{\text{Centr}}_G H \quad \text{and} \quad \underline{\text{Norm}}_G H$$

are representable by closed  $k$ -group subschemes of  $G$ .

**Corollary 7.3.**— *Let  $G$  be a  $k$ -group, let  $X$  be a geometrically reduced  $k$ -scheme, and let  $f : X \rightarrow G$  be a  $k$ -morphism.*

(i) *Let  $S$  be a  $k$ -scheme and let  $u$  be an endomorphism of the  $S$ -group  $G_S$ .*

(a) *If*

$$u(f_S(X_S)) \subset \Gamma_G(f)_S$$

*set-theoretically, then the morphism*

$$u : \Gamma_G(f)_S \longrightarrow G_S$$

*factors through  $\Gamma_G(f)_S$ .*

(b) *If  $u$  is an automorphism of the  $S$ -group  $G_S$  and*

$$u(f_S(X_S)) = f_S(X_S)$$

*set-theoretically, then  $u$  induces an automorphism of  $\Gamma_G(f)_S$ . In particular, if  $g \in G(S)$  satisfies*

$$\text{int}(g)(f_S(X_S)) = f_S(X_S)$$

*set-theoretically, then*

$$g \in \underline{\text{Norm}}_{G_S}(\Gamma_G(f)_S)(S).$$

(c) *If  $u \circ f_S = f_S$ , then the restriction of  $u$  to the subgroup  $\Gamma_G(f)_S$  of  $G_S$  is the identity. In particular, if  $g \in G(S)$  satisfies*

$$\text{int}(g) \circ f_S = f_S,$$

*then*

$$g \in \underline{\text{Centr}}_{G_S}(\Gamma_G(f)_S)(S).$$

(ii) *Let  $H$  be a group subscheme of  $G$ . Then  $H$  centralizes (respectively, normalizes)  $\Gamma_G(f)$  if and only if, for every  $k$ -scheme  $S$  and every  $h \in H(S)$ ,*

$$\text{int}(h) \circ f_S = f_S$$

*(respectively,*

$$\text{int}(h)(f_S(X_S)) \subset \Gamma_G(f)_S$$

*set-theoretically). Equivalently, for every  $S' \rightarrow S$  and every  $x \in X(S')$ , the elements  $h_{S'}$  and  $f(x)$  of  $G(S')$  commute (respectively,*

$$h_{S'} f(x) h_{S'}^{-1} \in \Gamma_G(f)(S').$$

*).*

- (iii) In particular, let  $Y$  be a second geometrically reduced  $k$ -scheme and let  $\varphi : Y \rightarrow G$  be a  $k$ -morphism. Suppose that, for every  $k$ -scheme  $S'$ , every  $x \in X(S')$ , and every  $y \in Y(S')$ , the elements  $\varphi(y)$  and  $f(x)$  of  $G(S')$  commute (respectively,

$$\text{int}(\varphi(y))(f(x)) \in \Gamma_G(f)(S').$$

). Then  $\Gamma_G(\varphi)$  is a  $k$ -subgroup of

$$\underline{\text{Centr}}_G(\Gamma_G(f)) \quad (\text{respectively, of}) \quad \underline{\text{Norm}}_G(\Gamma_G(f)).$$

- (iv) If  $g$  is a  $k$ -point of  $G$ , then the  $k$ -group  $\Gamma_G(\{g\})$  is commutative.  
 (v) Let  $A$  and  $B$  be two group subschemes of  $G$ , geometrically reduced over  $k$ . If  $A$  and  $B$  are invariant (respectively, characteristic), then so is  $(A, B)$ .

*Proof.* We first prove (i)(a). Put

$$\Gamma_S = \Gamma_G(f)_S.$$

Then  $u^{-1}(\Gamma_S)$  is a closed  $S$ -group subscheme of  $G_S$ . By 7.1(iii), it is therefore enough to show that  $f_S$  factors through  $u^{-1}(\Gamma_S)$ . Since

$$u(f_S(X_S)) \subset \Gamma_S$$

and  $X_S$  is reduced,  $u \circ f_S$  factors through  $\Gamma_S$ ; hence  $f_S$  factors through  $u^{-1}(\Gamma_S)$ . This proves (a).

The first assertion of (b) follows by applying (a) to  $u$  and  $u^{-1}$ . In fact, it is enough to assume

$$u(f_S(X_S)) \subset \Gamma_S \quad \text{and} \quad u^{-1}(f_S(X_S)) \subset \Gamma_S.$$

The second assertion is the special case  $u = \text{int}(g)$ .

We now prove (c). Consider the morphism of  $S$ -groups

$$G_S \longrightarrow G_S \times_S G_S, \quad g \longmapsto (u(g), g),$$

and let  $H$  be the inverse image of the diagonal. It is a group subscheme of  $G_S$ . Since  $G$  is separated over  $k$ , by VI<sub>A</sub>, 0.3,  $G_S$  is separated over  $S$ , and hence  $H$  is closed in  $G_S$ . By hypothesis,  $f_S$  factors through  $H$ ; therefore  $H$  contains  $\Gamma_G(f)_S$ , by 7.1(iii). This proves the first assertion of (c), and the second is the special case  $u = \text{int}(g)$ .

We next prove (ii). Put

$$\Gamma = \Gamma_G(f)$$

and let  $i : \Gamma \hookrightarrow G$  denote the inclusion. The group  $H$  is contained in

$$N = \underline{\text{Norm}}_G(\Gamma)$$

if and only if, for every  $k$ -scheme  $S$  and every  $h \in H(S)$ ,

$$\text{int}(h)(\Gamma_S) \subset \Gamma_S.$$

Applying this condition to  $h$  and  $h^{-1}$  shows that the inclusion is then an equality. By (i)(a), this condition holds if and only if

$$\text{int}(h)(f_S(X_S)) \subset \Gamma_S.$$

Similarly,  $H$  is contained in

$$C = \underline{\text{Centr}}_G(\Gamma)$$

if and only if, for every  $k$ -scheme  $S$  and every  $h \in H(S)$ ,

$$\text{int}(h) \circ i_S = i_S.$$

By (i)(c), this holds if and only if

$$\text{int}(h) \circ f_S = f_S.$$

This proves (ii).

We now prove (iii). In view of (ii), the hypothesis implies that  $\varphi$  factors through  $C$  (respectively,  $N$ ). The latter is a closed subgroup of  $G$ , by 6.2, and therefore contains  $\Gamma_G(\varphi)$ , by 7.1(i).\*

Assertion (iv) follows from (iii) by taking both  $f$  and  $\varphi$  to be the closed immersion

$$\text{Spec } k \hookrightarrow G$$

defined by  $g$ . Indeed, for every  $k$ -scheme  $S$ , the sets  $X(S)$  and  $Y(S)$  then each have a single element, which is sent by  $f$ , respectively by  $\varphi$ , to  $g$ .

Finally, we prove (v). Let

$$\nu : G \times_k G \longrightarrow G, \quad (g, h) \longmapsto ghg^{-1}h^{-1},$$

and let  $\nu'$  be its restriction to  $A \times_k B$ . By 7.2.2,

$$(A, B) = \Gamma_G(\nu').$$

Let  $S$  be a  $k$ -scheme, and let  $u$  be an inner automorphism, respectively an arbitrary  $S$ -group automorphism, of  $G_S$ . Then

$$u \circ \nu_S = \nu_S \circ (u \times_S u).$$

The hypothesis implies that  $u$  induces automorphisms  $u_1$  of  $A_S$  and  $u_2$  of  $B_S$ . Consequently,

$$u(\nu'_S(A_S \times_S B_S)) = \nu'_S(u_1(A_S) \times_S u_2(B_S)) = \nu'_S(A_S \times_S B_S).$$

By (i)(b),  $u$  therefore induces an automorphism of  $(A, B)_S$ . This proves (v).

**Proposition 7.4.**— *Let  $G$  be a  $k$ -group locally of finite type, let  $X$  be a finite-type  $k$ -scheme that is geometrically reduced and geometrically connected, and let  $f : X \rightarrow G$  be a  $k$ -morphism such that  $f(X)$  contains the unit element  $e$  of  $G$ . Then, with the notation of 7.1(ii), there exists an integer  $N$  such that, set-theoretically,*

$$f^N(X^N) = \Gamma_G(f).$$

*Proof.* By 7.1(iii) and EGA IV<sub>2</sub>, 2.6.1, we may assume that  $k$  is algebraically closed. By Corollary 7.2.1, we may assume that  $G = G^0$ . Finally, with the notation of 7.2(v), it is enough to show that there exists an integer  $N$  such that

$$f'^N(X'^N) = \Gamma_G(f).$$

*First case.* Suppose that  $X$  is irreducible. The closed irreducible subsets

$$\overline{f'^n(X'^n)}$$

---

\*Translator's note: the French source prints  $\Gamma_G(f)$  here. Since  $\varphi$ , rather than  $f$ , is the morphism shown to factor through  $C$  (respectively,  $N$ ), 7.1(i) gives  $\Gamma_G(\varphi)$ , as required by the statement.

then form an increasing sequence in the space  $G$ , which is Noetherian because  $G = G^0$  is of finite type over  $k$  (VI<sub>A</sub>, 2.4). Hence this sequence is stationary, and there exists an integer  $m$  such that

$$\overline{f'^m(X'^m)} = \Gamma_G(f).$$

Moreover, because  $X$  and  $G$  are of finite type over  $k$ , the morphisms  $f'^n$  are of finite type over  $k$ . Consequently,  $f'^m(X'^m)$  is constructible in  $G$  (EGA IV<sub>1</sub>, 1.8.5), and therefore contains an open subset  $U$  of its closure  $\Gamma_G(f)$  (EGA 0<sub>IV</sub>, 9.2.3). By VI<sub>A</sub>, 0.5, one then has

$$\Gamma_G(f) \subset U \cdot U \subset f'^{2m}(X'^{2m}) \subset \Gamma_G(f),$$

and hence

$$f'^{2m}(X'^{2m}) = \Gamma_G(f).$$

*Second case.* Suppose that  $X$  has exactly two irreducible components  $A_1$  and  $A_2$ . Since  $X$  is connected and  $k$  is algebraically closed, there exists

$$a \in (A_1 \cap A_2)(k).$$

The four irreducible subsets

$$A_i \times_k A_j \quad (i, j = 1, 2)$$

therefore cover  $X \times_k X$ , and the image of each under the morphism

$$f'^1 = \mu \circ (f \times_k (c \circ f))$$

contains  $e$ . Let  $f'_{ij}{}^1$  denote the restriction of  $f'^1$  to the reduced subscheme  $A_i \times_k A_j$ , and put

$$\begin{aligned} Y &= (A_1 \times_k A_1) \times_k (A_1 \times_k A_2) \times_k (A_2 \times_k A_2) \times_k (A_2 \times_k A_1), \\ g &= \mu \circ \left( \mu \circ (f'_{11}{}^1 \times_k f'_{12}{}^1) \times_k \mu \circ (f'_{22}{}^1 \times_k f'_{21}{}^1) \right). \end{aligned}$$

The scheme  $Y$  is irreducible, reduced, and of finite type. By the first case, there is therefore an integer  $m$  such that

$$g'^m(Y'^m) = \Gamma_G(g).$$

For every  $n \geq 1$ , however,\*

$$f'^n(X'^n) \subset g'^n(Y'^n) \subset f'^{4n}(X'^{4n}).$$

It follows that

$$\Gamma_G(f) = \Gamma_G(g) \quad \text{and} \quad f'^{4m}(X'^{4m}) = \Gamma_G(f).$$

*General case.* We argue by induction on the number of irreducible components of  $X$ ; this number is finite because  $X$ , being of finite type over  $k$ , is Noetherian. Suppose the proposition has been proved when  $X$  has at most  $r-1$  irreducible components, and suppose that  $X$  has  $r$  such components, say  $A_1, \dots, A_r$ . The unit  $e$  belongs to the image of one of the  $A_i$ ; suppose, for example, that

$$e \in f(A_1).$$

Put

$$X_r = A_1 \cup \dots \cup A_{r-1}, \quad Y = \Gamma_G(f_r),$$

---

\*Translator's note: the French source prints  $g'^n(Y'^n)$  in the middle term. Since the source of  $g'^n$  is  $Y'^n$ , the prime is restored here; compare the preceding equality  $g'^m(Y'^m) = \Gamma_G(g)$ .

where  $f_r$  denotes the restriction of  $f$  to the reduced subscheme  $X_r$ . We choose the numbering of the  $A_i$  so that  $X_r$  is connected, which is always possible. Then  $Y$  is a closed, reduced, irreducible subgroup of  $G$ , by Corollary 7.2.1.

Put

$$Z = \overline{f(A_r)}$$

and let  $T = Y \cup Z$ , endowed with its closed reduced subscheme structure; let  $i : T \hookrightarrow G$  be the inclusion. It follows immediately from 7.2(ii) that

$$\Gamma_G(i) = \Gamma_G(f).$$

Moreover,  $T$  is connected: since  $X$  is connected,  $X_r$  and  $A_r$  have a common point  $a$ , and  $Y$  and  $Z$  have the common point  $f(a)$ . Also,  $e \in T$ , and  $T$  has at most two irreducible components, because  $Y$  and  $Z$  are irreducible.

By the induction hypothesis, there exists an integer  $m$  such that

$$f_r'^m(X_r'^m) = \Gamma_G(f_r) = Y.$$

Since  $X = X_r \cup A_r$ , and since

$$Z = \overline{f(A_r)} \subset \overline{f'^m(X'^m)}$$

because  $e \in f(A_1)$ , we obtain

$$f(X) \subset Y \cup Z \subset \overline{f'^m(X'^m)}.$$

On the other hand, because  $T$  has at most two irreducible components, the second case gives an integer  $p$  such that

$$i'^p(T'^p) = \Gamma_G(i) = \Gamma_G(f).$$

Now

$$T \subset \overline{f'^m(X'^m)},$$

and hence

$$\overline{f'^{mp}(X'^{mp})} = \Gamma_G(f).$$

As in the first case, this last equality implies

$$f'^{2mp}(X'^{2mp}) = \Gamma_G(f).$$

This completes the proof.

**Lemma 7.5.**— *Let  $S$  be a scheme, let  $G$  be an  $S$ -group scheme, let  $X$  be an  $S$ -prescheme, and let  $f : X \rightarrow G$  be an  $S$ -morphism. Suppose that  $X$  and  $G$  are locally of finite presentation over  $S$ , and that, for every  $s \in S$  and every maximal point  $g$  of  $G_s$ , there exists a point  $x$  of  $X$  such that  $f(x) = g$  and  $f$  is flat at  $x$ . Then the morphism*

$$\mu \circ (f \times_S f) : X \times_S X \longrightarrow G$$

*is a covering for the (fppf) topology.*

<sup>75</sup> By EGA IV<sub>3</sub>, 11.3.1, the set  $V$  of points of  $X$  at which  $f$  is flat is open, and  $f|_V$  is an open morphism. Thus

$$U = f(V)$$

---

<sup>75</sup>Editor's note: the argument in the original has been expanded in what follows.

is an open subset of  $G$ . Moreover, by hypothesis,

$$U \cap G_s$$

is dense in  $G_s$  for every  $s \in S$ . Let  $\phi$  denote the restriction of  $\mu \circ (f \times_S f)$  to  $V \times_S V$ .

It is enough to show that  $\phi$  is a covering for the (fppf) topology, and for this it is enough to show that  $\phi$  is faithfully flat and of finite presentation (Exposé IV, 6.3.1(iv)). Now  $\phi$  is the composite

$$V \times_S V \xrightarrow{f|_V \times f|_V} U \times_S U \xrightarrow{\mu} G,$$

where the first morphism is faithfully flat and locally of finite presentation, since  $f|_V$  is. It therefore remains only to prove the same assertions for the restriction of  $\mu$  to  $U \times_S U$ .

Since  $G \rightarrow S$  is locally of finite presentation and flat, so is

$$\mu : G \times_S G \longrightarrow G.$$

Indeed, this morphism is isomorphic to the morphism obtained from  $G \rightarrow S$  by the base change  $G \rightarrow S$  (VI<sub>A</sub>, 0.1). The induced morphism

$$U \times_S U \longrightarrow G$$

therefore has the same properties. To prove that this last morphism is surjective, it is enough to check fiber by fiber. There the result follows from VI<sub>A</sub>, 0.5, since  $U \cap G_s$  is a dense open subset of  $G_s$  for every  $s \in S$ .

**Remark 7.6.0.**—<sup>76</sup> Let  $S$  be a scheme, let  $H$  be an  $S$ -group, and let  $f : X \rightarrow H$  be a morphism of  $S$ -schemes. The group subsheaf of  $H$  generated by the image of  $f$ , denoted

$$\langle \text{Im } f \rangle,$$

is the group subfunctor of  $H$  which assigns to every  $S$ -scheme  $S'$  the subgroup of  $H(S')$  generated by  $f(X(S'))$ . Since every element of this subgroup can be written as a finite product

$$f(x_1)^{\varepsilon_1} \cdots f(x_n)^{\varepsilon_n}, \quad n \in \mathbb{N}^*, \quad x_i \in X(S'), \quad \varepsilon_i = \pm 1,$$

it follows that, if one puts

$$X^1 = X \amalg X$$

and defines the morphisms  $f^n : X^n \rightarrow H$  as in 7.1, then  $\langle \text{Im } f \rangle$  is precisely the image presheaf of the morphism

$$f^\infty : X^\infty \longrightarrow H,$$

where

$$X^\infty = \coprod_{n \geq 1} X^n$$

and  $f^\infty : X^\infty \rightarrow H$  is the  $S$ -morphism whose restriction to each  $X^n$  is  $f^n$ .

**Proposition 7.6.**—<sup>77</sup> Let  $S$  be a scheme, let  $X$  be an  $S$ -scheme flat and locally of finite presentation over  $S$ , let  $H$  be an  $S$ -group locally of finite presentation over  $S$  and with reduced fibers, let  $f : X \rightarrow H$  be a morphism of  $S$ -schemes, and let  $f^\infty : X^\infty \rightarrow H$  be the  $S$ -morphism introduced above. The following conditions are equivalent:

<sup>76</sup>Editor's note: this definition has been added; in the original it was contained in the statement of Proposition 7.6.

<sup>77</sup>Editor's note: the original stated this result under the hypotheses of the special case, but the more general form was used implicitly in the proof of 10.12; the statement has been rewritten accordingly.

- (i)  $H$  represents the  $S$ -sheaf for the (fppf) topology associated with the presheaf  $\langle \text{Im } f \rangle$ .
- (ii)  $f^\infty$  is a covering for the (fppf) topology.
- (iii)  $f^\infty$  is surjective, that is,

$$H = \bigcup_{n \geq 1} f^n(X^n).$$

If, in addition,  $H$  is quasi-compact, these conditions are also equivalent to the following one:

- (iv) There exists an integer  $n$  such that  $f^n : X^n \rightarrow H$  is a covering for the (fppf) topology, and hence is surjective.

This applies in particular when  $X$  is a geometrically reduced  $k$ -scheme locally of finite type,  $G$  is a  $k$ -group locally of finite type,  $\varphi : X \rightarrow G$  is a morphism of  $k$ -schemes,  $H = \Gamma_G(\varphi)$ , and  $f : X \rightarrow H$  is the morphism induced by  $\varphi$ .

*Proof.* The sheaf considered in (i) is the image sheaf of  $X^\infty$  under  $f^\infty$ . Hence, by IV, 4.4.3, saying that  $H$  represents this sheaf is equivalent to saying that  $f^\infty$  is a covering; this in turn implies that  $f^\infty$  is surjective. Conversely, suppose that  $f^\infty$  is surjective. We shall show that it is then a covering for the (fppf) topology.

Let  $s \in S$ , let  $\eta$  be a maximal point of the fiber  $H_s$ , and let  $x \in X_s^\infty$  be such that

$$f^\infty(x) = \eta;$$

such an  $x$  exists because  $f^\infty$  is surjective. Since the fiber  $H_s$  is reduced,  $\mathcal{O}_{H_s, \eta}$  is a field, and therefore  $f_s^\infty$  is flat at  $x$ . Since  $X^\infty$  and  $H$  are locally of finite presentation over  $S$ , and  $X^\infty$  is flat over  $S$ , the fiberwise criterion for flatness (EGA IV<sub>3</sub>, 11.3.10) shows that  $f^\infty$  is flat at  $x$ . Thus, by Lemma 7.5, the morphism

$$\mu \circ (f^\infty \times_S f^\infty) : X^\infty \times_S X^\infty \longrightarrow H$$

is a covering for the (fppf) topology.

Now  $X^n \times_S X^m$  is canonically isomorphic to  $X^{n+m}$ , and, under this isomorphism,

$$\mu \circ (f^n \times_S f^m)$$

corresponds to  $f^{n+m}$ . It is therefore clear that

$$\mu \circ (f^\infty \times_S f^\infty)$$

factors through  $f^\infty$ , so  $f^\infty$  is a covering for the (fppf) topology. This proves (iii)  $\Rightarrow$  (ii), and hence the equivalence of conditions (i), (ii), and (iii).

Furthermore, since the morphisms  $X \rightarrow S$  and  $H \rightarrow S$  are locally of finite presentation, so is  $f : X \rightarrow H$  (EGA IV<sub>1</sub>, 1.4.3(v)). The multiplication

$$\mu : H \times_S H \longrightarrow H$$

is also locally of finite presentation (VI<sub>A</sub>, 0.1), and it follows that each  $f^n : X^n \rightarrow H$  is locally of finite presentation. Consequently, by EGA IV<sub>1</sub>, 1.9.5(viii), the sets

$$f^n(X^n)$$

are ind-constructible subsets of  $H$ .

Suppose, in addition, that  $H$  is quasi-compact; then  $S$  is also quasi-compact, since  $H \rightarrow S$  is surjective. By EGA IV<sub>1</sub>, 1.9.9, there exists an integer  $p$  such that

$$f^p(X^p) = H.$$

As above, the fact that the fibers of  $H$  are reduced, together with Lemma 7.5, shows that the morphism

$$\mu \circ (f^p \times_S f^p) : X^p \times_S X^p \longrightarrow H$$

is a covering for the (fppf) topology. Since this morphism is

$$f^{2p} : X^{2p} \longrightarrow H,$$

this completes the proof of Proposition 7.6.

**Remark 7.6.1.**— The equivalent conditions of 7.6 plainly imply that the sheaf  $F$  under consideration is representable. The converse is false in general:<sup>78</sup> for example, if  $k$  has characteristic 0, let  $G = \mathbb{G}_{a,k}$ , and let

$$f : \operatorname{Spec} k \longrightarrow G$$

be the morphism defined by the point 1 of  $G(k)$ . Then  $F$  is represented by the constant  $k$ -group  $\mathbb{Z}_k$ , whereas  $\Gamma_G(f) = G$ ; hence the monomorphism

$$\mathbb{Z}_k \hookrightarrow G$$

is not surjective.

For simplicity, suppose now that the hypotheses of the special case in 7.6 hold, and suppose that  $F$  is representable. By EGA IV<sub>3</sub>, 8.14.2, the sheaf  $F$  is locally of finite presentation over  $k$ . The question is therefore whether the dominant monomorphism

$$F \longrightarrow \Gamma_G(f)$$

is an isomorphism, or, equivalently, a closed immersion. By 1.4.2, this is so if  $F$  is quasi-compact; and by VI<sub>A</sub>, 0.5.1, it is so if  $F$  is connected. In particular, by 7.2.1, it is so when  $X$  is connected and  $f(X)$  contains the unit element of  $G$ .

**Lemma 7.7.**— *Let  $k$  be an algebraically closed field, let  $G$  be a  $k$ -group locally of finite type, let  $X$  be a geometrically reduced  $k$ -scheme locally of finite type, let  $f : X \rightarrow G$  be a  $k$ -morphism, and let  $H$  be a group subscheme of  $G$  such that  $H \subset f(X)$ . Put*

$$\Gamma' = \bigcup_{n \geq 1} f^n(X^n), \quad \Gamma'_0 = \Gamma' \cap G(k), \quad H_0 = H(k),$$

*and suppose that  $H_0$  has finite index in  $\Gamma'_0$ . Then there exists an integer  $m$  such that*

$$f^m(X^m) = \Gamma_G(f)$$

*(cf. 7.6), and  $\Gamma_G(f)$  is the union of finitely many translates of  $H$ .*

For every  $n \geq 1$ , the subset  $f^n(X^n)$  of  $G$  is ind-constructible (EGA IV<sub>1</sub>, 1.9.5(viii)). The same is therefore true of  $\Gamma'$ . Since  $G$  is a Jacobson scheme,  $\Gamma'_0$  is dense in  $\Gamma'$ . By hypothesis, there are finitely many points

$$a_1, \dots, a_r \in \Gamma'_0$$

<sup>78</sup>Editor's note: the argument in the original has been expanded in what follows.



such that

$$\Gamma'_0 = a_1 H_0 \cup \cdots \cup a_r H_0.$$

Consequently,

$$\begin{aligned} \Gamma_G(f) &= \overline{\Gamma'} = \overline{\Gamma'_0} = \overline{a_1 H_0 \cup \cdots \cup a_r H_0} \\ &= \overline{a_1 H_0} \cup \cdots \cup \overline{a_r H_0} \\ &= a_1 \overline{H_0} \cup \cdots \cup a_r \overline{H_0}, \end{aligned}$$

where the last equality follows because translation by each  $a_i$  is a homeomorphism of  $G$  onto itself. Thus

$$\Gamma_G(f) = a_1 H \cup \cdots \cup a_r H.$$

It is also clear that there exists an integer  $p$  such that every  $a_i$ , for  $1 \leq i \leq r$ , belongs to  $f^p(X^p)$ . Finally, since  $H \subset f(X)$ , one has

$$a_i H \subset f^{p+1}(X^{p+1})$$

for every  $i$ . Hence

$$f^{p+1}(X^{p+1}) = \Gamma_G(f).$$

**Proposition 7.8.**— *Let  $G$  be a  $k$ -group locally of finite type, and let  $A$  and  $B$  be two geometrically reduced  $k$ -group subschemes of  $G$  (hence smooth at the generic points of their irreducible components, and therefore smooth by 1.3). Suppose that one of the following conditions holds:*

- a)  *$A$  and  $B$  are invariant and of finite type over  $k$ ;*
- b)  *$A$  is connected and  $B$  is of finite type over  $k$ .*

*Then  $(A, B)$  is of finite type over  $k$ , and represents the sheaf associated, for the (fppf) topology (or the (fpqc) topology), with the group presheaf of commutators of  $A$  and  $B$  in  $G$ . Moreover,<sup>79</sup> the  $k$ -groups  $(A, B^0)$  and  $(A^0, B)$  are connected, and*

$$(A, B)^0 = (A, B^0) \cdot (A^0, B).$$

*Proof.* By 7.6, to show that  $(A, B)$  is the desired associated sheaf, it is enough to show that there exists an integer  $n$  such that

$$\nu^n((A \times_k B)^n) = (A, B),$$

with the notation of 7.2.2. To prove this and the other two assertions, we may suppose that  $k$  is algebraically closed.

Indeed, let  $\bar{k}$  be an algebraic closure of  $k$ . By 7.1(iii) and VI<sub>A</sub>, 2.4, one has, with the evident notation,

$$\begin{aligned} (A, B)_{\bar{k}} &= (A_{\bar{k}}, B_{\bar{k}}), \\ ((A, B)^0)_{\bar{k}} &= (A_{\bar{k}}, B_{\bar{k}})^0, \\ (A, B^0)_{\bar{k}} &= (A_{\bar{k}}, B_{\bar{k}}^0), \quad \text{and so on.} \end{aligned}$$

Consequently, suppose that  $(A_{\bar{k}}, B_{\bar{k}})$  is of finite type over  $\bar{k}$ ; respectively, suppose that  $(A_{\bar{k}}, B_{\bar{k}}^0)$  and  $(A_{\bar{k}}^0, B_{\bar{k}})$  are connected and that the morphism\*

$$(A_{\bar{k}}, B_{\bar{k}}^0) \times_{\bar{k}} (A_{\bar{k}}^0, B_{\bar{k}}) \longrightarrow (A_{\bar{k}}, B_{\bar{k}})^0$$

<sup>79</sup>Editor's note: the assertions that follow have been made explicit, and the reduction to the algebraically closed case has been expanded in the proof.

\* Translator's note: the French source prints  $(A_{\bar{k}}, B_{\bar{k}}^0)$  in both factors of this product. The second factor must be  $(A_{\bar{k}}^0, B_{\bar{k}})$ , as in the asserted identity and throughout the proof; the mixed product is restored here.

is surjective. The analogous assertions then hold over  $k$ , by EGA IV<sub>2</sub>, 2.7.1 and 2.6.1.

Let

$$B^1, \dots, B^p$$

be the connected components of  $B$  other than the identity component  $B^0$ . There are only finitely many, since  $B$  is of finite type over  $k$ , hence Noetherian. In case a), likewise let

$$A^1, \dots, A^q$$

be the connected components of  $A$  other than  $A^0$ . In case b), only  $A^0 = A$  is considered. Let  $\nu_{ij}$  be the restriction of  $\nu$  to

$$A^i \times_k B^j.$$

Each  $A^i$  and  $B^j$  is irreducible, by VI<sub>A</sub>, 2.4.1; hence so are  $A^0 \times_k B^j$  and  $A^i \times_k B^0$ . Since the unit element of  $G$  belongs to  $A^0$  and to  $B^0$ , it belongs to

$$\nu_{0j}(A^0 \times_k B^j) \quad \text{and} \quad \nu_{i0}(A^i \times_k B^0).$$

Thus every  $\Gamma_G(\nu_{0j})$  and every  $\Gamma_G(\nu_{i0})$  is connected, by 7.2.1.

Let  $u_{0j}$ , respectively  $u_{i0}$ , denote the inclusion of  $\Gamma_G(\nu_{0j})$ , respectively  $\Gamma_G(\nu_{i0})$ , in  $G$ . Then

$$(A^0, B) = \Gamma_G((u_{0j})_{j=0}^p), \quad (A, B^0) = \Gamma_G((u_{i0})_{i=0}^q),$$

and these groups are connected. Moreover, it follows from 7.4 and the preceding constructions that there is an integer  $r$  such that both  $(A^0, B)$  and  $(A, B^0)$  are contained in

$$\nu^r((A \times_k B)^r).$$

In case b), one has  $(A, B) = (A^0, B)$ , and the proof is complete.

We now consider case a).<sup>80</sup> We already know that  $(A^0, B)$  and  $(A, B^0)$  are smooth connected  $k$ -subgroups of  $G$ , hence are of finite type over  $k$  (VI<sub>A</sub>, 2.4). Since  $A^0$  is a characteristic subgroup of  $A$  (VI<sub>A</sub>, 2.6.5), it is an invariant subgroup of  $G$ ; hence 7.3(v) shows that  $(A^0, B)$  is an invariant subgroup of  $G$ , and the same holds for  $(A, B^0)$ .

By 7.1.1, the subgroup  $H$  of  $G$  generated by  $(A, B^0)$  and  $(A^0, B)$  is therefore

$$H = (A, B^0) \cdot (A^0, B).$$

In particular,

$$H \subset \nu^{2r}((A \times_k B)^{2r}).$$

If  $T$  is a subset of  $G$  stable under the group law (cf. 3.0), write  $T_0$  for the group of  $k$ -points of  $G$  belonging to  $T$ . Put

$$\Gamma' = \bigcup_{q \geq 1} \nu^q((A \times_k B)^q).$$

By Proposition 7.9 below,

$$(A^0, B)_0 = (A_0^0, B_0), \quad (A, B^0)_0 = (A_0, B_0^0), \quad \Gamma'_0 = (A_0, B_0).$$

Therefore

$$H_0 = (A_0^0, B_0) \cdot (A_0, B_0^0)$$

has finite index in  $\Gamma'_0$  (Bible, Exposé III, Appendix), since  $A_0$  and  $B_0$  are invariant and  $A_0^0$ , respectively  $B_0^0$ , has finite index in  $A_0$ , respectively  $B_0$ .

<sup>80</sup>Editor's note: the argument that follows has been expanded to take account of the added Corollary 7.1.1.

The hypotheses of Lemma 7.7 now hold. Since

$$H \subset \nu^{2r}((A \times_k B)^{2r}),$$

there exists an integer  $m$  such that

$$\nu^{2rm}((A \times_k B)^{2rm}) = (A, B),$$

and there are finitely many  $k$ -points  $a_1, \dots, a_n$  of  $G$  such that

$$(A, B) = a_1 H \cup \dots \cup a_n H.$$

Since  $H$  is of finite type over  $k$ , every  $a_i H$  is quasi-compact. Their union  $(A, B)$  is therefore quasi-compact, hence of finite type over  $k$ . Finally,  $H$  is irreducible, so each  $a_i H$  is irreducible; and because the unit  $e$  belongs to  $H$ , it follows that

$$H = (A, B)^0.$$

**Proposition 7.9.**—*Let  $k$  be an algebraically closed field, and let  $G$  be a  $k$ -group locally of finite type.*

- (i) *Let  $A$  and  $B$  be two ind-constructible subsets of  $G$ . Denote by  $A_0$  the set of rational points of  $G$  that belong to  $A$ . Then*

$$(A \cdot B)_0 = A_0 \cdot B_0,$$

*where the product on the right is taken in the group  $G(k)$ .*

- (ii) *Let  $X$  be a geometrically reduced  $k$ -scheme locally of finite type, and let  $f : X \rightarrow G$  be a  $k$ -morphism. Put*

$$\Gamma'_G(f) = \bigcup_{n \geq 1} f^n(X^n).$$

*Then  $\Gamma'_G(f)_0$  is the subgroup of  $G(k)$  generated by  $f(X)_0$ .*

- (iii) *In particular, let  $A$  and  $B$  be two smooth group subschemes of  $G$ , and put*

$$\Gamma' = \bigcup_{n \geq 1} \nu^n((A \times_k B)^n)$$

*with the notation of 7.2.2. Then  $\Gamma'_0$  is the commutator subgroup of  $A(k)$  and  $B(k)$  in  $G(k)$ .*

*Proof.* We first prove (i). It is clear that

$$A_0 \cdot B_0 \subset (A \cdot B)_0.$$

Conversely, let  $z \in (A \cdot B)_0$ . Then  $\mu^{-1}(z)$  is a closed subset of  $G \times_k G$ , while  $A \times_k B$  is an ind-constructible subset of  $G \times_k G$  (cf. 3.0). Consequently,

$$\mu^{-1}(z) \cap (A \times_k B)$$

is a nonempty ind-constructible subset of  $G \times_k G$ . By EGA IV<sub>3</sub>, 10.4.8, it therefore contains a rational point of  $G \times_k G$ . Its projections  $x$  and  $y$  are rational points of  $G$  such that  $x \in A$ ,  $y \in B$ , and  $x \cdot y = z$ . Hence \*

$$(A \cdot B)_0 = A_0 \cdot B_0.$$

---

\*Translator's note: the French proof prints  $(A, B)_0 = A_0 \cdot B_0$  here. The statement and the multiplication argument require  $(A \cdot B)_0 = A_0 \cdot B_0$ , which is restored in the English text.

To prove (ii), observe that, since  $f^n$  is locally of finite type,  $f^n(X^n)$  is an ind-constructible subset of  $G$  (EGA IV<sub>1</sub>, 1.9.5(viii))<sup>†</sup>. Assertion (i) then shows by induction that, if

$$A = f(X)_0,$$

one has

$$f^n(X^n)_0 = (A \cup A^{-1})^n.$$

Consequently,

$$\Gamma'_G(f)_0 = \bigcup_{n \geq 1} f^n(X^n)_0 = \bigcup_{n \geq 1} (A \cup A^{-1})^n,$$

which is the subgroup of  $G(k)$  generated by  $A = f(X)_0$ .

Finally, (iii) follows from (ii) and the definitions.

**Corollary 7.10.**—*Under the hypotheses of 7.8, if  $k$  is algebraically closed, then  $(A, B)(k)$  is the commutator subgroup of  $A(k)$  and  $B(k)$  in  $G(k)$ .*

Indeed, it is enough to apply 7.9(iii), 7.8, and 7.6.

## 8. Solvable or nilpotent group schemes

**8.1.**— Let  $\mathcal{C}$  be a category equipped with a topology  $T$  (cf. Exposé IV, §4). For every presheaf  $P$  on  $\mathcal{C}$ , write  $P^b$  for the associated sheaf.

Let  $G$  be a presheaf of groups on  $\mathcal{C}$ , let  $A$  and  $B$  be two group subsheaves of  $G$ , and let  $\text{Comm}(A, B)$  be the group presheaf of commutators of  $A$  and  $B$  in  $G$ . Thus, for every  $S \in \text{Ob } \mathcal{C}$ ,  $\text{Comm}(A, B)(S)$  is the subgroup of  $G(S)$  generated by the commutators

$$aba^{-1}b^{-1}, \quad a \in A(S), \quad b \in B(S).$$

We write

$$\text{Comm}_T(A, B) = \text{Comm}(A, B)^b.$$

In the proof of 8.2, we shall need the following results.<sup>81</sup>

**Lemma 8.1.1.**—*Let  $A \subset G$  be sheaves of groups, with  $A$  invariant in  $G$ .*

- (i)  *$\text{Comm}_T(G, A)$  is the smallest invariant group subsheaf  $C$  of  $G$  such that the sheaf  $(A/C)^b$ , associated with the quotient presheaf  $A/C$ , is central in  $(G/C)^b$ .*
- (ii) *In particular,  $\text{Comm}_T(G, G)$  is the smallest invariant group subsheaf  $C$  of  $G$  such that the quotient sheaf  $(G/C)^b$  is commutative.*

Clearly, (ii) is the special case  $A = G$  of (i), so it is enough to prove (i). Let  $C$  be a group subsheaf of  $A$ , invariant in  $G$ , such that the quotient sheaf  $(A/C)^b$  is central in  $(G/C)^b$ . By Lemma IV, 4.4.8.1, the presheaves  $A/C$  and  $G/C$  are separated. Hence, by IV, 4.3.11, all the morphisms in the following diagram are monomorphisms:

$$\begin{array}{ccc} A/C & \hookrightarrow & G/C \\ \downarrow & & \downarrow \\ (A/C)^b & \hookrightarrow & (G/C)^b \end{array}$$

<sup>†</sup> Translator's note: the French source prints EGA IV<sub>4</sub> here. The cited result 1.9.5(viii) belongs to EGA IV<sub>1</sub>, which is restored in the English text.

<sup>81</sup>Editor's note: these results were stated without proof in the original. We have highlighted them as Lemmas 8.1.1 and 8.1.2 and supplied the details of the proofs.

Since  $(A/C)^b$  is central in  $(G/C)^b$ , the presheaf  $A/C$  is central in  $G/C$ . It follows that

$$\mathrm{Comm}(G, A) \subset C,$$

and hence, by IV, 4.3.12,  $C$  contains  $\mathrm{Comm}_T(G, A)$ .

Conversely,  $\mathrm{Comm}(G, A)$  is an invariant, separated group subpresheaf of  $A$  (cf. IV, 4.3.1, editor's note 24). Therefore, by IV, 4.4.8.2(i) and IV, 4.3.11,

$$C = \mathrm{Comm}_T(G, A)$$

is an invariant group subsheaf of  $A$  containing  $\mathrm{Comm}(G, A)$ . Consequently, the presheaf  $A/C$  is central in  $G/C$ , and hence, by IV, 4.4.8.2(ii),  $(A/C)^b$  is central in  $(G/C)^b$ . This proves Lemma 8.1.1.

**Lemma 8.1.2.**— *Let  $G$  be a sheaf of groups, and let  $A, B$  be two group subpresheaves of  $G$ .*

(i) *The morphism*

$$\tau : \mathrm{Comm}(A, B) \longrightarrow \mathrm{Comm}(A^b, B^b)$$

*is a covering monomorphism.*

(ii) *Consequently, there is an isomorphism*

$$\mathrm{Comm}_T(A, B) \xrightarrow{\sim} \mathrm{Comm}_T(A^b, B^b).$$

*Proof.* (i) Since  $A$  (respectively,  $B$ ) is a subpresheaf of  $G$ , the associated sheaf  $A^b$  (respectively,  $B^b$ ) is a subpresheaf of  $G$  containing  $A$  (respectively,  $B$ ). It follows that  $\tau$  is a monomorphism.

We now show that  $\tau$  is a covering. Let  $S \in \mathrm{Ob} \mathcal{C}$  and let

$$g \in \mathrm{Comm}(A^b, B^b)(S).$$

There exist an integer  $n \geq 1$  and, for  $1 \leq i \leq n$ , elements

$$a'_i \in A^b(S), \quad b'_i \in B^b(S), \quad \varepsilon_i \in \{\pm 1\},$$

such that

$$g = (a'_1, b'_1)^{\varepsilon_1} \cdots (a'_n, b'_n)^{\varepsilon_n},$$

where  $(a, b)$  denotes the commutator  $aba^{-1}b^{-1}$ . There is a refinement  $R$  of  $S$  such that

$$a'_i \in A(R) \quad \text{and} \quad b'_i \in B(R)$$

for every  $i$ . Then  $g$  is the composite morphism

$$R \xrightarrow{(a'_1, \dots, b'_n)} (A \times B)^n \xrightarrow{\Phi^{\varepsilon_1, \dots, \varepsilon_n}} \mathrm{Comm}(A, B)$$

where  $\Phi = \Phi^{\varepsilon_1, \dots, \varepsilon_n}$  is the morphism defined set-theoretically by

$$\Phi(a_1, b_1, \dots, a_n, b_n) = (a_1, b_1)^{\varepsilon_1} \cdots (a_n, b_n)^{\varepsilon_n}$$

for every  $T \in \mathrm{Ob} \mathcal{C}$ , with  $a_i \in A(T)$  and  $b_i \in B(T)$ . This shows that

$$g \in \mathrm{Comm}(A, B)(R).$$

It follows, as in the proof of IV, 4.3.11(i), that

$$\mathrm{Comm}(A, B) \longrightarrow \mathrm{Comm}(A^b, B^b)$$

is a covering.

The morphism

$$\mathrm{Comm}(A^b, B^b) \longrightarrow \mathrm{Comm}(A^b, B^b)^b$$

is likewise a covering monomorphism (IV, 4.3.11(iv)). Hence the same is true of

$$\mathrm{Comm}(A, B) \longrightarrow \mathrm{Comm}(A^b, B^b)^b.$$

By IV, 4.3.12, we therefore obtain an isomorphism

$$\mathrm{Comm}(A, B)^b \xrightarrow{\sim} \mathrm{Comm}(A^b, B^b)^b.$$

This proves Lemma 8.1.2.

**Proposition 8.2.**— *Let  $\mathcal{C}$  be a category, let  $T$  be a topology on  $\mathcal{C}$ , let  $G$  be a sheaf of groups on  $\mathcal{C}$ , and let  $n$  be an integer with  $n \geq 0$ . The following conditions are equivalent:*

- (i) *Put  $K_0 = G$ , and, for  $p \geq 1$ , put*

$$K_p = \mathrm{Comm}(K_{p-1}, K_{p-1}) \quad (\text{respectively, } K_p = \mathrm{Comm}(G, K_{p-1})).$$

*Then  $K_n$  is the unit presheaf of groups.*

- (ii) *Put  $K'_0 = G$ , and, for  $p \geq 1$ , put*

$$K'_p = \mathrm{Comm}_T(K'_{p-1}, K'_{p-1}) \quad (\text{respectively, } K'_p = \mathrm{Comm}_T(G, K'_{p-1})).$$

*Then  $K'_n$  is the unit sheaf of groups.*

- (iii) *There is a sequence*

$$G = H_0 \supset H_1 \supset \cdots \supset H_n$$

*of invariant group subsheaves of  $G$  such that, for every  $i$ , the quotient sheaf  $H_i/H_{i+1}$  is commutative (respectively, central in  $G/H_{i+1}$ ), and  $H_n$  is the unit sheaf.*

It is clear that  $K_n \subset K'_n$ ; consequently, (ii) implies (i). We show that (i) implies (ii). One has

$$K'_1 = \mathrm{Comm}_T(G, G) = K_1^b,$$

and induction using Lemma 8.1.2 gives

$$K'_p = K_p^b$$

for every  $p$ . Consequently, if  $K_n$  is the unit presheaf, then  $K'_n = K_n^b$  is the unit sheaf.

Finally, conditions (ii) and (iii) are equivalent by Lemma 8.1.1.

**Definition 8.2.1.**— *When these conditions hold, the sheaf  $G$  is said to be solvable of class  $n$  (respectively, nilpotent of class  $n$ ). If there is an integer  $n$  for which these conditions hold,  $G$  is said to be solvable (respectively, nilpotent).*

Observe that, by condition (i), this does not depend on the topology  $T$ .

**Proposition 8.3.**— *Let  $k$  be a field, let  $S$  be a nonempty  $k$ -scheme, let  $\Omega$  be an algebraically closed extension of  $k$ , and let  $G$  be a smooth  $k$ -group of finite type. The following conditions are equivalent:*

- (i)  $G$  is solvable of class  $n$  (respectively, nilpotent of class  $n$ ).
- (ii)  $G \times_k S$  is solvable of class  $n$  (respectively, nilpotent of class  $n$ ).
- (iii) The group  $G(\Omega)$  is solvable of class  $n$  (respectively, nilpotent of class  $n$ ).
- (iv) Put  $K_0 = G$ , and, for  $p \geq 1$ , consider the  $k$ -groups

$$K_p = (K_{p-1}, K_{p-1}) \quad (\text{respectively, } K_p = (G, K_{p-1}))$$

(cf. 7.2.2). Then  $K_n$  is the unit  $k$ -group.

The equivalence of conditions (i) and (ii) follows from Proposition 8.2, since the formation of the group presheaf of commutators commutes with base change (IV, 4.1.3).

The equivalence of (i) and (iv) follows from the fact that, by 7.8, the  $k$ -group

$$K_p = (K_{p-1}, K_{p-1}) \quad (\text{respectively, } K_p = (G, K_{p-1}))$$

represents the sheaf

$$\text{Comm}_T(K_{p-1}, K_{p-1}) \quad (\text{respectively, } \text{Comm}_T(G, K_{p-1})),$$

where  $T$  is the (fppf) or (fpqc) topology.

To prove the equivalence of conditions (iii) and (iv), we may assume that  $k = \Omega$ ; the result then follows from 7.10.

**Proposition 8.4.**— *Let  $G$  be an  $S$ -group of finite presentation such that, for every  $s \in S$ , the fiber  $G_s$  is smooth over  $\kappa(s)$ . Let  $T$  be the set of points  $s \in S$  for which  $G_s$  is solvable (respectively, nilpotent).*

- (i) *Then  $T$  is locally constructible in  $S$ .*
- (ii) *If, in addition,  $G$  is assumed flat and separated over  $S$  (that is, when  $G$  is smooth, quasi-compact, and separated over  $S$ ), then  $T$  is closed in  $S$ .*

*Proof.* We may clearly suppose that  $S$  is affine, with ring  $A$ . By 10.1 and 10.10b),\* there exist a Noetherian subring  $A'$  of  $A$  and an  $A'$ -group  $G'$  of finite type such that

$$G' \otimes_{A'} A \simeq G.$$

By EGA IV<sub>3</sub>, 11.2.6 and 8.10.5,<sup>82</sup> if  $G$  is flat and separated over  $S$ , we may suppose that  $G'$  is flat and separated over  $S' = \text{Spec } A'$ .<sup>83</sup> Since  $G$  is of finite presentation over  $S$ , the set of points  $s \in S$  for which  $G_s$  is geometrically reduced—or, equivalently, smooth over  $\kappa(s)$ —is locally constructible (EGA IV<sub>3</sub>, 9.7.7). Hence, by EGA IV<sub>3</sub>, 9.3.3, we may suppose that  $G'_{s'}$  is smooth over  $\kappa(s')$  for every  $s' \in S'$ . On the other hand, if  $s'$  denotes the image of  $s$  in  $S'$ , then

$$G'_{s'} \otimes_{\kappa(s')} \kappa(s) \simeq G_s.$$

It follows from 8.3 that  $G_s$  is solvable (respectively, nilpotent) if and only if the same is true of  $G'_{s'}$ . We are therefore reduced to the case where  $S$  is a Noetherian affine scheme.

\* In the course of this proof we shall use results established in Section 10; like Section 9, they do not depend on the present Section 8.

<sup>82</sup>Editor's note: the reference 10.8.5 has been corrected to 8.10.5.

<sup>83</sup>Editor's note: what follows has been expanded.

We first show that  $T$  is constructible. Applying the criterion of EGA 0<sub>III</sub>, 9.2.3, and arguing as above, we are reduced to showing that, when  $S$  is Noetherian and integral, either  $T$  or  $S - T$  contains a nonempty open subset of  $S$ .

Suppose, then, that  $S$  is integral and Noetherian, with generic point  $\eta$ . For every  $s \in S$ , put  $K_s^0 = G_s$ , and put, for  $p \geq 1$ ,

$$K_s^p = (K_s^{p-1}, K_s^{p-1}) \quad (\text{respectively, } K_s^p = (G_s, K_s^{p-1})). \quad \dagger$$

We first show that the sequence of closed subschemes  $K_\eta^p$  is stationary. By 7.3(v), each  $K_\eta^p$  is invariant, and hence the sequence  $(K_\eta^p)_p$  is decreasing. It is therefore stationary, since  $G_\eta$  is Noetherian. Thus there is an integer  $n$  such that

$$K_\eta^p = K_\eta^n \quad \text{for every } p \geq n.$$

On the other hand, by 10.12.1 and 10.13, there exist a nonempty open subset  $S'$  of  $S$  and an  $S'$ -group  $D$  of finite presentation such that, for every  $s \in S'$ ,

$$D_s = K_s^n, \quad (D_s, D_s) = D_s \quad (\text{respectively, } (G_s, D_s) = D_s).$$

We may suppose that  $S' = S$ . Then, for every  $s \in S$  and every  $p \geq n$ , one has  $D_s = K_s^p$ . Consequently,  $G_s$  is solvable (respectively, nilpotent) if and only if  $D_s$  is isomorphic to the unit  $\kappa(s)$ -group.

By EGA IV<sub>3</sub>, 9.6.1(xi), the set of points  $s \in S$  for which the structural morphism

$$D_s \longrightarrow \text{Spec } \kappa(s)$$

is an isomorphism is constructible.<sup>84</sup> It is therefore either nowhere dense or contains a nonempty open subset of  $S$ . We have thus proved that  $T$  is locally constructible.

We now show that, if  $G$  is also flat and separated over  $S$ , then  $T$  is closed. Since  $T$  is locally constructible, it is closed if and only if it is stable under specialization (EGA IV<sub>1</sub>, 1.10.1).

Let  $s \in S$ , and let  $s'$  be a specialization of  $s$  in  $S$ . Since we have reduced to the case where  $S$  is Noetherian, EGA II, 7.1.9 provides a discrete valuation ring  $A$  and a morphism

$$\text{Spec } A \longrightarrow S$$

such that  $s$ , respectively  $s'$ , is the image of the generic point  $\alpha$ , respectively the closed point  $a$ , of  $\text{Spec } A$ . It is enough to show that, if

$$G' = G \times_S \text{Spec } A$$

and  $G'_\alpha$  is solvable (respectively, nilpotent), then the same is true of  $G'_a$ . Since  $G$  is flat and separated over  $S$ , the  $A$ -group  $G'$  is flat and separated.<sup>‡</sup> We are thus reduced to the case where  $S$  is the spectrum of a discrete valuation ring  $A$ .

Since  $G_\alpha$  is assumed solvable (respectively, nilpotent), there is an integer  $q$  such that  $K_\alpha^q$ , with the notation introduced above, is isomorphic to the unit  $\kappa(\alpha)$ -group. For every  $n$ ,

<sup>†</sup> Translator's note: in the nilpotent branch of this display, the French source prints  $G$ . Since  $K_s^{p-1}$  is a  $\kappa(s)$ -group and the subsequent formulas use  $G_s$ , the fiber subscript is restored here.

<sup>84</sup>Editor's note: here one uses the fact that  $S$  is assumed affine, and is therefore quasi-compact and quasi-separated (cf. EGA 0<sub>III</sub>, 9.1.12).

<sup>‡</sup> Translator's note: the French source prints  $G'_\alpha$  as being flat and separated over  $A$ . The intended object is the base-changed  $A$ -group  $G'$ ; its generic fiber  $G'_\alpha$  is a group over  $\kappa(\alpha)$ .



let  $\overline{K_\alpha^n}$  denote the schematic closure of  $K_\alpha^n$  in  $G$ , in the sense of EGA IV<sub>2</sub>, 2.8.5. We show by induction on  $p$  that

$$(\overline{K_\alpha^p})_a \supset K_a^p.$$

First, since  $G$  is flat over  $A$ , one has  $\overline{G_\alpha} = G$  by EGA IV<sub>2</sub>, 2.8.5, and hence

$$(\overline{K_\alpha^0})_a = K_a^0.$$

Let  $p \geq 0$ , and suppose that

$$K_a^p \subset (\overline{K_\alpha^p})_a.$$

Following 7.2.2, denote by  $\nu_a, \nu_\alpha, \bar{\nu}, \bar{\nu}_a$  the following morphisms. In the solvable case they are

$$\begin{aligned} \nu_a &: K_a^p \times_{\kappa(a)} K_a^p \longrightarrow G_a, \\ \nu_\alpha &: K_\alpha^p \times_{\kappa(\alpha)} K_\alpha^p \longrightarrow G_\alpha, \\ \bar{\nu} &: \overline{K_\alpha^p} \times_A \overline{K_\alpha^p} \longrightarrow G, \\ \bar{\nu}_a &: (\overline{K_\alpha^p})_a \times_{\kappa(a)} (\overline{K_\alpha^p})_a \longrightarrow G_a; \end{aligned}$$

respectively, in the nilpotent case they are

$$\begin{aligned} \nu_a &: G_a \times_{\kappa(a)} K_a^p \longrightarrow G_a, \\ \nu_\alpha &: G_\alpha \times_{\kappa(\alpha)} K_\alpha^p \longrightarrow G_\alpha, \\ \bar{\nu} &: G \times_A \overline{K_\alpha^p} \longrightarrow G, \\ \bar{\nu}_a &: G_a \times_{\kappa(a)} (\overline{K_\alpha^p})_a \longrightarrow G_a. \end{aligned}$$

Since  $\nu_\alpha$  factors through  $K_\alpha^{p+1}$ ,  $\bar{\nu}$  factors through  $\overline{K_\alpha^{p+1}}$ . The latter is clearly a group subscheme of  $G$ , and therefore contains  $\Gamma_G(\bar{\nu})$ . By 7.1(iii),

$$\Gamma_G(\bar{\nu})_a = \Gamma_{G_a}(\bar{\nu}_a).$$

Moreover, the induction hypothesis gives

$$\begin{aligned} K_a^p \times_{\kappa(a)} K_a^p &\subset (\overline{K_\alpha^p})_a \times_{\kappa(a)} (\overline{K_\alpha^p})_a && \text{in the solvable case,} \\ G_a \times_{\kappa(a)} K_a^p &\subset G_a \times_{\kappa(a)} (\overline{K_\alpha^p})_a && \text{in the nilpotent case.} \end{aligned}$$

Consequently,

$$K_a^{p+1} = \Gamma_{G_a}(\nu_a) \subset \Gamma_{G_a}(\bar{\nu}_a) = \Gamma_G(\bar{\nu})_a \subset (\overline{K_\alpha^{p+1}})_a.$$

Finally,  $K_\alpha^q$  is isomorphic to the unit  $\kappa(\alpha)$ -group. The unit  $A$ -group is flat over  $A$  and is isomorphic to a closed subscheme of  $G$ , since  $G$  is separated over  $A$  (cf. 5.1). It follows from EGA IV<sub>2</sub>, 2.8.5 that  $\overline{K_\alpha^q}$  is isomorphic to the unit  $A$ -group. Since we have proved that

$$K_a^q \subset (\overline{K_\alpha^q})_a,$$

the group  $K_a^q$  is isomorphic to the unit  $\kappa(a)$ -group. Thus  $G_a$  is solvable (respectively, nilpotent).

## 9. Quotient sheaves

The present section is essentially limited to recalling, in the particular case of homogeneous spaces of groups, well-known general facts concerning passage to the quotient by flat equivalence relations (cf. Exposé IV).

**Definition 9.1.**— *Given a monomorphism  $u : G' \rightarrow G$  of  $S$ -groups, we write  $G/G'$  (respectively,  $G' \backslash G$ ) and call it the right (respectively, left) quotient sheaf of  $G$  by  $G'$ , meaning the quotient sheaf, for the (fpqc) topology, of  $G$  by the equivalence relation defined by the monomorphism*

$$\begin{aligned} G \times_S G' &\xrightarrow{\delta \circ (\text{id}_G \times u)} G \times_S G, \\ &\text{(respectively,} \\ G' \times_S G &\xrightarrow{\gamma \circ (u \times \text{id}_G)} G \times_S G), \end{aligned}$$

where  $\delta$  (respectively,  $\gamma$ ) denotes the automorphism of  $G \times_S G$  defined, for  $g, h \in G(T)$ , by

$$(g, h) \mapsto (g, gh) \quad \text{(respectively, } (h, g) \mapsto (hg, g)).$$

**Proposition 9.2.**— *Let  $u : G' \rightarrow G$  be a monomorphism of  $S$ -groups. Suppose that  $G/G'$  is representable by an  $S$ -scheme  $G''$ . Then:*

- (i) *The canonical morphism  $p : G \rightarrow G''$  is a covering for the (fpqc) topology.*
- (ii) *If  $\varepsilon'' = p \circ \varepsilon$  (this morphism is called the unit section of  $G''$ ), the following diagrams are cartesian:*

$$\begin{array}{ccc} G \times_S G' & \xrightarrow{\mu \circ (\text{id}_G \times u)} & G \\ \text{pr}_1 \downarrow & & \downarrow p \\ G & \xrightarrow{p} & G'' \end{array} \quad , \quad \begin{array}{ccc} G' & \xrightarrow{u} & G \\ \pi' \downarrow & & \downarrow p \\ S & \xrightarrow{\varepsilon''} & G'' \end{array} .$$

*In particular,  $u$  is an immersion.*

- (iii) *There exists on  $G''$  a unique structure of an  $S$ -scheme with left operator group  $G$  such that  $p$  is a morphism of  $S$ -schemes with operator group  $G$ .*
- (iv) *If, in addition,  $G'$  is assumed invariant in  $G$ , there exists on  $G''$  a unique  $S$ -group structure such that  $p$  is a morphism of  $S$ -groups.*
- (v) *Let  $S_0$  be an  $S$ -scheme, and put  $G_0 = G \times_S S_0$  and  $G'_0 = G' \times_S S_0$ . Then  $G_0/G'_0$  is representable by  $G''_0 = G'' \times_S S_0$ .<sup>85</sup>*
- (vi) *Let  $\mathcal{P}$  be a property of an  $S$ -morphism. Suppose that  $\mathcal{P}$  is stable under base change. If  $p : G \rightarrow G''$  has property  $\mathcal{P}$ , then the structural morphism  $\pi' : G' \rightarrow S$  has property  $\mathcal{P}$  as well.*
- (vii) *Let  $\mathcal{P}$  be a property of an  $S$ -morphism. Suppose that  $\mathcal{P}$  is local for the (fpqc) topology (cf. 2.0 and 2.1.2). Then  $p : G \rightarrow G''$  has property  $\mathcal{P}$  if and only if  $\pi' : G' \rightarrow S$  has property  $\mathcal{P}$ .<sup>\*</sup>*

<sup>85</sup>Editor's note: that is, the quotient is universal; cf. Exposé IV, §3.

<sup>\*</sup> Translator's note: the French source prints  $S'$  as the target of  $\pi'$  in this clause. No  $S'$  is defined here, and clause (vi) identifies  $\pi'$  as the structural morphism  $G' \rightarrow S$ ; that target is used here.

(viii) Let  $\mathcal{P}$  be a property of an  $S$ -morphism. Suppose that  $\mathcal{P}$  is local for the (fpqc) topology and stable under composition. If the structural morphisms  $G'' \rightarrow S$  and  $G' \rightarrow S$  have property  $\mathcal{P}$ , then the structural morphism  $G \rightarrow S$  has property  $\mathcal{P}$  as well.

(ix) If  $G$  is reduced, then  $G''$  is reduced.

(x) For  $G''$  to be separated over  $S$ , it is necessary and sufficient that  $u$  (or, equivalently,  $\varepsilon''$ ) be a closed immersion.

(xi) For  $G'$  to be flat over  $S$ , it is necessary and sufficient that  $p$  be flat (or, equivalently, faithfully flat).

In this case, for  $G''$  to be flat over  $S$ , it is necessary and sufficient that  $G$  be flat over  $S$ .

(xii) For  $G'$  to be flat and locally of finite presentation over  $S$ , it is necessary and sufficient that  $p : G \rightarrow G''$  be faithfully flat and locally of finite presentation.

In this case, for  $G''$  to be locally of finite presentation (respectively, locally of finite type, of finite type, smooth, étale, unramified, locally quasi-finite, quasi-finite) over  $S$ , it is enough that  $G$  have the same property over  $S$  (and the condition is also necessary in the first two cases; cf. (viii)).

(xiii) Suppose that  $G'$  is flat and of finite presentation over  $S$ .

a) Then  $p$  is of finite presentation and faithfully flat.

b) Moreover, for  $G$  to be of finite presentation over  $S$ , it is necessary and sufficient that  $G''$  be so.

*Proof.* Recall that the equivalence relation under consideration is universally effective (IV, 4.4.9). Assertions (i), (iii), (iv), (v), and the first assertion of (ii) then follow, respectively, from IV, 4.4.3, 5.2.2, 5.2.4, 3.4.5, and 3.3.2(iii). The second assertion of (ii) follows from the first, as shown by the following cartesian diagram, since

$$(G \times_S G') \times_G S$$

is isomorphic to  $G'$ :

$$\begin{array}{ccccc} G' & \xrightarrow{(\varepsilon \circ \pi'), \text{id}_{G'}} & G \times_S G' & \xrightarrow{\mu \circ (\text{id}_G \times u)} & G \\ \pi' \downarrow & & \downarrow \text{pr}_1 & & \downarrow p \\ S & \xrightarrow{\varepsilon} & G & \xrightarrow{p} & G'' \end{array}$$

Indeed,  $\varepsilon''$  is an  $S$ -section of  $G''$ , hence an immersion (EGA I, 5.3.13); by the preceding cartesian diagram the same is true of  $u$ , which completes the proof of (ii). Moreover, (vi) is an immediate consequence of the second cartesian diagram in (ii).

We prove (vii). By (i),  $p$  is a covering for the (fpqc) topology. Hence, by (ii), in order to show that  $p$  has property  $\mathcal{P}$ , it is enough to show that the first projection

$$\text{pr}_1 : G \times_S G' \longrightarrow G$$

has property  $\mathcal{P}$ . This follows from the stability of  $\mathcal{P}$  under base change, since  $\text{pr}_1$  is obtained from  $\pi'$  by base change.

It is clear that (viii) follows from (vii), since

$$\pi = \pi'' \circ p,$$

where  $\pi'' : G'' \rightarrow S$  denotes the structural morphism.

We prove (ix). By (i),  $p$  is an epimorphism. Since  $G$  is reduced,  $p$  factors through the immersion

$$G''_{\text{red}} \longrightarrow G'',$$

which is therefore also an epimorphism, and hence an isomorphism (IV, 4.4.4).

We prove (x). If  $G''$  is separated over  $S$ , then  $\varepsilon''$  is a closed immersion by EGA I, 5.4.6. On the other hand, we saw in (ii) that  $\varepsilon''$  is a closed immersion if and only if  $u$  is. Finally, if  $u$  is a closed immersion, then so is

$$\delta \circ (\text{id}_G \times u) : G \times_S G' \longrightarrow G \times_S G.$$

It follows from Lemma 9.2.1 below that  $G''$  is separated over  $S$ .

Assertion (xi) follows from (vii) and EGA IV<sub>2</sub>, 2.2.13.

Assertion (xii) follows from (vii), EGA IV<sub>4</sub>, 17.7.5 and 17.7.7, and the fact that  $p$  is universally open: for every  $s \in S$ , if the underlying space of  $G_s$  is discrete, then the same is true of the underlying space of  $G''_s$ .

Finally, assertion (xiii) follows from (vii), (viii), and EGA IV<sub>4</sub>, 17.7.5.

**Lemma 9.2.1.**— *Let  $X$  be an  $S$ -scheme, and let  $R$  be an equivalence relation on  $X$  defined by the monomorphism  $v : R \rightarrow X \times_S X$ . Suppose that  $R$  is effective. Then:*

- (i) *The morphism  $v$  is an immersion; it is a closed immersion if  $Y = X/R$  is separated.*
- (ii) *Suppose, in addition, that  $Y$  represents the (fpqc) quotient sheaf of  $X$  by  $R$ ,<sup>86</sup> and that  $v$  is a closed immersion. Then  $Y = X/R$  is separated over  $S$ .*

Recall (IV, Definition 3.3.2) that the hypothesis “ $R$  is effective” means that there is a morphism of  $S$ -schemes

$$p : X \longrightarrow Y$$

such that the natural morphism

$$R \longrightarrow X \times_Y X$$

is an isomorphism. It follows (EGA I, 5.3.5) that the following diagram is cartesian:

$$\begin{array}{ccc} R & \xrightarrow{v} & X \times_S X \\ \downarrow & & \downarrow p \times p \\ Y & \xrightarrow{\Delta_{Y/S}} & Y \times_S Y \end{array}$$

Since  $\Delta_{Y/S}$  is an immersion (EGA I, 5.3.9), the same is true of  $v$ . Likewise, if  $Y$  is separated over  $S$ , then  $\Delta_{Y/S}$  is a closed immersion, and hence so is  $v$ .

Conversely, suppose that  $v$  is a closed immersion and that  $Y$  represents the (fpqc) quotient sheaf of  $X$  by  $R$ . Then  $p$  is a covering for the (fpqc) topology (IV, 4.4.3), and therefore  $p \times p$  is also a covering: by base change,  $p \times \text{id}_X$  and  $\text{id}_Y \times p$  are coverings, and so is their composite  $p \times p$ . Thus, by (fpqc) descent (cf. EGA IV<sub>2</sub>, 2.7.1),  $\Delta_{Y/S}$  is a closed immersion; in other words,  $Y$  is separated over  $S$ .

**Remark 9.2.2.**— *Under the general hypotheses of 9.2, if  $G'$  is flat and locally of finite presentation over  $S$ , then  $p$  is a covering for the (fppf) topology,<sup>87</sup> by 9.2(vii). Consequently, assertions 9.2(vii) and 9.2(viii) extend to properties  $\mathcal{P}$  that are local for the (fppf) topology.*

<sup>86</sup>Editor's note: as O. Gabber pointed out, this condition is used in the proof and must be included among the hypotheses.

<sup>87</sup>Editor's note: (fpqc) has been corrected to (fppf).

**Remark 9.3.**— a) The question whether a quotient  $G/G'$  is representable is often delicate; in this seminar we prove the representability of certain particular quotients.

In general, in order to conclude that  $G/G'$  is representable, it is not enough to suppose that  $G$  and  $G'$  are of finite presentation over  $S$  and that  $G'$  is flat over  $S$ . Indeed, suppose in addition that  $G$  is smooth with connected fibers. In that case, if  $G/G'$  is a scheme, it is separated by Corollary 5.4; therefore  $G' \hookrightarrow G$  is a closed immersion by 9.2(x). Consequently, if  $G'$  is not closed in  $G$ , then  $G/G'$  is not representable.

To obtain such a counterexample, take  $S$  to be the spectrum of a discrete valuation ring and put  $G = \mathbb{G}_{m,S}$ . Let  $n > 1$  be an integer invertible on  $S$ . Then

$$\mu_n = \ker(G \xrightarrow{n} G)$$

is a closed subgroup of  $G$ , étale over  $S$  (cf. Exposé VII<sub>A</sub><sup>88</sup>). Let  $G'$  be the open subgroup of  $\mu_n$  obtained by removing from  $\mu_n$  the closed subset of its closed fiber that is complementary to the origin. Then  $G'$  is not closed in  $G$ , and hence  $G/G'$  is not representable. One can also construct such examples in which  $G'$  is smooth with connected fibers.

b) It remains possible, however, that  $G/G'$  is representable when  $G$  and  $G'$  are of finite presentation over  $S$ , and  $G'$  is flat over  $S$  and closed in  $G^*$ <sup>89</sup>. Under these hypotheses,  $G/G'$  is known to be representable in the following particular cases:

- 1°—  $S$  is the spectrum of an Artinian ring (cf. VI<sub>A</sub>, 3.2 and 3.3.2).
- 2°—  $G'$  is proper over  $S$ , and  $G$  is quasi-projective over  $S$  (cf. V, 7.1).
- 3°—  $S$  is locally Noetherian of dimension 1 (cf. [An73], Theorem 4.C).

## 10. Passage to the projective limit in group schemes and schemes with an operator group

**10.0.**— Recall the essential result of EGA IV<sub>3</sub>, §8.8. Suppose that the following data are given: a quasi-compact and quasi-separated scheme  $S_0$ , an upward filtered preordered set  $I$ , a direct system  $(\mathcal{A}_i)_{i \in I}$  of quasi-coherent commutative  $\mathcal{O}_{S_0}$ -algebras,

$$\mathcal{A} = \varinjlim_i \mathcal{A}_i, \quad S_i = \operatorname{Spec} \mathcal{A}_i \quad (i \in I), \quad S = \operatorname{Spec} \mathcal{A}^{90}.$$

Then the category of  $S$ -schemes of finite presentation is determined, up to equivalence, by the categories of  $S_i$ -schemes of finite presentation, the functors between these categories

$$\rho_{ji} : X_i \longmapsto X_i \times_{S_i} S_j \quad (i \leq j),$$

and the transitivity isomorphisms

$$\rho_{kj} \circ \rho_{ji} \xrightarrow{\sim} \rho_{ki}.$$

More precisely, given  $j \in I$  and an  $S_j$ -scheme  $X_j$  of finite presentation, put, for every  $i \in I$  with  $i \geq j$ ,

$$X_i = X_j \times_{S_j} S_i, \quad X = X_j \times_{S_j} S.$$

Then EGA IV<sub>3</sub>, 8.8.2 gives:

<sup>88</sup>Editor's note: see Exposé VII<sub>A</sub>, 8.4, or Exposé VIII, 2.1.

<sup>\*</sup>This is too optimistic, as M. Raynaud shows in his thesis (loc. cit., X, 14).

<sup>89</sup>Editor's note: the starred remark refers to counterexample X, 14 in [Ray70a]. The base there is a regular local scheme of dimension 2.

<sup>90</sup>Editor's note: since  $S$  is affine over  $S_0$ , it is quasi-compact and quasi-separated; cf. Editor's note 92 below.

- (i) Given  $j \in I$  and two  $S_j$ -schemes  $X_j$  and  $Y_j$  of finite presentation, the canonical map

$$\varinjlim_{i \geq j} \operatorname{Hom}_{S_i}(X_i, Y_i) \longrightarrow \operatorname{Hom}_S(X, Y)$$

is bijective.

- (ii) For every  $S$ -scheme  $X$  of finite presentation, there are an index  $j \in I$ , an  $S_j$ -scheme  $X_j$  of finite presentation, and an  $S$ -isomorphism

$$X \xrightarrow{\sim} X_j \times_{S_j} S.$$

It follows from EGA IV<sub>3</sub>, 8.8.3 that whenever a diagram  $\mathcal{D}$  involves finitely many objects and arrows in the category of  $S$ -schemes of finite presentation, one can find an index  $i \in I$  and a diagram  $\mathcal{D}_i$  in the category of  $S_i$ -schemes of finite presentation such that  $\mathcal{D}$ , up to isomorphism, is obtained from  $\mathcal{D}_i$  by the base change  $S \rightarrow S_i$ . One can even find  $i$  and  $\mathcal{D}_i$  such that every cartesian square in  $\mathcal{D}$  is obtained from a cartesian square in  $\mathcal{D}_i$ .

**10.1.**— Moreover, many familiar properties of a morphism that are stable under base change have the following property:

*Let  $u : X \rightarrow Y$  be an  $S$ -morphism between  $S$ -schemes of finite presentation. By 10.0, it is obtained by base change from an  $S_j$ -morphism  $u_j : X_j \rightarrow Y_j$  between  $S_j$ -schemes of finite presentation. Then  $u$  has the property  $\mathcal{P}$  if and only if there is an  $i \geq j$  such that*

$$u_i = u_j \times_{S_j} S_i$$

*has the property  $\mathcal{P}$ .*

This applies when  $\mathcal{P}$  is any of the following properties of a morphism: separated, surjective, radicial, affine, quasi-affine, finite, quasi-finite, proper, projective, quasi-projective, an isomorphism, a monomorphism, an immersion, an open immersion, a closed immersion (EGA IV<sub>3</sub>, 8.10.5), flat (EGA IV<sub>3</sub>, 11.2.6), smooth, unramified, or étale (EGA IV<sub>4</sub>, 17.7.8).<sup>91</sup>

This also applies when  $\mathcal{P}$  is the property of being a covering for the (fppf) topology. Indeed, given two  $S$ -schemes  $X$  and  $Y$  of finite presentation and an  $S$ -morphism  $u : X \rightarrow Y$ , it follows from Exposé IV, 6.3.1(i)<sup>92</sup> that  $u$  is a covering for the (fppf) topology if and only if there are an  $S$ -scheme  $Z$  and a faithfully flat  $S$ -morphism  $v : Z \rightarrow Y$  of finite presentation that factors through  $u$ .

The purpose of this Section 10 is to give variants of results of this kind for the category of  $S$ -groups of finite presentation, for the category of  $S$ -schemes with an operator group, and for certain properties of group monomorphisms (being invariant, being central with representable quotient sheaf, and so on).

The first two preliminary results of this kind follow. In 10.2 through 10.9, we retain the notation introduced in 10.0.

**Lemma 10.2.**— *Let  $G_j$  and  $H_j$  be two  $S_j$ -groups of finite presentation. For every  $i \geq j$ , put*

$$G_i = G_j \times_{S_j} S_i, \quad G = G_j \times_{S_j} S,$$

*and define  $H_i$  and  $H$  similarly. Then the canonical map*

$$\varinjlim_{i \geq j} \operatorname{Hom}_{S_i\text{-gr}}(G_i, H_i) \longrightarrow \operatorname{Hom}_{S\text{-gr}}(G, H)$$

<sup>91</sup>Editor's note: the word "flat" has been added, and the reference 17.7.6 has been corrected to 17.7.8.

<sup>92</sup>Editor's note: one also uses the fact that  $Y$ , being of finite presentation over  $S$ , is quasi-compact.

is bijective.

**Lemma 10.3.**— *Let  $G$  be an  $S$ -group of finite presentation. Then there are an index  $j \in I$ , an  $S_j$ -group  $G_j$  of finite presentation, and an isomorphism of  $S$ -groups*

$$G \xrightarrow{\sim} G_j \times_{S_j} S.$$

Assertions 10.2 and 10.3 are easy consequences of 10.0 and 10.1, taking into account the interpretation<sup>93</sup> of the structure of an  $S$ -group given in EGA 0<sub>III</sub>, 8.2.5 and 8.2.6.

**Lemma 10.4.**— *Let  $u : G \rightarrow H$  be a morphism of  $S$ -groups between  $S$ -groups of finite presentation. By 10.3 and 10.2,  $u$  is obtained by base change from a morphism  $u_j : G_j \rightarrow H_j$  between  $S_j$ -groups of finite presentation. Then  $u$  is a central monomorphism (respectively, an invariant monomorphism) if and only if there is an  $i \geq j$  such that*

$$u_i = u_j \times_{S_j} S_i$$

*has the same property.*

This follows immediately from 10.0 and 10.1, in view of the characterization in 6.7 of central or invariant group monomorphisms.

**Corollary 10.5.**— *Let  $G_j$  be an  $S_j$ -group of finite presentation. Then  $G_j \times_{S_j} S$  is commutative if and only if there is an  $i \geq j$  such that  $G_j \times_{S_j} S_i$  is commutative.*

Indeed, to say that an  $S$ -group is commutative is the same as saying that it is central when regarded as a group subscheme of itself.

**Proposition 10.6.**— *Let  $G_j$  be an  $S_j$ -group of finite presentation, and let  $G'_j$  be a group subscheme of  $G_j$ , flat and of finite presentation over  $S_j$ . Then the quotient*

$$(G_j \times_{S_j} S)/(G'_j \times_{S_j} S)$$

*is representable for the (fpqc) topology if and only if there is an  $i \geq j$  such that*

$$(G_j \times_{S_j} S_i)/(G'_j \times_{S_j} S_i)$$

*is representable.*

This is a consequence of the following more general lemma.

**Lemma 10.7.**— *Let  $X_j$  be an  $S_j$ -scheme of finite presentation, and let  $R_j$  be an equivalence relation on  $X_j$ , flat and of finite presentation.<sup>94</sup> Then the quotient sheaf*

$$(X_j \times_{S_j} S)/(R_j \times_{S_j} S)$$

*for the topology  $\mathcal{T} = (\text{fppf})$  or  $(\text{fpqc})$  is representable if and only if there is an  $i \geq j$  such that the quotient sheaf*

$$(X_j \times_{S_j} S_i)/(R_j \times_{S_j} S_i)$$

*for the topology  $\mathcal{T}$  is representable.*

In view of the statements EGA IV<sub>2</sub>, 8.8.2, 8.8.3, 8.10.5, and 11.2.6 recalled in 10.0, this lemma follows from the next result.

<sup>93</sup>Editor's note: in terms of commutative diagrams of  $S$ -morphisms.

<sup>94</sup>Editor's note: that is, such that the composite

$$R_j \hookrightarrow X_j \times_{S_j} X_j \xrightarrow{\text{pr}_1} X_j$$

is flat and of finite presentation.

**Lemma 10.8.**— *Let  $\mathcal{T}$  be the (fppf) or (fpqc) topology. Let  $X$  be an  $S$ -scheme of finite presentation (respectively, locally of finite presentation), and let  $R$  be an equivalence relation on  $X$  defined by a monomorphism*

$$v : R \longrightarrow X \times_S X$$

*such that  $\mathrm{pr}_1 \circ v$  is flat and of finite presentation (respectively, flat and locally of finite presentation). Then the following conditions are equivalent:*

- (i) *The quotient sheaf  $X/R$  for the topology  $\mathcal{T}$  is representable.*
- (ii) *There are an  $S$ -scheme  $Y$  of finite presentation (respectively, locally of finite presentation) and a faithfully flat morphism  $p : X \rightarrow Y$  such that the diagram*

$$(D) \quad \begin{array}{ccc} R & \xrightarrow{\mathrm{pr}_1 \circ v} & X \\ \mathrm{pr}_2 \circ v \downarrow & & \downarrow p \\ X & \xrightarrow{p} & Y \end{array}$$

*is cartesian.*

First note that, by IV, 3.3.2 and 4.4.3, the sheaf  $X/R$  for the topology  $\mathcal{T}$  is representable by  $Y$  if and only if diagram (D) is cartesian and  $p$  is a covering for the topology  $\mathcal{T}$ .

Let us prove that (i) implies (ii). Hypothesis (i) implies that diagram (D) is cartesian. Thus  $\mathrm{pr}_1 \circ v$  is obtained from  $p$  by base change along  $p$ , and  $p$  is a covering for the (fpqc) topology. Consequently, by (fpqc) descent (EGA IV<sub>2</sub>, 2.7.1), since  $\mathrm{pr}_1 \circ v$  is faithfully flat and (locally) of finite presentation, the same is true of  $p$ . Then EGA IV<sub>4</sub>, 17.7.5 shows that, since  $X$  is (locally) of finite presentation over  $S$ , so is  $Y$ .

Let us prove that (ii) implies (i). It is enough to show that  $p$  is a covering for the (fppf) topology. But  $p$  is faithfully flat by hypothesis, and it is locally of finite presentation because  $X$  and  $Y$  are locally of finite presentation over  $S$  (EGA IV<sub>1</sub>, 1.4.3(v)).

**Lemma 10.9.**— *Let  $G_j$  be an  $S_j$ -group of finite presentation, and put*

$$G = G_j \times_{S_j} S.$$

*Then  $G^0$  is representable if and only if there is an  $i \geq j$  such that*

$$(G_i)^0 = (G_j \times_{S_j} S_i)^0$$

*is representable.*

The condition is sufficient, since the functor  $G \mapsto G^0$  commutes with base change, by 3.3.

Conversely, suppose that  $G^0$  is representable. Then, by 3.9,  $G^0$  is open in  $G$  and quasi-compact over  $S$ , and hence is of finite presentation over  $S$ , since  $G$  is. By 10.3 and 10.1, there are an  $i \geq j$  and an open group subscheme  $H_i$  of  $G_i$  such that

$$H_i \times_{S_i} S = G^0.$$

The structural morphism  $G^0 \rightarrow S$  has geometrically connected fibers (VI<sub>A</sub>, 2.1.1). Consequently, by EGA IV<sub>3</sub>, 9.3.3 and 9.7.7, after increasing  $i$  we may suppose that the structural morphism  $H_i \rightarrow S_i$  has geometrically connected fibers.\* It follows from 3.10.1 that the underlying space of  $H_i$  is precisely  $(G_i)^0$ ; hence  $H_i$  represents  $(G_i)^0$ .

\* Translator's note: the French source prints  $H_i \rightarrow S$ . Since  $H_i$  is an  $S_i$ -group and the argument spreads out the connectedness of  $G^0 \rightarrow S$ , the base here is  $S_i$ .



**10.10.**— Recall two very useful special cases of the situation stated in 10.0 (cf. EGA IV<sub>3</sub>, 8.1.2(a) and (c)):

a) Given a point  $x$  of a scheme  $X$ , put  $S_0 = \operatorname{Spec} \mathbb{Z}$ , and consider the decreasing filtered inverse system  $(S_i)_{i \in I}$  of affine open neighborhoods of  $x$ . Then

$$S = \operatorname{Spec} \mathcal{O}_{X,x}.$$

In particular, if  $x$  is the generic point of an integral scheme  $X$ , then

$$S = \operatorname{Spec} \kappa(x).$$

b) Put  $S_0 = \operatorname{Spec} \mathbb{Z}$ , and consider the family  $(\mathcal{A}_i)_{i \in I}$ , preordered by inclusion, of the finitely generated  $\mathbb{Z}$ -subalgebras of the ring of an affine scheme  $S$ . Since the  $\mathcal{A}_i$  are Noetherian rings, this makes it possible in many cases to pass from the Noetherian case to the general case.

We shall now give two results concerning the special case considered in (a).<sup>95</sup>

**Proposition 10.11.**—<sup>96</sup> *Let  $S$  be an integral scheme with generic point  $\eta$ . Let  $G$  be an  $S$ -group and let  $Y, Z$  be  $S$ -schemes, all of finite presentation. Let*

$$u, v, i : Y \longrightarrow G, \quad j : Z \longrightarrow G$$

*be morphisms of  $S$ -schemes. Suppose that*

$$i_\eta : Y_\eta \longrightarrow G_\eta, \quad j_\eta : Z_\eta \longrightarrow G_\eta$$

*are closed immersions.*

*Then there is a nonempty open subset  $S'$  of  $S$  such that the morphisms*

$$i' : Y' \longrightarrow G', \quad j' : Z' \longrightarrow G',$$

*obtained by base change, are closed immersions, and such that the functors*

$$\underline{\operatorname{Transp}}(u', v'), \quad \underline{\operatorname{Transp}}_{G'}(i'(Y'), j'(Z')), \\ \underline{\operatorname{Transp}}_{G'}^{\operatorname{str}}(i'(Y'), j'(Z'))$$

*respectively, the functors*

$$\underline{\operatorname{Centr}}(u'), \quad \underline{\operatorname{Norm}}_{G'} i'(Y'),$$

*are representable by closed  $S'$ -subschemes (respectively, closed  $S'$ -subgroups) of  $G'$ , of finite presentation over  $S'$ .*

*Proof.* We shall apply the results of 10.1, first in the situation of 10.10(a), and then in that of 10.10(b). The schemes  $G_\eta, Y_\eta, Z_\eta$  are flat over the field  $\kappa(\eta)$ ; the group  $G_\eta$  is separated over  $\kappa(\eta)$  (VI<sub>A</sub>, 0.3); and  $i_\eta, j_\eta$  are closed immersions. Therefore, by 10.1, there are an affine open subset  $S' = \operatorname{Spec}(A')$  of  $S$ , a Noetherian subring  $A$  of  $A'$ ,  $A$ -schemes  $G_A, Y_A, Z_A$ , flat and of finite presentation over  $A$ , and morphisms

$$u_A, v_A, i_A : Y_A \longrightarrow G_A, \quad j_A : Z_A \longrightarrow G_A,$$

such that  $G_A$  is an  $A$ -group separated over  $A$ ,  $i_A$  and  $j_A$  are closed immersions, and

$$G \times_S S' = G_A \otimes_A A',$$

<sup>95</sup>Editor's note: observe that the proof also uses case (b).

<sup>96</sup>Editor's note: the statement has been simplified, and the case of group subschemes has been treated separately in Corollary 10.11.1.

with analogous equalities for the remaining schemes and morphisms. Since the functors under consideration over  $S'$  are obtained by base change from the analogous functors over  $\mathrm{Spec}(A)$ , it is enough to prove the result for the latter.

By EGA IV<sub>2</sub>, 6.9.2, after replacing  $A$  by a localization  $A_f$  (and hence  $S'$  by the affine open subset  $S'_f$ ), we may suppose that  $G_A, Y_A, Z_A$  are essentially free over  $A$ , in the sense of 6.2.1.<sup>97</sup> It then follows from 6.2.4(b) and 6.2.4(e) that, under the hypotheses of the proposition, the functors under consideration are representable by closed  $A$ -subschemes of  $G_A$ . These subschemes are of finite presentation over  $A$ , since  $A$  is Noetherian and  $G_A$  is of finite presentation over  $A$ ; and in the cases of  $\underline{\mathrm{Centr}}(u_A)$  and  $\underline{\mathrm{Norm}}_{G_A} i_A(Y_A)$ , they are closed  $A$ -subgroups of  $G_A$ .

**Corollary 10.11.1.**— *Let  $S$  be an integral scheme with generic point  $\eta$ . Let  $G, H, K$  be  $S$ -groups of finite presentation, and let*

$$i : H \longrightarrow G, \quad j : K \longrightarrow G$$

*be two quasi-compact monomorphisms of  $S$ -groups. Then there is a nonempty open subset  $S'$  of  $S$  such that the morphisms*

$$i' : H' \longrightarrow G', \quad j' : K' \longrightarrow G',$$

*obtained by base change, are closed immersions, and such that the functors*

$$\underline{\mathrm{Transp}}_{G'}(H', K'), \quad \underline{\mathrm{Transp}}_{G'}^{\mathrm{str}}(H', K'),$$

*respectively, the functors*

$$\underline{\mathrm{Centr}}_{G'} H', \quad \underline{\mathrm{Norm}}_{G'} H',$$

*are representable by closed  $S'$ -subschemes (respectively, closed  $S'$ -subgroups) of  $G'$ , of finite presentation over  $S'$ .*

This follows from the preceding proposition, since the hypotheses imply, by 1.4.2, that  $i_\eta$  and  $j_\eta$  are closed immersions.

**Proposition 10.12.**— *Let  $S$  be an integral scheme, let  $G$  be an  $S$ -group of finite presentation, and let  $A$  and  $B$  be two group subschemes of  $G$ , of finite presentation over  $S$  and with smooth generic fibers. Suppose moreover that one of the following conditions holds:*

- a)  *$A$  has connected generic fiber;*
- b)  *$A$  and  $B$  are invariant in  $G$ .*

*Then there are a nonempty open subset  $S'$  of  $S$  and a closed group subscheme  $D'$  of*

$$G' = G|_{S'},$$

*of finite presentation over  $S'$  and with smooth fibers, that represents the (fppf) sheaf associated with the presheaf of groups of commutators of*

$$A' = A|_{S'} \quad \text{and} \quad B' = B|_{S'}$$

*in  $G'$ . Moreover,  $D'$  has connected fibers in case (a), and is invariant in  $G'$  in case (b).*

<sup>97</sup>Editor's note: indeed, since  $G_A$  is flat and of finite presentation over  $A$ , it is covered by affine open subsets  $G_1, \dots, G_n$  such that each  $\mathcal{O}(G_i)$  is a flat  $A$ -algebra of finite presentation. By EGA IV<sub>2</sub>, 6.9.2, there is an  $f_i \in A$  such that  $\mathcal{O}(G_i)_{f_i}$  is a free  $A_{f_i}$ -module. One may therefore replace  $\mathrm{Spec}(A)$  by the affine open subset  $D(f)$ , where  $f = f_1 \cdots f_n$ , and proceed in the same way for  $Y_A$  and  $Z_A$ .

In particular, for every  $s \in S'$ ,

$$D'_s = (A_s, B_s),$$

with the notation of 7.2.2.

*Proof.* Let  $\eta$  be the generic point of  $S$ , and put

$$H_\eta = (A_\eta, B_\eta).$$

Since  $A_\eta$  and  $B_\eta$  are smooth, 7.8 in case (a) and 7.3(v) in case (b) show that  $H_\eta$  is connected, respectively invariant in  $G_\eta$ .

We are in the situation of 10.0 corresponding to 10.10(a). Consequently, by 10.3 and 10.1, there are a nonempty open subset  $S'$  of  $S$  and an  $S'$ -group subscheme  $D'$ , of finite presentation and closed in  $G'$ , such that

$$D'_\eta = D' \otimes_{S'} \kappa(\eta) = H_\eta.$$

Moreover, by EGA IV<sub>3</sub>, 9.7.7 and 9.3.3, we may suppose that  $D'$  has geometrically reduced fibers. In case (a), another application of EGA IV<sub>3</sub>, 9.7.7 and 9.3.3 allows us to suppose that  $D'$  has connected fibers, hence geometrically connected fibers (cf. VI<sub>A</sub>, 2.1.1). In case (b), we may suppose, by 10.4, that  $D'$  is invariant in  $G'$ .

We also saw, in the course of the proof of 7.8, that there is an integer  $n$  such that

$$\nu_\eta^n((A_\eta \times_{\kappa(\eta)} B_\eta)^n) = D'_\eta,$$

where  $\nu_\eta$  and  $\nu_\eta^n$  are defined as in 7.2.2(a) and 7.1(ii). The same formulas define morphisms

$$\nu' : A' \times_{S'} B' \longrightarrow G', \quad \nu'^n : (A' \times_{S'} B')^n \longrightarrow G',$$

and

$$\nu'^n \otimes_{S'} \kappa(\eta) = \nu_\eta^n.$$

It follows from 10.1 that we may choose  $S'$  so that the morphism  $\nu'^n$  is flat and factors through  $D'$ , and so that the resulting morphism

$$\nu'^n : (A' \times_{S'} B')^n \longrightarrow D'$$

is surjective. By 7.5, the morphism<sup>98</sup>

$$\nu'^{2n} = \mu \circ (\nu'^n \times_{S'} \nu'^n) : (A' \times_{S'} B')^{2n} \longrightarrow D'$$

is a covering for the (fppf) topology. Therefore, by 7.6,  $D'$  represents the (fppf) sheaf associated with the presheaf of commutators of  $A'$  and  $B'$  in  $G'$ .

Furthermore, for every  $s \in S'$ ,  $\nu'^n$  induces a surjective morphism

$$\nu_s^n : (A_s \times_{\kappa(s)} B_s)^n \longrightarrow D'_s.$$

<sup>99</sup> Thus  $D'_s$  is a closed subgroup of  $G_s$  containing

$$\nu_s(A_s \times_{\kappa(s)} B_s),$$

and hence containing  $(A_s, B_s)$ . Since  $\nu_s^n$  is surjective,  $D'_s = (A_s, B_s)$ ; and by 7.6, it represents the (fppf) sheaf of commutators of  $A_s$  and  $B_s$  in  $G_s$ .

<sup>98</sup>Editor's note:  $n + 1$  has been corrected below to  $2n$ .

<sup>99</sup>Editor's note: the argument that follows has been expanded from the original.

**Corollary 10.12.1.**—<sup>100</sup> *Let  $S$  be an integral scheme with generic point  $\eta$ , and let  $G$  be an  $S$ -group of finite presentation with smooth fibers. Put*

$$K_\eta^0 = G_\eta, \quad K_\eta^p = (K_\eta^{p-1}, K_\eta^{p-1}) \quad (\text{respectively, } K_\eta^p = (G_\eta, K_\eta^{p-1}))$$

*for every  $p \in \mathbb{N}^*$ . \*Fix  $n \in \mathbb{N}^*$ . Then there are a nonempty open subset  $S'$  of  $S$  and a group subscheme  $D$ , invariant in  $G|_{S'}$ , of finite presentation and with smooth fibers, such that*

$$D_s = K_s^n$$

*for every  $s \in S'$ .*

This follows from 10.12, by induction on  $n$ .

**Corollary 10.13.**— *Let  $S$  be an integral scheme with generic point  $\eta$ . Let  $G$  be an  $S$ -group and let  $H$  be an invariant group subscheme of  $G$ . Suppose that  $G$  and  $H$  are of finite presentation over  $S$  and have smooth generic fibers.<sup>101</sup>*

*If*

$$(H_\eta, H_\eta) = H_\eta \quad (\text{respectively, } (G_\eta, H_\eta) = H_\eta),$$

*then there is a nonempty open subset  $S'$  of  $S$  such that, for every  $s \in S'$ ,*

$$(H_s, H_s) = H_s \quad (\text{respectively, } (G_s, H_s) = H_s).$$

Indeed, the proof of 10.12 gives a nonempty open subset  $S'$  of  $S$  and an  $S'$ -group subscheme  $D$  of  $G|_{S'}$ , of finite presentation and with smooth fibers, such that, for every  $s \in S'$ ,

$$D_s = (H_s, H_s) \quad (\text{respectively, } D_s = (G_s, H_s)).$$

On the other hand, since  $D_\eta = H_\eta$  and  $D$  and  $H$  are of finite presentation over  $S'$ , EGA IV<sub>3</sub>, 8.8.2.5 gives a nonempty open subset  $S''$  of  $S'$  such that

$$D|_{S''} = H|_{S''}.$$

Consequently, for every  $s \in S''$ ,

$$H_s = D_s = (H_s, H_s) \quad (\text{respectively, } H_s = D_s = (G_s, H_s)).$$

**10.14.**— Statements 10.2 and 10.3, concerning the category of  $S$ -groups of finite presentation, extend to the category of pairs consisting of an  $S$ -group of finite presentation and an  $S$ -scheme of finite presentation with operator group  $G$ . More precisely, in the situation recalled at the beginning of 10.0:

- (i) let  $j \in I$ , and let  $G_j$  and  $G'_j$  be two  $S_j$ -groups of finite presentation; let  $H_j$  (respectively,  $H'_j$ ) be an  $S_j$ -scheme of finite presentation with operator group  $G_j$  (respectively,  $G'_j$ ). For  $i \in I$ ,  $i \geq j$ , put

$$G_i = G_j \times_{S_j} S_i, \quad G = G_j \times_{S_j} S,$$

and define  $G'_i$ ,  $G'$ ,  $H_i$ ,  $H$ ,  $H'_i$ , and  $H'$  similarly. Denote by

$$\text{Dihom}_{S\text{-gr.}}((G, H), (G', H'))$$

<sup>100</sup>Editor's note: this corollary, which is used in the proof of 8.4, has been added.

\* Translator's note: the French re-edition prints  $G$  in the second recursion. Since  $K_\eta^{p-1}$  is a  $\kappa(\eta)$ -group subscheme of  $G_\eta$ , the commutator must be formed with  $G_\eta$ .

<sup>101</sup>Editor's note: the original assumed that  $G$  and  $H$  were of finite presentation and that  $H$  had smooth fibers; the hypothesis has been modified in order to apply 10.12. The proof has also been expanded.

the set of di-morphisms of  $S$ -groups and  $S$ -schemes with operator groups from the pair  $(G, H)$  to the pair  $(G', H')$ . Then the canonical map

$$\varinjlim_{i \geq j} \text{Dihom}_{S\text{-gr.}}((G_i, H_i), (G'_i, H'_i)) \longrightarrow \text{Dihom}_{S\text{-gr.}}((G, H), (G', H'))$$

is bijective.

- (ii) let  $G$  be an  $S$ -group of finite presentation and  $H$  an  $S$ -scheme of finite presentation with operator group  $G$ . Then there are an index  $j \in I$ , an  $S_j$ -group  $G_j$  of finite presentation, an  $S_j$ -scheme  $H_j$  of finite presentation with operator group  $G_j$ , and a di-isomorphism of  $S$ -groups and  $S$ -schemes with operator groups from

$$(G_j \times_{S_j} S, H_j \times_{S_j} S)$$

onto  $(G, H)$ .\*

**Definition 10.15.**—<sup>102</sup> Let  $\mathcal{T}$  be a topology on  $(\mathbf{Sch}/S)$  coarser than the canonical topology. Given an  $S$ -group scheme  $G$  and an  $S$ -scheme  $X$  with operator group  $G$ , one says that  $X$  is a formally homogeneous space under  $G$  (relative to the topology  $\mathcal{T}$ ) if the morphism

$$\Phi : G \times_S X \longrightarrow X \times_S X, \quad (g, x) \longmapsto (gx, x),$$

defined for every  $S' \rightarrow S$ ,  $g \in G(S')$ , and  $x \in X(S')$ , is an epimorphism in the category of sheaves for the topology  $\mathcal{T}$ ; equivalently,  $\Phi$  is a covering for the topology  $\mathcal{T}$  (cf. IV, 4.4.3).

One says that  $X$  is a homogeneous space if it is formally homogeneous and, in addition, the morphism  $X \rightarrow S$  is also a covering for the topology  $\mathcal{T}$ .

In particular, one says that  $X$  is a formally principal homogeneous space under  $G$  if  $\Phi$  is an isomorphism, and that  $X$  is a principal homogeneous bundle (or a  $G$ -torsor) if  $\Phi$  is an isomorphism and, in addition, the morphism  $X \rightarrow S$  is a covering for the topology  $\mathcal{T}$  (cf. IV, 5.1.5 and 5.1.6(ii)).

**Proposition 10.16.**— Retain the situation considered at the beginning of 10.0. Let  $j \in I$ , let  $G_j$  be an  $S_j$ -group, and let  $X_j$  be an  $S_j$ -scheme with operator group  $G_j$ . Suppose that  $G_j$  and  $X_j$  are of finite presentation over  $S_j$ .

For

$$X = X_j \times_{S_j} S$$

to be a homogeneous space (respectively, a principal homogeneous bundle) under

$$G = G_j \times_{S_j} S$$

for the (fppf) topology, it is necessary and sufficient that there be an index  $i \geq j$  such that

$$X_i = X_j \times_{S_j} S_i$$

is a homogeneous space (respectively, a principal homogeneous bundle) under

$$G_i = G_j \times_{S_j} S_i.$$

Taking into account 10.14 and EGA IV<sub>3</sub>, 8.8.2, 8.8.3, and 8.10.5, the assertion follows from the property of covering morphisms for the (fppf) topology recalled in 10.1.<sup>103</sup>

\* Translator's note: the French re-edition prints  $(G, X)$  here. Since this assertion begins with the pair  $(G, H)$  and  $X$  is not defined in 10.14,  $H$  is restored.

<sup>102</sup>Editor's note: the original has been corrected by distinguishing the notions of a "formally homogeneous" object and a "homogeneous" object; see IV, §§5.1 and 6.7.

<sup>103</sup>Editor's note: for limit-passage properties of torsors expressed in terms of Čech cohomology, see also SGA 4, VII, 5.7 and its corollaries. This is developed in detail in [Ma07].

## 11. Affine group schemes

**11.0. Reminders.**—<sup>104</sup> Let  $q : X \rightarrow S$  be a quasi-compact and quasi-separated morphism of schemes (cf. EGA IV<sub>1</sub>, 1.1 and 1.2), and let  $\mathcal{F}$  be a quasi-coherent  $\mathcal{O}_X$ -module. Recall that  $q_*(\mathcal{F})$  is a quasi-coherent  $\mathcal{O}_S$ -module (EGA I, 9.2.1). Moreover, by EGA III, 1.4.15 (completed by EGA IV<sub>1</sub>, 1.7.21), one has (c) below, and the proof in *loc. cit.* also gives (a) and (b):

(a) If  $\mathcal{F}$  is a filtered direct limit of quasi-coherent submodules  $\mathcal{F}_\alpha$ , then

$$q_*(\mathcal{F}) = \varinjlim_{\alpha} q_*(\mathcal{F}_\alpha).$$

(b) If  $\mathcal{E}$  is a flat  $\mathcal{O}_S$ -module, the canonical morphism

$$\mathcal{E} \otimes_{\mathcal{O}_S} q_*(\mathcal{O}_X) \longrightarrow q_*q^*(\mathcal{E})$$

is an isomorphism.

(c) Let  $p : S' \rightarrow S$  be a flat morphism, let  $q' : X' \rightarrow S'$  be obtained from  $q$  by base change, and let  $\mathcal{F}'$  be the inverse image of  $\mathcal{F}$  on  $X'$ . Then the canonical morphism

$$p^*q_*(\mathcal{F}) \longrightarrow q'_*(\mathcal{F}')$$

is an isomorphism.

Indeed, let  $U = \operatorname{Spec}(A)$  be an arbitrary affine open subset of  $S$ . By hypothesis,  $q^{-1}(U)$  is the union of affine open subsets

$$V_i = \operatorname{Spec}(B_i), \quad i = 1, \dots, n,$$

and every intersection  $V_i \cap V_j$  is the union of finitely many affine open subsets

$$W_{ijk} = \operatorname{Spec}(C_{ijk}).$$

Then  $\Gamma(U, q_*(\mathcal{F})) = \Gamma(q^{-1}(U), \mathcal{F})$  is the kernel of the morphism

$$\bigoplus_{i=1}^n \Gamma(V_i, \mathcal{F}) \longrightarrow \bigoplus_{i,j,k} \Gamma(W_{ijk}, \mathcal{F}).$$

Assertion (a) follows, since every term above commutes with filtered direct limits (the  $V_i$  and  $W_{ijk}$  being affine, hence quasi-compact).

Let us prove (b). The module  $E = \Gamma(U, \mathcal{E})$  is flat over  $A$ , and  $\Gamma(U, q_*q^*(\mathcal{E}))$  is the kernel  $K(E)$  of the morphism

$$\bigoplus_{i=1}^n B_i \otimes_A E \longrightarrow \bigoplus_{i,j,k} C_{ijk} \otimes_A E.$$

Since  $E$  is flat over  $A$ , this kernel is identified with

$$K(A) \otimes_A E = \mathcal{O}_X(q^{-1}(U)) \otimes_A E.$$

<sup>104</sup>Editor's note: this paragraph of reminders has been added.

Finally, if  $U'$  is an arbitrary affine open subset of  $S'$  lying over  $U$ , then  $A' = \mathcal{O}(U')$  is a flat  $A$ -algebra, and the same argument gives\*

$$\mathcal{F}(q^{-1}(U)) \otimes_A A' \xrightarrow{\sim} \mathcal{F}'(q'^{-1}(U')).$$

**Notation.**— Let  $S$  be a scheme, let  $X$  be an  $S$ -scheme, and let  $f : X \rightarrow S$  be the structural morphism. We shall put

$$\mathcal{A}(X) = f_*(\mathcal{O}_X).$$

**Lemma 11.1.**— *Let  $X$  and  $Y$  be two quasi-compact and quasi-separated  $S$ -schemes, and let  $f : X \rightarrow S$  and  $g : Y \rightarrow S$  be their structural morphisms. Then the canonical homomorphism*

$$\varphi : \mathcal{A}(X) \otimes_{\mathcal{O}_S} \mathcal{A}(Y) \longrightarrow \mathcal{A}(X \times_S Y)$$

*is an isomorphism in each of the following cases:*<sup>105</sup>

- (a)  $f$  and  $g$  are affine;
- (b)  $g$  (or  $f$ ) is flat and affine;
- (c)  $g$  is flat and  $f_*(\mathcal{O}_X)$  is a flat  $\mathcal{O}_S$ -module.

In case (b), we shall suppose that it is  $g$  that is flat and affine. Put

$$S' = \operatorname{Spec} \mathcal{A}(X), \quad Y' = Y \times_S S', \quad g' = g \times_S S',$$

and write  $v : S' \rightarrow S$  for the structural morphism:

$$\begin{array}{ccc} Y' & \xrightarrow{\quad} & Y \\ g' \downarrow & & \downarrow g \\ \operatorname{Spec} \mathcal{A}(X) = S' & \xrightarrow{\quad v \quad} & S \end{array}$$

In cases (a) and (b), the morphism  $g$  is affine, and hence EGA II, 1.5.2 gives

$$g'_*(\mathcal{O}_{Y'}) = v^* g_*(\mathcal{O}_Y) = \mathcal{A}(Y) \otimes_{\mathcal{O}_S} \mathcal{O}_{S'}. \quad (1)$$

The same equality holds in case (c), by 11.0(c), since  $S'$  is flat over  $S$  and  $g$  is quasi-compact and quasi-separated.

On the other hand, by EGA II, 1.2.7, the morphism  $f : X \rightarrow S$  factors through  $v$  by means of a morphism  $p : X \rightarrow S'$ , and

$$p_*(\mathcal{O}_X) = \mathcal{O}_{S'}, \quad X \times_S Y = X \times_{S'} Y'.$$

Since  $f$  is quasi-separated, so is  $p$  (EGA IV<sub>1</sub>, 1.2.2); and since  $f$  is quasi-compact and  $v$  is quasi-separated,  $p$  is also quasi-compact (EGA IV<sub>1</sub>, 1.2.4). Consider the cartesian square

---

\* Translator's note: the French re-edition prints  $q'^{-1}(U)$  in the right-hand side below. Since  $q' : X' \rightarrow S'$  and the affine open subset just introduced is  $U' \subset S'$ , the inverse image is  $q'^{-1}(U')$ .

<sup>105</sup>Editor's note: if  $k$  is a field and  $X$  is an infinite disjoint union of copies of  $S = \operatorname{Spec} k$  (so that  $X$  is not quasi-compact), then  $\mathcal{A}(X) = k^X$ , and the canonical morphism

$$k^X \otimes_k k^X \longrightarrow k^{X \times X}$$

is not surjective.

$$\begin{array}{ccc}
X \times_S Y & \xrightarrow{p'} & Y' \\
\downarrow & & \downarrow g' \\
X & \xrightarrow{p} & S'
\end{array}$$

In cases (b) and (c), the scheme  $Y$  is flat over  $S$ , and hence  $Y'$  is flat over  $S'$ . Applying 11.0(c) once again, we obtain

$$p'_*(\mathcal{O}_{X \times_S Y}) = g'^*p_*(\mathcal{O}_X) = g'^*(\mathcal{O}_{S'}) = \mathcal{O}_{Y'}. \quad (2)$$

The same equality holds in case (a), because  $p$  and  $p'$  are then isomorphisms.

Finally, since  $v$  is affine, EGA II, 1.4.7 gives

$$v_*(\mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{O}_{S'}) = \mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{A}(X)$$

for every quasi-coherent  $\mathcal{O}_S$ -module  $\mathcal{F}$ . Combined with (2) and (1), this gives

$$\begin{aligned}
\mathcal{A}(X \times_S Y) &= v_*g'_*p'_*(\mathcal{O}_{X \times_S Y}) \\
&\stackrel{(2)}{=} v_*g'_*(\mathcal{O}_{Y'}) \stackrel{(1)}{=} v_*(\mathcal{A}(Y) \otimes_{\mathcal{O}_S} \mathcal{O}_{S'}) \\
&= \mathcal{A}(Y) \otimes_{\mathcal{O}_S} \mathcal{A}(X).
\end{aligned}$$

**Corollary 11.2.**— *The functor*

$$X \longmapsto \operatorname{Spec} \mathcal{A}(X),$$

*from the full subcategory of  $(\mathbf{Sch}/S)$  formed by the  $S$ -schemes  $X$  that are flat, quasi-compact, and quasi-separated over  $S$ , and for which  $\mathcal{A}(X)$  is a flat  $\mathcal{O}_S$ -module, to the category of flat affine  $S$ -schemes, commutes with finite products and therefore takes  $S$ -groups to  $S$ -groups.*

**Definition 11.3.**— *Given a flat  $S$ -group  $G$ , quasi-compact and quasi-separated over  $S$ , such that  $\mathcal{A}(G)$  is flat over  $\mathcal{O}_S$ ,<sup>106</sup> we write  $G_{\text{af}}$ , and call the  $S$ -group*

$$G_{\text{af}} = \operatorname{Spec} \mathcal{A}(G)$$

*the affine envelope of  $G$ .*

**Proposition 11.3.1.**—<sup>107</sup> *The canonical morphism*

$$\tau_G : G \longrightarrow G_{\text{af}}$$

*is a morphism of  $S$ -groups. Moreover, it has the following universal property:*

- (i) *For every morphism of  $S$ -schemes  $\varphi : G \rightarrow H$ , where  $H$  is affine over  $S$ , there is a unique morphism of  $S$ -schemes  $\varphi' : G_{\text{af}} \rightarrow H$  such that*

$$\varphi = \varphi' \circ \tau_G.$$

- (ii) *If, moreover,  $H$  is an  $S$ -group and  $\varphi$  is a morphism of  $S$ -groups, then the same is true of  $\varphi'$ .*

<sup>106</sup>Editor's note: if  $S$  is a regular locally Noetherian scheme of dimension at most 2, and  $X$  is an  $S$ -scheme that is flat, quasi-compact, and quasi-separated, then  $\mathcal{A}(X)$  is a flat  $\mathcal{O}_S$ -module; cf. [Ray70a], VII, 3.2.

<sup>107</sup>Editor's note: this proposition has been added; see also the additional Section 12 below for a study of the morphism  $G \rightarrow G_{\text{af}}$  and of its kernel.



**11.4.**—<sup>108</sup> Let  $\mathcal{E}$  and  $\mathcal{F}$  be two quasi-coherent  $\mathcal{O}_S$ -modules. Consider the  $S$ -functor

$$\underline{\mathrm{Hom}}_{\mathcal{O}_S}(\mathrm{W}(\mathcal{E}), \mathrm{W}(\mathcal{F}))$$

(cf. I, 3.1.4), that is, for every  $S$ -scheme  $f : X \rightarrow S$ ,

$$\underline{\mathrm{Hom}}_{\mathcal{O}_S}(\mathrm{W}(\mathcal{E}), \mathrm{W}(\mathcal{F}))(X) = \mathrm{Hom}_{\mathcal{O}_X}(\mathrm{W}(\mathcal{E})_X, \mathrm{W}(\mathcal{F})_X).$$

Moreover, by I, 4.6.2, one has  $\mathrm{W}(\mathcal{E})_X = \mathrm{W}(f^*(\mathcal{E}))$  (and similarly for  $\mathcal{F}$ ), and

$$\mathrm{Hom}_{\mathcal{O}_X}(\mathrm{W}(f^*(\mathcal{E})), \mathrm{W}(f^*(\mathcal{F}))) = \mathrm{Hom}_{\mathcal{O}_X}(f^*(\mathcal{E}), f^*(\mathcal{F})).$$

We therefore obtain (using the adjunction formula for the last equality)

$$\underline{\mathrm{Hom}}_{\mathcal{O}_S}(\mathrm{W}(\mathcal{E}), \mathrm{W}(\mathcal{F}))(X) = \mathrm{Hom}_{\mathcal{O}_X}(f^*(\mathcal{E}), f^*(\mathcal{F})) = \mathrm{Hom}_{\mathcal{O}_S}(\mathcal{E}, f_*f^*(\mathcal{F})). \quad (\dagger)$$

**Proposition 11.5.**— *Let  $X$  be an  $S$ -scheme that is quasi-compact and quasi-separated over  $S$ , let  $f : X \rightarrow S$  be the structural morphism, and let  $\mathcal{E}$  and  $\mathcal{E}'$  be two quasi-coherent  $\mathcal{O}_S$ -modules. Suppose that one of the following two conditions holds:*

- a)  $f$  is affine;
- b)  $\mathcal{E}$  is flat over  $\mathcal{O}_S$ .

*Then the canonical morphism*

$$\mathcal{E} \otimes_{\mathcal{O}_S} \mathcal{A}(X) \longrightarrow f_*f^*(\mathcal{E})$$

*is an isomorphism, and hence*

$$\underline{\mathrm{Hom}}_{\mathcal{O}_S}(\mathrm{W}(\mathcal{E}'), \mathrm{W}(\mathcal{E}))(X) = \mathrm{Hom}_{\mathcal{O}_S}(\mathcal{E}', \mathcal{E} \otimes_{\mathcal{O}_S} \mathcal{A}(X)).$$

Indeed, the second assertion follows from 11.4 and the first; the latter follows from EGA II, 1.4.7 in case (a), and from 11.0(b) in case (b).<sup>109</sup>

**11.6.**—<sup>110</sup> Let  $G$  be an  $S$ -group and let  $\mathcal{E}$  be a quasi-coherent  $\mathcal{O}_S$ -module. Giving  $\mathcal{E}$  a  $G$ - $\mathcal{O}_S$ -module structure (that is, an  $\mathcal{O}_S$ -linear action of  $G$  on  $\mathrm{W}(\mathcal{E})$ , cf. I, 4.7.1) is equivalent to giving a morphism of  $S$ -functors in monoids

$$\rho : G \longrightarrow \underline{\mathrm{End}}_{\mathcal{O}_S}(\mathrm{W}(\mathcal{E}));$$

indeed, such a  $\rho$  necessarily takes  $G$  into  $\underline{\mathrm{Aut}}_{\mathcal{O}_S}(\mathrm{W}(\mathcal{E}))$ .

By 11.4, giving a morphism of  $S$ -functors

$$\rho : G \longrightarrow \underline{\mathrm{End}}_{\mathcal{O}_S}(\mathrm{W}(\mathcal{E}))$$

<sup>108</sup>Editor's note: in 11.4–11.6, we have considered  $\underline{\mathrm{Hom}}_{\mathcal{O}_S}(\mathrm{W}(\mathcal{E}), \mathrm{W}(\mathcal{F}))$  instead of  $\underline{\mathrm{Hom}}_{\mathcal{O}_S}(\mathrm{V}(\mathcal{F}), \mathrm{V}(\mathcal{E}))$  (cf. I, 4.6.3), and simplified the original by taking account of the addition 11.0(b).

<sup>109</sup>Editor's note: we have simplified the original, which used the isomorphism

$$\underline{\mathrm{Hom}}_{\mathcal{O}_S}(\mathrm{W}(\mathcal{E}'), \mathrm{W}(\mathcal{E})) \simeq \underline{\mathrm{Hom}}_{\mathcal{O}_S}(\mathrm{V}(\mathcal{E}), \mathrm{V}(\mathcal{E}')),$$

then the inclusion of the right-hand term in

$$\underline{\mathrm{Hom}}_S(\mathrm{V}(\mathcal{E}), \mathrm{V}(\mathcal{E}')) = \mathrm{Hom}_{\mathcal{O}_S}(\mathcal{E}', f_*f^*(\mathrm{Sym}(\mathcal{E}))),$$

and applied EGA III, 4.1.15 to  $\mathrm{V}(\mathcal{E}) = \mathrm{Spec}(\mathrm{Sym}(\mathcal{E}))$  to deduce 11.0(b).

<sup>110</sup>Editor's note: we have expanded 11.6 and displayed the results obtained in the form of Proposition 11.6.1.

is equivalent to giving an element  $\theta$  of

$$\mathrm{Hom}_{\mathcal{O}_G}(f^*(\mathcal{E}), f^*(\mathcal{E})),$$

which corresponds by adjunction to an element  $\delta$  of

$$\mathrm{Hom}_{\mathcal{O}_S}(\mathcal{E}, f_* f^*(\mathcal{E})),$$

where  $f$  denotes the projection  $G \rightarrow S$ .

Let  $m : G \times_S G \rightarrow G$  be multiplication, let  $\delta_G$  be the morphism

$$\mathcal{O}_G \longrightarrow m_*(\mathcal{O}_{G \times_S G}),$$

and let  $\varphi : G \times_S G \rightarrow S$  be the projection (which equals  $f \circ m$ ). It is convenient to write  $\boxtimes$  for the “external” tensor product. We thus obtain a morphism

$$\mathrm{id}_{\mathcal{E}} \boxtimes \delta_G : f^*(\mathcal{E}) = \mathcal{E} \boxtimes_{\mathcal{O}_S} \mathcal{O}_G \longrightarrow \mathcal{E} \boxtimes_{\mathcal{O}_S} m_*(\mathcal{O}_{G \times_S G}).$$

By abuse of notation, we again write  $\mathrm{id}_{\mathcal{E}} \boxtimes \delta_G$  for the composite of this morphism with the canonical morphism

$$\mathcal{E} \boxtimes_{\mathcal{O}_S} m_*(\mathcal{O}_{G \times_S G}) \longrightarrow m_* m^* f^*(\mathcal{E}) = m_* \varphi^*(\mathcal{E}).$$

On the other hand, let  $h : G \rightarrow S$  denote a second copy of  $f : G \rightarrow S$ , and consider the following commutative diagram, in which  $p, q$  denote the two projections:

$$\begin{array}{ccc} G \times_S G & \xrightarrow{q} & G \\ p \downarrow & \searrow \varphi & \downarrow h \\ G & \xrightarrow{f} & S \end{array}$$

Writing  $\delta$  again for the morphism  $\mathcal{E} \rightarrow h_* h^*(\mathcal{E})$ , we obtain the morphism

$$\delta \boxtimes \mathrm{id}_{\mathcal{O}_G} : f^*(\mathcal{E}) = \mathcal{E} \boxtimes_{\mathcal{O}_S} \mathcal{O}_G \longrightarrow h_* h^*(\mathcal{E}) \boxtimes_{\mathcal{O}_S} \mathcal{O}_G = f^* h_* h^*(\mathcal{E}).$$

By abuse of notation, we again write  $\delta \boxtimes \mathrm{id}_{\mathcal{O}_G}$  for the composite of this morphism with the canonical morphism

$$f^* h_* h^*(\mathcal{E}) \longrightarrow p_* \varphi^*(\mathcal{E}).$$

The condition that  $\rho$  be compatible with multiplication is then equivalent to saying that, for every open subset  $U$  of  $S$ , the following diagram is commutative:

$$(1) \quad \begin{array}{ccc} \Gamma(U, \mathcal{E}) & \xrightarrow{\delta} & \Gamma(U \times_S G, \mathcal{E} \boxtimes_{\mathcal{O}_S} \mathcal{O}_G) \\ \delta \downarrow & & \downarrow \mathrm{id}_{\mathcal{E}} \boxtimes \delta_G \\ \Gamma(U \times_S G, \mathcal{E} \boxtimes_{\mathcal{O}_S} \mathcal{O}_G) & \xrightarrow{\delta \boxtimes \mathrm{id}_{\mathcal{O}_G}} & \Gamma(U \times_S G \times_S G, \mathcal{E} \boxtimes_{\mathcal{O}_S} \mathcal{O}_G \boxtimes_{\mathcal{O}_S} \mathcal{O}_G) \end{array}$$

Furthermore, the unit section  $\varepsilon : S \rightarrow G$  induces a morphism  $u$  from  $\mathcal{O}_G$  to

$$\varepsilon_* \varepsilon^*(\mathcal{O}_G) = \varepsilon_*(\mathcal{O}_S),$$

and the condition that  $\rho$  preserve unit elements is equivalent to the commutativity of the diagram

$$(2) \quad \begin{array}{ccc} \Gamma(U, \mathcal{E}) & \xrightarrow{\delta} & \Gamma(U \times_S G, \mathcal{E} \boxtimes_{\mathcal{O}_S} \mathcal{O}_G) \\ & \searrow \sim & \swarrow \text{id}_{\mathcal{E}} \boxtimes u \\ & \Gamma(U \times_S S, \mathcal{E} \boxtimes_{\mathcal{O}_S} \mathcal{O}_S) & \end{array}$$

Thus, giving  $\mathcal{E}$  a  $G$ - $\mathcal{O}_S$ -module structure is equivalent to giving a morphism of  $\mathcal{O}_S$ -modules

$$\delta : \mathcal{E} \longrightarrow f_* f^*(\mathcal{E})$$

satisfying conditions (1) and (2) above. In that case, the morphism

$$\theta : f^*(\mathcal{E}) \longrightarrow f^*(\mathcal{E})$$

deduced from  $\delta$  by adjunction is an isomorphism, because it corresponds to the isomorphism

$$G \times_S W(\mathcal{E}) \xrightarrow{\sim} G \times_S W(\mathcal{E})$$

defined set-theoretically by  $(g, x) \mapsto (g, gx)$ ; see also I, 6.5.4.

Now suppose that  $G$  is flat, quasi-compact, and quasi-separated over  $S$ , and that  $\mathcal{A}(G)$  is a flat  $\mathcal{O}_S$ -module. Then, by 11.1(c), the canonical morphism

$$\mathcal{A}(G) \otimes_{\mathcal{O}_S} \mathcal{A}(G) \longrightarrow \mathcal{A}(G \times_S G)$$

is an isomorphism, and we write  $\Delta$  for the morphism

$$\delta_G : \mathcal{A}(G) \longrightarrow \mathcal{A}(G) \otimes_{\mathcal{O}_S} \mathcal{A}(G).$$

If, moreover,

$$\mathcal{E} \otimes_{\mathcal{O}_S} \mathcal{A}(G) = f_* f^*(\mathcal{E}),$$

as is the case by 11.5 if  $G \rightarrow S$  is affine or if  $\mathcal{E}$  is flat over  $\mathcal{O}_S$ , then conditions (1) and (2) are equivalent to the following conditions, which express that

$$\delta : \mathcal{E} \longrightarrow \mathcal{E} \otimes_{\mathcal{O}_S} \mathcal{A}(G)$$

makes  $\mathcal{E}$  a right  $\mathcal{A}(G)$ -comodule (cf. I, 4.7.2):

(CM 1) Put  $\mathcal{A} = \mathcal{A}(G)$ . The following diagram is commutative:

$$\begin{array}{ccc} \mathcal{E} & \xrightarrow{\delta} & \mathcal{E} \otimes \mathcal{A} \\ \delta \downarrow & & \downarrow \text{id}_{\mathcal{E}} \otimes \Delta \\ \mathcal{E} \otimes \mathcal{A} & \xrightarrow{\delta \otimes \text{id}_{\mathcal{A}}} & \mathcal{E} \otimes \mathcal{A} \otimes \mathcal{A} \end{array}$$

(CM 2) Write  $\eta : \mathcal{A} \rightarrow \mathcal{O}_S$  for the morphism  $\mathcal{A}(\varepsilon)$ . The following diagram is commutative:

$$\begin{array}{ccc} \mathcal{E} & \xrightarrow{\delta} & \mathcal{E} \otimes \mathcal{A} \\ & \searrow \sim & \swarrow \text{id}_{\mathcal{E}} \otimes \eta \\ & \mathcal{E} \otimes \mathcal{O}_S & \end{array}$$

**Remark 11.6.A.**— Recall that  $V(\mathcal{E})$  denotes the vector fibration over  $S$  representing the functor  $V(\mathcal{E})$ ; that is, for every  $S' \rightarrow S$ ,

$$V(\mathcal{E})(S') = \operatorname{Hom}_{\mathcal{O}_{S'}}(\mathcal{E}_{S'}, \mathcal{O}_{S'}).$$

Since I, 4.6.2 gives an anti-isomorphism of  $S$ -functors in monoids

$$\underline{\operatorname{End}}_{\mathcal{O}_S}(W(\mathcal{E})) \simeq \underline{\operatorname{End}}_{\mathcal{O}_S}(V(\mathcal{E})),$$

if  $\mathcal{E}$  is a left  $G$ - $\mathcal{O}_S$ -module, then there is a right action

$$\mu : V(\mathcal{E}) \times_S G \longrightarrow V(\mathcal{E})$$

of  $G$  on  $V(\mathcal{E})$ , defined set-theoretically by

$$(\phi g)(x) = \phi(gx)$$

for every  $g \in G(S')$ ,  $x \in \Gamma(S', \mathcal{E}_{S'})$ , and  $\phi \in \operatorname{Hom}_{\mathcal{O}_{S'}}(\mathcal{E}_{S'}, \mathcal{O}_{S'})$ . We therefore have the commutative diagrams

$$\begin{array}{ccc} V(\mathcal{E}) \times_S G \times_S G & \xrightarrow{\mu \times \operatorname{id}_G} & V(\mathcal{E}) \times_S G \\ \operatorname{id}_{V(\mathcal{E})} \times m \downarrow & & \downarrow \mu \\ V(\mathcal{E}) \times_S G & \xrightarrow{\mu} & V(\mathcal{E}) \end{array} \quad \begin{array}{ccc} V(\mathcal{E}) & \xleftarrow{\mu} & V(\mathcal{E}) \times_S G \\ & \nwarrow \tilde{\operatorname{id}}_{V(\mathcal{E})} \times \varepsilon & \uparrow \\ & & V(\mathcal{E}) \times_S S \end{array}$$

When  $G$  is flat, quasi-compact, and quasi-separated over  $S$ , when  $\mathcal{A}(G)$  is a flat  $\mathcal{O}_S$ -module, and when one of the conditions of 11.5 is satisfied, one similarly recovers conditions (CM 1) and (CM 2).

Consequently, we have proved:

**Proposition 11.6.1.**— *Let  $G$  be an  $S$ -group that is flat, quasi-compact, and quasi-separated over  $S$ , such that  $\mathcal{A}(G)$  is a flat  $\mathcal{O}_S$ -module, and let  $\mathcal{E}$  be a quasi-coherent  $\mathcal{O}_S$ -module.*

(i) *Giving an  $\mathcal{A}(G)$ -comodule structure*

$$\delta : \mathcal{E} \longrightarrow \mathcal{E} \otimes_{\mathcal{O}_S} \mathcal{A}(G)$$

*is equivalent to giving  $\mathcal{E}$  a  $G_{\text{af}}$ - $\mathcal{O}_S$ -module structure (that is, an  $\mathcal{O}_S$ -linear action of  $G_{\text{af}}$  on  $\mathcal{E}$ ). By composition with the morphism of  $S$ -groups  $G \rightarrow G_{\text{af}}$ , this defines a  $G$ - $\mathcal{O}_S$ -module structure on  $\mathcal{E}$ .*

(ii) *If, moreover,  $\mathcal{E}$  is flat, every  $\mathcal{O}_S$ -linear action of  $G$  on  $\mathcal{E}$  factors through  $G_{\text{af}}$  and corresponds to a unique  $\mathcal{A}(G)$ -comodule structure on  $\mathcal{E}$ .*

**Lemma 11.7.**— *Let  $G$  be an  $S$ -group that is flat, quasi-compact, and quasi-separated over  $S$ , such that  $\mathcal{A} = \mathcal{A}(G)$  is a flat  $\mathcal{O}_S$ -module. Let  $\mathcal{E}$  be a quasi-coherent  $\mathcal{O}_S$ -module, let*

$$\delta : \mathcal{E} \longrightarrow \mathcal{E} \otimes \mathcal{A}$$

*be an  $\mathcal{A}$ -comodule structure, and let*

$$\rho : G \longrightarrow \underline{\operatorname{Aut}}_{\mathcal{O}_S}(W(\mathcal{E}))$$

*be the associated action of  $G$  on  $\mathcal{E}$ . Let  $\mathcal{E}_0$  be a quasi-coherent  $\mathcal{O}_S$ -submodule of  $\mathcal{E}$  such that the restriction  $\delta_0$  of  $\delta$  to  $\mathcal{E}_0^*$  factors through*

$$\mathcal{E}_0 \longrightarrow \mathcal{E}_0 \otimes \mathcal{A};$$

*that is, suppose that the following diagram is commutative:*

---

\* Translator's note: the French source writes “the restriction  $\delta_0$  of  $\mu$ ” here. The surrounding notation and the displayed square show that  $\delta$  is intended.

$$\begin{array}{ccc}
\mathcal{E}_0 & \hookrightarrow & \mathcal{E} \\
\delta_0 \downarrow & & \downarrow \delta \\
\mathcal{E}_0 \otimes \mathcal{A} & \hookrightarrow & \mathcal{E} \otimes \mathcal{A}
\end{array}$$

(N.B. The morphism  $\mathcal{E}_0 \otimes \mathcal{A} \rightarrow \mathcal{E} \otimes \mathcal{A}$  is injective because  $\mathcal{A}$  is flat over  $\mathcal{O}_S$ .) Then  $\delta_0$  makes  $\mathcal{E}_0$  an  $\mathcal{A}(G)$ -comodule, and hence defines an action  $\rho_0$  of  $G$  on  $\mathcal{E}_0$ . This is called the action on  $\mathcal{E}_0$  induced by  $\rho$ , and we say that  $\mathcal{E}_0$  is stable under  $\rho$ .

This follows immediately from the definitions and from 11.6. Notice, however, that in general the canonical map

$$W(\mathcal{E}_0) \longrightarrow W(\mathcal{E})$$

is not a monomorphism.

**Remark 11.7.bis.**—<sup>111</sup> Let  $G$  be a flat  $S$ -group and let  $\mathcal{E}$  be a quasi-coherent  $G$ - $\mathcal{O}_S$ -module. Write  $f$  for the morphism  $G \rightarrow S$  and  $\delta$  for the morphism

$$\mathcal{E} \longrightarrow f_* f^*(\mathcal{E})$$

defined in 11.6. Let  $\mathcal{E}_0$  be a quasi-coherent  $\mathcal{O}_S$ -submodule of  $\mathcal{E}$ . Since  $f$  is flat,  $f_* f^*(\mathcal{E}_0)$  is an  $\mathcal{O}_S$ -submodule of  $f_* f^*(\mathcal{E})$ , and the same holds for  $\varphi = f \times f$ . Consequently, if the restriction  $\delta_0$  of  $\delta$  to  $\mathcal{E}_0$  factors through  $f_* f^*(\mathcal{E}_0)$ , then it makes  $\mathcal{E}_0$  a  $G$ - $\mathcal{O}_S$ -module. In this case, we say that  $\mathcal{E}_0$  is a  $G$ -stable submodule of  $\mathcal{E}$ .

**Definition 11.8.0.**—<sup>112</sup> Let  $S$  be a scheme. An  $\mathcal{O}_S$ -coalgebra is an  $\mathcal{O}_S$ -module  $\mathcal{C}$ , together with two morphisms of  $\mathcal{O}_S$ -modules

$$\Delta : \mathcal{C} \longrightarrow \mathcal{C} \otimes \mathcal{C}, \quad \varepsilon : \mathcal{C} \longrightarrow \mathcal{O}_S,$$

satisfying the following two axioms (cf. I, 4.2):

(CO 1)  $\Delta$  is coassociative: the following diagram is commutative:

$$\begin{array}{ccccc}
& & \mathcal{C} \otimes \mathcal{C} & & \\
& \nearrow \Delta & & \searrow \text{id} \otimes \Delta & \\
\mathcal{C} & & & & \mathcal{C} \otimes \mathcal{C} \otimes \mathcal{C} \\
& \searrow \Delta & & \nearrow \Delta \otimes \text{id} & \\
& & \mathcal{C} \otimes \mathcal{C} & & 
\end{array}$$

(CO 2)  $\varepsilon$  is a counit; that is, the following two composites are the identity:

$$\mathcal{C} \xrightarrow{\Delta} \mathcal{C} \otimes \mathcal{C} \xrightarrow{\text{id} \otimes \varepsilon} \mathcal{C} \otimes \mathcal{O}_S \xrightarrow{\sim} \mathcal{C},$$

$$\mathcal{C} \xrightarrow{\Delta} \mathcal{C} \otimes \mathcal{C} \xrightarrow{\varepsilon \otimes \text{id}} \mathcal{O}_S \otimes \mathcal{C} \xrightarrow{\sim} \mathcal{C}.$$

<sup>111</sup>Editor's note: we have added this remark, which generalizes 11.7 and will be useful in 11.10.bis.

<sup>112</sup>Editor's note: the statements 11.8 and 11.9 concern only the notion of a comodule over a coalgebra. We have therefore introduced Definition 11.8.0 and reformulated 11.8 and 11.9 accordingly.

A  $C$ -comodule (on the right) is an  $\mathcal{O}_S$ -module  $\mathcal{E}$ , together with a morphism of  $\mathcal{O}_S$ -modules

$$\mu : \mathcal{E} \longrightarrow \mathcal{E} \otimes C$$

satisfying axioms (CM 1) and (CM 2) of 11.6.

We say that  $C$  (respectively,  $\mathcal{E}$ ) is a *quasi-coherent coalgebra* (respectively, a *quasi-coherent comodule*) if it is a quasi-coherent  $\mathcal{O}_S$ -module.

Let  $A$  be a commutative ring and let  $C$  be an  $A$ -coalgebra. Then

$$C^\vee = \operatorname{Hom}_A(C, A)$$

is an  $A$ -algebra. We denote by

$$\operatorname{ev} : C \otimes_A C^\vee \longrightarrow A$$

the natural evaluation map.

**Lemma 11.8.**— *Let  $C$  be an  $A$ -coalgebra,  $V$  a  $C$ -comodule, and  $M$  an  $A$ -submodule of  $V$ . Suppose that  $C$  is a projective  $A$ -module.<sup>113</sup> Let  $c(M)$  be the image of the morphism of  $A$ -modules*

$$\theta : M \otimes_A C^\vee \xrightarrow{\mu \otimes \operatorname{id}} V \otimes_A C \otimes_A C^\vee \xrightarrow{\operatorname{id} \otimes \operatorname{ev}} V.$$

*Then  $c(M)$  is the smallest subcomodule of  $V$  containing  $M$ , and it is a finitely generated  $A$ -module if  $M$  is. We call  $c(M)$  the subcomodule generated by  $M$ .*

*Moreover, for every homomorphism of rings  $A \rightarrow A'$ , if  $M'$  denotes the image of  $M \otimes_A A'$  in  $V' = V \otimes_A A'$ , then  $c(M')$  is the image of  $c(M) \otimes_A A'$  in  $V'$ . Thus the formation of  $c(M)$  commutes with base change.*

First,  $M \subset c(M)$  by (CM 2). If  $N$  is a subcomodule of  $V$  containing  $M$ , then  $\mu(M) \subset N \otimes C$ , and hence  $c(M) \subset N$ .

By hypothesis,  $C$  is a direct summand of a free  $A$ -module  $L$  with basis  $(e_i)_{i \in I}$ . Let  $\varphi_i$  denote the restriction to  $C$  of the linear form  $e_i^*$ , defined by

$$e_i^*(e_j) = \delta_{ij}.$$

Let  $x \in M$ . We may write

$$\mu(x) = \sum_{i \in J} x_i \otimes e_i, \tag{1}$$

where  $x_i \in V$  and  $J$  is a finite subset of  $I$ . Then

$$x_i = \theta(x \otimes \varphi_i) \in c(Ax),$$

and consequently

$$c(Ax) = \sum_{i \in J} Ax_i.$$

Since  $C$  is a direct summand of  $L$ , say  $L = C \oplus R$ , whence

$$V \otimes L = (V \otimes C) \oplus (V \otimes R),$$

we obtain

$$(c(Ax) \otimes L) \cap (V \otimes C) = c(Ax) \otimes C.$$

<sup>113</sup>Editor's note: in the original,  $C$  is assumed to be a free  $A$ -module. The generalization to the case where  $C$  is a projective  $A$ -module, pointed out by J.-P. Serre, is mentioned in Remark 11.10.1. We have incorporated this generalization here and in 11.9, and have expanded the proofs accordingly.

Consequently,  $\mu(x)$  may also be written in the form

$$\mu(x) = \sum_{j \in J} x_j \otimes b_j, \quad (2)$$

with  $b_j \in C$ . Write

$$\Delta(b_j) = \sum_{i \in I} b_{ij} \otimes e_i, \quad b_{ij} \in C.$$

Applying  $\mu \otimes \text{id}$  to (1) and  $\text{id} \otimes \Delta$  to (2), and using axiom (CM 1), we obtain, for every  $i \in J$ ,

$$\mu(x_i) = \sum_{j \in J} x_j \otimes b_{ij} \in c(Ax) \otimes C.$$

This proves that  $c(M)$  is a subcomodule of  $V$ , and therefore that it is the smallest subcomodule of  $V$  containing  $M$ .

It is clear that  $c(M)$  is a finitely generated  $A$ -module whenever  $M$  is: if

$$M = Ax_1 + \cdots + Ax_n, \quad \mu(x_k) = \sum_i x_{ik} \otimes e_i,$$

then the  $x_{ik}$ , for  $k = 1, \dots, n$  and  $i$  ranging over a finite subset of  $I$ , generate  $c(M)$ .

Finally, let  $A \rightarrow A'$  be a homomorphism of rings, and let  $M'$  be the image of  $M \otimes_A A'$  in  $V' = V \otimes_A A'$ . Then  $c(M')$ , respectively the image of  $c(M) \otimes_A A'$  in  $V'$ , is the image of the morphism  $\theta'$  below, respectively of the composite  $\theta' \circ \tau$ :

$$M \otimes_A A' \otimes_A C^\vee \xrightarrow{\tau} M \otimes_A \text{Hom}_A(C, A') \xrightarrow{\theta'} V'.$$

These two morphisms have the same image. Indeed, let  $\psi \in \text{Hom}_A(C, A')$  and  $x \in M$ , and write

$$\mu(x) = \sum_{i \in J} x_i \otimes e_i.$$

Then

$$\theta'(x \otimes \psi) = \sum_{i \in J} \psi(e_i) x_i$$

is the image under  $\theta' \circ \tau$  of the element

$$\sum_{i \in J} x \otimes \psi(e_i) \otimes \varphi_i \in M \otimes_A A' \otimes_A C^\vee.$$

This proves the lemma.

We also have the following proposition.

**Proposition 11.8.bis.**—<sup>114</sup> *Let  $A$  be a Noetherian ring, let  $C$  be an  $A$ -coalgebra flat over  $A$ , let  $V$  be a  $C$ -comodule, and let  $M$  be a finitely generated  $A$ -submodule of  $V$ . Then there is a subcomodule  $W$  of  $V$ , finitely generated over  $A$ , that contains  $M$ .*

Indeed, since  $M$  is finitely generated, so is  $\Delta_V(M)$ . Hence there is a finitely generated  $A$ -submodule  $M'$  of  $V$  such that

$$\Delta_V(M) \subset M' \otimes_A C.$$

Let  $\pi : V \rightarrow V/M'$  be the projection, put

$$\overline{\Delta}_V = (\pi \otimes \text{id}_C) \Delta_V,$$

<sup>114</sup>Editor's note: this proposition, taken from [Se68], Section 1.5, Proposition 2, has been added.

and set

$$W = \{x \in V \mid \Delta_V(x) \in M' \otimes_A C\} = \ker \bar{\Delta}_V.$$

This is an  $A$ -submodule of  $V$  that contains  $M$  and is contained in  $M'$ , because

$$x = (\mathrm{id}_V \otimes \varepsilon)\Delta_V(x).$$

It is therefore finitely generated over  $A$ . Moreover,

$$(\bar{\Delta}_V \otimes \mathrm{id}_C)\Delta_V = (\pi \otimes \Delta_C)\Delta_V$$

vanishes on  $W$ ; in other words,  $\Delta_V(W)$  is contained in the kernel  $K$  of  $\bar{\Delta}_V \otimes \mathrm{id}_C$ . Since  $C$  is flat over  $A$ , one has

$$K = W \otimes_A C.$$

Thus  $W$  is a subcomodule of  $V$ .

**Lemma 11.8.1.**—<sup>115</sup> *Let  $C$  be an  $A$ -coalgebra, let  $V$  be a  $C$ -comodule, let  $M$  be an  $A$ -submodule of  $V$ , and let  $f : A \rightarrow A'$  be a faithfully flat homomorphism of rings. Suppose that  $C' = C \otimes_A A'$  is a projective  $A'$ -module.*

- (i) *There is a smallest subcomodule  $t(M)$  of  $V$  containing  $M$ , and  $t(M)$  is a finitely generated  $A$ -module if  $M$  is. Moreover, the formation of  $t(M)$  commutes with base change.*
- (ii) *More precisely,  $C$  is a projective  $A$ -module, and  $t(M) = c(M)$ .*

*Proof.* For (ii), [RG71] (see Proposition 11.8.2 below) shows that  $C$  is a projective  $A$ -module. We may therefore apply Lemma 11.8:  $c(M)$  is the smallest subcomodule of  $V$  containing  $M$ , it is a finitely generated  $A$ -module if  $M$  is, and its formation commutes with base change.

To avoid an anachronism ([RG71] postdates SGA 3), let us sketch a direct proof of (i). Since  $A \rightarrow A'$  is flat,

$$M' = M \otimes_A A'$$

is an  $A'$ -submodule of  $V' = V \otimes_A A'$ . Since  $C'$  is a projective  $A'$ -module,  $c(M')$  is the smallest subcomodule of  $V'$  containing  $M'$ . Denote by  $V' \otimes_A A'$  and  $A' \otimes_A V'$  the two  $A' \otimes_A A'$ -comodule structures on

$$V'' = V' \otimes_{A'} (A' \otimes_A A')$$

obtained by the two base changes

$$A' \rightrightarrows A' \otimes_A A', \quad a' \mapsto a' \otimes 1, \quad a' \mapsto 1 \otimes a'.$$

The  $A'$ -comodule  $V'$  carries an isomorphism of  $A' \otimes_A A'$ -comodules

$$\varphi : V' \otimes_A A' \xrightarrow{\sim} A' \otimes_A V', \quad (x \otimes a') \otimes b' \mapsto b' \otimes (x \otimes a'),$$

which is a descent datum, that is,

$$\varphi_{31} = \varphi_{32} \circ \varphi_{21}.$$

---

<sup>115</sup>Editor's note: this lemma, which is used in the proof of 11.9, has been added.



Since  $M' = M \otimes_A A'$ , the isomorphism  $\varphi$  takes  $M' \otimes_A A'$  onto  $A' \otimes_A M'$ , and hence takes  $c(M' \otimes_A A')$  onto  $c(A' \otimes_A M')$ . Because the formation of  $c(M')$  commutes with base change,

$$c(M' \otimes_A A') = c(M') \otimes_A A', \quad c(A' \otimes_A M') = A' \otimes_A c(M').$$

Thus

$$\varphi(c(M') \otimes_A A') = A' \otimes_A c(M'),$$

and  $\varphi$  equips  $c(M')$  with a descent datum. By fpqc descent, there is a unique subcomodule  $t(M)$  of  $V$  such that

$$c(M') = t(M) \otimes_A A'.$$

It contains  $M$ , since  $t(M) \otimes_A A'$  contains  $M'$ . Moreover, if  $N$  is a subcomodule of  $V$  containing  $M$ , then  $N$  contains  $t(M)$ , because  $N \otimes_A A'$  contains

$$c(M') = t(M) \otimes_A A'.$$

Hence  $t(M)$  is the smallest subcomodule of  $V$  containing  $M$ .

Finally, let  $A \rightarrow B$  be a homomorphism of rings. Put

$$B' = B \otimes_A A',$$

and let  $M_B$ , respectively  $M'_{B'}$ , be the image of  $M \otimes_A B$  in  $V_B = V \otimes_A B$ , respectively of  $M' \otimes_{A'} B'$  in  $V' \otimes_{A'} B' = V \otimes_A B'$ . Then

$$M_B \otimes_B B' = M'_{B'}.$$

On the one hand, the preceding construction, applied to  $C_B$  and  $B \rightarrow B'$ , gives

$$c(M_B \otimes_B B') = t(M_B) \otimes_B B' = t(M_B) \otimes_A A'.$$

On the other hand, because the formation of  $c(M')$  commutes with base change,  $c(M'_{B'})$  is the image in

$$V' \otimes_{A'} B' = V \otimes_A B \otimes_A A'$$

of

$$c(M') \otimes_{A'} B' = t(M) \otimes_A B \otimes_A A'.$$

It follows that  $t(M_B)$  is the image of  $t(M) \otimes_A B$  in  $V_B$ .

**Proposition 11.8.2 (Gruson–Raynaud).**— *Let  $f : A \rightarrow A'$  be a faithfully flat homomorphism. Then  $f$  “descends projectivity”: if  $M$  is an  $A$ -module and  $M \otimes_A A'$  is a projective  $A'$ -module, then  $M$  is a projective  $A$ -module.*

Indeed, by [RG71], II, 2.5.1,  $f$  descends the Mittag–Leffler condition, and therefore, by *loc. cit.*, II, 3.1.3,  $f$  descends projectivity.<sup>116</sup>

**Proposition 11.9.**—<sup>117</sup> *Let  $\mathcal{C}$  be an  $\mathcal{O}_S$ -coalgebra, let  $\mathcal{E}$  be a  $\mathcal{C}$ -comodule, and let  $\mathcal{F}$  be an  $\mathcal{O}_S$ -submodule of  $\mathcal{E}$ , all of them quasi-coherent. Suppose that  $S$  is covered by affine open subsets  $U_\alpha = \operatorname{Spec} A_\alpha$ , and that for each  $\alpha$  there is a faithfully flat homomorphism of rings  $A_\alpha \rightarrow A'_\alpha$  such that*

$$\Gamma(U_\alpha, \mathcal{C}) \otimes_{A_\alpha} A'_\alpha$$

<sup>116</sup>Editor’s note: let us also point out that assertion II, 2.5.2 of *loc. cit.*, which is more general than II, 2.5.1, is corrected in [Gr73]; this does not affect the case of faithfully flat homomorphisms.

<sup>117</sup>Editor’s note: as indicated in editor’s note 112, we have rewritten the statement for an  $\mathcal{O}_S$ -coalgebra  $\mathcal{C}$  (rather than for an  $S$ -group  $G$  satisfying the hypotheses in Corollary 11.10). We have also expanded the proof (the original merely said: “... the proposition follows from Lemma 11.8”).

is a projective  $A'_\alpha$ -module.<sup>(\*)</sup>

Then there is a smallest quasi-coherent subcomodule  $t(\mathcal{F})$  of  $\mathcal{E}$  containing  $\mathcal{F}$ , and  $t(\mathcal{F})$  is a finitely generated  $\mathcal{O}_S$ -module if  $\mathcal{F}$  is. Moreover, for every base change  $S' \rightarrow S$ , if  $\mathcal{F}'$  denotes the image of  $\mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{O}_{S'}$  in

$$\mathcal{E}' = \mathcal{E} \otimes_{\mathcal{O}_S} \mathcal{O}_{S'},$$

then  $t(\mathcal{F}')$  is the image of  $t(\mathcal{F}) \otimes_{\mathcal{O}_S} \mathcal{O}_{S'}$  in  $\mathcal{E}'$ . In other words, the formation of  $t(\mathcal{F})$  commutes with base change.

*Proof.* For each  $\alpha$ , let  $T_\alpha$  be the  $A_\alpha$ -module

$$T_\alpha = t(\Gamma(U_\alpha, \mathcal{F})).$$

By 11.8.1(i), the associated  $\mathcal{O}_{U_\alpha}$ -module  $\tilde{T}_\alpha$  is the smallest quasi-coherent subcomodule of  $\mathcal{E}|_{U_\alpha}$  containing  $\mathcal{F}|_{U_\alpha}$ ; and it is a finitely generated  $\mathcal{O}_{U_\alpha}$ -module if  $\mathcal{F}|_{U_\alpha}$  is.

For every  $\alpha, \beta$ , put

$$U_{\alpha\beta} = U_\alpha \cap U_\beta.$$

Since the construction of  $t(M)$  commutes with base change, there are canonical isomorphisms of  $\mathcal{O}_{U_{\alpha\beta}}$ -modules

$$\varphi_{\alpha\beta} : \tilde{T}_\beta \otimes_{A_\beta} \mathcal{O}_{U_{\alpha\beta}} \xrightarrow{\sim} \tilde{T}_\alpha \otimes_{A_\alpha} \mathcal{O}_{U_{\alpha\beta}}.$$

They satisfy the cocycle condition

$$\varphi'_{\alpha\gamma} = \varphi'_{\alpha\beta} \circ \varphi'_{\beta\gamma},$$

where  $\varphi'_{\alpha\gamma}$ , respectively the other maps, denotes the restriction of  $\varphi_{\alpha\gamma}$ , respectively the other maps, to  $U_\alpha \cap U_\beta \cap U_\gamma$ .

Consequently, the  $\tilde{T}_\alpha$  glue to a quasi-coherent subcomodule  $t(\mathcal{F})$  of  $\mathcal{E}$  containing  $\mathcal{F}$ . We leave it to the reader to verify that  $t(\mathcal{F})$  is the smallest quasi-coherent subcomodule of  $\mathcal{E}$  containing  $\mathcal{F}$ , and that its formation commutes with base change.

**Definition 11.9.1.**—<sup>118</sup> Let  $S$  be a scheme and let  $\mathcal{P}$  be a quasi-coherent  $\mathcal{O}_S$ -module. The following conditions are equivalent:

- (i) For every affine open subset  $U$  of  $S$ , the module  $\Gamma(U, \mathcal{P})$  is a projective  $\mathcal{O}_S(U)$ -module.
- (ii) There is a covering  $(U_\alpha)$  of  $S$  by affine open subsets such that every  $\Gamma(U_\alpha, \mathcal{P})$  is a projective  $\mathcal{O}_S(U_\alpha)$ -module.
- (iii) There is a covering  $(U_\alpha)$  of  $S$  by affine open subsets, and faithfully flat homomorphisms of rings

$$A_\alpha = \mathcal{O}_S(U_\alpha) \longrightarrow A'_\alpha,$$

such that, for every  $\alpha$ ,

$$\Gamma(U_\alpha, \mathcal{P}) \otimes_{A_\alpha} A'_\alpha$$

is a projective  $A'_\alpha$ -module.

---

<sup>\*</sup>This holds, for example, when  $\mathcal{C} = \mathcal{A}(G)$ , where  $G$  is a reductive  $S$ -group, as we shall see in Exposé XXII, 5.7.8.

<sup>118</sup>Editor's note: this definition, taken from the bottom of page 82 of [RG71], has been added. Thus the hypothesis in Proposition 11.9 is that the coalgebra  $\mathcal{C}$  be a locally projective  $\mathcal{O}_S$ -module, and we have used this terminology in Corollary 11.10.

It is clear that

$$(i) \implies (ii) \implies (iii).$$

Conversely, if (iii) holds, 11.8.2 implies that every  $\Gamma(U_\alpha, \mathcal{P})$  is a projective  $\mathcal{O}_S(U_\alpha)$ -module, and hence (ii) holds.

Finally, suppose that (ii) holds, and let

$$V = \operatorname{Spec} A$$

be an arbitrary affine open subset. It is covered by finitely many affine open subsets

$$V_1, \dots, V_n, \quad V_i = \operatorname{Spec} A_i,$$

each contained in at least one  $V \cap U_\alpha$ , so that  $\Gamma(V_i, \mathcal{P})$  is a projective  $A_i$ -module. Put

$$A' = A_1 \times \dots \times A_n.$$

Then  $A \rightarrow A'$  is faithfully flat and

$$\Gamma(V, \mathcal{P}) \otimes_A A'$$

is a projective  $A'$ -module. Therefore 11.8.2 implies that  $\Gamma(V, \mathcal{P})$  is a projective  $A$ -module.

When these equivalent conditions hold, we say that  $\mathcal{P}$  is a *locally projective*  $\mathcal{O}_S$ -module.

**Corollary 11.10.**— *Let  $S$  be a quasi-compact and quasi-separated scheme, let  $G$  be an  $S$ -group, and let  $\rho$  be a linear action of  $G$  on a quasi-coherent  $\mathcal{O}_S$ -module  $\mathcal{E}$ . Suppose that:*

(i)  *$G$  satisfies one of the following conditions:*

(a)  *$G$  is affine and flat over  $S$ ;*

(b)  *$G$  is flat, quasi-compact, and quasi-separated over  $S$ , and  $\mathcal{E}$  is flat;*

(ii)  *$\mathcal{A}(G)$  is a locally projective  $\mathcal{O}_S$ -module.*

*Then  $\mathcal{E}$  is the direct limit of an increasing filtered family of finitely generated quasi-coherent  $\mathcal{O}_S$ -submodules of  $\mathcal{E}$  that are stable under  $G$ .*

By hypothesis (i) and 11.6.1,  $\mathcal{E}$  is endowed with an  $\mathcal{A}(G)$ -comodule structure. On the other hand, since  $S$  is quasi-compact and quasi-separated,  $\mathcal{E}$  is the direct limit of its finitely generated quasi-coherent submodules (EGA I, 9.4.9 and EGA IV<sub>1</sub>, 1.7.7). Consequently, the corollary follows from Proposition 11.9, applied to the coalgebra  $\mathcal{A}(G)$ .

We also have the following proposition.

**Proposition 11.10.bis.**<sup>119</sup> *Let  $S$  be a Noetherian scheme, let  $G$  be an  $S$ -group that is flat, quasi-compact, and quasi-separated over  $S$ , let  $\mathcal{E}$  be a quasi-coherent  $G$ - $\mathcal{O}_S$ -module, and let  $\mathcal{M}$  be a coherent  $\mathcal{O}_S$ -submodule of  $\mathcal{E}$ . Then  $\mathcal{M}$  is contained in a coherent  $\mathcal{O}_S$ -submodule stable under  $G$ .*

*Proof.* Write  $f$  for the morphism  $G \rightarrow S$ , and  $\tau$  for the adjunction morphism

$$\mathcal{E} \longrightarrow f_* f^*(\mathcal{E}).$$

By 11.6, the  $G$ - $\mathcal{O}_S$ -module structure on  $\mathcal{E}$  is given by an automorphism  $\theta$  of the  $\mathcal{O}_G$ -module  $f^*(\mathcal{E})$ , such that the morphism

$$\delta = \theta \circ \tau$$

<sup>119</sup>Editor's note: this proposition has been added; it is a special case of [Th87], 1.4–1.5. The author there refers to an argument of Deligne (cf. [Kn71], III, Th. 1.1); one may also note its similarity with Serre's argument ([Se68], Prop. 2) recalled in 11.8.bis.

satisfies conditions (1) and (2) of 11.6. (The situation considered in [Th87] is more general: the author considers a  $G$ -scheme  $X$  and a  $G$ -equivariant  $\mathcal{O}_X$ -module  $\mathcal{E}$  (cf. Exposé I, Section 6); here  $X = S$ , endowed with the trivial action of  $G$ .)

Since  $S$  is Noetherian,  $\mathcal{E}$  is the filtered direct limit of its coherent submodules  $\mathcal{F}_\alpha$  (cf. EGA I, 9.4.9). Then  $f^*(\mathcal{E})$  is the filtered direct limit of the  $f^*(\mathcal{F}_\alpha)$ , which are submodules of  $f^*(\mathcal{E})$  because  $f$  is flat. Moreover, since  $f$  is quasi-compact and quasi-separated, by 11.0 the  $\mathcal{O}_S$ -module  $f_*f^*(\mathcal{E})$  is quasi-coherent and is the filtered direct limit of the quasi-coherent submodules  $f_*f^*(\mathcal{F}_\alpha)$ . Consequently,  $\mathcal{E}$  is the filtered direct limit of the quasi-coherent  $\mathcal{O}_S$ -submodules  $\mathcal{E}_\alpha$ , where  $\mathcal{E}_\alpha$  denotes the inverse image under  $\delta$  of  $f_*f^*(\mathcal{F}_\alpha)$ ; equivalently, it is the kernel of the composite

$$\bar{\delta}_\alpha : \mathcal{E} \xrightarrow{\delta} f_*f^*(\mathcal{E}) \longrightarrow f_*f^*(\mathcal{E}/\mathcal{F}_\alpha).$$

Since  $\mathcal{M}$  is coherent and  $S$  is Noetherian, every increasing sequence of submodules of  $\mathcal{M}$  is stationary; hence  $\mathcal{M}$  is contained in some  $\mathcal{E}_\alpha$ . We shall show that every  $\mathcal{E}_\alpha$  is coherent and  $G$ -stable.

Let

$$u : f^*(\mathcal{E}) \longrightarrow \varepsilon_*(\mathcal{E})$$

be the morphism corresponding by adjunction to the identity morphism from  $\mathcal{E}$  to  $f_*\varepsilon_*(\mathcal{E}) = \mathcal{E}$ . Then the composite

$$\mathcal{E} \xrightarrow{\tau} f_*f^*(\mathcal{E}) \xrightarrow{f_*(u)} f_*\varepsilon_*(\mathcal{E}) = \mathcal{E}$$

is the identity (cf. 11.6(2)). We have the commutative diagram

$$\begin{array}{ccccc} \mathcal{E}_\alpha & \xrightarrow{\delta} & f_*f^*(\mathcal{F}_\alpha) & \xrightarrow{f_*(u)} & \mathcal{F}_\alpha \\ i_\alpha \downarrow & & \downarrow & & \downarrow j_\alpha \\ \mathcal{E} & \xrightarrow{\tau} & f_*f^*(\mathcal{E}) & \xrightarrow{f_*(u)} & \mathcal{E} \end{array}$$

where  $i_\alpha$  (respectively  $j_\alpha$ ) denotes the inclusion of  $\mathcal{E}_\alpha$  (respectively  $\mathcal{F}_\alpha$ ) in  $\mathcal{E}$ . It follows that  $i_\alpha$  factors through  $j_\alpha$ ; thus  $\mathcal{E}_\alpha$  is a submodule of  $\mathcal{F}_\alpha$ , and is therefore coherent.

Finally, let us show that  $\mathcal{E}_\alpha$  is  $G$ -stable (cf. 11.7.bis). Let  $h : G \rightarrow S$  denote a second copy of  $f : G \rightarrow S$ , and consider the following commutative diagram:

$$\begin{array}{ccc} G \times_S G & \xrightarrow{q} & G \\ p \downarrow & \searrow \varphi & \downarrow h \\ G & \xrightarrow{f} & S \end{array}$$

The exact sequence

$$0 \longrightarrow \mathcal{E}_\alpha \longrightarrow \mathcal{E} \xrightarrow{\bar{\delta}_\alpha} h_*h^*(\mathcal{E}/\mathcal{F}_\alpha)$$

therefore gives, since  $f$  is flat, the exact sequence

$$0 \longrightarrow f_*f^*(\mathcal{E}_\alpha) \longrightarrow f_*f^*(\mathcal{E}) \xrightarrow{f_*f^*(\bar{\delta}_\alpha)} f_*f^*h_*h^*(\mathcal{E}/\mathcal{F}_\alpha). \quad (\dagger)$$

Moreover, since  $h$  is quasi-compact and quasi-separated and  $f$  is flat, the canonical morphism

$$f^*h_*h^*(\mathcal{E}/\mathcal{F}_\alpha) \longrightarrow p_*q^*h^*(\mathcal{E}/\mathcal{F}_\alpha) = p_*\varphi^*(\mathcal{E}/\mathcal{F}_\alpha)$$

is an isomorphism. Thus the right-hand term in (†) is  $\varphi_*\varphi^*(\mathcal{E}/\mathcal{F}_\alpha)$ . Using the notation of 11.6 and writing  $\pi_\alpha$  for the projection  $\mathcal{E} \rightarrow \mathcal{E}/\mathcal{F}_\alpha$ , we obtain the following commutative diagram, whose bottom row is exact:

$$\begin{array}{ccccc} \mathcal{E}_\alpha & \xrightarrow{\delta} & f_*f^*(\mathcal{E}) & & \\ \delta \downarrow & & \downarrow f_*(f^*(\pi_\alpha) \boxtimes \delta_G) & & \\ f_*f^*(\mathcal{E}_\alpha) & \longrightarrow & f_*f^*(\mathcal{E}) & \xrightarrow{f_*f^*(\bar{\delta}_\alpha)} & \varphi_*\varphi^*(\mathcal{E}/\mathcal{F}_\alpha) \end{array}$$

Since

$$f_*(f^*(\pi_\alpha) \boxtimes \delta_G) \circ \delta$$

vanishes on  $\mathcal{E}_\alpha$ , it follows that  $\delta$  carries  $\mathcal{E}_\alpha$  into  $f_*f^*(\mathcal{E}_\alpha)$ ; in other words,  $\mathcal{E}_\alpha$  is  $G$ -stable. This proves the proposition.

**Remark 11.10.1.**—<sup>120</sup> In 11.8, it is not enough to assume that  $C$  is a flat  $A$ -module, even when  $A$  is a principal ideal domain. Indeed, we have the following counterexamples, pointed out to us independently by O. Gabber and J.-P. Serre.

(a) Let  $(A, \mathfrak{m})$  be a discrete valuation ring, let  $K$  be its field of fractions, and let  $G$  be the  $A$ -group obtained by “extension by zero” from the  $K$ -group  $(\mathbb{Z}/2\mathbb{Z})_K$ . The constant group  $(\mathbb{Z}/2\mathbb{Z})_A$ , and hence also its subgroup  $G$ , acts on the free  $A$ -module  $V$  with basis  $v_1, v_2$  by interchanging  $v_1$  and  $v_2$ . Then  $Av_1$  is not a  $G$ -submodule of  $V$ , but it is the intersection of the  $G$ -submodules

$$Av_1 + \mathfrak{m}^n v_2, \quad n \in \mathbb{N}^*.$$

Thus there is no smallest  $G$ -submodule of  $V$  containing  $v_1$ .

(b) Let  $A$  be an integral domain distinct from its field of fractions  $K$ , and let  $G$  be the flat affine  $A$ -group corresponding to the Hopf algebra

$$\mathcal{A}(G) = \{P \in K[T] \mid P(0) \in A\}.$$

Its comultiplication, counit, and antipode are respectively defined by

$$\Delta(T) = T \otimes 1 + 1 \otimes T, \quad \varepsilon(T) = 0, \quad \tau(T) = -T.$$

(Thus  $G \otimes_A K = \mathbb{G}_{a,K}$ .)

Let  $V = Av_1 \oplus Av_2$  be free over  $A$ , and let  $u$  be the endomorphism of  $V$  defined by

$$u(v_1) = v_2, \quad u(v_2) = 0,$$

so that  $u^2 = 0$ . Then  $V$  has the  $\mathcal{A}(G)$ -comodule structure

$$\mu(m) = 1 \otimes m + T \otimes u(m).$$

The  $G$ -submodules of  $V$  containing  $v_1$  are precisely the  $A$ -submodules  $Av_1 \oplus Iv_2$ , where  $I$  is a nonzero ideal of  $A$ . Their intersection is  $Av_1$ , which is not a  $G$ -submodule. Hence there is no smallest  $G$ -submodule of  $V$  containing  $v_1$ . Moreover,

$$C = A \oplus K \cdot T$$

is a subcoalgebra of  $\mathcal{A}(G)$ , flat over  $A$ , and the coaction

$$\mu : V \longrightarrow V \otimes \mathcal{A}(G)$$

<sup>120</sup>Editor’s note: the first part of the original Remark 11.10.1 has been incorporated into 11.8 and 11.9 (with “free” replaced by “projective”); the counterexamples below correct its second part.

factors through  $V \otimes C$ . Thus the “very simple” coalgebra  $C$  also gives a counterexample.

Finally, the preceding two examples are special cases of the following construction. Let  $A$  be an integral domain distinct from its field of fractions  $K$ , and let  $B$  be a Hopf  $A$ -algebra, free over  $A$ . Write

$$\varepsilon : B \longrightarrow A, \quad I = \ker(\varepsilon)$$

for the augmentation and the augmentation ideal. Since  $B = A \cdot 1 \oplus I$ , one readily sees that

$$B' = \{b \in B \otimes_A K \mid \varepsilon(b) \in A\}$$

is a Hopf subalgebra of  $B_K$ . Suppose that  $V$  is a  $B$ -comodule, free with basis  $(v_1, \dots, v_n)$  as an  $A$ -module, and that  $\mu_V(v_1) \neq v_1 \otimes 1$ . Then  $Av_1$  is not a subcomodule of  $V$ , but it is the intersection of the subcomodules

$$Av_1 + J \cdot V$$

as  $J$  ranges over the nonzero ideals of  $A$ .<sup>\*</sup> Thus there is no smallest subcomodule of  $V$  containing  $v_1$ .

**Proposition 11.11.**— *Let  $G$  be an algebraic group over a field  $k$ . The following conditions are equivalent:*

- (i)  $G$  is affine.
- (ii)  $G$  is quasi-affine.
- (iii)  $G$  acts faithfully on a quasi-affine  $k$ -scheme  $X$ .
- (iv)  $G$  acts linearly and faithfully on a  $k$ -vector space (not necessarily finite-dimensional).
- (v)  $G$  is isomorphic to a closed subgroup of  $\mathrm{GL}(n)_k$ .

*Proof.* The implication (i)  $\Rightarrow$  (ii) is immediate, and (ii)  $\Rightarrow$  (iii) because  $G$  acts faithfully on itself by translations.

Suppose that  $G$  acts faithfully on the right on a quasi-affine  $k$ -scheme  $X$ .<sup>121</sup> Since  $X$  is quasi-affine, it is separated and quasi-compact.<sup>122</sup> Similarly,  $G$  is separated (VI<sub>A</sub>, 0.3) and quasi-compact (since it is of finite type over  $k$ ). Hence, by 11.1(c), there are canonical isomorphisms

$$\mathcal{O}(X \times G) = \mathcal{O}(X) \otimes \mathcal{O}(G), \quad \mathcal{O}(X \times G \times G) = \mathcal{O}(X) \otimes \mathcal{O}(G) \otimes \mathcal{O}(G).$$

It follows that the morphism

$$\mu : \mathcal{O}(X) \longrightarrow \mathcal{O}(X) \otimes \mathcal{O}(G)$$

induced by  $X \times G \rightarrow X$  gives  $\mathcal{O}(X)$  the structure of a right  $\mathcal{O}(G)$ -comodule; that is,  $G$  acts linearly on the left on the  $k$ -algebra  $\mathcal{O}(X)$ . Consequently,  $G$  also acts on the right on the affine envelope

$$X_{\mathrm{af}} = \mathrm{Spec} \mathcal{O}(X)$$

<sup>\*</sup> Translator’s note: the French re-edition uses  $I$  both for the fixed augmentation ideal above and for the ideals ranging in this final intersection. We use  $J$  for the latter to keep the two roles distinct.

<sup>121</sup>Editor’s note: the original has been expanded in what follows. We have also chosen a right action of  $G$  on  $X$ , so as to obtain a left linear action of  $G$  on  $\mathcal{O}(X)$ .

<sup>122</sup>Editor’s note: recall that a  $k$ -scheme  $X$  is called quasi-affine if it is isomorphic to a quasi-compact open subscheme of an affine  $k$ -scheme (EGA II, 5.1.1).

of  $X$ , and the canonical morphism  $X \rightarrow X_{\text{af}}$  is  $G$ -equivariant.

Since  $X$  is quasi-affine, the morphism  $X \rightarrow X_{\text{af}}$  is an open immersion (EGA II, 5.1.2). Hence, a fortiori,  $G$  acts faithfully on  $X_{\text{af}}$ , and therefore the left linear action of  $G$  on the  $k$ -algebra  $\mathcal{O}(X) = \mathcal{O}(X_{\text{af}})$  is faithful. This proves (iii)  $\Rightarrow$  (iv).

Now suppose that  $G$  acts faithfully on a  $k$ -vector space  $V$ . By 11.10,  $V$  is the direct limit of finite-dimensional vector subspaces  $V_i$  stable under  $G$ . Let  $K_i$  be the kernel of the induced action of  $G$  on  $V_i$ ; that is, the kernel of  $G \rightarrow \text{Aut}(V_i)$ . Then  $K_i$  is a closed subscheme of  $G$ , and the faithfulness of the action of  $G$  means that the intersection of the  $K_i$  is the identity subgroup of  $G$ . Since  $G$  is Noetherian, one of the  $K_i$  is already the identity subgroup. Thus  $G \rightarrow \text{Aut}(V_i)$  is a monomorphism, and hence, by 1.4.2, a closed immersion. This proves (iv)  $\Rightarrow$  (v). Finally, (v)  $\Rightarrow$  (i) is immediate, and the proposition follows.

**Remark 11.11.1.**— Proposition 11.11 admits the following generalization. Let  $S$  be a regular, locally Noetherian scheme of dimension at most 1, and let  $G$  be a group scheme that is flat, quasi-compact, and quasi-separated over  $S$ . (In this case,  $\mathcal{A}(G)$  is a torsion-free  $\mathcal{O}_S$ -module, and hence is flat.)

(a) The following conditions are equivalent:<sup>123</sup>

- (i)  $G$  is affine over  $S$ .
- (ii)  $G$  is quasi-affine over  $S$ .
- (iii)  $G$  acts faithfully on a quasi-affine flat  $S$ -scheme.
- (iv)  $G$  acts linearly and faithfully on a flat quasi-coherent  $\mathcal{O}_S$ -module.

(b) If, moreover,  $G$  is of finite type over  $S$  and  $S$  is Noetherian, these conditions imply that  $G$  is isomorphic to a closed subgroup of  $\text{Aut}(\mathcal{E})$ , where  $\mathcal{E}$  is a finite-rank locally free  $\mathcal{O}_S$ -module.<sup>124</sup>

**Lemma 11.12.**— Let  $k$  be a field and let  $G$  be an affine  $k$ -group. Put  $A = \mathcal{A}(G)$ . Given  $x \in A$ , there is a finitely generated  $k$ -subalgebra  $B$  of  $A$  such that

$$x \in B, \quad \Delta(B) \subset B \otimes_k B, \quad u(B) \subset B,$$

where  $u$  denotes the involution of  $A$  corresponding to the inversion morphism of  $G$ .<sup>125</sup>

We may suppose that  $x \neq 0$ . Then  $\Delta(x) \neq 0$ , since

$$(\varepsilon \otimes \text{id})\Delta(x) = x,$$

where  $\varepsilon$  denotes the augmentation (counit) of  $A$ . Write

$$\Delta(x) = \sum_{j=1}^n e_j \otimes a_j, \quad n \text{ minimal.} \tag{1}$$

The  $e_j$  (respectively the  $a_j$ ) are then linearly independent. Complete  $(e_1, \dots, e_n)$  to a basis  $(e_i)_{i \in I}$  of  $A$ , and, for

$$j \in J = \{1, \dots, n\},$$

<sup>123</sup>Editor's note: the equivalence of these conditions is proved in the added Section 12.

<sup>124</sup>Editor's note: this is proved, with various generalizations, in the added Section 13.

<sup>125</sup>Editor's note: on the one hand, this is generalized in Section 13 to the case where  $G$  is affine and flat over a regular base of dimension at most 2. On the other hand, the original has been expanded and corrected in what follows.

write

$$\Delta(e_j) = \sum_{i \in I} e_i \otimes b_{ij}.$$

Applying  $\Delta \otimes \text{id}$  and  $\text{id} \otimes \Delta$  to (1), the axiom (CO 1) of 11.8.0 (see also (HA 1) in I, 4.2) gives

$$\sum_{j \in J} e_j \otimes \Delta(a_j) = \sum_{\ell \in J} \Delta(e_\ell) \otimes a_\ell = \sum_{i \in I} e_i \otimes \left( \sum_{\ell \in J} b_{i\ell} \otimes a_\ell \right).$$

Since the  $e_i$  are linearly independent, it follows that

$$\Delta(a_j) = \sum_{\ell \in J} b_{j\ell} \otimes a_\ell \quad (j \in J). \quad (2)$$

Let  $B$  be the finitely generated  $k$ -subalgebra of  $A$  generated by the  $b_{ij}$  and the  $u(b_{ij})$ , for  $i, j \in J$ . It is clear that  $u(B) = B$ .

Applying  $\Delta \otimes \text{id}$  and  $\text{id} \otimes \Delta$  to (2), we obtain once again from (CO 1) the equalities

$$\sum_{\ell \in J} \Delta(b_{j\ell}) \otimes a_\ell = \sum_{i \in J} b_{ji} \otimes \Delta(a_i) = \sum_{i, \ell \in J} b_{ji} \otimes b_{i\ell} \otimes a_\ell.$$

Since the  $a_\ell$  are linearly independent, this yields

$$\Delta(b_{j\ell}) = \sum_{i \in J} b_{ji} \otimes b_{i\ell} \quad (j, \ell \in J). \quad (3)$$

Moreover,

$$\Delta \circ u = (u \otimes u) \circ v \circ \Delta, \quad v(a \otimes b) = b \otimes a,$$

and hence

$$\Delta(u(b_{j\ell})) = \sum_{i \in J} u(b_{i\ell}) \otimes u(b_{ji}) \quad (j, \ell \in J). \quad (4)$$

Since  $\Delta$  is an algebra homomorphism, (3) and (4) imply

$$\Delta(B) \subset B \otimes_k B.$$

Finally, axiom (CO 2) of 11.8.0 (see also (HA 2) in I, 4.2), applied to (2) and (1), gives\*

$$a_j = \sum_{i \in J} b_{ji} \varepsilon(a_i), \quad x = \sum_{j \in J} \varepsilon(e_j) a_j.$$

Thus  $x \in B$ , proving the lemma.

**Proposition 11.13.**— *Let  $k$  be a field and let  $G$  be an affine  $k$ -group with algebra  $A$ . Then  $G$  is the projective limit of an increasing filtered system of affine  $k$ -groups of finite type whose transition morphisms are faithfully flat.*

If  $B$  and  $B'$  are two finitely generated subalgebras of  $A$  stable under  $\Delta$  and  $u$ , the subalgebra generated by  $B$  and  $B'$  has the same properties. Hence, by Lemma 11.12,  $A$  is the direct limit of an increasing filtered family  $(B_i)_{i \in I}$  of finitely generated subalgebras stable under  $\Delta$  and  $u$ . Each  $B_i$ , equipped with the restriction of  $u$  and with the morphism

$$B_i \longrightarrow B_i \otimes_k B_i$$

---

\* Source-reading note: the French re-edition prints  $a_j = \sum_{i \in I} \varepsilon(a_i) b_{ij}$ . Applying  $\text{id} \otimes \varepsilon$  to (2) gives the first identity displayed here. The Reinhold comparison lineage repeats the printed formula.



induced by  $\Delta$ , is a Hopf algebra. Thus, by I, 4.2, it is the algebra of an affine  $k$ -group  $G_i$  of finite type over  $k$ . Finally, since

$$A = \varinjlim_i B_i,$$

we have

$$G = \varprojlim_i G_i$$

(cf. EGA IV<sub>3</sub>, 8.2.3). The transition morphisms are faithfully flat by the following lemma.

**Lemma 11.14.**— *Let  $k$  be a field, let  $u : G \rightarrow H$  be a morphism between affine  $k$ -groups of finite type, and let*

$$u^\natural : B \longrightarrow A$$

*be the corresponding morphism of  $k$ -algebras.<sup>126</sup> Then  $u$  is faithfully flat if and only if  $u^\natural$  is injective.*

The condition is plainly necessary (cf. EGA 0<sub>I</sub>, 6.6.1). Conversely, suppose that  $u^\natural$  is injective, and put

$$N = \text{Ker}(u).$$

By VI<sub>A</sub>, 3.3.2 and VI<sub>A</sub>, 5.4.1,  $G/N$  is a  $k$ -group of finite type and  $u$  factors as

$$G \xrightarrow{p} G/N \xrightarrow{v} H,$$

where  $p$  is faithfully flat and  $v$  is a closed immersion. Since  $H$  is affine,  $G/N$  is affine, and

$$v^\natural : B \longrightarrow \mathcal{O}(G/N)$$

is surjective (cf. EGA I, 4.2.3). But

$$u^\natural = p^\natural \circ v^\natural$$

is injective, so  $v^\natural$  is also injective. Hence  $v^\natural$ , and therefore  $v$ , is an isomorphism. Since  $p$  is faithfully flat, so is  $u$ .

**Definition 11.15.**—<sup>127</sup> *Let  $k$  be a field, let  $G$  be a quasi-compact  $k$ -group, and let  $V$  be a  $k$ -vector space endowed with a  $k$ -linear action of  $G$ , hence, by 11.6.1, with an  $\mathcal{A}(G)$ -comodule structure*

$$\delta : V \longrightarrow V \otimes \mathcal{A}(G).$$

*Let  $v \in V$  be nonzero. The following conditions are equivalent:*

- (i) *There is a  $\lambda \in \mathcal{A}(G)$ , necessarily unique, such that*

$$\delta(v) = v \otimes \lambda.$$

- (ii) *For every  $k$ -algebra  $R$  and every  $g \in G(R)$ , one has  $g \cdot v \in Rv$ ; that is, there is a necessarily unique  $f \in R$  such that, in  $V \otimes R$ ,*

$$g \cdot v = v \otimes f.$$

<sup>126</sup>Editor's note: we write  $u^\natural$  (instead of  $u^\circ$ ) for the morphism  $B \rightarrow A$  corresponding to  $u : \text{Spec}(A) \rightarrow \text{Spec}(B)$ .

<sup>127</sup>Editor's note: we have added the hypothesis that  $G$  be quasi-compact and have spelled out the equivalence of conditions (i) and (ii).

It is clear that (i) implies (ii). Conversely, if (ii) holds, apply it with

$$R = \mathcal{A}(G), \quad g = \mathrm{id}_{\mathcal{A}(G)}.$$

There is then a unique  $\lambda \in \mathcal{A}(G)$  such that  $\delta(v) = v \otimes \lambda$ .

When  $v$  satisfies these conditions, we call  $v$  a *semi-invariant vector under  $G$* , and  $\lambda$  the *weight* of  $v$ . We also say that  $v$  is a *semi-invariant of weight  $\lambda$* .

Let  $\Delta$  denote the comultiplication of  $\mathcal{A}(G)$ . The equality

$$v \otimes \lambda \otimes \lambda = (\delta \otimes \mathrm{id})(\delta(v)) = (\mathrm{id} \otimes \Delta)(\delta(v)) = v \otimes \Delta(\lambda)$$

implies

$$\Delta(\lambda) = \lambda \otimes \lambda.$$

Consequently,  $\lambda$  defines a morphism of Hopf algebras

$$\mathcal{A}(\mathbb{G}_{m,k}) = k[T, T^{-1}] \longrightarrow \mathcal{A}(G), \quad T \longmapsto \lambda,$$

and hence a morphism of  $k$ -groups

$$\lambda : G \longrightarrow \mathbb{G}_{m,k}.$$

Thus  $\lambda$  is a *character* of  $G$ , called the *character associated with the semi-invariant vector  $v$* .

**Lemma 11.16.0.**—<sup>128</sup> *Let  $k$  be a field, let  $H$  be an affine  $k$ -group, let  $V$  be an  $H$ -module of dimension  $n$ , and let  $U$  be a  $d$ -dimensional vector subspace of  $V$ . Consider the line*

$$D = \bigwedge^d U \subset \bigwedge^d V.$$

*Then  $U$  is stable under  $H$  if and only if  $D$  is stable under  $H$ .*

Necessity is clear. To prove sufficiency, we may assume  $d < n$ . Let  $(e_1, \dots, e_d)$  be a basis of  $U$ , and complete it to a basis  $(e_1, \dots, e_n)$  of  $V$ . For every  $k$ -algebra  $R$ , the module

$$V_R = V \otimes R$$

is free over  $R$ , and

$$U_R = \{v \in V_R \mid v \wedge (e_1 \wedge \cdots \wedge e_d) = 0\}.$$

Indeed, for  $i > d$ , the elements

$$e_i \wedge e_1 \wedge \cdots \wedge e_d$$

are linearly independent in  $\bigwedge_R^{d+1} V_R$ . Since  $H(R)$  acts on  $\bigwedge_R^\bullet V_R$  by

$$h(x_1 \wedge \cdots \wedge x_s) = h(x_1) \wedge \cdots \wedge h(x_s),$$

it follows that, if  $H(R)$  stabilizes

$$D_R = R(e_1 \wedge \cdots \wedge e_d),$$

then it also stabilizes  $U_R$ .

**Theorem 11.16 (Chevalley).**— *Let  $k$  be a field, let  $G$  be an affine algebraic  $k$ -group, and let  $H$  be a closed group subscheme of  $G$ .<sup>129</sup> Then there are a finite-dimensional  $G$ -module  $V$  and a line  $D$  in  $V$  such that*

$$H = \mathrm{Norm}_G(D);$$

<sup>128</sup>Editor's note: this lemma, taken from [DG70], Section II.2, 3.5, has been added.

<sup>129</sup>Editor's note: first, recall that every group subscheme of  $G$  is closed (1.4.2). Second, we have stated the result in its usual form, namely that “ $H$  is the stabilizer of a line in a representation of  $G$ ,” while retaining the original formulation in terms of a sequence  $a_1, \dots, a_n$  of semi-invariants in  $\mathcal{A}(G)$ .

that is, for every  $k$ -algebra  $R$ ,

$$H(R) = \{ g \in G(R) \mid g(D_R) = D_R \}.$$

Equivalently, there are finitely many elements  $a_1, \dots, a_n \in \mathcal{A}(G)$ , all semi-invariant of the same weight  $\lambda$  for the “right” action of  $H$ ; that is,

$$a_i(gh) = \lambda(h)a_i(g)$$

for every  $k$ -algebra  $R$ ,  $g \in G(R)$ , and  $h \in H(R)$ , such that  $H$  is the largest closed group subscheme of  $G$  under which the  $a_i$  are semi-invariant.

*Proof.* Let  $\Delta$  and  $\varepsilon$  denote, respectively, the comultiplication and the augmentation of

$$A = \mathcal{A}(G).$$

There is an ideal  $I$  of  $A$ , contained in  $\text{Ker } \varepsilon$ , such that

$$H = \text{Spec}(A/I), \quad \Delta(I) \subset I \otimes A + A \otimes I.$$

Put  $B = A/I$ , and let  $\pi : A \rightarrow B$  be the projection. Consider the left action of  $H$  on  $A$  given by

$$(h\phi)(g) = \phi(gh).$$

The corresponding  $B$ -comodule structure is

$$\overline{\Delta} : A \xrightarrow{\Delta} A \otimes A \xrightarrow{\text{id}_A \otimes \pi} A \otimes B.$$

Then  $I$  is an  $H$ -submodule of  $A$ , since

$$\overline{\Delta}(I) \subset I \otimes B.$$

The  $k$ -algebra  $A$  is finitely generated, and hence Noetherian, so  $I$  has a finite system of generators  $(x_1, \dots, x_r)$ . By 11.8, the  $x_i$  are contained in a finite-dimensional  $G$ -submodule  $V$  of  $A$ . Then

$$W = V \cap I$$

is a finite-dimensional  $H$ -module; write

$$d = \dim_k W.$$

Since  $V$  contains every  $x_i$ ,  $W$  generates the ideal  $I$ .

Put

$$E = \bigwedge^d V,$$

choose a basis  $(w_1, \dots, w_d)$  of  $W$ , and choose a basis  $(e_0, \dots, e_n)$  of  $E$  containing

$$e_0 = w_1 \wedge \dots \wedge w_d.$$

The action of  $G$  on  $V$  canonically induces an action of  $G$  on  $E$ , hence an  $A$ -comodule structure

$$\rho : E \longrightarrow E \otimes A.$$

For  $j = 0, \dots, n$ , write<sup>†</sup>

$$\rho(e_j) = \sum_{i=0}^n e_i \otimes b_{ij}.$$

---

<sup>†</sup> Source-reading note: the French re-edition and the Reinhold comparison lineage print the lower summation bound as  $i = 1$ , although the chosen basis begins with  $e_0$  and the ensuing identity sums from 0. The lower bound is therefore  $i = 0$  here.

As in the proof of 11.12, one has

$$\Delta(b_{ij}) = \sum_{\ell=0}^n b_{i\ell} \otimes b_{\ell j}. \quad (*)$$

Put  $a_i = b_{i0}$ . Equivalently, if  $(e_0^*, \dots, e_n^*)$  is the dual basis, then the  $a_i$  are the matrix coefficients

$$g \mapsto e_i^*(ge_0).$$

The action of  $H$  on  $E$  corresponds to

$$\bar{\rho}: E \xrightarrow{\rho} E \otimes A \xrightarrow{\text{id}_E \otimes \pi} E \otimes B.$$

Since  $W$  is stable under  $H$ , the vector  $e_0$  is semi-invariant under  $H$ . Hence

$$\bar{\rho}(e_0) = e_0 \otimes \pi(a_0), \quad a_i = b_{i0} \in I \quad (1 \leq i \leq n).$$

Substitution in  $(*)$  gives

$$\bar{\Delta}(a_i) = a_i \otimes \pi(a_0) \quad (0 \leq i \leq n).$$

Thus the  $a_i$  are semi-invariant under  $H$ , all of weight  $\pi(a_0)$ . Moreover, after replacing  $E$ , if necessary, by the  $G$ -submodule generated by  $e_0$  (cf. 11.8), we may suppose that the  $a_i$  are linearly independent.

Conversely, let

$$H' = \text{Spec } B', \quad B' = A/I',$$

be a closed group subscheme of  $G$  under which every  $a_i$  is semi-invariant, of weight  $\lambda_i \in B'$ . This includes, in particular, the case in which  $e_0$  is semi-invariant under  $H'$  of weight  $\lambda'$ . We shall prove that  $H' \subset H$ .<sup>‡</sup> Let  $\pi': A \rightarrow B'$  be the projection. The hypothesis gives

$$a_i \otimes \lambda_i = (\text{id}_A \otimes \pi')\Delta(b_{i0}) = \sum_{\ell=0}^n b_{i\ell} \otimes \pi'(a_\ell).$$

It follows that

$$\lambda_i = \pi'(a_0), \quad a_\ell \in I' \quad (1 \leq \ell \leq n),$$

and hence  $e_0$  is semi-invariant under  $H'$ . By Lemma 11.16.0,  $W$  is stable under  $H'$ . Since  $W$  generates  $I$ , the ideal  $I$  is also stable under  $H'$ , and therefore

$$\Delta(I) \subset I \otimes A + A \otimes I'.$$

Finally,

$$I \subset \text{Ker } \varepsilon, \quad (\varepsilon \otimes \text{id}_A) \circ \Delta = \text{id}_A,$$

so  $I \subset I'$ , whence  $H' \subset H$ . This completes the proof.

**Lemma 11.17.0.**<sup>130</sup> *Let  $k$  be an algebraically closed field, let  $G$  be a reduced affine algebraic  $k$ -group, let  $V$  be a finite-dimensional  $G$ -module over  $k$ , and let  $v \in V$ . Let  $E$  be*

<sup>‡</sup> Source-reading note: the French re-edition says at this point “let us show that  $H' = H$ ,” as does the Reinhold comparison lineage. The maximality assertion requires  $H' \subset H$ , and that is exactly the conclusion proved below.

<sup>130</sup>Editor’s note: we have added this lemma, drawn from the proof of Theorem 5.6 of [DG70], Section III.3. By an abuse of notation, the same letter  $E$  denotes a  $k$ -vector space and the  $k$ -module scheme  $W(E) = \text{Spec } S(E^*)$ .

the vector subspace of  $V$  spanned by the vectors  $gv$ , for  $g \in G(k)$ . Then  $E$  is the smallest  $G$ -submodule of  $V$  containing  $v$ . Consequently, the morphism

$$G \times E \longrightarrow V, \quad (g, x) \longmapsto gx,$$

factors through  $E$ .

*Proof.* By 11.8, there is a smallest  $G$ -submodule  $U$  of  $V$  containing  $v$ . Indeed, if

$$\mu : V \longrightarrow V \otimes \mathcal{A}(G)$$

denotes the comodule structure and

$$\mu(v) = \sum_{i=1}^n v_i \otimes a_i$$

with the  $a_i$  linearly independent, then

$$U = \text{Span}_k(v_1, \dots, v_n).$$

It is clear that  $U$  contains  $E$ , and that the morphism  $G \times U \rightarrow V$  factors through  $U$ .

Conversely, the inverse image of  $E$  under the morphism

$$\mu_v : G \longrightarrow V, \quad g \longmapsto gv,$$

is a closed subset of  $G$  containing all the rational points. Those points are dense in  $G$ , since  $G$  is of finite type over  $k$  (cf. EGA IV<sub>3</sub>, 10.4.8). Hence

$$\mu_v^{-1}(E) = G.$$

Since  $G$  is reduced,  $\mu_v$  therefore factors through  $E$ , whence  $E = U$ .

**Theorem 11.17 (Chevalley).**— *Let  $k$  be a field, let  $G$  be an affine  $k$ -group (not necessarily of finite type), and let  $N$  be a closed group subscheme of  $G$ , invariant in  $G$ . Then the (fpqc) sheaf quotient  $G/N$  is representable by an affine  $k$ -group.*<sup>131</sup>

*Proof.* Suppose first that  $G$  is of finite type. By VI<sub>A</sub>, 3.2 and 5.2, the (fpqc) sheaf quotient  $G/N$  is representable by a  $k$ -group  $Q$ ; it remains to prove that  $Q$  is affine. We proceed in several steps, first supposing that  $k$  is algebraically closed.<sup>132</sup>

(a) Suppose, in addition, that  $G$  is reduced and connected, and that  $N$  is reduced. By 11.16, there are a finite-dimensional  $G$ -module  $V$  over  $k$  and a line  $D = ke_0$  such that

$$\text{Norm}_G(D) = N.$$

In particular,  $N$  acts on  $D$  through a character

$$\chi : N \longrightarrow \mathbb{G}_{m,k}.$$

Fix  $h \in N(k)$ . For every  $g \in G(k)$ , one has

$$hge_0 = g(g^{-1}hg)e_0 = \chi(g^{-1}hg)ge_0.$$

<sup>131</sup>Editor's note: for another proof of this theorem that does not use the results of Exposé VI<sub>A</sub>, see [Ta72], Theorem 5.2 (see also Remark 11.18.5).

<sup>132</sup>Editor's note: in what follows, we have expanded the original and corrected the erroneous assertion  $(G/N)_{\text{red}} = G_{\text{red}}/N_{\text{red}}$ , drawing on [DG70], Section III.3, 5.6.

Thus  $\chi(g^{-1}hg)$  is an eigenvalue of  $h$ . The continuous map

$$\varphi : G(k) \longrightarrow k, \quad g \longmapsto \chi(g^{-1}hg),$$

therefore takes only finitely many values. Since  $G(k)$  is irreducible (it is dense in  $G$ ), it follows that

$$\varphi(g) = \varphi(e) = \chi(h) \quad \text{for every } g \in G(k).$$

Consequently,

$$\chi(g^{-1}hg) = \chi(h) \quad \text{for all } g \in G(k), h \in N(k).$$

Let  $E$  be the vector subspace of  $V$  spanned by the vectors  $ge_0$ , for  $g \in G(k)$ . By Lemma 11.17.0, this is the  $G$ -submodule of  $V$  generated by  $e_0$ .

By the preceding identity, the two morphisms

$$N \times E \longrightarrow E, \quad (h, x) \longmapsto hx \quad \text{and} \quad (h, x) \longmapsto \chi(h)x,$$

agree on

$$(N \times E)(k) = N(k) \times E(k),$$

which is dense in  $N \times E$ . Since  $N \times E$  is reduced (and  $E$  is separated), the two morphisms are equal. Thus  $N$  acts on  $E$  by scalar multiplications.

It follows that  $N$  is contained in the kernel  $K$  of the morphism

$$\rho : G \longrightarrow \mathrm{GL}(\mathrm{End}_k(E))$$

defined, for every  $k$ -algebra  $R$ , by

$$\rho(g)(u) = gug^{-1} \quad (g \in G(R), u \in \mathrm{End}_R(E \otimes_k R)).$$

Conversely, if  $g \in K(R)$ , then

$$g(Re_0) = Re_0,$$

and hence  $g \in N(R)$ . Thus  $N = K$ . By VI<sub>A</sub>, 5.4.1, the morphism

$$G/N \longrightarrow \mathrm{GL}(\mathrm{End}_k(E))$$

is a closed immersion. Therefore  $G/N$  is affine.

(b) Suppose now that  $G$  and  $N$  are reduced, without assuming that  $G$  is connected. Put

$$N' = N \cap G^0.$$

Then  $G^0/N'$  is affine by (a). Moreover,  $NG^0$  is an invariant subgroup of  $G$ , and  $G/NG^0$ , being a quotient of the finite constant group  $G/G^0$  (cf. VI<sub>A</sub>, 5.5.1), is itself a finite constant group. The scheme  $G/N$  is therefore the disjoint union of the fibers of

$$G/N \longrightarrow G/NG^0.$$

By VI<sub>A</sub>, 5.3.3, all these fibers are isomorphic to  $NG^0/N$ , hence to  $G^0/N'$ . Thus  $G/N$  is affine.

(c) Suppose that  $G$  is reduced, with  $N$  arbitrary. The morphism

$$G \times N \longrightarrow N, \quad (g, h) \longmapsto ghg^{-1},$$

induces a morphism

$$(G \times N)_{\mathrm{red}} \longrightarrow N_{\mathrm{red}}.$$

Since  $G$  is reduced and  $k$  is algebraically closed,

$$(G \times N)_{\text{red}} = G \times N_{\text{red}},$$

so  $N_{\text{red}}$  is an invariant subgroup of  $G$ . (This fails when  $G$  is not reduced; see VI<sub>A</sub>, 0.2.)

Hence, by (b),\*

$$G' = G/N_{\text{red}}$$

is affine. By VI<sub>A</sub>, 5.6.1,

$$N' = N/N_{\text{red}}$$

is a finite  $k$ -group. Hence, by Theorem 4.1 of Exposé V,

$$G/N = G'/N'$$

is affine.

(d) For arbitrary  $G$  and  $N$ , base change of the equivalence relation

$$G \times N \rightrightarrows G$$

along  $G_{\text{red}} \rightarrow G$  gives

$$G_{\text{red}} \times N' \rightrightarrows G_{\text{red}}, \quad N' = N \cap G_{\text{red}}.$$

Since the underlying spaces are the same (and the quotient is the quotient ringed space), the morphism

$$G_{\text{red}}/N' \longrightarrow G/N$$

is a homeomorphism. The quotient  $G_{\text{red}}/N'$  is reduced, because

$$p : G_{\text{red}} \longrightarrow G_{\text{red}}/N'$$

is faithfully flat. It follows that

$$(G/N)_{\text{red}} \simeq G_{\text{red}}/N'.$$

The scheme on the right is affine by (c). Since  $G/N$  is of finite type over  $k$  (cf. VI<sub>A</sub>, 3.3.2), EGA I, 5.1.10 implies that  $G/N$  is affine.

Finally, let  $k$  be arbitrary and let  $\bar{k}$  be an algebraic closure. By 9.2(v), one has

$$(G \otimes_k \bar{k})/(N \otimes_k \bar{k}) \simeq (G/N) \otimes_k \bar{k}.$$

The first scheme is affine, hence so is the second. Therefore  $G/N$  is affine by (fpqc) descent (cf. EGA IV<sub>2</sub>, 2.7.1). This proves Theorem 11.17 when  $G$  is of finite type. To pass to the general case, we need the following lemma.<sup>133</sup>

**Lemma 11.17.1.**— *Let  $(C_i \rightarrow A_i)_{i \in I}$  be a filtered direct system of ring homomorphisms, all faithfully flat. Then*

$$A = \varinjlim_i A_i$$

*is faithfully flat over*

$$C = \varinjlim_i C_i.$$

\* Source-reading note: the French re-edition prints “by (a)” here, and the Reinhold comparison lineage repeats that reference. At this stage  $G$  need not be connected; the applicable preceding case is (b).

<sup>133</sup>Editor’s note: we have added this lemma, taken from [DG70], Section III.3, 7.1.

*Proof.* By [BAC], Section I.3, Proposition 9, a ring homomorphism  $B \rightarrow B'$  is faithfully flat if and only if it is injective and  $B'/B$  is a flat  $B$ -module. Since each  $C_i \rightarrow A_i$  is faithfully flat, there are exact sequences

$$0 \longrightarrow C_i \longrightarrow A_i \longrightarrow A_i/C_i \longrightarrow 0,$$

and  $A_i/C_i$  is a flat  $C_i$ -module. Thus

$$(A_i/C_i) \otimes_{C_i} C$$

is a flat  $C$ -module. Since direct limits are exact and commute with tensor products, we obtain an exact sequence

$$0 \longrightarrow C \longrightarrow A \longrightarrow A/C \longrightarrow 0$$

and an isomorphism

$$\varinjlim_i ((A_i/C_i) \otimes_{C_i} C) = (A/C) \otimes_C C = A/C.$$

It follows that  $A/C$  is a flat  $C$ -module. Hence  $A$  is faithfully flat over  $C$ .

We now return to the proof of Theorem 11.17 in the general case, where  $G$  is not assumed to be of finite type. Put

$$\mathcal{A}(G) = A, \quad \mathcal{A}(N) = B = A/J.$$

By 11.13,  $A$  is the direct limit of an increasing filtered family  $(A_i)_{i \in I}$  of finitely generated sub-Hopf algebras. Thus

$$G = \varprojlim_i G_i, \quad G_i = \operatorname{Spec}(A_i),$$

where the  $G_i$  are affine algebraic  $k$ -groups. Let  $\Delta$  and  $\tau$  denote, respectively, the comultiplication and antipode of  $A$ , and put

$$\Delta^2 = (\Delta \otimes \operatorname{id}_A) \circ \Delta.$$

For each  $i$ ,

$$B_i = A_i/(J \cap A_i)$$

is a quotient Hopf algebra of  $A_i$ , so

$$N_i = \operatorname{Spec}(B_i)$$

is a closed subgroup of  $G_i$ . Since  $N$  is invariant in  $G$ , the morphism

$$G \times N \longrightarrow G, \quad (g, n) \longmapsto gng^{-1},$$

factors through  $N$ . Equivalently, the pair  $(A, J)$  satisfies

$$(m_{13} \circ (\Delta \otimes \tau) \circ \Delta)(J) \subset A \otimes_k J,$$

where

$$m_{13}(a_1 \otimes a_2 \otimes a_3) = a_1 a_3 \otimes a_2.$$

It follows that  $(A_i, A_i \cap J)$  satisfies the analogous property, so  $N_i$  is invariant in  $G_i$ . Moreover,

$$\varinjlim_i B_i = B, \quad \varprojlim_i N_i = N.$$



By the finite-type case already proved, each (fpqc) sheaf quotient  $G_i/N_i$  is represented by an affine  $k$ -group

$$Q_i = \operatorname{Spec}(C_i).$$

Put

$$C = \varinjlim_i C_i.$$

The  $Q_i$  form a filtered projective system of affine  $k$ -groups, and its projective limit

$$Q = \varprojlim_i Q_i$$

is the  $k$ -group  $\operatorname{Spec}(C)$  (cf. EGA IV<sub>3</sub>, 8.2.3). We have an exact sequence of  $k$ -groups

$$1 \longrightarrow N \longrightarrow G \longrightarrow Q.$$

We claim that  $Q$  represents the (fpqc) sheaf quotient of  $G$  by  $N$ . It is enough to verify that

$$G \longrightarrow Q$$

is a covering for the (fpqc) topology (cf. IV, 3.3.2.1 and 5.1.7.1). Every morphism

$$G_i \longrightarrow Q_i$$

is faithfully flat (cf. 9.2(xi)); equivalently,  $A_i$  is faithfully flat over  $C_i$ . Since

$$A = \varinjlim_i A_i, \quad C = \varinjlim_i C_i,$$

Lemma 11.17.1 shows that  $A$  is faithfully flat over  $C$ . Thus  $G \rightarrow Q$  is faithfully flat. It is also affine, hence quasi-compact, and is therefore an (fpqc) covering. This completes the proof of Theorem 11.17.

**11.18. Complements.**—<sup>134</sup> Theorem 11.17 and its proof also give the following results, taken from [DG70], Section III.3.7. Let  $k$  be a field. We begin with a lemma (cf. [An73], 2.3.3.2) that will be used later (see 12.10).

**Lemma 11.18.1.**— *Let  $u : G \rightarrow G'$  be a morphism of  $k$ -groups and put  $N = \operatorname{Ker}(u)$ . Suppose that  $G'$  is affine and that  $G$  is of finite type.*

- (i) *The morphism  $\tau : G/N \rightarrow G'$  is a closed immersion. In particular,  $G/N$  is affine.*
- (ii) *If, moreover, the morphism  $u^\sharp : \mathcal{O}(G') \rightarrow \mathcal{O}(G)$  is injective, then  $\tau$  is an isomorphism. Thus  $G'$  is of finite type and  $u$  is faithfully flat.*

Indeed, by 11.13,  $G'$  is the projective limit of a filtered system of affine algebraic  $k$ -groups  $G'_i$ . Let

$$u_i : G \longrightarrow G' \longrightarrow G'_i$$

be the composite, and let  $N_i$  be its kernel. The  $N_i$  form a filtered decreasing system of closed subgroups of  $G$ , whose intersection is  $N$ . Since  $G$  is Noetherian, there is an index  $i$  such that

$$N = N_i.$$

<sup>134</sup>Editor's note: we have added this subsection.

The groups  $G$  and  $G'_i$  are of finite type. Therefore, by VI<sub>A</sub>, 3.2 and 5.4.1, the quotient  $G/N$  is a  $k$ -group of finite type, and  $u_i$  is the composite of the faithfully flat projection

$$p : G \longrightarrow G/N$$

and a closed immersion

$$\tau_i : G/N \hookrightarrow G'_i.$$

Consider the commutative diagram

$$\begin{array}{ccc} G & \xrightarrow{u} & G' \\ p \downarrow & \nearrow \tau & \downarrow q_i \\ G/N & \xrightarrow{\tau_i} & G'_i \end{array}$$

Since

$$q_i \circ \tau = \tau_i$$

is a closed immersion and  $q_i$  is separated ( $G'$  is separated; cf. VI<sub>A</sub>, 0.3),  $\tau$  is a closed immersion (cf. EGA I, 5.4.4). Hence  $G/N$  is affine and

$$\tau^\sharp : \mathcal{O}(G') \longrightarrow \mathcal{O}(G/N)$$

is surjective, proving (i).

If  $u^\sharp : \mathcal{O}(G') \rightarrow \mathcal{O}(G)$  is also injective, then so is  $\tau^\sharp$ . It follows that  $\tau^\sharp$  is an isomorphism, and hence so is  $\tau$ , since  $G'$  and  $G/N$  are affine. This proves (ii).

**Theorem 11.18.2.**— *Let  $G$  and  $G'$  be affine  $k$ -groups, with algebras  $A$  and  $A'$ , and let  $u : G \rightarrow G'$  be a morphism of  $k$ -groups. Put  $N = \text{Ker}(u)$ , and let  $\varphi : A' \rightarrow A$  be the morphism induced by  $u$ .*

- (i) *If  $\varphi$  is injective, then  $u$  is faithfully flat and identifies  $G'$  with  $G/N$ .*
- (ii) *One has  $G/N = \text{Spec}(B)$ , where  $B = \varphi(A')$ . Thus  $u$  is the composite of the faithfully flat morphism  $G \rightarrow G/N$ , corresponding to the inclusion  $B \hookrightarrow A$ , and the closed immersion  $G/N \hookrightarrow G'$ , corresponding to the surjection  $A' \rightarrow B$ . Moreover,  $N$  is defined in  $G$  by the ideal  $AB_+$ , where  $B_+$  denotes the augmentation ideal of  $B$ .*
- (iii) *In particular, if  $u$  is a monomorphism, it is a closed immersion.*

*Proof.* (i) Suppose that  $\varphi$  is injective and identify  $A'$  with a sub-Hopf algebra of  $A$ . By 11.13,  $A$  is the filtered union of finitely generated sub-Hopf algebras

$$A_i = \mathcal{O}(G_i).$$

Put

$$A'_i = A' \cap A_i, \quad G'_i = \text{Spec}(A'_i),$$

and let  $N_i$  be the kernel of the morphism

$$G_i \longrightarrow G'_i$$

induced by  $A'_i \hookrightarrow A_i$ . By Lemma 11.18.1,

$$G_i/N_i \simeq G'_i.$$

Thus, for each  $i$ , there is an exact sequence

$$1 \longrightarrow N_i \longrightarrow G_i \xrightarrow{p_i} G'_i \longrightarrow 1,$$

where  $p_i$  is faithfully flat. Since

$$G = \varprojlim_i G_i, \quad G' = \varprojlim_i G'_i,$$

we obtain an exact sequence <sup>†</sup>

$$1 \longrightarrow N \longrightarrow G \xrightarrow{p} G',$$

where  $N = \varprojlim_i N_i$ . By Lemma 11.17.1,  $p$  is faithfully flat (and affine). Therefore  $G'$  represents the (fpqc) sheaf quotient of  $G$  by  $N$ , proving (i).

In general,  $B = \varphi(A')$  is a sub-Hopf algebra of  $A$ . Let

$$H = \operatorname{Spec}(B),$$

and let  $N'$  be the kernel of the morphism  $G \rightarrow H$  induced by  $B \hookrightarrow A$ . By (i),  $H$  is identified with  $G/N'$ , and  $u$  is the composite of the projection  $G \rightarrow G/N'$  and the closed immersion

$$G/N' \hookrightarrow G'$$

induced by the surjection  $A' \rightarrow B$ . Hence  $N' = N$ . Moreover, by 9.2(ii), the square

$$\begin{array}{ccc} N & \longrightarrow & G \\ \downarrow & & \downarrow p \\ \operatorname{Spec}(k) & \xrightarrow{\varepsilon} & G' \end{array}$$

is cartesian. Here  $\varepsilon$  is the unit section of  $G'$ , corresponding to the augmentation  $B \rightarrow k$ . It follows that  $N$  is defined in  $G$  by the ideal  $AB_+$ . This proves (ii), and (iii) follows.

**Remark 11.18.3.**— Let  $G$  be an affine  $k$ -group and  $N$  an invariant  $k$ -subgroup. Since

$$p : G \longrightarrow G/N$$

is faithfully flat and quasi-compact,  $\mathcal{O}(G/N)$  is, by IV, 3.3.3.2, the subalgebra of  $\mathcal{O}(G)$  formed by the functions  $\phi$  that are right  $N$ -invariant; that is,

$$\phi(gh) = \phi(g)$$

for every  $k$ -scheme  $S$ , every  $g \in G(S)$ , and every  $h \in N(S)$ . Let  $J$  be the ideal of

$$A = \mathcal{O}(G)$$

defining  $N$ . The preceding condition is equivalent to

$$\Delta(\phi) - \phi \otimes 1 \in \mathcal{O}(G) \otimes J,$$

where  $\Delta$  is the comultiplication of  $A$ .

The preceding theorem may therefore be reformulated in terms of Hopf algebras as follows.

**Corollary 11.18.4.**— *Let  $k$  be a field, let  $A$  be a commutative  $k$ -Hopf algebra, and put  $G = \operatorname{Spec}(A)$ .*

<sup>†</sup> Source-reading note: the French re-edition prints  $G/N$  as the final term of this limiting exact sequence, and the Reinhold comparison lineage repeats it. At this point the argument is proving that  $G'$  represents  $G/N$ ; taking the projective limits of the preceding sequences gives  $G'$  as the final term.

- (i) If  $B$  is a sub-Hopf algebra of  $A$ , then  $A$  is faithfully flat over  $B$ .
- (ii) The map

$$N \longmapsto \mathcal{O}(G/N)$$

is a bijection from the set of invariant subgroups of  $G$  to the set of sub-Hopf algebras of  $A$ . Its inverse is

$$B \longmapsto \operatorname{Spec}(A/AB_+).$$

Moreover, if  $J$  is the ideal of  $A$  defining  $N$ , then

$$\mathcal{O}(G/N) = \{x \in A \mid \Delta(x) - x \otimes 1 \in A \otimes J\}.$$

**Remarks 11.18.5.**—

- (a) One consequence of the preceding theorem is that the category of commutative affine  $k$ -groups is abelian. For this, as well as for other results on affine  $k$ -groups, we refer to [DG70], § III.3, 7.4–7.8.
- (b) Finally, let us point out that M. Takeuchi gave another proof of the results of 11.17–11.18.4; see [Ta72], § 5. He moreover strengthened 11.18.4(i) above by showing that  $A$  is even a projective  $B$ -module; see [Ta79], Theorem 5 (see also [MW94], Theorem 3.6).

## 12. Complements on $G_{\text{af}}$ and the “anti-affine” groups

<sup>135</sup> We begin with the following lemma, which extends 11.18.1 to the case where  $G$  is not assumed to be of finite type.<sup>136</sup>

**Lemma 12.1.**— *Let  $k$  be a field and let  $u : G \rightarrow H$  be a monomorphism of  $k$ -groups, with  $H$  affine. Suppose that  $u$  is quasi-compact. Then  $u$  is a closed immersion.*

By (fpqc) descent, we may suppose that  $k$  is algebraically closed. By VI<sub>A</sub>, 6.4, the closed image  $I$  of  $u$  is a closed group subscheme of  $H$ , hence is still affine. Replacing  $H$  by  $I$ , we may therefore suppose that  $u$  is schematically dominant. Since  $k$  is algebraically closed,  $H' = H_{\text{red}}$  is a group subscheme of  $H$ . Put  $G' = G \times_H H'$ . The morphism

$$u' : G' \longrightarrow H'$$

deduced from  $u$  by base change is a quasi-compact monomorphism and is dominant (the underlying continuous map being the same for  $u$  and  $u'$ ). Hence, by VI<sub>A</sub>, 6.2,  $u'$  is faithfully flat. It is therefore a quasi-compact faithfully flat monomorphism, and thus an isomorphism (cf. IV, 1.14).

Consequently  $u : G \rightarrow H$  is a homeomorphism, and hence is affine by 2.9.1. Thus  $G$  is affine, and  $u$  is a closed immersion by 11.18.2(iii).

**Theorem 12.2.**— *Let  $G$  be an algebraic  $k$ -group. Denote by  $\rho : G \rightarrow G_{\text{af}}$  the canonical morphism and by  $N$  its kernel.*

- (i) *The canonical morphism  $G/N \rightarrow G_{\text{af}}$  is an isomorphism. Consequently  $G_{\text{af}}$  is an affine algebraic group and  $\rho$  is faithfully flat.*
- (ii) *There is a canonical isomorphism  $(G/N)_{\text{af}} = G_{\text{af}}$ .*

<sup>135</sup>Editor’s note: this section has been added.

<sup>136</sup>Editor’s note: this lemma was communicated to us by M. Raynaud; it will be used in the proof of Proposition 12.9.

- (iii)  $N$  is a characteristic subgroup of  $G$ .
- (iv)  $\mathcal{O}(N) = k$ .
- (v)  $N$  is smooth, connected, and commutative.

Assertion (i) is a special case of 11.18.1, and (ii) follows from the universal properties of  $G_{\text{af}}$  and  $(G/N)_{\text{af}}$ .

Let us prove (iii). For an arbitrary  $k$ -scheme  $\pi : S \rightarrow \text{Spec } k$ , consider the cartesian square

$$\begin{array}{ccc} G_S & \longrightarrow & G \\ q \downarrow & & \downarrow p \\ S & \xrightarrow{\pi} & \text{Spec } k \end{array}$$

Since  $p$  is quasi-compact and separated and  $\pi$  is flat, EGA III, 1.4.15 and EGA IV<sub>1</sub>, 1.7.21 give

$$q_*(\mathcal{O}_{G_S}) = \pi^*(\mathcal{O}(G)) = \pi^*(\mathcal{O}(G_{\text{af}})).$$

Hence EGA II, 1.5.2 gives

$$(G_S)_{\text{af}} = (G_{\text{af}})_S.$$

Thus  $N_S$ , being the kernel of the canonical morphism  $G_S \rightarrow (G_S)_{\text{af}}$ , is invariant under every automorphism of  $G_S$ ; in other words,  $N$  is a characteristic subgroup of  $G$ .

To prove (iv), put

$$N' = \text{Ker}(N \rightarrow N_{\text{af}}).$$

By (ii), this is an invariant subgroup of  $G$ . Since  $N$  is algebraic (being a closed subgroup of  $G$ ), (i) gives  $N/N' \simeq N_{\text{af}}$ . Moreover, by VI<sub>A</sub>, 3.2 and 5.3.2, there is an isomorphism of  $k$ -groups

$$(G/N')/N_{\text{af}} \simeq (G/N')/(N/N') \simeq G/N.$$

Since  $N_{\text{af}}$  is affine, the projection  $G/N' \rightarrow G/N$  is affine by 9.2(vii); and since  $G/N = G_{\text{af}}$  is affine,  $G/N'$  is affine as well. The universal property of  $G_{\text{af}}$  therefore makes the projection

$$p' : G \rightarrow G/N'$$

factor through  $G_{\text{af}} = G/N$ . Hence  $N \subset N'$ , and consequently  $N = N'$ . Thus  $N_{\text{af}}$  is the trivial group, and  $\mathcal{O}(N) = k$ .

Finally, (v) follows from the next lemma.

**Lemma 12.3.**— *Let  $k$  be a field and let  $N$  be an algebraic  $k$ -group such that  $\mathcal{O}(N) = k$ . Then  $N$  is smooth, connected, and commutative.*

Indeed, we may suppose that  $k$  is algebraically closed. Then

$$H = N_{\text{red}}^0$$

is a  $k$ -subgroup of  $N$ , and the quotient  $k$ -scheme  $X = N/H$  is finite (hence affine) over  $k$ , by VI<sub>A</sub>, 5.5.1 and 5.6.1. Since  $p : N \rightarrow X$  is faithfully flat,

$$\mathcal{O}(X) \subset \mathcal{O}(N) = k.$$

It follows that  $N = N_{\text{red}}^0$ , and hence that  $N$  is smooth (VI<sub>A</sub>, 1.3.1) and connected.

Let  $Z$  be the center of  $N$ . By 6.2.6,  $N/Z$  is affine, and the preceding argument gives  $\mathcal{O}(N/Z) = k$ . Hence  $N = Z$ . This proves the lemma and completes the proof of 12.2.

We also record, without proof, the following theorem. Recall that an *abelian variety* over a field  $k$  is a proper, smooth, connected  $k$ -group scheme.

**Theorem 12.4 (Chevalley).**— *Let  $k$  be a perfect field and let  $G$  be a smooth, connected algebraic  $k$ -group. Then there exists a smooth, connected affine  $k$ -subgroup  $L$ , invariant in  $G$ , such that  $G/L$  is an abelian variety. Moreover,  $L$  is unique and its formation commutes with extension of the base field.*

**Remarks 12.5.**—

- (1) This theorem was announced in 1953 by C. Chevalley, who published his proof in 1960 ([Ch60]). Meanwhile, other proofs were obtained independently by I. Barsotti and M. Rosenlicht ([Ba55, Ro56]); see [Se99] for historical comments.
- (2) A modern version (that is, in the language of schemes) of Chevalley’s proof was given by B. Conrad ([Co02]). (In *loc. cit.*, “algebraic group” means a smooth connected  $k$ -group scheme.)
- (3) A modern version of Rosenlicht’s proof, on the other hand, was given by Ngô B.-C. in a course at Orsay in 2005–2006.
- (4) If the hypothesis that  $k$  be perfect is omitted, there still exists a smallest invariant connected affine subgroup  $L$ , not necessarily smooth, such that  $G/L$  is an abelian variety ([BLR], § 9.2, Theorem 1).
- (5) One may also omit the hypothesis that  $G$  be smooth over  $k$ . Indeed, by VII<sub>A</sub>, 8.3, there is an integer  $n \geq 1$  such that the quotient

$$G' = G/(\mathrm{Fr}^n G)$$

is smooth. Then  $G'$  contains a subgroup  $L'$  as in (4) above, and the inverse image of  $L'$  in  $G$  again has the same properties. Thus, for every connected algebraic group  $G$  over a field  $k$ , there is an exact sequence

$$1 \longrightarrow H \longrightarrow G \longrightarrow A \longrightarrow 1,$$

where  $H$  is an affine  $k$ -group and  $A$  is an abelian  $k$ -variety. Moreover, by [Per76], Corollary 4.2.9, such an exact sequence exists for every connected  $k$ -group  $G$ , not necessarily algebraic.

- (6) Let  $k$  be an algebraically closed field and let  $G$  be the semidirect product of an elliptic curve  $E$  by the constant  $k$ -group  $\{\pm 1\}_k$ , for the action defined by  $(-1) \cdot x = -x$ . In this case, if  $L$  is a closed invariant subgroup of  $G$  such that  $G/L$  is connected, then  $L = G$ .

**Remark 12.6.**— Following [Br09], a  $k$ -group  $N$  will be called *anti-affine* if  $\mathcal{O}(N) = k$ . By 12.3 and 12.4, if  $k$  is perfect, every anti-affine algebraic  $k$ -group is an extension of an abelian variety by a smooth, connected, commutative affine algebraic  $k$ -group. For the precise structure of anti-affine algebraic groups over a perfect field, and various consequences, see the recent articles of M. Brion and C. & F. Sancho de Salas ([Br09, SS09]).

To conclude this section, we shall prove two results due to M. Raynaud, the first being Remark 11.11.1, and the second Proposition 2.1 of Exposé XVII, Appendix III. We shall need the following lemma,<sup>137</sup> which improves, for a complete discrete valuation ring  $R$ , the flatness criteria given in [BAC], § III.5.

<sup>137</sup>Editor’s note: this is a version improved by O. Gabber of a statement communicated by M. Raynaud.

**Lemma 12.7.**— *Let  $R$  be a discrete valuation ring,  $K$  its fraction field, and  $\pi$  a uniformizer. Let  $A$  be a flat  $R$ -algebra and let  $M$  be an  $A$ -module flat over  $R$ . Suppose that*

- (i)  $M/\pi M$  is flat over  $\overline{A} = A/\pi A$ ,
- (ii)  $M \otimes_R K$  is flat over  $A \otimes_R K$ .

*Then  $M$  is flat over  $A$ .*

By the flatness criterion in the nilpotent case (cf. [BAC], III § 5.2, Theorem 1),

$$M/\pi^n M$$

is flat over  $A/\pi^n A$  for every  $n \in \mathbb{N}^*$ . It then follows from [RG71], II, Lemma 1.4.2.1, that  $M$  is flat over  $A$ . For the reader's convenience, let us briefly indicate the proof. Put

$$s = \pi^n, \quad P = M/sM.$$

Since  $P$  is flat over  $A/sA$ , and  $A/sA$  has projective dimension 1 over  $A$ , the spectral sequence of composite functors shows that  $P$  has Tor-dimension at most 1 over  $A$ . Since  $M$  is flat over  $R$ , and thus has no  $\pi$ -torsion,  $M_K/M$  is the direct limit of the  $A$ -modules

$$\pi^{-n}M/M \simeq M/\pi^n M.$$

Consequently  $M_K/M$  also has Tor-dimension at most 1. From the exact sequence

$$0 \longrightarrow M \longrightarrow M_K \longrightarrow M_K/M \longrightarrow 0$$

and the hypothesis that  $M_K$  is flat over  $A_K$ , hence over  $A$ , it follows that  $M$  is flat.

For completeness, let us also give the following simpler proof, pointed out by O. Gabber. Let  $I$  be a finitely generated ideal of  $A$ . We must show that

$$u : M \otimes_A I \longrightarrow M$$

is injective. By (ii),  $u \otimes_R K$  is injective, so  $\text{Ker}(u)$  is an  $R$ -module of  $\pi$ -torsion. It therefore suffices to show that the  $\pi$ -torsion submodule of  $M \otimes_A I$  is zero. This submodule is a quotient of  $\text{Tor}_1^A(M, I/\pi I)$ , as is seen by tensoring with  $M$  the exact sequence

$$0 \longrightarrow I \xrightarrow{\pi} I \longrightarrow I/\pi I \longrightarrow 0.$$

On the other hand, since  $M$  has no  $\pi$ -torsion (being flat over  $R$ ), one has

$$\text{Tor}_i^A(M, \overline{A}) = 0 \quad (i \geq 1).$$

Consequently, if  $P_\bullet$  is a projective resolution of the  $A$ -module  $M$ , then

$$P_\bullet \otimes_A \overline{A}$$

is a projective resolution of the  $\overline{A}$ -module  $\overline{M} = M/\pi M$ . Thus, for every  $\overline{A}$ -module  $Q$ ,

$$\text{Tor}_i^A(M, Q) = \text{Tor}_i^{\overline{A}}(\overline{M}, Q),$$

which vanishes for  $i \geq 1$ , since  $\overline{M}$  is flat over  $\overline{A}$ . Hence  $\text{Tor}_1^A(M, I/\pi I) = 0$ , proving the lemma.

**Remark 12.8.**— Let  $S$  be a regular locally noetherian scheme of dimension 1, and let  $X$  be an  $S$ -scheme that is flat, quasi-separated, and quasi-compact over  $S$ . Then  $\mathcal{A}(X)$  is a

flat  $\mathcal{O}_S$ -module. Indeed, we may suppose that  $S$  is local. Let  $s$  be its closed point, let  $i : X_s \hookrightarrow X$  be the inclusion, and let  $\pi$  be a uniformizer of  $R = \mathcal{O}(S)$ . Since  $X$  is flat over  $S$ , there is an exact sequence of sheaves

$$0 \longrightarrow \mathcal{O}_X \xrightarrow{\pi} \mathcal{O}_X \longrightarrow i_*(\mathcal{O}_{X_s}) \longrightarrow 0.$$

Taking global sections shows that  $\mathcal{A}(X)$  has no  $\pi$ -torsion as an  $R$ -module, and hence is flat.<sup>138</sup> Moreover, the morphism

$$\mathcal{A}((X_{\text{af}})_s) = \mathcal{A}(X)/\pi\mathcal{A}(X) \longrightarrow \mathcal{A}(X_s),$$

induced by  $X \rightarrow X_{\text{af}}$ , is injective.

We can now prove the following proposition (cf. Remark 11.11.1).

**Proposition 12.9.**— *Let  $S$  be a regular locally noetherian scheme of dimension at most 1, and let  $G$  be an  $S$ -group scheme that is flat, quasi-separated, and quasi-compact over  $S$ . Then the following conditions are equivalent:*

- (i)  $G$  is affine over  $S$ .
- (ii)  $G$  is quasi-affine over  $S$ .
- (iii)  $G$  acts faithfully on a quasi-affine and flat  $S$ -scheme  $X$ .
- (iv)  $G$  acts linearly and faithfully on a quasi-coherent module  $\mathcal{E}$  that is flat over  $S$ .
- (v) The morphism  $\rho : G \rightarrow G_{\text{af}}$  is a monomorphism.

The implication (i)  $\Rightarrow$  (ii) is clear, as is (ii)  $\Rightarrow$  (iii) (take  $X = G$ ).

Suppose (iii) holds. Since  $\mathcal{A}(G)$  and  $\mathcal{A}(X)$  are flat  $\mathcal{O}_S$ -modules, the argument of 11.11 shows that  $G$  acts faithfully on the right on  $X_{\text{af}}$ , and hence acts linearly and faithfully on the left on the flat quasi-coherent  $\mathcal{O}_S$ -module  $\mathcal{A}(X)$ .

If (iv) holds, 11.6.1(ii) shows that the monomorphism

$$G \longrightarrow \text{Aut}_{\mathcal{O}_S}(\mathcal{E})$$

factors through  $G_{\text{af}}$ . Hence  $G \rightarrow G_{\text{af}}$  is a monomorphism.

Finally, suppose (v) holds, and let us prove that  $\rho : G \rightarrow G_{\text{af}}$  is an isomorphism. Replacing  $S$  by one of its connected components, we may suppose that  $S$  is irreducible, with generic point  $\eta$ . Since the formation of  $G_{\text{af}}$  commutes with flat base change,

$$(G_{\text{af}})_{\eta} = (G_{\eta})_{\text{af}}.$$

Therefore  $G_{\eta} \rightarrow (G_{\eta})_{\text{af}}$  is a monomorphism, hence a closed immersion by 12.1, and consequently an isomorphism because

$$\mathcal{O}((G_{\eta})_{\text{af}}) = \mathcal{O}(G_{\eta}).$$

If  $S = \text{Spec}(\kappa(\eta))$ , the proof is complete; we may therefore suppose that  $\dim S = 1$ .

Let  $s$  be a closed point of  $S$ . We shall show that

$$G_s \longrightarrow (G_{\text{af}})_s$$

<sup>138</sup>Editor's note: this is true more generally when  $S$  is regular, locally noetherian, and of dimension at most 2; see [Ray70a], VII 3.2.



is an isomorphism and that  $\rho$  is flat at every point of  $G_s$ . We may suppose that  $S$  is local, with closed point  $s$ . The base-change morphism

$$\rho_s : G_s \longrightarrow (G_{\text{af}})_s$$

is a monomorphism, hence a closed immersion by 12.1. Thus the morphism

$$\mathcal{O}((G_{\text{af}})_s) \longrightarrow \mathcal{O}(G_s)$$

induced by  $\rho_s$  is surjective. By the preceding remark it is also injective, and hence is an isomorphism. In particular,  $\rho$  is surjective.

It follows from Lemma 12.7 that  $\rho : G \rightarrow G_{\text{af}}$  is faithfully flat. Since  $G$  is quasi-compact over  $S$  and  $G_{\text{af}}$  is separated over  $S$ ,  $\rho$  is also quasi-compact (cf. EGA I, 6.6.4). Therefore  $\rho$  is a quasi-compact faithfully flat monomorphism, and is thus an isomorphism. This proves the proposition.

Finally, we prove Proposition 2.1 of Exposé XVII, Appendix III. Taking [Per76], Corollary 4.2.5 into account, we have replaced “of finite type” in the hypotheses by “quasi-compact and quasi-separated”. (If  $G$  is assumed to be of finite type, 12.1 may be used in place of *loc. cit.*)

**Proposition 12.10.**— *Let  $S$  be a regular locally noetherian scheme of dimension at most 1, and let  $G$  be a flat, quasi-compact, and quasi-separated  $S$ -group scheme.*

- (i) *The canonical morphism  $\rho : G \rightarrow G_{\text{af}}$  is faithfully flat and quasi-compact. Consequently,  $G_{\text{af}}$  represents the (fpqc) sheaf quotient of  $G$  by  $N = \text{Ker}(\rho)$ .*
- (ii) *If moreover  $G$  is of finite type over  $S$ , then  $\rho$  is of finite presentation,  $G_{\text{af}}$  represents the (fppf) sheaf quotient of  $G$  by  $N$ , and  $G_{\text{af}}$  is of finite type over  $S$ .*
- (iii) *Suppose moreover that  $G_\eta$  is affine for every maximal point  $\eta$  of  $S$ . Then  $N$  is an étale  $S$ -group and is the unit group if  $G$  is separated over  $S$ .*

Since  $G_{\text{af}}$  is affine, and therefore separated over  $S$ , the morphism  $\rho$  is quasi-compact (cf. EGA I, 6.6.4), and its kernel

$$N = \text{Ker}(\rho)$$

is a closed subgroup of  $G$ . Replacing  $S$  by one of its connected components, we may suppose that  $S$  is irreducible, with generic point  $\eta$ .

To prove (i) and (ii), it is enough to show that  $\rho$  is faithfully flat. Indeed, Exposé IV, 5.1.7.1 then shows that  $G_{\text{af}}$  represents the (fpqc) sheaf quotient of  $G$  by  $N$ . If  $G$  is moreover of finite type over  $S$ , it is of finite presentation because  $S$  is locally noetherian; by 9.2(xiii),  $\rho$  is then of finite presentation, as is  $G_{\text{af}} \rightarrow S$ , and  $\rho$  is therefore a covering for the (fppf) topology.

We may thus suppose that

$$S = \text{Spec}(R),$$

where  $R$  is a discrete valuation ring if  $\dim S = 1$ , and is the field  $\kappa(\eta)$  if  $\dim S = 0$ . Let  $s$  be the closed point of  $S$ . Since formation of  $G_{\text{af}}$  commutes with flat base change, the canonical morphism  $G_\eta \rightarrow (G_\eta)_{\text{af}}$  is identified with

$$\rho_\eta : G_\eta \longrightarrow (G_{\text{af}})_\eta.$$

Since

$$\mathcal{O}((G_{\text{af}})_\eta) = \mathcal{O}((G_\eta)_{\text{af}}) = \mathcal{O}(G_\eta),$$

$\rho_\eta$  is faithfully flat by [Per76], 4.2.5 (see also the addition VI<sub>A</sub>, 6.6). If  $\dim S = 1$ , the same argument shows that  $\rho_s$  is faithfully flat, because the morphism

$$\mathcal{O}((G_{\text{af}})_s) \longrightarrow \mathcal{O}(G_s)$$

is injective by Remark 12.8. Lemma 12.7 now shows that  $\rho$  is faithfully flat. This proves (i) and (ii). In particular,  $N$  is flat over  $S$ .

Suppose now that  $G_\eta$  is affine. Since  $\rho_\eta$  coincides with the canonical morphism

$$G_\eta \longrightarrow (G_\eta)_{\text{af}},$$

its kernel  $N_\eta$  is the unit group. Let us prove that  $N$  is étale over  $S$ . Since  $N$  is flat over  $S$ , it remains to show that  $N_s$  is étale over  $\kappa(s)$  for every point  $s \neq \eta$  of  $S$ . There is nothing to prove when  $S = \text{Spec}(\kappa(\eta))$ , so we may suppose that  $S = \text{Spec}(R)$ , where  $R$  is a discrete valuation ring. Let  $s$  be the closed point of  $S$ , let  $K$  be the fraction field of  $R$ , and let  $\pi$  be a uniformizer.

Let  $x \in N_s$ , and let  $U_x$  be an affine open neighborhood of  $x$  in  $N$ . Since  $N$  is flat over  $S$ , the intersection  $U_x \cap N_\eta$  is nonempty, and hence equals  $\{\epsilon(\eta)\}$ , where  $\epsilon$  denotes the unit section. Thus

$$A = \mathcal{O}(U_x)$$

is a flat  $R$ -algebra such that  $A_K = K$  and  $\pi^{-1} \notin A$ , since  $\pi$  belongs to the maximal ideal of  $\mathcal{O}_{U_x, x}$ . It follows that  $A = R$ , and hence the projection  $U_x \rightarrow S$  is an isomorphism. Thus  $N$  is étale over  $S$ . If  $G$  is also separated over  $S$ , the inverse isomorphism  $S \rightarrow U_x$  is the unit section, since the two coincide on the dense open subset  $\{\eta\}$  of  $S = \text{Spec}(R)$ . Hence  $N$  is the unit group. This proves the proposition.

In particular, we obtain the following corollary, two other proofs of which may be found in [An73], Proposition 2.3.1, and [PY06], Proposition 3.1.

**Corollary 12.10.1.**— *Let  $R$  be a discrete valuation ring,  $K$  its fraction field, and  $G$  a separated, flat  $R$ -group scheme of finite type over  $R$ . If  $G_K$  is affine, then  $G$  is affine.*

**Remarks 12.10.2.**—

- (a) O. Gabber has pointed out to us examples in which  $G$  is a flat group of finite type over a discrete valuation ring, its generic fiber is an abelian variety, and the kernel  $N$  of  $G \rightarrow G_{\text{af}}$  is not smooth.
- (b) M. Raynaud, on the other hand, gave an example, with  $S$  the affine plane of dimension 2 over a field  $k$ , of a smooth quasi-affine  $S$ -group scheme with affine, connected fibers that is not affine over  $S$ ; see [Ray70a], § VII.3, p. 116.

### 13. Flat affine groups over a regular base of dimension at most 2

Let us begin by observing<sup>139</sup> that the familiar argument showing that every affine algebraic group over a field  $k$  is linear, as well as Lemma 11.12, extends to a group scheme  $G$  that is affine, flat, and of finite type over a Noetherian regular base scheme  $S$  of dimension at most 2. For  $S$  of dimension at most 1, this proves Remark 11.11.1(b). The extension to  $\dim S = 2$  rests on the following lemma, communicated to us by O. Gabber.

**Lemma 13.1.**— *Let  $S$  be a normal Noetherian scheme, let  $\mathcal{A}$  be a flat quasi-coherent  $\mathcal{O}_S$ -module, let  $\mathcal{F}$  be a finite-type  $\mathcal{O}_S$ -submodule of  $\mathcal{A}$ , let  $\mathcal{F}^{**}$  be its bidual, and let  $U$  be the flatness locus of  $\mathcal{F}$ , that is, the set of points  $s \in S$  such that  $\mathcal{F}_s$  is a flat  $\mathcal{O}_{S, s}$ -module.*

<sup>139</sup>Editor's note: this section has been added.

(i)  $U$  is open in  $S$ , and

$$\mathcal{F}^{**} = j_* j^*(\mathcal{F}),$$

where  $j : U \hookrightarrow S$  denotes the inclusion.\*

(ii) For every quasi-coherent  $\mathcal{O}_U$ -module  $\mathcal{E}$ , the canonical morphism

$$j_*(\mathcal{E}) \otimes_{\mathcal{O}_S} \mathcal{A} \longrightarrow j_*(\mathcal{E} \otimes_{\mathcal{O}_U} j^*(\mathcal{A}))$$

is an isomorphism.

(iii) In particular,

$$\mathcal{V} = j_* j^*(\mathcal{F}) \subset \mathcal{A} = j_* j^*(\mathcal{A}),$$

and the canonical morphism

$$\mathcal{V} \otimes_{\mathcal{O}_S} \mathcal{A} \longrightarrow j_* j^*(\mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{A})$$

is an isomorphism.

*Proof.* Replacing  $S$  by one of its connected components, we may suppose that  $S$  is integral. By EGA IV<sub>2</sub>, 2.1.12, the flatness locus of  $\mathcal{F}$ , namely the set of points  $s \in S$  such that  $\mathcal{F}_s$  is a flat  $\mathcal{O}_{S,s}$ -module, is an open subset  $U$  of  $S$ ; write  $j : U \hookrightarrow S$  for its inclusion. Since  $\mathcal{A}_s$  is flat, hence torsion-free, the same is true of  $\mathcal{F}_s$ . Thus  $U$  contains every point of codimension at most 1. Consequently, by [BAC], VII, § 4.2, Corollary to Theorem 1,

$$\mathcal{F}^{**} = j_* j^*(\mathcal{F}),$$

and hence there is a monomorphism

$$\mathcal{F}^{**} \longrightarrow j_* j^*(\mathcal{A}).$$

The proof of (ii) is analogous to that of EGA III, 1.4.15, recalled in 11.0. Since  $S$  is normal, the morphism

$$\mathcal{O}_S \longrightarrow j_* j^*(\mathcal{O}_S)$$

is an isomorphism (cf. EGA IV<sub>2</sub>, 5.8.6 and 5.10.5). Applying (ii) to  $\mathcal{E} = \mathcal{O}_U$  therefore gives

$$\mathcal{A} = j_* j^*(\mathcal{A}).$$

Finally, since

$$j^*(\mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{A}) = j^*(\mathcal{F}) \otimes_{\mathcal{O}_U} j^*(\mathcal{A}),$$

the last assertion of (iii) follows from (ii) applied to  $\mathcal{E} = j^*(\mathcal{F})$ . This proves the lemma.

Recall also that a finitely generated  $R$ -module  $M$  is called *reflexive* if the canonical morphism from  $M$  to its bidual  $M^{**}$  is an isomorphism. If  $R$  is a Noetherian regular ring of dimension at most 2, this implies that  $M$  is projective. Indeed, for every finitely generated  $R$ -module  $N$ , choose a resolution

$$L_1 \longrightarrow L_0 \longrightarrow N \longrightarrow 0,$$

where  $L_0$  and  $L_1$  are finite free  $R$ -modules. There is then an exact sequence

$$0 \longrightarrow N^* \longrightarrow L_0^* \longrightarrow L_1^* \longrightarrow Q \longrightarrow 0,$$

---

\* Source-reading note: the French re-edition prints  $j : U \hookrightarrow X$  in the statement. The ambient scheme throughout the lemma is  $S$ , and its proof explicitly defines  $j : U \hookrightarrow S$ ; the latter is used here. The Reinhold comparison lineage retains the printed  $X$ .

where  $Q$  is the cokernel of  $L_0^* \rightarrow L_1^*$ . Since  $R$  has homological dimension at most 2 ([BAC], X, § 4.2, Corollary 1 to Theorem 1), it follows that  $N^*$  is projective.

**Proposition 13.2.**— *Let  $S$  be a Noetherian regular scheme of dimension at most 2, let  $G$  be an affine flat  $S$ -group, and let  $\mathcal{A}(G)$  be its affine algebra.*

(i) *If  $G$  is of finite type over  $S$ , it is isomorphic to a closed subgroup of*

$$H = \operatorname{Aut}_{\mathcal{O}_S}(\mathcal{V})$$

*for some locally free  $\mathcal{O}_S$ -module  $\mathcal{V}$  of finite rank. If, moreover,  $S$  is affine, one may take  $\mathcal{V} = \mathcal{O}_S^{\oplus d}$  for some  $d$ , so that  $H = \operatorname{GL}_{d,S}$ .*

(ii)  *$\mathcal{A}(G)$  is the filtered direct limit of flat finite-type  $\mathcal{O}_S$ -sub-Hopf algebras.*

*Proof.* Let  $\mathcal{B}$  be a finite-type  $\mathcal{O}_S$ -subalgebra of  $\mathcal{A}(G)$ . Since every coherent module on an open subset of  $S$  extends to a coherent module on  $S$  (EGA I, 9.4.5), there is a coherent  $\mathcal{O}_S$ -submodule  $\mathcal{M}$  of  $\mathcal{B}$  that generates  $\mathcal{B}$  as an  $\mathcal{O}_S$ -algebra (*loc. cit.*, 9.6.5). By 11.10.bis,  $\mathcal{M}$  is contained in a coherent  $G$ -stable  $\mathcal{O}_S$ -submodule  $\mathcal{F}$ . (N.B. Since  $G$  is affine over  $S$ , the proof of *loc. cit.* is simpler here: one may replace  $f_*f^*(\mathcal{E})$  by  $\mathcal{E} \otimes_{\mathcal{O}_S} \mathcal{A}(G)$ , and so on.)

Let  $j : U \hookrightarrow S$  be the inclusion, where  $U$  is the flatness locus of  $\mathcal{F}$ . By Lemma 13.1 and the preceding observation,

$$\mathcal{V} = j_*j^*(\mathcal{F})$$

is a locally free  $\mathcal{O}_S$ -submodule of  $\mathcal{A}(G)$ , and the canonical morphism

$$\mathcal{V} \otimes_{\mathcal{O}_S} \mathcal{A}(G) \longrightarrow j_*j^*(\mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{A}(G))$$

is an isomorphism. Thus  $\mathcal{V}$  is a  $G$ -submodule of  $\mathcal{A}(G)$ . The action of  $G$  on  $\mathcal{V}$  induces a morphism of affine  $S$ -groups

$$\rho_{\mathcal{V}} : G \longrightarrow H = \operatorname{Aut}_{\mathcal{O}_S}(\mathcal{V})$$

and therefore a morphism of  $\mathcal{O}_S$ -Hopf algebras

$$\varphi_{\mathcal{V}} : \mathcal{A}(H) \longrightarrow \mathcal{A}(G).$$

Write  $\mathcal{A}_{\mathcal{V}}$  for the image of  $\varphi_{\mathcal{V}}$ . It is the affine algebra of a closed subgroup  $G_{\mathcal{V}}$  of  $H$ , the closed image of  $\rho_{\mathcal{V}}$ . We shall show that  $\mathcal{A}_{\mathcal{V}}$  contains  $\mathcal{B}$ .

The question is local on  $S$ , so we may suppose that  $S = \operatorname{Spec}(R)$  and that

$$V = \Gamma(S, \mathcal{V})$$

is a free  $R$ -module with basis  $v_1, \dots, v_n$ . Then  $H \simeq \operatorname{GL}_{n,R}$ , and  $\mathcal{A}(H)$  is generated as an  $R$ -algebra by the “matrix coefficients”  $c_{ij}$  and by  $d^{-1}$ , where  $d$  denotes the determinant. Let  $\Delta$  (respectively,  $\varepsilon$ ) denote the comultiplication (respectively, augmentation) of  $A = \Gamma(S, \mathcal{A}(G))$ . For  $j = 1, \dots, n$ , write

$$\Delta(v_j) = \sum_{i=1}^n v_i \otimes a_{ij}.$$

Then  $a_{ij} = \varphi(c_{ij})$  belongs to  $\operatorname{Im}(\varphi)$ . Since  $V$  is an  $R$ -submodule of  $A$ , the identity

$$(\varepsilon \otimes \operatorname{id}_A) \circ \Delta = \operatorname{id}_A$$

gives

$$v_j = \sum_i \varepsilon(v_i) a_{ij} \in \operatorname{Im}(\varphi).$$

As  $V$  contains a set of generators of  $B = \Gamma(S, \mathcal{B})$ , it follows that

$$B \subset \text{Im}(\varphi).$$

If  $G$  is of finite type over  $S$ , take  $\mathcal{B} = \mathcal{A}(G)$ . Then  $\varphi$  is surjective, so the morphism

$$G \longrightarrow H = \text{Aut}_{\mathcal{O}_S}(\mathcal{V})$$

is a closed immersion.

If  $S$  is also affine, there is a locally free  $\mathcal{O}_S$ -module  $\mathcal{V}'$  of finite rank such that

$$\mathcal{V} \oplus \mathcal{V}' = \mathcal{O}_S^{\oplus d}.$$

Regard  $\mathcal{V}'$  as a trivial  $G$ -module and replace  $\mathcal{V}$  by  $\mathcal{O}_S^{\oplus d}$ . This realizes  $G$  as a closed subgroup of  $\text{GL}_{d,S}$ .

Finally, return to an arbitrary affine flat  $S$ -group  $G$ . By EGA I, 9.4.9,  $\mathcal{A}(G)$  is the union of its coherent  $\mathcal{O}_S$ -submodules  $\mathcal{M}$ , hence also of the  $\mathcal{O}_S$ -sub-Hopf algebras  $\mathcal{A}_{\mathcal{V}}$  constructed above. This proves (ii).

**Example 13.3.**— *Let  $R$  be a discrete valuation ring with uniformizer  $\pi$  and fraction field  $K$ . Consider the filtered projective system of  $R$ -groups*

$$\cdots \longrightarrow \mathbb{G}_{a,R} \xrightarrow{\times\pi} \mathbb{G}_{a,R} \xrightarrow{\times\pi} \mathbb{G}_{a,R}$$

*corresponding to the direct system*

$$R[X_0] \longrightarrow R[X_1] \longrightarrow R[X_2] \longrightarrow \cdots,$$

*whose transition morphisms are given by  $X_n = \pi X_{n+1}$ . Its projective limit  $G$  is a flat affine  $R$ -group scheme, not of finite type, with trivial special fiber and generic fiber  $\mathbb{G}_{a,K}$ . The affine  $R$ -algebra  $\mathcal{A}(G)$  is the subring of  $K[X]$  consisting of the polynomials whose constant coefficient lies in  $R$ . (N.B. We already encountered this example in Remark 11.10.1.) Furthermore,  $G$  represents the functor assigning to every  $R$ -algebra  $B$  the set of sequences  $(x_n)_{n \in \mathbb{N}}$  of elements of  $B$  such that*

$$x_n = \pi x_{n+1} \quad \text{for every } n.$$

(In particular, every  $x_n$  is indefinitely  $\pi$ -divisible.)

Now let  $S$  be a Noetherian scheme such that every coherent  $\mathcal{O}_S$ -module is a quotient of a locally free  $\mathcal{O}_S$ -module of finite type. This holds, for example, when  $S$  is a separated Noetherian regular scheme; see SGA 6, Exposé II, 2.1.1 and 2.2.7. (One can show that every Noetherian regular scheme of dimension at most 1 also has this property. It does not hold, however, when  $S$  is the affine plane over a field  $k$  with the origin doubled; see *loc. cit.*, 2.2.7.2.)

**Definition 13.4.**— *Let  $G$  be an affine flat  $S$ -group. Following R. W. Thomason ([Th87], 2.1), we say that the pair  $(G, S)$  has the equivariant resolution property, or satisfies (RE), if, for every coherent  $G$ - $\mathcal{O}_S$ -module  $\mathcal{F}$ , there are a locally free  $G$ - $\mathcal{O}_S$ -module  $\mathcal{E}$  of finite rank and a  $G$ -equivariant epimorphism*

$$\mathcal{E} \longrightarrow \mathcal{F}.$$

In *loc. cit.*, Theorem 3.1, Thomason proves the result below under the hypothesis that  $G$  be essentially free over  $S$  (see the remark below). Gabber pointed out to us the simpler proof given here, which does not use that hypothesis.

**Proposition 13.5.**— *Let  $S$  be a Noetherian scheme, and let  $G$  be an affine  $S$ -group, flat and of finite type, with affine algebra  $\mathcal{A}(G)$ . Suppose that  $(G, S)$  satisfies (RE). Then:*

(i)  $G$  is isomorphic to a closed subgroup of

$$H = \operatorname{Aut}_{\mathcal{O}_S}(\mathcal{V})$$

for some locally free  $\mathcal{O}_S$ -module  $\mathcal{V}$  of finite rank.

(ii) If, moreover,  $S$  is affine, one may take  $\mathcal{V} = \mathcal{O}_S^{\oplus d}$  for some  $d$ , so that  $H = \operatorname{GL}_{d,S}$ .<sup>†</sup>

The proof is analogous to that of 13.2. As there, there is a coherent  $G$ -stable  $\mathcal{O}_S$ -submodule  $\mathcal{F}$  that generates  $\mathcal{A} = \mathcal{A}(G)$  as an  $\mathcal{O}_S$ -algebra. Replacing  $S$  by one of its connected components, we may suppose that  $S$  is connected. By (RE), there are a locally free  $G$ - $\mathcal{O}_S$ -module  $\mathcal{E}$  of rank  $n$  and an epimorphism of  $\mathcal{A}$ -comodules

$$\pi : \mathcal{E} \longrightarrow \mathcal{F}.$$

Put  $H = \operatorname{Aut}_{\mathcal{O}_S}(\mathcal{E})$ . This is an  $S$ -group scheme locally isomorphic to  $\operatorname{GL}_{n,S}$ . The action of  $G$  on  $\mathcal{E}$  induces a morphism of affine  $S$ -groups

$$\rho : G \longrightarrow H,$$

corresponding to a morphism of  $\mathcal{O}_S$ -Hopf algebras

$$\varphi : \mathcal{A}(H) \longrightarrow \mathcal{A}(G).$$

We shall show that  $\rho$  is a closed immersion.

The question is local on  $S$ , so suppose that  $S = \operatorname{Spec}(R)$  and that

$$V = \Gamma(S, \mathcal{E})$$

is a free  $R$ -module with basis  $v_1, \dots, v_n$ . Then  $H \simeq \operatorname{GL}_{n,R}$ , and

$$B = \Gamma(S, \mathcal{A}(H))$$

is generated as an  $R$ -algebra by the matrix coefficients  $c_{ij}$  and by  $d^{-1}$ , where  $d$  denotes the determinant. Let  $\Delta$  (respectively,  $\varepsilon$ ) be the comultiplication (respectively, augmentation) of

$$A = \Gamma(S, \mathcal{A}(G)),$$

let

$$\mu : V \longrightarrow V \otimes_R A$$

be the  $A$ -comodule structure on  $V$ , and put

$$F = \Gamma(S, \mathcal{F}).$$

For  $j = 1, \dots, n$ , write

$$\mu(v_j) = \sum_{i=1}^n v_i \otimes a_{ij}.$$

Then  $\varphi(c_{ij}) = a_{ij}$ . Since  $\pi : V \rightarrow F$  is a morphism of  $A$ -comodules, one has

$$\sum_{i=1}^n \pi(v_i) \otimes a_{ij} = (\pi \otimes_R \operatorname{id}_A)(\mu(v_j)) = \Delta(\pi(v_j)),$$

---

<sup>†</sup> Source-reading note: the French statement prints “for some  $n$ ” after  $\mathcal{O}_S^{\oplus d}$ , while its conclusion is  $H = \operatorname{GL}_{d,S}$ . The same rank is denoted by  $d$  here. The Reinhold comparison lineage likewise uses  $d$ .

and therefore

$$\begin{aligned}\pi(v_j) &= (\varepsilon \otimes_R \text{id}_A) \Delta(\pi(v_j)) \\ &= \sum_{i=1}^n \varepsilon(\pi(v_i)) a_{ij} \\ &= \varphi \left( \sum_{i=1}^n \varepsilon(\pi(v_i)) c_{ij} \right).\end{aligned}$$

Thus  $\varphi(B)$  contains  $\pi(V) = F$ , which generates  $A$  as an  $R$ -algebra. Hence  $\varphi$  is surjective, and  $\rho$  is a closed immersion. This proves (i).

If  $S$  is affine, there is a locally free  $\mathcal{O}_S$ -module  $\mathcal{V}'$  of finite rank such that

$$\mathcal{V} \oplus \mathcal{V}' = \mathcal{O}_S^{\oplus d}.$$

Regard  $\mathcal{V}'$  as a trivial  $G$ -module and replace  $\mathcal{V}$  by  $\mathcal{O}_S^{\oplus d}$ . This realizes  $G$  as a closed subgroup of  $\text{GL}_{d,S}$  and completes the proof.

**Remarks 13.6.**—

- (a) For completeness, let us briefly sketch Thomason's argument ([Th87], Theorem 3.1), retaining the preceding notation. The  $G$ -equivariant epimorphism

$$\pi : \mathcal{E} \longrightarrow \mathcal{F}$$

induces a closed immersion

$$\tau : G \longrightarrow V(\mathcal{E})$$

such that

$$\tau(g'g) = \tau(g') \cdot g.$$

(N.B.  $G$  acts on the right on  $V(\mathcal{E})$ .) There is an isomorphism

$$H \simeq \text{Aut}_{\mathcal{O}_S}(V(\mathcal{E}))^{\text{op}}$$

compatible with the left action of  $G$  on  $\mathcal{E}$  and its right action on  $V(\mathcal{E})$ . Let

$$N = \text{Transp}_H^{\text{str}}(\tau(G), \tau(G))$$

be the strict transporter. If  $G$  is essentially free over  $S$ , it follows from 6.2.4(e) that  $N$  is a closed subgroup scheme of  $H$ , hence affine over  $S$ . Moreover,  $\rho$  factors through a morphism of  $S$ -groups

$$\rho' : G \longrightarrow N.$$

For every  $S' \rightarrow S$  and  $h \in N(S')$ , put

$$\pi(h) = \tau(1) \cdot h,$$

where 1 is the unit element of  $G(S')$ . This defines a morphism of  $S$ -schemes

$$\pi : N \longrightarrow \tau(G)$$

that is a retraction of  $\rho'$ , after identifying  $G$  with  $\tau(G)$ . Since  $N$  is separated over  $S$ , it follows that  $\rho'$  is a closed immersion.

- (b) The proof of [Th87], Theorem 3.1, appears to require the hypothesis that  $G$  be essentially free over  $S$ , although that hypothesis is not stated there (the author invokes instead the fact that  $H$  is essentially free). It is, however, satisfied when  $G$  is reductive (cf. Exposé XXII, 5.7.8), and hence in all cases considered in *loc. cit.*, Corollary 3.2. In particular, Thomason proves in *loc. cit.*, 2.5, that if  $S$  is separated, Noetherian, and regular of dimension at most 2, if  $G$  is affine and of finite presentation, and if  $\mathcal{A}(G)$  is a locally projective  $\mathcal{O}_S$ -module, then  $(G, S)$  satisfies (RE). By 13.5, this gives 13.2 under a slightly more restrictive hypothesis.

To conclude, let us point out that the proof of [Th87], 2.5, can be simplified slightly as follows. (For brevity, we work in the affine situation  $S = \operatorname{Spec}(R)$ .)

**Proposition 13.7.**— *Let  $R$  be a Noetherian regular ring of dimension at most 2, let  $C$  be an  $R$ -coalgebra that is projective as an  $R$ -module, and let  $F$  be a  $C$ -comodule of finite type over  $R$ . Then  $F$  is a quotient of a  $C$ -comodule  $V$  that is projective and of finite type over  $R$ .*

*Proof.* Replacing  $\operatorname{Spec}(R)$  by one of its connected components, we may suppose that  $R$  is integral, with fraction field  $K$ . Let  $\Delta$  (respectively,  $\varepsilon$ ) denote the comultiplication (respectively, augmentation) of  $C$ , and let

$$\rho : F \longrightarrow F \otimes C$$

be the comodule structure on  $F$ . Let

$$\pi : W \longrightarrow F$$

be a surjective morphism, with  $W$  a finite free  $R$ -module. Give  $W \otimes C$  the comodule structure defined by  $\operatorname{id}_W \otimes \Delta$ , and do the same for  $F \otimes C$ . Then  $\rho : F \rightarrow F \otimes C$  is a morphism of  $C$ -comodules with retraction  $\operatorname{id}_F \otimes \varepsilon$ .

Let  $W'$  be the  $C$ -comodule defined by the following cartesian square:

$$\begin{array}{ccc} W' & \longrightarrow & F \\ \downarrow & & \downarrow \rho \\ W \otimes C & \xrightarrow{\pi \otimes \operatorname{id}_C} & F \otimes C \end{array}$$

Thus  $W'$  identifies with the kernel of the morphism

$$W \otimes C \longrightarrow (F \otimes C)/\rho(F),$$

and the projection

$$\pi' : W' \longrightarrow F, \quad x \longmapsto (\pi \otimes \varepsilon)(x),$$

is surjective. Since  $F$  is a finitely generated  $R$ -module, there is a  $C$ -subcomodule  $V'$  of  $W'$ , finitely generated over  $R$ , such that

$$\pi'(V') = F.$$

Since  $W \otimes C$  is  $R$ -torsion-free, so is  $V'$ . Replacing  $F$  by  $V'$ , we may therefore suppose from the outset that  $F$  is torsion-free.

Apply the preceding construction to this new  $F$ , obtaining  $V'$  as above. Consider the subcomodule  $V$  that is the kernel of the morphism

$$W \otimes C \longrightarrow E = \frac{W \otimes C \otimes K}{V' \otimes K}.$$



Then  $V$  contains  $V'$ , and

$$Q = (W \otimes C)/V$$

is an  $R$ -submodule of the  $K$ -vector space  $E$ . Put

$$Q' = E/Q.$$

Since  $W \otimes C$  and  $E$  are flat  $R$ -modules, for every  $R$ -module  $N$  one obtains

$$\mathrm{Tor}_1^R(V, N) \simeq \mathrm{Tor}_2^R(Q, N) \simeq \mathrm{Tor}_3^R(Q', N).$$

Because  $R$  is regular of dimension at most 2, the right-hand term vanishes. Hence  $V$  is a flat  $R$ -module.

It remains to prove that  $V$  is finitely generated. Put

$$M = W \otimes C.$$

By construction,

$$V/V' \simeq (M/V')_{\mathrm{tors}},$$

the  $R$ -torsion submodule of  $M/V'$ . Since  $M$  is a projective  $R$ -module, there is a projective  $R$ -module  $P$  such that

$$M \oplus P = L$$

is free. Then

$$(M/V')_{\mathrm{tors}} \simeq (L/V')_{\mathrm{tors}}.$$

Since  $V'$  is finitely generated, there is a direct summand

$$L' \simeq R^n$$

of  $L$  containing  $V'$ . Hence

$$(L/V')_{\mathrm{tors}} \simeq (L'/V')_{\mathrm{tors}},$$

and the latter is finitely generated because  $L'/V'$  is. Thus  $V$  is a finitely generated flat  $R$ -module, hence a finitely generated projective  $R$ -module, since  $R$  is Noetherian. This proves Proposition 13.7.

## Bibliography

140

- [An73] S. Anantharaman, *Schémas en groupes, espaces homogènes et espaces algébriques sur une base de dimension 1*, Mém. Soc. Math. France **33** (1973), 5–79.
- [Ba55] I. Barsotti, *Un teorema di struttura per le varietà gruppali*, Atti Acad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Nat. **18** (1955), 43–50.
- [BLR] S. Bosch, W. Lütkebohmert, and M. Raynaud, *Néron Models*, Springer-Verlag, 1990.
- [BAC] N. Bourbaki, *Algèbre commutative*, Chapters I–IV, V–VII, and X, Masson, 1985 and 1998.
- [BTop] N. Bourbaki, *Topologie générale*, Chapters I–IV, Hermann, 1971.

---

<sup>140</sup>Editor's note: additional references cited in this Exposé.

- [Br09] M. Brion, *Anti-affine algebraic groups*, J. Algebra **321** (2009), no. 3, 934–952.
- [Ch60] C. Chevalley, *Une démonstration d'un théorème sur les groupes algébriques*, J. Maths. Pure Appl. **39** (1960), 307–317.
- [Co02] B. Conrad, *A modern proof of Chevalley's theorem on algebraic groups*, J. Ramanujan Math. Soc. **17** (2002), 1–18.
- [DG70] M. Demazure and P. Gabriel, *Groupes algébriques*, Masson & North-Holland, 1970.
- [Gr73] L. Gruson, *Dimension homologique des modules plats sur un anneau noethérien*, Symposia Mathematica **XI** (1973), 243–254.
- [Kn71] D. Knutson, *Algebraic spaces*, Lecture Notes in Mathematics **203**, Springer-Verlag, 1971.
- [Ma07] B. Margaux, *Passage to the Limit in Non-Abelian Čech Cohomology*, J. Lie Theory **17** (2007), 591–596.
- [MW94] A. Masuoka and D. Wigner, *Faithful flatness of Hopf algebras*, J. Algebra **170** (1994), 156–184.
- [Per76] D. Perrin, *Approximation des schémas en groupes, quasi-compacts sur un corps*, Bull. Soc. Math. France **104** (1976), 323–335.
- [Pes66] C. Peskine, *Une généralisation du “Main Theorem” de Zariski*, Bull. Sci. Math. **90** (1966), 119–127.
- [PY06] G. Prasad and J.-K. Yu, *On quasi-reductive group schemes*, J. Algebraic Geom. **15** (2006), 507–549.
- [Ray70a] M. Raynaud, *Faisceaux amples sur les schémas en groupes et les espaces homogènes*, Lecture Notes in Mathematics **119**, Springer-Verlag, 1970.
- [Ray70b] M. Raynaud, *Anneaux locaux henséliens*, Lecture Notes in Mathematics **169**, Springer-Verlag, 1970.
- [RG71] M. Raynaud and L. Gruson, *Critères de platitude et de projectivité*, Invent. Math. **13** (1971), 1–89.
- [Ro56] M. Rosenlicht, *Some basic theorems on algebraic groups*, Amer. J. Math. **78** (1956), 401–443.
- [SS09] C. Sancho de Salas and F. Sancho de Salas, *Principal bundles, quasi-abelian varieties and structure of algebraic groups*, J. Algebra **322** (2009), no. 8, 2751–2772.
- [Se68] J.-P. Serre, *Groupes de Grothendieck des schémas en groupes réductifs déployés*, Publ. Math. I.H.É.S. **34** (1968), 37–52.
- [Se99] C. S. Seshadri, *Chevalley: some reminiscences*, Transform. Groups **4** (1999), nos. 2–3, 119–125.
- [Ta72] M. Takeuchi, *A correspondence between Hopf ideals and sub-Hopf algebras*, Manuscripta Math. **7** (1972), 251–270.
- [Ta79] M. Takeuchi, *Relative Hopf modules—Equivalences and freeness criteria*, J. Algebra **60** (1979), 452–471.

- [Th87] R. W. Thomason, *Equivariant resolution, linearization, and Hilbert's fourteenth problem over arbitrary base schemes*, Adv. Math. **65** (1987), 16–34.

# Working-state notice for Exposé VII

The complete local Exposé VII translation is included below because this reader is intended to preserve and expose all currently readable work. It is *not* a certified release state. The included r5 source repairs the known page-26 diagram/footnote collision, but its detector-neutral reference package and final independent release audits were still in progress when this reader was assembled. This caveat concerns release verification, not the existence of the translated body.

# Exposé VII<sub>A</sub>

## Infinitesimal Study of Group Schemes

by P. Gabriel

In Exposé II we restricted ourselves to the study of first-order differential invariants, and we did not address certain phenomena special to characteristic  $p > 0$  or to characteristic zero. Our purpose in Part A of this exposé is to fill that gap.

Moreover, the infinitesimal study of a group scheme to arbitrary order is related to that of the associated formal group; the purpose of the second part of this exposé is to present the first definitions and properties concerning formal groups.

### A) Differential Operators and Restricted $p$ -Lie Algebras\*

#### 1. Differential Operators

In this section, as well as in Sections 2 and 3,  $S$  denotes a fixed scheme, and all products under consideration are cartesian products in the category of  $S$ -schemes.<sup>1</sup> If  $X$  is an  $S$ -scheme, we write  $p_{X/S}$ ,  $p_X$ , or simply  $p$  for the structural morphism from  $X$  to  $S$ .

**1.1.** Let  $u : Y \rightarrow X$  be a morphism of  $S$ -schemes, and endow the direct image  $u_*(\mathcal{O}_Y)$  of the structure sheaf of  $Y$  with the  $\mathcal{O}_X$ -module structure induced by  $u$ . The sheaf

$$\mathcal{H} = \mathcal{H}om_{p_X^{-1}(\mathcal{O}_S)}(\mathcal{O}_X, u_*(\mathcal{O}_Y))$$

of homomorphisms of  $p_X^{-1}(\mathcal{O}_S)$ -modules from  $\mathcal{O}_X$  to  $u_*(\mathcal{O}_Y)$  is therefore naturally an  $\mathcal{O}_X$ -bimodule. If  $U$  is an open subset of  $X$ , and  $f$  and  $d$  are sections over  $U$  of  $\mathcal{O}_X$  and  $\mathcal{H}$ , respectively, then  $fd$  and  $df$  are the morphisms

$$g \longmapsto f d(g) \quad \text{and} \quad g \longmapsto d(fg)$$

from  $\mathcal{O}_X$  to  $u_*(\mathcal{O}_Y)$ . From now on we write  $(\text{ad } f)(d)$  in place of  $fd - df$ .

**Definition 1.1.1.** An  $S$ -deviation of order  $\leq n$  is a pair  $D = (u, d)$ , consisting of a morphism of  $S$ -schemes  $u : Y \rightarrow X$  and a morphism of  $p_X^{-1}(\mathcal{O}_S)$ -modules

$$d : \mathcal{O}_X \longrightarrow u_*(\mathcal{O}_Y),$$

---

\*Part A of the present exposé was not treated seriously in the oral seminars.

<sup>1</sup>In particular, if  $X$  and  $Y$  are two  $S$ -schemes,  $X \times_S Y$  is denoted simply by  $X \times Y$ . For the material in Sections 1 and 2, one may also consult *Groupes algébriques* by Demazure and Gabriel, II.4, nos. 5–6; see also Jantzen, *Representations of Algebraic Groups*, I.7.

such that, for every open subset  $U$  of  $X$  and every sequence of  $n+1$  sections  $f_0, \dots, f_n \in \mathcal{O}_X(U)$ , one has

$$(\operatorname{ad} f_0)(\operatorname{ad} f_1) \cdots (\operatorname{ad} f_n)(d) = 0. \quad (*_n)$$

2

In this case we also say that  $d$  is an  $S$ -deviation of  $u$  of order  $\leq n$ . In particular, an  $S$ -deviation of  $u$  of order  $\leq 0$  is a morphism of  $\mathcal{O}_X$ -modules from  $\mathcal{O}_X$  to  $u_*(\mathcal{O}_Y)$ , that is, an element of  $\Gamma(Y, \mathcal{O}_Y)$ .

**Definition 1.1.2.** A morphism of  $p_X^{-1}(\mathcal{O}_S)$ -modules

$$d : \mathcal{O}_X \longrightarrow u_*(\mathcal{O}_Y)$$

is an  $S$ -deviation of  $u$  if, for every point  $y \in Y$ , there exist an open neighborhood  $U$  of  $u(y)$  in  $X$  and an open neighborhood  $V$  of  $y$  in  $Y$  satisfying:

- a)  $u(V) \subset U$ ;
- b) if  $v : V \rightarrow U$  is induced by  $u$ , there is an integer  $n$  such that the morphism  $\mathcal{O}_U \rightarrow v_*(\mathcal{O}_V)$  induced by  $d$  is an  $S$ -deviation of  $v$  of order  $\leq n$ .

3

If  $d$  is an  $S$ -deviation of  $u$ , we also call the pair  $D = (u, d)$  an  $S$ -deviation, and we may write

$$Y \xrightarrow{D} X \quad \text{or} \quad Y \xrightarrow[u]{d} X.$$

When  $d$  is the homomorphism of algebras

$$u^\sharp : \mathcal{O}_X \longrightarrow u_*(\mathcal{O}_Y)$$

corresponding to the morphism  $u : Y \rightarrow X$ , we also write  $u$  in place of  $D$ .

**Remarks 1.1.3.**<sup>4</sup> Let  $\operatorname{Dev}(u)$  (respectively,  $\operatorname{Dev}_{\leq n}(u)$ ) be the set of  $S$ -deviations of  $u$  (respectively, of order  $\leq n$ ). It has a natural  $\mathcal{O}_Y(Y)$ -module structure: if  $\lambda \in \mathcal{O}_Y(Y)$ , then  $\lambda d$  is the deviation that sends  $f$  to  $\lambda d(f)$  for every local section  $f$  of  $\mathcal{O}_X$ .

For every open subset  $V$  of  $Y$ , set  $\mathcal{D}|_{\square}(u)(V) = \operatorname{Dev}(u|_V)$ . Thus  $\mathcal{D}|_{\square}(u)(V)$  is the set of

$$\begin{aligned} d_V &\in \operatorname{Hom}_{p_X^{-1}(\mathcal{O}_S)}(\mathcal{O}_X, (u|_V)_*(\mathcal{O}_V)) \\ &\simeq \operatorname{Hom}_{p_Y^{-1}(\mathcal{O}_S)}((u|_V)^{-1}\mathcal{O}_X, \mathcal{O}_V) \\ &\simeq \operatorname{Hom}_{p_Y^{-1}(\mathcal{O}_S)}(u^{-1}\mathcal{O}_X, \mathcal{O}_Y)(V) \end{aligned}$$

<sup>2</sup>Equivalently, for every  $x \in X$  and  $f_0, \dots, f_n, g \in \mathcal{O}_{X,x}$ ,

$$(\operatorname{ad} f_0) \cdots (\operatorname{ad} f_n)(d_x)(g) = 0.$$

Recall also that the adjunction isomorphism

$$\theta : \operatorname{Hom}_{p_X^{-1}(\mathcal{O}_S)}(\mathcal{O}_X, u_*(\mathcal{O}_Y)) \xrightarrow{\sim} \operatorname{Hom}_{p_Y^{-1}(\mathcal{O}_S)}(u^{-1}\mathcal{O}_X, \mathcal{O}_Y)$$

sends  $d$  to  $d' = \varepsilon \circ u^{-1}(d)$ , where  $\varepsilon : u^{-1}u_*(\mathcal{O}_Y) \rightarrow \mathcal{O}_Y$  is canonical. Conversely,  $\theta^{-1}(d') = u_*(d') \circ \eta$ , with  $\eta : \mathcal{O}_X \rightarrow u_*u^{-1}\mathcal{O}_X$  canonical. Hence  $d$  satisfies  $(*_n)$  if and only if  $d'$  satisfies the analogous identity for all local sections of  $u^{-1}\mathcal{O}_X$ .

<sup>3</sup>If  $X$  and  $u$  are quasi-compact, every  $S$ -deviation of  $u$  is therefore of order  $\leq n$  for some integer  $n$ .

<sup>4</sup>These remarks were added by the editors because they are useful in 1.3, 1.4, and 2.1.

such that, for every open  $U \subset X$ , the map

$$d_V(U) : \mathcal{O}_X(U) \longrightarrow \mathcal{O}_Y(u^{-1}(U) \cap V)$$

satisfies  $(*_n)$ . This defines a presheaf of  $\mathcal{O}_Y$ -modules on  $Y$ , and it is easily seen to be a sheaf (more precisely, a subsheaf of  $\mathcal{H}om_{p_Y^{-1}(\mathcal{O}_S)}(u^{-1}\mathcal{O}_X, \mathcal{O}_Y)$ ).

**1.2.** Consider two  $S$ -deviations  $D = (u, d)$  and  $E = (v, e)$ :

$$Z \xrightarrow[v]{e} Y \xrightarrow[u]{d} X.$$

As  $U$  ranges over the open subsets of  $X$ , the composites

$$\Gamma(U, \mathcal{O}_X) \xrightarrow{d(U)} \Gamma(u^{-1}U, \mathcal{O}_Y) \xrightarrow{e(u^{-1}U)} \Gamma(v^{-1}u^{-1}U, \mathcal{O}_Z)$$

define an  $S$ -deviation of  $uv$ , denoted by  $de$ . If  $d$  has order  $\leq m$  and  $e$  has order  $\leq n$ , then  $de$  has order  $\leq m + n$ . We also write

$$D \circ E = (uv, de) \tag{†}$$

<sup>5</sup> and call  $D \circ E$ , or  $DE$ , the *composite  $S$ -deviation*. When  $d = u^\natural$ , so that  $D = u$  under the convention of 1.1, we also say that  $DE$  is the *image of  $E$  under  $u$* .

The map  $(D, E) \mapsto D \circ E$  allows us to speak from now on of the *category of  $S$ -deviations*, whose objects are  $S$ -schemes and whose morphisms are  $S$ -deviations.<sup>6</sup>

**Definition 1.2.0.**<sup>7</sup> Let  $w : Z \rightarrow X$  be an  $S$ -morphism. An  $S$ -derivation of  $w$ , or an  $S$ -derivation of  $\mathcal{O}_X$  into  $w_*(\mathcal{O}_Z)$ , is a morphism of  $p^{-1}(\mathcal{O}_S)$ -modules

$$d : \mathcal{O}_X \longrightarrow w_*(\mathcal{O}_Z)$$

such that, for every open  $U \subset X$  and  $f, g \in \mathcal{O}_X(U)$ ,

$$d(fg) = w^\natural(f)d(g) + w^\natural(g)d(f).$$

Then  $d$  is a deviation of  $w$  of order  $\leq 1$ , vanishing on the unit section of  $\mathcal{O}_X$ . We denote the set of  $S$ -derivations of  $w$  by  $\text{Der}_S(w)$ ; it is an  $\mathcal{O}(Z)$ -module.

With the notation of 1.2, take

$$Y = I_Z = \text{Spec } \mathcal{O}_Z[t], \quad t^2 = 0,$$

let  $v$  be the zero section  $\tau : Z \rightarrow I_Z$ , defined by the homomorphism  $\mathcal{O}_Z[t] \rightarrow \mathcal{O}_Z$  sending  $t$  to zero, and let  $e$  be the morphism of  $\mathcal{O}_Z$ -modules  $\sigma : \mathcal{O}_Z[t] \rightarrow \mathcal{O}_Z$  defined by  $\sigma(1) = 0$  and  $\sigma(t) = 1$ , which it is convenient to denote by  $\partial_t$ .<sup>8</sup>

If  $u : I_Z \rightarrow X$  is a morphism satisfying  $w = u \circ \tau$ , then  $\sigma \circ u^\natural$  is an  $S$ -derivation from  $\mathcal{O}_X$  to  $w_*(\mathcal{O}_Z)$ . Conversely, an  $S$ -derivation  $d$  determines the morphism  $u : I_Z \rightarrow X$  which equals  $w$  on the underlying spaces and satisfies

$$u^\natural(f) = w^\natural(f) + d(f)t$$

<sup>5</sup>With this notation,  $de$  means the composite “ $d$  followed by  $e$ .”

<sup>6</sup>Often one considers only the  $S$ -deviations of  $\text{id}_X$ ; they form the algebra of  $S$ -differential operators on  $X$ , cf. 1.4 below. The more general framework provides a convenient functorial language for statements such as the following: if  $G$  is an  $S$ -group, the algebra of left-translation-invariant  $S$ -differential operators on  $G$  is isomorphic to the algebra of  $S$ -deviations of the unit section  $\varepsilon : S \rightarrow G$ ; cf. 2.1 and 2.4.

<sup>7</sup>The editors expanded this paragraph and assigned the number 1.2.0 to this definition and 1.2.1 to the lemma that follows.

<sup>8</sup>The editors added this notation  $\partial_t$ .

for every local section  $f$  of  $\mathcal{O}_X$ .

**Lemma 1.2.1.** *Let  $E = (\tau, \partial_t)$  be the deviation of  $\tau : Z \rightarrow I_Z$  defined above. For every  $S$ -morphism  $w : Z \rightarrow X$ , the map  $u \mapsto u \circ E$  is a bijection between the  $S$ -morphisms  $u : I_Z \rightarrow X$  such that  $u \circ \tau = w$ , and the  $S$ -derivations of  $w$ .*

**1.2.2.** Let  $d$  be an  $S$ -deviation of  $u : Y \rightarrow X$ . On the one hand,  $d$  is plainly an  $S'$ -deviation of  $u$  for every morphism  $s : S \rightarrow S'$ .

On the other hand, let  $t : T \rightarrow S$  be a morphism with target  $S$ , let  $u_T : Y_T \rightarrow X_T$  be obtained from  $u$  by base change, and let  $t_Y : Y_T \rightarrow Y$  and  $t_X : X_T \rightarrow X$  be the canonical projections. There exists a unique  $T$ -deviation of  $u_T$ , denoted  $d_T$  or  $d \times T$ , which satisfies

$$t_X d_T = dt_Y$$

in the sense of  $(\dagger)$ . Thus, for every open  $U \subset X$ , the following diagram is commutative:

$$\begin{array}{ccc} \mathcal{O}(U) & \xrightarrow{t_X^\sharp} & \mathcal{O}(U \times T) \\ d(U) \downarrow & & \downarrow d_T(U \times T) \\ \mathcal{O}(u^{-1}U) & \xrightarrow{t_Y^\sharp} & \mathcal{O}(u^{-1}U \times T). \end{array}$$

<sup>9</sup> If  $D = (u, d)$ , we also write  $D_T = (u_T, d_T)$ , and say that  $d_T$  and  $D_T$  are obtained from  $d$  and  $D$  by base change.

**1.2.3.** For example, let  $u : Y \rightarrow X$  and  $v : Z \rightarrow T$  be two  $S$ -morphisms, and let  $d$  and  $e$  be  $S$ -deviations of  $u$  and  $v$ . There is a commutative diagram

$$\begin{array}{ccc} X \times T & \xleftarrow{u_T} & Y \times T \\ v_X \uparrow & \swarrow u \times v & \uparrow v_Y \\ X \times Z & \xleftarrow{u_Z} & Y \times Z \end{array}$$

and we denote by  $d \times e$ , the *product of  $d$  and  $e$* , the  $S$ -deviation of  $u \times v$  equal to

$$d_T e_Y = e_X d_Z$$

under the convention of  $(\dagger)$ . More explicitly, for every open  $U \subset X \times T$ , set

$$W = v_Y^{-1} u_T^{-1} U = u_Z^{-1} v_X^{-1} U.$$

Then the following diagram is commutative:

$$\begin{array}{ccc} \mathcal{O}(U) & \xrightarrow{d_T(U)} & \mathcal{O}(u_T^{-1}U) \\ e_X(U) \downarrow & \searrow (d \times e)(U) & \downarrow e_Y(u_T^{-1}U) \\ \mathcal{O}(v_X^{-1}U) & \xrightarrow{d_Z(v_X^{-1}U)} & \mathcal{O}(W). \end{array}$$

<sup>9</sup>Explicitly, if  $V$  is an affine open subset of  $S$ , and  $U$  (respectively,  $U'$ ) is an affine open subset of  $X$  (respectively,  $T$ ) over  $V$ , then

$$\mathcal{O}_{X \times T}(U \times U') = \mathcal{O}_X(U) \otimes_{\mathcal{O}_S(V)} \mathcal{O}_T(U'),$$

and  $d_T(U \times U')$  is the composite

$$\mathcal{O}_X(U) \otimes_{\mathcal{O}_S(V)} \mathcal{O}_T(U') \xrightarrow{d(U) \otimes \text{id}} \mathcal{O}_Y(u^{-1}U) \otimes_{\mathcal{O}_S(V)} \mathcal{O}_T(U') \longrightarrow \mathcal{O}_{Y \times T}(u^{-1}U \times U').$$

The author left to the reader the verification that  $d_T$  is well-defined, and the editors do the same.



If  $D = (u, d)$  and  $E = (v, e)$ , we also write

$$D \times E = (u \times v, d \times e).$$

10

**1.3.**<sup>11</sup> Let  $u : Y \rightarrow X$  be a morphism of  $S$ -schemes. Recall that the adjunction isomorphism

$$\mathrm{Hom}_{p_X^{-1}(\mathcal{O}_S)}(\mathcal{O}_X, u_*(\mathcal{O}_Y)) \xrightarrow{\sim} \mathrm{Hom}_{p_Y^{-1}(\mathcal{O}_S)}(u^{-1}\mathcal{O}_X, \mathcal{O}_Y)$$

sends a morphism of  $p^{-1}(\mathcal{O}_S)$ -modules

$$d : \mathcal{O}_X \longrightarrow u_*(\mathcal{O}_Y)$$

to  $d' = \varepsilon \circ u^{-1}(d)$ , where  $\varepsilon : u^{-1}u_*(\mathcal{O}_Y) \rightarrow \mathcal{O}_Y$  is the canonical morphism.

Write  $\mathcal{J}_u$  (respectively,  $\mathcal{I}_u$ ) for the kernel of the homomorphism of algebras

$$u^\sharp : \mathcal{O}_X \longrightarrow u_*(\mathcal{O}_Y) \quad \left( \text{respectively, } u^{\sharp'} : u^{-1}\mathcal{O}_X \longrightarrow \mathcal{O}_Y \right),$$

and let  $d : \mathcal{O}_X \rightarrow u_*(\mathcal{O}_Y)$  be a morphism of  $p^{-1}(\mathcal{O}_S)$ -modules. If  $U \subset X$  is open and  $f_0, \dots, f_n, g \in \mathcal{O}_X(U)$ , induction on  $n$  shows that condition  $(*)_n$  is equivalent to

$$0 = \sum_{I \subset \{0, \dots, n\}} (-1)^{|I|} u^\sharp(f_{\{0, \dots, n\} \setminus I}) d(f_I g), \quad (**_n)$$

where  $f_I$  denotes the product of the  $f_i$  for  $i \in I$ . Consequently, if  $d$  satisfies  $(*)_n$ , then  $d$  vanishes on the ideal  $\mathcal{J}_u^{n+1}$ .

Now suppose that  $Y = S$ . Then  $u : S \rightarrow X$  is a section of  $p : X \rightarrow S$ , hence an immersion. On the one hand,

$$\varepsilon : u^{-1}u_*\mathcal{O}_S \longrightarrow \mathcal{O}_S$$

is an isomorphism, and therefore  $u^{-1}\mathcal{J}_u = \mathcal{I}_u$ . On the other hand, there is an isomorphism

$$u^{-1}\mathcal{O}_X \simeq \mathcal{O}_S \oplus \mathcal{I}_u. \quad (\star)$$

Suppose that  $d$  vanishes on  $\mathcal{J}_u^{n+1}$ . Then  $d' = \varepsilon \circ u^{-1}(d)$  vanishes on  $\mathcal{I}_u^{n+1}$ , and therefore satisfies the analogues  $(**'_n)$  and  $(*'_n)$  of  $(**_n)$  and  $(*)_n$  when

$$f_0, \dots, f_n \in \mathcal{I}_u(u^{-1}(U)).$$

Moreover,

$$(\mathrm{ad} a)(\varphi) = 0$$

for every  $a \in \mathcal{O}_S(u^{-1}U)$  and every morphism of  $\mathcal{O}_{u^{-1}U}$ -modules  $\varphi : u^{-1}\mathcal{O}_U \rightarrow \mathcal{O}_{u^{-1}U}$ . The splitting  $(\star)$  therefore implies that  $d'$  satisfies  $(*_n')$ , and hence that  $d$  satisfies  $(*)_n$ .

**Lemma.** *If  $u : S \rightarrow X$  is a section of  $p : X \rightarrow S$ , then  $d$  is an  $S$ -deviation of  $u$  of order  $\leq n$  if and only if  $d'$  vanishes on  $\mathcal{I}_u^{n+1}$ .*

This interpretation generalizes as follows. Let  $u : Y \rightarrow X$  be any  $S$ -morphism, and let  $\Gamma_u : Y \rightarrow Y \times X$  be its graph, with components  $\mathrm{id}_Y$  and  $u$ . Every  $S$ -deviation  $d$  of  $u$  of order  $\leq n$  gives, by composition,

$$Y \xrightarrow{\mathrm{diag}} Y \times Y \xrightarrow[u_Y]{d_Y} Y \times X,$$

<sup>10</sup>Source note: the French sentence at this point prints  $E = (v, d)$ ; the established notation is  $E = (v, e)$ .

<sup>11</sup>The editors expanded the original in this paragraph; see also their note 2 to Definition 1.1.1.

a  $Y$ -deviation of  $\Gamma_u$  of order  $\leq n$ , denoted  $\Gamma_d$ , the graph of  $d$ .

Conversely, a  $Y$ -deviation  $e$  of  $\Gamma_u$  gives the composite  $S$ -deviation

$$e_X = \text{pr}_2 \circ e : Y \xrightarrow[\Gamma_u]{e} Y \times X \xrightarrow{\text{pr}_2} X.$$

One has immediately  $(\Gamma_d)_X = d$ . The identity  $\Gamma_{e_X} = e$  follows from the  $\mathcal{O}_Y$ -linearity of  $e$ .<sup>12</sup> Thus there is an isomorphism of  $\mathcal{O}_Y(Y)$ -modules

$$\left\{ \begin{array}{c} S\text{-deviations of } u \\ \text{of order } \leq n \end{array} \right\} \xrightarrow{\sim} \left\{ \begin{array}{c} Y\text{-deviations of } \Gamma_u \\ \text{of order } \leq n \end{array} \right\}, \quad d \mapsto \Gamma_d.$$

Moreover,  $d$  is an  $S$ -derivation of  $u$  if and only if  $\Gamma_d$  is a  $Y$ -derivation of  $\Gamma_u$ .

Let  $\mathcal{I}_{\Gamma_u}$  denote the kernel of the homomorphism of algebras

$$\Gamma_u^{-1} \mathcal{O}_{Y \times X} \longrightarrow \mathcal{O}_Y$$

corresponding to  $\Gamma_u$ . The preceding lemma yields:

**Proposition.** *Let  $u : Y \rightarrow X$  be an  $S$ -morphism and  $\Gamma_u : Y \rightarrow Y \times X$  its graph. The  $S$ -deviations of  $u$  of order  $\leq n$  identify with the  $Y$ -deviations of  $\Gamma_u$  of order  $\leq n$ , and these are in bijection with*

$$\text{Hom}_{\mathcal{O}_Y} \left( \frac{\Gamma_u^{-1} \mathcal{O}_{Y \times X}}{\mathcal{I}_{\Gamma_u}^{n+1}}, \mathcal{O}_Y \right).$$

**1.3.1.**<sup>13</sup> Return to the case in which  $u : S \rightarrow X$  is a section of  $p : X \rightarrow S$ . The homomorphism

$$\varphi : u^{-1} \mathcal{O}_X \longrightarrow \mathcal{O}_S$$

has a section, denoted simply by  $g \mapsto g \cdot 1$ . Thus, with the notation of 1.3,

$$u^{-1} \mathcal{O}_X \simeq \mathcal{O}_S \oplus \mathcal{I}_u,$$

and, for every section  $f$  of  $u^{-1} \mathcal{O}_X$ ,

$$f - \varphi(f) \cdot 1 \in \mathcal{I}_u.$$

Let  $d$  be an  $S$ -deviation of  $u$  of order  $\leq 1$ , and let  $d' : u^{-1} \mathcal{O}_X \rightarrow \mathcal{O}_S$  be the  $\mathcal{O}_S$ -morphism corresponding to  $d$ . If  $a, b$  are sections of  $u^{-1} \mathcal{O}_X$ , then

$$\begin{aligned} 0 &= d'((a - \varphi(a) \cdot 1)(b - \varphi(b) \cdot 1)) \\ &= d'(ab) - \varphi(a)d'(b) - \varphi(b)d'(a) + \varphi(ab)d'(1). \end{aligned}$$

Consequently,  $d$  is an  $S$ -derivation of  $u$  if and only if  $d'(1) = 0$ .

**Lemma.** *The  $S$ -derivations of  $u$  are precisely the  $S$ -deviations of  $u$  of order 1 which vanish on the unit section of  $\mathcal{O}_X$ . They correspond to the  $\mathcal{O}_S(S)$ -module*

$$\text{Hom}_{\mathcal{O}_S}(\mathcal{I}_u/\mathcal{I}_u^2, \mathcal{O}_S),$$

<sup>12</sup>If  $\lambda$  and  $f$  are local sections of  $\mathcal{O}_Y$  and  $\mathcal{O}_X$ , then

$$(\Gamma_{e_X})(\lambda \otimes f) = \lambda e(1 \otimes f) = e(\lambda \otimes f).$$

The French note prints  $g$  in the two right-hand tensor expressions; the variables just introduced require  $f$ .

<sup>13</sup>This paragraph was added by the editors.

and there is an isomorphism of  $\mathcal{O}_S(S)$ -modules

$$\mathrm{Dev}_{\leq 1}(u) \simeq \mathcal{O}_S(S) \oplus \mathrm{Der}_S(u).$$

Returning to the general case gives:

**Corollary.** *Let  $u : Y \rightarrow X$  be an  $S$ -morphism and  $\Gamma_u : Y \rightarrow Y \times X$  its graph. There is a canonical isomorphism of  $\mathcal{O}_Y(Y)$ -modules*

$$\mathrm{Der}_S(u) \simeq \mathrm{Der}_Y(\Gamma_u) \simeq \mathrm{Hom}_{\mathcal{O}_Y}(\mathcal{I}_{\Gamma_u}/\mathcal{I}_{\Gamma_u}^2, \mathcal{O}_Y).$$

**Definition 1.4.** *Let  $X$  be an  $S$ -scheme. An  $S$ -differential operator (respectively, an  $S$ -differential operator of order  $\leq n$ ) on  $X$  is an  $S$ -deviation (respectively, an  $S$ -deviation of order  $\leq n$ ) of the identity morphism of  $X$ .*

By 1.1, an  $S$ -differential operator of order  $\leq n$  is therefore an endomorphism of the  $p^{-1}(\mathcal{O}_S)$ -module  $\mathcal{O}_X$  satisfying the identities  $(*_n)$  of 1.1. We write  $\mathrm{Dif}_{X/S}^n$  for the  $\Gamma(\mathcal{O}_S)$ -module<sup>14</sup> of  $S$ -differential operators of order  $\leq n$ , and  $\mathrm{Dif}_{X/S}$  for the module of all  $S$ -differential operators.

As observed in 1.2,  $S$ -deviations of  $\mathrm{id}_X$  can be composed. This makes  $\mathrm{Dif}_{X/S}$  a  $\Gamma(\mathcal{O}_S)$ -algebra, called the *algebra of differential operators of  $X/S$* .

Similarly, for every open subset  $V \subset X$ , set

$$\mathcal{D}\{_{X/S}(V) = \mathrm{Dif}_{V/S} = \mathrm{Dev}(\mathrm{id}_V).$$

By 1.1.3 this defines a sheaf of  $\mathcal{O}_X$ -modules, called the *sheaf of  $S$ -differential operators on  $X$* .<sup>15</sup>

**1.4.1.** As in 1.3, differential operators on  $X/S$  may be interpreted using the graph of the identity morphism of  $X$ , namely the diagonal

$$\Delta = \Delta_{X/S} : X \longrightarrow X \times X.$$

Endow  $\mathcal{O}_{X \times X}$  with the  $\mathrm{pr}_1^{-1}(\mathcal{O}_X)$ -algebra structure defined by  $\mathrm{pr}_1$ . Then  $\Delta^{-1}\mathcal{O}_{X \times X}$  is an algebra over

$$\mathcal{O}_X = \Delta^{-1} \mathrm{pr}_1^{-1} \mathcal{O}_X.$$

Let  $\mathcal{I}_{X/S}$  be the kernel of the homomorphism

$$\Delta^{-1}\mathcal{O}_{X \times X} \xrightarrow{m} \mathcal{O}_X$$

adjoint to  $\mathcal{O}_{X \times X} \rightarrow \Delta_*(\mathcal{O}_X)$ , and set

$$\mathcal{P}_{X/S}^m = \Delta^{-1}\mathcal{O}_{X \times X} / \mathcal{I}_{X/S}^{m+1}.$$

If  $V \subset S$  and  $U \subset X$  are affine open subsets with  $U$  above  $V$ , and if

$$k = \Gamma(V, \mathcal{O}_S), \quad A = \Gamma(U, \mathcal{O}_X),$$

then

$$\Gamma(U, \mathcal{P}_{X/S}^m) = (A \otimes_k A) / I^{m+1},$$

where  $I$  is generated by the elements  $a \otimes 1 - 1 \otimes a$ , for  $a \in A$ .

<sup>14</sup>In this exposé the ring  $\Gamma(S, \mathcal{O}_S) = \mathcal{O}_S(S)$  is denoted by  $\Gamma(\mathcal{O}_S)$ .

<sup>15</sup>The editors modified the original here. It mentioned the sheaf  $U \mapsto \mathrm{Dif}_{X_U/U}$ , where  $U$  ranges over open subsets of  $S$ ; that sheaf is the direct image of  $\mathcal{D}\{_{X/S}$  under  $p_X : X \rightarrow S$ .

By 1.3 there is an isomorphism of  $\mathcal{O}_X(X)$ -modules

$$j_X : \text{Dif}_{X/S}^m \xrightarrow{\sim} \text{Hom}_{\mathcal{O}_X}(\mathcal{P}_{X/S}^m, \mathcal{O}_X).$$

It is given explicitly as follows: if  $d \in \text{Dif}_{X/S}^m$  and a section  $c$  of  $\mathcal{P}_{X/S}^m$  over  $U$  has the form

$$c = a \otimes b + I^{m+1},$$

then  $j_X(d)(c) = a d(b)$ .<sup>16</sup>

**1.4.2.** Let  $d$  be a differential operator and  $u$  a section of  $X$  over  $S$ . The *value of  $d$  at  $u$*  is the composite  $S$ -deviation

$$S \xrightarrow{u} X \xrightarrow[\text{id}_X]{d} X.$$

By 1.3 and 1.4.1, if  $d$  has order  $\leq m$ , then  $du$  (respectively,  $d$ ) is canonically associated with a morphism of  $\mathcal{O}_S$ -modules

$$d' : u^{-1}\mathcal{O}_X/\mathcal{I}_u^{m+1} \longrightarrow \mathcal{O}_S$$

(respectively, a morphism of  $\mathcal{O}_X$ -modules  $d'' : \mathcal{P}_{X/S}^m \rightarrow \mathcal{O}_X$ ).

The morphism  $d'$  may be recovered from  $d''$  as follows. The square

$$\begin{array}{ccc} X \simeq S \times X & \xrightarrow{u \times X} & X \times X \\ p \downarrow & & \downarrow \text{pr}_1 \\ S & \xrightarrow{u} & X \end{array}$$

is cartesian. Hence  $X$  identifies with  $S \times_X (X \times X)$ ,  $u$  with  $S \times_X \Delta$ , and  $u^*(\mathcal{P}_{X/S}^m)$  with  $u^{-1}\mathcal{O}_X/\mathcal{I}_u^{m+1}$ . Under these identifications,  $u^*(d'')$  is the morphism

$$u^{-1}\mathcal{O}_X/\mathcal{I}_u^{m+1} \longrightarrow \mathcal{O}_S$$

which is precisely  $d'$ .

**1.5.** Set, as usual,

$$I_S = \text{Spec } \mathcal{O}_S[T]/(T^2).$$

Let  $\tau : S \rightarrow I_S$  be the zero section, and let  $\sigma$  be the canonical deviation of  $\tau$  defined in 1.2.0: it is the  $\mathcal{O}_S$ -module homomorphism which vanishes on the unit section of  $\mathcal{O}_S[T]/(T^2)$  and sends the class  $t$  of  $T$  modulo  $T^2$  to the unit section of  $\mathcal{O}_S$ .

Let  $X$  be an  $S$ -scheme. Every  $I_S$ -automorphism  $u$  of  $I_S \times X$  inducing the identity on  $X$  determines, by composition, a differential operator  $D_u$  on  $X$ :

$$X \simeq S \times X \xrightarrow{\sigma \times X} I_S \times X \xrightarrow{u} I_S \times X \xrightarrow{\text{pr}_2} X.$$

By Exposé II, 3.14, the map  $u \mapsto D_u$  is an isomorphism from the  $\Gamma(\mathcal{O}_S)$ -Lie algebra

$$\text{Lie}(\text{Aut } X) := \text{Lie}(\text{Aut } X)(S)$$

onto the  $\Gamma(\mathcal{O}_S)$ -Lie algebra of  $p^{-1}(\mathcal{O}_S)$ -derivations of  $\mathcal{O}_X$ . Its inverse sends a derivation  $D$  to the automorphism of  $I_S \times X$  corresponding to

$$a + bt \mapsto a + (Da + b)t$$

on  $\mathcal{O}_X[T]/(T^2)$ .

---

<sup>16</sup>Under this isomorphism, the  $X$ -derivations of  $\Delta_{X/S}$  correspond, by 1.3.1, to the  $S$ -derivations of  $\text{id}_X$ , that is, to the  $p^{-1}(\mathcal{O}_S)$ -derivations of  $\mathcal{O}_X$ .

## 2. Invariant Differential Operators on Group Schemes

**2.1.** Let  $G$  be an  $S$ -group scheme; we denote its unit section by  $\varepsilon$ , or by  $\varepsilon_G : S \rightarrow G$ .

**Definition.** Let  $U(G)$  be the  $\Gamma(\mathcal{O}_S)$ -module of  $S$ -deviations of  $\varepsilon_G$  (or  $S$ -deviations of the origin), as in 1.1.

If  $d, e \in U(G)$ , then  $d \times e$  is an  $S$ -deviation of

$$\varepsilon \times \varepsilon : S \simeq S \times S \longrightarrow G \times G.$$

The image of  $d \times e$  under the multiplication morphism  $m : G \times G \rightarrow G$ , as in 1.2, is called the product of  $d$  and  $e$  and is denoted  $d \cdot e$ .

The  $\Gamma(\mathcal{O}_S)$ -module  $U(G)$  is thereby made into an associative  $\Gamma(\mathcal{O}_S)$ -algebra with unit  $\varepsilon_G$ . We call  $U(G)$  the infinitesimal algebra of  $G$ .<sup>17</sup>

As  $T$  ranges over schemes above  $S$ , the infinitesimal algebra  $U(G_T)$  of the  $T$ -group  $G \times T$  plainly varies contravariantly with  $T$ ; thus one may speak of the *infinitesimal-algebra functor*.

Restricting  $T$  to the open subsets of  $S$  gives a presheaf  $T \mapsto U(G_T)$  of  $\mathcal{O}_S$ -algebras. By 1.1.3 it is a sheaf. We denote it by  $\mathcal{U}(G)$  and call it the *sheaf of infinitesimal algebras of  $G$* .

The algebra  $U(G)$  is also covariantly functorial in  $G$ . Indeed, if  $u : G \rightarrow H$  is a homomorphism of  $S$ -groups and  $d$  is an  $S$ -deviation of  $\varepsilon_G$ , its image under  $u$  is the element

$$U(u)(d) = ud$$

of  $U(H)$ . The resulting map  $U(u) : U(G) \rightarrow U(H)$  is a homomorphism of  $\Gamma(\mathcal{O}_S)$ -algebras. Likewise one obtains a homomorphism  $\mathcal{U}(u) : \mathcal{U}(G) \rightarrow \mathcal{U}(H)$ .

**2.2.** Let  $d \in U(G)$ , that is, let  $d$  be an  $S$ -deviation of the origin of  $G$ . The base change of  $d$  is the  $S$ -deviation

$$d \times G$$

of  $\varepsilon \times G : G \simeq S \times G \rightarrow G \times G$ , as in 1.2.2. Its image under multiplication  $m : G \times G \rightarrow G$  is an  $S$ -deviation of

$$m \circ (\varepsilon \times \text{id}_G) = \text{id}_G,$$

hence an element of  $\text{Dif}_{G/S}$ , which we denote by  $d^G$ .

The map  $d \mapsto d^G$  is plainly  $\Gamma(\mathcal{O}_S)$ -linear. The following commutative diagram shows that

$$(e \cdot d)^G = d^G \cdot e^G.$$

$$\begin{array}{ccccc}
 & & G \times G \times G & \xrightarrow{m \times G} & G \times G \\
 & \nearrow^{G \times d \times G} & & \searrow^{G \times m} & \\
 & G \times G & & G \times G & \\
 \nearrow^{e \times G} & & \nearrow^{d \times G} & & \nearrow^{m} \\
 G & \xrightarrow{\varepsilon \times G} & G & \xrightarrow{\varepsilon \times G} & G \\
 \searrow^{e^G} & & \searrow^{d^G} & & \searrow^{m} \\
 & G & & G & \\
 \xrightarrow{\text{id}_G} & & \xrightarrow{\text{id}_G} & & 
 \end{array}$$

<sup>17</sup>This is now usually called the algebra of distributions (at the origin) of  $G$ ; see Demazure–Gabriel, II.4, 6.1, and Jantzen, I.7.7.

18

The two lower triangles commute by the definitions of  $d^G$  and  $e^G$ . The composite  $S$ -deviation of  $e \times G$  and  $G \times d \times G$  is  $(e \times d) \times G$ , by 1.2.2. Its image under  $m \times G$  is  $(e \cdot d) \times G$ , and the image of the latter under  $m$  is therefore  $(e \cdot d)^G$ .

Thus one obtains an antihomomorphism

$$U(G) \longrightarrow \mathrm{Dif}_{G/S}$$

of  $\Gamma(\mathcal{O}_S)$ -algebras, called *right translation*.<sup>19</sup>

If  $\mathcal{D}\{_{G/S}$  denotes the sheaf of  $S$ -differential operators on  $G$ , as in 1.4, and  $p : G \rightarrow S$  is the structural morphism, one likewise defines a right translation

$$\mathcal{U}(G) \longrightarrow p_*(\mathcal{D})\{_{G/S}.$$

**2.3.** We now characterize the differential operators on  $G/S$  of the form  $d^G$ . Let  $g : S \rightarrow G$  be a section of the structural morphism and let  $g_G$  denote right translation by  $g$ , namely

$$g_G : G \simeq G \times S \xrightarrow{G \times g} G \times G \xrightarrow{m} G.$$

For every differential operator  $D$  on  $G/S$ , the composite

$$g_G^{-1} D g_G$$

is again an  $S$ -deviation of  $\mathrm{id}_G$ , hence an element of  $\mathrm{Dif}_{G/S}$ . We write

$$D^g = g_G^{-1} D g_G.$$

We call  $D$  *right-invariant* if for every base change  $t : T \rightarrow S$  and every section  $g : T \rightarrow G \times T$ , one has

$$(D_T)^g = D_T.$$

**Lemma.** *For every differential operator  $D$  on  $G/S$ , the following assertions are equivalent, where  $m$  is the multiplication morphism of  $G$ :*

- (i)  $D$  is right-invariant.
- (ii) The following two deviations of  $m$  are equal:

$$Dm = m(D \times G).$$

(ii)  $\Rightarrow$  (i). Since condition (ii) is stable under base change, it suffices to show that it implies  $D^g = D$  for every section  $g : S \rightarrow G$ . Let

$$h = G \times g : G \simeq G \times S \longrightarrow G \times G,$$

so that  $m \circ h = g_G$ . The equality  $D^g = D$  is equivalent to  $g_G \circ D = D \circ g_G$ , and this follows from the commutative diagram

<sup>18</sup>The editors corrected the original diagram by replacing  $d \times G \times G$  with  $G \times d \times G$ . The composite on the left side of the upper triangle is then  $(e \times d) \times G$ , and  $d \mapsto d^G$  becomes an anti-isomorphism from  $U(G)$  to the right-invariant differential operators (2.3–2.4). If instead  ${}^G d$  is defined as the image of  $G \times d$  under  $m$ , one similarly obtains an isomorphism onto the left-invariant differential operators. Sections 2.4 and 2.5 have been corrected accordingly.

<sup>19</sup>It would be preferable to call this a *left action*. For example, if  $d$  is an  $S$ -derivation of the origin, then by 1.2.1 it is the composite of  $(\tau, \partial_t) : S \rightarrow I_S$  with a morphism  $x : I_S \rightarrow G$  satisfying  $x \circ \tau = \varepsilon$ . The derivation  $d^G$  of  $\mathcal{O}_G$  sends a local section  $\varphi$  to the section  $g \mapsto \partial_t \varphi(xg)$ . With this terminology one could say that the left action commutes with right translations.

$$\begin{array}{ccccc}
G & \xleftarrow{m} & G \times G & \xleftarrow{h} & G. \\
D \downarrow \text{id}_G & & D \times G \downarrow \text{id}_{(G \times G)} & & D \downarrow \text{id}_G \\
G & \xleftarrow{m} & G \times G & \xleftarrow{h} & G
\end{array}$$

(i)  $\Rightarrow$  (ii). Take  $t : T \rightarrow S$  to be the structural morphism  $p : G \rightarrow S$ , and take the section  $g : T \rightarrow G \times T$  to be the diagonal  $\Delta : G \rightarrow G \times G$ . Right translation

$$\Delta_{G \times G} : G \times G \longrightarrow G \times G$$

then has components  $m$  and  $\text{pr}_2$ . The equality  $(D_G)^\Delta = D_G$  is equivalent to commutativity of the first square in

$$\begin{array}{ccccc}
G \times G & \xrightarrow{\Delta_{G \times G}} & G \times G & \xrightarrow{\text{pr}_1} & G. \\
D_G \downarrow \text{id}_{G \times G} & & D_G \downarrow \text{id}_{G \times G} & & D \downarrow \text{id}_G \\
G \times G & \xrightarrow{\Delta_{G \times G}} & G \times G & \xrightarrow{\text{pr}_1} & G
\end{array}$$

because  $m = \text{pr}_1 \circ \Delta_{G \times G}$ . This proves (ii).

Now consider  $d \in U(G)$ . Every square in the following diagram is commutative:

$$\begin{array}{ccccccc}
G \times G = S \times G \times G & \xrightarrow[\varepsilon \times G \times G]{d \times G \times G} & G \times G \times G & \xrightarrow{m \times G} & G \times G \\
m \downarrow & & S \times m \downarrow & & G \times m \downarrow & & m \downarrow \\
G & \xrightarrow[\varepsilon \times G]{d \times G} & G \times G & \xrightarrow{m} & G
\end{array}$$

Since

$$m \circ (d \times G) = d^G \quad \text{and} \quad (m \times G) \circ (d \times G \times G) = d^G \times G,$$

one also has

$$d^G \circ m = m \circ (d^G \times G).$$

Thus, for every  $S$ -deviation  $d$  of the origin,  $d^G$  is a right-invariant differential operator.

**Theorem 2.4.**

- (i) The map  $d \mapsto d^G$  is an anti-isomorphism of the infinitesimal algebra  $U(G)$  onto the subalgebra  $\text{Dif}_{G/S}^G$  of  $\text{Dif}_{G/S}$  formed by the right-invariant differential operators.
- (ii) Likewise, the map  $d \mapsto {}^G d$  is an isomorphism from  $U(G)$  onto the subalgebra of  $\text{Dif}_{G/S}$  formed by the left-invariant differential operators.

20

Let  $D$  be an arbitrary differential operator on  $G/S$ , and let  $D_0$  denote its *value at the origin*, that is, the composite deviation

$$S \xrightarrow{\varepsilon} G \xrightarrow[\text{id}_G]{D} G.$$

The right-invariant differential operator  $(D_0)^G$  is obtained by the composite

$$G \simeq S \times G \xrightarrow{\varepsilon \times G} G \times G \xrightarrow[\text{id}_G \times G]{D \times G} G \times G \xrightarrow{m} G.$$

<sup>20</sup>The editors corrected “isomorphism” to “anti-isomorphism” and added assertion (ii); see the preceding editorial note.

If  $D$  is right-invariant, then  $Dm = m(D \times G)$ , and hence

$$D = Dm(\varepsilon \times G) = m(D \times G)(\varepsilon \times G) = (D_0)^G.$$

Thus  $d \mapsto d^G$  is surjective.

Conversely, for every  $S$ -deviation  $d$  of the origin one has the commutative square

$$\begin{array}{ccc} G \times G & \xleftarrow{d \times G} & G \\ \uparrow G \times \varepsilon & & \uparrow \varepsilon \\ G \times S \simeq G & \xleftarrow{d} & S \end{array}$$

from which

$$d = m(G \times \varepsilon)d = m(d \times G)\varepsilon = (d^G)_0.$$

*A fortiori*,  $d \mapsto d^G$  is injective. This proves the theorem.

As  $S$  varies, Theorem 2.4 also shows that right translation

$$\mathcal{U}(G) \longrightarrow p_*(\mathcal{D})\{_{G/S}\}$$

is an anti-isomorphism from  $\mathcal{U}(G)$  onto the sheaf of  $\mathcal{O}_S$ -algebras

$$p_*(\mathcal{D})\{_{G/S}\}^G,$$

which assigns to every open  $V \subset S$  the algebra  $\text{Dif}_{G_V/V}^{G_V}$ .

**Remark 2.4.1.** Consider the commutative diagram

$$\begin{array}{ccc} G & \xleftarrow{\eta} & G \times G \\ \uparrow p \quad \uparrow \varepsilon & & \uparrow \text{pr}_1 \quad \uparrow \Delta \\ S & \xleftarrow{p} & G, \end{array}$$

where  $\eta$  denotes the morphism  $(x, y) \mapsto yx^{-1}$ .<sup>21</sup> It induces morphisms

$$\eta' : \eta^{-1}(\mathcal{O}_G) \longrightarrow \mathcal{O}_{G \times G}$$

and

$$\Delta^{-1}(\eta') : p^{-1}\varepsilon^{-1}(\mathcal{O}_G) \longrightarrow \Delta^{-1}(\mathcal{O}_{G \times G}).$$

For every integer  $n \geq 1$ , set

$$\mathfrak{p}_{G/S}^n = \varepsilon^{-1}(\mathcal{O}_G)/\mathcal{I}_\varepsilon^{n+1}.$$

<sup>22</sup> Since the square formed by  $\varepsilon, \eta, \Delta, p$  is cartesian,  $\Delta^{-1}(\eta')$  induces an isomorphism of  $\mathcal{O}_G$ -modules

$$p^*(\mathfrak{p}_{G/S}^n) \xrightarrow{\sim} \mathcal{P}_{G/S}^n.$$

The differential operators on  $G/S$  of order  $\leq n$  therefore correspond bijectively to the homomorphisms of  $\mathcal{O}_G$ -modules

$$p^*(\mathfrak{p}_{G/S}^n) \longrightarrow \mathcal{O}_G,$$

or equivalently to the homomorphisms of  $\mathcal{O}_S$ -modules

$$\mathfrak{p}_{G/S}^n \longrightarrow p_*(\mathcal{O}_G).$$

Under this bijection the right-invariant differential operators correspond to the composites

<sup>21</sup>Thus  $G$  acts on itself on the left by right translations.

<sup>22</sup>The editors corrected the original here: the relevant cartesian square is formed by  $\varepsilon, \eta, \Delta, p$ , not by  $p, p, \eta, \text{pr}_1$ .



$$\mathfrak{p}_{G/S}^n \longrightarrow \mathcal{O}_S \xrightarrow{\text{can.}} p_*(\mathcal{O}_G).$$

This recovers the isomorphism of Theorem 2.4.

**2.5.** <sup>23</sup> Let  $\text{Lie}(G)$  be the Lie algebra of  $G$ .<sup>24</sup> We shall define a homomorphism of  $\Gamma(\mathcal{O}_S)$ -Lie algebras

$$\alpha : \text{Lie}(G) \longrightarrow U(G).$$

Let  $s : S \rightarrow I_S$  be the zero section and let  $\sigma$  be the deviation of  $s$  defined in 1.2.0. Recall from II, 4.1 that  $\text{Lie}(G)$  is the set of morphisms  $x : I_S \rightarrow G$  such that  $x \circ s = \varepsilon_G$ . The composite

$$S \xrightarrow[\sigma]{\sigma} I_S \xrightarrow{x} G$$

is an  $S$ -deviation of  $\varepsilon_G$ , hence an element of  $U(G)$ ; in the notation of 1.2 it is denoted  $\sigma x$ . Moreover, by 1.2.1 the map

$$\alpha : x \longmapsto \sigma x$$

is an isomorphism of  $\mathcal{O}_S(S)$ -modules from  $\text{Lie}(G)$  onto the submodule  $\text{Der}(\varepsilon_G) \subset U(G)$  formed by the  $S$ -derivations of  $\varepsilon_G$ . We show that  $\alpha$  is a homomorphism of Lie algebras.<sup>25</sup>

Let

$$\rho' : U(G) \longrightarrow \text{Dif}_{G/S}$$

be the algebra homomorphism which sends an  $S$ -deviation  $d$  of  $\varepsilon_G$  to the left-invariant differential operator  ${}^G d$ , as in 2.2.

Let

$$\rho : G \longrightarrow \text{Aut}_S(G)$$

be the homomorphism of group functors which sends an  $S$ -morphism  $g : T \rightarrow G$  to right translation by  $g$  on  $G_T$ , namely

$$G_T \simeq T \times_T G_T \xrightarrow{G_T \times g} G_T \times_T G_T \xrightarrow{m_T} G_T.$$

Recall also from 1.5 and II, 3.14 that

$$\text{Lie}(\text{Aut } G) = \text{Lie}(\text{Aut}_S(G)/S)(S)$$

identifies with the infinitesimal automorphisms of  $G$ , that is, the automorphisms of  $I_S \times G$  which induce the identity on  $G$ .

Since  $\rho$  is a monomorphism, so is

$$\text{Lie}(\rho) : \text{Lie}(G/S) \longrightarrow \text{Lie}(\text{Aut}_S(G)/S)$$

(see, for example, Exposé II, editorial note 50). Consequently

$$\text{Lie}(\rho) : \text{Lie}(G) \longrightarrow \text{Lie}(\text{Aut } G)$$

is injective.

On the other hand, by 1.5 the map  $\beta$ , which assigns to every infinitesimal automorphism  $u$  of  $G$  the differential operator  $D_u$ ,

<sup>23</sup>The editors reordered this paragraph, beginning with the definition of  $\alpha : \text{Lie}(G) \rightarrow U(G)$ , and corrected the original as indicated in the note to 2.2.

<sup>24</sup>In this exposé, for an  $S$ -group scheme  $G$  (respectively, an  $S$ -scheme  $X$ ),  $\text{Lie}(G)$  (respectively,  $\text{Lie}(\text{Aut } X)$ ) denotes, in the notation of Exposé II,  $\text{Lie}(G/S)(S)$  (respectively,  $\text{Lie}(\text{Aut}_S(X)/S)(S)$ ). It is a  $\Gamma(\mathcal{O}_S)$ -Lie algebra by II, 4.11 and 3.14.

<sup>25</sup>See also Exposé II, 4.11.

$$G \simeq S \times G \xrightarrow{\sigma \times G} I_S \times G \xrightarrow{u} I_S \times G \xrightarrow{\text{pr}_2} G$$

is an isomorphism from  $\text{Lie}(\text{Aut } G)$  onto the Lie subalgebra of  $\text{Dif}_{G/S}$  formed by the  $p^{-1}(\mathcal{O}_S)$ -derivations of  $\mathcal{O}_G$ .

For every  $x \in \text{Lie}(G)$ , the following commutative square determines the image of  $x$  under  $\text{Lie}(\rho)$ :

$$\begin{array}{ccc} I_S \times G & \xrightarrow{\text{Lie}(\rho)(x)} & I_S \times G \\ x \times G \downarrow & & \downarrow \text{pr}_2 \\ G \times G & \xrightarrow{m} & G. \end{array}$$

It follows from this diagram that the image of  $\text{Lie}(\rho)(x)$  under  $\beta$  is the composite deviation

$$G \simeq S \times G \xrightarrow[\substack{\sigma \times G \\ s \times G}]{} I_S \times G \xrightarrow{G \times x} G \times G \xrightarrow{m} G$$

which, by the editorial note to 2.2, is precisely

$$G(\sigma x) = \rho'(\alpha(x)).$$

Thus one obtains the commutative diagram

$$\begin{array}{ccc} \text{Lie}(G) & \xleftarrow{\text{Lie}(\rho)} & \text{Lie}(\text{Aut } G) \\ \alpha \downarrow & & \downarrow \beta \\ U(G) & \xleftarrow{\rho'} & \text{Dif}_{G/S} \end{array}$$

in which  $\text{Lie}(\rho)$ ,  $\beta$ , and  $\rho'$  are homomorphisms of Lie algebras. Since  $\rho'$  is injective,  $\alpha$  is also a homomorphism of Lie algebras. Hence:

**Proposition.** *The map  $\alpha$  is an isomorphism of  $\mathcal{O}_S(S)$ -Lie algebras from  $\text{Lie}(G)$  onto the Lie algebra of  $S$ -derivations of  $\varepsilon_G$ ; the latter is itself isomorphic, via  $\text{Lie}(\rho)$ , to the Lie algebra of left-invariant  $S$ -derivations of  $G$ .<sup>26</sup>*

### 3. Coalgebras and Cartier Duality

**3.1.** Let  $S$  be a scheme (or, more generally, a ringed space). An  $\mathcal{O}_S$ -coalgebra<sup>27</sup> is a pair  $(\mathcal{U}, \Delta_{\mathcal{U}})$  consisting of an  $\mathcal{O}_S$ -module  $\mathcal{U}$  and a morphism of  $\mathcal{O}_S$ -modules

$$\Delta_{\mathcal{U}} : \mathcal{U} \longrightarrow \mathcal{U} \otimes_{\mathcal{O}_S} \mathcal{U},$$

called the *diagonal morphism* or *comultiplication*, such that:

(i)  $\sigma \circ \Delta_{\mathcal{U}} = \Delta_{\mathcal{U}}$ , where  $\sigma(a \otimes b) = b \otimes a$ .

(ii) The square

<sup>26</sup>There exist Lie algebras  $\mathfrak{g}$  over a ring  $A$  for which the canonical map  $\mathfrak{g} \rightarrow U(\mathfrak{g})$  is not injective; see Bourbaki, *Lie Groups and Lie Algebras*, I.2, Exercise 9. The result above shows, since  $\alpha$  factors through

$$\text{Lie}(G) \longrightarrow U(\text{Lie}(G)) \longrightarrow U(G),$$

that this cannot happen for “algebraic” Lie algebras, namely those of the form  $\text{Lie}(G)$  for an  $A$ -group scheme  $G$ .

<sup>27</sup>The term *cogebra* is also used; see Bourbaki, *Algebra*, III, §11.1. Notice also that in this exposé, as well as in Exposé VII<sub>B</sub>, we work in the category of cocommutative coalgebras (that is, coalgebras satisfying condition (i)). This is crucial for defining the product and the notion of a coalgebra in groups; see 3.1.0 and 3.2.

$$\begin{array}{ccc}
\mathcal{U} & \xrightarrow{\Delta_{\mathcal{U}}} & \mathcal{U} \otimes_{\mathcal{O}_S} \mathcal{U} \\
\Delta_{\mathcal{U}} \downarrow & & \downarrow \text{id}_{\mathcal{U}} \otimes \Delta_{\mathcal{U}} \\
\mathcal{U} \otimes_{\mathcal{O}_S} \mathcal{U} & \xrightarrow{\Delta_{\mathcal{U}} \otimes \text{id}_{\mathcal{U}}} & \mathcal{U} \otimes_{\mathcal{O}_S} \mathcal{U} \otimes_{\mathcal{O}_S} \mathcal{U}
\end{array}$$

is commutative.

(iii) There exists a morphism of  $\mathcal{O}_S$ -modules

$$\varepsilon_{\mathcal{U}} : \mathcal{U} \longrightarrow \mathcal{O}_S,$$

called the *augmentation*, such that the two composites

$$\begin{aligned}
\mathcal{U} &\xrightarrow{\Delta_{\mathcal{U}}} \mathcal{U} \otimes_{\mathcal{O}_S} \mathcal{U} \xrightarrow{\text{id}_{\mathcal{U}} \otimes \varepsilon_{\mathcal{U}}} \mathcal{U} \otimes_{\mathcal{O}_S} \mathcal{O}_S \simeq \mathcal{U} \\
\mathcal{U} &\xrightarrow{\Delta_{\mathcal{U}}} \mathcal{U} \otimes_{\mathcal{O}_S} \mathcal{U} \xrightarrow{\varepsilon_{\mathcal{U}} \otimes \text{id}_{\mathcal{U}}} \mathcal{O}_S \otimes_{\mathcal{O}_S} \mathcal{U} \simeq \mathcal{U}
\end{aligned}$$

are the identity morphism of  $\mathcal{U}$ .

If  $\varepsilon_{\mathcal{U}}$  and  $\varepsilon'_{\mathcal{U}}$  are two augmentations, then

$$\varepsilon_{\mathcal{U}} = (\varepsilon_{\mathcal{U}} \otimes \varepsilon'_{\mathcal{U}}) \circ \Delta_{\mathcal{U}} = \varepsilon'_{\mathcal{U}};$$

the augmentation is therefore uniquely determined by (iii).

If  $(\mathcal{U}, \Delta_{\mathcal{U}})$  and  $(\mathcal{V}, \Delta_{\mathcal{V}})$  are two  $\mathcal{O}_S$ -coalgebras, a *morphism* from the first to the second is a morphism of  $\mathcal{O}_S$ -modules  $f : \mathcal{U} \rightarrow \mathcal{V}$  for which the diagrams

$$\begin{array}{ccc}
\mathcal{U} & \xrightarrow{f} & \mathcal{V} \\
\Delta_{\mathcal{U}} \downarrow & & \downarrow \Delta_{\mathcal{V}} \\
\mathcal{U} \otimes \mathcal{U} & \xrightarrow{f \otimes f} & \mathcal{V} \otimes \mathcal{V}
\end{array}
\quad \text{and} \quad
\begin{array}{ccc}
\mathcal{U} & \xrightarrow{f} & \mathcal{V} \\
\varepsilon_{\mathcal{U}} \searrow & & \swarrow \varepsilon_{\mathcal{V}} \\
& \mathcal{O}_S &
\end{array}$$

are commutative. Morphisms of coalgebras compose like morphisms of  $\mathcal{O}_S$ -modules, so we may speak of the *category of  $\mathcal{O}_S$ -coalgebras*.

**3.1.0.** <sup>28</sup> This category has finite products: its final object is the  $\mathcal{O}_S$ -module  $\mathcal{O}_S$ , with the identity as comultiplication. The product of two coalgebras  $(\mathcal{U}, \Delta_{\mathcal{U}})$  and  $(\mathcal{V}, \Delta_{\mathcal{V}})$  is the tensor product  $\mathcal{U} \otimes_{\mathcal{O}_S} \mathcal{V}$ , whose comultiplication is the composite

$$\mathcal{U} \otimes \mathcal{V} \xrightarrow{\Delta_{\mathcal{U}} \otimes \Delta_{\mathcal{V}}} \mathcal{U} \otimes \mathcal{U} \otimes \mathcal{V} \otimes \mathcal{V} \xrightarrow{\text{id}_{\mathcal{U}} \otimes \sigma \otimes \text{id}_{\mathcal{V}}} \mathcal{U} \otimes \mathcal{V} \otimes \mathcal{U} \otimes \mathcal{V}$$

where  $\sigma(a \otimes b) = b \otimes a$ . The canonical projections of  $\mathcal{U} \otimes \mathcal{V}$  onto the factors  $\mathcal{U}$  and  $\mathcal{V}$  are  $\text{id}_{\mathcal{U}} \otimes \varepsilon_{\mathcal{V}}$  and  $\varepsilon_{\mathcal{U}} \otimes \text{id}_{\mathcal{V}}$ , respectively.<sup>29</sup> The “diagonal morphism”  $\mathcal{U} \rightarrow \mathcal{U} \otimes \mathcal{U}$ , corresponding to the pair  $(\text{id}_{\mathcal{U}}, \text{id}_{\mathcal{U}})$ , is precisely the comultiplication  $\Delta_{\mathcal{U}}$ .

**3.1.1.** Let  $\mathcal{A}$  be a commutative  $\mathcal{O}_S$ -algebra which, as an  $\mathcal{O}_S$ -module, is locally free and of finite type. Put

$$\mathcal{A}^* = \mathcal{H}om_{\mathcal{O}_S\text{-Mod.}}(\mathcal{A}, \mathcal{O}_S).$$

<sup>28</sup>The editors added the numbering 3.1.0 for later references.

<sup>29</sup>The preceding assertion was added by the editors. Recall also that, to show that  $\mathcal{U} \otimes \mathcal{V}$  is indeed the product of  $\mathcal{U}$  and  $\mathcal{V}$  in the category of cocommutative  $\mathcal{O}_S$ -cogbras, one argues as follows. Given an arbitrary  $\mathcal{O}_S$ -cogebra  $\mathcal{E}$  and cogebra morphisms  $f : \mathcal{E} \rightarrow \mathcal{U}$  and  $g : \mathcal{E} \rightarrow \mathcal{V}$ , any cogebra morphism  $\varphi : \mathcal{E} \rightarrow \mathcal{U} \otimes \mathcal{V}$  satisfying  $\text{pr}_{\mathcal{U}} \circ \varphi = f$  and  $\text{pr}_{\mathcal{V}} \circ \varphi = g$  must equal  $(f \otimes g) \circ \Delta_{\mathcal{E}}$ ; and this is a cogebra morphism if and only if it also equals  $(g \otimes f) \circ \Delta_{\mathcal{E}}$ .

The canonical morphism

$$\varphi : \mathcal{A}^* \otimes_{\mathcal{O}_S} \mathcal{A}^* \longrightarrow (\mathcal{A} \otimes_{\mathcal{O}_S} \mathcal{A})^*$$

is invertible. If  $m : \mathcal{A} \otimes \mathcal{A} \rightarrow \mathcal{A}$  is the morphism defining the multiplication of  $\mathcal{A}$ , composition gives a diagonal morphism

$$\Delta_{\mathcal{A}^*} : \mathcal{A}^* \xrightarrow{m^*} (\mathcal{A} \otimes \mathcal{A})^* \xrightarrow{\varphi^{-1}} \mathcal{A}^* \otimes \mathcal{A}^*.$$

This diagonal morphism plainly makes  $\mathcal{A}^*$  into an  $\mathcal{O}_S$ -coalgebra whose augmentation is the transpose of the morphism  $\mathcal{O}_S \rightarrow \mathcal{A}$  defined by the unit section of  $\mathcal{A}$ . Moreover, it is clear that:

*The functor  $\mathcal{A} \mapsto \mathcal{A}^*$  is an anti-equivalence from the category of  $\mathcal{O}_S$ -algebras which are locally free and of finite type as  $\mathcal{O}_S$ -modules onto the category of  $\mathcal{O}_S$ -coalgebras which are locally free and of finite type as  $\mathcal{O}_S$ -modules.*

**3.1.2.** To every  $\mathcal{O}_S$ -coalgebra  $\mathcal{U}$  there is canonically associated an  $S$ -functor

$$\mathrm{Spec}^* \mathcal{U} : (\mathbf{Sch}/S)^\circ \longrightarrow (\mathbf{Ens}).$$

Indeed, for every  $S$ -scheme  $q : T \rightarrow S$ ,

$$q^*(\mathcal{U} \otimes_{\mathcal{O}_S} \mathcal{U}) \simeq q^*(\mathcal{U}) \otimes_{\mathcal{O}_T} q^*(\mathcal{U}),$$

so  $q^*(\Delta_{\mathcal{U}})$  makes  $\mathcal{U}_T = q^*(\mathcal{U})$  into an  $\mathcal{O}_T$ -coalgebra. Thus, by definition and with an obvious abuse of notation,<sup>30</sup>

$$(\mathrm{Spec}^* \mathcal{U})(T) = \{x \in \Gamma(T, \mathcal{U}_T) \mid \varepsilon_{\mathcal{U}_T}(x) = 1, \quad \Delta_{\mathcal{U}_T}(x) = x \otimes x\}.$$

Sections  $x$  of  $\mathcal{U}_T$  plainly correspond to morphisms of  $\mathcal{O}_T$ -modules  $\xi : \mathcal{O}_T \rightarrow \mathcal{U}_T$ . The conditions  $\varepsilon(x) = 1$  and  $\Delta(x) = x \otimes x$  say exactly that  $\xi$  is a morphism of coalgebras. Hence also

$$(\mathrm{Spec}^* \mathcal{U})(T) = \mathrm{Hom}_{\mathcal{O}_T\text{-coalg.}}(\mathcal{O}_T, \mathcal{U}_T).$$

In particular:

**Proposition 3.1.2.1.** <sup>31</sup> *Let  $\mathcal{A}$  be a commutative  $\mathcal{O}_S$ -algebra which is locally free of finite type as an  $\mathcal{O}_S$ -module. Then the  $S$ -functor  $\mathrm{Spec}^* \mathcal{A}^*$  is represented by  $\mathrm{Spec} \mathcal{A}$ .*

Indeed, for every  $S$ -scheme  $T$ , there are canonical isomorphisms

$$\begin{aligned} (\mathrm{Spec}^* \mathcal{A}^*)(T) &= \mathrm{Hom}_{\mathcal{O}_T\text{-coalg.}}(\mathcal{O}_T, \mathcal{A}_T^*) \\ &\simeq \mathrm{Hom}_{\mathcal{O}_T\text{-alg.}}(\mathcal{A}_T, \mathcal{O}_T) \simeq (\mathrm{Spec} \mathcal{A})(T). \end{aligned}$$

**3.2.** An  $\mathcal{O}_S$ -coalgebra in groups, that is, a group in the category of  $\mathcal{O}_S$ -coalgebras, consists of an  $\mathcal{O}_S$ -coalgebra  $(\mathcal{U}, \Delta_{\mathcal{U}})$  together with three morphisms of  $\mathcal{O}_S$ -coalgebras

$$m_{\mathcal{U}} : \mathcal{U} \otimes \mathcal{U} \longrightarrow \mathcal{U}, \quad \eta_{\mathcal{U}} : \mathcal{O}_S \longrightarrow \mathcal{U}, \quad c_{\mathcal{U}} : \mathcal{U} \longrightarrow \mathcal{U},$$

satisfying conditions (ii)\*, (iii)\*, and (vi) below. The requirement that  $m_{\mathcal{U}}$  be a morphism of coalgebras is expressed by the commutativity of diagrams (iv) and (v):

<sup>30</sup>For  $x \otimes y \in \Gamma(T, \mathcal{U}_T) \otimes_{\mathcal{O}(T)} \Gamma(T, \mathcal{U}_T)$ , its image in  $\Gamma(T, \mathcal{U}_T \otimes_{\mathcal{O}_T} \mathcal{U}_T)$  is again denoted  $x \otimes y$ .

<sup>31</sup>The editors added the numbering 3.1.2.1 for later references. Notice also that the  $S$ -functor  $\mathrm{Spec}^* \mathcal{U}$  is a sheaf for the Zariski topology; it is even an fpqc sheaf if  $\mathcal{U}$  is a quasi-coherent  $\mathcal{O}_S$ -module.

$$\begin{array}{ccc}
\text{(iv)} & \mathcal{U} \otimes \mathcal{U} & \xrightarrow{m_{\mathcal{U}}} \mathcal{U} \\
& \Delta_{\mathcal{U}} \otimes \Delta_{\mathcal{U}} \downarrow & \downarrow \Delta_{\mathcal{U}} \\
& \mathcal{U} \otimes \mathcal{U} \otimes \mathcal{U} \otimes \mathcal{U} & \\
& \text{id}_{\mathcal{U}} \otimes \sigma \otimes \text{id}_{\mathcal{U}} \downarrow & \\
& \mathcal{U} \otimes \mathcal{U} \otimes \mathcal{U} \otimes \mathcal{U} & \xrightarrow{m_{\mathcal{U}} \otimes m_{\mathcal{U}}} \mathcal{U} \otimes \mathcal{U}
\end{array}$$

$$\begin{array}{ccc}
\text{(v)} & \mathcal{U} \otimes \mathcal{U} & \xrightarrow{m_{\mathcal{U}}} \mathcal{U} \\
& \searrow \varepsilon_{\mathcal{U}} \otimes \varepsilon_{\mathcal{U}} & \swarrow \varepsilon_{\mathcal{U}} \\
& \mathcal{O}_S &
\end{array}$$

(ii)\* The square

$$\begin{array}{ccc}
\mathcal{U} \otimes \mathcal{U} \otimes \mathcal{U} & \xrightarrow{\text{id}_{\mathcal{U}} \otimes m_{\mathcal{U}}} & \mathcal{U} \otimes \mathcal{U} \\
m_{\mathcal{U}} \otimes \text{id}_{\mathcal{U}} \downarrow & & \downarrow m_{\mathcal{U}} \\
\mathcal{U} \otimes \mathcal{U} & \xrightarrow{m_{\mathcal{U}}} & \mathcal{U}
\end{array}$$

is commutative.

(iii)\* The two composites

$$\mathcal{U} \simeq \mathcal{U} \otimes \mathcal{O}_S \xrightarrow{\text{id}_{\mathcal{U}} \otimes \eta_{\mathcal{U}}} \mathcal{U} \otimes \mathcal{U} \xrightarrow{m_{\mathcal{U}}} \mathcal{U},$$

$$\mathcal{U} \simeq \mathcal{O}_S \otimes \mathcal{U} \xrightarrow{\eta_{\mathcal{U}} \otimes \text{id}_{\mathcal{U}}} \mathcal{U} \otimes \mathcal{U} \xrightarrow{m_{\mathcal{U}}} \mathcal{U}.$$

are the identity morphism of  $\mathcal{U}$ .

(vi) The composite

$$\mathcal{U} \xrightarrow{\Delta_{\mathcal{U}}} \mathcal{U} \otimes \mathcal{U} \xrightarrow{c_{\mathcal{U}} \otimes \text{id}_{\mathcal{U}}} \mathcal{U} \otimes \mathcal{U} \xrightarrow{m_{\mathcal{U}}} \mathcal{U}.$$

is equal to  $\eta_{\mathcal{U}} \circ \varepsilon_{\mathcal{U}}$ .

**3.2.1.** The morphisms  $\eta_{\mathcal{U}}$  and  $c_{\mathcal{U}}$  are uniquely determined by  $m_{\mathcal{U}}$ . Conditions (ii)\* and (iii)\* simply say that  $m_{\mathcal{U}}$  makes  $\mathcal{U}$  into an  $\mathcal{O}_S$ -algebra whose unit section is the image under  $\eta_{\mathcal{U}}$  of the unit section of  $\mathcal{O}_S$ . Condition (iv) also says that the diagonal morphism  $\Delta_{\mathcal{U}}$  is compatible with multiplication. Indeed,

$$\Delta_{\mathcal{U}} : \mathcal{U} \longrightarrow \mathcal{U} \otimes \mathcal{U}$$

must be a homomorphism of coalgebras in groups, which also implies the commutativity of the triangle

$$\begin{array}{ccc}
& \mathcal{O}_S & \\
\eta_{\mathcal{U}} \swarrow & & \searrow \eta_{\mathcal{U}} \otimes \eta_{\mathcal{U}} \\
\mathcal{U} & \xrightarrow{\Delta_{\mathcal{U}}} & \mathcal{U} \otimes \mathcal{U}
\end{array}$$

On the other hand, as in every category, the antipode  $c_{\mathcal{U}}$  is an isomorphism from  $\mathcal{U}$  onto the “opposite” group object;<sup>32</sup> in particular,  $c_{\mathcal{U}}$  induces an algebra isomorphism from  $\mathcal{U}$  onto the opposite algebra  $\mathcal{U}^{\circ}$ .

<sup>32</sup>That is, the object endowed with the multiplication  $m'_{\mathcal{U}} = m_{\mathcal{U}} \circ \sigma$ .

**3.2.2.** Since the functor  $\mathcal{U} \mapsto \mathrm{Spec}^* \mathcal{U}$  commutes with finite products, it carries a coalgebra in groups to a group-valued  $S$ -functor. Indeed, for every  $S$ -scheme  $T$ , the elements

$$x \in \Gamma(T, \mathcal{U}_T)$$

belonging to  $(\mathrm{Spec}^* \mathcal{U})(T)$  form a group under the multiplication of the algebra  $\Gamma(T, \mathcal{U}_T)$ , and the inverse of  $x$  is  $c_{\mathcal{U}}(x)$ . By 3.1.2.1:

**Scholium 3.2.2.1.** <sup>33</sup> *Let  $\mathcal{U}$  be an  $\mathcal{O}_S$ -coalgebra in groups, finite and locally free as an  $\mathcal{O}_S$ -module. Then the group-valued  $S$ -functor  $\mathrm{Spec}^* \mathcal{U}$  is represented by the finite locally free  $S$ -group  $\mathrm{Spec} \mathcal{U}^*$ .*

**Remark 3.2.2.2.** Let  $\mathcal{L}$  be an  $\mathcal{O}_S$ -Lie algebra and  $\mathcal{U}(\mathcal{L})$  its enveloping algebra, that is, the sheaf on  $S$  associated with the presheaf assigning to every open subset  $V$  the enveloping algebra

$$U(\Gamma(V, \mathcal{L}))$$

of the Lie algebra  $\Gamma(V, \mathcal{L})$ .

Every homomorphism from  $\mathcal{L}$  to the Lie algebra underlying an associative  $\mathcal{O}_S$ -algebra factors uniquely through the canonical morphism  $\mathcal{L} \rightarrow \mathcal{U}(\mathcal{L})$ . Besides the functoriality of  $\mathcal{U}(\mathcal{L})$  in  $\mathcal{L}$ , this universal property also shows that the enveloping algebra of a product of Lie algebras is identified with the tensor product of their enveloping algebras.

In particular, the diagonal morphism  $\delta : \mathcal{L} \rightarrow \mathcal{L} \times \mathcal{L}$  induces an algebra homomorphism

$$\Delta : \mathcal{U}(\mathcal{L}) \longrightarrow \mathcal{U}(\mathcal{L} \times \mathcal{L}) \simeq \mathcal{U}(\mathcal{L}) \otimes \mathcal{U}(\mathcal{L}).$$

The zero morphism  $\mathcal{L} \rightarrow 0$  induces a homomorphism

$$\varepsilon : \mathcal{U}(\mathcal{L}) \longrightarrow \mathcal{U}(0) \simeq \mathcal{O}_S.$$

The isomorphism  $x \mapsto -x$  from  $\mathcal{L}$  onto the opposite Lie algebra  $\mathcal{L}^\circ$  induces an anti-isomorphism  $c$  of the algebra  $\mathcal{U}(\mathcal{L})$ . One then checks readily that the multiplication  $m$  of  $\mathcal{U}(\mathcal{L})$  makes  $(\mathcal{U}(\mathcal{L}), \Delta)$  into an  $\mathcal{O}_S$ -coalgebra in groups with augmentation  $\varepsilon$  and antipode  $c$ .<sup>34</sup>

**3.2.3.** <sup>35</sup> Let  $\mathcal{U}$  be an  $\mathcal{O}_S$ -coalgebra in groups. We shall see that the group-valued  $S$ -functor

$$G = \mathrm{Spec}^* \mathcal{U}$$

is *very good*, in the sense of Exposé II, 4.6 and 4.10.

Let  $\mathcal{M}$  be a free  $\mathcal{O}_S$ -module of rank  $r$ , and let  $T \rightarrow S$  be an  $S$ -scheme. Since

$$I_T(\mathcal{M}) = \mathrm{Spec}(\mathcal{O}_T \oplus \mathcal{M}_T),$$

the morphism  $\pi : I_T(\mathcal{M}) \rightarrow T$  is affine, and

$$\pi_*(\mathcal{U}_{I_T(\mathcal{M})}) = \mathcal{U}_T \otimes_{\mathcal{O}_T} \pi_*(\mathcal{O}_{I_T(\mathcal{M})}) = \mathcal{U}_T \otimes_{\mathcal{O}_T} (\mathcal{O}_T \oplus \mathcal{M}_T).$$

Consequently,

$$\Gamma(I_T(\mathcal{M}), \mathcal{U}_{I_T(\mathcal{M})}) \simeq \Gamma(T, \mathcal{U}_T) \otimes_{\mathcal{O}(T)} (\mathcal{O}(T) \oplus \Gamma(T, \mathcal{M}_T)). \quad (1)$$

<sup>33</sup>The editors added this scholium, which is implicit in the original.

<sup>34</sup>The group-valued  $S$ -functor  $\mathrm{Spec}^* \mathcal{U}(\mathcal{L})$  is not representable in general. Later, in 5.5, it will be shown that if  $S$  is a scheme of characteristic  $p$ , if  $\mathcal{L}$  is finite locally free over  $\mathcal{O}_S$ , and if  $\mathcal{U}_p(\mathcal{L})$  is its restricted enveloping algebra (see 5.3), then  $\mathrm{Spec}^* \mathcal{U}_p(\mathcal{L})$  is represented by a finite locally free  $S$ -group.

<sup>35</sup>The editors expanded this paragraph.

Let  $(d_1, \dots, d_r)$  be a basis of  $\mathcal{M}$ . An element

$$u_0 + \sum_i u_i d_i \in \Gamma(I_T(\mathcal{M}), \mathcal{U}_{I_T(\mathcal{M})})$$

belongs to  $G(I_T(\mathcal{M}))$  if and only if

$$1 = \varepsilon \left( u_0 + \sum_i u_i d_i \right) = \varepsilon(u_0) + \sum_i \varepsilon(u_i) d_i$$

and

$$\left( u_0 + \sum_i u_i d_i \right) \otimes \left( u_0 + \sum_i u_i d_i \right) = \Delta \left( u_0 + \sum_i u_i d_i \right) = \Delta(u_0) + \sum_i \Delta(u_i) d_i.$$

Equivalently,

$$\begin{cases} \varepsilon(u_0) = 1, & \Delta u_0 = u_0 \otimes u_0 & (\text{that is, } u_0 \in G(T)), \\ \varepsilon(u_i) = 0, & \Delta(u_i) = u_i \otimes u_0 + u_0 \otimes u_i & (i = 1, \dots, r). \end{cases} \quad (2)$$

Moreover, the morphism  $G(I_T(\mathcal{M})) \rightarrow G(T)$  corresponding to the zero section of  $I_T(\mathcal{M}) \rightarrow T$  is given by

$$u_0 + \sum_i u_i d_i \mapsto u_0.$$

Combining this with (1) and (2), one sees that if  $\mathcal{N}$  is a second free  $\mathcal{O}_S$ -module of finite rank, then the diagram of sets

$$\begin{array}{ccc} G(I_T(\mathcal{M} \oplus \mathcal{N})) & \longrightarrow & G(I_T(\mathcal{N})) \\ \downarrow & & \downarrow \\ G(I_T(\mathcal{M})) & \longrightarrow & G(T) \end{array}$$

is cartesian; that is,  $G$  satisfies condition (E) of Exposé II, 3.5.

Write

$$\text{Prim } \Gamma(T, \mathcal{U}_T)$$

for the  $\mathcal{O}(T)$ -submodule of  $\Gamma(T, \mathcal{U}_T)$  formed by the *primitive elements*, that is, the elements  $u$  satisfying, with the abuse of notation noted in 3.1.2,

$$\Delta u = u \otimes 1 + 1 \otimes u, \quad \varepsilon(u) = 0.$$

36

Since  $(\text{Lie } G)(T)$  is the set of elements  $u_0 + ud \in G(I_T)$  above the unit element  $u_0 = 1$  of  $G(T)$ , there is an isomorphism of  $\mathcal{O}(T)$ -modules, functorial in  $T$ ,<sup>37</sup>

$$(\text{Lie } G)(T) \simeq \text{Prim } \Gamma(T, \mathcal{U}_T).$$

On the other hand, (1) yields

$$\text{Prim } \Gamma(I_T(\mathcal{M}), \mathcal{U}_{I_T(\mathcal{M})}) \simeq \text{Prim } \Gamma(T, \mathcal{U}_T) \otimes_{\mathcal{O}(T)} \mathcal{O}(I_T(\mathcal{M})).$$

<sup>36</sup>The second condition follows from the first, since the first gives  $u = (\text{id} \otimes \varepsilon)\Delta(u) = u + \varepsilon(u)$ , and hence  $\varepsilon(u) = 0$ .

<sup>37</sup>The  $\mathcal{O}_S$ -module structure on  $\text{Lie } G$  is defined in Exposé II, Proposition 3.6.

It follows that the natural morphism of  $\mathcal{O}(I_T(\mathcal{M}))$ -modules

$$(\mathrm{Lie} G)(T) \otimes_{\mathcal{O}(T)} \mathcal{O}(I_T(\mathcal{M})) \longrightarrow (\mathrm{Lie} G)(I_T(\mathcal{M}))$$

is an isomorphism; thus  $\mathrm{Lie} G$  is a good  $\mathcal{O}_S$ -module (see Exposé II, Definition 4.4).

Therefore  $G$  is a good group-valued  $S$ -functor (see Exposé II, Definition 4.6), and by Exposé II, 4.7.2,  $\mathrm{Lie} G$  carries an  $\mathcal{O}_S$ -bilinear Lie bracket satisfying the Jacobi identity. It remains to show that  $G$  is very good, that is, that the bracket on  $(\mathrm{Lie} G)(T)$  satisfies  $[u, u] = 0$  for every  $u \in (\mathrm{Lie} G)(T)$ ; see Exposé II, 4.10.

Let  $u, v$  be two elements of  $(\mathrm{Lie} G)(T)$ , that is, two primitive elements of  $\Gamma(T, \mathcal{U}_T)$ . Put

$$I = \mathrm{Spec} \mathcal{O}_S[d]/(d^2), \quad I' = \mathrm{Spec} \mathcal{O}_S[d']/(d'^2).$$

Since the composition law on  $G(I \times I')$  is induced by the multiplication in  $\mathcal{U}_{I \times I'}$ , one has in  $G(I \times I')$

$$\begin{aligned} (1 + ud)(1 + vd')(1 + ud)^{-1}(1 + vd')^{-1} \\ = (1 + ud)(1 + vd')(1 - ud)(1 - vd') \\ = 1 + (uv - vu)dd'. \end{aligned}$$

*Source note.* The re-edition prints  $1 - vd$ , without a prime, in the last factor of the middle expression. Since that factor is the inverse of  $1 + vd'$ , and since the displayed result is proportional to  $dd'$ , the English reconstruction makes the required reading  $1 - vd'$  explicit. Reinhold's comparison Markdown reproduces the printed missing prime.

By the description of the bracket  $[u, v]$  given before Proposition 4.8 of Exposé II, it follows that

$$[u, v] = uv - vu,$$

where the right-hand side is the commutator of  $u$  and  $v$  in the algebra  $\Gamma(T, \mathcal{U}_T)$ . Hence  $[u, u] = 0$ , and we have proved:

**Proposition.** *Let  $\mathcal{U}$  be an  $\mathcal{O}_S$ -coalgebra in groups. The group-valued  $S$ -functor  $G = \mathrm{Spec}^* \mathcal{U}$  is very good, and there is an isomorphism*

$$\mathrm{Lie} G \simeq \mathrm{Prim} \mathcal{W}(\mathcal{U})$$

*of  $\mathcal{O}_S$ -Lie algebras, where  $\mathrm{Prim} \mathcal{W}(\mathcal{U})$  denotes the functor which assigns to every  $T \rightarrow S$  the  $\mathcal{O}(T)$ -Lie algebra of primitive elements of  $\mathcal{W}(\mathcal{U})(T) = \Gamma(T, \mathcal{U}_T)$ .*<sup>38</sup>

**3.3.** Suppose finally that  $\mathcal{U}$  is a coalgebra in commutative groups, that is, that the triangle

$$(i)^* \quad \begin{array}{ccc} \mathcal{U} \otimes \mathcal{U} & \xrightarrow{\sigma} & \mathcal{U} \otimes \mathcal{U} \\ & \searrow m_{\mathcal{U}} & \swarrow m_{\mathcal{U}} \\ & \mathcal{U} & \end{array}$$

is commutative, or equivalently that  $m_{\mathcal{U}}$  makes  $\mathcal{U}$  into a commutative  $\mathcal{O}_S$ -algebra. Conditions (i), (ii), (iii), (iv), (v), (vi), (i)\*, (ii)\*, (iii)\*, and (v)\* then also say that  $\mathcal{U}$  is a cogroup in the category of commutative  $\mathcal{O}_S$ -algebras. Thus, if in addition  $\mathcal{U}$  is a quasi-coherent  $\mathcal{O}_S$ -module, the affine  $S$ -scheme  $\mathrm{Spec} \mathcal{U}$  is a commutative  $S$ -group scheme.

In this case the diagonal morphism  $\Delta'$  of  $\mathcal{O}_S[T, T^{-1}]$  sends  $T$  to  $T \otimes T$ . Consequently, morphisms of  $S$ -groups from  $\mathrm{Spec} \mathcal{U}$  to  $\mathbb{G}_{m, S}$  (Exposé I, 4.3.2) correspond bijectively to morphisms of unital  $\mathcal{O}_S$ -algebras

$$\varphi : \mathcal{O}_S[T, T^{-1}] \longrightarrow \mathcal{U}$$

<sup>38</sup>The editors added this proposition to summarize the preceding discussion.



such that

$$(\varphi \otimes \varphi) \circ \Delta' = \Delta_{\mathcal{U}} \circ \varphi.$$

In this situation,  $\varepsilon_{\mathcal{U}} \circ \varphi$  is the neutral element of  $\mathbb{G}_{m,S}(S)$ , that is, the augmentation  $\varepsilon'$ . Such a morphism  $\varphi$  is determined by the image  $\varphi(T)$ , which must be an invertible element  $x \in \mathcal{U}$  satisfying

$$\Delta_{\mathcal{U}}x = x \otimes x, \quad \varepsilon_{\mathcal{U}}(x) = \varepsilon'(T) = 1.$$

Therefore

$$\mathrm{Hom}_{S\text{-gr.}}(\mathrm{Spec}\mathcal{U}, \mathbb{G}_{m,S}) \simeq (\mathrm{Spec}^*\mathcal{U})(S).$$

Since this formula remains valid after arbitrary base change, it gives

$$\mathrm{Spec}^*\mathcal{U} \simeq \mathrm{Hom}_{S\text{-gr.}}(\mathrm{Spec}\mathcal{U}, \mathbb{G}_{m,S}).$$

We have therefore proved:

**Proposition 3.3.0.** *If  $\mathcal{U}$  is an  $\mathcal{O}_S$ -coalgebra in commutative groups, quasi-coherent as an  $\mathcal{O}_S$ -module, then the affine  $S$ -scheme  $G = \mathrm{Spec}\mathcal{U}$  is a commutative  $S$ -group scheme, and there is an isomorphism of group-valued  $S$ -functors*

$$\mathrm{Spec}^*\mathcal{U} \simeq \mathrm{Hom}_{S\text{-gr.}}(G, \mathbb{G}_{m,S}).$$

If, moreover,  $\mathcal{U}$  is locally free of finite type as an  $\mathcal{O}_S$ -module, then by 3.1.2.1 the group-valued  $S$ -functor  $\mathrm{Spec}^*\mathcal{U}$  is represented by  $\mathrm{Spec}\mathcal{U}^*$ . Hence:

**Proposition 3.3.1 (Cartier duality).** *The functor*

$$\mathcal{A}(G) \longmapsto \mathcal{A}(G)^* = \mathrm{Hom}_{\mathcal{O}_S\text{-Mod.}}(\mathcal{A}(G), \mathcal{O}_S)$$

*induces a duality\* on the category of commutative finite locally free  $S$ -group schemes; it associates to  $G$  the  $S$ -group  $\mathrm{Hom}_{S\text{-gr.}}(G, \mathbb{G}_{m,S})$ .*

## 4. “Frobeniusseries”

Fix a prime number  $p$ , and let  $(\mathbf{Sch}/\mathbb{F}_p)$  be the category of schemes of characteristic  $p$ , that is, schemes over the prime field  $\mathbb{F}_p$ . Following the general conventions of this seminar, we identify  $(\mathbf{Sch}/\mathbb{F}_p)$  with a subcategory of  $(\mathbf{Sch}/\mathbb{F}_p)^\wedge$  by means of the functor  $h$  of Exposé I, 1.1. We likewise use the isomorphism

$$\mathrm{Hom}(h_X, F) \xrightarrow{\sim} F(X)$$

defined there to identify these sets whenever  $X$  is an  $\mathbb{F}_p$ -scheme and  $F$  is an object of  $(\mathbf{Sch}/\mathbb{F}_p)^\wedge$ .

**Notation 4.0.** <sup>39</sup> *If  $T$  is an  $\mathbb{F}_p$ -scheme, a  $T$ -functor is a morphism  $q : F \rightarrow T$  in  $(\mathbf{Sch}/\mathbb{F}_p)^\wedge$  with target  $T$ . For every  $T$ -scheme  $r : X \rightarrow T$ , the set of  $T$ -morphisms  $X \rightarrow F$ , that is, of  $\mathbb{F}_p$ -morphisms  $s : X \rightarrow F$  satisfying  $q \circ s = r$ , is denoted*

$$q(r), \quad q(X/T), \quad F(r), \quad \text{or} \quad F(X/T).$$

*It may also be denoted  $F(X)$  when no confusion with  $\mathrm{Hom}(h_X, F)$  can result.*

---

\*A duality of a category  $\mathcal{C}$  is a pair  $(D, \varphi)$  consisting of a contravariant functor  $D : \mathcal{C} \rightarrow \mathcal{C}$  and a natural isomorphism  $\varphi : \mathrm{Id}_{\mathcal{C}} \rightarrow DD$ , such that the isomorphisms  $\varphi D : D \rightarrow DDD$  and  $D\varphi^{-1} : DDD \rightarrow D$  are inverse to one another. The editors corrected  $D\varphi$  to  $D\varphi^{-1}$ .

<sup>39</sup>The editors added the number 4.0 for later references.

**4.1.** For every scheme  $S$  of characteristic  $p$ , let  $\mathrm{fr}(S)$ , or simply  $\mathrm{fr}$ , denote the endomorphism of  $S$  which is the identity on the underlying topological space and which sends a section  $x$  of  $\mathcal{O}_S$  over an open subset to  $x^p$ .

The assignment

$$\mathrm{fr} : S \longmapsto \mathrm{fr}(S)$$

is an endomorphism of the identity functor of  $(\mathbf{Sch}/\mathbb{F}_p)$ .<sup>40</sup> This has the following consequences. If  $E$  is an  $\mathbb{F}_p$ -functor, the assignment which sends every  $\mathbb{F}_p$ -scheme  $S$  to the endomorphism  $E(\mathrm{fr}(S))$  of  $E(S)$  is a functorial endomorphism of  $E$ , denoted  $\mathrm{fr}(E)$  or simply  $\mathrm{fr}$ . This notation is compatible with the identification of  $(\mathbf{Sch}/\mathbb{F}_p)$  with a subcategory of  $(\mathbf{Sch}/\mathbb{F}_p)^\wedge$ . Moreover,  $E \mapsto \mathrm{fr}(E)$  is itself an endomorphism of the identity functor of  $(\mathbf{Sch}/\mathbb{F}_p)^\wedge$ , again denoted  $\mathrm{fr}$ .<sup>41</sup>

For every  $\mathbb{F}_p$ -scheme  $S$  and every  $S$ -functor  $q : X \rightarrow S$ , let  $X^{(p/S)}$ , or  $X^{(p)}$ , denote the pullback of  $X$  along  $\mathrm{fr}(S)$ :

$$\begin{array}{ccc} X^{(p/S)} & \xrightarrow{\mathrm{pr}_X} & X \\ \downarrow & & \downarrow q \\ S & \xrightarrow{\mathrm{fr}(S)} & S. \end{array}$$

The commutative square

$$\begin{array}{ccc} X & \xrightarrow{\mathrm{fr}(X)} & X \\ q \downarrow & & \downarrow q \\ S & \xrightarrow{\mathrm{fr}(S)} & S \end{array}$$

induces an  $S$ -morphism

$$\mathrm{Fr}(X/S) : X \longrightarrow X^{(p/S)}$$

such that

$$\mathrm{fr}(X) = \mathrm{pr}_X \circ \mathrm{Fr}(X/S) :$$

$$\begin{array}{ccccc} X & & & & \\ & \searrow \mathrm{Fr}(X/S) & & \searrow \mathrm{fr}(X) & \\ & & X^{(p/S)} & \xrightarrow{\mathrm{pr}_X} & X \\ & \searrow q & \downarrow & & \downarrow q \\ & & S & \xrightarrow{\mathrm{fr}(S)} & S. \end{array}$$

We call  $\mathrm{Fr}(X/S)$  the *Frobenius morphism of  $X$  relative to  $S$* . Plainly the assignment

$$\mathrm{Fr} : X \longmapsto \mathrm{Fr}(X/S)$$

<sup>40</sup>Thus, for every morphism  $f : Y \rightarrow X$  of  $\mathbb{F}_p$ -schemes, the following square commutes:

$$\begin{array}{ccc} Y & \xrightarrow{f} & X \\ \mathrm{fr}(Y) \downarrow & & \downarrow \mathrm{fr}(X) \\ Y & \xrightarrow{f} & X. \end{array}$$

<sup>41</sup>The morphism  $\mathrm{fr}(X)$  is called the *absolute* Frobenius morphism of  $X$ , to distinguish it from the relative Frobenius morphism  $\mathrm{Fr}(X/S)$  introduced below.

is a functorial homomorphism.

<sup>42</sup> Let  $r : T \rightarrow S$  be an  $S$ -scheme. For every

$$\varphi \in X(r) = \operatorname{Hom}_S(T, X),$$

one has the commutative diagram

$$\begin{array}{ccccc} X & \xrightarrow{\operatorname{Fr}(X/S)} & X^{(p/S)} & \xrightarrow{\operatorname{pr}_X} & X \\ \varphi \uparrow & \searrow q & \downarrow q^{(p/S)} & & \downarrow q \\ T & \xrightarrow{r} & S & \xrightarrow{\operatorname{fr}(S)} & S. \end{array}$$

By the definition of  $X^{(p/S)}$  as a fiber product,  $\operatorname{pr}_X$  induces a bijection

$$X^{(p/S)}(r) = \operatorname{Hom}_S(T, X^{(p/S)}) \xrightarrow{\sim} \operatorname{Hom}_S(T, X) = X(\operatorname{fr}(S) \circ r).$$

On the other hand,

$$r \circ \operatorname{fr}(T) = \operatorname{fr}(S) \circ r,$$

because  $\operatorname{fr}$  is an endomorphism of the identity functor. Hence

$$\operatorname{Fr}(X/S)(r) : X(r) \longrightarrow X^{(p/S)}(r)$$

is characterized by commutativity of

$$\begin{array}{ccc} X(r) & \xrightarrow{\operatorname{Fr}(X/S)(r)} & X^{(p/S)}(r) \\ \downarrow X(\operatorname{fr}(T)) & & \downarrow \wr \\ X(r \circ \operatorname{fr}(T)) & \xlongequal{\quad} & X(\operatorname{fr}(S) \circ r). \end{array}$$

For example, if  $X$  is the subscheme of  $S$  defined by a quasi-coherent ideal  $\mathcal{I}$ , then  $X^{(p)}$  is the subscheme defined by the ideal  $\mathcal{I}^{\{p\}}$  generated by the  $p$ -th powers of local sections of  $\mathcal{I}$ . Moreover,  $\operatorname{Fr}(X/S)$  is the canonical immersion

$$\operatorname{Spec}(\mathcal{O}_S/\mathcal{I}) \longrightarrow \operatorname{Spec}(\mathcal{O}_S/\mathcal{I}^{\{p\}}).$$

<sup>43</sup>

**4.1.1.** <sup>44</sup> Let  $t : T \rightarrow S$  be a base change and put

$$X_T = X \times_{q,t} T.$$

Consider the pullback of  $X_T$  along  $\operatorname{fr}(T)$ :

$$\begin{array}{ccccc} (X_T)^{(p/T)} & \longrightarrow & X_T & \longrightarrow & X \\ \downarrow & & \downarrow & & \downarrow q \\ T & \xrightarrow{\operatorname{fr}(T)} & T & \xrightarrow{t} & S. \end{array}$$

<sup>42</sup>The editors expanded the original in the remainder of this paragraph.

<sup>43</sup>The French source prints  $\operatorname{Spec}(\mathcal{O}_X/\mathcal{I})$  for the source of this immersion. Since  $X \subset S$  is defined by  $\mathcal{I}$ , the intended source is  $\operatorname{Spec}(\mathcal{O}_S/\mathcal{I}) = X$ , as used here.

<sup>44</sup>The editors expanded the original in this paragraph.

Since

$$t \circ \text{fr}(T) = \text{fr}(S) \circ t,$$

$(X_T)^{(p/T)}$  identifies with the pullback of  $X^{(p/S)}$  along  $t$ . Equivalently, there is a canonical isomorphism

$$X_T^{(p/T)} \xrightarrow{\sim} (X^{(p/S)})_T.$$

Under this identification,  $\text{Fr}(X_T/T)$  identifies with the pullback  $\text{Fr}(X/S)_T$ .

**4.1.1.1.** In particular, if  $S = \text{Spec } \mathbb{F}_p$ , then  $X^{(p/S)} = X$  and  $\text{Fr}(X/S) = \text{fr}(X)$ . Consequently,  $X_T^{(p/T)}$  identifies with  $X_T$ , and  $\text{Fr}(X_T/T)$  with  $\text{fr}(X)_T$ .

For example, if  $E$  is a set and  $E_T$  the constant  $T$ -scheme of type  $E$ , then

$$E_T^{(p/T)} \simeq E_T, \quad \text{Fr}(E_T/T) \simeq \text{id}_{E_T}.$$

**4.1.2.** The functor  $X \mapsto X^{(p/S)}$  plainly commutes with products. It therefore sends an  $S$ -group  $G$  to an  $S$ -group  $G^{(p/S)}$ . Since  $\text{Fr}$  is functorial,

$$\text{Fr}(G/S) : G \longrightarrow G^{(p/S)}$$

is a homomorphism of  $S$ -groups. We denote its kernel by  $\text{Fr } G$ .

If  $r : T \rightarrow S$  is an  $S$ -scheme, the pointwise Frobenius diagram shows that the value of  $\text{Fr } G$  at  $r$  is the kernel of

$$G(\text{fr}(T)) : G(r) \longrightarrow G(r \circ \text{fr}(T)).$$

When  $T = I_R$  is the scheme of dual numbers over an  $S$ -scheme  $R$ , the absolute Frobenius of  $I_R$  factors as

$$I_R, \xrightarrow{\text{can.}} R \xrightarrow{\text{fr}(R)} R \xrightarrow{s} I_R$$

where  $s$  is the zero section. It follows that  $(\text{Fr } G)(I_R)$  contains the kernel  $\text{Lie}(G/S)(R)$  of

$$G(s) : G(I_R) \longrightarrow G(R),$$

and hence

$$\text{Lie}(G/S) = \text{Lie}(\text{Fr } G/S).$$

**4.1.3.**

More generally, for every  $S$ -functor  $X$ , define the  $S$ -functor  $X^{(p^n)}$  recursively in  $n$  by

$$X^{(p)} = X^{(p/S)} \quad \text{and} \quad X^{(p^n)} = (X^{(p^{n-1})})^{(p)}.$$

Likewise,  $\text{Fr}^n(X/S)$ , or simply  $\text{Fr}^n$ , denotes the composite functorial homomorphism

$$X \xrightarrow{\text{Fr}(X/S)} X^{(p)} \xrightarrow{\text{Fr}(X^{(p)}/S)} X^{(p^2)} \longrightarrow \dots \longrightarrow X^{(p^{n-1})} \xrightarrow{\text{Fr}(X^{(p^{n-1})}/S)} X^{(p^n)}.$$

By 4.1.1,  $\text{Fr}(X^{(p)}/S)$  coincides with  $\text{Fr}(X/S)^{(p)}$ ; in other words, the following diagram commutes:

$$\begin{array}{ccc} X^{(p)} & \longrightarrow & X \\ \text{Fr}(X^{(p)}/S) \downarrow & & \downarrow \text{Fr}(X/S) \\ X^{(p^2)} & \longrightarrow & X^{(p)}. \end{array}$$

If  $G$  is a group  $S$ -functor, then  $G^{(p^n)}$  is again a group  $S$ -functor and  $\mathrm{Fr}^n(G/S)$  is a homomorphism of group  $S$ -functors.

**Definition.** We denote the kernel of  $\mathrm{Fr}^n(G/S)$  by  $\mathrm{Fr}^n G$ , and say that  $G$  has height  $\leq n$  if  $\mathrm{Fr}^n(G/S)$  is zero, that is, if  $\mathrm{Fr}^n G = G$ .

**Lemma.** The group subfunctor  $\mathrm{Fr}^n G$  of  $G$  is characteristic: for every  $S$ -scheme  $T$ , every endomorphism  $\varphi$  of the group  $T$ -functor  $G_T$  induces an endomorphism of  $(\mathrm{Fr}^n G)_T$ .

*Proof.* By 4.1.1, the constructions of  $G^{(p^n)}$  and  $\mathrm{Fr}^n(G/S)$  commute with base change. We may therefore suppose that  $T = S$ . The assertion then follows from the fact that  $\mathrm{Fr}^n(G/S)$  is a functorial homomorphism.  $\square$

#### 4.1.4. Examples.

a) Let  $M$  first be an “abstract” abelian group, and let

$$G = D_S(M)$$

be the diagonalizable group of type  $M$  (I, 4.4). For every  $S$ -scheme  $T$ ,

$$G(T) = \mathrm{Hom}_{(\mathbf{Ab})}(M, \Gamma(T, \mathcal{O}_T)^\times).$$

Since  $G$  is the pullback of the diagonalizable group  $D(M)$  over  $\mathbb{F}_p$ ,  $G^{(p)}$  identifies with  $G$ , and  $\mathrm{Fr}(G/S)(T)$  identifies with the endomorphism  $x \mapsto x^p$  of  $G(T)$  (4.1.1). In particular, when  $M = \mathbb{Z}$ , one has  $D_S(M) = \mathbb{G}_{m,S}$ . Hence:

$\mathrm{Fr} \mathbb{G}_{m,S}$  is the  $S$ -group  $\mu_{p,S}$  which associates to every  $S$ -scheme  $T$  the group of  $p$ -th roots of unity in  $\Gamma(T, \mathcal{O}_T)^\times$ .

b) Now let  $S$  be a scheme of characteristic  $p$ , and let  $\mathcal{E}$  be a sheaf of modules on  $S$ . By I, 4.6.2 there is a canonical isomorphism

$$\mathcal{W}(\mathcal{E})^{(p)} \xrightarrow{\sim} \mathcal{W}(\mathcal{E}^{(p)}),$$

where  $\mathcal{E}^{(p)}$  is the pullback of  $\mathcal{E}$  along  $\mathrm{fr}(S)$ . For every  $S$ -scheme  $\pi : T \rightarrow S$ , the map  $\mathrm{Fr}(\mathcal{W}(\mathcal{E})/S)(\pi)$  is determined, by 4.1 (†), by the commutative triangle

$$\begin{array}{ccc} \Gamma(T, \pi^* \mathrm{fr}(S)^* \mathcal{E}) & \xrightarrow[\mathrm{can.}]{\sim} & \Gamma(T, \mathrm{fr}(T)^* \pi^* \mathcal{E}) \\ & \nwarrow \mathrm{Fr}(\mathcal{W}(\mathcal{E})/S)(\pi) \quad \nearrow f' & \\ & \Gamma(T, \pi^* \mathcal{E}) & \end{array},$$

where  $f'$  is the map induced by  $\mathrm{fr}(T)$ .

In particular, if  $\mathcal{E} = \mathcal{O}_S$ , then  $\mathcal{W}(\mathcal{E})$  identifies with the additive group  $\mathbb{G}_{a,S}$ . In this case  $\mathcal{E}^{(p)} = \mathcal{E} = \mathcal{O}_S$ , and the Frobenius morphism  $\mathrm{Fr}(\mathbb{G}_{a,S}/S)$  sends  $x \in \Gamma(T, \mathcal{O}_T)$  to  $x^p$ . Thus:

$\mathrm{Fr} \mathbb{G}_{a,S}$  is the  $S$ -group  $\alpha_{p,S}$  which associates to every  $S$ -scheme  $T$  the group

$$\{x \in \Gamma(T, \mathcal{O}_T) \mid x^p = 0\}.$$

c) Similarly, for every quasi-coherent  $\mathcal{O}_S$ -algebra  $\mathcal{A}$ , the scheme  $(\mathrm{Spec} \mathcal{A})^{(p)}$  identifies with  $\mathrm{Spec} \mathcal{A}^{(p)}$ , the spectrum of the pullback of  $\mathcal{A}$  along  $\mathrm{fr}(S)$ . If  $\pi$  denotes the endomorphism  $x \mapsto x^p$  of the sheaf of rings  $\mathcal{O}_S$ , then

$$\mathcal{A}^{(p)} = \mathcal{A} \otimes_{\pi} \mathcal{O}_S, \quad (44)$$

and  $\mathrm{Fr}((\mathrm{Spec} \mathcal{A})/S)$  is induced by the morphism of  $\mathcal{O}_S$ -algebras

$$\mathcal{A} \otimes_{\pi} \mathcal{O}_S \longrightarrow \mathcal{A}, \quad a \otimes_{\pi} x \longmapsto a^p x.$$

45

Finally, for every quasi-coherent  $\mathcal{O}_S$ -module  $\mathcal{E}$ , there are canonical isomorphisms

$$\mathcal{V}(\mathcal{E})^{(p)} \xrightarrow{\sim} \mathcal{V}(\mathcal{E}^{(p)}) \quad \text{and} \quad \mathcal{S}(\mathcal{E})^{(p)} \xrightarrow{\sim} \mathcal{S}(\mathcal{E}^{(p)}),$$

where  $\mathcal{S}(\mathcal{E})$  denotes the symmetric algebra of the  $\mathcal{O}_S$ -module  $\mathcal{E}$ .

d) Let  $\mathcal{U}$  be an  $\mathcal{O}_S$ -coalgebra (3.1), and let  $T$  be a scheme of characteristic  $p$ . If  $\mathcal{U}^{(p/S)}$ , or  $\mathcal{U}^{(p)}$ , denotes the pullback of the coalgebra  $\mathcal{U}$  along  $\mathrm{fr}(S)$ , then, as in b), there is a canonical isomorphism

$$(\mathrm{Spec}^* \mathcal{U})^{(p)} \xrightarrow{\sim} \mathrm{Spec}^* \mathcal{U}^{(p)}.$$

If  $\mathcal{U}$  is a group coalgebra, then for an  $S$ -scheme  $T$ , the value of  $\mathrm{Fr}(\mathrm{Spec}^* \mathcal{U})$ , that is, of the kernel of the Frobenius morphism

$$\mathrm{Spec}^* \mathcal{U} \longrightarrow (\mathrm{Spec}^* \mathcal{U})^{(p)},$$

is the set of elements  $\gamma$  of

$$(\mathrm{Spec}^* \mathcal{U})(T) = \{x \in \Gamma(T, \mathcal{U}_T) \mid \varepsilon_{\mathcal{U}_T}(x) = 1, \Delta_{\mathcal{U}_T}x = x \otimes x\}$$

such that the image in  $\Gamma(T, \mathcal{U}_T \otimes_{\mathrm{fr}(T)} \mathcal{O}_T)$  of the element

$$\gamma \otimes_{\mathrm{fr}(T)} 1 \in \Gamma(T, \mathcal{U}_T) \otimes_{\mathrm{fr}(T)} \mathcal{O}(T)$$

is equal to 1.

#### 4.2. <sup>46</sup>

We now consider a construction closely related to the preceding one. Let  $S$  be a scheme of characteristic  $p$ , let  $X$  be an  $S$ -scheme, and let  $X_S^p$  be the product, in  $(\mathbf{Sch}/S)$ , of  $p$  copies of  $X$ .

Denote by  $U^p(X)$  the open subscheme of  $X_S^p$  which is the union of the products  $U_S^p$  as  $U$  ranges over the affine open subsets of  $X$ . Thus a point  $x$  of  $X_S^p$  belongs to  $U^p(X)$  if and only if its projections  $\mathrm{pr}_i x$  on the factors of  $X_S^p$  belong to a common affine open subset of  $X$ . For example, if every finite subset of  $X$  is contained in an affine open subset, then

$$U^p(X) = X_S^p.$$

The symmetric group  $\mathfrak{S}_p$  on  $p$  letters acts on  $X_S^p$  by permuting the factors and leaves  $U^p(X)$  stable. We call the quotient of  $X_S^p$  by  $\mathfrak{S}_p$  in the category of ringed spaces the *p-fold symmetric product of X*, and denote it by  $\Sigma^p X$ . Let

$$q(X), \text{ or simply } q : X_S^p \longrightarrow \Sigma^p X$$

<sup>45</sup>  $\mathcal{A} \otimes_{\pi} \mathcal{O}_S$  denotes the  $\mathcal{O}_S$ -algebra obtained by extension of scalars along  $\pi : \mathcal{O}_S \rightarrow \mathcal{O}_S$ ; thus  $ax \otimes_{\pi} 1 = a \otimes_{\pi} x^p$  and  $x \cdot (a \otimes_{\pi} 1) = a \otimes_{\pi} x$ .

<sup>46</sup> For the material in 4.2 and 4.3, see also [DG70], IV, §3, nos. 4–6.

be the canonical projection.

The morphism  $q$  maps  $U^p(X)$  onto an open subset  $V^p(X)$  of the symmetric product, which may be described as follows (cf. V, 4.1). The structure sheaf of  $\Sigma^p X$  induces a scheme structure on  $V^p(X)$ , and the morphism

$$q'(X) : U^p(X) \longrightarrow V^p(X)$$

induced by  $q(X)$  is affine, indeed integral. As  $U$  ranges over the affine open subsets of  $X$  which map into a varying affine open  $V$  of  $S$ , the schemes  $\Sigma^p U$  form an affine open covering of  $V^p(X)$ . If  $R$  is the affine algebra of  $V$ , and  $A$  that of  $U$ , the affine algebra of  $\Sigma^p U$  is the subalgebra  $\Sigma^p A$  of  $\bigotimes_R^p A$  consisting of the symmetric tensors.

Now consider the diagonal morphism

$$\delta : X \longrightarrow U^p(X).$$

If  $V = \operatorname{Spec} R$  is an affine open subset of  $S$  and  $U = \operatorname{Spec} A$  is an affine open subset of  $X$  lying over  $V$ , the restriction of  $\delta$  to  $U$  is defined by the algebra homomorphism

$$\eta : \bigotimes_R^p A \longrightarrow A, \quad a_1 \otimes \cdots \otimes a_p \longmapsto a_1 a_2 \cdots a_p.$$

Consequently, if  $N$  denotes the symmetrization operator,

$$\begin{aligned} \eta(N(a_1 \otimes \cdots \otimes a_p)) &= \eta \left( \sum_{\sigma \in \mathfrak{S}_p} a_{\sigma(1)} \otimes \cdots \otimes a_{\sigma(p)} \right) \\ &= p! a_1 \cdots a_p = 0. \end{aligned}$$

In other words,  $\eta$  vanishes on the subspace

$$N \left( \bigotimes_R^p A \right) \subseteq \Sigma^p A$$

consisting of symmetrized tensors. Moreover, if  $f$  is a symmetric tensor, then plainly  $N(fa) = fN(a)$ . Hence  $N(\bigotimes_R^p A)$  is an ideal of  $\Sigma^p A$ . We put

$$U^{[p/S]} = \operatorname{Spec} \left( \Sigma^p A / N \left( \bigotimes_R^p A \right) \right).$$

This is a closed subscheme of  $\Sigma^p(U) = V^p(U)$ . As  $U$  ranges over the affine open subsets of  $X$  which map into a varying affine open  $V$  of  $S$ , the schemes  $U^{[p/S]}$  glue to a closed subscheme of  $V^p(X)$ , denoted  $X^{[p/S]}$ .

If  $i(X)$  denotes the inclusion of  $X^{[p/S]}$  in  $V^p(X)$ , the preceding calculation shows that  $q'(X) \circ \delta$  factors through  $X^{[p/S]}$ . We therefore obtain a morphism

$$F^{[p]}(X/S) : X \longrightarrow X^{[p/S]}.$$

<sup>47</sup> The factorization is displayed by the commutative diagram

<sup>47</sup>In the original, this morphism, respectively the relative Frobenius morphism, was denoted  $F'$ , respectively  $F$ .

$$\begin{array}{ccccc}
X_S^p & \supset & U^p(X) & \xleftarrow{\delta(X)} & X \\
\downarrow q(X) & & \downarrow q'(X) & & \downarrow F^{[p]}(X/S) \\
\Sigma^p(X) & \supset & V^p(X) & \xleftarrow{i(X)} & X^{[p/S]}.
\end{array}$$

It is clear that  $X^{[p/S]}$  is functorial in  $X$ , and that

$$F^{[p]} : X \mapsto F^{[p]}(X/S)$$

is a functorial homomorphism.

#### 4.2.1.

The schemes  $X^{[p/S]}$  and  $X^{(p/S)}$  are naturally related. Let  $V$  be an affine open subset of  $S$ , with affine algebra  $R$ , and let  $U$  be an affine open subset of  $X$  lying over  $V$ , with affine algebra  $A$ . If  $\pi$  denotes the endomorphism  $x \mapsto x^p$  of  $R$ , the affine algebra of  $U^{(p/S)}$  is  $A \otimes_\pi R$ . The map

$$a \otimes_\pi \lambda \mapsto \left( \lambda a \otimes \cdots \otimes a \bmod N \left( \bigotimes_R^p A \right) \right)$$

is a homomorphism of  $R$ -algebras

$$A \otimes_\pi R \longrightarrow \Sigma^p A / N \left( \bigotimes_R^p A \right).$$

It induces a morphism

$$\varphi(U) : U^{[p/S]} \longrightarrow U^{(p/S)}$$

satisfying

$$\varphi(U) \circ F^{[p]}(U/S) = \text{Fr}(U/S).$$

Gluing these morphisms gives the commutative triangle

$$\begin{array}{ccc}
& X & \\
F^{[p]}(X/S) \swarrow & & \searrow \text{Fr}(X/S) \\
X^{[p/S]} & \xrightarrow{\varphi(X)} & X^{(p/S)}.
\end{array}$$

For example, if  $X$  is the subscheme of  $S$  defined by a quasi-coherent ideal  $\mathcal{I}$ , then  $F^{[p]}(X/S)$  identifies with the identity morphism of  $X$ . Thus  $\varphi(X)$  is the canonical immersion

$$\text{Spec}(\mathcal{O}_S/\mathcal{I}) \longrightarrow \text{Spec}(\mathcal{O}_S/\mathcal{I}^{\{p\}}).$$

This shows that  $\varphi(X)$  need not be an isomorphism.

On the other hand, when  $M$  is a free  $R$ -module, the map

$$\begin{aligned}
M \otimes_\pi R &\longrightarrow \Sigma^p M / N \left( \bigotimes_R^p M \right), \\
m \otimes_\pi \lambda &\mapsto \left( \lambda m \otimes \cdots \otimes m \bmod N \left( \bigotimes_R^p M \right) \right)
\end{aligned}$$



is plainly bijective. It remains bijective when  $M$  is  $R$ -flat, because every flat module is a filtered direct limit of free modules.<sup>48</sup> It follows that

$$\varphi(X) : X^{[p/S]} \longrightarrow X^{(p/S)} \text{ is an isomorphism if } X \text{ is flat over } S.$$

#### 4.3.

Finally, let  $G$  be an  $S$ -group scheme in abelian groups. The composite morphism

$$\mu(G) : U^p(G) \hookrightarrow G_S^p \longrightarrow G,$$

defined by multiplication, factors through  $V^p(G)$ . Thus there is a morphism

$$\nu(G) : V^p(G) \longrightarrow G$$

such that

$$\nu(G) \circ q'(G) = \mu(G).$$

The resulting commutative diagram is

$$\begin{array}{ccccc} G & \xleftarrow{\mu(G)} & U^p(G) & \xleftarrow{\delta(G)} & G \\ & \nwarrow \nu(G) & \searrow q'(G) & & \nwarrow \text{Fr}(G/S) \\ & & V^p(G) & \xleftarrow{i(G)} & G^{[p/S]} \\ & & & \searrow \varphi(G) & \\ & & & & G^{(p/S)} \end{array}$$

Ver( $G/S$ )  $\nearrow$  (dotted arrow from  $G^{(p/S)}$  to  $G$ )

When  $G$  is flat over  $S$ ,  $\varphi(G)$  is an isomorphism, and we may define the morphism called the *Verschiebung*

$$\text{Ver}(G/S) : G^{(p/S)} \longrightarrow G$$

by

$$\text{Ver}(G/S) = \nu(G) \circ i(G) \circ \varphi(G)^{-1}.$$

As  $G$  ranges over the flat commutative  $S$ -group schemes, the assignment

$$\text{Ver} : G \longmapsto \text{Ver}(G/S)$$

is plainly a functorial homomorphism. Consequently,  $\text{Ver}(G/S)$  is a group homomorphism.

For every  $S$ -scheme  $T$ , the composite

$$G(T) \xrightarrow{\delta(G)(T)} U^p(G)(T) \xrightarrow{\mu(G)(T)} G(T)$$

sends  $x \in G(T)$  to  $p \cdot x$ . We may therefore write  $p \cdot \text{id}_G$  in place of  $\mu(G) \circ \delta(G)$ , obtaining the classical formula

$$\text{Ver}(G/S) \circ \text{Fr}(G/S) = p \cdot \text{id}_G. \quad (*)$$

#### Examples 4.3.1.

<sup>48</sup>D. Lazard, *Sur les modules plats*, C. R. Acad. Sci. Paris **258** (1964), 6313–6316. See also D. Lazard, Bull. Soc. Math. France **97** (1969), 81–128, or [BAlg], X, §1.6, Th. 1.

a) If  $G$  is a constant  $S$ -scheme in abelian groups, then  $\mathrm{Fr}(G/S)$  identifies with the identity morphism of  $G$  (cf. 4.1.1.1). Hence

$$\mathrm{Ver}(G/S) = p \cdot \mathrm{id}_G.$$

b) If  $G$  is the diagonalizable  $S$ -group of type  $M$ , then  $\mathrm{Fr}(G/S) = p \cdot \mathrm{id}_G$  by 4.1.4 a). It follows at once that  $\mathrm{Ver}(G/S)$  is the identity morphism of  $G$ .

c) If  $\mathcal{E}$  is a flat  $\mathcal{O}_S$ -module and  $G = \mathcal{V}(\mathcal{E})$ , then  $\mathrm{Ver}(G/S)$  is zero, as is  $p \cdot \mathrm{id}_G$ . In Exposé VII<sub>B</sub> we shall see that a commutative algebraic group  $G$  over a field  $k$  is “unipotent” if and only if, for some  $n$ , the composite homomorphism

$$G^{(p^n)} \xrightarrow{\mathrm{Ver}(G^{(p^{n-1})}/S)} G^{(p^{n-1})} \longrightarrow \cdots \longrightarrow G^{(p)} \xrightarrow{\mathrm{Ver}(G/S)} G$$

is zero. Here

$$G^{(p^n)} = (G^{(p^{n-1})})^{(p)}$$

as in 4.1.3.<sup>49</sup>

#### 4.3.2.

Since  $\mathrm{Ver} : G \mapsto \mathrm{Ver}(G/S)$  is a functorial homomorphism on the category of flat commutative  $S$ -group schemes, the square

$$\begin{array}{ccc} G^{(p)} & \xrightarrow{\mathrm{Ver}(G/S)} & G \\ \mathrm{Fr}(G/S)^{(p)} \downarrow & & \downarrow \mathrm{Fr}(G/S) \\ G^{(p^2)} & \xrightarrow{\mathrm{Ver}(G^{(p)}/S)} & G^{(p)} \end{array}$$

commutes. Here  $\mathrm{Fr}(G/S)^{(p)}$  denotes the pullback of  $\mathrm{Fr}(G/S)$  along the base change  $\mathrm{fr}(S)$ . By 4.1.1,

$$\mathrm{Fr}(G/S)^{(p)} = \mathrm{Fr}(G^{(p)}/S).$$

Applying formula (\*) of 4.3 to  $G^{(p)}$ , we obtain

$$\mathrm{Fr}(G/S) \circ \mathrm{Ver}(G/S) = \mathrm{Ver}(G^{(p)}/S) \circ \mathrm{Fr}(G^{(p)}/S) = p \cdot \mathrm{id}_{G^{(p)}}. \quad (**)$$

#### 4.3.3.

Suppose finally that  $G$  is a commutative  $S$ -group which is finite and locally free. Let  $\mathcal{A}$  be its affine  $\mathcal{O}_S$ -algebra, and let  $\pi$  be the endomorphism of the sheaf of rings  $\mathcal{O}_S$  which sends a section  $x$  to  $x^p$ .<sup>50</sup> Denote by  $\Sigma^p \mathcal{A}$  the subalgebra of  $\bigotimes_{\mathcal{O}_S}^p \mathcal{A}$  consisting of the sections invariant under the symmetric-group action, and by

$$i(\mathcal{A}) : \Sigma^p \mathcal{A} \longrightarrow \bigotimes_{\mathcal{O}_S}^p \mathcal{A}$$

its inclusion.

Let

$$\Delta^p(\mathcal{A}) : \mathcal{A} \longrightarrow \bigotimes_{\mathcal{O}_S}^p \mathcal{A}$$

<sup>49</sup>This criterion is not stated explicitly in Exposé VII<sub>B</sub>; see [DG70], IV, §3, Prop. 4.11.

<sup>50</sup>The editors changed the order of presentation, introducing first the objects which occur in the diagram below.

be the morphism obtained by iterating the diagonal morphism of the coalgebra  $\mathcal{A}$ . It corresponds to the multiplication morphism  $U^p(G) = G_S^p \rightarrow G$ . By the beginning of 4.3,  $\Delta^p(\mathcal{A})$  factors through  $\Sigma^p \mathcal{A}$ ; that is, it induces a morphism

$$a(\mathcal{A}) : \mathcal{A} \longrightarrow \Sigma^p \mathcal{A}$$

such that

$$i(\mathcal{A}) \circ a(\mathcal{A}) = \Delta^p(\mathcal{A}).$$

On the other hand, let  $S^p(\mathcal{A})$  be the homogeneous component of degree  $p$  in the symmetric algebra of  $\mathcal{A}$ , and let

$$q(\mathcal{A}) : \bigotimes_{\mathcal{O}_S}^p \mathcal{A} \longrightarrow S^p(\mathcal{A})$$

be the canonical projection. The multiplication

$$m^p(\mathcal{A}) : \bigotimes_{\mathcal{O}_S}^p \mathcal{A} \longrightarrow \mathcal{A}$$

factors through  $S^p(\mathcal{A})$ ; equivalently, it induces a map

$$b(\mathcal{A}) : S^p(\mathcal{A}) \longrightarrow \mathcal{A}$$

such that

$$b(\mathcal{A}) \circ q(\mathcal{A}) = m^p(\mathcal{A}).$$

Since  $\Sigma^p \mathcal{A}$  is the affine algebra of  $V^p(G)$ , the composite  $i(G) \circ \varphi(G)^{-1}$  induces, again by the beginning of 4.3, an algebra homomorphism

$$r(\mathcal{A}) : \Sigma^p \mathcal{A} \longrightarrow \mathcal{A} \otimes_{\pi} \mathcal{O}_S.$$

It vanishes on sections of the form

$$\sum_{\sigma \in \mathfrak{S}_p} a_{\sigma(1)} \otimes \cdots \otimes a_{\sigma(p)}$$

and sends  $a \otimes \cdots \otimes a$  to  $a \otimes_{\pi} 1$ . Similarly,  $j(\mathcal{A})$  is the morphism of  $\mathcal{O}_S$ -modules

$$a \otimes_{\pi} 1 \longmapsto q(a \otimes \cdots \otimes a).$$

These maps form the commutative diagram

$$\begin{array}{ccccc}
 \mathcal{A} & \xrightarrow{\Delta^p(\mathcal{A})} & \bigotimes_{\mathcal{O}_S}^p \mathcal{A} & \xrightarrow{m^p(\mathcal{A})} & \mathcal{A} \\
 \searrow a(\mathcal{A}) & & \nearrow i(\mathcal{A}) & \searrow q(\mathcal{A}) & \nearrow b(\mathcal{A}) \\
 & \Sigma^p \mathcal{A} & & S^p(\mathcal{A}) & \\
 & \searrow r(\mathcal{A}) & & \nearrow j(\mathcal{A}) & \\
 & & \mathcal{A} \otimes_{\pi} \mathcal{O}_S & & 
 \end{array}$$

The composite  $r(\mathcal{A}) \circ a(\mathcal{A})$  corresponds to the Verschiebung morphism  $\text{Ver}(G/S)$ , while  $b(\mathcal{A}) \circ j(\mathcal{A})$  corresponds to the Frobenius morphism  $\text{Fr}(G/S)$ .

The preceding commutative diagram  $(\mathcal{A})$  is self-dual. Indeed, let  $D$  be the functor which associates to each  $\mathcal{O}_S$ -module  $\mathcal{M}$  its dual

$$D\mathcal{M} = \text{Hom}_{\mathcal{O}_S}(\mathcal{M}, \mathcal{O}_S).$$

The image of diagram  $(\mathcal{A})$  under  $D$  is precisely diagram  $(D\mathcal{A})$ . The morphisms

$$Dr(\mathcal{A}), \quad Da(\mathcal{A}), \quad Dj(\mathcal{A}), \quad Db(\mathcal{A})$$

identify respectively with

$$j(D\mathcal{A}), \quad b(D\mathcal{A}), \quad r(D\mathcal{A}), \quad a(D\mathcal{A}).$$

By 3.3.1, it follows that:

*In the category of commutative  $S$ -groups which are finite and locally free, Cartier duality interchanges the Frobenius morphism and the Verschiebung.*

51

## 5. Restricted $p$ -Lie Algebras

We first recall some results from the *Séminaire Sophus Lie*.<sup>52</sup>

### 5.1.

Let  $p$  be a prime number, let  $R$  be a commutative ring of characteristic  $p$ , and let  $A$  be an associative  $R$ -algebra, not necessarily commutative. For  $a, b \in A$ , put

$$[a, b] = ab - ba, \quad a \cdot b = L_a(b) = R_b(a).$$

Then

$$\begin{aligned} (\text{ad } x^p)(y) &= [x^p, y] \\ &= (L_x^p - R_x^p)(y) \\ &= (L_x - R_x)^p(y) \\ &= (\text{ad } x)^p(y), \end{aligned}$$

which gives Jacobson's first formula:

$$\text{ad}(x^p) = (\text{ad } x)^p. \quad (\text{i})$$

If  $a_1, \dots, a_p$  are arbitrary elements of  $A$ , and  $N$  is the symmetrization operator (cf. 4.2), then

$$\begin{aligned} N(a_1 \otimes \cdots \otimes a_p) &= \sum_{\sigma \in \mathfrak{S}_p} a_{\sigma(1)} \cdots a_{\sigma(p)} \\ &= \sum_{\tau \in \mathfrak{S}_{p-1}} [a_{\tau(1)}, [a_{\tau(2)}, [\cdots, [a_{\tau(p-1)}, a_p] \cdots ]]]. \end{aligned} \quad (*)$$

Here  $\sigma$  ranges over the permutations of  $p$  letters and  $\tau$  over those of  $p-1$  letters.

<sup>51</sup>See also [DG70], IV, §3, 4.9.

<sup>52</sup>See P. Cartier, *Exemples d'hyperalgèbres*, Séminaire Sophus Lie 1955/56, Exp. 3.

Indeed, the final expression is equal to

$$\sum_{\tau} \sum_{r=0}^{p-1} \sum_{i_1 < \dots < i_r} (-1)^s a_{\tau(i_1)} \cdots a_{\tau(i_r)} a_p a_{\tau(j_s)} \cdots a_{\tau(j_1)},$$

where  $\tau$  ranges over the permutations of  $p-1$  letters;  $i_1, \dots, i_r$  ranges over the strictly increasing sequences in  $[1, p-1]$ ; and  $j_1, \dots, j_s$  is the strictly increasing sequence of those integers of  $[1, p-1]$  which are different from  $i_1, \dots, i_r$ . For fixed  $r$ , the sum of the terms

$$(-1)^s a_{\tau(i_1)} \cdots a_{\tau(i_r)} a_p a_{\tau(j_s)} \cdots a_{\tau(j_1)}$$

is plainly

$$(-1)^s \binom{p-1}{s} \sum_{\rho} a_{\rho(1)} \cdots a_{\rho(r)} a_p a_{\rho(r+1)} \cdots a_{\rho(p-1)},$$

where  $\rho$  ranges over the permutations of  $p-1$  letters. But

$$(-1)^s \binom{p-1}{s} = 1 \quad \text{in } \mathbb{F}_p,$$

because in  $\mathbb{F}_p[x]$ ,

$$(x-1)^p = x^p - 1 = (x-1)(x^{p-1} + \cdots + 1),$$

and hence

$$(x-1)^{p-1} = x^{p-1} + \cdots + 1.$$

This proves (\*).

On the other hand, if  $x_0, x_1 \in A$ , then

$$(x_0 + x_1)^p = x_0^p + x_1^p + \sum_z x_{z(1)} x_{z(2)} \cdots x_{z(p)},$$

where  $z$  ranges over the nonconstant maps  $[1, p] \rightarrow \{0, 1\}$ . It follows that

$$\begin{aligned} (x_0 + x_1)^p &= x_0^p + x_1^p \\ &+ \sum_{0 < r < p} \frac{1}{r!(p-r)!} N(\underbrace{x_0, \dots, x_0}_r, \underbrace{x_1, \dots, x_1}_{p-r}). \end{aligned}$$

<sup>53</sup> By (\*),

$$\begin{aligned} &N(\underbrace{x_0, \dots, x_0}_r, \underbrace{x_1, \dots, x_1}_{p-r}) \\ &= r!(p-1-r)! \sum_t [x_{t(1)}, [x_{t(2)}, [\cdots, [x_{t(p-1)}, x_1] \cdots ]]], \end{aligned}$$

where  $t$  ranges over the maps  $[1, p-1] \rightarrow \{0, 1\}$  which take the value 0 exactly  $r$  times. We obtain Jacobson's second formula:

$$\begin{aligned} (x_0 + x_1)^p &= x_0^p + x_1^p \\ &- \sum_{0 < r < p} \sum_t \frac{1}{r} [x_{t(1)}, [x_{t(2)}, [\cdots, [x_{t(p-1)}, x_1] \cdots ]]], \end{aligned} \quad (\text{ii})$$

where  $t$  has the same meaning.

---

<sup>53</sup>The editors inserted the following explanation from [DG70], III.7, 3.2.

**5.2.**

Let now  $\mathfrak{g}$  be an  $R$ -Lie algebra. A map

$$x \longmapsto x^{(p)}$$

from  $\mathfrak{g}$  to itself is said to make  $\mathfrak{g}$  a *restricted  $p$ -Lie algebra over  $R$*  if the following conditions hold:

$$\begin{aligned} (0) \quad & (\lambda x)^{(p)} = \lambda^p x^{(p)} \quad (\lambda \in R, x \in \mathfrak{g}), \\ (i) \quad & \operatorname{ad} x^{(p)} = (\operatorname{ad} x)^p \quad (x \in \mathfrak{g}), \end{aligned}$$

and

$$\begin{aligned} (ii) \quad & (x_0 + x_1)^{(p)} = x_0^{(p)} + x_1^{(p)} \\ & - \sum_{0 < r < p} \sum_t \frac{1}{r} [x_{t(1)}, [x_{t(2)}, [\cdots, [x_{t(p-1)}, x_1] \cdots ]]], \end{aligned}$$

where  $t$  ranges over the maps  $[1, p-1] \rightarrow \{0, 1\}$  which take the value 0 exactly  $r$  times, and  $x_0, x_1 \in \mathfrak{g}$ . The map  $x \mapsto x^{(p)}$  is called the *symbolic  $p$ -th power*.

For example, if  $A$  is an associative  $R$ -algebra, 5.1 shows that the  $R$ -module underlying  $A$ , endowed with

$$[x, y] = xy - yx, \quad x^{(p)} = x^p,$$

is a  $p$ -Lie algebra, denoted  $A_{\text{Lie}}$ . We call it the  $p$ -Lie algebra underlying  $A$ .

We shall chiefly consider restricted  $p$ -Lie subalgebras of algebras of the form  $A_{\text{Lie}}$ . Here is one example. Let  $S$  be a scheme of characteristic  $p > 0$ , and let  $X$  be an  $S$ -scheme. Recall that a derivation of  $X$  over  $S$  is an endomorphism  $D$  of the sheaf of abelian groups  $\mathcal{O}_X$  such that

$$D(\lambda s) = \lambda D(s), \quad D(st) = (Ds)t + s(Dt),$$

whenever  $\lambda$ , respectively  $s, t$ , are sections of  $\mathcal{O}_S$ , respectively  $\mathcal{O}_X$ , on open subsets for which the formulas make sense. Leibniz's formula

$$D^n(st) = \sum_{i=0}^n \binom{n}{i} (D^i s)(D^{n-i} t)$$

shows that  $D^p$  is again a derivation of  $X$  over  $S$ , since

$$\binom{p}{i} \equiv 0 \pmod{p} \quad (i \neq 0, p).$$

It follows that:

*The algebra  $\operatorname{Der}_{X/S}$  of derivations of  $X$  over  $S$  is a restricted  $p$ -Lie subalgebra of the  $\Gamma(S, \mathcal{O}_S)$ -algebra of differential operators of  $X$  over  $S$ .*

**5.2.1.** If  $\mathfrak{g}$  and  $\mathfrak{h}$  are two restricted  $p$ -Lie algebras, a *homomorphism*

$$h : \mathfrak{g} \longrightarrow \mathfrak{h}$$

is an  $R$ -linear map such that, for  $x, y \in \mathfrak{g}$ ,

$$h([x, y]) = [h(x), h(y)], \quad h(x^{(p)}) = h(x)^{(p)}.$$

The composite of two homomorphisms is again a homomorphism, so we may speak of the *category of restricted  $p$ -Lie algebras over  $R$* .

If  $(X, \mathcal{R})$  is a ringed space, we say that an  $\mathcal{R}$ -module  $\mathfrak{g}$  is endowed with the structure of a restricted  $p$ -Lie algebra over  $\mathcal{R}$  if, for every open subset  $U$ , the module  $\Gamma(U, \mathfrak{g})$  is a restricted  $p$ -Lie algebra over  $\Gamma(U, \mathcal{R})$ , and if all restriction maps are homomorphisms.

### 5.3.

We are now interested in the left adjoint of the functor

$$A \longmapsto A_{\text{Lie}}$$

of 5.2. Let  $\mathfrak{g}$  be a restricted  $p$ -Lie algebra over a ring  $R$  of characteristic  $p$ , let  $U(\mathfrak{g})$  be the enveloping algebra of the Lie algebra underlying  $\mathfrak{g}$  (cf. [BLie], I, §2.1), and let

$$i_{\mathfrak{g}} : \mathfrak{g} \longrightarrow U(\mathfrak{g})$$

(or simply  $i$ ) be the canonical map.

Let  $A$  be a unital associative  $R$ -algebra. For every Lie algebra homomorphism

$$\varphi : \mathfrak{g} \longrightarrow A_{\text{Lie}},$$

there is a unique homomorphism of unital  $R$ -algebras

$$\psi : U(\mathfrak{g}) \longrightarrow A$$

such that  $\psi \circ i = \varphi$ . Moreover,  $\varphi$  is a homomorphism of restricted  $p$ -Lie algebras if and only if  $\psi$  vanishes on all the elements

$$i(x)^p - i(x^{(p)}), \quad x \in \mathfrak{g}.$$

**Definition.** The quotient of  $U(\mathfrak{g})$  by the two-sided ideal generated by the elements

$$i(x)^p - i(x^{(p)})$$

is denoted  $U_p^R(\mathfrak{g})$ , or simply  $U_p(\mathfrak{g})$ . We denote by  $j_{\mathfrak{g}}$ , or simply  $j$ , the composite

$$\mathfrak{g} \xrightarrow{i} U(\mathfrak{g}) \longrightarrow U_p(\mathfrak{g}).$$

The algebra  $U_p(\mathfrak{g})$  is called the *restricted enveloping algebra* of  $\mathfrak{g}$ .

The preceding discussion gives the following result.

**Proposition.** *For every unital associative  $R$ -algebra  $A$  and every homomorphism of restricted  $p$ -Lie algebras*

$$\varphi : \mathfrak{g} \longrightarrow A_{\text{Lie}},$$

*there is a unique homomorphism of unital algebras*

$$\psi : U_p(\mathfrak{g}) \longrightarrow A$$

*such that  $\psi \circ j = \varphi$ . In other words, the functor  $\mathfrak{g} \mapsto U_p(\mathfrak{g})$  is left adjoint to the forgetful functor  $A \mapsto A_{\text{Lie}}$ .*

#### 5.3.1.

With the notation of 5.3, put

$$\beta(x) = i(x)^p - i(x^{(p)}).$$

For every  $y \in \mathfrak{g}$ , formulas 5.1 (i) and 5.2 (i) give

$$\begin{aligned}\beta(x)i(y) &= i(y)\beta(x) + [\beta(x), i(y)] \\ &= i(y)\beta(x) + (\operatorname{ad} i(x))^p i(y) - i((\operatorname{ad} x)^p y) \\ &= i(y)\beta(x).\end{aligned}$$

Thus  $\beta(x)$  belongs to the center of  $U(\mathfrak{g})$ ; in particular, the left ideal generated by the elements  $\beta(x)$  is already two-sided.

On the other hand,

$$\beta(\lambda x) = \lambda^p \beta(x) \quad (\lambda \in R),$$

and formulas 5.1 (ii) and 5.2 (ii) give

$$\beta(x + y) = \beta(x) + \beta(y) \quad (x, y \in \mathfrak{g}).$$

Consequently, if  $(x_\alpha)$  is a family of generators of the  $R$ -module  $\mathfrak{g}$ , the left ideal generated by all  $\beta(x)$  is already generated by the elements  $\beta(x_\alpha)$ .

**5.3.2. Proposition.**<sup>54</sup> *Let  $\mathfrak{g}$  be an  $R$ -Lie algebra whose underlying  $R$ -module is free with basis  $(x_\alpha)$ . Then the restricted  $p$ -Lie algebra structures on  $\mathfrak{g}$  correspond bijectively to the families  $(y_\alpha)$  of elements of  $\mathfrak{g}$  such that*

$$\operatorname{ad} y_\alpha = (\operatorname{ad} x_\alpha)^p.$$

Indeed, suppose that  $\mathfrak{g}$  is endowed with a restricted  $p$ -Lie algebra structure  $x \mapsto x^{(p)}$ . By 5.2 (i), (0), and (ii), the elements

$$y_\alpha = x_\alpha^{(p)}$$

satisfy

$$\operatorname{ad} y_\alpha = (\operatorname{ad} x_\alpha)^p,$$

and they determine the restricted  $p$ -Lie algebra structure.

We prove the converse. Since  $\mathfrak{g}$  is a free  $R$ -module, the canonical map

$$i : \mathfrak{g} \longrightarrow U(\mathfrak{g})$$

is injective by the Poincaré–Birkhoff–Witt theorem (cf. [BLie], I, §2.7). We may therefore identify  $\mathfrak{g}$  with an  $R$ -submodule of  $U(\mathfrak{g})$ . Suppose that  $(y_\alpha)$  is a family of elements of  $\mathfrak{g}$  such that

$$\operatorname{ad} y_\alpha = (\operatorname{ad} x_\alpha)^p.$$

Let

$$\pi : R \longrightarrow R, \quad r \longmapsto r^p,$$

and let  $\mathfrak{g} \otimes_\pi R$  be the  $R$ -Lie algebra obtained by extension of scalars along  $\pi$ .<sup>55</sup>

There is an  $R$ -linear map

$$\gamma : \mathfrak{g} \otimes_\pi R \longrightarrow U(\mathfrak{g})$$

which sends  $x_\alpha \otimes_\pi 1$  to  $x_\alpha^p - y_\alpha$ . Moreover, for every  $x \in \mathfrak{g}$ ,

$$(\operatorname{ad} x_\alpha^p)(x) = (\operatorname{ad} x_\alpha)^p(x) = (\operatorname{ad} y_\alpha)(x).$$

<sup>54</sup>In this paragraph, the editors changed the order, stating the result first and then giving the detailed proof.

<sup>55</sup>Thus  $xr \otimes_\pi 1 = x \otimes_\pi r^p$  and  $r \cdot (x \otimes_\pi 1) = x \otimes_\pi r$ , for  $x \in \mathfrak{g}$  and  $r \in R$ .



It follows that  $\gamma$  maps  $\mathfrak{g} \otimes_\pi R$  into the center of  $U(\mathfrak{g})$ . For  $x \in \mathfrak{g}$ , put

$$x^{(p)} = x^p - \gamma(x \otimes_\pi 1).$$

Then  $x_\alpha^{(p)} = y_\alpha$  for every  $\alpha$ . If

$$x = \sum_\alpha \lambda_\alpha x_\alpha,$$

formula 5.1 (ii), applied inductively to the number of indices  $\alpha$  for which  $\lambda_\alpha \neq 0$ , gives

$$x^p - \sum_\alpha \lambda_\alpha^p x_\alpha^p \in \mathfrak{g}.$$

Denote this element by  $z$ . Then

$$x^{(p)} = \sum_\alpha \lambda_\alpha^p y_\alpha + z,$$

and hence  $x^{(p)} \in \mathfrak{g}$ .

It is clear that

$$(\lambda x)^{(p)} = \lambda^p x^{(p)}.$$

Since  $\gamma(x \otimes_\pi 1)$  is central, we have  $\text{ad } x^{(p)} = \text{ad } x^p$ . Jacobson's first formula, 5.1 (i), therefore yields

$$\text{ad } x^{(p)} = (\text{ad } x)^p.$$

Finally, Jacobson's second formula, 5.1 (ii), shows that the map  $x \mapsto x^{(p)}$  satisfies condition (ii) of 5.2. It thus makes  $\mathfrak{g}$  into a restricted  $p$ -Lie algebra. This proves the proposition.

**5.3.3. Proposition.** *Let  $\mathfrak{g}$  be a restricted  $p$ -Lie algebra over  $R$  whose underlying module is free with basis  $(x_\alpha)$ . Then*

$$j : \mathfrak{g} \longrightarrow U_p(\mathfrak{g})$$

*is injective. If  $z_\alpha = j(x_\alpha)$ , then  $U_p(\mathfrak{g})$  has as a basis the monomials*

$$\prod_\alpha z_\alpha^{n_\alpha}, \quad 0 \leq n_\alpha < p.$$

*Here the  $n_\alpha$  are zero except for finitely many  $\alpha$ ; the basis is assumed to be totally ordered, and the products are taken in increasing order.*

Identify  $\mathfrak{g}$  with a submodule of the enveloping algebra  $U(\mathfrak{g})$  by means of the canonical map  $i$ . For every family

$$n = (n_\alpha)$$

of nonnegative integers, all but finitely many zero, put

$$|n| = \sum_\alpha n_\alpha, \quad x^n = \prod_\alpha x_\alpha^{n_\alpha}.$$

Write

$$n_\alpha = m_\alpha + p\ell_\alpha, \quad 0 \leq m_\alpha < p,$$

and put

$$T_n = \prod_\alpha x_\alpha^{m_\alpha} \beta(x_\alpha)^{\ell_\alpha},$$

where

$$\beta(x) = x^p - x^{(p)}$$

is the map  $\mathfrak{g} \rightarrow U(\mathfrak{g})$  defined in 5.3.1.

For  $r \in \mathbb{N}$ , let  $U^r$  be the  $R$ -submodule of  $U(\mathfrak{g})$  generated by the  $x^n$  for which  $|n| \leq r$ . Since the graded ring

$$\bigoplus_r U^r / U^{r-1}$$

is commutative (cf. [BLie], I, §2.6), for every  $n$  we have

$$T_n - \prod_{\alpha} x_{\alpha}^{n_{\alpha}} \in U^{|n|-1}.$$

For every  $s \in \mathbb{N}$ , the  $x^n$  with  $|n| = s$  form a basis of  $U^s / U^{s-1}$  by the Poincaré–Birkhoff–Witt theorem (*loc. cit.*, §2.7); the same is therefore true of the  $T_n$  with  $|n| = s$ .

As  $s = |n|$  varies, the  $T_n$  consequently form a basis of  $U(\mathfrak{g})$ . The kernel  $J$  of the canonical map

$$U(\mathfrak{g}) \longrightarrow U_p(\mathfrak{g})$$

is the left ideal of  $U(\mathfrak{g})$  generated by the central elements  $\beta(x_{\alpha})$ , by 5.3.1. The  $T_n$  for which  $\ell = (\ell_{\alpha}) \neq 0$  therefore form a basis of  $J$ , while the  $T_n$  for which  $n_{\alpha} < p$  for every  $\alpha$  form a basis of

$$U_p(\mathfrak{g}) = U(\mathfrak{g}) / J.$$

### 5.3.3 bis.

Let  $\mathfrak{g}$  be a restricted  $p$ -Lie algebra over  $R$ , and let  $f : R \rightarrow R'$  be an extension of the base ring. We claim that the  $R'$ -module

$$R' \otimes_R \mathfrak{g}$$

admits one and only one restricted  $p$ -Lie algebra structure such that

$$[\lambda \otimes x, \mu \otimes y] = \lambda \mu \otimes [x, y], \quad (\lambda \otimes x)^{(p)} = \lambda^p \otimes x^{(p)}. \quad (*)$$

In particular, it will follow that the functor

$$\mathfrak{g} \longmapsto R' \otimes_R \mathfrak{g}$$

is left adjoint to restriction of scalars from  $R'$  to  $R$ .

The uniqueness of the structure defined by  $(*)$  is clear; we prove its existence. If  $\mathfrak{g}$  is free with basis  $(x_{\alpha})$ , Proposition 5.3.2 gives a unique restricted  $p$ -Lie algebra structure on the Lie algebra  $R' \otimes_R \mathfrak{g}$  such that

$$(1 \otimes x_{\alpha})^{(p)} = 1 \otimes x_{\alpha}^{(p)}.$$

This is the structure sought.

Suppose now that  $\mathfrak{g}$  is arbitrary. There is a restricted  $p$ -Lie algebra  $L_0$ , free as an  $R$ -module, together with a surjective homomorphism

$$q_0 : L_0 \longrightarrow \mathfrak{g}.$$

For example, we may take

$$L_0 = R \otimes_{\mathbb{F}_p} \mathfrak{g}$$

and let  $q_0$  be the homomorphism

$$\lambda \otimes x \longmapsto \lambda x;$$

here  $\mathfrak{g}$  is free over the prime field  $\mathbb{F}_p$ . The kernel of  $q_0$  is a  $p$ -ideal of  $L_0$ , that is, an ideal of the Lie algebra  $L_0$  stable under  $x \mapsto x^{(p)}$ . Hence there is also a restricted  $p$ -Lie algebra  $L_1$ , free as an  $R$ -module, and a homomorphism  $q_1 : L_1 \rightarrow L_0$  whose image is  $\text{Ker } q_0$ . This gives the exact sequence

$$L_1 \xrightarrow{q_1} L_0 \xrightarrow{q_0} \mathfrak{g} \longrightarrow 0.$$

and therefore an exact sequence of  $R'$ -Lie algebras

$$R' \otimes_R L_1 \xrightarrow{R' \otimes_R q_1} R' \otimes_R L_0 \xrightarrow{R' \otimes_R q_0} R' \otimes_R \mathfrak{g} \longrightarrow 0.$$

Since  $R' \otimes_R q_1$  is manifestly a homomorphism of restricted  $p$ -Lie algebras, the kernel of  $R' \otimes_R q_0$  is a  $p$ -ideal. Thus the symbolic  $p$ -th power operation on  $R' \otimes_R L_0$  induces, by passage to the quotient, a map from  $R' \otimes_R \mathfrak{g}$  to itself (use formula (ii) of 5.2). This map equips  $R' \otimes_R \mathfrak{g}$  with the required restricted  $p$ -Lie algebra structure.

#### 5.3.4.

The canonical map

$$j_{\mathfrak{g}} : \mathfrak{g} \longrightarrow U_p(\mathfrak{g})$$

induces, for every extension  $R \rightarrow R'$  of the base ring, a homomorphism

$$R' \otimes_R j_{\mathfrak{g}} : R' \otimes_R \mathfrak{g} \longrightarrow R' \otimes_R U_p(\mathfrak{g}).$$

It therefore induces a homomorphism

$$h : U_p(R' \otimes_R \mathfrak{g}) \longrightarrow R' \otimes_R U_p(\mathfrak{g})$$

such that

$$h \circ j_{R' \otimes_R \mathfrak{g}} = R' \otimes_R j_{\mathfrak{g}}.$$

The universal properties of  $R' \otimes_R \mathfrak{g}$  and of the restricted enveloping algebra show immediately that  $h$  is an isomorphism. We may therefore identify

$$U_p(R' \otimes_R \mathfrak{g}) \simeq R' \otimes_R U_p(\mathfrak{g}).$$

In particular, let  $r \in R$ , and let  $R' = R_r$  be the localized ring. Then

$$\mathfrak{g}_r = R_r \otimes_R \mathfrak{g}$$

has a canonical restricted  $p$ -Lie algebra structure over  $R_r$ . Consequently, the sheaf  $\tilde{\mathfrak{g}}$  on  $\text{Spec } R$  is a quasi-coherent restricted  $p$ -Lie algebra on  $\text{Spec } R$ . Moreover,

$$U_p^{R_r}(\mathfrak{g}_r) \simeq U_p^R(\mathfrak{g})_r,$$

so the sheaf associated with the presheaf

$$V \longmapsto U_p(\Gamma(V, \tilde{\mathfrak{g}}))$$

is quasi-coherent.

**Definition.** More generally, let  $S$  be a scheme of characteristic  $p$ , and let  $\mathcal{G}$  be a quasi-coherent restricted  $p$ -Lie algebra over  $\mathcal{O}_S$ . The sheaf associated with the presheaf

$$V \longmapsto U_p(\Gamma(V, \mathcal{G}))$$

is quasi-coherent. It is denoted  $\mathcal{U}_p(\mathcal{G})$  and is called the *restricted enveloping algebra* of  $\mathcal{G}$ . If  $V$  is affine, then

$$U_p(\Gamma(V, \mathcal{G})) \simeq \Gamma(V, \mathcal{U}_p(\mathcal{G})).$$

#### 5.4.

The universal property of  $U_p(\mathfrak{g})$  implies that  $U_p(\mathfrak{g})$  is functorial in  $\mathfrak{g}$ . Every homomorphism of restricted  $p$ -Lie algebras

$$\varphi : \mathfrak{g} \longrightarrow \mathfrak{h}$$

induces a unique homomorphism of unital algebras  $U_p(\varphi)$  such that

$$j_{\mathfrak{h}} \circ \varphi = U_p(\varphi) \circ j_{\mathfrak{g}}.$$

Here are some examples.

a) If  $\mathfrak{h} = 0$ , then  $U_p(\mathfrak{h})$  identifies with the base ring, and  $U_p(\varphi)$  is an algebra homomorphism

$$\varepsilon_{\mathfrak{g}} : U_p(\mathfrak{g}) \longrightarrow R$$

called the *augmentation*.

b) Let now  $\mathfrak{h} = \mathfrak{g}^\circ$ , the Lie algebra opposite to  $\mathfrak{g}$ . Thus  $\mathfrak{g}^\circ$  has the same underlying module and the same symbolic  $p$ -th power as  $\mathfrak{g}$ , while its bracket is the negative of the bracket of  $\mathfrak{g}$ . We may identify  $U_p(\mathfrak{g}^\circ)$  with the algebra opposite to  $U_p(\mathfrak{g})$ . Moreover, the isomorphism

$$x \longmapsto -x$$

from  $\mathfrak{g}$  onto  $\mathfrak{g}^\circ$  induces an isomorphism

$$c_{\mathfrak{g}} : U_p(\mathfrak{g}) \longrightarrow U_p(\mathfrak{g}^\circ) \simeq U_p(\mathfrak{g})^\circ.$$

We call  $c_{\mathfrak{g}}$  the *antipode* of  $U_p(\mathfrak{g})$ .

c) Finally, let  $\mathfrak{f}$  and  $\mathfrak{g}$  be two restricted  $p$ -Lie algebras, and let  $\mathfrak{h} = \mathfrak{f} \times \mathfrak{g}$  be their product. Its underlying  $R$ -module is the direct product, and

$$[(x, y), (x', y')] = ([x, x'], [y, y']), \quad (x, y)^{(p)} = (x^{(p)}, y^{(p)}).$$

If

$$h_1 : \mathfrak{f} \rightarrow \mathfrak{k}, \quad h_2 : \mathfrak{g} \rightarrow \mathfrak{k}$$

are restricted  $p$ -Lie algebra homomorphisms satisfying

$$[h_1(x), h_2(y)] = 0 \quad (x \in \mathfrak{f}, y \in \mathfrak{g}),$$

then

$$h_1 + h_2 : (x, y) \longmapsto h_1(x) + h_2(y)$$

is a restricted  $p$ -Lie algebra homomorphism. Conversely, every homomorphism  $\mathfrak{f} \times \mathfrak{g} \rightarrow \mathfrak{k}$  is of this form. This characterizes  $\mathfrak{f} \times \mathfrak{g}$  by a universal property.

For example, the maps

$$h_1 : x \longmapsto j_{\mathfrak{f}}(x) \otimes 1, \quad h_2 : y \longmapsto 1 \otimes j_{\mathfrak{g}}(y)$$

induce a homomorphism  $h_1 + h_2$  from  $\mathfrak{f} \times \mathfrak{g}$  to the restricted  $p$ -Lie algebra underlying

$$U_p(\mathfrak{f}) \otimes U_p(\mathfrak{g}).$$

The universal properties of  $\mathfrak{f} \times \mathfrak{g}$  and of the restricted enveloping algebras show that this homomorphism extends to an isomorphism

$$\varphi : U_p(\mathfrak{f} \times \mathfrak{g}) \xrightarrow{\sim} U_p(\mathfrak{f}) \otimes U_p(\mathfrak{g}).$$

**Definition.** If  $\mathfrak{f} = \mathfrak{g}$ , the diagonal map

$$\delta : x \longmapsto (x, x)$$

from  $\mathfrak{g}$  to  $\mathfrak{g} \times \mathfrak{g}$  induces a homomorphism from  $U_p(\mathfrak{g})$  to  $U_p(\mathfrak{g} \times \mathfrak{g})$ . We denote by

$$\Delta_{\mathfrak{g}} : U_p(\mathfrak{g}) \longrightarrow U_p(\mathfrak{g}) \otimes U_p(\mathfrak{g})$$

the composite of this homomorphism with  $\varphi$ .<sup>56</sup>

It is then easy to see that  $\Delta_{\mathfrak{g}}$ , together with the multiplication of  $U_p(\mathfrak{g})$ , makes  $U_p(\mathfrak{g})$  an  $R$ -coalgebra in groups (cf. 3.2), with augmentation  $\varepsilon_{\mathfrak{g}}$  and antipode  $c_{\mathfrak{g}}$ .

### 5.5.<sup>57</sup>

Let  $S$  be a scheme of characteristic  $p$ . First let  $\mathcal{U}$  be an  $\mathcal{O}_S$ -coalgebra in groups and put

$$G = \mathrm{Spec}^* \mathcal{U}.$$

We saw in 3.2.3 that, for every  $T \rightarrow S$ ,  $(\mathrm{Lie} G)(T)$  is the Lie subalgebra of  $\Gamma(T, \mathcal{U}_T)$  consisting of primitive elements. If  $x$  is primitive, then

$$\Delta(x^p) = x^p \otimes 1 + 1 \otimes x^p,$$

since  $\binom{p}{i} \equiv 0 \pmod{p}$  for  $0 < i < p$ . Thus  $x^p$  is again primitive. By 5.1 and 5.2, the map  $x \mapsto x^p$  makes  $(\mathrm{Lie} G)(T)$  into an  $\mathcal{O}(T)$ -restricted  $p$ -Lie algebra.

Now let  $\mathcal{L}$  be a quasi-coherent restricted  $p$ -Lie algebra over  $\mathcal{O}_S$ . As  $V$  ranges over the open subsets of  $S$ , the group-coalgebra structures defined above on

$$U_p(\Gamma(V, \mathcal{L}))$$

induce on the associated sheaf  $\mathcal{U}_p(\mathcal{L})$  a structure of  $\mathcal{O}_S$ -coalgebra in groups. Moreover, for every  $S$ -scheme  $T$ , there is an isomorphism

$$\mathcal{U}_p(\mathcal{L}_T) \xrightarrow{\sim} \mathcal{U}_p(\mathcal{L})_T.$$

Let  $\mathrm{Prim} \mathcal{U}_p(\mathcal{L})$  denote the subpresheaf which assigns to every open subset  $V$  the primitive elements of  $\mathcal{U}_p(\mathcal{L})(V)$ ; this is readily seen to be a sheaf. As  $V$  ranges over the open subsets of  $S$ , the composites

$$\Gamma(V, \mathcal{L}) \xrightarrow{j} \mathrm{Prim} U_p(\Gamma(V, \mathcal{L})) \longrightarrow \mathrm{Prim} \mathcal{U}_p(\mathcal{L})(V)$$

define a morphism

$$\mathcal{L} \longrightarrow \mathrm{Prim} \mathcal{U}_p(\mathcal{L}),$$

again denoted  $j$  or  $j_{\mathcal{L}}$ . It in turn defines a morphism of  $\mathcal{O}_S$ -restricted  $p$ -Lie algebras

$$\mathcal{W}(\mathcal{L}) \longrightarrow \mathrm{Prim} \mathcal{W}(\mathcal{U}_p(\mathcal{L}))$$

(cf. 3.2.3).

**Proposition 5.5.1.** *Let  $\mathcal{L}$  be an  $\mathcal{O}_S$ -restricted  $p$ -Lie algebra, locally free as an  $\mathcal{O}_S$ -module. Then  $j_{\mathcal{L}}$  induces an isomorphism of  $\mathcal{O}_S$ -restricted  $p$ -Lie algebras*

$$\mathcal{W}(\mathcal{L}) \xrightarrow{\sim} \mathrm{Prim} \mathcal{W}(\mathcal{U}_p(\mathcal{L})).$$

<sup>56</sup>Thus  $\Delta_{\mathfrak{g}}(x) = x \otimes 1 + 1 \otimes x$  for every  $x \in \mathfrak{g}$ ; in particular,  $\Delta_{\mathfrak{g}}$  is cocommutative.

<sup>57</sup>The editors reorganized §5.4.1 of the original into this §5.5. Proposition 5.5.1 combines results from Section 5 and Proposition 3.2.3 and incorporates the original Lemma 7.3; the proof of Proposition 5.5.3(ii) expands the proof of (i)  $\Rightarrow$  (ii) in the original Theorem 7.4.

*Proof.* Let  $T$  be an  $S$ -scheme. In view of the identification

$$\mathcal{U}_p(\mathcal{L}_T) = \mathcal{U}_p(\mathcal{L})_T,$$

we must show that

$$\Gamma(T, \mathcal{L}_T) \longrightarrow \text{Prim } \Gamma(T, \mathcal{U}_p(\mathcal{L}_T))$$

is bijective. Replacing  $S$  by  $T$ , we reduce to  $T = S$ ; it then suffices to prove that the morphism of sheaves

$$j_{\mathcal{L}} : \mathcal{L} \longrightarrow \text{Prim } \mathcal{U}_p(\mathcal{L})$$

is an isomorphism. The question is local on  $S$ , so suppose that  $S = \text{Spec } R$  and that  $\mathcal{L}$  is the sheaf associated with an  $R$ -restricted  $p$ -Lie algebra  $L$  with a totally ordered basis  $(x_\alpha)$ .

As in 5.3.3, let  $z_\alpha$  be the image of  $x_\alpha$  in  $U = U_p(L)$ . For a family  $n = (n_\alpha)$ , with  $0 \leq n_\alpha \leq p-1$  and only finitely many nonzero terms, put

$$z^{(n)} = \prod_{\alpha} \frac{z_{\alpha}^{n_{\alpha}}}{n_{\alpha}!},$$

the product being taken in increasing order. Since

$$\Delta(z_{\alpha}) = z_{\alpha} \otimes 1 + 1 \otimes z_{\alpha},$$

one obtains

$$\Delta(z^{(n)}) = \sum_r z^{(n-r)} \otimes z^{(r)},$$

where the sum is over the finite set of  $r$  satisfying

$$0 \leq r_{\alpha} \leq n_{\alpha} \quad \text{for every } \alpha.$$

The  $z^{(n)}$ , respectively the  $z^{(n)} \otimes z^{(m)}$ , form a basis of  $U$ , respectively of  $U \otimes U$ . It follows that

$$\Delta(u) = u \otimes 1 + 1 \otimes u$$

holds for  $u \in U$  if and only if  $u$  is a linear combination of the  $z_{\alpha}$ . This proves the proposition.  $\square$

**Remark 5.5.2.** Recall from 3.2.2 and 3.2.3 that the group  $S$ -functor

$$G = \text{Spec}^* \mathcal{U}_p(\mathcal{L})$$

is very good and that

$$\text{Lie } G = \text{Prim } \mathcal{W}(\mathcal{U}_p(\mathcal{L})).$$

Thus Proposition 5.5.1 says that  $j_{\mathcal{L}}$  induces an isomorphism

$$\mathcal{W}(\mathcal{L}) \xrightarrow{\sim} \text{Lie } G.$$

If, moreover,  $\mathcal{L}$  is locally free of finite rank, then  $\mathcal{U}_p(\mathcal{L})$  is finite locally free over  $\mathcal{O}_S$  by 5.3.3. Hence  $\text{Spec}^* \mathcal{U}_p(\mathcal{L})$  is represented by the  $S$ -group

$$\mathfrak{G}_p(\mathcal{L}) = \text{Spec } \mathcal{U}_p(\mathcal{L})^*$$

(cf. 3.2.2.1). We obtain the following more precise statement.

**Proposition 5.5.3.** *Let  $\mathcal{L}$  be an  $\mathcal{O}_S$ -restricted  $p$ -Lie algebra, locally free of finite rank as an  $\mathcal{O}_S$ -module. Put*

$$\mathcal{A} = \mathcal{U}_p(\mathcal{L})^*, \quad G = \mathfrak{G}_p(\mathcal{L}) = \operatorname{Spec} \mathcal{A}.$$

*Then:*

(i)  $j_{\mathcal{L}}$  induces an isomorphism

$$\mathcal{W}(\mathcal{L}) \xrightarrow{\sim} \operatorname{Lie} G$$

*of  $\mathcal{O}_S$ -restricted  $p$ -Lie algebras.*

(ii) *Let  $\mathcal{I}$  be the augmentation ideal of  $\mathcal{A}$  and put  $\omega_G = \mathcal{I}/\mathcal{I}^2$  (cf. II, 4.11.4). Then*

$$\omega_G \simeq \mathcal{L}^* = \operatorname{Hom}_{\mathcal{O}_S}(\mathcal{L}, \mathcal{O}_S).$$

*In particular,  $\omega_G$  is locally free of finite rank, and  $\omega_{G/S}^* \simeq \mathcal{L}$ .*

*Proof.* Part (i) follows from Remark 5.5.2. We prove (ii). Let  $\eta_{\mathcal{U}}$  and  $\varepsilon_{\mathcal{U}}$  denote the unit and augmentation of  $\mathcal{U} = \mathcal{U}_p(\mathcal{L})$ , and let  $\eta_{\mathcal{A}}$  and  $\varepsilon_{\mathcal{A}}$  denote those of  $\mathcal{A}$ . Put

$$\mathcal{J} = \ker \varepsilon_{\mathcal{U}}.$$

Then

$$\mathcal{U} = \eta_{\mathcal{U}}(\mathcal{O}_S) \oplus \mathcal{J}. \quad (1)$$

Let  $\delta$  be the morphism defined by the diagram

$$\begin{array}{ccc} \mathcal{J} & \xrightarrow{\delta} & \mathcal{J} \otimes \mathcal{J} \\ \tau \downarrow & & \uparrow \pi \\ \mathcal{U} & \xrightarrow{\Delta} & \mathcal{U} \otimes \mathcal{U} \end{array}$$

where  $\tau$  and  $\pi$  are the inclusion and projection deduced from decomposition (1). There is an exact sequence

$$(*) \quad 0 \longrightarrow \mathcal{L}^* \xrightarrow{j_{\mathcal{L}}} \mathcal{J} \xrightarrow{\delta} \mathcal{J} \otimes \mathcal{J}.$$

By 5.3.3, the  $\mathcal{O}_S$ -module  $\mathcal{J}/\mathcal{L}$  is locally free, and by 5.5.1 the preceding sequence remains exact after every base change. Consequently,  $\delta$  induces an isomorphism from  $\mathcal{J}/\mathcal{L}$  onto a locally direct summand  $Q$  of  $\mathcal{J} \otimes \mathcal{J}$ . By duality, the exact sequence therefore becomes

$$(**) \quad 0 \longleftarrow \mathcal{L} \xleftarrow{t_{j_{\mathcal{L}}}} \mathcal{I} \xleftarrow{t_{\delta}} \mathcal{I} \otimes \mathcal{I}.$$

Decomposition (1) corresponds by duality to

$$\mathcal{A} = \eta_{\mathcal{A}}(\mathcal{O}_S) \oplus \mathcal{I}. \quad (2)$$

The transpose of  $\Delta$  is the multiplication

$$m_{\mathcal{A}} : \mathcal{A} \otimes \mathcal{A} \longrightarrow \mathcal{A}.$$

Since  $\mathcal{I}$  is an ideal of  $\mathcal{A}$ ,  $m_{\mathcal{A}}$  sends  $\mathcal{I} \otimes \mathcal{I}$  into  $\mathcal{I}$ . More precisely, decomposition (2) gives the commutative square

$$\begin{array}{ccc} \mathcal{I} & \xleftarrow{m'} & \mathcal{I} \otimes \mathcal{I} \\ t_{\tau} \uparrow & & \downarrow t_{\pi} \\ \mathcal{A} & \xleftarrow{m_{\mathcal{A}}} & \mathcal{A} \otimes \mathcal{A} \end{array}$$

This shows that the restriction  $m'$  of  $m_{\mathcal{A}}$  to  $\mathcal{I} \otimes \mathcal{I}$  is the transpose of  $\delta$ . The dual exact sequence therefore yields

$$\mathcal{I}/\mathcal{I}^2 \simeq \mathcal{L}^*,$$

which proves the proposition.  $\square$

## 6. The $p$ -Lie Algebra of an $S$ -Group Scheme

Let  $S$  be a scheme of characteristic  $p > 0$ . In 5.5 we associated with every quasi-coherent  $\mathcal{O}_S$ - $p$ -Lie algebra  $\mathcal{L}$  the group  $S$ -functor

$$\mathfrak{G}_p(\mathcal{L}) = \mathrm{Spec}^* \mathcal{U}_p(\mathcal{L}).$$

We shall now see that, for every  $S$ -group scheme  $G$ , the  $\mathcal{O}_S$ -Lie algebra  $\mathrm{Lie}(G/S)$  defined in II, 4.11 is naturally endowed with the structure of an  $\mathcal{O}_S$ - $p$ -Lie algebra.

### 6.1.

First identify  $\mathrm{Lie}(G/S)(S)$  and  $\mathrm{Lie}(\mathrm{Aut} G/S)(S)$ , respectively, with Lie subalgebras of  $\mathrm{U}(G)$  and  $\mathrm{Dif}_{G/S}$  by means of the injections  $\alpha$  and  $\beta$  of 2.5. Thus  $\mathrm{Lie}(\mathrm{Aut} G/S)(S)$  is identified with the  $\Gamma(S, \mathcal{O}_S)$ -Lie algebra of  $S$ -derivations of  $\mathcal{O}_G$ . By 5.2, the latter is a restricted  $p$ -Lie subalgebra of  $\mathrm{Dif}_{G/S}$ .

On the other hand, the image of

$$L = \mathrm{Lie}(G/S)(S)$$

under the injective algebra homomorphism

$$\ell : \mathrm{U}(G) \longrightarrow \mathrm{Dif}_{G/S}, \quad d \longmapsto {}_G d,$$

consists of the left-invariant derivations (cf. 2.2, editor's note (17), 2.4, and 2.5). If  $x \in L$ , then  $\ell(x)^p = \ell(x^p)$  by the cited passages. Since  $\ell(x)^p$  is again a derivation,  $x^p$  belongs to  $\mathrm{Lie}(G/S)(S)$ . Hence

*$\mathrm{Lie}(G/S)(S)$  is a restricted  $p$ -Lie subalgebra of the infinitesimal algebra  $\mathrm{U}(G)$ .*

58

#### 6.1.1.

Let  $\varphi : G \rightarrow H$  be a homomorphism of  $S$ -group schemes. The homomorphisms  $\mathrm{Lie}(\varphi/S)(S)$  and  $\mathrm{U}(\varphi)$  are compatible with the identifications of  $\mathrm{Lie}(G/S)(S)$  and  $\mathrm{Lie}(H/S)(S)$  with restricted  $p$ -Lie subalgebras of  $\mathrm{U}(G)$  and  $\mathrm{U}(H)$ . Since  $\mathrm{U}(\varphi)$  is an algebra homomorphism,  $\mathrm{Lie}(\varphi/S)(S)$  is a homomorphism of restricted  $p$ -Lie algebras.

Likewise, if  $s : T \rightarrow S$  is a base change, the map from  $\mathrm{Lie}(G/S)(S)$  to  $\mathrm{Lie}(G/S)(T)$  induced by  $s$  is a homomorphism of restricted  $p$ -Lie algebras. Equivalently, the functor  $\mathrm{Lie}(G/S)$  is endowed with a structure of  $\mathcal{O}_S$ - $p$ -Lie algebra. In particular, as  $T$  ranges over the open subsets of  $S$ , the sheaf  $\mathrm{Lie}(G/S)$  is endowed with a structure of  $\mathcal{O}_S$ - $p$ -Lie algebra.

### 6.2.

Following an idea of Demazure, we shall generalize the preceding construction to certain group  $S$ -functors which need not be representable. We first give another definition of the symbolic  $p$ -th power in the Lie algebra of an  $S$ -group scheme  $G$ .

<sup>58</sup>One may also prove directly, without passing through  $\mathrm{Dif}_{G/S}$ , that the Lie algebra of derivations of  $G$  at the origin (isomorphic to  $\mathrm{Lie}(G/S)(S)$  by 2.5) is stable under taking  $p$ -th powers in  $\mathrm{U}(G)$ . This is done in 6.2 below.



Let  $D$  be a derivation of  $G$  at the origin.<sup>59</sup> By 1.2.1,  $D$  is the composite of the  $S$ -derivation

$$\delta = (\tau, \partial_t)$$

of the zero section  $\tau : S \rightarrow I_S$ , and a morphism  $x : I_S \rightarrow G$  such that  $x \circ \tau = \varepsilon$ , that is,  $x \in \text{Lie}(G/S)(S)$ . By the definition in 2.1,  $D^p$  is the following composite deviation:

$$S \simeq \underbrace{S \times S \times \cdots \times S}_p \xrightarrow{\delta \times \cdots \times \delta} I_S \times \cdots \times I_S \xrightarrow{x \times \cdots \times x} G \times \cdots \times G \xrightarrow{m^{(p)}} G$$

Here  $m^{(p)}$  is induced by the multiplication  $m : G \times G \rightarrow G$ . Since  $I_S \times \cdots \times I_S$  is affine over  $S$ , with affine algebra

$$\mathcal{B} = \mathcal{O}_S[d_1, \dots, d_p]/(d_1^2, \dots, d_p^2),$$

the deviation  $\delta \times \cdots \times \delta$  is defined by the homomorphism of  $\mathcal{O}_S$ -modules

$$\phi : \mathcal{B} \longrightarrow \mathcal{O}_S$$

which sends the monomial  $d_1 d_2 \cdots d_p$  to 1, and all the other monomials  $d_{i_1} \cdots d_{i_r}$ , for  $0 \leq r < p$ , to 0.

Let  $\text{pr}_i$  denote the projection of  $I_S^p$  onto its  $i$ -th factor, and let  $x_i$  be the image of  $x$  in  $G(I_S^p)$  under  $G(\text{pr}_i)$ . Then

$$m^{(p)} \circ (x \times \cdots \times x)$$

is precisely the product  $x_1 x_2 \cdots x_p$ . Consequently,  $D^p$  is also the composite deviation

$$S \xrightarrow{\delta \times \cdots \times \delta} I_S \times \cdots \times I_S \xrightarrow{x_1 x_2 \cdots x_p} G.$$

This description gives another proof that  $D^p$  is a derivation of  $G$  at the origin. Since  $G$  is a very good group (II, 4.11), the images  $G(\text{pr}_1)(x)$  and  $G(\text{pr}_2)(x)$  of  $x$  in  $G(I_S \times I_S)$  commute. It follows that the elements  $x_i \in G(I_S^p)$  commute pairwise. Hence, for every permutation  $\gamma$  of the factors of  $I_S^p$ ,

$$(x_1 \cdots x_p) \circ \gamma = x_1 \cdots x_p.$$

Thus  $x_1 \cdots x_p$  factors through the canonical projection from  $I_S^p$  to the symmetric product  $\Sigma^p I_S$  (cf. 4.2).

The affine algebra of  $\Sigma^p I_S$  is the subalgebra  $\mathcal{A}$  of  $\mathcal{B}$  having as an  $\mathcal{O}_S$ -basis the elementary symmetric functions

$$1 = \sigma_0, \sigma_1, \dots, \sigma_p$$

in  $d_1, \dots, d_p$ . Let  $\kappa : \mathcal{A} \hookrightarrow \mathcal{B}$  be the inclusion, and let

$$\pi : \mathcal{A} \longrightarrow \mathcal{O}_S[t]/(t^2)$$

be the homomorphism of  $\mathcal{O}_S$ -algebras which kills  $\sigma_1, \dots, \sigma_{p-1}$  and sends  $\sigma_p$  to  $t$ . Then

$$\phi \circ \kappa = \partial_t \circ \pi,$$

where  $\partial_t : \mathcal{O}_S[t]/(t^2) \rightarrow \mathcal{O}_S$  kills 1 and sends  $t$  to 1. If

$$i : I_S \hookrightarrow \Sigma^p I_S$$

denotes the closed immersion defined by  $\pi$ , we obtain the commutative diagram

---

<sup>59</sup>The editors expanded the argument in the original text in what follows.

$$\begin{array}{ccccc}
& & D^p & & \\
& \nearrow & & \searrow & \\
S & \xrightarrow{\delta \times \cdots \times \delta} & I_S^p & \xrightarrow{x_1 \cdots x_p} & G \\
\downarrow \delta & & \downarrow \text{can.} & & \parallel \\
I_S & \xrightarrow{i} & \Sigma^p I_S & \xrightarrow{\quad} & G \\
& \searrow & & \nearrow & \\
& & y & & 
\end{array}$$

which shows that  $D^p$  has the form  $y \circ \delta$ , and is therefore indeed a derivation of  $G$  at the origin.

### 6.3.

Let  $\mathfrak{S}_p$  be the symmetric group of degree  $p$ , and let  $I_S^p \times \mathfrak{S}_p$  denote the disjoint union of a family of copies of  $I_S^p$  indexed by  $\mathfrak{S}_p$ . Let

$$\pi : I_S^p \times \mathfrak{S}_p \longrightarrow I_S^p$$

be the canonical projection, and let

$$\mu : I_S^p \times \mathfrak{S}_p \longrightarrow I_S^p$$

be the morphism defining the action of  $\mathfrak{S}_p$  on  $I_S^p$ : if  $\rho \in \mathfrak{S}_p$ , the restriction of  $\mu$  to  $I_S^p \times \rho$  has  $\text{pr}_{\rho(j)}$  as its  $j$ -th component.

**Definition.** A functor

$$X : (\text{Sch}/S)^\circ \longrightarrow (\text{Ens})$$

satisfies *condition (F)* if:

- a)  $X$  takes finite disjoint unions to direct products;
- b) for every  $S$ -scheme  $T$ , the following sequence is exact:

$$X(T \times \Sigma^p I_S) \longrightarrow X(T \times I_S^p) \xrightleftharpoons[X(\text{id}_T \times \mu)]{X(\text{id}_T \times \pi)} X(T \times I_S^p \times \mathfrak{S}_p).$$

Every  $S$ -scheme satisfies (F). If  $\mathcal{F}$  is an  $\mathcal{O}_S$ -module, then  $\mathcal{W}(\mathcal{F})$  satisfies (F). Every projective limit of functors satisfying (F) again satisfies (F). Finally, if  $Y$  satisfies (F) and  $X$  is an arbitrary  $S$ -functor, then  $\text{Hom}_S(X, Y)$  satisfies (F).

Let  $X$  be a very good group (II, 4.10) satisfying condition (F). Let  $x : I_S \rightarrow X$  be a morphism extending the unit section of  $X$ , and use the notation of 6.2. As above,

$$x_1 \cdots x_p : I_S^p \longrightarrow X$$

factors through  $\Sigma^p I_S$ :

$$\begin{array}{ccc}
I_S^p & \xrightarrow{x_1 \cdots x_p} & X \\
\searrow \text{can.} & & \nearrow \Sigma^p(x) \\
& \Sigma^p I_S & 
\end{array}$$

and therefore defines, by composition, a morphism

$$x^{(p)} : I_S \xrightarrow{i} \Sigma^p I_S \xrightarrow{\Sigma^p(x)} X$$

which we call the *symbolic  $p$ -th power* of  $x$ .

The endomorphism

$$x \longmapsto x^{(p)}$$

of  $\mathrm{Lie}(G/S)(S)$  is plainly compatible with base change and functorial in  $G$ . It would be interesting to know for which  $G$  this endomorphism makes  $\mathrm{Lie}(G/S)(S)$  into a restricted  $p$ -Lie algebra.

#### 6.4.

The preceding definition of the symbolic  $p$ -th power is particularly well suited to computation. Here are some examples.

##### 6.4.1.

Let  $M$  be an “abstract” abelian group, and let  $D_S(M)$  be the diagonalizable  $S$ -group of type  $M$  (I, 4.4.2). For every  $S$ -scheme  $T$ ,

$$D_S(M)(T) = \mathrm{Hom}_{(\mathrm{Ab})}(M, \mathcal{O}(T)^\times).$$

Let  $x \in \mathrm{Lie}(D_S(M)/S)(S)$ , that is, an abelian-group homomorphism

$$x : M \longrightarrow \Gamma(S, \mathcal{O}_S + d\mathcal{O}_S)^\times$$

of the form

$$m \longmapsto 1 + d\xi(m), \quad \xi \in \mathrm{Hom}_{(\mathrm{Ab})}(M, \mathcal{O}(S)).$$

With the notation of 6.2 and 6.3, the product  $x_1 \cdots x_p$  associates with  $m \in M$  the expression

$$(1 + d_1\xi(m)) \cdots (1 + d_p\xi(m)),$$

that is,

$$1 + \sigma_1\xi(m) + \sigma_2\xi(m)^2 + \cdots + \sigma_p\xi(m)^p.$$

This expression belongs to  $\mathcal{O}(\Sigma^p I_S)$ . Projecting it to  $\mathcal{O}(S)[d]/(d^2)$ , killing  $\sigma_1, \dots, \sigma_{p-1}$ , and sending  $\sigma_p$  to  $d$ , we see that  $x^{(p)}$  is the homomorphism

$$m \longmapsto 1 + d\xi(m)^p.$$

Thus, under the identification

$$\mathrm{Lie}(D_S(M)/S)(S) \simeq \mathrm{Hom}_{(\mathrm{Ab})}(M, \mathcal{O}(S))$$

of II, 5.1, the symbolic  $p$ -th power sends  $\xi$  to

$$\xi^{(p)} : m \longmapsto \xi(m)^p.$$

##### 6.4.2.

Let  $\mathcal{F}$  be an  $\mathcal{O}_S$ -module, and let  $G = \mathcal{W}(\mathcal{F})$  be the group  $S$ -functor in abelian groups (cf. I, 4.6). Let

$$y \in \mathcal{W}(\mathcal{F})(S) = \Gamma(S, \mathcal{F}),$$

and let  $y'$  be its image in  $\mathcal{W}(\mathcal{F})(I_S)$  under  $\mathcal{W}(\mathcal{F})(I_S \rightarrow S)$ .

By II, 4.4.2 and 4.5.1, the map

$$y \longmapsto dy'$$

is an isomorphism of  $\mathcal{O}(S)$ -modules from  $\mathcal{W}(\mathcal{F})(S)$  onto  $\mathrm{Lie}(\mathcal{W}(\mathcal{F})/S)(S)$ . Put  $x = dy'$ . Then the element  $x_i$  of 6.2 is  $d_i y''$ , where  $y''$  denotes the canonical image of  $y'^{60}$  in  $\mathcal{W}(\mathcal{F})(I_S^p)$ . Hence

$$x_1 + \cdots + x_p = (d_1 + \cdots + d_p)y'' = \sigma_1 y'',$$

and this belongs to  $\mathcal{W}(\mathcal{F})(\Sigma^p I_S)$ . Since the homomorphism

$$\mathcal{O}(\Sigma^p I_S) \longrightarrow \mathcal{O}(I_S)$$

defining the morphism  $i$  kills  $\sigma_1$ , we have  $x^{(p)} = 0$ . Therefore

*For every  $\mathcal{O}_S$ -module  $\mathcal{F}$ , the operation  $x \mapsto x^{(p)}$  on  $\mathrm{Lie} \mathcal{W}(\mathcal{F})$  is zero.*

### 6.4.3.

Let  $X$  be an  $S$ -scheme, let  $G = \mathrm{Aut}_S X$ , and let  $D$  be an  $S$ -derivation of the structure sheaf  $\mathcal{O}_X$ . By 1.5,  $D$  may be identified with an  $I_S$ -automorphism  $x$  of  $X_{I_S}$  inducing the identity on  $X$ . It can be described as follows. If  $f$  is a section of  $\mathcal{O}_S[d]/(d^2)$  of the form  $a + bd$ , put

$$D_{I_S} f = Da + (Db)d.$$

Thus  $D_{I_S}$  is obtained from  $D$  by the base change  $I_S \rightarrow S$ , and the automorphism of  $X_{I_S}$  is associated with the endomorphism

$$f \longmapsto f + (D_{I_S} f)d = a + (D(a) + b)d$$

of  $\mathcal{O}_S[d]/(d^2)$ .

Similarly, let  $D_{I_S^p}$  be the differential operator on  $X_{I_S^p}$  obtained from  $D$  by the base change  $I_S^p \rightarrow S$ . With the notation of 6.2, the automorphism  $x_i$  of  $X_{I_S^p}$  is associated with the endomorphism

$$f \longmapsto f + (D_{I_S^p} f)d_i$$

of

$$\mathcal{O}_S[d_1, \dots, d_p]/(d_1^2, \dots, d_p^2).$$

The product  $x_1 \cdots x_p$  is therefore associated with

$$(1 + d_1 D_{I_S^p})(1 + d_2 D_{I_S^p}) \cdots (1 + d_p D_{I_S^p}),$$

that is, with

$$1 + \sigma_1 D_{I_S^p} + \sigma_2 D_{I_S^p}^2 + \cdots + \sigma_p D_{I_S^p}^p.$$

The coefficient of  $\sigma_p$  is  $D_{I_S^p}^p$ . Consequently, the Lie-algebra isomorphism

$$\mathrm{Der}_S(\mathcal{O}_X) \xrightarrow{\sim} \mathrm{Lie}(\mathrm{Aut}_S X), \quad D \longmapsto x$$

(cf. 1.5) is also an isomorphism of restricted  $p$ -Lie algebras.

### 6.4.4.

The same method shows that, for every  $\mathcal{O}_S$ -module  $\mathcal{F}$ , the Lie-algebra isomorphism

$$\mathrm{Lie}(\mathrm{Aut}_{\mathcal{O}_S\text{-mod.}} \mathcal{W}(\mathcal{F})/S)(S) \xrightarrow{\sim} (\mathrm{End}_{\mathcal{O}_S\text{-mod.}} \mathcal{W}(\mathcal{F}))(S)$$

(cf. II, 4.8) is also an isomorphism of restricted  $p$ -Lie algebras.

---

<sup>60</sup>The editors corrected  $x$  to  $y'$ .

**6.4.5.**<sup>61</sup>

Let  $\mathcal{U}$  be an  $\mathcal{O}_S$ -coalgebra in groups, and let

$$G = \mathrm{Spec}^* \mathcal{U}$$

be the group  $S$ -functor; suppose that  $G$  is representable. For every  $T \rightarrow S$ , 5.5 and 6.1.1 define two restricted  $p$ -Lie algebra structures on

$$L(T) = \mathrm{Lie}(G)(T).$$

There is a commutative diagram

$$\begin{array}{ccc} L(T) & \xleftarrow{\tau} & \Gamma(T, \mathcal{U}_T) \\ \alpha \downarrow & \searrow i & \uparrow \psi \\ U(G_T) & \xleftarrow{\varphi} & U(L(T)) \end{array}$$

in which  $U(L(T))$  is the enveloping algebra of  $L(T)$ , and  $\varphi, \psi$  are the algebra homomorphisms induced by  $\alpha, \tau$ . It follows that the two restricted  $p$ -Lie algebra structures coincide: if  $x \in L(T)$  is identified with its image in  $U(G_T)$ , respectively in  $\Gamma(T, \mathcal{U}_T)$ , then  $x^{(p)}$  is the image under  $\varphi$ , respectively under  $\psi$ , of the element  $x^p \in U(L(T))$ .

## 7. Radicial Groups of Height 1

Let  $S$  be a scheme of characteristic  $p > 0$ .<sup>62</sup> We shall say that an  $\mathcal{O}_S$ -algebra  $\mathcal{A}$  (respectively, an  $\mathcal{O}_S$ - $p$ -Lie algebra  $\mathcal{L}$ ) is *finite locally free* if its underlying  $\mathcal{O}_S$ -module is locally free and of finite type. If  $\mathcal{L}$  is a finite locally free  $\mathcal{O}_S$ - $p$ -Lie algebra, we know (cf. 5.5.2) that the group  $S$ -functor

$$\mathrm{Spec}^* \mathcal{U}_p(\mathcal{L})$$

is represented by a finite locally free  $S$ -group scheme  $\mathfrak{G}_p(\mathcal{L})$ . We shall see that this  $S$ -group scheme solves a universal problem (7.2), and we shall characterize the  $S$ -group schemes of the form  $\mathfrak{G}_p(\mathcal{L})$  (7.4).

**Definition 7.0.**<sup>63</sup> *Let  $H = \mathrm{Spec} \mathcal{A}$  be a finite locally free  $S$ -group scheme. We say that  $H$  is infinitesimal if its unit section  $\varepsilon_H : S \rightarrow H$  is a homeomorphism; equivalently, the augmentation ideal of  $\mathcal{A}$  is locally nilpotent.*

**7.1.**<sup>64</sup>

Let  $\mathcal{L}$  be a finite locally free  $\mathcal{O}_S$ - $p$ -Lie algebra, and let  $\mathfrak{G}_p(\mathcal{L})$  be the affine  $S$ -group  $\mathrm{Spec} \mathcal{U}_p(\mathcal{L})$ . By 5.5,  $\mathcal{L}$  is identified with  $\mathrm{Lie} \mathfrak{G}_p(\mathcal{L})$ .

Now let  $G$  be a very good group  $S$ -functor satisfying condition (F) of 6.3, and let

$$\varphi : \mathfrak{G}_p(\mathcal{L}) \longrightarrow G$$

be a morphism of group  $S$ -functors. By 6.3, the morphism of  $\mathcal{O}_S$ -Lie algebras

$$\mathrm{Lie} \varphi : \mathrm{Lie} \mathfrak{G}_p(\mathcal{L}) \longrightarrow \mathrm{Lie} G$$

<sup>61</sup>The editors supplied the number 6.4.5 and expanded the original text.

<sup>62</sup>For the results of this section, one may also consult [DG70], § II.7, nos. 3–4.

<sup>63</sup>The editors added this definition (cf. [DG70], § II.4, 7.1), which will be used in 7.2.1.

<sup>64</sup>The editors simplified the original, taking into account the additions made in 5.5.

is compatible with the symbolic  $p$ -th power. If  $\mathrm{Hom}_p(\mathcal{L}, \mathrm{Lie} G)$  denotes the set of morphisms of  $\mathcal{O}_S$ -Lie algebras compatible with the symbolic  $p$ -th power, we therefore have a map

$$\mathrm{Lie} : \mathrm{Hom}_{S\text{-Gr.}}(\mathfrak{G}_p(\mathcal{L}), G) \longrightarrow \mathrm{Hom}_p(\mathcal{L}, \mathrm{Lie} G), \quad \varphi \longmapsto \mathrm{Lie} \varphi.$$

### 7.2. Theorem.

*If  $\mathcal{L}$  is a finite locally free  $\mathcal{O}_S$ - $p$ -Lie algebra, the map*

$$\mathrm{Hom}_{S\text{-gr.}}(\mathfrak{G}_p(\mathcal{L}), G) \longrightarrow \mathrm{Hom}_p(\mathcal{L}, \mathrm{Lie} G)$$

*is bijective in each of the following cases:*

- (i)  $G$  is an  $S$ -group scheme;
- (ii)  $G$  is of the form  $\mathrm{Aut}_S X$ , where  $X$  is an  $S$ -scheme;
- (iii)  $G$  is of one of the forms  $\mathcal{W}(\mathcal{F})$  or  $\mathrm{Aut}_{\mathcal{O}_S\text{-mod.}} \mathcal{W}(\mathcal{F})$ , where  $\mathcal{F}$  is a quasi-coherent  $\mathcal{O}_S$ -module.

The proof of the theorem rests on the following lemma.

#### Lemma.

*If  $\mathcal{L}$  is a finite locally free  $\mathcal{O}_S$ - $p$ -Lie algebra, the  $S$ -group*

$$G = \mathfrak{G}_p(\mathcal{L})$$

*is annihilated by the Frobenius morphism  $\mathrm{Fr} : G \rightarrow G^{(p)}$ . In particular,  $G$  is infinitesimal.*

Indeed, let  $\mathcal{U}$  be the restricted enveloping algebra of  $\mathcal{L}$ , let  $\mathcal{A} = \mathcal{U}^*$  be the affine algebra of  $G$ , and let  $\mathcal{I} = \ker \varepsilon_{\mathcal{A}}$  be the augmentation ideal of  $\mathcal{A}$ .<sup>65</sup> We have

$$\mathcal{A} = \mathcal{I} \oplus \eta_{\mathcal{A}}(\mathcal{O}_S), \tag{1}$$

where  $\eta_{\mathcal{A}}$  denotes the unit section of  $\mathcal{A}$ . Since  $\varepsilon_{\mathcal{A}}$  (respectively,  $\eta_{\mathcal{A}}$ ) is the transpose of  $\eta_{\mathcal{U}}$  (respectively,  $\varepsilon_{\mathcal{U}}$ ), this decomposition corresponds by duality to

$$\mathcal{U} = \mathcal{J} \oplus \eta_{\mathcal{U}}(\mathcal{O}_S), \tag{2}$$

where  $\mathcal{J}$  is the augmentation ideal of  $\mathcal{U}$ ; thus  $\mathcal{J} = \mathcal{I}^*$ .

Let  $\pi$  denote the endomorphism  $x \mapsto x^p$  of  $\mathcal{O}_S$ . We must show that  $\mathrm{Fr} : G \rightarrow G^{(p)}$  factors through the unit section of  $G^{(p)}$ . By 4.1.4 (c), this is equivalent to saying that the morphism

$$\Phi : \mathcal{I} \otimes_{\pi} \mathcal{O}_S \longrightarrow \mathcal{A}, \quad a \otimes_{\pi} x \longmapsto a^p x,$$

is zero. Since  $\mathcal{A}$  is finite locally free over  $\mathcal{O}_S$ , it suffices to show that the transpose  ${}^t\Phi$  is zero.

Now  $\Phi$  is precisely the following composite:

$$\mathcal{J} \otimes_{\pi} \mathcal{O}_S \xrightarrow{\tau} \mathcal{A} \otimes_{\pi} \mathcal{O}_S \xrightarrow{j(\mathcal{A})} S^p \mathcal{A} \xrightarrow{b(\mathcal{A})} \mathcal{A},$$

Here  $\tau$  is induced by the inclusion  $\mathcal{I} \hookrightarrow \mathcal{A}$ , while  $b(\mathcal{A})$  and  $j(\mathcal{A})$  are defined as in 4.3.3:  $b(\mathcal{A})$  is induced by multiplication in  $\mathcal{A}$ , and  $j(\mathcal{A})$  sends  $a \otimes_{\pi} 1$  to the image of  $a \otimes \cdots \otimes a$  in  $S^p \mathcal{A}$ . The  $\mathcal{O}_S$ -dual of  $S^p \mathcal{A}$  is the submodule  $\Sigma^p \mathcal{U}$  of  $\bigotimes^p \mathcal{U}$  consisting of the sections invariant under the symmetric group of degree  $p$ . Hence  ${}^t\Phi$  is the following composite:

<sup>65</sup>The editors supplied details in what follows. For another proof, see [DG70], § II.7, 3.9.

$$\mathcal{U} \xrightarrow{a(\mathcal{U})} \Sigma^p \mathcal{U} \xrightarrow{r(\mathcal{U})} \mathcal{U} \otimes_{\pi} \mathcal{O}_S \xrightarrow{q} \mathcal{J} \otimes_{\pi} \mathcal{O}_S,$$

Here  $q$  is induced by the projection  $\mathcal{U} \rightarrow \mathcal{J}$  with kernel  $\eta_{\mathcal{U}}(\mathcal{O}_S)$ ,  $a(\mathcal{U})$  is induced by the comultiplication of  $\mathcal{U}$ , and  $r(\mathcal{U})$  vanishes on symmetrized tensors and sends a section  $x \otimes \cdots \otimes x$  to  $x \otimes_{\pi} 1$  (cf. 4.3.3).

It is clear that  ${}^t\Phi \circ \eta_{\mathcal{U}} = 0$ ; by (2), it remains to show that  ${}^t\Phi$  annihilates the augmentation ideal  $\mathcal{J}$ . Since  ${}^t\Phi$  is an algebra morphism and  $\mathcal{J}$  is generated by  $\mathcal{L}$ , identified with its image in  $\mathcal{U}$ , it suffices to show that  ${}^t\Phi(x) = 0$  for every section  $x$  of  $\mathcal{L}$ . But

$$-a(\mathcal{U})(x) = (p-1)! a(\mathcal{U})(x)$$

is the symmetrization of  $x \otimes 1 \otimes \cdots \otimes 1$ , so its image under  $r(\mathcal{U})$  is zero. This proves the first assertion of the lemma.

The second follows. Every local section of  $\mathcal{I}$  has zero  $p$ -th power, and  $\mathcal{I}$  is an  $\mathcal{O}_S$ -module of finite type; hence  $\mathcal{I}$  is locally nilpotent. Explicitly, if  $V$  is an affine open subset of  $S$  such that

$$I = \Gamma(V, \mathcal{I})$$

is generated by  $r$  elements, then

$$I^{r(p-1)+1} = 0.$$

Thus  $G_{\text{red}} = S_{\text{red}}$ , and the unit section  $\varepsilon_G : S \rightarrow G$  is a homeomorphism.

### 7.2.1.<sup>66</sup>

We first prove assertion (ii) of Theorem 7.2. Let  $\pi : X \rightarrow S$  be an  $S$ -scheme, and first consider an arbitrary infinitesimal  $S$ -group  $H$ .

Morphisms  $\varphi : H \rightarrow \text{Aut } X$  correspond bijectively to left actions  $\mu : H \times X \rightarrow X$  of  $H$  on  $X$ . For such an action, if  $\varepsilon$  is the unit section of  $H$ , the composite

$$X \simeq S \times X \xrightarrow{\varepsilon \times X} H \times X \xrightarrow{\mu} X$$

must be the identity. Since  $(H \times X)_{\text{red}}$  is identified with  $X_{\text{red}}$ , the morphism  $\mu$  must induce the identity on the associated reduced schemes. In particular,  $\mu$  induces an action of  $H$  on every open subset of  $X$ . Thus, for every open  $U \subset X$  affine over  $S$ , we obtain a morphism of unital associative algebras

$$\mathcal{A}(U) \longrightarrow \mathcal{A}(H) \otimes \mathcal{A}(U)$$

which makes  $\mathcal{A}(U)$  a left  $\mathcal{A}(H)$ -comodule, in such a way that the restriction maps

$$\mathcal{A}(U) \longrightarrow \mathcal{A}(U'), \quad U' \subset U,$$

are comodule morphisms. Conversely, every datum of this kind comes from a unique left action  $\mu : H \times X \rightarrow X$ .

We also have the following lemma.

#### **Lemma.**

*Let  $X = \text{Spec } \mathcal{C}$  be an affine  $S$ -scheme, let  $H = \text{Spec } \mathcal{A}$  be an infinitesimal  $S$ -group, and put*

$$\mathcal{U} = \mathcal{A}^* = \text{Hom}_{\mathcal{O}_S}(\mathcal{A}, \mathcal{O}_S).$$

<sup>66</sup>The editors supplied details in what follows.

Left actions of  $H$  on  $X$  correspond bijectively to representations of the algebra  $\mathcal{U}$  in the  $\mathcal{O}_S$ -module  $\mathcal{C}$  such that

$$\begin{aligned} (a) \quad & u(1_{\mathcal{C}}) = \varepsilon(u) 1_{\mathcal{C}}, \\ (b) \quad & u(xy) = \sum_i v_i(x) w_i(y) \quad \text{if} \quad \Delta u = \sum_i v_i \otimes w_i. \end{aligned}$$

Here  $u$  is an arbitrary section of  $\mathcal{U}$  over an affine open subset  $V$  of  $S$ , while  $x$  and  $y$  are sections of  $\mathcal{C}$  over  $V$ . The symbol  $1_{\mathcal{C}}$  denotes the unit section of  $\mathcal{C}$ , and  $\varepsilon$  and  $\Delta$  denote the augmentation and the diagonal morphism of  $\mathcal{U}$ , respectively.

Indeed, a left action  $\mu$  of  $H$  on  $X$  is defined by a morphism of unital associative algebras

$$\lambda : \mathcal{C} \longrightarrow \mathcal{A} \otimes \mathcal{C}$$

which makes  $\mathcal{C}$  a left  $\mathcal{A}$ -comodule. Let  $\alpha$  be the composite

$$\mathcal{U} \otimes_{\mathcal{O}_S} \mathcal{C} \xrightarrow{\mathcal{U} \otimes \lambda} \mathcal{U} \otimes_{\mathcal{O}_S} \mathcal{A} \otimes_{\mathcal{O}_S} \mathcal{C} \xrightarrow{\gamma \otimes \mathcal{C}} \mathcal{O}_S \otimes_{\mathcal{O}_S} \mathcal{C} \simeq \mathcal{C}$$

where  $\gamma$  is the “contraction” from  $\mathcal{A}^* \otimes_{\mathcal{O}_S} \mathcal{A}$  to  $\mathcal{O}_S$ . Since  $\mathcal{A}$  is finite locally free over  $\mathcal{O}_S$ , the map

$$\lambda \longmapsto (\gamma \otimes \mathcal{C})(\mathcal{U} \otimes \lambda)$$

is a bijection from

$$\mathrm{Hom}_{\mathcal{O}_S}(\mathcal{C}, \mathcal{A} \otimes \mathcal{C}) \quad \text{onto} \quad \mathrm{Hom}_{\mathcal{O}_S}(\mathcal{U} \otimes \mathcal{C}, \mathcal{C}).$$

Moreover, by duality, the condition that  $\lambda$  define a left  $\mathcal{A}$ -comodule structure (respectively, be a morphism of unital associative algebras) is equivalent to the condition that  $\alpha$  be a representation of  $\mathcal{U}$  in  $\mathcal{C}$  (respectively, satisfy conditions (a) and (b)). This proves the lemma.

For any representation of  $\mathcal{U}$  in the  $\mathcal{O}_S$ -module  $\mathcal{C}$ , it is also clear that the sections  $u$  of  $\mathcal{U}$  satisfying conditions (a) and (b) form a subalgebra of  $\mathcal{U}$ .

In the particular case

$$H = \mathfrak{G}_p(\mathcal{L})$$

which concerns us, these conditions will therefore hold for every section  $u$  of  $\mathcal{U} = \mathcal{U}_p(\mathcal{L})$  if they hold for the sections  $u$  of  $\mathcal{L}$ , identified with its image in  $\mathcal{U}_p(\mathcal{L})$ .

If  $u$  is a section of  $\mathcal{L}$ , conditions (a) and (b) simply mean

$$u(1_{\mathcal{C}}) = 0, \quad u(xy) = u(x)y + xu(y);$$

that is,  $\alpha(u)$  is an  $\mathcal{O}_S$ -derivation of  $\mathcal{C} = \mathcal{A}(X)$ . Assertion (ii) of 7.2 follows. Indeed, every morphism

$$\varphi : \mathfrak{G}_p = \mathfrak{G}_p(\mathcal{L}) \longrightarrow \mathrm{Aut} X$$

defines a morphism of  $p$ -Lie algebras  $\mathrm{Lie} \varphi$  from  $\mathcal{L} = \mathrm{Lie} \mathfrak{G}_p$  to

$$\pi_*(\mathrm{Der}_{\mathcal{O}_S} \mathcal{O}_X),$$

and conversely every datum of this kind comes, by what precedes, from a unique action  $\mu : G \times X \rightarrow X$ .

### 7.2.2.



We now show how assertion (i) of Theorem 7.2 follows from (ii). Let  $G$  be an  $S$ -group scheme. If  $T$  is an  $S$ -scheme and  $x \in G(T)$ , let  $\ell_x^T$  (respectively,  $r_x^T$ ) denote the left (respectively, right) translation of  $G_T$  defined by  $x$ . The maps

$$\ell^T : x \mapsto \ell_x^T$$

therefore determine a homomorphism  $\ell : G \rightarrow \text{Aut } G$ .

On the other hand, let  $f$  be a  $T$ -automorphism of  $G_T$ , and define

$${}^x f = (r_x^T)^{-1} f r_x^T;$$

thus, for every  $T' \rightarrow T$  and  $g \in G(T')$ ,

$$({}^x f)(g) = f(gx)x^{-1}.$$

In this way  $G$  acts on the left on the group  $S$ -functor  $\text{Aut } G$ , and hence also on the functors

$$T \mapsto \text{Hom}_{T\text{-Gr.}}(\mathfrak{G}_p(\mathcal{L}_T), \text{Aut } G_T)$$

and

$$T \mapsto \text{Hom}_p(\mathcal{L}_T, \text{Lie}(\text{Aut } G_T/T)).$$

The morphism  $\ell^T : G_T \rightarrow \text{Aut } G_T$  identifies  $G_T$  with the group of automorphisms of the  $T$ -scheme  $G_T$  which commute with right translations. Its derived morphism  $\text{Lie}(\ell^T)$  identifies  $\text{Lie } G_T$  with the  $p$ -Lie algebra of  $\mathcal{O}_T$ -derivations of  $\mathcal{O}_{G_T}$  which commute with right translations (cf. II, 4.11.1). They therefore induce the commutative squares

$$\begin{array}{ccc} \text{Hom}_{T\text{-Gr.}}(\mathfrak{G}_p(\mathcal{L}_T), G_T) & \xrightarrow{\text{Lie}} & \text{Hom}_p(\mathcal{L}_T, \text{Lie}(G_T/T)) \\ \ell_T \downarrow & & \downarrow \text{Lie } \ell_T \\ \text{Hom}_{T\text{-Gr.}}(\mathfrak{G}_p(\mathcal{L}_T), \underline{\text{Aut}} G_T) & \xrightarrow{\text{Lie}} & \text{Hom}_p(\mathcal{L}_T, \text{Lie}(\underline{\text{Aut}} G_T/T)). \end{array}$$

The images of the two vertical arrows are the subfunctors formed by the invariants under the action of the  $S$ -group  $G$ . Since the lower horizontal arrow is invertible by 7.2.1 and is compatible with the action of  $G$ , the upper horizontal arrow is invertible as well. This proves 7.2 (i).

### 7.2.3.

Consider the case in which  $G = \text{Aut}_{\mathcal{O}_S\text{-mod.}} \mathcal{W}(\mathcal{F})$ .<sup>67</sup> Put

$$\mathcal{U} = \mathcal{U}_p(\mathcal{L}), \quad \mathcal{A} = \mathcal{U}^*, \quad H = \mathfrak{G}_p(\mathcal{L}) = \text{Spec } \mathcal{A}.$$

Since  $H$  is affine over  $S$ , VIB, 11.6.1 shows that a morphism of  $S$ -groups from  $H$  to  $\text{Aut}_{\mathcal{O}_S\text{-mod.}} \mathcal{W}(\mathcal{F})$  is the same as a right  $\mathcal{A}$ -comodule structure

$$\mu : \mathcal{F} \longrightarrow \mathcal{F} \otimes \mathcal{A}.$$

Since  $\mathcal{A}$  is finite locally free over  $\mathcal{O}_S$ , this is in turn equivalent to giving a representation

$$\alpha : \mathcal{U} \longrightarrow \text{End}_{\mathcal{O}_S}(\mathcal{F})$$

<sup>67</sup>In what follows, the editors expanded and simplified the original, taking VIB, 11.6.1 into account.

of  $\mathcal{U}$  in  $\mathcal{F}$ . Finally, by the universal property of  $\mathcal{U} = \mathcal{U}_p(\mathcal{L})$ , giving such a morphism  $\alpha$  is equivalent to giving its restriction  $\rho$  to  $\mathcal{L}$ , identified with its image in  $\mathcal{U}$ , and  $\rho$  is a morphism of  $p$ -Lie algebras from  $\mathcal{L}$  to  $\text{End}_{\mathcal{O}_S}(\mathcal{F})$ .

Finally, consider the case  $G = \mathcal{W}(\mathcal{F})$ , with the preceding notation.<sup>68</sup> To give a morphism of  $S$ -functors  $\varphi : H \rightarrow \mathcal{W}(\mathcal{F})$  is first of all equivalent to giving an element  $\theta$  of  $\Gamma(H, \mathcal{F} \otimes \mathcal{O}_H)$ . Since  $H$  is finite locally free over  $S$ , we have

$$\Gamma(H, \mathcal{F} \otimes \mathcal{O}_H) = \Gamma(S, \mathcal{F} \otimes \mathcal{A}) = \text{Hom}_{\mathcal{O}_S}(\mathcal{U}, \mathcal{F}).$$

The condition that  $\varphi$  be a group morphism means that  $\theta$ , regarded as a morphism of  $\mathcal{O}_S$ -modules  $\mathcal{U} \rightarrow \mathcal{F}$ , vanishes on  $1_{\mathcal{U}}$  and on  $\mathcal{J}^2$ , where  $\mathcal{J}$  is the augmentation ideal of  $\mathcal{U}$ . Consequently,

$$\text{Hom}_{S\text{-gr.}}(H, \mathcal{W}(\mathcal{F})) = \text{Hom}_{\mathcal{O}_S}(\mathcal{J}/\mathcal{J}^2, \mathcal{F}). \quad (1)$$

Next consider the quasi-coherent sheaf  $[\mathcal{L}, \mathcal{L}]$ , the image of the morphism

$$\mathcal{L} \otimes \mathcal{L} \longrightarrow \mathcal{L}, \quad x \otimes y \longmapsto [x, y].$$

For every affine open subset  $V$  of  $S$ , we have

$$[\mathcal{L}, \mathcal{L}](V) = [\mathcal{L}(V), \mathcal{L}(V)].$$

There is an exact sequence

$$(\dagger) \quad 0 \longrightarrow [\mathcal{L}, \mathcal{L}] \longrightarrow \mathcal{L} \xrightarrow{\pi} \mathcal{J}/\mathcal{J}^2 \longrightarrow 0,$$

in which  $\pi$  is the composite of the inclusion  $\mathcal{L} \hookrightarrow \mathcal{J}$  and the projection  $\mathcal{J} \rightarrow \mathcal{J}/\mathcal{J}^2$ .

Indeed, the question is local on  $S$ , so we may suppose that  $S$  is affine with ring  $R$ , and that

$$L = \mathcal{L}(S)$$

is free with basis  $(x_1, \dots, x_r)$ . Identify  $L$  with its image in  $U = U_p(L)$ , and let  $K$  be the sub- $R$ -module of  $U$  which is the direct sum of  $[L, L]$  and the submodule with basis the monomials

$$x_1^{n_1} \cdots x_r^{n_r} \quad \text{such that} \quad n_1 + \cdots + n_r \geq 2.$$

One checks that  $K$  is a two-sided ideal of  $U$ . Since  $K$  is contained in  $J^2$ , where  $J$  is the augmentation ideal of  $U$ , and contains all the products  $x_i x_j$ , which generate  $J^2$ , it follows that  $K = J^2$ . Hence

$$J^2 \cap L = [L, L],$$

and the displayed sequence is exact.

By 6.4.2,  $\text{Lie } \mathcal{W}(\mathcal{F})$  is simply  $\mathcal{W}(\mathcal{F})$ , with zero Lie bracket and zero symbolic  $p$ -th power. From this and the preceding discussion, we obtain

$$\text{Hom}_p(\mathcal{L}, \mathcal{F}) = \text{Hom}_{\mathcal{O}_S}(\mathcal{L}/[\mathcal{L}, \mathcal{L}], \mathcal{F}) = \text{Hom}_{\mathcal{O}_S}(\mathcal{J}/\mathcal{J}^2, \mathcal{F}). \quad (2)$$

Together with (1), this completes the proof of Theorem 7.2.

### 7.3. Lemma.

*If  $\mathcal{L}$  is a finite locally free  $\mathcal{O}_S$ - $p$ -Lie algebra, the morphism*

$$j_{\mathcal{L}} : \mathcal{L} \longrightarrow \text{Lie } \mathfrak{G}_p(\mathcal{L})$$

<sup>68</sup>The editors added what follows. The original said that “the case of  $\mathcal{W}(\mathcal{F})$  is analogous.”

of 5.5 is invertible.<sup>69</sup>

For the proof, see 5.5.1.

#### 7.4.

To conclude this section, we give a characterization of the  $S$ -group schemes of the form  $\mathfrak{G}_p(\mathcal{L})$ , where  $\mathcal{L}$  is a finite locally free  $\mathcal{O}_S$ - $p$ -Lie algebra.

Let  $G$  be an  $S$ -group scheme, let  $\varepsilon_G$  be its unit section, and let  $\mathcal{I}'$  be the kernel of the morphism

$$\varepsilon_G^{-1}(\mathcal{O}_G) \longrightarrow \mathcal{O}_S$$

corresponding to  $\varepsilon_G$ . By 2.5 and 1.3.1, the image of  $\mathrm{Lie}(G/S)(S)$  in  $U(G)$  is identified with the morphisms of  $\mathcal{O}_S$ -modules from  $\varepsilon_G^{-1}(\mathcal{O}_G)$  to  $\mathcal{O}_S$  which vanish on the unit section of  $\varepsilon_G^{-1}(\mathcal{O}_G)$  and on  $\mathcal{I}'^2$ .

We thereby recover the canonical isomorphism

$$\mathrm{Lie}(G/S)(S) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{O}_S}(\mathcal{I}'/\mathcal{I}'^2, \mathcal{O}_S)$$

of II, 3.3 and 4.11.4.<sup>70</sup> As in the cited passage, put

$$\omega_{G/S} = \mathcal{I}'/\mathcal{I}'^2.$$

Thus the sheaf  $\mathrm{Lie}(G/S)$  is identified with

$$\omega_{G/S}^* = \mathrm{Hom}_{\mathcal{O}_S}(\omega_{G/S}, \mathcal{O}_S).$$

Moreover,<sup>71</sup> if  $G = \mathfrak{G}_p(\mathcal{L})$ , where  $\mathcal{L}$  is a finite locally free  $\mathcal{O}_S$ - $p$ -Lie algebra, then 5.5.3 gives

$$\omega_{G/S} = \mathcal{L}^*.$$

#### Theorem.

If  $G$  is a group scheme over a scheme  $S$  of characteristic  $p > 0$ , the following assertions are equivalent:

- (i) There exists a finite locally free  $\mathcal{O}_S$ - $p$ -Lie algebra  $\mathcal{L}$  such that  $G \simeq \mathfrak{G}_p(\mathcal{L})$ .
- (i') The  $\mathcal{O}_S$ - $p$ -Lie algebra  $\mathrm{Lie}(G)$  is finite locally free, and  $G \simeq \mathfrak{G}_p(\mathrm{Lie}(G))$ .
- (ii)  $G$  is affine over  $S$ , the  $\mathcal{O}_S$ -module  $\omega_{G/S}$  is locally free of finite type, and the affine algebra of  $G$  is locally isomorphic to the quotient of the symmetric algebra  $S_{\mathcal{O}_S}(\omega_{G/S})$  by the ideal generated by the  $p$ -th powers of the sections of  $\omega_{G/S}$ .
- (iii)  $G$  is locally of finite presentation over  $S$ , is of height  $\leq 1$ , and  $\omega_{G/S}$  is locally free.
- (iii')  $G$  is locally of finite type over  $S$ , is of height  $\leq 1$ , and  $\omega_{G/S}$  is locally free.
- (iv)  $G$  is locally of finite presentation and flat over  $S$ , and is of height  $\leq 1$ .<sup>72</sup>

<sup>69</sup>To avoid changing the numbering, the editors retained statement 7.3, although they included it, together with its proof, in 5.5.1.

<sup>70</sup>If  $G$  is affine over  $S$  and  $\mathcal{I}$  denotes the augmentation ideal of  $\mathcal{A}(G)$ , then  $\mathcal{I}'/\mathcal{I}'^2$  is identified with  $\varepsilon_G^*(\mathcal{I}/\mathcal{I}^2)$ ; cf. loc. cit.

<sup>71</sup>The editors added the following sentence.

<sup>72</sup>The editors added assertion (i'), implicit in the original, and assertions (iii') and (iv), pointed out by O. Gabber. Assertion (iv) takes up a footnote in the original, which stated: "The condition on  $\omega_{G/S}$  is in fact unnecessary, as is easily seen by reducing to the case where  $S$  is local with residue field  $k$ , and applying the theorem to  $G_k$ ." As Gabber pointed out, this is incorrect without a flatness hypothesis. Indeed, let  $A$

**7.4.1.**

The equivalence (i)  $\Leftrightarrow$  (i') follows from 5.5.3 (i), and the implications (ii)  $\Rightarrow$  (iii)  $\Rightarrow$  (iii') are clear. We also have (i)  $\Rightarrow$  (iv), since  $\mathfrak{G}_p(\mathcal{L})$  is finite locally free and of height  $\leq 1$ , by 5.5.2 and the lemma in 7.2. We show that (i) implies (ii).

Let  $\mathcal{I}$  be the augmentation ideal of

$$\mathcal{A} = \mathcal{U}_p(\mathcal{L})^*.$$

We saw in 5.5.3 (ii) that  $\omega_{G/S} = \mathcal{I}/\mathcal{I}^2$  is identified with  $\mathcal{L}^*$ , and hence is finite locally free.

Now suppose that  $S$  is affine. There is a section

$$\sigma : \omega_{G/S} \longrightarrow \mathcal{I}$$

of the projection  $\mathcal{I} \rightarrow \mathcal{I}/\mathcal{I}^2$ . It induces an algebra morphism

$$\sigma' : S_{\mathcal{O}_S}(\omega_{G/S}) \longrightarrow \mathcal{A},$$

and, by the lemma in 7.2,  $\sigma'$  factors through a morphism

$$\varphi : S_{\mathcal{O}_S}(\omega_{G/S})/\mathcal{K} \longrightarrow \mathcal{A},$$

where  $\mathcal{K}$  denotes the ideal generated by the  $p$ -th powers of sections of  $\omega_{G/S}$ . Filter  $\mathcal{A}$  by powers of  $\mathcal{I}$ , and filter  $S_{\mathcal{O}_S}(\omega_{G/S})/\mathcal{K}$  by powers of the ideal generated by  $\omega_{G/S}$ . The morphism  $\varphi$  plainly induces an epimorphism on the associated graded modules. Hence  $\varphi$  is an epimorphism between locally free  $\mathcal{O}_S$ -modules of the same rank (cf. 5.3.3), and is therefore an isomorphism. This proves that (i) implies (ii).

**7.4.2.**

Suppose now that  $G$  is of height  $\leq 1$  and locally of finite presentation over  $S$ .<sup>73</sup> The Frobenius morphism  $\text{Fr} : G \rightarrow G^{(p)}$  is integral and factors through the unit section of  $G^{(p)}$ ; consequently,  $G$  is integral, and hence affine, over  $S$ . Put  $\mathcal{A} = \mathcal{A}(G)$ . Since  $G$  is locally of finite presentation over  $S$ , it follows that  $G$  is finite and of finite presentation over  $S$ , and hence that  $\mathcal{A}(G)$  is an  $\mathcal{O}_S$ -module of finite presentation (cf. EGA IV<sub>1</sub>, 1.4.7).

Let  $\mathcal{I}$  be the augmentation ideal of  $\mathcal{A}$ . Since

$$\mathcal{A} = \eta_{\mathcal{A}}(\mathcal{O}_S) \oplus \mathcal{I},$$

where  $\eta_{\mathcal{A}}$  is the unit section of  $\mathcal{A}$ , the module  $\mathcal{I}$  is of finite presentation over  $\mathcal{O}_S$ , as is  $\omega_G = \mathcal{I}/\mathcal{I}^2$ . If instead  $G$  is assumed to be of height  $\leq 1$  and locally of finite type over  $S$ , the same argument shows that  $\mathcal{A}(G)$ ,  $\mathcal{I}$ , and  $\omega_G = \mathcal{I}/\mathcal{I}^2$  are  $\mathcal{O}_S$ -modules of finite type.

Thus, under hypothesis (iii'),  $\omega_{G/S}$  is finite locally free over  $\mathcal{O}_S$ , as is

$$\mathcal{L} = \text{Lie}(G/S) = \omega_{G/S}^*.$$

Put

$$\mathcal{B} = \mathcal{U}_p(\mathcal{L})^*, \quad H = \mathfrak{G}_p(\mathcal{L}) = \text{Spec } \mathcal{B}.$$

be an Artinian local ring of characteristic  $p > 0$ , let  $J$  be a proper nonzero ideal of  $A$ , and let  $H$  be the subgroup

$$\text{Spec } A[x]/(x^p, Jx)$$

of  $\alpha_{p,A}$ ; that is, for every  $A$ -algebra  $R$ ,

$$H(R) = \{x \in R \mid x^p = 0 \text{ and } Jx = 0\}.$$

Then  $H$  is not flat over  $A$ , and hence is not of the form  $\mathfrak{G}_p(\mathcal{L})$  with  $\mathcal{L}$  a finite-rank free  $p$ -Lie algebra over  $A$ .

<sup>73</sup>The editors supplied details and simplified the original in what follows.

By Theorem 7.2, the identity map of  $\mathcal{L}$  corresponds to a group morphism from  $H = \mathfrak{G}_p(\mathcal{L})$  to  $G$ , and hence to a morphism of  $\mathcal{O}_S$ -algebras

$$\theta : \mathcal{A} \longrightarrow \mathcal{B}.$$

By definition,  $\theta$  induces an isomorphism from  $\omega_{G/S}$  to  $\omega_{H/S}$ ; we must show that  $\theta$  itself is an isomorphism.

We may restrict to the case in which  $S$  is affine. There is then a section  $\tau$  of the projection  $\mathcal{I} \rightarrow \omega_{G/S}$ , inducing an algebra morphism

$$\tau' : S_{\mathcal{O}_S}(\omega_{G/S}) \longrightarrow \mathcal{A}.$$

Every local section of  $\mathcal{I}$  has zero  $p$ -th power, because  $\text{Fr} : G \rightarrow G^{(p)}$  factors through the unit section of  $G^{(p)}$ . Thus  $\tau'$  induces a morphism of  $\mathcal{O}_S$ -algebras  $\psi$  fitting into the commutative diagram

$$\begin{array}{ccc} S_{\mathcal{O}_S}(\omega_{G/S})/\mathcal{K} & \xrightarrow{\psi} & \mathcal{A} \\ & \searrow \varphi & \downarrow \theta \\ & & \mathcal{B} \end{array}$$

where  $\mathcal{K}$  is the ideal generated by the  $p$ -th powers of sections of  $\omega_{G/S}$ . As in 7.4.1,  $\psi$  is an epimorphism of  $\mathcal{O}_S$ -modules. On the other hand, 7.4.1 shows that  $\varphi = \theta \circ \psi$  is an isomorphism. Hence  $\theta$  is an isomorphism as well. This proves (iii')  $\Rightarrow$  (i).

#### 7.4.3.<sup>74</sup>

We finally prove that (iv) implies (iii). It suffices to show that  $\omega_{G/S}$  is locally free, so we may suppose that  $S$  is affine with ring  $R$ . As observed at the beginning of 7.4.2, hypothesis (iv) then implies that  $G = \text{Spec } A$ , where  $A$  is an  $R$ -algebra which is an  $R$ -module of finite presentation, as is

$$\omega_{G/A} = I/I^2,$$

where  $I$  is the augmentation ideal of  $A$ . Since  $G$  is also assumed flat over  $S$ , the algebra  $A$  is a finite locally free  $R$ -module (cf. [BAC] II, § 5.2, Theorem 1 and Corollary 2). By the cited result, to show that  $\omega_{G/A}$  is locally free of finite rank, it suffices to show that

$$(\omega_{G/A})_{\mathfrak{m}}$$

is flat for every maximal ideal  $\mathfrak{m}$  of  $R$ . Thus we may suppose that  $R$  is local and that  $A$  is free of rank  $n+1$ ; then  $I$  is free of rank  $n$ . Let  $\mathfrak{m}$  be the maximal ideal of  $R$ , and put

$$k = R/\mathfrak{m}.$$

Let  $I_k$  be the augmentation ideal of  $A_k$ , and let  $r$  be the dimension of

$$\omega_{G_k/k} = I_k/I_k^2.$$

Choose a basis  $(e_1, \dots, e_n)$  of  $I_k$  such that  $(e_{r+1}, \dots, e_n)$  is a basis of  $I_k^2$ , and choose elements  $x_1, \dots, x_n$  of  $I$  lifting the  $e_i$ . Nakayama's lemma shows that  $(x_1, \dots, x_n)$  is a basis of  $I$  over  $R$ .

<sup>74</sup>The editors added this paragraph to prove (iv)  $\Rightarrow$  (iii); cf. editor's note (71).

Let  $N$  be the sub- $R$ -module of  $I$  with basis  $(x_1, \dots, x_r)$ , and let  $B$  be the quotient of the symmetric algebra of  $N$  by the ideal generated by the elements  $x^p$ , for  $x \in N$ . Since every element of  $I$  has zero  $p$ -th power, we obtain a morphism of  $R$ -algebras

$$\psi : B \longrightarrow A.$$

By 7.4.2,  $\psi \otimes k$  is an isomorphism. It follows that  $\text{Coker } \psi = 0$ , and, if  $K = \ker \psi$ , the morphism

$$\tau : K \otimes k \longrightarrow B \otimes k$$

is zero. But  $\psi$  is surjective and  $A$  is flat over  $R$ , so  $\tau$  is also injective. Hence  $K \otimes k = 0$ .

Since  $A$  is an  $R$ -module of finite presentation,  $K$  is an  $R$ -module of finite type (cf. [BAC] I, § 2.8, Lemma 9). Nakayama's lemma now gives  $K = 0$ . Thus  $\psi$  is an isomorphism of  $R$ -algebras. Since  $\psi^{-1}(I)$  contains the augmentation ideal  $J$  of  $B$ , it follows that  $\psi^{-1}(I) = J$ . Therefore  $\psi^{-1}$  induces an isomorphism of  $R$ -modules

$$I/I^2 \xrightarrow{\sim} J/J^2 = N.$$

This proves that  $\omega_{G/S}$  is finite locally free, and hence proves (iv)  $\Rightarrow$  (iii). The proof of Theorem 7.4 is complete.

**Remark 7.5.**<sup>75</sup>

It follows immediately from Theorems 7.2 and 7.4 that the functors

$$G \longmapsto \text{Lie}(G), \quad \mathcal{L} \longmapsto \mathfrak{G}_p(\mathcal{L})$$

induce quasi-inverse equivalences between the category of  $S$ -groups which are locally of finite presentation, flat, and of height  $\leq 1$ , and the full subcategory of the category of  $\mathcal{O}_S$ - $p$ -Lie algebras consisting of the finite locally free  $\mathcal{O}_S$ - $p$ -Lie algebras.

## 8. The Case of a Base Field

### 8.1.

We now summarize the results obtained when  $S$  is the spectrum of a field  $k$  of characteristic  $p > 0$ . We call a  $k$ -group scheme *algebraic* if its underlying scheme is of finite type over  $k$ . Theorem 7.2 then gives the following result.<sup>76</sup>

**Theorem 8.1.1.** *The functor  $\mathfrak{G}_p$ , which associates with every finite-dimensional restricted  $p$ -Lie algebra  $\mathcal{L}$  over  $k$  the  $k$ -group  $\mathfrak{G}_p(\mathcal{L})$ , is left adjoint to the functor which associates  $\text{Lie}(G)$  with every algebraic  $k$ -group  $G$ .*

Combining this with Theorem 7.4 gives:

**Theorem 8.1.2.** *The functors*

$$\mathfrak{G}_p : \mathcal{L} \longmapsto \mathfrak{G}_p(\mathcal{L}), \quad G \longmapsto \text{Lie}(G),$$

*induce quasi-inverse equivalences between the category of finite-dimensional restricted  $p$ -Lie algebras over  $k$  and the category of algebraic  $k$ -groups of height at most 1.*

Since  $\mathfrak{G}_p$  is a left adjoint, it commutes with inductive limits,<sup>77</sup> and in particular with cokernels. Let

$$\varphi : G \longrightarrow H, \quad \varphi' : G' \longrightarrow H$$

<sup>75</sup>The editors added this remark.

<sup>76</sup>The editors supplied the numbering 8.1.1 through 8.1.3 in order to highlight the results stated here.

<sup>77</sup>The editors expanded the argument which follows, as well as the proof of Corollary 8.1.3.

be two morphisms between algebraic  $k$ -groups of height at most 1. The fiber product  $G \times_H G'$  is again an algebraic  $k$ -group of height at most 1, because the Frobenius morphism

$$\mathrm{Fr} : G \longrightarrow G^{(p)}$$

commutes with fiber products. Thus the inclusion of the category of algebraic  $k$ -groups of height at most 1 into the category of all algebraic  $k$ -groups commutes with fiber products, and hence in particular with kernels. We obtain:

**Corollary 8.1.3.** *The functor  $\mathfrak{G}_p$  is exact in the following sense. Let*

$$\pi : \mathcal{L}_1 \longrightarrow \mathcal{L}_2$$

*be a surjective homomorphism between finite-dimensional restricted  $p$ -Lie algebras over  $k$ , and let  $i$  be the inclusion of  $\mathcal{L}_0 = \ker \pi$  into  $\mathcal{L}_1$ . Then the following sequence of algebraic  $k$ -groups is exact:*

$$1 \longrightarrow \mathfrak{G}_p(\mathcal{L}_0) \xrightarrow{\mathfrak{G}_p(i)} \mathfrak{G}_p(\mathcal{L}_1) \xrightarrow{\mathfrak{G}_p(\pi)} \mathfrak{G}_p(\mathcal{L}_2) \longrightarrow 1.$$

78

Indeed, by the preceding discussion,  $\mathfrak{G}_p(i)$  induces an isomorphism from  $\mathfrak{G}_p(\mathcal{L}_0)$  onto  $\ker(\mathfrak{G}_p(\pi))$ . This kernel is the same whether it is taken in the category of all algebraic  $k$ -groups or in the subcategory of those of height at most 1. Furthermore,

$$\mathfrak{G}_p(\pi) : \mathfrak{G}_p(\mathcal{L}_1) \longrightarrow \mathfrak{G}_p(\mathcal{L}_2)$$

identifies  $\mathfrak{G}_p(\mathcal{L}_2)$  with the quotient of  $\mathfrak{G}_p(\mathcal{L}_1)$  by  $\mathfrak{G}_p(\mathcal{L}_0)$  in the category of algebraic  $k$ -groups.

**Remark 8.1.4.**<sup>79</sup>

Let  $\varphi : G \rightarrow H$  be a morphism of  $k$ -groups and put  $K = \ker(\varphi)$ . Suppose that  $\varphi$  is covering for the (fpqc) topology. This is in particular the case if it is covering for a coarser topology, such as the (fppf) topology. Then  $\varphi$  is a  $K$ -torsor over  $H$  (cf. IV, 5.1.7.1). Moreover, by IV, 6.3.1, there exists a covering of  $H$  by affine open subsets  $S_i$ , and for each  $i$  an affine faithfully flat morphism  $T_i \rightarrow S_i$  factoring through  $\varphi$ . Now  $G \times_H T_i$  is  $T_i$ -isomorphic to  $K \times T_i$ , hence faithfully flat over  $T_i$ . By (fpqc) descent,  $G \times_H S_i \rightarrow S_i$  is faithfully flat, and therefore  $\varphi$  is faithfully flat.

Conversely, if  $\varphi$  is faithfully flat and quasi-compact, respectively faithfully flat and locally of finite presentation, then it is covering for the (fpqc), respectively (fppf), topology (cf. IV, 6.3.1). Finally, a morphism of sheaves is covering if and only if it is an epimorphism (cf. IV, 4.4.3). In particular, a quasi-compact morphism of  $k$ -groups is faithfully flat if and only if it is an epimorphism of (fpqc) sheaves.

**8.2. Proposition.** *Consider an exact sequence<sup>80</sup> of algebraic groups over a field  $k$  of characteristic  $p > 0$ :*

$$1 \longrightarrow G' \xrightarrow{\tau} G \xrightarrow{\pi} G'' \longrightarrow 1.$$

*Consider the following assertions:*

(i)  $\pi$  is smooth;

<sup>78</sup>Moreover, by VIA, 3.2,  $\mathfrak{G}_p(\mathcal{L}_2)$  represents the (fppf) quotient sheaf of  $\mathfrak{G}_p(\mathcal{L}_1)$  by  $\mathfrak{G}_p(\mathcal{L}_0)$ .

<sup>79</sup>The editors added this remark, communicated by O. Gabber, for use in 8.3.1.

<sup>80</sup>Thus  $\pi$  is faithfully flat and  $\tau$  identifies  $G'$  with  $\ker \pi$ , so  $G''$  represents the (fppf) quotient sheaf of  $G$  by  $G'$ ; cf. VIA, 3.2 and 5.2.

(ii)  $G'$  is smooth;

(iii) for every integer  $n > 0$ , the sequence induced by  $\tau$  and  $\pi$ ,

$$1 \longrightarrow \mathrm{Fr}^n G' \longrightarrow \mathrm{Fr}^n G \longrightarrow \mathrm{Fr}^n G'' \longrightarrow 1.$$

is exact;

(iv) the morphism

$$\mathrm{Fr} \pi : \mathrm{Fr} G \longrightarrow \mathrm{Fr} G''$$

is an epimorphism of (fppf) sheaves;

(v) the morphism

$$\mathrm{Lie}(\pi) : \mathrm{Lie}(G) \longrightarrow \mathrm{Lie}(G'')$$

is surjective.

Then

$$(i) \Longleftrightarrow (ii) \implies (iii) \implies (iv) \Longleftrightarrow (v),$$

and all five assertions are equivalent when  $G$  is smooth over  $k$ .

Indeed, (i) is equivalent to (ii) by VIB, 9.2 (vii), and it is clear that (iii) implies (iv). The equivalence of (iv) and (v) follows from 8.1.3.

The implication (ii)  $\Rightarrow$  (iii) follows from the diagram

$$\begin{array}{ccccccc} 1 & \longrightarrow & G' & \xrightarrow{\tau} & G & \xrightarrow{\pi} & G'' \longrightarrow 1 \\ & & \downarrow \mathrm{Fr}^n(G'/k) & & \downarrow \mathrm{Fr}^n(G/k) & & \downarrow \mathrm{Fr}^n(G''/k) \\ 1 & \longrightarrow & G'^{(p^n)} & \xrightarrow{\tau(p^n)} & G^{(p^n)} & \xrightarrow{\pi(p^n)} & G''^{(p^n)} \longrightarrow 1 \end{array}$$

whose two rows are exact. Since  $\mathrm{Fr}^n(G'/k)$  is an epimorphism of (fppf) sheaves by Corollary 8.3.1 below,  $\pi$  induces an epimorphism from  $\mathrm{Fr}^n G$  onto  $\mathrm{Fr}^n G''$ ; here one uses the snake lemma for sheaves of groups, not necessarily commutative.

Likewise, suppose that  $G$  is smooth over  $k$ . Then  $\mathrm{Fr}(G/k)$  is an epimorphism. If  $\mathrm{Fr} \pi$  is also an epimorphism, the same snake lemma, applied to the preceding diagram with  $n = 1$ , shows that  $\mathrm{Fr}(G'/k)$  is an epimorphism. Hence  $G'$  is smooth over  $k$  by 8.3.1 below.

**8.3. Proposition.**<sup>81</sup> *Let  $G$  be a group locally of finite type over a field  $k$  of characteristic  $p > 0$ . There exists an integer  $n_0$  such that*

$$G/(\mathrm{Fr}^n G)$$

*is smooth over  $k$  for every  $n \geq n_0$ .*

The construction of  $G/(\mathrm{Fr}^n G)$  commutes with extension of the base field (4.1.1 and VIA, 3.3.2), so we may suppose that  $k$  is perfect. Then  $G_{\mathrm{red}}$  is a  $k$ -group locally of finite type (cf. VIA, 0.2), and, putting

$$H = G_{\mathrm{red}} \backslash G,$$

we have the following commutative exact diagram:

---

<sup>81</sup>The editors replaced “algebraic” by “locally of finite type.”



$$\begin{array}{ccccccc}
1 & \longrightarrow & G_{\text{red}} & \longrightarrow & G & \longrightarrow & H \\
& & \downarrow \text{Fr}^n(G_{\text{red}}/k) & & \downarrow \text{Fr}^n(G/k) & & \downarrow \text{Fr}^n(H/k) \\
1 & \longrightarrow & G_{\text{red}}^{(p^n)} & \longrightarrow & G^{(p^n)} & \longrightarrow & H^{(p^n)}.
\end{array}$$

The scheme  $H$  is the spectrum of a finite local  $k$ -algebra with residue field  $k$  (cf. VIA, 5.6.1). There is therefore an integer  $n_0$  such that, for every  $n \geq n_0$ ,  $\text{Fr}^n(H/k)$  factors through the unit section of  $H^{(p^n)}$ . It follows that, for  $n \geq n_0$ ,  $\text{Fr}^n(G/k)$  factors through  $G_{\text{red}}^{(p^n)}$ , and VIA, 5.4.1 gives the commutative diagram

$$\begin{array}{ccc}
G & \xrightarrow{\text{Fr}^n(G/k)} & G_{\text{red}}^{(p^n)} \\
\pi \downarrow & \nearrow i & \\
G/(\text{Fr}^n G) & & 
\end{array}$$

where  $i$  is a closed immersion and  $\pi$  is the canonical projection. Moreover,  $i$  induces a homeomorphism of the underlying topological spaces, so it is an isomorphism. Since  $k$  is perfect,  $G_{\text{red}}^{(p^n)}$  is smooth over  $k$  (VIA, 1.3.1). Thus  $G/(\text{Fr}^n G)$  is smooth over  $k$  for every  $n \geq n_0$ .

**8.3.1. Corollary.** *Let  $G$  be a group locally of finite type over a field  $k$  of characteristic  $p > 0$ , and let  $n \geq 1$ .<sup>82</sup> The following conditions are equivalent:*

- (i)  $G$  is smooth over  $k$ ;
- (ii)  $\text{Fr}^n(G/k) : G \rightarrow G^{(p^n)}$  is an epimorphism of (fppf) sheaves;
- (iii)  $\text{Fr}^n(G/k) : G \rightarrow G^{(p^n)}$  is faithfully flat.

Since  $G$  is locally of finite type over  $k$ ,  $\text{Fr}^n(G/k)$  is of finite presentation, so the equivalence of (ii) and (iii) follows from 8.1.4. Suppose that  $G$  is smooth over  $k$ , and hence reduced. Since  $\text{Fr}^n(G/k)$  is surjective, it is faithfully flat (cf. VIA, 6.2 or VIB, 1.3).

Conversely, suppose that  $\text{Fr}^n(G/k)$  is faithfully flat. The morphism

$$\text{Fr}^n(G^{(p^n)}/k)$$

is obtained from  $\text{Fr}^n(G/k)$  by base change (cf. 4.1.3), so it too is faithfully flat, as is the composite

$$\text{Fr}^{2n}(G/k) : G \longrightarrow G^{(p^n)} \longrightarrow G^{(p^{2n})}.$$

It follows that, for every  $m \in \mathbb{N}$ ,

$$\text{Fr}^{mn}(G/k) : G \longrightarrow G^{(p^{mn})}$$

is faithfully flat. Hence it induces an isomorphism

$$G/(\text{Fr}^{mn} G) \simeq G^{(p^{mn})}$$

<sup>82</sup>The editors made explicit the equivalence of (ii) and (iii) and expanded the proof.

(cf. VIA, 5.4.1). By Proposition 8.3,  $G^{(p^{mn})}$  is smooth over  $k$  for  $m$  sufficiently large, and therefore  $G$  is smooth as well by (fpqc) descent (cf. EGA IV<sub>4</sub>, 17.7.1).

#### 8.4.

In the two statements which conclude this Exposé, we return to the case of a field  $k$  of arbitrary characteristic.

If  $k$  has characteristic 0, let  $n \geq 1$ . If  $k$  has characteristic  $p > 0$ , let  $n \geq 1$  be prime to  $p$ . In both cases we simply say that  $n$  is *prime to the characteristic of  $k$* . If  $G$  is a group scheme over  $k$ , we denote by

$$n_G : G \longrightarrow G$$

the morphism of  $k$ -schemes which sends  $x \in G(T)$  to  $x^n \in G(T)$ , for every  $k$ -scheme  $T$ .

**Proposition.** *Let  $G$  be an algebraic group over a field  $k$ , and let  $n$  be prime to the characteristic of  $k$ . Then*

$$n_G : G \longrightarrow G$$

*is an étale morphism.*

By VIB, 1.3, it suffices to show that  $n_G$  is étale at the origin.<sup>83</sup> Let  $A$  be the local ring of  $G$  at the origin and  $I$  its maximal ideal. By II, 3.9.4, the map

$$\mathrm{Lie}(n_G) : \mathrm{Lie}(G) \longrightarrow \mathrm{Lie}(G)$$

induced by  $n_G$  is multiplication by  $n$ . It is therefore an isomorphism, as is the endomorphism induced by  $n_G$  on

$$I/I^2 = \mathrm{Lie}(G)^*.$$

Suppose first that  $\mathrm{char} k = 0$ . The group  $G$  is smooth over  $k$  (VIB, 1.6.1; see also VIIB, 3.3.1), and the canonical homomorphism

$$\mathrm{Sym}(I/I^2) \longrightarrow \mathrm{gr}_I(A)$$

is an isomorphism, where  $\mathrm{gr}_I(A)$  is the graded ring associated with the  $I$ -adic filtration. Hence  $n_G$  induces an automorphism of  $\mathrm{gr}_I(A)$ , and therefore also of the completion  $\hat{A}$ . Thus  $n_G$  is étale at the origin (cf. EGA IV<sub>4</sub>, 17.6.3).

Suppose next that  $\mathrm{char} k = p > 0$  and  $G$  has height at most 1. Then  $A$  is isomorphic to the quotient of the symmetric algebra on

$$\omega_{G/k} = I/I^2$$

by the ideal generated by the  $p$ -th powers of the elements of  $\omega_{G/k}$  (cf. 7.4). The same argument as in characteristic 0 shows that  $n_G$  induces an automorphism of  $A$ .

Suppose that  $G$  has height at most  $r$ , and that the result has been proved for groups of height at most  $r - 1$ . Let  $B, A, A'$  be the affine algebras of  $\mathrm{Fr} G, G$ , and

$$G' = \mathrm{Fr} G \backslash G,$$

and let  $n_B, n_A, n_{A'}$  be the endomorphisms induced by  $n_{\mathrm{Fr} G}, n_G, n_{G'}$ , respectively. Put

$$I' = I \cap A',$$

the maximal ideal of  $A'$ . The cartesian square

<sup>83</sup>The editors changed “étale at the origin” to “étale” in the statement and added the preceding sentence.

$$\begin{array}{ccc} \mathrm{Fr} G & \longrightarrow & G \\ \downarrow & & \downarrow \\ e & \longrightarrow & G' \end{array}$$

gives

$$B = A/I'A.$$

<sup>84</sup> The endomorphism  $n_{A'}$ , respectively  $n_B$ , is the endomorphism induced by  $n_A$  on  $A'$ , respectively on  $B$ . By VIA, 3.2,  $A$  is a faithfully flat  $A'$ -module. Since  $A'$  is an Artinian local ring, because  $G'$  is an algebraic  $k$ -group of height at most  $r - 1$ ,  $A$  is a free  $A'$ -module. The restriction of  $n_A$  to  $A'$  is  $n_{A'}$ , which is an isomorphism by the induction hypothesis. Nakayama's lemma therefore shows that  $n_A$  is an automorphism as soon as the endomorphism it induces on  $A/I'A$  is an automorphism.

That endomorphism is  $n_B$ , which is an automorphism because  $\mathrm{Fr} G$  has height at most 1. Hence  $n_A$  is an automorphism.

Finally, let  $G$  be any algebraic group over a field of characteristic  $p > 0$ . The preceding argument shows that  $n_G$  induces automorphisms of the  $k$ -schemes  $\mathrm{Fr}^r G$ . These schemes are affine over  $k$ , and their algebras are the quotients of the local algebra  $A$  by the ideal

$$I^{\{p^r\}}$$

generated by the  $p^r$ -th powers of the elements of  $I$ . Since  $n_G$  induces automorphisms of all the algebras  $A/I^{\{p^r\}}$ , passage to the projective limit shows that it induces an automorphism of  $\hat{A}$ . Thus  $n_G$  is étale at the origin (EGA IV<sub>4</sub>, 17.6.3).

**8.5. Proposition.** *Let  $G$  be a finite algebraic group of rank  $n$  over the field  $k$ . Then*

$$n_G : G \longrightarrow G$$

*is the zero morphism of  $G$ .*

Combining 8.4 and 8.5 immediately gives the following corollary.

**Corollary 8.5.1.** *Let  $G$  be a finite algebraic group of rank  $n$  over the field  $k$ . If  $n$  is prime to the characteristic of  $k$ , then  $G$  is étale over  $k$ .<sup>85</sup>*

We now prove 8.5. Let  $H$  be a normal subgroup of  $G$ , of rank  $m$  over  $k$ , and let

$$\lambda : H \times G \longrightarrow G$$

be induced by the multiplication of  $G$ . With the notation of VIA, 3.2, there is a cartesian square

$$\begin{array}{ccc} H \times G & \xrightarrow{\lambda} & G \\ \mathrm{pr}_2 \downarrow & & \downarrow \pi \\ G & \xrightarrow{\pi} & H \backslash G \end{array}$$

<sup>84</sup>The editors expanded the original argument in what follows.

<sup>85</sup>The editors added this corollary, which the original indicated implicitly by “cf. 8.4.” For another proof which does not use 8.5, see [TO70], Lemma 5.

Since  $\pi : G \rightarrow H \backslash G$  is faithfully flat and quasi-compact (VIA, 3.2), while  $\text{pr}_2$  is locally free of rank  $m$ , EGA IV<sub>2</sub>, 2.5.2 shows that  $G \rightarrow H \backslash G$  is locally free of rank  $m$ . If

$$r = \text{rk}_k(H \backslash G),$$

then

$$n = \text{rk}_k(G) = rm.$$

On the other hand, for every  $k$ -scheme  $T$  there is an exact sequence of abstract groups

$$1 \longrightarrow H(T) \longrightarrow G(T) \longrightarrow (H \backslash G)(T).$$

It is therefore clear that  $n_G$  is zero whenever  $m_H$  and  $r_{H \backslash G}$  are zero. Taking  $H = G^0$ , the identity component, the quotient  $G^0 \backslash G$  is étale (cf. VIA, 5.5.1). We may consequently assume that  $G$  is étale over  $k$ , or that it is infinitesimal (cf. 7.0).

If  $G$  is étale, extension of the base field reduces us to the case where  $k$  is algebraically closed. Then  $G$  is a constant group (cf. I, 4.1), and the assertion is classical.

If  $G$  is nonzero and infinitesimal, then  $k$  necessarily has characteristic  $p > 0$  (cf. VIB, 1.6.1 or VIIB, 3.3.1). The subgroups  $\text{Fr}^r G$  form a composition series of  $G$  whose successive quotients have height at most 1.

We are thus reduced to the case where  $G$  has height at most 1. Let  $A$ , respectively  $L$ , be the affine algebra, respectively the Lie algebra, of  $G$ , and put

$$U = U_p(L).$$

By 7.4,  $G = \mathfrak{G}_p(L)$ , and hence  $A = U^*$ . Therefore, if  $\dim_k L = r$ , the rank of  $G$  over  $k$  is  $p^r$  (cf. 5.3.3). We shall study the morphism

$$p_G : G \longrightarrow G$$

defined by taking the  $p$ -th power. It induces an endomorphism  $p_A$  of  $A$ , and, by duality, an endomorphism  $p_U$  of  $U$ .

Let  $I$  be the augmentation ideal of  $A$ . We shall prove

$$p_A(I) \subset I^p. \quad (*)$$

Assuming this, one has

$$p_A^r(I) \subset I^{p^r}.$$

Moreover,

$$I^{r(p-1)+1} = 0,$$

because  $I$  is generated by  $r$  elements whose  $p$ -th powers vanish. Since  $p^r > r(p-1)$ , we obtain  $p_A^r(I) = 0$ , and therefore  $p_G^r$  is the zero morphism.

For every integer  $s \geq 1$ , let

$$m_A^{s-1} : A^{\otimes s} \longrightarrow A, \quad \Delta_U^{s-1} : U \longrightarrow U^{\otimes s},$$

denote the maps induced by the multiplication  $m_A$  of  $A$  and the comultiplication  $\Delta_U$  of  $U$ , respectively. Then  $p_A$  is the composite

$$A \xrightarrow{\Delta_A^{p-1}} A^{\otimes p} \xrightarrow{m_A^{p-1}} A.$$

Since the transpose of  $m_A$ , respectively of  $\Delta_A$ , is  $\Delta_U$ , respectively  $m_U$ , the endomorphism

$$p_U = {}^t p_A$$

is the composite

$$U \xrightarrow{\Delta_U^{p-1}} U^{\otimes p} \xrightarrow{m_U^{p-1}} U.$$

Let  $J$  be the augmentation ideal of  $U$ .<sup>86</sup> Then

$$U = k1_U \oplus J,$$

and we write  $\pi : U \rightarrow J$  for the projection with kernel  $k1_U$ . For  $s \geq 1$ , let  $(I^s)^\perp$  be the orthogonal of  $I^s$  in  $A^* = U$ . Thus  $(I^s)^\perp$  consists of the elements  $u \in U$  for which the composite

$$I^{\otimes s} \xrightarrow{m_A^{s-1}} I \xrightarrow{u} k$$

is zero. Since the transpose of  $m_A$  is  $\Delta_U$ ,  $(I^s)^\perp$  is the vector subspace  $P_{s-1}$  consisting of those  $u \in U$  for which  $\Delta_U^{s-1}(u)$  vanishes on  $I^{\otimes s}$ . If

$$\overline{\Delta}_U^{s-1} = \pi^{\otimes s} \circ \Delta_U^{s-1},$$

then

$$(I^s)^\perp = P_{s-1} = \ker \overline{\Delta}_U^{s-1}$$

(see also VIIB, 1.3.6).

To prove (\*), it is enough to show that the transpose  $p_U = {}^t p_A$  maps  $P_{p-1}$  into

$$I^\perp = k1_U.$$

Since  $p_U(1_U) = 1_U$ , it suffices to prove

$$p_U(P_{p-1}^+) = 0, \quad P_{p-1}^+ = J \cap P_{p-1}. \quad (**)$$

An easy induction on  $s$  shows that  $P_{s-1}^+$  is spanned by the products

$$x_1 \cdots x_t, \quad 1 \leq t \leq s-1, \quad x_i \in L$$

(cf. VIIB, 4.3).

For  $x_1, \dots, x_t \in L$ , one has

$$p_U(x_1 x_2 \cdots x_t) = m_U^{p-1} \left( \prod_{j=1}^t \sum_{i=1}^p 1 \otimes \cdots \otimes \underset{\text{position } i}{x_j} \otimes \cdots \otimes 1 \right).$$

The expression inside parentheses is a sum of  $p^t$  terms  $x_h$ , indexed by the maps

$$h : \{1, \dots, t\} \longrightarrow \{1, \dots, p\}.$$

Such a map determines an ordered partition  $\mathcal{P}_h$  of  $\{1, \dots, t\}$  into at most  $p$  parts.<sup>87</sup> Indeed, let

$$i_1 < \cdots < i_q$$

be the elements in the image of  $h$ , put

$$I_a = h^{-1}(i_a) \quad (1 \leq a \leq q), \quad x_{I_a} = \prod_{j \in I_a} x_j,$$

<sup>86</sup>The editors expanded the original argument in this paragraph.

<sup>87</sup>The editors expanded the original argument and replaced the notion of a preorder by the equivalent notion of an ordered partition.

where the product is taken in increasing order. The map  $h$  corresponds to the  $p$ -tensor

$$1 \otimes \cdots \otimes x_{I_1} \otimes \cdots \otimes x_{I_q} \otimes \cdots \otimes 1,$$

where  $x_{I_a}$  occupies position  $i_a$ . Its image under the iterated multiplication is the product

$$x_{I_1} \cdots x_{I_q}.$$

This depends only on the ordered partition

$$\mathcal{P} = (I_1, \dots, I_q);$$

denote it by  $x_{\mathcal{P}}$ . For a fixed  $\mathcal{P}$ , the same term is obtained for every choice of  $i_1 < \cdots < i_q$  in  $\{1, \dots, p\}$ . Consequently,

$$p_U(x_1 x_2 \cdots x_t) = \sum_{\mathcal{P}} \binom{p}{n(\mathcal{P})} x_{\mathcal{P}},$$

where  $\mathcal{P}$  ranges over the ordered partitions of  $\{1, \dots, t\}$  into at most  $p$  parts and  $n(\mathcal{P})$  is the number of parts. Here

$$1 \leq n(\mathcal{P}) \leq \min(t, p).$$

If  $t < p$ , all the coefficients  $\binom{p}{n(\mathcal{P})}$  vanish in  $k$ , and hence

$$p_U(x_1 \cdots x_t) = 0.$$

Thus  $p_U$  vanishes on  $P_{p-1}^+$ . This proves (\*\*), then (\*), and completes the proof of 8.5.

**Corollary 8.5.2.**<sup>88</sup> *Let  $S$  be a reduced scheme and  $G$  a finite locally free  $S$ -group of rank  $n$ . Then*

$$n_G : G \longrightarrow G$$

*is the zero morphism of  $G$ .*

Let  $S'$  be the disjoint union of the  $\mathrm{Spec} \mathcal{O}_{S,\eta}$ , where  $\eta$  ranges over the maximal points of  $S$ . Since  $S$  is reduced, the morphism  $S' \rightarrow S$  is schematically dominant. The same is therefore true of

$$f : G_{S'} \longrightarrow G,$$

because  $G$  is finite locally free over  $S$  (cf. EGA IV<sub>3</sub>, 11.10.5). Since  $G \rightarrow S$  is affine, and hence separated, the coincidence locus of  $n_G$  and the zero morphism is a closed subscheme of  $G$ . By 8.5 it contains the image of  $f$ , and therefore equals  $G$ . Thus  $n_G$  is the zero morphism.

**Remark 8.5.3.**

A theorem of P. Deligne (see [TO70], p. 4) also says that if  $G$  is a commutative finite locally free  $S$ -group of rank  $n$  over an arbitrary base  $S$ , then  $n_G = 0$ .

<sup>88</sup>The editors added this corollary, which is noted in Exposé VIII, Remark 7.3.1.

# Bibliography

- [BAI] N. Bourbaki, *Algèbre*, Chapters I–III, Hermann, 1974; Chapter X, Masson, 1980.
- [BAC] N. Bourbaki, *Algèbre commutative*, Chapters I–IV, Masson, 1985.
- [BLie] N. Bourbaki, *Groupes et algèbres de Lie*, Chapter I, Hermann, 1971.
- [DG70] M. Demazure and P. Gabriel, *Groupes algébriques*, Masson & North-Holland, 1970.
- [Ja03] J. C. Jantzen, *Representations of Algebraic Groups*, Academic Press, 1987; 2nd ed., American Mathematical Society, 2003.
- [TO70] J. Tate and F. Oort, “Group schemes of prime order,” *Annales scientifiques de l’École Normale Supérieure* **3** (1970), 1–21.

# Exposé VII<sub>B</sub>

## Infinitesimal Study of Group Schemes

by *P. Gabriel*

### B) Formal Groups

<sup>0</sup> The study of formal groups is usually extremely simple. If this does not appear clearly in the pages that follow, the responsibility lies with an arithmetician who claims to know formal groups over “something other than fields.”<sup>1</sup> We have therefore developed, for formal groups “locally free over projective limits of Artinian rings,” the generalities ordinarily stated for formal groups defined over a field. For a more detailed study of the latter, we refer to the 1964/65 Heidelberg–Strasbourg algebraic geometry seminar.<sup>2</sup>

### 0. Review of Pseudocompact Rings and Modules

This section contains a few technical preliminaries. We recall and complete here several results from [CA] (*Des catégories abéliennes*, Bull. Soc. Math. France **90** (1962)).

**0.1.** A *left pseudocompact ring* is a separated, complete topological ring with identity which has a basis of neighborhoods of 0 consisting of left ideals  $I$  of finite colength, that is,

$$\text{length}_A(A/I) < +\infty.$$

We shall assume here that  $A$  is commutative, so that no distinction need be made “between left and right.”

In particular, the quotients  $A/I$  are Artinian rings, and  $A$  identifies with the topological projective limit of these rings, each endowed with the discrete topology.

A complete Noetherian local ring  $(A, \mathfrak{m})$  is plainly pseudocompact.<sup>3</sup>

---

<sup>0</sup>Editors’ note: Version of 13 October 2024.

<sup>1</sup>The interest of formal groups over a complete Noetherian local ring appears, for example, in the work of Lubin and Tate (cf. [LT65]). The study of formal groups over an arbitrary base, and of lifting and deformation questions—in particular for Barsotti–Tate groups (“ $p$ -divisible groups”)—has given rise to an abundant literature; see, for example, [LT66, Ta67, Gr74, Me72, La75, Fo77, Il85, Br00]. In particular, the results of the present exposé are largely taken up again in Chapter I of [Fo77].

<sup>2</sup>The editors found only such a seminar dated 1965/66 and entitled “Linear Algebraic Groups,” in which the notion of a formal group does not appear; see instead [De72].

<sup>3</sup>When it is endowed with the  $\mathfrak{m}$ -adic topology.



**0.1.1.** Every closed ideal  $I$  of  $A$  is the intersection of the open ideals that contain it.<sup>4</sup> Every closed maximal ideal is therefore open. Moreover, if  $\mathfrak{l}$  is an open ideal of  $A$ , the maximal ideals of  $A/\mathfrak{l}$  correspond bijectively to the maximal ideals  $\mathfrak{m}$  of  $A$  containing  $\mathfrak{l}$ ; the latter are thus open and closed. Consequently, every closed maximal ideal is an open maximal ideal (and hence closed), the converse being clear. We denote the set of these ideals by  $\Upsilon(A)$ .

If  $\mathfrak{l}$  is an open ideal of  $A$  and  $\mathfrak{m} \in \Upsilon(A)$ , the localization  $(A/\mathfrak{l})_{\mathfrak{m}}$  is a local ring when  $\mathfrak{m}$  contains  $\mathfrak{l}$ , and is zero otherwise. Since  $A/\mathfrak{l}$  is Artinian, it is a direct product of finitely many local rings; thus

$$A/\mathfrak{l} \simeq \prod_{\mathfrak{m} \in \Upsilon(A)} (A/\mathfrak{l})_{\mathfrak{m}}.$$

It follows that there are “canonical” isomorphisms

$$A \simeq \varprojlim_{\mathfrak{l}} (A/\mathfrak{l}) \simeq \varprojlim_{\mathfrak{l}} \prod_{\mathfrak{m}} (A/\mathfrak{l})_{\mathfrak{m}} \simeq \prod_{\mathfrak{m}} \varprojlim_{\mathfrak{l}} (A/\mathfrak{l})_{\mathfrak{m}} \simeq \prod_{\mathfrak{m}} A_{\mathfrak{m}},$$

where

$$A_{\mathfrak{m}} = \varprojlim_{\mathfrak{l}} (A/\mathfrak{l})_{\mathfrak{m}}.$$

This *local component*  $A_{\mathfrak{m}}$  is a filtered projective limit of Artinian local rings endowed with the discrete topology. It is therefore a local ring which is pseudocompact for the projective-limit topology.<sup>5</sup>

**0.1.2.** Let  $r(A)$  be the intersection of the open maximal ideals of  $A$ , that is, the cartesian product of the ideals  $\mathfrak{m}A_{\mathfrak{m}}$  under the identification

$$A = \prod_{\mathfrak{m}} A_{\mathfrak{m}}.$$

For every open ideal  $\mathfrak{l}$  of  $A$ , the image of  $r(A)$  in  $A/\mathfrak{l}$  is contained in the radical of  $A/\mathfrak{l}$ . Some power of this image is therefore zero, so  $r(A)^n \subset \mathfrak{l}$  for all sufficiently large  $n$ . Thus the sequence  $r(A)^n$  tends to 0.

The same is true of  $x^n$  when  $x \in r(A)$ . In other words, every element of  $r(A)$  is topologically nilpotent, and the converse is clear. It follows that the sequence with general term

$$1 + x + \cdots + x^n$$

converges, and converges to  $1/(1-x)$ , whenever  $x \in r(A)$ . This shows that  $r(A)$  is the Jacobson radical of  $A$ , that is, the intersection of all maximal ideals of  $A$  (cf. Bourbaki, *Algèbre*, Chap. 8, §6, Th. 1).<sup>6</sup>

**Remarks.**<sup>7</sup>

a) If  $\mathfrak{p}$  is an open prime ideal of  $A$ , then, since  $A/\mathfrak{p}$  is Artinian,  $\mathfrak{p}$  is maximal. Consequently,  $\Upsilon(A)$  is the set of open prime ideals of  $A$ .

<sup>4</sup>Indeed, if  $x \notin I$ , there is an open ideal  $\mathfrak{l}$  such that  $(x + \mathfrak{l}) \cap I = \emptyset$ ; then  $I + \mathfrak{l}$  is an open ideal that does not contain  $x$ . In what follows, the editors have also made explicit the fact that every “closed maximal” ideal is maximal and open.

<sup>5</sup>Notice that  $A_{\mathfrak{m}}$  is the localization of  $A$  at the maximal ideal  $\mathfrak{m}$ . Indeed, the identity element  $e_{\mathfrak{m}}$  of  $A_{\mathfrak{m}}$  is an idempotent of  $A$  not belonging to  $\mathfrak{m}$ . Since  $A = A_{\mathfrak{m}} \times B$ , where  $B = A(1 - e_{\mathfrak{m}})$ , the ring  $A_{\mathfrak{m}}$  identifies with the localization  $A_{e_{\mathfrak{m}}}$ , and hence also with  $S^{-1}A$ , where  $S = A \setminus \mathfrak{m}$ . Furthermore,  $e_{\mathfrak{m}}(1 - e_{\mathfrak{m}}) = 0$ , so  $\mathfrak{m}$  contains  $1 - e_{\mathfrak{m}}$ , hence also  $B$ ; thus  $\mathfrak{m}$  identifies with  $\mathfrak{m}A_{\mathfrak{m}} \times B$ .

<sup>6</sup>Indeed, let  $x \in r(A)$ . If  $\mathfrak{m}$  were a maximal ideal not containing  $x$ , there would be  $y \in A$  such that  $1 - xy \in \mathfrak{m}$ . But  $xy \in r(A)$ , so  $1 - xy$  is invertible, a contradiction. Notice the following consequence: if  $\Upsilon(A) = \{\mathfrak{m}_1, \dots, \mathfrak{m}_r\}$  is finite, then the  $\mathfrak{m}_i$  are all the maximal ideals of  $A$ .

<sup>7</sup>The editors added these remarks in order to compare the definition of the formal spectrum  $\mathrm{Spf}(A)$  given in 1.1 with those in EGA I, 10.1.2 and 10.4.2.

b) Each  $\mathfrak{m}A_{\mathfrak{m}}$  is an ideal of definition of  $A_{\mathfrak{m}}$ , that is, an open ideal  $I$  for which  $I^n$  tends to 0 (cf. EGA 0<sub>I</sub>, 7.1.2). Consequently,

$$\mathrm{Spec}(A_{\mathfrak{m}}/\mathfrak{m}A_{\mathfrak{m}}),$$

endowed with the topological ring  $A_{\mathfrak{m}}$ , is an affine formal scheme in the sense of EGA I, 10.1.2.

c) The topological ring  $A$  is admissible in the sense of EGA 0<sub>I</sub>, 7.1.2 if and only if  $r(A)$  is open (and hence an ideal of definition), and this holds if and only if  $\Upsilon(A)$  is finite. In that case the affine formal scheme

$$\mathrm{Spf}(A) = \mathrm{Spec}(A/r(A))$$

(cf. EGA I, 10.1.2) has  $\Upsilon(A)$ , endowed with the discrete topology, as its underlying space; the sections of its structure sheaf over a subset  $E \subset \Upsilon(A)$  form the ring

$$\prod_{\mathfrak{m} \in E} A_{\mathfrak{m}}.$$

d) Let  $A$  be an arbitrary pseudocompact ring. In 1.1 below, the space  $\Upsilon(A)$  is endowed with the discrete topology and with the sheaf of rings whose ring of sections over any subset  $E$  is

$$\prod_{\mathfrak{m} \in E} A_{\mathfrak{m}}.$$

By b), every point then has an open neighborhood which is an affine formal scheme; this therefore defines a formal scheme (EGA I, 10.4.2), denoted by  $\mathrm{Spf}(A)$ . For this formal scheme to be affine, it must be quasi-compact, so  $\Upsilon(A)$  must be finite; in that case  $\mathrm{Spf}(A)$  agrees with the definition of EGA I, 10.1.2.

**0.1.3.** If  $A$  and  $B$  are pseudocompact rings, a *homomorphism* from  $A$  to  $B$  is, by definition, a continuous map compatible with addition, multiplication, and the identity elements. Such a homomorphism sends a topologically nilpotent element of  $A$  to a topologically nilpotent element of  $B$ ; it therefore maps the radical  $r(A)$  into the radical  $r(B)$ .

**0.2.** Let  $A$  be a commutative pseudocompact ring. A *pseudocompact  $A$ -module*  $M$  is a separated, complete topological  $A$ -module having a basis of neighborhoods of 0 consisting of submodules  $M'$  for which  $M/M'$  has finite length over  $A$ .

If  $M$  and  $N$  are pseudocompact  $A$ -modules, a *morphism* from  $M$  to  $N$  is, by definition, a continuous  $A$ -linear map. We write

$$\mathrm{Hom}_c(M, N)$$

for the group of these morphisms.

**Proposition 0.2.B.**<sup>8</sup>

- (i) The pseudocompact  $A$ -modules form an abelian category, denoted  $\mathrm{PC}(A)$ . In particular, for every morphism  $f : M \rightarrow N$ , the submodule  $\mathrm{Im}(f)$  is complete and hence closed in  $N$ .
- (ii) The pseudocompact  $A$ -modules of finite length, whose topology is therefore discrete, form a system of cogenerators of  $\mathrm{PC}(A)$ .

---

<sup>8</sup>The editors have stated the results of this paragraph separately in the following proposition, and have then indicated the steps of the proof; cf. [CA], § IV.3, or [DG70], § V.2.

- (iii) *Infinite products and filtered projective limits are exact; in other words,  $\text{PC}(A)$  satisfies axiom (AB5\*).<sup>9</sup>*

For the reader's convenience, we briefly indicate the steps of the proof. First we use the following lemma ([CA], § IV.3, Lemma 1; for the proof, see [BEns], III, § 7.4, Theorem 1 and Example 2).

**Lemma 0.2.C.** *Let  $B$  be a ring, let  $I$  be a filtered ordered set, and let  $(M_i)$  and  $(N_i)$  be two projective systems of left  $B$ -modules indexed by  $I$ . Let*

$$(s_i) : (M_i) \longrightarrow (N_i)$$

*be a morphism of projective systems such that  $s_i$  is surjective and has Artinian kernel for every  $i$ . Then the map*

$$\varprojlim_i s_i : \varprojlim_i M_i \longrightarrow \varprojlim_i N_i$$

*is surjective.*

**Corollary 0.2.D** ([CA], § IV.3, Propositions 10 and 11). *Let  $M$  be a pseudocompact  $A$ -module.*

- (i) *If  $K$  is a closed submodule of  $M$ , then  $M/K$ , endowed with the quotient topology, is a pseudocompact  $A$ -module.*
- (ii) *Let  $(M_i)$  be a filtered decreasing family of closed submodules of  $M$ .*
  - (a) *The canonical map*

$$M \longrightarrow \varprojlim_i M/M_i$$

*is surjective and has kernel  $\bigcap_i M_i$ .*

- (b) *For every closed submodule  $N$  of  $M$ , one has*

$$N + \bigcap_i M_i = \bigcap_i (N + M_i).$$

*Proof.* Let  $(L_j)$  be the filtered decreasing family of open submodules of  $M$ . We endow  $M/K$  with the quotient topology; that is, a basis of neighborhoods of 0 is formed by the open submodules  $(K + L_j)/K$ . Since  $K$  is closed, it is equal to the intersection of the  $K + L_j$ , and hence the map

$$\tau : M/K \longrightarrow \varprojlim_j M/(K + L_j)$$

is injective. It is also open, since the right-hand side is the inverse limit of the discrete modules  $M/(K + L_j)$ . Moreover, for each  $j$ , the map

$$t_j : M/L_j \longrightarrow M/(K + L_j)$$

is surjective and has Artinian kernel. Thus, by the preceding lemma, the map  $t$  in the following commutative diagram is surjective:

---

<sup>9</sup>Cf. [Gr57], I, § 1.5 and Proposition 1.8. One may also consult, for example, [Po73], § 2.8, or [We95], Appendix A.4.

$$\begin{array}{ccc}
M & \xrightarrow[p \sim]{} & \varprojlim_j M/L_j \\
\downarrow & & \downarrow t \\
M/K & \xrightarrow{\tau} & \varprojlim_j M/(K + L_j).
\end{array}$$

Since  $p$  is an isomorphism because  $M$  is complete, it follows that  $\tau$  is surjective, and hence is an isomorphism. This proves (i).

Let us prove (ii)(a). By the foregoing, for every  $i$  there is an isomorphism

$$M/M_i \xrightarrow{\sim} \varprojlim_j M/(M_i + L_j),$$

so the two horizontal arrows in the following commutative diagram are isomorphisms:

$$\begin{array}{ccc}
M & \xrightarrow[p \sim]{} & \varprojlim_j M/L_j \\
\downarrow g & & \downarrow s \\
\varprojlim_i M/M_i & \xrightarrow{\sim} & \varprojlim_{i,j} M/(M_i + L_j).
\end{array}$$

Moreover, for each  $j$ , the family of submodules  $(M_i + L_j)/L_j$  has a least element, since  $M/L_j$  is Artinian. Consequently, the morphism

$$s_j : M/L_j \longrightarrow \varprojlim_i M/(L_j + M_i)$$

is surjective; therefore, by the preceding lemma,  $s$  is surjective. It follows that  $g$  is surjective. Finally, the kernel of  $g$  is the inverse limit of the  $M_i$ , that is, their intersection. This proves (a).

Let us deduce (b) from this. Since  $N$  is a closed submodule, and hence separated and complete, it is a pseudocompact module for the topology induced by that of  $M$ . Therefore, by (a), the morphisms  $f$  and  $g$  in the following commutative exact diagram are surjective:

$$\begin{array}{ccccccc}
0 & \longrightarrow & N & \longrightarrow & M & \longrightarrow & M/N \longrightarrow 0 \\
& & \downarrow f & & \downarrow g & & \downarrow h \\
0 & \longrightarrow & \varprojlim_i N/(N \cap M_i) & \longrightarrow & \varprojlim_i M/M_i & \longrightarrow & \varprojlim_i M/(N + M_i).
\end{array}$$

The “snake lemma” then shows that the sequence

$$0 \longrightarrow \ker f \longrightarrow \ker g \longrightarrow \ker h \longrightarrow 0$$

is exact, and the equality

$$N + \bigcap_i M_i = \bigcap_i (N + M_i)$$

follows.

We can now prove Proposition 0.2.B. Let  $f : M \rightarrow N$  be a morphism of pseudocompact  $A$ -modules. Then  $K = \ker(f)$  is a closed submodule of  $M$ , and hence is separated and complete; thus  $K$ , with the topology induced by that of  $M$ , is a pseudocompact module. By 0.2.D (i), the module  $M/K$ , endowed with the quotient topology, is pseudocompact.

Let us show that the continuous bijective morphism

$$M/K \longrightarrow \operatorname{Im}(f)$$

is bicontinuous. After identifying  $M/K$  with  $\operatorname{Im}(f)$ , it is enough to show that the topology  $T$  induced by that of  $N$  is finer than the quotient topology  $Q$ .<sup>10</sup> Let  $(L_j)$ , respectively  $(N_i)$ , be the filtered decreasing families of open submodules of  $M$ , respectively  $N$ , and put

$$N'_i = N_i \cap \operatorname{Im}(f).$$

Let  $P = (K + L_j)/K$  be a submodule of  $M/K$  open for  $Q$ . Since  $M/(K + L_j)$  is Artinian, the family  $N'_i + P$  has a least element  $N'_{i_0} + P$ . The  $N'_i$  are open, and hence closed, for  $T$ , and therefore also closed for  $Q$ . It follows from 0.2.D (ii)(b) that

$$N'_{i_0} + P = \bigcap_i (N'_i + P) = P + \bigcap_i N'_i = P,$$

whence  $N'_{i_0} \subset P$ . Thus  $P$  is open for  $T$ , and  $M/K \rightarrow \operatorname{Im}(f)$  is consequently an isomorphism of pseudocompact modules.

In particular,  $\operatorname{Im}(f)$  is complete for  $T$ , and hence closed in  $N$ . By 0.2.D (i) once more,  $\operatorname{Coker}(f)$ , endowed with the quotient topology, is pseudocompact. This proves that  $\operatorname{PC}(A)$  is an abelian category.

Arbitrary inverse limits also exist in  $\operatorname{PC}(A)$ : if  $(M_i)$  is an inverse system of pseudocompact modules, its inverse limit has as underlying module the inverse limit of the underlying modules, endowed with the inverse-limit topology. Moreover, if one has a family of exact sequences in  $\operatorname{PC}(A)$ ,

$$0 \longrightarrow K_i \longrightarrow M_i \longrightarrow Q_i \longrightarrow 0,$$

then the sequence

$$0 \longrightarrow \prod_i K_i \longrightarrow \prod_i M_i \longrightarrow \prod_i Q_i \longrightarrow 0$$

is exact. Assertion (iii) of 0.2.B follows, since in every abelian category in which arbitrary products exist, conditions (a) and (b) of 0.2.D are equivalent, and are equivalent to the exactness of filtered inverse limits (cf. [Mi65], III, 1.2–1.9, or [Po73], Chap. 2, Th. 8.6).

Finally, every pseudocompact module  $M$  is a submodule of the product

$$\prod_L M/L,$$

where  $L$  ranges over the open submodules of  $M$ . Thus the objects of finite length form a system of cogenerators of  $\operatorname{PC}(A)$ . (Moreover, every object of length  $n$  is isomorphic to a quotient  $A^n/L$ , where  $L$  is an open submodule of  $A^n$  of colength  $n$ ; these quotients therefore form a *set* of cogenerators.) This completes the proof of 0.2.B.

Let  $(\operatorname{Ab})$ <sup>11</sup> be the category of abelian groups and let  $\operatorname{LF}(A)$  be the full subcategory of  $\operatorname{PC}(A)$  formed by the objects of finite length. For every object  $M$  of  $\operatorname{PC}(A)$ , let  $h_c^M$  denote the functor

$$\operatorname{LF}(A) \longrightarrow (\operatorname{Ab}), \quad N \longmapsto \operatorname{Hom}_c(M, N).$$

<sup>10</sup>The French re-edition says here that  $Q$  is finer than  $T$ ; the ensuing argument proves the reverse inclusion of open sets, which is the one needed for bicontinuity.

<sup>11</sup>The editors inserted Proposition 0.2.E and Corollary 0.2.F into this paragraph; they will be useful in 0.2.2. In the original, these results appeared in 0.3.

By [CA], § II.4, Theorem 1, Lemma 4, and Corollary 1, we have the following results.<sup>12</sup>

**Proposition 0.2.E.** *The functor  $M \mapsto h_c^M$  is an anti-equivalence from  $\mathrm{PC}(A)$  to the category*

$$\mathrm{Lex}(\mathrm{LF}(A), (\mathrm{Ab}))$$

*of left-exact functors  $\mathrm{LF}(A) \longrightarrow (\mathrm{Ab})$ .*

**Corollary 0.2.F.**

- (i) *An object  $P$  of  $\mathrm{PC}(A)$  is projective if and only if the functor  $h_c^P$  is exact; that is, if and only if the functor  $\mathrm{Hom}_c(P, -)$  is exact on  $\mathrm{LF}(A)$ .*
- (ii) *Let  $(M_i)$  be a filtered projective system<sup>13</sup> of objects of  $\mathrm{PC}(A)$ . For every object  $N$  of  $\mathrm{LF}(A)$ , there is an isomorphism, functorial in  $N$ ,*

$$\mathrm{Hom}_c\left(\varprojlim_i M_i, N\right) \simeq \varinjlim_i \mathrm{Hom}_c(M_i, N).$$

- (iii) *Every filtered projective limit and every product<sup>13</sup> of projective objects of  $\mathrm{PC}(A)$  is a projective object of  $\mathrm{PC}(A)$ .*

Finally, Corollary 0.2.F gives the following result.

**Corollary 0.2.G.** *Let  $(M_i)_{i \in I}$  be a family of objects of  $\mathrm{PC}(A)$ . Then  $\prod_{i \in I} M_i$  is projective if and only if each  $M_i$  is projective.*

Indeed, for every  $N \in \mathrm{Ob} \mathrm{LF}(A)$ , one has

$$\mathrm{Hom}_c\left(\prod_i M_i, N\right) \simeq \bigoplus_i \mathrm{Hom}_c(M_i, N).$$

**0.2.1.** Each local component  $A_{\mathfrak{m}}$  of  $A$  is a direct factor of  $A$ , hence a projective object of  $\mathrm{PC}(A)$  ( $A$  is manifestly projective). Moreover,

$$S_{\mathfrak{m}} = A_{\mathfrak{m}}/\mathfrak{m}A_{\mathfrak{m}}$$

is the unique simple quotient of  $A_{\mathfrak{m}}$ , so  $A_{\mathfrak{m}}$  is indecomposable. On the other hand, every simple object of  $\mathrm{PC}(A)$  is isomorphic to a unique  $S_{\mathfrak{m}}$ . By [CA], IV, § 3, Corollary 1 of Theorem 3,<sup>14</sup> we therefore have:

**Proposition.**

- (i) *Every projective object of  $\mathrm{PC}(A)$  is a direct product of indecomposable projective objects, uniquely determined up to isomorphism.*
- (ii) *Every indecomposable projective object is isomorphic to a unique  $A_{\mathfrak{m}}$ , where  $\mathfrak{m} \in \Upsilon(A)$ .*

**Definition.** A pseudocompact  $A$ -module  $M$  is said to be *topologically free* if it is isomorphic to the product of a family  $(A_i)$  of copies of  $A$ .

<sup>12</sup>Indeed,  $\mathrm{PC}(A)$  has a set of Artinian cogenerators, filtered projective limits are exact in it, and  $\mathrm{LF}(A)$  is the subcategory of Artinian objects. The dual category  $\mathrm{PC}(A)^\circ$  therefore has a set of Noetherian generators, and filtered inductive limits are exact in it. By the proof of [CA], § II.4, Theorem 1, the functor  $M \mapsto \mathrm{Hom}_c(M, -)$  is an anti-equivalence from  $\mathrm{PC}(A)$  to  $\mathrm{Lex}(\mathrm{LF}(A), (\mathrm{Ab}))$ . Likewise, Lemma 4 and Corollary 1 of *loc. cit.* state results “dual” to those of Corollary 0.2.F. For a “direct” proof, see [DG70], § V.2, Theorem 3.1, Lemma 3.5, and Corollaries 3.3 and 3.4.

<sup>13</sup>Since every infinite product is a filtered projective limit of finite products, every additive functor that commutes with filtered projective limits also commutes with infinite products.

<sup>14</sup>This refers to the “dual” statements established in *loc. cit.*, § 2, Theorem 2, and § 1, Proposition 2. For a “direct” proof, see [DG70], V, § 2, Theorem 4.5 and Example 4.6(b).

In this case, a family  $(m_i)$  of elements of  $M$  is called a *pseudobasis* of  $M$  if the  $A$ -linear maps  $A_i \rightarrow M$  sending the unit element of  $A_i$  to  $m_i$  extend to an isomorphism

$$\prod_i A_i \xrightarrow{\sim} M.$$

**0.2.2.** <sup>15</sup> If  $M$  is a pseudocompact  $A$ -module, we write  $M^\dagger$  for the  $A$ -module, without topology,

$$\mathrm{Hom}_c(M, A).$$

Conversely, if  $N$  is an  $A$ -module, we write

$$N^* = \mathrm{Hom}_A(N, A)$$

for its dual, endowed with the topology of pointwise convergence. Thus a basis of neighborhoods of 0 in  $N^*$  is formed by the submodules

$$\mathcal{V}(x, \mathfrak{l}) = \{f \in N^* \mid f(x) \in \mathfrak{l}\},$$

where  $x \in N$  and  $\mathfrak{l}$  is an open ideal of  $A$ .

This makes  $N^*$  a pseudocompact  $A$ -module. Indeed, first suppose that  $N = A$ . Then  $N^* = A$ , endowed with its topology as a pseudocompact ring. If  $N$  is a free module  $A^{(I)}$ , then  $N^*$  is the product  $A^I$ , endowed with the product topology. Furthermore, for every morphism  $\varphi : N_1 \rightarrow N_2$ , the transpose

$${}^t\varphi : N_2^* \rightarrow N_1^*$$

is continuous, because the inverse image under  ${}^t\varphi$  of a submodule  $\mathcal{V}(x_1, \mathfrak{l})$  of  $N_1^*$  is precisely the submodule  $\mathcal{V}(\varphi(x_1), \mathfrak{l})$  of  $N_2^*$ . For arbitrary  $N$ , choose a presentation

$$A^{(J)} \xrightarrow{\varphi} A^{(I)} \xrightarrow{\pi} N \rightarrow 0.$$

Then  $N^*$  is the kernel of the continuous morphism

$${}^t\varphi : A^I \rightarrow A^J,$$

and hence  $N^*$  is pseudocompact.

When  $A$  is Artinian (in which case one may take  $\mathfrak{l} = 0$  above), Corollary 0.2.F gives:

**Proposition.** *When  $A$  is Artinian, the functors*

$$P \mapsto P^\dagger \quad \text{and} \quad Q \mapsto Q^*,$$

where  $P$ , respectively  $Q$ , is a projective object of  $\mathrm{PC}(A)$ , respectively a projective  $A$ -module, establish an anti-equivalence between the category of projective pseudocompact  $A$ -modules and the category of projective  $A$ -modules.<sup>16</sup>

In particular, when  $A$  is a field  $k$ , the functor

$$P \mapsto P^\dagger$$

<sup>15</sup>The editors have spelled out the results of this paragraph, which play an important role below; cf. 1.2.3, 1.3.5, 2.2.1, and so on.

<sup>16</sup>Recall also that, over an Artinian ring, a module is projective if and only if it is flat; see, for example, Exposé VI<sub>B</sub>, editors' note (54), or 0.3.7 below.

is an anti-equivalence between the category of all pseudocompact  $k$ -modules (also called *linearly compact  $k$ -vector spaces*) and the category of  $k$ -vector spaces.<sup>17</sup>

**0.3.**<sup>18</sup> Let  $L$  and  $M$  be two pseudocompact  $A$ -modules. The functor

$$\mathrm{LF}(A) \longrightarrow (\mathrm{Ab}), \quad N \longmapsto \mathrm{Bil}_c(L \times M, N),$$

where  $\mathrm{Bil}_c(L \times M, N)$  denotes the abelian group of continuous  $A$ -bilinear maps from  $L \times M$  to  $N$ , is left exact.

By 0.2.E, there therefore exists a pseudocompact  $A$ -module  $L \widehat{\otimes}_A M$ , unique up to unique isomorphism, representing this functor; that is, for every object  $N$  of  $\mathrm{LF}(A)$ , there is a functorial isomorphism

$$\mathrm{Hom}_c(L \widehat{\otimes}_A M, N) \simeq \mathrm{Bil}_c(L \times M, N).$$

Moreover,  $L \widehat{\otimes}_A M$  identifies with the projective limit

$$P = \widehat{L \otimes_A M}$$

of the discrete  $A$ -modules

$$(L/L') \otimes_A (M/M'),$$

where  $L'$  and  $M'$  range respectively over the open submodules of  $L$  and  $M$ .

Indeed, let  $\varphi : L \times M \rightarrow N$  be a continuous bilinear map into a discrete  $A$ -module  $N$  of finite length. By Lemma 0.3.1 below, there exist open submodules  $L'$  and  $M'$  of  $L$  and  $M$  such that

$$\varphi(L' \times M) = \varphi(L \times M') = \{0\}.$$

This means that the map

$$\bar{\varphi} : L \otimes_A M \longrightarrow N$$

induced by  $\varphi$  has the form  $\varphi' \circ p$ , where  $p$  is the canonical projection from  $L \otimes_A M$  onto  $(L/L') \otimes_A (M/M')$ . If  $\widehat{\varphi}$  denotes the composite

$$P \longrightarrow (L/L') \otimes_A (M/M') \xrightarrow{\varphi'} N,$$

then the map  $\varphi \mapsto \widehat{\varphi}$  is a functorial bijection from  $\mathrm{Bil}_c(L \times M, N)$  onto  $\mathrm{Hom}_c(P, N)$ . Hence

$$P \simeq L \widehat{\otimes}_A M.$$

Thus the pseudocompact module  $L \widehat{\otimes}_A M$  is the separated completion of  $L \otimes_A M$  for the linear topology defined by the kernels of the canonical projections

$$L \otimes_A M \longrightarrow (L/L') \otimes_A (M/M'),$$

and it will be called the *completed tensor product* of  $L$  and  $M$ .

If  $x \in L$  and  $y \in M$ , the image of  $x \otimes_A y$  in  $L \widehat{\otimes}_A M$  will be denoted by  $x \widehat{\otimes}_A y$ .

**Lemma 0.3.1.** *Let  $L$ ,  $M$ , and  $N$  be pseudocompact  $A$ -modules, with  $N$  of finite length. If  $\varphi : L \times M \rightarrow N$  is a continuous  $A$ -bilinear map, there exist open submodules  $L'$  and  $M'$  of  $L$  and  $M$  such that*

$$\varphi(L' \times M) = \varphi(L \times M') = \{0\}.$$

<sup>17</sup>The direct sum in  $\mathrm{PC}(k)$  of a family  $(V_i)_{i \in I}$  of linearly compact  $k$ -vector spaces is  $\left(\prod_{i \in I} V_i^\dagger\right)^*$ .

<sup>18</sup>Editors' note: This paragraph has been modified to take account of the additions made in 0.2.



*Proof.* The inverse image  $\varphi^{-1}(0)$  is an open neighborhood of  $(0, 0)$ , and therefore contains an open set of the form  $L_1 \times M_1$ , where  $L_1$  and  $M_1$  are open submodules of  $L$  and  $M$ . Since  $L/L_1$  has finite length, there are elements  $x_1, \dots, x_r$  of  $L$  such that

$$L_1 + Ax_1 + \dots + Ax_r = L.$$

If  $M' \subset M_1$  is sufficiently small, then  $\varphi(x_i, M') = 0$  for every  $i$ , since the map  $y \mapsto \varphi(x_i, y)$  is continuous. It follows that  $\varphi(L, M') = \{0\}$ . Likewise,  $\varphi(L', M) = \{0\}$  when  $L'$  is sufficiently small.  $\square$

**Corollary 0.3.1.1.**<sup>19</sup> *Let  $M$  be a pseudocompact  $A$ -module.*

(i) *For every open submodule  $M'$ , there exists an open ideal  $\mathfrak{l}$  of  $A$  such that*

$$\overline{\mathfrak{l}M} \subset M'.$$

(ii) *Consequently,*

$$M \simeq \varprojlim_{\mathfrak{l}} M/\overline{\mathfrak{l}M},$$

*where  $\mathfrak{l}$  ranges over the filtered projective system of open ideals of  $A$ , and every  $M/\overline{\mathfrak{l}M}$  is endowed with the quotient topology, which makes it a pseudocompact module; cf. 0.2.D.*

*Proof.* Consider the map

$$\varphi : A \times M \longrightarrow M/M', \quad (a, m) \longmapsto am + M'.$$

By 0.3.1, there is an open ideal  $\mathfrak{l}$  of  $A$  such that  $\mathfrak{l}M \subset M'$ . Since  $M'$  is also closed, it contains  $\overline{\mathfrak{l}M}$ . Since the intersection of the open submodules of  $M$  is zero, it follows that

$$\bigcap_{\mathfrak{l}} \overline{\mathfrak{l}M} = (0).$$

On the other hand, by 0.2.D the map

$$\phi : M \longrightarrow \varprojlim_{\mathfrak{l}} M/\overline{\mathfrak{l}M}$$

is surjective. By the proof of 0.2.B,  $\phi$  therefore induces an isomorphism

$$M/\text{Ker}(\phi) \xrightarrow{\sim} \varprojlim_{\mathfrak{l}} M/\overline{\mathfrak{l}M}.$$

But we have just seen that

$$\text{Ker}(\phi) = \bigcap_{\mathfrak{l}} \overline{\mathfrak{l}M} = (0).$$

$\square$

**Remark 0.3.1.2.**<sup>20</sup> The completed tensor product satisfies the usual associativity condition: if  $L$ ,  $M$ , and  $P$  are pseudocompact  $A$ -modules, there is a canonical isomorphism

$$(L \widehat{\otimes}_A M) \widehat{\otimes}_A P \simeq L \widehat{\otimes}_A (M \widehat{\otimes}_A P).$$

<sup>19</sup>Editors' note: This corollary has been added.

<sup>20</sup>Editors' note: This remark has been added.

Indeed, both objects represent the functor which associates to every object  $N$  of  $\text{LF}(A)$  the abelian group of continuous  $A$ -trilinear maps from  $L \times M \times P$  to  $N$ .

**0.3.2.** Let

$$L' \xrightarrow{f} L \xrightarrow{g} L'' \longrightarrow 0$$

be an exact sequence, and let  $M$  be an object of  $\text{PC}(A)$ . It is clear that for every object  $N$  of  $\text{LF}(A)$ , the induced sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Bil}_c(L'' \times M, N) & \longrightarrow & \text{Bil}_c(L \times M, N) & \longrightarrow & \text{Bil}_c(L' \times M, N) \\ & & \parallel & & \parallel & & \parallel \\ 0 & \longrightarrow & \text{Hom}_c(L'' \widehat{\otimes}_A M, N) & \longrightarrow & \text{Hom}_c(L \widehat{\otimes}_A M, N) & \longrightarrow & \text{Hom}_c(L' \widehat{\otimes}_A M, N) \end{array}$$

are exact. By 0.2.E, this is equivalent to saying that the sequence

$$L' \widehat{\otimes}_A M \xrightarrow{f \widehat{\otimes} M} L \widehat{\otimes}_A M \xrightarrow{g \widehat{\otimes} M} L'' \widehat{\otimes}_A M \longrightarrow 0 \quad (*)$$

is exact. Hence:

**Corollary.** *For every pseudocompact  $A$ -module  $M$ , the functor  $-\widehat{\otimes}_A M$  is right exact.*

In particular, take  $L = A$ , let  $f$  be the inclusion of a closed ideal  $\mathfrak{a}$  into  $A$ , and let  $g$  be the canonical projection of  $A$  onto  $A/\mathfrak{a}$ . We may then identify  $A \widehat{\otimes}_A M$  with  $M$  by means of the map

$$x \widehat{\otimes}_A m \longmapsto xm.$$

Since the image of  $\mathfrak{a} \widehat{\otimes}_A M$  is closed in  $M$  (cf. 0.2.B), and the image of  $\mathfrak{a} \otimes_A M$  is dense in  $\mathfrak{a} \widehat{\otimes}_A M$ , the image of  $f \widehat{\otimes} M$  is precisely the closure  $\overline{\mathfrak{a}M}$  of  $\mathfrak{a}M$  in  $M$ . The exact sequence  $(*)$  therefore gives the isomorphism

$$(A/\mathfrak{a}) \widehat{\otimes}_A M \xrightarrow{\sim} M/\overline{\mathfrak{a}M}.$$

**0.3.3. Nakayama's Lemma.** *Let  $A$  be a pseudocompact ring,  $M$  a pseudocompact  $A$ -module, and  $\mathfrak{a}$  an ideal of  $A$  contained in the radical  $r(A)$ . Then*

$$\overline{\mathfrak{a}M} = M \implies M = 0.$$

*Proof.* Suppose that  $\overline{\mathfrak{a}M} = M$ .<sup>21</sup> Let  $M'$  be an open submodule of  $M$ , and put  $M'' = M/M'$ . Since  $M''$  is discrete,  $\mathfrak{a}M''$  is closed in  $M''$ , and hence equals  $\overline{\mathfrak{a}M''}$ . By 0.3.2, the canonical map

$$M/\overline{\mathfrak{a}M} \longrightarrow M''/\overline{\mathfrak{a}M''}$$

is surjective, so

$$\mathfrak{a}M'' = \overline{\mathfrak{a}M''} = M''.$$

Since  $M''$  is a finitely generated  $A$ -module and  $\mathfrak{a} \subset r(A)$ , the usual Nakayama lemma implies that  $M'' = 0$ . Consequently, every open submodule  $M'$  of  $M$  is equal to  $M$ , and therefore  $M = 0$ .<sup>22</sup>  $\square$

**0.3.4.** The usual consequences of Nakayama's lemma follow.

**Corollary.** *Let  $\mathfrak{a}$  be a closed ideal contained in  $r(A)$ , and let  $f : M \rightarrow N$  be a morphism of pseudocompact  $A$ -modules.*

<sup>21</sup>Editors' note: Replacing  $\mathfrak{a}$  by its closure if necessary, we may assume that  $\mathfrak{a}$  is closed.

<sup>22</sup>Editors' note: Indeed,  $M$  is the projective limit of the quotients  $M/M' = 0$ .

(i) The map  $f$  is surjective if the induced map

$$f' : M/\overline{\mathfrak{a}M} \longrightarrow N/\overline{\mathfrak{a}N}$$

is surjective.<sup>23</sup>

(ii) If  $N$  is projective, then  $f$  is an isomorphism whenever  $f'$  is an isomorphism.

*Proof.* Part (i) follows by applying Lemma 0.3.3 to  $\text{Coker}(f)$ . For (ii), suppose that  $f'$  is an isomorphism. By (i),  $f$  is then surjective, and hence has a section. Apply Lemma 0.3.3 to  $\text{Ker}(f)$ .  $\square$

When  $A$  is local with maximal ideal  $\mathfrak{m}$ , one may also deduce from 0.3.3 the following exchange theorem.

**Theorem.** *Let  $A$  be a local pseudocompact ring, let  $\mathfrak{m}$  be its maximal ideal, let  $M$  be a topologically free  $A$ -module with pseudobasis  $(m_i)_{i \in I}$  (0.2.1), and let  $N$  be a closed direct summand of  $M$ . There is a pseudobasis of  $M$  consisting of elements of  $N$  together with some of the  $m_i$ .*

*Proof.* This is clear when  $A$  is a field: use the duality of 0.2.2 and apply the usual exchange theorem. Consequently,

$$N/\overline{\mathfrak{m}N}$$

has a complementary topologically free  $A/\mathfrak{m}$ -module with pseudobasis  $(\overline{m}_i)_{i \in J}$ , where  $\overline{m}_i$  is the image of  $m_i$  in

$$M/\overline{\mathfrak{m}M},$$

and  $J$  is a subset of  $I$ .<sup>24</sup> If

$$P = \prod_{i \in J} Am_i,$$

then the canonical map

$$N \oplus P \longrightarrow M$$

is “bijective modulo  $\mathfrak{m}$ .” It is therefore bijective by what precedes. For another proof, see [CA], § IV.2, Proposition 8.  $\square$

**0.3.5.** Consider three pseudocompact  $A$ -modules  $L, M, N$ , where  $N$  has finite length. Give the  $A$ -module  $\text{Hom}_c(M, N)$  the discrete topology. Every element

$$\psi \in \text{Hom}_c(L, \text{Hom}_c(M, N))$$

defines a continuous bilinear map

$$\psi' : L \times M \longrightarrow N, \quad (\ell, m) \longmapsto \psi(\ell)(m).$$

This gives a natural isomorphism

$$\text{Hom}_c(L, \text{Hom}_c(M, N)) \xrightarrow{\sim} \text{Hom}_c(L \widehat{\otimes}_A M, N). \quad (1)$$

This is another characterization of  $\text{Hom}_c(L \widehat{\otimes}_A M, N)$ , which we now use when  $M$  is the projective limit of a filtered projective system  $(M_i)$  of pseudocompact  $A$ -modules. By (1)

<sup>23</sup>Editors’ note: Consequently, every pseudocompact  $A$ -module is a quotient of a topologically free  $A$ -module. One first reduces to the case in which  $A$  is local; cf. the proof of 0.3.7.

<sup>24</sup>Editors’ note: We have corrected  $N/\mathfrak{m}N$  to  $N/\overline{\mathfrak{m}N}$ , and likewise for  $M/\overline{\mathfrak{m}M}$  below.

and 0.2.F (ii), there are natural isomorphisms

$$\begin{aligned} \mathrm{Hom}_c \left( L \widehat{\otimes}_A \varprojlim_i M_i, N \right) &\simeq \mathrm{Hom}_c \left( L, \mathrm{Hom}_c(\varprojlim_i M_i, N) \right) \\ &\simeq \mathrm{Hom}_c \left( L, \varinjlim_i \mathrm{Hom}_c(M_i, N) \right). \end{aligned} \quad (2)$$

Moreover, because  $\varinjlim_i \mathrm{Hom}_c(M_i, N)$  is discrete, every continuous map from  $L$  into this module factors through a finite-length quotient of  $L$ . Hence the natural map

$$\varinjlim_i \mathrm{Hom}_c(L, \mathrm{Hom}_c(M_i, N)) \xrightarrow{\sim} \mathrm{Hom}_c \left( L, \varinjlim_i \mathrm{Hom}_c(M_i, N) \right) \quad (3)$$

is an isomorphism. Finally, (1) and 0.2.F (ii) again give natural isomorphisms

$$\begin{aligned} \varinjlim_i \mathrm{Hom}_c(L, \mathrm{Hom}_c(M_i, N)) &\simeq \varinjlim_i \mathrm{Hom}_c(L \widehat{\otimes}_A M_i, N) \\ &\simeq \mathrm{Hom}_c \left( \varprojlim_i (L \widehat{\otimes}_A M_i), N \right). \end{aligned} \quad (4)$$

Combining (2), (3), and (4) yields the following result.

**Proposition.** *Let  $(M_i)$  be a filtered projective system of objects of  $\mathrm{PC}(A)$ , let  $L \in \mathrm{PC}(A)$ , and let  $N \in \mathrm{LF}(A)$ . There is an isomorphism, functorial in  $N$ ,*

$$\mathrm{Hom}_c \left( L \widehat{\otimes}_A \varprojlim_i M_i, N \right) \simeq \mathrm{Hom}_c \left( \varprojlim_i (L \widehat{\otimes}_A M_i), N \right),$$

and consequently an isomorphism

$$L \widehat{\otimes}_A \varprojlim_i M_i \simeq \varprojlim_i (L \widehat{\otimes}_A M_i).$$

Thus the completed tensor product commutes with filtered projective limits.<sup>25</sup>

**0.3.6.** In particular,<sup>26</sup> the completed tensor product commutes with arbitrary products. For example, since  $A$  is the product of its local components  $A_{\mathfrak{m}}$  (0.1.1), every pseudocompact  $A$ -module

$$M \simeq A \widehat{\otimes}_A M$$

identifies with the product of the modules

$$M_{\mathfrak{m}} = A_{\mathfrak{m}} \widehat{\otimes}_A M,$$

which are called the local components of  $M$ .

Likewise, let  $M$  and  $N$  be pseudocompact  $A$ -modules. Recall from 0.2.2 that

$$M^{\dagger} = \mathrm{Hom}_c(M, A).$$

---

<sup>25</sup>This also recovers 0.3.1.1:

$$L = L \widehat{\otimes}_A A = L \widehat{\otimes}_A \varprojlim_{\mathfrak{l}} (A/\mathfrak{l}) \simeq \varprojlim_{\mathfrak{l}} L/\mathfrak{l}L,$$

where  $\mathfrak{l}$  ranges over the open ideals of  $A$ .

<sup>26</sup>Cf. editor's note (12).

Consider the map

$$\varphi : M^\dagger \otimes_A N^\dagger \longrightarrow (M \widehat{\otimes}_A N)^\dagger$$

which sends  $f \otimes g$  to the map

$$m \widehat{\otimes} n \longmapsto f(m)g(n)$$

from  $M \widehat{\otimes}_A N$  to  $A$ . The map  $\varphi$  is bijective when  $M \simeq A$ .

**Corollary.** *If  $A$  is Artinian, then  $\varphi$  is an isomorphism whenever  $M$  is topologically free, or more generally projective.*

Indeed, with  $N$  fixed, each of the functors

$$M \longmapsto (M \widehat{\otimes}_A N)^\dagger \quad \text{and} \quad M \longmapsto M^\dagger \otimes_A N^\dagger$$

takes arbitrary products to direct sums, by the preceding result and 0.2.F.

**Remark 0.3.6.A.**<sup>27</sup> *The same use of 0.2.F gives the following result. Let  $A$  be an Artinian ring, let  $M, Q \in \text{PC}(A)$ , and let  $N \in \text{LF}(A)$ . If  $Q$  is projective, then there are natural isomorphisms*

$$\text{Hom}_c(M, Q) \xrightarrow{\sim} \text{Hom}_A(Q^\dagger, M^\dagger), \quad Q^\dagger \otimes_A N \xrightarrow{\sim} \text{Hom}_c(Q, N).$$

**0.3.7.** For every  $\mathfrak{m} \in \Upsilon(A)$ , the functor

$$M \longmapsto A_{\mathfrak{m}} \widehat{\otimes}_A M$$

is plainly exact. Since every projective pseudocompact  $A$ -module  $P$  is a product of modules of the form  $A_{\mathfrak{m}}$ , it follows that

$$M \longmapsto P \widehat{\otimes}_A M$$

is exact whenever  $P$  is projective. The converse is also true.

**Proposition.** *Let  $A$  be a pseudocompact ring and  $P$  a pseudocompact  $A$ -module. The following conditions are equivalent:*

- (i)  $P$  is a projective object of  $\text{PC}(A)$ ;
- (ii) every local component  $P_{\mathfrak{m}}$  is a topologically free  $A_{\mathfrak{m}}$ -module;
- (iii) the functor  $M \mapsto P \widehat{\otimes}_A M$  is exact.

The equivalence of (i) and (ii) follows from 0.2.F (iii) and 0.2.1, and we have just proved (ii)  $\Rightarrow$  (iii). Suppose that the functor  $M \mapsto P \widehat{\otimes}_A M$  is exact. Since  $P \widehat{\otimes}_A M$  is the product of its local components,

$$(P \widehat{\otimes}_A M)_{\mathfrak{m}} \simeq P_{\mathfrak{m}} \widehat{\otimes}_{A_{\mathfrak{m}}} M_{\mathfrak{m}},$$

we reduce to the case in which  $A$  is local. We shall prove that  $P$  is then topologically free.

Let  $\mathfrak{m}$  be the maximal ideal of  $A$ . The quotient  $P/\overline{\mathfrak{m}}P$  is a linearly compact vector space over  $A/\mathfrak{m}$ , and hence a product of copies of  $A/\mathfrak{m}$  (0.2.2). Therefore there is a family  $(A_i)_{i \in I}$  of copies of  $A$  and an isomorphism

$$\phi : \prod_{i \in I} (A_i/\mathfrak{m}) \xrightarrow{\sim} P/\overline{\mathfrak{m}}P.$$

Since  $\prod_{i \in I} A_i$  is projective, there is a commutative square

---

<sup>27</sup>The editors added this remark, which will be useful in 2.3.1.

$$\begin{array}{ccc}
\prod_{i \in I} A_i & \xrightarrow{\psi} & P \\
p \downarrow & & \downarrow q \\
\prod_{i \in I} (A_i/\mathfrak{m}) & \xrightarrow[\sim]{\varphi} & P/\mathfrak{m}P
\end{array}$$

where  $p$  and  $q$  are the canonical projections. Apply Nakayama's lemma to  $\text{Coker } \psi$ . Since

$$(A/\mathfrak{m}) \widehat{\otimes}_A \psi$$

is precisely  $\phi$ , the map  $\psi$  is surjective.<sup>28</sup>

Put

$$B = \prod_{i \in I} A_i, \quad N = \ker \psi.$$

There is then the following commutative exact diagram:

$$\begin{array}{ccccccc}
& & & & 0 & & \\
& & & & \downarrow & & \\
& & \mathfrak{m} \widehat{\otimes}_A N & \longrightarrow & \mathfrak{m} \widehat{\otimes}_A B & \longrightarrow & \mathfrak{m} \widehat{\otimes}_A P \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & A \widehat{\otimes}_A N & \longrightarrow & A \widehat{\otimes}_A B & \xrightarrow{\psi} & A \widehat{\otimes}_A P \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
& & (A/\mathfrak{m}) \widehat{\otimes}_A N & \longrightarrow & (A/\mathfrak{m}) \widehat{\otimes}_A B & \xrightarrow{\varphi} & (A/\mathfrak{m}) \widehat{\otimes}_A P \longrightarrow 0
\end{array}$$

The snake lemma, applied to the first two rows, shows that in the bottom row the morphism

$$(A/\mathfrak{m}) \widehat{\otimes}_A N \longrightarrow (A/\mathfrak{m}) \widehat{\otimes}_A B$$

is a monomorphism. But  $\phi$  is an isomorphism, so  $(A/\mathfrak{m}) \widehat{\otimes}_A N = 0$ . Hence  $N = 0$  by 0.3.3, and  $\psi$  is an isomorphism.<sup>29</sup>

**0.3.8. Corollary.** *Let  $A$  be a complete Noetherian local ring and  $P$  a pseudocompact  $A$ -module. Then  $P$  is topologically free if and only if  $P$  is flat over  $A$ .*

Indeed, the canonical map

$$M \otimes_A P \longrightarrow M \widehat{\otimes}_A P$$

is bijective when  $M = A$ , and hence also when  $M$  is Noetherian: take a finite presentation of  $M$  and use the right exactness of both the tensor product and the completed tensor product.

Now  $P$  is flat if and only if

$$M \longmapsto M \otimes_A P$$

<sup>28</sup>This proves the result announced in editor's note (22): every pseudocompact  $A$ -module  $M$  is a quotient of a topologically free  $A$ -module. Since products are exact in  $\text{PC}(A)$ , one reduces to the local case treated here.

<sup>29</sup>A completely analogous proof, using the "nilpotent Nakayama lemma," shows that if  $A$  is Artinian and  $P$  is a flat  $A$ -module, then every local component of  $P$  is a free  $A$ -module (cf. [BAC], II, § 3.2, Corollary 2 to Proposition 5).

is exact on Noetherian modules. Likewise, the proof of Proposition 0.3.7 shows that  $P$  is topologically free provided the sequence

$$0 \longrightarrow \mathfrak{m} \hat{\otimes}_A P \longrightarrow A \hat{\otimes}_A P \longrightarrow (A/\mathfrak{m}) \hat{\otimes}_A P \longrightarrow 0$$

is exact. Thus  $P$  is topologically free if and only if the functor

$$M \longmapsto M \hat{\otimes}_A P$$

is exact on Noetherian modules. The corollary follows from the equality

$$M \otimes_A P = M \hat{\otimes}_A P$$

established above.

## 0.4. Profinite Algebras

Let  $k$  be a pseudocompact ring. A *topological  $k$ -algebra* is a commutative topological ring  $A$ , together with a morphism of topological rings  $k \rightarrow A$ . The algebra  $A$  is called a *profinite  $k$ -algebra* if its underlying topological  $k$ -module is pseudocompact.

Let  $\mathfrak{l}$  be an open  $k$ -submodule of such an  $A$ . The composite map

$$\varphi : A \times A \xrightarrow{\text{mult}} A \longrightarrow A/\mathfrak{l}$$

is continuous. Lemma 0.3.1 therefore gives an open  $k$ -submodule  $\mathfrak{n}$  of  $A$  such that  $\varphi(A \times \mathfrak{n}) = 0$ . Thus  $\mathfrak{l}$  contains the open ideal  $A\mathfrak{n}$ , and  $A$  is a pseudocompact ring.

Likewise, let  $M$  be a topological  $A$ -module whose underlying  $k$ -module is pseudocompact. If  $M'$  is an open  $k$ -submodule of  $M$ , applying Lemma 0.3.1 to

$$A \times M \xrightarrow{\text{mult}} M \longrightarrow M/M'$$

shows that  $M'$  contains an open  $A$ -submodule. Hence  $M$  is also a pseudocompact  $A$ -module.<sup>30</sup> Conversely:

**Lemma.** *Let  $A$  be a profinite  $k$ -algebra and  $M$  a pseudocompact  $A$ -module. The  $k$ -module  $M|_k$ , obtained by restriction of scalars, is pseudocompact.*

Indeed, every finite-length pseudocompact  $A$ -module is isomorphic to a quotient

$$A^n/L,$$

where  $L$  is an open submodule of  $A^n$ , and is therefore a pseudocompact  $k$ -module. Since  $M|_k$  is a projective limit of modules of this kind, it is a pseudocompact  $k$ -module.

**0.4.1.** If  $A$  and  $B$  are profinite  $k$ -algebras, a *morphism* from  $A$  to  $B$  is, by definition, a continuous homomorphism of  $k$ -algebras. We denote the category of profinite  $k$ -algebras by

$$\text{Alp}/k.$$

It plainly has projective limits: the algebra underlying a projective limit is the projective limit of the underlying algebras, endowed with the projective-limit topology.

It also has finite direct limits.<sup>31</sup> For example, if

$$f : A \longrightarrow B, \quad g : A \longrightarrow C$$

<sup>30</sup>The editors added the following lemma; cf. editor's note (36) in Proposition 0.5.

<sup>31</sup>In 0.4.2 we shall see that it has arbitrary direct limits.

are morphisms of profinite  $k$ -algebras, the amalgamated sum of  $B$  and  $C$  over  $A$  has

$$B\widehat{\otimes}_A C$$

as its underlying topological  $A$ -module. Here 0.4 makes  $B$  and  $C$  pseudocompact  $A$ -modules via  $f$  and  $g$ , and the multiplication is characterized by

$$(b\widehat{\otimes}c)(b'\widehat{\otimes}c') = bb'\widehat{\otimes}cc'$$

for  $b, b' \in B$  and  $c, c' \in C$ .

**Definition 0.4.2.** A profinite  $k$ -algebra  $C$  is said to have finite length if its underlying  $k$ -module has finite length, and is therefore discrete. We denote by  $\mathbf{Alf}/k$  the full subcategory of  $\mathbf{Alp}/k$  formed by the  $k$ -algebras of finite length.<sup>32</sup>

For every profinite  $k$ -algebra  $A$ , let  $h_A$  be the functor

$$h_A : \mathbf{Alf}/k \longrightarrow (\mathbf{Sets}), \quad C \longmapsto \mathrm{Hom}_{\mathbf{Alp}/k}(A, C).$$

The functor  $h_A$  is left exact,<sup>33</sup> and the canonical projections  $A \rightarrow A/\mathfrak{l}$ , where  $\mathfrak{l}$  ranges over the open ideals of  $A$ , induce for every  $C \in \mathbf{Alf}/k$  an isomorphism

$$\varinjlim_{\mathfrak{l}} \mathrm{Hom}_{\mathbf{Alf}/k}(A/\mathfrak{l}, C) \xrightarrow{\sim} \mathrm{Hom}_{\mathbf{Alp}/k}(A, C),$$

functorial in  $C$ . In other words,  $h_A$  is the direct limit of the representable functors  $h_{A/\mathfrak{l}}$ :

$$h_A \simeq \varinjlim_{\mathfrak{l}} h_{A/\mathfrak{l}}. \quad (*)$$

If  $B$  is another profinite  $k$ -algebra, the general properties of the bifunctor  $\mathrm{Hom}$ , together with

$$\mathrm{Hom}(h_C, h_B) = h_B(C)$$

for  $C$  of finite length, give isomorphisms

$$\begin{aligned} \mathrm{Hom}_{\mathbf{Alp}/k}(B, A) &\simeq \varprojlim_{\mathfrak{l}} \mathrm{Hom}_{\mathbf{Alp}/k}(B, A/\mathfrak{l}) \\ &\simeq \varprojlim_{\mathfrak{l}} \mathrm{Hom}(h_{A/\mathfrak{l}}, h_B) \\ &\simeq \mathrm{Hom}\left(\varinjlim_{\mathfrak{l}} h_{A/\mathfrak{l}}, h_B\right). \end{aligned}$$

Together with (\*), this proves that the contravariant functor  $A \mapsto h_A$  is fully faithful. In fact:

**Proposition.** The functor  $A \mapsto h_A$  is an anti-equivalence from  $\mathbf{Alp}/k$  onto the category of left-exact functors from  $\mathbf{Alf}/k$  to  $(\mathbf{Sets})$ .

By what precedes, it remains only to show that every left-exact functor

$$F : \mathbf{Alf}/k \longrightarrow (\mathbf{Sets})$$

is isomorphic to a functor  $h_A$ . The algebra  $A$  may be constructed as follows (cf. TDTE II, § 3).

<sup>32</sup>Caution:  $k$  is an object of  $\mathbf{Alf}/k$  only when  $k$  is Artinian; cf. 1.2.2 below.

<sup>33</sup>That is, it commutes with finite projective limits.



Because  $F$  is left exact, for every finite-length  $k$ -algebra  $C$  and every  $\xi \in F(C)$ , there is a smallest subalgebra  $C'$  of  $C$  such that  $\xi$  belongs to the image of  $F(C')$  in  $F(C)$ . If  $C' = C$ , the pair  $(C, \xi)$  is called *minimal*.

The minimal pairs form a category when a morphism from  $(C, \xi)$  to  $(D, \eta)$  is defined to be a homomorphism  $\phi : C \rightarrow D$  such that

$$(F\phi)(\xi) = \eta.$$

Such a  $\phi$  is necessarily surjective, and the category of minimal pairs is left filtered. Moreover, one may restrict to pairs  $(C, \xi)$  for which  $C$  belongs to a set containing one finite-length  $k$ -algebra of every isomorphism type.<sup>34</sup> The functor  $(C, \xi) \mapsto C$ , from the category of minimal pairs to the category of profinite  $k$ -algebras, therefore has a projective limit; take  $A$  to be this limit.

**Corollary.** *The category  $\text{Alp}/k$  has arbitrary direct limits.*

Indeed, the category of left-exact functors from  $\text{Alf}/k$  to **(Sets)** has projective limits, computed pointwise: if  $(F_i)$  is a projective system of such functors, then for every  $C \in \text{Alf}/k$ ,

$$(\varprojlim_i F_i)(C) = \varprojlim_i F_i(C).$$

35

## 0.5. Base Change

Let

$$\phi : k \longrightarrow \ell$$

be a homomorphism of pseudocompact rings (0.1.3). The construction of 0.3 admits the following generalization.<sup>36</sup>

**Definition 0.5.A.** *For every  $M \in \text{PC}(k)$  and  $N \in \text{PC}(\ell)$ , let*

$$M \widehat{\otimes}_k N$$

*be the separated completion of  $M \otimes_k N$  for the linear topology defined by the kernels of the projections*

$$M \otimes_k N \longrightarrow (M/M') \otimes_k (N/N'),$$

*where  $M'$ , respectively  $N'$ , ranges over the open submodules of  $M$ , respectively  $N$ . Then  $M \widehat{\otimes}_k N$  is a pseudocompact  $\ell$ -module. If  $m \in M$  and  $x \in N$ , the canonical image of  $m \otimes_k x$  in  $M \widehat{\otimes}_k N$  is denoted by*

$$m \widehat{\otimes}_k x.$$

This applies in particular to  $N = \ell$ . In that case

$$M \widehat{\otimes}_k \ell$$

<sup>34</sup>Indeed, every discrete  $k$ -module of length  $n$  is isomorphic to  $k^n/L$ , where  $L$  is an open submodule of  $k^n$  of colength  $n$ . One may then consider the set of isomorphism classes of profinite  $k$ -algebra structures on each such module.

<sup>35</sup>If  $(A_i)_{i \in I}$  is a direct system in  $\text{Alp}/k$ , its direct limit in  $\text{Alp}/k$  is the separated completion of the  $k$ -algebra  $B$ , the direct limit of the underlying  $k$ -algebras, for the topology defined by the ideals  $I$  such that  $I \cap A_i$  is open in  $A_i$  for every  $i$  and  $\text{length}_k(B/I) < \infty$ . Notice, for example, that if  $K/k$  is an algebraic extension of fields of infinite degree and  $(k_i)$  is the filtered direct system of finite-degree subextensions, then the direct limit of the system  $(k_i)$  in  $\text{Alp}/k$  is the zero ring.

<sup>36</sup>The editors expanded the details of this paragraph.

is called the pseudocompact  $\ell$ -module obtained from  $M$  by the base change  $k \rightarrow \ell$ .

**Remarks 0.5.B.**

(i) In such a base change,  $\ell$  need not be a profinite  $k$ -algebra. The typical example is that in which  $k$  is a field and  $\ell = K$  is an arbitrary field extension of  $k$ .

(ii) If, however, the  $k$ -module underlying  $N$  is pseudocompact—for example, if  $\ell$  is a profinite  $k$ -algebra—then 0.4 shows that every open  $k$ -submodule of  $N$  contains an open  $\ell$ -submodule. In that case  $M \widehat{\otimes}_k N$  agrees with the completed tensor product of 0.3 for the pseudocompact  $k$ -modules  $M$  and  $N$ , so the notation is unambiguous.

In every case, the  $k$ -module  $N|_k$  obtained by restriction of scalars is a topological  $k$ -module: the map

$$k \times N \longrightarrow N, \quad (t, n) \longmapsto \phi(t)n$$

is continuous. We write  $\text{Hom}_c(M, N|_k)$  for the abelian group of continuous  $k$ -linear maps from  $M$  to  $N|_k$ .

**Proposition 0.5.C.** *For every  $M \in \text{PC}(k)$  and  $N \in \text{PC}(\ell)$ , there is a functorial isomorphism*

$$\text{Hom}_{\text{PC}(\ell)}(M \widehat{\otimes}_k \ell, N) \simeq \text{Hom}_c(M, N|_k).$$

37

Indeed, let  $u : M \rightarrow N|_k$  be a continuous homomorphism. The map

$$u' : M \times \ell \longrightarrow N, \quad (m, \lambda) \longmapsto \lambda u(m)$$

is continuous and “bilinear,” meaning  $k$ -linear in the first factor and  $\ell$ -linear in the second. If  $N'$  is an open  $\ell$ -submodule of  $N$ , the argument of Lemma 0.3.1 gives an open submodule  $M' \subset M$  and an open ideal  $\ell' \subset \ell$  such that

$$u'(M' \times \ell) \subset N', \quad u'(M \times \ell') \subset N'.$$

Thus  $u$  induces a continuous  $\ell$ -linear map

$$U : M \widehat{\otimes}_k \ell \longrightarrow N$$

satisfying

$$U(m \widehat{\otimes}_k \lambda) = \lambda u(m).$$

Conversely, every morphism

$$f : M \widehat{\otimes}_k \ell \longrightarrow N$$

gives the morphism

$$f' : M \longrightarrow N|_k, \quad m \longmapsto f(m \widehat{\otimes}_k 1).$$

As in 0.3.2, 0.3.5, and 0.3.1.2, this yields:

**Corollary 0.5.D.** *The functor*

$$\text{PC}(k) \longrightarrow \text{PC}(\ell), \quad M \longmapsto M \widehat{\otimes}_k \ell$$

*is right exact and commutes with filtered projective limits. Thus, if  $(M_i)$  is a filtered projective system in  $\text{PC}(k)$ , there is a canonical isomorphism*

$$(\varprojlim_i M_i) \widehat{\otimes}_k \ell \simeq \varprojlim_i (M_i \widehat{\otimes}_k \ell).$$

---

<sup>37</sup>Consequently, when  $\ell$  is a profinite  $k$ -algebra, the functor  $M \mapsto M \widehat{\otimes}_k \ell$  is left adjoint to the restriction-of-scalars functor.

Moreover, if  $M$  and  $N$  are pseudocompact  $k$ -modules, there is a canonical isomorphism

$$(M \widehat{\otimes}_k N) \widehat{\otimes}_k \ell \simeq (M \widehat{\otimes}_k \ell) \widehat{\otimes}_\ell (N \widehat{\otimes}_k \ell).$$

**Definition 0.5.E.** Finally, if  $A$  is a profinite  $k$ -algebra, there is a unique profinite  $\ell$ -algebra structure on  $A \widehat{\otimes}_k \ell$  such that

$$(a \widehat{\otimes}_k \lambda)(b \widehat{\otimes}_k \mu) = ab \widehat{\otimes}_k (\lambda \mu)$$

for  $a, b \in A$  and  $\lambda, \mu \in \ell$ . The profinite  $\ell$ -algebra  $A \widehat{\otimes}_k \ell$  is said to be obtained from  $A$  by extension of scalars, or base change, along  $k \rightarrow \ell$ .

## 1. Formal Varieties over a Pseudocompact Ring

**1.1.** To every pseudocompact ring  $A$  one may associate a formal scheme (EGA I, 10.4.2) as follows. Its underlying topological space is the set  $\Upsilon(A)$  of open prime ideals of  $A$ , which are therefore maximal, endowed with the discrete topology. For every subset  $E \subset \Upsilon(A)$ , the space of sections of the structure sheaf over  $E$  is the Cartesian product

$$\prod_{\mathfrak{m} \in E} A_{\mathfrak{m}}.$$

The formal scheme so obtained is denoted by  $\mathrm{Spf}(A)$ , the *formal spectrum* of  $A$ .<sup>38</sup>

If  $A$  and  $B$  are two pseudocompact rings, a *morphism* from  $\mathrm{Spf}(B)$  to  $\mathrm{Spf}(A)$  consists of a map

$$f : \Upsilon(B) \longrightarrow \Upsilon(A)$$

and a family of continuous homomorphisms

$$f_y^\sharp : A_{f(y)} \longrightarrow B_y, \quad y \in \Upsilon(B).$$

Such a morphism induces a continuous homomorphism

$$f^\sharp : A = \prod_{x \in \Upsilon(A)} A_x \longrightarrow B = \prod_{y \in \Upsilon(B)} B_y.$$

The converse is true:

**Proposition.** *The contravariant functor  $A \mapsto \mathrm{Spf}(A)$  is fully faithful.*

Indeed, let  $\phi : A \rightarrow B$  be a continuous homomorphism. The inverse image  $\phi^{-1}(\mathfrak{n})$  of an open maximal ideal of  $B$  is an open prime ideal of  $A$ , and hence is maximal in  $A$ . We therefore obtain a map

$$\mathfrak{n} \longmapsto \phi^{-1}(\mathfrak{n})$$

from  $\Upsilon(B)$  to  $\Upsilon(A)$ , and  $\phi$  induces a continuous homomorphism

$$A_{\phi^{-1}(\mathfrak{n})} \longrightarrow B_{\mathfrak{n}}.$$

Thus  $\phi$  induces a morphism of formal schemes

$$\mathrm{Spf}(\phi) : \mathrm{Spf}(B) \longrightarrow \mathrm{Spf}(A).$$

---

<sup>38</sup>Cf. Remarks 0.1.2.

It is readily verified that

$$\mathrm{Spf}(\phi)^{\natural} = \phi, \quad \mathrm{Spf}(f^{\natural}) = f$$

for every morphism  $f : \mathrm{Spf}(B) \rightarrow \mathrm{Spf}(A)$ , which proves the proposition.

Although we shall speak here only of formal schemes of the form  $\mathrm{Spf}(A)$ , we shall use the language of formal schemes rather than that of pseudocompact rings, so as to ground our assertions in geometric intuition.

**1.2.** Let  $k$  be a pseudocompact ring.

**Definition 1.2.A.**<sup>39</sup> A *formal variety over  $k$*  is a formal scheme  $X$  over  $\mathrm{Spf}(k)$  which is isomorphic to a formal  $k$ -scheme  $\mathrm{Spf}(A)$  for some profinite  $k$ -algebra  $A$ . The algebra  $A$  is then isomorphic to the *affine algebra* of  $X$ , that is, the algebra of sections of the structure sheaf  $\mathcal{O}_X$  of  $X$ .

We denote by  $\mathrm{Vaf}/k$  the full subcategory of the category of formal schemes over  $\mathrm{Spf}(k)$  whose objects are the formal  $k$ -varieties.<sup>40</sup>

By 1.1, the functor  $A \mapsto \mathrm{Spf}(A)$  is an anti-equivalence from  $\mathrm{Alp}/k$  (0.4.1) onto  $\mathrm{Vaf}/k$ . Hence, by the corollary of 0.4.2:

**Proposition 1.2.B.** *The category  $\mathrm{Vaf}/k$  has projective and direct limits.*<sup>41</sup>

For example, let  $X \in \mathrm{Ob}(\mathrm{Vaf}/k)$ , let

$$f : Y \longrightarrow X, \quad g : Z \longrightarrow X$$

be two formal  $k$ -varieties over  $X$ , and let  $A, B, C$  be the affine algebras of  $X, Y, Z$ , respectively. By 0.4.1, the affine algebra of the fiber product  $Y \times_X Z$  identifies with  $B \widehat{\otimes}_A C$ ,<sup>42</sup> Consequently, the inclusion of  $\mathrm{Vaf}/k$  into the category of all formal  $k$ -schemes commutes with finite projective limits (cf. EGA I, 10.7).

Direct limits of formal  $k$ -varieties correspond to projective limits of their affine algebras.

**Example 1.2.C (Cokernels).** Let  $d, e : X \rightrightarrows Y$  be a double arrow in  $\mathrm{Vaf}/k$ . The affine algebra of  $\mathrm{Coker}(d, e)$  is isomorphic to the kernel of the homomorphisms induced on the affine algebras of  $X$  and  $Y$ . There is also the following construction of  $\mathrm{Coker}(d, e)$ . Its underlying topological space is the cokernel of the underlying spaces.<sup>43</sup> Let  $p$  be the canonical projection from the set underlying  $Y$  to this cokernel, and let  $z$  be a point of the cokernel. The local algebra of  $\mathrm{Coker}(d, e)$  at  $z$  is the kernel of the double arrow

$$d^{\natural}, e^{\natural} : \prod_{p(y)=z} \mathcal{O}_{Y,y} \rightrightarrows \prod_{q(x)=z} \mathcal{O}_{X,x}$$

where

$$q = p \circ d = p \circ e,$$

and  $d^{\natural}$  and  $e^{\natural}$  are induced by the homomorphisms  $d_x^{\natural}$  and  $e_x^{\natural}$ , in the notation of 1.1.

**Definition 1.2.D.** Let  $\phi : k \rightarrow \ell$  be a homomorphism of pseudocompact rings and let  $X$  be a formal  $k$ -variety with affine algebra  $A$ . The formal scheme

$$X \times_{\mathrm{Spf}(k)} \mathrm{Spf}(\ell),$$

<sup>39</sup>The editors added the numbering 1.2.A–1.2.E for later reference.

<sup>40</sup>By the definition of morphisms in 1.1, every  $X \in \mathrm{Ob}(\mathrm{Vaf}/k)$  is the direct sum in  $\mathrm{Vaf}/k$  of the one-point formal varieties  $\mathrm{Spf}(\mathcal{O}_{X,x})$ , for  $x \in X$ .

<sup>41</sup>In particular, cokernels exist in  $\mathrm{Vaf}/k$ ; see below. Moreover, a filtered projective limit in  $\mathrm{Vaf}/k$ , all of whose transition morphisms are surjective, can be empty; cf. editor's note (34).

<sup>42</sup>The formal variety  $Y \times_X Z$  is the direct sum, over  $x \in X$ , of the formal varieties  $B_x \widehat{\otimes}_{\mathcal{O}_{X,x}} C_x$ , where  $B_x$  is the product of the  $\mathcal{O}_{Y,y}$  for  $y \in Y$  such that  $f(y) = x$ , and  $C_x$  is defined analogously. This will be used in 2.5.1.

<sup>43</sup>That is, it is the quotient set of  $Y$  by the identifications  $d(x) = e(x)$ , for every  $x \in X$ , endowed with the quotient topology, which here is discrete.

obtained by base change, is a formal  $\ell$ -variety. We also denote it by

$$X \widehat{\otimes}_k \ell.$$

Its affine algebra is the completed tensor product  $A \widehat{\otimes}_k \ell$  (cf. 0.5 and EGA I, § 10).

**Remark 1.2.E.** Since every formal variety over  $k$  decomposes into formal varieties over the local components of  $k$ , *in certain proofs we shall assume that  $k$  is a local pseudocompact ring.*

We now give some examples while fixing our terminology.

**1.2.1.** A  $k$ -functor is, by definition, a covariant functor from  $\mathbf{Alf}/k$  to **(Sets)**. By 1.1 and 0.4.2, we may identify

$$\mathbf{Vaf}/k \simeq (\mathbf{Alp}/k)^\circ$$

with a full subcategory of the category of  $k$ -functors by associating to every formal  $k$ -variety  $X$  the functor

$$\mathbf{Alf}/k \longrightarrow (\mathbf{Sets}), \quad C \longmapsto X(C) = \mathrm{Hom}_{\mathbf{Vaf}/k}(\mathrm{Spf}(C), X).$$

We shall later encounter  $k$ -functors  $\mathbf{F}$  which associate to every object  $C$  of  $\mathbf{Alf}/k$  a  $C$ -module  $\mathbf{F}(C)$ , and to every morphism  $\phi : C \rightarrow D$  in  $\mathbf{Alf}/k$  a  $k$ -linear map

$$\mathbf{F}(\phi) : \mathbf{F}(C) \longrightarrow \mathbf{F}(D)$$

such that, for  $x \in \mathbf{F}(C)$  and  $\lambda \in C$ ,

$$\mathbf{F}(\phi)(\lambda x) = \phi(\lambda) \mathbf{F}(\phi)(x).$$

By Exposé I, 3.1, such an  $\mathbf{F}$  is endowed with the structure of an  $\mathbf{O}_k$ -module, where  $\mathbf{O}_k$  denotes the  $k$ -functor in rings which assigns to every object  $C$  of  $\mathbf{Alf}/k$  its underlying ring.

**Definitions.**

(i) An  $\mathbf{O}_k$ -module  $\mathbf{F}$  is called *admissible* if every morphism  $\phi : C \rightarrow D$  in  $\mathbf{Alf}/k$  induces a bijection from  $D \otimes_C \mathbf{F}(C)$  onto  $\mathbf{F}(D)$ .<sup>44</sup>

(ii) The module  $\mathbf{F}$  is called *flat* if it is admissible and if, for every object  $C$  of  $\mathbf{Alf}/k$ , the  $C$ -module  $\mathbf{F}(C)$  is flat.<sup>45</sup>

For example, if  $M$  is a  $k$ -module, not necessarily pseudocompact, we denote by  $\mathbf{W}(M)$ , as in Exposé I, 4.6.1, the  $\mathbf{O}_k$ -module

$$C \longmapsto C \otimes_k M.$$

Then  $\mathbf{W}(M)$  is flat whenever  $M$  is flat over  $k$ .

Moreover, the functor  $M \mapsto \mathbf{W}(M)$  has as right adjoint the functor which assigns to every  $\mathbf{O}_k$ -module  $\mathbf{F}$  the  $k$ -module

$$\varprojlim_{\mathfrak{l}} \mathbf{F}(k/\mathfrak{l}),$$

where  $\mathfrak{l}$  ranges over the open ideals of  $k$ .

**1.2.2.** In what follows, an  $\mathbf{O}_k$ -module will always be denoted by a boldface letter such as  $\mathbf{F}$ . When  $k$  is Artinian, we shall simply write  $F$  instead of  $\mathbf{F}(k)$ . In that case, it is clear that the functor

$$\mathbf{F} \longmapsto F$$

<sup>44</sup>The editors introduced this terminology; cf. editor's note (47) in 1.2.3.

<sup>45</sup>It is therefore projective, since  $C$  is Artinian.

induces an equivalence from the category of flat  $\mathbf{O}_k$ -modules onto the category of flat  $k$ -modules! The purpose of this terminology is solely to allow us to reason “as if  $k$  were always Artinian.”

In accordance with Exposé I, § 3.1, we shall use analogous conventions for other algebraic structures. Thus an  $\mathbf{O}_k$ -algebra, respectively an  $\mathbf{O}_k$ -coalgebra, an  $\mathbf{O}_k$ -Lie algebra, or an  $\mathbf{O}_k$ - $p$ -Lie algebra, consists of an  $\mathbf{O}_k$ -module  $\mathbf{M}$  together with, for every object  $C$  of  $\mathbf{Alf}/k$ , a structure of  $C$ -algebra, respectively  $C$ -coalgebra,  $C$ -Lie algebra, or  $C$ - $p$ -Lie algebra, on  $\mathbf{M}(C)$ . We further require that for every morphism  $\phi : C \rightarrow D$  in  $\mathbf{Alf}/k$ , the map

$$D \otimes_C \mathbf{M}(C) \longrightarrow \mathbf{M}(D)$$

induced by  $\mathbf{M}(\phi)$  be a homomorphism of  $D$ -algebras, respectively of  $D$ -coalgebras, and so on.

Finally, if  $\mathbf{F}$  and  $\mathbf{G}$  are two  $\mathbf{O}_k$ -modules, then  $\mathbf{F} \otimes_k \mathbf{G}$  denotes the  $\mathbf{O}_k$ -module

$$C \longmapsto \mathbf{F}(C) \otimes_C \mathbf{G}(C).$$

**1.2.3.**<sup>46</sup> We begin with the following lemma.

**Lemma 1.2.3.A.** *Let  $k \rightarrow K$  be a morphism of pseudocompact rings, let  $B$  be a pseudocompact  $K$ -module, and let  $M$  be a projective  $k$ -module, without topology. There is a canonical isomorphism of pseudocompact  $K$ -modules*

$$\theta : \mathrm{Hom}_k(M, k) \widehat{\otimes}_k B \xrightarrow{\sim} \mathrm{Hom}_k(M, B). \quad (2)$$

Here  $M^* = \mathrm{Hom}_k(M, k)$  is endowed with the topology defined in 0.2.2, which makes it a pseudocompact  $k$ -module. The topology on  $\mathrm{Hom}_k(M, B)$  is defined analogously: a basis of neighborhoods of 0 is formed by the following  $K$ -submodules, where  $x \in M$  and  $B'$  is an open  $K$ -submodule of  $B$ :

$$\mathcal{H}(x, B') = \{f \in \mathrm{Hom}_k(M, B) \mid f(x) \in B'\}.$$

Finally,  $M^* \widehat{\otimes}_k B$  is the pseudocompact  $K$ -module obtained from  $M^*$  by base change; cf. 0.5.A.

With these conventions,  $\theta$  is plainly an isomorphism when  $M = k$ . Moreover, both sides of (2), regarded as functors in  $M$ , carry direct sums to products; in particular, they commute with finite direct sums. It follows that (2) is an isomorphism first when  $M$  is a free  $k$ -module, and then when  $M$  is projective.

**Definition 1.2.3.B.** Let  $N$  be a pseudocompact  $k$ -module. We write  $\mathbf{V}_k^f(N)$ , or  $\mathbf{N}^\dagger$ ,<sup>47</sup> for the  $\mathbf{O}_k$ -module which assigns to every  $C \in \mathbf{Ob}(\mathbf{Alf}/k)$  the  $C$ -module

$$(C \widehat{\otimes}_k N)^\dagger,$$

the dual of  $C \widehat{\otimes}_k N$  in the sense of 0.2.2; that is, the  $C$ -module

$$\mathrm{Hom}_c(C \widehat{\otimes}_k N, C) \simeq \mathrm{Hom}_c(N, C|_k),$$

where  $C|_k$  denotes the  $k$ -module obtained from  $C$  by restriction of scalars. This  $\mathbf{O}_k$ -module  $\mathbf{V}_k^f(N)$  is called the *dual* of  $N$ .

<sup>46</sup>The editors added the following lemma and introduced the numbering 1.2.3.A through 1.2.3.F for later references.

<sup>47</sup>In fact, the second notation will not be used; see editors' note (47).

If  $k_{\mathfrak{m}}$  is a local component of  $k$ , then for every object  $C$  of  $\mathbf{Alf}/k$ ,

$$\mathrm{Hom}_c(k_{\mathfrak{m}}, C|_k) = C_{\mathfrak{m}} = C \otimes_k k_{\mathfrak{m}}$$

is a projective  $C$ -module. Moreover,

$$D \otimes_C C_{\mathfrak{m}} = D_{\mathfrak{m}}$$

for every morphism  $C \rightarrow D$  in  $\mathbf{Alf}/k$ . Thus  $\mathbf{V}_k^f(k_{\mathfrak{m}})$  is a flat  $\mathbf{O}_k$ -module; cf. 1.2.1. Since every projective pseudocompact  $k$ -module is a product of modules  $k_{\mathfrak{m}}$  (Proposition 0.2.1), Corollary 0.2.F gives:

**Corollary 1.2.3.C.** *The  $\mathbf{O}_k$ -module  $\mathbf{V}_k^f(N)$  is flat whenever  $N$  is a projective pseudocompact  $k$ -module.*

**Definition 1.2.3.D.** Conversely, let  $\mathbf{M}$  be an admissible  $\mathbf{O}_k$ -module (cf. 1.2.1).<sup>48</sup> We call the following pseudocompact  $k$ -module the *dual* of  $\mathbf{M}$ , and denote it by  $\Gamma^*(\mathbf{M})$ . As  $\mathfrak{l}$  ranges over the open ideals of  $k$ , endow each  $k/\mathfrak{l}$ -module

$$\mathbf{M}(k/\mathfrak{l})^* = \mathrm{Hom}_{k/\mathfrak{l}}(\mathbf{M}(k/\mathfrak{l}), k/\mathfrak{l})$$

with the topology described in 0.2.2; it is then a pseudocompact  $k$ -module. Since

$$\mathbf{M}(k/\mathfrak{l}) \simeq \mathbf{M}(k/\mathfrak{l}') \otimes_{k/\mathfrak{l}'} (k/\mathfrak{l}) \quad (\mathfrak{l} \supset \mathfrak{l}'),$$

there are transition morphisms

$$\begin{aligned} \mathrm{Hom}_{k/\mathfrak{l}'}(\mathbf{M}(k/\mathfrak{l}'), k/\mathfrak{l}') &\longrightarrow \mathrm{Hom}_{k/\mathfrak{l}'}(\mathbf{M}(k/\mathfrak{l}'), k/\mathfrak{l}) \\ &= \mathrm{Hom}_{k/\mathfrak{l}}(\mathbf{M}(k/\mathfrak{l}), k/\mathfrak{l}). \end{aligned}$$

By definition,  $\Gamma^*(\mathbf{M})$  is the projective limit of this filtered projective system of pseudocompact  $k$ -modules.

From now on, suppose in addition that  $\mathbf{M}$  is a flat  $\mathbf{O}_k$ -module (cf. 1.2.1). Each  $\mathbf{M}(k/\mathfrak{l}')$  is then a projective  $k/\mathfrak{l}'$ -module, so the transition morphisms above are surjective. Consequently the projections

$$\Gamma^*(\mathbf{M}) \longrightarrow \mathbf{M}(k/\mathfrak{l})^*$$

are also surjective, since filtered projective limits are exact in  $\mathrm{PC}(k)$ .

Let  $\tau : k \rightarrow K$  be a morphism of pseudocompact rings. We write  $\mathbf{M} \otimes_k K$ , or simply  $\mathbf{M}_K$ , for the  $\mathbf{O}_K$ -functor defined as follows. If  $C$  is a finite-length  $K$ -algebra, then the kernel of  $k \rightarrow C$  is an open ideal  $\mathfrak{l}$ , and we set

$$\mathbf{M}_K(C) = \mathbf{M}(k/\mathfrak{l}) \otimes_k C.$$

The same value is obtained from  $\mathbf{M}(k/\mathfrak{l}') \otimes_k C$  for every open ideal  $\mathfrak{l}' \subset \mathfrak{l}$ . We then define  $\Gamma_K^*(\mathbf{M}_K)$  as the projective limit, with  $I$  ranging over the open ideals of  $K$ , of the pseudocompact  $K$ -modules

$$\mathrm{Hom}_{K/I}(\mathbf{M}_K(K/I), K/I) = \mathrm{Hom}_{k/\mathfrak{l}}(\mathbf{M}(k/\mathfrak{l}), K/I),$$

where on the right  $\mathfrak{l}$  is any open ideal of  $k$  such that  $\tau(\mathfrak{l}) \subset I$ .

<sup>48</sup>The editors do not understand the definition that follows unless  $\mathbf{M}$  is assumed to be admissible, which is why this hypothesis has been added. They also write  $\Gamma^*(\mathbf{M})$ , rather than  $\mathbf{M}^*$ , to avoid confusion between  $\mathbf{M}^*$ , the module dual to the functor  $\mathbf{M}$ , and  $\mathbf{N}^\dagger$ , the functor dual to the module  $N$ ; cf. editors' note (46). Finally, the editors have spelled out the construction of  $\Gamma^*(\mathbf{M})$ .

By Lemma 1.2.3.A, the right-hand term identifies with

$$\mathbf{M}(k/\mathfrak{l})^* \widehat{\otimes}_k (K/I).$$

Since the projections  $\Gamma^*(\mathbf{M}) \longrightarrow \mathbf{M}(k/\mathfrak{l})^*$  are surjective, the projective limit of these latter modules is precisely the pseudocompact  $K$ -module

$$\Gamma^*(\mathbf{M}) \widehat{\otimes}_k K$$

(cf. 0.5.A). We have therefore proved that, for every flat  $\mathbf{O}_k$ -module  $\mathbf{M}$ , formation of  $\Gamma^*(\mathbf{M})$  commutes with extension of the base; that is,

$$\Gamma_K^*(\mathbf{M} \otimes_k K) \simeq \Gamma^*(\mathbf{M}) \widehat{\otimes}_k K. \quad (\star)$$

**Proposition 1.2.3.E.**

(i) *The functors*

$$N \longmapsto \mathbf{V}_k^f(N), \quad \mathbf{M} \longmapsto \Gamma^*(\mathbf{M})$$

*define an anti-equivalence between the category of projective pseudocompact  $k$ -modules and the category of flat  $\mathbf{O}_k$ -modules.*<sup>49</sup>

(ii) *Moreover, if  $k \longrightarrow K$  is a morphism of pseudocompact rings, the preceding anti-equivalence “commutes with base change” in the following sense: if  $N \simeq \Gamma^*(\mathbf{M})$ , then*

$$N \widehat{\otimes}_k K \simeq \Gamma_K^*(\mathbf{M} \otimes_k K).$$

*Proof.* First, there is a natural transformation

$$\phi_N : N \longrightarrow \Gamma^*(\mathbf{V}_k^f(N)).$$

The functor  $\Gamma^* \circ \mathbf{V}_k^f$  commutes with products by 0.3.5 and 0.2.F, so it is enough to verify that  $\phi_N$  is an isomorphism when  $N$  is a local component  $k_{\mathfrak{m}}$  of  $k$ . In this case, for every open ideal  $\mathfrak{l}$  of  $k$  contained in  $\mathfrak{m}$ , the natural morphism

$$(k/\mathfrak{l})_{\mathfrak{m}} \longrightarrow \mathrm{Hom}_{k/\mathfrak{l}}(\mathrm{Hom}_c(k_{\mathfrak{m}} \widehat{\otimes}_k k/\mathfrak{l}, k/\mathfrak{l}), k/\mathfrak{l})$$

is an isomorphism, which proves the assertion.

Conversely, let  $\mathbf{M}$  be a flat  $\mathbf{O}_k$ -module. We show that

$$\Gamma^*(\mathbf{M}) = \varprojlim_{\mathfrak{l}} \mathbf{M}(k/\mathfrak{l})^*$$

is a projective object of  $\mathrm{PC}(k)$ . Let  $N \longrightarrow N'$  be a surjective morphism between objects of  $\mathrm{LF}(k)$ . By 0.2.F (i) and (ii), it is enough to prove that the natural map

$$\varprojlim_{\mathfrak{l}} \mathrm{Hom}_c(\mathbf{M}(k/\mathfrak{l})^*, N) \longrightarrow \varprojlim_{\mathfrak{l}} \mathrm{Hom}_c(\mathbf{M}(k/\mathfrak{l})^*, N')$$

is surjective. This is clear:  $N$  and  $N'$  are  $k/\mathfrak{l}_0$ -modules for some open ideal  $\mathfrak{l}_0$ , and if  $\mathfrak{l} \subset \mathfrak{l}_0$ , every morphism  $\mathbf{M}(k/\mathfrak{l})^* \longrightarrow N'$  lifts to a morphism  $\mathbf{M}(k/\mathfrak{l})^* \longrightarrow N$ , since  $\mathbf{M}(k/\mathfrak{l})^*$  is a projective object of  $\mathrm{PC}(k/\mathfrak{l})$ .

<sup>49</sup>The editors added assertion (ii) below and the proof that follows. The original said: “If  $\mathbf{M}$  is a flat  $\mathbf{O}_k$ -module, it is clear that  $\mathbf{M}$  is the dual of  $\mathbf{M}^*$ ; hence the functors  $N \mapsto N^*$  and  $\mathbf{M} \mapsto \mathbf{M}^*$  define an anti-equivalence ...”



There is a morphism

$$\psi : \mathbf{M} \longrightarrow \mathbf{V}_k^f(\Gamma^*(\mathbf{M}))$$

of functors from  $\mathbf{Alf}/k$  to  $(\mathbf{Sets})$ . Let  $B$  be an object of  $\mathbf{Alf}/k$ . We claim that

$$\begin{aligned} \psi(B) : \mathbf{M}(B) &\longrightarrow \mathbf{V}_k^f(\Gamma^*(\mathbf{M}))(B) \\ &= \mathrm{Hom}_c \left( \varprojlim_{\mathfrak{l}} (\mathbf{M}(k/\mathfrak{l})^* \widehat{\otimes}_k B), B \right) \end{aligned} \quad (1)$$

is a bijection. In the equality above we used the fact that  $\widehat{\otimes}$  commutes with filtered projective limits. Let  $\mathfrak{l}_0$  be an open ideal of  $k$  contained in the kernel of  $k \rightarrow B$ . By Lemma 1.2.3.A, for every  $\mathfrak{l} \subset \mathfrak{l}_0$ , there is a canonical isomorphism of pseudocompact  $B$ -modules

$$\mathbf{M}(k/\mathfrak{l})^* \widehat{\otimes}_k B \xrightarrow{\sim} \mathrm{Hom}_{k/\mathfrak{l}}(\mathbf{M}(k/\mathfrak{l}), B).$$

Since

$$\mathbf{M}(B) = \mathbf{M}(k/\mathfrak{l}) \otimes_{k/\mathfrak{l}} B,$$

the right-hand side is  $\mathrm{Hom}_B(\mathbf{M}(B), B)$ . Thus the projective system in (1) is constant for  $\mathfrak{l} \subset \mathfrak{l}_0$ , and (1) reduces to the canonical morphism

$$\mathbf{M}(B) \longrightarrow \mathrm{Hom}_c(\mathrm{Hom}_B(\mathbf{M}(B), B), B).$$

This is an isomorphism by 0.2.2, since  $B$  is Artinian and  $\mathbf{M}(B)$  is a projective  $B$ -module. This proves assertion (i) of the proposition; assertion (ii) follows from the isomorphism  $(\star)$  established above.  $\square$

**Remark 1.2.3.F.** Let us return to the statement of the proposition and suppose, in addition, that  $N$  is a topologically free pseudocompact  $k$ -module. In this case one can choose, “coherently,” bases for the  $C$ -modules  $\mathbf{V}_k^f(N)(C)$ .

Indeed, let  $(n_i)$  be a pseudobasis of  $N$  (0.2.1), and let  $n_i^C$  be the canonical image of  $n_i$  in  $C \widehat{\otimes}_k N$ , for  $C \in \mathbf{Alf}/k$ . Define  $\delta_i^C \in (C \widehat{\otimes}_k N)^\dagger$  by

$$\delta_i^C(n_i^C) = 1, \quad \delta_i^C(n_j^C) = 0 \quad (i \neq j).$$

Then  $(\delta_i^C)$  is a basis of  $\mathbf{V}_k^f(N)(C)$ . Moreover, for every morphism  $\varphi : C \rightarrow D$  in  $\mathbf{Alf}/k$ , the map  $\mathbf{V}_k^f(N)(\varphi)$  sends  $\delta_i^C$  to  $\delta_i^D$ .

**1.2.4.**<sup>50</sup> For every pseudocompact  $k$ -module  $E$ , let  $\widehat{S}_k(E)$  denote the *completed symmetric algebra* of  $E$ , defined as follows. Let  $T_k(E)$  be the direct sum of the pseudocompact  $k$ -modules

$$\widehat{\bigotimes}_k^n E = E \widehat{\otimes}_k \cdots \widehat{\otimes}_k E \quad \left( n \geq 0; \quad \widehat{\bigotimes}_k^0 E = k \right).$$

Give  $T_k(E)$  its usual graded  $k$ -algebra multiplication. Let  $S_k(E)$  be the quotient of  $T_k(E)$  by the homogeneous ideal whose  $n$ -th component is the closed  $k$ -submodule of  $\widehat{\bigotimes}_k^n E$  generated by the elements

$$\begin{aligned} &x_1 \widehat{\otimes} \cdots \widehat{\otimes} x_i \widehat{\otimes} x_{i+1} \widehat{\otimes} \cdots \widehat{\otimes} x_n \\ &\quad - x_1 \widehat{\otimes} \cdots \widehat{\otimes} x_{i+1} \widehat{\otimes} x_i \widehat{\otimes} \cdots \widehat{\otimes} x_n. \end{aligned}$$

If this quotient is denoted by  $S_k^n(E)$ , then  $S_k(E)$  is a graded  $k$ -algebra whose  $n$ -th component is  $S_k^n(E)$ .

<sup>50</sup>Editors’ note: In this paragraph the order has been changed:  $\widehat{S}_k(E)$  is defined first, and it is then stated that  $\mathbf{V}_k^f(E)$  represents  $\mathbf{V}_k^f(E)$ .

Endow  $S_k(E)$  with the linear topology defined by the set  $\Upsilon$  of ideals  $\mathfrak{l}$  such that  $S_k(E)/\mathfrak{l}$  is a  $k$ -module of finite length and

$$S_k^n(E) \cap \mathfrak{l}$$

is an open submodule of  $S_k^n(E)$  for every  $n$ . The profinite algebra  $\widehat{S}_k(E)$  is the separated completion of  $S_k(E)$  for this topology.<sup>51</sup>

We write

$$\mathbf{V}_k^f(E) = \mathrm{Spf}(\widehat{S}_k(E))$$

for the corresponding formal  $k$ -variety. It represents the  $k$ -functor  $\mathbf{V}_k^f(E)$ ; that is, for every object  $C$  of  $\mathrm{Alf}/k$ , there is a canonical isomorphism

$$\begin{aligned} \mathbf{V}_k^f(E)(C) &= \mathrm{Hom}_c(E, C|_k) \xrightarrow{\sim} \mathrm{Hom}_{\mathrm{Alp}/k}(\widehat{S}_k(E), C) \\ &= \mathrm{Hom}_{\mathrm{Vaf}/k}(\mathrm{Spf}(C), \mathbf{V}_k^f(E)). \end{aligned}$$

In what follows, we identify the formal variety  $\mathbf{V}_k^f(E)$  with the  $k$ -functor  $\mathbf{V}_k^f(E)$ .

**1.2.5.** By 1.2.4, the zero morphism from  $E$  to  $k$  corresponds to a morphism of profinite algebras

$$\pi : \widehat{S}_k(E) \longrightarrow k.$$

The morphism  $\pi$  induces the zero map on  $S_k^n(E)$  for  $n \geq 1$ , and defines a section of the structure morphism

$$\mathbf{V}_k^f(E) \longrightarrow \mathrm{Spf}(k).$$

We denote by  $\mathbf{V}_k^{f,0}(E)$  the formal variety whose points are the images of the points of  $\mathrm{Spf}(k)$  under the section  $\mathrm{Spf}(\pi)$ , and whose local algebras at those points are the same as those of  $\mathbf{V}_k^f(E)$ .<sup>52</sup>

The affine algebra of  $\mathbf{V}_k^{f,0}(E)$  is thus the separated completion of  $S_k(E)$  for the topology defined by the ideals  $\mathfrak{l} \in \Upsilon$  (cf. 1.2.4) which contain  $S_k^n(E)$  for all sufficiently large  $n$ . Hence it is the infinite product

$$k[[E]] = k \times E \times S_k^2(E) \times S_k^3(E) \times \cdots.$$

Let  $C$  be an object of  $\mathrm{Alf}/k$ , let

$$u \in \mathbf{V}_k^f(E)(C) = \mathrm{Hom}_c(E, C),$$

and let

$$\phi : \widehat{S}_k(E) \longrightarrow C$$

<sup>51</sup>Editors' note: For example, let  $k$  be a field, let  $E$  be a pseudocompact  $k$ -vector space, and put  $V = \mathrm{Hom}_c(E, k)$ ; then  $E \simeq V^*$  (cf. 0.2.2). For every finite-dimensional subspace  $W$  of  $V$ , let  $\mathcal{F}(W)$  be the set of closed points of the  $k$ -scheme  $\mathbf{V}(W) = \mathrm{Spec} S_k(W^*)$ , and let  $\mathcal{F}(V) = \varinjlim_W \mathcal{F}(W)$ . Then  $\widehat{S}_k(E)$  is the product, over  $x \in \mathcal{F}(V)$ , of the pseudocompact  $k$ -algebras

$$\widehat{S}_k(E)_x = \varprojlim_W \widehat{\mathcal{O}}_{W,x},$$

where  $W$  ranges over the finite-dimensional subspaces of  $V$  such that  $x \in \mathcal{F}(W)$ , and  $\widehat{\mathcal{O}}_{W,x}$  is the separated completion of the local ring  $\mathcal{O}_{\mathbf{V}(W),x}$  for the topology defined by the ideals of finite codimension (which here coincides with the  $\mathfrak{m}$ -adic topology). If  $k$  is algebraically closed and  $(v_i)_{i \in I}$  is a basis of  $V$ , so that  $E$  has a pseudobasis  $(e_i)_{i \in I}$ , then every local component  $\widehat{S}_k(E)_x$  is isomorphic to the formal power-series ring  $k[[e_i, i \in I]]$ , endowed with the topology defined by the ideals of finite codimension.

<sup>52</sup>Editors' note: The remainder of this paragraph has been spelled out in detail.

be the corresponding morphism of profinite  $k$ -algebras. Then  $\text{Ker}(\phi)$  is an open ideal—that is,  $\phi$  belongs to  $\mathbf{V}_k^{f,0}(E)(C)$ —if and only if  $\text{Ker}(\phi)$  contains  $S_k^n(E)$  for all sufficiently large  $n$ , or equivalently if and only if  $u(E)$  is contained in the radical  $r(C)$  of  $C$ . Consequently, for every object  $C$  of  $\text{Alf}/k$ , there are canonical isomorphisms

$$\mathbf{V}_k^{f,0}(E)(C) \simeq \text{Hom}_{\text{Alp}/k}(k[[E]], C) \simeq \text{Hom}_c(E, r(C)).$$

**1.2.6.** A formal  $k$ -variety  $V$  is said to be of *finite length* if its affine algebra has finite length. Similarly, if  $S$  is a scheme, an  $S$ -scheme  $X$  is said to be of *finite length* if  $X$  is finite over  $S$  and the direct image of  $\mathcal{O}_X$  on  $S$  is an  $\mathcal{O}_S$ -module of finite length.<sup>53</sup> Thus giving an  $S$ -scheme  $X$  of finite length is the same as giving a finite set

$$\{s_1, \dots, s_n\}$$

of closed points of  $S$ , together with an  $\mathcal{O}_{S,s_i}$ -algebra of finite length at each such point.

It follows that the  $S$ -schemes of finite length identify with the formal varieties of finite length over the following formal scheme  $\widehat{S}$ . The underlying topological space of  $\widehat{S}$  is the set of closed points of  $S$ , endowed with the discrete topology. If  $s$  is such a closed point, the stalk  $\mathcal{O}_{\widehat{S},s}$  of the structure sheaf is the separated completion

$$\widehat{\mathcal{O}}_{S,s}$$

of  $\mathcal{O}_{S,s}$  for the linear topology defined by the ideals of finite colength. Thus

$$\widehat{S} = \text{Spf } \mathcal{A}(\widehat{S}),$$

where  $\mathcal{A}(\widehat{S})$  is the product of the  $\widehat{\mathcal{O}}_{S,s}$ , as  $s$  ranges over the closed points of  $S$ , endowed with the product topology.

**Definition.** If  $X$  is an  $S$ -scheme, let  $\widehat{X}/\widehat{S}$  denote the formal variety over

$$k = \mathcal{A}(\widehat{S})$$

defined as follows. Its underlying topological space consists of the points  $x \in X$  such that

$$[\kappa(x) : \kappa(s)] < \infty,$$

where  $s$  is the image of  $x$  in  $S$ . The local ring  $\mathcal{O}_{\widehat{X}/\widehat{S},x}$  is the separated completion of  $\mathcal{O}_{X,x}$  for the linear topology defined by the ideals  $I$  of  $\mathcal{O}_{X,x}$  for which  $\mathcal{O}_{X,x}/I$  has finite length as an  $\mathcal{O}_{S,s}$ -module. Since  $[\kappa(x) : \kappa(s)] < \infty$ , this is equivalent to requiring that  $\mathcal{O}_{X,x}/I$  have finite length as an  $\mathcal{O}_{X,x}$ -module.

Let  $\text{Vaf}^{\ell f}/\widehat{S}$  be the category of formal varieties of finite length over  $\widehat{S}$ , identified with the category of  $S$ -schemes of finite length. By 1.1 and 1.2.1, the category  $\text{Vaf}/\widehat{S}$  of formal varieties over  $\widehat{S}$  is equivalent to the category of left-exact contravariant functors

$$\text{Vaf}^{\ell f}/\widehat{S} \longrightarrow (\mathbf{Sets}).$$

In particular, for every  $S$ -scheme  $X$ , the functor

$$T \longmapsto \text{Hom}_{(\text{Sch}/S)}(T, X)$$

<sup>53</sup>Editors' note: The following sentence has been added.

from  $\mathbf{Vaf}^{\ell f}/\widehat{S}$  to **(Sets)** is left exact and therefore corresponds to a formal variety over  $\widehat{S}$ . This variety is precisely  $\widehat{X}/\widehat{S}$ .<sup>54</sup>

**Proposition.** *For every  $S$ -scheme  $X$ , the functors*

$$\mathrm{Hom}_{\mathbf{Vaf}/\widehat{S}}(-, \widehat{X}/\widehat{S}) \quad \text{and} \quad \mathrm{Hom}_{(\mathrm{Sch}/S)}(-, X)$$

*have the same restriction to  $\mathbf{Vaf}^{\ell f}/\widehat{S}$ . This defines a functor*

$$X \longmapsto \widehat{X}/\widehat{S}$$

*from  $(\mathrm{Sch}/S)$  to  $\mathbf{Vaf}/\widehat{S}$ , and this functor commutes with finite projective limits.*

*Proof.* It is immediate that the formal variety  $\widehat{X}/\widehat{S}$  has the required property and that the correspondence  $X \mapsto \widehat{X}/\widehat{S}$  is functorial. It remains to prove the second assertion.

Put

$$\mathcal{S} = (\mathrm{Sch}/S), \quad \mathcal{V} = \mathbf{Vaf}/\widehat{S},$$

and write  $\widehat{X}$  for  $\widehat{X}/\widehat{S}$ . By 1.2.B, the category  $\mathcal{V}$  has arbitrary projective limits. Let  $(X_i)_{i \in I}$  be a projective system in  $\mathcal{S}$ , and suppose that

$$X = \varprojlim_i X_i$$

exists in  $\mathcal{S}$ , as it does when  $I$  is finite. The functor which associates with every object  $Y$  of  $\mathcal{S}$  the functor

$$h_Y = \mathrm{Hom}_{\mathcal{S}}(-, Y)$$

and, respectively, with every object  $V$  of  $\mathcal{V}$  the functor

$$h_V = \mathrm{Hom}_{\mathcal{V}}(-, V)$$

commutes with projective limits. Hence, for every  $S$ -scheme  $T$  of finite length, there are functorial isomorphisms

$$\begin{aligned} \mathrm{Hom}_{\mathcal{S}}(T, X) &\simeq \varprojlim_i \mathrm{Hom}_{\mathcal{S}}(T, X_i) \\ &\simeq \varprojlim_i \mathrm{Hom}_{\mathcal{V}}(T, \widehat{X}_i) \simeq \mathrm{Hom}_{\mathcal{V}}\left(T, \varprojlim_i \widehat{X}_i\right). \end{aligned}$$

Consequently, the functor  $X \mapsto \widehat{X}/\widehat{S}$  commutes with projective limits whenever they exist in  $(\mathrm{Sch}/S)$ , and in particular with finite projective limits.  $\square$

### 1.3

**Proposition 1.3.** *Let  $f : X \rightarrow Y$  be a morphism of formal varieties over  $k$ , let  $A$  and  $B$  be the affine algebras of  $X$  and  $Y$ , and let  $g : B \rightarrow A$  be the morphism induced by  $f$ . Then  $f$  is a monomorphism in  $\mathbf{Vaf}/k$  if and only if  $g$  is surjective.*

By 1.1,<sup>55</sup> the functor  $A \mapsto \mathrm{Spf}(A)$  is an anti-equivalence from  $\mathbf{Alp}/k$  to  $\mathbf{Vaf}/k$ . Consequently,  $f$  is a monomorphism if and only if  $g$  is an epimorphism, which it certainly is when  $g$  is surjective.

<sup>54</sup>Editors' note: The following proposition has been expanded by adding the fact that the functor  $X \mapsto \widehat{X}/\widehat{S}$  commutes with finite projective limits. In the original this appeared in the proof of 1.3.4; the proof given here is more direct than the original one. This result will also be used in Section 2 and in 3.3.1.

<sup>55</sup>The editors expanded the beginning of the proof.

Conversely, suppose that  $g$  is an epimorphism. For every  $\mathfrak{n} \in \Upsilon(B)$ , put

$$A_{\mathfrak{n}} = A \widehat{\otimes}_B B_{\mathfrak{n}}.$$

By 0.4,  $A$  is a pseudocompact  $B$ -module, and hence is the product of the  $A_{\mathfrak{n}}$  (0.3.6). The morphism  $g$  is therefore the product of the morphisms

$$g_{\mathfrak{n}} : B_{\mathfrak{n}} \longrightarrow A_{\mathfrak{n}}$$

obtained from  $g$  by base change. These remain epimorphisms. We are thus reduced to the case in which  $B$  is local, with maximal ideal  $\mathfrak{n}$ ; set  $K = B/\mathfrak{n}$ .

By Nakayama's lemma 0.3.3, it is enough to prove that

$$g \widehat{\otimes}_B K$$

is surjective. This map is obtained from  $g$  by base change and is therefore an epimorphism in  $\text{Alp}/K$ . We may thus suppose that  $B = K$  is a field. Now  $f$  is a monomorphism if and only if the diagonal

$$X \longrightarrow X \times_Y X$$

is an isomorphism; equivalently, the homomorphism

$$A \widehat{\otimes}_K A \longrightarrow A, \quad x \widehat{\otimes}_K x' \longmapsto xx',$$

is an isomorphism. Since  $K$  is a field, this implies  $A = K$ .

**Remark.** *It follows from the proposition that every monomorphism  $f : X \rightarrow Y$  of formal varieties is an isomorphism from  $X$  onto a necessarily closed formal subvariety of  $Y$ .*<sup>56</sup>

### 1.3.1

The preceding proposition implies, in particular, that every monomorphism  $f : X \rightarrow Y$  in  $\text{Vaf}/k$  is effective (cf. IV 1.3).<sup>57</sup> The same is not true of epimorphisms, as one sees by slightly modifying the counterexample of Exposé V, § 2.c).<sup>58</sup> This is why we now consider a useful class of effective epimorphisms.

**Lemma.** <sup>59</sup> *Let  $f : X \rightarrow Y$  be a morphism of formal  $k$ -varieties, let  $A$  and  $B$  be the affine algebras of  $X$  and  $Y$ , and let  $f^{\natural} : B \rightarrow A$  be the morphism induced by  $f$ . The following conditions are equivalent:*

- (i) *For every  $x \in X$ , the homomorphism*

$$f_x^{\natural} : \mathcal{O}_{Y, f(x)} \longrightarrow \mathcal{O}_{X, x}$$

*makes  $\mathcal{O}_{X, x}$  a topologically free pseudocompact  $\mathcal{O}_{Y, f(x)}$ -module.*

<sup>56</sup>The editors added this remark.

<sup>57</sup>In the opposite category  $(\text{Vaf}/k)^{\circ} = \text{Alp}/k$ , the corresponding map  $g : B \rightarrow A$  is an effective epimorphism: since it is surjective, it induces an isomorphism of profinite  $k$ -algebras  $B/I \xrightarrow{\sim} A$ , where  $I = \ker g$ . Hence every morphism  $\phi : B \rightarrow C$  in  $\text{Alp}/k$  that vanishes on  $I$  descends to a morphism  $\bar{\phi} : A \rightarrow C$  such that  $\phi = \bar{\phi} \circ g$ .

<sup>58</sup>Let  $k$  be a field,  $X = \text{Spf}(k[[T]])$ , and  $Y = \text{Spf}(B)$ , where  $B$  is the  $k$ -subalgebra of  $A = k[[T]]$  topologically generated by  $T^3$  and  $T^4$ ; thus  $B$  consists of the series  $\sum a_n T^n$  with  $a_n = 0$  for  $n = 1, 2, 5$ . Then  $X \rightarrow Y$  is an epimorphism that is not effective: the cokernel of  $X \times_Y X \rightarrow X$  is  $\text{Spf}(B')$ , where

$$B' = \{a \in A \mid a \widehat{\otimes} 1 = 1 \widehat{\otimes} a\},$$

and  $B'$  contains  $T^5$ .

<sup>59</sup>The editors added this lemma, which explains the term “topologically flat.”

(ii) For every  $y \in Y$ , the local component

$$A_y = \prod_{f(x)=y} \mathcal{O}_{X,x}$$

is a topologically free pseudocompact  $B_y$ -module.

(iii) The map  $f^\sharp : B \rightarrow A$  makes  $A$  a projective pseudocompact  $B$ -module.

(iv) The functor

$$\mathrm{PC}(B) \longrightarrow \mathrm{PC}(A), \quad M \longmapsto M \widehat{\otimes}_B A,$$

is exact.

When these conditions hold,  $f$  is said to be topologically flat.

The implications

$$(i) \implies (ii) \iff (iii) \iff (iv)$$

follow from 0.2.F (iii) and 0.3.7. Conversely, suppose that (ii) holds, and let  $x \in X$ ,  $y = f(x)$ . Since  $\mathcal{O}_{X,x}$  is a direct factor of  $A_y$ , it is a projective pseudocompact  $B_y$ -module, and therefore is topologically free by 0.2.1, because  $B_y$  is local.

A morphism  $f : X \rightarrow Y$  of formal  $k$ -varieties is called *surjective* if it induces a surjection of the underlying sets.

**Proposition.** *Let  $f : X \rightarrow Y$  be a surjective, topologically flat morphism of formal  $k$ -varieties. Then  $f$  is an effective epimorphism (cf. IV 1.3).*

Let  $A$  and  $B$  be the affine algebras of  $X$  and  $Y$ , and let  $g : B \rightarrow A$  be the morphism induced by  $f$ . We must prove that  $Y$  identifies with the cokernel of

$$X \times_Y X \rightrightarrows X;$$

that is, for every  $\mathfrak{n} \in \Upsilon(B)$ ,  $B_{\mathfrak{n}}$  must identify with the subring of

$$A_{\mathfrak{n}} = A \widehat{\otimes}_B B_{\mathfrak{n}}$$

formed by the elements  $a$  such that

$$a \widehat{\otimes} 1 = 1 \widehat{\otimes} a.$$

We may suppose that  $B$  is local, with maximal ideal  $\mathfrak{n}$ . The hypothesis then says that  $g$  makes  $A$  a nonzero topologically free  $B$ -module. By Nakayama's lemma 0.3.3,  $A/\overline{\mathfrak{n}A} \neq 0$ , and hence the induced morphism

$$g' : B/\mathfrak{n} \longrightarrow A/\overline{\mathfrak{n}A}$$

is injective. By Lemma 1.3.2 below,  $B$  is a direct factor of  $A$  as a  $B$ -module; write  $A = B \oplus A'$ . It follows that  $B$  identifies with the set of  $a \in A$  such that

$$a \widehat{\otimes}_B 1 = 1 \widehat{\otimes}_B a.$$

## 1.3.2

**Lemma 1.3.2.** *Let  $B$  be a local pseudocompact ring, let  $\mathfrak{n}$  be its maximal ideal, let  $M, N$  be projective pseudocompact  $B$ -modules, and let  $g : M \rightarrow N$  be a morphism. If*

$$(B/\mathfrak{n})\widehat{\otimes}_B g$$

*is injective, then  $g$  is an isomorphism from  $M$  onto a direct factor of  $N$ .*

Indeed, suppose

$$g' = (B/\mathfrak{n})\widehat{\otimes}_B g$$

is injective. Since  $B/\mathfrak{n}$  is a field,  $g'$  has a retraction  $r'$ . Let  $p$  and  $q$  be the canonical projections from  $M$  and  $N$  onto

$$M/\overline{\mathfrak{n}M} \quad \text{and} \quad N/\overline{\mathfrak{n}N}.$$

Since  $N$  is projective, there exists  $r : N \rightarrow M$  such that

$$p \circ r = r' \circ q;$$

thus  $r'$  is obtained from  $r$  by passage to the quotient. Since  $r' \circ g'$  is an isomorphism, 0.3.4 shows that the same is true of  $r \circ g$ , because  $M$  is projective. If  $s$  denotes the inverse of  $r \circ g$ , then  $s \circ r$  is a retraction of  $g$ .

## 1.3.3

**Proposition 1.3.3.** *Let  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  be morphisms of formal  $k$ -varieties.*

- (i) *If  $f$  and  $g$  are topologically flat, so is  $g \circ f$ .*
- (ii) *If  $f$  and  $g \circ f$  are topologically flat and  $f$  is surjective, then  $g$  is topologically flat.*
- (iii) *If  $f$  is topologically flat, then so is  $f \times_Y Y'$  after every base change  $Y' \rightarrow Y$ .*

Assertions (i) and (iii) are clear. To prove (ii), let  $A, B, C$  be the affine algebras of  $X, Y, Z$ , and let  $f' : B \rightarrow A$ ,  $g' : C \rightarrow B$  be the homomorphisms induced by  $f, g$ . Since  $g \circ f$  is topologically flat,  $f' \circ g'$  makes  $A$  a projective pseudocompact  $C$ -module. Likewise,  $f'$  makes  $A$  a faithful projective pseudocompact  $B$ -module. Thus, as  $P$  ranges over pseudocompact  $C$ -modules and  $N$  over pseudocompact  $B$ -modules, the functors

$$P \mapsto P\widehat{\otimes}_C A, \quad N \mapsto N\widehat{\otimes}_B A$$

are exact; the second is also faithful. Hence the functor

$$P \mapsto P\widehat{\otimes}_C B$$

is exact, and 0.3.7 implies that  $B$  is a projective pseudocompact  $C$ -module.

## 1.3.4

**Proposition 1.3.4.** *Let  $S$  be a scheme, let  $Y$  be a locally Noetherian  $S$ -scheme, and let  $f : X \rightarrow Y$  be a faithfully flat  $S$ -morphism locally of finite type. Thus  $f$  is an effective epimorphism; equivalently, the following sequence is exact (cf. IV 6.3.1 (iv) and IV 1.3):*

$$(*) \quad X \times_Y X \xrightarrow[\text{pr}_2]{\text{pr}_1} X \xrightarrow{f} Y.$$

Then the morphism of formal  $\widehat{S}$ -varieties

$$\widehat{f}: \widehat{X}/\widehat{S} \longrightarrow \widehat{Y}/\widehat{S}$$

(cf. 1.2.6) is surjective and topologically flat, and the following sequence, obtained from the preceding one, is exact:

$$(*) \quad \widehat{X \times_Y X}/\widehat{S} \xrightarrow[\widehat{\text{pr}}_2]{\widehat{\text{pr}}_1} \widehat{X}/\widehat{S} \xrightarrow{\widehat{f}} \widehat{Y}/\widehat{S}.$$

Let  $y$  be a point of  $Y$ , with image  $s$  in  $S$ , such that  $\kappa(y)$  is a finite extension of the residue field  $\kappa(s)$ . Since  $f$  is surjective and locally of finite type,  $f^{-1}(y)$  is nonempty and locally of finite type over  $\kappa(y)$ . Its closed points are precisely the points of  $\widehat{X}/\widehat{S}$  mapping to  $y$ . This proves that  $\widehat{f}$  is surjective.

Let  $x$  be a closed point of  $f^{-1}(y)$ .<sup>60</sup> Since  $Y$  is locally Noetherian and  $f$  is locally of finite type, the local rings  $\mathcal{O}_{Y,y}$  and  $\mathcal{O}_{X,x}$  are Noetherian. The powers of their maximal ideals therefore have finite colength, and the local rings of  $\widehat{Y}/\widehat{S}$  at  $y$  and of  $\widehat{X}/\widehat{S}$  at  $x$  are the completions

$$\widehat{\mathcal{O}}_{Y,y} \quad \text{and} \quad \widehat{\mathcal{O}}_{X,x}$$

for their adic topologies. Since  $f$  is flat,  $\widehat{\mathcal{O}}_{X,x}$  is flat over  $\widehat{\mathcal{O}}_{Y,y}$  by SGA 1, IV 5.8. By 0.3.8 it is therefore a topologically free  $\widehat{\mathcal{O}}_{Y,y}$ -module. Thus  $\widehat{f}$  is topologically flat.

Proposition 1.3.1 now shows that  $\widehat{f}$  is an effective epimorphism. In other words, the following sequence is exact, where  $\widehat{X}$  is written for  $\widehat{X}/\widehat{S}$ :

$$\widehat{X} \times_{\widehat{Y}} \widehat{X} \xrightarrow[\widehat{\text{pr}}_2]{\widehat{\text{pr}}_1} \widehat{X} \xrightarrow{\widehat{f}} \widehat{Y}.$$

Moreover, 1.2.6 gives a natural isomorphism, compatible in particular with the projections onto  $\widehat{X}$ :

$$\widehat{X \times_Y X} \simeq \widehat{X} \times_{\widehat{Y}} \widehat{X}.$$

Consequently, the completed sequence displayed in the proposition is exact.

### 1.3.5

Let  $k$  be a pseudocompact ring. A formal variety  $X$  over  $k$  is said to be *topologically flat* if its affine algebra  $A$  is a *projective* pseudocompact  $k$ -module, that is, if the structure morphism

$$X \longrightarrow \text{Spf}(k)$$

is topologically flat.

Let us first note<sup>61</sup> that 0.2.2 and 0.3.6 imply the following result, analogous to VII<sub>A</sub>, 3.1.1.

**Lemma 1.3.5.A.** *Suppose that  $k$  is Artinian. The functors*

$$A \longmapsto A^\dagger = \text{Hom}_c(A, k), \quad C \longmapsto C^* = \text{Hom}_k(C, k)$$

<sup>60</sup>The editors expanded the argument that follows and then used the material added in 1.2.6.

<sup>61</sup>The editors added the following lemma, used implicitly in the original; they also introduced the numbering 1.3.5.A–1.3.5.D for later references.



define an anti-equivalence between the category of topologically flat profinite  $k$ -algebras and the category of flat  $k$ -coalgebras.

Indeed, if  $A$  is a topologically flat profinite  $k$ -algebra, then, by 0.3.6, there is an isomorphism of  $k$ -modules

$$A^\dagger \otimes_k A^\dagger \xrightarrow{\sim} (A \widehat{\otimes} A)^\dagger.$$

Thus the multiplication

$$A \widehat{\otimes} A \longrightarrow A$$

induces by duality a  $k$ -coalgebra structure on  $A^\dagger$ . The rest then follows from Proposition 0.2.2.

Let us return to the case of an arbitrary pseudocompact ring  $k$ .

**Definition 1.3.5.B.** To every formal  $k$ -variety  $X$  whose affine algebra  $A$  is a projective pseudocompact  $k$ -module, one associates a flat  $\mathbf{O}_k$ -coalgebra  $\mathbf{H}(X)$ , defined as follows.

Denote by  $\mathbf{H}(X)$  the  $\mathbf{O}_k$ -module  $\mathbf{V}_k^f(A)$ , the “dual of  $A$ ”. It is a flat  $\mathbf{O}_k$ -module, since the pseudocompact  $k$ -module underlying  $A$  is projective (cf. 1.2.3.C). Moreover, by 0.3.6, there is an isomorphism

$$\mathbf{V}_k^f(A \widehat{\otimes} A) \simeq \mathbf{V}_k^f(A) \otimes \mathbf{V}_k^f(A).$$

Consequently, the multiplication of  $A$  induces by transposition a diagonal morphism

$$\mathbf{H}(X) = \mathbf{V}_k^f(A) \longrightarrow \mathbf{V}_k^f(A \widehat{\otimes} A) = \mathbf{H}(X) \otimes \mathbf{H}(X),$$

which makes  $\mathbf{H}(X)$  a flat  $\mathbf{O}_k$ -coalgebra. We shall say that  $\mathbf{H}(X)$  is the *coalgebra of  $X$  over  $\mathbf{O}_k$* .

**Definition 1.3.5.C.** Conversely, to every  $\mathbf{O}_k$ -coalgebra  $\mathbf{C}$  one can associate a  $k$ -functor (cf. 1.2.1)  $\mathrm{Spf}^*(\mathbf{C})$ , defined as follows. For every object  $B$  of  $\mathrm{Alf}/k$ , set, with the notation of VII<sub>A</sub> 3.1,

$$\begin{aligned} \mathrm{Spf}^*(\mathbf{C})(B) &= \mathrm{Hom}_{B\text{-coalg.}}(B, \mathbf{C}(B)) \\ &= \{x \in \mathbf{C}(B) \mid \varepsilon_{\mathbf{C}(B)}(x) = 1 \text{ and } \Delta_{\mathbf{C}(B)}(x) = x \otimes x\}. \end{aligned} \quad (1)$$

Suppose in addition<sup>62</sup> that the  $\mathbf{O}_k$ -module underlying  $\mathbf{C}$  is admissible (cf. 1.2.1), and put

$$A = \Gamma^*(\mathbf{C}) = \varprojlim_{\mathfrak{l}} \mathbf{C}(k/\mathfrak{l})^*. \quad (2)$$

The algebra structures on the modules  $\mathbf{C}(k/\mathfrak{l})^*$  endow  $A$  with the structure of a profinite  $k$ -algebra. For every object  $B$  of  $\mathrm{Alf}/k$ , one has

$$\begin{aligned} \mathrm{Hom}_{\mathrm{Vaf}/k}(\mathrm{Spf}(B), \mathrm{Spf}(A)) &= \mathrm{Hom}_{\mathrm{Alp}/k}(A, B) \\ &= \mathrm{Hom}_{\mathrm{Alp}/B}(A \widehat{\otimes} B, B). \end{aligned} \quad (3)$$

Suppose finally that  $\mathbf{C}$  is a flat  $\mathbf{O}_k$ -module. Then, by 1.2.3.E,  $A = \Gamma^*(\mathbf{C})$  is a projective pseudocompact  $k$ -module. Moreover, we saw in the proof of *loc. cit.* that if  $\mathfrak{l}_0$  is an open ideal of  $k$  contained in the kernel of  $k \rightarrow B$ , there are isomorphisms

$$\begin{aligned} A \widehat{\otimes} B &= \varprojlim_{\mathfrak{l}} \mathbf{C}(k/\mathfrak{l})^* \widehat{\otimes}_k B \\ &\simeq \mathrm{Hom}_{k/\mathfrak{l}_0}(\mathbf{C}(k/\mathfrak{l}_0), B) \\ &\simeq \mathrm{Hom}_B(\mathbf{C}(B), B). \end{aligned} \quad (4)$$

<sup>62</sup>The editors supplied the details that follow.

We shall denote the term on the right by  $\mathbf{C}(B)^*$ . Finally, Lemma 1.3.5.A, applied to the Artinian ring  $B$ , gives a natural isomorphism

$$\mathrm{Hom}_{B\text{-}\mathrm{coalg.}}(B, \mathbf{C}(B)) \xrightarrow{\sim} \mathrm{Hom}_{\mathrm{Alp}/B}(\mathbf{C}(B)^*, B). \quad (5)$$

Combining (1), (5), (4), (3), and (2), we obtain, when  $\mathbf{C}$  is a flat  $\mathbf{O}_k$ -coalgebra, an isomorphism of functors

$$\mathrm{Spf}^*(\mathbf{C}) \simeq \mathrm{Spf}(A) = \mathrm{Spf}(\Gamma^*(\mathbf{C})). \quad (\star)$$

Consequently, if  $\mathcal{A}(X)$  denotes the affine algebra of a formal  $k$ -variety  $X$ , then, taking 1.2.3.E into account, we obtain the following.

**Proposition 1.3.5.D.**

(i) *The functors*

$$X \mapsto \mathbf{H}(X) = \mathbf{V}_k^f(\mathcal{A}(X)), \quad \mathbf{C} \mapsto \mathrm{Spf}^*(\mathbf{C}) = \mathrm{Spf}(\Gamma^*(\mathbf{C}))$$

*define an equivalence between the category of topologically flat formal  $k$ -varieties and the category of flat  $\mathbf{O}_k$ -coalgebras.*

(ii) *Moreover, this equivalence “commutes with base change”: if  $k \rightarrow K$  is a morphism of pseudocompact rings, then  $X \hat{\otimes}_k K$  corresponds to  $\mathbf{H}(X) \otimes_k K$ .*

### 1.3.6

In the remainder of this Exposé, we shall several times define a topologically flat formal  $k$ -variety  $X$  by exhibiting the coalgebra  $\mathbf{H}(X)$ . We shall then have to interpret, by means of  $\mathbf{H}(X)$ , certain geometric properties of  $X$ . Here is one example of this situation: suppose that a section  $\sigma$  of the structure morphism  $X \rightarrow \mathrm{Spf}(k)$  is given, and ask under what condition  $\sigma$  induces an isomorphism of the underlying topological spaces.<sup>63</sup>

*To begin, suppose that  $k$  is Artinian.* Let  $(H, \Delta, \varepsilon)$  be a flat  $k$ -coalgebra, put  $H^+ = \ker(\varepsilon)$ , and let  $A = H^*$  be the profinite  $k$ -algebra dual to  $H$ . Suppose that a morphism of  $k$ -coalgebras  $k \rightarrow H$  is given; that is, an element  $\phi \in H$  such that

$$\varepsilon(\phi) = 1, \quad \Delta(\phi) = \phi \otimes \phi.$$

On the one hand,  $\phi$  defines a continuous algebra homomorphism  $\Phi : A \rightarrow k$ , and hence a section

$$\sigma : \mathrm{Spf}(k) \rightarrow \mathrm{Spf}(A)$$

of the projection  $\mathrm{Spf}(A) \rightarrow \mathrm{Spf}(k)$ .

On the other hand, define  $k$ -submodules of  $H$  by setting  $H_0 = k\phi$  and, for  $n \geq 1$ ,

$$H_n = \{x \in H \mid \Delta(x) - x \otimes \phi \in H_{n-1} \otimes H^+\}.$$

This formula also applies when  $n = 0$  if  $H_{-1} = 0$ . Induction on  $n$  shows that  $H_{n-1} \subset H_n$ . We call

$$H_0 \subset H_1 \subset \cdots$$

the *filtration of  $H$  defined by  $\phi$* .

**Remark.** Since

$$\Delta(H_n) \subset H_n \otimes H_0 \oplus H_{n-1} \otimes H^+,$$

<sup>63</sup>In Lemma 1.3.6.A below, the editors supplied detailed proofs of parts (i) and (ii) and added part (iii).

we have  $\Delta(H_n) \subset H_n \otimes H$ . Since  $\Delta$  is cocommutative—that is,  $\tau \circ \Delta = \Delta$ , where  $\tau(a \otimes b) = b \otimes a$ —we likewise have  $\Delta(H_n) \subset H \otimes H_n$ . When  $H/H_n$  is flat over  $k$ , it follows that  $H_n$  is a subcoalgebra of  $H$  (see also 1.3.6.A (iii) below). In general, however, the map

$$\Delta : H_n \longrightarrow H_n \otimes H$$

does not factor through  $H_n \otimes H_n$ .<sup>64</sup>

**Lemma 1.3.6.A.** *Let  $k$  be an Artinian ring, let  $H$  be a flat  $k$ -coalgebra, and let  $A = H^*$  be its dual profinite  $k$ -algebra. Let  $\phi \in H$  satisfy  $\varepsilon(\phi) = 1$  and  $\Delta(\phi) = \phi \otimes \phi$ . Let  $\Phi : A \rightarrow k$  be the continuous algebra homomorphism, let  $\sigma : \mathrm{Spf}(k) \rightarrow \mathrm{Spf}(A)$  be the section of  $\mathrm{Spf}(A) \rightarrow \mathrm{Spf}(k)$ , and let  $(H_n)$  be the filtration of  $H$  defined by  $\phi$ . Put  $I = \ker \Phi$ .*

- (i) *For every  $n \geq 1$ ,  $H_{n-1}$  is the orthogonal in  $H$  of the closure  $\overline{I^n}$  of  $I^n$ .*
- (ii) *Consequently,  $\sigma$  induces a bijection of the underlying sets if and only if*

$$H = \bigcup_n H_n.$$

65

- (iii) *If, in addition, every  $H/H_n$  is flat over  $k$ , then for every  $n \geq 0$ ,*

$$\Delta(H_n) \subset \sum_{i=0}^n H_i \otimes H_{n-i}. \quad (*)$$

*In particular, every  $H_n$  is then a subcoalgebra of  $H$ .*

*Proof.* For every  $x \in H$ , the map

$$A \longrightarrow k, \quad f \longmapsto f(x),$$

is continuous. Thus, if  $I^n$  is annihilated by  $x$ , so is its closure  $\overline{I^n}$ . For  $n \geq 1$ , put

$$(\overline{I^n})^\perp = \{x \in H \mid f(x) = 0 \text{ for every } f \in \overline{I^n}\}.$$

Suppose first that

$$\sigma : \mathfrak{m} \longmapsto \Phi^{-1}(\mathfrak{m})$$

is a bijection from  $\mathrm{Spf}(k)$  onto  $\mathrm{Spf}(A)$ . Since  $I$  is contained in the intersection of the ideals  $\Phi^{-1}(\mathfrak{m})$ , it follows from 0.1.2 that the sequence of ideals  $(\overline{I^n})$  tends to zero. Let  $x \in H$ . The submodule

$$J(x) = \{f \in A \mid f(x) = 0\}$$

is open and closed in  $A$ ; hence it contains  $\overline{I^n}$  for all sufficiently large  $n$ . In other words,  $x \in (\overline{I^n})^\perp$  for all sufficiently large  $n$ .

Conversely, suppose that

$$H = \bigcup_n (\overline{I^n})^\perp.$$

<sup>64</sup>For example, let  $k_0$  be a field, let  $k = k_0[T]/(T^n)$ , where  $n \geq 4$ , and let  $H = k\phi \oplus kx$ , with  $\varepsilon(x) = 0$  and

$$\Delta(x) = x \otimes \phi + \phi \otimes x + tx \otimes x,$$

where  $t$  is the image of  $T$  in  $k$ . Then  $H_i = k\phi \oplus t^{n-i}x$  for  $i = 0, \dots, n$ , but for  $2 \leq i \leq n-2$ , the element  $\Delta(t^{n-i}x)$  does not belong to the image of  $H_i \otimes H_i$  in  $H \otimes H$ .

<sup>65</sup>In this case the coalgebra  $H$  is said to be *connected*; cf. Addition 2.9.

Let  $\mathfrak{p}$  be an open prime ideal of  $A$ . By the definition of the topology on  $A$  (0.2.2),  $\mathfrak{p}$  contains an open  $k$ -submodule of the form

$$\mathcal{V}(x_1, \dots, x_s) = \{f \in A \mid f(x_1) = \dots = f(x_s) = 0\}.$$

By hypothesis, there is an integer  $n$  such that

$$x_1, \dots, x_s \in (\overline{I^n})^\perp;$$

therefore  $\overline{I^n} \subset \mathfrak{p}$ . Moreover, since  $k$  is Artinian,

$$\mathrm{Spf}(k) = \{\mathfrak{m}_1, \dots, \mathfrak{m}_r\}$$

is finite, and there is an integer  $t \geq 1$  such that

$$\left(\prod_i \mathfrak{m}_i\right)^t = 0.$$

It follows that

$$\prod_i \Phi^{-1}(\mathfrak{m}_i)^t \subset I.$$

Thus  $\mathfrak{p}$  contains the product of the ideals  $\Phi^{-1}(\mathfrak{m}_i)^{tn}$ . Since  $\mathfrak{p}$  is prime, it contains, and hence equals, one of the  $\Phi^{-1}(\mathfrak{m}_i)$ . We have therefore proved that <sup>66</sup>

$$\sigma \text{ is a bijection} \iff H = \bigcup_n (\overline{I^n})^\perp. \quad (65)$$

We also have  $H = k\phi \oplus H^+$ ; let  $\pi : H \rightarrow H^+$  be the projection with kernel  $k\phi$ . For  $n \geq 0$ , let

$$\Delta^n : H \longrightarrow H^{\otimes(n+1)}$$

be the iterated comultiplication, let  $\overline{\Delta}^n$  be the composite of  $\Delta^n$  with the projection

$$\pi^{\otimes(n+1)} : H^{\otimes(n+1)} \longrightarrow (H^+)^{\otimes(n+1)},$$

and put

$$H'_n = \ker(\overline{\Delta}^n) = \left\{ x \in H \mid \Delta^n(x) \in \sum_{i=0}^n H^{\otimes(n-i)} \otimes H_0 \otimes H^{\otimes i} \right\}.$$

We set  $\overline{\Delta}^0 = \mathrm{id}_H$ , so that  $H'_0 = H_0$ . A straightforward induction on  $n$  gives

$$H_n \subset H'_n \subset (\overline{I^{n+1}})^\perp. \quad (*)$$

Up to this point we have not used the hypothesis that  $H$  is flat over  $k$ . Assume now that  $H$  is flat, and hence projective, over  $k$ . Then  $A^\dagger = H$  by 0.2.2. We shall prove that

$$H_n = (\overline{I^{n+1}})^\perp.$$

This is clear for  $n = 0$ . Suppose it has been proved for  $n < r$ . Then  $H_{r-1}$  is the kernel of the homomorphism

$$H \longrightarrow (\overline{I^r})^\dagger.$$

<sup>66</sup>This remains valid without assuming that  $H$  is cocommutative, with  $k$  still a commutative Artinian ring. In that case a basis of neighborhoods of zero in  $A = H^*$  is given by the two-sided ideals  $J$  such that  $A/J$  has finite length over  $k$ , and the preceding proof shows that  $H = \bigcup_n (\overline{I^n})^\perp$  if and only if the  $\Phi^{-1}(\mathfrak{m}_i)$  are the only open prime ideals of  $A$ .

Since  $H^+$  is flat, the homomorphism

$$(H/H_{r-1}) \otimes H^+ \longrightarrow (\overline{I}^r)^\dagger \otimes H^+$$

is injective. On the other hand, the hypothesis implies that  $I$  is a projective pseudocompact  $k$ -module, since it is a direct factor of  $A = H^*$ . Thus, by 0.3.6,

$$(\overline{I}^r \widehat{\otimes} I)^\dagger \simeq (\overline{I}^r)^\dagger \otimes I^\dagger = (\overline{I}^r)^\dagger \otimes H^+.$$

The exact sequence

$$\overline{I}^r \widehat{\otimes} I \longrightarrow A \longrightarrow A/\overline{I}^{r+1} \longrightarrow 0$$

therefore gives, by duality, the exact sequence

$$(\overline{I}^r)^\dagger \otimes H^+ \xleftarrow{\delta} H \longleftarrow (A/\overline{I}^{r+1})^\dagger \longleftarrow 0, \quad (1)$$

where  $\delta$  is obtained by composing  $\Delta_H$  with the projection

$$H \otimes H \longrightarrow H \otimes H^+ \longrightarrow (H/H_{r-1}) \otimes H^+ \hookrightarrow (\overline{I}^r)^\dagger \otimes H^+. \quad (2)$$

For every  $u \in H$ , the projection of  $\Delta(u)$  onto  $H \otimes H^+$  is  $\Delta(u) - u \otimes \phi$ . Equations (1) and (2) show that if

$$u \in (A/\overline{I}^{r+1})^\dagger = (\overline{I}^{r+1})^\perp,$$

then  $\Delta(u) - u \otimes \phi$  belongs to the kernel of

$$H \otimes H^+ \longrightarrow (H/H_{r-1}) \otimes H^+,$$

namely  $H_{r-1} \otimes H^+$ . Thus  $u \in H_r$ . This completes the proof of (i) and (ii), and also proves that

$$H_n = \ker(\overline{\Delta}^n).$$

We now prove (iii). For  $i \geq 0$ , put

$$H_i^+ = H_i \cap H^+.$$

Let  $n \geq 1$ . For every  $x \in H_n$ , the element  $\bar{x} = x - \varepsilon(x)\phi$  belongs to  $H_n^+$ , and

$$\Delta(x) = \varepsilon(x)\phi \otimes \phi + \bar{x} \otimes \phi + \phi \otimes \bar{x} + \overline{\Delta}(\bar{x}).$$

It is therefore enough to prove that

$$\overline{\Delta}(H_n^+) \subset \sum_{i=1}^{n-1} H_i^+ \otimes H_{n-i}^+.$$

For  $i = 0, \dots, n-1$ , the map  $\overline{\Delta}^n$  factors as follows:

$$\begin{array}{ccc} H^+ & \xrightarrow{\overline{\Delta}} & H^+ \otimes H^+ \xrightarrow{\overline{\Delta}^i \otimes \overline{\Delta}^{n-i-1}} (H^+)^{\otimes(i+1)} \otimes (H^+)^{\otimes(n-i)} \\ & \downarrow & \uparrow g \\ & \frac{H^+}{H_i^+} \otimes \frac{H^+}{H_{n-i-1}^+} \xrightarrow{f} \frac{H^+}{H_i^+} \otimes (H^+)^{\otimes(n-i)}. \end{array}$$

Moreover, since  $H^+/H_i^+$  and  $(H^+)^{\otimes(n-i)}$  are flat, the maps  $f$  and  $g$  in this diagram are injective. It follows that

$$\overline{\Delta}(H_n^+) \subset H_i^+ \otimes H^+ + H^+ \otimes H_{n-i-1}^+$$

for every  $i = 0, \dots, n-1$ . Part (iii) now follows from the next lemma, applied with  $M = H^+$ ,  $E_0 = 0$ , and  $E_i = H_{i-1}^+$  for  $i \geq 1$ .

**Lemma 1.3.6.B.** *Let  $k$  be a ring, and let*

$$0 = E_0 \subset E_1 \subset \dots \subset E_n \subset M$$

*be  $k$ -modules. Suppose that  $M/E_i$  is flat for every  $i$ . Then*<sup>67</sup>

$$\bigcap_{i=0}^n (E_i \otimes M + M \otimes E_{n-i}) = \sum_{i=1}^n E_i \otimes E_{n-i+1}.$$

*Proof.* Denote the left-hand and right-hand sides by  $K$  and  $S$ , respectively. Clearly  $S \subset K$ ; we prove the reverse inclusion. For  $i = 0, \dots, n$ , put

$$K_i = K \cap (E_i \otimes M).$$

For  $i = 1, \dots, n$ , both  $M/E_{n-i+1}$  and  $E_i/E_{i-1}$  are flat. Consequently, the homomorphism  $\tau_i$  below is injective, and the composite

$$(E_i/E_{i-1}) \otimes M \longrightarrow (E_i/E_{i-1}) \otimes (M/E_{n-i+1}) \xrightarrow{\tau_i} (M/E_{i-1}) \otimes (M/E_{n-i+1})$$

has kernel

$$(E_i/E_{i-1}) \otimes E_{n-i+1}.$$

The image of  $K_i$  in  $(E_i \otimes M)/(E_{i-1} \otimes M)$  is both contained in and contains this kernel. Hence

$$K_i = K_{i-1} + E_i \otimes E_{n-i+1},$$

which proves the lemma.

To conclude this paragraph, return to an arbitrary pseudocompact ring  $k$ .<sup>68</sup> Let  $(\mathbf{H}, \Delta, \varepsilon)$  be a flat  $\mathbf{O}_k$ -coalgebra, put  $\mathbf{H}^+ = \ker(\varepsilon)$ , and let

$$A = \Gamma^*(\mathbf{H})$$

be the dual profinite  $k$ -algebra. Put  $X = \mathrm{Spf}(A)$ , so that  $\mathbf{H} = \mathbf{H}(X)$  (cf. 1.3.5). Suppose that a morphism of  $\mathbf{O}_k$ -coalgebras

$$\phi : \mathbf{O}_k \longrightarrow \mathbf{H}$$

is given. It defines a continuous homomorphism of  $k$ -algebras  $A \rightarrow k$ , and hence a section  $\sigma$  of the structure morphism  $X \rightarrow \mathrm{Spf}(k)$ .

For every object  $B$  of  $\mathrm{Alf}/k$ , put

$$\mathbf{H}_0(B) = \phi(B) = B\phi_B, \quad \phi_B = \phi(1_B) \in \mathbf{H}(B).$$

<sup>67</sup>The French source prints  $M \otimes E_i$ , rather than  $M \otimes E_{n-i}$ , in the second summand on the left. The proof immediately below, in particular its use of  $M/E_{n-i+1}$ , requires the latter indexing, which is used here.

<sup>68</sup>The editors added what follows; the original treated only the case in which  $k$  is Artinian.

For  $n \geq 1$ , define  $\mathbf{O}_k$ -submodules  $\mathbf{H}_n$  of  $\mathbf{H}$  by

$$\mathbf{H}_n(B) = \{u \in \mathbf{H}(B) \mid \Delta(u) - u \otimes \phi_B \in \mathbf{H}_{n-1}(B) \otimes \mathbf{H}^+(B)\}.$$

This gives a filtration

$$\mathbf{H}_0 \subset \mathbf{H}_1 \subset \cdots$$

of  $\mathbf{H}(X)$ . The preceding result gives:

**Proposition.** *The section  $\sigma$  induces an isomorphism of the underlying topological spaces if and only if*

$$\mathbf{H}(X) = \bigcup_n \mathbf{H}_n.$$

## 1.4

**Theorem 1.4.** *Let  $k$  be a pseudocompact ring and let*

$$d_0, d_1 : X_1 \rightrightarrows X$$

*be an equivalence pair in  $\mathbf{Vaf}/k$  (cf. Exposé V, § 2.b) such that  $d_1$  is topologically flat.*

- (i) *The canonical projection from  $X$  onto  $X/X_1 = \text{Coker}(d_0, d_1)$  is surjective and topologically flat, and the morphism*

$$X_1 \longrightarrow X \times_{X/X_1} X$$

*with components  $d_0$  and  $d_1$  is an isomorphism.*

- (ii) *If  $X$  is topologically flat over  $k$ , then so is  $X/X_1$ .*

First note that (ii) follows from (i) by 1.3.3 (ii). The proof of (i) occupies 1.4.1, 1.4.2, and 1.4.3.

### 1.4.1

We first show that we may reduce to the case in which  $X$  has only one point. Since we are dealing with an equivalence pair, one sees, as in Exposé V, § 3.e), that an equivalence relation is defined on the underlying set of  $X$  by declaring two points  $x, y$  to be equivalent if there exists a point  $z \in X_1$  such that

$$d_0(z) = y \quad \text{and} \quad d_1(z) = x.$$

We may plainly suppose, without loss of generality, that  $X$  contains a single equivalence class for this relation; in other words, that  $X/X_1$  has only one point (see the construction of  $X/X_1$  given in 1.2).

In this case, let  $x$  be any point of  $X$ , and let  $U$  be the formal variety having  $x$  as its only point and the same local ring as  $X$  at  $x$ . As in Exposé V, § 6, one then sees that the equivalence relation induced by  $(d_0, d_1)$  on  $U$  still satisfies the hypotheses of the theorem and that it is enough to prove the theorem for this latter equivalence relation ( $U$  is a “quasi-section”).

Let us briefly recall the principle of the proof given in Exposé V, § 6. Put

$$V = d_0^{-1}(U) = U \times_{i, d_0} X_1,$$

where  $i$  is the inclusion of  $U$  into  $X$ , and let  $u$  and  $v$  be the morphisms with source  $V$  induced respectively by  $d_0$  and  $d_1$ :

$$X \xleftarrow{v} V \xrightarrow{u} U.$$

It is clear that  $u$  and  $v$  are topologically flat and that  $u$  is surjective; since  $X$  contains a single equivalence class,  $v$  is surjective. If  $(v_0, v_1)$  is the inverse image of the equivalence pair  $(d_0, d_1)$  under  $v$  (cf. V, 3.a)), it follows from V, 3.c) and 3.d) that  $X/X_1$  and the quotient of  $U$  by the equivalence relation induced by  $(d_0, d_1)$  both identify with  $\text{Coker}(v_0, v_1)$ . As in the proof of V, 6.1, one then sees that if the conclusion of Theorem 1.4 holds for  $U$ , it also holds for  $X$ .

#### 1.4.2

We are thus reduced to the case in which  $X$  has only one point.<sup>69</sup> Consider the following commutative diagram (cf. V, § 1, (0,1,2)):

$$\begin{array}{ccccc} X_2 & \xrightleftharpoons[d'_0]{d'_1} & X_1 & \xrightarrow{d_0} & X \\ d'_2 \downarrow & & \downarrow d_1 & & \downarrow \\ X_1 & \xrightleftharpoons[d_0]{d_1} & X & \longrightarrow & X/X_1 \end{array}$$

Here

$$X_2 = X_1 \times_{d_1, d_0} X_1,$$

and  $d'_0, d'_1, d'_2$  are respectively the morphisms<sup>70</sup>

$$(x, y, z) \mapsto (x, y), \quad (x, y, z) \mapsto (x, z), \quad (x, y, z) \mapsto (y, z).$$

If  $B, A, A_1, A_2$  denote respectively the affine algebras of  $X/X_1, X, X_1, X_2$ , the preceding diagram induces the commutative diagram

$$\begin{array}{ccccc} A_2 & \xrightleftharpoons[j_0]{j_1} & A_1 & \xleftarrow{i_0} & A \\ j_2 \uparrow & & \uparrow i_1 & & \uparrow i \\ A_1 & \xrightleftharpoons[i_0]{i_1} & A & \xleftarrow{i} & B \end{array}$$

in which both rows are exact and the squares determined by  $(i_0, j_0)$  and  $(i_1, j_1)$  are cocartesian. Since the morphism

$$X_1 \longrightarrow X \times X$$

with components  $d_0$  and  $d_1$  is a monomorphism by hypothesis, the morphism

$$A \widehat{\otimes}_k A \longrightarrow A_1$$

with components  $i_0$  and  $i_1$  is surjective by Proposition 1.3.

<sup>69</sup>Editors' note: Thus one may suppose that  $k$  is local.

<sup>70</sup>Editors' note: Cf. Exposé V, § 2.b).



This means that  $i_1$  makes  $A_1$  a pseudocompact  $A$ -module, topologically free by hypothesis, generated by  $i_0(A)$ . Since  $A$  is local, Lemma 1.4.3 below gives a topologically free  $k$ -module  $V$  and a morphism of pseudocompact  $k$ -modules  $f : V \rightarrow A$  such that the morphism

$$\alpha_1 : A \widehat{\otimes}_k V \longrightarrow A_1, \quad a \widehat{\otimes} v \longmapsto i_1(a) \cdot i_0(f(v))$$

is an isomorphism. Let

$$\alpha : B \widehat{\otimes}_k V \longrightarrow A \quad \text{and} \quad \alpha_2 : A_1 \widehat{\otimes}_k V \longrightarrow A_2$$

be the morphisms

$$b \widehat{\otimes} v \longmapsto i(b) \cdot f(v) \quad \text{and} \quad a_1 \widehat{\otimes} v \longmapsto j_2(a_1) \cdot j_0 i_0(f(v)).$$

In the following commutative diagram, both rows are exact and the two left-hand squares are cocartesian. Since  $\alpha_1$  is an isomorphism, the same is true of  $\alpha_2$ , and hence of  $\alpha$ .

$$\begin{array}{ccccc} A_2 & \xleftarrow{j_1} & A_1 & \xleftarrow{i_0} & A \\ & \searrow j_0 & \uparrow \alpha_1 & & \uparrow \alpha \\ A_1 \widehat{\otimes}_k V & \xleftarrow[i_0 \widehat{\otimes} V]{i_1 \widehat{\otimes} V} & A \widehat{\otimes}_k V & \xleftarrow{i \widehat{\otimes} V} & B \widehat{\otimes}_k V \end{array}$$

This shows, on the one hand, that  $A$  is topologically free over  $B$ , with pseudobasis  $f(V)$  (cf. 0.2.1), and that a pseudobasis of  $A_1$  over  $A$ , where  $A_1$  is regarded as an  $A$ -module through  $i_1$ , is obtained by taking the image under  $i_0$  of  $f(V)$ . It follows that the morphism

$$A \widehat{\otimes}_B A \longrightarrow A_1$$

with components  $i_1$  and  $i_0$  is an isomorphism:

$$A \widehat{\otimes}_B A \simeq A \widehat{\otimes}_B B \widehat{\otimes}_k V \simeq A \widehat{\otimes}_k V \simeq A_1.$$

This proves Theorem 1.4, modulo the following lemma.

### 1.4.3

**Lemma 1.4.3.** *Let  $k$  be a pseudocompact ring, let  $A$  be a local profinite  $k$ -algebra, let  $A_1$  be a topologically free  $A$ -module, and let*

$$i_0 : M \longrightarrow A_1$$

*be a morphism of pseudocompact  $k$ -modules. Suppose that the map*

$$A \widehat{\otimes}_k M \longrightarrow A_1, \quad a \widehat{\otimes} m \longmapsto a \cdot i_0(m),$$

*is surjective. Then there exist a topologically free  $k$ -module  $V$  and a morphism of pseudocompact  $k$ -modules*

$$f : V \longrightarrow M$$

*such that the map*

$$A \widehat{\otimes}_k V \longrightarrow A_1, \quad a \widehat{\otimes} v \longmapsto a \cdot i_0(f(v)),$$

*is bijective.*

Since every pseudocompact  $k$ -module is a quotient of a topologically free  $k$ -module (cf. Editors' note (27)), we may suppose, without loss of generality, that  $M$  is topologically free. Thus write

$$M = \prod_{i \in I} M_i,$$

where each  $M_i$  is a copy of  $k$ . In this case,

$$A \widehat{\otimes}_k M = \prod_{i \in I} A \widehat{\otimes}_k M_i.$$

Since the map

$$a \widehat{\otimes} m \longmapsto a \cdot i_0(m)$$

is surjective and  $A_1$  is projective, its kernel is a direct factor of  $A \widehat{\otimes}_k M$ . As  $A$  is local, the exchange theorem (0.3.4) shows that this kernel has as a complement a partial product

$$\prod_{i \in J} A \widehat{\otimes}_k M_i,$$

where  $J$  is a subset of  $I$ . One may therefore take

$$V = \prod_{i \in J} M_i.$$

## 1.5

Let  $k$  be a pseudocompact ring.

**Definition.** We shall say that a family of morphisms

$$f_i : X_i \longrightarrow X$$

in  $\mathbf{Vaf}/k$  is a *surjective topologically flat family* if the morphism

$$\coprod_i X_i \longrightarrow X$$

induced by the  $f_i$  is surjective and topologically flat. This means that every  $f_i$  is topologically flat and that every point of  $X$  lies in the image of at least one  $X_i$ .

It follows from 1.3.3 that surjective topologically flat families define a pretopology on  $\mathbf{Vaf}/k$  (IV, 4.2.5). The corresponding topology will be called the *flat topology* on  $\mathbf{Vaf}/k$ .

By IV, 4.3.5, a functor

$$F : (\mathbf{Vaf}/k)^{\text{op}} \longrightarrow (\mathbf{Sets})$$

is a sheaf for the flat topology if and only if  $F$  carries every direct sum to a direct product and the sequence

$$F(Y) \xrightarrow{F(f)} F(X) \xrightleftharpoons[F(\text{pr}_2)]{F(\text{pr}_1)} F(X \times_Y X)$$

is exact for every surjective topologically flat morphism  $f : X \rightarrow Y$ .

By IV, 4.5,<sup>71</sup> Proposition 1.3.1 implies that the flat topology is coarser than the canonical topology. In other words, for every object  $X$  of  $\mathbf{Vaf}/k$ , the functor

$$h_X : T \longmapsto \text{Hom}_{\mathbf{Vaf}/k}(T, X)$$

---

<sup>71</sup>The editors modified the original in what follows. In particular, they replaced the statement: “if  $d_0, d_1 : X_1 \rightrightarrows X$  is an equivalence relation such that  $d_1$  is topologically flat, formation of the quotient commutes with  $h$ ,” by the theorem below.

is a sheaf for the flat topology. In what follows, as usual (cf. Exposé I), we identify  $X$  with  $h_X$ .

By IV, 4.6.5, Theorem 1.4 may be reformulated as follows.

**Theorem.** *Let  $k$  be a pseudocompact ring, let  $d_0, d_1 : X_1 \rightrightarrows X$  be an equivalence pair in  $\mathbf{Vaf}/k$ , and let  $X/X_1$  be the quotient formal variety (that is,  $\mathrm{Coker}(d_0, d_1)$ , cf. 1.2). If  $d_1$  is topologically flat, then  $X/X_1$  represents the quotient sheaf for the flat topology.*

## 1.6

To complete these generalities on formal varieties, it remains to define briefly the formal varieties that are étale over  $k$ .<sup>72</sup>

**Recollections 1.6.A.** (i) Recall first (cf. EGA 0<sub>IV</sub>, 19.10.2) that a topological  $k$ -algebra  $A$  is said to be *formally étale over  $k$* —for the given topologies on  $k$  and  $A$ , respectively for the discrete topologies—if, for every discrete topological  $k$ -algebra  $C$  (not necessarily Artinian) and every nilpotent ideal  $J$  of  $C$ , every continuous morphism of  $k$ -algebras

$$A \longrightarrow C/J$$

lifts uniquely to a continuous morphism  $A \rightarrow C$ , where  $A$  is endowed with the given topology, respectively with the discrete topology.

This property is plainly preserved under base change. Thus, for every morphism  $k \rightarrow k'$  of pseudocompact rings,

$$A \widehat{\otimes}_k k'$$

is formally étale over  $k'$ . It is also enough to verify the lifting condition for square-zero ideals  $J$ ; cf. EGA IV<sub>4</sub>, 17.1.2 (ii).

The algebra  $A$  is said to be *étale over  $k$*  if it is formally étale over  $k$  for the discrete topologies and is, in addition, a  $k$ -algebra of finite presentation; cf. EGA IV<sub>4</sub>, 17.3.2 (ii). In what follows, when  $k$  is a pseudocompact ring and  $A$  a profinite  $k$ -algebra, “formally étale” refers to the given topologies unless the contrary is stated.

(ii) Recall also that if  $F \in k[T]$  is a separable monic polynomial of degree  $d \geq 1$ , meaning that the ideal generated by  $F$  and its derivative  $F'$  is all of  $k[T]$ , then

$$B = k[T]/(F),$$

which is free of rank  $d$  over  $k$  and is endowed with the product topology, is formally étale over  $k$ .

Indeed, let  $C$  be a discrete  $k$ -algebra, so that the kernel  $\mathfrak{l}$  of  $k \rightarrow C$  is an open ideal of  $k$ ; let  $J$  be a square-zero ideal of  $C$ ; and let  $\phi : B \rightarrow C/J$  be a continuous morphism of  $k$ -algebras. Since  $B$  is a free  $k$ -module of finite rank,  $\mathfrak{l}B$  is an open ideal of  $B$ , and hence every lift  $\Phi : B \rightarrow C$  of  $\phi$  is automatically continuous.

Let  $t$  be the image of  $T$  in  $B$ , and choose any lift  $u_0 \in C$  of  $\phi(t)$ . Then  $F(u_0) \in J$ , since  $\phi(t)$  is a root of  $F$ . There exist  $G, H \in k[T]$  such that

$$GF + HF' = 1.$$

<sup>72</sup>The original continued as follows: “A formal variety  $X$  over  $k$  is said to be étale if the diagonal morphism  $\Delta_X : X \rightarrow X \times X$  is a local isomorphism; that is, if  $\Delta_X$  induces an isomorphism from  $\mathcal{O}_{X \times X, \Delta_X(x)}$  onto  $\mathcal{O}_{X, x}$  for every point  $x$  of  $X$ . It follows readily from SGA I that this formulation is equivalent to each of the following two: the formal variety  $X$  is topologically flat and, for every point  $x \in X$  projecting to  $s \in \mathrm{Spf}(k)$ ,  $\mathcal{O}_{X, x} \widehat{\otimes}_k \kappa(s)$  is a finite separable extension of the residue field  $\kappa(s)$ ; or, if  $A$  denotes the affine algebra of  $X$ , the local components (0.1) of  $A \widehat{\otimes}_k (k/\mathfrak{l})$  are finite étale algebras over  $k/\mathfrak{l}$ , for every open ideal  $\mathfrak{l}$  of  $k$ .” In what follows, the editors corrected the omission of the flatness hypothesis from the first condition above and supplied details for the equivalence of the conditions.

Consequently,

$$H(u_0)F'(u_0) = 1 - G(u_0)F(u_0).$$

The right-hand side is invertible because  $F(u_0)$  is square-zero, and therefore  $F'(u_0)$  is invertible. We seek  $h \in J$  such that  $u = u_0 + h$  is a root of  $F$ . This is equivalent to

$$0 = F(u) = F(u_0) + F'(u_0)h,$$

whose unique solution is

$$h = -F'(u_0)^{-1}F(u_0) \in J.$$

The same proof, with no continuity assumptions on the morphisms  $k \rightarrow C$ ,  $\phi$ , and  $\Phi$ , also shows that  $B$  is an étale  $k$ -algebra.

(iii) Finally, if

$$A = A_1 \times \cdots \times A_n$$

is a finite product, then  $A$  is formally étale over  $k$  if and only if every  $A_i$  is.<sup>73</sup> It is enough to prove this for  $n = 2$ . Let  $e = 1_{A_1}$  be the idempotent such that  $A_1 = Ae$  and  $A_2 = A(1 - e)$ , and suppose that a continuous morphism

$$\phi : A \longrightarrow C/J$$

is given, where  $J$  is square-zero. The polynomial  $F = X^2 - X$  is separable, since

$$F' = 2X - 1 \quad \text{and} \quad (F')^2 - 4F = 1.$$

Therefore the idempotent  $\phi(e)$  of  $C/J$  lifts uniquely to an idempotent  $f$  of  $C$ . Hence

$$C = Cf \oplus C(1 - f),$$

and giving a lift  $\Phi$  is equivalent to giving two morphisms

$$\Phi_1 : A_1 \longrightarrow Cf, \quad \Phi_2 : A_2 \longrightarrow C(1 - f),$$

lifting the respective restrictions of  $\phi$ . The same argument shows that if  $e$  is an idempotent of  $k$  and  $A = Ae$ , then  $A$  is formally étale over  $k$  if and only if it is formally étale over the local component  $k_e$ , identified with  $ke$ .

Now let  $X$  be a formal  $k$ -variety with affine algebra  $A$ . If  $x$  is a point of  $X$ , that is, an open maximal ideal  $\mathfrak{m}$  of  $A$ , and  $s$  is its image in  $\mathrm{Spf}(k)$ , we denote by  $k_{\mathfrak{m}}$ , or  $k_s$ , the local component of  $k$  corresponding to  $s$ .

**Definition 1.6.B.** *The following conditions are equivalent:*

- (a)  $A$  is formally étale over  $k$ .
- (b) For every  $\mathfrak{m} \in \Upsilon(A)$ ,  $A_{\mathfrak{m}}$  is formally étale over  $k$ , or equivalently over  $k_{\mathfrak{m}}$ .
- (c) For every open ideal  $\mathfrak{l}$  of  $k$ ,

$$A \widehat{\otimes}_k (k/\mathfrak{l}) = A/\mathfrak{l}A$$

is formally étale over  $k/\mathfrak{l}$ .

- (d) For every  $\mathfrak{m} \in \Upsilon(A)$  and every open ideal  $\mathfrak{l}$  of  $k_{\mathfrak{m}}$ ,  $A_{\mathfrak{m}}/A_{\mathfrak{m}}\mathfrak{l}$  is formally étale over  $k_{\mathfrak{m}}/\mathfrak{l}$ .

<sup>73</sup>The editors point out that the proof given in EGA 0<sub>IV</sub>, 19.3.5 (v), is erroneous.

We say that  $X$  is *étale over  $k$*  if these conditions hold, and write

$$\mathbf{Vaf}^{\text{ét}}/k$$

for the full subcategory of  $\mathbf{Vaf}/k$  consisting of formal varieties that are étale over  $k$ .<sup>74</sup>

Indeed, suppose that

$$\phi : A \longrightarrow C/J$$

is a continuous morphism of  $k$ -algebras, where  $C$  is a discrete  $k$ -algebra and  $J$  is square-zero. Then  $I = \text{Ker}(\phi)$  is an open ideal of  $A$ , so  $A/I$  is Artinian. Thus  $I$  is contained in only finitely many open maximal ideals

$$\mathfrak{m}_1, \dots, \mathfrak{m}_r.$$

It therefore contains the product of the components  $A_{\mathfrak{m}}$  for  $\mathfrak{m} \notin \{\mathfrak{m}_1, \dots, \mathfrak{m}_r\}$ . This product is  $A(1 - e)$ , where  $e$  is the idempotent of  $A$  for which

$$Ae = \prod_{i=1}^r A_{\mathfrak{m}_i}.$$

Hence  $\phi(e) = 1_{C/J}$ , and giving a continuous lift of  $\phi$  is equivalent to giving a continuous lift of the induced morphism

$$Ae \simeq A/A(1 - e) \longrightarrow C/J.$$

On the other hand, by editors' note (24),

$$A \simeq \varprojlim_{\mathfrak{l}} A/\mathfrak{l}.$$

The equivalence of the stated conditions follows readily from these observations and the preceding recollections.

**Definitions 1.6.C.** Put

$$\kappa(k) = \prod_{s \in \text{Spf}(k)} \kappa(s),$$

endowed with the product topology. Thus the formal variety  $\text{Spf}(\kappa(k))$  is the direct sum of the schemes  $\text{Spec } \kappa(s)$ , for  $s \in \text{Spf}(k)$ . We write  $S_{\kappa(k)}$  for the scheme that is the direct sum of the same  $\text{Spec } \kappa(s)$ .

For every formal variety  $X$  over  $k$ , put

$$X_{\kappa} = X \widehat{\otimes}_k \kappa(k).$$

This is the formal variety over  $\kappa(k)$  obtained by base change. It has the same points as  $X$ , and if  $x \in X$  projects to  $s \in \text{Spf}(k)$ , then

$$\mathcal{O}_{X_{\kappa}, x} = \mathcal{O}_{X, x} \widehat{\otimes}_k \kappa(s).$$

The base-change functor

$$\mathbf{Vaf}/k \longrightarrow \mathbf{Vaf}/\kappa(k)$$

sends  $\mathbf{Vaf}^{\text{ét}}/k$  into  $\mathbf{Vaf}^{\text{ét}}/\kappa(k)$ , by 1.6.A (i).

One then has the following result (cf. SGA 1, I 6.2).

**Lemma 1.6.D.** *For every*

$$Y \in \text{Ob}(\mathbf{Vaf}^{\text{ét}}/k) \quad \text{and} \quad X \in \text{Ob}(\mathbf{Vaf}/k),$$

<sup>74</sup>The category  $\mathbf{Vaf}^{\text{ét}}/k$  is stable under finite projective limits.

the canonical map

$$\mathrm{Hom}_{\mathrm{Vaf}/k}(X, Y) \longrightarrow \mathrm{Hom}_{\mathrm{Vaf}/\kappa(k)}(X_\kappa, Y_\kappa)$$

is bijective. In particular, the functor

$$\mathrm{Vaf}^{\mathrm{\acute{e}t}}/k \longrightarrow \mathrm{Vaf}^{\mathrm{\acute{e}t}}/\kappa(k)$$

is fully faithful; below it will be shown to be an equivalence.

Indeed, let  $A$ , respectively  $B$ , be the affine algebra of  $X$ , respectively  $Y$ , and let  $r$  be the radical of  $k$ . Suppose that a morphism

$$B\widehat{\otimes}_k \kappa(k) \longrightarrow A\widehat{\otimes}_k \kappa(k)$$

is given, or equivalently a morphism

$$\phi : B \longrightarrow A/rA.$$

For every open ideal  $\mathfrak{l}$  of  $k$ , there is an integer  $n \geq 1$  such that

$$r^n \subset \mathfrak{l}.$$

Because the multiplication map

$$m^{n-1} : A^n \longrightarrow A$$

is continuous, it follows that

$$(rA)^n \subset \mathfrak{l}A.$$

Thus  $rA/\mathfrak{l}A$  is a nilpotent ideal of  $A/\mathfrak{l}A$ . Consequently,  $\phi$  lifts uniquely to a morphism

$$\phi_{\mathfrak{l}} : B \longrightarrow A/\mathfrak{l}A.$$

By uniqueness, these morphisms form a projective system and therefore give a continuous morphism

$$\Phi : B \longrightarrow \varprojlim_{\mathfrak{l}} A/\mathfrak{l}A = A.$$

Moreover,  $\Phi$  is unique: if  $\Phi'$  is another lift of  $\phi$ , then  $\Phi'$  and  $\Phi$  agree modulo  $\mathfrak{l}A$  for every  $\mathfrak{l}$ , and hence are equal.

**Proposition 1.6.E.** (a) Let  $X$  be a formal variety over  $k$ , with affine algebra  $A$ . The following conditions are equivalent:

- (i)  $X$  is étale over  $k$ .
- (ii)  $X$  is topologically flat over  $k$ , and the diagonal morphism

$$\Delta : X \longrightarrow X \times X$$

is a local isomorphism; that is, for every point  $x$  of  $X$ ,  $\Delta_X$  induces an isomorphism

$$\mathcal{O}_{X \times X, \Delta(x)} \xrightarrow{\sim} \mathcal{O}_{X, x}.$$

- (iii) For every  $x \in X$  projecting to  $s \in \mathrm{Spf}(k)$ , there is an isomorphism

$$\mathcal{O}_{X, x} \simeq k_s[T]/(F),$$

where  $F \in k_s[T]$  is a separable monic polynomial (cf. 1.6.A (ii)).

(iv)  $X$  is topologically flat over  $k$ , and, for every  $x \in X$  projecting to  $s \in \mathrm{Spf}(k)$ ,

$$\mathcal{O}_{X,x} \widehat{\otimes}_k \kappa(s)$$

is a finite separable extension of  $\kappa(s)$ .

(v) For every open ideal  $\mathfrak{l}$  of  $k$ , each local component of

$$A \widehat{\otimes}_k (k/\mathfrak{l})$$

is a finite étale algebra over the Artinian ring  $k/\mathfrak{l}$ .

(b) The category  $\mathrm{Vaf}^{\mathrm{ét}}/\kappa(k)$  identifies with the category of étale schemes over  $S_{\kappa(k)}$  (cf. 1.6.C), and the functor  $X \mapsto X_{\kappa}$  induces an equivalence

$$\mathrm{Vaf}^{\mathrm{ét}}/k \xrightarrow{\sim} \mathrm{Vaf}^{\mathrm{ét}}/\kappa(k)$$

(cf. SGA 1, I 6.1).

*Proof.* Let  $I$  be the kernel of the multiplication morphism

$$m : A \widehat{\otimes}_k A \longrightarrow A.$$

Suppose first that  $X$  is étale over  $k$ , so that  $A$  is formally étale over  $k$ . By EGA 0<sub>IV</sub>, 20.7.4, the separated completion of  $I/I^2$ , for the quotient topology induced from  $I$ , is zero; hence

$$I = I^2.$$

For every  $x \in X$ , the ideal  $I$  is contained in the maximal ideal  $\mathfrak{m}_{\Delta(x)}$  of  $A \widehat{\otimes}_k A$ . Thus its localization  $I_{\mathfrak{m}_{\Delta(x)}}$  lies in the maximal ideal of  $\mathcal{O}_{X \times X, \Delta(x)}$ . Nakayama's lemma 0.3 therefore gives

$$I_{\mathfrak{m}_{\Delta(x)}} = 0.$$

Consequently,  $\Delta : X \rightarrow X \times X$  is a local isomorphism.

Suppose next that  $\Delta$  is a local isomorphism and that  $k$  is a field  $\kappa$ . We show that every  $\mathcal{O}_{X,x}$  is a finite-dimensional étale  $\kappa$ -algebra. Replacing  $X$  by  $\mathrm{Spf}(\mathcal{O}_{X,x})$ , we may assume that

$$A = \mathcal{O}_{X,x}$$

is local. We proceed as in the proof of EGA IV<sub>4</sub>, 17.4.1, (b) $\Rightarrow$ (d"). Let  $K$  be a finite normal extension of  $\kappa$  containing the residue field  $\kappa(x)$ , and put

$$B = A \otimes_{\kappa} K = A \widehat{\otimes}_{\kappa} K, \quad X_K = \mathrm{Spf}(B) = X \widehat{\otimes}_{\kappa} K.$$

The two tensor products agree because  $[K : \kappa] < \infty$ .

The morphism

$$\Delta_K : X_K \longrightarrow X_K \times_K X_K$$

is again a local isomorphism. Hence, for every  $y \in X_K$ , multiplication induces an isomorphism

$$(B_y \widehat{\otimes}_K B_y)_{\mathfrak{m}_{\Delta_K(y)}} \xrightarrow{\sim} B_y.$$

The residue field of  $B_y$  is  $K$ ; cf., for example, VI<sub>A</sub>, 1.1.1, editors' note (11). Moreover,

$$C = B_y \widehat{\otimes}_K B_y$$

is already local. Indeed,

$$\mathfrak{n} = \mathfrak{m}_y \widehat{\otimes}_K B_y + B_y \widehat{\otimes}_K \mathfrak{m}_y$$

consists of topologically nilpotent elements and therefore lies in the radical of  $C$ , while

$$C/\mathfrak{n} = K \widehat{\otimes}_K K = K$$

is a field. It follows that multiplication itself is an isomorphism

$$B_y \widehat{\otimes}_K B_y \xrightarrow{\sim} B_y.$$

Choosing a pseudobasis of  $B_y$  over  $K$  that contains the unit element 1, we deduce that  $B_y = K$ . Since  $B$  is finite over  $A$ , the set  $X_K$  is finite. Thus

$$B = A \otimes_{\kappa} K$$

is a product of finitely many copies of  $K$ , and  $A$  is a finite étale  $\kappa$ -algebra.

We have therefore shown that, over a field  $\kappa$ , every profinite  $\kappa$ -algebra  $A$  that is étale over  $\kappa$  is a product of finite separable extensions  $K_i/\kappa$ , endowed with the product topology. Consequently,  $\mathrm{Spf}(A)$  is the direct sum of the schemes

$$\mathrm{Spf}(K_i) = \mathrm{Spec}(K_i),$$

and  $\mathrm{Vaf}^{\mathrm{ét}}/\kappa$  identifies with the category of étale  $\kappa$ -schemes.

The preceding argument proves (ii)  $\Rightarrow$  (iv), because  $\mathcal{O}_{X,x} \widehat{\otimes}_k \kappa(s)$  is formally étale over  $\kappa(s)$ , and also proves part (b). Indeed, return to an arbitrary pseudocompact ring  $k$ . The category  $\mathrm{Vaf}^{\mathrm{ét}}/\kappa(k)$  has just been identified with the category of étale schemes over  $S_{\kappa(k)}$ . By 1.6.D, to show that

$$\mathrm{Vaf}^{\mathrm{ét}}/k \longrightarrow \mathrm{Vaf}^{\mathrm{ét}}/\kappa(k)$$

is an equivalence, it remains only to prove essential surjectivity.

Let  $s \in \mathrm{Spf}(k)$ , and let  $K/\kappa(s)$  be a finite separable extension. Choose a primitive element  $\xi \in K$ , let  $n = [K : \kappa(s)]$ , and choose a monic polynomial

$$F \in k_s[T]$$

of degree  $n$  whose image  $\overline{F} \in \kappa(s)[T]$  is the minimal polynomial of  $\xi$ . Since  $\overline{F}'$  is invertible in  $\kappa(s)[T]/(\overline{F})$ , Nakayama's lemma shows that  $F'$  is invertible in  $k_s[T]/(F)$ . Thus  $F$  is separable, and 1.6.A (ii) shows that

$$A = k_s[T]/(F)$$

is an étale  $k_s$ -algebra satisfying

$$A \widehat{\otimes}_k \kappa(s) \simeq K.$$

It follows that every local profinite étale  $k_s$ -algebra is free of finite rank over  $k_s$ , and therefore is topologically free over  $k$ . Hence every profinite étale  $k$ -algebra is topologically free over  $k$ .

Condition (v) implies condition (d) of 1.6.B, and therefore implies (i). We have thus proved that (i), (iii), and (v) are equivalent and imply (ii), which in turn implies (iv).

Finally, let  $A$  be a profinite  $k$ -algebra satisfying (iv). We show that  $A$  is formally étale over  $k$ . We may suppose that  $A$  and  $k$  are local, and write  $\kappa$  for the residue field of  $k$ . By hypothesis,

$$K = A \widehat{\otimes}_k \kappa$$



is a finite separable extension of  $\kappa$ , say of degree  $n$ . By the preceding argument and Lemma 1.6.D, there is a  $k$ -algebra  $B$ , free of rank  $n$  and formally étale over  $k$ , together with a continuous morphism

$$\phi : B \longrightarrow A$$

such that

$$\phi \hat{\otimes}_k \kappa$$

is an isomorphism. Since  $A$  is topologically flat over  $k$ , this implies

$$\mathrm{Coker}(\phi) \hat{\otimes}_k \kappa = 0 \quad \text{and} \quad \mathrm{Ker}(\phi) \hat{\otimes}_k \kappa = 0.$$

Nakayama's lemma 0.3 gives

$$\mathrm{Coker}(\phi) = 0 = \mathrm{Ker}(\phi).$$

Thus  $\phi$  is an isomorphism (cf. the proof of 0.2.B), which completes the proof of 1.6.E.

Let  $X$  be a formal variety over  $k$ . If  $x \in X$  projects to  $s \in \mathrm{Spf}(k)$ , then the residue field  $\kappa(x)$  is a finite extension of  $\kappa(s)$ . We denote by  $\kappa_e(x)$  the separable closure of  $\kappa(s)$  in  $\kappa(x)$ .<sup>75</sup>

**Proposition 1.6.F.** (i) *The inclusion*

$$\mathrm{Vaf}^{\mathrm{ét}}/k \longrightarrow \mathrm{Vaf}/k$$

has a left adjoint  $X \mapsto X_e$ . The formal variety  $X_e$  has the same points as  $X$ . For each  $x \in X$  projecting to  $s \in \mathrm{Spf}(k)$ , choose a primitive element  $\xi$  of  $\kappa_e(x)/\kappa(s)$ , let  $n$  be its degree, choose any lift  $u \in \mathcal{O}_{X,x}$  of  $\xi$ , and let  $F \in k[T]$  be a monic polynomial of degree  $n$  that annihilates  $u$ . Then put

$$\mathcal{O}_{X_e,x} = k[T]/(F).$$

For every  $Y \in \mathrm{Ob}(\mathrm{Vaf}^{\mathrm{ét}}/k)$ , the canonical maps in the following diagram are bijective:

$$\begin{array}{ccc} \mathrm{Hom}_{\mathrm{Vaf}/k}(X, Y) & \xrightarrow{\sim} & \mathrm{Hom}_{\mathrm{Vaf}/\kappa(k)}(X_\kappa, Y_\kappa) \\ & & \downarrow \wr \\ \mathrm{Hom}_{\mathrm{Vaf}/k}(X_e, Y) & \xrightarrow{\sim} & \mathrm{Hom}_{\mathrm{Vaf}/\kappa(k)}((X_e)_\kappa, Y_\kappa), \end{array}$$

The vertical arrow is induced by the inclusions

$$\kappa_e(x) \hookrightarrow \kappa(x), \quad x \in X.$$

In particular, this defines a morphism

$$p : X \longrightarrow X_e,$$

and every morphism of formal  $k$ -varieties  $\phi : X \rightarrow Y$ , with  $Y$  étale over  $k$ , factors uniquely through  $p$ .

(ii) *The functor  $X \mapsto X_e$  commutes with finite products.*

*Proof.* Assertion (i) follows from 1.6.E (b) and 1.6.D. We prove (ii). By the equivalence of categories in 1.6.E (b), we may suppose that  $k$  is a field. In that case, if  $X$  is a semilocal formal  $k$ -variety, so that its affine algebra  $A$  is semilocal, the affine algebra of  $X_e$  is the largest subalgebra of  $A$  that is étale over  $k$ ; denote it by  $A_e$ .

<sup>75</sup>In what follows, the editors used the preceding additions to give details of the construction of the functor  $X \mapsto X_e$  and to prove that it commutes with finite products; this is used in 2.5.1.

It is therefore enough to show that, if  $K$  and  $L$  are two finite extensions of  $k$ , then

$$K_e \otimes_k L_e = (K \otimes_k L)_e.$$

Let  $p$  be the characteristic exponent of  $k$ , so that  $p = 1$  if  $\text{char}(k) = 0$ , and  $p = \text{char}(k)$  otherwise. For every  $x \in K \otimes_k L$ , there is an integer  $n \geq 1$  such that

$$x^{p^n} \in K_e \otimes_k L_e.$$

Thus  $K \otimes_k L$  is radicial over  $K_e \otimes_k L_e$ , and its largest étale  $k$ -subalgebra is precisely  $K_e \otimes_k L_e$ . Hence

$$K_e \otimes_k L_e = (K \otimes_k L)_e.$$

Keeping this notation, observe that  $\mathcal{O}_{X_e, x}$  is not necessarily a subalgebra of  $\mathcal{O}_{X, x}$ . It is a subalgebra when  $X$  is topologically flat over  $k$ , by the proposition that follows.

### 1.6.1

Let  $Y$  be an étale formal  $k$ -variety and let  $f : X \rightarrow Y$  be a morphism in  $\text{Vaf}/k$ . Then the following square is Cartesian, where  $\Gamma_f$  is the graph morphism  $X \rightarrow X \times Y$ , with components  $\text{id}_X$  and  $f$ :

$$\begin{array}{ccc} X & \xrightarrow{\Gamma_f} & X \times Y \\ f \downarrow & & \downarrow f \times \text{id}_Y \\ Y & \xrightarrow{\Delta_Y} & Y \times Y. \end{array}$$

It follows that  $\Gamma_f$  is a local isomorphism, and hence that

$$f = \text{pr}_Y \circ \Gamma_f$$

is topologically flat if  $\text{pr}_Y$  is; this holds, for example, if  $X$  is topologically flat over  $k$ .

Conversely, since  $Y \rightarrow k$  is topologically flat,  $X \rightarrow k$  will also be topologically flat if  $f$  is (cf. 1.3.3). Taking in particular for  $f$  the canonical morphism

$$p : X \rightarrow X_e$$

of 1.6, we obtain the following.

**Proposition.** *Let  $X$  be a formal variety over  $k$ . The morphism  $X \rightarrow X_e$  is topologically flat if and only if  $X$  is topologically flat over  $k$ .*

**Remark 1.6.2.**<sup>76</sup> When  $k$  is a perfect field, the functor

$$X \mapsto X_e$$

is also right adjoint to the inclusion

$$\text{Vaf}^{\text{ét}}/k \hookrightarrow \text{Vaf}/k.$$

Indeed, in this case

$$\mathcal{O}_{X_e, x} = \kappa(x)$$

for every  $x \in X$ , and the canonical projections

$$\mathcal{O}_{X, x} \rightarrow \kappa(x)$$

define a morphism  $s : X_e \rightarrow X$ , which is a section of  $p : X \rightarrow X_e$ . For every morphism  $f : X \rightarrow Y$ , the following diagrams are commutative:

<sup>76</sup>The editors inserted this remark here; in the original it appeared in 2.5.2.

$$\begin{array}{ccc}
\mathcal{O}_{Y,f(x)} & \longrightarrow & \mathcal{O}_{X,x} \\
\downarrow & & \downarrow \\
\kappa(f(x)) & \longrightarrow & \kappa(x)
\end{array}
\qquad
\begin{array}{ccc}
X & \xrightarrow{f} & Y \\
s_X \uparrow & & \uparrow s_Y \\
X_e & \xrightarrow{f_e} & Y_e
\end{array}$$

Thus  $s$  is functorial in  $X$ , and it follows that  $X \mapsto X_e$  is right adjoint to the inclusion

$$\mathrm{Vaf}^{\mathrm{\acute{e}t}}/k \hookrightarrow \mathrm{Vaf}/k.$$

Consequently, being a right adjoint, the functor  $X \mapsto X_e$  commutes with projective limits whenever they exist in  $\mathrm{Vaf}^{\mathrm{\acute{e}t}}/k$ ,<sup>77</sup> and hence, in particular, with finite products.

(This can also be verified directly: for every formal  $k$ -variety  $X$ , with affine algebra  $A$ , the affine algebra of  $X_e$  is the quotient of  $A$  by its radical  $r(A)$ ; and, since  $k$  is perfect, the quotient of  $\mathcal{O}_{X,x} \hat{\otimes} \mathcal{O}_{Y,y}$  by its radical is the algebra  $\kappa(x) \otimes_k \kappa(y)$ , because the latter is semisimple.)

## 2. Generalities on Formal Groups

### 2.1

Let  $k$  be a pseudocompact ring and let  $G$  be a formal  $k$ -group, that is, a group object in the category  $\mathrm{Vaf}/k$  of formal varieties over  $k$ . Let  $A$  be the affine algebra of  $G$ . The composition law of  $G$  plainly defines a diagonal morphism, namely a homomorphism of profinite  $k$ -algebras

$$\Delta_A : A \longrightarrow A \hat{\otimes}_k A.$$

This homomorphism has the following properties.

- (i) The diagram

$$\begin{array}{ccc}
A & \xrightarrow{\Delta_A} & A \hat{\otimes}_k A \\
\Delta_A \downarrow & & \downarrow \Delta_A \hat{\otimes} \mathrm{id}_A \\
A \hat{\otimes}_k A & \xrightarrow{\mathrm{id}_A \hat{\otimes} \Delta_A} & A \hat{\otimes}_k A \hat{\otimes}_k A
\end{array}$$

is commutative.

- (ii) There is a necessarily unique *augmentation*, that is, a homomorphism of profinite  $k$ -algebras

$$\varepsilon_A : A \longrightarrow k,$$

such that the two composites displayed below are the identity maps of  $A$ :

$$\begin{aligned}
A &\xrightarrow{\Delta_A} A \hat{\otimes}_k A \xrightarrow{\varepsilon_A \hat{\otimes} \mathrm{id}_A} k \hat{\otimes}_k A \simeq A \\
\text{and} \quad A &\xrightarrow{\Delta_A} A \hat{\otimes}_k A \xrightarrow{\mathrm{id}_A \hat{\otimes} \varepsilon_A} A \hat{\otimes}_k k \simeq A
\end{aligned}$$

- (iii) There is a necessarily unique *antipode*, that is, a homomorphism of profinite  $k$ -algebras

$$c_A : A \longrightarrow A,$$

such that the composite

---

<sup>77</sup>This is the case for finite projective limits.

$$A \xrightarrow{\Delta_A} A \widehat{\otimes}_k A \xrightarrow{c_A \widehat{\otimes} \text{id}_A} A \widehat{\otimes}_k A \xrightarrow{m_A} A$$

equals  $\eta_A \circ \varepsilon_A$ . Here  $m_A$  is the linear map that sends  $a \widehat{\otimes} b$  to  $ab$ , and  $\eta_A : k \rightarrow A$  sends  $\lambda$  to  $\lambda 1_A$ .

Conversely, data

$$(\Delta_A, \varepsilon_A, c_A)$$

satisfying (i)–(iii) endow  $G$  with the structure of a formal  $k$ -group.<sup>78</sup> Explicitly, for every profinite  $k$ -algebra  $B$ , the set

$$\text{Hom}_c(A, B)$$

of continuous morphisms of  $k$ -modules  $\phi : A \rightarrow B$  carries a group structure, functorial in  $B$ , defined by

$$\phi \cdot \phi' = m_B \circ (\phi \widehat{\otimes} \phi') \circ \Delta_A.$$

The identity element is  $\eta_B \circ \varepsilon_A$ , where  $m_B$  is the multiplication of  $B$  and  $\eta_B : k \rightarrow B$  sends  $\lambda$  to  $\lambda 1_B$ ; the inverse of  $\phi$  is  $\phi \circ c_A$ . The set

$$\text{Hom}_{\text{Alp}/k}(A, B)$$

of continuous morphisms of  $k$ -algebras  $A \rightarrow B$  is a subgroup, because  $B$  is commutative.

**Definition.** A *morphism of formal  $k$ -groups*  $\theta : K \rightarrow G$  is, by definition, a morphism of formal  $k$ -varieties that respects the group structures. Let  $B$ , respectively  $A$ , be the affine algebra of  $K$ , respectively  $G$ , and let  $f : A \rightarrow B$  be the morphism corresponding to  $\theta$ . The condition is equivalently that  $f$  be compatible with comultiplication:

$$(f \widehat{\otimes} f) \circ \Delta_A = \Delta_B \circ f.$$

The identities

$$\varepsilon_B \circ f = \varepsilon_A, \quad c_B \circ f = f \circ c_A$$

then follow automatically. We write  $\text{Grf}/k$  for the category of formal  $k$ -groups.

**Notation.** In what follows, the ideal

$$I_A = \text{Ker}(\varepsilon_A)$$

will be called the *augmentation ideal* of  $A$ , and we write

$$\omega_{G/k} = I_A / \overline{I_A^2}$$

for the pseudocompact  $k$ -module obtained by quotienting  $I_A$  by the closed ideal generated by the products  $xy$ , with  $x, y \in I_A$ .

<sup>78</sup>One also says that  $A$  is a *cogroup* in the category of profinite  $k$ -algebras. The editors expanded the material that follows, in particular by making explicit what a morphism of formal groups  $K \rightarrow G$  means; cf. Proposition 2.3.1.

## 2.2

Let  $\mathbf{H}$  be a group object in the category of coalgebras over  $\mathbf{O}_k$ . Thus, for every object  $C$  of  $\mathbf{Alf}/k$ , the object  $\mathbf{H}(C)$  carries the structure of a  $C$ -coalgebra in groups (cf. VII<sub>A</sub> 3.2). Following Manin, we shall say *bialgebra*<sup>79</sup> instead of “coalgebra in groups.” Moreover, for every morphism

$$\phi : C \longrightarrow D$$

in  $\mathbf{Alf}/k$ , the map

$$D \otimes_C \mathbf{H}(C) \longrightarrow \mathbf{H}(D)$$

is a homomorphism of  $D$ -bialgebras.

**Definition.** These properties will be summarized by saying that  $\mathbf{H}$  is a *bialgebra over  $\mathbf{O}_k$* .

The functor

$$\mathbf{H} \longmapsto \mathrm{Spf}^*(\mathbf{H})$$

of 1.3.5 plainly commutes with finite products. It therefore sends a bialgebra over  $\mathbf{O}_k$  to a  $k$ -group functor, that is, a covariant functor from  $\mathbf{Alf}/k$  to the category of groups.

Indeed, for every  $k$ -algebra  $C$  of finite length, the elements of

$$\begin{aligned} \mathrm{Spf}^*(\mathbf{H})(C) &= \mathrm{Spf}^*(\mathbf{H}(C)) \\ &= \{x \in \mathbf{H}(C) \mid \varepsilon(x) = 1 \text{ and } \Delta(x) = x \otimes x\} \end{aligned}$$

form a group under multiplication in the algebra  $\mathbf{H}(C)$ ; cf. VII<sub>A</sub> 3.2.2. Notice also that  $\Delta(x) = x \otimes x$  implies

$$\varepsilon(x) = \varepsilon(x)^2.$$

Thus  $\varepsilon(x) = 1$  whenever  $C$  is local and  $x \neq 0$ .<sup>80</sup>

### 2.2.1

A bialgebra  $\mathbf{H}$  over  $\mathbf{O}_k$  is said to be *flat* if its underlying module is flat (cf. 1.2.1).<sup>81</sup> If  $\mathbf{H}$  is flat, then, by 1.3.5,

$$A = \Gamma^*(\mathbf{H})$$

is a topologically flat profinite  $k$ -algebra, and  $\mathrm{Spf}^*(\mathbf{H})$ , as a functor from

$$(\mathbf{Alf}/k)^{\mathrm{op}} = \mathbf{Vaf}/k$$

to the category of sets, is isomorphic to the functor

$$\mathrm{Spf}(A) : C \longmapsto \mathrm{Hom}_{\mathbf{Vaf}/k}(\mathrm{Spf}(C), \mathrm{Spf}(A)).$$

The group structure on  $\mathrm{Spf}^*(\mathbf{H})$  therefore endows

$$\mathcal{G}(\mathbf{H}) = \mathrm{Spf}(A)$$

with the structure of a formal group, described explicitly as follows.

<sup>79</sup>The term “bigebra” is also used (cf. [BAlg], III § 11.4). Recall (cf. VII<sub>A</sub>, 3.1, editors’ note (26)) that all bialgebras considered here are assumed to be cocommutative and equipped with an antipode; they are therefore cocommutative Hopf algebras.

<sup>80</sup>Since  $\Delta(x) = x \otimes x$ , one has  $x = \varepsilon(x)x$ ; hence  $\varepsilon(x) = 0$  can occur only when  $x = 0$ .

<sup>81</sup>The editors have expanded the original in what follows.

For every object  $C$  of  $\mathbf{Alf}/k$ , the  $C$ -module underlying  $\mathbf{H}(C)$  is projective. Lemma 1.2.3.A therefore gives, by induction on  $n$ , natural isomorphisms

$$\begin{aligned} (\mathbf{H}(C)^*)^{\widehat{\otimes}(n+1)} &\simeq \mathrm{Hom}_C(\mathbf{H}(C), (\mathbf{H}(C)^*)^{\widehat{\otimes}n}) \\ &\simeq \mathrm{Hom}_C(\mathbf{H}(C), (\mathbf{H}(C)^{\otimes n})^*) \\ &\simeq (\mathbf{H}(C)^{\otimes(n+1)})^*. \end{aligned}$$

For  $n = 1, 2$ , it follows that the  $C$ -algebra structure on  $\mathbf{H}(C)$  endows  $\mathbf{H}(C)^*$  with a diagonal map satisfying conditions 2.1(i)–(iii), functorially in  $C$ .

Consequently,

$$A = \Gamma^*(\mathbf{H}) = \varprojlim_{\mathbf{l}} \mathbf{H}(k/\mathbf{l})^*$$

is endowed with the structure of a cogroup in  $\mathbf{Alp}/k$ , which defines the announced formal-group structure on  $\mathcal{G}(\mathbf{H})$ .

Conversely, let  $G$  be a topologically flat formal  $k$ -group with affine algebra  $A$ , and write

$$\mathbf{H}(G) = \mathbf{V}_k^f(A)$$

for its  $\mathbf{O}_k$ -coalgebra (cf. 1.2.3). The diagonal morphism

$$\Delta_A : A \longrightarrow A \widehat{\otimes}_k A$$

then induces, for every finite-length  $k$ -algebra  $C$ , a  $C$ -linear map

$$\mathbf{V}_k^f(A)(C) \otimes_C \mathbf{V}_k^f(A)(C) \longrightarrow \mathbf{V}_k^f(A)(C),$$

which makes the coalgebra  $\mathbf{V}_k^f(A)(C)$  into a  $C$ -bialgebra. We shall call  $\mathbf{H}(G)$  the *covariant bialgebra* of the formal group  $G$ .<sup>82</sup> Thus, by Proposition 1.3.5.D, we obtain the following.

**Proposition.**

(i) *The functors*

$$G \longmapsto \mathbf{H}(G), \quad \mathbf{H} \longmapsto \mathcal{G}(\mathbf{H})$$

*define an equivalence between the category of topologically flat formal  $k$ -groups and the category of flat  $\mathbf{O}_k$ -bialgebras.*<sup>83</sup>

(ii) *This equivalence “commutes with base change”: if  $k \rightarrow K$  is a morphism of pseudo-compact rings, then*

$$\mathbf{H}(G \widehat{\otimes}_k K) = \mathbf{H}(G) \otimes_k K$$

*and*

$$\mathcal{G}(\mathbf{H} \otimes_k K) = \mathcal{G}(\mathbf{H}) \widehat{\otimes}_k K.$$

When  $k$  is an Artinian ring and  $G$  is a topologically flat formal  $k$ -group, the functor  $\mathbf{H}(G)$  is plainly determined by its value

$$\mathbf{H}(G) = \mathbf{H}(G)(k)$$

<sup>82</sup>The editors added the adjective “covariant” to make clear that the functor  $G \mapsto \mathbf{H}(G)$  is covariant; this terminology is used in [Di73], I § 2.14.

<sup>83</sup>Recall that in this Exposé, “bialgebra” means “cocommutative Hopf algebra.”

at  $k$ . We shall also call  $\mathbf{H}(G)$  the (covariant) bialgebra of  $G$ .<sup>84</sup> Let  $\mathcal{H}_k$  denote the category of  $k$ -Hopf algebras, and let

$$\mathcal{H}_{\text{flat}/k}^{\text{cocom}}$$

be its full subcategory formed by the  $k$ -Hopf algebras that are flat over  $k$  and cocommutative. We then obtain the following.

**Corollary.** *Let  $k$  be an Artinian ring.*

(i) *The functors*

$$G \longmapsto \mathbf{H}(G), \quad \mathbf{H} \longmapsto \mathcal{G}(\mathbf{H}) = \text{Spf}^*(\mathbf{H})$$

*define an equivalence between the category of topologically flat formal  $k$ -groups and  $\mathcal{H}_{\text{flat}/k}^{\text{cocom}}$ .*

(ii) *This equivalence “commutes with base change”: if  $k \rightarrow K$  is a morphism of Artinian rings, then*

$$\mathbf{H}(G \widehat{\otimes}_k K) = \mathbf{H}(G) \otimes_k K$$

*and*

$$\mathcal{G}(\mathbf{H} \otimes_k K) = \mathcal{G}(\mathbf{H}) \widehat{\otimes}_k K.$$

On the other hand, let

$$\mathcal{H}_{\text{flat}/k}^{\text{com}}$$

denote the full subcategory of  $\mathcal{H}_k$  formed by the  $k$ -Hopf algebras that are flat over  $k$  and commutative. Recall that the functor

$$T \longmapsto \mathcal{O}(T)$$

is an anti-equivalence from the category of affine  $k$ -group schemes onto  $\mathcal{H}_{\text{flat}/k}^{\text{com}}$ .

## 2.2.2

Suppose, for simplicity, that  $k$  is Artinian, and let  $G$  be a topologically flat formal  $k$ -group.<sup>85</sup> Then  $G$  is commutative if and only if its affine algebra  $\mathcal{A}(G)$  has a cocommutative comultiplication, which is equivalent to saying that the bialgebra  $\mathbf{H}(G)$  has a commutative multiplication. In that case,  $\mathbf{H}(G)$  is a commutative and cocommutative Hopf algebra, flat over  $k$ . Thus, if

$$D'(G) = \text{Spec } \mathbf{H}(G),$$

then  $D'(G)$  is a commutative  $k$ -group scheme, affine and flat over  $k$ .

Conversely, if  $T$  is such a  $k$ -group scheme, its affine algebra  $\mathcal{O}(T)$  is a commutative group object in the category of cocommutative  $k$ -coalgebras flat over  $k$ . Hence 1.3.5.C gives a topologically flat formal  $k$ -group  $D(T)$ , defined by

$$D(T) = \text{Spf}^* \mathcal{O}(T) = \text{Spf}(\mathcal{A}), \quad \mathcal{A} = \mathcal{O}(T)^*.$$

By 1.3.5.D, there are canonical isomorphisms

$$G = \text{Spf}^* \mathbf{H}(G), \quad \mathbf{H}(D(T)) = \mathcal{O}(T).$$

<sup>84</sup>The editors have introduced here the notation  $\mathcal{H}_{\text{flat}/k}^{\text{cocom}}$ , which will be useful below.

<sup>85</sup>The editors have expanded the original in what follows in order to introduce the notation  $D'(G)$  and  $D(K)$ ; cf. [Ca62], § 14.

We therefore obtain canonical isomorphisms

$$\begin{aligned} D(D'(G)) &= \mathrm{Spf}^* \mathcal{O}(D'(G)) = \mathrm{Spf}^* \mathbf{H}(G) = G, \\ D'(D(T)) &= \mathrm{Spec} \mathbf{H}(D(T)) = \mathrm{Spec} \mathcal{O}(T) = T. \end{aligned}$$

Moreover, if  $k\text{-Gr.}$  denotes the category of  $k$ -group schemes, Corollary 2.2.1 gives functorial isomorphisms

$$\mathrm{Hom}_{\mathrm{Grf}/k}(G, D(T)) \simeq \mathrm{Hom}_{\mathcal{H}_k}(\mathbf{H}(G), \mathcal{O}(T)) \simeq \mathrm{Hom}_{k\text{-Gr.}}(T, D'(G)).$$

This proves the following.

**Proposition (Cartier duality).** *Let  $k$  be an Artinian ring.*

(i) *The functors*

$$G \longmapsto D'(G) = \mathrm{Spec} \mathbf{H}(G), \quad T \longmapsto D(T) = \mathrm{Spf}^* \mathcal{O}(T)$$

*induce an anti-equivalence between the category  $\mathcal{FC}_k$  of commutative topologically flat formal  $k$ -groups and the category  $\mathcal{AC}_k$  of commutative affine flat  $k$ -group schemes. In other words,  $G$  and  $D'(G)$ , respectively  $T$  and  $D(T)$ , are related by the equalities*<sup>86</sup>

$$\mathbf{H}(G) = \mathcal{O}(D'(G)), \quad \mathcal{O}(T) = \mathbf{H}(D(T)).$$

(ii) *This anti-equivalence “commutes with base change”: if  $k \rightarrow K$  is a morphism of Artinian rings, then*

$$D'(G \widehat{\otimes}_k K) = D'(G) \otimes_k K$$

*and*

$$D(T \otimes_k K) = D(T) \widehat{\otimes}_k K.$$

(iii) *In particular, if  $k$  is a field, one obtains an anti-equivalence between the category of commutative formal  $k$ -groups and the category of commutative affine  $k$ -group schemes, and this anti-equivalence commutes with extension of the base field.*

## 2.3

Let  $G$  now be an arbitrary formal  $k$ -group,<sup>87</sup> with affine algebra  $A$ . We continue to write

$$\mathbf{H}(G) = \mathbf{V}_k^f(A)$$

for the  $\mathbf{O}_k$ -module dual to  $A$ , and denote by

$$\phi_G : \mathbf{H}(G) \otimes_k \mathbf{H}(G) \longrightarrow \mathbf{H}(G \times G)$$

the functorial homomorphism induced, for every object  $C$  of  $\mathrm{Alf}/k$ , by the natural map of 0.3.6:

$$\begin{aligned} (A \widehat{\otimes}_k C)^\dagger \otimes_C (A \widehat{\otimes}_k C)^\dagger \\ \longrightarrow ((A \widehat{\otimes}_k C) \widehat{\otimes}_C (A \widehat{\otimes}_k C))^\dagger. \end{aligned}$$

<sup>86</sup>Let  $\mathcal{FC}_k^f$ , respectively  $\mathcal{AC}_k^f$ , denote the full subcategory of  $\mathcal{FC}_k$ , respectively  $\mathcal{AC}_k$ , consisting of the objects  $G$ , respectively  $T$ , for which  $\mathbf{H}(G)$ , respectively  $\mathcal{O}(T)$ , is a finite  $k$ -module (and hence finite locally free). Then  $\mathcal{FC}_k^f$  and  $\mathcal{AC}_k^f$  both have the same objects as the category  $\mathcal{C}$  of commutative and cocommutative Hopf  $k$ -algebras that are finite and flat over  $k$ . The correspondence  $G \mapsto \mathbf{H}(G)$ , respectively  $T \mapsto \mathcal{O}(T)$ , is covariant, respectively contravariant. Thus one recovers the “Cartier duality” of the category  $\mathcal{AC}_k$ , already seen in VII<sub>A</sub> 3.3.1.

<sup>87</sup>That is,  $G$  is not necessarily topologically flat over  $k$ .



If

$$m : G \times G \longrightarrow G$$

is the multiplication of  $G$ , the composite

$$\mathbf{H}(G) \otimes_k \mathbf{H}(G) \xrightarrow{\phi_G} \mathbf{H}(G \times G) \xrightarrow{\mathbf{H}(m)} \mathbf{H}(G)$$

endows  $\mathbf{H}(G)$  with an algebra structure over  $\mathbf{O}_k$ . For every  $C \in \text{Alf}/k$ , the unit element of

$$\mathbf{H}(G)(C) = (A \hat{\otimes}_k C)^\dagger$$

is the augmentation of  $A \hat{\otimes}_k C$ ; cf. 2.1.<sup>88</sup>

If  $G$  is not topologically flat over  $k$ , the morphism  $\phi_G$  need not be an isomorphism. Consequently, the morphism

$$\delta_G : \mathbf{H}(G) \longrightarrow \mathbf{H}(G \times G)$$

induced by the diagonal  $x \mapsto (x, x)$  of  $G$  need not factor through

$$\mathbf{H}(G) \otimes_k \mathbf{H}(G).$$

Thus  $\mathbf{H}(G)$  need not be a bialgebra over  $\mathbf{O}_k$ . For this reason, in the general case we simply call  $\mathbf{H}(G)$  the *covariant algebra* of the formal group  $G$ .

When  $G$  is topologically flat over  $k$ , of course,  $\phi_G$  is an isomorphism, and one recovers the  $\mathbf{O}_k$ -bialgebra structure on  $\mathbf{H}(G)$  defined in 2.2.1.

**Proposition 2.3.1.** *Let  $K$  and  $G$  be two formal  $k$ -groups, with affine algebras  $B$  and  $A$ , respectively. Suppose that  $K$  is topologically flat over  $k$ . Then there is a canonical bijection from*

$$\text{Hom}_{\text{Grf}/k}(K, G)$$

*onto the set of homomorphisms of unital  $\mathbf{O}_k$ -algebras*

$$h : \mathbf{H}(K) \longrightarrow \mathbf{H}(G)$$

*for which the diagram*

$$\begin{array}{ccccc} \mathbf{H}(K) \otimes_k \mathbf{H}(K) & \xrightarrow{h \otimes h} & \mathbf{H}(G) \otimes_k \mathbf{H}(G) & \xrightarrow{\varphi_G} & \mathbf{H}(G \times G) \\ \uparrow \Delta_{\mathbf{H}(K)} & & & \nearrow \delta_G & \\ \mathbf{H}(K) & \xrightarrow{h} & \mathbf{H}(G) & & \end{array}$$

*is commutative.*

Since  $K$  is topologically flat,  $\mathbf{H}(K)$  is endowed with a bialgebra structure (cf. 2.2), and  $\Delta_{\mathbf{H}(K)}$  is its diagonal morphism. In other words, with the notation of 2.3,

$$\Delta_{\mathbf{H}(K)} = \phi_K^{-1} \circ \delta_K.$$

When  $G$  too is topologically flat over  $k$ , the proposition follows from the equivalence of categories established in 2.2.1.

<sup>88</sup>The editors changed the order of the sentences that follow.

In the general case, we may suppose that  $k$  is Artinian and argue with the algebras

$$\mathbf{H}(K) = B^\dagger, \quad \mathbf{H}(G) = A^\dagger.$$

Let  $\text{Hom}_c(A, B)$  denote the set of continuous  $k$ -linear maps from  $A$  to  $B$ , and let  $\text{Hom}_k(B^\dagger, A^\dagger)$  denote the set of  $k$ -linear maps from  $B^\dagger$  to  $A^\dagger$ .<sup>89</sup>

By 0.3.6.A, if  $M$  and  $P$  are pseudocompact  $k$ -modules and  $P$  is projective, the canonical map

$$\text{Hom}_c(M, P) \longrightarrow \text{Hom}_k(P^\dagger, M^\dagger), \quad f \longmapsto {}^t f,$$

where  ${}^t f$  denotes the transpose of  $f$ , is bijective. We shall apply this either with

$$M = A \widehat{\otimes} A, \quad P = B,$$

or with

$$M = A, \quad P = B \widehat{\otimes} B.$$

Let  $f \in \text{Hom}_c(A, B)$ . Consider the following diagrams. The squares marked (0) are commutative, and the two unnamed vertical arrows in the second diagram are  ${}^t(f \widehat{\otimes} f)$ .

$$\begin{array}{ccccccc} A \widehat{\otimes} A & \xrightarrow{m_A} & A & \xrightarrow{\Delta_A} & A \widehat{\otimes} A \\ f \widehat{\otimes} f \downarrow & (1) & \downarrow f & (2) & \downarrow f \widehat{\otimes} f \\ B \widehat{\otimes} B & \xrightarrow{m_B} & B & \xrightarrow{\Delta_B} & B \widehat{\otimes} B \end{array}$$
  

$$\begin{array}{ccccccc} A^\dagger \otimes A^\dagger & \xrightarrow{\varphi_G} & (A \widehat{\otimes} A)^\dagger & \xleftarrow{\delta_G = m_A^t} & A^\dagger & \xleftarrow{\Delta_A^t} & (A \widehat{\otimes} A)^\dagger & \xleftarrow{\varphi_G} & A^\dagger \otimes A^\dagger \\ {}^t f \otimes {}^t f \uparrow & (0) & \uparrow & (1') & \uparrow {}^t f & (2') & \uparrow & (0) & \uparrow {}^t f \otimes {}^t f \\ B^\dagger \otimes B^\dagger & = & (B \widehat{\otimes} B)^\dagger & \xleftarrow{\delta_K = m_B^t} & B^\dagger & \xleftarrow{\Delta_B^t} & (B \widehat{\otimes} B)^\dagger & = & B^\dagger \otimes B^\dagger \end{array}$$

If  $f : A \rightarrow B$  corresponds to a morphism of formal groups  $K \rightarrow G$ , then the squares (1) and (2) are commutative and

$$\varepsilon_B \circ f = \varepsilon_A.$$

Consequently, the squares (1') and (2') are commutative, and  ${}^t f$  sends the unit of  $B^\dagger = \mathbf{H}(K)$  to that of  $A^\dagger = \mathbf{H}(G)$ . Thus  ${}^t f$  is a homomorphism of unital  $k$ -algebras

$$\mathbf{H}(K) \longrightarrow \mathbf{H}(G)$$

for which the diagram in the proposition is commutative.

Conversely, suppose that  ${}^t f$  satisfies these conditions. Then

$$\varepsilon_B \circ f = \varepsilon_A$$

and the squares (1') and (2') are commutative. For

$$M = A \widehat{\otimes} A, \quad P = B \quad (\text{respectively } M = A, \quad P = B \widehat{\otimes} B),$$

the map  $g \mapsto {}^t g$  is injective. Hence the squares (1) and (2) are commutative, so  $f$  is compatible with the multiplications and diagonal morphisms of  $A$  and  $B$ .

<sup>89</sup>The editors supplied the details of the remainder of the proof.

It remains to prove that  $f(1_A) = 1_B$ . What precedes gives

$$\varepsilon_B f(1) = 1, \quad \Delta_B f(1) = f(1) \widehat{\otimes} f(1), \quad f(1) \cdot f(1) = f(1).$$

By 2.1(iii), the first two conditions imply that  $f(1)$  has inverse  $c_B f(1)$  in  $B$ . The equality  $f(1) \cdot f(1) = f(1)$  therefore implies  $f(1) = 1$ . Thus

$$f : A \longrightarrow B$$

is a morphism in  $\mathbf{Alp}/k$ , compatible with the comultiplications of  $A$  and  $B$ .

### 2.3.2

Let us now suppose, for simplicity, that the ring  $k$  is Artinian. When  $G$  is topologically flat over  $k$ , the algebra

$$\mathbf{H}(G) = \mathbf{H}(G)(k)$$

may be characterized by a universal property due to Cartier. Recall (cf. 1.2.1) that, if  $U$  is a  $k$ -module, we write  $\mathbf{W}(U)$  for the functor that assigns to every finite-length  $k$ -algebra  $C$  the  $C$ -module

$$U \otimes_k C.$$

<sup>90</sup> If  $U$  is a  $k$ -algebra (associative and with a unit element), then so is  $U \otimes_k C$ . We shall write  $\mathbf{W}(U)^\times$  for the  $k$ -group functor that assigns to every  $C \in \mathbf{Ob}(\mathbf{Alf}/k)$  the multiplicative group of invertible elements of the algebra  $U \otimes_k C$ :

$$\mathbf{W}(U)^\times(C) = (U \otimes_k C)^\times.$$

Moreover, identify  $G$  with the  $k$ -group functor

$$C \longmapsto \mathrm{Hom}_{\mathbf{Vaf}/k}(\mathrm{Spf}(C), G),$$

and write

$$\mathrm{Hom}_{k\text{-Gr.}}(G, \mathbf{W}(U)^\times)$$

for the set of homomorphisms of  $k$ -group functors from  $G$  to  $\mathbf{W}(U)^\times$ . We have the following.

**Proposition.** *Let  $k$  be an Artinian ring. For every formal group  $G$  topologically flat over  $k$ , and every  $k$ -algebra  $U$ , there is a canonical isomorphism*

$$\mathrm{Hom}_{k\text{-Gr.}}(G, \mathbf{W}(U)^\times) \xrightarrow{\sim} \mathrm{Hom}_{k\text{-Alg.}}(\mathbf{H}(G), U).$$

Let  $A$  be the affine algebra of  $G$ . By hypothesis,  $A$  is a projective object of  $\mathbf{PC}(k)$ , and

$$\mathbf{H}(G) = A^\dagger.$$

For every object  $P$  of  $\mathbf{PC}(k)$ , write

$$U \widehat{\otimes}_k P$$

for the inverse limit of the  $k$ -modules

$$U \otimes_k (P/N),$$

---

<sup>90</sup>The editors added the preceding sentence and, in what follows, wrote  $\mathbf{W}(U)^\times$  in place of  $U_m$ . Recall also (cf. 1.2.1) that a “ $k$ -functor” is a covariant functor from  $\mathbf{Alf}/k$  to  $(\mathbf{Sets})$ .

where  $N$  runs through the open submodules of  $P$ . There are linear maps

$$U \otimes_k (P/N) \longrightarrow \operatorname{Hom}_k((P/N)^*, U)$$

that send  $u \otimes x$  to the  $k$ -linear map

$$f \longmapsto f(x)u$$

and form a filtered inverse system. Passing to the inverse limit therefore gives a morphism

$$U \widehat{\otimes}_k P \xrightarrow{\psi_P} \operatorname{Hom}_k(P^\dagger, U). \quad (1)$$

When  $P = k$ , the morphism  $\psi_k$  is plainly an isomorphism. Moreover, the two sides of (1), considered as functors of  $P$ , commute with infinite products (every product being a filtered inverse limit of finite products). Thus (1) is an isomorphism when  $P$  is a product of copies of  $k$ , and then when  $P$  is a projective object of  $\operatorname{PC}(k)$  (the two sides of (1) commute with finite direct sums).

Now let

$$\operatorname{Hom}_{\mathcal{F}}(G, \mathbf{W}(U))$$

denote the set of morphisms of  $k$ -functors from  $G$  to  $\mathbf{W}(U)$ . Since

$$G = \operatorname{Spf}(A) = \varprojlim_{\mathfrak{l}} \operatorname{Spf}(A/\mathfrak{l}),$$

where  $\mathfrak{l}$  runs through the open ideals of  $A$ , there are canonical isomorphisms

$$\begin{aligned} \operatorname{Hom}_{\mathcal{F}}(G, \mathbf{W}(U)) &= \varprojlim_{\mathfrak{l}} \operatorname{Hom}_{\mathcal{F}}(\operatorname{Spf}(A/\mathfrak{l}), \mathbf{W}(U)) \\ &= \varprojlim_{\mathfrak{l}} U \otimes_k (A/\mathfrak{l}) = U \widehat{\otimes}_k A. \end{aligned}$$

The preceding argument therefore gives a canonical isomorphism

$$\operatorname{Hom}_{\mathcal{F}}(G, \mathbf{W}(U)) = U \widehat{\otimes}_k A \xrightarrow{\psi_A} \operatorname{Hom}_k(\mathbf{H}(G), U). \quad (2)$$

For every finite-length  $k$ -algebra  $C$ , multiplication makes  $U \otimes_k C$  a monoid with a unit element, and every homomorphism of monoids with unit

$$G(C) \longrightarrow U \otimes_k C$$

is necessarily a group homomorphism

$$G(C) \longrightarrow (U \otimes_k C)^\times.$$

It follows that

$$\operatorname{Hom}_{k\text{-Gr.}}(G, \mathbf{W}(U)^\times)$$

is the part of  $\operatorname{Hom}_{\mathcal{F}}(G, \mathbf{W}(U))$  consisting of morphisms of  $k$ -functors into monoids with a unit element.

It remains to show that these morphisms correspond to the  $k$ -linear maps from  $\mathbf{H}(G)$  to  $U$  that preserve multiplication and unit elements.<sup>91</sup> To simplify the notation, write

$$H = \mathbf{H}(G) = A^\dagger$$

---

<sup>91</sup>The original stated that this “follows from the functoriality of  $\psi_A$ .” The editors gave the details that follow.

and write  $\widehat{\otimes}$  instead of  $\widehat{\otimes}_k$ . Denote by

$$\Delta_A, m_A, \varepsilon_A \quad \text{and} \quad \Delta_H, m_H, \varepsilon_H$$

the comultiplication, multiplication, and augmentation of  $A$  and  $H$ , respectively. Let

$$\theta \in \text{Hom}_{\mathcal{F}}(G, \mathbf{W}(U)),$$

let  $\gamma$  be its image in  $U \widehat{\otimes} A$ , and let

$$\phi : H \longrightarrow U$$

be the associated  $k$ -linear map. The morphism  $\theta$  sends the unit section  $s \in G(k)$  to an element  $u$  of  $U$ . Since  $s$  corresponds to the augmentation

$$\varepsilon : A \longrightarrow k,$$

which is the unit element  $1_H$  of  $H$ , we have

$$\theta(s) = 1_U \iff \phi(1_H) = 1_U.$$

The morphism

$$\theta \circ m_G : G \times G \longrightarrow \mathbf{W}(U)$$

corresponds to the element

$$(\text{id}_U \widehat{\otimes} \Delta_A)(\gamma),$$

which, by duality, corresponds to the map

$$\phi \circ m_H : H \otimes H \longrightarrow U.$$

On the other hand, the morphism

$$\theta \circ \text{pr}_1 : G \times G \longrightarrow \mathbf{W}(U)$$

corresponds to the element

$$\gamma \widehat{\otimes} 1_A \in U \widehat{\otimes} A \widehat{\otimes} A,$$

which, by duality, corresponds to

$$\phi \circ (\text{id}_H \otimes \varepsilon_H) : H \otimes H \longrightarrow U.$$

Similarly,  $\theta \circ \text{pr}_2$  corresponds to the element

$$\tau(\gamma \widehat{\otimes} 1_A) \in U \widehat{\otimes} A \widehat{\otimes} A,$$

where

$$\tau(u \widehat{\otimes} a \widehat{\otimes} b) = u \widehat{\otimes} b \widehat{\otimes} a,$$

and this element corresponds by duality to

$$\phi \circ (\varepsilon_H \otimes \text{id}_H) : H \otimes H \longrightarrow U.$$

Finally, the multiplication map

$$\mu = m_{U \widehat{\otimes} A \widehat{\otimes} A}$$

given by

$$\begin{aligned} (U \hat{\otimes} A \hat{\otimes} A) \times (U \hat{\otimes} A \hat{\otimes} A) &\longrightarrow U \hat{\otimes} A \hat{\otimes} A, \\ (u \hat{\otimes} a_1 \hat{\otimes} a_2, u' \hat{\otimes} a_3 \hat{\otimes} a_4) &\longmapsto uu' \hat{\otimes} a_1 a_3 \hat{\otimes} a_2 a_4 \end{aligned}$$

may be viewed as the composite of the endomorphism  $\sigma_{23}$  of

$$(U \otimes U) \hat{\otimes} A^{\hat{\otimes} 4},$$

which “exchanges the factors  $a_2$  and  $a_3$ ,” and the map

$$(U \otimes U) \hat{\otimes} A^{\hat{\otimes} 2} \hat{\otimes} A^{\hat{\otimes} 2} \xrightarrow{m_U \hat{\otimes} m_A \hat{\otimes} m_A} U \hat{\otimes} A \hat{\otimes} A.$$

It follows that the map

$$\mu \circ (\phi \circ \text{pr}_1, \phi \circ \text{pr}_2) : G \times G \longrightarrow \mathbf{W}(U)$$

corresponds to the composite map  $\beta$  below:

$$\begin{array}{ccc} & H \otimes H & \xrightarrow{\beta} U \\ & \swarrow \text{id}_H \otimes \sigma_{23} \otimes \text{id}_H & \uparrow m_U \\ H \otimes H \otimes H \otimes H & & U \otimes U \\ & \searrow (\text{id}_H \otimes \varepsilon_H) \otimes (\varepsilon_H \otimes \text{id}_H) & \uparrow \phi \otimes \phi \\ & H \otimes H & . \end{array}$$

Finally,  $\theta$  is compatible with the laws of  $G$  and  $\mathbf{W}(U)^\times$  if and only if

$$\theta \circ m_G = \mu \circ (\phi \circ \text{pr}_1, \phi \circ \text{pr}_2),$$

which, by the preceding argument, is equivalent to

$$\phi \circ m_H = \beta.$$

But plainly

$$(\phi \circ m_H)(x \otimes y) = \phi(xy),$$

whereas

$$\beta(x \otimes y) = \phi(x)\phi(y).$$

## 2.4

We now return to an arbitrary pseudocompact ring  $k$ , in order to apply to formal groups the results of 1.4–1.5 on passage to the quotient by a topologically flat equivalence relation.  
92

Let  $u : H \rightarrow G$  be a monomorphism of formal  $k$ -groups, let  $\mu : G \times G \rightarrow G$  be the multiplication morphism of  $G$ , and let  $\lambda$  be the composite

$$\lambda : G \times H \xrightarrow{\text{id}_G \times u} G \times G \xrightarrow{\mu} G.$$

Since  $u$  is a monomorphism, the pair

<sup>92</sup>The editors supplied the details of the original argument that follow.

$$G \times H \xrightarrow[\lambda]{\text{pr}_1} G$$

is an equivalence pair in  $\text{Vaf}/k$  (cf. V, 2.b). Recall (cf. 1.2.C) that the cokernel  $G/H$  of this pair is defined as follows.

Let  $\mathcal{O}(G)$  and  $\mathcal{O}(H)$  be the affine algebras of  $G$  and  $H$ , let

$$\Delta : \mathcal{O}(G) \longrightarrow \mathcal{O}(G) \hat{\otimes}_k \mathcal{O}(G)$$

be the diagonal morphism of  $\mathcal{O}(G)$ , and let  $I$  be the kernel of

$$u^\sharp : \mathcal{O}(G) \longrightarrow \mathcal{O}(H).$$

By Proposition 1.3, the morphism  $u^\sharp$  induces an isomorphism

$$\mathcal{O}(G)/I \xrightarrow{\sim} \mathcal{O}(H).$$

The affine algebra  $\mathcal{O}(G/H)$  of  $G/H$  is then the kernel of the pair

$$\mathcal{O}(G) \xrightarrow[\text{(id} \hat{\otimes} u^\sharp)_\Delta]{\tau_1} \mathcal{O}(G) \hat{\otimes}_k \mathcal{O}(H),$$

where  $\tau_1(x) = x \hat{\otimes} 1$ . Thus

$$\mathcal{O}(G/H) = \{x \in \mathcal{O}(G) \mid \Delta(x) - x \hat{\otimes} 1 \in \mathcal{O}(G) \hat{\otimes} I\}.$$

If, moreover,  $H$  is topologically flat over  $k$ , then  $\text{pr}_1$  is topologically flat, and Theorem 1.4 gives the following result.

**Theorem.** *Let  $u : H \rightarrow G$  be a monomorphism of formal  $k$ -groups. Suppose that  $H$  is topologically flat over  $k$ . Then the projection*

$$p : G \longrightarrow G/H$$

*is surjective and topologically flat, there is an isomorphism*

$$G \times H \xrightarrow{\sim} G \times_{G/H} G, \quad (*)$$

*and  $G/H$  represents the quotient sheaf for the flat topology.*

Consequently,  $G/H$  has a canonical structure of an object on which  $G$  acts, and  $p : G \rightarrow G/H$  is an equivariant morphism. If, in addition,  $u$  identifies  $H$  with a normal subgroup of  $G$ , then  $G/H$  has a canonical structure of formal  $k$ -group for which  $p : G \rightarrow G/H$  is a morphism of formal  $k$ -groups, and  $H$  is the kernel of  $p$ .

Indeed, the first assertion follows from 1.4, and the other two from Exposé IV, Corollaries 5.2.2 and 5.2.4.

**Corollary.** <sup>93</sup> *Let  $G$  be a formal  $k$ -group and  $H$  a formal subgroup of  $G$ . Let  $A$ ,  $A/J$ , and  $B$  be the affine algebras of  $G$ ,  $H$ , and  $G/H$ , respectively; let  $I_A$  be the augmentation ideal of  $A$ , and put  $I_B = B \cap I_A$ . Suppose that  $H$  is topologically flat over  $k$ . Then*

$$J = AI_B,$$

*the closed ideal generated by  $I_B$ .*

Indeed, the projection  $B \rightarrow B/I_B$  corresponds to the unit section

$$e : \text{Spf}(k) \longrightarrow G/H$$

of  $G/H$ . By (\*),  $H$  is identified with the fiber product

$$\text{Spf}(k) \times_{G/H} G.$$

Its affine algebra  $A/J$  is therefore identified with

$$(B/I_B) \hat{\otimes}_B A \simeq A/AI_B.$$

<sup>93</sup>The editors made this corollary explicit; it is used, for example, in 5.2.1 and 5.2.3.

**2.4.A**

Let  $G$  and  $Q$  be formal  $k$ -groups that are topologically flat. Suppose that there are homomorphisms

$$\sigma : Q \longrightarrow G \quad \text{and} \quad \pi : G \longrightarrow Q$$

such that

$$\pi \circ \sigma = \text{id}_Q.$$

<sup>94</sup> In particular,  $\sigma$  is a monomorphism, so  $Q$  is a formal subgroup of  $G$  (cf. Remark 1.3). Let

$$N = \text{Ker}(\pi)$$

and let  $\sigma'$  be the inclusion  $N \hookrightarrow G$ . Then  $G$  is the semidirect product of  $N$  by  $Q$  (cf. I, 2.3.8). In other words, for every

$$B \in \text{Ob}(\text{Alf}/k),$$

identify  $N(B)$  and  $Q(B)$  with their images in  $G(B)$  under  $\sigma'$  and  $\sigma$ . Then  $N(B)$  is a normal subgroup of  $G(B)$ , and the map

$$\mu : N(B) \times Q(B) \longrightarrow G(B), \quad (x, q) \longmapsto xq \tag{1}$$

is bijective.

The morphism of formal  $k$ -varieties

$$\theta : G(B) \longrightarrow N(B), \quad g \longmapsto g \cdot \sigma\pi(g^{-1}) \tag{2}$$

is then a retraction of  $\sigma'$ , and the inverse of  $\mu$  is

$$g \longmapsto (\theta(g), \pi(g)). \tag{3}$$

If  $p : G \rightarrow G/Q$  denotes the quotient projection, then

$$p \circ \sigma' : N \longrightarrow G/Q$$

is an isomorphism of formal  $k$ -varieties.

*Source note.* The Polo–Gille text prints  $\theta \circ \sigma' : N \rightarrow G/Q$  here. Since  $\theta : G \rightarrow N$ , that composite is ill-typed; the isomorphism with the displayed source and target is the quotient composite  $p \circ \sigma'$ .

In particular,  $N$  is topologically flat over  $k$ , by 1.4 (ii). Let

$$\alpha : Q \times N \longrightarrow N$$

and

$$\beta : G \times N = N \times Q \times N \longrightarrow N$$

be the maps defined on points by

$$\alpha(q, y) = qyq^{-1}, \quad \beta(x, q, y) = x\alpha(q, y).$$

Then the following diagram is commutative:

$$(4) \quad \begin{array}{ccccc} G \times G & \xrightarrow{\text{id} \times \theta} & G \times N & \xrightarrow{\theta \times \text{id}} & N \times N \\ m_G \downarrow & & \downarrow \beta & \swarrow m_N & \\ G & \xrightarrow{\theta} & N & & \end{array} .$$

<sup>94</sup>The editors added paragraphs 2.4.A and 2.4.B, which will be useful in 5.1.3 and 2.9, respectively.



This can be expressed as follows in terms of the affine algebras  $A$ ,  $A_0$ , and  $A'$  of  $G$ ,  $Q$ , and  $N$  (cf. 5.1.3 below). Let

$$\rho' : A \longrightarrow A', \quad \rho : A \longrightarrow A_0, \quad \tau : A_0 \longrightarrow A$$

be the homomorphisms of  $k$ -bialgebras corresponding to  $\sigma'$ ,  $\sigma$ , and  $\pi$ , and put

$$I = \text{Ker}(\rho).$$

By the preceding corollary,  $A'$  is identified with

$$A/\overline{A\tau(J_0)},$$

where  $J_0$  is the augmentation ideal of  $A_0$ .

On the other hand, let  $B$  be the affine algebra of  $G/Q$ . It is the kernel of the pair of morphisms

$$A \xrightarrow[\text{(id}_A \widehat{\otimes} \rho) \circ \Delta_A]{\tau_1} A \widehat{\otimes}_k A_0,$$

where  $\tau_1(x) = x \widehat{\otimes} 1$ . Thus

$$B = \{x \in A \mid \Delta_A(x) - x \widehat{\otimes} 1 \in A \widehat{\otimes} I\}.$$

Let

$$\gamma = \theta^\natural : A' \longrightarrow A$$

be the corresponding continuous homomorphism of  $k$ -algebras. It is a section of  $\rho'$ , and it is an isomorphism of profinite  $k$ -algebras from  $A'$  onto  $B$ . Moreover,

$$\tau = \pi^\natural$$

identifies  $A_0$  with a sub-bialgebra of  $A$ , namely the affine algebra of the quotient  $N \backslash G$ .

It follows from equations (1) and (3) that there is an isomorphism of profinite  $k$ -algebras

$$A' \widehat{\otimes}_k A_0 \xrightarrow{\sim} A, \quad a' \widehat{\otimes} a_0 \longmapsto \gamma(a')\tau(a_0), \quad (*)$$

whose inverse is the map

$$a \longmapsto (\rho' \widehat{\otimes} \rho) \Delta_A(a).$$

Finally, identify  $A'$  with its image in  $A$  under  $\gamma$ , so that the projection  $A \rightarrow A'$  is then  $\gamma\rho'$ . Writing  $\Delta_{A'}$  for the comultiplication of  $A'$ , one deduces from (4) that

$$\Delta_A(A') \subset A \widehat{\otimes} A'$$

and that the following diagram is commutative:

$$\begin{array}{ccc} A' & \xrightarrow{\Delta_A} & A \widehat{\otimes} A' \\ \parallel & & \downarrow \gamma\rho' \widehat{\otimes} \text{id} \\ A' & \xrightarrow{\Delta_{A'}} & A' \widehat{\otimes} A' \end{array}$$

Consequently, one also has

$$\gamma\rho' \circ c_A = c_{A'},$$

where  $c_A$  and  $c_{A'}$  are the antipodes of  $A$  and  $A'$ , respectively. On the other hand, writing  $m_A$  for multiplication in  $A$ , equation (2) gives, for every  $a \in A$ ,

$$\gamma\rho'(a) = (m_A \circ (\text{id}_A \widehat{\otimes} \tau\rho c_A) \circ \Delta_A)(a).$$

**2.4.B**

Suppose, for simplicity, that  $k$  is Artinian. The preceding discussion can then be expressed more simply in terms of the covariant bialgebras of  $G$ ,  $Q$ , and  $N$ . Indeed, since

$$G = N \times Q$$

as formal  $k$ -varieties, one has

$$\mathbf{H}(G) = \mathbf{H}(N) \otimes_k \mathbf{H}(Q)$$

as  $k$ -coalgebras. Moreover, since multiplication in  $G$  is given by

$$(x, q) \cdot (x', q') = (x\alpha(q, x'), qq'), \quad \alpha(q, x') = qx'q^{-1},$$

multiplication in  $\mathbf{H}(G)$  is given, for  $x \in \mathbf{H}(N)$  and  $q \in \mathbf{H}(Q)$ , by

$$(x \otimes q) \cdot (x' \otimes q') = x\phi(q, x') \otimes qq',$$

where

$$\phi : \mathbf{H}(Q) \otimes_k \mathbf{H}(N) \longrightarrow \mathbf{H}(N)$$

is the morphism of  $k$ -coalgebras induced by  $\alpha$ .

The morphism  $\alpha$  is the following composite. Here  $\delta_G$  and  $m_G$  denote, respectively, the diagonal and multiplication morphisms of  $G$ ,  $c_Q$  is inversion in  $Q$ , and

$$v(q \otimes q' \otimes x) = q \otimes x \otimes q'.$$

$$Q \times N \xrightarrow{v \circ (\delta_G \times \text{id})} Q \times N \times Q \xrightarrow{\text{id} \times \text{id} \times c_Q} Q \times N \times Q \xrightarrow{m_G} G,$$

Writing  $c_Q$  also for the antipode of  $\mathbf{H}(Q)$ , we obtain

$$\phi(q \otimes x') = \sum_i q_i x' c_Q(q'_i) \quad \text{if} \quad \Delta_{\mathbf{H}(Q)}(q) = \sum_i q_i \otimes q'_i. \quad (\star)$$

In particular, let  $M$  be an abstract group and suppose that  $\mathbf{H}(Q)$  is the group algebra  $kM$ , so that

$$Q = \text{Spf}^* kM$$

is the constant formal  $k$ -group  $M_k$ . Then for every  $\gamma \in M$  and  $x' \in \mathbf{H}(N)$ ,

$$\phi(\gamma \otimes x') = \gamma x' \gamma^{-1}.$$

This defines an action of  $M$  on  $\mathbf{H}(N)$  by Hopf-algebra automorphisms.

**Proposition 2.4.1.** *Let  $f : G \rightarrow K$  be a morphism of formal  $k$ -groups. If*

$$H = \ker(f)$$

*is topologically flat over  $k$ , the homomorphism*

$$f' : G/H \longrightarrow K$$

*induced by  $f$  is a monomorphism.*

This is a consequence of the results of Exposé IV,<sup>95</sup> but we give a direct proof. Let  $T$  be a formal variety of finite length over  $k$ , and let  $t \in (G/H)(T)$  be such that  $f' \circ t$  is the identity element of  $K(T)$ . We must show that  $t$  is the identity element of  $(G/H)(T)$ . Let

$$p : G \longrightarrow G/H$$

be the projection, and let

$$X = T \times_{G/H} G.$$

By 2.4,  $p$  is surjective and topologically flat. The same is therefore true of

$$\mathrm{pr}_1 : X \longrightarrow T.$$

By Proposition 1.3.1,  $\mathrm{pr}_1$  is consequently an epimorphism, so it is enough to show that  $t \circ \mathrm{pr}_1$  is the identity element of  $(G/H)(X)$ .

Let  $\mathrm{pr}_2 : X \rightarrow G$  be the second projection. We have

$$t \circ \mathrm{pr}_1 = p \circ \mathrm{pr}_2,$$

and hence

$$1 = f' \circ t \circ \mathrm{pr}_1 = f' \circ p \circ \mathrm{pr}_2 = f \circ \mathrm{pr}_2.$$

The exact sequence

$$1 \longrightarrow H \longrightarrow G \xrightarrow{f} K$$

therefore shows that  $\mathrm{pr}_2$  factors through  $H$ . Thus  $p \circ \mathrm{pr}_2$  is the zero morphism. Since

$$p \circ \mathrm{pr}_2 = t \circ \mathrm{pr}_1,$$

this proves the proposition.

The following corollary follows. Write  $\mathcal{O}(G)$ ,  $\mathcal{O}(K)$ , and  $\mathcal{O}(G/H)$  for the affine algebras of  $G$ ,  $K$ , and  $G/H$ , respectively. We saw in 2.4 that  $p$  induces an injection

$$\mathcal{O}(G/H) \hookrightarrow \mathcal{O}(G).$$

Moreover, since  $f' : G/H \rightarrow K$  is a monomorphism, Proposition 1.3 shows that

$$\mathcal{O}(K) \longrightarrow \mathcal{O}(G/H)$$

is surjective. Hence:

**Corollary.** *Let  $f : G \rightarrow K$  be a morphism of formal  $k$ -groups and let  $H = \ker(f)$ . If  $H$  is topologically flat over  $k$ , then*

$$\mathcal{O}(G/H)$$

*is the image of  $\mathcal{O}(K)$  in  $\mathcal{O}(G)$ .*

---

<sup>95</sup>See Exposé IV, 5.2.6.

## 2.4.2

Retain the preceding notation and suppose that  $H$  and  $G$  are topologically flat over  $k$ . Then  $G$  is topologically flat both over  $k$  and over  $G/H$ ; hence, by 1.3.3,  $G/H$  is topologically flat over  $k$ . Consequently, by 2.4, the canonical projection

$$q : K \longrightarrow \operatorname{Coker}(f')$$

is topologically flat, and  $f'$  is an isomorphism from  $G/H$  onto  $\ker(q)$ . It is also clear that

$$\operatorname{Coker}(f) = \operatorname{Coker}(f').$$

Thus, under the hypothesis that  $G$  and  $H = \ker(f)$  are topologically flat over  $k$ , we have obtained an isomorphism between  $\ker(q)$ , the image of  $f$ , and

$$G/\ker(f),$$

the coimage of  $f$ . This proves the following theorem.

**Theorem.** *Let  $k$  be a field. The commutative formal  $k$ -groups form an abelian category.*

**Corollary.** *Let  $k$  be a field. The commutative affine  $k$ -group schemes form an abelian category.*<sup>96</sup>

This follows from the theorem and the equivalence of categories in 2.2.2.

## 2.5

A formal  $k$ -group is said to be *étale* if its underlying formal variety is étale (cf. 1.6). Such formal groups have a very simple structure. Suppose that  $k$  is local, let  $\kappa$  be its residue field, let  $\kappa_s$  be a separable closure of  $\kappa$ , and let

$$\Gamma = \operatorname{Gal}(\kappa_s/\kappa)$$

be the topological Galois group. A  $\Gamma$ -set means a set  $E$  with a continuous action of  $\Gamma$ ; equivalently, the stabilizer of every element  $x \in E$  is an open subgroup of  $\Gamma$ .

For every formal  $k$ -variety  $X$ , put

$$X(\kappa_s) = \varinjlim_{\ell} X(\ell),$$

where  $\ell$  ranges over the finite extensions of  $\kappa$  contained in  $\kappa_s$ .<sup>97</sup> The group  $\Gamma$  acts continuously on every  $X(\ell)$ , and hence on  $X(\kappa_s)$ . Moreover, if

$$X_{\kappa} = X \widehat{\otimes}_k \kappa \quad (\text{cf. 1.6.C}),$$

then  $X(\ell) = X_{\kappa}(\ell)$  for every  $\ell$ , and therefore

$$X(\kappa_s) = X_{\kappa}(\kappa_s).$$

Now suppose that  $X$  is étale over  $k$ . Then  $X_{\kappa}$  is the formal  $\kappa$ -variety that is the disjoint union of the  $\operatorname{Spec} \kappa(x)$ , for  $x \in X$ . If  $X'_{\kappa}$  denotes the  $\kappa$ -scheme given by the same disjoint union, then

$$\begin{aligned} X_{\kappa}(\kappa_s) &= X'_{\kappa}(\kappa_s) \\ &= \operatorname{Hom}_{\operatorname{Sch}/\kappa}(\operatorname{Spec} \kappa_s, X'_{\kappa}). \end{aligned}$$

<sup>96</sup>See also the remarks following Exposé VI<sub>A</sub>, 5.4.3.

<sup>97</sup>If  $[\kappa_s : \kappa] = \infty$ , then  $\kappa_s$  is not a profinite  $\kappa$ -algebra, so the preceding notation is an abuse. The editors also supplied the details that follow.

Let  $\mathcal{C}$  be the full subcategory

$$\mathrm{Sch}^{\mathrm{\acute{e}t}}/\kappa \subset \mathrm{Sch}/\kappa$$

of étale  $\kappa$ -schemes. It is known that

$$X' \longmapsto X'(\kappa_s)$$

is an equivalence from  $\mathcal{C}$  onto the category  $\mathcal{C}'$  of  $\Gamma$ -sets (cf. SGA 1, V, §§7–8, or [DG70], §I.4, 6.4). It therefore induces an equivalence between the category of group objects in  $\mathcal{C}$  and the category of group objects in  $\mathcal{C}'$ . A group object in  $\mathcal{C}'$  is simply an abstract group endowed with a continuous action of  $\Gamma$  by group automorphisms; we call this a  $\Gamma$ -group.

Taking into account the equivalences of 1.6.E,

$$\mathrm{Vaf}^{\mathrm{\acute{e}t}}/k \xrightarrow{\sim} \mathrm{Vaf}^{\mathrm{\acute{e}t}}/\kappa \xrightarrow{\sim} \mathrm{Sch}^{\mathrm{\acute{e}t}}/\kappa,$$

we obtain the following.

**Proposition.** *Let  $k$  be a local pseudocompact ring,  $\kappa$  its residue field,  $\kappa_s$  a separable closure of  $\kappa$ , and  $\Gamma = \mathrm{Gal}(\kappa_s/\kappa)$ .*

- (i) *The functor  $X \mapsto X(\kappa_s)$  is an equivalence from the category of étale formal  $k$ -varieties onto the category of  $\Gamma$ -sets.*
- (ii) *It induces an equivalence from the category of étale formal  $k$ -groups onto the category of  $\Gamma$ -groups.*

**Remark 2.5.A.** <sup>98</sup> Let  $k$  be a field, let  $k_s$  be a separable closure of  $k$ , let  $G$  be an étale formal  $k$ -group, and put

$$M = G(k_s).$$

Choose a set  $X$  of representatives for the orbits of  $\Gamma = \mathrm{Gal}(k_s/k)$  in  $M$ . For each  $x \in X$ , let  $\Gamma_x$  be its stabilizer, an open subgroup of finite index in  $\Gamma$ , and put

$$L_x = k_s^{\Gamma_x}.$$

Then  $L_x/k$  has degree  $[\Gamma : \Gamma_x]$  (see, for example, [BA1g], V, §10.10).

By the preceding equivalence of categories, the affine algebra  $\mathcal{A}(G)$  of  $G$  is the product of the  $L_x$ , endowed with the product topology. The  $L_x$  are therefore exactly the simple quotients

$$\mathcal{A}(G)/\mathfrak{m},$$

where  $\mathfrak{m}$  ranges over the open maximal ideals of  $\mathcal{A}(G)$ . Since these quotients correspond by duality to the simple subcoalgebras of  $\mathbf{H}(G)$ , the coalgebra  $\mathbf{H}(G)$  is pointed—that is,

$$\dim_k C = 1$$

for every simple subcoalgebra  $C$ —if and only if  $L_x = k$  for every  $x$ . In that case,  $\mathcal{A}(G)$  is the topological algebra  $k^M$ , so  $\mathbf{H}(G)$  is the group algebra  $kM$ , and

$$M = \{x \in \mathbf{H}(G) \mid \varepsilon(x) = 1, \Delta(x) = x \otimes x\} = G(k).$$

---

<sup>98</sup>The editors added this remark.

## 2.5.1

Suppose again that  $k$  is an arbitrary pseudocompact ring. Since the functor

$$X \longmapsto X_e$$

of 1.6.F commutes with finite products, it sends every formal group  $G$  to an étale formal group  $G_e$ . The morphism  $p : X \rightarrow X_e$  of that result is functorial in  $X$ , so in this case

$$p : G \longrightarrow G_e$$

is a morphism of formal groups.

Consider its kernel  $\ker(p)$ .<sup>99</sup> It is the fiber product of the diagram

$$\begin{array}{ccc} & & G \\ & & \downarrow p \\ \mathrm{Spf}(k) & \xrightarrow{\varepsilon} & G_e \end{array}$$

where  $\varepsilon$  is the identity section of  $G_e$ . Since  $p$  induces a bijection on the underlying sets, it follows (cf. 1.2, editorial note 41) that the underlying set of  $\ker(p)$  is the image of  $\mathrm{Spf}(k)$  under  $\varepsilon$ , and that, for every point  $s$  of  $\mathrm{Spf}(k)$ , the local algebra of  $\ker(p)$  at  $\varepsilon(s)$  is

$$\mathcal{O}_{G, \varepsilon(s)} \hat{\otimes}_{\mathcal{O}_{G_e, \varepsilon(s)}} \kappa(s).$$

At each point  $\varepsilon(s)$ , the residue field of  $\mathcal{O}_{G, \varepsilon(s)}$  is  $\kappa(s)$ , and hence

$$\mathcal{O}_{G_e, \varepsilon(s)} = \kappa(s), \quad \mathcal{O}_{\ker(p), \varepsilon(s)} = \mathcal{O}_{G, \varepsilon(s)}.$$

For these reasons, we call  $\ker(p)$  the *infinitesimal neighborhood of the identity* of  $G$ , and write

$$\ker(p) = G_0.$$

We have thus obtained the exact sequence

$$1 \longrightarrow G_0 \xrightarrow{\text{incl.}} G \xrightarrow{p} G_e.$$

In what follows, we say that  $G$  is *infinitesimal* if

$$G = G_0.$$

<sup>100</sup> This is equivalent to saying that, for every  $s \in \mathrm{Spf}(k)$ , or equivalently for every continuous morphism  $k \rightarrow \kappa$ , where  $\kappa$  is a field with the discrete topology, the identity is the unique element of  $G(\kappa(s))$ , respectively of  $G(\kappa)$ .

Suppose in addition that  $G$  is topologically flat over  $k$ .<sup>101</sup> By 1.6.1, the morphism

$$p : G \longrightarrow G_e$$

is topologically flat. Since it is bijective, Proposition 1.3.1 shows that it is an effective epimorphism. Consequently,  $G_e$  is identified with the quotient  $G/G_0$ . We have proved:

**Corollary.** *Let  $G$  be a formal group topologically flat over  $k$ . Then  $G_e$  is identified with the quotient  $G/G_0$ ; in other words, there is an exact sequence of formal groups*

$$1 \longrightarrow G_0 \xrightarrow{\text{incl.}} G \xrightarrow{p} G_e \longrightarrow 1.$$

<sup>99</sup>The editors supplied the details of the original argument that follow.

<sup>100</sup>The editors added the sentence that follows.

<sup>101</sup>The editors supplied the details that follow.

**2.5.2**

Suppose that  $k$  is a perfect field. In this case, we saw in 1.6.2 that the morphism

$$p : X \mapsto X_e$$

has a section

$$s : X_e \longrightarrow X$$

that depends functorially on  $X$ . Consequently, when  $X$  is a formal group, this section is a morphism of formal groups. We thus obtain:

**Proposition.** *When  $k$  is a perfect field, every formal  $k$ -group  $G$  is canonically isomorphic to the semidirect product of an infinitesimal group  $G_0$  and an étale group  $G_e$  acting on  $G_0$ .<sup>102</sup>*

If, moreover,  $G$  is commutative, then  $G$  is the product of  $G_0$  and  $G_e$ . By 2.2.2, this canonical decomposition of commutative formal  $k$ -groups corresponds to an analogous decomposition of affine commutative  $k$ -group schemes; see 2.5.3 below.

**Remark 2.5.2.A.** <sup>103</sup> When  $k$  is not perfect, the exact sequence

$$1 \longrightarrow G_0 \longrightarrow G \longrightarrow G_e \longrightarrow 1$$

need not split. Consider the following example, taken from [DG70], §III.6, 8.6. Let  $k$  be a nonperfect field of characteristic  $p > 0$ , choose

$$\lambda \in k \setminus k^p,$$

and let  $L_\lambda$  be the abelian restricted  $p$ -Lie algebra with basis  $(x, y)$ , whose symbolic  $p$ -th power (cf. VII<sub>A</sub>, 5.2) is given by

$$x^{(p)} = x, \quad y^{(p)} = \lambda x.$$

Let

$$U_p(L_\lambda) = k[x, y]/(x^p - x, y^p - \lambda x)$$

be the restricted enveloping algebra of  $L_\lambda$  (cf. VII<sub>A</sub>, 5.3), and let  $G_\lambda$  be the commutative formal  $k$ -group whose affine algebra is  $U_p(L_\lambda)$ .

Then  $G_\lambda$  is a nonsplit extension of the constant étale  $k$ -group  $(\mathbb{Z}/p\mathbb{Z})_k$  by the infinitesimal  $k$ -group

$$\alpha_{p,k} = \mathrm{Spf} \, k[x]/(x^p).$$

*Source note.* The French re-edition prints  $\mathrm{Spf} \, k[x]/(x^p - x)$  here. That algebra is étale, whereas the asserted kernel is the standard infinitesimal group  $\alpha_{p,k} = \mathrm{Spf} \, k[x]/(x^p)$ . Indeed, the kernel fiber  $x = 0$  in  $U_p(L_\lambda)$  has coordinate algebra  $k[y]/(y^p)$ . The English text uses the standard infinitesimal reading while preserving the printed reading in this note. Thus there is a nonsplit exact sequence of commutative formal  $k$ -groups:<sup>104</sup>

$$0 \longrightarrow \alpha_{p,k} \longrightarrow G_\lambda \longrightarrow (\mathbb{Z}/p\mathbb{Z})_k \longrightarrow 0.$$

<sup>102</sup>The same argument also shows that, for an arbitrary field  $k$ , if all the residue fields of  $G$  are equal to  $k$ , then  $G_e$  is the constant  $k$ -group  $(M)_k$ , where

$$M = G(k) = \mathrm{Spf}^* \mathbf{H}(G),$$

and  $\mathbf{H}(G)$  is the semidirect product of  $\mathbf{H}(G_0)$  by the group algebra  $kM$ ; see the addition in 2.9 below.

<sup>103</sup>The editors added this remark.

<sup>104</sup>The groups  $\alpha_{p,k}$ ,  $(\mathbb{Z}/p\mathbb{Z})_k$ , and  $G_\lambda$  are also finite flat algebraic  $k$ -groups; see editorial note 84 in 2.2.2.

By duality, this corresponds to a nonsplit exact sequence of commutative algebraic  $k$ -groups:

$$0 \longrightarrow \mu_{p,k} \longrightarrow D'(G_\lambda) \longrightarrow \alpha_{p,k} \longrightarrow 0$$

where

$$\mu_{p,k} = \operatorname{Spec} k[t]/(t^p - 1).$$

In this way one obtains all extensions of  $\alpha_{p,k}$  by  $\mu_{p,k}$ ; see [DG70], §III.6, 8.6.

### 2.5.3

Let  $k$  be a field.<sup>105</sup>

**Definition 2.5.3.A.** A commutative  $k$ -group scheme is said to be *unipotent* if it is isomorphic to

$$\operatorname{Spec} \mathbf{H}(G),$$

where  $G$  is an infinitesimal commutative formal  $k$ -group.<sup>106</sup>

On the other hand, following Exposé IX, Definition 1.1,<sup>107</sup> a  $k$ -group scheme  $H$  is said to be of *multiplicative type* if there exists a scheme  $S$ , faithfully flat and quasi-compact over  $\operatorname{Spec} k$ , such that  $H_S$  is a diagonalizable  $S$ -group; that is,

$$H_S \simeq D_S(M)$$

for some “abstract” abelian group  $M$ . By fpqc descent, this implies that  $H$  is affine and commutative. Moreover, since  $S$  may be replaced by the residue field of one of its points,  $H$  is of multiplicative type if and only if there exists an extension  $K$  of  $k$  such that  $H_K$  is a diagonalizable  $K$ -group.

**Proposition 2.5.3.B.** *Let  $T$  be an affine commutative  $k$ -group scheme, and let  $k_s$  be a separable closure of  $k$ .*

- (i) *The scheme  $T$  is of multiplicative type if and only if its dual  $D(T)$  is an étale commutative formal  $k$ -group.*
- (ii) *Consequently,  $T$  is of multiplicative type if and only if*

$$T \otimes_k k_s$$

*is diagonalizable.*

*Proof.* Let  $\mathcal{A}$  denote the affine algebra of  $D(T)$ , and let  $\mathcal{A}_0$  be its local component at the identity. Then  $D(T)$  is étale over  $k$  if and only if

$$\mathcal{A}_0 = k.$$

If  $K$  is an extension of  $k$ , then  $\mathcal{A}_0 \widehat{\otimes}_k K$  is local, being an inverse limit of Artinian local rings, and hence coincides with the local component

$$(\mathcal{A} \widehat{\otimes}_k K)_0.$$

<sup>105</sup>The editors introduced the numbering 2.5.3 and 2.5.3.A–2.5.3.C.

<sup>106</sup>This agrees with the “usual” notion of a unipotent  $k$ -group scheme; cf. the addition in 2.8 at the end of Section 2.

<sup>107</sup>The original stated: “Let us say that a commutative  $k$ -group scheme is of multiplicative type if it is isomorphic to  $\operatorname{Spec} \mathbf{H}(G)$ , where  $G$  is an étale commutative formal  $k$ -group.” The editors have given here the “usual” definition, taken from Exposé IX, 1.1, and proved its equivalence with the preceding condition; see also [DG70], §IV.1, Theorem 2.2.



Moreover, since the formation of  $D(T)$  commutes with base change (cf. 2.2.2), one also has

$$\mathcal{A} \widehat{\otimes}_k K \simeq \mathcal{A}(D(T_K)).$$

*Source note.* The French re-edition prints  $\mathcal{A} \widehat{\otimes}_k K \simeq D(T_K)$ . The left-hand side is an algebra and the right-hand side a formal group. The English therefore uses the type-correct intended reading above and preserves the printed formula here.

Suppose that  $T$  is of multiplicative type. Then there exists an extension  $K$  of  $k$  such that

$$\mathcal{O}(T) \otimes_k K$$

is isomorphic, as a Hopf  $K$ -algebra, to the group algebra  $KM$  for some abelian group  $M$ . It follows that  $\mathcal{A} \widehat{\otimes}_k K$  is isomorphic to the algebra  $K^M$ , endowed with the product topology. Thus

$$K = (\mathcal{A} \widehat{\otimes}_k K)_0 = \mathcal{A}_0 \widehat{\otimes}_k K.$$

This implies that  $\mathcal{A}_0 = k$ , and hence that  $D(T)$  is étale.

Conversely, suppose that  $D(T)$  is étale. Then

$$D(T) \widehat{\otimes}_k k_s = D(T_{k_s})$$

is the constant  $k_s$ -group associated with a group  $M$ . Therefore its covariant bialgebra is the group algebra  $k_s M$  (cf. Remark 2.5.A), and consequently

$$\mathcal{O}(T) \otimes_k k_s = k_s M.$$

This proves that  $T$  is of multiplicative type and is split by the extension  $k \rightarrow k_s$ . The proposition follows.

By 2.2.2, 2.5.1, and 2.5, we obtain:

**Corollary 2.5.3.C.** *Let  $k$  be a field, and let  $G$  be an affine commutative  $k$ -group scheme.*

- (i) *The group  $G$  contains a subgroup  $G_m$  of multiplicative type such that  $G/G_m$  is unipotent.*
- (ii) *When  $k$  is perfect, there is in addition a canonical retraction of  $G$  onto  $G_m$ , so that  $G$  is the product of a unipotent group and a group of multiplicative type.*
- (iii) <sup>108</sup> *Let  $k_s$  be a separable closure of  $k$ , and put*

$$\Gamma = \text{Gal}(k_s/k).$$

*The category of  $k$ -group schemes of multiplicative type is anti-equivalent to the category of continuous  $\Gamma$ -modules.*

## 2.6

We shall now study infinitesimal formal groups, to which the sections that follow are devoted. In this study, Lie algebras play a fundamental role.

Suppose first that the base ring  $k$  is Artinian, and let  $G$  be a formal group over  $k$ . The Lie algebra of  $G$  admits three different interpretations, all of which we shall use:

a) Let

$$D = k[d]/(d^2)$$

---

<sup>108</sup>The editors added item (iii), a consequence of Proposition 2.5.

be the algebra of dual numbers over  $k$ , and let

$$\delta : D \longrightarrow k$$

be the homomorphism that annihilates  $d$ . For every formal group  $G$  over  $k$ ,  $\text{Lie}(G)$  is the kernel of  $G(\delta)$ . Thus, by definition, there is an exact sequence of groups

$$1 \longrightarrow \text{Lie}(G) \longrightarrow G(D) \xrightarrow{G(\delta)} G(k) \longrightarrow 1.$$

b) Let  $A$  be the affine algebra of  $G$ , and let

$$I_A = \ker \varepsilon_A$$

be its augmentation ideal. The elements of  $G(D)$  are the morphisms of profinite  $k$ -algebras

$$f : A \longrightarrow D.$$

The condition  $G(\delta)(f) = 1$  is equivalent to

$$\delta \circ f = \varepsilon_A.$$

Since

$$x - \varepsilon_A(x) \cdot 1_A \in I_A \quad \text{for every } x \in A,$$

this is equivalent to  $f(I_A) \subset k \cdot d$ . Hence  $\ker(f)$  contains  $I_A^2$ , and therefore also its closure  $\overline{I_A^2}$ . Thus  $f$  induces a continuous linear map

$$f' : I_A / \overline{I_A^2} = \omega_{G/k} \longrightarrow k$$

such that, for every  $x \in A$ ,

$$f(x) = \varepsilon_A(x) \cdot 1_D + f'(\bar{x}) \cdot d,$$

where  $\bar{x}$  denotes the image of  $x - \varepsilon_A(x) \cdot 1_A$  in  $I_A / \overline{I_A^2}$ . It follows that the map

$$f \longmapsto f'$$

defines a bijection from  $\text{Lie}(G)$  onto the topological dual

$$\omega_{G/k}^\dagger$$

of  $\omega_{G/k}$  (cf. 0.2.2).

This bijection respects the group structures. Indeed, let  $f$  and  $g$  be two elements of  $\text{Lie}(G)$ . Their product  $f \cdot g$  is the composite  $h \circ \Delta_A$ , where

$$h : A \widehat{\otimes} A \longrightarrow D$$

is defined by

$$h(b \widehat{\otimes} b') = f(b)g(b').$$

If  $a \in I_A$ , then, by 2.1(ii),

$$\Delta_A(a) - a \widehat{\otimes} 1 - 1 \widehat{\otimes} a \in I_A \widehat{\otimes} I_A.$$

Consequently,

$$(f \cdot g)(a) = f(a) + g(a)$$

(cf. also II, 3.10).

Henceforth we identify

$$\mathrm{Lie}(G) = \omega_{G/k}^\dagger$$

by means of the bijection  $f \mapsto f'$  just described. The group  $\mathrm{Lie}(G)$  is thereby endowed with a  $k$ -module structure.

c) Let  $A^\dagger$  and  $D^\dagger$  be the  $k$ -modules dual to  $A$  and  $D$ , and let

$$\{1_D^\dagger, d^\dagger\}$$

be the basis dual to the basis  $\{1_D, d\}$  of  $D$  over  $k$ ; thus  $1_D^\dagger = \delta$ .<sup>109</sup> Since  $D$  is free of finite rank over  $k$ , the canonical map

$$\mathrm{Hom}_c(A, D) \longrightarrow \mathrm{Hom}_k(D^\dagger, A^\dagger), \quad f \longmapsto {}^t f,$$

is bijective. On the other hand,  ${}^t f$  is determined by the values

$${}^t f(1_D^\dagger) \quad \text{and} \quad {}^t f(d^\dagger) = x.$$

The condition  $G(\delta)(f) = 1$  is equivalent to

$${}^t f(1_D^\dagger) = \varepsilon_A.$$

Moreover, one sees readily that  $f$  is compatible with multiplication if and only if (cf. 2.3)

$$\delta_G(x) = \phi_G(x \otimes 1 + 1 \otimes x). \quad (*)$$

Finally, it is clear that a continuous linear map

$$f : A \longrightarrow D$$

that is compatible with multiplication and satisfies  $\delta \circ f = \varepsilon_A$  sends the unit element of  $A$  to that of  $D$ .<sup>110</sup> The map  $f \mapsto x$  therefore allows us to identify  $\mathrm{Lie}(G)$  with the set of “primitive elements” of  $\mathbf{H}(G)$ , that is, the elements  $x \in \mathbf{H}(G)$  satisfying relation (\*). If  $x$  and  $y$  are two such elements, then

$$\begin{aligned} \delta_G(xy) &= \delta_G(x)\delta_G(y) \\ &= \phi_G((x \otimes 1 + 1 \otimes x)(y \otimes 1 + 1 \otimes y)) \\ &= \phi_G(xy \otimes 1 + x \otimes y + y \otimes x + 1 \otimes xy), \end{aligned}$$

whence

$$\delta_G(xy - yx) = \phi_G((xy - yx) \otimes 1 + 1 \otimes (xy - yx)).$$

This shows that the  $k$ -module  $\mathrm{Lie}(G)$  is identified with a Lie subalgebra of  $\mathbf{H}(G)$ ; we call  $\mathrm{Lie}(G)$  the *Lie algebra* of  $G$ .<sup>111</sup>

<sup>109</sup>The French authority prints  $1_D^*$  and  $d^*$  in the two occurrences immediately below, after defining  $1_D^\dagger$  and  $d^\dagger$  as the dual basis. The consistent  $\dagger$ -notation is used here.

<sup>110</sup>Indeed,  $\delta \circ f = \varepsilon_A$  implies  $f(1) = 1 + \lambda d$ , with  $\lambda \in k$ , and then

$$f(1) = f(1)^2 = 1 + 2\lambda d$$

gives  $\lambda = 0$ .

<sup>111</sup>Comparing with VII<sub>A</sub>, 2.5, we see that if  $K$  is a group scheme of finite type over  $k$  and  $G$  is the formal completion of  $K$  at the identity—that is, if  $\mathcal{A}(G)$  is the completion of the local ring  $\mathcal{O}_{K,e}$  for the  $\mathfrak{m}$ -adic topology, where

$$\mathfrak{m} = \ker(\varepsilon : \mathcal{O}_{K,e} \longrightarrow k)$$

—then  $\mathbf{H}(G)$  is identified with the algebra of distributions  $U(K)$ , and  $\mathrm{Lie}(G)$  with  $\mathrm{Lie}(K)$ . The hypothesis that  $K$  is of finite type over  $k$  is used to ensure that  $\mathfrak{m}/\mathfrak{m}^2$  is a  $k$ -module of finite length, and hence discrete, so that its topological dual coincides with its ordinary dual. In particular, when  $G$  is a finite formal  $k$ -group—that is, when  $\mathcal{A}(G)$  is a finite  $k$ -module—in which case  $G$  may also be regarded as the  $k$ -group scheme  $\mathrm{Spec} \mathcal{A}(G)$ , the two definitions of  $\mathrm{Lie}(G)$  coincide.

**2.6.1**

Let  $k$  now be an arbitrary pseudocompact ring, and let  $G$  be a formal group over  $k$ . We call the functor  $\mathrm{Lie}(G)$  that assigns to every object  $C$  of  $\mathbf{Alf}/k$  the  $C$ -Lie algebra of the  $C$ -formal group

$$G' = G \widehat{\otimes}_k C$$

the  $\mathbf{O}_k$ -Lie algebra of  $G$ .<sup>112</sup> Put

$$A' = A \widehat{\otimes}_k C.$$

Since  $I_A$  is a direct factor of  $A$ , we have

$$I_{A'} = I_A \widehat{\otimes}_k C = I_A \widehat{\otimes}_A A'.$$

Moreover,

$$\omega_{G/k} = I_A \widehat{\otimes}_A k, \quad \omega_{G'/C} = I_{A'} \widehat{\otimes}_{A'} C.$$

It follows that

$$\omega_{G'/C} = \omega_{G/k} \widehat{\otimes}_A C,$$

and hence

$$\begin{aligned} \mathrm{Lie}(G)(C) &= \mathrm{Hom}_c(\omega_{G/k} \widehat{\otimes}_A C, C), \\ \text{i.e.} \quad \mathrm{Lie}(G) &= \mathbf{V}_k^f(\omega_{G/k}). \end{aligned}$$

With the notation of 1.2.3.B, Proposition 1.2.3.E therefore shows that  $\mathrm{Lie}(G)$  is flat over  $\mathbf{O}_k$  if and only if  $\omega_{G/k}$  is a projective pseudocompact  $k$ -module.

**2.6.2**

Conversely, every Lie algebra  $\mathbf{L}$  over  $\mathbf{O}_k$  defines a  $k$ -group functor. Indeed, let  $U(\mathbf{L})$  denote the functor assigning to every object  $C$  of  $\mathbf{Alf}/k$  the enveloping algebra

$$U(\mathbf{L}(C))$$

of the  $C$ -Lie algebra  $\mathbf{L}(C)$ . By VII<sub>A</sub>, 3.2.2,  $U(\mathbf{L})$  is a bialgebra over  $\mathbf{O}_k$ , and hence, by 2.2, determines the  $k$ -group functor

$$\mathrm{Spf}^* U(\mathbf{L}),$$

which from now on we denote by  $\mathcal{G}(\mathbf{L})$ . Thus  $\mathcal{G}(\mathbf{L})(C)$  is the group of elements  $z \in U(\mathbf{L}(C))$  having augmentation 1 and satisfying

$$\Delta_{U(\mathbf{L}(C))}(z) = z \otimes z.$$

When  $\mathbf{L}$  is flat over  $\mathbf{O}_k$ , we have the following proposition.

**Proposition.**<sup>113</sup> *Let  $\mathbf{L}$  be a flat Lie algebra over  $\mathbf{O}_k$ .*

- (i)  $\mathcal{G}(\mathbf{L})$  is a formal group topologically flat over  $k$ , whose  $\mathbf{O}_k$ -bialgebra is  $U(\mathbf{L})$ .
- (ii)  $\mathcal{G}(\mathbf{L})$  is infinitesimal.
- (iii) For every object  $C$  of  $\mathbf{Alf}/k$ ,  $\mathrm{Lie}(\mathcal{G}(\mathbf{L}))(C)$  is identified with the set

$$\mathrm{Prim} U(\mathbf{L}(C)) = \{x \in U(\mathbf{L}(C)) \mid \varepsilon(x) = 0, \quad \Delta(x) = x \otimes 1 + 1 \otimes x\}$$

of primitive elements of  $U(\mathbf{L}(C))$ . In particular, there is a natural morphism of Lie algebras over  $\mathbf{O}_k$ ,

$$\tau_{\mathbf{L}} : \mathbf{L} \longrightarrow \mathrm{Lie}(\mathcal{G}(\mathbf{L})).$$

<sup>112</sup>The editors expanded the original in what follows.

<sup>113</sup>The editors set off parts (i) and (ii) and added part (iii), which will be used in 2.6.3 and 3.3.2.

*Proof.* The hypothesis that  $\mathbf{L}$  is flat over  $\mathbf{O}_k$  means that, for every morphism  $C \rightarrow D$  in  $\mathbf{Alf}/k$ ,

$$\mathbf{L}(D) = \mathbf{L}(C) \otimes_C D,$$

and that, for every local component  $C'$  of  $C$ , the  $C'$ -module  $\mathbf{L}(C')$  is free. The first condition, together with the universal properties of tensor products and enveloping algebras, gives

$$U(\mathbf{L}(D)) = U(\mathbf{L}(C)) \otimes_C D.$$

The second condition and the Poincaré–Birkhoff–Witt theorem (cf. Bourbaki, *Groupes et algèbres de Lie*, I, 2.7) show that  $U(\mathbf{L}(C'))$  is a free  $C'$ -module. Consequently, the bialgebra  $U(\mathbf{L})$  is flat over  $\mathbf{O}_k$ , proving (i).

To prove (ii) and (iii), we may suppose that  $k$  is Artinian. Put

$$L = \mathbf{L}(k), \quad U = U(L), \quad U_0 = k \cdot 1_U,$$

and let  $U^+$  be the two-sided ideal of  $U$  generated by the image of  $L$ . For every  $n \geq 1$ , put

$$U_n = \{u \in U \mid \Delta_U(u) - u \otimes 1 \in U_{n-1} \otimes U^+\}.$$

By 1.3.6, it is enough to show that  $U$  is the union of the  $U_n$ . Identify  $L$  with its canonical image in  $U$ . Then  $L \subset U_1$ . If  $x_1, \dots, x_n \in L$ , then

$$\Delta_U(x_1 \cdots x_n) = (x_1 \otimes 1 + 1 \otimes x_1) \cdots (x_n \otimes 1 + 1 \otimes x_n).$$

Induction on  $n$  therefore shows that  $x_1 \cdots x_n \in U_n$ , and hence

$$U = \bigcup_n U_n.$$

This proves (ii).

Finally, let

$$D = k[d]/(d^2)$$

be the algebra of dual numbers over  $k$ . By hypothesis,

$$\mathbf{L}(D) \simeq L \otimes_k D, \quad U(\mathbf{L}(D)) \simeq U \otimes_k D.$$

It follows that  $\text{Lie}(\mathcal{G}(\mathbf{L}))(k)$  is identified with the set of elements

$$z = 1 + xd \in U \oplus Ud \quad (x \in U)$$

such that  $\varepsilon(z) = 1$  and  $\Delta(z) = z \otimes z$ . These conditions are equivalent to

$$\varepsilon(x) = 0, \quad \Delta(x) = x \otimes 1 + 1 \otimes x,$$

that is, to  $x \in \text{Prim } U$ . In particular,

$$\tau_{\mathbf{L}} : x \longmapsto 1 + dx$$

is a morphism of Lie algebras over  $\mathbf{O}_k$  from  $\mathbf{L}$  to  $\text{Lie}(\mathcal{G}(\mathbf{L}))$ .

## 2.6.3

If  $\mathbf{L}$  is a flat Lie algebra over  $\mathbf{O}_k$ , the formal group  $\mathcal{G}(\mathbf{L})$  can be characterized by a universal property.<sup>114</sup> Indeed, every morphism

$$\phi : \mathcal{G}(\mathbf{L}) \longrightarrow G$$

to a formal group  $G$  induces a morphism

$$\mathrm{Lie}(\phi) : \mathrm{Lie}(\mathcal{G}(\mathbf{L})) \longrightarrow \mathrm{Lie}(G).$$

Composing it with

$$\tau_{\mathbf{L}} : \mathbf{L} \longrightarrow \mathrm{Lie}(\mathcal{G}(\mathbf{L}))$$

(cf. 2.6.2) gives a morphism

$$\phi' : \mathbf{L} \longrightarrow \mathrm{Lie}(G).$$

We then have:

**Proposition.** *Let  $G$  be a formal  $k$ -group and let  $\mathbf{L}$  be a flat Lie algebra over  $\mathbf{O}_k$ . The map  $\phi \mapsto \phi'$  defined above is a bijection*

$$\mathrm{Hom}_{\mathrm{Grf}/k}(\mathcal{G}(\mathbf{L}), G) \xrightarrow{\sim} \mathrm{Hom}_{\mathrm{Lie}}(\mathbf{L}, \mathrm{Lie}(G)),$$

where the right-hand side is the set of morphisms of Lie algebras over  $\mathbf{O}_k$  from  $\mathbf{L}$  to  $\mathrm{Lie}(G)$ .

*Proof.* We may immediately reduce to the case in which  $k$  is Artinian. Put  $L = \mathbf{L}(k)$ . By 2.3.1,

$$\mathrm{Hom}_{\mathrm{Grf}/k}(\mathcal{G}(\mathbf{L}), G)$$

is in bijection with the set of unital algebra morphisms

$$h : U(L) \longrightarrow \mathbf{H}(G)$$

for which the following diagram is commutative:

$$\begin{array}{ccccc} U(L) & \xrightarrow{h} & \mathbf{H}(G) & \xrightarrow{\delta_G} & \mathbf{H}(G \times G) \\ \Delta_{U(L)} \downarrow & & & \nearrow \varphi_G & \\ U(L) \otimes U(L) & \xrightarrow{h \otimes h} & \mathbf{H}(G) \otimes \mathbf{H}(G) & & \end{array} .$$

Now  $h$  is determined by its restriction to  $L$ , which is a morphism from  $L$  to the Lie algebra underlying  $\mathbf{H}(G)$ . Commutativity of the diagram says precisely that  $h$  sends  $L$  into the subset of  $\mathbf{H}(G)$  formed by the elements  $x$  satisfying

$$\delta_G(x) = \varphi_G(x \otimes 1 + 1 \otimes x).$$

This subset is exactly  $\mathrm{Lie}(G)$ ; cf. 2.6(c). The proposition follows.

<sup>114</sup>The editors modified what follows to take advantage of the addition made in 2.6.2.

## 2.7

We conclude these generalities with a statement that goes back to S. Lie and that will be used in 5.1. A *formal monoid* over  $k$  is, by definition, a pair  $(M, m)$  consisting of a formal variety  $M$  and a morphism

$$m : M \times M \longrightarrow M$$

such that  $m(C)$  makes  $M(C)$  a monoid for every object  $C$  of  $\mathbf{Alf}/k$ .<sup>115</sup> In particular, the “unit section,” which assigns to every object  $C$  the identity element of  $M(C)$ , defines a section

$$\varepsilon_M : \mathrm{Spf}(k) \longrightarrow M$$

of the canonical projection  $M \rightarrow \mathrm{Spf}(k)$ . We shall say that the formal monoid  $M$  is *infinitesimal* if  $\varepsilon_M$  induces a bijection of the underlying sets.

**Proposition.** *Every topologically flat infinitesimal formal  $k$ -monoid  $M$  is a formal  $k$ -group.*<sup>116</sup>

*Proof.* We must show that  $M(C)$  is a group for every object  $C$  of  $\mathbf{Alf}/k$ . We therefore reduce immediately to the case in which  $k$  is Artinian. Let

$$U = \mathbf{H}(M)$$

be the coalgebra of  $M$  (1.3.5). The multiplication

$$m : M \times M \longrightarrow M$$

induces a morphism of coalgebras

$$m_U : U \otimes U \longrightarrow U$$

that makes  $U$  an associative algebra over  $k$ . The identity element of this algebra is the image of the identity element of  $k$  under the map  $k \rightarrow U$  induced by the unit section  $\varepsilon_M$  of  $M$ . Likewise, the projection

$$M \longrightarrow \mathrm{Spf}(k)$$

induces a homomorphism

$$\varepsilon_U : U \longrightarrow k;$$

we write  $U^+$  for the kernel of  $\varepsilon_U$ .

We must show that there is an antipode, that is, a morphism of coalgebras

$$c_U : U \longrightarrow U$$

such that, for every  $u \in U$ ,

$$(m_U \circ (c_U \otimes \mathrm{id}_U) \circ \Delta_U)(u) = \varepsilon_U(u) 1_U. \quad (*)$$

<sup>115</sup>Here and in what follows, the editors have written “monoid” instead of “monoid with identity element” (recall that a monoid is, by definition, a set equipped with an associative composition law and possessing an identity element).

<sup>116</sup>By the equivalence of categories in 1.3.5.D, a monoid in the category of topologically flat formal  $k$ -varieties “is the same thing” as a monoid in the category of flat cocommutative  $\mathbf{O}_k$ -coalgebras, that is, a cocommutative  $\mathbf{O}_k$ -bialgebra (in the usual sense, hence not necessarily equipped with an antipode). Moreover, by 1.3.6, the hypothesis that  $M$  be infinitesimal is equivalent to saying that the corresponding bialgebra is connected. Thus, if  $k$  is an Artinian ring, the proposition is equivalent to the following statement: every connected cocommutative  $k$ -bialgebra that is flat over  $k$  is a  $k$ -Hopf algebra, that is, it possesses an antipode. In fact, the cocommutativity hypothesis is superfluous; cf. the proof.

*Source note.* In (\*), the French re-edition prints  $c_n$  in place of  $c_U$ , applies the left-hand side to  $x$ , and retains  $u$  on the right after quantifying over  $u$ . At the end of the proof it again prints  $m_U \circ (c_n \otimes \text{id}_U) \circ \Delta_U = \eta$  while calling this the right-inverse identity. The global identity above therefore uses  $c_U$  and the single variable  $u$ ; below,  $c_n$  denotes the coherent map at the  $n$ -th stage of the induction, and the final right-inverse identity has the tensor factors in the order required by that assertion.

Let  $(U_n)$  be the filtration of  $U$  defined in 1.3.6, and put

$$U_n^+ = U^+ \cap U_n.$$

Since  $M$  is infinitesimal,  $U^+$  is the union of the submodules  $U_n^+$ .<sup>117</sup> Set

$$c_0(1) = 1$$

and

$$c_1(x) = -x \quad \text{if } x \in U_1^+,$$

that is, if  $x$  is primitive. Suppose that

$$c_{n-1} : U_{n-1} \longrightarrow U$$

has been constructed so as to satisfy (\*), with  $c_{n-1}$  in place of  $c_U$ , for every  $x \in U_{n-1}$ , and let  $x \in U_n^+$ . By the proof of Lemma 1.3.6.A,

$$\Delta_U(x) - x \otimes 1 \in U_{n-1} \otimes U^+;$$

this is where the hypothesis that  $U$  is flat over  $k$  enters. We may therefore write

$$\Delta_U(x) = x \otimes 1 + \sum_i y_i \otimes z_i, \quad y_i \in U_{n-1},$$

and then set

$$c_n(x) = - \sum_i c_{n-1}(y_i) z_i.$$

Together with  $c_n(1) = 1$ , this prescription extends  $c_{n-1}$  coherently to  $U_n$ . In this way we obtain a  $k$ -linear map

$$c : U \longrightarrow U$$

that is the left inverse of  $\text{id}_U$  for the monoid law on  $\text{End}_k(U)$  defined by

$$f \cdot g = m_U \circ (f \otimes g) \circ \Delta_U;$$

the identity element is the map

$$\eta : u \longmapsto \varepsilon_U(u) 1_U.$$

It follows that  $c$  is uniquely determined and is also the right inverse of  $\text{id}_U$ ; that is,

$$m_U \circ (\text{id}_U \otimes c) \circ \Delta_U = \eta,$$

without any assumption that  $U$  be cocommutative. Thus  $c_U := c$  is the required antipode, and  $M$  is a formal  $k$ -group.

---

<sup>117</sup>The original continued as follows: “one then shows easily, by induction on  $n$ , that there exists one and only one linear map  $c_n : U_n \rightarrow U_n$  such that the composite

$$m_U \circ (c_n \otimes \text{id}_U) \circ \Delta_U : U_n^+ \longrightarrow U$$

is zero.” The editors have supplied the details of the proof, which rests on that of Lemma 1.3.6.A.



## 2.8. Unipotent group schemes over a field

118

Let  $k$  be a field. Recall that a  $k$ -group scheme  $G$  is called *unipotent* if it satisfies the following two conditions (cf. [DG70], § IV.2, Proposition 2.5):

- (a)  $G$  is affine.
- (b) Every simple  $\mathcal{O}(G)$ -comodule is trivial. Explicitly, if

$$\rho : V \longrightarrow V \otimes_k \mathcal{O}(G)$$

is an  $\mathcal{O}(G)$ -comodule structure on a nonzero  $k$ -vector space  $V$ , and there is no nonzero proper subspace  $W \subset V$  such that

$$\rho(W) \subset W \otimes_k \mathcal{O}(G),$$

then  $V$  is one-dimensional and

$$\rho(v) = v \otimes 1 \quad \text{for every } v \in V.$$

According to the cited result, when  $G$  is of finite type over  $k$ , this is equivalent to the definition given in Exposé XVII, § 1:  $G$  has a finite composition series whose successive quotients are isomorphic to  $k$ -subgroups of  $\mathbb{G}_{a,k}$ .

For every affine  $k$ -group scheme  $G$ , the comultiplication of  $\mathcal{O}(G)$  endows the pseudo-compact  $k$ -module

$$A = \mathcal{O}(G)^*$$

with the structure of a profinite  $k$ -algebra, not necessarily commutative; its unit  $1_A$  is the augmentation

$$\varepsilon : \mathcal{O}(G) \longrightarrow k.$$

Set

$$I = \{f \in A \mid f(1_{\mathcal{O}(G)}) = 0\}.$$

Then  $I$  is a two-sided ideal of  $A$ , and

$$A = k \cdot 1_A \oplus I;$$

cf. 1.3.6.

Let  $V$  be a finite-codimensional subspace of  $A$ , and consider the continuous  $k$ -bilinear map

$$\varphi : A \times A \longrightarrow A/V, \quad (a, b) \longmapsto ab + V.$$

By Lemma 0.3.1, there are finite-codimensional subspaces  $L_1, L_2 \subset A$  such that  $V$  contains  $AL_2$  and  $L_1A$ . Thus

$$L = L_1 \cap L_2$$

is finite-codimensional, and  $V$  contains the two-sided ideal  $ALA$ , which is itself finite-codimensional. Hence the finite-codimensional two-sided ideals form a neighborhood basis of zero in  $A$ .

It follows that an  $\mathcal{O}(G)$ -comodule is the same thing as a continuous  $A$ -module: an  $A$ -module  $V$ , equipped with the discrete topology, for which the map

$$A \times V \longrightarrow V$$

---

<sup>118</sup>The editors added this subsection.

is continuous. Every such module is the union of finite-dimensional  $A$ -submodules  $V_i$ , each of which is a module over a finite-dimensional quotient  $k$ -algebra  $A_i$  of  $A$ .

Consequently, a simple continuous  $A$ -module  $M$  is finite-dimensional over  $k$  and is a faithful simple module over the finite-dimensional algebra  $A/\text{Ann}(M)$ . The latter is therefore a finite-dimensional simple  $k$ -algebra, so  $\text{Ann}(M)$  is an open maximal ideal of  $A$ .

Conversely, let  $P$  be an open prime ideal of  $A$ .<sup>119</sup> Then  $A/P$  is a finite-dimensional  $k$ -algebra in which the zero ideal is prime, and hence is a finite-dimensional simple  $k$ -algebra. Up to isomorphism, there is therefore a unique simple continuous  $A$ -module with annihilator  $P$ . Thus

$$M \mapsto \text{Ann}(M)$$

is a bijection from the isomorphism classes of simple continuous  $A$ -modules to the open prime ideals of  $A$ .

In particular, the  $A$ -module  $A/I$ , which is one-dimensional over  $k$ , is called the *trivial module*. It corresponds to the trivial one-dimensional  $\mathcal{O}(G)$ -comodule  $V$ , for which

$$\rho(v) = v \otimes 1_{\mathcal{O}(G)} \quad \text{for every } v \in V.$$

We have proved:

**Proposition 2.8.1.** *Let  $k$  be a field, let  $G$  be an affine  $k$ -group scheme, and put  $A = \mathcal{O}(G)^*$ .*

- (i)  *$G$  is unipotent if and only if  $I$  is the unique open prime ideal of  $A$ .<sup>120</sup>*
- (ii) *In particular, suppose that  $G$  is commutative, so that*

$$\mathcal{O}(G) = \mathbf{H}(D(G)),$$

*where  $D(G) = \text{Spf}(A)$  is the Cartier dual of  $G$ . Then  $G$  is unipotent if and only if  $D(G)$  is infinitesimal.*

## 2.9. Pointed cocommutative Hopf algebras over a field

121

Let  $k$  be a field, let  $C$  be a  $k$ -coalgebra, and endow

$$A = C^*$$

with the structure of a profinite  $k$ -algebra, not necessarily commutative, described in 2.8. By 0.2.2,

$$C = A^\dagger = \text{Hom}_c(A, k).$$

The map

$$D \mapsto D^\perp = \{f \in A \mid f(D) = 0\}$$

is therefore a bijection from the subcoalgebras of  $C$  to the closed two-sided ideals of  $A$  (which we shall simply call ideals below). Its inverse is

$$I \mapsto I^\perp = \{x \in C = A^\dagger \mid x(I) = 0\}.$$

<sup>119</sup>Recall that a two-sided ideal  $P$  of a ring  $A$  is called prime if, in  $A/P$ , the product of any two nonzero two-sided ideals is nonzero.

<sup>120</sup>Equivalently,  $k \cdot 1_{\mathcal{O}(G)}$  is the unique simple  $k$ -subcoalgebra of  $\mathcal{O}(G)$ ; see, for example, [Ab80], 3.1.4. The editors also note a misprint in the cited work, p. 130, line 4:  $M \simeq C^*/\text{ann } M$  should be replaced by  $C^*/\text{ann } M \simeq \text{End}_k(M)$ .

<sup>121</sup>The editors added this subsection to express Proposition 2.5.2, together with the variant noted in editorial note 100, in the language of cocommutative Hopf algebras.

Since every maximal closed ideal is a maximal open ideal (cf. 0.2.1), every subcoalgebra contains a simple subcoalgebra, necessarily finite-dimensional.

Recall that a subcoalgebra  $D \subset C$  is called *irreducible* if it contains a unique simple subcoalgebra  $S_0$ . Equivalently,

$$\mathfrak{m}_0 = S_0^\perp$$

is the unique open maximal ideal containing  $D^\perp$ , or, equivalently,

$$D^\perp + \mathfrak{m} = A \quad \text{for every open maximal ideal } \mathfrak{m} \neq \mathfrak{m}_0.$$

This applies in particular when  $D = S_0$ .

Let  $\Sigma_0$  be the sum of all irreducible subcoalgebras  $C_i$  containing  $S_0$ . It is plainly a subcoalgebra, and it is irreducible because, for every  $\mathfrak{m} \neq \mathfrak{m}_0$ , 0.2.D gives

$$\mathfrak{m} + \bigcap_i C_i^\perp = \bigcap_i (\mathfrak{m} + C_i^\perp) = A.$$

The coalgebra  $\Sigma_0$  is called the *irreducible component of  $C$  corresponding to  $S_0$* . *The French prints  $C_0$  in this last phrase; the preceding definition makes  $S_0$  the intended simple subcoalgebra.*

A coalgebra  $C$  is called *pointed* if every simple  $k$ -subcoalgebra of  $C$  is one-dimensional. Equivalently,

$$A/\mathfrak{m} = k$$

for every open maximal ideal  $\mathfrak{m}$  of  $A$ . Recall also that  $C$  is called *connected* if it is irreducible and pointed. If  $C$  is a bialgebra, then it is connected if and only if it is irreducible, since  $k \cdot 1_C$  is a simple subcoalgebra.

Suppose from now on that  $C$  is cocommutative. Then  $A$  is commutative and hence is the product of its local components

$$A_{\mathfrak{m}}, \quad \mathfrak{m} \in \Upsilon(A);$$

cf. 0.1.1. Put

$$S_{\mathfrak{m}} = \mathfrak{m}^\perp \simeq (A/\mathfrak{m})^*,$$

and let  $\Sigma_{\mathfrak{m}}$  be its irreducible component. This component can be described explicitly. Let

$$J_{\mathfrak{m}} = \ker(A \longrightarrow A_{\mathfrak{m}}).$$

Then  $J_{\mathfrak{m}} \subset \mathfrak{m}$ , and it is the smallest closed ideal  $I \subset \mathfrak{m}$  satisfying

$$I + \mathfrak{m}' = A \quad \text{for every open maximal ideal } \mathfrak{m}' \neq \mathfrak{m}.$$

Indeed, any such  $I$  contains  $A_{\mathfrak{m}'}$  for every  $\mathfrak{m}' \neq \mathfrak{m}$ , and hence contains  $J_{\mathfrak{m}}$ . Since

$$A = J_{\mathfrak{m}} \oplus A_{\mathfrak{m}},$$

we obtain

$$\Sigma_{\mathfrak{m}} = J_{\mathfrak{m}}^\perp \simeq A_{\mathfrak{m}}^\dagger.$$

We can now prove:

**Proposition 2.9.1.** *Let  $k$  be a field.*

- (i) Let  $G$  be a formal  $k$ -group all of whose residue fields are equal to  $k$ . Then  $G_e$  is the constant  $k$ -group  $M_k$ , where

$$M = G(k) = \{x \in \mathbf{H}(G) \mid \varepsilon(x) = 1, \quad \Delta(x) = x \otimes x\},$$

and  $\mathbf{H}(G)$  is the semidirect product of  $\mathbf{H}(G_0)$  by the group algebra  $kM$  (cf. 2.4.B).

- (ii) Equivalently, let  $H$  be a pointed cocommutative Hopf  $k$ -algebra. Then  $H$  is the semidirect product of the irreducible component  $H_0$  of the unit  $1_H$  by  $kM$ , where

$$M = \{x \in H \mid \varepsilon(x) = 1, \quad \Delta(x) = x \otimes x\}.$$

*Proof.* We first prove (i). Since every residue field of  $G$  is equal to  $k$ , the projection

$$\pi : G \longrightarrow G_e$$

has a section  $s : G_e \rightarrow G$ , defined at every point  $g \in G$  by

$$\mathcal{O}_{G,g} \longrightarrow \kappa(g) = k.$$

Moreover, for all  $g, h \in G$ , the algebra

$$\mathcal{O}_{G,g} \widehat{\otimes}_k \mathcal{O}_{G,h}$$

is local with residue field  $k$ . Thus  $s \times s$  is a section of

$$\pi \times \pi : G \times G \longrightarrow (G \times G)_e = G_e \times G_e.$$

Since  $\pi$  is a group morphism,

$$\pi \circ m_G \circ (s \times s) = m_{G_e} = \pi \circ s \circ m_{G_e}.$$

As  $\pi$  is an epimorphism, it follows that  $s$  is a group morphism. Hence

$$G = G_0 \rtimes G_e,$$

and  $\mathbf{H}(G)$  is the semidirect product of  $\mathbf{H}(G_0)$  by  $\mathbf{H}(G_e)$ .

Since every residue field of  $G$  is  $k$ ,  $\mathbf{H}(G_e)$  is the group algebra  $kM$ , where

$$M = G_e(k);$$

cf. 2.5.A. Finally, because  $G_0$  is infinitesimal, the map

$$G(k) \longrightarrow G_e(k)$$

is injective. It is also surjective, since it has the section induced by  $s$ . Thus  $M = G(k)$ , proving (i).

For (ii), it remains only to show that

$$H_0 = \mathbf{H}(G_0).$$

The unit  $1_H$  of  $H$  is the augmentation

$$\varepsilon_{\mathcal{A}} : \mathcal{A} \longrightarrow k,$$

which is the identity section of  $G(k)$ . Therefore the local component of the affine algebra  $\mathcal{A}(G)$  corresponding to  $H_0$  is precisely  $\mathcal{A}(G_0)$ . By the description above,

$$H_0 = \mathcal{A}(G_0)^\dagger = \mathbf{H}(G_0).$$

This proves the proposition.

**Remarks 2.9.2.**

- (a) The preceding proposition, which is implicit in 2.5.2, was obtained independently by B. Kostant (cf. [Sw69], Preface). Combined with Cartier's Theorem 3.3 below (cf. [Ca62], § 12, Theorem 3), which was also obtained by Kostant (cf. [Sw69], loc. cit.), this result is often called the *Cartier–Gabriel–Kostant theorem*.
- (b) In the form (ii), Proposition 2.9.1 was extended by R. G. Heyneman and M. E. Sweedler to the case in which  $H$  is assumed to be pointed and the direct sum of its irreducible components, but is not necessarily cocommutative; see [HS69], Theorem 3.5.8.

### 3. Phenomena specific to characteristic 0

Throughout Section 3, we assume that the pseudocompact ring  $k$  contains the field of rational numbers  $\mathbb{Q}$ .

#### 3.1. Lemma

**Lemma 3.1.** *Let  $C$  be a commutative unital  $\mathbb{Q}$ -algebra, and let  $L$  be a Lie algebra over  $C$  whose underlying  $C$ -module is free. Then the canonical map*

$$L \longrightarrow U(L)$$

*identifies  $L$  with the set of primitive elements of  $U(L)$ .*

*Proof.* Identify  $L$  with its canonical image in  $U(L)$ . Let  $I$  be a totally ordered set, and let

$$(x_i)_{i \in I}$$

be a basis of  $L$  indexed by  $I$ . Denote by  $\mathbb{N}^{(I)}$  the set of families

$$n = (n_i)_{i \in I}$$

of nonnegative integers for which  $n_i = 0$  except possibly at finitely many indices

$$i_1 < i_2 < \cdots < i_s;$$

these indices depend on  $n$ . Put

$$x^n = x_{i_1}^{n_{i_1}} x_{i_2}^{n_{i_2}} \cdots x_{i_s}^{n_{i_s}}, \quad n! = (n_{i_1}!)(n_{i_2}!) \cdots (n_{i_s}!).$$

By the Poincaré–Birkhoff–Witt theorem, the elements  $x^n$  form a basis of  $U(L)$ . Moreover,

$$\Delta_{U(L)} \left( \frac{x^n}{n!} \right) = \sum_{0 \leq m \leq n} \frac{x^m}{m!} \otimes \frac{x^{n-m}}{(n-m)!}. \quad (*)$$

Here the sum ranges over all  $m \in \mathbb{N}^{(I)}$  such that  $0 \leq m \leq n$ , that is,

$$0 \leq m_i \leq n_i \quad \text{for every } i.$$

It follows immediately that

$$\Delta_{U(L)}(u) = u \otimes 1 + 1 \otimes u$$

if and only if  $u$  is a linear combination of the  $x_i$ . This proves the lemma.

## 3.2

Suppose now that  $C$  is Artinian, with radical  $\mathfrak{r}$ . For every  $C$ -algebra  $U$  (associative and unital), the ideal  $\mathfrak{r}U$  therefore consists of nilpotent elements. If  $x \in \mathfrak{r}U$ , put

$$\exp_U(x) = 1 + x + \frac{x^2}{2!} + \cdots.$$

<sup>122</sup> This gives a bijection from  $\mathfrak{r}U$  onto  $1 + \mathfrak{r}U$ ; its inverse sends  $1 + y \in 1 + \mathfrak{r}U$  to

$$\log_U(1 + y) = y - \frac{y^2}{2} + \frac{y^3}{3} - \cdots.$$

Moreover, the map  $\exp_U$  is plainly functorial in  $U$ .<sup>123</sup>

Still assuming  $C$  Artinian, suppose that  $U$  is endowed with a bialgebra structure over  $C$  (cf. 2.2). For every primitive element  $x \in \mathfrak{r}U$  (cf. VII<sub>A</sub>, 3.2.3), we then have

$$\begin{aligned} \Delta_U(\exp_U(x)) &= \exp_{U \otimes_C U}(\Delta_U(x)) \\ &= \exp_{U \otimes_C U}(x \otimes 1 + 1 \otimes x) \\ &= \exp_{U \otimes_C U}(x \otimes 1) \exp_{U \otimes_C U}(1 \otimes x) \\ &= (\exp_U(x) \otimes 1)(1 \otimes \exp_U(x)) \\ &= \exp_U(x) \otimes \exp_U(x). \end{aligned}$$

Thus the bijection  $\exp_U$  sends a primitive element of  $\mathfrak{r}U$  to an element  $z \in 1 + \mathfrak{r}U$  satisfying

$$\Delta_U(z) = z \otimes z.$$

Conversely, if  $z$  satisfies these conditions, then, on putting  $x = \log_U(z)$ , the preceding calculation shows that

$$\exp_{U \otimes_C U}(x \otimes 1 + 1 \otimes x) = z \otimes z = \Delta_U(\exp_U(x)) = \exp_{U \otimes_C U}(\Delta_U(x));$$

hence

$$x \otimes 1 + 1 \otimes x = \Delta_U(x).$$

<sup>124</sup> Furthermore, if  $z \in 1 + \mathfrak{r}U$  satisfies  $\Delta_U(z) = z \otimes z$ , then

$$\varepsilon_U(z)^2 = \varepsilon_U(z).$$

Since  $\varepsilon_U(z)$  is invertible (because  $z$  is invertible, and  $\mathfrak{r}$  is nilpotent), it follows that  $\varepsilon_U(z) = 1$ .

In particular, let  $L$  be a Lie algebra flat over  $C$ , take for  $U$  the enveloping algebra  $U(L)$  of  $L$  over  $C$ , and identify  $L$  with its canonical image in  $U$ . By Lemma 3.1,  $L$  is the set of primitive elements of  $U$ : indeed, on every local component of  $C$ ,  $L$  is a free module. Consider the  $C$ -formal group

$$\mathcal{G}(L) = \mathrm{Spf}^* U(L),$$

---

<sup>122</sup>If  $x, x' \in \mathfrak{r}U$  commute, then

$$\exp_U(x + x') = \exp_U(x) \exp_U(x').$$

<sup>123</sup>That is, for every homomorphism  $\phi : U \rightarrow V$  of  $C$ -algebras,

$$\phi(\exp_U(x)) = \exp_V(\phi(x)).$$

<sup>124</sup>The editors supplied the details of the preceding argument and added the sentence that follows.

whose covariant bialgebra is  $U = U(L)$  (cf. 2.6.2). Let  $\mathfrak{m}$  be a maximal ideal of  $C$ , and put

$$\overline{C} = C/\mathfrak{m}, \quad \overline{U} = U \otimes_C \overline{C}.$$

Since  $\mathcal{G}(L)$  is infinitesimal (loc. cit.), the unit element of  $\overline{U}$  is the only element  $\bar{z} \in \overline{U}$  such that<sup>125</sup>

$$\varepsilon_{\overline{U}}(\bar{z}) = 1, \quad \Delta_{\overline{U}}(\bar{z}) = \bar{z} \otimes \bar{z}.$$

It follows that every  $z \in U$  satisfying

$$\varepsilon_U(z) = 1, \quad \Delta_U(z) = z \otimes z$$

necessarily belongs to  $1 + \mathfrak{r}U$ .

Finally, since

$$L \cap \mathfrak{r}U = \mathfrak{r}L = L \otimes_C \mathfrak{r},$$

we conclude that  $\exp_U$  defines a bijection from  $L \otimes_C \mathfrak{r}$  onto the group  $\mathcal{G}(L)(C)$ . We summarize this as follows.

**Proposition 3.2.** *Let  $k$  be a pseudocompact ring containing  $\mathbb{Q}$ , and let  $\mathbf{L}$  be a flat Lie algebra over  $\mathbf{O}_k$ .*

(i) *For every object  $C$  of  $\mathbf{Alf}/k$ , let  $\mathfrak{r}(C)$  denote its radical. Then the map*

$$\exp_{U(\mathbf{L}(C))} : \mathbf{L}(C) \otimes_C \mathfrak{r}(C) \longrightarrow \mathcal{G}(\mathbf{L})(C)$$

*is bijective and functorial in  $C$  and  $\mathbf{L}$ .*

(ii) *The natural morphism*

$$\mathbf{L} \longrightarrow \mathrm{Lie}(\mathcal{G}(\mathbf{L}))$$

*(cf. 2.6.2) is an isomorphism of Lie algebras over  $\mathbf{O}_k$ .*

126

### 3.2.1

The bijection  $\exp_{U(L(C))}$  allows us, by transport of structure, to define a group law on the set

$$L(C) \otimes_C \mathfrak{r}(C),$$

which, as in 3.2, we identify with a subset of  $U(L(C))$ . If  $x$  and  $y$  are two elements of  $L(C) \otimes_C \mathfrak{r}(C)$ , this law is given by

$$\begin{aligned} x \cdot y &= \log((\exp x)(\exp y)) = \log \left( 1 + \sum_{p+q>0} \frac{x^p y^q}{p! q!} \right) \\ &= \sum_{m \geq 1} \sum_{p_i + q_i > 0} \frac{(-1)^{m-1}}{m} \frac{x^{p_1} y^{q_1}}{p_1! q_1!} \cdots \frac{x^{p_m} y^{q_m}}{p_m! q_m!} \\ &= \sum_{\ell \geq 1} P_\ell(x, y). \end{aligned}$$

<sup>125</sup>The editors added the condition  $\varepsilon_{\overline{U}}(\bar{z}) = 1$ , which was omitted in the original.

<sup>126</sup>The editors added part (ii), which follows immediately from 3.1.

Here  $P_\ell(x, y)$  denotes the sum of the monomials of total degree  $\ell$  in  $x$  and  $y$ . For example,

$$P_1(x, y) = x + y,$$

and

$$\begin{aligned} P_2(x, y) &= \underbrace{\frac{x^2}{2} + xy + \frac{y^2}{2}}_{m=1} - \underbrace{\frac{1}{2}(x^2 + xy + yx + y^2)}_{m=2} \\ &= \frac{1}{2}(xy - yx) = \frac{1}{2}[x, y]. \end{aligned}$$

Moreover,  $P_3(x, y)$  is the sum of the following three terms:

$$\begin{aligned} P_3(x, y) &= \underbrace{\frac{x^3}{6} + \frac{x^2y}{2} + \frac{xy^2}{2} + \frac{y^3}{6}}_{m=1} \\ &\quad - \underbrace{\frac{1}{2}\left(x^3 + \frac{3}{2}x^2y + \frac{1}{2}yx^2 + xyx + yxy + \frac{1}{2}y^2x + \frac{3}{2}xy^2 + y^3\right)}_{m=2} \\ &\quad + \underbrace{\frac{1}{3}(x^3 + x^2y + yx^2 + xyx + yxy + y^2x + xy^2 + y^3)}_{m=3}. \end{aligned}$$

Consequently,

$$P_3(x, y) = \frac{1}{12}(x^2y + yx^2 - 2xyx - 2yxy + y^2x + xy^2) = \frac{1}{12}([y, x], x) + [y, [y, x]].$$

More generally, one can prove the Campbell–Hausdorff formula:<sup>127</sup>

$$\begin{aligned} P_\ell(x, y) &= \sum_{m=1}^{\ell} \frac{(-1)^{m-1}}{m \cdot \ell} \sum_{\substack{p_1, \dots, p_m \\ q_1, \dots, q_{m-1}}} \left( \prod_{i=1}^{m-1} \frac{(\operatorname{ad} x)^{p_i}}{p_i!} \frac{(\operatorname{ad} y)^{q_i}}{q_i!} \right) \frac{(\operatorname{ad} x)^{p_m}}{p_m!}(y) \\ &\quad + \sum_{m=1}^{\ell} \frac{(-1)^{m-1}}{m \cdot \ell} \sum_{\substack{p_1, \dots, p_{m-1} \\ q_1, \dots, q_{m-1}}} \left( \prod_{i=1}^{m-1} \frac{(\operatorname{ad} x)^{p_i}}{p_i!} \frac{(\operatorname{ad} y)^{q_i}}{q_i!} \right) (x). \end{aligned}$$

Here the  $p_j, q_i \in \mathbb{N}$  satisfy

$$p_i + q_i \geq 1 \quad (1 \leq i \leq m-1), \quad p_m + \sum_{i=1}^{m-1} (p_i + q_i) = \ell - 1.$$

In other words, in the sums above every nonzero “Lie monomial” has total degree  $\ell$ . For a proof, see N. Jacobson, *Lie Algebras* (Interscience, 1962), §V.5, or [BLie], II, §6.4, Theorem 2.

### 3.3

If  $G$  is a formal  $k$ -group with affine algebra  $A$ , recall that  $I_A$  denotes the augmentation ideal of  $A$ , and that

$$\omega_{G/k} = I_A/I_A^2 \simeq I_A \hat{\otimes}_A k$$

<sup>127</sup>The editors corrected the formula given in the original, which was erroneous, and added the reference [BLie].



is a pseudocompact  $k$ -module.

**Theorem 3.3 (Cartier).** *Let  $k$  be a pseudocompact ring containing  $\mathbb{Q}$ ,<sup>128</sup> and let  $G$  be a formal  $k$ -group. The following assertions are equivalent:*

- (i) *There exists an  $\mathcal{O}_k$ -Lie algebra  $\mathbf{L}$ , flat over  $\mathcal{O}_k$ , such that  $G$  is isomorphic to  $\mathcal{G}(\mathbf{L})$ . In this case  $\mathbf{L} = \text{Lie}(G)$  by 3.2.*
- (ii) *There exists a projective pseudocompact  $k$ -module  $\omega$  such that the formal variety underlying  $G$  is isomorphic to*

$$\mathbf{V}_k^{f,0}(\omega)$$

(cf. 1.2.5), whose affine algebra is  $k[[\omega]]$ . In this case  $\omega \simeq \omega_{G/k}$ .

- (iii) *The group  $G$  is infinitesimal, and  $\omega_{G/k}$  is a projective pseudocompact  $k$ -module.*
- (iv) *The group  $G$  is infinitesimal and topologically flat over  $k$ .*

*Proof of (i)  $\Rightarrow$  (ii).* Let

$$\omega = \Gamma^*(\mathbf{L})$$

be the pseudocompact  $k$ -module dual to  $\mathbf{L}$  (cf. 1.2.3.D). For every  $C \in \text{Alf}/k$ , we must construct, functorially in  $C$ , an isomorphism

$$\mathbf{V}_k^{f,0}(\omega)(C) \xrightarrow{\sim} \mathcal{G}(\mathbf{L})(C).$$

By 1.2.5, the left-hand side identifies with

$$\text{Hom}_c(\omega, \mathfrak{r}(C)),$$

where  $\mathfrak{r}(C)$  is the radical of  $C$ . This set is naturally isomorphic to the set

$$\text{Hom}_c(\omega \widehat{\otimes}_k C, \mathfrak{r}(C))$$

of continuous  $C$ -linear maps. Since  $\omega \widehat{\otimes}_k C$  is a projective pseudocompact  $C$ -module, the canonical map

$$(\omega \widehat{\otimes}_k C)^\dagger \otimes_C \mathfrak{r}(C) \longrightarrow \text{Hom}_c(\omega \widehat{\otimes}_k C, \mathfrak{r}(C))$$

is bijective (cf. 0.3.6.A). By 1.2.3.E,

$$\mathbf{L}(C) \simeq \mathbf{V}_k^f(\Gamma^*(\mathbf{L}))(C) = (\omega \widehat{\otimes}_k C)^\dagger.$$

Thus  $\mathbf{V}_k^{f,0}(\omega)(C)$  is canonically isomorphic to

$$\mathbf{L}(C) \otimes_C \mathfrak{r}(C),$$

which Proposition 3.2 identifies canonically with  $\mathcal{G}(\mathbf{L})(C)$ . This proves (i)  $\Rightarrow$  (ii).

*Proof of (ii)  $\Rightarrow$  (iii).* Let  $\omega$  be a projective object of  $\text{PC}(k)$ , and let

$$h : k[[\omega]] \xrightarrow{\sim} A$$

be an isomorphism onto the affine algebra of  $G$ . The composite

$$\varepsilon_A \circ h : k[[\omega]] \longrightarrow k$$

<sup>128</sup>The editors removed the hypothesis that  $k$  be local; the proof reduces to that case.

is determined by its restriction  $\lambda : \omega \rightarrow \mathfrak{r}(k)$ . The map

$$x \mapsto x - \lambda(x)$$

from  $\omega$  to the radical of  $k[[\omega]]$  extends, by the universal property of  $k[[\omega]]$ , to an endomorphism  $\ell_\lambda$  of  $k[[\omega]]$ . Since

$$\ell_\lambda \circ \ell_{-\lambda} = \ell_{-\lambda} \circ \ell_\lambda = \text{id},$$

the map  $\ell_\lambda$  is an automorphism. Replacing  $h$  by  $h \circ \ell_\lambda$ , we may assume that  $\varepsilon_A \circ h$  vanishes on the closed ideal  $I$  of  $k[[\omega]]$  generated by  $\omega$ . Then  $h$  induces an isomorphism

$$I/I^2 \xrightarrow{\sim} I_A/I_A^2.$$

Since  $I/I^2 \simeq \omega$ , we obtain

$$\omega_{G/k} \simeq \omega,$$

so  $\omega_{G/k}$  is projective. The formal variety  $\mathbf{V}_k^{f,0}(\omega)$ , and hence  $G$ , is infinitesimal.

*Proof of (iii)  $\Rightarrow$  (i).* Assume that  $G$  is infinitesimal and  $\omega_{G/k}$  is projective. Let  $\mathbf{L}$  be the  $\mathcal{O}_k$ -Lie algebra of  $G$ . By 2.6.1, its underlying  $\mathcal{O}_k$ -module is

$$\mathbf{V}_k^f(\omega_{G/k}).$$

Consequently,  $\mathbf{L}$  is flat over  $\mathcal{O}_k$ , and Proposition 1.2.3.E gives

$$\Gamma^*(\mathbf{L}) \simeq \omega_{G/k}.$$

The proof of (i)  $\Rightarrow$  (ii) therefore identifies the affine algebra of  $\mathcal{G}(\mathbf{L})$  with  $k[[\omega_{G/k}]]$ . By 2.6.3, the identity morphism of  $\mathbf{L}$  corresponds canonically to a morphism

$$\mathcal{G}(\mathbf{L}) \longrightarrow G,$$

and hence to a continuous  $k$ -algebra morphism

$$\varphi : A \longrightarrow k[[\omega_{G/k}]].$$

Let  $I$  be the closed ideal generated by  $\omega_{G/k}$ . Filter  $k[[\omega_{G/k}]]$  by the closures of the powers  $I^n$ , and  $A$  by the closures of the powers  $I_A^n$ . By definition,  $\varphi$  induces the identity on

$$\omega_{G/k} = I_A/I_A^2 = I/I^2.$$

Because  $\omega_{G/k}$  is projective, the canonical projection

$$I_A \longrightarrow \omega_{G/k}$$

has a section  $\tau$ . By the universal property of  $k[[\omega_{G/k}]]$ , this section defines a continuous algebra morphism

$$\psi : k[[\omega_{G/k}]] \longrightarrow A.$$

The composite  $\varphi \circ \psi$  induces the identity on  $\omega_{G/k}$ , and hence on the associated graded algebra of  $k[[\omega_{G/k}]]$ . It follows from [CA], §V.7, Lemma 1 that  $\varphi \circ \psi$  is an isomorphism.<sup>129</sup>

Moreover,  $\psi$  induces an isomorphism

$$I/I^2 \xrightarrow{\sim} I_A/I_A^2,$$

<sup>129</sup>Compare 5.1.5 below, and also [BAC], III, §2.8, Theorem 1 and its corollaries. The editors supplied the details of the original argument that follow.

and hence a surjection on associated graded algebras. Since  $G$  is radicial,  $I_A$  is contained in the radical of  $A$ , so

$$\bigcap_n I_A^n = 0.$$

The cited lemma therefore shows that  $\psi$  is surjective. Since  $\varphi \circ \psi$  is an isomorphism and  $\psi$  is surjective, both  $\psi$  and  $\varphi$  are isomorphisms. This proves (iii)  $\Rightarrow$  (i).

Finally, (i) or (ii) plainly implies (iv), so it remains to prove (iv)  $\Rightarrow$  (ii). We may assume that  $k$  is local, with residue field  $k_0$ . Put

$$G_0 = G \widehat{\otimes}_k k_0, \quad \omega = \omega_{G/k}, \quad \omega_0 = \omega \widehat{\otimes}_k k_0.$$

<sup>130</sup> The pseudocompact  $k_0$ -module  $\omega_0$  has a pseudobasis  $(y_\lambda)_{\lambda \in \Lambda}$ . Let

$$\omega' = \prod_{\lambda \in \Lambda} k_\lambda$$

be the topologically free  $k$ -module with that index set, and lift each  $y_\lambda$  to an element  $x_\lambda \in \omega$ . This gives a continuous  $k$ -linear map

$$f : \omega' \longrightarrow \omega$$

such that

$$f_0 = f \widehat{\otimes}_k k_0$$

is invertible.<sup>131</sup> Since  $\omega'$  is projective,  $f$  lifts to a continuous  $k$ -linear map

$$g : \omega' \longrightarrow I_A$$

with  $\pi \circ g = f$ , where  $\pi : I_A \rightarrow \omega$  is the projection. The universal property of  $k[[\omega']]$  gives a morphism of topological algebras

$$\phi : k[[\omega']] \longrightarrow A.$$

Now

$$\omega' \widehat{\otimes}_k k_0 \simeq \omega \widehat{\otimes}_k k_0 = \omega_{G_0/k_0},$$

and consequently

$$k[[\omega']] \widehat{\otimes}_k k_0 \simeq k_0[[\omega_{G_0/k_0}]].$$

The pair  $k_0, G_0$  satisfies (iii), and the proof of (iii)  $\Rightarrow$  (i) shows that

$$\phi_0 = \phi \widehat{\otimes}_k k_0$$

is invertible. Both  $k[[\omega']]$  and  $A$  are projective pseudocompact  $k$ -modules, so 0.3.4 implies that  $\phi$  itself is invertible. In particular, if  $I$  is the augmentation ideal of  $k[[\omega']]$ , then  $\phi$  induces an isomorphism

$$\omega' = I/I^2 \xrightarrow{\sim} I_A/I_A^2 = \omega.$$

This completes the proof.

<sup>130</sup>The editors supplied the details of the original argument that follow.

<sup>131</sup>One does not know a priori that  $\omega$  is a projective pseudocompact  $k$ -module; this will follow from the argument. Compare the proof of (iv)  $\Rightarrow$  (iii) in VIIA 7.4.

### 3.3.1. Corollary

Let  $S$  be a locally Noetherian scheme over  $\mathbb{Q}$ , and let  $G$  be an  $S$ -group scheme that is flat and locally of finite type.<sup>132</sup> If  $\omega_{G/S}$  is a locally free  $\mathcal{O}_S$ -module, then  $G$  is smooth over  $S$  at every point of the unit section.

*Proof.* Let  $x$  be a point of the unit section, let  $s$  be its image in  $S$ , and denote the local rings at  $x$  and  $s$  by  $\mathcal{O}_x$  and  $\mathcal{O}_s$ , respectively.<sup>133</sup> By hypothesis,  $\mathcal{O}_s$  and  $\mathcal{O}_x$  are Noetherian local rings. The maximal-ideal-adic topology on each of them therefore agrees with the “pseudocompact” topology defined by the ideals of finite colength. Write

$$\widehat{\mathcal{O}}_x \quad \text{and} \quad \widehat{\mathcal{O}}_s$$

for their completions in this topology. By EGA IV<sub>4</sub>, 17.5.3,  $G$  is smooth over  $S$  at  $x$  if and only if  $\widehat{\mathcal{O}}_x$  is formally smooth over  $\widehat{\mathcal{O}}_s$ , both algebras being endowed with their maximal-ideal-adic topologies. It is therefore enough to prove this last assertion.

The rings  $\widehat{\mathcal{O}}_x$  and  $\widehat{\mathcal{O}}_s$  are the local rings at  $x$  and  $s$  of the formal varieties  $\widehat{G}/\widehat{S}$  and  $\widehat{S}$  defined in 1.2.6. Put

$$k = \widehat{\mathcal{O}}_s, \quad H = \mathrm{Spf}(\widehat{\mathcal{O}}_x).$$

Then  $H$  is an infinitesimal formal  $k$ -variety. Since the formation of  $\widehat{G}/\widehat{S}$  commutes with products (loc. cit.),  $H$  is in fact an infinitesimal formal  $k$ -group.

Let  $I$  be the augmentation ideal of  $\mathcal{O}_x$ . By hypothesis,

$$\omega_{G/S,x} = I/I^2$$

is a free  $\mathcal{O}_s$ -module of finite rank  $n$ . Since  $\mathcal{O}_s$  is Noetherian, the module  $\omega_{H/k}$ , which is the completion of  $\omega_{G/S,x}$ , is identified with

$$\omega_{G/S,x} \otimes_{\mathcal{O}_s} \widehat{\mathcal{O}}_s.$$

It is therefore a topologically free  $k$ -module of rank  $n$ . By the implication (iv) $\Rightarrow$ (ii) of Theorem 3.3,

$$\widehat{\mathcal{O}}_x \simeq k[[\omega_{H/k}]] \simeq k[[t_1, \dots, t_n]].$$

Finally, the latter formal power-series algebra is formally smooth over  $k$ , by EGA 0<sub>IV</sub>, 19.3.3. This proves the corollary.

Thus we recover a result obtained independently for group schemes locally of finite presentation over an arbitrary scheme  $S$ ; cf. VI<sub>B</sub>, 1.6.

### 3.3.2 Corollary

**Corollary 3.3.2.** *Let  $k$  be a field of characteristic 0. The functor*

$$\mathbf{L} \longmapsto \mathcal{G}(\mathbf{L})$$

*is an equivalence from the category of  $k$ -Lie algebras to the category of infinitesimal formal  $k$ -groups.*<sup>134</sup>

<sup>132</sup>The editors added the flatness hypothesis, which was omitted from the original.

<sup>133</sup>The editors supplied the details of the argument that follows.

<sup>134</sup>By 2.7, 2.2.1, and 1.3.6, an infinitesimal formal  $k$ -group  $G$  is “the same thing” as a connected cocommutative  $k$ -bialgebra  $H$  (cf. 2.9). The corollary is therefore equivalent to the theorem below, obtained independently by Kostant (cf. 2.9.2). J. W. Milnor and J. C. Moore had earlier proved it in [MM65] under the additional hypothesis that  $H$  be generated as an algebra by its primitive elements. Although published

Indeed, as  $G$  ranges over the infinitesimal formal  $k$ -groups, Theorem 3.3 identifies the functor

$$G \longmapsto \mathcal{G}(\mathrm{Lie} G)$$

with the identity functor. Likewise, 3.2(ii) identifies

$$\mathbf{L} \longmapsto \mathrm{Lie}(\mathcal{G}(\mathbf{L})) = \mathrm{Prim} U(\mathbf{L})$$

with the identity functor on  $k$ -Lie algebras.

### 3.3.3

Continue to suppose that  $k$  is a field of characteristic 0. Let  $\bar{k}$  be an algebraic closure of  $k$ , and let  $\Gamma$  be the topological Galois group of  $\bar{k}$  over  $k$ .

For every formal  $k$ -group  $G$ , denote by  $\mathbf{Aut}_k(G)$  the  $k$ -group functor assigning to every finite-dimensional commutative  $k$ -algebra  $C$  the group of automorphisms of the formal  $C$ -group

$$G \widehat{\otimes}_k C.$$

<sup>135</sup> Since  $k$  is a field,  $G$  is topologically flat over  $k$ . Equivalently, its affine algebra

$$A = \mathcal{A}(G)$$

is topologically flat over  $k$ , or its covariant bialgebra

$$H = \mathbf{H}(G)$$

is flat over  $k$ . The functor  $\mathbf{Aut}_k(G)$  is therefore a formal  $k$ -group.

Indeed,  $\mathbf{Aut}_k(G)$  is the fiber product of the following diagram (cf. Exposé I, 1.7.3):

$$\begin{array}{ccc} & \underline{\mathrm{End}}_k(G) \times \underline{\mathrm{End}}_k(G) & \\ & \downarrow & \\ \mathrm{Spf}(k) & \longrightarrow & \underline{\mathrm{End}}_k(G) \times \underline{\mathrm{End}}_k(G) \end{array}$$

The vertical arrow sends

$$(\phi, \psi) \longmapsto (\phi \circ \psi, \psi \circ \phi),$$

in 1965, their text had circulated before 1960 (see the review [Ja65]); the result is consequently often called the Cartier–Kostant–Milnor–Moore theorem.

**Cartier–Kostant–Milnor–Moore Theorem.** Let  $k$  be a field of characteristic 0. The functors

$$\mathbf{L} \longmapsto U(\mathbf{L}), \quad H \longmapsto \mathrm{Prim}(H),$$

define an equivalence between the category of  $k$ -Lie algebras and the category of connected cocommutative  $k$ -bialgebras.

If  $k$  is an Artinian ring containing  $\mathbb{Q}$ , then 3.2(ii) and 3.3, combined with 2.7, 2.2.1, and 1.3.6, likewise show that

$$\mathbf{L} \longmapsto \mathcal{G}(\mathbf{L})$$

(respectively,  $\mathbf{L} \mapsto U(\mathbf{L})$ ) is an equivalence from the category of  $k$ -Lie algebras flat over  $k$  to the category of topologically flat infinitesimal formal  $k$ -groups (respectively, to the category of connected cocommutative  $k$ -bialgebras flat over  $k$ ).

<sup>135</sup>The original stated: “This  $k$ -functor is manifestly left exact ( $H$  is topologically flat over  $k$ !).” The editors supplied the details that follow.

whereas the horizontal arrow is the unit section. It is therefore enough to show that the  $k$ -functor  $\mathbf{End}_k(G)$  is represented by an object of  $\mathbf{Vaf}/k$ ; by 1.1 and 0.4.2, this amounts to showing that it is left exact, or, equivalently, that it commutes with fiber products of  $k$ -algebras.

To give an element of  $\mathbf{End}_k(G)(C)$  is equivalent to giving a  $k$ -algebra morphism

$$\phi : H \longrightarrow H \otimes_k C$$

that also respects the coalgebra structure, that is, for which the usual compatibility diagrams commute. Since  $H$  is flat over  $k$ , the functor

$$H \otimes_k -$$

commutes with fiber products of  $k$ -algebras. Thus  $\mathbf{End}_k(G)$  is left exact and can be treated as a formal  $k$ -group.

If  $L$  is a  $k$ -Lie algebra and  $G = \mathcal{G}(L)$ , Theorem 3.3 shows that  $\mathbf{Aut}_k(G)$  is isomorphic to the  $k$ -group functor  $\mathbf{Aut}_k(L)$  which assigns to each finite-dimensional  $k$ -algebra  $C$  the group of automorphisms of the  $C$ -Lie algebra

$$C \otimes_k L.$$

Now let  $G$  be an arbitrary formal  $k$ -group. We saw in 2.5.2 that  $G$  decomposes canonically as a semidirect product of an étale formal group  $G_e$  and an infinitesimal formal group  $G_0$ . This semidirect product is determined by the data of

$$L = \mathrm{Lie}(G) = \mathrm{Lie}(G_0),$$

the  $\Gamma$ -group

$$M = G_e(\bar{k}) = G(\bar{k}),$$

and a morphism of formal  $k$ -groups

$$\Phi : G_e \longrightarrow \mathbf{Aut}_k(G_0) \simeq \mathbf{Aut}_k(L).$$

Such a morphism sends  $G_e$  into the étale part  $\mathbf{Aut}_k(L)_e$  of  $\mathbf{Aut}_k(L)$  (cf. 2.5.1). It is consequently determined by a morphism of  $\Gamma$ -groups

$$\phi = \Phi(\bar{k}) : M \longrightarrow (\mathbf{Aut}_k L)(\bar{k}) = \mathrm{Aut}_{\bar{k}}(L \otimes_k \bar{k}).$$

Let  $\Gamma$  act on  $L \otimes_k \bar{k}$  by

$$\gamma(\ell \otimes t) = \ell \otimes \gamma(t).$$

Then  $\phi$  makes  $M$  act on  $L \otimes_k \bar{k}$  by automorphisms of  $\bar{k}$ -Lie algebras in such a way that

$$\phi(\gamma(m)) = \gamma \circ \phi(m) \circ \gamma^{-1}.$$

Equivalently,

$$\gamma(m)(\ell \otimes t) = \gamma(m(\ell \otimes \gamma^{-1}(t)))$$

for every  $m \in M$ ,  $\gamma \in \Gamma$ , and  $\ell \otimes t \in L \otimes_k \bar{k}$ . We express this last condition by saying that  $M$  acts on  $L \otimes_k \bar{k}$  compatibly with  $\Gamma$ .

The situation can be summarized in a “highbrow” statement. A  $\Gamma$ -Lie algebra over  $k$  is a triple

$$(L, M, \phi)$$

consisting of a  $k$ -Lie algebra  $L$ , a  $\Gamma$ -group  $M$ , and an action  $\phi$  of  $M$  on  $L \otimes_k \bar{k}$  that is compatible with the actions of  $\Gamma$  on  $M$  and on  $L \otimes_k \bar{k}$ .

If  $(L, M, \phi)$  and  $(L', M', \phi')$  are two such  $\Gamma$ -Lie algebras, a morphism from the first to the second is a pair  $(f, \theta)$ , where

$$f : L \longrightarrow L'$$

is a morphism of  $k$ -Lie algebras and

$$\theta : M \longrightarrow M'$$

is a morphism of  $\Gamma$ -groups, such that

$$(f \otimes 1)(m \cdot x) = \theta(m) \cdot (f \otimes 1)(x)$$

for every  $x \in L \otimes_k \bar{k}$  and  $m \in M$ . We have therefore proved:

**Theorem.** *Let  $k$  be a field of characteristic 0. The functor that assigns to a formal  $k$ -group  $G$  the triple*

$$(\mathrm{Lie}(G), G(\bar{k}), \Phi(\bar{k}))$$

*is an equivalence from the category of formal  $k$ -groups to the category of  $\Gamma$ -Lie algebras.*<sup>136</sup>

## 4. Phenomena specific to characteristic $p > 0$

Throughout Section 4,  $p$  denotes a prime number,  $k$  a pseudocompact ring of characteristic  $p$ , and

$$\pi : k \longrightarrow k, \quad x \longmapsto x^p,$$

the continuous Frobenius endomorphism of  $k$ .

### 4.1

Let  $X$  be a formal  $k$ -variety with affine algebra  $A$ . We write  $X^{(p/k)}$ , or simply  $X^{(p)}$ , for the formal  $k$ -variety

$$X \widehat{\otimes}_{\pi} k$$

obtained by the base change  $\pi : k \rightarrow k$  (cf. 1.2.D). Its affine algebra is the completed tensor product

$$A \widehat{\otimes}_{\pi} k.$$

The continuous morphism

$$A \widehat{\otimes}_{\pi} k \longrightarrow A, \quad a \widehat{\otimes}_{\pi} \lambda \longmapsto a^p \lambda,$$

therefore induces a morphism of formal  $k$ -varieties

$$\mathrm{Fr}(X/k) : X \longrightarrow X^{(p/k)}.$$

We call  $\mathrm{Fr}(X/k)$  the *Frobenius morphism of  $X$  relative to  $k$* , and below we shall often write  $\mathrm{Fr}$  in place of  $\mathrm{Fr}(X/k)$ .

---

<sup>136</sup>In particular, when  $k = \bar{k}$ , this recovers the Cartier–Gabriel–Kostant theorem mentioned in 2.9.2.

## 4.1.1

137

Let  $S$  be a scheme over the prime field  $\mathbb{F}_p$ , and let  $\eta : X \rightarrow S$  be an  $S$ -scheme. Write  $\text{fr}(S)$  for the “absolute” Frobenius morphism of  $S$ : it induces the identity on the underlying topological space and sends every section  $f$  of the structure sheaf to  $f^p$ ; cf. VII<sub>A</sub>, 4.1. Let

$$X^{(p)} = X \times_{\text{fr}(S)} S$$

be the corresponding  $S$ -scheme (VII<sub>A</sub>, loc. cit.); thus its structural morphism to  $S$  is  $\text{fr}(S) \circ \eta$ .

Let  $\widehat{S}$  be the formal scheme whose underlying topological space is the set of closed points of  $S$ , endowed with the discrete topology, and whose local ring at a closed point  $s$  is the separated completion

$$\mathcal{O}_{\widehat{S},s} = \widehat{\mathcal{O}}_{S,s}$$

of  $\mathcal{O}_{S,s}$  for the linear topology defined by ideals of finite colength (cf. 1.2.6). Its affine algebra is therefore

$$k = \mathcal{A}(\widehat{S}) = \prod_{s \in S_{\text{cl}}} \widehat{\mathcal{O}}_{S,s}.$$

Recall (loc. cit.) that  $\widehat{X}/\widehat{S}$  denotes the formal  $k$ -variety consisting of those points  $x \in X$  for which

$$[\kappa(x) : \kappa(s)] < \infty,$$

where  $s$  is the image of  $x$  in  $S$ . Its local ring at  $x$  is the separated completion

$$\mathcal{O}_{\widehat{X}/\widehat{S},x} = \widehat{\mathcal{O}}_{X,x}$$

of  $\mathcal{O}_{X,x}$  for the linear topology defined by those ideals  $I$  such that  $\mathcal{O}_{X,x}/I$  has finite length as an  $\mathcal{O}_{X,x}$ -module, and hence also as an  $\mathcal{O}_{S,s}$ -module. By base change we can therefore form the formal  $k$ -variety

$$(\widehat{X}/\widehat{S})^{(p)} = (\widehat{X}/\widehat{S}) \widehat{\otimes}_{\pi} k.$$

We can also consider the formal  $k$ -variety  $\widehat{X^{(p)}}/\widehat{S}$ . Its underlying set consists of the points  $x \in X^{(p)} = X$  such that, if  $s$  denotes the image of  $x$  in  $S$ , the composite

$$\kappa(s) \xrightarrow{\text{fr}} \kappa(s) \xrightarrow{\eta^\sharp} \kappa(x)$$

makes  $\kappa(x)$  a finite extension of  $\kappa(s)$ . In that case the same is true of

$$\eta^\sharp : \kappa(s) \longrightarrow \kappa(x),$$

so  $x$  is a point of  $\widehat{X}/\widehat{S}$ , and

$$\mathcal{O}_{\widehat{X^{(p)}}/\widehat{S},x} = \widehat{\mathcal{O}}_{X,x} \widehat{\otimes}_{\pi} \widehat{\mathcal{O}}_{S,s} = \widehat{\mathcal{O}}_{X^{(p)},x}.$$

The second equality follows because the construction  $X \mapsto \widehat{X}/\widehat{S}$  commutes with products (cf. 1.2.6).

It follows that

$$\widehat{X^{(p)}}/\widehat{S} \subset (\widehat{X}/\widehat{S})^{(p)}$$

as a formal subvariety. Equality holds if and only if, for every closed point  $s$  of  $S$ , the field  $\kappa(s)$  has finite degree over  $\kappa(s)^p$ . This is the case, for example, if  $S$  is locally of finite type over a field  $\kappa$  such that

$$[\kappa : \kappa^p] < \infty.$$

<sup>137</sup>The editors corrected the original, which gave the inclusion

$$(\widehat{X}/\widehat{S})^{(p)} \subset \widehat{X^{(p)}}/\widehat{S}$$

instead of the reverse inclusion. They also note that this paragraph is not used below.



**4.1.2**

Let  $G$  be a formal  $k$ -group. Since the functor

$$X \longmapsto X^{(p)} = X \hat{\otimes}_{\pi} k$$

commutes with finite products, it carries formal  $k$ -groups to formal  $k$ -groups. Moreover,  $\mathrm{Fr}(X/k)$  is functorial in  $X$ , so

$$\mathrm{Fr} : G \longrightarrow G^{(p)}$$

is a homomorphism of formal  $k$ -groups.

Define recursively

$$G^{(p^{n+1})} = (G^{(p^n)})^{(p)}.$$

Then the same is true of the composite

$$\mathrm{Fr}^n = \mathrm{Fr}^n(G/k) : G \xrightarrow{\mathrm{Fr}} G^{(p)} \xrightarrow{\mathrm{Fr}} G^{(p^2)} \xrightarrow{\mathrm{Fr}} \cdots \xrightarrow{\mathrm{Fr}} G^{(p^n)}.$$

Let  $A$  be the affine algebra of  $G$ , and let  $I_A$  be its augmentation ideal.

**Definitions.**

- (a) The kernel of  $\mathrm{Fr}^n$  is denoted

$$\mathrm{Fr}_n G.$$

It follows directly from the definitions that  $\mathrm{Fr}_n G$  is infinitesimal and has affine algebra

$$A/I_A^{\{p^n\}},$$

where  $I_A^{\{p^n\}}$  is the closed ideal of  $A$  generated by the  $p^n$ -th powers of the elements of  $I_A$ .

- (b) The group  $G$  is said to have *height at most  $n$*  if

$$I_A^{\{p^n\}} = 0;$$

equivalently,

$$\mathrm{Fr}_n G = G.$$

**4.2**

It is clear that the Lie algebra  $\mathrm{Lie}(G)$  of a formal  $k$ -group  $G$  is a  $p$ -Lie subalgebra of  $\mathbf{H}(G)$  (cf. 2.3). Indeed, we may immediately reduce to the case in which  $k$  is Artinian. Then  $\mathrm{Lie}(G)$  is the subset of  $\mathbf{H}(G)$  consisting of the elements  $x$  such that

$$\varphi_G(x \otimes 1 + 1 \otimes x) = \Delta_G(x),$$

with the notation of 2.3 and 2.6(c). Hence

$$\begin{aligned} \varphi_G(x^p \otimes 1 + 1 \otimes x^p) &= \varphi_G((x \otimes 1 + 1 \otimes x)^p) \\ &= \varphi_G(x \otimes 1 + 1 \otimes x)^p \\ &= \Delta_G(x)^p = \Delta_G(x^p). \end{aligned}$$

## 4.2.1

Conversely, every  $p$ -Lie algebra  $\mathbf{L}$  over  $\mathbf{O}_k$  defines a  $k$ -group functor. Let  $U_p(\mathbf{L})$  denote the functor assigning to every object  $C$  of  $\mathbf{Alf}/k$  the restricted enveloping algebra

$$U_p(\mathbf{L}(C))$$

of the  $p$ -Lie algebra  $\mathbf{L}(C)$  over  $C$  (VII<sub>A</sub>, 5.3). By VII<sub>A</sub>, 5.4,  $U_p(\mathbf{L})$  is a bialgebra over  $\mathbf{O}_k$ , and by 2.2 it therefore determines the  $k$ -group functor

$$\mathrm{Spf}^* U_p(\mathbf{L}),$$

which we henceforth denote by  $\mathcal{G}_p(\mathbf{L})$ .

Thus  $\mathcal{G}_p(\mathbf{L})(C)$  is the group of elements

$$z \in U_p(\mathbf{L}(C))$$

having augmentation 1 and satisfying

$$\Delta_{U_p(\mathbf{L}(C))}(z) = z \otimes z.$$

## 4.2.2

Suppose that  $\mathbf{L}$  is a flat  $p$ -Lie algebra over  $\mathbf{O}_k$ . Taking VII<sub>A</sub>, 5.3.3 into account, the argument of part (i) of Proposition 2.6.2 shows that  $U_p(\mathbf{L})$  is flat over  $\mathbf{O}_k$ . Consequently,  $\mathcal{G}_p(\mathbf{L})$  is a formal  $k$ -group topologically flat over  $k$ , with  $U_p(\mathbf{L})$  as its covariant  $\mathbf{O}_k$ -bialgebra.

<sup>138</sup> By the proof of 2.6.2(iii), for every finite-length  $k$ -algebra  $C$ , the Lie algebra  $\mathrm{Lie}(\mathcal{G}_p(\mathbf{L}))(C)$  is the set of primitive elements of

$$U_p(\mathbf{L})(C) = U_p(\mathbf{L}(C));$$

see also VII<sub>A</sub>, 3.2.3. Moreover, by VII<sub>A</sub>, 5.5.3, the canonical morphism  $\mathbf{L} \rightarrow U_p(\mathbf{L})$  induces an isomorphism of  $p$ -Lie algebras

$$\tau_{p,\mathbf{L}} : \mathbf{L} \xrightarrow{\sim} \mathrm{Lie}(\mathcal{G}_p(\mathbf{L})).$$

Compare this with 3.1 in characteristic 0.

The formal group  $\mathcal{G}_p(\mathbf{L})$  can be characterized by a universal property. Every morphism

$$h : \mathcal{G}_p(\mathbf{L}) \longrightarrow G$$

to a formal group  $G$  induces a morphism

$$\mathrm{Lie}(h) : \mathrm{Lie}(\mathcal{G}_p(\mathbf{L})) \longrightarrow \mathrm{Lie}(G).$$

Composing this morphism with  $\tau_{p,\mathbf{L}}$  gives

$$h' : \mathbf{L} \longrightarrow \mathrm{Lie}(G).$$

**Proposition.** *Let  $k$  be a pseudocompact ring of characteristic  $p > 0$ , let  $G$  be a formal  $k$ -group, and let  $\mathbf{L}$  be a flat  $p$ -Lie algebra over  $\mathbf{O}_k$ . The map  $h \mapsto h'$  defined above is a bijection*

$$\mathrm{Hom}_{\mathrm{Grf}/k}(\mathcal{G}_p(\mathbf{L}), G) \xrightarrow{\sim} \mathrm{Hom}_{p\text{-Lie}}(\mathbf{L}, \mathrm{Lie}(G)),$$

<sup>138</sup>The editors supplied the details that follow.

where the right-hand side is the set of morphisms of  $p$ -Lie algebras from  $\mathbf{L}$  to  $\mathrm{Lie}(G)$ .

*Proof.*<sup>139</sup> We may immediately reduce to the case in which  $k$  is Artinian. Put

$$L = \mathbf{L}(k).$$

By 2.3.1,

$$\mathrm{Hom}_{\mathrm{Grf}/k}(\mathcal{G}_p(\mathbf{L}), G)$$

is in bijection with the set of unital algebra morphisms

$$h : U_p(L) \longrightarrow \mathbf{H}(G)$$

for which the following diagram commutes:

$$\begin{array}{ccccc} U_p(L) & \xrightarrow{h} & \mathbf{H}(G) & \xrightarrow{\delta_G} & \mathbf{H}(G \times G) \\ \Delta_{U_p(L)} \downarrow & & & & \uparrow \varphi_G \\ U_p(L) \otimes U_p(L) & \xrightarrow{h \otimes h} & \mathbf{H}(G) \otimes \mathbf{H}(G) & & \end{array} .$$

The morphism  $h$  is determined by its restriction to  $L$ , which is a morphism of  $p$ -Lie algebras from  $L$  to the underlying  $p$ -Lie algebra of  $\mathbf{H}(G)$ . The commutativity of the diagram says precisely that  $h$  sends  $L$  into the subset of  $\mathbf{H}(G)$  consisting of those  $x$  such that

$$\delta_G(x) = \varphi_G(x \otimes 1 + 1 \otimes x).$$

This subset is  $\mathrm{Lie}(G)$ , by 2.6(c). The proposition follows.

### 4.3

We now study the formal  $k$ -group  $\mathcal{G}_p(\mathbf{L})$  in greater detail when  $\mathbf{L}$  is a flat  $p$ -Lie algebra over  $\mathbf{O}_k$ .

For this purpose, first let  $C$  be a ring of characteristic  $p$ , and let  $L$  be a  $p$ -Lie algebra over  $C$  whose underlying module is free with basis

$$(x_i)_{i \in I}.$$

Let  $\mathcal{P}$  be the set of families

$$n = (n_i)_{i \in I}$$

of nonnegative integers such that

$$0 \leq n_i < p$$

and  $n_i = 0$  except possibly for finitely many indices. Endow  $I$  with a total order, and let

$$i_1 < i_2 < \cdots < i_r$$

be the indices for which  $n_i \neq 0$ . Put

$$|n| = n_{i_1} + \cdots + n_{i_r},$$

and

$$x^n = x_{i_1}^{n_{i_1}} x_{i_2}^{n_{i_2}} \cdots x_{i_r}^{n_{i_r}}, \quad n! = (n_{i_1}!) \cdots (n_{i_r}!).$$

<sup>139</sup>The proof is identical to that of 2.6.3.

By VII<sub>A</sub>, 5.3.3, the monomials  $x^n/n!$  form a basis of  $U_p(L)$ , and

$$\Delta\left(\frac{x^n}{n!}\right) = \sum_{m \leq n} \frac{x^m}{m!} \otimes \frac{x^{n-m}}{(n-m)!}. \quad (*)$$

The sum ranges over all  $m \in \mathcal{P}$  such that  $m \leq n$ , that is,

$$m_i \leq n_i \quad \text{for every } i \in I.$$

Formula (\*) gives an immediate description of the natural filtration of  $U_p(L)$  (cf. 1.3.6). Put

$$U = U_p(L),$$

let  $U^+$  be the two-sided ideal generated by  $L$ , and put

$$U_0 = C \cdot 1_U.$$

As in 1.3.6, define a filtration of  $U$  by  $C$ -subcoalgebras by setting, for  $n \geq 1$ ,

$$U_n = \{u \in U \mid \Delta_U(u) - u \otimes 1 \in U_{n-1} \otimes U^+\}.$$

Formula (\*) then shows that  $U_n$  is the free  $C$ -module with basis the monomials  $x^m$  satisfying

$$|m| \leq n.$$

#### 4.3.1

With the notation of 4.3, let us determine the elements  $\xi \in U = U_p(L)$  such that

$$\varepsilon_U(\xi) = 1, \quad \Delta_U(\xi) = \xi \otimes \xi.$$

Every element  $\xi \in U$  has a unique expression

$$\xi = \sum_{n \in \mathcal{P}} \xi_n \frac{x^n}{n!}, \quad \xi_n \in C.$$

The condition  $\varepsilon_U(\xi) = 1$  gives

$$\xi_0 = 1,$$

where 0 denotes the element of  $\mathcal{P}$  all of whose components vanish. The condition  $\Delta_{U_p(L)}(\xi) = \xi \otimes \xi$  gives

$$\sum_{\substack{n \in \mathcal{P} \\ m \leq n}} \xi_n \frac{x^m}{m!} \otimes \frac{x^{n-m}}{(n-m)!} = \sum_{q, r \in \mathcal{P}} \xi_q \xi_r \frac{x^q}{q!} \otimes \frac{x^r}{r!}.$$

Equivalently,

$$\xi_{q+r} = \xi_q \xi_r \quad \text{if } q_i + r_i < p \text{ for every } i,$$

and

$$\xi_q \xi_r = 0 \quad \text{otherwise.}$$

Let  $\xi_i$  denote the coefficient  $\xi_n$  for which  $n_i = 1$  and  $n_j = 0$  whenever  $j \neq i$ . The preceding conditions say precisely that

$$\xi_n = \prod_i \xi_i^{n_i} \quad (n \in \mathcal{P}), \quad \xi_i^p = 0 \quad (i \in I).$$

We obtain:

**Proposition.** *Let  $k$  be a local pseudocompact ring<sup>140</sup> of characteristic  $p > 0$ , let  $\mathbf{L}$  be a flat  $p$ -Lie algebra over  $\mathbf{O}_k$ , and let  $C$  be an object of  $\mathbf{Alf}/k$ . Write*

$$\sqrt[p]{0_C} = \{x \in C \mid x^p = 0\}.$$

*There is a bijection, “functorial in  $C$ ,”*

$$\mathbf{L}(C) \otimes_C \sqrt[p]{0_C} \xrightarrow{\sim} \mathcal{G}_p(\mathbf{L})(C).$$

*Proof.* By Remark 1.2.3.F, one may choose a basis

$$({}^C x_i)_{i \in I}$$

of  $\mathbf{L}(C)$  over  $C$  in such a way that, for every morphism  $\phi : C \rightarrow D$  in  $\mathbf{Alf}/k$ , the map  $\mathbf{L}(\phi)$  sends  ${}^C x_i$  to  ${}^D x_i$ . The preceding calculation shows that

$$\sum_i ({}^C x_i) \otimes \xi_i \mapsto \sum_{n \in \mathcal{P}} \left( \prod_i \xi_i^{n_i} \right) \frac{({}^C x)^n}{n!}$$

is a functorial bijection from  $\mathbf{L}(C) \otimes_C \sqrt[p]{0_C}$  onto  $\mathcal{G}_p(\mathbf{L})(C)$ .

#### 4.3.2

“There is no Campbell–Hausdorff formula in characteristic  $p > 0$ .” Let us explain. The functorial isomorphism of 4.3.1 depends on the choice of the bases  $({}^C x_i)_{i \in I}$ . Unlike the situation in 3.2, in characteristic 0, there is in general no bijection

$$\mathbf{L}(C) \otimes_C \sqrt[p]{0_C} \longrightarrow \mathcal{G}_p(\mathbf{L})(C)$$

that is “functorial both in  $C$  and in  $\mathbf{L}$ .” The following argument shows, for example, that no such isomorphism exists when  $k$  is a field containing all  $(p^2 - 1)$ -st roots of unity.

Choose  $\mathbf{L}$  so that, for every  $C \in \mathbf{Alf}/k$ , the algebra  $\mathbf{L}(C)$  is the abelian  $p$ -Lie algebra over  $C$  generated by an element  $x$  subject to

$$x^{(p^2)} = 0.$$

Thus

$$\mathbf{L}(C) = Cx \oplus Cx^{(p)}, \quad U_p(\mathbf{L}(C)) \simeq C[x]/(x^{p^2}).$$

*Source note.* The French prints  $k[x]/(x^{p^2})$  in the last display, although the enveloping algebra here is formed over  $C$ . A little later it also identifies  $\mathcal{G}_p(\mathbf{L})(C)$ , whose elements are group-like, with  $\text{Prim } U_p(\mathbf{L}(C))$ . The target uses the coefficient ring  $C$  and retains the group-like set defined in 4.2.1; primitive elements instead describe its Lie algebra in 4.2.2.

We shall show that the only functorial morphism

$$\chi_C : \mathbf{L}(C) \otimes_C \sqrt[p]{0_C} \longrightarrow U_p(\mathbf{L}(C))$$

which is compatible with the endomorphisms of  $\mathbf{L}$  and sends  $\mathbf{L}(C) \otimes_C \sqrt[p]{0_C}$  into  $\mathcal{G}_p(\mathbf{L})(C)$  is the constant morphism with value 1.

<sup>140</sup>The editors added the hypothesis that  $k$  be local, so that every topologically flat pseudocompact  $k$ -module is topologically free; see the proof.

Let

$$\psi_C : \mathbf{L}(C) \otimes_C \sqrt[p]{0_C} \xrightarrow{\sim} \mathcal{G}_p(\mathbf{L})(C)$$

be the bijection of 4.3.1, namely

$$x \otimes a + x^{(p)} \otimes b \mapsto \sum_{0 \leq i, j < p} a^i b^j x^{i+pj}.$$

Composing  $\chi_C$  with  $\psi_C^{-1}$  gives a morphism, functorial in  $C$ ,

$$\begin{aligned} \theta_C : \mathbf{L}(C) \otimes_C \sqrt[p]{0_C} &\longrightarrow \mathbf{L}(C) \otimes_C \sqrt[p]{0_C}, \\ x \otimes a + x^{(p)} \otimes b &\longmapsto x \otimes P(a, b) + x^{(p)} \otimes Q(a, b). \end{aligned}$$

Functoriality in  $C$  implies that  $P(a, b)$  and  $Q(a, b)$  are the values at  $(a, b)$  of two polynomials

$$P, Q \in k[X, Y]$$

which may be assumed to have degree less than  $p$  in each of  $X$  and  $Y$ .<sup>141</sup>

Let  $\alpha \in k$ , and let  $\ell_\alpha$  be the endomorphism of the  $p$ -Lie algebra  $\mathbf{L}$  sending

$$x \mapsto \alpha x, \quad x^{(p)} \mapsto \alpha^p x^{(p)}.$$

Then  $U_p(\ell_\alpha)$  sends  $x$  to  $\alpha x$ , and hence sends  $x^i$  to  $\alpha^i x^i$  for  $i < p^2$ . The following square commutes:

$$\begin{array}{ccc} \mathbf{L}(C) \otimes_C \sqrt[p]{0_C} & \xrightarrow{\psi_C} & U_p(\mathbf{L}(C)) \\ \downarrow \ell_\alpha(C) \otimes \text{id} & & \downarrow U_p(\ell_\alpha)(C) \\ \mathbf{L}(C) \otimes_C \sqrt[p]{0_C} & \xrightarrow{\psi_C} & U_p(\mathbf{L}(C)) \end{array}$$

The hypothesis

$$\chi_C \circ (\ell_\alpha(C) \otimes \text{id}) = U_p(\ell_\alpha)(C) \circ \chi_C$$

therefore gives

$$P(\alpha a, \alpha^p b) = \alpha P(a, b), \quad Q(\alpha a, \alpha^p b) = \alpha^p Q(a, b).$$

Write

$$P = \sum_{i, j < p} \lambda_{ij} X^i Y^j, \quad Q = \sum_{i, j < p} \mu_{ij} X^i Y^j,$$

and take

$$C = k[X, Y]/(X^p, Y^p).$$

If  $\lambda_{ij} \neq 0$ , then

$$\alpha^{i-1+pj} = 1;$$

if  $\mu_{ij} \neq 0$ , then

$$\alpha^{i+p(j-1)} = 1.$$

Taking  $\alpha$  to be a primitive root of unity of order  $p^2 - 1$ , we conclude that

$$P = \lambda X, \quad Q = \mu Y$$

<sup>141</sup>The editors expanded the original in the argument that follows.

for some  $\lambda, \mu \in k$ . Consequently,

$$\chi_C(x \otimes a + x^{(p)} \otimes b) = \sum_{0 \leq i, j < p} (\lambda a)^i (\mu b)^j x^{i+pj}.$$

Finally, consider the endomorphism  $f$  of  $\mathbf{L}$  sending

$$x \longmapsto x + x^{(p)}.$$

Take  $C = k[a]/(a^3)$  and compare the coefficients of  $x^p$  and  $x^{p+1}$  in

$$(\chi_C \circ f(C))(x \otimes a) \quad \text{and} \quad (U_p(f)(C) \circ \chi_C)(x \otimes a).$$

This gives

$$\lambda = \mu, \quad \lambda^2 a^2 = 0,$$

and hence  $\lambda = \mu = 0$ , as claimed.

#### 4.4 Theorem

**Theorem 4.4.** *Let  $k$  be a pseudocompact ring of characteristic  $p > 0$ , and let  $G$  be a formal  $k$ -group. The following assertions are equivalent:*

- (i) *There exists a  $p$ -Lie algebra  $\mathbf{L}$ , flat over  $\mathcal{O}_k$ , such that  $G$  is isomorphic to  $\mathcal{G}_p(\mathbf{L})$ . In this case,  $\mathbf{L} = \text{Lie}(G)$  by 4.2.2.*
- (ii) *There exists a projective pseudocompact  $k$ -module  $\omega$  such that the affine algebra of  $G$  is isomorphic to the quotient of  $k[[\omega]]$  by the closed ideal generated by the elements  $x^p$ ,  $x \in \omega$ . In this case,  $\omega \simeq \omega_{G/k}$ .*
- (iii) *The group  $G$  has height at most 1, and  $\omega_{G/k}$  is a projective pseudocompact  $k$ -module.*
- (iv) *The group  $G$  has height at most 1, and is topologically flat over  $k$ .*

*Proof of (i)  $\Rightarrow$  (ii).* Set

$$\omega = \Gamma^*(\mathbf{L})$$

(cf. 1.2.3.D), and let  $A$  be the quotient of  $k[[\omega]]$  by the closed ideal generated by the elements  $x^p$ ,  $x \in \omega$ . Every continuous morphism

$$h : A \longrightarrow C, \quad C \in \text{Alf}/k,$$

is determined by its restriction  $h'$  to  $\omega$ , and this restriction maps  $\omega$  into  $\sqrt[p]{0_C}$ . Thus there is a canonical bijection from

$$\text{Hom}_{\text{Alf}/k}(A, C)$$

onto the set

$$\text{Hom}_c(\omega, \sqrt[p]{0_C})$$

of continuous  $k$ -linear maps from  $\omega$  into  $\sqrt[p]{0_C}$ . The latter set is canonically isomorphic to

$$\mathbf{L}(C) \otimes_C \sqrt[p]{0_C}$$

(see the proof of 3.3). The implication (i)  $\Rightarrow$  (ii) therefore follows from the functorial bijection

$$\mathbf{L}(C) \otimes_C \sqrt[p]{0_C} \xrightarrow{\sim} \mathcal{G}_p(\mathbf{L})(C)$$

established in 4.3.1.

For the remaining implications, consult the analogous proofs of Theorems 3.3 and VIIA 7.4.1.

**Remark 4.4.A.** Let  $G$  be an infinitesimal formal  $k$ -group, with affine algebra  $A$ , such that

$$\omega_{G/k} = I_A/I_A^2$$

is a projective pseudocompact  $k$ -module. Then the projection

$$I_A \longrightarrow \omega_{G/k}$$

has a section

$$\tau : \omega_{G/k} \longrightarrow I_A,$$

and  $\tau$  induces a surjective continuous morphism of algebras

$$\psi : k[[\omega_{G/k}]] \longrightarrow A;$$

compare the proof of the implication (iii)  $\Rightarrow$  (i) in 3.3.<sup>142</sup>

#### 4.4.1 Corollary

**Corollary 4.4.1.** *If  $k$  is a field of characteristic  $p > 0$ , the functor*

$$\mathbf{L} \longmapsto \mathcal{G}_p(\mathbf{L})$$

*is an equivalence from the category of  $p$ -Lie algebras over  $k$  to the category of formal  $k$ -groups of height at most 1.*

Indeed, as  $G$  ranges over the formal groups of height at most 1, Theorem 4.4 identifies the functor

$$G \longmapsto \mathcal{G}_p(\mathrm{Lie} G)$$

with the identity functor. Likewise, 4.2.2 identifies the functor

$$\mathbf{L} \longmapsto \mathrm{Lie} \mathcal{G}_p(\mathbf{L})$$

with the identity functor (see also VIIA 5.5.3).<sup>143</sup>

#### 4.4.2

Continue to assume that  $k$  is a field of characteristic  $p$ . Let  $G$  be an infinitesimal formal  $k$ -group, with affine algebra  $A$ . Since  $G$  is infinitesimal, every open ideal of  $A$  contains

$$I_A^{\{p^n\}}$$

for all sufficiently large  $n$ . Thus  $A$  is the projective limit of the affine algebras

$$A/I_A^{\{p^n\}}$$

of the groups  $\mathrm{Fr}_n G$  (cf. 4.1.2). Equivalently, every infinitesimal formal  $k$ -group is an inductive limit of formal  $k$ -groups of finite height.

<sup>142</sup>The editors added this remark, which is used in 4.4.2.

<sup>143</sup>If  $k$  is an Artinian ring of characteristic  $p > 0$ , the same proof gives an equivalence between the category of  $p$ -Lie algebras flat over  $k$  and the category of formal  $k$ -groups of height at most 1 that are topologically flat over  $k$ .



Suppose that  $G$  has height at most  $n$ , and write

$$H = G / \mathrm{Fr} G.$$

<sup>144</sup> By 2.4 and 2.4.1, the morphism

$$\mathrm{Fr} : G \longrightarrow G^{(p)}$$

factors as an epimorphism

$$\pi : G \longrightarrow H$$

followed by a monomorphism

$$i : H \longrightarrow G^{(p)}.$$

We therefore have the following commutative diagram:

$$\begin{array}{ccccc} G & \xrightarrow{\pi} & H & \xrightarrow{\mathrm{Fr}^{n-1}} & H^{(p^{n-1})} \\ & \searrow \mathrm{Fr} & \downarrow i & & \downarrow i^{(p^{n-1})} \\ & & G^{(p)} & \xrightarrow{\mathrm{Fr}^{n-1}} & G^{(p^n)} \end{array}$$

Since the functor  $X \mapsto X^{(p)}$  commutes with fiber products, the morphism

$$i^{(p^{n-1})}$$

is again a monomorphism. Moreover,  $\mathrm{Fr}^n(G/k)$  is zero and  $\pi$  is an epimorphism. Hence

$$\mathrm{Fr}^{n-1}(G^{(p)}/k) \circ i = i^{(p^{n-1})} \circ \mathrm{Fr}^{n-1}(H/k)$$

is zero. Because  $i^{(p^{n-1})}$  is a monomorphism,

$$\mathrm{Fr}^{n-1}(H/k) = 0,$$

so  $H$  has height at most  $n - 1$ .

It follows that every formal  $k$ -group of finite height has a composition series whose quotients have height at most 1, and therefore can be described by  $p$ -Lie algebras over  $k$ .

Finally, the affine algebra  $A$  of  $G$  is a quotient of

$$k[[\omega_{G/k}]]$$

by Remark 4.4.A. If  $\omega_{G/k}$  is finite-dimensional over  $k$ , all the algebras

$$A/I_A^{\{p^n\}}$$

are finite-dimensional  $k$ -vector spaces. Consequently, every infinitesimal formal group  $G$  over a field of characteristic  $p > 0$  for which  $\omega_{G/k}$  is finite-dimensional over  $k$  is an inductive limit of finite formal groups (that is, groups of finite length; cf. 1.2.6).

<sup>144</sup>The editors supplied the details of the original argument in what follows.

## 5. Homogeneous spaces of infinitesimal formal groups over a field

### 5.0

For simplicity, suppose that  $k$  is a field.<sup>145</sup> Let  $G$  be a formal  $k$ -group with affine algebra

$$A = \mathcal{A}(G).$$

Let  $A^+$  be the augmentation ideal of  $A$ . For  $a \in A$ , write

$$\bar{a} = a - \varepsilon_A(a)1_A$$

for its projection onto  $A^+$ . Let  $F$  be a formal subgroup of  $G$ , defined by a closed ideal  $J$  of  $A$ , and let

$$\pi : A \longrightarrow A/J$$

be the projection. Write

$$\bar{\pi} : A \longrightarrow A^+/J$$

for the composite of the projections

$$A \longrightarrow A^+ \longrightarrow A^+/J.$$

For every  $a \in A$ , the projection of  $\Delta_A(a)$  onto  $A \hat{\otimes} A^+$  is

$$\Delta_A(a) - a \hat{\otimes} 1.$$

By Theorem 2.4, the quotient formal  $k$ -variety  $X = G/F$  exists. Its affine algebra is

$$\begin{aligned} B &= \text{Eq} \left( A \begin{array}{c} \xrightarrow{\tau_1} \\ (\text{id}_A \hat{\otimes} \pi) \Delta_A \end{array} A \hat{\otimes}_k (A/J) \right) \\ &= \{a \in A \mid (\text{id}_A \hat{\otimes} \pi) \Delta_A(a) = a \hat{\otimes} \pi(1)\} \\ &= \ker((\text{id}_A \hat{\otimes} \bar{\pi}) \Delta_A), \end{aligned}$$

where

$$\tau_1(a) = a \hat{\otimes} \pi(1).$$

Equivalently,  $B$  is the subalgebra of  $A$  consisting of the elements  $\varphi$  such that

$$\varphi(gh) = \varphi(g)$$

for every  $C \in \text{Alf}/k$ ,  $g \in G(C)$ , and  $h \in F(C)$ . For  $g, g' \in G(C)$  and  $h \in F(C)$ , one has

$$\varphi(gg'h) = \varphi(gg').$$

Thus  $\Delta_A(\varphi)$  belongs to the kernel of

$$\text{id}_A \hat{\otimes}_k ((\text{id}_A \hat{\otimes} \bar{\pi}) \Delta_A),$$

which equals  $A \hat{\otimes}_k B$ , since  $A$  is topologically flat over  $k$ . Consequently,

$$\Delta_A(B) \subset A \hat{\otimes}_k B;$$

<sup>145</sup>The editors added subsection 5.0 in order to express Proposition 5.1 below in the language of cocommutative Hopf algebras and to cite results subsequently obtained in this direction.

that is, the closed subalgebra  $B$  is also a left coideal.

Conversely,  $B$  determines  $F$ , because Corollary 2.4.1 gives

$$J = AB^+, \quad B^+ = B \cap A^+;$$

in other words,  $J$  is the closed ideal generated by  $B^+$ . We therefore obtain an injective map

$$\mathcal{Q} : \mathcal{F} \longrightarrow \mathcal{B}$$

from the set  $\mathcal{F}$  of formal subgroups of  $G$  to the set  $\mathcal{B}$  of closed subalgebras of  $A$  that are left coideals. Proposition 5.1 below shows that  $\mathcal{Q}$  is bijective when  $G$  is infinitesimal. In fact, it is bijective for every formal  $k$ -group  $G$ .

Recall from 2.2.1 that the functor

$$G \longmapsto \mathbf{H}(G)$$

is an equivalence from formal  $k$ -groups to cocommutative  $k$ -Hopf algebras. If  $F$  is defined by the closed ideal  $J \subset A$ , then the Hopf subalgebra

$$\mathbf{H}(F) \subset H = \mathbf{H}(G)$$

is the orthogonal complement of  $J$  for the duality

$$A = H^*, \quad H = A^\dagger$$

(cf. 0.2.2). Conversely, if  $B$  is a closed subalgebra of  $A$  that is also a left coideal, then

$$I = B^\perp$$

is a coideal and a left ideal of  $H$ ; explicitly,

$$\Delta_H(I) \subset I \otimes H + H \otimes I, \quad \varepsilon_H(I) = 0.$$

Let  $\mathcal{H}$  be the set of Hopf subalgebras of  $H$ , and let  $\mathcal{I}$  be the set of left ideals of  $H$  that are also coideals. For  $I \in \mathcal{I}$ , denote by

$$\pi_I : H \longrightarrow H/I$$

the projection, and by

$$\bar{\pi}_I : H \longrightarrow H^+/I$$

the composite

$$H \longrightarrow H^+ \longrightarrow H^+/I,$$

where  $H^+$  is the augmentation ideal of  $H$ .

Let  $K$  be a Hopf subalgebra of  $H$ , and put

$$K^+ = K \cap H^+.$$

If  $F$  is the formal subgroup corresponding to  $K$ , then

$$J = K^\perp$$

and  $A^+/J$  identifies with the dual of  $K^+$ . Since  $B = \mathcal{Q}(F)$  is the kernel of

$$A \xrightarrow{\Delta_A} A \hat{\otimes}_k A \xrightarrow{\text{id} \hat{\otimes} \bar{\pi}} A \hat{\otimes}_k (A^+/J),$$

duality identifies  $\mathcal{Q}$  with the map

$$\Phi : \mathcal{H} \longrightarrow \mathcal{I}, \quad K \longmapsto HK^+.$$

Indeed,  $HK^+$  is the image of

$$H \otimes_k K^+ \xrightarrow{\text{id} \otimes \text{can}} H \otimes_k H \xrightarrow{m_H} H,$$

and is both a left ideal and a coideal.

Similarly, the map that assigns  $AB^+$  to  $B \in \mathcal{B}$  corresponds by duality to

$$\Psi : \mathcal{I} \longrightarrow \mathcal{H},$$

where

$$\Psi(I) = \{x \in H \mid (\text{id}_H \otimes \pi_I)\Delta_H(x) = x \otimes \pi_I(1)\}.$$

**Theorem 5.0.1.** *Let  $k$  be a field and  $H$  a cocommutative Hopf algebra. The maps  $\Phi$  and  $\Psi$  above are mutually inverse bijections between the Hopf subalgebras of  $H$  and the left ideals of  $H$  that are coideals.*

This theorem was first proved by K. Newman ([Ne75], Theorem 4.1, where the word “cocommutative” was omitted). Newman’s proof uses the Cartier–Gabriel–Kostant theorem (cf. 2.9) to reduce to the case in which  $H$  is connected, and then uses the existence of a “Sweedler basis” in that case ([Sw67], Theorem 3), a result dual to the Dieudonné–Cartier Theorem 5.2.2 below. A second, shorter proof was given by H. J. Schneider ([Sch90], Theorem 4.15). A. Masuoka subsequently generalized the result to the case in which only the coradical  $H_0$  of  $H$ , the sum of its simple subcoalgebras, is assumed commutative ([Ma91], Theorem 1.3 (3)).

Finally, for a commutative  $k$ -Hopf algebra corresponding to an affine  $k$ -group scheme  $G$ , one cannot expect an analogue of Theorem 5.0.1 without additional hypotheses: for a  $k$ -subgroup  $F \subset G$ , the quotient  $G/F$  need not be affine. Nevertheless, M. Takeuchi established analogous bijections. In [Tak72], Theorem 4.3, invariant  $k$ -subgroups  $F$  correspond to subalgebras  $B \subset \mathcal{O}(G)$  satisfying

$$\Delta(B) \subset A \otimes B$$

and stable under the antipode. In [Tak79], Theorem 3, the  $k$ -subgroups  $F$  for which  $G/F$  is affine correspond to such subalgebras  $B$  for which  $B \rightarrow A$  is faithfully flat.

## 5.1

Let  $k$  be a pseudocompact ring.<sup>146</sup> Let  $G$  be a topologically flat infinitesimal formal  $k$ -group, let  $A$  be its affine algebra, and let  $B \subset A$  be a closed subalgebra. Put

$$X = \text{Spf}(B),$$

and let

$$r : G \longrightarrow X$$

be the epimorphism induced by the inclusion  $B \subset A$ . We shall determine when  $r$  makes  $X$  the quotient of  $G$  on the right by a subgroup  $H$  (cf. 2.4).<sup>147</sup>

<sup>146</sup>The original assumes in 5.1 that  $k$  is a field. In fact, that hypothesis can be replaced by flatness hypotheses; the editors modified 5.1–5.1.5 accordingly.

<sup>147</sup>The editors replaced “left” by “right” and modified the statement of Proposition 5.1 so as to display both the equivalent conditions (i), (ii) and the conclusion  $\text{Spf}(B) \simeq G/H$  more clearly.

**Proposition 5.1.** *Let  $G$  be a topologically flat infinitesimal formal  $k$ -group, let  $A$  be its affine algebra, let  $I_A$  be the augmentation ideal of  $A$ , and let  $B$  be a closed subalgebra of  $A$ . Put*

$$I_B = B \cap I_A, \quad J_B = AI_B.$$

*Assume that  $A$ , and also*

$$J_B^n/J_B^{n+1}$$

*for every  $n \geq 0$ , are topologically flat over  $k$ . Then the following two assertions are equivalent:*

(i) *For every  $x \in B$ ,*

$$\Delta_A(x) - x \hat{\otimes} 1 \in A \hat{\otimes}_k J_B.$$

(ii) *Let*

$$\pi : A \longrightarrow A/J_B$$

*be the projection, and put*

$$\tau_1(a) = a \hat{\otimes} 1.$$

*The sequence*

$$B \longrightarrow A \begin{array}{c} \xrightarrow{\tau_1} \\ \xrightarrow{(\text{id}_A \hat{\otimes} \pi) \Delta_A} \end{array} A \hat{\otimes}_k (A/J_B) \quad (*)$$

*is exact. Equivalently,  $B$  is the set of all  $x \in A$  for which*

$$\Delta_A(x) - x \hat{\otimes} 1 \in A \hat{\otimes}_k J_B.$$

*In this case,*

$$H = \text{Spf}(A/J_B)$$

*is a formal subgroup of  $G$ , and the sequence*

$$G \times H \begin{array}{c} \xrightarrow{\text{pr}_1} \\ \xrightarrow{\lambda} \end{array} G \longrightarrow \text{Spf}(B) \quad (**)$$

*is exact, where  $\lambda$  is the restriction of the multiplication of  $G$  to  $G \times H$ . Thus*

$$\text{Spf}(B) \simeq G/H.$$

*Source note.* In (\*), the French PDF labels the lower arrow  $(\pi \hat{\otimes} \text{id}_A) \Delta_A$ . The displayed equivalent condition immediately before and after the sequence, as well as the stated right-quotient convention, require  $(\text{id}_A \hat{\otimes} \pi) \Delta_A$ , which is used here.

*Proof.* Put

$$\bar{A} = A/J_B, \quad H = \text{Spf}(\bar{A}).$$

Then  $H$  is a formal subvariety of  $G$ . Since  $J_B \subset I_A$ , the augmentation  $\varepsilon_A$  induces a continuous  $k$ -algebra morphism

$$\bar{\varepsilon} : \bar{A} \longrightarrow k.$$

If (i) holds, then

$$\Delta_A(I_B) \subset I_B \hat{\otimes} 1 + A \hat{\otimes} J_B.$$

Therefore  $\Delta_A$  induces, after passage to the quotient, a diagonal morphism

$$\bar{\Delta} : \bar{A} \longrightarrow \bar{A} \hat{\otimes}_k \bar{A}.$$

The maps  $\bar{\Delta}$  and  $\bar{\varepsilon}$  make  $H$  a formal submonoid of  $G$ . Since  $G$  is infinitesimal, so is  $H$ ; Proposition 2.7 then shows that  $H$  is a formal subgroup of  $G$ . By the definition of  $G/H$  (cf. 2.4), assertion (ii) implies the final assertion of the proposition.

It is clear that (ii) implies (i). The proof of the converse occupies 5.1.1–5.1.5.

## 5.1.1

Consider the following category  $\mathcal{C}$ . An object of  $\mathcal{C}$  is a pair  $(A, J)$  consisting of a profinite  $k$ -algebra  $A$  and a closed ideal  $J \subset A$ . A morphism

$$\psi : (A, J) \longrightarrow (A', J')$$

in  $\mathcal{C}$  is a continuous  $k$ -algebra homomorphism

$$A \longrightarrow A'$$

that maps  $J$  into  $J'$ .

The assignment

$$(A, J) \longmapsto (\mathrm{Spf}(A/J), \mathrm{Spf}(A))$$

is an antiequivalence from  $\mathcal{C}$  to the category of pairs  $(Z, Y)$  consisting of a formal  $k$ -variety  $Y$  and a formal subvariety  $Z \subset Y$ . A morphism

$$\varphi : (Z, Y) \longrightarrow (Z', Y')$$

is a morphism of formal  $k$ -varieties

$$Y \longrightarrow Y'$$

that maps  $Z$  into  $Z'$ .

A cogroup structure on an object  $(A, J)$  of  $\mathcal{C}$  consists of a formal-group structure on  $\mathrm{Spf}(A)$  satisfying the following conditions (with the notation of 2.1):

- (1)  $\Delta_A(J) \subset J \widehat{\otimes}_k A + A \widehat{\otimes}_k J$ ,
- (2)  $\varepsilon_A(J) = 0$ ,
- (3)  $c_A(J) \subset J$ .

These conditions also say that

$$H = \mathrm{Spf}(A/J)$$

is a formal subgroup of

$$G = \mathrm{Spf}(A).$$

148

Suppose in addition that  $A$  is local; equivalently,  $\mathrm{Spf}(A)$  is an infinitesimal formal group. If  $J \neq A$ , then conditions (2) and (3) follow from (1). Indeed, a proper closed ideal  $J$  of  $A$  is contained in the augmentation ideal  $I_A$ , so (2) holds. Moreover,

$$M = \mathrm{Spf}(A/J)$$

is a formal submonoid of  $G$ . Since  $G$  is infinitesimal, Proposition 2.7 shows that  $M$  is a formal subgroup of  $G$ ; hence (3) also holds.

---

<sup>148</sup>If  $(A', J')$  is a second cogroup of  $\mathcal{C}$ , corresponding to a pair  $H' \subset G'$  of formal  $k$ -groups, then giving a cogroup morphism

$$(A', J') \longrightarrow (A, J)$$

is equivalent to giving a morphism of formal  $k$ -groups

$$G \longrightarrow G'$$

that maps  $H$  into  $H'$ .

## 5.1.2

Denote by  $\mathbf{Alpg}/k$  the category of *graded profinite  $k$ -algebras*. An object of this category consists of a sequence

$$A_0, A_1, \dots, A_n, \dots$$

of pseudocompact  $k$ -modules, together with a profinite algebra structure on

$$\prod_{n \geq 0} A_n$$

such that

$$A_n A_m \subset A_{m+n}.$$

Here  $A_n$  is identified with a subset of  $\prod_{i \geq 0} A_i$  by means of the canonical injection. A morphism

$$\psi : (A_n) \longrightarrow (B_n)$$

is a sequence of continuous linear maps

$$\psi_n : A_n \longrightarrow B_n$$

such that

$$\psi_{m+n}(aa') = \psi_m(a)\psi_n(a')$$

for  $a \in A_m$  and  $a' \in A_n$ .

*Definitions.* Two graded profinite  $k$ -algebras  $(A_n)$  and  $(B_n)$  clearly have a coproduct<sup>149</sup> in  $\mathbf{Alpg}/k$ . Its  $n$ -th component is

$$\prod_{i=0}^n A_i \widehat{\otimes}_k B_{n-i},$$

the product of the pseudocompact  $k$ -modules

$$A_i \widehat{\otimes}_k B_{n-i}.$$

This coproduct will be denoted by

$$(A_n) \widehat{\otimes}_k (B_n).$$

A cogroup structure on an object  $(A_n)$  of  $\mathbf{Alpg}/k$  is therefore the datum of continuous  $k$ -linear maps

$$\Delta_n : A_n \longrightarrow \prod_{i=0}^n A_i \widehat{\otimes}_k A_{n-i}, \quad \varepsilon : A_0 \longrightarrow k,$$

which induce on  $\prod_{n \geq 0} A_n$ , upon setting  $\varepsilon(A_i) = 0$  for  $i \geq 1$ , a cogroup structure in  $\mathbf{Alp}/k$ .

Finally, for every object  $(A, J)$  of  $\mathcal{C}$ , denote by

$$\mathrm{Gr}_J(A)$$

the graded profinite algebra associated with the filtration of  $A$  by the closures  $\overline{J^n}$  of the powers of  $J$ . Thus

$$\mathrm{Gr}_J(A)_n = \overline{J^n} / \overline{J^{n+1}},$$

and the multiplication of  $\mathrm{Gr}_J(A)$  is induced by that of  $A$ .

<sup>149</sup>The editors replaced “direct sum” by “coproduct”.

**Lemma.** <sup>150</sup> Let  $U, V$  be two pseudocompact  $k$ -modules, with  $U$  topologically flat, and let

$$U = U_0 \supset U_1 \supset \cdots, \quad V = V_0 \supset V_1 \supset \cdots$$

be two decreasing sequences of closed  $k$ -submodules. Filter the completed tensor product

$$W = U \widehat{\otimes}_k V$$

by means of the closed submodules

$$W_n = U_n \widehat{\otimes}_k V_0 + U_{n-1} \widehat{\otimes}_k V_1 + \cdots + U_0 \widehat{\otimes}_k V_n.$$

Suppose that every  $U_i/U_{i+1}$  is topologically flat over  $k$ , so that  $U/U_n$ , and hence also  $U_n$ , is topologically flat for every  $n$ . Then, for every  $n$ , there is an isomorphism

$$W_n/W_{n+1} \simeq \bigoplus_{i+j=n} (U_i/U_{i+1}) \widehat{\otimes}_k (V_j/V_{j+1}).$$

*Proof.* For every  $i, j \geq 0$ , put

$$W_{i,j} = U_i \widehat{\otimes}_k V_j, \quad \overline{W}_{i,j} = (U_i/U_{i+1}) \widehat{\otimes}_k (V_j/V_{j+1}).$$

We prove by induction on  $n$  that the natural map

$$\pi_n : W_n \longrightarrow \bigoplus_{i+j=n} \overline{W}_{i,j}$$

is surjective and that the inclusion

$$W_{n+1} \subset \ker(\pi_n)$$

is an equality. For  $n = 0$ , the projection

$$\pi_0 : U_0 \widehat{\otimes}_k V_0 \longrightarrow (U_0/U_1) \widehat{\otimes}_k (V_0/V_1)$$

is surjective. Since  $U_0$ ,  $U_0/U_1$ , and hence  $U_1$ , are topologically flat over  $k$ , we have

$$\ker(\pi_0) = U_0 \widehat{\otimes}_k V_1 + U_1 \widehat{\otimes}_k V_0,$$

and moreover

$$(U_0 \widehat{\otimes}_k V_1) \cap (U_1 \widehat{\otimes}_k V_0) = U_1 \widehat{\otimes}_k V_1.$$

Suppose now that  $n > 0$  and that the result has been proved for  $n - 1$ . Put

$$M_0 = U_0 \widehat{\otimes}_k V_n, \quad S_0 = \sum_{i=1}^n U_i \widehat{\otimes}_k V_{n-i}.$$

We have  $S_0 \subset U_1 \widehat{\otimes}_k V_0$ . Applying the preceding argument to  $V_0 \supset V_n$  in place of  $V_0 \supset V_1$ , we obtain

$$M_0 \cap S_0 \subset (U_0 \widehat{\otimes}_k V_n) \cap (U_1 \widehat{\otimes}_k V) = U_1 \widehat{\otimes}_k V_n.$$

It follows that

$$M_0 \cap S_0 = U_1 \widehat{\otimes}_k V_n.$$

Since  $W_n = M_0 + S_0$ , we obtain a commutative diagram with exact rows, where

$$U'_i = U_{i+1}, \quad W'_{i,n-1-i} = W_{i+1,n-1-i} \quad (0 \leq i \leq n-1) :$$

<sup>150</sup>In the original, the lemma is stated when  $k$  is a field, and in that case the proof is left to the reader.



$$\begin{array}{ccccccc}
0 & \longrightarrow & S_0 & \longrightarrow & W_n & \longrightarrow & (U_0/U_1) \hat{\otimes}_k V_n \longrightarrow 0 \\
& & \downarrow \pi'_{n-1} & & \downarrow \pi_n & & \downarrow p \\
0 & \longrightarrow & \bigoplus_{i=0}^{n-1} \overline{W}'_{i,n-1-i} & \longrightarrow & \bigoplus_{i=0}^n \overline{W}_{i,n-1} & \longrightarrow & \overline{W}_{0,n} \longrightarrow 0
\end{array}$$

*Source note.* The source-bound diagram is reproduced exactly. Its lower middle term is printed  $\bigoplus_{i=0}^n \overline{W}_{i,n-1}$ ; the natural map  $\pi_n$  defined immediately above instead has target  $\bigoplus_{i+j=n} \overline{W}_{i,j}$ , which would give  $\bigoplus_{i=0}^n \overline{W}_{i,n-i}$ . No silent change has been made inside the Loop 1 image.

Then  $p$  is surjective, with kernel

$$(U_0/U_1) \hat{\otimes}_k V_{n+1}.$$

Moreover, by the induction hypothesis applied to the sequence  $(U'_i)$ , the map  $\pi'_{n-1}$  is surjective, with kernel

$$W'_n = \sum_{i=1}^n W_{i,n+1-i}.$$

It follows that  $\pi_n$  is surjective and that

$$W_{n+1} \subset \ker(\pi_n)$$

is an equality. This proves the lemma.

Let us return to an object  $(A, J)$  of  $\mathcal{C}$ . By 0.2.G, the hypothesis that every

$$\overline{J^n}/\overline{J^{n+1}}$$

is topologically flat over  $k$  is equivalent to saying that  $\mathrm{Gr}_J(A)$  is topologically flat over  $k$ .

**Corollary.** *Let  $\mathcal{P}$  be the full subcategory of  $\mathcal{C}$  consisting of the objects  $(A, J)$  such that  $A$  and  $\mathrm{Gr}_J(A)$  are topologically flat over  $k$ . Then the functor*

$$\mathcal{P} \longrightarrow \mathbf{Alpg}/k, \quad (A, J) \longmapsto \mathrm{Gr}_J(A),$$

*commutes with finite coproducts and hence carries a cogroup of  $\mathcal{P}$  to a cogroup of  $\mathbf{Alpg}/k$ .*

*In particular, if  $k$  is a field, then for every cogroup  $(A, J)$  of  $\mathcal{C}$ , the algebra  $\mathrm{Gr}_J(A)$  is a cogroup of  $\mathbf{Alpg}/k$ .<sup>151</sup>*

### 5.1.3

Identify every profinite  $k$ -algebra  $\Gamma$  with the graded profinite  $k$ -algebra  $(\Gamma_n)_{n \geq 0}$  such that

$$\Gamma_0 = \Gamma, \quad \Gamma_n = 0 \quad \text{if } n > 0.$$

In particular, if  $(A_n)_{n \geq 0}$  is a graded profinite  $k$ -algebra, we shall regard  $A_0$ , without distinction, as either a profinite  $k$ -algebra or a graded profinite  $k$ -algebra. Denote by

$$\rho : (A_n) \longrightarrow A_0$$

<sup>151</sup>Thus, to a pair  $H \subset G$  of formal  $k$ -groups one associates the “formal completion  $\hat{G}_H$  of  $G$  along  $H$ ”, which is a formal  $k$ -group. Moreover, we shall see in 5.1.3–5.1.4 that the inclusion  $\sigma : H \hookrightarrow \hat{G}_H$  has a retraction  $\pi : \hat{G}_H \rightarrow H$ , and that the formal  $k$ -group  $N = \ker(\pi)$  identifies, as a formal variety, with the completion of the homogeneous space  $G/H$  along the unit section. This will be useful in 5.2.2.

the morphism of  $\mathbf{Alpg}/k$  such that

$$\rho_0 = \text{id}_{A_0}, \quad \rho_n = 0 \quad \text{if } n > 0.$$

Similarly,

$$\tau : A_0 \longrightarrow (A_n)$$

will denote the section of  $\rho$  such that

$$\tau_0 = \text{id}_{A_0}, \quad \tau_n = 0 \quad \text{if } n > 0.$$

Every cogroup structure on  $(A_n)_{n \in \mathbb{N}}$  induces a cogroup structure on  $A_0$  for which  $\rho$  and  $\tau$  are cogroup homomorphisms. In this case, let  $I_0$  be the augmentation ideal of  $A_0$ , and put

$$A'_n = A_n / I_0 A_n \quad (n \geq 0),$$

so that  $A'_0 = k$ .

Then  $(A'_n)_{n \in \mathbb{N}}$  is a cogroup in  $\mathbf{Alpg}/k$ . Indeed, since

$$A_0 = I_0 \oplus k \cdot 1,$$

the direct summand

$$I_0 \widehat{\otimes}_k A_n \simeq I_0 A_n$$

of  $A_n$  is defined for every  $n$ . Since  $\tau$  is a section of  $\rho$ , it follows from 2.4.A that the cogroup  $(A_n)_{n \in \mathbb{N}}$  is the “semidirect coproduct” of  $A_0$  and the cogroup  $(A'_n)_{n \in \mathbb{N}}$ .

More precisely,  $(A'_n)_{n \in \mathbb{N}}$  is isomorphic, as an object of  $\mathbf{Alpg}/k$ , to the kernel of the following pair:

$$(A_n) \xrightarrow[\text{(id} \otimes \rho) \Delta]{\tau_1} (A_n) \widehat{\otimes}_k A_0$$

Here

$$\Delta : (A_n) \longrightarrow (A_n) \widehat{\otimes} (A_n)$$

is the comultiplication of  $(A_n)$ , and

$$\tau_1(x) = x \widehat{\otimes} 1.$$

Identifying  $A'_n$  with its image in  $A_n$ , the map

$$A'_n \widehat{\otimes}_k A_0 \longrightarrow A_n, \quad a'_n \widehat{\otimes} a_0 \longmapsto a'_n a_0,$$

is an isomorphism in  $\mathbf{Alpg}/k$ .

This is not an isomorphism of cogroups. Rather,

$$\Delta(A') \subset A \widehat{\otimes} A',$$

and

$$(\gamma'_\rho \widehat{\otimes} \text{id}) \circ \Delta|_{A'} = \Delta',$$

where  $\Delta'$  is the comultiplication of  $A'$  and  $\gamma'_\rho : A \rightarrow A'$  is the projection (cf. 2.4.A).

## 5.1.4

Let  $(A, J)$  be an object of  $\mathcal{C}$ , and let

$$(A_n) = \mathrm{Gr}_J(A)$$

be the associated object of  $\mathbf{Alpg}/k$ ; thus

$$A_n = \overline{J^n}/\overline{J^{n+1}} \quad (n \geq 0).$$

It is clear that the algebra

$$\mathcal{A} = \prod_{n \geq 0} A_n$$

is generated by  $A_0$  and  $A_1$ . In other words, for  $n \geq 1$ , the multiplication map

$$A_1 \widehat{\otimes}_{A_0} A_1 \widehat{\otimes}_{A_0} \cdots \widehat{\otimes}_{A_0} A_1 \longrightarrow A_n \quad (1)$$

is surjective.

Suppose further that  $(A, J)$  is a cogroup of  $\mathcal{C}$  and that  $A$  and the quotients

$$\overline{J^n}/\overline{J^{n+1}}$$

are topologically flat over  $k$ . By Corollary 5.1.2,  $\mathrm{Gr}_J(A)$  is then a cogroup of  $\mathbf{Alpg}/k$ . Hence, by 5.1.3, if one puts

$$A'_n = \left\{ x \in \overline{J^n}/\overline{J^{n+1}} \mid \Delta(x) - x \widehat{\otimes} 1 \in \bigoplus_{i=1}^n (\overline{J^{n-i}}/\overline{J^{n-i+1}}) \widehat{\otimes}_k (\overline{J^i}/\overline{J^{i+1}}) \right\}, \quad (2)$$

then the map

$$A'_n \widehat{\otimes}_k A_0 \longrightarrow A_n, \quad a'_n \widehat{\otimes} a_0 \longmapsto a'_n a_0,$$

is an isomorphism in  $\mathbf{Alpg}/k$ .<sup>152</sup>

*Source note.* In the displayed element-wise description of this map, the French PDF prints  $a'_n \widehat{\otimes} a'_n \mapsto a'_n a_0$ . The domain and the product require  $a'_n \widehat{\otimes} a_0 \mapsto a'_n a_0$ , which is used here.

Let  $I_0$  denote the augmentation ideal of  $A_0$ . It follows from (1) and the following commutative diagram, in which

$$A'_1 \widehat{\otimes}^n = \underbrace{A'_1 \widehat{\otimes}_k \cdots \widehat{\otimes}_k A'_1}_{n \text{ factors}}$$

that:

$$\begin{array}{ccccc} A_1 \widehat{\otimes}_{A_0} \cdots \widehat{\otimes}_{A_0} A_1 & \xleftarrow{\sim} & A'_1 \widehat{\otimes}^n \widehat{\otimes}_k A_0 & \xleftarrow{\sim} & (A'_1 \widehat{\otimes}^n \widehat{\otimes}_k I_0) \oplus A'_1 \widehat{\otimes}^n \\ \downarrow m & & \downarrow m' \widehat{\otimes} \mathrm{id} & & \downarrow (m' \widehat{\otimes} \mathrm{id}) \oplus m' \\ A_n & \xleftarrow{\sim} & A'_n \widehat{\otimes}_k A_0 & \xleftarrow{\sim} & (A'_n \widehat{\otimes}_k I_0) \oplus A'_n \end{array}$$

<sup>152</sup>Let  $G$ ,  $H$ , and  $\widehat{G}_H$  be the formal  $k$ -groups corresponding to  $A$ ,  $A_0 = A/J$ , and  $\mathrm{Gr}_J(A)$ , respectively. Then  $\tau : A_0 \hookrightarrow \mathrm{Gr}_J(A)$  corresponds to a retraction  $\pi : \widehat{G}_H \rightarrow H$  of the inclusion  $H \hookrightarrow \widehat{G}_H$ , and the preceding assertion means that  $\widehat{G}_H$  is the semidirect product of  $N = \ker(\pi)$  by  $H$ .

the multiplication map

$$m' : A'_1 \widehat{\otimes}_k \cdots \widehat{\otimes}_k A'_1 \longrightarrow A'_n \quad (3)$$

is surjective. In other words, the profinite  $k$ -algebra

$$\mathcal{A}' = \prod_{n \geq 0} A'_n$$

is generated by its degree-one terms.

Let us now return to hypothesis (i) of Proposition 5.1.<sup>153</sup> Let  $G$  be a topologically flat infinitesimal formal  $k$ -group,  $A$  its affine algebra,  $I_A$  the augmentation ideal of  $A$ ,  $B$  a closed subalgebra of  $A$ , and

$$J = AI_B, \quad I_B = B \cap I_A.$$

Let  $H$  be the formal subgroup  $\mathrm{Spf}(A/J)$ , and let

$$\pi : A \longrightarrow A/J$$

be the projection. Suppose that

$$A/\overline{J^n}$$

is topologically flat over  $k$  for every  $n \geq 1$ , and that  $B$  is contained in the kernel  $\widetilde{B}$  of the following pair:

$$A \xrightarrow[\mathrm{id}_A \widehat{\otimes}_k \pi \Delta_A]{\tau_1} A \widehat{\otimes}_k (A/J).$$

Put

$$(A_n) = \mathrm{Gr}_J(A),$$

let

$$I_0 = I_A/J$$

be the augmentation ideal of  $A_0 = A/J$ , and define  $(A'_n)_{n \in \mathbb{N}}$  as in (2) above. Let  $(B_n)_{n \in \mathbb{N}}$ , respectively  $(\widetilde{B}_n)_{n \in \mathbb{N}}$ , be the object of  $\mathbf{Alg}/k$  associated with the filtration of  $B$ , respectively of  $\widetilde{B}$ , induced by that of  $A$ ; these filtrations are defined by the ideals

$$B \cap \overline{J^n}, \quad \text{respectively} \quad \widetilde{B} \cap \overline{J^n}.$$

It is clear that

$$B_n \subset \widetilde{B}_n \subset A'_n \quad \text{for every } n,$$

and that

$$\mathcal{B} = \prod_{n \geq 0} B_n \subset \widetilde{\mathcal{B}} = \prod_{n \geq 0} \widetilde{B}_n$$

are subalgebras of

$$\mathcal{A}' = \prod_{n \geq 0} A'_n.$$

On the other hand,  $J$ , respectively  $\overline{J^2}$ , is the image in  $A$  of

$$I_B \widehat{\otimes}_k A, \quad \text{respectively} \quad \overline{I_B^2} \widehat{\otimes}_k A.$$

Consequently, the map

$$(I_B/\overline{I_B^2}) \widehat{\otimes}_k A \longrightarrow J/\overline{J^2} = A_1 \simeq A'_1 \widehat{\otimes}_k A_0$$

<sup>153</sup>The editors expanded the original in what follows and moved to the end the “supplement”  $\overline{I_B^n} = B \cap \overline{J^n}$ , which is not needed to prove Proposition 5.1.

is surjective, and it factors through

$$(I_B/\overline{I_B^2}) \widehat{\otimes}_k A_0.$$

Since

$$A_0 = k \cdot 1 \oplus I_0,$$

the following commutative exact diagram shows that  $A'_1$  is the image of  $I_B/\overline{I_B^2}$ :

$$\begin{array}{ccc} (I_B/\overline{I_B^2}) \widehat{\otimes}_k A_0 & \xleftarrow{\sim} & ((I_B/\overline{I_B^2}) \widehat{\otimes}_k I_0) \oplus (I_B/\overline{I_B^2}) \\ \downarrow & & \downarrow \\ J/\overline{J^2} & \xleftarrow{\sim} & (A'_1 \widehat{\otimes}_k I_0) \oplus A'_1 \\ \downarrow & & \downarrow \\ 0 & & 0 \end{array}$$

Thus

$$B_1 = A'_1.$$

Since  $\mathcal{A}'$  is generated by  $A'_1$ , it follows that, for every  $n \geq 1$ ,

$$A'_n \subset B_n \subset \widetilde{B}_n \subset A'_n,$$

and hence

$$B_n = \widetilde{B}_n = A'_n.$$

154

Finally, because  $G$  is infinitesimal,

$$\mathfrak{r}(A) = I_A,$$

and hence the ideals  $\overline{I_A^n}$  tend to zero (cf. 0.1.2). A fortiori, the ideals  $\overline{J^n}$  tend to zero, so the induced filtrations on  $\widetilde{B}$  and  $B$  are separated. Moreover, since  $B$  is a closed subalgebra of  $A$ , it is complete for the topology defined by the ideals  $B \cap \overline{J^n}$ . It therefore follows from [CA], § V.7, Lemma 1 (see also 5.1.5 below) that

$$B = \widetilde{B}.$$

This completes the proof of Proposition 5.1.

There is also the following supplement. For every  $n$ ,

$$\overline{J^n} = A \overline{I_B^n}$$

is the image in  $A$  both of

$$A \widehat{\otimes}_B \overline{I_B^n}$$

and of

$$A \widehat{\otimes}_B (B \cap \overline{J^n}).$$

<sup>154</sup>With the notation of editorial note 149, this implies that the formal completion of  $G/H$  along the unit section, whose affine algebra is  $\prod_n B_n$ , is isomorphic as a formal variety to the formal  $k$ -group  $N$ .

By hypothesis, the affine algebra  $A/J$  of the formal subgroup  $H$  is topologically flat over  $k$ . Thus, by Theorem 2.4, the morphism

$$G = \mathrm{Spf}(A) \longrightarrow G/H = \mathrm{Spf}(B)$$

is surjective and topologically flat. Consequently,

$$A \widehat{\otimes}_B \overline{I}_B^n = \overline{J}^n = A \widehat{\otimes}_B (B \cap \overline{J}^n),$$

which implies

$$\overline{I}_B^n = B \cap \overline{J}^n \quad \text{for every } n.$$

This also follows from the surjectivity of the maps

$$\overline{I}_B^n / \overline{I}_B^{n+1} \longrightarrow A'_n = (B \cap \overline{J}^n) / (B \cap \overline{J}^{n+1})$$

and from 5.1.5(ii) below, applied to

$$B'_n = \overline{I}_B^n, \quad B_n = B \cap \overline{J}^n.$$

*Source note.* In the preceding  $n$ -indexed display, the French PDF prints  $A'_i$ . The surrounding numerators and denominators, and the application to  $B'_n$  and  $B_n$ , require  $A'_n$ , which is used here.

### 5.1.5

**Lemma 5.1.5.** <sup>155</sup>

(i) *Let  $M$  and  $N$  be two abelian groups filtered by decreasing sequences of subgroups*

$$(M_n)_{n \in \mathbb{Z}}, \quad (N_n)_{n \in \mathbb{Z}}.$$

*Suppose that*

$$\bigcup_{n \in \mathbb{Z}} M_n = M, \quad \bigcup_{n \in \mathbb{Z}} N_n = N,$$

*that*

$$\bigcap_{n \in \mathbb{Z}} M_n = 0, \quad \bigcap_{n \in \mathbb{Z}} N_n = 0,$$

*and that  $M$  is complete for the topology defined by the  $M_n$ . Let*

$$f : M \longrightarrow N$$

*be a homomorphism of filtered groups.*

a) *If  $f$  induces a surjection on the associated graded groups, then  $f$  is surjective and  $N$  is complete for the topology defined by the  $N_n$ .*

b) *If  $f$  induces an injection on the associated graded groups, then  $f$  is injective.*

<sup>155</sup>The original stated only part (ii). For the reader's convenience, the editors stated in part (i) Lemma 1 of [CA], § V.7.

(ii) Let  $B$  be an abelian group, and let

$$B = B'_0 \supset B'_1 \supset \cdots, \quad B = B_0 \supset B_1 \supset \cdots$$

be two separated filtrations of  $B$  by subgroups such that

$$B'_n \subset B_n \quad \text{for every } n.$$

Suppose that  $B$  is complete for the topology defined by the filtration  $(B'_n)$ . If the map

$$B'_i/B'_{i+1} \longrightarrow B_i/B_{i+1}$$

is surjective for every  $i$ , then

$$B'_n = B_n \quad \text{for every } n.$$

*Proof.* Part (i) is Lemma 1 of [CA], § V.7 (see also [BAC], III, § 2.8). Part (ii) follows by taking

$$M = B'_n \supset B'_{n+1} \supset \cdots, \quad N = B_n \supset B_{n+1} \supset \cdots.$$

## 5.2

Throughout the rest of Section 5,  $k$  denotes a perfect field of characteristic  $p > 0$ .

Put

$$\overline{\mathbb{N}} = \mathbb{N} \cup \{\infty\}.$$

If  $B$  is a profinite  $k$ -algebra and  $r \in \overline{\mathbb{N}}$ , denote by

$$((x^{p^r}))_{x \in \mathfrak{r}(B)}$$

the closed ideal of  $B$  generated by the elements  $x^{p^r}$ , where  $x$  ranges over the radical  $\mathfrak{r}(B)$  of  $B$ . If  $r = \infty$ , use the same notation with the convention that

$$((x^{p^\infty}))_{x \in \mathfrak{r}(B)}$$

is the zero ideal. In both cases,  $B_r$  denotes the quotient

$$B_r = B / ((x^{p^r}))_{x \in \mathfrak{r}(B)}.$$

We say that  $B$  is of height at most  $r$  if

$$((x^{p^r}))_{x \in \mathfrak{r}(B)}$$

is the zero ideal. If this holds and  $r$  is finite, we say that  $B$  is of finite height.

Consider in particular the case

$$B = k[[\omega]],$$

where  $\omega$  is a pseudocompact  $k$ -vector space (cf. 1.2.5).<sup>156</sup> We then say that  $B$  is a formal power series algebra and that  $B_r$  is a truncated formal power series algebra ( $r \in \overline{\mathbb{N}}$ ); thus, by convention,

$$B = B_\infty$$

<sup>156</sup>In the original, the author uses “linearly compact vector space,” which is equivalent to “pseudocompact vector space” (cf. [BAC], §III.2, Exercises 15 a), 19 a), and 20 d)). The editors preferred to retain the terminology “pseudocompact,” used up to this point.

is also called “truncated.” If  $B = k[[\omega]]$ , we also write

$$((x^{p^r}))_{x \in \omega}$$

instead of

$$((x^{p^r}))_{x \in \mathfrak{t}(B)}.$$

**Notation.** Let  $\omega$  be a pseudocompact  $k$ -vector space filtered by an increasing sequence of closed vector subspaces

$$0 = \omega_0 \subset \omega_1 \subset \omega_2 \subset \omega_3 \subset \cdots.$$

- (a) The closed ideal of  $k[[\omega]]$  generated by the elements  $x^{p^r}$ , where  $r$  ranges over  $\mathbb{N}$  and  $x$  ranges over  $\omega_r$ , will be denoted by

$$((x^{p^r}))_{x \in \omega_r}.$$

- (b) For  $r \in \mathbb{N}$ , denote by  ${}^r\omega$  the filtered pseudocompact vector space defined by

$${}^r\omega_i = \omega_i \quad \text{if } i < r, \quad {}^r\omega_i = \omega \quad \text{if } i \geq r.$$

*Source note.* In Notation (a), the French PDF prints the subscript  $r \in \omega_r$ . The preceding defining sentence and every subsequent use require  $x \in \omega_r$ , which is used here.

**Theorem (Dieudonné–Cartier).** *Let*

$$H \longrightarrow G$$

*be a monomorphism of infinitesimal formal groups over a perfect field  $k$  of characteristic  $p > 0$ . Let  $B$  be the affine algebra of the homogeneous space  $G/H$ , and suppose that one of the following three conditions holds:*<sup>157</sup>

- (i)  *$B$  has finite height (in particular, this holds if  $G$  has finite height).*
- (ii)  *$B$  is a complete Noetherian local ring.*
- (iii)  *$B$  is a reduced ring.*

*Then  $B$  is isomorphic to the completed tensor product of a finite family of truncated formal power series algebras.*

The proof of this theorem occupies 5.2.1–5.2.5.

### 5.2.1

Let  $A$  be the affine algebra of  $G$ , let  $I_A$  be its augmentation ideal, and put

$$I = B \cap I_A.$$

By 2.4,

$$H = \mathrm{Spf}(A/\overline{AI})$$

and

$$B = \{x \in A \mid \Delta(x) - x \hat{\otimes} 1 \in A \hat{\otimes} \overline{AI}\}.$$

<sup>157</sup>The editors replaced  $H \setminus G$  by  $G/H$ , and did the same in the proof; they also added condition (iii).



*Source note.* The French PDF prints  $\Delta(x) - 1 \hat{\otimes} x$  here. The right-quotient convention fixed in 5.0 and Proposition 5.1, together with the defining condition  $(\mathrm{id}_A \hat{\otimes} \pi)\Delta(x) = x \hat{\otimes} 1$ , requires  $\Delta(x) - x \hat{\otimes} 1$ , which is used above.

Put

$$\omega = I/\overline{I^2}.$$

For each  $r$ , denote by  $I_r$  the closed subideal of  $I$  consisting of the elements  $x$  such that

$$x^{p^r} = 0,$$

and denote by  $\omega_r$  the canonical image of  $I_r$  in  $\omega$ . We shall prove the following proposition.

**Proposition.** *If there exists a continuous section*

$$\sigma : \omega \longrightarrow I$$

*of the projection  $I \rightarrow I/\overline{I^2}$  such that*

$$\sigma(\omega_r) \subset I_r$$

*for every  $r$ , then  $B$  is isomorphic to*

$$k[[\omega]] / ((x^{p^r}))_{x \in \omega_r}.$$

Indeed, such a section extends to a continuous morphism

$$k[[\omega]] \longrightarrow B,$$

which factors through

$$B' = k[[\omega]] / ((x^{p^r}))_{x \in \omega_r}.$$

In 5.2.2–5.2.5 we prove that the resulting morphism

$$\varphi : B' = k[[\omega]] / ((x^{p^r}))_{x \in \omega_r} \longrightarrow B$$

is an isomorphism.

**5.2.1.A.** <sup>158</sup> For every  $r \in \mathbb{N}$ , put

$$\mathcal{I}_r = \overline{I^2} + I_r,$$

so that

$$\mathcal{I}_r / \overline{I^2} \simeq \omega_r.$$

There is then a commutative diagram with exact rows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{I}_{r-1} & \longrightarrow & \mathcal{I}_r & \longrightarrow & \mathcal{I}_r / \mathcal{I}_{r-1} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \wr \\ 0 & \longrightarrow & \omega_{r-1} & \longrightarrow & \omega_r & \longrightarrow & \omega_r / \omega_{r-1} \longrightarrow 0 \end{array}$$

---

<sup>158</sup>The editors added 5.2.1.A and 5.2.1.B.

Since  $k$  is a field, the rows split. Thus a pseudobasis  $\mathcal{B}_{r-1}$  of  $\mathcal{I}_{r-1}$  can be completed to a pseudobasis

$$\mathcal{B}_{r-1} \cup \mathcal{B}'_r$$

of  $\mathcal{I}_r$ . The closed subspace  $S_r$  with pseudobasis  $\mathcal{B}'_r$  is then a complement of  $\mathcal{I}_{r-1}$  in  $\mathcal{I}_r$ , and the projection

$$\pi : I \longrightarrow \omega$$

induces an isomorphism from  $S_r$  onto a complement  $\omega'_r$  of  $\omega_{r-1}$  in  $\omega_r$ .

Let  $\mathcal{I}_\infty$  be the closed ideal

$$\mathcal{I}_\infty = \overline{\bigcup_r \mathcal{I}_r}.$$

It likewise has a complement  $S_\infty$  in  $I$ , and  $\pi$  induces an isomorphism from

$$\mathcal{I}_\infty / \overline{I^2}$$

onto the closure

$$\omega_\infty = \overline{\bigcup_r \omega_r}$$

of the union of the  $\omega_r$ , and an isomorphism from  $S_\infty$  onto a complement  $\omega'_\infty$  of  $\omega_\infty$  in  $\omega$ . Denote the latter isomorphism by

$$\eta : S_\infty \xrightarrow{\sim} \omega'_\infty.$$

We thus obtain continuous linear maps:

$$\begin{array}{ccc} \overline{I^2} \times S_\infty \times \widehat{\bigoplus_r S_r} & \xrightarrow{\phi} & I \\ \eta \times \theta \downarrow & & \\ \omega = \omega'_\infty \times \omega_\infty & & \end{array}$$

Here

$$\widehat{\bigoplus_r S_r}$$

is the direct sum of the  $S_r$  in  $\text{PC}(k)$ , that is,

$$\left( \prod_r S_r^\dagger \right)^*$$

(cf. editorial note (16) to 0.2.2), and

$$\theta : \widehat{\bigoplus_r S_r} \longrightarrow \omega_\infty$$

is induced by the maps

$$S_r \xrightarrow{\sim} \omega'_r \hookrightarrow \omega_\infty.$$

It follows that a sufficient condition (but not a necessary one, as seen below) for obtaining a section

$$\sigma : \omega \longrightarrow I$$

of the desired kind is that  $\theta$  be an isomorphism. By duality (cf. 0.2.2), this is equivalent to saying that the linear map

$$\omega_\infty^\dagger \longrightarrow \prod_r S_r^\dagger$$

is bijective.

**5.2.1.B.** As above, put

$$\omega_\infty = \overline{\bigcup_{n \in \mathbb{N}} \omega_n}.$$

A second case in which a section

$$\sigma : \omega \longrightarrow I$$

of the desired kind exists is the case in which  $\omega_\infty$  has a pseudobasis  $\mathcal{B}_\infty$  that is the union, for  $n \in \mathbb{N}^*$ , of pseudobases of the spaces

$$\omega_n / \omega_{n-1}.$$

One can then complete it by a pseudobasis  $\mathcal{B}'_\infty$  of  $\omega / \omega_\infty$ , obtaining a pseudobasis of  $\omega$  compatible with the filtration.

Put

$$V = \omega_\infty^\dagger,$$

and let  $V_n$  be the orthogonal complement of  $\omega_n$  in  $V$ . The preceding condition is equivalent to saying that, in the category of ordinary  $k$ -vector spaces, the separated decreasing filtration

$$V = V_0 \supset V_1 \supset V_2 \supset \cdots$$

splits: namely,  $V$  is the direct sum, over  $n \in \mathbb{N}$ , of subspaces  $F_n$  such that

$$F_n \simeq V_n / V_{n+1}.$$

This need not hold. For example, let

$$V = \mathcal{S} = k^{\mathbb{N}}$$

be the space of sequences of elements of  $k$ , and let  $\mathcal{S}_n$  be the subspace of sequences  $(u_i)$  such that  $u_i = 0$  for  $i < n$ . Then

$$\dim \mathcal{S}_n / \mathcal{S}_{n+1} = 1,$$

but  $\mathcal{S}$  is not isomorphic to the direct sum of the  $\mathcal{S}_n / \mathcal{S}_{n+1}$ , because  $\mathcal{S}$  is not countable-dimensional. In this example, however,  $\mathcal{S}$  is the product of the  $\mathcal{S}_n / \mathcal{S}_{n+1}$  (cf. 5.2.1.A).

The splitting does hold when  $V$  is countable-dimensional.<sup>159</sup> Indeed, let

$$(e_n)_{n \in \mathbb{N}}$$

be a basis of  $V$ . We shall construct inductively an increasing function

$$g : \mathbb{N} \longrightarrow \mathbb{N}$$

and subspaces  $F_i$ , for  $i = 0, \dots, g(n)$ , such that

$$F_i \simeq V_i / V_{i+1}$$

and

$$F_{\leq g(n)} = \bigoplus_{i=0}^{g(n)} F_i$$

---

<sup>159</sup>This paragraph arose from discussions with J.-M. Fontaine and E. Bouscaren; in particular, Bouscaren indicated the proof that follows.

is a complement of  $V_{g(n)+1}$  containing

$$e_0, \dots, e_n.$$

It will then follow that

$$V = \bigoplus_{i \geq 0} F_i.$$

Let  $n+1 \in \mathbb{N}$ , and assume the assertion already proved for  $n$  (it is vacuous for  $n = -1$ ). If

$$e_{n+1} \in F_{\leq g(n)},$$

put  $g(n+1) = g(n)$ . Otherwise write

$$e_{n+1} = f + x, \quad f \in F_{\leq g(n)}, \quad 0 \neq x \in V_{g(n)+1}.$$

Let  $j$  be the least integer such that

$$x \in V_j \setminus V_{j+1}.$$

For

$$i = g(n) + 1, \dots, j,$$

choose a complement  $F_i$  of  $V_{i+1}$  in  $V_i$ , choosing  $F_j$  so that  $x \in F_j$ . Then

$$e_{n+1} \in F_{\leq j};$$

put  $g(n+1) = j$ .

*Source note.* At the last induction step, the French PDF prints  $e_{n+1} \in F_j$ . From  $e_{n+1} = f + x$ , with  $f \in F_{\leq g(n)}$ , the argument instead gives  $x \in F_j$  and  $e_{n+1} \in F_{\leq j}$ ; these are the relations used here.

**5.2.1.C.** <sup>160</sup> In particular, the two preceding conditions (5.2.1.A and 5.2.1.B) hold when the filtration of  $\omega$  is stationary, that is, when there exists an integer  $n_0$  such that

$$\omega_n = \omega_{n_0} \quad \text{for } n_0 \leq n < +\infty.$$

In this case,

$$k[[\omega]] / ((x^{p^r}))_{x \in \omega_r}$$

is isomorphic to the completed tensor product

$$\begin{aligned} & \frac{k[[\omega_1]]}{((x^p))_{x \in \omega_1}} \widehat{\otimes}_k \frac{k[[\omega'_2]]}{((x^{p^2}))_{x \in \omega'_2}} \widehat{\otimes}_k \cdots \\ & \widehat{\otimes}_k \frac{k[[\omega'_{n_0}]]}{((x^{p^{n_0}}))_{x \in \omega'_{n_0}}} \widehat{\otimes}_k k[[\omega'_\infty]], \end{aligned}$$

where  $\omega'_n$  is a complement of  $\omega_{n-1}$  in  $\omega_n$ , and  $\omega'_\infty$  is a complement of

$$\omega_\infty = \omega_{n_0}$$

in  $\omega$ .

The filtration of  $\omega$  is plainly stationary in case (i), where  $\omega_r = \omega$  for all sufficiently large  $r$ ; in case (ii), where  $\omega$  is finite-dimensional; and also in case (iii), where

$$I_r = 0$$

for every  $r$ . In the last case,  $B$  is isomorphic to the formal power series algebra  $k[[\omega]]$ . The preceding remarks therefore imply the theorem, subject to 5.2.2–5.2.5 below.

---

<sup>160</sup>Here the editors return to the original, shortened to take account of the preceding additions.

**5.2.2**

Suppose first that  $B$  has height at most 1; that is,

$$x^p = 0 \quad \text{for every } x \in I.$$

By 5.1.4, the graded algebra

$$\mathrm{Gr}_I(B)$$

associated with the filtration

$$I \supset I^2 \supset I^3 \supset \dots$$

of  $B$  is endowed with a cogroup structure in the category  $\mathbf{Alpg}/k$ . In other words, the profinite algebra

$$C = \prod_{n \geq 0} \mathrm{Gr}_I(B)_n$$

is the affine algebra of a formal  $k$ -group  $N$ . Clearly,

$$\omega_{N/k} = I/I^2,$$

and  $N$  is infinitesimal of height at most 1.

By 4.4, the identity map of  $\omega_{N/k}$  therefore induces an isomorphism

$$B' = k[[\omega_{N/k}]] / ((x^p))_{x \in \omega_{N/k}} \xrightarrow{\sim} C.$$

In particular, the map  $\phi$  of 5.2.1 induces an isomorphism between the associated graded algebras of  $B'$  and  $B$ , when both are filtered by the powers of their augmentation ideals. It follows from [CA], §V.7, Lemma 1 (see also 5.1.5) that  $\phi$  is an isomorphism.

**5.2.3**

Suppose now that  $B$  has finite height at most  $r$ . Let

$$\pi : k \xrightarrow{\sim} k, \quad x \mapsto x^p,$$

be the Frobenius automorphism. The linear map

$$B \widehat{\otimes}_{\pi} k \longrightarrow B, \quad b \widehat{\otimes}_{\pi} x \mapsto b^p x = (bx^{1/p})^p,$$

has closed image equal to the closed subalgebra

$$B^p = \{ b^p \mid b \in B \}$$

of  $B$ . Put

$$J = B^p \cap I = B^p \cap I_A.$$

<sup>161</sup> Let  $G_1$  be the kernel of the Frobenius morphism

$$\mathrm{Fr} : G \longrightarrow G^{(p)},$$

and let  $HG_1$  be the formal subgroup of  $G$  that is the inverse image of the formal subgroup  $H^{(p)} \subset G^{(p)}$ . Then  $HG_1$  is defined by the closed ideal generated by the  $p$ -th powers of the

---

<sup>161</sup>The editors added what follows.

elements of  $AJ$ ; this ideal is  $AJ$ . Moreover, the formation of  $G/H$  commutes with base change, because  $G/H$  represents the quotient sheaf for the flat topology (cf. 2.4). Hence

$$(G/H)^{(p)} = G^{(p)}/H^{(p)},$$

and we have the following commutative diagrams:

$$\begin{array}{ccc} G & \xrightarrow{\text{Fr}} & G^{(p)} \\ \downarrow & & \downarrow \\ G/H & \xrightarrow{\text{Fr}} & G^{(p)}/H^{(p)} \end{array} \quad \begin{array}{ccc} A & \xleftarrow{a \otimes 1 \mapsto a^p} & A \hat{\otimes}_{\pi} k \\ \uparrow & & \uparrow \\ B & \xleftarrow{b \otimes 1 \mapsto b^p} & B \hat{\otimes}_{\pi} k \end{array}$$

It follows that  $B^p$  is the affine algebra of the quotient  $G/HG_1$ .<sup>162</sup> Temporarily denote by  $C$  the affine algebra of the quotient  $HG_1/H$ . Since the formation of  $G/H$  commutes with base change, the square

$$\begin{array}{ccc} HG_1 & \longrightarrow & G \\ \downarrow & & \downarrow \\ HG_1/H & \longrightarrow & G/H \end{array}$$

is Cartesian. Consequently,

$$A \hat{\otimes}_B C \simeq A/AJ = A \hat{\otimes}_B (B/BJ).$$

Since  $A$  is topologically free over  $B$  (by 2.4, because  $A$  and  $B$  are local), the natural morphism

$$B/BJ \longrightarrow C$$

is an isomorphism. Thus  $B/BJ$  is the affine algebra of  $HG_1/H$ , and of course

$$B/BJ = B_1$$

has height at most 1, since

$$J = ((x^p))_{x \in \mathfrak{t}(B)}.$$

Let

$$B' = k[[\omega]] / ((x^{p^r}))_{x \in \omega_r},$$

let  $\phi : B' \rightarrow B$  be the morphism introduced in 5.2.1, let

$$B'^p = \{x^p \mid x \in B'\},$$

and let  $J'$  be the augmentation ideal of  $B'^p$ . We then have the commutative diagram

$$\begin{array}{ccc} B' & \xrightarrow{\psi} & B \\ \downarrow & & \downarrow \\ B'_1 = B'/\overline{B'J'} & \xrightarrow{\phi_1} & B_1 = B/\overline{BJ} \end{array}$$

<sup>162</sup>As indicated in the original, this also follows from Proposition 5.1. The editors preferred to give the preceding argument, which does not use the implication (i)  $\Rightarrow$  (ii) of that proposition.

and, by 5.2.2,  $\phi_1$  is an isomorphism.

On the other hand, by 2.4,  $A$  is topologically flat over

$$B = \mathcal{A}(G/H) \quad \text{and over} \quad B^p = \mathcal{A}(G/HG_1).$$

It follows from 1.3.3 that  $B$  is topologically flat over  $B^p$ . Moreover, 5.2.4 below shows that the morphism

$$B'^p \longrightarrow B^p$$

induced by  $\phi$  is an isomorphism. We may therefore apply 0.3.4 to the pseudocompact ring  $B'^p = B^p$  and the pseudocompact  $B^p$ -modules  $M = B'$ ,  $N = B$ . By what precedes,

$$\phi_1 = \phi \hat{\otimes}_{B^p} k$$

is an isomorphism, and it follows that  $\phi$  itself is an isomorphism. This proves 5.2 when  $B$  has finite height, subject only to 5.2.4 below.

### 5.2.4

For every pseudocompact vector space  $V$  over  $k$ , denote by

$${}_{\pi}V = V \hat{\otimes}_{\pi} k$$

the space obtained from  $V$  by extension of scalars along  $x \mapsto x^p$ .<sup>163</sup> We then have a commutative diagram with exact rows:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & {}_{\pi}\overline{I^2} & \xrightarrow{\alpha} & {}_{\pi}I & \xrightarrow{\beta} & {}_{\pi}\omega & \longrightarrow & 0 \\ & & \downarrow u & & \downarrow v & & \downarrow w & & \\ 0 & \longrightarrow & \overline{J^2} & \xrightarrow{\gamma} & J & \xrightarrow{\delta} & \overline{\omega} & \longrightarrow & 0, \end{array}$$

Here

$$\overline{\omega} = J/J^2,$$

and the maps  $u, v, w$  are induced by the linear map

$${}_{\pi}B \longrightarrow B^p, \quad x \hat{\otimes} a \longmapsto x^p a.$$

Since  $u$  and  $v$  are surjective, so is  $w$ , and its kernel is the image  ${}_{\pi}\omega_1$  of

$${}_{\pi}I_1 = \ker(v).$$

Put

$$J_n = \{x \in J \mid x^{p^n} = 0\}, \quad \overline{\omega}_n = \delta(J_n).$$

Then, for every  $n \geq 0$ ,

$$J_n = v({}_{\pi}I_{n+1}), \quad \overline{\omega}_n = w({}_{\pi}\omega_{n+1}).$$

*Source note.* The French PDF prints  $\overline{\omega}_n = \delta(I_n)$ , although the displayed exact row defines  $\delta$  on  $J$ , and the immediately preceding definition introduces  $J_n \subset J$ . The type-correct  $\delta(J_n)$  is used above; all subsequent identities agree with this reading.

<sup>163</sup>Since  $k$  is perfect, one may identify  ${}_{\pi}V$  with the abelian group  $V$  on which  $k$  acts by  $\lambda \cdot v = \lambda^{1/p}v$ .

The section

$$\pi\sigma : \pi\omega \longrightarrow \pi I$$

induced by the section  $\sigma$  of 5.2.1 therefore descends to a section

$$\tau : \bar{\omega} \longrightarrow J$$

compatible with the filtrations of  $J$  and  $\bar{\omega}$ . Since  $B^p$  has height at most  $r-1$ , the induction hypothesis shows that this section induces an isomorphism

$$\psi : B'' = k[[\bar{\omega}]] / ((x^{p^n}))_{x \in \bar{\omega}_n} \xrightarrow{\sim} B^p.$$

But  $B''$  identifies with  $B'^p$ , and  $\psi$  identifies with the morphism  $B'^p \rightarrow B^p$  induced by  $\phi$ . Thus the theorem is proved when  $B$  has finite height.

**Remark 5.2.4.A.** <sup>164</sup> Suppose that  $B$  has height at most  $r+1$ , with  $r \in \mathbb{N}^*$ . Then  $I_{r+1} = I$ , and there is an isomorphism

$$B \simeq \frac{k[[S_1]]}{((x_1^p))_{x_1 \in S_1}} \hat{\otimes}_k \cdots \hat{\otimes}_k \frac{k[[S_r]]}{((x_r^p))_{x_r \in S_r}} \hat{\otimes}_k \frac{k[[S_{r+1}]]}{((x_{r+1}^{p^{r+1}}))_{x_{r+1} \in S_{r+1}}}, \quad (1)$$

where each  $S_n$  is a complement of

$$I^2 + I_{n-1}$$

in  $I^2 + I_n$ . Thus  $\omega = I/I^2$  identifies with

$$\prod_{i=1}^{r+1} S_i,$$

and it is immediate that, for  $n = 1, \dots, r+1$ , the image  $\omega_n$  of  $I_n$  in  $\omega$  identifies with

$$\prod_{i=1}^n S_i.$$

This has the following consequence. Let

$$B_r = B/\check{J}_r, \quad \check{J}_r = ((x^{p^r}))_{x \in \mathfrak{t}(B)},$$

and let

$$\mathfrak{m} = I/\check{J}_r$$

be the augmentation ideal of  $B_r$ . Since  $\check{J}_r \subset I^2$ , the space

$$\omega(r) = \mathfrak{m}/\mathfrak{m}^2$$

identifies with  $\omega$ . For  $n = 1, \dots, r$ , denote by  $\omega(r)_n$  the image in  $\omega(r)$  of

$$\mathfrak{m}_n = \{y \in \mathfrak{m} \mid y^{p^n} = 0\}.$$

It is also the image in  $\omega$  of

$$\{x \in I \mid x^{p^n} \in \check{J}_r\};$$

hence  $\omega(r)_n$  contains  $\omega_n$ .

<sup>164</sup>The editors added this remark, which is used in 5.2.5.



On the other hand, isomorphism (1) gives

$$\check{J}_r = ((x_{r+1}^{p^r}))_{x_{r+1} \in S_{r+1}},$$

and therefore

$$B_r \simeq \frac{k[[S_1]]}{((x_1^p))_{x_1 \in S_1}} \widehat{\otimes}_k \cdots \widehat{\otimes}_k \frac{k[[S_r]]}{((x_r^{p^r}))_{x_r \in S_r}} \widehat{\otimes}_k \frac{k[[S_{r+1}]]}{((x_{r+1}^{p^r}))_{x_{r+1} \in S_{r+1}}}. \quad (2)$$

It follows that  $\omega(r)_n$  identifies with  $\prod_{i=1}^n S_i$  for  $n = 1, \dots, r-1$ . Thus the inclusion

$$\omega_n \subset \omega(r)_n$$

is an equality for  $n = 1, \dots, r-1$ .

### 5.2.5

It remains to consider the case in which  $B$  has infinite height and the projection  $I \rightarrow \omega$  admits a section  $\sigma$  compatible with the filtrations of  $\omega$  and  $I$ . Consider the morphism induced by  $\sigma$ :

$$\phi : B' = k[[\omega]] / ((x^{p^n}))_{x \in \omega_n} \longrightarrow B.$$

It is enough to prove that, for every  $r \in \overline{\mathbb{N}}$ , the map

$$\phi_r : B'_r \longrightarrow B_r$$

induced by  $\phi$  is an isomorphism.<sup>165</sup>

For every  $r \in \mathbb{N}^*$ , let  $G_r$  be the kernel of the iterated Frobenius morphism

$$G \longrightarrow G^{(p^r)},$$

and let  $HG_r$  be the formal subgroup of  $G$  that is the inverse image of  $H^{(p^r)} \subset G^{(p^r)}$ . Thus  $HG_r$  is defined by the closed ideal generated by the  $p^r$ -th powers of the elements of  $AI$ , namely

$$A\check{J}_r, \quad \check{J}_r = \{x^{p^r} \mid x \in I\}.$$

Exactly as in 5.2.3, it follows that

$$B_r = B/B\check{J}_r$$

is the affine algebra of  $HG_r/H$ ; in particular, it has height at most  $r$ .

Let

$$\mathfrak{m}(r) = I/B\check{J}_r$$

be the augmentation ideal of  $B_r$ . The canonical projection  $B \rightarrow B_r$  induces an isomorphism

$$\omega = I/I^2 \xrightarrow{\sim} \omega(r) = \mathfrak{m}(r)/\mathfrak{m}(r)^2,$$

which we use to identify the two spaces. Let  $\omega(r)_n$  be the image in  $\omega(r)$  of the closed ideal

$$\mathfrak{m}(r)_n = \{y \in \mathfrak{m}(r) \mid y^{p^n} = 0\}.$$

It is also the image in  $\omega$  of the closed ideal

$$I(r)_n = \{x \in I \mid x^{p^n} \in B\check{J}_r\}.$$

<sup>165</sup>Indeed,  $B'$ , respectively  $B$ , is the inverse limit of the  $B'_r$ , respectively the  $B_r$ . Moreover, the editors modified the original in what follows in order to take account of the addition made in 5.2.3.

166

Clearly,

$$\omega(r)_n = \omega \quad \text{if } n \geq r.$$

We shall prove that

$$\omega(r)_n = \omega_n \quad \text{if } n < r.$$

For every  $r, n$ , the sequence

$$0 \longrightarrow I(r)_n \cap I^2 \longrightarrow I(r)_n \longrightarrow \omega(r)_n \longrightarrow 0$$

is exact. For fixed  $n$ ,

$$\bigcap_r I(r)_n = I_n,$$

because  $\bigcap_r B\check{J}_r = 0$ . Filtered inverse limits are exact in  $\text{PC}(k)$  (cf. 0.2), so for every  $n$ ,

$$\omega_n = \bigcap_r \omega(r)_n. \quad (*)$$

On the other hand, Remark 5.2.4.A gives

$$\omega(r)_n = \omega(r+1)_n \quad \text{if } n < r.$$

Together with (\*), this implies

$$\omega(r)_n = \omega_n \quad \text{if } n < r.$$

Consequently, the vector space  $\omega$ , filtered by the subspaces  $(\omega(r)_n)_{n \geq 0}$ , is precisely  ${}^r\omega$  in the notation of 5.2. The composite

$$\sigma(r) : \omega \xrightarrow{\sigma} I \longrightarrow \mathfrak{m}(r)$$

is therefore compatible with the filtrations  $(\omega(r)_n)$  and  $(\mathfrak{m}(r)_n)$ . Since

$$k[[{}^r\omega]] / ((x^{p^n}))_{x \in {}^r\omega_n}$$

is exactly  $B'_r$ , and  $\phi_r : B'_r \rightarrow B_r$  is induced by  $\sigma(r)$ , the result already proved for finite-height algebras shows that  $\phi_r$  is an isomorphism.

### 5.2.6

**Definition 5.2.6.** <sup>167</sup> Let  $(A_\lambda)_{\lambda \in \Lambda}$  be a family of profinite  $k$ -algebras, each endowed with an augmentation

$$\varepsilon_\lambda : A_\lambda \longrightarrow k.$$

(This holds in particular if every  $A_\lambda$  is local with residue field  $k$ .) Define the infinite completed tensor product

$$\mathcal{A} = \widehat{\bigotimes_{\lambda \in \Lambda} A_\lambda}^c$$

to be the inverse limit in  $\text{Alp}/k$  of

$$\mathcal{A}_F = \widehat{\bigotimes_{\lambda \in F} A_\lambda}^c,$$

<sup>166</sup>The editors expanded the original in what follows.

<sup>167</sup>The editors added this number in order to define the infinite tensor products used in 5.3(a).

where  $F$  ranges through the finite subsets of  $\Lambda$ . For  $F' = F \cup \{\lambda\}$ , the transition morphism

$$\mathcal{A}_{F'} \longrightarrow \mathcal{A}_F$$

is

$$\mathrm{id} \hat{\otimes} \varepsilon_\lambda.$$

In particular, suppose  $\Lambda = \mathbb{N}^*$ , and write

$$X_n = \mathrm{Spf}(A_n).$$

Then  $\mathrm{Spf}(\mathcal{A})$  represents the functor that assigns to every  $C \in \mathrm{Alf}/k$  the set of *finite* sequences in  $\prod_{n \geq 1} X_n(C)$ ; that is, the sequences

$$(x_1, x_2, \dots) \in \prod_{n \geq 1} X_n(C)$$

such that

$$x_n = \varepsilon_n$$

for all sufficiently large  $n$ . Here, by abuse of notation,  $\varepsilon_n$  denotes the composite of

$$\varepsilon_n : A_n \longrightarrow k$$

with the structural morphism  $k \rightarrow C$ .

If, in addition, every  $A_n$  is a quotient of an algebra  $k[[\omega'_n]]$ , one may write 0 for the unique morphism

$$A_n \longrightarrow C$$

that vanishes on  $\omega'_n$ . The preceding description then becomes the set of sequences for which  $x_n = 0$  for all sufficiently large  $n$ .

### 5.3. Remarks

(a) Call a profinite  $k$ -algebra *stationary* if it is the completed tensor product of a family of truncated formal power series algebras.<sup>168</sup>

Let  $G$  be an infinitesimal formal  $k$ -group and let  $B$  be the affine algebra of a homogeneous space of  $G$ . It follows from Theorem 5.2 that, for every  $r \in \mathbb{N}$ , the algebra

$$B / ((x^{p^r}))_{x \in \mathfrak{r}(B)}$$

is stationary. In particular,  $B$  is an inverse limit of stationary algebras.<sup>169</sup>

<sup>168</sup>The author doubtless had in mind a completed tensor product

$$\mathcal{A} = \widehat{\bigotimes_{n \in \mathbb{N}^*}}^c \frac{k[[\omega'_n]]}{((x^{p^n}))_{x \in \omega'_n}},$$

where the  $\omega'_n$  are arbitrary pseudocompact vector spaces. Then  $\omega = \omega_{\mathcal{A}/k}$  identifies with  $\prod_n \omega'_n$ , and the filtration is  $\omega_n = \prod_{i=1}^n \omega'_i$ ; this is the case of 5.2.1.B. For this reason these algebras would be better called *stable*, rather than *stationary*; cf. [Di73], II, §2.9, p. 75. If, for example,  $\dim \omega'_n = 1$  for every  $n \in \mathbb{N}^*$ , then  $\mathcal{A}$  represents the functor that assigns to  $C \in \mathrm{Alf}/k$  the sequences  $(x_n)_{n \in \mathbb{N}^*}$  of elements of  $C$  such that  $x_n^{p^n} = 0$  and  $x_n = 0$  for all sufficiently large  $n$ . This case, where  $\omega = \prod_{n \in \mathbb{N}^*} \omega'_n$ , corresponds in the dual setting of connected cocommutative Hopf algebras to the case studied by M. E. Sweedler; cf. [Sw67], Theorem 3.

<sup>169</sup>Such an inverse limit need not itself be a stable profinite  $k$ -algebra in the sense of the preceding editor's note. For example, let  $S$  be the ordinary  $k$ -vector space of sequences  $(u_1, u_2, \dots)$  of elements of  $k$ , and put  $\omega = S^*$ . Then  $\omega$  is the direct sum in  $\mathrm{PC}(k)$  of copies  $kx_n$  of  $k$ , for  $n \in \mathbb{N}^*$ , where  $x_n(u) = u_n$ . Thus this is the case of 5.2.1.A. The algebra

$$\mathcal{A} = k[[\omega]] / ((x_n^{p^n}))_{n \in \mathbb{N}^*}$$

has  $\omega_{\mathcal{A}/k} = \omega$  and  $\omega_n = \prod_{i=1}^n kx_i$ , but it is not stable:  $\mathrm{Spf}(\mathcal{A})$  represents the functor that assigns to every  $C \in \mathrm{Alf}/k$  the infinite sequences  $(x_n)_{n \in \mathbb{N}^*}$  of elements of  $C$  such that  $x_n^{p^n} = 0$  for every  $n$ .

(b) I do not know whether, with the notation of 5.2.1,  $A$  and  $B$  can be chosen so that there exists no section

$$\sigma : \omega \longrightarrow I$$

compatible with the filtrations.<sup>170</sup> Nevertheless,  $\omega$  may be any pseudocompact vector space filtered by an increasing sequence of closed subspaces. Indeed, given

$$\omega_1 \subset \omega_2 \subset \cdots \subset \omega,$$

one can define on

$$B = A = k[[\omega]] / ((x^{p^r}))_{x \in \omega_r}$$

a diagonal morphism

$$\Delta_A : A \longrightarrow A \hat{\otimes}_k A$$

satisfying conditions (i), (ii), and (iii) of 2.1. It suffices to put

$$\Delta_A(y) = y \hat{\otimes} 1 + 1 \hat{\otimes} y$$

when  $y$  is the image in  $A$  of an element of  $\omega$ .

#### 5.4. Corollary

**Corollary 5.4.** *Let  $G$  be an algebraic group over a perfect field  $k$  of characteristic  $p > 0$ , let  $H$  be an algebraic subgroup of  $G$ , let  $e$  be the image of the identity element of  $G$  in  $G/H$ , and let  $A$  be the local ring of  $G/H$  at  $e$ . Then  $\hat{A}$  is isomorphic to an algebra of the form*

$$k[[X_1, \dots, X_r, \dots, X_s]] / (X_1^{p^{n_1}}, \dots, X_r^{p^{n_r}}).$$

*Proof.* Consider the infinitesimal formal groups

$$\hat{G} = \mathrm{Spf}(\hat{\mathcal{O}}_{G,e}), \quad \hat{H} = \mathrm{Spf}(\hat{\mathcal{O}}_{H,e}).$$

By 1.3.4, the completion  $\hat{A}$  of

$$A = \mathcal{O}_{G/H,e}$$

is isomorphic to

$$\mathcal{A}(\hat{G}/\hat{H}),$$

and the corollary follows from Theorem 5.2(ii).<sup>171</sup>

#### 5.5. Complements

<sup>172</sup>

Recall the following definitions. A Noetherian local ring  $A$  is called a *complete intersection* if its completion  $\hat{A}$  is the quotient of a complete regular Noetherian local ring  $B$  by an ideal  $I$  generated by a regular sequence of elements of  $B$  (cf. EGA IV<sub>4</sub>, 19.3.1).

On the other hand, let

$$\tau : Y \hookrightarrow X$$

<sup>170</sup>The editors do not know either, outside the cases considered in 5.2.1.A and 5.2.1.B.

<sup>171</sup>See also [DG70], §III.3, Theorem 6.1.

<sup>172</sup>The editors added this subsection in order to give some consequences of 5.4 mentioned in Exposés III and VIA.

be a closed immersion of schemes. If  $y \in Y$ , then  $\tau$  is called a *regular immersion at the point  $y$*  if the kernel of

$$\mathcal{O}_{X,y} \longrightarrow \mathcal{O}_{Y,y}$$

is generated by a regular sequence. If, moreover,  $X$  is locally Noetherian and  $\tau$  is a regular immersion at every point, then  $\tau$  is called a *regular immersion* (cf. *loc. cit.*, Proposition 16.9.10 and Definition 16.9.2).

**Corollary 5.5.1.** *If  $G$  is an algebraic group over a field  $k$ , then the local ring  $\mathcal{O}_{G,e}$  is a complete intersection.*

*Proof.* By EGA IV<sub>4</sub>, 19.3.4, we may assume that  $k$  is algebraically closed. If  $\text{char}(k) = 0$ , then  $G$  is already known to be smooth (cf. 3.3.1 or VIB, 1.6.1); hence  $\hat{\mathcal{O}}_{G,e}$  is a  $k$ -algebra of formal power series, by EGA IV<sub>4</sub>, 17.5.3 (d''). If  $\text{char}(k) = p > 0$ , then 5.4, applied with  $H = \{e\}$ , shows that  $\hat{\mathcal{O}}_{G,e}$  is a complete intersection.

**Remarks 5.5.2.** *Let  $k$  be a field, let  $G$  be a smooth algebraic  $k$ -group, and let  $H$  be a closed subgroup of  $G$ .*

(a) We saw in Exposé III, 4.15, that the immersion

$$H \hookrightarrow G$$

is regular. This can also be deduced from 5.4 as follows. As in *loc. cit.*, we may assume that  $k$  is algebraically closed, and it is enough to show that the kernel  $I$  of

$$\mathcal{O}_{G,e} \longrightarrow \mathcal{O}_{H,e}$$

is generated by a regular sequence. Put

$$A = \mathcal{O}_{G,e}, \quad \hat{G} = \text{Spf}(\hat{A}), \quad \hat{H} = \text{Spf}(\hat{\mathcal{O}}_{H,e}).$$

Since  $A$  is Noetherian, there is an exact sequence

$$0 \longrightarrow I \otimes_A \hat{A} \longrightarrow \hat{A} \xrightarrow{\pi} \mathcal{A}(\hat{H}) \longrightarrow 0,$$

and  $\hat{A}$  is faithfully flat over  $A$ . Therefore, by EGA IV<sub>4</sub>, 16.9.10(ii) and 19.1.5(ii), it is enough to show that the kernel

$$\hat{I} = I \otimes_A \hat{A}$$

of  $\pi$  is generated by a regular sequence of elements of  $\hat{A}$ .

Since  $G$  is smooth,  $\hat{A}$  is reduced. By 5.4, the subalgebra

$$B = \mathcal{A}(\hat{G}/\hat{H})$$

is therefore isomorphic to a formal power series algebra

$$k[[x_1, \dots, x_n]].$$

Consequently, the identity section of  $\hat{G}/\hat{H}$  is defined in  $B$  by the regular sequence  $(x_1, \dots, x_n)$ . Since  $\hat{A}$  is Noetherian, the ideal  $J$  of  $\hat{A}$  generated by  $x_1, \dots, x_n$  is closed, and hence equals  $\hat{I}$ , by Corollary 1.4. Moreover,  $\hat{A}$  is topologically flat, and hence flat over  $B$  (cf. 0.3.8). Thus  $(x_1, \dots, x_n)$  is a regular sequence in  $\hat{A}$ , by EGA IV<sub>4</sub>, 19.1.5(ii).

(b) The following more precise result can also be deduced from 5.2(ii). Suppose that  $k$  is perfect. Applying 5.2(ii) to the algebra

$$C = \mathcal{A}(H),$$

there are a basis

$$(y_1, \dots, y_{r+s})$$

of  $\omega_H$  and integers

$$1 \leq n_1 \leq \dots \leq n_r$$

such that  $\mathcal{A}(H)$  is isomorphic to the quotient of

$$k[[y_1, \dots, y_{r+s}]]$$

by the ideal generated by

$$y_i^{p^{n_i}}, \quad i = 1, \dots, r.$$

Lift the  $y_i$  to elements  $x_i$  of  $\omega_G$ , and extend  $(x_1, \dots, x_{r+s})$  to a basis  $(x_1, \dots, x_n)$  of  $\omega_G$ . Since  $\mathcal{A}(G)$  is reduced, the morphism

$$k[[x_1, \dots, x_n]] \longrightarrow \mathcal{A}(G)$$

is an isomorphism, by 5.2(iii).

Thus there exists a “coordinate system”  $(x_1, \dots, x_n)$  on  $G$ , that is, an isomorphism

$$\mathcal{A}(G) \simeq k[[x_1, \dots, x_n]],$$

such that  $H$  is defined by the equations

$$x_i^{p^{n_i}} = 0 \quad (i = 1, \dots, r), \quad x_i = 0 \quad (i > r + s).$$

This is “Dieudonné’s theorem”; compare [Di55], §19, Theorem 6, and [Di73], II, §3.2, Proposition 3 and the material preceding it.

# Bibliography

- [CA] P. Gabriel, *Des catégories abéliennes*, Bull. Soc. Math. France **90** (1962), 323–448.  
173
- [Ab80] E. Abe, *Hopf algebras*, Cambridge Univ. Press, 1980.
- [Br00] C. Breuil, *Groupes  $p$ -divisibles, groupes finis et modules filtrés*, Ann. of Math. **152** (2000), n° 2, 489–549.
- [BAlg] N. Bourbaki, *Algèbre*, Chap. I–III et IV–VII, Hermann, 1970, et Masson, 1981.
- [BAC] N. Bourbaki, *Algèbre commutative*, Chap. I–IV, Masson, 1985.
- [BEns] N. Bourbaki, *Théorie des ensembles*, Chap. I–IV, Hermann, 1970.
- [BLie] N. Bourbaki, *Groupes et algèbres de Lie*, Chap. II–III, Hermann, 1970.
- [Ca62] P. Cartier, *Groupes algébriques et groupes formels*, pp. 87–111 in: *Colloque sur la théorie des groupes algébriques* (Bruxelles, 1962), Gauthier-Villars, 1962.
- [DG70] M. Demazure, P. Gabriel, *Groupes algébriques*, Masson & North-Holland, 1970.
- [De72] M. Demazure, *Lectures on  $p$ -divisible groups*, Lect. Notes Math. **302**, Springer-Verlag, 1972.
- [Di55] J. Dieudonné, *Groupes de Lie et hyperalgèbres de Lie sur un corps de caractéristique  $p > 0$* , Comm. Math. Helv. **28** (1955), 87–118.
- [Di73] J. Dieudonné, *Introduction to the theory of formal groups*, Marcel Dekker, 1973.
- [Fo77] J.-M. Fontaine, *Groupes  $p$ -divisibles sur les corps locaux*, Astérisque **47–48**, Soc. Math. France, 1977.
- [Gr57] A. Grothendieck, *Sur quelques points d’algèbre homologique*, Tôhoku Math. J. **9** (1957), 119–221.
- [Gr74] A. Grothendieck, *Groupes de Barsotti-Tate et cristaux de Dieudonné*, Presses Univ. Montréal, 1974.
- [HS69] R. G. Heyneman, M. E. Sweedler, *Affine Hopf Algebras I*, J. Algebra **13** (1969), 194–241.
- [Il85] L. Illusie, *Déformations de groupes de Barsotti-Tate, d’après A. Grothendieck*, pp. 151–198 in: *Séminaire sur les pinceaux arithmétiques: la conjecture de Mordell*, Astérisque **127**, Soc. Math. France, 1985.

---

<sup>173</sup>The editors added to this reference, which appeared in the original, the references that follow.

- [Ja65] I. M. James, review of [MM65] in Math. Reviews, MR0174052.
- [La75] M. Lazard, *Commutative formal groups*, Lect. Notes Math. **443**, Springer-Verlag, 1975.
- [LT65] J. Lubin, J. Tate, *Formal complex multiplication in local fields*, Ann. of Math. **81** (1965), 380–387.
- [LT66] J. Lubin, J. Tate, *Formal moduli for one-parameter formal Lie groups*, Bull. Soc. Math. France **94** (1966), 49–60.
- [Ma91] A. Masuoka, *On Hopf algebras with cocommutative coradicals*, J. Algebra **144** (1991), 451–466.
- [Me72] W. Messing, *The crystals associated with Barsotti-Tate Groups: with applications to abelian schemes*, Lect. Notes Math. **264**, Springer-Verlag, 1972.
- [MM65] J. W. Milnor, J. C. Moore, *On the structure of Hopf algebras*, Ann. of Math. **81** (1965), 211–264.
- [Mi65] B. Mitchell, *Theory of categories*, Academic Press, 1965.
- [Ne75] K. Newman, *A correspondence between bi-ideals and sub-Hopf algebras in cocommutative Hopf algebras*, J. Algebra **36** (1975), 1–15.
- [Po73] N. Popescu, *Abelian categories with applications to rings and modules*, Academic Press, 1973.
- [Sch90] H. J. Schneider, *Principal homogeneous spaces for arbitrary Hopf algebras*, Israel J. Math. **72** (1990), n<sup>os</sup> 1–2, 167–195.
- [Sw67] M. Sweedler, *Hopf algebras with one grouplike element*, Trans. Amer. Math. Soc. **127** (1967), n<sup>o</sup> 3, 515–526.
- [Sw69] M. Sweedler, *Hopf algebras*, Benjamin, 1969.
- [Tak72] M. Takeuchi, *A correspondence between Hopf ideals and sub-Hopf algebras*, Manuscripta Math. **7** (1972), 251–270.
- [Tak79] M. Takeuchi, *Relative Hopf modules—Equivalence and freeness criteria*, J. Algebra **60** (1979), 452–471.
- [Ta67] J. Tate, *p-divisible groups*, pp. 158–183 in: *Proc. Conf. Local Fields* (Driebergen, 1966) (éd. T. A. Springer), Springer-Verlag, 1967.
- [We95] C. A. Weibel, *An introduction to homological algebra*, Cambridge Univ. Press, 1995.



# Exposé VIII

## Diagonalizable Groups

by A. Grothendieck<sup>0</sup>

### 1. Biduality

Let  $\mathcal{C}$  be a category, which as usual we identify with a full subcategory of

$$\widehat{\mathcal{C}} = \text{Hom}(\mathcal{C}^\circ, \mathbf{Set})$$

(cf. Exposé I). Let  $I$  be a commutative group functor, that is, an object of  $\widehat{\mathcal{C}}$  endowed with a commutative group structure (cf. I, 2.1).<sup>1</sup> For every  $X \in \text{Ob}(\widehat{\mathcal{C}})$ , the object  $\text{Hom}(X, I)$  carries the commutative group structure induced by that of  $I$ . For every group  $G$  in  $\widehat{\mathcal{C}}$ , put

$$D(G) = \text{Hom}_{\text{gr}}(G, I),$$

the subobject of  $\text{Hom}(G, I)$  defined, for every  $S \in \text{Ob}(\mathcal{C})$ , by

$$(x) \quad D(G)(S) = \text{Hom}_{S\text{-gr}}(G_S, I_S),$$

where  $G_S = G \times S$  and  $I_S = I \times S$  are regarded as  $S$ -groups, that is, as groups in  $\widehat{\mathcal{C}}/S$ . Thus  $D(G)$  is a  $\widehat{\mathcal{C}}$ -subgroup of  $\text{Hom}(G, I)$ . In this way we obtain a contravariant functor  $D$  from the category of  $\widehat{\mathcal{C}}$ -groups to the category of commutative  $\widehat{\mathcal{C}}$ -groups.

The right-hand side of (x) may also be interpreted as the subset of  $\text{Hom}(G \times S, I)$  consisting of morphisms  $G \times S \rightarrow I$  that are “multiplicative in the first argument  $G$ .” Moreover, the preceding formulas remain valid when  $S$  is an arbitrary object of  $\mathcal{C}$ , not necessarily one arising from  $\mathcal{C}$ .

Now take for  $S$  a group in  $\widehat{\mathcal{C}}$ , denoted  $G'$ . In the left-hand side  $\text{Hom}(G', D(G))$  of (x), consider the subset  $\text{Hom}_{\text{gr}}(G', D(G))$  consisting of morphisms that respect the group structures of  $G'$  and  $D(G)$ .

It corresponds to the subset of  $\text{Hom}(G \times G', I)$  consisting of morphisms that are multiplicative in each of the two arguments—which one may call *bilinear morphisms from  $G \times G'$  to  $I$* , or *pairings of  $G$  and  $G'$  with values in  $I$* . We therefore have

$$(xx) \quad \text{Hom}_{\text{gr}}(G', D(G)) \xrightarrow{\sim} \text{Hom}_{\text{bil}}(G \times G', I),$$

<sup>0</sup>Editor’s note: version 1.1 of November 8, 2009. Additions were made in 1.2, 1.4, 1.7, 3.1, 3.4, 4.5.1, 6.4, and 6.8; 1.5.1 and Section 7 remain to be revised.

<sup>1</sup>Editor’s note: the original has been expanded in what follows.

functorially in the pair  $(G, G')$ . Since the right-hand side is symmetric in  $G$  and  $G'$ , this gives a functorial bijection

$$(xxx) \quad \mathrm{Hom}_{\mathrm{gr}}(G', D(G)) \xrightarrow{\sim} \mathrm{Hom}_{\mathrm{gr}}(G, D(G')).$$

In other words, giving a group homomorphism  $G' \rightarrow D(G)$  “amounts to the same thing” as giving a group homomorphism  $G \rightarrow D(G')$ : both amount to giving a pairing  $G \times G' \rightarrow I$ .

Apply this with  $G' = D(G)$  and with the identity homomorphism  $G' \rightarrow D(G)$ . We obtain a canonical homomorphism

$$(xxxx) \quad G \longrightarrow D(D(G)).$$

**Definition 1.0.** <sup>2</sup> *The group  $G$  is called reflexive (relative to  $I$ ) if the preceding homomorphism is an isomorphism. This implies, in particular, that  $G$  is commutative.*

We therefore have:

**Proposition 1.0.1.** *The functor  $D$  induces an anti-equivalence from the category of reflexive  $\mathcal{C}$ -groups to itself.*

In particular, if  $G$  and  $H$  are reflexive groups,  $D$  induces an isomorphism

$$\mathrm{Hom}_{\mathrm{gr}}(G, H) \xrightarrow{\sim} \mathrm{Hom}_{\mathrm{gr}}(D(H), D(G)).$$

It is enough, in fact, that  $H$  be reflexive, as follows from (xxx).

**Definition 1.0.2.** *As usual, a  $\mathcal{C}$ -group  $G$  is called reflexive if it is reflexive as a  $\widehat{\mathcal{C}}$ -group, regardless of whether  $D(G)$  is representable.*

Thus  $D$  gives an anti-equivalence from the category of reflexive  $\mathcal{C}$ -groups  $G$  for which  $D(G)$  is representable to itself.

**Remark 1.0.3.** To conclude these generalities, note that formation of the dual  $D(G)$  commutes with base extension, which therefore carries reflexive groups to reflexive groups.

We shall henceforth work in the following setting. Let  $\mathcal{C} = (\mathbf{Sch})_S$  be the category of preschemes over  $S$ , and let  $I = \mathbf{G}_{m,S}$  be the “multiplicative group over  $S$ ” (cf. Exposé I). For every ordinary group  $M$ , consider the constant  $S$ -group  $M_S$ . For every group prescheme  $J$  over  $S$ , there is a canonical isomorphism, functorial in  $M$  and  $J$  and compatible with base extension,

$$\mathrm{Hom}_{S\text{-gr}}(M_S, J) = \mathrm{Hom}_{\mathrm{gr}}(M, J(S)).$$

Applying this with  $J = I = \mathbf{G}_{m,S}$  and with a variable  $S'$  over  $S$ , we obtain a functorial isomorphism

$$(1.0.4) \quad D(M_S)(S') \xrightarrow{\sim} \mathrm{Hom}_{\mathrm{gr}}(M, \mathbf{G}_m(S')).$$

We thus recover the functor already considered in I, 4.4, also denoted  $D_S(M)$ , which is representable when  $M$  is commutative, since

$$D_S(M) = D(M_S) = \mathrm{Spec} \mathcal{O}_S(M),$$

where  $\mathcal{O}_S(M)$  denotes the group algebra of  $M$  over  $\mathcal{O}_S$ . In general, moreover,  $D(M)$  is unchanged when  $M$  is replaced by its abelianization, so no information is lost by assuming that  $M$  is commutative.

---

<sup>2</sup>Editor’s note: the numbering 1.0, 1.0.1, and so on has been added to make the definitions and statements contained here explicit.

**Definition 1.1.** A group prescheme  $G$  over  $S$  is called diagonalizable if it is isomorphic to a scheme of the form

$$D_S(M) = D(M_S) = \operatorname{Hom}_{S\text{-gr}}(M_S, \mathbf{G}_m)$$

for a suitable commutative group  $M$ . It is called locally diagonalizable if every point of  $S$  has an open neighborhood  $U$  such that  $G|_U$  is diagonalizable.

**Theorem 1.2.** Let  $\Gamma$  be a constant commutative group scheme over  $S$ ; that is, suppose that it is isomorphic to a group scheme of the form  $M_S$ , where  $M$  is an ordinary commutative group. Then  $\Gamma$  is reflexive: the canonical homomorphism

$$\Gamma \longrightarrow D(D(\Gamma))$$

is an isomorphism.<sup>3</sup> Hence the diagonalizable group  $D(M_S)$  is also reflexive.

In view of the definitions, this follows from the next statement, applied to a prescheme  $S'$  over  $S$ .

**Corollary 1.3.** Let  $G = D(M_S)$ . Every homomorphism of  $S$ -groups

$$u : G \longrightarrow \mathbf{G}_{m,S}$$

is defined by a unique section of  $M_S$  over  $S$ , equivalently by a unique locally constant map from  $S$  to  $M$ .

*Proof.* By definition,

$$\mathbf{G}_{m,S} = \operatorname{GL}(1)_S = \operatorname{Aut}_{\mathcal{O}_S\text{-mod}}(\mathcal{O}_S).$$

Thus giving a group homomorphism  $G \rightarrow \mathbf{G}_{m,S}$  is equivalent to giving  $\mathcal{O}_S$  the structure of a  $G$ - $\mathcal{O}_S$ -module compatible with its natural  $\mathcal{O}_S$ -module structure (cf. I, 4.7). By I, 4.7.3, this is also equivalent to giving an  $M$ -grading on  $\mathcal{O}_S$ , that is, a direct sum decomposition

$$\mathcal{O}_S = \bigoplus_{m \in M} \mathcal{L}_m.$$

It is well known that a direct summand of a locally free module of finite type is locally free of finite type. Hence, near each point of  $S$ , every  $\mathcal{L}_m$  is either zero or free of rank one; in the latter case it identifies with  $\mathcal{O}_S$  on that neighborhood. Let  $S_m$  be the open subset of  $S$  on which the latter alternative occurs. The direct-sum decomposition shows that the  $S_m$  cover  $S$  and are pairwise disjoint.

Consequently, giving a group homomorphism  $G \rightarrow \mathbf{G}_{m,S}$  is equivalent to giving a decomposition of  $S$  as a union of pairwise disjoint open subsets  $S_m$ ,  $m \in M$ , hence to giving a locally constant map from  $S$  to  $M$ . This proves Corollary 1.3 and therefore Theorem 1.2.  $\square$

**Corollary 1.4.** A diagonalizable group is reflexive, and therefore so is a locally diagonalizable group. If  $M$  and  $N$  are ordinary commutative groups, the natural homomorphism

$$\operatorname{Hom}_{S\text{-gr}}(M_S, N_S) \longrightarrow \operatorname{Hom}_{S\text{-gr}}(D(N_S), D(M_S))$$

is bijective.

---

<sup>3</sup>Editor's note: the sentence that follows has been added.

<sup>4</sup> Since the preceding isomorphism is compatible with base extension, it yields an isomorphism of  $S$ -functors in groups

$$(1) \quad \underline{\mathrm{Hom}}_{S\text{-gr}}(M_S, N_S) \xrightarrow{\sim} \underline{\mathrm{Hom}}_{S\text{-gr}}(D(N_S), D(M_S)).$$

For every  $S$ -prescheme  $T$ ,

$$\underline{\mathrm{Hom}}_{S\text{-gr}}(M_S, N_S)(T) = \mathrm{Hom}_{\mathrm{gr}}(M, \Gamma(N_T/T)),$$

and by I, 1.8,  $\Gamma(N_T/T)$  is the abelian group of locally constant maps  $T \rightarrow N$ . On the other hand, let  $\mathrm{Hom}_{\mathrm{gr}}(M, N)_S$  be the constant  $S$ -group associated with the ordinary abelian group  $\mathrm{Hom}_{\mathrm{gr}}(M, N)$ . There is an evident homomorphism of  $S$ -functors in commutative groups

$$(2) \quad \mathrm{Hom}_{\mathrm{gr}}(M, N)_S \xrightarrow{\theta} \underline{\mathrm{Hom}}_{S\text{-gr}}(M_S, N_S),$$

which is always a monomorphism and is an isomorphism if  $M$  is of finite type.<sup>5</sup>

Part (a) of the following corollary follows from the preceding discussion; part (b) follows from it by the descent results recalled in 1.7.<sup>6</sup>

**Corollary 1.5.**

- (a) *Let  $M$  and  $N$  be ordinary commutative groups, with  $M$  of finite type. Then there is an isomorphism*

$$\mathrm{Hom}_{\mathrm{gr}}(M, N)_S \xrightarrow{\sim} \underline{\mathrm{Hom}}_{S\text{-gr}}(D(N_S), D(M_S));$$

*consequently,  $\underline{\mathrm{Hom}}_{S\text{-gr}}(D(N_S), D(M_S))$  is representable.*

- (b) *More generally, if  $G$  and  $H$  are locally diagonalizable and  $H$  is of finite type, then  $\underline{\mathrm{Hom}}_{S\text{-gr}}(G, H)$  is representable.*

<sup>7</sup> Corollary 1.5 gives:

**Corollary 1.6.** *Under the hypotheses of 1.5, if  $S$  is connected, then*

$$\mathrm{Hom}_{S\text{-gr}}(D_S(N), D_S(M)) \xrightarrow{\sim} \mathrm{Hom}_{\mathrm{gr}}(M, N)$$

and

$$\mathrm{Isom}_{S\text{-gr}}(D_S(N), D_S(M)) \xrightarrow{\sim} \mathrm{Isom}_{\mathrm{gr}}(M, N).$$

---

<sup>4</sup>Editor's note: the discussion that follows has been expanded.

<sup>5</sup>Editor's note: write  $F(M)$  and  $G(M)$  for the left- and right-hand sides of (2), respectively. If  $M = M_1 \oplus M_2$ , there are canonical isomorphisms  $F(M) = F(M_1) \oplus F(M_2)$  and  $G(M) = G(M_1) \oplus G(M_2)$ . It is therefore enough to check that  $\theta$  is an isomorphism for  $M = \mathbf{Z}/r\mathbf{Z}$ ,  $r \geq 0$ . In that case  $F(M) = ({}_rN)_S$ , where  ${}_rN$  is the kernel of  $r \mathrm{id}_N$ , and for every  $T \rightarrow S$  the homomorphism

$$F(M)(T) = \Gamma({}_rN_T/T) \longrightarrow G(M)(T) = {}_r\Gamma(N_T/T)$$

is bijective.

<sup>6</sup>Editor's note: the original stated after 1.5 that, more generally, if  $G$  and  $H$  are locally diagonalizable and  $H$  is of finite type, then  $\underline{\mathrm{Hom}}_{S\text{-gr}}(G, H)$  is representable. That assertion has been included in 1.5 and its proof made explicit in 1.7.

<sup>7</sup>Editor's note: a statement 1.5.1 treating the functor  $\underline{\mathrm{Isom}}_{S\text{-gr}}(G, H)$ , considered in X, 5.10 and 5.11, should be added.

**1.7. Descent of representability.** <sup>8</sup> We recall here several descent results that will be used frequently below.

**Scholium 1.7.1.** *Let  $S$  be a prescheme and let  $X, Y, T$  be  $S$ -preschemes. If  $(T_i)$  is an open covering of  $T$ , and if  $T_{ij} = T_i \cap T_j = T_i \times_T T_j$ , then, because giving a morphism of  $T$ -preschemes  $X_T \rightarrow Y_T$  is local on  $T$ , there is an exact sequence of sets*

$$(1) \quad \mathrm{Hom}_T(X_T, Y_T) \longrightarrow \prod_i \mathrm{Hom}_{T_i}(X_{T_i}, Y_{T_i}) \rightrightarrows \prod_{i,j} \mathrm{Hom}_{T_{ij}}(X_{T_{ij}}, Y_{T_{ij}}).$$

Thus  $\underline{\mathrm{Hom}}_S(X, Y)$  is a local  $S$ -functor, that is, a sheaf on  $(\mathbf{Sch})/S$  for the Zariski topology.

More generally, by IV, 4.5.13,  $\underline{\mathrm{Hom}}_S(X, Y)$  is a sheaf on  $(\mathbf{Sch})/S$  for every topology coarser than the canonical topology, for example the fpqc topology. If  $G$  and  $H$  are group preschemes over  $S$ , it follows that the subfunctor  $\underline{\mathrm{Hom}}_{S\text{-gr}}(G, H)$  is an fpqc sheaf, hence a fortiori a local functor.

**Lemma 1.7.2.** <sup>9</sup> *Let  $F$  be a local  $S$ -functor.*

- (i) *Suppose that there is an open covering  $(S_i)$  of  $S$  such that the restriction  $F_i = F \times_S S_i$  of  $F$  to each  $S_i$  is representable by an  $S_i$ -prescheme  $X_i$ . Then  $F$  is representable by an  $S$ -prescheme  $X$ .*
- (ii) *Suppose that  $F$  is an fpqc sheaf and that there is a faithfully flat quasi-compact morphism  $S' \rightarrow S$  such that  $F' = F \times_S S'$  is representable by an  $S'$ -prescheme  $X'$ . Then  $X'$  carries a descent datum relative to  $S' \rightarrow S$  (cf. IV, 2.1). If this descent datum is effective—as it is when  $X'$  is affine over  $S'$ —then  $F$  is representable by an  $S$ -prescheme  $X$ .*

*Proof.* (i) The hypothesis implies that  $X_i \times_S S_j$  and  $X_j \times_S S_i$  both represent the restriction of  $F$  to  $S_{ij} = S_i \times_S S_j$ . The Yoneda lemma therefore gives a unique isomorphism of  $S_{ij}$ -preschemes

$$c_{ji} : X_i \times_S S_j \xrightarrow{\sim} X_j \times_S S_i.$$

Over  $S_{ijk} = S_i \times_S S_j \times_S S_k$ , we obtain the commutative diagram of isomorphisms

$$\begin{array}{ccc} X_i \times_S S_j \times_S S_k \xrightarrow{c_{ji} \times \mathrm{id}_{S_k}} X_j \times_S S_i \times_S S_k & = & X_j \times_S S_k \times_S S_i \\ \parallel & & \downarrow c_{kj} \times \mathrm{id}_{S_i} \\ X_i \times_S S_k \times_S S_j \xrightarrow{c_{ki} \times \mathrm{id}_{S_j}} X_k \times_S S_i \times_S S_j & = & X_k \times_S S_j \times_S S_i. \end{array}$$

Indeed, all of its objects represent the restriction of  $F$  to  $S_{ijk}$ . Thus the  $c_{ji}$  satisfy the usual cocycle relation  $c_{kj} \circ c_{ji} = c_{ki}$ .

It follows that the  $X_i$  glue to an  $S$ -prescheme  $X$  with  $X \times_S S_i = X_i$  for every  $i$ . For every  $Y$  over  $S_i$ ,

$$(*) \quad F(Y) = F_i(Y) = \mathrm{Hom}_{S_i}(Y, X \times_S S_i) = \mathrm{Hom}_S(Y, X) = h_X(Y).$$

For an arbitrary  $Y \rightarrow S$ , the  $Y_i = Y \times_S S_i$  form an open covering of  $Y$ ; put  $Y_{ij} = Y_i \times_Y Y_j = Y \times_S S_{ij}$ . Since  $F$  is local by hypothesis and  $h_X$  is local because the Zariski topology is

<sup>8</sup>Editor's note: this paragraph has been added to make explicit the "local on  $S$ " character of the representability of certain sheaves on  $S$ , which is used repeatedly below and was used implicitly in the original.

<sup>9</sup>Editor's note: see also Remark XI, 3.4.

coarser than the canonical topology, both  $F(Y)$  and  $h_X(Y)$ , in view of  $(*)$ , identify with the kernels of the respective double arrows

$$\prod_i F(Y_i) \rightrightarrows \prod_{i,j} F(Y_{ij}), \quad \prod_i h_X(Y_i) \rightrightarrows \prod_{i,j} h_X(Y_{ij}).$$

This proves (i).

(ii) Put  $S'' = S' \times_S S'$ , and let  $S''_1$  and  $S''_2$  denote  $S''$  regarded as an  $S'$ -prescheme through the first and second projections. The functors  $F''_1 = F' \times_{S'} S''_1$  and  $F''_2 = F' \times_{S'} S''_2$  are represented respectively by  $X''_1 = X' \times_{S'} S''_1$  and  $X''_2 = X' \times_{S'} S''_2$ . Since

$$F''_1 = F \times_S S'' = F''_2,$$

there is a unique  $S''$ -isomorphism

$$c : X''_1 \xrightarrow{\sim} X''_2.$$

Let  $q_i$ , respectively  $p_{ji}$ , be the projection from  $S''' = S' \times_S S' \times_S S'$  onto its  $i$ -th factor, respectively onto its  $i$ -th and  $j$ -th factors. Let  $X'''_i = X' \times_{S'} S'''_i$ , where  $S'''_i$  means  $S'''$  viewed as an  $S'$ -prescheme through  $q_i$ , and let  $p_{ji}^*(c) : X'''_i \xrightarrow{\sim} X'''_j$  be obtained from  $c$  by base change. We obtain the diagram

$$\begin{array}{ccc} X'''_1 & \xrightarrow{p_{21}^*(c)} & X'''_2 \\ & \searrow p_{31}^*(c) & \downarrow p_{32}^*(c) \\ & & X'''_3. \end{array}$$

All three objects represent the restriction of  $F$  to  $S'''$ , so the diagram commutes:

$$p_{32}^*(c) \circ p_{21}^*(c) = p_{31}^*(c).$$

Thus  $c$  is a descent datum on  $X'$  relative to  $S' \rightarrow S$  (cf. IV, 2.1).

Suppose that this descent datum is effective, so there is an  $S$ -prescheme  $X$  such that  $X' \simeq X \times_S S'$ . This is the case when  $X'$  is affine over  $S'$ , by SGA 1, VIII, 2.1.<sup>10</sup> For every  $Y \rightarrow S'$ , one then has

$$(**) \quad F(Y) = F'(Y) = \operatorname{Hom}_{S'}(Y, X \times_S S') = \operatorname{Hom}_S(Y, X) = h_X(Y).$$

For arbitrary  $Y \rightarrow S$ , put  $Y' = Y \times_S S'$  and  $Y'' = Y' \times_Y Y' \simeq Y \times_S S''$ . Like  $S' \rightarrow S$ , the morphism  $Y' \rightarrow Y$  is faithfully flat and quasi-compact, hence an  $\mathcal{M}$ -effective epimorphism, where  $\mathcal{M}$  is the family of faithfully flat quasi-compact morphisms. Thus the equivalence relation

$$Y' \times_Y Y' \rightrightarrows Y'$$

is  $\mathcal{M}$ -effective and has quotient  $Y$ . Since  $F$  is an fpqc sheaf by hypothesis, while  $h_X$  is one because the fpqc topology is coarser than the canonical topology, both  $F(Y)$  and  $h_X(Y)$ , in view of  $(**)$ , identify with the respective kernels of

$$F(Y') \rightrightarrows F(Y' \times_Y Y'), \quad h_X(Y') \rightrightarrows h_X(Y' \times_Y Y').$$

This proves (ii). □

<sup>10</sup>Editor's note: for another effectivity criterion, see X, 5.4–5.6.

**Corollary 1.7.3.** *Let  $F$  be an fpqc sheaf on  $(\mathbf{Sch})/S$ . Suppose there is an open covering  $(S_i)$  of  $S$ , and for each  $i$  a faithfully flat quasi-compact morphism  $S'_i \rightarrow S_i$ , such that  $F'_i = F \times_S S'_i$  is representable by an  $S'_i$ -prescheme  $X'_i$  affine over  $S'_i$ . Then  $F$  is representable by an  $S$ -prescheme  $X$  affine over  $S$ , with  $X \times_S S'_i = X'_i$  for every  $i$ . If, moreover, every  $X'_i \rightarrow S'_i$  is a closed immersion (respectively, a finite étale morphism), then so is  $X \rightarrow S$ .*

The first assertion follows from 1.7.2. For the second, it is enough to check that every morphism  $X \times_S S_i \rightarrow S_i$  is a closed immersion (respectively, finite and étale); this follows from EGA IV<sub>2</sub>, 2.7.1 (respectively, together with IV<sub>4</sub>, 17.7.3).

**Remark 1.7.4.** Assertion 1.5(b) follows, as announced, from 1.7.1 and 1.7.2(i).

## 2. Scheme-theoretic Properties of Diagonalizable Groups

These properties are summarized in the following proposition.

**Proposition 2.1.** *Let  $S$  be a nonempty prescheme, let  $M$  be an ordinary commutative group, and let  $G = D(M_S)$  be the diagonalizable  $S$ -group defined by  $M$ . Then:*

- (a)  *$G$  is faithfully flat over  $S$  and affine over  $S$  (a fortiori, quasi-compact over  $S$ );*
- (b)  *$M$  is of finite type if and only if  $G$  is of finite type over  $S$ , if and only if  $G$  is of finite presentation over  $S$ ;*
- (c)  *$M$  is finite if and only if  $G$  is finite over  $S$ , if and only if  $G$  is of finite type over  $S$  and is annihilated by an integer  $n > 0$ . In that case,*

$$\deg(G/S) = \text{Card}(M);$$

- (c')  *$M$  is a torsion group if and only if  $G$  is integral over  $S$ ;*
- (d)  *$M = 0$  if and only if  $G$  is the unit  $S$ -group;*
- (e)  *$M$  is of finite type and the order of its torsion subgroup is prime to the residue characteristics of  $S$  if and only if  $G$  is smooth over  $S$ .*

The verification of (a) through (d) is immediate and is left to the reader. Let us prove (e). If  $G$  is smooth over  $S$ , it is locally of finite presentation over  $S$ , hence of finite presentation over  $S$ , since it is affine over  $S$ ; thus  $M$  is of finite type. We may therefore assume from the outset that  $M$  is of finite type, and hence that  $G$  is of finite presentation over  $S$ . Then<sup>11</sup>  $G$  is smooth over  $S$  if and only if its geometric fibers are smooth, reducing us to the case in which  $S$  is the spectrum of an algebraically closed field  $k$ .

Write  $M = T \times L$ , where  $T$  is the torsion subgroup and  $L$  is free, with  $L \simeq \mathbb{Z}^r$ . Then

$$D(M) = D(T) \times D(L), \quad D(L) \simeq \mathbb{G}_m^r,$$

and  $D(L)$  is smooth over  $k$ . Thus  $G = D(M)$  is smooth over  $k$  if and only if  $D(T)$  is smooth. Since  $D(T)$  is finite over  $k$  of degree equal to the order  $n$  of  $T$ , this means that  $D(T)(k)$  has  $n$  elements. Now  $T$  is isomorphic to a sum of groups  $\mathbb{Z}/n_i\mathbb{Z}$ , where  $n = \prod_i n_i$ ; consequently

$$D(T) = \prod_i D(\mathbb{Z}/n_i\mathbb{Z}) = \prod_i \mu_{n_i},$$

<sup>11</sup>Editor's note: this uses the fact that  $G$  is flat over  $S$ , by (a).

where  $\mu_{n_i}$  is the group scheme of  $n_i$ -th roots of unity. Hence

$$\text{card}(D(T)(k)) = \prod_i \text{card}(\mu_{n_i}(k)).$$

Here  $\text{card}(\mu_{n_i}(k))$ , the number of  $n_i$ -th roots of unity in  $k$ , is at most  $n_i$ , and equality holds if and only if  $n_i$  is prime to the characteristic  $p$  of  $k$ . [The source prints  $\mu_i(k)$  in both occurrences.] It follows that

$$\text{card}(D(T)(k)) = n, \quad n = \prod_i n_i,$$

if and only if every  $n_i$  is prime to  $p$ , that is, if and only if  $n$  is prime to  $p$ .  $\square$

### 3. Exactness Properties of the Functor $D_S$

**Theorem 3.1.** *Let  $S$  be a prescheme, and let*

$$0 \longrightarrow M' \xrightarrow{u} M \xrightarrow{v} M'' \longrightarrow 0$$

*be an exact sequence of ordinary commutative groups. Consider the sequence of transposed homomorphisms*

$$0 \longrightarrow D_S(M'') \xrightarrow{v^t} D_S(M) \xrightarrow{u^t} D_S(M') \longrightarrow 0.$$

- (i) *The homomorphism  $v^t$  induces an isomorphism from  $D_S(M'')$  onto the kernel of  $u^t$ , and  $u^t$  is faithfully flat and quasi-compact.*
- (ii) <sup>12</sup>  *$D_S(M')$  represents the quotient sheaf  $D_S(M)/D_S(M'')$  for the (fpqc) topology.*

Let  $\mathcal{M}$  denote the family of faithfully flat quasi-compact morphisms. First, (ii) follows from (i) (cf. IV, 4.6.5.1). Indeed, the equivalence relation on  $D_S(M)$  defined by  $u^t$  is the same as that defined by the subgroup

$$\text{Ker}(u^t) = D_S(M'').$$

Since  $u^t \in \mathcal{M}$ , this equivalence relation is  $\mathcal{M}$ -effective (cf. IV, 3.3.2.1); therefore  $D_S(M')$  represents the quotient sheaf for the (fpqc) topology (cf. IV, 4.6.5).

The first assertion of (i) is an immediate consequence of the definition of the functors  $D_S(-)$ . More generally, every exact sequence

$$M' \longrightarrow M \longrightarrow M'' \longrightarrow 0$$

with no zero on the left gives rise to an exact transposed sequence

$$0 \longrightarrow D_S(M'') \longrightarrow D_S(M) \longrightarrow D_S(M').$$

(This remains valid in the more general setting considered at the beginning of §1.) On the other hand, because  $D_S(M)$  and  $D_S(M')$  are affine over  $S$ , the morphism  $u^t$  is necessarily affine, and a fortiori quasi-compact, whatever the homomorphism  $u : M' \rightarrow M$ . The second assertion of (i) will therefore follow from part (a) of the following corollary.

**Corollary 3.2.** *Let  $S$  be a nonempty prescheme, let  $u : M' \rightarrow M$  be a homomorphism of ordinary commutative groups, and let  $u^t : G \rightarrow G'$  be the transposed homomorphism. Then:*

---

<sup>12</sup>Editor's note: what follows has been added, and the proof has been expanded accordingly.



- (a)  $u$  is a monomorphism if and only if  $u^t$  is faithfully flat;
- (b)  $u$  is an epimorphism if and only if  $u^t$  is a monomorphism; in that case  $u^t$  is in fact a closed immersion.

To prove (a), observe that if  $u$  is a monomorphism, then  $\mathcal{O}_S(M)$ , as a module over  $\mathcal{O}_S(M')$ , has a nonempty basis: namely, the system of sections defined by any system of representatives of  $M$  modulo  $M'$ . It is therefore faithfully flat. Conversely, if  $\mathcal{O}_S(M)$  is faithfully flat over  $\mathcal{O}_S(M')$ , then

$$u^t : \mathcal{O}_S(M') \longrightarrow \mathcal{O}_S(M)$$

is injective, which, when  $S \neq \emptyset$ , implies that  $u : M' \rightarrow M$  is injective.

To prove (b), observe that if  $u$  is an epimorphism, then  $\mathcal{O}_S(M') \rightarrow \mathcal{O}_S(M)$  is surjective, so  $u^t$  is a closed immersion and, a fortiori, a monomorphism. Conversely, if  $u^t$  is a monomorphism, then  $\text{Ker } u^t$  is the unit group. Putting  $M'' = \text{Coker } u$ , we have seen that

$$\text{Ker } u^t \simeq D_S(M'').$$

Thus 2.1(d) gives  $M'' = 0$ , and  $u$  is an epimorphism.

The usual argument now gives the following consequence of 3.1.

**Corollary 3.3.** *Let*

$$M' \xrightarrow{u} M \xrightarrow{v} M''$$

*be an exact sequence of ordinary commutative groups, and consider the transposed sequence*

$$G'' \xrightarrow{v^t} G \xrightarrow{u^t} G'.$$

*Then  $v^t$  induces a faithfully flat quasi-compact morphism from  $G''$  to  $\text{Ker } u^t$ , and this kernel is a diagonalizable group isomorphic to*

$$D_S(v(M)) = D_S(\text{Coker } u).$$

**Corollary 3.4.** *Let  $S$  be a prescheme, let  $u : G \rightarrow H$  be a homomorphism of  $S$ -group preschemes that are locally diagonalizable, with  $H$  of finite type over  $S$ , and put  $G' = \text{Ker } u$ . Then:*

- (a)  $G'$  is locally diagonalizable, and is of finite type over  $S$  if  $G$  is;
- (b) the quotient  $G/G'$  “exists”: more precisely, the equivalence relation on  $G$  defined by  $G'$  is  $\mathcal{M}$ -effective, where  $\mathcal{M}$  is the family of faithfully flat quasi-compact morphisms (cf. IV, 3.4). Moreover,  $G/G'$  is locally diagonalizable and of finite type over  $S$ ;
- (c) the homomorphism  $u : G \rightarrow H$  factors uniquely as

$$G \xrightarrow{v} G/G' \xrightarrow{w} H,$$

*where  $v$  is the canonical homomorphism, and is therefore faithfully flat and quasi-compact. Moreover,  $w$  is a closed immersion, and a fortiori a monomorphism.*

*Finally, the quotient*

$$H' = H / \text{Im } w = \text{Coker } w = \text{Coker } u$$

*exists: more precisely, the equivalence relation on  $H$  defined by  $G/G'$  is  $\mathcal{M}$ -effective, and  $H'$  is of finite type over  $S$ .*

The first assertion of (c) follows from (b), by the definition of the quotient  $G/G'$  (cf. IV, 3.2.3).<sup>13</sup> Let us show that the (fpqc) quotient sheaf  $\widetilde{G}/\widetilde{G'}$  is representable. This can be checked locally on  $S$ , as can all the other assertions; we may therefore suppose that  $G$  and  $H$  are diagonalizable, of the form  $D_S(M)$  and  $D_S(N)$ .

Since  $H$  is of finite type over  $S$ , the group  $N$  is of finite type by 2.1(b). Therefore, by 1.5,  $u$  is defined by a homomorphism  $u' : N \rightarrow M$ . It follows from 3.1 and 3.2 that

$$G' \simeq D_S(\text{Coker } u')$$

and that  $\widetilde{G}/\widetilde{G'}$  is represented by  $D_S(\text{Im } u')$ . Moreover, on considering the exact sequence

$$0 \longrightarrow \text{Ker } u' \longrightarrow N \xrightarrow{w'} \text{Im } u' \longrightarrow 0,$$

we find that  $w$  is a closed immersion and that

$$H' = H / \text{Im } w = D_S(\text{Ker } u').$$

The latter is of finite type over  $S$ , since  $N$ , and hence  $\text{Ker } u'$ , is of finite type.

**Remarks.** The existence result for quotients in 3.4 will be substantially generalized in §5.

On the other hand, note that in the present section and the preceding one, the hypothesis  $S \neq \emptyset$  was used only to ensure the validity of certain converses, allowing properties of the corresponding ordinary groups to be deduced from certain hypotheses on diagonalizable  $S$ -groups. The results in the “forward” direction hold without any restriction on  $S$ , and the proofs given here apply in general.

**Corollary 3.5.** *Let  $G$  be a diagonalizable group prescheme over  $S$ , and let  $n \neq 0$  be an integer. Then the subgroup  ${}_nG$  of  $G$ , the kernel of the homomorphism*

$$n \cdot \text{id}_G : G \longrightarrow G,$$

*is integral over  $S$ , and is finite over  $S$  if  $G$  is of finite type over  $S$ .*

Indeed, if  $G = D_S(M)$ , then

$${}_nG = D_S(M/nM)$$

by 3.1, and the result follows from 2.1(b), (c), and (c').

## 4. Torsors under a Diagonalizable Group

Let  $S$  be a prescheme and let  $G = D_S(M)$  be a diagonalizable group over  $S$ . We shall determine the  $G$ -torsors (or principal homogeneous  $G$ -bundles) over  $S$  for the faithfully flat quasi-compact topology; cf. Exposé IV, 5.1. Recall that a prescheme  $P$  over  $S$ , with  $S$ -operator group  $G$ , is called a *torsor* or a *principal homogeneous bundle* if every point of  $S$  has an open neighborhood  $U$  and a faithfully flat quasi-compact morphism  $S' \rightarrow U$  such that

$$P' = P \times_S S'$$

is isomorphic, as an object with operators, to

$$G' = G \times_S S',$$

<sup>13</sup>Editor's note: the original has been expanded in what follows.

where  $G'$  acts on itself by right translations. Since  $G$  is affine over  $S$ , SGA 1, VIII 5.6 implies that  $P$  is necessarily affine over  $S$ . Also, because  $G$  itself is faithfully flat and quasi-compact over  $S$ ,  $P$  is principal homogeneous under  $G$  if and only if it is formally principal homogeneous and is, in addition, faithfully flat and quasi-compact over  $S$  (cf. IV, 5.1.6).

Recall also (Exposé I, 4.7.3) that giving an  $S$ -prescheme  $P$ , affine over  $S$ , with  $S$ -operator group  $G = D_S(M)$ , amounts to giving a commutative quasi-coherent algebra graded by  $M$  over  $S$ . In other words, it amounts to giving a quasi-coherent algebra  $\mathcal{A}$  over  $S$ , together with a direct-sum decomposition as a module

$$\mathcal{A} = \coprod_{m \in M} \mathcal{A}_m,$$

such that

$$\mathcal{A}_m \cdot \mathcal{A}_{m'} \subseteq \mathcal{A}_{m+m'} \quad \text{for } m, m' \in M.$$

With this understood, the answer to the preceding problem is as follows.

**Proposition 4.1.** *Let  $P$  be the prescheme with  $S$ -operator group  $G = D_S(M)$  defined by the  $M$ -graded algebra  $\mathcal{A}$ . Then  $P$  is a principal homogeneous bundle under  $G$  if and only if  $\mathcal{A}$  satisfies the following conditions:*

- (a) *for every  $m \in M$ ,  $\mathcal{A}_m$  is an invertible module over  $S$ ;*
- (b) *for  $m, m' \in M$ , the homomorphism*

$$\mathcal{A}_m \otimes_{\mathcal{O}_S} \mathcal{A}_{m'} \longrightarrow \mathcal{A}_{m+m'}$$

*induced by multiplication in  $\mathcal{A}$  is an isomorphism.*

The necessity of these conditions follows immediately by descent, since they hold when  $P$  is the trivial principal homogeneous bundle, that is, when  $\mathcal{A} = \mathcal{O}_S(M)$ . For sufficiency, note that (a) already implies that  $P$  is faithfully flat over  $S$ ; it is in any event quasi-compact over  $S$ , since it is affine over  $S$ . It therefore remains to check that  $P$  is formally principal homogeneous under  $G$ , that is, that the familiar morphism

$$P \times_S G \longrightarrow P \times_S P$$

is an isomorphism. On affine algebras this morphism is given explicitly by the homomorphism

$$\mathcal{A} \otimes \mathcal{A} \longrightarrow \mathcal{A}(M) = \mathcal{A} \otimes \mathcal{O}_S(M).$$

In bidegree  $(m, n)$ , where  $m, n \in M$ , it is given by

$$x_m \otimes y_n \longmapsto x_m y_n \otimes e_n.$$

At the level of degrees, this homomorphism is compatible with the homomorphism  $M \times M \rightarrow M \times M$  given by

$$(m, n) \longmapsto (m + n, n),$$

which is an isomorphism. This shows that (b) expresses precisely, and independently of (a), that  $P$  is formally principal homogeneous, and proves 4.1.

Faithfully flat descent also gives:

**Corollary 4.2.** *The conditions of 4.1 imply that the homomorphism*

$$\mathcal{O}_S \longrightarrow \mathcal{A}_0$$

is an isomorphism.

For example, if  $M = \mathbb{Z}$ , then, under the conditions of 4.1, the algebra  $\mathcal{A}$  is essentially determined by  $\mathcal{A}_1 = \mathcal{L}$ , namely

$$\mathcal{A} \simeq \prod_{n \in \mathbb{Z}} \mathcal{L}^{\otimes n}$$

as graded algebras. Thus we recover the familiar result:

**Corollary 4.3.** *The category of principal homogeneous bundles  $P$  over  $S$  under  $\mathbb{G}_{m,S}$  is equivalent to the category of invertible modules  $\mathcal{L}$  over  $S$ , where the morphisms in both categories are the isomorphisms of the structures in question. Quasi-inverse functors are obtained by assigning to  $P$  the degree-1 component of its  $\mathbb{Z}$ -graded affine algebra, and by assigning to  $\mathcal{L}$  the spectrum of the  $\mathbb{Z}$ -graded algebra*

$$\prod_{n \in \mathbb{Z}} \mathcal{L}^{\otimes n}.$$

In particular:

**Corollary 4.4.** *The group of isomorphism classes of principal homogeneous bundles over  $S$  under  $\mathbb{G}_{m,S}$  is isomorphic to the group  $\text{Pic}(S)$  of isomorphism classes of invertible modules over  $S$ , that is, to*

$$H^1(S, \mathcal{O}_S^\times).$$

Since  $\mathbb{G}_{m,S}$  is the scheme of automorphisms of the module  $\mathcal{O}_S$ , Corollary 4.4 is equivalent to the following statement, one of the forms of Hilbert's "Theorem 90."

**Corollary 4.5.** *Every principal homogeneous bundle over  $S$  under  $\mathbb{G}_m$  is locally trivial for the Zariski topology.*

**Remark 4.5.1.** The preceding statement is no longer true in general for a group such as  $\mu_n$ , or for a "twisted form" of  $\mathbb{G}_m$ . For example, the unique twisted form of  $\mathbb{G}_m$  over the field  $\mathbb{R}$  of real numbers has a first cohomology group equal to  $\mathbb{Z}/2\mathbb{Z}$ .

<sup>14</sup> Indeed, let  $\mathbb{S}^1$  be the kernel of the norm morphism

$$N : \prod_{\mathbb{C}/\mathbb{R}} \mathbb{G}_{m,\mathbb{C}} \longrightarrow \mathbb{G}_{m,\mathbb{R}};$$

it is a  $\mathbb{C}/\mathbb{R}$ -twisted form of  $\mathbb{G}_{m,\mathbb{R}}$ . The equation  $N(z) = -1$  in  $\prod_{\mathbb{C}/\mathbb{R}} (\mathbb{G}_{m,\mathbb{C}})$  defines an  $\mathbb{S}^1$ -torsor  $X$  over  $\text{Spec}(\mathbb{R})$ , locally trivial for the étale topology, but not trivial, since  $X(\mathbb{R}) = \emptyset$ . Let us show that

$$H_{\text{ét}}^1(\mathbb{R}, \mathbb{S}^1) \simeq \mathbb{Z}/2\mathbb{Z}.$$

There is an exact sequence of smooth commutative algebraic  $\mathbb{R}$ -groups

$$1 \longrightarrow \mathbb{S}^1 \longrightarrow \prod_{\mathbb{C}/\mathbb{R}} \mathbb{G}_{m,\mathbb{C}} \longrightarrow \mathbb{G}_{m,\mathbb{R}} \longrightarrow 1,$$

which gives rise to a long exact sequence in étale cohomology (or in Galois cohomology):

$$\begin{aligned} 0 \longrightarrow \mathbb{S}^1(\mathbb{R}) \longrightarrow \mathbb{C}^\times &\xrightarrow{N} \mathbb{R}^\times \longrightarrow H_{\text{ét}}^1(\mathbb{R}, \mathbb{S}^1) \\ &\longrightarrow H_{\text{ét}}^1\left(\mathbb{R}, \prod_{\mathbb{C}/\mathbb{R}} \mathbb{G}_{m,\mathbb{C}}\right) \longrightarrow \cdots \end{aligned}$$

<sup>14</sup>Editor's note: what follows has been added.

Now (see, for example, XXIV, 8.4),

$$H_{\text{ét}}^1\left(\mathbb{R}, \prod_{\mathbb{C}/\mathbb{R}} \mathbb{G}_{m,\mathbb{C}}\right) \simeq H_{\text{ét}}^1(\mathbb{C}, \mathbb{G}_{m,\mathbb{C}}),$$

and the latter group is zero by 4.5 (or, in this case, because  $\mathbb{C}$  is algebraically closed). We therefore obtain an isomorphism

$$H_{\text{ét}}^1(\mathbb{R}, \mathbb{S}^1) \simeq \mathbb{R}^\times / N(\mathbb{C}^\times) \simeq \{\pm 1\}.$$

We shall need the following result in the next section.

**Proposition 4.6.** *Under the conditions of 4.1, conditions (a) and (b) are equivalent to the following conditions:*

- (a') *the homomorphism  $\mathcal{O}_S \rightarrow \mathcal{A}_0$  is an isomorphism;*
- (b') *for every  $m \in M$  (it is enough to require this for the members of a system of generators of  $M$ ), one has*

$$\mathcal{A}_m \cdot \mathcal{A}_{-m} = \mathcal{A}_0.$$

The necessity is clear in view of 4.2.<sup>15</sup> We are thus reduced to proving the following corollary.

**Corollary 4.7.** *Let*

$$A = \prod_{n \in \mathbb{Z}} A_n$$

*be a  $\mathbb{Z}$ -graded ring such that*

$$A_1 \cdot A_{-1} = A_0.$$

*Then the  $A_n$  are invertible  $A_0$ -modules, and, for  $n, n' \in \mathbb{Z}$ , the homomorphism*

$$A_n \otimes_{A_0} A_{n'} \longrightarrow A_{n+n'}$$

*induced by multiplication in  $A$  is an isomorphism.*

By hypothesis there exist  $f_i \in A_1$  and  $g_i \in A_{-1}$  such that

$$(*) \quad \sum_i f_i g_i = 1.$$

Since the assertion to be proved is local on  $\text{Spec}(A_0)$ ,<sup>16</sup> and since (\*) shows that  $\text{Spec}(A_0)$  is covered by the affine open subsets  $D(f_i g_i)$ , we are reduced to the case in which there is an element  $f \in A_1$  that is invertible in  $A$ . Then, for every  $n \in \mathbb{Z}$ ,  $f^n$  is an invertible element of  $A_n$ , and therefore defines an isomorphism

$$A_0 \xrightarrow{\sim} A_n, \quad h \longmapsto f^n h.$$

Moreover, this shows that one obtains an isomorphism

$$A_0[t, t^{-1}] \longrightarrow A$$

of graded  $A_0$ -algebras by sending  $t$  to  $f$ , which completes the proof of 4.7.

<sup>15</sup>Editor's note: "4.1" has been corrected to "4.2."

<sup>16</sup>Editor's note: the following passage has been expanded.

Under the hypotheses of 4.6, 4.7 already shows that the  $\mathcal{A}_m$ , for  $m \in M$ , are invertible. To prove condition 4.1(b), we may therefore suppose that  $\mathcal{A}_m$  and  $\mathcal{A}_{m'}$  have bases  $f_m$  and  $f_{m'}$ , with inverses

$$f_m^{-1} \in \Gamma(\mathcal{A}_{-m}) \quad \text{and} \quad f_{m'}^{-1} \in \Gamma(\mathcal{A}_{-m'}).$$

Multiplication by

$$f_m^{-1} f_{m'}^{-1} \in \Gamma(\mathcal{A}_{-m-m'})$$

then defines a homomorphism

$$\mathcal{A}_{m+m'} \longrightarrow \mathcal{A}_0 \simeq \mathcal{O}_S$$

that sends the image of  $f_m \otimes f_{m'}$  to the unit section of  $\mathcal{O}_S$ . In the diagram

$$\begin{array}{ccccc} & & w & & \\ & \nearrow & & \searrow & \\ \mathcal{A}_m \otimes \mathcal{A}_{m'} & \xrightarrow{u} & \mathcal{A}_{m+m'} & \xrightarrow{v} & \mathcal{A}_0 \simeq \mathcal{O}_S \end{array}$$

the maps  $w$  and  $v$  are therefore epimorphisms of invertible sheaves, hence isomorphisms; consequently  $u$  is an isomorphism.  $\square$

## 5. Quotient of an Affine Scheme by a Diagonalizable Group Acting Freely

<sup>17</sup> We denote by  $\mathcal{M}$  the set of faithfully flat quasi-compact morphisms, and recall that torsors are understood with respect to the (fpqc) topology.

**Theorem 5.1.** *Let  $S$  be a prescheme,  $M$  an ordinary commutative group,  $G = D_S(M)$  the diagonalizable group over  $S$  that it defines, and  $P$  an  $S$ -prescheme affine over  $S$ , on which  $G$  acts freely on the right.*

*Then the equivalence relation defined by  $G$  on  $P$  is  $\mathcal{M}$ -effective (cf. IV, 3.4); that is, the quotient  $X = P/G$  exists and  $P$  is a torsor over  $X$  under  $G_X = D_X(M)$ . Moreover,  $P/G$  is affine over  $S$ . More precisely, if  $P$  is defined by the  $M$ -graded algebra  $\mathcal{A}$ , then  $P/G$  is isomorphic to  $\text{Spec}(\mathcal{A}_0)$ , where  $\mathcal{A}_0 = \mathcal{A}^G$  is the degree-zero component of  $\mathcal{A}$ .*

*Proof.* Put  $X = \text{Spec}(\mathcal{A}_0)$ . The inclusion  $\mathcal{A}_0 \rightarrow \mathcal{A}$  gives a natural morphism  $P \rightarrow X$ , which is invariant under the action of  $G$ . Thus  $P$  becomes an  $X$ -prescheme, affine over  $X$ , with operator group  $G_X = D_X(M)$ ; the hypothesis that  $G$  acts freely on  $P/S$  implies that  $G_X$  acts freely on  $P/X$ . It remains only to show that  $P$  is a principal homogeneous bundle under  $G_X$ , using the fact that  $\mathcal{B}_0 = \mathcal{O}_X$ , where  $\mathcal{B}$  is the  $M$ -graded algebra over  $X$  defining  $P/X$ .

We may therefore suppose that  $X = S$  and that  $S$  is affine. Then  $P$  is affine, defined by an  $M$ -graded ring  $A$ ; write  $A_m$  for its homogeneous part of degree  $m$ , so that  $S = \text{Spec}(A_0)$ . In view of 4.6, it remains to verify

$$(x) \quad A_m \cdot A_{-m} = A_0 \quad \text{for every } m \in M.$$

A direct calculation also shows that (x) is equivalent to saying that

$$P \times_S G \longrightarrow P \times_S P$$

<sup>17</sup>Editor's note: We have added the following sentence.

is a closed immersion, rather than merely a monomorphism (as follows from the hypothesis that  $G$  acts freely); equivalently, the homomorphism of affine rings

$$\theta : A \otimes_{A_0} A \longrightarrow A(M)$$

is surjective.<sup>18</sup> This proves 5.1 if one assumes that the equivalence relation defined by  $G$  on  $P$  is closed. We shall show, however, that this hypothesis already follows from the fact that  $G$  acts freely. This is also implicit in Theorem 5.1: the scheme  $G \times_S P$  must be isomorphic to  $P \times_X P$ , which is closed in  $P \times_S P$  because  $X$  is affine, hence separated, over  $S$ .

Let  $R = P \times_S G$ . The hypothesis that  $G$  acts freely, that is, that  $R \rightarrow P \times_S P$  is a monomorphism, says that the diagonal morphism

$$R \longrightarrow R \times_{(P \times_S P)} R = R'$$

is an isomorphism. We have  $R = \text{Spec}(A(M))$  and

$$R' = \text{Spec}(A(M \times M)/K),$$

<sup>19</sup> where  $K$  is the ideal generated by the elements

$$x_m(e_{m,0} - e_{0,m}), \quad m \in M, \quad x_m \in A_m.$$

Let

$$\phi : A(M \times M) \longrightarrow A(M)$$

be the surjective ring homomorphism defined by

$$xe_{m,n} \longmapsto xe_{m+n} \quad (m, n \in M, x \in A),$$

where the  $e_m$ , respectively  $e_{m,n} = e_m \otimes e_n$ , are the elements of the canonical basis of  $A(M)$ , respectively  $A(M \times M)$ . The diagonal morphism  $R \rightarrow R'$  then corresponds to the homomorphism

$$\bar{\phi} : A(M \times M)/K \longrightarrow A(M)$$

obtained by passing to the quotient by  $K$ . The kernel of  $\bar{\phi}$  is the ideal  $K'$  generated by

$$d_m = e_{m,0} - e_{0,m}.$$

[The source prints  $\ker(\bar{\phi}) = K'$ ; after quotienting by  $K$ , the corresponding kernel is  $K'/K$ .] We have  $K \subseteq K'$ , and the hypothesis that  $G$  acts freely on  $P$ , that is, that  $\bar{\phi}$  is an isomorphism, is equivalent to  $K' = K$ , expressed by the relations

$$(xx) \quad d_m \in K = \sum_p A(M \times M) A_p d_p \quad \text{for every } m \in M.$$

Using the natural trigrading of  $A(M \times M)$ , and the fact that the first degree of  $d_m$  is zero, this says that  $d_m$  can be written as a sum of elements of the form

$$f e_{r,s}(e_{p,0} - e_{0,p}), \quad f \in A_{-p} \cdot A_p.$$

<sup>18</sup>Editor's note: Indeed, for every  $b \in A$  and  $a_m \in A_m$ , one has  $\theta(b \otimes a_m) = ba_m \otimes e_m$ ; hence the surjectivity of  $\theta$  is equivalent to (x).

<sup>19</sup>Editor's note: We have denoted by  $K$  the ideal denoted by  $J$  in the original, in order to distinguish it from the ideals  $J_p$  of  $A_0$  occurring in (xxx). We have also made explicit below that the relations (xx) are equivalent to the equality  $K = K'$ ; see what follows.

Since the total degree of  $d_m$  is  $m$ , it is enough to use terms satisfying

$$r + s + p = m.$$

Thus, for every  $m \in M$ , there must be an expression

$$(xxx) \quad \begin{cases} d_m = e_{m,0} - e_{0,m} = \sum_{r,s} \lambda_{r,s} (e_{m-s,s} - e_{r,m-r}), \\ \lambda_{r,s} \in J_p = A_p \cdot A_{-p} \subseteq A_0, \quad p = m - (r + s). \end{cases}$$

We must deduce (x), that is, the relations

$$(xxxx) \quad J_n = A_0 \quad \text{for every } n \in M.$$

For this it is enough to establish the same relation modulo every maximal ideal of  $A_0$ . Since all the hypotheses are preserved under such a reduction, we may assume that  $A_0$  is a field.

**Lemma 5.2.** *Under the preceding conditions, with  $A_0$  a field, if  $M \neq 0$ , there exists  $p \in M - \{0\}$  such that  $J_p = A_0$ .*

Indeed, otherwise the sum on the right-hand side of (xxx) would vanish for every  $m \in M$ , which is absurd.

**Corollary 5.3.** *Under the preceding conditions, but without assuming that  $A_0$  is a field, there are finitely many elements  $p_i \in M - \{0\}$  such that*

$$\sum_i J_{p_i} = A_0.$$

Indeed, apply 5.2 to the situations obtained from  $A/A_0$  by reduction modulo the maximal ideals of  $A_0$ .<sup>20</sup>

**Corollary 5.4.** *Suppose again that  $A_0$  is a field. For every subgroup  $N$  of  $M$  with  $N \neq M$ , there exists  $p \in M - N$  such that  $J_p = A_0$ .*

Indeed, let  $M' = M/N$ , and consider the  $M'$ -graded ring  $A'$ , whose underlying ring is  $A$  and whose grading is given by

$$A'_{m'} = \coprod_{m \in h^{-1}(m')} A_m,$$

where  $h : M \rightarrow M' = M/N$  is the canonical homomorphism. Geometrically, this construction amounts to considering the subgroup  $G' = D_S(M')$  of  $G$ , and the structure on  $P$  of a scheme with operator group  $G'$  induced by the action of  $G$ . It is then clear that  $G'$  acts freely on  $P$  [the source prints  $P'$ ]; that is, the pair  $(M', A')$  satisfies the hypotheses of 5.3. Thus

$$1 = \sum_i f_i g_i, \quad f_i \in A'_{m'_i}, \quad g_i \in A'_{-m'_i}, \quad m'_i \in M' - \{0\}.$$

Taking the components of the right-hand side along  $A_0$ , and using the fact that  $A_0$  is a field, immediately gives 5.4.

<sup>20</sup>Editor's note: It follows from 5.2 that  $\sum_{p \neq 0} J_p = A_0$ ; hence 1 can be written as a finite sum  $\sum_i x_i$ , with  $x_i \in J_{p_i}$ .



Now observe that

$$J_p J_q \subseteq J_{p+q} \quad \text{and} \quad J_p = J_{-p}.$$

Thus, if  $N$  denotes the set of  $m \in M$  such that  $J_m = A_0$  [the source prints  $J_p = A_0$ ], then  $N$  is a subgroup of  $M$ . By 5.4, it equals  $M$ . This completes the proof of Theorem 5.1.

As noted in the course of the proof, Theorem 5.1 implies:

**Corollary 5.5.** *Under the hypotheses of 5.1, the graph morphism*

$$P \times_S G \longrightarrow P \times_S P$$

*is a closed immersion.*

It follows immediately that:

**Corollary 5.6.** *Let  $\sigma$  be a section of  $P$  over  $S$ . Then the morphism  $g \mapsto \sigma \cdot g$  from  $G$  to  $P$  defined by  $\sigma$  is a closed immersion.*

**Corollary 5.7.** *Let  $G$  and  $H$  be two  $S$ -groups, with  $G$  diagonalizable and  $H$  affine over  $S$ , and let  $u : G \rightarrow H$  be a homomorphism of  $S$ -groups that is a monomorphism. Then  $u$  is a closed immersion,  $H/G = X$  exists,  $H$  is a principal homogeneous bundle over  $X$  under  $G_X$ , and  $X$  is affine over  $S$ .*

**Corollary 5.8.** *Under the hypotheses of 5.1, if  $P$  is of finite type (respectively, of finite presentation) over  $S$ , then the same is true of  $X = P/G$ .*

Indeed, the hypothesis implies that the fibers of  $G_X$  are of finite type. Hence  $G_X$  is of finite presentation over  $X$ , by 2.1(b); since  $P$  is a torsor under  $G_X$ , it is therefore of finite presentation over  $X$ .<sup>21</sup> It is also faithfully flat over  $X$ , so the conclusion follows from Exposé V, Proposition 9.1.

## 6. Essentially Free Morphisms and Representability of Certain Functors of the Form $\coprod_{Y/S} Z/Y^{(*)}$

**Definition 6.1.** Let  $f : X \rightarrow S$  be a morphism of preschemes. We say that  $f$  is *essentially free*, or that  $X$  is *essentially free over  $S$* , if there are an affine open covering  $(S_i)$  of  $S$ , for each  $i$  an  $S_i$ -prescheme  $S'_i$  that is affine and faithfully flat over  $S_i$ , and a covering  $(X'_{ij})_j$  of  $X'_i = X \times_S S'_i$  by affine opens  $X'_{ij}$ , such that, for every  $(i, j)$ , the ring of  $X'_{ij}$  is a free module over the ring of  $S'_i$ .<sup>22</sup>

**Proposition 6.2.**

- (a) *If  $X$  is essentially free over  $S$ , then it is flat over  $S$ . The converse holds if  $S$  is Artinian.*
- (b) *If  $S$  is the spectrum of a field, every  $S$ -prescheme is essentially free over  $S$ .*
- (c) *If  $X$  is essentially free over  $S$ , and if  $S' \rightarrow S$  is a base-change morphism, then  $X' = X \times_S S'$  is essentially free over  $S'$ . The converse holds if  $S' \rightarrow S$  is faithfully flat and quasi-compact.*

<sup>21</sup>Editor's note: Cf. EGA II, 2.7.1(vi).

(\*) This section is independent of the theory of diagonalizable groups; its natural place would be in Exposé VI<sub>B</sub>.

<sup>22</sup>Editor's note: "Free" may be replaced by "projective"; cf. 6.8 below. This notion should also be compared with that of an  $S$ -prescheme that is flat and pure, introduced and developed in [RG71]; see in particular *loc. cit.*, Part I, 3.3.12, and Part II, 3.1.4.1.

The proof is immediate, using for the converse in (a) the fact that a flat module over an Artinian local ring is free.<sup>23</sup>

**Proposition 6.3.** *Let  $H$  be a diagonalizable  $S$ -group prescheme (more generally, one that becomes diagonalizable after a suitable faithfully flat quasi-compact extension of every affine open of  $S$ ; that is,  $H$  is “of multiplicative type,” cf. IX, 1.1). Then  $H$  is essentially free over  $S$ .*

Indeed, if  $H$  is diagonalizable, it is affine over  $S$  and is defined by an algebra that is a free  $\mathcal{O}_S$ -module.

The introduction of Definition 6.1 is justified by the following theorem.

**Theorem 6.4.** *Let  $S$  be a prescheme,  $Z$  an  $S$ -prescheme essentially free over  $S$ , and  $Y$  a closed subprescheme of  $Z$ . Consider the functor<sup>24</sup>*

$$F = \coprod_{Z/S} Y/Z : (\mathbf{Sch}/S)^\circ \longrightarrow \mathbf{Set},$$

$$F(S') = \Gamma(Y_{S'}/Z_{S'}) = \begin{cases} \emptyset, & Z_{S'} \neq Y_{S'}, \\ \{\mathrm{id}_{Z_{S'}}\}, & Z_{S'} = Y_{S'}. \end{cases}$$

*This functor is representable by a closed subprescheme of  $S$ .<sup>25</sup>*

<sup>26</sup> First note that  $F$  is a sheaf for the (fpqc) topology. Since  $F(S')$  is either  $\emptyset$  or  $\{\mathrm{pt}\}$  for every  $S'$ , it is enough to check the following assertion: if  $(S_i)$  is an open covering of  $S$  (respectively, if  $S' \rightarrow S$  is faithfully flat and quasi-compact), and every  $Y_{S_i} \rightarrow Z_{S_i}$  is an isomorphism (respectively,  $Y_{S'} \rightarrow Z_{S'}$  is an isomorphism), then  $Y \rightarrow Z$  is an isomorphism. This is clear in the first case and follows from SGA 1, VIII, 5.4, or EGA IV<sub>2</sub>, 2.7.1, in the second.

Moreover, by SGA 1, VIII, 2.1 and 5.5, faithfully flat quasi-compact morphisms are morphisms of effective descent for the fibered category of arrows that are closed immersions. Thus, with the notation of 6.1, we may reduce to the case  $S = S'_i$ .

Let  $(Z_j)$  be a covering of  $Z$  by affine opens such that  $\mathcal{O}(Z_j)$  is a free module over  $A = \mathcal{O}(S)$ , put  $Y_j = Y \cap Z_j$ , and let

$$F_j : (\mathbf{Sch}/S)^\circ \longrightarrow \mathbf{Set}$$

be the functor defined from  $(Z_j, Y_j)$  as  $F$  is defined from  $(Z, Y)$ . It is a subfunctor of the terminal functor, and clearly

$$F = \bigcap_j F_j.$$

It therefore suffices to prove that every  $F_j$  is representable by a closed subscheme  $T_j$  of  $S$ , for then  $F$  is represented by the closed subscheme  $T = \bigcap_j T_j$ . We may consequently suppose that  $Z$  is affine as well, say  $Z = \mathrm{Spec}(B)$ , where  $B$  is a free  $A$ -module. Let  $J$  be a subset of  $B$  defining the closed subprescheme  $Y$  of  $Z$ , and let  $K$  be the ideal of  $A$

<sup>23</sup>Editor's note: Indeed, let  $(A, \mathfrak{m})$  be an Artinian local ring,  $k$  its residue field,  $M$  an arbitrary  $A$ -module, and  $(x_i)_{i \in I}$  elements of  $M$  whose images form a basis of  $M/\mathfrak{m}M$  over  $k$ . Let  $F$  be the free  $A$ -module with basis  $(e_i)_{i \in I}$ , and let  $\phi : F \rightarrow M$  be the  $A$ -morphism defined by  $\phi(e_i) = x_i$ . Then  $Q = \mathrm{Coker} \phi$  satisfies  $Q = \mathfrak{m}Q$ , whence  $Q = 0$ , since  $\mathfrak{m}$  is nilpotent. Suppose moreover that  $M$  is flat over  $A$ . Then  $K = \mathrm{Ker} \phi$  satisfies  $K \otimes_A k = 0$ , that is,  $K = \mathfrak{m}K$ , whence  $K = 0$ .

<sup>24</sup>Editor's note: Cf. II.1, where this functor is denoted  $\coprod_{Z/S} Y$ .

<sup>25</sup>Editor's note: We have corrected  $Z$  to  $S$ .

<sup>26</sup>Editor's note: We have expanded the original in what follows; see also 1.7.3.

generated by the  $u_i(J) \subseteq A$ , where the  $u_i : B \rightarrow A$  are the coordinate forms for the chosen basis. It is immediate that

$$T = \mathcal{V}(K) = \operatorname{Spec}(A/K)$$

has the required property. This completes the proof.

**Examples 6.5.** We give some important examples of functors that reduce to functors  $\prod_{Z/S} Y/Z$  of the type considered in 6.4, and for which criteria of representability will be useful below. Here  $S$  denotes a prescheme, and  $X, Y, Z, \dots$  denote preschemes over  $S$ .

(a) Suppose given an  $S$ -morphism

$$(x) \quad q : X \longrightarrow \underline{\operatorname{Hom}}_S(Y, Z),$$

(“ $X$  acts on  $Y$  with values in  $Z$ ”), that is, a morphism

$$(xx) \quad r : X \times_S Y \longrightarrow Z.$$

Let  $Z'$  be a subprescheme of  $Z$ . It gives a monomorphism

$$\underline{\operatorname{Hom}}_S(Y, Z') \longrightarrow \underline{\operatorname{Hom}}_S(Y, Z),$$

making the first functor a subfunctor of the second. Let  $X'$  be the inverse image of this subfunctor under (x). Thus  $X'$  is the subfunctor of  $X$  for which  $X'(T)$  is the set of  $x \in X(T)$  such that

$$q(x) : Y_T \longrightarrow Z_T$$

factors through  $Z'_T$ . This functor has the following description. Put  $P = X \times_S Y$ , and let  $P'$  be the inverse image of  $Z'$  under  $r : P \rightarrow Z$ . Then there is an evident isomorphism

$$(xxx) \quad X' \simeq \prod_{P/X} P'/P.$$

It follows that *if  $Y$  is essentially free over  $S$  and  $Z'$  is closed in  $Z$ , then the subfunctor  $X'$  of  $X$  is representable by a closed subprescheme of  $X$ .*

(b) Suppose given two actions of  $X$  on  $Y$  with values in  $Z$ , that is, two morphisms

$$q_1, q_2 : X \rightrightarrows \underline{\operatorname{Hom}}_S(Y, Z),$$

and put  $X' = \operatorname{Ker}(q_1, q_2)$ . This is the subfunctor of  $X$  for which  $X'(T)$  is the set of  $x \in X(T)$  such that the two morphisms

$$q_1(x), q_2(x) : Y_T \rightrightarrows Z_T$$

are equal. Giving  $q_1, q_2$  is equivalent to giving a morphism

$$q : X \longrightarrow \underline{\operatorname{Hom}}_S(Y, Z \times_S Z),$$

or, equivalently, a morphism

$$r : X \times_S Y \longrightarrow Z \times_S Z.$$

Put  $U = Z \times_S Z$ , and let  $U'$  be the diagonal subprescheme of  $Z \times_S Z$ . Then  $X'$  is the inverse image under  $q$  of the subfunctor

$$\underline{\operatorname{Hom}}_S(Y, U') \longrightarrow \underline{\operatorname{Hom}}_S(Y, U).$$

It can therefore be written in the form (xxx), with  $P = X \times_S Y$  and  $P'$  the inverse image of the diagonal under  $r$ , that is, the kernel of

$$X \times_S Y \xrightarrow[r_2]{r_1} Z.$$

Thus we are in the situation of (a).

Consequently, *if  $Y$  is essentially free over  $S$  and  $Z$  is separated over  $S$ , then the subfunctor  $X'$  of  $X$  is representable by a closed subprescheme of  $X$ .*

(c) Suppose given a morphism

$$q : X \longrightarrow \underline{\mathrm{Hom}}_S(Y, Y);$$

that is, “ $X$  acts on  $Y$ .” Let  $X'$  be the “kernel” of this morphism, namely the subfunctor of  $X$  for which  $X'(T)$  is the set of  $x \in X(T)$  such that  $q(x) : Y_T \rightarrow Y_T$  is the identity. This functor falls under (b), as one sees by introducing a second morphism

$$q' : X \longrightarrow \underline{\mathrm{Hom}}_S(Y, Y)$$

that “makes  $X$  act trivially on  $Y$ .” Hence, *if  $Y$  is essentially free and separated over  $S$ , then the kernel subfunctor of  $q$  is representable by a closed subprescheme of  $X$ .*

(d) Under the hypotheses of (c), consider the subfunctor  $Y'$  of  $Y$  of “invariants under  $X$ .” Thus  $Y'(T)$  is the set of  $y \in Y(T)$  such that the corresponding morphism  $\bar{q}(y) : X_T \rightarrow Y_T$  is the “ $T$ -constant morphism with value  $y$ .” Introducing  $q'$  as in (c), and writing  $\bar{q}$  and  $\bar{q}'$  for the corresponding morphisms,

$$\bar{q}, \bar{q}' : Y \rightrightarrows \underline{\mathrm{Hom}}_S(X, Y),$$

one sees that  $Y' = \mathrm{Ker}(\bar{q}, \bar{q}')$ . It therefore again falls under (b), with the roles of  $X$  and  $Y$  reversed and  $Z = Y$ . Consequently, *if  $X$  is essentially free over  $S$  and  $Y$  is separated over  $S$ , then the subfunctor  $Y'$  of  $Y$  of invariants under  $X$  is representable by a closed subprescheme of  $Y$ .*

(e) Constructions of the type described in the preceding examples occur especially often in group theory. Thus, when a group prescheme  $G$  over  $S$  acts on the  $S$ -prescheme  $X$ ,

$$q : G \longrightarrow \underline{\mathrm{Aut}}_S(X),$$

the kernel of  $q$ , the “subgroup of  $G$  acting trivially,” is a closed subprescheme of  $G$  provided that  $X$  is essentially free and separated over  $S$ , by (c). The subobject  $X^G$  of invariants is a closed subprescheme of  $X$ , provided that  $G$  is essentially free over  $S$  and  $X$  is separated over  $S$ ,<sup>27</sup> by (d).

Let  $Y, Z$  be subpreschemes of  $X$ , and consider the subfunctor  $\underline{\mathrm{Transp}}_G(Y, Z)$  of  $G$ , the “transporter of  $Y$  into  $Z$ .” Its points with values in an  $S$ -prescheme  $T$  are the  $g \in G(T)$  such that the corresponding automorphism of  $X_T$  satisfies  $g(Y_T) \subseteq Z_T$ , that is, induces a morphism  $Y_T \rightarrow Z_T$  factoring through  $Z_T$ . Hence, *if  $Y$  is essentially free over  $S$ , and  $Z$  is closed in  $X$ , then  $\underline{\mathrm{Transp}}_G(Y, Z)$  is a closed subprescheme of  $G$ , by (a).*

One may also consider the *strict transporter* of  $Y$  into  $Z$ ,<sup>28</sup> whose  $T$ -valued points are the  $g \in G(T)$  such that  $g(Y_T) = Z_T$ . It is precisely

$$\underline{\mathrm{Transp}}_G(Y, Z) \cap \sigma(\underline{\mathrm{Transp}}_G(Z, Y)),$$

<sup>27</sup>Editor’s note: We have corrected  $S_X$  to  $S$ .

<sup>28</sup>Editor’s note: It is denoted  $\underline{\mathrm{Transp}}_G^{\mathrm{str}}(Y, Z)$ .

where  $\sigma$  is the symmetry of  $G$ . Consequently, if  $Y$  and  $Z$  are essentially free over  $S$  and closed in  $X$ , then the strict transporter of  $Y$  into  $Z$  is a closed subprescheme of  $G$ .

An important case is  $X = G$ , with  $G$  acting on itself by inner automorphisms. If  $H$  is a subprescheme of  $G$ , the strict transporter of  $H$  into itself is also called the *normalizer of  $H$  in  $G$* , and is denoted  $\underline{\text{Norm}}_G H$ . Hence, if  $H$  is a closed subgroup prescheme of  $G$ , essentially free over  $S$ , then  $\underline{\text{Norm}}_G H$  is representable by a closed subgroup prescheme of  $G$ .

Finally, let  $Z$  be a subprescheme of  $G$ . Its *centralizer*  $\underline{\text{Centr}}_G(Z)$  in  $G$  is the group subfunctor of  $G$  defined by the procedure in (d), regarding “ $Z$  acts on  $G$ ” through the action induced by that of  $G$ . Thus, if  $Z$  is essentially free over  $S$  and  $G$  is separated over  $S$ , then  $\underline{\text{Centr}}_G(Z)$  is a closed subgroup prescheme of  $G$ . In particular, if  $G$  is essentially free and separated over  $S$ , then the center  $C$  of  $G$ , which is none other than  $\underline{\text{Centr}}_G(G)$ , is a closed subgroup prescheme of  $G$ .

When  $S$  is the spectrum of a field, 6.2(b) [the source prints 6.3(b)] shows that in examples (a)–(e) above the “essentially free” hypotheses are automatically satisfied; only separation hypotheses remain. Recalling that a group prescheme over a field is necessarily separated, one obtains, for example:

**Corollary 6.7.** <sup>29</sup> Let  $G$  be a group prescheme over a field  $k$ . Then:

- For every subprescheme  $Z$  of  $G$ , the centralizer of  $Z$  in  $G$  is a closed subgroup prescheme of  $G$ . In particular, this holds for the center  $\underline{\text{Centr}}_G(G)$  of  $G$ .
- More generally, if  $u, v : X \rightarrow G$  are morphisms of preschemes, then  $\underline{\text{Transp}}_G(u, v)$  is representable by a closed subprescheme of  $G$ .
- For two subpreschemes  $Y, Z$  of  $G$ , with  $Z$  closed,  $\underline{\text{Transp}}_G(Y, Z)$  is a closed subprescheme of  $G$ . If  $Y$  is also closed, the same conclusion holds for  $\underline{\text{Transp}}_G^{\text{str}}(Y, Z)$ .
- For every subgroup prescheme<sup>30</sup>  $H$  of  $G$ , the normalizer  $\underline{\text{Norm}}_G(H)$  is a closed subgroup prescheme of  $G$ .

**Remark 6.8.** <sup>31</sup> Let  $A$  be a commutative ring, let  $M$  be an  $A$ -module, and put

$$M^\vee = \text{Hom}_A(M, A).$$

Endow the  $A$ -module  $\text{End}_A(M)$  with the topology of discrete pointwise convergence. Thus a basis of neighborhoods of zero is formed by the following  $A$ -submodules, where  $n \in \mathbb{N}$  and  $m_1, \dots, m_n \in M$ :

$$K(m_1, \dots, m_n) = \{u \in \text{End}_A(M) \mid u(m_i) = 0 \text{ for } i = 1, \dots, n\}.$$

We say that  $M$  is a *quasi-free  $A$ -module* if the image of the canonical morphism

$$\Theta : M \otimes_A M^\vee \longrightarrow \text{End}_A(M)$$

contains  $\text{id}_M$  in its closure; that is, if the following condition holds:

<sup>29</sup>Editor’s note: We have retained the numbering of the original; there is no 6.6.

<sup>30</sup>Editor’s note: Indeed, over a field  $k$ , every subgroup prescheme of  $G$  is closed; cf. Exposé VI<sub>A</sub>, 0.6.1.

<sup>31</sup>Editor’s note: We have expanded the original in what follows; in particular, we have added Lemma 6.8.1.

$$(*) \quad \begin{cases} \text{for all } m_1, \dots, m_n \in M, \text{ there exist } x_1, \dots, x_r \in M \text{ and } f_1, \dots, f_r \in M^\vee \\ \text{such that } m_i = \Theta \left( \sum_{s=1}^r x_s \otimes f_s \right) (m_i) = \sum_{s=1}^r f_s(m_i) x_s \quad \text{for } i = 1, \dots, n. \end{cases}$$

(In this case  $\text{Im } \Theta$  is dense in  $\text{End}_A(M)$ , since for every  $u \in \text{End}_A(M)$  one has  $u(m_i) = \sum_{s=1}^r f_s(m_i) u(x_s) = \Theta(\sum_{s=1}^r u(x_s) \otimes f_s)(m_i)$ .)

First note that this property is stable under base change. Indeed, let  $\phi : A \rightarrow A'$  be a ring homomorphism, put  $M' = M \otimes_A A'$ , and let  $m'_1, \dots, m'_n \in M'$ . Write

$$m'_i = \sum_j m_{ij} \otimes b_{ij}, \quad m_{ij} \in M, \quad b_{ij} \in A'.$$

By hypothesis there exist  $x_1, \dots, x_r \in M$  and  $f_1, \dots, f_r \in \text{Hom}_A(M, A)$  such that

$$m_{ij} = \sum_s x_s f_s(m_{ij}) \quad \text{for all } i, j.$$

Write  $\phi \circ f_s$  for the image of  $f_s$  in

$$\text{Hom}_A(M, A') = \text{Hom}_{A'}(M', A').$$

Then, for  $i = 1, \dots, n$ ,

$$\begin{aligned} \left( \sum_s x_s \otimes \phi \circ f_s \right) (m'_i) &= \sum_{s,j} x_s \otimes \phi(f_s(m_{ij})) b_{ij} \\ &= \sum_j \left( \sum_s x_s f_s(m_{ij}) \right) \otimes b_{ij} \\ &= m'_i. \end{aligned}$$

This proves that  $M'$  is quasi-free over  $A'$ .

Also, every projective  $A$ -module  $P$  is quasi-free. Indeed, there are  $A$ -morphisms

$$A^{(I)} \xrightarrow{\pi} P \xrightarrow{\tau} A^{(I)} \quad \text{such that} \quad \pi \circ \tau = \text{id}_P.$$

Let  $(e_i)$  be the canonical basis of  $A^{(I)}$ , and put  $f_i = e_i^* \circ \tau$ , a linear form on  $P$ . Given  $m_1, \dots, m_n \in P$ , there is a finite subset  $J$  of  $I$  such that

$$m_k = \sum_{i \in J} f_i(m_k) \pi(e_i) \quad \text{for } k = 1, \dots, n.$$

Theorem 6.4 remains valid if, in the statement of Definition 6.1, “free” is replaced by “projective” or, more generally, by “quasi-free.” Proceeding as in the proof of 6.4, one reduces to the following lemma.

**Lemma 6.8.1.** *Let  $M$  be a quasi-free  $A$ -module,  $N$  a submodule, and  $F$  the covariant functor*

$$F : (A\text{-algebras}) \longrightarrow \mathbf{Set}$$

*such that  $F(B) = \{\text{pt}\}$  if  $M_B = (M/N)_B$ , and  $F(B) = \emptyset$  otherwise. Then there is an ideal  $K$  of  $A$  such that  $F(B) = \{\text{pt}\}$  if and only if the morphism  $A \rightarrow B$  factors through  $A/K$ ; that is, there is an isomorphism, functorial in  $B$ ,*

$$F(B) \simeq \text{Hom}_{A\text{-alg.}}(A/K, B).$$

*Proof.* Let  $(n_j)$  be a system of generators of  $N$ , and let  $F_j$  be the subfunctor of the terminal functor  $e$ , where  $e(B) = \{\text{pt}\}$  for every  $B$ , corresponding to the submodule  $An_j$ . We have  $F(B) = \{\text{pt}\}$  if and only if the image of every  $n_j$  in  $M_B$  is zero. Hence  $F$  is the intersection of the functors  $F_j$ , reducing us to the case in which  $N$  is generated by one element  $n$ .

Let  $B$  be an  $A$ -algebra, with structure map  $\phi : A \rightarrow B$ . If the image  $n \otimes 1$  of  $n$  in  $M_B$  is zero, then for every  $f \in M^\vee = \text{Hom}_A(M, A)$ ,

$$0 = f(n) \otimes 1 = \phi(f(n)).$$

Conversely, because  $M$  is quasi-free, there exist  $x_1, \dots, x_r \in M$  and  $f_1, \dots, f_r \in M^\vee$  such that

$$n = \sum_s f_s(n) x_s,$$

whence

$$n \otimes 1 = \sum_s x_s \otimes \phi(f_s(n)).$$

It follows that  $n \otimes 1 = 0$  if and only if  $\phi$  factors through  $A/K$ , where  $K$  is the ideal generated by the  $f_s(n)$ , for  $s = 1, \dots, r$ . This proves the lemma.

## 7. Appendix: On Monomorphisms of Group Preschemes

The result proved in this section is not needed in the remainder of the seminar, except in X, 8.8 and Exposés XV and XVI. It depends essentially on the existence theorem for quotient groups in Exposé VI<sub>A</sub>.<sup>32</sup>

Corollary 5.7 raises the question of when a monomorphism  $u : G \rightarrow H$  of  $S$ -groups can be asserted to be an immersion, or even a closed immersion.

We saw in VI<sub>B</sub>, 1.4.2 that this is so when  $S$  is the spectrum of a field  $k$ , provided that  $G$  is of finite type over  $k$  and  $H$  is locally of finite type over  $k$ . It follows readily that the same result remains valid when  $S$  is merely assumed Artinian.<sup>33</sup>

On the other hand, examples of bijective monomorphisms that are not immersions are easy to give, for instance over the affine line over a field or over the spectrum of a discrete valuation ring. Take

$$H = (\mathbf{Z}/2\mathbf{Z})_S, \quad G = G_1 \times_S G_2,$$

where  $G_1$  is the open subgroup of  $H$  complementary to the closed point  $x \neq 0$  in the fiber  $H_s \simeq \mathbf{Z}/2\mathbf{Z}$ , with  $s$  a fixed closed point of  $S$ , and  $G_2$  is the closed subscheme of  $H$  which is the union of the subscheme given by the identity section and the reduced closed subscheme defined by  $x$ . One checks immediately that  $G_2$  is stable under multiplication in  $H$ , and hence is a group scheme. The immersions  $G_i \rightarrow H$ ,  $i = 1, 2$ , then define a homomorphism of  $S$ -groups

$$G = G_1 \times_S G_2 \longrightarrow H$$

which is visibly a bijective monomorphism, and even a local immersion, but is not an immersion. Indeed,  $G$  and  $H$  are reduced;  $G$  has three disjoint irreducible components, whereas  $H$  has only two.

Notice also that  $G_1 \rightarrow H$  is an example of an open subgroup of  $H$  that is not closed, in contrast to the situation for algebraic groups over a field. The theory of degenerations

<sup>32</sup>Editor's note: cf. Exposé VI<sub>A</sub>, Theorems 3.2 and 3.3.2.

<sup>33</sup>Editor's note: the additions made in Exposé VI<sub>B</sub> should be taken into account.

of elliptic curves gives further examples of this phenomenon in which  $H$  is smooth over  $S$  of relative dimension one and  $G$  has connected fibers.

One might hope<sup>(\*)</sup> that once  $G$  is assumed flat over  $S$ , and, say,  $G$  and  $H$  are of finite presentation over  $S$ , every monomorphism  $u : G \rightarrow H$  of  $S$ -groups is automatically an immersion. We shall prove a result of this kind under additional hypotheses.

First, we may assume that  $S$  is affine and, by the finite-presentation hypothesis on  $G$  and  $H$ , that  $S$  is Noetherian: one reduces to the spectrum of a ring of finite type over  $\mathbf{Z}$ .<sup>34</sup> Then  $G$  and  $H$  are Noetherian. Saying that  $u : G \rightarrow H$  is a closed immersion (respectively, an immersion) is then equivalent to saying that  $u$  is a monomorphism—which holds by hypothesis—and is proper (respectively, proper at every point of  $u(G)$ ).<sup>35</sup>

The valuative criterion for properness shows that it is enough to check, after every base change  $S' \rightarrow S$  with  $S'$  the spectrum of a discrete valuation ring, complete if desired, that the morphism  $u' : G' \rightarrow H'$  has the corresponding properness property. The case of local properness was omitted from EGA II, 7.3 and was to appear as an erratum in EGA IV.<sup>(\*)</sup> We are therefore reduced to the case in which  $S$  itself is the spectrum of a complete discrete valuation ring, subject to the proviso that any additional hypotheses imposed on  $G, H, u$  be stable under base change.

Let  $s$ , respectively  $s'$ , be the closed, respectively generic, point of  $S$ . The induced homomorphisms on the fibers,

$$u_s : G_s \longrightarrow H_s, \quad u_{s'} : G_{s'} \longrightarrow H_{s'},$$

are closed immersions because they are monomorphisms of group schemes of finite type over fields (VI<sub>B</sub>, 1.4.2). We may therefore identify  $G_{s'}$  with a closed subscheme of  $H_{s'}$ . We shall use the following result.

**Lemma 7.1.** *Let  $S$  be the spectrum of a discrete valuation ring, with generic point  $s'$ . Let  $H$  be an  $S$ -prescheme and let  $L'$  be a closed subprescheme of the generic fiber  $H_{s'}$ , hence also a subprescheme of  $H$ . Its schematic closure  $L$  in  $H$ —the smallest closed subprescheme of  $H$  containing  $L'$ , cf. EGA I, 9.5—exists. It is also the unique closed subprescheme of  $H$ , flat over  $S$ , whose generic fiber is  $L'$ . Formation of  $L$  is functorial in the pair  $(H, L')$  and commutes with fiber products over  $S$ . In particular, if  $H$  is an  $S$ -group and  $L'$  is a subgroup of  $H_{s'}$ , then  $L$  is a subgroup of  $H$ .*

The proof is immediate and is left to the reader.<sup>(\*\*)</sup> Apply this to the pair  $(H, G_{s'})$ . The monomorphism  $u : G \rightarrow H$  factors as

$$G \xrightarrow{u'} L \longrightarrow H,$$

where  $L$  is a subgroup of  $H$ , a closed subprescheme flat over  $S$ , and  $u' : G \rightarrow L$  induces an isomorphism on generic fibers. Thus  $u : G \rightarrow H$  is an immersion (respectively, a closed immersion) if and only if  $u' : G \rightarrow L$  is one. We are reduced to the case in which  $H$  is flat over  $S$  and  $u_{s'} : G_{s'} \rightarrow H_{s'}$  is an isomorphism, again subject to the requirement that any additional hypotheses remain valid when  $H$  is replaced by a closed subgroup prescheme.

Since  $H$  is then the schematic closure of  $H_{s'}$ , if  $u$  is an immersion its image  $u(G)$  is a subprescheme<sup>36</sup> of  $H$  containing  $H_{s'}$ . Its schematic closure is therefore  $H$ , so  $u(G)$  is an

<sup>(\*)</sup>In fact, M. Raynaud constructed a counterexample with  $G$  smooth and with connected fibers; cf. XVI, 1.1(c). Without the connected fiber hypothesis, one may take  $S = \operatorname{Spec}(\mathbf{Z}_2)$ , let  $G = (\mathbf{Z}/2\mathbf{Z})_S$  with its nonidentity closed point removed, and let  $H = (\mu_2)_S$ .

<sup>34</sup>Editor's note: cf. EGA IV<sub>3</sub>, §8 and Exposé VI<sub>B</sub>, §10.

<sup>35</sup>Editor's note: cf. EGA IV<sub>3</sub>, 8.11.5 and 15.7.1.

<sup>(\*)</sup>Cf. EGA IV<sub>3</sub>, 15.7.

<sup>(\*\*)</sup>Cf. EGA IV<sub>2</sub>, 2.8.

<sup>36</sup>Editor's note: the word "closed" has been removed.



open subprescheme of  $H$ . We must consequently prove that  $u$  is an open immersion, or an isomorphism if the desired conclusion is that it is a closed immersion.

Because  $G$  and  $H$  are flat over  $S$ , this is equivalent to the induced maps on the fibers being open immersions, respectively isomorphisms (cf. SGA 1, I, 5.7). It already holds for  $u_{s'}$ , so it remains to prove that  $u_s$  is an open immersion, respectively an isomorphism.

Since  $G$  and  $H$  are flat over  $S$ , their fiber dimensions are constant (VI<sub>B</sub>, 4.3). Their generic fibers agree, hence these dimensions are the same for  $G$  and  $H$ . Thus  $u_s : G_s \rightarrow H_s$  is a monomorphism of algebraic groups of the same dimension. It follows that  $G_s$  is open in  $H_s$ , and set-theoretically is a union of connected components of  $H_s$ . Indeed, after passing to an algebraically closed base field and reducing  $G$  and  $H$ , both are smooth over the field and the assertion is immediate.

The complement  $H_s - G_s$  is closed in  $H_s$ , hence in  $H$ . Its complement  $H'$  in  $H$  is open and is stable under the group law. Upon replacing  $H$  by  $H'$ , we may therefore reduce, with the usual proviso, to the case in which  $u$  is bijective, that is,  $u_s : G_s \rightarrow H_s$  is bijective. In every case we are reduced to proving that  $u$ , equivalently  $u_s$ , is an isomorphism, possibly under this bijectivity hypothesis.

Assume first that  $u$  is bijective. If  $H_s$  is reduced, then  $u_s$  is an isomorphism, because  $G_s$  identifies with a closed subscheme of  $H_s$  having the same underlying set. In particular, if  $k = \kappa(s)$  has characteristic zero, every algebraic group over  $k$  is reduced by Cartier's theorem (cf. VI<sub>B</sub>, 1.6.1; VII<sub>B</sub>, 3.3.1; or EGA IV<sub>4</sub>, 16.12.2 and 17.12.5). We have proved:

**Proposition 7.2.** *Let  $u : G \rightarrow H$  be a homomorphism of group preschemes of finite presentation over  $S$ . If  $u$  is a monomorphism,  $G$  is flat over  $S$ , and every residue field of  $S$  has characteristic zero, then  $u$  is an immersion.*

Suppose now that  $\kappa(s)$  has characteristic  $p > 0$ , and restrict to the case in which  $G$  is commutative. Under the preceding reductions,  $H$  is also commutative, because it is the schematic closure of  $H_{s'} \simeq G_{s'}$ . For  $n \geq 0$ , set

$$S_n = \operatorname{Spec}(V/\mathfrak{m}^{n+1}), \quad G_n = G \times_S S_n, \quad H_n = H \times_S S_n,$$

where  $V$  is the valuation ring defining  $S$  and  $\mathfrak{m}$  its maximal ideal. For  $m > 0$ , denote by  ${}_mG$  and  ${}_mH$  the kernels of multiplication by  $m$  in  $G$  and  $H$ . Define similarly

$${}_m(G_n) = ({}_mG)_n,$$

which we abbreviate to  ${}_mG_n$ , and likewise for  $H$ .

By VI<sub>A</sub>, 3.2, the quotients

$$Q_n = H_n/G_n$$

exist. Each  $Q_n$  is a commutative group scheme over  $S_n$ , flat over  $S_n$ , and  $H_n \rightarrow Q_n$  is faithfully flat with kernel  $G_n$ . Formation of the quotient commutes with base extension,<sup>37</sup> so

$$Q_n \simeq Q_m \times_{S_m} S_n \quad (m \geq n).$$

In particular,  $Q_n \times_{S_n} S_0 = Q_0 = H_0/G_0$ . Since  $G_0$  and  $H_0$  have the same underlying set,  $Q_0$  is set-theoretically a single point: it is a purely infinitesimal group.

It follows that every  $Q_n$  is finite and flat over  $S_n$ . Let it be defined by an algebra  $C_n$  over  $V_n = V/\mathfrak{m}^{n+1}$ , finite free as a  $V_n$ -module. For  $m \geq n$ ,

$$C_n = C_m \otimes_{V_m} V_n,$$

<sup>37</sup>Editor's note: this should be made explicit in Exposés V and VI<sub>A</sub>.

compatibly also with the comultiplication maps. Hence

$$C = \varprojlim C_n \quad \text{is a finite free module over} \quad V = \varprojlim V_n,$$

and the comultiplications on the  $C_n$  induce one on  $C$ . Thus  $Q = \operatorname{Spec}(C)$  is a finite flat group scheme over  $S$  satisfying

$$Q \times_S S_n = Q_n$$

for every  $n$ .

**Lemma 7.3.** <sup>38</sup> *Let  $K$  be a field and let  $Q$  be a finite group scheme over  $K$ , of degree  $n$ . Then the  $n$ -th power morphism on  $Q$  is zero.*

See VII<sub>A</sub>, 8.5.

**Remark 7.3.1.** The statement of Lemma 7.3 still makes sense for a finite locally free group scheme  $Q/S$  over an arbitrary base prescheme  $S$ . It would be interesting to find a proof in that generality.

Lemma 7.3 (that is, VII<sub>A</sub>, 8.5) does show that the proposed statement holds whenever  $S$  is reduced: apply it to the fibers of  $Q$  at the maximal points of  $S$ , equivalently the generic points of its irreducible components.<sup>39</sup>

In the preceding situation, where  $S$  is the spectrum of a discrete valuation ring and  $Q$  is commutative, it follows that  $n \operatorname{id}_Q = 0$ . Here  $n$  is a power  $p^{\nu_0}$  of the residue characteristic. Thus:

**Corollary 7.4.** *With  $Q$  and the  $Q_n$  as above, suppose that their common degree is  $p^{\nu_0}$ . Then*

$$p^{\nu_0} \operatorname{id}_Q = 0, \quad p^{\nu_0} \operatorname{id}_{Q_n} = 0 \quad \text{for every } n.$$

**Corollary 7.5.** *Suppose in addition that  $G_0$  is smooth over  $k$  and that  $p \operatorname{id}_{G_0}$  is flat. Equivalently, by the structure theory of algebraic groups over an algebraically closed field  $\bar{k}$ ,  $G_{\bar{k}}$  contains no subgroup isomorphic to the additive group. Then, for every  $\nu \geq \nu_0$  and every  $n > 0$ , the group  $p^\nu H_n$  is flat over  $S_n$ .*

Indeed,  $p^{\nu_0} \operatorname{id}_{Q_n} = 0$  implies that  $p^\nu \operatorname{id}_{H_n}$  factors through  $G_n$ , giving the commutative diagram

$$\begin{array}{ccc} H_n & \xrightarrow{p^\nu \operatorname{id}_{H_n}} & H_n \\ \uparrow u_n & \searrow v_n & \uparrow u_n \\ G_n & \xrightarrow{p^\nu \operatorname{id}_{G_n}} & G_n. \end{array}$$

We claim that  $v_n$  is flat. Since  $H_n$  and  $G_n$  are flat over  $S_n$ , it is enough to check that  $v_0$  is flat (SGA 1, IV, 5.9). Thus we may take  $n = 0$ , so that  $S_n = \operatorname{Spec}(k)$ . Since  $p \operatorname{id}_{G_0}$ , and hence  $p^\nu \operatorname{id}_{G_0}$ , is flat, its image is an induced open subgroup  $G'_0$  of  $G_0$ . The map  $u_0 : G_0 \rightarrow H_0$  is surjective, so  $v_0$  takes its values in  $G'_0$  and may be regarded as a homomorphism to  $G'_0$ . Its composite with  $u_0$  is an epimorphism; hence  $v_0$  itself is an epimorphism, and therefore a flat homomorphism into  $G'_0$ , and thus into  $G_0$ . Consequently  $v_n$  is flat and  $\ker v_n = p^\nu H_n$  <sup>†</sup> is flat over  $S_n$ .  $\square$

<sup>38</sup>Editor's note: the hypothesis that  $Q$  be commutative has been removed here, and 7.3.1 modified accordingly. If  $Q$  is not commutative, the  $n$ -th power morphism need not be a group homomorphism.

<sup>39</sup>Editor's note: this should be added to Exposé VII<sub>A</sub>. On the other hand, the statement is true for every  $S$  when  $Q$  is commutative; cf. Deligne's theorem in [TO70], p. 4.

<sup>†</sup>English editor's note: the source prints  $p^\nu H$  over  $S$  here and  $p^\nu H$  over  $S_n$  in the proof of Lemma 7.7. The subscripts  $n$  are supplied in both places because those arguments concern the base-changed family over  $S_n$ .

**Remark.** We did not explicitly use the assumption that  $G_0$  is smooth over  $k$ . It is, however, an easy consequence of the flatness of  $\mathrm{pid}_{G_0}$  [the source prints  $\mathrm{pid}_G$ ]; this is why the condition was stated explicitly in Corollary 7.5.

**Lemma 7.6.** *Let  $u : G \rightarrow H$  be a surjective monomorphism of commutative algebraic groups over a field  $k$  of characteristic  $p > 0$ , and put  $Q = H/G$ , a purely infinitesimal group. There is an integer  $\nu_1$  such that, for every  $\nu \geq \nu_1$ , the sequence*

$$0 \longrightarrow p^\nu G \longrightarrow p^\nu H \longrightarrow Q \longrightarrow 0$$

*is exact.*

It is enough to establish exactness at  $Q$ , which in turn amounts to the injectivity of the homomorphism

$$(x) \quad \mathcal{O}_{Q,e} \longrightarrow \mathcal{O}_{p^\nu H,e}$$

of local rings at the identity elements. Recall that  $Q$  is set-theoretically supported at the single point  $e$ . There is a natural homomorphism

$$(xx) \quad \mathcal{O}_{p^\nu H,e} \longrightarrow \mathcal{O}_{H,e}/\mathfrak{m}^{p^\nu},$$

where  $\mathfrak{m}$  is the augmentation ideal, equivalently the maximal ideal, of  $\mathcal{O}_{H,e}$ . Indeed,  $p^\nu \mathrm{id}_H$  vanishes on the kernel of the iterated Frobenius homomorphism  $F^\nu$ . The composite of (x) and (xx) is the natural composite

$$(xxx) \quad \mathcal{O}_{Q,e} \longrightarrow \mathcal{O}_{H,e} \longrightarrow \mathcal{O}_{H,e}/\mathfrak{m}^{p^\nu}.$$

Now  $\mathcal{O}_{Q,e} \rightarrow \mathcal{O}_{H,e}$  is injective, because  $H \rightarrow Q$  is an epimorphism and therefore flat. Moreover,  $\mathcal{O}_{Q,e}$  is Artinian, while  $\bigcap_\nu \mathfrak{m}^{p^\nu} = 0$  in  $\mathcal{O}_{H,e}$ , so the intersection of the corresponding ideals in  $\mathcal{O}_{Q,e}$  is also zero. One of them is therefore zero. Thus (xxx) is injective, and a fortiori so is (x).

**Lemma 7.7.** *Under the hypotheses of 7.5, there is an integer  $\nu_1$  such that, for every  $\nu \geq \nu_1$  and every  $n$ , the sequence of  $S_n$ -groups*

$$0 \longrightarrow p^\nu G_n \longrightarrow p^\nu H_n \xrightarrow{w_n} Q_n \longrightarrow 0$$

*is exact. More precisely,  $w_n$  is faithfully flat and its kernel is  $p^\nu G_n$ .*

Take  $\nu_1$  as in Lemma 7.6, applied to  $u_0 : G_0 \rightarrow H_0$ , and require  $\nu_1 \geq \nu_0$ , where  $p^{\nu_0} = \mathrm{rank} Q_0$ . It remains only to check that  $w_n$  is faithfully flat. By Corollary 7.5,  $p^\nu H_n$  is flat over  $S_n$ , as is  $Q_n$ ; it is therefore enough to check that

$$w_0 : p^\nu H_0 \longrightarrow Q_0$$

is faithfully flat, equivalently an epimorphism. This follows from the choice of  $\nu_1$ .

**Corollary 7.8.** *Suppose in addition that  $H$  is separated over  $S$ , or more generally that  $p^\nu H$  is separated over  $S$  for every  $\nu$ , and that  $p^\nu G$  is finite over  $S$  for every  $\nu$ . Then, for  $\nu \geq \nu_1$ , the group schemes  $p^\nu G$  and  $p^\nu H$  are finite and flat over  $S$ , and there is an exact sequence*

$$0 \longrightarrow p^\nu G \longrightarrow p^\nu H \xrightarrow{w} Q \longrightarrow 0.$$

Because  $u : G \rightarrow H$  is surjective, the induced morphism  $p^\nu G \rightarrow p^\nu H$  is also surjective. The source is finite over  $S$  and the target is separated over  $S$ , hence the target is finite over  $S$ . To prove that  $p^\nu G$  and  $p^\nu H$  are flat over  $S$ , it is enough to prove this after every base

change to  $S_n$ , where it is contained in Corollary 7.5. Finally, the exact sequence above is obtained from the exact sequences in Lemma 7.7 as  $n$  varies.

Taking generic fibers in the sequence of Corollary 7.8, and recalling that  $u_{s'} : G_{s'} \rightarrow H_{s'}$  is an isomorphism, gives that  $Q_{s'}$  is the unit group. Since  $Q$  is flat over  $S$ ,  $Q$  itself is the unit group; therefore so is  $Q_s$ , and  $u_s : G_s \rightarrow H_s$  is an isomorphism. We have proved:

**Proposition 7.9.** *Let  $u : G \rightarrow H$  be a homomorphism of group preschemes of finite presentation over a prescheme  $S$ . Assume:*

- (a)  *$u$  is a monomorphism;*
- (b)  *$G$  is flat over  $S$ ;*
- (c) *for every  $s \in S$  whose residue field has characteristic  $p > 0$ , let  $S' = \operatorname{Spec}(\mathcal{O}_{S,s})$ , and let  $u' : G' \rightarrow H'$  be obtained from  $u$  by base change to  $S'$ . Then  $G'$  is commutative, the special fiber  $G'_0 = G_s$  is smooth over  $\kappa(s)$ , and, for every  $\nu > 0$ , the group  ${}_{p^\nu}G'$  is finite over  $S'$  while  ${}_{p^\nu}H'$  is separated over  $S'$ .*

*Under these assumptions,  $u$  is an immersion.*

Indeed, the finiteness of  ${}_pG'$  over  $S'$  implies the finiteness of its special fiber over  $\kappa(s)$ . Hence  $G'_0 \otimes_{\kappa(s)} \overline{\kappa(s)}$  contains no subgroup isomorphic to the additive group, and the hypotheses of Corollary 7.5 apply.

**Remark 7.10.** Examples of M. Raynaud (XVI, 1.1(a),(b)) show that in condition (c) one cannot omit either the finiteness of the  ${}_{p^\nu}G'$  over  $S'$  or the separatedness of the  ${}_{p^\nu}H'$  over  $S'$ .

We now seek conditions under which  $u$  is a closed immersion. Retain the hypotheses preceding Proposition 7.9, but no longer suppose that  $u$  is surjective; assume only that it is an isomorphism on generic fibers. We already know that  $u : G \rightarrow H$  is an open immersion. The same is therefore true of  $u_0 : G_0 \rightarrow H_0$ , whose image contains the identity component  $(H_0)^0$ . Since  $G_0$  is smooth over  $k$  and has no additive component after extending to  $\bar{k}$ , the same is true of  $H_0$ . We use:

**Lemma 7.11.** *Let  $H$  be a commutative algebraic group over a field  $k$ , and suppose that  $H \otimes_k \bar{k}$  contains no subgroup isomorphic to  $\mathbf{G}_a$ . If  $n = \deg(H/H^0)$ , then*

$${}_nH \longrightarrow H/H^0$$

*is surjective.*

We may assume that  $k$  is algebraically closed, in which case this follows from the familiar fact that  $H^0(k)$  is a divisible group.

Return to  $u : G \rightarrow H$ , and suppose that the  ${}_nG$ ,  $n > 0$ , are proper over  $S$ , while the  ${}_nH$  are separated over  $S$ . The  ${}_nH$  are also flat over  $S$ . Indeed, it is enough to check flatness at the points above  $s$ , and hence to check that  $n \operatorname{id}_{H_0}$  is flat. As noted above, this is equivalent to  $(H_0)^0$  being smooth<sup>40</sup> over  $k$  and having no  $\mathbf{G}_a$ -component over  $\bar{k}$ . Since  $(H_0)^0 = (G_0)^0$ , this follows from the corresponding hypothesis on  $G_0$ .

Because  $H$  is separated over  $S$ , so are the  ${}_nH$ . The morphism  ${}_nG \rightarrow {}_nH$  is proper and therefore has closed image. Its image contains the generic fiber of  ${}_nH$ , because  $u_{s'} : G_{s'} \rightarrow H_{s'}$  is an isomorphism. Since  ${}_nH$  is flat over  $S$ , it is the schematic closure of its generic fiber. Hence  ${}_nG \rightarrow {}_nH$ , and therefore  ${}_nG_0 \rightarrow {}_nH_0$ , is surjective. Together with

<sup>40</sup>Editor's note: the terminology has been updated, replacing "simple" by "smooth"; see, for example, A. Grothendieck's note (1) in SGA 1, Exposé II.

$G_0 \supset (H_0)^0$ , Lemma 7.11 now implies that  $G_0 \rightarrow H_0$  is surjective. Hence  $u$  is surjective. We obtain:

**Proposition 7.12.** *With the notation of 7.9, suppose that conditions (a) and (b) hold, together with the stronger condition (c') obtained from (c) by requiring, for every integer  $n > 0$ , not only those of the form  $p^\nu$ , that  ${}_nG'$  be finite over  $S'$  and  ${}_nH'$  separated over  $S'$ . Then  $u$  is a closed immersion.*

**Remark 7.13.**

- (a) The separatedness hypothesis on the  ${}_nH$  easily implies that  $H$  itself is separated over  $S$ . This is also formally contained in Proposition 7.12, by taking  $G$  to be the unit  $S$ -group. If  $S$  is locally Noetherian, in Proposition 7.9 it is enough to assume that  $H$  is locally of finite type over  $S$ , rather than of finite type; the proof applies verbatim. For Proposition 7.12, one also assumes that the fibers of  $H$  are of finite type.
- (b) Using Proposition 7.12, it is not difficult to prove that if  $u : G \rightarrow H$  is a monomorphism of group preschemes of finite presentation over  $S$ ,  $G$  is diagonalizable—or, more generally, of multiplicative type—and  $H$  is separated over  $S$ , then  $u$  is a closed immersion. This is a satisfactory generalization of the first conclusion in 5.7. If  $G$  is smooth over  $S$ , apply Proposition 7.12 directly; the general case reduces readily to this one. If  $H$  is not assumed separated over  $S$ , one can still show that  $u$  is an immersion; for a torus  $G$ , this also follows from the material that comes next.
- (c) If  $G$  is assumed to have connected fibers in Proposition 7.9, the separatedness of  ${}_{p^\nu}H'$  over  $S'$  may be omitted from condition (c). Indeed, after the reductions in the proof,  $H$  may be assumed flat over  $S$  and  $u$  bijective. Thus  $H$  has connected fibers, and Raynaud's theorem (VI<sub>B</sub>, 5.5) implies that  $H$  is separated over  $S$ .<sup>41</sup>

## Bibliography

42

- [RG71] M. Raynaud and L. Gruson, *Critères de platitude et de projectivité*, Invent. Math. **13** (1971), 1–89.
- [TO70] J. Tate and F. Oort, *Group schemes of prime order*, Ann. Sci. Éc. Norm. Supér. (4) **3** (1970), 1–21.

---

<sup>41</sup>Editor's note: the attribution, which appeared in SGAD 1965, should be made explicit in Exposé VI<sub>B</sub>, §5, together with the changes between SGAD and Lecture Notes 151.

<sup>42</sup>Editor's note: additional references cited in this exposé.

# Exposé IX

## Groups of Multiplicative Type: Homomorphisms into a Group Scheme

by A. Grothendieck<sup>0</sup>

### 1. Definitions

**Definition 1.1.** — *Let  $S$  be a prescheme and let  $G$  be a group prescheme over  $S$ . The group  $G$  is said to be of multiplicative type if it is locally diagonalizable for the faithfully flat quasi-compact topology (cf. IV 6.3); that is, if for every  $s \in S$ , there are an open neighborhood  $U$  of  $s$  and a faithfully flat quasi-compact morphism  $S' \rightarrow U$  such that*

$$G' = G \times_S S'$$

*is a diagonalizable  $S'$ -group (I 4.4 and VIII 1.1).*

*The group  $G$  is said to be of quasi-isotrivial multiplicative type if it is locally diagonalizable even for the étale topology (IV 6.3); that is, if in the preceding definition one can take  $S' \rightarrow U$  to be étale and surjective. Equivalently—as one sees by taking the disjoint union of the schemes  $S'$  attached to the various  $s \in S$ —there is an étale surjective morphism  $S' \rightarrow S$  such that  $G' = G \times_S S'$  is a locally diagonalizable  $S'$ -group.*

*If one can moreover choose  $S' \rightarrow S$  to be finite, étale, and surjective, then  $G$  is said to be of isotrivial multiplicative type.*

*Finally,  $G$  is said to be a group of locally trivial multiplicative type (respectively, of locally isotrivial multiplicative type) if every  $s \in S$  has an open neighborhood  $U$  such that  $G \times_S U$  is a diagonalizable  $U$ -group (respectively, is of isotrivial multiplicative type; in other words, there is a finite étale surjective morphism  $S' \rightarrow U$  such that  $G \times_S S'$  is a diagonalizable  $S'$ -group).*


**Comments 1.2.** — The five preceding notions are all obtained from the notion of a diagonalizable group by localization, with respect to five different “topologies” associated with (Sch).

It is understood in general that, when the word “locally” is not qualified by naming the topology under consideration, the Zariski topology is meant, in accordance with established terminology. In the terminology introduced here, “of locally trivial multiplicative

---

<sup>0</sup>Editor’s note: version 1.0 of November 8, 2009. Additions were made in the proof of 3.6 bis, in 4.4–7, 5.0–6, 6.1, 7.1, and 8.2. Sections 8.1 and 4.5 remain to be revised.

type” is equivalent to “locally diagonalizable” in VIII 1.1; likewise, “trivial” translates as “diagonalizable.” The five notions introduced above fit into the following implication diagram, which follows from the corresponding relations among the topologies involved:



For practical purposes, let us point out at once that every group of multiplicative type that we shall encounter will be quasi-isotrivial. Thus we shall see in the next exposé (X 4.5) that, when  $G$  is of finite type over  $S$ , it is automatically quasi-isotrivial; we shall nevertheless give examples in which it is not locally isotrivial. We shall likewise see that  $G$  may be locally trivial without being isotrivial, and hence *a fortiori* without being trivial. This readily implies that the inclusions in the preceding diagram are strict.

On the other hand, we shall also see there (X 5.16) that, when  $S$  is locally noetherian and normal—or, more generally, geometrically unibranch—every group of multiplicative type and finite type over  $S$  is necessarily isotrivial; moreover, it is trivial as soon as it is locally trivial (or even merely trivial over a dense open subset). This explains why most groups of multiplicative type encountered in practice will doubtless be isotrivial, all the more so because we shall later see that the maximal tori of semisimple group schemes are automatically isotrivial.

**Definition 1.3.** — *Let  $S$  and  $G$  be as in 1.1. The group  $G$  is called a torus if, for the faithfully flat quasi-compact topology, it is locally isomorphic to a group of the form  $\mathbf{G}_m^r$ , where  $r$  is an integer with  $r \geq 0$ .*

With the notation of 1.1, this says that  $S'$  may be chosen so that  $G'$  is isomorphic to a group of the form  $(\mathbf{G}_{m,S'})^r$ . The integer  $r$  depends on  $s \in S$ : it is the dimension of the fiber

$$G_s = G \otimes_S \kappa(s).$$

It is a locally constant function of  $s$ , as is seen at once. This observation should be generalized as follows.

**Definition 1.4.** — *Let  $G$  be a diagonalizable group scheme over a field  $k$ . Thus  $G$  is isomorphic to a group  $D_k(M)$ , where  $M$  is an ordinary commutative group determined up to isomorphism by this condition—more precisely,*

$$M \simeq \text{Hom}_{k\text{-gr}}(G, \mathbf{G}_{m,k})$$

(VIII 1.3). *The isomorphism class of  $M$  is called the type of the diagonalizable group  $G$ ; it is plainly invariant under extension of the base field.*

Now let  $G$  be a group scheme of multiplicative type over an arbitrary prescheme  $S$ . For every  $s \in S$ , there is an extension  $k$  of  $\kappa(s)$  such that  $G \times_S \text{Spec}(k)$  is a diagonalizable  $k$ -group.<sup>1</sup> Its type is independent of the chosen extension by the preceding observation and

<sup>1</sup>Editor’s note: indeed, by hypothesis there are an open neighborhood  $U$  of  $s$  and a faithfully flat quasi-compact morphism  $S' \rightarrow U$  such that  $G_{S'}$  is diagonalizable. For every  $s' \in S'$  above  $s$ , the field  $\kappa(s')$  is an extension of  $\kappa(s)$ , and  $G_{s'} = G \times_S \text{Spec}(\kappa(s'))$  is diagonalizable.

is called the type of  $G$  at  $s$ , or the type of  $G_s$ . In particular, if  $G$  itself is diagonalizable, and hence isomorphic to a group of the form  $D_S(M)$ , then for every  $s \in S$  the type of  $G$  at  $s$  is the isomorphism class of  $M$ .

More generally, if  $G$  is a group of multiplicative type over  $S$  and  $M$  is an ordinary commutative group, then  $G$  is said to be of type  $M$  if it is of type  $M$  at every point  $s \in S$ ; equivalently, if  $G$  is locally isomorphic to  $D_S(M)$  for the faithfully flat quasi-compact topology.

**Remark 1.4.1.** <sup>2</sup> a) For a group  $G$  of multiplicative type over  $S$ , the function

$$s \longmapsto \text{the type of } G_s$$

is locally constant on  $S$ . Indeed, with the notation of 1.1, if  $G'$  is of type  $M$ , then  $G$  is of type  $M$  over the neighborhood  $U$  of  $s$ . Consequently  $S$  has a canonical decomposition as a disjoint union of preschemes  $S_i$  such that, for every  $i$ ,  $G_{S_i}$  is of type  $M_i$ , where the commutative groups  $M_i$  are pairwise nonisomorphic.

b) In particular, if  $S$  is connected, the type of the fibers of  $G$  is constant; that is, there is a commutative group  $M$  such that  $G$  is of type  $M$ . Finally, if  $G$  is a torus, the type of  $G$  at  $s$  is determined by the integer  $\dim G_s$ : indeed,  $G_s$  is of type  $\mathbb{Z}^n$ , where  $n = \dim G_s$ .

**Remark 1.5.** — a) It follows immediately from Definitions 1.1 and 1.3 that these notions are “compatible with base extension.” Thus, if  $G$  is a group prescheme over  $S$  and  $S' \rightarrow S$  is a base-change morphism, then whenever  $G$  is of multiplicative type (respectively, of isotrivial multiplicative type, and so on), the same is true of  $G'$ .

b) If, moreover,  $S' \rightarrow S$  is faithfully flat and quasi-compact, then  $G'$  being of multiplicative type, respectively a torus, implies the same for  $G$ . If  $S' \rightarrow S$  is also étale (respectively, finite étale), then  $G'$  being quasi-isotrivial (respectively, isotrivial) implies the same for  $G$ .

c) Finally, return to an arbitrary base-change morphism  $f : S' \rightarrow S$ . If  $s' \in S'$  and  $s = f(s')$ , then the type of  $G'$  at  $s'$  is the same as that of  $G$  at  $s$ , since

$$G'_{s'} = G_s \otimes_{\kappa(s)} \kappa(s').$$

## 2. Extension of Certain Properties of Diagonalizable Groups to Groups of Multiplicative Type

The extensions in question are essentially trivial consequences of the results of the preceding Exposé, in view of the definitions in 1.1 and the “local” nature (for the faithfully flat and quasi-compact topology) of the results under consideration.

**Definition 2.0.** <sup>3</sup> We denote by  $\mathcal{M}$  the set of faithfully flat and quasi-compact morphisms.

**Proposition 2.1.** Let  $G$  be a group of multiplicative type over a prescheme  $S$ . Then:

- (a)  $G$  is faithfully flat over  $S$ , and affine over  $S$  (a fortiori, quasi-compact over  $S$ );
- (b)  $G$  is of finite type over  $S$  if and only if  $G$  is of finite presentation over  $S$ , if and only if, for every  $s \in S$ , the type of  $G$  at  $s$  is given by a commutative group of finite type;

<sup>2</sup>Editor’s note: the number 1.4.1 has been added to the remarks that follow, for use in later references.

<sup>3</sup>Editor’s note: This definition has been added; it occurs in Propositions 2.3 and 2.7.



- (c)  $G$  is finite over  $S$  if and only if, for every  $s \in S$ , the type of  $G$  at  $s$  is given by a finite commutative group, if and only if (when  $S$  is quasi-compact)  $G$  is of finite type over  $S$  and is annihilated by an integer  $n > 0$ ;
- (c')  $G$  is integral over  $S$  if and only if, for every  $s \in S$ , the type of  $G$  at  $s$  is given by a torsion commutative group;
- (d)  $G$  is the unit group if and only if, for every  $s \in S$ , the type of  $G$  at  $s$  is given by the zero commutative group;
- (e)  $G$  is smooth over  $S$  if and only if, for every  $s \in S$ , the type of  $G$  at  $s$  is given by a commutative group of finite type whose torsion subgroup has order prime to the characteristic of  $\kappa(s)$ .

These statements follow from VIII 2.1, taking into account that the properties in question descend under faithfully flat and quasi-compact morphisms (cf. SGA 1, VIII, or EGA IV<sub>2</sub>, §2).

Using VIII 3.5, one likewise obtains:

**Proposition 2.2.** *Let  $G$  be a group of multiplicative type and of finite type over  $S$ . For every integer  $n \neq 0$ , the kernel  ${}_nG = \ker(n \cdot \text{id}_G)$  is a group of multiplicative type, finite over  $S$ .*

**Proposition 2.3.** *Let  $G$  be a group of multiplicative type over the prescheme  $S$ , acting freely on the right on an  $S$ -prescheme  $X$  that is affine over  $S$ . Then:*

- (a) *the equivalence relation defined by  $G$  in  $X$  is  $\mathcal{M}$ -effective (IV 3.4), and  $Y = X/G$  is affine over  $S$ ;*
- (b) *if, moreover,  $X$  is of finite presentation (respectively, of finite type) over  $S$ , then the same is true of  $Y$ .*

The first assertion follows from VIII 5.1, which treats the case in which  $G$  is diagonalizable, and from IV 3.5.2, which allows one to reduce to that case. Indeed, faithfully flat and quasi-compact morphisms  $S' \rightarrow S$  are effective descent morphisms for the fibered category of affine schemes over schemes: if  $Y'$  is affine over  $S'$  and is equipped with a descent datum relative to  $S' \rightarrow S$ , then this datum is effective, that is,  $Y'$  comes from a scheme  $Y$  affine over  $S$  (cf. SGA 1, VIII 2.1).

For the second assertion, one is likewise reduced to the diagonalizable case of VIII 5.8, since the finiteness conditions under consideration descend under faithfully flat and quasi-compact morphisms (SGA 1, VIII 3.3 and 3.6<sup>4</sup>). Proceeding as in VIII, Corollaries 5.5–5.7, one deduces from 2.3:

**Corollary 2.4.** *Under the hypotheses of 2.3, the graph morphism*

$$X \times_S G \longrightarrow X \times_S X$$

*is a closed immersion. For every section  $\sigma$  of  $X$  over  $S$ , the corresponding morphism*

$$G \longrightarrow X, \quad g \longmapsto \sigma \cdot g,$$

*is a closed immersion.*

In particular:

---

<sup>4</sup>Editor's note: See also V, 9.1, or EGA IV<sub>2</sub>, 2.7.1.

**Corollary 2.5.** *Let  $u : G \rightarrow H$  be a monomorphism of  $S$ -preschemes in groups, with  $G$  of multiplicative type and  $H$  affine over  $S$ . Then  $u$  is a closed immersion,  $H/G = Y$  exists and is affine over  $S$ , and  $H$  is a principal homogeneous bundle over  $Y$  under  $G_Y$ .*

**Remark 2.6.** Let  $u : G \rightarrow H$  be a monomorphism of  $S$ -preschemes in groups, where  $G$  is of multiplicative type and of finite type over  $S$ , and  $H$  is of finite presentation and separated over  $S$ . One can then show that  $u$  is a closed immersion by using VIII 7.12 (see Remark VIII 7.13(b)).

**Proposition 2.7.** *Let  $u : G \rightarrow H$  be a homomorphism of  $S$ -groups of multiplicative type, with  $H$  of finite type over  $S$ . Then:*

- (i)  $G_0 = \ker u$  is an  $S$ -group of multiplicative type; the equivalence relation defined by  $G_0$  in  $G$  is  $\mathcal{M}$ -effective, and hence (IV 3.4)  $u$  factors as

$$G \longrightarrow G/G_0 = I \longrightarrow H,$$

where  $I \rightarrow H$  is a monomorphism<sup>5</sup> of  $S$ -groups. The group  $I$  is of multiplicative type, the equivalence relation in  $H$  defined by  $I$  is  $\mathcal{M}$ -effective, and consequently  $H_0 = H/I$  exists and is also of multiplicative type;

- (ii)  $H_0$  and  $I$  are of finite type, and so is  $G_0$  when  $G$  is.

*Proof.* Proceeding as for 2.3, one is reduced to the case in which  $G$  and  $H$  are diagonalizable; then 2.7 reduces to VIII 3.4.

We record the following corollaries.

**Corollary 2.8.**

- (a) *Let  $S$  be a prescheme. The category of  $S$ -groups of multiplicative type and of finite type is an abelian category.*
- (b) *Let  $u : G \rightarrow H$  be a homomorphism in this category. It is a monomorphism (respectively, an epimorphism, respectively, an isomorphism) in this category if and only if  $u$  is a monomorphism of preschemes (respectively,  $u$  is faithfully flat, respectively,  $u$  is an isomorphism of preschemes).*

It is enough to note that the product of two  $S$ -groups of multiplicative type is again of multiplicative type (which is immediate), and is plainly of finite type over  $S$  if both factors are. The rest of 2.8 follows immediately from 2.7; the details are left to the reader.

**Corollary 2.9.** *Let  $u : G \rightarrow H$  be a homomorphism of  $S$ -groups of multiplicative type and of finite type. Let  $U$  be the set of points  $s \in S$  for which  $u_s : G_s \rightarrow H_s$  is a monomorphism (respectively, faithfully flat, respectively, an isomorphism). Then  $U$  is both open and closed, and the induced homomorphism  $u_U : G_U \rightarrow H_U$  is a monomorphism (respectively, faithfully flat, respectively, an isomorphism).*

Let  $P$  (respectively,  $Q$ ) be the kernel (respectively, cokernel) of  $u$ . By 2.7,  $Q$  exists and  $P$  and  $Q$  are of multiplicative type; moreover, formation of  $P$  and  $Q$  commutes with every base change  $S' \rightarrow S$ , in particular with the formation of fibers. On the other hand,  $u$  is a monomorphism (respectively, faithfully flat, respectively, an isomorphism) if and only if  $P = 0$  (respectively,  $Q = 0$ , respectively,  $P = Q = 0$ ). These remarks reduce us to checking the following assertion: if  $R$  is an  $S$ -group of multiplicative type, then the set  $U$

<sup>5</sup>Editor's note: Indeed, even a closed immersion.

of points  $s \in S$  for which  $R_s = 0$  is open and closed, and  $R_U = 0$ . This is contained in Remark 1.4.1(a).

**Corollary 2.10.** *Let  $G$  be an  $S$ -group of multiplicative type and of finite type, and let  $H$  and  $H'$  be two subgroups of multiplicative type.*

- (a)  $H'' = H \cap H'$  is a subgroup of  $G$  of multiplicative type.
- (b) Let  $U$  be the set of  $s \in S$  for which  $H_s \subseteq H'_s$  (respectively,  $H_s = H'_s$ ). Then  $U$  is open and closed, and  $H_U \subseteq H'_U$  (respectively,  $H_U = H'_U$ ).

Here, of course, the intersection sign in  $H \cap H'$  denotes the intersection in the functorial sense, namely  $H \times_G H'$ , which is plainly an  $S$ -subgroup of  $G$ . Likewise, the inclusion sign (respectively, the equality sign) denotes the order relation (respectively, equality) between subfunctors of  $H$ , not inclusion (respectively, equality) of the underlying sets.

Applying 2.7 first to the inclusion morphism  $H \rightarrow G$ , one finds that  $H$  is of finite type; the same holds for  $H'$ . It then follows from 2.8 that  $H \cap H'$  is of multiplicative type and of finite type. (Note that the canonical functor from the category considered in 2.8 to the category **(Sch)**/ $S$  commutes with finite inverse limits.)

Formation of  $H \cap H'$  commutes with base extension, in particular with formation of the fibers. Moreover,  $H \subseteq H'$  (respectively,  $H = H'$ ) is equivalent to  $H'' = H$  (respectively, to  $H'' = H$  and  $H'' = H'$ ). Consider the canonical homomorphisms  $H'' \rightarrow H$  and  $H'' \rightarrow H'$ . Then  $U$  is the set of points  $s \in S$  for which the induced homomorphism  $H''_s \rightarrow H_s$  is an isomorphism (respectively, for which  $H''_s \rightarrow H_s$  and  $H''_s \rightarrow H'_s$  are isomorphisms).

By 2.9, it follows that  $U$  is open and closed, and that  $H''_U \rightarrow H_U$  (respectively,  $H''_U \rightarrow H_U$  and  $H''_U \rightarrow H'_U$ ) are isomorphisms. This gives the desired conclusion.

**Proposition 2.11.** *Let  $S$  be a prescheme, let  $G$  be an  $S$ -group of multiplicative type and of finite type over  $S$ , let  $H$  be a subgroup of  $G$  of multiplicative type, and put  $K = G/H$ , a quotient group of  $G$  of multiplicative type.*

- (i) Suppose  $G$  is trivial, that is, diagonalizable. There is a partition of  $S$  into open and closed subsets  $S_i$  such that, for every  $i$ ,  $H_{S_i}$  and  $K_{S_i}$  are diagonalizable. In particular, if  $S$  is connected,  $H$  and  $K$  are diagonalizable.
- (ii) The same assertion holds with “diagonalizable” replaced by “isotrivial,” provided that  $S$  is connected, locally connected, or quasi-compact.
- (iii) If  $G$  is locally trivial (respectively, locally isotrivial, respectively, quasi-isotrivial), then the same is true of  $K$  and  $H$ .

*Proof.* (i) By hypothesis,  $G = D_S(M)$ , where  $M$  is a commutative group of finite type. For each quotient group  $M_i$  of  $M$ , let  $H_i = D_S(M_i)$  be the corresponding diagonalizable subgroup of  $G$  (VIII 3.1). Let  $S_i$  be the set of points  $s \in S$  for which  $H_s = (H_i)_s$ . By 2.10,  $S_i$  is open and closed, and  $H_{S_i} = (H_i)_{S_i}$ ; thus  $H_{S_i}$  is diagonalizable, and hence so is  $K_{S_i}$  (VIII 3.1). The  $S_i$  are plainly pairwise disjoint, and we claim that they cover  $S$ . Indeed, for every  $s$ , the group  $H_s$ , being a subgroup of the diagonalizable group  $G_s$ , is diagonalizable (cf. 8.1 below<sup>6</sup>). Hence  $H_s$  has the form  $D_{\kappa(s)}(M_i)$ , by VIII 1.5 and 3.2(b). Restricting to the family of nonempty  $S_i$ , assertion (i) follows.

(ii) By hypothesis, there exists a finite surjective étale morphism  $S' \rightarrow S$  such that  $G_{S'}$  is diagonalizable. Every point of  $S'$  therefore has an open and closed neighborhood  $U'$

<sup>6</sup>Editor’s note: The erroneous reference to IV 4.7.5 has been corrected. Note that Section 8 of the present Exposé is independent of Sections 3–7.

such that  $H_{U'}$  and  $K_{U'}$  are diagonalizable. The image  $U$  of  $U'$  in  $S$  is open and closed, and  $S' \rightarrow S$  still induces a finite surjective étale morphism  $U' \rightarrow U$ . Thus every point  $s \in S$  has an open and closed neighborhood  $U$  such that  $H_U$  and  $K_U$  are isotrivial. Assertion (ii) follows at once.

(iii) This follows at once from (i), (ii), and the definitions. Moreover, the quasi-isotrivial case will also follow from the more general fact “finite type  $\implies$  quasi-isotrivial,” announced in 1.2 (cf. X 4.5).

### 3. Infinitesimal Properties: Lifting and Conjugation Theorems

The fundamental infinitesimal properties of groups of multiplicative type follow from the following theorem.

**Theorem 3.1.** *Let  $S$  be an affine scheme, let  $H$  be an  $S$ -group of multiplicative type, and let  $\mathcal{F}$  be a quasi-coherent sheaf on  $S$  on which  $H$  acts (I 4.7). Then*

$$H^i(H, \mathcal{F}) = 0 \quad \text{for } i > 0,$$

where  $H^i$  denotes the Hochschild cohomology studied in Exposé I, §5.

Indeed, by loc. cit., 5.3, Hochschild cohomology is expressed, in terms of the affine rings  $A$  and  $B$  of  $S$  and  $G$ , and the  $A$ -module  $M$  defining  $\mathcal{F}$ , as the cohomology of a complex of  $A$ -modules  $C^\bullet(H, M)$ , whose formation commutes with every base change  $A \rightarrow A'$ . Consequently, for a base change  $A \rightarrow A'$  with  $A'$  flat over  $A$ , one has

$$H^i(H', \mathcal{F}') = H^i(H, \mathcal{F}) \otimes_A A'.$$

If  $A'$  is moreover faithfully flat over  $A$ , it is therefore enough, in order to prove that  $H^i(H, \mathcal{F})$  vanishes, to prove that  $H^i(H', \mathcal{F}')$  vanishes. This observation reduces us to checking 3.1 in the case where  $G$  is diagonalizable; in that case it was proved in I 5.3.3.

Using the results of Exposé III, we shall derive from 3.1 several consequences of a geometric nature.

**Theorem 3.2.** *Let  $S$  be an affine scheme, let  $S_0$  be an affine subscheme defined by an ideal  $J$ , let  $H$  be a group of multiplicative type over  $S$ , and let  $G$  be an arbitrary group prescheme over  $S$ . Let  $u, v : H \rightarrow G$  be two group homomorphisms, and let  $u_0, v_0 : H_0 \rightarrow G_0$  be the homomorphisms obtained from  $u, v$  by the base change  $S_0 \rightarrow S$ . If  $J^2 = 0$  and  $u_0 = v_0$ , then there exists  $g \in G(S)$  such that*

$$v = \text{int}(g) \circ u \quad \text{and} \quad g_0 = e.$$

This follows from III 2.1(ii) and 3.1 (the vanishing of  $H^1$ ).

**Corollary 3.3.** *Under the preliminary hypotheses of 3.2, suppose in addition that  $G$  is smooth over  $S$ , but assume only that  $J$  is nilpotent (not necessarily square-zero). If there exists  $g_0 \in G_0(S_0)$  such that  $v_0 = \text{int}(g_0) \circ u_0$ , then  $g_0$  lifts to an element  $g \in G(S)$  such that  $v = \text{int}(g) \circ u$ .*

By induction on an integer  $n$  such that  $J^n = 0$ , one is reduced to the case in which  $J$  is square-zero. Since  $G$  is smooth over  $S$ , one may lift  $g_0$  to an element  $g'$  of  $G(S)$ . Replacing  $v$  by  $v' = \text{int}(g') \circ u$ , one is then reduced to the situation of 3.2.

**Corollary 3.4.** *Under the preliminary hypotheses of 3.2, with  $J$  nilpotent, suppose in addition that  $u$  is central (for example, that  $G$  is commutative). Then  $u_0 = v_0$  implies  $u = v$ .*

Indeed, the reduction to the case in which  $J$  is square-zero is again immediate, and then it is enough to apply 3.2. In particular:

**Corollary 3.5.** *Let  $S, S_0, H, G$  be as in 3.2, with  $J$  nilpotent. Let  $u : H \rightarrow G$  be a homomorphism of  $S$ -groups such that  $u_0 : H_0 \rightarrow G_0$  is the unit homomorphism. Then  $u$  is the unit homomorphism.*

**Theorem 3.6.** *Let  $S$  be an affine scheme, let  $S_0$  be a subscheme defined by a nilpotent ideal  $J$ , let  $H$  be a group of multiplicative type over  $S$ , let  $G$  be a group prescheme smooth over  $S$ , and let  $u_0 : H_0 \rightarrow G_0$  be a homomorphism of the  $S_0$ -groups obtained by base change  $S_0 \rightarrow S$ . Then there exists a homomorphism  $u : H \rightarrow G$  of  $S$ -groups that lifts  $u_0$ . Moreover, by 3.3, any two such liftings  $u, u'$  are conjugate by an element of  $G(S)$  whose reduction in  $G(S_0)$  is the identity element.*

This follows from III 2.1 and 2.3 and from 3.1: the vanishing of  $H^2$  gives the existence of the lifting of  $u_0$ , and the vanishing of  $H^1$  gives uniqueness up to conjugation.

One may prove in the same way the following variants of 3.2–3.6. In fact, these variants imply the preceding statements when applied to the graph subgroups of the homomorphisms considered there.

**Theorem 3.2 bis.** *Let  $S$  be an affine scheme, let  $S_0$  be a subscheme defined by an ideal  $J$ , let  $G$  be an  $S$ -group prescheme, and let  $H, H'$  be subgroup preschemes of multiplicative type. Let  $G_0, H_0, H'_0$  be the groups over  $S_0$  obtained by base change  $S_0 \rightarrow S$ . If  $J^2 = 0$  and  $H_0 = H'_0$ , then there exists  $g \in G(S)$  such that*

$$H' = \text{int}(g)(H) \quad \text{and} \quad g_0 = e.$$

**Corollary 3.3 bis.** *Under the preliminary hypotheses of 3.2 bis, suppose in addition that  $G$  is smooth over  $S$ , and that  $J$  is nilpotent (not necessarily square-zero). If  $g_0 \in G_0(S_0)$  satisfies  $H'_0 = \text{int}(g_0)(H_0)$ , then  $g_0$  lifts to an element  $g \in G(S)$  such that  $H' = \text{int}(g)(H)$ .*

**Corollary 3.4 bis.** *Under the preliminary hypotheses of 3.2 bis, suppose that  $J$  is nilpotent and that  $H$  is central (for example, that  $G$  is commutative). Then  $H_0 = H'_0$  implies  $H = H'$ .*

**Theorem 3.6 bis.** *Let  $S$  be an affine scheme, let  $S_0$  be a subscheme defined by a nilpotent ideal  $J$ , let  $G$  be an  $S$ -group prescheme smooth over  $S$ , and let  $H_0$  be a subgroup prescheme of  $G_0 = G \times_S S_0$ , of multiplicative type. Then:*

- (a) *there exists a subgroup prescheme  $H$  of  $G$ , flat over  $S$ , such that  $H \times_S S_0 = H_0$ ;*
- (b) *such an  $H$  is necessarily of multiplicative type;*
- (c) *finally, by 3.2 bis, any two such liftings  $H, H'$  of  $H_0$  are conjugate by an element  $g \in G(S)$  whose reduction in  $G(S_0)$  is the identity element.*

Let us indicate how one can prove 3.2 bis (of which 3.3 bis and 3.4 bis are immediate consequences) and assertion (a) of 3.6 bis. The latter follows from 3.1 (the vanishing of  $H^2$ ) and III 4.37(ii). Similarly, 3.2 bis follows from 3.1 (the vanishing of  $H^1$ ) and III 4.37(i), at least when  $G$  and  $H$  are smooth over  $S$ .

Using the results of the following Exposé, however, one can very simply deduce 3.2 bis and 3.6 bis, in the general form stated here, from 3.2 and 3.6, respectively. For 3.2 bis, it is enough to note that, by X 2.1,  $H$  and  $H'$  are isomorphic, which reduces us to 3.2.

For 3.6 bis, note that  $H_0$  is necessarily of finite type over  $S_0$ :<sup>7</sup> indeed, as a subgroup prescheme of  $G_0$ , which is smooth over  $S_0$ , the group  $H_0$  is locally of finite type over  $S_0$ . But  $H_0$  is affine over  $S_0$  by 2.1(a), and is therefore of finite type over  $S_0$ . By X 4.5,  $H_0$  is consequently quasi-isotrivial, and hence comes, by X 2.1, from a group  $H$  of multiplicative type over  $S$ .<sup>8</sup> By 3.6, there exists a homomorphism of  $S$ -groups  $u : H \rightarrow G$  lifting the immersion  $H_0 \hookrightarrow G_0$ . Since  $H$  and  $H_0$  (respectively,  $G$  and  $G_0$ ) have the same underlying topological space, and since, for every  $h \in H$ , the morphism

$$\mathcal{O}_{G,u(h)} \longrightarrow \mathcal{O}_{H,h}$$

is surjective (because it is so after reduction modulo the nilpotent ideal  $J$ ), the morphism  $u$  is also an immersion (cf. EGA I, 4.2.2).

Finally, every lifting  $H$  of  $H_0$  to a flat subgroup of  $G$  is necessarily of multiplicative type by X 2.3,<sup>9</sup> which also proves assertion (b) of 3.6 bis. The reader may verify that the results 3.2 bis–3.6 bis are not used in Exposé X, so there is no vicious circle.

**Proposition 3.8.** <sup>10</sup> *Let  $S$  be a prescheme, let  $S_0$  be a closed subscheme defined by a locally nilpotent ideal  $J$ , let  $G$  be an  $S$ -group of multiplicative type, and let  $X$  be an  $S$ -prescheme locally of finite presentation. Suppose that  $G$  acts on  $X$  in such a way that  $G_0 = G \times_S S_0$  acts trivially on  $X_0 = X \times_S S_0$ . Then  $G$  acts trivially on  $X$ .*

The proof is the proof of III 2.4(b). One may also reduce to loc. cit. by observing that 3.8 immediately reduces to the case in which  $G$  is diagonalizable, which is the case treated there.

**Corollary 3.9.** *Let  $S, S_0$  be as above, and let  $u : G \rightarrow H$  be a homomorphism of  $S$ -groups, with  $G$  of multiplicative type and  $H$  locally of finite presentation over  $S$ . Suppose that the homomorphism  $u_0 : G_0 \rightarrow H_0$  obtained by base change  $S_0 \rightarrow S$  is central. Then  $u$  is central.*

Indeed, it is enough to apply 3.8 with  $X = H$ , letting  $G$  act on  $H$  by

$$(g, h) \longmapsto \text{int}(u(g)) \cdot h = u(g)h u(g)^{-1}.$$

## 4. The Density Theorem

This theorem (cf. 4.7 below)\*, together with the theorem<sup>11</sup> of Section 7, will be the essential tool in the present exposé and the two following ones for passing from the infinitesimal properties of groups of multiplicative type developed above to their “finite” properties.

**Definition 4.1.** *Let  $X$  be a prescheme. A family  $(Z_i)_{i \in I}$  of subpreschemes of  $X$  is said to be schematically dense if, for every open subset  $U$  of  $X$  and every closed subprescheme  $U_0$  of  $U$  containing all the  $Z_i \cap U$ , one has  $U_0 = U$ .*

*A subprescheme  $Z$  of  $X$  is said to be schematically dense in  $X$  if the one-member family consisting of  $Z$  is schematically dense.*

<sup>7</sup>Editor’s note: The original said, “by 2.1(b), since its fibers are.” The editors did not understand this argument and have replaced it by the argument that follows.

<sup>8</sup>Editor’s note: The sentence that follows has been expanded.

<sup>9</sup>Editor’s note: X 2.3 depends essentially on Theorem 3.6 of the present Exposé.

<sup>10</sup>Editor’s note: The numbering of the original has been retained; there is no 3.7.

\*All the results from 4.1 through 4.6 are contained in EGA IV<sub>3</sub>, 11.10, to which we refer the reader for a systematic account of the notion of schematic density.

<sup>11</sup>Editor’s note: the “algebraicity” theorem 7.1.

It is immediate (cf. EGA IV<sub>3</sub>, 11.10.1) that the definition is equivalent to saying that, for every open subset  $U$  of  $X$ , every section  $f$  of  $\mathcal{O}_U$  that vanishes on each  $Z_i \cap U$  is zero. Equivalently, the intersection of the kernels of the canonical homomorphisms

$$u_i : \mathcal{O}_X \longrightarrow (v_i)_*(\mathcal{O}_{Z_i})$$

is zero, where  $v_i : Z_i \rightarrow X$  is the canonical immersion.<sup>12</sup> When  $X$  lies over a prescheme  $S$ , this is also equivalent to the following assertion: for every open subset  $U$  of  $X$ , every  $S$ -prescheme  $Y$  separated over  $S$ , and every pair of morphisms  $u, v : U \rightarrow Y$  that coincide on the  $Z_i \cap U$ , one has  $u = v$ . Indeed,  $u = v$  is equivalent to  $U_0 = U$ , where  $U_0$  is the inverse image of the diagonal in  $Y \times_S Y$  under the  $S$ -morphism  $U \rightarrow Y \times_S Y$  defined by  $(u, v)$ . The diagonal is a closed subprescheme of  $Y \times_S Y$ , so  $U_0$  is a closed subprescheme of  $U$  containing all the  $Z_i \cap U$ . Thus schematic density gives  $U_0 = U$ , hence  $u = v$ . Conversely, take  $Y = \operatorname{Spec} \mathcal{O}_S[T]$ .

In the terminology introduced at the end of EGA I, 9.5, saying that the subprescheme  $Z$  is schematically dense in  $X$  also means that  $X$  is the *closure subprescheme* of  $Z$  in  $X$ .

**Lemma 4.2.** *Let  $X$  be a flat  $S$ -prescheme, and let  $(Z_i)_{i \in I}$  be a family of subpreschemes of  $X$ , flat over  $S$ . Let  $S_0$  be a subprescheme of  $S$  defined by a nilpotent ideal  $\mathcal{J}$ , and suppose that the modules  $\mathcal{J}^n / \mathcal{J}^{n+1}$  are locally free over  $S_0$ . Let  $X_0, Z_{i0}$  be obtained from  $X, Z_i$  by the base change  $S_0 \rightarrow S$ . If  $(Z_{i0})_{i \in I}$  is schematically dense in  $X_0$ , then  $(Z_i)_{i \in I}$  is schematically dense in  $X$ .*

*Proof.* Suppose  $\mathcal{J}^{n+1} = 0$ , where  $n \geq 0$ , and argue by induction on  $n$ ; the assertion is trivial for  $n = 0$ . Define  $S_m, X_m, Z_{im}$  by reduction modulo  $\mathcal{J}^{m+1}$ , as usual. The induction hypothesis already implies that  $(Z_{i,n-1})_{i \in I}$  is schematically dense in  $X_{n-1}$ . After replacing  $X$  by an induced open subset, it is enough to prove that every section  $f$  of  $\mathcal{O}_X$  that vanishes on all the  $Z_i$  is zero.

The section  $f_{n-1}$  of

$$\mathcal{O}_{X_{n-1}} = \mathcal{O}_X / \mathcal{J}^n \mathcal{O}_X$$

defined by  $f$  vanishes on the  $Z_{i,n-1}$ , hence is zero. Consequently,  $f$  is a section of  $\mathcal{J}^n \mathcal{O}_X$ . Since  $X$  is flat over  $S$ , there is an isomorphism

$$\mathcal{J}^n \mathcal{O}_X \xleftarrow{\sim} \mathcal{E} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{X_0}, \quad \mathcal{E} = \mathcal{J}^n = \mathcal{J}^n / \mathcal{J}^{n+1}.$$

Similarly, since each  $Z_i$  is flat over  $S$ , the restriction  $f_i$  of  $f$  to  $Z_i$  may be regarded as a section of

$$\mathcal{J}^n \mathcal{O}_{Z_i} \xleftarrow{\sim} \mathcal{E} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{Z_{i0}}.$$

By hypothesis, all the  $f_i$  are zero. The module  $\mathcal{E}$  is locally free by assumption, and therefore so is

$$\mathcal{F} = \mathcal{E} \otimes_{\mathcal{O}_{S_0}} \mathcal{O}_{X_0}.$$

Thus  $f$  is a section of the locally free module  $\mathcal{F}$  on  $X_0$  whose restriction to every  $Z_{i0}$  is zero. Since the family  $(Z_{i0})_{i \in I}$  is schematically dense in  $X_0$ , it follows at once that  $f = 0$ .  $\square$

**Corollary 4.3.** *Let  $X$  be a locally Noetherian  $S$ -prescheme, flat over  $S$ , and let  $(Z_i)_{i \in I}$  be a family of subpreschemes of  $X$ , flat over  $S$ . Suppose that, for every  $s \in S$ , the family of fibers  $(Z_{is})_{i \in I}$  is schematically dense in  $X_s$ . Then  $(Z_i)_{i \in I}$  is schematically dense in  $X$ .*

*Proof.* After replacing  $X$  by an induced open subset, it is enough to prove that every section  $f$  of  $\mathcal{O}_X$  whose restriction to all the  $Z_i$  is zero is itself zero.<sup>13</sup> For this, it suffices to

<sup>12</sup>Editor's note: all the results on schematic density stated in EGA IV<sub>3</sub>, §11.10 follow from the results on "separating families of homomorphisms" proved in *loc. cit.*, §11.9, to which some of the editor's notes below will refer.

<sup>13</sup>Editor's note: the original has been expanded in what follows.

show that the image of  $f$  in  $\mathcal{O}_{X,x}$  is zero for every  $x \in X$ . Let  $\mathfrak{m}_x$  be the maximal ideal of  $\mathcal{O}_{X,x}$ , and let  $\mathfrak{m}_s$  be that of  $\mathcal{O}_{S,s}$ , where  $s$  is the image of  $x$  in  $S$ . Since  $\mathcal{O}_{X,x}$  is Noetherian,

$$\bigcap_{n \in \mathbb{N}} \mathfrak{m}_x^n = 0, \quad \text{and hence, a fortiori,} \quad \bigcap_{n \in \mathbb{N}} \mathcal{O}_{X,x} \mathfrak{m}_s^n = 0.$$

It is therefore enough to show, for every integer  $n$ , that the section induced by  $f$  on

$$X \times_S \operatorname{Spec}(\mathcal{O}_{S,s}/\mathfrak{m}_s^{n+1})$$

is zero. By hypothesis,

$$Z_{is} = Z_i \otimes_{\mathcal{O}_{S,s}} \kappa(s)$$

is a schematically dense family in

$$X_s = X \otimes_{\mathcal{O}_{S,s}} \kappa(s).$$

We are thus reduced to the case in which  $S$  is the spectrum of a local ring whose maximal ideal  $\mathcal{J}$  is nilpotent,  $S_0 = \operatorname{Spec} \kappa(s)$ , and the hypotheses of Lemma 4.2 hold. The conclusion follows.  $\square$

**Lemma 4.4.** *Let  $S$  be a locally Noetherian prescheme, let  $X$  be a locally Noetherian  $S$ -prescheme, flat over  $S$ , and let  $(Z_i)_{i \in I}$  be a family of subpreschemes of  $X$ , flat over  $S$ . Suppose that, for every  $s \in S$ , the family of fibers  $(Z_{is})_{i \in I}$  is schematically dense in  $X_s$ . Then, for every base change  $S' \rightarrow S$ , with  $S'$  not necessarily locally Noetherian, the family  $(Z'_i)_{i \in I}$  is schematically dense in  $X'$ . In other words,  $(Z_i)_{i \in I}$  is universally schematically dense in  $X$  relative to  $S$  (cf. EGA IV<sub>3</sub>, Definition 11.10.8).*

*Proof.*<sup>14</sup> Let  $f'$  be a section of  $\mathcal{O}_{X'}$  over an open subset  $U'$  of  $X'$ , vanishing on the  $Z'_i$ ; we must show that  $f' = 0$ . Fix  $s' \in S'$ . It is enough to prove that  $f'$  vanishes at every point of  $U'$  above  $s'$ . If  $s$  is the image of  $s'$  in  $S$ , then, after replacing  $S, S'$  by the spectra of their local rings at  $s, s'$ , we may suppose that  $S, S'$  are local and that  $s, s'$  are their closed points. We may also suppose that  $X$  is affine, so that

$$\begin{aligned} S &= \operatorname{Spec}(A), & S' &= \operatorname{Spec}(A'), \\ X &= \operatorname{Spec}(B), & X' &= \operatorname{Spec}(B'), & B' &= B \otimes_A A', \end{aligned}$$

where  $A, A'$  are local,  $A \rightarrow A'$  is a local homomorphism, and  $A, B$  are Noetherian. We may further suppose that  $U'$  is affine, of the form  $X'_{g'}$ , with  $g' \in B'$ .

For every  $A$ -subalgebra  $A''$  of  $A'$ , put

$$S'' = \operatorname{Spec}(A'')$$

and let  $X'', Z''_i$  be obtained from  $X, Z_i$  by the base change  $S'' \rightarrow S$ . We obtain the diagram

$$\begin{array}{ccccc} Z^i & \longleftarrow & Z''_n & \longleftarrow & Z^{i'}_n \\ \downarrow & & \downarrow & & \downarrow \\ X & \longleftarrow & X'' & \longleftarrow & X' \\ \downarrow & & \downarrow & & \downarrow \\ S & \longleftarrow & S'' & \longleftarrow & S'. \end{array}$$

<sup>14</sup>Editor's note: for another proof, reducing to the case in which  $S'$  is locally Noetherian and using 4.3 and 4.5 as here, see EGA IV<sub>3</sub>, 11.9.16 and 11.9.12. In the last line of the proof of 11.9.16, replace 11.9.5 by 11.9.12.



In what follows, we restrict to those  $A''$  that are localizations of  $A$ -subalgebras  $A''_1$  of  $A'$ , of finite type over  $A$ , at  $\mathfrak{m}' \cap A''_1$ , where  $\mathfrak{m}'$  is the maximal ideal of  $A'$ . Thus the homomorphisms  $A \rightarrow A'' \rightarrow A'$  are local. These  $A''$  form an increasing filtered family of subalgebras whose union, and hence whose direct limit, is  $A'$ . Likewise,

$$B' = \varinjlim_{A''} (B \otimes_A A'').$$

There therefore exist such an  $A''$  and an element  $g'' \in B'' = B \otimes_A A''$  whose image in  $B'$  is  $g'$ .

By Lemma 4.5 below, the property that  $(Z_{is})_{i \in I}$  be schematically dense in  $X_s$  is preserved by every base change  $S'' \rightarrow S$ .<sup>15</sup> Consequently, because each  $S''$  is locally Noetherian, replacing  $S$  by a suitable  $S''$  and  $X, Z_i$  by  $X'', Z''_i$  preserves the hypotheses of Lemma 4.4.

We may therefore suppose that  $g'$  comes from an element  $g \in B$ . After replacing  $X$  by the open subset  $X_g = U$ , we may suppose that  $U' = X'$ . In the same way, we may suppose that  $f'$  comes from a section  $f$  of  $\mathcal{O}_X$  over  $X$ .

Let  $Y = V(f)$  be the subscheme of  $X$  defined by  $f$ , and put

$$Y_i = Z_i \times_X Y.$$

Thus  $Y_i$  is the closed subscheme  $V(f_i)$  of  $Z_i$ , where  $f_i$  is the section of  $\mathcal{O}_{Z_i}$  induced by  $f$ . Denoting by  $Y', Y'_i$  the  $S'$ -schemes obtained by base change, we have

$$Y' = V(f'), \quad Y'_i = Z'_i \times_{X'} Y' = V(f'_i),$$

and analogous relations hold for  $Y'', Y''_i$ . The assertion that  $f'$  vanishes on the  $Z'_i$  is equivalent to

$$Y'_i = Z'_i \quad \text{for every } i.$$

16

Let  $\mathfrak{m}$  be the maximal ideal of  $A$ . For every  $n \geq 0$ , introduce the closed subscheme

$$S_n = \text{Spec}(A/\mathfrak{m}^{n+1})$$

of  $S$ , and let  $X_n, Y_n, Z_n^i, Y_n^i$  be obtained from  $X, Y, Z_i, Y_i$  by the base change  $S_n \rightarrow S$ . More generally, for every  $S$ -prescheme  $P$ , write  $P_n = P \times_S S_n$ .

For every  $n$  and  $i \in I$ , consider the functor

$$F_n^i = \prod_{Z_n^i/S_n} Y_n^i/Z_n^i : (\mathbf{Sch})_{/S_n}^\circ \longrightarrow \mathbf{Set},$$

defined, for every  $P$  over  $S_n$ , by

$$F_n^i(P) = \Gamma((Y_n^i)_P/(Z_n^i)_P) = \begin{cases} \emptyset, & (Y_n^i)_P \neq (Z_n^i)_P, \\ \{\text{id}\}, & (Y_n^i)_P = (Z_n^i)_P. \end{cases}$$

Since  $S_n$  is local Artinian and  $Z_n^i$  is flat, hence essentially free, over  $S_n$ , VIII, 6.4 shows that each  $F_n^i$  is represented by a closed subprescheme  $T_n^i$  of  $S_n$ .

<sup>15</sup>Editor's note: the preceding sentence has been added, and the original has been expanded in what follows.

<sup>16</sup>Editor's note: the original added "at least at every point  $x'$  of  $X'$  above  $s'$ ," but this restriction seems unnecessary.

The completion  $\widehat{A}$  of the local ring  $A$  is Noetherian because  $A$  is, and it is the projective limit of the rings  $A_n$ .<sup>17</sup> Let  $K_n^i$  be the ideal of

$$A_n = A/\mathfrak{m}^{n+1}$$

defining  $T_n^i$ , and let  $K^i = \varprojlim_n K_n^i$ ; this is an ideal of  $\widehat{A}$ .

For fixed  $i$ , integers  $m \geq n$ , and any  $S_n$ -prescheme  $P$ ,

$$(Y_n^i)_P = Y_i \times_S S_n \times_{S_n} P = Y_i \times_S P = (Y_m^i)_P,$$

and similarly  $(Z_n^i)_P = (Z_m^i)_P$ . It follows that  $F_n^i$  is the restriction  $F_m^i \times_{S_m} S_n$  of  $F_m^i$  to  $(\mathbf{Sch})_{/S_n}$ , and therefore

$$T_n^i = T_m^i \times_{S_m} S_n.$$

We thus have a commutative diagram with exact rows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & K_m^i & \longrightarrow & A_m & \longrightarrow & \mathcal{O}(T_m^i) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & K_n^i & \longrightarrow & A_n & \longrightarrow & \mathcal{O}(T_n^i) \longrightarrow 0. \end{array}$$

All the vertical arrows are surjective. It follows, first, that the projective system  $(K_n^i)_{n \in \mathbb{N}}$  satisfies the Mittag-Leffler condition, so that

$$\varprojlim_n \mathcal{O}(T_n^i) \simeq \widehat{A}/K^i$$

as topological rings (cf. EGA 0<sub>III</sub>, §13.2). Second, the map  $K^i \rightarrow K_n^i$  is surjective (cf. [BEns], III, §7.4, Proposition 5). Hence

$$K_n^i \simeq \frac{K^i + \mathfrak{m}^{n+1}\widehat{A}}{\mathfrak{m}^{n+1}\widehat{A}}, \quad \mathcal{O}(T_n^i) \simeq (\widehat{A}/K^i) \otimes_A (A/\mathfrak{m}^{n+1}).$$

In other words,  $(T_n^i)_n$  is an inductive system of affine Artinian schemes, and its inductive limit

$$T^i = \varinjlim_n T_n^i$$

is a closed formal subscheme of

$$\widehat{S} = \mathrm{Spf}(\widehat{A})$$

(cf. EGA I, §10), whose reduction modulo  $\mathfrak{m}^{n+1}$  is  $T_n^i$ .

Let  $T$  be the closed formal subscheme of  $\widehat{S}$  given by the intersection of the  $T^i$ , that is, defined by

$$K = \sum_i K^i.$$

Since  $\widehat{A}$  is Noetherian, there is a finite subset  $J$  of  $I$  such that

$$K = \sum_{i \in J} K^i, \quad \text{equivalently} \quad T = \bigcap_{i \in J} T^i.$$

<sup>17</sup>Editor's note: the original has been expanded and corrected to show that the projective limit of the rings  $\mathcal{O}(T_n^i)$  defines a closed formal subscheme of  $\mathrm{Spf}(\widehat{A})$ .

For every  $n$ , therefore,

$$K_n = \sum_{i \in J} K_n^i,$$

where  $K_n$  is the image of  $K$  in  $A_n$ .

Recall that  $f_i$  is the image of  $f \in B = \mathcal{O}(X)$  in  $\mathcal{O}(Z_i)$ , and that  $f'_i$  is its image in

$$\mathcal{O}(Z'_i) = \mathcal{O}(Z_i) \otimes_A A'.$$

The relation  $Y'_i = Z'_i$  is equivalent to  $f'_i = 0$ . Now  $\mathcal{O}(Z_i) \otimes_A A'$  is the direct limit of the subalgebras  $\mathcal{O}(Z_i) \otimes_A A''$ , for  $A''$  as above. Consequently, there is an  $A''$  for which  $Y''_i = Z''_i$ . A priori  $A''$  depends on  $i$ , but, because  $J$  is finite, one  $A''$  can be chosen that works for every  $i \in J$ . Put

$$S'' = \text{Spec}(A''), \quad S''_{(n)} = S'' \times_S S_n.$$

18

For every  $i \in J$ , the equality

$$Y''_i \times_{S''} S''_{(n)} = (Y''_i)_{S''_{(n)}} = Z''_i \times_{S''} S''_{(n)} = (Z''_i)_{S''_{(n)}}$$

shows, by the definition of  $T_n^i$ , that  $S''_{(n)} \rightarrow S_n$  factors through  $T_n^i$ . It therefore factors through

$$T_n = \bigcap_{i \in J} T_n^i,$$

and hence through  $T_n^i$  for every  $i \in I$ . Let  $f''$  be the image of  $f$  in  $B'' = B \otimes_A A''$ , and let  $f''_{(n)}$  be its image in

$$B''_{(n)} = B'' / \mathfrak{m}^{n+1} B''.$$

The preceding factorization means that, for every  $n$ , the section  $f''_{(n)}$  vanishes on all the  $Z_i \times_S S''_{(n)}$ .

Because  $A \rightarrow A'' \rightarrow A'$  are local homomorphisms, the image  $s''$  of  $s'$  in  $S''$  is the closed point of  $S''$ , corresponding to the maximal ideal  $\mathfrak{m}''$  of  $A''$ , and its image in  $S$  is  $s$ . *[The source prints  $B''$  here; since  $s'' \in S'' = \text{Spec}(A'')$ , the intended ring is  $A''$ .]* Fix  $n$ , and now let  $S''_n$  denote the  $n$ -th infinitesimal neighborhood of  $s''$  in  $S''$ :

$$S''_n = \text{Spec}(A'' / \mathfrak{m}''^{n+1}).$$

Put

$$B''_n = B'' / \mathfrak{m}''^{n+1} B'', \quad Z_n^{i''} = Z_i \times_S S''_n.$$

*[The source appends  $Z_n^{i''} = \text{Spec}(B''_n)$ ; the displayed equality would identify each  $Z_n^{i''}$  with  $X''_n$ , so the base-change definition is retained here.]* Then the image  $f''_n$  of  $f''_{(n)}$  in  $B''_n$  vanishes on every  $Z_n^{i''}$ . By Lemma 4.5 below, the family of fibers

$$Z_0^{i'''} = Z_n^{i''} \times_{S''_n} \kappa(s'') = Z_{is} \times_{\kappa(s)} \kappa(s'')$$

is schematically dense in

$$X_0''' = X''_n \times_{S''_n} \kappa(s'') = X_s \times_{\kappa(s)} \kappa(s'').$$

---

<sup>18</sup>Editor's note: the original has been expanded and corrected, distinguishing  $S''_{(n)}$  from the subscheme  $S''_n = S''_n(s'')$  introduced below.

Thus  $S''_n, X''_n, (Z''_n)_{i \in I}$ , and

$$S'''_0 = \operatorname{Spec} \kappa(s'')$$

satisfy the hypotheses of Lemma 4.2. It follows that  $f''_n = 0$ , that is,

$$f'' \in \mathfrak{m}''^{n+1} B'' \quad \text{for every } n.$$

Since  $B$  is Noetherian and  $B'' = B \otimes_A A''$  is a localization of a finite-type  $B$ -algebra,  $B''$  is Noetherian. Its local rings are therefore separated for their usual topology. Hence  $f''$  is zero at every point  $x''$  of  $X''$  for which  $\mathfrak{m}'' B''$  is contained in the maximal ideal of  $\mathcal{O}_{X'', x''}$ , that is, at every point of  $X''$  above  $s''$ . A fortiori,  $f'$  is zero at every point of  $X'$  above  $s'$ .  $\square$

**Lemma 4.5.0.**<sup>19</sup> *Let  $X$  be an  $S$ -prescheme, let  $(Z_i)_{i \in I}$  be a family of subpreschemes of  $X$ , and let  $S' \rightarrow S$  be faithfully flat. If  $(Z'_i)_{i \in I}$  is schematically dense in  $X'$ , then  $(Z_i)_{i \in I}$  is schematically dense in  $X$ .*

This follows from EGA IV<sub>3</sub>, 11.10.5(i) and 11.9.10(i). It remains to give the proof of the following lemma.

**Lemma 4.5.** *Let  $X$  be a locally Noetherian prescheme<sup>20</sup> over a field  $k$ , let  $(Z_i)_{i \in I}$  be a family of subpreschemes of  $X$ , let  $k'/k$  be a field extension, and let  $X', Z'_i$  be obtained from  $X, Z_i$  by base change. Then  $(Z_i)_{i \in I}$  is schematically dense in  $X$  if and only if  $(Z'_i)_{i \in I}$  is schematically dense in  $X'$ .*

*Proof.* The “if” direction follows from Lemma 4.5.0; let us prove the converse.<sup>21</sup> First, we may suppose  $X = \operatorname{Spec}(B)$  affine. For every  $i$ , let  $(Z_{ij})_{j \in J_i}$  be an affine open covering of  $Z_i$ . Replacing  $(Z_i)_{i \in I}$  by  $(Z_{ij})_{(i,j) \in J}$ , where

$$J = \coprod_{i \in I} J_i,$$

we may also suppose that every  $Z_i$  is affine.

Let

$$g \in B' = B \otimes_k k', \quad t' \in B'_g$$

be a section of  $\mathcal{O}_{X'}$  on the affine open  $U' = X'_g$  whose restriction to each  $Z'_i \cap U'$  is zero. Thus its image in every

$$(\mathcal{O}(Z_i) \otimes_k k')_g$$

is zero. There exists a finite-type  $k$ -subalgebra  $A$  of  $k'$  such that

$$g \in B_A = B \otimes_k A, \quad t' \in (B_A)_g.$$

The map

$$\mathcal{O}(Z_i) \otimes_k A \longrightarrow \mathcal{O}(Z_i) \otimes_k k'$$

is injective, and therefore so is

$$(\mathcal{O}(Z_i) \otimes_k A)_g \longrightarrow (\mathcal{O}(Z_i) \otimes_k k')_g.$$

<sup>19</sup>Editor’s note: this lemma has been inserted here because it is used in the proofs of 4.5 and 4.7.

<sup>20</sup>Editor’s note: this hypothesis is in fact superfluous; cf. EGA IV<sub>3</sub>, 11.10.6 and 11.9.13.

<sup>21</sup>Editor’s note: the original has been expanded in what follows.

The hypothesis consequently implies that the image of  $t'$  in every  $(\mathcal{O}(Z_i) \otimes_k A)_g$  is zero. It remains to show that  $(Z_{iA})_{i \in I}$  is schematically dense in  $X_A$ , which follows from EGA IV<sub>3</sub>, 11.9.10(b).<sup>22</sup>  $\square$

Let us record in passing the following result, which will be used in a later exposé.<sup>23</sup>

**Corollary 4.6.** *Let  $X$  be a flat  $S$ -prescheme and  $U$  an open subset of  $X$ . Suppose that  $U_s$  is schematically dense in  $X_s$  for every  $s \in S$ , and that either  $X$  is locally Noetherian or  $X$  is locally of finite presentation over  $S$ . Then  $U$  is schematically dense in  $X$ .*

*Proof.* For simplicity, suppose  $U$  is retrocompact in  $X$  (cf. EGA IV<sub>3</sub>, 11.10.10 and 11.9.17 for the general case). The locally Noetherian case is included only for reference; it is contained in 4.3.

In the second case, we may plainly suppose  $S$  and  $X$  affine. Then  $X$  and  $U$  are of finite presentation over  $S = \operatorname{Spec}(A)$ . The ring  $A$  is the direct limit of its finite-type  $\mathbb{Z}$ -subalgebras  $A_i$ . The “patented procedure” already used (cf. EGA IV<sub>3</sub>, 8.8.2 and 8.10.5(iii)) shows that there are an index  $i$ , an affine scheme  $X_i$  over

$$S_i = \operatorname{Spec}(A_i),$$

and an open subset  $U_i$  of  $X_i$ , from which  $X, U$  are obtained by the base change  $S \rightarrow S_i$ . Let  $E_i$  be the subset of  $S_i$  consisting of those  $s \in S_i$  for which  $(U_i)_s$  is schematically dense in  $(X_i)_s$ .

By EGA IV<sub>2</sub>, 5.9.9 and 5.10.2,  $E_i$  is the set of points  $s \in S_i$  for which  $(U_i)_s$  contains the set

$$\operatorname{Ass} \mathcal{O}_{(X_i)_s}$$

of points associated with the structure sheaf of  $(X_i)_s$ . By EGA IV<sub>3</sub>, 11.9.17.1, this is a constructible condition; hence  $E_i$  is constructible.<sup>24</sup>

By 4.5, the inverse image of  $E_i$  under  $S_j \rightarrow S_i$  is  $E_j$ , and its inverse image under  $S \rightarrow S_i$  is the set  $E$  of those  $s \in S$  for which  $U_s$  is schematically dense in  $X_s$ .<sup>24</sup> By hypothesis,  $E = S$ , which is also the inverse image of  $S_i$  under every  $S \rightarrow S_i$ . EGA IV<sub>3</sub>, 8.3.11 therefore implies that there is a  $j \geq i$  such that  $E_j = S_j$ ; that is,  $(U_j)_s$  is schematically dense in  $(X_j)_s$  for every  $s \in S_j$ .

Apply 4.4 to  $(S_j, X_j, U_j)$  and the base change  $S \rightarrow S_j$ . It follows that  $U$  is schematically dense in  $X$ . Notice that here 4.4 is used only for a family with a finite index set (in fact, a singleton); in this case its proof simplifies considerably.  $\square$

**Theorem 4.7.** *Let  $S$  be a prescheme, and let  $G$  be a group of multiplicative type and of finite type over  $S$ . For every integer  $n > 0$ , let  ${}_nG$  be the kernel of  $n \cdot \operatorname{id}_G$ . Then the family of subpreschemes  $({}_nG)_{n>0}$  is schematically dense in  $G$  in the sense of Definition 4.1.*

*Proof.* We distinguish two cases.

(a)  *$S$  locally Noetherian.* By 4.3, we are reduced to the case in which  $S$  is the spectrum of a field  $k$ . By 4.5.0, we may suppose that  $k$  is algebraically closed and that  $G$  is diagonalizable, say

$$G = D_k(M),$$

where  $M$  is a commutative group of finite type. Then

$$M \simeq \Gamma \times \mathbb{Z}^r$$

<sup>22</sup>Editor’s note: the original continued: “Using 4.3 for the  $X_{A'}$ , where  $A'$  is a quotient of  $A$  by a power of a maximal ideal, and Hilbert’s Nullstellensatz, one is reduced to the case where  $k'$  is a finite extension of  $k$ .” This reduction could be expanded and that argument pursued, since the case of a finite extension  $k'/k$  is somewhat simpler than the more general case treated in EGA IV<sub>3</sub>, 11.9.10(b).

<sup>23</sup>Editor’s note: this reference remains to be specified.

<sup>24</sup>Editor’s note: the original has been expanded in this paragraph and the next.

with  $\Gamma$  finite, and consequently

$$G \simeq G_1 \times G_2, \quad G_1 = D(\Gamma), \quad G_2 = \mathbf{G}_m^r.$$

For  $n$  sufficiently divisible—namely, for  $n$  a multiple of the order of  $\Gamma$ —one has

$${}_nG = G_1 \times {}_nG_2, \quad \text{since} \quad {}_nG_1 = G_1.$$

Applying 4.3 again to the projection  $G \rightarrow G_1$ , we are reduced to  $G_2$ , hence to the case  $G = \mathbf{G}_m^r$ . Since  $G$  is then reduced, schematic density of  $({}_nG)_{n>0}$  is equivalent to density of  $\bigcup_n {}_nG$  in the ordinary topology. As

$${}_nG = ({}_n\mathbf{G}_m)^r,$$

we are reduced to  $G = \mathbf{G}_m$ , an irreducible group of dimension one. The assertion now follows because  $\bigcup_n {}_nG$ , the set of roots of unity in  $k$ , is infinite.

(b) *The general case.* For every  $s \in S$ , there are an open neighborhood  $U$  of  $s$  and a faithfully flat quasi-compact morphism  $S' \rightarrow U$  such that  $G' = G_{S'}$  is diagonalizable, say  $G' = D_{S'}(M)$ . By 4.5.0 we may reduce to  $G'$ , and therefore suppose that  $G$  itself is diagonalizable. It is obtained by base change

$$S \longrightarrow \operatorname{Spec}(\mathbb{Z})$$

from the absolute group  $H = D_{\mathbb{Z}}(M)$ . Part (a) shows that, for every  $s \in \operatorname{Spec}(\mathbb{Z})$ , the family  $({}_nH_s)_{n>0}$  is schematically dense in  $H_s$ . It remains only to apply 4.4.<sup>25</sup>  $\square$

**Corollary 4.8.**

- (a) *Let  $u, v : H \rightarrow G$  be two homomorphisms of  $S$ -preschemes in groups, with  $H$  of multiplicative type<sup>26</sup> and of finite type. Suppose that, for every integer  $n > 0$ , the restrictions of  $u$  and  $v$  to  ${}_nH$  are equal. Then  $u = v$ .*
- (b) *Let  $H_1, H_2$  be two subgroups of multiplicative type and of finite type of a prescheme in groups  $G$ . If  ${}_nH_1 = {}_nH_2$  for every integer  $n > 0$ , then  $H_1 = H_2$ .*

*Proof.* Part (a) follows from part (b) by considering the subgroups  $H_1, H_2$  of  $H \times G$  that are the graphs of  $u, v$ . For part (b), put

$$H = H_1 \cap H_2.$$

This is a subgroup prescheme of  $H_i$ , for  $i = 1, 2$ , and we must show that it equals  $H_i$ . The hypothesis says precisely that  $H$  contains all the  ${}_nH_i$ . We are therefore reduced to the next corollary.  $\square$

**Corollary 4.9.** *Let  $G$  be an  $S$ -group of multiplicative type and of finite type, and let  $H$  be a subgroup prescheme containing  ${}_nG$  for every  $n > 0$ . Then  $H = G$ .*

*Proof.* By 4.7, it is enough to prove that  $H$  is closed, or equivalently that it equals  $G$  set-theoretically. This reduces us to the case in which  $S$  is the spectrum of a field. Every subgroup prescheme of  $G$  is then closed, by VI<sub>B</sub>, 1.4.2.  $\square$

**Remark 4.10.** Under the hypotheses of 4.7, let  $m > 0$  be an integer with the following properties: for every  $s \in S$ ,  $m$  is not a power of the characteristic of  $\kappa(s)$ ; and if  $G_s$

<sup>25</sup>Editor's note: this result will be needed in X, 4.3 for a non-locally-Noetherian base, namely  $S = \operatorname{Spec}(\hat{A} \otimes_A \hat{A})$ , where  $\hat{A}$  is the completion of the Noetherian local ring  $A$ .

<sup>26</sup>Editor's note: "diagonalizable" has been replaced by "of multiplicative type."

has type  $M$ , every prime divisor of the torsion subgroup of  $M$  divides  $m$ . The second condition is automatically satisfied if  $G$  is a torus. The proof shows that in Theorem 4.7 and Corollary 4.8 and Corollary 4.9 one may restrict to the subgroups

$$m^r G, \quad r > 0.$$

Moreover, if  $G$  is smooth over  $S$ , then for every  $s \in S$  there are an open neighborhood  $U$  of  $s$  and an integer  $m > 0$ , prime to every residue characteristic on  $U$ , that satisfy the preceding conditions for  $G_U$ . For instance, take  $m$  to be the order of the torsion subgroup of the type of  $G$  at  $s$  (cf. 1.4.1(a) and 2.1(e)). After restriction to  $U$ , the groups  $m^r G$  are then finite and étale over  $S$ , and their family is schematically dense. In certain applications—in particular those involving Theorem 3.2 and Theorem 3.2 bis (cf. XI, 6), though not those involving Theorem 3.6 and Theorem 3.6 bis—this observation allows one to avoid Theorem 3.1 and its use of Hochschild cohomology.

## 5. Central Homomorphisms of Groups of Multiplicative Type

**Lemma 5.0.**<sup>27</sup> *Let  $(A, \mathfrak{m})$  be a Noetherian local ring, put  $S = \operatorname{Spec}(A)$ , and let  $H$  be a finite scheme over  $S$ , say  $H = \operatorname{Spec}(B)$ , where  $B$  is finite over  $A$ . Let  $K$  be a subscheme of  $H$  such that reduction modulo  $\mathfrak{m}^{n+1}$  gives  $K_n = H_n$  for every  $n$ . Then  $K = H$ .*

*Proof.* Let  $s$  be the closed point of  $S$ . First,  $K$  is a closed subscheme of  $H$ . Indeed, a priori it is a closed subscheme of an open subset  $U$  of  $H$ . But  $K$ , and hence  $U$ , contains the fiber

$$K_s = H_s.$$

Since  $H \rightarrow S$  is finite, and therefore closed, the complement of  $U$  is empty; thus  $U = H$ . Hence  $K$  is defined by an ideal  $I$  of  $B$ . The hypothesis implies

$$I \subset \mathfrak{m}^n B \quad \text{for every } n.$$

Since  $B$  is a finite  $A$ -module, it is separated for the  $\mathfrak{m}$ -adic topology. Therefore  $I = 0$ .  $\square$

**Theorem 5.1.** *Let  $u, v : H \rightarrow G$  be two homomorphisms of  $S$ -preschemes in groups, where  $H$  is of multiplicative type and of finite type. Suppose either that  $S$  is locally Noetherian or that  $G$  is of finite presentation over  $S$ . Let  $s \in S$  satisfy  $u_s = v_s$ , and suppose that  $u_s$  is central. Then there is an open neighborhood  $U$  of  $s$  such that  $u_U = v_U$ .*

*Proof.* We distinguish the two cases.

(a)  $S$  locally Noetherian.<sup>28</sup> Let

$$K = \operatorname{Ker}(u, v)$$

be the inverse image of the diagonal of  $G \times_S G$  under  $(u, v) : H \rightarrow G \times_S G$ . It is a subgroup prescheme of  $H$ . We seek an open  $U$  for which  $K_U = H_U$ . Because  $S$  is locally Noetherian and  $H$  is of finite type over  $S$ , the prescheme  $H$  is locally Noetherian. Hence the immersion  $K \hookrightarrow H$  is of finite type (cf. EGA I, 6.3.5), so  $K$  is of finite type, and therefore of finite presentation, over  $S$ . By EGA IV<sub>3</sub>, 8.8.2.4, it is enough to prove that

$$K_{S_0} = H_{S_0}, \quad S_0 = \operatorname{Spec}(\mathcal{O}_{S,s}).$$

<sup>27</sup>Editor's note: this lemma has been made explicit because it is used several times below; it was implicit in the original.

<sup>28</sup>Editor's note: the original has been expanded in what follows.

We may thus suppose that  $S$  is local with closed point  $s$ . Replacing the Noetherian local ring  $A$  by its completion  $\widehat{A}$ , if desired, gives a faithfully flat quasi-compact base change  $\widehat{S} \rightarrow S$ , and allows us to suppose that  $A$  is complete.<sup>29</sup>

By 3.4, one has  $u_n = v_n$  for every  $n$ , where the subscript denotes reduction modulo  $\mathfrak{m}^{n+1}$ , with  $\mathfrak{m}$  the maximal ideal of  $A$ . For every  $m > 0$ , let

$${}_m u, {}_m v : {}_m H \longrightarrow G$$

be the homomorphisms induced by  $u, v$ . Then

$$({}_m u)_n = ({}_m v)_n \quad \text{for every } n.$$

The group  ${}_m H$  is finite over  $S$ , by 2.2, so 5.0 gives  ${}_m u = {}_m v$ . This holds for every  $m$ , and therefore 4.7 gives  $u = v$ .

(b) *G of finite presentation.*<sup>30</sup> The group  $H$  is also of finite presentation over  $S$ . By EGA IV<sub>3</sub>, 8.8.2, we may again suppose that  $S$  is local with closed point  $s$ , and it suffices to prove  $u = v$ . Let  $f : S' \rightarrow S$  be a faithfully flat quasi-compact morphism. Write

$$f_H : H' \rightarrow H, \quad f_G : G' \rightarrow G, \quad u', v' : H' \rightarrow G'$$

for the resulting morphisms. If  $u' = v'$ , then

$$u \circ f_H = f_G \circ u' = f_G \circ v' = v \circ f_H,$$

and  $u = v$  because  $f_H$  is an epimorphism. After a faithfully flat quasi-compact extension of the base, we may therefore suppose that  $H$  is diagonalizable:

$$H = D_S(M)$$

for a commutative group  $M$  of finite type.

As in the proof of 4.6, introduce the increasing filtered family of  $\mathbb{Z}$ -subalgebras  $A_i$  of  $A = \mathcal{O}_{S,s}$  that are of finite type, and put

$$S_i = \text{Spec}(A_i).$$

<sup>30</sup> For every  $i$ , the group  $H = D_S(M)$  comes from

$$H_i = D_{S_i}(M).$$

Since  $G$  is of finite presentation over  $S$ , EGA IV<sub>3</sub>, 8.8.2 (see also VI<sub>B</sub>, 10.2 and 10.3) gives an index  $i$ , an  $S_i$ -group prescheme  $G_i$  of finite presentation, and homomorphisms

$$u_i, v_i : H_i \longrightarrow G_i$$

whose base changes are  $u, v$ . Let  $s_i$  be the image of  $s$  in  $S_i$ , and define

$$\rho_i : H_i \times_{S_i} G_i \longrightarrow G_i, \quad \rho_i(h, g) = u_i(h)g u_i(h)^{-1}.$$

Because  $u_s$  is central,

$$\rho_s = \rho_i \times_{\kappa(s_i)} \kappa(s)$$

<sup>29</sup>Editor's note: this is because  $K = H$  if  $K_{\widehat{S}} = H_{\widehat{S}}$ ; cf. SGA 1, VIII, 5.4. The argument below does not use completeness of  $A$ .

<sup>30</sup>Editor's note: the original has been expanded in what follows.



is the second projection. The extension  $\kappa(s_i) \rightarrow \kappa(s)$  is faithfully flat, so  $\rho_i$  is also the second projection over  $s_i$ ; hence  $(u_i)_{s_i}$  is central. Likewise  $u_s = v_s$  implies

$$(u_i)_{s_i} = (v_i)_{s_i}.$$

Part (a), applied over  $S_i$ , now gives the conclusion.  $\square$

**Corollary 5.2.** *Let  $u : H \rightarrow G$  be a homomorphism of  $S$ -preschemes in groups, where  $H$  is of multiplicative type and of finite type, and suppose either that  $S$  is locally Noetherian or that  $G$  is of finite presentation over  $S$ . If  $s \in S$  and  $u_s$  is the unit homomorphism, then there is an open neighborhood  $U$  of  $s$  on which  $u_U$  is the unit homomorphism.*

**Corollary 5.3.** *Let  $u, H, G$  be as in 5.2, and suppose in addition that  $G$  is separated over  $S$ . Let  $U$  be the set of points  $s \in S$  for which  $u_s$  is the unit homomorphism. Then  $U$  is open and closed in  $S$ , and  $u_U : H_U \rightarrow G_U$  is the unit homomorphism.*

*Proof.* Only closedness remains to be proved. Put  $K = \text{Ker}(u)$ . Since  $G$  is separated over  $S$ ,  $K$  is a closed subprescheme of  $H$ , and  $U$  is the set of points  $s \in S$  for which  $K_s = H_s$ . It follows readily that  $U$  is closed, for example by VIII, 6.4, since  $H$  is essentially free over  $S$  by VIII, 6.3. Alternatively, we may suppose that  $S$  is reduced and that  $U$  is dense in  $S$ , hence schematically dense. Then  $H_U$  is schematically dense in  $H$  because  $H$  is flat over  $S$ ,<sup>31</sup> and  $u$  and the unit homomorphism, being equal on  $H_U$ , are equal on  $H$ .  $\square$

**Corollary 5.4.** *Let  $S$  be a prescheme; let  $H, G$  be  $S$ -groups, with  $H$  of multiplicative type and of finite type and  $G$  separated and of finite presentation over  $S$ ; and let  $\pi : S' \rightarrow S$  be a<sup>32</sup> universal effective epimorphism—for example, a faithfully flat quasi-compact morphism or a morphism admitting a section (cf. IV, 1.12)—whose fibers are geometrically connected.<sup>33</sup>*

*Let  $u' : H' \rightarrow G'$  be a central homomorphism of the  $S'$ -groups obtained from  $H, G$  by base change. Then there is a unique homomorphism  $u : H \rightarrow G$  of  $S$ -groups such that  $u \times_S S' = u'$ . If  $S'/S$  admits a section  $g$ , then  $u$  is the morphism obtained from  $u'$  by the base change  $g : S \rightarrow S'$ .*

*Proof.* Because  $\pi$  is a universal effective epimorphism, so is  $H' \rightarrow H$ ; uniqueness follows as at the beginning of the proof of 5.1(b). If  $\pi$  has a section  $g$ , then

$$u' = \pi^* u \implies u = g^* \pi^* u = g^* u'.$$

For existence, IV, 2.3 reduces us to showing that the two homomorphisms

$$u''_1, u''_2 : H'' \longrightarrow G''$$

of  $S''$ -groups obtained from  $u'$  by the two base changes

$$\text{pr}_1, \text{pr}_2 : S'' = S' \times_S S' \longrightarrow S'$$

are equal. Their pullbacks along the diagonal  $S' \rightarrow S''$  are equal, since both are  $u'$ .<sup>34</sup> Because  $u''_1, u''_2$  are central, we may apply 5.3 to

$$u''_1(u''_2)^{-1}.$$

<sup>31</sup>Editor's note: and of finite presentation over  $S$ ; cf. EGA IV<sub>3</sub>, 11.10.5(ii) and 11.9.10(ii).

<sup>32</sup>Editor's note: "covering morphism for the faithfully flat quasi-compact topology" has been replaced by "universal effective epimorphism," so that the result applies to the morphism  $K \rightarrow S$  in 5.5; the beginning of the proof has been modified accordingly.

<sup>33</sup>Editor's note: this holds, for example, when  $S = \text{Spec}(k)$  and  $S' = \text{Spec}(k')$ , where  $k$  is a field and  $k'/k$  is radical; cf. X, Proposition 1.4.

<sup>34</sup>Editor's note: the preceding sentence has been added.

There is therefore an open-and-closed subset  $U$  of  $S''$ , containing the diagonal, over which  $u_1''$  and  $u_2''$  coincide. The fibers of  $S'/S$  are geometrically connected, hence so are those of  $S''/S$ , and in particular they are connected. Since  $U$  contains the diagonals of these fibers, it contains every fiber and therefore equals  $S''$ . This proves existence.  $\square$

**Corollary 5.5.** *Let  $S$  be a prescheme, let  $K$  be an  $S$ -group prescheme locally of finite type<sup>35</sup> and with connected fibers, and let  $H$  be an invariant subgroup of  $K$ , of multiplicative type and of finite type. Then  $H$  is a central subgroup of  $K$ .*

*Proof.*<sup>36</sup> First,  $\pi : K \rightarrow S$  has geometrically connected fibers, because for a group scheme locally of finite type over a field, connectedness implies geometric connectedness (cf. VI<sub>A</sub>, 2.4). Apply 5.4 with  $G = H$  and  $S' = K$  to the homomorphism of  $K$ -groups

$$u' : H_K \longrightarrow H_K, \quad (h, k) \longmapsto (khk^{-1}, k).$$

It is central because  $H$  is commutative. Its pullback along the unit section  $\epsilon : S \rightarrow K$  is the identity homomorphism of  $H$ . [The source prints “of  $K$ ” here; the displayed endomorphism of  $H_K$  shows that  $H$  is intended.] Corollary 5.4 therefore implies that  $u'$  itself is the identity on  $H_K$ . Thus  $H$  is central in  $K$ .  $\square$

Let us state the variants of the preceding results for central subgroups of multiplicative type. Arguing as above, and using 3.2 bis, gives:

**Theorem 5.1 bis.** *Let  $G$  be an  $S$ -group prescheme, and let  $H_1, H_2$  be two subgroup preschemes of multiplicative type and of finite type, with  $(H_1)_s$  central. Suppose either that  $S$  is locally Noetherian or that  $G$  is of finite presentation over  $S$ . For every  $s \in S$  such that  $(H_1)_s = (H_2)_s$ , there is an open neighborhood  $U$  of  $s$  for which  $H_1|_U = H_2|_U$ .*

**Corollary 5.3 bis.** *Under the preceding hypotheses, suppose moreover that  $G$  is separated over  $S$ . Let  $U$  be the set of points  $s \in S$  for which  $(H_1)_s = (H_2)_s$ . Then  $U$  is open and closed in  $S$ , and  $H_1|_U = H_2|_U$ .*

**Corollary 5.4 bis.** *Let  $S$  be a prescheme, let  $G$  be a group prescheme separated and of finite presentation over  $S$ , let  $S' \rightarrow S$  be a covering morphism for the faithfully flat quasi-compact topology with geometrically connected fibers, and let  $H'$  be a subgroup of multiplicative type and of finite type of  $G' = G \times_S S'$ . Then there is a unique subgroup  $H$  of  $G$  such that  $H' = H \times_S S'$ . The subgroup  $H$  is of multiplicative type and of finite type over  $S$ , by 1.1 and 2.1(b). If  $S' \rightarrow S$  admits a section  $g$ , then  $H = g^*H'$ .*

**Theorem 5.6.** *Let  $S$  be a prescheme and let  $u : H \rightarrow G$  be a homomorphism of  $S$ -preschemes in groups, where  $H$  is of multiplicative type and of finite type and  $G$  is of finite presentation over  $S$ .*

(a) *Suppose that  $G$  has connected fibers. If  $s \in S$  and  $u_s : H_s \rightarrow G_s$  is central, then there is an open neighborhood  $U$  of  $s$  such that the homomorphism  $H|_U \rightarrow G|_U$  induced by  $u$  is central.*

(b) *If  $u_s : H_s \rightarrow G_s$  is central for every  $s \in S$ , then  $u$  is central.*

*Proof.* We first prove (a).<sup>37</sup> Arguing exactly as in the proof of 5.1(b), we may suppose first that  $S$  is local with closed point  $s$ , then that  $H = D_S(M)$  is diagonalizable, and finally that there exist a finite-type  $\mathbb{Z}$ -subalgebra  $A_i$  of  $A = \mathcal{O}_{S,s}$  and a homomorphism

$$u_i : D_{S_i}(M) \longrightarrow G_i$$

<sup>35</sup>Editor’s note: the hypothesis “locally of finite type” has been added to agree with VI<sub>A</sub>, 2.4. In fact, over a field every connected group scheme is geometrically connected.

<sup>36</sup>Editor’s note: the original has been expanded in what follows.

<sup>37</sup>Editor’s note: the original has been expanded in the proof of (a).

whose base change is  $u$  and for which  $(u_i)_{s_i}$  is central, where  $s_i$  is the image of  $s$  in  $S_i$ . We are thereby reduced to the case in which  $S$  is Noetherian local with closed point  $s$ , and we must prove that  $u$  is central.

Let  $K = \text{Ker}(v, w)$ , where

$$v, w : H \times_S G \rightrightarrows G$$

are defined by

$$v(h, g) = g, \quad w(h, g) = \text{int}(u(h)) \cdot g = u(h)g u(h)^{-1}.$$

Then  $K$  is a subgroup over  $G$  of

$$H_G = H \times_S G,$$

and we must prove that  $K = H_G$ . By 4.9, it is enough to show that  $K$  contains

$${}_m(H_G) = ({}_m H) \times_S G$$

for every  $m > 0$ . We are thus reduced to the case  $H = {}_m H$ , so that  $H$  is finite over  $S$ .

Let  $e$  be the unit element of the fiber  $G_s$ , and put

$$S^0 = \text{Spec}(\mathcal{O}_{G,e}), \quad S_n^0 = S^0 \times_S S_n, \quad S_n = \text{Spec}(A/\mathfrak{m}^{n+1}).$$

The scheme  $S^0$  is Noetherian local because  $G$  is of finite presentation over the Noetherian scheme  $S$ . The base change

$$K_{S^0} = K \times_G S^0$$

is a subscheme of

$$H_{S^0} = H \times_S S^0,$$

and 3.9 gives

$$K_{S_n^0} = H_{S_n^0} \quad \text{for every } n.$$

<sup>38</sup> Since  $H_{S^0}$  is finite over  $S^0$ , 5.0, applied to the Noetherian local ring  $\mathcal{O}_{G,e}$ , gives

$$K_{S^0} = H_{S^0}.$$

[The source prints the undefined local ring  $B$  here; the Noetherian local ring  $\mathcal{O}_{G,e}$  just introduced is intended.]

On the other hand,  $H \times_S G$  is Noetherian, being of finite presentation over the Noetherian scheme  $S$ . Thus the immersion

$$K \hookrightarrow H \times_S G$$

is of finite type (cf. EGA I, 6.3.5), and  $K$  is of finite presentation over  $G$ . The equality

$$K_{S^0} = H_G \times_G S^0$$

therefore implies, by EGA IV<sub>3</sub>, 8.8.2.4, that there is an open neighborhood  $W$  of  $e$  in  $G$  such that

$$K \times_G W = H_G \times_G W = H \times_S W.$$

Hence  $K$  contains the open neighborhood

$$V = H \times_S W$$

---

<sup>38</sup>Editor's note: the original has been expanded and corrected, replacing "the unit section of  $(G_H)_n$  over  $H_n$ " by  $H_{S_n^0}$ .

of the unit section of  $G_H = G \times_S H$  over  $H$ .

For every  $t \in H$ , the fiber  $G_t$  is an algebraic group over  $\kappa(t)$ , hence Cohen–Macaulay by VI<sub>A</sub>, 1.1.1 and therefore has no embedded components. It is also connected, hence irreducible by VI<sub>A</sub>, 2.4, so its generic point is its unique associated point. EGA IV<sub>2</sub>, 3.1.8 therefore shows that  $V_t$  is schematically dense in  $G_t$ . By 4.3,  $V$  is schematically dense in  $G_H$ .

Moreover, since  $K$  is an  $H$ -subgroup of  $G_H$ , it induces on each fiber  $G_h$  a subgroup  $K_h$ . This subgroup contains an open neighborhood of the unit element; because  $G_h$  is connected, it follows that  $K_h = G_h$ . Thus  $K = G_H$  set-theoretically. Finally,  $K$  is a closed subscheme of  $G_H$  containing the schematically dense subscheme  $V$ , and hence  $K = G_H$ . This proves (a).

Part (b) follows directly from 3.9: to verify that a subscheme  $K$  of  $P = G \times_S H$  equals  $P$ , it is enough to verify, after every base change  $S' \rightarrow S$  in which  $S'$  is the spectrum of an Artinian quotient of a local ring of  $S$ , that  $K' = P'$ .  $\square$

## 6. Monomorphisms of Groups of Multiplicative Type, and Canonical Factorization of a Homomorphism from Such a Group

**Lemma 6.1.** *Let  $S$  be a quasi-compact prescheme, and let  $G$  be a commutative  $S$ -group prescheme, of finite presentation and quasi-finite over  $S$ . Then there is an integer  $n > 0$  such that*

$$n \cdot \text{id}_G = 0, \quad \text{that is,} \quad G = {}_n G.$$

*Proof.* If  $S$  is the spectrum of a field  $k$ , then  $G$  is finite over  $k$ , and by VII<sub>A</sub> 8.5 it is enough to take

$$n = \deg(G/k).$$

Suppose now that  $S = \text{Spec}(A)$ , where  $A$  is local Artinian. Let  $k$  be the residue field of  $A$ , put

$$G_0 = G \otimes_A k, \quad n_0 = \deg(G_0/k),$$

and distinguish two cases.

(a)  *$k$  has characteristic zero.* Then  $G_0$  is separable over  $k$ , and hence  $G$  is unramified over  $S$ . The unit section of  $G$  is therefore an open immersion, so

$${}_n G = \text{Ker}(n \cdot \text{id}_G)$$

is an open subscheme of  $G$ . For it to equal  $G$ , it is consequently enough that it do so set-theoretically; thus one may take  $n = n_0$ .

(b)  *$k$  has characteristic  $p > 0$ .* Let  $\mathfrak{m}$  be the maximal ideal of  $A$ , and put

$$S_r = \text{Spec}(A/\mathfrak{m}^{r+1}) \quad (r \geq 0).$$

Choose an integer  $m$  such that  $\mathfrak{m}^{m+1} = 0$ . We claim that one may take  $n = p^m n_0$ . Indeed, arguing by induction on  $m$  and putting  $n' = p^{m-1} n_0$ , we may suppose that  $n' \cdot \text{id}_G$  induces the zero endomorphism on

$$G_{m-1} = G \times_S S_{m-1}.$$

<sup>39</sup> Let  $E$  be the set of endomorphisms  $u$  of  $G$  having this property, and let  $K$  be the kernel of the morphism of group functors

$$G \longrightarrow \prod_{S_{m-1}/S} G$$

(cf. III 0.1.2). Then

$$E = \operatorname{Hom}_{S\text{-gr}}(G, K);$$

in particular, the abelian group law on  $E$  is induced by that on  $K$ . By III 0.9,  $K$  is the  $S$ -group functor which, to every  $g : T \rightarrow S$ , assigns the  $k$ -vector space

$$\operatorname{Hom}_{\mathcal{O}_{T_0}}(g_0^*(\Omega_{G_0/S_0}^1), \mathfrak{m}^m \mathcal{O}_T).$$

Thus  $pu = 0$  for every  $u \in E$ . It follows that

$$pn' \cdot \operatorname{id}_G = 0, \quad \text{that is,} \quad p^m n_0 \cdot \operatorname{id}_G = 0.$$

Suppose now that  $S$  is Noetherian; this is the case to which the general assertion reduces by the usual reduction to the Noetherian case.<sup>40</sup> It is enough to combine the foregoing with the following lemma.

**Lemma 6.2.** *Let  $X$  be a quasi-finite prescheme over a Noetherian prescheme  $S$ , and let  $(X_i)_{i \in I}$  be an increasing filtered family of subpreschemes having the following property:*

$$\left\{ \begin{array}{l} \text{for every } s \in S \text{ and every } n \geq 0, \text{ putting} \\ S_{s,n} = \operatorname{Spec}(\mathcal{O}_{S,s}/\mathfrak{m}^{n+1}), \text{ there is an } i \in I \text{ such that} \\ X_i \times_S S_{s,n} = X \times_S S_{s,n}. \end{array} \right.$$

*Then there is an  $i \in I$  such that  $X_i = X$ .*

*Proof.* Since  $S$  is Noetherian, there is a maximal open subset  $U$  on which

$$X|_U = X_i|_U$$

for all sufficiently large  $i$ . We shall prove that  $U = S$ . In other words, if  $U \neq S$ , we shall find an open subset  $U'$  strictly larger than  $U$ , and an  $i$ , such that

$$X|_{U'} = X_i|_{U'}.$$

Localizing at a maximal point  $s$  of  $S - U$ , we reduce to the case in which  $S$  is local with closed point  $s$ , and

$$U = S - \{s\}.$$

Indeed, put  $S' = \operatorname{Spec}(\mathcal{O}_{S,s})$ . If

$$X \times_S S' = X_i \times_S S'$$

for some  $i$ ,<sup>41</sup> there is an open neighborhood  $V$  of  $s$  such that

$$X|_V = X_i|_V.$$

<sup>39</sup>Editor's note: the original has been expanded in what follows, with the results of Exposé III that are used made explicit.

<sup>40</sup>Editor's note: since  $S$  is quasi-compact, it is covered by finitely many affine open subsets, so one reduces to  $S = \operatorname{Spec}(A)$ . Since  $G$  is of finite presentation over  $S$ , EGA IV<sub>3</sub>, 8.8.2 may then be applied.

<sup>41</sup>Editor's note:  $X - \{s\}$  has been corrected above to  $S - \{s\}$ . Recall also that a quasi-finite morphism is assumed to be of finite type; cf. EGA II, 6.2.3, and EGA III<sub>2</sub>, Err<sub>III</sub> 20 for the definition of "locally quasi-finite." Thus here, since  $S$  is Noetherian,  $X$  is of finite presentation over  $S$ ; every immersion  $X_i \hookrightarrow X$  is of finite type by EGA I, 6.3.5, so  $X_i$  is of finite presentation over  $S$ , and EGA IV<sub>3</sub>, 8.8.2 applies.

Taking  $i$  large enough that  $X_i|_U = X|_U$ , we then have

$$X|_W = X_i|_W, \quad W = U \cup V.$$

For all sufficiently large  $i$ , the equality  $X|_U = X_i|_U$  shows that  $X_i$  is a closed subscheme of  $X$ , defined by an ideal  $\mathcal{J}^{(i)}$  whose support is contained in

$$X_s = X_0 = X \times_S S_0,$$

where

$$S_n = \operatorname{Spec}(A/\mathfrak{m}^{n+1}), \quad S = \operatorname{Spec}(A).$$

Since  $X_0$  is quasi-finite over  $S_0$ , it is a finite closed subset of the Noetherian prescheme  $X$ . Hence  $\mathcal{J}^{(i)}$  is a module of finite length. There is therefore an integer  $n \geq 0$  such that

$$\mathcal{J}^{(i)} \cap \mathfrak{m}^{n+1} \mathcal{O}_X = 0.$$

On the other hand, by the hypothesis of the lemma, after increasing  $i$  if necessary we may suppose that the image of  $\mathcal{J}^{(i)}$  in

$$\mathcal{O}_{X_n} = \mathcal{O}_X / \mathfrak{m}^{n+1} \mathcal{O}_X$$

is zero. It follows that  $\mathcal{J}^{(i)} = 0$ , and therefore  $X_i = X$ .  $\square$

**Lemma 6.3.** *Let  $S$  be a prescheme, let  $K$  be a group prescheme over  $S$ , locally of finite presentation, and let  $s \in S$ . Suppose that  $K_s$  is quasi-finite (respectively, unramified) over  $\kappa(s)$  at the unit element. Then there is an open neighborhood  $U$  of  $s$  such that  $K|_U$  is locally quasi-finite (respectively, unramified) over  $U$ .\**

*Proof.* Let  $V$  be the set of points  $x \in K$  having the following property: if  $t$  is the image of  $x$  in  $S$ , then  $K_t$  is quasi-finite (respectively, unramified) over  $\kappa(t)$  at  $x$ ; that is,  $x$  is isolated in  $K_t$  (respectively, it is isolated and its local ring in  $K_t$  is a separable extension of  $\kappa(t)$ ). The set  $V$  is open because  $K$  is locally of finite presentation over  $S$ .<sup>42</sup> If  $\epsilon$  denotes the unit section of  $K$ , then  $\epsilon^{-1}(V)$  is open. By hypothesis it contains  $s$ , and is therefore an open neighborhood  $U$  of  $s$ . This  $U$  has the required property. Indeed, if  $t \in U$ , then  $K_t$ , being a group locally of finite type over  $\kappa(t)$ , is locally quasi-finite (respectively, unramified) over  $\kappa(t)$  because it has that property at  $\epsilon(t)$ ; cf. VI<sub>B</sub> 1.3.  $\square$

Combining 6.1 and 6.3, one obtains:

**Theorem 6.4.** *Let  $S$  be a prescheme, let  $H$  be an  $S$ -group of multiplicative type and of finite type, and let  $K$  be a closed group subscheme of  $H$ , of finite presentation over  $S$ . Let  $s \in S$  be such that  $K_s$  is finite. Then there are an open neighborhood  $U$  of  $s$  and an integer  $n > 0$  such that*

$$K|_U \subset {}_n H|_U.$$

*In particular, since  ${}_n H$  is finite over  $S$ , the group  $K|_U$  is finite over  $U$ .*

Nakayama's lemma now gives:

**Corollary 6.5.** *With the preceding notation, suppose that  $K_s$  is the unit group. Then there is an open neighborhood  $U$  of  $s$  such that  $K|_U$  is the unit group.*

**Corollary 6.6.** *Let  $u : H \rightarrow G$  be a homomorphism of  $S$ -group preschemes, where  $H$  is of multiplicative type and of finite type and  $G$  is separated over  $S$ . Suppose in addition*

\*Cf. VI<sub>B</sub> 2.5(ii) for a more general statement.

<sup>42</sup>Editor's note: cf. EGA IV<sub>3</sub>, 13.1.4 and EGA IV<sub>4</sub>, 17.4.1. In addition,  $t$ , rather than  $s$ , has been used for the image of  $x$ .

that either  $S$  is locally Noetherian or  $G$  is locally of finite presentation over  $S$ .<sup>43</sup> If  $s \in S$  and

$$u_s : H_s \longrightarrow G_s$$

is a monomorphism, then there is an open neighborhood  $U$  of  $s$  such that  $u$  induces a monomorphism

$$H|_U \longrightarrow G|_U.$$

*Proof.* Put  $K = \text{Ker}(u)$ . The hypothesis on  $u_s$  says that  $K_s$  is the unit group, and the desired conclusion says that  $K|_U$  is the unit group for a suitable  $U$ . Since  $G$  is separated over  $S$ , the group  $K$  is a closed subgroup of  $H$ . If  $S$  is not assumed locally Noetherian, but  $G$  is assumed locally of finite presentation over  $S$ , then  $K$  is locally of finite presentation over  $S$ .<sup>43</sup> It is therefore of finite presentation over  $S$ , since it is separated over  $S$  (as  $H$  is) and quasi-compact over  $S$  (being closed in  $H$ , which is quasi-compact over  $S$ ). Corollary 6.5 now applies.  $\square$

**Remark 6.6.1.**<sup>44</sup> Under the hypotheses of 6.6, note that if, in addition,  $G$  is affine over  $S$  (respectively, of finite presentation over  $S$ ), then

$$H|_U \longrightarrow G|_U$$

is even a closed immersion, as was observed in 2.5 (respectively, 2.6).

**Corollary 6.7.** Let  $u : H \rightarrow G$  be a homomorphism of  $S$ -group preschemes, where  $H$  is of multiplicative type and of finite type. Suppose that either  $S$  is locally Noetherian or  $G$  is locally of finite presentation over  $S$ . If every homomorphism on fibers

$$u_s : H_s \longrightarrow G_s$$

is a monomorphism, then  $u$  is a monomorphism.

*Proof.* The argument is the same as in 6.6. Here the hypothesis that  $G$  be separated over  $S$  is unnecessary in order to ensure that  $K = \text{Ker}(u)$  is closed in  $H$ , since the assumption that all the  $u_s$  are monomorphisms implies that  $K$  is set-theoretically reduced to the unit section.  $\square$

**Theorem 6.8.** Let  $u : H \rightarrow G$  be a homomorphism of  $S$ -group preschemes, where  $H$  is of multiplicative type and of finite type and  $G$  is separated over  $S$ . Suppose in addition that either  $S$  is locally Noetherian or  $G$  is of finite presentation over  $S$ . Then

$$K = \text{Ker}(u)$$

is a subgroup of  $H$  of multiplicative type and of finite type, and  $u$  factors as

$$H \xrightarrow{u'} H/K \xrightarrow{u''} G,$$

where  $H/K$  is of multiplicative type and of finite type,  $u'$  is the canonical homomorphism and is faithfully flat (and affine), and  $u''$  is a monomorphism.

*N.B.* As noted in 6.6.1, the homomorphism  $u''$  is a closed immersion if  $G$  is affine over  $S$ , or if  $G$  is of finite presentation over  $S$ .

<sup>43</sup>Editor's note: it is enough to suppose that  $G$  is locally of finite type over  $S$ . By EGA IV<sub>4</sub>, 1.4.3(v), this implies, since  $H \rightarrow S$  is of finite presentation, that  $H \rightarrow G$  is locally of finite presentation; the same is then true of  $K \rightarrow S$ , which is obtained from it by base change.

<sup>44</sup>Editor's note: the number 6.6.1 has been added for use in later references.

*Proof.* It is enough to prove that  $K$  is of multiplicative type; the rest then follows from 2.7 and IV 5.2.6.

Suppose first that  $G$  is of finite presentation over  $S$ . Since this hypothesis is stable under base change, in proving that  $K$  is of multiplicative type we may reduce to the case in which  $H$  is diagonalizable:

$$H = D_S(M),$$

where  $M$  is a commutative group of finite type. Let  $s \in S$ . The group  $K_s$  is a closed subgroup of

$$H_s = D_{\kappa(s)}(M).$$

By 8.1,<sup>45</sup> it is therefore of the form  $D_{\kappa(s)}(N)$ , where  $N$  is a quotient of  $M$ . Put

$$K' = D_S(N).$$

Then  $K'$  is a subgroup of  $H$  of multiplicative type. Let

$$v' : K' \longrightarrow G$$

be the homomorphism induced by  $u$ .<sup>46</sup> By construction,  $v'_s$  is the unit homomorphism. Corollary 5.2 therefore gives an open neighborhood  $U$  of  $s$  such that

$$v'|_U : K'|_U \longrightarrow G|_U$$

is the unit homomorphism. Replacing  $S$  by  $U$ , we may suppose that  $v'$  itself is the unit homomorphism, and hence that  $u$  factors as

$$H \xrightarrow{u'} H/K' \xrightarrow{u''} G.$$

The homomorphism  $u''_s$  is obtained from  $u_s$  by factoring through

$$H_s \longrightarrow H_s / \text{Ker}(u_s),$$

and is therefore a monomorphism by IV 5.2.6. By 6.6, after restricting to a suitable open neighborhood  $U$  of  $s$ , the homomorphism

$$u''|_U : (H/K')|_U \longrightarrow G|_U$$

is a monomorphism. Thus, after this restriction,

$$\text{Ker}(u) = \text{Ker}(u') = K',$$

which proves that  $\text{Ker}(u)$  is of multiplicative type.

The same proof applies if, instead of assuming  $G$  of finite presentation over  $S$ , one assumes  $S$  locally Noetherian, at least when  $H$  is diagonalizable. Without this additional hypothesis on  $H$ , one must show that there is a covering base change

$$S' \longrightarrow S,$$

with  $S'$  locally Noetherian, which trivializes  $H$ . This will be proved in the next exposé, X 4.6. □

---

<sup>45</sup>Editor's note: the erroneous reference to IV 4.7.5 has been corrected. Note that Section 8 of the present exposé is independent of Sections 2 through 7.

<sup>46</sup>Editor's note: "induced by  $G$ " has been corrected to "induced by  $u$ ."



## 7. Algebraicity of Formal Homomorphisms into an Affine Group

**Theorem 7.1.** *Let  $A$  be a Noetherian ring equipped with an ideal  $I$  for which  $A$  is separated and complete in the  $I$ -adic topology. Put*

$$S = \operatorname{Spec}(A), \quad S_n = \operatorname{Spec}(A/I^{n+1}),$$

*and let  $H, G$  be  $S$ -group preschemes, with  $H$  of isotrivial multiplicative type and  $G$  affine. Then the canonical map*

$$(x) \quad \theta : \operatorname{Hom}_{S\text{-gr}}(H, G) \longrightarrow \varprojlim_n \operatorname{Hom}_{S_n\text{-gr}}(H_n, G_n)$$

*is bijective, where  $H_n, G_n$  are the  $S_n$ -groups obtained from  $H, G$  by the base change  $S_n \rightarrow S$ .*

*Proof.* Suppose first that  $H$  is diagonalizable, hence of the form

$$H = \operatorname{Spec} A(M),$$

where  $B = A(M)$  is the algebra of the commutative group  $M$  with coefficients in  $A$ . We may also write

$$G = \operatorname{Spec}(C),$$

where  $C$  is an  $A$ -algebra equipped with a diagonal map satisfying the usual axioms. Homomorphisms of  $S$ -groups  $H \rightarrow G$  correspond bijectively to homomorphisms of  $A$ -algebras

$$\varphi : C \longrightarrow B$$

compatible with the diagonal maps; that is, such that, for every  $f \in C$ ,

$$\Delta_H(\varphi(f)) = (\varphi \otimes \varphi)(\Delta_G(f)).$$

There is an analogous description of the homomorphisms of  $S_n$ -groups  $H_n \rightarrow G_n$ , given by homomorphisms of  $A_n$ -algebras

$$\varphi_n : C_n \longrightarrow B_n,$$

where

$$A_n = A/I^{n+1}, \quad B_n = B \otimes_A A_n, \quad C_n = C \otimes_A A_n.$$

[The source prints  $C = C \otimes_A A_n$  in the last equality;  $C_n$  is intended.] Put

$$\widehat{B} = \varprojlim_n B_n, \quad \widehat{C} = \varprojlim_n C_n.$$

Then

$$\widehat{B} = \varprojlim_n A_n(M)$$

identifies with a submodule of the product  $A^M$ , namely the submodule of families  $(a_m)_{m \in M}$  of elements of  $A$  that tend to zero for the  $I$ -adic topology along the filter of complements of finite subsets of  $M$ . In particular, the canonical homomorphism

$$B \longrightarrow \widehat{B}$$

is injective.<sup>47</sup> The projective system

$$\theta(\varphi) = (\varphi_n)_{n \geq 0}$$

defines a homomorphism

$$\widehat{\varphi} : \widehat{C} \longrightarrow \widehat{B}$$

such that the following diagram is commutative:

$$\begin{array}{ccc} C & \xrightarrow{\varphi} & B \\ \downarrow & & \downarrow \\ \widehat{C} & \xrightarrow{\widehat{\varphi}} & \widehat{B}. \end{array}$$

We prove that  $\theta$  is surjective. Let  $(\varphi_n)_{n \geq 0}$  be a projective system; we must show that it is obtained by reduction from some  $\varphi$  in the left-hand side of (x). A priori, the system  $(\varphi_n)$  defines a homomorphism on the completed algebras,

$$\widehat{\varphi} : \widehat{C} \longrightarrow \widehat{B}.$$

It is enough to prove that the composite

$$\Phi : C \longrightarrow \widehat{C} \xrightarrow{\widehat{\varphi}} \widehat{B}$$

sends  $C$  into  $B$ . Indeed, in that case it gives a homomorphism of  $A$ -algebras

$$\varphi : C \longrightarrow B$$

whose reductions are the  $\varphi_n$ . This homomorphism is immediately seen to be compatible with the diagonal maps, since the  $\varphi_n$  are so, and since

$$B \otimes_A B \longrightarrow \widehat{B \otimes_A B}$$

is injective, as follows from the same argument after replacing  $M$  by  $M \times M$ .

Passing to the completions, the diagonal homomorphism  $\Delta_H$  defines

$$\widehat{\Delta}_H : \widehat{B} \longrightarrow \widehat{B \otimes_A B}.$$

Taking the projective limit of the analogous diagrams defined by the  $\varphi_n$ , one obtains the commutative diagram

$$\begin{array}{ccc} C & \xrightarrow{\Phi} & \widehat{B} \\ \Delta_G \downarrow & & \downarrow \widehat{\Delta}_H \\ C \otimes_A C & \xrightarrow{\Psi} & \widehat{B \otimes_A B}, \end{array}$$

where  $\Psi$  is the composite

$$C \otimes_A C \xrightarrow{\Phi \otimes \Phi} \widehat{B} \otimes_A \widehat{B} \longrightarrow \widehat{B \otimes_A B};$$

the last arrow is the evident canonical homomorphism. Consequently, for every  $f \in C$ , the element  $\Phi(f) \in \widehat{B}$  has the property that its image under  $\widehat{\Delta}_H$  is a “decomposable” element of  $\widehat{B \otimes_A B}$ , namely an element in the image of  $\widehat{B} \otimes_A \widehat{B}$ .

<sup>47</sup>Editor’s note: the argument that follows has been expanded.

Let  $(e_m)_{m \in M}$  be the canonical basis of  $A(M)$ , and let  $(e_{m,m'})$  be that of

$$A(M \times M) = A(M) \otimes_A A(M).$$

Since

$$\Delta_H(e_m) = e_m \otimes e_m = e_{m,m} \quad (m \in M),$$

it remains only to apply the following lemma.

**Lemma 7.2.** *Let  $A$  be a Noetherian ring, let  $M$  be a set, and let  $(a_{m,m'})$  be a family of elements of  $A$ , indexed by  $M \times M$ , satisfying:*

(i)  $a_{m,m'} = 0$  if  $m \neq m'$ ; that is, the support of the family is contained in the diagonal of  $M \times M$ ;

(ii) one has

$$a_{m,m'} = \sum_i b_m^i c_{m'}^i,$$

where the  $b^i, c^i$  are finitely many elements of  $A^M$ ; in other words,  $(a_{m,m'})$  belongs to the image of the canonical homomorphism

$$A^M \otimes_A A^M \longrightarrow A^{M \times M}.$$

Then the support of  $(a_{m,m'})$  is finite.

*Proof.* By (i), the family  $(a_{m,m'})$  is determined by the elements

$$a_m = a_{m,m}.$$

For  $x = (x_n)_{n \in M} \in A(M)$ , put

$$(u \cdot x)_m = \sum_{m'} a_{m,m'} x_{m'}.$$

Thus  $(a_{m,m'})$  is viewed as the matrix of a homomorphism

$$u : A(M) \longrightarrow A^M.$$

Condition (i) gives simply

$$(u \cdot x)_m = a_m x_m.$$

On the other hand, condition (ii) gives

$$u \cdot x \in \sum_i A \cdot b^i.$$

Hence  $u(A(M))$  lies in a finitely generated  $A$ -module. If  $(e_m)_{m \in M}$  denotes the canonical basis of

$$A(M) \subset A^M,$$

then all the elements  $a_m e_m$  lie in a finitely generated  $A$ -module. Since  $A$  is Noetherian, the module generated by these elements is itself finitely generated. Because the  $e_m$  are linearly independent, all but finitely many of the  $a_m$  are zero. This proves the lemma, and hence Theorem 7.1 when  $H$  is diagonalizable.  $\square$

We now prove the general case of 7.1, in which  $H$  is only assumed isotrivial. There is then a finite étale surjective morphism

$$S' \longrightarrow S$$

such that

$$H' = H \times_S S'$$

is diagonalizable. We shall use only the fact that  $S' \rightarrow S$  is finite and a covering for the faithfully flat quasi-compact topology, or simply for the canonical topology on **(Sch)**. Thus “étale” could be replaced by “flat.”

Put

$$S'' = S' \times_S S',$$

and similarly introduce

$$S'_n \quad \text{and} \quad S''_n = S'' \times_S S_n = S'_n \times_{S_n} S'_n,$$

together with  $H', G', H'', G'', H'_n, G'_n, H''_n, G''_n$ , obtained from  $H, G$  by the evident base changes. The groups  $H'$  and  $H''$  are diagonalizable. Moreover,  $S'$ , and hence  $S''$ , is affine. If

$$S' = \text{Spec}(A'), \quad S'' = \text{Spec}(A''),$$

then  $A'$  and  $A''$  are separated and complete for the topologies defined by  $IA'$  and  $IA''$ , respectively, because both are finite over  $A$ .

Since  $S' \rightarrow S$  and  $S'_n \rightarrow S_n$  are covering morphisms, we obtain a commutative diagram of maps of sets whose two rows are exact:

$$\begin{array}{ccccc} \text{Hom}_{S\text{-gr}}(H, G) & \longrightarrow & \text{Hom}_{S'\text{-gr}}(H', G') & \rightrightarrows & \text{Hom}_{S''\text{-gr}}(H'', G'') \\ \downarrow u & & \downarrow u' & & \downarrow u'' \\ \varprojlim_n \text{Hom}_{S_n\text{-gr}}(H_n, G_n) & \longrightarrow & \varprojlim_n \text{Hom}_{S'_n\text{-gr}}(H'_n, G'_n) & \rightrightarrows & \varprojlim_n \text{Hom}_{S''_n\text{-gr}}(H''_n, G''_n). \end{array}$$

The second row is exact because it is the projective limit of the exact diagrams associated with the various  $S'_n \rightarrow S_n$ . By the diagonalizable case already proved,  $u'$  and  $u''$  are bijective. Thus  $u$  is bijective as well, completing the proof of 7.1.  $\square$

**Corollary 7.3.** *Under the hypotheses of 7.1, suppose in addition that  $G$  is smooth over  $S$ , and let*

$$u_0 : H_0 \longrightarrow G_0$$

*be a homomorphism of  $S_0$ -groups. Then there is a homomorphism of  $S$ -groups*

$$u : H \longrightarrow G$$

*lifting  $u_0$ . Any two such liftings  $u, u'$  are conjugate by an element  $g \in G(S)$  whose reduction is the unit element of  $G_0(S_0)$ .*

*Proof.* This follows by combining 3.6 and 7.1. To construct  $u$ , construct the  $u_n$  successively, as is possible by 3.6; Theorem 7.1 then shows that the system  $(u_n)$  comes from an  $S$ -homomorphism  $u$ . Given two liftings  $u, u'$ , construct successively elements  $g_n$  such that

$$g_0 = 1, \quad g_n \text{ is the reduction of } g_{n+1}, \quad u'_n = \text{int}(g_n)u_n.$$

Again, this is possible by 3.6. Since  $A$  is separated and complete, the  $g_n$  come from an element  $g \in G(S)$ . Finally, the injectivity in 7.1 gives

$$u' = \text{int}(g)u.$$

□

**Remark 7.4.** Compare 7.1 with EGA III, 5.4.1, which implies that the statement of 7.1 remains true if, instead of assuming  $H$  of multiplicative type and  $G$  affine, one assumes that  $H$  is proper over  $S$ , and that  $G$  is separated and locally of finite type over  $S$ . A statement such as 7.1 without a properness hypothesis is rather exceptional, and should be understood here as one aspect of the great “rigidity” of the structure of a group of multiplicative type.

The analogous statement with

$$G = H = \mathbf{G}_a$$

is false in general. Indeed, take  $A$  of characteristic  $p > 0$ , and define the  $u_n$ , by reduction modulo  $I^{n+1}$ , from an additive formal power series

$$u(T) = \sum_{n \geq 0} a_n T^{p^n},$$

where the  $a_n \in A$  tend to zero for the  $I$ -adic topology, but not for the discrete topology.

The statement of 7.1 also becomes false if the hypothesis that  $G$  be affine is omitted, even when  $H = \mathbf{G}_m$ . An example, with  $A$  a complete discrete valuation ring, is obtained from an elliptic curve over the fraction field  $K$  of  $A$  whose reduction, in the Néron–Kodaira theory, say, is the group  $\mathbf{G}_m$  over the residue field  $k$ .<sup>48</sup> One then has a smooth commutative group scheme  $G$  over  $S$ , with special fiber  $\mathbf{G}_{m,k}$ . By 3.6 this permits the construction of a projective system of isomorphisms

$$u_n : H_n \xrightarrow{\sim} G_n, \quad H = \mathbf{G}_{m,A},$$

while the generic fiber of  $G$  is an abelian variety. Consequently, there is no homomorphism of  $S$ -groups  $H \rightarrow G$  other than the zero homomorphism.

## 8. Subgroups, Quotient Groups, and Extensions of Groups of Multiplicative Type over a Field\*

**Scholium 8.0.**<sup>49</sup> — Let  $k$  be a field and let  $H$  be an affine  $k$ -group scheme. Write  $k[H]$  for its affine algebra and

$$X(H) = \text{Hom}_{k\text{-gr}}(H, \mathbf{G}_{m,k})$$

for the group of characters of  $H$ . By the independence lemma for characters,  $X(H)$  is a linearly independent subset of  $k[H]$ . Consequently,  $H$  is diagonalizable if and only if  $X(H)$  spans  $k[H]$  as a  $k$ -vector space.

**Proposition 8.1.** — Let  $G$  be a diagonalizable group scheme (respectively, a group scheme of multiplicative type) over a field  $k$ .<sup>50</sup> Then, for every subgroup scheme  $H$  of  $G$ , the schemes  $H$  and  $G/H$  are diagonalizable (respectively, of multiplicative type).

<sup>48</sup>Editor’s note: such an example could be made explicit.

\*Added in July 1969. This afterthought section is independent of Sections 3 through 7.

<sup>49</sup>Editor’s note: this scholium has been added.

<sup>50</sup>Editor’s note: the words “over a field  $k$ ,” implicit in the original, have been added. In addition, “algebraic group” has been replaced by “group scheme,” since  $G$  is not assumed to be of finite type.

Definition 1.1 immediately reduces the assertion to the case without the parenthetical qualification. An easy passage to the limit, using VI<sub>B</sub> 11.13 and VIII 3.1, reduces us to the case in which  $G$  is of finite type over  $k$ .<sup>51</sup> By VI<sub>B</sub> 11.16,<sup>52</sup> one can find a finite family of nonzero elements

$$(8.1.1) \quad f_i = \sum_m a_{im} m, \quad a_{im} \in k,$$

of the affine ring  $k(M)$  of  $G = D_k(M)$ , such that the points of  $H$ , with values in an arbitrary  $k$ -algebra  $k'$ , are precisely the points  $g \in G(k')$  for which

$$(8.1.2) \quad \tau_g f_i = \lambda_i(g) f_i, \quad \lambda_i(g) \in k',$$

where  $\tau_g$  denotes translation by  $g$ . But

$$(8.1.3) \quad \tau_g f_i = \sum_m a_{im} \chi_m(g) m,$$

where  $\chi_m : G \rightarrow \mathbf{G}_{m,k}$  is the character associated with  $m$ . Thus (8.1.2) is equivalent to

$$(8.1.4) \quad \chi_m(g) = \chi_{m'}(g) \quad \text{if } m, m' \in Z_i,$$

where  $Z_i$  denotes the set of  $m \in M$  for which  $a_{im} \neq 0$ . The same relation may be written

$$\chi_{m'-m}(g) = 1 \quad \text{if } m, m' \in Z_i.$$

Let  $N$  be the subgroup of  $M$  generated by all the elements  $m' - m$ , where  $i$  varies and  $m, m' \in Z_i$ . It follows from the definition of  $D_k(M) \simeq G$  that  $H$  identifies with  $D_k(M/N)$ . Hence VIII 3.1 identifies  $G/H$  with  $D_k(N)$ .  $\square$

**Proposition 8.2.** <sup>53</sup> — *Let  $k$  be a field; let  $H$  and  $K$  be  $k$ -group schemes of multiplicative type and finite type; and let  $G$  be a  $k$ -group scheme fitting into an exact sequence*

$$1 \longrightarrow H \longrightarrow G \longrightarrow K \longrightarrow 1$$

(which implies that  $G$  is of finite type over  $k$ ).

(i) *If  $G$  is commutative or  $K$  is connected, then  $G$  is of multiplicative type.*

(ii) *If  $K$  and  $H$  are diagonalizable and  $K$  is a torus, then  $G$  is diagonalizable.*

For the proof of (i), the reader is referred to XVII 7.1.1, of which 8.2(i) is a special case. The case of a field is treated in part (i) of the proof of XVII 7.1.1, without using the results of the later exposés.

For (ii), now suppose that  $K$  and  $H$  are diagonalizable:

$$K \simeq D_k(M) \quad \text{and} \quad H \simeq D_k(N),$$

<sup>51</sup>Editor's note: this "passage to the limit" should be spelled out. It uses 8.0 and the equality

$$G/H = \varprojlim_i G_i/H_i$$

(cf. VI<sub>B</sub>, proof of 11.17). To see that  $H = \varprojlim_i (H \cap G_i)$ , does one use the fact that  $H$  is closed in  $G$ ? Using this fact, one can give a direct proof.

<sup>52</sup>Editor's note: VI<sub>B</sub> 11.16 should doubtless be rewritten in the usual, more agreeable form. In particular, a single  $f_i$  suffices below.

<sup>53</sup>Editor's note: both the statement and its proof have been expanded.

and suppose that  $G$  is of multiplicative type, as follows from (i) when  $K$  is a torus. By X 1.4,  $G$  is isotrivial over  $k$ ; that is, there is a finite separable extension  $k'/k$  such that

$$G' = G \times_k k'$$

is diagonalizable. Hence  $G' = D_{k'}(E)$  for a commutative group  $E$ , and there is an exact sequence

$$(1) \quad 0 \longrightarrow D_{k'}(N) \longrightarrow D_{k'}(E) \longrightarrow D_{k'}(M) \longrightarrow 0.$$

By VIII 3.1 and 3.2,  $M$  is therefore a subgroup of  $E$ , and one has an exact sequence

$$(2) \quad 0 \longrightarrow M \longrightarrow E \longrightarrow N \longrightarrow 0.$$

For a given extension  $E_0$  of  $N$  by  $M$ , consider the diagonalizable  $k$ -group  $G_0 = D_k(E_0)$ . The  $k$ -group functor  $A$  of automorphisms of the extension

$$(3) \quad 1 \longrightarrow H \longrightarrow G_0 \longrightarrow K \longrightarrow 1,$$

that is, the group subfunctor of  $\text{Aut}_{k\text{-gr}}(G_0)$  whose points on a  $k$ -prescheme  $T$  are the

$$\phi \in \text{Aut}_{T\text{-gr}}(G_T)$$

inducing the identity on  $H_T$  and  $K_T$ , identifies with  $\text{Hom}_{k\text{-gr}}(K, H)$ . By VIII 1.5, this is the constant  $k$ -group with value

$$L = \text{Hom}_{\text{gr}}(N, M).$$

It follows that the extensions  $G$  of  $K$  by  $H$  which become isomorphic to (3) over a separable closure  $k_s$  of  $k$  are classified in the same way as the  $k$ -torsors, for the étale topology, under the constant group  $L_k$ . Put  $\Gamma = \text{Gal}(k_s/k)$ . Since  $L$  is a trivial  $\Gamma$ -module, these torsors are classified by

$$H_{\text{ét}}^1(k, L_k) = H^1(\Gamma, L) = \text{Hom}_{\text{gr. top.}}(\Gamma, L).$$

If  $K$  is moreover a torus, then  $M$  is torsion-free, and hence so is  $L$ . Therefore

$$\text{Hom}_{\text{gr. top.}}(\Gamma, L) = 0,$$

and every extension of  $K$  by  $H$  is already diagonalizable.

**Remark 8.3.** — *As already noted in the proof of 8.2, the first assertion is stated and proved over an arbitrary base in XVII 7.1.1. The second assertion generalizes, with essentially the same proof, to the case of an integral normal base scheme—or, more generally, a geometrically unibranch one—by using X 5.13 below.*

*Finally, 8.1 may also be regarded as a corollary of the distinctly less trivial result 6.8, which is likewise valid over an arbitrary base scheme.*<sup>54</sup>

## Bibliography

55

[BEns] N. Bourbaki, *Théorie des ensembles*, Chapters I–IV, Hermann, 1970.

<sup>54</sup>Editor's note: the proof of 6.8 nevertheless uses 8.1.

<sup>55</sup>Editor's note: additional references cited in this exposé.

# Working-state notice for Exposé X

The complete Exposé X Loop-1 English text and equations are included below. All 44 authority pages were reviewed by the producing top-level session and the standalone reader completed three clean build passes. Four raster diagram placeholders remain and are explicitly deferred to Loop 2. This is readable current-progress text, not a claim of native-diagram or exhaustive release certification.



# Exposé X

## Characterization and Classification of Groups of Multiplicative Type

by A. Grothendieck

### 1. Classification of Isotrivial Groups. Case of a Base Field

Recall (IX 1.1) that a group of multiplicative type  $H$  over the prescheme  $S$  is said to be *isotrivial* if there exists a finite étale surjective morphism  $S' \rightarrow S$  such that  $H' = H \times_S S'$  is diagonalizable. When  $S$  is connected,  $S'$  decomposes as a finite disjoint sum of connected components  $S'_i$ , and hence, after replacing  $S'$  by one of the  $S'_i$ , we may suppose that  $S'$  is connected. Finally,  $S'$  can be dominated by a connected finite étale  $S'_1$  that is Galois, that is, a principal homogeneous bundle over  $S$  with group  $\Gamma_S$ , where  $\Gamma$  is an ordinary finite group (cf. SGA 1, V, nos. 7 and 3 when  $S$  is locally noetherian, and EGA IV<sub>4</sub>, 18.2.9 in general).<sup>1</sup> We suppose  $S'$  chosen in this way and propose to determine the groups of multiplicative type  $H$  over  $S$  that are “trivialized” by  $S'$ , that is, those for which  $H' = H \times_S S'$  is diagonalizable.<sup>2</sup> By descent theory (cf. SGA 1, VIII, 5.4), the category of such  $H$  is equivalent to the category of diagonalizable groups  $H'$  over  $S'$ , equipped with actions of  $\Gamma$  on  $H'$  compatible with the actions of  $\Gamma$  on  $S'$ . (N.B. Since the groups under consideration are affine over the base, every descent datum here is effective; cf. SGA 1, VIII, 2.1.) Since  $S'$  is connected, the contravariant functor

$$M \longmapsto D_{S'}(M)$$

is an anti-equivalence from the category of ordinary commutative groups to the category of diagonalizable groups over  $S'$  (cf. VIII 1.6); a quasi-inverse is<sup>3</sup>

$$H \longmapsto \mathrm{Hom}_{S'\text{-gr}}(H, \mathbb{G}_{m,S'}).$$

**Proposition 1.1.** — *Let  $S$  be a connected prescheme and let  $S'$  be a connected principal cover of  $S$  with finite group  $\Gamma$ . Then the category of groups of multiplicative type over  $S$*

<sup>0</sup>xy version of November 6, 2009: Addenda placed in Section 9; reviewed through 8.8.

<sup>1</sup>Editor’s note: More precisely, this follows from the “axiomatic conditions of a Galois theory,” cf. SGA 1, V, no. 4 g); when  $S$  is locally noetherian, the axioms are verified in *loc. cit.*, nos. 7 and 3, especially 3.7, which rests on SGA 1, I, 10.9. The latter result is proved without noetherian hypotheses in EGA IV<sub>4</sub>, 18.2.9.

<sup>2</sup>Editor’s note: In what follows, an  $S$ -group of multiplicative type will be called “split” if it is “trivial” in the sense of IX 1.2, that is, if it is a diagonalizable  $S$ -group.

<sup>3</sup>Editor’s note:  $S$  has been corrected to  $S'$ .

split by  $S'^4$  is anti-equivalent to the category of  $\Gamma$ -modules, that is, of ordinary commutative groups  $M$  equipped with a homomorphism

$$\Gamma \longrightarrow \mathrm{Aut}_{\mathrm{gr}}(M).$$

The following standard consequence results.

**Corollary 1.2.** — *Let  $S$  be a connected prescheme, and let  $\xi : \mathrm{Spec}(\Omega) \rightarrow S$  be a “geometric point” of  $S$ , that is, a morphism into  $S$  from the spectrum of an algebraically closed field  $\Omega$ . Consider the fundamental group of  $S$  at  $\xi$  (cf. SGA 1, V, no. 7):*

$$\pi = \pi_1(S, \xi).$$

*Then the category of isotrivial groups of multiplicative type  $H$  over  $S$  is anti-equivalent to the category of “Galois modules” under  $\pi$ , that is, of  $\pi$ -modules  $M$  such that the stabilizer in  $\pi$  of every element of  $M$  is an open subgroup.*

*Under this correspondence, the group associated with an isotrivial group of multiplicative type  $H$  is*

$$M = \mathrm{Hom}_{\Omega\text{-gr}}(H_\xi, \mathbb{G}_{m, \Omega}),$$

*where  $H_\xi$  is the fiber of  $H$  at  $\xi$ ; this group is naturally endowed with an action of  $\pi_1(S, \xi)$ .*

**Remark 1.3.** — We shall see below (cf. 5.16) that if  $S$  is normal, or more generally geometrically unibranch, then every group of multiplicative type and of finite type over  $S$  is necessarily isotrivial. Consequently, the classification principle of 1.2 applies to groups of multiplicative type and of finite type over  $S$ ; they correspond to Galois  $\pi$ -modules that are of finite type over  $\mathbb{Z}$ . For the moment, let us confine ourselves to the following result.

**Proposition 1.4.** — *Let  $k$  be a field and  $H$  a group of multiplicative type and of finite type over  $k$ . Then  $H$  is isotrivial; that is, there exists a finite separable extension  $k'$  of  $k$  that splits  $H$ .*

*Consequently, by 1.2, if  $\pi$  is the topological Galois group of an algebraic closure  $\Omega$  of  $k$ , the category of groups  $H$  of multiplicative type and of finite type over  $k$  is anti-equivalent to the category of Galois modules  $M$  under  $\pi$  that are of finite type as  $\mathbb{Z}$ -modules.*

Since  $H$  is of finite type over  $k$ , the “principle of finite extension” (cf. EGA IV<sub>3</sub>, 9.1.4) first gives a finite extension  $k'$  of  $k$  that splits  $H$ . Recall the argument. By hypothesis, there exist a diagonalizable group  $G$  of finite type over  $k$ , a faithfully flat  $S'$  over  $S = \mathrm{Spec}(k)$ , and an isomorphism of  $S'$ -groups

$$H_{S'} \simeq G_{S'}.$$

After replacing  $S'$  by the residue field of one of its points, we may suppose that  $S'$  is the spectrum of an extension  $K$  of  $k$ . The field  $K$  is the filtered direct limit of its finitely generated  $k$ -subalgebras  $A_i$ ; it follows readily (cf. EGA IV<sub>3</sub>, 8.8.2.4) that the given isomorphism  $u$  comes, for sufficiently large  $i$ , from an  $A_i$ -isomorphism

$$u_i : H_{A_i} \simeq G_{A_i}.$$

By the Nullstellensatz,  $A_i$  has a quotient ring  $k'$  that is a finite extension of  $k$ . Thus  $k'$  splits  $H$ .

Now  $k'$  is a radical extension of a separable extension  $k'_s$  of  $k$ . By IX 5.4, the isomorphism

$$u' : H_{k'} \simeq G_{k'}$$

---

<sup>4</sup>Editor’s note: Here and below, “splittés” has been replaced in the French text by “déployés,” and “splitte” by “déploie,” etc. Both are rendered here by “split.”


comes from an isomorphism  $H_{k'_s} \simeq G_{k'_s}$ . Thus  $k'_s$  splits  $H$ , proving 1.4.

**Remark 1.5.** — Statement 1.2 gives, in particular, a characterization of the isotrivial tori over  $S$  of relative dimension  $n$ : setting  $\pi = \pi_1(S, \xi)$ , they correspond to classes, up to “equivalence,” of representations

$$\pi \longrightarrow \mathrm{GL}(n, \mathbb{Z})$$

whose kernel is an open subgroup of  $\pi$ .

**1.6.** Even when  $S$  is an algebraic curve, there may exist tori over  $S$  (of relative dimension 2) that are not locally isotrivial (and *a fortiori* not isotrivial); there may also exist locally trivial tori that are not isotrivial. (Observe, however, that such phenomena can occur only when  $S$  is not normal, as already noted in 1.3.) For example, let  $S$  be an irreducible algebraic curve (over an algebraically closed field, to fix ideas) with an ordinary double point  $a$ , let  $S'$  be its normalization, and let  $b$  and  $c$  be the two points of  $S'$  above  $a$ . A connected principal homogeneous bundle  $P$  over  $S$ , with structural group  $\mathbb{Z}$ , is obtained by attaching infinitely many copies of  $S'$  according to the diagram



(N.B. This is a principal bundle for the étale topology.) There is a homomorphism

$$\varphi : \mathbb{Z} \longrightarrow \mathrm{GL}(2, \mathbb{Z}), \quad \varphi(n) = \begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix},$$

which makes it possible to construct a torus  $T$  over  $S$ , of relative dimension 2, from the trivial torus  $\mathbb{G}_m^2$  over  $P$  and the descent datum on it deduced from  $\varphi$ . (N.B. The projection  $P \rightarrow S$  is a covering for the étale topology and *a fortiori* for the canonical topology of (Sch); the descent datum in question is necessarily effective because  $\mathbb{G}_{m,P}^2$  is affine over  $P$ .) It is not difficult to prove that  $T$  is not isotrivial in a neighborhood of  $a^5$  (although it is trivial on  $S - \{a\}$ ).

A variant of this construction is obtained by taking for  $S$  a curve with two irreducible components  $S_1$  and  $S_2$  meeting in two points  $a'$  and  $a''$ . This makes it possible to construct a connected, locally trivial principal homogeneous bundle  $P$  over  $S$  with group  $\mathbb{Z}_S$ , and hence an associated torus  $T$  that is locally trivial but not isotrivial.

## 2. Infinitesimal Variations of Structure

<sup>6</sup> Let us begin by recalling the following result (cf. SGA 1, I, 8.3 when  $S$  is locally noetherian, and EGA IV<sub>4</sub>, 18.1.2 in general).

**Recall 2.0.** — Let  $S$  be a prescheme and  $S_0$  a subprescheme with the same underlying set. Then the functor

$$X \longmapsto X_0 = X \times_S S_0$$

is an equivalence between the category of étale preschemes over  $S$  and the analogous category over  $S_0$ .

<sup>5</sup>Editor's note: See 7.3 below.

<sup>6</sup>Editor's note: This “recall” has been added for later references.

**Proposition 2.1.** — *Let  $S$  be a prescheme and  $S_0$  a subprescheme with the same underlying set (that is, defined by a nilideal  $\mathcal{I}$ ). Then the functor*

$$H \mapsto H_0 = H \times_S S_0$$

*from the category of group preschemes of multiplicative type over  $S$  to the analogous category over  $S_0$  is fully faithful.*

*Moreover, it induces an equivalence between the category of quasi-isotrivial groups of multiplicative type over  $S$  and the analogous category over  $S_0$ .*

We first prove full faithfulness, that is, if  $H$  and  $G$  over  $S$  are of multiplicative type, then

$$\mathrm{Hom}_{S\text{-gr}}(H, G) \longrightarrow \mathrm{Hom}_{S_0\text{-gr}}(H_0, G_0)$$

is bijective. The question is local on  $S$ , so we may suppose that  $S$  is affine. There then exists a faithfully flat quasi-compact morphism  $S' \rightarrow S$  that splits  $H$  and  $G$ . Let  $S'' = S' \times_S S'$ ; write  $H', G'$ , respectively  $H'', G''$ , for the groups obtained from  $H, G$  by the base changes  $S' \rightarrow S$ , respectively  $S'' \rightarrow S$ . Define  $S'_0$  and  $S''_0$  similarly; the latter is also isomorphic to  $S'_0 \times_{S_0} S'_0$ . We obtain a commutative diagram with exact rows:

$$\begin{array}{ccccc} \mathrm{Hom}_{S\text{-gr}}(H, G) & \longrightarrow & \mathrm{Hom}_{S'\text{-gr}}(H', G') & \rightrightarrows & \mathrm{Hom}_{S''\text{-gr}}(H'', G'') \\ \downarrow u & & \downarrow u' & & \downarrow u'' \\ \mathrm{Hom}_{S_0\text{-gr}}(H_0, G_0) & \longrightarrow & \mathrm{Hom}_{S'_0\text{-gr}}(H'_0, G'_0) & \rightrightarrows & \mathrm{Hom}_{S''_0\text{-gr}}(H''_0, G''_0), \end{array}$$

Thus, to prove that  $u$  is bijective, it is enough to prove that  $u'$  and  $u''$  are bijective. This reduces us to the case where  $H$  and  $G$  are diagonalizable, hence of the form  $D_S(M)$  and  $D_S(N)$ , where  $M$  and  $N$  are ordinary commutative groups. Likewise,

$$H_0 = D_{S_0}(M), \quad G_0 = D_{S_0}(N).$$

We then have a commutative diagram<sup>7</sup>

$$\begin{array}{ccc} \mathrm{Hom}_{S\text{-gr}}(N_S, M_S) & \xrightarrow{\sim} & \mathrm{Hom}_{S\text{-gr}}(D_S(M), D_S(N)) \\ \downarrow & & \downarrow \\ \mathrm{Hom}_{S_0\text{-gr}}(N_{S_0}, M_{S_0}) & \xrightarrow{\sim} & \mathrm{Hom}_{S_0\text{-gr}}(D_{S_0}(M), D_{S_0}(N)), \end{array}$$

whose horizontal arrows are isomorphisms by VIII 1.4. We are therefore reduced to proving that the homomorphism

$$\mathrm{Hom}_{S\text{-gr}}(N_S, M_S) \longrightarrow \mathrm{Hom}_{S_0\text{-gr}}(N_{S_0}, M_{S_0}) \tag{x}$$

is bijective; in other words, the functor  $M_S \mapsto M_{S_0}$ , from constant commutative group preschemes over  $S$  to constant commutative group preschemes over  $S_0$ , is fully faithful. Now (x) is also identified with the natural map

$$\mathrm{Hom}_{\mathrm{gr}}(N, \Gamma(M_S)) \longrightarrow \mathrm{Hom}_{\mathrm{gr}}(N, \Gamma(M_{S_0}))$$

<sup>7</sup>Editor's note: The original has been corrected by interchanging  $M$  and  $N$  in the right-hand terms.

deduced from  $\Gamma(M_S) \rightarrow \Gamma(M_{S_0})$ . The latter map is plainly bijective, since  $\Gamma(M_S)$ , the set of locally constant maps from  $S$  to  $M$ , depends only on the underlying topological space of  $S$ . This proves the desired assertion.

To prove the second assertion of 2.1, it remains to show that every quasi-isotrivial group  $H_0$  of multiplicative type over  $S_0$  comes from a quasi-isotrivial group  $H$  of multiplicative type over  $S$ . Let  $S'_0 \rightarrow S_0$  be an étale surjective morphism that splits  $H_0$ .

By 2.0, there exist an étale morphism  $S' \rightarrow S$  and an  $S_0$ -isomorphism

$$S' \times_S S_0 \simeq S'_0.$$

Thus we may suppose that  $S'_0$  is obtained from  $S'$  by reduction. Since  $H'_0$  is diagonalizable, it is immediately seen to be isomorphic to the group obtained by the base change  $S'_0 \rightarrow S'$  from a diagonalizable group  $H'$  over  $S'$ . (N.B. If  $H'_0 = D_{S'_0}(M)$ , take  $H' = D_{S'}(M)$ .) As usual, set

$$S'' = S' \times_S S', \quad S''' = S' \times_S S' \times_S S',$$

and similarly define  $S''_0$  and  $S'''_0$  by base change from  $S$  to  $S_0$ . They are also isomorphic to the fibered square and fibered cube, respectively, of  $S'_0$  over  $S_0$ . Applying the full faithfulness already proved to  $(S'', S''_0)$  and  $(S''', S'''_0)$ , we see that the natural descent datum on  $H'_0$  relative to  $S'_0 \rightarrow S_0$  (cf. IV 2.1) comes from a uniquely determined descent datum on  $H'$  relative to  $S' \rightarrow S$ . This descent datum is effective because  $H'$  is affine over  $S'$  (SGA 1, VIII, 2.1). Hence there exists an  $S$ -group  $H$  such that

$$H \times_S S' = H' = D_{S'}(M),$$

and  $H$  is therefore a quasi-isotrivial group of multiplicative type.

Using the full-faithfulness result for  $(S', S'_0)$ , one now checks readily that the given isomorphism between  $H'_0$  and  $H' \times_{S'} S'_0$  comes from an isomorphism between  $H_0$  and  $H \times_S S_0$ . (For a more formal presentation of results of this kind, see Giraud's article in preparation on descent theory.\*)

**Corollary 2.2.** — *Let  $H$  be an  $S$ -group of multiplicative type, and set  $H_0 = H \times_S S_0$ . Then  $H$  is quasi-isotrivial (respectively, locally isotrivial, isotrivial, locally trivial, or trivial) if and only if  $H_0$  has the corresponding property.*

The “only if” direction is immediate. The “if” direction has already been proved in the quasi-isotrivial case: by full faithfulness, it is enough to know that every quasi-isotrivial group over  $S_0$  lifts to a quasi-isotrivial group over  $S$ . The same argument applies to “trivial.” For “isotrivial,” repeat the argument establishing the second assertion of 2.1, but take  $S'_0 \rightarrow S_0$  to be finite étale and surjective. The locally isotrivial and locally trivial cases follow at once from the isotrivial and trivial cases.

When  $\mathcal{I}$  is nilpotent, 2.2 can be generalized slightly without assuming *a priori* that  $H$  is of multiplicative type.<sup>8</sup>

**Corollary 2.3.** — *Suppose that the ideal  $\mathcal{I}$  defining  $S_0$  is locally nilpotent. Let  $H$  be a group prescheme over  $S$ , flat over  $S$ , and set  $H_0 = H \times_S S_0$ . Then  $H$  is of quasi-isotrivial multiplicative type if and only if  $H_0$  is so.*

Suppose that  $H_0$  is of quasi-isotrivial multiplicative type. We prove that the same is true of  $H$ . The question is local for the étale topology, and by 2.0 the category of étale preschemes over  $S$  is equivalent to the category of étale preschemes over  $S_0$  by the functor

\*Cf. J. Giraud, “Méthode de la descente,” *Bull. Soc. Math. France*, Mémoire 2 (1964).

<sup>8</sup>Editor's note: Recall that 2.3 is used in proving Theorem IX 3.6 bis.

$S' \mapsto S' \times_S S_0$ . We are therefore immediately reduced to the case where  $H_0$  is diagonalizable, hence isomorphic to  $D_{S_0}(M)$ . Let  $G = D_S(M)$ . We then have an isomorphism

$$u_0 : H_0 \xrightarrow{\sim} G_0.$$

I claim that  $u_0$  comes from a unique homomorphism  $u : H \rightarrow G$ , which is necessarily an isomorphism because  $u_0$  is one,  $H$  and  $G$  are flat over  $S$ , and  $\mathcal{I}$  is locally nilpotent.<sup>9</sup> This will prove 2.3. We have (cf. VIII 1 (xxx))

$$\mathrm{Hom}_{S\text{-gr}}(H, G) \simeq \mathrm{Hom}_{S\text{-gr}}(M_S, \underline{\mathrm{Hom}}_{S\text{-gr}}(H, \mathbb{G}_{m,S})),$$

and the right-hand side is also identified with

$$\mathrm{Hom}_{\mathrm{gr}}(M, \mathrm{Hom}_{S\text{-gr}}(H, \mathbb{G}_{m,S})).$$

Thus the homomorphism

$$\mathrm{Hom}_{S\text{-gr}}(H, G) \longrightarrow \mathrm{Hom}_{S_0\text{-gr}}(H_0, G_0) \quad (\mathrm{x})$$

is isomorphic to the homomorphism

$$\mathrm{Hom}_{\mathrm{gr}}(M, \mathrm{Hom}_{S\text{-gr}}(H, \mathbb{G}_{m,S})) \longrightarrow \mathrm{Hom}_{\mathrm{gr}}(M, \mathrm{Hom}_{S_0\text{-gr}}(H_0, \mathbb{G}_{m,S_0}))$$

deduced from the restriction homomorphism

$$\mathrm{Hom}_{S\text{-gr}}(H, \mathbb{G}_{m,S}) \longrightarrow \mathrm{Hom}_{S_0\text{-gr}}(H_0, \mathbb{G}_{m,S_0}). \quad (\mathrm{xx})$$

It is therefore enough to prove that (xx) is bijective. The question is local on  $S$ , so we may suppose that  $S$  is affine. Since  $\mathbb{G}_{m,S}$  is commutative and smooth over  $S$ , IX 3.6 applies and completes the proof.

**Corollary 2.4.** — *Let  $A$  be an artinian local ring with residue field  $k$ ,  $S = \mathrm{Spec}(A)$ , and  $S_0 = \mathrm{Spec}(k)$ .*

- (i) <sup>10</sup> *Let  $H$  be an  $S$ -group prescheme, flat and locally of finite type, such that  $H_0 = H \times_S S_0$  is of multiplicative type. Then  $H$  is of multiplicative type, of finite type, and isotrivial. In particular, every  $S$ -group prescheme  $H$  of multiplicative type and of finite type is isotrivial.*
- (ii) *The functor  $H \mapsto H_0$  is an equivalence between the category of groups of multiplicative type and of finite type over  $A$  and the analogous category over  $k$ .*

Indeed, let  $H$  be as in (i). Then  $H_0$  is of multiplicative type and locally of finite type, hence of finite type, and therefore isotrivial by 1.4. By 2.3 and 2.2,  $H$  is consequently of multiplicative type (and hence of finite type) and isotrivial. Assertion (ii) now follows from 2.1.

**Corollary 2.5.** — *Let  $S' \rightarrow S$  be a faithfully flat radicial morphism.*

<sup>9</sup>Editor's note: Since  $u_0$  is an isomorphism, it is enough to see that for every  $h \in H$ , the morphism  $\varphi : \mathcal{O}_{G,u(h)} \rightarrow \mathcal{O}_{H,h}$  is bijective. It is bijective after reduction modulo  $\mathcal{I} = \mathcal{I}_s$ , where  $s$  is the image of  $h$  in  $S$ ; hence its cokernel  $C$  satisfies  $C = \mathcal{I}C$ , so  $C = 0$  because  $\mathcal{I}$  is nilpotent. Since  $\mathcal{O}_{H,h}$  is flat over  $\mathcal{O}_{S,s}$ , the kernel  $K$  of  $\varphi$  likewise satisfies  $K = \mathcal{I}K$ , and hence  $K = 0$ .

<sup>10</sup>Editor's note: The statement has been rewritten, retaining only the nontrivial implication in order to highlight it.

(i) *The functor*

$$H \longmapsto H' = H \times_S S'$$

*from the category of groups of multiplicative type over  $S$  to the analogous category over  $S'$  is fully faithful. Moreover, it induces an equivalence between the subcategories of quasi-isotrivial groups of multiplicative type.*

(ii) *If  $H$  is of multiplicative type, then  $H$  is quasi-isotrivial (respectively, locally isotrivial, isotrivial, locally trivial, or trivial) if and only if  $H'$  has the corresponding property.*

Indeed, let

$$S'' = S' \times_S S', \quad S''' = S' \times_S S' \times_S S'.$$

The assumption that  $S' \rightarrow S$  is radicial implies that the diagonal immersions  $S' \rightarrow S''$  and  $S' \rightarrow S'''$  are surjective, so base change by either immersion is governed by 2.1 and 2.2. Taking into account that  $S' \rightarrow S$  is a morphism of effective descent for the fibered category of groups of multiplicative type over preschemes (because it is faithfully flat and quasi-compact<sup>11</sup>), the assertion follows formally from 2.1 and 2.2 (cf. the already cited article of Giraud for a formal argument).

### 3. Finite Variations of Structure: Complete Base Ring

**Lemma 3.1.** — *Let  $A$  be a noetherian ring equipped with an ideal  $I$  such that  $A$  is separated and complete for the  $I$ -adic topology; set  $S = \operatorname{Spec}(A)$  and  $S_0 = \operatorname{Spec}(A/I)$ . Let  $G$  and  $H$  be two  $S$ -group preschemes, with  $G$  of multiplicative type and isotrivial, and with  $H$  affine over  $S$ , flat over  $S$  at the points of  $H_0 = H \times_S S_0$ , where  $H_0$  is of quasi-isotrivial multiplicative type. Then the natural map*

$$\operatorname{Hom}_{S\text{-gr}}(G, H) \longrightarrow \operatorname{Hom}_{S_0\text{-gr}}(G_0, H_0)$$

*is bijective.*

For every integer  $n \geq 0$ , let

$$S_n = \operatorname{Spec}(A/I^{n+1}),$$

and let  $G_n, H_n$  be obtained from  $G, H$  by the base change  $S_n \rightarrow S$ . Since  $G/S$  is of isotrivial multiplicative type and  $H$  is affine over  $S$ , IX 7.1 shows that the natural homomorphism

$$\operatorname{Hom}_{S\text{-gr}}(G, H) \longrightarrow \varprojlim_n \operatorname{Hom}_{S_n\text{-gr}}(G_n, H_n)$$

is bijective. On the other hand, 2.3 shows that the  $H_n$  are of quasi-isotrivial multiplicative type, and 2.1 shows that all transition homomorphisms in the inverse system

$$(\operatorname{Hom}_{S_n\text{-gr}}(G_n, H_n))_n$$

are isomorphisms. Lemma 3.1 follows at once.

**Theorem 3.2.** — *Let  $A$  be a noetherian ring equipped with an ideal  $I$  such that  $A$  is separated and complete for the  $I$ -adic topology, and set  $S = \operatorname{Spec}(A)$  and  $S_0 = \operatorname{Spec}(A/I)$ . Then:*

---

<sup>11</sup>Editor's note: Spell out this point ...

(i) *The functor*

$$H \longmapsto H_0 = H \times_S S_0$$

*is an equivalence between the category of isotrivial groups of multiplicative type over  $S$  and the analogous category over  $S_0$ .*

(ii) *Let  $H$  be an  $S$ -group of multiplicative type and of finite type. Then  $H$  is isotrivial if and only if  $H_0$  is isotrivial.*

*Proof.* For (i), one may either repeat the proof of 2.1 or use 1.2. In either case one uses the fact that the functor

$$S' \longmapsto S'_0 = S' \times_S S_0$$

from the category of finite étale schemes over  $S$  to the category of finite étale schemes over  $S_0$  is an equivalence (SGA 1, I, 8.4). Equivalently, after reducing to the case where  $S$  is connected (that is,  $S_0$  is connected) and choosing a geometric point  $\xi$  of  $S_0$ , the canonical homomorphism

$$\pi_1(S_0, \xi) \longrightarrow \pi_1(S, \xi)$$

is an isomorphism.

We prove (ii), namely that if  $H_0$  is isotrivial, then  $H$  is isotrivial. By (i), there exist an isotrivial group  $G$  of multiplicative type over  $S$  and an  $S_0$ -isomorphism

$$u_0 : G_0 \xrightarrow{\sim} H_0.$$

<sup>12</sup> Since  $H$  is of finite type, so are  $H_0$  and  $G_0$ . Therefore, by IX 2.1 b), the type of  $G$  at every point of  $S$  is a finitely generated abelian group, and hence  $G$  is of finite type over  $S$ . Moreover, by 3.1,  $u_0$  comes from a homomorphism of  $S$ -groups

$$u : G \longrightarrow H.$$

Finally,  $G$  and  $H$  are groups of multiplicative type and of finite type over  $S$ , and  $u_0$  is an isomorphism. It follows from IX 2.9 that  $u$  is an isomorphism, since every neighborhood of  $S_0$  in  $S$  is equal to  $S$ .

**Corollary 3.3.** — *Let  $A$  be a complete noetherian local ring with residue field  $k$ .*

(i) *Every group of multiplicative type and of finite type over  $A$  is isotrivial.*

(ii) *The functor*

$$H \longmapsto H \otimes_A k = H_0$$

*is an equivalence between the categories of groups of multiplicative type and of finite type over  $A$  and over  $k$ .*

<sup>12</sup> First, (i) follows from 3.2 (ii) and 1.4. Then (ii) follows from 3.2 (i), taking into account that  $H$  is of finite type if and only if  $H_0$  is of finite type (cf. the proof of 3.2 (ii)).

**Remark 3.3.1.** — By 1.4, statement 3.3 gives a complete classification of groups of multiplicative type and of finite type over  $A$  in terms of the topological Galois group of an algebraic closure  $\Omega$  of  $k$ .

**Remarks 3.4.** — Under the hypotheses of 3.2 (that is,  $A$  noetherian, separated and complete for the  $I$ -adic topology, with no further hypothesis on  $A/I$ ), it will follow from no. 5 that the functor

$$H \longmapsto H_0$$

---

<sup>12</sup>Editor's note: The original has been expanded in what follows.



from the category of groups of multiplicative type and of finite type over  $S$  to the analogous category over  $S_0$  is still fully faithful, without any isotriviality hypothesis.<sup>13</sup>

It is not, however, essentially surjective in general. Indeed, there may exist a group  $H_0$  of multiplicative type and of finite type over  $S_0$ , even locally trivial if desired (but not isotrivial), that does not arise by reduction from a group  $H$  of multiplicative type over  $S$ .

To see this, take either example from 1.6 of a non-isotrivial group of multiplicative type over a nonnormal curve. The curve may plainly be taken affine, say  $S_0$ , and embedded in affine 2-space; it is therefore defined by an ideal  $J$  in  $k[X, Y]$ . Take  $A$  to be the completion of  $k[X, Y]$  for the  $J$ -adic topology. Then  $A$  is a regular ring of dimension 2, and *a fortiori* normal. We shall see in 5.16 that every group of multiplicative type and of finite type over  $S = \text{Spec}(A)$  is consequently isotrivial. Thus  $H_0$ , being of finite type and non-isotrivial, cannot come from a group  $H$  of multiplicative type over  $S$ , since such an  $H$  would necessarily be of finite type and hence isotrivial.

**Lemma 3.5.\*** — *Let  $S$  be a prescheme and  $u : G \rightarrow H$  a homomorphism of  $S$ -group preschemes that are locally of finite presentation and flat over  $S$ . Let  $U$  be the set of  $s \in S$  such that the homomorphism induced on the fibers,*

$$u_s : G_s \longrightarrow H_s,$$

*is flat (respectively, smooth, unramified, étale, or quasi-finite). Then  $U$  is open, and the restriction*

$$u|_U : G|_U \longrightarrow H|_U$$

*is flat (respectively, smooth, unramified, étale, or quasi-finite).*

Let  $V$  be the set of points at which  $u$  is flat (respectively, ...). It is known that  $V$  is open (cf. SGA 1, I through IV in the locally noetherian case, and EGA IV in general<sup>14</sup>), and for  $x \in G$  above  $s \in S$ , one has  $x \in V$  if and only if  $u_s : G_s \rightarrow H_s$  is flat (respectively, ...) at  $x$  (same reference; in the flat, smooth, and étale cases, the flatness of  $G$  and  $H$  over  $S$  is used here). Since  $u_s$  is a homomorphism of locally algebraic groups, this is also equivalent to  $u_s$  being flat (respectively, ...) everywhere (VI<sub>B</sub>, 1.3), that is, to  $s \in U$ . Therefore  $U$  is the inverse image of the open set  $V$  under the unit section of  $G$ , and hence is open. Moreover,  $V = G|_U$ , so  $G|_U \rightarrow H|_U$  is flat (respectively, ...), completing the proof. (N.B. In the unramified and quasi-finite cases, the flatness hypothesis on  $G$  and  $H$  is unnecessary.)

**Lemma 3.6.** — *Let  $H$  be a commutative algebraic group over a field  $k$  that admits an open subgroup  $G$  of multiplicative type. Then the family of subschemes  ${}_nH$ ,  $n > 0$ , is schematically dense in  $H$ . In particular, if  ${}_nH = {}_nG$  for every  $n > 0$ , then  $H = G$ .*

Here  ${}_nH$  denotes the kernel of  $n \cdot \text{id}_H$ .<sup>15</sup> Let  $\bar{k}$  be an algebraic closure of  $k$ . It is enough to show that the family  $({}_nH_{\bar{k}})_{n>0}$  is schematically dense in  $H_{\bar{k}}$ , for then  $({}_nH)_{n>0}$  is schematically dense in  $H$  (cf. IX 4.5). We may therefore suppose that  $k$  is algebraically closed, so that  $G = D_k(M)$ , where  $M$  is an ordinary finitely generated commutative group.

Let

$$M_0 = M / \text{Torsion}(M).$$

<sup>13</sup>Editor's note: Specify this reference, possibly adding a corollary in no. 5 ...

\*Cf. also VI<sub>B</sub>, 2.5, for more systematic developments of this kind.

<sup>14</sup>Editor's note: Specify these references. Since  $G, H$  are locally of finite presentation,  $u$  is locally of finite presentation (IV<sub>1</sub>, 1.4.3 (v)); one then applies IV<sub>3</sub>, 11.3.10 and 13.1.4 for flat and quasi-finite, and IV<sub>4</sub>, 17.4.1, 17.5.1, and 17.6.1 for unramified, smooth, and étale; compare VI<sub>B</sub>, 2.5.

<sup>15</sup>Editor's note: The preceding sentence has been added.

Then  $T = D_k(M_0)$  is the largest torus contained in  $G$ , and  $H/T$  is finite. Hence  $H(k)/T(k)$  is annihilated by an integer  $\nu > 0$ . There exist finitely many  $g_i \in H(k)$  such that

$$H = \coprod_i g_i \cdot G,$$

and  $g'_i \in T(k)$ . Since  $k$  is algebraically closed, the map  $\nu \cdot \text{id}_T$  is surjective on

$$T(k) \simeq (k^\times)^d.$$

After replacing  $g_i$  by  $g_i t_i^{-1}$ , where  $t_i \in T(k)$  satisfies  $t_i^\nu = g'_i$ , we may therefore suppose that  $g_i^\nu = 1$ . If  $n$  is now a multiple of  $\nu$ , then

$${}_n H \supset g_i \cdot {}_n G.$$

As  $n > 0$  varies, the family  ${}_n G$  is schematically dense in  $G$  by IX 4.7, and the conclusion of 3.6 follows.

**Theorem 3.7.** — *Let  $A$  be a noetherian ring equipped with an ideal  $I$  such that  $A$  is separated and complete for the  $I$ -adic topology; set  $S = \text{Spec}(A)$  and  $S_0 = \text{Spec}(A/I)$ . Let  $H$  be an affine  $S$ -group scheme of finite type, flat over  $S$  at the points of  $H_0$ , such that  $H_0 = H \times_S S_0$  is of multiplicative type and isotrivial. Then there exists an open and closed subgroup  $G$  of  $H$ , of isotrivial multiplicative type (and of finite type), such that  $G_0 = H_0$ .*

By 3.2 (i), there exist an isotrivial group  $G$  of multiplicative type over  $S$  and an isomorphism

$$u_0 : G_0 \xrightarrow{\sim} H_0.$$

By 3.1,  $u_0$  comes from a unique homomorphism of  $S$ -groups

$$u : G \longrightarrow H.$$

Using IX 6.6, we see that  $u$  is a monomorphism: if  $U$  is the set of  $s \in S$  such that  $u_s : G_s \rightarrow H_s$  is a monomorphism, then  $U$  is an open neighborhood of  $S_0$ , hence equals  $S$ , and  $G|_U \rightarrow H|_U$  is a monomorphism. By IX 2.5,  $u$  is even a closed immersion.

Thus  $G$  is of finite type, and therefore of finite presentation, over  $S$ .<sup>16</sup> Applying Lemma 3.5 in the étale case, there is an open neighborhood  $U$  of  $S_0$ , hence  $U = S$ , such that  $G|_U \rightarrow H|_U$  is étale. Thus  $u$  is étale and therefore an open immersion, since it is an étale monomorphism.<sup>17</sup> This completes the proof.

**Corollary 3.8.** — *Let  $A$  be a noetherian ring equipped with an ideal  $I$  such that  $A$  is separated and complete for the  $I$ -adic topology; set  $S = \text{Spec}(A)$  and  $S_0 = \text{Spec}(A/I)$ . Let  $H$  be an  $S$ -group prescheme of finite type, affine and flat over  $S$ .*

*Then  $H$  is of multiplicative type and isotrivial if and only if  $H_0$  is so and  $H$  satisfies either one of the following conditions (a) and (b), which are therefore equivalent under the preceding hypotheses:*

- (a) *The fibers of  $H$  are of multiplicative type and have constant type on each connected component of  $S$ .*
- (b)  *$H$  is commutative and the  ${}_n H$ ,  $n > 0$ , are finite over  $S$ .*

*Both of these conditions are implied by the following one:*

<sup>16</sup>Editor's note: The preceding sentence has been added.

<sup>17</sup>Editor's note: Cf. EGA IV<sub>4</sub>, 17.9.1.

(c) *The fibers of  $H$  are connected.*

Of course, if  $H$  is of multiplicative type (and isotrivial), then (a) and (b) hold by IX 1.4.1 a) and 2.1 c). Thus only the “if” direction requires proof. We use the subgroup  $G$  provided by 3.7. When (c) holds, plainly  $G = H$ , and there is nothing to prove.

In case (b), the immersion  $u : G \rightarrow H$  induces an open immersion

$${}_nu : {}_nG \longrightarrow {}_nH$$

and an isomorphism

$$({}_nG)_0 \xrightarrow{\sim} ({}_nH)_0.$$

Since  ${}_nH$  is finite over  $S$ , it follows at once that  ${}_nu$  is an isomorphism: the complement of its image is finite over  $S$  and reduces to the empty set. By 3.6, the induced morphisms on fibers

$$u_s : G_s \longrightarrow H_s$$

are isomorphisms. Hence  $u$  is surjective and therefore an isomorphism.

Finally, in case (a), we may suppose that  $S$  is connected.<sup>18</sup> It follows that, for every  $s \in S$ ,

$$u_s : G_s \longrightarrow H_s$$

is a monomorphism of algebraic groups of multiplicative type and of the same type over  $\kappa(s)$ .<sup>19</sup> Such a homomorphism is necessarily an isomorphism, which again completes the proof.<sup>20</sup> After extending the base field, we may suppose that both groups over  $\kappa(s)$  are diagonalizable. The assertion then follows from VIII 3.2 b) and the fact that a surjective homomorphism between isomorphic finitely generated  $\mathbb{Z}$ -modules

$$M \longrightarrow N$$

is necessarily bijective.

**Corollary 3.9.** — *Let  $A$  be a noetherian ring equipped with an ideal  $I$  such that  $A$  is separated and complete for the  $I$ -adic topology, set  $S = \operatorname{Spec}(A)$ , and let  $H$  be an  $S$ -group prescheme of finite type, affine and flat over  $S$ , with connected fibers. Then  $H$  is an isotrivial torus if and only if  $H_0 = H \times_S \operatorname{Spec}(A/I)$  is an isotrivial torus.*

## 4. Case of an arbitrary base. Quasi-isotriviality theorem

Let  $A$  be a local ring. Recall, following Nagata, that  $A$  is said to be *henselian* if every algebra  $B$  finite over  $A$  is a product of local algebras  $B_i$  finite over  $A$ .

**Recall 4.0.**<sup>21</sup> *Let  $A$  be a henselian local ring,  $k$  its residue field,  $S = \operatorname{Spec}(A)$ ,  $S_0 = \operatorname{Spec}(k)$ , and  $\xi$  a geometric point of  $S_0$ . Then the functor*

$$X \longmapsto X_0 = X \times_S S_0$$

*is an equivalence between the category of étale coverings of  $S$  and the analogous category over  $S_0$  (cf. EGA IV<sub>4</sub>, § 18.5). Consequently (cf. SGA 1, V),*

$$\pi_1(S_0, \xi) = \pi_1(S, \xi).$$

<sup>18</sup>Editor’s note: The original said “that is,  $S_0$  connected.” Since  $A$  is separated for the  $I$ -adic topology,  $S_0$  meets every connected component of  $S$ ; and since  $A$  is complete, the connected components of  $S_0$  and  $S$  are in bijection.

<sup>19</sup>Editor’s note: This is because  $G_s$  and  $H_s$  have the same type for every  $s \in S_0$ , and hence for every  $s$ .

<sup>20</sup>Editor’s note: Indeed,  $u$  will again be a surjective open immersion.

<sup>21</sup>Editors’ note. This recall has been spelled out, and Remark 4.0.1 has been added.

Suppose in addition that  $A$  is noetherian, and write  $A'$  for its completion and  $S' = \operatorname{Spec}(A')$ . Then  $A'$  is a complete noetherian local ring, hence henselian (*loc. cit.*, 18.5.14), and the functor

$$X \mapsto X' = X \times_S S'$$

from the category of étale coverings of  $S$  to the analogous category over  $S'$  fits into a commutative diagram

$$\begin{array}{ccc} \operatorname{Rev.ét.}(S) & \xrightarrow{\quad} & \operatorname{Rev.ét.}(S') \\ & \searrow \simeq & \swarrow \simeq \\ & \operatorname{Rev.ét.}(S_0) & \end{array}$$

and is therefore also an equivalence of categories, whence  $\pi_1(S_0, \xi) = \pi_1(S', \xi)$ .

**Remark 4.0.1.** Since  $S$  is connected ( $A$  being local), it follows from 1.2 that the category of isotrivial groups of multiplicative type over  $S$  is equivalent to the analogous category over  $S_0$  (and also over  $S'$  if  $A$  is noetherian).

**Lemma 4.1.** *Let  $A$  be a henselian local ring with residue field  $k$ ,  $S = \operatorname{Spec}(A)$ , and  $S_0 = \operatorname{Spec}(k)$ .*

(i) *The functor*

$$H \mapsto H_0 = H \times_S S_0$$

*is an equivalence between the category of finite groups of multiplicative type over  $S$  and the analogous category over  $S_0$ .*

(ii) *If, in addition,  $A$  is noetherian, writing  $A'$  for its completion and  $S' = \operatorname{Spec}(A')$ , the functor*

$$H \mapsto H' = H \times_S S'$$

*is an equivalence between the category of finite groups of multiplicative type over  $S$  and the analogous category over  $S'$ .*

As in 4.0, the second assertion follows from the first; let us prove the first. We already know that the functor under consideration is essentially surjective: every group of multiplicative type  $H_0$  over  $S_0 = \operatorname{Spec}(k)$ , being finite and hence of finite type over  $k$ , is isotrivial by 1.4, and therefore comes from an isotrivial group of multiplicative type over  $S$ , by 4.0.1.

It remains to prove full faithfulness, that is, that for two finite groups  $G, H$  of multiplicative type over  $S$ , the following map is bijective:

$$\operatorname{Hom}_{S\text{-gr.}}(G, H) \longrightarrow \operatorname{Hom}_{S_0\text{-gr.}}(G_0, H_0).$$

In other words,<sup>22</sup> if  $F$  denotes the  $S$ -functor  $\operatorname{Hom}_{S\text{-gr.}}(G, H)$ , the natural map

$$\operatorname{Hom}_S(S, F) \longrightarrow \operatorname{Hom}_{S_0}(S_0, F_0)$$

induced by the base change  $S_0 \rightarrow S$  is bijective. In view of Recall 4.0, it is enough to prove the following lemma.

**Lemma 4.2.** *Let  $G, H$  be two finite groups of multiplicative type over a prescheme  $S$ . Then  $F = \operatorname{Hom}_{S\text{-gr.}}(G, H)$  is representable by a finite étale scheme over  $S$ .*

<sup>23</sup> Let  $f : S' \rightarrow S$  be a faithfully flat quasi-compact morphism such that  $G'$  and  $H'$  are diagonalizable. It is enough to show that  $F_{S'}$  is represented by a prescheme  $X'$  étale and finite (hence affine) over  $S'$ . Indeed, the descent datum on  $X'$  relative to  $f$  (cf. VIII 1.7.2) is

<sup>22</sup> *Editors' note.* What follows has been added.

<sup>23</sup> *Editors' note.* The original has been expanded in what follows.

then effective (by SGA 1, VIII 2.1), so there is an  $S$ -prescheme  $X$  such that  $X \times_S S' = X'$ , representing  $F$ , and  $X$  is étale and finite over  $S$  (cf. SGA 1, V 5.7 and EGA IV<sub>4</sub>, 17.7.3 (ii)).

We may therefore suppose that  $G = D_S(M)$  and  $H = D_S(N)$ , where  $M$  and  $N$  are finite commutative groups (cf. VIII 2.1 c)). Then  $K = \text{Hom}_{\text{gr.}}(N, M)$  is a finite abelian group and, by VIII 1.5, there is an isomorphism

$$\text{Hom}_{S\text{-gr.}}(G, H) \simeq K_S,$$

which completes the proof of 4.2 and hence of 4.1.

**Proposition 4.3.0.**<sup>24</sup> *Let  $S$  be a henselian local scheme,  $s$  its closed point,  $g : X \rightarrow S$  a morphism locally of finite type, and  $x$  an isolated point of the fiber  $X_s$ .*

(i) *Then  $\mathcal{O}_{X,x}$  is finite over  $\mathcal{O}_{S,s}$ . (In particular, if the residue-field extension  $\kappa(x)/\kappa(s)$  is trivial, then  $\mathcal{O}_{S,s} \rightarrow \mathcal{O}_{X,x}$  is surjective, by Nakayama's lemma.)*

(ii) *If  $g$  is also separated, then  $X' = \text{Spec}(\mathcal{O}_{X,x})$  is an open and closed subscheme of  $X$ ; that is, there is a decomposition into a disjoint sum  $X = X' \amalg X''$ .*

*Proof.* By the local form of Zariski's Main Theorem found in [Pes66], or in [Ray70, Ch. IV, Th. 1],  $x$  has an affine open neighborhood  $U = \text{Spec}(B)$ , of finite type and quasi-finite over  $A = \mathcal{O}_{S,s}$ , and there is an open immersion  $U \hookrightarrow Y = \text{Spec}(C)$ , where  $C$  is a finite  $A$ -algebra. (N.B. To obtain this conclusion one may also use Chevalley's semicontinuity theorem (EGA IV<sub>3</sub>, 13.1.4), followed by the form of Zariski's Main Theorem given in *loc. cit.*, 8.12.8.)

Since  $A$  is henselian,  $Y$  is the disjoint sum of local schemes  $Y_1, \dots, Y_n$ , each finite over  $S$ , and the points of  $Y$  above  $s$  are their closed points  $y_1, \dots, y_n$ . Thus  $x = y_i$  for some  $i$ , and  $\mathcal{O}_{X,x} = \mathcal{O}_{U,x} = C_i$  is finite over  $A$ . Moreover,  $X' = \text{Spec}(C_i)$  is an open subscheme of  $U$ , hence of  $X$ .

Suppose in addition that  $g$  is separated. Since  $X' \rightarrow S$  is finite ( $C_i$  being finite over  $A$ ), so is the immersion  $X' \rightarrow X$  (cf. EGA II, 6.1.5 (v)); therefore  $X'$  is also closed in  $X$ .  $\square$

**Lemma 4.3.** *Let  $A$  be a noetherian henselian local ring,  $A'$  its completion,  $S = \text{Spec}(A)$ ,  $S' = \text{Spec}(A')$ , and  $s$  the closed point of  $S$ . Let  $G, H$  be two  $S$ -preschemes in groups, with  $G$  of multiplicative type and of finite type over  $S$ , and with  $H$  locally of finite type and separated over  $S$ ,  $H_s$  of multiplicative type, and  $H$  flat over  $S$  at the points of  $H_s$ .*

*Let  $G', H'$  be obtained from  $G, H$  by the base change  $S' \rightarrow S$ . Then the following natural map is bijective:*

$$\text{Hom}_{S\text{-gr.}}(G, H) \xrightarrow{\sim} \text{Hom}_{S'\text{-gr.}}(G', H').$$

Since  $S'$  is faithfully flat and quasi-compact over  $S$ , SGA 1, VIII 5.2 shows that this map identifies the first member with the subset of the second formed by the  $u' : G' \rightarrow H'$  for which the two inverse images  $u'_1, u'_2 : G'' \rightarrow H''$  of  $u'$  on  $S'' = S' \times_S S'$ , under the projections  $\text{pr}_1, \text{pr}_2 : S'' \rightarrow S'$ , are equal.

It therefore remains to prove that  $u'_1 = u'_2$  for every homomorphism of  $S'$ -groups  $u' : G' \rightarrow H'$ . By the density theorem IX 4.8, it is enough to prove that  $u'_1$  and  $u'_2$  agree on  ${}_n G''$  for every integer  $n > 0$ . (N.B. Here the density theorem is needed essentially in a case where the base prescheme, namely  $S''$ , is not locally noetherian.)

Replacing  $G$  by  ${}_n G$ , this reduces us to the case in which  $n \cdot \text{id}_G = 0$  for some integer  $n > 0$ , and hence  $G$  is finite over  $S$ . Likewise let  ${}_n H$  be the kernel of the  $n$ -th power morphism  $\varphi_n$  on  $H$ . (N.B. We have not assumed  $H$  commutative, so  $\varphi_n$ , respectively  ${}_n H$ ,

<sup>24</sup> *Editors' note.* This recall, which is used in the proof of 4.3, has been added (the reference to EGA IV<sub>4</sub>, 18.5.11 was not entirely satisfactory).

need not be a group homomorphism, respectively a subgroup of  $H$ .) Since  ${}_nH$  is defined as the fiber product of  $\varphi_n : H \rightarrow H$  and the unit section  $\varepsilon : S \rightarrow H$ , its formation commutes with every base change  $T \rightarrow S$ :

$$({}_nH)_T = {}_n(H_T).$$

<sup>25</sup> We use a right subscript  $m$  for reduction modulo  $\mathfrak{m}^{m+1}$ , where  $\mathfrak{m}$  is the maximal ideal of  $A$ . Then  $H_m$  is flat over  $S_m$  (because  $H$  is flat over  $S$  at the points of  $H_0$ ); hence, by 2.4,  $H_m$  is of multiplicative type and of finite type, since  $H_0$  is. Thus every  $({}_nH)_m = {}_n(H_m)$  is a subgroup of  $H_m$ , of multiplicative type, finite and flat over  $S_m$ .

In particular,  $({}_nH)_0 = {}_n(H_0)$  is finite over  $S_0$ . Since  $S$  is henselian local and  ${}_nH$ , like  $H$ , is locally of finite type and separated over  $S$ , Proposition 4.3.0 shows that the local rings of  ${}_nH$  at the points of  $({}_nH)_0$  are finite over  $S$ , and that there is a decomposition as a disjoint sum of open subsets

$${}_nH = {}_nH^+ \amalg {}_nH^-, \quad (*)$$

where  $Z = {}_nH^+$  is finite over  $S$ , while  ${}_nH^-$  lies above  $S - \{s\}$ .

Observe that for every finite  $S$ -scheme  $Y$  (such as  $G$ ), every  $S$ -morphism  $Y \rightarrow {}_nH$  factors through  $Z$ , and that the formation of the decomposition  $(*)$  commutes with the base change  $S' \rightarrow S$ , where  $S' = \text{Spec}(A')$  and  $A'$  is the completion of  $A$ .

Then  $Z' = Z \times_S S'$  is finite over  $S'$ , as is  $P' = Z' \times_{S'} Z'$ . Let  $\nu$  denote the restriction to  $P'$  of the multiplication on  $H'$ , and let  $\sigma$  be the automorphism of  $P'$  interchanging the two factors. Since  $P'$  is finite over  $S'$ , while  $H'$  is separated and locally of finite type over  $S'$ , EGA II 5.4.3 and IV<sub>1</sub> 1.1.3 show that  $Y = \nu(P')$  is a closed subscheme of  $H'$ , universally closed and quasi-compact, hence of finite type and therefore proper over  $S'$ . Moreover,  $Y \rightarrow S'$  has finite fibers (because  $P' \rightarrow S'$  does). Since  $S'$  is noetherian,  $Y \rightarrow S'$  is therefore finite (cf. EGA III, 4.4.2). Since  $Z' \subseteq Y$  and  $Z'_m = Y_m$  for every  $m$ , Lemma IX 5.0 gives  $Z' = Y$ ; thus  $Z'$  is a subgroup of  $H'$ . Similarly, the kernel  $K = \text{Ker}(\nu, \nu \circ \sigma)$  is a closed subscheme of  $P'$  such that  $K_m = P'_m$  for every  $m$  (because  $Z'_m = {}_nH_m$  is commutative). Hence  $K = P'$ , that is,  $Z'$  is a commutative subgroup of  $H'$ .

On the other hand, every reduction  $Z'_m = {}_nH_m$  is flat over  $S_m$ . The “local criterion for flatness” (cf. EGA 0<sub>III</sub>, 10.2.2, or [BAC], III § 5, Example 1 and Th. 1) therefore shows that  $Z'$  is flat over  $S'$ . Moreover,  $Z'_0 = {}_nH_0$  is a finite group of multiplicative type over  $S_0$ , and hence isotrivial by 1.4. It follows from 3.8 b) that  $Z'$  is of multiplicative type (and isotrivial) over  $S'$ . Since  $S' \rightarrow S$  is faithfully flat and quasi-compact, the multiplication  $Z \times_S Z \rightarrow H$  factors through  $Z$  and makes  $Z$  a subgroup of  $H$ , finite over  $S$  and of multiplicative type (because  $Z'$  is).

Finally, by the observations made above concerning  $(*)$ ,  $Z'$  is the “finite part” of  ${}_nH'$ , and therefore  $u' : G' \rightarrow {}_nH'$  takes its values in  $Z'$ . Since  $Z$ , as well as  $G = {}_nG$ , is finite over  $S$  and of multiplicative type, Lemma 4.1 shows that  $u'$  comes from a  $u : G \rightarrow Z$ , and hence  $u'_1 = u'_2$ . This completes the proof of 4.3.

**Lemma 4.4.0.**<sup>26</sup> *Let  $A$  be a ring,  $S = \text{Spec}(A)$ , and let  $H$  be an  $S$ -group scheme of multiplicative type and of finite type, quasi-isotrivial (respectively isotrivial). Let  $(A_i)$  be a filtered family of subrings of  $A$  whose inductive limit is  $A$ , and put  $S_i = \text{Spec}(A_i)$ .*

*Then there are an index  $i$  and an  $S_i$ -group scheme  $H_i$ , of multiplicative type and of finite type, quasi-isotrivial (respectively isotrivial), such that  $H = H_i \times_{S_i} S$ .*

**Theorem 4.4.** *Let  $S$  be a prescheme, let  $H$  be an affine  $S$ -prescheme in groups of finite presentation over  $S$ , and let  $s \in S$ . Suppose:*

*α)  $H$  is flat over  $S$  at the points of  $H_s$ ;*

<sup>25</sup> Editors' note. The original has been expanded in what follows, on the basis of indications by M. Raynaud.

<sup>26</sup> Editors' note. This lemma, used several times below, has been added.

$\beta$ )  $H_s$  is of multiplicative type.

Then there exist an étale morphism  $S' \rightarrow S$ , a point  $s' \in S'$  above  $s$  such that  $\kappa(s')/\kappa(s)$  is trivial, and an open and closed subgroup  $G'$  of  $H' = H \times_S S'$ , of multiplicative type and finite type, and isotrivial, such that  $G'_{s'} = H'_{s'}$ .

<sup>27</sup> **a)** Write provisionally  $T = \operatorname{Spec}(\mathcal{O}_{S,s})$ , and let us show, by “descent of the conclusions”, that one may reduce to the case  $S = T$ . Suppose indeed that we have found an étale morphism  $g : T' \rightarrow T$ , a point  $s' \in T'$ , and a subgroup  $G'$  of  $H' = H_{T'}$  satisfying the conditions of the statement. Replacing  $T'$  by an affine open neighborhood of  $s'$ , we may assume that  $T'$  is of finite presentation over  $T$ .

Since  $G'$  is isotrivial, there is a finite étale surjective morphism  $f : \tilde{T} \rightarrow T'$  such that  $\tilde{G} = G' \times_{T'} \tilde{T}$  is isomorphic to  $D_{\tilde{T}}(M)$ , where  $M$  is a finitely generated abelian group. Since  $T'$ ,  $H$ ,  $\tilde{T}$ , and  $G'$  are of finite presentation over  $T = \operatorname{Spec}(\mathcal{O}_{S,s})$ , and  $\mathcal{O}_{S,s}$  is the inductive limit of the subalgebras  $A_i = \mathcal{O}_S(S_i)$ , where  $S_i$  runs through the affine open neighborhoods of  $s$  in  $S$ , EGA IV<sub>3</sub>, 8.8.2 (and Exposé VI<sub>B</sub>, 10.2 and 10.3) gives an index  $i$ ,  $S_i$ -preschemes  $S'_i, \tilde{T}_i$  of finite presentation (respectively an  $S_i$ -prescheme in groups  $G'_i$  of finite presentation), and morphisms

$$g_i : S'_i \rightarrow S_i, \quad f_i : \tilde{T}_i \rightarrow S'_i, \quad u_i : G'_i \rightarrow H \times_S S'_i,$$

from which  $g, f$ , respectively  $u : G' \rightarrow H'$ , are obtained by base change. Taking  $i$  large enough,  $g_i$  is étale,  $f_i$  is finite étale surjective, and  $u_i$  is an open and closed immersion (cf. EGA IV, 8.10.5 and 17.7.8).

Now  $\tilde{G}$  comes from both  $\tilde{G}_i = G'_i \times_{S'_i} \tilde{T}_i$  and  $D_{\tilde{T}_i}(M)$ . By EGA IV<sub>3</sub>, 8.8.2 (i) (and VI<sub>B</sub>, 10.2), there is an index  $j \geq i$  such that  $\tilde{G}_j \simeq D_{\tilde{T}_j}(M)$ , so  $G'_j$  is an isotrivial group of multiplicative type. Let  $s'_j$  be the image of  $s'$  in  $S'_j$ . Then the étale morphism  $S'_j \rightarrow S_j \hookrightarrow S$ , the point  $s'_j$ , and the open and closed subgroup  $G'_j$  of  $H \times_S S'_j$  satisfy the conditions of the statement. Thus we may assume that  $S = \operatorname{Spec}(A)$  is local, with closed point  $s$ .

**b)** The ring  $A$  is the inductive limit of local rings  $A_i$  which are localizations of finitely generated  $\mathbb{Z}$ -algebras; put  $S_i = \operatorname{Spec}(A_i)$ . Let us show that hypotheses  $\alpha$  and  $\beta$  “descend” to some  $A_i$ . Since  $H \rightarrow S$  is of finite presentation, the set of  $x \in H$  at which  $H$  is flat over  $S$  is an open set  $W$ . It contains  $H_s$  by hypothesis, and hence contains a quasi-compact open set  $V$  containing  $H_s$  (since the affine scheme  $H_s$  can be covered by finitely many affine open subsets contained in  $W$ ). The open immersion  $\tau : V \hookrightarrow H$  is of finite presentation, so  $V \rightarrow S$  is also of finite presentation.

By EGA IV<sub>3</sub>, 8.8.2 (and Exposé VI<sub>B</sub>, 10.2 and 10.3), there are an index  $i$ , an  $S_i$ -prescheme  $V_i$  of finite presentation (respectively an  $S_i$ -prescheme in groups  $H_i$  of finite presentation), and an  $S_i$ -morphism  $\tau_i : V_i \rightarrow H_i$ , from which  $V, H, \tau$  are obtained by base change  $S \rightarrow S_i$ . Taking  $i$  large enough,  $\tau_i$  is an open immersion and  $V_i$  is flat over  $S_i$ , by EGA IV<sub>3</sub>, 8.10.5 and 11.2.6. Let  $s_i$  be the image of  $s$  in  $S_i$ . Since the open immersion  $(V_i)_{s_i} \hookrightarrow (H_i)_{s_i}$ , after base change  $\kappa(s_i) \rightarrow \kappa(s)$ , gives the equality  $V_s = H_s$ , one already has  $(V_i)_{s_i} = (H_i)_{s_i}$ . Thus  $H_i$  is flat over  $S_i$  at the points of  $(H_i)_{s_i}$ . Finally,  $(H_i)_{s_i}$  is of multiplicative type, because

$$H_s = (H_i)_{s_i} \otimes_{\kappa(s_i)} \kappa(s)$$

is. Hence  $(S_i, H_i, s_i)$  satisfies the hypotheses of 4.4, and if the desired assertion holds for this triple, it also holds for  $(S, H, s)$  by base change. We are reduced to the case in which  $A$  is local and noetherian. We now distinguish two cases.

<sup>27</sup> *Editors' note.* The reductions that follow have been spelled out (the original said: “One reduces at once to the case where  $S$  is the spectrum of a noetherian local ring  $A$ , and where  $s$  is the closed point of  $S$ .”)

1°  $A$  is local noetherian and henselian. Let  $\widehat{A}$  be its completion,  $\widehat{S} = \operatorname{Spec}(\widehat{A})$ , and  $\widehat{H} = H \times_S \widehat{S}$ . Applying Theorem 3.7, we find an isotrivial  $\widehat{S}$ -group  $\widehat{G}$  of multiplicative type and of finite type, together with a homomorphism

$$\widehat{u} : \widehat{G} \longrightarrow \widehat{H}$$

which is both an open and a closed immersion, and which induces an isomorphism

$$\widehat{u}_0 : \widehat{G}_0 \xrightarrow{\sim} \widehat{H}_0.$$

By Remark 4.0.1, base change along  $\widehat{S} \rightarrow S$  induces an equivalence between the categories of isotrivial groups of multiplicative type over  $S$  and over  $\widehat{S}$ . In particular,  $\widehat{G}$  comes from an isotrivial  $S$ -group  $G$  of multiplicative type and of finite type. By 4.3,  $\widehat{u}$  comes from a homomorphism

$$u : G \longrightarrow H.$$

Moreover,  $u$  is an open and closed immersion and induces an isomorphism  $u_0 : G_0 \rightarrow H_0$ , since this is true after the faithfully flat quasi-compact base change  $\widehat{S} \rightarrow S$ . This proves 4.4 in this case (taking  $S' = S$  and  $s' = s$  in its conclusion).

2°  $A$  is local noetherian. The reduction to case 1° is immediate, by applying that case to the henselization  $A^h$  of  $A$ . More precisely, one sees easily (using SGA 1, I § 5 \* ) that the local rings  $\mathcal{O}_{S',s'}$  of étale  $S$ -preschemes  $S'$ , equipped with a point  $s'$  above  $s$  for which  $\kappa(s')/\kappa(s)$  is trivial, form a filtered inductive system whose inductive limit is a noetherian henselian local ring  $A^h$  (called the henselization of  $A$ ); for details of this construction, due to Nagata in the normal case, see SGA 4, VIII § 4. The standard arguments of EGA IV<sub>3</sub>, § 8 then allow one, as in part a), to deduce from the known result over the inductive limit  $A^h$  of the  $\mathcal{O}_{S',s'}$  an analogous result over one of the pairs  $(S', s')$ . This proves 4.4.

**Corollary 4.5.** *Let  $S$  be a prescheme and  $H$  an  $S$ -prescheme in groups of multiplicative type and of finite type. Then  $H$  is quasi-isotrivial; that is,  $H$  is split by an étale surjective morphism  $S' \rightarrow S$ .*

Indeed, let  $s \in S$ . By 4.4, there are an étale morphism  $S' \rightarrow S$ , a point  $s' \in S'$  above  $s$  with  $\kappa(s) = \kappa(s')$ , and an isotrivial subgroup  $G'$  of  $H'$ , of multiplicative type and of finite type, such that  $G'_{s'} = H'_{s'}$ . Since  $G'$  and  $H'$  are of multiplicative type and finite type, IX 2.9 gives an open neighborhood  $U'$  of  $s'$  such that  $G'|_{U'} = H'|_{U'}$ .

<sup>28</sup> Suppose in addition that  $S$  is henselian local, with closed point  $s$ . By EGA IV<sub>4</sub>, 18.5.11 b), there is a section  $\sigma$  of  $S' \rightarrow S$  such that  $\sigma(s) = s'$ . (N.B. One can see directly that  $B = \mathcal{O}_{S',s'}$  equals  $A = \mathcal{O}_{S,s}$  as follows. By 4.3.0 (i),  $B \simeq A/I$ . Since  $B$  is a flat  $A$ -algebra of finite presentation,  $I$  is a finitely generated ideal (cf. EGA IV<sub>1</sub>, 1.4.7), and  $I = I\mathfrak{m}$ , where  $\mathfrak{m}$  is the maximal ideal of  $A$ ; hence  $I = 0$ .) Thus  $H$  is already isotrivial. We obtain:

**Corollary 4.6.** *Let  $A$  be a henselian local ring,  $k$  its residue field, and  $\pi$  the topological Galois group of an algebraic closure of  $k$ .*

- (i) *Every group of multiplicative type and of finite type over  $S = \operatorname{Spec}(A)$  is isotrivial.*
- (ii) *The category of these groups over  $S$  is equivalent, through the functor  $H \mapsto H \otimes_A k$ , to the analogous category over  $S_0 = \operatorname{Spec}(k)$ . It is therefore anti-equivalent, by 1.4, to the category of Galois modules under  $\pi$  which are finitely generated as  $\mathbb{Z}$ -modules.*

**Corollary 4.7.** *Under the hypotheses of 4.4, suppose in addition that one of the following conditions holds:*

---

\*Or EGA IV<sub>4</sub>, § 18.6.

<sup>28</sup>Editors' note. The original has been expanded in what precedes and what follows.



- a) for every generization  $t$  of  $s$ ,  $H_t$  is of multiplicative type and has the same type as  $H_s$ ;
- b)  $H$  is commutative and the  ${}_nH$ ,  $n > 0$ , are finite over  $S$  in a neighborhood of  $s$ ;
- c) for every generization  $t$  of  $s$ , the fiber  $H_t$  is connected.

Then there is an open neighborhood  $U$  of  $s$  such that  $H|_U$  is of multiplicative type.

Since an étale morphism is open, it is enough to show, with the notation of the conclusion of 4.4, that there is an open neighborhood  $U'$  of  $s'$  such that  $G'|_{U'} = H'|_{U'}$ . Put  $S'' = \text{Spec}(\mathcal{O}_{S',s'})$ . Since  $G'$  and  $H'$  are of finite presentation over  $S'$ , EGA IV<sub>3</sub>, 8.8.2 reduces the matter to showing that  $G'' = H''$ . We may therefore assume that  $S = S''$ ; then conditions a), b), c) become those of the following lemma, whose proof is the same as that of 3.8.

**Lemma 4.7.1.** *Let  $S$  be a local scheme with closed point  $s$ , let  $H$  be an  $S$ -group scheme of finite type, and let  $G$  be an open and closed subgroup of  $H$ , of multiplicative type, such that  $G_s = H_s$ . Suppose in addition that one of the following conditions holds:*

- a) the fibers of  $H$  are of multiplicative type and have the same type as  $H_s$ ;
- b)  $H$  is commutative and the  ${}_nH$ ,  $n > 0$ , are finite over  $S$ ;
- c) the fibers of  $H$  are connected.

Then  $G = H$ .

**Corollary 4.8.** *Let  $S$  be a prescheme and let  $H$  be an affine  $S$ -prescheme in groups, flat and of finite presentation over  $S$ , with fibers of multiplicative type.*

*For  $H$  to be of multiplicative type, it is necessary and sufficient that it satisfy one of the following two conditions (which are equivalent under the preceding hypotheses):*

- a) the type of  $H_s$  (cf. IX 1.4) is a locally constant function of  $s \in S$ ;
- b)  $H$  is commutative and the  ${}_nH$ ,  $n > 0$ , are finite over  $S$ .

Moreover, a) and b) are implied by:

- c) the fibers of  $H$  are connected.

In particular:

**Corollary 4.9.** *Let  $S$  be a prescheme and  $H$  an  $S$ -prescheme in groups, flat and of finite presentation over  $S$ . Suppose in addition that  $H$  is affine over  $S$  and has connected fibers.<sup>29</sup> If  $s \in S$  is such that  $H_s$  is a torus, there is an open neighborhood  $U$  of  $s$  such that  $H|_U$  is a torus.*

*In particular, if all the fibers of  $H$  are tori,  $H$  is a torus.*

## 5. Scheme of homomorphisms from one group of multiplicative type to another. Twisted constant groups and groups of multiplicative type

**Definition 5.1.** a) *Let  $S$  be a prescheme and  $R$  a prescheme in groups over  $S$ . We say that  $R$  is a twisted constant group over  $S$  if it is locally isomorphic, for the faithfully flat quasi-compact topology, to a constant group scheme, that is, one of the form  $M_S$  for an ordinary group  $M$ .*

b) *A twisted constant group  $R$  over  $S$  is said to be quasi-isotrivial, respectively isotrivial, respectively locally isotrivial, respectively locally trivial, respectively trivial, if in the preceding definition one can replace the faithfully flat quasi-compact topology by the étale topology,*

<sup>29</sup>Editors' note. This result is generalized in 8.1: it is enough to suppose that the fibers of  $H$  are affine and connected.

respectively the global finite étale topology, respectively the finite étale topology, respectively the Zariski topology, respectively the coarsest (or “chaotic”) topology; cf. IX 1.1 and 1.2, and IV 6.6.

Thus, to say that  $R$  is quasi-isotrivial (respectively isotrivial) means that there is an étale surjective (respectively also finite) morphism  $S' \rightarrow S$  such that  $R' = R \times_S S'$  is a constant group over  $S'$ ; to say that  $R$  is trivial means that  $R$  is a constant group.

c) As in VIII 1.4, one defines the type of a twisted constant group  $R$  over  $S$  at a point  $s \in S$ . It is an isomorphism class of ordinary groups and, as  $s$  varies, is a locally constant function of  $s$ , hence a constant function if  $S$  is connected. We also say that  $R$  is “of type  $M$ ” if all the fibers of  $R$  are of type  $M$ .

One should beware that  $R$  is quasi-compact over  $S$  only if it is finite over  $S$ , that is, only if its type at every  $s \in S$  is a finite group.<sup>30</sup>

d) The case of greatest interest to us is that in which  $R$  is commutative. We then say that  $R$  is finitely generated if its type at every point  $s \in S$  is given by a finitely generated  $\mathbb{Z}$ -module. This notion must not be confused with the scheme-theoretic notion “ $R$  of finite type over  $S$ ” (cf. above).

**Remark 5.2.** We shall also have to consider  $S$ -preschemes  $X$  which are locally isomorphic, for the faithfully flat quasi-compact topology, to constant preschemes, independently of any group structure. We then call  $X$  a *twisted constant bundle* over  $S$ , and extend to these preschemes the terminology introduced in 5.1. Of course, when  $X$  is endowed with an  $S$ -group structure, the meaning of the expressions “twisted constant”, “isotrivial”, and so on changes according as one does or does not take the group structure over  $S$  into account. The same applies when  $X$  carries any other kind of algebraic structure (for example, that of a Galois principal bundle, to be considered in the next section).

**Proposition 5.3.** *Let  $R$  be a commutative twisted constant group over  $S$ .*

- (i) *The functor  $H = D_S(R)$  (cf. VIII 1) is representable and is a group of multiplicative type over  $S$ .*
- (ii) *For every  $s \in S$ , the type of  $R$  at  $s$  equals the type of  $H$  at  $s$ .*
- (iii) *The group  $R$  is quasi-isotrivial (respectively isotrivial, respectively trivial, respectively locally isotrivial, respectively locally trivial) if and only if  $H$  is so.*

**Remark 5.3.1.**<sup>31</sup> One may be concerned that in 5.3 the word “type” is being used in two different senses, according as it refers to  $R$  or to  $H$ . Fortunately, when an  $S$ -prescheme in groups  $G$  is both a twisted constant group and a group of multiplicative type, its type in either sense is the same, thanks to the fact that an ordinary finite commutative group is isomorphic to its dual.

*Proof.* Covering families for the faithfully flat quasi-compact topology are families of effective descent for the fibered category of affine preschemes in groups over variable base preschemes (SGA 1, VIII 2.1). Consequently  $H$  is representable (and affine over  $S$ ), since this is true “locally”<sup>32</sup> (when  $R$  is constant,  $H$  is a diagonalizable group).

That  $H$  is of multiplicative type is then immediate from the definition, as is the equality of the types of  $R$  and  $H$  at  $s \in S$ . Finally, since

$$H_{S'} = D_{S'}(R_{S'}),$$

the last assertion reduces to the “trivial” case, namely to checking that  $R$  is trivial if and only if  $H$  is trivial; this follows at once from the biduality theorem VIII 1.2.  $\square$

<sup>30</sup> *Editors’ note.* See Lemma 5.12 below.

<sup>31</sup> *Editors’ note.* This remark, which appeared after the proof in the original, has been moved here.

<sup>32</sup> *Editors’ note.* Cf. VIII § 1.7.

To complete the description of the correspondence between twisted constant groups and groups of multiplicative type, one must start with a group of multiplicative type  $H$  and study  $R = D_S(H)$ . If  $R$  is representable, it is plainly a twisted constant group and  $H \simeq D_S(R)$ . In other words:

**Scholium 5.4.0.** *The functor  $R \mapsto D_S(R)$  is an anti-equivalence<sup>33</sup> between the category of twisted constant groups over  $S$  and the category of groups of multiplicative type  $H$  over  $S$  for which  $D_S(H)$  is representable.*

I do not know whether this condition on  $H$  is always satisfied. We shall see, however, that it is satisfied when  $H$  is quasi-isotrivial, and in particular when  $H$  is of finite type.

**Lemma 5.4.** *Let  $S' \rightarrow S$  be a faithfully flat morphism locally of finite presentation, and let  $X'$  be an  $S'$ -prescheme separated, locally of finite presentation, and locally quasi-finite over  $S'$ .*

*Then every descent datum on  $X'$  relative to  $S' \rightarrow S$  is effective: there are an  $S$ -prescheme  $X$  and an  $S'$ -isomorphism  $X \times_S S' \simeq X'$  compatible with the descent datum.*

We have already observed that the hypothesis on  $S' \rightarrow S$  implies that it is a covering morphism for the faithfully flat quasi-compact topology (IV 6.3.1 (iv)), and hence *a fortiori* a morphism of effective descent.

When  $X' \rightarrow S'$  is quasi-compact, and hence of finite presentation and quasi-finite, it is a quasi-affine morphism (cf. SGA 1, VIII 6.2<sup>34</sup>), and effectivity follows in this case from SGA 1, VIII 7.9. In the general case, one reduces at once to the case where  $S$  and  $S'$  are affine. Cover  $X'$  by affine open subsets  $U'_i$ , and let  $V'_i$  be the saturation of  $U'_i$  under the equivalence relation on  $X'$  defined by the descent datum:

$$V'_i = q_2(q_1^{-1}(U'_i)).$$

Here  $q_1, q_2$  are the two projections to  $X'$  of

$$X''_1 = X' \times_{S'} (S'', p_1) \quad \text{and} \quad X''_2 = X' \times_{S'} (S'', p_2),$$

respectively:  $q_1 = \text{pr}_1$ , and  $q_2$  is obtained from the first projection of  $X''_2$  by means of the given descent isomorphism  $X''_1 \simeq X''_2$ . Since  $S' \rightarrow S$  is faithfully flat, quasi-compact, and locally of finite presentation, the same is true of

$$p_1 : S'' = S' \times_S S' \longrightarrow S',$$

and hence of  $q_1, q_2$ . These are consequently open morphisms (SGA 1, IV 6.6<sup>35</sup>). Therefore  $V'_i$  is an open quasi-compact part of  $X'$ . By the case already proved, the induced descent data on the  $V'_i$  are effective; hence so is the descent datum on  $X'$  (SGA 1, VIII 7.2).

**Corollary 5.5.** *A faithfully flat morphism of finite presentation  $S' \rightarrow S$  is a morphism of effective descent for the fibered category of twisted constant groups over variable base preschemes.*

Indeed, this amounts to the effectivity assertion of 5.4 when  $X'$  is a constant  $S'$ -prescheme.

**Theorem 5.6.** *Let  $S$  be a prescheme, and let  $G, H$  be two quasi-isotrivial  $S$ -preschemes in groups of multiplicative type, with  $G$ <sup>36</sup> of finite type.*

<sup>33</sup>Editors' note. "Equivalence" has been corrected to "anti-equivalence".

<sup>34</sup>Editors' note. When  $S'$  is locally noetherian; EGA IV<sub>3</sub>, 8.11.2 in general.

<sup>35</sup>Editors' note. When  $S'$  is locally noetherian; EGA IV<sub>2</sub>, 2.4.6 in general.

<sup>36</sup>Editors' note.  $H$  has been corrected to  $G$ .

Then  $\mathrm{Hom}_{S\text{-gr.}}(H, G)$  is representable and is a quasi-isotrivial twisted constant group over  $S$ ,<sup>37</sup> for every  $s \in S$ , if the type of  $G$  at  $s$  is  $M$  and the type of  $H$  at  $s$  is  $N$ , then the type of  $\mathrm{Hom}_{S\text{-gr.}}(H, G)$  is  $\mathrm{Hom}_{\mathrm{gr.}}(M, N)$ .

The proof proceeds as in 4.2, using the fact that the assertion is known (VIII 1.5) when  $G$  and  $H$  are trivial. The required effectivity criterion is provided by 5.5, applied to an étale surjective morphism  $S' \rightarrow S$ .

Taking  $G = \mathbb{G}_{m,S}$ , in particular, gives:

**Corollary 5.7.** (i) Let  $H$  be a quasi-isotrivial  $S$ -group of multiplicative type. Then the  $S$ -group  $D_S(H)$  is representable and is a quasi-isotrivial twisted constant group over  $S$ .

(ii) The functors  $R \mapsto D_S(R)$  and  $H \mapsto D_S(H)$  are mutually quasi-inverse anti-equivalences between the category of quasi-isotrivial twisted constant  $S$ -groups and the category of quasi-isotrivial  $S$ -groups of multiplicative type.

(iii) These functors induce anti-equivalences between the subcategories of isotrivial, respectively locally isotrivial, respectively locally trivial, respectively trivial groups.

The last assertion is included only for reference, since it is already contained in 5.3.

Moreover, by 4.5 every  $S$ -group of multiplicative type and of finite type is quasi-isotrivial. Theorem 5.6 therefore gives:

**Corollary 5.8.** Let  $S$  be a prescheme, and let  $G, H$  be two  $S$ -preschemes in groups of multiplicative type and of finite type. Then  $\mathrm{Hom}_{S\text{-gr.}}(H, G)$  is representable and is a finitely generated quasi-isotrivial twisted constant  $S$ -group.

Also, in 5.3,  $R$  is finitely generated if and only if  $H = D_S(R)$  is of finite type (IX 2.1 b)). By 4.5,  $H$  is then quasi-isotrivial, and hence so is  $R$ . Thus:

**Corollary 5.9.** The functors of 5.7 induce mutually quasi-inverse anti-equivalences between the category of  $S$ -groups  $H$  of multiplicative type and of finite type and the category of finitely generated twisted constant  $S$ -groups  $R$ . Moreover, every such group  $R$  is quasi-isotrivial.

**Corollary 5.10.** Let  $H, G$  be two  $S$ -preschemes in groups of multiplicative type and of finite type.

(i)  $\mathrm{Isom}_{S\text{-gr.}}(H, G)$  is representable by an open and closed subprescheme of  $\mathrm{Hom}_{S\text{-gr.}}(H, G)$ , and is a twisted constant  $S$ -prescheme.

(ii) In particular,  $\mathrm{Aut}_{S\text{-gr.}}(H)$  is representable and is a twisted constant  $S$ -group, in general noncommutative.

This follows from 5.8 and VIII 1.6.<sup>38</sup>

**Recall 5.11.0.**<sup>39</sup> Recall that if  $X$  is a locally noetherian prescheme, its connected components (which are always closed) are open. Indeed, let  $C$  be a connected component of  $X$ , let  $x \in C$ , and let  $U$  be a noetherian open neighborhood of  $x$ . The prescheme  $U$  has only finitely many connected components, so the connected component  $U_x$  of  $x$  in  $U$  is open in  $U$ , hence in  $X$ . Since  $U_x \subseteq C$ , this shows that  $C$  is open in  $X$ .

**Proposition 5.11.** Let  $S$  be a prescheme, let  $R$  be a commutative twisted constant group over  $S$ , and let  $H = D_S(R)$  be the group of multiplicative type that it defines. Consider the following conditions:

(i)  $H$  is isotrivial (that is,  $R$  is isotrivial);

<sup>37</sup>Editors' note. What follows has been added.

<sup>38</sup>Editors' note. This reference should be corrected by treating the functor  $\mathrm{Isom}_{S\text{-gr.}}(H, G)$  in an addition 1.5.1 to VIII ...

<sup>39</sup>Editors' note. This recall, used several times below, has been added. See also EGA I, 6.1.9; note, however, that the proof in *loc. cit.* seems unnecessarily complicated.

(ii)  $R$  is a union of open and closed subpreschemes  $R_i$  which are quasi-compact over  $S$  (and then necessarily finite over  $S$ );

(iii) the connected components of  $R$  are finite over  $S$ .

a) Then (i)  $\Rightarrow$  (ii)  $\Rightarrow$  (iii).<sup>40</sup>

b) If  $S$  is locally noetherian, then (i)  $\Leftrightarrow$  (ii)  $\Leftrightarrow$  (iii).

c) Finally, if  $R$  is finitely generated (that is, if  $H$  is of finite type over  $S$ ), then (i)  $\Leftrightarrow$  (ii), at least when  $S$  is quasi-compact or when its connected components are open.

First decompose  $S$  as a prescheme sum of preschemes  $S_i$  on each of which  $H$  has constant type (cf. IX 1.4.1). We are reduced to the case where  $H$ , and therefore  $R$ , has constant type  $M$ . We shall use:

**Lemma 5.12.** *Let  $S$  be a prescheme and  $R$  a twisted constant prescheme over  $S$ . Every closed subprescheme  $Z$  of  $R$  which is quasi-compact over  $S$  is finite over  $S$ .*

Indeed, we reduce to the case where  $R$  is constant, hence of the form  $I_S$  for a set  $I$ . It is the filtered increasing union of the  $J_S$ , where  $J$  runs through the finite subsets of  $I$ . We may also assume that  $S$  is affine. Then  $Z$  is quasi-compact, and is therefore contained in some  $J_S$ . Since  $J_S$  is finite over  $S$ , so is  $Z$ .

Lemma 5.12 already proves the parenthetical assertion in 5.11 (ii).<sup>41</sup> The implication (ii)  $\Rightarrow$  (iii) is then clear, since the connected components of  $R$  are closed. If  $S$  is locally noetherian, they are open and closed by 5.11.0 ( $R$  being étale and hence locally of finite type over  $S$ ); this gives (iii)  $\Rightarrow$  (ii) in that case.

Let us prove (i)  $\Rightarrow$  (ii). Choose a finite étale surjective morphism  $S' \rightarrow S$  which splits  $H$ , and hence  $R$  (cf. 5.3), so that

$$R' \simeq M_{S'} = \coprod_{m \in M} R'_m,$$

where the  $R'_m$  are disjoint open subsets of  $R'$ , each  $S'$ -isomorphic to  $S'$ . Let  $g : R' \rightarrow R$  be the projection. It is a finite étale surjective morphism, and hence both open and closed. The subsets  $R_m = g(R'_m)$  are therefore open and closed parts of  $R$ , and are plainly quasi-compact over  $S$ , since the  $R'_m$  are.

Finally, suppose that  $H$  is of finite type over  $S$ , and let us prove (ii)  $\Rightarrow$  (i). If the connected components of  $S$  are open, the matter reduces at once to the case where  $S$  is connected; thus we may assume that  $S$  is quasi-compact or connected. Since  $M$  is finitely generated, write

$$M = \mathbb{Z}^r \times N,$$

where  $r \geq 0$  and  $N$  is a finite abelian group. Let  $G = D_S(M)$ , and consider the preschemes

$$P = \text{Isom}_{S\text{-gr.}}(H, G) \subset \text{Hom}_{S\text{-gr.}}(H, G) = Q$$

(cf. 5.8 and 5.10). There are isomorphisms

$$\begin{aligned} Q &\simeq \text{Hom}_{S\text{-gr.}}(M_S, R) \\ &\simeq \text{Hom}_{S\text{-gr.}}(\mathbb{Z}_S^r, R) \times \text{Hom}_{S\text{-gr.}}(N_S, R) \\ &\simeq R^r \times E, \end{aligned}$$

where

$$E = \text{Hom}_{S\text{-gr.}}(N_S, R)$$

<sup>40</sup> *Editors' note.* Compare the examples of 1.6 ...

<sup>41</sup> *Editors' note.* The original has been expanded in what follows.

is finite over  $S$ .<sup>42</sup> It follows that  $Q$  is a union of open and closed subpreschemes  $Q_i$  finite over  $S$ . Hence  $P$  is a union of the open and closed subpreschemes  $P_i = P \cap Q_i$ , finite over  $S$ . Since they are étale over  $S$ , their images in  $S$  are open and closed parts  $S_i$ , and these cover  $S$ . If  $S$  is connected or quasi-compact, there is a finite set of indices  $i$  for which the  $S_i$  cover  $S$ . Let  $S'$  be the union of the corresponding  $P_i$ . Then  $S' \rightarrow S$  is finite étale surjective. Putting

$$P' = P \times_S S' = \text{Isom}_{S'-\text{gr.}}(H', G'),$$

we see that  $P'$  has a section over  $S'$ ; hence there is an isomorphism of  $S'$ -groups

$$H' = H \times_S S' \xrightarrow{\sim} G' = G \times_S S' = D_{S'}(M).$$

Thus  $S'$  splits  $H$ , completing the proof of 5.11.<sup>43</sup>

**Remark 5.11 bis.** If  $H$  is of finite type over  $S$  and of constant type, one can also give the following isotriviality criterion, with no restriction on  $S$ :  $H = D_S(R)$  is isotrivial if and only if  $R$  is the union of a sequence of open and closed parts finite over  $S$ .<sup>44</sup>

**Lemma 5.13.** *Let  $S$  be a locally noetherian connected prescheme, let  $P$  be a quasi-isotrivial twisted constant  $S$ -prescheme, and let  $Z$  be an open and closed part of  $P$ . If  $Z_s$  is finite for some  $s \in S$ , then  $Z$  is finite over  $S$ .*

First, do not assume  $S$  connected. Let  $U$  be the set of  $s \in S$  for which  $Z_s$  is finite. We shall prove that  $U$  is both open and closed, and that  $Z|_U$  is finite over  $U$ . This statement is essentially equivalent to 5.13, but has the advantage of being local in  $S$  for the étale topology, say. We are therefore reduced to the case where  $P$  is constant, of the form  $I_S$  for a set  $I$ . (N.B. The locally noetherian hypothesis is preserved by étale base change; this is where the quasi-isotriviality of  $P$  over  $S$  is used.)

We may also suppose that  $S$  is connected, since its connected components are open ( $S$  being locally noetherian). Then necessarily  $Z = J_S$  for a subset  $J \subseteq I$ , and consequently  $U = \emptyset$  or  $U = S$  according as  $J$  is infinite or finite. This proves the assertion.

**Recalls 5.14.0.**<sup>45</sup> Let  $S$  be a locally noetherian prescheme and let  $\tilde{S}$  be the normalization of  $S_{\text{red}}$  (cf. EGA II, 6.3.8). Recall that  $S$  is said to be *geometrically unibranch* (cf. EGA 0<sub>IV</sub>, § 23.2 and IV<sub>2</sub>, § 6.15) if the canonical morphism  $\tilde{S} \rightarrow S$  is radicial, and hence a universal homeomorphism. In particular, the connected components of  $S$  are irreducible.

Suppose now that  $S$  is connected, and hence irreducible. Let  $\eta$  be its generic point and let  $f : P \rightarrow S$  be a flat, locally quasi-finite morphism. Let  $P_i$  be the irreducible components of  $P$ , with generic points  $\xi_i$ . Since  $P$  is flat over  $S$ , every  $\xi_i$  lies above  $\eta$  (cf. EGA IV<sub>2</sub>, 2.3.4), and

$$(P_i)_\eta = P_i \cap P_\eta$$

is the closure of  $\xi_i$  in  $P_\eta$ . Since the fiber  $P_\eta$  is discrete, one has  $(P_i)_\eta = \{\xi_i\}$ . This applies in particular when  $f$  is étale. In that case  $P$  is also locally noetherian and geometrically

<sup>42</sup>*Editors' note.* Indeed, it is a twisted constant group of type  $\text{End}_{\text{gr.}}(N)$ ; cf. 5.6 and 5.8.

<sup>43</sup>*Editors' note.* This is modulo verifying (ii)  $\Rightarrow$  (i) when  $S$  is locally noetherian and  $R$  is not finitely generated ...

<sup>44</sup>*Editors' note.* To spell this out: since  $M$  is a finitely generated  $\mathbb{Z}$ -module, it is countable, and the proof of 5.11 (i)  $\Rightarrow$  (ii) shows that  $R$  is the union of a countable family of open and closed parts finite over  $S$ . The converse would have to be checked ...

<sup>45</sup>*Editors' note.* The original treated in 5.14 the case where  $S$  is locally noetherian and normal, and observed in Remark 5.15 that the argument applies more generally when  $S$  is merely geometrically unibranch rather than normal. The statement of 5.14 (and also 5.16) has accordingly been modified, and these "recalls", drawn from EGA IV<sub>4</sub>, 18.10.6 and 18.10.7, have been added; they show that the proof of 5.14 applies verbatim to the geometrically unibranch case.

unibranch (cf. EGA IV<sub>4</sub>, 17.5.7), so its irreducible components are its connected components and are open (and closed).

**Corollary 5.14.** *Let  $S$  be a locally noetherian geometrically unibranch prescheme, and let  $P$  be a quasi-isotrivial twisted constant  $S$ -prescheme. Then the connected components of  $P$  are finite over  $S$ .*

We may plainly suppose that  $S$  is connected and hence irreducible, and write  $\eta$  for its generic point. By the preceding recalls, every connected component  $P_i$  of  $P$  is open and closed and meets the fiber  $P_\eta$  in a single point. Lemma 5.13 therefore applies and shows that every  $P_i$  is finite over  $S$ .

**Theorem 5.16.**<sup>46</sup> *Let  $S$  be a locally noetherian geometrically unibranch prescheme. Then every  $S$ -group of multiplicative type and of finite type is isotrivial.*

Indeed, we may suppose  $S$  connected, and hence  $H$  of constant type  $M$ . It is enough to apply 5.14 to

$$P = \text{Isom}_{S\text{-gr.}}(H, G), \quad G = D_S(M),$$

and then argue as in the proof of 5.11 (ii)  $\Rightarrow$  (i). Alternatively, one may apply 5.14 to  $P = R = D_S(H)$  (cf. 5.9), and then use 5.11.

## 6. Infinite Galois principal bundles and the enlarged fundamental group

(The results of the present section and of the next will not be used again in the remainder of this Seminar.)

Let  $S$  be a prescheme. We propose to determine the principal homogeneous bundles  $P$  over  $S$  whose structural group is of the form  $G_S$ , the constant  $S$ -group defined by an ordinary group  $G$ , not necessarily finite. We shall also call them *Galois principal bundles over  $S$  with group  $G$* . We use “principal bundle” in the sense of the faithfully flat quasi-compact topology (cf. Exposé IV, Def. 5.1.5). For such a  $P$ , however, the structural morphism  $P \rightarrow S$  is necessarily étale and surjective, hence a covering for the étale topology; consequently  $P$  is also locally trivial for the étale topology (cf. IV, Prop. 5.1.6).<sup>47</sup>

We assume that  $S$  is a sum of connected preschemes, that is, that its connected components are open; this reduces us at once to the case in which  $S$  is connected. Choose a “geometric point”  $\xi$  of  $S$ , that is, an  $S$ -scheme  $\xi$  which is the spectrum of an algebraically closed field  $\Omega = \kappa(\xi)$ . For every Galois principal bundle  $P$  over  $S$  with group  $G$ ,  $P_\xi$  is a Galois principal bundle over the algebraically closed field  $\kappa(\xi)$ , and is therefore trivial. We refine the initial problem by asking for the category of Galois principal bundles over  $S$  *pointed* above  $\xi$ , that is, endowed with an  $S$ -morphism  $\xi \rightarrow P$ , or equivalently with a trivialization of  $P_\xi$ . For fixed  $G$ , let  $\bar{\pi}^1(S, \xi; G)$  denote the set of classes of such bundles, up to an isomorphism respecting the base point. Then the set  $\bar{\pi}^1(S; G)$  of isomorphism classes of Galois principal bundles over  $S$  with group  $G$ , without a specified base point, is the set of orbits of  $G$  in  $\bar{\pi}^1(S, \xi; G)$ . Here  $G$  acts naturally by translating the marked point of a pointed Galois principal bundle:

$$\bar{\pi}^1(S; G) = \bar{\pi}^1(S, \xi; G)/G.$$

<sup>46</sup> *Editors' note.* Remark 5.15, rendered obsolete by the addition of 5.14.0 (cf. the preceding editors' note), has been deleted, and in 5.16 “normal” has been replaced by “geometrically unibranch”.

<sup>47</sup> *Editors' note.* Notice that  $P$  is assumed to be a prescheme. In the *a priori* more general case of an fpqc sheaf  $P$  which is a  $G_S$ -torsor, is  $P$  necessarily representable?

Let  $S' \rightarrow S$  be any morphism which is a morphism of universal effective descent for the fibered category of twisted constant preschemes over a variable base (for example,  $S' \rightarrow S$  faithfully flat and locally of finite presentation, cf. 5.4; for other examples, cf. SGA 1 IX). We propose to determine the subsets

$$\bar{\pi}^1(S'/S, \xi; G) \quad \text{and} \quad \bar{\pi}^1(S'/S; G)$$

of the preceding sets, formed by the Galois principal bundles over  $S$  which become trivial over  $S'$ , or, as one says, are “split” by  $S'$ . In fact, we shall determine the category of Galois principal bundles  $P$  over  $S$  split by  $S'$ . Of course,

$$\bar{\pi}^1(S, \xi; G) = \varinjlim_{S'} \bar{\pi}^1(S'/S, \xi; G),$$

where  $S'$  runs through a cofinal system among the  $S'/S$  which cover for the étale topology (for example, if  $S$  is quasi-compact, among the quasi-compact  $S'$  over  $S$  whose structural morphism is étale and surjective). Likewise, the category of Galois principal bundles over  $S$  is the inductive limit of the subcategories defined by the  $S'$ , formed by the bundles split over  $S'$ .

By the assumption on  $S'/S$ , the category of Galois principal bundles over  $S$  split by  $S'$  is equivalent to the category of trivial Galois principal bundles over  $S'$ , hence of the form  $G_{S'}$ , with  $G$  acting by right translations, equipped with descent data relative to  $S' \rightarrow S$ . In terms of the corresponding trivial bundle  $P'$  over  $S'$  and its descent datum, a base point on a Galois principal bundle  $P$  over  $S$  split by  $S'$  amounts to a trivialization of  $P' \times_{S'} S'_\xi$  compatible with the induced descent datum relative to  $S'_\xi \rightarrow \xi$ , where  $S'_\xi = S' \times_S \xi$ ; equivalently, it is a section  $\sigma$  of  $P'_\xi$  over  $S'_\xi$  compatible with the descent datum.

It is therefore useful, for an arbitrary  $S$ -prescheme  $S'$ , without now assuming that  $S' \rightarrow S$  is a morphism of universal effective descent for the fibered category of twisted constant bundles, to define  $\bar{\pi}^1(S'/S; G)$  and  $\bar{\pi}^1(S'/S, \xi; G)$  as the sets of isomorphism classes of the structures with descent data just specified. These functors in  $G$  then have a very simple simplicial description in terms of the fibered square and cube  $S''$  and  $S'''$  of  $S'$  over  $S$ ; we shall sketch it below (cf. 6.3).

The important conclusion is the following.

**Proposition 6.1.** *Suppose that the connected components of  $S'$  and  $S''$  are open, and, for example, that the quotient of  $\pi_0(S')$  by the equivalence relation induced by the two projections  $\pi_0(S'') \rightarrow \pi_0(S')$  is a single point.<sup>48</sup>*

(i) *The functor*

$$G \longmapsto \bar{\pi}^1(S'/S, \xi; G)$$

*from the category of groups to the category of sets is representable by a group, denoted  $\pi_1(S'/S, \xi)$  and called the fundamental group of  $S$  at  $\xi$  relative to  $S' \rightarrow S$ . Thus there is a functorial bijection*

$$\bar{\pi}^1(S'/S, \xi; G) \simeq \text{Hom}_{\text{gr.}}(\pi_1(S'/S, \xi), G).$$

(ii) *This group has a set of generators in bijection with  $\pi_0(S'')$ , and is described in terms of those generators by relations in bijection with the elements of  $\pi_0(S''')$ .<sup>49</sup> In particular,*

<sup>48</sup>*Editors' note.* The hypothesis that the connected components of  $S'$  be open, as well as the second hypothesis, has been added. The latter says that the simplicial set  $K_\bullet = \pi_0(S_\bullet)$  defined in 6.3 is connected. In fact a weaker hypothesis suffices: the cone  $\tilde{K}_\bullet$  of a certain morphism of simplicial sets  $k_\bullet \rightarrow K_\bullet$  need only be connected (cf. *loc. cit.*).

<sup>49</sup>*Editors' note.* The superfluous hypothesis that the connected components of  $S'''$  be open has been deleted. On the other hand, the later description (cf. . . .) gives the natural set of generators  $\pi_0(S'') \amalg \pi_0(S'_\xi)$ ; the relations between these generators arising from the 2-cells then reduce it to  $\pi_0(S'')$ . See, for example, [Kan58], § 19 for a finer description.



$\pi_1(S'/S, \xi)$  is finitely generated (respectively finitely presented) if  $\pi_0(S'')$  is finite (respectively if both  $\pi_0(S'')$  and  $\pi_0(S''')$  are finite).

(iii) The category of Galois principal bundles over  $S$  split by  $S'$ , with base point above  $\xi$ , is equivalent to the category of ordinary groups  $G$  equipped with a homomorphism  $\pi_1(S'/S, \xi) \rightarrow G$ .

The proof is given below; cf. ...

**6.2.** When  $S$  is a connected locally noetherian prescheme, every étale prescheme  $S'$  over  $S$  is locally noetherian and therefore has open connected components. It follows from the preceding discussion<sup>50</sup> that the functor

$$G \longmapsto \bar{\pi}^1(S, \xi; G)$$

from ordinary groups to sets is *strictly pro-representable* (cf. *Séminaire Bourbaki*, February 1960, No. 195, §§ A.2 and A.3). Thus there is a projective system

$$\Pi = \Pi_1(S; \xi) = (\pi_i)_{i \in I}$$

of ordinary groups over an increasing filtered index set  $I$ , which is “strict”, meaning that its transition morphisms  $\pi_j \rightarrow \pi_i$  are surjective, together with an isomorphism of functors in  $G$

$$\bar{\pi}^1(S, \xi; G) \simeq \varprojlim_i \text{Hom}_{\text{gr.}}(\pi_i, G).$$

The second member is also denoted  $\text{Hom}_{\text{pro-gr.}}(\Pi, G)$  (cf. *loc. cit.*).

If the projective limit

$$\pi = \varprojlim_i \pi_i$$

is “large enough”, more precisely if the canonical homomorphisms  $\pi \rightarrow \pi_i$  are surjective, it is natural to equip  $\pi$  with the projective-limit topology of the discrete topologies on the  $\pi_i$ . The preceding isomorphism can then also be written

$$\bar{\pi}^1(S, \xi; G) \simeq \text{Hom}_{\text{gr. top.}}(\pi, G),$$

where the right-hand side is the set of homomorphisms of topological groups,  $G$  being given the discrete topology.

The hypothesis just formulated on the projective system  $\Pi$  holds, as is well known, when the  $\pi_i$  are finite groups (cf. [BEns], III § 7.4, Th. 1). This last condition plainly also means that every Galois principal bundle over  $S$  is isotrivial, that is, split by a finite étale surjective morphism. This is the case when  $S$  is geometrically unibranch (for example normal), as follows at once from 5.14.<sup>51</sup> When the  $\pi_i$  are finite,  $\pi$  also coincides with the fundamental group  $\pi_1(S, \xi)$  introduced in SGA 1, V.

Thus, in the favorable case in which the maps  $\pi \rightarrow \pi_i$  are surjective, one might call  $\pi$  the *enlarged fundamental group of  $S$  at  $\xi$* . Outside this favorable case,  $\pi$  itself is of little interest, and the role of the usual fundamental group is played by the projective system  $\Pi$  itself, which we call the *enlarged fundamental pro-group of  $S$  at  $\xi$* . (Any better terminological suggestion than “enlarged” is welcome!<sup>52</sup>) Knowledge of this pro-group is more precise than knowledge of the usual fundamental group  $\pi_1(S, \xi)$  of SGA 1, V; more precisely, the latter is the projective limit of the projective system formed by the finite quotients of the  $\pi_i$ .

<sup>50</sup> *Editors' note.* This deduction could be spelled out:  $I$  is in bijection with a cofinal set of morphisms  $S' \rightarrow S$  covering for the étale topology ...

<sup>51</sup> *Editors' note.* 5.13 has been corrected to 5.14.

<sup>52</sup> *Editors' note.* Some authors speak of the “true” fundamental group.

**6.3.** Let us indicate briefly the “calculation” of  $\pi_1(S'/S, \xi)$ . Let  $S_i$  be the  $(i+1)$ -st fibered power of  $S'$  over  $S$ , so that  $S_0 = S'$ ,  $S_1 = S''$ , and so on. The evident simplicial operations between the  $S_i$  make  $(S_i)_{i \in \mathbb{N}}$  a simplicial object of  $(\text{Sch})/S$ .

Applying to this simplicial object the functor “set of connected components”

$$\pi_0 : (\text{Sch})/S \longrightarrow (\text{Ens}),$$

we obtain a simplicial set  $K_\bullet = (K_i)_{i \in \mathbb{N}}$ , with

$$K_i = \pi_0(S_i).$$

Likewise, the  $S_{i,\xi}$ , the  $(i+1)$ -st fibered powers of  $S'_\xi$  over  $\xi$ , form a simplicial object of  $(\text{Sch})/\xi$ , hence of  $(\text{Sch})/S$ , together with a natural morphism of simplicial objects into  $(S_i)_{i \in \mathbb{N}}$ . Thus we obtain a simplicial set  $k_\bullet$ , with

$$k_i = \pi_0(S_{i,\xi}),$$

and a canonical morphism

$$k_\bullet \longrightarrow K_\bullet.$$

Taking the cone of this morphism (cf. 9.5.1), we form a new simplicial set

$$\tilde{K}_\bullet = \text{Cone}(k_\bullet \longrightarrow K_\bullet).$$

<sup>53</sup> In this way we obtain a “pointed simplicial set”  $\tilde{K}_\bullet$ , that is, a simplicial set equipped with a morphism

$$\tilde{\xi} : e_\bullet \longrightarrow \tilde{K}_\bullet,$$

where  $e_\bullet$  is the final simplicial set. We may form its familiar combinatorial invariants

$$\pi_0(\tilde{K}_\bullet, \tilde{\xi}) \quad \text{and} \quad \pi_1(\tilde{K}_\bullet, \tilde{\xi}),$$

whose constructions involve only the components of degree at most 1 and at most 2, respectively. These invariants are defined without any restriction on  $S$  or  $S'$ . When the connected components of  $S_0$  and  $S_1$  are open and  $\tilde{K}_\bullet$  is connected,<sup>54</sup> one checks without difficulty that  $\pi_1(\tilde{K}_\bullet, \tilde{\xi})$  represents the functor  $G \mapsto \bar{\pi}^1(S'/S, \xi; G)$ ; that is,

$$\pi_1(S'/S; \xi) \simeq \pi_1(\tilde{K}_\bullet, \tilde{\xi}).$$

We also note that if  $S' \rightarrow S$  is “universally submersive” (cf. SGA 1, IV 2.1) and the connected components of  $S'$  are open, then the simplicial set  $K_\bullet$ , and hence also  $\tilde{K}_\bullet$ , is connected.<sup>55</sup>

**Examples 6.4.** It remains to give examples of enlarged fundamental groups. Return to the examples of 1.6. Let  $k$  be an algebraically closed field; let  $C_1$  be a complete rational curve over  $k$  with exactly one singular point, that point being an ordinary double point; and let  $C_2$  be a curve which is the union of two irreducible components, each isomorphic to

<sup>53</sup> *Editors' note.* The original, which takes  $\tilde{K}_\bullet = K_\bullet/k_\bullet$ , has been corrected: on the one hand  $k_\bullet$  is already contractible (cf. 9.7.1), and on the other hand  $k_\bullet \rightarrow K_\bullet$  is not in general injective. It is an epimorphism if  $S'/S$  is finite étale (cf. 9.7.3).

<sup>54</sup> *Editors' note.* The hypothesis that  $\tilde{K}_\bullet$  be connected has been added; for the proof, see the addendum below (Section 9).

<sup>55</sup> *Editors' note.* The sequel has been modified to take account of the correction made above; cf. Editors' note (53). (The original read: “ $\pi_0(\tilde{K}_\bullet, \tilde{\xi})$  is also canonically isomorphic to the set  $\pi_0(S, \xi)$  of connected components of  $S$ , pointed by the connected component of  $\xi$  in  $S$ .”)

$\mathbb{P}_k^1$ , meeting in exactly two points, which are ordinary double points of  $C_2$ . In either case, the enlarged fundamental group of the curve is isomorphic to  $\mathbb{Z}$ .<sup>56</sup>

More generally, one should take up, simplifying and correcting them, the results of SGA 1, IX 5 in the setting of the enlarged fundamental group. The examples of *loc. cit.*, 5.5 would give as many examples of enlarged fundamental pro-groups which are not profinite. Thus, if in the first example the ordinary double point were replaced by a double point with  $n$  distinct branches,<sup>57</sup> the enlarged fundamental group would be the free (discrete!) group on  $n - 1$  generators.

## 7. Classification of twisted constant preschemes and of groups of multiplicative type of finite type in terms of the enlarged fundamental group

### 7.0.

Let  $S$  be a prescheme, which we continue to assume locally noetherian, in order to ensure that  $S$ , and certain preschemes over  $S$  that we shall consider (notably those étale over  $S$ , and more generally those locally of finite type over  $S$ ), are locally connected.

**Proposition 7.0.1.**— *Every twisted constant prescheme  $X$  over  $S$  that is locally trivial for the (fppf) topology (that is, that is split by a faithfully flat morphism  $S' \rightarrow S$  locally of finite presentation) is quasi-isotrivial (that is, one may even choose  $S' \rightarrow S$  étale and surjective).*

Indeed, we may suppose that  $S$  is connected, and hence that  $X$  is of type  $I$ , where  $I$  is a fixed set. Thus

$$X' = X \times_S S'$$

is isomorphic to  $I_{S'}$ , so  $I_{S'}$  is endowed with descent data relative to  $S' \rightarrow S$ ; in other words, there is an isomorphism

$$I_{S''} \xrightarrow{\sim} I_{S''}$$

satisfying the usual transitivity condition. Now  $S'' = S' \times_S S'$  is locally noetherian and hence locally connected. It follows that the automorphisms of  $I_{S''}$  correspond to the sections of  $G_{S''}$ , where  $G = \text{Aut}(I)$  is the group of permutations of  $I$ .

In this way one obtains descent data on  $G_{S'}$  (regarded as a trivial Galois principal bundle) relative to  $S' \rightarrow S$ . By 5.4 these descent data are effective, and therefore give a Galois principal bundle  $P$  over  $S$  with group  $G$ . By construction,  $P$  represents the functor  $\text{Isom}_S(I_S, X)$  in the category of preschemes over  $S$  that are locally noetherian. Consequently, the étale surjective base change  $P \rightarrow S$  splits  $X$ , so  $X$  is indeed quasi-isotrivial.

**Remark 7.0.2.**— One should bear in mind that, even when  $S$  is the spectrum of a field, it is not true in general that every twisted constant bundle over  $S$  is quasi-isotrivial. For example, one may take  $X$  to be the disjoint union of a sequence of schemes of the form  $\text{Spec}(k_i)$ , where the  $k_i$  are separable extensions of  $k$  of strictly increasing degrees.

<sup>56</sup> *Editors' note.* This could be spelled out. First, by 5.14, the enlarged fundamental group of the projective line  $\mathbb{P}_k^1$  is zero; that is,  $\mathbb{P}_k^1$  is “truly” simply connected. One would then have to extend Corollary 5.4 of SGA 1, IX, and the discussion following it, to the enlarged fundamental group and to the category of principal bundles not necessarily finite. Let  $\Gamma_1, \Gamma_2$  be two copies of  $\mathbb{P}_k^1$ , and let  $a_i, b_i$  be two distinct points of  $\Gamma_i$ . Let  $C'_2$  be the disjoint union of  $\Gamma_1$  and  $\Gamma_2$ , and let  $C'_2$  be the curve obtained by identifying  $a_1$  with  $a_2$ ; then  $C_2$  is obtained from  $C'_2$  by also identifying  $b_1$  with  $b_2$ . The discussion following *loc. cit.*, extended to the enlarged case, shows that the enlarged fundamental group of  $C'_2$ , respectively  $C_2$ , is zero, respectively  $\mathbb{Z}$ .

<sup>57</sup> *Editors' note.* This means an  $n$ -fold point with  $n$  distinct tangents, for example the curve  $X^n - Y^n = X^{n+1}$ .

The proof just given shows at the same time that the classification of quasi-isotrivial twisted constant bundles  $X$  over  $S$  of type  $I$  is equivalent to that of Galois principal bundles over  $S$  with group  $G = \text{Aut}(I)$ . Indeed, this is an equivalence of categories.

It can be put into a more convenient form, as in SGA 1, V. To this end, suppose that  $S$  is connected and equipped with a geometric point  $\xi$ . The enlarged fundamental pro-group  $\Pi = \Pi_1(S, \xi)$  is then defined. For every quasi-isotrivial twisted constant bundle  $X$  over  $S$ , let  $I = X(\xi)$  be its set-theoretic fiber at  $\xi$ . Thus  $X$  is of type  $I$ , and is consequently associated, as above, with a Galois principal bundle

$$P = \text{Isom}_S(I_S, X)$$

over  $S$ , with group  $G = \text{Aut}(I)$ .

By the definition of  $\Pi$ , one therefore obtains a canonical homomorphism from  $\Pi$  to  $G$ , that is, from one of the  $\Pi_i$  to  $G$ . Since  $G$  is the group of permutations of  $I = X(\xi)$ , this says that  $\Pi$  “acts continuously on  $I = X(\xi)$ ”, meaning that the  $\pi_i$ , for  $i$  sufficiently large, act on  $I$  compatibly with the transition morphisms.

We leave it to the reader to verify that every  $S$ -morphism  $X \rightarrow Y$  between quasi-isotrivial twisted constant bundles over  $S$  induces a map  $X(\xi) \rightarrow Y(\xi)$  compatible with the actions of  $\Pi$ , and that the resulting functor is an equivalence of categories:

**Proposition 7.0.3.**— *Let  $S$  be a locally noetherian connected prescheme, let  $\xi$  be a geometric point of  $S$ , and let  $\Pi = \Pi_1(S, \xi)$  be the enlarged fundamental pro-group of  $S$  at  $\xi$ . Then the functor*

$$X \longmapsto X(\xi)$$

*is an equivalence between the category of quasi-isotrivial twisted constant bundles over  $S$  and the category of sets on which  $\Pi$  acts continuously.*

This functor is compatible with finite sums and finite inverse limits. It follows, for example, that quasi-isotrivial twisted constant groups (or rings, and so forth) over  $S$  correspond to ordinary groups (respectively rings, and so forth) on which the pro-group  $\Pi$  acts continuously. In particular:

**Corollary 7.0.4.**— *The category of quasi-isotrivial twisted constant commutative groups over  $S$  is equivalent to the category of “ $\Pi$ -modules”, that is, commutative groups  $M$  on which  $\Pi$  acts continuously.*

Using 5.7, we now obtain:

**Theorem 7.1.**— *Let  $S$  be a locally noetherian connected prescheme, let  $\xi$  be a geometric point of  $S$ , and let  $\Pi = \Pi_1(S, \xi)$  be the enlarged fundamental pro-group of  $S$  at  $\xi$ . Then the functor*

$$G \longmapsto \text{Hom}_{\kappa(\xi)\text{-gr}}(G_\xi, \mathbf{G}_{m, \xi})$$

*induces an anti-equivalence between the category of quasi-isotrivial groups of multiplicative type over  $S$  and the category of  $\Pi$ -modules.*

Using 4.5, we obtain:

**Corollary 7.2.**— *The preceding functor induces an anti-equivalence between the category of groups of multiplicative type and of finite type over  $S$ , and the category of  $\Pi$ -modules that are of finite type over  $\mathbb{Z}$ .*

**Example 7.3.**— Take for  $S$ , for example, a complete rational curve over an algebraically closed field, having exactly one multiple point with  $n + 1$  distinct branches. By 6.4, the enlarged fundamental group  $\Pi(S, \xi)$  is a free group on  $n$  generators. Thus, by 7.2, the classification of tori of relative dimension  $m$  over  $S$  is equivalent to the classification of

systems of  $n$  endomorphisms  $A_1, \dots, A_n$  of the  $\mathbb{Z}$ -module  $\mathbb{Z}^m$ , up to automorphism of  $\mathbb{Z}^m$ . Except when  $n \leq 1$  or  $m \leq 1$ , an explicit classification of such systems seems hopeless. One can at least define a multitude of nontrivial invariants for such a system, such as the characteristic polynomials of the  $A_i$ .<sup>58</sup>

**Remark 7.4.**— Without any hypothesis on  $S$ , it remains true that, for a given ordinary commutative group  $M$  of finite type over  $\mathbb{Z}$ , the category of groups of multiplicative type of type  $M$  over  $S$  is anti-equivalent to the category of Galois principal bundles over  $S$  with group  $G = \text{Aut}_{\text{gr}}(M)$ . This follows easily from 5.9 and 5.10.

**Remark 7.5.**— The theory of the fundamental pro-group sketched in the present two sections is more advantageously formulated in the setting of general sites. In that form it applies equally well, for example, to ordinary topological spaces, and gives a satisfactory theory at least for a locally connected topological space (not necessarily locally simply connected). Here too it seems that one cannot make do with a fundamental group, and that a pro-group is needed. Finally, once one has at one's disposal the language of topologies and descent (which really lies at the heart of these questions), the account sketched here is also technically simpler than that of SGA 1, V, and should therefore in principle replace it.

## 8. Appendix: Elimination of certain affineness hypotheses

Our aim is to prove the following generalization of 4.9.

**Theorem 8.1.**— *Let  $S$  be a prescheme, and let  $H$  be a flat  $S$ -prescheme in groups of finite presentation, with connected affine fibers.<sup>59</sup> Let  $s \in S$ , and suppose that  $H_s$  is a torus. Then there exists an open neighborhood  $U$  of  $s$  such that  $H|_U$  is a torus.*

We immediately obtain:

**Corollary 8.2.**— *Let  $H$  be an  $S$ -prescheme in groups, flat and of finite presentation over  $S$ . Then  $H$  is a torus if and only if its fibers are tori.*

**Remark 8.3.**— Even when  $S$  is the spectrum of a discrete valuation ring, the hypothesis in 8.1 that the fibers of  $H$  (here, the generic fiber) be affine cannot be dropped: there are examples of smooth groups over  $S$  whose generic fiber is an elliptic curve and whose special fiber is  $\mathbf{G}_m$ .

*Proof of 8.1.* We may clearly suppose that  $S = \text{Spec}(A)$  is affine. By the standard procedure (cf. EGA IV<sub>2</sub>, 8.8.2), this reduces us to the case in which  $S$  is moreover noetherian. We begin by proving 8.2 in this case.

By 4.9, we are reduced to proving that  $H$  is affine over  $S$ . We may therefore suppose<sup>60</sup> that  $A$  is noetherian and local. Since its completion  $\hat{A}$  is faithfully flat over  $A$ , descent reduces us to the case in which  $A$  is a complete noetherian local ring. Let  $S'$  be the normalization of  $S_{\text{red}}$ . Nagata's theorem says that it is finite over  $S$  (EGA 0<sub>IV</sub>, 23.1.5). Moreover,  $S' \rightarrow S$  is surjective, so

$$H' = H \times_S S' \longrightarrow H$$

is finite and surjective. Hence, to prove that  $H$  is affine, it suffices to show that  $H'$  is affine (EGA II, 6.7.1).

<sup>58</sup> *Editors' note.* In particular, this shows that the tori of relative dimension 2 considered in 1.6 are non-isotrivial.

<sup>59</sup> *Editors' note.* Observe that the hypotheses imply that  $H$  is separated over  $S$ , by VI<sub>B</sub>, Theorem 5.3 or Corollary 5.5.

<sup>60</sup> *Editors' note.* By EGA IV<sub>2</sub>, 8.8.2 once again.

Replacing  $S$  by a connected component of  $S'$ , we are reduced to the case in which  $A$  is a noetherian (semilocal), normal, integral ring.<sup>61</sup> Moreover, after possibly replacing  $A$  by its normalization in a finite separable extension of its field of fractions, we may suppose that the generic fiber  $H_\eta$  of  $H$  is diagonalizable, that is, that there is an isomorphism

$$u_\eta : H_\eta \xrightarrow{\sim} T_\eta, \quad T = \mathbf{G}_{m,S}^r.$$

We now have:

**Lemma 8.4.**— *Let  $S$  be a locally noetherian normal irreducible prescheme with generic point  $\eta$ . Let  $H$  be a smooth prescheme in groups over  $S$ , with connected fibers, let  $T$  be a prescheme in groups of multiplicative type and of finite type over  $S$ , and let  $u_\eta : H_\eta \rightarrow T_\eta$  be a homomorphism of algebraic groups over  $\kappa(\eta)$ . Then  $u_\eta$  extends to a homomorphism of groups  $u : H \rightarrow T$ .*

After possibly replacing  $S$  by its normalization in a finite separable extension of its function field, we may suppose that  $T$  is diagonalizable (which is already the case in the application we have in mind). Then  $T$  is a closed subgroup of a group of the form  $\mathbf{G}_{m,S}^r$ , reducing us to the case  $T = \mathbf{G}_{m,S}$ .

It remains only to prove that  $u_\eta$ , regarded as a rational map from  $H$  to  $T = \mathbf{G}_{m,S}$ , is everywhere defined (for then the morphism  $u : H \rightarrow T$  extending it is necessarily a group homomorphism). We may regard  $u_\eta$  as an invertible section  $f$  of the structure sheaf of  $H_\eta$ , and must show that it extends to an invertible section of the structure sheaf of  $H$ . Since  $H$  is smooth over the normal prescheme  $S$ , it is normal (SGA 1, II 3.1). It therefore suffices to find a closed subset  $Z$  of  $H$ , of codimension at least 2, such that  $f$  extends to an invertible section of the structure sheaf of  $H - Z$ . By localizing at the codimension-one points of  $S$ , this at once reduces us to the case in which  $S$  is the spectrum of a discrete valuation ring  $A$ .

Let  $t$  be a uniformizer of  $A$ , and let  $t'$  be the section of  $\mathcal{O}_H$  that it defines, so that the special fiber  $H_0$  equals  $V(t')$ . By hypothesis,  $H_0$  is smooth and connected over the residue field  $k$ . The function  $f$  is then a rational function on  $H$  with neither zeros nor poles in  $H - H_0$ . Since  $H_0 = V(t')$  is an irreducible divisor, there is an integer  $n \in \mathbb{Z}$  such that

$$t'^n f = t^n f$$

has neither zeros nor poles, and hence is an invertible section of  $\mathcal{O}_H$ . It therefore defines a morphism  $v : H \rightarrow \mathbf{G}_{m,S}$ . Since  $v_\eta = t'^n u_\eta$ , and  $u_\eta$  is a group homomorphism,  $v$  sends the identity section of  $H$  to a section of  $\mathbf{G}_{m,S}$  whose value at the generic point of  $S$  is  $t^n$ . Because this is a section of  $\mathbf{G}_{m,S}$ ,  $t^n$  must be a unit; thus  $n = 0$ . Hence  $v$  extends  $u_\eta$ , completing the proof of 8.4.

Applying this lemma in the present situation, we obtain a group homomorphism

$$u : H \longrightarrow T = \mathbf{G}_{m,S}^r$$

that induces an isomorphism on the generic fibers. We shall prove that  $u$  is an isomorphism.

**Lemma 8.5.**— *For every integer  $n > 0$  prime to the residual characteristic of  $A$ , the group*

$${}_n H = \ker(n \cdot \text{id}_H)$$

*is finite over  $S$ .*

If  $n$  is prime to the residual characteristic of  $A$ , then it is prime to all the residual characteristics of the points of  $S$ . Consequently,  $n \cdot \text{id}_H$  induces an étale morphism on every

<sup>61</sup> *Editors' note.* The original has been expanded in what follows.

fiber of  $H$ , and hence is itself étale (SGA 1, I 5.9); its kernel  ${}_nH$  is therefore étale over  $S$ . On the other hand,  ${}_nH$  is separated over  $S$ , because  $H$  is. \*<sup>62</sup> Moreover, all its fibers have the same rank  $n^r$ , since the fibers of  $H$  are tori, all of the same dimension  $r$  (as  $H$  is smooth over  $S$ ). We conclude that  ${}_nH$  is finite over  $S$  (SGA 1, I 10.9, or EGA IV<sub>4</sub>, 18.2.9).

By the preceding lemma,  $u({}_nH)$  is therefore a closed subset of  $T$ . On the generic fiber  $T_\eta$ , this subset is identical to  ${}_n(T_\eta)$ , so it contains the closure of that subset, namely  ${}_nT$ . For every  $s \in S$ , the groups  ${}_n(T_s)$ , as  $n$  varies, are dense in  $T_s$ ; since  $u_s(H_s)$  is a closed subset (VI<sub>B</sub>, 1.4.2) containing them, we have  $u_s(H_s) = T_s$ . Thus  $u$  is surjective. A surjective homomorphism between tori of the same dimension over a field is flat,<sup>63</sup> and hence  $u$  induces a flat morphism on every fiber. Therefore  $u$  is flat (SGA 1, I 5.9). Consequently  $K = \ker(u)$  is flat over  $S$ ,<sup>64</sup> and hence equals the closure of its generic fiber  $K_\eta$ . By construction,  $K_\eta$  is the identity group. Since  $K$  is separated over  $S$  (because  $H$  is), its identity section is closed, and it follows that  $K$  is the identity group. Thus  $u$  is a monomorphism; since we have seen that it is faithfully flat, it is an isomorphism (cf. SGA 1, I 5.1, or EGA IV<sub>4</sub>, 17.9.1). This proves that  $H$  is a torus, and completes the proof of 8.2.<sup>65</sup>

**Remark 8.5.1.**— Instead of invoking 8.5, one can also invoke Zariski's "Main Theorem", which directly implies that  $u$  is an open immersion and hence an isomorphism.<sup>66</sup>

To prove 8.1, the quasi-isotriviality theorem reduces us to the case in which  $S$  is local and  $s$  is its closed point,<sup>67</sup> and it remains to prove that, under the other hypotheses,  $H$  is then a torus. By 8.2, already proved, we are reduced to showing that the fibers of  $H$  are tori. We may suppose that  $S$  is the spectrum of a complete noetherian local ring  $A$ . By 3.3, for every  $n > 0$  there exists a group of multiplicative type  $T_n$ , finite over  $S$ , and an isomorphism

$$(T_n)_s \simeq {}_n(H_s),$$

where  $s$  is the closed point of  $S$ . Arguing as in 3.1 and using the fact that  $T_n$  is finite over  $S$ ,<sup>68</sup> we see that the preceding isomorphism comes from a homomorphism  $u_n : T_n \rightarrow H$ , which is moreover uniquely determined. (Pass to the limit over the artinian quotients of  $A$ .)

Moreover, by the uniqueness properties, when  $m$  is a multiple of  $n$ , the homomorphism  $u_n : T_n \rightarrow H$  is obtained from  $u_m : T_m \rightarrow H$  by restriction to  ${}_n(T_m) \simeq T_n$ . It follows from IX 6.6 that the  $u_n$  are monomorphisms, so the  $T_n$  are subgroups of  $H$ , and for  $m$  a multiple of  $n$  one has

$$T_n = {}_n(T_m).$$

Thus, for every  $t \in S$ , the groups  $(T_n)_t$  are subgroups of  $H_t$ , of type  $(\mathbb{Z}/n\mathbb{Z})^r$ , where  $r = \dim H_s = \dim H_t$ , and when  $m$  is a multiple of  $n$  one has

$$(T_n)_t = {}_n(T_m)_t.$$

The fact that  $H_t$  is a torus now follows from:

**Lemma 8.6.**— *Let  $H$  be a smooth affine algebraic group over an algebraically closed field  $k$ , and let  $(T_n)_{n>0}$  be a family of subgroups of multiplicative type of  $H$ , such that for every*

\*By Raynaud's theorem, VI<sub>B</sub>, 5.3.

<sup>62</sup>*Editors' note.* This point should be expanded in connection with the modifications in VI<sub>B</sub>, §5.

<sup>63</sup>*Editors' note.* This follows, for example, from VIII 3.2(a). More generally, if  $f : G \rightarrow H$  is a surjective morphism of algebraic groups over a field  $k$ , and  $H$  is reduced, then  $f$  is flat (cf. VI<sub>B</sub>, 1.3).

<sup>64</sup>*Editors' note.* The original has been expanded in what follows.

<sup>65</sup>*Editors' note.* At least in the case considered up to this point, namely  $S$  locally noetherian.

<sup>66</sup>*Editors' note.* The editors did not understand this remark, since they did not see why  $u$  would *a priori* have finite and surjective fibers.

<sup>67</sup>*Editors' note.* Expand this reduction.

<sup>68</sup>*Editors' note.* Expand this point.

integer  $n > 0$ ,  $T_n$  is of type  $(\mathbb{Z}/n\mathbb{Z})^r$ , and for every multiple  $m$  of  $n$ , one has  $T_n = {}_n(T_m)$ . Under these conditions,  $H$  contains a torus of dimension at least  $r$  containing the  $T_n$ . Consequently, if  $H$  is connected and has dimension at most  $r$ , then  $H$  is a torus of dimension  $r$ .

This is an exercise on affine algebraic groups, which we shall handle by repeated reference to *Bible*. We restrict attention to the  $T_n$  for  $n$  prime to the characteristic. Let  $K$  be the closure of the union of the  $T_n$  in  $H$ , endowed with the induced reduced structure. Standard arguments then show that  $K$  is a commutative algebraic subgroup of  $H$ . By *Bible*, §4.5, Theorem 4,  $K$  is isomorphic to a product  $K_u \times K_s$ , with  $K_u$  “unipotent” and  $K_s$  diagonalizable. Every diagonalizable subgroup of  $K$  is contained in  $K_s$ , so the  $T_n$  are subgroups of  $K_s$ , and therefore  $K = K_s$ . Write  $K = D(M)$ , where  $M$  is an ordinary commutative group of finite type over  $\mathbb{Z}$ . The inclusion  $T_n \subseteq K$  means that  $M$  has a quotient group isomorphic to  $(\mathbb{Z}/n\mathbb{Z})^r$ . Since this holds for every integer  $n$  prime to the characteristic of  $k$  (powers of a single fixed prime would suffice),  $M$  has rank at least  $r$ . Thus  $K$  contains a torus  $T$  of dimension  $r$ . When  $H$  is connected of dimension  $r$ , it follows that  $T = H$ , completing the proof of 8.6. Theorem 8.1 is now proved.

**Remarks 8.7.**— Using 8.1, it should not be difficult to give analogous generalizations of 4.7 and 4.8. A more interesting problem would be to study the situation of 8.1 after dropping the hypothesis that the fibers of  $H$  be affine. One can show that there is then an open neighborhood  $U$  of  $s$  such that  $H|_U$  is commutative and, for every  $t \in U$ , the geometric fiber  $H_t$  is an extension of an abelian variety by a torus.<sup>69</sup> Of course, in questions of this kind one may restrict to the case in which  $S$  is the spectrum of a discrete valuation ring,  $s$  its closed point, and  $t$  its generic point.

This result can be generalized as follows. For every smooth connected algebraic group  $G$  over an algebraically closed field  $k$ , a well-known theorem of Chevalley says that  $G$  is, in a unique way, an extension of an abelian variety  $A$  by a smooth connected affine group  $V$ . Define the *abelian rank* (respectively, *reductive rank*, *nilpotent rank*, and *semisimple dimension*) of  $G$ , denoted  $\rho_{\text{ab}}(G)$  (respectively,  $\rho_r(G)$ ,  $\rho_n(G)$ , and  $d_s(G)$ ), to be the dimension of  $A$  (respectively, the dimension of the maximal tori of  $G$ , the dimension of the Cartan subgroups<sup>70</sup> of  $G$ , and the dimension of the quotient of  $G$ , or equivalently of  $V$ , by its radical); see *Bible* for all these notions. Also define the *unipotent rank* by

$$\rho_u(G) = \rho_u(V) = \rho_n(G) - \rho_r(G) - \rho_{\text{ab}}(G).$$

When the ground field is not algebraically closed, the same names and notations designate the corresponding invariants of  $G_{\bar{k}}$ , where  $\bar{k}$  is an algebraic closure of  $k$ .

Now let  $G$  be a smooth group scheme over the spectrum  $S$  of a discrete valuation ring. Let  $\rho_{\text{ab}}$ , and so on (respectively,  $\rho'_{\text{ab}}$ , and so on), denote the invariants associated with the special fiber (respectively, the generic fiber). Then one has the inequalities

$$\begin{cases} \rho_{\text{ab}} \leq \rho'_{\text{ab}}, \\ \rho_r + \rho_{\text{ab}} \leq \rho'_r + \rho'_{\text{ab}}, \\ d_s \leq d'_s, \end{cases} \quad \begin{cases} \rho_n \geq \rho'_n, \\ \rho_u \geq \rho'_u. \end{cases}$$

Equivalently, if  $G$  is smooth of finite type over an arbitrary base  $S$ , the functions

$$s \longmapsto \rho_{\text{ab}}(s), \quad s \longmapsto \rho_{\text{ab}}(s) + \rho_r(s), \quad s \longmapsto d_s(s)$$

<sup>69</sup> *Editors' note.* Give a reference here?

<sup>70</sup> *Editors' note.* Recall the definition of “Cartan subgroup”.



are lower semicontinuous, while the functions

$$s \longmapsto \rho_n(s), \quad s \longmapsto \rho_u(s)$$

are upper semicontinuous.<sup>71</sup>

The same results probably remain valid without assuming  $G$  smooth over  $S$ , but assuming only that it is flat and of finite presentation over  $S$ , provided that, for an algebraic group  $G$  over an algebraically closed field  $k$ ,  $\rho_{\text{ab}}(G)$ , and so on, are understood to mean the corresponding invariants of  $G_{\text{red}}^0$ .

In this Seminar we present some results of this kind when  $G$  is affine over  $S$ , or more generally has affine fibers. In this case we shall verify the semicontinuity properties for  $\rho_r$  and  $\rho_n$ , hence for  $\rho_u = \rho_n - \rho_r$ , and the continuity of  $\rho_s$  in a neighborhood of a point  $s$  whose fiber is a reductive group.<sup>71</sup>

When  $G$  is already assumed commutative, 8.2 can be generalized as follows.

**Theorem 8.8.**—<sup>72</sup>Let  $G$  be a commutative  $S$ -prescheme in groups, flat and of finite presentation over  $S$ , with affine connected fibers. Let  $s \in S$ , and suppose that:

- a) if  $\bar{k}$  denotes an algebraic closure of  $\kappa(s)$ , then  $(G_{\bar{k}})_{\text{red}}$  is a torus;
- b) there is a generization  $t$  of  $s$  such that  $G_t$  is smooth over  $\kappa(t)$ .

Under these conditions, there exists an open neighborhood  $U$  of  $s$  such that  $G|_U$  is a torus.

(Observe moreover that, if one assumes only that  $G_t$  is affine, respectively connected, for every generization  $t$  of  $s$ , it follows easily that the same conclusion holds for  $t$  in an open neighborhood of  $s$ .)

*Proof of 8.8.* It suffices to prove that  $G_s$  is smooth over  $\kappa(s)$ . Indeed, since  $G$  is flat and of finite presentation over  $S$ , it then follows that  $G$  is smooth over  $S$  above a neighborhood of  $s$  (cf. 3.5), and we are then in the situation of 8.1.

To prove that  $G_s$  is smooth over  $\kappa(s)$ , the usual procedure reduces us to the case in which  $S$  is affine and noetherian. Choose a morphism  $S' \rightarrow S$ , where  $S'$  is the spectrum of a discrete valuation ring, sending the closed point to  $s$  and the generic point  $\eta$  to  $t$ . We are thereby reduced to the case in which  $S$  is itself the spectrum of a discrete valuation ring; we may moreover suppose that it is complete and has algebraically closed residue field, and that  $s$  and  $t = \eta$  are respectively the closed point and the generic point of  $S$ .

Thus  $G$  is flat, separated, and of finite type over  $S$ ; the generic fiber  $G_\eta$  is smooth and connected, and the special fiber  $G_0$  is such that

$$T_0 = (G_0)_{\text{red}}$$

is a torus. Let  $m > 0$  be an integer prime to the residual characteristic at  $s$ , and hence also to that at  $\eta$ . Then  $m \cdot \text{id}_G$  is étale on every fiber (VII<sub>A</sub>, 8.4), and is therefore étale because  $G$  is flat over  $S$  (SGA 1, I 5.9). Its kernel  ${}_mG$  is thus étale over  $S$ ; since  $G$  is separated<sup>\*</sup> and of finite type over  $S$ , the same holds for  ${}_mG$ . Since

$$({}_mG)_0 = m(T_0),$$

<sup>71</sup>Editors' note. Give a reference for these results?

<sup>72</sup>Editors' note. G. Prasad and J.-K. Yu generalized this result (subject to some additional hypotheses) without assuming  $G$  commutative, and with "torus" replaced by "reductive group" in hypothesis (a) and in the conclusion; cf. [PY06], Theorem 6.2.

<sup>\*</sup>By Raynaud's theorem, VI<sub>B</sub>, 5.3.

<sup>73</sup>Editors' note. See Editors' note (62) in 8.5.

its degree is  $m^r$ , where

$$r = \dim T_0 = \dim G_0 = \dim G.$$

It follows that the rank of  $({}_mG)_\eta = {}_m(G_\eta)$  is at least  $m^r$ , which already proves, using 8.6, that  $G_\eta$  is a torus of dimension  $r$ , since the two fibers of  ${}_mG$  have the same rank, and hence, as in 8.5, that  ${}_mG$  is finite over  $S$ .<sup>74</sup>

Since  $S$  is complete with algebraically closed residue field, the finite étale cover  ${}_mG$  splits completely. Thus, through every point of  ${}_mG_0$  there passes a section of  $G$  over  $S$ ; in particular, the set of points of  $G_0$  through which there passes a section of  $G$  over  $S$  is everywhere dense. We now use the following lemma.

**Lemma 8.9.**— *Let  $S$  be a locally noetherian regular prescheme of dimension 1, and let  $G$  be an  $S$ -prescheme in groups, flat and locally of finite type, such that  $G_\eta$  is smooth over  $\kappa(\eta)$  for every maximal point  $\eta$  of  $S$  (which implies that  $G$  is reduced). Suppose that the normalization  $X$  of  $G$  is finite over  $G$  (this is the case, by Nagata, when  $S$  is the spectrum of a complete discrete valuation ring; cf. EGA IV<sub>2</sub>, 7.7.4), and let  $G'$  be the open subset of  $X$  formed by the points at which  $X$  is smooth over  $S$ . With these notations:*

- a) *If the projection  $G' \rightarrow S$  is surjective, then  $G'$  carries a unique structure of an  $S$ -prescheme in groups for which the canonical morphism  $G' \rightarrow G$  is a group homomorphism.*
- b) *Suppose that, for every closed point  $s$  of  $S$ , the set of points of  $G'_s$  through which an étale quasi-section passes is dense in  $G'_s$  for the Zariski topology. Then  $X$  is regular, the hypotheses of (a) hold, and the map*

$$\Gamma(G'/S) \longrightarrow \Gamma(G/S)$$

*is bijective.*

In the case that concerns us, this lemma applies and gives a group homomorphism  $u : G' \rightarrow G$ , where  $G'$  is smooth over  $S$  and

$$u_\eta : G'_\eta \xrightarrow{\sim} G_\eta$$

is an isomorphism. After possibly replacing  $G'$  by an open subgroup, we may suppose that  $G'_0$  is connected. Since  $u_0 : G'_0 \rightarrow G_0$  induces a surjective morphism  $G'_0 \rightarrow T_0$ , where  $T_0$  is a torus of the same dimension as  $G'_0$ , it follows easily that  $G'_0$  is a torus (for example, by using 8.6, or by referring to *Bible*, §7.3, Theorem 3(a)). Consequently, by 8.2,  $G'$  is a torus. By IX 6.8,  $\ker u$  is then a subgroup of multiplicative type of  $G'$ ; since its generic fiber is the identity group, it is itself the identity group, and therefore  $u$  is a monomorphism. It follows from VIII 7.9 that  $u$  is an immersion. Since it is surjective and  $G$  is reduced,  $u$  is an isomorphism, completing the proof of 8.8.

It remains to prove 8.9. For part (a), the uniqueness of the group law on  $G'$  for which  $u : G' \rightarrow G$  is a group homomorphism is clear, since the group law on the generic fiber of  $G'$  is known (we may suppose  $S$  irreducible). For existence, we reduce easily to the case in which  $S$  is local, the spectrum of a discrete valuation ring  $A$ . By uniqueness, and because integral closure commutes with an étale extension of the base, we may make étale base extensions on  $S$ . This reduces us to the case in which  $A$  is “strictly local”, that is, henselian with separably closed residue field. The same reduction applies to part (b); under the hypothesis made there, “étale quasi-section” can now be replaced by “section”.

<sup>74</sup> *Editors' note.* Revise the preceding sentence.

Since  $G'$  is smooth over  $S$  and  $X$  is normal, the prescheme  $G' \times_S X$  is normal (being smooth over the normal prescheme  $X$ ). Thus the composite morphism

$$G' \times_S X \longrightarrow G \times_S G \longrightarrow G$$

factors as

$$p : G' \times_S X \longrightarrow X.$$

We shall prove that the restriction of this morphism to the open subset  $G' \times_S G'$  of  $G' \times_S X$  induces a morphism

$$G' \times_S G' \longrightarrow G'.$$

It is therefore enough to show that  $G'_0 \times_k G'_0$  maps into the open subset  $G'_0$  of  $X_0$ . It suffices to check that, for every  $k$ -valued point  $g'_0$  of  $G'_0$ , the morphism

$$h'_0 \longmapsto p_0(g'_0, h'_0) \quad (\times)$$

from  $G'_0$  to  $X_0$  takes its values in  $G'_0$ . Since  $G'$  is smooth over  $S$  and  $S$  is henselian, every such  $g'_0$  is induced by a section  $g'$  of  $G'$  over  $S$ . The morphism  $(\times)$  is then induced by the morphism

$$h' \longmapsto p(g', h')$$

from  $G'$  to  $X$ , which itself is obtained by transport of structure from the automorphism  $h \mapsto g \cdot h$  of  $G$ , namely left translation by the section  $g$  of  $G$  that is the image of  $g'$ . Thus  $h' \mapsto p(g', h')$  is itself an automorphism of  $X$ , and hence maps  $G'$  into  $G'$ . This proves the assertion.

It remains to prove that the composition law thereby obtained on  $G'$  is a group law. Associativity follows at once from associativity on the generic fiber (which is isomorphic to that of  $G$ ). Moreover, the inversion automorphism of the  $S$ -prescheme  $G$  induces an automorphism of  $X$ , which therefore preserves  $G'$  and induces an automorphism  $\sigma$  of  $G'$ . One verifies that  $\sigma$  has the properties of an inverse for the composition law on  $G'$ . Indeed, this again amounts to verifying the commutativity of certain diagrams involving fiber powers of  $G'$  over  $S$ ; since these are smooth over  $S$ , it suffices to check commutativity on the generic fiber, where it is clear. This proves part (a) of 8.9.

We prove part (b). Let  $Z'$  be the set of  $x \in X$  for which  $\mathcal{O}_{X,x}$  is not regular. It is closed by a theorem of Nagata (EGA IV<sub>2</sub>, 6.12.6), and is clearly contained in  $X_0$ . Let  $Z$  be its image in  $G$ , which is therefore a closed subset of  $G_0$ . Then  $Z$  is nowhere dense in  $G_0$ , that is, it contains no maximal point  $y$  of  $G_0$ . Indeed,  $G_0$  is defined in  $G$  by an equation  $t = 0$ , where  $t$  is a uniformizer of the discrete valuation ring  $A$ . By the Hauptidealsatz,  $\mathcal{O}_{G,y}$  has dimension 1. Hence, for every  $x \in X$  above  $y$ ,  $\mathcal{O}_{X,x}$  has dimension 1, and is therefore a discrete valuation ring, since  $X$  is normal and hence regular in codimension 1.

On the other hand, for every section  $g$  of  $G$  over  $S$ , the set  $Z'$  is plainly stable under the automorphism of  $X$  obtained by transport of structure from left translation by  $g$  on  $G$ . Thus  $Z$  is stable under the left translation on  $G_0$  defined by  $g_0$ . By hypothesis, the set of these  $g_0$  is dense in  $G_0^0$ . Since  $Z$  is stable under translation by these  $g_0$  and is a nowhere-dense closed subset, it follows at once that  $Z = \emptyset$ , whence  $Z' = \emptyset$ , so  $X$  is regular. Consequently  $X$  is smooth over  $S$  at every point through which a section passes. Every section of  $G$  over  $S$  lifts uniquely to a section of  $X$  over  $S$ , and hence to a section of  $G'$  over  $S$ . This applies in particular to the identity section, proving that the image of  $G'$  in  $S$  is all of  $S$ , that is, that the hypothesis of (a) holds. This completes the proof of 8.9, and hence of 8.8.

## 9. Addenda

75

### 0.9.1 Simplicial sets, topoi, groupoids, and topological spaces

**Notations 9.1.1.**— (1) Let  $E$  be a simplicial set. The following objects can be associated with it:

- (2) a topos  $T = E^\sim$ , obtained from the topoi naturally associated with the sets  $E_i$ , by the procedure described in [Del74], 6.3.1 (see also [Ill72], VI.5.2 and SGA 4, VI.7);
- (3) a groupoid  $G = \Pi(E)$ , whose objects are the elements of  $E_0$  (the “vertices”) and whose arrows are defined in [GZ67], II.7;
- (4) a topological space  $X = |E|$ , a cellular complex called the “geometric realization” (or “topological realization”) (cf. loc. cit., III.1).

Observe that a sheaf on  $T$  is nothing other than a simplicial set over  $E$ .

### 0.9.2 Locally constant sheaves; descent data

**Definitions 9.2.1.**— A locally constant sheaf on

- (1) a simplicial set  $E$  is a morphism of simplicial sets  $E' \rightarrow E$  such that, for every  $i \in \mathbb{N}$  and every  $e \in E_i$ , the face operators  $d$  induce isomorphisms

$$E'_e \xrightarrow{\sim} E'_{d(e)}$$

between the fibers (cf. [AM69], §10);

- (2) a topos  $T$  is an object  $F$  of  $T$  such that there exist an epimorphism  $U \rightarrow 1$  and an isomorphism

$$F \times U \simeq f^* I \times U,$$

where  $I$  is a set and  $f : T \rightarrow \mathbf{Ens}$  is the final morphism (cf. SGA 4, IX.2);

- (3) a groupoid  $G$  is a presheaf on  $G$ , that is, a contravariant functor from  $G$  to the category of sets (cf. [GZ67], Appendix I.1.2);
- (4) a topological space  $X$  is a sheaf of sets on  $X$  that is locally constant in the usual sense.

Finally:

- (5) a *descent datum* on a simplicial set  $E$  consists of a sheaf  $F$  on  $E_0^\sim$  (that is, a set-valued function on  $E_0$ ) and an isomorphism

$$\alpha : p_1^* F \xrightarrow{\sim} p_2^* F,$$

where  $p_1, p_2 : E_1^\sim \rightarrow E_0^\sim$  are the morphisms induced by the faces, satisfying the usual cocycle relation (cf. Exposé IV, 2.1<sup>1</sup> and below).

The morphisms between these five kinds of objects, as well as the associated inverse-image functors, are defined in the evident way.

<sup>75</sup> *Editors' note.* This additional section was written by Fabrice Orgogozo (following indications by Ofer Gabber).

<sup>1</sup>PP: I have replaced “FGA, *Technique de descente* I, 1.6” by “Exposé IV, 2.1”.

### 0.9.3 Some equivalences of categories

Let  $E$  be a simplicial set, and let  $T$ ,  $G$ , and  $X$  be, respectively, the associated topoi, groupoid, and topological space.

**Proposition 9.3.1.**— *The categories of locally constant sheaves on  $E$ ,  $T$ ,  $G$ , and  $X$ , as well as the category of descent data on  $E$ , are equivalent.*

*Sketch of proof.* Denote by (1) through (5) the categories of objects defined in the preceding paragraph.

- (1)  $\iff$  (5). This is a special case of [AM69], 10.6 (see also [GZ67], Appendix I.2.3, [Fri82], ?5.6, or [Ill72], VI.8.1.6). An equivalence of categories is given by the functor that associates with the object  $E' \rightarrow E$  the pair  $(E'_0, \alpha)$ , where  $E'_0$  is regarded as a sheaf on  $E_0$  and  $\alpha$  is the unique isomorphism

$$p_1^* E'_0 \xrightarrow{\sim} p_2^* E'_0$$

whose fiber at each  $y \in E_1$ , with images denoted  $x_1$  and  $x_2$  under the two projections, is given by the isomorphisms

$$(E'_0)_{x_1} \xleftarrow{\sim} (E'_1)_y \xrightarrow{\sim} (E'_0)_{x_2}.$$

- (5)  $\iff$  (3). This is evident: one of the two relations defining the morphisms in the groupoid  $G$  associated with  $E$  is a cocycle relation.

- (1)  $\iff$  (4). Cf. [GZ67], Appendix I.3.2.1.

- (1)  $\iff$  (2). One must show that an object over  $E$  is a locally constant sheaf in the simplicial sense if and only if it is one as a sheaf on  $T = E^\sim$ . This follows from the fact that the locally constant sheaves  $F_\bullet$  on  $T$  are precisely the cartesian sheaves, that is, those for which the arrows

$$([n] \rightarrow [m])^* F_m \longrightarrow F_n$$

are isomorphisms. This last assertion is a special case of a general fact about simplicial topoi, together with the fact that every sheaf on  $E_0^\sim$  is locally constant. (See also [Ill72], VI.8.1.6.)

Q.E.D.

For every group  $H$ , the preceding equivalences of categories induce equivalences between the categories of  $H$ -torsors. Here these are locally constant sheaves equipped with an action of  $H$ , whose fibers are isomorphic to  $H$  acting on itself by translation.

### 0.9.4 Fundamental groups and groupoids

It follows from the preceding equivalences of categories that, for every group  $H$  and every simplicial set  $E$ , the sets of isomorphism classes of  $H$ -torsors

$$H^1(E, H), \quad H^1(E^\sim, H), \quad \pi_0(\mathrm{Hom}(\Pi(E), BH)), \quad H^1(|E|, H)$$

are naturally in bijection. Recall that  $BH$  denotes the one-object groupoid associated with  $H$ , and that  $\pi_0$  is the functor that associates with a category the set of isomorphism classes of its objects.

Likewise, if  $e$  is a point of  $E$ , the preceding equivalences induce bijections between the sets of isomorphism classes of  $H$ -torsors trivialized over  $e$ , denoted respectively by

$$H^1(E \text{ rel } e, H), \quad H^1(E^\sim \text{ rel } e^\sim, H), \quad \mathrm{Hom}(\pi_1(\Pi(E), e), H), \quad H^1(|E| \text{ rel } |e|, H).$$

Recall that

$$\pi_1(\Pi(E), e) = \text{Isom}_{\Pi(E)}(e, e).$$

For variable  $H$ , these functors are represented, in the connected case, by a group denoted  $\pi_1(E, e)$ . The group  $\pi_1(E, e)$  is isomorphic to  $\pi_1(\Pi(E), e)$  and to  $\pi_1(|E|, |e|)$ , and hence also to the fundamental group of a simplicial set as defined by Kan (cf., for example, [May67], 16.1, or [Ill72], I.2.1.1). Recall also that  $H^1(E, H)$  is in turn isomorphic to the set  $H^1(\pi_1(E, e), H)$  of homomorphisms to  $H$  modulo conjugation, also denoted

$$\text{Hom ext}(\pi_1(E, e), H).$$

### 0.9.5 Cones

**Definitions 9.5.1.**— (1) Let  $f : E' \rightarrow E$  be a morphism of simplicial sets. Recall (cf., for example, [Del74], 6.3.1) that  $C(f)$  denotes the pointed simplicial set whose underlying set in degree  $n \geq 0$  is

$$E_n \amalg \left( \coprod_{i < n} E'_i \right) \amalg \star,$$

where  $\star$  is a singleton. We leave it to the reader to define the simplicial maps; the faces in degrees at most two are recalled below. (See also [GZ67], VI.2, for a pointed variant.) The category of locally constant sheaves on  $C(f)$  is equivalent to the category of locally constant sheaves on  $E$  equipped with a trivialization of the inverse image on  $E'$ . The simplicial set  $C(f)$  is naturally pointed by  $\star \rightarrow C(f)$ .

(2) Let  $f : T' \rightarrow T$  be a morphism of topoi. Denote by  $C(f)$  the topos whose objects are quintuples

$$(F, F', A, \alpha : f^*F \rightarrow F', \beta : A \rightarrow \Gamma(T', F')),$$

where  $F$  (respectively  $F'$ ) is an object of  $T$  (respectively  $T'$ ),  $A$  is a set, and  $\alpha$  (respectively  $\beta$ ) is a morphism in  $T'$  (respectively in **Ens**). (See also [Del80], 4.3.4 and [Ill72], III.4 for a variant of this construction.) The category of locally constant sheaves on  $C(f)$  is equivalent to the category of locally constant sheaves on  $T$  equipped with a trivialization of the inverse image on  $T'$ . The topos  $C(f)$  is naturally pointed by the fiber functor that sends the quintuple above to the set  $A$ .

(3) Let  $f : G' \rightarrow G$  be a morphism of groupoids. Denote by  $C(f)$  the colimit of the diagram

$$G \longleftarrow G' \longrightarrow B1,$$

where  $B1$  is the one-object category (one object, one arrow). The category of locally constant sheaves on  $C(f)$  is equivalent to the category of locally constant sheaves on  $G$  equipped with a trivialization of the inverse image on  $G'$ .

(4) Let  $f : X' \rightarrow X$  be a morphism of topological spaces. Denote by  $C(f)$  the colimit of the diagram

$$\star \longleftarrow X' \times \{0\} \longrightarrow X' \times [0, 1] \longleftarrow X' \times \{1\} \longrightarrow X.$$

The category of locally constant sheaves on  $C(f)$  is equivalent to the category of locally constant sheaves on  $X$  equipped with a trivialization of the inverse image on  $X'$ . The topological space  $C(f)$  is naturally pointed by  $\star \rightarrow C(f)$ .

**9.5.2. Descent data on the cone of a simplicial map**

Let  $f : E' \rightarrow E$  be a morphism of simplicial sets, and let  $C(f)$  be its cone (cf. above, (1)). Use  $p$  (respectively  $q, r$ ) to denote the face maps of  $E'$  (respectively  $E, C(f)$ ). Before stating the proposition below, we describe the faces  $r$  in degrees at most two in terms of  $p$  and  $q$ . By convention,  $p_{ij}$  (respectively  $p_i$ ) is the face map

$$E'_2 = E'_{\{1,2,3\}} \longrightarrow E'_1 = E'_{\{1,2\}} \quad \left( \text{respectively } E'_1 = E'_{\{1,2\}} \longrightarrow E'_0 = E'_{\{1\}} \right)$$

corresponding to the increasing map

$$\{1, 2\} \longrightarrow \{1, 2, 3\} \quad (\text{respectively } \{1\} \longrightarrow \{1, 2\})$$

whose image is  $\{i, j\}$  (respectively  $\{i\}$ ).<sup>2</sup> The same convention is adopted for the faces of  $E$  and  $C(f)$ .

The morphism

$$r_1 : C(f)_1 = E_1 \amalg E'_0 \amalg \star \longrightarrow C(f)_0 = E_0 \amalg \star$$

(respectively  $r_2$ ) is:

- $q_1 : E_1 \rightarrow E_0$  (respectively  $q_2 : E_1 \rightarrow E_0$ ) on  $E_1$ ;
- $E'_0 \rightarrow \star$  (respectively  $f_0 : E'_0 \rightarrow E_0$ ) on  $E'_0$ ;
- $\star \rightarrow \star$  on  $\star$ .

Likewise, the morphism

$$r_{21} : C(f)_2 = E_2 \amalg E'_1 \amalg E'_0 \amalg \star \longrightarrow C(f)_1 = E_1 \amalg E'_0 \amalg \star$$

(respectively  $r_{32}, r_{31}$ ) is:

- $q_{21} : E_2 \rightarrow E_1$  (respectively  $q_{32}, q_{31}$ ) on  $E_2$ ;
- $p_1 : E'_1 \rightarrow E'_0$  (respectively  $f_1 : E'_1 \rightarrow E_1, p_2 : E'_1 \rightarrow E'_0$ ) on  $E'_1$ ;
- $E'_0 \rightarrow \star$  (respectively  $\text{Id} : E'_0 \rightarrow E'_0, \text{Id} : E'_0 \rightarrow E'_0$ ) on  $E'_0$ .

Let  $H$  be a descent datum on  $C(f)$ , that is, a sheaf  $H_0$  on

$$C(f)_0 = E_0 \amalg \star$$

equipped with an isomorphism

$$\gamma : r_1^* H_0 \xrightarrow{\sim} r_2^* H_0$$

between its inverse images on

$$C(f)_1 = E_1 \amalg E'_0 \amalg \star,$$

satisfying the cocycle relation

$$r_{31}^* \gamma = r_{32}^* \gamma \circ r_{21}^* \gamma.$$

The restriction of  $\gamma$  to  $E_1$  (respectively  $E'_0$ ) is a descent datum

$$\beta : q_1^* G_0 \xrightarrow{\sim} q_2^* G_0,$$

---

<sup>2</sup>This convention departs from the present simplicial convention but is closer to the notation used by Grothendieck in his theory of descent.

where  $G_0$  is the restriction of  $H_0$  to  $E_0$  (respectively, a trivialization

$$t : H_0(\star) \xrightarrow{\sim} f^*G_0 =: F_0,$$

where  $H_0(\star)$  is the constant sheaf with stalk  $H_0(\star)$ ). The restriction of the cocycle relation satisfied by  $\gamma$  to  $E_2$  (respectively  $E'_1$ ) is the cocycle relation for  $\beta$  (respectively,

$$p_2^*(t) = f_1^*(\beta) \circ p_1^*(t).$$

This latter relation says that the trivialization  $t$  is compatible with the descent datum  $f_1^*(\beta)$  induced by  $\beta$  on  $F_0$ . It follows that:

**Proposition 9.5.3.**— *Let  $f : E' \rightarrow E$  be a morphism of simplicial sets, and let  $C(f)$  be its cone, pointed by  $\star$ . The category of descent data on  $C(f)$  trivialized over  $\star$  is equivalent to the category of descent data on  $E$  equipped with a trivialization of the descent datum induced on  $E'$ .*

### 0.9.6 Representability of the functor $\bar{\pi}^1(S'/S, \xi; -)$

The notation is that of page 90, and from now on we suppose that the connected components of

$$S_0 = S' \quad \text{and} \quad S_1 = S' \times_S S'$$

are open. Under this hypothesis we have the following explicit combinatorial description, which follows immediately from the preceding recall.

**Proposition 9.6.1.**— *For every group  $G$ , the set  $\bar{\pi}^1(S'/S, \xi; G)$  of isomorphism classes of descent data equipped with a trivialization over  $\xi$  is naturally in bijection with the relative nonabelian cohomology set*

$$H^1(K \text{ rel } k, G) := H^1(\tilde{K} \text{ rel } \star, G).$$

It follows that this functor is representable by a group, denoted  $\pi_1(K, k)$ , the *relative fundamental group*, as soon as the simplicial set

$$\tilde{K} = C(k \rightarrow K)$$

is connected. One verifies immediately that this holds if and only if the map  $\pi_0(k) \rightarrow \pi_0(K)$  is surjective. It is enough, in particular, that the simplicial set  $K$  be connected. As indicated in the text, this is the case when the scheme  $S$  is connected and the morphism  $S' \rightarrow S$  is universally submersive.

### 0.9.7 Contractibility of $k$ ; counterexample to the injectivity of $k_\bullet \rightarrow K_\bullet$ .

The following two propositions make Editors' note (53) more precise.

**Proposition 9.7.1.**— *Let  $\kappa$  be an algebraically closed field, and let  $X$  be a connected  $\kappa$ -scheme. For every integer  $i \geq 0$ , let  $X_i$  denote the  $(i+1)$ -st fiber power of  $X$  over  $\kappa$ . For every integer  $i \geq 0$ , the canonical map*

$$\pi_0(X_i) \longrightarrow \pi_0(X)^{i+1}$$

*is a bijection. In particular, the simplicial set  $\pi_0(X_\bullet)$  is contractible.*

It suffices to prove the following lemma, which follows by passage to the limit from the Künneth formula. [Expand?]



**Lemma 9.7.2.**— *Let  $\kappa$  be an algebraically closed field, and let  $X$  and  $Y$  be two connected  $\kappa$ -schemes. Then the fiber product  $X \times_{\kappa} Y$  is connected.*

**Proposition 9.7.3.**— *Let  $X$  be a connected scheme, let  $X' \rightarrow X$  be a finite étale morphism, and let  $\bar{x}$  be a geometric point of  $X$  localized at a point  $x$ . The canonical morphism*

$$\pi_0(X'_{\bar{x}}) \longrightarrow \pi_0(X')$$

*is surjective.*

Indeed, the image of every connected component of  $X'$  is open and closed in  $X$ , and therefore meets the fiber  $X'_x$ . Thus

$$\pi_0(X'_x) \longrightarrow \pi_0(X')$$

is surjective, as is

$$\pi_0(X'_{\bar{x}}) \longrightarrow \pi_0(X'_x).$$

## Bibliography

76

- [BAC] N. Bourbaki, *Algèbre commutative*, Chap. I–IV, Hermann, 1961; Masson, 1985; Springer-Verlag, 2006.
- [BEns] N. Bourbaki, *Théorie des ensembles*, Chap. I–IV, Hermann, 1970.
- [Pes66] C. Peskine, *Une généralisation du “Main Theorem” de Zariski*, Bull. Sci. Math. **90** (1966), 119–127.
- [PY06] G. Prasad and J.-K. Yu, *On quasi-reductive group schemes*, J. Alg. Geom. **15** (2006), 507–549.
- [Ray70] M. Raynaud, *Anneaux locaux henséliens*, Lect. Notes Maths. **169**, Springer-Verlag, 1970.
- [AM69] M. Artin and B. Mazur, *Étale homotopy*, Lect. Notes Maths. **100**, Springer, 1969.
- [Del74] P. Deligne, *Théorie de Hodge III*, Publ. Math. IHÉS **44** (1974), 5–77.
- [Del80] P. Deligne, *La conjecture de Weil II*, Publ. Math. IHÉS **52** (1980), 137–252.
- [Fri82] E. M. Friedlander, *Étale homotopy of simplicial schemes*, Princeton Univ. Press, 1982.
- [GZ67] P. Gabriel and M. Zisman, *Calculus of fractions and homotopy theory*, Springer, 1967.
- [Ill72] L. Illusie, *Complexe cotangent et déformations I & II*, Lect. Notes Maths. **239** & **283**, Springer, 1971 & 1972.
- [Kan58] D. Kan, *A combinatorial definition of homotopy groups*, Ann. of Math. **67** (1958), 282–312.
- [May67] J. P. May, *Simplicial objects in algebraic topology*, Univ. of Chicago Press, 1967.

---

<sup>76</sup>*Editors’ note.* Additional references cited in this Exposé; the references [AM69] and following concern the Addenda.

# Exposé XI

## Representability Criteria

### Applications to

### Multiplicative-Type Subgroups

### of Affine Group Schemes

by A. Grothendieck<sup>0</sup>

#### 0. Introduction

As we have already seen in Exposé X, nos. 4 and 5, the representability of certain functors—for example, functors of the form  $\mathrm{Hom}_S(X, Y)$  and various related constructions—plays an important role in many questions concerning group preschemes.

Among the results that are particularly useful in this theory, besides the representability of quotients studied in Exposés V and VI<sub>B</sub> and in Exposé VIII, no. 5, let us mention the representability of functors of the form

$$\prod_{X/S} Y/X,$$

where  $Y$  is a subobject of  $X$ . This was studied in a very elementary case in Exposé VIII, no. 6; variants will be given in no. 6 of the present exposé. These results provide the representability of various centralizers, normalizers, and transporters.

Less elementary representability criteria, using results that will appear in EGA VI, are indicated in 6.12 and in Exposés XV and XVI. There we shall give a criterion for the representability of quotients  $G/H$  in cases not covered by the preceding exposés; this criterion was not developed in the oral lectures.

Our principal purpose in the present exposé is to prove Theorems 4.1 and 4.2, which provide a typical example of a nonprojective construction technique, close to the technique that will be developed in EGA VI. Since the oral lecture and the drafting of the present text, it has moreover become apparent that the affine hypotheses in 4.1 and 4.2 can be removed to a large extent (cf. Exposé XV), and that the main part of the theory developed in the following exposé can dispense with 4.1 and 4.2. Finally, it might be interesting to prove analogues of these results for a general reductive group prescheme (for example, a semisimple one) in place of a group of multiplicative type; in that case 4.1 and 4.2 would doubtless be the key result in the proof.

---

<sup>0</sup>Editor's note: xy version of December 5, 2008.

## 1. Review of Smooth, Étale, and Unramified Morphisms

The reader is referred to EGA IV, §§17 and 18, and, pending its publication, to SGA 1, Exposés I, II, and III. Certain noetherian hypotheses used there, which are inconvenient in applications, should however be replaced by finite-presentation hypotheses.

**Definition 1.1.** — *Let  $S$  be a prescheme and let*

$$F : (\mathbf{Sch})^\circ / S \longrightarrow (\mathbf{Ens})$$

*be a functor. The functor  $F$  is said to be formally smooth (respectively, formally unramified<sup>1</sup>, respectively, formally étale) if, for every  $S$ -prescheme  $S'$  that is affine in the absolute sense and every subscheme  $S'_0 \subset S'$  defined by a nilpotent ideal  $\mathcal{J}$ , the map*

$$F(S') \longrightarrow F(S'_0)$$

*is surjective (respectively, injective, respectively, bijective).*

*An  $S$ -prescheme  $X$  is said to be formally smooth over  $S$  (respectively, formally unramified over  $S$ , respectively, formally étale over  $S$ ) if the corresponding functor has that property. It is said to be smooth over  $S$  (respectively, unramified over  $S$ , respectively, étale over  $S$ ) if it satisfies the preceding condition and is, in addition, locally of finite presentation over  $S$ .*

The usefulness of these definitions for  $X$  lies, on the one hand, in their remarkably simple expression in terms of the functor represented by  $X$  (and in practice  $X$  is often given as the object over  $S$  representing an explicitly described functor), and, on the other hand, in the remarkable local structural properties that they express. We recall those properties in the following statements; for their proofs, see the references just cited.

**Proposition 1.2.** — *Let  $X$  be a prescheme locally of finite presentation over  $S$ . Then:*

- (i)  *$X$  is smooth over  $S$  if and only if it is flat over  $S$  and its geometric fibers*

$$X \otimes_S \mathrm{Spec} \kappa(s)$$

*are regular schemes. More generally,  $X$  is smooth over  $S$  in a neighborhood of  $x \in X$  (one then says that  $X$  is smooth over  $S$  at  $x$ ) if and only if  $X$  is flat over  $S$  at  $x$  and  $X_s$  is smooth over  $\kappa(s)$  at  $x$ ; that is,*

$$X_s \otimes_{\kappa(s)} \overline{\kappa(s)}$$

*is regular at the points (or merely at one point) lying over  $x$ .*

- (ii) *Suppose that  $S$ , and hence  $X$ , is locally noetherian. Let  $x \in X$ , and let  $s \in S$  be its image. The smoothness of  $X$  over  $S$  at  $x$  can then be detected from the local homomorphism*

$$A = \mathcal{O}_{S,s} \longrightarrow B = \mathcal{O}_{X,x}$$

*of noetherian local rings (or even from the local homomorphism  $\hat{A} \rightarrow \hat{B}$  of their completions), by the following characteristic property:  $B$  is flat over  $A$ , and  $B \otimes_A k$  is geometrically regular over  $k$ , the residue field of  $A$ . Equivalently, for every finite extension  $k'$  of  $k$ ,*

$$(B \otimes_A k) \otimes_k k' = B \otimes_A k'$$

<sup>1</sup>The source notes that one would now rather say “net” in place of “unramified.”

is a regular semilocal ring. When the residue-field extension  $k(B)/k(A)$  is trivial, these conditions are also equivalent to the following one:  $\widehat{B}$  is isomorphic, as an  $\widehat{A}$ -algebra, to an algebra of formal power series

$$\widehat{A}[[t_1, \dots, t_n]].$$

Thus, from the “formal” point of view, the structure of  $X$  over  $S$  is that of the standard affine space  $S[t_1, \dots, t_n]$  over  $S$ .

**Proposition 1.3.** — *Let  $X$  be a prescheme locally of finite presentation over  $S$ . Then:*

(i) *The following conditions are equivalent:*

- (a)  *$X$  is unramified over  $S$ .*
- (b) *The diagonal morphism  $X \rightarrow X \times_S X$  is an open immersion.*
- (c)  $\Omega_{X/S}^1 = 0$ .
- (d) *The geometric fibers of  $X/S$  are discrete and reduced; that is, they are isomorphic to disjoint sums of copies of the base field.*
- (e) *For every  $s \in S$ , the fiber*

$$X \otimes_S \operatorname{Spec} \kappa(s) = X_s$$

*is unramified over  $\kappa(s)$ , or equivalently is isomorphic to a disjoint sum of spectra of finite separable extensions of  $\kappa(s)$ .*

(ii) *There are analogous pointwise conditions for  $X$  to be unramified over  $S$  at a given point  $x$ , that is, in a neighborhood of that point. For example, it is necessary and sufficient that  $\mathcal{O}_{X_s, x}$  be a finite separable extension of  $\kappa(s)$ . In terms of the local homomorphism*

$$A = \mathcal{O}_{S, s} \longrightarrow B = \mathcal{O}_{X, x},$$

*this says that  $B \otimes_A k$  is a finite separable extension of the residue field  $k$  of  $A$ ; equivalently,*

$$\mathfrak{r}(A)B = \mathfrak{r}(B) \quad \text{and} \quad k(B)/k(A) \text{ is finite separable.}$$

*When  $A$ , and hence  $B$ , is noetherian and  $k(A) \xrightarrow{\sim} k(B)$ , this also means that  $\widehat{A} \rightarrow \widehat{B}$  is surjective.*

Thus, from the “formal” point of view, saying that  $X$  is unramified over  $S$  means that  $X$  is essentially a subprescheme of  $S$ . Observe also that conditions (d) and (e) are expressed solely in terms of the fibers of  $X/S$ .

**Proposition 1.4.** — *Let  $X$  be a prescheme locally of finite presentation over  $S$ . Then  $X$  is étale over  $S$  if and only if it is smooth and unramified over  $S$  (which is immediate from the definition), so the criteria in 1.2 and 1.3 apply. In particular:*

- (i)  *$X$  is étale over  $S$  if and only if it is flat and unramified over  $S$ . There is an analogous local criterion for  $X$  to be étale over  $S$  at a given point  $x$ .*

- (ii) Suppose in addition that  $S$ , and hence  $X$ , is locally noetherian. Whether  $X$  is étale over  $S$  at  $x$  can then be detected from the local homomorphism

$$A = \mathcal{O}_{S,s} \longrightarrow B = \mathcal{O}_{X,x}$$

<sup>2</sup> (and even from  $\hat{A} \rightarrow \hat{B}$ ) by the following characteristic property:  $B$  is flat over  $A$ , and  $B \otimes_A k$  is a finite separable extension of  $k$ , the residue field<sup>3</sup> of  $A$ . When  $k(A) \xrightarrow{\sim} k(B)$ , this condition simply says that  $\hat{A} \rightarrow \hat{B}$  is an isomorphism.

Thus, from the “formal” point of view, saying that  $X$  is étale over  $S$  simply means that  $X$  is locally isomorphic to  $S$ .

**Remark 1.5.** — When  $X$  is locally of finite presentation over  $S$  and a point  $x \in X$  is given, the fact that  $X$  is smooth (respectively, unramified, respectively, étale) over  $S$  at  $x$ , that is, in a neighborhood of  $x$ , can also be detected from the functor

$$X : (\mathbf{Sch})^\circ / S \longrightarrow (\mathbf{Ens})$$

by the following property. For every  $S$ -scheme  $S'$  that is the spectrum of a local ring, every subscheme  $S'_0 \subset S'$  defined by a nilpotent ideal, and every  $S$ -morphism  $u_0 : S'_0 \rightarrow X$  sending the closed point  $s'$  of  $S'$  to  $x$ , there exists at least one (respectively, at most one, respectively, exactly one)  $S$ -morphism  $u : S' \rightarrow X$  extending  $u_0$ .

This statement shows in particular that, in Definition 1.1, one may restrict to local schemes  $S'$ , provided the functor under consideration is representable by a prescheme locally of finite presentation over  $S$ . Moreover, when  $S$  is locally noetherian, one may even restrict the preceding pointwise criterion to local artinian schemes  $S'$ , and the same restriction may therefore be made in Definition 1.1, provided that  $S$  is locally noetherian and that the functor in question is represented by a prescheme locally of finite type over  $S$ .

**Remark 1.6.** — Of course, given a morphism  $f : X \rightarrow Y$  of preschemes, one says that  $f$  is smooth (respectively, unramified, respectively, étale) if  $f$  makes  $X$  a smooth (respectively, unramified, respectively, étale)  $Y$ -prescheme. When  $X$  and  $Y$  are  $S$ -preschemes and  $f$  is an  $S$ -morphism of finite presentation, these properties of  $f$  can be expressed immediately in terms of the morphism of functors

$$F, G : (\mathbf{Sch})^\circ / S \longrightarrow (\mathbf{Ens})$$

defined by  $f$ . Namely,  $f$  is smooth (respectively, unramified, respectively, étale) if, for every  $S$ -prescheme  $S'$  that is affine in the absolute sense and every subscheme  $S'_0 \subset S'$  defined by a nilpotent ideal  $\mathcal{J}$ , the map

$$F(S') \longrightarrow F(S'_0) \times_{G(S'_0)} G(S')$$

deduced from the commutative square

$$\begin{array}{ccc} F(S') & \longrightarrow & G(S') \\ \downarrow & & \downarrow \\ F(S'_0) & \longrightarrow & G(S'_0) \end{array}$$

<sup>2</sup>The source prints  $\mathcal{O}_{X,s}$ ; the local ring at the point of  $X$  is  $\mathcal{O}_{X,x}$ .

<sup>3</sup>Editor’s note: “residue extension” was replaced by “residue field.”

is surjective (respectively, injective, respectively, bijective). Equivalently, for every commutative square of morphisms of functors over  $S$ , where  $S'$  and  $S'_0$  are as above and  $i : S'_0 \rightarrow S'$  is the canonical immersion,

$$(Q) \quad \begin{array}{ccc} S'_0 & \xrightarrow{i} & S' \\ u_0 \downarrow & & \downarrow v \\ F & \xrightarrow{f} & G, \end{array}$$

there exists at least one (respectively, at most one, respectively, exactly one) morphism

$$\begin{array}{ccc} & & S' \\ & \swarrow u & \\ F & & \end{array}$$

that makes the two corresponding triangles commute:

$$u_0 = u \circ i \quad \text{and} \quad v = f \circ u.$$

This property for a homomorphism of functors  $f : F \rightarrow G$  over  $S$  retains its meaning even when the functors under consideration are not representable. If it holds, the homomorphism  $f : F \rightarrow G$  is said to be *formally smooth* (respectively, *formally unramified*, respectively, *formally étale*). Notice that this property depends only on the homomorphism of the underlying functors

$$(\mathbf{Sch})^\circ \longrightarrow (\mathbf{Ens})$$

defined by  $F$  and  $G$  (cf. Exposé I, 1.4.1), and not on the structure morphisms  $F \rightarrow S$  and  $G \rightarrow S$ . An equivalent formulation of the preceding definition, somewhat easier to use in applications, is the following: a morphism of functors  $F \rightarrow G$  over  $S$  is formally smooth (respectively, formally unramified, respectively, formally étale) if, for every  $S'$  over  $S$  and every morphism  $S' \rightarrow G$ , the functor

$$F' = F \times_G S'$$

over  $S'$  is formally smooth (respectively, formally unramified, respectively, formally étale).

**Remark 1.7.** — Under the hypotheses of 1.6, suppose that we are given a “point of  $F$  with values in a field  $k$ ,” that is, an element  $x \in F(\text{Spec } k)$ , or equivalently a morphism

$$x : \text{Spec } k \longrightarrow F.$$

We likewise say that  $f$  is *formally smooth* (respectively, *formally unramified*, respectively, *formally étale*) at  $x$  if the condition in the preceding definition is satisfied whenever  $S'$  is a local scheme and the morphism  $S'_0 \rightarrow F$  in diagram (Q) above is “compatible with the marked points” in the following sense. Let  $k'$  denote the residue field of  $S'$  and  $S'_0$  at their closed point. Then the diagram

$$\begin{array}{ccc} \text{Spec}(k') & & \text{Spec}(k) \\ j \downarrow & & \downarrow x \\ S'_0 & \xrightarrow{u_0} & F \end{array}$$

in which  $j : \operatorname{Spec}(k') \rightarrow S'_0$  is the canonical immersion, must admit a completion to a commutative diagram

$$\begin{array}{ccc}
 & \operatorname{Spec}(k'') & \\
 \swarrow & & \searrow \\
 \operatorname{Spec}(k') & & \operatorname{Spec}(k) \\
 \downarrow & & \downarrow \\
 S'_0 & \xrightarrow{\quad} & F,
 \end{array}$$

where  $k''$  is a field.<sup>4</sup>

When  $F$  and  $G$  are represented by  $S$ -preschemes  $X$  and  $Y$ , and the  $S$ -morphism  $f : X \rightarrow Y$  is locally of finite presentation, this condition says precisely, by 1.5, that  $f : X \rightarrow Y$  is smooth at the point of  $X$  that is the image of  $\operatorname{Spec}(k)$  under  $x : \operatorname{Spec}(k) \rightarrow X$ .

**Remark 1.8.** — When the condition in the preceding definition is satisfied after restricting to local artinian schemes  $S'$ , we say that  $F \rightarrow G$  is *infinitesimally smooth* (respectively, *infinitesimally unramified*, respectively, *infinitesimally étale*) at  $x$ . We say that  $F \rightarrow G$  is infinitesimally smooth (respectively, infinitesimally unramified, respectively, infinitesimally étale) if it has the corresponding property at every point  $x$ ; equivalently, if the condition considered in 1.6 is satisfied whenever  $S'$  is local artinian.

This variant of the preceding notions is technically useful. Since it is weaker, it is often easier to verify, while it is nevertheless frequently sufficient to imply the stronger condition; this is the case, for example, when  $F \rightarrow G$  is a morphism locally of finite presentation and  $G$  is represented by a locally noetherian prescheme.

**Proposition 1.9.** — *Let  $f : X \rightarrow S$  be a smooth morphism of preschemes. Then  $\Omega_{X/S}^1$  is a locally free module of finite type on  $X$ , and its rank at a point  $x \in X$  is equal to the dimension of the fiber  $X_s$ , where  $s = f(x)$ , in a neighborhood of  $x$ .*

This dimension is called the *relative dimension of  $X$  over  $S$  at  $x$* . Notice that it is zero (when  $f$  is smooth at  $x$ ) if and only if  $f$  is étale at  $x$ . It can also be computed directly in terms of the functor  $F$  represented by  $X$ , as follows. Let  $\xi$  be a point of  $F$  with values in a field  $k$ , “localized at  $x$ ,” that is, a morphism

$$\operatorname{Spec}(k) \longrightarrow F = X$$

whose image is  $x$ . Consider the algebra

$$D(k) = k[t]/(t^2)$$

of dual numbers over  $k$ , regarded as an  $S$ -prescheme, and the map

$$F(D(k)) \longrightarrow F(k)$$

induced by the augmentation  $D(k) \rightarrow k$ . Finally, let  $F(D(k), \xi)$  be the inverse image of  $\xi \in F(k)$  under this map. This set is naturally endowed with the structure of a vector space over  $k$ . In fact, it is the vector-space dual of

$$\Omega_{X/S}^1 \otimes_{\mathcal{O}_{X,x}} k,$$

<sup>4</sup>The source says that  $k''$  is “the spectrum of a field”; the displayed object  $\operatorname{Spec}(k'')$  makes clear that  $k''$  itself is the field.

and its dimension is the relative dimension of  $X$  over  $S$  at  $x$ .

To describe the vector-space law on  $F(D(k), \xi)$  explicitly, it is useful to introduce, more generally and as in Exposé II, for every vector space  $V$  over  $k$ , the algebra

$$D_k(V) = k + V,$$

where  $V$  is an ideal of square zero, and to regard  $F(D_k(V), \xi)$ , the inverse image of  $\xi$  under

$$F(D_k(V)) \longrightarrow F(k),$$

as a covariant functor of  $V$  with values in **(Ens)**. It is then enough that this functor commute with products of two factors. This means that  $F$  carries certain amalgamated sums of a very special kind to fiber products; compare Exposé II. This condition always holds when  $F$  is representable. It follows that the sets  $F(D_k(V), \xi)$ , and in particular  $F(D(k), \xi)$ , are endowed with vector-space structures over  $k$ . Thus one can define the relative dimension of  $F$  over  $X$  at the “point”  $\xi$  under conditions appreciably more general than the representability of  $F$ .

In the present Exposé, the fact that certain functors to be described explicitly are representable by preschemes smooth over  $S$  will be used chiefly through the following result. It will be our technical means of passing from constructions over the completion of the local ring  $A$  of a point  $s$  of a noetherian scheme  $S$  to a local ring  $A'$  over  $A$  that is étale over  $A$ , and hence, in particular, of passing from  $s$  to neighboring points.

**Proposition 1.10.** — *Let  $f : X \rightarrow S$  be a smooth morphism of preschemes, let  $s \in S$ , and let  $x \in X$  lie over  $s$ , with  $\kappa(x)$  a finite separable extension of  $\kappa(s)$ . Then there exists a subscheme  $S' \subset X$ , étale over  $S$ , passing through  $x$ .<sup>5</sup> Thus one can find an étale morphism  $S' \rightarrow S$ , a point  $s' \in S'$  above  $s$  whose residue-field extension is  $\kappa(x)/\kappa(s)$ , and an  $S$ -morphism  $S' \rightarrow X$  carrying  $s'$  to  $x$ .*

To construct  $S'$ , simply choose sections  $g_1, \dots, g_n$  of  $\mathcal{O}_X$  on a neighborhood of  $x$  whose images at  $x$  form a regular system of parameters for the local ring of the fiber  $X_s$  at  $x$ . The subscheme  $S'$  defined by the  $g_i$  is then étale over  $S$  at  $x$ , and, after shrinking  $S'$ , it is étale over  $S$ .

We shall use 1.10 when  $\kappa(x) = \kappa(s)$ , that is, when  $x$  is rational over  $\kappa(s)$ , or equivalently when  $x$  may be regarded as a section of  $X_s$  over  $\text{Spec } \kappa(s)$ . In that case, 1.10 has the character of an extension theorem for sections, after an étale extension of the base. It takes a particularly simple form in the following special case.

**Corollary 1.11.** — *Under the hypotheses of 1.10, suppose that  $S$  is the spectrum of a henselian local ring and that  $\kappa(x) = \kappa(s)$ . Then there exists a section of  $X$  over  $S$  passing through  $x$ . It is uniquely determined if  $X$  is in fact étale over  $S$  at  $x$ .*

Indeed, since  $S$  is henselian, under the hypotheses of 1.10 the prescheme  $S'$  contains an open subscheme that is finite over  $S$  and whose fiber at  $s$  is reduced to  $x$ . Since this open subscheme is étale over  $S$ , it is isomorphic to  $S$ , which proves the assertion. Notice that when  $S$  is the spectrum of a complete local ring, 1.10 or 1.11 is more or less equivalent to the classical “Hensel’s lemma,” and one sometimes refers to the result by that name.

<sup>5</sup>The French re-edition prints “a subscheme  $S'$  of  $S$ , étale over  $S$ , passing through  $x$ .” Since  $x \in X$ , and both the equivalent formulation and the construction immediately below use  $S' \rightarrow X$  and sections of  $\mathcal{O}_X$ , the intended reading is  $S' \subset X$ .



## 2. Examples of Formally Smooth Functors from the Theory of Groups of Multiplicative Type

We shall interpret, in the language introduced in the preceding number, the results stated in Exposé IX, no. 3, concerning infinitesimal extensions of a homomorphism from a group of multiplicative type. Those results are consequences of the vanishing of the Hochschild cohomology of such a group, established in Exposé I.

**Proposition 2.1.** — *Let  $S$  be a prescheme, let  $H$  be a group of multiplicative type over  $S$ , and let  $G$  be an  $S$ -group prescheme smooth over  $S$ . Consider the functor over  $S$*

$$M_H = \mathrm{Hom}_{S\text{-gr}}(H, G).$$

*Its value at an  $S'$  over  $S$  is the set of homomorphisms of  $S'$ -groups from  $H_{S'}$  to  $G_{S'}$ . This functor is formally smooth over  $S$ .*

Cf. Exposé IX, 3.6. More generally:

**Corollary 2.2.** — *Let  $S$  and  $G$  be as above, and consider a homomorphism  $u : H_1 \rightarrow H_2$  of group schemes of multiplicative type over  $S$ . With the preceding notation this induces a morphism of functors over  $S$*

$$M_{H_2} \longrightarrow M_{H_1}, \quad M_{H_i} = \mathrm{Hom}_{S\text{-gr}}(H_i, G),$$

*given by  $w \mapsto w \circ u$ . This morphism is formally smooth.*

Indeed, by the definitions in 1.6, this is equivalent to the following assertion. Suppose that  $S$  is affine and that  $S_0 \subset S$  is a subscheme defined by a nilpotent ideal. Let

$$v : H_1 \longrightarrow G$$

be a homomorphism of  $S$ -groups, and let

$$w_0 : (H_2)_{S_0} \longrightarrow G_{S_0}$$

be a homomorphism of  $S_0$ -groups such that

$$w_0 u_{S_0} = v_{S_0}.$$

Then there exists a homomorphism of  $S$ -groups

$$w : H_2 \longrightarrow G$$

extending  $w_0$  and satisfying  $wu = v$ .

To see this, first extend  $w_0$  to a homomorphism of  $S$ -groups

$$w' : H_2 \longrightarrow G,$$

which is possible by 2.1. Set

$$v' = w'u : H_1 \longrightarrow G.$$

By the hypothesis on  $w_0$ , one has  $v'_{S_0} = v_{S_0}$ . Consequently, by Exposé VIII, 3.6, there exists an element  $g \in G(S)$  whose image in  $G(S_0)$  is the identity element and such that

$$v = \mathrm{int}(g)v'.$$

It follows that

$$v = (\mathrm{int}(g)w')u,$$

and it is therefore enough to take

$$w = \text{int}(g)w'.$$

**Corollary 2.3.** — *With the notation of 2.1, regard  $M_H = M$  as a functor with operator group  $G$ , where  $G$  acts by*

$$(v, g) \mapsto \text{int}(g) \circ v.$$

*Then the corresponding morphism*

$$R : G \times_S M \longrightarrow M \times_S M,$$

*defined by*

$$(g, v) \mapsto (\text{int}(g) \circ v, v),$$

*is formally smooth.*

After a base change  $S' \rightarrow S$ , this is equivalent to the following assertion.

**Corollary 2.4.** — *With the notation of 2.1, let  $v_1, v_2 : H \rightarrow G$  be two morphisms of  $S$ -groups, and let  $\text{Transp}(v_1, v_2)$  be the subfunctor of  $G$  consisting of the  $g$  such that*

$$\text{int}(g) \circ v_1 = v_2.$$

*Then this functor is formally smooth over  $S$ . In particular, if  $v_1 = v_2 = v$ , the functor  $\text{Centr}(v)$ , the subgroup of  $G$  consisting of the  $h$  such that  $\text{int}(h)v = v$ , is formally smooth over  $S$ .*

*N.B.* The pair  $(v_1, v_2)$  may be regarded as a section over  $S$  of the second factor in the codomain of the morphism  $R$  in 2.3, and  $\text{Transp}(v_1, v_2)$  as the inverse-image functor of that section under  $R$ .

The assertion in 2.4 is itself equivalent to the following. Suppose that  $S$  is affine and that  $S_0 \subset S$  is a subscheme defined by a nilpotent ideal. If  $g_0 \in G(S_0)$  satisfies

$$\text{int}(g_0)(v_1)_{S_0} = (v_2)_{S_0},$$

then  $g_0$  extends to an element  $g \in G(S)$  such that

$$\text{int}(g)v_1 = v_2.$$

To prove this, first extend  $g_0$  to a section  $g'$  of  $G$  over  $S$ , which is possible because  $G$  is smooth over  $S$ , and put

$$v'_2 = \text{int}(g')v_1.$$

The morphisms  $v_2$  and  $v'_2$  have the same restriction over  $S_0$ . Hence, by Exposé IX, 3.6, there exists  $g'' \in G(S)$  inducing the identity section over  $S_0$  and such that

$$v_2 = \text{int}(g'')v'_2.$$

Therefore

$$v_2 = \text{int}(g'') \text{int}(g')v_1 = \text{int}(g''g')v_1,$$

and it is enough to take  $g = g''g'$ .

**Proposition 2.1 bis.** — *Let  $S$  be a prescheme and let  $G$  be an  $S$ -group prescheme smooth over  $S$ . Consider the functor*

$$M : (\mathbf{Sch})^\circ / S \longrightarrow (\mathbf{Ens}),$$

defined by

$$M(S') = \left\{ \begin{array}{c} \text{multiplicative-type subgroups} \\ \text{of } G_{S'} \end{array} \right\}.$$

Then  $M$  is formally smooth over  $S$ .

Cf. Exposé IX, 3.6 bis.

**Corollary 2.2 bis.** — Let  $n$  be an integer, and consider the morphism of functors

$$\varphi_n : M \longrightarrow M$$

defined by

$$\varphi_n(H) = {}_nH = \ker(n \cdot \text{id}_H).$$

Then  $\varphi_n$  is formally smooth. For every integer  $p > 0$ , let  $M_p$  denote the subfunctor of  $M$  for which  $M_p(S')$  is the set of multiplicative-type subgroups  $H$  of  $G_{S'}$  satisfying  ${}_pH = H$ . Then the morphism induced by  $\varphi_n$

$$M_{np} \longrightarrow M_n$$

is formally smooth.

The second assertion is trivially contained in the first and is included only for the convenience of a later reference. The proof of the first assertion is analogous to that of 2.2, now invoking Exposé IX, 3.6 bis.

**Corollary 2.3 bis.** — With the notation of 2.1 bis, regard  $M$  as a functor with operator group  $G$ , where  $G$  acts by

$$(g, H) \longmapsto \text{int}(g)(H).$$

Then the corresponding morphism

$$R : G \times_S M \longrightarrow M \times_S M,$$

defined by

$$(g, v) \longmapsto (\text{int}(g)v, v),$$

is formally smooth.

This is equivalent to the following assertion.

**Corollary 2.4 bis.** — With the notation of 2.1 bis, let  $H_1$  and  $H_2$  be two multiplicative-type subgroups of  $G$ , and let  $\text{Transp}_G(H_1, H_2)$  be the subfunctor of  $G$  consisting of the  $g$  such that

$$\text{int}(g)H_1 = H_2.$$

Then this functor is formally smooth over  $S$ . In particular, if  $H_1 = H_2 = H$ , the subfunctor  $\text{Norm}_G(H)$  of  $G$ , the normalizer of  $H$ , is formally smooth over  $S$ .

The proof is analogous to that of 2.4, again invoking Exposé IX, 3.6 bis.

**Proposition 2.5.** — Let  $S$  be a prescheme, let  $G$  be an  $S$ -group prescheme smooth over  $S$ , let  $K$  be an  $S$ -group subprescheme that is smooth over  $S$  or of multiplicative type, let  $H$  be an  $S$ -group prescheme of multiplicative type, and let  $u : H \rightarrow G$  be a homomorphism of  $S$ -groups. Denote by  $\text{Transp}_G(u, K)$  the subfunctor of  $G$  whose value at an  $S'$  over  $S$  consists of the elements  $g \in G(S')$  for which

$$\text{int}(g)u_{S'} : H_{S'} \longrightarrow G_{S'}$$

factors through  $K_{S'}$ . Then this functor is formally smooth over  $S$ .

The proof is analogous to that of 2.2, invoking Exposé IX, 3.6 and Exposé X, 2.1, the latter in the case where  $K$  is of multiplicative type.

### 3. Auxiliary Representability Results

**Proposition 3.1.** — *Let  $F \rightarrow S$  be a functor over the prescheme  $S$ . The following conditions are equivalent:*

- (i)  *$F$  is representable and  $F \rightarrow S$  is an open immersion (one also says simply that  $F \rightarrow S$  is an open immersion).*
- (ii)  *$F$  is a sheaf for the faithfully flat quasi-compact topology, “commutes with filtered direct limits of rings,” and  $F \rightarrow S$  is a monomorphism. Moreover, the following condition holds: for every local prescheme  $S'$  over  $S$ , with residue field  $k$ , and every  $S$ -morphism*

$$\mathrm{Spec}(k) = S'_0 \longrightarrow F,$$

*there exists an  $S$ -morphism  $S' \rightarrow F$  extending it.*<sup>6</sup>

- (iii) *Suppose that  $S$  is locally noetherian. Then  $F$  is a sheaf for the faithfully flat quasi-compact topology, commutes with filtered direct limits of rings, commutes with adic inverse limits of local artinian rings, and  $F \rightarrow S$  is a monomorphism and is infinitesimally étale (cf. 1.8).*

Let us first make two points of terminology precise.

**Remark 3.2.** — A functor<sup>7</sup>  $F$  over  $S$  is said to *commute with direct limits of rings* (understood to mean filtered direct limits) if the following holds. Let  $(S'_i)_{i \in I}$  be any filtered inverse system of affine  $S$ -schemes lying over an affine open subscheme of  $S$ , and let  $A'_i$  be their rings. Then the natural homomorphism

$$(*) \quad \varinjlim_i F(S'_i) \longrightarrow F(S'), \quad S' = \mathrm{Spec}(A'), \quad A' = \varinjlim_i A'_i$$

is bijective.

Notice that  $S'$  is precisely the inverse limit of the  $S'_i$  in the category of preschemes, and even in the category of all ringed spaces. Thus the condition under consideration has the character of a right-exactness condition—commutation with certain direct limits in  $(\mathbf{Sch})^\circ/S$ —just as does the condition of being a sheaf for a topology.

Care must be taken that this condition is essentially relative: it involves the morphism  $F \rightarrow S$ , not merely the functor

$$F : (\mathbf{Sch})^\circ \longrightarrow (\mathbf{Ens}).$$

More precisely, in  $(*)$ ,  $F(S'_i)$  and  $F(S')$  mean  $\mathrm{Hom}_S(S'_i, F)$  and  $\mathrm{Hom}_S(S', F)$ . Consequently, when  $F$  is representable, the condition says that  $F$  is locally of finite presentation over  $S$ . We used several times in the preceding two exposés the fact that a functor represented by an  $S$ -prescheme locally of finite presentation commutes with direct limits of rings.

**Remark 3.3.** — A functor  $F$  over  $S$  is said to *commute with adic inverse limits of local artinian rings* if the following condition holds. For every  $S'$  over  $S$  that is the spectrum of a complete noetherian local ring  $A'$ , set

$$S'_n = \mathrm{Spec}(A' / \mathrm{rad}(A')^{n+1}).$$

<sup>6</sup>The French re-edition prints  $S' \rightarrow S$ . Since this morphism must extend the displayed morphism  $S'_0 \rightarrow F$ , its intended target is  $F$ .

<sup>7</sup>Editor’s note: here and in Remark 3.3, the functor is contravariant.

Then the natural map

$$(**) \quad F(S') \longrightarrow \varinjlim_n F(S'_n)$$

is required to be bijective.

This condition has the character of a left-exactness condition and is satisfied whenever  $F$  is representable. In contrast to commutation with direct limits of rings, it is intrinsic to  $F$  as an object of  $\widehat{\mathbf{Sch}}$ ; that is, it does not involve the morphism  $F \rightarrow S$ .

**Remark 3.4.** — Let  $F$  be a functor over  $S$  that is a sheaf for the Zariski topology, or, as one also says, is “of local nature.” It is enough for this that  $F$  be a sheaf for a finer topology, such as the faithfully flat quasi-compact topology. Let  $(S_i)$  be an open covering of  $S$ . A straightforward gluing argument shows that  $F$  is representable if and only if each

$$F_i = F \times_S S_i$$

is representable; this allows one, for example, to reduce to the case where  $S$  is affine.

Suppose that the functor  $F$ , of local nature, commutes with direct limits of rings. Then  $F$  is representable if and only if its restriction to the category of preschemes locally of finite presentation over  $S$  is representable. Necessity was noted in 3.2. Sufficiency amounts to the following assertion: if  $X$  is a prescheme locally of finite presentation over  $S$ , and  $X \rightarrow F$  is a morphism such that, for every  $S'$  locally of finite presentation over  $S$ , the induced map

$$\mathrm{Hom}_S(S', X) \longrightarrow \mathrm{Hom}_S(S', F)$$

is bijective, then  $X \rightarrow F$  is an isomorphism. This follows readily from the fact that  $X$  and  $F$  are both functors of local nature that commute with direct limits of rings.

*Proof of Proposition 3.1.* The implications (i)  $\Rightarrow$  (ii) and (i)  $\Rightarrow$  (iii) are immediate. We prove the converse implications.

Assume (ii). Let  $U$  be the set of points  $s \in S$  for which the canonical monomorphism

$$\mathrm{Spec} \kappa(s) \longrightarrow S$$

factors through  $F$ . By the last condition in (ii), for every  $s \in U$ , the canonical map  $\mathrm{Spec} \mathcal{O}_{S,s} \rightarrow S$ , equivalently its factorization through  $U$ , factors through  $F$ . The local ring  $\mathcal{O}_{S,s}$  is the direct limit of the rings of affine neighborhoods of  $s$ . Since  $F$  commutes with direct limits of rings, for every  $s \in U$ <sup>8</sup> there is an open neighborhood  $U^s$  such that the canonical immersion  $U^s \rightarrow S$  factors through  $F$ . It follows that  $U^s \subset U$ , and hence that  $U$  is open.

Because  $F \rightarrow S$  is a monomorphism and  $F$  is of local nature, the  $S$ -morphisms  $U^s \rightarrow F$  agree on the intersections  $U^s \cap U^{s'}$ , for  $s, s' \in U$ , and therefore glue to an  $S$ -morphism  $U \rightarrow F$ . It remains to prove that this morphism is an isomorphism, or equivalently that every  $S$ -morphism  $S' \rightarrow F$  factors uniquely through  $U$ , where  $S'$  is an  $S$ -prescheme. Since both  $F \rightarrow S$  and  $U \rightarrow S$  are monomorphisms, it is equivalent to say that the structure morphism  $S' \rightarrow S$  factors through  $U$ .

---

<sup>8</sup>The French re-edition prints  $s \in S$  at this occurrence; the argument and the conclusion  $U^s \subset U$  require  $s \in U$ .

By locality, we are reduced to the case in which  $S'$  is the spectrum of a field and hence has a single point  $s'$ . Let  $s$  be the point of  $S$  below  $s'$ . We claim that the given  $S$ -morphism  $S' \rightarrow F$  factors through

$$\mathrm{Spec} \kappa(s) = S_0 \longrightarrow F.$$

This implies  $s \in U$  and proves the assertion. Since  $S' \rightarrow S_0$  is an fpqc covering and  $F$  is a sheaf for the fpqc topology, it is enough that the two composites

$$S'' = S' \times_{S_0} S' \rightrightarrows S' \longrightarrow F$$

coincide. They do coincide because  $F \rightarrow S$  is a monomorphism. This proves (ii)  $\Rightarrow$  (i).

Now assume (iii), with  $S$  locally noetherian. It is enough to prove the last condition of (ii); by the preceding proof, it is enough to do so when

$$S' = \mathrm{Spec} \mathcal{O}_{S,s}$$

for some  $s \in S$ . Put

$$A = \mathcal{O}_{S,s}, \quad A_n = A/\mathfrak{m}^{n+1}, \quad S_n = \mathrm{Spec}(A_n).$$

Because  $F \rightarrow S$  is infinitesimally smooth, the given morphism  $S'_0 \rightarrow F$  extends to morphisms  $S_n \rightarrow F$ . Since  $F \rightarrow S$  is a monomorphism, these morphisms form an element of  $\varprojlim_n F(S_n)$ . Since  $F$  commutes with adic inverse limits of local artinian rings, they arise from a morphism

$$\mathrm{Spec}(\widehat{A}) = \widehat{S}' \longrightarrow F.$$

Finally,  $F \rightarrow S$  is a monomorphism,  $F$  is an fpqc sheaf, and  $\widehat{S}' \rightarrow S'$  is an fpqc covering. The last morphism therefore descends to  $S' \rightarrow F$ , completing the proof of (iii)  $\Rightarrow$  (ii).  $\square$

**Proposition 3.5.** — *Let  $S$  be a locally noetherian prescheme, let  $F \rightarrow S$  be a functor over  $S$ , and let  $(X_i, u_i)_{i \in I}$  be a family of  $S$ -morphisms  $u_i : X_i \rightarrow F$ , where the  $X_i$  are preschemes locally of finite type over  $S$ . Suppose that the following conditions hold:*

- (a)  *$F$  is a sheaf for the faithfully flat quasi-compact topology, commutes with direct limits of rings, and commutes with adic inverse limits of local artinian rings.*
- (b) *The morphisms  $u_i : X_i \rightarrow F$  are monomorphisms and are infinitesimally étale (cf. 1.8).*
- (c) *The family  $(u_i)$  is “set-theoretically surjective.”*

*Under these conditions,  $F$  is representable by a prescheme locally of finite type over  $S$ . Moreover, the  $u_i$  are open immersions and the family  $(X_i)$  is an open covering of  $F$ .*

**Remark 3.6.** — Proceeding as at the end of Remark 1.7, one defines, for every functor

$$F : (\mathbf{Sch})^\circ \longrightarrow (\mathbf{Ens}),$$

an “underlying set”  $\mathrm{ens}(F)$ : it is the quotient of the set of field-valued points of  $F$  by the equivalence relation described in 1.7. When  $F$  is represented by  $X$ , this recovers the underlying set of  $X$ .

Plainly,  $\mathrm{ens}(F)$  depends functorially on  $F$ . Thus, if  $G \rightarrow F$  is a morphism of functors, we say that it is *set-theoretically surjective* if the induced map

$$\mathrm{ens}(G) \longrightarrow \mathrm{ens}(F)$$

is surjective. Equivalently, every point of  $F$  with values in a field  $k$  “comes from” a point of  $G$  with values in a suitable extension of  $k$ . This definition extends immediately to a family of morphisms  $G_i \rightarrow F$ , and thereby specifies the meaning of condition (c).

*Proof of Proposition 3.5.* For  $(i, j) \in I \times I$ , put

$$X_{i,j} = X_i \times_F X_j$$

and consider the projections

$$v_{i,j} : X_{i,j} \longrightarrow X_i, \quad w_{i,j} : X_{i,j} \longrightarrow X_j.$$

We claim that these morphisms are representable by open immersions. Apply criterion 3.1(iii). The functor  $X_{i,j}$  satisfies the three exactness conditions: it is a sheaf, it commutes with direct limits of rings, and it commutes with adic inverse limits of local artinian rings. Indeed,  $F$ ,  $X_i$ , and  $X_j$  satisfy these conditions, and the conditions are stable under finite inverse limits, in particular under fiber products. Since every  $X_i \rightarrow F$  is a monomorphism, both projections are monomorphisms, each being obtained from one of these maps by base change. Finally, being infinitesimally étale is also preserved by base change. Thus the hypotheses of 3.1(iii) are satisfied.

We may now use the  $X_i$ ,  $X_{i,j}$ ,  $v_{i,j}$ , and  $w_{i,j}$  to construct in the usual way an  $S$ -prescheme  $X$ , in which the  $X_i$  are identified with open subschemes, the  $X_{i,j}$  with the intersections  $X_i \cap X_j$ , and  $v_{i,j}$ ,  $w_{i,j}$  with the canonical immersions.

Put<sup>9</sup>

$$Y = \coprod_i X_i.$$

Then  $X$  is also the quotient of  $Y$  by the equivalence relation

$$R = \coprod_{i,j} X_{i,j} \rightrightarrows Y,$$

whose two projections  $v, w$  are defined by the  $v_{i,j}$  and  $w_{i,j}$ , respectively.

More precisely, because  $F$  is an fpqc sheaf, the morphisms  $u_i : X_i \rightarrow F$  induce a morphism  $u : Y \rightarrow F$ , and

$$R = Y \times_F Y$$

is exactly the equivalence relation defined by  $u$ . The quotient  $X = Y/R$  is also a quotient in the category of fpqc sheaves, and even in the category of Zariski sheaves; this follows directly from the definitions of quotient and sheaf. Consequently,  $u$  factors uniquely through a morphism

$$\bar{u} : X \longrightarrow F,$$

and this morphism is a monomorphism.

It remains to show that  $\bar{u}$  is an isomorphism. Since  $F$  is of local nature, we may assume that  $S$  is affine. Since  $F$  also commutes with direct limits of rings, it is enough, by 3.4, to verify that, for every affine  $T$  of finite type over  $S$ , every morphism  $T \rightarrow F$  factors through  $X$ . Consider

---

<sup>9</sup>The French re-edition labels this coproduct  $X$  and later writes  $R = X \times_F X$ , while simultaneously using  $R \rightrightarrows Y$  and  $X = Y/R$ . The coherent notation, adopted here, is  $Y = \coprod_i X_i$  and  $R = Y \times_F Y$ .

$$G = X \times_F T \longrightarrow T.$$

It is enough to prove that this morphism is an isomorphism. Since  $T$  is noetherian, the argument above shows that it is an open immersion. Here one uses that  $X \rightarrow F$  is infinitesimally étale, which follows immediately because the induced morphisms  $u_i : X_i \rightarrow F$  are infinitesimally étale. By hypothesis,  $X \rightarrow F$  is set-theoretically surjective, and this condition is stable under base change. Hence  $G \rightarrow T$  is set-theoretically surjective. Being also an open immersion, it is an isomorphism.  $\square$

**Proposition 3.7.** — *Let  $S$  be a locally noetherian prescheme, let  $I$  be a directed partially ordered index set, let  $(T_i)_{i \in I}$  be a projective system of  $S$ -preschemes locally of finite type, let*

$$T = \varprojlim_{i \in I} T_i$$

*be the projective-limit functor, let  $F$  be a functor over  $S$ , and let  $u : F \rightarrow T$  be an  $S$ -morphism. Suppose that the following conditions hold:*

- (a)  *$F$  is a sheaf for the faithfully flat quasi-compact topology, commutes with inductive limits of rings, and commutes with adic projective limits of local artinian rings.*
- (b) *The morphism  $u : F \rightarrow T$  is a monomorphism.*
- b' The morphism  $u : F \rightarrow T$  is infinitesimally étale.*
- (c) *For every point  $\xi$  of  $F$  with values in the spectrum of a field  $k$ , let  $\xi_i \in T_i(\text{Spec } k)$  be its image and let  $t_i$  be the corresponding point of  $T_i$ . There exists  $i \in I$  such that, for every  $j \geq i$ , the transition morphism*

$$p_{i,j} : T_j \longrightarrow T_i$$

*is étale at  $t_j$ .*

- (d) *For every prescheme  $X$  locally of finite type over  $S$  and every  $S$ -morphism  $X \rightarrow F$ , the set of points  $x \in X$  at which this morphism is infinitesimally étale is open.*

*Under these conditions,  $F$  is representable by a prescheme locally of finite type over  $S$ .*

Let us note at once that, in the case to be considered in the next section, conditions (c) and (d) will be verified by means of the following corollary.

**Corollary 3.8.** — *Assuming (a), (b), and (b'), conditions (c) and (d) are implied by the following conditions:*

- c' The  $T_i$  are smooth over  $S$ , and the transition morphisms  $p_{i,j} : T_j \rightarrow T_i$  are smooth.*
- d' For every point  $\xi$  of  $F$  with values in the spectrum of a field  $k$ , let  $t_i(\xi)$  be the point of  $T_i$  defined by  $\xi$ , let  $d_i(\xi)$  be the relative dimension of  $T_i$  over  $S$  at  $t_i(\xi)$ , and put*

$$d(\xi) = \sup_i d_i(\xi).$$

*Then:*

- 1° for every  $\xi$  as above,  $d(\xi) < +\infty$ ; and*



2° for every prescheme  $X$  locally of finite type over  $S$  and every  $S$ -monomorphism  $v : X \rightarrow F$ , the function

$$x \longmapsto d(\xi_x)$$

on  $X$  is locally constant, where, for  $x \in X$ ,  $\xi_x$  denotes the point of  $F$  with values in  $\kappa(x)$  induced by  $v$ .

*Proof of Proposition 3.7.* Work under condition (c). Let  $t$  be the element of  $\text{ens}(F)$  defined by  $\xi$  (cf. 3.6), and put

$$\mathcal{O}_t = \varinjlim_i \mathcal{O}_{T_i, t_i}.$$

Using condition (c), one sees readily that  $\mathcal{O}_t$  is a noetherian local ring (EGA 0<sub>IV</sub>, 10.3.1.3). Its residue field  $\kappa(t)$  is the inductive limit of the residue fields  $\kappa(t_i)$ , and  $k$  is an extension of it. If  $\eta$  denotes  $\text{Spec } \kappa(t)$ , and  $\xi$  denotes  $\text{Spec } k$ , there is a commutative diagram

$$\begin{array}{ccc} \xi & \longrightarrow & F \\ \downarrow & & \downarrow \\ \eta & \longrightarrow & T, \end{array}$$

whose definition is evident and is left to the reader. Since  $\xi \rightarrow \eta$  is a covering for the fpqc topology,  $F$  is a sheaf for that topology by (a), and  $F \rightarrow T$  is a monomorphism by (b), the morphism  $\eta \rightarrow T$  factors uniquely through a morphism  $\eta \rightarrow F$ . We now record the following lemma.

**Lemma 3.9.** — *Under the hypotheses of 3.7, let*

$$T' = \text{Spec}(A) \quad \text{and} \quad T'_0 = \text{Spec}(k(A))$$

*with  $T'$  a noetherian local scheme, and let  $i : T'_0 \rightarrow T'$  be the canonical immersion. Suppose that a commutative square of morphisms is given:*

$$\begin{array}{ccc} T'_0 & \longrightarrow & F \\ i \downarrow & & \downarrow \\ T' & \longrightarrow & T. \end{array}$$

*Then there exists a unique  $T$ -morphism  $T' \rightarrow F$ .*

The proof is the proof of 3.1, (iii)  $\Rightarrow$  (ii), in which the representability of the  $S$  occurring in that statement was not used. One uses the facts that  $F$  is a sheaf for the fpqc topology, that it commutes with adic projective limits of local artinian rings, here the rings

$$A/\text{rad}(A)^{n+1},$$

and that  $F \rightarrow T$  is both a monomorphism and infinitesimally étale.

We apply Lemma 3.9 with

$$T' = \text{Spec}(\mathcal{O}_t), \quad T'_0 = \text{Spec}(\kappa(t)).$$

We have just constructed  $T'_0 \rightarrow F$ , and the canonical homomorphisms

$$\mathcal{O}_{T_i, t_i} \longrightarrow \mathcal{O}_t$$

define a projective system of morphisms  $\mathrm{Spec}(\mathcal{O}_t) \rightarrow T_i$ , hence a canonical morphism  $\mathrm{Spec}(\mathcal{O}_t) \rightarrow T$ . The corresponding square (x) plainly commutes: by the definition of  $T$ , it is enough to check this after replacing  $T$  by  $T_i$ . We therefore obtain a unique  $T$ -morphism

$$v' : T' = \mathrm{Spec}(\mathcal{O}_t) \longrightarrow F.$$

Since  $F$  commutes with inductive limits of rings, this morphism factors, for sufficiently large  $i$ , through

$$v'_i : T'_i = \mathrm{Spec}(\mathcal{O}_{T_i, t_i}) \longrightarrow F.$$

For such an  $i$ , one has

$$T' \xrightarrow{\sim} T'_i, \quad \text{that is,} \quad \mathcal{O}_{T_i, t_i} \xrightarrow{\sim} \mathcal{O}_t.$$

Indeed,  $T' \rightarrow T'_i$  is faithfully flat and quasi-compact, hence an effective epimorphism, so it is enough to show that it is a monomorphism. Suppose that two morphisms  $R \rightrightarrows T'$ , with  $R$  representable, have the same composite with  $T' \rightarrow T'_i$ . Their composites with  $T' \rightarrow T$ , which factors through  $T' \rightarrow T'_i$ , are then equal. Hence their composites with  $T' \rightarrow T_j$ , and therefore with  $T' \rightarrow T'_j$ , are equal for every  $j$ . The two original morphisms are equal because  $T'$  is the projective limit of the  $T'_j$  in the category **(Sch)**.

Consider the diagram

$$\begin{array}{ccc} T' & \xrightarrow{v'} & F \\ \downarrow & \nearrow v'_i & \downarrow \\ T'_i & \xrightarrow{\quad} & T_i, \end{array}$$

where the unlabeled arrows are the evident ones. The square and the upper triangle commute. Since  $T' \rightarrow T'_i$  is an epimorphism, the lower triangle also commutes.

The ring  $\mathcal{O}_{T_i, t_i}$  is the inductive limit of the rings of affine open neighborhoods of  $t_i$  in  $T_i$ . Since  $F$  commutes with inductive limits of rings,  $v'_i$  comes from a morphism

$$v_t : U_t \longrightarrow F$$

defined on an open neighborhood  $U_t$  of  $t_i$  in  $T_i$ . After shrinking this neighborhood if necessary,  $v_t$  is a  $T_i$ -morphism. Indeed,  $T_i$ , being locally of finite type over  $S$ , also commutes with inductive limits of rings. The composite of  $v_t$  with  $F \rightarrow T_i$  is a monomorphism, and therefore  $v_t$  is a monomorphism.

We claim that  $v_t$  is infinitesimally étale at  $t_i$ . Since  $F \rightarrow T$  is infinitesimally étale, this is equivalent to the composite

$$T'_i \xrightarrow{v'_i} F \longrightarrow T$$

being infinitesimally étale at  $t_i$ , or to the induced morphism  $T'_i \rightarrow T$  being infinitesimally étale. Since  $T' \xrightarrow{\sim} T'_i$ , it is in turn equivalent to  $T' \rightarrow T$  being infinitesimally étale. This is immediate, because that morphism is the projective limit of the infinitesimally étale morphisms  $T'_j \rightarrow T_j$ .

Apply condition (d), which has not yet been used. It follows that  $v_t$  is infinitesimally étale in a neighborhood of  $t_i$ . After replacing  $U_t$  by a smaller open, we may therefore suppose that  $v_t$  is a monomorphism that is infinitesimally étale.

As  $t$  varies in  $\mathrm{ens}(F)$ , the family of morphisms  $v_t$  satisfies Proposition 3.5, which gives the conclusion of Proposition 3.7.

*Proof of Corollary 3.8.* Assume conditions (c') and (d'). With the notation of (d'),  $d_i(\xi)$  is constant for all sufficiently large  $i$ . Consequently, the relative dimension of  $T_j$  over  $T_i$  at  $t_j$ , which is

$$d_j(\xi) - d_i(\xi),$$

is zero. Thus  $T_j \rightarrow T_i$  is étale at  $t_j$ , which proves condition (c).

The preceding proof now gives, for each  $t \in \text{ens}(F)$ , an index  $i$ , an open neighborhood  $U_t$  of  $t_i$ , and a morphism  $U_t \rightarrow F$  that is a monomorphism and is infinitesimally étale at  $t_i$ . It remains to show that this morphism is infinitesimally étale throughout a neighborhood of  $t_i$ . In condition (d'),  $2^\circ$ , take  $X = U_t$ . Replacing  $U_t$ , if necessary, by the connected component of  $t_i$  in  $U_t$ , we may suppose that

$$d(\xi_x) = d(\xi_{t_i}) \quad \text{for every } x \in U_t.$$

Now  $d(\xi_{t_i})$  is the relative dimension of  $T_i$  over  $S$  at  $t_i$ . Since that relative dimension remains constant on  $U_t$ , we also have, for every  $x \in U_t$ ,

$$(*) \quad d(\xi_x) = \text{the relative dimension of } U_t \text{ over } S \text{ at } x.$$

On the other hand, using the notation of the proof of 3.7, consider any local noetherian prescheme

$$R' = \text{Spec}(A), \quad R'_0 = \text{Spec}(k(A)).$$

Every morphism  $R' \rightarrow F$  whose induced point  $R'_0 \rightarrow F$  represents  $t \in \text{ens}(F)$ , that is, sends the closed point of  $R'$  to  $t$ , factors uniquely through  $T'$ . The proof is the proof of 3.1 or of Lemma 3.9: one uses that  $T' \rightarrow F$  is a monomorphism infinitesimally étale at  $t$ , and that  $T'$  is an fpqc sheaf commuting with adic projective limits.

Apply this result to the morphism

$$\text{Spec}(\mathcal{O}_{U_t, x}) = R' \longrightarrow F$$

induced by  $v_t$ , and to the point  $t_x \in \text{ens}(F)$  that is the image of the closed point of  $R'$ , or equivalently the image of  $x$  under  $v_t$ . We obtain a factorization

$$\text{Spec}(\mathcal{O}_{U_t, x}) \longrightarrow \text{Spec}(\mathcal{O}_{t_x}) \longrightarrow F.$$

The second arrow is infinitesimally étale at  $t_x$ . To prove that the composite is infinitesimally étale at  $x$ , it is therefore enough to prove the same for the first arrow at  $x$ .

By (x) above and formula (\*), the first arrow is induced by a local  $S$ -homomorphism between the localizations, at  $x$  and  $t_x$ , of the smooth  $S$ -preschemes  $U_t$  and  $U_{t_x}$ , which have the same relative dimension  $d$  over  $S$  at those points. By EGA I, 6.5.1, this local homomorphism is induced by an  $S$ -morphism

$$w : U \longrightarrow V,$$

where  $U$  is an open neighborhood of  $x$  in  $U_t$  and  $V = U_{t_x}$ . It remains to prove that  $w$  is étale at  $x$ . Moreover,  $U$  and  $V$  carry monomorphisms to  $F$ , and, after shrinking  $U$  once more,  $w$  is an  $F$ -morphism. Hence  $w$  is a monomorphism. It is now enough to prove the following lemma.

**Lemma 3.10.** — *Let  $U$  and  $V$  be two smooth  $S$ -preschemes of the same relative dimension  $d$  over  $S$ , and let  $w : U \rightarrow V$  be an  $S$ -morphism that is a monomorphism. Then  $w$  is an open immersion, and hence is étale.*

By SGA 1, I, 5.7, we reduce to the case in which  $S$  is the spectrum of a field, which may be assumed algebraically closed. By SGA 1, I, 5.1, it is enough to prove that  $w$  is étale, and it is enough to do so at the closed points of  $U$ . Let  $x$  be such a point and put  $y = w(x)$ . Choose a regular system of parameters  $f_1, \dots, f_d$  of  $\mathcal{O}_{V,y}$ . Then

$$A = \mathcal{O}_{U,x} / \sum_i f_i \mathcal{O}_{U,x}$$

is the trivial extension of

$$k = \mathcal{O}_{V,y} / \sum_i f_i \mathcal{O}_{V,y};$$

that is,  $A \simeq k$ . Indeed, since  $w$  is a monomorphism, so is the structural morphism

$$\mathrm{Spec}(A) \longrightarrow \mathrm{Spec}(k)$$

obtained from it by base change. Since  $\mathcal{O}_{U,x}$  is a regular local ring of dimension  $d$ , the  $f_i$  form a regular system of parameters of that ring. It follows immediately that  $w$  is étale at  $x$ . This proves Lemma 3.10, and hence Corollary 3.8.

**Corollary 3.11.** — *Under the hypotheses of 3.8, for every quasi-compact open subfunctor  $U$  of  $F$  that is separated over  $S$ , there exists  $i \in I$  such that, for every  $j \geq i$ , the morphism*

$$u_j|_U : U \longrightarrow T_j$$

*is an open immersion. In particular, if the  $T_i$  are quasi-affine over  $S$ , then every open subfunctor  $U$  of  $F$  that is quasi-compact over  $S$ , that is, of finite type over  $S$ , is quasi-affine over  $S$ .*

The proof of Proposition 3.7 shows that for every  $t \in \mathrm{ens}(F)$  there exists  $i \in I$  such that  $u_i : F \rightarrow T_i$  is a local isomorphism at  $t$ , and then  $u_j$  is a local isomorphism at that point for every  $j \geq i$ . By quasi-compactness,  $i$  may be chosen independently of  $x \in U$ . It remains to prove that, for sufficiently large  $i$ ,

$$u_i|_U : U \longrightarrow T_i$$

is a monomorphism.

Since  $U \rightarrow T$  is a monomorphism, the intersection of the equivalence relations

$$U \times_{T_i} U \subset U \times_S U$$

is the diagonal. The prescheme  $U \times_S U$  is noetherian, and the  $U \times_{T_i} U$  are closed subpreschemes. It follows that one of them is already the diagonal; equivalently,  $u_i|_U$  is a monomorphism. This proves the first assertion of Corollary 3.11, and the second follows immediately.

**Proposition 3.12.** — *Let  $S$  be a prescheme and  $G$  an affine  $S$ -group prescheme.*

(a) *Let*

$$F : (\mathbf{Sch})^\circ / S \longrightarrow (\mathbf{Ens})$$

*be the functor such that, for every  $T$  over  $S$ ,*

$$F(T) = \left\{ \begin{array}{c} \text{multiplicative-type subgroups of } G_T \\ \text{that are finite over } T \end{array} \right\}.$$

*Suppose that  $S$  is locally noetherian or that  $G$  is of finite presentation over  $S$ . Then  $F$  is representable and affine over  $S$ . If  $G$  is of finite presentation over  $S$ , then  $F$  is locally of finite presentation over  $S$ .*

- (b) Let  $H$  be a finite  $S$ -group prescheme of multiplicative type. Then  $\mathrm{Hom}_{S\text{-gr}}(H, G)$  is representable and affine over  $S$ . If  $G$  is of finite type (respectively, of finite presentation) over  $S$ , then the same is true of  $\mathrm{Hom}_{S\text{-gr}}(H, G)$ .

**Remark 3.13.** — Apart from the assertion that  $\mathrm{Hom}_{S\text{-gr}}$  is affine, and from the case in which  $G$  is of finite presentation over  $S$ , which is the case that will suffice for us, 3.12 is an immediate consequence of the theory of Hilbert schemes (A. Grothendieck, *Techniques de construction et théorèmes d'existence en Géométrie Algébrique: IV. Les Schémas de Hilbert*, Séminaire Bourbaki, May 1961, no. 221). It is enough, in fact, that  $G$  be quasi-projective over  $S$ . In case (a) one can also represent the larger functor

$$F_0(T) = \left\{ \begin{array}{l} \text{group subpreschemes of } G_T \text{ that are flat, proper,} \\ \text{and of finite presentation over } T \end{array} \right\}.$$

The canonical monomorphism  $F \rightarrow F_0$  is an “open immersion,” as follows from the criterion in 3.1 and Exposé X, 4.7 (b). Thus the representability of  $F_0$  implies that  $F$  is representable by an open subprescheme of  $F_0$ . In case (b) it is enough to assume that  $H$  is projective and of finite presentation over  $S$ . In both cases one obtains a functor locally of finite presentation over  $S$ .

In the present Exposé, 3.12 is only a technical lemma used to prove a key result in the next section. We shall therefore sketch an easy direct proof that does not use Hilbert schemes.

We first prove (b). It is enough to prove that  $\mathrm{Hom}_S(H, G)$  is representable, independently of any group structure on  $H$  and  $G$ , and that it has the additional properties asserted for  $\mathrm{Hom}_{S\text{-gr}}(H, G)$ . Indeed, applying the same result to  $\mathrm{Hom}_S(H \times_S H, G)$ , the subfunctor  $\mathrm{Hom}_{S\text{-gr}}(H, G)$  of  $\mathrm{Hom}_S(H, G)$  can be expressed by a finite inverse limit—in fact, by fiber products—involving  $G$ ,  $\mathrm{Hom}_S(H, G)$ , and  $\mathrm{Hom}_S(H \times_S H, G)$ . We leave this expression to the reader.

Write

$$H = \mathrm{Spec}(B),$$

where  $B$  is a sheaf of algebras on  $S$  that is locally free as a sheaf of modules; this is the only hypothesis on  $H$  that the argument uses. Since the representability question is local on  $S$ , we may suppose that  $S$  is affine, with ring  $A$ . We may also write  $G = \mathrm{Spec}(C)$ , where  $C$  is an  $A$ -algebra.

When  $G = S[t] = G_0$ , with  $t$  an indeterminate, the Hom functor is simply

$$T \longmapsto \Gamma(T, B_T).$$

Since  $B$  is locally free, this functor is represented by the vector bundle  $V(B^\vee)$ , where  $B^\vee$  is the dual module sheaf. When  $G = S[(t_i)_{i \in I}] = G_0^I$ , for a not necessarily finite family of indeterminates, it follows that<sup>10</sup>

$$\mathrm{Hom}_S(H, G) = \mathrm{Hom}_S(H, G_0)^I = V(B^\vee)^I = V(B^{\vee(I)}).$$

In general  $G$  is isomorphic to a closed subscheme of a scheme of the form  $S[(t_i)_{i \in I}]$ ; equivalently,  $C$  is a quotient of an  $A$ -algebra  $A[(t_i)_{i \in I}]$ . Let  $(F_j)$  be a family of generators

<sup>10</sup>The source display prints  $\mathrm{Hom}_S(H, G)^I = V(B^\vee)^I = V(B^{\vee(I)})$ . Since the preceding sentence sets  $G = G_0^I$ , the displayed equality requires the explicit factor  $\mathrm{Hom}_S(H, G) = \mathrm{Hom}_S(H, G_0)^I$ .

of the ideal by which one takes the quotient. After covering  $S$  by smaller affine open subsets, we may assume that  $B$  is free of rank  $n$ . Choose a basis  $(e_k)_{1 \leq k \leq n}$  of  $B$ . Writing

$$x_i = \sum_{k=1}^n x_{ik} e_k$$

with indeterminate coefficients  $x_{ik}$  in an unspecified  $A$ -algebra  $A'$ , expand  $F_j((x_i))$  in this basis. For every  $F_j$  this gives  $n$  polynomials

$$F_{j,k} \quad (1 \leq k \leq n)$$

in the variables  $(x_{i,k})_{i \in I, 1 \leq k \leq n}$ , with coefficients in  $A$ . It follows at once that  $\mathrm{Hom}_S(H, G)$  is represented by the spectrum of the quotient of the polynomial ring  $A[(x_{i,k})]$  by the ideal generated by the  $F_{j,k}$ . This proves (b).

We now prove (a). For every ordinary finite commutative group  $M$ , let  $F_M$  be the subfunctor of  $F$  obtained by restricting to those subgroups of  $G_T$  that are of multiplicative type and of type  $M$ . A gluing argument like the one used in 3.5 (which we should have stated after decomposing the problem a little further) shows that it is enough to prove that each  $F_M$  is representable. Then  $F$  is represented by the disjoint sum of the  $F_M$ , with  $M$  ranging over the isomorphism classes of finite commutative groups. Indeed,  $F$  is the corresponding sum in the category of sheaves.

Fix  $M$  and write  $F$  in place of  $F_M$ . Put  $H = D_S(M)$ , and let

$$F' = \mathrm{Imm}_{S\text{-gr}}(H, G) \subset \mathrm{Hom}_{S\text{-gr}}(H, G)$$

be the subfunctor whose value at  $T$  is the set of homomorphisms of  $T$ -groups  $H_T \rightarrow G_T$  that are closed immersions. By (b),  $\mathrm{Hom}_{S\text{-gr}}(H, G)$  is represented by an affine  $S$ -prescheme. Using Exposé IX, 6.8, one sees immediately that  $F'$  is represented by an open-and-closed subscheme of it; in particular,  $F'$  is affine over  $S$ .

Consider the canonical morphism  $F' \rightarrow F$ , which assigns to every monomorphism  $H_T \rightarrow G_T$  its image group. Directly from the definitions,  $F'$  is thereby a principal homogeneous bundle over  $F$ , in the category of fpqc sheaves, under

$$\Gamma_F = \Gamma_S \times_S F, \quad \Gamma = \mathrm{Aut}_{\mathrm{gr}}(M).$$

It follows by descent that  $F' \rightarrow F$  is representable: for every morphism  $T \rightarrow F$  with  $T$  representable, the prescheme  $T \times_F F'$  is representable over  $T$ , in fact by a principal Galois bundle under  $\Gamma$ .

By Exposé IV, 4.6.6,  $F$  is therefore representable if and only if the quotient  $F'/F''$ , where  $F'' = F' \times_F F'$ , exists in **(Sch)** and is universally effective for faithfully flat quasi-compact morphisms. Equivalently, the quotient  $F'/\Gamma$  must exist and be universally effective for such morphisms. Since  $F'$  is affine over  $S$ , Exposé V, 4.1 shows that this condition is satisfied. This proves the representability of  $F$  in (a).

Finally, suppose that  $G$  is of finite presentation over  $S$ . The additional assertion follows immediately from the preceding proof: by (b),  $F'$  is then locally of finite presentation over  $S$ , so that Exposé V may be applied.<sup>11</sup>

<sup>11</sup>Editor's note: the source marks this final reference to Exposé V as needing verification or further specification.

## 4. The Scheme of Multiplicative-Type Subgroups of a Smooth Affine Group

The principal result of the present Exposé is the following.

**Theorem 4.1.** — *Let  $S$  be a prescheme, let  $G$  be an  $S$ -group prescheme smooth and affine over  $S$ , and let*

$$F: (\mathbf{Sch})^\circ/S \longrightarrow (\mathbf{Ens})$$

*be the functor defined by*

$$F(T) = \{\text{multiplicative-type subgroups of } G_T\}.$$

*Then  $F$  is representable, smooth, and separated over  $S$ .*

Let us record at once the following variant.

**Corollary 4.2.** — *Let  $G$  and  $H$  be two  $S$ -group preschemes, with  $G$  smooth and affine over  $S$ , and  $H$  of multiplicative type and of finite type. Then  $\mathrm{Hom}_{S\text{-gr}}(H, G)$  is representable, smooth, and separated over  $S$ .*

Indeed, when  $H$  is smooth over  $S$ , so is  $H \times_S G$ , and one may apply 4.1 to the latter. By considering graph subgroups,  $\mathrm{Hom}_{S\text{-gr}}(H, G)$  becomes a subfunctor  $F'$  of the functor  $F$  of multiplicative-type subgroups of  $H \times_S G$ : namely,  $F'(T)$  is the set of multiplicative-type subgroups  $K$  of

$$(H \times_S G)_T = H_T \times_T G_T$$

such that the homomorphism induced by the first projection,

$$K \longrightarrow H_T,$$

is an isomorphism. By Exposé IX, 2.9, the canonical morphism  $F' \rightarrow F$  is an open immersion. Since  $F$  is representable, the same is true of  $F'$ , which is represented by an open subprescheme of  $F$ ; and since  $F$  is separated and smooth over  $S$ , so is  $F'$ .

When  $H$  is finite over  $S$ , it is enough to apply 3.12(b) to obtain the representability of  $\mathrm{Hom}_{S\text{-gr}}(H, G)$ . When  $H = H_1 \times_S H_2$ , with  $H_1$  smooth over  $S$  and  $H_2$  finite over  $S$ , then  $\mathrm{Hom}_{T\text{-gr}}(H_T, G_T)$  is identified with the subset of

$$\mathrm{Hom}_{T\text{-gr}}((H_1)_T, G_T) \times \mathrm{Hom}_{T\text{-gr}}((H_2)_T, G_T)$$

formed by the pairs  $(u_1, u_2)$  such that  $u_1 \cdot u_2 = u_2 \cdot u_1$ . It follows that  $\mathrm{Hom}_{S\text{-gr}}(H, G)$  is represented by a closed subprescheme of

$$\mathrm{Hom}_{S\text{-gr}}(H_1, G) \times_S \mathrm{Hom}_{S\text{-gr}}(H_2, G) = X.$$

Indeed, apply Exposé VIII, 6.5(b), with  $Y = H$ ,  $Z = G$ , and with  $q_1$  and  $q_2$  defined respectively by

$$(u_1, u_2) \longmapsto u_1 \cdot u_2 \quad \text{and} \quad (u_1, u_2) \longmapsto u_2 \cdot u_1.$$

In the general case, the question is local on  $S$  for the Zariski topology, so we may suppose that  $S$  is affine and that  $H$  has constant type over  $S$ . Since  $H$  is quasi-isotrivial (Exposé X, 4.5), there is a surjective étale morphism  $S' \rightarrow S$ , with  $S'$  affine, that splits  $H$ , that is, such that  $H' = H_{S'}$  is diagonalizable. The preceding result then applies, because a diagonalizable group is the product of a diagonalizable torus and a finite diagonalizable group. It remains to see that the descent datum obtained on the  $S'$ -prescheme

$$\mathrm{Hom}_{S'\text{-gr}}(H', G') = X'$$

is effective. This follows from the argument of Exposé X, 5.4, noting that, there, the hypothesis that  $X' \rightarrow S'$  be separated and locally quasi-finite was used only to ensure that every open part of  $X'$  quasi-compact over  $S'$  was quasi-affine over  $S'$ . That property still holds here, as follows readily from the fact that it holds for the functor  $F$  of 4.1 (as will be seen during the proof of 4.1). This argument—or any similar variant of this small dévissage—establishes the representability of  $\mathrm{Hom}_{S\text{-gr}}(H, G)$ , and at the same time shows that it is separated over  $S$ . It is locally of finite presentation over  $S$ , as follows, for example, from the fact already noted in 3.2 that this functor “commutes with filtered direct limits of rings.” Finally, the functor is formally smooth over  $S$  by 2.1, and is therefore smooth over  $S$ .

**Remark 4.3.** — We have deduced 4.2 from 4.1; this is genuinely immediate only when  $H$  is also smooth over  $S$ . To make the deduction without contortions in the general case, one would need the representability assertion in 4.1 without assuming that  $G$  is smooth over  $S$ , assuming only that it is affine and of finite presentation over  $S$ . Of course,  $F$  would then no longer be smooth over  $S$  in general. There is little doubt that 4.1 remains true under these more general hypotheses, but the proof appears likely to be more delicate, because one can no longer invoke 3.8.<sup>1213</sup>

Let us nevertheless point out that when  $G$  is a closed subgroup of an affine smooth group  $G_0$  over  $S$ , the functor  $F$  of multiplicative-type subgroups of  $G$  is represented by a closed subprescheme of the prescheme representing the analogous functor  $F_0$  for  $G_0$ , which is covered by 4.1; this follows readily by applying Exposé VIII, 6.4. This also raises the question whether an affine  $S$ -group scheme  $G$ , affine and of finite presentation over  $S$ , is isomorphic to a subgroup scheme of  $\mathrm{GL}(n)_S$  for some suitable  $n$ . This is true when  $S$  is the spectrum of a field (cf. Exposé VI<sub>B</sub>, 11.11), but is unfortunately false in general, even for tori; see 4.6. Finally, one could also prove 4.2 directly by exactly the same method as 4.1.

We now prove 4.1. Since the functor  $F$  is evidently local in nature, we may suppose that  $S$  is affine, say  $S = \mathrm{Spec}(A)$ , where  $A$  is a ring. Regarding  $A$  as the filtered direct limit of its subrings of finite type over  $\mathbb{Z}$ , and noting that  $G$  descends from a smooth affine group over one such subring (EGA IV, 8), we are reduced to the case in which  $S$  is noetherian.

For every integer  $n > 0$ , let  $T_n$  be the functor defined like  $F$ , but restricted to multiplicative-type subgroups  $H$  of  $G_T$  such that  $n \cdot \mathrm{id}_H = 0$ , that is, such that  ${}_nH = H$ . Order the set  $I$  of positive integers by divisibility. When  $m$  is a multiple of  $n$ , define

$$p_{n,m}: T_m \longrightarrow T_n$$

by

$$p_{n,m}(H) = {}_nH = \mathrm{Ker}(n \cdot \mathrm{id}_H).$$

In this way the  $T_n$  form an inverse system of functors over  $S$ . By 3.12(a), the functors  $T_n$  are representable, affine, and of finite type over  $S$ . Similarly define morphisms

$$u_n: F \longrightarrow T_n$$

<sup>12</sup>This is indeed proved for  $G$  flat and quasi-affine over  $S$ , with connected fibres, provided one restricts to the central subtori of  $G$  (Exposé XV, 8.8).

<sup>13</sup>Editor’s note: when  $H$  is smooth (not necessarily affine) over a normal locally noetherian base  $S$ , M. Raynaud showed that the largest representable open subfunctor of the functor of subtori of  $H$  is a disjoint sum of opens smooth and affine over  $S$ . This is Theorem IX.9.26 in *Faisceaux amples sur les schémas en groupes et sur les espaces homogènes*, Lecture Notes in Mathematics 119 (1970).



by  $u_n(H) = {}_nH$ . Thus one obtains a morphism

$$u: F \longrightarrow T = \varprojlim_n T_n,$$

where the inverse limit is taken in the category of functors over  $S$ . Since the  $T_n$  are representable and affine over  $S$ , so is  $T$ : it is the spectrum of the filtered direct limit of the quasi-coherent algebras on  $S$  defining the  $T_n$ . Of course,  $T$  is not in general of finite type over  $S$ .

We shall apply 3.7. It is enough to verify conditions (a)–(d) there; these will imply that  $F$  is represented by a prescheme locally of finite type over  $S$ . It then follows from 2.1 bis that  $F$  is even smooth over  $S$ . Moreover, since  $F$  is a subfunctor of  $T$ , which is affine over  $S$ ,  $F$  is separated over  $S$ : it is separated over  $T$ , and  $T$  is separated over  $S$ . We can also prove at once the supplementary assertion used above: every open  $U \subset F$  that is quasi-compact over  $S$  is quasi-affine over  $S$ . This follows from 3.11 and the fact that the  $T_n$  are affine over  $S$ .

Let us now verify the conditions of 3.7.

(a)  $F$  is a sheaf for the faithfully flat quasi-compact topology, by the descent theory of SGA 1, Exposé VIII, which applies here because groups of multiplicative type over  $T$  are affine over  $T$ .<sup>14</sup> It commutes with filtered direct limits of rings by the standard machinery of EGA IV, 8. Let us show that it commutes with adic inverse limits of local artinian rings. For the analogous functor considered in 4.2, this is precisely Exposé IX, 7.1, in the special case where  $A$  is a complete noetherian local ring endowed with an ideal of definition for its usual topology. Notice that this is exactly where the hypothesis that  $G$  be affine enters essentially. In the present case we are reduced to the following lemma.

**Lemma 4.4.** — *Let  $A$  be a complete noetherian local ring with maximal ideal  $\mathfrak{m}$ , let  $G$  be a group scheme affine over  $S = \operatorname{Spec}(A)$ , and, for every integer  $n \geq 0$ , put*

$$S_n = \operatorname{Spec}(A/\mathfrak{m}^{n+1}), \quad G_n = G \times_S S_n.$$

*For every  $n$ , let  $H_n$  be a multiplicative-type subgroup of finite type of  $G_n$ , and suppose that, whenever  $m \geq n$ ,  $H_n$  is obtained from  $H_m$  by reduction. Then there exists a unique multiplicative-type subgroup  $H$  of  $G$  whose reductions are the  $H_n$ .*

By Exposé X, 3.2, there exists a multiplicative-type group  $H$  over  $S$ , necessarily of finite type and isotrivial and determined up to unique isomorphism, together with an isomorphism

$$H \times_S S_0 \simeq H_0.$$

Here one uses the fact that  $H_0$  is isotrivial, since it is of finite type over a field; see Exposé X, 1.4. By Exposé X, 2.1, for every  $n$  the isomorphism  $H \times_S S_0 \simeq H_0$  lifts to a unique isomorphism  $H \times_S S_n \simeq H_n$ . Exposé IX, 7.1, already cited, now shows that the homomorphisms  $H_n \rightarrow G_n$  arise from a unique homomorphism of  $S$ -groups

$$u: H \longrightarrow G.$$

By Exposé IX, 6.6, this homomorphism is a monomorphism because  $u_0: H_0 \rightarrow G_0$  is one. This proves 4.4.

(b) The morphism  $u: F \rightarrow T$  is a monomorphism. This follows from the density theorem, Exposé IX, 4.7, in the form of Corollary 4.8(b). It is essential for the present application

<sup>14</sup>The printed source says “affine over  $S$ ” here; the descent assertion concerns groups over  $T$ , so the required base is  $T$ .

that this result is available over a base that need not be noetherian. Since the functor  $T$  does not commute with filtered direct limits of rings, one cannot reduce to the noetherian case a priori.

(b') The morphism  $u: F \rightarrow T$  is infinitesimally étale; in other words, we have the following lemma.

**Lemma 4.5.** — *Let  $A$  be an artinian local ring with residue field  $k$ , let  $S = \operatorname{Spec}(A)$ , let  $I \subset \operatorname{rad}(A)$  be an ideal, and put  $S_0 = \operatorname{Spec}(A/I)$ . Let  $G$  be an  $S$ -group prescheme smooth over  $S$ , and put  $G_0 = G \times_S S_0$ . Suppose given a multiplicative-type subgroup  $H_0$  of  $G_0$  and, for every integer  $n > 0$ , a multiplicative-type subgroup  $H(n)$  of  $G$ , in such a way that*

1° *for every multiple  $m$  of  $n$ ,  $H(n) = {}_nH(m)$ ; and*

2°  *$H(n)_0 = {}_nH_0$ .*

*Then there exists a unique multiplicative-type subgroup  $H$  of  $G$  such that  ${}_nH = H(n)$  for every  $n$ .*

Uniqueness is already contained in (b). For existence, an immediate induction reduces us to the case in which  $I\mathfrak{m} = 0$ , where  $\mathfrak{m}$  is the maximal ideal of  $A$ .<sup>15</sup> Put

$$k = A/\mathfrak{m}, \quad S_0 = \operatorname{Spec}(k), \quad G_0 = G \times_S S_0,$$

and let  $\mathfrak{g}_0$  and  $\mathfrak{h}_0$  be the Lie algebras of  $G_0$  and  $H_0$ , respectively. There is a canonical isomorphism of groups

$$\mathfrak{g}_0 \otimes_k I \simeq \operatorname{Ker}(G(S) \rightarrow G(S_0));$$

see Exposé III. By Exposé IX, 3.6 bis and 3.7, there exists a multiplicative-type subgroup  $H$  of  $G$  reducing to  $H_0$ , and such an  $H$  is determined, up to an inner automorphism by an element of

$$N = \operatorname{Ker}(G(S) \rightarrow G(S_0)).$$

Thus the set  $P$  of liftings  $H$  of  $H_0$  is a principal homogeneous set under  $N/M$ , where

$$M = \{g \in N \mid \operatorname{int}(g) \cdot H = H\}.$$

One sees readily (Exposé III) that this subgroup is precisely the vector subspace of  $\mathfrak{g}_0 \otimes_k I$  consisting of the invariants under  $H_0$ , where  $H_0$  acts on  $\mathfrak{g}_0 \otimes_k I$  through the representation induced by the adjoint representation of  $G_0$ . Likewise, the set  $P(n)$  of liftings of  $H(n)_0 = {}_nH_0$  is a principal homogeneous set under  $N/M(n)$ , where  $M(n)$  is the vector subspace of  $\mathfrak{g}_0 \otimes_k I = N$  consisting of the invariants under  ${}_nH_0$ . The density theorem, Exposé IX, 4.7, shows readily that  $M = M(n)$  for sufficiently large  $n$ , where “large” is understood with respect to the divisibility order on the positive integers. Consequently, the natural map

$$H \mapsto {}_nH$$

from  $P$  to  $P(n)$ , which is compatible with the actions of  $N$ , and hence with the homomorphism

$$N/M \rightarrow N/M(n)$$

between the acting groups, is bijective for sufficiently large  $n$ . The conclusion of 4.5 follows immediately.

<sup>15</sup>The printed source abruptly writes  $J\mathfrak{m} = 0$  here and uses  $J$  throughout the ensuing argument, although only  $I$  was introduced in the statement. We retain  $I$ , the same ideal, throughout.

To verify conditions (c) and (d) of 3.7, we use 3.8, which reduces us to checking (c') and (d') below.

(c') The  $T_n$  are smooth over  $S$ , and the transition morphisms  $p_{n,m}: T_m \rightarrow T_n$  are smooth. This is precisely 2.2 bis.

(d') With the notation of 3.8, a point  $\xi$  of  $F$  with values in a field  $k$  over  $S$  is just a multiplicative-type subgroup  $H_0$  of  $G_0 = G_k$ . Repeating the argument in (b') above, we see that the integer  $d(\xi)$  considered in 3.8 is

$$d(\xi) = \dim_k(\mathfrak{g}_0/\mathfrak{g}_0^{H_0}),$$

where  $\mathfrak{g}_0$  is the Lie algebra of  $G_0$  and  $\mathfrak{g}_0^{H_0}$  is the vector subspace of invariants under  $H_0$ . This is a finite integer, so condition (d')<sup>1°</sup> of 3.8 holds.

With the notation of (d')<sup>2°</sup>, giving a morphism  $v: X \rightarrow F$  amounts to giving a multiplicative-type subgroup  $H$  of  $G_X$ . For  $x \in X$ , the integer  $d(\xi_x)$ <sup>16</sup> is then the dimension of

$$(\mathfrak{g} \otimes \kappa(x))/(\mathfrak{g} \otimes \kappa(x))^{H_x}.$$

where  $\mathfrak{g}$  is the sheaf of Lie algebras of  $G_X$ . It is locally free of finite type over  $X$ , because  $G_X$  is smooth over  $X$ , and  $\mathfrak{g} \otimes \kappa(x)$  is the Lie algebra of the fibre  $G_x$  of  $G_X$  at  $x$ ; here  $H_x$  is the fibre of  $H$  at  $x$ . Now  $\mathfrak{g}$  is a locally free module on  $X$ <sup>17</sup> on which the multiplicative-type group  $H$  acts, and the invariant subfunctor  $\mathfrak{g}^H$  is therefore given by a locally direct-summand, hence locally free, subsheaf of  $\mathfrak{g}$ . By descent, one reduces to the case in which  $H$  is diagonalizable, and then applies Exposé I, 4.7.3, noting that the invariant subsheaf corresponds to the degree-zero component. Consequently,

$$d(\xi_x) = \text{rank}_x(\mathfrak{g}/\mathfrak{g}^H),$$

so this is a locally constant function of  $x$ . This proves condition (d').

We have thus verified the conditions of 3.7, completing the proof of 4.1.

**Remark 4.6.** — When  $G = \text{GL}(n)_S$ , one can give a much simpler and more explicit direct proof of 4.1, using Exposé I, 4.7.3. The proof moreover shows that, in this case, the moduli scheme is a disjoint sum of affine  $S$ -schemes. Proceeding as indicated in 4.3, one obtains the same result whenever  $G$  is a closed subgroup of a group of the form  $\text{GL}(n)_S$ .

One must not, however, suppose that the preschemes representing the functors in 4.1 or 4.2 are always disjoint sums of affine  $S$ -schemes. For example, let  $H$  be a multiplicative-type group of finite type over a locally noetherian prescheme  $S$ . By Exposé X, 5.11,  $H$  is isotrivial if and only if

$$\text{Hom}_{S\text{-gr}}(H, \mathbb{G}_m)$$

is a disjoint sum of preschemes affine over  $S$ . But, as noted in Exposé X, 1.6, there may exist nonisotrivial tori  $H$  of relative dimension 2. For such a torus,  $\text{Hom}_{S\text{-gr}}(H, \mathbb{G}_m)$  is not a disjoint sum of  $S$ -preschemes affine over  $S$ , and likewise the “dual” twisted constant group

$$\text{Hom}_{S\text{-gr}}(\mathbb{G}_m, H)$$

<sup>16</sup>The printed source gives only  $(\mathfrak{g} \otimes \kappa(x))^{H_x}$  here and later  $\text{rank}_x \mathfrak{g}^H$ , omitting the quotient. That reading contradicts both the immediately preceding formula  $d(\xi) = \dim(\mathfrak{g}_0/\mathfrak{g}_0^{H_0})$  and the lifting torsor  $N/M$ . The quotient is restored in both places.

<sup>17</sup>The printed source says “over  $S$ ” here, although  $\mathfrak{g}$  is the Lie-algebra sheaf of  $G_X$ ; the base is therefore  $X$ .

is not such a sum. Indeed, if two commutative twisted constant groups  $R, R'$  of finite presentation are dual to one another,

$$R' = \operatorname{Hom}_{S\text{-gr}}(R, \mathbb{Z}_S),$$

then one readily sees that one is isotrivial if and only if the other is.

This last point also shows that such an  $H$  is not isomorphic to a subgroup of a group of the form  $\operatorname{GL}(n)_S$ . More precisely, one can show that a multiplicative-type group of finite type over a connected locally noetherian  $S$  is isomorphic to a subgroup of a group of the form  $\operatorname{Aut}_{\operatorname{Mod}}(E)$ , where  $E$  is a locally free module of finite type over  $S$ , if and only if it is isotrivial. Finally, taking  $G = H \times \mathbb{G}_m$  in the two preceding examples gives an example in which the prescheme representing the functor  $F$  of 4.1 is not a disjoint sum of affine  $S$ -preschemes; if desired,  $G$  may be taken to be a torus of relative dimension 3.

## 5. First Corollaries of the Representability Theorem

Let  $H$  and  $G$  be as in 4.2. Put

$$M = \operatorname{Hom}_{S\text{-gr}}(H, G),$$

which, by 4.2, is a smooth  $S$ -prescheme separated over  $S$ . Notice that  $G$  acts on the functor  $M = \operatorname{Hom}_{S\text{-gr}}(H, G)$  by

$$(g, u) \longmapsto \operatorname{int}(g) \circ u.$$

Hence there is a canonical morphism

$$(x) \quad G \times_S M \longrightarrow M \times_S M,$$

whose components are the preceding morphism  $G \times_S M \rightarrow M$  and the second projection  $G \times_S M \rightarrow M$ .

**Corollary 5.1.** — *The preceding morphism (x) is smooth.*

This follows from 4.2 and 2.3. This statement is equivalent to the following one.

**Corollary 5.2.** — *Let  $u_1, u_2 : H \rightrightarrows G$  be two homomorphisms of  $S$ -groups. Then the subfunctor  $\operatorname{Transp}(u_1, u_2)$  of  $G$  (cf. 2.4) is representable by a closed subprescheme of  $G$ , smooth over  $S$ .*

It remains only to prove that  $\operatorname{Transp}(u_1, u_2) \rightarrow G$  is indeed a closed immersion. This follows because it is the equalizer of a pair of morphisms  $G \rightrightarrows M$ , namely

$$g \longmapsto \operatorname{int}(g)u_1 \quad \text{and} \quad g \longmapsto u_2,$$

the latter being the “constant” morphism, and because  $M$  is separated over  $S$ . In particular:

**Corollary 5.3.** — *Let  $u : H \rightarrow G$  be a morphism of  $S$ -groups. Then*

$$\operatorname{Centr}_G(u) = \operatorname{Transp}(u, u)$$

*is representable by a closed group subprescheme of  $G$ , smooth over  $S$ . Moreover, the quotient of  $G$  by  $\operatorname{Centr}_G(u)$  is representable by an open subprescheme of  $M$ .*

It remains to prove the last assertion. The morphism

$$g \longmapsto \operatorname{int}(g) \circ u$$

from  $G$  to  $M$  is smooth and of finite type by 5.2, hence is an open morphism (EGA IV, 6.6). Let  $U$  denote its image, equipped with the structure induced from  $M$ . The induced morphism  $G \rightarrow U$  is smooth, surjective, and of finite type, and is therefore a covering for the faithfully flat quasi-compact topology. Moreover, it is clear that the preceding morphism  $G \rightarrow M$  makes  $G$  a formally principal homogeneous sheaf under  $\text{Centr}_G(u)_M$ . It follows that the sheaf  $G/\text{Centr}_G(u)$  is represented by  $U$ .

**Corollary 5.4.** — *Let  $u_1, u_2 : H \rightrightarrows G$  be two homomorphisms of  $S$ -groups. The following conditions are equivalent:*

- (i)  $u_1$  and  $u_2$  are conjugate locally for the étale topology.
- (i bis)  $u_1$  and  $u_2$  are conjugate locally for the faithfully flat quasi-compact topology.
- (ii) For every  $s \in S$ , let  $\bar{s}$  denote the spectrum of an algebraic closure of  $\kappa(s)$ . The morphisms
 
$$u_{1,\bar{s}}, u_{2,\bar{s}} : H_{\bar{s}} \rightrightarrows G_{\bar{s}}$$
 are conjugate by an element of  $G(\kappa(\bar{s}))$ .
- (ii bis) The structural morphism  $\text{Transp}(u_1, u_2) \rightarrow S$  is surjective.
- (iii)  $\text{Transp}(u_1, u_2)$  is a torsor under the action of the smooth  $S$ -group prescheme of finite type  $\text{Centr}_G(u_1)$ .

The implications (i)  $\Rightarrow$  (i bis) and (ii)  $\Rightarrow$  (ii bis) are immediate, the latter by the Nullstellensatz. The implication (i bis)  $\Rightarrow$  (ii) follows from the “principle of finite extension” (EGA IV, 9). Moreover, (ii bis)  $\Rightarrow$  (iii): since  $\text{Transp}(u_1, u_2)$  is smooth over  $S$ , it is flat over  $S$ , and since it is of finite type, it is quasi-compact over  $S$ . Thus it is faithfully flat and quasi-compact over  $S$  if and only if its structural morphism is surjective. On the other hand, it is formally principal homogeneous under  $\text{Centr}_G(u_1)$ , which is faithfully flat and quasi-compact over  $S$ . The last condition is therefore also equivalent to saying that  $\text{Transp}(u_1, u_2)$  is a torsor under  $\text{Centr}_G(u_1)$ , in the faithfully flat quasi-compact topology. Finally, (iii)  $\Rightarrow$  (i) follows from the smoothness of  $\text{Transp}(u_1, u_2)$  over  $S$  and from Hensel’s lemma in the form 1.10.

**Remark 5.5.** — Fix  $u_1 = u : H \rightarrow G$ . Consider the functor

$$(\text{Sch})^\circ/S \longrightarrow (\text{Ens})$$

that assigns to every  $T$  over  $S$  the set of homomorphisms of  $T$ -groups  $u_2 : H_T \rightarrow G_T$  that are locally conjugate, for the étale topology, to  $u_T : H_T \rightarrow G_T$ . This functor is represented precisely by the open subscheme of  $M$ , isomorphic to  $G/\text{Centr}_G(u)$ , considered in 5.3.

Let us sketch the variants of the preceding results obtained by applying 4.1 instead of 4.2. Thus let  $G$  be a group prescheme smooth and affine over  $S$ , and now let  $M$  denote the smooth  $S$ -prescheme separated over  $S$  that represents the functor considered in 4.1. Again  $G$  acts on  $M$ :

$$(g, H) \longmapsto \text{int}(g)(H).$$

Hence, as above, there is a morphism

$$(x \text{ bis}) \quad G \times_S M \longrightarrow M \times_S M.$$

**Corollary 5.1 bis.** — *The preceding morphism is smooth.*

This follows from 4.1 and 2.3 bis. We again conclude:

**Corollary 5.2 bis.** — *Let  $H_1, H_2$  be two multiplicative-type subgroups of  $G$ . Then the subfunctor  $\mathrm{Transp}_G(H_1, H_2)$  of  $G$  is representable by a closed subprescheme of  $G$ , smooth over  $S$ .*

In particular:

**Corollary 5.3 bis.** — *Let  $H$  be a multiplicative-type subgroup of  $G$ . Then the group subfunctor  $\mathrm{Norm}_G(H)$  of  $G$  is representable by a closed subprescheme of  $G$ , smooth over  $S$ . Moreover, the quotient  $G/\mathrm{Norm}_G(H)$  is representable by an open subprescheme of  $M$ .*

**Corollary 5.4 bis.** — *Let  $H_1, H_2$  be two multiplicative-type subgroups of  $G$ . The following conditions are equivalent:*

- (i)  $H_1$  and  $H_2$  are conjugate locally for the étale topology.
- (i bis)  $H_1$  and  $H_2$  are conjugate locally for the faithfully flat quasi-compact topology.
- (ii) For every  $s \in S$ , let  $\bar{s}$  denote the spectrum of an algebraic closure of  $\kappa(s)$ . The subgroups  $H_{1, \bar{s}}$  and  $H_{2, \bar{s}}$  of  $G_{\bar{s}}$  are conjugate by an element of  $G(\kappa(\bar{s}))$ .
- (ii bis) The structural morphism  $\mathrm{Transp}(H_1, H_2) \rightarrow S$  is surjective.
- (iii)  $\mathrm{Transp}(H_1, H_2)$  is a principal homogeneous bundle under the action of the smooth  $S$ -group prescheme of finite type  $\mathrm{Norm}_G(H_1)$ .

**Remark 5.5 bis.** — Remark 5.5 carries over to the present case as well.

**Remark 5.6.** — To prove 5.2, and hence also the first assertion of 5.3 and the assertions of 5.4, the reference to 4.2 may be replaced by a reference to VIII, 6.4, whose proof is much easier. This also shows that the hypothesis that  $G$  be affine over  $S$  is unnecessary there. Moreover, in Section 6 we shall show how a variant of this method extends these results to certain groups  $H$  more general than groups of multiplicative type. The same observations apply to the variants 5.2 bis and following. By contrast, the result 5.8 below makes essential use of all the hypotheses imposed—in particular, that  $G$  be affine and smooth over  $S$ , and that  $H$  be of multiplicative type—and the lecturer knows no proof other than one using the representability theorems 4.1 or 4.2.<sup>18</sup>

**5.7.** Since the morphism (x), respectively (x bis), is smooth and hence open, its image is open. Let  $R$  be this image, equipped with the structure induced from  $M \times_S M$ . One checks readily that  $R$  is an equivalence relation on  $M$ , namely the relation made explicit in 5.4, respectively 5.4 bis. It would be interesting to know whether the quotient sheaf  $M/R$ , which is formally étale over  $S$ , is representable; if so, it is represented by a prescheme étale over  $S$ . This is the case, for example, when  $S$  is the spectrum of a field. Notice also that  $R$  need not be closed in  $M \times_S M$ , even when  $G$  is quasi-finite over  $S$ . Thus, when  $M/R$  is representable, it may be nonseparated over  $S$ .

**Theorem 5.8.** — *Let  $S$  be a prescheme, let  $G$  be a smooth affine  $S$ -prescheme, and let  $s \in S$ . Then:*

- (a) *For every multiplicative-type subgroup  $H_0$  of  $G_s$ , there exist an étale morphism  $S' \rightarrow S$ , a point  $s' \in S'$  lying over  $s$  such that the residue-field extension  $\kappa(s')/\kappa(s)$  is trivial, and a multiplicative-type subgroup  $H'$  of  $G' = G \times_S S'$ , such that*

$$H'_{s'} = H_0 \otimes_{\kappa(s)} \kappa(s').$$

<sup>18</sup>The situation has changed since this text was written; cf. Exposés XV and XIX, no. 6.

- (b) For every group homomorphism  $u_0 : H_s \rightarrow G_s$ , where  $H$  is a multiplicative-type  $S$ -group prescheme of finite type, there exist an étale morphism  $S' \rightarrow S$ , a point  $s' \in S'$  lying over  $s$  such that  $\kappa(s')/\kappa(s)$  is trivial, and a group homomorphism

$$u' : H' \longrightarrow G', \quad H' = H \times_S S', \quad G' = G \times_S S',$$

such that

$$u'_{s'} : H'_{s'} \longrightarrow G'_{s'} \quad \text{equals} \quad u_0 \otimes_{\kappa(s)} \kappa(s').$$

This follows from 4.1, respectively 4.2, and from Hensel's lemma in the form 1.10.

**Proposition 5.9.** — *Let  $S$  be a prescheme, let  $G$  be a smooth affine  $S$ -prescheme, and let  $H$  be a multiplicative-type subgroup of  $G$ . Consider*

$$\mathrm{Centr}_G(H) \hookrightarrow \mathrm{Norm}_G(H),$$

*which, by 5.3 and 5.3 bis, are closed group subpreschemes of  $G$ , smooth over  $S$ . The first is then an open and closed subprescheme of the second, and the quotient sheaf*

$$W_G(H) = \mathrm{Norm}_G(H) / \mathrm{Centr}_G(H)$$

*is represented by an open group subprescheme of  $\mathrm{Aut}_{S\text{-gr}}(H)$ . It is therefore a quasi-finite, étale, and separated  $S$ -group prescheme.*

Indeed, consider the evident homomorphism

$$\theta : \mathrm{Norm}_G(H) \longrightarrow \mathrm{Aut}_{S\text{-gr}}(H),$$

whose kernel is, by definition,  $\mathrm{Centr}_G(H)$ . Since  $\mathrm{Aut}_{S\text{-gr}}(H)$  is represented by an étale  $S$ -group prescheme separated over  $S$  (X, 5.10), its unit section is an open and closed immersion. Its inverse image under  $\theta$  is therefore an open and closed subgroup of  $\mathrm{Norm}_G(H)$ .

Moreover,  $\theta$  is smooth. This follows formally from the definitions and from the facts that  $\mathrm{Norm}_G(H)$  is smooth over  $S$  and that  $\mathrm{Aut}_{S\text{-gr}}(H)$  is étale over  $S$ . As in 5.3, the image of  $\theta$  is consequently an open subprescheme  $U$  of  $\mathrm{Aut}_{S\text{-gr}}(H)$ , and, with the induced structure,  $U$  represents the quotient sheaf  $\mathrm{Norm}_G(H) / \mathrm{Centr}_G(H)$ . The latter is therefore étale and separated over  $S$ . It is also quasi-finite over  $S$ , since it is quasi-compact over  $S$ , being the image of  $\mathrm{Norm}_G(H)$ , which is quasi-compact. This proves 5.9.

**Corollary 5.10.** — *For every  $s \in S$ , put*

$$w(s) = \mathrm{rank}(\mathrm{Norm}_{G_s}(H_s) / \mathrm{Centr}_{G_s}(H_s)).$$

*This is also the index of  $\mathrm{Centr}_{G(\bar{k})}(H(\bar{k}))$  in  $\mathrm{Norm}_{G(\bar{k})}(H(\bar{k}))$ , where  $\bar{k}$  is an algebraic closure of  $\kappa(s)$ . Then the function  $s \mapsto w(s)$  is lower semicontinuous. It is constant in a neighborhood of  $s$  if and only if  $W_G(H)$  is finite over  $S$  in a neighborhood of  $s$ .*

Indeed, for every  $S$ -prescheme  $W$  that is étale, of finite type, and separated over  $S$ , the function

$$s \longmapsto \mathrm{rank}[W_s : \kappa(s)]$$

is lower semicontinuous, and it is constant in a neighborhood of  $s$  if and only if  $W$  is finite over  $S$  in a neighborhood of  $s$ . This fact was noted in SGA 1, I, 10.9; its proof, which presents no difficulty, is given in EGA IV.<sup>19</sup>

<sup>19</sup>EGA IV, 15.5.1 and 18.10.7.

**Remark 5.11.**<sup>20</sup> — Let  $G$  be a group prescheme affine and smooth over  $S$ , and let  $H$  be a multiplicative-type subgroup. Then 5.3 and 5.3 bis imply that the quotients

$$G/\mathrm{Centr}_G(H) \quad \text{and} \quad G/\mathrm{Norm}_G(H)$$

are preschemes smooth and quasi-affine over  $S$ . Indeed, in both cases the moduli prescheme  $M$  in which the quotient is embedded has the property that every open  $U \subset M$  quasi-compact over  $S$  is quasi-affine over  $S$ , by 3.11.

We shall moreover see in the next Exposé<sup>21</sup> that for every such  $U$ , the schematic closure  $\overline{U}$  of  $U$  in  $M$  is even affine over  $S$ . Thus the quotients under consideration are affine over  $S$ , provided that the open subprescheme  $U$  of the corresponding moduli prescheme—of group homomorphisms from  $H$  to  $G$ , respectively of multiplicative-type subgroups of  $G$ —is closed in  $M$ . This is the case, for example, when  $H$  is a “maximal torus” in the second case, or when  $S$  is the spectrum of a field; cf. XII.5.4.<sup>22</sup> I do not know whether the quotients under consideration are affine over  $S$  in general.

## 6. A Rigidity Property for Homomorphisms of Certain Group Schemes, and the Representability of Certain Transporters<sup>23</sup>

**Proposition 6.1.** — *Let  $S$  be a locally noetherian prescheme, let  $H$  be a commutative  $S$ -group prescheme of finite type over  $S$ , and let  $E$  be a set of positive integers stable under multiplication. Suppose that the following conditions hold:*

(a) *for every  $n \in E$ , the subgroup*

$${}_nH = \mathrm{Ker}(n \cdot \mathrm{id}_H)$$

*is finite and flat over  $S$ ;*

(b) *for every  $s \in S$ , the family of the  ${}_nH_s$ ,  $n \in E$ , is schematically dense in the fibre  $H_s$ .*<sup>24</sup>

*Let  $Y$  be a closed subprescheme of  $H$ . Then the subfunctor  $\prod_{H/S}(Y/H)$  of  $S$  (compare Exposé VIII, 6) is representable by a closed subprescheme  $T$  of  $S$ . If  $S$  is noetherian, there exists  $n \in E$  such that<sup>25</sup>*

$$(x) \quad \prod_{H/S}(Y/H) = \prod_{{}_nH/S}((Y \cap {}_nH)/{}_nH).$$

<sup>20</sup>Editor’s note: For a generalization to the nonaffine case, see M. Raynaud, *Faisceaux amples sur les schémas en groupes et les espaces homogènes*, Lecture Notes in Mathematics 119 (1970), IX.2.8.

<sup>21</sup>Editor’s note: Cf. Remark XII.5.7. We mention here the following generalization. Without assuming  $G$  affine, if  $H$  is a torus and  $S$  is locally noetherian,  $U$  is affine and smooth by a theorem of Raynaud (*loc. cit.*, IX.2.6 and IX.2.8). Indeed,  $G/\mathrm{Norm}_G(H)$  is an open subprescheme of  $M$ , and the largest representable open subfunctor of  $M$  is a disjoint sum of open subpreschemes smooth and affine over  $S$ .

<sup>22</sup>Editor’s note: This assertion assumes that  $G$  has locally constant reductive rank.

<sup>23</sup>The present section does not use the results of Sections 3, 4, and 5; its natural place would be in Exposé VI<sub>B</sub>.

<sup>24</sup>The printed French text has  $H$  here; the fibrewise condition and its use in the proof require  $H_s$ .

<sup>25</sup>The printed formula writes the right-hand quotient as  $(Y \cap {}_nH)/S$ . Since the product functor is taken over  ${}_nH/S$ , and as the proof requires, the quotient is  $(Y \cap {}_nH)/{}_nH$ .



Indeed, by Exposé VIII, 6.4, condition (a) implies that  ${}_nH$  is finite and flat, and hence a fortiori “essentially free” over  $S$ . Thus the right-hand side of (x) is representable by a closed subprescheme  $T_n$  of  $S$ .<sup>26</sup> Order  $E$  by divisibility. The  $T_n$  then form a decreasing family of closed subpreschemes of  $S$ ; consequently, when  $S$  is noetherian—as we may now suppose—this family is stationary for sufficiently large  $n$ . Let  $T$  be the stable value of  $T_n$ . We claim that  $T$  represents the left-hand side of (x), which will complete the proof.

It is enough to prove the following. Let  $S'$  be a prescheme over  $S$  such that

$$(Y \cap {}_nH)_{S'} = ({}_nH)_{S'} \quad \text{for every } n \in E,$$

or, equivalently, such that  $Y_{S'} \supset ({}_nH)_{S'}$  for every  $n \in E$ . Then  $Y_{S'} = H_{S'}$ . This follows at once from Exposé IX, 4.4: in view of conditions (a) and (b), the family of the  $({}_nH)_{S'}$  is schematically dense in  $H_{S'}$ .

**Theorem 6.2.** — *Let  $S$  be a prescheme, let  $H, E$  be as in 6.1 and satisfy conditions (a) and (b), let  $u : H \rightarrow G$  be a homomorphism of  $S$ -groups from  $H$  to an  $S$ -group prescheme locally of finite type over  $S$ , and let  $K$  be a closed subgroup prescheme of  $G$ .<sup>27</sup> Consider the subfunctor  $\text{Transp}_G(u, K)$  of  $G$  (cf. 2.5). It is representable by a closed subprescheme of  $G$ . If  $G$  is noetherian—for example, if  $S$  is noetherian and  $G$  is of finite type over  $S$ —there exists  $n \in E$  such that*

$$\text{Transp}_G(u, K) = \text{Transp}_G(u|_{{}_nH}, K).$$

*Suppose, finally, that  $G$  is smooth over  $S$ , that  $K$  is smooth over  $S$  or is of multiplicative type, and that  $H, E$  satisfy the following condition, stronger than (a):*

*a' for every  $n \in E$ , the subgroup  ${}_nH$  of  $H$  is of multiplicative type.*

*Then  $\text{Transp}_G(u, K)$  is smooth over  $S$ .*

Consider the  $G$ -group

$$H_G = H \times_S G,$$

which plainly satisfies the conditions of 6.1, with  $S$  replaced by  $G$  and  $H$  by  $H_G$ . Let  $Y$  be the closed subprescheme of  $H_G$  that is the inverse image of  $K \subset G$  under the map

$$H_G = H \times_S G \longrightarrow G, \quad (h, g) \longmapsto \text{int}(g) \cdot u(h).$$

Then

$$\text{Transp}_G(u, K) = \coprod_{H_G/G} (Y/H_G)$$

(compare Exposé VIII, Example 6.5(e)). The first assertions therefore follow from 6.1. Moreover, for every quasi-compact open  $U \subset G$ , there exists  $n \in E$  for which the restrictions to  $U$  of the two functors

$$\text{Transp}_G(u, K) \quad \text{and} \quad \text{Transp}_G(u|_{{}_nH}, K)$$

coincide. To verify the last assertion of 6.2, we may therefore replace  $H$  by some  ${}_nH$ , after which it is enough to apply 2.5, since  ${}_nH$  is assumed to be of multiplicative type over  $S$ .

**Remark 6.3.** — The preceding proof uses only the very elementary result of Exposé VIII, 6.4, together, for its last part, with 2.5; when  $K$  is smooth over  $S$ , the latter is

<sup>26</sup>The printed French text has “ $T_n$  of  $T$ ” here; the next sentence makes the  $T_n$  closed subpreschemes of  $S$ , and  $T$  is defined only afterward as their stable value.

<sup>27</sup>The printed French text has  $H$  here; the proof immediately uses  $K \subset G$ , as required by the transporter.

the infinitesimal result of Exposé IX, 3.6, and hence the vanishing of the cohomology of multiplicative-type groups. In the most important cases (cf. 6.7), the integers  $n \in E$  may even be taken prime to the residue characteristics of  $S$ , that is, invertible in  $\mathcal{O}_S$ . The  ${}_nH$  are then finite étale over  $S$ , and the cohomological result invoked is practically trivial. Thus, in that situation, 6.2 is independent of the theory of multiplicative-type groups.

**6.4.** — As usual, one sees that 6.1 extends to the following situation. Let  $S'$  be an  $S$ -prescheme (not necessarily locally noetherian), put  $H' = H_{S'}$ , and let  $Y'$  be a closed subprescheme of  $H'$ , provided that  $Y' \rightarrow H'$  is of finite presentation, that is, that the ideal defining  $Y'$  is of finite type. Then  $\prod_{H'/S'}(Y'/H')$  is representable by a closed subprescheme  $T'$  of  $S'$ , the morphism  $T' \rightarrow S'$  is of finite presentation, and, if  $S'$  is quasi-compact, there exists  $n \in E$  for which the analogue of (x) holds. It follows as well that the first assertion of 6.2 remains valid without assuming that  $G$  is locally of finite type over  $S$ , provided that the immersion  $K \rightarrow G$  is of finite presentation.

**6.5.** — As announced in 5.6, Theorem 6.2 makes it possible to extend to group preschemes satisfying (a') and (b) above certain results proved by another method, under more restrictive hypotheses, for groups of multiplicative type. This applies to 5.2, the beginning of 5.3, and 5.4 and their corresponding “bis” variants, and also to 5.9 and 5.10. It also applies to the results of Exposé IX, Sections 5 and 6, with the exception of Exposé IX, 6.8. The latter assertion is already false for a homomorphism of abelian schemes  $u : H \rightarrow G$  over the spectrum  $S$  of a discrete valuation ring of residue characteristic  $p > 0$ : it can happen that  $\text{Ker } u$  has the trivial group as its generic fibre and, as its special fibre, a radicial group that is not reduced to the trivial group.

**6.6.** — We have just given an example of a rigidity property of multiplicative-type groups that is not shared by abelian schemes. Another, extremely important, example is that the existence theorem for infinitesimal extensions of homomorphisms in Exposé IX, 3.6 no longer holds when  $H$  is an abelian scheme. Thus a nonzero abelian scheme over a field admits nontrivial infinitesimal variations, in contrast to a multiplicative-type group. This is the infinitesimal aspect of the fact that there is a theory of moduli—one that, moreover, is far from complete—for abelian varieties, whereas the theory of moduli of multiplicative-type groups is empty.

Another, “global,” aspect of this infinitesimal difference is the following. Let  $H$  be an abelian scheme over a locally noetherian prescheme  $S$ , and let  $G$  be a commutative  $S$ -group prescheme locally of finite type over  $S$ . One can show that  $\text{Hom}_{S\text{-gr}}(H, G)$  is representable by a prescheme locally of finite type over  $S$ . Unlike the corresponding prescheme when  $H$  is of multiplicative type, however, it is not étale over  $S$ , but only unramified. Thus, for example, if  $S$  is the spectrum of a complete discrete valuation ring and  $H, G$  are abelian schemes over  $S$ , there may exist homomorphisms  $H_0 \rightarrow G_0$  on the special fibres that do not arise “by specialization” from a homomorphism on the generic fibres.

**6.7.** — Theorem 6.2 applies whenever  $H$  is an abelian scheme over  $S$ , or, more generally, an extension of such a scheme by a torus. Indeed, the question is local on  $S$ , so one may suppose that there is a prime number  $\ell$  prime to the residue characteristics of  $S$ . Taking  $E$  to be the set of powers of  $\ell$  then suffices to satisfy conditions (a') and (b).

An argument analogous to that of 6.1 gives the following result.

**Theorem 6.8.** — *Let  $X$  be a prescheme smooth over  $S$ , with nonempty connected geometric fibres. For every closed subprescheme  $Y$  of  $X$ , the functor  $\prod_{X/S}(Y/X)$  is representable by a closed subprescheme  $T$  of  $S$ . If  $Y$  is of finite presentation over  $X$ , then  $T$  is of finite presentation over  $S$ .*

Since  $f : X \rightarrow S$  is faithfully flat and locally of finite presentation, it is a covering for the fpqc topology. On the other hand,  $T = \prod_{X/S}(Y/X)$  is plainly a subsheaf of  $S$  for the fpqc topology. Hence both the representability of  $T$  by a closed subscheme of  $S$  and the question whether  $T$  is of finite presentation over  $S$  are local on  $S$  for the fpqc topology. After the base change  $S' \rightarrow S$  with  $S' = X$ , we are reduced to the case in which  $X$  admits a section  $e$  over  $S$ . We may also suppose that  $S$  is affine, and hence in particular quasi-compact. We then have:

**Corollary 6.9.** — *Under the hypotheses of 6.8, suppose that  $S$  is quasi-compact and that  $X$  admits a section  $e$  over  $S$ . For every integer  $n \geq 0$ , let  $X_n$  be the order- $n$  infinitesimal neighbourhood of the section  $e$  in  $X$ . Suppose that  $Y$  is of finite presentation over  $X$ . Then there exists an integer  $n \geq 0$  such that*

$$\prod_{X/S}(Y/X) = \prod_{X_n/S}(Y_n/X_n), \quad Y_n = Y \cap X_n.$$

This corollary implies 6.8 when  $Y$  is of finite presentation over  $X$ , by Exposé VIII, 6.4. Indeed, since  $X$  is smooth over  $S$ ,  $X_n$  is finite and locally free over  $S$ , and a fortiori “essentially free” over  $S$  in the sense of Exposé VIII, 6.1. Thus  $\prod_{X_n/S}(Y_n/X_n)$  is representable by a closed subscheme of  $S$ . Furthermore, either the proof of the cited result or reduction to the noetherian case shows at once that this closed subscheme is of finite presentation over  $S$ .

Let us first prove 6.9, and hence 6.8, when  $S$  is noetherian. Put

$$T_n = \prod_{X_n/S}(Y_n/X_n).$$

The  $T_n$  form a decreasing sequence of closed subschemes of  $S$ , and this sequence is stationary because  $S$  is noetherian. Let

$$R = \bigcap_{n \geq 0} \prod_{X_n/S}(Y_n/X_n)$$

be their common value for large  $n$ . Plainly  $T \subset R$ , and it remains to show that  $R \subset T$ . After the base change  $R \rightarrow S$ , we are reduced to the case  $R = S$ , that is,  $Y_n = X_n$  for every  $n$ , or equivalently  $Y \supset X_n$  for every  $n$ . We must then prove  $T = S$ , that is,  $Y = X$ .

The inclusions  $Y \supset X_n$  for all  $n$ , together with the fact that  $X$  is locally noetherian, imply that in a neighbourhood of each point of  $e(S)$ ,  $Y$  is an induced open subscheme of  $X$ . Hence there is an induced open subscheme  $U$  of  $X$  such that  $e(S) \subset U \subset Y$ . By Exposé IX, 4.3, since the fibres of  $X/S$  are integral,  $U$  is schematically dense in  $X$ . As  $Y$  is a closed subscheme containing  $U$ , it follows that  $Y = X$ . This proves 6.9, and hence 6.8, in the noetherian case.

The general case is reduced to the preceding one. For every  $s \in S$ , there is an affine open neighbourhood  $U$  of  $s$  and an affine open neighbourhood  $V$  of  $e(s)$  such that  $f(V) \subset U$ . Then  $f(V)$  is an open neighbourhood of  $s$  contained in  $U$ . Choose an affine open neighbourhood  $S'$  of  $s$  contained in  $e^{-1}(V) \cap f(V)$ , and put  $X' = V \cap f^{-1}(S')$ . Then  $X'$  and  $S'$  are affine open subschemes of  $X$  and  $S$ , respectively, and  $X'/S'$  admits a section. By the local nature of 6.8 and 6.9, we may suppose that  $S = S'$ .

We claim that

$$\prod_{X/S}(Y/X) = \prod_{X'/S'}(Y'/X'), \quad Y' = Y \cap X'.$$

Indeed, by Exposé IX, 4.6,  $X'$  is schematically dense in  $X$ , at least when  $X$  is quasi-separated over  $S$ , so that  $X'$  is retrocompact in  $X$ . In fact, one can readily prove that Exposé IX, 4.6 remains valid without the retrocompactness hypothesis. Likewise, after every base change  $S_1 \rightarrow S$ ,  $X' \times_S S_1$  is schematically dense in  $X \times_S S_1$ , and the claimed equality follows immediately. We are thus reduced to the case  $X = X'$ , so we may suppose that both  $S$  and  $X$  are affine.

Now let  $X = \operatorname{Spec}(B)$ , and let  $J$  be the ideal of  $B$  defining  $Y$ . The ideal  $J$  is the filtered colimit of its finite-type subideals. Consequently,  $Y$  is the intersection of closed subpreschemes  $Y_i$  of  $X$  that are of finite presentation over  $S$ , and

$$\prod_{X/S} (Y/X) = \bigcap_i \prod_{X/S} (Y_i/X).$$

To prove 6.8, it is therefore enough to treat the case in which  $Y$  is of finite presentation over  $X$ . It remains to prove 6.9 with  $S$  and  $X$  affine. But  $X$  and  $Y$  over  $S$  then arise by base change  $S \rightarrow S_0$  from an analogous situation  $X_0, Y_0$  over a noetherian prescheme  $S_0$ . This reduces us to the already treated case in which  $S$  is noetherian, and completes the proof of 6.8 and 6.9.

**Corollary 6.10.** — *Let  $X$  be an  $S$ -group prescheme, smooth and of finite presentation, with connected fibres. Let  $Y$  be a group prescheme of finite presentation over  $S$ , and let  $i : Y \rightarrow X$  be a monomorphism of  $S$ -group preschemes, so that  $Y$  is a subgroup of  $X$ . Then  $\prod_{X/S} (Y/X)$  is representable by a closed subprescheme of  $S$  of finite presentation. If  $S$  is quasi-compact, let  $X_n$  denote, for every  $n \geq 0$ , the order- $n$  infinitesimal neighbourhood of the unit section of  $X$ , and put  $Y_n = X_n \cap Y$ . Then, for all sufficiently large  $n$ ,*

$$\prod_{X/S} (Y/X) = \prod_{X_n/S} (Y_n/X_n).$$

The proof is essentially that of 6.9. The unit sections of  $X$  and  $Y$  induce bijective immersions  $S \rightarrow X_n$  and  $S \rightarrow Y_n$ , and hence isomorphisms

$$S_{\text{red}} \xrightarrow{\sim} (X_n)_{\text{red}}, \quad S_{\text{red}} \xrightarrow{\sim} (Y_n)_{\text{red}}.$$

It follows that the injection  $Y_n \rightarrow X_n$  is proper. Since it is a monomorphism of finite presentation, it is therefore a closed immersion. Exposé VIII, 6.4, already used above, now shows that  $\prod_{X_n/S} (Y_n/X_n)$  is representable by a closed subprescheme of  $S$  of finite presentation over  $S$ . It remains to prove the last assertion of 6.10 under the additional assumption that  $S$  is affine.

We reduce once more to the case in which  $S$  is noetherian. With the notation of the proof of 6.9, it is enough to prove  $R = T$ , or, equivalently, that  $Y \supset X_n$  for every  $n$  implies  $Y = X$ . This hypothesis implies that  $i : Y \rightarrow X$  is étale at the points of the unit section of  $Y$  over  $S$ , and therefore that  $Y$  is smooth over  $S$  at those points. Hence the open subprescheme  $Y'$  consisting of the points where  $Y$  is smooth over  $S$  is an induced open subgroup of  $Y$ . By Exposé X, 3.5, the morphism  $Y' \rightarrow X$  is an étale monomorphism and hence an open immersion. Since the fibres of  $X$  are connected and every open subgroup of an algebraic group is also closed, this open immersion is surjective, and therefore an isomorphism. Thus  $Y' = X$ , and a fortiori  $Y = X$ , completing the proof of 6.10.

Arguing as in Exposé VIII, 6.5, we deduce:

**Corollary 6.11.** — *Let  $G$  be an  $S$ -group prescheme locally of finite type and quasi-separated over  $S$ , let  $H$  be a group prescheme smooth and of finite presentation over  $S$ , with connected fibres, and let  $i : H \rightarrow G$  be a monomorphism of  $S$ -groups, so that  $H$  is a subgroup of  $G$ . Then:*

- (a)  $\text{Centr}_G(H)$  and  $\text{Norm}_G(H)$  are representable by closed subpreschemes of  $G$  of finite presentation over  $G$ . Likewise, for every monomorphism  $j : K \rightarrow G$  of finite presentation between  $S$ -group preschemes,  $\text{Transp}_G(H, K)$  is representable by a closed subprescheme of  $G$  of finite presentation over  $G$ .
- (b) If  $G$  is quasi-compact, then in each of the cases in (a) there exists an integer  $n \geq 0$  such that, if  $H_n$  denotes the order- $n$  infinitesimal neighbourhood of the unit section of  $H$  (and  $K_n$  is defined similarly in the transporter case),

$$\begin{aligned}\text{Centr}_G(H) &= \text{Centr}_G(H_n), \\ \text{Norm}_G(H) &= \text{Norm}_G(H_n), \\ \text{Transp}_G(H, K) &= \text{Transp}_G(H_n, K) = \text{Transp}_G(H_n, K_n).\end{aligned}$$

Apply 6.10 over the base prescheme  $G$  to the group prescheme

$$X = H_G = H \times_S G.$$

For the centralizer, take as  $Y$  the inverse image of the diagonal subgroup of  $(G \times_S G)_G$  under a suitable homomorphism of  $G$ -groups from  $X$  to  $(G \times_S G)_G$ . For the transporter, take the inverse image of  $K_G$  under a suitable homomorphism of  $G$ -groups from  $X$  to  $G_G$ . The normalizer case reduces to the transporter case by taking  $K = H$ . The hypothesis on  $G$  ensures that  $H \rightarrow G$  is of finite presentation, and hence that  $Y \rightarrow X$  is of finite presentation. In the centralizer case, the same hypothesis ensures that the diagonal subgroup of  $G \times_S G$  is of finite presentation over  $G \times_S G$ ; again it follows that  $Y$  is of finite presentation over  $X$ .

**Remark 6.12.** — One can prove, using rather delicate results of EGA VI,<sup>28</sup> that if  $X$  is a prescheme flat and of finite presentation over  $S$ , and is either proper over  $S$  or has nonempty connected geometric fibres, then for every closed subprescheme  $Y$  of  $X$  of finite presentation over  $S$ , the functor  $\prod_{X/S}(Y/X)$  is representable by a closed subprescheme of  $S$  of finite presentation over  $S$ . Likewise, if  $X$  is an  $S$ -group prescheme flat and of finite presentation with connected fibres, and  $i : Y \rightarrow X$  is a monomorphism of  $S$ -groups, with  $Y$  an  $S$ -group prescheme of finite presentation, then  $\prod_{X/S}(Y/X)$  is representable by a closed subprescheme of  $S$  of finite presentation over  $S$ . In particular, 6.11(a) remains valid if the hypothesis “ $H$  smooth over  $S$ ” is replaced by “ $H$  flat over  $S$ .”

---

<sup>28</sup>Cf. also J. P. Murre, “Representation of unramified functors. Applications,” *Séminaire Bourbaki*, no. 294 (May 1965), Theorem 3, p. 13.

# Exposé XII

**This exposé is not included in this current-progress reader.**

No complete Exposé XII English reader was available to this integration.  
The next included scope is the complete working Exposé XIII translation.

# Exposé XIII

## Regular Elements of Algebraic Groups and Lie Algebras

by A. Grothendieck

### 0.10 An auxiliary lemma on varieties with operators

Let  $S$  be a prescheme, let  $G$  be an  $S$ -group prescheme acting on the left on an  $S$ -prescheme  $V$ , let  $W$  be a closed  $S$ -subprescheme of  $V$ , and let  $N$  be its stabilizer in  $G$ . Thus  $N$  is the subgroup of  $G$  whose points with values in an  $S'$  over  $S$  are the elements  $g \in G(S')$  such that

$$g \cdot W_{S'} = W_{S'}.$$

We endow  $(\text{Sch}/S)$  with the faithfully flat quasi-compact topology and identify  $G$ ,  $V$ , and  $W$  with the corresponding sheaves.<sup>29</sup> We shall therefore reason in the category of sheaves on  $(\text{Sch}/S)$ ; throughout this section, “locally” refers to the topology just specified on  $(\text{Sch}/S)$ .

Since  $N$  is a sheaf, consider the quotient sheaf  $G/N$ . It is immediately seen to be isomorphic to the following functor: to every  $S'$  over  $S$ , it assigns the set of subsheaves  $W'$  of  $V_{S'}$  that are locally conjugate to  $W_{S'}$  under  $G$ . Let  $X$  be the subsheaf of  $G/N \times_S V$  whose value on an  $S'$  over  $S$  is the set of pairs  $(W', v)$ , where  $W'$  is as above and  $v$  is a section of  $W'$  over  $S'$  (hence also a section of  $V_{S'}$  over  $S'$ ). Let  $Z$  be the inverse image of  $X$  in  $G \times_S V$ . We then have the cartesian diagram

$$(x) \quad \begin{array}{ccc} Z & \longrightarrow & X \\ \downarrow & & \downarrow i \\ G \times_S V & \longrightarrow & G/N \times_S V, \end{array}$$

where  $i$  is the canonical immersion, while the lower horizontal arrow is induced by the canonical morphism  $G \rightarrow G/N$ .

This morphism sends  $g \in G(S')$  to the subsheaf  $g \cdot W_{S'}$  of  $V_{S'}$ . Consequently,  $Z(S')$  is the set of pairs  $(g, v) \in G(S') \times V(S')$  such that  $v \in (g \cdot W_{S'})(S')$ . It follows that  $Z$  is isomorphic to the sheaf  $G \times_S W$ , via the isomorphism

$$G \times_S W \xrightarrow{\sim} Z, \quad (g, w) \longmapsto (g, g \cdot w).$$

<sup>29</sup>The French source prints  $M$  in this list; the surrounding sentence and all subsequent uses require  $W$ .

Thus the preceding cartesian diagram yields the cartesian diagram

$$(xx) \quad \begin{array}{ccc} G \times_S W & \xrightarrow{q} & X \\ \lambda \downarrow & & \downarrow i \\ G \times_S V & \longrightarrow & G/N \times_S V \end{array}$$

where

$$\lambda(g, w) = (g, g \cdot w), \quad q(g, w) = (\bar{g}, g \cdot w),$$

where  $\bar{g}$  denotes the image of  $g$  under the canonical map  $G(S') \rightarrow (G/N)(S')$ .

Diagram (x) also shows that  $Z \rightarrow X$  makes  $Z$  a principal bundle over  $X$ , with  $N$  acting on the right by

$$(g, v) \cdot n = (gn, v).$$

Accordingly, in diagram (xx), the morphism  $q : G \times_S W \rightarrow X$  makes  $G \times_S W$  a principal bundle over  $X$ , with right  $N$ -action

$$(g, w) \cdot n = (gn, n^{-1} \cdot w).$$

The principal morphisms just introduced are summarized by

$$(D) \quad \begin{array}{ccccc} & & G \times_S W & & \\ & & \downarrow q & \searrow \varphi & \\ G/N & \xleftarrow{p} & X & \xrightarrow{\psi} & V \\ & \swarrow \text{pr}_1 & \downarrow i & \searrow \text{pr}_2 & \\ & & G/N \times_S V & & \end{array}$$

where

$$\psi = \text{pr}_2 \circ i, \quad \varphi = \psi \circ q, \quad \varphi(g, w) = g \cdot w.$$

If  $v$  is a section of  $V$  over  $S$ , let  $X_v$  be its inverse image in  $X$  under  $\psi$ . For every  $S'$  over  $S$ ,  $X_v(S')$  is the set of subsheaves  $W'$  of  $V_{S'}$  that are locally conjugate to  $W_{S'}$  under  $G$  and contain the section  $v_{S'}$  of  $V_{S'}$ . The inverse image of  $v$  in  $G \times_S W$  under  $\varphi$  is isomorphic to the subsheaf  $M_v$  of  $G$  defined by

$$M_v(S') = \{g \in G(S') \mid v_{S'} \in (g \cdot W_{S'})(S')\} = \{g \in G(S') \mid g^{-1}v_{S'} \in W(S')\}.$$

If  $v$  is a section of  $W$ , and not merely of  $V$ , then  $M_v$  evidently contains  $N$ .

None of these explicit descriptions used the representability of  $G$ ,  $V$ , or  $W$  (nor even the fact that the site in question is defined in terms of preschemes). Suppose now, however, that  $N$  is representable, faithfully flat, and quasi-compact over  $S$ , and that  $G/N$  is representable. When  $S$  is the spectrum of a field and  $G$  is of finite type over that field, this hypothesis is necessarily satisfied (VIII 6 and VI<sub>B</sub>.11.18). Using the cartesian diagram (xx), faithfully flat quasi-compact descent, and the fact that  $Z \rightarrow G \times_S V$  is a closed immersion, one sees that  $X$  is representable: it is obtained by descending the closed subprescheme  $Z$  of  $G \times_S V$  along the faithfully flat quasi-compact morphism

$$G \times_S V \longrightarrow G/N \times_S V.$$



Thus diagram (D) is a diagram of morphisms of preschemes over  $S$ .

We henceforth assume that  $S$  is the spectrum of a field  $k$ , and that  $G$ ,  $V$ , and  $W$  are of finite type over  $k$ . Let  $\mathfrak{n}$  be the Lie algebra of  $N$ . Then  $\dim N \leq \operatorname{rank}_k \mathfrak{n}$ , with equality if and only if  $N$  is smooth over  $k$  (Exposé VI<sup>30</sup>). Let  $a \in W(k)$ , and consider the subscheme  $M_a$  of  $G$  defined above; it contains  $N$  and is isomorphic to  $\varphi^{-1}(a)$ . We denote by  $\mathfrak{m}_a$  its Zariski tangent space at the identity element  $e$  of  $N$ . Thus<sup>31</sup>

$$(1) \quad \mathfrak{n} \subset \mathfrak{m}_a, \quad \dim N \leq \operatorname{rank}_k \mathfrak{n} \leq \operatorname{rank}_k \mathfrak{m}_a.$$

**Lemma 1.1.** — *With the preceding notation:*

a) Consider the following conditions:

- (i)  $\mathfrak{n} = \mathfrak{m}_a$ , and  $N$  is smooth over  $k$ .
- (i bis)  $\dim N = \operatorname{rank}_k \mathfrak{m}_a$ .
- (ii) The morphism  $\psi : X \rightarrow V$  is unramified at  $(\bar{e}, a)$ .
- (iii)  $M_a$  and  $N$  coincide in a neighborhood of  $e$ .

Then

$$(i) \iff (i \text{ bis}) \implies (ii) \iff (iii).$$

b) Suppose that  $\varphi : G \times_S W \rightarrow V$  is smooth at  $(e, a)$ . Then  $M_a$  is smooth over  $k$  at  $e$ , and  $\psi$  is smooth at  $(\bar{e}, a)$ .

*Proof.* a) The equivalence of (i) and (i bis) follows immediately from (1) and from the fact noted above that  $N$  is smooth over  $k$  if and only if  $\dim N = \operatorname{rank}_k \mathfrak{n}$ . Consider the inclusion morphism  $N \rightarrow M_a$ . It is well known<sup>32</sup> that if  $N$  is smooth over  $k$  at  $e$  and the tangent map at  $e$  is surjective, then  $N \rightarrow M_a$  is smooth at  $e$ ; since it is an immersion, it is therefore an isomorphism at  $e$ . This proves that (i) implies (iii).

To prove the equivalence of (ii) and (iii), let  $X_a = \psi^{-1}(a)$ , as above. From the isomorphisms

$$M_a \simeq \varphi^{-1}(a) \simeq q^{-1}(X_a)$$

we obtain a morphism  $p_a : M_a \rightarrow X_a$  making  $M_a$  a principal homogeneous bundle with group  $N_{X_a}$ . Consider the diagram

$$\begin{array}{ccc} M_a & \xleftarrow{j_a} & N \\ p_a \downarrow & & \downarrow \\ X_a & \xrightarrow{j'_a} & S = \operatorname{Spec}(k). \end{array}$$

Here  $\operatorname{Spec}(k) \rightarrow X_a$ <sup>33</sup> is defined by the point  $\bar{a} = (\bar{e}, a)$  of  $X$ , and  $j_a : N \rightarrow M_a$  is the canonical immersion. To say that  $\psi$  is unramified at  $(\bar{e}, a)$  means that  $j'_a$  is an open

<sup>30</sup>Editor's note in the French source: since this reference could not be identified, see Theorem II.5.2.1 of M. Demazure and P. Gabriel, *Groupes algébriques I*, Masson (1970).

<sup>31</sup>The French source omits  $\operatorname{rank}_k$  before  $\mathfrak{m}_a$  in (1); it is restored here, as required by the surrounding text and Lemma 1.1.

<sup>32</sup>Editor's note in the French source: see, for example, EGA IV<sub>4</sub>, Theorem 17.11.1(d).

<sup>33</sup>The printed diagram directs  $j'_a : X_a \rightarrow S$ , whereas the immediately following prose defines the point morphism  $S \rightarrow X_a$  and calls it  $j'_a$ . Both the printed diagram direction and the verbal definition are retained here. The source also prints  $X_n$  here and  $X_{a,b}$  in the two local rings below. They are corrected to  $X_a$  and  $X_{a,\bar{a}}$ , in accordance with the fiber notation and the point  $\bar{a}$ .

immersion, or equivalently that it induces an isomorphism

$$S \xrightarrow{\sim} \operatorname{Spec}(\mathcal{O}_{X_a, \bar{a}}).$$

Since  $p_a$  is flat, this is equivalent to saying that the morphism obtained from the preceding one by the base change

$$\operatorname{Spec}(\mathcal{O}_{M_a, e}) \longrightarrow \operatorname{Spec}(\mathcal{O}_{X_a, \bar{a}})$$

is an isomorphism. The resulting morphism is precisely

$$\operatorname{Spec}(\mathcal{O}_{N, e}) \longrightarrow \operatorname{Spec}(\mathcal{O}_{M_a, e}),$$

which proves the equivalence of (ii) and (iii).

Notice moreover that the proof shows that (ii) and (iii) imply the following condition, apparently stronger than (iii):

(iii bis)  $N$  is an open and closed subscheme of  $M_a$ .

b) The first assertion follows from the isomorphism  $M_a \simeq \varphi^{-1}(a)$ , and the second from the fact that  $q$  is flat and  $q(e, a) = (\bar{e}, a)$ .

## 0.11 Density theorem and theory of the regular points of $G$

We apply the constructions and notation of the preceding section in the following situation. Let  $G$  be a smooth connected algebraic group over  $k$ ; take  $V = G$ , with  $G$  acting on  $V$  by inner automorphisms; and let  $W = H$ , where  $H$  is a smooth connected algebraic subgroup of  $G$ . We denote the Lie algebras of  $G$  and  $H$  by  $\mathfrak{g}$  and  $\mathfrak{h}$ , the normalizer of  $H$  in  $G$  by  $N$ , and its Lie algebra by  $\mathfrak{n}$ . For  $a \in G(k)$ , we again denote by  $M_a$ , as in Section 1, the inverse of the transporter of  $a$  into  $H$ . Thus, if  $a \in H(k)$ , then  $N \subset M_a$ ; in that case  $\mathfrak{m}_a$  denotes the Zariski tangent space of  $M_a$  at the identity element  $e$  of  $G$ . Note that

$$\mathfrak{h} \subset \mathfrak{n} \subset \mathfrak{m}_a \subset \mathfrak{g} \quad \text{for } a \in H(k).$$

We shall use the following lemma.

**Lemma 2.0.** — *The equality  $H = N^0$  holds if and only if*

$$(\mathfrak{g}/\mathfrak{h})^H = 0,$$

*where the left-hand side denotes the subspace of invariants for the action of  $H$  induced by the adjoint representation. When this condition holds,  $N$  is smooth and  $\dim X = \dim G$ . In all cases,  $\dim X \leq \dim G$ , and equality holds if and only if  $H$  has finite index in  $N$ .*

Indeed, by II 5.2.3(i),  $\mathfrak{n}$  is the inverse image of  $(\mathfrak{g}/\mathfrak{h})^H$  under  $\mathfrak{g} \rightarrow \mathfrak{g}/\mathfrak{h}$ . Hence  $(\mathfrak{g}/\mathfrak{h})^H = 0$  is equivalent to  $\mathfrak{h} = \mathfrak{n}$ , which in turn is equivalent to  $H = N^0$ , since  $H$  is a smooth connected algebraic subgroup of the algebraic group  $N$  (cf. VI.2). This also shows that  $N$  is smooth. Furthermore,

$$\dim X = (\dim G - \dim N) + \dim H = \dim G - (\dim N - \dim H),$$

and therefore

$$\dim X \leq \dim G,$$

with equality if and only if  $\dim H = \dim N$ , that is, if and only if  $H$  has finite index in  $N$ . This holds in particular when  $H = N^0$ , and proves the lemma.

**Theorem 2.1.** — *Let  $G$  be a smooth connected algebraic group over an algebraically closed field  $k$ , let  $H$  be a smooth connected algebraic subgroup, and let  $N$  be its normalizer. Write  $\mathfrak{g}, \mathfrak{h}, \mathfrak{n}$  for their Lie algebras, let  $X = G \times^N H$  be the scheme introduced in Section 1 (fibered over  $G/N$  with typical fiber  $H$ ), and let  $\psi : X \rightarrow G$  be the canonical morphism. Its image is also the image of  $\varphi : G \times H \rightarrow G$ , where*

$$\varphi(g, h) = \text{int}(g)h = ghg^{-1}.$$

*The following conditions are equivalent:*

- (i)  $H$  contains a Cartan subgroup  $C$  of  $G$  (XII 1).
- (i bis)  $H$  has the same reductive rank and the same nilpotent rank (XII 1) as  $G$ .
- (ii)  $H$  contains a maximal torus  $T$  of  $G$ , and  $(\mathfrak{g}/\mathfrak{h})^T = 0$ .
- (iii) The set of conjugates of  $H$  containing a given maximal torus is finite and nonempty, and  $H$  has finite index in its normalizer.
- (iv) There exists  $a \in H(k)$  that belongs to only finitely many conjugates of  $H$  (or merely such that  $\psi^{-1}(a)$  has an isolated point), and  $H$  has finite index in its normalizer.
- (iv bis) The morphism  $\psi : X \rightarrow G$  is generically quasi-finite (that is, there is a dense open subset of  $X$  on which  $\psi$  is quasi-finite), and  $H$  has finite index in its normalizer.
- (v) There is a dense open subset  $U$  of  $G$  such that, for every  $x \in U(k)$ , the set of conjugates of  $H$  containing  $x$  is finite and nonempty; equivalently,  $\psi : X \rightarrow G$  is dominant and generically quasi-finite.
- (vi) There is a dense open subset  $U$  of  $G$  such that every  $x \in U(k)$  is contained in a conjugate of  $H$ ; equivalently,  $\psi : X \rightarrow G$  is dominant.
- (vii) There exists  $a \in H(k)$  such that the fixed subspace of  $\mathfrak{g}/\mathfrak{h}$  under  $\text{ad}_{\mathfrak{g}/\mathfrak{h}}(a)$  is zero.

*Moreover, these conditions imply that  $H$  is its own connected normalizer:  $N$  is smooth and  $\dim H = \dim N$ . They also imply that  $\psi : X \rightarrow G$  is generically étale.*

*Proof.* By 2.0,  $\dim X \leq \dim G$ , with equality if and only if  $\dim H = \dim N$ , or equivalently if and only if  $H$  has finite index in  $N$ . The inequality  $\dim X \leq \dim G$  shows that  $\psi$  is dominant if and only if it is dominant and generically quasi-finite, and also if and only if it is generically quasi-finite and  $\dim X = \dim G$ . Since the last equality is equivalent, by 2.0, to  $H$  having finite index in  $N$ , conditions (vi), (v), and (iv bis) are equivalent. The equivalence of (iv) and (iv bis) is immediate.

The equivalence of (i) and (i bis) follows directly from the definitions and is left to the reader. If  $H$  contains a Cartan subgroup  $C$  of  $G$ , it contains the maximal torus  $T$  of  $C$ , which is a maximal torus of  $G$ . Since  $C$  is the centralizer of  $T$ , its Lie algebra  $\mathfrak{c}$  is

$$\mathfrak{c} = \mathfrak{g}^T$$

(II 5.2.3(ii)). Hence  $\mathfrak{g}^T = \mathfrak{c} \subset \mathfrak{h}$ , and I 4.7.3 gives the relation<sup>34</sup>

$$(x) \quad (\mathfrak{g}/\mathfrak{h})^T = 0.$$

<sup>34</sup>The French source has two evident typographical slips in this sentence, writing  $\mathfrak{n}$  for  $\mathfrak{c}$  and reversing the inclusion involving  $\mathfrak{h}$  and  $\mathfrak{g}^T$ . The mathematically required relation is restored here.

Conversely, suppose that  $H$  contains the maximal torus  $T$  and that (x) holds, so that  $\mathfrak{h} \supset \mathfrak{c}$ . We claim that  $H$  contains the centralizer  $C$  of  $T$ , which will prove the equivalence of (i) and (ii). This follows from the next lemma.

**Lemma 2.1.1.** — *Let  $G$  be a smooth algebraic group over a field  $k$ , let  $T$  be a subgroup of multiplicative type of  $G$ , let  $C$  be its connected centralizer (equal to the centralizer of  $T$  when  $G$  is connected and  $T$  is a torus, by XII 6.6(b)), and let  $H$  be a smooth subgroup of  $G$  containing  $T$ . Then  $H$  contains  $C$  if and only if its Lie algebra  $\mathfrak{h}$  contains the Lie algebra  $\mathfrak{c}$  of  $C$ .*

Indeed,  $\text{Centr}_G(T)$  is smooth over  $k$  by XI 2.4, and hence  $C$  is smooth over  $k$ . Likewise,  $\text{Centr}_H(T)$  is smooth, and

$$\text{Centr}_H(T) = \text{Centr}_G(T) \cap H$$

has Lie algebra  $\mathfrak{h} \cap \mathfrak{c}$ . The hypothesis therefore implies that this smooth subgroup has the same Lie algebra as the smooth group  $\text{Centr}_G(T)$ , so it contains the connected component  $C$  of the latter. Thus  $H$  contains  $C$ .  $\square$

We now prove the equivalence of (i) and (iii). If  $H$  contains the maximal torus  $T$  of  $G$ , we must show that  $H \supset C$  (equivalently, that (x) holds) if and only if  $H$  has finite index in its normalizer and the set of conjugates of  $H$  containing  $T$  is finite. If  $H \supset C$ , then (x) implies, a fortiori,

$$(xx) \quad (\mathfrak{g}/\mathfrak{h})^H = 0.$$

The Lie algebra  $\mathfrak{n}$  is the inverse image of the left-hand side under the canonical homomorphism  $\mathfrak{g} \rightarrow \mathfrak{g}/\mathfrak{h}$  (II 5.2.3(i)). Consequently, by 2.0, (xx) says that  $H = N^0$ ; in particular,  $H$  has finite index in its normalizer. Consider the diagram of subgroups

$$\begin{array}{ccccc} T & \longrightarrow & N(T) \cap H & \longrightarrow & H \\ & & \downarrow & & \downarrow \\ & & N(T) \cap N(H) & \longrightarrow & N(H) \\ & & \downarrow & & \\ & & N(T) & & \end{array}$$

By the conjugacy theorem for maximal tori in  $H$  (XII 6.6(a)), every conjugate of  $H$  containing  $T$  is obtained from  $H$  by conjugation by an element of  $N(T)(k)$ . Thus the set of such conjugates is in bijection with

$$(N(T)/(N(T) \cap N(H)))(k).$$

Since  $H \supset C$ , we have  $N(T) \cap H \supset C$ , so the preceding set is a quotient of  $(N(T)/C)(k)$ , which is finite. This proves (i)  $\Rightarrow$  (iii).

Conversely, suppose (iii), so that  $N(T)/(N(T) \cap N(H))$  and  $N(H)/H$  are finite. The conjugacy theorem in  $H$  shows that the homomorphism

$$(N(T) \cap N(H))/(N(T) \cap H) \longrightarrow N(H)/H$$

induced by the preceding diagram is bijective on  $k$ -valued points (indeed, it is an isomorphism). Since the latter quotient is finite, so is the former. Hence  $N(T) \cap H$  has finite

index in  $N(T)$  and therefore contains  $C = N(T)^0$ . Thus  $H \supset C$ , and (i), (i bis), (ii), and (iii) are equivalent.

We next prove (ii)  $\Rightarrow$  (vii). Both conditions are invariant under an extension  $k \rightarrow k'$  with  $k'$  algebraically closed. We may therefore assume that  $k$  has infinite transcendence degree over its prime subfield. There is then an element  $a \in T(k)$  such that the subgroup of  $T(k)$  generated by  $a$  is Zariski dense in  $T$ . It follows that

$$(\mathfrak{g}/\mathfrak{h})^T = (\mathfrak{g}/\mathfrak{h})^{\text{ad}(a)}.$$

The left-hand side is zero by hypothesis, which proves (vii).

The implication (vii)  $\Rightarrow$  (vi) is contained in the following result, which sharpens 2.1.

**Corollary 2.2.** — *Let  $G$  be a smooth algebraic group over a field  $k$ , let  $H$  be a smooth algebraic subgroup, let  $N$  be its normalizer in  $G$ , and let*

$$\varphi : G \times H \longrightarrow G, \quad \varphi(g, h) = \text{ad}(g)h = ghg^{-1}.$$

*Let  $\psi : X = G \times^N H \rightarrow G$  be the morphism obtained from  $\varphi$  by passage to the quotient (cf. Section 1), and let  $a \in H(k)$ . The following conditions are equivalent:*

- (i)  $\varphi$  is smooth at  $(e, a)$ .
- (ii)  $\psi$  is étale at  $(\bar{e}, a)$ , and  $N$  is smooth over  $k$ .
- (iii)  $(\mathfrak{g}/\mathfrak{h})^{\text{ad}(a)} = 0$ , where  $\mathfrak{g}$  and  $\mathfrak{h}$  are the Lie algebras of  $G$  and  $H$ .

*These conditions imply  $H^0 = N^0$ .*

The smoothness at a  $k$ -rational point of  $\varphi$ , a morphism between smooth  $k$ -preschemes, is equivalent to the surjectivity of the tangent map at that point. Under the usual identifications of the tangent spaces of  $G$  and  $H$  with their Lie algebras, a direct calculation gives

$$d\varphi(\xi, \eta) = (\text{id} - \text{ad}(a))\xi + \eta,$$

as a map from  $\mathfrak{g} \times \mathfrak{h}$  to  $\mathfrak{g}$ . Its surjectivity is therefore equivalent to the surjectivity of  $\text{id} - \text{ad}(a)$  on  $\mathfrak{g}/\mathfrak{h}$ , which is precisely (iii).

Condition (iii) implies, a fortiori,  $(\mathfrak{g}/\mathfrak{h})^H = 0$ , and hence  $H^0 = N^0$  by 2.0 (the connectedness hypothesis made at the beginning of the section is not needed there). It follows that  $N$  is smooth and that  $\dim H = \dim N$ , whence  $\dim X = \dim G$ . Since  $q : G \times H \rightarrow X$  is flat and  $\psi \circ q = \varphi^{35}$  is smooth at  $(e, a)$ , the morphism  $\psi$  is smooth at  $q(e, a) = (\bar{e}, a)$ , and hence is étale there by dimension. We have proved (i)  $\iff$  (iii)  $\implies$  (ii). Conversely, (ii) implies (i), because the smoothness of  $N$  implies that of  $q$ .  $\square$

In particular, under condition (vii) of 2.1 the morphism  $\varphi$  is smooth at  $(e, a)$ , so its image contains a nonempty open subset of  $G$ . Thus  $\psi$  is dominant, proving (vii)  $\Rightarrow$  (vi).

It remains to prove (vi)  $\Rightarrow$  (i). Suppose first that  $G$  is affine. Let  $U$  be a nonempty open subset of  $G$  such that every  $x \in U(k)$  is contained in a conjugate of  $H$ , and let  $C$  be a Cartan subgroup of  $G$ . Applying the already proved implication (i)  $\Rightarrow$  (iv) with  $C$  in place of  $H$  (Borel's "density theorem"), we find a conjugate of  $C$  meeting  $U$ . We may therefore assume that  $U \cap C \neq \emptyset$ ; equivalently, there is a nonempty open subset  $V$  of  $C$  such that every  $x \in V(k)$  belongs to a conjugate of  $H$ .

Write

$$C = T \cdot C_u,$$

<sup>35</sup>The source reverses this composition, printing  $\psi = \varphi \circ q$ . The displayed morphisms and the definition of  $\varphi$  require  $\varphi = \psi \circ q$ .

where  $T$  is the maximal torus of  $C$ , hence a maximal torus of  $G$ , and  $C_u$  is the unipotent part of  $C$ ;  $T$  lies in the center of  $C$  (*BIBLE* 6, Theorem 2). We may again assume that  $k$  has infinite transcendence degree over its prime subfield. Choose  $t \in T(k)$  in the projection of  $V$  onto  $T$ , a nonempty open subset of  $T$ , so that  $tC_u \cap V \neq \emptyset$ , and choose it so that it generates a Zariski-dense subgroup of  $T$ . Every algebraic subgroup of  $G$  containing a product  $tu$ , with  $t \in T(k)$  and  $u \in C_u(k)$ , contains both factors (*BIBLE* 4, Theorem 3). Taking  $tu \in V(k)$ , we conclude that a conjugate of  $H$  contains  $t$ , and therefore contains  $T$ . After replacing  $H$  by that conjugate, we may assume

$$T \subset H.$$

Let  $W$  be the inverse image in  $C_u$  of  $V$  under  $u \mapsto tu$ . For every  $x \in (T \cdot W)(k)$ , there is then a conjugate of  $H$  containing both  $T$  and  $x$ . As observed above, every such conjugate is of the form  $\text{int}(g)H$  for some  $g \in N(T)(k)$ . Consider

$$f : N(T) \times H \longrightarrow G, \quad f(g, h) = \text{int}(g)h = ghg^{-1}.$$

The image of  $f$  contains  $TW$ . Since  $C$  has finite index in  $N(T)$ , the latter is a finite union of translates  $Cg_i$ , with  $g_i \in N(T)(k)$ . It follows that some dense open subset  $V'$  of  $C = TC_u$  is contained in the image under  $f$  of  $(Cg_i) \times H$ . Replacing  $H$  by  $\text{int}(g_i)H$ , we may assume  $g_i = e$ , so that

$$f(C \times H) \supset V'.$$

Thus for every  $u \in V'(k)$  there are  $v \in C(k)$  and  $h \in H(k)$  such that

$$v^{-1}hv = u, \quad \text{hence} \quad vuv^{-1} \in H(k).$$

Set

$$C' = C \cap H = \text{Centr}_H(T).$$

Since  $vuv^{-1}$  also belongs to  $C$ , it lies in  $C'$ , and hence  $u \in \text{int}(v^{-1})C'$ . Therefore the union of the conjugates of  $C'$  in  $C$ , by elements of  $C(k)$ , is dense. As in the preceding argument for the pair  $(G, H)$ , now applied to  $(C, C')$ , this implies that  $C'$  has finite index in its normalizer in  $C$ . By the lemma of *BIBLE* 7, Section 1, we obtain  $C' = C$ , and consequently  $C \subset H$ .<sup>36</sup> This proves (vi)  $\Rightarrow$  (i) when  $G$  is affine.

In the general case we argue by induction on  $n = \dim G$ , the assertion being trivial for  $n = 0$ . Let  $Z$  be the center of  $G$ . There are two cases.

- 1° If  $\dim(Z \cap H) > 0$ , set  $G' = G/(Z \cap H)$ . Then  $\dim G' < n$ , and condition (vi) for  $H$  implies the same condition for its image  $H'$  in  $G'$ . By induction,  $H'$  contains a Cartan subgroup  $C'$  of  $G'$ . Hence  $H$  contains the inverse image  $C$  of  $C'$ , which is a Cartan subgroup by XII 6.6(e).
- 2° If  $\dim(Z \cap H) = 0$ , the canonical morphism  $H \rightarrow G/Z$  is finite. Since  $G/Z$  is affine by XII 6.1,  $H$  is affine. Every homomorphism from  $H$  to an abelian variety is therefore zero (indeed, every morphism of preschemes from  $H$  to an abelian variety is constant). This follows from the fact that a smooth connected affine algebraic group over an algebraically closed field is a rational variety, or simply from the fact that it is the union of its Borel subgroups (*BIBLE* 6, Theorem 5(b)); the structure theorems *BIBLE* 6.2 and 6.3 show readily that a smooth connected solvable affine group is a rational variety.

<sup>36</sup>The source prints  $C = H$ ; only the inclusion  $C \subset H$  follows and is used.

Apply Chevalley's structure theorem to  $G$ : the group  $G$  is an extension of an abelian variety  $A$  by a smooth affine group. Since  $H$  is affine, its image in  $A$  is zero. On the other hand, that image must be all of  $A$ , because the union of its conjugates in  $A$  is dense. Hence  $A = 0$ , so  $G$  is affine, reducing us to the case already proved.

This completes the proof of 2.1.

**Corollary 2.3.** — *Assume that the equivalent conditions of 2.1 hold.*

- a) Let  $k(X)$  and  $k(G)$  be the fields of rational functions of  $X$  and  $G$ , respectively. Then  $k(X)$  is a finite separable extension of  $k(G)$ . Denote its degree by  $d$ .
- b) Let  $T$  be a maximal torus of  $G$  contained in  $H$ , which exists by 2.1(ii), and let  $C$  be the corresponding Cartan subgroup of  $G$ . Then  $C \subset H$ . Moreover,  $\text{Norm}_G(T)$  is a smooth subgroup of  $G$ , and

$$\text{Norm}_G(T) \cap \text{Norm}_G(H) = \text{Norm}_{\text{Norm}_G(H)}(T)$$

is a smooth subgroup of it, of finite index equal to the degree  $d$  defined in (a). The number of conjugates of  $H$  containing a given maximal torus, or a given Cartan subgroup, is  $d$ .

- c) Let  $U$  be the largest open subset of  $G$  such that  $\psi : X \rightarrow G$  induces a finite étale morphism

$$\psi^{-1}(U) \longrightarrow U.$$

Then  $U$  is dense. For  $g \in G(k)$ , the following are equivalent:

- (a)  $g \in U(k)$ ;
- (b) exactly  $d$  conjugates of  $H$  contain  $g$ ;
- (c) there are at least  $d$  distinct conjugates  $H_i$  of  $H$  containing  $g$ , and for every  $i$ ,

$$(\mathfrak{g}/\mathfrak{h}_i)^{\text{ad}(g)} = 0, \quad \mathfrak{h}_i = \text{Lie}(H_i).$$

Assertion (a) follows from the generic étaleness of  $\psi$ , stated at the end of 2.1. This also shows that the open set  $U$  in (c) is nonempty, hence dense, and gives the two stated characterizations of its  $k$ -points: here one uses that  $\psi$  is separated, that  $X$  is integral, that  $G$  is integral and normal (SGA 1, I 10.11), and that  $(\mathfrak{g}/\mathfrak{h}_i)^{\text{ad}(g)} = 0$  means that  $\psi$  is étale at the point  $x_i \in \psi^{-1}(g)$  corresponding to  $H_i$ .

If  $H$  contains the maximal torus  $T$  of  $G$ , the centralizers of  $T$  in  $H$  and  $G$  have the same dimension and are smooth and connected (XII 6.6(b)); they are therefore equal, proving that  $C \subset H$ .<sup>37</sup> The normalizer of  $T$  in any smooth group containing it is smooth (XI 2.4 bis), so  $\text{Norm}_G(T)$  and  $\text{Norm}_{\text{Norm}_G(H)}(T)$  are smooth. Recall also that  $N = \text{Norm}_G(H)$  is smooth by the last assertion of 2.1. Furthermore,  $\text{Norm}_N(T)$  contains  $C$ , which has finite index in  $\text{Norm}_G(T)$ ; hence  $\text{Norm}_N(T)$  itself has finite index in  $\text{Norm}_G(T)$ .

The conjugacy theorem for maximal tori in  $H$  shows that this index is the number of conjugates of  $H$  containing  $T$ , equivalently the number containing  $C$ . The union of the conjugates of  $C$  in  $G$  is dense, by 2.1(i)  $\Rightarrow$  (vi) applied to  $C$  in place of  $H$ , and  $U$  is stable under inner automorphisms. Hence  $C \cap U \neq \emptyset$ . Arguing as in the proof of 2.1(vi)  $\Rightarrow$  (ii), and making a harmless extension of the base field if necessary, one finds  $g \in (C \cap U)(k)$  such that every conjugate of  $H$  containing  $g$  contains  $T$ , and hence  $C$ . Thus the conjugates

<sup>37</sup>The source prints the tautological  $C \subset G$ ; equality of the two centralizers gives  $C \subset H$ .

of  $H$  containing  $C$  are precisely those containing  $g$ ; since  $g \in U(k)$ , their number is  $d$ . This proves (b).

In fact, the argument shows that the set of conjugates of  $H$  containing  $T$  is a homogeneous set under the group of rational points of

$$W_G(T) = \text{Norm}_G(T) / \text{Centr}_G(T).$$

In particular,

$$d \leq \text{the order of the Weyl group of } G.$$

**Corollary 2.4.** — *With the notation of 2.1, the following conditions are equivalent:*

- (i)  $\psi : X \rightarrow G$  is birational.
- (ii) Exactly one conjugate of  $H$  contains a given Cartan subgroup of  $G$ .
- (iii)  $H$  contains a Cartan subgroup  $C$  of  $G$ , and  $\text{Norm}_G(H) \supset \text{Norm}_G(C)$ .
- (iv) There is a nonempty open subset  $V$  of  $G$  such that every  $g \in V(k)$  belongs to exactly one conjugate of  $H$ .

This follows immediately from 2.1 and 2.3.

**Corollary 2.5.** — *Assume that the conditions of 2.4 hold, and let  $g \in G(k)$ . The following conditions are equivalent:*

- (i)  $g \in U(k)$ , where  $U$  is defined in 2.3(c); that is,  $g$  is contained in exactly one conjugate of  $H$ .
- (ii) The set of conjugates of  $H$  containing  $g$  is finite and nonempty.
- (iii) The scheme  $\psi^{-1}(g)$ , the “scheme of conjugates of  $H$  containing  $g$ ,” has an isolated point.
- (iv) There is a conjugate  $H'$  of  $H$  containing  $g$  such that

$$(\mathfrak{g}/\mathfrak{h}')^{\text{ad}(g)} = 0, \quad \mathfrak{h}' = \text{Lie}(H').$$

Finally,  $U$  is also the largest open subset of  $G$  on which  $\psi$  induces an isomorphism

$$\psi^{-1}(U) \xrightarrow{\sim} U.$$

The equivalence of (i), (ii), and (iii), together with the last assertion, follows from the “Main Theorem”<sup>38</sup> applied to the birational morphism  $\psi : X \rightarrow G$ , since  $G$  is normal. Their equivalence with (iv) follows immediately from Corollary 2.2, which characterizes the points of  $X$  at which  $\psi$  is étale.<sup>39</sup>

**Theorem 2.6.** — *Let  $G$  be a smooth connected algebraic group over an algebraically closed field  $k$ , let  $C$  be a Cartan subgroup associated with a maximal torus  $T$ , and put*

$$N = \text{Norm}_G(C) = \text{Norm}_G(T) \quad (\text{cf. XII 8.4}).$$

*Let  $X = G \times^N C$ , where  $N$  acts on the left factor  $G$  by right translations and on the right factor  $C$  by inner automorphisms, and let  $\psi : X \rightarrow G$  be the canonical morphism.*

<sup>38</sup>Editor’s note in the French source: Zariski’s Main Theorem.

<sup>39</sup>The source cites the final assertion of 2.1 here; the pointwise étaleness criterion is Corollary 2.2.



- a)  $\psi$  is birational.  
 b) Let  $U$  be the largest open subset of  $G$  such that  $\psi$  induces an isomorphism

$$\psi^{-1}(U) \xrightarrow{\sim} U$$

(cf. 2.5), and let

$$\rho = \rho_\nu(G) = \dim C$$

be the nilpotent rank of  $G$ . For every  $g \in G(k)$ , the multiplicity of the eigenvalue 1 in the action of  $\text{ad}(g)$  on  $\mathfrak{g}$  is at least  $\rho$ ; it equals  $\rho$  if and only if  $g \in U(k)$ .

*Proof.* Condition 2.1(i) holds, so 2.4(iii)  $\Rightarrow$  (i) proves (a). In *BIBLE* 7, when  $G$  is affine, the points of  $U(k)$  are called the *regular points* of  $G(k)$ . We shall use the same terminology and call  $U$  the *open subset of regular points* of  $G$ . Notice that the proof in *BIBLE* that this set is itself open is incorrect; the present Exposé establishes its openness under more general hypotheses.

To prove (b), for each  $g \in G(k)$  introduce the characteristic polynomial

$$P(\text{ad}(g), t) = t^n + c_1(g)t^{n-1} + \cdots + c_n(g).$$

Replacing  $k$  by an arbitrary  $k$ -algebra shows immediately that the functions  $c_i(g)$  arise from uniquely determined sections

$$c_i \in \Gamma(G, \mathcal{O}_G).$$

Suppose that  $g \in G(k)$  belongs to a Cartan subgroup, which we may take to be  $C$ ; this includes the case of a regular element. By 2.5(iv),

$$(\mathfrak{g}/\mathfrak{c})^{\text{ad}(g)} = 0$$

if and only if  $g$  is regular, where  $\mathfrak{c} = \text{Lie}(C)$ . Since  $C$  is nilpotent,  $\text{ad}_{\mathfrak{c}}(g)$  has only the eigenvalue 1. It follows that the multiplicity of the eigenvalue 1 in  $\text{ad}_{\mathfrak{g}}(g)$  is at least  $\rho$ , and is exactly  $\dim C = \rho$  if and only if  $g$  is regular.

In particular, the characteristic polynomial above is divisible by  $(t-1)^\rho$ . Divisibility by  $(t-1)^\rho$  is expressed by linear relations, with integer coefficients, among the coefficients of the polynomial. These relations hold for  $g \in U(k)$ , where  $U$  is dense; since  $G$  is reduced, they therefore hold everywhere. Thus<sup>40</sup>

$$(\dagger) \quad t^n + c_1 t^{n-1} + \cdots + c_n = (t-1)^\rho (t^{n-\rho} + b_1 t^{n-\rho-1} + \cdots + b_{n-\rho})$$

in the polynomial ring over  $\Gamma(G, \mathcal{O}_G)$ . In particular, for every field extension  $K/k$  and every  $g \in G(K)$ , the eigenvalue 1 of  $\text{ad}(g)$  has multiplicity at least  $\rho$ .

We have already seen that equality holds when  $g$  is regular; let us prove the converse. Suppose first that  $G$  is affine, and write

$$g = g_s g_u$$

as the product of its semisimple and unipotent parts (*BIBLE* 4, Section 4). Then

$$\text{ad}(g) = \text{ad}(g_s) \text{ad}(g_u)$$

<sup>40</sup>The French source prints  $c_0$  in place of the omitted tail  $+\cdots + c_n$  on the left-hand side of  $(\dagger)$ . The characteristic polynomial displayed immediately above determines the correction made here.

is the corresponding decomposition of  $\mathrm{ad}(g)$  (loc. cit., corollary to Theorem 3). Hence  $\mathrm{ad}(g)$  and  $\mathrm{ad}(g_s)$  have the same eigenvalues, counted with multiplicity; in particular, the eigenvalue 1 has the same multiplicity in both.

By *BIBLE* 7, Theorem 2, Corollary 1, the element  $g$  is regular if and only if  $g_s$  is regular. We may therefore assume that  $g = g_s$  is semisimple. It is then contained in a maximal torus by *BIBLE* 6, Theorem 5(c), and hence in a Cartan subgroup, a case already treated. This proves (b) when  $G$  is affine.

In general, let  $Z = \mathrm{Centr}(G)_{\mathrm{red}}$ . By XII 6.6(e), the Cartan subgroups of  $G$  are the inverse images of those of  $G' = G/Z$ . Thus  $g$  is regular in  $G$  if and only if its image  $g'$  is regular in  $G'$ . Since  $Z$  is smooth, the Lie algebra of  $G'$  is  $\mathfrak{g}' = \mathfrak{g}/\mathfrak{z}$ , where  $\mathfrak{z} = \mathrm{Lie}(Z)$ , and  $\mathrm{ad}(g')$  is the endomorphism induced by  $\mathrm{ad}(g)$  on  $\mathfrak{g}/\mathfrak{z}$ . Consequently, the multiplicity of the eigenvalue 1 in  $\mathrm{ad}(g)$  is

$$d = \dim Z$$

plus its multiplicity in  $\mathrm{ad}(g')$ . The former therefore equals the nilpotent rank of  $G$  if and only if the latter equals the nilpotent rank of  $G'$ . We are reduced to  $G'$ , which is affine by XII 6.1, and that case has already been proved. This completes the proof of 2.6.

**Corollary 2.7.** — *With the notation of the preceding proof,<sup>41</sup> let*

$$b = 1 + b_1 + \cdots + b_{n-\rho} \in \Gamma(G, \mathcal{O}_G).$$

*Then the open subset of regular points of  $G$  is*

$$U = G_b,$$

*the set of points of  $G$  at which  $b$  is invertible. In particular,  $U$  is an affine open subset when  $G$  is affine.*

**Corollary 2.8.** — *Let  $H$  be a smooth connected algebraic subgroup of  $G$  containing a Cartan subgroup of  $G$ .*

- a) Let  $C$  be an algebraic subgroup of  $H$ . Then  $C$  is a Cartan subgroup of  $H$  if and only if it is a Cartan subgroup of  $G$ .*
- b) Let  $g \in G(k)$ , and let  $d$  be the integer introduced in 2.3. Then  $g$  is a regular point of  $G$  if and only if there are exactly  $d$  conjugates  $H_i$  of  $H$  containing  $g$ , and  $g$  is a regular element of every  $H_i$ . Equivalently, there are at most  $d$  conjugates of  $H$  containing  $g$ , and  $g$  is regular in one of them. When these conditions hold, if  $C$  is the unique Cartan subgroup of  $G$  containing  $g$ , then the conjugates of  $H$  containing  $g$  are precisely the conjugates of  $H$  containing  $C$ .*
- c) Let  $g \in H(k)$ . Then  $g$  is regular in  $G$  if and only if it is regular in  $H$  and*

$$(\mathfrak{g}/\mathfrak{h})^{\mathrm{ad}(g)} = 0.$$

*Proof.* *a)* Under either hypothesis on  $C$ , its unique maximal torus  $T$  is a maximal torus of both  $H$  and  $G$ , since  $H$  has the same reductive rank as  $G$ . The smooth connected groups  $\mathrm{Centr}_H(T) \subset \mathrm{Centr}_G(T)$  have the same dimension, and therefore are equal. Hence  $C$  is equal to one of these two centralizers if and only if it is equal to the other, proving (a).

<sup>41</sup>Editor's note in the French source: cf. (†) above.

b) Suppose first that  $g$  is regular in  $G$ , and let  $C$  be the unique Cartan subgroup of  $G$  containing  $g$ . By 2.3(b), exactly  $d$  conjugates  $H_i$  of  $H$  contain  $C$ . Since

$$(\mathfrak{g}/\mathfrak{c})^{\mathrm{ad}(g)} = 0,$$

the endomorphism  $\mathrm{ad}(g)$  has no eigenvalue 1 on  $\mathfrak{g}/\mathfrak{c}$ , and a fortiori

$$(\mathfrak{g}/\mathfrak{h}_i)^{\mathrm{ad}(g)} = 0.$$

By 2.3(c), exactly  $d$  conjugates of  $H$  contain  $g$ , namely the  $H_i$ . A Cartan subgroup of such an  $H_i$  containing  $g$  is a Cartan subgroup of  $G$  by (a), and hence is  $C$ . Thus  $g$  is regular in every  $H_i$ .

Conversely, suppose that at most  $d$  conjugates  $H_i$  of  $H$  contain  $g$ , and that  $g$  is regular in one of them, which we may take to be  $H$ . Since  $g$  is regular in  $H$ , it belongs to a unique Cartan subgroup  $C$  of  $H$ ; by (a),  $C$  is a Cartan subgroup of  $G$ . Let  $C'$  be any Cartan subgroup of  $G$  containing  $g$ . By 2.3(b), exactly  $d$  conjugates of  $H$  contain  $C'$ . They all contain  $g$ , so they are precisely the  $H_i$ ; in particular,  $H$  contains  $C'$ . Thus  $C$  and  $C'$  are two Cartan subgroups of  $H$  containing the same regular element  $g$  of  $H$ , and therefore  $C = C'$ . Hence  $g$  is regular in  $G$ .<sup>42</sup>  $\square$

c) For an endomorphism  $u$  of a finite-dimensional vector space, let  $\nu(u)$  denote the nullity<sup>43</sup> of  $\mathrm{id} - u$ . Then

$$\nu(\mathrm{ad}(g)_{\mathfrak{g}}) = \nu(\mathrm{ad}(g)_{\mathfrak{h}}) + \nu(\mathrm{ad}(g)_{\mathfrak{g}/\mathfrak{h}}).$$

The two terms on the right are respectively at least the nilpotent rank of  $H$ , which equals the nilpotent rank  $\rho$  of  $G$  by (a), and at least 0. Hence  $\nu(\mathrm{ad}(g)_{\mathfrak{g}}) = \rho$  if and only if

$$\nu(\mathrm{ad}(g)_{\mathfrak{h}}) = \rho, \quad \nu(\mathrm{ad}(g)_{\mathfrak{g}/\mathfrak{h}}) = 0.$$

This says precisely that  $g$  is regular in  $G$  if and only if it is regular in  $H$  and  $\mathrm{ad}(g)_{\mathfrak{g}/\mathfrak{h}}$  has no nonzero invariants.  $\square$

**Remarks 2.9.** — In the statement of 2.1, condition (iii) cannot be weakened by assuming only that  $H$  contains a maximal torus and has finite index in its normalizer, even if one also requires that normalizer to be smooth, that is, that  $H = N^0$ , and even when  $G$  is affine and solvable. An example is the group  $G$  of matrices of the form

$$g = \begin{pmatrix} t & a & c \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix},$$

and the subgroup  $H$  formed by the matrices above for which  $b = c = 0$ . The Cartan subgroup of  $G$  here consists of the matrices  $g$  for which  $a = c = 0$ .

**Remarks 2.10.** — Let  $G$  be a smooth algebraic group over  $k$ , and let  $H$  be a smooth algebraic subgroup; we no longer assume that  $G$  and  $H$  are connected. Suppose that  $H^0$  contains a Cartan subgroup of  $G^0$ . Then

$$(\mathfrak{g}/\mathfrak{h})^H \subset (\mathfrak{g}/\mathfrak{h})^{H^0} = 0,$$

so  $H^0 = N^0$ , where  $N$  is the normalizer of  $H$ ; in particular,  $N$  is smooth.

<sup>42</sup>The source prints “regular in  $C$ ” at this conclusion; the argument proves regularity in  $G$ .

<sup>43</sup>Editor’s note in the French source: that is, the dimension of the nullspace of  $\mathrm{id} - u$ .

Nevertheless, even with  $G$  connected, examples are easily constructed in which  $H$  has a connected component  $H_i$  such that

$$(\mathfrak{g}/\mathfrak{h})^{\mathrm{ad}(h)} \neq 0 \quad \text{for every } h \in H_i(k).$$

Equivalently, the morphism

$$G \times H_i \longrightarrow G, \quad (g, h) \longmapsto \mathrm{ad}(g) \cdot h$$

is nowhere étale, or even quasi-finite. One may take, for example,  $H$  to be the normalizer of a maximal torus in  $\mathrm{SL}(2)_k$ .

Similarly, even if the image  $H''$  of  $H$  in the finite group  $G'' = G/G^0$  is all of  $G''$ , the union of the conjugates of  $H$  in  $G$  need not be dense. For example, take  $G$  to be the semidirect product of  $\mathbb{Z}/2\mathbb{Z}$  with  $G^0 = \mathrm{SL}(2)_k$ , on which it acts by “symmetry,” and take  $H$  to be the semidirect product  $\mathbb{Z}/2\mathbb{Z} \cdot T$ , where  $T$  is a maximal torus of  $G^0$ .

On the other hand, if one does not assume in advance that  $H^0$  contains a Cartan subgroup of  $G^0$ , but assumes that the union of the conjugates of  $H$  in  $G$  is dense, then  $H^0$  necessarily contains a Cartan subgroup of  $G^0$ . To see this, one may plainly assume that  $G$  is connected and repeat the proof of 2.1(vi)  $\Rightarrow$  (i), which does not require  $H$  to be connected.

## 0.12 The case of an arbitrary base prescheme

Suppose first that the base is a field  $k$ , not necessarily algebraically closed. Since conditions 2.1 (i bis), (iv bis), (v), and (vi) are invariant under extension of the base field, passage to the algebraic closure  $\bar{k}$  of  $k$  shows that they are equivalent to one another, and equivalent to the assertion that  $H_{\bar{k}}$  contains a Cartan subgroup of  $G_{\bar{k}}$ . When this condition is satisfied, it is still true (with the notation of 2.3) that  $k(X)$  is a finite separable extension of  $k(G)$ , of degree  $d$  independent of every extension of the base. If  $U$  is the largest open subset of  $G$  such that  $\psi$  induces a finite étale morphism

$$\psi^{-1}(U) \longrightarrow U,$$

then the formation of  $U$  commutes with extension of the base field. If  $\psi$  is birational, then  $U$  is also the largest open subset of  $G$  such that  $\psi$  induces an isomorphism

$$\psi^{-1}(U) \xrightarrow{\sim} U;$$

and if  $g \in U(k)$ , there is one and only one subgroup  $H'$  of  $G$ , conjugate to  $H$  over  $\bar{k}$ , such that  $H' \ni g$ .

A point  $g \in G(k)$  is called *regular* if it is regular as an element of

$$G(\bar{k}) = G_{\bar{k}}(\bar{k}).$$

More generally, the construction of 2.7 gives an open subset of  $G$ , whose formation commutes with every extension of the base field, called the *open subset of regular points* of  $G$ . It is also characterized by the following property: for every algebraically closed extension  $K$  of  $k$  and every  $g \in G(K)$ , the point  $g$  is regular in  $G_K$  if and only if  $g \in U(K)$ . If  $g \in U(k)$ <sup>44</sup>, then  $g$  is contained in one and only one Cartan subgroup of  $G$ , as we shall prove below under more general hypotheses.

<sup>44</sup>Editor’s note:  $G(k)$  has been corrected to  $U(k)$ .

Let  $G$  be a smooth, separated group prescheme of finite type, with connected fibers over the prescheme  $S$ , and consider the functor

$$\mathcal{C} : (\mathrm{Sch})^\circ_{/S} \longrightarrow (\mathrm{Ens})$$

defined by

$$\mathcal{C}(S') = \{ \text{Cartan subgroups (XII 3.1) of } G_{S'} \}.$$

Suppose that this functor is representable by a prescheme smooth over  $S$ . In Exposé XV we give a necessary and sufficient condition for this to hold, but we already know that the hypothesis is satisfied if  $G$  is affine over  $S$  with locally constant reductive rank (XII 3.3), or, more generally, if  $G$  locally admits a maximal torus for the fpqc topology (XII 7.1 a)); this is the case, for example, if  $S$  is the spectrum of a field.

Let  $X$  be the Cartan subgroup of the  $\mathcal{C}$ -group prescheme  $G_{\mathcal{C}}$ , the “universal Cartan subgroup” of  $G$ . Thus, as a prescheme over  $S$ ,  $X$  represents the functor

$$X(S') = \left\{ (C, g) \mid \begin{array}{l} C \text{ is a Cartan subgroup of } G_{S'}, \\ g \text{ is a section of } C \text{ over } S' \end{array} \right\}.$$

Consider the canonical projection morphism  $(C, g) \mapsto g$ ,

$$\psi : X \longrightarrow G.$$

We then have the following theorem.

**Theorem 0.12.1.** *Under the preceding hypotheses on  $G$ , and with the preceding notation, let  $U$  be the set of  $g \in G$  such that  $g$  is a regular element of its fiber  $G_s$ . Then  $U$  is open, and it is also the largest open subset  $U$  of  $G$  such that  $\psi$  induces an isomorphism*

$$\psi^{-1}(U) \xrightarrow{\sim} U.$$

Let us first prove that  $U$  is open. By the representability of  $\mathcal{C}$  as a prescheme smooth over  $S$ , and since its structure morphism is plainly surjective, it follows at once that  $G$  locally admits a Cartan subgroup for the étale topology and that the nilpotent rank of the fibers of  $G$  is locally constant. The same holds for the dimension of the fibers of  $G$ . After localizing on  $S$ , we may therefore suppose that these two integers are constant, say  $\rho$  and  $n$ . Consider the Killing polynomial

$$P_G(t) = t^n + c_1 t^{n-1} + \cdots + c_n \in A[t], \quad A = \Gamma(G, \mathcal{O}_G).$$

The restriction of this polynomial to the fibers  $G_s$  of  $G$ , and in particular to the fibers at the maximal points of  $S$ , is divisible by  $(t-1)^\rho$ . This says that certain linear combinations of the  $c_i$ , with integer coefficients, vanish on the fibers  $G_s$ . When  $S$  is reduced, as we may assume in order to prove that  $U$  is open, it follows that these combinations themselves vanish. Hence the Killing polynomial itself is divisible by  $(t-1)^\rho$ , say<sup>45</sup>

$$P_G(t) = (t-1)^\rho (t^{n-\rho} + b_1 t^{n-\rho-1} + \cdots + b_{n-\rho}).$$

Let  $b$  be the sum of the coefficients  $b_0 = 1, b_1, \dots, b_{n-\rho}$  of the second factor. By 2.7, applied to the fibers of  $G$ , we have

$$U = G_b,$$

<sup>45</sup>The source prints the undefined exponent  $n - r - 1$ . It is corrected to  $n - \rho - 1$ , as required by the degree  $n - \rho$  of the second factor.

which proves that  $U$  is open.

To prove that  $\psi^{-1}(U) \rightarrow U$  is an isomorphism, SGA 1, I 5.7 reduces us to checking this fiber by fiber. We are thereby reduced to the case of a base field, which may be assumed algebraically closed. There is then a Cartan subgroup  $C$  of  $G$ . If  $N$  is its normalizer, the conjugacy theorem XII 7.1 a) and b) identifies  $\mathcal{C}$  with  $G/N$ , and the morphism  $\psi : X \rightarrow G$  considered here is precisely the one defined in Section 2. The conclusion follows from 2.6 b). The same argument also shows that  $U$  is the largest open subset of  $G$  for which  $\psi$  induces an isomorphism  $\psi^{-1}(U) \rightarrow U$ .

**Corollary 0.12.2.** *Under the hypotheses of 3.1, let  $g$  be a regular section of  $G$ , that is, suppose that  $g(s)$  is a regular point of  $G_s$  for every  $s \in S$ . Then there is one and only one Cartan subgroup  $C$  of  $G$  such that  $g$  is a section of  $C$ .*

Indeed, the hypothesis on  $g$  says that  $g$  is a section of  $U$ , while the conclusion says that it admits a unique lift to a section of  $X$ . This is simply another way of saying that  $\psi^{-1}(U) \rightarrow U$  is an isomorphism.

Observe now that the open subset  $\psi^{-1}(U)$  of the Cartan subgroup  $X$  of  $G_{\mathcal{C}}$  is precisely the open subset of  $X$  consisting of the points regular in  $G_{\mathcal{C}}$ , by which we mean regular in their fibers. We thus obtain a natural “fibration” of the dense open subset  $U$  of regular points of  $G$  over the prescheme  $\mathcal{C}$ ; its fibers are dense open subsets of Cartan subgroups of the fibers of  $G_{\mathcal{C}}$ , namely their open subsets of points regular in  $G$ . For example, we obtain the following result, which will be considerably sharpened in the next Exposé.

**Corollary 0.12.3.** *Let  $G$  be a smooth connected algebraic group over the field  $k$ , and let  $\mathcal{T}$  be the scheme of maximal tori of  $G$ , which is isomorphic to the scheme of Cartan subgroups of  $G$ . Then the function field  $k(G)$  of  $G$  is isomorphic to the function field of a smooth connected affine nilpotent algebraic group  $C$  over the function field  $k(\mathcal{T})$  of  $\mathcal{T}$ , namely*

$$C = \text{“the generic Cartan subgroup of } G\text{”}.$$

*If  $G$  is affine and has unipotent rank zero, that is, if the Cartan subgroups of  $G_{\bar{k}}$ <sup>46</sup> are tori, then  $k(G)$  is a unirational extension of  $k(\mathcal{T})$ .*

Of course, by the generic Cartan subgroup of  $G$  we mean, by an abuse of language, the Cartan subgroup of  $G_{k(\mathcal{T})}$  that is the generic fiber of  $X$  over  $\mathcal{T}$ . It remains only to prove the last assertion of 3.3, which is contained in the following well-known result, due to Chevalley.

**Lemma 0.12.4.** *Let  $k$  be a field,  $T$  a torus over  $k$ , and  $k(T)$  the field of rational functions on  $T$ . Then  $k(T)$  is a unirational extension of  $k$ , that is, it is contained in a purely transcendental extension of  $k$ .*

Indeed, let  $k'$  be a finite separable extension of  $k$  that splits<sup>47</sup>  $T$  (X 1.4). Then  $T \otimes_k k'$  is a rational variety: it contains a dense open subset isomorphic to a dense open subset of affine  $n$ -space  $\mathbb{A}_{k'}^n$ . Hence

$$T' = \coprod_{\text{Spec}(k')/\text{Spec}(k)} T_{k'}/\text{Spec}(k')$$

is a rational variety, since it has a dense open subset isomorphic to a dense open subset of

$$\coprod_{\text{Spec}(k')/\text{Spec}(k)} \mathbb{A}_{k'}^n,$$

<sup>46</sup>The source prints the undefined base field  $\bar{K}$ ; it is corrected to  $\bar{k}$ .

<sup>47</sup>Editor’s note: “splits” (?), or: “over which  $T$  is split”.

which is isomorphic to affine  $mn$ -space over  $k$ , where  $m = [k' : k]$ . Consider the norm homomorphism from  $T'$  to  $T$ , which is defined whenever  $T$  is a commutative group scheme over  $k$ . The composite

$$T \longrightarrow T' \longrightarrow T$$

is the  $m$ -th power map on  $T$ , and is therefore dominant. Consequently  $T' \rightarrow T$  is dominant, proving that  $T$  is unirational.

Return to the hypotheses of 3.1, but suppose in addition that  $G$  locally admits a maximal torus for the fpqc topology (XII 7.1). Let  $Y$  be the maximal torus of the Cartan subgroup  $X$  of  $G_{\mathcal{C}}$ . The morphism  $\psi : X \rightarrow G$  then induces a morphism  $Y \rightarrow G$  whose set-theoretic image consists of the semisimple elements of the fibers of  $G$  (XII 8). Finally, 3.1 shows that the restriction of  $\psi$  to the open subset  $Y^{\text{reg}}$  of regular points of  $Y$  induces a closed immersion

$$Y^{\text{reg}} \longrightarrow U = G^{\text{reg}}.$$

Making explicit the functorial meaning of  $Z = Y^{\text{reg}}$ , we obtain the following.

**Corollary 0.12.5.** *Let  $G$  be a smooth, separated  $S$ -group prescheme of finite type, with connected fibers over the prescheme  $S$ , and suppose that  $G$  locally admits a maximal torus for the fpqc topology. Let*

$$Z : (\text{Sch})_S^{\circ} \longrightarrow (\text{Ens})$$

*be the functor defined by*

$$Z(S') = \left\{ \begin{array}{l} \text{regular sections of } G_{S'} \text{ over } S' \text{ that are} \\ \text{contained in a maximal torus of } G_{S'} \end{array} \right\}.$$

*Then  $Z$  is representable by a closed subscheme, smooth over  $S$ , of the open subset  $U = G^{\text{reg}}$  of  $G$  introduced in 3.1.*

To conclude, let us record the following result, which sharpens the density theorem 2.1 (i)  $\Rightarrow$  (vi).

**Corollary 0.12.6.** *Under the hypotheses of 3.5, let  $C$  be a Cartan subgroup of  $G$ , and consider the morphism*

$$\varphi : Z \times C \longrightarrow G$$

*defined by*

$$\varphi(g, h) = \text{ad}(g)h = ghg^{-1}.$$

*Then  $\varphi$  is dominant.*

It plainly suffices to prove the assertion fiber by fiber, which reduces us to the case where  $S$  is the spectrum of an algebraically closed field  $k$ . Let  $T$  be the maximal torus of  $C$ , let  $t_0 \in T(k)$  be regular in  $G$ , and let  $c_0 \in C(k)$  be regular in  $G$ . Consider  $\varphi^{-1}(\varphi(t_0, c_0))$ . Its  $k$ -rational points are the pairs  $(t, c)$ , with  $t \in Z(k)$  and  $c \in C(k)$ , such that

$$\text{ad}(t)c = \text{ad}(t_0)c_0, \quad \text{that is,} \quad c = \text{ad}(t^{-1}t_0)c_0.$$

They are therefore in bijection with the  $t \in Z(k)$  such that  $\text{ad}(t^{-1}t_0)c_0 \in C$ , or equivalently, since  $c_0$  is regular, such that  $t^{-1}t_0 \in N$ , where  $N$  is the normalizer of  $C$ ; in other words,  $t \in N$ . Restricting to those  $t \in Z(k)$  that belong to  $C(k)$  gives an open and closed part of this fiber. We have therefore found a connected component of  $\varphi^{-1}(\varphi(t_0, c_0))$  isomorphic to  $T$ . The preceding set-theoretic argument indeed gives an isomorphism of schemes: one simply replaces  $k$ -valued points by points with values in an arbitrary  $k$ -prescheme.

It follows that the generic fiber of  $\varphi$  has dimension at most  $\dim T$ , and therefore

$$\dim \operatorname{Im}(\varphi) \geq \dim(Z \times C) - \dim T = \dim Z + \dim C - \dim T.$$

But

$$\dim Z = \dim Y = \dim \mathcal{C} + \dim T = \dim G - \dim C + \dim T.$$

Thus  $\dim \operatorname{Im}(\varphi) \geq \dim G$ , and  $\varphi$  is dominant.

Q.E.D.

*Remarks 0.12.7.* The argument shows in addition that the connected component at  $(t_0, c_0)$  of the fiber  $\varphi^{-1}(\varphi(t_0, c_0))$  is isomorphic to  $T$ . In particular, it is smooth over  $k$  and has the same dimension as the generic fiber. It follows that  $\varphi$  is in fact smooth at  $(t_0, c_0)$ , something that one should also be able to verify by calculating the tangent map. Consequently, under the hypotheses of 3.6, the induced morphism

$$Z \times_S C^{\operatorname{reg}} \longrightarrow G^{\operatorname{reg}}, \quad C^{\operatorname{reg}} = C \cap G^{\operatorname{reg}},$$

is smooth. Likewise, the analogous morphism

$$Z \times T^{\operatorname{reg}} \longrightarrow Z,$$

where  $T$  is a maximal torus of  $G$ , is smooth. More generally, for every smooth connected normal algebraic subgroup  $H$  of  $C$  containing an element  $c_0 \in G(k)$  that is regular in  $G$ , the image of  $Z \times H \rightarrow G$  is dense in the image of  $G \times H \rightarrow G$ .

### 0.13 Lie algebras over a field: rank, regular elements, Cartan subalgebras

In the remainder of this Exposé we take up the theory developed by Chevalley in his book *Théorie des Groupes de Lie III* (Act. Sc. Ind. 1226, Paris 1955); the methods of schemes allow us to dispense with the hypothesis of characteristic zero. We begin by recalling in this section several well-known notions and results.

Let  $\mathfrak{g}$  be a Lie algebra over a ring  $k$ . For every  $a \in \mathfrak{g}$ , denote by  $\operatorname{ad}(a)$  the endomorphism

$$\operatorname{ad}(a) \cdot x = [a, x]$$

of  $\mathfrak{g}$ ; it is a derivation of the Lie algebra  $\mathfrak{g}$ . For every derivation  $D$  of  $\mathfrak{g}$ , the nil-space of  $D$ , that is, the union of the kernels of the iterates of  $D$ , is a Lie subalgebra of  $\mathfrak{g}$ , as follows from the Leibniz formula

$$D^m([x, y]) = \sum_{0 \leq p \leq m} \binom{m}{p} [D^p x, D^{m-p} y].$$

We set

$$\operatorname{Nil}(a, \mathfrak{g}) = \text{nil-space of } \operatorname{ad}(a) = \bigcup_{m \geq 0} \operatorname{Ker} \operatorname{ad}(a)^m.$$

When no confusion can arise, we write simply  $\operatorname{Nil}(a)$ , and call it the *nil-space of  $a$  in  $\mathfrak{g}$* .

**Proposition 0.13.1.** *For every  $a \in \mathfrak{g}$ , its nil-space  $\operatorname{Nil}(a, \mathfrak{g})$  is a Lie subalgebra of  $\mathfrak{g}$ , equal to its own normalizer.*



It remains only to prove that it is its own normalizer. In other words, every element of  $\mathfrak{g}/\text{Nil}(a, \mathfrak{g})$  annihilated by the adjoint representation of  $\text{Nil}(a, \mathfrak{g})$  on this quotient must be zero. This is immediate, since every element of the quotient annihilated by  $\text{ad}(a)$ <sup>48</sup> is zero.

For the remainder of this section, suppose that  $k$  is a field and that  $\mathfrak{g}$  is finite-dimensional over  $k$ . We denote by  $W(\mathfrak{g})$  the scheme over  $k$  defined by  $\mathfrak{g}$ ; its points in a  $k$ -algebra  $A$  are the elements of  $\mathfrak{g} \otimes_k A$ . If  $a \in \mathfrak{g}$ , the characteristic polynomial of  $\text{ad}(a)$  is also called the *characteristic polynomial*, or the *Killing polynomial*, of  $a$  in  $\mathfrak{g}$ :

$$P_{\mathfrak{g}}(a, t) = t^n + c_1(a)t^{n-1} + \cdots + c_n(a),$$

where  $n = \text{rank}_k \mathfrak{g}$  and  $c_i(a) \in k$ . Taking the same polynomial for  $a \in \mathfrak{g} \otimes_k A$ <sup>49</sup>, where  $A$  is any  $k$ -algebra, we see that the  $c_i(a)$  arise from uniquely determined sections  $c_i$  of the structure sheaf of  $W(\mathfrak{g})$ , that is, from elements of the symmetric algebra

$$A = \text{Sym}_k(\mathfrak{g}^\vee),$$

where  $\mathfrak{g}^\vee$  is the dual of the  $k$ -module  $\mathfrak{g}$ . When  $k$  is infinite, the  $c_i$  are determined by the corresponding polynomial functions  $\mathfrak{g} \rightarrow k$ , but this is no longer true when  $k$  is finite.

Let  $r$  be the largest integer such that the Killing polynomial

$$P_{\mathfrak{g}}(t) = t^n + c_1 t^{n-1} + \cdots + c_n \in A[t]$$

is divisible by  $t^r$ ; thus

$$P_{\mathfrak{g}}(t) = t^n + c_1 t^{n-1} + \cdots + c_{n-r} t^r, \quad c_{n-r} \neq 0.$$

The integer  $r$  is called the *nilpotent rank* of the Lie algebra  $\mathfrak{g}$ . It is invariant under extension of the base field.

**Proposition 0.13.2.** *Let  $r$  be the nilpotent rank of  $\mathfrak{g}$ , and let  $a \in \mathfrak{g}$ . Then*

$$\text{rank}_k \text{Nil}(a, \mathfrak{g}) \geq r,$$

*with equality if and only if*

$$c_{n-r}(a) \neq 0.$$

*In this case  $\text{Nil}(a, \mathfrak{g})$  is a nilpotent Lie algebra. We shall see the converse in 5.7 b) when  $\mathfrak{g}$  is the Lie algebra of an algebraic group  $G$  smooth over  $k$ .*

The first assertion is immediate, since by definition

$$\text{rank}_k \text{Nil}(a, \mathfrak{g}) = \text{the multiplicity of the zero root in } P_{\mathfrak{g}}(a, t).$$

Suppose now that  $c_{n-r}(a) \neq 0$ . We prove that  $\text{Nil}(a)$  is nilpotent, which is equivalent to saying that  $\text{ad}_{\text{Nil}(a)}(x)$  is a nilpotent endomorphism for every  $x \in \text{Nil}(a)$ . We may suppose that  $k$  is algebraically closed. Since

$$\text{ad}(a)_{\mathfrak{g}/\text{Nil}(a)}$$

is injective, there is a nonempty open subset  $U \subset W(\text{Nil}(a))$  such that, for every  $x \in U(k)$ , the endomorphism  $\text{ad}(x)_{\mathfrak{g}/\text{Nil}(a)}$  is injective and hence  $\text{Nil}(x) \subset \text{Nil}(a)$ . We may further

<sup>48</sup>The source prints  $\text{Ad}(a)$ ; because  $a \in \mathfrak{g}$ , the endomorphism here is  $\text{ad}(a)$ .

<sup>49</sup>Editor's note:  $W \otimes_k A$  has been corrected to  $\mathfrak{g} \otimes_k A$ .

suppose that  $U$  is contained in the open subset on which  $c_{n-r}$  does not vanish; that open subset is nonempty because  $c_{n-r}(a) \neq 0$ . Then  $\text{Nil}(x)$  and  $\text{Nil}(a)$  have the same dimension, so

$$\text{Nil}(x) = \text{Nil}(a).$$

It follows that  $\text{ad}(x)_{\text{Nil}(a)}$  is nilpotent for every  $x \in U(k)$ . By the principle of extension of algebraic identities, this remains true for every  $x \in \text{Nil}(a)$ , and hence  $\text{Nil}(a)$  is nilpotent.

An element  $a \in \mathfrak{g}$  is called *regular* if  $c_{n-r}(a) \neq 0$ , that is, if

$$\text{rank}_k \text{Nil}(a, \mathfrak{g}) = r.$$

When  $k$  is infinite, this also says that  $\text{rank}_k \text{Nil}(a, \mathfrak{g})$  is as small as possible as  $a$  varies in  $\mathfrak{g}$ . In every case, the notion of a regular element of  $\mathfrak{g}$  is invariant under extension of the base field. The set of regular points of  $W(\mathfrak{g})$ , meaning those that arise from regular points with values in a suitable extension of  $k$ , is open, since it is

$$W(\mathfrak{g})_{c_{n-r}},$$

the set of points at which  $c_{n-r}$  is invertible.

**Corollary 0.13.3.** *Let  $a$  be a regular element of  $\mathfrak{g}$ , and let  $\mathfrak{h}$  be a Lie subalgebra of  $\mathfrak{g}$  containing  $a$ . Then  $\mathfrak{h}$  is nilpotent if and only if*

$$\mathfrak{h} \subset \text{Nil}(a, \mathfrak{g}).$$

*In particular,  $\text{Nil}(a, \mathfrak{g})$  is a maximal nilpotent subalgebra of  $\mathfrak{g}$ .*

Since  $\text{Nil}(a)$  is nilpotent, the inclusion  $\mathfrak{h} \subset \text{Nil}(a)$  implies that  $\mathfrak{h}$  is nilpotent. Conversely, if  $\mathfrak{h}$  is nilpotent, it is contained in the nil-space of its element  $a$ ; that is,  $\mathfrak{h} \subset \text{Nil}(a)$ .

**Proposition 0.13.4.** *Suppose that  $k$  is infinite. Let  $\mathfrak{d}$  be a Lie subalgebra of  $\mathfrak{g}$ . Consider the following conditions:*

- (i)  $\mathfrak{d}$  is maximal nilpotent and contains a regular element of  $\mathfrak{g}$ .
- (i bis)  $\mathfrak{d} = \text{Nil}(a, \mathfrak{g})$  for some regular element  $a$  of  $\mathfrak{g}$ .
- (ii)  $\mathfrak{d}$  is nilpotent and  $\mathfrak{d} = \text{Nil}(a, \mathfrak{g})$  for some  $a \in \mathfrak{g}$ .
- (ii bis)  $\mathfrak{d}$  is nilpotent, and there exists  $a \in \mathfrak{d}$  such that  $\text{ad}(a)_{\mathfrak{g}/\mathfrak{d}}$  is injective.
- (iii)  $\mathfrak{d}$  is nilpotent and equal to its own normalizer.

Then

$$(i) \iff (i \text{ bis}) \implies (ii) \iff (ii \text{ bis}) \iff (iii).$$

We shall see in 5.7 a) that if  $\mathfrak{g}$  is the Lie algebra of a smooth algebraic group, then all the preceding conditions are equivalent.

The equivalence of (i) and (i bis) follows immediately from 4.3, and these conditions plainly imply (ii). The equivalence of (ii) and (ii bis) is also immediate, as is (ii bis)  $\implies$  (iii) (cf. 4.1). It remains to prove (iii)  $\implies$  (ii bis). This is the only implication that uses the infinitude of  $k$ , and it follows at once from the next lemma.

**Lemma 0.13.5.** *Let  $\mathfrak{d}$  be a nilpotent Lie algebra over an infinite field  $k$ , acting on a finite-dimensional vector space  $V$ . Suppose that the endomorphism  $u(x)$  is not injective for every  $x \in \mathfrak{d}$ . Then there is a nonzero element  $v \in V$  annihilated by  $\mathfrak{d}$ .*

We may suppose that  $k$  is algebraically closed and that  $\mathfrak{d}$  is finite-dimensional. Then  $V$  is known to be a direct sum of finitely many nonzero stable vector subspaces  $V_i$ ,  $1 \leq i \leq n$ , such that, for every  $i$  and every  $x \in \mathfrak{d}$ , the restriction  $u(x)|_{V_i}$  has a single eigenvalue  $\lambda_i(x)$  (cf. Bourbaki, *Groupes et Algèbres de Lie*, Chap. I, §4, Exercise 22). Let  $c_i(x)$  be the constant term of the characteristic polynomial of  $u(x)|_{V_i}$ , so that

$$\lambda_i(x) = 0 \iff c_i(x) = 0.$$

Each  $c_i$  is a polynomial function on  $\mathfrak{d}$ , and the hypothesis says that  $\mathfrak{d}$  is the union of their zero sets. Thus one of the  $c_i$  is identically zero. Replacing  $V$  by  $V_i$ , we are reduced to the case in which all  $u(x)$ ,  $x \in \mathfrak{d}$ , are nilpotent. Engel's theorem (Bourbaki, loc. cit., Th. 1) then implies that some nonzero  $v \in V$  is annihilated by  $\mathfrak{d}$ .

Q.E.D.

It is easy to see, still assuming that  $k$  is infinite, that conditions (i) and (i bis) of 4.4 are invariant under every extension of the base field. When they hold,  $\mathfrak{d}$  is called a *Cartan subalgebra* of  $\mathfrak{g}$ . In general, with  $k$  not necessarily infinite,  $\mathfrak{d}$  is called a Cartan subalgebra of  $\mathfrak{g}$  if it becomes one after one, and hence every, extension  $k \rightarrow k'$  with  $k'$  infinite. It follows in particular that  $\mathfrak{d}$  is nilpotent and equal to its own normalizer.

**Proposition 0.13.6.** a) *Let  $a \in \mathfrak{g}$ . If  $a$  is regular, then it is contained in one and only one Cartan subalgebra of  $\mathfrak{g}$ . We shall see the converse in 6.1 d), when  $k$  is algebraically closed and  $\mathfrak{g}$  is the Lie algebra of a smooth algebraic group.*

b) *Let  $\mathfrak{d}$  be a Cartan subalgebra of  $\mathfrak{g}$ , and let  $a \in \mathfrak{d}$ . Then  $a$  is regular in  $\mathfrak{g}$  if and only if  $\text{ad}(a)_{\mathfrak{g}/\mathfrak{d}}$  is injective.*

For a), if  $a$  is regular, then  $\text{Nil}(a, \mathfrak{g})$  is a Cartan subalgebra of  $\mathfrak{g}$ , since this is true after an infinite extension  $k'$  of  $k$ . It follows at once from 4.3 that every Cartan subalgebra of  $\mathfrak{g}$  containing  $a$  is equal to this one. For b), observe that the nullity<sup>50</sup> of  $\text{ad}(a)_{\mathfrak{g}}$  is the sum of the nullities of  $\text{ad}(a)_{\mathfrak{d}}$  and  $\text{ad}(a)_{\mathfrak{g}/\mathfrak{d}}$ . Since the first is  $r$ , their sum is  $r$  if and only if  $\text{ad}(a)_{\mathfrak{g}/\mathfrak{d}}$  is injective.

Q.E.D.

**Corollary 0.13.7.** *Let  $a$  be a regular element of  $\mathfrak{g}$ , let  $\mathfrak{d}$  be a Cartan subalgebra of  $\mathfrak{g}$  containing  $a$ , let  $A$  be a  $k$ -algebra, and let  $\mathfrak{g}_A$  and  $\mathfrak{d}_A \subset \mathfrak{g}_A$  be the  $A$ -Lie algebras obtained from  $\mathfrak{g}$  and  $\mathfrak{d}$  by base change; write  $a_A$  for the image of  $a$  in  $\mathfrak{g}_A$ . Let  $u$  be an automorphism of  $\mathfrak{g}_A$ . Then*

$$u(\mathfrak{d}_A) = \mathfrak{d}_A$$

*if and only if*

$$u(a_A) \in \mathfrak{d}_A.$$

The condition is plainly necessary; let us prove that it is sufficient. Suppose it holds. Then

$$\mathfrak{d}' = u^{-1}(\mathfrak{d}_A)$$

is a Lie subalgebra containing  $a_A$ , and every  $b \in \mathfrak{d}'$  is such that  $\text{ad}(b)_{\mathfrak{d}'}$  is nilpotent.<sup>51</sup> Indeed,  $\mathfrak{d}'$  is isomorphic to  $\mathfrak{d}_A$ , which has this property, as follows immediately from the definition of “nilpotent” in Bourbaki, *Groupes et Algèbres de Lie*, Chap. I, §4, Def. 1. Taking  $b = a_A$ , we see that  $\text{Nil}(b, \mathfrak{g}_A)$  contains  $\mathfrak{d}'$ . On the other hand,

$$\text{Nil}(b, \mathfrak{g}_A) = \mathfrak{d}_A.$$

<sup>50</sup>Editor's note: that is, the dimension of the nil-space.

<sup>51</sup>The source prints  $\mathfrak{d}' = u(\mathfrak{d}_A)$  and  $\text{ad}(b)_{\mathfrak{d}}$ . Under the hypothesis  $u(a_A) \in \mathfrak{d}_A$ , the inverse image  $u^{-1}(\mathfrak{d}_A)$  is the subalgebra that contains  $a_A$ , and the nilpotence assertion concerns its restriction  $\text{ad}(b)_{\mathfrak{d}'}$ .

Moreover,  $\mathfrak{d}'$  is locally a direct summand of  $\mathfrak{g}_A$ , since  $\mathfrak{d}$  is, and hence is locally a direct summand of  $\mathfrak{d}_A$ . It is a projective module of the same rank  $r$  as  $\mathfrak{d}_A$ , and therefore equals  $\mathfrak{d}_A$ .

Q.E.D.

**Proposition 0.13.8.** *Let  $\mathfrak{h}$  be a Lie subalgebra of  $\mathfrak{g}$ .*

a) *If  $k$  is infinite, the following conditions are equivalent:*

(i)  *$\mathfrak{h}$  contains a Cartan subalgebra  $\mathfrak{d}$  of  $\mathfrak{g}$ .*

(ii)  *$\mathfrak{h}$  contains a regular element  $a$  of  $\mathfrak{g}$ , and an element  $b$  such that  $\text{ad}(b)_{\mathfrak{g}/\mathfrak{h}}$  is injective.*

(iii)  *$\mathfrak{h}$  has the same nilpotent rank as  $\mathfrak{g}$ , and contains a regular element of  $\mathfrak{g}$ .*

*These conditions are invariant under extension of the base field  $k$ .*

b) *Suppose that these conditions hold after a suitable infinite extension  $k'$  of  $k$ . For  $a \in \mathfrak{h}$ , the element  $a$  is regular in  $\mathfrak{g}$  if and only if it is regular in  $\mathfrak{h}$  and  $\text{ad}(a)_{\mathfrak{g}/\mathfrak{h}}$  is injective; equivalently, if and only if*

$$\text{Nil}(a, \mathfrak{h}) = \text{Nil}(a, \mathfrak{g})$$

*and this common Lie algebra is a Cartan subalgebra of  $\mathfrak{h}$ .*

c) *Under the hypotheses of b), let  $\mathfrak{d}$  be a Lie subalgebra of  $\mathfrak{h}$ . In order that  $\mathfrak{d}$  be a Cartan subalgebra of  $\mathfrak{h}$ , it is sufficient that it be a Cartan subalgebra of  $\mathfrak{g}$ . We shall see in 5.8 that this condition is also necessary when  $\mathfrak{g}$  is the Lie algebra of a smooth algebraic group  $G$ , and  $\mathfrak{h}$  is the Lie algebra of a smooth algebraic subgroup  $H$  of  $G$ .*

Conditions (ii) and (iii) of a) are visibly invariant under extension of the base field  $k$ , assumed infinite. In b) and c), we may suppose that  $k$  is infinite, and shall do so. If  $\mathfrak{h}$  contains the Cartan subalgebra

$$\mathfrak{d} = \text{Nil}(a, \mathfrak{g}),$$

then  $a$  is a regular element of  $\mathfrak{g}$  such that  $\text{ad}(a)_{\mathfrak{g}/\mathfrak{h}}$  is injective. Thus (i)  $\Rightarrow$  (ii). Conversely, if (ii) holds, then a “sufficiently general” element  $a$  of  $\mathfrak{h}$  simultaneously satisfies both conditions in (ii). Hence  $\text{Nil}(a, \mathfrak{g})$  is a Cartan subalgebra of  $\mathfrak{g}$  contained in  $\mathfrak{h}$ , proving (i). Conditions (i) and (ii) are therefore equivalent.

Suppose they hold, and let  $a$  vary in  $\mathfrak{h}$ . Then

$$(x) \quad \text{rank}_k \text{Nil}(a, \mathfrak{g}) = \text{rank}_k \text{Nil}(a, \mathfrak{h}) + \text{rank}_k \text{Nil}(\text{ad}(a)_{\mathfrak{g}/\mathfrak{h}}).$$

The two terms on the right are respectively at least  $r'$ , the nilpotent rank of  $\mathfrak{h}$ , and at least 0; moreover, both equalities are attained<sup>52</sup> for a sufficiently general element of  $\mathfrak{h}$ . We also have

$$\text{rank}_k \text{Nil}(a, \mathfrak{g}) \geq r,$$

where  $r$  is the nilpotent rank of  $\mathfrak{g}$ . Equality is attained for a sufficiently general element of  $\mathfrak{h}$ , and holds if and only if  $a$  is regular in  $\mathfrak{g}$ .

It follows that  $r = r'$ , and that  $a$  is regular if and only if the two terms on the right-hand side of (x) are respectively  $r'$  and 0; that is, if and only if  $a$  is regular in  $\mathfrak{h}$  and  $\text{ad}(a)_{\mathfrak{g}/\mathfrak{h}}$  is injective. This proves b), and c) follows immediately by taking an element  $a \in \mathfrak{d}$  regular in  $\mathfrak{g}$ , so that  $\text{Nil}(a, \mathfrak{g}) = \mathfrak{d}$ .

The preceding result also proves (i)  $\Rightarrow$  (iii). Finally, (iii)  $\Rightarrow$  (i): under (iii), a sufficiently general  $a \in \mathfrak{h}$  is regular both in  $\mathfrak{h}$  and in  $\mathfrak{g}$ . Thus

$$\text{Nil}(a, \mathfrak{h}) \subset \text{Nil}(a, \mathfrak{g})$$

---

<sup>52</sup>Editor’s note: “the equality” has been replaced by “the equalities”.

are respectively Cartan subalgebras of  $\mathfrak{h}$  and  $\mathfrak{g}$ . Since they have the same rank over  $k$ , they are equal, proving (i). This completes the proof of 4.8.

## 0.14 The case of the Lie algebra of a smooth algebraic group: the density theorem

Let  $G$  be a smooth algebraic group over the field  $k$ , and let  $\mathfrak{g}$  be its Lie algebra. Let  $\mathfrak{h}$  be a Lie subalgebra of  $\mathfrak{g}$ . Let  $G$  act on  $W(\mathfrak{g})$  through the adjoint representation, and consider the subscheme  $W(\mathfrak{h})$ . The construction of Section 1 leads us to introduce

$$N = \text{Norm}_G(\mathfrak{h}) = \text{Norm}_G(W(\mathfrak{h})),$$

which is an algebraic subgroup of  $G$  (not necessarily smooth), and

$$\mathfrak{n} = \text{Lie}(N),$$

together with the scheme

$$X = G \times^N W(\mathfrak{h}),$$

the quotient of  $\tilde{X} = G \times W(\mathfrak{h})$  by the right action of  $N$  given by

$$(g, x) \cdot n = (gn, \text{Ad}(n^{-1})x).$$

We consider the canonical morphisms

$$\begin{array}{ccc} G \times W(\mathfrak{h}) & & \\ q \downarrow & \searrow \varphi & \\ X & \xrightarrow{\psi} & W(\mathfrak{g}). \end{array}$$

**Theorem 0.14.1.** *With the preceding notation, suppose that  $k$  is infinite. Consider the following conditions:*

- (i)  $\mathfrak{h}$  contains a Cartan subalgebra  $\mathfrak{d}$  of  $\mathfrak{g}$ .
- (ii) There exists  $a \in \mathfrak{h}$  such that  $\text{ad}(a)_{\mathfrak{g}/\mathfrak{h}}$  is injective.
- (iii) The morphism  $\varphi : G \times W(\mathfrak{h}) \rightarrow W(\mathfrak{g})$  is generically smooth.
- (iv) The preceding morphism  $\varphi$  (or, equivalently,  $\psi : X \rightarrow W(\mathfrak{g})$ ) is dominant, and  $\mathfrak{h}$  has the same nilpotent rank as  $\mathfrak{g}$ .
- (v) The morphism  $\psi : X \rightarrow W(\mathfrak{g})$  is generically smooth and  $\mathfrak{h} = \mathfrak{n}$ .
- (vi) The morphism  $\psi : X \rightarrow W(\mathfrak{g})$  is dominant, and  $\mathfrak{h} = \mathfrak{n}$ .
- (vii) The morphism  $\psi : X \rightarrow W(\mathfrak{g})$  is dominant.
- (viii) The morphism  $\psi : X \rightarrow W(\mathfrak{g})$  is dominant, and  $N$  is smooth.
- (ix) The morphism  $\psi : X \rightarrow W(\mathfrak{g})$  is dominant,  $\mathfrak{h} = \mathfrak{n}$ , and  $\mathfrak{h}$  is the Lie algebra of a smooth algebraic subgroup  $H$  of  $G$ .
- (x) The morphism  $\psi : X \rightarrow W(\mathfrak{g})$  is generically quasi-finite, and  $\mathfrak{n} = \mathfrak{h}$ .

- (xi) The morphism  $\psi : X \rightarrow W(\mathfrak{g})$  is generically étale, and  $\mathfrak{n} = \mathfrak{h}$ .
- (xii) There exists a smooth algebraic subgroup  $H$  of  $G$ , with Lie algebra  $\mathfrak{h}$ , such that  $\mathfrak{h}$  contains a Cartan subalgebra  $\mathfrak{d}$  of  $\mathfrak{g}$ .
- (xiii) There exists  $a \in \mathfrak{h}$  such that  $N$  and the transporter  $M_a$  of  $a$  into  $\mathfrak{h}$  (cf. Section 1) coincide in a neighborhood of  $e$ , and  $\mathfrak{n} = \mathfrak{h}$ .

One then has the following diagram of implications:

$$\begin{array}{ccccccccccc}
 \text{(xi)} & \longleftrightarrow & \text{(xii)} & \longleftrightarrow & \text{(xiii)} & \implies & \text{(viii)} & \longleftrightarrow & \text{(ix)} & \longleftrightarrow & \text{(x)} \\
 & & \Downarrow & & & & & & \Downarrow & & \\
 \text{(i)} & \longleftrightarrow & \text{(ii)} & \longleftrightarrow & \text{(iii)} & \longleftrightarrow & \text{(iv)} & \implies & \text{(v)} & \implies & \text{(vi)} \implies \text{(vii)}.
 \end{array}$$

When  $k$  has characteristic zero, all the conditions under consideration are equivalent. Finally,

$$\text{(xi)} \longleftrightarrow [(\text{i}) \text{ and } (\text{viii})] \longleftrightarrow [(\text{v}) \text{ and } (\text{viii})].$$

Let us first record the trivial implications

$$(\text{v}) \implies (\text{vi}) \implies (\text{vii}), \quad (\text{ix}) \implies (\text{vi}), \quad (\text{xi}) \longleftrightarrow [(\text{x}) \text{ and } (\text{v})].$$

We prove the equivalence of conditions (i) through (iv), and that they imply (v). The implication (i)  $\implies$  (ii) is immediate. On the other hand, when  $k$  is algebraically closed, condition (iii) says that there is a  $k$ -rational point of  $G \times W(\mathfrak{h})$  at which the tangent map of  $\varphi$  is surjective. It is clear that, after translating by a suitable element of  $G(k)$ , this point may be taken to have the form  $(e, a)$ , with  $a \in \mathfrak{h}$ . It follows that when  $k$  is infinite, though not necessarily algebraically closed, this condition—which is plainly sufficient for generic smoothness—is also necessary.

The tangent map is readily computed. Upon identifying the tangent space of  $W(\mathfrak{h})$  at  $a$  with  $\mathfrak{h}$ , it is the map

$$(\xi, x) \longmapsto [\xi, a] + x$$

from  $\mathfrak{g} \times \mathfrak{h}$  to  $\mathfrak{g}$ . Its surjectivity is equivalent to the surjectivity of  $\text{ad}(a)_{\mathfrak{g}/\mathfrak{h}}$ , or, what amounts to the same thing, to its injectivity. This proves the equivalence of (ii) and (iii). Moreover, (ii) plainly implies  $\mathfrak{h} = \mathfrak{n}$ , and (iii) implies that  $\psi$  is generically smooth: if  $\varphi$  is smooth at a point  $u$ , then, since  $q$  is flat,  $\psi$  is smooth at  $q(u)$ . Thus (ii) and (iii) imply (v).

We now show that they imply (i). Since  $\psi$  is dominant and the set of regular points of  $\mathfrak{g}$  is dense and open,  $\mathfrak{h}$  contains regular elements of  $\mathfrak{g}$ . Hence a sufficiently general element  $b$  of  $\mathfrak{h}$  is regular in  $\mathfrak{g}$  and makes  $\text{ad}(b)_{\mathfrak{g}/\mathfrak{h}}$  injective. Consequently  $\text{Nil}(b, \mathfrak{g}) \subset \mathfrak{h}$ , so  $\mathfrak{h}$  contains the Cartan subalgebra  $\mathfrak{d} = \text{Nil}(b, \mathfrak{g})$ . Conditions (i), (ii), and (iii) are therefore equivalent. Finally, (i)  $\iff$  (iv): we have already observed that if  $\mathfrak{h}$  contains a Cartan subalgebra, then it has the same nilpotent rank as  $\mathfrak{g}$  (4.6), so (i) implies (iv). Conversely, if (iv) holds, then  $\mathfrak{h}$  contains a regular element of  $\mathfrak{g}$ , and since it has the same rank as  $\mathfrak{g}$ , it contains a Cartan subalgebra by 4.6.

We next prove the equivalence of conditions (viii) through (x). We first note the following facts.

**Lemma 0.14.2.** *a) If  $N$  is smooth, then*

$$\dim X \leq \dim G,$$

*with equality if and only if  $\mathfrak{h} = \mathfrak{n}$ .*

*b) If  $\mathfrak{h} = \mathfrak{n}$ , then*

$$\dim X \geq \dim G,$$

*with equality if and only if  $N$  is smooth.*

These assertions follow at once from the formula

$$\dim X = \dim G - \dim N + \dim_k \mathfrak{h}$$

and from the fact that  $\dim N = \dim_k \mathfrak{n}$  is equivalent to the smoothness of  $N$ .

Now (viii) implies (x): by dominance, (viii) gives  $\dim X \geq \dim G$ , so Lemma 5.2(a) gives equality and  $\mathfrak{h} = \mathfrak{n}$ , hence (x). The same argument in reverse, using Lemma 5.2(b), shows that (x) implies (viii). Moreover, (viii) implies (ix), since it implies  $\mathfrak{h} = \mathfrak{n}$ , so  $\mathfrak{h}$  is the Lie algebra of the smooth algebraic subgroup  $N$  of  $G$ . Conversely, (ix) implies (viii): since  $H$  normalizes its Lie algebra  $\mathfrak{h}$ , it is contained in  $N$ ; and since  $H$  is smooth and has the same Lie algebra as  $N$ , it follows that  $N$  is smooth.

Finally, we prove the equivalence of (xi), (xii), and (xiii), and show that they imply (iii); this will complete the implication diagram. We have (xi)  $\iff$  (xiii). Indeed, if  $\mathfrak{n} = \mathfrak{h}$ , then Lemma 5.2(b) gives  $\dim X \geq \dim G$ . Since  $W(\mathfrak{g})$  is normal, condition (xi) is therefore equivalent to  $\psi$  being generically unramified, and this in turn is equivalent to (xiii) by 1.1, (ii)  $\iff$  (iii), arguing as above in the proof of (ii)  $\iff$  (iii).

Since (xi)  $\Rightarrow$  (x)  $\Rightarrow$  (viii), what we have already proved shows that (xi) implies that  $N$  is smooth. Thus  $q : G \times W(\mathfrak{h}) \rightarrow X$  is smooth, and the composite  $\varphi = \psi \circ q$  is generically smooth; this is (iii). Since (iii) implies (i), it also follows that (xi)  $\Rightarrow$  (xii).<sup>53</sup> Finally, (xii) implies (xi). Indeed, (xii) plainly implies (i), and we have seen that (i)  $\Rightarrow$  (iii)  $\Rightarrow$  (v); hence (xii) implies (v). It also implies (ix), and (ix) implies (x). Thus (xii) implies both (v) and (x), and therefore (xi), since

$$\text{generically étale} = \text{generically smooth} + \text{generically quasi-finite}.$$

If  $k$  has characteristic zero, then (vii) implies (viii), because Cartier's theorem makes  $N$  automatically smooth (VI<sub>B</sub> 1.6.1). Moreover, [(viii) and (x)] implies (xi), because in characteristic zero, for a morphism of integral preschemes,

$$\text{generically étale} = \text{dominant and generically quasi-finite}.$$

Thus in this case all conditions (i) through (xiii) are equivalent.

**Corollary 0.14.3.** *Under the equivalent conditions (viii) through (x), there exists a unique smooth connected algebraic subgroup  $H$  of  $G$  whose Lie algebra is  $\mathfrak{h}$ , and*

$$\text{Norm}_G(H) = \text{Norm}_G(\mathfrak{h}) = N, \quad H = N^0.$$

Indeed,  $H = N^0$  has the required properties. Conversely, if  $H$  has these properties, then  $H \subset N$ , since  $H$  normalizes its Lie algebra  $\mathfrak{h}$ . This is an inclusion of smooth groups with the same Lie algebra, and  $H$  is connected, so  $H = N^0$ . To prove  $\text{Norm}_G(H) =$

<sup>53</sup>Editor's note: because  $H = N$  will do.

$\text{Norm}_G(\mathfrak{h})$ , we may assume that  $k$  is algebraically closed. What we have just proved then shows immediately that the two groups have the same  $k$ -points. On the other hand, the inclusions

$$H \subset \text{Norm}_G(H) \subset N$$

show that  $\text{Norm}_G(H)$  and  $N$  have the same Lie algebra; hence they are equal.

**Corollary 0.14.4.** *Under the equivalent conditions (i) through (iv), let  $a \in \mathfrak{h}$ . Then the following conditions are equivalent, and they hold whenever  $a$  is regular in  $\mathfrak{g}$ :*

- (i)  $\varphi$  is smooth at  $(e, a)$ .
- (ii)  $M_a = \text{Transp}_G(a, \mathfrak{h})$  is smooth at  $e$ , and  $\dim_e(M_a) = \text{rank}_k \mathfrak{h}$ .
- (iii)  $\text{ad}(a)_{\mathfrak{g}/\mathfrak{h}}$  is injective (equivalently, bijective).

*Under the equivalent conditions (xi) through (xiii), let  $H$  be the algebraic subgroup of  $G$  considered in Corollary 5.3. The preceding conditions are also equivalent to:*

- (iv)  $\psi$  is étale at  $(\bar{e}, a)$ .
- (v) If  $M_a^0$  denotes the connected component of  $e$  in the transporter  $M_a$  of  $a$  into  $\mathfrak{h}$ , endowed with the structure induced by  $M_a$ , then

$$H = M_a^0.$$

Plainly (i) implies (ii), since  $M_a$  is isomorphic to the fiber  $\varphi^{-1}(a)$ , with  $e$  corresponding to  $(e, a)$ . Conversely, (ii) implies (i): condition (ii) says that  $\varphi$  is “equidimensional” at  $(e, a)$ , in the sense that the dimension of the fiber through this point is the dimension of the generic fiber. Since  $G \times W(\mathfrak{h})$  and  $W(\mathfrak{g})$  are regular, this implies that  $\varphi$  is flat at  $(e, a)$ , hence smooth there because its fiber is smooth at that point. The equivalence of (i) and (iii) was obtained in the proof of Theorem 5.1 by the elementary calculation of the tangent map. We also saw in 4.8(b) that regularity of  $a$  in  $\mathfrak{g}$  implies (iii).

Under conditions (xi) through (xiii), the morphism  $q : G \times W(\mathfrak{h}) \rightarrow X$  is smooth, since  $N$  is smooth. Thus (i) is equivalent to the smoothness of  $\psi$  at  $(\bar{e}, a)$ ; and since  $\psi$  is generically étale, this is equivalent to (iv). Finally, as was noted at the end of the proof of 1.1, condition (iv) implies that  $N$  is the prescheme induced on  $M_a$  by an open and closed subset of  $M_a$ , which gives (v). Conversely, (v) trivially implies (ii). Alternatively, (v) implies (iv) by 1.1, since  $\psi$  is dominant and  $W(\mathfrak{g})$  is normal, and “unramified” is equivalent here to “étale”. This completes the proof of Corollary 5.4.

**Corollary 0.14.5.** *Let  $G$  be a smooth algebraic group over a field  $k$ , and let  $H$  be a smooth algebraic subgroup such that its Lie algebra  $\mathfrak{h}$  contains, at least after a suitable extension of  $k$ , a Cartan subalgebra of the Lie algebra  $\mathfrak{g}$  of  $G$ . Let  $K$  be a connected algebraic subgroup of  $G$ , not necessarily smooth, with Lie algebra  $\mathfrak{k}$ , and suppose that  $\mathfrak{k}$  contains a regular element  $a$  of  $\mathfrak{g}$ , again at least after a suitable extension of  $k$ . Then  $H$  contains  $K$  if and only if  $\mathfrak{h}$  contains  $\mathfrak{k}$ .*

Indeed, Corollary 5.4, (iii) $\Rightarrow$ (v), applied to the connected subgroup supplied by Corollary 5.3, gives  $H^0 = M_a^0$ .<sup>54</sup> This identity is invariant under extension of the base field, so it remains valid without assuming that the base field is infinite. Moreover,  $\mathfrak{k} \subset \mathfrak{h}$  plainly implies  $K \subset M_a$ . Since  $K$  is connected,  $K \subset M_a^0 \subset H$ .  $\square$

<sup>54</sup>The French source prints  $H = M_a^0$  here. In the stated generality the argument gives  $H^0 = M_a^0$ , which is all that is needed.



**Corollary 0.14.6.** *Let  $G$  and  $H$  be as in Corollary 5.5, and let  $K$  be an algebraic subgroup of  $G$ . Suppose that  $H$  is connected and  $K$  is smooth. Then  $K$  contains  $H$  if and only if  $\mathfrak{k}$  contains  $\mathfrak{h}$ .*

Indeed, if  $\mathfrak{k} \supset \mathfrak{h}$ , then  $K$  satisfies, with respect to  $H$ , the hypothesis imposed on  $H$  in Corollary 5.5. Meanwhile  $H$  plainly satisfies, with respect to  $K$ , the condition imposed on  $K$  there. The conclusion follows from Corollary 5.5.

**Corollary 0.14.7.** *Let  $\mathfrak{g}$  be the Lie algebra of a smooth algebraic group  $G$  over a field  $k$ . Then:*

- a) A Lie subalgebra  $\mathfrak{d}$  of  $\mathfrak{g}$  is a Cartan subalgebra if and only if it is nilpotent and equal to its own normalizer.*
- b) An element  $a \in \mathfrak{g}$  is regular if and only if  $\text{Nil}(a, \mathfrak{g})$  is nilpotent.*

After extending the base field, we may assume that  $k$  is infinite. In view of 4.4, to prove (a) it remains only to show that if  $\mathfrak{d}$  is nilpotent and contains an element  $a$  for which  $\text{ad}(a)_{\mathfrak{g}/\mathfrak{d}}$  is injective, then  $\mathfrak{d}$  is a Cartan subalgebra. By Theorem 5.1, (ii) $\Rightarrow$ (i),  $\mathfrak{d}$  contains a Cartan subalgebra

$$\mathfrak{d}' = \text{Nil}(a, \mathfrak{g}).$$

Since  $\mathfrak{d}$  is nilpotent, 4.3 now gives  $\mathfrak{d} = \mathfrak{d}'$ . For (b), one notes that  $\text{Nil}(a, \mathfrak{g})$  is a Cartan subalgebra of  $\mathfrak{g}$  by (a), and therefore  $a$  is regular.

**Corollary 0.14.8.** *Let  $G$  be a smooth algebraic group over a field  $k$ , and let  $H$  be a smooth algebraic subgroup, with Lie algebras  $\mathfrak{g}$  and  $\mathfrak{h}$ , respectively. Suppose that after a suitable extension of the base field,  $\mathfrak{h}$  contains a Cartan subalgebra of  $\mathfrak{g}$ . Let  $\mathfrak{d}$  be a Lie subalgebra of  $\mathfrak{h}$ . Then  $\mathfrak{d}$  is a Cartan subalgebra of  $\mathfrak{h}$  if and only if it is a Cartan subalgebra of  $\mathfrak{g}$ .*

In view of 4.8(c), it remains to show that if  $\mathfrak{d}$  is a Cartan subalgebra of  $\mathfrak{h}$ , then it is one of  $\mathfrak{g}$ . It is enough to show that  $\mathfrak{d}$  contains an element  $a$  regular in  $\mathfrak{g}$ , and we may assume that  $k$  is algebraically closed. But there is a dense open subset of  $\mathfrak{h}$  consisting of points regular in  $\mathfrak{g}$ , so the assertion follows from Theorem 5.1, (i) $\Rightarrow$ (vii), applied to  $(\mathfrak{h}, \mathfrak{d})$ .

## 0.15 Cartan subalgebras and subgroups of type (C) associated with a smooth algebraic group

For simplicity, in the following theorem we restrict ourselves to an algebraically closed base field; the case of an arbitrary base prescheme will be treated in the next Exposé.

**Theorem 0.15.1.** *Let  $G$  be a smooth algebraic group over an algebraically closed field  $k$ , and let  $\mathfrak{g}$  be its Lie algebra. Then:*

- a) The Cartan subalgebras of  $\mathfrak{g}$  are conjugate.*
- b) Let  $\mathfrak{d}$  be a Cartan subalgebra of  $\mathfrak{g}$ . Its normalizer  $N$  in  $G$  is smooth, and  $D = N^0$  is the unique smooth connected subgroup of  $G$  whose Lie algebra is  $\mathfrak{d}$ . One has*

$$\text{Norm}_G(\mathfrak{d}) = \text{Norm}_G(D) = N, \quad \text{and hence} \quad D = \text{Norm}_G(D)^0.$$

c) With  $\mathfrak{d}$  as in (b), set, as in Section 5,

$$X = G \times^N W(\mathfrak{d}),$$

and consider the canonical morphism

$$\psi : X \longrightarrow W(\mathfrak{g}).$$

Its fiber over  $a \in \mathfrak{g}$  has as its  $k$ -valued points the set of Cartan subalgebras of  $\mathfrak{g}$  containing  $a$ . Then  $\psi$  is birational.

d) With the notation of (c), let  $U$  be the largest open subset of  $W(\mathfrak{g})$  for which  $\psi$  induces an isomorphism

$$\psi^{-1}(U) \xrightarrow{\sim} U.$$

Then, for  $a \in \mathfrak{g}$ , the following conditions are equivalent:

- (i)  $a \in U(k)$ .
- (ii) The element  $a$  belongs to one and only one Cartan subalgebra of  $\mathfrak{g}$ .
- (iii) The set of Cartan subalgebras of  $\mathfrak{g}$  containing  $a$  is finite and nonempty.
- (iv) The fiber  $\psi^{-1}(a)$  has an isolated point.
- (v) If  $a \in \mathfrak{d}$ , the morphism  $\psi$  is étale—or merely quasi-finite—at the point  $(\bar{e}, a)$ .
- (vi) The element  $a$  is regular in  $\mathfrak{g}$ .
- (vii) With the notation of Corollary 5.4(iii), one has  $N = M_a$ .

*Proof.* Apply Theorem 5.1 with the Cartan subalgebra  $\mathfrak{d}$  of  $\mathfrak{g}$  in place of  $\mathfrak{h}$ . We show that the strongest conditions, (xi) through (xiii), hold. This is clear either in the form (xiii), by 4.7—which says that if a regular element  $a \in \mathfrak{g}$  lies in  $\mathfrak{d}$ , then  $N = M_a$ —or in the form

$$(xi) = (x) + (i).$$

Indeed, (i) is automatic, while (x) follows from the fact that a regular point of  $\mathfrak{g}$  belongs to a unique Cartan subalgebra of  $\mathfrak{g}$  (4.6(a)), and therefore, a fortiori, to a unique conjugate of  $\mathfrak{d}$ .

Part (b) now follows from Corollary 5.3. Part (c) follows because  $\psi$  is generically étale and because a sufficiently general point of  $\mathfrak{g}$ —more precisely, a regular point—belongs to a unique Cartan subalgebra of  $\mathfrak{g}$ . Under these conditions, the equivalence of conditions (i) through (v) on  $a$  follows immediately from Zariski's Main Theorem, applied to the separated birational morphism  $\psi$ , since  $W(\mathfrak{g})$  is normal and  $X$  is integral. The equivalence of (v) and (vi) is the special case of Corollary 5.4, (iii)  $\iff$  (iv), obtained after carrying  $a$  into  $\mathfrak{d}$  by a suitable element  $g \in G(k)$ , in view of 4.6(b). Moreover, Corollary 5.4 shows that (v) and (vi) are also equivalent to

$$N \supset M_a^0.$$

By 4.7 this even implies  $N = M_a$ .<sup>55</sup> This proves (d).

Of course, (b), (c), and the equivalence of (i), (iv), (v), (vi), and (vii) remain valid over an arbitrary base field. We now prove (a), using the assumption that  $k$  is algebraically closed. By Theorem 5.1, (i)  $\implies$  (vii), the morphism  $\psi : X \rightarrow W(\mathfrak{g})$  is dominant. Hence

<sup>55</sup>The French source prints  $N = M$  at this point; condition (vii) and the surrounding notation require  $N = M_a$ .

there is a dense open subset  $V$  of  $W(\mathfrak{g})$  such that every  $a \in V(k)$  is the image of a point of  $X(k)$ , or equivalently, belongs to a conjugate of  $\mathfrak{d}$ . Apply the same result to a second Cartan subalgebra  $\mathfrak{d}'$  of  $\mathfrak{g}$ . We may choose  $V$  so that every element of  $V(k)$  is conjugate both to an element of  $\mathfrak{d}$  and to an element of  $\mathfrak{d}'$ . Choose a regular element of  $V(k)$ . Some conjugate  $\mathfrak{d}''$  of  $\mathfrak{d}'$  then contains an element of  $\mathfrak{d}$  regular in  $\mathfrak{g}$ , and 4.6(a) gives  $\mathfrak{d}'' = \mathfrak{d}$ . This proves (a) and completes the proof.  $\square$

**Definition 0.15.2.** Let  $G$  be a smooth algebraic group over a field  $k$ . A *subgroup of type (C)* of  $G$  is a smooth connected algebraic subgroup whose Lie algebra is a Cartan subalgebra of  $\mathfrak{g}$ . The *infinitesimal rank* of  $G$  is the nilpotent rank of its Lie algebra  $\mathfrak{g}$ ; it is equal to the dimension of every subgroup of type (C) of  $G$ .

Theorem 6.1(b) immediately gives:

**Corollary 0.15.3.** *Under the hypotheses of Definition 6.2, the map*

$$D \longmapsto \mathfrak{d} = \text{Lie}(D)$$

*establishes a bijection between the subgroups of type (C) of  $G$  and the Cartan subalgebras  $\mathfrak{d}$  of  $\mathfrak{g}$ . If  $D$  is a subgroup of type (C) of  $G$ , then  $D$  is its own connected normalizer:*

$$D = \text{Norm}_G(D)^0.$$

Combining Theorem 6.1(a) and Corollary 6.3 gives:

**Corollary 0.15.4.** *If  $k$  is algebraically closed, the subgroups of type (C) of  $G$  are conjugate to one another.*

**Corollary 0.15.5.** *Let  $G$  be a smooth algebraic group over an algebraically closed field  $k$ , and let  $H$  be a smooth algebraic subgroup of  $G$ , with Lie algebras  $\mathfrak{g}$  and  $\mathfrak{h}$ , respectively. Then  $\mathfrak{h}$  contains a Cartan subalgebra of  $\mathfrak{g}$  if and only if  $H$  contains a subgroup  $D$  of type (C) of  $G$ .*

The condition is plainly sufficient. It is also necessary, since by Corollary 5.5,

$$H \supset D \iff \mathfrak{h} \supset \mathfrak{d}.$$

**Remarks 0.15.6.**

- a) The connected subgroups  $H$  of  $G$  described in Corollary 6.5 correspond bijectively to the Lie subalgebras  $\mathfrak{h}$  of  $\mathfrak{g}$  satisfying the strongest condition, (xii), of Theorem 5.1. By Corollary 5.5, inclusion relations among such subgroups can already be read from their Lie algebras.
- b) Continue to suppose that  $k$  is algebraically closed, and let  $D$  be a subgroup of type (C) of  $G$ . It is then easy to show that  $D$  contains a Cartan subgroup  $C$  of  $G$ . Indeed, put  $V = \mathfrak{g}/\mathfrak{d}$ . For a general element  $a \in \mathfrak{d}$ , the endomorphism  $\text{ad}(a)$  of  $V$  is injective; it suffices to take  $a$  regular in  $\mathfrak{g}$ . From this one readily deduces that, for a general  $g \in D(k)$ , the action of  $\text{Ad}(g)$  on  $V$  has no nonzero invariant vector. Thus 2.1, (vii) $\Rightarrow$ (i), applies. In fact, in the next Exposé we shall prove the more precise result that every Cartan subgroup  $C$  of  $G$  is contained in one and only one subgroup of type (C) of  $G$ .

- c) In general one must not confuse Cartan subgroups with subgroups of type  $(C)$ . The subgroups of type  $(C)$  of  $G$  are Cartan subgroups if and only if they are nilpotent, since Cartan subgroups are maximal nilpotent subgroups. It may happen that  $\mathfrak{g}$  is nilpotent although  $G$  is not; an example is  $\mathrm{SL}(2)_k$  when  $k$  has characteristic 2. Then the Cartan subalgebras of  $\mathfrak{g}$  are all equal to  $\mathfrak{g}$ . Thus, if  $G$  is connected,  $G$  is a subgroup of type  $(C)$  of itself, but it is not a Cartan subgroup of itself.

On the other hand, we shall see in Exposé XIV that if  $G$  is an adjoint semisimple group, its subgroups of type  $(C)$  are precisely its Cartan subgroups. The same conclusion holds in characteristic zero for every smooth algebraic group  $G$  over  $k$ . This follows at once from the fact that, in characteristic zero, a connected algebraic group is nilpotent if and only if its Lie algebra is nilpotent. In turn, this follows from the fact that in characteristic zero the center  $Z$  of a connected algebraic group  $G$  has Lie algebra equal to the center of  $\mathfrak{g}$ . Indeed, one obtains a priori the space of invariants in  $\mathfrak{g}$  under the adjoint representation of  $G$ . Since  $G$  is connected and  $k$  has characteristic zero, these are also the infinitesimal invariants of the adjoint representation of  $\mathfrak{g}$ , namely the center of  $\mathfrak{g}$ . Finally, Cartier's theorem (cf. VI<sub>B</sub> 1.6.1) shows that  $Z$  is smooth.

# Current translation boundary

This reader ends after the complete current Exposé XIII working translation. Exposés XIV–XXVI are not included. Later additive successors may replace this reader as further exposés become readable; exhaustive reference, diagram, privacy, rights, and release certification remains tracked separately from this preservation-oriented reading surface.