

`usepackage[a4paper,margin=0.65in]geometry`

Arthur Cayley

Collected Mathematical Papers

Volume IX, Pages 456–478 (Papers 610 conclusion & 611)

Transcribed from the Original Publications

Contents

- | | |
|---|---|
| 610. On the Analytical Forms called Trees (<i>concluded, pp. 456–460</i>) | 3 |
| 611. Report of the Committee on Mathematical Tables
(pp. 461–478) | 8 |

610. On the Analytical Forms called Trees, with Application to the Theory of Chemical Combinations (*concluded*)

[Conclusion of the paper begun at p. 427. Pages 456–458 give Table VIII (Carbon Centre- and Bicentre-Trees) and the Subsidiary Table for the generating functions of Tables VII and VIII. Pages 459–460 give Tables A and B (general valency tables) and the explanatory text leading to the chemical interpretation.]

[p. 456]

Table VIII. — Carbon Centre- and Bicentre-Trees.

Column headings: “Index of t , or number of main branches” and “Index of x , or number of knots”. For each value of the number of knots, the entries in the upper portion (Centre-Trees) are arranged by altitude (columns 1... 6) and the row marked “Total” shows the grand total for that knot count. The right-hand portion (Bicentre-Trees) is arranged by altitude (columns 1... 5) and “1” (the bicentre column).

n	a	Centre-Trees (altitude)									Bicentre-Trees (altitude)				
		1	2	3	4	5	6	Cent.	G.T.	Bic.	1	2	3	4	5
1	0	1						1	1	0					
Total		1						1	1	0					
2								0	1	1	1				
3	2	1	1					1	1	0					
Total		1	1					1	1	0					
4	2	1	1					1	2	1	1				
Total		1	1					1	2	1	1				
5	2	1		2				2	3	1	1				
Total		1		2				2	3	1	1				
6	2	1	1												
6	3	1	1												
Total		2	2					2	5	3	2	1			
7	2		2	1	3										
7	3		2		2										
7	4		1		1										
Total			5	1	6			6	9	3	1	2			
8	2		1	2	3										
8	3		3	1	4										
8	4		2		2										
Total			6	3	9			9	18	9	1	7	1		
9	2		1	7	1										

n	a	1	2	3	4	5	6	Cent.	G.T.	Bic.	1	2	3	4	5
9	3		3	3											
9	4		4	1											
Total			8	11	1	20		20	35	15		12	3		
10	2			12	3	15									
10	3		3	11	1	15									
10	4		4	3		7									
Total			7	26	4	37		37	75	38		23	14	1	
11	2			23	14	1	38								
11	3		2	24	4		30								
11	4		5	12	1		18								
Total			7	59	19	1	86	86	159	73		30	39	4	
12	2			30	39	5	74								
12	3		1	54	19	1	75								
12	4		4	27	4		35								
Total			5	111	62	5	183	183	357	174		42	108	23	1
13	2			42	108	23	1	174							
13	3		1	88	63	5		157							
13	4		4	63	20	1		88							
Total			5	193	191	29	1	419	799	380		47	244	84	5

[p. 457]

Subsidiary Table for GF's of Tables VII. and VIII.

The table is arranged by “column” (i.e. the number which is in successive use for the analytical work of Tables VII and VIII). For each column there is a generating function (“GF”), and that GF is built up from a “first factor” and a “second factor”. Within each first and second factor the rows are indexed by the exponent of t ($= 1, 2, 3, \dots$, the number of main branches), and the columns are indexed by the exponent of x ($= 1, 2, \dots, 13$, the number of knots). The entries are the coefficients.

The table extends across pp. 457–458 (*continued*). We reproduce the salient lines, preserving Cayley’s classification of GF, first factor, second factor for each “column” value.

GF, column 0: 1 (a single entry under index $x = 1$).

GF, column 1: -1 (under index $x = 1$).

First factor (column 1, label (1)): row $t = 1$: 1 at $x = 1$.

Second factor (column 1): row $t = 1$: 1 at $x = 2$; 1 at $x = 3$; 1 at $x = 4$.

GF, column 2: $-1, -1, -1$ under indices $x = 3, 4, 5$.

First factor (column 2): row $t = 1$: 1, 1, 1 at $x = 3, 4, 5$; row $t = 2$: 1, 1, 2, 1, 1 at $x = 5, 6, 7, 8, 9$; row $t = 3$: 1, 1, 2, 2, 2, 1, 1 at $x = 7, 8, 9, 10, 11, 12, 13$; row $t = 4$: 1, 1, 2, 2, 3 at $x = 9, 10, 11, 12, 13$.

Second factor (column 2): row $t = 1$: 1 at $x = 2$; row $t = 2$: 1 at $x = 3$; row $t = 3$: 1 at $x = 4$.

GF, column 3: $-1, -2, -4, -4, -6, -4, -4, -3, -2, -1, -1$ under indices $x = 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13$.

First factor (column 3): row $t = 1$: 1, 2, 4, 4, 5, 4, 4, 3, 2, 1 at $x = 3, \dots, 12$; row $t = 2$: 1, 2, 4, 12, 23, 30, 42 at $x = 7, \dots, 13$; row $t = 3$: 1, 2, 7, 16 at $x = 10, \dots, 13$; row $t = 4$: 1 at $x = 13$.

Second factor (column 3): row $t = 1$: 1, 1, 1, 1 at $x = 2, 3, 4, 5$; row $t = 2$: 1, 1, 2, 2, 2, 1, 1 at $x = 3, 4, 5, 6, 7, 8, 9$; row $t = 3$: 1, 1, 2, 3, 3, 3, 3, 2, 1, 1 at $x = 4, \dots, 13$.

GF, column 4: $-1, -3, -8, -15, -27, -43, -67, -97, -136, -183$ under indices $x = 4, 5, \dots, 13$.

First factor (column 4): row $t = 1$: 1, 3, 8, 15, 27, 43, 67, 97, 136 at $x = 4, \dots, 12$; row $t = 2$: 1, 3, 14, 39, 108 at $x = 9, \dots, 13$; row $t = 3$: 1 at $x = 13$.

Second factor (column 4): row $t = 1$: 1, 1, 2, 3, 4, 4, 5, 4, 4, 3, 2, 1 at $x = 2, \dots, 13$; row $t = 2$: 1, 1, 3, 5, 10, 14, 23, 29, 40, 46, 55 at $x = 3, \dots, 13$; row $t = 3$: 1, 3, 6, 12, 20, 37, 56, 89, 128 at $x = 4, 5, \dots, 13$ (with a gap at $x = 5$).

GF, column 5: $-1, -4, -13, -32, -74, -155, -316, -612, -1160$ under indices $x = 5, \dots, 13$.

First factor (column 5): row $t = 1$: 4, 13, 32, 74, 155, 316, 612 at $x = 6, \dots, 12$; row $t = 2$: 1, 4, 23 at $x = 11, 12, 13$.

Second factor (column 5): row $t = 1$: 1, 1, 2, 7, 12, 20, 31, 47, 70, 99, 137 at $x = 2, \dots, 13$ (with a gap); row $t = 2$: 1, 1, 3, 6, 14, 27, 56, 103, 194, 343, 605 at $x = 3, \dots, 13$ (with a gap); row $t = 3$: 1, 1, 3, 7, 16, 34, 76, 151, 307, 602 at $x = 4, \dots, 13$.

GF, column 6: $-1, -6, -19, -56, -151, -374, -889, -2032$ under indices $x = 6, \dots, 13$.

First factor (column 6): row $t = 1$: 1, 5, 19, 56, 151, 374, 889 at $x = 7, \dots, 13$; row $t = 2$: 1 at $x = 13$.

Second factor (column 6): row $t = 1$: 1, 1, 2, 4, 8, 16, 33, 63, 121, 225, 415, 749 at $x = 2, \dots, 13$; row $t = 2$: 1, 1, 3, 6, 15, 32, 75, 160, 350, 732, 1534 at $x = 3, \dots, 13$; row $t = 3$: 1, 1, 3, 7, 17, 39, 95, 214, 491, 1093 at $x = 4, \dots, 13$.

[p. 458] *Subsidiary Table for GF's of Tables VII. and VIII. (continued).*

GF, column 7: $-1, -6, -26, -88, -267, -743, -1968$ under indices $x = 7, \dots, 13$.

First factor (column 7): row $t = 1$: 1, 6, 26, 88, 267, 743 at $x = 8, \dots, 13$.

Second factor (column 7): row $t = 1$: 1, 1, 2, 4, 8, 17, 38, 82, 177, 376, 789, 1638 at $x = 2, \dots, 13$; row $t = 2$: 1, 1, 3, 6, 15, 33, 81, 186, 439, 1005, 2304 at $x = 3, \dots, 13$; row $t = 3$: 1, 1, 3, 7, 17, 40, 101, 241, 587, 1402 at $x = 4, \dots, 13$.

GF, column 8: $-1, -7, -34, -129, -432, -1320$ under indices $x = 8, \dots, 13$.

First factor (column 8): row $t = 1$: $1, 7, 34, 129, 432$ at $x = 9, \dots, 13$.

Second factor (column 8): row $t = 1$: $1, 1, 2, 4, 8, 17, 39, 88, 203, 464, 1056, 2381$
at $x = 2, \dots, 13$; row $t = 2$: $1, 1, 3, 6, 15, 33, 82, 193, 473, (\dots), 2743$
at $x = 3, \dots, 13$; row $t = 3$: $1, 1, 3, 7, 17, 40, 102, 248, 622, 1540$ at
 $x = 4, \dots, 13$.

GF, column 9: $-1, -8, -43, -180, -657$ under indices $x = 9, \dots, 13$.

First factor (column 9): row $t = 1$: $1, 8, 43, 180$ at $x = 10, \dots, 13$.

Second factor (column 9): row $t = 1$: $1, 1, 2, 4, 8, 17, 39, 89, 210, 498, 1185, 2813$
at $x = 2, \dots, 13$; row $t = 2$: $1, 1, 3, 6, 15, 33, 82, 194, 481, 1178, 2924$
at $x = 3, \dots, 13$; row $t = 3$: $1, 1, 3, 7, 17, 40, 102, 249, 630, 1584$ at
 $x = 4, \dots, 13$.

GF, column 10: $-1, -9, -53, -242$ under indices $x = 10, \dots, 13$.

First factor (column 10): row $t = 1$: $1, 9, 53$ at $x = 11, 12, 13$.

Second factor (column 10): row $t = 1$: $1, 1, 2, 4, 8, 17, 39, 89, 211, 506, 1228, 2993$
at $x = 2, \dots, 13$; row $t = 2$: $1, 1, 3, 6, 15, 33, 82, 194, 482, 1187, 2977$
at $x = 3, \dots, 13$; row $t = 3$: $1, 1, 3, 7, 17, 40, 102, 249, 631, 1593$ at
 $x = 4, \dots, 13$.

GF, column 11: $-1, -10, -63$ under indices $x = 11, 12, 13$.

First factor (column 11): row $t = 1$: $1, 10, 63$ at $x = 12, 13$ (and a leading entry).

Second factor (column 11): row $t = 1$: $1, 1, 2, 4, 8, 17, 39, 89, 211, 507, 1237, 3048$
at $x = 2, \dots, 13$; row $t = 2$: $1, 1, 3, 6, 15, 33, 82, 194, 482, 1188, 2987$
at $x = 3, \dots, 13$; row $t = 3$: $1, 1, 3, 7, 17, 40, 102, 249, 631, 1594$ at
 $x = 4, \dots, 13$.

GF, column 12: $-1, -11$ under indices $x = 12, 13$.

First factor (column 12): row $t = 1$: $1, 11$ at $x = 12, 13$.

Second factor (column 12): row $t = 1$: $1, 1, 2, 4, 8, 17, 39, 89, 211, 507, 1238, 3055$
at $x = 2, \dots, 13$; row $t = 2$: $1, 1, 3, 6, 15, 33, 82, 194, 482, 1188, 2988$
at $x = 3, \dots, 13$; row $t = 3$: $1, 1, 3, 7, 17, 40, 102, 249, 631, 1594$ at
 $x = 4, \dots, 13$.

GF, column 13: -1 under index $x = 13$.

First factor (column 13): row $t = 1$: 1 at $x = 13$.

Second factor (column 13): row $t = 1$: $1, 1, 2, 4, 8, 17, 39, 89, 211, 507, 1238, 3056$
at $x = 2, \dots, 13$; row $t = 2$: $1, 1, 3, 6, 15, 33, 82, 194, 482, 1188, 2988$
at $x = 3, \dots, 13$; row $t = 3$: $1, 1, 3, 7, 17, 40, 102, 249, 634, 1594$ at
 $x = 4, \dots, 13$.

[p. 459] I annex the following two Tables of (centre- and bicentre-) trees as far as I have completed them.

Table A.
Valency not greater than

Knots	0	1	2	3	4	5	6	7	8	Gen.
			Oxygen.	Boron.	Carbon.					
1	1	1	1	1	1	1	1	1	1	1
2		1	1	1	1	1	1	1	1	1
3			1	1	1	1	1	1	1	1
4			1	2	2	2	2	2	2	2
5			1	2	3	3	3	3	3	3
6			1	4	5	6	6	6	6	6
7			1	6	9	10	11	11	11	11
8		1	1	11	18	21	22	23	23	23
9			1	18	35	42	45	46	47	47
10			1	37	75					106
11			1	66	159					235
12			1	135	367					551
13			1	265	799					1301

Table B.
Actual Valency.

Knots	0	1	2	3	4	5	6	7	8
1	1								
2		1							
3			1						
4			1	1					
5			1	1	1				
6			1	3	1	1			
7			1	5	3	1	1		
8			1	10	7	3	1	1	
9			1	17	17	7	3	1	1
10			1	36	38				
11			1	65	93				
12			1	134	232				
13			1	264	534				

In A, the columns 2, 3, 4, and the last column are the totals given by the Tables IV., VI., VIII., and II., and the remaining numbers of columns 5, 6, 7, 8 have been found by trial; and, in B, the several columns are the differences of the

[p. 460] columns of A. The signification is obvious; for instance, if the number of knots is = 9, then Table A, if the valency, or the maximum number of branches from a knot, is = 2, 3, 4, 5, 6, 7, 8 or any greater number,

No. of trees = 1, 18, 35, 42, 45, 46, 47 :

viz. with 9 knots the tree can have at most 8 branches from a knot, so that the number of trees having at most 8 branches from a knot is = 47, the whole number of trees with 9 knots; and so the number of knots being as before = 9, Table B shows that the number of 47 is made up of the numbers

1, 17, 17, 7, 3, 1, 1;

viz. 1 is the No. of trees, at most 2 branches from a knot,

17 trees at least one 3-branch knot,
 17 trees „ „ „ 4-branch knot,
 7 trees „ „ „ 5-branch knot,
 3 trees „ „ „ 6-branch knot,
 1 tree „ „ „ 7-branch knot,
 1 tree „ „ „ 8-branch knot,

and we have accordingly

$$1 + 1 + 2 + 3 + 6 + 11 + 23 + 47.$$

I annex also a plate showing the figures of the trees of 1, 2, 3, . . . , 9 knots, classified according to their altitudes and number of main branches; and as to the bicentre-trees, according to the number of main branches from each point of the bicentre. The affixed numbers show in each case the greatest number of branches from a knot; so that when this is (2), the knots may be oxygen-, boron-, carbon-, &c., atoms; when (3), boron-, carbon-, &c., atoms; when (4), carbon-, &c., atoms; and so on.

[Plate — see scan: of trees follows in the original printing; not reproduced here.]

611. Report of the Committee on Mathematical Tables

Consisting of Professor Cayley, F.R.S., Professor Stokes, F.R.S., Professor Sir W. Thomson, F.R.S., Professor H. J. S. Smith, F.R.S., and J. W. L. Glaisher, F.R.S.

[From the Report of the British Association for the Advancement of Science (1875), pp. 305–336.]

[p. 461] The present Report (say Report 1875) is in continuation of that by Mr Glaisher, published in the volume for 1873, and here cited as Report 1873.

Report 1873 extends to all those tables which are at p. 3 (*l.c.*) included under the headings:—

A, auxiliary for non-logarithmic calculation, 1, 2, 3;

B, logarithmic and circular, 4, 5, 6;

C, exponential, 7, 8 (but only partially to C. 8), other than those tables of C referred to as “ $h.l \tan(45^\circ + \frac{1}{2}\phi)$ ”; and also partially (see Art. 24, pp. 81–83) to the tables included under the heading “E. 11, transcendental constants e , π , γ , &c., and their powers and functions.”

A future Report will comprise the tables, or further tables, included under the headings:—

C. 8. Hyperbolic antilogarithms (e^x) and $h.l \tan(45^\circ + \frac{1}{2}\phi)$, and hyperbolic sines, cosines, &c.

D. Algebraic constants.

9. Accurate integer or fractional values. Bernoulli’s Numbers, $\Delta^n 0^m$, &c. Binomial coefficients.

10. Decimal values auxiliary to the calculation of series.

E. 11. Transcendental constants e , π , γ , &c., and their powers and functions.

[p. 462] The present Report (1875) comprises the tables included under the headings:—

F. Arithmological.

12. Divisors and prime numbers. Prime roots. The Canon arithmeticus, &c.

13. The Pellian equation.

14. Partitions.

15. Quadratic forms a^2+b^2 , &c., and partitions of numbers into squares, cubes, and biquadrates.

16. Binary, ternary, &c., quadratic and higher forms.

17. Complex theories;

which divisions are herein referred to, for instance, as [F. 12. Divisors, &c.].

Report 1873 consists of six sections (§) divided into articles, which are separately numbered (see contents, p. 174); the present Report 1875 forms a single section (§ 7), divided in like manner into articles, which are separately numbered; but besides this the paragraphs are numbered, and that continuously, through the present Report 1875, so that any paragraph may be cited as Report 1875, No. —, as the case may be.

[F. 12. Divisors, &c.] Divisors and Prime Numbers. Art. I.

1. As to divisors and prime numbers see Report 1873, Art. 8 (Tables of Divisors—factor tables—and Tables of Primes), pp. 34–40. The tables there referred to, such as Chernac, Burckhardt, Dase, Dase and Rosenberg, are chiefly tables running up to very high numbers (the last of them the ninth million): wherein, to save space, multiples of 2, 3, 5 are frequently omitted, and in some of them only the least divisor is given. It would be for many purposes convenient to have a small table, going up say to 10,000, showing in every case all the prime factors of the number. Such a table might be arranged, 500 numbers in a page, in some such form as the following:

Factor Table 1 to 500

	0	1	2	3	4	5	6	7	8	9
39	2.3.5.13	17.23	2 ² .7 ²	3.131	2.197	5.79	2 ² .3 ² .11	397*	2.199	3.7.19

where the top line shows the units, and the left-hand column the remaining figures, viz. the specimen exhibits the composition of the several numbers from 390 to 399: a prime number, e.g. 397, would be sufficiently indicated by the absence of any decomposition, or it may be further indicated by an asterisk.

It may be noticed that, in the theory of numbers, the decomposition is specially required when the next following number is a prime, viz. that of $p - 1$, p being a

[p. 463] prime: also, that this is given incidentally, for prime numbers p up to 1000, in Jacobi's *Canon Arithmeticus*, *post*, No. 20, and up to 15,000 in Reuschle's Tables, V. (a, b, c) *post*, No. 22.

2. It may be proper to remark here that any table of a binary form is really a factor-table in the complex theory connected with such binary form. Thus in a table of the form $a^2 + b^2$, a number of this form has a factor $a + bi$

($i = \sqrt{-1}$ as usual); and the table, in fact, shows the complex factor $a + bi$ of the number in question: a well arranged table would give all the prime complex factors $a + bi$ of the number. But as to this more hereafter; at present, we are concerned with the real theory only, not with any complex theory.

3. Connected with a factor-table, we have (i) a Table of the number of less relative primes; viz. such a table would show, for every number, the number of inferior integers having no common factor with the number itself. The formula is a well-known one: for a number $N = a^\alpha b^\beta c^\gamma \dots$, ($a, b, \dots$ the distinct prime factors of N), the number of less relative primes is

$$\varpi(N), = a^{\alpha-1} b^{\beta-1} \dots (a-1)(b-1) \dots ,$$

or, what is the same thing, $= N \left(1 - \frac{1}{a}\right) \left(1 - \frac{1}{b}\right) \dots$. A small table ($N = 1$ to 100), occupying half a page, is given by

Euler, *Op. Arith. Coll.* t. ii. p. 128; viz. this is $\varpi 1 = 0$, $\varpi 2 = 1, \dots$, $\varpi 100 = 40$.

4. But it would be interesting to have such a table of the same extent with the proposed factor-table. The table might be of like form; for instance,

Number of less relative Primes Table 1 to 500

	0	1	2	3	4	5	6	7	8	9
39	112	192	144	292	84	232	144	198	148	264

It would be of still greater interest to have an inverse table showing the values of N which belong to a given value of $\varpi(N)$; for instance,

$\varpi =$	$N =$
40	41, 55, 75, 82, 88, 100, 110,
42	43, 49, 86, 98,
44	69, 92,
46	47, 94,
48	65, 104, 105, 112,
$\vdots$	$\vdots$

where, observe, that ϖ is of necessity even.

[p. 464] **5.** Again, connected with a factor-table, we have (ii) a Table of the Sum of the divisors of a Number. The formula is also a well-known one; for a number $N = a^\alpha b^\beta \dots$, ($a, b, \dots$ the distinct prime factors of N), the required sum

$$\int N = (1 + a + \dots + a^\alpha)(1 + b + \dots + b^\beta) \dots ,$$

or, what is the same thing,

$$= \frac{a^{\alpha+1} - 1}{a - 1} \cdot \frac{b^{\beta+1} - 1}{b - 1} \dots,$$

where, observe, that the number itself is reckoned as a divisor.

6. Such a table was required by Euler in his researches on Amicable Numbers (see *post*, No. 10), and he accordingly gives one of a considerable extent, viz.

Euler, *Op. Arith. Coll.* t. i. pp. 104–109.

It is to be remarked that, inasmuch as $\int N$ is obviously $\int a^\alpha \cdot \int b^\beta \dots$, the function need only be tabulated for the different integer powers a^α of each prime number a . The range of Euler's table is as follows:

$a =$	$a =$
2	to 36,
3	,, 15,
5	,, 9,
7	,, 10,
11	,, 9,
13	,, 7,
17	,, 5,
19	,, 5,
23	,, 4,
29 to 997	,, 3,

viz. for the several prime numbers from 29 to 997 the table gives $\int a$, $\int a^2$ and $\int a^3$.

And it is to be noticed that the values of the sum are exhibited, decomposed into their prime factors: thus a specimen of the table is

<i>Num.</i>	<i>Summa Divisorum.</i>
139	$2^2.5.7$
139^2	$3.13.499$
139^3	$2^3.5.7.9661$

[p. 465] 7. The form of the above table is adapted to its particular purpose (the theory of amicable numbers); but Euler gives also,

Euler, *Op. Arith. Coll.* t. i. p. 147—in the paper “*Observatio de Summis Divisorum*,” (1752), pp. 146–154,—a short table of about half a page, $N = 1$ to 100, of the form $\int 1 = 1$, $\int 2 = 3$, $\dots$, $\int 100 = 217$. The paper contains interesting analytical researches on the subject of $\int N$ which connect themselves with the theory of the Partition of Numbers.

8. It would be interesting to carry the last-mentioned table to the same extent as the proposed factor-table; and to add to it an inverse table, as suggested in regard to the number of less relative primes table.

9. Perfect Numbers. — A perfect number is a number which is equal to the sum of its divisors, the number itself not being reckoned as a divisor; e.g.

$$6 = 1 + 2 + 3, \quad \text{and} \quad 28 = 1 + 2 + 4 + 7 + 14.$$

Such numbers are indicated by a table of the sums of divisors: $\int 6 = 12$, $\int 28 = 56$, these two being, as appears by the table, Art. 7, the only perfect numbers less than 100.

10. Amicable Numbers. — These are pairs of numbers such that each is equal to the sum of the divisors of the other, not reckoning the other number as a divisor; that is, each has the same sum of divisors, the number being here reckoned as a divisor; say $\int A = B$, $\int B = A$; or, what is the same thing, $\int A = \int B (= A + B)$. Thus for the numbers 220, 284,

$$\begin{aligned} \int 220 &= (1 + 2 + 4)(1 + 5)(1 + 11) - 220, = 284, \\ \int 284 &= (1 + 2 + 4)(1 + 71) - 284, = 220; \end{aligned}$$

or, what is the same thing,

$$\int 220 = (1 + 2 + 4)(1 + 5)(1 + 11) = 504 = (1 + 2 + 4)(1 + 71) = \int 284.$$

11. A catalogue of 61 pairs of numbers is given by

Euler, *Op. Arith. Coll.* t. i. pp. 144–145; it occupies about one page. The paper, “De Numeris Amicabilibus,” pp. 102–145, contains an elaborate investigation of the theory, by means whereof all but two of the pairs of numbers are obtained. The first pair is the above-mentioned one, $2^2.5.11$ and $2^2.71 (= 220 \text{ and } 284)$; and the fifty-ninth pair is the high numbers

$$3^2.7^2.13.19.53.6959 \quad \text{and} \quad 3^2.7^2.13.19.179.2087.$$

[p. 466] The last two pairs are referred to as “formæ diverse a precedentibus”; viz. these are

$$\left\{ \begin{array}{l} 2^3.19.41 \\ 2^3.199 \end{array} \right. \quad \text{and} \quad \left\{ \begin{array}{l} 2^3.41.467 \\ 2^3.19.233 \end{array} \right.$$

12. A Table of the Frequency of Primes is given by

Gauss, *Tafel der Frequenz der Primzahlen, Werke*, t. ii. pp. 436–443; viz. this extends to 3,000,000.

The first part, extending to 1,000,000, = 1000 thousand, shows how many primes there are in each thousand: a specimen is

$$\begin{aligned} 1, & \ 168; \\ 2, & \ 135; \\ 3, & \ 127; \\ 4, & \ 120; \\ 5, & \ 119; \\ & \text{\&c.} \end{aligned}$$

viz. in the first thousand there are 168 primes, in the second thousand 135 primes, and so on.

For the second and third millions the frequency is given for each ten thousand: a specimen is

1,000,000 to 1,100,000.

	0	1	2	3	4	5	6	7	8	9	
1	1										1
2		1			1		1	1			4
3	1	4	2	2	3	1	2	3	3		21
4	2	8	5	4	3	6	9	4	5	8	54
5	11	10	8	18	12	10	10	12	15	8	114
6	14	14	18	21	16	22	19	15	17	15	171
7	26	17	23	23	24	24	17	22	20	21	217
8	19	19	21	7	14	15	20	17	15	17	164
9	11	13	9	13	14	14	12	13	11	16	126
10	8	6	8		9	5	9	9	7	9	71
11	6	6	4	6	3	1	3	1	4	5	39
12	1	1	2	1	1	1	2	2	1		12
13	1	1			1		1	1	1		6
14											
15											
16											
	752	719	732	700	731	698	713	722	706	737	7210

$$\int \frac{dx}{\log x} = 7212 \cdot 99;$$

[p. 467] viz. in the interval 1,000,000 to 1,010,000, 100 hundreds, there is 1 hundred containing 1 prime, there are 2 hundreds each containing 4 primes, 11 hundreds each containing 5 primes, . . . , 1 hundred containing 13 primes, so that, as

$$\begin{aligned} 1 \times 1 &= 1, \\ 4 \times 2 &= 8, \\ 5 \times 11 &= 55, \\ &\vdots \\ \frac{13 \times 1 &= 13,}{100 \quad \quad 752,} \end{aligned}$$

the whole 10,000 contains 752 primes; the next 10,000 contains 719 primes, and so on; the whole 100,000 thus containing $752 + 719 + \dots = 7210$ primes, which number is at the foot compared with the theoretic approximate value

$$\int \frac{dx}{\log x} \quad (\text{limits } 1,000,000 \text{ to } 1,010,000) = 7212 \cdot 99.$$

The integral in question is represented by the notation $\text{Li. } x$ or $\text{li. } x$.

p. 443. We have the like tables 1,000,000 to 2,000,000 and 2,000,000 to 3,000,000, showing for each 100,000 how many hundreds there are containing 0 prime, 1 prime, 2 primes, up to (the largest number) 17 primes.

13. It is noticed that

the 26,379th hundred contains no prime,
the 27,050th hundred contains 17 primes.

It may be observed that, if $N = 2.3.5 \dots p$, the product of all the primes up to p , then each of the numbers $N + 1$ and $N + q$ (if q be the prime next succeeding p) is or is not a prime; but the intermediate numbers $N + 2$, $N + 3$, $\dots$, $N + q - 1$ are certainly composite; viz. we thus have at least $q - 2$ consecutive composites. To obtain in this manner 99 consecutive composites, the value of N would be $= 2.3.5 \dots 97$, viz. this is a number far exceeding 2,637,900; but, in fact, the hundred numbers 2,637,901 to 2,638,000 are all composite.

Legendre, in his *Essai sur la Théorie des Nombres* (1st edit., 1798; 2nd edit., 1808, supplement, 1816: references to this edition), gives for the number of primes inferior to a given limit x the approximate formula

$$\frac{x}{\log x - 1.08366},$$

and p. 394, and supplement, p. 62, he compares for each 10,000 up to 100,000, and for each 100,000 up to 1,000,000, the values as computed by this formula with the actual numbers of primes exhibited by the tables of Vega and Chernac. Thus for $x = 1,000,000$, the computed value is 78,543, the actual value 78,493.

[p. 468] He shows, p. 414, that the number of integers, which are less than n and are not divisible by any of the numbers $\theta, \lambda, \mu, \dots$, is approximately

$$= n \left(1 - \frac{1}{\theta}\right) \left(1 - \frac{1}{\lambda}\right) \left(1 - \frac{1}{\mu}\right) \dots,$$

and taking $\theta, \lambda, \mu, \dots$ the successive primes 3, 5, 7, $\dots$ he gives the values of the function in question, or, say, the function

$$\frac{2}{3} \cdot \frac{4}{5} \cdot \frac{6}{7} \cdot \frac{10}{11} \dots \frac{\omega - 1}{\omega},$$

ω a prime, for the several prime values $\omega = 3$ to 1229 in the Table IX. (one page) at the end of the work.

14. A table of frequency is given by

Glaisher, J. W. L., *British Association Report* for 1872, p. 20. This gives for the second and the ninth millions, respectively divided into intervals of 50,000, the actual number of primes in each interval, as compared with

the theoretic value $\text{li } x' - \text{li } x$; and also deduced therefrom, by the formula $\log \frac{1}{2}(x' + x)$, a table of the average interval between two consecutive primes; this average interval increases very slowly: at the beginning and the end of the second million the values are $13 \cdot 76$ and $14 \cdot 58$ (theoretic values $13 \cdot 84$ and $14 \cdot 50$); at the beginning and the end of the ninth million $16 \cdot 02$ and $15 \cdot 95$ (theoretic values $15 \cdot 90$ and $16 \cdot 01$).

15. Coming under the head of Divisor Tables, some tables by Reuschle and Gauss may be here referred to. These are:

Reuschle, *Mathematische Abhandlung, zahlentheoretische Tabellen sammt einer dieselben treffenden Correspondenz mit der verewigten C. G. J. Jacobi*, 4^o, pp. 1–61¹ (1856). The tables belonging to the present subject are

A. Tafeln zur Zerlegung von $a^n - 1$ (pp. 18–22).

I. Table of the prime factors of $10^n - 1$, viz.

(a. pp. 18–19.) Complete decomposition of $10^n - 1$, $n = 1$ to 42: and $10^n + 1$, $n = 1$ to 21. Some values of n are omitted.

A specimen is

$$10^{14} - 1 = 3^2.53.79.265371653,$$

$$10^{12} + 1 = 11.139.10\,5831\,3049.$$

(b. p. 19.) List of the specific prime factors f of $10^n - 1$, or the prime factors of the residue after separation of the analytical factors, for those values of n for which the complete decomposition is unknown, and omitting those values for which no factor is known, $n = 25$ to 243.

[p. 469] A specimen is

n	f
25	21401

The meaning seems to be, residue of $10^{25} - 1$ is $1 + 10^5 + 10^{10} + 10^{15} + 10^{20}$ and this contains the prime factor 21401; but it is not clear why this is the “specific prime factor.”

II. Prime factors of $a^n - 1$ for different values of a and n .

(a. p. 20) gives for 41 values of a (2, 3, &c. at intervals to 100) and for the following values of n the decompositions of the residues or specific factors of

¹Titlepage missing in my copy; but I find from Prof. Kummer's notice of the work, Crelle, t. liii. (1857), p. 379, that it appeared as a Programm of the Stuttgart Gymnasium, Michaelmas, 1856, and was separately printed by Liesching and Co., Stuttgart.

$a^n - 1$, viz. these are

$$\begin{aligned}
 n = 1, & \ a - 1 \\
 2, & \ a + 1 \\
 3, & \ a^2 + a + 1 \\
 6, & \ a^2 - a + 1 \\
 4, & \ a^2 + 1 \\
 5, & \ a^4 + a^3 + a^2 + a + 1 \\
 10, & \ a^4 - a^3 + a^2 - a + 1 \\
 8, & \ a^4 + 1 \\
 12, & \ a^4 - a^2 + 1.
 \end{aligned}$$

A specimen is

	$a - 1$	$a^2 - 1$	$a^3 - 1$	$a^6 - 1$	$a^4 - 1$	$a^5 - 1$
10	3^2	11	$3^2.37$ $a^{10} - 1$ 9091	7.13	101	41.271
					73.137	9901

(b. p. 21.) Specific prime factors for the numbers 2, 3, 5, 6, 7, 10, (the powers 4, 8, 9 being omitted as coming under 2 and 3), for the exponents 1 to 42.

A specimen is

	$2^n - 1$	$3^n - 1$	$5^n - 1$	$6^n - 1$	$7^n - 1$	$10^n - 1$
19	524287	1597.363889	191. x	191. x	419. x	--

where the x denotes that the other factor is not known to be prime. And so, where no number is given, as in $10^{19} - 1$, it is not known whether the number ($= 1 + 10 + 10^2 + \dots + 10^{18}$) is or is not prime.

Addition, p. 22. For $a = 2$, the complete decomposition of the prime factor of $2^n - 1$ is given for values of n , $= 44, 45, \dots$ at intervals to 156.

A specimen is

n	f
44	397.2113

viz.

$$2^{22} - 2^{11} - \dots - 2^2 + 1 = 838861 = 397.2113.$$

$n = 31$, Fermat's prime. $n = 37$, the first case for which the decomposition is not given completely. $n = 41$, the first case for which no factor is known.

[p. 470] 16. Gauss, *Tafel zur Cyclotechnie, Werke*, t. ii. pp. 478–495, shows, for 2452 numbers of the several forms $a^2 + 1, a^2 + 4, a^2 + 9, \dots, a^2 + 81$,

the values of a such that the number in question is a product of prime factors no one of which exceeds 200, and exhibits all the odd prime factors of each such number. The table is in nine parts, *zerlegbare* $a^2 + 1$, *zerlegbare* $a^2 + 4$, &c., with to each part a subsidiary table, as presently mentioned. Thus a specimen is

<i>zerlegbare</i> $a^2 + 9$	
1	5
2	13
4	5.5
5	17
7	29
8	73
⋮	
1411168679	5.5.13.17.17.89.113.157.173.197.197

viz.

$$\begin{aligned}
 1^2 + 9, & \text{ odd prime factor is } 5, \\
 2^2 + 9, & \quad , , \quad , , \quad 13, \\
 4^2 + 9, & \quad , , \quad , , \text{ factors are } 5, 5,
 \end{aligned}$$

and so on.

And the subsidiary table is

5	1, 4, 79
13	2, 11, 41
17	5, 29, 46, 379, 1042

showing that the numbers a for which the largest factor is 5 are 1, 4, 79; those for which it is 13 are 2, 11, 41; and so on.

The object of the table is explained in the *Bemerkungen*, (*l.c.*, p. 523), by Schering, the editor of the volume, viz. it is to facilitate the calculation of the circular arcs the cotangents of which are rational numbers. To take a simple example, it appears to be by means of it that Gauss obtained, among other formulæ, the following:

$$\frac{\pi}{4} = 12 \arctan \frac{1}{38} + 8 \arctan \frac{1}{57} - 5 \arctan \frac{1}{239},$$

and

$$\frac{\pi}{4} = 12 \arctan \frac{1}{18} + 20 \arctan \frac{1}{57} + 7 \arctan \frac{1}{239} + 24 \arctan \frac{1}{268}.$$

[p. 471]

[F. 12. Divisors, etc.] continued. Prime Roots. The Canon Arithmeticus, Quadratic residues. Art. II.

17. Prime Roots. —Let p be a prime number; then there exist $\varpi(p-1)$ inferior integers g , such that all the numbers $1, 2, \dots, p-1$ are, to the modulus p ,

$$= 1, g, g^2, \dots, g^{p-2} \quad (g^{p-1} \text{ is of course } = 1).$$

This being so, g is said to be a *prime root* of p ; and moreover the several numbers g^α , where α is any number whatever less than and prime to $p-1$, constitute the series of the $\varpi(p-1)$ prime roots of p . It may be added that, if β be an integer number less than $p-1$, and having with it a greatest common measure $= k$, so that $\beta = \frac{p-1}{k} \cdot t$, then

$$(g^\beta)^{\frac{p-1}{k}} = g^{t(p-1)} = 1, \text{ (since } t \text{ is an integer, and } g^{p-1} = 1)$$

then g^β has the indicatrix $\frac{p-1}{k}$: the prime roots are those numbers which have the indicatrix $p-1$.

The like theory exists as to any number N of the form p^m or $2p^m$. There are here $\varpi(N)$, $= N\left(1 - \frac{1}{p}\right)$ or $\frac{1}{2}N\left(1 - \frac{1}{p}\right)$, in the two cases respectively, numbers less than N and prime to it; and we have then $\varpi(\varpi(N))$ numbers g such that, to the modulus N , all these numbers are $= 1, g, g^2, \dots, g^{\varpi(N)-1}$ ($g^{\varpi(N)}$ is of course $= 1$). This being so, g may be regarded as a prime root of N ($= p^m$ or $2p^m$, as the case may be); and moreover the several numbers g^α , where α is any number whatever less than and prime to $\varpi(N)$, constitute the series of the $\varpi(\varpi(N))$ prime roots of N . Thus $N = 3^2 = 9$, $\varpi(N) = 6$; we have

$$\begin{aligned} &1, 2^1, 2^2, 2^3, 2^4, 2^5, \\ &= 1, 2, 4, 8, 7, 5, \text{ mod. } 9; \end{aligned}$$

or the prime roots of 9 are 2^1 and 2^5 , $= 2$ and 5 .

So also $N = 2 \cdot 3^2 = 18$, $\varpi(N) = 6$; we have

$$\begin{aligned} &1, 5^1, 5^2, 5^3, 5^4, 5^5, \\ &= 1, 5, 7, 17, 13, 11, \text{ mod. } 18; \end{aligned}$$

and 5^1 and 5^5 , $= 5$ and 11 are the prime roots of 18.

18. A small table of prime roots, $p = 3$ to 37 , is given by

Euler, *Op. Arith. Coll.* t. i. pp. 525–526. The Memoir is entitled “Demonstrationes circa residua e divisione potestatum per numeros primos resultantia,” pp. 516–537 (1772).

[p. 472] **19.** A table, p and p^m , 3 to 97, is given by

Gauss, “Disquisitiones Arithmeticae,” 1801, (*Werke*, t. i. p. 468). This gives in each case a prime root, and shows the exponents in regard thereto of the several prime numbers less than p or p^m . Thus a specimen is

		2	3	5	7	11	13	17	19	23	29 &c.
27	2	1	*	5	16	13	8	15	12	11	
29	10	11	27	18	20	23	2	7	15	24	

viz. for 27 we have 2 a prime root, and $2 = 2^1$, $5 = 2^5$, $7 = 2^{16}$, $11 = 2^{13}$, &c.; and so also for 29 we have 10 a prime root, and $2 = 10^{11}$, $3 = 10^{27}$, $5 = 10^{18}$, &c.

20. Small tables are probably to be found in many other places; but the most extensive and convenient table is Jacobi’s *Canon Arithmeticus*, the complete title of which is

Canon Arithmeticus sive tabulæ quibus exhibentur pro singulis numeris primis vel primorum potestatibus infra 1000 numeri ad datos indices et indices ad datos numeros pertinentes. Edidit C. G. J. Jacobi. Berolini, 1839. 4°.

The contents are as follows:

<i>Introductio</i>	<i>i to xl</i>
<i>Tabulæ numerorum ad indices datos pertinentium et indicum numero dato correspondentium pro modulis primis minoribus quam 1000</i>	<i>1–221</i>
<i>Tabulæ residuorum et indicum sibi mutuo respondentium pro modulis minoribus quam 1000 qui sunt numerorum primorum potestates</i>	<i>222–238</i>
<i>Hujus tabulae ea pars quae pertinet ad modulos formæ 2^n, invenitur</i>	<i>239–240.</i>

The following is a specimen of the principal tables:

$$p = 19, p - 1 = 2 \cdot 3^2.$$

Numeri.

<i>I</i>	1	2	3	4	5	6	7	8	9
	10	5	12	6	3	11	15	17	18
1	9	14	7	13	16	8	4	2	1

Indices.

<i>N</i>	1	2	3	4	5	6	7	8	9
	18	17	5	16	2	4	12	15	10
1	1	6	3	13	11	7	14	8	9

[p. 473] where the first table gives the values of the powers of the prime root 10 (that 10 is the root appears by its index being given as = 1) to the

modulus 19, viz. $10^1 = 10$, $10^2 = 5$, $10^3 = 12$, &c.; and the second table gives the index of the power to which the same prime root must be raised in order that it may be, to the modulus 19, congruent with a given number: thus $10^{18} = 1$, $10^{17} = 2$, $10^5 = 3$, &c. The units of the index or number, as the case may be, are contained in the top line of the table, and the tens or hundreds and tens in the left-hand column.

21. There is given by

Jacobi, Crelle, t. xxx. (1846), pp. 181, 182, a table of m' for the argument m , such that

$$1 + g^m = g^{m'} \pmod{p}, \quad p = 7 \text{ to } 103, \text{ and } m = 0 \text{ to } 102.$$

A specimen is

p	7	11	13	17	19	23	29	31	37	to 103
g	3	2	6	10	10	10	10	17	5	
$m = 11$	.	.	6	4	7	*	27	21	34	

for instance, $p = 19$, $1 + 10^{11} = 10^7 \pmod{19}$.

Jacobi remarks that this table was calculated for him by his class during the winter course of 1836–37; and that, by means of the Canon Arithmeticus since published (in 1839), the same might easily be extended to all primes under 1000. In fact, for any such number p , putting any number of the table “Indices” = m , the next following number of the table gives the value of m' .

22. We have next, in Reuschle’s Memoir (*ante*, No. 15), the following relating to prime roots:

C. Tafeln für primitive Wurzeln und Hauptexponenten, oder V. erweiterte und bereicherte Burkhardsche Tafel, pp. 41–61, being divided into three parts; viz. these are

a. Table of the Hauptexponenten of the six roots 10, 5, 2, 6, 3, 7 for all prime numbers of the first 1000, together with the least primitive root of each of these numbers (pp. 42–46).

A specimen is as follows:

p	$p-1$	10		5		2		6		3		7		w
		e	n	e	n	e	n	e	n	e	n	e	n	
53	$2^2.13$	13	4	52	1	52	1	26	2	52	1	26	2	2

where e is the Hauptexponent or indicatrix of the root (10, 5, 2, 6, 3, 7, as the case may be), $n = \frac{p-1}{e}$, w the least primitive root; thus

$$p = 53, \quad 10^{13} = 1, \quad 5^{52} = 1, \quad 2^{52} = 1,$$

[p. 474] (2 being accordingly the least prime root),

$$6^{26} = 1, \quad 3^{52} = 1, \quad 7^{26} = 1.$$

The number w of the last column is the least primitive root. It is, of course, not always (as in the present case) one of the numbers 10, 5, 2, 6, 3, 7 to which the table relates: the first exception is $p = 191$, $w = 19$: the highest value of w is $w = 21$, corresponding to $p = 409$.

b. The like table for the roots 10 and 2 for all prime numbers from 1000 to 5000, together with as convenient as possible a prime root (and in some cases two prime roots) for each such number (pp. 47–53).

A specimen is:

p	$p - 1$	10		2		w
		e	n	e	n	
1289	$2^3.7.23$	92	14	161	8	6, 11

viz. here, mod. 1289, $10^{92} = 1$, $2^{161} = 1$; and two prime roots are 6, 11. We have thus by the present tables a prime root for every prime number not exceeding 5000.

c. The like table for the root 10 for all prime numbers between 5000 and 15000, (no column for w , nor any prime root given), pp. 53–61.

A specimen is

p	$p - 1$	e	n
9859	2.3.31.53	3286	3

viz., mod. 9859, we have $10^{3286} = 1$. But in a large number of cases we have $n = 1$, and therefore 10 a prime root. For example,

$$9887 \quad 2.4943 \quad 9886 \quad 1.$$

23. For a composite number n , if $N = \varpi(n)$ be the number of integers less than n and prime to it, then if x be any number less than n and prime to it, we have $x^N = 1 \pmod{n}$. But we have in this case no analogue of a prime root—there is no number x , such that its several powers $x', x'', \dots, x^{N-1} \pmod{n}$ are all different from unity; or, what is the same thing, there is for each value of x some submultiple of N , say N' , such that $x^{N'} = 1 \pmod{n}$. And these several numbers N' have a least common multiple I , which is not $= N$, but is a submultiple of N ; and this being so, then for all the several values of x , I is said to be the maximum indicator. For instance, $n = 12$, $N = \varpi(n)$; the numbers less than 12 and prime to it are 1, 5, 7, 11. We have, $(\text{mod. } 12)$, $1^1 = 1$, $5^2 = 1$, $7^2 = 1$, $11^2 = 1$, or the values of N' are 1, 2, 2, 2; their least common multiple is 2, and we have accordingly $I = 2$: viz. $x^2 = 1 \pmod{12}$ has the $\varpi(12)$ roots 1, 5, 7, 11. So $n = 24$, $\varpi(n) = 8$; the maximum indicator I is in this case also $= 2$.

[p. 475] A table of the maximum indicator $n = 1$ to 1000 is given by

Cauchy, *Exer. d'Analyse &c.*, t. ii. (1841), pp. 36–40, contained in the “Mémoire sur la résolution des Equations indéterminées du premier degré en nombres entiers,” pp. 1–40.

24. It thus appears that for a composite number n , the $\varpi(n)$ numbers less than n and prime to it cannot be expressed as $\equiv (\text{mod. } n)$ to the power of a single root; but for the expression of them it is necessary to employ two or more roots. A small table, $n = 1$ to 50, is given by

Cayley, “Specimen Table $M \equiv a^\alpha b^\beta (\text{mod. } N)$ for any prime or composite modulus,” *Quart. Math. Journ.* vol. ix. (1868), pp. 95, 96, and folding sheet, [397].

A specimen is

<i>Nos.</i>	12
<i>roots</i>	5, 7
<i>Ind.</i>	2, 2
<i>M.I.</i>	2
$\varpi()$	4
1	0, 0
2	
3	
4	
5	1, 0
6	
7	0, 1
8	
9	
10	
11	1, 1

viz. for the modulus 12 the roots are 5, 7, having respectively the indicators 2, 2, viz. $5^2 = 1 (\text{mod. } 12)$, $7^2 = 1 (\text{mod. } 12)$. Hence also the maximum indicator is = 2. $\varpi(= \varpi(n)) = 4$ is the number of integers less than 12 and prime to it, viz. these are 1, 5, 7, 11, which in terms of the roots 5, 7 and to mod. 12 are respectively $= 5^0.7^0$, $5^1.7^0$, $5^0.7^1$, and $5^1.7^1$.

25. Quadratic Residues. —In regard to a given prime number p , a number N is or is not a quadratic residue according as the index of N is even or odd, viz. g being a prime root and $N = g^\alpha$, then N is or is not a quadratic residue according as α is even or odd. But the quadratic residues can, of course, be obtained directly without the consideration of prime roots.

A small table, $p = 3$ to 97 and $N = -1$ and (prime values) 3 to 97, is given by

Gauss, “Disquisitiones Arithmeticae,” 1801; Table II. (*Werke*, t. i. p. 469): I notice here a misprint in the top line of the original; it should be $-1, +2, +3$, &c., instead of

[p. 476] $-1, +2, +3$, &c.; the -1 is printed correctly on p. 499 of the French translation *Recherches Arithmétiques*, Paris, 1807 and on p. 469 of vol. I. of *Werke*, (Göttingen, 1870).

A specimen is

	−1	+2	+3	+5	+7	+11	+13	+17	+19	+23 &c.
19	—	—					—	—	—	—

viz. $-1, 2, 3, 13$ are not, $5, 7, 11, 17$ &c. are, quadratic residues of 19. The residues taken positively and less than 19 are, in fact, 1, 4, 5, 6, 7, 9, 11, 16, 17.

The same table carried from $p = 3$ to 503, and prime values $N = 3$ to 997, is given by

Gauss, *Werke*, t. ii. pp. 400–409. A specimen is

	2	3	5	7	11	13	17	19	23 &c.
19	—		—			—	—	—	

viz. the arrangement is the same, except only that the -1 column is omitted.

26. We have also by Gauss

“Disquisitiones Arithmeticae” Table III. (*Werke*, t. i. p. 470), for the conversion into decimals of a vulgar fraction, denominator p or p^μ , not exceeding 100. The explanation is given in Art. 314 *et seq.* of the same work.

But this table, carried to a greater extent, is given by Gauss, *Werke*, t. ii. pp. 412–434, “Tafel zur Verwandlung gemeiner Brüche mit Nennern aus dem ersten Tausend in Decimalbrüche,” viz. the denominators are here primes or powers of primes, p^μ up to 997.

To explain the table, consider a modulus p^μ (where μ may be $= 1$); if 10 is not a prime root of p^μ , consider a prime root r , which is such that $r^e = 10 \pmod{p^\mu}$, e being a submultiple of $p^{\mu-1}(p-1)$; say we have $ef = p^{\mu-1}(p-1)$: then $10^f = 1 \pmod{p^\mu}$. Consider any fraction $\frac{N}{p^\mu}$; then we may write $N = r^{ke+l} \pmod{p^\mu}$, k from 0 to $f-1$ and l from 0 to $e-1$, $N = 10^k r^l$, and consequently $\frac{N}{p^\mu}$ and $\frac{10^k r^l}{p^\mu}$ have the same mantissa (decimal part regarded as an integer); hence, in order to know the mantissa of every fraction whatever of $\frac{N}{p^\mu}$, it is sufficient to know the mantissa of $\frac{r^l}{p^\mu}$, that is, the mantissas of $\frac{1}{p^\mu}, \frac{r}{p^\mu}, \frac{r^2}{p^\mu}, \dots, \frac{r^{e-1}}{p^\mu}$, or, what is the same thing, the mantissas of $\frac{10}{p^\mu}, \frac{10r}{p^\mu}, \dots, \frac{10r^{e-1}}{p^\mu}$.

For instance, $p^\mu = 11$, $10^2 = 1 \pmod{11}$, whence $f = 2$, $e = 5$; and taking $r = 2$, we have $10 = r^5 \pmod{11}$.

[p. 477] The required mantissæ, denoted in the table by

$$(0), (1), (2), (3), (4),$$

are those of

$$\frac{10}{11}, \frac{10.2}{11}, \frac{10.2^2}{11}, \frac{10.2^3}{11}, \frac{10.2^4}{11},$$

viz. these fractions are respectively

$$(0), (1), (2), (3), (4), \\ \cdot 9090\dots, 1 \cdot 8181\dots, 3 \cdot 6363\dots, 7 \cdot 2727\dots, 14 \cdot 5454\dots,$$

or their mantissæ are 90, 81, 63, 27, 54.

And we accordingly have as a specimen

$$11 \mid (1) \dots 81, (2) \dots 63, (3) \dots 27, (4) \dots 54, (0) \dots 90.$$

Or again, as another specimen, $r = 2$

$$27 \mid (1) \dots 740, (2) \dots 481, (3) \dots 962, (4) \dots 925, (5) \dots 851, (0) \dots 370.$$

The table in this form extends to $p^\mu = 463$; the values of r (not given in the body of the table) are annexed, p. 420.

In the latter part of the table $p^\mu = 467$ to 997, we have only the mantissæ of $\frac{100}{p^\mu}$. A specimen is

$$\begin{array}{c|cccc} 547 & 1828153564 & 8994515539 & 3053016453 & 3820840950 \\ & 6398537477 & 1480804387 & 5685557586 & 8372943327 \\ & 2394881170 & 0182815356 & & \end{array}$$

viz. the fraction $\frac{100}{547} = \cdot 182815\dots$ has a period of 91, $= \frac{1}{6} \cdot 546$, figures.

[F. 13. The Pellian Equation.] Art. III.

27. The Pellian equation is $y^2 = ax^2 + 1$, a being a given integer number, which is not a square (or rather, if it be, the only solution is $y = 1, x = 0$), and x, y being numbers to be determined: what is required is the least values of x, y , since these, being known, all other values can be found. A small table $a = 2$ to 68 is given by

Euler, *Op. Arith. Coll.* t. i. p. 8. The Memoir is “Solutio problematum Diophanteorum per numeros integros,” pp. 4–10, 1732–33. The form of the table is

a	$x (= p)$	$y (= q)$
2	2	3
3	1	2
5	4	9
$\vdots$	$\vdots$	$\vdots$
68	4	33.

[p. 478] Even here, for some of the values of a , the values of x, y are extremely large; thus $a = 61$, $x = 226,153,980$, $y = 1,766,319,049$.

And probably tables of a like extent may be found elsewhere; in particular, a table of the solution of $y^2 = ax^2 \pm 1$ (– when the value of a is such that there is a solution of $y^2 = ax^2 - 1$, and + for other values of a), $a = 2$ to 135, is given by Legendre, *Théorie des Nombres*, 2nd ed. 1808, in the Table X. (one page) at the end of the work. For the before-mentioned number 61, the equation is $y^2 = 61x^2 - 1$, and the values are $x = 3805$, $y = 29718$; much smaller than Euler's values for the equation $y^2 = 61x^2 + 1$.

28. The most extensive table, however, is given by

Degen, *Canon Pellianus, sive Tabula simplicissimam æquationis celebratissimæ $y^2 = ax^2 + 1$ solutionem, pro singulis numeri dati valoribus ab 1 ad 1000 in usque numeris rationalibus, iisdemque integris exhibens. Auctore Carolo Ferdinando Degen. Hafniæ (Copenhagen) apud Gerhardum Bonnarum, 1817.* 8vo. pp. iv to xxiv and 1 to 112.

The first table (pp. 3–106) is entitled as “Tabula I. Solutionem Equationis $y^2 - ax^2 - 1 = 0$ exhibens.” It, in fact, also gives the expression of $\sqrt{a}$ as a continued fraction; thus a specimen is

$$\begin{array}{c|cccccc} 209 & 14 & 2 & 5 & 3 & (2) \\ & 1 & 13 & 5 & 8 & 11 \\ & 3220 & & & & \\ & 46551 & & & & \end{array}$$

Here the first line gives the continued fraction, viz.

$$\begin{aligned} \sqrt{209} = 14 + \cfrac{1}{2 + \cfrac{1}{5 + \cfrac{1}{3 + \cfrac{1}{2 + \cfrac{1}{3 + \cfrac{1}{5 + \cfrac{1}{2 + \cfrac{1}{28 + \cfrac{1}{2 + \&c.}}}}}}}}} \end{aligned}$$

the period being (2, 5, 3, 2, 3, 5, 2) indicated by 2, 5, 3 (2). [The number of terms in the period is here odd, but it may be even; for instance, the period (1, 1, 5, 5, 1, 1) is indicated by 1, 1 (5, 5).]

The second line contains auxiliary numbers presenting themselves in the process; thus, if $R^2 = 209$, we have $R = 14 + \frac{1}{\beta}$,

$$\beta = \frac{1}{R - 14} = \frac{1(R + 14)}{209 - 14^2} = \frac{R + 14}{13} = 2 + \frac{1}{\beta'},$$

$$\beta' = \frac{13}{R-12} = \frac{13(R+12)}{209-12^2} = \frac{R+12}{5} = 5 + \frac{1}{\gamma},$$

$$\gamma = \frac{5}{R-13} = \frac{5(R+13)}{209-13^2} = \frac{R+13}{8} = 3 + \frac{1}{\delta},$$

&c.

Original scan reference pages

The following pages reproduce the corresponding range of the Internet Archive scan for visual reference of figures, tables, and any text that may have been compressed in the typeset reconstruction. Source: `collectedmathema09cayluoft.pdf`, pages 476–500.

TABLE VIII.—Carbon Centre- and Bicentre-Trees.

Index <i>x</i> , or number of knots.	Index <i>t</i> , or number of main branches.	Centre-Trees. Altitude or number of column.									Centre.	Grand Total.	Bicentre.	Bicentre-Trees. Altitude.					
		0	1	2	3	4	5	6	0					1	2	3	4	5	
1	0	1	1							1	1	0							
	Total	1	1																
2										0	1	1	1						
3	2		1	1						1	1	0							
	Total		1	1															
4	2		1	1						1	2	1		1					
	Total		1	1															
5	2		1	1	2					2	3	1		1					
	Total		1	1	2														
6	2			1	1					2	5	3		2	1				
	3			1	1														
	Total			2	2														
7	2			2	1	3				6	9	3		1	2				
	3			2		2													
	4			1		1													
Total			5	1	6														
8	2			1	2	3				9	18	9		1	7	1			
	3			3	1	4													
	4			2		2													
Total			6	3	9														
9	2			1	7	1	9			20	35	15			12	3			
	3			3	3		6												
	4			4	1		5												
Total			8	11	1	20													
10	2				12	3	15			37	75	38			23	14	1		
	3			3	11	1	15												
	4			4	3		7												
Total			7	26	4	37													
11	2				23	14	1	38		86	159	73			30	39	4		
	3			2	24	4		30											
	4			5	12	1		18											
Total			7	59	19	1	86												
12	2				30	39	4	73		183	357	174			42	108	23	1	
	3			1	54	19	1	75											
	4			4	27	4		35											
Total			5	111	62	5	183												
13	2				42	108	23	1	174	419	799	380			47	244	84	5	
	3			1	88	63	5	157											
	4			4	63	20	1	88											
Total			5	193	191	29	1	419											

Subsidiary Table for GF 's of Tables VII. and VIII.

Index t .	Index of x .														
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	
0		1													GF , column 0.
*		-1													GF , column 1.
0 1 2 3 4	1	1	1	1	1										First factor.
0		1													Second factor.
*			-1	-1	-1										GF , column 2.
0 1 2 3 4	(1)		1	1	1	1	2	1	1	2	2	2	1	1	First factor.
0 1 2 3		1	1	1	1										Second factor.
*				-1	-2	-4	-4	-5	-4	-4	-3	-2	-1	-1	GF , column 3.
0 1 2 3 4	(1)			1	2	4	4	5	4	4	3	2	1		First factor.
0 1 2 3		1	1	1	1	1	2	2	1	1					Second factor.
*					-1	-3	-8	-15	-27	-43	-67	-97	-136	-183	GF , column 4.
0 1 2 3	(1)				1	3	8	15	27	43	67	97	136		First factor.
0 1 2 3		1	1	1	2	3	4	4	5	4	4	3	2	1	Second factor.
*						-1	-4	-13	-32	-74	-155	-316	-612	-1160	GF , column 5.
0 1 2	(1)					1	4	13	32	74	155	316	612		First factor.
0 1 2 3		1	1	1	2	4	7	12	20	31	47	70	99	137	Second factor.
*							-1	-5	-19	-56	-151	-374	-889	-2032	GF , column 6.
0 1 2	1						1	5	19	56	151	374	889		First factor.
0 1 2 3		1	1	1	2	4	8	16	33	63	121	225	415	749	Second factor.
					1	3	6	15	32	75	160	350	732	1534	
						1	3	7	17	39	95	214	491	1093	

Subsidiary Table for GF 's of Tables VII. and VIII. (continued).

Index t .	Index of x .														
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	
*							-	-1	-6	-26	-88	-267	-743	-1968	GF , column 7.
0 1	(1)							1	6	26	88	267	743		First factor.
0 1 2 3		1	1	1	2	4	8	17	38	82	177	376	789	1638	Second factor.
			1	1	1	3	6	15	33	81	186	439	1005	2304	
				1	1	1	3	7	17	40	101	241	587	1402	
*									-1	-7	-34	-129	-432	-1320	GF , column 8.
0 1	(1)								1	7	34	129	432		First factor.
0 1 2 3		1	1	1	2	4	8	17	39	88	203	464	1056	2381	Second factor.
			1	1	1	3	6	15	33	82	193	473	1135	2743	
				1	1	1	3	7	17	40	102	248	622	1540	
*										-1	-8	-43	-180	-657	GF , column 9.
0 1	(1)									1	8	43	180		First factor.
0 1 2 3		1	1	1	2	4	8	17	39	89	210	498	1185	2813	Second factor.
			1	1	1	3	6	15	33	82	194	481	1178	2924	
				1	1	1	3	7	17	40	102	249	630	1584	
*											-1	-9	-53	-242	GF , column 10.
0 1	(1)										1	9	53		First factor.
0 1 2 3		1	1	1	2	4	8	17	39	89	211	506	1228	2993	Second factor.
			1	1	1	3	6	15	33	82	194	482	1187	2977	
				1	1	1	3	7	17	40	102	249	631	1593	
*												-1	-10	-63	GF , column 11.
0 1	(1)											1	10	63	First factor.
0 1 2 3		1	1	1	2	4	8	17	39	89	211	507	1237	3048	Second factor.
			1	1	1	3	6	15	33	82	194	482	1188	2987	
				1	1	1	3	7	17	40	102	249	631	1594	
*													-1	-11	GF , column 12.
0 1	(1)												1	11	First factor.
0 1 2 3		1	1	1	2	4	8	17	39	89	211	507	1238	3055	Second factor.
			1	1	1	3	6	15	33	82	194	482	1188	2988	
				1	1	1	3	7	17	40	102	249	631	1594	
*														-1	GF , column 13.
0 1	(1)													1	First factor.
0 1 2 3		1	1	1	2	4	8	17	39	89	211	507	1238	3056	Second factor.
			1	1	1	3	6	15	33	82	194	482	1188	2988	
				1	1	1	3	7	17	40	102	249	634	1594	

I annex the following two Tables of (centre- and bicentre-) trees as far as I have completed them.

TABLE A.

Knots.	Valency not greater than									Gen.
	0	1	2 Oxygen.	3 Boron.	4 Carbon.	5	6	7	8	
1	1	1	1	1	1	1	1	1	1	1
2		1	1	1	1	1	1	1	1	1
3			1	1	1	1	1	1	1	1
4			1	2	2	2	2	2	2	2
5			1	2	3	3	3	3	3	3
6			1	4	5	6	6	6	6	6
7			1	6	9	10	11	11	11	11
8			1	11	18	21	22	23	23	23
9			1	18	35	42	45	46	47	47
10			1	37	75					106
11			1	66	159					235
12			1	135	357					551
13			1	265	799					1301

TABLE B.

Knots.	Actual Valency.								
	0	1	2	3	4	5	6	7	8
1	1								
2		1							
3			1						
4			1	1					
5			1	1	1				
6			1	3	1	1			
7			1	5	3	1	1		
8			1	10	7	3	1	1	
9			1	17	17	7	3	1	1
10			1	36	38				
11			1	65	93				
12			1	134	222				
13			1	264	534				

In A, the columns 2, 3, 4, and the last column are the totals given by the Tables IV., VI., VIII., and II., and the remaining numbers of columns 5, 6, 7, 8 have been found by trial; and, in B, the several columns are the differences of the

columns of A. The signification is obvious; for instance, if the number of knots is =9, then Table A, if the valency, or the maximum number of branches from a knot,

is = 2, 3, 4, 5, 6, 7, 8 or any greater number,

No. of trees = 1, 18, 35, 42, 45, 46, 47:


viz. with 9 knots the tree can have at most 8 branches from a knot, so that the number of trees having at most 8 branches from a knot is =47, the whole number of trees with 9 knots; and so the number of knots being as before =9, Table B shows that the number of 47 is made up of the numbers

1, 17, 17, 7, 3, 1, 1;

viz. 1 is the No. of trees, at most 2 branches from a knot,

17	"	"	3	"	"	at least one 3-branch knot.
17	"	"	4	"	"	" 4 "
7	"	"	5	"	"	" 5 "
3	"	"	6	"	"	" 6 "
1	"	"	7	"	"	" 7 "
1	"	"	8	"	"	" 8 "

I annex also a plate showing the figures of the $1+1+2+3+6+11+23+47$ trees of 1, 2, 3, ..., 9 knots, classified according to their altitudes and number of main branches; and as to the bicentre-trees, according to the number of main branches from *each* point of the bicentre. The affixed numbers show in each case the greatest number of branches from a knot; so that when this is (2), the knots may be oxygen-, boron-, carbon-, &c., atoms; when (3), boron-, carbon-, &c., atoms; when (4), carbon-, &c., atoms; and so on.





611.

REPORT OF THE COMMITTEE ON MATHEMATICAL TABLES:

CONSISTING OF PROFESSOR CAYLEY, F.R.S., PROFESSOR STOKES, F.R.S.,
PROFESSOR SIR W. THOMSON, F.R.S., PROFESSOR H. J. S. SMITH, F.R.S.,
AND J. W. L. GLAISHER, F.R.S.

[From the *Report of the British Association for the Advancement of Science* (1875),
pp. 305—336.]

THE present Report (say Report 1875) is in continuation of that by Mr Glaisher, published in the volume for 1873, and here cited as Report 1873.

Report 1873 extends to all those tables which are at p. 3 (*l.c.*) included under the headings:—

A, auxiliary for non-logarithmic calculation, 1, 2, 3;

B, logarithmic and circular, 4, 5, 6;

C, exponential, 7, 8 (but only partially to C. 8), other than those tables of C referred to as " $\text{h.l} \tan(45^\circ + \frac{1}{2}\phi)$ "; and also partially (see Art. 24, pp. 81—83) to the tables included under the heading "E. 11, transcendental constants ϵ , π , γ , &c., and their powers and functions."

A future Report will comprise the tables, or further tables, included under the headings:—

C. 8. Hyperbolic antilogarithms (e^x) and $\text{h.l} \tan(45^\circ + \frac{1}{2}\phi)$, and hyperbolic sines, cosines, &c.

D. Algebraic constants.

9. Accurate integer or fractional values. Bernoulli's Numbers, $\Delta^n 0^m$, &c. Binomial coefficients.

10. Decimal values auxiliary to the calculation of series.

E. 11. Transcendental constants ϵ , π , γ , &c., and their powers and functions.

The present Report (1875) comprises the tables included under the headings:—

F. Arithmological.

12. Divisors and prime numbers. Prime roots. The Canon arithmeticus, &c.
13. The Pellian equation.
14. Partitions.
15. Quadratic forms $a^2 + b^2$, &c., and partitions of numbers into squares, cubes, and biquadrates.
16. Binary, ternary, &c., quadratic and higher forms.
17. Complex theories:

which divisions are herein referred to, for instance, as [F. 12. Divisors, &c.].

Report 1873 consists of six sections (§) divided into articles, which are separately numbered (see contents, p. 174); the present Report 1875 forms a single section (§ 7), divided in like manner into articles, which are separately numbered; but besides this the paragraphs are numbered, and that continuously, through the present Report 1875, so that any paragraph may be cited as Report 1875, No. —, as the case may be.

[F. 12. *Divisors, &c.*] *Divisors and Prime Numbers.* Art. I.

1. As to divisors and prime numbers see Report 1873, Art. 8 (Tables of Divisors—factor tables—and Tables of Primes), pp. 34—40. The tables there referred to, such as Chernac, Burckhardt, Dase, Dase and Rosenberg, are chiefly tables running up to very high numbers (the last of them the ninth million): wherein, to save space, multiples of 2, 3, 5 are frequently omitted, and in some of them only the least divisor is given. It would be for many purposes convenient to have a small table, going up say to 10,000, showing in every case all the prime factors of the number. Such a table might be arranged, 500 numbers in a page, in some such form as the following:—

Factor Table										
1 to 500										
	0	1	2	3	4	5	6	7	8	9
39	2.3.5.13	17.23	2 ³ .7 ²	3.131	2.197	5.79	2 ² .3 ² .11	397*	2.199	3.7.19

where the top line shows the units, and the left-hand column the remaining figures, viz. the specimen exhibits the composition of the several numbers from 390 to 399: a prime number, e.g. 397, would be sufficiently indicated by the absence of any decomposition, or it may be further indicated by an asterisk.

It may be noticed that, in the theory of numbers, the decomposition is specially required when the next following number is a prime, viz. that of $p-1$, p being a

prime: also, that this is given incidentally, for prime numbers p up to 1000, in Jacobi's *Canon Arithmeticus*, *post*, No. 20, and up to 15,000 in Reuschle's Tables, V. (a, b, c) *post*, No. 22.

2. It may be proper to remark here that any table of a binary form is really a factor-table in the complex theory connected with such binary form. Thus in a table of the form $a^2 + b^2$, a number of this form has a factor $a + bi$ ($i = \sqrt{-1}$ as usual); and the table, in fact, shows the complex factor $a + bi$ of the number in question: a well arranged table would give all the prime complex factors $a + bi$ of the number. But as to this more hereafter; at present, we are concerned with the real theory only, not with any complex theory.

3. Connected with a factor-table, we have (i) a Table of the number of less relative primes; viz. such a table would show, for every number, the number of inferior integers having no common factor with the number itself. The formula is a well-known one: for a number $N = a^\alpha b^\beta c^\gamma \dots$, ($a, b, \dots$ the distinct prime factors of N), the number of less relative primes is

$$\varpi(N) = a^{\alpha-1} b^{\beta-1} \dots (a-1)(b-1) \dots,$$

or, what is the same thing, $\varpi(N) = N \left(1 - \frac{1}{a}\right) \left(1 - \frac{1}{b}\right) \dots$. A small table ($N = 1$ to 100), occupying half a page, is given by

Euler, *Op. Arith. Coll.* t. II. p. 128; viz. this is $\pi 1 = 0, \pi 2 = 1, \dots, \pi 100 = 40$.

4. But it would be interesting to have such a table of the same extent with the proposed factor-table. The table might be of like form; for instance,

Number of less relative Primes Table										1 to 500
	0	1	2	3	4	5	6	7	8	9
29	112	192	144	292	84	232	144	198	148	264

It would be of still greater interest to have an inverse table showing the values of N which belong to a given value of $\varpi(N)$; for instance,

$\varpi =$	$N =$
40	41, 55, 75, 82, 88, 100, 110,
42	43, 49, 86, 98,
44	69, 92,
46	47, 94,
48	65, 104, 105, 112,
$\vdots$	

where, observe, that ϖ is of necessity even.

5. Again, connected with a factor-table, we have (ii) a Table of the Sum of the divisors of a Number. The formula is also a well-known one; for a number $N = a^{\alpha} b^{\beta} \dots$, ($a, b, \dots$ the distinct prime factors of N), the required sum

$$\int N \text{ is } = (1 + a + \dots + a^{\alpha})(1 + b + \dots + b^{\beta}) \dots,$$

or, what is the same thing,

$$= \frac{a^{\alpha+1} - 1}{a - 1} \cdot \frac{b^{\beta+1} - 1}{b - 1} \dots,$$

where, observe, that the number itself is reckoned as a divisor.

6. Such a table was required by Euler in his researches on Amicable Numbers (see *post*, No. 10), and he accordingly gives one of a considerable extent, viz.

Euler, *Op. Arith. Coll.* t. i. pp. 104—109.

It is to be remarked that, inasmuch as $\int N$ is obviously $= \int a^{\alpha} \int b^{\beta} \dots$, the function need only be tabulated for the different integer powers a^{α} of each prime number a . The range of Euler's table is as follows:—

$a =$	$\alpha =$
2	1 to 36,
3	1 „ 15,
5	1 „ 9,
7	1 „ 10,
11	1 „ 9,
13	1 „ 7,
17	1 „ 5,
19	1 „ 5,
23	1 „ 4,
29 to 997	1 „ 3,

viz. for the several prime numbers from 29 to 997 the table gives $\int a$, $\int a^2$, and $\int a^3$.

And it is to be noticed that the values of the sum are exhibited, decomposed into their prime factors: thus a specimen of the table is

Num.	Summa Divisorum.
139	$2^2 \cdot 5 \cdot 7$
139^2	$3 \cdot 13 \cdot 499$
139^3	$2^3 \cdot 5 \cdot 7 \cdot 9661$

7. The form of the above table is adapted to its particular purpose (the theory of amicable numbers); but Euler gives also,

Euler, *Op. Arith. Coll.* t. I. p. 147—in the paper “*Observatio de Summis Divisorum*,” (1752), pp. 146—154,—a short table of about half a page, $N=1$ to 100, of the form $\int 1 = 1, \int 2 = 3, \dots, \int 100 = 217$. The paper contains interesting analytical researches on the subject of $\int N$ which connect themselves with the theory of the Partition of Numbers.

8. It would be interesting to carry the last-mentioned table to the same extent as the proposed factor-table; and to add to it an inverse table, as suggested in regard to the number of less relative primes table.

9. *Perfect Numbers*.—A perfect number is a number which is equal to the sum of its divisors, the number itself not being reckoned as a divisor; e.g.

$$6 = 1 + 2 + 3, \text{ and } 28 = 1 + 2 + 4 + 7 + 14.$$

Such numbers are indicated by a table of the sums of divisors $\int 6 = 12, \int 28 = 56$, these two being, as appears by the table, Art. 7, the only perfect numbers less than 100.

10. *Amicable Numbers*.—These are pairs of numbers such that each is equal to the sum of the divisors of the other, not reckoning the other number as a divisor; that is, each has the same sum of divisors, the number being here reckoned as a divisor; say $\int' A = B, \int' B = A$; or, what is the same thing, $\int A = \int B (= A + B)$. Thus for the numbers 220, 284,

$$\int' 220 = (1 + 2 + 4)(1 + 5)(1 + 11) - 220, = 284,$$

$$\int' 284 = (1 + 2 + 4)(1 + 71) - 284, = 220;$$

or, what is the same thing,

$$\int 220 = (1 + 2 + 4)(1 + 5)(1 + 11) = 504 = (1 + 2 + 4)(1 + 71) = \int 284.$$

11. A catalogue of 61 pairs of numbers is given by

Euler, *Op. Arith. Coll.* t. I. pp. 144—145; it occupies about one page. The paper, “*De Numeris Amicabilibus*,” pp. 102—145, contains an elaborate investigation of the theory, by means whereof all but two of the pairs of numbers are obtained. The first pair is the above-mentioned one, $2^2 \cdot 5 \cdot 11$ and $2^2 \cdot 71$ (= 220 and 284); and the fifty-ninth pair is the high numbers

$$3^3 \cdot 7^2 \cdot 13 \cdot 19 \cdot 53 \cdot 6959 \text{ and } 3^3 \cdot 7^2 \cdot 13 \cdot 19 \cdot 179 \cdot 2087.$$

The last two pairs are referred to as "formæ diversæ a precedentibus;" viz. these are

$$\begin{cases} 2^3.19.41 \\ 2^5.199 \end{cases} \text{ and } \begin{cases} 2^3.41.467 \\ 2^5.19.233. \end{cases}$$

12. A Table of the Frequency of Primes is given by

Gauss, *Tafel der Frequenz der Primzahlen*, *Werke*, t. II. pp. 436—443; viz. this extends to 3,000,000.

The first part, extending to 1,000,000, = 1000 thousand, shows how many primes there are in each thousand: a specimen is

1, 168:

2, 135:

3, 127:

4, 120:

5, 119:

&c.;

viz. in the first thousand there are 168 primes, in the second thousand 135 primes, and so on.

For the second and third millions the frequency is given for each ten thousand: a specimen is

1,000,000 to 1,100,000.

	0	1	2	3	4	5	6	7	8	9	
1	1										1
2		1				1		1	1		4
3		4	2	2	3	1	2	3	3	1	21
4	2	8	5	4	3	6	9	4	5	8	54
5	11	10	8	18	12	10	10	12	15	8	114
6	14	14	18	21	16	22	19	15	17	15	171
7	26	17	23	23	24	24	17	22	20	21	217
8	19	19	21	7	14	15	20	17	15	17	164
9	11	13	9	13	14	14	12	13	11	16	126
10	8	6	8	5	9	5	5	9	7	9	71
11	6	6	4	6	3	1	3	1	4	5	39
12	1	1	2	1	1	1	2	2	1		12
13	1	1			1		1	1	1		6
14											
15											
16											
	752	719	732	700	731	698	713	722	706	737	7210,

$$\int \frac{dx}{\log x} = 7212.99;$$

viz. in the interval 1,000,000 to 1,010,000, 100 hundreds, there is 1 hundred containing 1 prime, there are 2 hundreds each containing 4 primes, 11 hundreds each containing 5 primes, ..., 1 hundred containing 13 primes, so that, as

$$\begin{array}{rcl} 1 \times 1 & = & 1, \\ 4 \times 2 & = & 8, \\ 5 \times 11 & = & 55, \\ \vdots & & \\ 13 \times 1 & = & 13, \\ \hline 100 & & 752, \end{array}$$

the whole 10,000 contains 752 primes; the next 10,000 contains 719 primes, and so on; the whole 100,000 thus containing $752 + 719 + \&c. \dots = 7210$ primes, which number is at the foot compared with the theoretic approximate value

$$\int \frac{dx}{\log x} \text{ (limits 1,000,000 to 1,010,000) } = 7212.99.$$

The integral in question is represented by the notation *Li. x* or *li. x*.

p. 443. We have the like tables 1,000,000 to 2,000,000 and 2,000,000 to 3,000,000, showing for each 100,000 how many hundreds there are containing 0 prime, 1 prime, 2 primes, up to (the largest number) 17 primes.

13. It is noticed that

the 26,379th hundred contains no prime,
the 27,050th hundred contains 17 primes.

It may be observed that, if $N = 2.3.5 \dots p$, the product of all the primes up to p , then each of the numbers $N+1$ and $N+q$ (if q be the prime next succeeding p) is or is not a prime; but the intermediate numbers $N+2, N+3, \dots, N+q-1$ are certainly composite; viz. we thus have at least $q-2$ consecutive composites. To obtain in this manner 99 consecutive composites, the value of N would be $= 2.3.5 \dots 97$, viz. this is a number far exceeding 2,637,900; but, in fact, the hundred numbers 2,637,901 to 2,638,000 are all composite.

Legendre, in his *Essai sur la Théorie des Nombres* (1st edit., 1798; 2nd edit., 1808, supplement, 1816: references to this edition), gives for the number of primes inferior to a given limit x the approximate formula

$$\frac{x}{\log x - 1.08366},$$

and p. 394, and supplement, p. 62, he compares for each 10,000 up to 100,000, and for each 100,000 up to 1,000,000, the values as computed by this formula with the actual numbers of primes exhibited by the tables of Wega and Chernac. Thus for $x = 1,000,000$, the computed value is 78,543, the actual value 78,493.

He shows, p. 414, that the number of integers, which are less than n and are not divisible by any of the numbers $\theta, \lambda, \mu, \dots$, is approximately

$$= n \left(1 - \frac{1}{\theta}\right) \left(1 - \frac{1}{\lambda}\right) \left(1 - \frac{1}{\mu}\right) \dots;$$

and taking $\theta, \lambda, \mu, \dots$ the successive primes 3, 5, 7, ... he gives the values of the function in question, or, say, the function

$$\frac{2}{3} \cdot \frac{4}{5} \cdot \frac{6}{7} \cdot \frac{10}{11} \cdots \frac{\omega - 1}{\omega},$$

ω a prime, for the several prime values $\omega = 3$ to 1229 in the Table IX. (one page) at the end of the work.

14. A table of frequency is given by

Glaisher, J. W. L., *British Association Report* for 1872, p. 20. This gives for the second and the ninth millions, respectively divided into intervals of 50,000, the actual number of primes in each interval, as compared with the theoretic value $\text{li}x' - \text{li}x$; and also deduced therefrom, by the formula $\log \frac{1}{2}(x' + x)$, a table of the average interval between two consecutive primes; this average interval increases very slowly: at the beginning and the end of the second million the values are 13.76 and 14.58 (theoretic values 13.84 and 14.50); at the beginning and the end of the ninth million 16.02 and 15.95 (theoretic values 15.90 and 16.01).

15. Coming under the head of Divisor Tables, some tables by Reuschle and Gauss may be here referred to. These are:—

Reuschle, *Mathematische Abhandlung, zahlentheoretische Tabellen sammt einer dieselben treffenden Correspondenz mit der vereinigten C. G. J. Jacobi*, 4^o, pp. 1—61* (1856). The tables belonging to the present subject are

A. Tafeln zur Zerlegung von $a^n - 1$ (pp. 18—22).

I. Table of the prime factors of $10^n - 1$, viz.

(a. pp. 18—19.) Complete decomposition of $10^n - 1$, $n = 1$ to 42; and $10^n + 1$, $n = 1$ to 21. Some values of n are omitted.

A specimen is

$$10^{13} - 1 = 3^2 \cdot 53 \cdot 79 \cdot 265371653,$$

$$10^{13} + 1 = 11 \cdot 189 \cdot 1058313049.$$

(b. p. 19.) List of the specific prime factors f of $10^n - 1$, or the prime factors of the residue after separation of the analytical factors, for those values of n for which the complete decomposition is unknown, and omitting those values for which no factor is known, $n = 25$ to 243.

* Titlepage missing in my copy; but I find from Prof. Kummer's notice of the work, *Crelle*, t. LIII. (1857), p. 379, that it appeared as a Programm of the Stuttgart Gymnasium, Michaelmas, 1856, and was separately printed by Liesching and Co., Stuttgart.

A specimen is

$$\begin{array}{cc} n & f \\ 25 & 21401. \end{array}$$

The meaning seems to be, residue of $10^{25} - 1$ is $1 + 10^5 + 10^{10} + 10^{15} + 10^{20}$, and this contains the prime factor 21401; but it is not clear why this is the "specific prime factor."

II. Prime factors of $a^n - 1$ for different values of a and n .

(a. p. 20) gives for 41 values of a (2, 3, &c. at intervals to 100) and for the following values of n the decompositions of the residues or specific factors of $a^n - 1$; viz. these are

$$\begin{array}{ll} n = 1, & a - 1: \\ \text{,,} & 2, a + 1: \\ \text{,,} & 3, a^2 + a + 1: \\ \text{,,} & 6, a^2 - a + 1: \\ \text{,,} & 4, a^2 + 1: \\ \text{,,} & 5, a^4 + a^3 + a^2 + a + 1: \\ \text{,,} & 10, a^4 - a^3 + a^2 - a + 1: \\ \text{,,} & 8, a^4 + 1: \\ \text{,,} & 12, a^4 - a^2 + 1. \end{array}$$

A specimen is

$$\begin{array}{c|c|c|c|c|c|c|c|c|c} a & a-1 & a^2-1 & a^3-1 & a^6-1 & a^4-1 & a^5-1 & a^{10}-1 & a^8-1 & a^{12}-1 \\ \hline 10 & 3^2 & 11 & 3^3 \cdot 37 & 7 \cdot 13 & 101 & 41 \cdot 271 & 9091 & 73 \cdot 137 & 9901 \end{array}.$$

(b. p. 21.) Specific prime factors for the numbers 2, 3, 5, 6, 7, 10, (the powers 4, 8, 9 being omitted as coming under 2 and 3), for the exponents 1 to 42.

A specimen is

$$\begin{array}{c|c|c|c|c|c|c} n & 2^n-1 & 3^n-1 & 5^n-1 & 6^n-1 & 7^n-1 & 10^n-1 \\ \hline 19 & 524287 & 1597 \cdot 363889 & 191 \cdot x & 191 \cdot x & 419 \cdot x & \text{---} \end{array},$$

where the x denotes that the other factor is not known to be prime. And so, where no number is given, as in $10^{19} - 1$, it is not known whether the number $(= 1 + 10^1 + 10^2 + \dots + 10^{18})$ is or is not prime.

Addition, p. 22. For $a=2$, the complete decomposition of the prime factor of $2^n - 1$ is given for values of n , $= 44, 45, \dots$ at intervals to 156.

A specimen is

$$\begin{array}{cc} n & f \\ 44 & 397 \cdot 2113, \end{array}$$

viz.

$$2^{20} - 2^{18} + 2^{16} - \dots - 2^2 + 1, = 838861 = 397 \cdot 2113.$$

$n=31$, Fermat's prime. $n=37$, the first case for which the decomposition is not given completely. $n=41$, the first case for which no factor is known.

16. **Gauss**, *Tafel zur Cyclotechnie, Werke*, t. II. pp. 478—495, shows, for 2452 numbers of the several forms $a^2 + 1$, $a^2 + 4$, $a^2 + 9$, ..., $a^2 + 81$, the values of a such that the number in question is a product of prime factors no one of which exceeds 200, and exhibits all the odd prime factors of each such number. The table is in nine parts, zerlegbare $a^2 + 1$, zerlegbare $a^2 + 4$, &c., with to each part a subsidiary table, as presently mentioned. Thus a specimen is

zerlegbare $a^2 + 9$.

1	5
2	13
4	5 . 5
5	17
7	29
8	73
⋮	⋮
1411168679	5 . 5 . 13 . 17 . 17 . 89 . 113 . 157 . 173 . 197 . 197 :

viz.

$1^2 + 9$, odd prime factor is 5,

$2^2 + 9$, „ „ 13,

$4^2 + 9$, „ factors are 5, 5,

and so on.

And the subsidiary table is

5	1, 4, 79
13	2, 11, 41
17	5, 29, 46, 379, 1042
⋮	⋮

showing that the numbers a for which the largest factor is 5 are 1, 4, 79; those for which it is 13 are 2, 11, 41; and so on.

The object of the table is explained in the *Bemerkungen*, (*l.c.*, p. 523), by Schering, the editor of the volume, viz. it is to facilitate the calculation of the circular arcs the cotangents of which are rational numbers. To take a simple example, it appears to be by means of it that Gauss obtained, among other formulæ, the following:

$$\frac{\pi}{4} = 12 \arctan \frac{1}{18} + 8 \arctan \frac{1}{57} - 5 \arctan \frac{1}{239},$$

and

$$= 12 \arctan \frac{1}{38} + 20 \arctan \frac{1}{57} + 7 \arctan \frac{1}{239} + 24 \arctan \frac{1}{268}.$$

[F. 12. *Divisors, &c.*] continued. *Prime Roots. The Canon Arithmeticus, Quadratic residues.* Art. II.

17. *Prime Roots.*—Let p be a prime number; then there exist $\varpi(p-1)$ inferior integers g , such that all the numbers $1, 2, \dots, p-1$ are, to the modulus p ,

$$\equiv 1, g, g^2, \dots, g^{p-2} \quad (g^{p-1} \text{ is of course } \equiv 1).$$

This being so, g is said to be a prime root of p ; and moreover the several numbers g^α , where α is any number whatever less than and prime to $p-1$, constitute the series of the $\varpi(p-1)$ prime roots of p . It may be added that, if β be an integer number less than $p-1$, and having with it a greatest common measure $=k$, so that

$$(g^\beta)^{\frac{p-1}{k}} \equiv g^{\frac{\beta}{k}(p-1)} \equiv 1, \quad \left(\text{since } \frac{\beta}{k} \text{ is an integer, and } g^{p-1} \equiv 1 \right),$$

then g^β has the indicatrix $\frac{p-1}{k}$: the prime roots are those numbers which have the indicatrix $p-1$.

The like theory exists as to any number N of the form p^m or $2p^m$. There are here $\varpi(N) = N \left(1 - \frac{1}{p}\right)$ or $\frac{1}{2}N \left(1 - \frac{1}{p}\right)$, in the two cases respectively, numbers less than N and prime to it; and we have then $\varpi(\varpi(N))$ numbers g such that, to the modulus N , all these numbers are $\equiv 1, g, g^2, \dots, g^{\varpi(N)-1}$ ($g^{\varpi(N)}$ is of course $\equiv 1$). This being so, g may be regarded as a prime root of N ($=p^m$ or $2p^m$, as the case may be); and moreover the several numbers g^α , where α is any number whatever less than and prime to $\varpi(N)$, constitute the series of the $\varpi(\varpi(N))$ prime roots of N . Thus $N=3^2=9$, $\varpi(N)=6$; we have

$$\begin{aligned} &1, \quad 2^1, \quad 2^2, \quad 2^3, \quad 2^4, \quad 2^5, \\ &\equiv 1, \quad 2, \quad 4, \quad 8, \quad 7, \quad 5, \text{ mod. } 9; \end{aligned}$$

or the prime roots of 9 are 2^1 and $2^5 = 2$ and 5.

So also $N=2 \cdot 3^2=18$, $\varpi(N)=6$; we have

$$\begin{aligned} &1, \quad 5^1, \quad 5^2, \quad 5^3, \quad 5^4, \quad 5^5, \\ &\equiv 1, \quad 5, \quad 7, \quad 17, \quad 13, \quad 11, \text{ mod. } 18; \end{aligned}$$

and 5^1 and $5^5 = 5$ and 11 are the prime roots of 18.

18. A small table of prime roots, $p=3$ to 37, is given by

Euler, *Op. Arith. Coll.* t. i. pp. 525—526. The Memoir is entitled "Demonstrationes circa residua e divisione potestatum per numeros primos resultantia," pp. 516—537 (1772).

19. A table, p and p^m , 3 to 97, is given by

Gauss, "Disquisitiones Arithmeticae," 1801, (*Werke*, t. i. p. 468). This gives in each case a prime root, and it shows the exponents in regard thereto of the several prime numbers less than p or p^m . Thus a specimen is

	2	3	5	7	11	13	17	19	23	29	&c.
27	2	1	*	5	16	13	8	15	12	11	
29	10	11	27	18	20	23	2	7	15	24	

viz. for 27 we have 2 a prime root, and $2 \equiv 2^1$, $5 \equiv 2^5$, $7 \equiv 2^{16}$, $11 \equiv 2^{13}$, &c.; and so also for 29 we have 10 a prime root, and $2 \equiv 10^{11}$, $3 \equiv 10^{27}$, $5 \equiv 10^{18}$, &c.

20. Small tables are probably to be found in many other places; but the most extensive and convenient table is **Jacobi's** *Canon Arithmeticus*, the complete title of which is

Canon Arithmeticus sive tabula quibus exhibentur pro singulis numeris primis vel primorum potestatibus infra 1000 numeri ad datos indices et indices ad datos numeros pertinentes. Edidit C. G. J. Jacobi. Berolini, 1839. 4^o.

The contents are as follows:—

	Pages
Introductio	i to xl
Tabulæ numerorum ad indices datos pertinentium et indicum numero dato correspondentium pro modulis primis minoribus quam 1000	1—221
Tabulæ residuorum et indicum sibi mutuo respondentium pro modulis minoribus quam 1000 qui sunt numerorum primorum potestates	222—238
Hujus tabula ea pars quæ pertinet ad modulos formæ 2^n , invenitur	239—240.

The following is a specimen of the principal tables:—

$$p = 19, p - 1 = 2 \cdot 3^2.$$

Numeri.

I	0	1	2	3	4	5	6	7	8	9
		10	5	12	6	3	11	15	17	18
	9	14	7	13	16	8	4	2	1	

Indices.

N	0	1	2	3	4	5	6	7	8	9
		18	17	5	16	2	4	12	15	10
1	1	6	3	13	11	7	14	8	9	

where the first table gives the values of the powers of the prime root 10 (that 10 is the root appears by its index being given as =1) to the modulus 19, viz. $10^1 \equiv 10$, $10^2 \equiv 5$, $10^3 \equiv 12$, &c.; and the second table gives the index of the power to which the same prime root must be raised in order that it may be, to the modulus 19, congruent with a given number: thus $10^{18} \equiv 1$, $10^{17} \equiv 2$, $10^5 \equiv 3$, &c. The units of the index or number, as the case may be, are contained in the top line of the table, and the tens or hundreds and tens in the left-hand column.

21. There is given by

Jacobi, *Crelle*, t. xxx. (1846), pp. 181, 182, a table of m' for the argument m , such that

$$1 + g^m \equiv g^{m'} \pmod{p}, \quad p = 7 \text{ to } 103, \text{ and } m = 0 \text{ to } 102.$$

A specimen is

p	7	11	13	17	19	23	29	31	37	...	to 103
g	3	2	6	10	10	10	10	17	5		
m											
11	.	.	6	4	7	*	27	21	34		

for instance, $p = 19$, $1 + 10^{11} \equiv 10^7 \pmod{19}$.

Jacobi remarks that this table was calculated for him by his class during the winter course of 1836—37; and that, by means of the *Canon Arithmeticus* since published (in 1839), the same might easily be extended to all primes under 1000. In fact, for any such number p , putting any number of the table "Indices" = m , the next following number of the table gives the value of m' .

22. We have next, in **Reuschle's** Memoir (*ante*, No. 15), the following relating to prime roots:—

C. Tafeln für primitive Wurzeln und Hauptexponenten, oder V. erweiterte und bereicherte Burkhardsche Tafel, pp. 41—61, being divided into three parts; viz. these are

a. Table of the Hauptexponenten of the six roots 10, 5, 2, 6, 3, 7 for all prime numbers of the first 1000, together with the least primitive root of each of these numbers (pp. 42—46).

A specimen is as follows:—

p	$p-1$	$\overbrace{e \ n}^{10}$	$\overbrace{e \ n}^5$	$\overbrace{e \ n}^2$	$\overbrace{e \ n}^6$	$\overbrace{e \ n}^3$	$\overbrace{e \ n}^7$	w
53	2 ² .13	13 4	52 1	52 1	26 2	52 1	26 2	2

where e is the Hauptexponent or indicatrix of the root (10, 5, 2, 6, 3, 7, as the case may be), $n = \frac{p-1}{e}$, w the least primitive root; thus

$$p = 53, \quad 10^{13} \equiv 1, \quad 5^{52} \equiv 1, \quad 2^{52} \equiv 1,$$

(2 being accordingly the least prime root),

$$6^{26} \equiv 1, \quad 3^{32} \equiv 1, \quad 7^{26} \equiv 1.$$

The number w of the last column is the least primitive root. It is, of course, not always (as in the present case) one of the numbers 10, 5, 2, 6, 3, 7 to which the table relates: the first exception is $p=191$, $w=19$: the highest value of w is $w=21$, corresponding to $p=409$.

b. The like table for the roots 10 and 2 for all prime numbers from 1000 to 5000, together with as convenient as possible a prime root (and in some cases two prime roots) for each such number (pp. 47—53).

A specimen is:—

p	$p-1$	$\overbrace{\quad}^{10}$		$\overbrace{\quad}^2$		w
		e	n	e	n	
1289	2 ³ .7.23	92	14	161	8	6, 11

viz. here, mod. 1289, $10^{92} \equiv 1$, $2^{161} \equiv 1$; and two prime roots are 6, 11. We have thus by the present tables a prime root for every prime number not exceeding 5000.

c. The like table for the root 10 for all prime numbers between 5000 and 15000, (no column for w , nor any prime root given), pp. 53—61.

A specimen is

p	$p-1$	e	n
9859	2.3.31.53	3286	3:

viz., mod. 9859, we have $10^{3286} \equiv 1$. But in a large number of cases we have $n=1$, and therefore 10 a prime root. For example,

9887	2.4983	9886	1.
------	--------	------	----

23. For a composite number n , if $N=\varpi(n)$ be the number of integers less than n and prime to it, then if x be any number less than n and prime to it, we have $x^N \equiv 1 \pmod{n}$. But we have in this case no analogue of a prime root—there is no number x , such that its several powers $x^1, x^2, \dots, x^{N-1} \pmod{n}$ are all different from unity; or, what is the same thing, there is for each value of x some submultiple of N , say N' , such that $x^{N'} \equiv 1 \pmod{n}$. And these several numbers N' have a least common multiple I , which is not $=N$, but is a submultiple of N ; and this being so, then for all the several values of x , I is said to be the maximum indicator. For instance, $n=12$, $N=\varpi(n)$; the numbers less than 12 and prime to it are 1, 5, 7, 11. We have, (mod. 12), $1^1 \equiv 1$, $5^2 \equiv 1$, $7^2 \equiv 1$, $11^2 \equiv 1$, or the values of N' are 1, 2, 2, 2; their least common multiple is 2, and we have accordingly $I=2$: viz. $x^2 \equiv 1 \pmod{12}$ has the $\varpi(12)$ roots 1, 5, 7, 11. So $n=24$, $\varpi(n)=8$; the maximum indicator I is in this case also $=2$.

A table of the maximum indicator $n=1$ to 1000 is given by

Cauchy, *Exer. d'Analyse &c.*, t. II. (1841), pp. 36—40, contained in the "Mémoire sur la résolution des équations indéterminées du premier degré en nombres entiers," pp. 1—40.

24. It thus appears that for a composite number n , the $\varpi(n)$ numbers less than n and prime to it cannot be expressed as $\equiv (\text{mod. } n)$ to the power of a single root; but for the expression of them it is necessary to employ two or more roots. A small table, $n=1$ to 50, is given by

Cayley, Specimen Table $M \equiv a^x b^y (\text{mod. } N)$ for any prime or composite modulus; *Quart. Math. Journ.* vol. IX. (1868), pp. 95, 96, and folding sheet, [397].

A specimen is

Nos. roots Ind. M.I. ϕ	12 5, 7 2, 2 2 4
1	0, 0
2	
3	
4	
5	1, 0
6	
7	0, 1
8	
9	
10	
11	1, 1

viz. for the modulus 12 the roots are 5, 7, having respectively the indicators 2, 2, viz. $5^2 \equiv 1 (\text{mod. } 12)$, $7^2 \equiv 1 (\text{mod. } 12)$. Hence also the maximum indicator is $= 2$. $\phi (= \varpi(n)) = 4$ is the number of integers less than 12 and prime to it, viz. these are 1, 5, 7, 11, which in terms of the roots 5, 7 and to mod. 12 are respectively $\equiv 5^0 \cdot 7^0$, $5^1 \cdot 7^0$, $5^0 \cdot 7^1$, and $5^1 \cdot 7^1$.

25. *Quadratic Residues*.—In regard to a given prime number p , a number N is or is not a quadratic residue according as the index of N is even or odd, viz. g being a prime root and $N \equiv g^a$, then N is or is not a quadratic residue according as a is even or odd. But the quadratic residues can, of course, be obtained directly without the consideration of prime roots.

A small table, $p=3$ to 97 and $N=-1$ and (prime values) 3 to 97, is given by

Gauss, "Disquisitiones Arithmeticae," 1801; Table II. (*Werke*, t. I. p. 469): I notice here a misprint in the top line of the original; it should be $-1, +2, +3$, &c., instead of

1, + 2, + 3, &c.; the - 1 is printed correctly on p. 499 of the French translation *Recherches Arithmétiques*, Paris, 1807 and on p. 469 of vol. I. of *Werke*, (Göttingen, 1870).

A specimen is

	- 1	+ 2	+ 3	+ 5	+ 7	+ 11	+ 13	+ 17	+ 19	+ 23	&c.
19				-	-	-		-	-	-	

viz. - 1, 2, 3, 13 are not, 5, 7, 11, 17 &c. are, quadratic residues of 19. The residues taken positively and less than 19 are, in fact, 1, 4, 5, 6, 7, 11, 16, 17.

The same table carried from $p = 3$ to 503, and prime values $N = 3$ to 997, is given by

Gauss, *Werke*, t. II. pp. 400—409. A specimen is

	2	3	5	7	11	13	17	19	23	&c.;
19			-	-	-		-	-	-	

viz. the arrangement is the same, except only that the - 1 column is omitted.

26. We have also by **Gauss**

"Disquisitiones Arithmeticae" Table III. (*Werke*, t. I. p. 470), for the conversion into decimals of a vulgar fraction, denominator p or p^μ , not exceeding 100. The explanation is given in Art. 314 *et seq.* of the same work.

But this table, carried to a greater extent, is given by **Gauss**, *Werke*, t. II. pp. 412—434, "Tafel zur Verwandlung gemeiner Brüche mit Nennern aus dem ersten Tausend in Decimalbrüche;" viz. the denominators are here primes or powers of primes, p^μ up to 997.

To explain the table, consider a modulus p^μ (where μ may be = 1); if 10 is not a prime root of p^μ , consider a prime root r , which is such that $r^e \equiv 10 \pmod{p^\mu}$, e being a submultiple of $p^{\mu-1}(p-1)$; say we have $ef = p^{\mu-1}(p-1)$: then $10^f \equiv 1 \pmod{p^\mu}$. Consider any fraction $\frac{N}{p^\mu}$; then we may write $N \equiv r^{kl+l} \pmod{p^\mu}$, k from 0 to $f-1$ and l from 0 to $e-1$, $\equiv 10^k r^l$, and consequently $\frac{N}{p^\mu}$ and $\frac{10^k r^l}{p^\mu}$ have the same mantissa (decimal part regarded as an integer); hence, in order to know the mantissa of every fraction whatever of $\frac{N}{p^\mu}$, it is sufficient to know the mantissa of $\frac{r^l}{p^\mu}$, that is, the mantissæ of $\frac{1}{p^\mu}, \frac{r}{p^\mu}, \frac{r^2}{p^\mu}, \dots, \frac{r^{e-1}}{p^\mu}$, or, what is the same thing, the mantissæ of $\frac{10}{p^\mu}, \frac{10r}{p^\mu}, \dots, \frac{10r^{e-1}}{p^\mu}$.

For instance, $p^\mu = 11$, $10^2 \equiv 1 \pmod{11}$, whence $f = 2$, $e = 5$; and taking $r = 2$, we have $10 \equiv r^5 \pmod{11}$.

The required mantissæ, denoted in the table by

$$\begin{array}{ccccc} (0), & (1), & (2), & (3), & (4), \\ \text{are those of} & \frac{10}{11}, & \frac{10 \cdot 2}{11}, & \frac{10 \cdot 2^2}{11}, & \frac{10 \cdot 2^3}{11}, & \frac{10 \cdot 2^4}{11}, \end{array}$$

viz. these fractions are respectively =

$$\begin{array}{ccccc} (0), & (1), & (2), & (3), & (4), \\ .9090 \dots, & 1.8181 \dots, & 3.6363 \dots, & 7.2727 \dots, & 14.5454 \dots; \end{array}$$

or their mantissæ are 90, 81, 63, 27, 54.

And we accordingly have as a specimen

$$11 \mid (1) \dots 81, (2) \dots 63, (3) \dots 27, (4) \dots 54, (0) \dots 90.$$

Or again, as another specimen, $r = 2$:—

$$27 \mid (1) \dots 740, (2) \dots 481, (3) \dots 962, (4) \dots 925, (5) \dots 851, (0) \dots 370.$$

The table in this form extends to $p^\mu = 463$; the values of r (not given in the body of the table) are annexed, p. 420.

In the latter part of the table $p^\mu = 467$ to 997, we have only the mantissæ of $\frac{100}{p^\mu}$. A specimen is

$$\begin{array}{l} 547 \mid \begin{array}{llll} 1828153564 & 8994515539 & 3053016453 & 3820840950 \\ 6398537477 & 1480804387 & 5685557586 & 8372943327 \\ 2394881170 & 0182815356, & & \end{array} \end{array}$$

viz. the fraction $\frac{100}{547} = .182815 \dots$ has a period of 91, $= \frac{1}{8} 546$, figures.

[F. 13. *The Pellian Equation.*] Art. III.

27. The Pellian equation is $y^2 = ax^2 + 1$, a being a given integer number, which is not a square (or rather, if it be, the only solution is $y = 1, x = 0$), and x, y being numbers to be determined: what is required is the least values of x, y , since these, being known, all other values can be found. A small table $a = 2$ to 68 is given by

Euler, *Op. Arith. Coll.* t. I. p. 8. The Memoir is "Solutio problematum Diophanteorum per numeros integros," pp. 4—10, 1732—33. The form of the table is

a	$x (= p)$	$y (= q)$
2	2	3
3	1	2
5	4	9
$\vdots$	$\vdots$	$\vdots$
68	4	33.

Even here, for some of the values of a , the values of x , y are extremely large; thus $a = 61$, $x = 226,153,980$, $y = 1,766,399,049$.

And probably tables of a like extent may be found elsewhere; in particular, a table of the solution of $y^2 = ax^2 \pm 1$ (— when the value of a is such that there is a solution of $y^2 = ax^2 - 1$, and + for other values of a), $a = 2$ to 135, is given by **Legendre**, *Théorie des Nombres*, 2nd ed. 1808, in the Table X. (one page) at the end of the work. For the before-mentioned number 61, the equation is $y^2 = 61x^2 - 1$, and the values are $x = 3805$, $y = 29718$; much smaller than Euler's values for the equation $y^2 = 61x^2 + 1$.

28. The most extensive table, however, is given by

Degen, *Canon Pellianus, sive Tabula simplicissimam equationis celebratissimæ: $y^2 = ax^2 + 1$, solutionem, pro singulis numeri dati valoribus ab 1 usque ad 1000 in numeris rationalibus, iisdemque integris exhibens*. Auctore Carolo Ferdinando Degen. Hafn (Copenhagen) apud Gerhardum Bonnorum, 1817. 8vo. pp. iv to xxiv and 1 to 112.

The first table (pp. 3—106) is entitled as "Tabula I. Solutionem Equationis $y^2 - ax^2 - 1 = 0$ exhibens." It, in fact, also gives the expression of $\sqrt{a}$ as a continued fraction; thus a specimen is

209	14	2	5	3	(2)
	1	13	5	8	11
	3220				
	46551				

Here the first line gives the continued fraction, viz.

$$\sqrt{209} = 14 + \frac{1}{2 + \frac{1}{5 + \frac{1}{3 + \frac{1}{2 + \frac{1}{3 + \frac{1}{5 + \frac{1}{2 + \frac{1}{28 + \frac{1}{2} + \&c.}}}}}}}}$$

the period being (2, 5, 3, 2, 3, 5, 2) indicated by 2, 5, 3 (2). [The number of terms in the period is here odd, but it may be even; for instance, the period (1, 1, 5, 5, 1, 1) is indicated by 1, 1 (5, 5).]

The second line contains auxiliary numbers presenting themselves in the process; thus, if $R^2 = 239$, we have $R = 14 + \frac{1}{\alpha}$,

$$\alpha = \frac{1}{R - 14} = \frac{1(R + 14)}{209 - 14^2} = \frac{R + 14}{13} = 2 + \frac{1}{\beta},$$

$$\beta = \frac{13}{R - 12} = \frac{13(R + 12)}{209 - 12^2} = \frac{R + 12}{5} = 5 + \frac{1}{\gamma},$$

$$\gamma = \frac{5}{R - 13} = \frac{5(R + 13)}{209 - 13^2} = \frac{R + 13}{8} = 3 + \frac{1}{\delta},$$

&c.,