

James Joseph Sylvester
Collected Mathematical Papers, Vol. I
Book pp. 1–268

1.

ANALYTICAL DEVELOPMENT OF FRESNEL'S OPTICAL THEORY OF CRYSTALS

[*Philosophical Magazine*, XI. (1837), pp. 461-469, 537-541; XII. (1838), pp. 73-83, 341-345]

p. 1

The following is, I believe, the first successful attempt to obtain the full development of Fresnel's Theory of Crystals by direct geometrical methods. Hitherto little has been done beyond finding and investigating the properties of the wave surface, a subject certainly curious and interesting, but not of chief importance for ordinary practical purposes. Mr Kelland, in a most valuable contribution to the *Cambridge Philosophical Transactions*¹, has incidentally obtained the difference of the squares of the velocities of a plane front in terms of the angles made by it with the optic axes. I have obtained each of the velocities separately, and in a form precisely the same for biaxal as for uniaxal crystals.

I have also assigned in my last proposition the place of the lines of vibration in terms of the like quantities, and that in a shape remarkably convenient for determining the plane of polarization when the ray is given. For at first sight there appears to be some ambiguity in selecting which of the two lines of vibration is to be chosen when the front is known. If p be the perpendicular from the centre of the surface of elasticity let fall upon the front, ι_1, ι_2 the angles made by the front with the optic planes, $\varepsilon_1, \varepsilon_2$ the angles between its due line of vibration and the optic axes, I have shown that

$$\cos \varepsilon_1 = \sqrt{\frac{b^2 - p^2}{a^2 - c^2} \cdot \frac{\sin \iota_1}{\sin \iota_2}}, \quad \cos \varepsilon_2 = \sqrt{\frac{b^2 - p^2}{a^2 - c^2} \cdot \frac{\sin \iota_2}{\sin \iota_1}},$$

so that all doubt is completely removed. The equation preparatory to obtaining the wave surface is found in Prop. 6 by common algebra, without any use of the properties of maxima and minima, and various other curious relations are discussed.

Without the most careful attention to preserve pure symmetry, the expressions could never have been reduced to their present simple forms.

p. 2

ANALYTICAL REDUCTION OF FRESNEL'S OPTICAL THEORY OF CRYSTALS.

Index of Contents.

In Proposition 1, a plane front within a crystal being given, the two lines of vibration are investigated.

¹See *Lond. and Edinb. Phil. Mag.* Vol. x. p. 336.

In Proposition 2 it is shown that the product of the cosines of the inclinations of one of the axes of elasticity to the two lines of vibration, is to the same for either other axis of elasticity in a constant ratio for the same crystal; and the two lines of vibration are proved to be perpendicular to each other.

In Proposition 3, a line of vibration being given, the front to which it belongs is determined; and it is proved that there is only one such, and consequently any line of vibration has but one other line conjugate to it.

In Proposition 4, certain relations are instituted between the positions of, and velocities due to, conjugate lines.

In Proposition 5, the angles made by the front with the planes of elasticity are found in terms of the velocities only.

In Proposition 6, the above is reversed.

In Proposition 7, the position of the planes in which the two velocities are equal (viz. the optic planes) is determined.

In Proposition 8, the position of a front in respect to the optic axes is expressed in terms of the velocities.

In Proposition 9, the problem is reversed, and it is shown that if v_1, v_2 be the two normal velocities with which any front can move perpendicular to itself, and ι_1, ι_2 the angles which it makes with the optic planes, then

$$v_1^2 = a^2 \left(\sin \frac{\iota_1 + \iota_2}{2} \right)^2 + c^2 \left(\cos \frac{\iota_1 + \iota_2}{2} \right)^2,$$

$$v_2^2 = a^2 \left(\sin \frac{\iota_1 - \iota_2}{2} \right)^2 + c^2 \left(\cos \frac{\iota_1 - \iota_2}{2} \right)^2.$$

In the 10th the angle made by a line of vibration with the axes of elasticity is expressed in terms of the two velocities of the front to which it belongs.

In the 11th Proposition the velocity due to any line of vibration is expressed in terms of the angles which it makes with the optic axes, viz.

$$v^2 - b^2 = (a^2 - c^2) \cos \varepsilon_1 \cos \varepsilon_2.$$

In the 12th Proposition $\varepsilon_1, \varepsilon_2$ are separately expressed in terms of ι_1, ι_2 .

In the Appendix I have given the polar or rather radio-angular equation to the wave surface, from which the celebrated proposition of the ray flows as an immediate consequence.

p. 3

PROPOSITION 1.

If

$$lx + my + nz = 0 \tag{a}$$

be the equation to a given front, to determine the lines of vibration therein.

It is clear that if x, y, z be any point in one of these lines, the force acting on a particle placed there when resolved into the plane must tend to the centre. Consequently the line of force at x, y, z must meet the perpendicular drawn upon the front from the origin. Now the equation to this perpendicular is

$$\frac{X}{l} = \frac{Y}{m} = \frac{Z}{n} \quad (1)$$

and the forces acting at x, y, z are a^2x, b^2y, c^2z parallel to x, y, z , so that the equation to the line of force is

$$\frac{X - x}{a^2x} = \frac{Y - y}{b^2y} = \frac{Z - z}{c^2z}. \quad (2)$$

From (2) we obtain

$$b^2yX - a^2xY = (b^2 - a^2)xy, \quad (3)$$

$$c^2zY - b^2yZ = (c^2 - b^2)yz, \quad (4)$$

$$a^2xZ - c^2zX = (a^2 - c^2)zx. \quad (5)$$

Hence

$$\begin{aligned} (b^2 - a^2)xyn + (c^2 - b^2)yzl + (a^2 - c^2)zxm \\ = b^2y(nX - lZ) + c^2z(lY - mX) + a^2x(mZ - nY); \end{aligned}$$

but by equations (1)

$$lZ - nX = 0, \quad mX - lY = 0, \quad nY - mZ = 0$$

therefore

$$(b^2 - a^2)\frac{n}{z} + (c^2 - b^2)\frac{l}{x} + (a^2 - c^2)\frac{m}{y} = 0. \quad (b)$$

Also we have

$$nz + lx + my = 0 \quad (a)$$

therefore

$$(b^2 - a^2)n^2 + (c^2 - b^2)l^2 + nl \left\{ (c^2 - b^2)\frac{z}{x} + (b^2 - a^2)\frac{x}{z} \right\} = (a^2 - c^2)m^2$$

or

$$(c^2 - b^2) \left(\frac{z}{x} \right)^2 + \frac{1}{nl} \left\{ (c^2 - b^2)l^2 + (b^2 - a^2)n^2 - (a^2 - c^2)m^2 \right\} \frac{z}{x} + (b^2 - a^2) = 0.$$

And in like manner interchanging b, y, m with c, z, n

$$(b^2 - c^2) \left(\frac{y}{x} \right)^2 + \frac{1}{ml} \left\{ (b^2 - c^2)l^2 + (c^2 - a^2)m^2 - (a^2 - b^2)n^2 \right\} \frac{y}{x} + (c^2 - a^2) = 0.$$

Hence if $\left(\frac{y_1}{x_1}, \frac{z_1}{x_1}\right), \left(\frac{y_2}{x_2}, \frac{z_2}{x_2}\right)$ be the two systems of values of $\frac{y}{x}, \frac{z}{x}$, then

$$\left(\frac{Y}{X} = \frac{y_1}{x_1}, \frac{Z}{X} = \frac{z_1}{x_1}\right) \left(\frac{Y}{X} = \frac{y_2}{x_2}, \frac{Z}{X} = \frac{z_2}{x_2}\right)$$

are the two lines of vibration required.

PROPOSITION 2.

By last proposition it appears that

$$\frac{y_1 y_2}{x_1 x_2} = \frac{c^2 - a^2}{b^2 - c^2} \quad (c)$$

and

$$\frac{z_1 z_2}{x_1 x_2} = \frac{b^2 - a^2}{c^2 - b^2} \quad (d)$$

therefore

$$\frac{y_1 y_2 + z_1 z_2}{x_1 x_2} = \frac{c^2 - b^2}{b^2 - c^2} = -1$$

therefore

$$x_1 x_2 + y_1 y_2 + z_1 z_2 = 0.$$

And therefore the two lines of vibration are perpendicular to each other.

N.B. Equations (c) and (d) must not be overlooked.

PROPOSITION 3.

A line of vibration is given (that is $\frac{y_1}{x_1}, \frac{z_1}{x_1}$ are given) and the position of the front is to be determined.

Let $lx + my + nz = 0$ be the front required, then $lx_1 + my_1 + nz_1 = 0$, and

$$(b^2 - c^2) \frac{l}{x_1} + (c^2 - a^2) \frac{m}{y_1} + (a^2 - b^2) \frac{n}{z_1} = 0.$$

Eliminating n we get

$$l \left\{ (a^2 - b^2) \frac{x_1}{z_1} - (b^2 - c^2) \frac{z_1}{x_1} \right\} + m \left\{ (a^2 - b^2) \frac{y_1}{z_1} - (c^2 - a^2) \frac{z_1}{y_1} \right\} = 0$$

therefore

$$\begin{aligned} \frac{l}{m} &= \frac{x_1}{y_1} \frac{(a^2 - b^2)y_1^2 - (c^2 - a^2)z_1^2}{(b^2 - c^2)z_1^2 - (a^2 - b^2)x_1^2} \\ &= \frac{x_1}{y_1} \frac{a^2(x_1^2 + y_1^2 + z_1^2) - (a^2x_1^2 + b^2y_1^2 + c^2z_1^2)}{b^2(x_1^2 + y_1^2 + z_1^2) - (a^2x_1^2 + b^2y_1^2 + c^2z_1^2)}. \end{aligned}$$

If now we make $x_1^2 + y_1^2 + z_1^2 = 1$

$$a^2 x_1^2 + b^2 y_1^2 + c^2 z_1^2 = v_1^2$$

and therefore

$$\frac{l}{m} = \frac{x_1}{y_1} \cdot \frac{a^2 - v_1^2}{b^2 - v_1^2}$$

and in like manner

$$\frac{l}{n} = \frac{x_1}{z_1} \cdot \frac{a^2 - v_1^2}{c^2 - v_1^2};$$

therefore

$$(a^2 - v_1^2)x_1x + (b^2 - v_1^2)y_1y + (c^2 - v_1^2)z_1z = 0$$

is the equation required.

PROPOSITION 4.

$\frac{l}{m}, \frac{l}{n}$ having each only one value, shows that only one front corresponds to the given line of vibration. Let x_2, y_2, z_2, v_2 correspond to x_1, y_1, z_1, v_1 for the conjugate line of vibration, then the equation to the front may be expressed likewise by

$$(a^2 - v_2^2)x_2x + (b^2 - v_2^2)y_2y + (c^2 - v_2^2)z_2z = 0,$$

so that

$$\frac{(a^2 - v_1^2)x_1}{(a^2 - v_2^2)x_2} = \frac{(b^2 - v_1^2)y_1}{(b^2 - v_2^2)y_2} = \frac{(c^2 - v_1^2)z_1}{(c^2 - v_2^2)z_2}.$$

PROPOSITION 5.

To find ω, ϕ, ψ , the angles made by the front with the planes of elasticity in terms of v_1, v_2 .

By the last proposition

$$\begin{aligned} (\cos \omega)^2 &= \frac{(a^2 - v_1^2)^2 x_1^2}{(a^2 - v_1^2)^2 x_1^2 + (b^2 - v_1^2)^2 y_1^2 + (c^2 - v_1^2)^2 z_1^2} \\ &= \frac{(a^2 - v_1^2)(a^2 - v_2^2)x_1x_2}{(a^2 - v_1^2)(a^2 - v_2^2)x_1x_2 + (b^2 - v_1^2)(b^2 - v_2^2)y_1y_2 + (c^2 - v_1^2)(c^2 - v_2^2)z_1z_2}. \end{aligned}$$

Now, by Proposition 2,

$$\frac{x_1x_2}{c^2 - b^2} = \frac{y_1y_2}{a^2 - c^2} = \frac{z_1z_2}{b^2 - a^2}$$

therefore $(\cos \omega)^2$

$$\begin{aligned}
&= \frac{(a^2 - v_1^2)(a^2 - v_2^2)(c^2 - b^2)}{(a^2 - v_1^2)(a^2 - v_2^2)(c^2 - b^2) + (b^2 - v_1^2)(b^2 - v_2^2)(a^2 - c^2) + (c^2 - v_1^2)(c^2 - v_2^2)(b^2 - a^2)} \\
&= \frac{(a^2 - v_1^2)(a^2 - v_2^2)(c^2 - b^2)}{a^4(c^2 - b^2) + b^4(a^2 - c^2) + c^4(a^2 - b^2)} \\
&= \frac{(a^2 - v_1^2)(a^2 - v_2^2)}{(a^2 - b^2)(a^2 - c^2)}.
\end{aligned}$$

Similarly,

$$\begin{aligned}
(\cos \phi)^2 &= \frac{(b^2 - v_1^2)(b^2 - v_2^2)}{(b^2 - a^2)(b^2 - c^2)}, \\
(\cos \psi)^2 &= \frac{(c^2 - v_1^2)(c^2 - v_2^2)}{(c^2 - a^2)(c^2 - b^2)}.
\end{aligned}$$

PROPOSITION 6.

To find v_1, v_2 in terms of ω, ϕ, ψ .

By the last proposition

$$\begin{aligned}
\frac{(\cos \omega)^2}{a^2 - v_1^2} &= \frac{a^2}{(a^2 - b^2)(a^2 - c^2)} - v_2^2 \cdot \frac{1}{(a^2 - b^2)(a^2 - c^2)}, \\
\frac{(\cos \phi)^2}{b^2 - v_1^2} &= \frac{b^2}{(b^2 - a^2)(b^2 - c^2)} - v_2^2 \cdot \frac{1}{(b^2 - a^2)(b^2 - c^2)}, \\
\frac{(\cos \psi)^2}{c^2 - v_1^2} &= \frac{c^2}{(c^2 - b^2)(c^2 - a^2)} - v_2^2 \cdot \frac{1}{(c^2 - a^2)(c^2 - b^2)},
\end{aligned}$$

therefore

$$\frac{(\cos \omega)^2}{a^2 - v_1^2} + \frac{(\cos \phi)^2}{b^2 - v_1^2} + \frac{(\cos \psi)^2}{c^2 - v_1^2} = 0.$$

Just in the same way

$$\frac{(\cos \omega)^2}{a^2 - v_2^2} + \frac{(\cos \phi)^2}{b^2 - v_2^2} + \frac{(\cos \psi)^2}{c^2 - v_2^2} = 0,$$

so that v_1^2, v_2^2 are the two roots of the equation

$$\frac{(\cos \omega)^2}{a^2 - v^2} + \frac{(\cos \phi)^2}{b^2 - v^2} + \frac{(\cos \psi)^2}{c^2 - v^2} = 0.$$

COR. Hence the equation to the wave surface may be obtained by making

$$(\cos \omega)x + (\cos \phi)y + (\cos \psi)z = v,$$

or if we please to apply Prop. 5, we may make

$$\sqrt{\frac{(a^2 - v_1^2)(a^2 - v_2^2)}{(a^2 - b^2)(a^2 - c^2)}} x + \sqrt{\frac{(b^2 - v_1^2)(b^2 - v_2^2)}{(b^2 - a^2)(b^2 - c^2)}} y + \sqrt{\frac{(c^2 - v_1^2)(c^2 - v_2^2)}{(c^2 - a^2)(c^2 - b^2)}} z = v_1,$$

or, if we please²,

$$\sqrt{\frac{(a^2 u^2 - 1)(a^2 - v^2)}{(a^2 - b^2)(a^2 - c^2)}} x + \sqrt{\frac{(b^2 u^2 - 1)(b^2 - v^2)}{(b^2 - a^2)(b^2 - c^2)}} y + \sqrt{\frac{(c^2 u^2 - 1)(c^2 - v^2)}{(c^2 - a^2)(c^2 - b^2)}} z = 1.$$

PROPOSITION 7.

To find when $v_1 = v_2$.

By Prop. 4,

$$\frac{x_1(v_1^2 - a^2)}{x_2(v_2^2 - a^2)} = \frac{y_1(v_1^2 - b^2)}{y_2(v_2^2 - b^2)} = \frac{z_1(v_1^2 - c^2)}{z_2(v_2^2 - c^2)}. \quad (\theta)$$

Hence when $v_1 = v_2$ we have, generally speaking,

$$\frac{x_1}{x_2} = \frac{y_1}{y_2} = \frac{z_1}{z_2}.$$

Now

$$x_1 x_2 + y_1 y_2 + z_1 z_2 = 0;$$

therefore $x_1^2 + y_1^2 + z_1^2$ would be 0, which is absurd.

The only case therefore when $v_1 = v_2$ is when one of those terms of equation (θ) becomes $\frac{0}{0}$: thus suppose $v_1 = b$, then we have $\frac{x_1}{x_2} = \frac{z_1}{z_2} = \frac{0}{0}$, and we can no longer infer $\frac{x_1}{x_2} = \frac{y_1}{y_2}$.

Let now $(\omega_1, \phi_1, \psi_1)(\omega_2, \phi_2, \psi_2)$ be the two systems of values which ω, ϕ, ψ assume when $v_1 = v_2 = b$, then applying the equation of Prop. 5 we have

$$\begin{aligned} \cos \omega_1 &= \sqrt{\frac{a^2 - b^2}{a^2 - c^2}} & \cos \omega_2 &= \sqrt{\frac{a^2 - b^2}{a^2 - c^2}} \\ \cos \phi_1 &= 0 & \cos \phi_2 &= 0 \\ \cos \psi_1 &= \sqrt{\frac{b^2 - c^2}{a^2 - c^2}} & \cos \psi_2 &= \sqrt{\frac{b^2 - c^2}{a^2 - c^2}}, \end{aligned}$$

²See below, p. 27. ED.

so that b must correspond to the mean axis.

p. 8

PROPOSITION 8.

ι_1, ι_2 being the angles made by the front with the optic planes, to find ι_1, ι_2 in terms of v_1, v_2 .

By analytical geometry

$$\begin{aligned}
 \cos \iota_1 &= \cos \omega \cdot \cos \omega_1 + \cos \phi \cdot \cos \phi_1 + \cos \psi \cdot \cos \psi_1 \\
 &= \sqrt{\frac{(v_1^2 - a^2)(v_2^2 - a^2)}{(a^2 - b^2)(a^2 - c^2)}} \cdot \sqrt{\frac{a^2 - b^2}{a^2 - c^2}} \\
 &\quad + \sqrt{\frac{(v_1^2 - c^2)(v_2^2 - c^2)}{(c^2 - a^2)(c^2 - b^2)}} \cdot \sqrt{\frac{c^2 - b^2}{c^2 - a^2}} \\
 &= \frac{\sqrt{\{(v_1^2 - a^2)(v_2^2 - a^2)\}} + \sqrt{\{(v_1^2 - c^2)(v_2^2 - c^2)\}}}{a^2 - c^2}
 \end{aligned}$$

and similarly

$$\begin{aligned}
 \cos \iota_2 &= \cos \omega \cdot \cos \omega_2 + \cos \phi \cdot \cos \phi_2 + \cos \psi \cdot \cos \psi_2 \\
 &= \frac{\sqrt{\{(v_1^2 - a^2)(v_2^2 - a^2)\}} - \sqrt{\{(v_1^2 - c^2)(v_2^2 - c^2)\}}}{a^2 - c^2}.
 \end{aligned}$$

PROPOSITION 9.

To find v_1, v_2 in terms of ι_1, ι_2 .

By the last proposition

$$\begin{aligned}
 \cos \iota_1 \cdot \cos \iota_2 &= \frac{(v_1^2 - a^2)(v_2^2 - a^2) - (v_1^2 - c^2)(v_2^2 - c^2)}{(a^2 - c^2)^2} \\
 &= \frac{(a^4 - c^4) - (a^2 - c^2)(v_1^2 + v_2^2)}{(a^2 - c^2)^2} \\
 &= \frac{(a^2 + c^2) - (v_1^2 + v_2^2)}{a^2 - c^2}
 \end{aligned}$$

therefore

$$v_1^2 + v_2^2 = a^2 + c^2 - (a^2 - c^2) \cos \iota_1 \cos \iota_2.$$

Again,

$$\begin{aligned}
 (\sin \iota_1)^2 \cdot (\sin \iota_2)^2 &= 1 - (\cos \iota_1)^2 - (\cos \iota_2)^2 + (\cos \iota_1)^2 (\cos \iota_2)^2 \\
 &= 1 - 2 \cdot \frac{(v_1^2 - a^2)(v_2^2 - a^2) + (v_1^2 - c^2)(v_2^2 - c^2)}{(a^2 - c^2)^2} \\
 &\quad + \frac{(a^2 + c^2)^2 - 2(a^2 + c^2)(v_1^2 + v_2^2) + (v_1^2 + v_2^2)^2}{(a^2 - c^2)^2} \\
 &= \frac{v_1^4 - 2v_1^2 v_2^2 + v_2^4}{(a^2 - c^2)^2}
 \end{aligned}$$

therefore

$$v_1^2 - v_2^2 = (a^2 - c^2) \sin \iota_1 \cdot \sin \iota_2$$

but

$$v_1^2 + v_2^2 = (a^2 + c^2) - (a^2 - c^2) \cos \iota_1 \cos \iota_2$$

therefore

$$\begin{aligned}
 v_1^2 &= \frac{a^2 + c^2}{2} - \frac{a^2 - c^2}{2} \cos(\iota_1 + \iota_2) \\
 &= a^2 \left(\sin \frac{\iota_1 + \iota_2}{2} \right)^2 + c^2 \left(\cos \frac{\iota_1 + \iota_2}{2} \right)^2
 \end{aligned}$$

$$\begin{aligned}
 v_2^2 &= \frac{a^2 + c^2}{2} - \frac{a^2 - c^2}{2} \cos(\iota_1 - \iota_2) \\
 &= a^2 \left(\sin \frac{\iota_1 - \iota_2}{2} \right)^2 + c^2 \left(\cos \frac{\iota_1 - \iota_2}{2} \right)^2.
 \end{aligned}$$

Thus for uniaxal crystals where $\iota_1 + \iota_2 = 180^\circ$

$$v_1^2 = a^2$$

$$v_2^2 = a^2 (\cos \iota)^2 + c^2 (\sin \iota)^2.$$

COR. Hence we may reduce the discovery of the two fronts into which a plane front is refracted on entering a crystal to the following trigonometrical problem.




Fig. 1.

Let a sphere be described about any point in the line in which the air front intersects the plane of incidence. Let the great circle PI denote the latter plane, IF the former, OA, OC also great circles, the planes of single velocity. Suppose IGH to be one of the refracted fronts intersecting OA, OC in G and H , then

$$\frac{(a^2 + c^2) - (a^2 - c^2) \cos(G + H)}{2(\text{vel. in air})^2} = \frac{(\sin PIF)^2}{(\sin PIGH)^2}.$$

The double sign will give rise to two positions of the refracted front IGH .

The propositions which follow are perhaps more curious than immediately useful.

p. 10

PROPOSITION 10.

To determine the portion of a line of vibration in terms of the two velocities of its corresponding front.

We have here to determine the quantities $\frac{y_1}{x_1}, \frac{z_1}{x_1}$ (of Prop. 1) in terms of v_1, v_2 , or on putting $x_1^2 + y_1^2 + z_1^2 = 1$, x_1, y_1, z_1 are to be found in terms of v_1, v_2 .

By Prop. 3

$$x_1 : y_1 : z_1 :: \frac{l}{a^2 - v_1^2} : \frac{m}{b^2 - v_1^2} : \frac{n}{c^2 - v_1^2}$$

and by Prop. 5

$$\begin{aligned} l^2 : m^2 : n^2 &:: (b^2 - c^2)(a^2 - v_1^2)(a^2 - v_2^2) \\ &: (c^2 - a^2)(b^2 - v_1^2)(b^2 - v_2^2) \\ &: (a^2 - b^2)(c^2 - v_1^2)(c^2 - v_2^2); \end{aligned}$$

therefore

$$\begin{aligned} x_1^2 : y_1^2 : z_1^2 \\ &:: (b^2 - c^2) \frac{a^2 - v_2^2}{a^2 - v_1^2} : (c^2 - a^2) \frac{b^2 - v_2^2}{b^2 - v_1^2} : (a^2 - b^2) \frac{c^2 - v_2^2}{c^2 - v_1^2}. \end{aligned}$$

Let α, β, γ be the angles made by the given line of vibration with the elastic axes, then

$$\begin{aligned} (\cos \alpha)^2 &= \frac{x_1^2}{x_1^2 + y_1^2 + z_1^2} \\ &= (b^2 - c^2)(a^2 - v_2^2)(b^2 - v_1^2)(c^2 - v_1^2) \end{aligned}$$

divided by

$$\begin{aligned} (b^2 - c^2)(a^2 - v_2^2)(b^2 - v_1^2)(c^2 - v_1^2) &+ (c^2 - a^2)(b^2 - v_2^2)(c^2 - v_1^2)(a^2 - v_1^2) \\ &+ (a^2 - b^2)(c^2 - v_2^2)(a^2 - v_1^2)(b^2 - v_1^2) \end{aligned}$$

and therefore

$$= \frac{(b^2 - c^2)(a^2 - v_2^2)(b^2 - v_1^2)(c^2 - v_1^2)}{(v_1^2 - v_2^2)(a^2 - b^2)(b^2 - c^2)(c^2 - a^2)}$$

(where it is to be observed that the reduction of the denominator is simply the effect of a vast heap of terms disappearing under the influence of contact with the magic circuit $(a^2 - b^2), (b^2 - c^2), (c^2 - a^2)$, a simpler instance of which was seen in Proposition 5).

In fact the coefficient of $v^4 \cdot v^2$

$$\begin{aligned} &= (b^2 - c^2) + (c^2 - a^2) + (a^2 - b^2) \\ &= 0 \end{aligned}$$

p. 11

that of $v_1^2 \cdot v_2^2$

$$\begin{aligned} &= (c^2 + b^2)(c^2 - b^2) \\ &\quad + (a^2 + c^2)(a^2 - c^2) \\ &\quad + (b^2 + a^2)(b^2 - a^2) \\ &= (c^4 - b^4) + (a^4 - c^4) + (b^4 - a^4) \\ &= 0. \end{aligned}$$

The term in which neither v_1 nor v_2 enters

$$\begin{aligned} &= a^2 b^2 c^2 \{ (b^2 - c^2) + (c^2 - a^2) + (a^2 - b^2) \} \\ &= 0. \end{aligned}$$

The coefficient of

$$-v_1^2 = a^2 \cdot (b^4 - c^4) + b^2 \cdot (c^4 - a^4) + c^2 \cdot (a^4 - b^4)$$

and that of

$$v_2^2 = b^2 c^2 \cdot (c^2 - b^2) + c^2 a^2 \cdot (a^2 - c^2) + a^2 b^2 \cdot (b^2 - a^2)$$

each of which

$$= (a^2 - b^2) \cdot (b^2 - c^2) \cdot (c^2 - a^2).$$

Hence

$$(\cos \alpha)^2 = \frac{v_1^2 - b^2}{v_1^2 - v_2^2} \cdot \frac{(a^2 - v_2^2)(c^2 - v_1^2)}{(a^2 - b^2)(a^2 - c^2)},$$

in like manner $(\cos \beta)^2 = \&c.$ and

$$(\cos \gamma)^2 = \frac{v_1^2 - b^2}{v_1^2 - v_2^2} \cdot \frac{(c^2 - v_2^2)(a^2 - v_1^2)}{(c^2 - b^2)(c^2 - a^2)}.$$

PROPOSITION 11.

$\varepsilon_1, \varepsilon_2$ being the angles between any line of vibration and the optic axes, required the velocity due to that line in terms of $\varepsilon_1, \varepsilon_2$.

By analytical geometry,

$$\cos \varepsilon_1 = \cos \alpha \cdot \cos \phi_1 + \cos \gamma \cdot \cos \psi_1$$

$$\cos \varepsilon_2 = \cos \alpha \cdot \cos \phi_1 - \cos \gamma \cdot \cos \psi_1$$

therefore

$$\begin{aligned} \cos \varepsilon_1 \cdot \cos \varepsilon_2 &= (\cos \alpha)^2 (\cos \phi_1)^2 - (\cos \gamma)^2 (\cos \psi_1)^2 \\ &= \frac{v_1^2 - b^2}{v_1^2 - v_2^2} \cdot \left\{ \frac{(a^2 - v_2^2)(c^2 - v_1^2) - (c^2 - v_2^2)(a^2 - v_1^2)}{(a^2 - c^2)^2} \right\} \\ &= \frac{v_1^2 - b^2}{v_1^2 - v_2^2} \cdot \frac{(a^2 - c^2)(v_2^2 - v_1^2)}{(a^2 - c^2)^2} \\ &= \frac{b^2 - v_1^2}{a^2 - c^2}. \end{aligned}$$

Hence

$$v_1^2 = b^2 - (a^2 - c^2) \cos \varepsilon_1 \cos \varepsilon_2$$

and in like manner, for the conjugate line of vibration

$$v_2^2 = b^2 - (a^2 - c^2) \cos \varepsilon'_1 \cos \varepsilon'_2.$$

p. 12

PROPOSITION 12.

To find $\varepsilon_1, \varepsilon_2$ in terms of ι_1, ι_2 .

$$\begin{aligned} (\cos \varepsilon_1)^2 + (\cos \varepsilon_2)^2 &= 2(\cos \alpha)^2 \cdot (\cos \phi_1)^2 + 2(\cos \gamma)^2 \cdot (\cos \psi_1)^2 \\ &= 2 \frac{v_1^2 - b^2}{v_1^2 - v_2^2} \left\{ \frac{(a^2 - v_2^2)(c^2 - v_1^2) + (c^2 - v_2^2)(a^2 - v_1^2)}{(a^2 - c^2)^2} \right\} \end{aligned}$$

but by Prop. 9

$$\begin{aligned} v_1^2 &= a^2 \left(\sin \frac{\iota_1 + \iota_2}{2} \right)^2 + c^2 \left(\cos \frac{\iota_1 + \iota_2}{2} \right)^2 \\ v_2^2 &= a^2 \left(\sin \frac{\iota_1 - \iota_2}{2} \right)^2 + c^2 \left(\cos \frac{\iota_1 - \iota_2}{2} \right)^2 \end{aligned}$$

therefore

$$\begin{aligned}
(\cos \varepsilon_1)^2 + (\cos \varepsilon_2)^2 &= \frac{b^2 - v_1^2}{(a^2 - c^2) \sin \iota_1 \cdot \sin \iota_2} \quad \text{multiplied by} \\
&\quad \frac{2(a^2 - c^2)^2 \left[\left(\cos \frac{\iota_1 + \iota_2}{2} \right)^2 \left(\sin \frac{\iota_1 - \iota_2}{2} \right)^2 + \left(\cos \frac{\iota_1 - \iota_2}{2} \right)^2 \left(\sin \frac{\iota_1 + \iota_2}{2} \right)^2 \right]}{(a^2 - c^2)^2} \\
&= \frac{b^2 - v_1^2}{(a^2 - c^2) \sin \iota_1 \cdot \sin \iota_2} \{ (\sin \iota_1)^2 + (\sin \iota_2)^2 \}
\end{aligned}$$

and we have seen that

$$\cos \varepsilon_1 \cos \varepsilon_2 = \frac{b^2 - v_1^2}{a^2 - c^2}$$

therefore

$$\begin{aligned}
\cos \varepsilon_1 + \cos \varepsilon_2 &= \sqrt{\frac{b^2 - v_1^2}{a^2 - c^2}} \cdot \frac{\sin \iota_1 + \sin \iota_2}{\sqrt{\sin \iota_1 \cdot \sin \iota_2}}, \\
\cos \varepsilon_1 - \cos \varepsilon_2 &= \sqrt{\frac{b^2 - v_1^2}{a^2 - c^2}} \cdot \frac{\sin \iota_1 - \sin \iota_2}{\sqrt{\sin \iota_1 \cdot \sin \iota_2}},
\end{aligned}$$

therefore

$$\begin{aligned}
\cos \varepsilon_1 &= \sqrt{\left\{ \frac{b^2 - v_1^2}{a^2 - c^2} \cdot \frac{\sin \iota_1}{\sin \iota_2} \right\}} \\
\cos \varepsilon_2 &= \sqrt{\left\{ \frac{b^2 - v_1^2}{a^2 - c^2} \cdot \frac{\sin \iota_2}{\sin \iota_1} \right\}}
\end{aligned}$$

and in like manner

$$\begin{aligned}
\cos \varepsilon'_1 &= \sqrt{\left\{ \frac{b^2 - v_2^2}{a^2 - c^2} \cdot \frac{\sin \iota_1}{\sin \iota_2} \right\}} \\
\cos \varepsilon'_2 &= \sqrt{\left\{ \frac{b^2 - v_2^2}{a^2 - c^2} \cdot \frac{\sin \iota_2}{\sin \iota_1} \right\}}
\end{aligned}$$

where v_1, v_2 for the sake of neatness are left unexpressed in terms of ι_1, ι_2 . p. 13

This is the simplest form by which the position of the lines of vibration can be denoted.

COR. From the last proposition it appears that

$$\frac{\cos \varepsilon_1}{\cos \varepsilon_2} = \frac{\sin \iota_1}{\sin \iota_2}.$$

Hence we may construct geometrically for the two planes of polarization.




Fig. 2.

Let I, K be the projections of the two optic axes on a sphere, E the projection of the normal to the front, P the projection of one line of vibration; then

$$\frac{\cos PK}{\cos PI} = \frac{\sin KE}{\sin IE}.$$

Draw FEG the circle of which P is the pole, meeting PK, PI produced in G and F .

Then

$$\cos PK = \sin KG,$$

and

$$\cos PI = \sin IF,$$

therefore

$$\frac{\sin KG}{\sin FI} = \frac{\sin KE}{\sin IE}$$

therefore

$$\frac{\sin KG}{\sin KE} = \frac{\sin IF}{\sin IE}$$

therefore

$$\sin KEG = \sin IEF$$

therefore $KEG = IEF$ or $180^\circ - IEF$. But $PEF = PEG$, therefore EP bisects either the angle IEK or the supplement to it.

These two positions of EP give the two planes of polarization. The construction is the same as that given in Mr Airy's tracts, and originally proposed, I believe, by Mr MacCullagh.

p. 14

ADDENDUM.

If in the equation of Prop. 6, viz.

$$\frac{(\cos \omega)^2}{a^2 - v^2} + \frac{(\cos \phi)^2}{b^2 - v^2} + \frac{(\cos \psi)^2}{c^2 - v^2} = 0$$

we change a, b, c, v into $\frac{1}{a}, \frac{1}{b}, \frac{1}{c}, \frac{1}{v}$, and consider v to be the length of a line drawn perpendicular to the plane

$$\cos \omega \cdot x + \cos \phi \cdot y + \cos \psi \cdot z = 0,$$

the equation to the extremity thereof must be

$$\frac{a^2 r^2 (\cos \omega)^2}{a^2 - r^2} + \frac{b^2 r^2 (\cos \phi)^2}{b^2 - r^2} + \frac{c^2 r^2 (\cos \psi)^2}{c^2 - r^2} = 0,$$

where ω, ϕ, ψ denote the angles between the radius vector r , and the axes of x, y, z , so that the equation may be written

$$\frac{a^2 x^2}{a^2 - r^2} + \frac{b^2 y^2}{b^2 - r^2} + \frac{c^2 z^2}{c^2 - r^2} = 0,$$

which is that of the wave surface.

But we have seen that

$$v^2 = c^2 \left\{ \cos \left(\frac{\iota_1 \pm \iota_2}{2} \right) \right\}^2 + a^2 \left\{ \sin \left(\frac{\iota_1 \pm \iota_2}{2} \right) \right\}^2,$$

therefore the equation to the wave surface may be written

$$\frac{1}{r^2} = \frac{(\cos \frac{\iota_1 \pm \iota_2}{2})^2}{c^2} + \frac{(\sin \frac{\iota_1 \pm \iota_2}{2})^2}{a^2},$$

where ι_1, ι_2 denote the angles between the radius vector v and the two lines which would be the optic axes if a, b, c were changed into $\frac{1}{a}, \frac{1}{b}, \frac{1}{c}$ so that if e be the inclination of either to the mean axis of elasticity

$$\begin{aligned} \cos e &= \sqrt{\frac{\frac{1}{a^2} - \frac{1}{b^2}}{\frac{1}{a^2} - \frac{1}{c^2}}} = \frac{c}{b} \sqrt{\frac{a^2 - b^2}{a^2 - c^2}}, \\ \sin e &= \sqrt{\frac{\frac{1}{b^2} - \frac{1}{c^2}}{\frac{1}{a^2} - \frac{1}{c^2}}} = \frac{a}{b} \sqrt{\frac{b^2 - c^2}{a^2 - c^2}}. \end{aligned}$$

These lines I shall call by way of distinction the prime radii³.

p. 15

Cor. 1. If r_1, r_2 be the two values of r corresponding to the same values of ι_1, ι_2 we have

$$\begin{aligned} \frac{1}{r_1^2} - \frac{1}{r_2^2} &= \frac{1}{c^2} \left\{ \left(\cos \frac{\iota_1 - \iota_2}{2} \right)^2 - \left(\cos \frac{\iota_1 + \iota_2}{2} \right)^2 \right\} \\ &\quad + \frac{1}{a^2} \left\{ \left(\sin \frac{\iota_1 - \iota_2}{2} \right)^2 - \left(\sin \frac{\iota_1 + \iota_2}{2} \right)^2 \right\} \\ &= \left(\frac{1}{c^2} - \frac{1}{a^2} \right) \sin \iota_1 \cdot \sin \iota_2, \end{aligned}$$

³Upon the authority of Professor Airy I have appropriated the term optic axes to the lines normal to the fronts of single velocity.

which proves the celebrated problem of *two rays* having a common direction in a crystal.

Cor. 2. The intersection of any concentric sphere with the wave surface is found by making r constant. Hence $\iota_1 \pm \iota_2$ becomes constant, and therefore $r\iota_1 \pm r\iota_2 = \text{constant}$. Hence the curve of intersection is the locus of points, the sum or difference of whose distances from two poles when measured by the arcs of great circles is constant; the poles being the points in which the prime radii pierce the sphere.

In three cases these spherico-ellipses or spherico-hyperbolas become great circles:

(1) When $\iota_1 \pm \iota_2 = \text{the angle between the two poles}$, in which case the curve of intersection is the great circle which comprises the two poles.

(2) When $\iota_1 - \iota_2 = 0$, when the locus is a great circle perpendicular to the former and bisecting the angle between the optic axes.

(3) When $\iota_1 + \iota_2 = 180^\circ$, when the locus is a great circle perpendicular to the two above, and bisecting the supplemental angle between the two axes.

Various other properties may be with the greatest simplicity deduced from the radio-angular equation. The hurry of the press leaves me time only to subjoin the following

PROPOSITION.

To find the inclination of the radius vector to the tangent plane, in terms of the angles which the radius vector makes with the prime radii.

Let O be the centre of the wave surface, OA, OB the two prime radii, OP any radius vector. Let $OP = v$, $POA = \iota_1$, $POB = \iota_2$, and let the inclination of the planes $POA, POB = \mu$; then

$$\frac{1}{r^2} = \frac{(\sin \frac{\iota_1 + \iota_2}{2})^2}{a^2} + \frac{(\cos \frac{\iota_1 + \iota_2}{2})^2}{c^2},$$

(taking only the positive sign for the sake of brevity).

p. 16

Let OQ, OR be the two adjacent radii vectores, so assumed that

$$QOA = POA, \quad QOB = POB + \delta\iota_2,$$

$$ROB = POB, \quad ROA = POA + \delta\iota_1,$$

and let p, q, r, a, b be the projections of P, Q, R, A, B on a sphere of which O is the centre, then it is clear that

$$qpa = 90^\circ, \quad rpb = 90^\circ;$$

draw qm perpendicular to pb , then $pm = \delta\iota_2$, and therefore

$$pq = \frac{pm}{\sin pqm} = \frac{pm}{\sin apb} = \frac{\delta\iota_2}{\sin \mu}.$$

In like manner

$$pr = \frac{\delta \iota_1}{\sin \mu}.$$

Now the angle QPO




Fig. 3.

$$= \tan^{-1} \frac{r \cdot POQ}{OQ - OP} = \tan^{-1} \frac{r \cdot pq}{\frac{dr}{d\iota_2} \delta \iota_2};$$

also

$$\begin{aligned} \frac{d(1/r^2)}{d\iota_2} &= \frac{d}{d\iota_2} \left\{ \left(\frac{1}{c^2} - \frac{1}{a^2} \right) \left(\cos \frac{\iota_1 + \iota_2}{2} \right)^2 \right\} \\ &= - \left(\frac{1}{c^2} - \frac{1}{a^2} \right) \left(\sin \frac{\iota_1 + \iota_2}{2} \right) \left(\cos \frac{\iota_1 + \iota_2}{2} \right); \end{aligned}$$

therefore

$$\frac{dr}{r d\iota_2} = \frac{1}{4} r^2 \left(\frac{1}{c^2} - \frac{1}{a^2} \right) \sin(\iota_1 + \iota_2),$$

therefore

$$\cot QPO = \frac{r^2}{4} \left(\frac{1}{c^2} - \frac{1}{a^2} \right) \sin(\iota_1 + \iota_2) \sin \mu.$$

In like manner

$$\cot RPO = \frac{r^2}{4} \left(\frac{1}{c^2} - \frac{1}{a^2} \right) \sin(\iota_1 + \iota_2) \sin \mu,$$

therefore

$$QPO = RPO.$$

Also it is clear that $rpq = apb = \mu$. And to find the inclination of OP to RPQ , we have only to describe a sphere of which P is the centre, and intersecting PQ, PR, PO in Q', R', O' .

Then $R'O'Q' = \mu$, and

$$O'Q' = O'R' = \cot^{-1} \left\{ \frac{r^2}{4} \left(\frac{1}{c^2} - \frac{1}{a^2} \right) \sin(\iota_1 + \iota_2) \sin \mu \right\}.$$




Fig. 4.

p. 17

Draw $O'N$ perpendicular to $R'Q'$, then $O'N$ measures the inclination of the radius vector to the tangent plane⁴.

And

$$Q'O'N = \frac{\mu}{2},$$

therefore

$$\cos \frac{\mu}{2} = \tan O'N \cdot \cot O'Q',$$

therefore

$$\cot O'N = \frac{\cot O'Q'}{\cos \frac{\mu}{2}},$$

and therefore

$$\cot O'N = \frac{1}{2}r^2 \left(\frac{1}{a^2} - \frac{1}{c^2} \right) \sin \frac{\mu}{2} \cdot \sin(\iota_1 + \iota_2).$$

Let AOB the angle between the optic axes $= 2e$, then by mere trigonometry

$$\sin \frac{\mu}{2} = \sqrt{\frac{\sin(e + \frac{\iota_1 - \iota_2}{2}) \sin(e - \frac{\iota_1 - \iota_2}{2})}{\sin \iota_1 \cdot \sin \iota_2}},$$

therefore the tangent of the inclination between the radius vector and the normal

$$= \frac{1}{2}r^2 \left(\frac{1}{a^2} - \frac{1}{c^2} \right) \sin(\iota_1 + \iota_2) \sqrt{\frac{\sin(e + \frac{\iota_1 - \iota_2}{2}) \sin(e - \frac{\iota_1 - \iota_2}{2})}{\sin \iota_1 \cdot \sin \iota_2}}.$$

In like manner the tangent of the inclination between the same radius vector and the normal at the other point of the wave-surface pierced by it

$$= \frac{1}{2}(r_1)^2 \left(\frac{1}{a^2} - \frac{1}{c^2} \right) \sin(\iota_1 - \iota_2) \sqrt{\frac{\sin(e + \frac{\iota_1 + \iota_2}{2}) \sin(e - \frac{\iota_1 + \iota_2}{2})}{\sin \iota_1 \cdot \sin \iota_2}}.$$

⁴ O' is the projection of the ray and $R'O'$ of the tangent plane. Therefore $O'N$ being perpendicular to $R'Q'$ represents their inclination.

We may, in the same way, find the inclination of the tangent plane to either of the prime radii, and to the plane which contains them both, in terms of ι_1 and ι_2 ; the former by a remarkably elegant construction; but the final expressions do not present themselves under the same simple aspect.

If we call ϕ the angle between the ray and the front, we may still further reduce by substituting for r^2 its values in terms of ι_1, ι_2 and we shall obtain

$$\cot \phi = \frac{2(c^2 - a^2)}{c^2 \tan \frac{\iota_1 + \iota_2}{2} + a^2 \cot \frac{\iota_1 + \iota_2}{2}} \\ \times \sqrt{\left\{ \sin \left(e + \frac{\iota_1 + \iota_2}{2} \right) \sin \left(e - \frac{\iota_1 + \iota_2}{2} \right) \cdot \operatorname{cosec} \iota_1 \cdot \operatorname{cosec} \iota_2 \right\}}.$$

And if π_1, π_2 be the inclinations of the normal to the two prime radii, it may be shown that p. 18

$$\cos \pi_1 = \cos \phi \sin \iota_1 \mp \sin \phi \cos \iota_1 \sin \frac{\mu}{2}, \\ \cos \pi_2 = \cos \phi \sin \iota_2 \pm \sin \phi \cos \iota_2 \sin \frac{\mu}{2}.$$

Cor. 1. For uniaxal crystals $\frac{\mu}{2} = 90^\circ$ and $\iota_1 + \iota_2 = 180^\circ$, so that the tangent of the inclination of normal to radius vector

$$= \frac{1}{2}r^2 \cdot \left(\frac{1}{a^2} - \frac{1}{c^2} \right) \sin 2\iota \quad \text{for one point,}$$

and

$$= 0 \quad \text{for the other.}$$

Cor. 2. For every point in the circular section which passes through the poles $\sin \frac{\mu}{2} = 0$, and for the other two circular sections $\iota_1 \pm \iota_2 = 0$ or 180° .

Therefore every point in the three circular sections is an apse.

Cor. 3. When a nearly $= c$, $\frac{1}{a^2} - \frac{1}{c^2}$ is very small; and therefore the normal and radius vector very nearly coincide.

Cor. 4. Referring to fig. 4 we see that $O'N$ bisects the angle $R'O'Q'$. Now $R'O, Q'O$ are respectively perpendicular to the planes passing through O' and the optic axes; and therefore the meridian plane as we may term it, that is, the plane containing both the ray and the normal, always bisects the angle formed by the two planes drawn through the ray and the two optic axes.

Cor. 5. When

$$\iota_1 \text{ or } \iota_2 = 0,$$

$$\iota_2 \text{ or } \iota_1 = e.$$

And therefore ϕ assumes the form $\frac{0}{0}$, which indicates that the extremities of the four prime radii are singular points.

In concluding for the present it behoves me to state that one step has been omitted in the foregoing paper⁵, viz. the actual performance of the eliminations which lead to the rectilinear equation to the wave-surface. But Mr Archibald Smith's elegant and brief Memoir in the *Cambridge Philosophical Transactions*⁶ of last year leaves nothing to be desired further on that head. p. 19

That I have not exhibited it in its proper place (Prop. 6) arises only from my respect to the principle of literary propriety. With this important blank supplied the Analytical Theory may be pronounced to be complete.

For all errors and imperfections in what precedes my excuse must be press of time and a total want of the materials to be derived from consulting works of reference.

Since writing the above I have had an opportunity of reading the paper of our living Laplace inserted as part of the Third Supplement to his System of Rays in the *Transactions* of the Royal Irish Academy, in which the principal foregoing results are obtained by aid of a more refined and transcendental analysis.

The nature of the four singular points is there discussed and the existence of four circles of plane contact demonstrated.

The former may be very easily shown thus: when ι_1 is very small $\iota_2 = 2e - \iota_1 \cos \psi$ very nearly, ψ denoting the inclination of the plane in which e is reckoned to the plane in which ι_2 is reckoned.

Hence

$$\begin{aligned} \left(\frac{1}{r}\right)^2 &= \frac{1}{2} \left(\frac{1}{a^2} + \frac{1}{c^2}\right) - \frac{1}{2} \left(\frac{1}{a^2} - \frac{1}{c^2}\right) \cos\{2e - \iota_1(\cos \psi \pm 1)\} \\ &= \frac{1}{2} \left(\frac{1}{a^2} + \frac{1}{c^2}\right) - \frac{1}{2} \left(\frac{1}{a^2} - \frac{1}{c^2}\right) \cos 2e - \frac{1}{2} \left(\frac{1}{a^2} - \frac{1}{c^2}\right) \sin 2e (\cos \psi \pm 1) \iota_1 \\ &= \frac{1}{b^2} - \frac{1}{b^2 ac} \sqrt{\{(a^2 - b^2)(b^2 - c^2)\}} (\cos \psi \pm 1) \iota_1, \end{aligned}$$

therefore

$$r = b \left\{ 1 + \frac{1}{2} (\cos \psi \pm 1) \left(1 - \frac{b^2}{a^2}\right)^{1/2} \left(\frac{b^2}{c^2} - 1\right)^{1/2} \iota_1 \right\}.$$

Take ψ constant and let the abscissae and ordinates be reckoned respectively along and perpendicular to the prime ray.

Then

$$\iota_1 = \frac{y}{x} \quad \text{nearly, and} \quad r = \sqrt{y^2 + x^2} = x,$$

⁵See below, p. 27. Ed.

⁶Vol. vi. Also *Phil. Mag.* April, 1838, p. 335. Ed.

or, if we change the origin to the other extremity of the prime ray,

$$\iota_1 = \frac{y}{b}, \quad r = b - x,$$

so that the equation becomes

$$-\frac{x}{y} = \frac{1}{2}(\cos \psi \pm 1) \sqrt{\left\{ \left(1 - \frac{b^2}{a^2}\right) \left(\frac{b^2}{c^2} - 1\right) \right\}}.$$

Hence at each singular point the surface is touched by a cone, the equation to the generating line of which is given by the above, the extreme angle between it and the prime ray being

p. 20

$$\cot^{-1} \left[\sqrt{\left\{ \left(1 - \frac{b^2}{a^2}\right) \left(\frac{b^2}{c^2} - 1\right) \right\}} \right].$$

When $b = a$, ψ always $= \frac{\pi}{2}$ and the cone returns into a plane.

Again, let us suppose that the position of any perpendicular from the centre is given, and that of the corresponding radius vector required.

Let OA, OB^7 denote what we have termed the optic axes, but which it will be more agreeable to analogy to term the prime perpendiculars from centre, and let OP be the given normal. Take OQ, OR contiguous perpendiculars from centre in planes POQ, ROP , perpendicular to POA, POB respectively, then the inclination of the two former will be the same as that of the two latter, and may be termed μ .

Let ι_1, ι_2 now denote the angles POA, POB respectively, then

$$QOA = \iota_1, \quad QOB = \iota_2 + \delta\iota_2,$$

$$ROA = \iota_1 + \delta\iota_1, \quad ROB = \iota_2.$$

The ray will be found by joining O with the intersection of three planes drawn at P, Q, R , perpendicular to OP, OQ, OR , respectively.

Now from Prop. 9 it appears that

$$OP = \sqrt{\left\{ a^2 \left(\sin \frac{\iota_1 + \iota_2}{2} \right)^2 + c^2 \left(\cos \frac{\iota_1 + \iota_2}{2} \right)^2 \right\}},$$

using only one sign for the sake of simplicity, which we may do by throwing the ambiguity upon the *way* in which ι_1 or ι_2 is measured, also

$$OQ = OP + \frac{d \cdot OP}{d\iota_2} \delta\iota_2,$$




Fig. 5.

$$OR = OP + \frac{d \cdot OP}{du_1} \delta \iota_1.$$

Let $\delta \iota_1 = \delta \iota_2$, then it is clear that $OQ = OR$, and the intersection of the two planes perpendicular to OQ, OR is therefore a line perpendicular to the plane QOR , and to the line which bisects the angle QOR .

In fact if we draw QT, RT perpendicular to OQ, OR respectively in the plane QOR , the intersection in question passes through T and is perpendicular to OT ; also

$$OT = OQ \cdot \sec \left(\frac{1}{2} ROQ \right) = OQ$$

to the first order of smallness.

p. 21

Now it is easy to see (just as on p. 16) that

$$ROP = \frac{\delta \iota_1}{\sin \mu},$$

and also

$$QOP = \frac{\delta \iota_2}{\sin \mu},$$

therefore $ROP = QOP$ and therefore POT is perpendicular to QOR .

Hence the problem is reduced to finding L the intersection of two lines TL, PL drawn in the same plane POT .

Now because OTL, OPL are each right angles, a circle may be made to pass through L, T, P, O .

Hence the angle

$$\begin{aligned} PLO = PTO &= \tan^{-1} \frac{OP \times POT}{OT - OP} \\ &= \tan^{-1} \frac{OP \times POR \cdot \cos \frac{1}{2} \mu}{\frac{d \cdot OP}{du_2} \delta \iota_2} = \tan^{-1} \frac{OP \times \frac{\delta \iota_2}{\sin \mu} \cos \frac{1}{2} \mu}{\frac{d \cdot OP}{du_2} \delta \iota_2}, \end{aligned}$$

⁷ OA, OB are not expressed in the figure.

and

$$OL = OP \cdot \sec POL.$$

Also the position of the plane POL is known, and therefore the radius is completely determined in magnitude and position.

It may be worth while also to remark that the above constructions enable us to form a series of equations between the magnitude of the radius and its inclinations to the two prime perpendiculars.

In fact, if we call π_1, π_2 the two inclinations in question

$$\cos \pi_1 = \cos POL \cos \iota_1 \pm \sin POL \sin \iota_1 \cdot \sin \frac{\mu}{2},$$

$$\cos \pi_2 = \cos POL \cos \iota_2 \mp \sin POL \sin \iota_2 \cdot \sin \frac{\mu}{2},$$

and of course if we call the angle between the two prime normals $2E$

$$\sin \frac{\mu}{2} = \sqrt{\frac{\sin(E + \frac{\iota_1 + \iota_2}{2}) \sin(E - \frac{\iota_1 + \iota_2}{2})}{\sin \iota_1 \sin \iota_2}}.$$

Cor. 1. When ι_1 or $\iota_2 = 0$, $\tan POL$ assumes the form $\frac{0}{0}$ which may be interpreted analogously to the method used in the reverse problem, but may be more elegantly illustrated by

Cor. 2. Which is that the meridian plane POT (that is, the plane in which both normal and radius lie) bisects the angle formed by ROP, QOP , and therefore p. 22

that formed by the planes drawn through the normal and the two prime normals to which these two are perpendicular.

Now we have found (Cor. 4, page 18), that it also bisects the angle formed by the two planes passing through the radius and the two prime radii. Hence when the ray is given, we may find by the easiest geometry the normal and the tangent plane, and *vice versâ*.

Thus suppose $(N, N')(R, R')$ to be the projections of the prime perpendiculars and prime radii on a sphere concentric with the wave surface.

Let n be the projection of any given perpendicular on the same sphere; join nN, nN' ; bisect NnN' by nM , which will be the meridian plane.

Draw from R', R' perpendicular to nM and make $R'T = TV$. Produce RV to meet Mn in r , then $RrM = R'rM$, and therefore r is the projection of the radius. Just in the same way when r is given we may find n .

Now suppose n to come to N , then the position of the meridian plane nM becomes indeterminate, and r from a point becomes a locus, subject to the condition that $R'rN = RrN$. From r draw rD perpendicular to rN .

Then it is clear that because rN bisects RrR'

$$\frac{\sin RD}{\sin R'D} = \frac{\sin Rr}{\sin R'r} = \frac{\sin RN}{\sin R'N'},$$




Fig. 6.




Fig. 7.

and therefore D is a fixed point and ND a fixed length, and

$$\cos rND = \tan rN \cdot \cot ND;$$

therefore the projection of the locus of r upon a plane drawn at N perpendicular to the line joining N with the centre O is given by the equation

$$\rho = ON \cdot \cot ND \cdot \cos \theta,$$

N being the origin and the projection of ND the prime radius; which is the equation to a circle passing through N , and whose diameter $= ON \cot ND$.

Hence at the extremity of each prime perpendicular the tangent plane meets the surface in a circle passing through that extremity and whose radius $= \frac{1}{2}b \cot a$, a being to be found from the equation

$$\frac{\sin(2E + a)}{\sin a} = \frac{\sin(E + e)}{\sin(E - e)},$$

that is

$$\tan(E + a) = (\tan E)^2 \cot e.$$

Just in the same way it may be shown that the trace of the perpendiculars to the tangent planes of the surface at the point where it is pierced by any prime radius upon a plane perpendicular to that radius at its extremity, is also a circle passing through it, and curved in an opposite direction from the circle of plane contact nearest to it.

Hence the enveloping cone at these points may be described as being perpendicular to the circular cone, formed by drawing lines from the centre to the above described circle; that is every generating line of the one will be perpendicular to the generating line which it meets of the other.

More generally it easily appears from fig. 6 that if a series of great circles (representing meridian planes) be taken intersecting the great circle $NRR'N'$ in a fixed point, a plane perpendicular to the radius passing through that point, will intersect the cone of rays as well as the cone of perpendiculars corresponding to those meridian planes, in two *circles*. So that there exist an indefinite number of *circular* cones of rays corresponding to *circular* cones of perpendiculars touching each other in a line lying in the plane containing the extreme axes, and having their circular sections perpendicular to that line.

The cusps are explained by the cone of *rays* degenerating into a right line, and the circles of plane contact by the cone of perpendiculars so degenerating.

Furthermore I observe in conclusion that when a ray is given it follows from the general geometrical construction above that there will be two meridian planes according as we take R with R' , or with a point 180° from R' , and consequently these two planes will be perpendicular to each other.

And similarly when a *normal* is given there will be two meridian planes perpendicular to each other.

Thus the planes passing through any radius and the two normals at the points where it pierces the wave surface, are *perpendicular* to each other, as are also the two planes passing through any normal and *its* two corresponding radii.

Moreover a glance at fig. 2 will show that the two lines of vibration corresponding to any front lie respectively in the two meridian planes passing through the perpendicular to that front or, in other words, the intersection of a plane drawn through either ray belonging to a front perpendicular thereunto is *always* a line of vibration in that front.

This has been noticed, I think, by Sir William Hamilton for the particular case of the singular points.

As two fronts belong to every ray, so two rays pertain to every front. And from what has been said above it appears that the two lines of vibration in any front are the projections of its two rays upon its own plane.

p. 24

NOTE 1.

In the paper above, it is shown that the meridian plane, that is, the plane

containing the ray and normal, always passes through a line of vibration in the corresponding point. Now the line of force called into action by a displacement in the line of vibration clearly lies in this very plane; for the resolved part of it lies in the line of vibration itself.

Harmony and analogy concur in suggesting that as two of these four lines are perpendicular to each other, so are also the other two, or in other words, that the ray is always perpendicular to the direction of unresolved force.

The following investigation verifies this conjecture.

Let x, y, z be the coordinates of a point taken at distance unity from the origin and in any line of vibration; then the cosines of the angles made by the line of force with the axes are as $a^2x : b^2y : c^2z$ respectively.

Let ω be the inclination between the line of vibration and the line of force, then

$$\cos \omega = \frac{a^2x \cdot x + b^2y \cdot y + c^2z \cdot z}{\sqrt{\{(a^4x^2 + b^4y^2 + c^4z^2)(x^2 + y^2 + z^2)\}}} = \frac{a^2x^2 + b^2y^2 + c^2z^2}{\sqrt{a^4x^2 + b^4y^2 + c^4z^2}}.$$

Let

$$\sqrt{a^4x^2 + b^4y^2 + c^4z^2} = P,$$

then

$$P^2 = v^4(\sec \omega)^2.$$

Now let α, β, γ be the angles of inclination between the coordinate planes and the front in which the line of vibration lies, and λ some quantity to be determined. I have shown in Prop. 3 that if

$$\lambda \cos \alpha = (a^2 - v^2)x,$$

then will

$$\lambda \cos \beta = (b^2 - v^2)y,$$

and

$$\lambda \cos \gamma = (c^2 - v^2)z;$$

therefore

$$\lambda^2 = a^4x^2 + b^4y^2 + c^4z^2 - 2v^2(a^2x^2 + b^2y^2 + c^2z^2) + v^4 = P^2 - v^4.$$

Again,

$$\lambda^2 \left\{ \frac{(\cos \alpha)^2}{(a^2 - v^2)^2} + \frac{(\cos \beta)^2}{(b^2 - v^2)^2} + \frac{(\cos \gamma)^2}{(c^2 - v^2)^2} \right\} = x^2 + y^2 + z^2 = 1;$$

therefore

$$\frac{1}{P^2 - v^4} = \frac{(\cos \alpha)^2}{(a^2 - v^2)^2} + \frac{(\cos \beta)^2}{(b^2 - v^2)^2} + \frac{(\cos \gamma)^2}{(c^2 - v^2)^2}.$$

Now

$$\frac{1}{P^2 - v^4} = \frac{1}{v^4(\sec \omega)^2 - v^4} = \frac{1}{v^4}(\cot \omega)^2.$$

And in Mr Smith's investigation of the form of the wave surface (already alluded to⁸) by great good fortune I find ready to my hand p. 25

$$\frac{(\cos \alpha)^2}{(a^2 - v^2)^2} + \frac{(\cos \beta)^2}{(b^2 - v^2)^2} + \frac{(\cos \gamma)^2}{(c^2 - v^2)^2} = \frac{1}{v^2(r^2 - v^2)},$$

r being the radius vector to the point whose tangent plane is parallel to the point in question.

Hence

$$(\cot \omega)^2 = \frac{v^4}{v^2(r^2 - v^2)} = \frac{v^2}{r^2 - v^2} = \frac{v^2}{r^2 - p^2},$$

p being the length of the perpendicular from the centre upon the tangent plane, for $p = v$.

Hence $(\cot \omega)^2 =$ the square of the cotangent of the angle between radius vector and normal.

Or, in other words, the line of force is as much inclined to the line of vibration as the ray is to the normal.

Now the normal is perpendicular to the line of vibration, and all four lines lie in one plane.

Therefore the ray is perpendicular to the line of force. Q. E. D.

I may be allowed to conclude this long paper with a summary of some of the most remarkable consequences which I have extricated from Fresnel's hypothesis.

(1) The two meridian planes corresponding to any given radius are perpendicular to each other⁹.

(2) So are the two corresponding to any given normal.

(3) Every meridian plane bisects the angle formed by two planes drawn through the radius and the two prime radii.

(4) It also bisects the angle formed by two planes drawn through the normal and the two prime normals.

(5) Each meridian plane contains one line of vibration and the corresponding line of force.

(6) The ray is perpendicular to the line of force.

All these conclusions, except the fourth, are, I believe, original. p. 26

The theory of external and internal conical refraction follows immediately as a particular consequence from the third and fourth combined as already shown;

⁸See above, p. 18.

⁹I have defined the meridian plane to be that which contains radius vector and normal belonging to the same point.

the same propositions also enable us to draw a tangent plane to any point of the wave surface by mere Euclidean geometry. May not some of these conclusions serve to suggest to physical inquirers the question, Has the theory been started from the most natural point of view?

NOTE 2. INVESTIGATION¹⁰ OF THE WAVE SURFACE.

Since the appearance of the preceding parts, I have succeeded in completing the self-sufficiency of my method by deducing the equation to the wave surface from the expressions given in Prop. 5 for the angles between a front and the principal planes in terms of its two velocities. If these angles be ω, ϕ, ψ , and the two velocities v_1, v_2 we found

$$\cos \omega = \sqrt{\frac{(a^2 - v_1^2)(a^2 - v_2^2)}{(a^2 - b^2)(a^2 - c^2)}},$$

$$\cos \phi = \sqrt{\frac{(b^2 - v_1^2)(b^2 - v_2^2)}{(b^2 - a^2)(b^2 - c^2)}},$$

$$\cos \psi = \sqrt{\frac{(c^2 - v_1^2)(c^2 - v_2^2)}{(c^2 - a^2)(c^2 - b^2)}}.$$

Let the tangent plane to the wave surface be written

$$\frac{\cos \omega}{v_1} \cdot x + \frac{\cos \phi}{v_1} \cdot y + \frac{\cos \psi}{v_1} \cdot z = 1,^{11} \quad (\alpha)$$

then

$$\frac{d \frac{\cos \omega}{v_1}}{d \left(\frac{1}{v_1} \right)^2} x + \frac{d \frac{\cos \phi}{v_1}}{d \left(\frac{1}{v_1} \right)^2} y + \frac{d \frac{\cos \psi}{v_1}}{d \left(\frac{1}{v_1} \right)^2} z = 0, \quad (\beta)$$

$$\frac{d \cos \omega}{d(v_2)^2} x + \frac{d \cos \phi}{d(v_2)^2} y + \frac{d \cos \psi}{d(v_2)^2} z = 0. \quad (\gamma)$$

Let¹²

$$\frac{1}{v_1} \sqrt{\frac{a^2 - v_1^2}{a^2 - v_2^2}} = \xi, \quad \sqrt{(a^2 - b^2)(a^2 - c^2)} = \frac{1}{A},$$

$$\frac{1}{v_1} \sqrt{\frac{b^2 - v_1^2}{b^2 - v_2^2}} = \eta, \quad \sqrt{(b^2 - a^2)(b^2 - c^2)} = \frac{1}{B},$$

$$\frac{1}{v_1} \sqrt{\frac{c^2 - v_1^2}{c^2 - v_2^2}} = \zeta, \quad \sqrt{(c^2 - a^2)(c^2 - b^2)} = \frac{1}{C},$$

then equation (γ) becomes

$$A\xi x + B\eta y + C\zeta z = 0, \quad (1)$$

and equation (β)

$$\frac{Aa^2}{\xi}x + \frac{Bb^2}{\eta}y + \frac{Cc^2}{\zeta}z = 0, \quad (2)$$

and equation (α) may be written under two forms, viz.

$$(a^2 - v_2^2)A\xi x + (b^2 - v_2^2)B\eta y + (c^2 - v_2^2)C\zeta z = 1, \quad (3)$$

or

$$\left(\frac{a^2}{v_1^2} - 1\right)\frac{A}{\xi}x + \left(\frac{b^2}{v_1^2} - 1\right)\frac{B}{\eta}y + \left(\frac{c^2}{v_1^2} - 1\right)\frac{C}{\zeta}z = 1. \quad (4)$$

From (1)

$$A\xi x + B\eta y = -C\zeta z. \quad (5)$$

From (2)

$$\frac{Aa^2}{\xi}x + \frac{Bb^2}{\eta}y = -\frac{Cc^2}{\zeta}z. \quad (6)$$

From (3) and (1)

$$A(a^2 - c^2)\xi x + B(b^2 - c^2)\eta y = 1. \quad (7)$$

From (2) and (4)

$$A(a^2 - c^2)\frac{x}{\xi} + B(b^2 - c^2)\frac{y}{\eta} = c^2. \quad (8)$$

From (5) and (6)

$$C^2c^2z^2 - B^2b^2y^2 - A^2a^2x^2 = ABxy\left(a^2\frac{\eta}{\xi} + b^2\frac{\xi}{\eta}\right). \quad (9)$$

From (7) and (8)

$$c^2 - B^2(b^2 - c^2)^2y^2 - A^2(a^2 - c^2)^2x^2 = ABxy\left(\frac{\eta}{\xi} + \frac{\xi}{\eta}\right)(a^2 - c^2)(b^2 - c^2). \quad (10)$$

¹⁰This investigation supplies the step which Mr Tovey was desirous should appear in the *Magazine*. [*Phil. Mag.* March, 1838, p. 261. Ed.]

¹¹In lieu of v_1 we might write v_2 in the denominator without affecting the result.

¹²Observe, that $\frac{\cos \omega}{v_1} = \frac{\sqrt{\left\{\left(\frac{a^2}{v_1^2} - 1\right)(a^2 - v_2^2)\right\}}}{\sqrt{\{(a^2 - b^2)(a^2 - c^2)\}}}$, and so on for the rest.

From (9) and (10)

$$\begin{aligned}
& AB(a^2 - b^2)(a^2 - c^2)(b^2 - c^2)xy\frac{\xi}{\eta} \\
&= a^2c^2 - (a^2 - c^2)(b^2 - c^2)C^2c^2z^2 \\
&\quad - \{a^2(b^2 - c^2)^2 - b^2(a^2 - c^2)(b^2 - c^2)\}B^2y^2 \\
&\quad - \{a^2(a^2 - c^2)^2 - a^2(a^2 - c^2)(b^2 - c^2)\}A^2x^2 \\
&= a^2c^2 - c^2z^2 - c^2y^2 - a^2x^2.
\end{aligned} \tag{11}$$

From (11), interchanging (a, x, ξ) with (b, y, η) we have

$$AB(b^2 - a^2)(b^2 - c^2)(a^2 - c^2)xy\frac{\eta}{\xi} = b^2c^2 - c^2z^2 - c^2x^2 - b^2y^2. \tag{12}$$

Finally, from (11) and (12) we have

$$\begin{aligned}
& \{a^2c^2 - (a^2 - c^2)x^2 - c^2(x^2 + y^2 + z^2)\}\{b^2c^2 - (b^2 - c^2)y^2 - c^2(x^2 + y^2 + z^2)\} \\
&= (a^2 - c^2)(b^2 - c^2)x^2y^2,
\end{aligned}$$

that is

$$\begin{aligned}
& (x^2 + y^2 + z^2)(a^2x^2 + b^2y^2 + c^2z^2) - a^2(b^2 + c^2)x^2 \\
& \quad - b^2(a^2 + c^2)y^2 - c^2(b^2 + a^2)z^2 + a^2b^2c^2 = 0
\end{aligned}$$

the equation required.

2.

ON THE MOTION AND REST OF FLUIDS

[*Philosophical Magazine*, XIII. (1838), pp. 449–453]

p. 28

M. Ostrogradsky's memoir on this subject inserted in the *Scientific Memoirs* seems to have excited much attention, and has been made the occasion of some annotations¹³ by a distinguished writer in the *Philosophical Magazine*. Mr Ivory's recent papers in the same periodical must still more tend to invest with a new interest all such speculations. It seems to me desirable therefore to present the theory of fluids in all the simplicity of which it is susceptible.

I consider a fluid as a collection of particles subject to some law of relative position other than that of rigidity. These particles by their mutual actions maintain the connections of the system. As to the law of force between them we know nothing; but I assume it is a general principle of nature, that for each instant of time the sum of the internal actions (reckoned by the product of each particle into the square of the space due to the internal force acting on it) is a minimum. This in fact is Gauss's principle of least restraint. We may if we please split this principle into two parts; that is to say, assume that the internal system of forces is always such as if acting alone would keep the fluid at rest; and then again assume that any equilibrating system of forces must be subject to the law of virtual velocities. I say *assume*, because it is impossible *à priori* to prove this.

Lagrange's so-called demonstration is unworthy of his name, and (albeit sanctioned by the powerful oral authority of an ex-Cambridge Professor) contrary alike to sense and honesty. It is better therefore at once to proceed upon Gauss's principle. It might easily be shown that this is in effect tantamount in all cases to D'Alembert's and Lagrange's principles combined.

Before entering upon the investigation I may call attention to one point of great analytical interest, and relating to the difficult subject of the algebraical sign, viz. that if the density of a point (x, y) in any circumscribed space be expressed by the quantity $\frac{du}{dx} + \frac{dv}{dy}$ so that the mass is

$$\iint dx dy \left(\frac{du}{dx} \right) + \iint dx dy \left(\frac{dv}{dy} \right),$$

that is not equivalent to

$$\int (u dy + v dx),$$

p. 29

that is if we please

$$\int \left(u \frac{dy}{ds} + v \frac{dx}{ds} \right) ds,$$

¹³*Phil. Mag.* May, 1838, p. 385. ED.

(where s is for clearness' sake and to avoid double limits taken an element of the bounding curve) as at first sight it might appear to be, but is in fact equal to

$$\int \left(u \frac{dy}{ds} - v \frac{dx}{ds} \right) ds.$$

I shall demonstrate this point in the next number¹⁴ of the *Magazine*. It at first caused me some trouble in conducting the annexed inquiry. I shall also take occasion at some other time to revert to a new species (as I believe) of partial differential equations; that is to say, where there are fewer of them than of the principal variables, which may be called therefore Indeterminate Partial Differential Equations. A complete solution of one of these appears in the subjoined

Investigation.

For the sake of simplicity I take an incompressible fluid. The method is nowise different for a fluid of varying density.

Let Δx , Δy , Δz be any displacement undergone by a particle at the point x, y, z parallel to the axes x, y, z respectively; it is easily shown that to satisfy the condition of invariability of mass we must have

$$\frac{d\Delta x}{dx} + \frac{d\Delta y}{dy} + \frac{d\Delta z}{dz} = 0. \quad (\alpha)$$

One relation between u, v, w the velocities parallel to x, y, z is obtained immediately by putting $u\delta t, v\delta t, w\delta t$, for $\Delta x, \Delta y, \Delta z$, which gives

$$\frac{du}{dx} + \frac{dv}{dy} + \frac{dw}{dz} = 0, \quad (1)$$

as usual.

Again, if X, Y, Z be the impressed forces, and X_1, Y_1, Z_1 the internal forces acting on any particle parallel to the axes, we have

$$X_1 + X = \frac{du}{dt} + \frac{du}{dx}u + \frac{du}{dy}v + \frac{du}{dz}w, \quad (2)$$

$$Y_1 + Y = \frac{dv}{dt} + \frac{dv}{dx}u + \frac{dv}{dy}v + \frac{dv}{dz}w, \quad (3)$$

$$Z_1 + Z = \frac{dw}{dt} + \frac{dw}{dx}u + \frac{dw}{dy}v + \frac{dw}{dz}w, \quad (4)$$

from the mere geometry of the question.

p. 30

¹⁴p. 36, below. ED.

Finally, Gauss's principle teaches us that

$$\iiint dx dy dz \{X_1 \Delta X_1 + Y_1 \Delta Y_1 + Z_1 \Delta Z_1\} = 0. \quad (\beta)$$

Now

$$\begin{aligned} & \frac{d(X + X_1)}{dx} + \frac{d(Y + Y_1)}{dy} + \frac{d(Z + Z_1)}{dz} \\ &= \left(\frac{du}{dx}\right)^2 + \left(\frac{dv}{dy}\right)^2 + \left(\frac{dw}{dz}\right)^2 + 2 \left\{ \frac{dv}{dz} \frac{dw}{dy} + \frac{dw}{dx} \frac{du}{dz} + \frac{du}{dy} \frac{dv}{dx} \right\}, \end{aligned}$$

as appears from the equations (1), (2), (3), (4); and hence

$$\frac{d\Delta X_1}{dx} + \frac{d\Delta Y_1}{dy} + \frac{d\Delta Z_1}{dz} = 0,$$

the complete solution of which, free from the sign of integration, is

$$\begin{aligned} \Delta X_1 &= \frac{d\psi}{dy} - \frac{d\phi}{dz}, \\ \Delta Y_1 &= \frac{d\omega}{dz} - \frac{d\psi}{dx}, \\ \Delta Z_1 &= \frac{d\phi}{dx} - \frac{d\omega}{dy}, \end{aligned}$$

ω, ϕ, ψ being any three independent functions of x, y, z .

On substituting these values in equation (β) we obtain

$$\begin{aligned} & \iiint dx dy dz \left\{ X_1 \frac{d\psi}{dy} - Y_1 \frac{d\psi}{dx} \right\} + \iiint dx dy dz \left\{ Y_1 \frac{d\omega}{dz} - Z_1 \frac{d\omega}{dy} \right\} \\ & + \iiint dx dy dz \left\{ Z_1 \frac{d\phi}{dx} - X_1 \frac{d\phi}{dz} \right\} = 0. \end{aligned}$$

This may be put under the form

$$\begin{aligned} & \int dz \iint dx dy \left\{ \frac{d}{dy}(\psi X_1) - \frac{d}{dx}(\psi Y_1) \right\} \\ & + \int dx \iint dy dz \left\{ \frac{d}{dz}(\omega Y_1) - \frac{d}{dy}(\omega Z_1) \right\} \\ & + \int dy \iint dz dx \left\{ \frac{d}{dx}(\phi Z_1) - \frac{d}{dz}(\phi X_1) \right\} \\ & - \iiint dx dy dz \cdot \psi \left(\frac{dX_1}{dy} - \frac{dY_1}{dx} \right) \\ & - \iiint dx dy dz \cdot \omega \left(\frac{dY_1}{dz} - \frac{dZ_1}{dy} \right) \\ & - \iiint dx dy dz \cdot \phi \left(\frac{dZ_1}{dx} - \frac{dX_1}{dz} \right) = 0. \end{aligned}$$

Here it must be remembered that ω, ϕ, ψ are perfectly independent of each other. Also the values of the three first written quantities depend upon the values of X_1, Y_1, Z_1 at the bounding surface; the values of the three last-written depend upon the general values of X_1, Y_1, Z_1 . It is clear therefore that each system of three equations and each member of each system must be separately zero.

The three latter equations give

$$\left. \begin{aligned} \frac{dX_1}{dy} - \frac{dY_1}{dx} &= 0, \\ \frac{dY_1}{dz} - \frac{dZ_1}{dy} &= 0, \\ \frac{dZ_1}{dx} - \frac{dX_1}{dz} &= 0, \end{aligned} \right\} \quad (\gamma)$$

The three former require that for each section of the surface parallel to the plane xy

$$\int \psi(X_1 dx + Y_1 dy) = 0,$$

for each section parallel to yz

$$\left. \begin{aligned} \int \omega(Y_1 dy + Z_1 dz) &= 0, \\ \int \phi(Z_1 dz + X_1 dx) &= 0 \end{aligned} \right\}^* \quad (\delta)$$

for each section parallel to zx and these equations are to hold good whatever ψ, ϕ, ω may be. From the equations (γ) we derive

$$X_1 dx + Y_1 dy + Z_1 dz = df, \quad (5)$$

from equations (δ) we obtain

f = constant for all points in any section of the bounding surface parallel to the plane of xy ,

f = constant for all points in any section of the bounding surface parallel to the plane of yz ,

f = constant for all points in any section of the bounding surface parallel to the plane of zx .

Now by drawing through all the points in a plane parallel to xy , planes parallel to yz , we may cover the whole surface; hence f is constant all over the surface bounding the fluid.

* See remark at introduction.

Therefore

$$X_1 dx + Y_1 dy + Z_1 dz = 0, \quad (6)$$

for all variations of dx, dy, dz taken upon the surface.

The equations (1, 2, 3, 4, 5, 6) are coincident with those obtained by the usual method; with this difference, that X_1, Y_1, Z_1 here take the place of

$$-\frac{dp}{dx}, \quad -\frac{dp}{dy}, \quad -\frac{dp}{dz}.$$

Thus then we have obtained all the conditions requisite for determining the motion of fluids from the universal principle of least constraint conjoined with the specific character of the system in question.

General Remarks.

In the case of equilibrium, that is in the case where no particle moves, we have $X_1 + X = 0, Y_1 + Y = 0, Z_1 + Z = 0$. Hence $X dx + Y dy + Z dz$ is a complete differential always and zero for the surface.

The above results have been obtained upon the principles of the differential calculus, and the continuity of the forces has been tacitly assumed. If now we were to suppose forces of finite magnitude (as compared with the *whole sum* acting upon the entire system) to be applied to a layer of single particles or to a layer of a thickness of the same order of magnitude as the distances between the particles themselves, (which has been treated as an infinitesimal) it would appear that our results would be no longer applicable, just in the same manner as it would be erroneous to apply the principle of *vis-viva* (for example) without modification, to the case of impulsive forces, because we had deduced it by the calculus in the case of the motion being continuous. Hence the above equations ought not strictly to apply to the motion or rest of a fluid *contained between physical surfaces*; for the pressure afforded by these surfaces, whatever its actual value may be, we know *à priori* is commensurable with the whole amount of force acting on the fluid; but the immediate application of this pressure (*alias* repulsive force) is confined to the bounding layer of fluid particles, or at most extends to a distance bearing a low ratio to the distances between the particles themselves.

Accordingly, to the non-applicability of the equations for free fluids to the case of fluids confined at the boundaries, and to an independent investigation upon the minimum principle for this class of problems, it is, that I look for the true explanation of the phenomena of capillary attraction (vulgarly so called).

3.

ON THE MOTION AND REST OF RIGID BODIES

[*Philosophical Magazine*, XIV. (1839), pp. 188–190]

p. 33

In the subjoined investigation, which, as far as I know, is my own, I apply the same method to rigid as in the preceding paper I applied to fluid systems.

Let x, y, z be the coordinates of any particle in a rigid body; x', y', z' the coordinates of some other particle, and let

$$x' = x + h, \quad y' = y + k, \quad z' = z + l.$$

Call $\Delta x, \Delta y, \Delta z$ the increments which x, y, z receive after the lapse of a small interval of time; so that terms in which they enter in two or more dimensions may be neglected.

Then

$$\begin{aligned} \Delta(x') &= \Delta x + \frac{d\Delta x}{dx}h + \frac{d\Delta x}{dy}k + \frac{d\Delta x}{dz}l + P, \\ \Delta(y') &= \Delta y + \frac{d\Delta y}{dx}h + \frac{d\Delta y}{dy}k + \frac{d\Delta y}{dz}l + Q, \\ \Delta(z') &= \Delta z + \frac{d\Delta z}{dx}h + \frac{d\Delta z}{dy}k + \frac{d\Delta z}{dz}l + R, \end{aligned}$$

P, Q, R containing binary and higher combinations of h, k, l , which we shall have no occasion to express.

At the commencement of the interval the squared distance of the two particles was $(x' - x)^2 + (y' - y)^2 + (z' - z)^2$; at the end of the interval the distance squared is

$$(x' - x + \Delta(x') - \Delta x)^2 + (y' - y + \Delta(y') - \Delta y)^2 + (z' - z + \Delta(z') - \Delta z)^2,$$

and these two expressions must be the same by the conditions of rigidity whatever h, k , and l may be; that is

$$\begin{aligned} h^2 + k^2 + l^2 &= \left(h + \frac{d\Delta x}{dx}h + \frac{d\Delta x}{dy}k + \frac{d\Delta x}{dz}l + P \right)^2 \\ &\quad + \left(k + \frac{d\Delta y}{dx}h + \frac{d\Delta y}{dy}k + \frac{d\Delta y}{dz}l + Q \right)^2 \\ &\quad + \left(l + \frac{d\Delta z}{dx}h + \frac{d\Delta z}{dy}k + \frac{d\Delta z}{dz}l + R \right)^2, \end{aligned}$$

for all values of h, k , and l .

p. 34

Hence rejecting infinitesimals of the second order and equating to zero separately the coefficients of h^2 , k^2 , l^2 , and of kl , lh , hk , we have

$$\begin{aligned} \frac{d\Delta x}{dx} = 0, & \quad (a) \quad \frac{d\Delta y}{dz} + \frac{d\Delta z}{dy} = 0, & (d) \\ \frac{d\Delta y}{dy} = 0, & \quad (b) \quad \frac{d\Delta z}{dx} + \frac{d\Delta x}{dz} = 0, & (e) \\ \frac{d\Delta z}{dz} = 0, & \quad (c) \quad \frac{d\Delta x}{dy} + \frac{d\Delta y}{dx} = 0. & (f) \end{aligned}$$

By differentiating (d), (e), (f) with respect to z, x, y respectively, and substituting from (a), (b), (c), we obtain

$$\frac{d^2\Delta y}{dz^2} = 0, \quad \frac{d^2\Delta z}{dx^2} = 0, \quad \frac{d^2\Delta x}{dy^2} = 0.$$

By differentiating the same with respect to y, z, x respectively, and proceeding as before, we have

$$\frac{d^2\Delta z}{dy^2} = 0, \quad \frac{d^2\Delta x}{dz^2} = 0, \quad \frac{d^2\Delta y}{dx^2} = 0.$$

Thus, then, we have

$$\begin{aligned} \frac{d\Delta x}{dx} = 0, \quad \frac{d^2\Delta x}{dy^2} = 0, \quad \frac{d^2\Delta x}{dz^2} = 0, \\ \frac{d\Delta y}{dy} = 0, \quad \frac{d^2\Delta y}{dz^2} = 0, \quad \frac{d^2\Delta y}{dx^2} = 0, \\ \frac{d\Delta z}{dz} = 0, \quad \frac{d^2\Delta z}{dx^2} = 0, \quad \frac{d^2\Delta z}{dy^2} = 0, \end{aligned}$$

therefore

$$\Delta x = A + By + Cz, \tag{o}$$

$$\Delta y = D + Ez + Fx, \tag{p}$$

$$\Delta z = G + Hx + Ky, \tag{q}$$

A, B, C, D, E, F being constant for a *given instant* of time; between which by virtue of the equations (d), (e), (f), we have the relations

$$E + K = 0, \quad H + C = 0, \quad B + F = 0.$$

If we call u, v, w the three component velocities of the particles at x, y, z parallel to the three axes, and X_1, Y_1, Z_1 , the three internal forces, it is at once seen that

u, v, w , as also $\Delta X_1, \Delta Y_1, \Delta Z_1$ must be subject to the same equations as limit $\Delta x, \Delta y, \Delta z$; so that

$$u = a + \gamma y - \beta z, \quad (1)$$

$$v = b + \alpha z - \gamma x, \quad (2)$$

$$w = c + \beta x - \alpha y, \quad (3)$$

$$\Delta X_1 = a_1 + \gamma_1 y - \beta_1 z, \quad (h)$$

$$\Delta Y_1 = b_1 + \alpha_1 z - \gamma_1 x, \quad (j)$$

$$\Delta Z_1 = c_1 + \beta_1 x - \alpha_1 y. \quad (k)$$

p. 35

Also if X, Y, Z be the impressed forces, we have

$$X_1 + X = \frac{du}{dt}, \quad (4)$$

$$Y_1 + Y = \frac{dv}{dt}, \quad (5)$$

$$Z_1 + Z = \frac{dw}{dt}. \quad (6)$$

And by Gauss's principle, calling m the mass of the particle at x, y, z ,

$$\Delta \sum m(X_1^2 + Y_1^2 + Z_1^2) = 0.$$

Hence equating separately to zero the coefficients of a_1, b_1, c_1 and of $\alpha_1, \beta_1, \gamma_1$ in the quantity $\sum m(X_1 \Delta X_1 + Y_1 \Delta Y_1 + Z_1 \Delta Z_1)$ we have

$$\left. \begin{aligned} \sum m X_1 &= 0, \\ \sum m Y_1 &= 0, \\ \sum m Z_1 &= 0, \\ \sum m (Z_1 y - Y_1 z) &= 0, \\ \sum m (X_1 z - Z_1 x) &= 0, \\ \sum m (Y_1 x - X_1 y) &= 0. \end{aligned} \right\} \quad (7-12)$$

Lastly, we have the equations

$$u = \frac{dx}{dt}, \quad (13)$$

$$v = \frac{dy}{dt}, \quad (14)$$

$$w = \frac{dz}{dt}. \quad (15)$$

From the fifteen equations marked (1) to (15), the motion may be determined by assigning the position of each particle at the end of the time t in terms of its

three initial coordinates, its three initial velocities, and the initial values of the nine quantities

$$\begin{array}{lll} \sum mx, & \sum myz, & \sum mx^2, \\ \sum my, & \sum mzx, & \sum my^2, \\ \sum mz, & \sum mxy, & \sum mz^2. \end{array}$$

In the case of rest $X_1 = -X$, $Y_1 = -Y$, $Z_1 = -Z$, and the equations (7) to (12) inclusively taken, express the conditions of equilibrium.

The equations (o) , (p) , (q) , which have been obtained from conditions *purely geometrical*, establish the well-known but interesting and *not obvious* fact, that any *small* motion of a rigid body may be conceived as made up of a motion of translation and a motion about *one* axis.

4.

ON DEFINITE DOUBLE INTEGRATION, SUPPLEMENTARY TO A FORMER PAPER ON THE MOTION AND REST OF FLUIDS

[*Philosophical Magazine*, XIV. (1839), pp. 298–300]

p. 36

In a paper on Fluids which appeared in the December Number of this *Magazine*, I had occasion to remark, that the mass of an area having at the point (x, y) a density $\frac{du}{dx} + \frac{dv}{dy}$ could be expressed by the simple formula

$$\int_l^0 \left\{ u \frac{dy}{ds} - v \frac{dx}{ds} \right\} ds;$$

l being the length, and ds an element of the bounding curve: this may be thought to require some explanation.

(1) Let $APBq$ represent any oval; PpL , QqM any two contiguous ordinates cutting the curve in Pp , Qq respectively, AC , BD the two extreme tangents parallel to Oy , and ρ the density at any point (x, y) . The expression $\iint \rho dx dy$ will serve to denote the mass of the oval area $APBq$, and the limits may be twice taken, that is, (i) the two values of y corresponding to any one of x ; and (ii) the two values of x corresponding to C and D . This method is in fact tantamount to taking the sum of the columns Pp , qQ ; but this is not necessary, for $APBq$ may be considered as the algebraical sum of the mixtilinear area $APQBDC$, and the mixtilinear area $BDCApq$, or (if any line $O'C'D'$ be drawn parallel to $OCLMD$) of $APQBD'C'$ and $BD'C'Apq$.




Fig. 1.

Thus then the mass = $\int dx (\int \rho dy)$, $\int \rho dy$ being left indeterminate, and the extremity of x travelled round from C to D , and back again from D to C . p. 37

This will be better expressed by transforming the variable, and summing with respect to some quantity, such as the arc of the curve, which continuously

increases, or if we please, with respect to θ , the angle subtending any point taken within the curve.

The mass is then

$$= \pm \int_{2\pi}^0 d\theta \left\{ \left(\int \rho dy \right) \frac{dx}{d\theta} \right\};$$

always remembering that no constant need be added to $\int \rho dy$, and that the doubtful sign arises from the choice of ways in which θ may be measured round. If the area be not included by one line; but by several, as for example, by a curve and a right line, the above integral, if broken up into as many parts as there are breaches of continuity, will still apply.

(2) Let us suppose that we have two areas exactly coinciding with, and overlapping one another; but the density of the one at (x, y) to be ρ , and of the other ρ' .

Let the mass of the first be treated as the sum of columns parallel to Oy , and that of the second as the sum of columns parallel to Ox .

The one will be represented by

$$\pm \int_{2\pi}^0 d\theta \left(\int \rho dy \right) \frac{dx}{d\theta},$$

the other will be represented by

$$\pm \int_{2\pi}^0 d\theta \left(\int \rho' dx \right) \frac{dy}{d\theta},$$

and the sum of the two, or the joint mass, by

$$\pm \int_{2\pi}^0 d\theta \left(\int \rho dy \right) \frac{dx}{d\theta} \pm \int_{2\pi}^0 d\theta \left(\int \rho' dx \right) \frac{dy}{d\theta}.$$

So long as these two operations are performed separately, the doubtful signs may be preserved in each term, because s need not be travelled round in the same direction for the two summations; but if we perform the second integration conjointly for the two masses, their sum

$$= \pm \int_{2\pi}^0 d\theta \left\{ \left(\int \rho dy \right) \frac{dx}{d\theta} \pm? \left(\int \rho' dx \right) \frac{dy}{d\theta} \right\},$$

the mark of interrogation denoting that *one or the other*, but not *either* of the signs $\pm$ must be used, and the question is, which?

This will be answered by taking different points in the bounding line which may be continuous or not. Now every line returning into itself, whether continuous or not, will naturally divide with respect of any given

p. 38

system of axes, into at most four parts, or sets of parts; two in which dx and dy both increase or both decrease, and two in which one increases and the other decreases.

Take P_1, P_2, P_3, P_4 , any points in the four quadrants respectively, it will be observed that,




Fig. 2.

At P_1 the ρ column enters additively, and the ρ' column subtractively.

At P_2 both columns are additive.

At P_3 the ρ' column is additive and the ρ column subtractive.

At P_4 both columns enter subtractively.

Again, reckoning round in the direction of the arrows,

At P_1 , x and y are both increasing.

At P_2 , x is increasing and y decreasing.

At P_3 , x and y both decrease.

At P_4 , x is decreasing and y increasing.

Thus when $\int \rho dy$ and $\int \rho' dx$ are affected with the same signs, dx and dy are of opposite signs; and when $\int \rho dy$, $\int \rho' dx$ are of opposite signs, dx and dy are of the same sign.

Hence it appears that the mass of the area, whose density at (x, y) is $\rho + \rho'$, is capable of being represented by

$$\pm \int_{2\pi}^0 d\theta \left\{ \left(\int \rho dy \right) \frac{dx}{d\theta} - \left(\int \rho' dx \right) \frac{dy}{d\theta} \right\}.$$

5.

ON AN EXTENSION OF SIR JOHN WILSON'S THEOREM TO ALL NUMBERS WHATEVER

[*Philosophical Magazine*, XIII. (1838), p. 454]

p. 39

The annexed original theorem in numbers will serve as a pendant to the elegant discovery announced by the ever-to-be-lamented and commemorated Horner¹⁵ with his dying voice, in your valued pages¹⁶.

THEOREM.

If N be any number whatever and

$$p_1, p_2, p_3 \dots p_c$$

be all the numbers less than N and prime to it, then either

$$p_1 \cdot p_2 \cdot p_3 \dots p_c + 1,$$

or else

$$p_1 \cdot p_2 \cdot p_3 \dots p_c - 1,$$

is a multiple of N .

¹⁵Horner's proof is highly valuable as a novel and highly ingenious form of reasoning, but his theorem may be deduced with infinitely more ease and brevity from Fermat's than he seems to have been aware of.

¹⁶*Phil. Mag.* Vol. XI. p. 456. ED.

6.

NOTE TO THE FOREGOING

[*Philosophical Magazine*, XIV. (1839), pp. 47, 48]

p. 39

I have to apologize for calling “original” (in the last Number of the *Magazine*) the theorem of numbers which I termed “a pendant to Horner’s theorem.” This Mr Ivory has done me the honour to inform me may be found in Gauss’s *Disquisitiones Arithmeticae*, p. 76. As Horner’s extension of Fermat’s theorem suggested this extension of Sir John Wilson’s to me, so I concluded that had this extension of Wilson’s been known to the world it would naturally have suggested his to Horner. No acknowledgment of this kind having been made, I took it for granted that the theorem I gave was new. Undoubtedly had Mr Horner been aware of Gauss’s theorem he would have made mention of it.

I take this opportunity of adding that my acquaintance with Gauss’s principle¹⁷ has not been derived from the study of his works, but from a casual statement of it in an English work, *Dynamics*, by Mr Earnshaw, of St John’s College, Cambridge.

¹⁷See p. 28 above. ED.

7.

ON RATIONAL DERIVATION FROM EQUATIONS OF COEXISTENCE, THAT IS TO SAY, A NEW AND EXTENDED THEORY OF ELIMINATION. PART I

[Philosophical Magazine, XV. (1839), pp. 428-435]

p. 40

Any number of equations existing at the same time and having the same quantities repeated, may be termed equations of coexistence: in the present paper we consider only the case of two algebraical equations:

$$x^m + a_1x^{m-1} + a_2x^{m-2} + \cdots + a_m = 0,$$

$$x^n + b_1x^{n-1} + b_2x^{n-2} + \cdots + b_n = 0.$$

The above being “equations of coexistence,” x is called “the repeating term.”

If we suppose the equation

$$c_0x^r + c_1x^{r-1} + c_2x^{r-2} + \cdots + c_r = 0$$

to be capable of being deduced from the two above, and, therefore, necessarily implied by them, this will be called “a Particular Derivative” from the equations of coexistence, of the r th degree, (r being supposed less than m and n ¹⁸), and the coefficients being rational functions of the coefficients of the equations of coexistence.

There will be an indefinite number in general of such derivatives, and the form involving arbitrary quantities which includes them all is called “the general derivative of the r th degree.”

Any “Particular Derivative,” in which the terms are all integral, numerically as well as literally speaking, is called an “Integral Derivative.”

That “Integral Derivative” of any given degree in which the literal parts of the coefficients are of the lowest possible dimensions¹⁹, and the numerical parts as low as they can be made, is called the “Prime Derivative”²⁰

p. 41

of that degree. So that there is nothing left ambiguous in the prime derivative save the sign.

The “Derivative by succession” is that particular derivative which is obtained by performing upon the equations of coexistence the process commonly employed

¹⁸This restriction upon the value of r is not essentially requisite, and is only introduced to keep the attention fixed upon the particular objects of this first Part.

¹⁹Of course the dimensions of the coefficients in the equations of coexistence are to be understood as denoted by the indices subscribed.

²⁰The results of this and some following papers were repeated, with demonstrations, in the paper “On a Theory of the Syzygetic Relations of two rational integral functions comprising an application to the Theory of Sturm’s Functions, and that of the greatest Algebraical Common Measure,” *Phil. Trans. Royal Soc.* Vol. CXLIII., Part I. pp. 407-548, 1853. See below Section II. Art. (16) of that paper. Ed.

for the discovery of the greatest common measure, and equating the successive remainders to zero.

To express the product of the sums formed by adding each of one row of quantities to each of another row, we simply write the one row above the other; a notation clearly capable of extension to any number of rows, which would not be the case if we spoke of differences instead of sums²¹.

THEOREM 1.

Let $h_1, h_2, \dots, h_m$, be the roots of one equation of coexistence, $k_1, k_2, \dots, k_n$, the roots of the other. The general derivative of the r th degree is represented by

$$\Sigma \left(SR(h_1, h_2, h_3 \dots h_r) \{ (x - h_1)(x - h_2) \dots (x - h_r) \} \times \left\{ \begin{matrix} h_{r+1}, h_{r+2} \dots h_m \\ -k_1, -k_2 \dots -k_n \end{matrix} \right\} \right) = 0,$$

$SR(h_1, h_2, h_3 \dots h_r)$ denoting any symmetrical rational (integral or fractional) function of $h_1, h_2 \dots h_r$;

$$\left\{ \begin{matrix} h_{r+1}, h_{r+2} \dots h_m \\ -k_1, -k_2 \dots -k_n \end{matrix} \right\}$$

being to be interpreted as above explained, and Σ of course including as many terms as there are ways of putting n things r and r together²².

A form tantamount to the above, and which may be substituted for it, is its analogue,

$$\Sigma \left(SR(k_1, k_2 \dots k_r) \{ (x - k_1)(x - k_2) \dots (x - k_r) \} \times \left\{ \begin{matrix} k_{r+1}, k_{r+2} \dots k_n \\ -h_1, -h_2 \dots -h_m \end{matrix} \right\} \right) = 0.$$

When $r = 0$ the theorem gives simply

$$\left\{ \begin{matrix} h_1, h_2 \dots h_m \\ -k_1, -k_2 \dots -k_n \end{matrix} \right\} = 0,$$

and is coincident with that given by Bezout in his Theory of *Elimination*.

p. 42

Subsidiary Theorem (A).

²¹The wider views which I have attained since writing the above, and which will be developed in a future paper, lead me to request that this notation may be considered only as temporary. It would have been more in accordance with these views to have used the two rows to denote products of differences than of sums. But a change now in the text would be very apt to cause errors in printing.

²²The general derivative may clearly be expressed also by the sum of any two particular derivatives affected respectively with arbitrary rational coefficients. The equivalency of an arbitrary function to two arbitrary multipliers is very remarkable, and analogous to what occurs in the solution of certain differential equations.

If $h_1, h_2 \dots h_m$ be the roots of the equation

$$x^m + a_1 x^{m-1} + a_2 x^{m-2} + \dots + a_m = 0,$$

and if

$$e^m + a_1 e^{m-1} + a_2 e^{m-2} + \dots + a_m - u = 0,$$

then

$$\Sigma \frac{h_1^r}{(h_1 - h_2)(h_1 - h_3) \dots (h_1 - h_m)} = \frac{1}{r+1} \frac{d}{du} \Sigma(e^{r+1}),$$

u being made zero after differentiation.

COR. If $R(h_1)$ denote any integral rational function of h_1 , then

$$\Sigma \frac{R(h_1)}{(h_1 - h_2)(h_1 - h_3) \dots (h_1 - h_m)}$$

is always integral and is zero when the dimensions of $R(h_1)$ fall short of $(m-1)$.

Subsidiary Theorem (B).

$$\Sigma \frac{SR(h_1, h_2 \dots h_r)}{\left\{ \begin{array}{c} h_1, h_2 \dots h_r \\ -h_{r+1}, -h_{r+2} \dots -h_m \end{array} \right\}}$$

can be expressed by the sum of terms, each of which is the product of series of the form

$$\Sigma \frac{R(h_1)}{(h_1 - h_2)(h_1 - h_3) \dots (h_1 - h_m)},$$

it is always integral, and when the dimensions of the numerator fall short of $(m-r)r$ it vanishes²³.

Subsidiary Theorem (C).

The *only* modes of satisfying the equation

$$\Sigma \{f(h_1, h_2 \dots h_r) \times SR(h_1, h_2 \dots h_r)\} = 0,$$

for all forms of the latter factors short of $(m-r)(n-r)$ dimensions, are to put $f(h_1, h_2 \dots h_r) = 0$, or else

$$f(h_1, h_2 \dots h_r) = \frac{\text{constant}}{\left\{ \begin{array}{c} h_1, h_2 \dots h_r \\ -h_{r+1}, -h_{r+2} \dots -h_m \end{array} \right\}}.$$

p. 43

²³It may be remarked also in passing, that any term in the numerator which contains any one power not greater than $m-2r$ may be neglected and thrown out of calculation. Moreover, an analogous proposition may be stated of fractions in the denominators of which any number of rows are written one under the other; see the first note, page 41.

THEOREM 2.

By virtue of the subsidiary theorem (B), the two equations

$$\pm \Sigma \left((x - h_1)(x - h_2) \cdots (x - h_r) \times \frac{\left\{ \begin{matrix} h_{r+1}, h_{r+2} \dots h_m \\ -k_1, -k_2 \dots -k_n \end{matrix} \right\}}{\left\{ \begin{matrix} h_{r+1}, h_{r+2} \dots h_m \\ -h_1, -h_2 \dots -h_r \end{matrix} \right\}} \right) = 0,$$

$$\pm \Sigma \left((x - k_1)(x - k_2) \cdots (x - k_r) \times \frac{\left\{ \begin{matrix} k_{r+1}, k_{r+2} \dots k_n \\ -h_1, -h_2 \dots -h_m \end{matrix} \right\}}{\left\{ \begin{matrix} k_{r+1}, k_{r+2} \dots k_n \\ -k_1, -k_2 \dots -k_r \end{matrix} \right\}} \right) = 0,$$

are each integer derivatives of the r th degree.

THEOREM 3.

And by virtue of the subsidiary theorem (C), the two above equations are the “Prime Integer Derivatives,” and are exactly identical with each other.

COR. 1.. The leading coefficient of the “prime derivative” of the r th degree is always of $(m - r)(n - r)$ dimensions.

COR. 2.. If P_r be the prime derivative of the r th degree and if $(X = 0, Y = 0)$ be the two equations of coexistence, and λ_r, μ_r the two “prime constituents of multiplication” to the said derivative, that is if λ_r and μ_r satisfy the equation $\lambda_r X + \mu_r Y = P_r$, then the coefficient of the leading terms in λ_r and in μ_r is of $(m - r - 1)(n - r - 1)$ dimensions.

THEOREM 4.

The “Prime Derivative” of any given degree is an exact factor of the “derivative by succession,” of the same degree. The quotient resulting from striking out this factor is called “the quotient of succession.”

THEOREM 5.

If L_1, L_2, L_3 , &c., be the leading coefficients of the derivatives occurring first, second, third, &c., in order after the equations of coexistence, and if Q_1, Q_2, Q_3 , &c., represent the first, second, third, “quotients of succession” reckoned in the same order, then

$$Q_1 = 1, \quad Q_2 = \frac{1}{L_1^2}, \quad Q_3 = \frac{L_1^4}{L_2^2}, \quad Q_4 = \frac{L_2^4}{L_1^4 L_3^2},$$

$$Q_{2n} = \frac{L_2^2 L_4^4 \cdots L_{2n-4}^4 L_{2n-2}^4}{L_1^4 L_3^4 \cdots L_{2n-3}^4 L_{2n-1}^2},$$

$$Q_{2n+1} = \frac{L_1^4 L_3^4 \cdots L_{2n-3}^4 L_{2n-1}^4}{L_2^4 L_4^4 \cdots L_{2n-2}^4 L_{2n}^2}.$$

COR. Hence, in place of Sturm's auxiliary functions, we may substitute the functions derived from the equations of coexistence $\left(fx = 0, \frac{dfx}{dx} = 0 \right)$ according to Theorem 2, due regard being had to the sign.

Scholium. Hitherto it has been supposed that the values of the coefficients in the equations of coexistence are independent of one another, but particular relations may be supposed to exist which shall cause the leading terms given by Theorem 2 to vanish, giving rise to anormal or singular primes, as they may be called, of the degree r of fewer than $(m-r)(n-r)$ dimensions. The theory of this, the failing case (so to say), is highly interesting, and I have already discovered the law of formation for the quotients of succession on the supposition of *any number* of primes vanishing consecutively; but I forbear to vex the patience of my reader further, the more so, as I hope soon to be able to present a complete memoir, with all the steps here indicated filled up, and numerous important additions, (the perfect image of which this is but a rough mould), as homage to the learned and illustrious society which has lately done me the honour of admitting me into its ranks.

Why this has not already been done must be excused, by the fact of the theory having suggested itself abroad in the intervals of sickness²⁴. Yet thus much will I add in general terms, namely, that as many primes as vanish consecutively, so many units must be added to the index 2 of the accessions^{25 26}

received in the numerator and denominator of the subsequent quotient; and in the quotient after that, it is not the square of the leading term of the penultimate

²⁴The reflections which Sturm's memorable theorem had originally excited, were revived by happening to be present at a sitting of the French Institute, where a letter was read from the Minister of Public Instruction, requesting an opinion upon the expediency of forming tables of elimination between two equations as high as the 5th or 6th degree containing one repeating term. The offer was rejected, on the ground of the excessive labour that would be required. I think that this has been very much overrated; and probably many will be of the same opinion who have dwelt upon the fact that no numerical quantity will occur in the result higher than the highest index of the repeating term. Would it not redound to the honour of British science that some painstaking ingenious person should gird himself to the task? and would not this be a proper object to meet with encouragement from the Scientific Association of Great Britain?

²⁵That the appearance of the index 4 may not startle, let my reader bear in mind that there are what may be termed secondary derivatives of succession for every degree appearing in the process of successive division.

²⁶The prime derivatives must be capable of yielding an internal evidence of the truth of Sturm's theorem. In fact, for the case of all the roots being possible, a little consideration will serve to show that the leading term of each prime derivative of the equation $\left(fx \frac{dfx}{dx} \right) = 0$ will consist of a series of fractions, each of which fractions is, numerically speaking, of the same sign.

prime,—but the product of this term by the leading term of that anormal prime of the same degree which has the lowest dimensions,—that finds its way into the numerator. The rest of the formation remaining undisturbed, unless and until a new failure have taken place.

NOTE ON STURM’S THEOREM.

When one of the equations of coexistence is the differential coefficient with respect to the repeated term of the other, the prime derivatives given in Theorem 2 which coincide in this case with Sturm’s auxiliary functions reduced to their lowest terms, may be exhibited under an integral aspect.

Let *SPD* intimate that the squared product of the differences is to be taken of the quantities which follow it.

Let *S*₁ indicate the sum of the quantities to which it is prefixed.

*S*₂ the sum of the binary products.

*S*₃ the sum of the ternary products, and so on

Let *h*₁, *h*₂ . . . *h*_{*n*} be the roots of any equation.

Then Sturm’s last auxiliary function may be replaced by

$$SPD(h_1, h_2 \dots h_n).$$

The last but one may be replaced by

$$\Sigma SPD(h_1, h_2 \dots h_{n-1})x + \Sigma S_1(h_2, h_3 \dots h_{n-1})SPD(h_1, h_2 \dots h_{n-1}).$$

The one preceding by

$$\begin{aligned} \Sigma SPD(h_1, h_2 \dots h_{n-2})x^2 + \Sigma S_1(h_1, h_2 \dots h_{n-2})SPD(h_1, h_2 \dots h_{n-2})x \\ + \Sigma S_2(h_1, h_2 \dots h_{n-2})SPD(h_1, h_2 \dots h_{n-2}), \end{aligned}$$

and so on.

Thus then Sturm’s rule for determining the absolute number of real roots in an equation is based wholly and solely upon the following

ALGEBRAICAL PROPOSITION.

If there be *n* quantities, real and imaginary, the imaginary ones entering in pairs, as many changes of sign as there are in the terms

$$\begin{aligned} \Sigma SPD(h_1, h_2), \\ \Sigma SPD(h_1, h_2, h_3), \\ \dots\dots\dots \\ \Sigma SPD(h_1, h_2 \dots h_{n-1}), \\ \Sigma SPD(h_1, h_2 \dots h_n), \end{aligned}$$

so many in number are these pairs.

Query (1). Is there no proposition applicable to any n quantities whatever?

Query (2). Is there no faintly analogous proposition applicable to higher powers than the squares?

Query (3). Seeing that in forming the coefficients in the equation of the squares of the differences, we pass from n functions of the roots to $n\frac{n-1}{2}$ and not n functions, of their squared differences, does not a natural passage to the former lie through n functions of the squared differences?

In other words, may not the quantities $\Sigma SPD(h_1, h_2 \dots h_n)$, &c., serve as natural and valuable intermediaries between the coefficients of an equation involving simple quantities and the coefficients of the equation involving the squares of their differences?

P.S. In the next part I trust to be able to present the readers of this *Magazine* with a direct and symmetrical method of eliminating any number of unknown quantities between any number of equations of any degree, by a newly invented process of symbolical multiplication, and the use of compound symbols of notation.

I must not omit to state that the constituents of multiplication λ_r and μ_r explained in Cor. 2 to Theorem 3 are equal to the expression

$$\Sigma(x - k_1)(x - k_2) \dots (x - k_{n-r-1}) \frac{\left\{ \begin{array}{c} k_1, k_2 \dots k_{n-r-1} \\ -h_1, -h_2 \dots -h_m \end{array} \right\}}{\left\{ \begin{array}{c} k_1, k_2 \dots k_{n-r-1} \\ -k_{n-r}, \dots -k_n \end{array} \right\}},$$

and its analogue respectively.

8.

ON DERIVATION OF COEXISTENCE. PART II. BEING THE THEORY OF SIMULTANEOUS SIMPLE HOMOGENEOUS EQUATIONS

[Philosophical Magazine, XVI. (1840), pp. 37-43]

p. 47

Art. (1). We shall have constant occasion in this paper to denote different quantities by the same letter affected with different subscribed numerical indices.

Such a letter is to be termed a "Base."

Every character consisting of a base and an inferior index, this index is called an argument of the base, namely, the first, second, or n th argument, according as 1, 2, or in general n , be the number subscribed.

Art. (2). I use the symbol PD to denote the product of the differences of the quantities to which it is prefixed (each being to be subtracted from each that follows); thus

$$PD(a, b, c) \text{ indicates } (b - a)(c - a)(c - b).$$

$$PD(0, a, b, c) \text{ indicates } abc(b - a)(c - a)(c - b).$$

$$PD(0, a, b, c \dots l) \text{ indicates } abc \dots l \times PD(a, b, c \dots l).$$

Art. (3). For want of a better symbol I use the Greek letter ζ to denote that the product of factors to which it is prefixed is to be effected after a certain symbolical manner. This I shall distinguish as the zeta-ic product.

The symbol ζ will never be prefixed except to factors, each of which is made up of one or more terms, consisting solely of linear arguments of different bases, that is, characters bearing indices below but none above.

I am thereby enabled to give this short rule for zeta-ic multiplication: "Imagine all the inferior indices to become superior, so that each argument is transformed into a power of its base; multiply according to the rules of ordinary algebra; after the multiplication has been done fully out depress all the indices into their original position; the result is the zeta-ic product²⁷."

p. 48

Thus for example $\zeta(a_r, b_s)$ is the same as simply $a_r b_s$, but $\zeta(a_r, a_s)$ represents not $a_r a_s$ but a_{r+s} .

So in like manner

$$\zeta\{(a_h - b_k)(a_l - b_m)\} = a_{h+l} - a_h b_m - b_k a_l + b_{m+k},$$

$$\zeta\{(a_1 - b_1)(a_1 - c_1)(b_1 - c_1)\}$$

²⁷It is scarcely necessary to add that an analogous interpretation may be extended to any zeta-ic function whatever. Thus $\zeta(a_1 + b_1)^2 = a_2 + 2a_1 b_1 + b_2$, $\zeta \cos(a_1) = 1 - \frac{a_2}{1 \cdot 2} + \frac{a_4}{1 \cdot 2 \cdot 3 \cdot 4}$, &c.

= the depressed product of $(a - b)(a - c)(b - c)$
= the depressed value of $a^2(b - c) + b^2(c - a) + c^2(a - b)$,

that is,

$$= a_2b_1 - a_2c_1 + b_2c_1 - b_2a_1 + c_2a_1 - c_2b_1.$$

Art. (4). We shall have occasion in this part to combine the two symbols ζ, PD : thus we shall use

$$\zeta PD(a_1b_1) \text{ to denote } \zeta(b_1 - a_1),$$

$$\zeta PD(a_1b_1c_1) \text{ to denote } \zeta\{(b_1 - a_1)(c_1 - a_1)(c_1 - b_1)\}.$$

Art. (5). For the sake of elegance of diction I shall in future sometimes omit to insert the inferior index when it is unity; but the reader must always bear in mind that it is to be *understood* though not expressed.

I shall thus be able to speak of the zeta-ic product of such and such bases mentioned by name.

Art. (6). We are not yet come to the limit of the powers of our notation. The zeta-ic product of the sum of arguments will consist of the sum of products of arguments, each argument being (as I have defined) made up of a base and an inferior index. Now we may imagine each index of every term of the zeta-ic product after it is fully expanded to be increased or diminished by unity, or each at the same time to be increased or diminished by 2, or each in general to be increased or diminished by r . I shall denote this alteration by affixing an r with the positive or negative sign to the ζ . Thus

$$\zeta(a_1 - b_1)(a_1 - c_1) \text{ being equal to } a_2 - a_1c_1 + b_1c_1 - b_1a_1,$$

$$\zeta_{+1}(a_1 - b_1)(a_1 - c_1) \text{ is equal to } a_3 - a_2c_2 + b_2c_2 - b_2a_2,$$

$$\zeta_{-1}(a_1 - b_1)(a_1 - c_1) \text{ is equal to } a_1 - a_0c_0 + b_0c_0 - b_0a_0.$$

In like manner $\zeta PD(a, b, c)$ indicating

$$b_2a_1 - b_2c_1 + c_2b_1 - c_2a_1 + a_2c_1 - a_2b_1,$$

$\zeta_{\pm r} PD(a, b, c)$ indicates

$$b_{2\pm r}a_{1\pm r} - b_{2\pm r}c_{1\pm r} + c_{2\pm r}b_{1\pm r} - c_{2\pm r}a_{1\pm r} + a_{2\pm r}c_{1\pm r} - a_{2\pm r}b_{1\pm r}.$$

I shall in general denote $\zeta_{+r} PD(a, b, c \dots l)$ actually expanded as the zeta-ic product of $a, b, c, \dots l$ in its r th phase.

p. 49

Art. (7). *General Properties of Zeta-ic Products of Differences.*

If there be made one interchange in the order of the bases to which ζ is prefixed, the zeta-ic product, in whatever phase it be taken, remains unaltered in magnitude, but changes its sign.

If in any phase of a zeta-ic product two of the bases be made to coincide, the expansion vanishes.

Let f_1 be used, agreeably to the ordinary notation, to denote the sum of the quantities to which it is prefixed, f_2 to denote the sum of the binary products, f_3 of the ternary ones, and so on.

Thus let $f_1(a_1b_1c_1)$ or $f_1(a, b, c)$ indicate $a_1 + b_1 + c_1$,

and $f_2(a_1b_1c_1)$ or $f_2(a, b, c)$ indicate $a_1b_1 + a_1c_1 + b_1c_1$,

and $f_3(a_1b_1c_1)$ or $f_3(a, b, c)$ indicate $a_1b_1c_1$,

we shall be able now to state the following remarkable proposition connecting the several phases of certain the same zeta-ic products.

Art. (8). Let $a, b, c, \dots l$, denote any number of independent bases, say $(n - 1)$; but let the arguments of each base be periodic, and the number of terms in each period the same for every base, namely n , so that

$$\begin{aligned} a_r &= a_{r+n} = a_{r-n}, & a_n &= a_0 = a_{-n}, \\ b_r &= b_{r+n} = b_{r-n}, & b_n &= b_0 = b_{-n}, \\ c_r &= c_{r+n} = c_{r-n}, & c_n &= c_0 = c_{-n}, \\ &\dots\dots\dots, & & \\ l_r &= l_{r+n} = l_{r-n}, & l_n &= l_0 = l_{-n}, \end{aligned}$$

r being any number whatever. Then

$$\begin{aligned} \zeta_{-1}PD(0, a, b, c \dots l) &= \zeta\{f_1(a, b, c \dots l) \zeta PD(0, a, b, c \dots l)\}, \\ \zeta_{-2}PD(0, a, b, c \dots l) &= \zeta\{f_2(a, b, c \dots l) \zeta PD(0, a, b, c \dots l)\}, \\ &\dots\dots\dots, \\ \zeta_{-r}PD(0, a, b, c \dots l) &= \zeta\{f_r(a, b, c \dots l) \zeta PD(0, a, b, c \dots l)\}. \end{aligned}$$

This proposition admits of a great generalization²⁸, but we have now all that is requisite for enabling us to arrive at a proposition exhibiting under one coup d'oeil every combination and every effect of every combination that can possibly be made with any number of coexisting equations of the first degree, containing any number of repeated, or to use the ordinary language of analysts, (variable or) unknown quantities.

p. 50

For the sake of symmetry I make every equation homogeneous; so that to eliminate n repeated terms, no more than n equations will be required.

²⁸See the Postscript to this paper for one specimen.

In like manner the problem of determining n quantities from n equations will be here represented by the case in which we have to determine the ratios of $(n + 1)$ quantities from n equations.

Art. (9). *Statement of the Equations of Coexistence.*

Let there be any number of bases $(a, b, c \dots l)$, and as many repeated terms $(x, y, z \dots t)$, and let the number of equations be any whatever, say n . The system may be represented by the type equation

$$a_r x + b_r y + c_r z + \dots + l_r t = 0,$$

in which r can take up all integer values from $-\infty$ to $+\infty$. The specific number of equations given will be represented by making the arguments of each base periodic, so that

$$a_r = a_{\mu n + r}, \quad b_r = b_{\mu n + r}, \quad c_r = c_{\mu n + r}, \dots \quad l_r = l_{\mu n + r},$$

μ being any integer whatever.

Art. (10). *Combination of the given Equations.—Leading Theorem.*

Take $f, g, \dots k$ as the arbitrary bases of new and absolutely independent but periodic arguments, having the same index of periodicity (n) as $a, b, c \dots l$, and being in number $(n - 1)$, that is, one fewer than there are units in that index.

The number of differing arbitrary constants thus manufactured is $n(n - 1)$.

Let $Ax + By + Cz + \dots + Lt = 0$ be the general prime derivative from the given equations, then we may make

$$\begin{aligned} A &= \zeta PD(0, a, f, g \dots k), \\ B &= \zeta PD(0, b, f, g \dots k), \\ C &= \zeta PD(0, c, f, g \dots k), \\ &\dots\dots\dots \\ L &= \zeta PD(0, l, f, g \dots k). \end{aligned}$$

Art. (11). COR. 1.. *Inferences from the Leading Theorem.*

Let the number of equations, or, which is the same thing, the index of periodicity (n) , be the same as the number of repeated terms $(x, y, z \dots t)$, then one relation exists between the coefficients: this is found by making the $(n - 1)$ new bases coincide with $(n - 1)$ out of the old bases. We get accordingly, as the result of elimination,

$$\zeta PD(0, a, b, c \dots l) = 0.$$

p. 51

Art. (11). COR. 2.. Let the number of equations be one more than that of the given bases, there will then be two equations of condition. These are

represented by preserving one new arbitrary base, as λ . The result of elimination being in this case

$$\zeta PD(0, a, b, c \dots l, \lambda) = 0.$$

Example. The result of eliminating between

$$a_1x + b_1y = 0,$$

$$a_2x + b_2y = 0,$$

$$a_3x + b_3y = 0,$$

is $\zeta PD(0, a, b, \lambda) = 0$, that is

$$\lambda_3 b_2 a_1 - \lambda_3 b_1 a_2 + \lambda_1 b_3 a_2 - \lambda_1 b_2 a_3 + \lambda_2 b_1 a_3 - \lambda_2 b_3 a_1 = 0,$$

from which we infer, seeing that $\lambda_3, \lambda_2, \lambda_1$ are independent,

$$b_2 a_1 - b_1 a_2 = 0,$$

$$b_3 a_2 - b_2 a_3 = 0,$$

$$b_1 a_3 - b_3 a_1 = 0,$$

any *two* of which imply the third.

In like manner, in general, if the number of equations exceed in any manner the number of bases or repeated terms, the rule is to introduce so many new and arbitrary bases as together with the old bases shall make up the number of equations, and then equate the zeta-ic product of the differences of zero, the old bases and the new bases, to nothing.

Art. (12). COR. 3.. Let the number of equations be one fewer than the number (n) of bases or repeated terms; the number of introduced bases in the general theorem is here ($n - 2$). Make these ($n - 2$) bases equal severally to the bases which in the type equation are affixed to $z, u \dots t$, then

$$C = 0, \quad D = 0, \quad \dots, \quad L = 0,$$

and we have left simply

$$\zeta PD(0, a, c, d \dots kl)x + \zeta PD(0, b, c, d \dots kl)y = 0.$$

In like manner we may make to vanish all but A and C , and thus get

$$\zeta PD(0, a, b, d \dots kl)x + \zeta PD(0, c, b, d \dots kl)z = 0,$$

and similarly

$$\zeta PD(0, a, b \dots k)x + \zeta PD(0, b, c \dots l)t = 0.$$

Hence

$$\left\{ \begin{array}{c} x \\ y \\ z \\ . \\ . \\ t \end{array} \right\} \text{ are severally as } \left\{ \begin{array}{c} \zeta PD(0, b, c \dots l) \\ \zeta PD(a, 0, c \dots l) \\ \zeta PD(a, b, 0 \dots l) \\ \dots\dots\dots \\ \dots\dots\dots \\ \zeta PD(a, b, c \dots 0) \end{array} \right\}.$$

This is the symbolical representation as a formula of the remarkable method discovered by Cramer, perfected by Bezout and demonstrated by Laplace for the solution of simultaneous simple equations.

Art. (13). COR. 4.. In like manner if the number of repeated terms be two greater than the number of equations, we have for the relation between any three of them, taken at pleasure, for instance, x, y, z ,

$$\zeta PD(0, a, d \dots l)x + \zeta PD(0, b, d \dots l)y + \zeta PD(0, c, d \dots l)z = 0.$$

And in like manner we may proceed, however much in excess the number of repeated terms (unknown quantities) is over the number of equations.

Art. (14). *Subcorollary to Corollary 3.*

If there be any number of bases $(a, b, c \dots l)$, and any other two fewer in number $(f, g \dots k)$

$$\left. \begin{array}{l} \zeta PD(a, f, g \dots k) \times \zeta PD(b, c \dots l) \\ + \zeta PD(b, f, g \dots k) \times \zeta PD(a, c \dots l) \\ + \zeta PD(c, f, g \dots k) \times \zeta PD(b, c \dots l) \\ \dots\dots\dots \\ + \zeta PD(l, f, g \dots k) \times \zeta PD(a, b, c \dots) \end{array} \right\} = 0,$$

a formula that from its very nature suggests and proves a wide extension of itself²⁹.

In conclusion I feel myself bound to state that the principal substance of Corollaries (1), (2) and (3) may be found in Garnier's *Analyse Algébrique*, in the chapter headed "Développement de la Théorie donnée par M. Laplace, &c." But I am not aware of having been anticipated either in the fertile notation which serves to express them nor in the general theorems to which it has given birth.

P.S. I shall content myself for the present with barely enunciating a theorem, one of a class destined it seems to the author to play no secondary part in the development of some of the most curious and interesting points of analysis.

²⁹The cross is used to denote ordinary algebraical multiplication.

Let there be $(n - 1)$ bases $a, b, c \dots l$, and let the arguments of each be “recurrents of the n th order,”³⁰ that is to say let

$$a_t = \phi \left(\cos \frac{2\pi t}{n} \right), \quad b_t = \psi \left(\cos \frac{2\pi t}{n} \right), \quad c_t = \chi \left(\cos \frac{2\pi t}{n} \right), \dots \quad l_t = \omega \left(\cos \frac{2\pi t}{n} \right).$$

Let R_r denote that any symmetrical function of the r th degree is to be taken of the quantities in a parenthesis which come after it, and let $\mathfrak{S}$ indicate any function whatever. Then the zeta-ic product

$$\zeta \{ R_r(a, b, c \dots l) \times \mathfrak{S}_\rho \zeta PD(0, a, b, c \dots l) \}$$

is equal to the product of the number

$$R_r \left\{ \left(\cos \frac{2\pi}{n} + \sqrt{-1} \sin \frac{2\pi}{n} \right), \left(\cos \frac{4\pi}{n} + \sqrt{-1} \sin \frac{4\pi}{n} \right), \right. \\ \left. \left(\cos \frac{6\pi}{n} + \sqrt{-1} \sin \frac{6\pi}{n} \right), \dots, \left(\cos \frac{2(n-1)\pi}{n} + \sqrt{-1} \sin \frac{2(n-1)\pi}{n} \right) \right\},$$

multiplied by the zeta-ic phase

$$\zeta_{\rho-r} \mathfrak{S} PD(0, a, b, c \dots l)!!$$

³⁰I am indebted for this term to Professor De Morgan, whose pupil I may boast to have been. I have the sanction also of his authority, and that of another profound analyst, my colleague Mr Graves, for the use of the arbitrary terms zeta-ic, zeta-ically. I take this opportunity of retracting the symbol *SPD* used in my last paper, the letter *S* having no meaning except for English readers. I substitute for it *QDP*, where *Q* represents the Latin word *Quadratus*. On some future occasion I shall enlarge upon a new method of notation, whereby the language of analysis may be rendered much more expressive, depending essentially upon the use of similar figures inserted within one another, and containing numbers or letters, according as quantities or operations are to be denoted. This system to be carried out would require special but very simple printing types to be founded for the purpose. In the next part of this paper an easy and symmetrical mode will be given of representing any polynomial either in its developable or expanded form.

9.

A METHOD OF DETERMINING BY MERE INSPECTION THE DERIVATIVES FROM TWO EQUATIONS OF ANY DEGREE

[Philosophical Magazine, XVI. (1840), pp. 132–135]

p. 54

Let there be two equations, one of the n th, the other of the m th degree in x ; let the coefficients of the first equation be $a_n, a_{n-1}, a_{n-2} \dots a_0$, each power of x having a coefficient attached to it, a_n belonging to x^n and a_0 to the constant term.

In like manner let $b_m, b_{m-1} \dots b_0$ be the coefficients of the second equation.

I begin with

A Rule for absolutely eliminating x .

Form out of the (a) progression of coefficients m lines, and in like manner out of the (b) progression of coefficients form n lines in the following manner:

1. (a) . Attach $(m-1)$ zeros all to the *right* of the terms in the (a) progression; next attach $(m-2)$ zeros to the right and carry over to the left; next attach $(m-3)$ zeros to the right and carry over 2 to the left. Proceed in like manner until all the $(m-1)$ zeros are carried over to the left and none remain on the right.

The m lines thus formed are to be written under one another.

1. (b) Proceed in like manner to form n lines out of the (b) progression by scattering $(n-1)$ zeros between the right and left.

2. If we write these n lines under the m lines last obtained, we shall have a solid square $(m+n)$ terms deep and $(m+n)$ terms broad.

3. Denote the lines of this square by arbitrary characters, which write down in vertical order and permute in every possible way, but separate the permutations that can be derived from one another by an even number of interchanges (effected between contiguous terms) from the rest; there will thus be half of one kind and half of another.

p. 55

4. Now arrange the $(m+n)$ lines accordingly, so as to obtain

$$\frac{1}{2}\{(m+n)(m+n-1)\cdots 2\cdot 1\}$$

squares of one kind which shall be called positive squares, and an equal number of the opposite kind which shall be called negative.

Draw diagonals in the same direction in all the squares; multiply the coefficients that stand in any diagonal line together: take the sum of the diagonal products of the positive squares, and the sum of the diagonal products of the negative squares; the difference between these two sums is the prime derivative of the

zero degree, that is, is the result of elimination between the two given equations reduced to its ultimate state of simplicity, there will be no irrelevant factors to reject, and no terms which mutually destroy.

Example. To eliminate between

$$ax^2 + bx + c = 0, \quad lx^2 + mx + n = 0,$$

I write down

$$\begin{array}{llll} a, & b, & c, & 0, & (1) \\ 0, & a, & b, & c, & (2) \\ l, & m, & n, & 0, & (3) \\ 0, & l, & m, & n. & (4) \end{array}$$

I permute the four characters (1), (2), (3), (4), distinguishing them into positive and negative; thus I write together

Positive Permutations.

1	2	3	1	2	3	2	1	3	4	4	4
2	3	1	4	4	4	1	3	2	2	1	3
3	1	2	2	3	1	4	4	4	1	3	2
4	4	4	3	1	2	3	2	1	3	2	1

and again

Negative Permutations.

1	2	3	4	4	4	2	1	3	2	1	3
2	3	1	1	2	3	4	4	4	1	3	2
4	4	4	2	3	1	1	3	2	3	2	1
3	1	2	3	1	2	3	2	1	4	4	4

I reject from the permutations of each species all those where 1 or 3 appear in the fourth place, and also those where 2 or 4 appear in the first place, for these will be presently seen to give rise to diagonal products which are zero.

The permutations remaining are

Positive effectual permutations.

1	3	3	1
2	1	4	3
3	2	1	4
4	4	2	2

Negative effectual permutations.

3	1	1	3
1	4	3	2
4	3	2	1
2	2	4	4

I now accordingly form four positive squares, which are

a	b	c	0	l	m	n	0	l	m	n	0	a	b	c	0
0	a	b	c	a	b	c	0	0	l	m	n	l	m	n	0
l	m	n	0	0	a	b	c	a	b	c	0	0	l	m	n
0	l	m	n	0	l	m	n	0	a	b	c	0	a	b	c

Drawing diagonal lines from left to right, and taking the sum of the diagonal products, I obtain $a^2n^2 + lb^2n + l^2c^2 + am^2c$. Again, the four negative squares

l	m	n	0	a	b	c	0	a	b	c	0	l	m	n	0
a	b	c	0	0	l	m	n	l	m	n	0	0	a	b	c
0	l	m	n	l	m	n	0	0	a	b	c	a	b	c	0
0	a	b	c	0	a	b	c	0	l	m	n	0	l	m	n

give as the sum of the diagonal products

$$lbmc + alnc + ambn + lacn,$$

that is,

$$lbmc + ambn + 2acln.$$

Thus the result of eliminating between

$$ax^2 + bx + c = 0, \quad lx^2 + mx + n = 0,$$

ought to be, and is

$$a^2n^2 + l^2c^2 - 2acln + lb^2n + am^2c - lbmc - ambn = 0.$$

p. 57

Rule for finding the prime derivative of the first degree, which is of the form $Ax - B$.

Begin as before, only attach one zero less to each progression; we shall thus obtain not a square, but an oblong broader than it is deep, containing $(m + n - 2)$ rows, and $(m + n - 1)$ terms in each row: in a word, $(m + n - 2)$ rows, and $(m + n - 1)$ columns.

To find A reject the column at the extreme right, we thus recover a square arrangement $(m + n - 2)$ terms broad and deep.

Proceed with this new square as with the former one; the difference between the sums of the positive and negative diagonal products will give A .

To find B , do just the same thing, with the exception of striking off not the last column, but the last but *one*.

Rule for finding the prime derivative of any degree, say the r th, namely,

$$A_r x^r - A_{r-1} x^{r-1} + \cdots \pm A_0.$$

Begin with adding zeros as before, but the number to be added to the (a) progression is $(m - r)$ and to the (b) progression $(n - r)$.

There will thus be formed an oblong containing $(m + n - 2r)$ rows, and $(m + n - r)$ terms in each row, and therefore the same number of columns.

To find any coefficient as A_s , strike off all the last $(r + 1)$ columns except that which is (s) places distant from the extreme right, and proceed with the resulting squares as before.

Through the well-known ingenuity and kindly proffered help of a distinguished friend, I trust to be able to get a machine made for working Sturm's theorem, and indeed all problems of derivation, after the method here expounded; on which subject I have a great deal more yet to say, than can be inferred from this or my preceding papers.

10.

NOTE ON ELIMINATION

[*Philosophical Magazine*, XVII. (1840), pp. 379, 380]

p. 58

The object of this brief note is to generalise Theorem 2 in my paper on Elimination³¹ which appeared in the last December number of this *Magazine*. The theorem so generalised presents a symmetry which before was wanting. Here, as in so many other instances, the whole occupies in the memory a less space than the part.

To avoid the ill-looking and slippery negative symbols, I warn my reader that I now use two rows of quantities written one over the other, to denote the product of the terms resulting from taking away each quantity in the under from each in the upper row.

Let $h_1, h_2 \dots h_m$ be the roots of one equation of coexistence,

$k_1, k_2 \dots k_n$ of the other,

and let the prime derivative of the degree r be required. Take any two integers p and q , such that $p + q = r$. The derivative in question may be written

$$\Sigma \left\{ (x - h_1) \dots (x - h_p)(x - k_1) \dots (x - k_q) \frac{\begin{pmatrix} h_1 h_2 \dots h_p \\ k_1 k_2 \dots k_q \end{pmatrix} \cdot \begin{pmatrix} h_{p+1} h_{p+2} \dots h_m \\ k_{q+1} k_{q+2} \dots k_n \end{pmatrix}}{\begin{pmatrix} h_1 & h_2 & \dots & h_p \\ h_{p+1} h_{p+2} \dots h_m \end{pmatrix} \cdot \begin{pmatrix} k_1 & k_2 & \dots & k_q \\ k_{q+1} k_{q+2} \dots k_n \end{pmatrix}} \right\}.$$

N.B. Whatever p and q be taken, so long only as $p + q = r$, the above expression changes nothing but its sign; which, therefore, upon transcendental grounds, it is easy to see is of one name or another, according as p is odd or even.

In the original paper, I asserted this theorem only for the case of $p = 0$, or $q = 0$.

³¹p. 43 above. Ed.

11.

ON THE RELATION OF STURM'S AUXILIARY FUNCTIONS TO THE ROOTS OF AN ALGEBRAIC EQUATION

[*Plymouth British Association Report 1841, (Pt. II.), pp. 23, 24*]

p. 59

The author availed himself of the present meeting of the British Association to bring under the more general notice of mathematicians his discovery, made in the year 1839, of the real nature and constitution of the auxiliary functions (so-called) which Sturm makes use of in locating the roots of an equation: these are obtained by proceeding with the left-hand side of the equation and its first differential coefficient as if it were our object to obtain their greatest common factor; the successive remainders, with their signs alternately changed and preserved, constitute the functions in question. Each of these may be put under the form of a fraction, the denominator of which is a perfect square, or in fact the product of many: likewise the numerator contains a huge heap of factors of a similar form.

These therefore, as well as the denominator, since they cannot influence the series of signs, may be rejected; and furthermore we may, if we please, again make every other function, beginning from the last but one, change its sign, if we consent to use changes wherever Sturm speaks of continuations of sign, and *vice versâ*.

The functions of Sturm, thus modified and purged of irrelevancy, the author, by way of distinction, and still to attribute honour where it is really most due, proposes to call "Sturm's Determinators"; and he proceeds to lay bare the internal anatomy of these remarkable forms.

He uses the Greek letter " ζ " to indicate that the squared product of the differences of the letters before which it is prefixed is to be taken.

Let the roots of the equation be called respectively $a, b, c, e \dots l$, the determinators taken in the inverse order are as follows:

$$\zeta(a, b, c, e \dots l).$$

$$\Sigma\{\zeta(b, c, e \dots l)x - \Sigma a\zeta(b, c, e \dots l)\}.$$

$$\Sigma\{\zeta(c, e \dots l)x^2 - \Sigma(a + b)\zeta(c, e \dots l)x + \Sigma ab \cdot \zeta(c, e \dots l)\}.$$

$$\Sigma\{\zeta(k, l)(x - a)(x - b)(x - c)(x - e) \dots (x - h)\}.$$

p. 60

It may be here remarked, that the work of assigning the total number of real and of imaginary roots falls exclusively upon the coefficients of the leading terms, which the author proposes to call "Sturm's Superiors"; these superiors are only partial symmetric functions of the squared differences, but complete symmetric functions of the roots themselves, differing in the former respect from those other

(at first sight similar-looking) functions of the squared differences of the roots, in which, from the time of Waring downwards, the conditions of reality have been sought for. It seems to have escaped observation, that the series of terms constituting any one of the coefficients in the equation of the squares of the differences (with the exception of the first and last) each admit of being separated and classified into various subordinate groups in such a way, that instead of being treated as a single symmetric function of the roots, they ought to be viewed as aggregates of many. In fact, Sturm's superior No. 1 is identical with Waring's coefficient No. 1; Sturm's superior No. 2 is a part of Waring's coefficient No. 3; Sturm's superior No. 3 is a part of Waring's coefficient No. 6; and so forth till we come to Sturm's final superior, which is again coextensive and identical with the last coefficient in the equation of the squares of the differences. The theory of symmetric functions of forms which are themselves symmetric functions of simple letters, or even of other forms, the author states his belief is here for the first time shadowed forth, but would be beside his present object to enter further into. He would conclude by calling attention to the importance to the general interests of algebraical and arithmetical science that a searching investigation should be instituted for showing, *a priori*, how, when a set of quantities is known to be made up partly of possible and partly of pairs of impossible values, symmetrical functions of these, one less in number than the quantities themselves, may be formed, from the signs of the ratios of which to unity and to one another the respective amounts of possible and impossible quantities may at once be inferred; in short, we ought not to rest satisfied, until, from the very form of Sturm's Determinators, without caring to know how they may have been obtained, we are able to pronounce upon the uses to which they may be applied.

12.

EXAMPLES OF THE DIALYTIC METHOD OF ELIMINATION AS APPLIED TO TERNARY SYSTEMS OF EQUATIONS

[Cambridge Mathematical Journal, II. (1841), pp. 232–236]

p. 61

This method is of universal application, and at once enables us to reduce any case of elimination to the form of a problem, where that operation is to be effected between quantities linearly involved in the equations which contain them.

As applied to a binary system, $fx = 0$, $\phi x = 0$, the method furnishes a rule by which we may unfailingly arrive at the determinant, free from every species of irrelevancy, whether of a linear, factorial, or numerical kind.

The rule itself is given in the *Philosophical Magazine* (London and Edinburgh, Dec. 1840). The principle of the rule will be found correctly stated by Professor Richelot, of Königsberg, in a late number of Crelle's Journal, at the commencement of a memoir in Latin bordering on the same subject ("Nota ad Eliminationem pertinens").

My object at present is to supply a few instances of its application to ternary systems of equations.

Ex. 1. To eliminate x, y, z , between the three homogeneous equations

$$Ay^2 - 2C'xy + Bx^2 = 0, \quad (1)$$

$$Bz^2 - 2A'yz + Cy^2 = 0, \quad (2)$$

$$Cx^2 - 2B'zx + Az^2 = 0. \quad (3)$$

Multiply the equations in order by $-z^2, x^2, y^2$, add together, and divide out by $2xy$; we obtain

$$C'z^2 + Cxy - A'xz - B'yz = 0. \quad (4)$$

By similar processes we obtain

$$A'x^2 + Ayz - B'yx - C'zx = 0, \quad (5)$$

$$B'y^2 + Bzx - C'zy - A'xy = 0. \quad (6)$$

p. 62

Between these six, treated as simple equations, the six functions of x, y, z , namely, $x^2, y^2, z^2, xy, xz, yz$, treated as independent of each other, may be eliminated; the results may be seen, by mere inspection, to come out

$$ABC(ABC - AB'^2 - BC'^2 - CA'^2 + 2A'B'C') = 0,$$

or rejecting the special (N.B. not *irrelevant*) factor ABC , we obtain

$$ABC - AB'^2 - BC'^2 - CA'^2 + 2A'B'C' = 0.$$

I may remark, that the equations (1), (2), (3), or (4), (5), (6), express the condition of

$$Ax^2 + By^2 + Cz^2 + 2A'yz + 2B'zx + 2C'xy,$$

having a factor $\lambda x + \mu y + \nu z$; a general symbolical formula of which I am in possession for determining in general the condition of any polynomial of any degree having a factor, furnishes me at once with either of the two systems indifferently. The aversion I felt to reject *either*, led me to employ both, and thus was the occasion of the Dialectic Principle of Solution manifesting itself.

Ex. 2.

$$Ax^2 + ayz + bzx + cxy = 0, \quad (1)$$

$$My^2 + lyz + mzx + nxy = 0, \quad (2)$$

$$Rz^2 + pyz + qzx + rxy = 0. \quad (3)$$

Multiply equation (1) by $\beta y + \gamma z$, equations (2) and (3) by νz and κy respectively, and add the products together, we obtain terms of which y^2z and yz^2 are the only two into which x does not enter.

Make now the coefficients of each of these zero, and we have

$$a\gamma + l\nu + R\kappa = 0, \quad a\beta + M\nu + p\kappa = 0.$$

Let $\nu = a$, $\kappa = a$, then $\gamma = -(l + R)$, $\beta = -(M + p)$.

Hence, multiplying as directed, and then dividing out by x , we obtain

$$(m\nu + b\gamma)z^2 + (r\kappa + c\beta)y^2 + (b\beta + c\gamma + n\nu + q\kappa)yz + A\beta xy + A\gamma xz = 0,$$

or by substitution,

$$\{ra - c(M + p)\}y^2 + \{ma - b(l + R)\}z^2 + \{an + aq - b(M + p) - c(l + R)\}yz - A(M + p)xy - A(l + R)xz = 0. \quad (4)$$

Similarly, by preparing the equations so as to admit in turn of y and z as a divisor, we obtain

$$\{ma - l(R + b)\}z^2 + \{mr - n(A + q)\}x^2 + \{mc + mp - n(R + b) - l(A + \gamma)\}xz - M(R + b)yz - A(A + q)xy = 0, \quad (5)$$

$$\{rm - q(A + n)\}x^2 + \{ra - p(M + c)\}y^2 + \{rl + rb - p(A + n) - q(M + c)\}xy - R(A + n)xz - R(M + c)yz = 0. \quad (6)$$

Between the six equations (1), (2), (3), (4), (5), (6), $x^2, y^2, z^2, xy, xz, yz$, may be eliminated; the result will be a function of nine letters {three out of each

p. 63

equation (1), (2), (3)} equated to zero. *Perhaps* the determinant may be found to contain a special factor of three letters; and if so, may be replaced by a simpler function of six letters only.

Ex. 3. To eliminate between the three general equations

$$\begin{aligned} Ax^2 + By^2 + Cz^2 + 2Dyz + 2Ezx + 2Fxy &= 0, \\ Lx^2 + My^2 + Nz^2 + 2Pyz + 2Qzx + 2Rxy &= 0, \\ fx + gy + hz &= 0. \end{aligned}$$

By virtue of one of the two canons which limit the forms in which the letters can appear combined in the determinant of a general system of equations, we know that the determinant in this case (freed of irrelevant factors) ought to be made up in every term of eight letters (powers being counted as repetitions), namely, (A, B, C, D, E, F) must enter in binary combinations, (L, M, N, P, Q, R) the same, whereas f, g, h must enter in quaternary combinations.

To obtain the determinant, write

$$Ax^2 + By^2 + Cz^2 + Dyz + Ezx + Fxy = 0, \quad (1)$$

$$Lx^2 + My^2 + Nz^2 + Pyz + Qzx + Rxy = 0, \quad (2)$$

$$fx^2 + gyx + hzx = 0, \quad (3)$$

$$fxy + gy^2 + hzy = 0, \quad (4)$$

$$fxz + gyz + hz^2 = 0. \quad (5)$$

We want one equation more of *three letters* between $x^2, y^2, z^2, xy, xz, yz$. To obtain this, write

$$\begin{aligned} (Ax + Ez + Fy)x_1 + (By + Fx + Dz)y_1 + (Cz + Dy + Ex)z_1 &= 0, \\ (Lx + Qz + Ry)x_1 + (My + Rx + Pz)y_1 + (Nz + Py + Qx)z_1 &= 0, \\ fx_1 + gy_1 + hz_1 &= 0. \end{aligned}$$

Forget that $x_1 = x, y_1 = y, z_1 = z$, and eliminate x_1, y_1, z_1 , we obtain

$$\begin{aligned} &h \left\{ \begin{array}{l} (Ax + Ez + Fy)(My + Rx + Pz) \\ -(By + Fx + Dz)(Lx + Qz + Ry) \end{array} \right\} \\ &+ g \left\{ \begin{array}{l} (Cz + Dy + Ex)(Lx + Qz + Ry) \\ -(Nz + Py + Qx)(Ax + Ez + Fy) \end{array} \right\} \\ &+ f \left\{ \begin{array}{l} (Nz + Py + Qx)(By + Fx + Dz) \\ -(Cz + Dy + Ex)(My + Rx + Pz) \end{array} \right\} = 0. \end{aligned}$$

This may be put under the form

$$\alpha x^2 + \beta y^2 + \gamma z^2 + \alpha' yz + \beta' zx + \gamma' xy = 0, \quad (6)$$

where the coefficients are of the first order in respect to $f, g, h, L, M, N, P, Q, R, A, B, C, D, E$ in all of the third order.

Between the equations marked from (1) to (6), the process of linear elimination being gone through, we obtain as equated to zero a function of $5 + 3$, or of eight letters, two belonging to the first equation, two to the second, and four to the third; so that the determinant is clear of all factorial irrelevancy.

Ex. 4. To eliminate x, y, z between the three equations

$$\begin{aligned} Ax^2 + By^2 + Cz^2 + 2A'yz + 2B'zx + 2C'xy &= 0, \\ Lx^2 + My^2 + Nz^2 + 2L'yz + 2M'zx + 2N'xy &= 0, \\ Px^2 + Qy^2 + Rz^2 + 2P'yz + 2Q'zx + 2R'xy &= 0. \end{aligned}$$

Call these three equations $U = 0$, $V = 0$, $W = 0$, respectively. Write

$$\begin{aligned} xU &= 0, & yU &= 0, & zU &= 0, \\ xV &= 0, & yV &= 0, & zV &= 0, \\ xW &= 0, & yW &= 0, & zW &= 0. \end{aligned} \tag{1-9}$$

We have here nine unilateral equations: one more is wanted to enable us to eliminate linearly the ten quantities

$$x^3, y^3, z^3, x^2y, x^2z, xy^2, xz^2, xyz, y^2z, yz^2.$$

This tenth may be found by eliminating x, y, z between the three equations

$$\begin{aligned} x(Ax + B'z + C'y) + y(By + C'x + A'z) + z(Cz + A'y + B'x) &= 0, \\ x(Lx + M'z + N'y) + y(My + N'x + L'z) + z(Nz + L'y + M'x) &= 0, \\ x(Px + Q'z + R'y) + y(Qy + R'x + P'z) + z(Rz + P'y + Q'x) &= 0; \end{aligned}$$

for, by forgetting the relations between the bracketed and unbracketed letters, we obtain

$$(Ax + B'z + C'y) \left\{ \begin{array}{l} (My + N'x + L'z)(Rz + P'y + Q'x) \\ - (Qy + R'x + P'z)(Nz + L'y + M'x) \end{array} \right\} + \&c. + \&c. = 0,$$

which may be put under the form

$$\alpha x^3 + \beta y^3 + \gamma z^3 + \delta x^2y + \dots = 0^*. \tag{10}$$

p. 65

³¹We might dispense with a 10th equation, using the nine above given, to determine the ratios of the ten quantities involved to one another; and then by means of any such relations as $x^3y \times xy^3 = x^2y^2 \times x^2y^2$, or $x^3 \times y^3 = x^2y \times xy^2$, &c. obtain a determinant. But it is easy to see that this would be made up of terms, each containing literal combinations of the 18th order. Again, we might use five out of the nine equations to obtain a new equation free from y^3, y^2z, yz^2, z^3 ; that is, containing x in every term: which being divided by x , and multiplied

By eliminating linearly between the equations marked from (1) to (10), we obtain as zero a quantity of the twelfth order in all, being of the fourth order in respect to the coefficients of each of the three equations, which is therefore the determinant in its simplest form.

I have purposely, in this brief paper, avoided discussing any theoretical question. I may take some other opportunity of enlarging upon several points which have hitherto been little considered in the theory of elimination, such as the Canons of Form, the Doctrine of Special Factors, the Method of Multipliers as extended to a system of any order, the Connexion between the Method of Multipliers and the Dialytic Process, the Idea of Derivations and of Prime Derivatives extended to ultra-binary Systems. For the present I conclude with the expression of my best wishes for the continued success of this valuable Journal.

³¹by y , or by z , would furnish a 10th equation no longer linearly involved in the 9 already found. The determinant, however, found in this way, would consist of 14-ary combinations of letters. Finally, we might, instead of a system of ten equations, employ a system of 15, obtained by multiplying each of the given three by any 5 out of the 6 quantities $x^2, y^2, z^2, xy, xz, yz$; but the determinant, besides being not totally symmetrical, would contain combinations of the 15th order. I may take this opportunity of just adverting to the fact, that the method in the text does in fact contain a solution of the equation

$$x^r \lambda + y^s \mu + z^t \nu = 0,$$

where $r + s + t = 1$, and λ, μ, ν are functions of the second degree in regard to x, y, z to be determined.

13.

INTRODUCTION TO AN ESSAY ON THE AMOUNT AND DISTRIBUTION OF THE MULTIPLICITY OF THE ROOTS OF AN ALGEBRAIC EQUATION

[*Philosophical Magazine*, XVIII. (1841), pp. 136–139]

p. 66

I use the word *multiplicity* to denote a number, and distinguish between the total and partial multiplicities of the roots of an algebraic equation.

There may be r different roots repeated respectively $h_1, h_2 \dots h_r$ times.

r is the index of distribution.

$h_1, h_2 \dots h_r$ are the partial multiplicities, and if $h = h_1 + h_2 + \dots + h_r$, h is the *total* multiplicity.

The total multiplicity it is clear may be defined as the difference between the index of the equation and the number of its roots distinguishable from one another.

In this Introduction, I propose merely to consider how existing methods may be applied to determine the amount and distribution of multiplicity in a given equation, and conversely, how equations of condition can be formed which shall imply a given distribution and amount.

Let the greatest common factor between fx (the argument of the proposed equation) and $\frac{dfx}{dx}$ be called f_1x .

And in like manner, let the greatest common factor of f_1x and $\frac{df_1x}{dx}$ be called f_2x , and so on, till in the end we come to f_rx , which has no common factor with $\frac{df_rx}{dx}$.

Let $k_1, k_2 \dots k_r$ denote the degrees in x of $fx, f_1x \dots f_rx$ respectively.

It is easy to see that

$k_1 - k_2$, partial multiplicities, are less than 2, that is, are each units,
 $k_2 - k_3$, partial multiplicities, will be less than 3, and therefore either 1 or 2,
in value respectively, and so on till we come to
 $k_{r-1} - k_r$, which will severally be between zero and $r - 1$, and
 $k_r - 0$, of values intermediate between zero and r .

p. 67

Hence there will be

$k_1 - 2k_2 + k_3$	multiplicities each of the value	1,
$k_2 - 2k_3 + k_4$		2,
.....		
$k_{r-1} - 2k_r$	of the value	$r - 1$,
k_r	of the value	r .

In place of fx with $\frac{dfx}{dx}$ we might employ $\frac{dfx}{dx}$ with $\frac{d^2fx}{dx^2}$ and so on for the rest; the values of $k_2, k_3 \dots k_r$ will remain unaffected by this change; but the former method would be more expeditious in practice.

The total multiplicity is, of course, $= k_1$.

Suppose now that we propose to ourselves the converse problem to determine the conditions that an algebraic equation may have a given amount of multiplicity distributed in a given manner.

If $h_1, h_2, h_3 \dots h_r$ be used to denote the given number of partial multiplicities which are respectively of the values $1, 2, 3 \dots r$, it is easy to see that the quantities derived above by $k_1, k_2 \dots k_r$ are respectively equal to

$$\begin{aligned} &h_1 + 2h_2 + \dots + rh_r, \\ &h_2 + 2h_3 + \dots + rh_{r-1}, \\ &h_3 + 2h_4 + \dots + rh_{r-2}, \\ &\dots\dots\dots \\ &h_r. \end{aligned}$$

Now from $\frac{dfx}{dx}$ having a factor of the degree k_1 common with fx we obtain k_1 conditions, from $\frac{df_1x}{dx}$ having a factor of the degree k_2 common with f_1x we obtain k_2 more, and so on. So that altogether we obtain in this way

$$k_1 + k_2 + \dots + k_r \text{ conditions.}$$

But it may easily be seen that the total multiplicity being k_1 , the number of conditions need never to exceed k_1 in number, no matter what its distribution may be. Hence, besides the enormous labour of the process, and the extreme complexity of the results, we obtain by this method more equations by far than are necessary, and it requires some caution to know which to reject.

In my forthcoming paper (to appear in *Philosophical Magazine* of next month) I shall show, by a most simple means, how without the use of derived or other subsidiary functions, to obtain the simplest equations of condition which correspond to a given distribution of a given amount of multiplicity.

The total multiplicity, say m , being given in as many ways as that number can be broken into parts, so many different systems of m equations can be formed differing each from the other in the dimensions of the terms.

p. 68

These systems may be arranged in order so that each in the series shall imply all those that follow it, and be implied in all those that go before, without the converse being satisfied.

The subject of the unreciprocal implication of systems of equations is a very curious one, upon which the limits assigned to me prevent me from enlarging at present. It is closely connected with a part of the theory of elimination, which, as

far as I am aware, has either been overlooked, or has not met with the attention which it deserves; I mean the theory of *Special Factors*.

An *example* may make what I mean by these clear.

Let C be a function (if my reader please) void of x , which equivalent to zero implies two given equations in x having a common root.

Let C be rid of all irrelevant factors, that is, let C be the simplest form of the determinant, when the coefficients of the two equations are perfectly independent qualities. Now suppose, as is quite possible in a variety of ways, that such relations are instituted between the coefficients alluded to as make C split up into factors, so that $C = L \times M \times N = 0$.

Only one of the factors L, M, N will satisfy the condition of the coexistence of the two given equations: the others are clearly, however, not to be confounded with factors of solution, or irrelevant factors, as they are termed, but are of quite a different nature, and enjoy remarkable properties, which point to an enlarged theory of elimination, and constitute what I call special or singular factors.

I shall feel much obliged to any of the readers of your widely circulated Journal, interested in the subject of this paper, who would do me the honour of communicating with me upon it, and especially if they would (between now and the next coming out of the *Magazine*) inform me whether anything, and if so how much, different from what is here stated has been done in the matter of determining the relations between the coefficients of an equation corresponding to a given amount and distribution of multiplicity in its roots.

I ought to add, that my method enables me not merely to determine the conditions of multiplicity, but also to decompose the equations containing multiple roots into others free of multiplicity, that is, to find, *à priori*, the values of the several quantities

$$\frac{fx f_2x}{(f_1x)^2}, \quad \frac{f_1x f_3x}{(f_2x)^2}, \quad \dots, \quad \frac{f_{r-1}x}{(f_rx)^2}, \quad f_rx.$$

Moreover, other decompositions, not necessary to be enlarged upon in this place, may be obtained with equal facility.

14.

A NEW AND MORE GENERAL THEORY OF MULTIPLE ROOTS

[*Philosophical Magazine, XVIII. (1841), pp. 249-254*]

p. 69

I shall begin with developing the theory of polynomials containing perfect square factors, one or more.

First, let us proceed to determine the relations which must exist between the coefficients of such polynomials, and afterwards show how they may be broken up into others of an inferior degree.

A parallelogram filled with letters standing in one row is intended to express the product of the squared difference of the quantities contained. Thus $\overline{ab}$ indicates $(a - b)^2$, $\overline{abc}$ is used to indicate $(a - b)^2(a - c)^2(b - c)^2$, and so forth.

Suppose now that two of the roots $e_1, e_2 \dots e_n$ belonging to the equation $fx = 0$ are equal to one another, it is clear that $\overline{e_1, e_2 \dots e_n} = 0$; and moreover is a symmetric function, and can be calculated in terms of the coefficients of fx .

Next let us suppose that we have two couples of equals (as for instance a and b , two of the roots equal, as also c and d two others), it is clear, that on leaving any one of the roots out, the $(n - 1)$ that are left will still contain one equality, and therefore we have

$$\overline{e_2, e_3 \dots e_n} = 0, \quad \overline{e_1, e_3 \dots e_n} = 0 \dots \overline{e_1, e_2 \dots e_{n-1}} = 0.$$

None of the parallelogrammatic functions above taken singly, are symmetric functions of the coefficients, but their sum is; so also is the sum of the product of each into the quantity left out.

Now in general, suppose that the polynomial fx contains r perfect square factors, so that we have r couples of equal roots belonging to the equation $fx = 0$, it is clear that $\overline{e_r, e_{r+1} \dots e_n}$ and all the other

$$\frac{n(n-1) \dots (n-r+2)}{1 \cdot 2 \dots (r-1)}$$

functions of which it is the type are severally zero. Moreover, the sum of

p. 70

these or the sum of the products of each by any symmetrical function of the $(r - 1)$ letters left out will be a symmetrical function of the coefficients of the powers of x in fx . To express now the affirmative³² conditions corresponding to the case of there being r pairs of equal roots, we might employ the r equations,

$$\begin{aligned} \overline{e_1, e_2 \dots e_n} &= 0, \\ \overline{\Sigma e_2, e_3 \dots e_n} &= 0, \\ &\dots\dots\dots \\ \overline{\Sigma e_r, e_{r+1} \dots e_n} &= 0. \end{aligned}$$

³²The importance of the restriction hinted at by the use of the word affirmative will appear hereafter.

But these, except the last, are not the simplest that can be employed; that is to say, we can write down r others, the terms of which shall be of lower dimensions in respect to the roots.

Let f_μ denote that any rational symmetrical function of the μ th degree is to be taken of the quantities which it precedes.

Then the r equations in question are all contained in the general equation

$$\Sigma\{f_\mu(e_1, e_2 \dots e_{r-1}) \times \overline{e_r, e_{r+1} \dots e_n}\} = 0.$$

μ being taken from 0 up to $(r - 1)$ we obtain r equations, which in respect to the roots are respectively of all degrees between

$$\frac{n(n-1) \dots (n-r+2)}{1 \cdot 2 \dots (r-1)} \quad \text{and} \quad \frac{n(n-1) \dots (n-r+2)}{1 \cdot 2 \dots (r-1)} + (r-1)$$

reckoned inclusively.

Now at this stage it is important to remark that the above r equations, although necessary, are not *sufficient*; and indeed, no mere affirmations of equality can be sufficient to ensure there being r pairs of equal roots.

To make this manifest, suppose $r = 2$. Then in order that an equation may have two pairs of equal roots, we must have by the above formula

$$\Sigma \overline{e_2, e_3 \dots e_n} = 0, \quad \Sigma\{e_1 \overline{e_2, e_3 \dots e_n}\} = 0.$$

But if instead of there being two perfect square factors there be one perfect cube factor in fx , it may be shown by the same reasoning as above, that the very same two equations apply. In fact, it may be shown in general that no such equations as those given above can be affirmed in consequence of there being an amount r of multiplicity consisting of unit parts which may not be affirmed with equal truth as necessary consequences of the same

p. 71

amount distributed in any other manner whatever. How to obtain affirmative equations sufficient as well as necessary (under certain limitations) will appear at the close of this present paper.

It is worthy of being remarked, that if we make f_μ denote the sum of the products of the quantities to which it is prefixed, taken μ and μ together, the equations of affirmation become identical with those obtained by eliminating between fx and $\frac{dfx}{dx}$ ³³.

It can scarcely be doubted that the illustrious Lagrange, had he chosen to perfect the incomplete theory of equal roots given in the *Résolution Numérique*, by applying to it his own favourite engine of symmetric functions, could scarcely have failed of stumbling by a back passage upon Sturm's memorable theorem.

³³See my note on Sturm's Theorem, *Phil. Mag.*, December, 1839 [p. 45 above. Ed.].

Let us now proceed to show how a polynomial known to contain one or more perfect square factors may be decomposed.

Let us begin with supposing that it contains but one such factor; so that $fx = \phi x(x - a)^2$.

I shall show how to obtain the equations

$$C(x - a) = 0, \quad D\phi x(x - a) = 0, \quad E(x - a)^2 = 0, \quad F(\phi x) = 0,$$

each in its lowest terms.

1. To form the equation $Lx + M = 0$, where $x = a$, it is easy to see that if we write down in general the expression $(x - e_1)\overline{e_2, e_3 \dots e_n}$ this will become zero whenever the root e_1 left out is not one of the equal roots (a); so that in fact (calling the two equal roots e_1, e_2 respectively)

$$\Sigma\{(x - e_1)\overline{e_2, e_3 \dots e_n}\} = (x - e_1)\overline{e_2, e_3 \dots e_n} + (x - e_2)\overline{e_1, e_3 \dots e_n},$$

or simply

$$= 2(x - a)\overline{e_2, e_3 \dots e_n}.$$

Hence by making

$$x\Sigma\overline{e_2, e_3 \dots e_n} - \Sigma\{e_1\overline{e_2, e_3 \dots e_n}\} = 0,$$

we have an equation for finding the equal roots e_1, e_2 .

Again, it is easily seen upon the same hypothesis, that

$$\Sigma\{(x - e_2)(x - e_3) \dots (x - e_n)\overline{e_2, e_3 \dots e_n}\} = 2(x - e_2)(x - e_3) \dots (x - e_n)\overline{e_2, e_3 \dots e_n}.$$

p. 72

Hence, to form the equation having the same roots as $(x - a)\phi x$, we have only to make

$$\begin{aligned} x^{n-1}\Sigma\overline{e_2, e_3 \dots e_n} - x^{n-2}\Sigma\{(e_2 + e_3 + \dots + e_n)\overline{e_2, e_3 \dots e_n}\} + \dots \\ \pm \Sigma\{(e_2 e_3 \dots e_n)\overline{e_2, e_3 \dots e_n}\} = 0. \end{aligned}$$

Suppose now in general that we have r perfect square factors, so that

$$fx = \phi x(x - a_1)^2(x - a_2)^2 \dots (x - a_r)^2.$$

To form the equation $G(x - a_1)(x - a_2) \dots (x - a_r) = 0$, we have only to make

$$\Sigma\{(x - e_1)(x - e_2) \dots (x - e_r)\overline{e_{r+1}, e_{r+2} \dots e_n}\} = 0.$$

And to obtain

$$D\phi x \times (x - a_1)(x - a_2) \dots (x - a_r) = 0,$$

we must make

$$\Sigma\{(x - e_{r+1})(x - e_{r+2}) \cdots (x - e_n) \overline{e_{r+1}, e_{r+2} \dots e_n}\} = 0.$$

The theory of perfect square factors is not yet complete until it has been shown how to obtain constructively ϕx , and, as analogy suggests, the complementary part $D(x - a_1)^2(x - a_2)^2 \cdots (x - a_r)^2$, each in its lowest terms. To effect the latter it might be said that it is only necessary to take the square of $C(x - a_1)(x - a_2) \cdots (x - a_r)$. It is true the polynomial so formed would contain every pair of equal factors, but not in the lowest terms as regards the coefficients (as we shall presently show).

To solve this last part of the problem, let it be agreed that two rows of letters inclosed in a parenthesis shall indicate the product of the squares of the differences got by subtracting each in the row from each in the other, so that

$$\begin{pmatrix} a \\ b \end{pmatrix} = (a-b)^2, \quad \begin{pmatrix} a \\ b, c \end{pmatrix} = (a-b)^2(a-c)^2, \quad \begin{pmatrix} a, b \\ c, d \end{pmatrix} = (a-c)^2(a-d)^2(b-c)^2(b-d)^2.$$

Let us begin with supposing that fx has one pair only of equal roots; to form the simplest quadratic equation containing this pair, write down

$$\Sigma \left\{ (x - e_1)(x - e_2) \overline{e_3, e_4 \dots e_n} \begin{pmatrix} e_1, e_2 \\ e_3, e_4 \dots e_n \end{pmatrix} \right\}.$$

Now if e_1 and e_2 are the two equal roots in question neither of the multipliers of $(x - e_1)(x - e_2)$ vanishes.

If e_1 and e_2 are neither of them equal roots $\overline{e_3, e_4 \dots e_n} = 0$.

If one of the two only belong to the pair of equal roots

$$\begin{pmatrix} e_1, e_2 \\ e_3, e_4 \dots e_n \end{pmatrix} = 0.$$

Hence it is clear that

$$\Sigma \left\{ (x - e_1)(x - e_2) \overline{e_3, e_4 \dots e_n} \begin{pmatrix} e_1, e_2 \\ e_3, e_4 \dots e_n \end{pmatrix} \right\} = 0$$

is the equation desired.

In like manner if there be r pairs of equal roots the equation of the $(2r)$ th degree which contains them all may be written

$$\Sigma \left\{ (x - e_1)(x - e_2) \cdots (x - e_{2r}) \overline{e_{2r+1} \dots e_n} \begin{pmatrix} e_1, e_2, \dots, e_{2r} \\ e_{2r+1}, \dots, e_n \end{pmatrix} \right\} = 0.$$

The coefficient of x^{2r} in this equation is clearly of

$$(n - 2r)(n - 2r - 1) + 4r(n - 2r),$$

that is, of $(n + 2r - 1)(n - 2r)$ dimensions. The coefficient of x^r in the equation which contains the r equal roots unyoked together is of $(n - r)(n - r - 1)$ dimensions, and consequently the coefficient of x^{2r} in the square of this equation would be of $2(n - r)(n - r - 1)$ dimensions, that is, would be $n^2 + 6r^2 - (4r + 1)n$ dimensions higher than needful.

Finally, to obtain an equation clear of *simple* as well as double appearances of the equal roots, we have only to write the complementary form

$$\Sigma \left\{ (x - e_{2r+1})(x - e_{2r+2}) \cdots (x - e_n) \overline{e_{2r+1} \cdots e_n} \begin{pmatrix} e_1, e_2, \dots, e_{2r} \\ e_{r+1}, \dots, e_n \end{pmatrix} \right\} = 0.$$

Let us, now that we are more familiarized with the notation essential to this method, revert to the question with which we set out, and endeavour to obtain r such equations as shall imply unambiguously the existence of r pairs of equal roots.

The existence of r such pairs enables us to assert the following disjunctive proposition, which cannot be asserted when the same amount of multiplicity is distributed in any other way.

To wit, on selecting any r roots out of the entire number, either these r will all be found again in those that are left, or those that are left will contain *inter se*, one repetition at least; so that except on the latter supposition any $(r - 1)$ may be absolutely sunk out of those that are left, and there will still be one root common to the $(n - 2r + 1)$ remaining, and to the r originally selected to be left out.

Wherefore calling the roots $e_1, e_2 \dots e_n$, and giving μ any value whatever, we have

$$\Sigma \left\{ f_\mu(e_1, e_2 \dots e_r) \times \overline{e_{r+1}, e_{r+2} \dots e_n} \times \Sigma \begin{pmatrix} e_1, e_2 \dots e_r \\ e_{2r}, e_{r+1} \dots e_n \end{pmatrix} \right\} = 0.$$

p. 74

Hence the simplest distinctive equations indicative of the existence of r pairs of equal roots are to be found by putting μ equal in succession to all values from 0 up to $(r - 1)$.

For instance, if we require that an equation of the seventh degree shall have three pairs of equal roots, we need only to call the seven roots respectively a, b, c, d, e, f, g , and then our type equation becomes

$$\Sigma \left\{ f_\mu(a, b, c) \times \overline{d, e, f, g} \times \left[\begin{pmatrix} d, e \\ a, b, c \end{pmatrix} + \begin{pmatrix} d, f \\ a, b, c \end{pmatrix} + \begin{pmatrix} d, g \\ a, b, c \end{pmatrix} \right. \right. \\ \left. \left. + \begin{pmatrix} e, f \\ a, b, c \end{pmatrix} + \begin{pmatrix} e, g \\ a, b, c \end{pmatrix} + \begin{pmatrix} f, g \\ a, b, c \end{pmatrix} \right] \right\} = 0.$$

From this it appears that the r distinctive equations for r pairs of equal roots are of different dimensions from the r general or overlying ones corresponding to

the multiples r , anyhow distributed; the lowest of the latter being of $(n - r + 1)(n - r)$, the lowest of the former of

$$(n - r)(n - r - 1) + 2r(n - 2r + 1),$$

that is, of $n(n - 1) - 3r(n - 1)$ dimensions. In general we shall find that the more unequally distributed the multiplicity may be the lower are the dimensions of the distinctive equations, and are accordingly lowest when the multiplicity is absolutely undistributed³⁴.

³⁴It must not, however, be overlooked, that the equations above given, although decisive as to the existence of r pairs of equal roots when the multiplicity is known to be not greater than r , do not enable us to affirm with certainty their existence when this limitation is absent; for should the multiplicity exceed r , then inevitably (no matter how it may be distributed) $\overline{e_{r+1}, e_{r+2} \dots e_n}$ is always zero, and consequently nullifies each term of every one of the equations in question. In fact (repugnant as it may appear to be to the ordinary assumptions of analytical reasoning), it is not possible to express with absolute unambiguity the conditions of there being a multiplicity (r) distributed in any assigned manner by means of r affirmative equations alone.

15.

ON A LINEAR METHOD OF ELIMINATING BETWEEN DOUBLE, TREBLE, AND OTHER SYSTEMS OF ALGEBRAIC EQUATIONS

[*Philosophical Magazine, XVIII. (1841), pp. 425-435*]

p. 75

PART I. BINARY SYSTEMS.

Let U and V be two integer complete homogeneous functions of x and y , one of the m th, the other of the n th degree; and let it be required to express the condition of the coexistence of the two equations $U = 0$, $V = 0$ by means of the equation $C = 0$, where C is free from all appearances of x or y .

This equation, according to the system of notation developed in a preceding paper, and which has been since adopted and sanctioned by the high authority of M. Cauchy, I call the final derivative: the quantity C is designated the final derivee: and it is our present object to show how this may be obtained in a *prime* form, that is to say, divested of irrelevant factors: in this state it must consist of terms, each containing $m + n$ letters, of which n belong to the coefficients of U , and m to those of V .

Of course in applying this rule it is to be understood that every combination of powers in U or V has a single letter prefixed for its coefficient, and that in the final derivee powers are represented by repetitions of the same character.

Every term in U or V being of the form Cx^py^q , x^py^q is called an argument, C its prefix.

Assume two integer positive numbers r and r' , and also two others s and s' , such that $r + r' = n - 1$, $s + s' = m - 1$, and form from $U = 0$, $V = 0$ two new equations,

$$x^ry^{r'}U = 0, \quad x^sy^{s'}V = 0.$$

Such equations are termed the augmentatives of the two given ones respectively; also $x^ry^{r'}U$ and its fellow are termed the augmentees of U and V .

p. 76

r and r' are termed the indices of augmentation belonging to U , s and s' the same belonging to V .

Finally, it will be useful hereafter to call the given polynomials U and V themselves the proposees, and the given equations which assert their nullity, the propositive equations, or, briefly, the propositives.

Now as many augmentees of either proposee can be formed as there are ways of stowing away between two lockers (vacancies admissible) a number of things equal to the index of the other³⁵; hence we shall have n augmentees of U , and m

³⁵“Tot augmenta utriusvis ex æquationibus propositis formari possunt quot modi sint inter duo receptacula (utriusvis vel ambobus omnino vacare licet) rerum, quarum numerus indicem alterius æquat, distributionem faciendi.”

of V : thus there will be $m + n$ augmentatives each of the degree $m + n - 1$, and the number of arguments is clearly $m + n$ also, so that they can be eliminated linearly, and the final derivee thus found, containing $m + n$ letters (properly aggregated) in each term, will be in its prime form, that is, incapable of further reduction, and void of irrelevant factors.

It is worthy of remark, that the final derivee obtained by arranging in square battalion the prefixes of the augmentees, permuting the rows or columns, and reading off diagonal products, affected each with the proper sign (according to the well known rule of Duality), will not only be free from factorial irrelevancy, but also of linear redundancy, which latter term I use to signify the reappearance of the same combination of prefixes, sometimes with positive and sometimes with negative signs: furthermore, it follows obviously from the nature of the process that no numerical quantity in the final derivee will be greater than the higher of the indices of the two given polynomials.

PART II. TERNARY SYSTEMS.

CASE A. INDICES ALL EQUAL.

Method 1.

Let there be now three proposees, U, V, W , integer complete homogeneous functions of x, y, z , each of the degree n : let

$$r + r' + r'' = n - 1, \quad s + s' + s'' = n - 1, \quad t + t' + t'' = n - 1,$$

$$x^r y^{r'} z^{r''} U, \quad x^s y^{s'} z^{s''} V, \quad x^t y^{t'} z^{t''} W,$$

will, as above, be called the augmentees of U, V, W , and every other part of the notation previously described is to be preserved.

p. 77

Suppose now

$$U = 0, \quad V = 0, \quad W = 0,$$

we shall have as many augmentative equations formed from each proposee as there are ways of stowing away n things between three lockers (vacancies admissible)³⁶, that is, $\frac{n(n+1)}{2}$ of each kind; in all, therefore, $3\frac{n(n+1)}{2}$, and every one of these will be of the degree $2n - 1$, so that the number of arguments to be eliminated is equal to the number of ways of stowing away $2n - 1$ things between three lockers (empty ones counting), that is

$$\frac{2n(2n+1)}{2}.$$

³⁶See for Latin translation the preceding note.

As yet, then, we have not *enough* equations for eliminating these linearly.

Make, however,

$$\alpha + \beta + \gamma = n + 1,$$

and write

$$U = x^\alpha F + y^\beta F' + z^\gamma F'' = 0,$$

$$V = x^\alpha G + y^\beta G' + z^\gamma G'' = 0,$$

$$W = x^\alpha H + y^\beta H' + z^\gamma H'' = 0,$$

it will always be possible to make the multipliers of x^α , y^β , z^γ integer functions: for if we look to any argument in U , V , or W , it is of the form $x^a y^b z^c$, and one of the letters a, b, c must be *not less* than its correspondent α, β, γ , for otherwise $a + b + c$ would be not greater than $\alpha + \beta + \gamma - 3$, that is, n would be not greater than $(n + 1) - 3$, or $n - 2$, which is absurd: if now any one, as a , be equal to or greater than α , it may be made to supply an integer part to the multiplier of x^α .

Here it may be asked what is to be done with such terms as $Kx^a y^b z^c$, when two letters a, b are each not less than their correspondents α, β : the answer is, such terms may be made to enter under the multiplier of x^α , or of y^β , or to supply a part to both in any proportion at pleasure³⁷.

From the equations above we get, by linear elimination,

$$FG'H'' + GH'F'' + HF'G'' - GF'H'' - HG'F'' - FH'G'' = 0.$$

This may be denoted thus: $\Pi(\alpha, \beta, \gamma) = 0$, which equation I call a secondary derivative, and the left side of it a secondary derivate; α, β, γ may likewise be termed the indices of derivation (as r, s, t , &c. are of augmentation).

Now since $\alpha + \beta + \gamma = n + 1$, it is clear that the index of $\Pi(\alpha, \beta, \gamma)$ is always $n + n + n - (n + 1)$; that is, $2n - 1$.

p. 78

1st. Let any two of the indices of derivation be taken zero, then it is easily seen that all the terms in $\Pi(\alpha, \beta, \gamma)$ vanish, and consequently the secondary derivative equations obtained upon this hypothesis become mere identities, and are of no use.

2nd. Let any *one* of them become zero.

It is manifest, from the doctrine of simple equations, that $\Pi(\alpha, \beta, \gamma)$ may be made equal to

$$\{\lambda U + \mu V + \nu W\} \frac{1}{x^\alpha},$$

or

$$\{\lambda' U + \mu' V + \nu' W\} \frac{1}{y^\beta},$$

or

$$\{\lambda'' U + \mu'' V + \nu'' W\} \frac{1}{z^\gamma},$$

³⁷The prefixes of any such terms (say K) may be conceived as made up of two parts, an arbitrary constant, as e and $(K - e)$; e will disappear spontaneously from the final derivate.

upon the understanding that

$$\begin{aligned}\lambda &= G'H'' - G''H', & \mu &= H'F'' - H''F', & \nu &= F'G'' - F''G', \\ \lambda' &= G''H - GH'', & \mu' &= H''F - HF'', & \nu' &= F''G - FG'', \\ \lambda'' &= GH' - G'H, & \mu'' &= HF' - H'F, & \nu'' &= FG' - F'G.\end{aligned}$$

The three rows of coefficients will be respectively of the degrees

$$(n - \beta) + (n - \gamma), \quad (n - \gamma) + (n - \alpha), \quad (n - \alpha) + (n - \beta).$$

Thus if any one of the indices α, β, γ be zero, $\Pi(\alpha, \beta, \gamma)$ becomes identical with $\lambda'U + \mu'V + \nu'W$, where the multipliers of U, V, W are of $2n - (\alpha + \beta + \gamma)$ dimensions, that is of $(n - 1)$ dimensions, and may accordingly be put under the form

$$\Sigma A x^r y^{r'} z^{r''} U + \Sigma B x^s y^{s'} z^{s''} V + \Sigma C x^t y^{t'} z^{t''} W,$$

that is to say, becomes a *linear function* of the augmentatives, and therefore if combined with them in the process of linear elimination would give rise to the identity $0 = 0$.

Hence we must reject all such secondary derivatives as have zero for one of the indices of derivation. But all others, it may be shown, will be *linearly* independent of one another, and of the augmentees previously found. Hence, besides $3 \frac{n(n+1)}{2}$ equations of augment of the degree $2n - 1$, we shall have of the same degree so many equations of derivation as there are ways of stowing away between three lockers $(n + 1)$ things, under the condition that no locker shall ever be left empty, that is $\frac{n(n-1)}{2}$ ³⁸.

Thus, then, in all we have

$$\frac{n(n-1)}{2} + 3 \frac{n(n+1)}{2} = \frac{2n(2n+1)}{2}$$

equations, which is exactly equal to the number of arguments to be eliminated. Hence

p. 79

the final derivate can be obtained by the usual explicit rule of permutation, and moreover will be *its lowest form*, for it will contain in each term $\frac{n(n+1)}{2}$ prefixes belonging to the augmentatives of U , and a like number pertaining to those of V and of W , as well as $\frac{n(n-1)}{2}$ belonging to the secondary derivatives, each prefix in any one of which is trilateral, containing a prefix drawn out of those belonging to each of the proposees.

Thus every member containing $n \frac{n+1}{2} + n \frac{n-1}{2}$, that is n^2 of the original prefixes belonging to U, V, W , singly and respectively, the final derivate evolved by this process will be in its lowest terms; as was to be proved.

³⁸Vide page 76 for the Latin version.

CASE A. INDICES ALL EQUAL.

Method 2.

It is remarkable that we may vary the method just given by making

$$r + r' + r'' = n - 2, \quad s + s' + s'' = n - 2, \quad t + t' + t'' = n - 2.$$

The augmentatives will thus be of the degree $2n - 2$.

Furthermore, we must make $\alpha + \beta + \gamma = n + 2$. It will still be possible to satisfy by integer multipliers the equations

$$\begin{aligned} U &= x^\alpha F + y^\beta F' + z^\gamma F'', \\ V &= x^\alpha G + y^\beta G' + z^\gamma G'', \\ W &= x^\alpha H + y^\beta H' + z^\gamma H'', \end{aligned}$$

[these it will be useful in future to term the *equations*, x^α , y^β , z^γ being the *arguments*, and F, G, H , &c. the *factors* of decomposition] for otherwise calling the indices of x, y, z in any original argument a, b, c , their sum or n would be not greater than $(n + 2) - 3$, that is $(n - 1)$, which is absurd.

For the same reasons as in the last case no index of augmentation must be made zero: the degree of each will be $(n - \alpha) + (n - \beta) + (n - \gamma)$, that is $(2n - 2)$, and their number $\frac{(n + 1)n}{2}$; the number of augmentatives will be $3\frac{(n - 1)n}{2}$ linearly uninvolved, each of the degree $2n - 2$, and therefore containing $\frac{(2n - 1)2n}{2}$ arguments.

Now

$$\frac{(n + 1)n}{2} + 3\frac{(n - 1)n}{2} = \frac{(2n - 1)2n}{2}.$$

p. 80

Hence the final derivee may be found, and it will be in its lowest terms, for every member will contain $\frac{3(n - 1)n}{2}$ letters due to the augmentative, and $\frac{3(n + 1)n}{2}$ due to the partial derivative equations; in all then there will be $3n^2$ letters in each term.

This second method being applied to three quadratic equations of the most general form, leads to the problem of eliminating between six simple equations which lies within the limits of practical feasibility, and it is my intention to register the final derivee upon the pages of some one of our scientific Transactions as a standing monument for the guidance of hereafter coming explorers³⁹.

³⁹Elimination between *two* quadratics leads to a final derivee made up of *seven* terms only; the final derivee of *three* quadratics is made up of at least several *thousand*; nay, I believe I may safely say, several *myriads* of terms!

SCHOLIUM TO CASE A.

If we attempt to carry forward these processes to quaternary systems, it becomes necessary to make

$$\alpha + \beta + \gamma + \delta = (r - 2)n + 1$$

or else

$$\alpha + \beta + \gamma + \delta = (r - 2)n + 2,$$

where r is the number of proposees.

Now if the factors in the equations of decomposition are all integer, one of the indices of derivation must be not greater than the corresponding index in any of the original arguments, which may easily be shown to be always impossible for a system of equations, complete in all their terms, whenever their number r is greater than three, if $\alpha + \beta + \gamma + \delta = (r - 2)n + 2$; but if $\alpha + \beta + \gamma + \delta = (r - 2)n + 1$ only possible for the case of $n = 2$.

PARTICULAR METHOD APPLICABLE TO FOUR QUADRATICS.

Let $U = 0$, $V = 0$, $W = 0$, $Z = 0$, be four quadratic equations existing between x, y, z, t .

Make

$$\begin{aligned} xU &= 0, & xV &= 0, & xW &= 0, & xZ &= 0, \\ yU &= 0, & yV &= 0, & yW &= 0, & yZ &= 0, \\ zU &= 0, & zV &= 0, & zW &= 0, & zZ &= 0, \\ tU &= 0, & tV &= 0, & tW &= 0, & tZ &= 0. \end{aligned}$$

Also write

$$\begin{aligned} U &= x^2F + yF' + zF'' + tF''' = 0, \\ V &= x^2G + yG' + zG'' + tG''' = 0, \\ W &= x^2H + yH' + zH'' + tH''' = 0, \\ Z &= x^2K + yK' + zK'' + tK''' = 0. \end{aligned}$$

By eliminating linearly we get

$$\Sigma\{F\Sigma G'(H''K''' - H'''K'')\} = 0,$$

which will be of the third degree, since the factors represented by the unmarked letters F, G, H, K are of zero, and all the rest of unit dimensions.

Similarly we may obtain other equations, so that besides the sixteen augmentatives already written down, we have four secondary derivatives, namely,

$$\Pi(2\ 111) = 0, \quad \Pi(12\ 11) = 0, \quad \Pi(11\ 21) = 0, \quad \Pi(111\ 2) = 0.$$

Thus we have twenty equations and as many arguments to eliminate, since a perfect cubic function of four letters contains twenty terms.

The final derivee will contain $16 + 4 \cdot 4$ letters, that is 32 , 8 or 2^3 belonging to each system of original prefixes in each member, and will therefore be in its lowest terms: for one of the canons of form teaches us, *à priori*, that every member of the derivee deduced from any number of assumed equations must contain in each member as many prefixes belonging to one equation of the system as there are units in the product of the indices of all the rest taken together.

COROLLARY TO CASE A.

Either of the two methods given as applicable to this case enables us to determine integer values of X, Y, Z , which shall satisfy the equation

$$XU + YV + ZW = Fx^p y^q z^r,$$

where F is the final derivee and $p + q + r = 3n - 2$. For by the doctrine of simple equations we know how to express F in terms of the linear functions, out of which it is obtained by permutation, that is we are able to assign values of A, B, C , and their antitypes, as also of L and its antitype, which shall satisfy the equation

$$\begin{aligned} \Sigma(Ax^r y^{r'} z^{r''} U) + \Sigma(Bx^s y^{s'} z^{s''} V) + \Sigma(Cx^t y^{t'} z^{t''} W) \\ + \Sigma\{L\Pi(\alpha, \beta, \gamma)\} = Fx^f y^g z^h, \end{aligned} \quad (1)$$

where A, B, C , as well as L and all the quantities formed after them, are made up of integer combinations of the original prefixes.

Now the functions $\Pi(\alpha, \beta, \gamma)$ may be expressed in three ways in terms of U, V, W , as has been already shown. p. 82

We may therefore suppose these functions to be divided into three groups, and make

$$\Sigma L\Pi(\alpha\beta\gamma) = \Sigma \frac{QU + Q'V + Q''W}{x^\alpha} + \Sigma \frac{RU + R'V + R''W}{y^\beta} + \Sigma \frac{SU + S'V + S''W}{z^\gamma}. \quad (2)$$

And it is evident that the equations (1) and (2) lead immediately to the equation

$$XU + YV + ZW = Fx^{a+f} y^{b+g} z^{c+h},$$

if we call a, b, c the *greatest values* attributed respectively to α, β, γ .

Now if we suppose the first method to be followed,

$$f + g + h = 2n - 1.$$

And it will always be possible to make a, b, c of what values we please subject to the condition of $a + b + c = n - 1$; for *one* at least of the indices of derivation in

$\Pi(\alpha, \beta, \gamma)$ must be not greater than its correspondent among a, b, c ; otherwise $\alpha + \beta + \gamma$ would be not less than $(a + b + c) + 3$; but

$$\alpha + \beta + \gamma = n + 1, \quad a + b + c = n - 1,$$

which is absurd.

Hence we can satisfy $XU + YV + ZW = Fx^p y^q z^r$, p, q, r being subject to the condition of $p + q + r = 3n - 2$, but otherwise arbitrary.

Moreover, we can *not* do so if $p + q + r$ be less than $3n - 2$, for that would require $a + b + c$ to be less than $n - 1$. Now if two of the indices of derivation, as α and β , be made equal to $a + 1, b + 1$ respectively, the third $\gamma = (n + 1) - (a + b + 2) = (n - 1) - (a + b)$, and is therefore greater than c : so that $\alpha + \beta + \gamma$ for this case becomes greater than $a + b + c$, and the method falls to the ground.

In fact, I have discovered a theorem which lets me know this, *à priori*, a law which serves as a staff to guide my feet from falling into error in devising linear methods of solution, and the importance of which all candid judges who have studied the general theory of elimination cannot fail to recognize. To wit, if $X_1, X_2, X_3 \dots X_n$ be n integer complete polynomial functions of n letters $x_1, x_2 \dots x_n$, and severally of the degree $b_1, b_2, b_3 \dots b_n$; then it is always possible to satisfy the identity

$$P_1 X_1 + P_2 X_2 + P_3 X_3 + \dots + P_n X_n = F x_1^{a_1} x_2^{a_2} x_3^{a_3} \dots x_n^{a_n},$$

p. 83

if $a_1 + a_2 + a_3 + \dots + a_n$ be equal to or greater than $b_1 + b_2 + b_3 + \dots + b_n - n + 1$, but otherwise *not*⁴⁰.

This again is founded immediately upon a simple proposition, of which I have obtained a very interesting and instructive demonstration, shortly to appear, and which may be enumerated thus: “*The number of augmentees of the same degree that can be formed, linearly independent of one another, out of any number of polynomial functions of as many variables, may be either equal to or less than the number of distinct arguments contained in such augmentees, but never greater. The latter will be the case when the index of the augmentees diminished by unity*

⁴⁰Hence it is apparent, that in applying the method of multipliers, a curious and important distinction exists between the cases of there being two equations, and there being a greater number to eliminate from: for in the first case the element of arbitrariness needs never to appear; in the latter it cannot possibly be excluded from appearing in the multipliers.

This will explain how it comes to pass that the method of the text may be employed to give *various* solutions of the $XU + YV + ZW = Fx^p y^q z^r$; thus not only can p, q and r be variously made up of $(f + a), (g + b), (h + c)$, but also $\Pi(\alpha, \beta, \gamma)$ when two of the indices (α, β suppose) are each not greater than the assigned greatest values a, b may be made to figure indifferently either under the form

$$\frac{\lambda U + \mu V + \nu W}{x^\alpha} \quad \text{or that of} \quad \frac{\lambda' U + \mu' V + \nu' W}{y^\beta}.$$

is less than the sum of the indices of the original unaugmented polynomials each so diminished; the former, when the aforesaid index is equal to or greater than the aforesaid sum."

To return to the particular case of finding X, Y, Z to satisfy

$$XU + YV + ZW = Fx^p y^q z^r.$$

This has been already done according to the first method; if we employ the *second* method of elimination we shall have

$$f + g + h = 2n - 2.$$

But, now since $\alpha + \beta + \gamma = n + 2$, we shall easily see by the same method as above, that the least value of $a + b + c$ {where a, b, c denote respectively the greatest values of α, β, γ , appearing in the denominator of the fractional forms used to express $\Pi(\alpha, \beta, \gamma)$ }, will be one greater than before, or n ; so that $f + g + h + a + b + c$ will still be equal to $3n - 2$, as we might, *à priori*, by virtue of our rule, have been assured.

TERNARY SYSTEMS.

CASE B. TWO OF THE INDICES EQUAL; THE THIRD LESS BY A UNIT.

Let $U = 0, V = 0, W = 0$, be the three given equations severally of the degree $n, n, (n - 1)$.

p. 84

Make

$$r + r' + r'' = n - 2, \quad s + s' + s'' = n - 2, \quad t + t' + t'' = n - 1,$$

by multiplying U into $x^r y^{r'} z^{r''}$, V into $x^s y^{s'} z^{s''}$, W into $x^t y^{t'} z^{t''}$, we obtain augmentees each of the same, namely, the $(2n - 2)$ th degree.

The number of these is

$$\frac{(n - 1)n}{2} + \frac{(n - 1)n}{2} + \frac{n(n + 1)}{2}.$$

Again, make

$$\alpha + \beta + \gamma = n + 1.$$

It will still be possible, as before, to form equations of decomposition in which $x^\alpha, y^\beta, z^\gamma$ are the arguments, and affected with integer factors. For if we look to W even, all its arguments are of the form $x^a y^b z^c$, where $a + b + c = (n - 1)$, and each of these cannot be less than its correspondent, for that would be to say that $(n - 1)$ is not greater $(n + 1) - 3$, *à fortiori*, U and V can be decomposed in the manner described. Thus, then, we shall obtain as many secondary derivees

as in the last case (Method 1), that is, $\frac{n(n-1)}{2}$ {since $\alpha + \beta + \gamma$ is still equal to $(n+1)$ }, as before. Moreover, each of these will be of $(n-\alpha) + (n-\beta) + (n-1-\gamma)$, that is of $2n-2$ dimensions.

Altogether, therefore, we have

$$\left\{ \frac{(n-1)n}{2} + \frac{(n-1)n}{2} + \frac{n(n+1)}{2} \right\} + \frac{(n-1)n}{2}$$

linear independent equations of the degree $2n-2$, and the number of arguments to eliminate is $\frac{(2n-1)2n}{2}$. Now these two numbers are equal.

Thus we obtain a final derivate containing of U 's coefficients $\frac{(n-1)n}{2} + \frac{(n-1)n}{2}$, an equal number of V 's, but of W 's $\frac{n(n+1)}{2} + \frac{(n-1)n}{2}$; now $n(n-1)$, $n(n-1)$ and n^2 exactly express the number that ought to appear of each of these respectively: hence the final derivate is clear of *irrelevant factors*.

TERNARY SYSTEMS.

CASE C. TWO OF THE INDICES EQUAL; THE THIRD ONE GREATER BY A UNIT.

Here, calling n the highest index, the augmentees must each be made of the degree $(2n-3)$, their number will evidently be

$$\frac{(n-2)(n-1)}{2} + \frac{(n-1)n}{2} + \frac{(n-1)n}{2},$$

making the sum of the indices of derivation now, as before, equal to $(n+1)$; it will be still possible to form integer equations of decomposition, which will give rise to augmentatives of the degree $(n-\alpha) + (n-1) - \beta + (n-1) - \gamma$, that is, of $(2n-3)$ dimensions. The total number of equations, what with augmentatives and secondary derivatives, will be

$$\begin{aligned} & \left\{ \frac{(n-2)(n-1)}{2} + \frac{(n-1)n}{2} + \frac{(n-1)n}{2} \right\} + \frac{n(n-1)}{2} \\ &= \frac{4n^2 - 4n + 2}{2} = \frac{(2n-2)(2n-1)}{2}, \end{aligned}$$

that is, is equal to the exact number of distinct arguments contained between them.

Also the final derivative will contain in each member

$$\frac{(n-2)(n-1)}{2} + \frac{n(n-1)}{2},$$

p. 85

that is, $(n-1)(n-1)$, letters belonging to the first equation, and

$$\frac{(n-1)n}{2} + \frac{n(n-1)}{2},$$

that is, $n(n-1)$ belonging to those of the second and of the third, and will therefore be in its *lowest terms*.

COROLLARY TO CASES B AND C.

It is not necessary, after all that has been already said, to do more than just point out that the processes applicable to these cases enable us to determine X, Y, Z , which satisfy the equation

$$XU + YV + ZW = Fx^f y^g z^h,$$

where

$$f + g + h = 3n - 3 \quad \text{for Case B,}$$

and

$$f + g + h = 3n - 4 \quad \text{for Case C.}$$

16.

MEMOIR ON THE DIALYTIC METHOD OF ELIMINATION. PART I

[*Philosophical Magazine*, XXI. (1842), pp. 534-539; reprinted from *Proc. Roy. Irish Acad.*, Vol. II. (1840-1844), p. 130]

p. 86

The author confines himself in this part to the treatment of two equations, the final and other derivees of which form the subject of investigation.

The author was led to reconsider his former labours in this department of the general theory by finding certain results announced by M. Cauchy in *L'Institut*, March Number of the present year, which flow as obvious and immediate consequences from Mr Sylvester's own previously published principles and method.

Let there be two equations in x ,

$$\begin{aligned} U &= ax^n + bx^{n-1} + cx^{n-2} + ex^{n-3} + \&c. = 0, \\ V &= \alpha x^m + \beta x^{m-1} + \gamma x^{m-2} + \&c. = 0, \end{aligned}$$

and let $n = m + \iota$, where ι is zero or any positive value (as may be).

Let any such quantities as $x^r U$, $x^s V$, be termed augmentatives of U or V .

To obtain the derivee of a degree s units lower than V , we must join s augmentatives of U with $s + \iota$ of V . Then out of $2s + \iota$ equations

$$\begin{aligned} x^0 U &= 0, & x^1 U &= 0, & x^2 U &= 0, & \dots & x^{s-1} U &= 0, \\ x^0 V &= 0, & x^1 V &= 0, & x^2 V &= 0, & \dots & x^{\iota+s-1} V &= 0, \end{aligned}$$

we may eliminate linearly $2s + \iota - 1$ quantities.

Now these equations contain no power of x higher than $m + \iota + s - 1$; accordingly, all powers of x , superior to $m - s$, may be eliminated, and the derivee of the degree $(m - s)$ obtained in its prime form.

Thus to obtain the final derivee (which is the derivee of the degree zero), we take m augmentatives of U with n of V , and eliminate $(m + n - 1)$ quantities, namely,

$$x, \quad x^2, x^3, \dots \text{ up to } x^{m+n-1}.$$

p. 87

This process, founded upon the dialytic principle, admits of a very simple modification. Let us begin with the case where $\iota = 0$, or $m = n$. Let the augmentatives of U be termed $U_0, U_1, U_2, U_3, \dots$ and of V , $V_0, V_1, V_2, V_3, \dots$, the equations themselves being written

$$\begin{aligned} U &= ax^n + bx^{n-1} + cx^{n-2} + \&c. \\ V &= a'x^n + b'x^{n-1} + c'x^{n-2} + \&c. \end{aligned}$$

It will readily be seen that

$$\begin{aligned} & a'U_0 - aV_0, \\ & (b'U_0 - bV_0) + (a'U_1 - aV_1), \\ & (c'U_0 - cV_0) + (b'U_1 - bV_1) + (a'U_2 - aV_2), \text{ \&c.} \end{aligned}$$

will be each linearly independent functions of $x, x^2, \dots x^{m-1}$, no higher power of x remaining. Whence it follows, that to obtain a derivate of the degree $(m - s)$ in its prime form, we have only to employ the s of those which occur first in order, and amongst them eliminate $x^{m-1}, x^{m-2}, \dots x^{m-s+1}$. Thus, to obtain the *final* derivate, we must make use of n , that is, the entire number of them.

Now, let us suppose that ι is not zero, but $m = n - \iota$. The equation V may be conceived to be of n instead of m dimensions, if we write it under the form

$$0x^n + 0x^{n-1} + 0x^{n-2} + \dots + 0x^{m+1} + \alpha x^m + \beta x^{m-1} + \text{\&c.} = 0,$$

and we are able to apply the same method as above; but as the first ι of the coefficients in the equation above written are zero, the first ι of the quantities

$$(a'U_0 - aV_0), \quad (b'U_0 - bV_0) + (a'U_1 - aV_1), \quad \text{\&c.}$$

may be read simply

$$-aV_0, \quad -bV_0 - aV_1, \quad -cV_0 - bV_1 - aV_2, \quad \text{\&c.}$$

and evidently their office can be supplied by the simple augmentatives themselves,

$$V_0 = 0, \quad V_1 = 0, \quad V_2 = 0 \dots V_{l-1} = 0;$$

and thus ι letters, which otherwise would be *irrelevant*, fall out of the several derivees.

The author then proceeds with remarks upon the general theory of simple equations, and shows how by virtue of that theory his method contains a solution of the identity

$$X_r U + Y_r V = D_r,$$

where D_r is a derivate of the r th degree of U and V , and accordingly, X_r of the form

$$\lambda + \mu x + \nu x^2 + \dots + \theta x^{m-r-1},$$

and Y_r of the form

$$l + mx + \dots + tx^{n-r-1},$$

p. 88

and accounts *à priori* for the fact of not more than $(n - r)$ simple equations being required for the determination of the $(m + n - 2r)$ quantities λ, μ, ν , &c. l, m, n , &c., by exhibiting these latter as *known* linear functions of no more than $(n - r)$ unknown quantities left to be determined.

Upon this remarkable relation may be constructed a method well adapted for the expeditious computation of numerical values of the different derives.

He next, as a point of curiosity, exhibits the values of the secondary functions,

$$\begin{aligned} & a'U_0 - aV_0, \\ & b'U_0 - bV_0 + a'U_1 - aV_1, \\ & c'U_0 - cV_0 + b'U_1 - bV_1 + a'U_2 - aV_2, \quad \&c. \end{aligned}$$

under the form of symmetric functions of the roots of the equations $U = 0$, $V = 0$, by aid of the theorems developed in the London and Edinburgh *Philosophical Magazine*, December 1839⁴¹, and afterwards proceeds to a more close examination of the final derivee resulting from two equations each of the same (any given) degree.

He conceives a number of cubic blocks each of which has two numbers, termed its characteristics, inscribed upon one of its faces, upon which the value of such a block (itself called an *element*) depends.

For instance, the value of the *element*, whose characteristics are r, s , is the difference between two products: the one of the coefficient r th in order occurring in the polynomial U , by that which comes s th in order in V ; the other product is that of the coefficient s th in order of the polynomial U , by that r th in order of V ; so that if the degree of each equation be n , there will be altogether $\frac{1}{2}n(n+1)$ such elements.

The blocks are formed into squares or flats (*plafonds*) of which the number is $\frac{n}{2}$ or $\frac{n+1}{2}$, according as n is even or odd. The first of these contains n blanks in a side, the next $(n-2)$, the next $(n-4)$, till finally we reach a square of four blocks or of one, according as n is even or odd. These flats are laid upon one another so as to form a regularly ascending pyramid, of which the two diagonal planes are termed the planes of separation and symmetry respectively. The former divides the pyramid into two halves, such that no element on the one side of it is the same as that of any block in the other. The plane of symmetry, as the name denotes, divides the pyramid into two exactly *similar* parts; it being a rule, that *all elements lying in any given line of a square (plafond) parallel to the plane of separation are identical*; moreover, the sum of the characteristics is the same, for *all elements lying anywhere in a plane* parallel to that of separation. p. 89

All the terms in the final derivee are made up by multiplying n elements of the pile together, under the sole restriction, that no two or more terms of the said product shall lie in any one plane out of the two sets of planes perpendicular to the sides of the squares. The *sign* of any such product is determined by the places of either set of planes parallel to a side of the squares and to one another, in which the elements composing it may be conceived to lie.

⁴¹p. 40 above. ED.

The author then enters into a disquisition relating to the *number* of terms which will appear in the final derieve, and concludes this first part with the statement of two general canons, each of which affords as many tests for determining whether a prepared combination of coefficients can enter into the final derieve of any number of equations as there are units in that number, but so connected as together only to afford double that number, less one, of independent conditions.

The first of these canons refers simply to the number of letters drawn out of *each of the given equations* (supposed homogeneous); the second to what he proposes to call the *weight* of every term in the derieve in respect to *each of the variables* which are to be eliminated.

The author subjoins, for the purpose of conveying a more accurate conception of his Pyramid of derivation, examples of the mode in which it is constructed.

When $n = 2$ there is one flat, viz.

When $n = 1$ there is one flat, viz.

1, 2

2, 3	2, 4
2, 4	3, 4

Let $n = 3$, there will be two flats:

Let $n = 4$, there will still be two flats only:

2, 3

2, 3	2, 4
2, 4	3, 4

1, 2	1, 3	1, 4
1, 3	1, 4	2, 4
1, 4	2, 4	3, 4

1, 2	1, 3	1, 4	1, 5
1, 3	1, 4	1, 5	2, 5
1, 4	1, 5	2, 5	3, 5
1, 5	2, 5	3, 5	4, 5

Let $n = 5$, there will be three flats:

p. 90

3, 4	2, 3	2, 4	2, 5		
	2, 4	2, 5	3, 5		
	2, 5	3, 5	4, 5		
	1, 2	1, 3	1, 4	1, 5	1, 6
	1, 3	1, 4	1, 5	1, 6	2, 6
	1, 4	1, 5	1, 6	2, 6	3, 6
	1, 5	1, 6	2, 6	3, 6	4, 6
	1, 6	2, 6	3, 6	4, 6	5, 6

Let $n = 6$, there will be three flats:

3, 4	3, 5	2, 3	2, 4	2, 5	2, 6
3, 5	4, 5	2, 4	2, 5	2, 6	3, 6
		2, 5	2, 6	3, 6	4, 6
		2, 6	3, 6	4, 6	5, 6
1, 2	1, 3	1, 4	1, 5	1, 6	1, 7
1, 3	1, 4	1, 5	1, 6	1, 7	2, 7
1, 4	1, 5	1, 6	1, 7	2, 7	3, 7
1, 5	1, 6	1, 7	2, 7	3, 7	4, 7
1, 6	1, 7	2, 7	3, 7	4, 7	5, 7
1, 7	2, 7	3, 7	4, 7	5, 7	6, 7

Thus the work of computation reduces itself merely to calculating $n\frac{n+1}{2}$ elements, or the $n(n+1)$ cross-products out of which they are constituted, and combining them factorially after that law of the pyramid, to which allusion has been already made.

17.

ELEMENTARY RESEARCHES IN THE ANALYSIS OF COMBINATORIAL AGGREGATION

[*Philosophical Magazine*, XXIV. (1844), pp. 285–296]

p. 91

The ensuing inquiries will be found to relate to combination-systems, that is, to combinations viewed in an aggregative capacity, whose species being given, we shall have to discover rules for ranging or evolving them in classes amenable to certain prescribed conditions. The question of numerical amount will only appear incidentally, and never be made the primary object of investigation⁴².

The number of things combined will be termed the modulus of the system to which they belong. The elements taken singly, or combined in twos, threes, &c., will be denominated accordingly the monadic, duadic, triadic elements, or simply the monads, duads, or triads of the system.

Let us agree to denote by the word *synthème*⁴³ any aggregate of combinations in which all the monads of a given system appear once, and once only.

It is manifest that many such *synthèmes* totally diverse in every term may be obtained for a given system to any modulus, and for any order of combination.

Let us begin with considering the case of duad *synthèmes*. Take the modulus 4 and call the elements *a, b, c, d*.

$$(ab, cd), \quad (ac, bd), \quad (ad, cb)$$

constitute three perfectly independent *synthèmes*, and these three *synthèmes* include between them all the duad elements, so that no more independent *synthèmes* can be obtained from them.

p. 92

Again, let *a, b, c, d, e, f* be the monads; we can write down five independent *synthèmes*, to wit,

$$\left. \begin{array}{lll} ab, & cd, & ef \\ ad, & cf, & eb \\ ac, & de, & fb \\ af, & bd, & ce \\ ae, & df, & bc \end{array} \right\}.$$

We can write no more than these without repeating duads which have already appeared⁴⁴.

⁴²The present theory may be considered as belonging to a part of mathematics which bears to the combinatorial analysis much the same relation as the geometry of position to that of measure, or the theory of numbers to computative arithmetic; number, place, and combination (as it seems to the author of this paper) being the three intersecting but distinct spheres of thought to which all mathematical ideas admit of being referred.

⁴³From $\sigma\nu\nu$ and $\tau i\theta\eta\mu\iota$.

⁴⁴Such an aggregate of *synthèmes* may be therefore termed a Total.

We propose to ourselves this problem:—*A system to any even⁴⁵ modulus being given, to arrange the whole of its duads⁴⁶ in the form of syntheses; or in other words, to evolve a Total of duad syntheses to any given even modulus⁴⁷.*

When the modulus is odd, as before remarked, the formation of a duad synthesis is of course impossible, for any number of duads must necessarily contain an even number of monadic elements; but there is nothing to prevent us from forming in all cases what may be termed a bisynthesis or diplothesis, that is, an aggregate of combinations, where each element occurs twice and no more.

For instance, if the elements be called after the letters of the alphabet, we have

$$\left(\begin{array}{l} ab, bc, cd, de, ea \\ ac, ce, eb, bd, da \end{array} \right),$$

the bisynthetic total to modulus 5; and in like manner

$$\left. \begin{array}{l} ab, bc, cd, de, ef, fg, ga \\ ac, ce, eg, gb, bd, df, fa \\ ad, dg, gc, cf, fb, be, ea \end{array} \right\} \text{ the total to modulus 7.}$$

p. 93

In general, if n be the modulus, the number of duads is $n \frac{n-1}{2}$; n being even, $\frac{n}{2}$ duads go to each synthesis, and therefore the total contains $(n-1)$ of these.

⁴⁵The modulus must be even, as otherwise it is manifest no single synthesis can be formed. We shall before long extend the scope of our inquiry so as to take in the case of odd moduli.

⁴⁶Triadic systems will be treated of hereafter.

⁴⁷It is scarcely necessary to advert here to the fact of the problem being in general indeterminate and admitting of a great variety of solutions.

When the modulus is four there is only one synthetic arrangement possible, and there is no indeterminateness of any kind; from this we can infer, *a priori*, the reducibility of a biquadratic equation; for using Φ, ϕ, F to denote rational symmetrical forms of function, it follows that

$$F \left\{ \begin{array}{l} f\{\phi(a, b), \phi(c, d)\} \\ f\{\phi(a, c), \phi(b, d)\} \\ f\{\phi(a, d), \phi(b, c)\} \end{array} \right\}$$

is itself a rational symmetric function of a, b, c, d . Whence it follows that if a, b, c, d be the roots of a biquadratic equation, $f\{\phi(a, b), \phi(c, d)\}$ can be found by the solution of a cubic: for instance, $(a+b) \times (c+d)$ can be thus determined, whence immediately the sum of any two of the roots comes out from a quadratic equation.

To the modulus 6 there are fifteen different syntheses capable of being constructed; at first sight it might be supposed that these could be classed in natural families of three or of five each, on which supposition the equation of the sixth degree could be depressed; but on inquiry this hope will prove to be futile, not but what natural affinities do exist between the totals; but in order to separate them into families each will have to be taken twice over, or in other words, the fifteen syntheses to modulus 6 being reduplicated subdivide into six natural families of five each. Again, it is true that the triads to modulus 6 (just like the duads to modulus 4) admit of being thrown into but one synthetic total, but then this will contain ten syntheses, a number greater than the modulus itself.

If n be odd, then, since always n duads go to a bisyntheme, the number of such in the total is $\frac{n-1}{2}$.

Before proceeding to the solution of the problem first proposed, let us investigate the theory of diplothematic arrangement. Here we shall find another term convenient to employ. By a cyclotheme, I designate a fixed arrangement of the elements in one or more circles, in which, although for typographical purposes they are written out in a straight line, the last term is to be viewed as contiguous and antecedent to the first; the recurrence may be denoted by laying a dot upon the two opened ends of the circle; $\dot{a}.b.c.d.\dot{e}$ will thus denote a cyclotheme to modulus 5; $\dot{a}.b.c.d.e.f.g.h.\dot{k}$ the same to modulus 9; so also is $\dot{a}.b.c, \dot{d}.e.f, \dot{g}.h.k$ a cyclotheme of another species to the same modulus. In general the number of terms will be alike in each division of a cyclotheme.

Now it is evident that every cyclotheme, on taking together the elements that lie in conjunction, may be developed into a diplotheme. Thus

$$\begin{aligned}\dot{1}.2.\dot{3} &= 12, 23, 31, \\ \dot{1}.2.3.\dot{4} &= 12, 23, 34, 41, \\ (\dot{1}.2.\dot{3}; \dot{4}.5.\dot{6}; \dot{7}.8.\dot{9}) &= \begin{pmatrix} 12, & 23, & 31 \\ 45, & 56, & 64 \\ 78, & 89, & 97 \end{pmatrix}.\end{aligned}$$

Hence we shall derive a rule for throwing the duads of any system into bisynthemes.

Let $m = 3$, we have simply $\dot{a}\dot{b}\dot{c}$;

$$m = 5, \text{ we write } \begin{array}{l} \dot{a}.b.c.d.\dot{e}, \\ \dot{a}.c.e.b.\dot{d}, \end{array}$$

the second being derived from the first by omitting every alternate term; similarly below, the lines are derived each from its antecedent.

$$m = 7, \text{ we have } \begin{array}{l} \dot{a}.b.c.d.e.f.\dot{g}, \\ \dot{a}.c.e.g.b.d.\dot{f}, \\ \dot{a}.e.b.f.c.g.\dot{d}. \end{array}$$

A very little consideration will serve to prove that in this way, m being a prime number, $\frac{m-1}{2}$ cyclothemes may be formed, such that no element will ever be found more than once in contact on either side with any other; whence the rule for obtaining the diplothematic total to any prime-number modulus is apparent.

For example, to modulus 7 the total reads thus:—

$$\begin{array}{lllllll} \text{1st.} & ab, & bc, & cd, & de, & ef, & fg, & ga \\ \text{2nd.} & ac, & ce, & eg, & gb, & bd, & df, & fa \\ \text{3rd.} & ae, & eb, & bf, & fc, & cg, & gd, & da \end{array}$$

and no more remains to be said on this special case.

Let us now return to the theory of even moduli, and show how to apply what has been just done to constructing a synthematic total to a modulus which is the double of a prime number.

Suppose the modulus to be six, the number of synthemes is five. Let the six elements, a, b, c, d, e, f , be taken in three parts, so that each part contains two of them; let these parts be called A, B, C , where A denotes ab , B, cd , and C, ef .

Now the duads will evidently admit of a distinction into two classes, those that lie in one part, and those that lie between two; thus ab, cd, ef will be each unipartite duads, the rest will be bipartite.

The unipartite duads may be conveniently formed into a syntheme by themselves; it only remains to form the four remaining bipartite duad synthemes.

Write the parts in cyclothematic order, as below:

$$\dot{A}\dot{B}\dot{C}.$$

It will be observed that each part may be written in two positions; thus

$$\begin{array}{rcll} A & \text{may be expressed by} & \begin{array}{c} a \\ b \end{array} & \text{or by} \quad \begin{array}{c} b \\ a' \end{array} \\ B & & \begin{array}{c} c \\ d \end{array} & \begin{array}{c} d \\ c' \end{array} \\ C & & \begin{array}{c} e \\ f \end{array} & \begin{array}{c} f \\ e' \end{array} \end{array}$$

Now we may form a cyclic table of positions as below:

$$\begin{array}{ccc} \dot{A} & B & \dot{C} \\ \hline 1 & 1 & 1 \\ 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{array}$$

Here the numbers in each horizontal line denote the synchronic positions of the parts. p. 95

On inspection it will be discovered that A will be found in each of its two positions, with B in each of its two; similarly B with C , and C with A . In fact the four permutations, 11, 12, 21, 22, occur, though in different orders, in any two assigned vertical columns.

Now developpe the preceding table, and we have

$$\begin{array}{cccc} ace & adf & bcf & bde, \\ bdf & bce & ade & acf; \end{array}$$

and these being read off (the *superior* of each antecedent with the *inferior* of each consequent⁴⁸) must manifestly give the four independent bipartite syntheses which we were in quest of, *videlicet*

$$(ad, cf, eb), \quad (ac, de, fb), \quad (bd, ce, fa), \quad (bc, df, ea) :$$

these four, together with the synthesis first described (ab, cd, ef) , constitute a duad synthemetic total to modulus 6.

Before proceeding further let us take occasion to remark that the foregoing table of positions may evidently be extended to any odd number of terms by repetition of the second and third places, as seen in the annexed tables of position.

$$\begin{array}{ccccccccc} \dot{1} & 1 & 1 & \dot{1} & \dot{1} & 1 & 1 & 1 & 1 & \dot{1} \\ \dot{1} & 2 & 2 & \dot{2} & \dot{1} & 2 & 2 & 2 & 2 & \dot{2} \\ \dot{2} & 1 & 2 & \dot{1} & \dot{2} & 1 & 2 & 1 & 2 & \dot{2} \\ \dot{2} & 2 & 1 & \dot{1} & \dot{2} & 2 & 1 & 2 & 1 & \dot{2} \end{array}$$

Now let 10 be the modulus.

As before divide the elements into five parts, which call A, B, C, D, E .

The unipartite duads fall into a single synthesis; the eight remaining bipartite syntheses may be found as follows:—

Arrange in cyclothemes $\left(\frac{n-1}{2} \text{ in number}\right)$ the odd modulus system A, B, C, D, E .

We have thus

$$\dot{A}BCD\dot{E}, \quad \dot{A}CEB\dot{D}.$$

p. 96

Let each cyclotheme be taken in the four positions given in the table above, we have thus 2×4 , that is, eight arguments.

$$\begin{array}{cccc} \dot{a}bcd\dot{e}, & \dot{a}\beta\gamma\delta\dot{\epsilon}, & \dot{a}b\gamma d\dot{e}, & \dot{a}\beta c\delta\dot{e}, \\ \alpha\beta\gamma\delta\epsilon, & abcde, & a\beta cde, & ab\gamma de, \\ \dot{a}ceb\dot{d}, & \dot{a}\gamma e\beta\dot{\delta}, & \dot{a}ceb\dot{\delta}, & \dot{a}\gamma e\beta\dot{d}, \\ \alpha\gamma e\beta\delta, & acebd, & a\gamma e\beta d, & acebd. \end{array}$$

And each of these arguments will furnish one bipartite synthesis, by reading off, as before, the superior of each antecedent with the inferior of each consequent; and the least reflection will serve to show that the same duad can never appear in two distinct arguments.

In like manner, if the modulus be 14 and seven parts be taken, the bipartite syntheses, twelve in number, may be expressed symbolically thus:

$$\left(\begin{array}{c} \dot{1}.1.1.1.1.1.\dot{1} \\ +\dot{1}.2.2.2.2.2.\dot{2} \\ +\dot{2}.1.2.1.2.1.\dot{2} \\ +\dot{2}.2.1.2.1.2.\dot{1} \end{array} \right) \times \left(\begin{array}{c} \dot{A}.B.C.D.E.F.\dot{G} \\ +\dot{A}.C.E.G.B.D.\dot{F} \\ +\dot{A}.E.B.F.C.G.\dot{D} \end{array} \right).$$

⁴⁸Any other fixed order of successive conjunction would answer equally well.

Nay more, from the above table, if we agree to name the elements

$$\begin{array}{cc} A_1 & B_1 \\ A_2 & B_2 \end{array}, \quad \&c.,$$

we can at once proceed to calculate each of the twelve synthemes in question by an easy algorithm. For instance,

$$(\dot{1}.2.2.2.2.\dot{2}) \times (\dot{A}.C.E.G.B.D.\dot{F}) = (A_1C_1, C_2E_1, E_2G_1, G_2B_1, B_2D_1, D_2F_1, F_2A_2).$$

And again

$$(\dot{2}.1.2.1.2.1.\dot{2}) \times (\dot{A}.E.B.F.C.G.\dot{D}) = A_2E_2, E_1B_1, B_2F_2, F_1C_1, C_2G_2, G_1D_1, D_2A_1;$$

each figure occurring once unchanged as an antecedent and once changed as a consequent.

If it were thought worth while it would not be difficult, by using numbers instead of letters, to obtain a general analytical formula, from which all similarly constituted synthemes to any modulus might be evolved.

But the rule of proceeding must be now sufficiently obvious; the modulus being $2p$, we divide the elements into p classes; these may be arranged into $\frac{p-1}{2}$ distinct forms of cyclothematic arrangement, and each of the cyclothemes taken in four positions, thus giving $4 \times \frac{p-1}{2}$, that is, $2p-2$ bipartite synthemes, the whole number that can be formed to the given modulus $2p$.

p. 97

I shall now proceed to the theory of bipartite synthemes to the modulus $2m \times p$, by which it is to be understood that we have p parts each containing $2m$ terms, and p is at present supposed to be a prime number; the total number of synthemes to the modulus $2mp$ being $2mp-1$, and $2m-1$ of these evidently being capable of being made unipartite; the remainder, $2mp-2m$, that is, $(p-1)2m$, will be the number of bipartites to be obtained⁴⁹:

$$2m(p-1) = \frac{p-1}{2} \times 4m;$$

$\frac{p-1}{2}$ denotes the total number of cyclothemes to modulus p ; $4m$, as will be presently shown, the number of lines or syzygies in the Table of position.

To fix our ideas let the modulus be 4×3 , and let A, B, C be three parts:

$$\left. \begin{array}{cccc} a_1 & a_2 & a_3 & a_4, \\ b_1 & b_2 & b_3 & b_4, \\ c_1 & c_2 & c_3 & c_4 \end{array} \right\} \text{ their constituents respectively.}$$

⁴⁹In general, if there be π parts of μ terms each, and $\mu\pi$ be even, the number of bipartite synthemes is $(\pi-1)\mu$, as is easily shown from dividing the whole number of bipartite duads by the semi-modulus.

Give a fixed order to the constituents of each part, then each of them may be taken in four positions; thus A may be written

$$\begin{array}{cccc} a_1 & a_2 & a_3 & a_4, \\ a_2 & a_3 & a_4 & a_1, \\ a_3 & a_4 & a_1 & a_2, \\ a_4 & a_1 & a_2 & a_3. \end{array}$$

Assume some particular position for each, as, for instance,

$$\begin{array}{ccc} a_1 & b_1 & c_1, \\ a_2 & b_2 & c_2, \\ a_3 & b_3 & c_3, \\ a_4 & b_4 & c_4, \end{array}$$

and read off by coupling the first and third vertical places of each antecedent with the second and fourth respectively of each consequent; we have accordingly,

$$a_1b_2, b_1c_2, c_1a_2, \quad a_3b_4, b_3c_4, c_3a_4.$$

It is apparent that the same combinations will recur if any two contiguous parts revolve simultaneously through two steps; or in other words, that $A_rB_s = A_{r+2}B_{s+2}$, where μ is any number, odd or even.

p. 98

Symbolically speaking, therefore, as regards our table of position,

$$r : s = r + 2 : s + 2,$$

or more generally,

$$= r + 2 \pm 4i : s + 2 \pm 4i.$$

So that

$$\begin{array}{ll} 1 : 1 = 3 : 3, & 2 : 1 = 4 : 3, \\ 1 : 2 = 3 : 4, & 2 : 2 = 4 : 4, \\ 1 : 3 = 3 : 1, & 2 : 3 = 4 : 1, \\ 1 : 4 = 3 : 2, & 2 : 4 = 4 : 2. \end{array}$$

There are therefore no more than eight independent unequivalent permutations to every pair of parts. Now inspect the following table of position:—

$$\begin{array}{ll} 1.1.1, & 2.1.2, \\ 1.2.3, & 2.2.4, \\ 1.3.2, & 2.3.1, \\ 1.4.4, & 2.4.3. \end{array}$$

It will be seen that in the first and second, second and third, third and first places, all the eight independent permutations occur under different names; the law of formation of such and similar tables will be explained in due time; enough

for our present object to see how, by means of this table, we are able to obtain the bipartite syntheses to the given modulus 4×3 ; the number according to our formula is $2 \times 4 \times \frac{3-1}{2} = 8$, and they may be denoted symbolically as follows:—

$$(\dot{A}.B.\dot{C}) \begin{pmatrix} 1.1.1 + 1.2.3 + 1.3.2 + 1.4.4 \\ +2.1.2 + 2.2.4 + 2.3.1 + 2.4.3 \end{pmatrix}.$$

Each of the eight terms connected by the sign of + gives a distinct syntheme; for example, let us operate on

$$\dot{A}.B.\dot{C} \times (2.3.1).$$

2.3.1 denotes 2.3, 3.1, 1.2.

$$2.3 \text{ gives rise to } 2(3+1) + (2+2).(3+3) = 2.4 + 4.2.$$

$$3.1 \text{ gives rise to } 3(1+1) + (3+2).(1+3) = 3.2 + 1.4.$$

$$1.2 \text{ gives rise to } 1(2+1) + (1+2).(2+3) = 1.3 + 3.1.$$

The syntheme in question is therefore

$$A_2B_4, A_4B_2, B_3C_2, B_1C_4, C_1A_3, C_3A_1,$$

and so on for all the rest, the rule being that

$$r : s = r(s+1) + (r+2)(s+3).$$

p. 99

Now, as before, it is evident that if we look only to contiguous terms, the above table of position may be extended to any number of odd terms, simply by repetition of the second and third figures in each syzygy; and hence the rule for obtaining the bipartite syntheses to the modulus $4 \times p$ is apparent. For instance, let $p = 7$, there will be $8 \times \frac{7-1}{2}$, that is, 8×3 of them denoted as follows:—

$$\left\{ \begin{array}{l} \dot{A}.B.C.D.E.F.\dot{G} \\ +\dot{A}.C.E.G.B.D.\dot{F} \\ +\dot{A}.E.B.F.C.G.\dot{D} \end{array} \right\} \times \left\{ \begin{array}{l} 1.1.1.1.1.1.1 + 2.1.2.1.2.1.2 \\ +1.2.3.2.3.2.3 + 2.2.4.2.4.2.4 \\ +1.3.2.3.2.3.2 + 2.3.1.3.1.3.1 \\ +1.4.4.4.4.4.4 + 2.4.3.4.3.4.3 \end{array} \right\}.$$

As an example of the mode of development, let us take the term

$$\dot{A}.E.B.F.C.G.\dot{D} \times \dot{2}.4.3.4.3.4.\dot{3},$$

$$\begin{aligned} \dot{2}.4.3.4.3.4.\dot{3} &= (2 : 4, 4 : 3, 3 : 4, 4 : 3, 3 : 4, 4 : 3, 3 : 2) \\ &= \begin{pmatrix} 2.1 & 4.4 & 3.1 & 4.4 & 3.1 & 4.4 & 3.3 \\ +4.3 & +2.2 & +1.3 & +2.2 & +1.3 & +2.2 & +1.1 \end{pmatrix}, \end{aligned}$$

$$\dot{A}.E.B.F.C.G.\dot{D} = AE, EB, BF, FC, CG, GD, DA,$$

and the product

$$= \begin{pmatrix} A_2E_1, & E_4B_4, & B_3F_1, & F_4C_4, & C_3G_1, & G_4D_4, & D_3A_3 \\ A_4E_3, & E_2B_2, & B_1F_3, & F_2C_2, & C_1G_3, & G_2D_2, & D_1A_1 \end{pmatrix}.$$

Let the modulus be 6×3 ; as before, give a fixed cyclic order to the constituents of each part, and each will admit of being exhibited in six positions.

Write similarly as before,

$$\begin{array}{lll} a_1 & b_1 & c_1, \\ a_2 & b_2 & c_2, \\ a_3 & b_3 & c_3, \\ a_4 & b_4 & c_4, \\ a_5 & b_5 & c_5, \\ a_6 & b_6 & c_6, \end{array}$$

and take the odd places of each antecedent with the even places of each consequent; it will now be seen that

$$r : s = r + 2 : s + 2 = r + 4 : s + 4,$$

and the number of independent permutations is $\frac{6.6}{3} = 2.6$; and so in general, if there be $2m$ constituents in a part, the number of independent permutations is

$$\frac{2m.2m}{m} = 4m.$$

The rule for the formation of the table will be apparent on inspection. I suppose only three parts, as the rule may always be extended to any number by reiteration of the second and third terms. The table will be found to resolve itself naturally into four parts, each containing m lines. p. 100

Let $m = 1$, we have

$$\begin{array}{ll} 1.1.1 & 2.1.2 \\ 1.2.2 & 2.2.1 \end{array}$$

$m = 2$, we have

$$\begin{array}{ll} 1.1.1 & 2.1.2 \\ 1.2.3 & 2.2.4 \\ 1.3.2 & 2.3.1 \\ 1.4.4 & 2.4.3 \end{array}$$

$m = 3$, we have

$$\begin{array}{ll} 1.1.1 & 2.1.2 \\ 1.2.3 & 2.2.4 \\ 1.3.5 & 2.3.6 \\ 1.4.2 & 2.4.1 \\ 1.5.4 & 2.5.3 \\ 1.6.6 & 2.6.5 \end{array}$$

$m = 4$, we have

1.1.1	2.1.2
1.2.3	2.2.4
1.3.5	2.3.6
1.4.7	2.4.8
1.5.2	2.5.1
1.6.4	2.6.3
1.7.6	2.7.5
1.8.8	2.8.7

So that x , going through all its values from 1 to m , the general expression for the four parts is

$$\Sigma \left\{ \begin{array}{l} 1.x(2x-1) + 1(m+x)2x \\ +2.x.2x + 2(m+x)(2x-1) \end{array} \right\}.$$

To show the use of this formula, let us suppose that we have seven parts, each containing ten terms, the general expression for the bipartite duad synthemes is

$$\left\{ \begin{array}{l} \dot{A}.B.C.D.E.F.\dot{G} \\ +\dot{A}.C.E.G.B.D.\dot{F} \\ +\dot{A}.E.B.F.C.G.\dot{D} \end{array} \right\} \times \Sigma \left\{ \begin{array}{l} 1.x(2x-1)x(2x-1)x(2x-1) \\ +2.x.2x.x.2x.x.2x \\ +1(5+x)2x(5+x)2x(5+x)2x \\ +2(5+x)(2x-1)(5+x)(2x-1)(5+x)(2x-1) \end{array} \right\}.$$

Make, for example, $x = 3$, one of the synthemes in question out of the twelve corresponding to this value will be

$$\dot{A}.C.E.G.B.D.\dot{F} \times \dot{2}.3.6.3.6.3.\dot{6}.$$

Here

$$\dot{A}.C.E.G.B.D.\dot{F} = AC, CE, EG, GB, BD, DF, FA,$$

$$\dot{2}.3.6.3.6.3.\dot{6} = \left(\begin{array}{ccccccc} 2.4 & 3.7 & 6.4 & 3.7 & 6.4 & 3.7 & 6.3 \\ +4.6 & +5.9 & +8.6 & +5.9 & +8.6 & +5.9 & +8.5 \\ +6.8 & +7.1 & +10.8 & +7.1 & +10.8 & +7.1 & +10.7 \\ +8.10 & +9.3 & +2.10 & +9.3 & +2.10 & +9.3 & +2.9 \\ +10.2 & +1.5 & +4.2 & +1.5 & +4.2 & +1.5 & +4.1 \end{array} \right),$$

and the product

$$\begin{array}{ccccccc} = A_2C_4, & C_3E_7, & E_6G_4, & G_3B_7, & B_6D_4, & D_3F_7, & F_6A_3 \\ A_4C_6, & C_5E_9, & E_8G_6, & G_5B_9, & B_8D_6, & D_5F_9, & F_8A_5, \\ & \&c. & & \&c. & & \&c. \end{array}$$

To prove the rule for the table of formation, it will be sufficient to show that no two contiguous duads ever contain the same or equivalent permutations; the equation of equivalence it will be remembered is

$$r : s = r + 2i \pm 2m : s + 2i \pm 2m.$$

Now, as regards the first and second terms, it is manifest that $1 : x$ cannot be equivalent, either to $1 : x'$ nor to $2 : x$, nor to $2 : x'$, where x' is any number differing from x .

Similarly, as regards the last and first terms, $x : 1$ cannot be equivalent to $x' : 1$, nor to $x : 2$, nor to $x' : 2$; therefore there is no danger as far as the first term is concerned, either as antecedent or consequent.

Again, it is clear that $x : (2x - 1)$ cannot interfere with $x' : 2x'$, nor $(m + x) : 2x$ with $(m + x') : (2x' - 1)$; neither can $(2x - 1) : x$ with $2x' : x'$, nor $2x : (m + x)$ with $(2x' - 1) : (m + x')$.

Again, if possible, let

$$x : (2x - 1) = (m + x') : (2x' - 1);$$

then

$$m + x' - x = 2i,$$

and

$$2x' - 2x = 2i,$$

therefore

$$2m = 2i,$$

or

$$m = i,$$

which is impossible, since $\pm i$ is the difference between two indices, each less than m .

p. 102

Similarly,

$$m + x : 2x \text{ cannot } = x' : 2x',$$

and *vice versa* with the terms changed

$$2x : (m + x) \text{ cannot } = 2x' : x',$$

and

$$(2x - 1) : x \text{ cannot } = (2x' - 1) : (m + x'),$$

which proves the rule for the table of formation.

So much for the bipartite duad synthemes. As regards the unipartite synthemes little need be said, for every part may be treated as a separate system, and as each will produce an equal number of synthemes, these being taken one with another, will furnish just as many unipartite synthemes of the whole system as there are synthemes due to each part. Thus then the synthemetic resolution of the modulus $2m \times p$ may be made to depend on the synthemization of $2m$ and the cyclothematization of p . This has been already shown (whatever m may be) for the case of p being a prime number; but I proceed now to extend the rule to the more general case of p being any number whatever.

18.

ON THE EXISTENCE OF ABSOLUTE CRITERIA FOR DETERMINING THE ROOTS OF NUMERICAL EQUATIONS

[*Philosophical Magazine*, XXV. (1844), pp. 442-445]

p. 103

I wish to indicate in this brief notice a fact which I believe has escaped observation hitherto, that there exist, certainly in some cases, and probably in all, infallible criteria for determining whether a given equation has all its roots rational or not.

In the equation of the second degree it is enough, in order that this may be the case, that the expression for the square of the difference of the roots shall be a perfect square; in other words, if $x^2 - px + q = 0$ have its roots rational, $p^2 - 4q$ must be not only a positive number (the condition of the roots being real), but that number must also be a complete square. In this case it is further evident that p must be either prime to q , or if not, the greatest common measure of p^2 and q must be a perfect square; but this condition is contained in the former, which is a sufficient criterion in itself.

If we now consider the equation of the third degree,

$$x^3 - px^2 + qx - r = 0,$$

one condition is, that the product of the squared differences shall be a perfect square; in other words, the equation cannot have all its roots rational unless

$$p^2q^2 - 4q^3 - 18pqr - 4p^3r - 27r^2$$

be a positive square number.

This remark is made at the end of the second supplement of Legendre's *Theory of Numbers*, and is indeed self-evident; and in like manner one condition may be obtained for an equation of any degree which is to have all its roots rational; but this is far from being the sole condition required.

p. 104

In the equation of the third degree, however, one other condition, conjoined with that above expressed, will serve to determine infallibly whether all the roots are rational or not.

To obtain this condition, let us suppose that by making $3x = y + p$ we obtain the equation

$$y^3 - Qy - R = 0.$$

Calling the three roots of this new equation α, β, γ (all of which it is evident must be rational if those of the first equation are so), we have

$$\alpha + \beta + \gamma = 0,$$

$$Q = -(\alpha\beta + \alpha\gamma + \beta\gamma) = \alpha^2 + \alpha\beta + \beta^2,$$

$$R = \alpha\beta\gamma.$$

From the last two equations it is easily seen that if k be any prime factor common to Q and R , k^2 will be contained in Q , and k^3 in R ; or, in other words, k will be a common measure of α, β, γ .

We have therefore a *second condition*, that $9q - 3p^2$ shall be a negative quantity, which is either prime to $2p^3 - 9qp + 27r$, or else so related to it, that the greatest common measure of the cube of the first and the square of the second is a perfect sixth power.

I now proceed to show the converse, that if these two conditions be both satisfied (and it will appear in the course of the inquiry that the first does *not* involve the second), the roots cannot help being all rational.

It is evident that the two conditions in question are tantamount to supposing that the roots of the proposed equation are linearly connected with those of another $z^3 - Qz - R = 0$ (by virtue of the assumption $3x = kz + p$), where Q may be considered as prime to R ; and where $4Q^3 - 27R^2$ is a perfect square.

Let now $4Q^3 - 27R^2 = D^2$, then $D^2 + 27R^2 = 4Q^3$, or $D^2 + 3(3R)^2 = 4Q^3$.

Here, as Q is prime to R , D can have no common measure but 3, with $3R$.

Firstly, let Q be prime to $3R$.

Then putting $f^2 + 3g^2 = Q^3$, the complete solution of the equation immediately preceding is contained in the two systems:

$$\text{1st. } D = 2f, \quad 3R = 2g.$$

$$\text{2nd. } D = (f \pm 3g), \quad 3R = f \mp g,$$

and for both systems,

$$f \pm g\sqrt{-3} = \{h \pm 3k\sqrt{-3}\}^3.$$

p. 105

The second system must therefore be rejected, for g evidently contains 3, and therefore $f = 3R \pm g$ will contain 3, and therefore D and therefore Q will do the same, contrary to supposition.

Hence

$$\begin{aligned} \sqrt[3]{\left[\frac{R}{2} \pm \sqrt{\left\{-\left(\frac{Q^3}{27} - \frac{R^2}{4}\right)\right\}}\right]} &= \sqrt[3]{\left\{\frac{R}{2} \pm \frac{D}{2} \sqrt{\left(-\frac{1}{27}\right)}\right\}} \\ &= \sqrt[3]{\left\{\frac{g}{3} \pm f \sqrt{\left(-\frac{1}{27}\right)}\right\}} \\ &= \mp \frac{1}{3\sqrt{-3}} \sqrt[3]{\{f \pm g\sqrt{-3}\}} \\ &= -K \pm \frac{h}{3\sqrt{-3}} = \lambda \pm \mu\sqrt{-3}; \end{aligned}$$

and the three roots of the equation being

$$\begin{cases} \{\lambda + \mu\sqrt{-3}\} + \{\lambda - \mu\sqrt{-3}\}, \\ \frac{1 \pm \sqrt{-3}}{2} \{\lambda + \mu\sqrt{-3}\} + \frac{1 \mp \sqrt{-3}}{2} \{\lambda - \mu\sqrt{-3}\}, \end{cases}$$

will evidently be all rational, which of course includes the necessity of their being also integer.

Again, secondly, if we suppose that Q does contain 3, D^2 will contain 27, and consequently D will contain 9; and we shall have

$$R^2 + 3 \left(\frac{D}{9} \right)^2 = 4 \left(\frac{Q}{3} \right)^3.$$

Here R being prime to $\frac{D}{9}$, it may be shown, as in the last case, that the complete solution is

$$\frac{R}{2} \pm \frac{D}{18} \sqrt{-3} = \{h \pm k\sqrt{-3}\}^3;$$

consequently

$$\sqrt[3]{\left\{ \frac{R}{2} \pm \sqrt{\left(\frac{R^2}{4} - \frac{Q^3}{27} \right)} \right\}} = h \pm k\sqrt{-3};$$

and the three roots of the equation are

$$2h, \quad h - 3k, \quad h + 3k$$

respectively, and are therefore all rational.

Here it may be observed that the condition of R being an even number, which we know, *à priori*, is the case when all the roots are rational, is p. 106

involved in the two more general conditions already expressed. It will now be evident that the first condition by no means involves the second, as it is perfectly easy to satisfy the equation $f^2 + 3g^2 = Q^3$ without supposing anything relative to k , the common measure of f, g, Q , except that it be itself of the form $\lambda^2 + 3\mu^2$, which will give

$$\left(\frac{f}{k} \right)^2 + 3 \left(\frac{g}{k} \right)^2 = (\lambda^2 + 3\mu^2)(r^2 + 3s^2)^3,$$

an equation which can be solved in rational terms for all values of λ, μ, r, s ; and consequently the product of the squares of the differences of the roots may be a square, and at the same time the roots themselves may be irrational⁵⁰.

⁵⁰Thus then it appears that the total rationality of the roots of the equation $x^3 - qx - r = 0$ may be determined by a direct method without having recourse to the method of divisors to determine the roots themselves; the two conditions being that $4q^3 - 27r^2$ shall be a perfect square, and the greatest common measure of q^3 and r^2 a perfect sixth power.

I believe it will be found on inquiry that the equation $x^n - qx + r = 0$ will always have two rational roots if

$$(n-1)^{n-1} \cdot q^n - n^n \cdot r^{n-1}$$

be a complete square, provided that q be prime to r .

Furthermore, viewing the striking analogy of the general nature of the conditions of rationality already obtained, to those which serve to determine the reality of the roots of equations, I am strongly of opinion that a theorem remains to be discovered, which will enable us to pronounce on the existence of integer, as Sturm's theorem on that of possible roots of a complete equation of any degree: the analogy of the two cases fails however in this respect, that while imaginary roots enter an equation in pairs, irrational roots are limited to entering in groups, each containing two or MORE.

19.

AN ACCOUNT OF A DISCOVERY IN THE THEORY OF NUMBERS RELATIVE TO THE EQUATION $Ax^3 + By^3 + Cz^3 = Dxyz$

[Philosophical Magazine, XXXI. (1847), pp. 189–191]

FIRST GENERAL THEOREM OF TRANSFORMATION.

p. 107

If in the equation

$$Ax^3 + By^3 + Cz^3 = Dxyz, \quad (1)$$

A and B are equal, or in the ratio of two cube numbers to one another, and if $27ABC - D^3$ (which I shall call the Determinant) is free from all single or square prime positive factors of the form $6n + 1$, but without exclusion of cubic factors of such form, and if A and B are each odd, and C the double or quadruple of an odd number, or if A and B are each even and C odd, then, I say, the given equation may be made to depend upon another of the form

$$A'u^3 + B'v^3 + C'w^3 = D'uvw;$$

where

$$A'B'C' = ABC, \quad D' = D, \quad uvw = \text{some factor of } z.$$

The following are some of the consequences which I deduce from the above theorem. In stating them it will be convenient to use the term Pure Factorial to designate any number into the composition of which no single or square prime positive factor of the form $6n + 1$ enters.

The equations

$$x^3 + y^3 + 2z^3 = Dxyz, \quad x^3 + y^3 + 4z^3 = Dxyz, \quad 2x^3 + 2y^3 + z^3 = Dxyz,$$

are insoluble in integer numbers, provided that the Determinant in each case is a Pure Factorial.

p. 108

The equation

$$x^3 + y^3 + Az^3 = 9Bxyz$$

is insoluble in integer numbers, provided that the Determinant, for which in this case we may substitute $A - 27B^3$, is a pure factorial whenever A is of the form $9n \pm 1$, and equal to $2p^{3i \pm 1}$ or $4p^{3i \pm 1}$, p being any prime number whatever.

I wish however to limit my assertion as to the insolubility of the equations above given. The theorem from which this conclusion is deduced does not preclude the possibility of two of the three quantities x, y, z being taken positive or negative *units*, either in the given equation itself or in one or the other of those into which it may admit of being transformed. Should such values of two

of the variables afford a particular solution, then instead of affirming that the equations are insoluble, I should affirm that the *general solution* can be obtained by equations in finite differences⁵¹.

SECOND GENERAL THEOREM OF TRANSFORMATION.

The equation

$$f^3x^3 + g^3y^3 + h^3z^3 = Kxyz \quad (2)$$

may always be made to depend upon an equation of the form

$$Au^3 + Bv^3 + Cw^3 = Duvw,$$

where

$$ABC = R^3 - S^3, \quad D = 3R;$$

and $uvw = \text{some factor of } fx + gy + hz.$

$$R \quad \text{representing } K + 6fgh, \quad S \quad K - 3fgh.$$

p. 109

I have not leisure to show the consequences of this theorem of transformation in connexion with the one first given, but shall content myself with a single numerical example of its applications:

$$x^3 + y^3 + z^3 = -6xyz$$

may be made to depend on the equation

$$u^3 + v^3 + w^3 = 0,$$

and is therefore insoluble.

It is moreover apparent that the Determinant of equation (2) transformed is in general $-27R^3$, and is therefore always a Pure Factorial, and consequently the equation

$$f^3x^3 + g^3y^3 + h^3z^3 = Kxyz$$

will be itself insoluble, being convertible into an insoluble form, provided that $K + 6fgh$ is divisible by 9, and provided further that $(K + 6fgh)^3 - (K - 3fgh)^3$

⁵¹Take for instance the equation $x^3 + y^3 + 2z^3 = 3xyz$. The Determinant 27.25 is a Pure Factorial: consequently if the solution be possible, since in this case the transformed must be identical with the given equation, this latter must be capable of being satisfied by making x and y positive or negative units. Upon trial we find that $x = 1, y = 1, z = 2$ will satisfy the equation. I believe, but have not fully gone through the work of verification, that these are the only possible values (prime to one another) which will satisfy the equation. Should they not be so, my method will infallibly enable me to discover and to give the law for the formation of all the others.

Here, then, under any circumstances, is an example, the first on record, of the complete resolution of a numerical equation of the third degree between three variables.

belongs to the form m^3Q , where Q is of the form $9n \pm 1$, and also of one or the other of the two forms $2p^{3i \pm 1}, 4p^{3i \pm 1}$, p being any prime number whatever.

Pressing avocations prevent me from entering into further developments or simplifications at this present time.

It remains for me to state my reasons for putting forward these discoveries in so imperfect a shape. They occurred to me in the course of a rapid tour on the continent, and the results were communicated by me to my illustrious friend M. Sturm in Paris, who kindly undertook to make them known on my part to the Institute.

Unfortunately, in the heat of invention I got confused about the law of oddness and evenness, to which the coefficients of the given equation are in the first theorem *generally* (in order for the successful application of my method as far as it is yet developed) required to be subject. I stated this law erroneously, and consequently drew erroneous conclusions from my Theorems of Transformation, which I am very anxious to seize the earliest opportunity of correcting. I venture to flatter myself that as opening out a new field in connexion with Fermat's renowned Last Theorem, and as breaking ground in the solution of equations of the third degree, these results will be generally allowed to constitute an important and substantial accession to our knowledge of the Theory of Numbers.

20.

ON THE EQUATION IN NUMBERS $Ax^3 + By^3 + Cz^3 = Dxyz$, AND ITS ASSOCIATE SYSTEM OF EQUATIONS

[*Philosophical Magazine*, XXXI. (1847), pp. 293–296]

p. 110

In the last Number of this *Magazine* I gave an account of a remarkable transformation to which the equation

$$Ax^3 + By^3 + Cz^3 = Dxyz$$

is subject when certain conditions between the coefficients A, B, C, D are satisfied; which conditions I shall begin by expressing with more generality and precision than I was enabled to do in my former communication.

1. Two of the quantities A, B, C are to be to one another in the ratio of two cubes.

2. $27ABC - D^3$ must contain no positive prime factor whatever of the form $6n + 1$. I erred in my former communication in not excluding cubic factors of this form.

3. If 2^m is the highest power of 2 which enters into ABC , and 2^n the highest power of 2 which enters into D , then either m must be of the form $3n \pm 1$, or if not, then m must be greater than $3n$.

These three conditions being satisfied, the given equation can always be transformed into another,

$$A'u^3 + B'v^3 + C'w^3 = D'uvw,$$

where

$$A'B'C' = ABC, \quad D' = D, \quad uvw = \text{a factor of } z.$$

The consequence of this is, as stated in my former paper, that wherever A, B, C, D , besides satisfying the conditions above stated, are taken so as likewise to satisfy the condition,—firstly, of ABC being equal to $2^{3m \pm 1}$, or secondly, of ABC being equal to $2^{3m \pm 1} \cdot p^{3n \pm 1}$, provided in the second case that ABC is of the form $9m \pm 1$, and that D is divisible by 9, p being in both cases a prime, then the given equation will be *generally* insoluble.

p. 111

And I am now enabled to add that the *only* solution of which it will in any case admit, is the solitary one found by making two of the terms Ax^3, By^3, Cz^3 equal to one another; so that, for instance, if the given equation should be of the form

$$x^3 + y^3 + ABCz^3 = Dxyz,$$

then the above conditions being satisfied, the one solitary solution of which the equation can possibly admit, is $x = 1, y = 1$,

$$Az^3 - Dz + 2 = 0,$$

which may or may not have possible roots. I call this a *solitary* or singular solution, because it exists alone and no other solution can be deduced from it; whereas in general I shall show that any one solution of the equation

$$Ax^3 + By^3 + Cz^3 = Dxyz$$

can be made to furnish an infinity of other solutions independent of the one supposed given, that is, not reducible thereto by expelling a common factor from the new system of values of x, y, z deduced from the given system.

The following is the Theorem of Derivation in question:

Let

$$A\alpha^3 + B\beta^3 + C\gamma^3 = D\alpha\beta\gamma.$$

Then if we write

$$F = A\alpha^3, \quad G = B\beta^3, \quad H = C\gamma^3,$$

and make

$$x = F^2G + G^2H + H^2F - 3FGH, \quad y = FG^2 + GH^2 + HF^2 - 3FGH,$$

$$z = \frac{1}{D}\{F^3 + G^3 + H^3 - 3FGH\},$$

or

$$= \alpha\beta\gamma\{F^2 + G^2 + H^2 - FG - FH - GH\},$$

we shall have

$$x^3 + y^3 + ABCz^3 = Dxyz.$$

I am hence enabled to show that whenever $x^3 + y^3 + Az^3 = Dxyz$ is insoluble, there will be a whole family of allied equations equally insoluble. For instance, because $x^3 + y^3 + z^3 = 0$ is insoluble in integer numbers, I know likewise that

$$x^6 + y^6 + z^6 = x^3y^3 + x^3z^3 + y^3z^3$$

$$x^6 + y^6 + z^6 = x^3y^3 + x^3z^3 - 2y^3z^3$$

are each equally insoluble.

p. 112

In fact

$$\begin{aligned} & (x^3 + y^3 + z^3) \times (x^6 + y^6 + z^6 - x^3y^3 - x^3z^3 - y^3z^3) \\ & \times (x^6 + y^6 + z^6 - x^3y^3 - x^3z^3 + 2y^3z^3) \\ & \times (x^6 + y^6 + z^6 - y^3z^3 - y^3x^3 + 2x^3z^3) \\ & \times (x^6 + y^6 + z^6 - x^3z^3 - z^3y^3 + 2y^3x^3) \\ & = u^3 + v^3 + w^3, \end{aligned}$$

where u, v, w are rational integral functions of x, y, z .

Hence each of the factors must be incapable of becoming zero⁵².

As a particular instance of my general theory of transformation and elevation, take the equation

$$x^3 + y^3 + 2z^3 = Mxyz.$$

Then, with the exception of the singular or solitary solution $x = 1, y = 1$, of which I take no account, I am able to affirm that for all values of M between 7 and -6 , both inclusive, with the exception of $M = -2$, the equation is insoluble in integer numbers.

Take now the equation where $M = -2$, namely

$$x^3 + y^3 + 2z^3 + 2xyz = 0.$$

One particular solution of this is

$$x = 1, \quad y = -1, \quad z = 1.$$

Another, which I shall call the second⁵³, is

$$x = 1, \quad y = 3, \quad z = -2.$$

From the first solution I can deduce in succession the following:

$$\begin{array}{lll} x = 11, & y = 5, & z = -7, \\ x = -793269121, & y = 1179490001, & z = -1189735855, \\ \&c. & \&c. & \&c. \end{array}$$

From the second,

$$\begin{array}{lll} x = -10085, & y = 8921, & z = -8442, \\ x = \&c. & y = \&c. & z = \&c. \end{array}$$

As another example, take the equation

$$x^3 + y^3 + 6z^3 = 6xyz.$$

p. 113

One solution of the transformed equation

$$u^3 + 2v^3 + 3w^3 = 6uvw$$

is evidently

$$u = 1, \quad v = 1, \quad w = 1.$$

⁵²It is however sufficiently evident from their intrinsic form, which may be reduced to $\frac{1}{4}(M^2 + 3N^2)$, that this impossibility exists for all the factors except the first.

⁵³See Postscript.

Hence I can deduce an infinite series of solutions of the given equation, of which the first in order of ascent will be

$$x = 5, \quad y = 7, \quad z = 3.$$

Again, the lowest possible solution in integers of the equation

$$x^3 + y^3 + 6z^3 = 0$$

will be

$$x = 17, \quad y = 37, \quad z = -21.$$

The equation

$$x^3 + y^3 + 9z^3 = 0$$

admits of the solutions

$$\begin{aligned} x &= 1, & y &= 2, & z &= -1, \\ x &= -271, & y &= 919, & z &= -438. \end{aligned}$$

I trust that my readers will do me the justice to believe that I am in possession of a strict demonstration of all that has been here advanced without proof. Certain of the writer's friends on the continent have, in their comments upon one of his former papers which appeared in this *Magazine*, complimented his powers of divination at the expense of his judgment, in rather gratuitously assuming that the author of the Theory of Elimination was unprovided with the demonstrations, which he was too inert or too beset with worldly cares and distractions to present to the public in a sufficiently digested form. The proof of whatever has been here advanced exists not merely as a conception of the author's mind, but fairly drawn out in writing, and in a form fit for publication.

P.S. It must not be supposed that the two primary or basic solutions above given of the equation

$$x^3 + y^3 + 2z^3 + 2xyz = 0,$$

namely,

$$\begin{aligned} x &= 1, & y &= -1, & z &= 1, \\ x &= 1, & y &= 3, & z &= -2, \end{aligned}$$

are independent of one another. The second may be derived from the first, as I shall show in a future communication. In fact there exist *three* independent processes, by combining which together, one particular solution may be made to give rise to an infinite series of infinite series of infinite series of correlated solutions, which it may possibly be discovered contain between them the *general* complete solution of the equation

$$x^3 + y^3 + Az^3 = Dxyz.$$

21.

ON THE GENERAL SOLUTION (IN CERTAIN CASES) OF THE EQUATION
 $x^3 + y^3 + Az^3 = Mxyz$, &c.

[*Philosophical Magazine*, XXXI. (1847), pp. 467-471]

p. 114

I shall restrict the enunciation of the proposition I am about to advance to much narrower limits than I believe are necessary to the truth, with a view to avoid making any statement which I may hereafter have occasion to modify. Let us then suppose in the equation

$$x^3 + y^3 + Az^3 = Mxyz$$

that A is a *prime* number, and that $27A - M^3$ is *positive*, but exempt from positive prime factors of the form $6i + 1$. Then I say, and have succeeded in demonstrating, that all the possible solutions in integer numbers of the given equation may be obtained by explicit processes from one particular solution or system of values of x, y, z , which may be called the Primitive system.

This system of roots or of values of x, y, z is that system in which the value of the greatest of the three terms $x, y, A^{\frac{1}{3}} \cdot z$ (which may be called the Dominant) is the least possible of all such dominants. I believe that in general the system of the least Dominant is identical with the system of the least Content, meaning by the latter term the product of the three terms out of which the Dominant is elected. I proceed to show the law of derivation.

To express this simply, I must premise that I shall have to employ such an expression as $S' = \phi(S)$ to indicate, not that a certain quantity, S' , is a function of S , but that a certain system of quantities disconnected from one another, denoted by S' , are severally functions of a certain other system of quantities denoted by S ; and, as usual, I shall denote $\phi\phi S$ by $\phi^2 S$, $\phi\phi^2 S$ by $\phi^3 S$, and so forth.

Let now P be the Primitive system of solution of the equation

$$x^3 + y^3 + Az^3 = Mxyz,$$

P denoting a certain system of values of and written in the order of the letters x, y, z , which may always be found by a limited number of trials (provided that the equation admits of any solution).

p. 115

That this is the case is obvious, since we have only to give the Dominant every possible value from the integer next greatest to $A^{\frac{1}{3}}$ upwards, and combine the values of x^3, y^3, Az^3 so that none shall ever exceed at each step the cube of such dominant, and we must at last, if there *exist any solution*, arrive at the System of the Least Dominant.

Now, every system of solution is of one or the other of two characters. Either x and y must be odd and z even, or x and y must be one odd and the other even and z odd. That all three should be odd is inconsistent with the given conditions as to A being odd and M even; and if all three were even, by driving out the common factor we should revert to one or the other of the foregoing cases.

The systems of solution where z is even may be termed Reducible, those where z is odd Irreducible. Let ϕ denote a certain symbol of transformation hereafter to be explained.

Then the Reducible systems of the first order may be expressed by

$$\phi P, \phi^2 P, \phi^3 P, \text{ ad infinitum};$$

or in general by $\phi^{n_1} P$, n_1 being absolutely arbitrary. I will anticipate by stating that the function ϕ involves no *variable* constants; that is to say, $\phi(S)$ may be found explicitly from S without any reference to the particular equation to which S belongs. Let now ψ denote another symbol of transformation, also hereafter to be defined, and differing from ϕ insofar as it does involve as *constants* the three values of x, y, z contained in P : then the general representations of Irreducible systems of the first order will be denoted by $\psi\phi^{n_1} P$.

It is proper to state here that the symbol ψ is ambiguous; and $\psi\phi^{n_1} P$, when P and n_1 are given, will have two values, according to the way in which the terms represented by P are compared with x, y, z in the given equation

$$x^3 + y^3 + Az^3 = Mxyz;$$

for it is obvious that if $x = a, y = b, z = c$ satisfies the equation, so likewise will

$$x = b, \quad y = a, \quad z = c.$$

Each however of these values of $\psi\phi^{n_1} P$ gives a solution of the kind above designated.

Proceeding in like manner as before, the Reducible system of the second order may be designated by $\phi^{n_2} \cdot \psi\phi^{n_1} \cdot P$, the Irreducible by $\psi\phi^{n_2} \cdot \psi\phi^{n_1} \cdot P$; and in general *every possible* system of values of x, y, z satisfying the proposed equation, in which z is even, is comprised under the form

$$\phi^{n_r} \cdot \psi\phi^{n_{r-1}} \cdot \psi \dots \psi\phi^{n_1} \cdot P;$$

p. 116

and every possible system of such values, in which z is odd, is comprised under the form

$$\psi\phi^{n_r} \cdot \psi\phi^{n_{r-1}} \cdot \psi \dots \psi\phi^{n_1} \cdot P :$$

the quantities $n_1, n_2 \dots n_r$ being of course all independent of one another, and unlimited in number and value.

Thus then we may be said to have the general solution of the given equation in the same sense as an arbitrary sum of terms, each of a certain form, is in certain cases accepted as the complete solution of a partial differential equation.

As regards the value of the symbols ψ and ϕ , ϕ indicates the process by which a, b, c becomes transformed into α, β, γ , the relations between the two sets of elements being contained in the following equations:

$$\begin{aligned} a' &= a^3, & b' &= b^3, & c' &= Ac^3, \\ \alpha &= a'^2b' + b'^2c' + c'^2a' - 3a'b'c', \\ \beta &= a'b'^2 + b'c'^2 + c'a'^2 - 3a'b'c', \\ \gamma &= abc\{a'^2 + b'^2 + c'^2 - a'b' - a'c' - b'c'\}. \end{aligned}$$

Next, as to the effect of the Duplex symbol ψ . Let e, g, ι be the elements of the Primitive system P : ι being the value of z and e, g of x and y taken in either mode of combination, each with each, which satisfy the proposed equation

$$x^3 + y^3 + Az^3 = Mxyz.$$

Let l, m, n represent any system S ,

$$\lambda, \mu, \nu \text{ represent any system } \psi(S),$$

ψS has two values, which we may denote by $\psi'S, '\psi S$ respectively, and accentuating the elements λ, μ, ν accordingly to correspond, we shall have

$$\begin{aligned} \lambda' &= 3gm(gl - em) + 3A\iota n(\iota l - en) - M(gl^2 - e^2lm), \\ \mu' &= 3A\iota n(\iota m - gl) + 3el(em - gl) - M(e\iota m^2 - g^2lm), \\ \nu' &= 3el(en - \iota l) + 3gm(gn - \iota m) - M(egn^2 - \iota^2lm) : \end{aligned}$$

we have then

$$\psi'S = \lambda', \mu', \nu',$$

and in like manner

$$' \psi S = '\lambda, '\mu, '\nu,$$

$' \psi S$ being derived from $\psi'S$ by the mere interchange of e and g one with the other.

p. 117

I have stated that every possible solution of the proposed equation comes under one or the other of the orders, infinite in number and infinite to the power of infinity in variety of degree, above given: this is not strictly true, unless we understand that all systems of solution are considered to be equivalent which differ only in a multiplier common to all three terms of each; that is to say, which may be rendered identical by the expulsion of a common factor. So that $m\alpha, m\beta, m\gamma$ as a system is treated as identical with α, β, γ , which of course

substantially it is; and it should be remarked that there is nothing to prevent the operations denoted by ϕ and ψ introducing a common factor into the systems which they serve to generate, and the latter in particular will have a strong tendency so to do.

I believe that this theorem may be extended with scarcely any modification to the case where A , instead of being a prime, is any power of the same, and to suppositions still more general. I believe also that, subject to certain very limited restrictions, the theorem may prove to apply to the case where the determinant $27A - M^3$ becomes negative.

The peculiarity of this case which distinguishes it from the former, is that it admits of all the three variables x, y, z in the equation

$$x^3 + y^3 + Az^3 = Mxyz$$

having the same sign, which is impossible when the determinant is positive; or in other words, the curve of the third degree represented by the equation

$$Y^3 + X^3 + 1 = \frac{M}{A^{\frac{1}{3}}}XY$$

(in which I call the coefficient of XY the characteristic), which, as long as the quantity last named is less than 3, is a single continuous curve extending on both sides to infinity, as soon as the characteristic becomes equal to 3 assumes to itself an isolated point, the germ of an oval or closed branch, which continues to swell out (always lying apart from the infinite branch) as the characteristic continues indefinitely to increase.

I ought not to omit to call attention to the fact that the theorem above detailed is always applicable to the case of the equation

$$x^3 + y^3 + Az^3 = 0,$$

when A is *any* power of a prime number *not* of the form $6i + 1$; in other words, the above always belongs to the class of equations having Monogenous solutions, which for the sake of brevity may be termed themselves Monogenous Equations⁵⁴. p. 118

On the probable existence of such a class of equations I hazarded a conjecture at the conclusion of my last communication to this *Magazine*. As I hope shortly to bring out a paper on this subject in a more complete form, I shall content myself at this time with merely stating a theorem of much importance to the completion of the theory of insoluble and of Monogenous equations of the third degree; to wit, that the equation in integers

$$a(x^3 + y^3 + z^3) + c(x^2y + y^2z + z^2x + xy^2 + yz^2 + zx^2) + exyz = 0$$

⁵⁴Thus the equation $x^3 + y^3 + 9z^3 = 0$ alluded to by Legendre is Monogenous, and the Primitive system of solution is $x = 1, y = 2, z = -1$, from which every other possible solution in Integers may be deduced.

may always be transformed so as to depend upon the equation

$$fu^3 + gv^3 + hw^3 = (6a - e)uvw,$$

wherein

$$fgh = ae^2 - (c^2 + 3a^2)e + 9a^2 - 3ac^2 - 2c^3.$$

By means of the above theorem, among other and more remarkable consequences, we are enabled to give a theory of the irresoluble and monogenous cases of the equation

$$x^3 + y^3 + m^3z^3 = Mxyz,$$

when m is some power of 2, or of certain other numbers.

22.

ON THE INTERSECTIONS, CONTACTS, AND OTHER CORRELATIONS OF TWO CONICS EXPRESSED BY INDETERMINATE COORDINATES

[Cambridge and Dublin Mathematical Journal, v. (1850), pp. 262–282]

p. 119

Let $U = 0$, $V = 0$ be two homogeneous equations of the second degree with real coefficients, between the same three variables ξ, η, ζ .

The direct and most general mode of determining the intersections of the conics expressed by these equations would be to make

$$a\xi + b\eta + c\zeta = t, \quad a'\xi + b'\eta + c'\zeta = u :$$

eliminating ξ, η, ζ between the four equations in which they appear, there results a biquadratic equation between t and u . The nature of the intersections will depend upon the nature of the roots of this biquadratic; and thus the conditions may be expressed analytically, which will represent the several cases of all the intersections being real or all imaginary, or one pair real and the other imaginary. These analytical conditions will depend upon the signs of certain functions of the coefficients of the given and the assumed equations being of an assigned character; my endeavour has been to obtain conditions of a character perfectly symmetrical and free from the coefficients arbitrarily introduced.

In this research I have only partially succeeded, but the method employed, and some of the collateral results, will, I think, be found of sufficient interest to justify their appearance in the pages of this *Journal*.

Adopting Mr Cayley's excellent designation, let the four points of intersection of the two conics be called a quadrangle. This quadrangle will have three pairs of sides; the intersections of each pair, from principles of analogy, I call the vertices of the quadrangle. Then, inasmuch as the four

p. 120

sets of ratios $\xi : \eta : \zeta$, corresponding with the four sets of the ratio $t : u$, must be so related that we may always make

$$\begin{aligned} \frac{\xi_1}{\zeta_1} &= a + b\sqrt{-1}, & \frac{\eta_1}{\zeta_1} &= c + d\sqrt{-1}, \\ \frac{\xi_2}{\zeta_2} &= a - b\sqrt{-1}, & \frac{\eta_2}{\zeta_2} &= c - d\sqrt{-1}, \\ \frac{\xi_3}{\zeta_3} &= \alpha + \beta\sqrt{-1}, & \frac{\eta_3}{\zeta_3} &= \gamma + \delta\sqrt{-1}, \\ \frac{\xi_4}{\zeta_4} &= \alpha - \beta\sqrt{-1}, & \frac{\eta_4}{\zeta_4} &= \gamma - \delta\sqrt{-1}, \end{aligned}$$

we may easily draw the following conclusions.

If all the four points of the quadrangle of intersection are real, the three vertices and the three pairs of sides are all real. If only two points of the quadrangle are real, one vertex and one of the three pairs of sides will be real; the other two vertices and two pairs of sides being imaginary. If all four points of the quadrangle are unreal, one pair of sides will be real and the other two pairs imaginary, as in the last case; but all the three vertices will remain real, as in the first case. Hence we have a direct and simple criterion for distinguishing the case of *mixed* intersection from intersection wholly real or wholly imaginary; namely, that the cubic equation of the roots of which the coordinates of the vertices are real linear functions shall have a *pair of imaginary roots*. This is the sole and unequivocal condition required.

The equation in question is, or ought to be, well known to be the determinant in respect to ξ, η, ζ of $\lambda U + \mu V$. In fact, if we write

$$\begin{aligned} U &= a\xi^2 + b\eta^2 + c\zeta^2 + 2a'\eta\zeta + 2b'\zeta\xi + 2c'\xi\eta, \\ V &= \alpha\xi^2 + \beta\eta^2 + \gamma\zeta^2 + 2\alpha'\eta\zeta + 2\beta'\zeta\xi + 2\gamma'\xi\eta, \\ \lambda U + \mu V &= (a\lambda + \alpha\mu)\xi^2 + \&zc. \\ &= A\xi^2 + B\eta^2 + C\zeta^2 + 2A'\eta\zeta + 2B'\zeta\xi + 2C'\xi\eta, \end{aligned}$$

the ratios of the coordinates ξ, η, ζ of the vertex of $\lambda U + \mu V$ may easily be shown to be identical with

$$AB - C'^2 : C'A' - B'B : B'C' - A'A,$$

and will be real or imaginary as $\lambda : \mu$ is one or the other.

If then the cubic equation in $\lambda : \mu$, namely, $\square_{\xi\eta\zeta}(\lambda U + \mu V) = 0$, has a pair of imaginary roots, that is, if

$$\square_{\lambda\mu}\square_{\xi\eta\zeta}(\lambda U + \mu V)$$

is a positive quantity, the intersections of U and V are of a mixed kind, that is, the two conics have two real points in common.

p. 121

I may remark here, *en passant*, that if we form the biquadratic equation in t and u , $\phi(t, u) = 0$ from the equations

$$U = 0, \quad V = 0, \quad a\xi + b\eta + c\zeta = t, \quad a'\xi + b'\eta + c'\zeta = u,$$

and if any reducing cubic of this equation be $P(\theta, \omega) = 0$, the determinant of $P(\theta, \omega)$ must, from what has been shown above, be identical with

$$\square_{\lambda\mu}\square_{\xi\eta\zeta}(\lambda U + \mu V)$$

multiplied by some squared function of the extraneous coefficients

$$a, b, c; \quad a', b', c'.$$

If $\square\square(\lambda U + \mu V)$ is a negative quantity, it remains to distinguish between the cases of the conics intersecting really in four points or not at all.

The most obvious mode of proceeding to distinguish between purely real and purely imaginary intersections would be as follows. Let $\lambda_1, \mu_1; \lambda_2, \mu_2; \lambda_3, \mu_3$ be the three sets of values of λ, μ which satisfy the equation

$$\square(\lambda U + \mu V) = 0$$

and make

$$\begin{aligned} A_1 &= a\lambda_1 + \alpha\mu_1, & A_2 &= a\lambda_2 + \alpha\mu_2, & A_3 &= a\lambda_3 + \alpha\mu_3, \\ C_1 &= c\lambda_1 + \gamma\mu_1, & C_2 &= c\lambda_2 + \gamma\mu_2, & C_3 &= c\lambda_3 + \gamma\mu_3, \\ B'_1 &= b'\lambda_1 + \beta'\mu_1, & B'_2 &= b'\lambda_2 + \beta'\mu_2, & B'_3 &= b'\lambda_3 + \beta'\mu_3, \\ A_1C_1 - B'^2_1 &= e_1, & A_2C_2 - B'^2_2 &= e_2, & A_3C_3 - B'^2_3 &= e_3. \end{aligned}$$

Now if the equation

$$A\xi^2 + B\eta^2 + C\zeta^2 + 2A'\eta\zeta + 2B'\zeta\xi + 2C'\xi\eta = 0$$

represent a pair of straight lines, it may be thrown into the form

$$Au^2 + \frac{AC - B'^2}{A}v^2 = 0,$$

where u and v are linear functions of ξ, η, ζ , and the straight lines will be real or imaginary, according as $B'^2 - AC$ is positive or negative; hence one or else all of the quantities e_1, e_2, e_3 will be necessarily negative, and the intersections will be all real or all imaginary, according as all three are negative or only one is so. A cubic equation in e may be formed containing e_1, e_2, e_3 as its roots by eliminating between the equations

$$e = AC - B'^2, \quad \square(\lambda U + \mu V) = 0,$$

and the conditions for the reality of the intersections will be that all four coefficients of this cubic shall be of the same sign, which in reality amount only to two, since the first and last must in all cases have the same sign.

p. 122

The same objection however of want of symmetry and consequent irrelevancy and complexity attaches to this as much as to the method originally proposed. The following treatment of the question relieves the objection of want of symmetry as far as the coefficients of the same equation are concerned, but in its practical application necessitates an arbitrary and therefore unsymmetrical election to be made between the two sets of coefficients appertaining to the two equations. It is however, I think, too curious and suggestive to be suppressed.

I observe that if the four intersections are all real, an imaginary conic cannot be drawn through them; for the equation to an imaginary conic may always

be reduced to the form $Ax^2 + By^2 + Cz^2 = 0$, where A, B, C are all positive and can therefore have at utmost one real point. Consequently the case of total non-intersection is distinguishable from that of complete intersection by the peculiarity that in the one case μ may be so taken that $U + \mu V = 0$ shall represent an imaginary conic, that is, $U + \mu V$ will be a function whose sign never changes for real values of ξ, η, ζ , whereas in the latter case no value of μ will make $U + \mu V = 0$ the equation to an imaginary conic, and therefore $U + \mu V$ will have values on both sides of zero. On the other hand, it is obvious that an infinite number of real as well as unreal conics may be drawn through four imaginary points of intersection. Consequently if we make $U + \mu V = 0$ (supposing the intersections of U and V to be imaginary), there will be a range or ranges of values of μ consistent, and another range or ranges of values of μ inconsistent with real values of ξ, η, ζ ; in other words, $U + \mu V = 0$ treated as an equation between the four variables ξ, η, ζ, μ , will give one or more maxima or minima values of μ , in the case supposed, but no such values when the intersections are two or all of them real.

To determine these values of μ , let $d\mu = 0$; then we have

$$\frac{d}{d\xi}(U - \mu V) = 0, \quad \frac{d}{d\eta}(U - \mu V) = 0, \quad \frac{d}{d\zeta}(U - \mu V) = 0,$$

that is

$$\square_{\xi\eta\zeta}(U - \mu V) = 0.$$

In order that any value of μ found from this equation may be a maximum or minimum, Lagrange's condition requires that

$$\left(h \frac{d}{d\xi} + k \frac{d}{d\eta} + l \frac{d}{d\zeta}\right)^2 \mu$$

may be a function of unchangeable sign.

p. 123

Now

$$\frac{dU}{d\xi} = \mu \frac{dV}{d\xi} + V \frac{d\mu}{d\xi};$$

therefore since $d\mu = 0$,

$$\frac{d^2U}{d\xi^2} = \mu \frac{d^2V}{d\xi^2} + V \frac{d^2\mu}{d\xi^2}.$$

Hence

$$\frac{d^2\mu}{d\xi^2} = \frac{1}{V} \left(\frac{d}{d\xi}\right)^2 \{U - \mu V\};$$

similarly

$$\frac{d}{d\xi} \frac{d}{d\eta} = \frac{1}{V} \frac{d}{d\xi} \frac{d}{d\eta} \{U - \mu V\},$$

and so on. Making now as before

$$U = a\xi^2 + b\eta^2 + \&c., \quad V = \alpha\xi^2 + \beta\eta^2 + \&c.,$$

$$a - \mu\alpha = A, \quad b - \mu\beta = B, \quad \&c.,$$

the condition for μ , a root of $\square\{U - \mu V\} = 0$, giving μ a maximum or minimum, may be expressed by saying that

$$Ah^2 + Bk^2 + Cl^2 + 2A'kl + 2B'hl + 2C'hk$$

shall be unchangeable in sign for all real values of h, k, l .

The above quantity, by virtue of the equation $\square = 0$, is always the product of two linear functions. Hence we see, as above indicated, that if all these pairs are real, that is, if all the points of intersection of U and V are real, there is no maximum or minimum value of μ ; but if only one pair be real and the other two pairs be imaginary, that is, if all the four intersections are imaginary, then two of the values of μ , namely those corresponding to the imaginary pairs, are real maxima or minima values of μ , but the third is illusory.

Now I shall show that if $V = 0$ is a real conic, but the intersections of U and V are all unreal, the value of μ which makes $U + \mu V$ the product of real linear functions of ξ, η, ζ , is always one or the other extreme of the three values of μ which satisfy the equation

$$\square(U - \mu V) = 0.$$

Assume as the three axes of coordinates the three lines joining the vertices of the quadrangle each with each, the two non-intersecting conics may evidently be written under the form

$$U = c(x^2 + y^2) - e(y^2 + z^2) = 0, \quad V = -\gamma(x^2 + y^2) + \epsilon(y^2 + z^2) = 0;$$

p. 124

these equations being only other modes of writing

$$U = Ax^2 + By^2 + Cz^2, \quad V = A'x^2 + B'y^2 + C'z^2,$$

in which $A, B, C; A', B', C'$ will be real, because by hypothesis $\square(U + \mu V) = 0$ has all its roots real.

Hence x, y, z are linear functions of ξ, η, ζ and consequently, by a simple inference from a theorem of Prof. Boole⁵⁵, the roots of $\square_{\xi\eta\zeta}\{U + \mu V\}$ are identical with those of

$$\square_{xyz}\{U + \mu V\} = 0.$$

These latter are evidently

$$\frac{c}{\gamma}, \quad \frac{e}{\epsilon}, \quad \frac{c - e}{\gamma - \epsilon},$$

the third of which is the one which makes $U + \mu V$ the product of two real linears, for we have

$$\gamma U + cV = (c\epsilon - \gamma e)(y^2 + z^2),$$

⁵⁵See Postscript.

$$\begin{aligned}\epsilon U + eV &= (\epsilon c - e\gamma)(x^2 + y^2), \\ (\gamma - \epsilon)U + (c - e)V &= (c\epsilon - e\gamma)(z^2 - x^2)^{56}.\end{aligned}$$

Now

$$\frac{c}{\gamma} - \frac{c-e}{\gamma-\epsilon} = \frac{e\gamma - c\epsilon}{\gamma(\gamma-\epsilon)}, \quad \frac{e}{\epsilon} - \frac{c-e}{\gamma-\epsilon} = \frac{e\gamma - c\epsilon}{\epsilon(\gamma-\epsilon)};$$

and ϵ, γ are supposed to have the same sign, as otherwise V would be an unreal conic; hence the ascending or descending order of magnitudes of the three values of λ follows the scale

$$\frac{c}{\gamma}, \quad \frac{e}{\epsilon}, \quad \frac{c-e}{\gamma-\epsilon},$$

as was to be shown.

Imagine now lengths reckoned on a line corresponding to all values of μ from $-\infty$ to $+\infty$, and mark off upon this line by the letters A, B, C , the lengths corresponding with the three roots of $\square(U + \mu V) = 0$. Then observing that when $\mu = \pm\infty$, $U + \mu V$ is of the same nature as V , and is therefore a possible conic by hypothesis, and agreeing to understand by a possible and impossible region of μ , a range of values for which $U + \mu V$ corresponds to a possible and impossible conic respectively, one or the other of the annexed schemes will represent the circumstances of the case supposed:

$-\infty$ Poss. Reg. A Imposs. Reg. B Poss. Reg. C Poss. Reg. $+\infty$,

$-\infty$ Poss. Reg. A Poss. Reg. B Imposs. Reg. C Poss. Reg. $+\infty$.

But in either scheme it is essential to observe that the *middle* root of $\square(U + \mu V) = 0$ divides a possible from an impossible region; and therefore

p. 125

if we can find u, v , any two values lying between the first and second and second and third roots of the above equation arranged in order of their magnitude, one of the two equations $U + vV = 0$, $U + uV = 0$, will represent a possible and the other an impossible conic: one such couple of values may always be found by taking the roots of the quadratic equation

$$\frac{d}{d\mu}\square\{U + \mu V\} = 0.$$

Hence calling the two roots thereof m and M , we see (which is in itself a theorem) that one at least of the conics $U + mV = 0$, $U + MV = 0$, must be a possible conic, provided only that $V = 0$ be a possible conic: if both $U + mV$ and $U + MV$ are possible conics, the intersections of U and V are all real, and if not, not⁵⁷.

⁵⁶ $z^2 - x^2 = 0$ of course represents a real pair of lines.

⁵⁷ It must be well observed however that the possibility of the conics $U + mV$ and of $U + MV$ does not imply the reality of the intersections unless the conic V is known to be possible. For if V be impossible ϵ and γ have opposite signs, and therefore $\frac{c-\epsilon}{\gamma-\epsilon}$ is intermediate between $\frac{c}{\epsilon}$ and

The criteria for distinguishing possible from impossible conics being well known need not be stated in this place.

We may of course proceed analogously by forming the two conics $lU + V$, $LU + V$, where l and L are roots of

$$\frac{d}{d\lambda} \square \{\lambda U + V\} = 0$$

upon the supposition of $U = 0$ being a possible conic.

If either of the two U and V be not possible, their intersections are of course impossible, and the question is already decided.

It will be seen as pre-indicated that this method only fails in symmetry because of the choice between the couples m, M , and l, L . But moreover a perfect method for the discrimination of the two cases of unmixed intersection one from the other should (perhaps?) require the application of only a single test (in lieu of the two conditions which the above method supposes), over and above the condition which expresses the fact of the intersections being so unmixed. Such more perfect method I have not yet been able to achieve.

Another interesting question of intersections remains to be discussed, namely, supposing the two conics are known to be non-intersecting, how are we to ascertain if they are external to one another, or if one contains the other? In order to settle this point we must first establish a criterion for determining whether a given *point* is internal or external to a given conic; the point being in general said to be external when two real tangents can be drawn from it to the curve, and internal when this cannot be done.

p. 126

Let now

$$\phi(x, y, z) = ax^2 + by^2 + cz^2 + 2a'yz + 2b'zx + 2c'xy = 0,$$

be the equation to any conic: l, m, n the coordinates of any point. Let

$$\begin{aligned} A &= bc - a'^2, & B &= ca - b'^2, & C &= ab - c'^2, \\ A' &= aa' - b'c', & B' &= bb' - c'a', & C' &= cc' - a'b'. \end{aligned}$$

Then the reciprocal equation to the conic is

$$A\xi^2 + B\eta^2 + C\zeta^2 + 2A'\eta\zeta + 2B'\zeta\xi + 2C'\xi\eta = 0,$$

and in making $l\xi + m\eta + n\zeta = 0$, the ratios of ξ, η, ζ must be real if the tangents

$\frac{\gamma}{\epsilon}$, and the scheme for μ will be as here annexed:

$$-\infty \quad \text{Impossible.} \quad A \quad \text{Possible.} \quad B \quad \text{Possible.} \quad C \quad \text{Impossible.} \quad +\infty,$$

so that $U + mV$ and $U + MV$ will both represent possible conics.

drawn from l, m, n are real: this will be found to imply that the determinant

$$\begin{vmatrix} A & C' & B' & l \\ C' & B & A' & m \\ B' & A' & C & n \\ l & m & n & 0 \end{vmatrix}$$

shall be negative⁵⁸. This determinant may be shown⁵⁹ to be equal to the product of the determinant

$$\begin{vmatrix} a & c' & b' \\ c' & b & a' \\ b' & a' & c \end{vmatrix}$$

by the quantity

$$al^2 + bm^2 + cn^2 - 2a'mn - 2b'ln - 2c'lm,$$

that is, equal to $\phi(l, m, n) \times \square$.

Hence l, m, n is internal or external to $\phi(x, y, z)$ according as $\phi(l, m, n)$ and $\square\phi$ have the same or contrary sign.

If $\phi(l, m, n) = 0$, the point lies on the conic, and the point is neither internal nor external; if $\square\phi = 0$, the conic becomes a pair of straight lines, and no point can be said either to be within or without such a system. Hence our criterion fails, as it ought to do, just in the very two cases where the distinction vanishes. I believe that this criterion is here given for the first time. p. 127

To return to the two non-intersecting conics. Let us again throw them under the form

$$U = (x^2 + y^2) - e^2(z^2 + y^2), \quad V = k(x^2 + y^2) - k\epsilon^2(z^2 + y^2),$$

e and ϵ being real, that is, U and V being both functions corresponding to possible conics. Suppose U external to V ; then *any point* in U is an external point to V .

Take in U either of the two points represented by the equations $y = 0$, $x^2 = e^2z^2$; substituting these values of y and x , V becomes $k(e^2 - \epsilon^2)z^2$, and $\square V$ becomes $-k^3\epsilon^2(1 - \epsilon^2)$; therefore $(1 - \epsilon^2)(e^2 - \epsilon^2)$ must be positive, that is, ϵ^2 must be one of the extremes of the three values $1, e^2, \epsilon^2$. In like manner, if V

⁵⁸See theorem of the "Diminished Determinant" in Postscript to this paper.

⁵⁹As we know *à priori* by virtue of a theorem given by M. Cauchy, and which is included as a particular case in a theorem of my own, relating to Compound Determinants, that is, Determinants of Determinants, which will take its place as an immediate consequence of my fundamental Theorem given in a Memoir about to appear. The well-known rule for the multiplication of Determinants is also a direct and simple consequence from my theorem on Compound Determinants, which indeed comprises, I believe, in one glance, all the heretofore existing Doctrine of Determinants.

is external to U , e will be also one of the extremes of the same three quantities; and hence, if the two conics are mutually external, unity will be the middle magnitude of the group $e^2, 1, \epsilon^2$.

Now the three roots of $\square(V + \lambda U) = 0$, are

$$\lambda = -k, \quad \lambda = -k \frac{\epsilon^2}{e^2}, \quad \lambda = -k \frac{1 - \epsilon^2}{1 - e^2}.$$

Hence if U and V be without one another, or, as it may be termed, are extra-spatial, the third value of λ will be of a different sign from the first two; but if the two conics be co-spatial, that is, if one includes the other, all the three values of λ will have the same sign. Hence we have the following elegant criterion of co-spatiality of two possible conics expressed by the equations $U = 0, V = 0$, between indeterminate coordinates ξ, η, ζ ; the coefficients of the cubic function $\square_{\xi\eta\zeta}(\lambda U + \mu V)$ must give only changes or only continuations of sign.

If this test be not satisfied, it will remain to determine which of the two conics contains, and which is contained by the other. Let U contain V , then the order of magnitudes will be $1, e^2, \epsilon^2$; therefore $k \frac{1 - \epsilon^2}{1 - e^2}$ is greater than k , and therefore $k \frac{1 - \epsilon^2}{1 - e^2}$, which is that root of the equation $\square(V + \lambda U) = 0$ which is always one or the other of the extremes, is the *greatest* of the three. Hence the scheme for the impossible and possible regions of λ will be as below:

$$-\infty \quad \text{Poss.} \quad A \quad \text{Imposs.} \quad B \quad \text{Poss.} \quad C \quad \text{Poss.} \quad +\infty.$$

Hence if the two roots of $\frac{d}{d\lambda}\{V + \lambda U\} = 0$ be l and L , and of the two conics $V + lU = 0, V + LU = 0$, the former be the possible, and the latter the impossible one, U contains V or is contained in it according as l is greater or less than L . p. 128

Observe that if U and V be non-cospatial, so that the three values of μ in $\square(U + \mu V) = 0$ have not all the same sign and consequently zero lies between the greatest and least of them, it will not be necessary to make trial of the characters of the two curves $U + mV = 0$, and $U + MV = 0$, in order to ascertain whether U and V intersect or not; for it will be sufficient to find which of the two quantities m and M substituted for μ in $\square(U + \mu V)$ causes it to have the opposite sign to $\square(U + 0V)$, that is, $\square U$, and this one of the two it is, if either, which will make $U + \mu V$ an impossible conic, and will thus alone serve to determine whether the intersections of U and V are unreal, or the contrary.

It might be a curious question to consider whether, in a certain sense, conics not both possible may not be said to lie one within or without the other. Upon general logical grounds, I think it not improbable that two impossible conics might be discovered each to contain the other; but this is an inquiry which I have not had leisure to enter upon.

I have thus far supposed the roots of $\square(\lambda U + V) = 0$ to be all distinct from one another. I now approach the discussion of the contact of two conics, in which

event two or more of the roots will be equal. The condition for simple contact is evidently

$$\square_{\lambda\mu}\square_{\xi\eta\zeta}(\lambda U + \mu V) = 0.$$

The unpaired value of λ in $\square(\lambda U + V)$ makes $\lambda U + V$ an impossible pair of lines, and therefore, in the scheme for λ drawn as above, will separate the possible from the impossible region.

Whether the conics intersect in two real or two unreal points, besides the point of contact, will be known at once by ascertaining whether $U + \mu V = 0$ represents two real or two imaginary lines. If the latter, the two curves lie *dos-à-dos* or one within the other, according as the successions of sign in $\square(\lambda U + V)$ are all of the same kind or not; if they be all of the same kind, one will include the other, namely, U will include V if the equal roots are greater, and be included in it if they be less than the unequal one. This last conclusion however, it should be observed, is inferred upon the principle of continuity, by making two values of λ approach indefinitely near to one another, but cannot be strictly deduced from the equations given for U and V applicable to the general case, in which the axes of coordinates are the three axes joining the vertices; since these latter, in the case supposed, reduce to two only, and consequently such representation of U and V becomes illusory.

If all three values of λ are equal, the three vertices come together, and hence the two conics will have three consecutive points in common, that is, will have the same circle of curvature. On this supposition the two curves cut at the point of contact, and all four points of intersection are of course real.

p. 129

The classification of contacts between two conics may be stated as follows:

Simple contact = one case.

Second degree contact = two cases, namely, common curvature or double contact.

Third degree contact = one case, namely, contact in four consecutive points.

These four cases of course correspond to the several suppositions of there being two equal roots, three equal roots, two pairs of equal roots, or four equal roots in the biquadratic equation obtained between two variables by elimination performed in any manner between the given equations in the two conics.

The first species and the first case of the second species have been already disposed of. I proceed to assign the conditions appertaining to the second case of the second species, when U and V have a double contact.

Let A, A', B, B' be the two pairs of coincident points in which the conics are supposed to meet; either pair of lines $AB, A'B'$, and $AB', A'B$, becomes a coincident pair. Hence such a value of μ can be found as will make $U + \mu V$ the square of a linear function of ξ, η, ζ . If therefore we make $U + \mu V = W$, and

form the determinant

$$\begin{vmatrix}
 \frac{d^2W}{d\xi^2} & \frac{d^2W}{d\xi d\eta} & \frac{d^2W}{d\xi d\zeta} & p \\
 \frac{d^2W}{d\eta d\xi} & \frac{d^2W}{d\eta^2} & \frac{d^2W}{d\eta d\zeta} & q \\
 \frac{d^2W}{d\zeta d\xi} & \frac{d^2W}{d\zeta d\eta} & \frac{d^2W}{d\zeta^2} & r \\
 p & q & r & 0
 \end{vmatrix}$$

$$= Ap^2 + Bq^2 + Cr^2 + 2Fqr + 2Grp + 2Hpq,$$

where all the coefficients are quadratic functions of μ , and make

$$A = 0, \quad B = 0, \quad C = 0, \quad F = 0, \quad G = 0, \quad H = 0,$$

each of these six equations in μ will have one and the same root in common.

It is, however, enough to select any three; if these vanish together for any value of μ , the remaining three must also vanish. This is a simple application of a general law⁶⁰ which will appear in a forthcoming memoir on “Determinants and Quadratic Forms,” of which this paper is to be considered as an accidental episode.

p. 130

Take now any three of the six equations which for the sake of generality call $P = 0$, $Q = 0$, $R = 0$. The hypothesis of double contact requires that P and Q , Q and R , R and P shall have a factor in common; but these conditions are not sufficiently explicit for our present object, since P, Q, R might be of the form

$$\kappa(\lambda - a)(\lambda - b), \quad \kappa'(\lambda - b)(\lambda - c), \quad \kappa''(\lambda - c)(\lambda - a),$$

and would thus satisfy the conditions above stated, without P, Q, R having a common factor. A sufficient criterion is that $fQ + gR$ and P shall have a common factor for all values of f and g .

Let then the resultant of $fQ + gR$ and P be

$$Lf^2 + Mfg + Ng^2;$$

we must have

$$L = 0, \quad M = 0, \quad N = 0,$$

where L is the resultant of P and Q , N is the resultant of R and Q , and M is a new function, which if we call $Q = \phi(\lambda)$, $R = \psi(\lambda)$, and suppose a and b to be the two roots of $P = 0$, is easily seen to be equal to

$$\phi a \cdot \psi b + \phi b \cdot \psi a.$$

⁶⁰For statement of this law called the Homaloidal Law, see *Philosophical Magazine* of this month “On Certain Additions, &c.” [p. 150 below. Ed.]

This I call the connective of $P \cdot Q$ and $P \cdot R$.

L, M, N may conveniently be denoted by the forms

$$P \cdot Q, \quad P \cdot R, \quad Q \cdot P \cdot R.$$

We may now take more generally

$$aP + bQ + cR, \quad \alpha P + \beta Q + \gamma R,$$

which will have a factor in common for all values of $a, b, c, \alpha, \beta, \gamma$.

I am indebted to Mr Cayley for the remark that the resultant of these two functions is a new quadratic function, which, according to my notation just given, may be put under the form

$$\begin{aligned} &PQ(a\beta - b\alpha)^2 + QR(b\gamma - c\beta)^2 + RP(c\alpha - a\gamma)^2 \\ &+ PRQ(b\gamma - c\beta)(c\alpha - a\gamma) + QPR(c\alpha - a\gamma)(a\beta - b\alpha) \\ &+ RQP(a\beta - b\alpha)(b\gamma - c\beta). \end{aligned}$$

Ternary systems of the six coefficients formed upon the type of (PQ, QR, RP) , I call *complete* systems, because the three functions included in such a system equated severally to zero, imply that the remaining three coefficients are all zero. Such a system as (PQ, QR, RP) I term an *incomplete* ternary system as not drawing with it the like implication. Probably (?) we should find on investigation that PRQ, QPR, RQP , would also be an

p. 131

incomplete system, but that systems formed after the type of PRQ, RQ, RQP are complete. This however is only matter of conjecture, as I have been too much occupied with other things to enter upon the inquiry. The distinct types of ternary systems are altogether six in number, namely, four of a symmetrical species,

$$\begin{aligned} &PQ, \quad QR, \quad RP, \\ &PRQ, \quad QPR, \quad RQP, \\ &PQ, \quad PQR, \quad QR, \\ &PRQ, \quad RQ, \quad RQP; \end{aligned}$$

and two of an unsymmetrical species, namely,

$$\begin{aligned} &PQ, \quad PQR, \quad PR, \\ &PRQ, \quad RQ, \quad QPR.^{61} \end{aligned}$$

If instead of confining ourselves to three out of the six original quantities, $A, B, C; F, G, H$, we take them all into account, and write down the resultant of

$$aA + bB + cC + fF + gG + hH,$$

⁶¹ PQ, QR, RP , may be compared in a general way with the angles, and PRQ, QPR, RQP , with the sides of a triangle.

$$\alpha A + \beta B + \gamma C + \phi F + \chi G + \eta H;$$

we shall obtain a quadratic function of 15 variables (not however all independent) having 120 coefficients, all of which must be zero. It would be extremely interesting to determine how many *complete* ternary groups can be formed out of these 120 terms.

It will be recollected that we have assigned as the condition of contact in three consecutive points, that a certain cubic equation shall have all its roots real. Now, as well remarked by Mr Cayley, we cannot express this fact by less than three equations in integral terms of the coefficients. Thus if the cubic be written

$$a\lambda^3 + 3b\lambda^2 + 3c\lambda + d = 0,$$

we have as one of such ternary systems,

$$U = ac - b^2 = 0, \quad V = bd - c^2 = 0, \quad W = bc - ad = 0.$$

The significant parts of these equations are of course, however, capable of being connected by integral multipliers U', V', W' , such that

$$U'U + V'V + W'W = 0.$$

p. 132

Any number of functions U, V, W so related, I call *syzygetic* functions, and U', V', W' I term the *syzygetic multipliers*⁶². These in the case supposed are c, a, b , respectively.

In like manner it is evident that the members of any group of functions, more than two in number, whose nullity is implied in the relation of double contact, whether such group form a complete system or not, must be in syzygy.

Thus PQ, PQR, QR , must form a syzygy; nor is there any difficulty in assigning a system of multipliers to exhibit such syzygy. Calling $P = \phi(\lambda)$, $R = \psi(\lambda)$, a and b the two roots of $Q = 0$, I have found that

$$\{(\psi a)^2 + (\psi b)^2\}PQ - (\phi a \cdot \psi a + \phi b \cdot \psi b)PQR + \{(\phi a)^2 + (\phi b)^2\}QR = 0.$$

Again, if we take the *incomplete* system

$$(PQ), \quad (QR), \quad (RP),$$

it will be found that

$$L(QR) + M(RP) + N(PQ) = 0,$$

⁶²There will be in general various such systems of multipliers.

provided that, calling a, b, c, d, e, f , the roots of $P = 0, Q = 0, R = 0$, respectively, we make

$$\begin{aligned}
L &= (k_0 + k_1a + k_2a^2 + k_3a^3 + k_4a^4) \frac{(a-c)(a-d)(a-e)(a-f)}{a-b} \\
&\quad + (k_0 + k_1b + k_2b^2 + k_3b^3 + k_4b^4) \frac{(b-c)(b-d)(b-e)(b-f)}{b-a}, \\
M &= (k_0 + k_1c + k_2c^2 + k_3c^3 + k_4c^4) \frac{(c-a)(c-b)(c-d)(c-e)(c-f)}{c-d} \\
&\quad + (k_0 + k_1d + k_2d^2 + k_3d^3 + k_4d^4) \frac{(d-a)(d-b)(d-c)(d-e)(d-f)}{d-c}, \\
N &= (k_0 + k_1e + k_2e^2 + k_3e^3 + k_4e^4) \frac{(e-a)(e-b)(e-c)(e-d)}{e-f} \\
&\quad + (k_0 + k_1f + k_2f^2 + k_3f^3 + k_4f^4) \frac{(f-a)(f-b)(f-c)(f-d)}{f-e};
\end{aligned}$$

k_0, k_1, k_2, k_3, k_4 being quite arbitrary, and L, M, N , although presented in a fractional form, being essentially integral.

This fact of L, M, N constituting a system of multipliers to the syzygy QR, RP, PQ , is easily demonstrated; for

$$QR = (c-e)(c-f)(d-e)(d-f),$$

$$RP = (e-a)(e-b)(f-a)(f-b),$$

$$PQ = (a-c)(a-d)(b-c)(b-d).$$

p. 133

Hence

$$\begin{aligned}
&L(QR) + M(RP) + N(PQ) \\
&= (a-c)(a-d)(a-e)(a-f)(b-c)(b-d)(b-e)(b-f)(c-e)(c-f)(d-e)(d-f) \\
&\quad \times \sum \frac{k_0 + k_1a + k_2a^2 + k_3a^3 + k_4a^4}{(a-b)(a-c)(a-d)(a-e)(a-f)} = 0.
\end{aligned}$$

My theory of elimination enables me to explain exactly the nature of L, M, N , and the *reason* of their appearance as syzygetic factors.

Let L_r, M_r, N_r signify what L, M, N become, when all the k 's except k_r are taken zero. Then the theory given by me in the *Philosophical Magazine* for the year 1838, or thereabouts⁶³, shows that $L_0\lambda + L_1$ is the *prime derivate* of the first

⁶³This cannot be obtained directly from what is stated in the paper referred to, although contained in the general theory of derivation there given. The arbitrary functions which enter into the expression for the general derivees have been in that paper evaluated only for the prime derivees, which however are only particular phenomena, with reference to the general results of Dialytic Elimination. Hereafter I may give a more general exposition of this remarkable, although ignored or neglected theory. The prime derivees of fx and $f'x$ are Sturm's Functions, cleared of quadratic factors, and are expressed by virtue of the general theorems there laid down as functions of x and of symmetrical functions of the roots of fx . [p. 40 above. Ed.]

degree between the two equations P and $Q \times R$, or, in other words, will be the remainder integralized of

$$\frac{QR}{P}.$$

In like manner $M_0\lambda + M_1$, $N_0\lambda + N_1$ are the integralized remainders of

$$\frac{RP}{Q} \quad \text{and of} \quad \frac{PQ}{R}$$

respectively.

If now the resultant of P, Q and of Q, R are each zero, but the resultant of P and R is not zero, it will be evident that P, Q, R must be of the form

$$f(\lambda + a)(\lambda + c), \quad g(\lambda + c)(\lambda + d), \quad h(\lambda + d)(\lambda + b);$$

and therefore $P \times R$ will contain Q , and consequently we must have

$$M_0 = 0, \quad M_1 = 0.$$

More generally, if we write

$$Q = 0, \quad \lambda Q = 0, \quad \lambda^2 Q = 0, \quad P \times R = 0,$$

and eliminate dialytically, that is, treating $\lambda^4, \lambda^3, \lambda^2, \lambda$ as distinct quantities, we shall obtain

$$\lambda^4 : \lambda^3 : \lambda^2 : \lambda : 1 :: M_4 : M_3 : M_2 : M_1 : M_0;$$

and therefore when $P \times R$ contains Q ,

$$M_0 = 0, \quad M_1 = 0, \quad M_2 = 0, \quad M_3 = 0, \quad M_4 = 0.$$

p. 134

In like manner, when $Q \times P$ contains R ,

$$N_0 = 0, \quad N_1 = 0, \quad N_2 = 0, \quad N_3 = 0, \quad N_4 = 0;$$

and when $R \times Q$ contains P ,

$$L_0 = 0, \quad L_1 = 0, \quad L_2 = 0, \quad L_3 = 0, \quad L_4 = 0.$$

Accordingly, we see from the equation

$$L(QR) + M(RP) + N(PQ) = 0,$$

that if $QR = 0$, $RP = 0$; but PQ not $= 0$, then $N = 0$; and therefore

$$N_0 = 0, \quad N_1 = 0, \quad N_2 = 0, \quad N_3 = 0, \quad N_4 = 0,$$

and so in like manner for the remaining corresponding two suppositions⁶⁴.

Before proceeding to consider the remaining case of the highest species of contact, I must observe that besides the equations involved in the condition that $A, B, C; F, G, H$, or, which is the same thing, that any three of them shall all have a factor in common, we must have $\square(U + \lambda V)$ containing the square of such common factor. In the memoir before adverted to a general theorem will be given and proved, which shows that this latter condition is involved in the former one; in fact, more generally (but still only as a particular case) that when U and V are quadratic functions of n letters, but $U + eV$ admits of being represented as a complete function of $(n - 2)$ quantities only, which are themselves linear functions of the n letters, then $\square(U + \lambda V)$, which is of course a function of λ of the n th degree, will contain the factor $(\lambda - e)^2$.

When the two conics have four consecutive points in common, the characters of double-point contact and of contact in three consecutive points must exist simultaneously; and consequently the factor common to $A, B, C; F, G, H$, will enter not as a binary but as a ternary factor into $\square(U + \lambda V)$. This gives the extra condition required. As an example take the two conics,

$$U = \frac{y^2}{1-k} + x^2 - z^2 = 0,$$

$$V = y^2 + x^2 - 2kxz + (2k-1)z^2 = 0,$$

$$U + \lambda V = \left(\frac{1}{1-k} + \lambda \right) y^2 + (1 + \lambda)x^2 - \{1 + \lambda(1-2k)\}z^2 - 2k\lambda xz.$$

p. 135

The complete determinant of $U + \lambda V$ is then

$$-\frac{1}{1-k} \{1 + (1-k)\lambda\} \{(1+\lambda)^2 - 2k\lambda(1+\lambda) + k^2\lambda^2\} = -\frac{1}{1-k} \{1 + (1-k)\lambda\}^3.$$

A, B, C are the determinants of $U + \lambda V$, when $x = 0, y = 0, z = 0$, respectively. Thus

$$A = \left(\frac{1}{1-k} + \lambda \right) (1 + \lambda),$$

$$B = \left(\frac{1}{1-k} + \lambda \right) \{1 + \lambda(1-2k)\},$$

⁶⁴Since we are able to assign the values of the syzygetic multipliers in the equations

$$\begin{aligned} L(PQ) + M(QR) + N(RP) &= 0, \\ L'(PQ) + M'(PQR) + N'(QR) &= 0, \\ L''(QR) + M''(QRP) + N''(RP) &= 0, \\ L'''(RP) + M'''(RPQ) + N'''(PQ) &= 0, \end{aligned}$$

it follows that we may eliminate between these four equations any three of the six quantities $(PQ), (PRQ), \&c.$, and thus express any one of them in terms of any two others: this method, however, is not practically convenient. I may probably hereafter return to this subject.

$$C = k^2\lambda^2 - (1 + \lambda)\{1 + \lambda(1 - 2k)\} = \lambda^2(1 - k)^2 - 2\lambda(1 - k) - 1;$$

$$\lambda = -\frac{1}{1 - k}$$

makes $A = 0$, $B = 0$, $C = 0$, and the factor $\lambda + \frac{1}{1-k}$ enters cubed into $\square(U + \lambda V)$.

Hence the two conics have a contact of the third order.

This is easily verified; for if we pass from general to Cartesian and rectangular coordinates, and make z unity; $U = 0$ will represent an ellipse with centre at the origin, eccentricity $\sqrt{k}$, and mean focal distance 1, and $V = 0$ the circle of curvature at the extremity of the axis major⁶⁵.

I had intended to have added some other remarks connected with the present discussion, and also to have appended an *à posteriori* proof of the propositions relative to the reality and otherwise of the vertices and chordal pairs of intersection which I have, at the commencement of this paper, deduced quite legitimately, but in a manner not at first sight perhaps easily intelligible, from the general principles of conjugate forms; but this discussion has run on already to a length so much greater than I had anticipated and than the importance of the inquiry may seem to justify, that I must reserve for a future number of the *Journal* what further matter I may have to communicate concerning it.

POSTSCRIPT

As I have alluded to Professor Boole's theorem relative to Linear Transformations, it may be proper to mention my theorem on the subject, which is of a much more general character, and includes Mr Boole's (so far as it refers to Quadratic Functions) as a corollary to a particular case. The demonstration will be given in the forthcoming memoir above alluded to.

Let U be a quadratic function of any number of letters $x_1, x_2 \dots x_n$, and let any number r of linear equations of the general form

$$^1a_rx_1 + ^2a_rx_2 + \dots + ^na_rx_n = 0$$

p. 136

be instituted between them: and by means of these equations let U be expressed as a function of any $(n - r)$ of the given letters, say of $x_{r+1}, x_{r+2} \dots x_n$, and let U , so expressed, be called M . Let

$$^1a_rx_1 + ^2a_rx_2 + \dots + ^na_rx_n$$

be called L_r . Then the determinant of M in respect to the $(n - r)$ letters above given is equal to the determinant of

$$U + L_1x_{n+1} + L_2x_{n+2} + \dots + L_rx_{n+r},$$

⁶⁵We have thus discussed all the four cases of biconical contact: for an exactly parallel discussion of the theory of contact of a plane with the curve of double curvature in which two surfaces of the second order intersect, see the paper in the *Philosophical Magazine* for this month, before referred to. [p. 148 below. Ed.]

considered as a function of the $(n + r)$ letters

$$x_1 x_2 \dots x_{n+r},$$

divided by the square of the determinant

$$\begin{vmatrix} {}^1a_1 & {}^2a_1 & \dots & {}^ra_1 \\ {}^1a_2 & {}^2a_2 & \dots & {}^ra_2 \\ \dots & \dots & \dots & \dots \\ {}^1a_r & {}^2a_r & \dots & {}^ra_r \end{vmatrix}.$$

This I call the theorem of Diminished Determinants.

If now we have U a function of r letters, and V of r other letters, and V is derived from U by linear transformations, that is, by r equations connecting the $2r$ letters; then, since U may be considered as a function of *all the* $2r$ letters with abortive coefficients for all the terms where any of the second set of r letters enter, we may apply our theorem of diminished determinants to the question so considered, and the result may be found to represent Mr Boole's theorem in a form rather more general and symmetrical, but substantially identical with that given by Mr Boole.

Thus suppose $\frac{1}{2}ax^2 + bxy + \frac{1}{2}cy^2$ say P , and $\frac{1}{2}\alpha u^2 + \beta uv + \frac{1}{2}\gamma v^2$ say Q , are mutually transformable by virtue of the linear equations

$$lx + my = \lambda u + \mu v, \quad l'x + m'y = \lambda' u + \mu' v,$$

P may be considered as a function of x, y, u, v , and Q as the value of P , when we eliminate x and y by virtue of the two linear equations

$$L_1 = lx + my - \lambda u - \mu v = 0, \quad L_2 = l'x + m'y - \lambda' u - \mu' v = 0;$$

we have therefore by our theorem the determinant of Q equal to the squared reciprocal of the determinant

$$\begin{vmatrix} l & m \\ l' & m' \end{vmatrix}$$

multiplied by the determinant

$$\begin{vmatrix} a & b & 0 & 0 & l & l' \\ b & c & 0 & 0 & m & m' \\ 0 & 0 & 0 & 0 & -\lambda & -\lambda' \\ 0 & 0 & 0 & 0 & -\mu & -\mu' \\ l & m & -\lambda & -\mu & 0 & 0 \\ l' & m' & -\lambda' & -\mu' & 0 & 0 \end{vmatrix}.$$

which last determinant is evidently equal to the determinant of P multiplied by the square of the determinant

$$\begin{vmatrix} \lambda & \mu \\ \lambda' & \mu' \end{vmatrix}.$$

Whence we see that the determinant of Q divided by the square of

$$\begin{vmatrix} \lambda & \mu \\ \lambda' & \mu' \end{vmatrix}$$

is equal to the determinant of P divided by the square of

$$\begin{vmatrix} l & m \\ l' & m' \end{vmatrix}.$$

There is also another way more simple, but less direct, by means of which the theorem of diminished determinants may be made to yield Mr Boole's theorem of transformation⁶⁶. Some unavowed use has been made in the foregoing pages of this former theorem, one of the highest importance in the analytical and geometrical theory of quadratic functions. It has been nearly a year in my possession, and I trust and believe that I am committing no act of involuntary misappropriation in announcing it as a result of my own researches.

⁶⁶Namely, by considering P and Q as each derived from some common function of x, y, u, v, w , by means of the equations $L_1 = 0, L_2 = 0$; the law of Diminished Determinants will then indicate the determinants of P and Q , each under the form of fractions having the same numerator, but whose denominators will be $\begin{vmatrix} \lambda & \mu \\ \lambda' & \mu' \end{vmatrix}$ and $\begin{vmatrix} l & m \\ l' & m' \end{vmatrix}$ respectively.

23.

AN INSTANTANEOUS DEMONSTRATION OF PASCAL'S THEOREM BY THE METHOD OF INDETERMINATE COORDINATES

[*Philosophical Magazine*, XXXVII. (1850), p. 212]

p. 138

The new analytical geometry consists essentially of two parts—the one determinate, the other indeterminate.

The determinate analysis comprehends that class of questions in which it is necessary to assume *independent* linear coordinates, or else to take cognizance of the equations by which they are connected if they are not independent. The indeterminate analysis assumes at will any number of coordinates, and leaves the relations which connect them more or less indefinite, and reasons chiefly through the medium of the general properties of algebraic forms, and their correspondencies with the objects of geometrical speculation. Pascal's theorem of the mystic hexagon, and the annexed demonstration of its fundamental property, belong to this branch of the subject, and afford an instructive and striking example of the application of the pure method of indeterminate coordinates.

Let x, y, z, t, u, v be the sides of a hexagon inscribed in the conic U . Let the hexagon be divided by a new line ϕ in any manner into two quadrilaterals, say $xyz\phi, tuv\phi$. Then

$$ay\phi + bvx = U = au\phi + \beta tv;$$

therefore

$$(ay - \alpha u)\phi = \beta tv - bvx;$$

therefore $ay - \alpha u$ and ϕ are the diagonals of the quadrilateral $txvz$.

By construction, ϕ is the diagonal joining x, v (that is, the intersection of x and v) with z, t ; and thus we see that $ay - \alpha u$ is the line joining t, x with v, z ; but this line passes through y, u . Therefore x, t ; y, u ; z, v lie in one and the same right line. Q.E.D.

24.

ON A NEW CLASS OF THEOREMS IN ELIMINATION BETWEEN QUADRATIC FUNCTIONS

[Philosophical Magazine, XXXVII. (1850), pp. 213–218]

p. 139

In a forthcoming memoir on determinants and quadratic functions, I have demonstrated the following remarkable theorem as a particular case of one much more general, also there given and demonstrated.

Let U and V be respectively quadratic functions of the same $2n$ letters, and let it be supposed possible to institute n such linear equations between these letters as shall make U and V both simultaneously become identically zero⁶⁷. Then the determinant of $\lambda U + \mu V$, which is of course a function of λ and of μ of the $2n$ th degree, will become a *perfect square* of a function of λ and μ of the n th degree; and conversely, if this determinant be a perfect square, U and V may be made to vanish simultaneously by the institution of n linear equations between the $2n$ letters⁶⁸.

Let now P and Q be respectively quadratic functions of three letters only, say x, y, z ; and let

$$U = P + (lx + my + nz)t,$$

$$V = Q + k(lx + my + nz)t.$$

The determinant of $\lambda U + \mu V$ in respect to x, y, z, t is easily seen to be $(\lambda + k\mu)^2$ times the determinant of

$$\lambda P + \mu Q + (lx + my + nz)t$$

in respect to x, y, z, t . Hence if we call

$$\lambda P + \mu Q + (lx + my + nz)t = W,$$

and make $\square_{xyzt}W$ a squared function of λ, μ or which is the same thing, if

$$\square_{\lambda\mu}\square_{xyzt}\{W\} = 0,$$

p. 140

U and V will vanish simultaneously when two linear relations are instituted between the quantities (all or some of them) x, y, z, t .

In order that this may be the case, it will be seen to be sufficient that

$$P = 0, \quad Q = 0, \quad lx + my + nz = 0,$$

⁶⁷In the more general theorem above alluded to, the number of letters is any number m , the number of linear equations being any number not exceeding $\frac{m}{2}$.

⁶⁸When $n = 1$, we obtain a theorem of elimination between two quadratics, which has been already given by Professor Boole.

shall coexist; for then two equations between x, y, z of which $lx + my + nz = 0$ will be one, will suffice to make U and V each identically zero. Hence we have the following theorem:

$$\square_{\lambda\mu}\square_{xyzt}\{\lambda U + \mu V + (lx + my + nz)t\}$$

is a factor of the resultant of

$$P = 0, \quad Q = 0, \quad lx + my + nz = 0.$$

A comparison of the orders of the resultant and the determinant shows that they must be identical, à-ci-près, of a numerical factor, which, if the resultant be taken in its *general* lowest terms, may no doubt be easily shown to be unity.

As an illustration of our theorem, let

$$P = xy + yz + zx, \quad Q = cxy + ayz + bzx.$$

Then

$$\begin{aligned} \square_{xyzt}\{\lambda P + \mu Q + (lx + my + nz)t\} &= \begin{vmatrix} 0 & \lambda + c\mu & \lambda + b\mu & l \\ \lambda + c\mu & 0 & \lambda + a\mu & m \\ \lambda + b\mu & \lambda + a\mu & 0 & n \\ l & m & n & 0 \end{vmatrix} \\ &= n^2(\lambda + c\mu)^2 + m^2(\lambda + b\mu)^2 + l^2(\lambda + a\mu)^2 \\ &\quad - 2lm(\lambda + b\mu)(\lambda + a\mu) - 2mn(\lambda + c\mu)(\lambda + b\mu) \\ &\quad - 2nl(\lambda + a\mu)(\lambda + c\mu) \\ &= \lambda^2\{n^2 + m^2 + l^2 - 2lm - 2mn - 2nl\} \\ &\quad + 2\lambda\mu\{cn^2 + bm^2 + al^2 - lm(a + b) - mn(b + c) - nl(c + a)\} \\ &\quad + \mu^2\{c^2n^2 + b^2m^2 + a^2l^2 - 2ablm - 2bcmn - 2canl\}. \end{aligned}$$

And we thus obtain, finally,

$$\begin{aligned} &\square_{\lambda\mu}\square_{xyzt}\{\lambda P + \mu Q + (lx + my + nz)t\} \\ &= (n^2 + m^2 + l^2 - 2lm - 2mn - 2nl) \\ &\quad \times (c^2n^2 + b^2m^2 + a^2l^2 - 2ablm - 2bcmn - 2canl) \\ &\quad - \{cn^2 + bm^2 + al^2 - lm(a + b) - mn(b + c) - nl(c + a)\}^2 \\ &= -4lmn\{(a - b)(a - c)l + (b - a)(b - c)m + (c - a)(c - b)n\}. \end{aligned}$$

p. 141

Now to obtain the resultant of

$$xy + yz + zx = 0, \quad cxy + azy + bxz = 0, \quad lx + my + nz = 0,$$

we need only find the four systems *in their lowest terms* of $x : y : z$, which satisfy the first two equations, and multiply the four linear functions obtained

by substituting these values of x, y, z in the fourth: the product will contain the resultant of the system affected with some numerical factor. In the present case, the four systems of x, y, z are

$$x = 0, \quad y = 0, \quad z = 1,$$

$$y = 0, \quad z = 0, \quad x = 1,$$

$$z = 0, \quad x = 0, \quad y = 1,$$

$$x = (a - b)(a - c), \quad y = (b - a)(b - c), \quad z = (c - a)(c - b),$$

and accordingly the product of

$$lx_1 + my_1 + nz_1,$$

$$lx_2 + my_2 + nz_2,$$

$$lx_3 + my_3 + nz_3,$$

$$lx_4 + my_4 + nz_4,$$

becomes

$$lmn\{(a - b)(a - c)l + (b - a)(b - c)m + (c - a)(c - b)n\},$$

agreeing with the result obtained by my theorem,—a special numerical factor 4, arising from the peculiar form of the equations, having disappeared from the resultant.

A geometrical demonstration may be given of the theorem which is instructive in itself, and will suggest a remarkable extension of it to functions containing more than three letters; the equation

$$\square_{xyzt}\{\lambda U + \mu V + (lx + my + nz)t\} = 0,$$

which is a quadratic equation in $\lambda : \mu$, may easily be shown to imply that the conic $\lambda U + \mu V$ is touched by the straight line

$$lx + my + nz = 0.$$

And we thus see that in general two conics,

$$\lambda U + \mu V = 0,$$

passing through the intersections of two given conics,

$$U = 0, \quad V = 0,$$

may be drawn to touch a given line. If, however, the given line passes through any of the four points of intersection, in such case only one conic can be drawn to touch it; accordingly

$$\square\square\{\lambda U + \mu V + (lx + my + nz)t\}$$

must be zero when l, m, n are so taken as to satisfy this condition, that is, if

$$lx_1 + my_1 + nz_1 = 0,$$

or

$$lx_2 + my_2 + nz_2 = 0,$$

or

$$lx_3 + my_3 + nz_3 = 0,$$

or

$$lx_4 + my_4 + nz_4 = 0,$$

whence the theorem.

Now suppose U and V to be each functions of four letters, x, y, z, t ; when

$$\square_{xyztu}\{\lambda U + \mu V + (lx + my + nz + pt)u\} = 0,$$

the conoid $\lambda U + \mu V$ touches the plane

$$lx + my + nz + pt = 0;$$

and $\square = 0$ being a cubic equation, in general three such conoids can be drawn.

Considerations of analogy make it obvious to the intuition, that in the particular case of two of these becoming coincident, the given plane

$$lx + my + nz + pt$$

must be a tangent plane to those two coincident conoids at one of the points where it meets the intersections of $U = 0, V = 0$; that is

$$lx + my + nz + pt = 0$$

will pass through a tangent line to, or in other words, may be termed a tangent plane to the intersections. Hence the following analytical theorem, derived from supposing q, r, s, t to be proportional to the areas of the triangular faces of the pyramid cut out of space by the four coordinate planes to which x, y, z, t refer. As these planes are left indefinite, q, r, s, t are perfectly arbitrary.

Theorem. The resultant of

1. $U = 0$
 2. $V = 0$
- where U and V are functions of x, y, z, t ;

and

$$3. \quad lx + my + nz + pt = 0;$$

together with

$$4. \quad \begin{vmatrix} \frac{dU}{dx} & \frac{dU}{dy} & \frac{dU}{dz} & \frac{dU}{dt} \\ \frac{dV}{dx} & \frac{dV}{dy} & \frac{dV}{dz} & \frac{dV}{dt} \\ l & m & n & p \\ q & r & s & t \end{vmatrix} = 0;$$

which system, it will be observed, consists of three quadratic functions, and one linear function of x, y, z, t , contains the factor

p. 143

$$\square_{\lambda\mu}\square_{xyzt}\{\lambda U + \mu V + (lx + my + nz + pt)u\}.$$

This last quantity is of the 4×3 th, that is, the 12th order in respect of the coefficients in U and V combined; of the 4×2 th, that is, the 8th order in respect of l, m, n, p ; and of the zero order in respect of q, r, s, t .

The resultant which contains it is of the $(4 + 4 + 2 \cdot 4)$ th, that is, 16th order in respect to the coefficients in U and V ; of the $(4 + 8)$ th, that is, the 12th, in respect of l, m, n, p ; and of the 4th in respect of q, r, s, t . Hence the special (and, as far as the geometry of the question is concerned, the unnecessary, I may not say extraneous or irrelevant) factor which enters into the resultant is of the 4th order in respect to the combined coefficients of U and V ⁶⁹; and of the same order in respect to l, m, n, p , and in respect to q, r, s, t .

I have not yet succeeded in divining its general value.

In the very particular example, of the system,

$$\alpha x^2 + \beta y^2 = 0,$$

$$cz^2 + dt^2 = 0,$$

$$lx + my + nz + pt = 0,$$

$$\begin{vmatrix} \alpha x & \beta y & 0 & 0 \\ 0 & 0 & cz & dt \\ l & m & n & p \\ q & 0 & 0 & 0 \end{vmatrix} = 0,$$

I find that the double determinant is

$$c^2 d^2 \alpha^2 \beta^2 (cp^2 + dn^2)^2 (m^2 \alpha + l^2 \beta)^2,$$

and the resultant is

$$q^4 c^2 d^2 \alpha^2 \beta^4 (cp^2 + dn^2)^4 (m^2 \alpha + l^2 \beta)^2,$$

⁶⁹And consequently of the second in respect to the separate coefficients of each.

giving as the special factor

$$q^4\beta^2(cp^2 + dn^2)^2.$$

I believe that the theorem which I have here given for determining the condition that $lx + my + nz + pt$ shall be a tangent plane to the intersection of two conoids U and V , namely, that the determinant of

$$\lambda U + \mu V + (lx + my + nz + pt)u$$

shall have two equal roots, is altogether novel.

p. 144

What is the meaning of all three roots of this determinant becoming equal, that is, of only one conoid being capable of being drawn through the intersection of U and V to touch the plane

$$lx + my + nz + pt?$$

Evidently (*ex vi analogiae*) that this plane shall pass through three consecutive points of the curve of intersection, that is, that it shall be the osculating plane to the curve.

If we return to the intersection of two co-planar conics, and if we suppose a line to be drawn through two of the points of intersection, the conics capable of being drawn through the four points of intersection to touch the line, besides becoming coincident, evidently degenerate each into a pair of right lines. It would seem, therefore, by analogy, that if a plane be drawn including any two tangent lines to the curve of intersection of two surfaces of the second degree, this should be touched by two coincident cones drawn through the curve of intersection, and consequently every such double tangent plane to the intersection of two conoids (and it is evident that one or more of these can be taken at every point of the curve) must pass through one of the vertices of the four cones in which the intersection may also be considered to lie; and it would appear from this, that in general four double tangent planes admit of being drawn to the curve, which is the intersection of two conoids, at each point thereof. At particular points a tangent plane may be drawn passing through more than one of the vertices, and then of course the number of double tangent planes that can be drawn will be lessened. These results, indicated by analogy, become immediately apparent on considering the curve in question as traced upon any one of the four containing cones. For the plane drawn through a tangent at any point, and the vertex of the cone being a tangent plane to the cone, must evidently touch the curve again where it meets it. We thus have an additional confirmation of the analogy between a point of intersection of two curves and the tangent at any point of the intersection of two surfaces.

I might extend the analytical theorems which have been established for functions of three and four to functions of a greater number of variables; but

enough has been done to point out the path to a new and interesting class of theorems at once in elimination and in geometry, which is all that I have at present leisure or the disposition to undertake.

25.

ADDITIONS TO THE ARTICLES "ON A NEW CLASS OF THEOREMS," AND "ON PASCAL'S THEOREM"

[Philosophical Magazine, XXXVII. (1850), pp. 363-370]

p. 145

First addition.—I have alluded in the second of the above articles to a more general theorem, comprising, as a particular case, the theorem there given for the simultaneous evanescence of two quadratic functions of $2n$ letters, on n linear equations becoming instituted between the letters.

In order to make this generalization intelligible, I must premise a few words on the Theory of Orders, a term which I have invented with particular reference to quadratic functions, although obviously admitting of a more extended application. A linear function of all the letters entering into a function or system of functions under consideration I call an order of the letters, or simply an order. Now it is clear that we may always consider a function of any number of letters as a function of as many orders as there are letters; but in certain cases a function may be expressed in terms of a fewer number of orders than it has letters, as when the general characteristic function of a conic becomes that of a pair of crossing lines or a pair of coincident lines, in which event it loses respectively one and two orders, and so for the characteristic of a conoid becoming that of a cone, a pair of planes or two coincident planes, in which several events, a function of four letters becomes that of only three orders, or two orders, or one order, respectively. When a function may be expressed by means of r orders less than it contains letters, I call it a function minus r orders. I now proceed to state my theorem.

Let U and V be functions each of the same m letters, and suppose that the determinant in respect of those letters of $U + \mu V$ contains i pairs of p. 146

equal linear factors of μ ; then it is possible, by means of i linear equations instituted between the letters, to make U and V each become functions of *the same* $m - 2i$ orders; and conversely, if by i equations between the letters U and V may be made functions of the *same* $m - 2i$ orders, the determinant of $U + \mu V$ considered as a function of μ will contain i square factors.

Thus when $m = 2n$ and $i = n$, U and V will each become functions of zero orders, that is, will both disappear, provided that on the institution of a certain system of n linear equations, among the letters of which U and V are functions, the determinant of $(U + \mu V)$ is a perfect square,—which is the theorem given in the article referred to.

So for example if U and V be quadratic functions of four letters, and therefore the characteristics of two conoids, $\square(U + \mu V)$ being a perfect square, expresses that these conoids have a straight line in common lying upon each of their surfaces.

If U and V be quadratic functions of three letters only, and admit therefore of being considered as the characteristics of two conics, $\square(U + \mu V)$ containing a square factor, is indicative of these conics having a common tangent at a common point, that is, of their touching each other at some point; for it is easily shown that the disappearance of two orders from any quadratic function by virtue of one linear function of its letters being zero, indicates that the line, plane, &c. of which the linear function is the characteristic is a tangent to the curve, surface, &c. of which the quadratic function is the characteristic.

I pass now to a generalization of the theorem which shows how to express, under the form of a double determinant, the resultant of one linear and two quadratic homogeneous functions of three letters (which I should have given in the original paper, had I not there been more intent upon developing an ascending scale than of expatiating upon a superficial ramification of analogies), and which constitutes my *Second addition* to that paper, to wit—

If U and V be homogeneous quadratic, and $L_1, L_2 \dots L_n$ homogeneous linear functions of $(n + 2)$ letters $x_1, x_2 \dots x_{n+2}$, the determinant of the entire system of $n + 2$ functions is equal to

$$\square_{\lambda, \mu} \square_{x_1, x_2, \dots, x_{n+2}, t_1, t_2, \dots, t_n} \{ \lambda U + \mu V + L_1 t_1 + L_2 t_2 + \dots + L_n t_n \};$$

the demonstration is precisely similar to the analytical one given in the September Number⁷⁰ for the particular case of $n = 1$.

When $n = 0$, we revert to Mr Boole's theorem of elimination between U and V already adverted to. The proof, it will be easily recognized, does not require the application of the more general theorem relative to the simultaneous

p. 147

depression of orders of two quadratic functions, but only the limited one before given, which supplies the conditions of their simultaneous disparition. I now proceed to develop more particularly certain analogies between the theory of the mutual contacts of two conics, and that of the tangencies to the intersection of two conoids.

But here again I must anticipate some of the results which will be given in my forthcoming memoir on Determinants and Quadratic Functions, by explaining what is to be understood by minor determinants, and the relation in which they stand to the complete determinant in which they are included. This preliminary explanation, and the statement of the analogies above alluded to, will constitute my *Third* and last addition.

Imagine any determinant set out under the form of a square array of terms. This square may be considered as divisible into lines and columns. Now conceive any one line and any one column to be struck out, we get in this way a square, one term less in breadth and depth than the original square; and by varying in every possible manner the selection of the line and column excluded, we obtain,

⁷⁰p. 140 above.

supposing the original square to consist of n lines and n columns, n^2 such minor squares, each of which will represent what I term a First Minor Determinant relative to the principal or complete determinant. Now suppose two lines and two columns struck out from the original square, we shall obtain a system of

$$\left\{ \frac{n(n-1)}{2} \right\}^2$$

squares, each two terms lower than the principal square, and representing a determinant of one lower order than those above referred to. These constitute what I term a system of Second Minor Determinants; and so in general we can form a system of r th minor determinants by the exclusion of r lines and r columns, and such system *in general* will contain

$$\left\{ \frac{n(n-1) \cdots (n-r+1)}{1 \cdot 2 \cdots r} \right\}^2$$

distinct determinants.

I say “*in general*”; because if the principal determinant be totally or partially symmetrical in respect to either or each of its diagonals, the number of distinct determinants appertaining to each system of minors will undergo a material diminution, which is easily calculable.

Now I have established the following law:—

The whole of a system of r th minors being zero, implies only $(r+1)^2$ equations, that is, by making $(r+1)^2$ of these minors zero, all will become zero; and this is true, no matter what may be the dimensions or form of the complete determinant. But furthermore, if the complete determinant be formed from a quadratic function, so as to be symmetrical about one of its diagonals, then $\frac{1}{2}(r+1)(r+2)$ only of the r th minors being zero, will serve

p. 148

to imply that all these minors are zero. Of course, in applying these theorems, care must be taken that the $(r+1)^2$ or $\frac{1}{2}(r+1)(r+2)$ selected equations must be mutually non-implicative, and shall constitute independent conditions.

In the application I am about to make of these principles, we shall have only to deal with a system of *first* minors and of a *symmetrical* determinant. If three of these properly selected be zero, from the foregoing it appears that all must be zero.

Now let U and V be characteristics of two conics, that is, let each be a function of only three letters, it may be shown (see my paper⁷¹ in the *Cambridge and Dublin Mathematical Journal* for November, 1850) that the different species of contacts between these two conics will correspond to peculiar properties of the compound characteristic $U + \mu V$.

If the determinant of this function have two equal roots, the conics simply touch; if it have three equal roots, the conics have a single contact of a higher order,

⁷¹p. 119 above.

that is, the same curvature; if its six first minors become zero simultaneously for the same value of μ , the conics have a double contact. If the same value of μ , which makes all these first minors zero, be at the same time not merely a double root (as of analytical necessity it always must be) but a treble root of

$$\square(U + \mu V) = 0,$$

then the conics have a single contact of the highest possible order short of absolute coincidence, that is, they meet in four consecutive points.

The parallelism between this theory and that of two quadratic functions P, Q , and one linear function L ⁷² of four letters, say x, y, z, t , is exact⁷³. For let $P + Lu + \mu Q$ be now taken as our compound characteristic (a function, it will be observed, of five letters, x, y, z, t, u); if its determinant have two equal roots, L has two consecutive points in common with the intersection of P and Q , that is, passes through a tangent to that intersection; if it have three equal roots, L has three consecutive points in common with the said intersection, that is, is an osculating plane thereto; if its fifteen first minors admit of all being made simultaneously zero, L has a double contact with the intersection of P and Q , that is, it is a tangent plane to some one of the four cones of the second order containing this intersection;

p. 149

if the same linear function of μ which enters into all these first minors be contained cubically in the complete determinant, then the plane L passes through four consecutive points of the intersection of P and Q , and the points where it meets the curve will be points of contrary plane flexure; and, as it seems to me, at such points the tangential direction of the curve must point to the summit of one or other of the four cones above alluded to⁷⁴. In assigning the conditions for L being a double tangent plane to the intersection of P and Q , we may take any three independent minors at pleasure equal to zero. One of these may be selected so as to be clear of the coefficients of L ; in fact, the determinant of $P + \mu Q$ will be a first minor of $P + \mu Q + Lu$; μ may thus be determined by a biquadratic equation; and then, by properly selecting the two other minors, we may obtain two equations in which only the first powers of the coefficients of x, y, z, t in L appear, and may consequently obtain L under the form of

$$(ae + \alpha)x + (be + \beta)y + (ce + \gamma)z + (de + \delta)t,$$

⁷²Observe that $P = 0, Q = 0, L = 0$ now express the equations to two conoids and a plane respectively.

⁷³This parallelism may be easily shown analytically to imply, and be implied, in the geometrical fact, that the contact of the plane L with the intersection of the two surfaces P and Q , is of exactly the same kind as the contact (which must exist) between the two conics which are the intersections of P and Q respectively with the plane L .

⁷⁴If this be so, then we have the following geometrical theorem:—"The summit of one of the four cones of the second degree which contain the intersections of two surfaces of the second order drawn in any manner respectively through two given conics lying in the same plane, and having with one another a contact of the third degree, will always be found in the same right line, namely, in the tangent line to the two given conics at the point of contact."

where a, α ; b, β ; c, γ ; d, δ will be known functions of any one of the four values of μ . The point of contact being given will then serve to determine e , and we shall thus have the equation to each of the four double tangent planes at any given point fully determined.

In the foregoing discussions I have freely employed the word *characteristic* without previously defining its meaning, trusting to that being apparent from the mode of its use. It is a term of exceeding value for its significance and brevity. The characteristic of a geometrical figure⁷⁵ is the function which, equated to zero, constitutes the equation to such figure. Pluecker, I think, somewhere calls it the line or surface function, as the case may be. Geometry, analytically considered, resolves itself into a system of rules for the construction and interpretation of characteristics. One more remark, and I have done. A very comprehensive theorem has been given at the commencement of this commentary, for interpreting the effect of a complete determinant of a linear function of two quadratic functions ($U + \mu V$), having

p. 150

one or more pairs of equal factors ($e + \epsilon\mu$). But here a far wider theory presents itself, of which the aim should be to determine the effect and meaning of this determinant, having any amount and distribution of multiplicity whatsoever among its roots. Nor must our investigations end at that point; but we must be able to determine the meaning and effect of common factors, one or more entering into the successive systems of *minor* determinants derived from the complete determinant of $U + \mu V$.

Nor are we necessarily confined to two, but may take several quadratic functions simultaneously into account.

Aspiring to these wide generalizations, the analysis of quadratic functions soars to a pitch from whence it may look proudly down on the feeble and vain attempts of geometry proper to rise to its level or to emulate it in its flights.

The law which I have stated for assigning the number of independent, or to speak more accurately, non-coevanescent determinants belonging to a given system of minors, I call the Homaloidal law, because it is a corollary to a proposition which represents analytically the indefinite extension of a property common to lines and surfaces to all loci (whether in ordinary or transcendental space) of the first order, all of which loci may, by an abstraction derived from the idea of levelness common to straight lines and planes, be called Homaloids. The property in question is, that neither two straight lines nor two planes can have

⁷⁵More generally, the characteristic of any fact or existence is the function which, equated to zero, expresses the condition of the actuality of such fact or existence. Perhaps the most important pervading principle of modern analysis, but which has never hitherto been articulately expressed, is *that*, according to which we infer, that when one fact of whatever kind is implied in another, the characteristic of the first must contain as a factor the characteristic of the second; and that when two facts are mutually involved, their characteristics will be powers of the same integral function. The doctrine of characteristics, applied to dependent systems of facts, admits of a wide development, logical and analytical.

a common segment; in other words, if n independent relations of rectilinearity or of coplanarity, as the case may be, exist between triadic groups of a series of $n + 2$, or between tetradic groups of a series of $n + 3$ points respectively, then every triad or tetrad of the series, according to the respective suppositions made, will be in rectilinear or in plane order. So, too, if n independent relations of *coincidence* exist between the duads formed out of $n + 1$ points, every duad will constitute a coincidence.

This homaloidal law has not been stated in the above commentary in its form of greatest generality. For this purpose we must commence, not with a square, but with an oblong arrangement of terms consisting, suppose, of m lines and n columns. This will not in itself represent a determinant, but is, as it were, a Matrix out of which we may form various systems of determinants by fixing upon a number p , and selecting at will p lines and p columns, the squares corresponding to which may be termed determinants of the p th order. We have, then, the following proposition. The number of uncoevanescent determinants constituting a system of the p th order derived from a given matrix, n terms broad and m terms deep, may equal, but can never exceed the number

$$(n - p + 1)(m - p + 1).$$

REMARK ON PASCAL'S AND BRIANCHON'S THEOREMS

p. 151

I omitted to state, in the September Number of the *Journal*⁷⁶, that the demonstration there given by me for Pascal's, applied equally to Brianchon's theorem. This remark is of the more importance, because the fault of the analytical demonstrations hitherto given of these theorems has been, that they make Brianchon's consequence of Pascal's, instead of causing the two to flow simultaneously from the application of the same principles. No demonstration can be held valid in *method*, or as touching the essence of the subject-matter, in which the indifference of the duadic law is departed from. Until these recent times, the analytic method of geometry, as given by Descartes, had been suffered to go on halting as it were on one foot. To Pluecker was reserved the honour of setting it firmly on its two equal supports by supplying the complementary system of coordinates. This invention, however, had become inevitable, after the profound views promulgated by Steiner, in the introduction to his Geometry, had once taken hold of the minds of mathematicians. To make the demonstration in the article referred to apply, *totidem literis*, to Brianchon's theorem (recourse being had to the correlative system of coordinates), it is only needful to consider U as the characteristic of the tangential envelope of the conic, x, y, z, t, u, v as the characteristics of the six points of the circumscribed hexagon, ϕ the characteristic of the point in which the line x, v meets the line z, t ; $ay - au$ will then be shown to characterize the point in which t, x meets v, z ; and thus we see that $y, u; t, x; v, z$,

⁷⁶p. 138 above.

the three pairs of opposite sides of the hexagon, will meet in one and the same point, which is Brianchon's theorem.

26.

ON THE SOLUTION OF A SYSTEM OF EQUATIONS IN WHICH THREE
HOMOGENEOUS QUADRATIC FUNCTIONS OF THREE UNKNOWN
QUANTITIES ARE RESPECTIVELY EQUATED TO NUMERICAL MULTIPLES
OF A FOURTH NON-HOMOGENEOUS FUNCTION OF THE SAME

[Philosophical Magazine, xxxvii. (1850), pp. 370–373]

p. 152

Let U, V, W be three homogeneous quadratic functions of x, y, z , and let ω be any function of x, y, z of the n th degree, and suppose that there is given for solution the system of equations

$$U = A\omega, \quad V = B\omega, \quad W = C\omega.$$

Theorem. The above system can be solved by the solution of a cubic equation, and an equation of the n th degree.

For let D be the determinant in respect to x, y, z of

$$fU + gV + hW,$$

then D is a cubic function of f, g, h . Now make

$$D = 0, \quad Af + Bg + Ch = 0;$$

the ratios of $f : g : h$ which satisfy the last two equations can be determined by the solution of a cubic equation, and there will accordingly be three systems of f, g, h which satisfy the same, as

$$\begin{array}{ccc} f_1, & g_1, & h_1, \\ f_2, & g_2, & h_2, \\ f_3, & g_3, & h_3. \end{array}$$

Now $D = 0$ implies that $fU + gV + hW$ breaks up into two linear factors; accordingly we shall find

$$\begin{array}{rcl} (l_1x + m_1y + n_1z)(\lambda_1x + \mu_1y + \nu_1z) & = & 0, \\ (l_2x + m_2y + n_2z)(\lambda_2x + \mu_2y + \nu_2z) & = & 0, \\ (l_3x + m_3y + n_3z)(\lambda_3x + \mu_3y + \nu_3z) & = & 0, \end{array}$$

in which the several sets of $l, m, n; \lambda, \mu, \nu$ can be expressed without difficulty in terms of the several values of $\sqrt{f}, \sqrt{g}, \sqrt{h}$. p. 153

Let the above equations be written under the form

$$PP' = 0, \quad QQ' = 0, \quad RR' = 0.$$

Since the given equations are perfectly general, it is readily seen that the equations

$$(P = 0, P' = 0), \quad (Q = 0, Q' = 0), \quad (R = 0, R' = 0),$$

will severally represent pairs of opposite sides of a quadrangle expressed by general coordinates x, y, z ; so that one of the two functions R, R' will be a linear function of P and Q and also of P' and Q' , and the other will be a linear function of P and Q' and also of P' and Q .

In order to solve the equations, we need only consider two such pairs: as $PP' = 0, QQ' = 0$; we then make

$$\begin{aligned} P &= 0, & Q &= 0, \\ P &= 0, & Q' &= 0, \\ P' &= 0, & Q &= 0, \\ P' &= 0, & Q' &= 0. \end{aligned}$$

Any one of these four systems will give the ratios of $x : y : z$; and then, by substitution in any one of the given equations, we obtain the values of x, y, z by the solution of an ordinary equation of the n th degree. The number of systems x, y, z is therefore always $4n$.

The equations connected with the solution of Malfatti's celebrated problem, "In a given triangle to inscribe three circles such that each circle touches the remaining two circles and also two sides of the triangle," given by Mr Cayley in the November Number for 1849 of the *Cambridge and Dublin Mathematical Journal*, to wit,

$$\begin{aligned} by^2 + cz^2 + 2fyz &= \theta^2 a(bc - f^2) = A, \\ cz^2 + ax^2 + 2gzx &= \theta^2 b(ca - g^2) = B, \\ ax^2 + by^2 + 2hxy &= \theta^2 c(ab - h^2) = C, \end{aligned}$$

come under the general form which has just been solved. It so happens, however, that in this particular case

$$\left. \begin{array}{ccc} f_1, & g_1, & h_1 \\ f_2, & g_2, & h_2 \\ f_3, & g_3, & h_3 \end{array} \right\}$$

77

p. 154

become respectively

$$\left. \begin{array}{ccc} 0, & \frac{1}{B}, & -\frac{1}{C} \\ -\frac{1}{B}, & 0, & \frac{1}{C} \\ -\frac{1}{C}, & \frac{1}{B}, & 0 \end{array} \right\},$$

⁷⁷Were it not for this being the case, the number of solutions would be n times the number of ways of obtaining duads out of three sets of two things, excluding the duads forming the sets, that is, the number of solutions would be $12n$ in place of $4n$, the true number.

and the cubic equation is resolved without extraction of roots.

It follows from my theorem that the eight intersections of three concentric surfaces of the second order can be found by the solution of one cubic and one quadratic equation; and in general, if we have ϕ, ψ, θ any three quadratic functions of x, y, z , and $\phi = 0, \psi = 0, \theta = 0$ be the system of equations to be solved, provided that we can by linear transformations express ϕ, ψ, θ under the form of

$$U - aw, \quad V - bw, \quad W - cw,$$

U, V, W being homogeneous functions, and w a non-homogeneous function of three new variables, x', y', z' , we can find the eight points of intersection of the three surfaces, of which U, V, W are the characteristics, by the solution of one cubic and one quadratic. But (as I am indebted to Mr Cayley for remarking to me) that this may be possible, implies the coincidence of the vertices of one cone of each of the systems of four cones in which the intersections of the three surfaces taken two and two are contained.

I may perhaps enter further hereafter into the discussion of this elegant little theory. At present I shall only remark, that a somewhat analogous mode of solution is applicable to two equations,

$$U = aP^2, \quad V = bP^2,$$

in which U, V are homogeneous quadratic functions, and P some non-homogeneous function of x, y .

We have only to make the determinant of $fU + gV$ equal to zero, and we shall obtain two systems of values of f, g , wherefrom we derive

$$l_1x + m_1y = \pm\sqrt{af_1 + bg_1} P, \quad l_2x + m_2y = \pm\sqrt{af_2 + bg_2} P,$$

from which x and y may be determined.

27.

ON A PORISMATIC PROPERTY OF TWO CONICS HAVING WITH ONE ANOTHER A CONTACT OF THE THIRD ORDER

[*Philosophical Magazine*, xxxvii. (1850), pp. 438, 439]

p. 155

If two conics have with one another a contact of the third order, that is, if they intersect in four consecutive points, it will easily be seen that their characteristics referred to coordinate axes in the plane containing them must be of the relative forms $x^2 + yz$, $k(y^2 + x^2 + yz)$ respectively, y characterizing their common tangent at the point of contact⁷⁸.

Hence if we take planes of reference in space, and call t the characteristic of the plane of the conics, the equations to any two conoids drawn through them respectively will be of the relative forms

$$U = x^2 + yz + tu = 0, \quad V = y^2 + x^2 + yz + tv = 0.$$

Using W to denote $V - U$, and (W) to denote what W becomes when ey is substituted for t , we see that W and (W) are of the respective forms $y^2 + tw$ and $y\theta$; showing that the former is the characteristic of a cone which will be cut by any plane $t - ey$ drawn through the line (t, y) in a pair of right lines; or, in other words, that one of the cones containing the intersection of the two variable conoids (V and U) will have its vertex in the invariable line which is the common tangent to the two fixed conics: this proves the theorem stated by me hypothetically in a foot-note in one of my papers in the last number of the *Magazine*⁷⁹. The steps of the geometrical proof there hinted at are as follows. p. 156

The four consecutive points in which the two conics intersect will be consecutive points in the curve of intersection of the two variable conoids. This curve lies in each of four cones of the second degree. Every double tangent plane to it passes through the vertex of one amongst these. The plane containing four, that is, two (consecutive) pairs of consecutive points, is a double tangent plane, and will therefore pass through a vertex; but four consecutive points of a curve of the fourth order described upon a cone, and lying in one tangent plane thereto, can only be conceived generally as disposed in the form of an $\int$, of which the belly part will point to the vertex; or, in other words, at any point where two consecutive osculating planes coincide so that the spherical curvature vanishes,

⁷⁸These relative or conjugate forms are taken from a table which I shall publish in a future number of this *Magazine*, exhibiting the conjugate characteristics in their simplest forms, correspondent to all the various species of contacts possible between lines and surfaces of the second degree. This table is as important to the geometer as the fundamental trigonometrical formula to the analyst, or the multiplication table to the arithmetician; and it is surprising that no one has hitherto thought of constructing such.

⁷⁹P. 149 above.

the linear curvature will also vanish, that is, there will be a point of inflexion at which, of course, the tangent line must pass through the vertex of the cone. This is the assumption felt to be true, but stated by me hypothetically in the paper referred to, because a ready demonstration did not at the moment occur to me. The legitimacy of this inference is now vindicated by the above analytical demonstration.

The methods of general and correlative coordinates and of determinants combined possess a perfectly irresistible force (to which I can only compare that of the steam-hammer in the physical world) for bringing under the grasp of intuitive perception the most complicated and refractory forms of geometrical truth.

ON THE ROTATION OF A RIGID BODY ABOUT A FIXED POINT

[*Philosophical Magazine*, xxxvii. (1850), pp. 440–444]

p. 157

In the *Cambridge and Dublin Mathematical Journal* for March 1848, an article by Professor Stokes, of the University of Cambridge, is ushered in with the words following:—

“The most general instantaneous⁸⁰ motion of a rigid body moveable in all directions about a fixed point consists in a motion of rotation about an axis passing through that point. This elementary proposition is sometimes assumed as self-evident, and sometimes deduced as the result of an analytical process. It ought hardly perhaps to be assumed, but it does not seem desirable to refer to a long algebraical process for the demonstration of a theorem so simple. Yet I am not aware of a geometrical proof anywhere published which might be referred to.”

The learned and ingenious professor is indubitably right, and might have trusted himself to assert less hesitatingly the necessity of demonstrating this proposition, which possesses none of the characters of a self-evident truth; but it is to be regretted that he should have stated it in such a form as naturally to lead the incautious reader to mistake the nature and grounds of its existence, which consist in this fact—that any kind of displacement of a body moveable about a fixed axis, whether instantaneous and infinitesimal, or secular and finite, is capable of being effected by a single rotation about a single axis.

The annexed simple proof of this capital law has the advantage of affording a rule for compounding into one any two (and therefore any number of) rotations given in direction, magnitude and order of succession.

p. 158


It will somewhat conduce to simplicity if we fix our attention upon a spherical surface rigidly connected with the rotating body, and having its centre at the fixed point thereof. When the positions of two points in this are given, the position of the body is completely determined.

Now evidently two points A, B may be brought respectively to $A'B'$ (if $AB = A'B'$) by two rotations; the first taking place about a pole situated anywhere in the great circle bisecting AA' at right angles, the second about A' , the position into which it is brought by the first rotation. This view leads us to consider the effect of two rotations taking place successively about two axes fixed in the rotating body. Or again, we may make the plane $A'B'$ revolve into the position AB round a pole taken at the node in which the two planes intersect,

⁸⁰The italics do not exist in the original.

and then the points A, B swing into their new positions A', B' by means of a rotation about the pole of the great circle, of which $A'B'$ forms a part. This mode of effecting the displacement naturally suggests the consideration of the effect of rotations taking place successively about two axes fixed in space.

First, then, let us study the effect of the combination of a rotation (α) having P for its pole, followed by another (β), of which Q is the pole, P and Q being points in the surface of the revolving sphere.



In drawing the annexed figure, I have supposed that the two rotations are of the same kind, each tending, when a spectator is standing with his head to the respective poles and his feet to the centre, to make a point at his right-hand pass in front of his face towards his left-hand. Let now PQ revolve through $\frac{\alpha}{2}$ positively into the position of PR , and through $\frac{\beta}{2}$ negatively into that of QR .

Then I say that the two impressed rotations α and β about P and Q will be equivalent to a single rotation about R , equal to twice the acute angle between QR, RP .

Let the first rotation about P bring Q to Q' and R to R' ; it is clear that $QPR, Q'PR, Q'PR'$ are all equal triangles. Therefore $R'Q'R = 2PQR = \beta$. Consequently the positive rotation β about Q' (the new position of Q) will carry R' back again to R , its original position. Hence the actual motion which results from the successive rotations combined being consistent with R remaining at rest, must be equivalent to a single rotation about R .

To find its magnitude, let the second rotation carry P to P'^{81} ; then the angular displacement PRP' (which is the required rotation of the whole body) is equal p. 159 to twice the acute angle between $Q'R, RP$, which is the same as that between QR, RP , as was to be shown. Thus we see that the semi-rotations about three poles (considered as the angular points of a spherical triangle), which, taken in order, would bring the sphere back to its first, undisturbed position, are equal to the included angles at such poles respectively.

If in our figure the order of the rotations had been reversed, PQr, QPr would

⁸¹The reader is requested to fill in the point P' and join $P'R$.

have been taken respectively equal to PQR, QPR , but on the opposite side of PQ , and r would have been the resultant pole, the resultant rotation remaining in amount the same as before.

If either of the rotations had been negative, the resultant pole would be found in QR produced, namely, at the intersection of rQ or rP with PQ .

Calling the resultant rotation γ , we have always

$$\sin \frac{\alpha}{2} : \sin \frac{\beta}{2} : \sin \frac{\gamma}{2} :: \sin QR : \sin RP : \sin PQ.$$

When the component rotations are infinitesimal in amount, R and r will come together in QP ; the order of succession of the rotations will be indifferent, and we shall have

$$\alpha : \beta : \gamma :: \sin \frac{\alpha}{2} : \sin \frac{\beta}{2} : \sin \frac{\gamma}{2} :: \sin QR : \sin RP : \sin PQ,$$

which gives the rule for the parallelogrammatic composition of two simultaneously impressed rotations⁸².

If, next, we consider the effect of rotations about two poles, P and Q , fixed in space (supposing, as above, that they take place first about P and then about Q), we must take QPr equal to half the contrary of the rotation about P , and PQr to half the direct rotation about Q (the angle being now taken positive which was on the first supposition negative, and vice versa⁸³); so that, retaining the original figure, the first rotation will bring r to R , and the second carry R back to r ; showing that r is the resultant pole, and that⁸³ $P'rP$, the resultant rotation, will be double the acute angle between Qr, rP , as in the former case.

To popular apprehension the important doctrine of uniaxial rotation may be made intelligible by the following mode of statement. Take a pocket-globe, open the case and roll about the sphere within it in any manner whatever; then closing the case, there will unavoidably remain two points on the terrestrial surface touching the same two points on the celestial surface as they were in apposition with before the sphere was so turned about in its case.

p. 160

It is right to bear in mind that the whole of this doctrine is comprised in, and convertible with, the following easy geometrical proposition relative to arcs of great circles on any spherical surface, including the plane as an extreme case.

“The arcs joining the extremities (each with each in either order) of two other equal arcs, subtend equal angles at either of the points of intersection of two great circles bisecting at right angles the first-named connecting arcs⁸⁴.”

⁸²Compare Mr Airy's *Tracts*, Art. “On Precession and Nutation.”

⁸³ P' is not expressed in the figure given.

⁸⁴This proposition will be seen to be immediately demonstrable, by the comparison of equal triangles, when viewed as the converse of this other. “The arcs (or right lines) joining the correspondent extremities of the bases of two similar isosceles spherical (or plane) triangles having a common vertex, are equal to each other.”

The spherico-triangular mode of compounding rotations given in the above simple disquisition may easily be made the parent of a whole brood of geometrical consequences, which, however, I must leave to the ingenuity and care of those who have a turn for this kind of invention.

But I ought not to omit to invite attention to a remarkable form, which may be imparted to the theorems above stated for the composition of finite rotations, or rather to a theorem which may be derived from them by an obvious process of inference.

Let $P, Q, R \dots X, Z$ be any number of points on a sphere capable of moving about its centre, joined together by arcs of great circles so as to form a spherical polygon. Imagine any number of rotations to take place about these points in succession as poles. It matters not which is considered the first pole of rotation, but the order of the circulation must be supposed given, as, for instance, $PQR \dots XZ$, or $QR \dots XZP$, or $R \dots XZPQ$, &c. This will be one order; the reverse order would be $PZX \dots RQ$, or $QPZX \dots R$, &c.

I shall suppose the circulation to be of the kind first above written. Now we may make two hypotheses:—

1. That the poles are fixed in space.
2. That they are fixed in the rotating body.

In the first case, let the rotations about the given poles $P, Q, R, S \dots X, Z$ be double the amounts which would serve to transport PQ to QR , QR to $RS \dots XZ$ to ZP respectively.

In the second case, let the rotations be double the amounts which would carry PZ to $ZX \dots SR$ to RQ , RQ to QP respectively. Then, on either supposition, the sum of the combined rotations is zero; or, to use a more convenient and suggestive form of expression, if the poles of rotation form a closed spherical polygon whose angles are respectively equal to the semi-rotations about the poles, the resultant rotation is zero.

p. 161

This proposition is immediately derivable from the fundamental one relative to three poles, given above, by dividing the polygon into triangles by arcs, joining any one of the poles with all the rest, or (as pointed out to me by my eminent friend Prof. W. Thomson) it becomes apparent as a particular case of a more general proposition, on representing the motion about the successive axes as effected by two equal pyramids having a common vertex at the centre of motion, of which the one is fixed in space, and the other is fixed in the revolving body and rolls over the first, so that the corresponding equal faces are successively brought into coincident apposition.

P.S. To find the pole of rotation whereby PQ may be brought into the position $P'Q'$, we may use the following simple construction.

Measure off from O the node of the great circles (or right lines) containing PQ and $P'Q'$, two distances in the proper direction upon each (four distinct

assumptions may be made), say OR and OS equal to one another and to the difference between OP and OP' , then the pole of rotation required, say E , is the centre of the circle described about ROS , and the amount of rotation is the angle subtended by OR or OS at E . The writer of this paper suggests that axis of displacement would be a convenient term for designating the line whereby any finite change in the position of a body moveable about a fixed centre may be brought about; a geometrical theory of rotation leading to the investigation of a very curious species of correlation, now opens upon the view, the general object of which may be stated as follows:

“Given upon a sphere or plane any curve considered as the locus of successive poles of instantaneous rotation, and the ratio of the rotation about each pole to its distance from the one that follows⁸⁵, to construct the curve of the poles of displacement, and to determine the amount of rotation corresponding to each such pole.”

The discussion of this question offers a fine field for the exercise of geometrical taste and skill.

⁸⁵Which by analogy may be termed the “density of rotation.”

29.

ON THE INTERSECTIONS OF TWO CONICS

[Cambridge and Dublin Mathematical Journal, vi. (1851), pp. 18–20]

p. 162

Let the two conics be written

$$\begin{aligned} U &= ax^2 + by^2 + cz^2 + 2a'yz + 2b'zx + 2c'xy = 0, \\ V &= \alpha x^2 + \beta y^2 + \gamma z^2 + 2\alpha'yz + 2\beta'zx + 2\gamma'xy = 0, \end{aligned}$$

and make

$$U + \lambda V = Ax^2 + By^2 + Cz^2 + 2A'yz + 2B'zx + 2C'xy.$$

In my paper in the last number of the *Journal*⁸⁶, I showed that the case of intersection of the two conics in two points was distinguishable from all other cases by the equation $\square(U + \lambda V) = 0$ having two imaginary roots. When all the roots are real, the curves either intersect in four points or not at all.

On the former supposition,

$$-C'^2 + AB, \quad -A'^2 + BC, \quad -B'^2 + CA,$$

which are quadratic functions of λ , will be negative for all three values of λ . On the contrary supposition, one value of λ will make all these three quantities negative, but the other two values with each make them all three positive.

Hence we obtain a symmetrical criterion (which I strangely omitted to state in my former paper) by forming the quantity

$$A^2 + B^2 + C^2 - AB - AC - BC.$$

A cubic equation

$$Ly^3 + My^2 + Ny + P = 0$$

may be then constructed, of which the three values of the above function corresponding to three values of λ will be the roots.

The condition for real intersection is that L, M, N, P should be all of the same sign. The conics being supposed real, L and P are necessarily in both cases of the same sign. The condition is therefore satisfied if either L, M, N , or M, N, P be of the same sign, and is consequently equivalent to the condition that $\frac{M}{L}$ and $\frac{N}{L}$ shall be both positive, or $\frac{N}{P}$ and $\frac{M}{P}$ both positive. It does not appear to be possible in the nature of the question to find a criterion for distinguishing between the two cases, dependent on the sign of one single function of the coefficients.

⁸⁶P. 119 above.

The case of double contact, abstraction being made of binary intersection, is a sort of intermediary state between intersection in four points and non-intersection; and accordingly, as shown in my former paper for this case, the two equal values of λ will make the three quantities

$$AB - C'^2, \quad BC - A'^2, \quad CA - B'^2$$

all real; so that two of the values of y corresponding to the equal values of λ are zero, and the criterion becomes nugatory as it ought to do.

Again, when the two conics do not intersect, I distinguished two cases according as they lie each without, or one within the other, that is, according as they have four common tangents or none.

But, as Mr Cayley has well remarked to me, a similar distinction exists when the conics intersect in four points; in that case also they may have four common tangents or not any: when they intersect in two points they have necessarily two and only two common tangents. There is no difficulty in separating these four cases.

Let the conics be written

$$(U) = \xi^2 + \eta^2 - \zeta^2, \quad (V) = A\xi^2 + B\eta^2 - C\zeta^2,$$

(U) and (V) being what U and V become when the coordinates are changed from x, y, z to ξ, η, ζ .

A, B, C are the three values of λ in the equation

$$\square(V - \lambda U) = 0.$$

If the curves intersect $A - C, B - C$ must have different signs, that is, C must be an intermediary quantity between A and B .

Again, the tangential equations to the conics expressed by the correlative system of coordinates will be

$$\xi_1^2 + \eta_1^2 - \zeta_1^2 = 0, \quad \frac{\xi_1^2}{A} + \frac{\eta_1^2}{B} - \frac{\zeta_1^2}{C} = 0;$$

and that these may have four real systems of roots,

$$\frac{1}{A} - \frac{1}{C}, \quad \frac{1}{C} - \frac{1}{B}$$

must have the same sign; and consequently, as $A - C$ and $C - B$ are supposed p. 164 to have the same sign, A and B , and therefore all three A, B, C , have the same sign. We have therefore the following rule:

Let the equation in λ , namely, $\square(U + \lambda V) = 0$, be called $\theta = 0$, and the equation in y , above given, $\omega = 0$. By an equation being congruent or incongruent, understand that its roots have all the same sign or not all the same sign.

Then ω congruent, θ congruent, implies that the intersections and common tangents are both real; ω congruent, θ incongruent, implies that the intersections are real, but the common tangents imaginary; ω incongruent, θ congruent, implies that the intersections and common tangents are both imaginary; ω incongruent, θ incongruent, implies that the intersections are imaginary, but the common tangents real.

In like manner, as the cases of contact of lines are limiting cases to those which relate to the relative configurations of their points of intersection, so the cases of contact of surfaces are limiting cases in which the characters which usually separate the different forms of their curve of intersection exist blended and indistinguishable. The first step therefore to the study of the particular species of the curve of the fourth degree⁸⁷ in which two surfaces of the second degree intersect, is to obtain the analytical and geometrical characters of their various species of contact. Accordingly I have made an enumeration of these different species, no less than 12 in number, many of them highly curious and I believe unsuspected, which the reader may consult in the *Philosophical Magazine* for February, 1851⁸⁸.

By the aid of these landmarks, I have little doubt, should time and leisure permit, of mapping out a natural arrangement of the principal distinctions of form between that class at least of lines in space of the fourth order which admit of being considered the complete intersection of two surfaces.

⁸⁷I have found that the 16 points of spherical flexure in this curve are the four sets of four points in which it meets the four faces of the pyramid whose summits are the vertices of the four cones of the second degree in which the curve is completely contained, which 16 points reduce to 4 when the two surfaces have an ordinary contact, and to 1 when they have a cuspidal contact: of course in the case of contact the pyramid above described in a manner folds up and vanishes, as there are no longer 4 distinct vertices. I have found also that when the factors of $D(U + \lambda V)$, (U and V being the characteristics of the two surfaces) are all unreal, the points of flexure are all unreal. When two factors are real and two imaginary, two of the faces of the pyramid (namely, its two real faces) will each contain one (and only one) pair of real points of flexure, and the other two planes none; and lastly, when the factors of $\square(U + \lambda V)$ are all real, then either all the points of flexure are imaginary, or else all the eight contained in a certain two of the pyramidal faces are real: and these two cases admit of being distinguished by a method analogous in its general features to that whereby I have shown in the text above how to distinguish between the cases of 4 real and 4 imaginary points of intersection of two conics. Where the two surfaces have an ordinary contact, the curve of intersection, it is well known, has a double point; and where the surfaces have a higher contact, the curve has a cusp. Thus in the fact of the 16 flexures reducing to 4 and to 1 in these respective cases, we see a beautiful analogy to what takes place with the 9 flexures of a plane curve of the third degree, which contract to 3 and 1, according as the curve has a double point or a cusp.

⁸⁸P. 219 below.

ON CERTAIN GENERAL PROPERTIES OF HOMOGENEOUS FUNCTIONS

[*Cambridge and Dublin Mathematical Journal*, vi. (1851), pp. 1-17]

p. 165

Let χ denote the operation

$$x_1 \frac{d}{da_1} + x_2 \frac{d}{da_2} + \cdots + x_n \frac{d}{da_n},$$

and A the operation

$$a_1 \frac{d}{dx_1} + a_2 \frac{d}{dx_2} + \cdots + a_n \frac{d}{dx_n};$$

and now suppose that ω , a homogeneous function of ι dimensions of $a_1, a_2 \dots a_n$, and not of any of the quantities $x_1, x_2 \dots x_n$, is subjected to the successive operations indicated by $A^s \chi^r$.

We have

$$\begin{aligned} A^s \chi^r \omega &= A^{s-1} A \chi^r \omega, \\ A \chi^r \omega &= \left(a_1 \frac{d}{dx_1} + a_2 \frac{d}{dx_2} + \cdots + a_n \frac{d}{dx_n} \right) \left(x_1 \frac{d}{da_1} + x_2 \frac{d}{da_2} + \cdots + x_n \frac{d}{da_n} \right)^r \omega \\ &= r \left(a_1 \frac{d}{da_1} + a_2 \frac{d}{da_2} + \cdots + a_n \frac{d}{da_n} \right) \chi^{r-1} \omega \\ &= r(\iota - r + 1) \chi^{r-1} \omega, \end{aligned}$$

for $\chi^{r-1} \omega$ is of $(r-1)$ dimensions, lower than ω (which is of ι dimensions) in $a_1, a_2 \dots a_n$.

Hence

$$\begin{aligned} A^s \chi^r \omega &= r(\iota - r + 1) A^{s-1} \chi^{r-1} \omega \\ &= \&c. \\ &= \{r(r-1) \cdots (r-s+1)\} \{(\iota - r + 1)(\iota - r + 2) \cdots (\iota - r + s)\} \chi^{r-s} \omega. \end{aligned} \tag{1} \text{ p. 166}$$

Now in the expression

$$\chi^r \omega(a_1, a_2 \dots a_n),$$

suppose that we write

$$\begin{aligned} x_1 &= u_1 + a_1 \varepsilon, \\ x_2 &= u_2 + a_2 \varepsilon, \\ &\dots\dots\dots \\ x_n &= u_n + a_n \varepsilon, \end{aligned}$$

we have, by Taylor's theorem,

$$\chi^r \omega = U^r \omega + A U^r \omega \varepsilon + A^2 U^r \omega \frac{\varepsilon^2}{1 \cdot 2} + \cdots + A^r U^r \omega \frac{\varepsilon^r}{1 \cdot 2 \cdot 3 \cdots r},$$

where $U^r\omega$ denotes what $\chi^r\omega$ becomes, on substituting u 's for x 's, and A now represents

$$u_1 \frac{d}{da_1} + u_2 \frac{d}{da_2} + \cdots + u_n \frac{d}{da_n}.$$

This expansion stops spontaneously at the $(r+1)$ th term, because $\chi^r\omega$ is only of r dimensions in $x_1, x_2 \dots x_n$.

Applying now theorem (1), we obtain

$$\begin{aligned} \chi^r\omega = & U^r\omega + r(\iota - r + 1)U^{r-1}\omega\varepsilon + \frac{1}{2}r(r-1)\{(\iota - r + 1)(\iota - r + 2)\}U^{r-2}\omega\varepsilon^2 \\ & + \cdots + \{(\iota - r + 1)(\iota - r + 2) \cdots \iota\}\omega\varepsilon^r. \end{aligned} \quad (2)$$

In using this theorem in the course of the ensuing pages, it will be found convenient to assign to ε a specific value, and I shall suppose it equal to $\frac{x_n}{a_n}$; this gives

$$u_1 = x_1 - \frac{a_1}{a_n}x_n, \quad u_2 = x_2 - \frac{a_2}{a_n}x_n, \quad \cdots \quad u_n = x_n - \frac{a_n}{a_n}x_n = 0.$$

And inasmuch as the U symbol now contains $a_1, a_2, \dots, a_n$, so that UU^r no longer equals U^{r+1} , I shall write U_r for U^r . Theorem (2) will thus assume the form

$$\begin{aligned} \chi^r\omega = & U_r\omega + r(\iota - r + 1)U_{r-1}\omega\frac{x_n}{a_n} + \frac{1}{2}r(r-1)(\iota - r + 1)(\iota - r + 2)U_{r-2}\omega\left(\frac{x_n}{a_n}\right)^2 \\ & + \cdots + \{(\iota - r + 1) \cdots \iota\}\omega\left(\frac{x_n}{a_n}\right)^r. \end{aligned} \quad (3)$$

p. 167

where U_r for all values of r denotes what

$$\left(x_1 \frac{d}{da_1} + x_2 \frac{d}{da_2} + \cdots + x_{n-1} \frac{d}{da_{n-1}}\right)^r \omega$$

becomes, on substituting $u_1, u_2, \dots, u_{n-1}$ for $x_1, x_2, \dots, x_{n-1}$, after the processes of derivation have been completed: this it is essential to observe, because $u_1, u_2, \dots, u_{n-1}$ now involve $a_1, a_2, \dots, a_{n-1}, a_n$. The term $x_n \frac{d}{da_n}$ is omitted from the symbol of linear derivation, because in the substitutions x_n will be replaced by zero.

As an example of this last theorem, take

$$\omega = a^3 + b^3 + c^3 + kabc;$$

then

$$\begin{aligned} \chi\omega &= 3a^2x + 3b^2y + 3c^2z + kbcx + kca y + kabz, \\ \chi^2\omega &= 6ax^2 + 6by^2 + 6cz^2 + 2kctxy + 2kayz + 2kbzx, \\ \chi^3\omega &= 6x^3 + 6y^3 + 6z^3 + 6kxyz. \end{aligned}$$

$$\begin{aligned}
U_1\omega &= 3a^2 \left(x - \frac{az}{c}\right) + 3b^2 \left(y - \frac{bz}{c}\right) + kbc \left(x - \frac{az}{c}\right) + kca \left(y - \frac{bz}{c}\right), \\
U_2\omega &= 6a \left(x - \frac{az}{c}\right)^2 + 6b \left(y - \frac{bz}{c}\right)^2 + 2kc \left(x - \frac{az}{c}\right) \left(y - \frac{bz}{c}\right), \\
U_3\omega &= 6 \left(x - \frac{az}{c}\right)^3 + 6 \left(y - \frac{bz}{c}\right)^3,
\end{aligned}$$

and it will be found that the equations given by theorem (3) are satisfied, namely

$$\begin{aligned}
\chi\omega &= U\omega + 3\frac{z}{c}\omega, \\
\chi^2\omega &= U_2\omega + 2 \cdot 2\frac{z}{c}U\omega + 2 \cdot 3\frac{z^2}{c^2}\omega, \\
\chi^3\omega &= U_3\omega + 3\frac{z}{c}U_2\omega + 3 \cdot 1 \cdot 2\frac{z^2}{c^2}U\omega + 1 \cdot 2 \cdot 3\frac{z^3}{c^3}\omega.
\end{aligned}$$

Probably, as this theorem is of rather a novel character, the annexed sketch of a somewhat different course of demonstration may be not unacceptable to my readers.

We have

$$\chi\omega = \left(x_1 \frac{d}{da_1} + x_2 \frac{d}{da_2} + \cdots + x_n \frac{d}{da_n}\right)\omega;$$

and by the well-known law for homogeneous functions,

$$\iota\omega = \left(a_1 \frac{d}{da_1} + a_2 \frac{d}{da_2} + \cdots + a_n \frac{d}{da_n}\right)\omega.$$

p. 168

Hence

$$\left(\chi - \iota \frac{x_n}{a_n}\right)\omega = \left(u_1 \frac{d}{da_1} + u_2 \frac{d}{da_2} + \cdots + u_{n-1} \frac{d}{da_{n-1}}\right)\omega = U\omega.$$

Hence

$$\begin{aligned}
\chi\omega &= \left(U + \iota \frac{x_n}{a_n}\right)\omega, \\
\chi^2\omega &= \left\{U + (\iota - 1)\frac{x_n}{a_n}\right\} \left(U + \iota \frac{x_n}{a_n}\right)\omega, \\
\chi^3\omega &= \left\{U + (\iota - 2)\frac{x_n}{a_n}\right\} \left\{U + (\iota - 1)\frac{x_n}{a_n}\right\} \left(U + \iota \frac{x_n}{a_n}\right)\omega, \\
&\&c. = \&c.
\end{aligned}$$

But in performing the process indicated by the several factors it must be carefully borne in mind that UU_r is not $= U_{r+1}$; this would be the case were it not for the terms $-\frac{a_1}{a_n}x_n$, $-\frac{a_2}{a_n}x_n$, &c., which enter into $u_1, u_2 \dots u_{n-1}$. But on account of these terms, we have

$$\begin{aligned}
UU_r\omega &= \left(u_1 \frac{d}{da_1} + u_2 \frac{d}{da_2} + \cdots + u_{n-1} \frac{d}{da_{n-1}}\right) \left(u_1 \frac{d}{da_1} + u_2 \frac{d}{da_2} + \cdots + u_{n-1} \frac{d}{da_{n-1}}\right)^r \omega \\
&= U_{r+1}\omega - r \frac{x_n}{a_n} \left(u_1 \frac{d}{da_1} + u_2 \frac{d}{da_2} + \cdots + u_{n-1} \frac{d}{da_{n-1}}\right)^{r-1} \omega,
\end{aligned}$$

for

$$\frac{d}{da_1}u_1 = \frac{d}{da_2}u_2 = \cdots = \frac{d}{da_{n-1}}u_{n-1} = -\frac{x_n}{a_n}.$$

Hence

$$UU_r\omega = U_{r+1}\omega - r\frac{x_n}{a_n}U_r\omega.$$

Let $\frac{x_n}{a_n}$ be called ε ; we find

$$\begin{aligned}\chi &= U + \iota\varepsilon, \\ \chi^2 &= \{U + (\iota - 1)\varepsilon\}(U + \iota\varepsilon) \\ &= UU + (2\iota - 1)\varepsilon U + (\iota - 1)\iota\varepsilon^2 \\ &= U_2 + 2(\iota - 1)\varepsilon U + (\iota - 1)\iota\varepsilon^2; \\ \chi^3 &= \{U + (\iota - 2)\varepsilon\}\chi^2 \\ &= UU_2 + 2(\iota - 1)\varepsilon UU + (\iota - 1)\iota\varepsilon^2 U \\ &\quad + (\iota - 2)\varepsilon U_2 + 2(\iota - 2)(\iota - 1)\varepsilon^2 U + (\iota - 2)(\iota - 1)\iota\varepsilon^3 \\ &= U_3 + 3(\iota - 2)\varepsilon U_2 + 3(\iota - 2)(\iota - 1)\varepsilon^2 U + (\iota - 2)(\iota - 1)\iota\varepsilon^3.\end{aligned}$$

The same process being continued will lead to results identical with those previously obtained and expressed in theorem (3).

The expansion of χ^r , treated according to this second method, appears to require the solution of the partial equation in differences

$$a_{r+1,s+1} = a_{r,s+1} + (\iota - 2r)a_{r,s},$$

$a_{0,s}$ being given as unity for $s = 1$ and as zero for all other values of s .

It is probable however that the solution of this equation might be evaded by some artifice peculiar to the particular case to be dealt with. I do not propose to dwell upon this inquiry, which would be foreign to the object of my present research. It may however not be out of place to make the passing remark, that the equations expressing χ^r in terms of powers of U admit easily of being reverted, as indeed may the more general form

$$\chi_r = u_r + \varepsilon_r u_{r-1} + \frac{1}{1 \cdot 2} \varepsilon_r \varepsilon_{r-1} u_{r-2} + \&c.,$$

which becomes the equation of formula (3), on making

$$\varepsilon_r = r(\iota + 1 - r)\frac{x_n}{a_n}, \quad \chi_r = \chi^r\omega, \quad u_r = U_r\omega;$$

for let

$$u_r = \varepsilon_1 \varepsilon_2 \cdots \varepsilon_r v_r, \quad \chi_r = \varepsilon_1 \varepsilon_2 \cdots \varepsilon_r y_r,$$

then

$$y_r = v_r + v_{r-1} + \frac{v_{r-2}}{1 \cdot 2} + \frac{v_{r-3}}{1 \cdot 2 \cdot 3} + \&c.;$$

whence

$$v_r = e^{-\frac{d}{dr}} y_r = y_r - y_{r-1} + \frac{y_{r-2}}{1 \cdot 2} - \frac{y_{r-3}}{1 \cdot 2 \cdot 3} + \&c.;$$

and therefore

$$u_r = \chi_r - \varepsilon_r \chi_{r-1} + \frac{1}{2} \varepsilon_r \varepsilon_{r-1} \chi_{r-2} + \&c.$$

Thus we obtain, from equation (3),

$$U_r \omega = \chi^r \omega - r(\iota - r + 1) \chi^{r-1} \omega \frac{x_n}{a_n} + \&c.$$

As a first application of theorem (3), I shall proceed to show how Joachimsthal's equation to the surface drawn from a given point $(\alpha, \beta, \gamma, \delta)$ through the intersection of two surfaces $\phi(x, y, z, t) = 0$, $\theta(x, y, z, t) = 0$, may be expressed under the explicit form of the equation to a cone.

The equation in question is obtained by eliminating λ between

$$\phi \lambda^m + \chi \phi \lambda^{m-1} + \frac{1}{1 \cdot 2} \chi^2 \phi \lambda^{m-2} + \&c. = 0,$$

$$\theta \lambda^n + \chi \theta \lambda^{n-1} + \frac{1}{1 \cdot 2} \chi^2 \theta \lambda^{n-2} + \frac{1}{1 \cdot 2 \cdot 3} \chi^3 \theta \lambda^{n-3} + \&c. = 0,$$

where

p. 170

$$\phi = \phi(\alpha, \beta, \gamma, \delta), \quad \theta = \theta(\alpha, \beta, \gamma, \delta), \quad \chi = x \frac{d}{d\alpha} + y \frac{d}{d\beta} + z \frac{d}{d\gamma} + t \frac{d}{d\delta}.$$

By theorem (3), these two equations, on writing $\frac{x_n}{a_n} = \varepsilon$, become

$$\begin{aligned} & \phi \lambda^m + \{U\phi + m\phi\varepsilon\} \lambda^{m-1} + \{U^2\phi + 2(m-1)U\phi\varepsilon + (m-1)m\phi\varepsilon^2\} \frac{\lambda^{m-2}}{1 \cdot 2} + \&c. = 0, \\ & \theta \lambda^n + \{U\theta + n\theta\varepsilon\} \lambda^{n-1} + \{U^2\theta + \&c.\} \frac{\lambda^{n-2}}{1 \cdot 2} \\ & + \{U^3\theta + 3(n-2)U^2\theta\varepsilon + 3(n-2)(n-1)U\theta\varepsilon^2 + (n-2)(n-1)n\varepsilon^3\} \frac{\lambda^{n-3}}{1 \cdot 2 \cdot 3} + \&c. = 0. \end{aligned}$$

Now on writing $\lambda = \mu - \varepsilon$, these equations take the forms

$$\phi \mu^m + U\phi \mu^{m-1} + U^2\phi \frac{\mu^{m-2}}{1 \cdot 2} + \&c. = 0,$$

$$\theta \mu^n + U\theta \mu^{n-1} + U^2\theta \frac{\mu^{n-2}}{1 \cdot 2} + \&c. = 0,$$

as is easily seen by substituting back $\lambda + \varepsilon$ in place of μ . Consequently ε no longer appears in the coefficients of the terms of the equations between which the elimination is to be performed, and the resultant will accordingly come out as a function only of $\phi, U\phi, U^2\phi$, &c., that is, of $\alpha, \beta, \gamma, \delta$, and of

$$x - \frac{\alpha}{\delta} t, \quad y - \frac{\beta}{\delta} t, \quad z - \frac{\gamma}{\delta} t,$$

showing that the equation in x, y, z, t , is of the form of that to a cone, as we know *à priori* it ought to be. Precisely a similar method may be applied to the elucidation of the corresponding theorem for a system of rays drawn from a given point through the locus of the intersection of two curves.

Before entering upon some further and more interesting applications of theorem (3), it will be convenient to explain a nomenclature which has been employed by me on another occasion, and which is almost indispensable in inquiries of the nature we are now engaged upon. Homogeneous functions may be characterized by their degree, by the number of letters which enter into them, and lastly, by the lowest number of linear functions of the letters which may be introduced in place of the letters to represent such functions. Any such linear function I designate as an order, and am now able to discriminate between the number of letters and the number of orders which enter into a given function. The latter number, generally speaking, is the same as the former; it can never exceed it, but may be any number of units less than it.

p. 171

I need scarcely observe that a pair of points becoming coincident, a conic becoming a pair of lines, a conoid becoming a cone, and so forth, for the higher realms of space, will be expressed by the homogeneous function of the second order which characterizes such loci⁸⁹, losing one order, that is, having an order less than the number of letters entering therein. Calling such characteristic $\phi(x, y, z \dots t)$, it is well known that the condition of such loss of an order is the vanishing of the determinant

$$\begin{vmatrix} \frac{d^2\phi}{dx^2} & \frac{d^2\phi}{dxdy} & \dots & \frac{d^2\phi}{dxdt} \\ \frac{d^2\phi}{dydx} & \frac{d^2\phi}{dy^2} & \dots & \frac{d^2\phi}{dydt} \\ \dots & \dots & \dots & \dots \\ \frac{d^2\phi}{dtdx} & \frac{d^2\phi}{dtdy} & \dots & \frac{d^2\phi}{dt^2} \end{vmatrix}.$$

A conoid becoming a pair of planes, a cone becoming a pair of coincident lines, a pair of points becoming indeterminate, will, in like manner, be denoted by their characteristic losing two orders, and so forth, for the higher degrees of degradation. In like manner, in general, a homogeneous function of three letters of any degree losing an order, typifies that the locus to which it is the characteristic will break up into a system of right lines.

Now let ω be a homogeneous function of $\alpha, \beta, \gamma \dots \delta$, and suppose that we have the equations $\omega = 0$, $\chi\omega = 0$, $\chi^2\omega = 0$, where χ as above

$$= x \frac{d}{d\alpha} + y \frac{d}{d\beta} + z \frac{d}{d\gamma} + \dots + t \frac{d}{d\delta}.$$

⁸⁹If $U = 0$ is the equation to any locus, U may be said to characterize the same, or to be its characteristic.

I say that on eliminating any of the variables $x, y, z \dots t$ between the second and third of the above equations, the resulting equation will be of one order less than the number of letters, that is, the expulsion of one letter will be attended by the expulsion of two orders.

For we have, by theorem (3),

$$\chi\omega = U\omega + 2\frac{x_n}{a_n}\omega = 0,$$

$$\chi^2\omega = U_2\omega + 2\frac{x_n}{a_n}U\omega + 2\left(\frac{x_n}{a_n}\right)^2\omega = 0,$$

and by hypothesis

$$\omega = 0.$$

Hence we have also

$$U\omega = 0, \quad U_2\omega = 0;$$

and since $U\omega, U_2\omega$ contain one order less than the number of letters in ω , the resultant of the elimination between them will contain two orders less than the number of letters in ω ; and consequently, whichever of the letters $x, y, z \dots t$ we eliminate between $\chi\omega = 0$ and $\chi^2\omega = 0$, provided that $\omega = 0$, the resultant equation will contain one order less than the number of letters remaining. p. 172

Thus we see how it is that the tangent line to a conic meets it in two coincident points, the tangent plane to a conoid in two intersecting lines, and so forth, for the higher regions of space⁹⁰. For instance, if we take $\omega(x, y, z, t) = 0$, the equation to a conoid, and $\alpha, \beta, \gamma, \delta$, the coordinates to any point therein, we shall have $\omega(\alpha, \beta, \gamma, \delta) = 0$,

$$\left(x\frac{d}{d\alpha} + y\frac{d}{d\beta} + z\frac{d}{d\gamma} + t\frac{d}{d\delta}\right)\omega, \quad \text{that is, } \chi\omega = 0,$$

and $\omega(x, y, z, t)$, that is, $\chi^2\omega = 0$, x, y, z, t representing the coordinates of any point in the intersection of the conoid by the tangent plane.

Consequently, by what has been shown above, on eliminating any one of the four letters x, y, z, t , the resultant function of three letters will contain only two orders, and will thus represent a pair of lines, real or imaginary, intersecting one another at $\alpha, \beta, \gamma, \delta$.

The fact which has just been demonstrated (that the resultant of $\chi\omega = 0$, $\chi^2\omega = 0$, loses an order if $\omega = 0$), indicates that on expressing one of the quantities $x, y, z \dots t$ in terms of the others, by means of the first equation, and then substituting this value in the second, the determinant of the equation so obtained must be zero.

⁹⁰Thus a tangential section of a hyperlocus of the second degree at any point cuts it in two cones.

Now by virtue of a theorem which was given by me in a note⁹¹ to my paper in the preceding number of this *Journal*, this determinant will be equal to the squared reciprocal of the coefficient in the equation $\chi\omega = 0$ of the letter eliminated multiplied by the determinant in respect to $x, y, z \dots t, \lambda$ of

$$\chi^2\omega + \lambda\chi\omega.$$

This latter determinant is therefore zero; but this determinant is the resultant of the equations

$$\left\{ \begin{array}{l} \frac{d}{dx} \left(x \frac{d}{da} + y \frac{d}{db} + \&c. \right)^2 \omega + \frac{d}{dx} \left(x \frac{d}{da} + y \frac{d}{db} + \dots \right) \omega = 0, \\ \frac{d}{dy} \left(x \frac{d}{da} + y \frac{d}{db} + \&c. \right)^2 \omega + \frac{d}{dy} \left(x \frac{d}{da} + y \frac{d}{db} + \dots \right) \omega = 0, \\ \qquad \qquad \qquad \&c. \qquad \&c. \qquad \&c. \qquad \&c., \\ \chi\omega = 0, \quad \text{that is,} \quad \left(x \frac{d}{da} + y \frac{d}{db} + \dots \right) \omega = 0. \end{array} \right.$$

p. 173

Thus we obtain the singular law, that the symmetrical determinant

$$\left| \begin{array}{ccccc} \frac{d}{da} \frac{d}{da} \omega & \frac{d}{da} \frac{d}{db} \omega & \dots & \frac{d}{da} \frac{d}{dl} \omega & \frac{d}{da} \omega \\ \frac{d}{db} \frac{d}{da} \omega & \frac{d}{db} \frac{d}{db} \omega & \dots & \frac{d}{db} \frac{d}{dl} \omega & \frac{d}{db} \omega \\ \frac{d}{dc} \frac{d}{da} \omega & \frac{d}{dc} \frac{d}{db} \omega & \dots & \frac{d}{dc} \frac{d}{dl} \omega & \frac{d}{dc} \omega \\ \dots \frac{d}{dl} \frac{d}{da} \omega & \dots \frac{d}{dl} \frac{d}{db} \omega & \dots & \dots \frac{d}{dl} \frac{d}{dl} \omega & \dots \frac{d}{dl} \omega \\ \frac{d}{da} \omega & \frac{d}{db} \omega & \dots & \frac{d}{dl} \omega & 0 \end{array} \right|$$

is zero when ω is zero.

This is easily shown independently by means of a remarkable and I believe novel theorem, relative to homogeneous functions.

If ω be any homogeneous function of ι dimensions of $a, b, c \dots l$, we have (by Euler's theorem already repeatedly applied), remembering that $\frac{d\omega}{da}, \frac{d\omega}{db} \dots \frac{d\omega}{dl}$ are all homogeneous,

$$\begin{aligned} -\iota\omega + \left(a \frac{d}{da} + b \frac{d}{db} + \dots + l \frac{d}{dl} \right) \omega &= 0, \\ -(\iota - 1) \frac{d\omega}{da} + \left(a \frac{d}{da} \frac{d}{da} + b \frac{d}{da} \frac{d}{db} + \dots + l \frac{d}{da} \frac{d}{dl} \right) \omega &= 0, \\ -(\iota - 1) \frac{d\omega}{db} + \left(a \frac{d}{db} \frac{d}{da} + \dots + l \frac{d}{db} \frac{d}{dl} \right) \omega &= 0, \end{aligned}$$

⁹¹P. 135 above.

$$\begin{array}{ccc} \&c. & \&c. & \&c. \\ -(\iota-1)\frac{d\omega}{dl} + \left(a\frac{d}{dl}\frac{d}{da} + \cdots + l\frac{d}{dl}\frac{d}{dl}\right)\omega = 0. \end{array}$$

Between these equations we may eliminate all the letters, $a, b, c \dots l$, and we obtain

$$\begin{vmatrix} \frac{d}{da}\frac{d}{da}\omega & \frac{d}{da}\frac{d}{db}\omega & \cdots & \frac{d}{da}\frac{d}{dl}\omega & \frac{d}{da}\omega \\ \frac{d}{db}\frac{d}{da}\omega & \frac{d}{db}\frac{d}{db}\omega & \cdots & \frac{d}{db}\frac{d}{dl}\omega & \frac{d}{db}\omega \\ \frac{d}{dc}\frac{d}{da}\omega & \frac{d}{dc}\frac{d}{db}\omega & \cdots & \frac{d}{dc}\frac{d}{dl}\omega & \frac{d}{dc}\omega \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \frac{d}{dl}\frac{d}{da}\omega & \frac{d}{dl}\frac{d}{db}\omega & \cdots & \frac{d}{dl}\frac{d}{dl}\omega & \frac{d}{dl}\omega \\ \frac{d}{da}\omega & \frac{d}{db}\omega & \cdots & \frac{d}{dl}\omega & \frac{\iota}{\iota-1}\omega \end{vmatrix} = 0.$$

p. 174

As a corollary to this theorem, we see that if $\omega = 0$ the determinant obtained in the previous investigation becomes zero, agreeing with what has been already shown; in fact the last-named determinant is always equal to

$$\frac{\iota-1}{\iota}\omega \times \begin{vmatrix} \frac{d}{da}\frac{d}{da}\omega & \cdots & \frac{d}{da}\frac{d}{dl}\omega \\ \vdots & \ddots & \vdots \\ \frac{d}{dl}\frac{d}{da}\omega & \cdots & \frac{d}{dl}\frac{d}{dl}\omega \end{vmatrix}.$$

This remarkable theorem, which I have communicated to friends nearly a twelve-month back, is here, I believe, published for the first time⁹²

Suppose next that $\omega(x, y, z)$ is the characteristic of a line of any degree, to which a tangent is drawn at the point α, β, γ , using U in a manner correspondent to its previous signification to denote

$$\left(x - \frac{\alpha z}{\gamma}\right) \frac{d}{d\alpha} + \left(y - \frac{\beta z}{\gamma}\right) \frac{d}{d\beta},$$

and understanding $\omega(\alpha, \beta, \gamma)$ by ω , we have for determining the point of intersection, $\omega = 0$, $\chi\omega = 0$, $\chi^2\omega = 0$; and consequently, by aid of our theorem (3), we shall obtain

$$\omega = 0, \quad U\omega = 0, \quad U_n\omega + n\frac{z}{\gamma}U_{n-1}\omega + \cdots = 0.$$

⁹²Thus let z be a homogeneous function in x and y of ι dimensions, and let

$$\frac{dz}{dx}, \quad \frac{dz}{dy}, \quad \frac{d^2z}{dx^2}, \quad \frac{d^2z}{dxdy}, \quad \frac{d^2z}{dy^2}$$

be called p, q, r, s, t ; we shall find

$$\begin{vmatrix} r & s & p \\ s & t & q \\ p & q & \frac{\iota}{\iota-1}\omega \end{vmatrix} = 0,$$

that is,

$$\omega = \frac{\iota-1}{\iota} \frac{rq^2 - 2pqs + tp^2}{rt - s^2}.$$

By means of the two latter equations, we obtain

$$\left(x - \frac{\alpha z}{\gamma}\right)^2 F\left(x - \frac{\alpha z}{\gamma}\right) = 0,$$

$$\left(y - \frac{\beta z}{\gamma}\right)^2 G\left(y - \frac{\beta z}{\gamma}\right) = 0,$$

where F and G are each of only $(n - 2)$ dimensions, and serve to determine the intersections of the tangent with the curve, extraneous to the two coincident ones at the point of contact. p. 175

Again, suppose that ω is a function of any degree of any number of letters α, β, γ , &c., and that we have given $\omega = 0, \chi\omega = 0, \chi^2\omega = 0, \dots \chi^m\omega = 0$; it is evident from our fundamental theorem that these equations may be replaced by

$$\omega = 0, \quad U_1\omega = 0, \quad U_2\omega = 0, \quad \dots \quad U_m\omega = 0;$$

and consequently that the expulsion of $(m - 1)$ letters, by aid of the last m of the given equations, will be attended by the disappearance of m orders, or, in other words, the resultant will be minus an order, that is, will have one order less than the number of letters remaining in it.

In applying to space conceptions the preceding theorem, it will be convenient to use a general nomenclature for geometrical species of various dimensions.

Thus we may call a line a monotheme, a surface a ditheme, the species beyond a tritheme, and so on, *ad infinitum*.

A system of points according to the same system of nomenclature would be called a kenotheme.

An n -theme has for its characteristic a homogeneous function of $(n + 2)$ letters.

Again, it will be convenient to give a general name to all themes expressed by equations of the first degree. Right lines and planes agree in conveying an idea of levelness and uniformity; they may both be said to be homalous. I shall therefore employ the word homaloid to signify in general any theme of the first degree.

Now let $\omega(x, y, z \dots t)$ be the characteristic to an n -theme of the n th degree.

The number of letters $x, y, z \dots t$ is $(n + 2)$.

As usual, let ω represent $\omega(\alpha, \beta, \gamma \dots \delta)$, and suppose

$$\omega = 0, \quad \chi\omega = 0, \quad \chi^2\omega = 0 \dots \chi^n\omega = 0,$$

and consequently

$$U_1\omega = 0, \quad U_2\omega = 0 \dots U_n\omega = 0.$$

On eliminating $(n - 1)$ letters between the n last equations, the resulting function will be of three letters but of only two orders, and of the $1 \cdot 2 \cdot 3 \dots n$ degree.

Hence we see that at every point of an n -theme of the n th degree, and lying p. 176 in the tangent homaloid thereto, $1 \cdot 2 \cdots n$ right lines may be drawn coinciding throughout with the n -theme.

Thus one right line can be drawn at each point of a line of the first order lying on the line; two right lines at each point of a surface of the second order lying on the surface; six right lines at each point of a hyperlocus of the third degree, and so forth.

It is obvious that a surface may be treated as the homaloidal section of a tritheme, just as a plane curve may be regarded as a section of a surface. I shall proceed to show upon this view, how we may obtain a theorem given by Mr Salmon for surfaces of the third degree of a particular character from the law just laid down, according to which a tritheme of the third degree admits of six right lines being drawn upon it at every point⁹³.

Let $\omega(x, y, z, t, u)$ be the characteristic of any tritheme of the third degree; $\alpha, \beta, \gamma, \delta, \epsilon$, coordinates to any point in the same. Then $\omega(\alpha, \beta, \gamma, \delta, \epsilon) = 0$, and the equation to the tangent homaloid will be $\chi\omega(\alpha, \beta, \gamma, \delta, \epsilon) = 0$, and the equation to the polar of the second degree to the given tritheme in relation to the assumed point as origin, (that is, the infinite system of homaloids that may be drawn from the point to touch the tritheme), will be $\chi^2\omega(\alpha, \beta, \gamma, \delta, \epsilon) = 0$.

But the section of any polar through its origin is the polar of the section to the same origin; hence the polar to the intersection of $\omega(x, y, z, t, u) = 0$, with $\chi\omega(\alpha, \beta, \gamma, \delta, \epsilon) = 0$, is the intersection of $\chi\omega = 0$ with $\chi^2\omega = 0$.

The projections of these intersections upon the space x, y, z, t will be found by eliminating u , and getting the correspondent two equations between x, y, z, t . Hence we see that the projection of the latter intersection upon any space x, y, z, t is a cone; or, in other words, this intersection itself, that is, the polar to the intersection of the tritheme with its tangent homaloid, is a cone; that is to say, the surface of the third degree formed by cutting a tritheme of the third degree by any tangent homaloid has a conical point at the point of contact; so that every surface of the third degree with a conical point may be considered as the intersection of a tritheme of the third degree with any tangent homaloid thereto⁹⁴.

p. 177

⁹³The reduction of any equation of the sixth degree to depend upon one of the fifth may be shown by Mr Jerrard's method to be equivalent to drawing a straight line upon a tritheme of the third degree, just as the reduction of the equation of the fifth degree to a trinomial form may be shown to be dependent upon our being able to draw a straight line upon a ditheme of the second degree. Now at every point of a tritheme straight lines may be drawn, but as they keep together in groups of sixes they cannot be found in general at a given point without solving an equation of the sixth degree.

⁹⁴So in like manner a surface of the third degree with more than one conical point may be generated by the intersection of the tritheme with a pluri-tangent plane; and so too we may get other varieties by taking homaloidal sections of trithemes whose characteristics are minus one or more orders.

Hence then we see, as an instantaneous deduction from our general theorem, that at any conical point (when one exists) of a surface of the third degree six right lines may be drawn lying completely upon it. This theorem is thus brought into an immediate and natural connexion with the well-known one, that at every point in a surface of the second degree, two right lines can be drawn lying wholly upon the surface⁹⁵.

The last geometrical application of the theorem (3) which I shall make, refers to the equations employed by Mr Salmon in No. xxi. (New Series) of this *Journal*, to obtain the locus of the points on any surface at which four consecutive tangent lines can be drawn passing through points. I remark in passing that these equations may be obtained by rather simpler considerations than Mr Salmon has employed so to do, and without any reference to Joachimsthal's theorem; for if we take ξ, η, ζ, θ , as the coordinates of any point in one of the tangent lines above described, and if we take the first polar to the surface with this point as origin, three out of the four original points will be found in such polar consecutive but distinct; and consequently in the second polar, referred to the same origin, two will continue consecutive but distinct, and consequently one will remain over in the third polar.

Hence writing the equation to the surface $\omega(x, y, z, t) = 0$, and using D to denote $\xi \frac{d}{dx} + \eta \frac{d}{dy} + \zeta \frac{d}{dz} + \theta \frac{d}{dt}$, we shall evidently have

$$\omega = 0, \quad D\omega = 0, \quad D^2\omega = 0, \quad D^3\omega = 0,$$

as obtained by Mr Salmon. And the same kind of reasoning precisely applies to the theory of points of inflexion in curves; three consecutive points in a right line in this case corresponding to four such in the case above considered.

If now we make

$$\xi - \frac{x}{t}\theta = u, \quad \eta - \frac{y}{t}\theta = v, \quad \zeta - \frac{z}{t}\theta = w,$$

the equations (2), (3), (4), by our theorem, may be expressed in terms of p. 178
 u, v, w , which being eliminated we obtain an equation between x, y, z, t , which will express the surface whose intersection with the given surface $\omega = 0$ serves to determine the locus of the points in question.

Hence if we proceed in the ordinary manner to eliminate two of the four letters, as ξ and η , between the equations (2), (3), (4), the resultant will be

⁹⁵If we have an indeterminate system of algebraical equations consisting of one quadratic and another n^c function of three variables, this may be completely resolved by considering the first as an equation to a surface of the second degree, finding at any point thereof the two lines which lie upon the surface, and determining their respective intersections with the surface represented by the second equation. This will require therefore the solution only of a quadratic and an n^c equation. In like manner an indeterminate system of two equations of four variables, one of the third and the other of the n th degree, may be completely resolved (with the aid of the theorem in the text) by means of two equations, one of the sixth and the other of the n th degree.

of the form $M \times \phi(\zeta, \theta)$, where M does not contain ξ, η, ζ or θ , and where by the general laws of elimination $\phi(\zeta, \theta)$ will be an integral function of the sixth degree in respect to ζ, θ : and it is manifest that $M \times \phi(\zeta, \theta)$ will be identical with the resultant of (2), (3), (4) expressed in terms of u, v, w , when u and v are eliminated *cy-près* of an integralizing factor, showing that $\phi(\zeta, \theta)$ is w^6 integralized, that is, is equal to $(t\zeta - z\theta)^6$. Consequently as $M\phi$ is of the order $(n-1)2 \cdot 3 + (n-2)1 \cdot 3 + (n-3)1 \cdot 2$, that is, $11n - 18$ in respect to x, y, z, t , it follows that $M = 0$, the equation to the second surface spoken of above, will be of the order $11n - 24$, agreeable to Mr Salmon's showing.

I shall conclude this paper by showing the application of our theorem to the subject propounded by Mr Jerrard and Sir William Hamilton, of systems of equations containing a sufficient number of variable letters for effecting the solution without elevation of degree.

If we have n homogeneous equations containing a sufficient number of letters $a_1, a_2 \dots a_m$ to enable us to express the solution of $(n-1)$ of the equations under the form

$$a_1 = \alpha_1 + \lambda\beta_1, \quad a_2 = \alpha_2 + \lambda\beta_2, \quad \dots \quad a_m = \alpha_m + \lambda\beta_m,$$

where $\alpha_1, \alpha_2 \dots \alpha_m, \beta_1, \beta_2 \dots \beta_m$ are supposed known, and λ is indeterminate, it is evident that by substituting these values in the n th equation, λ may be found by solving an equation of the same degree as that equation contains dimensions of $a_1, a_2 \dots a_m$.

Let us then propose this question: how many letters $a_1, a_2 \dots a_r$ are needed to obtain a linear solution of a system of n equations

$$\phi_1 = 0, \quad \phi_2 = 0, \quad \dots \quad \phi_n = 0,$$

of the several degrees $\iota_1, \iota_2 \dots \iota_n$, without elevation of degree; by a linear solution being understood a solution under the form

$$a_1 = \alpha_1 + \lambda\beta_1, \quad a_2 = \alpha_2 + \lambda\beta_2, \quad \dots \quad a_r = \alpha_r + \lambda\beta_r,$$

where λ is left indeterminate.

p. 179

Let us suppose that $\alpha_1, \alpha_2 \dots \alpha_r$, substituted respectively for $a_1, a_2 \dots a_r$, satisfy the given system of equations. The determination of these values without elevation of degree will, from what has been said before, depend upon the linear solution of a system of equations differing from the given system by the omission of any one of them at pleasure.

Now make

$$D = a_1 \frac{d}{da_1} + a_2 \frac{d}{da_2} + \dots + a_r \frac{d}{da_r},$$

and then write

$$\left. \begin{aligned} D\phi_1 &= 0, \\ D^2\phi_1 &= 0 \dots D^{\iota_1}\phi_1 = 0, \\ D\phi_2 &= 0, \quad D^2\phi_2 = 0 \dots D^{\iota_2}\phi_2 = 0, \\ &\dots\dots\dots \\ D\phi_n &= 0, \quad D^2\phi_n = 0 \dots D^{\iota_n}\phi_n = 0 \end{aligned} \right\} \quad (\theta)$$

The values of $a_1, a_2 \dots a_r$ derived from this system, say $(a)_1, (a)_2 \dots (a)_r$, give

$$a_1 = \alpha_1 + \lambda(a)_1, \quad a_2 = \alpha_2 + \lambda(a)_2, \dots a_r = \alpha_r + \lambda(a)_r,$$

a solution under the required form, where λ is left indeterminate.

The solution of this new system without elevation of degree depends on the linear solution of all but one of them; this excepted one may be taken the one whose dimensions ι_r are the highest or as high as any of the quantities $\iota_1, \iota_2 \dots \iota_n$.

Consequently, if we use the symbol $(k_1, k_2 \dots k_r)$ to denote the number of letters required for the linear solution (without elevation of degree) of k_1 equations of the first degree, k_2 of the second, k_3 of the third, $\dots$, k_r of the r th, it would at first sight appear from the preceding reduction that we must have

$$(k_1, k_2 \dots k_r) = \{K_1, K_2 \dots K_{r-1}, K'_r\},$$

where

$$\begin{aligned} K_1 &= k_1 + k_2 + \dots + k_{r-1} + k_r, \\ K_2 &= k_2 + \dots + k_{r-1} + k_r, \\ &\dots\dots\dots, \\ K_{r-1} &= k_{r-1} + k_r, \\ K'_r &= k_r - 1. \end{aligned}$$

But now steps in our theorem (3), and shows that the system (θ) may be superseded by another, in which the variables, instead of being $a_1, a_2 \dots a_n$, will be

$$a_1 - \frac{\alpha_1}{a_n}a_n, \quad a_2 - \frac{\alpha_2}{a_n}a_n, \dots a_{n-1} - \frac{\alpha_{n-1}}{a_n}a_n;$$

consequently the number of really independent variables is only $(n-1)$; we must therefore have

$$(k_1, k_2 \dots k_r) = 1 + \{K_1, K_2 \dots K'_r\}.$$

p. 180

Since the introduction of a new simple equation is equivalent to the requirement of one more disposable letter, we may write the above more symmetrically under the form

$$(k_1, k_2 \dots k_r) = (K_1, K_2 \dots K_{r-1}, K'_r),$$

where

$$K_1 = 1 + k_1 + k_2 + \dots + k_r, \quad K'_r = k_r - 1.$$

By means of this formula of reduction $(k_1, k_2 \dots k_r)$ may be finally brought down to the form (L) , and the value of (L) being the number of letters required for the linear solution of a system of L linear equations is evidently $L + 2$.

Thus, to determine the number of letters required for the linear solution of a single quadratic, we write

$$(0, 1) = (2) = 4.$$

For two quadratics, we write

$$(0, 2) = (3, 1) = (5) = 7;$$

for a quadratic and a cubic,

$$(0, 1, 1) = (3, 2) = (6, 1) = (8) = 10;$$

for two cubics,

$$(0, 0, 2) = (3, 2, 1) = (7, 3) = (11, 2) = (14, 1) = (16) = 18.$$

These results coincide with those obtained by Sir William Hamilton in his Report on Mr Jerrard's Transformation of the Equation of the Fifth Degree in the *Transactions* of the British Association. I have much more to say on the subject of the linear solution of a system of indeterminate equations, and am, I believe, able to present the subject in a more general light than has hitherto been done; but my observations on this matter must be deferred until a subsequent communication.

31.

REPLY TO PROFESSOR BOOLE'S OBSERVATIONS ON A THEOREM CONTAINED IN THE LAST NOVEMBER NUMBER OF THE JOURNAL

[Cambridge and Dublin Mathematical Journal, VI. (1851), pp. 171–174]

p. 181

The restricted space that can be spared for discussion in these pages, necessitates me to compress within the narrowest limit the remarks which I feel bound to make on Mr Boole's extraordinary observations⁹⁶ in the February number of this *Journal*, on my theorem contained in the antecedent number thereof⁹⁷, which statements I cannot, in the interests of truth and honesty, suffer to pass unchallenged. The object of that theorem was to show how the determinant of the quadratic function resulting from the elimination of any set of the variables between a given quadratic function and a number of linear functions of the same variables, could be represented *without performing* the actual elimination by a fraction, of which the numerator would be constant whichever set of the variables might be selected for elimination, and the denominator the square of the determinant corresponding to the coefficients of the variables so eliminated. The numerator itself is a determinant, obtained by forming the square corresponding to the determinant of the given quadratic function, and bordering it horizontally and vertically with the lines and columns corresponding to the coefficients of all the variables in the given linear equations. An *immediate corollary* from this theorem leads to Mr Boole's. Conversely upon the principle that "tout est dans tout" Mr Boole devotes a page and a half of close print merely to indicate the steps of a method by which from his theorem mine is capable of being deduced, ending with the announcement, that the numerator in question is equal to the quantity

$$\phi_1 \phi_2 \cdots \phi_r \theta(Q),$$

(the symbols above employed being Mr Boole's own), and concludes with assuring his readers that "he has ascertained that Mr Sylvester's result is reducible to the above form." Mr Sylvester would be very sorry to put his

p. 182

result under any such form. Mr Boole could scarcely have reflected upon the effect of his words when he indulged in the remark which follows—"there cannot be a doubt that for the discovery of the actual relation in question, the above theorem is far more convenient than Mr Sylvester's." Of the value to be attached to this assertion the annexed comparison of results is submitted as a specimen.

Let the quadratic function be

$$ax^2 + by^2 + cz^2 + dt^2 + 2exy + 2\epsilon zt + 2gxz + 2\gamma yt + 2hyz + 2\eta xt,$$

⁹⁶*Cambr. and Dublin Math. Jour.* VI. (1851), pp. 90, 284.

⁹⁷p. 135 above.

and the linear functions (taken two in number)

$$lx + my + nz + pt, \quad l'x + m'y + n'z + p't.$$

My numerator will be the determinant (hereinafter cited as the *extended determinant*),

$$\begin{vmatrix} a & e & g & \eta & l & l' \\ e & b & h & \gamma & m & m' \\ g & h & c & \varepsilon & n & n' \\ \eta & \gamma & \varepsilon & d & p & p' \\ l & m & n & p & 0 & 0 \\ l' & m' & n' & p' & 0 & 0 \end{vmatrix}.$$

To find the numerator of Mr Boole's fraction, we must form the symbolical operator

$$\begin{aligned} & \left(l^2 \frac{d}{da} + m^2 \frac{d}{db} + n^2 \frac{d}{dc} + p^2 \frac{d}{dd} + 2lm \frac{d}{de} + 2np \frac{d}{d\varepsilon} + 2ln \frac{d}{dg} \right. \\ & \quad \left. + 2mp \frac{d}{d\gamma} + 2lp \frac{d}{dh} + 2mn \frac{d}{d\eta} \right) \\ & \times \left(l'^2 \frac{d}{da} + m'^2 \frac{d}{db} + n'^2 \frac{d}{dc} + p'^2 \frac{d}{dd} + 2l'm' \frac{d}{de} + 2n'p' \frac{d}{d\varepsilon} + 2l'n' \frac{d}{dg} \right. \\ & \quad \left. + 2m'p' \frac{d}{d\gamma} + 2l'p' \frac{d}{dh} + 2m'n' \frac{d}{d\eta} \right), \end{aligned}$$

and after expanding the determinant hereunder written

$$\begin{vmatrix} a & e & g & \eta \\ e & b & h & \gamma \\ g & h & c & \varepsilon \\ \eta & \gamma & \varepsilon & d \end{vmatrix},$$

perform the operations above indicated upon the result so obtained.

These are the operations and processes which, on Professor Boole's authority, we are to accept "as without doubt far more convenient" than the one simple process of forming, and when necessary, calculating the

p. 183

extended determinant above given. Here for the present I leave the case between Mr Boole and myself to the judgment of the readers of this *Journal*.

In the April Number of the *Philosophical Magazine*⁹⁸, I have shown that the extended determinant serves, not only to represent the full and complete determinant of the reduced quadratic function, but likewise all the minor determinants thereof; the last set of which will be evidently no other than the coefficients themselves. For instance, in the example above given, if we wish to find the

⁹⁸p. 241 below.

coefficient of x^2 after z and t have been eliminated, we have only to strike out the line and column $e b h \gamma m m'$ from the extended determinant; if we wish to find the coefficient of y^2 , we must strike out the line and column $a e g \eta l l'$; to find the coefficient of xy , we must strike out the line $a e g \eta l l'$ and the column $e b h \gamma m m'$, or *vice versa*.

In each of these cases the determinant so obtained is the numerator of the equivalent fraction; the denominator remaining always the same function of the coefficients of transformation as in the original theorem.

Again, if there be taken only one linear equation, and by aid of it x is supposed to be eliminated; and if the reduced quadratic function be called

$$Ly^2 + Mz^2 + Nt^2 + 2Pzt + 2Qyt + 2Rzy,$$

the same extended determinant as before given will serve, when stripped of its outer border, consisting of the line and column $l' m' n' p'$, to produce the various equivalent fractions: thus form the square

$$\begin{array}{ccc} L & R & Q \\ R & M & P \\ Q & P & N \end{array}.$$

The numerator of the fraction equivalent to

$$\begin{vmatrix} L & R \\ R & M \end{vmatrix}, \quad \text{that is, to } LM - R^2,$$

may be found by striking out from the form of the extended determinant the line and column $\eta \gamma \varepsilon d p$; that corresponding to

$$\begin{vmatrix} L & Q \\ R & P \end{vmatrix}, \quad \text{that is, } LP - RQ,$$

will be found by striking out the line $g h c \varepsilon n$ and the column $\eta \gamma \varepsilon d p$, or *vice versa*; and so forth for all the first minor determinants; and similarly the second minors, that is, L, M, N, P, Q, R , may be obtained by striking out in each case a correspondent pair of lines and pair of columns. Thus, to find the numerator of L the same pair of lines and columns, namely, $(g h c \varepsilon n)$, $(\eta \gamma \varepsilon d p)$, must be elided. To find the numerator of R , the pair of lines $(g h c \varepsilon n)$, $(\eta \gamma \varepsilon d p)$, and the pair of columns $(e b h \gamma m)$, $(\eta \gamma \varepsilon d p)$, or *vice versa*, will have to be elided; and so forth for the remaining second minors. I may conclude with observing, that the theorem contested by Mr Boole is an immediate corollary from the general Theory of Relative Determinants alluded⁹⁹ to in the “Sketch” inserted in the present number of the *Journal*.

⁹⁹p. 188 below.

32.

SKETCH OF A MEMOIR ON ELIMINATION, TRANSFORMATION, AND CANONICAL FORMS

[Cambridge and Dublin Mathematical Journal, VI. (1851), pp. 186–200]

p. 184

There exists a peculiar system of analytical logic, founded upon the properties of zero, whereby, from dependencies of equations, transition may be made to the relations of functional forms, and *vice versâ*: this I call the logic of characteristics.

The resultant of a given system of homogeneous equations of as many variables, is the function whose nullity implies and is implied by the possibility of their coexistence, that is, is the characteristic of such possibility; but inasmuch as any numerical product of any power of a characteristic is itself an equivalent characteristic, in order to give definiteness to the notion of a resultant, it must further be restricted to signify the characteristic taken in the *lowest form* of which it *in general* admits.

The following very general and important proposition for the change of the independent variables in the process of elimination, is an immediate consequence of the doctrine of characteristics.

Let there be two sets of homogeneous forms of function;

the 1st,

$$\phi_1, \quad \phi_2 \dots \phi_n,$$

the 2nd,

$$\psi_1, \quad \psi_2 \dots \psi_n.$$

Let the results of applying these forms to any sets of n variables be called

$$(\phi_1), \quad (\phi_2) \dots (\phi_n),$$

$$(\psi_1), \quad (\psi_2) \dots (\psi_n);$$

then will the resultant (in respect to those variables) of

$$\phi_1 \{(\psi_1), (\psi_2) \dots (\psi_n)\},$$

$$\phi_2 \{(\psi_1), (\psi_2) \dots (\psi_n)\},$$

.....

$$\phi_n \{(\psi_1), (\psi_2) \dots (\psi_n)\},$$

be the product of powers (assignable by the law of homogeneity) of the separate resultants of the two systems,

p. 185

$$\{(\phi_1), (\phi_2) \dots (\phi_n)\}, \quad \{(\psi_1), (\psi_2) \dots (\psi_n)\}.$$

By means of the doctrine of characteristics the following general problem may be resolved.

Given any number of functions of as many letters, and an inferior number of functions of the same inferior number of letters, obtained by combining, *inter se*, in a known manner, the given functions, to determine the factor by which, the resultant of the reduced system being divided, the resultant of the original system may be obtained.

If in the theorem for the change of the independent variables both sets of forms of functions be taken linear, we obtain the common rule for the multiplication of determinants: if we take one set linear and the other not, we deduce two rules, namely, That the resultant of a given set of functional forms of a given set of variables, enters as a factor into the resultant,

1st, of linear functions of the given functions of the given variables;

2nd, of the given functions of linear functions of the given variables:

the extraneous factor in each case being a power of what may be conveniently termed the *modulus of transformation*, that is, the resultant of the imported linear forms of functions.

From the second of these rules we obtain the law first stated I believe for functions beyond the second degree by Mr Boole, to wit, that the determinant of any homogeneous algebraical function (meaning thereby the resultant of its first partial differential coefficients) is unaltered by any linear transformations of the variables, except so far as regards the introduction of a power of the modulus of transformation. This is also abundantly apparent from the fact, that the nullity of such determinant implies an immutable, that is, a fixed and inherent, property of a certain corresponding geometrical locus.

There exist (as is now well known) other functions besides the determinant, called by their discoverer (Mr Cayley) hyperdeterminants, gifted with a similar property of immutability. I have discovered a process for finding hyperdeterminants of functions of any degree of any number of letters, by means of a process of Compound Permutation. All Mr Cayley's forms for functions of two letters may be obtained in this manner by the aid of one of the two processes (to wit, that one which will hereafter be called the derivational process), for passing from immutable constants to immutable forms. Such constants and forms, derived from given forms, may be best

p. 186

termed adjunctive; a term slightly varied from that employed by M. Hermite in a more restricted sense.

The two processes alluded to may be termed respectively appositional and derivational. The appositional is founded upon the properties of the binary function $x\xi + y\eta + z\zeta + \dots$; in which, whether we substitute linear functions of $x, y, z, \&c.$, or linear functions of $\xi, \eta, \zeta, \&c.$, in place of $x, y, z, \&c.$, or $\xi, \eta, \zeta, \&c.$, the result is the same.

Consequently, if we apply the form ϕ to $\xi, \eta \dots \zeta$, and take any constant (in respect to $\xi, \eta \dots \zeta$) adjunctive to

$$\phi(\xi, \eta \dots \zeta) + (x\xi + y\eta + \dots + z\zeta + kt)^{n-1},$$

calling this quantity $\psi(x, y \dots z, t)$, the form ψ is evidently adjunctive to the form ϕ : and if we expand so as to obtain

$$\psi(x, y \dots z, t) = \psi_1(x, y \dots z)t^a + \psi_2(x, y \dots z)t^\beta + \&c.,$$

it is evident $\psi_1, \psi_2, \&c.$ will be each separately adjunctive to ϕ . These forms, when ψ is obtained by finding the determinant in respect to $\xi, \eta \dots \zeta$ of S , are, in fact, identical with Hermite's "formes adjointes."

The derivational mode of generating forms from constants depends upon the property of the operative symbol

$$\chi = \xi \frac{d}{dx} + \eta \frac{d}{dy} + \dots + \zeta \frac{d}{dz},$$

applied to ϕ a function of $x, y \dots z$; namely, that if in ϕ , in place of these letters, we write linear functions thereof, to wit $x', y' \dots z'$, we may write

$$\chi = \xi' \frac{d}{dx'} + \eta' \frac{d}{dy'} + \dots + \zeta' \frac{d}{dz'},$$

where $\xi', \eta' \dots \zeta'$ will be the same functions of $\xi, \eta \dots \zeta$ that $x', y' \dots z'$ are of $x, y \dots z$.

Suppose now, in the first place, that in regard to $\xi, \eta \dots \zeta$, $\psi(x, y \dots z)$ is adjunctive to $\chi^r \phi(x, y \dots z)$; then is the form ψ adjunctive to the form ϕ , for on changing $x, y \dots z$ to $x', y' \dots z'$,

$$\left(\xi \frac{d}{dx} + \eta \frac{d}{dy} + \dots + \zeta \frac{d}{dz} \right)^r \phi(x, y \dots z)$$

becomes

$$\left(\xi' \frac{d}{dx'} + \eta' \frac{d}{dy'} + \dots + \zeta' \frac{d}{dz'} \right) \phi(x', y' \dots z');$$

and consequently $\psi(x, y \dots z)$ becomes $\psi(x', y' \dots z')$, multiplied by a power of the modulus of transformation, the modulus of that transformation, be it well observed, whereby $x', y' \dots z'$ would be replaced by $x, y \dots z$, and not as in the appositional mode of that converse transformation according to which

p. 187

$x, y \dots z$ would be replaced by $x', y' \dots z'$. It is on account of this converseness of the modes of transformation that the appositional and derivational modes of generating forms cannot except for a certain class of *restricted* linear transformations be combined in a single process. More generally, if instead of a single function $\chi^r \phi(x, y \dots z)$, we take as many such with different indices to χ as there

are variables, and form either the resultant in respect to $\xi, \eta \dots \zeta$, or any other immutable constant in regard to those variables, (presuming in extension of the hyperdeterminant theory and as no doubt is the case, that such exist), every such resultant or other constant will give a form of function of $x, y \dots z$ adjunctive to the given form ϕ .

It may be shown that every such resultant so formed will contain ϕ as a factor.

Again, in the former more available determinant mode of generation, if we take the determinant in respect to $\xi, \eta \dots \zeta$, it may be shown that all the adjunctive functions so obtained will be algebraical derivees of the partial differential coefficients of ϕ in respect to $x, y \dots z$; that is to say, if these be respectively zero, all such adjunctive functions so derived, as last aforesaid, will be zero, or in other words, each such adjunctive is a syzygetic function of the partial differential coefficients of the primitive function.

To Mr Boole is due the high praise of discovering and announcing, under a somewhat different and more qualified form and mode of statement, this marvel-working process of derivational generation of adjunctive forms. I was led back to it, in ignorance of what Mr Boole had done, by the necessity which I felt to exist of combining Hesse's so-called functional determinant, under a common point of view with the common constant determinant of a function; under pressure of which sense of necessity, it was not long before I perceived that they formed the two ends of a chain of which Hesse's end exists for all homogeneous functions, but the other only when such functions are algebraical.

In fact, if we give to r every value from 2 upwards, the successive determinants in respect to $\xi, \eta \dots \zeta$ of

$$\left(\xi \frac{d}{dx} + \eta \frac{d}{dy} + \dots + \zeta \frac{d}{dz} \right)^r \phi(x, y, z),$$

will produce the chain in question, which, when ϕ is algebraical and of n dimensions, comes to a natural termination when $r = n - 1$. The last member of and the number of terms in this chain are identical with the last member of and the number of terms in Sturm's auxiliary functions, when the variables are reduced to two. There is some reason to anticipate that this chain of functions may be made available in superseding Sturm's chain of auxiliaries; and if so, then the fatal hindrance to progress, arising from the unsymmetrical nature of the latter, is overcome, and we shall be

p. 188

able to pass from Sturm's theorem, which relates to the theory of Kenothemes, or Point-systems, to certain corresponding but much higher theories for lines, surfaces, and n -themes generally.

The restriction of space allowed to me in the present number of the *Journal* will permit me only to allude in the briefest terms to the theory of Relative Determinants, which, as it will be seen, plays an important part in the effectuation of the reductions of the higher algebraical functions to their simplest forms. Nor

can the effect of the processes to be indicated be correctly appreciated without a knowledge of the circumstances under which the resultant of a *given* system of equations can sink in degree below the resultant of the *general type* of such system. Abstracting from the case when the equations separately, or in combination, subdivide into factors, this lowering of degree, as may be shown by the doctrine of characteristics, can only happen in one of two ways. Either the particular resultant obtained is a rational root of the general resultant, or the general resultant becomes zero for the case supposed, and the particular resultant is of a distinct character from the general resultant, being in fact the characteristic of the possibility not of the given system of equations being merely able to coexist (for that is already supposed), but of their being able to coexist for a certain system of values *other than* a given system or given systems. Such a resultant may be termed a Sub-resultant; the lowest resultant in the former case may be termed a Reduced-resultant. The theory of Sub-resultants is one altogether remaining to be constructed, and is well worthy equally of the attention of geometers and of analysts.

As to the theory of Relative Determinants, the object of this theory is to obtain the determinant resulting from eliminating as many variables as can be eliminated, chosen at pleasure from a set of variables greater in number than the equations containing them; and the mode of effecting this object is through the method of the indeterminate multiplier. To avoid the discussion of the theory of sub-resultants and other particularities, I shall content myself with giving the rule applicable to the case (the only one of which as yet a practical application has offered itself to me in the course of my present inquiries) when all but one of the functions are linear.

If $U, L_1, L_2 \dots L_m$ be the first an n^c and the others linear functions of n variables, and it be desired to find the determinant of the resultant arising from the elimination of any m out of the n variables, the following is the rule:

Find the determinant, that is, the resultant of the partial differential coefficients in respect to the given variables, and of $\lambda_1, \lambda_2 \dots \lambda_m$ of

$$U + L_1\lambda_1 + L_2\lambda_2 + \dots + L_m\lambda_m.$$

p. 189

This resultant, in its lowest form, will be always a rational $(n - 1)$ th root of the resultant of the homogeneous system of equations to which the system above given can be referred as its type; and this reduced resultant divided by a power (determinable by the law of homogeneity) of the resultant of $L_1, L_2 \dots L_m$, when all but the selected variables are made zero, will be the resultant determinant required¹⁰⁰. As regards what has been said concerning the reducibility of the general typical resultant in the case before us, this is a consequence of, and may be

¹⁰⁰The same method applies not only to the Final or Constant Determinant, but likewise to all the Functional Determinants in the chain above described, extending upwards from this to the Hessian, or as it ought to be termed, the first Boolean Determinant.

brought into connexion with, the following theorem, which is easily demonstrable by the theory of characteristics. If $Q_1, Q_2 \dots Q_m$ be m homogeneous functions of m variables of the same degree, r of which enter in each equation only as simple powers uncombined with any of the other variables, then the degree of the reduced resultant is equal to the number of the equations multiplied by the $(m - r - 1)$ th power of the number of units in the degree of each, subject to the obvious exception that when r is m , (there being in fact but *one* step from $r = m - 2$ to $r = m$), instead of r , $(r - 1)$ must be employed in the above formula. As an example of a sub-resultant as distinguished from a reduced-resultant, I instance the case of three quadratics U, V, W , functions of x, y, z , in each of which no squared power of z is supposed to enter: it may easily be shown by my dialytic method that instead of six equations, between which to eliminate $x^2, y^2, z^2, xy, xz, yz$, we shall have only 5, the three original ones and two instead of three auxiliaries between which to eliminate x^2, y^2, xy, xz, yz , the *apparent* resultant is accordingly of the 9th instead of the 12th degree. But this is not the true characteristic of the possibility of the coexistence of the given systems, which in fact is zero, as is evidenced by the fact that they always *do* coexist, since they are always satisfiable by only *two* relations between the variables, to wit $x = 0, y = 0$. The apparent resultant is then something different, and what has been termed by the above a Sub-resultant.

I take this opportunity of entering my simple protest against the appropriation of my method of finding the resultant of any set of three equations of degrees equal or differing only by a unit, one from those of the other two, by Dr Hesse, so far as regards quadratic functions, without acknowledgment, four years after the publication of my memoir in the *Philosophical Magazine*: the fundamental idea of Dr Hesse's partial method is identical with that of my general one. Still more unjustifiable is the subsequent use of the *dialytic* principle, by the same author, equally without acknowledgment, and in cases where there is no peculiarity of form of procedure to give even a plausible ground for evading such acknowledgment. It is capable of moral proof that

p. 190

what I had written on the matter was sufficiently known in Berlin and at Königsberg, at each epoch of Dr Hesse's use of the method.

I now proceed to the consideration of the more peculiar branch of my inquiry, which is as to the mode of reducing Algebraical Functions to their simplest and most symmetrical, or as my admirable friend M. Hermite well proposes to call them, their Canonical forms. Every quadratic function of any number of variables may always be linearly transformed into any other quadratic functions of the same, and that too in an infinite variety of ways; but in every other instance there will be only a limited number of ways, whereby, when possible, one form will admit of being transmuted into any other: and with the sole exception of a cubic function of two letters, such transmutation will never be possible, unless a certain condition, or certain conditions, be satisfied between

the constants of the forms proposed for transmutation. The number of such conditions is the number of parameters entering into the canonical form, and is of course equal to the number of terms in the general form of the function diminished by the square of the number of letters. Thus there is one parameter in the canonical form for the biquadratic function of two and the cubic function of three letters, and no parameter in the cubic function of two letters. Hitherto no canonical forms have been studied beyond the cases above cited, but I have succeeded, as will presently be shown, in obtaining methods for reducing to their canonical forms functions with *two* and *four* parameters respectively. Owing to what has been remarked above, the theory of quadratic functions is a theory apart. Simultaneous transformation gives definiteness to that theory, but has no existence for any useful purpose for functions of the higher degrees. Where the theory of simultaneous transformation ends, that of canonical forms properly begins; and in what follows, the case of quadratic forms is to be understood as entirely excluded. Such exclusion being understood, there is no difficulty in assigning the canonical, that is, the simplest and most symmetrical general, form to which every function of two letters admits of being reduced by linear transformations. If the degree be odd, say $2m + 1$, the canonical form will be

$$u_1^{2m+1} + u_2^{2m+1} + \dots + u_{m+1}^{2m+1};$$

if the degree be even, say $2m$, the canonical form will be

$$u_1^{2m} + u_2^{2m} + \dots + u_m^{2m} + K(u_1 u_2 \dots u_m)^2,$$

all the u 's being linear functions of the two given variables. It is easy to extend an analogous mode of representation to functions of any number of letters. From the above we see that for cubic, biquadratic, and quintic functions of two letters, the canonical forms will be respectively

$$u^3 + v^3, \quad u^4 + v^4 + K u^2 v^2, \quad u^5 + v^5 + w^5,$$

with a linear relation in the last-named case between u, v, w .

p. 191

First as to the reduction of any 4^c function to Cayley's form

$$u^4 + v^4 + K u^2 v^2.$$

This may be effected in a great variety of ways, of which the following is not the simplest as regards the calculations required, but the most obvious. Let the modulus of transformation, whereby the given biquadratic function, say $F(x, y)$, becomes transmuted into its canonical form, be called M ; let the determinant of F be called D_1 , and the determinant of the determinant in respect to ξ and η of

$$\left(\xi \frac{d}{dx} + \eta \frac{d}{dy} \right)^2 F(x, y),$$

which latter, for brevity's sake, may be termed the Hessian of F , (although in stricter justice the Boolean would be the more proper designation), be called D_2 . Then, by examining the canonical form itself (which is as it were the very *palpitating heart* of the function laid bare to inspection), we shall obtain without difficulty the two equations

$$(1 - 9m^2)^2 = M^{12} D_1 \frac{1}{4^6},$$

$$m^2(1 - 9m^2)^2(m^2 - 1)^2 = M^{24} D_2 \frac{1}{12^2 4^4}.$$

Eliminating the unknown quantity M , we obtain

$$\frac{m^2(m^2 - 1)^2}{(1 - 9m^2)^2} = c, \quad \text{or} \quad \frac{m^3 - m}{1 - 9m^2} = c^{1/2},$$

where c is a known quantity.

This cubic equation for finding m is of a peculiar form; it being easy to show *à priori*, by going back to the canonical form, that its three roots are $m, \theta(m), \theta^2(m)$, where

$$\theta(m) = \frac{m - 1}{3m + 1},$$

θ being a periodical form of function such that $\theta^3(m) = m$.

This it is which accounts for the simple expression for m , that may be obtained by solving the cubic above given. A better practical mode is to take, instead of the determinant of the given function and its Hessian, the two hyperdeterminants and eliminate as before: a cubic equation having precisely the same properties, and in fact virtually identical with the former, will result. When m and consequently M are found, there is no difficulty whatever, calling the given function F and its Hessian $H(F)$, in forming linear functions of the two, as

$$\left. \begin{array}{l} \phi(m)F + \psi(m)H(F) \\ \phi_1(m)F + \psi_1(m)H(F) \end{array} \right\}$$

which shall be equal to, that is, identical with, $(u^2 + v^2)^2$ and u^2v^2 , whence u and v are completely determined.

p. 192

Another and interesting mode of solution is to take, besides the given function F and its Hessian, either the *second* Hessian or the post-Hessian of the given function, by the post-Hessian understanding the determinant in respect of ξ and η of

$$\left(\xi \frac{d}{dx} + \eta \frac{d}{dy} \right)^3 F :$$

any three of the four functions will be linearly related, and it may be shown that, calling either the second Hessian (that is, the Hessian of the Hessian) or

the post-Hessian H' , we shall have

$$H'(F) + aH(F) + bF = 0,$$

where a and b will be *rational* and *integer* functions of the coefficients of F , and numerical multiples of two quantities R and S , such that the determinant of F will be equal $R^3 + S^2$; and this, be it observed, without any previous knowledge of the existence of these hyperdeterminants R and S .

If now we go to Hesse's form for a cubic function of three letters, we shall find that precisely similar modes of investigation apply step for step. Calling the function F and its Hessian $H(F)$, and the post-Hessian or second Hessian at choice $H'(F)$, we shall find

$$H'(F) + mSH(F) + nR^2F = 0,$$

where m and n are numerical quantities and $R^3 + S^2$ equal the determinant of F . It is interesting to contrast this equation with the one previously mentioned as applicable to the 4^c functions of two letters, namely,

$$H'(F) + mRH(F) + nSF = 0.$$

In both instances there is no difficulty in assigning the relations between the original R and S , and the R and S of any adjunctive form. All Aronhold's results may be thus obtained and further extended without the slightest difficulty. As regards the equation for finding the parameter in Hesse's canonical form for the cubic of three letters, this will be of the 4th degree in respect to the cube of the parameter, and the roots will be functionally representable as

$$x; \quad \theta(x); \quad \phi(x); \quad \psi(x),$$

where

$$\theta^2(x) = \phi^2(x) = \psi^2(x) = x;$$

$$\theta\phi(x) = \phi\theta(x) = \psi(x),$$

$$\phi\psi(x) = \psi\phi(x) = \theta(x),$$

$$\psi\theta(x) = \theta\psi(x) = \phi(x);$$

owing to which property the equation is soluble under the peculiar form observed by Aronhold.

p. 193

I pass on now to a brief account of the method, or rather of a method (for I doubt not of being able to discover others more practical), of reducing a function of the 5th degree of two letters (say of x and y) to its canonical form $u^5 + v^5 + w^5$, subject to the linear relation $au + bv + cw = 0$, where the ratios $a : b : c$, and the linear relations between u, v, w and the two given variables are the objects of research. Here I have found great aid from the method of Relative Determinants;

and I may notice that the successful application of more compendious methods to the question would be greatly facilitated were there in existence a theory of Relative Hyperdeterminants, which is still all to form, but which I little doubt, with the blessing of God, to be able to accomplish. It may some little facilitate the comprehension of what follows, if c be considered as representing unity.

Calling as before the given quintic function F , the modulus of transformation M , the Hessian and post-Hessian of F , H and H' , and its ordinary or constant determinant D , we shall find

$$a^2v^3w^3 + b^2w^3u^3 + c^2u^3v^3 = M^2H,$$

and

$$P_1P_2P_3P_4 = M^6H',$$

where

$$P_1 = a^{3/2}vw + b^{3/2}wu + c^{3/2}uv,$$

$$P_2 = a^{3/2}vw - b^{3/2}wu - c^{3/2}uv,$$

$$P_3 = -a^{3/2}vw + b^{3/2}wu - c^{3/2}uv,$$

$$P_4 = -a^{3/2}vw - b^{3/2}wu + c^{3/2}uv;$$

also $D = M^{20}$ multiplied by the product of the sixteen values of

$$a^{5/4} + b^{5/4}(1)^{1/4} + c^{5/4}(1)^{1/4}.$$

From the above equations it may be shown that H' (a known function of the 8th degree of the given variables x, y) must be capable of being thrown under the form

$$L\{(x - a_1y)(x - a_2y) \times (x - a_3y)(x - a_4y) \\ \times (x - a_5y)(x - a_6y) \times (x - a_7y)(x - a_8y)\},$$

where

$$(a_1 - a_2)^2 \times (a_3 - a_4)^2 \times (a_5 - a_6)^2 \times (a_7 - a_8)^2 = \frac{D}{L^2} = K,$$

so that K is a known quantity¹⁰¹. Accordingly the said equation of the 8th degree, considered as an algebraical equation in $\frac{x}{y}$, may by known methods be p. 194

found by means of equations not exceeding the 4th or even the 3rd degree: in fact, to do this it is only necessary to form the equation to the squares of the differences of the roots of $\frac{x}{y}$ in the equation $H' \div y^8 = 0$, which new equation will be of the 28th degree. If we then form two other equations of the 378th degree, one having its roots equal to $\sqrt{K}$ multiplied by the binary products of

¹⁰¹Or in other words, the post-Hessian determinant of a given function in two letters of the second degree, may be divided into four quadratic factors in such a way that the product of the determinants of these several factors shall be equal to the determinant of the given function.

the twenty-eight roots of the equation last named, the other to $\sqrt{K}$ multiplied by the reciprocal of such binary products, the left-hand members of these two equations expressed under the usual form will have a factor in common, which may be found by the process of common measure and will be of the 6th degree, whose roots consisting of three pairs of reciprocals may be found by the solution of cubics only.

In this way, by means of cubics and quadratics,

$$(a_1 - a_2)^2, \quad (a_3 - a_4)^2, \quad (a_5 - a_6)^2, \quad (a_7 - a_8)^2,$$

can be found, which being known,

$$a_1a_2, \quad a_3a_4, \quad a_5a_6, \quad a_7a_8,$$

can be determined in pairs by means of quadratics from the equation $H' \div y^8 = 0$. This being supposed to be done, we have

$$P_1 = fL_1, \quad P_2 = gL_2, \quad P_3 = hL_3, \quad P_4 = kL_4,$$

where L_1, L_2, L_3, L_4 are *known* quadratic functions of x and y . To determine the ratios of f, g, h, k , we have three equations¹⁰² obtained from the identity

$$fL_1 + gL_2 + hL_3 + kL_4 = (P_1 + P_2 + P_3 + P_4) = 0;$$

$f : g : h : k$ being known, $fL_1 : gL_2 : hL_3 : kL_4$ are known ratios.

But

$$P_1 + P_2 = 2a^{3/2}vw, \quad P_1 + P_3 = 2b^{3/2}wu, \quad P_1 + P_4 = 2c^{3/2}uv.$$

Hence

$$a^{3/2}vw = \lambda P, \quad b^{3/2}wu = \lambda Q, \quad c^{3/2}uv = \lambda R,$$

where P, Q, R are known quadratic functions of x, y .

p. 195

Hence $a : b : c$ may be found by means of the identical equation

$$a^2w^3v^3 + b^2u^3w^3 + c^2v^3u^3 = H(F),$$

whereby the ratios $a^{-5/2} : b^{-5/2} : c^{-5/2}$ can be obtained without any further extraction of roots, showing that there is but one single true system of ratios $a^5 : b^5 : c^5$ applicable to the problem; $a : b : c$ being thus found, λ is easily determined, and thus finally u, v, w are found in terms of x and y ¹⁰³.

¹⁰²For we must have the coefficients of x^2, xy and y^2 in $fL_1 + gL_2 + hL_3 + kL_4$, of all them zero.

¹⁰³The problem thus solved may be stated as consisting in reducing the general function $ax^5 + bx^4y + cx^3y^2 + dx^2y^3 + exy^4 + fy^5$ to the form

$$(lx + my)^5 + (l'x + m'y)^5 + (l''x + m''y)^5.$$

I have little doubt that a more expeditious mode of solution than the foregoing¹⁰⁴ will be afforded by an examination of the properties and relations of the *quadratic and cubic forms*, adjunctive to the general quintic functions, and indeed to every $(4m + 1)^c$ function of two letters hereinbefore adverted to.

Sufficient space does not remain for detailing the steps whereby the general cubic function of *four* letters may, by aid of equations *not transcending the fifth degree*, be reduced to its canonical form $u^3 + v^3 + w^3 + p^3 + q^3$, wherein u, v, w, p, q are connected by a linear equation

$$au + bv + cw + dp + eq = 0;$$

the four ratios of whose coefficients $a : b : c : d : e$ give the necessary number

$$\frac{4 \cdot 5 \cdot 6}{1 \cdot 2 \cdot 3} - 4^2$$

parameters furnished by the general rule. Suffice it for the present to say, that the analytical mode of solution depends upon a circumstance capable of the following geometrical statement: Every surface of the 4th degree represented by a function which is the Hessian to any given cubic function whatever of four letters, has lying upon it ten straight lines meeting three and three in ten points, and these ten points are the only points which enjoy the following property in respect to the surface of the 3rd degree denoted by equating to zero the cubical function in question, to wit, that the cone drawn from any one of them as vertex to envelop the surface, will meet it not in a continuous double curve of the 6th degree, but in two curves each of the 3rd degree, lying in *planes* which intersect in the ten lines respectively above named; so that to each of the ten points corresponds one of the ten lines: these ten points and lines are the intersections taken respectively three with three, and two with two, of a *single and unique system* of five principal planes appurtenant to every surface of the 3rd degree, and these planes are no other than those denoted by

$$u = 0, \quad v = 0, \quad w = 0, \quad p = 0, \quad q = 0.$$

p. 196

I have found also by the theory of Sub-resultants, that the analogy between lines and surfaces of the third degree, in regard to the existence of double and conical points, is preserved in this wise: that in the same way as a double point on a curve of the 3rd degree commands the existence of a double point on its Hessian, so does a conical point in a surface of the 3rd degree command over and above the 10 necessary, and so to speak natural conical points, at least one extra, that is to say an 11th conical point on its Hessian. And here for the present I must quit my brief and imperfect notice of this subject, composed amidst the interruptions and distractions of an official and professional life.

¹⁰⁴The coefficients in the reducing recurrent equation of the 6th degree in the process above detailed may rise to be of 541632 dimensions in respect to the original coefficients in F .

Observation. It may be somewhat interesting and instructive to my readers, to have a table of the successive scalar¹⁰⁵ determinants of a quintic function of two letters presented to them at a single glance. Preserving the notation above [page 193], we have the following expressions:

$$\text{The given function} = u^5 + v^5 + w^5,$$

$$\text{its Hessian} = M^2(a^2v^3w^3 + b^2w^3u^3 + c^2u^3v^3),$$

its post-Hessian = $M^6 \times$ the product of the four forms of

$$a^{3/2}vw + b^{3/2}(1)^{1/2}wu + c^{3/2}(1)^{1/2}uv;$$

its praeter-post-Hessian = $M^{12} \times$ the product of the nine forms of

$$a^{4/3}v^{1/3}w^{1/3} + b^{4/3}(1)^{1/3}w^{1/3}u^{1/3} + c^{4/3}(1)^{1/3}u^{1/3}v^{1/3};$$

and the final determinant = $M^{20} \times$ the product of the sixteen forms of

$$a^{5/4} + (1)^{1/4}b^{5/4} + (1)^{1/4}c^{5/4}.$$

The success of the method applied depends (as above shown) upon the fact of a certain function of the roots of the post-Hessian (which is an octavic function of the variables) being known, which fact *hinges* upon the circumstance that

$$(M^6)^2 \times (M^2)^4 = M^{20}.$$

P.S. I have much pleasure in subjoining the cubical hyperdeterminant of the 12th degree function of two letters, worked out upon the principle of Compound Permutation hinted at in the foregoing pages, for which I am indebted to the kindness and skill of my friend Mr Spottiswoode.

p. 197

The function being called

$$ax^{12} + 12bx^{11}y + \frac{12 \cdot 11}{2}cx^{10}y^2 + \&c. \dots + ly^{12},$$

the following is¹⁰⁶ its cubical hyperdeterminant:

$$\begin{aligned} & agm - 6ahl + 15aik + 10aj^2 - 6bfm, \\ & - 24bhk + 30bal + 20bij - 24cfl + 114cgk, \\ & - 145ci^2 + 50chj + 15cem + 20cgi + 20ch^2, \\ & - 400dgj + 280dhi + 20del + 50dfe + 10d^2k, \\ & + 385egi - 135e^2k - 290eh^2 + 705fgh, \\ & - 330f^2i - 50g^3. \end{aligned}$$

¹⁰⁵By which I mean the determinants in respect to ξ, η of

$$\left(\xi \frac{d}{dx} + \eta \frac{d}{dy} \right)^r F(x, y).$$

¹⁰⁶See below, p. 202.

Mr Spottiswoode will I hope publish the work itself in the next number of the *Journal*, in which I shall also show how the hyperdeterminants of the cubical function of three letters, Aronhold's S and T , may be similarly obtained.

33.

ON THE GENERAL THEORY OF ASSOCIATED ALGEBRAICAL FORMS

[*Cambridge and Dublin Mathematical Journal*, VI. (1851), pp. 289–293]

p. 198

The following brief exposition of the general theory of Associated Forms, as far as it has been as yet developed by the labours or genius of mathematicians, is intended as elucidatory and, to a certain extent, emendative of some of the statements in my paper¹⁰⁷ on Linear Transformations, in the preceding number of the *Journal*.

In the first place, let a linear equivalent of any given homogeneous function be understood to mean what the function becomes when linear functions of the variables are substituted in place of the variables themselves, subject to the condition of the modulus of transformation (that is, the value of the determinant formed by the coefficients of transformation) being unity.

Secondly, let two square arrays of terms (the determinants corresponding to each of which are unity) be said to be complementary when each term in the one square is equal to the value of what the determinant represented by the other square becomes when the corresponding term itself is taken unity, but all the other terms in the same line and column with it are taken zero. This relation between the two squares is well known to be reciprocal. Thus, for instance,

$$\begin{vmatrix} a & b & c \\ a' & b' & c' \\ a'' & b'' & c'' \end{vmatrix} \quad \text{and} \quad \begin{vmatrix} \alpha & \beta & \gamma \\ \alpha' & \beta' & \gamma' \\ \alpha'' & \beta'' & \gamma'' \end{vmatrix}$$

will be said to be reciprocally complementary to one another when the two determinants which they represent are each unity, and when we have

p. 199

$$\begin{aligned} a &= \begin{vmatrix} 1 & 0 & 0 \\ 0 & \beta' & \gamma' \\ 0 & \beta'' & \gamma'' \end{vmatrix}, & \alpha &= \begin{vmatrix} 1 & 0 & 0 \\ 0 & b' & c' \\ 0 & b'' & c'' \end{vmatrix}, \\ b &= \begin{vmatrix} 0 & 1 & 0 \\ \alpha' & 0 & \gamma' \\ \alpha'' & 0 & \gamma'' \end{vmatrix}, & \beta &= \begin{vmatrix} 0 & 1 & 0 \\ a' & 0 & c' \\ a'' & 0 & c'' \end{vmatrix}, \\ b' &= \begin{vmatrix} \alpha & 0 & \gamma \\ 0 & 1 & 0 \\ \alpha'' & 0 & \gamma'' \end{vmatrix}, & \beta' &= \begin{vmatrix} a & 0 & c \\ 0 & 1 & 0 \\ a'' & 0 & c'' \end{vmatrix}, \\ & \&c. & \&c. \end{aligned}$$

¹⁰⁷p. 184 above.

Accordingly, two transformations, say of $F(x, y, z)$ and $G(u, v, w)$ respectively, may be said to be concurrent when in F for x, y, z , we write

$$\begin{aligned} & ax + by + cz, \\ & a'x + b'y + c'z, \\ & a''x + b''y + c''z; \end{aligned}$$

and in G for u, v, w , we write

$$\begin{aligned} & au + bv + cw, \\ & a'u + b'v + c'w, \\ & a''u + b''v + c''w; \end{aligned}$$

but complementary when for u, v, w , we write

$$\begin{aligned} & \alpha u + \beta v + \gamma w, \\ & \alpha' u + \beta' v + \gamma' w, \\ & \alpha'' u + \beta'' v + \gamma'' w; \end{aligned}$$

$a, b, c, \&c., \alpha, \beta, \gamma, \&c.$ being related in the manner antecedently explained.

Two forms, each of the same number of variables, may be said to be associate forms when the coefficients of the one are functions of those of the other; and when it happens, that the coefficients of the first are all explicit functions of those of the second, the latter may be termed the originant and the former the derivant.

If now all the linear equivalents of one or of two associated forms are similarly related to corresponding linear equivalents of the other, so that each may be derived from each by the same law, the forms so associated will be said to be concomitant each to the other. This concomitance may be of two kinds, and very probably, in the nature of things, only of the two kinds about to be described. p. 200

The first species of concomitance is defined by the corresponding equivalents of the two associated forms being deduced by precisely similar, or, as we have expressed it, concurrent transformations or substitutions, each from its given primitive. The second species of concomitance is defined by the corresponding equivalents being deduced not by similar but by contrary, that is, reciprocal or complementary substitutions. Concomitants of the first kind may be called covariants; concomitants of the second kind may be called contravariants. When of the two associated forms one is a constant, the distinction between co- and contra-variants disappears, and the constant may be termed an invariant of the form with which it is associated¹⁰⁸. It follows readily from these definitions

¹⁰⁸Accordingly an invariant to a given form may be defined to be such a function of the coefficients of the form, as remains absolutely unaltered when instead of the given form any linear equivalent thereto is substituted. Of course if the determinant of the coefficients of the transformations correspondent to the respective equivalents be not taken unity as supposed in this definition, the effect will be merely to introduce as a multiplier some power of the determinant formed by the coefficients of transformation.

that a covariant of a covariant and a contravariant of a contravariant are each of them covariants; but a covariant of a contravariant and a contravariant of a covariant are each of them contravariants; and also that an invariant, whether of a covariant or of a contravariant, is an invariant of the original function¹⁰⁹.

It will also readily be seen that as regards functions of two letters a contravariant becomes a covariant by the simple interchange of x, y with $-y, x$, respectively. Covariants are Mr Cayley's hyperdeterminants; contravariants include, but are not coincident with, M. Hermite's formes-adjointes, if we understand by the last-named term such forms as may be derived by the process described by M. Hermite in the third of his letters to M. Jacobi, "Sur différents objets de la Théorie des Nombres," (which process is an extension of that employed for determining the polar reciprocal of an algebraical locus¹¹⁰. M. Hermite appears, however, elsewhere to have used

p. 201

the term forme-adjointe in a sense as wide as that in which I employ contravariants. For instance, he has given a most remarkable theorem, which admits of being stated as follows:

If we have a function of any number of letters, say of x, y, z , as

$$ax^m + mbx^{m-1}y + mcx^{m-1}z + \frac{m(m-1)}{2}dx^{m-2}y^2 + \&c.,$$

and if I be any invariant of this function, then will

$$\left(x^m \frac{d}{da} + x^{m-1}y \frac{d}{db} + x^{m-1}z \frac{d}{dc} + x^{m-2}y^2 \frac{d}{dd} + \&c.\right)^r I$$

be a "forme-adjointe" of the given function. It is perfectly true and admits of being very easily proved, as I shall show in your next number, that this is

¹⁰⁹It may likewise be shown that linear equivalents of covariants and contravariants are themselves related to one another as covariants and contravariants respectively, the transformations by which the equivalents are obtained being taken concurrent in the one case and contrary or reciprocal in the other; and of course any algebraic function of any number of covariants is a covariant and of contravariants a contravariant.

¹¹⁰This has been further generalized by me in the theorem given in the last number of this *Journal*, where I have shown in effect that any invariant in respect to $\xi, \eta \dots \theta$ of

$$f(\xi, \eta \dots \theta) + (x\xi + y\eta + \dots + t\theta + \rho)^{n-1},$$

(f being supposed to be of the degree n) is a contravariant of $f(x, y \dots t)$. When this invariant is the determinant of f , it may be shown that we obtain M. Hermite's theorem. It is somewhat remarkable that contravariants should have been in use among mathematicians as well in geometry as the theory of numbers (although their character as such was not recognized) before covariants had ever made their appearance. Invariants of course first came up with the theory of the equation to the squares of the differences of the roots of equations, the last term in such equation being an invariant. I believe that I am correct in saying that covariants first made their appearance in one of Mr Boole's papers, in this *Journal*; but Hesse's brilliant application of one from among the infinite variety of these forms to the discovery of the points of inflexion in a curve of the third order, in other words, to the Canonical Reduction of the Cubic Function of Three Letters, appears to have been the first occasion of their being turned to practical account. [p. 186 above.]

a contravariant of the given function¹¹¹; but it is not (as far as I can see) a forme-adjointe in the sense in which the use of that word is restricted in the letter alluded to. If, however, we adopt as the *definition* of formes-adjointes generally, that property in regard to their transformées which M. Hermite has demonstrated of the particular class treated of by him in the letter alluded to, then his formes-adjointes become coincident with my contravariants. It will thus be seen that covariants and contravariants form two distinct and coextensive species of associated forms, which divide between them the wide and fertile empire of linear transformations so far as its provinces have been as yet laid open by the researches of analysts. In your next number I propose to enter much more largely into the subject generally. More particularly I shall describe the new method of Permutants, including the theory of Intermutants and Commutants (which latter are a species of the former, but embrace Determinants as a particular case), and their application to the theory of Invariants. I shall also exhibit the connexion between the theory of Invariants and that of Symmetrical Functions, and some remarkable theorems on Relative Invariants¹¹².

Some of your readers may like to be informed that a Supplement to my last paper, under the title of “An Essay on Canonical Forms,” has been since published¹¹³; and that I have there given a much simpler method of solution of the problem of the reduction of quintic functions to their canonical form than in the original memoir, and extended the method successfully to the

p. 202

reduction of all odd-degreed functions to their canonical form. I may take this occasion to state that the Lemma given in Note (B) of the Supplement, upon which this method of reduction is based, is an immediate deduction from the well-known theorem for the multiplication of Determinants.

There is a numerical error in “The Cubical Hyperdeterminant of the Twelfth Degree,” worked out after the method of commutants by Mr Spottiswoode, given at the end of my paper in the May Number. The correct result will be stated in the next number of the *Journal*, where I hope also to be able to fix the number of distinct solutions of the problem of reducing a Sextic Function to its canonical form

$$u^6 + v^6 + w^6 + mu^2v^2w^2.$$

For odd-degreed functions there is never more than one solution possible, as shown in the Supplement referred to.

P.S. Since the above was sent to press, I have discovered an uniform mode of solution for the canonical reduction of functions, whether of odd or even degrees. The canonical form however, except for the fourth and eighth degrees, requires

¹¹¹This is also true if *I* be taken any *covariant* instead of an invariant of the function.

¹¹²It will be readily apprehended that the definitions and conceptions above stated, respecting covariants and contravariants of two single functions, may be extended so as to comprehend systems of functions covariantive or contravariantive to one another.

¹¹³By Mr George Bell, University Bookseller, Fleet Street. [p. 203 below.]

to be varied from that assumed in my previous paper. Thus, for the sixth degree the canonical form will be

$$au^6 + bv^6 + cw^6 + muvw(v - w)(w - u)(u - v),$$

where u, v, w are supposed to be connected by the identical equation $u + v + w = 0$. And there will be only *two* solutions—a remarkable and most unexpected discovery. For functions of the eighth degree there are five distinct solutions, and in general there is the strongest reason for believing (indeed it may be positively affirmed) that *when the canonical form has been rightly assumed* for a function of the even degree n , the number of solutions will be $\frac{1}{2}(n + 2)$ when $\frac{1}{2}n$ is even, but $\frac{1}{4}(n + 2)$ when $\frac{1}{2}n$ is odd. It turns out therefore that the theory for functions of the sixth degree is in some respects simpler than for those of the fourth. The investigation into canonical forms here referred to has led me to the discovery of a most unexpected theorem for finding all the invariants of a certain class, belonging to functions of two letters of an even degree.

34.

AN ESSAY ON CANONICAL FORMS, SUPPLEMENT TO A SKETCH OF A MEMOIR ON ELIMINATION, TRANSFORMATION AND CANONICAL FORMS

[George Bell, Fleet Street, 1851]

p. 203

Since the above paper was in print I have succeeded in obtaining a canonical representation of the quadratic and cubic functions adjunctive to the general quintic (5th degreed) functions of two letters.

Let F the quintic function of x, y ,

$$F = u^5 + v^5 + w^5,$$

and

$$au + bv + cw = 0,$$

M being the modulus of the transformation, whereby transition is made from x, y to u, v . Then the quadratic adjunctive is

$$\frac{M^4}{c^4} \{a^4 vw + b^4 wu + c^4 uv\};$$

and the cubic adjunctive is simply

$$\frac{1}{c^3} M^6 (abc)^2 uvw^{114}.$$

Hence we can, in accordance with what I ventured to predict in the preceding sketch, find u, v, w , by means of a simple and practical co-process. To wit, call

$$F = lx^5 + 5mx^4y + 10nx^3y^2 + 10px^2y^3 + 5qxy^4 + ry^5.$$

p. 204

¹¹⁴The knowledge of the existence of these *lower* adjunctive forms is mainly a consequence of Mr Cayley's splendid discovery of hyperdeterminant constants. In fact, they are respectively the quadratic and cubic hyperdeterminants in respect to ξ and η of

$$\frac{1}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5} \left(\xi \frac{d}{dx} + \eta \frac{d}{dy} \right)^4 F,$$

x and y being treated as constants.

The fortunate proclaimer of a new outlying planet has been justly rewarded by the offer of a baronetcy and a national pension, which the writer of this wishes him long life and health to enjoy. In the meanwhile, what has been done in honour of the discoverer of a new and inexhaustible region of exquisite analysis?

¹¹⁴p. 184 above. See p. 201, note ‡.

Form the determinant

$$\begin{vmatrix} lx + my & mx + ny & nx + py \\ mx + ny & nx + py & px + qy \\ nx + py & px + qy & qx + ry \end{vmatrix}.$$

Let this cubic function, by solving it as a cubic equation, be made equal to

$$L(x + fy)(x + gy)(x + hy),$$

then

$$u = k(x + fy), \quad v = l(x + gy), \quad w = m(x + hy).$$

By means of the identity, $F = u^5 + v^5 + w^5$, k^5, l^5, m^5 , are known by the solution of linear equations, and thus u, v, w , are determined by solving a cubic equation instead of one of the eighth degree, as in the method first given, and the process of canonising a quintic function is rendered *practically* possible.

For brevity sake let c represent unity. The constant determinant of the cubic adjunctive will be found to be

$$3M^{30}(abc)^{10}.$$

Calling, then, the cubic adjunctive of F , $C(F)$, we have the remarkable equation

$$uvw = \frac{C(F)}{\sqrt[5]{\left\{\frac{1}{3} \square C(F)\right\}}}.$$

It may also be shown that if we call the Hessian of F , $H(F)$, we shall have the following equally remarkable equation:

$$\square H(F) = \frac{1}{3} \square F \times \square C(F).$$

Again, calling the quadratic adjunctive of F , $Q(F)$, we shall easily find

$$\square Q(F) = M^{10} \left\{ \begin{array}{l} a^{5/2} + b^{5/2} + c^{5/2} \\ a^{5/2} + b^{5/2} - c^{5/2} \\ a^{5/2} - b^{5/2} + c^{5/2} \\ a^{5/2} - b^{5/2} - c^{5/2} \end{array} \right\},$$

or, if we please,

$$= M^{10} \{a^{10} + b^{10} + c^{10} - 2a^5b^5 - 2a^5c^5 - 2b^5c^5\}.$$

When u, v, w are known, a, b, c , which are the resultants of v, w ; w, u ; u, v respectively are known. But their ratios, or, if we please to say so, the ratios of $a^5 : b^5 : c^5$, may be found independently and very elegantly as follows:—

Let

$$\begin{aligned} M^{10} &\times \text{ product of the 4 forms of } a^{5/2} + 1^{1/2}b^{5/2} + 1^{1/2}c^{5/2} = A, \\ M^{20} &\times \text{ product of the 16 forms of } a^{5/4} + 1^{1/4}b^{5/4} + 1^{1/4}c^{5/4} = B, \\ M^{30} &\times a^{10} \cdot b^{10} \cdot c^{10} = C. \end{aligned}$$

A, B, C are known quantities, being respectively what we have called $\square Q(F)$, $\square(F)^*$, $\frac{1}{3}\square C(F)$. p. 205

It may easily be shown that

$$B - A^2 = 128M^{20}a^5b^5c^5(a^5 + b^5 + c^5).$$

Hence M^5a^5 , M^5b^5 , M^5c^5 are the roots of ρ in the cubic equation

$$\rho^3 + \frac{B - A^2}{2^7C^{2/3}}\rho^2 + \frac{1}{4} \left\{ \frac{(B - A^2)^2}{2^{14}C} - A \right\} \rho + C^{1/3} = 0.$$

A, B, C , it will be observed, are independent and, as they may be termed, prime or radical adjunctive constants. Hitherto much mystery and uncertainty have attached to the theory of hyperdeterminants, from its having been tacitly assumed that they were always either of lower dimensions than the ordinary determinant, or else algebraical functions of such, and of the determinant. Whereas we now see that, whilst the determinant of a function in two letters of the fifth degree is of eight dimensions, one of its radical or primitive hyperdeterminants is of four, but the other of twelve dimensions. This is a most valuable consequence, and would seem to indicate that the number of radical hyperdeterminants to a function, over and above the common determinant, is always equal to the number of parameters entering into its canonical form. The importance of this ascertainment of an unsuspected third *radical* constant, adjunctive to a quintic function of two letters, in making to march the theory of hyperdeterminants, can hardly be over-estimated.

From the equation last given we are enabled to assign the conditions in order that two functions of the fifth degree may be capable of being linearly transformed either into the other. For if we call F and F' two such linearly equivalent quintic functions, they must be capable each of being thrown under the same form $u^5 + v^5 + (lu + mv)^5$, where l and m shall be the same for each. Consequently we must have the roots of ρ in the same ratio for F and F' , which conditions may be expressed by means of the two equations

$$\begin{aligned} \frac{B - A^2}{C^{2/3}} &= \frac{B' - A'^2}{C'^{2/3}}, \\ \frac{(B - A^2)^2 - 2^{14}AC}{C^{4/3}} &= \frac{(B' - A'^2)^2 - 2^{14}A'C'}{C'^{4/3}}, \end{aligned}$$

A', B', C' , of course representing the same functions of the coefficients of F' as A, B, C , respectively of F .

The two conditions required in their simplest form are accordingly

$$\frac{A}{C^{1/3}} = \frac{A'}{C'^{1/3}}, \quad \frac{B}{C^{2/3}} = \frac{B'}{C'^{2/3}},$$

or

$$A^3 : B^2 : C :: A'^3 : B'^2 : C',$$

that is to say, *all quintic functions of two letters of which the determinant is to the subduplicate power of the radical hyperdeterminant of the twelfth order and to the sesquiduplicate power of the radical hyperdeterminant of the fourth order in given ratios, are mutually convertible.*

So for the quartic (that is, biquadratic) function of two letters, calling R and S the radical adjunctive constants of the second and third orders, the condition of convertibility between different forms of the same is, that $R^3 : S^2$ shall be a given ratio. And, in general, we may infer that the condition of convertibility between different functions of any degree is, that the several radical adjunctive constants of each raised respectively to such powers as will make them of like dimensions, shall be to one another in given ratios. Of course all cubic functions of two letters, according to this rule, are mutually convertible without any condition, they having but one radical adjunctive constant; and in fact all such functions, being representable as the sum of two cubes of new variables linearly related to those given, are necessarily convertible.

I have further succeeded in obtaining the canonical form of the *quadratic* adjunctive to *any odd degeed* function of two letters, which presents a wonderful analogy to the theory of relative determinants of *quadratic functions of any number* of letters, and constitutes an important step towards the construction of the theory of relative hyperdeterminants.

Let a function of two letters of the odd degree $m (= 2n - 1)$ be thrown under its canonical form,

$$u_1^m + u_2^m + \cdots + u_n^m,$$

¹¹⁵More strictly speaking (and this correction should be supplied throughout in the "Sketch"), B is the negative determinant of $\frac{1}{5}F$. After finding, by the method of characteristics, or any special artifices, the algebraic part of the value of a resultant or determinant, a process frequently of some complexity remains over in assigning its numerical multiplier; this part of the operation being analogous to that which occurs in the Integral Calculus, of determining the constant to be added after the general form of an integral has been determined. In the "Sketch," a correction for the numerical multiplier remains also to be applied to the expressions given for the successive Hessian determinants.

the biquadratic adjunctive constant will be

$$\frac{M^{14}}{(cd' - c'd)^{14}}$$

multiplied by the determinant

$$\begin{vmatrix} 0 & (ab' - a'b)^6 & (ac' - a'c)^6 & (ad' - a'd)^6 & a & a' \\ (ba' - b'a)^6 & 0 & (bc' - b'c)^6 & (bd' - b'd)^6 & b & b' \\ (ca' - c'a)^6 & (cb' - c'b)^6 & 0 & (cd' - c'd)^6 & c & c' \\ (da' - d'a)^6 & (db' - d'b)^6 & (dc' - d'c)^6 & 0 & d & d' \\ a & b & c & d & 0 & 0 \\ a' & b' & c' & d' & 0 & 0 \end{vmatrix}.$$

p. 208

The determinants of the Hessian, the post-Hessian, and the praeter-post-Hessian of F will be found (in the case of the quintic function) to be always multiples of powers of the determinant of the given function, and of its cubic adjunctive; and I believe that in general for a function of two letters of any degree the determinants of all the derived forms in the Hessian scale¹¹⁷, will be necessarily algebraical functions of any two of them.

I hope very shortly to accomplish the reduction of functions, as high as the seventh degree of two letters, to their canonical form, and also to present a complete theory of the failing or singular cases of canonical forms.

Since the above was in print I have discovered the following

GENERAL THEOREM

for reducing a function of two letters of any odd degree to its canonical form.

Let the degree of the function be $(2n - 1)$; then its canonical form is

$$u_1^{2n-1} + u_2^{2n-1} + \dots + u_n^{2n-1},$$

with $(n - 2)$ linear relations between $u_1, u_2, \dots u_n$.

To find $u_1, u_2, \dots u_n$, proceed as follows. Let the given function of the $(2n - 1)$ th degree be supposed to be

$$a_1 x^{2n-1} + (2n - 1)a_2 x^{2n-2}y + (2n - 1)\frac{2n - 2}{2}a_3 x^{2n-3}y^2 + \dots + a_{2n}y^{2n-1}.$$

¹¹⁶See Note (A) of Appendix.

¹¹⁷I use the term Hessian (more properly speaking the Boolean) Scale, to denote the *determinants* in respect of ξ and η of $(\xi \frac{d}{dx} + \eta \frac{d}{dy} + \&c.) F$.

Neither Hesse, however, nor any other writer up to the present time, had thought of constructing, and still less of turning to account, the functions (the first only excepted) which figure in this scale.

Form the determinant

$$\begin{vmatrix} a_1x + a_2y & a_2x + a_3y & a_3x + a_4y & \cdots & a_nx + a_{n+1}y \\ a_2x + a_3y & a_3x + a_4y & a_4x + a_5y & \cdots & a_{n+1}x + a_{n+2}y \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ a_nx + a_{n+1}y & a_{n+1}x + a_{n+2}y & a_{n+2}x + a_{n+3}y & \cdots & a_{2n-1}x + a_{2n}y \end{vmatrix}.$$

This determinant is a function of x and y of the n th degree, and by resolving an equation of the n th degree, may be decomposed into n factors, say

$$(l_1x + m_1y)(l_2x + m_2y) \cdots (l_nx + m_ny);$$

p. 209

we shall then have

$$u_1 = p_1(l_1x + m_1y),$$

$$u_2 = p_2(l_2x + m_2y),$$

$$\vdots$$

$$u_n = p_n(l_nx + m_ny),$$

where the l 's and m 's are known, and the $(2n - 1)$ th powers of the p 's may be found linearly, by means of the identical equation $\Sigma u^{2n-1} = F(x, y)$. Thus for example a function of the seventh degree of two letters may be reduced to its canonical form

$$(lx + my)^7 + (l'x + m'y)^7 + (l''x + m''y)^7 + (l'''x + m'''y)^7,$$

by the resolution of a biquadratic equation. My demonstration of this extraordinary and unexpected consequence rests upon the following lemma¹¹⁸, itself a very beautiful and striking theorem (no doubt capable of much generalisation) in the theory of determinants. Form the rectangular matrix consisting of n rows and $(n + 1)$ columns

$$\begin{array}{ccccc} T_1 & T_2 & T_3 & \cdots & T_{n+1} \\ T_2 & T_3 & T_4 & \cdots & T_{n+2} \\ T_3 & T_4 & T_5 & \cdots & T_{n+3} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ T_n & T_{n+1} & T_{n+2} & \cdots & T_{2n} \end{array}$$

where

$$T_i = a_1^{r-i}b_1^{s+i} + a_2^{r-i}b_2^{s+i} + \cdots + a_{n-1}^{r-i}b_{n-1}^{s+i}.$$

Then all the $n + 1$ determinants that can be formed by rejecting any one column at pleasure out of this matrix are identically zero.

In order the better to realise the proof, suppose

$$n = 4, \quad \text{so that} \quad 2n - 1 = 7.$$

¹¹⁸See Note (B) of Appendix.

Let

$$F(x, y) = a_1x^7 + 7a_2x^6y + 21a_3x^5y^2 + 35a_4x^4y^3 + 35a_5x^3y^4 \\ + 21a_6x^2y^5 + 7a_7xy^6 + a_8y^7.$$

Suppose

$$t^7 + u^7 + v^7 + w^7 = F(x, y) = G(u, v),$$

$$at + bu = v, \quad a't + b'u = w.$$

Then, if M is the modulus of transition from x, y to u, v the hyper-

p. 210

determinant, or, to adopt my new expression, the permutant P_4 (meaning thereby)

$$\begin{vmatrix} a_1x + a_2y & a_2x + a_3y & a_3x + a_4y & a_4x + a_5y \\ a_2x + a_3y & a_3x + a_4y & a_4x + a_5y & a_5x + a_6y \\ a_3x + a_4y & a_4x + a_5y & a_5x + a_6y & a_6x + a_7y \\ a_4x + a_5y & a_5x + a_6y & a_6x + a_7y & a_7x + a_8y \end{vmatrix},$$

which is a constant adjunctive in respect to ξ and η of $\left(\xi \frac{d}{dx} + \eta \frac{d}{dy}\right)^6 F$, will, according to the principles laid down in the preceding "Sketch," be the product of a power of M multiplied by the corresponding adjunctive constant of $\left(\xi \frac{d}{du} + \eta \frac{d}{dv}\right)^6 G(u, v)$, and is therefore a multiple of the determinant

$$\begin{vmatrix} (1 + A_1)t + A_2u & A_2t + A_3u & A_3t + A_4u & A_4t + A_5u \\ A_2t + A_3u & A_3t + A_4u & A_4t + A_5u & A_5t + A_6u \\ A_3t + A_4u & A_4t + A_5u & A_5t + A_6u & A_6t + A_7u \\ A_4t + A_5u & A_5t + A_6u & A_6t + A_7u & A_7t + (1 + A_8)u \end{vmatrix},$$

where

$$A_1 = a^7 + a'^7, \quad A_2 = a^6b + a'^6b', \quad A_3 = a^5b^2 + a'^5b'^2, \quad \dots \quad A_8 = b^7 + b'^7.$$

In this determinant the coefficient of u^4 is

$$\begin{vmatrix} A_2 & A_3 & A_4 & A_5 \\ A_3 & A_4 & A_5 & A_6 \\ A_4 & A_5 & A_6 & A_7 \\ A_5 & A_6 & A_7 & 1 + A_8 \end{vmatrix}$$

which is numerically equal to

$$A_5 \begin{vmatrix} A_3 & A_4 & A_5 \\ A_4 & A_5 & A_6 \\ A_5 & A_6 & A_7 \end{vmatrix} - A_6 \begin{vmatrix} A_2 & A_4 & A_5 \\ A_3 & A_5 & A_6 \\ A_4 & A_6 & A_7 \end{vmatrix} \\ + A_7 \begin{vmatrix} A_2 & A_3 & A_5 \\ A_3 & A_4 & A_6 \\ A_4 & A_5 & A_7 \end{vmatrix} - (1 + A_8) \begin{vmatrix} A_2 & A_3 & A_4 \\ A_3 & A_4 & A_5 \\ A_4 & A_5 & A_6 \end{vmatrix} = 0,$$

because the second factors of the products are all zero by the lemma. Hence the permutant P_4 vanishes when $t = 0$, and consequently it contains t as a factor, and in like manner it may be proved to contain u, v, w .

Hence t, u, v, w are the algebraical factors of P_4 , and precisely the same proof applies to show in the case of a function in x and y , say F_{2n-1} of any p. 211

odd degree $(2n - 1)$ whatever, that the corresponding permutant P_n will contain the factors $u_1, u_2 \dots u_n$ linear functions of x, y , such that

$$u_1^{2n-1} + u_2^{2n-1} + \dots + u_n^{2n-1} = F_{2n-1}$$

as was to be shown.

Whenever P_n has equal roots, this will denote either (which is the more general case) that the usual canonical form fails and gives place to a singular form, (owing to some of the coefficients of transformation becoming infinite), or, which is the more special supposition, that the canonical form becomes catalectic by one or more of the linear roots¹¹⁹ disappearing. Thus in the cubic function, if P_2 has equal roots, and consequently its determinant (which is coincident with that of the function itself) vanish, then the canonical form in general fails; so that, for example, $ax^3 + bx^2y$ cannot in general be exhibited as the sum of two cubes: if, however, certain further relations obtain between the coefficients of F , the canonical form reappears catalectically, the function becoming in fact representable as a single cube. So, again, for the quintic function (referring back to the notation above [page 205]), if P_3 have equal roots, that is if $C = 0$, the canonical form fails, unless at the same time $B - A^2 = 0$, in which case the function becomes the sum of two fifth powers; but if furthermore $A = 0$, then this catalectic form again gives place to a singular form, which, on the satisfaction of a further condition between the coefficients, again in its turn gives way before a (bicatalectic, that is) doubly catalectic form, namely, a single fifth power.

It is remarkable, that the form to which Mr Jerrard's method reduces the function of the fifth degree, expressed homogeneously as $ax^5 + bxy^4 + cy^5$, is a singular form, being incapable of being exhibited as the sum of three cubes; such, however, is not the case with the form $ax^5 + bx^3y^2 + cy^5$. It may further be remarked, that although the singly catalectic form of the quintic function is expressible by two conditions only, namely, $C = 0$, $B - A^2 = 0$, it will be indicated by P_3 (which being a cubic function of x and y contains four terms) completely disappearing, so that apparently four conditions would appear to be required or implied. But of course these must be capable of being shown to be non-independent, and to be merely tantamount to the two independent ones, $C = 0$, $B - A^2 = 0$. The theory of the catalectic forms of functions of the higher degrees of two variables presents many strong points of resemblance and of contrast to that of the catalectic forms of quadratic functions of several variables.

¹¹⁹ $u_1, u_2 \dots u_n$ may be termed the linear roots of the form F_{2n-1} .

One important and immediate corollary from the General Theorem is, that the constants which enter into the linear functions appurtenant to the canonical form of any function of an odd degree form a single and unique system; or, in other words, the canonical forms for such functions are void of

multiplicity, a result contrary to what might have been anticipated, and to what we know is the case for the canonical forms of functions of an even degree.

It may further be shown that if we have the $(n-2)$ equations

$$\begin{array}{l} a_1u_1 + a_2u_2 + \cdots + a_nu_n = 0, \\ b_1u_1 + b_2u_2 + \cdots + b_nu_n = 0, \\ \dots\dots\dots \\ l_1u_1 + l_2u_2 + \cdots + l_nu_n = 0, \end{array}$$

and call M the modulus of transformation in respect to u_1, u_2 , and if we make

$$P_n = Ku_1u_2 \cdots u_n,$$

then

$$\left| \begin{array}{cc} a_3, & a_4 \cdots a_n \\ b_3, & b_4 \cdots b_n \\ & \dots \\ l_3, & l_4 \cdots l_n \end{array} \right| \begin{array}{c} \frac{1}{2}n(n-1) \\ K \end{array}$$

is equal to the product of the $\frac{1}{2}n(n-1)$ factors of the form

$$\begin{vmatrix} a_{\theta_1} & a_{\theta_2} & \cdots & a_{\theta_{n-2}} \\ b_{\theta_1} & b_{\theta_2} & \cdots & b_{\theta_{n-2}} \\ \cdots & \cdots & \cdots & \cdots \\ l_{\theta_1} & l_{\theta_2} & \cdots & l_{\theta_{n-2}} \end{vmatrix},$$

$\theta_1, \theta_2 \dots \theta_{n-2}$ being any $(n-2)$ numbers out of the n numbers $1, 2, 3 \dots n$.

It may hence be shown that

$$u_1 u_2 \cdots u_n = \frac{P_n}{\left(\frac{1}{m} \square_{x,y} P_n\right)^{1/(2n-1)}}^{120},$$

m being a number which is a function of n , and which may be shown to be equal to $\square_{x,y}(x^{n-1}y + xy^{n-1}) \div$ product of the squared differences of the roots of $l^{n-2} = 1$, that is

$$m = \left\{ \frac{(n-1)^2 - 1}{-(n-2)} \right\}^{n-2} = (-n)^{n-2},$$

and thus

$$u_1 u_2 \cdots u_n = \frac{P_n}{\sqrt[2n-1]{\{(-n)^{-(n-2)} \square_{x,y} P_n\}}}.$$

As an example of the mode of finding $u_1, u_2 \dots u_n$, let

$$F = 3x^5 + 20x^3y^2 + 10xy^4,$$

then

$$P_3 = \begin{vmatrix} 3x & 2y & 2x \\ 2y & 2x & 2y \\ 2x & 2y & 2x \end{vmatrix} = 4x^3 - 4y^2x.$$

Hence

$$u = fx, \quad v = g(x + y), \quad w = h(x - y).$$

To find f, g, h , we have $u^5 + v^5 + w^5 = F$, hence

$$f^5 + g^5 + h^5 = 3; \quad g^5 + h^5 = 2; \quad g^5 - h^5 = 1;$$

whence we have

$$F = x^5 + (x + y)^5 + (x - y)^5.$$

Again, we find

$$\square(4x^3 - 4y^2x) = -4^4 \times 12,$$

$$\left(\frac{1}{(-3)} \square P_3 \right)^{1/5} = 4,$$

and accordingly

$$x(x + y)(x - y) = \frac{P_3}{\{(-3)^{-1} \cdot \square P_3\}^{1/5}},$$

according to the general formula above given.

As a second example let

$$F = 3x^7 + 42x^5y^2 + 70x^3y^4 + 14xy^6 + y^7;$$

then

$$P_4 = \begin{vmatrix} 3x & 2y & 2x & 2y \\ 2y & 2x & 2y & 2x \\ 2x & 2y & 2x & 2y \\ 2y & 2x & 2y & 2x + y \end{vmatrix} = 4(x^3y - xy^3) = 4xy(x - y)(x + y),$$

and accordingly we shall find

$$x^7 + y^7 + (x - y)^7 + (x + y)^7 = F.$$

Moreover

$$\square(4x^3y - 4xy^3) = 4^9,$$

¹²⁰ $\square_{x,y}$ means the determinant in respect to x and y .

and

$$4^{4-2} = 4^2.$$

Thus

$$\frac{P_4}{tuvw} = \sqrt[7]{\frac{\square P_4}{4^{4-2}}},$$

agreeable to the general formula.

p. 214

As a corollary to our general proposition, it may be remarked, that if F_{2n-1} be a symmetrical function of x, y of the $(2n-1)$ th degree, $P_n(F_{2n-1})$ will be also a symmetrical function of x and y , and may therefore be resolved into its factors by solving a *recurring* equation of the n th degree, which may, by well-known methods, be made to depend on the solution of an equation of the $\frac{1}{2}n$ th or $\frac{1}{2}(n-1)$ th degree, according as n is even or odd.

Hence the reduction of a function of two letters of the degree $4m \pm 1$ to its canonical form as the sum of powers may be made to depend on the solution of an equation of the m th degree; so that, for example, a symmetrical function of x, y , as high as the fifteenth or seventeenth degree, may be reduced by means of a biquadratic equation only.

In a short time I hope to present to the public a complete solution of the canonical forms of functions of two letters of even degrees, and possibly to exhibit some important applications of the principles of the method to the theory of numbers.

APPENDIX.

NOTE (A).

The permutants (meaning, in Mr Cayley's language, the hyperdeterminants) of $F_{2n+1}(x, y)$ of the fourth dimension in respect to the coefficients of F , may be all obtained by taking the quadratic permutant in respect to x and y of the quadratic permutant in respect of ξ and η of

$$\left(\xi \frac{d}{dx} + \eta \frac{d}{dy} \right)^{2l} F_{2n+1}(x, y),$$

l having any integer value from 1 to n .

In extension of a theorem in the foregoing Supplement, which applies only to the case of $l = n$, I am able to state the following more general theorem, in which the same notation is preserved as above [page 207]. The quadratic permutant in respect to ξ and η of

$$\frac{1}{(2n+1)2n \cdots (2n-2l+2)} \left(\xi \frac{d}{dx} + \eta \frac{d}{dy} \right)^{2l} F_{2n+1}(x, y),$$

is equal to

$$\frac{M^{2l}}{(1, 2)^{2l}} \sum \{(\theta_r, \theta_s)^{2l} (u_r . u_s)^{2n+1-2l}\}.$$

If now we proceed to form the quadratic permutant of the above sum in respect to x and y , we know *à priori*, by reason of Mr Cayley's invaluable researches, that we shall not get radically distinct results for all values, but only for certain periodically changing values of l . p. 215

I have not yet had leisure to seek for an explicit demonstration of this remarkable law, founded upon the above given canonical representation.

NOTE (B).

The lemma, upon which the general method for reducing odd degreed functions to their canonical form is founded, may be stated rather more simply and more generally as follows:—

The determinant

$$\begin{vmatrix} T_{r_1} & T_{r_2} & \cdots & T_{r_n} \\ T_{r_1+t_1} & T_{r_2+t_1} & \cdots & T_{r_n+t_1} \\ T_{r_1+t_2} & T_{r_2+t_2} & \cdots & T_{r_n+t_2} \\ \cdots & \cdots & \cdots & \cdots \\ T_{r_1+t_{n-1}} & T_{r_2+t_{n-1}} & \cdots & T_{r_n+t_{n-1}} \end{vmatrix},$$

where T_θ denotes $A_1 a_1^\theta + A_2 a_2^\theta + \cdots + A_m a_m^\theta$ provided that m is less than n , is identically zero. In the theorem, as thus stated, there is no substantial loss of generality arising from the omission of the b 's.

Thus stated the theorem and its extensions evidently repose upon the same or the like basis as the theory of partial fractions.

NOTE (C), REFERRING TO THE ORIGINAL "SKETCH."

The Boolo-Hessian scale of determinants furnishes a very pretty general theorem of geometrical reciprocity in connexion with the doctrine of successive polars. Let $F(x, y, z)$, a cubic homogeneous function of x, y, z equated to zero, express in general a curve of the third degree; then

$$\left(a \frac{d}{dx} + b \frac{d}{dy} + c \frac{d}{dz}\right) F$$

will express its first polar in respect to the point a, b, c , that is, the conic which passes through the six points in which the tangents drawn from a, b, c to touch the given curve meet the same.

Again, if we take l, m, n the coordinates of any new point,

$$\left(l \frac{d}{dx} + m \frac{d}{dy} + n \frac{d}{dz}\right) \left(a \frac{d}{dx} + b \frac{d}{dy} + c \frac{d}{dz}\right) F$$

will express the polar, that is the chord of contact of the above conic, in respect to the last named point. If now we eliminate l, m, n between the three equations

$$\begin{aligned} \left(l \frac{d}{dx} + m \frac{d}{dy} + n \frac{d}{dz}\right) \left(a \frac{d}{dx} + b \frac{d}{dy} + c \frac{d}{dz}\right) F &= 0, \\ \left(l \frac{d}{dx} + m \frac{d}{dy} + n \frac{d}{dz}\right) \left(a' \frac{d}{dx} + b' \frac{d}{dy} + c' \frac{d}{dz}\right) F &= 0, \\ \left(l \frac{d}{dx} + m \frac{d}{dy} + n \frac{d}{dz}\right) \left(a'' \frac{d}{dx} + b'' \frac{d}{dy} + c'' \frac{d}{dz}\right) F &= 0, \end{aligned}$$

it is easily seen that the resultant of the elimination is the square of the determinant

$$\begin{vmatrix} a & b & c \\ a' & b' & c' \\ a'' & b'' & c'' \end{vmatrix},$$

multiplied by the Hessian of the given function. And, moreover, that if we eliminate x, y, z we shall obtain precisely the same result with the letters l, m, n substituted for x, y, z . Hence it follows, that if we take the doubly infinite system of first polars to a given curve of the third degree, in respect to all the points lying in its plane, and then from any point in the *Hessian* to the given curve, draw pairs of tangents to each conic of the system so generated, then all the chords of contact will meet in one and the same point, which will itself be also a point situated upon the Hessian and conjugate to the former.

So, in general, for a function of any degree of any number of letters, viewed with relation to the doctrine of successive polars, the determinants of the Boolo-Hessian scale take one another up in pairs; namely the first takes up the last but one, the second the last but two, and so on; and consequently, if the degree of the function be odd, that function which (making abstraction of the constant determinant at the end) lies in the middle of the scale pairs with itself, and, in a sense analogous to that above exhibited for a function of the third degree, may be said to be always its own reciprocal.

P.S. I have just discovered the method of reducing functions of two letters of *even degrees* to their canonical form, which will shortly be published in a second Supplement.

At present I offer the annexed theorem (which strikingly contrasts with the law of uniqueness demonstrated of functions of an odd degree) as a foretaste of the enchanting developments with which I hope shortly to present my readers:—

If a given homogeneous function of x and y of the degree $2n$ be supposed to be thrown under its canonical form,

$$u_1^{2n} + u_2^{2n} + \cdots + u_n^{2n} + K(u_1 u_2 \cdots u_n)^2,$$

then will K^n have $n^2 - 1$ in general distinct values, to each of which will correspond

a single distinct system of the linear functions of x and y ,

$$1^{1/n}u_1, \quad 1^{1/n}u_2, \quad \dots \quad 1^{1/n}u_n.$$

35.

EXPLANATION OF THE COINCIDENCE OF A THEOREM GIVEN BY MR SYLVESTER IN THE DECEMBER NUMBER OF THIS JOURNAL, WITH ONE STATED BY PROFESSOR DONKIN IN THE JUNE NUMBER OF THE SAME

[*Philosophical Magazine, Fourth Series, I. (1851), pp. 44-46*]

p. 217

I wish to state, without loss of time, that in the theorem given by me¹²¹ for the composition of two successive rotations about different axes, I have been anticipated by Prof. Donkin in the June Number of your *Journal*.

To my shame I must confess, that, although an occasional contributor to, I am not invariably a constant reader of your valuable miscellany, otherwise I should not have introduced the theorem in question without due acknowledgment of Professor Donkin's claims to whatever merit may attach to the priority of publication. The fact is, that I made out the theorem for myself nine years ago, and had some communication on the subject with Professor De Morgan, who was then writing the seventeenth chapter of his *Differential Calculus*. A recent conversation with this gentleman has brought back to my mind a vivid recollection of the course of that communication. I brought under Professor De Morgan's notice the analytical memoir of Sr Gabrio Pola on the subject in the Memoirs of the Italian Society of Modena, and satisfied myself of the existence of the single axis of displacement by compounding the two rotations in the manner given in my paper, which, for the case of two axes fixed in space, is the same as Professor Donkin's, and for two axes fixed in the rotating body is materially, although not formally the same.

It then occurred to me that a more simple demonstration ought to be deducible from the possibility of always *finding* the point on a sphere, by revolution about which, as a pole, one equal arc could actually be shown to be transportable into the place of another. But in proceeding to work out this idea I fell into a remarkable blunder, in which I have since been followed by more than one able friend to whom I have proposed the question. The

p. 218

blunder was of this kind:—Two arcs have to be drawn, bisecting at right angles the arcs joining the extremities of two equal arcs; the point of intersection of the two bisecting arcs must in all cases fall outside the quadrilateral formed by the equal and joining arcs. I supposed it to fall inside. There appears to be a fatal tendency to do so in all who take the subject in hand. In consequence of this error, the cause of which I did not at the moment perceive, I was driven to deny and admit in one breath the same proposition. Mr De Morgan sent me the correct proof after this method (the same as that given by him at page 489 of his *Calculus*), I am inclined to think after I had myself detected my error; but of this I cannot feel certain.

¹²¹p. 158 above.

This is the method alluded to by me in the words “it is right to bear in mind, &c.,” at the time of writing which all recollection of the same thing having been published by Mr De Morgan had vanished from my memory.

The proof of the triangle of rotations is so simple, that, as Professor Donkin states (in a letter which he has done me the favour of addressing me on the subject) was the case with himself, I thought it incredible that it should not have appeared in some elementary work, and I was therefore at no pains to publish it as my own; nor should I have written at all on the subject, had it not been for the surprise occasioned to my mind by falling in with Professor Stokes’s article in the *Cambridge and Dublin Mathematical Journal*, to demonstrate the existence of an instantaneous axis, which proceeds in apparent unconsciousness of the so simply demonstrable law, that any number of rotations of any kind (and therefore those that take place in an instant of time) are representable by a single rotation about a single axis. I shall feel obliged by the early insertion of this explanation, more in justice to myself than to Professor Donkin, whose high and worthily earned reputation, not to speak of the disinterested love of truth for its own sake, apart from personal considerations, which animates the labours of the genuine votary of science, must make him indifferent to whatever credit might be supposed to result from the first authorship or publication of the very simple (however important) theorem in question.

36.

AN ENUMERATION OF THE CONTACTS OF LINES AND SURFACES OF THE SECOND ORDER

[Philosophical Magazine, Fourth Series, I. (1851), pp. 119–140]

p. 219

It is well known that in general any two homogeneous quadratic functions of the same system of variables may be simultaneously transformed, so as to be expressed each of them as pure quadratic functions of a new system of variables equal in number and linearly connected with the original ones; a pure quadratic function meaning one in which only the squares of the variables are retained.

Every homogeneous quadratic function may be treated as the characteristic¹²² of a locus of the second degree: if the function be of two letters, the locus is a binary system of points in a line wherein the distances of two fixed points from either point of the given system or given multiples of such distances correspond to the variables; if of three letters, the locus is a conic, the distances or given multiples of the distances of every point in which from three given lines in the plane of the conic are represented by the variables; if of four letters, the locus is a surface of the second order, the coordinates being the distances or multiples of the distances of any point therein from four planes drawn in the space in which the surface is contained, and so on for loci of four and higher dimensions.

I propose, however, in the present paper to restrict myself to the theory of the contacts of loci not transcending the limits of vulgar space, by which I mean the space cognizable through the senses¹²³, and shall accordingly be

p. 220

almost exclusively concerned in determining the singular cases of conjugate systems of quadratic forms of two, three, and four letters respectively.

In order that the reduction of any such system, say U and V , to a pure quadratic form may be possible (as it generally is), it is necessary that none of the roots of the complete determinant of $U + \lambda V$ shall be equal; if any relation of equality exist between these roots, the general reduction is generally no longer possible; under peculiar conditions, however, as will hereafter appear, in spite

¹²²According to the definition stated by me in a previous paper, the characteristic of a locus is the function which, equated to zero, constitutes the equation thereto.

¹²³If the impressions of outward objects came only through the sight, and there were no sense of touch or resistance, would not space of three dimensions have been physically inconceivable? The geometry of three dimensions in ordinary parlance would then have been called transcendental. But in very truth the distinction is vain and futile. Geometry, to be properly understood, must be studied under a universal point of view; every (even the most elementary) proposition must be regarded as a fact, and but as a single specimen of an infinite series of homologous facts.

In this way only (discarding as but the transient outward form of a limited portion of an infinite system of ideas, all notion of extension as essential to the conception of geometry, however useful as a suggestive element) we may hope to see accomplished an organic and vital development of the science.

of the equality of certain of the roots, the irreducibility in its turn will cease, and the ordinary reduction be capable of being effected. It is easily seen, that to every relation of equality between the roots of the determinant of $U + \lambda V$ must correspond a particular species of contact between the loci which U and V characterize. But we should make a great mistake were we to suppose that every such relation of equality corresponded with but one species of contact; for instance, the characteristics of U and V of two conics are functions of three letters, and $\Delta(U + \lambda V)$ will be a cubic function of λ . Such a function may have two roots, or all its roots equal: this would seem to give but two species of contact, whereas we well know that there are no less than four species of contact possible between two conics. Accordingly we shall find, that, in order to determine the distinctive characters of each species of contact, we must look beyond the complete determinant, and examine into the relations (in themselves and to one another) of the several systems of minor determinants that can be formed from $U + \lambda V$.

By pursuing this method, we may assign *à priori* all the possible species of contact between any two loci of the second degree. How important this method is will be apparent from the fact, that not only have the distinctive characters of the various contacts possible between surfaces of the second order never been determined, but their number and the nature of certain of them have remained until this hour unknown and unsuspected.

The method which we shall pursue is an exhaustive one, and will conduct us by a natural order to a systematic arrangement of all the different modes and gradations of such contacts.

In a paper¹²⁴ in this Magazine for November 1850, I explained the decline of minor determinants, and stated a law, called the homaloidal law, concerning them.

If U and V be characteristics of the two loci whose contacts are to be considered, $U + \lambda V$ will be the function, the properties of whose complete determinant, and of the minor systems of determinants belonging to it, will serve to specify the nature of the contact.

It will be remembered, that, whatever be the number of variable letters in any quadratic function U , three of its first minor determinants being zero, p. 221

makes all the first minors zero; six of its second minors being zero, makes all the second minors zero; and so on for the third, fourth, &c. minor systems according to the progression of the triangular numbers.

It is well known that whatever linear transformations be applied to a quadratic function W , the complete determinant thereof will remain unaltered, except by a multiplier depending upon the coefficients introduced into the equations of transformation; consequently the roots of λ in the equation obtained by making

¹²⁴p. 150 above.

the determinant of $U + \lambda V$ zero remain unaffected by such transformation; and any relation or relations of equality among the roots of the equation $\square(U + \lambda V) = 0$ is an immutable property of the system U, V , which is unaffected by linear transformations. Another and more general kind of immutable property (comprehending the above as a particular case), to which I shall have occasion to refer, is the following.

Suppose all the minors of any order of $U + \lambda V$ have a factor $\lambda + e$ in common; this factor will continue common to the same system of minors when U and V are simultaneously transformed. This is a very important proposition, and easily demonstrated; for if $\lambda + e$ be a common factor to all the r th minors of $U + \lambda V$, $(U - eV)$ will have its r th minors zero, and therefore, as explained by me in the paper above referred to, $U - eV$ will be degraded r orders below U or V . This is clearly a property independent of linear transformation, consequently $\lambda + e$ will remain a factor of the transformed r th minors.

In like manner it is demonstrable that any number of distinct factors $\lambda + e_1, \lambda + e_2, \dots$ common to the r th minors of one form of $U + \lambda V$, will remain common factors of any other linearly derived form of the same. It is consequently necessary that each r th minor of one form of any quadratic function W shall be a syzygetic¹²⁵ function of all the r th minors of any other form of the same; and consequently a function of λ of any degree, whether all its factors be or be not distinct, which is common to the r th minors of one form of $U + \lambda V$, will remain so to the r th minors of any other form of the same.

The law exhibiting the connexion of each r th minor of one form of W (any homogeneous quadratic function) with all the r th minors of any other form of W , will form the subject of a distinct communication.

Finally, to fully comprehend the annexed discussion, the following principle must be apprehended.

p. 222

If any factor K^e enter into all the r th minors of W , and if K^i be the highest power of K common to all the $(r + 1)$ th minors, then K^{2e-i} will be a common factor to all the $(r - 1)$ th minors.

Let r be taken unity; it is easily proved¹²⁶ that the complete determinant of any square matrix may be expressed by the difference between two products¹²⁷, each

¹²⁵If $A = pL + qM + rN + \&c.$, where $p, q, r \dots$ do not any of them become infinite when $L, M, N \dots$ or any of them become zero, A may be termed a syzygetic function of $L, M, N \dots$

In the theorem above alluded to, it will be shown (as might be expected) that the syzygy in the case concerned is of the simplest kind, that is, that each r th minor of a quadratic function of any number of letters is a homogeneous linear function of all the r th minors of the same quadratic function linearly transformed.

¹²⁶This will appear in my promised paper on Determinants and Quadratic Functions.

¹²⁷When the matrix is symmetrical about one of its diagonals (as it is in the case which we are concerned with), one of these products becomes a square. I may take this occasion of hinting, that the theory of quadratic functions merges in a larger theory of binary functions, consisting of the sum of the multiples of binary products formed by combining each of one set of quantities, $x, y, z \dots$ with each of the same number of quantities of another set, as $x', y', z' \dots$

of two first minor determinants divided by a certain second minor determinant. The proposition is therefore demonstrated for this case, and thereby in fact implicitly for every case, inasmuch as the first minors of every r th minor are $(r + 1)$ th minors of the original matrix. Hence it follows, that if any system of r th minor determinants have a common factor ε^i , the complete determinant must contain at lowest the factor $\varepsilon^{(r+1)i}$, and any system of $(r - s)$ th minor determinants thereunto will contain at lowest the factor $\varepsilon^{(s+1)i}$.

I now proceed to apply these principles to the determination of the relative forms of conjugate quadratic functions representing geometrical loci of the second order. I shall begin with two binary systems of points in a right line.

The general characteristics U and V of two such systems may be thrown under the form

$$\left. \begin{aligned} U &= x^2 + y^2, \\ V &= ax^2 + by^2 \end{aligned} \right\}.$$

When $\square(V + \lambda U) = 0$ has its two roots equal, these systems have a point in common. The above forms cease to be applicable, and convert into

$$\left. \begin{aligned} U &= xy, \\ V &= ax^2 + bxy \end{aligned} \right\}$$

where $x = 0$ represents the common point.

p. 223

Let U and V now represent two conics. When there is no contact, we have as the types of their characteristics

$$U = x^2 + y^2 + z^2, \quad V = ax^2 + by^2 + cz^2.$$

The three roots of $\square(V + \lambda U) = 0$ are

$$\lambda = -a, \quad \lambda = -b, \quad \lambda = -c,$$

showing that there are three distinct pairs of lines in which the intersections of U and V are contained, the equations to three pairs being respectively

$$\begin{aligned} (b - a)y^2 + (c - a)z^2 &= 0, \\ (c - b)z^2 + (a - b)x^2 &= 0, \\ (a - c)x^2 + (b - c)y^2 &= 0; \end{aligned}$$

For instance,

$$axx' + bxy' + cxz' + a'yx' + b'yy' + c'yz' + a''zx' + b''zy' + c''zz'$$

would be a binary function, and its determinant (no longer, as in a quadratic function, symmetrical about either diagonal) would correspond to the square matrix

$$\begin{vmatrix} a & b & c \\ a' & b' & c' \\ a'' & b'' & c'' \end{vmatrix}.$$

Almost all the properties of quadratic apply, with slight modifications, to binary functions.

the four points of the intersection being defined by the equations corresponding to the proportions

$$x : y : z :: \sqrt{(b-c)} : \sqrt{(c-a)} : \sqrt{(a-b)}.$$

Now let $\square(U + \lambda V)$ have two equal roots; the characteristics assume the form

$$\left. \begin{aligned} U &= x^2 + y^2 + xz, \\ V &= ax^2 + by^2 + cxz \end{aligned} \right\}.$$

Two of the pairs of lines become identical, that is, two of the four points of intersection coincide.¹²⁸ p. 224

This may be termed "Simple Contact." The tangent at the point of contact is $x = 0$; this equation making U and V each become of only one order.

The intersections are

$$\begin{aligned} x &= 0, & y &= 0, & (1) \\ x &= 0, & y &= 0, & (2) \\ \sqrt{(a-c)}x + \sqrt{(b-c)}y &= 0, & z &= 0, & (3) \\ \sqrt{(a-c)}x - \sqrt{(b-c)}y &= 0, & z &= 0. & (4) \end{aligned}$$

These are obtained by making $V - aU = 0$, which gives $x = 0$ or $z = 0$.

¹²⁸We may if we please make $a = b$; for it may be shown that the equations, in their present forms, contain an arbitrariness of 10 degrees; namely, 9 on account of x, y, z being arbitrary linears of ξ, η, θ ; 2 on account of the ratios $a : b : c$; together 11 reduced by one degree on account of x, y, z , changed into lx, ly, lz , leaving $U = 0, V = 0$ unaffected. Now the degrees of arbitrariness in two conics, subject to satisfy only one condition, is $2 \times 5 - 1$ or 9. Hence there is one degree of arbitrariness to spare. In fact, on making $a = b$, the axis z becomes the line joining the two points of intersection distinct from the point of contact; x remaining the tangent at the point of contact, and y , strange to say, still arbitrary, subject only to passing through the point of contact; if, however, y be made to pass through the point of contact, and either one of the distinct intersections, this form,

$$\begin{aligned} U &= x^2 + y^2 + xz, \\ V &= ax^2 + ay^2 + cxz, \end{aligned}$$

becomes no longer tenable, but gives place to

$$\begin{aligned} U &= y^2 + yx + xz, \\ V &= ay^2 + ayx + cxz, \end{aligned}$$

where x is the tangent at the point of contact, z the line joining the two intersections with one another, and $x, x + y$ respectively the lines joining either of them with the point of contact; if the multiplier of yx in V in the above be made b instead of a , x remains the tangent as before, y becomes any line through the point of contact, and z any line through one of the distinct intersections. A systematic view of similar modulations of form and the study of the laws of arbitrariness connected with them, as applicable to the general subject-matter of this paper, must be deferred to a subsequent occasion.

$x = 0$ gives $y^2 = 0$, that is, $y = 0$ twice over, and $z = 0$ gives

$$(a - c)x^2 + (b - c)y^2 = 0.$$

The number of conditions to be satisfied in this case is one only.

Next let $\square(U + \lambda V)$ have all its roots equal. This condition will be satisfied (still leaving U and V as general as they can remain consistent with these conditions) by making

$$U = x^2 + yz + yx, \quad V = ax^2 + ayz + byx.$$

Here only one distinct pair of lines can be drawn to contain the intersections, showing that three out of the four points come together.

This may be termed "Proximal Contact." The number of affirmative conditions to be satisfied is two, and the contact is therefore entitled of the second degree.

The tangent at the point of contact is $y = 0$, and the four intersections become

$$\begin{aligned} x = 0, \quad y = 0, \\ x = 0, \quad y = 0, \\ x = 0, \quad y = 0, \\ x = 0, \quad z = 0. \end{aligned}$$

These may be obtained from the equation $V - aU = 0$, which gives $y = 0$ or $z = 0$; the former implying concurrently with itself $x^2 = 0$, and the latter $yz = 0$.

Thus we obtain three systems,

$$x = 0, \quad y = 0,$$

and one

$$x = 0, \quad z = 0,$$

corresponding to three consecutive points and the single distinct one.

p. 225

The determinant of $U + \lambda V$ being only of the third degree in λ , we have exhausted the singularities of the system U, V dependent on the form of the complete determinant of $U + \lambda V$.

Let now the first minors of $U + \lambda V$ have a factor in common; this will indicate that $U + \lambda V$ may be made to lose two orders by rightly assigning λ , in other words, that the intersections of U and V are contained upon a pair of coincident lines. Here it is remarkable that the original forms of U and V reappear, but with a special relation of equality between the coefficients: we shall have, in fact,

$$U = x^2 + y^2 + z^2, \quad V = ax^2 + ay^2 + bz^2.$$

This gives the law for double, or, as I prefer to call it, diploidal contact¹²⁹. By virtue of the Homaloidal law, we know that if three first minors of $U + \lambda V$ be zero, all are zero; we have therefore to express that three quadratic functions of λ have a root in common. This implies the existence of two affirmative conditions; the contact of the two conics taken collectively may therefore be still entitled of the second degree, although the contact at each of the two points where it takes place is simple, or of the first degree.

These points are evidently defined by the equation

$$\begin{cases} x + \sqrt{-1}y = 0, & z = 0, \\ x - \sqrt{-1}y = 0, & z = 0, \end{cases}$$

and the ordinary algebraical solution of the equations $U = 0, V = 0$ would naturally lead to the four systems

$$\begin{aligned} x + \sqrt{-1}y &= 0, & z &= 0, \\ x + \sqrt{-1}y &= 0, & z &= 0, \\ x - \sqrt{-1}y &= 0, & z &= 0, \\ x - \sqrt{-1}y &= 0, & z &= 0; \end{aligned}$$

the two tangents at the point of contact are $x + \sqrt{-1}y = 0, x - \sqrt{-1}y = 0$, and the coincident pair of lines containing the intersections is $z^2 = 0$.

p. 226

It may at first view appear strange, that whilst no condition is required in order that U and V may be simultaneously metamorphosed into the forms of $x^2 + y^2 + z^2, ax^2 + by^2 + cz^2$, a, b and c being all unequal, for this metamorphosis to be possible when any two become equal, not one but two conditions must be satisfied. The reason of this is, that the coefficients of transformation, which, as well as a, b, c , are functions of the coefficients of the given quadratic functions, become infinite on constituting between the said coefficients such relations as are necessary for satisfying the equation $a = b$, or $a = c$, or $b = c$, except upon the assumption of some further particular relations between them over and above that implied in such equality.

¹²⁹See my remarks on the conditions which express double contact in the *Cambridge Journal*, Nov. 1850; see p. 129 above. If n functions, being all zero, be the condition of a fact, but r independent syzygetic equations admit of being formed between these functions, the number of affirmative conditions required is not n , but $(n - r)$; because the fact may be expressed by affirming $(n - r)$ equations and denying certain others. Thus if $P = 0, Q = 0, R = 0, S = 0$ express a fact, and

$$\begin{aligned} PP' + QQ' + RR' + SS' &= 0, \\ PP'' + QQ'' + RR'' + SS'' &= 0, \end{aligned}$$

the fact is expressible by affirming $P = 0, Q = 0$, and denying $R'S'' - R''S' = 0$, for then $P = 0, Q = 0$ will imply $R = 0, S = 0$; or, in like manner, by affirming any other two out of the four necessary equations, and denying the other equations. Observe, however, that all the required equations may coexist in the absence of such right of denial.

¹²⁹p. 129 above.

In the ordinary case of diploidal contact, the first minors having a factor in common, this factor will enter twice into the complete determinant of $U + \lambda V$, but it may enter three times: this will indicate, that not only do the four intersections lie on a coincident pair of lines, but furthermore, that there is but one pair of lines of any kind on which they lie.

In the ordinary case of diploidal contact, it will be observed that this latter condition does not obtain; the four intersections lie on a coincident pair of lines; but they lie also on a crossing pair, namely, in the two tangents at the points of contact. In this higher species of diploidal contact, it is clear that the two points of contact, which are ordinarily distinct, come together, and that all four intersections coincide.

This I call confluent contact; the forms of U and V corresponding thereto will be

$$U = x^2 + y^2 + xz, \quad V = ay^2 + axz;$$

the common tangent at the point of contact being $x = 0$, and the four coincident points $x^2 = 0, y^2 = 0$.

The number of affirmative conditions to be satisfied being three, the contact is to be entitled of the third degree.

Observe, that it is of no use to descend below the first minors in this case; because the second minors, being linear functions of λ , could not have a factor in common, unless $V : U$ becomes a numerical ratio, which would imply that the conics coincided¹³⁰.

Fortified by the successful application of our general principles to the preceding more familiar cases of contact, we are now in a condition to apply with greater confidence the same *à priori* method to the exhaustion and characterization of all the varied species of contact possible between surfaces

p. 227

of the second order; a portion of the subject comparatively unexplored, and never before thought susceptible of reduction to a systematic arrangement.

When there is no contact, we may write

$$U = x^2 + y^2 + z^2 + t^2, \quad V = ax^2 + by^2 + cz^2 + dt^2,$$

and the intersection of the surfaces will lie in each of the four cones,

$$\begin{aligned} (a-d)x^2 + (b-d)y^2 + (c-d)z^2 &= 0, \\ (a-b)x^2 + (c-b)z^2 + (d-b)t^2 &= 0, \\ (a-c)x^2 + (b-c)y^2 + (d-c)t^2 &= 0, \\ (b-a)y^2 + (c-a)z^2 + (d-a)t^2 &= 0. \end{aligned}$$

Whenever the surfaces are in contact, certain of these cones will coincide with certain others, so that their number will be always less than four. Also, as

¹³⁰No-contact and complete coincidence may be conceived as the two extreme cases in the scale of relative conjugate forms.

we shall find in such event, they may degenerate into pairs of intersecting or coincident planes.

Let us begin with considering the cases of contact for which the first minors (and consequently *à fortiori* the minors inferior to the first) have no factor in common.

Here $\square(V + \lambda U)$ is a biquadratic function.

If λ have all its roots unequal, we have U and V as above given.

If two roots are equal, the characteristics assume the form

$$\left. \begin{aligned} U &= x^2 + y^2 + z^2 + xt, \\ V &= ax^2 + by^2 + cz^2 + dxt \end{aligned} \right\}.$$

The touching plane is $x = 0$; the point of contact is $x = 0, y = 0, z = 0$; the curve of intersection is one of the fourth degree, with a double point at the point of contact.

There is but one condition to be satisfied, and the contact may be entitled “simple” and of the first degree.

Next let λ have three equal values, the equations become

$$U = x^2 + yz + t^2 + xy, \quad V = x^2 + yz + at^2 + bxy.$$

The tangent plane at the point of contact $y = 0$, and the point itself $x = 0, y = 0, t = 0$. The curve of intersection is a curve of the fourth order, with a cusp at the point of contact. The number of affirmative conditions to be satisfied is two; the contact is of the second degree, and may be termed “proximal” or cuspidal. p. 228

Next let $\square(U + \lambda V)$ have two pairs of equal roots, we shall find

$$U = x^2 + xy + zt, \quad V = ayz + bxy + czt.$$

The line $x = 0, z = 0$ will be common to both surfaces. The curve of intersection will therefore break up into a right line and a line of the third order.

The former will meet the latter in two points, which will be each of them points of contact. The contact is therefore diploidal; but as there is another species of diploidal contact to which we shall presently come, it will be expedient to characterize each of them by the nature of the intersections of the two surfaces; accordingly this may be termed unilinear-intersection contact, or more briefly, unilinear contact.

The number of affirmative conditions to be satisfied being two, it may be said to be collectively of the second degree, but (obviously?) the contact at each of the two points is of the nature of simple contact.

Lastly, let us suppose that all four roots of $U + \lambda V$ are equal; we shall find, as the most simple expressions of the most general forms of the two surfaces,

$$U = x^2 + xy + yz + zt, \quad V = axy + bz^2 + azt.$$

In this case the two points of intersection of the curve of the third degree, and the right line on which the surfaces intersect, come together, so that the right line becomes a tangent to the curve. The number of conditions to be satisfied is three: there is but one point of contact which may be considered as the union of two which have coalesced, and the species may be defined as confluent-unilinear contact.

If we throw the equations to the conoids having an unilinear contact into the form

$$x(x + y) + zt = 0, \quad xy + z(y + ct) = 0,$$

we obtain

$$(x + y)(y + ct) - yt = 0,$$

which last equation is no longer satisfied by $x = 0, z = 0$, these systems of roots having been made to disappear by the process of elimination.

The curve of the third degree, in which the two given conoids intersect, may thus be defined as their common intersection with the new conical surface defined by the third of the above equations.

p. 229

More generally, it is apparent that the three conoids,

$$\left. \begin{aligned} xu - yt &= 0, \\ yv - zu &= 0, \\ zt - xv &= 0 \end{aligned} \right\},$$

in which x, y, z, t, u, v may any of them be considered as a homogeneous linear function of four others, intersect in the same line of the third degree. Besides which, the first and second intersect in the right line y, u ; the second and third in z, v ; the third and first in x, t ; each of which lines it is evident is a chord of the common curve of intersection. For instance, $y = 0, u = 0$ may be satisfied concurrently with all the above three equations by satisfying the equation $zt - xv = 0$, which, as two linear relations exist originally between the six letters, and two more have been thrown in, becomes a quadratic equation between any two of the letters.

The only case of exception to this reasoning is, when $y = 0, u = 0$ can be satisfied concurrently with $z = 0, v = 0$, and with $x = 0, t = 0$; but in this case the surfaces all become cones; and as there is no longer a curve of the third degree, “*Cadit quaestio*.” Even here, however, the intersection of any two of the surfaces becomes a conic, and two coincident generating lines on the two cones; so that if we take one of these and the conic to represent a degenerate form of a line of the third degree, the remaining straight line passes through a double point of such degenerate form, and the case passes into that of confluent-unilinear contact.

The two double points in the intersection of the two conoids

$$U = x(x + y) + zt = 0, \quad V = xy + z(y + ct) = 0,$$

by which I mean the points of intersection of the conic with the right line common to them, are found by making $x = 0$, $z = 0$, and substituting in the derived equation

$$(x + y)(y + ct) - ty = 0,$$

which gives $y = 0$, or $y + (c - 1)t = 0$; so that the two points required are

$$x = 0, \quad y = 0, \quad z = 0,$$

$$x = 0, \quad y = (1 - c)t, \quad z = 0.$$

It appears also that the entire intersection is contained in each of the two cones,

$$U - V, \quad \text{that is,} \quad x^2 + z\{(1 - c)t - y\}$$

and

$$cU - V, \quad \text{that is,} \quad cx^2 + y\{(c - 1)x - z\},$$

the respective vertices of which are at the points above determined.

p. 230

The equations for confluent-unilinear contact,

$$x(x + y) + z(y + t) = 0,$$

$$xy + z(cz + t) = 0,$$

give

$$(x + y)(cz + t) - (y + t)y = 0;$$

which, on making $x = 0$, $z = 0$, is satisfied by $y^2 = 0$; showing that the confluence takes place at the point

$$x = 0, \quad y = 0, \quad z = 0.$$

The number of terms in the two equations for ordinary unilinear contact being six, and in those given for confluent unilinears seven, and the empirical rule in all other cases being that the terms tend to diminish and never increase in number as the degree of the contact (expressed by the number of conditions to be satisfied) rises, I am led to suspect that the conjugate system for the latter species of contact may admit of being reduced to some more simple form.

I must state here once for all, that all the distinct systems of (at least consecutive) conjugate forms that have been, and will be given, are mutually untransformable. This it is which distinguishes singular from particular forms.

A particular form is included in its primitive; but a singular form is one, which, while it responds to the same conditions as some other more general form, is incapable of being expressed as a particular case of the latter, on account of the additional condition or conditions which attach to it.

I pass now to the singularities which arise from the first minor determinants of $U + \lambda V$ having a factor in common, the second minors being supposed to be still without a common factor.

When this common factor is linear in respect to λ , let it be supposed to enter not more than twice (twice, we know, by the general principle enunciated at the commencement of this paper, it must enter) into the complete determinant.

Two of the cones containing the intersection of U and V then become coincident, and degenerate each into the same pair of crossing planes. This may be termed biplanar-contact. The characteristics of such contact are

$$U = x^2 + y^2 + z^2 + t^2, \quad V = ax^2 + ay^2 + bz^2 + ct^2;$$

the points of contact are two in number, being at the intersection of the two plane conics into which the curve of intersection breaks up. The two planes p. 231

in which these lie are given by the equation $(b - a)z^2 + (c - a)t^2 = 0$; these intersect in the right line $z = 0, t = 0$, which meets both surfaces in the same two points,

$$z = 0, \quad t = 0, \quad x + \sqrt{-1}y = 0,$$

$$z = 0, \quad t = 0, \quad x - \sqrt{-1}y = 0,$$

the two common tangent planes at these points being

$$x + \sqrt{-1}y = 0, \quad x - \sqrt{-1}y = 0$$

respectively.

This, then, is another species of double contact between two conoids, and, as far as I know, the only kind hitherto recognized as such. The number of conditions to be satisfied remains two, as in the former species.

Next suppose that the common factor of the first minor enters three times into the complete determinant instead of twice only, as in the last case.

The corresponding characteristics will be found to be

$$U = x^2 + zt + y^2 + z^2, \quad V = ax^2 + azt + by^2 + cz^2.$$

The intersection of U, V still lies in two planes,

$$(b - a)y^2 + (c - a)z^2 = 0;$$

but the intersection of these two planes,

$$y = 0, \quad z = 0,$$

meets the surfaces in the two coincident points,

$$y = 0, \quad z = 0, \quad x^2 = 0.$$

This, therefore, I call confluent-biplanar contact; the two conics constituting the complete intersection, instead of cutting, touch and at their point of contact the two conoids have a contact of a superior order. The conditions to be satisfied for this case are three in number.

Next suppose that the common factor of the first minors enters only twice into the complete determinant, but that the remaining two factors become equal.

Here the analytical characters of unilinear and biplanar contact are blended; in fact, the intersection consists of a conic and a pair of right lines meeting one another and the conic. The characteristics are

$$U = x^2 + y^2 + z^2 + zt, \quad V = ax^2 + ay^2 + bz^2 + czt.$$

p. 232

The intersection is contained in the two planes

$$z = 0, \quad (b - a)z + (c - a)t = 0,$$

and consists of the two lines $z = 0, x^2 + y^2 = 0$, lying in the common tangent plane $z = 0$, and the conic

$$\left. \begin{aligned} (b - a)z + (c - a)t &= 0, \\ (a - c)x^2 + (a - c)y^2 + (b - c)z^2 &= 0 \end{aligned} \right\}.$$

There are three points of contact, namely, the point $x = 0, y = 0, z = 0$, where the two right lines cut, and $x^2 + y^2 = 0, t = 0, z = 0$, where these lines meet the conic. This, then, is a case of triple contact. I distinguish it by the name of bilinear-contact. The number of conditions is still three.

Now all else remaining as before, let the two pairs of equal roots in the complete determinant become identical, or, in other words, let the common factor of the first minors be contained four times in the complete determinant. The characteristics become

$$U = xz + xt + y^2 + z^2, \quad V = axz + bxt + by^2 + bz^2.$$

The intersection becomes the two right lines

$$x = 0, \quad y^2 + z^2 = 0,$$

and the conic

$$z = 0, \quad x^2 + y^2 = 0.$$

All these meet in the same point,

$$x = 0, \quad y = 0, \quad z = 0;$$

so that instead of contact in three points, the contact takes place about one only, in which the three may be conceived as merging. This I call confluent-bilinear contact. It requires the satisfaction of four conditions.

Next let us suppose that the two distinct factors are common to each of the first minors. This will imply the existence of four affirmative conditions.

The complete determinant will of necessity contain each of these factors twice, so that no additional singularity can enter through this determinant. The characteristics assume the form

$$U = x^2 + y^2 + z^2 + t^2, \quad V = ax^2 + ay^2 + bz^2 + bt^2.$$

The two surfaces will meet in four straight lines, forming a wry quadrilateral, whose equations are

$$x \pm \sqrt{-1}y = 0, \quad z \pm \sqrt{-1}t = 0.$$

p. 233

These intersect each other in the four points

$$x = 0, \quad y = 0, \quad z^2 + t^2 = 0,$$

$$z = 0, \quad t = 0, \quad x^2 + y^2 = 0,$$

each of which will be a distinct point. This I term quadrilinear contact.

Now let the two factors common to each of the first minors become identical; so that a squared function, instead of an ordinary quadratic function of λ , is now their common measure.

The factor which enters twice into each of the first minors will enter four times into the complete determinant; the number of conditions to be satisfied is one more than in the preceding case, namely five, and the characteristics become

$$U = x^2 + y^2 + xz + yt, \quad V = ax^2 + by^2 + cxz + cyt.$$

Here arises a singularity of form in the intersections utterly unlike anything which has been remarked in the preceding cases. For it will not fail to have been observed, that the intersection in the nine preceding cases was always a line or system of lines of the fourth degree, so as to be cut by any plane in four points.

But in this case, the fact of the first minors having a factor in common, shows that the intersection is contained in two planes (which is of course to be viewed as a degenerate species of cone); and the fact of the complete determinant having all its roots equal, shows that there is but one system of a pair of planes in which the intersection is contained, and no more.

So that the two pairs of planes, into which the wry quadrilateral was divisible in the case immediately preceding, now become a single pair. This can only be explained by two of the opposite sides of the quadrilateral becoming indefinitely near to one another, but still not coinciding in the same planes; so that the actual visible or quasi-visible¹³¹ intersection will be in three right lines, of which the middle one meets each of the two others.

¹³¹I use the term quasi-visible, because the intersection may become in part or whole imaginary.

This will further appear by proceeding regularly to solve the equations

$$U = 0, \quad V = 0.$$

$V - cU = 0$ gives $y = \pm kx$, where $k = \sqrt{\frac{a-c}{c-b}}$, and therefore $xz + kxt = 0$, or $xz - kxt = 0$; whence we see that the complete intersection is represented by the lines

$$\begin{aligned} (x = 0, y = 0); \quad (z + kt = 0, y - kx = 0), \\ (x = 0, y = 0); \quad (z - kt = 0, y + kx = 0), \end{aligned}$$

showing that there are but three physically distinct lines, as already premised. p. 234

This, then, may be considered as derived from the preceding case of a wry quadrilateral intersection, by conceiving two opposite sides of the quadrilateral to come indefinitely near, but without coinciding.

Let these two lines be called P and P' ; take any point in P and any two points in P' indefinitely near to one another and the point first taken, then this indefinitely small plane will be common to both surfaces, and consequently they ought to touch along every point in the line P . This is again confirmed by the forms given to U and V . For at any point where the coordinates are $0, 0, \xi, \theta$ the equations to the tangent planes to the two surfaces respectively are

$$\xi x + \theta y = 0, \quad c\xi x + c\theta y = 0,$$

that is to say, are identical.

Whilst, therefore, certain grounds of geometrical, and still stronger grounds of analytical analogy, might seem to justify this species of contact taking the name of confluent quadrilinear, yet as, in fact, the intersection is trilinear, and as, moreover, the two indefinitely proximate lines must be considered, not as coincident, but as turned away from one another through an indefinitely small angle and out of the same plane, I prefer to take advantage of this striking property of contact at every point along a line (a property entirely distinct from any that we have yet considered), and confer upon the species of contact we have been considering the designation of unilinear-indefinite contact.

Where the line of indefinite contact meets the two other lines of the intersection, the contact is of course of a higher order; thus offering a parallel to what takes place in ordinary unilinear contact, in which there is no contact, except only at two points of the right line forming part of the complete intersection.

I believe that this kind of contact, which forms a natural family with two others about to be described, and which will close the list, has never before been imagined, and would at first sight have been rejected as impossible.

Having now exhausted the cases of the first class, in which the minors have no factor in common, and the two sections of the second class, in which the second minors have no common factor, but the first minors of $U + \lambda V$ a linear

or quadratic function of λ in common, I descend to the third class, in which the second minors, which are quadratic functions of λ , are supposed to have a common factor.

This common factor must enter twice into each of the first minors by virtue of the law previously indicated, and cannot enter more than twice, as p. 235

otherwise the first minors of $U + \lambda V$ could only differ from one another by a numerical multiplier, which is obviously impossible, except when $U + \lambda V$ is of the form $(k + \lambda)U$, that is, when the two surfaces coincide.

Again, the common factor of the first minor must enter three times into the complete determinant; but there is no reason why it may not enter four times, and thus two cases arise. In the first, the characteristics take the form

$$U = x^2 + y^2 + z^2 + t^2, \quad V = ax^2 + ay^2 + az^2 + bt^2.$$

The second determinant having a factor in common, shows that the intersection U, V is contained in a pair of coincident planes; but the complete determinant, having two distinct factors, evidences that these plane intersections, viewed as indefinitely near but still distinct, lie in the same cone, which will be a cone enveloping both the surfaces U and V all along their mutual intersections. This is also seen easily from the forms of U and V ; for we have $V - aU = (b - a)t^2$, which proves that the intersection lies in the coincident, or, to speak more strictly, consecutive planes $t^2 = 0$; and at any point $x = \xi, y = \eta, z = \zeta$, the tangent plane to each surface becomes

$$\xi x + \eta y + \zeta z = 0.$$

As there are six independent, that is, non-necessarily co-evanescent second minors, that the second minor systems shall all have a common factor, implies the satisfaction of five conditions. This species of contact I call curvilineo-indefinite; it is, I believe, the only kind of indefinite contact between two surfaces of the second order hitherto taken account of.

There is still, however, a higher species of contact, *videlicet*, when all the four roots of the complete determinant of $U + \lambda V$ are identical with the root common to each of its second minors. In this case the common enveloping cone becomes identical with the plane (considered as a coincident pair of planes) in which the surfaces intersect.

The characteristics take the form

$$U = x^2 + xy + zt, \quad V = xy + zt.$$

The intersection is contained completely in the common tangent plane $x = 0$, and consists of the two right lines,

$$(x = 0, \quad z = 0),$$

$$(x = 0, \quad t = 0).$$

This, the highest and crowning species of contact, I call bilineo-indefinite. It is defined by six conditions.

At each point of the two lines of intersection of U and V there is contact, and a very peculiar species of contact at the intersection of these two lines themselves. p. 236

To form a distinct idea of this, let the physical visible or quasi-visible intersection of U, V take place along the two lines L, M ; the rational intersection must be conceived as made up of the wry quadrilateral, $L, M; L', M'$, in which L is indefinitely near to L' , and M to M' . It follows, therefore, that there is contact at the four angles of the quadrilateral; but as there is nothing to fix the relative directions of the diagonal joining the intersection of L and M to that of L' and M' , because there is nothing to restrict the position of the latter point, except that it shall lie upon either surface¹³², it appears that not only is there contact at the junction of the two lines constituting the complete intersection of the two surfaces, but that these surfaces continue to touch at consecutive points taken all round this first, and indefinitely near to it in any direction¹³³.

Bilineo-indefinite (the highest) contact for two conoids is strictly analogous to confluence, the highest species of contact between conics. For this latter may be conceived as an intersection made up of two coincident pairs of coincident points; and the former, as an intersection made up of two coincident pairs of crossing right lines; and a pair of crossing lines is to a plane locus of the second degree what a coincident pair of points is to a rectilinear locus of the same degree.

In the subjoined table I have brought under one point of view the characters and algebraic forms which I call the condensed forms corresponding to each species of contact above detailed.

¹³²This will be better seen by reference to the analogy presented by the case when the two conoids touch all along a curve. The rational intersection is made up of this curve and another indefinitely near it. The two curves, whatever be the position of their node, will lie in the same enveloping cone, so that the position of the node is indeterminate.

¹³³As the two surfaces jut one close into the other at this point, it would perhaps be not improper to designate the contact at such point as umbilical.

A. *Quadratic loci in a right line.*

Simple contact.	One condition.	$\begin{cases} xy \\ x^2 + xy \end{cases}$
-----------------	----------------	--

B. *Quadratic loci in a plane.*

1st Class.

Simple contact.	One condition.	$\begin{cases} x^2 + y^2 + xz \\ ax^2 + by^2 + cxz \end{cases}$
Proximal contact.	Two conditions.	$\begin{cases} x^2 + yx + yz \\ ax^2 + byx + ayz \end{cases}$

2nd Class.

Diploidal contact.	Two conditions.	$\begin{cases} x^2 + y^2 + z^2 \\ ax^2 + ay^2 + bz^2 \end{cases}$
Confluent contact.	Three conditions.	$\begin{cases} x^2 + y^2 + xz \\ y^2 + xz \end{cases}$

p. 237

C. *Quadratic loci in space.*

1st Class.

Simple contact.	One condition.	$\begin{cases} x^2 + y^2 + z^2 + xt \\ ax^2 + by^2 + cz^2 + dxt \end{cases}$	
Proximal contact.	Two conditions.	$\begin{cases} x^2 + y^2 + xt + zt \\ ax^2 + by^2 + cxt + azt \end{cases}$	
Unilinear contact. 1st species of diploidal.	Two conditions.	$\begin{cases} x^2 + xy + zt \\ ayz + bxy + czt \end{cases}$	
Confluent-unilinear, or triple contact.	Three conditions.	$\begin{cases} x^2 + yz + xy + zt \\ az^2 + bxy + bzt \end{cases}$	
2nd Class, 1st Section.			
Biplanar contact. 2nd species of diploidal.	Two conditions.	$\begin{cases} x^2 + y^2 + z^2 + t^2 \\ ax^2 + ay^2 + bz^2 + ct^2 \end{cases}$	
Confluent-biplanar contact.	Three conditions.	$\begin{cases} x^2 + zt + y^2 + z^2 \\ ax^2 + azt + by^2 + cz^2 \end{cases}$	
Bilinear contact.	Three conditions.	$\begin{cases} x^2 + y^2 + z^2 + zt \\ ax^2 + ay^2 + bz^2 + czt \end{cases}$	or $\begin{cases} xz + yt \\ axt + byz \end{cases}$
Confluent-bilinear contact.	Four conditions.	$\begin{cases} xz + xt + y^2 + z^2 \\ axz + bxt + by^2 + bz^2 \end{cases}$	

2nd Class, 2nd Section.

Quadrilinear, or quadruple contact.	Four conditions.	$\begin{cases} x^2 + y^2 + z^2 + t^2 \\ ax^2 + ay^2 + bz^2 + bt^2 \end{cases}$	or $\begin{cases} xy + zt \\ axy + bzt \end{cases}$
Unilineo-indefinite contact.	Five conditions.	$\begin{cases} x^2 + y^2 + xz + yt \\ ax^2 + by^2 + cxz + cyt \end{cases}$	

3rd Class.

Curvilineo-indefinite contact.	Five conditions.	$\begin{cases} x^2 + y^2 + z^2 + t^2 \\ ax^2 + ay^2 + az^2 + bt^2 \end{cases}$	
Bilineo-indefinite contact.	Six conditions.	$\begin{cases} x^2 + xy + zt \\ xy + zt \end{cases}$	

p. 238

Another (and, in a physical sense, more) natural mode of grouping the twelve species of conoidal contact, which, without observing the same lines of demarcation, leaves intact the sequence of the species, is into the three families. The first, or definite-continuous, for which the surfaces touch in a single point, and intersect in an unbroken curve, comprises simple and cuspidal contact.

The second definite-discontinuous, for which the surfaces touch in one, two, three or four points, but intersect in a curve more or less broken up into distinct parts, comprises all the species from the third to the ninth inclusive. The third natural family is that of indefinite contact, and comprises the three last species. It will of course be observed that there are five species of single contact, that is, contact at one point, namely, simple, cuspidal, and the three confluent species, two of double, one of treble, one of quadruple, and three of indefinite contact; the last being distinguishable *inter se* lineo-indefinite as being special at two points, curvilineo-indefinite as having no speciality, and bilineo-indefinite as being special at one point only.

I might now proceed to discuss more particularly the nature of the contact taken, not collectively, but with reference to each single point where it exists. This, however, must be reserved for a future communication; as also, among other important and curious matter, the ascertainment of the singular forms of quadratic conjugate functions of five or more letters. At present I shall content myself with stating the following general proposition, which naturally suggests itself from a consideration of the cases already considered.

In a conjugate quadratic system of any number of letters, the lowest and also the highest degree of singularity will be always unique; the conditions to be satisfied in the former case being only one in number, and, in the latter $\frac{1}{2}r(r-1)$, where r denotes the number of the letters. The first part of this proposition is self-apparent, the latter part may be inferred from the homaloidal law; for the $(r-2)$ nd minors will be quadratic functions, and the highest degree of contact will correspond to those having a factor in common, which would involve the satisfaction of $\frac{1}{2}r(r-1) - 1$ conditions only; but over and above this, that the complete determinant, instead of containing this common factor, as it needs must, $(r-1)$ times, shall contain it r times: this gives one condition more, making up the entire number to $\frac{1}{2}r(r-1)$.

The total number of different species of singularity for conjugate functions of a given number of letters can only be expressed by aid of formulae containing expressions for the number of various ways in which numbers admit of being broken up into a given number of parts.

The computation of this number in particular cases, upon the principle of the foregoing method, is attended with no difficulty. p. 239

We have seen that this number for two, three and four letters, is respectively one, four, twelve.

I have found that for five letters the number is twenty-four, for six letters fifty,

for seven letters a hundred, and (subject to further examination) for eight letters one hundred and ninety-three. The series, therefore, as far as I have yet traced it, is 1, 4, 12, 24, 50, 100, 193. The last number must not be relied upon at present.

It will be observed, that the foregoing table for the contacts of surfaces of the second order contains no form corresponding to a complete intersection in two non-intersecting lines and an undegenerated conic. In fact, if two such lines form part of the intersection, at least one other right line intersecting them both, must go to make up the remaining part. This is easily verified; for it is readily seen that the most general representation of two conoids intersecting in two non-meeting lines will be

$$U = xy + zt, \quad V = axy + bzt + cxt + eyz,$$

where the two lines in question are

$$(x = 0, \quad z = 0), \quad (y = 0, \quad t = 0).$$

Now it will be found that the first minors of $V + \lambda U$ formed from the above equation will all contain the common factor $(a + \lambda)(b + \lambda) - ce$, showing that the contact is quadrilinear or linear-indefinite, that is bilinear, according as the roots of

$$\lambda^2 + (a + b)\lambda + ab - ce = 0$$

are distinct or equal; which explains how it is that only one species of bilinear contact (that is to say, the case corresponding to the two conoids agreeing in the two right lines in which each is cut by a common tangent plane) comes to find a place in the preceding enumeration.

It may not be uninteresting, under an euristic point of view, to state that the above theory, which, as well in what it accomplishes as in what it suggests (the author cannot but feel conscious), constitutes a substantial accession to analytical science, arose out of a theorem which occurred to him as likely to be true, in the act of reviewing for the press his paper "On Certain Additions" in the last November Number¹³⁴ of this Magazine, and which he had only then time to throw into a foot-note as a probable conjecture.

Wishing to subject it to an analytical test, he found it necessary to obtain the condensed forms which serve to characterize the confluent contact of p. 240

conies. In this way he became aware of the great utility of these condensed forms, and of the desideratum to be supplied in obtaining a complete list of them applicable to all varieties of contact. The happy thought then occurred to him of inverting the process which he had applied in the treatment of the contacts of conies, in the November Number¹³⁵ of the *Cambridge and Dublin Mathematical Journal*; for whereas the nature of the contacts was there assumed and translated

¹³⁴p. 148 above.

¹³⁵p. 119 above.

into the language of determinants, he soon discovered that it was the more easy and secure course to assume the relations of every possible immutable kind that could exist between the complete and minor determinants corresponding to the characteristics, by aid of these relations to construct the characteristics, and from the characteristics so obtained, determine the geometrical character of each resulting species of contact. Thus he has been able to effect the very results stated by himself as desiderata at the close of the paper in this Magazine above referred to.

NOTE.

It is proper to remark, that all the condensed forms given in this paper have actually been obtained by the author in the way above pointed out. The limits imposed by the objects to which the Magazine is devoted have restricted him from exhibiting the method at full; but any of his readers will be able without difficulty to make it out for himself.

The process consists in finding $U + \lambda V$ by means of solving for each case a problem of position (a kind of chess-board problem) on a square table, containing three places in length and breadth for conies, four places by four for surfaces, and so on (if need be) according to the number of variable letters involved. $U + \lambda V$ being thus determined in form, U and V become readily cognizable. It is right also to add, that some of the condensed forms here set forth have been incidentally noticed and employed by previous authors, as Plücker and Mr Cayley.

The conditions in each case to which the position-problem is subject are immediately deducible from the laws which the complete determinant, and the successive minor systems of determinants of $U + \lambda V$, are required to satisfy.

37.

ON THE RELATION BETWEEN THE MINOR DETERMINANTS OF LINEARLY EQUIVALENT QUADRATIC FUNCTIONS

[*Philosophical Magazine, I. (1851), pp. 295–305*]

p. 241

I showed in the preliminary part of my paper on Contacts in the February Number of this *Magazine*¹³⁶, by *a priori* reasoning, that if a quadratic function (U) be linearly converted into another (V), any minor determinant of any order of V must be a syzygetic function of all the minor determinants of U of the same order.

The object of my present communication is to exhibit the syzygy in question, which, as I indicated, is linear; by which I mean that a determinant of the one function is equal to the sum of the pari-ordinal determinants of the other affected respectively with multipliers formed exclusively out of the coefficients of the equations of transformation. In order that a clear enunciation of the theorem in view may be possible, it is necessary to premise a new but simple, and, as experience has proved to me, a most powerful, because natural, method of notation applicable to all questions concerning determinants.

Every determinant is obtained by operating upon a square array of quantities, which, according to the ordinary method, might be denoted as follows:

$$\begin{array}{cccc} a_{1,1}, & a_{1,2} & \cdots & a_{1,n}, \\ a_{2,1}, & a_{2,2} & \cdots & a_{2,n}, \\ a_{3,1}, & a_{3,2} & \cdots & a_{3,n}, \\ \cdots & \cdots & & \cdots \\ a_{n,1}, & a_{n,2} & \cdots & a_{n,n}. \end{array}$$

My method consists in expressing the same quantities biliterally as below:

$$\begin{array}{cccc} a_1\alpha_1, & a_1\alpha_2 & \cdots & a_1\alpha_n, \\ a_2\alpha_1, & a_2\alpha_2 & \cdots & a_2\alpha_n, \\ \cdots & \cdots & & \cdots \\ a_n\alpha_1, & a_n\alpha_2 & \cdots & a_n\alpha_n. \end{array}$$

p. 242

where of course, whenever desirable, instead of $a_1, a_2 \dots a_n$ and $\alpha_1, \alpha_2 \dots \alpha_n$, we may write simply $a, b \dots l$, and $\alpha, \beta \dots \lambda$ respectively. Each quantity is now represented by two letters; the letters themselves, taken separately, being symbols neither of quantity nor of operation, but mere umbrae or ideal elements of quantitative symbols. We have now a means of representing the determinant above given in a compact form; for this purpose we need but to write one set of

¹³⁶p. 221 above.

umbræ over the other as follows: $\left\{ \begin{matrix} a_1, a_2 \dots a_n \\ \alpha_1, \alpha_2 \dots \alpha_n \end{matrix} \right\}$. If we now wish to obtain the algebraic value of this determinant, it is only necessary to take $a_1, a_2 \dots a_n$ in all its $1, 2, 3 \dots n$ different positions, and we shall have

$$\left\{ \begin{matrix} a_1, a_2 \dots a_n \\ \alpha_1, \alpha_2 \dots \alpha_n \end{matrix} \right\} = \sum \pm \{a_1 \alpha_{\theta_1} \times a_2 \alpha_{\theta_2} \times \dots \times a_n \alpha_{\theta_n}\},$$

in which expression $\theta_1, \theta_2 \dots \theta_n$ represents some order of the numbers $1, 2 \dots n$, and the positive or negative sign is to be taken according to the well-known dichotomous law. Thus, for example,

$$\left\{ \begin{matrix} abc \\ \alpha\beta\gamma \end{matrix} \right\} \quad \text{will represent} \quad \begin{cases} aa \times b\beta \times c\gamma \\ + a\beta \times b\gamma \times ca \\ + a\gamma \times ba \times c\beta \\ - a\beta \times ba \times c\gamma \\ - aa \times b\gamma \times c\beta \\ - a\gamma \times b\beta \times ca. \end{cases}$$

Although not necessary for our immediate object, it may not be inopportune to observe how readily this notation lends itself to a further natural extension of its application.

$$\left\{ \begin{matrix} \overline{ab} & \overline{cd} \\ \alpha\beta & \gamma\delta \end{matrix} \right\} \quad \text{will naturally denote} \quad \left\{ \begin{matrix} ab \\ \alpha\beta \end{matrix} \right\} \times \left\{ \begin{matrix} cd \\ \gamma\delta \end{matrix} \right\} - \left\{ \begin{matrix} ab \\ \gamma\delta \end{matrix} \right\} \times \left\{ \begin{matrix} cd \\ \alpha\beta \end{matrix} \right\};$$

that is

$$\left\{ \begin{matrix} aa \times b\beta \\ -(a\beta \times ba) \end{matrix} \right\} \times \left\{ \begin{matrix} c\gamma \times d\delta \\ -(c\delta \times d\gamma) \end{matrix} \right\} - \left\{ \begin{matrix} a\gamma \times b\delta \\ -(a\delta \times b\gamma) \end{matrix} \right\} \times \left\{ \begin{matrix} ca \times d\beta \\ -(c\beta \times da) \end{matrix} \right\}.$$

And in general the compound determinant

$$\overline{\left\{ \begin{matrix} a_1, b_1 \dots l_1, & a_2, b_2 \dots l_2 & \dots & a_r, b_r \dots l_r \\ \alpha_1, \beta_1 \dots \lambda_1, & \alpha_2, \beta_2 \dots \lambda_2 & \dots & \alpha_r, \beta_r \dots \lambda_r \end{matrix} \right\}}$$

will denote

$$\sum \pm \left\{ \begin{matrix} a_1, b_1 \dots l_1 \\ a_{\theta_1}, \beta_{\theta_1} \dots \lambda_{\theta_1} \end{matrix} \right\} \times \left\{ \begin{matrix} a_2, b_2 \dots l_2 \\ a_{\theta_2}, \beta_{\theta_2} \dots \lambda_{\theta_2} \end{matrix} \right\} \times \dots \times \left\{ \begin{matrix} a_r, b_r \dots l_r \\ a_{\theta_r}, \beta_{\theta_r} \dots \lambda_{\theta_r} \end{matrix} \right\},$$

where, as before, we have the disjunctive equation

$$\theta_1, \theta_2 \dots \theta_r = 1, 2 \dots r.$$

As an example of the power of this notation, I will content myself with stating the following remarkable theorem in compound determinants, one of the most prolific in results of any with which I am acquainted, but which is derived from a more particular case of another vastly more general. The theorem is contained in the annexed equation

$$\overline{\left\{ \begin{array}{ll} a_1, a_2 \dots a_r, a_{r+1}; & a_1, a_2 \dots a_r, a_{r+2} \dots a_1, a_2 \dots a_r, a_{r+s} \\ \alpha_1, \alpha_2 \dots \alpha_r, \alpha_{r+1}; & \alpha_1, \alpha_2 \dots \alpha_r, \alpha_{r+2} \dots \alpha_1, \alpha_2 \dots \alpha_r, \alpha_{r+s} \end{array} \right\}} \quad (1)$$

$$= \left\{ \begin{array}{l} a_1, a_2 \dots a_r \\ \alpha_1, \alpha_2 \dots \alpha_r \end{array} \right\}^{s-1} \times \left\{ \begin{array}{l} a_1, a_2 \dots a_r, a_{r+1}, a_{r+2} \dots a_{r+s} \\ \alpha_1, \alpha_2 \dots \alpha_r, \alpha_{r+1}, \alpha_{r+2} \dots \alpha_{r+s} \end{array} \right\}.$$

It is obvious, that, without the aid of my system of umbral or biliteral notation, this important theorem could not be made the subject of statement without an enormous periphrasis, and could never have been made the object of distinct contemplation or proof.

To return to the more immediate object of this communication, suppose that we have any binary function of two sets of quantities, $x_1, x_2 \dots x_n$; $\xi_1, \xi_2 \dots \xi_n$, of which the general term will be of the form $c_{r,s} \times x_r \xi_s$; according to the principles of notation above laid down, nothing can be more natural than to represent $c_{r,s}$ by the biliteral group $a_r \alpha_s$; the function in question will then take the form

$$\sum a_r \alpha_s \cdot x_r \xi_s;$$

the x 's and ξ 's denoting quantities, but the a 's and α 's mere umbrae. The function may then be thrown under the convenient symbolical form

$$(a_1 x_1 + a_2 x_2 + \dots + a_n x_n)(\alpha_1 \xi_1 + \alpha_2 \xi_2 + \dots + \alpha_n \xi_n).$$

So if we confine ourselves to quadratic functions, for which $x_1, x_2 \dots x_n$; $\xi_1, \xi_2 \dots \xi_n$ become respectively identical, the general symbolical representation of any such will be

$$(a_1 x_1 + a_2 x_2 + \dots + a_n x_n)^2.$$

The complete determinant will be denoted by

$$\left\{ \begin{array}{l} a_1, a_2 \dots a_n \\ \alpha_1, \alpha_2 \dots \alpha_n \end{array} \right\},$$

and any minor determinant of the r th order by

$$\left\{ \begin{array}{l} a_1, a_2 \dots a_r \\ a_{\theta_1}, a_{\theta_2} \dots a_{\theta_r} \end{array} \right\},$$

where $\theta_1, \theta_2 \dots \theta_r$ are some certain r distinct numbers taken out of the series $1, 2, 3 \dots r$. Suppose now that we have

$$U = (a_1x_1 + a_2x_2 + \dots + a_nx_n)^2$$

linearly transformable into

$$V = (b_1y_1 + b_2y_2 + \dots + b_ny_n)^2,$$

by means of the n equations

$$x_1 = a_1b_1 \cdot y_1 + a_1b_2 \cdot y_2 + \dots + a_1b_n \cdot y_n,$$

$$x_2 = a_2b_1 \cdot y_1 + a_2b_2 \cdot y_2 + \dots + a_2b_n \cdot y_n,$$

$$\dots\dots\dots$$

$$x_n = a_nb_1 \cdot y_1 + a_nb_2 \cdot y_2 + \dots + a_nb_n \cdot y_n,$$

in which equations, be it observed, each coefficient a_rb_s is a single quantity, perfectly independent of the quantities denoted generally by a_ra_s , b_rb_s , which enter into U and V . Our object is to be able to express the minor determinant

$$\left\{ \begin{matrix} b_{k_1}, b_{k_2} \dots b_{k_r} \\ b_{l_1}, b_{l_2} \dots b_{l_r} \end{matrix} \right\},$$

in which the one group of distinct numbers $k_1, k_2 \dots k_r$ may either differ wholly from, or agree wholly or in part with the other group of distinct numbers $l_1, l_2 \dots l_r$, under the form of

$$\sum \left\{ \left\{ \begin{matrix} a_{\theta_1}, a_{\theta_2} \dots a_{\theta_r} \\ b_{\phi_1}, b_{\phi_2} \dots b_{\phi_r} \end{matrix} \right\} \times Q \right\}.$$

The particular value of Q corresponding to each double group, $\left(\begin{matrix} \theta_1, \theta_2 \dots \theta_r \\ \phi_1, \phi_2 \dots \phi_r \end{matrix} \right)$, may be denoted by $Q \left(\begin{matrix} \theta_1, \theta_2 \dots \theta_r \\ \phi_1, \phi_2 \dots \phi_r \end{matrix} \right)$; so that our problem consists in determining the value of $Q \left(\begin{matrix} \theta_1, \theta_2 \dots \theta_r \\ \phi_1, \phi_2 \dots \phi_r \end{matrix} \right)$ in the equation

$$\left\{ \begin{matrix} b_{k_1}, b_{k_2} \dots b_{k_r} \\ b_{l_1}, b_{l_2} \dots b_{l_r} \end{matrix} \right\} = \sum \left\{ Q \left(\begin{matrix} \theta_1, \theta_2 \dots \theta_r \\ \phi_1, \phi_2 \dots \phi_r \end{matrix} \right) \times \left\{ \begin{matrix} a_{\theta_1}, a_{\theta_2} \dots a_{\theta_r} \\ a_{\phi_1}, a_{\phi_2} \dots a_{\phi_r} \end{matrix} \right\} \right\}.$$

Accordingly I enunciate that

$$\begin{aligned} Q \left(\begin{matrix} \theta_1, \theta_2 \dots \theta_r \\ \phi_1, \phi_2 \dots \phi_r \end{matrix} \right) &= \left\{ \begin{matrix} a_{k_1}, a_{k_2} \dots a_{k_r} \\ b_{\theta_1}, b_{\theta_2} \dots b_{\theta_r} \end{matrix} \right\} \times \left\{ \begin{matrix} a_{l_1}, a_{l_2} \dots a_{l_r} \\ b_{\phi_1}, b_{\phi_2} \dots b_{\phi_r} \end{matrix} \right\} \\ &+ \left\{ \begin{matrix} a_{l_1}, a_{l_2} \dots a_{l_r} \\ b_{\theta_1}, b_{\theta_2} \dots b_{\theta_r} \end{matrix} \right\} \times \left\{ \begin{matrix} a_{k_1}, a_{k_2} \dots a_{k_r} \\ b_{\phi_1}, b_{\phi_2} \dots b_{\phi_r} \end{matrix} \right\}, \end{aligned} \tag{2}$$

subject to one sole exception in the case of $\theta_1, \theta_2 \dots \theta_r$ being identical with $\phi_1, \phi_2 \dots \phi_r$; namely, that for the terms (for such case) of the form p. 245

$$Q \begin{pmatrix} \theta_1, \theta_2 \dots \theta_r \\ \theta_1, \theta_2 \dots \theta_r \end{pmatrix},$$

the value to be taken is not that which the general formula would give, namely,

$$2 \left\{ \frac{a_{k_1}, a_{k_2} \dots a_{k_r}}{b_{\theta_1}, b_{\theta_2} \dots b_{\theta_r}} \right\}^2,$$

but the half of this, that is simply the square of

$$\left\{ \frac{a_{k_1}, a_{k_2} \dots a_{k_r}}{a_{\theta_1}, a_{\theta_2} \dots a_{\theta_r}} \right\}.$$

The value of $Q \begin{pmatrix} \theta_1, \theta_2 \dots \theta_r \\ \phi_1, \phi_2 \dots \phi_r \end{pmatrix}$, it is obvious, contains only quantities of the form $a_r \cdot b_s$, which are coefficients in the equations of transformation, but none of the form $a_r \cdot a_s$ or $b_r \cdot b_s$; showing that the syzygetic connexion between the minor determinants of U and V of the same order is linear, as has been already anticipatively announced.

The problem which I have treated above is only a particular case of a more general one, which may be stated as follows: given

$$U = (a_1x_1 + a_2x_2 + \dots + a_nx_n)^2,$$

and supposing m linear equations to be instituted between $x_1, x_2 \dots x_n$, so that U may be made a function of $(n - m)$ letters only, to express any minor determinant of the reduced form of U without performing the process of elimination between the given equations. Let the given equations be written under the form

$$\begin{aligned} a_1a_{n+1}x_1 + a_2a_{n+1}x_2 + \dots + a_na_{n+1}x_n &= 0, \\ a_1a_{n+2}x_1 + a_2a_{n+2}x_2 + \dots + a_na_{n+2}x_n &= 0, \\ &\dots\dots\dots \\ a_1a_{n+m}x_1 + a_2a_{n+m}x_2 + \dots + a_na_{n+m}x_n &= 0, \end{aligned}$$

and let it be convened (which takes nothing away from the generality of these equations) that $a_{n+r}a_{n+s}$ shall signify zero for all values of r and s concurrently greater than zero. Suppose that $x_1, x_2 \dots x_m$, being eliminated, U becomes of the form

$$(b_{m+1}x_{m+1} + b_{m+2}x_{m+2} + \dots + b_nx_n)^2;$$

and suppose that we wish to determine the value of the complete determinant of this last function; it will be found to be

$$\left\{ \frac{b_{m+1}, b_{m+2} \dots b_n}{b_{m+1}, b_{m+2} \dots b_n} \right\} = \left\{ \frac{a_1, a_2 \dots a_n, a_{n+1} \dots a_{n+m}}{a_1, a_2 \dots a_n, a_{n+1} \dots a_{n+m}} \right\} \div \left\{ \frac{a_1, a_2 \dots a_m}{a_{n+1}, a_{n+2} \dots a_{n+m}} \right\}^2,$$

the squared divisor being, as is obvious, a function only of the coefficients of the transforming equations, and depending for its value upon the particular p. 246

m quantities selected for elimination. The dividend, on the contrary, is independent of this selection, but involves the coefficients of the function combined with the coefficients of transformation. This is the symbolical representation of the theorem given by me in the postscript to my paper in the *Cambridge and Dublin Mathematical Journal* for November 1850¹³⁷.

Suppose, now, more generally that we wish to find any minor determinant. The solution is given¹³⁸ by the equation

$$\left\{ \begin{matrix} b_{\theta_{m+1}}, b_{\theta_{m+2}} \dots b_{\theta_{m+s}} \\ b_{\phi_{m+1}}, b_{\phi_{m+2}} \dots b_{\phi_{m+s}} \end{matrix} \right\} = \left\{ \begin{matrix} a_{\theta_1}, a_{\theta_2} \dots a_{\theta_n}, a_{\theta_{m+1}}, a_{\theta_{m+2}} \dots a_{\theta_{m+s}} \\ a_{\phi_1}, a_{\phi_2} \dots a_{\phi_m}, a_{\phi_{m+1}}, a_{\phi_{m+2}} \dots a_{\phi_{m+s}} \end{matrix} \right\} \div \left\{ \begin{matrix} a_1, a_2 \dots a_m \\ a_{n+1}, a_{n+2} \dots a_{n+m} \end{matrix} \right\} \quad (3)$$

If we make $n = 2\gamma$ and $m = \gamma$, and $a_{\gamma+r}a_{\gamma+s} = 0$ for all positive values of either r or s , and $a_{\gamma-i}a_{n+e} = 0$ for all values of i and e differing from one another, and for equal values $a_{\gamma-e}a_{\gamma+e} = -1$, it will readily be seen that this last theorem reduces to the one first considered; and on careful inspection it will be found, that the solution given of the general question includes within it that presented for the particular case in question. Such inclusion, however, I ought in fairness to state is far from being obvious; and to demonstrate it exactly, and in general terms, requires the aid of methods which my readers would probably find to exceed their existing degree of knowledge or familiarity with the subject.

The theorem above enunciated was in part suggested in the course of a conversation with Mr Cayley (to whom I am indebted for my restoration to the enjoyment of mathematical life) on the subject of one of the preliminary theorems in my paper on Contacts in this *Magazine*.

It is wonderful that a theory so purely analytical should originate in a geometrical speculation. My friend M. Hermite has pointed out to me, that some faint indications of the same theory may be found in the *Recherches Arithmétiques* of Gauss. The notation which I have employed for determinants is very similar to that of Vandermonde, with which I have become acquainted since writing the above, in Mr Spottiswoode's valuable treatise *On the Elementary Theorems of Determinants*. Vandermonde was evidently on the right road. I do not hesitate to affirm, that the superiority of his and my notation over that in use in the ordinary methods is as great and almost as important to the progress of analysis, as the superiority of the notation of the differential calculus over that of the fluxional system. For what is the theory of determinants? It is an algebra upon algebra; a p. 247

calculus which enables us to combine and foretell the results of algebraical operations, in the same way as algebra itself enables us to dispense with the

¹³⁷p. 136 above.

¹³⁸see p. 251 below.

performance of the special operations of arithmetic. All analysis must ultimately clothe itself under this form¹³⁹.

I have in previous papers defined a “Matrix” as a rectangular array of terms, out of which different systems of determinants may be engendered, as from the womb of a common parent; these cognate determinants being by no means isolated in their relations to one another, but subject to certain simple laws of mutual dependence and simultaneous deperition. The condensed representation of any such Matrix, according to my improved Vandermondian notation, will be

$$\left\{ \begin{array}{c} a_1, a_2 \dots a_n \\ a_1, a_2 \dots a_m \end{array} \right\}.$$

To return to the theorems of the text. Theorem (2) admits of being presented in a more convenient form for the purposes of analytical operation, so as to become relieved from all cases of exception appertaining to particular terms.

The limitation to the generality of the expression for Q arises from our treating

$$\left\{ \begin{array}{c} a_{\theta_1}, a_{\theta_2} \dots a_{\theta_r} \\ a_{\phi_1}, a_{\phi_2} \dots a_{\phi_r} \end{array} \right\}$$

as identical with its equal,

$$\left\{ \begin{array}{c} a_{\phi_1}, a_{\phi_2} \dots a_{\phi_r} \\ a_{\theta_1}, a_{\theta_2} \dots a_{\theta_r} \end{array} \right\}.$$

If, however, we now convene to treat these two forms as distinct, so that in theorem (2)

$$\sum \left\{ Q \left(\begin{array}{c} \theta_1, \theta_2 \dots \theta_r \\ \phi_1, \phi_2 \dots \phi_r \end{array} \right) \times \left\{ \begin{array}{c} a_{\theta_1}, a_{\theta_2} \dots a_{\theta_r} \\ a_{\phi_1}, a_{\phi_2} \dots a_{\phi_r} \end{array} \right\} \right\}$$

will contain

$$\left\{ \frac{n(n-1) \dots (n-r+1)}{1 \cdot 2 \dots r} \right\}^2$$

terms, then we may write simply

$$Q \left(\begin{array}{c} \theta_1, \theta_2 \dots \theta_r \\ \phi_1, \phi_2 \dots \phi_r \end{array} \right) = \left\{ \begin{array}{c} a_{k_1}, a_{k_2} \dots a_{k_r} \\ b_{\theta_1}, b_{\theta_2} \dots b_{\theta_r} \end{array} \right\} \times \left\{ \begin{array}{c} a_{l_1}, a_{l_2} \dots a_{l_r} \\ b_{\phi_1}, b_{\phi_2} \dots b_{\phi_r} \end{array} \right\},$$

p. 248

¹³⁹Perhaps the most remarkable indirect question to which the method of determinants has been hitherto applied is Hesse’s problem of reducing a cubic function of 3 letters to another consisting only of 4 terms by linear substitutions—a problem which appears to set at defiance all the processes and artifices of common algebra. I have succeeded in applying a method founded upon this calculus to the linear reduction of a biquadratic function of two letters to Cayley’s form $x^4 + mx^2y^2 + y^4$, and of a 5^c function of two letters to the new form $x^5 + y^5 + (ax + by)^5$. This last reduction is effected by means of the properties of a certain other function of the 8th degree connected with the given function of the 5th degree. See a paper on this subject in the forthcoming May Number of the *Cambridge and Dublin Mathematical Journal*. [p. 191 above.]

which equation is subject to no exception for the case of the θ 's and ϕ 's becoming identical. As regards this theorem, it will not fail to strike the reader that it ought to admit of verification; for that U may be derived from V in the same manner as V from U if we express $y_1, y_2 \dots y_n$ in terms of $x_1, x_2 \dots x_n$, by solving the system of equations (2), which there is no difficulty in doing. In fact, if we write

$$\begin{aligned} y_1 &= \alpha_1 \beta_1 x_1 + \alpha_1 \beta_2 x_2 + \dots + \alpha_1 \beta_n x_n, \\ y_2 &= \alpha_2 \beta_1 x_1 + \alpha_2 \beta_2 x_2 + \dots + \alpha_2 \beta_n x_n, \\ &\dots\dots\dots \\ y_n &= \alpha_n \beta_1 x_1 + \alpha_n \beta_2 x_2 + \dots + \alpha_n \beta_n x_n, \end{aligned}$$

we shall obtain

$$\alpha_r \beta_s = \left\{ \begin{matrix} a_1, a_2 \dots a_{r-1}, a_{r+1}, a_{r+2} \dots a_n \\ b_1, b_2 \dots b_{s-1}, b_{s+1}, b_{s+2} \dots b_n \end{matrix} \right\} \div \left\{ \begin{matrix} a_1, a_2 \dots a_n \\ b_1, b_2 \dots b_n \end{matrix} \right\}.$$

Accordingly we shall find

$$\left\{ \begin{matrix} a_{m_1}, a_{m_2} \dots a_{m_r} \\ a_{p_1}, a_{p_2} \dots a_{p_r} \end{matrix} \right\} = \sum \left\{ Q \left(\begin{matrix} \psi_1, \psi_2 \dots \psi_r \\ \omega_1, \omega_2 \dots \omega_r \end{matrix} \right) \times \left\{ \begin{matrix} b_{\psi_1}, b_{\psi_2} \dots b_{\psi_r} \\ b_{\omega_1}, b_{\omega_2} \dots b_{\omega_r} \end{matrix} \right\} \right\},$$

and

$$Q \left(\begin{matrix} \psi_1, \psi_2 \dots \psi_r \\ \omega_1, \omega_2 \dots \omega_r \end{matrix} \right) = \left\{ \begin{matrix} a_{m_1}, a_{m_2} \dots a_{m_r} \\ \beta_{\psi_1}, \beta_{\psi_2} \dots \beta_{\psi_r} \end{matrix} \right\} \times \left\{ \begin{matrix} a_{p_1}, a_{p_2} \dots a_{p_r} \\ \beta_{\omega_1}, \beta_{\omega_2} \dots \beta_{\omega_r} \end{matrix} \right\};$$

substituting for the α 's and β 's their symbolical equivalents given above, and applying the theorem given below, we shall easily obtain

$$\begin{aligned} Q \left(\begin{matrix} \psi_1, \psi_2 \dots \psi_r \\ \omega_1, \omega_2 \dots \omega_r \end{matrix} \right) &= \left\{ \begin{matrix} a_{m+1}, a_{m+2} \dots a_{m_r} \\ b_{\psi_{r+1}}, b_{\psi_{r+2}} \dots b_{\psi_n} \end{matrix} \right\} \times \left\{ \begin{matrix} a_{p_{r+1}}, a_{p_{r+2}} \dots a_n \\ \beta_{p_{r+1}}, \beta_{p_{r+2}} \dots \beta_{p_n} \end{matrix} \right\} \\ &\div \left\{ \begin{matrix} a_1, a_2 \dots a_n \\ b_1, b_2 \dots b_n \end{matrix} \right\}^2. \end{aligned}$$

If, now, in the expression

$$\left\{ \begin{matrix} b_{k_1}, b_{k_2} \dots b_{k_r} \\ b_{l_1}, b_{l_2} \dots b_{l_r} \end{matrix} \right\} = \sum \left\{ \left\{ \begin{matrix} a_{k_1}, a_{k_2} \dots a_{k_r} \\ b_{\theta_1}, b_{\theta_2} \dots b_{\theta_r} \end{matrix} \right\} \left\{ \begin{matrix} a_{l_1}, a_{l_2} \dots a_{l_r} \\ b_{\phi_1}, b_{\phi_2} \dots b_{\phi_r} \end{matrix} \right\} \left\{ \begin{matrix} a_{\theta_1}, a_{\theta_2} \dots a_{\theta_r} \\ a_{\phi_1}, a_{\phi_2} \dots a_{\phi_r} \end{matrix} \right\} \right\},$$

we resubstitute for $\left\{ \begin{matrix} a_{\theta_1}, a_{\theta_2} \dots a_{\theta_r} \\ a_{\phi_1}, a_{\phi_2} \dots a_{\phi_r} \end{matrix} \right\}$ its value in the form of

$$\sum \left\{ \left\{ \begin{matrix} b_{\omega_1}, b_{\omega_2} \dots b_{\omega_r} \\ b_{\psi_1}, b_{\psi_2} \dots b_{\psi_r} \end{matrix} \right\} Q \right\},$$

we shall obtain $\left\{ \begin{matrix} b_{k_1}, b_{k_2} \dots b_{k_r} \\ b_{l_1}, b_{l_2} \dots b_{l_r} \end{matrix} \right\}$ under the form of

$$\sum \left\{ R \left(\begin{matrix} \omega_1, \omega_2 \dots \omega_r \\ \psi_1, \psi_2 \dots \psi_r \end{matrix} \right) \times \left\{ \begin{matrix} b_{\omega_1}, b_{\omega_2} \dots b_{\omega_r} \\ b_{\psi_1}, b_{\psi_2} \dots b_{\psi_r} \end{matrix} \right\} \right\};$$

and $R \left(\begin{matrix} \omega_1, \omega_2 \dots \omega_r \\ \psi_1, \psi_2 \dots \psi_r \end{matrix} \right)$ must = 0, except for the case of $\omega_1, \omega_2 \dots \omega_r; \psi_1, \psi_2 \dots \psi_r$ being respectively identical with $k_1, k_2 \dots k_r; l_1, l_2 \dots l_r$, for which case $R \left(\begin{matrix} k_1, k_2 \dots k_r \\ l_1, l_2 \dots l_r \end{matrix} \right)$ must be unity. I have gone through this calculation and verified the result; in order to effect which, however, the following important generalization of theorem (1) must be apprehended.

Suppose two sets of umbrae,

$$a_1, a_2 \dots a_{m+n}, \quad b_1, b_2 \dots b_{m+n},$$

and let r be any number less than m , and let any r -ary combination of the m numbers $1, 2, 3 \dots m$ be expressed by ${}^q\theta_1, {}^q\theta_2 \dots {}^q\theta_m$, where q goes through all the values intermediate between 1 and μ , μ being

$$\frac{m(m-1) \dots (m-r+1)}{1 \cdot 2 \dots r};$$

then I say that the compound determinant,

$$\begin{array}{ccc} \overline{a_{1\theta_1}, a_{1\theta_2} \dots a_{1\theta_m},} & \overline{a_{m+1}, a_{m+2} \dots a_{m+n}} & \overline{a_{2\theta_1}, a_{2\theta_2} \dots a_{2\theta_m},} & \overline{a_{m+1}, a_{m+2} \dots a_{m+n}} \\ b_{1\theta_1}, b_{1\theta_2} \dots b_{1\theta_m}, & b_{m+1}, b_{m+2} \dots b_{m+n} & b_{2\theta_1}, b_{2\theta_2} \dots b_{2\theta_m}, & b_{m+1}, b_{m+2} \dots b_{m+n} \\ & \dots\dots\dots & & \\ & & \overline{a_{\mu\theta_1}, a_{\mu\theta_2} \dots a_{\mu\theta_m},} & \overline{a_{m+1}, a_{m+2} \dots a_{m+n}} \\ & & b_{\mu\theta_1}, b_{\mu\theta_2} \dots b_{\mu\theta_m}, & b_{m+1}, b_{m+2} \dots b_{m+n} \end{array}$$

is equal to the following product,

$$\left\{ \begin{matrix} a_{m+1}, a_{m+2} \dots a_{m+n} \\ b_{m+1}, b_{m+2} \dots b_{m+n} \end{matrix} \right\}^{\mu'} \left\{ \begin{matrix} a_1, a_2 \dots a_{m+n} \\ b_1, b_2 \dots b_{m+n} \end{matrix} \right\}^{\mu''}, \quad (4)$$

where

$$\mu'' = \frac{(m-1)(m-2) \dots (m-r+1)}{1 \cdot 2 \dots (r-1)}, \quad \mu' = \frac{(m-1)(m-2) \dots (m-r)}{1 \cdot 2 \dots r};$$

when $r = 1$, we have the case already given in theorem (2), and of course μ'' is to be taken unity.

This very general theorem is itself several degrees removed from my still unpublished Fundamental Theorem which is a theorem for the expansion of the products of determinants.

Obs. The analogy upon which the extension of the Vandermondian notation from simple to compound determinants is grounded, would be better apprehended if the bilateral symbols of simple quantities were written with the umbral elements disposed vertically, as $\begin{smallmatrix} a \\ b \end{smallmatrix}$, instead of horizontally, as ab ; which latter is the method for the purposes of typographical uniformity adopted in the text above. The other mode is, however, much to be preferred, and is what I propose hereafter to adhere to. For my two general umbrae, a, b , Vandermonde uses two *numbers*, one set a-cock upon the other, as 5^4 . The objection to the use of numbers is apparent as soon as it becomes necessary to treat of the mutual relations of diverse systems of determinants, and his mode of writing the umbrae militates against the perception of the most valuable algebraical analogies. The one important point in which Vandermonde has anticipated me, consists in expressing a simple determinant by two horizontal rows of umbrae one over the other. But the idea upon which this depends is so simple and natural, that it was sure to reappear in any well-constructed system of notation.

NOTE ON QUADRATIC FUNCTIONS AND HYPER-DETERMINANTS

[*Philosophical Magazine, I. (1851), p. 415*]

p. 251

Permit me to correct an error of transcription in the MS. of my paper “On Linearly Equivalent Quadratic Functions” in the last number of the *Magazine*. The theorem [p. 246 above] marked (3), should read as follows:—

$$\left\{ \begin{matrix} b_{\theta_{m+1}}, b_{\theta_{m+2}} \dots b_{\theta_{m+s}} \\ b_{\phi_{m+1}}, b_{\phi_{m+2}} \dots b_{\phi_{m+s}} \end{matrix} \right\} = \left\{ \begin{matrix} a_1, a_2 \dots a_m, & a_{\theta_{m+1}}, a_{\theta_{m+2}} \dots a_{\theta_{m+s}}, & a_{n+1}, a_{n+2} \dots a_{n+m} \\ a_1, a_2 \dots a_m, & a_{\phi_{m+1}}, a_{\phi_{m+2}} \dots a_{\phi_{m+s}}, & a_{n+1}, a_{n+2} \dots a_{n+m} \end{matrix} \right\} \\ \div \left\{ \begin{matrix} a_1, a_2 \dots a_m \\ a_{n+1}, a_{n+2} \dots a_{n+m} \end{matrix} \right\}^2.$$

I may take this opportunity of mentioning, that by extending to algebraical functions generally a multilateral system of umbral notation, analogous to the biliteral system explained in the paper above referred to as applicable to quadratic functions, I have succeeded in reducing to a mechanical method of compound permutation the process for the discovery of those memorable forms invented by Mr Cayley, and named by him hyper-determinants, which have attracted the notice and just admiration of analysts all over Europe, and which will remain a perpetual memorial, as long as the name of algebra survives, of the penetration and sagacity of their author.

ON A CERTAIN FUNDAMENTAL THEOREM OF DETERMINANTS

[*Philosophical Magazine, II. (1851), pp. 142–145*]

p. 252

The subjoined theorem, which is one susceptible of great extension and generalization, appears to me, and indeed from use and acquaintance (it having been long in my possession) I know to be so important and fundamental, as to induce me to extract it from a mass of memoranda on the same subject; and as an act of duty to my fellow-labourers in the theory of determinants, more or less forestall time (the sure discoverer of truth) by placing it without further delay on record in the pages of this *Magazine*. Its developments and applications must be reserved for a more convenient occasion, when the interest in the New Algebra (for such, truly, it is the office of the theory of determinants to establish), and the number of its disciples in this country, shall have received their destined augmentation. In a recent letter to me, M. Hermite well alludes to the theory of determinants as “That vast theory, transcendental in point of difficulty, elementary in regard to its being the basis of researches in the higher arithmetic and in analytical geometry.”

The theorem is as follows:—Suppose that there are two determinants of the ordinary kind, each expressed by a square array of terms made up of n lines and n columns, so that in each square there are n^2 terms. Now let n be broken up in any given manner into two parts p and q , so that $p + q = n$. Let, firstly, one of the two given squares be divided in a given definite manner into two parts, one containing p of the n given lines, and the other part q of the same; and secondly, let the other of the two given squares be divided *in every possible way* into two parts, consisting of q and p lines respectively, so that on tacking on the part containing q lines of the second square to the part containing p lines of the first square, and the part containing p lines of the second square to the part containing q of the first, we

p. 253

get back a new couple of squares, each denoting a determinant different from the two given determinants; the number of such new couples will evidently be

$$\frac{n(n-1) \cdots (n-p+1)}{1 \cdot 2 \cdots p};$$

and my theorem is, that *the product of the given couple of determinants is equal to the sum of the products (affected with the proper algebraical sign) of each of the new couples formed as above described*. Analytically the theorem may be stated as follows.

Let

$$\left\{ \begin{matrix} a_1, a_2 \dots a_n \\ b_1, b_2 \dots b_n \end{matrix} \right\}, \quad \left\{ \begin{matrix} a_1, a_2 \dots a_n \\ \beta_1, \beta_2 \dots \beta_n \end{matrix} \right\},$$

according to the notation heretofore¹⁴⁰ employed by me in the preceding numbers of this *Magazine*, denote any two common determinants, each of the n th order, and let the numbers $\theta_1, \theta_2 \dots \theta_n$ be disjunctively equal to the numbers $1, 2 \dots n$ and $p + q = n$; then will

$$\begin{aligned} & \begin{Bmatrix} a_1, a_2 \dots a_n \\ b_1, b_2 \dots b_n \end{Bmatrix} \times \begin{Bmatrix} a_1, a_2 \dots a_n \\ \beta_1, \beta_2 \dots \beta_n \end{Bmatrix} \\ &= \sum \pm \begin{Bmatrix} a_1, a_2 \dots a_p, & a_{p+1} \dots a_n \\ b_1, b_2 \dots b_p, & \beta_{\theta_{p+1}}, \beta_{\theta_{p+2}} \dots \beta_{\theta_n} \end{Bmatrix} \times \begin{Bmatrix} a_1, a_2 \dots a_q, & a_{q+1} \dots a_n \\ \beta_{\theta_1}, \beta_{\theta_2} \dots \beta_{\theta_q}, & b_{p+1}, b_{p+2} \dots b_n \end{Bmatrix}. \end{aligned}$$

The general term under the sign of summation may be represented by aid of the disjunctive equations

$$\phi_1, \phi_2 \dots \phi_n = 1, 2 \dots n, \quad \psi_1, \psi_2 \dots \psi_n = 1, 2 \dots n,$$

under the form of

$$\begin{aligned} & (a_{\phi_1} b_1 \times a_{\phi_2} b_2 \times \dots \times a_{\phi_p} b_p) (a_{\psi_{p+1}} b_{p+1} \times a_{\psi_{p+2}} b_{p+2} \times \dots \times a_{\psi_n} b_n) \\ & \times (a_{\phi_{p+1}} \beta_{\theta_{p+1}} \times a_{\phi_{p+2}} \beta_{\theta_{p+2}} \times \dots \times a_{\phi_n} \beta_{\theta_n}) (a_{\psi_1} \beta_{\theta_1} \times a_{\psi_2} \beta_{\theta_2} \times \dots \times a_{\psi_p} \beta_{\theta_p}). \end{aligned}$$

1st. When $\phi_1, \phi_2 \dots \phi_p = \psi_1, \psi_2 \dots \psi_p$, it will readily be seen, that for given values of $\phi_1, \phi_2 \dots \phi_p$, the product of the third and fourth factors becomes substantially identical with the general term of the determinant

$$\begin{Bmatrix} a_1, a_2 \dots a_n \\ \beta_1, \beta_2 \dots \beta_n \end{Bmatrix},$$

and consequently, making the system $\phi_1, \phi_2 \dots \phi_p$ (or, which is the same thing, its equivalent $\psi_1, \psi_2 \dots \psi_p$) go through all its values, we get back for the sum of the terms corresponding to the equation

$$\phi_1, \phi_2 \dots \phi_p = \psi_1, \psi_2 \dots \psi_p,$$

p. 254

the product of the determinants

$$\begin{Bmatrix} a_1, a_2 \dots a_n \\ b_1, b_2 \dots b_n \end{Bmatrix} \quad \text{and} \quad \begin{Bmatrix} a_1, a_2 \dots a_n \\ \beta_1, \beta_2 \dots \beta_n \end{Bmatrix}.$$

2nd. When we have not the equality above supposed between the ϕ 's and the ψ 's, let

$$\phi_{p-\eta} = \psi_{p+k} \quad \text{and} \quad \phi_{p+\eta} = \psi_{p-\zeta};$$

the corresponding term included under the Σ will contain the factor

$$a_{\phi_{p+\eta}} \beta_{\theta_{p+\eta}} \times a_{\psi_{p-\zeta}} \beta_{\theta_{p-\zeta}}.$$

¹⁴⁰p. 242 above.

Now leaving $\phi_1, \phi_2 \dots \phi_p$, and $\psi_1, \psi_2 \dots \psi_p$ unaltered, we may take a system of values $\theta'_1, \theta'_2 \dots \theta'_n$, such that

$$\theta'_{p+\eta} = \theta_{p-\zeta}, \quad \theta'_{p-\zeta} = \theta_{p+\eta},$$

and for all other values of q except $p + \eta$, or $p - \zeta$, $\theta'_q = \theta_q$. The corresponding new value of the general term so formed by the substitution of the θ' for the θ series, will be identical with that of the term first spoken of, but will have the contrary algebraical sign, because the θ' arrangement of the figures $1, 2, 3 \dots p$ is deducible by a single interchange from the θ arrangement of the same, the rule for the imposition of the algebraical sign plus or minus being understood to be, that the term in which

$$\beta_{\theta_{p+1}}, \beta_{\theta_{p+2}} \dots \beta_{\theta_n}; \quad \beta_{\theta_1}, \beta_{\theta_2} \dots \beta_{\theta_p}$$

enter into the symbolical forms of the respective derived couples of determinants, has the same sign as, or the contrary sign to, that in which

$$\beta_{\theta'_{p+1}}, \beta_{\theta'_{p+2}} \dots \beta_{\theta'_n}; \quad \beta_{\theta'_1}, \beta_{\theta'_2} \dots \beta_{\theta'_p}$$

so enter, according as an odd or an even number of interchanges is required to transform the arrangement

$$\theta_{p+1}, \theta_{p+2} \dots \theta_n; \quad \theta_1, \theta_2 \dots \theta_p$$

into the arrangement

$$\theta'_{p+1}, \theta'_{p+2} \dots \theta'_n; \quad \theta'_1, \theta'_2 \dots \theta'_p.$$

I have therefore shown that all the terms arising from the expansion of the products included under the sign of summation, for which the disjunctive identity $\phi_1, \phi_2 \dots \phi_p = \psi_1, \psi_2 \dots \psi_p$ does not exist, enter into the final sum in pairs, equal in quantity and differing in sign, which consequently mutually destroy, and that the terms for which the said identity does exist together make up the sum

$$\left\{ \begin{matrix} a_1, a_2 \dots a_n \\ b_1, b_2 \dots b_n \end{matrix} \right\} \text{imes} \left\{ \begin{matrix} a_1, a_2 \dots a_n \\ \beta_1, \beta_2 \dots \beta_n \end{matrix} \right\};$$

which proves, upon first principles drawn direct from that notion of polar dichotomy of permutation systems which rests at the bottom of the whole theory of the subject, the fundamental, and, as I believe, perfectly new theorem, which it is the object of this communication to establish.

In applying the theorem thus analytically formulized, it is of course to be understood that, under the sign Σ , permutations within the separate parts of a given arrangement,

$$\theta_{p+1}, \theta_{p+2} \dots \theta_n; \quad \theta_1, \theta_2 \dots \theta_p,$$

are inadmissible, the total number of terms so included being restricted to

$$\frac{n(n-1)\cdots(n-p+1)}{1\cdot 2\cdots p}.$$

The theorem may be extended so as to become a theorem for the expansion of the product of any number of determinants, and adapted so as to take in that far more general class of functions known to Mr Cayley and myself under the new name of commutants, of which determinants present only a particular, and that the most limited instance.

ON EXTENSIONS OF THE DIALYTIC METHOD OF ELIMINATION

[*Philosophical Magazine, II. (1851), pp. 221–230*]

p. 256

The theory about to be described is a natural extension of the method of elimination presented by me ten years ago (in June, 1841) in the pages of this *Magazine*, which I have been induced to review in consequence of the flattering interest recently expressed in the subject by my friend M. Terquem, and some other continental mathematicians, and because of the importance of the geometrical and other applications of which it admits, and of the inquiries to which it indirectly gives rise. We shall be concerned in the following discussion with systems of homogeneous rational integral functions of a peculiar form, to which for present purposes I propose to give the name of aggregative functions, consisting of ordinary homogeneous functions of the same variables but of different degrees, brought together into one sum made homogeneous by means of powers of new variables entering factorially.

Thus if $F, G, H \dots L$ be any number of functions of any number of letters $x, y \dots t$ of the degrees $m, m - \iota, m - \iota' \dots m - (\iota)$ respectively,

$$F + G\lambda^\iota + H\mu^{\iota'} + \dots + L\theta^{(\iota)}$$

will be an aggregative function of the variables entering into F, G , &c. and of $\lambda, \mu \dots \theta$. I shall further call such a function binary, ternary, quaternary, and so forth, according to the number of variables contained in the functions F, G, H , &c. thus brought into coalition.

It will be convenient to recall the attention of the reader to the meaning of some of the terms employed by me in the paper above referred to.

If F be any homogeneous function of $x, y, z \dots t$, the term augmentative of F denotes any function obtained from F of the form

$$x^\alpha y^\beta z^\gamma \dots t^\delta \times F.$$

Again, if we have any number of such functions $F, G, H \dots K$ of as many p. 257

variables $x, y, z \dots t$, and we decompose $F, G, H \dots K$ in any manner so as to obtain the equations

$$\begin{aligned} F &= x^a P_1 + y^b P_2 + z^c P_3 + \&c. \dots + t^d(P), \\ G &= x^a Q_1 + y^b Q_2 + z^c Q_3 + \&c. \dots + t^d(Q), \\ H &= x^a R_1 + y^b R_2 + z^c R_3 + \&c. \dots + t^d(R), \\ &\dots\dots\dots \\ K &= x^a S_1 + y^b S_2 + z^c S_3 + \&c. \dots + t^d(S), \end{aligned}$$

and then form the determinant

$$\begin{vmatrix} P_1 & P_2 & P_3 & \cdots & (P) \\ Q_1 & Q_2 & Q_3 & \cdots & (Q) \\ R_1 & R_2 & R_3 & \cdots & (R) \\ \cdots & \cdots & \cdots & & \cdots \\ S_1 & S_2 & S_3 & \cdots & (S) \end{vmatrix},$$

this determinant, expressed as a function of $x, y, z \dots t$, is what, in the paper referred to, I called a secondary derivate, but which for the future I shall cite by the more concise and expressive name of a connective of the system of functions $F, G, H \dots K$ from which it is obtained. One prevailing principle regulates all the cases treated of in this and the antecedent memoir, namely that of forming linearly independent systems of augmentatives or connectives, or both, of the given system whose resultant is to be found, of the same degree one with the other, and equal in number (when this admits of being done) to the number of distinct terms in the functions thus formed. The resultant of these functions, treated as linear functions of the several combinations of powers of the variables in each term, will then be the resultant of the given system clear of all irrelevant factors. If the number of terms to be eliminated exceed the number of the functions, the elimination of course cannot be executed. If the contrary be the case, but the equality is restored by the rejection of a certain number of the equations, the resultant so obtained will vary according to the choice of the equations retained for the purpose of the elimination. The true resultant will not then coincide with any of the resultants so obtained, but will enter as a common factor into them all.

The following simple arithmetical principles will be found applicable and useful for quotation in the sequel:—

(a) The number of terms in a homogeneous function of p letters of the m th degree is

$$\frac{m(m+1) \cdots (m+p-1)}{1 \cdot 2 \cdots p}.$$

(b) The number of augmentatives of the $(m+n)$ th degree belonging to a function of p letters of the m th degree is

$$\frac{(n+1)(n+2) \cdots (n+p-1)}{1 \cdot 2 \cdots p}.$$

(c) The number of solutions in integers (excluding zeros) of the equation $a_1 + a_2 + \cdots + a_p = k$ is

$$\frac{(k-1)(k-2) \cdots (k-p+1)}{1 \cdot 2 \cdots (p-1)}.$$

p. 258

To begin with the case of binary aggregatives. Let

$$\left. \begin{aligned} &F_m(x, y) + F_{m-\iota}(x, y)\lambda^\iota + F_{m-\iota'}(x, y)\mu^{\iota'} + \&c. \cdots + F_{m-(\iota)}(x, y)\theta^{(\iota)} \\ &G_n(x, y) + G_{n-\iota}(x, y)\lambda^\iota + G_{n-\iota'}(x, y)\mu^{\iota'} + \&c. \cdots + G_{n-(\iota)}(x, y)\theta^{(\iota)} \\ &\dots\dots\dots \\ &K_p(x, y) + K_{p-\iota}(x, y)\lambda^\iota + K_{p-\iota'}(x, y)\mu^{\iota'} + \&c. \cdots + K_{p-(\iota)}(x, y)\theta^{(\iota)} \end{aligned} \right\} \quad (A)$$

be a system of functions (whose Resultant it is proposed to determine) equal in number to the variables $x, y, \lambda, \mu \dots \theta$, and similarly aggregative, that is having only the same powers of λ, μ , &c. entering into them, but of any degrees equal or unequal $m, n \dots p$. Let the number of the functions be r . Raise each of the given functions by augmentation to the degree s , where

$$s = (m + n + \cdots + p) - (\iota + \iota' + \cdots + (\iota)) - 1,$$

the number of augmentatives of the several functions will be

$$(s + 1) - m, \quad (s + 1) - n, \quad \cdots, \quad (s + 1) - p,$$

and the total number will therefore be

$$r(s + 1) - (m + n + \cdots + p),$$

which

$$= (r - 1)(m + n + \cdots + p) - r(\iota + \iota' + \cdots + (\iota)).$$

Again, the number of terms to be eliminated will be the sum of the numbers of terms in functions respectively of the s th, $(s - \iota)$ th, $(s - \iota')$ th, $\dots (s - (\iota))$ th degrees, which are respectively

$$s + 1, \quad s + 1 - \iota, \quad s + 1 - \iota', \quad \cdots, \quad s + 1 - (\iota),$$

p. 259

and the number of these partial functions is $r - 1$. Hence the number of terms to be eliminated is

$$\begin{aligned} &(r - 1)\{m + n + \&c. + p - (\iota + \iota' + \&c. + (\iota))\} - (\iota + \iota' + \&c. + (\iota)) \\ &= (r - 1)(m + n + \&c. + p) - r(\iota + \iota' + \cdots + (\iota)), \end{aligned}$$

which is exactly equal to the number of the augmentative functions. Hence the Resultant¹⁴¹ of the given functions can be found dialytically by linear elimination, and the exponent of its dimensions in respect to the coefficients of the given functions will be the number

$$(r - 1)\Sigma m - r\Sigma \iota,$$

¹⁴¹The Resultant of a system of functions means in general the same thing as the left-hand side of the final equation (clear of extraneous factors) resulting from the elimination of the variables between the equations formed by equating the said functions severally to zero.

as above found.

The method above given may be replaced by another more compendious, and analogous to that known by the name of Bezout's abridged method for ordinary functions of two letters. As the method is precisely the same whatever the number of the functions employed may be, I shall for the sake of greater simplicity restrict the demonstration to the case of three functions, U, V, W , whose degrees (if unequal, written in ascending order of magnitude) are m, n, p respectively. Let

$$U = F_m(x, y) + F_{m-\iota}(x, y)z^\iota,$$

$$V = G_n(x, y) + G_{n-\iota}(x, y)z^\iota,$$

$$W = H_p(x, y) + H_{p-\iota}(x, y)z^\iota.$$

Let θ, ω be taken any two numbers which satisfy in integers greater than zero the equation $\theta + \omega = m + 1$, and let

$$F_m(x, y) = \phi_{m-\theta} \cdot x^\theta + \phi_{m-\omega} \cdot y^\omega,$$

$$G_n(x, y) = \gamma_{n-\theta} \cdot x^\theta + \gamma_{n-\omega} \cdot y^\omega,$$

$$H_p(x, y) = \eta_{p-\theta} \cdot x^\theta + \eta_{p-\omega} \cdot y^\omega,$$

where the ϕ 's, γ 's, η 's may be always considered rational integer functions of x and y ; for every term in each of the functions F_m, G_n, H_p must either contain x^θ or y^ω , since, if not, its dimensions in x and y would not exceed

$$(\theta - 1) + (\omega - 1),$$

that is $m - 1$, whereas each term is of m conjoined dimensions, at least, in x and y . Hence from the equations

$$U = 0, \quad V = 0, \quad W = 0,$$

p. 260

by eliminating x^ω, y^θ and z^ι we obtain the connective determinant

$$\begin{vmatrix} \phi_{m-\theta} & \phi_{m-\omega} & F_{m-\iota} \\ \gamma_{n-\theta} & \gamma_{n-\omega} & G_{n-\iota} \\ \eta_{p-\theta} & \eta_{p-\omega} & H_{p-\iota} \end{vmatrix},$$

which will be of the degree

$$m + n + p - (\theta + \omega + \iota),$$

that is of the degree $(n + p - \iota - 1)$ in x and y ; and the number of such connectives by principle (c) is p .

Again, by augmentation we can raise each of the functions U, V, W to the same degree as the connectives, and by principle (b) the number of such will be

$$n + p - m - \iota, \quad p - \iota, \quad n - \iota,$$

from U, V, W respectively, together making up the number

$$2n + 2p - m - 3\iota.$$

Hence in all we have $2n + 2p - 3\iota$ equations; and the number of terms to be eliminated will be, $n + p - \iota$ arising from F_m, G_n, H_p , and $n + p - 2\iota$ from $F_{m-\iota}, G_{n-\iota}, H_{p-\iota}$; together making up the proper number $2n + 2p - 3\iota$.

Each connective contains ternary combinations of the coefficients, namely one of the coefficients belonging to that part of U, V, W which contains z^ι , and two coefficients from the other part: the dimensions of the resultant in respect of the coefficients of the former will hence be readily seen to be equal to the number of connectives + the number of terms in the augmentatives into which z^ι enters, that is, will equal $m + n + p - 2\iota$; the total dimensions of the resultant in respect to all the coefficients of U, V, W will be

$$3m + (2n + 2p - m - 3\iota),$$

that is,

$$2m + 2n + 2p - 3\iota;$$

and consequently, in respect to the coefficients of $F_m; G_n; H_p$, will be of

$$(2m + 2n + 2p - 3\iota) - (m + n + p - 2\iota),$$

that is, of $m + n + p - \iota$ dimensions. This result, which is of considerable importance, may be generalized as follows.

Returning to the general system (A), for which we have proved that the total dimensions of the resultant are

$$(r - 1)(m + n + \dots + p) - r(\iota + \iota' + \dots + (\iota)),$$

p. 261

let the coefficients of the column of partial functions

$$F_m, \quad G_n, \quad \vdots, \quad K_p,$$

be called the first set; the coefficients of the column

$$F_{m-\iota}, \quad G_{n-\iota}, \quad \vdots, \quad K_{p-\iota},$$

the second set, and so forth; then the dimensions in respect of the 1st, 2nd $\dots (r - 1)$ th sets respectively are $s, s - \iota, s - \iota' \dots s - (\iota)$, where

$$s = m + n + \&c. + p - (\iota + \iota' + \&c. + (\iota)).$$

The important observation remains to be made, that all the above results remain good although any one or more of the indices of dimension of the partial functions

in the system (A), as $m - \iota, m - \iota', n - \iota$, &c., should become negative, provided that the terms in which such negative indices occur be taken zero, as will be apparent on reviewing the processes already indicated upon this supposition. If we take

$$m = n = \dots = p, \quad \iota = \iota' = \&c. = (\iota) = m - \epsilon,$$

the exponent of the total dimensions of the resultant becomes

$$(r - 1)rm - r(r - 2)(m - \epsilon) = rm + r(r - 2)\epsilon,$$

when $\epsilon = 0$, this becomes mr , which is made up of $2m$ units of dimension belonging to the coefficients of the first column, and of m belonging to each of the $(r - 2)$ remaining columns. Consequently, if we have

$$\begin{aligned} F_m(x, y) + \xi\lambda + \xi'\lambda' &= 0, \\ G_m(x, y) + \eta\lambda + \eta'\lambda' &= 0, \\ H_m(x, y) + \zeta\lambda + \zeta'\lambda' &= 0, \\ K_m(x, y) + \theta\lambda + \theta'\lambda' &= 0, \end{aligned}$$

or any other number of equations similarly formed, the result of the elimination is always of m dimensions only in respect of ξ, η, ζ, θ , or of $\xi', \eta', \zeta', \theta'$, and of $2m$ in respect of the coefficients in F, G, H, K .

I now proceed to state and to explain some seeming paradoxes connected with the degree of the resultant of such systems of defective functions as have been previously treated of in this memoir, as compared with the degree

p. 262

of the general resultant of a corresponding system of complete functions of the same number of variables.

In order to fix our ideas, let us take a system of only three equations of the form

$$\left. \begin{aligned} F_m(x, y) + F_{m-\iota}(x, y)z^\iota &= 0, \\ G_n(x, y) + G_{n-\iota}(x, y)z^\iota &= 0, \\ H_p(x, y) + H_{p-\iota}(x, y)z^\iota &= 0 \end{aligned} \right\} \quad (B)$$

The resultant of this system found by the preceding method is in all of $2m + 2n + 2p - 3\iota$ dimensions. But in general, the resultant of three equations of the degrees m, n, p is of $mn + mp + np$ dimensions.

Now in order to reason firmly and validly upon the doctrine of elimination, nothing is so necessary as to have a clear and precise notion, never to be let go from the mind's grasp, of the proposition that every system of n homogeneous functions of n variables has a single and invariable Resultant. The meaning of this proposition is, that a function of the coefficients of the given functions can be found, such that, whenever it becomes zero, and never except when it becomes zero, the functions may be simultaneously made zero for some certain system of ratios between the variables. The function so found, which is sufficient

and necessary to condition the possibility of the coexistence of the equality to zero of each of the given functions, is their resultant, and by analogy they may be termed its components. It follows that if R be a resultant of a given system of functions, any numerical multiple of any power of R or of any root of R when (upon certain relations being supposed to be instituted between the coefficients of its components) R breaks up into equal factors, will also be a resultant. This is just what happens in system (B) when $m = n = p = \iota$; the resultant found by the method in the text is of the degree $3m$; the general resultant of the system of three equations to which it belongs is of the degree $3m^2$; the fact being, that the latter resultant becomes a perfect m th power for the particular values of the coefficients which cause its components to take the form of the functions in system (B).

Suppose, however, that we have still $m = n = p$, but ι less than m , $6m - 3\iota$ will express the degree of the resultant of system (B); but this is no longer in general an aliquot part of $3m^2$, and consequently the resultant of system (B) that we have found is no longer capable in general of being a root of the general resultant. The truth is, that on this supposition the general resultant is zero; as it evidently should be, because the values

$$\frac{x}{z} = 0, \quad \frac{y}{z} = 0$$

satisfy the equations in system (B), except for the case of $m = \iota$; consequently the resultant furnished in the text, although found by the same process, is something of a different nature from an ordinary resultant; it

p. 263

expresses, not that the system of equations (B) may be capable of coexisting, but that they may be capable of coexisting for values of $\frac{x}{z}, \frac{y}{z}$ other than 0 and 0. This is what I have elsewhere termed a sub-resultant. But there is yet a further case, to which neither of the above considerations will apply. This is when m, n, p are not equal, but $p - \iota = 0$.

On this supposition the degree of the resultant of (B) becomes $2m + 2n - p$, which in general will not be a factor of $mn + mp + np$; and in this case it will no longer be true that the values $\frac{x}{z} = 0, \frac{y}{z} = 0$ will satisfy the system (B), inasmuch as the last equation therein cannot so be satisfied. Now, calling the general resultant R and the particular resultant R' , if R' should break up into factors so as to become equal to $(r')^a \times (s')^b \cdots (t')^\nu$, it might be the case that R should equal $(r')^\alpha \cdot (s')^\beta \cdots (t')^\gamma$, and there would be nothing in this fact which would be inconsistent with the theory of the resultant as above set forth; but suppose that R' is indecomposable into factors, then it is evident that we must have $R = R' \cdot R''$, and consequently that the existence of such a particular resultant as R' will argue the necessity of the existence of another resultant R'' ; in other words, the resultant so found cannot be in a strict sense the true and complete resultant for the particular case assumed, and yet the process employed appears to give the complete resultant, or at least it is difficult to see how the wanting

factor escapes detection. To make this matter more clear, take a particular and a very simple case, where $m = 2$, $n = 2$, $p = \iota = 1$, so as to form the system of equations

$$\left. \begin{aligned} Ax^2 + Bxy + Cy^2 + (Dx + Ey)z &= 0 \\ A'x^2 + B'xy + C'y^2 + (D'x + E'y)z &= 0 \\ lx + my + nz &= 0 \end{aligned} \right\} \quad (C)$$

By virtue of my theorem, the degree of the resultant R' is

$$2(2 + 2 + 1) - 3 \cdot 1 = 7,$$

but the resultant R of the system

$$\left. \begin{aligned} Ax^2 + Bxy + Cy^2 + (Dx + Ey)z + Fz^2 &= 0 \\ A'x^2 + B'xy + C'y^2 + (D'x + E'y)z + F'z^2 &= 0 \\ lx + my + nz &= 0 \end{aligned} \right\} \quad (D)$$

which becomes identical with the former when $F = 0$, $F' = 0$ is of

$$2 \times 2 + 2 \times 1 + 2 \times 1,$$

that is, of 8 dimensions. Hence it is evident that when $F = 0$, $F' = 0$, R must become $R' \times R''$.

p. 264

It will be found in fact¹⁴², that on the supposition of $F = 0$, $F' = 0$, R becomes equal to $N \times R'$; and accordingly, besides the portion R' of the resultant of system (C), found by the method in the text, there is another portion N which has dropped through; but it may be asked, is N truly a relevant factor? were it not so, the theory of the resultant would be completely invalidated; but in truth it is; for $N = 0$ will make the equations in system (C), considered as a particular case of system (D), capable of co-existing; the peculiarity, which at first sight prevents this from being obvious, consisting in the fact that the values of $\frac{x}{z}$, $\frac{y}{z}$ which satisfy the three equations when $N = 0$ become infinite.

Thus, finally, we have arrived at a clear and complete view of the relation of the particular to the general resultant.

The general resultant may be zero, in which case the particular resultant is something altogether different from an ordinary resultant; or the particular resultant may be a root of the general resultant, or it may be more generally the product of powers of the simple factors, which enter into the composition of the general resultant; or lastly, it may be an incomplete resultant, the factors wanting to make it complete being such as when equated to zero, will enable the components of the resultant to coexist, but not for other than infinite values of certain of the ratios existing between the variables.

¹⁴²See the Author's remarks below, p. 283.

Without for the present further enlarging on the hitherto unexplored and highly interesting theory of Particular Resultants, I will content myself with stating one beautiful and general theorem relating to them; to wit, “if $F = 0$, $G = 0$, &c. be a given system of equations with the coefficients left general, and R be the resultant of F, G , &c., and if now the coefficients in F, G be so taken that R comes to contain as a factor or be coincident with R^m , then will $R' = 0$ indicate that (when the coefficients are so taken as above supposed) $F = 0$, $G = 0$, &c. will be capable of being satisfied, not, as in general, by one only, but by m distinct systems of values of the variables in F, G , &c., subject of course to the possibility, in special cases, of certain of the systems becoming multiple coincident systems.”

I pass on now¹⁴³ to the more recondite and interesting theory of the resultant of Ternary Aggregative Functions, that is to say, functions of the form

$$F_m(x, y, z) + F_{m-\iota}(x, y, z)t^\iota + \&c. \cdots + F_{m-(\iota)}(x, y, z)t^{(\iota)},$$

which will be seen to admit of some remarkable applications to the theory of reciprocal polars.

¹⁴³See the Author's remarks below, p. 283.

41.

ON A REMARKABLE DISCOVERY IN THE THEORY OF CANONICAL FORMS AND OF HYPERDETERMINANTS

[*Philosophical Magazine, II. (1851), pp. 391-410*]

p. 265

In a recently printed continuation¹⁴⁴ of a paper which appeared in the *Cambridge and Dublin Mathematical Journal*, I published a complete solution of the following problem. A homogeneous function of x, y of the degree $2n + 1$ being given, required to represent it as the sum of $n + 1$ powers of linear functions of x, y . I shall prepare the way for the more remarkable investigations which form the proper object of this paper, by giving a new and more simple solution of this linear transformation.

Let the given function be

$$a_0 x^{2n+1} + (2n+1)a_1 x^{2n}y + \frac{1}{2}(2n+1)(2n)a_2 x^{2n-1}y^2 + \cdots + a_{2n+1}y^{2n+1},$$

and suppose that this is identical with

$$(p_1x + q_1y)^{2n+1} + (p_2x + q_2y)^{2n+1} + \&c. + (p_{n+1}x + q_{n+1}y)^{2n+1}.$$

The problem is evidently possible and definite, there being $2n + 2$ equations to be satisfied, and $(2n + 2)$ quantities $p_1, q_1, \&c.$ for satisfying the same.

In order to effect the solution, let

$$q_1 = p_1\lambda_1, \quad q_2 = p_2\lambda_2, \quad \&c. = \&c., \quad q_{n+1} = p_{n+1}\lambda_{n+1},$$

p. 266

we have then

$$\begin{aligned} p_1^{2n+1} + p_2^{2n+1} + \cdots + p_{n+1}^{2n+1} &= a_0, \\ p_1^{2n+1}\lambda_1 + p_2^{2n+1}\lambda_2 + \cdots + p_{n+1}^{2n+1}\lambda_{n+1} &= a_1, \\ p_1^{2n+1}\lambda_1^2 + p_2^{2n+1}\lambda_2^2 + \cdots + p_{n+1}^{2n+1}\lambda_{n+1}^2 &= a_2, \\ &\dots\dots\dots \\ p_1^{2n+1}\lambda_1^n + p_2^{2n+1}\lambda_2^n + \cdots + p_{n+1}^{2n+1}\lambda_{n+1}^n &= a_n, \\ p_1^{2n+1}\lambda_1^{n+1} + p_2^{2n+1}\lambda_2^{n+1} + \cdots + p_{n+1}^{2n+1}\lambda_{n+1}^{n+1} &= a_{n+1}, \\ &\dots\dots\dots \\ p_1^{2n+1}\lambda_1^{2n+1} + p_2^{2n+1}\lambda_2^{2n+1} + \cdots + p_{n+1}^{2n+1}\lambda_{n+1}^{2n+1} &= a_{2n+1}. \end{aligned}$$

Eliminate $p_1, p_2 \dots p_{n+1}$ between the 1st, 2nd, 3rd $\dots (n + 2)$ th equations, and it is easily seen that we obtain

$$a_{n+1} - a_n \Sigma \lambda_1 + a_{n-1} \Sigma \lambda_1 \lambda_2 \cdots \pm a_0 \lambda_1 \lambda_2 \cdots \lambda_{n+1} = 0.$$

¹⁴⁴p. 203 above.

Again, eliminating in like manner $p_1^{2n+1}\lambda_1, p_2^{2n+1}\lambda_2 \dots p_{n+1}^{2n+1}\lambda_{n+1}$ between the 2nd, 3rd ... $(n+3)$ th equations, we obtain

$$a_{n+2} - a_{n+1}\Sigma\lambda_1 + \dots \mp a_1\lambda_1\lambda_2 \dots \lambda_{n+1} = 0;$$

and proceeding in the same way until we come to the combination of $(n+1)$ th ... $(2n+2)$ th equations, and writing

$$\Sigma\lambda_1 = s_1, \quad \Sigma\lambda_1\lambda_2 = s_2, \quad \dots, \quad \lambda_1\lambda_2 \dots \lambda_{n+1} = s_{n+1},$$

we find

$$\begin{aligned} a_{n+1} - a_n s_1 + a_{n-1} s_2 \dots + a_0 s_{n+1} &= 0, \\ a_{n+2} - a_{n+1} s_1 + a_n s_2 \dots \mp a_1 s_{n+1} &= 0, \\ a_{n+3} - a_{n+2} s_1 + a_{n+1} s_2 \dots + a_2 s_{n+1} &= 0, \\ &\dots\dots\dots \\ a_{2n+1} - a_{2n} s_1 + a_{2n-1} s_2 \dots + a_n s_{n+1} &= 0. \end{aligned}$$

Hence it is obvious that

$$(x + \lambda_1 y)(x + \lambda_2 y) \dots (x + \lambda_{n+1} y)$$

is a constant multiple of the determinant

$$\begin{vmatrix} x^{n+1}, & -x^n y, & x^{n-1} y^2, & \dots, & \pm y^{n+1} \\ a_{n+1}, & a_n, & a_{n-1}, & \dots, & a_0 \\ a_{n+2}, & a_{n+1}, & a_n, & \dots, & a_1 \\ \dots & \dots & \dots & \dots & \dots \\ a_{2n+1}, & a_{2n}, & a_{2n-1}, & \dots, & a_n \end{vmatrix}.$$

p. 267

Hence $\lambda_1, \lambda_2 \dots \lambda_{n+1}$ are known, and consequently

$$p_1, p_2 \dots p_{n+1}, \quad q_1, q_2 \dots q_{n+1}$$

are known, by the solution of an equation of the $(n+1)$ th degree.

Thus suppose the given function to be

$$\begin{aligned} F &= ax^5 + 5bx^4y + 10cx^3y^2 + 10dx^2y^3 + 5exy^4 + 10fy^5 \\ &= (p_1x + q_1y)^5 + (p_2x + q_2y)^5 + (p_3x + q_3y)^5, \end{aligned}$$

we shall have, by an easy inference from what has preceded,

$$(p_1x + q_1y)(p_2x + q_2y)(p_3x + q_3y)$$

equal to a numerical multiple of the determinant

$$\begin{vmatrix} x^3, & -x^2y, & xy^2, & -y^3 \\ d, & c, & b, & a \\ e, & d, & c, & b \\ f, & e, & d, & c \end{vmatrix}.$$

The solution of the problem given by me in the paper before alluded to presents itself under an *apparently* different and rather less simple form. Thus, in the case in question, we shall find according to that solution,

$$(p_1x + q_1y)(p_2x + q_2y)(p_3x + q_3y)$$

equal to a numerical multiple of the determinant

$$\begin{vmatrix} ax + by & bx + cy & cx + dy \\ bx + cy & cx + dy & dx + ey \\ cx + dy & dx + ey & ex + fy \end{vmatrix}.$$

The two determinants, however, are in fact identical, as is easily verified, for the coefficients of x^3 and y^3 are manifestly alike; and the coefficient of x^2y in the second form will be made up of the three determinants,

$$\begin{vmatrix} a & b & d \\ b & c & e \\ c & d & f \end{vmatrix}, \quad \begin{vmatrix} a & c & c \\ b & d & d \\ c & e & e \end{vmatrix}, \quad \begin{vmatrix} b & b & c \\ c & c & d \\ d & d & e \end{vmatrix},$$

of which the latter two vanish, and the first is identical with the coefficient of x^2y in the first solution. The same thing is obviously true in regard of the coefficients of xy^2 in the two forms, and a like method may be applied to show that in all cases the determinant above given is identical with the determinant of my former paper, namely

$$\begin{vmatrix} a_0x + a_1y & a_1x + a_2y & \cdots & a_nx + a_{n+1}y \\ a_1x + a_2y & a_2x + a_3y & \cdots & a_{n+1}x + a_{n+2}y \\ \cdots & \cdots & \cdots & \cdots \\ a_nx + a_{n+1}y & a_{n+1}x + a_{n+2}y & \cdots & a_{2n}x + a_{2n+1}y \end{vmatrix}.$$

Thus, then, we see that for odd-degreed functions, the reduction to their canonical form of the sum of $(n + 1)$ powers depends upon the solution of one single equation of the $(n + 1)$ th degree, and can never be effected in more than one way.

This new form of the resolving determinant affords a beautiful criterion for a function of x, y of the degree $2n + 1$ being composed of n instead of, as in general, $(n + 1)$ powers. In order that this may be the case, it is obvious that two conditions must be satisfied; but I pointed out in my supplemental paper on canonical forms, that all the coefficients of the resolving determinant must vanish, which appears to give far too many conditions. Thus, suppose we have

$$ax^7 + 7bx^6y + 21cx^5y^2 + 35dx^4y^3 + 35ex^3y^4 + 21fx^2y^5 + 7gxy^6 + hy^7.$$

The conditions of catalecticism, that is, of its being expressible under form of the sum of three (instead of, as in general, four) seventh powers, requires that all

the coefficients of the different powers of x and y must vanish in the determinant

$$\begin{vmatrix} y^4 & -y^3x & y^2x^2 & -yx^3 & x^4 \\ a & b & c & d & e \\ b & c & d & e & f \\ c & d & e & f & g \\ d & e & f & g & h \end{vmatrix};$$

in other words, we must have five determinants,

$$\begin{vmatrix} a & b & c & d \\ b & c & d & e \\ c & d & e & f \\ d & e & f & g \end{vmatrix}, \quad \begin{vmatrix} a & c & d & e \\ b & d & e & f \\ c & e & f & g \\ d & f & g & h \end{vmatrix}, \quad \begin{vmatrix} a & b & c & e \\ b & c & d & f \\ c & d & e & g \\ d & e & f & h \end{vmatrix},$$

$$\begin{vmatrix} a & b & d & e \\ b & c & e & f \\ c & d & f & g \\ d & e & g & h \end{vmatrix}, \quad \begin{vmatrix} b & c & d & e \\ c & d & e & f \\ d & e & f & g \\ e & f & g & h \end{vmatrix},$$

all separately zero. But by my homaloidal law¹⁴⁵, all these five equations amount only to $(5-4)(5-3)$, that is, to 2. I may notice here, that a theorem substantially identical with this law, and another absolutely identical with the theorem of compound determinants given by me in this *Magazine*, and afterwards generalized in a paper also published¹⁴⁶ in this *Magazine*, entitled

¹⁴⁵p. 150 above.

¹⁴⁶p. 241 above.