

Exposé III. Functoriality of Categories of Sheaves

Opening and Section 1: Continuous Functors

J.-L. Verdier

Translator's status note. This is a working English translation of SGA 4, Exposé III, from the opening paragraph through Example 1.9.3. It corresponds to source lines 34–611 of the Orgogozo–Laszlo TeX snapshot. I write $\widehat{C}$ for the category of presheaves on C , and $\widetilde{C}$ for the category of sheaves on C . Terminology is normalized to current English usage: *sieve*, *covering morphism*, *bicovering morphism*, *associated sheaf* or *sheafification*, and *base-changeable* for French *quarrable*.

In Exposé I, Section 5, the behavior of categories of presheaves with respect to functors between categories of arguments was studied. In the present exposé we extend that study to sites and to categories of sheaves (Sections 1 and 2). After introducing the induced topology (Section 3), we take up in Section 4 the comparison lemma, which will play an important role in Exposé IV. In Section 5, for the patient reader, we study certain commutative diagrams related to localized categories, of the type C/X . In Theorem 5.1 of Exposé I smallness hypotheses were imposed on the categories under consideration. These hypotheses are, in general, not satisfied by the sites encountered in practice, namely $\mathcal{U}$ -sites. This forces some circumlocutions in Sections 4.2, 4.3, and 4.4.

1 Continuous functors

Definition 1.1. Let C and C' be two $\mathcal{U}$ -sites, and let $u : C \rightarrow C'$ be a functor between the underlying categories. We say that u is *continuous* if, for every sheaf of sets F on C' , the presheaf

$$X \mapsto F \circ u(X)$$

on C is a sheaf.

This notion of continuity depends, a priori, on the universe $\mathcal{U}$ for which the two sites are $\mathcal{U}$ -sites. Proposition 1.5 shows that it does not depend on it.

1.1.1

Let

$$i : \widetilde{C} \longrightarrow \widehat{C}, \quad i' : \widetilde{C'} \longrightarrow \widehat{C'}$$

be the canonical inclusion functors from sheaves into presheaves. By Definition 1.1, the functor u is continuous if and only if there exists a functor

$$u_s : \widetilde{C'} \longrightarrow \widetilde{C}$$

such that

$$iu_s = u^*i', \quad u^*F = F \circ u$$

in the notation of Exposé I, 5.0.

Proposition 1.2. Let C be a small site, let C' be a $\mathcal{U}$ -site, and let $u : C \rightarrow C'$ be a functor between the underlying categories. The following properties are equivalent:

- i) The functor u is continuous.
- ii) For every object X of C , and every covering sieve $R \hookrightarrow X$, the morphism

$$u_! R \hookrightarrow u(X)$$

is bicobering in $\widehat{C'}$ (Exposé II, 5.2 and Exposé I, 5.1).¹

- iii) For every bicobering family $H_i \rightarrow K$, $i \in I$, in $\widehat{C}$, the family

$$u_! H_i \longrightarrow u_! K, \quad i \in I,$$

is bicobering in $\widehat{C'}$.

- iv) There exists a functor

$$u^s : \widetilde{C} \longrightarrow \widetilde{C'}$$

which commutes with inductive limits and “extends u ”, i.e. such that the diagram of canonical functors

$$\begin{array}{ccc} C & \xrightarrow{u} & C' \\ \downarrow \varepsilon_C & & \downarrow \varepsilon_{C'} \\ \widetilde{C} & \xrightarrow{u^s} & \widetilde{C'} \end{array}$$

is commutative up to isomorphism.

Moreover, when C is no longer necessarily a small site but is a $\mathcal{U}$ -site, one still has i) $\Leftrightarrow$ iv), and the functor u^s in iv) is necessarily a left adjoint of the functor

$$u_s : \widetilde{C'} \longrightarrow \widetilde{C},$$

and is therefore determined up to a unique isomorphism.

Proof. The proof of i) $\Leftrightarrow$ iv), when C is a $\mathcal{U}$ -site, will be given in Section 4.2. Suppose first that C is small. It is clear that iii) $\Rightarrow$ ii).

i) $\Rightarrow$ iii). Let $H_i \rightarrow K$, $i \in I$, be a bicobering family in $\widehat{C}$. For every sheaf F on C' , the presheaf $u^* F$ is a sheaf on C . Hence, by Exposé II, 5.3, the canonical map

$$\mathrm{Hom}(K, u^* F) \longrightarrow \prod_i \mathrm{Hom}(H_i, u^* F)$$

is bijective. By adjunction (Exposé I, 5.1), it follows that the canonical map

$$\mathrm{Hom}(u_! K, F) \longrightarrow \prod_i \mathrm{Hom}(u_! H_i, F)$$

is bijective. Hence, again by Exposé II, 5.3, the family $u_! H_i \rightarrow u_! K$, $i \in I$, is bicobering.

ii) $\Rightarrow$ i). Let X be an object of C , let $R \hookrightarrow X$ be a covering sieve, and let F be a sheaf on C' . By Exposé II, 5.3, the map

$$\mathrm{Hom}(u_! X, F) \longrightarrow \mathrm{Hom}(u_! R, F)$$

¹It would not necessarily be enough to say “covering” instead of “bicobering”; cf. Example 1.9.3 below.

is bijective. By adjunction (Exposé I, 5.1), it follows that

$$\mathrm{Hom}(X, u^*F) \longrightarrow \mathrm{Hom}(R, u^*F)$$

is bijective. Hence u^*F is a sheaf.

i) $\Rightarrow$ iv). Use the usual notation $\underline{a}$ and i , respectively $\underline{a}'$ and i' , for the associated-sheaf functors and the inclusions into presheaves. For every sheaf G on C , put

$$u^s G = \underline{a}' u_! i G.$$

For every sheaf F on C' , one has the following sequence of natural isomorphisms:

$$\begin{aligned} \mathrm{Hom}(G, u_s F) &\simeq \mathrm{Hom}(iG, i u_s F) \simeq \mathrm{Hom}(iG, u^* i' F) \\ &\simeq \mathrm{Hom}(u_! iG, i' F) \simeq \mathrm{Hom}(\underline{a}' u_! iG, F). \end{aligned}$$

The first isomorphism comes from the full faithfulness of i ; the second comes from the definition of u_s ; the third and fourth are obtained by adjunction. Therefore u^s is left adjoint to u_s . In particular, u^s commutes with inductive limits. For every presheaf K on C and every sheaf F on C' , one also has natural isomorphisms

$$\begin{aligned} \mathrm{Hom}(u^s \underline{a} K, F) &\simeq \mathrm{Hom}(\underline{a} K, u_s F) \simeq \mathrm{Hom}(K, i u_s F) \\ &\simeq \mathrm{Hom}(K, u^* i' F) \simeq \mathrm{Hom}(u_! K, i' F) \simeq \mathrm{Hom}(\underline{a}' u_! K, F), \end{aligned}$$

where the third isomorphism comes from the definition of u_s , and the others from adjunction. Hence there is a functorial isomorphism

$$u^s \underline{a} K \simeq \underline{a}' u_! K.$$

In particular, applying this isomorphism when K is representable and using Exposé I, 5.4(3), we obtain a commutative diagram up to isomorphism:

$$\begin{array}{ccc} C & \xrightarrow{u} & C' \\ \downarrow \varepsilon_C & & \downarrow \varepsilon_{C'} \\ \tilde{C} & \xrightarrow{u^s} & \tilde{C}'. \end{array}$$

iv) $\Rightarrow$ ii). Let

$$h : C \rightarrow \widehat{C}, \quad h' : C' \rightarrow \widehat{C}'$$

be the Yoneda functors. Consider the diagram

$$\begin{array}{ccc} C & \xrightarrow{u} & C' \\ \downarrow h & & \downarrow h' \\ \widehat{C} & \xrightarrow{u_!} & \widehat{C}' \\ \downarrow \underline{a} & & \downarrow \underline{a}' \\ \tilde{C} & \xrightarrow{u^s} & \tilde{C}'. \end{array}$$

The upper square is commutative (Remark 1.4(3)), and $\underline{a}h = \varepsilon_C$, $\underline{a}'h' = \varepsilon_{C'}$. For every object K of $\widehat{C}$ there is a canonical isomorphism (Exposé I, 3.4)

$$\varinjlim_{(h(X) \rightarrow K) \in \mathrm{Ob}(C/K)} h(X) \xrightarrow{\sim} K.$$

Since the functors $u_!$, u^s , $\underline{a}$, and $\underline{a}'$ commute with inductive limits, we obtain isomorphisms

$$\begin{aligned} \varinjlim_{(h(X) \rightarrow K) \in \text{Ob}(C/K)} u^s \underline{a} h(X) &\xrightarrow{\sim} u^s \underline{a} K, \\ \varinjlim_{(h(X) \rightarrow K) \in \text{Ob}(C/K)} \underline{a}' u_! h(X) &\xrightarrow{\sim} \underline{a}' u_! K. \end{aligned}$$

We have $u_! h = h' u$, and by hypothesis we have an isomorphism

$$\underline{a}' h' u \simeq u^s \underline{a} h.$$

Hence there is an isomorphism

$$u^s \underline{a} K \simeq \underline{a}' u_! K,$$

which is immediately checked to be functorial in K . Therefore the diagram

$$\begin{array}{ccc} \widehat{C} & \xrightarrow{u_!} & \widehat{C}' \\ \downarrow \underline{a} & & \downarrow \underline{a}' \\ \widetilde{C} & \xrightarrow{u^s} & \widetilde{C}' \end{array}$$

is commutative up to isomorphism. Since $\underline{a} i$ is isomorphic to the identity functor, one has an isomorphism

$$u^s \simeq \underline{a}' u_! i,$$

whence the uniqueness of u^s . Let $v : H \rightarrow K$ be a bicoverying morphism of $\widehat{C}$. Then $\underline{a}(v)$ is an isomorphism (Exposé II, 5.3), and hence $\underline{a}' u_!(v)$, being isomorphic to $u^s \underline{a}(v)$, is an isomorphism. Thus $u_!(v)$ is taken by $\underline{a}'$ to an isomorphism, and consequently $u_!(v)$ is bicoverying (Exposé II, 5.3). This proves ii).

The additional assertion in the case where C is small was proved during the proof. $\square$

1.2.1

The functor $u^s : \widetilde{C} \rightarrow \widetilde{C}'$ of Proposition 1.2(iv) can often be interpreted as an “inverse image” functor for a “morphism of topoi” $\widehat{C}' \rightarrow \widehat{C}$, cf. Exposé IV, 4.9. Its properties are summarized in the following proposition.

Proposition 1.3. Let $u : C \rightarrow C'$ be a continuous functor from a small site C to a $\mathcal{U}$ -site C' .

1) The functor u^s is left adjoint to the functor u_s .

2) There is a canonical isomorphism

$$u^s \simeq \underline{a}' u_! i.$$

3) There is a canonical isomorphism

$$u^s \underline{a} \simeq \underline{a}' u_!.$$

4) The functor u^s commutes with inductive limits.

5) When the functor $u_!$ is left exact (cf. Exposé I, 5.4(4)), the functor u^s is also left exact. More generally, u^s commutes with every type of finite projective limit with which $u_!$ commutes.

Proof. Assertions (1), (2), (3), and (4) were proved in the course of the proof of Proposition 1.2. Assertion (5) follows from the isomorphism in (2), taking into account that i commutes with projective limits and that $\underline{a}'$ commutes with finite projective limits (Exposé II, 4.1).

Remark 1.4. Combining Exposé I, 5.4(3), with Proposition 1.3(3), one obtains a diagram

$$\begin{array}{ccc} C & \xrightarrow{u} & C' \\ \downarrow h & & \downarrow h' \\ \widehat{C} & \xrightarrow{u_!} & \widehat{C'} \\ \downarrow \underline{a} & & \downarrow \underline{a'} \\ \widetilde{C} & \rightarrow & \widetilde{C'}, \end{array}$$

in which the upper square is commutative and the lower square is commutative up to isomorphism. But the functors $\underline{a}$, $\underline{a}'$, $u_!$, and u^s are defined only up to isomorphism. The reader may verify that the functors $\underline{a}$, $\underline{a}'$, and u^s may be chosen so that:

- 1) the composite functors $\underline{a}h$ and $\underline{a}'h'$ are injective on the sets of objects;
- 2) the diagram

$$\begin{array}{ccc} C & \xrightarrow{u} & C' \\ \downarrow \underline{a}h & & \downarrow \underline{a}'h' \\ \widetilde{C} & \xrightarrow{u^s} & \widetilde{C'} \end{array}$$

is commutative.

More precisely, one may choose $\underline{a}$ and $\underline{a}'$ so as to satisfy condition (1). Once $\underline{a}$ and $\underline{a}'$ have been chosen, one may choose u^s so as to satisfy condition (2).

Proposition 1.5. Let $\mathcal{U} \subset \mathcal{V}$ be two universes, let C and C' be two $\mathcal{U}$ -sites, and let $u : C \rightarrow C'$ be a functor between the underlying categories. Then u is continuous relative to $\mathcal{U}$ if and only if it is continuous relative to $\mathcal{V}$. Moreover, when u is continuous, let

$$u_{\mathcal{U}}^s \quad (\text{respectively } u_{\mathcal{V}}^s)$$

denote the functor between categories of $\mathcal{U}$ -sheaves (respectively, $\mathcal{V}$ -sheaves) introduced in Proposition 1.2(iv). Then the diagram

$$\begin{array}{ccc} \widetilde{C}_{\mathcal{U}} & \xrightarrow{u_{\mathcal{U}}^s} & \widetilde{C'}_{\mathcal{U}} \\ \downarrow & & \downarrow \\ \widetilde{C}_{\mathcal{V}} & \xrightarrow{u_{\mathcal{V}}^s} & \widetilde{C'}_{\mathcal{V}} \end{array} \tag{1.5.1}$$

where the vertical functors are the canonical inclusion functors, is commutative up to canonical isomorphism.

Proof. We give the proof only in the case where C is $\mathcal{U}$ -small. The general case will be proved in Section 4.3. It is clear that if the functor u^* takes every $\mathcal{V}$ -sheaf on C' to a $\mathcal{V}$ -sheaf on C , then it takes every $\mathcal{U}$ -sheaf on C' to a $\mathcal{U}$ -sheaf on C . Conversely, suppose that u^* takes every $\mathcal{U}$ -sheaf on C' to a $\mathcal{U}$ -sheaf on C . Write

$$\widehat{C}_{\mathcal{U}} \hookrightarrow \widehat{C}_{\mathcal{V}}, \quad \widehat{C'}_{\mathcal{U}} \hookrightarrow \widehat{C'}_{\mathcal{V}}$$

for the categories of $\mathcal{U}$ -presheaves and $\mathcal{V}$ -presheaves, and let $u_{\mathcal{U}!}$ and $u_{\mathcal{V}!}$ be the corresponding “inverse image” functors for $\mathcal{U}$ -presheaves and $\mathcal{V}$ -presheaves. There is a commutative diagram (cf. the explicit construction of $u_!$ in Exposé I, 5.1)

$$\begin{array}{ccc} \widehat{C}_{\mathcal{U}} & \xrightarrow{u_{\mathcal{U}!}} & \widehat{C'}_{\mathcal{U}} \\ \downarrow & & \downarrow \\ \widehat{C}_{\mathcal{V}} & \xrightarrow{u_{\mathcal{V}!}} & \widehat{C'}_{\mathcal{V}}. \end{array} \quad (*)$$

The inclusion functors from $\mathcal{U}$ -presheaves into $\mathcal{V}$ -presheaves are fully faithful and commute with projective limits. It then follows from Exposé II, 5.1(i), that a bicovering morphism of $\mathcal{U}$ -presheaves is a bicovering morphism of $\mathcal{V}$ -presheaves. Let $R \hookrightarrow X$ be a covering sieve of an object X of C . By Proposition 1.2(ii), the morphism $u_{\mathcal{U}!}R \rightarrow u_{\mathcal{U}!}X$ is bicovering. Using the commutativity of $(*)$ and what was just proved, we deduce that $u_{\mathcal{V}!}R \rightarrow u_{\mathcal{V}!}X$ is bicovering. Hence, by Proposition 1.2, the functor u^* takes $\mathcal{V}$ -sheaves on C' to $\mathcal{V}$ -sheaves on C . The commutativity of (1.5.1) follows from the commutativity of $(*)$, Exposé II, 3.6, and Proposition 1.3(2).

Proposition 1.6. Let C and C' be two $\mathcal{U}$ -sites, and let $u : C \rightarrow C'$ be a functor between the underlying categories. Consider the following conditions:

- i) u is continuous (cf. Definition 1.1 and Proposition 1.2).
- ii) For every covering family $(X_i \rightarrow X)$, $i \in I$, of C , the family $(uX_i \rightarrow uX)$, $i \in I$, of C' , is covering.

Then i) $\Rightarrow$ ii).

Suppose that the topology of C can be defined by a pretopology T (Exposé II, 1.3), and that the functor u commutes with fiber products.² Then i) and ii) are equivalent, and are also equivalent to the following condition:

- iii) The functor u takes the covering families of T to covering families of C' .

In particular, suppose that fiber products are representable in C , and that u commutes with fiber products. Then i) and ii) are equivalent.

Proof. We first show that i) $\Rightarrow$ ii). Let $(X_i \rightarrow X)$, $i \in I$, be a covering family of C . For every sheaf F on C' , u^*F is a sheaf. Hence the map

$$\mathrm{Hom}(X, u^*F) \longrightarrow \prod_i \mathrm{Hom}(X_i, u^*F)$$

is injective (Exposé II, 5.1). Therefore the map

$$\mathrm{Hom}(u_!X, F) \longrightarrow \prod_i \mathrm{Hom}(u_!X_i, F)$$

is injective. Thus the family $u_!X_i \rightarrow u_!X$, $i \in I$, is covering in C' (Exposé II, 5.1).

Every covering family of the pretopology T is a covering family of C , hence ii) $\Rightarrow$ iii). We prove iii) $\Rightarrow$ i). Let X be an object of C , and let $(X_i \rightarrow X)$, $i \in I$, be a covering family in

²In fact, it is enough that u commute with the fiber products involved in base changes of morphisms coming from covering families for T .

$\text{Cov}(X)$ (Exposé II, 1.3). The morphisms $X_i \rightarrow X$ are base-changeable, and u commutes with fiber products. Hence the fiber products

$$u(X_i) \times_{u(X)} u(X_j)$$

are representable and canonically isomorphic to $u(X_i \times_X X_j)$. Let F be a sheaf on C' . Then the diagram of sets

$$F(u(X)) \longrightarrow \prod_i F(u(X_i)) \rightrightarrows \prod_{i,j} F(u(X_i \times_X X_j))$$

is exact (Exposé II, 2.1; Exposé I, 3.5 and 2.12). Hence the diagram of sets

$$u^*F(X) \longrightarrow \prod_i u^*F(X_i) \rightrightarrows \prod_{i,j} u^*F(X_i \times_X X_j)$$

is exact. Thus u^*F is a sheaf (Exposé II, 2.4), proving i). The last assertion of Proposition 1.6 follows from the preceding argument and from the fact that, when fiber products are representable in C , all covering families of C^3 define on C a pretopology whose associated topology is the topology of C .

Proposition 1.7. Let C be a small site, let C' be a $\mathcal{U}$ -site, and let $u : C \rightarrow C'$ be a continuous functor. Let γ be an algebraic structure defined by finite projective limits, such that in the category of γ -objects of $\mathcal{U}\text{-Set}$, $\mathcal{U}$ -inductive limits are representable.⁴ We use the notation of Proposition II, 6.4. The functor u_s commutes with projective limits, and therefore defines a functor

$$u_s^\gamma : \widetilde{C}_\gamma \longrightarrow \widetilde{C'}_\gamma.$$

The functor u_s^γ admits a left adjoint u_γ^s , which has the following properties:

- 1) There is a canonical isomorphism

$$u_\gamma^s \xrightarrow{\sim} \underline{a}'_\gamma u_{\gamma!} i_\gamma,$$

and u_γ^s commutes with every finite projective limit with which $u_{\gamma!}$ commutes.

- 2) There is a canonical isomorphism

$$u_\gamma^s \underline{a}_\gamma \xrightarrow{\sim} \underline{a}'_\gamma u_{\gamma!}'.$$

- 3) The functor u_γ^s commutes with inductive limits.

- 4) If u^s is left exact, then the diagram (notation of Exposé II, 6.3.0)

$$\begin{array}{ccc} \widetilde{C}_\gamma & \xrightarrow{u_\gamma^s} & \widetilde{C'}_\gamma \\ \downarrow j^\sim & & \downarrow j'^\sim \\ \widetilde{C} & \xrightarrow{u^s} & \widetilde{C'} \end{array}$$

is commutative up to isomorphism, and the functor u_γ^s is exact.

³indexed by sets belonging to a suitable universe

⁴In fact, one can show that this condition is always satisfied.

- 5) Suppose that the functor $j^\sim$, respectively $j'^\sim$, has a left adjoint $\text{Lib}^\sim$, respectively $\text{Lib}'^\sim$ (Exposé II, 6.5). There is a canonical isomorphism

$$u_\gamma^s \text{Lib}^\sim \xrightarrow{\sim} \text{Lib}'^\sim u^s.$$

Moreover, when C is not necessarily a small site but is a $\mathcal{U}$ -site, the functor u_γ^γ admits a left adjoint u_γ^s . Finally, if $\mathcal{V}$ is a universe containing $\mathcal{U}$, and if $u_{\gamma\mathcal{U}}^s$, respectively $u_{\gamma\mathcal{V}}^s$, denotes the left adjoint of u_s relative to the universe $\mathcal{U}$, respectively $\mathcal{V}$, then the following diagram, analogous to diagram (1.5.1),

$$\begin{array}{ccc} \widetilde{C}_{\gamma\mathcal{U}} & \xrightarrow{u_{\gamma\mathcal{U}}^s} & \widetilde{C}'_{\gamma\mathcal{U}} \\ \downarrow & & \downarrow \\ \widetilde{C}_{\gamma\mathcal{V}} & \xrightarrow{u_{\gamma\mathcal{V}}^s} & \widetilde{C}'_{\gamma\mathcal{V}} \end{array} \quad (1.7.1)$$

is commutative up to canonical isomorphism.

Proof. In the case where C is a small site, the proof is left to the reader. It will be completed in Section 4.3 for the case where C is a $\mathcal{U}$ -site.

Notation 1.8. From now on we shall use the notation u_s also for the functor u_s^γ . This notation entails no risk of confusion, because the functor u_s , defined for sheaves of sets, commutes with finite projective limits.

Similarly, when u^s is left exact, we shall use the notation u^s also for the functor u_γ^s . This abuse of notation is justified by Proposition 1.7(4).

Example 1.9.1. Let X be a small topological space. Denote by $\text{Op}(X)$ the category of open subsets of X ; its objects are the open subsets of X , and its morphisms are inclusions. Give it the canonical topology. A family $(U_i \subset U)$, $i \in I$, is covering if and only if the open sets U_i cover U (Exposé II, 2.6). Thus sheaves of sets on $\text{Op}(X)$ are sheaves of sets in the sense of [TF]. Let $f : X \rightarrow Y$ be a continuous map. It gives a functor

$$f^{-1} : \text{Op}(Y) \longrightarrow \text{Op}(X),$$

which sends an open set U of Y to its inverse image under f . The functor f^{-1} is continuous (Proposition 1.6). The functor

$$f_s^{-1} : \text{Op}(X)^\sim \longrightarrow \text{Op}(Y)^\sim$$

is the direct image functor for sheaves in the sense of [TF]. The functor

$$(f^{-1})^s : \text{Op}(Y)^\sim \longrightarrow \text{Op}(X)^\sim$$

is the inverse image functor for sheaves in the sense of [TF]. The functor $(f^{-1})^s$ commutes with finite projective limits, by Proposition 1.3(5).

Example 1.9.2. Let X be a topological space and let $Y \hookrightarrow X$ be an open subset of X . Every open subset of Y is an open subset of X , hence there is a functor

$$\Phi : \text{Op}(Y) \longrightarrow \text{Op}(X).$$

The functor Φ is continuous (Proposition 1.6). The functor

$$\Phi_s : \mathrm{Op}(X)^\sim \longrightarrow \mathrm{Op}(Y)^\sim$$

is restriction to Y . The functor

$$\Phi_{\mathrm{ab}}^s : \mathrm{Op}(Y)_{\mathrm{ab}}^\sim \longrightarrow \mathrm{Op}(X)_{\mathrm{ab}}^\sim$$

(Exposé II, 6.3.3 and Exposé I, 1.7) is the functor “extension by zero outside Y ” introduced in [TF]. The functor

$$\Phi^s : \mathrm{Op}(Y)^\sim \longrightarrow \mathrm{Op}(X)^\sim$$

(Proposition 1.3) is analogous to extension by zero: for every sheaf H on Y and every point $y \in Y$, the stalk of $\Phi^s H$ at y is canonically isomorphic to the stalk of H at y ; for every point $x \in X$, $x \notin Y$, the stalk of $\Phi^s H$ at x is empty. The functor Φ^s is called “extension by the empty set outside Y ”. It commutes with fiber products and with finite products over a nonempty index set, but it does not take the final object of $\mathrm{Op}(Y)^\sim$ to the final object of $\mathrm{Op}(X)^\sim$.

Example 1.9.3. Let C and C' be two small sites, and let $u : C \rightarrow C'$ be a functor between the underlying categories which takes every covering family of C to a covering family of C' . In general, u is not continuous (cf. nevertheless Proposition 1.6). Here is a counterexample. Let S be an infinite set belonging to the universe $\mathcal{U}$, and let C be the full subcategory of the category of sets whose objects are the finite nonempty subsets of S . Give C the canonical topology. Observe that the covering families of C are the surjective families of maps, and that covering families are nonempty. Also observe that, up to isomorphism, there are only two constant sheaves on C : the sheaf with value the empty set and the sheaf with value a one-element set. Put $C = C'$, and let $u : C \rightarrow C'$ be a constant functor. Then u takes covering families of C to covering families of C' , but u is not continuous. Indeed, since the representable presheaves on C' are sheaves, for every object X of C' there exists a sheaf F on C' such that the cardinality of $F(X)$ is at least 2; consequently the presheaf u^*F is not a sheaf.

Continuation anchor. The next untranslated point is Exposé III, Section 2, “Cocontinuous functors”, beginning at source line 613 of 03/03.tex.