

[Scan page 526 is blank.]

478.

ON THE GEODESIC LINES ON AN ELLIPSOID.

[From the *Memoirs of the Royal Astronomical Society*, vol. XXXIX. (1872), pp. 31–53. Read January 13, 1871.]

[p. 493]

The fundamental equations, in regard to the geodesic lines on an ellipsoid, were established by Jacobi, viz., representing by a, b, c , the squares of the semiaxes, that is, taking the ellipsoid to be

$$\frac{x^2}{a} + \frac{y^2}{b} + \frac{z^2}{c} = 1$$

(where $a > b > c$), if we introduce the elliptic coordinates h, k , and write

$$\begin{aligned} \frac{x^2}{a+h} + \frac{y^2}{b+h} + \frac{z^2}{c+h} &= 1, \\ \frac{x^2}{a+k} + \frac{y^2}{b+k} + \frac{z^2}{c+k} &= 1, \end{aligned}$$

or, what is the same thing,

$$x^2 = \frac{a(a+h)(a+k)}{(a-b)(a-c)}, \quad y^2 = \frac{b(b+h)(b+k)}{(b-c)(b-a)}, \quad z^2 = \frac{c(c+h)(c+k)}{(c-a)(c-b)};$$

then, if β be an arbitrary constant, the differential equation of a geodesic line is

$$(1) \quad \text{const.} = \int dh \sqrt{\frac{h}{(a+h)(b+h)(c+h)(\beta+h)}} + \int dk \sqrt{\frac{k}{(a+k)(b+k)(c+k)(\beta+k)}};$$

and the expression for the length of any arc of the curve is given by

$$(2) \quad s = \int dh \sqrt{\frac{h(\beta+h)}{(a+h)(b+h)(c+h)}} + \int dk \sqrt{\frac{k(\beta+k)}{(a+k)(b+k)(c+k)}}.$$

I propose in the present Memoir to develop the theory to the extent of showing how we can, by means of the first of these equations, explain the course of the geodesic lines; and for given numerical values of a, b, c , calculate, construct, and exhibit in a drawing the course of these lines: I attend more particularly to the series of geodesic lines through an umbilicus (which lines pass also through the opposite umbilicus), and to the case where the semiaxes are connected by the equation $ac - b^2 = 0$, a relation which simplifies the formulae.

General Considerations as to the Course of the Lines.


1. It will be observed that h and k enter into the formulae symmetrically: it will be convenient to distinguish between these coordinates by considering h as extending between the values $-a, -b$; and k as extending between the values $-b, -c$. Thus:

$h = \text{const.}$ denotes a curve of curvature of the one kind, viz.

$h = -a$, the principal section ABA' (or major-mean section), $h = -b$, the curves UU' and $U''U'''$ (or portions of the umbilicar section $ACA'C'$); similarly,

$k = \text{const.}$ denotes a curve of curvature of the other kind, viz.

$k = -c$, the principal section CBC' (or minor-mean section), $k = -b$, the curves UU''' and $U'U''$ (remaining portions of the umbilicar section $ACA'C'$).



2. To any given (admissible) values of h, k there correspond eight points, situate in the eight octants of the surface respectively; but, unless the contrary is expressed, it is assumed that the coordinates x, y, z , are positive, and that the point is situate in the octant ABC .

3. The constant β may have any value from $+a$ to $+c$; viz., if it has a value between a and b , or say, if $-\beta$ has an h -value, then the geodesic lines wholly

[p. 494]

between the two ovals of the curve of curvature $h = -\beta$ (being in general an indefinite undulating curve touching each oval an indefinite number of times). Similarly, if β has any value between b and c , or say, if $-\beta$ has a k -value, then the geodesic line lies wholly between the two ovals of the curve of curvature $k = -\beta$ (being in general an indefinite undulating curve touching each oval an indefinite number of times). The intermediate case is when $\beta = b$, or say when $-\beta$ has the umbilicar value: here the geodesic line is in general an indefinite undulating curve passing an infinite number of times through the opposite umbilici U, U'' , or U', U''' ; to fix the ideas, say through U, U'' .

Lines through an Umbilicus.

4. I attend in particular to the last-mentioned case, and thus write $\beta = b$. We may in the formula (1) fix at pleasure a limit of each integral; and writing for convenience

$$\Pi(h) = \int_{-a}^h \frac{-dh}{b+h} \sqrt{\frac{h}{(a+h)(c+h)}}, \quad \Psi(k) = \int_b^k \frac{-dk}{b+k} \sqrt{\frac{k}{(a+k)(c+k)}},$$

the equation (1) becomes

$$\text{Const.} = \Pi(h) + \Psi(k).$$

5. It is to be observed, in regard to these integrals, that writing $h = -a + u$, we have

$$\Pi(h) = \int_0^u \frac{du}{a-b-u} \sqrt{\frac{a-u}{u(a-c-u)}},$$

which, for u small, is

$$= \frac{1}{a-b} \sqrt{\frac{a}{a-c}} \int_0^u \frac{du}{\sqrt{u}} = \frac{2\sqrt{u}}{a-b} \sqrt{\frac{a}{a-c}}.$$

By the assistance of this formula the value of the integral may be calculated by quadratures; viz., the formula gives the integral for any small value of u , and we can then proceed by the method of quadratures. The integral becomes infinite for $h = -b$: suppose that we have by quadratures calculated it up to $h = -b - m$ (m small), then to calculate it up to any value $-b - m + u$ nearer to $-b$, we have

$$\begin{aligned} \Pi(h) &= \Pi(-b-m) + \int_m^{m-u} \frac{du}{m-u} \sqrt{\frac{b+m-u}{(a-b-m+u)(b-c+m-u)}} \\ &= \Pi(-b-m) + \sqrt{\frac{b}{(a-b)(b-c)}} \left[\log \frac{m}{m-u} \right] \\ &= \Pi(-b-m) - \sqrt{\frac{b}{(a-b)(b-c)}} \log \left(1 - \frac{u}{m} \right), (1) \end{aligned}$$

where the second term is positive, and the value thus increases slowly with u , becoming as it should do $= \infty$ for $u = m$ or $h = -b$.

¹ Except when the contrary is stated, the symbol “log” denotes throughout the hyperbolic logarithm.

[p. 495]

6. Similarly in the second integral writing $k = -c - v$, we have

$$\Psi(k) = \int_0^v \frac{dv}{b-c-v} \sqrt{\frac{c+v}{(a-c-v)v}},$$

which, for v small, is

$$= \frac{1}{b-c} \sqrt{\frac{c}{a-c}} \int_0^v \frac{dv}{\sqrt{v}} = \frac{2\sqrt{v}}{b-c} \sqrt{\frac{c}{a-c}},$$

which is of the like assistance in regard to the calculation by quadratures. And if we have by quadratures calculated the integral up to $k = -b + n$ (n small), then, to calculate it up to any value $-b + n - v$ nearer to $-b$, we have

$$\begin{aligned} \Psi(k) &= \Psi(-b + n) + \int_n^{n-v} \frac{dv}{n-v} \sqrt{\frac{b-n+v}{(a-b+n-v)(b-c-n+v)}} \\ &= \Psi(-b + n) + \sqrt{\frac{b}{(a-b)(b-c)}} \left[\log \frac{n}{n-v} \right] \\ &= \Psi(-b + n) - \sqrt{\frac{b}{(a-b)(b-c)}} \log \left(1 - \frac{v}{n} \right), \end{aligned}$$

where the second term is positive, and the value thus increases slowly with v , becoming as it should do $= \infty$ for $v = n$, or $k = -b$.

7. It may be remarked that in $\Pi(h)$ and $\Psi(k)$ respectively the coefficient of the logarithmic term has in each case the same value $= \sqrt{\frac{b}{(a-b)(b-c)}}$. As regards the initial terms $\sqrt{u}$ and $\sqrt{v}$, the coefficients are $\frac{1}{a-b} \sqrt{\frac{a}{a-c}}$ and $\frac{1}{b-c} \sqrt{\frac{c}{a-c}}$ respectively, which are equal if $\frac{a}{(a-b)^2} = \frac{c}{(b-c)^2}$, or $ac - b^2 = 0$.

8. We may consider the two geodesic lines $\Pi(h) \pm \Psi(k) = \text{const.}$; suppose that these each of them pass through the point P , coordinates (h_0, k_0) in the ABC octant of the ellipsoid; then for one of them we have $\Pi(h) - \Psi(k) = \Pi(h_0) - \Psi(k_0)$, and for the other of them we have $\Pi(h) + \Psi(k) = \Pi(h_0) + \Psi(k_0)$: I attend first to the former: of these, say $\Pi(h) - \Psi(k) = C$ (where C is $= \Pi(h_0) - \Psi(k_0)$); and I say that this denotes the curve UPU'' . In fact, by reason of the equation $\Pi(h)$ and $\Psi(k)$ must both increase or both diminish; they both increase as h passes from h_0 to $-b$, and as k passes from k_0 to $-b$: we may have $h = -b + u$, $k = -b + v$ where u and v are both indefinitely small, the functions Π and Ψ being then indefinitely large, but $\Pi - \Psi = C$; and we have thus a series of points nearer and nearer to the umbilicus U ; that is, we have the portion PU of the curve. Tracing the curve in the opposite direction, or considering h as passing from h_0 to $-a$, and k as passing from k_0 to $-c$, then if C be positive, k will attain the value $-c$, before h attains the value $-a$, say that we have simultaneously $h = h_1$, $k = -c$; the equation is $\Pi(h_1) - \Psi(-c) = C$, that is, $\Pi(h_1) = C$; and the geodesic line then arrives at a point P_1 on the arc CB of the minor-mean

[p. 496]

principal section. The function Ψ then changes its sign, viz., considering it as always positive, the equation is now $\Pi(h) + \Psi(k) = C$, k passing from the value $-c$ towards $-b$, that is, $\Psi(k)$ increasing, and therefore $\Pi(h)$ diminishing, or h passing from h_1 towards the value $-a$; until at last, say for $k = k_2$, we have $h = -a$, that is, $C = \Pi(-a) + \Psi(k_2)$, or $C = \Psi(k_2)$; the geodesic line here arrives at a point P_2 on the arc BA' of the major mean principal section. The function Π then changes its sign, viz., Π denoting a positive function as before, the equation is $-\Pi(h) + \Psi(k) = C$; h passes from $-a$ towards $-b$, that is $\Pi(h)$ increases, and therefore $\Psi(k)$ must also increase, or k pass from k_2 towards $-b$: we have at length $h = -b - u$, $k = -b + v$, u and v being each indefinitely small; and therefore Π and Ψ each indefinitely large (but $-\Pi + \Psi = C$): that is, we arrive at the umbilicus U'' , completing the geodesic line UPU'' .

9. If instead of $C = +$ we have $C = -$, everything is similar, but the geodesic line proceeding from U in the direction UP will first cut the arc BA of the major mean section at a point P_1 ; then the arc BC of the minor mean section at a point P_2 ; and, finally, arrive as before at the umbilicus U'' .

10. The intermediate case is when $C = 0$, viz., we have here $\Pi(h) - \Psi(k) = 0$; the geodesic line here passes from U in the direction UP to B (extremity of the mean axis, $h = -a$, $k = -c$); Π and Ψ then each change their sign, so that, considering them as positive, the equation still is $\Pi(h) - \Psi(k) = 0$, and the geodesic line at last arrives at the umbilicus U'' . It will be easily understood how in the like manner $\Pi(h) + \Psi(k) = 0$ refers to the line $U'PU'''$.

11. Reverting to the equation $\Pi(h) - \Psi(k) = C$, or as I will now write it

$$\Pi(h) - \Psi(k) = \Pi(h_0) - \Psi(k_0),$$

which belongs to the portion UP of the geodesic line UPU'' , we require when h is $= -b - u$, and $k = -b + v$ (u and v indefinitely small) to know the ratio of the increments u, v ; this in fact serves to determine the direction at U of the geodesic line through the given point (h_0, k_0) .

12. For this purpose writing $h = -b - u$, we find

$$\Pi(h) = \int_u^{b-c} \frac{du}{u} \sqrt{\frac{b+u}{(a-b-u)(b-c+u)}},$$

which is

$$= \int_u^{b-c} \frac{du}{u} \left\{ \sqrt{\frac{b+u}{(a-b-u)(b-c+u)}} - \sqrt{\frac{b}{(a-b)(b-c)}} \right\} + \sqrt{\frac{b}{(a-b)(b-c)}} \log \frac{b-c}{u},$$

and, when u is indefinitely small, this is

$$= \int_0^{b-c} \frac{du}{u} \left\{ \sqrt{\frac{b+u}{(a-b-u)(b-c+u)}} - \sqrt{\frac{b}{(a-b)(b-c)}} \right\} + \sqrt{\frac{b}{(a-b)(b-c)}} \log \frac{b-c}{u}.$$

[p. 498]

Similarly, when $k = -b + v$, where v is indefinitely small

$$\Psi(k) = \int_0^{b-c} \frac{dv}{v} \left\{ \sqrt{\frac{b-v}{(a-b+v)(b-c-v)}} - \sqrt{\frac{b}{(a-b)(b-c)}} \right\} + \sqrt{\frac{b}{(a-b)(b-c)}} \log \frac{b-c}{v}.$$

13. Each of the integrals is of the dimension $-\frac{1}{2}$ in a, b, c , and the difference of the integrals may be represented by

$$M \sqrt{\frac{b}{(a-b)(b-c)}};$$

we have therefore

$$\Pi(h) - \Psi(k) = \sqrt{\frac{b}{(a-b)(b-c)}} \left\{ M + \log \frac{u}{v} \right\},$$

where

$$\begin{aligned} \sqrt{\frac{b}{(a-b)(b-c)}} M = & \int_0^{b-c} \frac{du}{u} \left\{ \sqrt{\frac{b+u}{(a-b-u)(b-c+u)}} - \sqrt{\frac{b}{(a-b)(b-c)}} \right\} \\ & - \int_0^{b-c} \frac{dv}{v} \left\{ \sqrt{\frac{b-v}{(a-b+v)(b-c-v)}} - \sqrt{\frac{b}{(a-b)(b-c)}} \right\}. \end{aligned}$$

14. Suppose the inferior limits replaced by the indefinitely small positive quantities ϵ, ϵ' respectively; and for the variable in the second integral write $-u$; then

$$M = \int_{-(b-c)}^{-\epsilon'} \frac{du}{u} + \int_{\epsilon}^{b-c} \frac{du}{u} \left\{ \sqrt{\frac{b+u}{(a-b-u)(b-c+u)}} - \sqrt{\frac{b}{(a-b)(b-c)}} \right\};$$

it being understood that the values $u = -\epsilon'$ to $u = +\epsilon$ are omitted from the integration: this is

$$= \int_{-(b-c)}^{b-c} \frac{du}{u} \left\{ \sqrt{\frac{b+u}{(a-b-u)(b-c+u)}} - \sqrt{\frac{b}{(a-b)(b-c)}} \right\} + \sqrt{\frac{b}{(a-b)(b-c)}} \log \frac{\epsilon}{\epsilon'} \cdot \frac{a-b}{b-c}.$$

with the same convention as to the integral; or if $\epsilon' = \epsilon$, then

$$M = M' - \sqrt{\frac{b}{(a-b)(b-c)}} \log \frac{a-b}{b-c},$$

where

$$\sqrt{\frac{b}{(a-b)(b-c)}} M' = \int_{-(b-c)}^{b-c} \frac{du}{u} \sqrt{\frac{b+u}{(a-b-u)(b-c+u)}} = \int_{-a}^{-c} \frac{dh}{b+h} \sqrt{\frac{h}{(a+h)(c+h)}},$$

the omitted elements being from $u = -\epsilon$ to $u = +\epsilon$; that is (in the language of Cauchy) we take for the integral its principal value. And hence

$$\Pi(h) - \Psi(k) = \sqrt{\frac{b}{(a-b)(b-c)}} \left\{ M' + \log \frac{u}{v} \right\}.$$

15. By what precedes this is $= \Pi(h_0) - \Psi(k_0)$; or if we write simply (h, k) instead of (h_0, k_0) , that is, consider the geodesic line UP , which is drawn from the

[p. 499]

point P , coordinates (h, k) , to the umbilicus U , the coordinates of a point consecutive to the umbilicus are $-b-u, -b+v$, where u, v are connected by the last-mentioned equation, in which M' is a transcendental function depending on (a, b, c) but independent of the particular geodesic line.

16. If for the geodesic line through the point B , or say for the B -geodesic


$$\frac{u}{u_0} = \frac{v}{v_0}, \text{ then } M' = -\log \frac{u_0}{v_0},$$

and we have in general

$$\Pi(h) - \Psi(k) = \sqrt{\frac{b}{(a-b)(b-c)}} \log \frac{v u_0}{u v_0},$$

a result which I proceed to further transform as follows:

If x_0, y_0, z_0 refer to the umbilicus U , then considering first the consecutive point P on the geodesic line (coordinates $-b-u, -b+v$) and next the consecutive



point Q on the umbilicar section, we have for these two points respectively,

$$dx_0 = \frac{\frac{1}{2}\sqrt{a}(v-u)}{\sqrt{(a-b)(a-c)}},$$

$$dy_0 = \frac{\sqrt{b}\sqrt{uv}}{\sqrt{(a-b)(b-c)}},$$

$$dz_0 = \frac{\frac{1}{2}\sqrt{c}(v-u)}{\sqrt{(b-c)(a-c)}};$$

$$\delta x_0 = \frac{\frac{1}{2}\sqrt{a}}{\sqrt{(a-b)(a-c)}},$$

$$\delta y_0 = 0,$$

$$\delta z_0 = \frac{-\frac{1}{2}\sqrt{c}}{\sqrt{(b-c)(a-c)}};$$

[p. 500]

say these are α, β, γ , and α', β', γ' ; and then

$$\begin{aligned}\alpha\alpha' + \beta\beta' + \gamma\gamma' &= \frac{1}{4} \left\{ \frac{a}{(a-b)(a-c)} + \frac{c}{(b-c)(a-c)} \right\} (v-u) \\ &= \frac{1}{4} \cdot \frac{(v-u)}{a-c} \left\{ \frac{a}{a-b} + \frac{c}{b-c} \right\} = \frac{1}{4} \cdot \frac{b(v-u)}{(a-b)(b-c)}, \\ \alpha^2 + \beta^2 + \gamma^2 &= \frac{1}{4} \left\{ \frac{a}{(a-b)(a-c)} + \frac{c}{(b-c)(a-c)} \right\} (v-u)^2 + \frac{buv}{(a-b)(b-c)} \\ &= \frac{1}{4} \cdot \frac{b(v-u)^2}{(a-b)(b-c)} + \frac{buv}{(a-b)(b-c)} \\ &= \frac{b}{(a-b)(b-c)} \cdot \frac{1}{4} (u+v)^2, \\ \alpha'^2 + \beta'^2 + \gamma'^2 &= \frac{b}{(a-b)(b-c)} \cdot \frac{1}{4},\end{aligned}$$

whence

$$\sqrt{\alpha^2 + \beta^2 + \gamma^2} \sqrt{\alpha'^2 + \beta'^2 + \gamma'^2} = \frac{\frac{1}{4}(u+v)b}{(a-b)(b-c)},$$

and hence

$$\cos \phi = \frac{\alpha\alpha' + \beta\beta' + \gamma\gamma'}{\sqrt{\alpha^2 + \beta^2 + \gamma^2} \sqrt{\alpha'^2 + \beta'^2 + \gamma'^2}} = \frac{v-u}{u+v},$$

that is,

$$\cos(180^\circ - \phi) = \frac{u-v}{u+v}, \quad \text{or} \quad \tan^2 \frac{1}{2}\phi = \frac{v}{u},$$

where it U is the umbilicus, P the consecutive point $-b-u$, $-b+v$, and UQ the element of the umbilicar principal section, $\phi = \angle PUQ$, $180^\circ - \phi = \angle PUQ'$. For the B -geodesic we have

$$2 \log \tan \frac{1}{2}\phi_0 = \log \frac{v_0}{u_0} = M'.$$

17. The foregoing equation for $\Pi(h) - \Psi(k)$ now becomes

$$\Pi(h) - \Psi(k) = \sqrt{\frac{b}{(a-b)(b-c)}} \log \frac{\tan^2 \frac{1}{2}\phi_0}{\tan^2 \frac{1}{2}\phi},$$

viz., ϕ_0 is the south azimuth of the B -geodesic at the umbilicus, a mere function of (a, b, c) and ϕ is the south azimuth at the umbilicus, of the geodesic line under consideration, so that we may consider the geodesic line to be determined by the south azimuth ϕ as its parameter.

[p. 501]

Formulæ for the case $ac - b^2 = 0$.

18. I annex the following investigation in regard to the case $ac - b^2 = 0$.

We have in general

$$\begin{aligned} \frac{1}{\sqrt{b(a-b)(b-c)}} \frac{d}{dx} \log \frac{\sqrt{-x(a-b)(b-c)} + \sqrt{-b(a+x)(c+x)}}{\sqrt{-x(a-b)(b-c)} - \sqrt{-b(a+x)(c+x)}} &= -\frac{1}{b} \cdot \frac{1}{\sqrt{x(a+x)(c+x)}} \\ &+ \frac{1}{b} \cdot \frac{1}{b+x} \sqrt{\frac{x}{(a+x)(c+x)}} \\ &- \frac{1}{bx+ac} \sqrt{\frac{x}{(a+x)(c+x)}}. \end{aligned}$$

In fact, denoting the logarithm by $\log \frac{P+Q}{P-Q}$, we have

$$\frac{d}{dx} \log \frac{P+Q}{P-Q} = \frac{2(PQ' - P'Q)}{P^2 - Q^2},$$

where

$$\begin{aligned} 2(PQ' - P'Q) &= 2PQ \left(\frac{Q'}{Q} - \frac{P'}{P} \right) = \sqrt{x(a+x)(c+x)} b(a-b)(b-c) \left(\frac{Q'}{Q} - \frac{P'}{P} \right) \\ &= \frac{\sqrt{b(a-b)(b-c)}}{\sqrt{x(a+x)(c+x)}} (x^2 - ac), \end{aligned}$$

and

$$P^2 - Q^2 = -x(a-b)(b-c) + b(a+x)(c+x) = (bx+ac)(b+x);$$

that is

$$\begin{aligned} \frac{2(PQ' - P'Q)}{P^2 - Q^2} &= \frac{\sqrt{b(a-b)(b-c)}}{\sqrt{x(a+x)(c+x)}} \cdot \frac{x^2 - ac}{(bx+ac)(b+x)} \\ &= \frac{\sqrt{b(a-b)(b-c)}}{\sqrt{x(a+x)(c+x)}} \left\{ -\frac{1}{b} + \frac{x}{b(b+x)} + \frac{x}{bx+ac} \right\}, \end{aligned}$$

which proves the theorem.

19. Hence in the particular case $ac = b^2$ we have

$$\begin{aligned} \frac{1}{\sqrt{b(a-b)(b-c)}} \log \frac{\sqrt{-h(a-b)(b-c)} + \sqrt{-b(a+h)(c+h)}}{\sqrt{-h(a-b)(b-c)} - \sqrt{-b(a+h)(c+h)}} &= -\frac{1}{b} \int_{-a}^h \frac{dh}{\sqrt{h(a+h)(c+h)}} \\ &+ \frac{2}{b} \int_{-a}^h \frac{dh}{b+h} \sqrt{\frac{h}{(a+h)(c+h)}}, \end{aligned}$$

[p. 502]

that is

$$\Pi(h) = \frac{1}{2} \int_{-a}^h \frac{dh}{\sqrt{h(a+h)(c+h)}} + \frac{1}{2} \sqrt{\frac{b}{(a-b)(b-c)}} \log \frac{\sqrt{-h(a-b)(b-c)} + \sqrt{-b(a+h)(c+h)}}{\sqrt{-h(a-b)(b-c)} - \sqrt{-b(a+h)(c+h)}}$$

or say

$$\Pi(h) = \frac{1}{2} \int_{-a}^h \frac{dh}{\sqrt{h(a+h)(c+h)}} + \frac{1}{2} \sqrt{\frac{b}{(a-b)(b-c)}} \log \frac{1+H}{1-H},$$

where

$$H^2 = \frac{b}{(a-b)(b-c)} \cdot \frac{(a+h)(c+h)}{h},$$

viz. we see that $\Pi(h)$ depends on the more simple integral

$$\int \frac{dh}{\sqrt{h(a+h)(c+h)}}.$$

20. Similarly

$$\begin{aligned} \frac{1}{\sqrt{b(a-b)(b-c)}} \log \frac{\sqrt{-k(a-b)(b-c)} + \sqrt{-b(a+k)(c+k)}}{\sqrt{-k(a-b)(b-c)} - \sqrt{-b(a+k)(c+k)}} &= -\frac{1}{b} \int_b^{-c} \frac{dk}{\sqrt{k(a+k)(c+k)}} \\ &+ \frac{2}{b} \int_b^{-c} \frac{dk}{b+k} \sqrt{\frac{k}{(a+k)(c+k)}}, \end{aligned}$$

that is

$$\Psi(k) = \frac{1}{2} \int_b^{-c} \frac{dk}{\sqrt{k(a+k)(c+k)}} + \frac{1}{2} \sqrt{\frac{b}{(a-b)(b-c)}} \log \frac{\sqrt{-k(a-b)(b-c)} + \sqrt{-b(a+k)(c+k)}}{\sqrt{-k(a-b)(b-c)} - \sqrt{-b(a+k)(c+k)}},$$

or say

$$\Psi(k) = \frac{1}{2} \int_b^{-c} \frac{dk}{\sqrt{k(a+k)(c+k)}} + \frac{1}{2} \sqrt{\frac{b}{(a-b)(b-c)}} \log \frac{1+K}{1-K},$$

where

$$K^2 = \frac{b}{(a-b)(b-c)} \cdot \frac{(a+k)(c+k)}{k},$$

that is, $\Psi(k)$ depends on the more simple integral,

$$\int \frac{dk}{\sqrt{k(a+k)(c+k)}}.$$

Write $h = -b - u$, $k = -b + v$, where u and v are indefinitely small, then

$$\Pi(h) - \Psi(k) = \frac{1}{2} \int_{-a}^{-c} \frac{dh}{\sqrt{h(a+h)(c+h)}} + \frac{1}{2} \sqrt{\frac{b}{(a-b)(b-c)}} \log \frac{1+U}{1-U} \cdot \frac{1-V}{1+V},$$

[p. 503]

where

$$U^2 = \frac{\left(1 + \frac{u}{a-b}\right) \left(1 + \frac{u}{b-c}\right)}{1 + \frac{u}{b}} = 1 - \frac{bu^2}{(a-b)(b-c)(b+u)} \quad (\text{attending to } ac = b^2),$$

and

$$V^2 = \frac{\left(1 + \frac{v}{a-b}\right) \left(1 - \frac{v}{b-c}\right)}{1 - \frac{v}{b}} = 1 - \frac{bv^2}{(a-b)(b-c)(b-v)}.$$

21. Comparing with the result obtained for the general case the two agree, if only

$$\int_{-a}^{-c} \frac{dh + b \sqrt{1 + \frac{(a+h)(c+h)}{h}}}{b+h} = \frac{1}{2} \int_{-a}^{-c} \frac{dh}{\sqrt{h(a+h)(c+h)}},$$

where on the left-hand side the integral has its principal value: a result which must therefore hold good when $ac = b^2$.

Calculation of the Umbilicar Geodesics for Ellipsoid $a : b : c = 4 : 2 : 1$.

22. As a specimen of the way in which we may, on a given ellipsoid, calculate the course of a geodesic line, I take the semiaxes to be as $2 : \sqrt{2} : 1$, or, for convenience, $a = 1000$, $b = 500$, $c = 250$; and, considering the geodesic lines through the umbilicus, I calculate by quadratures the functions

$$\Pi(h) = 100,000 \int_{-1000}^h \frac{-dh}{500+h} \sqrt{\frac{h}{(1000+h)(250+h)}},$$

$$\Psi(k) = 100,000 \int_b^k \frac{-dk}{500+k} \sqrt{\frac{k}{(1000+k)(250+k)}}.$$

The results do not pretend to minute accuracy: I have not attempted to estimate or correct for any error occasioned by the intervals (10 units) being too large; and there may possibly be accidental errors.

[p. 504]

TABLE I.

$-h$	Π'	$\Pi(h)$	$-h$	Π'	$\Pi(h)$	$-h$	Π'	$\Pi(h)$
1000	∞	0	840	27.6	6746	630	51.5	13972
999	231.4	462	830	27.8	7023	620	54.1	14499
998	164	659	820	27.9	7301	610	59.2	15066
997	134.2	809	810	28.1	7582	600	65.5	15689
996	116.5	934	800	28.4	7865	590	72.1	16377
995	104.4	1044	790	28.7	8151	580	80.9	17142
990	74.6	1492	780	29.2	8440	570	93.0	18011
980	54.0	2135	770	29.7	8735	560	106.8	19010
970	45.1	2630	760	30.3	9035	550	127.6	20183
960	40.1	3056	750	31.0	9341	540	159.1	21616
950	36.6	3439	740	31.8	9655	530	211.5	23469
940	34.2	3794	730	32.6	9977	520	316.7	26111
930	32.5	4127	720	33.6	10308	510	632.7	30858
920	31.2	4446	710	34.7	10650	505	1265	35602
910	30.2	4753	700	36.0	11004	504	1581.2	37014
900	29.4	5051	690	37.4	11371	503	2107.7	38834
890	28.8	5342	680	39.0	11754	502	3162.3	41398
880	28.4	5628	670	40.8	12153	501	6324.1	45792
870	28.1	5911	660	42.0	12567	500	∞	∞
860	27.9	6190	650	45.4	13005			
850	27.8	6469	640	48.2	13473			

[p. 505]

TABLE II.

$-k$	Ψ'	$\Psi(k)$	$-k$	Ψ'	$\Psi(k)$	$-k$	Ψ'	$\Psi(k)$
250	∞	0	320	45.5	4655	440	107.1	12207
251	232.2	462	330	46.1	5114	450	127.9	13383
252	165.5	661	340	47.3	5581	460	159.2	14818
253	136.0	811	350	48.9	6062	470	211.6	16673
254	118.6	939	360	51.1	6562	480	316.7	19314
255	106.8	1051	370	53.8	7086	490	632.7	24062
260	78.1	1514	380	57.2	7641	495	1265.0	28806
270	60.5	2207	390	61.4	8235	496	1581.2	30218
280	51.7	2768	400	66.7	8875	497	2108.1	32037
290	48.2	3268	410	73.2	9575	498	3162.3	34602
300	46.3	3741	420	81.6	10349	499	6234.1	38995
310	45.5	4200	430	91.4	11214	500	∞	∞

23. But it is obviously convenient to revert these Tables so as to have for the common arguments a series of uniformly increasing values of Π or Ψ , viz., we obtain by interpolation the values of h and k belonging to the given values of Π or Ψ , and thus obtain the following Table. Here, in any line of the Table the values of h , k are such that $\Pi(h) - \Psi(k) = 0$, viz., the values in question belong to successive points of the B -geodesic. And to obtain the values for any other geodesic line $\Pi(h) - \Psi(k) = \pm 500m$, we have only to take each value of k from the line m lines above or below the line from which h is taken; and similarly the table gives at once the values belonging to a geodesic line $\Pi(h) + \Psi(k) = 500m$.

[p. 506]

TABLE III.

$\Pi = \Psi =$	h	D.	k	D.	$\Pi = \Psi =$	h	D.	k	D.
0	1000		250		13000	650.1		446.7	
		1.2		1.4			20.6		7.8
500	998.8		251.4		14000	629.5		454.5	
		3.4		3.1			18.3		6.5
1000	995.4		254.5		15000	611.2		461.0	
		5.5		5.2			15.7		5.3
1500	989.9		259.7		16000	595.5		466.3	
		7.8		7.3			13.6		4.9
2000	982.1		267.0		17000	581.9		471.2	
		9.5		8.2			11.8		3.8
2500	972.6		275.2		18000	570.1		475.0	
		11.5		9.4			10.0		3.8
3000	961.1		284.6		19000	560.1		478.8	
		12.8		10.3			8.5		2.6
3500	948.3		294.9		20000	551.6		481.4	
		14.5		10.7			7.3		2.2
4000	933.8		305.6		21000	544.3		483.6	
		15.6		11.0			6.2		2.0
4500	918.2		316.6		22000	538.1		485.6	
		16.5		10.9			4.9		2.2
5000	901.7		327.5		23000	533.2		487.8	
		17.2		10.8			4.6		2.1
5500	884.5		338.3		24000	528.6		489.9	
		17.7		10.4			8.1		2.1
6000	866.8		348.7		26000	520.5		492.0	
		17.9		10.1			4.5		2.1
6500	848.9		358.8		28000	516.0		494.1	
		18.1		9.5			4.2		1.7
7000	830.8		368.3		30000	511.8		495.8	
		17.9		9.2			3.0		1.2
7500	812.9		377.5		32000	508.8		497.0	
		17.6		8.6			2.1		0.8
8000	795.3		386.1		34000	506.7		497.8	
		17.3		8.0			2.0		0.5
8500	778.0		394.1		36000	504.7		498.3	
		16.8		7.7			1.2		0.5
9000	761.2		401.8		38000	503.5		498.8	
		16.3		7.1			1.0		
9500	744.9		408.9		39000			499.0	
		15.6		6.6					
10000	729.3		415.5		40000	502.5			
		14.9		6.2			0.6		
10500	714.4		421.7		42000	501.9			
		14.3		5.9			0.6		
11000	700.1		427.6		44000	501.4			
		13.5		5.3			0.4		
11500	686.6		432.9		45800	501.0			
		12.8		5.0					
12000	673.8		437.9		∞	500		500	
		23.7		8.8					

[p. 507]

Graphical Construction: Projection on the Umbilicar Plane.

24. The most convenient mode of delineation of the geodesic lines is obtained by projecting them orthogonally on the umbilicar plane: the contour of the figure is here the umbilicar section, or ellipse

$$\frac{x^2}{a} + \frac{z^2}{c} = 1;$$

and the curves of curvature of each series are projected into elliptic arcs lying within the ellipse in question, the one set cutting at right angles the axes AA' , the other cutting at right angles the axes CC' ; the equations of the complete ellipses being

$$x^2 \frac{a-b}{a(a+h)} + z^2 \frac{c-b}{c(c+h)} - 1 = 0$$

and

$$x^2 \frac{a-b}{a(a+k)} + z^2 \frac{c-b}{c(c+k)} - 1 = 0.$$

25. I constructed, by means of the table, a drawing of this kind for the ellipsoid $a, b, c = 1000, 500, 250$, the lengths $\sqrt{a}$ and $\sqrt{c}$ being taken to be 12 inches and 6 inches respectively: the process consists in taking from the table for a series of values $\Pi = \Psi$ (say $\Pi = \Psi = 1000, = 2000$ &c.), the values of h and k , laying down for such values the elliptic arcs which represent the two curves of curvature respectively, thus dividing the bounding ellipse into a series of curvilinear rectangles, and then obtaining the geodesic lines by drawing the diagonals of these rectangles, and of course rounding off the corners so as to form continuous curves. The Plate shows on a reduced scale so much of the drawing as is comprised within a quadrant of the bounding ellipse (viz. it is a representation of an octant of the ellipsoid).

Elliptic-Function Formulæ.

26. I have in all that precedes abstained from the use of elliptic functions, since obviously the form $\sqrt{1-k^2\sin^2}$ of the radical of an elliptic function is in nowise specially appropriate to the present question. But (more particularly in the above-mentioned case $ac-b^2=0$, where the radical is $\sqrt{h(a+h)(c+h)}$ without any exterior factor $b+h$ in the denominator) the formulæ are expressible easily and elegantly by elliptic functions, and it is desirable to make the transformation. Reverting to the formulæ which, in the case in question (viz. when $ac-b^2=0$), give the values of $\Pi(h)$ and $\Psi(k)$; and writing therein $h = -a + (a-c)\sin^2\phi$, $k = -a + (a-c)\sin^2\psi$, also

$$\kappa = \sqrt{1 - \frac{c}{a}}, \quad \text{or} \quad \frac{c}{a} = 1 - \kappa^2 = \kappa'^2,$$

we have

$$\begin{aligned} \int_{-a}^h \frac{dh}{\sqrt{h(a+h)(c+h)}} &= 2 \int_0^\phi \frac{d\phi}{\sqrt{a - (a-c)\sin^2\phi}} = \frac{2}{\sqrt{a}} F(\kappa, \phi), \\ \int_b^k \frac{dk}{\sqrt{k(a+k)(c+k)}} &= 2 \int_{\frac{1}{2}\pi}^\psi \frac{d\psi}{\sqrt{a - (a-c)\sin^2\psi}} = \frac{2}{\sqrt{a}} \{F_1(\kappa) - F(\kappa, \psi)\}. \end{aligned}$$

[p. 508]

27. Hence

$$\Pi(h) = \frac{1}{\sqrt{a}} F(\kappa, \phi) + \frac{1}{2\sqrt{a(1-\kappa')}} \log \frac{1+H}{1-H},$$

where

$$H = \frac{(1-\kappa') \sin \phi \cos \phi}{\sqrt{1-\kappa'^2 \sin^2 \phi}};$$

(observe, as h passes from $-a$ to $-b$, ϕ passes from $\phi = 0$ to $\sin^2 \phi = \frac{1}{1+\kappa'}$, and H from $H = 0$ to $H = 1$).

Similarly

$$\Psi(k) = \frac{1}{\sqrt{a}} \{F_1(\kappa) - F(\kappa, \psi)\} + \frac{1}{2\sqrt{a(1-\kappa')}} \log \frac{1+K}{1-K},$$

where

$$K = \frac{(1+\kappa') \sin \psi \cos \psi}{\sqrt{1-\kappa'^2 \sin^2 \psi}};$$

and as k passes from $-c$ to $-b$, ψ passes from $\frac{1}{2}\pi$ to $\sin^2 \psi = \frac{1}{1+\kappa'}$, and K from 0 to 1.

28. The before-mentioned identical equation

$$\int_{-a}^{-c} \frac{dh}{b+h} \sqrt{\frac{h}{(a+h)(c+h)}} = \frac{1}{2} \int_{-a}^{-c} \frac{dh}{\sqrt{h(a+h)(c+h)}}$$

is by the same transformation converted into

$$\int_0^{\frac{1}{2}\pi} \frac{1 - (1-\kappa') \sin^2 \phi}{1 - (1+\kappa') \sin^2 \phi} \cdot \frac{d\phi}{\sqrt{1-\kappa'^2 \sin^2 \phi}} = 0.$$

To prove this, I remark that the equation is

$$0 = \int_0^{\frac{1}{2}\pi} d\phi \frac{\frac{1-\kappa'}{1+\kappa'} + \frac{2\kappa'}{1+\kappa'} \cdot \frac{1}{1 - (1+\kappa') \sin^2 \phi}}{\Delta \phi},$$

viz. this is

$$0 = \frac{1-\kappa'}{1+\kappa'} F_1 + \frac{2\kappa'}{1+\kappa'} \Pi_1(-1-\kappa'),$$

or, what is the same thing,

$$\Pi_1(-1-\kappa') = -\frac{1-\kappa'}{2\kappa'} F_1,$$

where $\Pi_1(-1-\kappa')$ denotes the principal value of the integral

$$\int_0^{\frac{1}{2}\pi} d\phi \frac{1}{1 - (1+\kappa') \sin^2 \phi} \cdot \frac{1}{\Delta \phi}.$$

[p. 509]

Now (Leg. *Fonct. Ellip.*, t. i., p. 71), we have

$$\Pi_1(-\kappa^2 \sin^2 \theta) + \Pi_1\left(-\frac{1}{\sin^2 \theta}\right) = F_1,$$

where, upon examination, it will appear that $\Pi_1\left(-\frac{1}{\sin^2 \theta}\right)$ in fact represents the principal value of the integral.

Writing herein $\sin^2 \theta = \frac{1}{1+\kappa'}$, and therefore $\cos^2 \theta = \frac{\kappa'}{1+\kappa'}$, or $\tan^2 \theta = \kappa'$, this is

$$\Pi_1(-1+\kappa') + \Pi_1(-1-\kappa') = F_1,$$

and the formula (p'), p. 141, attributing therein to θ the foregoing value, becomes

$$\Pi_1(-1 - \kappa') = \theta_1 + \frac{1}{\kappa'} \{F_1 E(\theta) - E_1 F(\theta)\}.$$

But θ is the value for the bisection of the function F_1 , viz., we have

$$\begin{aligned} 2F(\theta) &= F_1, \\ 2E(\theta) &= E_1 + 1 - \kappa', \end{aligned}$$

whence

$$F_1 E(\theta) - E_1 F(\theta) = \frac{1}{2}(1 - \kappa') F_1,$$

or the formula in question gives

$$\Pi_1(-1 + \kappa') = \frac{1 + \kappa'}{2\kappa'} F_1,$$

whence

$$\Pi_1(-1 - \kappa') = -\frac{1 - \kappa'}{2\kappa'} F_1,$$

the result which was to be proved.

29. The value of M' (observing that $\sqrt{(a-b)(b-c)} = \frac{b}{\sqrt{a-c}} = \frac{b}{\sqrt{a}\kappa}$) is

$$\frac{1}{\sqrt{a(1-\kappa')}} M' = \frac{1}{2} \int_{-a}^{-c} \frac{dh}{\sqrt{h(a+h)(c+h)}},$$

which is

$$= \frac{1}{2} \cdot \frac{2}{\sqrt{a}} F_1(\kappa),$$

that is we have

$$M' = (1 - \kappa') F_1(\kappa),$$

or, what is the same thing,

$$\log \tan \frac{1}{2} \phi_0 = \frac{1 - \kappa'}{2} F_1(\kappa),$$

that is

$$\tan \frac{1}{2} \phi_0 = e^{\frac{1-\kappa'}{2} F_1(\kappa)}$$

(ϕ_0 the South azimuth of the B -geodesic at the umbilicus).

30. I purposely calculated the Table by quadratures as being a method available where the equation $ac - b^2 = 0$ is not satisfied; but in the present case, where this

[p. 510]

equation is satisfied, the table might have been calculated from Legendre's Tables of Elliptic Integrals. Observe that $a = 1000$, $b = 500$, $c = 250$, gives $\kappa = \frac{1}{2}\sqrt{3}$ or angle of modulus $= 60^\circ$. As an instance of the comparison ⁽¹⁾, suppose $h = -800$, then $\sin^2 \phi = \frac{200}{750}$, $\log \sin \phi = 9.71298$, $\phi = 31^\circ 5'$.

$$\sqrt{\frac{b}{(a-b)(b-c)}} = \sqrt{\frac{500}{500 \cdot 250}} = \frac{\sqrt{10}}{50} = .06325,$$

$$H^2 = \frac{500 \cdot 200 \cdot 550}{800 \cdot 500 \cdot 250} = \frac{110}{200}, \quad \log = \bar{1}.87018,$$

$$H = .7416 \frac{1+H}{1-H} = \frac{1.7416}{.2584} = 6.7582,$$

$$\Pi(h) = .03163 F(31^\circ 5') + .03163 \text{ h. l. } 6.7582,$$

$$\begin{aligned}
 F\,31^\circ &= .56166 \\
 &163 \\
 F\,31^\circ 5' &= .56329 \\
 \text{h. l. } 6.7582 &= 1.91075 \\
 &2.47404 \\
 \times \text{ by } &.03163 \\
 &.0782043
 \end{aligned}$$

or multiplying by 100,000 (factor introduced into my Table) this is = 7820.43. The value $\Pi(-800) = 7864$ given by my Table agrees sufficiently well with this, the correct value.

31. I calculate also the angle ϕ_0 , viz. we have

$$\begin{aligned}
 \text{h. l. } \tan \frac{1}{2}\phi_0 &= \frac{1 - \kappa'}{2} F_1 \kappa = \frac{1}{4} F_1(60^\circ). \quad \text{Leg. vol. iii. Table viii.} \\
 &= \frac{1}{4} \cdot 2.15651 = .53913,
 \end{aligned}$$

whence by Leg. Table iv.

$$\begin{aligned}
 \frac{1}{2}\phi_0 &= 45^\circ + \frac{1}{2} \cdot 29^\circ 29' .64 \\
 &= 59^\circ 44' .82
 \end{aligned}$$

or

$$\phi_0 = 119^\circ 29' .64.$$

This exceeds 90° , and since at the umbilicus the tangent plane is at right angles to the plane of projection, the B -geodesic should in the drawing proceed (as it in fact does) from U in the sense UC , touching the bounding ellipse at the point U .

¹ In the present calculation, $\log$ denotes an ordinary logarithm, the hyperbolic logarithm being distinguished as **h. l.**

[Plate VII: graphical projection of geodesic lines on an ellipsoid $a : b : c = 4 : 2 : 1$ within a quadrant of the umbilicar ellipse, showing the network of curves of curvature and their diagonal geodesics; numerical labels along the bounding ellipse mark the values $\Pi = \Psi = 1000, 2000, \dots$ Cayley's Papers VII.]

[p. 511]

479.

THE SECOND PART OF A MEMOIR ON THE DEVELOPMENT OF THE DISTURBING FUNCTION IN THE LUNAR AND PLANETARY THEORIES.

[From the *Memoirs of the Royal Astronomical Society*, vol. XXXIX. (1872), pp. 55–74. Read January 12, 1872.]

The present communication is a sequel to my paper, “The First Part of a Memoir on the Development of the Disturbing Function in the Lunar and Planetary Theories,” *Memoirs R.A.S.*, vol. XXVIII. (1859), pp. 187–215, [214], and I have therefore entitled it as above, but it, in fact, relates only to the Planetary Theory. In the First Part, I gave in effect, but not explicitly, an expression for the general coefficient $D(j, j')$ in terms of the coefficients of the multiple cosines of θ in the expansions of the several powers $(r^2 + r'^2 - 2rr' \cos \theta)^{-s-\frac{1}{2}}$, or say $(a^2 + a'^2 - 2aa' \cos \theta)^{-s-\frac{1}{2}}$; viz., at the foot of page 208 I speak of the term involving $\cos(jU + j'U')$ as having a certain given value; the term in question is $D(j, j') \cos(jU + j'U')$; and consequently the expression for $D(j, j')$ is

$$D(j, j') = \Sigma \frac{\Pi_{\frac{1}{2}}(x - \frac{1}{2})}{\Pi x} \eta^{2x} \Sigma M_x^\xi R_x^\xi;$$

the omission was, however, a material one, inasmuch as this expression for the general coefficient serves to connect my formulae with Leverrier’s development, *Annales de l’Observ. de Paris*, t. i. (1855), pp. 275–330 and 358–383, and I resume the question for the purpose of applying it.

Formula for the general Coefficient $D(j, j')$.

In the First Part, the reciprocal of the distance of the two planets, or function

$$\{r^2 + r'^2 - 2rr'(\cos U \cos U' + \sin U \sin U' \cos \Phi)\}^{-\frac{1}{2}}$$

[p. 512]

is taken to be developed in multiple cosines of U, U' , the general term being

$$D(j, j') \cos(jU + j'U'),$$

where j, j' have each of them any integer value from $-\infty$ to $+\infty$ (zero not excluded), but so that j, j' are simultaneously even or simultaneously odd. We have $D(-j, -j') = D(j, j')$ and $D(j', j) = D(j, j')$; and it hence appears that the really distinct values of the coefficient may be taken to be those for which j is not negative, and as regards absolute magnitude is not less than j' ; and for such values of j, j' we have the above-mentioned expression

$$D(j, j') = \Sigma \frac{\Pi_{\frac{1}{2}}(x - \frac{1}{2})}{\Pi x} \eta^{2x} \Sigma M_x^\xi R_x^\xi,$$

which I proceed to explain and develope.

$\Pi_{\frac{1}{2}}(x - \frac{1}{2})$ and Πx (x being a positive integer) denote respectively $\frac{1}{2} \cdot \frac{3}{2} \dots (x - \frac{1}{2})$, and $1 \cdot 2 \cdot 3 \dots x$; in particular for $x = 0$, the value of each factorial is = 1.

η denotes $\sin \frac{1}{2}\Phi$.

The coefficients R_x^ξ are those of the multiple cosines in certain developments, viz. we have

$$r^s r'^s (r^2 + r'^2 - 2rr' \cos(U - U'))^{-s-\frac{1}{2}} = \Sigma R_x^\xi \cos i(U - U'),$$

where, as usual, i extends from $-\infty$ to ∞ and $R_x^{-i} = R_x^i$. Writing with Leverrier

$$\begin{aligned}(a^2 + a'^2 - 2aa' \cos H)^{-\frac{1}{2}} &= \frac{1}{2} \Sigma A^i \cos iH, \\ aa'(a^2 + a'^2 - 2aa' \cos H)^{-\frac{3}{2}} &= \frac{1}{2} \Sigma B^i \cos iH, \\ a^2 a'^2 (a^2 + a'^2 - 2aa' \cos H)^{-\frac{5}{2}} &= \frac{1}{2} \Sigma C^i \cos iH, \\ a^3 a'^3 (a^2 + a'^2 - 2aa' \cos H)^{-\frac{7}{2}} &= \frac{1}{2} \Sigma D^i \cos iH,\end{aligned}$$

then $2R_0^i, 2R_1^i, 2R_2^i, 2R_3^i$ are the same functions of r, r' that A^i, B^i, C^i, D^i respectively are of a, a' .

The expression of M_x^ξ is

$$M_x^\xi = (-)^{\frac{1}{2}(j+j'-\xi)} \frac{\Pi(x)}{\Pi_{\frac{1}{2}}(x-j-\xi) \Pi_{\frac{1}{2}}(x+j'+\xi)} \cdot \frac{\Pi(x)}{\Pi_{\frac{1}{2}}(x-j+\xi) \Pi_{\frac{1}{2}}(x+j'-\xi)},$$

and, finally, in the expression for $D(j, j')$, x has every integer value from 0 to ∞ and, for any given value of x , ξ extends by steps of two units from the inferior value $-(x-j')$ to the superior value $x-j$.

It is convenient to write $x = \frac{1}{2}(j+j') + s$; we have then ξ extending from $-\frac{1}{2}(j-j') - s$ to $-\frac{1}{2}(j-j') + s$, or writing $\xi = -\frac{1}{2}(j-j') + \vartheta$, ϑ has the $s+1$ values $s, s-2, s-4, \dots, -s$, viz. for $s = 2p+1$ the values are $\pm 1, \pm 3, \dots, \pm (2p+1)$, and for $s = 2p$ they are $0, \pm 2, \pm 4, \dots, \pm 2p$.

[p. 513]

Making these changes we have

$$D(j, j') = \Sigma \frac{\Pi_{\frac{1}{2}}\{\frac{1}{2}(j+j') + s - \frac{1}{2}\}}{\Pi\{\frac{1}{2}(j+j') + s\}} \eta^{j+j'+2s} \Sigma M_{\frac{1}{2}(j+j')+s}^{-\frac{1}{2}(j-j')+\vartheta} R_{\frac{1}{2}(j+j')+s}^{-\frac{1}{2}(j-j')+\vartheta},$$

where

$$M_{\frac{1}{2}(j+j')+s}^{-\frac{1}{2}(j-j')+\vartheta} = (-)^\vartheta \frac{\Pi\{\frac{1}{2}(j+j') + s\}}{\Pi_{\frac{1}{2}}(s-\vartheta) \Pi_{\frac{1}{2}}(j+j'+s+\vartheta)} \cdot \frac{\Pi\{\frac{1}{2}(j+j') + s\}}{\Pi_{\frac{1}{2}}(s+\vartheta) \Pi_{\frac{1}{2}}(j+j'+s-\vartheta)},$$

viz. this is $(-)^{\vartheta}$ into the product of two binomial coefficients, each belonging to the exponent $\frac{1}{2}(j+j') + s$.

Particular Cases, $j + j' = 0, 2, 4, 6$, being those required in the Planetary Theory.

Considering successively the cases $j + j' = 0, 2, 4, 6$, we have, first,

$$D(j, -j) = \Sigma \frac{\Pi_{\frac{1}{2}}(s - \frac{1}{2})}{\Pi s} \eta^{2s} \Sigma (-)^\vartheta \left\{ \frac{\Pi s}{\Pi_{\frac{1}{2}}(s-\vartheta) \Pi_{\frac{1}{2}}(s+\vartheta)} \right\}^2 R_s^{-j+\vartheta},$$

which, developed as far as η^6 , is

$$\begin{aligned} (*) \quad D(j, -j) &= \frac{1}{2} A^{-j} \\ &\quad - \frac{1}{2} \eta^2 (B^{-j-1} + B^{-j+1}) \\ &\quad + \frac{1 \cdot 3}{2 \cdot 4 \cdot 6} \eta^4 (\dots) \\ &\quad + \dots,\end{aligned}$$

where, and in what immediately follows, A, B, C, D are used to denote functions (not of (a, a') , but) of r, r' .

Secondly,

$$D(j, -j+2) = \Sigma \frac{\Pi_{\frac{1}{2}}(s - \frac{1}{2})}{\Pi(s+1)} \eta^{2s+2} \Sigma (-)^\vartheta \frac{\Pi(s+1)}{\Pi_{\frac{1}{2}}(s-\vartheta) \Pi_{\frac{1}{2}}(s+\vartheta)+2} \cdot \frac{\Pi(s+1)}{\Pi_{\frac{1}{2}}(s+\vartheta) \Pi_{\frac{1}{2}}(s-\vartheta)+2} R_{s+1}^{-j+1+\vartheta},$$

which, developed to η^6 , is

$$\begin{aligned} (*) \quad D(j, -j+2) &= \eta^2 \left\{ \frac{1}{2} B^{-j+1} \right. \\ &\quad \left. - \frac{1 \cdot 3}{2 \cdot 4 \cdot 6} \eta^4 (\dots) \right\}.\end{aligned}$$

[p. 514]

Thirdly,

$$D(j, -j+4) = \Sigma \frac{\Pi_{\frac{1}{2}}(s + \frac{3}{2})}{\Pi(s+2)} \eta^{2s+4} \Sigma(-)^{\vartheta} \left\{ \frac{\Pi(s+2)}{\Pi_{\frac{1}{2}}(s-\vartheta) \Pi_{\frac{1}{2}}(s+\vartheta) + 2} \cdot \frac{\Pi(s+2)}{\Pi_{\frac{1}{2}}(s+\vartheta) \Pi_{\frac{1}{2}}(s-\vartheta) + 2} \right\} R_{s+2}^{-j+2+\vartheta},$$

which, developed to η^6 , is

$$(*) \quad D(j, -j+4) = \eta^4 \left\{ \frac{1 \cdot 3}{2 \cdot 4} C^{-j+2} - \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \eta^2 \frac{1}{4} (3D^{-j+1} + 3D^{-j+3}) \right\};$$

and, fourthly,

$$D(j, -j+6) = \Sigma \frac{\Pi_{\frac{1}{2}}(s + \frac{5}{2})}{\Pi(s+3)} \eta^{2s+6} \Sigma(-)^{\vartheta} \left\{ \frac{\Pi(s+3)}{\Pi_{\frac{1}{2}}(s-\vartheta) \Pi_{\frac{1}{2}}(s+\vartheta) + 3} \cdot \frac{\Pi(s+3)}{\Pi_{\frac{1}{2}}(s+\vartheta) \Pi_{\frac{1}{2}}(s-\vartheta) + 3} \right\} R_{s+3}^{-j+3+\vartheta},$$

which, developed to η^6 , is simply

$$(*) \quad D(j, -j+6) = \eta^6 \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} D^{-j+3}.$$

The foregoing formulae, although obtained on the supposition $j = 0$, or positive, apply without alteration to the case $j = \text{negative}$, and the entire series of terms of an order not exceeding 6 as regards η may be written,

$$\begin{aligned} & D(j, -j) \cos(jU - jU') \\ & + 2D(j, -j+2) \cos(jU + (-j+2)U') \\ & + 2D(j, -j+4) \cos(jU + (-j+4)U') \\ & + 2D(j, -j+6) \cos(jU + (-j+6)U'), \end{aligned}$$

where j has every integer value from $-\infty$ to $+\infty$.

Comparison with Leverrier.

This is in fact what Leverrier's expression becomes on putting therein $e = e' = 0$. To verify this, observe that Leverrier having defined his A^i , B^i , C^i , D^i , as above, writes further

$$\begin{aligned} E^i &= \frac{1}{2}(B^{i-1} + B^{i+1}), \\ \mathfrak{E}^i &= \frac{1}{4}(C^{i-2} + 4C^i + C^{i+2}), \\ \mathfrak{F}^i &= \frac{1}{8}(D^{i-3} + 9D^{i-1} + 9D^{i+1} + D^{i+3}), \\ F^i &= \frac{1}{2}(C^{i-1} + C^{i+1}), \\ \mathfrak{G}^i &= \frac{1}{8}(D^{i-2} + 2D^i + D^{i+2}). \end{aligned}$$