Carl Friedrich Gauss

# Arithmetical Papers

English translation

Volume II: arithmetical papers, reviews and manuscript studies

AI-translated into English using OpenAI GPT-6.1 Sol.

This selection contains the historical title matter and first five complete papers of Volume II: A New Proof of an Arithmetical Theorem (Articles 1–7); Summation of Certain Special Series (Articles 1–36); New Proofs and Extensions of the Fundamental Theorem in the Theory of Quadratic Residues, with its introduction, fifth proof (Articles 1–3), sixth proof (Articles 1–6) and new algorithm (Articles 1–6); and both papers on the Theory of Biquadratic Residues (Articles 1–23 and 24–76). Five self-reviews from 1808–1831 and seven reviews of other writings from 1809–1831 follow. The latter concern irrational radicals, the factor tables of Chernac and Burckhardt, Erchinger's construction of the regular seventeen-sided polygon, and Seeber's positive ternary quadratic forms.

The manuscript studies include Solution of a Pure Congruence, Analysis of Residues, Chapter VI, first part (Articles 237–252); General Investigations of Congruences, Chapter VIII (Articles 330–375 as printed, including the unfinished ending of Article 375), with R. Dedekind's historical remarks; and Further Development of the Investigations of Pure Equations, complete as printed (Articles 1–19), with Dedekind's remarks. The fragment Proof of Some Theorems Concerning the Periods of Classes of Binary Quadratic Forms follows, with both theorems, the unfinished ending of the second proof, and Stern's historical remark.

The texts marked [I.] and [II.] of On the Connection between the Number of Classes into Which Binary Quadratic Forms Are Distributed and Their Determinant, First Paper, follow in full as printed, each with its title and Articles 1–8. The text marked [II.] was presented in 1837. Fragments III–X follow, with their historical editorial notes. Historical remarks on [I.]–[X.] follow. Geometrical Aspect of Ternary Forms follows with its two diagrams, crystallographic tables and Schering's historical remarks.

Originally German title matter, reviews, the manuscript division title, Fragments III–IV and VIII–X, the geometric notes and the historical remarks are rendered in modern German in the German reader. Their historical originals are supplied separately. The posthumous On the Theory of Biquadratic Residues, text [I.], is included in full as printed, Articles 1–19, with all five diagrams. Text [II.], Preparations for the General Theory of Biquadratic Residues, follows in full as printed, with both preparatory sections and Articles 1–12. Text [III.], Theory of Biquadratic Residues, follows in full as printed, with its title and Articles 1–14, including the joint heading 7–8. Text [IV.] follows in full as printed, Articles 1–5; its second method ends after the fourth step. Text [V.] follows in full as printed, Articles 1–17, with its editable lattice diagram. Text [VI.] follows in full as printed, Articles 1–6. Schering's complete historical remarks on texts [I.]–[VI.] follow. The originally German prose of these texts is rendered in modern German in the German reader; the historical originals are supplied separately. On the Theory of Complex Numbers follows in full as printed: six fragments [I.]–[VI.] on the decomposition of cubes, the nearest integer, measures of numbers, units and congruences, followed by Schering's historical remarks. The originally German passages are rendered in modern German in the German reader, and their historical originals are supplied separately. The complete historical table of the quadratic character of the primes from 2 to 997 as residues with respect to the primes from 3 to 503 as divisors follows. The complete historical table for converting common fractions with denominators from the first thousand into decimal fractions follows. The complete historical table of the frequency of prime numbers follows, including its matrices and closing observations. The complete letter from Gauss to Enke of 24 December 1849 follows, including its tables and discussion of prime-number frequency. The German reader renders the originally German letter in modern German; its historical original is supplied separately. The complete historical table of the number of classes of binary quadratic forms follows, with its negative and positive determinants and all printed summaries. Its historical values and abbreviations are retained. The German title is rendered in modern German in the German reader; the historical source is supplied separately.

The complete historical Table for Circle Division follows, with all nine factor tables and their printed index lists. Its originally German title and headings are rendered in modern German in the German reader; the historical source is supplied separately. A New Quadrature of the Circle follows with its four complete printed calculations and the arc table, including its historical blanks. On Calculating Common Logarithms follows in full as printed, with its tables and bracketed historical editorial account. Its German original is supplied separately; the German reader gives a modern German rendering. The First Ten Thousand Squares, Indices of Primes in the Higher Number Domain, and the Auxiliary Table for Solving the Indeterminate Equation by the Method of Exclusion follow in full as printed. The original German texts are supplied separately; the German reader gives modern German renderings. The Eighth Section, A Fuller Consideration of Certain Investigations Concerning Circle Division, follows through its whole printed ending in Article 373, including the interruption in Article 371. Both letters from Gauss to Dirichlet, dated 13 September 1826 and 30 May 1828, follow in full with their numerical tables. Their German originals are supplied separately; the German reader gives modern German renderings. Other volumes of Gauss's works lie outside this reader. Complete Volume I accompanies it in the collected reader.

Historical source: Carl Friedrich Gauss, Werke, Volume II, second printing, Göttingen, 1876.

AI-translated into English using OpenAI GPT-6.1 Sol.

Schering's complete historical closing remarks, the historical contents and the Göttingen imprint follow. Originally German passages are rendered in modern German in the German reader; their historical originals are supplied separately.

# Translation sources and terminology

Historical text: Carl Friedrich Gauss, *Werke*, Volume II, second printing, Göttingen, 1876. Historical editorial contributions retain their own authorship and signatures. Originally German passages are rendered in modern German in the German reader; their historical originals are supplied separately.
These English and German editions are AI-translated using OpenAI GPT-6.1 Sol, with AI-produced typesetting. The following named human translations and writings are scholarly references for the terminology and usage specified below. Gauss's original-language text remains the mathematical source.

**H. Maser.**

German translator and editor, *Carl Friedrich Gauss' Untersuchungen über höhere Arithmetik*, Berlin: Julius Springer, 1889. Corresponding arithmetical passages and German mathematical terminology.
`https://archive.org/details/carlfriedrichga00gausgoog`

**James Caddall Morehead and Adam Miller Hiltebeitel.**

Translators of Gauss's *General Investigations of Curved Surfaces of 1827 and 1825*, Princeton University Library, 1902; Project Gutenberg 36856, TeX edition. Related English geometrical terminology and usage.
`https://www.gutenberg.org/ebooks/36856`

**Paul Taylor; Bernard Leak.**

Taylor's English translation of Gauss's second proof of the fundamental theorem of algebra, with corrections by Leak, 1983. Related English algebraic terminology.
The consulted online edition identifies the translation as December 1983, its LaTeX version as August 2003, and the underlying source as last modified on 8 June 2007. Its acknowledgements credit Mark Wainwright for returning Taylor's notes and Nikita Danilov for typo corrections.
`https://www.paultaylor.eu/misc/gauss.html`

**Ulrich Görtz.**

*Algebra*, WS 2021/22, 3.1.1–3.1.3 and opening of 3.1.4; *Lineare Algebra II*, SS 2021, 18.4. Contemporary German arithmetic and algebraic usage.
`https://math.ug/a-ws2122/sect0006.html`
`https://math.ug/la2-ss21/sec-quotient-ring.html`

**Nikolay Moshchevitin.**

*Eine Bemerkung über positiv definite quadratische Formen und rationale Punkte*, arXiv:1510.06214v3. Related contemporary German terminology for quadratic forms.
`https://arxiv.org/abs/1510.06214v3`

**Ian O. Angell and Moreton Moore.**

*Projections of cubic crystals*, IUCr, Table 2. English crystallographic nomenclature.
IUCr: Projections of cubic crystals, Table 2

**Th. Hahn, H. Klapper, U. Müller and M. I. Aroyo.**

*Point groups and crystal classes*, *International Tables for Crystallography A*, 2016, chapter 3.2. Crystal-form names.
https://doi.org/10.1107/97809553602060000930


# CARL FRIEDRICH GAUSS COLLECTED WORKS

VOLUME II.

CARL FRIEDRICH GAUSS

# COLLECTED WORKS

SECOND VOLUME.



SECOND PRINTING

PUBLISHED
BY THE
ROYAL SOCIETY OF SCIENCES
AT
GÖTTINGEN
1876.

# A NEW PROOF

## OF AN ARITHMETICAL THEOREM

BY

CARL FRIEDRICH GAUSS

PRESENTED TO THE ROYAL SOCIETY OF SCIENCES JANUARY 15, 1808.

---

Commentationes societatis regiae scientiarum Gottingensis. Vol. XVI.
Gottingae MDCCCVIII.

---

# A NEW PROOF

## OF AN ARITHMETICAL THEOREM.

---

**1.**

Questions in higher arithmetic very often present a singular phenomenon, which occurs much more rarely in analysis and contributes greatly to their charm. In analytic investigations we usually cannot reach new truths unless we have first thoroughly mastered the principles on which they rest and which must, as it were, open the way to them. In arithmetic, by contrast, most elegant new truths very frequently spring forth through induction and a kind of unexpected good fortune; their proofs lie so deeply hidden and are wrapped in such darkness that they elude every attempt and deny access to the keenest investigations. Moreover, so great and so wonderful is the connection among arithmetical truths that at first sight seem utterly different, that quite often, while we are seeking something quite different, we at last most happily reach the proof so greatly desired and previously sought in vain through long reflection, by a path far different from the one expected. Usually, however, truths of this kind are of such a nature that they can be approached by several very different paths, and the paths that first present themselves are not always the shortest. It must therefore certainly be highly valued if, after such a truth has long been investigated in vain, and has then indeed been proved, but by rather obscure detours, we finally succeed in discovering the simplest and genuine path.

**2.**

Among the questions discussed in the preceding article, a distinguished place is held by the theorem containing almost the whole theory of quadratic residues, which in the *Arithmetical Investigations* (Section IV) is distinguished by the name of the *fundamental theorem*. The illustrious LEGENDRE must undoubtedly be regarded as the *first* discoverer of this most elegant theorem, after the great geometers EULER and LAGRANGE had, long before, already discovered several of its special cases by induction. I do not pause here to enumerate these men's attempts at a proof; those who wish may consult the work just mentioned. Let me add only, in confirmation of what was said in the preceding article, what concerns my own attempts. I came upon the theorem itself independently in the year 1795, when I was entirely ignorant of everything that had already been worked out in higher arithmetic and wholly cut off from access to the literature. But for an entire year it tormented me and eluded my most strenuous efforts, until at last I obtained the proof given in the fourth section of that work. Afterwards three others, resting on wholly

different principles, presented themselves to me; I gave one in the fifth section, and the others, not inferior to it in elegance, I shall make public on another occasion. But all these proofs, although they seem to leave nothing to be desired in rigour, are derived from principles that are too different in kind, perhaps excepting the first, which nevertheless proceeds by more laborious reasoning and is burdened by rather lengthy operations. I therefore do not hesitate to declare that a *genuine* proof has so far been lacking. Let the experts now judge whether the one that I recently succeeded in discovering, and that the following pages present, deserves to be honoured with this name.

**3.**

THEOREM. *Let $p$ be a positive prime number; $k$ any integer not divisible by $p$; use the following definitions:*

$A$ *collection of the numbers* $1,\ 2,\ 3\ \ldots.\ \frac{1}{2}(p-1)$

$B$ *collection of these* $\frac{1}{2}(p+1),\ \frac{1}{2}(p+3),\ \frac{1}{2}(p+5)\ \ldots.\ p-1$

*Take the products of $k$ by the individual numbers in $A$, and their least positive residues modulo $p$, which are evidently all distinct and belong partly to $A$, partly to $B$. Now if altogether $B$ contains $\mu$ of the residues, $k$ will be either a quadratic residue or a quadratic nonresidue of $p$, according as $\mu$ is even or odd.*

*Proof.* Let the residues belonging to $A$ be $a,\ a'\ a''\ \ldots.$, and the remaining ones, belonging to $B$, be $b, b', b'' \ldots.$. It is clear that the complements of the latter, $p-b, p-b', p-b'' \ldots.$, are all distinct from the numbers $a,\ a',\ a''\ \ldots.$, but together with these fill out the collection $A$. Thus we have

$$1\,.\,2\,.\,3\ \ldots.\ \tfrac{1}{2}(p-1) = aa'a''\ \ldots.\ (p-b)(p-b')(p-b'')\ \ldots.$$

But the latter product evidently becomes

$$\equiv (-1)^{\mu} aa'a''\ \ldots.\ bb'b''\ \ldots.\ \equiv (-1)^{\mu} k\,.\,2k\,.\,3k\ \ldots.\ \tfrac{1}{2}(p-1)k$$
$$\equiv (-1)^{\mu} k^{\frac{1}{2}(p-1)}\,1\,.\,2\,.\,3\ \ldots.\ \tfrac{1}{2}(p-1) (\mathrm{mod}.p)$$

Hence we have

$$1 \equiv (-1)^{\mu} k^{\frac{1}{2}(p-1)}$$

or $k^{\frac{1}{2}(p-1)} \equiv \pm 1$, according as $\mu$ is even or odd; our theorem follows immediately.

**4.**

We can greatly shorten the following reasoning by introducing some suitable notation. Thus the symbol $(k, p)$ will denote for us the number of products among

$$k,\ 2k,\ 3k\ \ldots.\ \tfrac{1}{2}(p-1)k,$$

whose least positive residues modulo $p$ exceed half of it. Further, when $x$ is any noninteger quantity, we shall use the symbol $[x]$ to denote the integer immediately smaller than $x$ itself, so that $x - [x]$ is always a positive quantity lying between the limits 0 and 1. The following relations are now obtained with little trouble:

I. $[x] + [-x] = -1$.

II. $[x] + h = [x + h]$, whenever $h$ is an integer.

III. $[x] + [h - x] = h - 1$.

IV. If $x - [x]$ is a fraction smaller than $\frac{1}{2}$, then $[2x] - 2[x] = 0$; but if $x - [x]$ is greater than $\frac{1}{2}$, then $[2x] - 2[x] = 1$.

V. Thus, when the least positive residue of the integer $h$ modulo $p$ lies below $\frac{1}{2}p$, we have $\left[\frac{2h}{p}\right] - 2\left[\frac{h}{p}\right] = 0$; but when that residue lies above $\frac{1}{2}p$, we have $\left[\frac{2h}{p}\right] - 2\left[\frac{h}{p}\right] = 1$.

VI. It follows at once that $(k, p) =$

$$\left[\frac{2k}{p}\right] + \left[\frac{4k}{p}\right] + \left[\frac{6k}{p}\right] \ldots. + \left[\frac{(p-1)k}{p}\right]$$
$$- 2\left[\frac{k}{p}\right] - 2\left[\frac{2k}{p}\right] - 2\left[\frac{3k}{p}\right] \ldots. - 2\left[\frac{\frac{1}{2}(p-1)k}{p}\right].$$

VII. From VI and I we obtain without difficulty

$$(k, p) + (-k, p) = \tfrac{1}{2}(p-1)$$

It follows that $-k$ has either the same or the opposite relation to $p$ (insofar as it is a quadratic residue or nonresidue of it) as $+k$, according as $p$ is of the form $4n + 1$ or of the form $4n + 3$. In the former case $-1$ is evidently a residue, in the latter a nonresidue, of $p$.

VIII. We shall transform the formula given in VI as follows. By III we have

$$\left[\frac{(p-1)k}{p}\right] = k - 1 - \left[\frac{k}{p}\right],$$
$$\left[\frac{(p-3)k}{p}\right] = k - 1 - \left[\frac{3k}{p}\right],$$
$$\left[\frac{(p-5)k}{p}\right] = k - 1 - \left[\frac{5k}{p}\right] \ldots.$$

Applying these substitutions to the last $\frac{p\mp 1}{4}$ terms of the upper series in that expression, we have

*first*, whenever $p$ is of the form $4n+1$

$$(k,p) = \tfrac{1}{4}(k-1)(p-1)$$
$$- 2\left\{\left[\frac{k}{p}\right] + \left[\frac{3k}{p}\right] + \left[\frac{5k}{p}\right] \ldots. + \left[\frac{\frac{1}{2}(p-3)k}{p}\right]\right\}$$
$$- \left\{\left[\frac{k}{p}\right] + \left[\frac{2k}{p}\right] + \left[\frac{3k}{p}\right] \ldots. + \left[\frac{\frac{1}{2}(p-1)k}{p}\right]\right\}$$

*second*, whenever $p$ is of the form $4n+3$

$$(k,p) = \tfrac{1}{4}(k-1)(p+1)$$
$$- 2\left\{\left[\frac{k}{p}\right] + \left[\frac{3k}{p}\right] + \left[\frac{5k}{p}\right] \ldots. + \left[\frac{\frac{1}{2}(p-1)k}{p}\right]\right\}$$
$$- \left\{\left[\frac{k}{p}\right] + \left[\frac{2k}{p}\right] + \left[\frac{3k}{p}\right] \ldots. + \left[\frac{\frac{1}{2}(p-1)k}{p}\right]\right\}$$

IX. For the special case $k = +2$, the formulas just given yield $(2,p) = \frac{1}{4}(p\mp 1)$, taking the upper or lower sign according as $p$ is of the form $4n+1$ or $4n+3$. Thus $(2,p)$ is even, and hence $2Rp$, whenever $p$ is of the form $8n+1$ or $8n+7$; on the other hand, $(2,p)$ is odd and $2Np$, whenever $p$ is of the form $8n+3$ or $8n+5$.

**5.**

THEOREM. *Let $x$ be a positive noninteger quantity, among whose multiples $x$, $2x$, $3x \ldots$. up to $nx$ none is an integer; put $[nx] = h$, from which it is easily concluded that among the multiples of the reciprocal quantity $\frac{1}{x}$, $\frac{2}{x}$, $\frac{3}{x} \ldots\ldots$ up to $\frac{h}{x}$ an integer is likewise not found. Then I say that we have*

$$\left.\begin{array}{l} [x] + [2x] + [3x] \,.\, + [nx] \\ + \left[\frac{1}{x}\right] + \left[\frac{2}{x}\right] + \left[\frac{3}{x}\right] \ldots. + \left[\frac{h}{x}\right] \end{array}\right\} = nh.$$

*Proof.* In the series $[x] + [2x] + [3x] \ldots. + [nx]$, which we shall put $= \Omega$, the first terms up to and including the term in position $\left[\frac{1}{x}\right]$ are evidently all $= 0$; the following ones up to the term in position $\left[\frac{2}{x}\right]$ are all $= 1$; the following ones up to the term in position $\left[\frac{3}{x}\right]$ are all $= 2$, and so on. Hence we have

$$\left.\begin{array}{l} \Omega = 0 \times \left[\frac{1}{x}\right] \\ \quad + 1 \times \left\{\left[\frac{2}{x}\right] - \left[\frac{1}{x}\right]\right\} \\ \quad + 2 \times \left\{\left[\frac{3}{x}\right] - \left[\frac{2}{x}\right]\right\} \\ \quad + 3 \times \left\{\left[\frac{4}{x}\right] - \left[\frac{3}{x}\right]\right\} \\ \qquad \text{etc.} \\ \quad + (h-1) \left\{\left[\frac{h}{x}\right] - \left[\frac{h-1}{x}\right]\right\} \\ \quad + h \left\{n - \left[\frac{h}{x}\right]\right\} \end{array}\right\} = hn - \left[\frac{1}{x}\right] - \left[\frac{2}{x}\right] - \left[\frac{3}{x}\right] \ldots. - \left[\frac{h}{x}\right]$$

Q. E. D.

**6.**

THEOREM. *If $k, p$ denote any positive odd numbers relatively prime to each other, we have*

$$\left.\begin{aligned}&\left[\frac{k}{p}\right]+\left[\frac{2k}{p}\right]+\left[\frac{3k}{p}\right]\ldots.+\left[\frac{\frac{1}{2}(p-1)k}{p}\right]\\&+\left[\frac{p}{k}\right]+\left[\frac{2p}{k}\right]+\left[\frac{3p}{k}\right]\ldots.+\left[\frac{\frac{1}{2}(k-1)p}{k}\right]\end{aligned}\right\}=\tfrac{1}{4}(k-1)(p-1).$$

*Proof.* Supposing, as we may, that $k<p$, we have $\frac{\frac{1}{2}(p-1)k}{p}$ less than $\frac{1}{2}k$, but greater than $\frac{1}{2}(k-1)$, and therefore $\left[\frac{\frac{1}{2}(p-1)k}{p}\right]=\frac{1}{2}(k-1)$. Thus the present theorem evidently follows immediately from the preceding one by putting there $\frac{k}{p}=x$, $\frac{1}{2}(p-1)=n$, and hence $\frac{1}{2}(k-1)=h$.

Moreover, it can be proved in a similar manner that, if $k$ is an *even* number relatively prime to $p$, we have

$$\left.\begin{aligned}&\left[\frac{k}{p}\right]+\left[\frac{2k}{p}\right]+\left[\frac{3k}{p}\right]\ldots.+\left[\frac{\frac{1}{2}(p-1)k}{p}\right]\\&+\left[\frac{p}{k}\right]+\left[\frac{2p}{k}\right]+\left[\frac{3p}{k}\right]\ldots.+\left[\frac{\frac{1}{2}kp}{k}\right]\end{aligned}\right\}=\tfrac{1}{4}k(p-1)$$

But we do not dwell on this proposition, which is not needed for our purpose.

**7.**

The fundamental theorem now follows immediately by combining the preceding theorem with proposition VIII of Article 4. Namely, if $k, p$ denote any unequal positive prime numbers, and we put

$$(k,p) + \left[\frac{k}{p}\right] + \left[\frac{2k}{p}\right] + \left[\frac{3k}{p}\right] \ldots . + \left[\frac{\frac{1}{2}(p-1)k}{p}\right] = L$$

$$(p,k) + \left[\frac{p}{k}\right] + \left[\frac{2p}{k}\right] + \left[\frac{3p}{k}\right] \ldots . + \left[\frac{\frac{1}{2}(k-1)p}{k}\right] = M$$

it is clear from VIII of Article 4 that $L$ and $M$ are always even numbers. But by the theorem of Article 6 we have

$$L + M = (k,p) + (p,k) + \tfrac{1}{4}(k-1)(p-1)$$

Thus whenever $\frac{1}{4}(k-1)(p-1)$ is even, which happens if either both $k, p$ or at least one of them is of the form $4n+1$, $(k,p)$ and $(p,k)$ must necessarily be either both even or both odd. But whenever $\frac{1}{4}(k-1)(p-1)$ is odd, which happens if both $k, p$ are of the form $4n+3$, one of the numbers $(k,p)$, $(p,k)$ must necessarily be even and the other odd. Thus in the former case the relation of $k$ to $p$ and the relation of $p$ to $k$ (insofar as one is a residue or nonresidue of the other) are identical, and in the latter case opposite.

Q. E. D.

# SUMMATION

# OF CERTAIN

# SPECIAL SERIES

BY

CARL FRIEDRICH GAUSS

PRESENTED TO THE SOCIETY AUGUST XXIV, MDCCCVIII.

---

Commentationes societatis regiae scientiarum Gottingensis recentiores. Vol. I.
Gottingae MDCCCXI.

---

# SUMMATION

## OF CERTAIN SPECIAL SERIES.

---

**1.**

Among the more notable truths to which the theory of circle division has opened the way, the summation proposed in Disquiss. Arithmet., Article 356, holds a distinguished place. This is due both to its particular elegance and remarkable fruitfulness, which another investigation will later give us occasion to discuss more fully, and to the unusual difficulties that beset its rigorous and complete proof. These difficulties should have been all the less expected because they concern a certain restriction of the theorem more than the theorem itself: once this restriction is disregarded, a proof is immediately available and follows very easily from the theory developed in that work. The theorem was given there in the following form. Suppose that $n$ is a prime number. Represent all the quadratic residues of $n$ between 1 and $n-1$ inclusive by $a$, and all the nonresidues between the same limits by $b$; finally, let $\omega$ be the arc $\frac{360^0}{n}$, and $k$ any fixed integer not divisible by $n$. Then we have

I. for a value of $n$ of the form $4m+1$,

$$\begin{aligned}
\Sigma \cos ak\omega &= -\tfrac{1}{2} \pm \tfrac{1}{2}\sqrt{n} \\
\Sigma \cos bk\omega &= -\tfrac{1}{2} \mp \tfrac{1}{2}\sqrt{n}, \text{ and hence} \\
\Sigma \cos ak\omega - \Sigma \cos bk\omega &= \pm\sqrt{n} \\
\Sigma \sin ak\omega &= 0 \\
\Sigma \sin bk\omega &= 0
\end{aligned}$$

II. for a value of $n$ of the form $4m+3$,

$$\begin{aligned}
\Sigma \cos ak\omega &= -\tfrac{1}{2} \\
\Sigma \cos bk\omega &= -\tfrac{1}{2} \\
\Sigma \sin ak\omega &= \pm\tfrac{1}{2}\sqrt{n} \\
\Sigma \sin bk\omega &= \mp\tfrac{1}{2}\sqrt{n} \\
\Sigma \sin ak\omega - \Sigma \sin bk\omega &= \pm\sqrt{n}
\end{aligned}$$

These summations were proved with full rigor in the place cited, and the only difficulty that remains here is to determine the *sign* to be placed before the radical. It can indeed be shown without difficulty that this sign depends on the number $k$ to the following extent: the *same* sign must hold for all values of $k$ that are quadratic residues of $n$, and the opposite sign for all values of $k$

that are quadratic nonresidues of $n$. Thus the whole question turns on the value $k = 1$: as soon as the sign for this value is known, the signs for all the other values of $k$ are immediately available as well. Yet this very question, which at first seems to be among the easier ones, presents us with unforeseen difficulties, and the method that had guided us this far without impediment utterly refuses any further help.

**2.**

Before proceeding further, it will be useful to work out some examples of our summation numerically; we should, however, first make a few general observations.

I. In the case where $n$ is a prime number of the form $4m + 1$, let all the quadratic residues of $n$ between 1 and $\frac{1}{2}(n-1)$ inclusive be represented by $a'$, and all the nonresidues between the same limits by $b'$. All the $n - a'$ are known to be among the $a$, and all the $n - b'$ among the $b$. Since all the $a'$, $b'$, $n-a'$, $n-b'$ together evidently make up the entire collection of numbers 1, 2, 3 .... $n-1$, all the $a'$ together with all the $n - a'$ comprise all the $a$; likewise all the $b'$ together with all the $n - b'$ comprise all the $b$. Hence

$$\begin{aligned}
\Sigma \cos ak\omega &= \Sigma \cos a'k\omega + \Sigma \cos(n-a')k\omega \\
\Sigma \cos bk\omega &= \Sigma \cos b'k\omega + \Sigma \cos(n-b')k\omega \\
\Sigma \sin ak\omega &= \Sigma \sin a'k\omega + \Sigma \sin(n-a')k\omega \\
\Sigma \sin bk\omega &= \Sigma \sin b'k\omega + \Sigma \sin(n-b')k\omega
\end{aligned}$$

Now, since $\cos(n-a')k\omega = \cos a'k\omega$, $\cos(n-b')k\omega = \cos b'k\omega$, $\sin(n-a')k\omega = -\sin a'k\omega$, $\sin(n-b')k\omega = -\sin b'k\omega$, it follows at once that

$$\begin{aligned}
\Sigma \sin ak\omega &= \Sigma \sin a'k\omega - \Sigma \sin a'k\omega = 0 \\
\Sigma \sin bk\omega &= \Sigma \sin b'k\omega - \Sigma \sin b'k\omega = 0
\end{aligned}$$

The summation of the cosines takes the form

$$\begin{aligned}
\Sigma \cos ak\omega &= 2\Sigma \cos a'k\omega \\
\Sigma \cos bk\omega &= 2\Sigma \cos b'k\omega
\end{aligned}$$

so that we must have

$$\begin{aligned}
1 + 4\Sigma \cos a'k\omega &= \pm\sqrt{n} \\
1 + 4\Sigma \cos b'k\omega &= \mp\sqrt{n} \\
2\Sigma \cos a'k\omega - 2\Sigma \cos b'k\omega &= \pm\sqrt{n}
\end{aligned}$$

II. In the case where $n$ is of the form $4m + 3$, the complement of any residue $a$ to $n$ is a nonresidue, and the complement of any $b$ is a residue. Thus all the $n - a$ coincide with all the $b$, and all the $n - b$ with all the $a$. This gives

$$\Sigma \cos ak\omega = \Sigma \cos(n - b)k\omega = \Sigma \cos bk\omega$$

Since all the $a$ and $b$ together make up all the numbers 1, 2, $3 \ldots . n - 1$, and hence $\Sigma \cos ak\omega + \Sigma \cos bk\omega = \cos k\omega + \cos 2k\omega + \cos 3k\omega + \text{etc.} + \cos(n - 1)k\omega = -1$, the summations

$$\Sigma \cos ak\omega = -\tfrac{1}{2}$$
$$\Sigma \cos bk\omega = -\tfrac{1}{2}$$

follow immediately. Likewise,

$$\Sigma \sin ak\omega = \Sigma \sin(n - b)k\omega = -\Sigma \sin bk\omega$$

which shows how the two summations

$$2\Sigma \sin ak\omega = \pm\sqrt{n}$$
$$2\Sigma \sin bk\omega = \mp\sqrt{n}$$

depend on one another.

**3.**

Here are the numerical calculations for several examples:

I. For $n = 5$ there is one value of $a'$, namely $a' = 1$, and one value of $b'$, namely $b' = 2$. We have

$$\cos \omega = +0{,}3090169944 \qquad \cos 2\omega = -0{,}8090169944$$

and hence $1 + 4 \cos \omega = +\sqrt{5}$, $1 + 4 \cos 2\omega = -\sqrt{5}$.

II. For $n = 13$ there are three values of $a'$, namely 1, 3, 4, and as many values of $b'$, namely 2, 5, 6. We calculate

| | | | |
|---|---|---|---|
| $\cos \omega$ | $= +0{,}8854560257$ | $\cos 2\omega$ | $= +0{,}5680647467$ |
| $\cos 3\omega$ | $= +0{,}1205366803$ | $\cos 5\omega$ | $= -0{,}7485107482$ |
| $\cos 4\omega$ | $= -0{,}3546048870$ | $\cos 6\omega$ | $= -0{,}9709418174$ |
| Sum | $= +0{,}6513878190$ | Sum | $= -1{,}1513878189$ |

Hence $1 + 4\Sigma \cos a'\omega = +\sqrt{13}$, $1 + 4\Sigma \cos b'\omega = -\sqrt{13}$.

III. For $n = 17$ we have four values of $a'$, namely 1, 2, 4, 8, and as many values of $b'$, namely 3, 5, 6, 7. The cosines are therefore calculated as follows:

$$\begin{array}{rcl} \cos\omega &=& +0{,}9324722294 \\ \cos 2\omega &=& +0{,}7390089172 \\ \cos 4\omega &=& +0{,}0922683595 \\ \cos 8\omega &=& -0{,}9829730997 \\ \hline \text{Sum} &=& +0{,}7807764064 \end{array} \qquad \begin{array}{rcl} \cos 3\omega &=& +0{,}4457383558 \\ \cos 5\omega &=& -0{,}2736629901 \\ \cos 6\omega &=& -0{,}6026346364 \\ \cos 7\omega &=& -0{,}8502171357 \\ \hline \text{Sum} &=& -1{,}2807764065 \end{array}$$

Hence $1 + 4\Sigma \cos a'\omega = +\sqrt{17}, \quad 1 + 4\Sigma \cos b'\omega = -\sqrt{17}$.

IV. For $n = 3$ there is just one value of $a$, namely $a = 1$, to which corresponds

$$\sin\omega = +0{,}8660254038$$

Hence $2 \sin\omega = +\sqrt{3}$.

V. For $n = 7$ there are three values of $a$, namely 1, 2, 4. These give the sines

$$\begin{array}{rcl} \sin\omega &=& +0{,}7818314825 \\ \sin 2\omega &=& +0{,}9749279122 \\ \sin 4\omega &=& -0{,}4338837391 \\ \hline \text{Sum} &=& +1{,}3228756556, \end{array}$$ and hence $2\Sigma \sin a\omega = +\sqrt{7}$.

VI. For $n = 11$ the values of $a$ are 1, 3, 4, 5, 9, to which correspond the sines

$$\begin{array}{rcl} \sin\omega &=& +0{,}5406408175 \\ \sin 3\omega &=& +0{,}9898214419 \\ \sin 4\omega &=& +0{,}7557495744 \\ \sin 5\omega &=& +0{,}2817325568 \\ \sin 9\omega &=& -0{,}9096319954 \\ \hline \text{Sum} &=& +1{,}6583123952, \end{array}$$ and hence $2\Sigma \sin a\omega = +\sqrt{11}$.

VII. For $n = 19$ the values of $a$ are 1, 4, 5, 6, 7, 9, 11, 16, 17, to which correspond the sines

$$\begin{aligned}
\sin\omega &= +0{,}3246994692\\
\sin 4\omega &= +0{,}9694002659\\
\sin 5\omega &= +0{,}9965844930\\
\sin 6\omega &= +0{,}9157733267\\
\sin 7\omega &= +0{,}7357239107\\
\sin 9\omega &= +0{,}1645945903\\
\sin 11\omega &= -0{,}4759473930\\
\sin 16\omega &= -0{,}8371664783\\
\sin 17\omega &= -0{,}6142127127\\
\text{Sum} &= +2{,}1794494718,
\end{aligned}$$

and hence $2\Sigma \sin a\omega = +\sqrt{19}$.

**4.**

In all these examples the radical takes the positive sign. This is also easily confirmed for the larger values $n = 23$, $n = 29$, etc., and already makes it highly probable that the same holds in general. But a proof of this phenomenon cannot be drawn from the principles set out in the place cited, and it must with full justification be regarded as a matter for deeper investigation. The purpose of this paper is therefore to present a rigorous proof of this most elegant theorem, attempted in vain in various ways over several years and finally achieved through unusual and quite subtle considerations. At the same time we shall bring the theorem to a far greater degree of generality, preserving, or rather increasing, its elegance. Finally, as a crowning addition, we shall show the remarkable and very close connection between this summation and another arithmetical theorem of the greatest importance. We hope that these investigations will themselves be welcome to geometers, and that the methods by which we have accomplished all this, and which may also be useful on other occasions, will be considered worthy of their attention.

**5.**

Our proof comes from considering a special kind of series whose terms depend on expressions such as

$$\frac{(1-x^m)\,(1-x^{m-1})\,(1-x^{m-2})\ldots.(1-x^{m-\mu+1})}{(1-x)\,(1-xx)\,(1-x^3)\ldots..(1-x^\mu)}$$

For brevity we shall denote such a fraction by $(m, \mu)$, and first make a few general observations about functions of this kind.

I. Whenever $m$ is a positive integer less than $\mu$, the function $(m, \mu)$ evidently vanishes, since the numerator contains the factor $1 - x^0$. For $m = \mu$, the factors in the numerator are identical, in reverse order, to those in the denominator, so that $(\mu, \mu) = 1$. Finally, for the case where $m$ is a positive integer greater than $\mu$, we have the formulas

$$(\mu+1,\mu) = \frac{1-x^{\mu+1}}{1-x} = (\mu+1,1)$$
$$(\mu+2,\mu) = \frac{(1-x^{\mu+2})\,(1-x^{\mu+1})}{(1-x)\qquad(1-xx)} = (\mu+2,2)$$
$$(\mu+3,\mu) = \frac{(1-x^{\mu+3})(1-x^{\mu+2})(1-x^{\mu+1})}{(1-x)\qquad(1-xx)\qquad(1-x^3)} = (\mu+3,3) \text{ etc.}$$

or, in general,

$$(m,\mu) = (m, m-\mu)$$

II. Further, it is easily confirmed that in general

$$(m,\mu+1) = (m-1,\mu+1) + x^{m-\mu-1}(m-1,\mu)$$

and since we likewise have

$$(m-1,\mu+1) = (m-2,\mu+1) + x^{m-\mu-2}(m-2,\mu)$$
$$(m-2,\mu+1) = (m-3,\mu+1) + x^{m-\mu-3}(m-3,\mu)$$
$$(m-3,\mu+1) = (m-4,\mu+1) + x^{m-\mu-4}(m-4,\mu), \text{ etc.}$$

and can continue this series as far as

$$\begin{aligned}(\mu+2,\mu+1) &= (\mu+1,\mu+1) + x(\mu+1,\mu)\\ &= (\mu,\mu) + x(\mu+1,\mu)\end{aligned}$$

provided that $m$ is a positive integer greater than $\mu+1$, we obtain

$$\begin{aligned}(m,\mu+1) = (\mu,\mu) &+ x(\mu+1,\mu) + xx(\mu+2,\mu)\\ &+ x^3(\mu+3,\mu) + \text{etc.}\\ &+ x^{m-\mu-1}(m-1,\mu)\end{aligned}$$

This shows that if, for some fixed value of $\mu$, every function $(m, \mu)$ is a polynomial whenever $m$ is a positive integer, then every function $(m, \mu + 1)$ must also be a polynomial. Since that supposition holds for $\mu = 1$, it therefore also holds for $\mu = 2$, and hence for $\mu = 3$, etc. That is, in general, for any positive integral value of $m$, $(m, \mu)$ is a polynomial, or equivalently the product

$$(1 - x^m)(1 - x^{m-1})(1 - x^{m-2}) \dots (1 - x^{m-\mu+1})$$

is divisible by

$$(1 - x)(1 - x^2)(1 - x^3) \dots (1 - x^\mu)$$

**6.**

We shall now consider two series, both of which can lead to our goal. The first series is

$$1 - \frac{1-x^m}{1-x} + \frac{(1-x^m)\,(1-x^{m-1})}{(1-x)\,(1-xx)} - \frac{(1-x^m)(1-x^{m-1})(1-x^{m-2})}{(1-x)\quad(1-xx)\quad(1-x^3)} + \text{etc.}$$

or

$$1 - (m,1) + (m,2) - (m,3) + (m,4) - \text{etc.}$$

which for brevity we shall denote by $f(x,m)$. First, it is immediately evident that whenever $m$ is a positive integer this series *terminates* after its term in position $m+1$ (with $= \pm 1$), and that in this case its sum must therefore be a finite polynomial in $x$. Further, Article 5, II, shows that in general, for any value of $m$,

$$\begin{aligned} 1 &= 1 \\ -(m,1) &= -(m-1,1) - x^{m-1} \\ +(m,2) &= +(m-1,2) + x^{m-2}(m-1,1) \\ -(m,3) &= -(m-1,3) - x^{m-3}(m-1,2), \text{ etc.} \end{aligned}$$

and hence

$$\begin{aligned} f(x,m) = 1 - x^{m-1} &- (1-x^{m-2})(m-1,1) \\ &+ (1-x^{m-3})(m-1,2) \\ &- (1-x^{m-4})(m-1,3) + \text{etc.} \end{aligned}$$

But evidently

$$\begin{aligned} (1-x^{m-2})(m-1,1) &= (1-x^{m-1})(m-2,1) \\ (1-x^{m-3})(m-1,2) &= (1-x^{m-1})(m-2,2) \\ (1-x^{m-4})(m-1,3) &= (1-x^{m-1})(m-2,3), \text{ etc.} \end{aligned}$$

from which we deduce the equation

$$f(x,m) = (1-x^{m-1})f(x,m-2) \ldots\ldots\ldots [1]$$

**7.**

Since for $m = 0$ we have $f(x, m) = 1$, the formula just found gives

$$f(x,2) = 1 - x$$
$$f(x,4) = (1-x)(1-x^3)$$
$$f(x,6) = (1-x)(1-x^3)(1-x^5)$$
$$f(x,8) = (1-x)(1-x^3)(1-x^5)(1-x^7), \text{ etc.}$$

or, in general, for any even value of $m$,

$$f(x,m) = (1-x)(1-x^3)(1-x^5)\ldots.(1-x^{m-1}) \ldots [2].$$

On the other hand, since for $m = 1$ we have $f(x, m) = 0$, we also have

$$f(x,3) = 0$$
$$f(x,5) = 0$$
$$f(x,7) = 0 \text{ etc.}$$

or, in general, for any odd value of $m$,

$$f(x,m) = 0$$

The latter summation could, moreover, already have been derived from the fact that in the series

$$1 - (m,1) + (m,2) - (m,3) + \text{etc.} + (m,m-1) - (m,m)$$

the last term cancels the first, the next to last cancels the second, etc.

**8.**

For our purpose the case where $m$ is a positive odd integer is sufficient. But the unusual nature of the matter makes it worthwhile to add a few remarks about the cases where $m$ is fractional or negative. Our series then evidently no longer terminates, but continues to infinity; it is also easily seen that it becomes divergent whenever $x$ is given a value less than 1, so that its summation must be restricted to values of $x$ greater than 1.

By formula [1] of Article 6 we have

$$f(x,-2) = \frac{1}{1-\frac{1}{x}}$$

$$f(x,-4) = \frac{1}{1-\frac{1}{x}} \cdot \frac{1}{1-\frac{1}{x^3}}$$

$$f(x,-6) = \frac{1}{1-\frac{1}{x}} \cdot \frac{1}{1-\frac{1}{x^3}} \cdot \frac{1}{1-\frac{1}{x^5}}, \text{ etc.}$$

so that the value of the function $f(x,m)$ can also be given in finite terms for a negative even integral value of $m$. For the remaining values of $m$, however, we shall convert the function $f(x,m)$ into an *infinite product* as follows.

As $m$ tends to a negative *infinite* value, the function $f(x,m)$ becomes

$$1 + \frac{1}{x-1} + \frac{1}{x-1} \cdot \frac{1}{xx-1} + \frac{1}{x-1} \cdot \frac{1}{xx-1} \cdot \frac{1}{x^3-1} + \text{etc.}$$

This series is therefore equal to the infinite product

$$\frac{1}{1-\frac{1}{x}} \cdot \frac{1}{1-\frac{1}{x^3}} \cdot \frac{1}{1-\frac{1}{x^5}} \cdot \frac{1}{1-\frac{1}{x^7}} \text{ etc. to infinity.}$$

Further, since in general

$$f(x,m) = f(x,m-2\lambda) \,.\, (1-x^{m-1})(1-x^{m-3}) \\ (1-x^{m-5}) \,..\, (1-x^{m-2\lambda+1})$$

we have

$$f(x,m) = f(x,-\infty) \,.\, (1-x^{m-1})(1-x^{m-3})(1-x^{m-5}) \\ \text{etc. to infinity.} \\ = \frac{1-x^{m-1}}{1-x^{-1}} \cdot \frac{1-x^{m-3}}{1-x^{-3}} \cdot \frac{1-x^{m-5}}{1-x^{-5}} \cdot \frac{1-x^{m-7}}{1-x^{-7}} \\ \text{etc. to infinity.}$$

whose factors evidently converge ever more closely to unity as they continue.

The case $m = -1$ deserves particular attention. Here

$$f(x,-1) = 1 + x^{-1} + x^{-3} + x^{-6} + x^{-10} + \text{etc.}$$

This series is therefore equal to the infinite product

$$\frac{1-x^{-2}}{1-x^{-1}} \cdot \frac{1-x^{-4}}{1-x^{-3}} \cdot \frac{1-x^{-6}}{1-x^{-5}} \text{ etc.}$$

or, writing $x$ for $x^{-1}$, we obtain

$$1+x+x^3+x^6+\text{etc.} = \frac{1-xx}{1-x} \cdot \frac{1-x^4}{1-x^3} \cdot \frac{1-x^6}{1-x^5} \cdot \frac{1-x^8}{1-x^7} \text{ etc.}$$

This equality between two rather intricate expressions, to which we shall return on another occasion, is certainly very remarkable.

**9.**

Secondly, we shall consider the series

$$1 + x^{\frac{1}{2}}\frac{1-x^m}{1-x} + x\frac{(1-x^m)(1-x^{m-1})}{(1-x)(1-xx)} + x^{\frac{3}{2}}\frac{(1-x^m)(1-x^{m-1})(1-x^{m-2})}{(1-x)(1-xx)(1-x^3)} + \text{etc.}$$

or

$$1 + x^{\frac{1}{2}}(m,1) + x(m,2) + x^{\frac{3}{2}}(m,3) + xx(m,4) + \text{etc.}$$

which we shall denote by $F(x,m)$. We restrict this investigation to the case where $m$ is a positive integer, so that this series too always ends at its term in position $m+1$ (with $= x^{\frac{1}{2}m}(m,m)$). Since

$$(m,m) = 1, \quad (m,m-1) = (m,1), \quad (m,m-2) = (m,2), \text{ etc.}$$

the series can also be written as follows:

$$F(x,m) = x^{\frac{1}{2}m} + x^{\frac{1}{2}(m-1)}(m,1) + x^{\frac{1}{2}(m-2)}(m,2) + x^{\frac{1}{2}(m-3)}(m,3) + \text{etc.}$$

This gives

$$\begin{aligned}(1+x^{\frac{1}{2}m+\frac{1}{2}})F(x,m) = 1 &+ x^{\frac{1}{2}}(m,1) + x(m,2) \\ &+ x^{\frac{3}{2}}(m,3) + \text{etc.} \\ &+ x^{\frac{1}{2}} \,.\, x^m + x \,.\, x^{m-1}(m,1) \\ &+ x^{\frac{3}{2}} \,.\, x^{m-2}(m,2) + \text{etc.}\end{aligned}$$

Since we have (Article 5, II)

$$\begin{aligned}(m,1)+x^m &= (m+1,1)\\ (m,2)+x^{m-1}(m,1) &= (m+1,2)\\ (m,3)+x^{m-2}(m,2) &= (m+1,3),\ \text{etc.}\end{aligned}$$

we obtain

$$(1+x^{\frac{1}{2}m+\frac{1}{2}})F(x,m) = F(x,m+1)\ \ldots\ldots\ [3]$$

But $F(x,0)=1$, so that

$$\begin{aligned}F(x,1) &= 1+x^{\frac{1}{2}}\\ F(x,2) &= (1+x^{\frac{1}{2}})(1+x)\\ F(x,3) &= (1+x^{\frac{1}{2}})(1+x)(1+x^{\frac{3}{2}}),\ \text{etc.}\end{aligned}$$

or, in general,

$$\begin{aligned}F(x,m) = (1+x^{\frac{1}{2}})&(1+x)(1+x^{\frac{3}{2}})\\ &\ldots.(1+x^{\frac{1}{2}m})\ \ldots.\quad [4]\end{aligned}$$

**10.**

With these preliminary investigations in place, let us now come closer to our aim. For a prime value of $n$, the squares 1, 4, $9\ldots.(\frac{1}{2}(n-1))^2$ are all incongruent to one another modulo $n$; their least residues for this modulus must therefore be identical to the numbers $a$, and hence

$$\begin{aligned}\Sigma\cos ak\omega &= \cos k\omega+\cos 4k\omega+\cos 9k\omega\\ &\quad+\text{etc.}+\cos(\tfrac{1}{2}(n-1))^2k\omega\\ \Sigma\sin ak\omega &= \sin k\omega+\sin 4k\omega+\sin 9k\omega\\ &\quad+\text{etc.}+\sin(\tfrac{1}{2}(n-1))^2k\omega\end{aligned}$$

Likewise, since the same squares 1, 4, $9\ldots.(\frac{1}{2}(n-1))^2$, in reverse order, are congruent to $(\frac{1}{2}(n+1))^2$, $(\frac{1}{2}(n+3))^2$, $(\frac{1}{2}(n+5))^2\ldots.(n-1)^2$, we also have

$$\begin{aligned}\Sigma\cos ak\omega &= \cos(\tfrac{1}{2}(n+1))^2k\omega\\ &\quad+\cos(\tfrac{1}{2}(n+3))^2k\omega+\text{etc.}+\cos(n-1)^2k\omega\\ \Sigma\sin ak\omega &= \sin(\tfrac{1}{2}(n+1))^2k\omega\\ &\quad+\sin(\tfrac{1}{2}(n+3))^2k\omega+\text{etc.}+\sin(n-1)^2k\omega\end{aligned}$$

Thus, putting

$$\begin{aligned} T &= 1 + \cos k\omega + \cos 4k\omega + \cos 9k\omega \\ &\quad + \text{etc.} + \cos(n-1)^2 k\omega \\ U &= \quad \sin k\omega + \sin 4k\omega + \sin 9k\omega \\ &\quad + \text{etc.} + \sin(n-1)^2 k\omega \end{aligned}$$

we obtain

$$\begin{aligned} 1 + 2\Sigma \cos ak\omega &= T \\ 2\Sigma \sin ak\omega &= U \end{aligned}$$

This shows that the summations proposed in Article 1 depend on summing the series $T$ and $U$. We shall therefore leave the former aside and direct our investigation to these, carrying it out generally enough to include both prime values of $n$ and any composite values. We shall, however, suppose that the number $k$ is relatively prime to $n$: the case where $k$ and $n$ have a common divisor can be reduced to this one without difficulty.

**11.**

Let us denote the imaginary quantity $\sqrt{-1}$ by $i$ and put

$$\cos k\omega + i \sin k\omega = r$$

Then $r^n = 1$, or $r$ is a root of the equation $r^n - 1 = 0$. It is easy to see that all the numbers $k$, $2k$, $3k \ldots . (n-1)k$ are not divisible by $n$ and are incongruent to one another modulo $n$. Thus the powers of $r$

$$1,\ r,\ rr,\ r^3 \ldots . r^{n-1}$$

are all unequal, while each also satisfies the equation $x^n - 1 = 0$. For this reason these powers represent *all* the roots of the equation $x^n - 1 = 0$.

These conclusions would not hold if $k$ had a common divisor with $n$. For if $\nu$ were such a common divisor, $k \,.\, \dfrac{n}{\nu}$ would be divisible by $n$, and hence a power lower than $r^n$, namely $r^{\frac{n}{\nu}}$, would equal unity. In this case, therefore, the powers of $r$ represent at most $\dfrac{n}{\nu}$ roots of the equation $x^n - 1 = 0$, and indeed give that many distinct roots if $\nu$ is the *greatest* common divisor of the numbers $k$, $n$. In our case, where $k$ and $n$ are supposed relatively prime, $r$ can conveniently be called a *proper root* of the equation $x^n - 1 = 0$. In the other case, where $k$ and $n$ have a (greatest) common divisor $\nu$, $r$ would be called an *improper root* of that equation. Then, however, the same $r$ would evidently be a proper root of the equation $x^{\frac{n}{\nu}} - 1 = 0$. The simplest improper root is unity, and when $n$ is a prime number there are no other improper roots at all.

**12.**

Now, if we put

$$W = 1 + r + r^4 + r^9 + \text{etc.} + \text{etc.} + r^{(n-1)^2}$$

we evidently have $W = T + iU$, so that $T$ is the real part of $W$, and $U$ is obtained from the imaginary part of $W$ by removing the factor $i$. The whole question is therefore reduced to finding the sum $W$. For this purpose we can use either the series considered in Article 6 or the one we showed how to sum in Article 9; the former, however, is less suitable when $n$ is even. Nevertheless, we hope that readers will welcome our treating the case where $n$ is odd by two methods.

First, then, suppose that $n$ is an odd number, that $r$ denotes any proper root of the equation $x^n - 1 = 0$, and that in the function $f(x, m)$ we put $x = r$ and $m = n - 1$. We evidently obtain

$$\begin{aligned}
\frac{1 - x^m}{1 - x} &= \frac{1 - r^{-1}}{1 - r} = -r^{-1} \\
\frac{1 - x^{m-1}}{1 - xx} &= \frac{1 - r^{-2}}{1 - rr} = -r^{-2} \\
\frac{1 - x^{m-2}}{1 - x^3} &= \frac{1 - r^{-3}}{1 - r^3} = -r^{-3}, \text{ etc.}
\end{aligned}$$

as far as

$$\frac{1 - x}{1 - x^m} = \frac{1 - r^{-m}}{1 - r^m} = -r^{-m}$$

(It is worth pointing out that these equations hold only insofar as $r$ is supposed to be a proper root: if $r$ were an improper root, the numerator and denominator of some of these fractions would vanish together, and the fractions would therefore become indeterminate).

From this we deduce the following equation

$$\begin{aligned}
f(r, n-1) &= 1 + r^{-1} + r^{-3} + r^{-6} + \text{etc.} + r^{-\frac{1}{2}(n-1)n} \\
&= (1 - r)(1 - r^3)(1 - r^5) \ldots . (1 - r^{n-2})
\end{aligned}$$

The same equation still holds if we replace $r$ by $r^\lambda$, where $\lambda$ denotes any integer relatively prime to $n$: for then $r^\lambda$ too is a proper root of the equation $x^n - 1 = 0$. Let us therefore replace $r$ by $r^{n-2}$, or equivalently $r^{-2}$. We obtain

$$\begin{aligned}
&1 + r^2 + r^6 + r^{12} + \text{etc.} + r^{(n-1)n} \\
&\quad = (1 - r^{-2})(1 - r^{-6})(1 - r^{-10}) \ldots (1 - r^{-2(n-2)})
\end{aligned}$$

Let us multiply both sides of this equation by

$$r \,.\, r^3 \,.\, r^5 \,.\,.\,.\,. \, r^{(n-2)} = r^{\frac{1}{4}(n-1)^2}$$

and, since

$$\begin{aligned}
r^{2+\frac{1}{4}(n-1)^2} &= r^{\frac{1}{4}(n-3)^2}, \\
r^{(n-1)n+\frac{1}{4}n-1)^2} &= r^{\frac{1}{4}(n+1)^2} \\
r^{6+\frac{1}{4}(n-1)^2} &= r^{\frac{1}{4}(n-5)^2}, \\
r^{(n-2)(n-1)+\frac{1}{4}(n-1)^2} &= r^{\frac{1}{4}(n+3)^2} \\
r^{12+\frac{1}{4}(n-1)^2} &= r^{\frac{1}{4}(n-7)^2}, \\
r^{(n-3)(n-2)+\frac{1}{4}(n-1)^2} &= r^{\frac{1}{4}(n+5)^2}, \text{ etc.}
\end{aligned}$$

the following equation results:

$$\begin{aligned}
&r^{\frac{1}{4}(n-1)^2} + r^{\frac{1}{4}(n-3)^2} + r^{\frac{1}{4}(n-5)^2} + \text{etc.} + r + 1 \\
&+ r^{\frac{1}{4}(n+1)^2} + r^{\frac{1}{4}(n+3)^2} + r^{\frac{1}{4}(n+5)^2} \\
&\qquad + \text{etc.} + r^{\frac{1}{4}(2n-2)^2} \\
&\qquad = (r - r^{-1})(r^3 - r^{-3})(r^5 - r^{-5}) \,.\,.\,.\,. \, (r^{n-2} - r^{-n+2})
\end{aligned}$$

or, arranging the terms on the left side differently,

$$\begin{aligned}
&1 + r + r^4 + \text{etc.} + r^{(n-1)^2} \\
&\qquad = (r - r^{-1})(r^3 - r^{-3}) \,.\,.\, (r^{n-2} - r^{-n+2}) \quad .. \quad [5]
\end{aligned}$$

**13.**

The factors on the right side of equation [5] can also be written as follows:

$$r - r^{-1} = -(r^{n-1} - r^{-n+1})$$
$$r^3 - r^{-3} = -(r^{n-3} - r^{-n+3})$$
$$r^5 - r^{-5} = -(r^{n-5} - r^{-n+5}), \text{ etc.}$$

up to

$$r^{n-2} - r^{-n+2} = -(r^2 - r^{-2})$$

The equation then takes this form:

$$W = (-1)^{\frac{1}{2}(n-1)}(r^2 - r^{-2})(r^4 - r^{-4})(r^6 - r^{-6}) \\ \ldots . (r^{n-1} - r^{-n+1})$$

Multiplying this equation by [5] in its original form gives

$$W^2 = (-1)^{\frac{1}{2}(n-1)}(r - r^{-1})(r^2 - r^{-2})(r^3 - r^{-3}) \\ \ldots . (r^{n-1} - r^{-n+1})$$

where $(-1)^{\frac{1}{2}(n-1)}$ is either $= +1$ or $= -1$, according as $n$ is of the form $4\mu + 1$ or of the form $4\mu + 3$. Hence

$$W^2 = \pm r^{\frac{1}{2}n(n-1)}(1 - r^{-2})(1 - r^{-4})(1 - r^{-6}) \\ \ldots . (1 - r^{-2(n-1)})$$

But it is easy to see that $r^{-2}$, $r^{-4}$, $r^{-6} \ldots . r^{-2n+2}$ give all the roots of the equation $x^n - 1 = 0$ except the root $x = 1$. The following identity must therefore hold for an indeterminate:

$$(x - r^{-2})(x - r^{-4})(x - r^{-6}) \ldots . (x - r^{-2n+2}) \\ = x^{n-1} + x^{n-2} + x^{n-3} + \text{etc.} + x + 1$$

Thus, putting $x = 1$, we obtain

$$(1 - r^{-2})(1 - r^{-4})(1 - r^{-6}) \ldots . (1 - r^{-2n+2}) = n$$

and since evidently $r^{\frac{1}{2}n(n-1)} = 1$, our equation becomes

$$W^2 = \pm n \ldots . \ldots . \ldots . [6]$$

Thus, in the case where $n$ is of the form $4\mu + 1$, we have

$$W = \pm\sqrt{n}, \quad \text{and hence} \quad T = \pm\sqrt{n}, \quad U = 0$$

In the other case, where $n$ is of the form $4\mu + 3$, we instead have

$$W = \pm i\sqrt{n}, \quad \text{and so} \quad T = 0, \quad U = \pm\sqrt{n}$$

**14.**

The method of the preceding article gives only the absolute value of the sums $T$, $U$, and leaves it uncertain whether $T$ in the first case and $U$ in the second should be set $= +\sqrt{n}$ or $= -\sqrt{n}$. But, at least for the case where $k = 1$, this can be decided from equation [5] as follows. Since, for $k = 1$,

$$\begin{aligned} r - r^{-1} &= 2i \sin \omega \\ r^3 - r^{-3} &= 2i \sin 3\omega \\ r^5 - r^{-5} &= 2i \sin 5\omega \text{ etc.} \end{aligned}$$

that equation becomes

$$W = (2i)^{\frac{1}{2}(n-1)} \sin\omega \sin 3\omega \sin 5\omega \ldots \sin(n-2)\omega$$

Now, in the case where $n$ is of the form $4\mu+1$, the sequence of odd numbers

$$1,\ 3,\ 5,\ 7 \ldots . \tfrac{1}{2}(n-3), \quad \tfrac{1}{2}(n+1) \ldots . (n-2)$$

contains $\frac{1}{4}(n-1)$ numbers smaller than $\frac{1}{2}n$, to which positive sines evidently correspond. The remaining $\frac{1}{4}(n-1)$, on the other hand, are greater than $\frac{1}{2}n$, and negative sines correspond to them. The product of all the sines must therefore equal a positive quantity multiplied by $(-1)^{\frac{1}{4}(n-1)}$. Thus $W$ will equal a positive real quantity multiplied by $i^{n-1}$, or by 1, since $i^4 = 1$ and $n-1$ is divisible by 4: that is, $W$ will be real and positive. Hence we must have

$$W = +\sqrt{n}, \quad T = +\sqrt{n}$$

In the other case, where $n$ is of the form $4\mu + 3$, in the sequence of odd numbers

$$1,\ 3,\ 5,\ 7 \ldots . \tfrac{1}{2}(n-1), \quad \tfrac{1}{2}(n+3) \ldots . (n-2)$$

the first $\frac{1}{4}(n+1)$ will be smaller than $\frac{1}{2}n$, but the remaining $\frac{1}{4}(n-3)$ will be greater. Thus, among the sines of the arcs $\omega$, $3\omega$, $5\omega \ldots . (n-2)\omega$, there will be $\frac{1}{4}(n-3)$ negative ones, and so $W$ will be the product of $i^{\frac{1}{2}(n-1)}$, a positive real quantity and $(-1)^{\frac{1}{4}(n-3)}$. The third factor is $= i^{\frac{1}{2}(n-3)}$, which, multiplied by the first, gives $i^{n-2} = i$, since $i^{n-3} = 1$. We must therefore have

$$W = +i\sqrt{n}, \quad \text{and} \quad U = +\sqrt{n}$$

**15.**

We shall now show how the same conclusions can be derived from the series considered in Article 9. In equation [4], replace $x^{\frac{1}{2}}$ by $-y^{-1}$; we then obtain

$$1 - y^{-1}\frac{1-y^{-2m}}{1-y^{-2}} + y^{-2}\frac{(1-y^{-2m})(1-y^{-2m+2})}{(1-y^{-2})(1-y^{-4})} - y^{-3}\frac{(1-y^{-2m})(1-y^{-2m+2})(1-y^{-2m+4})}{(1-y^{-2})(1-y^{-4})(1-y^{-6})} + \text{etc.}$$

up to the term in position $m+1$

$$= (1-y^{-1})(1+y^{-2})(1-y^{-3})(1+y^{-4}) \ldots (1 \pm y^{-m}) \ldots . \text{ [7]}$$

If we take for $y$ a proper root of the equation $y^n - 1 = 0$, say $r$, and at the same time put $m = n-1$, we obtain

$$\frac{1-y^{-2m}}{1-y^{-2}} = \frac{1-r^2}{1-r^{-2}} = -r^2$$
$$\frac{1-y^{-2m+2}}{1-y^{-4}} = \frac{1-r^4}{1-r^{-4}} = -r^4$$
$$\frac{1-y^{-2m+4}}{1-y^{-6}} = \frac{1-r^6}{1-r^{-6}} = -r^6$$

up to

$$\frac{1-y^{-2}}{1-y^{-2m}} = \frac{1-r^{2n-2}}{1-r^{-2n+2}} = -r^{2n-2}$$

Here we should note that none of the denominators $1-r^{-2}$, $1-r^{-4}$, etc. becomes $= 0$. Equation [7] therefore takes this form:

$$1 + r + r^4 + r^9 + \text{etc.} + r^{(n-1)^2} = (1-r^{-1})(1+r^{-2})(1-r^{-3}) \ldots . (1+r^{-n+1})$$

Multiplying, on the right side of this equation, the first term by the last, the second by the next to last, etc., we obtain

$$(1-r^{-1})(1+r^{-n+1}) = r - r^{-1}$$
$$(1+r^{-2})(1-r^{-n+2}) = r^{n-2} - r^{-n+2}$$
$$(1-r^{-3})(1+r^{-n+3}) = r^3 - r^{-3}$$
$$(1+r^{-4})(1-r^{-n+4}) = r^{n-4} - r^{-n+4} \text{ etc.}$$

It is easy to see that these partial products combine to form the product

$$(r - r^{-1})(r^3 - r^{-3})(r^5 - r^{-5}) \\ \ldots . (r^{n-4} - r^{-n+4})(r^{n-2} - r^{-n+2})$$

which will therefore be

$$= 1 + r + r^4 + r^9 + \text{etc.} + r^{(n-1)^2} = W$$

This equation is identical with equation [5], derived in Article 12 from the first series; the remaining arguments are then established in the same way as in Articles 13 and 14.

**16.**

We turn to the other case, where $n$ is an even number. First let $n$ be of the form $4\mu + 2$, or twice an odd number. It is clear that the numbers $\frac{1}{4}nn$, $(\frac{1}{2}n + 1)^2 - 1$, $(\frac{1}{2}n + 2)^2 - 4$, etc., or in general $(\frac{1}{2}n + \lambda)^2 - \lambda\lambda$, give odd quotients when divided by $\frac{1}{2}n$, and are therefore congruent, modulo $n$, to $\frac{1}{2}n$. It follows that, if $r$ is a proper root of the equation $x^n - 1 = 0$, and hence $r^{\frac{1}{2}n} = -1$, we have

$$\begin{aligned} r^{(\frac{1}{2}n)^2} &= -1 \\ r^{(\frac{1}{2}n+1)^2} &= -r \\ r^{(\frac{1}{2}n+2)^2} &= -r^4 \\ r^{(\frac{1}{2}n+2)^2} &= -r^9 \text{ etc.} \end{aligned}$$

Thus, in the series

$$1 + r + r^4 + r^9 + \text{etc.} + r^{(n-1)^2}$$

the term $r^{(\frac{1}{2}n)^2}$ cancels the first, the next cancels the second, etc., so that

$$W = 0, \quad T = 0, \quad U = 0$$

**17.**

The case remains where $n$ is of the form $4\mu$, or divisible by four. Here, in general, $(\frac{1}{2}n+\lambda)^2 - \lambda\lambda$ is divisible by $n$, and therefore

$$r^{(\frac{1}{2}n+\lambda)^2} = r^{\lambda\lambda}$$

Thus, in the series

$$1 + r + r^4 + r^9 + \text{etc.} + r^{(n-1)^2}$$

the term $r^{(\frac{1}{2}n)^2}$ equals the first, the next equals the second, etc., so that

$$W = 2(1 + r + r^4 + r^9 + \text{etc.} + r^{(\frac{1}{2}n-1)^2})$$

Now suppose that in equation [7] of Article 15 we put $m = \frac{1}{2}n - 1$, and take for $y$ a proper root of the equation $y^n - 1 = 0$, say $r$. Then, just as in Article 15, the equation takes the following form:

$$1 + r + r^4 + \text{etc.} + r^{(\frac{1}{2}n-1)^2} = (1 - r^{-1})(1 + r^{-2})(1 - r^{-3}) \ldots . (1 - r^{-\frac{1}{2}n+1})$$

or

$$W = 2(1 - r^{-1})(1 + r^{-2})(1 - r^{-3})(1 + r^{-4}) \ldots (1 - r^{-\frac{1}{2}n+1}) \quad .. \quad [8]$$

Further, since $r^{\frac{1}{2}n} = -1$, and therefore

$$1 + r^{-2} = -r^{\frac{1}{2}n-2}(1 - r^{-\frac{1}{2}n+2})$$
$$1 + r^{-4} = -r^{\frac{1}{2}n-4}(1 - r^{-\frac{1}{2}n+4})$$
$$1 + r^{-6} = -r^{\frac{1}{2}n-6}(1 - r^{-\frac{1}{2}n+6}) \text{ etc.}$$

and the product of the factors $-r^{\frac{1}{2}n-2}$, $-r^{\frac{1}{2}n-4}$, $-r^{\frac{1}{2}n-6}$, etc., up to $-r^2$ is $= (-1)^{\frac{1}{4}n-1} r^{\frac{1}{16}nn-\frac{1}{4}n}$, the preceding equation can also be written as

$$W = 2(-1)^{\frac{1}{4}n-1} r^{\frac{1}{16}nn-\frac{1}{4}n}(1 - r^{-1})(1 - r^{-2}) (1 - r^{-3})(1 - r^{-4}) \ldots . (1 - r^{-\frac{1}{2}n+1})$$

Since we have

$$1 - r^{-1} = -r^{-1}(1 - r^{-n+1})$$
$$1 - r^{-2} = -r^{-2}(1 - r^{-n+2})$$
$$1 - r^{-3} = -r^{-3}(1 - r^{-n+3}) \text{ etc.}$$

we obtain

$$(1 - r^{-1})(1 - r^{-2})(1 - r^{-3}) \dots (1 - r^{-\frac{1}{2}n+1})$$
$$= (-1)^{\frac{1}{2}n-1} r^{-\frac{1}{8}nn+\frac{1}{4}n}(1 - r^{-\frac{1}{2}n-1})$$
$$(1 - r^{-\frac{1}{2}n-2})(1 - r^{-\frac{1}{2}n-3}) \dots (1 - r^{-n+1})$$

and hence

$$W = 2(-1)^{\frac{3}{4}n-2} r^{-\frac{1}{16}nn}(1 - r^{-\frac{1}{2}n-1})$$
$$(1 - r^{-\frac{1}{2}n-2})(1 - r^{-\frac{1}{2}n-3}) \dots . (1 - r^{-n+1})$$

Multiplying this value of $W$ by the one found earlier, and introducing the factor $1 - r^{-\frac{1}{2}n}$ on both sides, gives

$$(1 - r^{-\frac{1}{2}n})W^2 = 4(-1)^{n-3} r^{-\frac{1}{4}n}(1 - r^{-1}.)$$
$$(1 - r^{-2})(1 - r^{-3}) \dots . (1 - r^{-n+1})$$

But we have

$$1 - r^{-\frac{1}{2}n} = 2$$
$$(-1)^{n-3} = -1$$
$$r^{-\frac{1}{4}n} = -r^{\frac{1}{4}n}$$

$$(1 - r^{-1})(1 - r^{-2})(1 - r^{-3}) \dots . (1 - r^{-n+1}) = n$$

From this we finally conclude

$$W^2 = 2r^{\frac{1}{4}n} n \text{ . . . . . [9]}$$

Now it is easy to see that $r^{\frac{1}{4}n}$ is either $= +i$ or $= -i$, according as $k$ is of the form $4\mu + 1$ or of the form $4\mu + 3$. And since

$$2i = (1 + i)^2, \quad -2i = (1 - i)^2$$

in the case where $k$ is of the form $4\mu + 1$, we have

$$W = \pm(1 + i)\sqrt{n}, \quad \text{and so} \quad T = U = \pm\sqrt{n}$$

but in the other case, where $k$ is of the form $4\mu + 3$,

$$W = \pm(1-i)\sqrt{n}, \quad \text{and so} \quad T = -U = \pm\sqrt{n}$$

**18.**

The method of the preceding article supplied the absolute values of the functions $T$, $U$, and specified the conditions under which they must be given equal or opposite signs; but it does not yet determine the signs themselves. For the case where $k = 1$, we shall supply what is missing as follows.

Let us put $\rho = \cos\frac{1}{2}\omega + i\sin\frac{1}{2}\omega$, so that $r = \rho\rho$. Since $\rho^n = -1$, it is clear that equation [8] can be written as

$$W = 2(1+\rho^{n-2})(1+\rho^{-4})(1+\rho^{n-6})(1+\rho^{-8})$$
$$\ldots(1+\rho^{-n+4})(1+\rho^2)$$

or, with the factors arranged in a different order,

$$W = 2(1+\rho^2)(1+\rho^{-4})(1+\rho^6)(1+\rho^{-8})$$
$$\ldots(1+\rho^{-n+4})(1+\rho^{n-2})$$

Now we have

$$1+\rho^2 = 2\rho\cos\frac{1}{2}\omega$$
$$1+\rho^{-4} = 2\rho^{-2}\cos\omega$$
$$1+\rho^{-6} = 2\rho^3\cos\frac{3}{2}\omega$$
$$1+\rho^{-8} = 2\rho^{-4}\cos 2\omega, \text{ etc.}$$

up to

$$1+\rho^{-n+4} = 2\rho^{-\frac{1}{2}n+2}, \cos(\frac{1}{4}n-1)\omega$$
$$1+\rho^{n-2} = 2\rho^{\frac{1}{2}n-1}\cos(\frac{1}{4}n-\frac{1}{2})\omega$$

Thus we have

$$W = 2^{\frac{1}{2}n}\rho^{\frac{1}{4}n}\cos\frac{1}{2}\omega\cos\omega\,\frac{3}{2}\omega$$
$$\ldots\cos(\frac{1}{4}n-\frac{1}{2})\omega$$

The cosines entering this product are evidently all positive, while the factor $\rho^{\frac{1}{4}n}$ is $= \cos 45^0 + i \sin 45^0 = (1+i)\sqrt{\frac{1}{2}}$. From this we conclude that $W$ is the product of $1+i$ and a positive real quantity, so that necessarily

$$W = (1+i)\sqrt{n}, \quad T = +\sqrt{n}, \quad U = +\sqrt{n}$$

**19.**

It will be worthwhile to collect all the summations developed so far into a single overview here. In general, we have

| $T =$ | $U =$ | for $n$ of the form |
|---|---|---|
| $\pm\sqrt{n}$ | $\pm\sqrt{n}$ | $4\mu$ |
| $\pm\sqrt{n}$ | $0$ | $4\mu+1$ |
| $0$ | $0$ | $4\mu+2$ |
| $0$ | $\pm\sqrt{n}$ | $4\mu+3$ |

and, when we set $k = 1$, the radical must be given the positive sign. What we observed by induction in Article 3 for prime values of $n$ has therefore now been proved with full rigor. Nothing remains except to show how to determine the signs in every case, for any values of $k$. But before we can tackle this task in full generality, we must first consider more closely the cases where $n$ is a prime number or a power of a prime number.

**20.**

First let $n$ be an odd prime number. From what we set out in Article 10, it is clear that $W = 1 + 2\Sigma r^a = 1 + 2\Sigma R^{ak}$ if we put $R = \cos\omega + i\sin\omega$, with $a$, as there, denoting indefinitely all the quadratic residues of $n$ lying between 1 and $n-1$. If we likewise use $b$ to denote indefinitely all the quadratic nonresidues between the same limits, we readily see that all the numbers $ak$ are congruent modulo $n$ either to all the $a$ or to all the $b$ (without regard to order), according as $k$ is a residue or a nonresidue. Thus, in the first case, we have

$$W = 1 + 2\Sigma R^a = 1 + R + R^4 + R^9 + \text{etc.} + R^{(n-1)^2}$$

and hence $W = +\sqrt{n}$ if $n$ is of the form $4\mu+1$, and $W = +i\sqrt{n}$ if $n$ is of the form $4\mu+3$.

In the other case, where $k$ is a nonresidue of $n$, we have

$$W = 1 + 2\Sigma R^b$$

Since all the $a$, $b$ evidently make up the complete collection of numbers 1, 2, 3 …, and therefore

$$\Sigma R^a + \Sigma R^b = R + R^2 + R^3 + \text{etc.} + R^{n-1} = -1$$

we obtain

$$W = -1 - 2\Sigma R^a = -(1 + R + R^4 + R^9 + \text{etc.} + R^{(n-1)^2})$$

and hence $W = -\sqrt{n}$ if $n$ is of the form $4\mu + 1$, and $W = -i\sqrt{n}$ if $n$ is of the form $4\mu + 3$.

From this we conclude:

*first*, if $n$ is of the form $4\mu + 1$ and $k$ is a quadratic residue of $n$,

$$T = +\sqrt{n}, \quad U = 0$$

*second*, if $n$ is of the form $4\mu + 1$ and $k$ is a nonresidue of $n$,

$$T = -\sqrt{n}, \quad U = 0$$

*third*, if $n$ is of the form $4\mu + 3$ and $k$ is a residue of $n$,

$$T = 0, \quad U = +\sqrt{n}$$

*fourth*, if $n$ is of the form $4\mu + 3$ and $k$ is a nonresidue of $n$,

$$T = 0, \quad U = -\sqrt{n}$$

**21.**

Second, let $n$ be the square or a higher power of an odd prime number $p$, and put $n = p^{2\varkappa}q$, with $q$ either $= 1$ or $= p$. Here we should first note that, if $\lambda$ is any integer not divisible by $p^{\varkappa}$, we have

$$\begin{aligned} & r^{\lambda\lambda} + r^{(\lambda+p^{\varkappa}q)^2} + r^{(\lambda+2p^{\varkappa}q)^2} + r^{(\lambda+3p^{\varkappa}q)^2} \\ & \quad + \text{etc.} + r^{(\lambda+n-p^{\varkappa}q)^3} \\ & = r^{\lambda\lambda}\{1 + r^{2\lambda p^{\varkappa}q} + r^{4\lambda p^{\varkappa}q} \\ & \quad + r^{6\lambda p^{\varkappa}q} + \text{etc.} + r^{2\lambda(n-p^{\varkappa}q)}\} \\ & = \frac{r^{\lambda\lambda}(1 - r^{2\lambda n})}{1 - r^{2\lambda p^{\varkappa}q}} = 0 \end{aligned}$$

It is then easy to see that

$$W = 1 + r^{p^{2\varkappa}} + r^{4p^{2\varkappa}} + r^{9p^{2\varkappa}} + \text{etc.} + r^{(n-p^{\varkappa})^2}$$

For the remaining terms of the series

$$1 + r + r^4 + r^9 + \text{etc.} + r^{(n-1)^2}$$

can be divided into $(p^\varkappa - 1)q$ partial series, each with $p^\varkappa$ terms, whose sums vanish under the transformation just given.

It follows that in the case where $q = 1$, or where $n$ is a power of a prime number with an even exponent, we have

$$W = p^\varkappa = +\sqrt{n}, \text{ and so } T = +\sqrt{n}, \quad U = 0$$

In the other case, where $q = p$, or where $n$ is a power of a prime number with an odd exponent, we shall put $r^{p^{2\varkappa}} = \rho$. Then $\rho$ is a proper root of the equation $x^p - 1 = 0$, namely $\rho = \cos \frac{k}{p} 360^0 + i \sin \frac{k}{p} 360^0$, and we obtain

$$\begin{aligned} W &= 1 + \rho + \rho^4 + \rho^9 + \text{etc.} + \rho^{(p^{\varkappa+1}-1)^2} \\ &= p^\varkappa (1 + \rho + \rho^4 + \rho^9 + \text{etc.} + \rho^{(p-1)^2}) \end{aligned}$$

But the sum of the series $1 + \rho + \rho^4 + \rho^9 + \text{etc.} + \rho^{(p-1)^2}$ is determined by the preceding article. It follows directly that

$$\begin{aligned} W &= \pm\sqrt{n} = T, \quad \text{if } p \text{ is of the form } 4\mu + 1 \\ W &= \pm i\sqrt{n} = iU, \quad \text{if } p \text{ is of the form } 4\mu + 3 \end{aligned}$$

with the positive or negative sign, according as $k$ is a residue or a nonresidue of $p$.

**22.**

What we set out in Articles 20 and 21 also readily gives the following proposition, which will be of considerable use to us below. Put

$$W' = 1 + r^h + r^{4h} + r^{9h} + \text{etc.} + r^{h(n-1)^2}$$

with $h$ denoting any integer not divisible by $p$. Then, in the case where $n = p$, or where $n$ is a power of $p$ with an odd exponent, we have

$$\begin{aligned} W' &= W, \text{ if } h \text{ is a quadratic residue of } p \\ W' &= -W, \text{ if } h \text{ is a quadratic nonresidue of } p \end{aligned}$$

For $W'$ evidently arises from $W$ if, for $k$, we substitute $kh$. In the first case, $k$ and $kh$ are alike, and in the second they are unlike, with respect to being residues or nonresidues of $p$.

But in the case where $n$ is a power of $p$ with an even exponent, we evidently have $W' = +\sqrt{n}$, and therefore always $W' = W$.

**23.**

In Articles 20, 21 and 22 we considered odd prime numbers and powers of such numbers. The case therefore remains where $n$ is a power of two.

For $n = 2$, we evidently have $W = 1 + r = 0$.

For $n = 4$, we obtain $W = 1 + r + r^4 + r^9 = 2 + 2r$. Thus $W = 2 + 2i$ whenever $k$ is of the form $4\mu + 1$, and $W = 2 - 2i$ whenever $k$ is of the form $4\mu + 3$.

For $n = 8$, we have $W = 1+r+r^4+r^9+r^{16}+r^{25}+r^{36}+r^{49} = 2+4r+2r^4 = 4r$. Hence

$$\begin{aligned} W &= (1+i)\sqrt{8}, \quad \text{whenever } k \text{ is of the form } 8\mu+1 \\ W &= (-1+i)\sqrt{8}, \quad \text{whenever } k \text{ is of the form } 8\mu+3 \\ W &= (-1-i)\sqrt{8}, \quad \text{whenever } k \text{ is of the form } 8\mu+5 \\ W &= (1-i)\sqrt{8}, \quad \text{whenever } k \text{ is of the form } 8\mu+7 \end{aligned}$$

If $n$ is a higher power of two, put $n = 2^{2\varkappa}q$, with $q$ either $= 1$ or $= 2$, and $\varkappa$ greater than 1. Here we must first observe that, if $\lambda$ is any integer not divisible by $2^{\varkappa-1}$, we have

$$\begin{aligned} & r^{\lambda\lambda} + r^{(\lambda+2^{\varkappa}q)^2} + r^{(\lambda+2.2^{\varkappa}q)^2} + r^{(\lambda+3.2^{\varkappa}q)^2} \\ & \quad + \text{etc.} + r^{(\lambda+n-2^{\varkappa}q)^2} \\ & = r^{\lambda\lambda}\{1 + r^{2^{\varkappa+1}\lambda q} + r^{2.2^{\varkappa+1}\lambda q} \\ & \quad + r^{3.2^{\varkappa+1}\lambda q} + \text{etc.} + r^{(2n-2^{\varkappa+1}q)\lambda}\} \\ & = \frac{r^{\lambda\lambda}(1 - r^{2\lambda n})}{1 - r^{2^{\varkappa+1}\lambda q}} = 0 \end{aligned}$$

It is then easy to see that

$$W = 1 + r^{2^{2\varkappa-2}} + r^{4.2^{2\varkappa-2}} + r^{9.2^{2\varkappa-2}} + \text{etc.} + r^{(n-2^{\varkappa-1})^2}$$

Put $r^{2^{2\varkappa-2}} = \rho$. Then $\rho$ is a root of the equation $x^{4q} - 1 = 0$, namely $\rho = \cos\frac{k}{4q}360^0 + i\sin\frac{k}{4q}360^0$; we then obtain

$$\begin{aligned} W &= 1 + \rho + \rho^4 + \rho^9 + \text{etc.} + \rho^{(2^{\varkappa+1}q-1)^2} \\ &= 2^{\varkappa-1}(1 + \rho + \rho^4 + \rho^9 + \text{etc.} + \rho^{(4q-1)^2}) \end{aligned}$$

But the sum of the series $1 + \rho + \rho^4 + \rho^9 + \text{etc.} + \rho^{(4q-1)^2}$ is determined by what we explained for the cases $n = 4$, $n = 8$. From this we conclude

that in the case where $q = 1$, or where $n$ is a power of the number 4, we have

$$W = (1+i)2^{\varkappa} = (1+i)\sqrt{n}, \quad \text{if } k \text{ is of the form } 4\mu + 1$$
$$W = (1-i)2^{\varkappa} = (1-i)\sqrt{n}, \quad \text{if } k \text{ is of the form } 4\mu + 3$$

these are exactly the formulas given for $n = 4$;

but in the case where $q = 2$, or where $n$ is a power of two with an odd exponent greater than 3, we have

$$W = (1+i)2^{\varkappa}\sqrt{2} = (1+i)\sqrt{n}, \quad \text{if } k \text{ is of the form } 8\mu + 1$$
$$W = (-1+i)2^{\varkappa}\sqrt{2} = (-1+i)\sqrt{n}, \quad \text{if } k \text{ is of the form } 8\mu + 3$$
$$W = (-1-i)2^{\varkappa}\sqrt{2} = (-1-i)\sqrt{n}, \quad \text{if } k \text{ is of the form } 8\mu + 5$$
$$W = (1-i)2^{\varkappa}\sqrt{2} = (1-i)\sqrt{n}, \quad \text{if } k \text{ is of the form } 8\mu + 7$$

these too agree exactly with those we gave for $n = 8$.

**24.**

Here too it will be worthwhile to determine the ratio of the sum of the series

$$W' = 1 + r^h + r^{4h} + r^{9h} + \text{etc.} + r^{h(n-1)^2}$$

to $W$, where $h$ denotes any odd integer. Since $W'$ arises from $W$ by changing $k$ into $kh$, the value of $W'$ depends on the form of the number $kh$ in the same way as $W$ depends on the form of $k$. Put $\frac{W'}{W} = l$. It is clear that

I. in the case where $n = 4$, or it is a higher power of two with an even exponent, we have

$$l = 1, \quad \text{if } h \text{ is of the form } 4\mu + 1$$
$$l = -i, \quad \text{if } h \text{ is of the form } 4\mu + 3, \text{ and } k \text{ is of the form } 4\mu + 1$$
$$l = +i, \quad \text{if } h \text{ is of the form } 4\mu + 3, \text{ and } k \text{ is of the same form.}$$

II. in the case where $n = 8$, or it is a higher power of two with an odd exponent, we have

$l = 1$, if $h$ is of the form $8\mu + 1$,
$l = -1$, if $h$ is of the form $8\mu + 5$,
$l = +i$, if either $h$ is of the form $8\mu + 3$, and $k$ is of the form $4\mu + 1$,
or $h$ is of the form $8\mu + 7$, and $k$ is of the form $4\mu + 3$,
$l = -i$, if either $h$ is of the form $8\mu + 3$, and $k$ is of the form $4\mu + 3$,
or $h$ is of the form $8\mu + 7$, and $k$ is of the form $4\mu + 1$.

The preceding articles have completely determined the sum $W$ for the cases where $n$ is a prime number or a power of a prime number. It therefore remains to complete the cases where $n$ is composed of several prime numbers; the following theorem will prepare the way for us:

**25.**

THEOREM. *Let $n$ be the product of two coprime positive integers $a$, $b$, and put*

$$P = 1 + r^{aa} + r^{4aa} + r^{9aa} + \text{etc.} + r^{(b-1)^2 aa}$$
$$Q = 1 + r^{bb} + r^{4bb} + r^{9bb} + \text{etc.} + r^{(a-1)^2 bb}$$

*Then I claim that $W = PQ$.*

*Proof.* Let $\alpha$ denote indefinitely the numbers $0, 1, 2, 3 \ldots . a - 1$, $\beta$ the numbers $0, 1, 2, 3 \ldots . b - 1$, and $\nu$ the numbers $0, 1, 2, 3 \ldots . n - 1$. Then evidently

$$P = \Sigma r^{aa\beta\beta}, \quad Q = \Sigma r^{bb\alpha\alpha}, \quad W = \Sigma r^{\nu\nu}$$

Thus $PQ = \Sigma r^{aa\beta\beta + bb\alpha\alpha}$, on substituting all the values of $\alpha$ and $\beta$, combined with one another in every way. Further, since $2ab\alpha\beta = 2\alpha\beta n$, we have $PQ = \Sigma r^{(a\beta + b\alpha)^2}$. But it is easy to see that the individual values of $a\beta + b\alpha$ are different from one another and equal to some value of $\nu$. Hence $PQ = \Sigma r^{\nu\nu} = W$.

It should also be noted that $r^{aa}$ is a proper root of the equation $x^b - 1 = 0$, and $r^{bb}$ a proper root of the equation $x^a - 1 = 0$.

**26.**

Now let $n$ be the product of three pairwise coprime numbers $a, b, c$. It is clear that, on putting $bc = b'$, $a$ and $b'$ will also be coprime; hence $W$ is the product of the two factors

$$1 + r^{aa} + r^{4aa} + r^{9aa} + \text{etc.} + r^{(b'-1)^2 aa}$$
$$1 + r^{b'b'} + r^{4b'b'} + r^{9b'b'} + \text{etc.} + r^{(a-1)^2 b'b'}$$

But since $r^{aa}$ is a proper root of the equation $x^{bc} - 1 = 0$, the first factor itself will be the product of

$$1 + \rho^{bb} + \rho^{4bb} + \rho^{9bb} + \text{etc.} + \rho^{(c-1)^2 bb}$$
$$1 + \rho^{cc} + \rho^{4cc} + \rho^{9cc} + \text{etc.} + \rho^{(b-1)^2 cc}$$

if we put $r^{aa} = \rho$. It is therefore clear that $W$ is the product of the three factors

$$1 + r^{bbcc} + r^{4bbcc} + r^{9bbcc} + \text{etc.} + r^{(a-1)^2 bbcc}$$
$$1 + r^{aacc} + r^{4aacc} + r^{9aacc} + \text{etc.} + r^{(b-1)^2 aacc}$$
$$1 + r^{aabb} + r^{4aabb} + r^{9aabb} + \text{etc.} + r^{(c-1)^2 aabb}$$

where $r^{bbcc}$, $r^{aacc}$, $r^{aabb}$ are respectively proper roots of the equations $x^a - 1 = 0$, $x^b - 1 = 0$, $x^c - 1 = 0$.

**27.**

It is easy to conclude in general that, if $n$ is the product of any number of pairwise coprime factors $a, b, c$, etc., $W$ is the product of the same number of factors, namely

$$1 + r^{\frac{nn}{aa}} + r^{\frac{4nn}{aa}} + r^{\frac{9nn}{aa}} + \text{etc.} + r^{\frac{(a-1)^2 nn}{aa}}$$
$$1 + r^{\frac{nn}{bb}} + r^{\frac{4nn}{bb}} + r^{\frac{9nn}{bb}} + \text{etc.} + r^{\frac{(b-1)^2 nn}{bb}}$$
$$1 + r^{\frac{nn}{cc}} + r^{\frac{4nn}{cc}} + r^{\frac{9nn}{cc}} + \text{etc.} + r^{\frac{(c-1)^2 nn}{cc}}, \text{ etc.}$$

where $r^{\frac{nn}{aa}}$, $r^{\frac{nn}{bb}}$, $r^{\frac{nn}{cc}}$, etc. are proper roots of the equations $x^a - 1 = 0$, $x^b - 1 = 0$. $x^c - 1 = 0$, etc.

**28.**

These principles now make the transition to a complete determination of $W$ for any value of $n$ immediately clear. Specifically, resolve $n$ into factors $a, b, c$, etc., that are either distinct prime numbers or powers of distinct prime numbers, and put $r^{\frac{nn}{aa}} = A$, $r^{\frac{nn}{bb}} = B$, $r^{\frac{nn}{cc}} = C$, etc. Then $A, B, C$, etc. will be proper roots of the equations $x^a - 1 = 0$, $x^b - 1 = 0$, $x^c - 1 = 0$, etc., and $W$ the product of the factors

$$1 + A + A^4 + A^9 + \text{etc.} + A^{(a-1)^2}$$
$$1 + B + B^4 + B^9 + \text{etc.} + B^{(b-1)^2}$$
$$1 + C + C^4 + C^9 + \text{etc.} + C^{(c-1)^2}$$

Each of these factors can be determined from what we showed in Articles 20, 21, 23, so the value of the product will also become known. It will be useful to collect the rules for determining these factors in a single overview here. Since the root $A$ is $= \frac{kn}{a} \cdot \frac{360^0}{a}$, the sum $1 + A + A^4 + A^9 + \text{etc.} + A^{(a-1)^2}$, which we shall denote by $L$, will be determined by the number $\frac{kn}{a}$ in the same way as $W$ is determined by $k$ in our general investigation. Twelve cases must now be distinguished.

I. If $a$ is a prime number of the form $4\mu + 1$, say $= p$, or a power of such a prime number with an odd exponent, and $\frac{kn}{a}$ is a quadratic residue of $p$, then $L = +\sqrt{a}$.

II. If the other conditions remain unchanged and $\frac{kn}{a}$ is a quadratic nonresidue of $p$, then $L = -\sqrt{a}$.

III. If $a$ is a prime number of the form $4\mu + 3$, say $= p$, or a power of such a prime number with an odd exponent, and $\frac{kn}{a}$ is a quadratic residue of $p$, then $L = +i\sqrt{a}$.

IV. If, with the other conditions as in III, $\frac{kn}{a}$ is a quadratic nonresidue of $p$, then $L = -i\sqrt{a}$.

V. If $a$ is the square, or a higher power with an even exponent, of an (odd) prime number, then $L = +\sqrt{a}$.

VI. If $a = 2$, then $L = 0$.

VII. If $a = 4$, or it is a higher power of two with an even exponent, and $\frac{kn}{a}$ is of the form $4\mu + 1$, then $L = (1 + i)\sqrt{a}$.

VIII. If, with the other conditions as in VII, $\frac{kn}{a}$ is of the form $4\mu + 3$, then $L = (1 - i)\sqrt{a}$.

IX. If $a = 8$, or it is a higher power of two with an odd exponent, and $\frac{kn}{a}$ is of the form $8\mu + 1$, then $L = (1 + i)\sqrt{a}$.

X. If, with the other conditions as in IX, $\frac{kn}{a}$ is of the form $8\mu + 3$, then $L = (-1 + i)\sqrt{a}$.

XI. If the other conditions remain unchanged and $\frac{kn}{a}$ is of the form $8\mu + 5$, then $L = (-1 - i)\sqrt{a}$.

XII. If the other conditions remain unchanged and $\frac{kn}{a}$ is of the form $8\mu+7$, then $L = (1-i)\sqrt{a}$.

**29.**

For example, let $n = 2520 = 8 \,.\, 9 \,.\, 5 \,.\, 7$ and $k = 13$. Here we obtain

for $a = 8$, by case XII, $L = (1-i)\sqrt{8}$
for the factor 9, by case V, the corresponding sum is $= \sqrt{9}$
for the factor 5, by case II, the corresponding sum is $= -\sqrt{5}$
for the factor 7, by case III, the corresponding sum is $= +i\sqrt{7}$.

Thus $W = (1-i) \,.\, (-i) \,.\, \sqrt{2520} = (-1-i)\sqrt{2520}$.
For the same value of $n$, let $k = 1$; then the corresponding sums are

for the factor 8 the sum $(-1+i)\sqrt{8}$
for the factor 9 the sum $\sqrt{9}$
for the factor 5 the sum $\sqrt{5}$
for the factor 7 the sum $-i\sqrt{7}$

This gives the product $W = (1+i)\sqrt{2520}$.

**30.**

Another method for determining the sum $W$ in general follows from what we set out in Articles 22. 24. Put $\cos\omega + i\sin\omega = \rho$, and

$$\rho^{\frac{nn}{aa}} = \alpha, \quad \rho^{\frac{nn}{bb}} = \beta, \quad \rho^{\frac{nn}{cc}} = \gamma \text{ etc.}$$

so that $r = \rho^k$, $A = \alpha^k$, $B = \beta^k$, $C = \gamma^k$ etc. Then

$$1 + \rho + \rho^4 + \rho^9 + \text{etc.} + \rho^{(n-1)^2}$$

is the product of the factors

$$1 + \alpha + \alpha^4 + \alpha^9 + \text{etc.} + \alpha^{(a-1)^2}$$
$$1 + \beta + \beta^4 + \beta^9 + \text{etc.} + \beta^{(b-1)^2}$$
$$1 + \gamma + \gamma^4 + \gamma^9 + \text{etc.} + \gamma^{(c-1)^2}, \text{ etc.}$$

and therefore $W$ is the product of the factors

$$w = 1 + \rho + \rho^4 + \rho^9 + \text{etc.} + \rho^{(n-1)^2}$$

$$\mathfrak{A} = \frac{1 + A + A^4 + A^9 + \text{etc.} + A^{(a-1)^2}}{1 + \alpha + \alpha^4 + \alpha^9 + \text{etc.} + \alpha^{(a-1)^2}}$$

$$\mathfrak{B} = \frac{1 + B + B^4 + B^9 + \text{etc.} + B^{(b-1)^2}}{1 + \beta + \beta^4 + \beta^9 + \text{etc.} + \beta^{(b-1)^2}}$$

$$\mathfrak{C} = \frac{1 + C + C^4 + C^9 + \text{etc.} + C^{(c-1)^2}}{1 + \gamma + \gamma^4 + \gamma^9 + \text{etc.} + \gamma^{(c-1)^2}}$$

The first factor $w$ has already been determined by the investigations given above (Article 19); the remaining factors $\mathfrak{A}, \mathfrak{B}, \mathfrak{C}$ etc. follow from the formulas of Articles 22. 24, which we collect here again to have everything together[*]. Twelve cases must be distinguished here, namely

I. If $a$ is an (odd) prime number $= p$, or a power of such a number with an odd exponent, and $k$ is a quadratic residue of $p$, the corresponding factor is $\mathfrak{A} = +1$.

II. If the other conditions remain unchanged and $k$ is a quadratic non-residue of $p$, then $\mathfrak{A} = -1$.

III. If $a$ is the square of an odd prime number, or a higher power of it with an even exponent, then $\mathfrak{A} = +1$.

IV. If $a$ is $= 4$, or a higher power of two with an even exponent, and $k$ is of the form $4\mu + 1$, then $\mathfrak{A} = +1$.

V. If, with the other conditions as in IV, $k$ is of the form $4\mu + 3$ and $\frac{n}{a}$ of the form $4\mu + 1$, then $\mathfrak{A} = -i$.

VI. If, with the other conditions as in IV, $k$ is of the form $4\mu + 3$ and $\frac{n}{a}$ of the form $4\mu + 3$, then $\mathfrak{A} = +i$.

VII. If $a$ is $= 8$, or a higher power of two with an odd exponent, and $k$ is of the form $8\mu + 1$, then $\mathfrak{A} = +1$.

VIII. If, with the other conditions as in VII, $k$ is of the form $8\mu + 5$, then $\mathfrak{A} = -1$.

IX. If, with the other conditions as in VII, $k$ is of the form $8\mu + 3$ and $\frac{n}{a}$ of the form $4\mu + 1$, then $\mathfrak{A} = +i$.

X. If, with the other conditions as in VII, $k$ is of the form $8\mu + 3$ and $\frac{n}{a}$ of the form $4\mu + 3$, then $\mathfrak{A} = -i$.

XI. If, with the other conditions as in VII, $k$ is of the form $8\mu + 7$ and $\frac{n}{a}$ of the form $4\mu + 1$, then $\mathfrak{A} = -i$.

XII. If, with the other conditions as in VII, $k$ is of the form $8\mu + 7$ and $\frac{n}{a}$ of the form $4\mu + 3$, then $\mathfrak{A} = +i$.

[*] Clearly, what were $k$ and $h$ there are here $\frac{n}{a}$ and $k$ for the second factor, $\frac{n}{b}$ and $k$ for the third factor, etc.

We omit the case where $a = 2$; here $\mathfrak{A}$ would indeed be $= \frac{0}{0}$, or indeterminate, but then we always have $W = 0$.

The remaining factors $\mathfrak{B}, \mathfrak{C}$ etc. depend on $b, c$ etc. in the same way as $\mathfrak{A}$ depends on $a$, insofar as these enter into their determination.

**31.**

By this second method, the first example of Article 29 runs as follows:

The factor $w$ is $= (1+i)\sqrt{2520}$
For $a = 8$ the corresponding factor $\mathfrak{A}$ is, by case VIII, $= -1$
For the second factor of $n$, 9, the corresponding factor is $+1$ (by case III.)
For the factor 5 the corresponding factor is $-1$ (by case II.)
For the factor 7 the corresponding factor is $-1$ (by case II.)

This gives the product $W = (-1-i)\sqrt{2520}$, as in Article 29.

**32.**

Since the value of $W$ can be determined by *two* methods, one of which rests on the relations of the numbers $\frac{nk}{a}$, $\frac{nk}{b}$, $\frac{nk}{c}$ etc. to the numbers $a, b, c$ etc., while the other depends on the relations of $k$ to the numbers $a, b, c$ etc., there must be a conditional connection among all these relations, so that any one can be determined from the others. Suppose that all the numbers $a, b, c$ etc. are odd prime numbers, and take $k = 1$. Divide the factors $a, b, c$ etc. into two classes: one contains those of the form $4\mu + 1$, denoted by $p, p', p''$ etc.; the other consists of those of the form $4\mu+3$, denoted by $q, q', q''$ etc. We shall denote the number of the latter by $m$. With this done, we first observe that $n$ is of the form $4\mu + 1$ if $m$ is even (this also includes the case where there are no factors at all in the second class, or $m = 0$), whereas $n$ is of the form $4\mu+3$ if $m$ is odd. Now the determination of $W$ by the first method is carried out as follows. Let the numbers $P, P'\ P''$ etc., $Q, Q', Q''$ etc. depend on the relations of the numbers $\frac{n}{p}$, $\frac{n}{p'}$, $\frac{n}{p''}$ etc., $\frac{n}{q}$, $\frac{n}{q'}$, $\frac{n}{q''}$ etc. to the numbers $p, p', p''$ etc., $q, q', q''$ etc., respectively, in such a way that we put

$$P = +1, \text{ if } \frac{n}{p} \text{ is a quadratic residue of } p$$
$$P = -1, \text{ if } \frac{n}{p} \text{ is a quadratic nonresidue of } p$$

and similarly for the others. Then $W$ will be the product of the factors $P\sqrt{p}$, $P'\sqrt{p'}$, $P''\sqrt{p''}$ etc. $iQ\sqrt{q}$, $iQ'\sqrt{q'}$, $iQ''\sqrt{q''}$ etc., and therefore

$$W = PP'P'' \ldots QQ'Q'' \ldots i^m\sqrt{n}$$

By the second method, or rather directly from the rules of Article 19, we have

$$W = +\sqrt{n}, \text{ if } n \text{ is of the form } 4\mu + 1,$$
$$\text{or equivalently, if } m \text{ is even}$$
$$W = +i\sqrt{n}, \text{ if } n \text{ is of the form } 4\mu + 3,$$
$$\text{or if } m \text{ is odd.}$$

Both cases can be combined in the following formula:

$$W = i^{mm}\sqrt{n}$$

It therefore follows that

$$PP'P'' \ldots QQ'Q'' \ldots = i^{mm-m}$$

But $i^{mm-m}$ is $= 1$ whenever $m$ is of the form $4\mu$ or $4\mu + 1$, and $= -1$ whenever $m$ is of the form $4\mu+2$ or $4\mu+3$. From this we deduce the following most elegant

THEOREM. *Let $a, b, c$ etc. denote distinct positive odd prime numbers, and put their product $= n$. Let $m$ of them be of the form $4\mu + 3$, and the rest of the form $4\mu + 1$. The number of those among $a, b, c$ etc. for which $\frac{n}{a}$, $\frac{n}{b}$, $\frac{n}{c}$ etc., respectively, are nonresidues will be even whenever $m$ is of the form $4\mu$ or $4\mu + 1$, but odd whenever $m$ is of the form $4\mu + 2$ or $4\mu + 3$.*

Thus, for example, putting $a = 3$, $b = 5$, $c = 7$, $d = 11$, we have three numbers of the form $4\mu + 3$, namely 3, 7 and 11; and $5 \,.\, 7 \,.\, 11\, R3$; $3 \,.\, 7 \,.\, 11\, R5$; $3 \,.\, 5 \,.\, 11\, R7$; $3 \,.\, 5 \,.\, 7\, N11$, so that only $\frac{n}{d}$ is a nonresidue of $d$.

**33.**

The celebrated *fundamental theorem* about quadratic residues is nothing other than a special case of the theorem just developed. Specifically, by limiting the number of the numbers $a, b, c$ etc. to *two*, it is clear that, if just one of them, or neither, is of the form $4\mu + 3$, we must have either both $aRb$, $bRa$, or both $aNb$, $bNa$; but if both are of the form $4\mu + 3$, one must be a nonresidue of the other, and the latter a residue of the former. Here, then, is a *fourth* proof of this most important theorem, whose first and second proofs we gave in the Disquisitiones Arithmeticae, and the third recently in a separate paper (*Commentt. T. XVI*); we shall later set out two others, again resting on entirely different principles. It is certainly most remarkable that this most beautiful theorem, which at first so stubbornly eluded every attempt, could later be reached by so many paths, worlds apart from one another.

**34.**

The remaining theorems, which form a kind of supplement to the fundamental theorem, namely those by which we distinguish the prime numbers for which $-1, +2$ and $-2$ are residues or nonresidues, can also be derived from the same principles. We shall begin with the residue $+2$.

Putting $n = 8a$, with $a$ a prime number, and $k = 1$, the method of Article 28 makes $W$ the product of two factors. One of them will be $+\sqrt{a}$, or $+i\sqrt{a}$, if 8, or equivalently 2, is a quadratic residue of $a$; but $-\sqrt{a}$ or $-i\sqrt{a}$ if 2 is a nonresidue of $a$. The second factor is

$$\begin{aligned}
&(1+i)\sqrt{8}, \quad \text{if } a \text{ is of the form } 8\mu+1\\
&(-1+i)\sqrt{8}, \quad \text{if } a \text{ is of the form } 8\mu+3\\
&(-1-i)\sqrt{8}, \quad \text{if } a \text{ is of the form } 8\mu+5\\
&(1-i)\sqrt{8}, \quad \text{if } a \text{ is of the form } 8\mu+7
\end{aligned}$$

But by Article 18 we always have $W = (1+i)\sqrt{n}$; dividing this value by the four values of the second factor shows that the first factor must be

$$\begin{aligned}
&+\sqrt{a}, \quad \text{if } a \text{ is of the form } 8\mu+1\\
&-i\sqrt{a}, \quad \text{if } a \text{ is of the form } 8\mu+3\\
&-\sqrt{a}, \quad \text{if } a \text{ is of the form } 8\mu+5\\
&+i\sqrt{a}, \quad \text{if } a \text{ is of the form } 8\mu+7
\end{aligned}$$

It follows immediately that 2 must be a residue of $a$ in the first and fourth cases, but a nonresidue in the second and third.

**35.**

The prime numbers for which $-1$ is a residue or a nonresidue are easily distinguished with the aid of the following theorem, which is also quite remarkable in itself.

THEOREM. *The product of the two factors*

$$W' = 1 + r^{-1} + r^{-4} + \text{etc.} + r^{-(n-1)^2}$$
$$W = 1 + r + r^4 + \text{etc.} + r^{(n-1)^2}$$

*is* $= n$ *if* $n$ *is odd; or* $= 0$ *if* $n$ *is twice an odd number; or* $= 2n$ *if* $n$ *is divisible by four.*

*Proof.* Since we evidently have

$$\begin{aligned} W &= r + r^4 + r^9 + \text{etc.} + r^{nn} \\ &= r^4 + r^9 + \text{etc.} + r^{(n+1)^2} \\ &= r^9 + \text{etc.} + r^{(n+2)^2} \quad \text{etc.} \end{aligned}$$

the product $WW'$ can also be written as follows:

$$\begin{aligned} & 1 + r + r^4 + r^9 + \text{etc.} + r^{(n-1)^2} \\ & + r^{-1}(r + r^4 + r^9 + r^{16} + \text{etc.} + r^{nn}) \\ & + r^{-4}(r^4 + r^9 + r^{16} + r^{25} + \text{etc.} + r^{(n+1)^2}) \\ & + r^{-9}(r^9 + r^{16} + r^{25} + r^{36} + \text{etc.} + r^{(n+2)^2}) \\ & \text{etc.} \\ & + r^{-(n-1)^2}(r^{(n-1)^2} + r^{nn} + r^{(n+1)^2} \\ & \qquad + r^{(n+2)^2} + \text{etc.} + r^{(2n-2)^2}) \end{aligned}$$

Summing this expression vertically gives

$$\begin{aligned} & n \\ & + r(1 + rr + r^4 + r^6 + \text{etc.} + r^{2n-2}) \\ & + r^4(1 + r^4 + r^8 + r^{12} + \text{etc.} + r^{4n-4}) \\ & + r^9(1 + r^6 + r^{12} + r^{18} + \text{etc.} + r^{6n-6}) \\ & + \text{etc.} \\ & + r^{(n-1)^2}(1 + r^{2n-2} + r^{4n-4} \\ & \qquad + r^{6n-6} + \text{etc.} + r^{2(n-1)^2}) \end{aligned}$$

Now, if $n$ is odd, each part of this expression, except the first, $n$, will be $= 0$; for the second is evidently $\frac{r(1-r^{2n})}{1-rr}$, the third $\frac{r^4(1-r^{4n})}{1-r^4}$ etc. Whenever $n$ is even, however, we must also except the part

$$r^{\frac{1}{4}nn}(1 + r^n + r^{2n} + r^{3n} + \text{etc.} + r^{nn-n})$$

which is $= nr^{\frac{1}{4}nn}$. In the first case, therefore, $WW' = n$, but in the second $= n + nr^{\frac{1}{4}nn}$; now $r^{\frac{1}{4}nn}$ is $= +1$ if $n$ is divisible by four, and then we obtain $WW' = 2n$; on the other hand, $r^{\frac{1}{4}nn} = -1$ if $n$ is twice an odd number, in which case $WW' = 0$. Q. E. D.

**36.**

Article 22 establishes that, if $n$ is an odd prime number, $\frac{W'}{W}$ is $= +1$ or $= -1$ according as $-1$ is a residue or a nonresidue of $n$. Hence in the first case we must have $W^2 = +n$, and in the second $W^2 = -n$; by Article 13 we therefore conclude that the first case can occur only when $n$ is of the form $4\mu + 1$, and the second when $n$ is of the form $4\mu + 3$.

Finally, combining the conditions found for the residues $+2$ and $-1$ immediately shows that $-2$ is a residue of every prime number of the form $8\mu + 1$ or $8\mu + 3$, and a nonresidue of every prime number of the form $8\mu + 5$ or $8\mu + 7$.

---

# NEW PROOFS AND EXTENSIONS OF THE FUNDAMENTAL THEOREM IN THE THEORY OF QUADRATIC RESIDUES

BY

CARL FRIEDRICH GAUSS

PRESENTED TO THE ROYAL SOCIETY OF SCIENCES ON FEBRUARY 10, 1817.

---

Commentationes societatis regiae scientiarum Gottingensis recentiores. Vol. IV.
Gottingae MDCCCXVIII.

---

# NEW PROOFS AND EXTENSIONS OF THE FUNDAMENTAL THEOREM IN THE THEORY OF QUADRATIC RESIDUES.

---

The fundamental theorem about quadratic residues, which ranks among the most beautiful truths of higher arithmetic, was readily discovered by induction, but proved with far greater difficulty. It often happens in this field that the proofs of the simplest truths, which seem to offer themselves to the investigator through induction of their own accord, lie deeply hidden and can finally be brought to light only after many unsuccessful attempts, perhaps by a path very different from the one along which they were sought. Then it often happens that, once one path has been found, *several* others gradually open up, leading to the same goal: some more briefly and directly, others as if by a roundabout way, starting from very different principles between which and the question at hand one would scarcely have suspected any connection. This remarkable connection among more hidden truths not only gives these investigations a particular charm; it also deserves careful study and unravelling, since it often yields new resources or advances for the science itself.

Although the arithmetical theorem considered here might therefore seem, through the earlier efforts that supplied four entirely different proofs[*)], to have been fully dealt with, I nevertheless return to the same subject and add two further proofs, which will certainly shed new light on it. The first is related to the third in a certain respect, since it starts from the same lemma; it then follows a different path, however, so that it can rightly be regarded as a new proof, which in elegance will seem, if not superior to that third proof, at least by no means inferior. The sixth proof rests on an entirely different and subtler principle, and gives a new example of the remarkable connection between arithmetical truths that at first sight seem very far apart. These two proofs are accompanied by a very simple new algorithm for deciding whether a given integer is a quadratic residue or a nonresidue of a given prime number.

[*)] Two are given in the fourth and fifth sections of the *Disquisitiones Arithmeticae*; the third in a separate paper (*Commentt. Soc. Gotting. Vol. XVI*), and the fourth in the paper *Summatio quarundam serierum singularium* (*Commentt. Recentiores, Vol. I*).

There was another reason that led me to publish now, in particular, the new proofs promised nine years ago. When I began in 1805 to investigate the theory of cubic and biquadratic residues, a far more difficult subject, I experienced much the same course of events as before in the theory of quadratic residues. The theorems that completely settle these questions, and display a remarkable analogy with the theorems about quadratic residues, were immediately discovered by induction as soon as they were sought along a suitable path; but all attempts to obtain proofs complete in every respect remained unsuccessful for a long time. This was precisely what prompted me to strive so eagerly to add more and more proofs to those already known for quadratic residues, sustained by the hope that, among many different methods, one or another might help to illuminate the related subject. This hope was by no means in vain, and tireless work was finally followed by good results. Soon it will be possible to bring the fruits of my nights of work before the public; but before I undertake this arduous task, I have resolved to return once more to the theory of quadratic residues, complete everything that still remains to be done in it, and thus, as it were, take leave of this part of higher arithmetic.

## *FIFTH PROOF OF THE FUNDAMENTAL THEOREM IN THE THEORY OF QUADRATIC RESIDUES.*

### 1.

We have already stated in the introduction that the fifth and third proofs start from the same lemma. For convenience, we have thought it appropriate to repeat it here in notation adapted to the present investigation.

LEMMA. *Let $m$ be a prime number (positive and odd), and $M$ an integer not divisible by $m$. Take the least positive residues of the numbers*

$$M,\ 2M,\ 3M,\ 4M\ \ldots\ldots\ \tfrac{1}{2}(m-1)M$$

*modulo $m$. Some will be less than $\frac{1}{2}m$, and others greater; let the number of the latter be $= n$. Then $M$ is a quadratic residue or a nonresidue of $m$, according as $n$ is even or odd.*

PROOF. Let those of these residues which are less than $\frac{1}{2}m$ be $a, b, c, d$ etc., and the remaining ones, greater than $\frac{1}{2}m$, be $a'$, $b'$, $c'$, $d'$ etc. The complements of the latter to $m$, namely $m-a'$, $m-b'$, $m-c'$, $m-d'$ etc., are evidently all less than $\frac{1}{2}m$ and distinct both from one another and from the residues $a$, $b$, $c$, $d$ etc. Thus, taken together with these, they are precisely all the numbers $1,\ 2,\ 3,\ 4\ldots.\frac{1}{2}(m-1)$, though in a different order. Putting the product

$$1\,.\,2\,.\,3\,.\,4\ldots..\tfrac{1}{2}(m-1) = P$$

we therefore have

$$P = abcd\ldots. \times (m-a')(m-b')(m-c')(m-d')\ldots.$$

and hence

$$(-1)^n P = abcd\ldots. \times (a'-m)(b'-m)(c'-m)(d'-m)\ldots.$$

Furthermore, modulo $m$, we have

$$\begin{aligned} PM^{\frac{1}{2}(m-1)} &\equiv abcd\,..\times a'b'c'd'\,..\\ &\equiv abcd\,..\times (a'-m)(b'-m)(c'-m)(d'-m)\,.. \end{aligned}$$

and hence

$$PM^{\frac{1}{2}(m-1)} \equiv P(-1)^n$$

It follows that $M^{\frac{1}{2}(m-1)} \equiv \pm 1$, taking the upper or lower sign according as $n$ is even or odd. With the aid of the theorem proved in Article 106 of the *Disquisitiones Arithmeticae*, the lemma follows immediately.

**2.**

THEOREM. *Let $m, M$ be positive odd integers coprime to one another, and let $n$ be the number of those among the least positive residues of the numbers*

$$M,\ 2M,\ 3M \ldots\ldots \tfrac{1}{2}(m-1)M$$

*modulo $m$ which are greater than $\frac{1}{2}m$. Likewise, let $N$ be the number of those among the least positive residues of the numbers*

$$m,\ 2m,\ 3m \ldots\ldots \tfrac{1}{2}(M-1)m$$

*modulo $M$ which are greater than $\frac{1}{2}M$. Then the three numbers $n, N$, $\frac{1}{4}(m-1)(M-1)$ are either all even, or one is even and the other two odd.*

PROOF. Let us denote

by $f$ the collection of numbers $1,\ 2,\ 3 \ldots\ldots \frac{1}{2}(m-1)$.
by $f'$ the collection of numbers $m-1,\ m-2,\ m-3 \ldots\ldots \frac{1}{2}(m+1)$
by $F$ the collection of numbers $1,\ 2,\ 3 \ldots\ldots \frac{1}{2}(M-1)$
by $F'$ the collection of numbers $M-1,\ M-2,\ M-3 \ldots\ldots \frac{1}{2}(M+1)$

Thus $n$ indicates how many numbers $Mf$ have their least positive residues modulo $m$ in the collection $f'$. Likewise, $N$ indicates how many numbers $mF$ have their least positive residues modulo $M$ in the collection $F'$. Finally, let

$\varphi$ denote the collection of numbers $1,\ 2,\ 3 \ldots\ldots \frac{1}{2}(mM-1)$
$\varphi'$ denote the collection of numbers $mM-1,\ mM-2,\ mM-3 \ldots\ldots \frac{1}{2}(mM+1)$

Every integer not divisible by $m$ must be congruent modulo $m$ either to a residue from $f$ or to one from $f'$. Likewise, every integer not divisible by $M$ is congruent modulo $M$ either to a residue from $F$ or to one from $F'$. Thus all the numbers in $\varphi$, among which evidently none is divisible by both $m$ and $M$, can be divided into eight classes as follows.

I. The first class contains the numbers congruent modulo $m$ to a number from $f$, and modulo $M$ to a number from $F$. We shall denote their number by $\alpha$.

II. The numbers congruent modulo $m, M$, respectively, to numbers from $f, F'$; we put their number $= \beta$.

III. The numbers congruent modulo $m, M$, respectively, to numbers from $f', F$; we put their number $= \gamma$.

IV. The numbers congruent modulo $m, M$, respectively, to numbers from $f', F'$; let their number be $= \delta$.

V. The numbers divisible by $m$, but congruent modulo $M$ to residues from $F$.

VI. The numbers divisible by $m$, but congruent modulo $M$ to residues from $F'$.

VII. The numbers divisible by $M$, but congruent modulo $m$ to residues from $f$.

VIII. The numbers divisible by $M$, but congruent modulo $m$ to residues from $f'$.

Evidently classes V and VI together contain all the numbers $mF$. The number contained in VI is $= N$, so the number contained in V is $\frac{1}{2}(M-1)-N$. Likewise, classes VII and VIII together contain all the numbers $Mf$; class VIII contains $n$ numbers, and class VII contains $\frac{1}{2}(m-1)-n$.

In just the same way, all the numbers in $\varphi'$ are divided into eight classes IX . . XVI. If we keep the same order in doing this, it is easy to see that the numbers in the classes

$$\text{IX, X, XI, XII, XIII, XIV, XV, XVI}$$

are, respectively, the complements to $mM$ of the numbers in the classes

$$\text{IV, III, II, I, VI, V, VIII, VII}$$

so that class IX contains $\delta$ numbers, class X contains $\gamma$, and so on. It is now clear that, if all the numbers of the first class are combined with all those of the ninth, we obtain all the numbers below $mM$ which are congruent modulo $m$ to a number from $f$, and modulo $M$ to a number from $F$. It is easy to see that their number equals the number of all combinations of each individual $f$ with each individual $F$. We therefore have

$$\alpha + \delta = \tfrac{1}{4}(m-1)(M-1)$$

and by a similar argument we also have

$$\beta + \gamma = \tfrac{1}{4}(m-1)(M-1)$$

Combining all the numbers of classes II, IV and VI evidently gives all the numbers below $\frac{1}{2}mM$ which are congruent to a residue from $F'$ modulo $M$. These same numbers can also be written as follows:

$$F',\ M+F',\ 2M+F',\ 3M+F' \ldots .\ \tfrac{1}{2}(m-3)M+F'$$

Their total number is therefore $= \frac{1}{4}(m-1)(M-1)$, or we have

$$\beta + \delta + N = \tfrac{1}{4}(m-1)(M-1)$$

Likewise, combining all of classes III, IV and VIII gives

$$\gamma + \delta + n = \tfrac{1}{4}(m-1)(M-1)$$

These four equations give the following:

$$2\alpha = \tfrac{1}{4}(m-1)(M-1) + n + N$$
$$2\beta = \tfrac{1}{4}(m-1)(M-1) + n - N$$
$$2\gamma = \tfrac{1}{4}(m-1)(M-1) - n + N$$
$$2\delta = \tfrac{1}{4}(m-1)(M-1) - n - N$$

each of which establishes the theorem.

**3.**

If we now suppose that $m$ and $M$ are prime numbers, the fundamental theorem follows immediately by combining the preceding theorem with the lemma of Article 1. For it is clear that

I. whenever the numbers $m, M$ both have the form $4k+1$, or just one of them does, the number $\frac{1}{4}(m-1)(M-1)$ is even. Thus $n$ and $N$ are either both even or both odd, and consequently either each of $m$ and $M$ is a quadratic residue of the other, or each is a quadratic nonresidue of the other.

II. Whenever both numbers $m, M$ are of the form $4k+3$, however, $\frac{1}{4}(m-1)(M-1)$ is odd. Hence one of the numbers $n, N$ is even and the other odd, and consequently one of the numbers $m, M$ is a quadratic residue of the other, and the other a quadratic nonresidue of the first. Q. E. D.

## *SIXTH PROOF OF THE FUNDAMENTAL THEOREM IN THE THEORY OF QUADRATIC RESIDUES.*

### 1.

THEOREM. *Let $p$ denote a prime number (positive and odd), $n$ a positive integer not divisible by $p$, and $x$ an indeterminate quantity. Then the function*

$$1 + x^n + x^{2n} + x^{3n} + \text{etc.} + x^{np-n}$$

*is divisible by*

$$1 + x + xx + x^3 + \text{etc.} + x^{p-1}$$

PROOF. Take a positive integer $g$ such that $gn \equiv 1 \pmod{p}$, and put $gn = 1 + hp$. Then we have

$$\begin{aligned}\frac{1 + x^n + x^{2n} + x^{3n} + \text{etc.}\, x^{np-n}}{1 + x + xx + x^3 + \text{etc.}\, x^{p-1}} &= \frac{(1 - x^{np})(1 - x)}{(1 - x^n)(1 - x^p)} \\ &= \frac{(1 - x^{np})(1 - x^{gn} - x + x^{hp+1})}{(1 - x^n)(1 - x^p)} \\ &= \frac{1 - x^{np}}{1 - x^p} \cdot \frac{1 - x^{gn}}{1 - x^n} - \frac{x(1 - x^{np})}{1 - x^n} \cdot \frac{1 - x^{hp}}{1 - x^p}\end{aligned}$$

which is evidently a polynomial. Q. E. D.

Thus every polynomial in $x$ divisible by $\frac{1-x^{np}}{1-x^n}$ is also divisible by $\frac{1-x^p}{1-x}$.

### 2.

Let $\alpha$ denote a positive primitive root modulo $p$; that is, let $\alpha$ be a positive integer such that the least positive residues of the powers $1, \alpha, \alpha\alpha, \alpha^3 \ldots\ldots \alpha^{p-2}$ modulo $p$ are precisely the numbers $1, 2, 3, 4 \ldots\ldots p-1$, without regard to order. Further, denoting by $fx$ the function

$$x + x^\alpha + x^{\alpha\alpha} + x^{\alpha^3} + \text{etc.} + x^{\alpha^{p-2}} + 1$$

it is clear that $fx - 1 - x - xx - x^3 - \text{etc.} - x^{p-1}$ is divisible by $1 - x^p$, and therefore all the more by $\frac{1-x^p}{1-x} = 1 + x + xx + x^3 + \text{etc.} + x^{p-1}$. This last function therefore also divides $fx$ itself. Since $x$ represents an indeterminate quantity, it follows that $f(x^n)$ is also divisible by $\frac{1-x^{np}}{1-x^n}$, and hence, by the preceding article, by $\frac{1-x^p}{1-x}$, whenever $n$ is an integer not divisible by $p$. If, on the other hand, $n$ is an integer divisible by $p$, the individual terms of the function $f(x^n)$, each decreased by one, are divisible by $1 - x^p$. In this case, therefore, $f(x^n) - p$ is also divisible by $1 - x^p$, and hence by $\frac{1-x^p}{1-x}$.

**3.**

THEOREM. *Putting*

$$x - x^{\alpha} + x^{\alpha\alpha} - x^{\alpha^3} + x^{\alpha^4} - \text{etc.} - x^{\alpha^{p-2}} = \xi$$

*then* $\xi\xi \mp p$ *is divisible by* $\frac{1-x^p}{1-x}$, *taking the upper sign whenever* $p$ *is of the form* $4k+1$, *and the lower whenever* $p$ *is of the form* $4k+3$.

PROOF. It is easy to see that, of the following $p-1$ functions

$$\begin{aligned}
&+ x\xi - xx + x^{\alpha+1} - x^{\alpha\alpha+1} \\
&\qquad + \text{etc.} + x^{\alpha^{p-2}+1} \\
&- x^{\alpha}\xi - x^{2\alpha} + x^{\alpha\alpha+\alpha} - x^{\alpha^3+\alpha} \\
&\qquad + \text{etc.} + x^{\alpha^{p-1}+\alpha} \\
&+ x^{\alpha\alpha}\xi - x^{2\alpha\alpha} + x^{\alpha^3+\alpha\alpha} - x^{\alpha^4+\alpha\alpha} \\
&\qquad + \text{etc.} + x^{\alpha^p+\alpha\alpha} \\
&- x^{\alpha^3}\xi - x^{2\alpha^3} + x^{\alpha^4+\alpha^3} - x^{\alpha^5+\alpha^3} \\
&\qquad + \text{etc.} + x^{\alpha^{p+1}+\alpha^3}
\end{aligned}$$

etc., as far as

$$\begin{aligned}
&- x^{\alpha^{p-2}}\xi - x^{2\alpha^{p-2}} + x^{\alpha^{p-1}+\alpha^{p-2}} - x^{\alpha^p+\alpha^{p-2}} \\
&\qquad\qquad + \text{etc.} + x^{\alpha^{2p-4}+\alpha^{p-2}}
\end{aligned}$$

the first is $= 0$, while each of the others is divisible by $1 - x^p$. Their sum is therefore also divisible by $1 - x^p$, and is found to be

$$\begin{aligned}
&= \xi\xi - (f(xx) - 1) + (f(x^{\alpha+1}) - 1) - (f(x^{\alpha\alpha+1}) - 1) \\
&\qquad + (f(x^{\alpha^3+1}) - 1) - \text{etc.} \\
&\qquad + (f(x^{\alpha^{p-2}+1}) - 1) \\
&= \xi\xi - f(xx) + f(x^{\alpha+1}) - f(x^{\alpha\alpha+1}) \\
&\qquad + f(x^{\alpha^3+1}) - \text{etc.} + f(x^{\alpha^{p-2}+1}) = \Omega
\end{aligned}$$

Thus this expression $\Omega$ is also divisible by $\frac{1-x^p}{1-x}$. Now, among the exponents $2,\ \alpha+1,\ \alpha\alpha+1,\ \alpha^3+1 \ldots\ldots \alpha^{p-2}+1$, only one is divisible by $p$, namely $\alpha^{\frac{1}{2}(p-1)}+1$. Hence, by the preceding article, the following individual parts of the expression $\Omega$

$$f(xx), \quad f(x^{\alpha+1}), \quad f(x^{\alpha\alpha+1}), \quad (fx^{\alpha^3+1}) \quad \text{etc.}$$

with the sole exception of the term $f(x^{\alpha^{\frac{1}{2}(p-1)}+1})$, are divisible by $\frac{1-x^p}{1-x}$. Those parts may therefore be removed, leaving the following function still divisible by $\frac{1-x^p}{1-x}$:

$$\xi\xi \mp f(x^{\alpha^{\frac{1}{2}(p-1)}+1})$$

Here the upper or lower sign applies according as $p$ is of the form $4k+1$ or of the form $4k+3$. Since, moreover, $f(x^{\alpha^{\frac{1}{2}(p-1)}+1})-p$ is divisible by $\frac{1-x^p}{1-x}$, $\xi\xi \mp p$ is also divisible by $\frac{1-x^p}{1-x}$. Q. E. D.

To prevent any ambiguity from the double sign, we shall use $\varepsilon$ to denote the number $+1$ or $-1$, according as $p$ is of the form $4k+1$ or $4k+3$. Thus $\frac{(1-x)(\xi\xi-\varepsilon p)}{1-x^p}$ is a polynomial in $x$, which we shall denote by $Z$.

**4.**

Let $q$ be a positive odd number, so that $\frac{1}{2}(q-1)$ is an integer. Then $(\xi\xi)^{\frac{1}{2}(q-1)} - (\varepsilon p)^{\frac{1}{2}(q-1)}$ is divisible by $\xi\xi - \varepsilon p$, and hence also by $\frac{1-x^p}{1-x}$. Let us put $\varepsilon^{\frac{1}{2}(q-1)} = \delta$, and

$$\xi^{q-1} - \delta p^{\frac{1}{2}(q-1)} = \frac{1-x^p}{1-x} \cdot Y$$

Then $Y$ is a polynomial in $x$, and $\delta = .+1$ whenever one of the numbers $p, q$, or indeed both, is of the form $4k+1$. On the other hand, $\delta = -1$ whenever both $p, q$ are of the form $4k+3$.

**5.**

Let us now suppose that $q$ too is a prime number, different from $p$. It is clear from the theorem proved in Article 51 of the *Disquisitiones Arithmeticae* that

$$. \quad \xi^q - (x^q - x^{q\alpha} + x^{q\alpha\alpha} - x^{q\alpha^3} + \text{etc.} - x^{q\alpha^{p-2}})$$

is divisible by $q$, or is of the form $qX$, where $X$ is a polynomial in $x$ whose numerical coefficients are also integers (as is also to be understood of the other polynomials $Z, Y, W$ occurring here). For the modulus $p$ and primitive root $\alpha$, let us denote the index of the number $q$ by $\mu$; that is, let $q \equiv \alpha^\mu \pmod{p}$. The numbers $q,\ q\alpha,\ q\alpha\alpha,\ q\alpha^3 \ldots\ldots q\alpha^{p-2}$ are therefore congruent modulo $p$, respectively, to the numbers $\alpha^\mu,\ \alpha^{\mu+1},\ \alpha^{\mu+2} \ldots\ldots \alpha^{p-2},\ 1,\ \alpha,\ \alpha\alpha \ldots\ldots \alpha^{\mu-1}$, so that

$$\begin{array}{l}
x^q - x^{\alpha^\mu} \\
x^{q\alpha} - x^{\alpha^{1+1}} \\
x^{q\alpha\alpha} - x^{\alpha^{\mu+2}} \\
x^{q\alpha^3} - x^{\alpha^{\mu+3}} \\
\quad . \\
\quad . \\
\quad . \\
\quad . \\
x^{q\alpha^{p-\mu-2}} - x^{\alpha^{p-2}} \\
x^{q\alpha^{p-\mu-1}} - x \\
x^{q\alpha^{p-\mu}} - x^\alpha \\
x^{q\alpha^{p-\mu+1}} - x^{\alpha\alpha} \\
\quad . \\
\quad . \\
\quad . \\
\quad . \\
x^{q\alpha^{p-2}} - x^{\alpha^{\mu-1}}
\end{array}$$

are divisible by $1 - x^p$. Taking these quantities alternately with positive and negative signs and adding them shows that $1 - x^p$ divides the function

$$x^q - x^{q\alpha} + x^{q\alpha\alpha} - x^{q\alpha^3} + \text{etc.} - x^{q\alpha^{p-2}} \mp \xi$$

with the upper or lower sign applying according as $\mu$ is even or odd; that is, according as $q$ is a quadratic residue or a nonresidue of $p$. We shall therefore put

$$x^q - x^{q\alpha} + x^{q\alpha\alpha} - x^{q\alpha^3} + \text{etc.} - x^{q\alpha^{p-2}} - \gamma\xi = (1 - x^p)W$$

taking $\gamma = +1$ or $\gamma = -1$, according as $q$ is a quadratic residue or a nonresidue of $p$. It is clear that $W$ is a polynomial.

**6.**

With these preparations, we combine the preceding equations to obtain

$$\begin{aligned} q\xi X = \varepsilon p(\delta p^{\frac{1}{2}(q-1)} - \gamma) & \\ & + \frac{1 - x^p}{1 - x} \cdot (Z(\delta p^{\frac{1}{2}(q-1)} - \gamma) \\ & + Y\xi\xi - W\xi(1 - x)) \end{aligned}$$

Let us suppose that dividing the function $\xi X$ by

$$x^{p-1} + x^{p-2} + x^{p-3} + \text{etc.} + x + 1$$

yields the quotient $U$ with remainder $T$, or that we have

$$\xi X = \frac{1 - x^p}{1 - x} \cdot U + T$$

where $U, T$ are polynomials whose numerical coefficients are also integers, and $T$ is certainly of lower degree than the divisor. We therefore have

$$\begin{aligned} qT - \varepsilon p(\delta p^{\frac{1}{2}(q-1)} - \gamma) & \\ & = \frac{1 - x^p}{1 - x} \cdot (Z(\delta p^{\frac{1}{2}(q-1)} - \gamma) \\ & \quad + Y\xi\xi - W\xi(1 - x) - qU) \end{aligned}$$

This equation evidently cannot hold unless the left side and the right side each vanish separately. Thus $\varepsilon p(\delta p^{\frac{1}{2}(q-1)} - \gamma)$ is divisible by $q$, as is $\delta p^{\frac{1}{2}(q-1)} - \gamma$; and therefore, since $\delta\delta = 1$, the number $p^{\frac{1}{2}(q-1)} - \gamma\delta$ is also divisible by $q$.

Now, if $\beta$ denotes unity taken positively or negatively, according as $p$ is a quadratic residue or a quadratic nonresidue of the number $q$, then $p^{\frac{1}{2}(q-1)} - \beta$ is divisible by $q$, and hence so is $\beta - \gamma\delta$. This cannot happen unless $\beta = \gamma\delta$. The fundamental theorem follows immediately. Namely,

I. Whenever the numbers $p, q$ both have the form $4k + 1$, or just one of them does, so that $\delta = +1$, we have $\beta = \gamma$. Consequently either $q$ is a quadratic residue of $p$ and at the same time $p$ a quadratic residue of $q$; or $q$ is a nonresidue of $p$ and at the same time $p$ a nonresidue of $q$.

II. Whenever both $p, q$ are of the form $4k + 3$, so that $\delta = -1$, we have $\beta = -\gamma$. Consequently either $q$ is a quadratic residue of $p$ and at the same time $p$ a nonresidue of $q$; or $q$ is a nonresidue of $p$ and at the same time $p$ a residue of $q$.

Q. E. D.

*A new algorithm for deciding whether a given positive integer is a quadratic residue or a nonresidue of a given positive prime number.*

**1.**

Before presenting a new solution to this problem, we shall briefly recall the solution given in the *Disquisitiones Arithmeticae*. It can be carried out quite readily with the help of the fundamental theorem and the following known theorems:

I. The relation of the number $a$ to the number $b$ (whether the former is a quadratic residue or a nonresidue of the latter) is the same as that of the number $c$ to $b$, if $a \equiv c \pmod{b}$.

II. If $a$ is the product of the factors $\alpha, \beta, \gamma, \delta$ etc., and $b$ is prime, the relation of $a$ to $b$ depends on the relations of these factors to $b$ as follows: $a$ is a quadratic residue of $b$ or a nonresidue according as an even or an odd number of those factors are nonresidues of $b$. Thus a factor that is a square can be disregarded entirely in this examination; if a factor is an integer raised to an odd power, that integer itself can take its place.

III. The number 2 is a quadratic residue of every prime of the form $8m+1$ or $8m+7$, and a nonresidue of every prime of the form $8m+3$ or $8m+5$.

Suppose, then, that a number $a$ is given whose relation to a prime $b$ is sought. If $a$ is greater than $b$, first replace it by its least positive residue modulo $b$. Once this residue has been resolved into prime factors, theorem II reduces the question to finding the relation of each factor to $b$. The relation of the factor 2 (if it occurs once, three times, five times, etc.) is known from theorem III; by the fundamental theorem, the relation of each remaining factor depends on the relation of $b$ itself to that factor. Thus, in place of the single relation of the given number to the prime $b$, we must now investigate several relations of the number $b$ to other odd primes smaller than $b$. These problems are reduced in the same way to smaller moduli, and these successive reductions must evidently come to an end.

**2.**

To illustrate this solution, suppose that the relation of the number 103 to 379 is sought. Since 103 is already less than 379 and is itself prime, the fundamental theorem applies immediately: it tells us that the required relation is opposite to that of the number 379 to 103. This in turn is the same as the relation of the number 70 to 103, which itself depends on the relations of 2, 5, 7 to 103. The first of these is known from theorem III. By the fundamental theorem, the second depends on the relation of the number 103 to 5, which by theorem I is the same as that of 3 to 5; this in turn depends, by the fundamental theorem, on the relation of 5 to 3, which by theorem I is the same as that of 2 to 3, known from theorem III. Likewise, by the

fundamental theorem, the relation of 7 to 103 depends on that of 103 to 7, which by theorem I is the same as that of 5 to 7; this in turn depends, by the fundamental theorem, on the relation of 7 to 5, which by theorem I is the same as that of 2 to 5, known from theorem III. If we now wish to turn this analysis into a synthesis, the decision is reduced to fourteen steps. We give them all here, so that the greater elegance of the new solution may stand out more clearly.

1. The number 2 is a quadratic residue of the number 103 (theorem III).
2. The number 2 is a quadratic nonresidue of the number 3 (theorem III).
3. The number 5 is a quadratic nonresidue of the number 3 (from I and 2).
4. The number 3 is a quadratic nonresidue of the number 5 (the fundamental theorem and 3).
5. The number 103 is a quadratic nonresidue of the number 5 (I and 4).
6. The number 5 is a quadratic nonresidue of the number 103 (the fundamental theorem and 5).
7. The number 2 is a quadratic nonresidue of the number 5 (theorem III).
8. The number 7 is a quadratic nonresidue of the number 5 (I and 7).
9. The number 5 is a quadratic nonresidue of the number 7 (the fundamental theorem and 8).
10. The number 103 is a quadratic nonresidue of the number 7 (I and 9).
11. The number 7 is a quadratic residue of the number 103 (the fundamental theorem and 10).
12. The number 70 is a quadratic nonresidue of the number 103 (II, 1, 6, 11).
13. The number 379 is a quadratic nonresidue of the number 103 (I and 12).
14. The number 103 is a quadratic residue of the number 379 (the fundamental theorem and 13).

For brevity, in what follows we shall use the notation introduced in *Comment. Gotting. Vol. XVI.* By $[x]$ we shall mean the quantity $x$ itself when $x$ is an integer, and the greatest integer below $x$ when $x$ is not an integer. Thus $x - [x]$ is always nonnegative and less than unity.

**3.**

Problem. *Let $a$, $b$ be positive integers coprime to one another, and put $[\frac{1}{2}a] = a'$. Find the sum*

$$\left[\frac{b}{a}\right] + \left[\frac{2b}{a}\right] + \left[\frac{3b}{a}\right] + \left[\frac{4b}{a}\right] + \text{etc.} + \left[\frac{a'b}{a}\right]$$

SOLUTION. For brevity, let us denote such a sum by $\varphi(a, b)$, so that we also have

$$\varphi(b, a) = \left[\frac{a}{b}\right] + \left[\frac{2a}{b}\right] + \left[\frac{3a}{b}\right] + \text{etc.} + \left[\frac{b'a}{b}\right]$$

if we put $[\frac{1}{2}b] = b'$. In the third proof of the fundamental theorem it was shown that, when $a$ and $b$ are odd,

$$\varphi(a, b) + \varphi(b, a) = a'b'$$

By following the same method, the truth of this proposition extends easily to the case in which one of the numbers $a,\ b$ is odd, as we already indicated there. Divide $a$ by $b$, as in the method for finding the greatest common divisor of two integers, and let $\beta$ be the quotient and $c$ the remainder; then divide $b$ by $c$, and so on, obtaining the equations

$$\begin{aligned} a &= \beta b + c \\ b &= \gamma c + d \\ c &= \delta d + e \\ d &= \varepsilon e + f \text{ etc.} \end{aligned}$$

In this way the continually decreasing sequence of numbers $b, c, d, e, f$ etc. will eventually reach unity, since by hypothesis $a$ and $b$ are coprime. The last equation is therefore

$$k = \lambda l + 1$$

Since evidently

$$\begin{aligned} \left[\frac{a}{b}\right] &= \left[\beta + \frac{c}{b}\right] = \beta + \left[\frac{c}{b}\right] \\ \left[\frac{2a}{b}\right] &= \left[2\beta + \frac{2c}{b}\right] = 2\beta + \left[\frac{2c}{b}\right] \\ \left[\frac{3a}{b}\right] &= \left[3\beta + \frac{3c}{b}\right] = 3\beta + \left[\frac{3c}{b}\right] \end{aligned}$$

etc.. we have

$$\varphi(b, a) = \varphi(b, c) + \tfrac{1}{2}\beta(b'b' + b')$$

and consequently

$$\varphi(a, b) = a'b' - \tfrac{1}{2}\beta(b'b' + b') - \varphi(b, c)$$

By similar reasoning, on putting $[\frac{1}{2}c] = c'$, $[\frac{1}{2}d] = d'$ $[\frac{1}{2}e] = e'$ etc., we obtain

$$\begin{aligned}\varphi(b,c) &= b'c' - \tfrac{1}{2}\gamma(c'c' + c') - \varphi(c,d)\\ \varphi(c,d) &= c'd' - \tfrac{1}{2}\delta(d'd' + d') - \varphi(d,e).\\ \varphi(d,e) &= d'e' - \tfrac{1}{2}\varepsilon(e'e' + e') - \varphi(e,f)\end{aligned}$$

etc., as far as

$$\varphi(k,l) = k'l' - \tfrac{1}{2}\lambda(l'l' + l') - \varphi(l,1)$$

Since evidently $\varphi(l,1) = 0$, we obtain from this the formula

$$\begin{aligned}\varphi(a,b) = {} & a'b' - b'c' + c'd' - d'e' + \text{etc.} \pm k'l'\\ & - \tfrac{1}{2}\beta(b'b' + b') + \tfrac{1}{2}\gamma(c'c' + c') - \tfrac{1}{2}\delta(d'd' + d')\\ & + \tfrac{1}{2}\varepsilon(e'e' + e') - \text{etc.} \mp \tfrac{1}{2}\lambda(l'l' + l')\end{aligned}$$

**4.**

It now follows easily from the third proof that the relation of the number $b$ to $a$, whenever $a$ is prime, can be read directly from the value of the sum $\varphi(a, 2b)$. According as this sum is even or odd, $b$ is a quadratic residue of $a$ or a nonresidue. The sum $\varphi(a,b)$ itself can also be used for the same purpose, provided that we distinguish the case in which $b$ is odd from that in which it is even. Namely,

I. When $b$ is odd, $b$ is a quadratic residue or a nonresidue of $a$ according as $\varphi(a,b)$ is even or odd.

II. When $b$ is even, the same rule holds if, in addition, $a$ is of the form $8n+1$ or $8n+7$; but if, for an even value of $b$, the modulus $a$ is of the form $8n+3$ or $8n+5$, the opposite rule must be applied: $b$ is a quadratic residue of $a$ if $\varphi(a,b)$ is odd, and a nonresidue if $\varphi(a,b)$ is even.

All this follows very easily from Article 4 of the third proof.

**5.**

*Example.* To find the relation of the number 103 to the prime 379, we have, for calculating the sum $\varphi(379, 103)$,

$$\begin{array}{r|r|l} a = 379 & a' = 189 & \\ b = 103 & b' = 51 & \beta = 3 \\ c = 70 & c' = 35 & \gamma = 1 \\ d = 33 & d' = 16 & \delta = 2 \\ e = 4 & e' = 2 & \varepsilon = 8 \end{array}$$

and hence

$$\begin{aligned} \varphi(379, 103) &= 9639 - 1785 + 560 - 32 - 3978 + 630 - 272 + 24 \\ &= 4786 \end{aligned}$$

so 103 is a quadratic residue of the number 379. If we prefer to use the sum $(379, 206)$ for the same purpose, we have the following scheme:

$$\begin{array}{r|r|l} 379 & 189 & \\ 206 & 103 & 1 \\ 173 & 86 & 1 \\ 33 & 16 & 5 \\ 8 & 4 & 4 \end{array}$$

from which we obtain

$$\begin{aligned} \varphi(379, 206) &= 19467 - 8858 + 1376 - 64 - 5356 + 3741 - 680 + 40 \\ &= 9666 \end{aligned}$$

Thus 103 is a quadratic residue of the number 379.

**6.**

To determine the relation of the number $b$ to $a$, it is not necessary to calculate the individual terms of the sum $\varphi(a, b)$; it is enough to know how many of them are odd. Our rule can therefore also be stated as follows:

As above, take $a = \beta b + c,\ b = \gamma c + d,\ c = \delta d + e$ etc., until the sequence of numbers $a,\ b,\ c,\ d,\ e$ etc. reaches unity. Put $[\frac{1}{2}a] = a',\ [\frac{1}{2}b] = b',\ [\frac{1}{2}c] = c'$ etc., and let $\mu$ count the odd numbers in the sequence $a', b', c'$ etc. that are immediately followed by an odd number. Further, let $\nu$ count the odd numbers in the sequence $\beta, \gamma, \delta$ etc. whose corresponding numbers in the sequence $b', c', d'$ etc., respectively, have the form $4n + 1$ or $4n + 2$. Then $b$ is a quadratic residue or a nonresidue of $a$ according as $\mu + \nu$ is even or odd, with the sole exception that, when $b$ is even and $a$ is of the form $8n + 3$ or $8n + 5$, the opposite rule holds.

In our example the sequence $a', b', c', d', e'$ has two instances in which an odd entry is immediately followed by an odd entry, so $\mu = 2$; the sequence $\beta', \gamma', \delta', \varepsilon'$ does contain two odd numbers, but their corresponding numbers in the sequence $b', c', d', e'$ have the form $4n+3$, so $\nu = 0$. Thus $\mu+\nu$ is even, and consequently 103 is a quadratic residue of the number 379.

---

# THEORY

# OF BIQUADRATIC RESIDUES

## FIRST PAPER

BY

CARL FRIEDRICH GAUSS

PRESENTED TO THE ROYAL SOCIETY ON APRIL 5, 1825.

---

Commentationes societatis regiae scientiarum Gottingensis recentiores. Vol. VI.
Göttingen MDCCCXXVIII.

---

# THEORY OF BIQUADRATIC RESIDUES.

## FIRST PAPER.

---

**1.**

The theory of quadratic residues reduces to a few fundamental theorems that rank among the most beautiful jewels of higher arithmetic. As is well known, these were first easily discovered through inductive observation and then proved in many different ways, leaving nothing further to be desired.

The theory of cubic and biquadratic residues, however, requires much deeper investigation. When we began to explore it in 1805, besides the results that lie, as it were, at the threshold, some special theorems presented themselves, remarkable both for their simplicity and for the difficulty of their proofs. But we soon found that the principles of arithmetic used up to then were wholly insufficient to establish the general theory. Rather, the theory necessarily requires the field of higher arithmetic to be enlarged, as it were, infinitely many times over. What this means will become entirely clear as these investigations continue. As soon as we entered this new field, inductive observation immediately opened a path to knowledge of the simplest theorems encompassing the entire theory. Their proofs, however, lay so deeply hidden that only after many unsuccessful attempts could they finally be brought to light.

Now that we are preparing to publish these studies, we shall begin with the theory of biquadratic residues. In this first paper we shall present the investigations that could already be completed on this side of the enlarged field of arithmetic. They pave the way to that field, as it were, and also contribute some new developments to the theory of the division of the circle.

**2.**

We introduced the notion of a biquadratic residue in Article 115 of the *Disquisitiones Arithmeticae*: an integer $a$, positive or negative, is called a biquadratic residue of the integer $p$ if $a$ can be congruent to a fourth power modulo $p$, and a biquadratic nonresidue if no such congruence exists. Throughout the following investigations, unless we expressly state otherwise, we shall assume that the modulus $p$ is an odd positive prime and that $a$ is not divisible by $p$, since all the remaining cases can very easily be reduced to this one.

**3.**

Clearly, every biquadratic residue of $p$ is also a quadratic residue of the same number, and hence every quadratic nonresidue is also a biquadratic nonresidue. The converse also holds whenever $p$ is a prime of the form $4n+3$.

For if, in this case, $a$ is a quadratic residue of $p$, let us put $a \equiv bb \pmod{p}$. Here $b$ is either a quadratic residue of $p$ or a nonresidue. In the first case, let us put $b \equiv cc$, giving $a \equiv c^4$; that is, $a$ is a biquadratic residue of $p$. In the second case, $-b$ is a quadratic residue of $p$ (since $-1$ is a nonresidue of every prime of the form $4n + 3$). Putting $-b \equiv cc$ again gives $a \equiv c^4$, and $a$ is a biquadratic residue of $p$. It is also easy to see that in this case the congruence $x^4 \equiv a \pmod{p}$ has no solutions besides $x \equiv c$ and $x \equiv -c$. Since these evident propositions exhaust the entire theory of biquadratic residues for prime moduli of the form $4n+3$, we shall exclude these moduli altogether from our investigation, restricting it to prime moduli of the form $4n + 1$.

**4.**

Thus, when $p$ is a prime of the form $4n+1$, the converse of the proposition in the preceding article does not hold: there can be quadratic residues that are not also biquadratic residues. This happens whenever a quadratic residue is congruent to the square of a quadratic nonresidue. For put $a \equiv bb$, with $b$ a quadratic nonresidue of $p$. If the congruence $x^4 \equiv a$ could be satisfied by a value $x \equiv c$, we would have $c^4 \equiv bb$, so that the product $(cc-b)(cc+b)$ would be divisible by $p$. Then $p$ would have to divide either the factor $cc - b$ or the other factor $cc + b$; that is, either $+b$ or $-b$ would be a quadratic residue of $p$, and therefore both would be (since $-1$ is a quadratic residue), contrary to the hypothesis.

All integers not divisible by $p$ could therefore be divided into three classes: the first would contain the biquadratic residues, the second the biquadratic nonresidues that are also quadratic residues, and the third the quadratic nonresidues. Clearly it suffices to classify only the numbers $1, 2, 3 \ldots . p - 1$. Half would belong to the third class, while the other half would be divided between the first and second classes.

**5.**

It will be better, however, to establish four classes as follows.

Let $A$ be the collection of all biquadratic residues of $p$ between 1 and $p-1$, inclusive, and let $e$ be any chosen quadratic nonresidue of $p$. Further, let $B$ be the collection of the least positive residues of the products $eA$ modulo $p$, and let $C, D$ respectively be the collections of the least positive residues of the products $eeA$, $e^3A$ modulo $p$. It is then easy to see that the individual numbers in $B$ are distinct, as are those in $C$ and those in $D$; zero cannot occur among any of these numbers. Also, all numbers in $A$ and $C$ are quadratic residues of $p$, while all those in $B$ and $D$ are quadratic nonresidues. Thus neither of the collections $A, C$ can have a number in common with $B$ or $D$. But neither can $A$ have a number in common with $C$, nor $B$ with $D$. For suppose

I. that some number in $A$, say $a$, also occurs in $C$, where it arises from the congruent product $eea'$, with $a'$ a number in $A$. Put $a \equiv \alpha^4$, $a' \equiv \alpha'^4$, and choose an integer $\Theta$ such that $\Theta\alpha' \equiv 1$. We then have $ee\alpha'^4 \equiv \alpha^4$, and multiplying by $\Theta^4$ gives

$$ee \equiv \alpha^4\Theta^4$$

That is, $ee$ is a biquadratic residue, and hence $e$ a quadratic residue, contrary to the hypothesis.

II. Similarly, suppose that a number belongs to both collections $B, D$, arising from the products $ea$, $e^3a'$, with $a, a'$ numbers in $A$. The congruence $ea \equiv e^3a'$ would imply $a \equiv eea'$. We would thus have a number arising from the product $eea'$ that belonged to both $C$ and $A$, which we have just proved impossible.

It is also easy to prove that *all* quadratic residues of $p$ between 1 and $p-1$, inclusive, must occur either in $A$ or in $C$, and all quadratic nonresidues of $p$ between those limits must occur either in $B$ or in $D$. For

I. Every such quadratic residue that is also a biquadratic residue belongs to $A$ by hypothesis.

II. Write a quadratic residue $h$ (less than $p$) that is also a biquadratic nonresidue as $\equiv gg$, where $g$ is a quadratic nonresidue. Choose an integer $\gamma$ such that $e\gamma \equiv g$. Then $\gamma$ is a quadratic residue of $p$, which we write as $\equiv kk$. Hence

$$h \equiv gg \equiv ee\gamma\gamma \equiv eek^4$$

Since the least residue of $k^4$ belongs to $A$, the number $h$, which arises by multiplying it by $ee$, must belong to $C$.

III. Let $h$ denote a quadratic nonresidue of $p$ between 1 and $p-1$. Find an integer $g$ between the same limits such that $eg \equiv h$. Then $g$ is a quadratic

residue and so belongs to either $A$ or $C$. In the first case, $h$ evidently belongs to $B$, and in the second to $D$.

All this shows that the numbers $1, 2, 3 \ldots . p-1$ are distributed among the four lists $A, B, C, D$ so that each occurs in one of them; each list must therefore contain $\frac{1}{4}(p-1)$ numbers. In this classification, the numbers belonging to $A$ and $C$ are intrinsic to those classes. The distinction between $B$ and $D$, however, is arbitrary insofar as it depends on the choice of $e$, which itself always belongs to $B$. Thus, if a number from $D$ is chosen instead, the classes $B, D$ are interchanged.

**6.**

Since $-1$ is a quadratic residue of $p$, let us put $-1 \equiv ff \pmod{p}$. The four roots of the congruence $x^4 \equiv 1$ are then $1, f, -1, -f$. Thus if a is a biquadratic residue of $p$, say $\equiv \alpha^4$, the four roots of the congruence $x^4 \equiv a$ are $\alpha, f\alpha, -\alpha, -f\alpha$, which are easily seen to be pairwise incongruent. Hence, if we collect the least positive residues of the fourth powers $1, 16, 81, 256 \ldots . (p-1)^4$, they are always equal in groups of four, giving $\frac{1}{4}(p-1)$ distinct biquadratic residues that form the collection $A$. If we collect the least residues of the fourth powers only as far as $(\frac{1}{2}p - \frac{1}{2})^4$, each occurs twice.

**7.**

The product of two biquadratic residues is clearly a biquadratic residue: multiplying two numbers from class $A$ always gives a product whose least positive residue belongs to the same class. Similarly, the product of a number from $B$ and one from $D$, or of a number from $C$ and one from $C$, has its least residue in $A$.

In $B$ lie the residues of the products $A.B$ and $C.D$; in $C$ those of $A.C$, $B.B$ and $D.D$; and finally in $D$ those of $A.D$ and $B.C$.

The proofs are so evident that it suffices to indicate one. Let, for example, $c$ and $d$ be numbers from $C$ and $D$, and put $c \equiv eea$, $d \equiv e^3a'$, with $a, a'$ numbers from $A$. Then $e^4aa'$ is a biquadratic residue, so its least residue belongs to $A$. For the product we have $cd \equiv e.e^4aa'$, so its least residue belongs to $B$.

We can now also easily decide which class contains a product of several factors. Assign to the classes $A, B, C, D$ the characters $0, 1, 2, 3$, respectively. The character of a product is then either the sum of the characters of its individual factors or the least residue of that sum modulo 4.

**8.**

It seemed worthwhile to develop these elementary propositions without using the theory of power residues. With its help, everything can be proved much more easily still.

Let $g$ be a primitive root for the modulus $p$: a number such that in the sequence of powers $g, gg, g^3 \ldots$., none before $g^{p-1}$ is congruent to unity modulo $p$. Then the least positive residues of the numbers $1, g, gg, g^3 \ldots . g^{p-2}$ agree, apart from their order, with $1, 2, 3 \ldots . . p-1$, and are distributed among four classes as follows:

| to | least residues of the numbers |
|---|---|
| $A$ | $1,\ g^4, g^8, g^{12} \ldots . . g^{p-5}$ |
| $B$ | $g,\ g^5, g^9, g^{13} \ldots . . g^{p-4}$ |
| $C$ | $gg, g^6, g^{10}, g^{14} \ldots . . g^{p-3}$ |
| $D$ | $g^3, g^7, g^{11}, g^{15} \ldots . . g^{p-2}$ |

All the preceding propositions follow directly from this.

Just as the numbers $1, 2, 3 \ldots . p-1$ have here been divided among four classes, whose collections we denote by $A, B, C, D$, we may assign *any* integer not divisible by $p$ to one of these classes according to its least residue modulo $p$.

**9.**

We denote by $f$ the least residue of the power $g^{\frac{1}{4}(p-1)}$ modulo $p$. Since $ff \equiv g^{\frac{1}{2}(p-1)} \equiv -1$ (*Disquis. Arithm.*, Article 62), the symbol $f$ clearly has the same meaning here as in Article 6. Thus the power $g^{\frac{1}{4}\lambda(p-1)}$, with $\lambda$ a positive integer, is congruent modulo $p$ to $1, f, -1, -f$, according as $\lambda$ has the form $4m, 4m+1, 4m+2, 4m+3$, respectively, or according as the least residue of $g^\lambda$ belongs to $A, B, C, D$, respectively. This gives us a very simple criterion for deciding which class contains a given number $h$ not divisible by $p$: $h$ belongs to $A, B, C$ or $D$ according as the power $h^{\frac{1}{4}(p-1)}$ is congruent modulo $p$ to $1, f, -1$ or $-f$.

As a corollary, $-1$ always belongs to class $A$ when $p$ has the form $8n+1$, and to class $C$ when $p$ has the form $8n+5$. A proof of this theorem independent of the theory of power residues can easily be obtained from what we established in Article 115, III of the *Disquisitiones Arithmeticae*.

**10.**

Since *all* primitive roots for the modulus $p$ arise as residues of the powers $g^\lambda$, on taking for $\lambda$ all numbers coprime to $p-1$, it is easy to see that these roots are equally divided between $B$ and $D$, with the base $g$ always in $B$. If we replace $g$ as base by another primitive root from $B$, the classification remains the same; if we use a primitive root from $D$ as base, the classes $B$ and $D$ are interchanged.

If the classification is based on the criterion in the preceding article, the distinction between $B$ and $D$ depends on which root of the congruence $xx \equiv -1 \pmod{p}$ we choose as the characteristic number $f$.

**11.**

To make it easier to illustrate by examples the subtler investigations that we are about to undertake, we give here the construction of the classes for all moduli below 100. For each modulus we have chosen the least primitive root.

$$p = 5$$

$$g = 2,\ f = 2$$

| | |
|---|---|
| $A$ | $1$ |
| $B$ | $2$ |
| $C$ | $4$ |
| $D$ | $3$ |

$$p = 13$$

$$g = 2,\ f = 8$$

| | |
|---|---|
| $A$ | $1, 3, 9$ |
| $B$ | $2, 5, 6$ |
| $C$ | $4, 10, 12$ |
| $D$ | $7, 8, 11$ |

$$p = 17$$

$$g = 3,\ f = 13$$

| | |
|---|---|
| $A$ | $1, 4, 13, 16$ |
| $B$ | $3, 5, 12, 14$ |
| $C$ | $2, 8, 9, 15$ |
| $D$ | $6, 7, 10, 11$ |

$$p = 29$$

$$g = 2,\ f = 12$$

| | |
|---|---|
| $A$ | $1, 7, 16, 20, 23, 24, 25$ |
| $B$ | $2, 3, 11, 14, 17, 19, 21$ |
| $C$ | $4, 5, 6, 9, 13, 22, 28$ |
| $D$ | $8, 10, 12, 15, 18, 26, 27$ |

$$p = 37$$

$$g = 2,\ f = 31$$

| | |
|---|---|
| $A$ | $1, 7, 9, 10, 12, 16, 26, 33, 34$ |
| $B$ | $2, 14, 15, 18, 20, 24, 29, 31, 32$ |
| $C$ | $3, 4, 11, 21, 25, 27, 28, 30, 36$ |
| $D$ | $5, 6, 8, 13, 17, 19, 22, 23, 35$ |

$$p = 41$$

$$g = 6,\ \ f = 32$$

| | |
|---|---|
| $A$ | $1, 4, 10, 16, 18, 23, 25, 31, 37, 40$ |
| $B$ | $6, 14, 15, 17, 19, 22, 24, 26, 27, 35$ |
| $C$ | $2, 5, 8, 9, 20, 21, 32, 33, 36, 39$ |
| $D$ | $3, 7, 11, 12, 13, 28, 29, 30, 34, 38$ |

$$p = 53$$

$$g = 2,\ \ f = 30$$

| | |
|---|---|
| $A$ | $1, 10, 13, 15, 16, 24, 28, 36, 42, 44, 46, 47, 49$ |
| $B$ | $2, 3, 19, 20, 26, 30, 31, 32, 35, 39, 41, 45, 48$ |
| $C$ | $4, 6, 7, 9, 11, 17, 25, 29, 37, 38, 40, 43, 52$ |
| $D$ | $5, 8, 12, 14, 18, 21, 22, 23, 27, 33, 34, 50, 51$ |

$$p = 61$$

$$g = 2,\ \ f = 11$$

| | |
|---|---|
| $A$ | $1, 9, 12, 13, 15, 16, 20, 22, 25, 34, 42, 47, 56, 57, 58$ |
| $B$ | $2, 7, 18, 23, 24, 26, 30, 32, 33, 40, 44, 50, 51, 53, 55$ |
| $C$ | $3, 4, 5, 14, 19, 27, 36, 39, 41, 45, 46, 48, 49, 52, 60$ |
| $D$ | $6, 8, 10, 11, 17, 21, 28, 29, 31, 35, 37, 38, 43, 54, 59$ |

$$p = 73$$

$$g = 5,\ \ f = 27$$

| | |
|---|---|
| $A$ | $1, 2, 4, 8, 9, 16, 18, 32, 36, 37, 41, 55, 57, 64, 65, 69, 71, 72$ |
| $B$ | $5, 7, 10, 14, 17, 20, 28, 33, 34, 39, 40, 45, 53, 56, 59, 63, 66, 68$ |
| $C$ | $3, 6, 12, 19, 23, 24, 25, 27, 35, 38, 46, 48, 49, 50, 54, 61, 67, 70$ |
| $D$ | $11, 13, 15, 21, 22, 26, 29, 30, 31, 42, 43, 44, 47, 51, 52, 58, 60, 62$ |

$$p = 89$$

$$g = 3,\ \ f = 34$$

| | |
|---|---|
| $A$ | $1, 2, 4, 8, 11, 16, 22, 25, 32, 39, 44.\ 45,$ |
| | $50, 57, 64, 67, 73, 78, 81, 85, 87, 88$ |
| $B$ | $3, 6, 7, 12, 14, 23, 24, 28, 33, 41, 43, 46,$ |
| | $48, 56, 61, 65, 66, 75, 77, 82, 83, 86$ |
| $C$ | $5, 9, 10, 17, 18, 20, 21, 34, 36, 40, 42, 47,$ |
| | $49, 53, 55, 68, 69, 71, 72, 79, 80, 84$ |
| $D$ | $13, 15.\ 19, 26, 27, 29.\ 30, 31, 35, 37, 38.\ 51,$ |
| | $52, 54, 58, 59, 60, 62, 63, 70, 74, 76$ |

$$p = 97$$

$$g = 5,\ \ f = 22$$

| | |
|---|---|
| $A$ | $1, 4, 6, 9, 16, 22, 24, 33, 35, 36, 43, 47,$ |
| | $50, 54, 61, 62, 64, 73, 75, 81, 88, 91, 93, 96$ |
| $B$ | $5, 13, 14, 17, 19, 20, 21, 23, 29, 30, 41, 45,$ |
| | $52, 56, 67, 68, 74, 76, 77, 78, 80, 83, 84, 92$ |
| $C$ | $2, 3, 8, 11, 12, 18, 25, 27, 31, 32, 44, 48,$ |
| | $49, 53, 65.\ 66, 70, 72, 79, 85, 86, 89, 94, 95$ |
| $D$ | $7, 10, 15, 26, 28, 34, 37, 38, 39.\ 40, 42, 46,$ |
| | $51, 55, 57, 58, 59, 60, 63, 69, 71, 82, 87, 90$ |

**12.**

Since 2 is a quadratic residue of every prime of the form $8n + 1$, but a nonresidue of every prime of the form $8n + 5$, for prime moduli of the first form 2 belongs to class $A$ or $C$, and for moduli of the second form to $B$ or $D$. The distinction between $B$ and $D$ is not intrinsic, depending only on the choice of $f$, so we shall set moduli of the form $8n + 5$ aside for a while. An *inductive investigation* of moduli of the form $8n + 1$ gives 2 in $A$ for $p = 73, 89, 113, 233, 257, 281, 337, 353$, etc.; in contrast, 2 belongs to $C$ for $p = 17, 41, 97, 137, 193, 241, 313, 401, 409, 433, 449, 457$, etc.

Since, moreover, for a prime modulus of the form $8n+1$, $-1$ is a biquadratic residue, $-2$ clearly always belongs to the same class as $+2$.

**13.**

When we compare the examples in the preceding article, at least at first sight no simple criterion seems to offer itself for distinguishing the first moduli from the second. Nevertheless, there are *two* such criteria, remarkable for their elegance and simplicity. The following considerations will prepare the way to one of them.

The modulus $p$, being a prime of the form $8n+1$, can be expressed in the form $aa+2bb$ in exactly one way (*Disquiss. Arithm.*, Article 182, II). We assume that the roots $a, b$ are taken positive. Clearly $a$ is odd and $b$ even; let us put $b = 2^\lambda c$, with $c$ odd. We now observe:

I. Since $p \equiv aa \pmod{c}$, $p$ is a quadratic residue of $c$, and hence also of each prime factor into which $c$ is resolved. Conversely, by the fundamental theorem, each of these prime factors is a quadratic residue of $p$, and so their product $c$ is also a quadratic residue of $p$. Since this also holds for 2, $b$ is clearly a quadratic residue of $p$, and hence both $bb$ and $-bb$ are biquadratic residues.

II. Thus $-2bb$ must belong to the same class as 2. Since $aa \equiv -2bb$, 2 clearly belongs to class $A$ or $C$ according as $a$ is a quadratic residue of $p$ or a quadratic nonresidue.

III. Now suppose that $a$ has been resolved into prime factors. Denote those of the form $8m+1$ or $8m+7$ by $\alpha, \alpha', \alpha''$, etc., and those of the form $8m+3$ or $8m+5$ by $\beta, \beta', \beta''$, etc.; let the number of the latter be $= \mu$. Since $p \equiv 2bb \pmod{a}$, $p$ is a quadratic residue of those prime factors of $a$ of which 2 is a quadratic residue, that is, of $\alpha, \alpha', \alpha''$, etc.; it is a quadratic nonresidue of those factors of which 2 is a quadratic nonresidue, that is, of $\beta, \beta', \beta''$, etc. Conversely, by the fundamental theorem, each of $\alpha, \alpha', \alpha''$, etc., is a quadratic residue of $p$, while each of $\beta, \beta', \beta''$, etc., is a quadratic nonresidue. We therefore conclude that their product $a$ is a quadratic residue of $p$ or a nonresidue according as $\mu$ is even or odd.

IV. It is easy to verify that the product of all $\alpha, \alpha', \alpha''$, etc., has the form $8m+1$ or $8m+7$, and that the same holds for the product of all $\beta, \beta', \beta''$, etc., when their number is even. In this case the product $a$ must therefore also have the form $8m+1$ or $8m+7$. In contrast, the product of all $\beta, \beta', \beta''$, etc., when their number is odd, has the form $8m+3$ or $8m+5$, and the same therefore holds for the product $a$ in this case.

From all this we obtain the elegant theorem:

*Whenever $a$ has the form $8m+1$ or $8m+7$, 2 belongs to $A$; whenever $a$ has the form $8m+3$ or $8m+5$, 2 belongs to $C$.*

The examples listed in the preceding article confirm this. The first moduli decompose as follows: $73 = 1 + 2 \,.\, 36$, $89 = 81 + 2 \,.\, 4$, $113 = 81 + 2 \,.\, 16$, $233 = 225 + 2 \,.\, 4$, $257 = 225 + 2 \,.\, 16$, $281 = 81 + 2 \,.\, 100$, $337 = 49 + 2 \,.\, 144$, $353 = 225 + 2 \,.\, 64$; the second as follows: $17 = 9 + 2 \,.\, 4$, $41 = 9 + 2 \,.\, 16$, $97 = 25 + 2 \,.\, 36$, $137 = 9 + 2 \,.\, 64$, $193 = 121 + 2 \,.\, 36$, $241 = 169 + 2 \,.\, 36$, $313 = 25 + 2 \,.\, 144$, $401 = 9 + 2 \,.\, 196$, $409 = 121 + 2 \,.\, 144$, $433 = 361 + 2 \,.\, 36$, $449 = 441 + 2 \,.\, 4$, $457 = 169 + 2 \,.\, 144$.

**14.**

The decomposition of $p$ into a square and twice a square has revealed such a remarkable connection with the classification of 2 that it seems worthwhile to ask whether a decomposition into two squares, which $p$ is also known to admit, might give a similar result. Here are the decompositions of the numbers $p$ for which 2 belongs to class

| $A$ | $C$ |
|---|---|
| $9 + 64$ | $1 + 16$ |
| $25 + 64$ | $25 + 16$ |
| $49 + 64$ | $81 + 16$ |
| $169 + 64$ | $121 + 16$ |
| $1 + 256$ | $49 + 144$ |
| $25 + 256$ | $225 + 16$ |
| $81 + 256$ | $169 + 144$ |
| $289 + 64$ | $1 + 400$ |
| | $9 + 400$ |
| | $289 + 144$ |
| | $49 + 400$ |
| | $441 + 16$ |

First we observe that, of the two squares into which $p$ decomposes, one must be odd, which we put $= aa$, and the other even, which we put $= bb$. Since $aa$ has the form $8n+1$, values of $b$ that are twice an odd number clearly correspond to values of $p$ of the form $8n + 5$, which are excluded from our inductive investigation here, since they would have 2 in class $B$ or $D$. For values of $p$ of the form $8n + 1$, however, $b$ must be divisible by four. If we may trust the inductive evidence presented by this table, 2 belongs to class $A$ for every modulus for which $b$ has the form $8n$, and to class $C$ for every modulus for which $b$ has the form $8n + 4$. This theorem, however, requires much deeper investigation than the one obtained in the preceding article. Its proof must be preceded by several preliminary investigations concerning the order in which the numbers in $A, B, C, D$ follow one another.

**15.**

Let us denote the number of numbers in $A$ immediately followed by a number in $A, B, C, D$, respectively, by $(00), (01), (02), (03)$; likewise the number of numbers in $B$ followed by a number in $A, B, C, D$, respectively, by $(10), (11), (12), (13)$. Similarly, let $C$ contain respectively $(20), (21), (22), (23)$ numbers, and $D$ contain $(30), (31), (32), (33)$ numbers, followed by a number in $A, B, C, D$. We propose to determine these sixteen counts a priori. To help readers compare the general reasoning with examples, it seemed useful to give here the numerical values of the entries in the scheme $(S)$

$$\begin{array}{llll} (00), & (01), & (02), & (03) \\ (10), & (11), & (12), & (13) \\ (20). & (21), & (22), & (23) \\ (30), & (31), & (32), & (33) \end{array}$$

for each modulus whose classification we gave in Article 11.

| $p = 5$ | $p = 13$ | $p = 17$ | $p = 29$ |
|---|---|---|---|
| $0, 1, 0.\ 0$ | $0, 1, 2, 0$ | $0, 2, 1, 0$ | $2.\ 3, 0.\ 2$ |
| $0, 0, 0, 1$ | $1, 1, 0, 1$ | $2, 0, 1, 1$ | $1, 1, 2, 3$ |
| $0, 0, 0, 0$ | $0, 1, 0, 1$ | $1, 1, 1, 1$ | $2, 1, 2.\ 1$ |
| $0, 0, 1, 0$ | $1, 0, 1, 1$ | $0, 1, 1, 2$ | $1, 2, 3, 1$ |

| $p = 37$ | $p = 41$ | $p = 53$ | $p = 61$ |
|---|---|---|---|
| $2, 1, 2, 4$ | $0, 4, 3, 2$ | $2, 3, 6, 2$ | $4, 3, 2, 6$ |
| $2, 2, 4, 1$ | $4, 2, 2, 2$ | $4, 4, 2, 3$ | $3, 3, 6, 3$ |
| $2, 2, 2, 2$ | $3, 2, 3, 2$ | $2, 4, 2, 4$ | $4, 3, 4, 3$ |
| $2, 4, 1, 2$ | $2, 2, 2, 4$ | $4, 2, 3, 4$ | $3, 6, 3, 3$ |

| $p = 73$ | $p = 89$ | $p = 97$ |
|---|---|---|
| $5, 6, 4, 2$ | $3, 8, 6, 4$ | $2, 6, 7, 8$ |
| $6, 2, 5, 5$ | $8, 4, 5, 5$ | $6, 8, 5, 5$ |
| $4, 5, 4, 5$ | $6, 5, 6, 5$ | $7, 5, 7, 5$ |
| $2, 5, 5, 6$ | $4, 5, 5, 8$ | $8, 5, 5, 6$ |

Since moduli of the forms $8n + 1$ and $8n + 5$ behave differently, we must treat them separately. We begin with the first.

**16.**

The symbol $(00)$ indicates how many different ways the equation $\alpha + 1 = \alpha'$ can be satisfied, with $\alpha, \alpha'$ denoting arbitrary numbers in $A$. For a modulus of the form $8n + 1$, as understood here, $\alpha'$ and $p - \alpha'$ belong to the same

collection. We may therefore say more elegantly that (00) expresses the number of different ways of satisfying the equation $1 + \alpha + \alpha' = p$. Clearly the congruence $1 + \alpha + \alpha' \equiv 0 \pmod{p}$ can also take the place of this equation.

Likewise,

(01) indicates the number of solutions of the congruence $1+\alpha+\beta \equiv 0 \pmod{p}$
(02) the number of solutions of the congruence $1 + \alpha + \gamma \equiv 0$
(03) the number of solutions of the congruence $1 + \alpha + \delta \equiv 0$
(11) the number of solutions of the congruence $1 + \beta + \beta' \equiv 0$, etc.

where $\beta$ and $\beta'$ denote arbitrary numbers in $B$, $\gamma$ numbers in $C$, and $\delta$ numbers in $D$. We immediately obtain the following six equations:

$$(01) = (10), \quad (02) = (20), \quad (03) = (30),$$
$$(12) = (21), \quad (13) = (31), \quad (23) = (32)$$

From any given solution of the congruence $1 + \alpha + \beta \equiv 0$ we obtain a solution of $1 + \delta + \delta' \equiv 0$ by taking for $\delta$ the number between $1 \ldots . p - 1$ that satisfies $\beta\delta \equiv 1$ (which clearly belongs to $D$), and for $\delta'$ the least positive residue of the product $\alpha\delta$ (which likewise belongs to $D$). The reverse passage from a given solution of $1 + \delta + \delta' \equiv 0$ to a solution of $1 + \alpha + \beta \equiv 0$ is equally clear: choose $\beta$ so that $\beta\delta \equiv 1$, and at the same time put $\alpha \equiv \beta\delta'$. Thus the two congruences have the same number of solutions, or $(01) = (33)$.

Similarly, from the congruence $1 + \alpha + \gamma \equiv 0$ we derive $\gamma' + \gamma'' + 1 \equiv 0$ by choosing $\gamma'$ in $C$ so that $\gamma\gamma' \equiv 1$, and $\gamma''$ in the same collection congruent to the product $\alpha\gamma'$. It follows easily that these two congruences admit the same number of solutions, or $(02) = (22)$.

Likewise, from $1 + \alpha + \delta \equiv 0$ we derive $\beta + \beta' + 1 \equiv 0$ by choosing $\beta, \beta'$ so that $\beta\delta \equiv 1$, $\beta\alpha \equiv \beta'$, giving $(03) = (11)$.

Finally, from $1 + \beta + \gamma \equiv 0$ we similarly derive both $\delta + 1 + \beta' \equiv 0$ and $\gamma' + \delta' + 1 \equiv 0$, and conclude that $(12) = (13) = (23)$.

We have thus obtained eleven equations among our sixteen unknowns, reducing them to five. The scheme $S$ can therefore be written as

$$\begin{array}{cccc} h, & i, & k, & l \\ i, & l, & m, & m \\ k, & m, & k, & m \\ l, & m, & m, & i \end{array}$$

Three further equations of condition can easily be added. Every number in $A$, except the last, $p - 1$, must be followed by a number in one of the collections $A, B, C$ or $D$. Thus

$$(00) + (01) + (02) + (03) = 2n - 1$$

and likewise

$$(10)+(11)+(12)+(13)=2n$$
$$(20)+(21)+(22)+(23)=2n$$
$$(30)+(31)+(32)+(33)=2n$$

In the notation just introduced, the first three equations give

$$h+i+k+l=2n-1$$
$$i+l+2m=2n$$
$$k+m=n$$

The fourth is identical to the second. These equations allow us to eliminate three unknowns, so that all sixteen have now been reduced to two.

**17.**

To determine them completely, it will be useful to investigate the number of solutions of the congruence

$$1+\alpha+\beta+\gamma\equiv 0 \pmod{p}$$

where $\alpha,\beta,\gamma$ denote arbitrary numbers in $A,B,C$. Clearly $\alpha=p-1$ is inadmissible, since $\beta+\gamma\equiv 0$ cannot hold. Substituting the remaining values for $\alpha$ in turn gives $h,i,k,l$ values of $1+\alpha$ belonging to $A,B,C,D$, respectively. For each *given* value of $1+\alpha$ belonging to $A$, say $1+\alpha=\alpha^0$, the congruence $\alpha^0+\beta+\gamma\equiv 0$ admits as many solutions as $1+\beta'+\gamma'\equiv 0$ (put $\beta\equiv\alpha^0\beta'$, $\gamma\equiv\alpha^0\gamma'$), that is, $(12)=m$ solutions. Likewise, for each given value of $1+\alpha$ belonging to $B$, say $1+\alpha=\beta^0$, the congruence $\beta^0+\beta+\gamma\equiv 0$ has as many solutions as $1+\alpha'+\beta'\equiv 0$ (put $\beta\equiv\beta^0\alpha'$, $\gamma\equiv\beta^0\beta'$), that is, $(01)=i$ solutions. Similarly, for each given value of $1+\alpha$ belonging to $C$, say $1+\alpha=\gamma^0$, the congruence $\gamma^0+\beta+\gamma\equiv 0$ can be solved in as many different ways as $1+\delta+\alpha'\equiv 0$ (put $\beta\equiv\gamma^0\delta$, $\gamma\equiv\gamma^0\alpha'$), so the number of solutions is $(03)=l$. Finally, for each given value of $1+\alpha$ belonging to $D$, say $1+\alpha=\delta^0$, the congruence $\delta^0+\beta+\gamma\equiv 0$ has as many solutions as $1+\gamma'+\delta'\equiv 0$ (put $\beta\equiv\delta^0\gamma'$, $\gamma\equiv\delta^0\delta'$), that is, $(23)=m$ solutions. Adding all these counts, the congruence $1+\alpha+\beta+\gamma\equiv 0$ clearly admits

$$hm+ii+kl+lm$$

different solutions.

In exactly the same way we find that, on substituting for $\beta$ the numbers in $B$ in turn, the sum $1+\beta$ takes respectively $(10),(11),(12),(13)$, or $i,l,m,m$, values belonging to $A,B,C,D$. For each *given* value of $1+\beta$ belonging to these collections, the congruence $1+\beta+\alpha+\gamma\equiv 0$ admits respectively

$(02), (31), (20), (13)$, or $k, m, k, m$, different solutions. The total number of solutions is therefore

$$= ik + lm + km + mm$$

We obtain the same value by basing the calculation on the values of the sum $1 + \gamma$.

**18.**

These two expressions for the same count give the equation

$$0 = hm + ii + kl - ik - km - mm$$

and hence, eliminating $h$ by means of $h = 2m - k - 1$,

$$0 = (k - m)^2 + ii + kl - ik - kk - m$$

The last two equations of Article 16 give $k = \frac{1}{2}(l + i)$. Substituting this value turns $ii + kl - ik - kk$ into $\frac{1}{4}(l - i)^2$, so the preceding equation, multiplied by 4, becomes

$$0 = 4(k - m)^2 + (l - i)^2 - 4m$$

Since $4m = 2(k + m) - 2(k - m) = 2n - 2(k - m)$, it follows that

$$2n = 4(k - m)^2 + 2(k - m) + (l - i)^2$$

or

$$8n + 1 = (4(k - m) + 1)^2 + 4(l - i)^2$$

Thus, putting

$$4(k - m) + 1 = a, \quad 2l - 2i = b$$

we have

$$p = aa + bb$$

It is known that $p$ can be decomposed into two squares in exactly one way, with the odd square taken as $aa$ and the even square as $bb$, so $aa, bb$ are completely determined numbers. But $a$ itself is also completely determined: the square root must be taken positive or negative according as the positive root has the form $4M+1$ or $4M+3$. We shall shortly discuss how to determine the sign of $b$.

Combining these new equations with the last three equations of Article 16 completely determines the five numbers $h, i, k, l, m$ in terms of $a, b$ and $n$, as follows:

$$\begin{aligned} 8h &= 4n - 3a - 5 \\ 8i &= 4n + a - 2b - 1 \\ 8k &= 4n + a - 1 \\ 8l &= 4n + a + 2b - 1 \\ 8m &= 4n - a + 1 \end{aligned}$$

If, in place of $n$, we prefer to use the modulus $p$, the scheme $S$, with every entry multiplied by 16 to avoid fractions, is

| | | | |
|---|---|---|---|
| $p-6a-11$ | $p+2a-4b-3$ | $p+2a-3$ | $p+2a+4b-3$ |
| $p+2a-4b-3$ | $p+2a+4b-3$ | $p-2a+1$ | $p-2a+1$ |
| $p+2a-3$ | $p-2a+1$ | $p+2a-3$ | $p-2a+1$ |
| $p+2a+4b-3$ | $p-2a+1$ | $p-2a+1$ | $p+2a-4b-3$ |

**19.**

It remains to show how to choose the sign of $b$. In Article 10 we pointed out that the distinction between $B$ and $D$, which is not intrinsic, depends on the choice of $f$, for which either root of $xx \equiv -1$ must be taken, and that the two collections are interchanged when one root is replaced by the other. Inspection of the scheme just given shows that a similar interchange accompanies a change in the sign of $b$. We can therefore foresee that there must be a connection between the sign of $b$ and $f$. To find it, we first observe that, if $\mu$ denotes a nonnegative integer and $z$ takes all the values $1, 2, 3 \ldots . p-1$, then modulo $p$ we have either $\Sigma z^\mu \equiv 0$ or $\Sigma z^\mu \equiv -1$, according as $\mu$ is not divisible by $p-1$ or is divisible by it. The latter part of the theorem follows from the fact that for a value of $\mu$ divisible by $p-1$, $z^\mu \equiv 1$. We prove the former part as follows. Let $g$ denote a primitive root. All the values of $z$ coincide with the least residues of all $g^y$ when $y$ takes all the values $0, 1, 2, 3 \ldots . p-2$, and hence $\Sigma z^\mu \equiv \Sigma g^{\mu . y}$. Now

$$\Sigma g^{\mu y} = \frac{g^{\mu(p-1)} - 1}{g^\mu - 1}, \quad \text{and hence}$$
$$(g^\mu - 1)\Sigma z^\mu \equiv g^{\mu(p-1)} - 1 \equiv 0$$

For a value of $\mu$ not divisible by $p-1$, $g^\mu$ cannot be congruent to 1, or equivalently $g^\mu - 1$ cannot be divisible by $p$. Hence $\Sigma z^\mu \equiv 0$. Q. E. D.

Now expand the power $(z^4+1)^{\frac{1}{4}(p-1)}$ by the binomial theorem. The preceding lemma gives

$$\Sigma(z^4+1)^{\frac{1}{4}(p-1)} \equiv -2 \text{ (mod. } p)$$

The least residues of all $z^4$ give all the numbers in $A$, each occurring four times. Among the least residues of $z^4+1$ we therefore have

$$\begin{array}{ll} 4(00) & \text{to } A \\ 4(01) & \text{to } B \\ 4(02) & \text{to } C \\ 4(03) & \text{to } D \end{array}$$

belonging to those classes, and four are $= 0$ (namely, when $z^4 \equiv p-1$). Using the criteria for the collections $A, B, C, D$, we obtain

$$\begin{aligned} \Sigma(z^4+1)^{\frac{1}{4}(p-1)} \equiv\ & 4(00) + 4f(01) \\ & - 4(02) - 4f(03) \end{aligned}$$

and hence

$$-2 \equiv 4(00) + 4f(01) - 4(02) - 4f(03)$$

or, substituting for $(00), (01)$, etc., the values found in the preceding article,

$$-2 \equiv -2a - 2 - 2bf$$

We therefore conclude that we must always have $a + bf \equiv 0$, or, multiplying by $f$,

$$b \equiv af$$

This congruence determines the sign of $b$ when $f$ has already been chosen, or determines $f$ when the sign of $b$ is prescribed in some other way.

**20.**

Having completely solved our problem for moduli of the form $8n+1$, we turn to the other case, where $p$ has the form $8n+5$. We can treat it more briefly, since all the arguments differ little from the preceding ones.

Since, for such a modulus, $-1$ belongs to class $C$, the numbers in the collections $A, B, C, D$ have their complements to the sum $p$ in the classes $C, D, A, B$, respectively. It follows readily that

| symbol | denotes the number of solutions of the congruence |
|---|---|
| (00) | $1+\alpha+\gamma \equiv 0$ |
| (01) | $1+\alpha+\delta \equiv 0$ |
| (02) | $1+\alpha+\alpha' \equiv 0$ |
| (03) | $1+\alpha+\beta \equiv 0$ |
| (10) | $1+\beta+\gamma \equiv 0$ |
| (11) | $1+\beta+\delta \equiv 0$ |
| (12) | $1+\beta+\alpha \equiv 0$ |
| (13) | $1+\beta+\beta' \equiv 0$ |
| (20) | $1+\gamma+\gamma' \equiv 0$ |
| (21) | $1+\gamma+\delta \equiv 0$ |
| (22) | $1+\gamma+\alpha \equiv 0$ |
| (23) | $1+\gamma+\beta \equiv 0$ |
| (30) | $1+\delta+\gamma \equiv 0$ |
| (31) | $1+\delta+\delta' \equiv 0$ |
| (32) | $1+\delta+\alpha \equiv 0$ |
| (33) | $1+\delta+\beta \equiv 0$ |

from which we immediately obtain six equations:

$$(00)=(22), \quad (01)=(32), \quad (03)=(12),$$
$$(10)=(23), \quad (11)=(33), \quad (21)=(30)$$

Multiplying the congruence $1+\alpha+\gamma \equiv 0$ by a number $\gamma'$ from the collection $C$, chosen so that $\gamma\gamma' \equiv 1$, and taking for $\gamma''$ the least residue of the product $\alpha\gamma'$, which must evidently also belong to the collection $C$, gives $\gamma'+\gamma''+1 \equiv 0$, from which we conclude that $(00)=(20)$.

In exactly the same way we obtain the equations $(01)=(13)$, $(03)=(31)$, $(10)=(11)=(21)$.

With these eleven equations we can reduce our sixteen unknowns to five and write the scheme $S$ as follows:

$$\begin{array}{cccc} h, & i, & k, & l \\ m, & m, & l, & i \\ h, & m, & h, & m \\ m, & l, & i, & m \end{array}$$

We also have the equations

$$\begin{aligned}(00)+(01)+(02)+(03)&=2n+1\\(10)+(11)+(12)+(13)&=2n+1\\(20)+(21)+(22)+(23)&=2n\\(30)+(31)+(32)+(33)&=2n+1\end{aligned}$$

or, using the notation just introduced, these three (I):

$$\begin{aligned}h+i+k+l&=2n+1\\2m+i+l&=2n+1\\h+m&=n\end{aligned}$$

With their help we can therefore now reduce our unknowns to two.

We shall derive the remaining equations by considering the number of solutions of the congruence $1+\alpha+\beta+\gamma\equiv 0$ (here too, $\alpha,\beta,\gamma$ denote unspecified numbers from the collections $A,B,C$, respectively). Consider *first* that $1+\alpha$ gives $h,i,k,l$ numbers belonging to $A,B,C,D$, respectively, and that for each given value of $\alpha$ in these four cases there are $m,l,i,m$ solutions, respectively. The total number of solutions will be

$$=hm+il+ik+lm$$

*Second*, since $1+\beta$ gives $m,m,l,i$ numbers belonging to $A,B,C,D$, and for each *given* value of $\beta$ in these four cases there are $h,m,h,m$ solutions, the total number of solutions will be

$$=hm+mm+hl+im$$

From this we derive the equation

$$0=mm+hl+im-il-ik-lm$$

which, using the equation $k=2m-h$ obtained from (I), becomes

$$0=mm+hl+hi-il-im-lm$$

Now from equations I we also have $l+i=1+2h$, and hence

$$\begin{aligned}2i&=1+2h+(i-l)\\2l&=1+2h-(i-l)\end{aligned}$$

Substituting these values into the preceding equation gives

$$0=4mm-4m-1-8hm+4hh+(i-l)^2$$

If, finally, we replace $4m$ here with $2(h+m)-2(h-m)$, or, by the last equation in I, with $2n-2(h-m)$, we obtain

$$0 = 4(h-m)^2 - 2n + 2(h-m) - 1 + (i-l)^2$$

and hence

$$8n + 5 = (4(h-m)+1)^2 + 4(i-l)^2$$

Thus, putting

$$4(h-m)+1 = a, \quad 2i-2l = b$$

gives

$$p = aa + bb$$

Since here too $p$ can be split in only one way into two squares, one even and the other odd, $aa$ and $bb$ will be completely determined numbers: evidently $aa$ must be set equal to the odd square and $bb$ to the even one. Moreover, the *sign* of $a$ must be chosen so that $a \equiv 1 \pmod 4$, and the sign of $b$ so that $b \equiv af \pmod p$, as is readily proved by arguments entirely similar to those used in the preceding article.

With these preliminaries, the five numbers $h, i, k, l, m$ are determined in terms of $a, b$ and $n$ as follows:

$$\begin{aligned} 8h &= 4n + a - 1 \\ 8i &= 4n + a + 2b + 3 \\ 8k &= 4n - 3a + 3 \\ 8l &= 4n + a - 2b + 3 \\ 8m &= 4n - a + 1 \end{aligned}$$

Or, if we prefer expressions in terms of $p$, the terms of the scheme $S$, multiplied by 16, will be as follows:

| | | | |
|---|---|---|---|
| $p+2a-7$ | $p+2a+4b+1$ | $p-2a+1$ | $p+2a-4b+1$ |
| $p-2a-3$ | $p-2a-3$ | $p+2a-4b+1$ | $p+2a+4b+1$ |
| $p+2a-7$ | $p-2a-3$ | $p+2a-7$ | $p-2a-3$ |
| $p-2a-3$ | $p+2a-4b+1$ | $p+2a+4b+1$ | $p-2a-3$ |

**21.**

Having solved our problem, we return to the main investigation, now undertaking the complete determination of the collection to which the number 2 belongs.

I. Whenever $p$ has the form $8n+1$, we already know that the number 2 lies either in the collection $A$ or in the collection $C$. In the former case we readily see that the numbers $\frac{1}{2}(p-1)$, $\frac{1}{2}(p+1)$ also belong to $A$, and in the latter to $C$. Now consider that, if $\alpha$ and $\alpha+1$ are consecutive numbers in the collection $A$, then $p-\alpha-1$, $p-\alpha$ are also such numbers; or, equivalently, numbers in the collection $A$ that are followed by a number in the same collection are always paired ($\alpha$ and $p-1-\alpha$). Their number, (00), will therefore always be even, unless one is paired with itself, that is, unless $\frac{1}{2}(p-1)$ belongs to $A$, in which case their number will be odd. From this we conclude that (00) is odd whenever 2 belongs to the collection $A$, and even whenever 2 belongs to $C$. But we have

$$16(00) = aa + bb - 6a - 11$$

or, putting $a = 4q+1$, $b = 4r$ (see Article 14),

$$(00) = qq - q + rr - 1$$

Since $qq-q$ is evidently always even, (00) will be odd or even according as $r$ is even or odd. Thus 2 will belong to $A$ or to $C$ according as $b$ has the form $8m$ or the form $8m+4$. This is precisely the theorem found by inductive investigation in Article 14.

II. We can treat the other case, where $p$ has the form $8n+5$, just as completely. Here the number 2 belongs either to $B$ or to $D$. We readily see that in the former case $\frac{1}{2}(p-1)$ belongs to $B$ and $\frac{1}{2}(p+1)$ to $D$, whereas in the latter $\frac{1}{2}(p-1)$ belongs to $D$ and $\frac{1}{2}(p+1)$ to $B$. Now consider that, if $\beta$ is a number from $B$ followed by a number from $D$, then $p-\beta-1$ will also be a number from $B$ and $p-\beta$ from $D$; that is, numbers with this property always occur in pairs. Their number, (13), will therefore be even, except when one of them is paired with itself, that is, when $\frac{1}{2}(p-1)$ belongs to $B$ and $\frac{1}{2}(p+1)$ to $D$; in this case (13) will be odd. From this we conclude that (13) is even whenever 2 belongs to $D$, and odd whenever 2 belongs to $B$. But we have

$$16(13) = aa + bb + 2a + 4b + 1$$

or, putting $a = 4q+1$, $b = 4r+2$,

$$(13) = qq + q + rr + 2r + 1$$

Thus (13) will be odd whenever $r$ is even; conversely, (13) will be even whenever $r$ is odd. From this we conclude that 2 belongs to $B$ whenever $b$ has the form $8m+2$, and to $D$ whenever $b$ has the form $8m+6$.

The result of these investigations can be stated as follows:

*The number* 2 *belongs to the collection* $A, B, C$ *or* $D$ *according as the number* $\frac{1}{2}b$ *has the form* $4m$, $4m+1$, $4m+2$ *or* $4m+3$.

**22.**

In the *Disquisitiones Arithmeticae* we explained the general theory of circle division and of the solution of the equation $x^p - 1 = 0$. Among other things, we showed that, if $\mu$ is a divisor of the number $p-1$, the function $\frac{x^p-1}{x-1}$ can be split into $\mu$ factors of degree $\frac{p-1}{\mu}$ with the help of an auxiliary equation of degree $\mu$. Besides the general theory of this factorization, we separately considered the special cases where $\mu = 2$ or $\mu = 3$ in Articles 356–358 of that work, and showed how to give the auxiliary equation a priori, that is, without working out the scheme of least residues of the powers of a primitive root for the modulus $p$. Even without our pointing it out, attentive readers will readily see the very close connection between the next case of that theory, namely $\mu = 4$, and the investigations developed here in Articles 15–20, with whose help that case too can be treated completely without difficulty. But we reserve this treatment for another occasion, and accordingly have preferred to carry out the investigation in this paper in a purely arithmetical form, without bringing in the theory of the equation $x^p - 1 = 0$ at all. Instead, as a finishing touch, we shall add some further new, purely arithmetical theorems very closely connected with the subject treated so far.

**23.**

If the power $(x^4+1)^{\frac{1}{2}(p-1)}$ is expanded by the binomial theorem, there will be three terms in which the exponent of $x$ is divisible by $p-1$, namely

$$x^{2(p-1)}, \quad Px^{p-1} \quad \text{and } 1$$

where $P$ denotes the middle coefficient

$$\frac{\frac{1}{2}(p-1) \,.\, \frac{1}{2}(p-3) \,.\, \frac{1}{2}(p-5) \,.\, \ldots \,.\, \frac{1}{4}(p+3)}{1 \,.\, 2 \,.\, 3 \,.\, \ldots \,.\, \frac{1}{4}(p-1)}$$

Thus, substituting for $x$ the numbers $1, 2, 3 \ldots p-1$ in turn, we obtain by the lemma in Article 19

$$\Sigma(x^4+1)^{\frac{1}{2}(p-1)} \equiv -2 - P$$

But, considering what we explained in Article 19, and also that the numbers in the collections $A, B, C, D$, raised to the power with exponent $\frac{1}{2}(p-1)$, are congruent modulo $p$ to the numbers $+1, -1, +1, -1$, respectively, we readily see that

$$\Sigma(x^4+1)^{\frac{1}{2}(p-1)} \equiv 4(00) - 4(01) + 4(02) - 4(03)$$

and hence, by the schemes given at the ends of Articles 18, 20,

$$\Sigma(x^4+1)^{\frac{1}{2}(p-1)} \equiv -2a - 2$$

Comparing these two values gives a particularly elegant theorem: we have

$$P \equiv 2a \pmod{p}$$

Denoting the four products

$$\begin{array}{l} 1 \,.\, 2 \,.\, 3 \,.\, \ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots \,.\, \frac{1}{4}(p-1) \\ \frac{1}{4}(p+3) \,.\, \frac{1}{4}(p+7) \,.\, \frac{1}{4}(p+11) \,.\, \ldots\ldots \,.\, \frac{1}{2}(p-1) \\ \frac{1}{2}(p+1) \,.\, \frac{1}{2}(p+3) \,.\, \frac{1}{2}(p+5) \,.\, \ldots\ldots \,.\, \frac{3}{4}(p-1) \\ \frac{1}{4}(3p+1) \,.\, \frac{1}{4}(3p+5) \,.\, \frac{1}{4}(3p+9) \,.\, \ldots\ldots \,.\, (p-1) \end{array}$$

by $q, r, s, t$, respectively, we can write the preceding theorem as

$$2a \equiv \frac{r}{q} \pmod{p}$$

Since each factor of $q$ has its complement to $p$ in $t$, we have $q \equiv t \pmod{p}$ whenever the number of factors is even, that is, whenever $p$ has the form $8m+1$, and $q \equiv -t$ whenever the number of factors is odd, or $p$ has the form $8m+5$. Likewise, in the former case $r \equiv s$, and in the latter $r \equiv -s$. In both cases $qr \equiv st$, and since we know that $qrst \equiv -1$, we have $qqrr \equiv -1$, and hence $qr \equiv \pm f \pmod{p}$. Combining this congruence with the theorem just found gives $rr \equiv \pm 2af$, and thus, by Articles 19, 20,

$$2b \equiv \pm rr \pmod{p}^{*)}$$

It is very remarkable that the splitting of the number $p$ into two squares can be found by entirely direct operations: the root of the odd square will be the least absolute residue of $\frac{r}{2q}$, and the root of the even square the least absolute residue of $\frac{1}{2}rr$ modulo $p$. The expression $\frac{r}{2q}$, whose value for $p = 5$ is $= 1$, can also be written as follows for larger values of $p$:

$$\frac{6 \,.\, 10 \,.\, 14 \,.\, 18 \,.\, \ldots \,.\, (p-3)}{2 \,.\, 3 \,.\, 4 \,.\, 5 \,.\, \ldots \,.\, \frac{1}{4}(p-1)}$$

But since we also know the sign with which this formula gives the root of the odd square, namely the sign that always makes it of the form $4m+1$, it deserves particular attention that a similar general criterion for the sign of the root of the even square has not yet been found. If anyone finds such a criterion and communicates it to us, we shall be very grateful. Meanwhile, it seemed appropriate to include here the values of the numbers $a, b, f$ obtained, for values of $p$ below 200, from the least residues of the expressions $\frac{r}{2q}$, $\frac{1}{2}rr$, $qr$.

*) and $\{(a \mp b)q\}^2 \equiv a \equiv \left(\frac{r-qrr}{2}\right)^2$

| $p$ | $a$ | $b$ | $f$ |
|---|---|---|---|
| 5 | +1 | +2 | 2 |
| 13 | −3 | −2 | 5 |
| 17 | +1 | −4 | 13 |
| 29 | +5 | +2 | 12 |
| 37 | +1 | −6 | 31 |
| 41 | +5 | +4 | 9 |
| 53 | −7 | −2 | 23 |
| 61 | +5 | −6 | 11 |
| 73 | −3 | −8 | 27 |
| 89 | +5 | −8 | 34 |
| 97 | +9 | +4 | 22 |
| 101 | +1 | −10 | 91 |
| 109 | −3 | +10 | 33 |
| 113 | −7 | +8 | 15 |
| 137 | −11 | +4 | 37 |
| 149 | −7 | −10 | 44 |
| 157 | −11 | −6 | 129 |
| 173 | +13 | +2 | 80 |
| 181 | +9 | +10 | 162 |
| 193 | −7 | +12 | 81 |
| 197 | +1 | −14 | 183 |

# THEORY

# OF BIQUADRATIC RESIDUES

## SECOND PAPER

BY

CARL FRIEDRICH GAUSS

PRESENTED TO THE ROYAL SOCIETY ON APRIL 15, 1831.

---

Commentationes societatis regiae scientiarum Gottingensis recentiores. Vol. VII.
Göttingen MDCCCXXXII.

---

# THEORY OF BIQUADRATIC RESIDUES.

SECOND PAPER.

---

## 24.

In the first paper we completely dealt with everything required for the biquadratic classification of the number $+2$. Namely, we think of all numbers not divisible by the modulus $p$ (assumed to be a prime number of the form $4n+1$) as distributed among four collections $A, B, C, D$, according as each, raised to the power with exponent $\frac{1}{4}(p-1)$, becomes congruent modulo $p$ to $+1$, $+f$, $-1$, $-f$, with $f$ denoting either root of the congruence $ff \equiv -1 \pmod{p}$. We found that the decision as to which collection the number $+2$ belongs to depends on splitting the number $p$ into two squares: if we put $p = aa + bb$, with $aa$ denoting the odd square and $bb$ the even one, and further assume that the *signs* of $a, b$ are chosen so that $a \equiv 1 \pmod{4}$, $b \equiv af \pmod{p}$, then the number $+2$ must belong to the collection $A, B, C, D$ according as $\frac{1}{2}b$ has the form $4n$, $4n+1$, $4n+2$, $4n+3$, respectively.

A rule for classifying the number $-2$ also follows immediately. Since $-1$ belongs to class $A$ when $\frac{1}{2}b$ is even, and to class $C$ when it is odd, the theorem in Article 7 shows that the number $-2$ belongs to class $A, B, C, D$ according as $\frac{1}{2}b$ has the form $4n$, $4n+3$, $4n+2$, $4n+1$, respectively.

These theorems can also be expressed as follows:

| Belongs | $+2$ | $-2$ |
|---|---|---|
| to the collection | if $b$, modulo 8, is congruent to | |
| $A$ | 0 | 0 |
| $B$ | $2a$ | $6a$ |
| $C$ | $4a$ | $4a$ |
| $D$ | $6a$ | $2a$ |

We readily see that, stated in this way, the theorems no longer depend on the condition $a \equiv 1 \pmod{4}$, but still hold if $a \equiv 3 \pmod{4}$, provided that the other condition, $af \equiv b \pmod{p}$, is maintained.

It is equally easy to see that the content of these theorems can be elegantly condensed into a single formula, namely:

*if $a$ and $b$ are taken positive, we always have*

$$b^{\frac{1}{2}ab} \equiv a^{\frac{1}{2}ab}\, 2^{\frac{1}{4}(p-1)} \pmod{p}$$

**25.**

Let us now see how far inductive investigation points towards the classification of the number 3. The table in Article 11, continued further (always taking the least primitive root), shows that +3 belongs

to the collection

| $A$ for | | | $B$ for | | | $C$ for | | | $D$ for | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| $p$ | $a$ | $b$ | $p$ | $a$ | $b$ | $p$ | $a$ | $b$ | $p$ | $a$ | $b$ |
| 13 | −3 | +2 | 17 | +1 | −4 | 37 | +1 | −6 | 5 | +1 | +2 |
| 109 | −3 | +10 | 29 | +5 | +2 | 61 | +5 | −6 | 41 | +5 | −4 |
| 181 | +9 | +10 | 53 | −7 | +2 | 73 | −3 | −8 | 149 | −7 | +10 |
| 193 | −7 | −12 | 89 | +5 | −8 | 97 | +9 | +4 | 173 | +13 | +2 |
| 229 | −15 | +2 | 101 | +1 | +10 | 157 | −11 | −6 | | | |
| 277 | +9 | +14 | 113 | −7 | −8 | 241 | −15 | −4 | | | |
| | | | 137 | −11 | −4 | | | | | | |
| | | | 197 | +1 | −14 | | | | | | |
| | | | 233 | +13 | +8 | | | | | | |
| | | | 257 | +1 | −16 | | | | | | |
| | | | 269 | +13 | +10 | | | | | | |
| | | | 281 | +5 | +16 | | | | | | |
| | | | 293 | +17 | +2 | | | | | | |

At least at first sight, we see no simple connection between the values of the numbers $a, b$ to which the same collection corresponds. But, considering that a similar decision in the theory of quadratic residues is made by a simpler rule for the number −3 than for the number +3, we may hope for equally favourable success in the theory of biquadratic residues. We find that −3 belongs to the collection

| $A$ for | | | $B$ for | | | $C$ for | | | $D$ for | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| $p$ | $a$ | $b$ | $p$ | $a$ | $b$ | $p$ | $a$ | $b$ | $p$ | $a$ | $b$ |
| 37 | +1 | −6 | 5 | +1 | +2 | 13 | −3 | +2 | 29 | +5 | +2 |
| 61 | +5 | −6 | 17 | +1 | −4 | 73 | −3 | −8 | 41 | +5 | −4 |
| 157 | −11 | −6 | 89 | +5 | −8 | 97 | +9 | +4 | 53 | −7 | +2 |
| 193 | −7 | −12 | 113 | −7 | −8 | 109 | −3 | +10 | 101 | +1 | +10 |
| | | | 137 | −11 | −4 | 181 | +9 | +10 | 197 | +1 | −14 |
| | | | 149 | −7 | +10 | 229 | −15 | +2 | 269 | +13 | +10 |
| | | | 173 | +13 | +2 | 241 | −15 | −4 | 293 | +17 | +2 |
| | | | 233 | +13 | +8 | 277 | +9 | +14 | | | |
| | | | 257 | +1 | −16 | | | | | | |
| | | | 281 | +5 | +16 | | | | | | |

where the law suggested by inductive investigation presents itself at once. Namely, $-3$ belongs to the collection

$A$, whenever $b$ is divisible by 3, equivalently $b \equiv 0 \pmod 3$
$B$, whenever $a + b$ is divisible by 3, equivalently $b \equiv 2a \pmod 3$
$C$, whenever $a$ is divisible by 3, equivalently $a \equiv 0 \pmod 3$
$D$, whenever $a - b$ is divisible by 3, equivalently $b \equiv a \pmod 3$

**26.**

We find that the number $+5$ is to be assigned to the collection

| | |
|---|---|
| $A$ for $p =$ | 101, 109, 149, 181, 269 |
| $B$ for $p =$ | 13, 17, 73, 97, 157, 193, 197, 233, 277, 293 |
| $C$ for $p =$ | 29, 41, 61, 89, 229, 241, 281 |
| $D$ for $p =$ | 37, 53, 113, 137, 173, 257 |

Considering the values of the numbers $a, b$ corresponding to each $p$, we find the law here as readily as for the classification of the number $-3$. Namely, we arrive at the collection

$A$, whenever $b \equiv 0 \pmod 5$
$B$, whenever $b \equiv a$
$C$, whenever $a \equiv 0$
$D$, whenever $b \equiv 4a$

These rules evidently cover all cases, since for $b \equiv 2a$ or $b \equiv 3a \pmod 5$ we would have $aa + bb \equiv 0$, Q. E. A., since by hypothesis $p$ is a prime number different from 5.

**27.**

Likewise, inductive investigation applied to the numbers $-7$, $-11$, $+13$, $+17$, $-19$, $-23$ and carried far enough points to the following rules:

For the number $-7$

| | |
|---|---|
| $A$ | $a \equiv 0$, or $b \equiv 0 \pmod 7$ |
| $B$ | $b \equiv 4a$, or $b \equiv 5a$ |
| $C$ | $b \equiv a$, or $b \equiv 6a$ |
| $D$ | $b \equiv 2a$, or $b \equiv 3a$ |

For the number $-11$.

| | |
|---|---|
| $A$ | $b \equiv 0,\ 5a,$ or $6a \pmod{11}$ |
| $B$ | $b \equiv a,\ 3a$ or $4a$ |
| $C$ | $a \equiv 0,$ or $b \equiv 2a$ or $9a$ |
| $D$ | $b \equiv 7a,\ 8a$ or $10a$ |

For the number $+13$.

| | |
|---|---|
| $A$ | $b \equiv 0,\ 4a,\ 9a \pmod{13}$ |
| $B$ | $b \equiv 6a,\ 11a,\ 12a$ |
| $C$ | $a \equiv 0;\ b \equiv 3a,\ 10a$ |
| $D$ | $b \equiv a,\ 2a,\ 7a$ |

For the number $+17$.

| | |
|---|---|
| $A$ | $a \equiv 0;\ b \equiv 0,\ a,\ 16a \pmod{17}$ |
| $B$ | $b \equiv 2a,\ 6a,\ 8a,\ 14a$ |
| $C$ | $b \equiv 5a,\ 7a,\ 10a,\ 12a$ |
| $D$ | $b \equiv 3a,\ 9a,\ 11a,\ 15a$ |

For the number $-19$.

| | |
|---|---|
| $A$ | $b \equiv 0,\ 2a,\ 5a,\ 14a,\ 17a \pmod{19}$ |
| $B$ | $b \equiv 3a,\ 7a,\ 11a,\ 13a,\ 18a$ |
| $C$ | $a \equiv 0;\ b \equiv 4a,\ 9a,\ 10a,\ 15a$ |
| $D$ | $b \equiv a,\ 6a,\ 8a,\ 12a,\ 16a$ |

For the number $-23$.

| | |
|---|---|
| $A$ | $a \equiv 0;\ b \equiv 0,\ 7a,\ 10a,\ 13a,\ 16a \pmod{23}$ |
| $B$ | $b \equiv 2a,\ 3a,\ 4a,\ 11a,\ 15a,\ 17a$ |
| $C$ | $b \equiv a,\ 5a,\ 9a,\ 14a,\ 18a,\ 22a$ |
| $D$ | $b \equiv 6a,\ 8a,\ 12a,\ 19a,\ 20a,\ 21a$ |

**28.**

The special theorems obtained in this way by inductive investigation are found to be confirmed as far as that investigation is continued, and reveal a particularly beautiful form of the criteria. But when they are compared with one another to draw general conclusions, the following observations immediately present themselves.

The criteria for deciding which collection the prime number $\pm q$ belongs to (taking the upper or lower sign according as $q$ has the form $4n+1$ or $4n+3$) depend on the forms of the numbers $a, b$ compared with one another with respect to the modulus $q$. Namely,

I. Whenever $a \equiv 0 \pmod{q}$, $\pm q$ belongs to a definite collection, which is $A$ for $q = 7,\ 17,\ 23$, and $C$ for $q = 3,\ 11,\ 13,\ 19$. This suggests the conjecture that the former case holds generally whenever $q$ has the form $8n \pm 1$, and the latter whenever $q$ has the form $8n \pm 3$. Moreover, the collections $B$ and $D$ can already be excluded without inductive investigation when the value of $a$ is divisible by $q$, in which case $p \equiv bb \pmod{q}$, that is, $p$ is a quadratic residue of $q$; hence, by the fundamental theorem, $\pm q$ must be a quadratic residue of $p$.

II. When $a$ is not divisible by $q$, the criterion depends on the value of the expression $\frac{b}{a} \pmod{q}$. This expression does admit $q$ different values, namely $0,\ 1,\ 2,\ 3 \ldots . q - 1$; but whenever $q$ has the form $4n + 1$, the two values of the expression $\sqrt{-1} \pmod{q}$ must be excluded. They evidently cannot be values of the expression $\frac{b}{a} \pmod{q}$, since $p = aa + bb$ is always assumed to be a prime number different from $q$. The number of admissible values of the expression $\frac{b}{a} \pmod{q}$ is therefore $= q - 2$ for $q \equiv 1 \pmod{4}$, while it remains $= q$ for $q \equiv 3 \pmod{4}$.

These values are now distributed among four classes: some, to be denoted without further specification by $\alpha$, correspond to the collection $A$; others, denoted by $\beta$, to the collection $B$; others, $\gamma$, to the collection $C$; and finally the rest, $\delta$, to the collection $D$. Thus $\pm q$ is to be assigned to the collection $A, B, C, D$ according as $b \equiv \alpha a$, $b \equiv \beta a$, $b \equiv \gamma a$, $b \equiv \delta a \pmod{q}$.

But the *law* of this distribution seems more deeply hidden, although some general features are readily noticed. The number of values in three of the classes is found to be the same, namely $= \frac{1}{4}(q - 1)$ or $\frac{1}{4}(q + 1)$, while in one (specifically, the one corresponding to the collection with the criterion $a \equiv 0$) it is smaller by one. Thus the total number of different criteria for each collection is the same, namely $= \frac{1}{4}(q - 1)$ or $\frac{1}{4}(q + 1)$. We further observe that 0 is always found in the first class (among the $\alpha$), and that the complements of the numbers $\alpha, \beta, \gamma, \delta$ to $q$, namely $q - \alpha$, $q - \beta$, $q - \gamma$, $q - \delta$, are found in the first, fourth, third and second classes, respectively. Finally, we see that the values of the expressions $\frac{1}{\alpha}$, $\frac{1}{\beta}$, $\frac{1}{\gamma}$, $\frac{1}{\delta} \pmod{q}$ belong to the first, fourth, third and second classes whenever the criterion $a \equiv 0$ corresponds to the collection $A$, but to the third, second, first and fourth classes, respectively, whenever the criterion $a \equiv 0$ refers to the collection $C$. But this is about the limit of what can be reached by inductive investigation, unless we rather boldly presume to anticipate what will later be drawn from the genuine sources.

**29.**

Before going further, we should note that the criteria for prime numbers (taken positively when they have the form $4n + 1$, negatively when they have the form $4n + 3$) suffice to decide the question for all other numbers, provided that the theorem of Article 7 and the criteria for $-1$ and $\pm 2$ are also used.

Thus, for example, if criteria for the number $+3$ are wanted, the criteria given in Article 25, which concern $-3$, will also hold for $+3$ whenever $\frac{1}{2}b$ is even. The collections $A, B, C, D$ must instead be exchanged with the collections $C, D, A, B$ whenever $\frac{1}{2}b$ is odd, giving the following rules:

$+3$ belongs

| to the collection | if |
|---|---|
| $A$ | $b \equiv 0 \pmod{12}$; or simultaneously $a \equiv 0 \pmod 3,\ b \equiv 2 \pmod 4$ |
| $B$ | $b \equiv 8a$ or $10a \pmod{12}$ |
| $C$ | $b \equiv 6a \pmod{12}$; or simultaneously $a \equiv 0 \pmod 3,\ b \equiv 0 \pmod 4$ |
| $D$ | $b \equiv 2a$ or $4a \pmod{12}$ |

Likewise, the criteria for $\pm 6$ are obtained by combining the criteria for $\mp 2$ and $-3$; namely,

$+6$ belongs

| to the collection | if |
|---|---|
| $A$ | $b \equiv 0,\quad 2a, 22a \pmod{24}$; or simultaneously $a \equiv 0 \pmod 3,\ b \equiv 4a \pmod 8$ |
| $B$ | $b \equiv 4a,\quad 6a,\quad 8a \pmod{24}$; or simultaneously $a \equiv 0 \pmod 3,\ b \equiv 2a \pmod 8$ |
| $C$ | $b \equiv 10a, 12a, 14a \pmod{24}$; or simultaneously $a \equiv 0 \pmod 3,\ b \equiv 0 \pmod 8$ |
| $D$ | $b \equiv 16a, 18a, 20a \pmod{24}$; or simultaneously $a \equiv 0 \pmod 3,\ b \equiv 6a \pmod 8$ |

while $-6$ belongs

| to the collection | if |
|---|---|
| $A$ | $b \equiv 0,\quad 10a, 14a \pmod{24}$; or simultaneously $a \equiv 0 \pmod 3,\ b \equiv 4a \pmod 8$ |
| $B$ | $b \equiv 4a,\quad 8a, 18a \pmod{24}$; or simultaneously $a \equiv 0 \pmod 3,\ b \equiv 6a \pmod 8$ |
| $C$ | $b \equiv 2a, 12a, 22a \pmod{24}$; or simultaneously $a \equiv 0 \pmod 3,\ b \equiv 0 \pmod 8$ |
| $D$ | $b \equiv 6a, 16a, 20a \pmod{24}$; or simultaneously $a \equiv 0 \pmod 3,\ b \equiv 2a \pmod 8$ |

In the same way, the criteria for the number $+21$ will be assembled from the criteria for $-3$ and $-7$; the criteria for $-105$ from those for $-1$, $-3$, $+5$, $-7$, etc.

**30.**

Inductive investigation thus opens up a very rich harvest of particular theorems related to the theorem for the number 2. But a common bond is lacking, rigorous proofs are lacking, since the method by which we completed the treatment of the number 2 in the first paper admits no further application. There are indeed various methods by which proofs could be obtained for particular cases, especially those concerning the distribution of quadratic residues between the collections $A, C$. We do not dwell on them, however, since a general theory embracing *all* cases must be our aim. Having begun to devote our thoughts to this matter in 1805, we soon became convinced that the true source of the general theory must be sought in an extension of the domain of arithmetic, as we already indicated in Article 1.

While higher arithmetic, in the questions treated so far, deals only with real integers, the theorems on biquadratic residues shine in their full simplicity and genuine beauty only when the domain of arithmetic is extended to *imaginary* quantities. Its objects are then, without restriction, numbers of the form $a + bi$, where $i$ denotes, as usual, the imaginary quantity $\sqrt{-1}$, and $a, b$ range without restriction over all real integers between $-\infty$ and $+\infty$. We shall call such numbers *complex integers*, treating real numbers as a species contained within the complex numbers rather than as opposed to them. The present paper will set out both the elementary theory of complex numbers and the first beginnings of the theory of biquadratic residues, which we shall undertake to complete in every respect in a subsequent continuation[*)].

[*)] We should at least mention here in passing that the domain thus defined is particularly suited to the theory of biquadratic residues. The theory of cubic residues must similarly be built on the consideration of numbers of the form $a + bh$, where $h$ is an imaginary root of the equation $h^3 - 1 = 0$, say $h = -\frac{1}{2} + \sqrt{\frac{3}{4}} \,.\, i$; and likewise the theory of residues of higher powers will require the introduction of other imaginary quantities.

**31.**

First we introduce some names that will help us to be brief and clear. The domain of complex numbers $a + bi$ contains

I. real numbers, where $b = 0$, and among these, according to the nature of $a$,

1) zero,

2) positive numbers,

3) negative numbers;

II. imaginary numbers, where $b$ is unequal to zero. Here we distinguish in turn

1) imaginary numbers without a real part, that is, where $a = 0$;

2) imaginary numbers with a real part, where $b$ is nonzero and $a = 0$ also does not hold.

The former may, if desired, be called pure imaginary numbers, and the latter mixed imaginary numbers.

In this theory we use four units, $+1$, $-1$, $+i$, $-i$, which will be called simply positive, negative, positive imaginary and negative imaginary.

We shall call the three products of any complex number by $-1$, $+i$, $-i$ its *companions* or *numbers associated with it*. Thus, except for zero (which is associated with itself), four *distinct* numbers are always associated.

By contrast, we call the number obtained from a complex number by exchanging $i$ with $-i$ its *conjugate*. Among imaginary numbers, therefore, two *distinct* numbers are always conjugate, while real numbers are their own conjugates, if we choose to extend the name to them.

We call the product of a complex number and its conjugate the *norm* of each. The square of a real number is therefore to be taken as its norm.

In general we have eight related numbers, namely,

$$\begin{array}{r|r} a+bi & a-bi \\ -b+ai & -b-ai \\ -a-bi & -a+bi \\ b-ai & b+ai \end{array}$$

in which we see two groups of four associated numbers and four pairs of conjugates; their common norm is $aa + bb$. But the eight numbers reduce to four distinct ones whenever either $a = \pm b$, or $a, b = 0$ for one of the two numbers.

The following results follow immediately from these definitions:

The conjugate of the product of two complex numbers is the product of their conjugates.

The same holds for a product of more factors, and also for quotients.

The norm of the product of two complex numbers is equal to the product of their norms.

This theorem also extends to products of any number of factors and to quotients.

The norm of any complex number (except zero, an exception that from now on we shall generally leave understood) is a *positive* number.

Moreover, nothing prevents us from extending our definitions to fractional or even irrational values of $a, b$; but $a + bi$ will be called a complex integer only when *each* of $a, b$ is an integer, and rational only when *each* of $a, b$ is rational.

**32.**

The rules for arithmetic operations on complex numbers are well known: introducing the norm reduces division to multiplication, since

$$\frac{a+bi}{c+di} = (a+bi)\,.\,\frac{c-di}{cc+dd} = \frac{ac+bd}{cc+dd} + \frac{bc-ad}{cc+dd}\,.\,i$$

The square root is extracted with the help of the formula

$$\surd(a+bi) = \pm\left(\surd\frac{\surd(aa+bb)+a}{2} + i\surd\frac{\surd(aa+bb)-a}{2}\right)$$

if $b$ is positive, or of the formula

$$\surd(a+bi) = \pm\left(\surd\frac{\surd(aa+bb)+a}{2} - i\surd\frac{\surd(aa+bb)-a}{2}\right)$$

if $b$ is negative. There is no need to dwell here on how transforming the complex quantity $a + bi$ into $r(\cos\varphi + i\sin\varphi)$ makes calculations easier.

**33.**

We call a complex integer that can be resolved into two factors different from the units[*)] a composite complex number; a number that admits no such factorization will be called a complex prime. It follows immediately that every real composite number is also a composite complex number. But a real prime may be a composite complex number, and this holds for the number 2 and all positive real primes of the form $4n+1$ (except the number 1), since they are known to be decomposable into two positive squares. For example, $2 = (1+i)(1-i)$, $5 = (1+2i)(1-2i)$, $13 = (3+2i)(3-2i)$, $17 = (1+4i)(1-4i)$, etc.

On the other hand, positive real primes of the form $4n+3$ are always complex primes. For if such a number $q$ were $= (a+bi)(\alpha+\beta i)$, we should also have $q = (a-bi)(\alpha-\beta i)$, and therefore $qq = (aa+bb)(\alpha\alpha+\beta\beta)$. But $qq$ can be resolved into positive factors greater than unity in only one way, namely into $q \times q$, so we should have to have $q = aa+bb = \alpha\alpha+\beta\beta$, Q. E. A.; for the sum of two squares cannot have the form $4n+3$.

Negative real numbers evidently retain the same names as positive ones, and the same holds for pure imaginary numbers.

It therefore remains to show how to distinguish composites from primes among mixed imaginary numbers. This is done by the following

THEOREM. *Every mixed imaginary integer $a+bi$ is a complex prime or a composite number according as its norm is a real prime or a composite number.*

*Proof.* I. Since the norm of a composite complex number is always composite, a complex number whose norm is a real prime must evidently be a complex prime. Q. E. P.

II. If, however, the norm $aa+bb$ is composite, let $p$ be a positive real prime dividing it. We must now distinguish two cases.

1) If $p$ has the form $4n+3$, we know that $aa+bb$ cannot be divisible by $p$ unless $p$ divides both $a, b$, so that $a+bi$ will be composite.

[*)] Or, equivalently, factors whose norms are greater than unity.

2) If $p$ does not have the form $4n+3$, it can certainly be decomposed into two squares. We therefore put $p = \alpha\alpha + \beta\beta$. Since

$$(a\alpha + b\beta)(a\alpha - b\beta) = aa(\alpha\alpha + \beta\beta) - \beta\beta(aa + bb)$$

and this is therefore divisible by $p$, $p$ must divide one of the factors $a\alpha + b\beta$, $a\alpha - b\beta$. Moreover, since

$$(a\alpha + b\beta)^2 + (b\alpha - a\beta)^2 = (a\alpha - b\beta)^2 + (b\alpha + a\beta)^2 = (aa + bb)(\alpha\alpha + \beta\beta)$$

and this is therefore divisible by $pp$, in the first case $b\alpha - a\beta$, and in the second $b\alpha + a\beta$, must also be divisible by $p$. Thus in the first case

$$\frac{a + bi}{\alpha + \beta i} = \frac{a\alpha + b\beta}{p} + \frac{b\alpha - a\beta}{p} \,.\, i$$

will be a complex integer, while in the second

$$\frac{a + bi}{\alpha - \beta i} = \frac{a\alpha - b\beta}{p} + \frac{b\alpha + a\beta}{p} \,.\, i$$

will be an integer. Since the given number is therefore divisible by $\alpha + \beta i$ or by $\alpha - \beta i$, and the norm of the quotient $= \frac{aa+bb}{p}$ differs from unity by hypothesis, $a + bi$ is evidently a composite complex number in either case. Q. E. S.

**34.**

The following four kinds therefore exhaust the entire range of complex primes:

1) the four units, $1, +i, -1, -i$, which, however, we shall usually leave understood to be excluded when discussing prime numbers;

2) the number $1 + i$ with its three associates $-1 + i$, $-1 - i$, $1 - i$;

3) the positive real primes of the form $4n + 3$ with their three associates each;

4) the complex numbers whose norms are real primes of the form $4n + 1$ greater than unity. To each given norm of this kind there always correspond eight complex primes and no more, since such a norm can be decomposed into two squares in only one way.

**35.**

Just as real integers are divided into even and odd, and the former in turn into those divisible by four and those that are twice an odd number, an equally essential distinction presents itself among complex numbers. They are either

not divisible by $1+i$, namely numbers $a+bi$ in which one of the numbers $a, b$ is odd and the other even;

or divisible by $1+i$ but not by 2, whenever both $a, b$ are odd;

or divisible by 2, whenever both $a, b$ are even.

The numbers of the first class may suitably be called *odd complex numbers*, those of the second *half-even*, and those of the third *even*.

A product of several complex factors will always be odd when all the factors are odd; half-even when one factor is half-even and the others odd; and even when the factors include at least two half-even ones or at least one even one.

The norm of any odd complex number has the form $4n+1$; the norm of a half-even number has the form $8n+2$; finally, the norm of an even number is the product of a number of the form $4n+1$ and the number 4 or a higher power of two.

**36.**

Since the relation between four associated complex numbers is analogous to that between two opposite real numbers (that is, numbers equal in absolute value and bearing opposite signs), and the positive one of these is usually and rightly regarded as primary, the question arises whether a similar distinction can be established among four associated complex numbers, and whether it should be considered useful. To decide this, we must consider that the principle of distinction must be such that the product of two numbers regarded as primary among their respective associates always becomes primary among its own associates. But we soon become convinced that no such principle exists at all unless the distinction is restricted to integers; indeed, a *useful* distinction must be limited to odd numbers. For these, however, the proposed aim can be reached in two ways. Namely,

I. The product of two numbers $a+bi, a'+b'i$ such that $a, a'$ have the form $4n+1$ and $b, b'$ are even will have the same property: the real part will be $\equiv 1 \pmod 4$, and the imaginary part even. And it is easy to see that among four associated odd numbers only one has that form.

II. If a number $a+bi$ is such that $a-1$ and $b$ are either both divisible by four or both twice an odd number, its product with a complex number of the same form will have the same form; and it is easy to see that only one of four associated odd numbers has this form.

Of these two principles, which are almost equally suitable, we shall adopt the latter: among four associated odd complex numbers we shall regard as primary the one congruent to the positive unit modulo $2+2i$. This will let us state several notable theorems more concisely. For example, the complex primes $-1+2i$, $-1-2i$, $+3+2i$, $+3-2i$, $+1+4i$, $+1-4i$, etc., are primary, as are the real numbers $-3$, $-7$, $-11$, $-19$, etc., which must evidently always be taken with a negative sign. The conjugate of an odd primary complex number will also be primary.

For half-even and even numbers in general, a similar distinction would be too arbitrary and of little use. Of the associated primes $1+i$, $1-i$, $-1+i$, $-1-i$, we can indeed choose one as primary in preference to the others, but we shall not extend such a distinction to composites.

**37.**

If some factors of a composite complex number are themselves composite, and these in turn are resolved into their factors, we evidently arrive eventually at prime factors; that is, every composite number can be resolved into prime factors. If any of these are not primary, replace each by the product of its primary associate and $i, -1$ or $-i$. In this way, every composite complex number $M$ can evidently be reduced to the form

$$M = i^{\mu} A^{\alpha} B^{\beta} C^{\gamma} \dots .$$

where $A, B, C$, etc., are distinct primary complex primes, and $\mu = 0, 1, 2$ or 3. Concerning this factorization, we have the theorem that it can be carried out in only one way. On a passing consideration this theorem might seem evident in itself, but it certainly needs a proof. The following paves the way to that proof.

THEOREM. *The product $M = A^{\alpha} B^{\beta} C^{\gamma} \dots$, where $A, B, C$, etc., denote distinct primary complex primes, cannot be divisible by any primary complex prime that does not occur among $A, B, C$, etc.*

*Proof.* Let $P$ be a primary complex prime not contained among $A, B, C$, etc., and let $p, a, b, c$, etc., be the norms of the numbers $P, A, B, C$, etc. It follows readily that the norm of $M$ will be $= a^{\alpha} b^{\beta} c^{\gamma}$, etc.; hence, if $M$ were divisible by $P$, this number would have to be divisible by $p$. Since the individual norms are either real primes (from the sequence $2, 5, 13, 17$, etc.) or squares of real primes (from the sequence $9, 49, 121$, etc.), this evidently cannot happen unless $p$ is identical with one of the norms $a, b, c$, etc. We therefore suppose $p = a$. But since $P, A$ are distinct primary complex primes by hypothesis, it is easy to see that these conditions cannot hold together unless $P, A$ are conjugate imaginary complex numbers, and consequently $p = a$ is an odd real prime (not the square of a prime). We therefore suppose $A = k+li$, $P = k-li$.

Hence, extending the notion and notation of congruence to complex integers, we have $A \equiv 2k \pmod{P}$, from which it follows readily that

$$M \equiv 2^\alpha k^\alpha B^\beta C^\gamma \dots . \pmod{P}$$

Therefore, if $M$ is assumed divisible by $P$, then

$$2^\alpha k^\alpha B^\beta C^\gamma \dots .$$

will also be divisible by $P$, and consequently the norm of this number, which is

$$= 2^{2\alpha} k^{2\alpha} b^\beta c^\gamma \dots .$$

will be divisible by $p$. But since 2 and $k$ are certainly not divisible by $p$, it follows that $p$ must be identical with one of the numbers $b, c$, etc.; let, for example, $p = b$. From this, however, we conclude that either $B = k+li$ or $B = k-li$, that is, either $B = A$ or $B = P$, both contrary to the hypothesis.

The other theorem, that factorization into primes can be carried out in only one way, follows very easily from this theorem, by reasoning entirely analogous to that which we used for real numbers in the *Disquisitiones Arithmeticae* (Article 16). It would therefore be superfluous to dwell on it here.

**38.**

We now turn to the congruence of numbers with respect to complex moduli. At the outset of this investigation, however, we should indicate how the domain of complex quantities can be made visible to intuition.

Every real quantity can be expressed by a segment of a line infinite in both directions, taken from an arbitrary starting point and measured against an arbitrary segment adopted as the unit. It can thus be represented by the other endpoint, with points on one side of the starting point representing positive quantities and those on the other negative quantities. In the same way, any complex quantity can be represented by a point in an infinite plane in which a fixed line corresponds to the real quantities: the complex quantity $x+iy$ by the point with abscissa $= x$ and ordinate, taken positively on one side of the line of abscissas and negatively on the other, $= y$. In this way we may say that any complex quantity measures the difference between the position of the point to which it corresponds and that of the starting point. The positive unit denotes an arbitrary fixed displacement in an arbitrary fixed direction; the negative unit an equally large displacement in the opposite direction; and the imaginary units equally large displacements in the two perpendicular lateral directions.

This sheds considerable light on the metaphysics of the quantities we call imaginary. If the starting point is denoted by (0), and two complex quantities $m, m'$ correspond to points $M, M'$, whose positions relative to (0) they

express, the difference $m-m'$ is nothing other than the position of $M$ relative to $M'$. On the other hand, if the product $mm'$ represents the position of $N$ relative to (0), it is easy to see that this position is determined by the position of $M$ relative to (0) in the same way as the position of $M'$ is determined by the position of the point corresponding to the positive unit. Thus we may aptly say that the positions of the points corresponding to the complex quantities $mm'$ $m, m', 1$ form a *proportion*. But we reserve a fuller treatment of this matter for another occasion. The difficulties in which the theory of imaginary quantities is thought to be entangled have largely arisen from unsuitable names (some have even used the discordant name of impossible quantities). Had we started from the concepts offered by two-dimensional manifolds, as seen in their greatest purity in our intuitions of space, and called positive quantities direct, negative quantities inverse and imaginary quantities lateral, simplicity would have taken the place of complications, and clarity the place of darkness.

**39.**

What was set out in the preceding article concerns continuous complex quantities. In arithmetic, which deals only with integers, the representation of complex numbers will be a system of equally spaced points arranged on equally spaced lines so as to divide the infinite plane into infinitely many equal squares. All numbers divisible by a given complex number $a + bi = m$ will likewise form infinitely many squares with sides $= \sqrt{(aa + bb)}$, or areas $= aa + bb$. The latter squares will be inclined to the former whenever neither of the numbers $a, b$ is $= 0$. Each number not divisible by the modulus $m$ will correspond to a point either inside such a square or on a side shared by two squares; the latter case, however, cannot occur unless $a, b$ have a common divisor. It is also clear that numbers congruent modulo $m$ occupy corresponding positions in their squares. Hence, if we collect all the numbers lying inside a specified square, together with all those that may lie on two of its non-opposite sides, and finally add a number divisible by $m$, we readily obtain a complete system of incongruent residues modulo $m$: every integer must be congruent to one of them, and to only one. Nor would it be difficult to show that the number of these residues equals the norm of the modulus, namely $= aa + bb$. But it seems advisable to prove this very important theorem in another, purely arithmetical way.

**40.**

THEOREM. *For a given complex modulus $m = a + bi$, whose norm is $aa + bb = p$ and for which $a, b$ are relatively prime, every complex integer will be congruent to a residue in the sequence $0, 1, 2, 3 \ldots p - 1$, and to no more than one.*

*Proof.* I. Let $\alpha, \beta$ be integers such that $\alpha a + \beta b = 1$. Then

$$i = \alpha b - \beta a + m(\beta + \alpha i)$$

Thus, for a given complex integer $A + Bi$, we have

$$A + Bi = A + (\alpha b - \beta a)B + m(\beta B + \alpha B i)$$

Denote by $h$ the least positive residue of $A + (\alpha b - \beta a)B$ modulo $p$, and put

$$A + (\alpha b - \beta a)B = h + kp = h + m(ak - bki)$$

Then

$$A + Bi = h + m(\beta B + ak + (\alpha B - bk)i)$$

or

$$A + Bi \equiv h (\text{mod. } m). \qquad \text{Q. E. P.}$$

II. Whenever two real numbers $h, h'$ are congruent to the same complex number modulo $m$, they are also congruent to each other. We therefore put $h - h' = m(c + di)$, giving

$$(h - h')(a - bi) = p(c + di)$$

and hence

$$(h - h')a = pc, \quad (h - h')b = -pd$$

and also, since $a\alpha + b\beta = 1$,

$$h - h' = p(c\alpha - d\beta), \text{ i. e. } h \equiv h' (\text{mod. } p)$$

Thus $h$ and $h'$, if distinct, cannot both belong to the collection of numbers $0, 1, 2, 3 \ldots . p - 1$. Q. E. S.

**41.**

THEOREM. *For a complex modulus $m = .a+bi$, whose norm is $aa+bb = p$ and for which $a, b$ are not relatively prime but have greatest common divisor $\lambda$ (which we suppose taken positively), every complex number is congruent to a residue $x + yi$ such that $x$ is one of the numbers $0, 1, 2, 3 \ldots . \frac{p}{\lambda} - 1$ and $y$ one of the numbers $0, 1, 2, 3 \ldots . \lambda - 1$, and to exactly one of all the $p$ residues of this form.*

*Proof.* I. Taking integers $\alpha, \beta$ such that $\alpha a + \beta b = \lambda$, we have $\lambda i = \alpha b - \beta a + m(\beta + \alpha i)$. Now let $A + Bi$ be a given complex number, $y$ the least positive residue of $B$ modulo $\lambda$, and $x$ the least positive residue of $A + (\alpha b - \beta a) . \frac{B-y}{\lambda}$ modulo $\frac{p}{\lambda}$, and put

$$A + (\alpha b - \beta a) . \frac{B-y}{\lambda} = x + \frac{p}{\lambda} . k$$

Then

$$\begin{aligned} A + Bi - (x + yi) &= \frac{p}{\lambda} . k + (B - y)i - (\alpha b - \beta a)\frac{B-y}{\lambda} \\ &= \frac{p}{\lambda} . k + \frac{B-y}{\lambda} . m(\beta + \alpha i) \\ &= \left(\frac{a}{\lambda} - \frac{b}{\lambda} . i\right) km + \frac{B-y}{\lambda}(\beta + \alpha i)m \end{aligned}$$

which is divisible by $m$; equivalently, $A + Bi \equiv x + yi (\text{mod. } m)$ Q. E. P.

II. Suppose that, modulo $m$, two numbers $x + yi, x' + y'i$ are congruent to the same complex number, and therefore also to each other modulo $m$. All the more, they are congruent modulo $\lambda$, so $y \equiv y' (\text{mod.} \lambda)$. If both $y, y'$ are therefore assumed to belong to the numbers $0, 1, 2, 3 \ldots . \lambda - 1$, we must have $y = y'$. We then also have $x \equiv x' (\text{mod. } m)$: $x - x'$ is divisible by $m$, and consequently the integer $\frac{x-x'}{\lambda}$ is divisible by $\frac{a}{\lambda} + \frac{b}{\lambda} . i$, or

$$\frac{x - x'}{\lambda} \equiv 0 \left(\text{mod.} \frac{a}{\lambda} + \frac{b}{\lambda} . i\right)$$

But since $\frac{a}{\lambda}, \frac{b}{\lambda}$ are relatively prime, the second part of the theorem in the preceding article shows that $\frac{x-x'}{\lambda}$ is also divisible by the norm of $\frac{a}{\lambda} + \frac{b}{\lambda} . i$, that is, by the number $\frac{p}{\lambda\lambda}$, and consequently $x - x'$ by $\frac{p}{\lambda}$. Thus, if both $x, x'$ are also assumed to belong to the collection of numbers $0, 1, 2, 3 \ldots . \frac{p}{\lambda} - 1$, we must have $x = x'$, or the residues $x + yi, x' + y'i$ are identical. Q. E. S.

It is also evident that this includes the case where the modulus is real, namely $b = 0$ and consequently $\lambda = \pm a$, as well as the case where it is purely imaginary, namely $a = 0$ and consequently $\lambda = \pm b$. In either case we have $\frac{p}{\lambda} = \lambda$.

**42.**

Thus, assigning complex numbers congruent to one another modulo a given modulus to the same class, and incongruent numbers to different classes, we obtain altogether $p$ classes exhausting the whole range of integers, where $p$ denotes the norm of the modulus. A collection of that many numbers, one taken from each class, gives a complete system of incongruent residues, such as we specified in Articles 40, 41. In that system, the choice of residues representing their classes was based on the principle of adopting in each class a residue $x + yi$ for which $y$ has its least value and, among all those with that same least value of $y$, the one for which $x$ has its least value, excluding negative values for both $x$ and $y$. But other purposes will call for other principles. In particular, we should note the method of choosing residues that give the simplest quotients when divided by the modulus. If $\alpha + \beta i, \alpha' + \beta' i, \alpha'' + \beta'' i$, etc., are quotients obtained by dividing congruent numbers by the modulus, the differences among $\alpha, \alpha', \alpha''$, etc., are evidently integers, as are the differences among $\beta, \beta', \beta''$, etc. It is clear that there is always one residue for which $\alpha$ and $\beta$ lie between the bounds 0 and 1, the former included and the latter excluded. We simply call such a residue a least residue. If preferred, the bounds $-\frac{1}{2}$ and $+\frac{1}{2}$ may be adopted instead, one included and the other excluded; we shall call a residue satisfying this restriction *absolutely least*.

Concerning · these least residues, the following problems arise.

**43.**

The least residue of a given complex number $A+Bi$ modulo $a+bi$, of norm $= p$, is found as follows. If $x+yi$ is the required least residue, $(x+yi)(a-bi)$ will be the least residue of the product $(A+Bi)(a-bi)$ modulo $(a+bi)(a-bi)$, that is, modulo $p$. Put

$$aA + bB = Fp + f, \quad aB - bA = Gp + g$$

so that $f, g$ are the least residues of $aA + bB, aB - bA$ modulo $p$. Then

$$x + yi = \frac{f + gi}{a - bi}$$

or

$$x = \frac{af - bg}{p} = A - aF + bG$$
$$y = \frac{ag + bf}{p} = B - aG - bF$$

Evidently, the least residues $f, g$ must be taken either between the bounds 0 and $p - 1$ or between $-\frac{1}{2}p$ and $+\frac{1}{2}p$, according as the complex number's simply least or absolutely least residue is required.

**44.**

A complete system of least residues for a given modulus can be constructed in several ways. The first method begins by determining the bounds within which the real terms must lie, and then, for each value within those bounds, assigning bounds to the imaginary parts. The general criterion for a least residue $x+yi$ modulo $a+bi$ is that both $ax+by = \xi$ and $ay-bx = \eta$ lie between the bounds 0 and $aa+bb$ when simply least residues are considered, or between $-\frac{1}{2}(aa+bb)$ and $+\frac{1}{2}(aa+bb)$ when absolutely least residues are required, one bound being excluded. Special rules would require distinguishing the cases arising from the different signs of the numbers $a, b$. Since developing that distinction presents no difficulty, we refrain from dwelling on it here; it will suffice to show the nature of the method by a single example.

For the modulus $5+2i$, the simply least residues $x+yi$ must be such that both $5x+2y = \xi$ and $5y-2x = \eta$ equal one of the numbers $0, 1, 2, 3 \ldots . 28$. The equation $29x = 5\xi - 2\eta$ shows that positive values of $x$ cannot exceed $\frac{5.28}{29}$, and negative values, disregarding their sign, cannot exceed $\frac{2.28}{29}$. All admissible values of $x$ are therefore $-1, 0, 1, 2, 3, 4$. For $x = -1$, $2y$ must equal one of the numbers $5, 6, 7 \ldots . 33$, and $5y$ one of the numbers $-2, -1, 0, 1 \ldots . 26$; hence the least value of $y$ is $+3$ and the greatest $+5$. Treating the remaining values of $x$ in the same way gives the following table of all the least residues:

| $x$ | $y$ |
|---|---|
| $-1$ | $3, 4, 5$ |
| $0$ | $0, 1, 2, 3, 4, 5$ |
| $+1$ | $1, 2, 3 \,.\, 4, 5 \,.\, 6$ |
| $+2$ | $1, 2, 3, 4, 5, 6$ |
| $+3$ | $2, 3, 4 \,.\, 5, 6$ |
| $+4$ | $2, 3, 4$ |

Similarly, for absolutely least residues, $\xi$ and $\eta$ must equal one of the numbers $-14, -13, -12 \ldots . + 14$. Hence $29x$ cannot lie outside the bounds $-7 \,.\, 14$ and $+7 \,.\, 14$, so $x$ must equal one of the numbers $-3, -2, -1, 0, 1, 2, 3$. For $x = -3$, $2y = \xi - 5x = \xi + 15$ will equal one of the numbers $1, 2, 3 \ldots . 29$, while $5y = \eta + 2x = \eta - 6$ will equal one of the numbers $-20, -19, -18 \ldots . + 8$; this gives for $y$ the single value $+1$. Treating the remaining values of $x$ in the same way, we have the table of all the absolutely least residues:

| $x$ | $y$ |
|---|---|
| $-3$ | $+1$ |
| $-2$ | $-2, -1, 0, +1, +2$ |
| $-1$ | $-3, -2, -1, 0, +1, +2$ |
| $0$ | $-2, -1, 0, +1, +2$ |
| $+1$ | $-2, -1, 0, +1, +2, +3$ |
| $+2$ | $-2, -1, 0, +1, +2$ |
| $+3$ | $-1$ |

**45.**

In applying the second method, it will be useful to distinguish two cases.

In the first case, where $a$ and $b$ have no common divisor, put $a\alpha + \beta b = 1$, and let $k$ be the least positive residue of $\beta a - \alpha b$ modulo $p$. The identities

$$a(\beta a - \alpha b) = \beta p - b(\alpha a + \beta b),$$
$$b(\beta a - \alpha b) = -\alpha p + a(\alpha a + \beta b)$$

then show that $ak \equiv -b, bk \equiv a (\text{mod. } p)$. Thus, putting as above $ax + by = \xi$, $ay - bx = \eta$, we have $\eta \equiv k\xi$, $\xi \equiv -k\eta (\text{mod. } p)$. All the numbers $\xi + \eta i$ to which simply least residues $x + yi$ correspond are therefore obtained by taking either for $\xi$, successively, the values $0, 1, 2, 3 \ldots . p - 1$, and for $\eta$ the least positive residues of the products $k\xi$ modulo $p$, or, in another order, for $\eta$ those values and for $\xi$ the least residues of the products $-k\eta$. From each $\xi + \eta i$ the corresponding $x + yi$ is then found by the formula

$$x + yi = \frac{\xi + \eta i}{a - bi} = \frac{a\xi - b\eta}{p} + \frac{a\eta + b\xi}{p} \,.\, i$$

It is also clear that $\eta$, as $\xi$ increases by one, either increases by $k$ or decreases by $p - k$, and consequently $x + yi$ undergoes

$$\text{either the change } \frac{a - kb}{p} + \frac{ak + b}{p} \,.\, i$$
$$\text{or this one } \frac{a - kb}{p} + b + \left( \frac{ak + b}{p} - a \right) i$$

This observation serves to make the construction easier.

Finally, if absolutely least residues $x + yi$ are required, these instructions evidently change only in that $\xi$ must successively be given values between the bounds $-\frac{1}{2}p$ and $+\frac{1}{2}p$, while for $\eta$ we must take the absolutely least residues of the products $k\xi$. Here is a table of least residues for the modulus $5 + 2i$, arranged in this way:

Simply least residues.

| $\xi + \eta i$ | $x + yi$ | $\xi + \eta i$ | $x + yi$ | $\xi + \eta i$ | $x + yi$ |
|---|---|---|---|---|---|
| $0$ | $0$ | $10 + 25i$ | $+5i$ | $20 + 21i$ | $+2 + 5i$ |
| $1 + 17i$ | $-1 + 3i$ | $11 + 13i$ | $+1 + 3i$ | $21 + 9i$ | $+3 + 3i$ |
| $2 + 5i$ | $+i$ | $12 + i$ | $+2 + i$ | $22 + 26i$ | $+2 + 6i$ |
| $3 + 22i$ | $+1 + 4i$ | $13 + 18i$ | $+1 + 4i$ | $23 + 14i$ | $+3 + 4i$ |
| $4 + 10i$ | $+2i$ | $14 + 6i$ | $+2 + 2i$ | $24 + 2i$ | $+4 + 2i$ |
| $5 + 27i$ | $-1 + 5i$ | $15 + 23i$ | $+1 + 5i$ | $25 + 19i$ | $+3 + 5i$ |
| $6 + 15i$ | $+3i$ | $16 + 11i$ | $+2 + 3i$ | $26 + 7i$ | $+4 + 3i$ |
| $7 + 3i$ | $+1 + i$ | $17 + 28i$ | $+1 + 6i$ | $27 + 24i$ | $+3 + 6i$ |
| $8 + 20i$ | $+4i$ | $18 + 16i$ | $+2 + 4i$ | $28 + 12i$ | $+4 + 4i$ |
| $9 + 8i$ | $+1 + 2i$ | $19 + 4i$ | $+3 + 2i$ | | |

Absolutely least residues.

| $\xi + \eta i$ | $x + yi$ | $\xi + \eta i$ | $x + yi$ | $\xi + \eta i$ | $x + yi$ |
|---|---|---|---|---|---|
| $-14 - 6i$ | $-2 - 2i$ | $-4 - 10i$ | $-2i$ | $+5 - 2i$ | $+1$ |
| $-13 + 11i$ | $-3 + i$ | $-3 + 7i$ | $-1 + i$ | $+6 - 14i$ | $+2 - 2i$ |
| $-12 - i$ | $-2 - i$ | $-2 - 5i$ | $-i$ | $+7 + 3i$ | $+1 + i$ |
| $-11 - 13i$ | $-1 - 3i$ | $-1 + 12i$ | $-1 + 2i$ | $+8 - 9i$ | $+2 - i$ |
| $-10 + 4i$ | $-2$ | $0$ | $0$ | $+9 + 8i$ | $+1 + 2i$ |
| $-9 - 8i$ | $-1 - 2i$ | $+1 - 12i$ | $+1 - 2i$ | $+10 - 4i$ | $+2$ |
| $-8 + 9i$ | $-2 + i$ | $+2 + 5i$ | $+i$ | $+11 + 13i$ | $+1 + 3i$ |
| $-7 - 3i$ | $-1 - i$ | $+3 - 7i$ | $+1 - i$ | $+12 + i$ | $+2 + i$ |
| $-6 + 14i$ | $-2 + 2i$ | $+4 + 10i$ | $+2i$ | $+13 - 11i$ | $+3 - i$ |
| $-5 + 2i$ | $-1$ | | | $+14 + 6i$ | $+2 + 2i$ |

The second case, where $a, b$ are not relatively prime, can easily be reduced to the preceding one. Let $\lambda$ be the greatest common divisor of $a, b$, and let $a = \lambda a'$, $b = \lambda b'$. Let $F$ denote any least residue modulo $\lambda$, with the modulus regarded as a complex number: that is, let it represent any number $x + yi$ such that $x, y$ lie either between the bounds 0 and $\lambda$ or between $-\frac{1}{2}\lambda$ and $+\frac{1}{2}\lambda$, according as simply least or absolutely least residues are considered. Further, let $F'$ denote any least residue modulo $a' + b'i$. Then $(a' + b'i)F + F'$ will represent any least residue modulo $a + bi$, and a complete system of these residues results when all values of $F$ are combined with all values of $F'$.

**46.**

Two complex numbers are called relatively prime if they have no common divisors other than the units. Whenever such common divisors do exist, those whose norm is greatest are called greatest common divisors.

If the factorization of two given numbers into prime factors is available, the greatest common divisor is determined in exactly the same way as for real numbers (*Disquiss. Ar.* Article 18). This also shows that every common divisor of two given numbers must divide their greatest common divisor found in this way. Since it is already evident that its three associates are also common divisors, there will always be four numbers, and no more, that are to be called greatest common divisors; their norm will be a multiple of the norm of any other common divisor.

If the factorization of two given numbers into simple factors is not available, the greatest common divisor is found by an algorithm similar to that for real numbers. Let $m, m'$ be the two given numbers, and form by repeated division the sequence $m'', m'''$, etc., so that $m''$ is the absolutely least residue of $m$ modulo $m'$, then $m'''$ the absolutely least residue of $m'$ modulo $m''$, and so on. Denoting the norms of $m, m', m'', m'''$, etc., respectively by $p, p', p'', p'''$, etc., $\frac{p''}{p'}$ will be the norm of the quotient $\frac{m''}{m'}$ and thus, by the definition of an absolutely least residue, certainly no greater than $\frac{1}{2}$; the same holds for $\frac{p'''}{p''}$, etc. The positive real integers $p', p'', p'''$, etc., will therefore form a steadily decreasing sequence, so that a term 0 must eventually be reached. Equivalently, in the sequence $m, m', m'', m'''$, etc., we shall eventually reach a term that divides the preceding term without a remainder. Let this be $m^{(n+1)}$, and put

$$\begin{aligned} m &= km' + m'' \\ m' &= k'm'' + m''' \\ m'' &= k''m''' + m'''' \end{aligned}$$

and so on, as far as

$$m^{(n)} = k^{(n)} m^{(n+1)}$$

Going through these equations in reverse order shows that $m^{(n+1)}$ divides each preceding term $m^{(n)} \ldots . m'', m', m$; going through the same equations in forward order makes it clear that every common divisor of $m, m'$ also divides each subsequent term. The first conclusion shows that $m^{(n+1)}$ is a common divisor of $m, m'$; the second, that this divisor is greatest.

Whenever the last residue $m^{(n+1)}$ equals one of the four units $1, -1, i, -i$, this will show that $m$ and $m'$ are relatively prime.

**47.**

If the equations of the preceding article, omitting the last, are combined so as to eliminate $m'', m''', m'''' \dots . m^{(n)}$, an equation of the following form results:

$$m^{(n+1)} = hm + h'm'$$

where $h, h'$ are integers; more specifically, if we use the notation introduced in *Disquiss. Ar.* Article 27,

$$\begin{aligned} h &= \pm[k', k'', k''' \dots . k^{(n-1)}] \\ &= \pm[k^{(n-1)}, k^{(n-2)} \dots . k'', k'] \\ h' &= \mp[k, k', k'', k''' \dots . k^{(n-1)}] \\ &= \mp[k^{(n-1)}, k^{(n-2)} \dots . k'', k', k] \end{aligned}$$

the upper or lower signs applying according as $n$ is even or odd. We state this theorem as follows:

*A greatest common divisor of two complex numbers $m, m'$ can be put in the form $hm + h'm'$, with $h, h'$ integers.*

This evidently holds for the greatest common divisor to which the algorithm of the preceding article led, and also for its three associates. For these, the coefficients $h, h'$ must be replaced either by $hi, h'i$, by $-h, -h'$ or by $-hi, -h'i$.

Thus, whenever the numbers $m, m'$ are relatively prime, the following equation can be satisfied:

$$1 = hm + h'm'$$

For example, let the given numbers be $31 + 6i = m$, $11 - 20i = m'$. Here we find

$$\begin{aligned} k &= i, & m'' &= +11 - 5i \\ k' &= +1 - i, & m''' &= +5 - 4i \\ k'' &= +2, & m'''' &= +1 + 3i \\ k''' &= -1 - 2i, & m''''' &= +i \\ k'''' &= +3 - i \end{aligned}$$

and hence

$$\begin{aligned} [k', k'', k'''] &= -6 - 5i \\ [k, k', k'', k'''] &= +4 - 10i \end{aligned}$$

and therefore

$$m''''' = i = (6 + 5i)m + (4 - 10i)m'$$

as well as

$$1 = (5 - 6i)m + (-10 - 4i)m'$$

which is confirmed by carrying out the calculation.

**48.**

The foregoing has prepared everything required for the theory of congruences of the first degree in the arithmetic of complex numbers. But since this theory does not differ essentially from the one that applies to the arithmetic of real numbers and has been set out at length in the *Disquisitiones Arithmeticae*, it will suffice to set down the main points here.

I. The congruence $mt \equiv 1 \pmod{m'}$ is equivalent to the indeterminate equation $mt + m'u = 1$. If this equation is satisfied by the values $t = h, u = h'$, the general solution of the congruence is expressed by $t \equiv h \pmod{m'}$. The condition for solvability is that the modulus $m'$ and the coefficient $m$ have no common divisor.

II. The solution of the congruence $ax + b \equiv c \pmod{M}$, in the case where $a, M$ are relatively prime, depends on the solution of

$$at \equiv 1 \pmod{M}$$

If $t = h$ satisfies this congruence, the general solution of the former is contained in the formula

$$x \equiv (c - b)h \pmod{M}$$

III, The congruence $ax + b \equiv c(\text{mod. } M)$, in the case where $a, M$ have a common divisor $\lambda$, is equivalent to

$$\frac{a}{\lambda} \,.\, x \equiv \frac{c-b}{\lambda} \quad \left(\text{mod.}\frac{M}{\lambda}\right)$$

Thus, when $\lambda$ is taken to be the greatest common divisor of $a, M$, the solution of the given congruence is reduced to the preceding case. It is clear that the necessary and sufficient condition for solvability is that $\lambda$ also divide the difference $c - b$.

**49.**

So far we have touched only on elementary matters, which nevertheless could not be omitted for the sake of the connection. In deeper investigations, the arithmetic of complex numbers resembles the arithmetic of real numbers in that more elegant and simpler theorems result when we admit only moduli that are prime numbers. Indeed, extending them to composite moduli usually involves more length than difficulty, and more work than ingenuity. The following discussion will therefore deal chiefly with prime moduli.

**50.**

Let $X$ denote a function of the indeterminate $x$ of the following form:

$$Ax^n + Bx^{n-1} + Cx^{n-2} + \text{etc.} + Mx + N;$$

where $n$ is a positive real integer, $A, B, C$, etc., real or imaginary integers, and $m$ a complex integer. Here too we shall call any integer a *root* of the congruence $X \equiv 0 \pmod{m}$ if, when substituted for $x$, it gives $X$ a value divisible by the modulus $m$. We shall not regard solutions given by roots congruent modulo the modulus as different.

Whenever the modulus is a prime number, here too such a congruence of degree $n$ cannot have more than $n$ different solutions. Let $\alpha$ denote any fixed integer (complex). $X$ can be put by division by $x-\alpha$, with the indeterminate left free, in the form $X = (x - \alpha)X' + h$, where $h$ is a fixed integer and $X'$ a function of degree $n - 1$ with integer coefficients. Whenever $\alpha$ is a root of the congruence $X \equiv 0 \pmod{m}$, $h$ is evidently divisible by $m$; equivalently, with the indeterminate left free, we have $X \equiv (x - \alpha)X' \pmod{m}$.

Similarly, if $\beta$ denotes a fixed integer and $X'$ is put in the form $(x - \beta)X'' + h'$, $X''$ will be a function of degree $n - 2$ with integer coefficients. If $\beta$ is supposed to be a root of the congruence $X \equiv 0$, it must also satisfy $(\beta-\alpha)X' \equiv 0$, and also $X' \equiv 0$ if the roots $\alpha, \beta$ are incongruent. We conclude that $h'$ must also be divisible by $m$; equivalently, with the indeterminate left free, $X \equiv (x - \alpha)(x - \beta)X'' \pmod{m}$.

In the same way, on adding a third root $\gamma$, incongruent to the preceding ones, we shall have, with the indeterminate left free, $X \equiv (x-\alpha)(x-\beta)(x-\gamma)X'''$, where $X'''$ is a function of degree $n-3$ with integer coefficients. We may continue in the same way. It is also clear that the coefficient of the highest term in each function is $= A$, which we may suppose not divisible by $m$, since otherwise the congruence $X \equiv 0$ would essentially have to be assigned a lower degree. Thus, whenever there are $n$ incongruent roots, say $\alpha, \beta, \gamma \ldots . \nu$, we have, with the indeterminate left free,

$$X \equiv A(x-\alpha)(x-\beta)(x-\gamma)\ldots.(x-\nu) \pmod{m}$$

Consequently, substituting a new value incongruent to each of $\alpha, \beta, \gamma \ldots . \nu$ would certainly give $X$ a value not divisible by $m$, from which the theorem follows at once.

This proof essentially agrees with the one we gave in *Disq. Ar.* Article 43, every step of which applies to complex numbers just as it does to real numbers.

**51.**

Most of what was given in the third section of the *Disquisitiones Arithmeticae* concerning residues of powers also holds in the arithmetic of complex numbers, with slight changes; indeed, the proofs of the theorems could generally be retained. So that nothing is missing, however, we shall present the principal theorems supported by concise proofs. Throughout, the modulus is to be understood to be a prime number.

THEOREM. *If $k$ denotes an integer not divisible by the modulus $m$, of norm $= p$, then $k^{p-1} \equiv 1 \pmod{m}$.*

*Proof.* Let $a, b, c$, etc., form a complete system of incongruent residues modulo $m$, but with the residue divisible by $m$ omitted, so that the number of these numbers, whose collection we denote by $C$, is $= p-1$. Further, let $C'$ be the collection of products $ka, kb, kc$, etc. By hypothesis, none of these products is divisible by $m$, so each has a congruent residue in the collection $C$: we may thus have $ak \equiv a', bk \equiv b', ck \equiv c'$, etc., (mod. $m$), where the numbers $a', b', c'$, etc., themselves occur in the collection $C$. Denote the collection of $a', b', c'$, etc., by $C''$. Let $P, P', P''$ be the products of the individual numbers in the collections $C, C', C''$, respectively; that is,

$$\begin{aligned} P &= abc\ldots. \\ P' &= k^{p-1}abc\ldots. = k^{p-1}P \\ P'' &= a'b'c'\ldots. \end{aligned}$$

Since the numbers in $C''$ are congruent, in succession, to the numbers in $C'$, we have $P'' \equiv P'$, or $P'' \equiv k^{p-1}P$. But any two numbers in $C''$ are readily seen to be incongruent, and therefore all are distinct. The numbers in $C''$ must consequently coincide exactly with those in $C$, with only their order changed, giving $P'' = P$. Thus $(k^{p-1} - 1)P$ is divisible by $m$. Since $m$ is a prime number dividing none of the individual factors of $P$, $k^{p-1} - 1$ must necessarily be divisible by $m$. Q. E. D.

**52.**

THEOREM. *If $k$, as in the preceding article, denotes an integer not divisible by the modulus $m$, and $t$ the least exponent (apart from 0) for which $k^t \equiv 1 \pmod{m}$, then $t$ divides every other exponent $u$ for which $k^u \equiv 1 \pmod{m}$.*

*Proof.* If $t$ did not divide $u$, let $gt$ be the multiple of $u$ next greater than $u$, so that $gt - u$ is a positive integer less than $t$. From $k^t \equiv 1, k^u \equiv 1$ it follows that $0 \equiv k^{gt} - k^u \equiv k^u(k^{gt-u} - 1)$, and hence $k^{gt-u} \equiv 1$: there is a power of $k$ with an exponent less than $t$ that is congruent to unity, contrary to the hypothesis.

It follows as a corollary that $t$ certainly divides $p - 1$.

Here too we shall call numbers $k$ for which $t = p - 1$ *primitive roots* for. the modulus $m$. We shall now show that such numbers really exist.

**53.**

Factor the number $p - 1$ into its prime factors, so that

$$p - 1 = a^\alpha b^\beta c^\gamma \ldots .$$

where $a, b, c$, etc., denote distinct positive real prime numbers. Let $A, B, C$, etc., be integers (complex) not divisible by $m$, and respectively *not* satisfying the congruences

$$x^{\frac{p-1}{a}} \equiv 1, \quad x^{\frac{p-1}{b}} \equiv 1, \quad x^{\frac{p-1}{c}} \equiv 1 \quad \text{etc.}$$

modulo $m$. The existence of such numbers is clear from the theorem of Article 50. Finally, let $h$ be congruent modulo $m$ to the product

$$A^{\frac{p-1}{a^\alpha}} B^{\frac{p-1}{b^\beta}} C^{\frac{p-1}{c^\gamma}} \ldots .$$

Then I say that $h$ will be a primitive root.

*Proof.* Denoting by $t$ the exponent of the lowest power $h^t$ congruent to unity, if $h$ were not a primitive root, $t$ would be a proper divisor of $p - 1$; that is, $\frac{p-1}{t}$ would be an integer greater than unity. This integer evidently has its real prime factors among $a, b, c$, etc. Let us therefore suppose, as we may, that $\frac{p-1}{t}$ is divisible by $a$, and put $p - 1 = atu$. Since $h^t \equiv 1$, we then also have $h^{tu} \equiv 1$, or

$$A^{\frac{p-1}{a^\alpha} \cdot \frac{p-1}{a}} B^{\frac{p-1}{b^\beta} \cdot \frac{p-1}{a}} C^{\frac{p-1}{c^\gamma} \cdot \frac{p-1}{a}} \ldots . \equiv 1$$

But $\frac{p-1}{ab^\beta}$ is evidently an integer, so

$$B^{\frac{p-1}{b^\beta}\cdot\frac{p-1}{a}} = (B^{p-1})^{\frac{p-1}{ab^\beta}} \equiv 1$$

and similarly $C^{\frac{p-1}{c^\gamma}\cdot\frac{p-1}{a}} \equiv 1$, and so on; consequently we must have $A^{\frac{p-1}{a^\alpha}\cdot\frac{p-1}{a}} \equiv 1$ Now determine a positive integer $\lambda$ such that

$$\lambda b^\beta c^\gamma \ldots . \equiv 1 \pmod{a}$$

This can be done, since the prime number $a$ does not divide $b^\beta c^\gamma \ldots .$; put $\lambda b^\beta c^\gamma \ldots . = 1 + a\mu$. Evidently $A^{\lambda\cdot\frac{p-1}{a^\alpha}\cdot\frac{p-1}{a}} \equiv 1$; or, since $\lambda \,.\, \frac{p-1}{a^\alpha} \,.\, \frac{p-1}{a} = (1+a\mu)\frac{p-1}{a} = (p-1)\mu + \frac{p-1}{a}$, we have $A^{(p-1)\mu} \,.\, A^{\frac{p-1}{a}} \equiv 1$. Hence, since $A^{(p-1)\mu} \equiv 1$ holds of itself, we also have $A^{\frac{p-1}{a}} \equiv 1$, contrary to the hypothesis. The supposition that $t$ is a proper divisor of $p-1$ therefore cannot stand, and $h$ must necessarily be a primitive root.

**54.**

Let $h$ be a primitive root for the modulus $m$, of norm $= p$. The terms of the progression

$$1, h, hh, h^3 \ldots . h^{p-2}$$

will be incongruent to one another. It follows easily that every integer not divisible by the modulus must be congruent to one of them: this sequence gives a complete system of incongruent residues with zero excluded. The exponent of the power to which a given number is congruent may be called its *index*, while $h$ is regarded as the *base*. Here are some examples, in which we have placed the absolutely least residue beside each index.

*First example.*

$$m = 5 + 4i, \quad p = 41, \quad h = 1 + 2i$$

| Ind. | Residue | Ind. | Residue | Ind. | Residue | Ind. | Residue | Ind. | Residue |
|---|---|---|---|---|---|---|---|---|---|
| 0 | $+1$ | 8 | $-4$ | 16 | $-2+2i$ | 24 | $+2i$ | 32 | $+1+i$ |
| 1 | $+1+2i$ | 9 | $-3+i$ | 17 | $-1+2i$ | 25 | $-3i$ | 33 | $+1+3i$ |
| 2 | $+1-i$ | 10 | $-i$ | 18 | $+4i$ | 26 | $+2+2i$ | 34 | $+2$ |
| 3 | $+3+i$ | 11 | $+2-i$ | 19 | $+1+3i$ | 27 | $+2+i$ | 35 | $-3$ |
| 4 | $-2i$ | 12 | $-1-i$ | 20 | $-1$ | 28 | $+4$ | 36 | $+2-2i$ |
| 5 | $+3i$ | 13 | $+1-3i$ | 21 | $-1-2i$ | 29 | $+3-i$ | 37 | $+1-2i$ |
| 6 | $-2-2i$ | 14 | $-2$ | 22 | $-1+i$ | 30 | $+i$ | 38 | $-4i$ |
| 7 | $-2-i$ | 15 | $+3$ | 23 | $-3-i$ | 31 | $-2+i$ | 39 | $-1-3i$ |

*Second example.*

$$m = 7, \quad p = 49, \quad h = 1 + 2i$$

| Ind. | Residue | Ind. | Residue | Ind. | Residue | Ind. | Residue | Ind. | Residue |
|---|---|---|---|---|---|---|---|---|---|
| 0 | $+1$ | 10 | $-1-i$ | 20 | $+2i$ | 30 | $+2-2i$ | 40 | $+3$ |
| 1 | $+1+2i$ | 11 | $+1-3i$ | 21 | $+3+2i$ | 31 | $-1+2i$ | 41 | $+3-i$ |
| 2 | $-3-3i$ | 12 | $-i$ | 22 | $-1+i$ | 32 | $+2$ | 42 | $-2-2i$ |
| 3 | $+3-2i$ | 13 | $+2-i$ | 23 | $-3-i$ | 33 | $+2-3i$ | 43 | $+2+i$ |
| 4 | $-3i$ | 14 | $-3+3i$ | 24 | $-1$ | 34 | $+1+i$ | 44 | $-2i$ |
| 5 | $-1-3i$ | 15 | $-2-3i$ | 25 | $-1-2i$ | 35 | $-1+3i$ | 45 | $-3-2i$ |
| 6 | $-2+2i$ | 16 | $-3$ | 26 | $+3+3i$ | 36 | $+i$ | 46 | $+1-i$ |
| 7 | $+1-2i$ | 17 | $-3+i$ | 27 | $-3+2i$ | 37 | $-2+i$ | 47 | $+3+i$ |
| 8 | $-2$ | 18 | $+2+2i$ | 28 | $+3i$ | 38 | $+3-3i$ | | |
| 9 | $-2+3i$ | 19 | $-2-i$ | 29 | $+1+3i$ | 39 | $+2+3i$ | | |

**55.**

We add some observations about primitive roots and the calculation of indices, omitting their proofs because they are easy.

I. In a given system, indices congruent modulo $p-1$ correspond to residues congruent modulo $m$, and conversely.

II. Residues corresponding to indices relatively prime to $p-1$ are themselves primitive roots, and conversely.

III. Suppose that, with a primitive root $h$ taken as the base, the index of another primitive root $h'$ is $t$, and, conversely, $t'$ is the index of $h$ when $h'$ is taken as the base. Then $tt' \equiv 1 \pmod{p-1}$. Under the same assumptions, if the indices of some other number in these two systems are respectively $u, u'$, then $tu' \equiv u, t'u \equiv u' \pmod{p-1}$.

IV. Excluding the numbers $1, 1+i$ and their three associates each, as being too trivial, from the moduli we consider leaves the prime numbers that we placed third and fourth in Article 34. The norms of the latter are real prime numbers of the form $4n+1$; those of the former, however, are squares of odd real prime numbers. Thus, in either case, $p-1$ is divisible by 4.

V. Denote the index of $-1$ by $u$. Then $2u \equiv 0 \pmod{p-1}$, so either $u \equiv 0$ or $u \equiv \frac{1}{2}(p-1)$. But since the index 0 corresponds to the residue $+1$, the index of $-1$ must be $\frac{1}{2}(p-1)$.

VI. Similarly, denoting by $u$ the index of $i$, we have $2u \equiv \frac{1}{2}(p-1) \pmod{p-1}$, and therefore either $u \equiv \frac{1}{4}(p-1)$ or $u \equiv \frac{3}{4}(p-1)$. Here, however, the ambiguity depends on the choice of primitive root. Specifically, if, with the primitive root $h$ taken as the base, the index of $i$ is $\frac{1}{4}(p-1)$, the index becomes $\frac{3}{4}(p-1)$ when $h^\mu$ is taken as the base, where $\mu$ denotes a positive integer of the form $4n+3$ relatively prime to $p-1$, for example the number $p-2$ itself; and conversely. Thus one half of the primitive roots gives $i$ the index $\frac{1}{4}(p-1)$, and the other half the index $\frac{3}{4}(p-1)$. Evidently, for the former bases $-i$ has the index $\frac{3}{4}(p-1)$, and for the latter the index $\frac{1}{4}(p-1)$.

VII. Whenever the modulus is a positive real prime number of the form $4n+3$, say $= q$, so that $p = qq$, the indices of all real numbers are divisible by $q+1$. For, if $t$ denotes the index of the real number $k$, then, because $k^{q-1} \equiv \pmod{q}$, we have $(q-1)t \equiv 0 \pmod{qq-1}$, and consequently $\frac{t}{q+1}$ is an integer. Similarly, the indices of purely imaginary numbers such as $ki$ are divisible by $\frac{1}{2}(q+1)$. Thus primitive roots for such moduli must evidently be sought only among mixed numbers.

VIII. For a modulus $m$ that is a mixed complex prime number, however, whose norm $p$ is consequently a real prime number of the form $4n+1$, any primitive roots may also be chosen among real numbers, among which a complete system of incongruent residues can indeed be exhibited (Article 40). Every real number that is a primitive root for the complex modulus $m$ is evidently also, in the arithmetic of real numbers, a primitive root for the modulus $p$, and conversely.

**56.**

Although the theory of quadratic residues and nonresidues in the arithmetic of complex numbers is contained within the theory of biquadratic residues itself, we shall present its principal theorems separately here before turning to the latter. For brevity, however, we shall discuss only the principal case, where the modulus is an odd complex prime number.

Let $m$ be such a modulus, and $p$ its norm. Evidently, any integer not divisible by $m$, a condition always to be understood here, can or cannot be congruent to a square modulo $m$, according as its index, with some primitive root taken as the base, is even or odd. In the former case that integer is called a quadratic residue of $m$, and in the latter a nonresidue. It follows that, among $p-1$ numbers forming a complete system of incongruent residues not divisible by $m$, one half belongs to the quadratic residues and the other half to the quadratic nonresidues. Any other number outside this system must be assigned the same character in this respect as the number in the system congruent to it.

It also follows that the product of two quadratic residues, as well as the product of two nonresidues, is a quadratic residue, whereas the product of a quadratic residue and a nonresidue is a nonresidue. In general, a product of any number of factors is a quadratic residue or a nonresidue according as the number of nonresidues among its factors is even or odd.

The following general criterion for distinguishing quadratic residues from nonresidues presents itself at once:

A number $k$ not divisible by the modulus is a quadratic residue or a nonresidue of it according as $k^{\frac{1}{2}(p-1)} \equiv 1$ or $k^{\frac{1}{2}(p-1)} \equiv -1 \pmod{m}$.

This theorem follows at once from the fact that, with any primitive root taken as the base, the index of the power $k^{\frac{1}{2}(p-1)}$ becomes either $\equiv 0$ or $\equiv \frac{1}{2}(p-1)$, according as the index of $k$ is even or odd.

**57.**

For a given modulus, it is easy to divide a complete system of incongruent residues into two classes, quadratic residues and nonresidues. This also assigns all the remaining numbers to their respective classes. But a much deeper investigation is needed to find criteria distinguishing the moduli for which a given number is a quadratic residue from those for which it is a nonresidue.

As for the real units $+1$ and $-1$, these are actual squares in the arithmetic of complex numbers, and therefore quadratic residues for *every* modulus. It follows just as easily from the criterion in the preceding article that $i$, and likewise $-i$, is a quadratic residue of every modulus whose norm $p$ has the form $8n+1$, but a nonresidue of every modulus whose norm has the form $8n+5$. Since it evidently makes no difference whether $m$ or one of its associates

$im, -m, -im$ is taken as the modulus, we may suppose that the modulus is the primary one among its associates (Article 36, II). Thus, putting the modulus $= a + bi$, we may suppose $a$ odd and $b$ even. We then always have $aa \equiv 1 \pmod 8$, while $bb$ is either $\equiv 0$ or $\equiv 4 \pmod 8$, according as $b$ is divisible by four or twice an odd number. Thus $+i$ and $-i$ are evidently quadratic residues of the modulus in the former case and nonresidues in the latter.

**58.**

Since the character of a composite number, whether it is a quadratic residue or a nonresidue, depends on the characters of its factors, it evidently suffices to restrict the development of criteria distinguishing the moduli for which a given number $k$ is a quadratic residue from those for which it is a nonresidue to values of $k$ that are prime numbers and, moreover, primary among their associates. In this investigation, *examining examples* immediately supplies very elegant theorems.

Let us begin with $1 + i$, which is found to be a quadratic residue of the moduli

$$-1 + 2i,\ +3 - 2i,\ -5 - 2i,\ -1 - 6i,\ +5 + 4i,\ +5 - 4i,$$
$$-7,\ +7 + 2i,\ -5 + 6i, \quad \text{etc.}$$

but a quadratic nonresidue of the following:

$$-1 - 2i,\ -3,\ +3 + 2i,\ +1 + 4i,\ +1 - 4i,\ -5 + 2i,$$
$$-1 + 6i,\ +7 - 2i,\ -5 - 6i,\ -3 + 8i,\ -3 - 8i,$$
$$+5 + 8i,\ +5 - 8i,\ +9 + 4i,\ +9 - 4i \quad \text{etc.}$$

Examining this list carefully, in which we have always entered the primary one of each four associated moduli, we readily notice that all the moduli $a+bi$ in the first class satisfy $a+b \equiv +1 \pmod 8$, while those in the second satisfy $a + b \equiv -3 \pmod 8$. If, in place of the primary modulus $m$, we take its associate $-m$, this criterion must evidently change so that $a + b \equiv -1$ for moduli in the first class, and $\equiv +3 \pmod 8$ for those in the second. Thus, provided our examination of examples has not misled us, in general, if $a + bi$ denotes a prime number in which $a$ is odd. $b$ even, $1+i$ is a quadratic residue or a quadratic nonresidue of it according as $a + b \equiv \pm 1$ or $\equiv \pm 3 \pmod 8$.

The same rule holds for $-1-i$ as for $1+i$. On the other hand, considering $1 - i$ as the product of $-i$ and $1 + i$, $1 - i$ evidently has the same character as must be assigned to $1 + i$ whenever $b$ is divisible by four, but the opposite character whenever $b$ is twice an odd number. It follows easily that $1 - i$ is a quadratic residue of the prime number $a + bi$ whenever $a - b \equiv \pm 1$, but a nonresidue whenever $a - b \equiv \pm 3 \pmod 8$, always supposing $a$ odd and $b$ even.

This second proposition can also be deduced from the first with the help of a more general theorem, which we state as follows:

In the theory of quadratic residues, the character of $\alpha+\beta i$ with respect to the modulus $a+bi$ is the same as that of $\alpha-\beta i$ with respect to the modulus $a-bi$.

The proof of this theorem follows from the fact that both moduli have the same norm $p$, and that whenever $(\alpha+\beta i)^{\frac{1}{2}(p-1)}-1$ is divisible by $a+bi$, $(\alpha-\beta i)^{\frac{1}{2}(p-1)}-1$ is also divisible by $a-bi$, while whenever $(\alpha+\beta i)^{\frac{1}{2}(p-1)}+1$ is divisible by $a+bi$, $(\alpha-\beta i)^{\frac{1}{2}(p-1)}+1$ must also be divisible by $a-bi$.

**59.**

Let us proceed to odd prime numbers.

We find that $-1+2i$ is a quadratic residue of the moduli $+3+2i$, $+1-4i$, $-5+2i$, $-5-2i$, $-1-6i$, $+7-2i$, $-3+8i$, $+5+8i$, $+5-8i$, $+9+4i$, etc.,

but a nonresidue of the moduli $-1-2i$, $-3$, $+3-2i$, $+1+4i$, $-1+6i$, $+5+4i$, $+5-4i$, $-7$, $+7+2i$, $-5+6i$, $-5-6i$, $-3-8i$, $+9-4i$, etc.

Reducing the moduli of the first class to their absolutely least residues modulo $-1+2i$, we find only $+1$ and $-1$, namely $+3+2i\equiv-1$, $+1-4i\equiv-1$, $-5+2i\equiv+1$, $-5-2i\equiv-1$, etc.

On the other hand, all moduli of the second class are found to be congruent modulo $-1+2i$ either to $+i$ or to $-i$.

But $+1,-1$ are themselves quadratic residues of the modulus $-1+2i$, while $+i$ and $-i$ are its nonresidues. Thus, as far as the examination of examples can be trusted, the following theorem emerges: $-1+2i$ is a quadratic residue or a quadratic nonresidue of the prime number $a+bi$ according as the latter is a quadratic residue or a quadratic nonresidue of $-1+2i$, provided that $a+bi$ is the primary one of its four associates, or rather, that $a$ is odd and $b$ even.

Analogous theorems concerning $+1-2i$, $-1-2i$, $+1+2i$ follow directly from this theorem.

**60.**

Examining examples similarly for $-3$ or $+3$, we find that each is a quadratic residue of the moduli $+3+2i$, $+3-2i$, $-1+6i$, $-1-6i$, $-7$, $-5+6i$, $-5-6i$, $-3+8i$, $-3-8i$, $+9+4i$, $+9-4i$, etc.,

but a nonresidue of $-1+2i$, $-1-2i$, $+1+4i$, $+1-4i$, $-5+2i$, $-5-2i$, $+5+4i$, $+5-4i$, $+7+2i$, $+7-2i$, $+5+8i$, $+5-8i$, etc.

The former are congruent modulo 3 to one of the four numbers $+1,-1,+i,-i$, the latter to one of $+1+i,+1-i,-1+i,-1-i$. The former are precisely the quadratic residues of 3, and the latter its nonresidues.

Thus this examination of examples shows that the prime number $a+bi$, supposing $a$ odd and $b$ even, has the same relation to $-3$, and also to $+3$,

as the latter has to it, in respect of whether one is a quadratic residue or a nonresidue of the other.

Extending a similar examination to other prime numbers, we find this most elegant law of reciprocity confirmed everywhere, and are led to the following fundamental theorem about quadratic residues in the arithmetic of complex numbers:

*If $a + bi, A + Bi$ denote prime numbers such that $a, A$ are odd and $b, B$ even, either each is a quadratic residue of the other, or each is a nonresidue of the other.*

Yet despite the theorem's great simplicity, its proof is beset by great difficulties. We shall not dwell on these here, however, since the theorem itself is only a special case of a more general theorem that, as it were, exhausts the substance of the theory of biquadratic residues. Let us now turn to that theory.

**61.**

We extend to the arithmetic of complex numbers what was said in Article 2 of the preceding paper about the notion of a biquadratic residue and nonresidue. Here, as there, we restrict the investigation to moduli that are prime numbers. It will also usually be tacitly understood that the modulus is chosen to be primary among its associates, that is, $\equiv 1$ modulo $2+2i$, and that the numbers whose character, as biquadratic residues or nonresidues, is under consideration are not divisible by the modulus.

For a given modulus, therefore, the numbers not divisible by it could be divided into three classes: the first containing biquadratic residues; the second, those biquadratic nonresidues that are quadratic residues; and the third, quadratic nonresidues.

But here too it is better to establish two classes in place of the third, so that there are four in all.

With any primitive root taken as the base, biquadratic residues will have indices divisible by 4, or of the form $4n$; those nonresidues that are quadratic residues will have indices of the form $4n+2$; finally, the indices of quadratic nonresidues will be partly of the form $4n+1$, partly of the form $4n+3$. Four classes do arise in this way, but the distinction between the latter two would not be absolute: it would depend on the choice of the primitive root taken as the base. For it is easy to see that one half of the primitive roots gives a given quadratic nonresidue an index of the form $4n+1$, and the other half an index of the form $4n+3$. To remove this ambiguity, we shall always suppose that a primitive root is chosen for which the index $\frac{1}{4}(p-1)$ belongs to the number $+i$ (compare Article 55, VI). This gives a classification that we can state more concisely, independently of primitive roots, as follows.

Let the *first* class contain those numbers $k$ for which $k^{\frac{1}{4}(p-1)} \equiv 1$; these numbers are biquadratic residues of the modulus.

Let the *second* class contain those for which $k^{\frac{1}{4}(p-1)} \equiv i$.

The *third*, those for which $k^{\frac{1}{4}(p-1)} \equiv -1$.

Finally, the *fourth*, those for which $k^{\frac{1}{4}(p-1)} \equiv -i$.

The third class will contain those biquadratic nonresidues that are quadratic residues; quadratic nonresidues will be distributed between the second and fourth classes.

We shall assign to the numbers in these classes the *biquadratic characters* $0, 1, 2, 3$, respectively. If we define the character $\lambda$ of a number $k$ modulo $m$ as the exponent of that power of $i$ to which $k^{\frac{1}{4}(p-1)}$ is congruent, then evidently characters congruent modulo 4 must be regarded as equivalent. For the time being, this notion is restricted to moduli that are prime numbers: in a continuation of these investigations we shall show how it can also be applied to composite moduli.

**62.**

To make an extensive examination of examples concerning the characters of numbers easier, we append a compact table here. With its help, the character of any given number with respect to a modulus whose norm does not exceed 157 is obtained with little work, provided that the following observations are taken into account.

Since the character of a composite number is equal to, or congruent modulo 4 to, the sum of the characters of its individual factors, it suffices if we can assign the characters of prime numbers for a given modulus. Further, since the characters of the units $-1, i, -i$ are evidently congruent to the numbers $\frac{1}{2}(p-1), \frac{1}{4}(p-1), \frac{3}{4}(p-1)$ modulo 4, it will also suffice to give the characters of numbers that are primary among their associates. Finally, since moduli congruent modulo $m$ have the same character, it suffices to enter in the table the characters of numbers contained in a system of absolutely least residues. In addition, reasoning similar to that in Article 58 shows that, if for the modulus $a+bi$ the character of $A+Bi$ is $\lambda$, while for the modulus $a-bi$ $\lambda'$ is the character of $A-Bi$, then we always have $\lambda \equiv -\lambda' \pmod 4$, or, equivalently, $\lambda+\lambda'$ is divisible by 4. It therefore suffices to enter in the table moduli in which $b$ is either 0 or positive.

Thus, for example, if the character of $11-6i$ with respect to the modulus $-5-6i$ is sought, we substitute the numbers $11+6i, -5+6i$ for them. We then determine (Article 43) the absolutely least residue of $11+6i$ modulo $-5+6i$, which is $-1-4i = -1\times(1+4i)$. Since for the modulus $-5+6i$ the character of $-1$ is 30, while the character of $1+4i$, from the table, is 2, we obtain 32, or 0, as the character of $11+6i$ for the modulus $-5+6i$; and, by the last observation, this is also the character of $11-6i$ for the modulus $-5-6i$. Likewise, if the character of $-5+6i$ with respect to the modulus $11+6i$ is sought, its absolutely least residue $1-5i$ is factored into $-i\ 1+i,\ 3-2i$, to which the characters $117, 0, 1$ correspond. Hence the character sought will be 118, or 2; the same character must also be assigned to $-5-6i$ with respect to the modulus $11-6i$.

| Modulus. | Character. | Numbers. |
|---|---|---|
| $-3$ | 3 | $1+i$ |
| $+3+2i$ | 3 | $1+i$ |
| $+1+4i$ | 1 | $-1+2i$ |
| | 3 | $1+i$ |
| $-5+2i$ | 0 | $-1-2i$ |
| | 1 | $1+i$ |
| | 2 | $-1+2i$ |
| $-1+6i$ | 0 | $-3$ |
| | 1 | $1+i,\ -1+2i$ |

| Modulus. | Character. | Numbers. |
|---|---|---|
| $-1+6i$ | 2 | $-1-2i$ |
| $+5+4i$ | 0 | $1+i$ |
| | 1 | $-3$ |
| | 3 | $-1+2i,\ -1-2i$ |
| $-7$ | 0 | $-3$ |
| | 1 | $-1+2i,\ 3-2i$ |
| | 2 | $1+i$ |
| | 3 | $-1-2i,\ 3+2i$ |
| $+7+2i$ | 0 | $1+i,\ 3+2i,\ 3-2i,\ 1-4i$ |
| | 1 | $-3$ |
| | 2 | $-1-2i,\ 1+4i$ |
| | 3 | $-1+2i$ |
| $-5+6i$ | 0 | $1+i,\ -3,\ 3+2i,\ 3-2i$ |
| | 1 | $1-4i$ |
| | 2 | $1+4i$ |
| | 3 | $-1+2i,\ -1-2i$ |
| $-3+8i$ | 0 | $-1+2i,\ 3-2i,\ 1-4i$ |
| | 1 | $1+i,\ 3+2i$ |
| | 2 | $-3$ |
| | 3 | $-1-2i,\ 1+4i,\ -5+2i$ |
| $+5+8i$ | 0 | $-1-2i$ |
| | 1 | $-5-2i,\ -1+6i$ |
| | 2 | $-1+2i,\ 3-2i$ |
| | 3 | $1+i,\ -3,\ 3+2i,\ 1+4i,\ 1-4i$ |
| $+9+4i$ | 0 | $-1+2i,\ 3+2i$ |
| | 1 | $1+i,\ -1-2i,\ 3-2i$ |
| | 2 | $-3,\ 1+4i$ |
| | 3 | $1-4i,\ -5+2i$ |
| $-1+10i$ | 0 | $1+i,\ -1+2i,\ -1-2i,\ 3+2i$ |
| | 1 | $-3$ |
| | 2 | $3-2i,\ -5+2i,\ 5-4i$ |
| | 3 | $1+4i,\ 1-4i$ |

| Modulus. | Character. | Numbers. |
|---|---|---|
| $+3+10i$ | 1 | $1+i,\ -1-2i,\ 1-4i$ |
| | 2 | $-3,\ 3+2i,\ 1+4i,\ -5-2i$ |
| | 3 | $-1+2i,\ 3-2i$ |
| $-7+8i$ | 0 | $1+i,\ -7$ |
| | 1 | $3+2i,\ 3-2i,\ 1-4i,\ -5-2i$ |
| | 2 | $-1-2i,\ 1+4i,\ -5+2i,\ -1-6i$ |
| | 3 | $-1+2i,\ -3,\ -1+6i$ |
| $-11$ | 0 | $-3$ |
| | 1 | $1+i,\ 3-2i,\ 1+4i,\ -5+2i,\ 5+4i$ |
| | 2 | $-1+2i,\ -1-2i$ |
| | 3 | $3+2i,\ 1-4i,\ -5-2i,\ 5-4i$ |
| $-11+4i$ | 0 | $1+i,\ -1+2i,\ 3+2i,\ 5+4i$ |
| | 1 | $-1-2i,\ -1+6i$ |
| | 2 | $-5+2i$ |
| | 3 | $-3,\ 3-2i,\ 1+4i,\ 1-4i,\ -5-2i$ |
| $+7+10i$ | 0 | $1+4i,\ 1-4i,\ -1+6i,\ -1-6i$ |
| | 1 | $-1+2i,\ 3+2i,\ -5+2i$ |
| | 2 | $1+i,\ 3-2i$ |
| | 3 | $-1-2i,\ -3,\ -5-2i$ |
| $+11+6i$ | 0 | $1+i,\ -1+2i,\ -3,\ 1+4i,\ 1-4i,\ -7$ |
| | 1 | $-1-2i,\ 3+2i,\ 3-2i$ |
| | 2 | $-5-2i,\ -1+6i,\ 5-4i$ |
| | 3 | $-5+2i,\ 5+4i,\ 7-2i.$ |

**63.**

We shall now seek, by examining examples, common criteria for the moduli for which a given prime number has the same character. We always suppose the moduli to be primary among their associates, that is, numbers $a+bi$ for which either $a \equiv 1, b \equiv 0$ or $a \equiv 3, b \equiv 2 \pmod 4$.

For the number $1+i$, with which we begin, the rule suggested by the examples is more easily grasped if we separate moduli of the first kind, for which $a \equiv 1, b \equiv 0$, from moduli of the second kind, for which $a \equiv 3, b \equiv 2$. With the help of the table in the preceding article, we find the following correspondences:

| Character | Moduli of the first kind. |
|---|---|
| 0 | $5+4i,\ -7+8i,\ -7-8i,\ -11+4i$ |
| 1 | $1-4i,\ -3+8i,\ -3-8i,\ 9+4i,\ -11$ |
| 2 | $5-4i,\ -7,\ -11-4i$ |
| 3 | $-3,\ 1+4i,\ 5+8i,\ 5-8i,\ 9-4i$ |

If we consider these seventeen examples carefully, in all of them we find the character $\equiv \frac{1}{4}(a-b-1) \pmod 4$.

Likewise, the correspondences are

| Character | Moduli of the second kind. |
|---|---|
| 0 | $3-2i,\ -1-6i,\ 7+2i,\ -5+6i,\ -1+10i,\ 11+6i$ |
| 1 | $-5+2i,\ -1+6i,\ 7-2i,\ -1-10i,\ 3+10i$ |
| 2 | $-1+2i,\ -5-2i,\ 3-10i,\ 7+10i$ |
| 3 | $-1-2i,\ 3+2i,\ -5-6i,\ 7-10i,\ 11-6i$ |

In all these twenty examples, with a little attention, the character is found to be $\equiv \frac{1}{4}(a-b-5) \pmod 4$.

These two rules can easily be combined into one valid for both kinds of moduli, if we consider that $\frac{1}{4}bb$ is $\equiv 0$ for moduli of the first kind and $\equiv 1 \pmod 4$ for those of the second. Thus the character of $1+i$ with respect to any prime modulus that is primary among its associates is $\equiv \frac{1}{4}(a-b-1-bb) \pmod 4$.

We should note in passing that, since $(b+1)^2$ always has the form $8n+1$, or, equivalently, $\frac{1}{4}(2b+bb)$ is even, this character is always even or odd according as $\frac{1}{4}(a+b-1)$ is even or odd. This agrees with the rule for the quadratic character stated in Article 58.

Since $\frac{1}{4}(a-b-1), \frac{1}{4}(a-b+3)$ are integers, one even and the other odd, their product will be even, or, equivalently, $\frac{1}{8}(a-b-1)(a-b+3) \equiv 0 \pmod 4$. Hence the following expression can also be used in place of the one given for the biquadratic character:

$$\frac{1}{4}(a-b-1-bb) - \frac{1}{8}(a-b-1)(a-b+3) = \frac{1}{8}(-aa+2ab-3bb+1)$$

This form also has the advantage that it is not restricted to primary moduli, but supposes only that $a$ is odd and $b$ even. For under this supposition either $a+bi$ or $-a-bi$ will evidently be primary among its associates, and the value of that formula is the same for either modulus.

**64.**

Starting from the last rule obtained in the preceding article, we find

| Number | Character $\equiv$ |
|---|---|
| $-1+i$ | $\frac{1}{8}(aa+2ab-bb-1)$ |
| $-1-i$ | $\frac{1}{8}(-aa+2ab+bb+1)$ |
| $+1-i$ | $\frac{1}{8}(aa+2ab+3bb-1)$ |

This follows at once from the fact that the character of $i$ is $\frac{1}{4}(aa+bb-1)$, while that of $-1$ is $\frac{1}{2}(aa+bb-1) \equiv \frac{1}{2}bb$, since $aa-1$ always has the form $8n$. These four rules, although so far obtained by examining examples, are evidently connected in such a way that, as soon as the proof of one has been completed, the other three will have been proved at the same time. It is scarcely necessary to point out that these rules too suppose only that $a$ is odd and $b$ even.

If we are willing to use formulas restricted to primary moduli, we can use the following form:

| Number | Character $\equiv$ |
|---|---|
| $-1+i$ | $\frac{1}{4}(-a-b+1-bb)$ |
| $-1-i$ | $\frac{1}{4}(a-b-1+bb)$ |
| $+1-i$ | $\frac{1}{4}(-a-b+1+bb)$ |

The simplest formulas result if, as we did at the beginning of our examination of examples, we distinguish moduli of the first and second kinds. The characters are

| Number | For moduli of the first kind | For moduli of the second kind |
|---|---|---|
| $-1+i$ | $\frac{1}{4}(-a-b+1)$ | $\frac{1}{4}(-a-b-3)$ |
| $-1-i$ | $\frac{1}{4}(a-b-1)$ | $\frac{1}{4}(a-b+3)$ |
| $+1-i$ | $\frac{1}{4}(-a-b+1)$ | $\frac{1}{4}(-a-b+5)$ |

**65.**

For the number $-1+2i$, to which we now proceed, we shall likewise use the same distinction between those moduli $a+bi$ for which $a \equiv 1, b \equiv 0$ and those for which $a \equiv 3, b \equiv 2$. The table in Article 62 shows that, with respect to that number, the following correspondences hold:

| Character | Moduli of the first kind |
|---|---|
| 0 | $-3+8i,\ +5-8i,\ +9+4i,\ -11+4i$ |
| 1 | $+1+4i,\ +5-4i,\ -7,\ -3-8i$ |
| 2 | $+1-4i,\ +5+8i,\ -7-8i,\ -11$ |
| 3 | $-3,\ +5+4i,\ +9-4i,\ -7+8i,\ -11-4i$ |

Reducing each of these moduli to its absolutely least residue modulo $-1+2i$, we observe that all those to which character 0 corresponds are $\equiv 1$; those to which character 1 corresponds are $\equiv i$; those whose character is 2 become $\equiv -1$; finally, all those whose character is 3 become $\equiv -i$. But the characters of $1, i, -1, -i$ for the modulus $-1+2i$ are themselves $0, 1, 2, 3$, respectively. Thus, in all these 17 examples, the character of $-1+2i$ with respect to a modulus $a+bi$ of the first kind is identical with the character of that number with respect to the modulus $-1+2i$.

Likewise, with the help of the table, we find the correspondences

| Character | Moduli of the second kind |
|---|---|
| 0 | $+3+2i,\ -5-2i,\ -1+10i,\ -1-10i,\ +11+6i$ |
| 1 | $+3-2i,\ -1+6i,\ -5-6i,\ +7+10i,\ +7-10i$ |
| 2 | $-5+2i,\ -1-6i,\ +7-2i$ |
| 3 | $-1-2i,\ +7+2i,\ -5+6i,\ +3+10i,\ +3-10i,\ +11-6i$ |

Reducing these moduli to their least residues modulo $-1+2i$, all those to which the characters $0, 1, 2, 3$ respectively correspond are found congruent to $-1, -i, +1, +i$. But when, conversely, $-1+2i$ is taken as the modulus, these very numbers have the characters $2, 3, 0, 1$, respectively. Consequently, in all these 19 examples, the character of $-1+2i$ with respect to a modulus of the second kind differs by two units from the character of that number with respect to $-1+2i$ taken as the modulus.

Moreover, it is readily seen that entirely similar results will hold for $-1-2i$.

**66.**

For the number $-3$, we omit the distinction between moduli of the first and second kinds, since the result shows that it is superfluous here. Thus the correspondences are

| Character | Moduli |
|---|---|
| 0 | $-1+6i,\ -1-6i,\ -7,\ -5+6i,$ $-5-6i,\ -11,\ 11+6i,\ 11-6i$ |
| 1 | $-1-2i,\ 1-4i,\ -5+2i,\ 5+4i,\ 7+2i,$ $5-8i,\ -1+10i,\ -7-8i,\ -11-4i,\ 7-10i$ |
| 2 | $3+2i,\ 3-2i,\ -3+8i,\ -3-8i,$ $9+4i,\ 3+10i,\ 3-10i$ |
| 3 | $-1+2i,\ 1+4i,\ -5-2i,\ 5-4i,\ 7-2i,$ $5+8i,\ -1-10i,\ -7+8i,\ -11+4i,\ 7+10i$ |

Reducing these moduli to their least residues modulo 3, we see that those to which character 0 corresponds are partly $\equiv 1$, partly $\equiv -1$; those whose character is 1 become either $\equiv 1 - i$ or $\equiv -1 + i$; those whose character is 2 become either $\equiv i$ or $\equiv -i$; finally, those to which character 3 belongs are either $\equiv 1 + i$ or $\equiv -1 - i$. From this examination of examples we therefore conclude that the character of $-3$ for a modulus that is a prime number primary among its associates is identical with the character of that very number when 3, or equivalently $-3$, is considered as the modulus.

**67.**

Examining examples similarly for other prime numbers, we find that $3 \pm 2i, -1 \pm 6i, 7 \pm 2i, -5 \pm 6i$, etc., give theorems similar to the one reached in Article 65 for $-1+2i$, whereas $1 \pm 4i, 5 \pm 4i, -3 \pm 8i, 5 \pm 8i, 9 \pm 4i$, etc., behave in the same way as $-3$. The examination of examples thus leads to a most elegant theorem,. which, by analogy with the theory of quadratic residues in the arithmetic of real numbers, may be called the FUNDAMENTAL THEOREM of the theory of biquadratic residues, namely:

*Let $a + bi, a' + b'i$ denote distinct prime numbers, primary among their respective associates, that is, congruent to unity modulo $2+2i$. The biquadratic character of $a+bi$ with respect to the modulus $a'+b'i$ will be identical with the character of $a' + b'i$ with respect to the modulus $a + bi$, if either both numbers $a + bi, a' + b'i$, or at least one of them, belong to the first kind, that is, are congruent to unity modulo 4. On the other hand, those characters will differ from one another by two units if neither number $a + bi, a' + b'i$ belongs to the first kind, that is, if each is congruent modulo 4 to $3 + 2i$.*

Yet despite the great simplicity of this theorem, its proof must be counted among the most deeply hidden mysteries of higher arithmetic. At least as matters now stand, it can be unravelled only through the most subtle investigations, which would go far beyond the limits of the present paper. We therefore reserve the publication of this proof, as well as the development of the connection between this theorem and those that we had begun to establish by examining examples at the beginning of this paper, for a third paper. Nevertheless, by way of conclusion, we shall give here what is required to prove the theorems proposed in Articles 63. 64.

**68.**

We begin with prime numbers $a + bi$ for which $b = 0$ (the third species in Article 34). Here, for the number to be primary among its associates, $a$ must therefore be a negative real prime number of the form $-(4n + 3)$, for which we shall write $-q$, such as $-3, -7, -11, -19$, etc. Denoting by $\lambda$ the character of $1 + i$, with that number taken as the modulus, we must have

$$i^{\lambda} \equiv (1+i)^{\frac{1}{4}(qq-1)} \equiv 2^{\frac{1}{8}(qq-1)} \,.\, i^{\frac{1}{8}(qq-1)} \pmod{q}$$

But it is known that 2 is a quadratic residue or a quadratic nonresidue of $q$ according as $q$ has the form $8n+7$ or the form $8n+3$. From this we conclude that, in general,

$$2^{\frac{1}{2}(q-1)} \equiv (-1)^{\frac{1}{4}(q+1)} \equiv i^{\frac{1}{2}(q+1)} \pmod{q}$$

and consequently, raising to the power with exponent $\frac{1}{4}(q+1)$,

$$2^{\frac{1}{8}(qq-1)} \equiv i^{\frac{1}{8}(q+1)^2} \pmod{q}$$

The preceding equation therefore takes the form

$$i^{\lambda} \equiv i^{\frac{1}{8}(q+1)^2 + \frac{1}{8}(qq-1)} \equiv i^{\frac{1}{4}(qq+q)} \pmod{q}$$

whence it follows that

$$\lambda \equiv \frac{1}{4}(qq+q) \equiv \frac{1}{4}(q+1)^2 - \frac{1}{4}(q+1) \pmod{4}$$

or, since $\frac{1}{4}(q+1)^2 \equiv 0 \pmod{4}$, $\lambda \equiv -\frac{1}{4}(q+1) \equiv \frac{1}{4}(a-1) \pmod{4}$. This is precisely the theorem of Article 63 for the case $b = 0$.

**69.**

Moduli $a+bi$ for which $b=0$ does not hold (numbers of the fourth species in Art. 34) are much harder to deal with, however, and several investigations must come first. We shall denote the norm $aa+bb$, which will be a real prime of the form $4n+1$, by $p$.

Let $S$ denote the collection of all simply least residues for the modulus $a+bi=m$, with zero excluded, so that the count of the numbers contained in $S$ is $=p-1$. Let $x+yi$ denote any number of this system, and put $ax+by=\xi, ay-bx=\eta$. Then $\xi,\eta$ will be integers lying between the bounds 0 and $p$, *excluding the bounds*. For in the present case, where $a,b$ are relatively prime, the formulas of Art. 45, namely $\eta\equiv k\xi, \xi\equiv -k\eta \pmod p$, show that neither of the numbers $\xi,\eta$ can be $=0$ unless the other vanishes at the same time, so that $x=0, y=0$, a combination we have already excluded. Thus the criterion for a number $x+yi$ to be contained in $S$ is that the four numbers $\xi,\eta,p-\xi,p-\eta$ be positive.

We further observe that $\xi=\eta$ cannot hold for any such number. For it would follow that $p(x+y)=a(\xi+\eta)+b(\xi-\eta)=2a\xi$, which is absurd, since none of the factors $2,a,\xi$ is divisible by $p$. Similarly, the equation $p(x-y+a+b)=2a\xi+(a+b)(p-\xi-\eta)$ shows that $\xi+\eta=p$ cannot hold. Since the numbers $\xi-\eta, p-\xi-\eta$ must therefore be either positive or negative, we obtain from this a subdivision of the system $S$ into four collections $C, C', C'', C'''$, assigning the numbers as follows:

| Into collection | Numbers for which |
|---|---|
| $C$ | $\xi-\eta$ positive, $p-\xi-\eta$ positive |
| $C'$ | $\xi-\eta$ positive, $p-\xi-\eta$ negative |
| $C''$ | $\xi-\eta$ negative, $p-\xi-\eta$ negative |
| $C'''$ | $\xi-\eta$ negative, $p-\xi-\eta$ positive |

The criterion for a number of the collection $C$ is thus, strictly speaking, sixfold: the six numbers $\xi,\eta,p-\xi,p-\eta,\xi-\eta,p-\xi-\eta$ must be positive. But conditions 2, 5 and 6 evidently already imply the others. The same applies to the collections $C', C'', C'''$, so that the complete criteria are threefold, namely:

| For collection | The following numbers must be positive |
|---|---|
| $C$ | $\eta,\quad \xi-\eta,\quad p-\xi-\eta$ |
| $C'$ | $p-\xi,\quad \xi-\eta,\quad \xi+\eta-p$ |
| $C''$ | $p-\eta,\quad \eta-\xi,\quad \xi+\eta-p$ |
| $C'''$ | $\xi,\quad \eta-\xi,\quad p-\xi-\eta$ |

Even without our pointing it out, anyone will readily understand that, in the geometric representation of complex numbers (see Art. 39), the numbers of the system $S$ lie inside the square whose sides join the points representing the numbers $0, a+bi, (1+i)(a+bi), i(a+bi)$, and that the subdivision of the system $S$ corresponds to the division of the square by its diagonals. Here, however, we have preferred to use purely arithmetical reasoning, leaving illustration by a geometric representation to the experienced reader for the sake of brevity.

**70.**

If four complex numbers $r = x + yi, r' = x' + y'i, r'' = x'' + y''i, r''' = x''' + y'''i$ are related in such a way that $r' = m+ir, r'' = m+ir' = (1+i)m-r, r''' = m + ir'' = im - ir$, and the first, $r$, is supposed to belong to the collection $C$, then the others $r', r'', r'''$ will belong respectively to the collections $C', C'', C'''$. For, putting $\xi = ax + by, \eta = ay - bx, \xi' = ax' + by', \eta' = ay' - bx', \xi'' = ax'' + by'', \eta'' = ay'' - bx'', \xi''' = ax''' + by''', \eta''' = ay''' - bx'''$, we find

$$\begin{aligned} \eta = p - \xi' = p - \eta'' = \xi''' \\ \xi - \eta = \xi' + \eta' - p = \eta'' - \xi'' = p - \xi''' - \eta''' \\ p - \xi - \eta = \xi' - \eta' = \xi'' + \eta'' - p = \eta''' - \xi''' \end{aligned}$$

from which the truth of the theorem follows at once by the criteria. Since $r = m + ir'''$ holds again, it is easy to see that, if $r$ is supposed to belong to $C'$, the numbers $r', r'', r'''$ belong respectively to $C'', C''', C$; if it belongs to $C''$, they belong to $C''', C, C'$; and finally, if it belongs to $C'''$, they belong to $C, C', C''$.

At the same time, we conclude that the individual collections $C, C', C'', C'''$ contain equally many numbers, namely $\frac{1}{4}(p-1)$.

**37.**

THEOREM. *Let $k$ denote an integer not divisible by $m$. If each number of the collection $C$ is multiplied by $k$, and the simply least residues of the products modulo $m$ are distributed among the collections $C, C', C'', C'''$, with the counts of the residues belonging to these individual collections denoted respectively by $c, c', c'', c'''$, then the character of the number $k$ with respect to the modulus $m$ will be $\equiv c' + 2c'' + 3c'''$ (mod 4).*

*Proof.* Let those $c$ least residues belonging to $C$ be $\alpha, \beta, \gamma, \delta$, etc.; next, let the $c'$ residues belonging to $C'$ be $m+i\alpha', m+i\beta', m+i\gamma', m+i\delta'$, etc.; further, let the $c''$ residues belonging to $C''$ be $(1+i)m-\alpha'', (1+i)m-\beta'', (1+i)m-\gamma'', (1+i)m-\delta''$, etc.; and finally, let the $c'''$ residues belonging to $C'''$ be $im-i\alpha''', im-i\beta''', im-i\gamma''', im-i\delta'''$, etc. Now consider four products, namely:

1) the product of all $\frac{1}{4}(p-1)$ numbers constituting the collection $C$:

2) the product of the products obtained by multiplying each of these numbers by $k$;

3) the product of the least residues of these products, namely of the numbers $\alpha, \beta, \gamma, \delta$, etc., $m+i\alpha', m+i\beta'$, etc., etc.

4) the product of all $c+c'+c''+c'''$ numbers $\alpha, \beta, \gamma, \delta$, etc., $\alpha', \beta', \gamma', \delta'$, etc., $\alpha'', \beta'', \gamma'', \delta''$, etc.

Denoting these four products in order by $P, P', P'', P'''$, we evidently have

$$P' = k^{\frac{1}{4}(p-1)}P, \quad P' \equiv P'', \quad P'' \equiv P'''i^{c'+2c''+3c'''} \pmod{m}$$

and therefore

$$Pk^{\frac{1}{4}(p-1)} \equiv P'''i^{c'+2c''+3c'''} \pmod{m}$$

It is easy to see, however, that the numbers $\alpha', \beta', \gamma', \delta'$, etc., $\alpha'', \beta'', \gamma'', \delta''$, etc., $\alpha''', \beta''', \gamma''', \delta'''$, etc., all belong to the collection $C$, and are distinct both from one another and from the numbers $\alpha, \beta, \gamma, \delta$, etc., just as the latter themselves are distinct from one another. All these numbers taken together, disregarding their order, must therefore be exactly the same as all the numbers constituting the collection $C$. Hence we conclude that $P = P'''$, and consequently

$$Pk^{\frac{1}{4}(p-1)} \equiv Pi^{c'+2c''+3c'''} \pmod{m}$$

Finally, since none of the individual factors of the product $P$ is divisible by $m$, we conclude that

$$k^{\frac{1}{4}(p-1)} \equiv i^{c'+2c''+3c'''} \pmod{m}$$

so that $c'+2c''+3c'''$ will be the character of the number $k$ with respect to the modulus $m$. Q. E. D.

**72.**

To apply the general theorem of the preceding article to the number $1+i$, we must subdivide the collection $C$ again into two smaller collections $G$ and $G'$. We shall assign to the collection $G$ those numbers $x+yi$ for which $ax+by=\xi$ is less than $\frac{1}{2}p$, and to the other collection $G'$ those for which $\xi$ is greater than $\frac{1}{2}p$. We shall denote the counts of the numbers contained in the collections $G, G'$ respectively by $g, g'$, so that $g+g'=\frac{1}{4}(p-1)$.

The complete criterion for numbers belonging to $G$ is therefore that the three numbers $\eta, \xi-\eta, p-2\xi$ be positive. For the third condition for the collection $C$, requiring $p-\xi-\eta$ to be positive, is already implicitly contained in these conditions, since $p-\xi-\eta=(\xi-\eta)+(p-2\xi)$. Similarly, the complete criterion for numbers belonging to $G'$ consists in positive values of the three numbers $\eta, p-\xi-\eta, 2\xi-p$.

It follows easily that the product of any number of the collection $G$ by the number $1+i$ belongs to the collection $C'''$. For, putting

$$(x+yi)(1+i)=x'+y'i, \quad \text{and} \quad ax'+by'=\xi', \ ay'-bx'=\eta', \ \text{we find}$$
$$\xi'=\xi-\eta, \quad \eta'-\xi'=2\eta, \quad p-\xi'-\eta'=p-2\xi$$

that is, the criterion for the number $x+yi$ to belong to the collection $G$ is identical with the criterion for the number $x'+y'i$ to belong to the collection $C'''$.

In precisely the same way, it is shown that the product of any number of the collection $G'$ by $1+i$ belongs to the collection $C''$.

Thus, if in the preceding article we give $k$ the value $1+i$, we have $c=0, c'=0, c''=g', c'''=g$, and the character of the number $1+i$ becomes $3g+2g'=\frac{1}{2}(p-1)+g$. Since the characters of the numbers $i, -1$ are $\frac{1}{4}(p-1), \frac{1}{2}(p-1)$, the characters of the numbers $-1+i, -1-i, 1-i$ will respectively be $\frac{3}{4}(p-1)+g, g, \frac{1}{4}(p-1)+g$. The whole matter now turns, therefore, on determining the number $g$.

**73.**

What we have set out in Arts. 69—72 is, strictly speaking, independent of the assumption that $m$ is a primary number. From now on, however, we shall at least suppose that $a$ is odd and $b$ even, and further that $a, b$ and $a-b$ are positive numbers. First of all, we must establish the bounds on the values of $x$ in the collection $G$.

Putting $ay - bx = \eta, (a+b)x - (a-b)y = \zeta, p - 2ax - 2by = \theta$, the criterion for numbers $x+yi$ to belong to the collection $G$ consists in the three conditions that $\eta, \zeta, \theta$ be positive numbers. Since $px = (a-b)\eta + a\zeta, p(a - 2x) = a\theta + 2b\eta$, it is evident that $x$ and $2a - x$ must be positive numbers, or that $x$ must equal one of the numbers $1, 2, 3 \ldots \frac{1}{2}(a-1)$. Further, since $(a-b)\theta = 2b\zeta + p(a-b-2x)$, it is clear that, as long as $x$ is less than $\frac{1}{2}(a-b)$, the second condition (requiring $\zeta$ to be positive) already implies the third (that $\theta$ must be positive). Conversely, whenever $x$ is greater than $\frac{1}{2}(a-b)$, the second condition is already contained in the third. Thus, for the values $x$ takes in $1, 2, 3 \ldots \frac{1}{2}(a-b-1)$, we need only ensure that $\eta$ and $\zeta$ are positive, or that $y$ is greater than $\frac{bx}{a}$ and less than $\frac{(a+b)x}{a-b}$. For such a given value of $x$, the total count of numbers $x+yi$ is therefore

$$\left[\frac{(a+b)x}{a-b}\right] - \left[\frac{bx}{a}\right]$$

if we use the brackets in the same sense in which we have already used them repeatedly elsewhere (compare *Theorematis arithm. dem. nova*, Art. 4, and *Theorematis fund. in doctr. de residuis quadr.* etc., *Algorithm. nov.*, Art. 3). For the values $x$ takes in $\frac{1}{2}(a-b+1), \frac{1}{2}(a-b+3) \ldots . \frac{1}{2}(a-1)$, on the other hand, it suffices to give $\eta$ and $\theta$ positive values, or to make $y$ greater than $\frac{bx}{a}$ and less than $\frac{p-2ax}{2b}$, or $\frac{1}{2}b + \frac{aa-2ax}{2b}$. Thus, for such a given value of $x$, the total count of numbers $x+yi$ is

$$\left[\frac{1}{2}b + \frac{aa-2ax}{2b}\right] - \left[\frac{bx}{a}\right]$$

Hence we conclude that the count of numbers in the collection $G$ is

$$g = \Sigma\left[\frac{(a+b)x}{a-b}\right] + \Sigma\left[\frac{1}{2}b + \frac{aa-2ax}{2b}\right] - \Sigma\left[\frac{bx}{a}\right]$$

where the first sum extends over all integer values of $x$ from 1 to $\frac{1}{2}(a-b-1)$, the second from $\frac{1}{2}(a-b+1)$ to $\frac{1}{2}(a-1)$, and the third from 1 to $\frac{1}{2}(a-1)$.

If we use the symbol $\varphi$ in the same sense as in the place cited (*Theorematis fund.* etc., *Algor. nov.*, Art. 3), namely so that

$$\varphi(t,u) = \left[\frac{u}{t}\right] + \left[\frac{2u}{t}\right] + \left[\frac{3u}{t}\right] \ldots + \left[\frac{t'u}{t}\right]$$

where $t, u$ denote any positive numbers and $t'$ the number $[\frac{1}{2}t]$, then the first term becomes $= \varphi(a-b, a+b)$, and the third $= -\varphi(a,b)$; the second, however, becomes

$$= \frac{1}{4}bb + \Sigma\left[\frac{aa-2ax}{2b}\right]$$

But, writing the terms in reverse order, we have

$$\begin{aligned}\Sigma\left[\frac{aa-2ax}{2b}\right] &= \left[\frac{a}{2b}\right] + \left[\frac{3a}{2b}\right] + \left[\frac{5a}{5b}\right] + \ldots + \left[\frac{(b-1)a}{2b}\right] \\ &= \varphi(2b,a) - \varphi(b,a)\end{aligned}$$

Our formula therefore takes the following form:

$$g = \varphi(a-b, a+b) + \varphi(2b,a) - \varphi(a,b) - \varphi(b,a) + \frac{1}{4}bb$$

Let us first consider the term $\varphi(a-b, a+b)$, which immediately changes into $\varphi(a-b, 2b) + 1 + 2 + 3 +$ etc. $+ \frac{1}{2}(a-b-1)$, or into

$$\varphi(a-b,2b) + \frac{1}{8}((a-b)^2 - 1)$$

Next, since the general theorem gives $\varphi(t,u) + \varphi(u,t) = [\frac{1}{2}t] \,.\, [\frac{1}{2}u]$ when $t, u$ are relatively prime positive integers, we have

$$\varphi(a-b,2b) = \frac{1}{2}b(a-b-1) - \varphi(2b, a-b)$$

and consequently

$$\varphi(a-b,a+b) = \frac{1}{8}(aa+2ab-3bb-4b-1) - \varphi(2b,a-b)$$

Let us arrange the parts of $\varphi(2b, a-b)$ as follows:

$$\begin{aligned}&\left[\frac{a-b}{2b}\right] + \left[\frac{3(a-b)}{2b}\right] + \left[\frac{5(a-b)}{2b}\right] + \text{ etc. } + \left[\frac{(b-1)(a-b)}{2b}\right] \\ &+ \left[\frac{a-b}{b}\right] + \left[\frac{2(a-b)}{b}\right] + \left[\frac{3(a-b)}{b}\right] + \text{ etc. } + \left[\frac{\frac{1}{2}b(a-b)}{b}\right]\end{aligned}$$

The second series evidently becomes

$$= \varphi(b, a-b) = \varphi(b,a) - 1 - 2 - 3 - \text{ etc. } - \frac{1}{2}b$$
$$= \varphi(b,a) - \frac{1}{8}(bb + 2b)$$

We write the first series, with the order of its terms reversed, as follows:

$$\left[\frac{1}{2}(a+1-b) - \frac{a}{2b}\right] + \left[\frac{1}{2}(a+3-b) - \frac{3a}{2b}\right]$$
$$+ \left[\frac{1}{2}(a+5-b) - \frac{5a}{2b}\right] + \text{etc.} + \left[\frac{1}{2}(a-1) - \frac{(b-1)a}{2b}\right]$$

This expression. Since, with $t$ denoting an integer and $u$ a fraction, we generally have $[t-u] = t - 1 - [u]$, it changes into the following:

$$\frac{1}{8}b(2a-4-b) - \left[\frac{a}{2b}\right] - \left[\frac{3a}{2b}\right] - \left[\frac{5a}{2b}\right] - \text{ etc. } - \left[\frac{(b-1)a}{2b}\right]$$
$$= \frac{1}{8}b(2a-4-b) - \varphi(2b,a) + \varphi(b,a)$$

Hence

$$\varphi(2b, a-b) = 2\varphi(b,a) - \varphi(2b,a) + \frac{1}{4}b(a-3-b)$$

and therefore

$$\varphi(a-b, a+b) = \varphi(2b,a) - 2\varphi(b,a) + \frac{1}{8}(aa - bb + 2b - 1)$$

Substituting this value in the formula for $g$ given above, and also substituting $\varphi(a,b) + \varphi(b,a) = \frac{1}{4}b(a-1)$, we obtain

$$g = 2\varphi(2b,a) - 2\varphi(b,a) + \frac{1}{8}(aa - 2ab + bb + 4b - 1)$$

**74.**

Precisely similar reasoning deals with the case where $a$, $b$ remain positive but $a-b$ is negative, or $b-a$ positive. The equations $p(a-2x) = 2b\eta + a\theta$, $p(b-a+2x) = 2b\zeta + (b-a)\theta$ show that $\frac{1}{2}a - x$ and $x + \frac{1}{2}(b-a)$ must be positive, and hence that $x$ must equal one of the numbers $-\frac{1}{2}(b-a-1), -\frac{1}{2}(b-a-3), -\frac{1}{2}(b-a-5) \ldots . + \frac{1}{2}(a-1)$. Further, the equation $px + (b-a)\eta = a\zeta$ shows that, for negative values of $x$, the condition requiring $\eta$ to be positive is already contained in the condition requiring $\zeta$ to be positive, whereas the reverse holds whenever $x$ is given a positive value. Thus the values of $y$, for a fixed negative value of $x$, must lie between $\frac{(a+b)x}{a-b}$ and $\frac{p-2ax}{2b}$, whereas for a positive value of $x$ they must lie between $\frac{bx}{a}$ and $\frac{p-2ax}{2b}$. For $x = 0$, these bounds are evidently 0 and $\frac{p-2ax}{2b}$, with the value $y = 0$ itself excluded. Hence

$$g = -\Sigma\left[\frac{(a+b)x}{a-b}\right] + \Sigma\left[\frac{1}{2}b + \frac{aa-2ax}{2b}\right] - \Sigma\left[\frac{bx}{a}\right]$$

where the first sum extends over all negative values of. $x$ from $-1$ to $-\frac{1}{2}(b-a-1)$; the second over all values of $x$ from $-\frac{1}{2}(b-a-1)$ to $\frac{1}{2}(a-1)$; and the third over all positive values of $x$ from $+1$ to $\frac{1}{2}(a-1)$. In this way the first sum gives $-\varphi(b-a, b+a)$, the second, as in the preceding article, $\frac{1}{4}bb + \varphi(2b,a) - \varphi(b,a)$, and finally the third $-\varphi(a,b)$; thus we have

$$g = -\varphi(b-a, b+a) + \varphi(2b,a) - \varphi(b,a) - \varphi(a,b) + \frac{1}{4}bb$$

Now, in the same way as in the preceding article, we obtain

$$\begin{aligned}\varphi(b-a, b+a) &= \varphi(b-a, 2b) - \frac{1}{8}((b-a)^2 - 1) \\ &= \frac{1}{8}(3bb - 2ab - aa - 4b + 1) - \varphi(2b, b-a)\end{aligned}$$

as well as

$$\varphi(2b, b-a) = \varphi(2b,a) - 2\varphi(b,a) + \frac{1}{4}b(b-1-a)$$

and consequently

$$\varphi(b-a, b+a) = 2\varphi(b,a) - \varphi(2b,a) + \frac{1}{8}(bb - aa - 2b + 1)$$

and finally

$$g = 2\varphi(2b,a) - 2\varphi(b,a) + \frac{1}{8}(aa - 2ab + bb + 4b - 1)$$

It is therefore proved that the same formula for $g$ holds whether $a-b$ is positive or negative, provided that $a$, $b$ are positive.

**75.**

To obtain a further reduction, we put

$$L = \left[\frac{a}{2b}\right] + \left[\frac{2a}{2b}\right] + \left[\frac{3a}{2b}\right] + \text{etc.} + \left[\frac{\frac{1}{2}ba}{2b}\right]$$

$$M = \left[\frac{(\frac{1}{2}b+1)a}{2b}\right] + \left[\frac{(\frac{1}{2}b+2)a}{2b}\right] + \left[\frac{(\frac{1}{2}b+3)a}{2b}\right] + \text{etc.} + \left[\frac{ba}{2b}\right]$$

$$N = \left[\frac{a+b}{2b}\right] + \left[\frac{2a+b}{2b}\right] + \left[\frac{3a+b}{2b}\right] + \text{etc.} + \left[\frac{\frac{1}{2}ba+b}{2b}\right]$$

It is easy to see that $[u]+[u+\frac{1}{2}] = [2u]$ holds in general, whatever real quantity $u$ denotes. Thus $L + N = \varphi(b, a)$, and since evidently $L + M = \varphi(2b, a)$, we have

$$\varphi(2b, a) - \varphi(b, a) = M - N$$

Further, it is evident that the sum of the first term of the series $N$ and the next-to-last term of the series $M$, namely $\left[\frac{a+b}{2b}\right] + \left[\frac{(b-1)a}{2b}\right]$, is $= \frac{1}{2}(a-1)$, and that the same sum is formed by the second term of the series $N$ and the third-to-last term of the series $M$, and so on. Since the last term of the series $M$ is also $= \frac{1}{2}(a-1)$, while the last term of the series $N$ is $= \left[\frac{a+2}{4}\right] = \frac{1}{4}(a \mp 1)$, with the upper or lower sign applying according as $a$ is of the form $4n+1$ or $4n-1$, we have

$$M + N = \frac{1}{4}(a-1)b + \frac{1}{4}(a \mp 1)$$

and therefore

$$\varphi(2b, a) - \varphi(b, a) = \frac{1}{4}(a-1)b + \frac{1}{4}(a \mp 1) - 2N$$

Thus the formula for $g$ found in Arts. 73 and 74 becomes

$$g = \frac{1}{8}((a+b)^2 - 1) + 2n - 4N$$

putting $a \mp 1 = 4n$, where $n$ will be an integer. Since this gives $1 = 16nn - 8an + aa$, this formula can also be written as follows:

$$g = \frac{1}{8}(-aa + 2ab + bb + 1) + 4(\frac{1}{2}(a+1)n - nn - N)$$

Since $g$ is the character of the number $-1-i$ for the modulus $a+bi$, this character is therefore $\equiv \frac{1}{8}(-aa + 2ab + bb + 1) \pmod 4$. This is precisely the theorem found above (Art. 64) by examining examples, and the theorems concerning the characters of the numbers $1+i$, $1-i$, $-1+i$ follow from it at once. Thus these four theorems have now been rigorously proved for the case where $a$ and $b$ are positive.

**76.**

If $a$ remains positive and $b$ is negative, put $b = -b'$, so that $b'$ is positive. It is now proved that, for the modulus $a + b'i$, the character of the number $-1 - i$ is $\equiv \frac{1}{8}(-aa + 2ab' + b'b' + 1) \pmod 4$. Hence the character of the number $-1 + i$ for the modulus $a - b'i$, by the theorem stated in Art. 62, is $\equiv \frac{1}{8}(aa - 2ab' - b'b' - 1)$, that is, the character of the number $-1 + i$ for the modulus $a + bi$ becomes $\equiv \frac{1}{8}(aa + 2ab - bb - 1)$. This is precisely the theorem stated in Art. 64, from which the other three concerning the characters of the numbers $1 + i$, $1 - i$, $-1 - i$ follow at once. Thus these theorems are also proved for the case where $b$ is negative, that is, for all cases where $a$ is positive.

Finally, if $a$ is negative, put $a = -a'$, $b = -b'$. By what has already been proved, the character of the number $1 + i$ with respect to the modulus $a' + b'i$ is $\equiv \frac{1}{8}(-a'a' + 2a'b' - 3b'b' + 1) \pmod 4$, and it makes no difference whether we take the number $a' + b'i$ or its opposite $-a' - b'i$ as the modulus. Thus the character of the number $1 + i$ with respect to the modulus $a + bi$ is evidently $\equiv \frac{1}{8}(-aa + 2ab - 3bb + 1)$, and the same applies to the characters of the numbers $1 - i$, $-1 + i$, $-1 - i$.

We therefore conclude that the proof of the theorems concerning the characters of the numbers $1 + i$, $1 - i$, $-1 + i$, $-1 - i$ (Arts. 63. 64) is no longer subject to any restriction.

# REVIEWS

## OF HIS OWN

# WRITINGS.

Göttingische gelehrte Anzeigen. 1808 May 12.

A paper submitted by Professor Gauss to the Royal Society of Sciences on January 15 of this year,

*Theorematis arithmetici demonstratio nova*,

an account of whose contents we still owe here, concerns the famous fundamental theorem in the theory of quadratic residues, which plays so important a role both throughout *higher arithmetic* and in the neighbouring parts of analysis. As is well known, an integer $a$ is called a *quadratic residue* of the integer $b$ if there are numbers of the form $xx - a$ divisible by $b$, while in the opposite case $a$ is called a *quadratic nonresidue* of $b$. The number $a$ may be positive or negative, whereas $b$ is always regarded as positive. Higher arithmetic teaches that all prime numbers $b$ for which a given number $a$ is a quadratic residue are included in certain linear forms, while other linear forms contain all the primes of which $a$ is a nonresidue. Thus, for example, $-1$ is a quadratic residue of all primes of the form $4n + 1$ and a quadratic nonresidue of all primes of the form $4n + 3$; further, $+2$ is a quadratic residue of all primes of the forms $8n + 1$, $8n + 7$, but a quadratic nonresidue of all primes of the forms $8n + 3$, $8n + 5$. There are infinitely many similar special theorems, but all can be derived by combining the two just mentioned with the following general theorem: Two distinct positive (odd) primes, $p$, $q$, always bear the *same* relation to each other (that is, one is a quadratic residue or nonresidue of the other according as the other is a residue or nonresidue of the first) if either both have the form $4n + 1$, or at least one does. Their mutual relation is opposite, however (that is, one is a nonresidue of the other if the latter is a residue of the former, and conversely), whenever *both* have the form $4n + 3$. This is the fundamental theorem mentioned above, which can be expressed in more than one form. The form chosen here is the one in which it is newly proved in Professor Gauss's paper.

The most beautiful theorems of higher arithmetic, especially those discussed here, have the peculiarity that they are easily discovered by examining examples, whereas their proofs lie extremely well hidden and can be found only through very deep investigations. This is precisely what gives higher arithmetic the magical appeal that has made it the favourite field of the foremost mathematicians, quite apart from its inexhaustible wealth, in which it so far surpasses every other part of pure mathematics. The two special theorems mentioned above were already known to Fermat, who claimed also to possess their proofs. Whether he was mistaken in this we cannot decide, since he never published anything about them; but we can certainly regard

it as possible, since there are several examples of self-deception among other great mathematicians, notably EULER, LEGENDRE, and also FERMAT himself. EULER gave the first proof of the first of these theorems. But this great mathematician did not succeed in proving the other, despite his devoted efforts over many years; it was left to LAGRANGE to fill that gap. Both mathematicians also proved various other special theorems, but a greater number, which they found by examining examples, always eluded their attempts at proof. It is, however, a remarkable twist of chance that neither mathematician discovered in this way the general fundamental theorem, which admits of so simple a statement. It was first presented, though in a somewhat different form, by LEGENDRE in the *Histoire de l'Académie des Sciences de Paris* 1785. Both there and later in his *Essai d'une théorie des nombres*, this excellent analyst sought to base the proof on very ingenious investigations, which nevertheless failed to reach the desired goal; and, unless we are mistaken, that goal could not have been reached by this route either.

The author of the paper to which this review is devoted entered the field of higher arithmetic at a time when all earlier work by other mathematicians in this subject was entirely unknown to him. It is chiefly to this circumstance that we must attribute his having followed a wholly distinctive course throughout. He found that fundamental theorem very early by examining examples, but only a whole year later, after many difficulties and unsuccessful attempts, did he succeed in finding the first completely rigorous proof, developed in the fourth section of his *Disquisitiones Arithmeticae*. This proof, however, rests on very laborious and lengthy arguments. He subsequently found three other proofs, which are free of that inconvenience but instead require other very deep investigations, wholly different in content. One of these proofs is also given in the cited work, Art. 262; the other two will be made known in due course. The wish therefore remained that it might be possible to discover a shorter proof, independent of investigations of an unrelated kind. The author thus hopes that friends of higher arithmetic will take pleasure in a fifth proof, presented in the present paper in fewer than five pages, which seems to leave nothing to be desired in any respect. Given the concentrated brevity of this proof, we can admittedly give only an imperfect idea of its course here. Anything more would be all the more unnecessary, since Volume XVI of the *Commentationes*, in which it has already been printed, will appear shortly.

The proof rests on the following new theorem: Let $p$ denote a (positive odd) prime and $k$ any integer not divisible by $p$. Suppose further that, among the remainders obtained by dividing the $\frac{1}{2}(p-1)$ products $k$, $2k$, $3k \ldots . \frac{1}{2}(p-1)k$ by $p$, there are in all $\mu$ remainders greater than $\frac{1}{2}p$ (and hence $\frac{1}{2}(p-1)-\mu$ less than $\frac{1}{2}p$). Then $k$ will be a quadratic residue of $p$ if $\mu$ is even, but a quadratic nonresidue if $\mu$ is odd. The number $\mu$, which depends only on $k$ and $p$, may be represented by the symbol $(k, p)$. A sequence of deductions that cannot be abridged now shows that, if $k$ and $p$ are two relatively prime odd numbers, $(k, p) + (p, k) + \frac{1}{4}(k-1)(p-1)$ is always an *even* number. It follows that, whenever $k$ and $p$ both have the form $4n+3$, one of the numbers $(k, p)$, $(p, k)$ must be even and the other odd; in all other cases, that is, whenever both numbers $k$ and $p$, or at least one of them, have the form $4n+1$, either $.(k, p)$, $(p, k)$ must both be even or both must be odd. Combined with the theorem above, this immediately gives the truth of the fundamental theorem.—The same route by which these results are obtained also gives, in the paper, a new proof of the two special theorems mentioned above. It is easy to show that $(-1, p) = \frac{1}{2}(p-1)$, hence even or odd according as $p$ has the form $4n+1$ or $4n+3$; similarly, $(2, p) = \frac{1}{4}(p-1)$ if $p$ has the form $4n+1$, and $(2, p) = \frac{1}{4}(p+1)$ if $p$ has the form $4n+3$. Thus $(2, p)$ is even whenever $p$ has the form $8n+1$ or $8n+7$, but odd whenever $p$ has the form $8n+3$ or $8n+5$.

---

Göttingische Gelehrte Anzeigen. 1808 September 19.

A lecture submitted by Professor Gauss to the Royal Society of Sciences,

*Summatio quarumdam serierum singularium*,

aims to develop more fully and in greater generality a remarkable investigation concerning the division of the circle, whose foundation had already been laid in the *Disquisitionibus Arithmeticis*, to provide it with complete proofs, and to show its unexpected connection with other important truths. Let $n$ denote a prime, $k$ any integer not divisible by $n$, and $\omega$ the arc $\frac{1}{n}360^\circ$. Among the numbers $1, 2, 3, 4, \ldots . n-1$, let the distinct quadratic residues of $n$ be represented by $a, a', a''$, and so on, while the numbers remaining after these have been excluded, or the quadratic nonresidues of $n$, are represented by $b, b', b''$, and so on. It is proved in the cited work, Art. 356, that, when $n$ has the form $4n+1$,

$$\left.\begin{array}{l}\cos ak\omega + \cos a'k\omega + \cos a''k\omega + \text{etc.} \\ -\cos bk\omega - \cos b'k\omega - \cos b''k\omega - \text{etc.}\end{array}\right\} = \pm\sqrt{n}.$$

and

$$\left.\begin{array}{l}\sin ak\omega + \sin a'k\omega + \sin a''k\omega + \text{etc.} \\ -\sin bk\omega - \sin b'k\omega - \sin b''k\omega - \text{etc.}\end{array}\right\} = 0$$

whereas, when $n$ has the form $4m+3$, the sum of the first series becomes $= 0$ and that of the second $= \pm\sqrt{n}$. The sign to be placed before the radical depends on the value of the number $k$, or rather on its relation to $n$, and is easily determined for *all* values of $k$ at a given value of $n$ once it is determined for *one*. For it can be shown that *the same* sign applies to all values of $k$ that are quadratic residues of $n$, and the opposite sign to all those that are quadratic nonresidues of $n$. Since the investigation had already been carried this far in the cited work, and only the determination of the sign for any one value of $k$ remained, one would have thought that, with the main matter settled, this further determination could easily be supplied. This seemed all the more likely because examining examples immediately gives an extremely simple result: for $k = 1$, or for all values that are quadratic residues of $n$, the radical in the formulas above must always be taken *positively*. Yet in seeking a proof of this observation we encounter wholly unexpected difficulties, and the procedure that led so satisfactorily to the absolute value of those series proves entirely inadequate when the complete determination of the signs is required. The *metaphysical* reason for this phenomenon (to use the expression customary among French mathematicians) must be sought in the

fact that, in dividing the circle, analysis makes no distinction between the arcs $\omega, 2\omega, 3\omega \ldots (n-1)\omega$, but treats them all alike. As this gives the investigation a new interest, Professor GAUSS found in it, as it were, an invitation to leave nothing untried in overcoming the difficulty. Only after many and varied unsuccessful attempts did he succeed, by a route remarkable in itself. He starts from the summation of certain series whose terms have the following form:

$$\frac{(1-x^m)(1-x^{m-1})(1-x^{m-2})\ldots(1-x^{m-\mu+1})}{(1-x)\ (1-xx)\ (1-x^3)\ldots.(1-x^\mu)}$$

If, for brevity, we denote such a function by $(m, \mu)$, which, as the paper shows, is always a *polynomial* in $x$, then the series

$$1-(m,1)+(m,2)-(m,3)+\text{etc.}$$

$$1+x^{\frac{1}{2}}(m,1)+x(m,2)+x^{\frac{3}{2}}(m,3)+\text{etc.}$$

terminate after the term in position $m+1$, provided that $m$ denotes a positive integer. For even values of $m$ the sum of the first series becomes

$$=(1-x)(1-x^3)(1-x^5)\ldots(1-x^{m-1})$$

and $=0$ for odd values of $m$, whereas the sum of the second series always becomes

$$=(1+x^{\frac{1}{2}})(1+x)(1+x^{\frac{3}{2}})\ldots(1+x^{\frac{1}{2}m})$$

For fractional and negative values of $m$, too, the summation of these series leads to interesting results, though they are not needed for the present purpose. We content ourselves with giving just one of them here. The infinite series

$$1+x+x^3+x^6+x^{10}+\text{etc.}$$

whose exponents are the triangular numbers is the product of the factors

$$\frac{1-xx}{1-x}\times\frac{1-x^4}{1-x^3}\times\frac{1-x^6}{1-x^5}\times\frac{1-x^8}{1-x^7}\ \text{etc.}$$

or, if preferred, of

$$(1+x)^2(1+xx)^2(1+x^3)^2(1+x^4)^2\ \text{etc.}$$

multiplied by

$$(1-x)(1-xx)1-x^3)(1-x^4)\ \text{etc.}$$

An account of how these summations are applied to the main subject would take us too far here. We may refer readers to the paper itself all the more

readily since it will soon appear in print. The summations quoted above are only a special application of the summation of the following series:

$$1 + \cos k\omega + \cos 4k\omega + \cos 9k\omega + \text{etc.} + \cos(n-1)^2 k\omega = T$$

$$\sin k\omega + \sin 4k\omega + \sin 9k\omega + \text{etc.} + \sin(n-1)^2 k\omega = U$$

which the paper gives for all values of $k$, and without the restriction that $n$ be prime. It is shown that

$$T = \pm\sqrt{n}, \quad T = \pm\sqrt{n}, \quad T = 0, \quad T = 0$$

and

$$U = \pm\sqrt{n}, \quad U = 0, \quad U = 0, \quad U = \pm\sqrt{n}$$

according as $n$ has the forms $4m, 4m+1, 4m+2, 4m+3$, respectively. Here again the sign of the radical depends on $k$. Its determination requires many individual cases to be distinguished, and is developed and proved by two different routes in such a way that nothing will remain to be desired. Comparing these two routes also leads to the following very remarkable theorem: Let $n$ be the product of any number of distinct odd primes $a, b, c, d$, and so on, of which altogether $\mu$ have the form $4m+3$. Suppose further that among those factors there are altogether $\nu$ for each of which the product of the others (respectively $\frac{n}{a}, \frac{n}{b}, \frac{n}{c}, \frac{n}{d}$, and so on) is a quadratic nonresidue. Then $\nu$ will be even whenever $\mu$ has the form $4m$ or $4m+1$, but odd whenever $\mu$ has the form $4m+2$ or $4m+3$. The known fundamental theorem concerning quadratic residues is only a special case of this theorem; conversely, the theorem just stated is easily derived from that fundamental theorem. Thus these investigations also provide a fourth proof of this important theorem, which the author first proved by two entirely different routes in the *Disquisitionibus Arithmeticis*, and recently by a third, equally different route in a separate paper.

---

Göttingische gelehrte Anzeigen. 1817 March 10.

On 10 February Court Councillor Gauss submitted a lecture to the Royal Society, entitled:

*Theorematis fundamentalis in doctrina de residuis quadraticis demonstrationes et ampliationes novae.*

A distinctive feature of higher arithmetic is that so many of its most beautiful theorems can be discovered with the greatest ease by examining examples, while their proofs are far from readily apparent. Often they are found only after many unsuccessful attempts, through deeply probing investigations and fortunate combinations. This remarkable phenomenon arises from the often wonderful interconnection between the different branches of this part of mathematics. For the same reason, theorems for which a proof was initially sought in vain for years can often later be proved in several entirely different ways. Once a new theorem has been discovered through examples, finding *some* proof must of course be the first requirement; but after this has succeeded, in higher arithmetic we must not always regard the investigation as finished and the search for other proofs as a needless luxury. For one thing, we do not usually find the most beautiful and simplest proofs first; for another, insight into the wonderful interconnection of the truth of higher arithmetic is precisely one of the chief attractions of this study, and not infrequently leads in turn to the discovery of new truths. For these reasons, finding new proofs of truths already known must here often be considered at least as important as discovering the truth itself. These reflections are familiar to those who know higher arithmetic: much of Euler's contribution to it is known to consist in finding proofs of theorems which Fermat had already discovered, apparently through examples.

The theory of quadratic residues provides a clear illustration of these remarks. It rests chiefly on the so-called fundamental theorem, which states that the mutual relations of two (odd positive) primes to one another (insofar as one is a quadratic residue or nonresidue of the other) are the same whenever one or both primes have the form $4k + 1$, but opposite whenever both primes have the form $4k + 3$. For readers less familiar with higher arithmetic, we recall that an integer is called a quadratic residue of another if adding a multiple of the latter to the former can give a square; it is called a nonresidue if this is impossible. We do not intend to repeat the whole history of this beautiful theorem, so easily discovered through examples. We merely observe that the author of the present paper, after investigations initially pursued in vain for quite some time, has already given, one after another, four

entirely different proofs. Two are contained in the *Disquisitionibus Arithmeticis*; the third is the subject of a separate paper in the sixteenth volume of the Commentationes; and the fourth is interwoven with the paper *summatio quarumdam serierum singularium* in the first volume of the *Commentationes recentiores*. For these two papers readers may consult our reviews of 1808, May 12 and September 19, which also give fuller historical references. That the author did not stop at these four proofs, although each leaves nothing to be desired as regards rigour, needs no justification among the friends of higher arithmetic. Still, he would probably not have worked so eagerly to add further proofs if a particular circumstance had not prompted him to do so, and this must be mentioned here. In the year 1805 he began to study the theories of cubic and biquadratic residues, which are much richer and more interesting still than the theory of quadratic residues. These investigations revealed the same phenomena as the latter theory, only, as it were, on a larger scale. As soon as the right path was taken, examining examples immediately yielded a number of extremely simple theorems which exhaust those theories, bear a surprising resemblance to the theorems for quadratic residues, and in particular also provide a counterpart to the fundamental theorem. But the difficulty of finding entirely satisfactory proofs of these theorems proved greater still, and only after many attempts, continued over a considerable number of years, did the author finally reach his goal. The close analogy between the theorems for quadratic and higher residues suggested that there must also be analogous proofs for the two. Yet the methods of proof first found for quadratic residues could not be applied to the higher residues at all; this was precisely what led him to keep seeking further new proofs for the former. The author therefore wishes the present paper, which opens up some further resources for the theory of quadratic residues, to be regarded as a precursor to the theory of cubic and biquadratic residues. He intends to publish that theory in the future; its subjects are among the most difficult in higher arithmetic.

The present paper consists of three independent parts. It contains the fifth and sixth proofs of the fundamental theorem, and a new method, connected with the third proof, · for deciding whether a given integer is a quadratic residue or nonresidue of a given prime. Of the first four proofs, the third was unquestionably the one that combined the greatest simplicity with independence from investigations of a different kind; LEGENDRE therefore also included it in the new edition of his *Essai d'une théorie des nombres*. The *fifth* proof seems at least to equal the third in both respects. The two proofs are related insofar as they start from the same theorem, but in their subsequent development they are completely different from one another.

The theorem is as follows. Let $m$ be a (positive odd) prime and $M$ an integer not divisible by $m$. If, among the remainders obtained by dividing the products

$$M,\ 2M,\ 3M,\ 4M \ldots\ldots \tfrac{1}{2}(m-1)M$$

by $m$, the number of those greater than $\frac{1}{2}m$ is denoted by $n$, then $M$ is a quadratic residue or nonresidue of $m$ according as $n$ is even or odd. To obtain a proof of the fundamental theorem, it is now assumed that $M$, too, is an odd positive prime, and that $N$ has the same meaning with respect to $M$ and $m$ as $n$ has with respect to $m$ and $M$. Thus whether $N$ is even or odd determines whether $m$ is a quadratic residue or nonresidue of $M$. By a very short chain of arguments the author shows that the number of all positive integers which are less than $\frac{1}{2}mM$ and at the same time on division by $m$ leave a remainder less than $\frac{1}{2}m$, and on division by $M$ a remainder less than $\frac{1}{2}M$,

$$= \tfrac{1}{8}(m-1)(M-1) + \tfrac{1}{2}n + \tfrac{1}{2}N$$

and consequently that

$$\tfrac{1}{4}(m-1)(M-1) + n + N$$

is always an even number. Thus, whenever at least one of the numbers $m, M$ has the form $4k+1$, so that $\frac{1}{4}(m-1)(M-1)$ is even, $n+N$ will also be even; hence $n$ and $N$ are either both even or both odd. If, on the other hand, both $m$ and $M$ have the form $4k+3$, then $n+N$ must be odd, so one of the numbers $n, N$ is even and the other odd. Together with the theorem above, this gives the fundamental theorem at once.

The *sixth* proof is as short and elegant as the fifth, but rests on somewhat more intricate combinations. The limited space of these pages allows us only to touch on its main point, omitting the details. Let the following symbols denote:

$p, q$ two (distinct positive odd) primes,

$\alpha$ a so-called *primitive root* for the modulus $p$, that is, an integer not divisible by $p$ (here taken positive), such that no power lower than $\alpha^{p-1}$ is congruent to one modulo $p$

$x$ an indeterminate quantity

$\xi$ the function

$$x - x^{\alpha} + x^{\zeta} - x^{\eta} + x^{\theta} - \text{etc.} - x^{\lambda}$$

where, for convenience in printing, $\zeta, \eta, \theta \ldots \lambda$ stand for the numbers $\alpha\alpha, \alpha^3, \alpha^4 \ldots \alpha^{p-2}$;

$\varepsilon$ one, taken positively if $p$ has the form $4k+1$, negatively if $p$ has the form $4k+3$;

$\delta$ one, taken positively if at least one of the numbers $p, q$ has the form $4k+1$, negatively if both have the form $4k+3$;

$\gamma$ one, taken positively if $q$ is a quadratic residue of $p$, negatively if $q$ is a quadratic nonresidue of $p$;

$\beta$ one, taken positively if $p$ is a quadratic residue of $q$, negatively if $p$ is a quadratic nonresidue of $q$.

With these preparations it follows readily from Article 51 of the *Disquisitiones Arithmeticae* that the function

$$\xi^q - x^q + x^{q\alpha} - x^{q\zeta} + x^{q\eta} - x^{q\theta} + \text{etc.} + x^{q\lambda}$$

has, when expanded, coefficients all divisible by $q$. Hence, if this function is set $= qX$, then $X$ is a *polynomial* whose coefficients, too, are integers. By arguments which it would take too much space to discuss here, the paper proves that dividing the function $qX\xi$ by $x^{p-1}+x^{p-2}+x^{p-3}+x^{p-4}+\text{etc.}+x+1$ leaves the remainder

$$\varepsilon p(\delta p^{\frac{1}{2}(q-1)} - \gamma)$$

and therefore that dividing the function $X\xi$ by the same divisor leaves the remainder

$$\frac{\varepsilon p(\delta p^{\frac{1}{2}(q-1)} - \gamma)}{q}$$

This quantity must therefore be an integer. Since $\delta\delta = 1$, it follows readily that

$$p^{\frac{1}{2}(q-1)} - \gamma\delta$$

must be divisible by $q$. A known theorem also gives that $p^{\frac{1}{2}(q-1)} - \beta$ is divisible by $q$, so necessarily $\beta = \gamma\delta$, from which the fundamental theorem again follows at once.

The fundamental theorem, together with some known theorems, can indeed provide a fairly short way of deciding whether a given positive integer is a quadratic residue or nonresidue of a given prime, as the paper shows in detail. But further reflection on the third proof of the fundamental theorem led the author to a much more flexible solution. This forms the paper's third part; here we give only its final rule, omitting the reasoning behind it for brevity. To decide whether the positive integer $b$, not divisible by the prime $a$, is a quadratic residue or nonresidue of that prime, form, in just the same way as when finding the greatest common divisor of $a$ and $b$, the equations

$$\begin{aligned} a &= \beta b + c \\ b &= \gamma c + d \\ c &= \delta d + e \\ d &= \varepsilon e + f \text{ etc.} \end{aligned}$$

until the sequence of numbers $a, b, c, d, e, f$ and so on reaches .the unit. Let. the numbers $\frac{1}{2}a, \frac{1}{2}b, \frac{1}{2}c, \frac{1}{2}d$ and so on, with their fractional part $\frac{1}{2}$ discarded whenever any of the numbers $a, b, c, d$ and so on is odd, be denoted by $a', b', c', d'$ and so on. Let $\mu$ be the number of occurrences of two odd numbers immediately following one another in the sequence $a', b', c', d'$ and so on. Finally, let $\nu$ be the number of odd numbers in the sequence $\beta, \gamma, \delta, \varepsilon$ and so on to which, in the sequence $b', c', d', e'$ and so on, a number of the form $4k + 1$ or $4k + 2$ respectively corresponds. Then $b$ is a quadratic residue or nonresidue of $a$ according as $\mu + \nu$ is even or odd, except in the single case where $b$ is even and $a$ has the form $8k + 3$ or $8k + 5$. In that case the rule is reversed: an even $\mu + \nu$ indicates that $b$ is a quadratic nonresidue of $a$, while an odd $\mu + \nu$ indicates that $b$ is a quadratic residue of $a$.

---

.Göttingische gelehrta Anzeigen. 1825 April 11.

On 5 April Court Councillor Gauss presented a lecture to the Royal Society, entitled:

*Theoria Residuorum Biquadraticorum, Commentatio prima.*

The theory of quadratic residues is known to form one of the most interesting parts of higher arithmetic. After repeated investigations it can now be regarded as complete and closed. Accounts of its history can be found in these pages for 1808 May 12 and September 19, and 1817 March 10. The last of these also gave some preliminary information about the investigations which the author of the present paper had been pursuing since the year 1805 into the related theory of cubic and biquadratic residues, equally fruitful and interesting but far more difficult. Although he already possessed the essential elements of this theory at that time, other work has so far prevented him from publishing anything about it. Only now has he been able to begin writing up part of these investigations. He has begun with the theory of biquadratic residues, which is more closely related to the theory of quadratic residues than is that of cubic residues. The present paper, however, is by no means intended to exhaust this exceptionally rich subject. The development of the *general* theory, which requires a quite distinctive extension of the domain of higher arithmetic, is reserved for a future continuation. This first paper contains those investigations which could be set out completely without such an extension. Only some of the results can be selected for this review.

An integer $a$ is called a biquadratic residue of the integer $p$ if there are numbers of the form $x^4 - a$ divisible by $p$; a biquadratic nonresidue if no numbers of that form can be divisible by $p$. Clearly all biquadratic residues of $p$ are also quadratic residues of the same number, and hence all quadratic nonresidues are also biquadratic nonresidues. But not all quadratic residues are also biquadratic residues. It suffices to restrict the investigations to the case where $p$ is a prime of the form $4n + 1$ and $a$ is not divisible by $p$, since all other cases can easily be reduced to this one.

The investigations of this subject divide into two parts, according as $p$ or $a$ is regarded as given. The first is much less difficult than the second and, compared with it, is to be regarded as entirely elementary. The paper contains a complete account of everything essential to be said about it.

From the second part, by contrast, only a few special cases are treated here. These could be dealt with without too much preparation and can serve as preliminaries to the general theory to be given in the future. They are the cases where $a = -1$ and $a = \pm 2$ are set.

The first presents no difficulty at all: the *Disquisitiones Arithmeticae* had already shown that $-1$ is a biquadratic residue of $p$ whenever $p$ has the form $8n+1$, but only a quadratic residue and a biquadratic nonresidue of $p$ when $p$ has the form $8n+5$. The case $a = \pm 2$ is quite different. It has long been known that $+2$ and $-2$ are quadratic, and therefore also biquadratic, nonresidues of $p$ when $p$ has the form $8n+5$, and at least quadratic residues when $p$ has the form $8n+1$. It is also known that for this form of $p$, $+2$ and $-2$ are either both biquadratic residues or both biquadratic nonresidues: but deciding which of these two cases occurs is an investigation of a much higher order. The paper develops two different criteria for this purpose.

The first criterion is connected with the decomposition of the number $p$ into one square plus twice another, which, as is known (since, as already remarked, $p$ is assumed to be prime), is always possible and possible in only one way. If we set $p = gg + 2hh$, then $\pm 2$ is a biquadratic residue of $p$ when $g$ has the form $8n+1$ or $8n+7$, but a biquadratic nonresidue when $g$ has the form $8n+3$ or $8n+5$.

The second criterion is connected with the decomposition of the number $p$ into two squares, which is also known to be always possible and possible in only one way. If we set $p = ee + ff$ and suppose that $ee$ denotes the odd square and $ff$ the even square, then the assumed form $p = 8n+1$ already implies that $\frac{1}{2}f$, too, is even. Thus $f$ has either the form $8m$ or the form $8m+4$: in the first case $\pm 2$ is a biquadratic residue, in the other a biquadratic nonresidue of $p$.

We only touch here on an observation for which higher arithmetic so often provides an occasion: it is not so much the beauty and simplicity of the theorems themselves as the difficulty of proving them that makes them especially remarkable. Once there is reason to suspect a connection between the behaviour of the number $\pm 2$ and the two decompositions of the number $p$ just described, it is extremely easy to discover that connection by examining examples. But even for the first criterion its proof is not quite easy; for the second it lies much deeper, closely bound up with other subtle auxiliary investigations, which in turn lead to a remarkable extension of the theory of circle division. It is chiefly this wonderful interconnection of truths. that, as has often been observed, gives higher arithmetic such a distinctive appeal. Naturally these proofs themselves cannot be excerpted here; readers must consult the paper itself. But a couple of other new arithmetical theorems, also closely connected with the proof of the second criterion, deserve to be singled out here because of their simplicity.

Let $p$ be a prime of the form $4k+1$, and put this number $= ee + ff$ so that $ee$ denotes the odd square and $ff$ the even square. If we further set

$$1 \,.\, 2 \,.\, 3 \ldots\ldots \; k = q$$

$$(k+1)(k+2)(k+3) \ldots\ldots 2k = r$$

then $\pm e$ is always the least remainder obtained by dividing $\frac{r}{2q}$ by $p$, and $\pm f$ the least remainder obtained by dividing $\frac{1}{2}rr$ by $p$ (where the least remainder is always understood to be taken between the bounds $-\frac{1}{2}p$ and $+\frac{1}{2}p$). The number $\frac{r}{2q}$, whose value for $p = 5$ is 1, can for larger values of $p$ also be put in the following form:

$$\frac{6 \,.\, 10 \,.\, 14 \,.\, 18 \ldots\ldots (p-3)}{2 \,.\, 3 \,.\, 4 \,.\, 5 \ldots\ldots k}$$

It is very remarkable that the decomposition of the number $p$ into two squares can thus be obtained quite directly. But an accompanying circumstance is almost more remarkable still. This procedure always gives the root of the odd square, $e$, with a positive sign if $e$, taken positively, has the form $4m+1$, and with a negative sign if $e$, taken positively, has the form $4m+3$. By contrast, no general rule whatever has yet been found for the sign with which the root of the even square, $f$, emerges from this operation, either a priori or by examining examples. At the end of the paper the author therefore recommends this subject to the friends of higher arithmetic for further investigation, convinced that success would also open up a fruitful source of new extensions of this beautiful part of mathematics.

---

Göttingische gelehrte Anzeigen. 1831, April 23.

A paper presented to the Royal Society on April 15 by Court Councillor Gauss:

*Theoria residuorum biquadraticorum, commentatio secunda,*

continues the paper already printed in the sixth volume of the *Commentationes novae*, which was also reviewed in these pages at the time, in 1825, on April 11. Even this continuation, although more than twice as long as the first paper, does not yet exhaust this exceptionally rich subject, and completion of the whole will remain reserved for a future third paper.

Although the basic concepts of these theories and the contents of the first paper may be assumed known to everyone who has made higher arithmetic a subject of study, we nevertheless wish to recall those concepts briefly here for the convenience of readers interested in this part of mathematics who do not have the first paper immediately to hand. With respect to any integer $p$, another integer $k$ is called a biquadratic residue if there are numbers of the form $x^4 - k$ that are divisible by $p$; in the opposite case it is called a biquadratic nonresidue of $p$. It is sufficient here to restrict attention to the case where $p$ is a prime of the form $4n + 1$, and $k$ is not divisible by that prime, since all other cases are either clear in themselves or can be reduced to this one.

For such a *given* value of $p$, all numbers not divisible by $p$ fall into *four* classes. One contains the biquadratic residues, a second contains those biquadratic nonresidues that are quadratic residues of $p$, and the biquadratic nonresidues that are also quadratic nonresidues are divided between the remaining two. The principle of this division is that either $k^n - 1$, or $k^n + 1$, or $k^n - f$, or $k^n + f$ will always be divisible by $p$, where $f$ denotes an integer that makes $ff + 1$ divisible by $p$. Anyone familiar with the elementary terminology will see how these verbal definitions can be expressed in it.

The theory of this classification is completed in the first paper, both for the immediately evident case $k = -1$ and for the cases $k = \pm 2$, which require subtle auxiliary investigations. The beginning of the present paper now proceeds to larger values of $k$. At first, however, only values that are themselves primes need be considered, and the outcome shows that the results take their simplest form when the values are taken positively or negatively according as their absolute values have the form $4m + 1$ or $4m + 3$. Examining examples immediately yields a rich harvest of new theorems with great ease; we give only a few of them here. The numbering of the classes as $1, 2, 3, 4$ refers to the cases where $k^n$ is congruent to $1, f, -1, -f$, respectively. At the same

time, the value always chosen for $f$ is the one that makes $a + bf$ divisible by $p$, where $aa + bb$ represents the decomposition of $p$ into an odd and an even square. Thus examining examples shows that $-3$ always belongs to class $1, 2, 3, 4$ according as $b, a + b, a, a - b$, respectively, is divisible by 3; that $+5$ belongs to these classes in order according as $b, a - b, a, a + b$, respectively, is divisible by 5; and that $-7$ falls into class 1 when $a$ or $b$; into class 2 when $a - 2b$ or $a - 3b$; into class 3 when $a - b$ or $a + b$; and into class 4 when $a + 2b$ or $a + 3b$ is divisible by 7, divisibility being the condition in each of these cases. Similar theorems result for $-11, +13, +17, -19, -23$, and so on. Yet, easy as it is to discover all such particular theorems by examining examples, it seems difficult to find a general law for these forms by that route, even though some common features soon become apparent; and finding proofs of these theorems is much more difficult still. The methods used for $+2$ and $-2$ in the first paper no longer apply here. Although other methods might also settle what concerns the first and third classes, they prove inadequate for establishing *complete* proofs.

One therefore soon recognizes that this rich region of higher arithmetic can be entered only by entirely new routes. In the first paper, the author had already indicated that a particular extension of the whole field of higher arithmetic was essential for this purpose, without then explaining more closely what it consisted in. The present paper is intended to bring this subject to light.

This consists in nothing other than the following: to give the theory of biquadratic residues its true foundation, the field of higher arithmetic, which had otherwise been extended only to the real integers, must also be extended to the imaginary ones, and these must be granted entirely equal rights of citizenship with the former. Once this is understood, that theory appears in an entirely new light, and its results acquire a most surprising simplicity.

Before the theory of biquadratic residues itself can be developed in this extended domain of numbers, however, the preceding parts of higher arithmetic, which have so far been treated only in relation to real numbers, must share in this extension. We can mention only a few of these preliminary investigations here. The author calls a quantity $a + bi$, where $a$ and $b$ denote real quantities and $i$ is written for $\sqrt{-1}$ for brevity, a complex integer when $a$ and $b$ are both integers. Complex quantities thus do not stand opposed to real ones: they include them as the special case where $b = 0$. For convenient use, several concepts relating to complex quantities needed particular names, but we shall try to avoid those names in this review.

Just as the arithmetic of real numbers speaks of only two units, the positive and the negative, so in the arithmetic of complex numbers we have four units, $+1$, $-1$, $+i$, $-i$. A complex integer is called *composite* if it is the product of two integer factors that are not units; a complex number that does not

admit *such* a factorization is called a complex prime. For example, the real number 3 is prime even when considered as a complex number, whereas 5 is composite as a complex number $= (1 + 2i)(1 - 2i)$. Just as in the higher arithmetic of real numbers, primes play a principal role in this extended field of the science.

If a complex integer $a + bi$ is taken as modulus, one can give $aa + bb$ pairwise incongruent complex numbers, and no more, to one of which any given complex integer must be congruent; these can be called a complete system of incongruent residues. The so-called least and absolutely least residues in the arithmetic of real numbers have their exact counterparts here as well. For example, for the modulus $1+2i$, the complete system of absolutely least residues consists of 0, 1, $i$, $-1$ and $-i$. Almost all the investigations in the first four sections of the *Disquisitiones Arithmeticae* also have their place, with some modifications, in the extended arithmetic. The celebrated theorem of FERMAT, for example, takes the following form here: if $a + bi$ is a complex prime, and $k$ a complex number not divisible by it, then always $k^{aa+bb-1} \equiv 1$ for the modulus $a + bi$. Especially remarkable, however, is that the Fundamental Theorem for quadratic residues has its exact counterpart in the arithmetic of complex numbers, in an even simpler form: if $a + bi$, $A + Bi$ are complex primes, with $a$ and $A$ odd, $b$ and $B$ even, then the first is a quadratic residue of the second if the second is a quadratic residue of the first, whereas the first is a quadratic nonresidue of the second if the second is a quadratic nonresidue of the first.

After these preliminary investigations, the paper turns to the theory of biquadratic residues itself. In place of the mere distinction between biquadratic residues and nonresidues, it first establishes a division of the numbers not divisible by the modulus into four classes. If the modulus is a complex prime $a + bi$, with $a$ always assumed odd and $b$ even, and $p$ written for $aa + bb$ for brevity, and if $k$ is a complex number not divisible by $a + bi$, then $k^{\frac{1}{4}(p-1)}$ will always be congruent to one of $+1, +i, -1, -i$. This establishes a division of all numbers not divisible by $a + bi$ into four classes, to which the biquadratic characters $0, 1, 2, 3$ are assigned in order. Clearly, character 0 refers to the biquadratic residues, the others to the biquadratic nonresidues, with character 2 also corresponding to quadratic $\cdot$ residues, and characters 1 and 3 to quadratic nonresidues.

It is easy to see that the main task is to determine this character only for values of $k$ that are themselves complex primes, and here examining examples immediately leads to exceedingly simple results.

If first $k = 1 + i$, the character of this number is always found to be $\equiv \frac{1}{8}(-aa + 2ab - 3bb + 1) (\text{mod. } 4)$, and similar expressions are found for the cases $k = 1 - i$, $k = -1 + i$, $k = -1 - i$.

If, on the other hand, $k = \alpha + \beta i$ is a prime with $\alpha$ odd and $\beta$ even, examining examples very easily yields a law of reciprocity entirely analogous to the Fundamental Theorem for quadratic residues. It can be expressed most simply as follows:

If both $\alpha + \beta - 1$ and $a + b - 1$ are divisible by 4 (all other cases can easily be reduced to this one), and the character of $\alpha + \beta i$ with respect to the modulus $a + bi$ is denoted by $\lambda$, whereas the character of $a + bi$ with respect to the modulus $\alpha + \beta i$ is denoted by $l$, then $\lambda = l$ if one of $\beta, b$ (or both) is divisible by 4. whereas $\lambda = l \pm 2$ if neither of $\beta, b$ is divisible by 4.

These theorems contain essentially everything of substance in the theory of biquadratic residues. Yet, easy as it was to discover them by examining examples, it is difficult to give rigorous proofs of them, especially of the second, the Fundamental Theorem of biquadratic residues. Because of the great length the present paper has already reached, the author was compelled to leave the presentation of the proof of the latter theorem, which he has possessed for 20 years, for a future third paper. The present paper does, however, give the complete proof of the former theorem concerning $1 + i$ (on which the other theorems for $1 - i$, $-1 + i$, $-1 - i$ depend), and this can already give some idea of the subject's complexity.

We still have a few general remarks to add. Moving the theory of biquadratic residues into the domain of complex numbers might seem objectionable and unnatural to someone less familiar with the nature of imaginary quantities and caught up in false ideas about them. It might suggest that the investigation is thereby left, as it were, hanging in the air, becomes unstable, and moves entirely beyond intuitive understanding. Nothing could be less well founded than such an opinion. On the contrary, the arithmetic of complex numbers admits the clearest visual representation. Although the author has followed a purely arithmetical treatment in this presentation, he has also given the necessary indications for this visual representation, which makes understanding more vivid and is therefore highly to be recommended. these will suffice for readers who think for themselves. Just as absolute integers are represented by a row of equally spaced points on a straight line, in which the starting point represents 0, the next point 1, and so on; and just as representing negative numbers then requires only an unbounded extension of this row on the opposite side of the starting point: so representing complex integers requires only the addition that this row be regarded as lying in a fixed unbounded plane, and that an unlimited number of similar rows be taken parallel to it on both sides, at equal distances from one another. In place of a row of points, we then have a system of points that can be arranged as rows of rows in two ways, and that serves to divide the whole plane into equal squares. The point nearest 0 in the first adjacent row on one side of the row representing the real numbers then corresponds to $i$, just as the point

nearest 0 in the first adjacent row on the other side corresponds to $-i$, and so on. With this representation, the execution of arithmetical operations on complex quantities, congruence, the formation of a complete system of incongruent numbers for a given modulus, and so on, admit a visual representation that leaves nothing to be desired.

On the other hand, this places the true metaphysics of imaginary quantities in a new, bright light.

Our general arithmetic, whose scope far surpasses the geometry of the ancients, is entirely a creation of modern times. Starting originally from the concept of absolute integers, it has extended its domain step by step: fractions were added to integers, irrational numbers to rational numbers, negative numbers to positive numbers, imaginary numbers to real numbers. Yet each advance was initially made with timid, hesitant steps. The first algebraists still called the negative roots of equations false roots, and they are indeed such when the problem to which they relate is presented in a form where the nature of the quantity sought admits no opposite. But just as there is no hesitation about including fractions in *general* arithmetic, although there are so many countable things for which a fraction has no meaning, so negative numbers could not be denied equal rights with positive ones in that arithmetic because countless things admit no opposite: the reality of negative numbers is sufficiently justified, since they find an adequate substrate in countless other cases. All this has, of course, long been understood. But imaginary quantities, set against real ones—formerly. and here and there even now, though inappropriately, called *impossible*—are still more tolerated than naturalized as citizens. They thus appear more like a play of signs empty of content in itself, to which any conceivable substrate is unconditionally denied, without anyone wishing to spurn the rich tribute this play of signs ultimately contributes to the treasury of relations among real quantities.

For many years the author has viewed this highly important part of mathematics from a different standpoint, from which imaginary quantities can be given an object just as well as negative ones. Until now, however, there has been no occasion to state it explicitly in public, although attentive readers will readily find traces of it in the paper on equations published in 1799 and in the prize essay on the transformation of surfaces. The present paper briefly indicates its main features, which are as follows.

Positive and negative numbers can have an application only where what is counted has an opposite that, conceived as combined with it, is equivalent to cancellation. Strictly considered, this condition holds only where what is counted consists of relations between pairs of objects, rather than substances (objects conceivable in themselves). It is postulated that these objects are arranged in a row in a definite way, for example $A, B, C, D \ldots$., and that the relation of $A$ to $B$ can be regarded as equal to the relation of $B$ to $C$, and so

on. Here the concept of opposition requires nothing more than *interchanging* the members of the relation, so that if the relation (or transition) from $A$ to $B$ counts as $+1$, the relation from $B$ to $A$ must be represented by $-1$. Thus, insofar as such a row is unbounded on both sides, every real integer represents the relation of a member chosen arbitrarily as starting point to a definite member of the row.

But if the objects are of a kind that cannot be arranged in a single row, even an unbounded one, and can only be arranged in rows of rows, or, equivalently, form a manifold of two dimensions; and if the relations of one row to another, or the transitions from one to another, behave similarly to the earlier transitions from one member of a row to another member of the same row, then measuring the transition from one member of the system to another clearly requires, besides the former units $+1$ and $-1$, two further units $+i$ and $-i$, likewise opposite to one another. But it must also be postulated that the unit $i$ always denotes the transition from a given member of a row to a *definite* member of the immediately adjacent row. In this way the system can therefore be arranged as rows of rows in two ways.

The mathematician abstracts entirely from the nature of the objects and the content of their relations; the sole concern is to count the relations and compare them with one another. In this respect, just as the relations denoted by $+1$ and $-1$, considered in themselves, are taken to be of the same kind, the mathematician is entitled to extend this to all four elements $+1$, $-1$, $+i$ and $-i$.

These relations can be brought within intuition only by a representation in space. The simplest case is where there is no reason to arrange the symbols of the objects in any way other than a square pattern: an unbounded plane is divided into squares by two systems of parallel lines crossing at right angles, and the intersections are chosen as the symbols. Each such point $A$ has four neighbours. If the relation of $A$ to one neighbouring point is denoted by $+1$, the relation to be denoted by $-1$ is thereby determined, whereas either of the other two may be chosen for $+i$, or the point corresponding to $+i$ may be taken at will to the *right* or to the *left*. Once forward and backward *in* the plane, and above and below with respect to its two sides, have been fixed (at will), this distinction between right and left is entirely determined *in itself*, although we can communicate our intuition of this distinction to others *only* by pointing to actually existing material things[*)]. Even after we have decided the latter point, however, we see that it depended on our arbitrary choice which of the two rows crossing at a point we wished to regard as the

[*)] KANT has already made both observations, but one cannot understand how this perceptive philosopher could believe he found in the first a proof of his opinion that space is *only* the form of our outer intuition, since the second so clearly proves the opposite, and that space must have a real meaning independent of our mode of intuition.

principal row, and which direction in it as corresponding to positive numbers. We also see that if we wish to take the relation previously treated as $+i$ for $+1$, we must necessarily take the relation previously denoted by $-1$ for $+i$. In mathematical language, this says that $+i$ is a mean proportional between $+1$ and $-1$, or corresponds to the sign $\sqrt{-1}$: we deliberately do not say *the* mean proportional, for $-i$ clearly has an equal claim. The possibility of giving $\sqrt{-1}$ an intuitive meaning is thus fully justified here, and nothing more is required to admit this quantity into the domain of the objects of arithmetic.

We believed that this brief presentation of the main points of a new theory of so-called imaginary quantities would serve readers interested in mathematics. If this subject has hitherto been viewed from a mistaken standpoint and mysterious obscurity has been found in it, this is largely attributable to the unsuitable names. Had $+1$, $-1$, $\sqrt{-1}$ been called direct, inverse and lateral units, for example, instead of positive, negative and imaginary (or even impossible) units, such obscurity could scarcely have been spoken of. The author has reserved a fuller treatment of the subject, which is really touched on only incidentally in the present paper, for the future. The question why relations among things that form a manifold of more than two dimensions cannot yield further kinds of quantities admissible in general arithmetic will then also receive its answer.

---

# REVIEWS

## OF OTHER AUTHORS'

# WRITINGS.

Göttingische gelehrte Anzeigen. 1809, March 11.

*Recherches sur l'irréductibilité Arithmétique et Géométrique des nombres et de leurs puissances.* 1808. (No place of publication. 25 pages in large quarto.)

A work whose aim is to represent irrational radical quantities in the form of rational quantities. We must content ourselves with having drawn the attention of readers interested in mathematics to this little work, since the limits of these pages do not allow us to give a fuller account and examination here of the author's distinctive standpoint and treatment of radical quantities, which departs entirely from the usual one.

Göttingische Gelehrte Anzeigen. 1812, March 23.

*Cribrum Arithmeticum, sive tabula continens numeros primos a compositis segregatos, occurrentes in serie numerorum ab unitate progredientium usque ad decies centena milia et ultra haec ad viginti millia* (1020000). *Numeris compositis, per* 2, 3, 5 *non dividuis, adscripti sunt divisores simplices, non minimi tantum, sed omnino omnes. Confecit* Ladislaus Chernac, *Pannonius, A. L. M. Philos. et Medic. Doctor, in almo lyceo Daventriensi philosophiae professor. Daventriae* 1811. (At the author's expense, printed by J. H. Lange. XXII and 1022 pages, large quarto.)

The full title of this important and highly meritorious work already sufficiently indicates its contents: it is a table of all the prime factors of all numbers not divisible by 2, 3 and 5, from 1 to 1020000, calculated through several years of work as careful as it was laborious, and printed neatly and, as far as we have found by occasional checks, very correctly. Anyone who frequently carries out calculations with larger numbers can readily judge the value of such a gift to arithmetic. The author deserves double thanks: both for the exceedingly laborious work itself, by which he has placed his name beside the unforgettable names of Rhaeticus, Pitiscus, Brigg, Vlacq, Wolfram, Taylor and others, and for the certainly very considerable expense incurred in printing it, for which a publisher would otherwise scarcely have been found. Such tables have often been calculated before, though mostly on a smaller scale, but have either remained entirely in manuscript or have not been completely printed. As is well known, Lambert formerly did all he could to encourage the continuation of Pell's often reprinted table, which extends to 100000. According to a report by Bernoulli in Lambert's correspondence,

Oberreit had extended it to 500000, and a copy had come into Schulze's hands. Anton Felkel, as reported in the Monatl. Correspondenz, volume 2, page 223, had completed it in manuscript up to two million and intended later to provide it up to 2460000; but what had already been printed in Vienna at public expense was used for cartridges in the Turkish war because no buyers could be found! Thus many years of meritorious work were lost to the public: all the more, therefore, we considered it our duty to announce the publication of the present work here. The first million is now available for everyone's use; anyone with the opportunity and enthusiasm for this subject may therefore direct their efforts to what follows.

---

Göttingische gelehrte Anzeigen. 1814, November 3.

---

*Tables des diviseurs pour tous les nombres du deuxième million, ou plus exactement depuis* 1020000 *à* 2028000, *avec les nombres premiers qui s'y trouvent. Par* J. Ch. Burckhardt, *membre de l'institut impérial, du bureau des longitudes de France, et de plusieurs autres sociétés savantes. Paris*, 1814. *M^me V^e Courcier.* (VIII. and 112 pages in folio.)

Sooner than we would have dared to hope when reviewing Chernac's factor table covering the first million, we can already report the completion and publication of a similar table for the second million. Through this laborious work the distinguished author, whose name itself guarantees the greatest care and accuracy, has placed all those interested in arithmetic deeply in his debt. Chernac's table for the first million gives all the prime factors; Burckhardt's for the second, however, gives only the least divisor in each case. The complete factorization of a number in the second million therefore requires division by the least divisor and looking up the quotient in Chernac's table. But this small effort is of no significance compared with the great advantage of having the table occupy so much less space, while the prospect remains of seeing it extended in time to ten million. The author made it possible to fit the table into this small volume partly by restricting it to the least divisor and partly by printing it as economically as possible. If $a$ stands for any of the eighty numbers below 300 that are not divisible by 2, 3 and 5, then every number not divisible by 2, 3 and 5 is included in the form $300n + a$. All eighty numbers for which $n$ has the same value are found in a single vertical column, and each page contains thirty such columns. Thus each page includes, out of nine thousand numbers proceeding in their natural order, all those not divisible by 2, 3 or 5.

The method by which Mr. BURCKHARDT constructed his table deserves special mention here. He had a grid engraved on copper, in which 81 horizontal and 78 vertical lines formed a rectangle divided into $80 \times 77$, that is, 6160 small squares, and had the necessary number of impressions made from it. At the side, the eighty values of $a$ could also be engraved immediately; the values of $300n$, in continuous order, were written with a pen above the 77 vertical columns. Thus each sheet represents all the numbers not divisible by 2, 3 and 5 found among each set of 23100 numbers proceeding in their natural order, and 44 sheets suffice to cover a whole million. It is easy to see that the numbers whose least divisor is 7 or 11 recur in the same order on every following sheet. These divisors could therefore be engraved directly on the copperplate and consequently appeared automatically in their proper places on every sheet. To enter the following divisors, for example 13, Mr. B. took only 13 columns across the width of a spare sheet and, treating it as the beginning of his table, cut out all the squares that had to contain the divisor 13. He therefore needed only to lay this stencil over the first thirteen columns of the first sheet, then over the next thirteen, and so on, to see at once all the places that had to be filled, insofar as they did not already contain 7 or 11, with 13. The same procedure was then used for the divisor 17, and so on. The spare sheets sufficed in this way up to the divisor 73; for the larger divisors 79, 83, and so on, Mr. B. appears to have assembled the frame from two or more pieces. For divisors exceeding 500, however, Mr. B. preferred to find the multiples by addition, needing to take only primes for the other factor. We find this whole procedure highly suitable, and would recommend it for imitation to anyone inclined to extend the table further. For the third and fourth millions, the author himself has meanwhile already carried out a large part of the calculations, so we have good reason to hope to see these too made available in print soon.

Göttingische gelehrte Anzeigen. 1816, November 7.

*Tables des diviseurs pour tous les nombres du troisième million, ou plus exactement, depuis* 2028000 *à* 3036000, *avec les nombres premiers qui s'y trouvent; par* J. Chr. Burckhardt, *membre de l'académie royale des sciences, du bureau des longitudes de France et de plusieurs autres sociétés savantes. Paris* 1816. *M^me V^e Courcier.* (112 pages in folio.)

Since in our review of the table of factors for the second million we have already described in detail both the method of calculation used by the distinguished author and the arrangement of the table itself, we can here content ourselves with announcing the publication of the table for the third million. Shortly we may also expect from the author the table for the *first* million, presented in the same way, so that the whole table extending beyond three million will then occupy only a moderately sized volume. For this the author deserves the thanks of all those interested in arithmetic, who see this laborious work meeting a need on a scale far exceeding anything one could have dared to hope for only a few years ago.

Göttingische gelehrte Anzeigen. 1817, August 9.

*Tables des diviseurs, pour tous les nombres du premier million, ou plus exactement depuis* 1 *à* 1020000, *avec les nombres premiers qui s'y trouvent; par* J. Chr. Burckhardt, *membre de l'académie des sciences dans l'institut royal, du bureau des longitudes de France, et de plusieurs autres sociétés savantes. Paris* 1817. *M^me V^e Courcier.* (114 pages in folio.)

Referring here to our reviews of the tables for the second and third millions, we now merely announce the actual · publication of these factor tables for the first million. We thus now possess a connected whole for the first three millions. For the present first million, the author used partly Chernac's *Cribrum Arithmeticum* and partly a manuscript table by Schenmark held by the library of the Royal Institute. The latter had not, however, been constructed with quite all the care one could wish for, and deciding which was right in cases where the two disagreed was often rather laborious. Only a very small number of errors were found in Chernac's table, which Mr. Burckhardt has reported here.

The author also already has most of the materials ready for the fourth, fifth and sixth millions, and offers to supply this continuation if the publisher is encouraged by sufficient sales of the first three millions. It would indeed be very regrettable if the fruits of such laborious and useful work were withheld from the world.

Göttingische gelehrte Anzeigen. December 19, 1825.

A short paper has been presented to the Royal Society on behalf of Mr. Erchinger of Thuningen in the Kingdom of Würtemberg, on the

*Geometric Construction of the Regular Seventeen-sided Polygon.*

As is well known, the general theory of regular polygons has acquired a new form and a wider scope through its close connection with higher arithmetic. The theory of those polygons that can be constructed geometrically is one part of it, though a comparatively small one. Since the time of the Greeks it had been known that the triangle, pentagon, fifteen-sided polygon and all polygons obtained from these by doubling or repeatedly doubling the number of sides have this property, and it was believed, and sometimes expressly asserted, that these were the only ones. Higher arithmetic has shown this to be an error: by laying open the true sources of the entirely general theory, it revealed of itself that, besides the polygons mentioned, there are countless others that can be constructed geometrically, of which the seventeen-sided polygon is the simplest. This shows in the clearest light the superiority of analysis, which encompasses the most general and the particular with equal ease, over geometry, which must always remain with the particular, is held back by ever greater complications as it advances from simpler cases to more complex ones, and would hardly ever have reached, without help from elsewhere, that next case after the known ones. Nevertheless, it remains important, interesting and desirable that purely geometric treatments should also continue to be cultivated, and that geometry should take possession of at least part of the new fields conquered by analysis. The reviewer knows of no one who has so far publicly treated the construction of the seventeen-sided polygon except Mr. Pauker, in the writings of the Courland Society and in his geometry. Mr. Erchinger's construction differs from this and is carried out more in the purely geometric spirit. It is as follows. (Anyone can draw the accompanying figure: a straight line on which the points $DBGAIFCE$ lie in that order.) Take any straight line $AB$ and extend it backwards and forwards to $C$ and $D$, so that $AC \times BC = AD \times BD = 4AB \times AB$; next determine the points $E$, $G$ on the two sides of the extended line $CA$, so that $AE \times EC = AG \times CG = AB \times AB$, and the point $F$ on the $A$ side of the extended line $BA$, so that $AF \times DF = AB \times AB$; finally divide $AE$ at $I$, so that $AI \times EI = AB \times AF$, where $AI$ is the smaller and $EI$ the larger part of $AE$. Then construct a triangle in which two sides are each $= AB$, and the third $= AI$. If a circle is drawn around this triangle, $AI$ will be the side of the regular seventeen-sided polygon inscribed in that circle.

If one checks the correctness of this construction by comparing it with the theory of the seventeen-sided polygon given as an example in Article 354 of the *Disquisitiones Arithmeticae*, one readily sees that it is simply the geometric translation of the equations to which the application of the general theory leads. Indeed, the distances of the points $C$, $D$, $E$, $F$, $G$, $I$ from $A$ are precisely the quantities denoted there by (8 . 1), (8 . 3), (4 . 1), (4 . 3), (4 . 9), (2 . 1), if positive and negative signs are expressed by position, and the distance of the point $B$ from $A$, taken in the same sense, is set $= -1$. But the real merit of Mr. ERCHINGER's paper lies less in setting out the construction itself, since analysis had already marked out the simplest path, than in giving a purely geometric proof of its correctness. This is carried out with such exemplary, painstaking care to avoid everything that is not purely elementary that it does the author credit, and prompts the wish that his indeed uncommon mathematical talent may receive every encouragement.

Göttingische gelehrte Anzeigen. July 9, 1831.

*Investigations into the Properties of Positive Ternary Quadratic Forms, by* Ludwig August Seeber, *Doctor of Philosophy, Full Professor of Physics at the University in Freiburg. Freiburg im Breisgau* 1831. (248 pages in 4.)

Functions of two indeterminate quantities $x$ and $y$ of the form $axx+2bxy+cyy$, where $a, b, c$ denote specified integers, are known as *quadratic forms*, or, when a further distinction is needed, *binary quadratic forms*. They form one of the most interesting and richest subjects of higher arithmetic. The problems that arise first are these: to decide whether a given form of this kind contains another, $a'x'x' + 2b'x'y' + c'y'y'$, that is, whether it can be transformed into that form by a substitution $x = \alpha x' + \beta y'$, $y = \gamma x' + \delta y'$, in which $\alpha, \beta, \gamma, \delta$ are integers; whether such a relation between two forms is mutual, in which case the forms are called equivalent; further, in both cases, to give all possible transformations of one into the other; and finally, to find all possible representations of a given integer by a given form with integer values of the indeterminate quantities. These problems are completely solved in the *Disquisitiones Arithmeticae*, but constitute by far the smaller part of that work's section on quadratic forms. The finer investigations that follow require in part a preliminary treatment of a field one level higher and presenting much greater difficulties: the theory of similar functions of three indeterminate quantities $x, y, z$, which therefore have the form $axx + byy + czz+2a'yz+2b'xz+2c'xy$ and are called ternary quadratic forms. The solution of the principal problems concerning these ternary forms is developed in the work mentioned, though only as far as was necessary for the stated purpose. After an interval of thirty years, the author of the present work has now been the first to take up these investigations again and, for one principal kind of ternary form, namely the positive forms, has brought to completion what was left unfinished in the *Disquisitiones Arithmeticae*. For those who have studied higher arithmetic more deeply, we could describe in a few words what is newly achieved in the present work. But to be understood by others too, we must allow ourselves somewhat more detail, and we do this all the more willingly because these investigations have an interest of their own outside higher arithmetic as well.

The properties of a binary form $axx + 2bxy + cyy$ depend chiefly on the number $bb - ac$, which is therefore called the determinant of that form. Two equivalent forms always have equal determinants. But not all forms having a given determinant are therefore equivalent. Rather, such forms fall into a smaller or larger, but always finite, number of classes, with the forms belonging to one class equivalent to one another, and those belonging to different classes not equivalent. Forms whose determinant is positive can represent

positive and negative numbers alike; forms with a negative determinant, however, can represent only numbers having the same sign as $a$ and $c$, so that positive and negative forms are distinguished here. The simplest forms in each class have definite criteria, are called reduced forms, and can be regarded as representatives of the whole class.

Similar relations concerning ternary forms are established in the *Disquisitiones Arithmeticae*. The determinant of the ternary form

$$axx + byy + czz + 2a'yz + 2b'xz + 2c'xy$$

is the number

$$aa'a' + bb'b' + cc'c' - abc - 2a'b'c'$$

Here too, equality of determinants is necessary for two forms to be equivalent, but is not sufficient. Rather, all forms with a specified determinant fall into a finite number of classes, in each of which the simplest forms may be called reduced and may, as it were, represent all the others. The distinction between positive and negative forms, however, behaves differently here than for binary forms. For every given determinant, whether positive or negative, there are some forms that can represent positive and negative numbers alike (indifferent forms), and others that can represent either only positive or only negative numbers (positive or negative forms). But positive forms exist only for negative determinants, and negative forms only for positive ones. It is evident, moreover, that a form's character as indifferent, positive or negative belongs at the same time to the whole class to which it belongs. The present work is confined to positive forms, whose determinants must therefore be negative. Everything that holds for these evidently transfers of itself to negative forms, whereas the indifferent forms, entirely excluded from the work, require a quite different treatment.

As already mentioned, the theory of ternary forms in the *Disquisitiones Arithmeticae* was developed only as far as was needed for its purpose there. Consequently, the problem of deciding whether two given ternary forms are equivalent had not yet been solved in full generality. It was indeed shown there how to find, for any given form, an equivalent form of the simplest kind, and that there can be only finitely many such reduced forms for any given determinant. But since each class contains several such reduced forms, which do not in all cases turn out *immediately* to be equivalent, a criterion was still lacking by which their equivalence or non-equivalence could be determined with certainty. The author of the present work has now met this need for positive forms completely and with exemplary thoroughness. His procedure is, however, presented somewhat differently from the account we have just given, and from how the matter would stand if the concept of reduced positive forms included only the essential conditions of greatest simplicity. For positive

forms, these conditions are that the numbers $a, b, c$, which are positive by their nature, must be no smaller than, respectively, $b'$ or $c'$, $a'$ or $c'$, and $a'$ or $b'$, without regard to signs. Mr. SEEBER has added further qualifications to the concept of reduced forms such that each class can contain only one form of this kind, and must contain one. In view of a beautiful theorem found by Mr. SEEBER from observed cases, to be mentioned further below, we give here the principal conditions that Mr. S. has incorporated into the concept of reduced forms: 1) among the numbers $a', b', c'$, no two may have opposite signs; 2) without regard to sign, $2b'$ and $2c'$ must be no greater than $a$, further $a$ and $2a'$ no greater than $b$, and $b$ no greater than $c$; 3) when $a', b', c'$ are all negative, twice the sum of these numbers must be no greater than $a + b$. We may omit here the remaining qualifications that must be added for certain special cases.

The principal content of the work consists first in solving the problem of finding, for any given positive form, an equivalent form having the character of a reduced form according to the established definition; and then in the rigorous proof of the theorem that two non-identical reduced forms cannot be equivalent, or, what is the same thing, that each class contains only one reduced form. We must do full justice to the spirit of thoroughness with which these matters are carried through. And although we must regret the very great length of treatment involved, perhaps discouraging to some, since the solution of the problem takes 41 pages and the proof of the theorem 91 pages, we would by no means have this taken as a criticism. When a difficult problem or theorem is to be solved or proved, the first step, always to be acknowledged with due thanks, is that a solution or proof should be found at all. The question whether this could have been done more easily and simply remains idle so long as the possibility is not at the same time established by actually doing it. We therefore consider it premature to dwell on this question here. — The remainder of the work chiefly contains equally thorough solutions of the following problems: to decide whether a given form contains another given form not equivalent to it; to find all possible transformations of a given form into a given equivalent form or one merely contained in it; and finally, for a given determinant, to give all possible classes of positive ternary forms.

We must also note that Mr. SEEBER has expressed ternary forms somewhat differently from the *Disquisitiones Arithmeticae*, where the coefficients of the products $yz$, $xz$, $xy$ were deliberately assumed to be even integers, whereas Mr. S. also allows odd ones, and therefore denotes by $a', b', c'$ what was denoted above by $2a'$, $2b'$, $2c'$. The greater generality obtained in this way is evidently only apparent, or at least unnecessary, since everything that can be said about such forms with odd coefficients also follows of itself if their doubles are considered instead. We therefore cannot approve this change,

which also entails some loss of simplicity. One consequence has been that what Mr. SEEBER calls the determinant is always four times the number bearing that name in the *Disquisitiones Arithmeticae*. In the present review we have retained the terminology of the *Disquisitiones Arithmeticae*.

For the last-mentioned problem (to give all possible reduced forms for each given determinant), Mr. SEEBER used a theorem to obtain bounds for the first three coefficients. According to this theorem, their product $abc$ cannot exceed three times the determinant. Mr. SEEBER rigorously proves this theorem. In the preface, however, he remarks that among more than 600 cases he examined, he found not a single one in which that product exceeded twice the determinant. He therefore considers it highly probable that this tighter bound holds universally, but says that he did not succeed in finding a rigorous proof. Since this theorem, found by Mr. SEEBER from observed cases, is both remarkable in itself and important for shortening the solution of the stated problem, we wish to add a very simple proof here, so that we too may contribute in this review to the further development of the theory. Two cases must be distinguished.

I. If none of the numbers $a', b', c'$ is negative, put

$$\begin{aligned} b - 2a' &= d, \quad c - 2b' = e, \quad a - 2c' = f \\ c - 2a' &= g, \quad a - 2b' = h, \quad b - 2c' = i \end{aligned}$$

The definition of reduced positive forms immediately implies that if

$$axx + byy + czz + 2a'yz + 2b'xz + 2c'xy$$

is such a form, none of those six numbers is negative; it is also evident that $a, b, c$ are positive. If we now denote the form's (negative) determinant by $-D$, we have the identity, readily checked by expansion,

$$2D - abc = aa'd + bb'e + cc'f + a'hi + b'gi + c'gh + ghi$$

in which none of the seven terms on the right can be negative, and consequently $abc$ is no greater than $2D$. The same conclusion follows in the same way from the identity

$$2D - abc = aa'g + bb'h + cc'i + a'ef + b'df + c'de + def$$

II. If none of the numbers $a', b', c'$ is positive, put

$$\begin{aligned} b + 2a' &= d, \quad c + 2b' = e, \quad a + 2c' = f \\ c + 2a' &= g, \quad a + 2b' = h, \quad b + 2c' = i \end{aligned}$$

$$\begin{aligned} b + c + 2a' + 2b' + 2c' &= k \\ a + c + 2a' + 2b' + 2c' &= l \\ a + b + 2a' + 2b' + 2c' &= m \end{aligned}$$

and take the determinant of the form, as before, $= -D$. By the definition of reduced positive forms, none of the nine numbers $d, e, f, g, h, i, k, l, m$ can be negative. Thus the identity

$$\begin{aligned} 6D - 3abc = & - aa'(d + 2k) - bb'(e + 2l) - cc'(f + 2m) \\ & - a'hi - b'gi - c'gh + def + 2ghi \end{aligned}$$

in which all terms on the right are positive or zero, since $a', b', c'$ are not positive but negative or zero, shows that $3abc$ cannot exceed $6D$, or $abc$ exceed $2D$. The same conclusion follows likewise from the identity

$$\begin{aligned} 6D - 3abc = & - aa'(g + 2k) - bb'(h + 2l) - cc'(i + 2m) \\ & - a'ef - b'df - c'de + 2def + ghi \end{aligned}$$

Both equations are symmetric. If full symmetry is given up, the proof can be carried out with an even smaller number of terms, for example by the identity

$$\begin{aligned} 8D - 4abc = & - 2aa'(g + k) - 2bb'(e + l) - 4cc'm \\ & + (c + e)df + (c + g)hi \end{aligned}$$

We now wish to add a few remarks on the meaning of positive binary and ternary quadratic forms outside higher arithmetic. It is unnecessary to treat negative forms separately, while indifferent forms lie wholly outside this treatment.

The positive binary form $axx+2bxy+cyy$ represents, in general, the square of the distance between two unspecified points in a plane whose coordinates, relative to two axes inclined to one another at an angle with cosine $= \frac{b}{\sqrt{ac}}$, differ by $x\sqrt{a}$, $y\sqrt{c}$. If $x$ and $y$ are to mean only *integers*, the form therefore refers to a system of points arranged in parallelograms, lying at the intersections of two systems of parallel lines. The lines in each system are equally spaced: the spacings in one, measured parallel to the lines in the second, are $= \sqrt{a}$, and the spacings in the other, measured parallel to the lines in the first, $= \sqrt{c}$; the inclination of the two systems is the one given above. The plane thus appears divided entirely into equal parallelograms whose vertices

make up the system of points, with none of these points lying inside a parallelogram. The determinant taken with positive sign, thus $ac - bb$, is the square of the area of an elementary parallelogram. One and the same system of such points can be divided into parallelograms in infinitely many different ways and thus referred to equally many different forms. All these different forms are, however, what is technically called equivalent, and the area of an elementary parallelogram always remains the same. Two forms that are not equivalent, but of which one contains the other, refer to the same system of points: the first form to the whole system, the second to a part. Two forms that are technically called improperly equivalent (improprie aequivalentes) refer to two equal but oppositely placed systems of points, the plane being thought of as turned over, and so on.

In the same way, the positive ternary form

$$axx + byy + czz + 2a'yz + 2b'xz + 2c'xy$$

represents, in general, the square of the distance between two unspecified points in space whose coordinates relative to three axes (1), (2), (3) give the differences $x\sqrt{a}$, $y\sqrt{b}$, $z\sqrt{c}$: the cosines of the angles between axes (2) and (3), (1) and (3), and (1) and (2) are respectively $\frac{a'}{\sqrt{bc}}$, $\frac{b'}{\sqrt{ac}}$, $\frac{c'}{\sqrt{ab}}$. If $x, y, z$ are to mean only integers, the form refers to a system of points arranged in parallelepipeds, that is, points arising as the intersections of three systems of parallel, equally spaced planes. The whole space thus appears divided into equal parallelepipeds whose vertices make up that system of points, and the square of the volume of an elementary parallelepiped equals the determinant of the ternary form taken with positive sign·. Equivalent forms represent one and the same system of points, referred only to other axes or fundamental planes. In the same way, all the other principal aspects of the theory of ternary forms find their geometric meaning here: one form being contained in another; the representation of a specified number or an indeterminate binary form by a ternary form; the theory of the adjoint ternary forms (formae adiunctae); the disappearance of the distinction between proper and improper equivalence; the nature of reduced forms; and so on. We must, however, confine ourselves to the indications above, especially since the present work, which considers ternary forms solely from a purely arithmetic standpoint, has given only indirect occasion for them. One will at least see from this what a rich field is opened here to investigations that have great theoretical interest in themselves and can also be used for a treatment of all relations among crystal forms that is both convenient and general. This is not the place to enter into the details of this application. We must nevertheless observe that, although $a, b, c, a', b', c'$ were originally assumed to denote integers, the greater part of the theory of ternary forms, and in particular what is required for that application, remains valid independently of this assumption. Indeed, HAÜY's

specifications lead, for most kinds of crystal, to very simple integer values of the coefficients in the ternary forms referring to the corresponding arrangement of the system of points. But the more accurate later measurements of Wollaston, Malus, Biot, Kupffer and others contradict this and make it doubtful whether rational ratios of those coefficients agree with nature in every case. In any case, however, if one does not wish to drop from the theory the restriction to integer values of the coefficients, integers can always be found that come as close as one wishes to the results of measurement, since what matters here is their ratios to one another, rather than their absolute values.

Finally, we wish to give the geometric meaning of Seeber's theorem mentioned above. If a parallelepiped is such that none of its twelve edges (of which four at a time are equal) is greater than any of the twelve face diagonals (which are equal in pairs) or any of the four diagonals of the parallelepiped, then its volume multiplied by $\sqrt{2}$ is no smaller than the volume of a rectangular parallelepiped formed from the same edges.

---

MANUSCRIPT

# PAPERS.

# SOLUTION OF THE CONGRUENCE $X^m - 1 \equiv 0$.
## ANALYSIS OF RESIDUES. CHAPTER SIX. FIRST PART.

---

**237.**

In Chapter III we showed that the congruence $x^n \equiv 1$, when a prime number $p$ is taken as modulus, has $\mu$ roots, where $\mu$ is the greatest common divisor of the numbers $n$ and $p-1$, and that these roots agree exactly with the roots of the congruence $x^\mu \equiv 1$. It therefore suffices to consider the case in which $n$ is a divisor of $p-1$. That the solution not only of the congruence $x^n - 1 \equiv 0$ but of any other congruence for arbitrary moduli can be derived from its solution for prime moduli has already been shown at various places and will be developed more fully below (Chapter VIII).

**238.**

But there is no need to stop even here. In the same Chapter III we explained that the solution of the congruence $x^n \equiv 1$ depends on the solution of similar congruences $x^a \equiv 1$, $x^b \equiv 1$, and so on, where $a, b$, and so on, are primes or powers of primes, and $n$ is the product of these numbers. Namely, if $A, B$, and so on, are respectively any roots of the congruences $x^a \equiv 1$, $x^b \equiv 1$, and so on, their product $AB\ldots$ will be one of the roots of the congruence $x^n \equiv 1$. Our investigations will therefore be confined to solving the congruence $x^n \equiv 1 \pmod{p}$, where $p$ is a prime, and $n$ a prime or a power of a prime and also a divisor of $p-1$.

**239.**

Furthermore, Chapter III shows that among the roots of the congruence $x^n \equiv 1$ there are always some whose powers can represent all the others. Thus, if $r$ denotes a root of this kind (we called it *primitive* above when $n = p-1$, and shall retain this expression here, though in a broader sense), all the roots of the proposed congruence will be

$$1,\ r,\ rr,\ r^3 \quad \ldots\ldots \quad r^{n-1}$$

We must therefore direct every effort towards finding roots of this kind, since once they have been found the others will become evident of themselves. For brevity we denote any power of $r$ by its exponent enclosed in parentheses, so that (0) denotes unity, (1) any primitive root of the congruence $x^n \equiv 1$, (2) the square of (1), and so on. Thus the sequence (0), (1), (2), (3), $\ldots$ $(n-1)$ includes all the roots. It is also known that $(k)$ will always be such a primitive root whenever $k$ is coprime to $n$; that is, in our case, where $n$ is a power of the prime $t$, $= t^\nu$, whenever $t$ does not divide $k$. The symbols (1), (2), and so

on, are evidently undetermined in themselves; but as soon as a definite value is assigned to (1), all the others become determined.

**240.**

Since our aim is to investigate primitive roots before the others, we must first separate them from the rest. This is done by removing from the sequence $(0), (1), (2) \ldots (n-1)$ all terms $(k)$ for which $k$ is divisible by $t$. If $n$ is prime, or $\nu = 1$, only (0) is to be removed. Before proceeding to investigate the remaining roots, however, we earnestly advise the reader to work out several examples, so as to see concretely everything that might otherwise seem too general. We give one example; but this will not make it unnecessary to work out others independently.

Let $p = 29$; $n = 7$. The seven roots of the congruence $x^7 \equiv 1 \pmod{29}$ will be $1, 7, 16, 20, 23, 24, 25$. Since $n$ is prime, all these roots except 1 will be primitive. Thus, if $7 = (1)$, the symbols will have the following meanings:

| (0) | (1) | (2) | (3) | (4) | (5) | (6) |
|---|---|---|---|---|---|---|
| 1 | 7 | 20 | 24 | 23 | 16 | 25 |

Everyone will also remember that the symbols $(n)$ and (0), $(n+1)$ and (1), and so on, and in general $(a)$ and $(b)$, are equivalent whenever $a \equiv b \pmod{n}$.

**241.**

For our purpose, however, we must proceed in yet another way. We retain only those terms $(k)$ for which $k$ is not divisible by $t$, whose number is $\frac{t-1}{t}.n = \lambda$. All these numbers, or numbers congruent to them modulo $n$, can be represented by successive powers of some number. Let this be $= \rho$; then all the primitive roots of the congruence $x^n \equiv 1$ will be denoted as follows:

$$(1) \quad (\rho) \quad (\rho^2) \quad (\rho^3) \quad \ldots. \quad (\rho^{\lambda-1})$$

By this device we completely exclude all nonprimitive roots; the reasons and advantages will become clearer below. In our example we may therefore put $\rho = 3$, and the primitive roots of the congruence $x^7 \equiv 1$ are arranged as follows:

| | | | | | | |
|---|---|---|---|---|---|---|
| | (1) | (3) | $(3^2)$ | $(3^3)$ | $(3^4)$ | $(3^5)$ |
| or | (1) | (3) | (2) | (6) | (4) | (5) |
| which are | 7 | 24 | 20 | 25 | 23 | 16 |

**242.**

To let the reader know the aim of the following investigations, it will be useful to state the theorem that we propose to prove and clarify.

*If the number $\lambda$, which is $= t^{\nu-1}.t-1$, has prime factors $a, b, c, d$, and so on, and $\lambda = a^\alpha b^\beta c^\gamma \ldots$, the solution of the congruence $x^n - 1 \equiv 0$ depends on solving $\alpha + \beta + ..$ congruences of lower degree, of which $\alpha$ have degree $a$, $\beta$ degree $b$, $\gamma$ degree $c$, and so on.*

Thus, in our example, solving the congruence $x^7 \equiv 1$ depends on a congruence of degree two and another of degree three; and it is apparent in general that the degrees of these congruences never depend on the modulus $p$. To reach the proof of this theorem, however, we must first give some propositions concerning the connection between congruences and their roots, although these investigations properly belong to Chapter eight, where they are to be pursued further.

**243.**

THEOREM. *Suppose the congruence*

$$x^m + Ax^{m-1} + Bx^{m-2} + \ldots. + N \equiv 0 \text{ (prime modulus)}$$

*is such that, when the product of the m factors $x-r$, $x-r'$, $x-r''$, $x-r'''\ldots$ is expanded to give $x^m + ax^{m-1} + bx^{m-2} \ldots + n$, we have $A \equiv a$, $B \equiv b$, $C \equiv c$, and so on, modulo $p$. Then the quantities $r, r', r'' \ldots$ will be roots of the proposed congruence, and it will have no others.*

*Proof.* I. We always have

$$\begin{aligned} &x^m + Ax^{m-1} + Bx^{m-2} + \ldots \\ &\quad \equiv x^m + ax^{m-1} + bx^{m-2} + \ldots \pmod{p} \end{aligned}$$

But the latter side of the congruence becomes $= 0$ on setting $x = r$, $x = r'$, $x = r''$, and so on; therefore, for these values of $x$, the former side becomes $\equiv 0 \pmod{p}$. This proves the first part.

II. If another value $\rho$, congruent to none of $r, r'$, and so on, also satisfied the proposed congruence, we would have

$$\begin{aligned} 0 &\equiv \rho^m + A\rho^{m-1} + B\rho^{m-2} + \ldots \\ &\equiv \rho^m + a\rho^{m-1} + b\rho^{m-2} + \ldots \\ &\equiv (\rho - r)(\rho - r')(\rho - r'')(\rho - r''')\ldots \end{aligned}$$

But since none of the factors $\rho - r$, $\rho - r'$, $\rho - r''$, and so on, is $\equiv 0$, their product being $\equiv 0$ is impossible, because $p$ is prime. There are therefore no roots besides $r, r'$, and so on. This proves the second part.

**244.**

PROBLEM. *Let $r, r', r'' \ldots$ be unknown quantities, whose number is $= m$, whose sum is $= \alpha$, sum of squares $= \beta$, sum of cubes $= \gamma \ldots$, and sum of powers with exponent $m$ is $= \mu$. Suppose that these numbers, also $= m$ in number, are not themselves given, but instead other numbers $\alpha', \beta', \gamma'$, and so on, congruent to them individually modulo $p$, where this is a prime and $> m$. Find a congruence of degree $m$ whose roots are $r, r', r''$, and so on.*

*Solution.* Regard $r, r', r''$, and so on, as the roots of an equation

$$x^m + Ax^{m-1} + Bx^{m-2} + Cx^{m-3} + \ldots = 0$$

and determine its coefficients $A, B, C$, and so on, by the known method, using congruence in place of equality; namely, put

$$\begin{aligned}
-A &\equiv \alpha' \\
-2B &\equiv \beta' + A\alpha' \\
-3C &\equiv \gamma' + A\beta' + B\alpha' \\
-4D &\equiv \delta' + A\gamma' + B\beta' + C\alpha' \\
&\text{etc.} \\
-mN &\equiv \mu' + A\lambda' + \text{etc.}
\end{aligned}$$

These coefficients cannot be undetermined, since all the numbers $1, 2, 3 \ldots m < p$. I say that the congruence

$$x^m + Ax^{m-1} + Bx^{m-2} + \ldots + N \equiv 0$$

is the one required.

*Proof.* Let the equation whose roots are $r, r', r'', r'''$, and so on, be

$$x^m + ax^{m-1} + bx^{m-2} + \ldots = 0$$

Then

$$\begin{aligned}
-a &= \alpha \\
-2b &= \beta + a\alpha \\
-3c &= \gamma + a\beta + b\alpha \\
-4d &= \delta + a\gamma + b\beta + c\alpha \\
&\text{etc.}
\end{aligned}$$

It will be evident to everyone from this that

$$a \equiv A, \quad b \equiv B, \quad c \equiv C \text{ etc. } (\text{mod. } p)$$

so, by the preceding section, the numbers $r, r', r''$, and so on, which are roots of the *equation*

$$x^m + ax^{m-1} + bx^{m-2} + \ldots = 0$$

will also be roots of the *congruence*

$$x^m + Ax^{m-1} + Bx^{m-2} + \ldots \equiv 0 \qquad \text{Q. E. D.}$$

We leave the construction of examples to the readers.

**245.**

We return to our purpose. Retaining the symbols used in sections 242 and earlier, we undertake to show that, if $\lambda$ is the product of any factors $efg$, and so on, the primitive roots of the congruence $x^n \equiv 1$, whose number is $\lambda$, can be divided into $e$ classes in such a way that the sums of the roots assigned to each class are given by a congruence of degree $e$. Once these are taken as known, each class can be subdivided into $f$ groups so that the sum of each group is given by a congruence of degree $f$. These groups can be subdivided again, and so on, until the individual roots are reached.

**246.**

*Definition.* We shall call the collection of *all* terms contained in a form $(\rho^{ke+\alpha})$ of this kind (section 241) a *complete period*, or simply a *period*. Here $e$ denotes any divisor of $\lambda$; $\alpha$ any given number; and $k$ all integers from 0 to $\frac{\lambda}{e} - 1$. For brevity we denote such a period by $(e * \alpha)$. Thus, in our example, the terms

| | | | |
|---|---|---|---|
| | $(1), (2), (4)$ | form the periods | $(2 * 0)$ , |
| | $(3), (6), (5)$ | | $(2 * 1)$ |
| and | $(1), (6)$ | form these | $(3 * 0)$ |
| | $(3), (4)$ | | $(3 * 1)$ |
| | $(2), (5)$ | | $(3 * 2)$ |

*Now, if all terms are distributed into periods in any way, and each period is again divided into smaller periods, and so on, we say that what we promised in the preceding section is achieved.*

Before entering into this exposition itself, however, we shall show that, although the formation of such a period depends on the two somewhat arbitrary quantities $r, \rho$, it contains nothing indeterminate: however these quantities are chosen, the same terms always come together in the same period, provided that the number of terms the period is to contain has been prescribed.

The criterion that two terms $A, B$ belong to the same period is that each is contained in a form $(\rho^{ke+\alpha})$ of this kind, or that $A \equiv r^{\rho^{ke+\alpha}}$, $B \equiv r^{\rho^{k'e+\alpha}} \pmod{p}$. Here $r$ is a primitive root of the congruence $x^n \equiv 1 \pmod{p}$, and $\rho$ a primitive root of the congruence $x^\lambda \equiv 1 \pmod{n}$; see above.

We must show that, if, in place of the numbers $r, \rho$, other numbers, say $s, \sigma$, are chosen, then $A$ and $B$ are contained in similar forms $s^{\sigma^{le+\beta}}$, $s^{\sigma^{l'e+\beta}}$.

Let $s^m \equiv r \pmod{p}$; $\sigma^\mu \equiv \rho \pmod{n}$ and $m \equiv \sigma^\zeta \pmod{n}$, which is possible because $r, \rho$ are primitive roots. Moreover, $m$ will be coprime to $n$, and $\mu$ to $\lambda$ (Chapter III). By the appropriate substitutions we obtain

$$A \equiv s^{\sigma^{\mu ke+\mu\alpha+\zeta}}, \qquad B = s^{\sigma^{\mu k'e+\mu\alpha+\zeta}} \qquad \text{Q. E. D.}$$

**247.**

THEOREM. *The product of two similar periods can be formed, independently of the number p, by adding similar periods and given numbers.*

(We call periods similar when they contain equally many terms, or when the number $e$ is the same.)

Example. Let $n = 7$. The product of the periods $(1)+(6)$ and $(2)+(5)$ is, because $(a)\times(b) = (a+b)$, $(3)+(6)+(8)+(11)$, or it consists of the periods $(3)+(4)$ and $(1)+(6)$.

*Proof.* Let $\frac{\lambda}{e} = f$, and let the given periods be $(e*\alpha)$ and $(e*\beta)$, or the sums

$$(\rho^\alpha)+(\rho^{\alpha+e})+(\rho^{\alpha+2e})+\ldots+(\rho^{\alpha+(f-1)e}) \;\ldots\ldots\; P$$
$$(\rho^\beta)+(\rho^{\beta+e})+(\rho^{\beta+2e})+\ldots+(\rho^{\beta+(f-1)e}) \;\ldots\ldots\; Q$$

The product $PQ$ consists of $f^2$ terms. Arrange them as follows. Form $f$ rows, each consisting of $f$ terms. Let the first contain the product of $P$ with $(\rho^\beta)$, the second the product $P\,.\,(\rho^{\beta+e})$, and so on, and so on. In the first row let the product arising from the term $(\rho^\alpha)$ occupy the first place, that from $(\rho^{\alpha+e})$ the second, and the others follow in succession. In the second row assign the first place to the product arising from $(\rho^{\alpha+e})$, the second to that from $(\rho^{\alpha+2e})$, and so on, and finally the last place to that from $(\rho^\alpha)$. Let the third begin with the product arising from $(\rho^{\alpha+2e})$, and continue in this way, following the product from the last term with the products from the first and second terms, and so on, and so on. Thus, denoting the successive terms of the period $P$ by $1, 2, 3 \ldots z$ and those of the period $Q$ by $\mathrm{I}, \mathrm{II}, \mathrm{III}, \ldots . Z$, arrange the terms of $PQ$ as

$$1\,.\,\mathrm{I}+2\,.\,\mathrm{I}+3\,.\,\mathrm{I}+4\,.\,\mathrm{I}+\ldots+z\,.\,\mathrm{I}$$
$$2\,.\,\mathrm{II}+3\,.\,\mathrm{II}+4\,.\,\mathrm{II}+\ldots.+1\,.\,\mathrm{II}$$
$$3\,.\,\mathrm{III}+4\,.\,\mathrm{III}+\ldots+1\,.\,\mathrm{III}+2\,.\,\mathrm{III}$$
$$\text{etc.} \quad \text{etc.}$$

Now collect all terms occupying the same place in the individual rows into $f$ groups; and I say
1° if any term is $\equiv 1$, all the other terms in the same group will also be $\equiv 1$
2° every group in which no term is $\equiv 1$ forms a period. — Once these statements have been proved, we will evidently have achieved our purpose.

The general form of such a group is

$$(\rho^{\alpha+ke}+\rho^{\beta}),\quad (\rho^{\alpha+(k+1)e}+\rho^{\beta+e}),\quad (\rho^{\alpha+(k+2)e}+\rho^{\beta+2e}),$$
$$\dots\ (\rho^{\alpha+(k+f-1)e}+\rho^{\beta+(f-1)e})$$

Indeed, in place of $\rho^{\alpha+(k-1)e}$ one can also write $\rho^{\alpha+(k+f-1)e}$, since $ef = \lambda$ and $\rho^{\lambda} \equiv 1 \pmod{n}$, and similarly for the preceding terms. Put $\rho^{\alpha+ke}+\rho^{\beta} \equiv \rho^{\chi} \pmod{n}$, which is permissible unless $\rho^{\alpha+ke}+\rho^{\beta}$ happens to be divisible by $n$[*]. The group can then be represented as $(\rho^{\chi})$, $(\rho^{\chi+e})$, $(\rho^{\chi+2e})\dots(\rho^{\chi+(f-1)e})$, which is evidently the period $(e*\chi)$. But if $\rho^{\alpha+ke}+\rho^{\beta}$ is divisible by $n$, all terms of the group will be $\equiv (0)$, that is, $\equiv 1$. Q. E. D.

*Note.* This proof also gives a very easy method of expanding the product. We shall give another below which lacks this advantage, but seems worthy of consideration for its simplicity.

**248.**

We call all the smaller periods that make up a larger period a system of periods. Thus this name will be given to the periods

$$(ef*\alpha),\quad (ef*f+\alpha),\quad (ef*2f+\alpha)\ \dots.\ (ef*(e-1)f+\alpha)$$

that make up the period $(f*\alpha)$. It will be *properly ordered* if the numbers placed after the sign $*$, here $\alpha$, $f+\alpha$, $2f+\alpha$, follow an arithmetic progression with difference $f$. Finally, systems are *similar* if both their smaller and their larger periods are similar.

THEOREM. *If the periods of two similar, properly ordered systems are multiplied together, namely the first by the first, the second by the second, the third by the third, and so on, the sum of all the products can be formed from given numbers and periods similar to the larger period.*

*Proof.* Let the systems be

$$(ef*\alpha),\quad (ef*\alpha+f),\quad (ef*\alpha+2f)\dots$$
$$(ef*\beta),\quad (ef*\beta+f),\quad (ef*\beta+2f)\dots$$

The products of the individual periods of the first system with the corresponding periods of the second consist, by the preceding section, of integers

[*] The proposition will have to be expressed somewhat differently if $n$ denotes a power of a prime in general; when it is a prime number, however, nothing needs changing.

and similar periods. A little attention to how these periods arise shows that, if $(ef*\alpha)\times(ef*\beta)$ consists of an integer $N$ and periods $(ef*A)$, $(ef*B)$, $(ef*C)$, and so on, the products are then formed as follows:

$$\begin{aligned}&(ef*\alpha+f)\times(ef*\beta+f) \text{ from } N \text{ and periods}\\ &\qquad (ef*A+f),\quad (ef*B+f),\quad (ef*C+f) \text{ etc.}\\ &(ef*\alpha+2f)\times(ef*\beta+2f) \text{ from } N \text{ and periods}\\ &\qquad (ef*A+2f),\quad (ef*B+2f),\quad (ef*C+2f) \text{ etc.}\end{aligned}$$

and in general

$$\begin{aligned}&(ef*\alpha+\mu f)\times(ef*\beta+\mu f) \text{ from } N \text{ and periods}\\ &\qquad (ef*A+\mu f),\quad (ef*B+\mu f),\quad (ef*C+\mu f) \text{ etc.}\end{aligned}$$

It follows directly that the sum of all the periods will be

$$eN+(f*A)+(f*B)+(f*C) \text{ etc.} \qquad \text{Q. E. D.}$$

This proof also supplies a method for finding that sum.

**249.**

It is easy to make this theorem still more general. Namely, if there are any number of similar, properly ordered systems, and the products of all the first periods, all the second periods, and so on, are formed, the sum of all these products consists of numbers and larger periods. If all these systems are taken to be equal, the sum of powers of all the periods, whatever the exponent, will consist of numbers and periods similar to the larger one. The aim of these results is now apparent. Let $\lambda = efgh\ldots$. Divide all the primitive roots into $e$ periods $A, A', A''$, and so on, each of these again into $f$: $B, B', B''$, and so on, and each of these into $g$: $C, C', C''$, and so on. The sum of all the periods is given: it is $\equiv -1$. But according to what we have just said, we shall also have

$$\begin{gathered}(A)^2+(A')^2+(A'')^2+(A''')^2+\text{etc.}\\ (A)^3+(A')^3+(A'')^3+(A''')^3+\text{etc.}\\ \text{etc. etc.}\end{gathered}$$

Hence, by section 244, a congruence of degree $e$ can be found whose roots are $A, A', A''$, and so on. Taking these as known, let each period be divided into smaller ones:

$$\begin{array}{lllll} A & \text{into} & B, & B', & B''\ldots\\ A' & \text{into} & B^{(n)}, & B^{(n+1)}, & B^{(n+2)}\ldots\\ & \text{etc.} & & & \end{array}$$

Thus $B + B' + B'' + \ldots \equiv A$ is given. But

$$(B)^2 + (B')^2 + (B'')^2 + \ldots$$
$$(B)^3 + (B')^3 + (B'')^3 + \ldots$$
$$\text{etc.}$$

consists of units and the periods $A, A', A''$, and so on. Therefore $B, B', B''$, and so on, will be given by a congruence of degree $f$, from which they can be found; and similarly the periods making up $A', A''$, and so on, can be determined. Anyone will now see that every period can be subdivided into smaller ones by precisely the same method, until the roots themselves are reached.

**250.**

But a difficulty occurs in applying these rules, and we must remove it. Since each congruence has several roots, we must decide which label to assign to each, so that they can be properly distinguished from one another. The labels of the periods depend on the numbers $r, \rho$, which can be chosen at will, so their labelling must also involve something arbitrary. The number $\rho$ must, however, be fixed at the outset. The main feature of our method is that we derive smaller periods from larger ones. But this cannot be done without the proper order of the periods, which we have obtained through their *labels*. We must therefore ensure that all periods are distinguished by their labels as soon as they have been found.

Let the period $A$ be denoted by $(e * \alpha)$ and divided into $f$ periods $B, C, D$, and so on, which must be labelled. Each is clearly contained in a form $(ef * ke + \alpha)$ of this kind. But I say that, for one of them, $B$, the number $k$ can be chosen at will, and the placement of the others derived from it.

Let $R$ be any primitive root of the congruence $x^n \equiv 1$, and let $B$ consist of the terms $R^\mu + R^\nu + \text{etc.}$. Put $\frac{1}{\mu}\rho^{ke+\alpha} \equiv \frac{1}{\upsilon} \pmod{n}$, and, since the value of $r$ is arbitrary, provided only that $A$ receives the label $(e * \alpha)$, which evidently happens of itself, put $r \equiv R^\upsilon \pmod{p}$. Then the first term of $B$ will be $r^{\rho^{ke+\alpha}}$, and $B$ may be denoted by $(ef * ke + \alpha)$. If, in place of $R^\mu$, we had considered the term $R^\nu$, we would have obtained a different value of $r$. But it is readily seen that, for any root $\rho$, the root $r$ can have $\frac{\lambda}{ef}$ different values.

**251.**

Let us now see how, once one period is labelled, the others are distinguished by their labels. For this purpose, however, we must seek another method of finding the remaining periods: insofar as they, like $A$ itself, are roots of a congruence, no order is discernible among them. Suppose that $A$ is denoted by $(ef*0)$. It follows from the preceding discussion that

$$\begin{aligned} A^2 &\text{ has the form } M + N(ef*0) + O(ef*1) + P(ef*2) + \dots \\ A^3 &\text{ has the form } M' + N'(ef*0) + O'(ef*1) + \dots \\ &\text{etc.} \\ A^{ef-1} &\text{ has the form } M^* + N^*(ef*0) + O^*(ef*1) + \dots \end{aligned}$$

To these is added the congruence

$$(ef*0) + (ef*1) + \dots + (ef*ef-1) \equiv -1$$

We thus have $cf - 1$ linear congruences and the same number of unknown quantities, which can therefore be determined by elimination.

*Note.* A case may occur in which the unknown quantities are given by expressions of the form $\frac{V}{Wp}$. We shall show below how this difficulty can be remedied. Since this case can occur only very rarely, we do not wish to dwell on it here.

**252.**

Let this suffice in general for solving pure congruences. Much more will be said about them at various places below; in particular, many things can be taken over from the solution of pure equations, which we shall not neglect to note in their proper places. We add another example: comparing it with the rules will make everything clearer to those with less experience.

Let $n = 31$, $p = 311$; that is, we seek the roots of the congruence $x^{31} - 1 \equiv 0 \pmod{311}$. We must first find a primitive root of the congruence $y^{30} - 1 \equiv \pmod{31}$, such as $y \equiv 3$. Thus put $\rho \equiv 3$, and first divide all the primitive roots of the proposed congruence into 5 periods, namely

$$\begin{array}{ll} (5*0) & \dots\dots \quad (1) + (26) + (25) + (30) + (5) + (6) \\ (5*1) & \dots\dots \quad (3) + (16) + (13) + (28) + (15) + (18) \\ (5*2) & \dots\dots \quad (9) + (17) + (8) + (22) + (14) + (23) \\ (5*3) & \dots\dots \quad (27) + (20) + (24) + (4) + (11) + (7) \\ (5*4) & \dots\dots \quad (19) + (29) + (10) + (12) + (2) + (21) \end{array}$$

The required calculations give the sum of the periods $\equiv -1$, of their squares $\equiv 25$, cubes $\equiv 26$, fourth powers $\equiv 249$, and fifth powers $\equiv 564$.

Thus the periods will be the roots of the congruence

$$x^5 + x^4 - 12x^3 - 21x^2 + x + 5 \equiv 0$$

Further, one finds

$$(5*0)^2 \equiv 6 + 2(5*0) + 2(5*3) + (5*4)$$
$$(5*0)^3 \equiv 12 + 15(5*0) + 4(5*1) + 3(5*2) + 6(5*3) + 6(5*4)$$
$$(5*0)^4 \equiv 90 + 60(5*0) + 28(5*1) + 26(5*2) + 49(5*3) + 38(5*4)$$

and hence, by elimination,

$$5(5*1) \equiv 3(5*0)^4 - (5*0)^3 - 33(5*0)^2 - 24(5*0) + 15$$
$$5(5*2) \equiv -2(5*0)^4 - (5*0)^3 + 22(5*0)^2 + 31(5*0)$$
$$5(5*3) \equiv (5*0)^4 - 2(5*0)^3$$
$$5(5*4) \equiv -2(5*0)^4 + 4(5*0)^3$$

One root of the congruence found above is $\equiv 17$. Thus, if $(5*0) \equiv 17$, then $(5*1) \equiv 183$, $(5*2) \equiv 263$, $(5*3) \equiv 91$, $(5*4) \equiv 67$.

Now divide each of the periods found into three again, namely

| | |
|---|---|
| $(5*0)$ | into $(15*0),\ (15*5),\ (15*10)$ |
| | or into $(1)+(30),\ (26)+(5),\ (25)+(6)$ |
| $(5*1)$ | into $(15*1),\ (15*6),\ (15*11)$ |
| | or into $(3)+(28),\ (16)+(15),\ (13)+(18)$ |
| | etc. etc. |

Suppose that the periods into which the following are divided are roots of the respective congruences:

| | | |
|---|---|---|
| $(5*0)$ | are roots of | $x^3 + Ax^2 + Bx + C \equiv 0$ |
| $(5*1)$ | | $x^3 + A'x^2 + B'x + C' \equiv 0$ |
| $(5*2)$ | | $x^3 + A''x^2 + B''x + C'' \equiv 0$ |
| | etc. | |

Then

| | | |
|---|---|---|
| $A \equiv -(5*0),$ | $B \equiv (5*0) + (5*3),$ | $C \equiv -2 - (5*4)$ |
| $A' \equiv -(5*1),$ | $B' \equiv (5*1) + (5*4),$ | $C' \equiv -2 - (5*0)$ |
| etc. | etc. | etc. |

Therefore

| | | |
|---|---|---|
| $(15*0),\ (15*5),\ (15*10)$ | will be roots of congruences | $x^3 - 17x^2 + 108x - 60 \equiv 0$ |
| $(15*1),\ (15*6),\ (15*11)$ | | |
| $(15*2),\ (15*7),\ (15*12)$ | | |
| $(15*3),\ (15*8),\ (15*13)$ | | |
| $(15*4),\ (15*9),\ (15*14)$ | | |

Here, however, we have

$$(15*0)^3 - 3(15*0) \equiv (15*1)$$
$$(15*1)^3 - 3(15*1) \equiv (15*2)$$
$$\text{etc.}$$

Hence, if one of the roots of the first congruence, 10, is taken as $(15*0)$, we have

| | | | |
|---|---|---|---|
| $(15*0) \equiv$ | $10$ | $(15*5) \equiv$ | $(15*10) \equiv$ |
| $(15*1) \equiv$ | $37$ | $(15*6) \equiv$ | $(15*11) \equiv$ |
| $(15*2) \equiv$ | $-151$ | $(15*7) \equiv$ | $(15*12) \equiv$ |
| $(15*3) \equiv$ | $-39$ | $(15*8) \equiv$ | $(15*13) \equiv$ |
| $(15*4) \equiv$ | $-112$ | $(15*9) \equiv$ | $(15*14) \equiv$ |

Finally, take the terms making up each of these periods. Then

| | | |
|---|---|---|
| (1), (30) | roots of congruences | $x^2 - (15*0)x + 1 \equiv 0$ |
| (3), (28) | | $x^2 - (15*1)x + 1 \equiv 0$ |
| | etc. | |

The roots of the first congruence are 126 and 195, which will therefore be primitive roots of the congruence $x^{31} \equiv 1$, and from these the remaining roots can readily be derived.

---

# GENERAL INVESTIGATIONS OF CONGRUENCES.

## ANALYSIS OF RESIDUES, CHAPTER EIGHT.

---

**330.**

The results on congruences presented in the preceding sections concern only the simplest cases and were mostly obtained by particular methods. In this section we shall try to bring the theory of congruences under higher principles, as far as is yet possible, much as the theory of *equations* is usually treated. As we have observed several times, there is a remarkable analogy between the two. Since all algebraic congruences involving a single unknown can be reduced to the form

$$X \equiv 0$$

where $X$ is an algebraic function of the unknown $x$ involving no fractions, functions of this kind must be considered first.

**331.**

If $P, Q$ are functions of the indeterminate $x$ of the forms

$$A + Bx + Cxx + Dx^3 + \ldots$$
$$H + Ix + Kxx + Lx^3 + \ldots$$

(which we shall henceforth always mean when we simply say *functions*), and the coefficients of like powers of $x$ in the two functions are congruent for any given modulus, the *functions will be called congruent for that modulus.* Congruent functions clearly take congruent values when equal or congruent values are substituted for the indeterminate. Most of what we proved about *numbers* in Chapters I and II also holds for functions. Thus, if $P \equiv P'$, $Q \equiv Q'$, $R \equiv R'$, and so on, we clearly have $P + Q + R$ etc. $\equiv P' + Q' + R' +$ etc.; $P - Q \equiv P' - Q'$; $PQ \equiv P'Q'$; $PQR$ etc. $\equiv P'Q'R'$ etc. The proofs are very easy and can be given in the same way as in Chapter I.

If $PQ \equiv R$, we shall denote the function $Q$ by $\frac{R}{P}$, with the modulus indicated, and say that $Q$ is the quotient when $R$ is divided by $P$ for this modulus. All functions congruent to $Q$ can clearly be taken in its place; we shall regard them all as a *single* value. Below, however, we shall show in which cases such a quotient can have several values, that is, incongruent ones.

**332.**

If the modulus is a prime number and the divisor $Q$ contains just one term $Hx^h$ whose coefficient $H$ is not divisible by the modulus, that is, provided only that $H$ is not $\equiv 0$, the quotient cannot have several values. For if $QA \equiv P$ and $QB \equiv P$, we would have $Q(A-B) \equiv 0$. Now let

$$Q \equiv .. + Hx^h + Ix^{h+1} + \text{etc.}$$

where $H$ is not divisible by $p$, and

$$A - B \equiv Lx^l + Mx^{l+1} + \text{etc.}$$

where $L$ is not divisible by $p$ (the function $A-B$ will have this form because we assume that $A$ is not $\equiv B$). We would then have $Q(A-B) \equiv HLx^{h+l} + \text{etc.} \equiv 0$, which is absurd, since $HL$ is not $\equiv 0$.

Rules for dividing the function $P$ by $Q$, when this is possible, are now easily given. Let

$$P \equiv ax^\alpha + bx^{\alpha+1} + cx^{\alpha+2} + \text{etc.} + kx^\kappa$$
$$Q \equiv mx^\mu + nx^{\mu+1} + qx^{\mu+2} + \text{etc.} + tx^\tau$$

where $a, k, m, t$ are not divisible by the modulus. We must have $\alpha$ not $< \mu$, and $\kappa$ not $< \tau$. Division can be carried out as in ordinary algebraic calculation, provided that an integer is always taken for the quotient; namely, the quotient will always have the form $\frac{r}{m}$, which must be determined for the modulus. Now, if a remainder is left after $\kappa + \mu - \alpha - \tau + 1$ terms have been found, it will have the form

$$Ax^{\kappa+\mu-\tau+1} + Bx^{\kappa+\mu-\tau+2} + .. + Cx^\kappa$$

and if not all the coefficients $A, B, C..$ are $\equiv 0$, then $P$ cannot be divided by $Q$.

It is also clear that division could have begun with the terms of highest degree, $kx^\kappa$, $tx^\tau$. The calculation will be easier if $Q$ is put in the form

$$mx^\mu(1 + qx + rxx + \text{etc.})$$

which, on putting $mv \equiv 1$, gives

$$\frac{P}{Q} \equiv \frac{vP : x^\mu}{1 + qx + \text{etc.}}$$

Division can then be completed by the usual methods.

**333.**

THEOREM. *If $x \equiv a$ is a root of the congruence $\xi \equiv 0$, then $\xi$ can be divided by $x - a$ for the modulus of the congruence.*

*Proof.* For if it could not be divided, we would have $\xi \equiv (x-a)\xi' + b$, where $b$ could not be divided by the modulus. Now, if $x$ is set $\equiv a$, then $\xi$ becomes $\equiv 0$ by hypothesis, so $(x-a)\xi' + b \equiv 0$. But then also $(x-a)\xi' \equiv 0$, and therefore $b$ must be $\equiv 0$.

**334.**

PROBLEM. *Given two functions, find their common divisor (of greatest degree) for a given modulus.*

*Solution.* Let the functions be $A, B$, and let the degree of $A$ be at least that of $B$. Divide $A$ by $B$. If this is possible without a remainder, $B$ will be the common divisor sought. If a remainder $C$ is left, it will have lower degree than $B$. Thus let

$$A \equiv aB + C, \quad B \equiv bC + D, \quad C \equiv cD + E, \quad \text{etc.}$$

where $A, B, C, D, a, b, c$, and so on, are functions, and the degrees of the functions $A, B, C, D$, and so on, form a decreasing sequence. If some division eventually succeeds, for example $D \equiv dE$, the last divisor will be the common divisor sought. If none succeeds, a remainder will eventually be reached which has no degree, that is, a *number*; in this case, however, the functions $A, B$ have no common divisor.

*Proof.* If the divisor $E$ divides the preceding function without a remainder, it is easily seen to divide all the preceding ones. Hence $E$ will be a common divisor of the functions $A, B$, proving the first assertion. But if there were a divisor of higher degree, say $E'$, then, because $C \equiv A - aB$, it would also divide $C$, and by the same argument $D$, and so on, and therefore $E$: a function of higher degree would divide one of lower degree. This is absurd, proving the second assertion. It is also clear from this that, if a common divisor of any degree exists, a remainder with no degree cannot be reached. Otherwise a function with no degree would be divided by a function of some degree, which is absurd.

**335.**

THEOREM. *If $A, B$ are coprime functions for the modulus $p$, and $A$ has degree $\alpha$, while $B$ has degree $\beta$, then functions $P, Q$ can be found whose degrees are respectively $< \beta$, $< \alpha$, such that*

$$PA + QB \equiv 1 \ (mod.\, p)$$

*Proof.* For in this case we have

$$A \equiv aB + C, \quad B \equiv bC + D, \quad \text{etc.} \quad K \equiv kL + M$$

where the degrees of the functions $A, B, C, D, \ldots K, L, M$ continually decrease, and $M$ has no degree. Now form the sequences

$$\begin{array}{cccccc} a, & a', & a'', & a''', & \ldots & a^{(\kappa)} \\ 1, & b, & b', & b'', & \ldots & b^{(\kappa-1)} \end{array}$$

so that

$$\begin{array}{llll} a' \equiv ba + 1 & a'' \equiv ca' + a & a''' \equiv da'' + a' & \text{etc.} \\ b' \equiv cb + 1 & b'' \equiv db' + b & b''' \equiv eb'' + b' & \text{etc.} \end{array}$$

We then have

$$A - aB \equiv +C, \quad bA - a'B \equiv -D, \quad b'A - a''B \equiv +E, \quad \text{etc.}$$

as is readily seen, and hence, finally,

$$b^{(\kappa-1)}A - a^{(\kappa)}B \equiv \pm M$$

Now let $\frac{1}{\pm M} \equiv \mu$. Then, on putting $P \equiv \mu b^{(\kappa-1)}$, $Q \equiv -\mu a^{(\kappa)}$, we obtain

$$PA + QB \equiv 1$$

Further, it is clear that

$$\begin{array}{c} \text{degree of } B + \text{degree of } a = \text{degree of } A. \\ \text{degree of } C + \text{degree of } b = .\text{degree of } B \\ \text{etc.} \\ \text{degree of } L + \text{degree of } k = \text{degree of } K \end{array}$$

Therefore

$$\text{degree of } L + \text{sum of the degrees of } a, b, \ldots k = \text{degree of } A$$

The degree of $a^{(\kappa)}$ is evident, and hence also

$$\text{degree of } Q = \text{sum of the degrees of } a, b, c, \ldots \text{ that is, } = \alpha - \text{degree of } L$$

Thus

$$\text{degree of } P = \beta - \text{degree of } L \qquad \text{Q. E. D.}$$

**336.**

It follows that, if $M$ is the common divisor of greatest degree of the functions $A, B$, we can always put

$$AP + BQ \equiv M$$

For brevity I omit examples of the preceding theorem, but readers will not neglect to gain skill in handling problems of this kind by working through examples. It is also worth pointing out that the preceding theorem holds for functions considered without a modulus, provided their coefficients are rational numbers. This will be evident from the method of proof itself. We cannot, however, dwell on this matter. The reader will observe similar things in what follows even without being prompted.

If $A$ has no common divisor of any degree with either $B$ or $C$, it will also have no common divisor with the product $BC$. For let

$$PA + QB \equiv 1, \quad \text{then} \quad PAC + QBC \equiv C$$

Now, if $A$ had with $BC$ a common divisor $M$, this would also divide $C$, contrary to the hypothesis. Thus, in general, if a function $A$ is coprime to $B, C, D$, and so on, it will also be coprime to the product of them all.

If $A, B, C, D$, and so on, have no divisor common to them all, we can obtain

$$PA + QB + RC + SD + \text{etc.} \equiv 1$$

Let the common divisor of greatest degree of $A$ and $B$ be $M$; that of $M$ and $C$ be $M'$; that of $M'$ and $D$ be $M''$, and so on. The last term of this sequence clearly has no degree, by hypothesis. We can therefore put

$$aA + bB \equiv M, \quad mM + cC \equiv M', \quad m'M' + dD \equiv M'', \quad \text{etc.}$$

and after substitution the truth of the theorem is apparent.

**337.**

THEOREM. *If $A, B, C$, and so on, are functions coprime to one another (any two of them having no common divisor) for the modulus $p$, and the function $M$ is divisible by each of them for the same modulus, it will also be divisible by the product of them all.*

*Proof.* For we can put $PA + QB \equiv 1$, so that

$$\frac{M}{A}Q + \frac{M}{B}P \equiv \frac{M}{AB}$$

Now, since $C$ is coprime to $AB$, the function $M$ will also be divisible by $ABC$, and by the same reasoning by $ABCD$, and so on.

**338.**

If the congruence $\xi \equiv 0$ has roots $x \equiv a$, $x \equiv b$, $x \equiv c$, and so on, then $\xi$ can be divided by the product of $(x-a)$, $(x-b)$, $(x-c)$, and so on. For, since $a, b, c.$, and so on, are assumed incongruent, the functions $x-a, x-b, x-c$, and so on, will be coprime to one another. Since $\xi$ is divisible by each of them, it will also be divisible by their product. Thus the number of roots cannot exceed the degree of the congruence: this is the proof of the theorem that we promised.

At the same time, however, this shows how the solution of congruences forms only one part of a much deeper investigation, namely the factorization of functions. The congruence $\xi \equiv 0$ clearly has no real roots if $\xi$ has no factors of degree one. Yet nothing prevents $\xi$ from being resolved into factors of degree two, three or more, from which roots that are, as it were, *imaginary* could be attributed to it. Indeed, had we wished to take a freedom similar to that taken by recent mathematicians and to introduce such imaginary quantities, we could have shortened all our following investigations incomparably. Nevertheless, we preferred to derive everything from principles[*)].

**339.**

*Functions* are called *prime* for a given modulus when they cannot be divided by any functions of lower degree for that modulus.

Thus all functions of degree one will be prime, while functions of degree two will either be prime or be composed of two of degree one. Hence $\xi$ will be a prime function of degree two if the congruence $\xi \equiv 0$ admits no real roots. For example, $xx + x + 1$ is prime for the modulus 5, because

$$xx + x + 1 \equiv (x-2)^2 - 3 \pmod 5$$

and 3 is a quadratic nonresidue of 5.

These prime functions demand our attention above all. Although those other than functions of degree one cannot serve to find real roots, a fuller study of them recommends itself both for their remarkable properties and for other excellent truths to be deduced from them.

[*)] Perhaps on another occasion we shall explain our opinion on this matter more fully.

**340.**

THEOREM. *Every function is either prime or composed of prime functions; in the latter case it can be composed of prime functions in only one way.*

*Proof.* For unless the proposed function $A$ is prime, it will be divisible by another function $B$ of lower degree. If $B$ is not a prime function, it will be divisible by another function $C$ of lower degree. Continuing in this way, a prime function must eventually be reached. Otherwise this sequence would be infinite, which is absurd, since the degrees continually decrease. Now, if the last prime function is $L$, it will divide all the preceding ones. Hence $A \equiv LA'$, and $A'$ has lower degree than $A$. Repeating this, $A' \equiv L'A''$, and so on, a prime function must clearly eventually be reached. Thus $A$ will be $\equiv$ a product of prime functions $L, L', L''$, and so on. . This proves the first assertion.

Now, if we also had $A \equiv MM'M''$, and so on, and the functions $L, L', L''$, and so on, were not all the same as $M, M', M''$, and so on, remove those common to both sequences. Let $\lambda, \lambda', \lambda''$ . .; $\mu, \mu', \mu'', \ldots$ remain. Then $\mu$ will be coprime to $\lambda, \lambda', \lambda''$, and so on, and hence also to the product $\lambda\lambda'\lambda''$, and so on. Yet we must have

$$\lambda\lambda'\lambda'' \ldots \equiv \mu\mu'\mu'' \ldots \text{ that is, } \frac{\lambda\lambda'\lambda'' \ldots}{\mu} \equiv \mu'\mu'' \ldots \quad \text{a contradiction}$$

**341.**

The first main task of these investigations will be to determine the number of prime functions of any given degree. For a given modulus, the number of all different (incongruent) functions of any given degree is finite. Some of these are composed of prime functions of lower degrees, while others are prime; hence the number of the latter is also finite. A rigorous development of this matter is rather delicate; we shall begin with simpler cases.

With the modulus set $= p$, the number of all different functions of degree $n$ of the form

$$x^n + Ax^{n-1} + Bx^{n-2} + Cx^{n-3} + \text{etc.}$$

will be $p^n$. For the number of coefficients $.A, B\,.\,C$, and so on, is $n$, and each can independently of the others be $\equiv 0, 1, 2, 3 \ldots (p-1) (\text{mod.}\, p)$. The theory of combinations therefore gives $p^n$ different combinations, which determine the collection of all different functions of this degree.

Thus there will be $p$ functions of degree one, namely $x, x+1, x+2$ up to $x+p-1$; there will be $pp$ functions of degree two, and so on.

**342.**

We have already pointed out that all functions of degree one must be regarded as prime. If, therefore, as is sufficient for our purpose, we restrict ourselves to functions whose highest term has coefficient 1, there will be $p$ functions of the first degree, or of one dimension.

All functions of degree two are either composed of two of degree one or are prime. The theory of combinations shows that $p$ different things, with repetitions allowed, can be combined in $\frac{p.p+1}{1.2}$ different ways. Thus there will be that many functions composed of two prime functions of degree one, and hence $pp - \frac{p.p+1}{1.2} = \frac{1}{2}(pp - p)$ prime functions of degree two.

Similarly, from all functions of degree three, whose number is $p^3$, we must exclude those composed of three prime functions of degree one, whose number is $\frac{p.p+1.p+2}{1.2.3}$, and also those composed of a prime function of degree one and another of degree two, whose number is $p \,.\, \frac{1}{2}(pp - p)$. Removing these leaves $\frac{1}{3}(p^3 - p)$; this, then, is the number of prime functions of degree three. Clearly we can always continue in this way.

**343.**

To carry out these operations more easily and at the same time prepare the way for developing the general law, we shall consider the matter generally. For brevity we denote by (1) the number of prime functions of degree one, by (2) the number of prime functions of degree two, and further by $(1^2)$ the number of functions composed of two prime functions of degree one, and so on, and so on. In general, $(1^\alpha 2^\beta 3^\gamma \ldots)$ denotes the number of all functions composed of prime functions, namely $\alpha$ of degree one, $\beta$ of degree two, $\gamma$ of degree three, and so on. Their degree will therefore be $\alpha + 2\beta + 3\gamma + \text{etc.}$ It follows from the preceding results and the theory of combinations that

$$(1^\alpha 2^\beta 3^\gamma 4^\delta \ldots) = (1^\alpha)(2^\beta)(3^\gamma)(4^\delta)\,..$$

$$(1^\alpha) = \frac{(1)\,.\,(1) + 1\,.\,(1) + 2\,.\,(1) + 3 \ldots (1) + \alpha - 1}{1\,.\,2\,.\,3\,.\,4 \ldots \alpha}$$

or, in general,

$$(a^\alpha) = \frac{(a)\,.\,(a) + 1\,.\,(a) + 2\,.\,(a) + 3 \ldots (a) + \alpha - 1}{1\,.\,2\,.\,3\,.\,4 \ldots \alpha}$$

Finally, if all the different ways of composing the number $n$ by addition from the numbers $1, 2, 3, \ldots$ are collected, and these ways are denoted by $\alpha \,.\, 1 + \beta \,.\, 2 + \gamma \,.\, 3 + \text{etc.}$, then the sum of all these expressions $(1^\alpha 2^\beta 3^\gamma \,..)$ is clearly equal to the number of all functions of degree $n$, that is, $= p^n$. Thus

$$\begin{aligned} p &= (1) \\ pp &= (1^2) + (2) \\ p^3 &= (1^3) + (1 \,.\, 2) + (3) \\ p^4 &= (1^4) + (1^2 \,.\, 2) + (1 \,.\, 3) + (2^2) + (4) \\ &\text{etc.} \end{aligned}$$

The expression $p^n$ clearly contains, besides the quantities $(1), (2), (3)$, and so on, also $(n)$, which shows how all these quantities are to be determined from the preceding ones. Thus we find

$$\begin{array}{lll} (1) = p & (4) = \frac{1}{4}(p^4 - pp) & (7) = \frac{1}{7}(p^7 - p) \\ (2) = \frac{1}{2}(pp - p) & (5) = \frac{1}{5}(p^5 - p) & (8) = \frac{1}{8}(p^8 - p^4) \\ (3) = \frac{1}{3}(p^3 - p) & (6) = \frac{1}{6}(p^6 - p^3 - pp + p) & \text{etc.} \end{array}$$

**344–346.**

From the beginning of this sequence we observe that the highest term of the expression $(n)$ is $\frac{1}{n}p^n$. If $n$ is prime, $-\frac{1}{n}p$ is added to it; if $n$ is composite, however, the law is less evident. But if we consider the matter more closely, we see that

$$\begin{array}{ll} p = (1) & p^5 = 5(5) + (1) \\ pp = 2(2) + (1) & p^6 = 6(6) + 3(3) + 2(2) + (1) \\ p^3 = 3(3) + (1) & p^7 = 7(7) + (1) \\ p^4 = 4(4) + 2(2) + (1) & p^8 = 8(8) + 4(4) + 2(2) + (1) \text{ etc.} \end{array}$$

Here the law of progression is apparent: if all the divisors of the number $n$ are $\alpha, \beta, \gamma, \delta$, and so on, then

$$p^n = \alpha(\alpha) + \beta(\beta) + \gamma(\gamma) + \delta(\delta) + \text{etc.}$$

We now set out to prove that this observation holds in general.

We showed that the sum of all expressions of the form $(1^\alpha)(2^\beta)(3^\gamma)\ldots$, where always $\alpha + 2\beta + 3\gamma + \ldots = n$, exhausts all functions of degree $n$ and is therefore $= p^n$. It follows that, — — —. If

$$\left(\frac{1}{1-x}\right)^{(1)}\left(\frac{1}{1-xx}\right)^{(2)}\left(\frac{1}{1-x^3}\right)^{(3)}\ldots$$

is expanded into the series $1 + Ax + Bx^2 \ldots = P$, then

$$A = p, \quad B = p^2, \quad C = p^3 \text{ etc.}$$

$$\frac{x\,\mathrm{d}P}{P\,\mathrm{d}x} = \frac{(1)x}{1-x} + \frac{2(2)x^3}{1-x^2} + \frac{3(3)x^3}{1-x^3}\ldots$$

— — — —

[From this, on substituting $\frac{px}{1-px}$ for $\frac{x\,\mathrm{d}P}{P\,\mathrm{d}x}$ and expanding the individual fractions into infinite series, the truth of the theorem follows of itself.]

**347.**

This theorem can also be expressed in another way. If all the divisors of the number $n$ are $n, 1, \delta, \delta', \delta'', \delta'''$, and so on, the theorem states that

$$p^n = n(n) + (1) + \delta(\delta) + \delta'(\delta') + \text{etc.}$$

Now the product of the $(n)$ prime functions of degree $n$ clearly has degree $n(n)$, and likewise for the others. Hence

*The product of all prime functions of degree one and of degrees $n, \delta, \delta'$, and so on, will have degree $p^n$.*

It is now easy to deduce the value of the expression $(n)$ itself from this theorem, but for brevity we omit the analysis, which is not difficult. Let $n = a^\alpha b^\beta c^\gamma$, and so on, where $a, b, c$, and so on, are distinct prime numbers. Then

$$n(n) = p^n - \Sigma p^{\frac{n}{a}} + \Sigma p^{\frac{n}{ab}} - \Sigma p^{\frac{n}{abc}} \text{ etc.}$$

Here $\Sigma p^{\frac{n}{abc..}}$ denotes the collection of all expressions similar to $p^{\frac{n}{abc..}}$ when the quantities $a, b, c..$ are permuted among themselves in every way. Thus for $n = 36$ we have $36(36) = p^{36} - p^{18} - p^{12} + p^6$.

Let us add one more observation. If $n$ has the form $a^\alpha$, and $a$ is prime, then $n(n) = p^n - p^{\frac{n}{a}}$. Since $(n)$ must be an integer, we therefore have, whatever $p$ may be,

$$p^n \equiv p^{\frac{n}{a}} \pmod{n}$$

Thus, if $p$ is coprime to $a$,

$$p^{n-\frac{n}{a}} \equiv 1 \pmod{n}$$

and for $\alpha = 1$,

$$p^{a-1} \equiv 1 \pmod{a}$$

It is remarkable that these theorems can be obtained in such different ways.

**348.**

PROBLEM. *Given the equation*

$$x^m + Ax^{m-1} + Bx^{m-2} + Cx^{m-3} + \text{etc.} + M = 0$$

*whose roots are $x = a$, $x = b$, $x = c$, and so on, find an equation whose roots are $x = a^\tau$, $x = b^\tau$, $x = c^\tau$, and so on.*

*First solution.* Find, by the known theorem, the sums of the roots of the given equation, of their squares, cubes, and so on, through the power with exponent $m\tau$. This also gives the sums of the roots of the required equation, and of their squares, and so on, namely $\Sigma a^\tau$, $\Sigma a^{2\tau}$, and so on. The coefficients can then be determined by the same theorem.

For practical calculation this solution is easier; but the following is better suited to our purpose, and also to showing that the coefficients of the required equation will be integers if the coefficients of the given equation are integers.

*Second solution.* Let $\theta$ be a primitive root of the equation $x^\tau = 1$, and form the product of

$$\begin{aligned}
&x^m + Ax^{m-1} + Bx^{m-2} + \text{etc.}\\
&x^m + A\theta x^{m-1} + B\theta\theta x^{m-2} + \text{etc.}\\
&x^m + A\theta\theta x^{m-1} + B\theta^4 x^{m-2} + \text{etc.}\\
&\text{etc.}\\
&x^m + A\theta^{\tau-1} x^{m-1} + B\theta^{2\tau-2} x^{m-2} + \text{etc.}
\end{aligned}$$

The roots of this product will therefore be

$$\begin{array}{llll}
a, & \theta a, & \theta\theta a & \text{etc.}\\
b, & \theta b, & \theta\theta b & \text{etc.}\\
c, & \theta c, & \theta\theta c & \text{etc.}
\end{array}$$

that is, the product will equal

$$(x^\tau - a^\tau)(x^\tau - b^\tau)(x^\tau - c^\tau)\ldots$$

and hence have the form

$$x^{\tau m} + A' x^{\tau(m-1)} + B' x^{\tau(m-2)} + \text{etc.}$$

Now, replacing $x^\tau$ by $x$, we have

$$x^m + A'x^{m-1} + B'x^{m-2} + \text{etc.} = (x - a^\tau)(x - b^\tau)(x - c^\tau)\dots$$

and hence

$$x^m + A'x^{m-1} + B'x^{m-2} + \text{etc.} = 0$$

is the required equation. That $A', B'$, and so on, are not only rational numbers but also integers follows easily from the theory of the equation $x^\tau = 1$.

Since we shall often use this operation below, by $(P, \rho^\tau)$ we shall denote the function which, on being set equal to zero, gives an equation whose roots are the powers with exponent $\tau$ of the roots of the equation $P = 0$.

If $P \equiv Q$ for any modulus, then also $(P, \rho^\tau) \equiv (Q, \rho^\tau)$ for the same modulus.

**349.**

THEOREM. *The coefficient of the term $x^n$ in $(P, \rho^\tau)$ is congruent, modulo $\tau$, to the coefficient of the term $x^{\tau n}$ in $P^\tau$, provided that $\tau$ is a prime number* (which, in this case, gives a third solution of the preceding problem).

*Proof.* It follows from Chapter Six that every coefficient of the product

$$(x^m + Ax^{m-1} + \text{etc.})(x^m + A\theta x^{m-1} + \text{etc.})\dots$$

has the following form after $\theta^\tau$ is replaced by unity:

$$E + (1 + \theta + \theta\theta + \text{etc.} + \theta^{\tau-1})F$$

Now, if $\theta$ is regarded as a primitive root of the equation $x^\tau = 1$, the whole product becomes $E$; but if $\theta = 1$, the whole product becomes $P^\tau = E + \tau F$. Thus the coefficient of the term $x^{n\tau}$ in $P^\tau$ is congruent, modulo $\tau$, to the coefficient of the term $x^{n\tau}$ in $E$, that is, to the coefficient of the term $x^n$ in $(P, \rho^\tau)$.

**350.**

THEOREM. *If $\tau$ is a prime number, then*

$$(P, \rho^\tau) \equiv P \ (mod.\, \tau)$$

*Proof.* Let the coefficient of the term $x^n$ in $(P, \rho^\tau)$ be $= N'$, and let the coefficient of the same term in $P$ be $= N$. Then, on putting

$$P = x^m + Ax^{m-1} + \text{etc.} + Nx^n + \text{etc.}$$

we have

$$P^\tau \equiv x^{m\tau} + A^\tau x^{(m-1)\tau} + \text{etc.} + N^\tau x^{n\tau} + \text{etc.} \ (\text{mod.}\, \tau)$$

and hence, by the preceding article, $N' \equiv N^\tau \ (\text{mod.}\, \tau)$. Thus, since $N^\tau \equiv N$, we obtain $N' \equiv N$. Q. E. D.

It is also clear from this that $(P, \rho^{\alpha}) \equiv (P, \rho^{\alpha\tau})$ and $(P, \rho^{\tau}) \equiv (P, \rho^{\tau\tau})$, whence, in general,

$$(P, \rho^{\alpha}) \equiv (P, \rho^{\alpha\tau^k}) \,(\mathrm{mod.}\, \tau)$$

**351.**

THEOREM. *There is a value of the number $\nu$ less than $p^m$ such that the function $x^{\nu} - 1$ can be divided by the given function $P$ of degree $m$, whose lowest term does not involve the indeterminate $x$, modulo $p$.*

*Proof.* Divide by $P$ the sequence of functions $1, x, xx \ldots$ through $x^{p^m-1}$ as soon as their degree exceeds $m$. Since none can be divided by $P$ without a remainder, all the remainders can be put in the form

$$Ax^{m-1} + Bx^{m-2} + \ldots + N$$

with all coefficients positive and $< p$. But since they can never all be $= 0$, there are clearly only $p^m - 1$ functions, one of which each remainder must equal. Since, through the power of $x$ whose exponent is $p^m - 1$, there are $p^m$ remainders, at least two must necessarily be the same. Let the same remainder result when $x^a$ and $x^{a+\nu}$ are divided by $P$, with $a + \nu < p^m$. Thus $x^{a+\nu} - x^a$ can be divided by $P$. Since, by hypothesis, $x$, and hence also $x^a$, is a function coprime to $P$, it follows that $x^{\nu} - 1$ can also be divided by $P$. Q. E. D.

*Corollary.* If $x^{\nu} - 1$ can be divided by $P$, then $x^{k\nu} - 1$ can also be divided by $P$, where $k$ denotes any integer.

**352.**

THEOREM. *Keeping the notation of the preceding article, if $P$ is a prime function and $x^{\nu}$ is the lowest power which, after unity is subtracted, can be divided by $P$, then $\nu$ is either $= p^m - 1$ or a divisor of this number, with the sole exception of the case $P = x$.*

*Proof.* Since $P$ is a prime function of degree $m$, there are $p^m - 1$ different functions of degree less than $m$ (excluding the function 0 from the total), all of which are coprime to $P$. Now, since $x^\nu$ is assumed to be the lowest power which leaves unity as remainder on division by $P$, it is clear that dividing all lower powers, from $1, x, ..$ through $x^{\nu-1}$, by $P$ gives $\nu$ different remainders, which we denote generally by $A$. If these exhaust all possible remainders, the theorem is proved; but if some are still absent from their number, let $B$ be any one of them. It is clear that the function $Bx^\nu$, divided by $P$, gives remainder $B$, and that, in general, $Bx^{\nu+k} \equiv Bx^k (\mathrm{mod.}\, P)$. But all the functions from $B$ through $Bx^{\nu-1}$ give remainders different from one another and from the remainders $A$. For if $Bx^\lambda \equiv Bx^{\lambda+\delta} (\mathrm{mod.}\, P)$, we would also have $1 \equiv x^\delta (\mathrm{mod.}\, P)$, with $\delta < \nu$, contrary to the hypothesis; while if $Bx^\lambda \equiv x^\mu (\mathrm{mod.}\, P)$, we would have $B \equiv x^{\mu+\nu-\lambda} (\mathrm{mod.}\, P)$, and hence $B$ would be one of the remainders $A$, contrary to the hypothesis. We thus clearly have $\nu$ further new remainders. We may proceed in the same way (exactly as above, §. .), and it will become apparent that the number $p^m - 1$ of all possible remainders is either $= \nu$, or $= 2\nu$, or $= 3\nu$, or, in general, a multiple of the number $\nu$. Q. E. D.

**353.**

It follows from the preceding theorem and the corollary to §. 351 that every prime function of degree $n$ divides the function $x^{p^n-1} - 1$ modulo $p$. Thus all functions of degree one, except the single one which is $\equiv x$, divide $x^{p-1} - 1$, which is FERMAT'S THEOREM; all prime functions of degree two, that is, of the form $xx + Ax + B$, divide the function $x^{pp-1} - 1$, and so on. Now let all divisors of the number $n$ be $n, \delta, \delta', \delta''$, and so on, . 1. It is clear that $p^n - 1$ is also divisible by $p^\delta - 1$, $p^{\delta'} - 1$, $p^{\delta''} - 1$, and so on, $p - 1$. Hence the function $x^{p^n-1} - 1$ can be divided by *all prime functions of degrees $n, \delta, \delta', \delta''$, and so on, down to the prime functions of degree one (excluding the function $x$)*, and therefore also, since all these functions are prime in themselves and hence coprime to one another, by *the product of them all.* But this same product has degree $p^n - 1$ (§. 347), because the one function $x$ is absent. Thus it is clear that *this product itself must be congruent to* $x^{p^n-1} - 1 (\mathrm{mod.}\, p)$.

**354.**

THEOREM. *If the function $x^\nu - 1$ is divisible by the function $P$, then*

$$(P, \rho^{k\nu+t}) \equiv (P, \rho^t)$$

*where $k, t$ denote any integers.*

*Proof.* Let

$$P = x^m + Ax^{m-1} + Bx^{m-2} + \text{etc.}$$

It is known that, if

$$\frac{mx^{m-1} + (m-1)Ax^{m-2} + \text{etc.}}{x^m + Ax^{m-1} + \text{etc.}}$$

is expanded in an infinite series of the form

$$m\frac{1}{x} + \alpha\frac{1}{xx} + \beta\frac{1}{x^3} + \gamma\frac{1}{x^4} + \text{etc.}$$

then $\alpha$ is the sum of the roots of the equation $P = 0$, $\beta$ the sum of their squares, and so on. From this it follows without difficulty that the sums of the powers with exponents $\nu + 1$, $\nu + 2$, and so on, are congruent to the sum of the roots, of their squares, and so on. Hence, unless the modulus is equal to or less than the degree of the function $P$, we have

$$(P, \rho^{\nu+1}) \equiv P, \quad (P, \rho^{\nu+2}) \equiv (P, \rho^2), \quad (P, \rho^{\nu+3}) \equiv (P, \rho^3) \text{ etc.}$$

We shall consider that case below.

**355.**

THEOREM. *If, in the sequence*

$$(P,\rho^0),\quad (P,\rho)\quad (P,\rho^2),\quad (P,\rho^3)\text{ etc.}$$

*the terms following the term in position $\nu$ are successively congruent to the first terms, then $x^\nu - 1$ can be divided by $P$, provided that $P$ contains no factor more than once.*

*Proof.* On putting $\frac{\mathrm{d}P}{\mathrm{d}x} = Q$, $Q$ will be a function coprime to $P$. Let

$$\frac{Q}{P} \equiv \frac{A}{x} + \frac{B}{xx} + \frac{C}{x^3} + \text{etc.}$$

Then, after the term $\frac{N}{x^\nu}$, there follows, by hypothesis,

$$\frac{A}{x^{\nu+1}} + \frac{B}{x^{\nu+2}} + \frac{C}{x^{\nu+3}} + \text{etc.}$$

Thus

$$\frac{Q}{P} \equiv \frac{Ax^{\nu-1} + Bx^{\nu-2} + \text{etc.}}{x^\nu - 1}$$

which shows that the function $x^\nu - 1$ can be divided by $P$. Q. E. D.

**356.**

THEOREM. *If $P$ is a prime function of $x$ of degree $m$, and $X$ is a function of $x, x^p, x^{pp}, x^{p^3} \,..\, x^{p^{m-1}}$ in which all these quantities occur in the same way, that is, which remains unchanged however they are permuted among themselves, then dividing the function $X$ by $P$ gives a remainder which is a number.*

*Proof.* Let the remainder be

$$Ax^{m-1} + Bx^{m-2} + \,..\, + N \equiv \xi$$

All coefficients $A, B, C \ldots$ up to, but excluding, $N$ will be $\equiv 0$. This is proved as follows. Since $X - \xi$ is divisible by $P$, $X^p - \xi^p$ can also be divided by $P$. But it is easy to see that $X^p$ is what $X$ becomes when $x$ is replaced by $x^p$, $x^p$ by $x^{pp}$, and so on, . and $x^{p^{m-1}}$ by $x^{p^m}$, or, equivalently, $x$. Thus it is clear that $X^p \equiv X \pmod{P}$; hence, since $X^p \equiv \xi^p$ and $X \equiv \xi (\mathrm{mod.}\, P)$, we also have $\xi^p \equiv \xi (\mathrm{mod.}\, P$, or

$$\xi^p - \xi \equiv 0 (\mathrm{mod.}\, P)$$

But $\xi^p - \xi$ is congruent, modulo $p$, to the product of $\xi, \xi+1, \xi+2, ..$ through $\xi + p - 1$. These factors are all coprime to $P$ unless $\xi$ is simply a number. Hence $\xi^p - \xi$ cannot be divisible by $P$ in any other way. Q. E. D.

Functions of this kind are the sum of all the quantities, the sum of their squares, cubes, and so on, and the sums of their products taken two at a time, three at a time, and so on. We shall determine what that number is in the following article:

**357.**

THEOREM. *Let the prime function of the preceding article be*

$$P \equiv x^m - Ax^{m-1} + Bx^{m-2} - Cx^{m-3} + \text{etc.}$$

*Then the remainder when the sum of the quantities* $x, x^p$, *and so on,* $x^{p^{m-1}}$ *is divided by* $P$ *is* $\equiv A$*; for the sum of their products taken two at a time it is* $\equiv B$*; for the sum of their products taken three at a time it is* $\equiv C$*, and so on.*

*Proof.* Let those functions be $X, Y, Z$, and so on, and let their remainders be the numbers $A', B', C'$, and so on, in their respective order. It is easy to see that $x, x^p, x^{pp}$, and so on, are roots of the equation

$$z^m - Xz^{m-1} + Yz^{m-2} - Zz^{m-3} + \text{etc.} = 0$$

Thus, on putting $z = x$, we have

$$x^m - Xx^{m-1} + Yx^{m-2} - Zx^{m-3} + \text{etc.} = 0$$

But the functions $X - A'$, $Y - B'$, $Z - C'$, and so on, are divisible by $P$, and hence so is the function

$$x^m - A'x^{m-1} + B'x^{m-2} - C'x^{m-3} + \text{etc.}$$

This cannot happen unless $A' \equiv A$, $B' \equiv B$, $C' \equiv C$, and so on. Q. E. D.

It is also known that any other function $X$ of $x, x^p$, and so on, [in which all these quantities occur in the same way,] can always be derived from these. Thus

$$x^2 + x^{2p} + x^{3pp} + \text{etc.} \equiv AA - 2B(\text{mod.}\, P) \text{ etc. etc.}$$

Example. Let $p = 5$ and $P \equiv x^2 + 2x + 3$. Then the function $x + x^5$, divided by $P$, gives $\equiv -2$, and $x^6 \equiv 3$, and so on, and so on.

**358. 359.**

THEOREM. *Let $P$ be a prime function and $x^\nu$ the lowest power of $x$ which, divided by $P$, gives remainder 1. Further, let $P \equiv (P, \rho^n)$. Then $n$ is congruent to some power of the number $p$ modulo $\nu$.*

*Proof.* We showed above that, if $P$

$$= x^m + Ax^{m-1} + Bx^{m-2} + \text{etc.}$$

then

$$z^m + Az^{m-1} + Bz^{m-2} + \text{etc.} - (z-x)(z-x^p)\ldots(z-x^{p^{m-1}})$$

is divisible by $P$. In the same way it would follow that

$$z^m + Az^{m-1} + Bz^{m-3} + \text{etc.} - (z-x^n)(z-x^{np})\ldots(z-x^{np^{m-1}})$$

is divisible by $P$. Since these factors are coprime to one another, the individual factors must necessarily be congruent to the corresponding individual factors modulo $P, p$. Thus $z - x^n$ must be $\equiv z - x^{p^\kappa}$, that is, $p^\kappa \equiv n (\text{mod.}\, \nu)$. Q. E. D.*)

*Finding the prime divisors of the function $x^\nu - 1$ for a prime modulus.*

**360.**

If $\nu$ is divisible by the modulus $p$ or by one of its powers, put $\nu = p^k\lambda$. Then

$$x^\nu - 1 \equiv (x^\lambda - 1)^{p^k} (\text{mod.}\, p)$$

This makes it clear that only the case where $\nu$ is not divisible by $p$ needs to be considered.

*) If $(P, \rho^a) \equiv (P, \rho^b) (\text{mod.}\, p)$, then $a \equiv p^\kappa b (\text{mod.}\, \nu)$.

*Proof.* Let $z^m + Az^{m-1} + Bz^{m-2} + \ldots = \Pi$. Then $(\Pi, \rho^a) \equiv (\Pi, \rho^b) (\text{mod.}\, P)$; but

$$(\Pi, \rho^a) \equiv (z-x^a)(z-x^{ap})(z-x^{app})\ldots(z-x^{ap^{m-1}}),$$
$$(\Pi, \rho^b) \equiv (z-x^b)(z-x^{bp})(z-x^{bpp})\ldots(z-x^{bpm-1})$$

which establishes the proposition.

The product of $\Pi, (\Pi, \rho^2), (\Pi, \rho^3)$, and so on, $(\Pi, \rho^\nu)$ is $\equiv (z^\nu - 1)^m (\text{mod.}\, P)$, for

$$(z-x)(z-x^2)(z-x^3)\ldots(z-x^\nu)$$
$$\equiv (z-x^p)(z-x^{2p})(z-x^{3p})\ldots(z-x^{\nu p}) \equiv \text{etc.} \equiv z^\nu - 1$$

In the sequence $P, (P, \rho^2), (P, \rho^3)$, and so on, $\ldots (P, \rho^\nu)$, all prime divisors of the function $x^\nu - 1$ occur, each $m$ times. It follows that the product of them all is $\equiv (x^\nu - 1)^m$.

If $p^m \equiv 1 (\mathrm{mod.}\, \nu)$, with $m$ as small as possible, then $x^{p^m-1} - 1$ is clearly divisible by $x^\nu - 1$. Thus $x^\nu - 1$ cannot have any divisors other than those of $x^{p^m-1} - 1$. But this expression has prime divisors of degree $m$ and others whose degrees divide the number $m$. Thus $x^\nu - 1$ also has divisors of these kinds. We shall show how many it has of each kind by an example, from which the general law can easily be deduced.

Let $\nu = 63$ and $p = 13$. Then $m = 6$. Thus $x^{63} - 1$ has prime factors of degrees six, three, two and one modulo 13. It is clear that the product of the factors of degree one is the common divisor of greatest degree of the functions $x^{63} - 1$ and $x^{12} - 1$, that is, $x^3 - 1$; hence there are three prime factors of degree one. The product of all prime factors of degrees two and one is the common divisor of the functions $x^{63} - 1$ and $x^{168} - 1$, that is, $x^{21} - 1$; hence there are $\frac{21-3}{2}$, or 9, factors of degree two. The product of the prime factors of degrees three and one is the common divisor of the functions $x^{63} - 1$ and $x^{2196} - 1$, that is, $x^9 - 1$; hence there are $\frac{9-3}{3}$, that is, 2, divisors of degree three. Finally, the remaining ones have degree six, and their number is therefore $= \frac{63-6-18-3}{6}$, that is, 6.

Careful consideration of this matter easily gives the following general rule:

*Let $\delta$ be a divisor of $m$, and let all divisors of the number $\delta$ less than $\delta$ be $\delta', \delta'', \delta'''$, and so on. Let the greatest common divisors of $\nu$ with $p^\delta - 1$, $p^{\delta'} - 1$, $p^{\delta''} - 1$, and so on, be $\mu, \mu', \mu''$, and so on, respectively. Put $\frac{\mu}{\mu'}; \frac{\mu}{\mu''}, \frac{\mu}{\mu'''}$, and so on, $= \lambda', \lambda'', \lambda'''$, and so on. Then $x^\nu - 1$ has $\frac{1}{\delta}$ times as many prime divisors of degree $\delta$ as there are numbers below $\mu$ divisible by none of the numbers $\lambda', \lambda'', \lambda'''$, and so on.*

**361.**

THEOREM. *If a function $X$ of the indeterminate $x$ is divisible by another function $\xi$, and $X$, when $x$ is replaced by $x^k$, becomes $X'$, then $X'$ is divisible by $(\xi, \rho^{\frac{1}{k}})$.*

*Proof.* Let $X \equiv \xi\upsilon$, and let $\xi, \upsilon$ become $\xi', \upsilon'$ when $x$ is replaced by $x^k$. It is clear that $X' \equiv \xi'\upsilon'$. But $\xi'$ is divisible by $(\xi, \rho^{\frac{1}{k}})$. Hence so is $X'$. Q. E. D.

**362.**

With these principles established, we can easily determine the prime divisors of the function $x^\nu - 1$. We assume that all those divisors which also divide some function $x^{\nu'} - 1$, where $\nu' < \nu$, have already been found, and propose to investigate the remaining ones. They can all be included in the expression $(P, \rho^k)$, if $P$ is one of them and $k$ takes all values less than $\nu$ and coprime to it.

In Chapter VI we showed how the primitive roots of the equation $x^\nu = 1$ can be divided into classes so that, when all are expressed as powers of one of them, the same division into classes results whichever primitive root is taken as the base; we called such collections of roots *periods*. It is clear that the functions $x, x^\alpha, x^\beta, x^\gamma$, and so on, where $\alpha, \beta, \gamma$, and so on, denote all numbers coprime to $\nu$, can be divided into periods in the same way, and each larger period again into smaller ones, until periods of the form $x^k, x^{kp}, x^{kpp} \ldots x^{kp^{m-1}}$ are finally reached. Once this is done, the following is clear:

1. Since every period is composed of smallest periods of the form $x^k + x^{kp} +$ etc., division by any prime function of degree $m$ gives a remainder which is a number.

2. Since all terms of a period can always be put in the form $x^{\kappa . a^\alpha b^\beta c^\gamma \ldots}$, where $\kappa.\ a, b, c \ldots$ are fixed numbers, while all values may be substituted for $\alpha, \beta, \gamma \ldots$, it is clear that the period becomes itself when $x$ is replaced by $x^k$ and $k$ has the form $a^\alpha b^\beta c^\gamma \ldots (\mathrm{mod.}\, \nu)$. From this it is easy to see that all the functions $P, (P\ \rho^k)$, and so on, with $k$ denoting a number of this kind, give the same remainder when the period is divided by them.

3. Thus, after such a remainder has been subtracted, the period can be divided by the product of all the functions $(P, \rho^k)$.

**363.**

The main task is to determine these remainders. First, find the remainder that the largest period gives on division by the product of all the appropriate prime functions. If this product is

$$\equiv x^\lambda - Ax^{\lambda-1} + \text{etc.}$$

then this remainder is $\equiv A$. The form of this product is readily found, and Chapter VI shows that $A = 0$ if $\nu$ is divisible by a square; otherwise $A$ is either $= +1$ or $= -1$, according as the number of prime factors of $\nu$ is even or odd.

Now split this largest period into smaller periods, and represent the terms of any one period by $x^{kp^\pi u}$, so that $k$ is a number fixed within each period but varying between periods, while $\pi$ and $u$ vary within each period and can take in the other periods the same values that they take in any one period. For the moment, take some prime function $P$ as the base, and let the

periods $\Sigma x^{p^\pi u}$, $\Sigma x^{k' p^\pi u}$, etc., on division by it, give the remainders $A, A'$, etc., respectively. Then $\Sigma x^{p^\pi u} - A$ is divisible by the product of all the functions $(P, \rho^u)$; $\Sigma x^{k' p^\pi u} - A'$ by the product of all the functions $(P, \rho^{k' u})$; etc., etc. It is readily seen that the quantities $A, A'$, etc., are roots of a known congruence. Namely, let the periods of the roots of the equation $x^\nu = 1$ corresponding to the preceding periods be roots of the equation $Q = 0$; then $A, A'$, etc., will be roots of the congruence $Q \equiv 0$. For we have

$$A + A' + \text{etc.} \equiv \text{the sum of the periods},$$
$$AA + A'A' + \text{etc.} \equiv \text{the sum of the squares of the periods}$$

etc., etc. The calculation is entirely similar to the one we set out in Chapter VI, if in place of $\rho$ we substitute $x$, since here too we can put unity for $x^\nu$, as we did there for $\rho^\nu$.

Once the roots $A, A'$, etc., have been found, choose one as the remainder of the period $\Sigma x^{p^\pi u}$, and use it to order the remainders of the others in the same way as in Chapter VI. This choice is left free here too, since the function $P$ is still entirely undetermined. The subsequent calculation is wholly analogous to the one we treated in Chapter VI; setting out every detail would take us too far. Finally, once we have reached $\Sigma x^{p^\pi}$, the main task is complete. For, on putting

$$P \equiv x^m + ax^{m-1} + bx^{m-2} + \text{etc.}$$

we have $-a \equiv \Sigma x^{p^\pi}$. The second coefficient of the other functions $(P, \rho^k)$ is obtained in the same way, and from this the remaining coefficients of $P$ can be determined. It may often happen that we reach identical congruences from which nothing seems to follow. How this difficulty can be met will be shown below.

**364.**

An example will make all this much clearer. The proposed task is to resolve the function $x^{15} - 1$ into factors modulo 17. We have $m = 4$, and since the product of all the elementary functions is

$$\equiv \frac{x^{15} - 1 \,.\, x - 1}{x^3 - 1 \,.\, x^5 - 1} = x^8 - x^7 + x^5 - x^4 + x^3 - x + 1$$

there will therefore be only two prime factors of degree four, $P$ and $P'$. Now distribute $x, xx, x^4, x^7, x^8, x^{11}, x^{13}, x^{14}$ into these two periods:

$$\Sigma x^{17^\alpha} \equiv x + xx + x^4 + x^8,$$
$$\Sigma x^{7.17^\alpha} \equiv x^7 + x^{11} + x^{13} + x^{14}$$

Modulo one of the functions $P, P'$, let

$$\Sigma x^{17^\alpha} \equiv A, \quad \Sigma x^{7.17^\alpha} \equiv A'$$

Then

$$\begin{aligned} A + A' &\equiv 1 \\ AA &\equiv \Sigma x^{2.17^\alpha} + \Sigma x^{3.17^\alpha} + \Sigma x^{5.17^\alpha} + \Sigma x^{9.17^\alpha} \\ A'A' &\equiv \Sigma x^{14.17^\alpha} + \Sigma x^{6.17^\alpha} + \Sigma x^{5.17^\alpha} + \Sigma x^{3.17^\alpha} \end{aligned}$$

and hence

$$\begin{aligned} AA + A'A' &\equiv \Sigma x^{17^\alpha} + \Sigma x^{7.17^\alpha} + 4\Sigma x^{3.17^\alpha} + 2\Sigma x^{5.17^\alpha} \\ &\equiv 1 - 4 - 4 \equiv -7 \end{aligned}$$

Thus $A$ and $A'$ will be the roots of the congruence

$$xx - x + 4 \equiv 0 \pmod{17}$$

which are $6, 12$. It follows that $P$ divides

$$x^8 + x^4 + xx + x - 6$$

and is

$$\equiv x^4 - 6x^3 - 2xx - 12x + 1$$

Further, $P'$ will be $\equiv (P, \rho^7)$, and will be

$$\equiv x^4 - 12x^3 - 2xx - 6x + 1$$

**365.**

It is enough for us here to have shown the possibility of these solutions. For brevity, we pass over many devices that can ease these operations. But we cannot omit some very important consequences.

What precedes proves that all the auxiliary equations for solving the equation $x^\nu = 1$, if converted into congruences, have possible roots when the period

$$x + x^p + x^{pp} + \ldots + x^{p^{m-1}}$$

has not yet been split. Let us stay with the case where $\nu$ is prime; then $m$ is a divisor of $\nu - 1$. Here, therefore, the auxiliary congruences have real roots if the number of periods found through them is a divisor of $\frac{\nu-1}{m}$. Thus, if $\frac{\nu-1}{m}$ is even, that is, if $m$ is a divisor of $\frac{\nu-1}{2}$, or equivalently if $p^{\frac{\nu-1}{2}} \equiv 1 \pmod{\nu}$, or equivalently if $p$ is a quadratic residue of the prime $\nu$, the quadratic equation that divides the roots into two periods will have real roots modulo $p$. But in Chapter VI we showed that, on putting $\nu = 4n \pm 1$, this equation is always $xx + x \mp n = 0$. Hence we have the notable

THEOREM. *If the prime $p$ is a quadratic residue of the prime $4n \pm 1$, the congruence*

$$xx + x \mp n \equiv 0 (\text{mod. } p)$$

*will have real roots, and therefore so will the congruence*

$$4xx + 4x \mp 4n \equiv 0 \quad \text{or} \quad (2x+1)^2 \mp \nu \equiv 0$$

*that is, $\pm\nu$ will be a quadratic residue of $p$.*

**366.**

This, then, is the third complete proof of the fundamental theorem of Chapter IV, all the more worthy of attention because the principles from which it is drawn are entirely different from those we used for the earlier proofs. .But from this same source, by the opposite route, let us derive a fourth. Namely, let $\nu$ be a prime of the form $4n \pm 1$, let $p$ be any other prime, and let $\pm\nu$ be a quadratic residue of the prime $p$; we shall prove that $p$ is a quadratic residue of $\nu$.

Let $p^m$ be the least power of $p$ that is $\equiv 1 (\text{mod. } \nu)$. The elementary divisors of the function $\frac{x^\nu - 1}{x-1}$ modulo $p$ will have degree $m$, so their total number will be $= \frac{\nu-1}{m}$. Now, since $\pm\nu Rp$, the congruence

$$xx + x \mp n \equiv 0 (\text{mod. } p)$$

will be solvable; let its roots be $A, A'$. Distribute the functions $x, xx, .. x^{\nu-1}$ into two classes, denoted by $\xi, \xi'$. We have

$$\xi + \xi' \equiv A + A' + (1 + x + xx + .. + x^{\nu-1})$$
$$\xi\xi' \equiv AA' + \lambda(1 + x + xx + .. + x^{\nu-1})$$

so that

$$(z - \xi)(z - \xi') - (z - A)(z - A')$$

will be divisible by every elementary divisor of the function $\frac{x^\nu - 1}{x-1}$. Each of these elementary divisors will therefore divide either $\xi - A$ and $\xi' - A'$, or $\xi - A'$ and $\xi' - A$. It follows (since $A$ is not $\equiv A'$) that, if in place of $x$ we substitute $x^p$, then $\xi$ and $\xi'$ do not change. For if $\xi$ passed into $\xi'$, and conversely, then $\xi - A$ and $\xi - A'$ would be divisible by the same prime function. Q. E. A. Finally, it follows that $\frac{\nu-1}{2}$ is divisible by $m$, or equivalently that $p^{\frac{\nu-1}{2}} - 1$ is divisible by $\nu$. Thus $p$ is a quadratic residue of $\nu$. Q. E. D.

It is easy to derive all cases of the fundamental theorem from the two theorems.

**367.**

Although we have restricted ourselves here to the case where $\nu$ is prime, analogous theorems can also be determined without much difficulty when $\nu$ is composite; brevity does not now allow us to set this out more fully.

It is evident that similar observations can also be made about a greater number of periods. Thus, if $\frac{\nu-1}{m}$ is divisible by 3, that is, if $p$ is a cubic residue of the prime $\nu$, the equation that distributes the roots of the equation $x^\nu = 1$ into three periods, and whose determination a priori we taught in Chapter VI, will be solvable modulo $p$, and conversely. For example, the congruence $x^3 + xx - 2x - 1 \equiv 0$ can be solved modulo any prime of the form $7n \pm 1$, but cannot if it has another form.

It would not be difficult for us to enrich this chapter with many other observations, if the limits within which we must keep did not prevent it. For those who wish to go further, these principles can at least point the way.

**368.**

We say that a congruence $S \equiv 0$ has *equal* roots, or more generally equal divisors, if it is divisible by a power of some function.

Whether a given congruence has equal divisors is decided in the same way as in the theory of equations. Put

$$X \equiv \xi^m P$$

Clearly,

$$\frac{\mathrm{d}X}{\mathrm{d}x} \equiv \xi^{m-1}\left(mP\frac{\mathrm{d}\xi}{\mathrm{d}x} + \xi\frac{\mathrm{d}P}{\mathrm{d}x}\right)$$

so $\frac{\mathrm{d}X}{\mathrm{d}x}$ is divisible by $\xi^{m-1}$. More generally, let

$$X \equiv A^a B^b C^c \text{ etc.}$$

where $A, B, C$, etc., denote distinct prime functions. Then

$$\frac{\mathrm{d}X}{\mathrm{d}x} \equiv X\left(\frac{a\,\mathrm{d}A}{A\,\mathrm{d}x} + \frac{b\,\mathrm{d}B}{B\,\mathrm{d}x} + \frac{c\,\mathrm{d}C}{C\,\mathrm{d}x} + \text{etc.}\right)$$

It follows that, unless one of the numbers $a, b, c$, etc., is divisible by the modulus, $\frac{\mathrm{d}X}{\mathrm{d}x}$ is divisible by $A^{a-1}B^{b-1}C^{c-1}$, etc., but not by $A^a, B^b, C^c$, etc. Hence the following

THEOREM. *If the common divisor of greatest degree of the functions $X$ and $\frac{\mathrm{d}X}{\mathrm{d}x}$ is $\xi$, all the prime factors of $\xi$ will also occur in $X$, each with a multiplicity $+1$ greater than in $\xi$. Thus, if $X$ and $\frac{\mathrm{d}X}{\mathrm{d}x}$ are coprime functions, $X$ will have no equal factors.*

**369.**

Example I. We ask whether the function

$$x^5 + 3x^4 - 6x^3 + 3x - 4 \ldots (X)$$

has equal divisors modulo 17. We have

$$\frac{\mathrm{d}X}{\mathrm{d}x} \equiv 5x^4 - 5x^3 - xx + 3$$

From this we find that the functions $X$ and $\frac{\mathrm{d}X}{\mathrm{d}x}$ are coprime, so $X$ has no equal divisors.

Example II. Let

$$X \equiv x^5 + 6x^4 - 3x^3 - 4xx + 2x - 3 (\mathrm{mod.}\, 13)$$

Then

$$\frac{\mathrm{d}X}{\mathrm{d}x} \equiv 5x^4 - 2x^3 + 4xx + 5x + 2$$

The greatest common divisor of the functions $X$, $\frac{\mathrm{d}X}{\mathrm{d}x}$ is $\equiv 5xx + 7x + 7$, or, on multiplication by 8: $xx + 4x + 4$. Since this divisor is $\equiv (x+2)^2$, the function $X$ is divisible by $(x+2)^3$, and the quotient (which is $xx + 11$) contains no further double divisor.

**370. 371.**

If, by the preceding sections, the function $X$ has been expressed as $A^a B^b C^c$, etc., with $A, B, C$, etc., coprime and the numbers $a, b, c$, etc., unequal, the resolution can be carried further. Let $X$, then, be a function containing no further equal divisors. We saw above that $x^p - x$ is the product of all prime functions of degree one. Let $\xi$ be the common divisor of greatest degree of the functions $X$ and $x^p - x$; then $\xi$ is the product of all divisors of $X$ of degree one, and $\frac{X}{\xi}$ has no further divisors of this kind. If, however, the functions $X$ and $x^p - x$ are found to be coprime, $X$ will have no divisor of degree one, and hence the congruence $X \equiv 0$ will have no real roots. Further, since $x^{pp} - x$ is the product of all prime functions of degrees two and one, the common divisor of greatest degree of the functions $x^{pp} - x$ and $\frac{X}{\xi}$, namely $\xi'$, contains all divisors of $X$ of degree two. Continuing in this way, we see that $X$ is thus resolved into factors $\xi, \xi', \xi''$, etc., which contain all divisors of degrees one, two, three, etc., respectively.

**372.**

But if a product of several prime functions of the same degree is given, the individual functions must be found by trial. This problem is closely analogous to that of finding the factors of composite numbers. Here, however, it is determined a priori whether the given function can still be split into factors.

Since here too the number of all possible factors is finite, we can use an aid similar to the one above. But we do not wish to dwell on this matter, for an experienced calculator who has thoroughly grasped the principles will readily find particular devices when needed.

We proceed to another topic, namely the consideration of congruences when the modulus is not a prime number, as we have always assumed up to now. Here the case where the modulus is a power of a prime number especially deserves attention, both in itself and because it is needed to remove certain doubts (§.§. . .).

**373.**

PROBLEM. *If the function $X$ has been resolved modulo $p$ into coprime factors $\xi, \xi', \xi''$, etc., resolve $X$ modulo $pp$ into similar factors $\Xi, \Xi', \Xi''$, etc., so that*

$$\xi \equiv \Xi, \quad \xi' \equiv \Xi', \quad \xi'' \equiv \Xi'', \text{ etc. } (\mathrm{mod.}\, p)$$

*Solution.* Let $X \equiv \xi\psi (\mathrm{mod.}\, p)$, or equivalently $X = \xi\psi + p\Sigma$. Put

$$\Xi = \xi + p\varphi, \quad \Psi = \psi + p\omega$$

Then

$$\Xi\Psi = X - p\Sigma + (\varphi\psi + \xi\omega)p + pp\varphi\omega$$

Thus, if $\Xi\Psi$ is to be $\equiv X (\mathrm{mod.}\, pp)$, $\varphi\psi + \xi\omega - \Sigma$ must necessarily be divisible by $p$. But since $\psi$ and $\xi$ are coprime functions modulo $p$, $\varphi$ and $\omega$ can be determined so that this condition is fulfilled (§. 336), and moreover so that the degrees of $\varphi$ and $\omega$ are respectively one less than the degrees of the functions $\xi, \psi$. Hence $X \equiv \Xi\Psi (\mathrm{mod.}\, pp)$. It is clear that $\Psi$ can again be split in the same way into factors $\Xi'\Omega$, with the factor $\Xi'$ being $\equiv \xi' (\mathrm{mod.}\, p)$, and so on, until finally

$$X \equiv \Xi\, \Xi'\, \Xi'' \text{ etc. } (\mathrm{mod.}\, pp). \quad \text{Q. E. Fac.}$$

**374.**

From this it is easy to prove that the function $X$ can also be resolved into factors modulo $p^3, p^4$, etc. More generally, let

$$X \equiv PQ(\mathrm{mod.}\, p^m) \quad \text{or} \quad X = PQ + p^m R$$

and let the function $P$ be coprime to $Q$ modulo $p$. On putting

$$P' = P + Ap^m, \quad Q' = Q + Bp^m$$

we have

$$P'Q' = X - p^m R + (AQ + BP)p^m + ABp^{2m}$$

Hence, for any modulus $p^\nu$ (with $\nu$ being $> m$ and $< 2m + 1$), we have

$$P'Q' \equiv X, \quad \text{if} \quad R \equiv AQ + BP(\mathrm{mod.}\, p^{\nu - m})$$

From this we see that, if the function $X$ has no equal divisors modulo $p$, it can be split into factors modulo $p^k$ in the same way as modulo $p$. But if $X$ has equal divisors, the matter becomes much more complicated and cannot even be fully dealt with by the preceding principles. Since we cannot communicate everything belonging to this subject, we shall consider only one case, which occurs very often and whose resolution is needed to settle certain doubts in what precedes. This is the case where only equal factors of degree one are considered. It can also be used directly to find the roots of congruences. We shall treat the matter in general on another occasion.

**375.**

Let, then, $X \equiv X'(x - a)^m(\mathrm{mod.}\, p)$, and let the function $X'$ be coprime to $x - a$: we seek all divisors of degree one congruent to $x - a$ modulo $p$ that divide $X$ modulo $pp, p^3$, etc. (We assume that the function $X$ is not divisible by $x - a$ without a modulus; otherwise $x - a$ would divide the function $X$ modulo every modulus.) If we substitute $z + a$ for $x$, we obtain

$$Z \equiv Z'z^m(\mathrm{mod.}\, p) \quad \text{or} \quad Z = Z'z^m + pA$$

Now, if $Z$ is divisible modulo $pp$ by some divisor of the form $z + \alpha p$, then $A$ must be of the form $zZ'' + pB$. Unless this is so, the investigation is already finished. Let us therefore put

$$Z \equiv Z'z^m + pZ''z(\mathrm{mod.}\, pp)$$
$$\text{or} \quad Z = Z'z^m + pZ''z + ppB$$

It is clear that $Z$ is divisible by $z$ and by every other divisor congruent to it modulo $p$.

To fix our attention, we put $m = 4$; it is easy to see that every other case can be treated in a similar way. Now, if $Z$ is divisible modulo $p^3$ by some divisor of the form $z + \alpha p$, we have

$$0 \equiv -\alpha pp Z'' + ppB(\text{modd. } z + \alpha p, p^3)$$
$$\text{or} \quad \alpha Z'' \equiv B(\text{modd. } z, p)$$

There can now be three cases

1) If $Z'' \equiv 0(\text{modd. } z, p)$ and $B$ is not $\equiv 0$, then clearly no value of $\alpha$ satisfies the congruence, and hence $Z$ has, modulo $p^3$, no divisor of the form $z + \alpha p$. The investigation is therefore finished

2) If neither $Z''$ nor $B$ is $\equiv 0(\text{modd. } z, p)$, then $\alpha$ has a unique value, namely

$$\alpha \equiv \frac{B}{Z''}(\text{modd. } z, p).$$

Thus there is a unique divisor $\equiv z + \alpha p(\text{mod. } pp)$ of $Z$ modulo $p^3$, and we have

$$Z \equiv V(z + \alpha p) + p^3 W$$

Now let the divisor of $Z(\text{mod. } p^4)$ be $z + \alpha p + \beta pp$; then

$$0 \equiv$$

# REMARKS ON THE ANALYSIS RESIDUORUM.

The two preceding treatises are taken from an extensive manuscript entitled Analysis Residuorum, which probably dates from 1797 or 1798; a complete revision of it later gave rise to the Disquisitiones Arithmeticae. The full title of Chapter Six reads:

Solution of the congruence $x^m - 1 \equiv 0$ and the equation $x^m - 1 = 0$; with elucidations on the theory of regular polygons.

The essential content of its second part (§§. 253–278) passed into the seventh section of the Disqq. Arithm.

Chapter Seven is also partly preserved: Various Applications of Certain Preceding Investigations (§§. 279–302). It is divided into the following subsections:

On Transformations of Common Fractions (§§. 279–281).

On the Conversion of Common Fractions into Decimals (§§. 282–292).

On the Solution of the Indeterminate Equation $xx = a + by$ (§§. 293–297).

On the Solution of the Indeterminate Equation $axx + byy = c$ (§§. 298–301).

On Finding the Divisors of Numbers (§. 302; the following sheets are missing).

All this was incorporated almost word for word into the sixth section of the Disqq. Arithm.

The two sections given here treat the subjects which, as appears from the preface and Articles 11, 44, 61, 62, 65, 84 of the Disqq. Arithm., were intended to form the eighth section of that work. It should, however, be noted that this plan was later changed again: among the manuscripts is a fragment headed Section Eight: A Fuller Consideration of Certain Investigations Concerning the Division of the Circle. This fragment begins with Art. 367 and was therefore intended to continue the Disqq. Arithm.; but the content of the few articles still preserved later passed into the treatise Summatio quarundam serierum singularium, and for this reason the fragment is excluded from the present edition.

In the preceding printing of the two parts of the Analysis Residuorum, the original text has essentially been retained faithfully, although its form cannot be called ready for printing; the following remarks indicate the most important alterations and add some elucidations.

§. 237. Compare Disqq. Arithm. artt. 61, 62.

§. 239. Compare Disqq. Arithm. artt. 53, 54, 65.

§. 241. If $n = 2^\nu$ and $\nu \geqq 3$, there is indeed no number $\rho$ of the stated kind, but this does not substantially change the investigation as a whole.

§. 251. The difficulty noted here was presumably to be removed by introducing higher powers of $p$ as moduli. Compare §§. 363, 372, 373.

§. 332. The assumption that the modulus is a prime number is retained through §. 372 inclusive.

§. 338. The incomplete citation may refer to Disqq. Arithm. art. 44.

§§. 344–346. Of the two proofs present in the manuscript, the first, introduced by the words "we now set about proving" and based on a closer examination of the expressions $(1^\alpha 2^\beta 3^\gamma \ldots)$, has here been omitted entirely in accordance with the author's own instruction ("The whole preceding proof, together with another proof of the preceding theorem which we intended to add, is to be suppressed, because we have discovered another that is infinitely simpler. It rests on this foundation"); moreover, in the printing above the second proof has been shortened by considering the expansion of $\frac{x\,\mathrm{d}P}{P\,\mathrm{d}x}$ instead of that of $\frac{x\,\mathrm{d}P}{\mathrm{d}x}$, thus also avoiding a reference in the original to the suppressed first proof.

§. 348. The expression radix prima is to be understood here in the same sense as radix propria in the treatise Summatio quarundam serierum singularum, art. 11. — The assertion that the coefficients $A'.B'\ldots$ of the expanded product are rational integers refers to Chapter Six, where, however, the theory of the equation $x^\tau - 1 = 0$ is treated only for the case where $\tau$ is prime; the form of the proof in §. 349 first leads to the following supplement. If the expanded product is put into the form (valid for all roots of the equation $\theta^\tau = 1$)

$$S = E + F\theta + \ldots + N\theta^{\tau-1}$$

then the coefficients $E, F \ldots N$ are polynomials in $x$ with rational integer coefficients. Further, since the product remains unchanged when $\theta$ is replaced by $\theta^k$, where $k$ is any number coprime to $\tau$, the same holds for the expression $S$. It follows without difficulty that all powers occurring in $S$ of $\theta$ whose exponents $s$ have one and the same greatest common divisor with $\tau$ must also have identical coefficients. Finally, since every sum of such powers $\theta^s$ is always an integer, it is evident that the expression $S$, and hence also the product in question, is a polynomial in $x$ with integer coefficients, as was to be shown. This consideration also establishes the correctness of the remark at the end of the paragraph. Other reasons nevertheless suggest that the author already knew at that time the general theorem on the transformation of symmetric functions (Demonstratio nova altera theorematis omnem functionem etc. art. 4), from which the propositions above follow immediately.

§. 352. The notation $R \equiv S(\mathrm{mod.}\, P)$, or also $R \equiv S(\mathrm{mod.}\, P, p)$, means here and in what follows that the difference $R - S$ has, modulo $p$, the divisor $P$. — The incomplete citation may refer to Disqq. Arithm. art. 49.

§. 354. Multiplication by $x^\nu - 1$ shows that the sums of equal powers of the roots of the two equations $(P, \rho^{k\nu+t}) = 0$, $(P, \rho^t) = 0$ are congruent to one another (mod. $p$), and from this follows the congruence $(P, \rho^{k\nu+t}) \equiv (P, \rho^t)(\mathrm{mod.}\, p)$ as soon as $m < p$ (compare §. 244). But if $m \geqq p$, the coefficient of the power $x^{m-p}$ in an equation can no longer be determined modulo $p$ from the given power sums of its roots, because it carries the factor $p$ in the formulas of NEWTON used for this purpose. Indeed, from the congruence of each pair of equal power sums of the roots of the equations $A = 0$, $B = 0$, one may in general infer only that $A \equiv \mathfrak{A}^p\mathfrak{C}$, $B = \mathfrak{B}^p\mathfrak{C}(\mathrm{mod.}\, p)$, where $\mathfrak{C}$ denotes the greatest common divisor of the two functions $A, B$ modulo the prime number $p$, while $\mathfrak{A}$ and $\mathfrak{B}$ are entirely undetermined functions. Presumably the author derived the general validity of the proposition from the theory of the transformation of symmetric functions, and specifically from the following proposition: if, with respect to any modulus $p$, the difference $R(x) - S(x)$ is divisible by the function $P(x)$, and $a, b, c \ldots$ are the roots of the equation $P(x) = 0$, then the functions

$$(x - R(a))(x - R(b))(x - R(c)) \ldots \quad \text{and}$$
$$(x - S(a))(x - S(b))(x - S(c)) \ldots$$

are congruent to one another modulo $p$.

§. 355. It is shown in §. 368 that $P$ and $\frac{\mathrm{d}P}{\mathrm{d}x}$ have no common divisor if $P$ contains no factor more than once.

§§. 358, 359. The note placed below the text is taken from a separate sheet, which was probably intended to replace §. 359, already crossed out in the manuscript.

§. 360. An inaccuracy in the manuscript has been corrected in the statement of the theorem.

§. 361. Here the exponent $\frac{1}{k}$ in the notation $(\xi, \rho^{\frac{1}{k}})$ means any positive integer $k'$ such that $kk' \equiv 1(\mathrm{mod.}\, \nu)$, where $\nu$ is the least positive integer for which $x^\nu - 1$ is divisible by $\xi$ modulo $p$; it must be assumed here that $\xi$ is not divisible by $x$ modulo $p$, and also that $k$ is coprime to $\nu$. The correctness of the assertion that $\xi'$ is divisible by $(\xi, \rho^{\frac{1}{k}})$ (mod. $p$) follows from §. 354.

§. 363. The concluding remark presumably concerns the introduction of moduli that are powers of the prime number $p$; compare §§. 251, 372, 373.

§. 367. The roots of the equation $x^3 + xx - 2x - 1 = 0$ are the periods of two terms into which the roots of the equation $\frac{x^7-1}{x-1} = 0$ divide. The same example also occurs on a separate sheet, where the principal result of §§. 362, 363 is stated under the title "the golden theorem".

§. 371. This paragraph was intended to contain an example, but it was not worked out.

R. Dedekind.

# FURTHER DEVELOPMENT OF THE INVESTIGATIONS OF PURE EQUATIONS.

---

**1.**

The method by which, in Article 360 of the *Disquiss. Arithm.*, we showed how to solve the equation $x^n - 1 = 0$ forms a most fruitful and important theory, of which we could touch on only the first elements in that work. We therefore hope that mathematicians will welcome our taking up this subject here again, treating more fully what was only briefly sketched, in part with the proofs omitted, and pursuing in greater depth the developments made since that time.

The exponent $n$ is assumed to be a prime number, and the number $n-1$ to be resolved into the factors $\alpha \times \beta \times \gamma$; further, we denote by $g$ any primitive root modulo $n$. Let $r$ denote an arbitrary root of the equation $x^n - 1 = 0$, and $R$ an arbitrary root of the equation $x^\beta - 1 = 0$. Thus, denoting the circumference of the circle whose radius $= 1$ by $P$, and the imaginary quantity $\sqrt{-1}$ by $i$, all the roots of the equation $x^\beta - 1 = 0$, or all the values of $R$, will be represented by the formula

$$\cos \frac{kP}{\theta} + i \sin \frac{kP}{\theta}$$

where $k$ ranges over the integers $0, 1, 2, 3 \ldots \beta - 1$. Further, all powers of any root $R$ are evidently themselves roots, and if $R$ is the root corresponding to a value of $k$ coprime to $\beta$, all the powers $R^0, R, R^2, R^3 \ldots R^{\beta-1}$ are distinct from one another and thus exhaust the entire collection of roots. In this case we shall call $R$ a *proper* root of the equation $x^\beta - 1 = 0$; conversely, a root $R$ corresponding to a value of $k$ that is not coprime to $\beta$ will be called *improper*. It is readily seen that, if $\delta$ is the greatest common divisor of the numbers $k$ and $\beta$, then $R^{\frac{\beta}{\delta}\cdot} = 1$, while all the powers $R^0, R, R^2, R^3 \ldots R^{\frac{\beta}{\delta}-1}$ are distinct from one another, so that $R$ is a proper root of the equation $x^{\frac{\beta}{\delta}} - 1 = 0$. The same holds for the equation $x^n - 1 = 0$, but all its roots are necessarily proper except for the root 1.

**2.**

With these preliminaries, our investigation will chiefly concern functions of the following form, consisting of $\beta\gamma$ terms,

$$r + Rr^{g^\alpha} + R^2 r^{g^{2\alpha}} + R^3 r^{g^{3\alpha}} \ldots + R^{\beta\gamma-1} r^{g^{\alpha\beta\gamma-\alpha}}$$

which for brevity we shall denote by the symbol $[r, R]$. The individual terms of such an expression are products of powers of $r$ and powers of $R$; the exponents of the former form a geometric progression, and those of the latter an arithmetic progression. The exponents

$$1, \quad g^\alpha, \quad g^{2\alpha}, \quad g^{3\alpha} \ldots\ldots g^{\alpha\beta\gamma-\alpha}$$

are all incongruent to one another modulo $n$, and hence those powers of $r$ are distinct from one another; continued further, however, they would begin the same series again, since $g^{\alpha\beta\gamma} \equiv 1 (\mathrm{mod.}\, n)$, and hence $r^{g^{\alpha\beta\gamma}} = r$. The other factors,

$$1, \quad R, \quad R^2, \quad R^3 \ldots . R^{\beta\gamma-1}$$

form $\gamma$ equal periods, since $R^\beta = 1$, $R^{\beta-1} = R$, etc. It follows that the function $[r, R]$ can also be represented as

$$\begin{aligned}
&r + r^{g^{\alpha\beta}} + r^{g^{2\alpha\beta}} \ldots\ldots + r^{g^{\alpha\beta\gamma-\alpha\beta}} \\
&+ R\big(r^{g^\alpha} + r^{g^{\alpha\beta+\alpha}} + r^{g^{2\alpha\beta+\alpha}} \ldots\ldots + r^{g^{\alpha\beta\gamma-\alpha\beta+\alpha}}\big) \\
&+ R^2\big(r^{g^{2\alpha}} + r^{g^{\alpha\beta+2\alpha}} + r^{g^{2\alpha\beta+2\alpha}} \ldots\ldots + r^{g^{\alpha\beta\gamma-\alpha\beta+2\alpha}}\big) \\
&+ \text{etc.} \\
&+ R^{\beta-1}\big(r^{g^{\alpha\beta-\alpha}} + r^{g^{2\alpha\beta-\alpha}} + r^{g^{3\alpha\beta-\alpha}} \ldots\ldots + r^{g^{\alpha\beta\gamma-\alpha}}\big).
\end{aligned}$$

or, introducing the notation of Article 343 of the Disq. Ar.,

$$[r, R] = (\gamma, 1) + R(\gamma, g^\alpha) + R^2(\gamma, g^{2\alpha}) \ldots + R^{\beta-1}(\gamma, g^{\alpha\beta-\alpha})$$

**3.**

If we take unity for the root $r$, we have

$$[1, R] = 1 + R + R^2 + R^3 \ldots + R^{\beta\gamma-1} = \gamma(1 + R + R^2 + R^3 \ldots + R^{\beta-1})$$

whose value will be $= \beta\gamma$ if $R$ is also taken to be the root 1, but $= 0$ for every other value of $R$. Conversely, leaving $r$ indeterminate and putting $R = 1$, we obtain $[r, 1] = r + r^{g^\alpha} + r^{g^{2\alpha}} + r^{g^{3\alpha}} \ldots + r^{g^{\alpha\beta\gamma-\alpha}}$, or, using the notation introduced in the Disq. Ar., $[r, 1] = (\beta\gamma, 1)$; that is, it will consist of a period of $\beta\gamma$ roots, one of which is $r$ itself. Whenever $\alpha = 1$, this period will contain all the roots $r, r^2, r^3 \ldots r^{n-1}$, with only their order changed.

Note also the following relations, whose reason is evident:

$$[r, R] = R[r^{g^\alpha}, R] = R^2[r^{g^{2\alpha}}, R] \quad \text{or generally} \quad = R^k[r^{g^{\alpha k}}, R]$$

where $k$ denotes any positive integer. It follows that the function $[r^m, R]$ either is $= [1, R]$, namely if $m$ is divisible by $n$, or can be reduced in the remaining cases to the form $R^\mu[r^{g^\nu}, R]$, with $\nu < \alpha$. For if $m$ is not divisible by $n$, it will be congruent modulo $n$ to some power of $g$, whose exponent is conveniently expressed by $\operatorname{ind}. m$, as in the Disq. Ar.; thus, putting $\operatorname{ind}. m = \lambda\alpha + \nu$, which can evidently be done with $\nu < \alpha$, we obtain $[r^m, R] = [r^{g^{\lambda\alpha+\nu}}, R] = R^{-\lambda}[r^{g^\nu}, R]$. We must therefore set $\mu = -\lambda$ or, if you want a positive exponent, $\mu \equiv -\lambda (\operatorname{mod}. \beta)$.

**4.**

THEOREM. *If $r'$, like $r$, denotes an arbitrary root of the equation $x^n - 1 = 0$, and $R'$, like $R$, an arbitrary root of the equation $x^\beta - 1 = 0$, the product will be*

$$\begin{aligned}[r, R] \times [r', R'] = \\ [rr', RR'] + R[r^{g^\alpha} r', RR'] + R^2[r^{g^{2\alpha}} r', RR'] \\ + R^3[r^{g^{3\alpha}} r', RR'] \ldots + R^{\beta\gamma-1}[r^{g^{\alpha\beta\gamma-\alpha}} r', RR']\end{aligned}$$

*Proof.* By multiplying $[r, R]$ by the individual terms of $[r', R']$, the product can be represented in the form

$$\begin{aligned}&[r, R]r' + RR'[r^{g^\alpha}, R]r'^{g^\alpha} + R^2R'^2[r^{g^{2\alpha}}, R]r'^{g^{2\alpha}} \\ &+ R^3R'^3[r^{g^{3\alpha}}, R]r'^{g^{3\alpha}} \ldots \\ &+ R^{\beta\gamma-1}R'^{\beta\gamma-1}[r^{g^{\alpha\beta\gamma-\alpha}}, R]r'^{g^{\alpha\beta\gamma-\alpha}}\end{aligned}$$

Then collecting the first terms of the individual parts, after duly expanding them, gives $[rr', RR']$; similarly, collecting the second terms gives $R[r^{g^\alpha} r', RR']$, and so on, finally producing the stated form of the product. Q. E. D.

Moreover, simply interchanging $r, R$ with $r', R'$ shows that the same product can also be put in the form

$$[rr', RR'] + R'[rr'^{g^\alpha}, RR'] + R'^2[rr'^{g^{2\alpha}}, RR'] + R'^3[rr'^{g^{3\alpha}}, RR'] \ldots + R'^{\beta\gamma-1}[rr'^{g^{\alpha\beta\gamma-\alpha}}, RR']$$

It follows further that, if $r'', r'''$, etc., also denote arbitrary roots of the equation $x^n - 1 = 0$, and $R'', R'''$, etc., arbitrary roots of the equation $x^\beta - 1 = 0$, the product of the functions $[r, R], [r', R'], [r'', R''], [r''', R'''],$ etc., however many there are, will equal the sum

$$\Sigma R'^{k'} R''^{k''} R'''^{k'''} \text{ etc.}$$
$$[rr'^{g^{\alpha k'}} r''^{g^{\alpha k''}} r'''^{g^{\alpha k'''}} \text{ etc.},\ RR'R''R''' \text{ etc.}]$$

For $k', k'', k'''$, etc., we substitute all the numbers $0, 1, 2, 3 \ldots \beta\gamma - 1$, combined with one another in every possible different way. This gives altogether $\beta^{\mu-1}\gamma^{\mu-1}$ terms, if $\mu$ denotes the number of those functions multiplied together.

**5.**

The formula by which we expressed the product of any number of functions in the preceding article is general and assumes no relation between the roots $r, r', r'', r'''$, etc., or between $R, R', R'', R'''$, etc. We can readily deduce that, if the roots $r', r'', r'''$, etc., may be regarded as powers of $r$, and the roots $R', R'', R'''$, etc., as powers of $R$, each term of the product will have the form $R^M[r^m, R^\lambda]$, where the exponent $\lambda$ is the same for every term, namely $R^\lambda = RR'R''R'''$, etc. Thus, by what we noted in Article 3, such a product reduces to the following form:

$$A[1, R^\lambda] + B[r, R^\lambda] + B'[r^g, R^\lambda] + B''[r^{g^2}, R^\lambda] + B'''[r^{g^3}, R^\lambda] + \text{etc.} + B^{(\alpha-1)}[r^{g^{\alpha-1}}, R^\lambda]$$

where each of the coefficients $A, B, B', B'', B'''$, etc., will have the form

$$h + h'R + h''R^2 + h'''R^3 + \text{etc.} + h^{(\beta-1)}R^{\beta-1}$$

and $h, h', h'', h'''$, etc., denote fixed integers.

The simplest case is that in which we put $r = r' = r'' = r'''$, etc., and also $R = R' = R'' = R'''$, etc.; our product then becomes the power $[r, R]^\lambda$, which therefore always reduces to the form given above.

**6.**

Thus, putting $\lambda = \beta$, the power $[r, R]^\beta$ takes the form

$$\begin{aligned} &A[1,1] + B[r,1] + B'[r^g,1] + \text{etc.} + B^{\alpha-1)}[r^{g^{\alpha-1}},1] \\ &= \beta\gamma A + B(\beta\gamma,1) + B'(\beta\gamma,g) + B''(\beta\gamma,g^2) + \text{etc.} \\ &\quad + B^{(\alpha-1)}(\beta\gamma,g^{\alpha-1}) = \theta' \end{aligned}$$

If, then, not only the value of the root $R$ (and hence the values of the coefficients $A, B, B'$, etc.) but also the values of the individual sums of $\beta\gamma$ terms $(\beta\gamma,1),(\beta\gamma,g)$, etc., are assumed known, the value of $\theta'$ will be known at once, and from it $[r, R]$ can be found by the formula $\sqrt[\beta]{\theta'}$. This expression admits $\beta$ distinct values, so it might seem doubtful which one should be taken; but it is easily shown that this choice is entirely arbitrary whenever $R$ is a *proper* root of the equation $x^\beta - 1 = 0$. For in this case those $\beta$ values of the radical expression $\sqrt[\beta]{\theta'}$ are evidently

$$[r,R], \quad [r^{g^\alpha},R], \quad [r^{g^{2\alpha}},R] \ldots . [r^{g^{\alpha\beta-\alpha}},R]$$

since, by Article 3, the powers of these functions with exponent $\beta$ are equal to one another, while the functions themselves are proportional to the $\beta$ distinct roots of the equation $x^\beta - 1 = 0$. But as long as only the sums of $\beta\gamma$ terms $(\beta\gamma,1),(\beta\gamma,g)$, etc., are known, the root $r$ itself is determined only to the extent that it must lie in the collection $(\beta\gamma,1)$, and it remains arbitrary which member of this collection we take for $r$. These roots are $r, r^{g^\alpha}, r^{g^{2\alpha}}$, etc., and therefore we can also take any of the functions $[r,R],[r^{g^\alpha},R],[r^{g^{2\alpha}},R]$, etc., for $[r, R]$.

These conclusions would not hold if $R$ were not a proper root of the equation $x^\beta - 1 = 0$. For if $R$ is assumed to be a proper root of the equation $x^{\beta'} - 1 = 0$, with $\beta'$ a divisor of $\beta$, it is readily seen that

$$[r,R] = [r^{g^{\alpha\beta'}},R], \quad [r^{g^\alpha},R] = [r^{g^{\alpha\beta'+\alpha}},R] \text{ etc.}$$

so that in the collection of $\beta$ functions $[r,R],[r^{g^\alpha},R] \ldots [r^{g^{\alpha\beta-\alpha}},R]$ only $\beta'$ distinct functions occur, and therefore among the values of the expression $\sqrt[\beta]{\theta'}$ no more than $\beta'$ are admissible, while the remaining $\beta - \beta'$ are extraneous. It is readily seen, however, that in this case we need not go as far as the power with exponent $\beta$ of the function $[r, R]$ since already the power $[r, R]^{\beta'}$ reduces to our form

$$\beta\gamma A + B(\beta\gamma,1) + B'(\beta\gamma,g) + B''(\beta\gamma,g^2) \text{ etc.}$$

Thus we shall obtain $[r, R]$ by an expression such as $\sqrt[\beta']{\theta}$, and it will make no difference which value of this expression we take.

**7.**

In the same way as $[r, R]$, we can also determine the functions $[r, R^2]$, $[r, R^3]$, etc., or in general $[r, R^k]$. For if substituting in $\theta'$, in place of $R$, the powers $R^2, R^3$, etc., $R^k$, is assumed to give the functions $\theta'', \theta'''$, etc., $\theta^{(k)}$, then evidently $[r, R^2]^\beta = \theta''$, $[r, R^3]^\beta = \theta'''$, etc., and in general $[r, R^k]^\beta = \theta^{(k)}$; hence these functions too can be expressed by radicals, $[r, R^2] = \sqrt[\beta]{\theta''}$, etc. But it is not suitable to use these radical expressions when a quantity is to be expressed by a function of $[r, R]$, $[r, R^2]$, etc. For since their individual values are not fully determined, it would remain doubtful which values could be combined with one another; yet this is evidently by no means arbitrary. It is readily seen that, as soon as a definite value is taken for $[r, R]$, all of $[r, R^2]$, $[r, R^3]$, etc., must also acquire fully determined values, which the radical expressions do not indicate. We must therefore set these expressions aside and seek others by which $[r, R^2]$, $[r, R^3]$, etc., can be expressed *rationally* in terms of $[r, R]$ and known quantities. We readily accomplish this as follows.

By the theorem of Article 4 and what we showed in Article 5, the product $[r, R^k] \times [r, R]^{\beta-k}$ also reduces to a form such as

$$\beta\gamma A + B(\beta\gamma, 1) + B'(\beta\gamma, g) + B''(\beta\gamma, g^2) + \text{etc.} \\ + B^{(\alpha-1)}(\beta\gamma, g^{\alpha-1})$$

where $A, B, B', B''$, etc., are rational functions of $R$. Thus, putting the products

$$[r, R^2] \times [r, R]^{\beta-2} = \vartheta'' \\ [r, R^3] \times [r, R]^{\beta-3} = \vartheta''' \\ [r, R^4] \times [r, R]^{\beta-4} = \beta'''' \\ \text{etc.}$$

the quantities $\vartheta'', \vartheta''', \vartheta''''$, etc., can also be determined rationally, and

$$[r, R^2] = \frac{\vartheta''}{\theta'}[r, R]^2 \\ [r, R^3] = \frac{\vartheta'''}{\theta'}[r, R]^3 \\ [r, R^4] = \frac{\vartheta''''}{\theta'}[r, R]^4 \\ \text{etc.}$$

These expressions therefore give the values of the functions $[r, R^2]$, $[r, R^3]$, etc., rationally, provided that we do not have $[r.R] = 0$; if this equality held, they would become indeterminate. But we can rigorously prove that $[r, R] = 0$ can never occur whenever $r$ denotes a root different from 1, although we must reserve the presentation of this proof for another occasion, so as not to become too lengthy here.

**8.**

What we set out in the preceding articles is useful when the aim is to descend from periods of $\beta\gamma$ terms to periods of $\gamma$ terms. For it is readily seen that, if $R$ denotes a proper root, we have

$$\begin{aligned}
\beta(\gamma,1) &= (\beta\gamma,1) + [r,R] + [r,R^2] + [r,R^3] + \text{etc.} + [r,R^{\beta-1}] \\
\beta(\gamma,g^\alpha) &= (\beta\gamma,1) + R^{\beta-1}[r,R] + R^{\beta-2}[r,R^2] \\
&\quad + R^{\beta-3}[r,R^3] + \text{etc.} + R[r,R^{\beta-1}] \\
\beta(\gamma,g^{2\alpha}) &= (\beta\gamma,1) + R^{2\beta-2}[r,R] + R^{2\beta-4}[r,R^2] \\
&\quad + R^{2\beta-6}[r,R^3] + \text{etc.} + R^2[r,R^{\beta-1}] \\
&\text{etc.}
\end{aligned}$$

If here, for the individual $[r,R]$, $[r,R^2]$, etc., we took the radical expressions $\sqrt[\beta]{\theta'}$, $\sqrt[\beta]{\theta''}$, etc., the value of each series would be uncertain among $\beta^{\beta-1}$ values; by contrast, when the rational expressions for $[r,R^2]$, etc., are taken, it will be subject to no ambiguity other than that which is unavoidable by the nature of the matter. This observation seems to have escaped the attention of the illustrious LAGRANGE, who thought he had *simplified* our method given in Article 360 of the Disquis. arithm., where we had deliberately passed over radical expressions and proposed only rational ones, by substituting the former for the latter (Traité de la résolution numérique des équations; edition 2, p. 311).

Moreover, it hardly needs mentioning here that, as soon as the values of the periods $(\gamma,1)$, $(\gamma,g^\alpha)$, etc., or of just one of them, have been found, the values of all the remaining periods of $\gamma$ terms can be deduced rationally from them. Thus the descent from periods of $\beta\gamma$ terms to periods of $\gamma$ terms requires solving the equations $x^\beta = 1$, $x^\beta = \theta'$, and the remaining operations are performed rationally.

**9.**

All this had already been treated in almost the same way in the Disquis. Ar.; some things, however, had been added there with their proof omitted, and we think it appropriate to supply that proof here. We stated there that finding the value of the radical quantity $\sqrt[\beta]{\theta'}$, which, since $\theta'$ is an imaginary quantity, seems to require dividing both a ratio and an angle into $\beta$ parts, depends on the latter alone, and that the former can always be reduced to extracting a single square root. We prove this as follows.

Denoting the imaginary quantity $\sqrt{-1}$ by $i$, as above, and writing $\theta' = P + iQ$ and $\sqrt[\beta]{\theta'} = p + iq$, using any one value of the radical and taking $P, Q, p, q$ to be real, it is known that, if the positive quantities $E, e$ and the angles $F, f$ are determined so that $P = E\cos F$, $Q = E\sin F$, $p = e\cos f$, $q = e\sin f$, then $e = \sqrt[\beta]{E}$, and $f$ equals one of the angles

$$\frac{1}{\beta}F, \quad \frac{1}{\beta}(F + 360^0), \quad \frac{1}{\beta}(F + 720^0) \ldots . \frac{1}{\beta}(F + (\beta - 1)360^0)$$

Thus $f$· ger is determined by division of the angle $F$ into $\beta$ parts, but we can avoid extracting the root $\sqrt[\beta]{E}$ as follows. Every product $r^k R^K$ has the same real part as $r^{-k}R^{-K}$, while the imaginary parts containing the factor $i$ in these products are equal and opposite. It follows at once that $[r^{-1}, R^{-1}] = p - iq = e(\cos f - i\sin f)$, and hence

$$[r, R] \times [r^{-1}, R^{-1}] = e^2$$

But by the theorem of Article 4 this product becomes

$$\begin{aligned} &= [1,1] + R[r^{g^\alpha - 1}, 1] + R^2[r^{g^{2\alpha}-1}, 1] + \text{etc.} \\ &\quad + R^{\beta\gamma-1}[r^{g^{\alpha\beta\gamma-\alpha}-1}, 1] \\ &= \beta\gamma + R(\beta\gamma, g^\alpha - 1) + R^2(\beta\gamma, g^{2\alpha} - 1) + \text{etc.} \\ &\quad + R^{\beta\gamma-1}(\beta\gamma, g^{\alpha\beta\gamma-\alpha} - 1) \end{aligned}$$

This quantity can be determined if $R$ and all the periods of $\beta\gamma$ terms are assumed known. Determining $e$ therefore requires only the extraction of a square root.

In the special case $\alpha = 1$, the individual periods $(\beta\gamma, g^\alpha - 1)$, $(\beta\gamma, g^{2\alpha} - 1)$, etc., are evidently $= r + r^2 + r^3 + r^4 + \text{etc.} + r^{n-1}$, and hence

$$\begin{aligned} ee &= \beta\gamma + (R + R^2 + R^3 + \text{etc.} + R^{\beta\gamma-1})(r + r^2 + r^3 + \text{etc.} + r^{n-1}) \\ &= \beta\gamma + 1 = n \end{aligned}$$

provided that $r$ and $R$ are assumed to denote roots different from 1, and therefore we always have $e = \sqrt{n}$ (Disq. arithm., end of Article 360).

**10.**

So far we have conducted our investigation in the greatest generality, to cover all values of the numbers $\alpha, \beta, \gamma$; from now on, however, we shall pass to the more restricted case $\alpha = 1$, which will open the way to most fruitful and elegant investigations. Thus the symbol $[r, R]$ will denote the function

$$r + Rr^g + R^2 r^{g^2} + R^3 r^{g^3} + \text{etc.} + R^{n-2} r^{g^{n-2}}$$

where $n$ is a prime number, $r$ an arbitrary root of the equation $x^n - 1 = 0$ (the root 1 not excluded), $R$ an arbitrary root of the equation $x^\beta - 1 = 0$, with $\beta$ denoting a given divisor of $n - 1$, and finally $g$ an integer that is a fixed primitive root modulo $n$. For brevity we shall also write

$$1 + r + r^2 + r^3 + \text{etc.} + r^{n-1} = s$$
$$1 + R + R^2 + R^3 + \text{etc.} + R^{n-2} = S$$

from which it is evident that $s$ becomes $= n$ for $r = 1$, but $s = 0$ for every other value of $r$, and similarly $S = n - 1$ for $R = 1$, but $S = 0$ for every other value of $R$.

By Article 3 we therefore have $[1, R] = S$, $[r, 1] = s - 1$; further, for every integer value of $m$ not divisible by $n$, $[r^m, R] = R^{-\operatorname{ind} m}[r, R]$, or more generally $[r^m, R^M] = R^{-M \operatorname{ind} m}[r, R^M]$, where $\operatorname{ind} m$ is the exponent of the power of $g$ congruent modulo $n$ to $m$. Applying this transformation to what we showed in Article 5, we find that a product of two or more functions such as $[r^h, R^H]$ reduces to the form

$$A[1, R^\lambda] + B[r, R^\lambda]$$

where $A$ and $B$ are rational functions of $R$ with integer coefficients, and $\lambda$ is the sum of all the values of $H$. It will be of great importance to put such transformations into an efficient algorithm; for this purpose we must first consider more closely the nature of the product of two functions.

**11.**

The product $[r, R^{\mu}] \times [r, R^{\nu}]$ becomes, by the theorem of Article 4, =

$$[r^2, R^{\mu+\nu}] + R^{\mu}[r^{g+1}, R^{\mu+\nu}] + R^{2\mu}[r^{g^2+1}, R^{\mu+\nu}] + R^{3\mu}[r^{g^3+1}, R^{\mu+\nu}] + \text{etc.} + R^{(n-2)\mu}[r^{g^{n-2}+1}, R^{\mu+\nu}]$$

Among the $n-1$ exponents $2, g+1, g^2+1, g^3+1$, etc., $g^{n-2}+1$, exactly one is divisible by $n$, namely $g^{\frac{1}{2}(n-1)}+1$, so that the corresponding term of our sum will be $R^{\frac{1}{2}(n-1)\mu}[1, R^{\mu+\nu}]$. This term will be $= 0$ whenever we do not have $R^{\mu+\nu} = 1$, and $= (n-1)R^{\frac{1}{2}(n-1)\mu} = \pm(n-1)$ for $R^{\mu+\nu} = 1$. The remaining parts of our sum, whose sum we shall put $= \Omega$, are transformed as follows:

$$\begin{aligned}
[r^2, R^{\mu+\nu}] &= R^{-(\mu+\nu)\operatorname{ind} 2}[r, R^{\mu+\nu}] \\
R^{\mu}[r^{g+1}, R^{\mu+\nu}] &= R^{\mu-(\mu+\nu)\operatorname{ind}(g+1)}[r, R^{\mu+\nu}] \\
R^{2\mu}[r^{g^2+1}, R^{\mu+\nu}] &= R^{2\mu-(\mu+\nu)\operatorname{ind}(g^2+1)}[r, R^{\mu+\nu}] \\
R^{3\mu}[r^{g^3+1}, R^{\mu+\nu}] &= R^{3\mu-(\mu+\nu)\operatorname{ind}(g^3+1)}[r.R^{\mu+\nu}] \\
&\text{etc.}
\end{aligned}$$

From this we obtain

$$\text{I.} \qquad \Omega = [r, R^{\mu+\nu}] \times \Sigma R^{\mu \operatorname{ind} x - (\mu+\nu)\operatorname{ind}(x+1)}$$

if for $x$ we successively substitute the values $1, g, g^2, g^3 \ldots . g^{n-2}$, except for $g^{\frac{1}{2}(n-1)}$, or, which evidently amounts to the same thing, if for $x$ we substitute the values $1, 2, 3, 4 \ldots . n-2$, since these values are congruent to those modulo $n$, though their order is changed.

Taking the integer $y$ inverse to $x$ modulo $n$, that is, determined so that $xy \equiv 1 \pmod{n}$, we have $\operatorname{ind} x \equiv -\operatorname{ind} y \pmod{n-1}$ and $\operatorname{ind}(x+1) + \operatorname{ind} y \equiv \operatorname{ind}(xy+y) \equiv \operatorname{ind}(1+y) \pmod{n-1}$; hence

$$\begin{aligned}
\mu \operatorname{ind} x - (\mu+\nu)\operatorname{ind}(x+1) &\equiv -\mu \operatorname{ind} y - (\mu+\nu)\{\operatorname{ind}(y+1) - \operatorname{ind} y\} \\
&\equiv \nu \operatorname{ind} y - (\mu+\nu)\operatorname{ind}(y+1)
\end{aligned}$$

Therefore, since the inverses of $1, 2, 3 \ldots n-2$ coincide with these same numbers, with only their order changed, we also have

$$\text{II.} \qquad \Omega = [r, R^{\mu+\nu}] \times \Sigma R^{\nu \operatorname{ind} y - (\mu+\nu)\operatorname{ind}(y+1)}$$

substituting for $y$ successively the numbers $1, 2, 3 \ldots n-2$. The same formula follows directly from I, since the numbers $\mu, \nu$ can evidently be interchanged.

Finally, taking the integer $z$ inverse to $x+1$ modulo $n$, or $xz+z \equiv 1 \pmod{n}$, we have $\operatorname{ind}(1-z) \equiv \operatorname{ind} x + \operatorname{ind} z \pmod{n-1}$, $\operatorname{ind}(x+1) \equiv -\operatorname{ind} z \pmod{n-1}$, and hence

$$\begin{aligned}\mu \operatorname{ind} x - (\mu+\nu)\operatorname{ind}(x+1) &\equiv \mu(\operatorname{ind}(1-z) - \operatorname{ind} z) + (\mu+\nu)\operatorname{ind} z \\ &\equiv \mu \operatorname{ind}(1-z) + \nu \operatorname{ind} z\end{aligned}$$

Thus, as $x$ runs through the values $1, 2, 3 \ldots n-2$, the number $z$ must run through the values $2, 3, 4 \ldots n-1$, though in another order, and we obtain the third expression

$$\text{III.} \qquad \Omega = [r, R^{\mu+\nu}] \times \Sigma R^{\mu \operatorname{ind}(1-z)+\nu \operatorname{ind} z}$$

substituting for $z$ successively the values $2, 3, 4 \ldots n-1$, or, if you prefer,

$$\begin{aligned}\text{IV.} \qquad \Omega &= [r, R^{\mu+\nu}] \times \Sigma R^{\mu \operatorname{ind}(n+1-z)+\nu \operatorname{ind} z} \\ &= [r, R^{\mu+\nu}] \times \Sigma R^{\mu \operatorname{ind} z+\nu \operatorname{ind}(n+1-z)}\end{aligned}$$

Since $\operatorname{ind}(1-z) = \frac{1}{2}(n-1) + \operatorname{ind}(z-1)$, our product can also be represented as follows:

$$\begin{aligned}[r, R^{\mu}] \times [r, R^{\nu}] &= R^{\frac{1}{2}(n-1)\mu}\{[1, R^{\mu+\nu}] + [r, R^{\mu+\nu}] \times \Sigma R^{\mu \operatorname{ind}(z-1)+\nu \operatorname{ind} z}\} \\ &= R^{\frac{1}{2}(n-1)\nu}\{[1, R^{\mu+\nu}] + [r, R^{\mu+\nu}] \times \Sigma R^{\mu \operatorname{ind} z+\nu \operatorname{ind}(z-1)}\}\end{aligned}$$

where $z$ is always understood to take the values $2, 3, 4 \ldots . n-1$.

Moreover, in all these formulas the numbers

$$\mu \operatorname{ind} x - (\mu+\nu)\operatorname{ind}(x+1), \quad \nu \operatorname{ind} y - (\mu+\nu)\operatorname{ind}(y+1), \quad \mu \operatorname{ind}(1-z) + \nu \operatorname{ind} z$$

etc., can evidently be replaced by their least residues modulo $\beta$.

If $\mu + \nu \equiv 0 \pmod{\beta}$, we have

$$\begin{aligned}[r, R^{\mu}][r, R^{\nu}] = (n-1)&R^{\frac{1}{2}(n-1)\mu} \\ &+ (r + r^2 + r^3 + \cdots + r^{n-1}) \\ &\times (1 + R^{\mu} + R^{2\mu} + R^{3\mu} + \cdots + R^{(n-2)\mu} - R^{\frac{1}{2}(n-1)\mu})\end{aligned}$$

**12.**

The product $[1, R^\mu] \times [r, R^\nu]$ becomes, by the theorem of Article 4,

$$= [r, R^{\mu+\nu}] + R^\mu [r, R^{\mu+\nu}] + R^{2\mu} [r, R^{\mu+\nu}] + \text{etc.} + R^{(n-2)\mu} [r, R^{\mu+\nu}]$$
$$= [r, R^{\mu+\nu}] \times (1 + R^\mu + R^{2\mu} + R^{3\mu} + \text{etc.} + R^{(n-2)\mu})$$
$$= [r, R^{\mu+\nu}] \times \frac{n-1}{\beta} (1 + R^\mu + R^{2\mu} + R^{3\mu} + \text{etc.} + R^{(\beta-1)\mu})$$

From this the product $[1, R^\mu] \times [1, R^\nu]$ expands into

$$\frac{n-1}{\beta} [1, R^{\mu+\nu}] \times (1 + R^\mu + R^{2\mu} + R^{3\mu} + \text{etc.} + R^{(\beta-1)\mu})$$

The general product $[r^m, R^\mu] \,.\, [r^{m'}, R^{\mu'}]$ can now be found without difficulty. For $[r^m, R^\mu] = R^{-\mu \operatorname{ind} m} [r, R^\mu]$ when the value of $m$ is not divisible by $n$, and $= [1, R^\mu]$ when it is divisible; and since a similar transformation holds for the other factor $[n^{m'}, R^{\mu'}]$, the multiplication reduces either to the problem of the preceding article or to the cases considered in this article.

**13.**

Now that we have shown how to expand a product of two factors, expanding a product of more factors will present no difficulty. Once the product $[r, R^\mu] \times [r, R^\nu]$ has been reduced to the form $A[1, R^{\mu+\nu}] + B[r, R^{\mu+\nu}]$, it is evident that, on introducing a third factor $[r, R^\pi]$, the product becomes $= C[1, R^{\mu+\nu+\pi}] + D[r, R^{\mu+\nu+\pi}]$ if we put

$$[r, R^{\mu+\nu}][r, R^\pi] = c[1, R^{\mu+\nu+\pi}] + d[r, R^{\mu+\nu+\pi}]$$

and

$$C = Bc$$
$$D = Bd + A\{1 + R^{\mu+\nu} + R^{2\mu+2\nu} + \text{etc.} + R^{(n-2)(\mu+\nu)}\}$$

Thus $[r, R]^\lambda$ can easily be reduced to the form $A[1, R^\lambda] + B[r, R^\lambda]$.

As an example, we shall expand the powers of the function $[r, R]$ for $n = 11$, $\beta = 5$, setting $g = 2$. The following indices then correspond to the numbers:

| | |
|---|---|
| numbers | 1.2.3.4.5.6.7.8.9.10 |
| indices | 0.1.8.2.4.9.7.3.6. 5 |

Thus, to expand the square $[r, R]^2$ by formula I of Article 11, we have

$$\mu = 1, \quad \nu = 1$$

| | |
|---|---|
| values of $x$ ... | 1. 2.3.4. 5. 6.7. 8. 9 |
| $\operatorname{ind} x$ ... | 0. 1.8.2. 4. 9.7. 3. 6 |
| $2 \operatorname{ind}(x+1)$ ... | 2.16.4.8.18.14.6.12.10 |
| Least residues of | $\operatorname{ind} x - 2 \operatorname{ind}(x+1)$ |
| modulo 5 ... | 3. 0.4.4. 1. 0.1. 1. 1 |

from which we deduce

$$\Omega = [r, R^2] \times \{2 + 4R + R^3 + 2R^4\}$$

and

$$1^0. \qquad [r, R]^2 = [1, R^2] + [r, R^2] \times \{2 + 4R + R^3 + 2R^4\}$$

The same expression follows from formula III of Article 11, namely

| | |
|---|---|
| values of $z$ ... | 2.3.4.5.6.7.8.9. 10 |
| $\operatorname{ind} z$ ... | 1.8.2.4.9.7.3.6. 5 |
| $\operatorname{ind}(n+1-z)$ ... | 5.6.3.7.9.4.2.8. 1 |
| least residues of | $\operatorname{ind} z + \operatorname{ind}(n+1-z)$ |
| modulo 5 ... | 1.4.0.1.3.1.0.4. 1 |

In exactly the same way we find

$$2^0. \qquad [r, R^2] \,.\, [r, R] = [1, R^3] + [r, R^3] \times \{2 + R + 4R^2 + 2R^3\}$$
$$3^0. \qquad [r, R^3] \,.\, [r, R] = [1, R^4] + [r, R^4] \times \{2 + 4R + R^3 + 2R^4\}$$

Finally, we obtain

$$4^0. \qquad [r, R^4] \,.\, [r, R] = [1, 1] + [r, 1] \times \{1 + 2R + 2R^2 + 2R^3 + 2R^4\}$$

Multiplying equation $1^0$ by $[r, R]$ and substituting for $[r, R^2] \,.\, [r, R]$ its value from $2^0$, and using

$$[1, R^2] \,.\, [r, R] = [r, R^3] \,.\, \{2 + 2R + 2R^2 + 2R^3 + 2R^4\}$$

we deduce

$$\begin{aligned}[r, R]^3 = {} & [1, R^3] \times \{2 + 4R + R^3 + 2R^4\} \\ & + [r, R^3] \times \{12 + 22R + 18R^2 + 24R^3 + 15R^4\}\end{aligned}$$

and in the same way

$$\begin{aligned}[r, R]^4 = {} & [1, R^4] \times \{12 + 22R + 18R^2 + 24R^3 + 15R^4\} \\ & + [r, R^4] \times \{164 + 170R + 205R^2 + 180R^3 + 190R^4\} \\ [r, R]^5 = {} & [1, 1] \times \{164 + 170R + 205R^2 + 180R^3 + 190R^4\} \\ & + [r, 1] \times \{1836 + 1830R + 1795R^2 + 1820R^3 + 1810R^4\} \\ = {} & 1640 + 1700R + 2050R^2 + 1800R^3 + 1900R^4 \\ & + (1836 + 1830R + 1795R^2 + 1820R^3 + 1810R^4)(s - 1) \\ = {} & 918Ss - 98S - (6R + 41R^2 + 16R^3 + 26R^4)s \\ & + 66R + 451R^2 + 176R^3 + 286R^4\end{aligned}$$

**14.**

The calculation in the preceding articles' was carried out so that it could be extended to all values of $r$ and $R$. It becomes considerably shorter if we regard $R$ from the outset as a proper root of the equation $x^\beta - 1 = 0$. Under this assumption the product $[r, R^\mu] \times [r, R^\nu]$ reduces to the form $B[r, R^{\mu+\nu}]$ whenever $\mu + \nu$ is not divisible by $\beta$; but when $\mu + \nu$ is divisible by $\beta$, that product becomes $= (n-1)R^{\frac{1}{2}(n-1)\mu} + [r, 1]\Sigma R^{\mu \operatorname{ind} x}$, substituting for $\operatorname{ind} x$ all the numbers $0, 1, 2, 3 \ldots . n-2$ except $\frac{1}{2}(n-1)$. It follows easily that in this case, if $\mu$ and therefore also $\nu$ are not divisible by $\beta$, we have

$$[r, R^\mu] \,.\, [r, R^\nu] = R^{\frac{1}{2}(n-1)\mu}\{n - 1 - [r, 1]\}$$

and hence $= 0$ for $r = 1$, and $= nR^{\frac{1}{2}(n-1)\mu}$ for every other value of $r$. Moreover, since $R^{\frac{1}{2}(n-1)\mu}$ becomes $= +1$ or $= -1$ according as $\frac{n-1}{\beta} \,.\, \mu$ is even or odd, our product becomes $= n$ in the former case and $= -n$ in the latter.

It further follows that we can put

$$\begin{aligned} [r, R]^2 &= A'[r, R^2] \\ [r, R^2] \,.\, [r, R] &= A''[r, R^3] \\ [r, R^3] \,.\, [r, R] &= A'''[r, R^4] \end{aligned}$$

and so on, up to

$$[r, R^{\beta-2}] \,.\, [r, R] = A^{(\beta-2)}[r, R^{\beta-1}]$$

from which we have

$$\begin{aligned} [r, R]^2 &= A'[r, R^2] \\ [r, R]^3 &= A'A''[r, R^3] \\ [r, R]^4 &= A'A''A'''[r, R^4] \end{aligned}$$

and so on. Finally,

$$[r, R]^\beta = \pm nA'A''A''' \ldots A^{(\beta-2)}$$

where the upper or lower sign is to be taken according as $\frac{n-1}{\beta}$ is even or odd.

It is therefore evident that, once the value of $[r, R]$ has been found, the remaining functions

$$[r, R^2] = \frac{[r, R]^2}{A'}, \quad [r, R^3] = \frac{[r, R]^3}{A'A''} \quad \text{etc.}$$

can be determined here much more readily than in the cases where the condition $\alpha = 1$ does not hold, as we already noted in the *Disq. Ar.* (Article 360, III). A fuller consideration of the nature of the functions $A', A''$, etc., makes these operations still easier.

**15.**

In Article 9 we showed that the value of the function $[r, R]$ can be reduced to the form $\sqrt{n}(\cos f + i \sin f)$, and in the same way the functions $[r, R^2]$, $[r, R^3]$, etc., up to $[r, R^{\beta-1}]$, can be reduced to a similar form. Let us put

$$\begin{aligned} [r, R] &= \sqrt{n}(\cos f' + i \sin f') \\ [r, R^2] &= \sqrt{n}(\cos f'' + i \sin f'') \\ [r, R^3] &= \sqrt{n}(\cos f''' + i \sin f''') \\ &\text{etc.} \end{aligned}$$

Then

$$\begin{aligned} A' &= \sqrt{n}(\cos(2f' - f'') + i \sin(2f' - f'')) \\ A'' &= \sqrt{n}(\cos(f' + f'' - f''') + i \sin(f' + f'' - f''')) \\ A''' &= \sqrt{n}(\cos(f' + f''' - f'''') + i \sin(f' + f''' - f'''')) \\ &\text{etc.} \end{aligned}$$

It is evident from this that, if the functions $A', A'', A'''$, etc., are reduced to the forms

$$\begin{aligned} A' &= a'(\cos b' + i \sin b') \\ A'' &= a''(\cos b'' + i \sin b'') \\ A''' &= a'''(\cos b''' + i \sin b''') \\ &\text{etc.} \end{aligned}$$

with all of $a', a'', a'''$, etc., positive, we shall have

$$\begin{aligned} a' &= a'' = a''' \text{ etc.} = \sqrt{n} \\ f' &= \frac{1}{\beta}(b' + b'' + b''' + \text{etc.} + b^{(\beta-2)}) \end{aligned}$$

if $\frac{n-1}{\beta}$ is even, or

$$f' = \frac{1}{\beta}(180^0 + b' + b'' + \text{etc.} + b^{(\beta-2)})$$

if $\frac{n-1}{\beta}$ is odd, and then

$$\begin{aligned} [r, R] &= \sqrt{n}(\cos f' + i \sin f') \\ [r, R^2] &= \sqrt{n}(\cos(2f' - b') + i \sin(2f' - b')) \\ [r, R^3] &= \sqrt{n}(\cos(3f' - b' - b'') + i \sin(3f' - b' - b'')) \\ &\text{etc.} \end{aligned}$$

Finally, by the formulas of Article 8,

$$\left(\frac{n-1}{\beta},1\right) = -\frac{1}{\beta} + \frac{\sqrt{n}}{\beta}\{\cos f'' + \cos(2f' - b') + \cos(3f' - b' - b'') + \text{etc.}$$
$$+ \cos((\beta-1)f' - b' - b'' - b''' - \text{etc.} - b^{(\beta-2)})\}$$
$$+ \frac{i\sqrt{n}}{\beta}\{\sin f' + \sin(2f' - b') + \sin(3f' - b' - b'') + \text{etc.}\}$$

and similarly we obtain the values of the functions $\left(\frac{n-1}{\beta},g\right)$, $\left(\frac{n-1}{\beta},g^2\right)$, $\left(\frac{n-1}{\beta},g^3\right)$, etc., if in this formula $f'$ is replaced respectively by $f' - \frac{360^0 k}{\beta}$, $f' - 2\frac{360^0 k}{\beta}$, $f' - 3\frac{360^0 k}{\beta}$, etc., assuming $R = \cos\frac{360^0 k}{\beta} + i\sin\frac{360^0 k}{\beta}$.

**16.**

A further simplification follows from the observation below. Since by Article 14 we have

$$\pm[r,R][r,R^{\beta-1}] = [r,R^2][r,R^{\beta-2}] = \pm[r,R^3][r,R^{\beta-3}] \text{ etc.} = n$$

taking [in the first, third, etc., product] the upper or lower sign according as $\frac{n-1}{\beta}$ is even or odd, in the former case we must have

$$\cos(f' + f^{(\beta-1)}) = \cos(f'' + f^{(\beta-2)}) = \cos(f''' + f^{(\beta-3)}) \text{ etc.} = 1$$

in the latter

$$-\cos(f' + f^{(\beta-1)}) = \cos(f'' + f^{(\beta-2)}) = -\cos(f''' + f^{(\beta-3)}) \text{ etc.} = 1$$

and in both cases

$$\sin(f' + f^{(\beta-1)}) = \sin(f'' + f^{(\beta-2)}) = \sin(f''' + f^{(\beta-3)}) \text{ etc.} = 0$$

We may therefore put, in the former case,

$$f^{(\beta-1)} = -f', \quad f^{(\beta-2)} = -f'', \quad f^{(\beta-3)} = -f''' \quad \text{etc.}$$

in the latter

$$f^{(\beta-1)} = 180^0 - f', \quad f^{(\beta-2)} = -f'', \quad f^{(\beta-3)} = 180^0 - f''' \quad \text{etc.}$$

From this it follows that, in the former case,

$$b^{(\beta-2)} = b', \quad b^{(\beta-3)} = b'', \quad b^{(\beta-4)} = b''' \quad \text{etc.}$$
$$A^{(\beta-2)} = A', \quad A^{(\beta-3)} = A'', \quad A^{(\beta-4)} = A''' \quad \text{etc.}$$

and in the latter

$$b^{(\beta-2)} = b' - 180^0, \quad b^{(\beta-3)} = b'' + 180^0, \quad b^{(\beta-4)} = b''' - 180^0 \quad \text{etc.}$$
$$A^{(\beta-2)} = -A', \qquad A^{(\beta-3)} = -A'', \qquad A^{(\beta-4)} = -A''' \quad \text{etc.}$$

so that the number of functions $A', A'', A'''$, etc., is reduced by half. It further follows that, in the former case,

$$f' = \frac{1}{\beta}(2b' + 2b'' + \text{etc.} + 2b^{(\frac{1}{2}\beta-1)})$$

$$\begin{aligned}\left(\frac{n-1}{\beta}, 1\right) = -\frac{1}{\beta} + \frac{\sqrt{n}}{\beta}\{&2\cos f' + 2\cos(2f' - b') \\ &+ 2\cos(3f' - b' - b'') + \text{etc.} \\ &+ 2\cos((\tfrac{1}{2}\beta - 1)f' - b' - b'' - \text{etc.} - b^{(\frac{1}{8}\beta-2)}) \\ &+ \cos(\tfrac{1}{2}\beta f' - b' - b'' - \text{etc.} - b^{(\frac{1}{2}\beta-1)})\}\end{aligned}$$

(where the last term is evidently $= \cos 0 = 1$) or

$$f' = \frac{1}{\beta}(2b' + 2b'' + \text{etc.} + 2b^{(\frac{1}{2}(\beta-3))} + b^{(\frac{1}{2}(\beta-1))})$$

$$\begin{aligned}\left(\frac{n-1}{\beta}, 1\right) = -\frac{1}{\beta} + \frac{\sqrt{n}}{\beta}\{&2\cos f' + 2\cos(2f' - b') \\ &+ 2\cos(3f' - b' - b'') + \text{etc.} \\ &+ 2\cos(\tfrac{1}{2}(\beta - 1)f' - b' - b'' - \text{etc.} - b^{(\frac{1}{2}(\beta-3))})\}\end{aligned}$$

according as $\beta$ is even or odd; and in the latter case,

$$f' = \frac{1}{\beta}(2b' + 2b'' + \text{etc.} + 2b^{(\frac{1}{2}\beta-1)})$$

$$\begin{aligned}\left(\frac{n-1}{\beta}, 1\right) = -\frac{1}{\beta} + \frac{\sqrt{n}}{\beta}\{&2\cos(2f' - b') + 2\cos(4f' - b' - b'' - b''') + \text{etc.} \\ &+ 2\cos((\tfrac{1}{2}\beta - 2)f' - b' - b'' - \text{etc.} - b^{(\frac{1}{2}\beta-3)}) \\ &+ \cos(\tfrac{1}{2}\beta f' - b' - b'' - \text{etc.} - b^{(\frac{1}{2}\beta-1)})\} \\ + i\frac{\sqrt{n}}{\beta}\{&2\sin f' + 2\sin(3f' - b' - b'') + \text{etc.} \\ &+ 2\sin((\tfrac{1}{2}\beta - 1)f' - b' - b'' - \text{etc.} - b^{(\frac{1}{2}\beta-2)})\}\end{aligned}$$

or

$$f' = \frac{1}{\beta}(2b' + 2b'' + \text{etc.} + 2b^{(\frac{1}{2}\beta-1)} + 180^0)$$

$$\begin{aligned}\left(\frac{n-1}{\beta}, 1\right) = -\frac{1}{\beta} + \frac{\sqrt{n}}{\beta}\{&2\cos(2f' - b') + 2\cos(4f' - b' - b'' - b''') + \text{etc.}\\ &+ 2\cos((\tfrac{1}{2}\beta - 1)f' - b' - b'' - \text{etc.} - b^{(\frac{1}{2}\beta-2)})\}\\ + i\frac{\sqrt{n}}{\beta}\{&2\sin f' + 2\sin(3f' - b' - b'') + \text{etc.}\\ &+ 2\sin((\tfrac{1}{2}\beta - 2)f' - b' - b'' - \text{etc.} - b^{(\frac{1}{2}\beta-3)})\\ &+ \sin(\tfrac{1}{2}\beta f' - b' - b'' - \text{etc.} - b^{(\frac{1}{2}\beta-1)})\}\end{aligned}$$

according as $\frac{1}{2}\beta$ is even or odd. What we noted above also holds for the remaining periods of $\frac{n-1}{\beta}$ terms. In general, therefore, we conclude that determining these periods requires the division of the whole circle into $\beta$ parts, on which the construction of the angles $b', b'', b'''$, etc., depends rationally; then the division of the angle $b' + b'' + b''' + \text{etc.}$ into $\beta$ parts; and finally the square root $\sqrt{n}$. If we set $\beta = \frac{1}{2}(n-1)$ at once, those periods evidently coincide with twice the cosines of the angles $\frac{360^0}{n}$, $2\frac{360^0}{n}$, $3\frac{360^0}{n}$, etc., up to $\frac{1}{2}(n-1)\frac{360^0}{n}$, so that dividing the circle into $n$ parts depends on dividing the whole circle into $\frac{1}{2}(n-1)$ parts, dividing an angle that can be constructed after that division into $\frac{1}{2}(n-1)$ parts, and the radical quantity $\sqrt{n}$. If we intend to proceed as far as the sines of the angles $\frac{360^0}{n}$, etc., one further operation will be needed.

**17.**

For a clearer illustration, let us return to the example of Article 13, where we found

$$A' = A''' = 2 + 4R + R^3 + 2R^4 = 2R - 2R^2 - R^3$$
$$A'' = 2 + R + 4R^2 + 2R^3 = -R + 2R^2 - 2R^4$$

Taking $R$ to have the value $\cos 72^0 + i \sin 72^0$, we obtain

$$A' = A''' = 2\cos 72^0 - 3\cos 144^0 + i(2\sin 72^0 - \sin 144^0)$$
$$A'' = -3\cos 72^0 + 2\cos 144^0 + i(\sin 72^0 + 2\sin 144^0)$$

The angles $b', b''$ will therefore be determined by the equations

$$1) \qquad \sin b' = \frac{2\sin 72^0 - \sin 144^0}{\sqrt{11}}$$
$$2) \qquad \cos b' = \frac{2\cos 72^0 - 3\cos 144^0}{\sqrt{11}}$$
$$3) \qquad \operatorname{tang} b' = \frac{2\sin 72^0 - \sin 144^0}{2\cos 72^0 - 3\cos 144^0}$$
$$4) \qquad \sin b'' = \frac{\sin 72^0 + 2\sin 144^0}{\sqrt{11}}$$
$$5) \qquad \cos b'' = \frac{-3\cos 72^0 + 2\cos 144^0}{\sqrt{11}}$$
$$6) \qquad \operatorname{tang} b'' = \frac{\sin 72^0 + 2\sin 144^0}{-3\cos 72^0 + 2\cos 144^0}$$

Any one of the equations $1, 2, 3$ suffices to determine the angle $b'$, provided the quadrant in which it is to be taken is known; this must be decided from the signs of the quantities $2\sin 72^0 - \sin 144^0$, $2\cos 72^0 - 3\cos 144^0$. The same holds for the angle $b''$. In our case $b'$ is to be taken between 0 and $90^0$, and $b''$ between $90^0$ and $180^0$. If the numerator and denominator of equation 3 are multiplied by $-3\cos 72^0 + 2\cos 144^0$, it becomes

$$\operatorname{tang} b' = \frac{2}{31}\{-\sin 72^0 + 13\sin 144^0\}$$

Similarly, from equation 6, on multiplying the numerator and denominator by $2\cos 72^0 - 3\cos 144^0$, we obtain

$$\operatorname{tang} b'' = \frac{2}{31}\{-13\sin 72^0 - \sin 144^0\}$$

Numerically, this gives

$$\begin{aligned}\operatorname{tang} b' &= +0{,}4316226944, \quad \log\operatorname{tang} b' = 9{,}6351042715\\ b' &= 23^0 20' 46'' 04603\\ \operatorname{tang} b'' &= -0{,}8355819332, \quad \log\operatorname{tang} b'' = 9{,}9219890411\text{n}\\ b'' &= 140^0 7' 6'' 52441\end{aligned}$$

from which we derive

$$5f' = 186^0 48' 38'' 61647, \qquad f' = 37^0 21' 43'' 723294$$

.We therefore have

$$\begin{aligned}(2,1) &= -\frac{1}{5} + \frac{\sqrt{11}}{5}\{2\cos 37^0 21' 43'' 723294 + 2\cos 51^0 22' 41'' 400558\}\\ (2,2) &= -\frac{1}{5} + \frac{\sqrt{11}}{5}\{2\cos 325^0 21' 43'' 723294 + 2\cos 267^0 22' 41'' 400558\}\\ (2,4) &= -\frac{1}{5} + \frac{\sqrt{11}}{5}\{2\cos 253^0 21' 43'' 723294 + 2\cos 123^0 22' 41'' 400558\}\\ (2,8) &= -\frac{1}{5} + \frac{\sqrt{11}}{5}\{2\cos 181^0 21' 43'' 723294 + 2\cos 339^0 22' 41'' 400558\}\\ (2,5) &= -\frac{1}{5} + \frac{\sqrt{11}}{5}\{2\cos 109^0 21' 43'' 723294 + 2\cos 195^0 22' 41'' 400558\}\end{aligned}$$

from which we find

$$\begin{array}{lcll}(2,1) & = & +1{,}6825070652 & = 2\cos\frac{360^0}{11}\\ (2,2) & = & +0{,}8308299 & = 2\cos\frac{720^0}{11}\\ (2,4) & = & & = 2\cos\frac{1440^0}{11}\\ (2,8) & = & & = 2\cos\frac{2880^0}{11}\\ (2,5) & = & & = 2\cos\frac{1800^0}{11}\end{array}$$

**18.**

The equation $x^{17} - 1 = 0$, which we already treated by a different method in the *Disquis. Arithm.*, will provide another example. We shall therefore set $n = 17$, $\beta = 8$, $g = 3$; the following indices then correspond to the numbers:

| | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| numbers | 1. | 2. | 3. | 4. | 5. | 6. | 7. | 8. | 9. | 10. | 11. | 12. | 13. | 14. | 15. | 16 |
| indices | 0. | 14. | 1. | 12. | 5. | 15. | 11. | 10. | 2. | 3. | 7. | 13. | 4. | 9. | 6. | 8 |

From this we find

$$\begin{aligned}A' = A'''''' &= 2R + 2R^2 + 3R^4 + 4R^5 + 2R^6 + 2R^7\\ A'' = A'''' &= 2 + 3R + R^3 + R^4 + 3R^5 + 4R^6 + R^7\\ A''' = A'''' &= 3 + 3R + 2R^2 + 3R^3 + R^5 + 2R^6 + R^7\end{aligned}$$

or, since in this case $R^4 + 1 = 0$,

$$A' = A''''' = -3 - 2R - 2R^3$$
$$A'' = A'''' = 1 - 4R^2$$
$$A''' = A'''' = 3 + 2R + 2R^3$$

Putting $R = \cos 45^0 + i \sin 45^0$, we therefore have

$$A' = A''''' = -3 - 2i\sqrt{2},$$
$$A'' = A'''' = 1 - 4i,$$
$$A''' = A'''' = 3 + 2i\sqrt{2}$$

Thus $b', b'', b'''$ will be found from the equations

$$\begin{array}{lll} \sin b' = -\sqrt{\frac{8}{17}} & \sin b'' = -\sqrt{\frac{16}{17}} & \sin b''' = +\sqrt{\frac{8}{17}} \\ \cos b' = -\sqrt{\frac{9}{17}} & \cos b'' = +\sqrt{\frac{1}{17}} & \cos b''' = +\sqrt{\frac{9}{17}} \\ \operatorname{tang} b' = +\sqrt{\frac{8}{9}} & \operatorname{tang} b'' = -4 & \operatorname{tang} b''' = +\sqrt{\frac{8}{9}} \end{array}$$

from which we deduce

$$b' = 223^0 18' 49'', \quad b'' = 284^0 2' 10, \quad b''' = 43^0 18' 49'' = b' - 180^0$$

$$4f' = 550^0 39' 48'', \qquad f' = 137^0 39' 57''$$

$$(2,1) = -\frac{1}{8} + \frac{\sqrt{17}}{8}\{2\cos 137^0 39' 57'' + 2\cos 52^0 1' 5'' + 2\cos 265^0 38' 52'' + 1\}$$
$$(2,4) = -\frac{1}{8} + \frac{\sqrt{17}}{8}\{2\cos 92^0 39' 57'' + 2\cos 322^0 1' 5'' + 2\cos 130^0 38' 52'' - 1\}$$
$$(2,9) = -\frac{1}{8} + \frac{\sqrt{17}}{8}\{2\cos 47^0 39' 57'' + 2\cos 232^0 1' 5'' + 2\cos 355^0 38' 52'' + 1\}$$
$$(2,10) = -\frac{1}{8} + \frac{\sqrt{17}}{8}\{2\cos 2^0 39' 57'' + 2\cos 142^0 1' 5'' + 2\cos 220^0 38' 52'' - 1\}$$
$$(2,13) = -\frac{1}{8} + \frac{\sqrt{17}}{8}\{2\cos 317^0 39' 57'' + 2\cos 52^0 1' 5'' + 2\cos 85^0 38' 52'' + 1\}$$
$$(2,5) = -\frac{1}{8} + \frac{\sqrt{17}}{8}\{2\cos 272^0 39' 57'' + 2\cos 322^0 1' 5'' + 2\cos 310^0 38' 52'' - 1\}$$
$$(2,15) = -\frac{1}{8} + \frac{\sqrt{17}}{8}\{2\cos 227^0 39' 57'' + 2\cos 232^0 1' 5'' + 2\cos 175^0 38' 52'' + 1\}$$
$$(2,11) = -\frac{1}{8} + \frac{\sqrt{17}}{8}\{2\cos 182^0 39' 57'' + 2\cos 142^0 1' 5'' + 2\cos 40^0 38' 52'' - 1\}$$

$$\begin{array}{rcccl}
\frac{1}{2}(2,1) & = & +0{,}092268 & = & \cos\frac{4}{17}360^0 \\
\frac{1}{2}(2,3) & = & & = & \cos\frac{5}{17}360^0 \\
\frac{1}{2}(2,9) & = & & = & \cos\frac{2}{17}360^0 \\
\frac{1}{2}(2,10) & = & & = & \cos\frac{6}{17}360^0 \\
\frac{1}{2}(2,13) & = & +0{,}93247 & = & \cos\frac{1}{17}360^0 \\
\frac{1}{2}(2,5) & = & & = & \cos\frac{3}{17}360^0 \\
\frac{1}{2}(2,15) & = & & = & \cos\frac{8}{17}360^0 \\
\frac{1}{2}(2,11) & = & & = & \cos\frac{7}{17}360^0
\end{array}$$

* *

From these more general investigations of the functions $[r, R]$, which further clarify and extend the second theory of pure equations begun in Article 360 of the *Disquiss. Ar.*, we proceed to consider certain special cases more closely, namely those in which definite values are taken for $\beta$. Several investigations, as fruitful as they are elegant, will emerge from this; some had already been treated in the *Disq. Ar.* (Articles . . .), though by a different method, while others are to be regarded as entirely new. We reserve for another paper, to be published soon, the development of the remarkable connection between these investigations and higher arithmetic, which thereby makes very great and hitherto unexpected advances. — Moreover, throughout the following investigation we shall assume that $r$ is taken to be a *proper* root of the equation $x^n - 1 = 0$, and $R$ a proper root of the equation $R^\beta - 1 = 0$.

**19.**

We begin with the value $\beta = 2$, so that $R$ must take the value $-1$. Our function $[r, R]$ therefore becomes

$$= r - r^g + r^{g^2} - r^{g^3} \ldots . - r^{g^{n-2}}$$

and we have

$$[r, R] = -[r^g, R] = +[r^{g^2}, R] = -[r^{g^3}, R] \quad \text{etc.}$$

and, in general, if $\lambda$ denotes any integer not divisible by $n$,

$$[r^\lambda, R] = +[r, R] \quad \text{if } \lambda \text{ is a quadratic residue of } n,$$
$$[r^\lambda, R] = -[r, R] \quad \text{if } \lambda \text{ is a quadratic nonresidue of } n.$$

Further, if the quadratic residues of $n$ among $1, 2, 3 \ldots n-1$ are denoted generically by $a$, and the nonresidues of $n$ within the same limits by $b$, it is evident that the numbers

$$1, g^2, g^4 \ldots . g^{n-3}$$

with their order disregarded, are congruent modulo $n$ to the numbers $a$, and similarly the numbers

$$g, g^3, g^5 \ldots . g^{n-2}$$

are congruent to $b$, so that $[r, R] = \Sigma r^a - \Sigma r^b$.

If, therefore, we put $\frac{360^0}{n} = \omega$ and $r = \cos k\omega + i \sin k\omega$, then $[r, R] = \Sigma \cos ak\omega - \Sigma \cos bk\omega + i\Sigma \sin ak\omega - i\Sigma \sin bk\omega$. Now, by Article 14, the square of the function $[r, R]$ will be $= +n$ or $= -n$ according as $n$ has the form $4z+1$ or $4z-1$, and hence in the former case $[r, R] = \pm\sqrt{n}$, in the latter $[r.R] = \pm i\sqrt{n}$; the sign placed before the radical quantity, however, remains ambiguous. From this we obtain the following sums:

I. If $n$ has the form $4z+1$,

$$\Sigma \cos ak\omega - \Sigma \cos bk\omega = \pm\sqrt{n}$$
$$\Sigma \sin ak\omega - \Sigma \sin bk\omega = 0$$

II. If $n$ has the form $4z-1$,

$$\Sigma \cos ak\omega - \Sigma \cos bk\omega = 0$$
$$\Sigma \sin ak\omega - \Sigma \sin bk\omega = \pm\sqrt{n}$$

Moreover, since the whole collection of numbers $a, b$ evidently coincides with $1, 2, 3 \ldots n-1$, we have $\Sigma r^a + \Sigma r^b = r + r^2 + r^3 + \text{etc.} + r^{n-1} = -1$, and therefore $\Sigma \cos ak\omega + \Sigma \cos bk\omega = -1$, $\Sigma \sin ak\omega + \Sigma \sin bk\omega = 0$. These identities and the preceding sums give:

I. For the former case,

$$\Sigma \cos ak\omega = -\tfrac{1}{2} \pm \tfrac{1}{2}\sqrt{n}$$
$$\Sigma \cos bk\omega = -\tfrac{1}{2} \mp \tfrac{1}{2}\sqrt{n}$$
$$\Sigma \sin ak\omega = \Sigma \sin bk\omega = 0$$

II. For the latter case,

$$\Sigma \cos ak\omega = \Sigma \cos bk\omega = -\tfrac{1}{2}$$
$$\Sigma \sin ak\omega = \pm\tfrac{1}{2}\sqrt{n}$$
$$\Sigma \sin bk\omega = \mp\tfrac{1}{2}\sqrt{n}$$

These sums were already obtained by a method not very different in Article 356 of the *Disquiss. Arr.*; neither method can remove the ambiguity in the sign to be placed before the radical quantity. We have, however, recently overcome this shortcoming in a separate paper, where it was proved that, for the value $k = 1$, the upper signs must be taken in all the formulas given.

---

## Remarks.

Only a few articles remain of the original continuation of this paper from Article 19 onward, which was devoted to special cases. They deal with the quadratic equation whose roots are the two periods of $\frac{n-1}{2}$ terms; the manuscript stops at the beginning of the investigation that was to determine the sign of the square root occurring in its solution. The agreement of this surviving beginning with the paper Summatio quarumdam serierum singularium shows that the author changed his plan to make the determination of the sign just mentioned the subject of a separate paper. Comparison with the citation in Article 8, where the manuscript mistakenly gave the third edition of Lagrange's work instead of the second, shows that this manuscript dates from 1808. That publication of the preceding material had not been abandoned is shown by a leaf evidently inserted later at Article 19, on which a different continuation begins and which already refers to the paper Summatio, etc., for the determination of the sign. This second continuation, which also soon stops, is presented here. In preparing the edition, the text of the manuscript, which was entirely ready for printing, was faithfully retained; only in Article 16 did the formulas for the second case have to be added.

R. Dedekind.

---

# PROOF OF SOME THEOREMS CONCERNING THE PERIODS OF CLASSES OF BINARY QUADRATIC FORMS.

THEOREM. I. *The number of (properly primitive) classes of the same determinant which, when raised to the $P^{\text{th}}$ power, where $P$ is either a prime number or a prime power $= p^\pi$, produce the principal class $K$, is equal either to* 1 *or to a power of this same prime number $p$.*

*Proof.* Let $(\mathfrak{Q})$ be the whole group of all the classes in question, and $n$ their number. Since the principal class $K$ is necessarily contained in $(\mathfrak{Q})$, the theorem is evident if it is the only one there. But if there are others, the number of classes contained in the period of each will be a power of $p$; let one of them be $A$, and suppose that its period $(\mathfrak{A})$ contains $p^\alpha$ classes, all of which will be included in $(\mathfrak{Q})$. Now, if the classes of this period $(\mathfrak{A})$ exhaust $(\mathfrak{Q})$, we shall have $p^\alpha = n$, and the theorem will be proved; otherwise, let $B$ be any class of $(\mathfrak{Q})$ not contained in $(\mathfrak{A})$, and suppose its period is developed until we reach a class $bB$ which is also among the classes of $(\mathfrak{A})$. This must necessarily happen, because at least the principal class is common to this period and to $(\mathfrak{A})$. Now, supposing that $bB$ is the *first* class in the period of $B$ common to $(\mathfrak{A})$, or that $b$ is as small as possible, I claim

$1^0$. That $b$ will be a power of $p$. For it is evident that, on putting $b = p^\beta h$, $bB = iA$ and $hk \equiv 1 \pmod{p^\pi}$ (which will be possible), we shall have $kbB = p^\beta hkB = p^\beta B = ikA$, that is, $p^\beta B$ will also be among the classes of $(\mathfrak{A})$, from which it follows that $h = 1$ and $b = p^\beta$.

$2^0$. That, if the classes $K, B, 2B \ldots . (b-1)B$ are denoted by $(\mathfrak{B})$, all the compositions of a class of $(\mathfrak{A})$ with a class of $(\mathfrak{B})$ will give $p^{\alpha+\beta}$ different classes. For, supposing $mA+nB = m'A+n'B$ and $n = n'$, we shall necessarily have $m = m'$; if $n > n'$, we shall have $(n-n')B = (m'-m)A$, which is impossible unless $n = n'$.

$3^0$. That these $p^{\alpha+\beta}$ different classes will be included in $(\mathfrak{Q})$, which is evident.

Now, if these $p^{\alpha+\beta}$ classes exhaust $(\mathfrak{Q})$, the theorem is proved; otherwise, we shall choose another class of $(\mathfrak{Q})$ not contained among them, namely $C$; we shall continue its period until we reach a class already included among the classes composed from $(\mathfrak{A})$ and $(\mathfrak{B})$. By reasoning similar to the preceding, we shall prove that the exponent of this class must be a power of $p$, $= p^\gamma$, and that the composition of the first $p^\gamma$ classes of the period of $C$ with the $p^{\alpha+\beta}$ classes already found will give $p^{\alpha+\beta+\gamma}$ different classes, all included in $(\mathfrak{Q})$. If these classes still do not exhaust $(\mathfrak{Q})$, we shall treat a fourth class $D$, etc., in the same way; and it is evident that, since $(\mathfrak{Q})$ consists of a finite number

of classes, these operations will also come to an end, and $n$ will be equal to a power of $p$. Q. E. D.

THEOREM. II. *If the number of all the classes of the principal genus is expressed by $a^\alpha b^\beta c^\gamma$, etc., where $a, b, c$ denote different prime numbers, there will be in this genus $a^\alpha, b^\beta, c^\gamma$, etc., classes which, when raised respectively to the powers $a^\alpha, b^\beta, c^\gamma$, etc., produce the principal class.*

*Proof* Let $A, A', A''$, etc., be all the classes which, when raised to the power $a^\alpha$, produce $K$, and let $(\mathfrak{A})$ be their totality; likewise $B, B', B''$, etc., $(\mathfrak{B})$, $C, C', C''$, $(\mathfrak{C})$, etc., etc. I claim that the composition of all the classes of $(\mathfrak{A})$ with all the classes of $(\mathfrak{B})$ with all the classes of $(\mathfrak{C})$, etc., will produce classes different from one another. For if $A + B + C \cdots = A' + B' + C' \ldots$, etc., then, on putting $A - A' = A''$, $B - B' = B''$, etc., we shall have

$$A'' + B'' + C'' \text{ etc.} = K$$

and hence, on raising to the power $b^\beta c^\gamma$, etc., $(b^\beta c^\gamma \ldots)A'' = K$, from which it follows readily that $A'' = K$ and $A = A'$, and in the same way we shall have $B = B'$, $C = C'$, etc. Let the totality of these classes be $= (S)$. Further, it is clear that all these classes will belong to the principal genus. Finally, there can be no class in the principal genus which is not included in $(S)$. Let . . .

---

REMARK.

This fragment, written in 1801, relates to Disq. Arithm., Article 306, IX. The word *dignité* is used here in a sense that is not otherwise customary.

STERN.

---

[I.]

# ON THE CONNECTION BETWEEN THE NUMBER OF CLASSES INTO WHICH BINARY QUADRATIC FORMS ARE DISTRIBUTED AND THEIR DETERMINANT.

## FIRST PAPER

## PRESENTED TO THE ROYAL SOCIETY 1834 . . . .

---

**1.**

Thirty-three years have now passed since we discovered the principles of the remarkable connection to which this paper is devoted, as was already announced at the end of the *Arithmetical Investigations*. Other occupations, however, had long drawn us away from this inquiry, until more recently we were able to return to it and extend it through renewed efforts. Yet, since this new part of higher arithmetic exceeds the bounds of a single paper, this first paper will be devoted to forms with negative determinants; forms with positive determinants, which require an entirely distinct treatment, must remain reserved for another paper.

**2.**

The basis of the whole subject is a particular investigation of the number of all combinations of integer values taken by two indeterminate integers $x, y$ within a prescribed region. Evidently this problem can also be presented geometrically, by finding the number of *complex numbers* whose representations fall within a prescribed figure. The nature of the figure will depend on that of the line surrounding it, and thus either on a single equation between the coordinates $x, y$ (when the boundary is a curve returning to itself) or on several such equations (when it .consists of several curved or straight parts); it will be at our discretion whether to include in the count, or exclude from it, points corresponding to complex integers that may lie on the boundary itself.

In the analytic presentation of the problem, these conditions of restriction can always be expressed by requiring one or more given functions of the variables $x, y$, namely $P, Q, R$, etc., to take positive or nonnegative values (according as the value 0 is excluded or admitted).

Thus, for example, if the prescribed figure is a circle with radius $= \sqrt{A}$, while its centre lies at a point corresponding to a complex integer, the analytic condition will be that $A - xx - yy$ is not negative, provided that, as we shall always suppose, we choose to retain points lying on the boundary itself.

If the figure is a triangle, the three linear functions $ax+by+c$, $a'x+b'y+c'$, $a''x+b''y+c''$ must have nonnegative values, and similarly in other cases.

**3.**

Generally speaking, an *exact* solution of the problem must proceed by first using the nature of the conditions to confine one variable, for example $x$, within limits between which the individual integer values are then taken successively. We must find how many integer values of the other variable $y$ correspond to each of them, and then add these numbers together. In special cases there will usually be particular devices for shortening the work.

For example, if the figure is, as above, a circle with radius $=\sqrt{A}$, let $r$ be the integer immediately below $\sqrt{A}$, or $\sqrt{A}$ itself if $A$ is a square. Likewise let $r', r'', r'''$, etc., $r^{(r)}$ be the integers immediately below $\sqrt{(A-1)}$, $\sqrt{(A-4)}$, $\sqrt{(A-9)}$, etc., as far as $\sqrt{(A-rr)}$. The required number will then be

$$\begin{aligned} &= 2r+1+2(2r'+1)+2(2r''+1)+2(2r'''+1)+\text{etc.} \\ &= 1+4r+4r'+4r''+4r'''+\text{etc.}+4r^{(r)} \end{aligned}$$

The following method is shorter in this example. Let $q$ be the integer immediately below $\sqrt{\frac{1}{2}A}$ (or equal to it when it is an integer), and let $r^{(q+1)}$, $r^{(q+2)}$, $r^{(q+3)}$, etc., be the integers immediately below $\sqrt{(A-(q+1)^2)}$, $\sqrt{(A-(q+2)^2)}$, $\sqrt{(A-(q+3)^2)}$, etc., as far as $\sqrt{(A-rr)}$. The required number will then be

$$= 4qq+1+4r+8(r^{(q+1)}+r^{(q+2)}+r^{(q+3)}+\text{etc.}+r^{(r)})$$

This formula gave the following numbers:

| $A$ | | $A$ | | $A$ | |
|---|---|---|---|---|---|
| 100 | 317 | 1000 | 3149 | 10000 | 31417 |
| 200 | 633 | 2000 | 6293 | 20000 | 62845 |
| 300 | 949 | 3000 | 9425 | 30000 | 94237 |
| 400 | 1257 | 4000 | 12581 | 40000 | 125629 |
| 500 | 1581 | 5000 | 15705 | 50000 | 157093 |
| 600 | 1885 | 6000 | 18853 | 60000 | 188453 |
| 700 | 2209 | 7000 | 21993 | 70000 | 219901 |
| 800 | 2521 | 8000 | 25137 | 80000 | 251305 |
| 900 | 2821 | 9000 | 28269 | 90000 | 282697 |
| 1000 | 3149 | 10000 | 31417 | 100000 | 314197 |

**4.**

Our purpose does not require an exact determination, but rather the search for an expression that can approach the exact number as closely as desired as the limits are extended to infinity. But first of all, since this involves something vague, the matter must be stated more exactly.

We shall therefore suppose that the functions $P, Q, R$, etc., besides the variables $x, y$, involve a constant element $k$, so that each of $P, Q, R$, etc., is a homogeneous function of the three quantities $x, y, k$. In this way the figure determined by the equations $P = 0, Q = 0, R = 0$, etc., will depend on $k$, so that different values of $k$ correspond to similar figures similarly situated with respect to the origin of coordinates; corresponding linear dimensions will be proportional to the values of $k$, and areas to the values of $kk$. Let the number of points within the figure now be denoted by $M$, and the area by $V$. It is evident that $M$ and $V$ must increase as $k$ increases; as $k$ increases to infinity, $M$ and $V$ will approach a ratio of equality as closely as desired. Or, if you require elementary clarity, given a quantity $\lambda$, however small, a threshold can always be assigned such that, for every value of $k$ exceeding it, $\frac{M}{V}$ must certainly lie between $1 - \lambda$ and $1 + \lambda$. In the customary manner this may be expressed by saying that $M = V$ for an infinite value of $k$.

In our example the required condition holds on putting $k = \sqrt{A}$, and the curve becomes a circle whose area $= \pi A$, where $\pi$ denotes the half-circumference of a circle of radius $= 1$. The numbers given above clearly indicate the convergence.

Moreover, if it were worth the effort, we could easily complete the proof of this theorem with the rigour of the ancients; here, however, we have preferred to omit it, hastening on to more difficult matters.

**5.**

In this paper the boundary will be expressed by a *single* equation of the form $axx + 2bxy + cyy = A$, where $a, b, c$ are integers and $bb - ac$ is a negative number, which we shall put $= -D$. Evidently the curve defining the figure will be an ellipse, and it is readily seen that the squares of the semiaxes are the roots of the equation

$$(ac - bb)qq - (a + c)Aq + AA = 0 \quad \text{or} \quad = A\left(\frac{a + c \pm \sqrt{(4bb + (a - c)^2)}}{2(ac - bb)}\right)$$

The product of these roots is $\frac{AA}{ac-bb} = \frac{AA}{D}$, and hence the area of the ellipse $= \frac{\pi A}{\sqrt{D}}$. We therefore infer that the number of all combinations of integer values of $x, y$ for which $axx + 2bxy + cyy$ does not exceed the value $A$, as $A$ increases, comes ever closer to $\frac{\pi A}{\sqrt{D}}$, and for infinite $A$ must be set equal to that value. Moreover, it is evident that in this respect it makes no difference whether the combination $x = 0, y = 0$ is counted with the others or excluded.

In this way the required number (in the latter sense) is simply the sum of the numbers of representations of the individual numbers $1, 2, 3 \ldots A$ by the binary quadratic form $axx + 2bxy + cyy$; and since some of those numbers cannot be represented by this form at all, while others admit more or fewer representations, the quantity $\frac{\pi}{\sqrt{D}}$ is to be regarded as the mean value of the number of representations of an indeterminate positive number by any form whose determinant $= -D$.

**6.**

Before investigating the consequences in general, we have thought it useful to work out some special cases, so that the manner of argument may be understood more easily. Let us therefore first return to the form $xx + yy$, for which the number of representations of an indeterminate number has the mean value $= \pi$. The number of actual representations of a given number is readily determined from the general principles established in the Arithmetical Investigations. Let $fA$ denote the number of representations of the number $A$; this will be $= 4$ if $A = 1$ or 2 or a power of two; $= 8$ if $A$ is a prime number of the form $4n + 1$, or the product of such a prime and a power of two; $= 0$ if $A$ is a prime number of the form $4n + 3$, or is divisible by such a prime but not by its square; finally, *in general*,

$$\begin{aligned} \text{either} \quad & = 4(\alpha + 1)(\beta + 1)(\gamma + 1) \ldots \\ \text{or} \quad & = 0 \end{aligned}$$

Here the number $A$ is put in the form $2^{\mu} S a^{\alpha} b^{\beta} c^{\gamma} \ldots$, where $a, b, c$, etc., denote different prime numbers of the form $4n + 1$, and $S$ the product of the primes of the form $4n + 3$ that occur once or more among the factors of the number $A$, if there are any. The first or second value applies according as the number $S$ is a square or is not a square. It is therefore evident that $fA$ depends solely on the way in which the prime numbers $3, 5, 7, 11, 13$, etc., occur among the factors of the number $A$, so that in general we must put

$$fA = 4(3) \,.\, (5) \,.\, (7) \,.\, (11) \,.\, (13) \ldots .$$

if we suppose the values of the symbols $(3), (5), (7)$, etc., to be chosen so that, with $p$ denoting a prime number, we have

**first** $(p) = 1$ if $p$ does not divide $A$

**second** $(p) = \alpha + 1$ if $p$ has the form $4n + 1$, and $p^\alpha$ is the highest power dividing $A$

**third** $(p) = 0$ if $p$ has the form $4n+3$, and the exponent of the highest power of $p$ dividing $A$ is odd; finally,

**fourth** $(p) = 1$ if $p$ has the form $4n + 3$, and the exponent of the highest power of $p$ dividing $A$ is even.

Evidently the first case is included in the second and fourth.

The terms of the sequence $f1, f2, f3, f4$, etc., thus proceed very irregularly, although the more terms are taken, the more accurately the mean value $= \pi$ must emerge from them. We shall denote the sum $f1 + f2 + f3 + \ldots + fA$ by $FA$.

**7.**

Let us now put, in general, $fm + f3m = f'm$; it is easily seen that

$$f'A = 4(5) \,.\, (7) \,.\, (11) \,.\, (13) \ldots$$

that is, $f'A$ will be independent of the relation of $A$ to the divisor 3, so that the irregularity of the sequence $f'1, f'2, f'3, f'4, f'5, f'6$, etc., will both begin later and be much smaller. Further, if we put

$$f'1 + f'2 + f'3 + f'4 + \text{etc.} + f'm = F'm$$

we shall have

$$\begin{aligned} F'3A &= F3A + f3 + f6 + f9 + \ldots + f9A \\ &= F3A + FA \end{aligned}$$

From this it follows readily that, as $A$ increases to infinity, we must put

$$F'3A = 4\pi A$$

or that the mean value of the terms of the sequence $f'1, f'2, f'3, f'4$, etc., is

$$= \tfrac{4}{3}\pi$$

In the same way, on putting in general $-f'm + f'5m = f''m$, we obtain

$$f''A = 4(7)(11)(13) \ldots$$

that is, the new sequence $f''1$, $f''2$, etc., loses the fluctuations that depend on the relation to the number 5. On setting the sum

$$f''1 + f''2 + f''3 + \ldots + f''m = F''m$$

we obtain

$$F''5m = -F'm + F'5m$$

from which it follows that, as $m$ increases to infinity, we must put

$$F''5m = \tfrac{4}{3}\pi \,.\, 4m$$

or that the mean value of the terms of the sequence is $= \frac{4}{3} \,.\, \frac{4}{5}\pi$.

If we proceed further in the same way, forming new sequences as we successively remove the factors $(7), (11), (13), (17)$, etc., these will approach constancy ever more closely, and their mean values will successively acquire the new factors $\frac{8}{7}, \frac{12}{11}, \frac{12}{13}, \frac{16}{17}, \frac{20}{19}$, etc. The denominators will be the prime numbers in their natural order, and the numerators will be greater or smaller by one according as those primes have the form $4n - 1$ or $\cdot 4n + 1$. Hence, since, on continuing this process to infinity, the constant value 4 must approach the mean value ever more closely, we have

$$4 = \pi \,.\, \tfrac{4}{3} \cdot \tfrac{4}{5} \cdot \tfrac{8}{7} \cdot \tfrac{12}{11} \cdot \tfrac{12}{13} \ldots \text{ to infinity}$$

or

$$\pi = 4 \,.\, \frac{3}{3+1} \cdot \frac{5}{5-1} \cdot \frac{7}{7+1} \cdot \frac{11}{11+1} \cdot \frac{13}{13-1} \cdots$$

If the individual fractions are expanded into infinite series,

$$\begin{aligned}
\frac{3}{3+1} &= 1 - \frac{1}{3} + \frac{1}{9} - \frac{1}{27} + \ldots \\
\frac{5}{5-1} &= 1 + \frac{1}{5} + \frac{1}{25} + \frac{1}{125} + \ldots \\
\frac{7}{7+1} &= 1 - \frac{1}{7} + \frac{1}{49} - \frac{1}{343} + \ldots \\
&\text{etc.}
\end{aligned}$$

the product readily expands into

$$1 - \tfrac{1}{3} + \tfrac{1}{5} - \tfrac{1}{7} + \tfrac{1}{9} - \tfrac{1}{11} + \tfrac{1}{13} \ldots .$$

and the sum of this series is generally known to be $= \frac{1}{4}\pi$. In fact, the illustrious Euler had long ago derived from this, in the reverse direction, the equality between $\frac{1}{4}\pi$ and the infinite product $\frac{4}{3} \,.\, \frac{4}{5} \,.\, \frac{8}{7} \,..$ (*Introd. in analys. inf.*, Vol. I, Chap. xv, Article 285).

**8.**

Let us consider, secondly, the form $xx + 2yy$, for which the number of representations of an indeterminate number will have the mean value $= \frac{\pi}{\sqrt{2}}$. If $fA$ denotes the number of representations of a given number $A$ by this form, this will be $= 2$ for $A = 1$ or $A = 2$, or whenever $A$ is a power of two; further, $fA = 4$ whenever $A$ is one of the primes of which $-2$ is a quadratic residue, that is, those of the forms $8k + 1, 8k + 3$, for example $A = 3, 11, 17, 19, 41, 43$, etc.; finally, $fA = 0$ whenever $A$ is a prime of which $-2$ is a quadratic nonresidue, for example one of the sequence $5, 7, 13, 23, 29, 31$, etc., that is, of either the form $8k + 5$ or the form $8k + 7$. In general, however, we must put

$$\begin{aligned} \text{either} \quad & fA = 2(\alpha + 1)(\beta + 1)(\gamma + 1) \dots \\ \text{or} \quad & fA = 0 \end{aligned}$$

Here the number $A$ is put in the form $2^\mu S a^\alpha b^\beta c^\gamma \dots$, where $a, b, c$, etc., denote different prime numbers of the forms $8k + 1, 8k + 3$, while $S$ is the product of the remaining numbers (of the forms $8k + 5, 8k + 7$), if any occur among the factors of the number $A$. The first or second value applies according as $S$ is a square or is not a square. From this, by reasoning entirely similar to that in the preceding article, we shall pass from the sequence $f1, f2, f3, f4, f5$, etc., namely $2, 2, 4, 2, 0, 2$, etc., successively to others that remain *constant* for increasingly long stretches. Their *mean* values will be, successively, $\frac{\pi}{\sqrt{2}}$, $\frac{\pi}{\sqrt{2}} \cdot \frac{2}{3}$, $\frac{\pi}{\sqrt{2}} \cdot \frac{2}{3} \cdot \frac{6}{5}$, $\frac{\pi}{\sqrt{2}} \cdot \frac{2}{3} \cdot \frac{6}{5} \cdot \frac{8}{7}$, etc.; we shall proceed in such a way as to arrive at the equation

$$2 = \frac{\pi}{\sqrt{2}} \cdot \frac{2}{3} \cdot \frac{6}{5} \cdot \frac{8}{7} \cdot \frac{10}{11} \cdot \frac{14}{13} \cdot \frac{16}{17} \cdot \frac{18}{19} \dots$$

where the denominators form the natural sequence of prime numbers, while the numerators are smaller by one whenever the denominators have the form $8k + 1$ or $8k + 3$, and larger by one whenever the denominators have the form $8k + 5$ or $8k + 7$.

[II.]

# ON THE CONNECTION BETWEEN THE NUMBER OF CLASSES INTO WHICH BINARY QUADRATIC FORMS ARE DISTRIBUTED AND THEIR DETERMINANT.

## FIRST PAPER

## PRESENTED TO THE ROYAL SOCIETY 1837 . . .

---

**1.**

Thirty-six years have passed since the principles of the remarkable connection to be treated in this paper were discovered, as was already announced at the end of the *Arithmetical Investigations*. Other occupations, however, had long drawn attention away from this inquiry, until more recently it became possible to return to it and extend it through renewed efforts. Yet, since this new part of higher arithmetic exceeds the bounds of a single paper, this first paper will be devoted to forms with negative determinants; forms with positive determinants, which require an entirely distinct treatment, will remain reserved for another paper.

**2.**

For our purpose we shall need a theorem which is itself arithmetical, but whose nature can be presented more conveniently and clearly through considerations expressed in geometrical form.

Given a figure in an unbounded plane, enclosed by any line, its area can be determined approximately by dividing the plane into squares and counting both those lying wholly inside the figure and those cut by its boundary. Evidently, the resulting area will be smaller or larger than the true area according as the latter squares are omitted or added to the former. If, however, we choose, by any rule, to exclude some and include others of the latter squares lying on the boundary, the error may be positive or negative, but must be smaller than the total area of all the boundary squares. The smaller the squares are taken, the more accurately the area will be determined in this way. We may carry such an approximation indefinitely further, or take the squares so small that the error becomes smaller than any given quantity. Although this may already seem evident in itself, we shall nevertheless not refuse to support it with a rigorous proof.

Two squares can have one common vertex, two, or none; in the first and second cases they will be called adjacent, and in the third disjoint. Evidently, at most four squares can all be adjacent to one another, and thus among any five distinct squares at least two must be disjoint. Now the distance between two points lying in disjoint squares cannot be smaller than the side of the squares, which we shall denote by $a$. Hence, if a point starts anywhere in one square, then passes through a second, third and fourth, and finally reaches a fifth, the length of its path certainly cannot be smaller than $a$. By the same reasoning, if the line continues through further squares, the part between the fifth and ninth, and likewise between the ninth and thirteenth, etc., cannot be smaller than $a$. We therefore readily conclude that any line returning to itself which has touched $n$ distinct squares in all certainly cannot be shorter than $\frac{(n-4)a}{4}$. Conversely, a closed line whose length is $= l$ certainly cannot have touched more than $4 + \frac{4l}{a}$ distinct squares. Since their area $= 4aa + 4al$ can become smaller than any given quantity as $a$ decreases indefinitely, the same holds all the more for the error in the determination of area discussed above.

**3.**

The rule for including or excluding squares lying on the boundary of the figure could be chosen in many different ways. The simplest, however, seems to be to consider only the position of the centre of each square, so that squares whose centres lie inside the figure are included, those whose centres lie outside it are excluded, and it is left to our choice whether centres that happen to lie on the boundary itself are counted with the interior or the exterior. Instead of the centres, we could also take any other points situated in the same relative position in each square.

Thus the matter comes down to imagining equally spaced points distributed along equally spaced straight lines in the plane so as to form squares. By the theorem of the preceding article, we can then assert that the number of points contained in the figure, multiplied by the square of the distance between two nearest points, becomes as nearly equal to the area of the figure as desired, provided that distance is taken sufficiently small; or, in the customary manner of speaking, that this product gives the area if the distance is infinitely small.

**4.**

The curve expressed by the following equation between the rectangular coordinates $p, q$,

$$app + 2bpq + cqq = 1$$

is a conic section, and specifically an ellipse if $a, c$ and $ac - bb$ are positive quantities. The area enclosed by this ellipse is found to be $. = \frac{\pi}{\sqrt{(ac-bb)}}$.

The value of $app + 2bpq + cqq$ is everywhere greater than 1 outside the ellipse, less than 1 inside it, and nowhere negative.

Suppose a .system of points distributed in the plane containing the ellipse so as to form squares whose sides $= \lambda$ are parallel to the coordinate axes. It makes no difference whether the origin of the coordinates, or centre of the ellipse, coincides with one of these points. Let the number of points inside the ellipse, including any on its boundary, be $= m$. By the theorem of the preceding article, $\frac{\pi}{\sqrt{(ac-bb)}}$ will then be the limit of $m\lambda\lambda$, which approaches it as closely as desired as $\lambda$ decreases indefinitely.

If we suppose that the origin of the coordinates coincides with a point of the system, then, on putting $p = \lambda x, q = \lambda y$, evidently $x$ and $y$ will be integers for each point of the system; conversely, every combination of integer values of $x, y$ will correspond to a point of the system. Thus the number $m$ is simply the number of all combinations of integer values of $x, y$ for which $F$ is no greater than $M$, if, for brevity, we denote the function, or quadratic form, $axx + 2bxy + cyy$ by $F$, and the quantity $\frac{1}{\lambda\lambda}$ by $M$. The determinant of this form is $bb - ac$, for which we shall write $-D$. Our theorem can now be stated as follows.

THEOREM I. *The number $m$ of all combinations of integer values of the indeterminates $x, y$ for which the value of a form with negative determinant $-D$ does not exceed the bound $M$ becomes $= \frac{\pi M}{\sqrt{D}}$, approximately, indeed, but with the approximation improving indefinitely as $M$ increases indefinitely. It scarcely needs to be mentioned that indefinite approximation here, and likewise in what follows, is not to be understood as saying that the difference between $\frac{\pi M}{\sqrt{D}}$ and $m$ itself decreases indefinitely. Rather, the ratio between these quantities will approach equality indefinitely; that is, $\frac{\pi M}{m\sqrt{D}} - 1$ will decrease indefinitely.*

**5.**

To carry out the count, we may proceed by computing, for each integer value of $x$ between the bounds $-\sqrt{\frac{cM}{D}}$ and $+\sqrt{\frac{cM}{D}}$, the two values of $y$ satisfying $F = M$; the number of integers lying between them is then obtained directly. Since this number is the same for opposite values of $x$, almost half the work is saved. The count can also be carried out by counting the values of $x$ corresponding to each value of $y$ between the bounds $-\sqrt{\frac{aM}{D}}$ and $+\sqrt{\frac{aM}{D}}$. A suitable combination of the two methods can reduce the work further, but we do not pursue this here at greater length; it will suffice to add a few observations on the simplest case.

Let the form be $F = xx + yy$, so that the curve is a circle, and let $r, r', r'', r''' \ldots r^{(r)}$ denote the integers immediately below

$$\sqrt{M}, \quad \sqrt{(M-1)}, \quad \sqrt{(M-4)}, \quad \sqrt{(M-9)} \ldots \sqrt{(M-rr)}$$

or these quantities themselves, if any of them are integers. The required number will then be

$$\begin{aligned} m &= 2r + 1 + 2(2r' + 1) + 2(2r'' + 1) + 2(2r''' + 1) + \text{etc.} + 2(2r^{(r)} + 1) \\ &= 1 + 4r + 4r' + 4r'' + 4r''' + \text{etc.} + 4r^{(r)} \end{aligned}$$

We obtain the same result more quickly, however, by denoting by $q$ the integer immediately below $\sqrt{\frac{1}{2}M}$ (or this quantity itself, if it is an integer), with the aid of the formula

$$m = 4qq + 1 + 4r + 8(r^{(q+1)} + r^{(q+2)} + r^{(q+3)} + \text{etc.} + r^{(r)})$$

The following numbers were obtained in this way.

| $M$ | $m$ | $M$ | $m$ | $M$ | $m$ |
|---|---|---|---|---|---|
| 100 | 317 | 1000 | 3149 | 10000 | 31417 |
| 200 | 633 | 2000 | 6293 | 20000 | 62845 |
| 300 | 949 | 3000 | 9425 | 30000 | 94237 |
| 400 | 1257 | 4000 | 12581 | 40000 | 125629 |
| 500 | 1581 | 5000 | 15705 | 50000 | 157093 |
| 600 | 1885 | 6000 | 18853 | 60000 | 188453 |
| 700 | 2209 | 7000 | 21993 | 70000 | 219901 |
| 800 | 2521 | 8000 | 25137 | 80000 | 251305 |
| 900 | 2821 | 9000 | 28269 | 90000 | 282697 |
| 1000 | 3149 | 10000 | 31417 | 100000 | 314197 |

**6.**

We give the theorem of Article 4 greater generality in the following way.

THEOREM II. *If we are to collect, not all combinations of integer values of $x, y$ for which $F$ does not exceed the value $M$, but only those taken at intervals, namely those for which $x$ is congruent to a given number $G$ to a given modulus $g$, and $y$ is congruent to a given number $H$ to a given modulus $h$, then the number $m'$ of these combinations will be expressed approximately by $\frac{\pi M}{gh\sqrt{D}}$, with the approximation improving indefinitely as $M$ increases indefinitely.*

Indeed, on putting $x = gx' + G, y = hy' + H$, it is evident that $m'$ is the number of all combinations of integer values of $x', y'$ for which

$$agg(x' + \tfrac{G}{g})^2 + 2bgh(x' + \tfrac{G}{g})(y' + \tfrac{H}{h}) + chh(y' + \tfrac{H}{h})^2$$

does not exceed the value $M$. Thus, if we suppose a system of points distributed in the plane as in Article 4, but so that the point with coordinates $p = \frac{G\lambda}{g}, q = \frac{H\lambda}{h}$, rather than the origin of the coordinates, coincides with a point of the system, then $m'$ will express the number of points lying inside the ellipse whose equation is

$$aggpp + 2bghpq + chhqq = 1$$

with any points on the boundary itself always included. The area of this ellipse $= \frac{\pi}{gh\sqrt{(ac-bb)}} = \frac{\pi}{gh\sqrt{D}}$ will be the limit which the product $m'\lambda\lambda = \frac{m'}{M}$ approaches indefinitely as $\lambda$ decreases or $M$ increases indefinitely.

It is also evident that our theorem includes the case where only one of the indeterminates $x, y$ must proceed by intervals, while the value of the other is subject to no condition. Clearly, this is the same as putting either $h$ or $g = 1$.

**7.**

What has been set out so far is independent of the nature of the coefficients of the form $axx + 2bxy + cyy$; from now on, however, we shall suppose these coefficients to be integers. Thus every combination of integer values of $x, y$ gives the form itself an integer value, or corresponds to a representation of some integer by this form. Hence the collection of all combinations of integer values of $x, y$ for which the form $F = axx + 2bxy + cyy$ takes a value no greater than the bound $M$ is the same as the collection of all representations of integers not exceeding the bound $M$, or up to and including this bound if it is itself an integer. If, for brevity, we denote the number of distinct representations of a specified integer $n$ by the form $F$ by $F(n)$, or simply by $Fn$ when there is no risk of ambiguity, then the number previously expressed by $m$ will be $= F0 + F1 + F2 + F3 + \text{etc.} + FM$, and the first theorem takes the following form.

THEOREM III. *The sum $F0 + F1 + F2 + etc. + FM$ is expressed approximately by $\frac{\pi M}{\sqrt{D}}$, with the approximation improving indefinitely as $M$ increases indefinitely.*

**8.**

To the third theorem, which concerns representations of *all* numbers, it is useful to add another concerning odd numbers alone. Evidently, odd numbers cannot be represented by the form $F$ if $a$ and $c$ are both even numbers; the investigation will therefore be restricted to the three remaining cases.

I. Whenever $a$ is odd and $c$ is even, an odd number is represented by giving $x$ an odd value, while the value of $y$ remains arbitrary. Theorem II, on putting $g = 2, G = 1, h = 1$, therefore shows that the number of all combinations of such values of $x, y$ as give the form an odd value no greater than the bound $M$ is expressed, with indefinite approximation, by $\frac{\pi M}{2\sqrt{D}}$, as $M$ increases indefinitely.

II. Whenever $a$ is even and $c$ is odd, the representation of an odd number requires $y$ to be odd. Thus, on putting $g = 1, h = 2, H = 1$, we reach the same conclusion.

III. Whenever both $a$ and $c$ are odd, either an odd value of $x$ must be combined with an even value of $y$, or an even value of $x$ with an odd value of $y$, for the formula to produce an odd value. The number of all combinations of each kind for which the value of the form does not exceed the bound $M$ is expressed, with indefinite approximation, by $\frac{\pi M}{4\sqrt{D}}$. Therefore the number of all combinations which give the form odd values not exceeding the bound $M$ is again expressed, with indefinite approximation, by $\frac{\pi M}{2\sqrt{D}}$.

Now the collection of all such combinations is simply the collection of all representations of all the numbers $1, 3, 5, 7 \ldots M$ when $M$ is an odd integer, or $1, 3, 5, 7 \ldots . M - 1$ when $M$ is even. We therefore have

THEOREM IV. *The sum*

$$F1 + F3 + F5 + F7 . . + FM \quad \text{or} \quad F1 + F3 + F5 + F7 . . + F(M-1)$$

*(according as $M$ is odd or even) is expressed, with indefinite approximation, by $\frac{\pi M}{2\sqrt{D}}$, provided that $F$ is a form in which one or both of the coefficients $a, c$ are odd.*

[III.]

Let $C$ be the collection of representatives of all classes of properly primitive forms for the determinant $-D$. We denote by $(n)$ the number of all representations of the number $n$ by forms in the collection $C$. Let $p$ be an odd prime. Then

$$\left.\begin{array}{ll} 1. & (pn) = (n) \quad \text{if } p \text{ divides } D \\ 2. & (pn) = (n) + (h) \\ 3. & (pn) = -(n) + (h) \end{array}\right\} \text{ if } p \text{ does not divide } D$$

$$\begin{cases} \text{divides } xx + D \\ \text{does not divide } xx + D \end{cases}$$

where $n = hp^\mu$, $\mu$ is arbitrary, and $h$ is not divisible by $p$.

In case 1. we have $(h) = (ph) = (pph) = (p^3h)$ etc.

2. $(ph) = 2(h),\ (pph) = 3(h),\ (p^3h) = 4(h)$ etc.

3. $(ph) = 0,\ (pph) = (h),\ (p^3h) = 0,\ (p^4h) = (h)$ etc.

* *
*

From each properly primitive class for the determinant $= -D$, the number of these classes being $= \lambda$, choose one form, and let $L$ be the collection of these forms.

Let $fA$ denote the number of all representations of the number $A$ by forms in $L$.

Further, let $f(A;p) = f\frac{A}{p^\alpha}$ when $p^\alpha$ is the highest power of the prime $p$ that divides $A$; also $f(A;p,q) = f\frac{A}{p^\alpha q^\beta}$ when $q$ is another prime whose highest power dividing $A$ is $= q^\beta$; and similarly $f(A;p,q,r) = f\frac{A}{p^\alpha q^\beta r^\gamma}$ when $r$ is a third prime whose highest power dividing $A$ is $r^\gamma$, and so on.

---

[IV.]

Let $(n)$ denote the number of values $x$ in the collection

$$0, 1, 2, 3, 4 \ldots . p^n - 1$$

for which $xx - D = xx - ap^\mu$ is divisible by $p^n$.

| 1) $\mu$ odd, e.g. $= 7$. | 2) $\mu$ even, e.g. $= 6$ | |
|---|---|---|
| | $aNp$ | $aRp$ |
| $(1) = 1$ | $(1) = 1$ | $(1) = 1$ |
| $(2) = p$ | $(2) = p$ | $(2) = p$ |
| $(3) = p$ | $(3) = p$ | $(3) = p$ |
| $(4) = pp$ | $(4) = pp$ | $(4) = pp$ |
| $(5) = pp$ | | |

| | | |
|---|---|---|
| $(6) = p^3$ | $(5) = pp$ | $(5) = pp$ |
| $(7) = p^3$ | $(6) = p^3$ | $(6) = p^3$ |
| $(8) = 0$ | $(7) = 0$ | $(7) = 2p^3$ |
| $(9) = 0$ | $(8) = 0$ | $(8) = 2p^3$ |
| etc. | etc. | etc. |

| Now put | Then |
|---|---|
| $(1) - \frac{(2)}{p} = (1)'$ | $fp = (1)'$ |
| $(2) - \frac{(3)}{p} = (2)'$ | $fpp = 1 + (2)'$ |
| $(3) - \frac{(4)}{p} = (3)'$ | $fp^3 = (1)' + (3)'$ |
| $(4) - \frac{(5)}{p} = (4)'$ | $fp^4 = 1 + (2)' + (4)'$ |
| etc. | etc. |

Consequently, putting $\frac{p-1}{p}(1 + \frac{fp}{p} + \frac{fpp}{pp} + \frac{fp^3}{p^3} + \text{etc.}) = T$, we have

$$\frac{p+1}{p}T = 1 + \frac{(1)'}{p} + \frac{(2)'}{pp} + \frac{(3)'}{p^3} + \frac{(4)'}{p^4} + \text{etc.}$$
$$= 1 + \frac{(1)}{p} = 1 + \frac{1}{p}$$

Thus $T = 1$

---

[V.]

The average number of classes[*)] near the negative determinant $-D$ is approximately

$$= \frac{\pi\sqrt{D}}{4(1 + \frac{1}{27} + \frac{1}{125} + \text{etc.})}$$

The actual number is expressed by the following formulas, where, for brevity, $m$ stands for the average number and $M$ for the actual number; $p, q$ denote all odd primes that do not divide $D$, the former being divisors and the latter nondivisors of $\square + D$; $r$ denotes the primes[**)] that divide $D$:

[*)] [Compare *Disquiss. Arithm.*, Article 302; the formula there differs by a constant $\delta$ from the one occurring here in the text.]

[**)] [odd.]

$$\text{I.} \quad M = m \text{ Product of } \frac{p^3+p^2}{p^3-1} \cdot \frac{q^3-q^2}{q^3-1} \cdot \frac{r^3-r}{r^3-1}$$

$$\text{II.} \quad M = \frac{\pi\sqrt{D}}{4} \text{ Product of } \frac{p+1}{p} \cdot \frac{q-1}{q} \cdot \frac{rr-1}{rr}$$

$$\text{III. NB.} \quad M = \frac{2\sqrt{D}}{\pi} \text{ Product of } \frac{p}{p-1} \cdot \frac{q}{q+1}$$

$$\text{IV.} \quad M = \sqrt{\left\{ \frac{D}{2} \cdot \text{Product of } \frac{p+1}{p-1} \cdot \frac{q-1}{q+1} \cdot \frac{rr-1}{rr} \right\}}$$

$$\text{V.} \quad M = \frac{2\sqrt{D}}{\pi}\{1 \pm \tfrac{1}{3} \pm \tfrac{1}{5} \pm \tfrac{1}{7} \pm \tfrac{1}{9} \text{ etc.}\}$$

NB. Formula III follows directly by comparing the two ways of counting the representable numbers up to a given bound.

---

[VI.]

THEOREM. The number of classes, into which all properly primitive binary forms with negative determinant $-D$[*], is equal to

$$\frac{\pi}{4} \times \sqrt{D} \times \text{Product of } \frac{p-1}{p}\frac{q+1}{q} \times \frac{rr-1}{rr}$$

where

$p$ denotes all primes[**] for which $-D$ is a nonresidue,

$q$ all primes[**] for which $-D$ is a residue,

$r$ all primes[**] that divide $D$;

$$= \frac{\frac{\pi}{4}\sqrt{D} \text{ Product of } \frac{rr-1}{rr}}{1 \pm \frac{1}{3} \pm \frac{1}{5} \text{ etc.}}$$

in the denominator, the positive sign is put before fractions whose denominators have the form of nondivisors, and the negative sign before those whose denominators have the form of divisors of $xx + D$; fractions whose denominators are not coprime to $D$ are omitted altogether[***].

[*] [are divided.]

[**] [odd.]

[***] [If $m$ denotes all positive integers coprime to $2D$, and we use LEGENDRE's symbol as generalized by JACOBI, the above rule for determining the signs must be corrected as follows: in the preceding formula the denominator is

$$1 \pm \tfrac{1}{3} \pm \tfrac{1}{5} \text{ etc.} = \Sigma \pm \left(\frac{-D}{m}\right)\frac{1}{m}$$

$$= \frac{2\sqrt{D}(1 \pm \frac{1}{3} \pm \frac{1}{5} \ldots)}{\pi}$$
$$= \frac{\operatorname{cotg}\theta \pm \operatorname{cotg} 3\theta \pm \operatorname{cotg} 5\theta \ldots \pm \operatorname{cotg} n\theta}{N : \sqrt{D}}$$

putting $\theta = \frac{\pi}{N}$, $N = \{\frac{1}{4}\} D$, and taking for $n$ all numbers coprime to $D$, with the sign determined as above*).

For a positive determinant the number of classes will be**)

$$= \frac{2\sqrt{D}(1 \pm \frac{1}{3} \pm \frac{1}{5} \ldots)}{\log .T + U\sqrt{D}}$$

Here $T, U$ denote the smallest values of the quantities $t, u$ that satisfy the equation $tt - Duu = 1$

$$= \frac{\log \sin \frac{1}{2}\theta \pm \log \sin \frac{3}{2}\theta \pm \log \sin \frac{5}{2}\theta \text{ etc.}}{\log T + U\sqrt{D}}$$

---

[VII.]

For a negative determinant $-p$, which***) is a prime of the form $4n+1$, the number of classes is†) $= (\alpha - \beta)$, where $\alpha$ is the number of quadratic residues in the first quarter

$$1 \,.\, 2 \,.\, 3 \ldots . \tfrac{1}{4}(p-1)$$

and $\beta$ the number of nonresidues.

---

where the upper or lower sign is to be taken according as the number $m$ is a product of an even or odd number of primes (equal or unequal); in contrast, in the numerator of the following formula we have

$$1 \pm \tfrac{1}{3} \pm \tfrac{1}{5} \ldots = \Sigma \left(\frac{-D}{m}\right) \frac{1}{m}$$

]

*)[See the note on this fragment further below.]

**)[In the following formula $D$ denotes the positive determinant, and

$$1 \pm \tfrac{1}{3} \pm \tfrac{1}{5} \ldots = \Sigma \left(\frac{D}{m}\right) \frac{1}{m}$$

]

***)[that is, when $p$ is a positive prime of the form $4n + 1$.]

†)[the number of classes is $= 2(\alpha - \beta)$.]

[VIII.]

$$b \equiv 2m + a - 1 \pmod 8$$

where $m$ is [half] the number of classes for the determinant $-p$

| $p$ | $m$ | $a$ | $b$ | $f$ | $\frac{2m+a-1-b}{8}$ | $\alpha$ | $\beta$ |
|---|---|---|---|---|---|---|---|
| 17 | 2 | +1 | −4 | −4 | +1 | 3 | 2 |
| 41 | 4 | +5 | +4 | +9 | +1 | 3 | 4 |
| 73 | 2 | −3 | −8 | +27 | +1 | 1 | 6 |
| 89 | 6 | +5 | −8 | +34 | +3 | 9 | 2 |
| 97 | 2 | +9 | +4 | +22 | +1 | 5 | 6 |
| 113 | 4 | −7 | +8 | +15 | −1 | 9 | 4 |
| 137 | 4 | −11 | +4 | +37 | −1 | 3 | 8 |
| 193 | 2 | −7 | +12 | +81 | −2 | 11 | 6 |
| 233 | 6 | +13 | +8 | +144 | +2 | 15 | 2 |
| 241 | 6 | +15 | +4 | +64 | −1 | 13 | 6 |
| 257 | 8 | +1 | +16 | +16 | 0 | 15 | 4 |
| 281 | 10 | +5 | −16 | +53 | +5 | 9 | 10 |
| 313 | 4 | +13 | −12 | −25 | +1 | 5 | 12 |
| 337 | 4 | +9 | +16 | −148 | 0 | 7 | 12 |
| 353 | 8 | +17 | +8 | +42 | +3 | 15 | 8 |
| 5 | .1 | +1 | +2 | +2 | 0 | | |
| 13 | 1 | −3 | −2 | +5 | 0 | | |
| 29 | 3 | +5 | +2 | +12 | +1 | | |
| 37 | 1 | +1 | −6 | −6 | +1 | | |
| 53 | 3 | −7 | −2 | +23 | 0 | | |
| 61 | 3 | +5 | −6 | +11 | +2 | | |
| 101 | 7 | +1 | −10 | −10 | +3 | | |
| 109 | 3 | −3 | +10 | +33 | −1 | | |
| 149 | 7 | −7 | −10 | +44 | +2 | | |
| 157 | 3 | −11 | −6 | −28 | 0 | | |
| 173 | 7 | +13 | +2 | +80 | +3 | | |
| 181 | 5 | +9 | +10 | −19 | +1 | | |
| 197 | 5 | +1 | −14 | −14 | +3 | | |
| 229 | 5 | −15 | +2 | −107 | −1 | | |
| 269 | 11 | +13 | +10 | −82 | +3 | | |
| 277 | 3 | +9 | +14 | −60 | 0 | | |
| 293 | 9 | +17 | +2 | +138 | +4 | | |
| 317 | 5 | −11 | +14 | +114 | −2 | | |
| 349 | 7 | +5 | +18 | −136 | 0 | | |
| 373 | 5 | −7 | +18 | +104 | −2 | | |
| 389 | 11 | +17 | −10 | −115 | +6 | | |
| 397 | 3 | −19 | −6 | +63 | −1 | | |

[IX.]

*Distribution of Quadratic Residues in Eighths.*

$p$ a prime; $(r)$ the number of quadratic residues of $p$ that lie between $(r-1)\frac{p}{8}$ and $r\frac{p}{8}$.

First case; $p = 8n + 1$.

$2t$ number of classes for the determinant $-p$;

$2u$ number of classes for the determinant $-2p$.

$$(1) = (8) = \tfrac{1}{4}(2n + t + u)$$
$$(2) = (4) = (5) = (7) = \tfrac{1}{4}(2n + t - u)$$
$$(3) = (6) = \tfrac{1}{4}(2n - 3t + u)$$

| $p$ | $2n$ | $t$ | $u$ | (1) | (2) | (3) |
|---|---|---|---|---|---|---|
| 17 | 4 | 2 | 2 | 2 | 1 | 0 |
| 41 | 10 | 4 | 2 | 4 | 3 | 0 |
| 73 | 18 | 2 | 8 | 7 | 3 | 5 |
| 89 | 22 | 6 | 4 | 8 | 6 | 2 |
| 97 | 24 | 2 | 10 | 9 | 4 | 7 |
| 113 | 28 | 4 | 4 | 9 | 7 | 5 |
| 137 | 34 | 4 | 6 | 11 | 8 | 7 |
| 193 | 48 | 2 | 10 | 15 | 10 | 13 |
| 233 | 58 | 6 | 4 | 17 | 15 | 11 |
| 241 | 60 | 6 | 10 | 19 | 14 | 13 |
| 257 | 64 | 8 | 8 | 20 | 16 | 12 |
| 281 | 70 | 10 | 4 | 21 | 19 | 11 |
| 313 | 78 | 4 | 18 | 25 | 16 | 21 |
| 337 | 84 | 4 | 12 | 25 | 19 | 21 |
| 353 | 88 | 8 | 12 | 27 | 21 | 19 |
| 401 | 100 | 10 | 6 | 29 | 26 | 19 |

Second case; $p = 8n + 5$.

$2t$ number of classes for the determinant $-p$;
$2u$ number of classes for the determinant $-2p$.

$$(1) = (3) = (6) = (8) = \tfrac{1}{4}(2n - t + u)$$
$$(2) = (7) = \tfrac{1}{4}(2n + 3t - u + 2)$$
$$(4) = (5) = \tfrac{1}{4}(2n - t - u + 2)$$

| $p$ | $2n$ | $t$ | $u$ | (1) | (2) | (4) |
|---|---|---|---|---|---|---|
| 5 | 0 | 1 | 1 | 0 | 1 | 0 |
| 13 | 2 | 1 | 3 | 1 | 1 | 0 |
| 29 | 6 | 3 | 1 | 1 | 4 | 1 |
| 37 | 8 | 1 | 5 | 3 | 2 | 1 |
| 53 | 12 | 3 | 3 | 3 | 5 | 2 |
| 61 | 14 | 3 | 5 | 4 | 5 | 2 |
| 101 | 24 | 7 | 3 | 5 | 11 | 4 |
| 109 | 26 | 3 | 5 | 7 | 8 | 5 |
| 149 | 36 | 7 | 3 | 8 | 14 | 7 |
| 157 | 38 | 3 | 13 | 12 | 9 | 6 |
| 173 | 42 | 7 | 5 | 10 | 15 | 8 |
| 181 | 44 | 5 | 9 | 12 | 13 | 8 |
| 197 | 58 | 5 | 5 | 12 | 15 | 10 |
| 229 | 56 | 5 | 13 | 16 | 15 | 10 |
| 269 | 66 | 11 | 5 | 15 | 24 | 13 |
| 277 | 68 | 3 | 11 | 19 | 17 | 14 |
| 293 | 72 | 9 | 9 | 18 | 23 | 14 |
| 317 | 78 | 5 | 7 | 20 | 22 | 17 |
| 349 | 86 | 7 | 13 | 23 | 24 | 17 |
| 373 | 92 | 5 | 13 | 25 | 24 | 19 |
| 389 | 96 | 11 | 7 | 23 | 31 | 20 |
| 397 | 98 | 3 | 21 | 29 | 22 | 19 |

Third case; $p = 8n + 3$.

$t$ number of classes for the determinant $-p$

$2u$ number of classes for the determinant $-2p$.

$$(1) = (4) = (7) = \tfrac{1}{4}(2n + t - u)$$
$$(2) = (5) = (8) = \tfrac{1}{4}(2n - t + u)$$
$$(3) = \tfrac{1}{4}(2n + t + u + 2)$$
$$(6) = \tfrac{1}{4}(2n - t - u + 2)$$

| $p$ | $2n$ | $t$ | $u$ | (1) | (2) | (3) | (6) |
|---|---|---|---|---|---|---|---|
| 3 | 0 | 1 | 1 | 0 | 0 | 1 | 0 |
| 11 | 2 | 3 | 1 | 1 | 0 | 2 | 0 |
| 19 | 4 | 3 | 3 | 1 | 1 | 3 | 0 |
| 43 | 10 | 3 | 5 | 2 | 3 | 5 | 1 |
| 59 | 14 | 9 | 3 | 5 | 2 | 7 | 1 |
| 67 | 16 | 3 | 7 | 3 | 5 | 7 | 2 |
| 83 | 20 | 9 | 5 | 6 | 4 | 9 | 2 |
| 107 | 26 | 9 | 3 | 8 | 5 | 10 | 4 |
| 131 | 32 | 15 | 3 | 11 | 5 | 13 | 4 |
| 139 | 34 | 9 | 7 | 9 | 8 | 13 | 5 |
| 163 | 40 | 3 | 11 | 8 | 12 | 14 | 7 |
| 179 | 44 | 15 | 3 | 14 | 8 | 16 | 7 |
| 211 | 52 | 9 | 5 | 14 | 12 | 17 | 10 |
| 227 | 56 | 15 | 7 | 16 | 12 | 20 | 9 |
| 251 | 62 | 21 | 7 | 19 | 12 | 23 | 9 |
| 283 | 70 | 9 | 15 | 16 | 19 | 24 | 12 |
| 307 | 76 | 9 | 17 | 17 | 21 | 26 | 13 |
| 331 | 82 | 9 | 11 | 20 | 21 | 26 | 16 |
| 347 | 86 | 15 | 5 | 24 | 19 | 27 | 17 |
| 379 | 94 | 9 | 11 | 23 | 24 | 29 | 19 |

Fourth case; $p = 8n + 7$.

$t$ number of classes for the determinant $-p$;
$2u$ number of classes for the determinant $-2p$.

$$(1) = \tfrac{1}{4}(2n + 2t - u)$$
$$(2) = (3) = (5) = \tfrac{1}{4}(2n + u + 2)$$
$$(4) = (6) = (7) = \tfrac{1}{4}(2n - u + 2)$$
$$(8) = \tfrac{1}{4}(2n - 2t + u)$$

| $p$ | $2n$ | $t$ | $u$ | (1) | (2) | (4) | (8) |
|---|---|---|---|---|---|---|---|
| 7 | 0 | 1 | 2 | 0 | 1 | 0 | 0 |
| 23 | 4 | 3 | 2 | 2 | 2 | 1 | 0 |
| 31 | 6 | 3 | 4 | 2 | 3 | 1 | 1 |
| 47 | 10 | 5 | 4 | 4 | 4 | 2 | 1 |
| 71 | 16 | 7 | 2 | 7 | 5 | 4 | 1 |
| 79 | 18 | 5 | 4 | 6 | 6 | 4 | 3 |
| 103 | 24 | 5 | 10 | 6 | 9 | 4 | 6 |
| 127 | 30 | 5 | 8 | 8 | 10 | 6 | 7 |
| 151 | 36 | 7 | 6 | 11 | 11 | 8 | 7 |
| 167 | 40 | 11 | 6 | 14 | 12 | 9 | 6 |
| 191 | 46 | 13 | 4 | 17 | 13 | 11 | 6 |
| 199 | 48 | 9 | 10 | 14 | 15 | 10 | 10 |
| 223 | 54 | 7 | 16 | 13 | 18 | 10 | 14 |
| 239 | 58 | 15 | 4 | 21 | 16 | 14 | 8 |
| 263 | 64 | 13 | 6 | 21 | 18 | 15 | 11 |
| 271 | 66 | 11 | 12 | 19 | 20 | 14 | 14 |
| 311 | 76 | 19 | 6 | 27 | 21 | 18 | 11 |
| 359 | 88 | 19 | 6 | 30 | 24 | 21 | 14 |
| 367 | 90 | 9 | 20 | 22 | 28 | 18 | 23 |
| 383 | 94 | 17 | 12 | 29 | 27 | 21 | 18 |

[X.]

*Distribution of Quadratic Residues in Twelfths.*

$p$ a prime; $(r)$ the number of quadratic residues of $p$ that lie between $\frac{r-1}{12}p$ and $\frac{r}{12}p$.

First case; $p = 24n + 1$.

$2t$ number of classes for the determinant $-p$

$4u$ number of classes for the determinant $-3p$

$$(1) = (12) = \tfrac{1}{6}(6n + 3t + 2u)$$
$$(2) = (4) = (6) = (7) = (9) = (11) = \tfrac{1}{6}(6n - 3t + 2u)$$
$$(3) = (5) = (8) = (10) = \tfrac{1}{6}(6n + 3t - 4u)$$

| $p$ | $n$ | $t$ | $u$ | (1) | (2) | (3) |
|---|---|---|---|---|---|---|
| 73 | 3 | 2 | 3 | 5 | 3 | 2 |
| 97 | 4 | 2 | 3 | 6 | 4 | 3 |
| 193 | 8 | 2 | 6 | 11 | 9 | 5 |
| 241 | 10 | 6 | 3 | 14 | 8 | 11 |

Second case; $p = 24n + 13$.

$2t$ number of classes for the determinant $-p$;

$4u$ number of classes for the determinant $-3p$.

$$(1) = (3) = (10) = (12) = \tfrac{1}{2}(2n + 1 + t)$$
$$(2) = (6) = (7) = (1.1) = \tfrac{1}{2}(2n + 1 - t)$$
$$(4) = (9) = \tfrac{1}{2}(2n + 1 - t + 2u)$$
$$(5) = (8) = \tfrac{1}{2}(2n + 1 + t - 2u)$$

| $p$ | $n$ | $t$ | $u$ | (1) | (2) | (4) | (5) |
|---|---|---|---|---|---|---|---|
| 13 | 0 | 1 | 1 | 1 | 0 | 1 | 0 |
| 37 | 1 | 1 | 2 | 2 | 1 | 3 | 0 |
| 61 | 2 | 3 | 2 | 4 | 1 | 3 | 2 |
| 109 | 4 | 3 | 3 | 6 | 3 | 6 | 3 |
| 157 | 6 | 3 | 4 | 8 | 5 | 9 | 4 |
| 181 | 7 | 5 | 3 | 10 | 5 | 8 | 7 |
| 229 | 9 | 5 | 3 | 12 | 7 | 10 | 9 |

Third case; $p = 24n + 5$.

$2t$ number of classes for the determinant $-p$;
$2u$ number of classes for the determinant $-3p$.

$$(1) = (2) = (6) = (7) = (11) = (12) = n$$
$$(3) = (10) = \tfrac{1}{2}(2n + 1 + t)$$
$$(4) = (9) = \tfrac{1}{2}(2n - t + u)$$
$$(5) = (8) = \tfrac{1}{2}(2n + 1 - u)$$

| $p$ | $n$ | $t$ | $u$ | (1) | (3) | (4) | (5) |
|---|---|---|---|---|---|---|---|
| 5 | 0 | 1 | 1 | 0 | 1 | 0 | 0 |
| 29 | 1 | 3 | 3 | 1 | 3 | 1 | 0 |
| 53 | 2 | 3 | 5 | 2 | 4 | 3 | 0 |
| 101 | 4 | 7 | 5 | 4 | 8 | 3 | 2 |
| 149 | 6 | 7 | 7 | 6 | 10 | 6 | 3 |
| 173 | 7 | 7 | 9 | 7 | 11 | 8 | 3 |
| 197 | 8 | 5 | 11 | 8 | 11 | 11 | 3 |
| 269 | 11 | 11 | 7 | 11 | 17 | 9 | 8 |

Fourth case; $p = 24n + 17$.

$2t$ number of classes for the determinant $-p$;
$2u$ number of classes for the determinant $-3p$.

$$(1) = (2) = (6) = (7) = (11) = (12) = \tfrac{1}{6}(6n + 3 + u)$$
$$(3) = (10) = \tfrac{1}{6}(6n + 6 + 3t - 2u)$$
$$(4) = (9) = \tfrac{1}{6}(6n + 3 - 3t + u)$$
$$(5) = (8) = \tfrac{1}{6}(6n + 6 - 2u)$$

| $p$ | $n$ | $t$ | $u$ | (1) | (3) | (4) | (5) |
|---|---|---|---|---|---|---|---|
| 17 | 0 | 2 | 3 | 1 | 1 | 0 | 0 |
| 41 | 1 | 4 | 3 | 2 | 3 | 0 | 1 |
| 89 | 3 | 6 | 3 | 4 | 6 | 1 | 3 |
| 113 | 4 | 4 | 9 | 6 | 4 | 4 | 2 |
| 137 | 5 | 4 | 9 | 7 | 5 | 5 | 3 |
| 233 | 9 | 6 | 15 | 12 | 8 | 9 | 5 |
| 257 | 10 | 8 | 9 | 12 | 12 | 8 | 8 |

## Remarks on the Paper

## On the Connection between the Number of Classes into Which Binary Quadratic Forms Are Distributed and Their Determinant.

---

### On I and II.

The second formula for the number of points inside the circle (I, Article 3, and II, Article 5) follows by considering the square inscribed in the circle, whose sides are parallel to the coordinate axes; comparing the two formulas leads to the theorem

$$r' + r'' + \ldots + r^{(q)} = qq + r^{(q+1)} + r^{(q+2)} + \ldots + r^{(r)}$$

which is also easy to prove arithmetically. This in turn gives the correctness of the first of the following two rules, which were found on a separate sheet:

"Solutions of the equation $xx + yy \leqq A$; formula

$$1 + 4\sqrt{A} + 4\sqrt{\tfrac{1}{2}A} + 8\Sigma(\sqrt{(A - nn)} - n)$$

where the fractional part of each square root is dropped, and the sum is taken from $n = 1$ to $n = \sqrt{A}$ '(should read $\sqrt{\frac{1}{2}A}$)'.

Another formula

$$1 + 4\{A - \tfrac{A}{3} + \tfrac{A}{5} - \tfrac{A}{7} + \tfrac{A}{9} - \tfrac{A}{11}\ldots\}$$

where the fractional part of each term is dropped."

The latter formula follows from the theorem applied later (I, Article 6) concerning the number of all distinct representations of a given number by the form $xx + yy$ (compare Disqq. Arithm., Article 182, note). This theorem can easily be put in the following form: the number of distinct representations of a positive integer $m$ by the form $xx + yy$ is $= 4(a - b)$, where $a, b$ denote the numbers of divisors of $m$ of the respective forms $4n+1$, $4n+3$. By comparing this arithmetical formula with the average number of representations obtained by geometrical considerations (in I, Article 5, or II, Article 4), one easily obtains, with full rigor, the familiar result

$$\frac{\pi}{4} = 1 - \tfrac{1}{3} + \tfrac{1}{5} - \tfrac{1}{7} + \ldots$$

which is derived in the paper (I, Article 7) by a similar comparison, but with the help of infinite products.

---

On III and IV.

Let $C$ be the collection of all positive, properly primitive forms of negative determinant $-D$ that are not properly equivalent. Assign to the variables of each form two values relatively prime to one another. The total number of representations of a positive integer $m$ is then $\epsilon\psi(m)$, where $\epsilon$ denotes the number of solutions of the equation $tt + Duu = 1$, and $\psi(m)$ the number of roots $n$ of the congruence $nn + D \equiv 0 \pmod{m}$ for which the three numbers $m,\ 2n$ and $\frac{nn+D}{m}$ have no common divisor (Disqq. Arithm., art. 180). For the factor, $\epsilon = 4$ when $D = 1$, and $= 2$ in every other case. Further, if $m = p^\pi p'^{\pi'} p''^{\pi''} \ldots$, where $p, p', p'' \ldots$ are distinct primes, then $\psi(m) = \psi(p^\pi)\psi(p'^{\pi'})\psi(p''^{\pi''}) \ldots$; if $\mathfrak{A}(m)$ denotes the number of roots $n$ of the congruence $nn + D \equiv 0 \pmod{m}$, and one uses the symbol introduced by Legendre and generalized by Jacobi, then $\psi(p^\pi) = \mathfrak{A}(p^\pi) = 1 + \left(\frac{-D}{p}\right)$ when $p$ does not divide $2D$, but otherwise $= \mathfrak{A}(p^\pi - \frac{1}{p}\mathfrak{A}(p^{\pi+1})$; the number $\mathfrak{A}(p^\pi)$ is always easy to determine (Disqq. Arithm., art. 104). For what follows, however, it suffices to observe that $\mathfrak{A}(p^\pi)$ always becomes independent of $\pi$ once $\pi$ exceeds a certain size.

If the variables of the forms in the collection $C$ are assigned all integer values without exception (Disqq. Arithm., art. 181), the total number $(m)$ of representations of the number $m$ becomes $\epsilon f(m)$, where $f(m) = \Sigma\psi\left(\frac{m}{\mu\mu}\right)$ and the sum extends over all square divisors $\mu\mu$ of the number $m$. This immediately gives

$$f(m) = f(p^\pi p'^{\pi'} p''^{\pi''} \ldots) = f(p^\pi) f(p'^{\pi'}) f(p''^{\pi''}) \ldots$$

and

$$f(p^\pi) = \psi(p^\pi) + \psi(p^{\pi-2}) + \psi(p^{\pi-4}) + \ldots$$

the series being continued as long as the exponents $\pi,\ \pi - 2,\ \pi - 4 \ldots$ do not become negative. When $p$ does not divide $2D$, this gives

$$f(p^\pi) = 1 + \left(\frac{-D}{p}\right) + \left(\frac{-D}{pp}\right) + \ldots + \left(\frac{-D}{p^\pi}\right)$$

and, in general, when $m$ is relatively prime to $2D$,

$$f(m) = \Sigma\left(\frac{-D}{n}\right)$$

where the sum extends over all divisors $n$ of the number $m$.

These observations immediately establish the statements in the text (III, 1, 2, 3) about the number $(m)$, provided that the first is supplemented by the condition that $D$ must not be divisible by $pp$ (the determination of the number of classes was already reduced in Disqq. Arithm., art. 256, to the case in which $D$ is divisible by no square). At the same time, even without this restriction, the infinite series

$$1 + \frac{f(p)}{p} + \frac{f(pp)}{pp} + \frac{f(p^3)}{p^3} + \dots$$

is found to have the value

$$\frac{1}{1 - \frac{1}{p}} \quad \text{or} \quad \frac{1}{1 - .\frac{1}{p}} \cdot \frac{1}{1 - \left(\frac{-D}{p}\right)\frac{1}{p}}$$

according as $2D$ is or is not divisible by the prime $p$.

---

On V.

The remark appended to formula III indicates how the author arrived at the number $k$ of forms in the collection $C$. Geometrical considerations (compare I, art. 5, and II, art. 4) give the following limit for the quotient

$$\frac{(1) + (2) + (3) + \dots + (m)}{m}$$

as $m$ grows without bound, that is, the average number of representations of an unspecified positive integer,

$$= k\frac{\pi}{\sqrt{D}}$$

A second expression for this same limit can be derived in several ways from the nature of the number $(m) = \epsilon f(m)$ of representations of the number $m$, determined above. The surviving fragments (I, arts. 7, 8; III and IV) suggest that the author's first approach was as follows.

Let $\theta(m)$ be any function of the positive integer $m$, and $p$ any prime. From $\theta(m)$ one can always derive a new function $\theta'(m)$ whose value is independent of whether, and how often, $p$ occurs as a factor in $m$, and which, whenever $p$ does not divide $m$, agrees with $\theta(m)$. Such a function is obtained by putting $\theta'(m) = \theta\left(\frac{m}{p^\pi}\right)$, where $p^\pi$ denotes the highest power dividing $m$ with base $p$; one may say that the function $\theta'(m)$ arises from $\theta(m)$ by elimination of the prime $p$. If in this way from $f(m)$ one forms a new function $f'(m)$ by eliminating the prime 2, then forms $f''(m)$ from it by eliminating 3, and so on, each successive function will have a more regular course than its predecessors.

If the primes are eliminated one after another in increasing order, such a function $\theta(m)$ will, for infinitely many values of $m$, have the value $f(1) = 1$, in particular for all values of $m$ smaller than the last prime eliminated. Continuing this process indefinitely brings one ever closer to the function $f^{\infty}(m)$, which for all values of $m$ has the value 1, and whose average value therefore also $= 1$. If it is possible to express the average value of any function $\theta(m)$ in terms of that of its successor $\theta'(m)$, it will also be possible to find the average value of $f(m)$ through an infinite chain of operations.

If $p$ is the prime whose elimination produces $\theta'(m)$ from $\theta(m)$, then $\theta(m) = \theta'(m)f(p^{\pi})$, where $p^{\pi}$ again denotes the highest power dividing $m$ with base $p$. When $p$ does not divide $2D$, this readily gives

$$\theta'(m) = \theta(mp) - \left(\frac{-D}{p}\right)\theta(m)$$

Writing for brevity

$$\Theta(m) = \theta(1) + \theta(2) + \ldots + \theta(m)$$
$$\Theta'(m) = \theta'(1) + \theta'(2) + \ldots + \theta(m)$$

one obtains

$$\Theta'(mp) = \Theta(mp) - \left(\frac{-D}{p}\right)\Theta(m)$$

and hence, using $\omega, \omega'$ for the respective average values of the functions $\theta(m), \theta'(m)$,

$$\omega = \frac{\omega'}{1 - \left(\frac{-D}{p}\right)\frac{1}{p}}$$

If, however, the prime $p$ divides $2D$, the functions $\theta(m)$ and $\theta'(m)$ no longer have so simple a relation in general; nevertheless, a similar argument readily gives $\omega = \omega'$ in this case. The relation between $\omega$ and $\omega'$ can also be derived in both cases as follows. Put

$$\vartheta(m) = \Sigma\theta(\mu)$$

where the sum extends over all numbers $\mu$ that are not divisible by $p$ and are no greater than $m$. Denote by $m', m'', m''' \ldots$, respectively, the greatest integers contained in $\frac{m}{p}, \frac{m'}{p}, \frac{m''}{p} \ldots$. Then

$$\Theta(m) = \vartheta(m) + \vartheta(m')f(p) + \vartheta(m'')f(pp) + \vartheta(m''')f(p^3) + \ldots$$
$$\Theta'(m) = \vartheta(m) + \vartheta(m') + \vartheta(m'') + \vartheta(m''') + \ldots$$

and this gives

$$\frac{\omega}{\omega'} = (1 - \tfrac{1}{p})\{1 + \frac{f(p)}{p} + \frac{f(pp)}{pp} + \frac{f(p^3)}{p^8} + \ldots\}$$

in agreement with the result found above (compare the note on III and IV).

The average value of the function $f(m)$ is therefore equal to the infinite product

$$\Pi \frac{1}{1 - \left(\frac{-D}{p}\right)\frac{1}{p}}$$

in which $p$ must run through all primes not dividing $2D$, and hence

$$k = \frac{\epsilon\sqrt{D}}{\pi}\Pi \frac{1}{1 - \left(\frac{-D}{p}\right)\frac{1}{p}}$$

There remains, however, a concern about the rigor of this deduction, concerning the method of determining the average value of $f(m)$ by successively eliminating all primes. For it is evident that after elimination of the first $n$ primes, the resulting function $f^{(n)}(m)$ agrees in value with the function $f^{\infty}(m) = 1$ as long as $m$ remains smaller than the last prime eliminated. Thus, by choosing $n$ large enough, this agreement can be extended up to any prescribed size of the number $m$. But this alone by no means proves that, as $n$ grows without bound, the average value of $f^{(n)}(m)$ approaches without limit the average value of $f^{\infty}(m)$, that is, the value 1. How the author intended to fill this gap cannot be determined with certainty from the surviving papers. Yet the formula already given above (in the note on I),

$$1 + 4\{A - \frac{A}{3} + \frac{A}{5} - \frac{A}{7} + \frac{A}{9} - \frac{A}{11}\ldots\}$$

for the number of pairs of numbers whose sum of squares does not exceed $A$, suggests that the author bypassed the infinite product and directly found, for the average value of $f(m)$, the infinite series

$$\Sigma \left(\frac{-D}{n}\right)\frac{1}{n}$$

in which $n$ must run in increasing order through all positive integers relatively prime to $2D$. The simplest way of indicating this transition seems to be the following.

Let $\mu$ be the greatest divisor of a number $m$ among those relatively prime to $2D$, and put $\theta(m) = f(\mu)$. Then $\theta(m)$ is the function obtained by eliminating all primes dividing $2D$ from $f(m)$; according to the preceding discussion, its average value agrees with that of $f(m)$. Since $\theta(m) = \Sigma\left(\frac{-D}{n}\right)$, where $n$ runs through all divisors of $\mu$, that is, all divisors of $m$ relatively prime to $2D$, one obtains the formula analogous to the one above,

$$\Theta(m) = \theta(1) + \theta(2) + \ldots + \theta(m) = \Sigma\left(\frac{-D}{n}\right)\frac{m}{n}$$

where, in the sum on the right, the letter $n$ runs through all numbers relatively prime to $2D$, and only the greatest integer contained in the quotient $\frac{m}{n}$ is to be retained. Arrange the terms of this series so that the numbers $n$ follow in increasing order. The factor $\frac{m}{n}$ then continually decreases, or at least never increases, and the series terminates as soon as $n > m$. Furthermore, the fundamental theorem in the theory of quadratic residues and its generalization imply that the sum of every $\varphi(4D)$ consecutive values of the factor $\left(\frac{-D}{n}\right)$ vanishes. It follows that the sum of any number of consecutive values of this factor never exceeds, in absolute value, the finite bound depending only on the determinant $D$, namely $\Delta = \varphi(2D)$. Combining these two observations, one readily finds that the sum of all terms following the term $\left(\frac{-D}{n}\right)\frac{m}{n}$ is smaller in absolute value than $\Delta\frac{m}{n}$, and consequently that the quotient $\Theta(m) : m$, as $m$ grows without bound, has as its limit the convergent infinite series

$$\Sigma\left(\frac{-D}{n}\right)\frac{1}{n}$$

ordered in the way specified. Having thus found the common average value of the functions $\theta(m)$ and $f(m)$, one immediately obtains

$$k = \frac{\epsilon\sqrt{D}}{\pi}\Sigma\left(\frac{-D}{n}\right)\frac{1}{n}$$

It should also be observed that Articles 6 and 8 of paper II point to an approach to the problem that is simpler in some respects and also easy to carry out, in which only representations of odd numbers, or even only of numbers relatively prime to $2D$, are considered.

---

## On VI and VII.

The way in which the author summed the series $\Sigma\left(\frac{-D}{n}\right)\frac{1}{n}$ can be seen from several particular examples found on separate sheets.

If $D \equiv 3 \pmod 4$, the fundamental theorem in the theory of quadratic residues, together with the series

$$\operatorname{cotang} u = \frac{1}{u} + \frac{1}{u-\pi} + \frac{1}{u+\pi} + \frac{1}{u-2\pi} + \frac{1}{u+2\pi} + \dots$$

gives

$$\Sigma\left(\frac{-D}{n}\right)\frac{1}{n} = \Sigma\left(\frac{n}{D}\right)\frac{1}{n} = \frac{\pi}{2D}\Sigma\left(\frac{\nu}{D}\right)\operatorname{cotang}\frac{\nu\pi}{2D}$$

where $\nu$ runs through all numbers relatively prime to $2D$ that are smaller than $D$. Put

$$\sqrt{-1} = i, \quad \cos\frac{2\pi}{D} + i\sin\frac{2\pi}{D} = r$$

and let $\mu$ denote all numbers relatively prime to $D$ that are no greater than $D$. The preceding sum can then readily be transformed into

$$\Sigma\left(\frac{-D}{n}\right)\frac{1}{n} = \frac{\pi i}{4D}\left(\frac{2}{D}\right)\Sigma\left(\frac{\mu}{D}\right)\frac{r^\mu - 1}{r^\mu + 1}$$

Now apply the formula valid for every root $\omega$ of the equation $\omega^D = 1$,

$$\frac{\omega - 1}{\omega + 1} = \Sigma(-1)^{\alpha-1}\omega^\alpha$$

in which $\alpha$ must run through the numbers $1, 2, 3 \dots (D-1)$. Reversing the order of summation gives

$$\Sigma\left(\frac{-D}{n}\right)\frac{1}{n} = \frac{\pi i}{4D}\left(\frac{2}{D}\right)\Sigma(-1)^{\alpha-1}\Sigma\left(\frac{\mu}{D}\right)r^{\alpha\mu}$$

The sum over $\mu$ can, as is known, be evaluated by means of the theorems proved in the paper *Summatio quarundam serierum singularium.* Restricting attention to the case in which $D$ is divisible by no square, one finds in general

$$\Sigma\left(\frac{\mu}{D}\right) r^{\alpha\mu} = \left(\frac{\alpha}{D}\right) i^{\left(\frac{D-1}{2}\right)^2} \sqrt{D}$$

where one must put $\left(\frac{\alpha}{D}\right) = 0$ if $\alpha$ is not relatively prime to $D$. In the case $D \equiv 3 \pmod 4$ one therefore obtains

$$\Sigma\left(\frac{-D}{n}\right)\frac{1}{n} = \frac{\pi}{4\sqrt{D}}\left(\frac{2}{D}\right)\Sigma(-1)^{\alpha}\left(\frac{\alpha}{D}\right) = \frac{\pi}{2\sqrt{D}}\Sigma\left(\frac{\alpha'}{D}\right)$$

where $\alpha'$ runs through all numbers relatively prime to $D$ that are smaller than $\frac{1}{2}D$. Finally, since $\epsilon = 2$, the number of classes becomes

$$k = \Sigma\left(\frac{\alpha'}{D}\right)$$

If, on the other hand, $D \equiv 1 \pmod 4$, the series

$$\operatorname{cosec} u = \frac{1}{u} - \frac{1}{u-\pi} - \frac{1}{u+\pi} + \frac{1}{u-2\pi} + \frac{1}{u+2\pi} - \dots$$

gives in a similar way

$$\begin{aligned}\Sigma\left(\frac{-D}{n}\right)\frac{1}{n} &= \Sigma(-1)^{\frac{n-1}{2}}\left(\frac{n}{D}\right)\frac{1}{n} = \frac{\pi}{2D}\Sigma(-1)^{\frac{\nu-1}{2}}\left(\frac{\nu}{D}\right)\operatorname{cosec}\frac{\nu\pi}{2D} \\ &= \frac{\pi}{2D}\Sigma\left(\frac{\mu}{D}\right)\frac{r^{\mu}}{r^{2\mu}+1}\end{aligned}$$

where the letters $\nu$ and $\mu$ have their earlier meanings. Excluding the evident case $D = 1$, apply the formula valid for every root $\omega$ of the equation $\omega^D = 1$ (except $\omega = 1$),

$$\frac{\omega}{\omega\omega+1} = 1 + \Sigma\omega^{4\alpha''} + \Sigma\omega^{D-4\alpha''}$$

in which $\alpha''$ must run through the numbers $1, 2, 3 \dots \frac{1}{4}(D-1)$. Again restricting $D$ to be divisible by no square, one obtains

$$\Sigma\left(\frac{-D}{n}\right)\frac{1}{n} = \frac{\pi}{\sqrt{D}}\Sigma\left(\frac{\alpha''}{D}\right)$$

and hence, since $\epsilon = 2$,

$$k = 2\Sigma\left(\frac{\alpha''}{D}\right)$$

The cases in which $D$ is even could be treated in a very similar way. —

As for determining the number of classes for *positive* determinants $D$, apart from the final formula given in the text, only a few geometrical figures survive. They depict hyperbolic sectors of finite dimensions, with inequalities beside them that confine the points whose coordinates are the variables of the quadratic forms to the interior of such a sector. These hyperbolic sectors take the place of the ellipses corresponding to quadratic forms of negative determinant. Determining their area again gives the average number of representations, provided that only representations whose variables satisfy those inequalities are admitted. On the other hand, the inequalities serve to isolate just one representation, excluding all others, from the infinitely many representations of a number $m$ that belong to one and the same root $n$ of the congruence $nn - D \equiv 0 \pmod{\frac{m}{\mu\mu}}$ and correspond to all the solutions of the equation $tt - Duu = 1$ (compare Disqq. Arithm., art. 205). The total number of admitted representations of the number $m$ by the collection of all properly primitive forms that are not properly equivalent is then equal to the value of the function $f(m)$, in which only $-D$ is to be replaced by $D$. As before for negative determinants, considering its properties gives a second expression for the average number of representations; comparing this with the value previously derived by geometrical considerations then immediately determines the number of classes.

## On VIII.

Here $p$ denotes a positive prime of the form $4n + 1$; the notation agrees with that used in the paper *Theoria residuorum biquadraticorum* I, art. 23. Thus

$$f \equiv 1 \,.\, 2 \,.\, 3 \,.\, \ldots \,.\, \tfrac{1}{2}(p-3) \,.\, \tfrac{1}{2}(p-1) \pmod{p}$$
$$p = aa + bb; \quad a \equiv 1 \pmod{4}; \quad b \equiv af \pmod{p}$$

The numbers denoted by $\alpha, \beta$ are determined by the decomposition $p = \alpha\alpha + 2\beta\beta$. The column $f$ has been added to the two tables found; some gaps in them have also been filled.

The theorem stated in the text is connected with the biquadratic character of the number 2. For, since (compare *Theoria resid. biqu.* I, art. 21)

$$2^{\frac{p-1}{4}} \equiv f^{\frac{1}{2}b} \pmod{p}$$

the congruence

$$b \equiv 2m + a - 1 \pmod{8}$$

implies

$$2^{\frac{p-1}{4}} \equiv f^{m+\frac{a-1}{2}} \pmod{p}$$

and conversely the former follows from the latter. The latter congruence is easily proved as follows. Let $\mu$ be the number of quadratic residues $\alpha_1$ lying between 0 and $\frac{1}{4}p$. Then, by VII,

$$m = 2\mu - \tfrac{1}{4}(p-1)$$

and the number of quadratic residues $\alpha_2$ lying between $\frac{1}{4}p$ and $\frac{1}{2}p$ is $= \frac{1}{4}(p-1) - \mu$. If now $p \equiv 1 \pmod{8}$, so that the number 2 is a quadratic residue, the numbers $2\alpha_1$ and $p - 2\alpha_2$, taken together, form the same collection as the numbers $\alpha_1$ and $\alpha_2$. Denoting the product of these numbers by $A$, one obtains

$$2^{\frac{p-1}{4}} A \equiv (-1)^{\frac{1}{4}(p-1)-\mu} A \pmod{p}$$

and consequently

$$2^{\frac{p-1}{4}} \equiv (-1)^{\mu} \equiv f^{2\mu} \equiv f^{m+\frac{1}{4}(p-1)} \pmod{p}$$

Further, in this case $b \equiv 0 \pmod{4}$, and consequently

$$\frac{p-1}{4} = (a+1)\frac{a-1}{4} + \frac{bb}{4} \equiv 2\frac{a-1}{4} \equiv \frac{a-1}{2} \pmod{4}$$

which gives the congruence to be proved,

$$2^{\frac{p-1}{4}} \equiv f^{m+\frac{a-1}{2}} \pmod{p}$$

If, on the other hand, $p \equiv 5 \pmod 8$, so that the number 2 is a quadratic nonresidue, the numbers $2\alpha_1$ and $p - 2\alpha_2$ comprise all the quadratic nonresidues between 0 and $\frac{1}{2}p$. Denote their product by $B$, and again denote the product of the numbers $\alpha_1$ and $\alpha_2$ by $A$. Then

$$f \equiv AB, \quad (-1)^{\frac{p-1}{4}-\mu} 2^{\frac{p-1}{4}} A \equiv B \pmod p$$

Squaring these two congruences, taking into account that

$$ff \equiv -1, \quad 2^{\frac{p-1}{2}} \equiv -1 \pmod p$$

gives

$$-1 \equiv AABB, \quad -AA \equiv BB$$

and hence $A^4 \equiv +1$. Now $A$ is a product of quadratic residues, so $AA$ is a product of biquadratic residues and is therefore itself a biquadratic residue. Thus $AA \equiv +1$ must hold, because $-1$ is a biquadratic nonresidue. It follows that

$$(-1)^{\frac{p-1}{4}-\mu} 2^{\frac{p-1}{4}} \equiv AB \equiv f \pmod p$$

and

$$2^{\frac{p-1}{4}} \equiv (-1)^{\mu-1} f \equiv f^{2\mu-1} \equiv f^{m+\frac{p-5}{4}} \pmod p$$

Finally, in this case $b \equiv 2 \pmod 4$, and consequently

$$\frac{p-5}{4} = (a+1)\frac{a-1}{4} + \frac{bb-4}{4} \equiv 2\frac{a-1}{4} \equiv \frac{a-1}{2} \pmod 4$$

which again gives the congruence to be proved,

$$2^{\frac{p-1}{4}} \equiv f^{m+\frac{a-1}{2}} \pmod p$$

---

## On IX.

Let $p$ be a positive odd number divisible by no square, and put

$$S_r = \Sigma\left(\frac{s_r}{p}\right)$$

where $s_r$ must run through all numbers relatively prime to $p$ that lie between $(r-1)\frac{p}{8}$ and $r\frac{p}{8}$. Denote the numbers of properly primitive forms that are not properly equivalent, for the determinants $-p$ and $-2p$, by $C_1$ and $C_2$, respectively. Then (compare DIRICHLET, Recherches sur diverses applications etc., § 11, in CRELLE's Journal XXI)

$$C_1 = 2(S_1 + S_2), \quad C_2 = 2(S_1 - S_4)$$

or

$$C_1 = S_1 + S_2 + S_3 + S_4, \quad C_2 = 2(S_2 + S_3)$$

according as $p \equiv 1$ or $\equiv 3 \pmod 4$. Further, the numbers $s_1$ and $s_2$, taken together, form the same collection as the numbers $2s_1$ and $p-2s_4$; likewise the numbers $s_3$ and $s_4$ form the same collection as the numbers $2s_2$ and $p-2s_3$. In the case $p \equiv 1 \pmod 4$, the sum $S_1+S_2+S_3+S_4 = 0$. These observations give two further relations between the four sums $S_1, S_2, S_3, S_4$ in both cases, so that each can be expressed in terms of $C_1$ and $C_2$. In this way, when $p \equiv 1 \pmod 4$, one obtains

$$S_1 = S_8 = \tfrac{1}{4}\left(\frac{2}{p}\right) C_1 + \tfrac{1}{4}C_2$$

$$S_2 = S_7 = \tfrac{1}{4}\left(2 - \left(\frac{2}{p}\right)\right) C_1 - \tfrac{1}{4}C_2$$

$$S_3 = S_6 = -\tfrac{1}{4}\left(2 + \left(\frac{2}{p}\right)\right) C_1 + \tfrac{1}{4}C_2$$

$$S_4 = S_5 = \tfrac{1}{4}\left(\frac{2}{p}\right) C_1 - \tfrac{1}{4}C_2$$

and, when $p \equiv 3 \pmod 4$,

$$S_1 = -S_8 = \tfrac{1}{4}\left(3 + \left(\frac{2}{p}\right)\right) C_1 - \tfrac{1}{4}C_2$$

$$S_2 = -S_7 = -\tfrac{1}{4}\left(1 - \left(\frac{2}{p}\right)\right) C_1 + \tfrac{1}{4}C_2$$

$$S_3 = -S_6 = \tfrac{1}{4}\left(1 - \left(\frac{2}{p}\right)\right) C_1 + \tfrac{1}{4}C_2$$

$$S_4 = -S_5 = \tfrac{1}{4}\left(1 - \left(\frac{2}{p}\right)\right) C_1 - \tfrac{1}{4}C_2$$

If $p$ is a prime, the formulas given in the text for the numbers of quadratic residues in the individual eighths follow immediately.

---

## On X.

Let $p$ be a positive number divisible by no square, of the form $6n \pm 1$, and put

$$S_r = \Sigma \left( \frac{s_r}{p} \right)$$

where $s_r$ must run through all numbers relatively prime to $p$ that lie between $(r-1)\frac{p}{12}$ and $r\frac{p}{12}$. Denote the numbers of properly primitive forms that are not properly equivalent, for the determinants $-p$ and $-3p$, by $C_1$, $C_3$. One readily finds (compare DIRICHLET, Recherches etc., § 11, or the note on VI and VII)

$$C_1 = 2(S_1 + S_2 + S_3), \quad C_3 = 2(S_1 + S_2 - S_5 - S_6)$$

or

$$C_1 = S_1 + S_2 + S_3 + S_4 + S_5 + S_6, \quad C_3 = 2(S_2 + S_3 + S_4 + S_5)$$

according as $p \equiv 1$ or $\equiv 3 \pmod 4$. Taking into account also that

| | | | | | | |
|---|---|---|---|---|---|---|
| the | numbers | $s_1$ and $s_2$ | coincide with | the | numbers | $2s_1$ and $p - 2s_6$ |
| „ | „ | $s_3$ and $s_4$ | „ | „ | „ | $2s_2$ and $p - 2s_5$ |
| „ | „ | $s_5$ and $s_6$ | „ | „ | „ | $2s_3$ and $p - 2s_4$ |

and likewise

| | | | | | | |
|---|---|---|---|---|---|---|
| the | numbers | $s_1, s_2, s_3$ | coincide with | the | numbers | $3s_1,\ 3s_5 - p,\ p - 3s_4$ |
| „ | „ | $s_4, s_5, s_6$ | „ | „ | „ | $3s_2,\ 3s_6 - p,\ p - 3s_3$ |

and that, in the case $p \equiv 1 \pmod 4$, the sum $S_1 + S_2 + S_3 + S_4 + S_5 + S_6 = 0$, one obtains four further relations, besides the two above, between the six sums $S_1, S_2 \ldots S_6$, so that all of them can be determined from $C_1$ and $C_3$. In this way, when $p \equiv 1 \pmod 4$, one obtains

$$S_1 = S_{12} = \tfrac{1}{4}\left(1 + \left(\frac{3}{p}\right)\right) C_1 + \tfrac{1}{12}\left(1 + \left(\frac{2}{p}\right)\right) C_3$$

$$S_2 = S_{11} = -\tfrac{1}{4}\left(1 + \left(\frac{3}{p}\right)\right) C_1 + \tfrac{1}{12}\left(1 + \left(\frac{2}{p}\right)\right) C_3$$

$$S_3 = S_{10} = \tfrac{1}{2} C_1 - \tfrac{1}{6}\left(1 + \left(\frac{2}{p}\right)\right) C_3$$

$$S_4 = S_9 = -\tfrac{1}{2} C_1 - \tfrac{1}{6}\left(2 - \left(\frac{2}{p}\right)\right) C_3$$

$$S_5 = S_8 = \tfrac{1}{4}\left(1 + \left(\frac{3}{p}\right)\right) C_1 - \tfrac{1}{12}\left(5 - \left(\frac{2}{p}\right)\right) C_3$$

$$S_6 = S_7 = -\tfrac{1}{4}\left(1 + \left(\frac{3}{p}\right)\right) C_1 + \tfrac{1}{12}\left(1 + \left(\frac{2}{p}\right)\right) C_3$$

and, when $p \equiv 3 \pmod 4$,

$$S_1 = -S_{12} = \tfrac{1}{12}\left(9 + 3\left(\frac{2}{p}\right) - \left(\frac{3}{p}\right) + \left(\frac{6}{p}\right)\right) C_1 - \tfrac{1}{4} C_3$$

$$S_2 = -S_{11} = \tfrac{1}{4}\left(-1 + \left(\frac{2}{p}\right) + \left(\frac{3}{p}\right) - \left(\frac{6}{p}\right)\right) C_1 + \tfrac{1}{4} C_3$$

$$S_3 = -S_{10} = \tfrac{1}{6}\left(-\left(\frac{3}{p}\right) + \left(\frac{6}{p}\right)\right) C_1$$

$$S_4 = -S_9 = \tfrac{1}{6}\left(3 - 2\left(\frac{3}{p}\right) - \left(\frac{6}{p}\right)\right) C_1$$

$$S_5 = -S_8 = \tfrac{1}{4}\left(-1 - \left(\frac{2}{p}\right) + \left(\frac{3}{p}\right) + \left(\frac{6}{p}\right)\right) C_1 + \tfrac{1}{4} C_3$$

$$S_6 = -S_7 = \tfrac{1}{12}\left(3 - 3\left(\frac{2}{p}\right) + \left(\frac{3}{p}\right) - \left(\frac{6}{p}\right)\right) C_1 - \tfrac{1}{4} C_3$$

If $p$ is a prime, the formulas given in the text follow from the first system; for the other cases, similar formulas are obtained from the second system.

R. Dedekind.

# GEOMETRICAL ASPECT OF TERNARY FORMS.

Let a point in space (0) be chosen as the origin. Let the displacements from it to three other points $P, P', P''$, which do not lie in one plane with it, be $t, t', t''$, respectively; whenever confusion is impossible, the points $P, P', P''$ themselves may be denoted by $(t)$. $(t')$, $(t'')$.

Further, let $(t, t')$ generally denote the product of the lengths of the two lines $t, t'$ and the cosine of their inclination, etc.

In general, $(\alpha t + \alpha' t' + \alpha'' t'' + \ldots \quad \beta u + \beta' u' + \beta'' u'' + \ldots)$ is obtained by carrying out the multiplication

$$(\alpha t + \alpha' t' + \alpha'' t'' + \ldots) \times (\beta u + \beta' u' + \beta'' u'' + \ldots)$$

and replacing $tu, tu', tu'', t'u', t'u'' \ldots$ and so on by $(t, u), (t, u'), (t, u''), (t', u')$, $(t', u'')$ and so on.

Every point in space can be represented by a trinomial

$$(xt + x't' + x''t'')$$

For all points in a given plane, an equation

$$\lambda x + \lambda' x' + \lambda'' x'' = L$$

will then hold, where $\lambda, \lambda', \lambda'', L$ are fixed numbers. For a plane through the three points $\mu t, \mu' t', \mu'' t''$,

$$\lambda\mu = \lambda'\mu' = \lambda''\mu'' = L$$

Writing

$$(t, t) = a, \quad (t', t') = a', \quad (t'', t'') = a'',$$
$$(t', t'') = b, \quad (t, t'') = b', \quad (t, t') = b''$$

and

$$a'a'' - bb = A, \quad aa'' - b'b' = A', \quad aa' - b''b'' = A''$$
$$b'b'' - ab = B, \quad bb'' - a'b' = B', \quad bb' - a''b'' = B''$$
$$D = aa'a'' + 2bb'b'' - abb - a'b'b' - a''b''b''$$

gives

| | | | | |
|---|---|---|---|---|
| $T$ | $=$ | $At + B''t' + B't''$ | perpendicular to | $t'$ and $t''$ |
| $T'$ | $=$ | $B''t + A't' + Bt''$ | | $t$ and $t''$ |
| $T''$ | $=$ | $B't + Bt' + A''t''$ | | $t$ and $t'$ |

and, in general, if

$$\lambda x + \lambda' x' + \lambda'' x'' = L$$

is the equation of a plane, the line

$$\lambda T + \lambda' T' + \lambda'' T''$$

will be perpendicular to that plane.

Further,

$$\begin{aligned} aT + b''T' + b'T'' &= Dt \\ b''T + a'T' + bT'' &= Dt' \\ b'T + bT' + a''T'' &= Dt'' \end{aligned}$$

and the lines $t, t', t''$ are perpendicular to the planes whose equations

$$\begin{aligned} ax + b''x' + b'x'' &= \text{Const} \\ b''x + a'x' + bx'' &= \text{Const} \\ b'x + bx' + a''x'' &= \text{Const} \end{aligned}$$

Twice the area of the triangle through the points $mt, m't', m''t''$ equals the square root of the value of the form

$$F \ldots \begin{pmatrix} A, & A', & A'' \\ B, & B', & B'' \end{pmatrix}$$

on substituting $X = m'm'', X' = mm'', X'' = mm'$, while six times the volume of the pyramid formed with the point 0 becomes $= mm'm''\sqrt{D}$; hence the altitude is

$$= \sqrt{\frac{D}{F(\frac{1}{m}, \frac{1}{m'}, \frac{1}{m''})}}$$

$T, T', T''$ relate to the form $\begin{pmatrix} AD, & A'D, & A''D \\ BD, & B'D, & B''D \end{pmatrix}$ as $t, t', t''$ do to $\begin{pmatrix} a, & a', & a'' \\ b, & b', & b'' \end{pmatrix}$

The three roots of the equation

$$0 = p^3 - pp(a + a' + a'') + p(A + A' + A'') - D$$

give the squares of the three principal axes of an ellipsoid inscribed in the parallelepiped to which the positive ternary form

$$\begin{pmatrix} a, & a', & a'' \\ b, & b', & b'' \end{pmatrix} \quad \text{with adjoint} \quad \begin{pmatrix} A, & A', & A'' \\ B, & B', & B'' \end{pmatrix} \quad \text{and det.} = -D$$

relates.

---

## Expressing spatial relations relative to a given tetrahedron.

Let $(0), (1), (2), (3)$ denote the four vertices, opposite faces and altitudes. Each point in space $P$, relative to an arbitrary origin $M$, then has four coordinates $x, x', x'', x'''$, subject, however, to the relation

$$x + x' + x'' + x''' = 0$$

Here $x$ denotes the quotient obtained by dividing the distance of the point $P$ from a plane drawn through $M$ parallel to the plane $(0)$ by the altitude $(0)$, and so on.

In general,

$$\begin{aligned} -(PM)^2 = {} & xx'(01)^2 + xx''(02)^2 + xx'''(03)^2 \\ & + x'x''(12)^2 + x'x'''(13)^2 + x''x'''(23)^2 \end{aligned}$$

---

*The fundamental law of crystallization* can be stated most briefly as follows: any five planes occurring in it satisfy the following relation:

If their normals on the sphere are $(0), (1), (2), (3), (4)$, the products $\sin 102 . \sin 304$, $\sin 103 . \sin 204$, $\sin 203 . \sin 104$ are always in a rational ratio; if this is $\alpha : \beta : \gamma$, then $\beta = \alpha + \gamma$.

If the coordinates of the 5 points on the sphere are

$$\left.\begin{array}{ccc} a & b & c \\ a' & b' & c' \\ a'' & b'' & c'' \\ a''' & b''' & c''' \\ 0 & 0 & 1 \end{array}\right\} \quad \text{then the products must} \quad \begin{array}{l} (ab' - ba') . (a''b''' - b''a''') \\ (ab'' - ba'') . (a'''b' - b'''a') \\ (ab''' - ba''') . (a'b'' - b'a'') \end{array}$$

be in a rational ratio.

In general, let $1, 2, 3, 4, 5$ be the 5 points on the sphere and 0 its centre; then, if 12 denotes the volume of the tetrahedron 0345,

$$\left.\begin{array}{l} 23 . 45 \\ 24 . 53 \\ 25 . 34 \end{array}\right\} \quad \text{are in a rational ratio}$$

and likewise

$$\left.\begin{array}{l} 12 . 34 \\ 13 . 42 \\ 14 . 23 \end{array}\right\} \quad \text{and so on}$$

**Transformations of the form** $\left(\begin{smallmatrix}5,&5,&5\\-1,&-1,&-1\end{smallmatrix}\right)$ **Det.** $= 108$

| | | | | | |
|---|---|---|---|---|---|
| | | | | | Chaux carbonatée |
| $\left\{\begin{smallmatrix}+1&+1&-1\\-1&+1&+1\\+1&-1&+1\end{smallmatrix}\right\}$ | $\left(\begin{smallmatrix}17&17&17\\-7&-7&-7\end{smallmatrix}\right)$ | | $\left(\begin{smallmatrix}240&240&240\\+168&+168&+168\end{smallmatrix}\right)$ | $\left(\begin{smallmatrix}10&10&10\\+7&+7&+7\end{smallmatrix}\right)$ | equiaxe |
| | 36 . 108 | | 256 . 11664 | 216 | |
| $\left\{\begin{smallmatrix}+1&+1&0\\0&+1&+1\\+1&0&+1\end{smallmatrix}\right\}$ | $\left(\begin{smallmatrix}8&8&8\\+2&+2&+2\end{smallmatrix}\right)$ | $\left(\begin{smallmatrix}4&4&4\\+1&+1&+1\end{smallmatrix}\right)$ | $\left(\begin{smallmatrix}15&15&15\\-3&-3&-3\end{smallmatrix}\right)$ | $\left(\begin{smallmatrix}5&5&5\\-1&-1&-1\end{smallmatrix}\right)$ | inverse |
| | 4 . 108 | 54 | 27 . 108 | 54 | |
| $\left\{\begin{smallmatrix}+2&+1&+1\\+1&+2&+1\\+1&+1&+2\end{smallmatrix}\right\}$ | $\left(\begin{smallmatrix}20&20&20\\+14&+14&+14\end{smallmatrix}\right)$ | $\left(\begin{smallmatrix}10&10&10\\+7&+7&+7\end{smallmatrix}\right)$ | $\left(\begin{smallmatrix}51&51&51\\-21&-21&-21\end{smallmatrix}\right)$ | $\left(\begin{smallmatrix}17&17&17\\-7&-7&-7\end{smallmatrix}\right)$ | contrastante |
| | 16 . 108 | 216 | 432 . 108 | 16 . 108 | |
| $\left\{\begin{smallmatrix}+1&+2&+2\\+2&+1&+2\\+2&+2&+1\end{smallmatrix}\right\}$ | $\left(\begin{smallmatrix}29&29&29\\+23&+23&+23\end{smallmatrix}\right)$ | | $\left(\begin{smallmatrix}312&312&312\\-138&-138&-138\end{smallmatrix}\right)$ | $\left(\begin{smallmatrix}52&52&52\\-23&-23&-23\end{smallmatrix}\right)$ | mixte |
| | 25 . 108 | | 67500 . 108 | 33750 | |

Writing the original form generally as $\left(\begin{smallmatrix}t,&t,&t\\u,&u,&u\end{smallmatrix}\right)$ and a derived form as $\left(\begin{smallmatrix}T,&T,&T\\U,&U,&U\end{smallmatrix}\right)$ gives

$$\begin{aligned}
1,&\quad T = 3t - 2u \quad U = -t + 2u\\
2,&\quad T = 2t + 2u \quad U = t + 3u\\
3,&\quad T = 6t + 10u \quad U = 5t + 11u\\
4,&\quad T = 9t + 16u \quad U = 8t + 17u
\end{aligned}$$

The form $\left(\begin{smallmatrix}1,&3,&k\\0,&0,&0\end{smallmatrix}\right)$ is transformed by the substitution

$$\begin{aligned}
x &= u + u' - 2u'' \quad \text{inversely} \quad 6u = x + 3y + 2z\\
y &= u - u' \qquad\qquad\qquad\quad 6u' = x - 3y + 2z\\
z &= u + u' + u'' \qquad\qquad\quad 6u'' = -2x + 2z
\end{aligned}$$

$$x \equiv z' \pmod 3, \quad x \equiv y \pmod 2$$

into $\left(\begin{smallmatrix}4+k,&4+k,&4+k\\k-2,&k-2,&k-2\end{smallmatrix}\right)$

To obtain calcite, set $k = 0{,}973103$.

Hind the complex values of the orthographic projection of three straight lines of equal length and perpendicular to one another be $a, b, c$, then $aa + bb + cc = 0$; in general, with $p$ and $q$ denoting arbitrary complex numbers, one may set


$$a = (p - q)(q - pii), \quad b = (q - qi)(pii - pi),$$
$$c = (qi - p)(pi - q)$$

## Hexakisoctahedron.




Equation: $px + qy + rz = 1$

*Coordinates.*




| | | | |
|---|---|---|---|
| 1. | $\frac{1}{\gamma}$ | 0 | 0 |
| 2. | $\frac{1}{\beta+\gamma}$ | $\frac{1}{\beta+\gamma}$ | 0 |
| 3. | $\frac{1}{\alpha+\beta+\gamma}$ | $\frac{1}{\alpha+\beta+\gamma}$ | $\frac{1}{\alpha+\beta+\gamma}$ |

$\alpha < \beta < \gamma$

Six times the volume of an elementary pyramid

$$= \frac{1}{\gamma \,.\, (\beta+\gamma)(\alpha+\beta+\gamma)}$$

All [faces] are circumscribed about a sphere whose radius

$$= \frac{1}{\sqrt{(\alpha\alpha+\beta\beta+\gamma\gamma)}}$$

Twice the area of a triangle

$$= \frac{\sqrt{(\alpha\alpha+\beta\beta+\gamma\gamma)}}{\gamma(\beta+\gamma)(\alpha+\beta+\gamma)}$$

Edge

$$1\,.\,2 = \frac{\sqrt{(\beta\beta+\gamma\gamma)}}{\gamma(\beta+\gamma)},$$

$$1\,.\,3 = \frac{\sqrt{((\alpha+\beta)^2+2\gamma\gamma)}}{\gamma(\alpha+\beta+\gamma)},$$

$$2\,.\,3 = \frac{\sqrt{(2\alpha\alpha+(\beta+\gamma)^2)}}{(\beta+\gamma)(\alpha+\beta+\gamma)}$$

Cosine of the angle between the edges $3\,.\,1\,.\,2$

$$= \frac{\alpha\beta+\beta\beta+\gamma\gamma}{\sqrt{(\beta\beta+\gamma\gamma)((\alpha+\beta)^3+2\gamma\gamma)}}$$

Sine

$$= \frac{\gamma \,.\, \sqrt{(\alpha\alpha+\beta\beta+\gamma\gamma)}}{\sqrt{(\beta\beta+\gamma\gamma)((\alpha+\beta)^2+2\gamma\gamma)}}$$

*Values occurring.*

| | $\alpha$ | $\beta$ | $\gamma$ | |
|---|---|---|---|---|
| 7. | 0. | 0. | 1 | Hexahedron |
| 3. | 0. | 1. | 1 | Rhombic dodecahedron |
| 6. | 0. | 1. | 2 | Tetrakishexahedron |
| | 0. | 1. | 3 | |
| | 0. | 2. | 3 | |
| 1. | 1. | 1. | 1 | Octahedron |
| 5. | 1. | 1. | 2 | Deltoidal icositetrahedron |
| | 1. | 1. | 3 | |

| | $\alpha$ | $\beta$ | $\gamma$ | |
|---|---|---|---|---|
| 2. | 1. | 2. | 2. | Triakisoctahedron |
| 4. | 1. | 2. | 3. | Hexakisoctahedron |
| | 1. | 2. | 4. | |
| 2. | 1. | 3. | 3. | Triakisoctahedron |
| 4. | 1. | 3. | 5. | Hexakisoctahedron |
| 2. | 2. | 3. | 3 | Triakisoctahedron |
| | 2. | 3. | 4? | |
| | 3. | 5. | 11 | |

# REMARKS.

Besides the preceding notes, which in part develop the ideas given in the review of SEEBER's investigations of ternary forms, the manuscript records several crystal measurements made by Gauss himself with an eight-inch REICHENBACH theodolite. Each record gives the date of the observation, showing that this investigation belongs to July 1831.

Of the theory of indefinite ternary quadratic forms, the manuscripts left behind contain only the following theorem, probably written down while the Disqu. Arr. were being prepared:

'All transformations of the ternary form

$$\begin{pmatrix} 1, & 1, & -1 \\ 0, & 0, & 0 \end{pmatrix}$$

into itself are given by the formula

$$\begin{array}{lll} \alpha\delta+\beta\gamma & \alpha\beta-\gamma\delta & \alpha\beta+\gamma\delta \\ \alpha\gamma-\beta\delta & \frac{1}{2}(\alpha\alpha+\delta\delta-\beta\beta-\gamma\gamma) & \frac{1}{2}(\alpha\alpha+\gamma\gamma-\beta\beta-\delta\delta) \\ \alpha\gamma+\beta\delta & \frac{1}{2}(\alpha\alpha+\beta\beta-\gamma\gamma-\delta\delta) & \frac{1}{2}(\alpha\alpha+\beta\beta+\gamma\gamma+\delta\delta) \end{array}$$

with $\alpha, \beta, \gamma, \delta$ chosen so that $\alpha\delta - \beta\gamma = 1$.'

All transformations in which the nine coefficients are integers are obtained by taking for $\alpha, \beta, \gamma, \delta$ both all integers satisfying the condition equation, specifically two even and two odd, and all odd multiples of $\sqrt{\frac{1}{2}}$ satisfying the same condition equation $\alpha\delta - \beta\gamma = 1$.

On page 309. Chaux carbonatée équiaxe, inverse, contrastante and mixte are the names used by HAUY (Traité de Minéralogie 1801, volume II, pp. 132, 137).

The table of transformations of the form $\begin{pmatrix} 5, & 5, & 5 \\ 1, & 1, & 1 \end{pmatrix}$ gives, in the first vertical column, the coefficients of the substitution; in the second, the resulting new form; in the third, the primitive form corresponding to the latter, if it is not already primitive; and in the fourth, its adjoint.

SCHERING.

# ON THE THEORY OF BIQUADRATIC RESIDUES.

---

## [I.]

**1.**

We extend the domain of higher arithmetic by including complex quantities as well. In this investigation, we call every quantity $x + iy$ a complex integer when $x, y$ are real integers.

**2.**

The infinitely many complex integers are most conveniently represented visually by points in an unbounded plane; we call the point whose horizontal coordinate is $x$ and vertical coordinate $y$ the point $x + iy$; all points representing integers will be called integer points.

**3.**

To fix a definite convention, the horizontal coordinates shall always be positive to the left, and the vertical coordinates positive upwards.

**4.**

The straight line drawn from the point $x + iy$ to the point $x' + iy'$ will be called the straight line $(x + iy, x' + iy')$; wherever it matters, we also take its direction into account, and therefore distinguish the straight line $x+iy, x'+iy'$ from $x' + iy', x + iy$.

**5.**

For brevity, we shall also denote complex quantities such as $x + iy$ by a single letter, such as $z$.

**6.**

We call the figure bounded by the straight lines $zz', z'z'', z''z''' \dots z^{n-1}z^n$, $z^n z$ the figure $zz'z''z''' \dots z^n$. We do not exclude the case where some of these lines intersect.

**7.**

By $S(z, z', z'' \dots z^n)$ we generally denote a sum of as many real integers as there are integer points inside the figure, assigning the number $\pm 1$, $\pm 2$, $\pm 3$, etc., to each point around which the boundary of the figure passes once, twice, three times, and so on; the upper signs apply when the boundary surrounds the point so that it lies on the right side of the figure, and the lower signs in the opposite case. Thus, if no sides of the figure intersect, $S(z, z', z'' \dots)$ is simply the number of points inside the figure, taken positively or negatively.

**8.**

It is evident that always

$$\begin{aligned} S(z, z', z'' \dots z^n) &= S(z', z'', z''' \dots z^n, z) \\ &= S(z'', z''' \dots z') \text{ etc.} \\ &= -S(z^n, z^{n-1} \dots z'', z', z) \\ &= -S(z^{n-1} \,.\, z^{n-2} \dots z', z, z^n) \text{ etc.} \end{aligned}$$

**9.**

How points on the boundary itself are to be treated must still be specified more closely. In many cases no integer points can lie on the boundary at all; no convention is then needed. But if such points lie on the boundary $zz'$, inserting between $z$ and $z'$ a $+$ indicates that they are to be treated as if they lay to the right of the boundary, and a $-$ as if they lay to the left. We shall also sometimes insert a 0 or $\frac{1}{2}$ to indicate that they are to be counted either not at all or only at half their value on each side. If any of the points $z, z', z''$, etc., is itself an integer point, it is not counted at all unless the contrary is expressly stated, except insofar as it may also occur as a point that is not a vertex.

**10.**

Theorems. If all $z, z', z''$, etc., are increased by the same integer, $S$ remains unchanged.

If $i$ is replaced by $-i$ and every connecting sign by its opposite, $S$ changes only its sign.

$$\begin{aligned} S(z, z', z'' \ldots z^n) &= S(z, u, u' \,..\, u^n, z', z'', z^n) \\ &\quad - S(z, u, u' \,..\, u^n, z') \\ &= S(z, u, u', u'' \,..\, u^n, z^m, z^{m+1} \,..\, z^n) \\ &\quad - S(z, u, u', u'' \,..\, u^n, z^m, z^{m-1} \,..\, z', z) \end{aligned}$$

Here the connecting signs must correspond, but must be reversed between the terms that run backwards.

If $\zeta$ is an integer $= a + bi$, then, when opposite connecting signs are opposite,

$$S(z, z', z' + \zeta, z + \zeta) = [bx' - ay'] - [bx - ay]$$

Here, if $bx' - ay'$ is itself an integer, that integer must be taken for $[bx' - ay']$ when the connecting sign between $z'$ and $z' + \zeta$ is $+$, but 1 or $\frac{1}{2}$ less when this sign is $-$ or $\frac{1}{2}$; the reverse applies to $bx - ay$.

The formula holds only when $a$ and $b$ have no common divisor; if their greatest common divisor is $= h$, the expression to use instead is

$$h\left[\frac{bx' - ay'}{h}\right] - h\left[\frac{bx - ay}{h}\right]$$

**11.**

Let us now turn more closely to our subject itself. If, for the modulus $m = a + bi$, the numbers $f, f', f''$, etc., are such that, first, they are all incongruent to one another modulo $m$, and, second, every integer must necessarily be congruent to one of them, we call the collection of numbers $f, f', f''$, etc., the system of primitive residues of $m$. Their number is always $= aa + bb$.

**12.**

The system of primitive residues can be formed in many ways. The simplest is to choose the points inside the square $0, m, (1 + i)m, im$, but the following must also be included:

I. the point or quantity 0;


II. all points on two boundary lines that are not opposite one another.


Instead of taking all points on one of the 4 boundary lines, one can also take them on several at once.

This choice of points on the boundary lines, if any lie on them, can be made in several ways that satisfy the above conditions. The following method is the simplest.

On the boundary line $0, m$, take all points between 0 and $\frac{1}{2}m$, inclusive, and on the boundary line $0, im$ all points from $\frac{1}{2}im$ to $im$, exclusive; proceed similarly on the other two.

These two methods can be represented visually as follows:




**13.**

If the following are excluded from the primitive points:

I. only the point 0, when $a$ is even and $b$ odd, or the reverse;

II. the points 0 and $\frac{1}{2}(1+i)m$, when $a$ and $b$ are both odd;

III. the four points $0, \frac{1}{2}m, \frac{1}{2}(1+i)m, \frac{1}{2}im$, when $a$ and $b$ are both even,

we call the remaining points proper primitive points and the excluded ones improper. The number of the former is therefore

$$\begin{array}{rrcl} \text{in case} & \text{I} & = & aa+bb-1 \\ & \text{II} & = & aa+bb-2 \\ & \text{III} & = & aa+bb-4 \end{array}$$

and is thus always divisible by 4.

**14.**

These proper primitive points can be divided into 4 classes $F, F', F'', F'''$ so that

$$\begin{array}{llll} iF \equiv F' & iF' \equiv F'' & iF'' \equiv F''' & iF''' \equiv F \\ -F \equiv F'' & -F' \equiv F''' & -F'' \equiv F & -F''' \equiv F' \\ -iF \equiv F''' & -iF' \equiv F & -iF'' \equiv F' & -iF''' \equiv F'' \end{array}$$

The following extremely important theorem now holds.


Let $M$ be a number with no factor in common with $m$. Of the numbers $MF$, the number belonging to the class $F$ is $n$,

$$\begin{array}{ll} F' & n' \\ F'' & n'' \\ F''' & n''' \end{array}$$

and let the least residue of $n' + 2n'' + 3n''$ modulo 4 be $= N$, so that $N$ equals one of the 4 numbers $0, 1, 2, 3$. Under this assumption, $N$ is *independent* of the way the primitive residues are distributed into classes. We call it the decident of the biquadratic relation of the number $M$ to $m$.

**15.**

The simplest distribution is certainly the following:




In particular cases, however, another distribution may be more advantageous.

**16.**


If $f, f', f''$, etc., are all the primitive residues of the modulus $m$, then


$$\begin{aligned}
&S(z + \tfrac{f}{m},\ z' + \tfrac{f}{m},\ z'' + \tfrac{f}{m},\ \text{etc.})\\
&+ S(z + \tfrac{f'}{m},\ z' + \tfrac{f'}{m},\ z'' + \tfrac{f'}{m},\ \text{etc.})\\
&+ S(z + \tfrac{f''}{m},\ z' + \tfrac{f''}{m},\ z'' + \tfrac{f''}{m},\ \text{etc.})\\
&+ \text{etc.}\\
&= S(mz, mz', mz'',\ \text{etc.}
\end{aligned}$$

**17.**

*Theory of the biquadratic residue* $1 + i$.

The modulus must have no divisor in common with the residue. We therefore assume that one of the numbers $a$ and $b$ is even and the other odd. The following diagram shows the distribution of the proper primitive residues into the four classes:



Count towards $n$ all numbers on

$$\text{the line } 0 \ldots \tfrac{1}{2}m \qquad \text{number } = g$$

Towards $n'$, all numbers on

$$\text{the line } 0 \ldots (\tfrac{1}{4} + \tfrac{1}{4}i)m \qquad \text{number } = g'$$

Towards $n''$, all numbers inside

$$\text{the triangle } \tfrac{1}{2}m, m, (\tfrac{1}{2} + \tfrac{1}{2}i)m \qquad \text{number } = h$$

Towards $n'''$, all numbers inside

$$\text{the triangle } 0, \tfrac{1}{2}m, (\tfrac{1}{2} + \tfrac{1}{2}i)m \qquad \text{number } = h'$$

and also all numbers on

$$\text{the line } (\tfrac{1}{4} + \tfrac{1}{4}i)m \,..\, (\tfrac{1}{2} + \tfrac{1}{2}i)m \qquad \text{number } = g''$$

We always have $g' + g'' = g$, $aa + bb = p$, $\frac{1}{4}(p-1) = g + g' + g'' + h + h'$. The decident is therefore

$$D = S(0_{(+)}, \tfrac{1}{2}m, (\tfrac{1}{2} + \tfrac{1}{2}i)m_{(-)}) + \tfrac{1}{2}(p-1) + 2g''$$

Now suppose that, for the modulus $m + 1 + i$,

| | | |
|---|---|---|
| $g$ | changes to | $G$ |
| $g'$ | | $G'$ |
| $g''$ | | $G''$ |
| $h$ | | $H$ |
| $h'$ | | $H'$ |

then we have

$$\begin{aligned}
&\Delta S(0_{(+)}, \tfrac{1}{2}m, (\tfrac{1}{2} + \tfrac{1}{2}i)m_{(-)}) \\
&\quad = +S(0_{(+)}, (\tfrac{1}{2} + \tfrac{1}{2}i)m, (\tfrac{1}{2} + \tfrac{1}{2}i)m + i_{(-)}) \\
&\qquad - S(0_{(+)}, \tfrac{1}{2}m, \tfrac{1}{2}m + \tfrac{1}{2} + \tfrac{1}{2}i_{(-)}) \\
&\qquad - S(\tfrac{1}{2}m, (\tfrac{1}{2} + \tfrac{1}{2}i)m, (\tfrac{1}{2} + \tfrac{1}{2}i)m + i,\ \tfrac{1}{2}m + \tfrac{1}{2} + \tfrac{1}{2}i)
\end{aligned}$$

The last of these $S$ is

$$= \left[\tfrac{1}{2}(a-b)\right] - \left[\tfrac{1}{2}a\right] - S((\tfrac{1}{2} + \tfrac{1}{2}i)m, \tfrac{1}{2}m, \tfrac{1}{2}m + \tfrac{1}{2} - \tfrac{1}{2}i)$$

whether $a$ is odd or even

$$\begin{aligned}
&= \left[\tfrac{1}{2}(a-b)\right] - \left[\tfrac{1}{2}a\right] - S(-\tfrac{1}{2} + \tfrac{1}{2}i, \tfrac{1}{2}m - \tfrac{1}{2} + \tfrac{1}{2}i, \tfrac{1}{2}m + i) \\
&= \left[\tfrac{1}{2}(a-b)\right] - \left[\tfrac{1}{2}a\right] + S(0_{(+)}, \tfrac{1}{2}m, \tfrac{1}{2}m + \tfrac{1}{2} + \tfrac{1}{2}i_{(-)}) \\
&\quad - S(0_{(+)}, (\tfrac{1}{2} + \tfrac{1}{2}i)m, (\tfrac{1}{2} + \tfrac{1}{2}i)m + i_{(-)})
\end{aligned}$$

Thus

$$\begin{aligned}\Delta D = & -\left[\tfrac{1}{2}(a-b)\right]+\left[\tfrac{1}{2}a\right]\\ & +2S(0_{(+)},(\tfrac{1}{2}+\tfrac{1}{2}i)m,(\tfrac{1}{2}+\tfrac{1}{2}i)m+i_{(-)})+a+b+1\\ & -2S(0_{(+)},\tfrac{1}{2}m,\tfrac{1}{2}m+\tfrac{1}{2}+\tfrac{1}{2}i_{(-)})-2g''+2G''\end{aligned}$$

All the connecting signs apply to the case where $a-b$ is positive; otherwise take the opposite signs.

We further split $S(0_{(+)},(\frac{1}{2}+\frac{1}{2}i)m,(\frac{1}{2}+\frac{1}{2}i)m+i_{(-)})$ into

$$\begin{aligned}& S(0_{(+)},(\tfrac{1}{4}+\tfrac{1}{4}i)m,(\tfrac{1}{4}+\tfrac{1}{4}m)+\tfrac{1}{2}i_{(-)})\\ & +\left[\tfrac{1}{2}(a-b)\right]-\left[\tfrac{1}{4}(a-b)\right]\\ & -S((\tfrac{1}{2}+\tfrac{1}{2}i)m_{(-)},(\tfrac{1}{4}+\tfrac{1}{4}i)m,(\tfrac{1}{4}+\tfrac{1}{4}i)m-\tfrac{1}{2}i_{(-)})\end{aligned}$$

The last part

$$\begin{aligned}&= -S(\tfrac{1}{2}+\tfrac{1}{2}i_{(-)},(-\tfrac{1}{4}-\tfrac{1}{4}i)m+\tfrac{1}{2}+\tfrac{1}{2}i,\ (-\tfrac{1}{4}-\tfrac{1}{4}i)m+\tfrac{1}{2}{}_{(+)})\\ &= -S(-\tfrac{1}{2}-\tfrac{1}{2}i_{(-)},(\tfrac{1}{4}+\tfrac{1}{4}i)m-\tfrac{1}{2}-\tfrac{1}{2}i,\ (\tfrac{1}{4}+\tfrac{1}{4}i)m-\tfrac{1}{2}{}_{(+)})\\ &= -S(-\tfrac{1}{2}-\tfrac{1}{2}i_{(+)},(\tfrac{1}{4}+\tfrac{1}{4}i)m-\tfrac{1}{2}-\tfrac{1}{2}i,\ (\tfrac{1}{4}+\tfrac{1}{4}i)m-\tfrac{1}{2}{}_{(-)})+g''-G''\\ &= S(0_{(+)},(\tfrac{1}{4}+\tfrac{1}{4}i)m,(\tfrac{1}{4}+\tfrac{1}{4}i)m+\tfrac{1}{2}i_{(-)})\\ &\quad -S(0_{(+)},\tfrac{1}{2}m,\tfrac{1}{2}m+\tfrac{1}{2}+\tfrac{1}{2}i_{(-)})+g''-G''\end{aligned}$$

This therefore gives

$$\begin{aligned}\Delta D = & +\left[\tfrac{1}{2}(a-b)\right]+\left[\tfrac{1}{2}a\right]\\ & -4S(-\tfrac{1}{2}-\tfrac{1}{2}i_{(+)},\tfrac{1}{4}+\tfrac{1}{4}im-\tfrac{1}{2}-\tfrac{1}{2}i,\ (\tfrac{1}{4}+\tfrac{1}{4}i)m-\tfrac{1}{2}{}_{(-)})\\ & -2\left[\tfrac{1}{4}(a-b)\right]+a+b+1\end{aligned}$$

Increasing the modulus by $1-i,-1+i,-1-i$ requires no separate investigation, because the moduli $m, im, -m, -im$ evidently have equal decidents. We therefore have the following theorems:

If the modulus and its decident are $a+bi, = D$, then the following moduli have the indicated decidents:

| | |
|---|---|
| $a+1+(b+1)i$ | $D+a+b+1+[\frac{1}{2}a]+[\frac{1}{2}(a-b)]-2[\frac{1}{4}(a-b)]$ |
| $a+1+(b-1)i$ | $D+a-b+1+[-\frac{1}{2}b]+[-\frac{1}{2}(a+b)]-2[-\frac{1}{4}(a+b)]$ |
| $a-1+(b+1)i$ | $D-a+b+1+[\frac{1}{2}b]+[\frac{1}{2}(a+b)]-2[\frac{1}{4}(a+b)]$ |
| $a-1+(b-1)i$ | $D-a-b+1+[-\frac{1}{2}a]+[\frac{1}{2}(b-a)]-2[\frac{1}{4}(b-a)]$ |

This also gives

| | | or when $a$ is odd |
|---|---|---|
| $a+2+bi$ | $D-\frac{a-b-3}{2}+2[\frac{1}{2}a]-2[\frac{a-b}{4}]-2[\frac{-a-b-2}{4}]$ | $D+\frac{b-a-1}{2}$ |
| $a-2+bi$ | $D+\frac{a-b+3}{2}+2[-\frac{1}{2}a]-2[\frac{b-a}{4}]-2[\frac{a+b-2}{4}]$ | $D$ |
| $a+(b+2)i$ | $D-\frac{a+b-3}{2}+2[\frac{1}{2}b]-2[\frac{a+b}{4}]-2[\frac{a-b-2}{4}]$ | $D+\frac{a+b-3}{2}$ |
| $a+(b-2)i$ | $D+\frac{a+b+3}{2}+2[-\frac{1}{2}b]-2[\frac{-a-b}{4}]-2[\frac{b-a-2}{4}]$ | |

| | | |
|---|---|---|
| $a+2+(b+2)i$ | $2a-b+1+D$ | or $D+b-1$ |
| $a+2+(b-2)i$ | $-a-2b+1+D$ | $D+a-1$ |
| $a-2+(b+2)i$ | $a+2b+1+D$ | $D-a-1$ |
| $a-2+(b-2)i$ | $-2a+b+1+D$ | $D-b-1$ |

| | | | |
|---|---|---|---|
| $a+4+bi$ | $D+a+b$ | $a+8+bi$ | $D+2$ |
| $a+(b+4)i$ | $D-a+b$ | | |
| $a-4+bi$ | $D-a-b$ | | |
| $a+(b-4)i$ | $D+a-b$ | | |

| | |
|---|---|
| $a+4+(b+4)i$ | $D+2b$ |
| $a+4+(b-4)i$ | $D+2a$ |
| $a-4+(b+4)i$ | $D+2a$ |
| $a-4+(b-4)i$ | $D+2b$ |

The result of the preceding investigations is therefore as follows: For the modulus $m=a+bi$, where $a$ is odd and $b$ even, we have

$$D\frac{1+i}{m}=\tfrac{1}{8}(-aa+2ab+bb-8b+1)$$
(and if $a+bi=1+(2+2i)(\alpha+\beta i)$)
$$\text{or } \tfrac{1}{8}(-aa+2ab-3bb+1)\equiv-(\alpha-\beta)^2-\beta$$

$$D\frac{1-i}{m}=\tfrac{1}{8}(+aa+2ab-bb-8b-1)$$
$$\text{or } \tfrac{1}{8}(+aa+2ab+3bb-1)$$

$$D\frac{-1-i}{m}=\tfrac{1}{8}(-aa+2ab+bb+1)$$
$$=\beta+\alpha\alpha+2\alpha\beta-\beta\beta\equiv-\beta+(\alpha+\beta)^2$$

$$D\frac{-1+i}{m}=\tfrac{1}{8}(+aa+2ab-bb-1)$$
$$=\alpha+\alpha\alpha-2\alpha\beta-\beta\beta\equiv-\alpha-(\alpha+\beta)^2$$

$$D\frac{2}{m} = \tfrac{1}{2}ab$$
$$D\frac{i}{m} = \tfrac{1}{4}(aa + bb - 1)$$
$$D\frac{-1}{m} = \tfrac{1}{2}(aa + bb - 1)$$

*General theorem on decidents.*

Let $A, B, C$, etc., be distinct (odd complex) prime numbers, none of which divides the number $M$; then

$$D\frac{M}{A^\alpha B^\beta C^\gamma D^\delta} = \alpha D\frac{M}{A} + \beta D\frac{M}{B} + \gamma D\frac{M}{C} + \text{etc.}$$

$$M^{\frac{1}{4}(aa+bb-1)} \equiv i^{D\frac{M}{a+bi}} \pmod{(a+bi)} \quad \text{if } a+bi \text{ is a prime number}$$

$$\begin{aligned} D\frac{1+i}{m} &= \tfrac{1}{8}(-aa + 2ab - 3bb + 1) \\ &= -\tfrac{1}{8}(3(a-b) \mp 1)(a - b \mp 1) \quad \text{if } a \equiv \tfrac{1}{3} \\ D\frac{1-i}{m} &= \tfrac{1}{8}(+aa + 2ab + 3bb - 1) \\ D\frac{-1-i}{m} &= \tfrac{1}{8}(-aa + 2ab + bb + 1) \\ &= \tfrac{1}{8}(a - b \mp 1)(a - b \mp 3) \quad \text{if } a \equiv \pm 1 \\ D\frac{-1+i}{m} &= \tfrac{1}{8}(+aa + 2ab - bb - 1) \end{aligned}$$

In general, $m \equiv 1 \bmod.(1+i)$

$$D\frac{1+i}{m} = -\text{Real part } \frac{(1+i)(m^4-1)}{16} = \text{Imag. coeff. } \frac{m^4-1}{8+8i}$$
$$D\frac{1-i}{m} = +\text{Real part } \frac{(1-i)(m^4-1)}{16} = \text{Imag. coeff. } \frac{m^4-1}{8-8i}$$

$1 + i \pmod{16}$


| $a$ | $b$ | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| | 0 | 2 | 4 | 6 | 8 | 10 | 12 | 14 |
| 1 | 0 | 3 | 3 | 0 | 2 | 1 | 1 | 2 |
| 3 | 3 | 3 | 0 | 2 | 1 | 1 | 2 | 0 |
| 5 | 1 | 2 | 0 | 3 | 3 | 0 | 2 | 1 |
| 7 | 2 | 0 | 3 | 3 | 0 | 2 | 1 | 1 |
| 9 | 2 | 1 | 1 | 2 | 0 | 3 | 3 | 0 |
| 11 | 1 | 1 | 2 | 0 | 3 | 3 | 0 | 2 |
| 13 | 3 | 0 | 2 | 1 | 1 | 2 | 0 | 3 |
| 15 | 0 | 2 | 1 | 1 | 2 | 0 | 3 | 3 |

## 18.

*Theory of the biquadratic residue* $-1-2i$.

Let the modulus $= m = a + bi$ be such that $a$ is odd and $b$ even; we also assume that it is a prime number.

The following diagrams represent the decident; their identity is easily seen:




For brevity, we denote $S(x, x+\alpha, x+\alpha+\beta, x+\beta)$ by $[x, \alpha, \beta]$, so that

$$\begin{aligned}[x,\alpha,\beta] &= -[x,\beta,\alpha] = [x+\alpha,\beta,-\alpha] \\ &= -[x+\alpha,-\alpha,\beta] \\ &= [x+\alpha+\beta,-\alpha,-\beta] \\ &= -[x+\alpha+\beta,-\beta,-\alpha]\end{aligned}$$

If we also put

$$\frac{m}{-2-4i} = \frac{-a-2b}{10} + \frac{-b+2a}{10}i = Q$$

then the decident consists of the following eight parts:

$$\begin{aligned}\text{I} &= [0, \tfrac{1}{2}, -iQ] \\ -2\text{II} &= -2[0, \tfrac{1}{2}, Q] \\ \text{III} &= +[Q, \tfrac{1}{2}, -iQ] \\ \text{IV} &= +[-iQ, \tfrac{1}{2}, Q] \\ -3\text{V} &= -3[Q, \tfrac{1}{2}, Q] \\ -3\text{VI} &= -3[2Q, \tfrac{1}{2}, -iQ] \\ +2\text{VII} &= +2[(1-i)Q, \tfrac{1}{2}, Q] \\ +\text{VIII} &= +[0, \tfrac{1}{2}, \tfrac{1}{2}im]\end{aligned}$$

If $F$ denotes an arbitrary elementary residue of the modulus $-1-2i$, then

$$\Sigma[2FQ, \tfrac{1}{2}, iQ] = [0, -\tfrac{1}{2} - i, \tfrac{1}{2}im]$$

Thus, for $F$ substitute $0, 1, i, -1, -i$; we have

$$\begin{aligned} 0 = \quad & [0, \tfrac{1}{2}, iQ] && = \text{IX} \\ & +[2Q, \tfrac{1}{2}, iQ] && = \text{X}. \\ & +[2iQ, \tfrac{1}{2}, iQ] && = \text{XI} \\ & +[-2Q, \tfrac{1}{2}, iQ] && = \text{XII} \\ & +[-2iQ, \tfrac{1}{2}, iQ] && = \text{XIII} \\ & -[0, -\tfrac{1}{2} - i, \tfrac{1}{2}im] && = -\text{XIV} \end{aligned}$$

Add this to the preceding value of the decident. Combining them gives the following results:

(1) Since $(1 + 2i)Q + \frac{1}{2}$ is an integer, we obtain

$$\begin{aligned} \text{XI} &= [-Q - \tfrac{1}{2}, \tfrac{1}{2}, iQ] = -[-Q, -\tfrac{1}{2}, iQ] \\ &= -[Q, \tfrac{1}{2}, -iQ] \end{aligned}$$

and hence

$$\text{III} + \text{XI} = 0$$

(2) We combine $\text{IV}, -3\text{V}, \text{X}, \text{XIII}$ as follows:

$$\begin{aligned} \text{IV} &= +[-2Q + \tfrac{1}{2}i, \tfrac{1}{2}, Q] = [-2iQ - \tfrac{1}{2}, \tfrac{1}{2}i, iQ] \\ \text{V}' &= -[2Q, +\tfrac{1}{2}, -Q] = -[-2iQ, -\tfrac{1}{2}i, iQ] \\ \text{X} &= [+iQ - \tfrac{1}{2}i, \tfrac{1}{2}, iQ] \\ &= [-iQ + \tfrac{1}{2}i, -\tfrac{1}{2}, -iQ] = [-2iQ - \tfrac{1}{2} + \tfrac{1}{2}i, \tfrac{1}{2}, iQ] \\ \text{XIII} &= [-2iQ, \tfrac{1}{2}, iQ] \end{aligned}$$

Thus the entire result from these parts is

$$\begin{aligned} & -4\text{V} \\ & + R(-2iQ) - R(-iQ) - I(-2iQ) + I(-iQ) \\ & - \text{Sq. } [-2iQ - \tfrac{1}{4} + \tfrac{1}{4}i] \\ & + \text{Sq. } [-2iQ + \tfrac{1}{4} - \tfrac{1}{4}i] \end{aligned}$$

(3) Combining $\text{I}, -2\text{II}, -3\text{VI}, +2\text{VII}, +\text{IX}, +\text{XII}$ gives

$$\begin{aligned} \text{I} &= [0, \tfrac{1}{2}, -iQ] \\ \text{II} &= [0, -\tfrac{1}{2}i, -iQ] \\ \text{VI} &= [iQ - \tfrac{1}{2}i, \tfrac{1}{2}, -iQ] = [\tfrac{1}{2} - \tfrac{1}{2}i, -\tfrac{1}{2}, iQ] \\ &= [-\tfrac{1}{2} + \tfrac{1}{2}i, \tfrac{1}{2}, -iQ] \\ \text{VII} &= [-Q - \tfrac{1}{2}i, \tfrac{1}{2}, Q] = [-\tfrac{1}{2} + \tfrac{1}{2}i, -\tfrac{1}{2}, -Q] \\ &= [+\tfrac{1}{2} + \tfrac{1}{2}i, -\tfrac{1}{2}i, -iQ] \\ \text{IX} &= [0, -\tfrac{1}{2}, -iQ] \\ \text{XII} &= [-iQ + \tfrac{1}{2}i, \tfrac{1}{2}, iQ] = [\tfrac{1}{2} + \tfrac{1}{2}i, -\tfrac{1}{2}, -iQ] \end{aligned}$$

$$
\begin{aligned}
&= +4\mathrm{I} - 4\mathrm{II} - 4\mathrm{VI} + 4\mathrm{XII} \\
&\quad - I(\tfrac{1}{2}i) + I(-iQ + \tfrac{1}{2}i) + I0 - I(-iQ) - 2R0 + 2R(-iQ) \\
&\quad + 2\mathrm{Sq.}\ (-iQ + \tfrac{1}{4} + \tfrac{1}{4}i) \\
&= 4\mathrm{I} - 4\mathrm{II} - 4\mathrm{VI} + 4\mathrm{XII} \\
&\quad + R(-1-2i)Q - R(-2iQ) - I(-iQ) + 2R(-iQ) \\
&\quad + 2\mathrm{Sq.}\ (-2iQ + \tfrac{1}{4} + \tfrac{1}{4}i)
\end{aligned}
$$

All this together therefore gives


$$
\begin{aligned}
&+4\mathrm{I} - 4\mathrm{II} - 4\mathrm{V} - 4\mathrm{VI} + 4\mathrm{XII} + \tfrac{1}{2}(a-1) - \tfrac{1}{2}b \\
&+ \mathrm{Sq.}\ (-2iQ + \tfrac{1}{4} - \tfrac{1}{4}i) \\
&+ 2\mathrm{Sq.}\ (-2iQ + \tfrac{1}{4} + \tfrac{1}{4}i) \\
&- \mathrm{Sq.}\ (-2iQ - \tfrac{1}{4} + \tfrac{1}{4}i)
\end{aligned}
$$

Finally, $\mathrm{VIII} - \mathrm{XIV} = \frac{1}{2}(a-1) + \frac{1}{2}b$. Thus, since the three square parts equal the decident of $\frac{m}{-1-2i}$, we obtain

$$\text{Dec. } \frac{-1-2i}{m} \equiv a - 1 + \text{Dec. } \frac{m}{-1-2i} \quad \text{Q. E. D.}$$

The proof can probably be simplified considerably further by the fact that

$$\text{Dec. } \frac{1+2i}{m} =$$




**19.**

The following has been found by induction:

$$
\begin{aligned}
\text{Dec. } \frac{a-bi}{a+bi} &\equiv \frac{aa+2ab-1}{4}, && a \equiv 1 \pmod 4 \\
&\equiv \frac{aa+2ab+2bb-1}{4}, && a+bi \equiv 1 \pmod{2+2i}
\end{aligned}
$$

The following is connected with this: Let $aa + bb = p$ (a prime number), $a \equiv 1 \pmod 4$

$$
\begin{aligned}
1 \,.\, 2 \,.\, 3 \,.\, \ldots\ldots\ldots\ldots \tfrac{1}{4}(p-1) &\equiv \alpha \bmod p \\
\tfrac{1}{4}(p+3) \,.\, \tfrac{1}{4}(p+7) \ldots \tfrac{1}{2}(p-1) &\equiv \beta \\
\tfrac{1}{2}(p+1) \,.\, \tfrac{1}{2}(p+3) \ldots \tfrac{3}{4}(p-1) &\equiv \gamma \\
\tfrac{1}{4}(3p+1) \ldots\ldots\ldots\ldots\ p-1 &\equiv \delta
\end{aligned}
$$

then

$$
\begin{aligned}
&\alpha \equiv \delta, \quad \beta \equiv \gamma \quad \text{if } \tfrac{1}{2}b \text{ is even} \\
&\alpha \equiv -\delta, \quad \beta \equiv -\gamma \quad \text{if } \tfrac{1}{2}b \text{ is odd}
\end{aligned}
$$

$$\pm\alpha\beta \equiv i, \quad \pm\beta\beta \equiv 2b, \quad \frac{\beta}{\alpha} \equiv 2a,$$
$$\frac{\beta}{1+\alpha\beta} \equiv \sqrt{a}, \quad \tfrac{1}{2}\beta(1-\alpha\beta) \equiv \sqrt{a}$$

It therefore remains only to determine the decidents for real residues.

$$a^{\frac{1}{4}bb} \,.\, b^{\frac{1}{4}(aa-1)} \equiv 1 \text{ (mod. } (aa+bb))$$
$$\text{if } a \equiv 1)(\text{mod. } 4) \quad b \text{ even} \quad aa+bb \text{ prime}$$

If only real numbers are to be used, the matter comes down to the following main theorem. Let $a-1$ be divisible by 4, and $b$ by 2; let $a$ and $b$ have no common divisor, and let $k$ denote the numbers $1, 2, 3 \ldots aa+bb-1$. Let

$\alpha$ the number of all values of $k$ for which the least residues of $ak, bk, \quad aak, \quad abk$ all lie between 0 and $\frac{1}{2}(aa+bb)$

$\beta$ $ak, bk, \quad aak, -abk$

$\gamma$ $ak, bk, -aak, -abk$

$\delta$ $ak, bk, -aak, \quad abk$

Then $\beta + 2\gamma + 3\delta - \frac{1}{4}(aa-1)$ is divisible by 4.

---

# [II.]

## PREPARATIONS FOR THE GENERAL THEORY OF BIQUADRATIC RESIDUES.

(1.)

Let $P = x + iy$, where neither $x$ nor $y$ is an integer. We denote the number $+1$ by $LP, L'P, L''P, L'''P$, according as $P$ lies in the first, second, third or fourth quadrant (in the first and second quadrants $[y]$ is even, in the third and fourth odd; in the first and fourth $[x]$ is even, in the second and third odd). In all cases where these symbols are not $= 1$, they are assumed to be $= 0$. We then have the following 24 relations:

$$\begin{array}{lll} L(P \pm 1) = L'P & L(P \pm i) = L'''P & L(P \pm 1 \pm i) = L''P \\ L'(P \pm 1) = LP & L'(P \pm i) = L''P & L'(P \pm 1 \pm i) = L'''P \\ L''(P \pm 1) = L'''P & L''(P \pm i) = L'P & L''(P \pm 1 \pm i) = LP \\ L'''(P \pm 1) = L''P & L'''(P \pm i) = LP & L'''(P \pm 1 \pm i) = L'P \end{array}$$

$$\begin{array}{lll} LiP = L'''P & L(-P) = L''P & L(-iP) = L'P \\ L'iP = LP & L'(-P) = L'''P & L'(-iP) = L''P \\ L''iP = L'P & L''(-P) = LP & L''(-iP) = L'''P \\ L'''iP = L''P & L'''(-P) = L'P & L'''(-iP) = LP \end{array}$$

(2.)

By $PP'$ or $z$ we denote a line beginning at $P$ and ending at $P'$. It need not be straight. We assign a positive and a negative side to every straight line drawn from $2x + 2iy$ to $2x + (2y+1)i$ (where $x, y$ range over all integers); we choose the right side as positive and the left as negative. By $Tz$ we denote the number of all intersections

of the line $z$ with the lines just described, counting positively those where $z$ passes from the negative side to the positive, and negatively the others. We also put

$$Tz - T(z-1) = Sz$$

($z-1$ is a line parallel to $z$, going from the point $P-1$ to $P'-1$). Evidently we need only add to the above system the lines drawn from $2x+1+2yi$ to $2x+1+(2y+1)i$, regarding their left sides as positive and their right sides as negative, to recognize in $Sz$ the number of all intersections of $z$ with this double system of straight lines. We further have

$$\begin{aligned}
T(-z) &= -T(z+i) \\
T(z)+T(z+i) &= [\tfrac{1}{2}x]-[\tfrac{1}{2}x'] \\
S(z+1) &= -Sz \\
S(z+i) &= -Sz+LP+L'''P-LP'-L'''P' \\
S(z+1+i) &= Sz-LP+L'''P+LP'+L'''P' \\
Siz &= Sz-LP+LP' \\
S(-z) &= Sz-LP-L'''P+LP'+L'''P' \\
S(-iz) &= Sz+L'P-L'P'
\end{aligned}$$

**1.**

We consider two kinds of points in the plane: first, those corresponding to integers; second, those determined by multiplying integers by the quantity $Q=\frac{m}{2M}$. We can distinguish them by calling them points of the first order and points of the second order.

**2.**

Joining each point of the second order to its four neighbours by straight lines, which we shall call *ligatures*, divides the whole plane into infinitely many squares. The points of the first order lie partly inside these squares, partly on the ligatures between their

endpoints, and partly at the endpoints of the ligatures, the last case occurring when they are also points of the second order. If $kQ$ is such a point, then, provided $m, M$ have no common divisor and are both odd, $k$ must be divisible by $M$.

**3.**

For ligatures we can also distinguish the starting point from the endpoint, and thus $PQ$ from $QP$, or in some cases set this distinction aside. We call two such ligatures opposite. In general, the most convenient way to denote ligatures is by their starting points and endpoints, which may if desired be enclosed in brackets. The ligature opposite to a given one shall be indicated by a double overline: $QP=\overline{\overline{PQ}}$.

**4.**

Each of the squares in question is enclosed by four such ligatures

$$\{kQ, (k+1)Q\},\ \{(k+1)Q, (k+1+i)Q\},$$
$$\{(k+1+i)Q, (k+i)Q\},\ \{(k+i)Q, kQ\} \ldots . \Omega$$

to whose right it lies. Here it is important to consider the form of the number $k$. We distinguish four kinds of squares according as $k \equiv 0, 1, 1+i, i \pmod 2$, and use the numbers $0, 1, 2, 3$, which we call the respective intensors of the squares.

**5.**

We assign to the ligatures the same intensors as those of the squares lying to their right.

**6.**

We must now consider another, larger square $\Omega'$, namely the one obtained when the square indicated in 4 for $k = 0$ is multiplied by $M$: it is therefore bounded by the straight lines $\mu, \mu', \mu'', \mu'''$

$$\{0, \tfrac{1}{2}m\},\ \{\tfrac{1}{2}m, \tfrac{1}{2}(1+i)m\},\ \{\tfrac{1}{2}(1+i)m, \tfrac{1}{2}im\},\ \{\tfrac{1}{2}im, 0\}$$

It consists of complete squares $\Omega$ and pieces of such squares. Add up

all points of the first order inside it, taking for each point the intensor of the square $\Omega$ in which it lies. This sum, or its least residue modulo 4, is called the decident of $M$ for the modulus $m$, and determines the biquadratic modality of $M$ with respect to this modulus.

**7.**

We divide the square $\Omega'$ into 5 pieces as follows. Join the point 0 to $\frac{1}{4}(1+i)(m-1)$ by the line $\lambda$, which is to run entirely along ligatures inside $\Omega'$. Let

$$\tfrac{1}{2}m + i\lambda = \lambda', \quad \tfrac{1}{2}(1+i)m - \lambda = \lambda'', \quad \tfrac{1}{2}im - i\lambda = \lambda'''$$

These 4 lines therefore run from the corners of the square $\Omega'$ into its interior and end at the four corners of the innermost square. Its intensor will be 0 if $m \equiv 1 \pmod{2+2i}$; let the ligatures of this square be $\nu, \nu', \nu'', \nu'''$.

The 5 pieces will therefore be bounded by

I. $\ldots$ $\mu,\ \lambda',\ \overline{\overline{\nu}},\ \overline{\overline{\lambda}}$
II. $\ldots$ $\mu',\ \lambda'',\ \overline{\overline{\nu'}},\ \overline{\overline{\lambda'}}$
III. $\ldots$ $\mu'',\ \lambda''',\ \overline{\overline{\nu''}},\ \overline{\overline{\lambda''}}$
IV. $\ldots$ $\mu''',\ \lambda,\ \overline{\overline{\nu'''}},\ \overline{\overline{\lambda'''}}$
V. the inner square $\nu,\ \nu',\ \nu'',\ \nu'''$

The decident is therefore the enumeration of all points of the first order in I. II. III. IV.

**8.**

For brevity, the intensor of any point shall mean the intensor of the square in which it lies, and shall be expressed by a prefixed $\Upsilon$.

**9.**

The decident is therefore

$$\Sigma\Upsilon P + \Sigma\Upsilon P' + \Sigma\Upsilon P'' + \Sigma\Upsilon P'''$$

where $P$ denotes all points in I., and so on.

**10.**

We now also consider the region VI $= -i$IV, which lies outside $\Omega'$, but adjoins I along $\mu$ and together with it forms the region $\omega$, consisting of $AA + BB$ complete squares. If $\Pi$ denotes all integer points, and $\Pi'$ all integer points increased by $\frac{1}{2}i$, in this region, it is easy to prove that the decident

$$\begin{aligned} &= \Sigma\Upsilon\Pi - \Sigma\Upsilon\Pi' + \text{number of all integer points inside VI} \\ &\quad - \text{number of all half-points inside VI.} \end{aligned}$$

**11.**

Imagine straight lines drawn from each integer point $k$ to $k + \frac{1}{2}i$, their right sides regarded as positive and their left sides as negative. Let $l$ be a line, and let $Sl$ denote the sum of all intersections of $l$ with this system of lines, counting positively those where $l$ passes from the negative side to the positive, and negatively the opposite intersections. The decident then has the following expression:

$$\Sigma(\Upsilon l \,.\, Sl) + \Sigma Sl' - S\mu$$

Here $l$ denotes all ligatures of the squares in $\omega$ (always taken so that the squares lie to their right), and $l'$ denotes the ligatures on the boundary of the figure $\omega$ between 0 and $\frac{1}{2}m$, hence outside $\Omega'$.

All ligatures $l$ consist of

1) $l'$;

2) $l''$, the boundary ligatures inside $\Omega'$, hence $\lambda', \overline{\overline{\nu}}, \overline{\overline{\lambda}}$;

3) $l'''$, which lie in the interior of $\omega$.

If $l$ were allowed to range over all ligatures inside $\omega$ or on the boundary of this figure, insofar as they start at points $\frac{km}{2M}$ such that $k$ is divisible by $1+i$, the decident would be

$$= \Sigma\alpha \,.\, Sl - S\mu$$

where $\alpha = 1$ for all ligatures in the interior of $\omega$;

$\alpha = \Upsilon l + 1$ for all boundary ligatures outside $\Omega'$ whose direction follows the boundary running from 0 to $\frac{1}{2}m$;

$\alpha = -(\Upsilon\overline{\overline{\lambda}} + 1) = -\Upsilon l$ for all those on this boundary running in the opposite direction;

$\alpha = \Upsilon l$ for all boundary ligatures inside $\Omega'$ directed towards 0;

$\alpha = -\Upsilon l + 1$ for all boundary ligatures inside $\Omega'$ directed away from 0.

**12.**

We can now combine all the $l$ that occur (in the last manner) in pairs, namely $l$ with $\overline{\overline{\frac{1}{2}m - l}}$, which we shall call associated ligatures. A single one is excepted, standing isolated or identical with its associated ligature, namely the one running from

$$\tfrac{1}{2}(M-1) \,.\, \frac{m}{2M} \quad \text{to} \quad \tfrac{1}{2}(M+1)\frac{m}{2M}$$

For associated ligatures, $\alpha$ is always the same.

[III.]

## Theory of Biquadratic Residues.

---

**1.**

The least residues for the modulus $m = a + bi$ are the complex integers $\mu = \alpha + \beta i$ for which $\frac{\mu}{m} = x + yi$ has $x$ and $y$ positive and less than 1. The residue 0 is included as well*). Their number is $= aa + bb$.

**2.**

If $aa + bb$ is odd, $aa + bb$ is of the form $4n + 1$. On excluding the least residue 0, the others fall into four classes. In the first class $f$ we place those for which $x$ and $y$ are less than $\frac{1}{2}$,

| | | | |
|---|---|---|---|
| the second | $f'$ | where $x > \frac{1}{2}$, | $y < \frac{1}{2}$ |
| third | $f''$ | $x > \frac{1}{2}$, | $y > \frac{1}{2}$ |
| fourth | $f'''$ | $x < \frac{1}{2}$, | $y > \frac{1}{2}$ |

All the residues are obtained

| | | |
|---|---|---|
| $f'$ | from | $if + m$ |
| $f''$ | from | $-f + (1 + i)m$ |
| $f'''$ | from | $-if + im$ |

**3.**

Let $M$ be another number with no factor in common with $m$. Then

$$M^{aa+bb-1} \equiv 1 \text{ (mod. } m)$$

holds; consequently $M^{\frac{1}{4}(aa+bb-1)}$ is either $\equiv 1$, or $\equiv i$, or $\equiv -1$, or $\equiv -i$, that is, $\equiv i^{\varepsilon}$, where $\varepsilon$ represents one of the four numbers $0, 1, 2, 3$. In the first case $M$ is a biquadratic residue of $m$, and hence also a quadratic residue. In the third case $M$ is a quadratic residue but not a biquadratic residue; in the second and fourth cases it is both a quadratic and a biquadratic nonresidue. We call this $\varepsilon$, on which the biquadratic character of the number $M$ with respect to the modulus $m$ depends, the decident of $M$ for the modulus $m$. Induction gives the following beautiful theorem. "If $M$ and $m$ are odd primes of the form $1 + (2 + 2i)\mu$, where $\mu$ is an integer, the difference between the two decidents, that of $M$ for the modulus $m$ and that of $m$ for the modulus $M$, is either $= 0$ or $= 2$; the former when at least one of the numbers $m, M$ is of the form $1 + 4N$, the latter when both are of the form $1 + 2i + 4N$." This reciprocity theorem is analogous to the one for quadratic residues among real numbers alone.

*) and, if $a$ and $b$ happen to have a common divisor $e$, the numbers $\frac{m}{e}, \frac{2m}{e}, \frac{3m}{e} \ldots \frac{(e-1)m}{e}$. For now, however, we shall leave this case out of the investigation.

**4.**

Multiply all the numbers $f$ by $M$ and find their least residues for the modulus $m$. If among them there are

$$\begin{array}{ll} \alpha & \text{belonging to } f \\ \beta & f' \\ \gamma & f'' \\ \delta & f''' \end{array}$$

then $\varepsilon \equiv \beta + 2\gamma + 3\delta \pmod 4$.

Proof. Let the collection of those numbers in $f$ whose products with $M$ give residues belonging to $f$ be $g$; let the collection of those whose products give residues from $f'$ be $g'$, and similarly $g''$ . $g'''$. Then the least residues of

$$-ig'M, \quad -g''M, \quad ig'''M$$

are all contained in $f$ and are distinct both from one another and from the least residues of the products $gM$. Consequently the product of all

$$gM, \quad -ig'M, \quad -g''M, \quad +ig'''M$$

is congruent to the product of all $f$, and hence also to the product of all $g, g', g'', g'''$. But that product equals the product of all $g, g', g'', g'''$ multiplied by

$$M^\alpha \,.\, (-iM)^\beta \,.\, (-M)^\gamma \,.\, (iM)^\delta$$

so this last product is $\equiv 1$

$$\begin{array}{rl} \text{therefore} & M^{\alpha+\beta+\gamma+\delta}(-i)^\beta(-1)^\gamma i^\delta \equiv 1 \\ \text{or} & M^{\alpha+\beta+\gamma+\delta} \equiv i^\beta(-1)^\gamma(-i)^\delta \equiv i^{\beta+2\gamma+3\delta} \end{array}$$

from which the theorem follows directly.

**5.**

It is easy to decide whether the least residue of a number $N$ for the modulus $m$ belongs to the class $f, f', f''$ or $f'''$. If $\omega$ is the integer contained in $\frac{N}{m}$, that residue is $= N - \omega m$ and therefore belongs to $f, f', f''$, according as, on putting

$$\frac{N}{m} - \omega = x + iy$$

we have

$$\begin{array}{ll} x < \frac{1}{2}, & y < \frac{1}{2} \\ x > \frac{1}{2}, & y < \frac{1}{2} \\ x > \frac{1}{2}, & y > \frac{1}{2} \\ x < \frac{1}{2}, & y > \frac{1}{2} \end{array}$$

In these 4 cases the integer contained in $\frac{2N}{m}$ is, in the same order, as follows:

$$\begin{array}{l} 2\omega \\ 2\omega+1 \\ 2\omega+1+i \\ 2\omega+i \end{array}$$

It follows that the least residue of $N$ for the modulus $m$ belongs to $f, f', f'', f'''$ according as, on writing the integer contained in $\frac{2N}{m}$ as $= \xi + \eta i$, we have

$$\begin{array}{ll} \xi \text{ even} & \eta \text{ even} \\ \xi \text{ odd} & \eta \text{ even} \\ \xi \text{ odd} & \eta \text{ odd} \\ \xi \text{ even} & \eta \text{ odd} \end{array}$$

**6.**

The decident of $M$ for the modulus $m$ is therefore found as follows. Find the integers contained in each of the individual $\frac{2fM}{m}$. Writing these generally as $x + yi$, disregard entirely those for which $x$ and $y$ are both even; count one for each for which $x$ is odd and $y$ even, take two for each for which $x$ and $y$ are both odd, and three for each for which $x$ is even and $y$ odd. Take the least residue modulo 4 of the sum of all these numbers; it is the required decident. We express this as

$$\text{Dec.}\frac{M}{m} = \Sigma n$$

where $[\frac{2fM}{m}] = x + yi$, and we put $n = 0$ when $x$ is even and $y$ even

$$\begin{array}{lll} 1 & x \text{ odd} & y \text{ even} \\ 2 & x \text{ odd} & y \text{ odd} \\ 3 & x \text{ even} & y \text{ odd} \end{array}$$

For brevity we shall denote $n$ by the characteristic $\Theta$, $n = \Theta\frac{2fM}{m}$ *).

*) To decide which class $M$ belongs to with respect to $m$, choose those values of $k$ (among the numbers $1, 2, 3 \ldots p-1$) for which $[\frac{2km'M}{p}]$ is even, and add $-\Sigma[\frac{2km'}{p}]^2$. If $k$ is taken only up to $\frac{1}{2}p$, the sum to be taken is

$$-\Sigma\{[\frac{2km'M}{p}]^2 + [\frac{2km'}{p}]^2\}$$

for those values of $[\frac{2km'M}{p}]$ that are divisible by $1+i$.

**7. 8.**

This rule is general, whatever number $M$ denotes. For the case that will be the immediate subject of our investigation, where $M$ is assumed to be odd and of the form $1 + (2 + 2i)N$, a slightly modified rule is more convenient.

Suppose the numbers $f$ are again divided into 4 classes. In the first we place the $(h)$, whose doubles are also found in $f$; in the second, $h'$, we count those whose doubles $2h'$ belong to $f'$; similarly, $h''$ and $h'''$ denote those whose doubles belong to $f''$ and $f'''$. Thus the decident $\varepsilon$ is

$$\varepsilon = \Sigma\Theta\frac{2hM}{m} + \Sigma\Theta\frac{2h'M}{m} + \Sigma\Theta\frac{2h''M}{m} + \Sigma\Theta\frac{2h'''M}{m}$$

We call the collection of all $2h$ and $-2h'' + (1 + i)m$ $H$

and the collection of all $-i(2h' - m)$ and $i(2h''' - im)$ $H'$

Thus $H$ and $H'$ comprise all $f$; the former are the even ones, the latter the odd ones.

The following relations must also be applied

$$\begin{aligned}
\Theta iN &= 1 + \Theta N \\
\Theta(-N) &= 2 + \Theta N \\
\Theta(-iN) &= 3 + \Theta N \\
\Theta(N + 1) &= 1 - \Theta N \\
\Theta(N + 1 + i) &= 2 + \Theta N \\
\Theta(N + i) &= 3 - \Theta N
\end{aligned}$$

consequently

$$\begin{aligned}
\Theta\frac{(-2h'i + mi)M}{m} &= 3 - \Theta\frac{-2h'iM}{m} = -\Theta\frac{2h'M}{m} \\
\Theta\frac{(-2h'' + m(1 + i))M}{m} &= 2 + \Theta\frac{-2h''M}{m} = \Theta\frac{2h''M}{m} \\
\Theta\frac{(2h'''i + m)M}{m} &= 1 - \Theta\frac{2h'''iM}{m} = -\Theta\frac{2h'''M}{m}
\end{aligned}$$

and

$$\begin{aligned}
\varepsilon &= \Sigma\Theta\frac{2hM}{m} - \Theta\frac{(-2h'i + mi)M}{m} \\
&\quad + \Theta\frac{(-2h'' + m(1 + i))M}{m} - \Theta\frac{(2h'''i + m)M}{m} \\
&= \Sigma\Theta\frac{HM}{m} - \Sigma\Theta\frac{H'M}{m}
\end{aligned}$$

or

$$\varepsilon = \Sigma \pm \Theta\frac{fM}{m}$$

where the upper sign is to be taken for even $f$, the lower for odd.

**9.**

Let, in general, $f = \xi + \eta i$. The numbers $\xi, \eta$ are restricted within certain bounds by the condition that $f$ be a least residue of $m$, or that, on putting $\frac{f}{m} = x + yi$, $x$ and $y$ must lie between the bounds 0 and $\frac{1}{2}$. Definite rules for these bounds could easily be given by distinguishing the different cases. If we give $\eta$ a particular value, $\xi$ will in turn have its particular bounds. For example, if we assume that $a$ is negative and $b$ positive, then, since

$$x = \frac{a\xi + b\eta}{aa + bb}$$
$$y = \frac{a\eta - b\xi}{aa + bb}$$

the following conditions must hold:

| | | |
|---|---|---|
| I. | for $x$ to be positive | $\xi < -\frac{b}{a}\eta$ |
| II. | for $y$ to be positive | $\xi < \frac{a}{b}\eta$ |
| III. | for $x < \frac{1}{2}$ to hold | $\xi > \frac{aa+bb-2b\eta}{2a}$ |
| IV. | for $y < \frac{1}{2}$ to hold | $\xi > \frac{2a\eta-aa-bb}{2b}$ |

For positive $\eta$, the second condition already includes the first; for negative $\eta$, the reverse holds. Likewise, the third condition is already contained in the fourth,

| | |
|---|---|
| if | $\eta < \frac{1}{2}(a+b)$ |
| and conversely if | $\eta > \frac{1}{2}(a+b)$ |

We do not, however, need to consider separately all eight cases that can occur here. For a particular value of $\eta$, we merely denote the smaller bound of $\xi$ by $\xi^0$, the larger by $\xi^{00}$, and remark only that at these boundary values one always has either $x = 0, y = 0, x = \frac{1}{2}, y = \frac{1}{2}$, and specifically that the following must hold:

| if | at the *upper* bound | at the *lower* bound |
|---|---|---|
| $a$ pos. $b$ positive | $x = \frac{1}{2}$ or $y = 0$ | $x = 0$ or $y = \frac{1}{2}$ |
| $a$ neg. $b$ positive | $x = 0$ or $y = 0$ | $x = \frac{1}{2}$ or $y = \frac{1}{2}$ |
| $a$ neg. $b$ negative | $x = 0$ or $y = \frac{1}{2}$ | $x = \frac{1}{2}$ or $y = 0$ |
| $a$ pos. $b$ negative | $x = \frac{1}{2}$ or $y = \frac{1}{2}$ | $x = 0$ or $y = 0$ |

For brevity we shall distinguish these four cases by saying that, in the first, $m$ belongs to the first quadrant, in the second to the second, and so on.

**10.**

We now wish to examine more closely the sum of all $\pm\Theta\frac{fM}{m}$ for which $\eta$ has a particular value. As $\xi$ gradually increases continuously from its smallest value $\xi^0$ to its largest $\xi^{00}$,

$$\frac{(\xi+\eta i)M}{m} = X + Yi$$

will also change according to the law of continuity. If $\frac{M}{m}$ lies in the first quadrant, both $X$ and $Y$ will keep increasing; if $\frac{M}{m}$ lies in the second quadrant, $X$ will keep decreasing and $Y$ increasing. In the third quadrant the reverse of the first case occurs, and in the fourth the reverse of the second. But the integer contained in $X+iY$ will change in jumps, as either $[X]$ or $[Y]$ changes by one unit. Let the values of $\xi$ at which such a transition occurs, that is, at which either $X$ or $Y$ becomes an integer, be, in order,

$$\xi',\ \xi'',\ \xi'''\ldots.\xi^n$$

Here it must be remarked that neither these values nor $\xi^0$ and $\xi^{00}$ can be integers, except for $\eta = 0$, when either $\xi^0$ or $\xi^{00} = 0$. Now let

$$\Theta\frac{(\xi'+\eta i)M}{m} - \Theta\frac{(\xi^0+\eta i)M}{m} = \delta' \quad \text{(to be expressed differently)}$$
$$\Theta\frac{(\xi''+\eta i)M}{m} - \Theta\frac{(\xi'+\eta i)M}{m} = \delta''$$
$$\text{etc.}$$
$$\Theta\frac{(\xi^{00}+\eta i)M}{m} - \Theta\frac{(\xi^{n-1}+\eta i)M}{m} = \delta^n$$

Then it is easy to see, since between $\xi^0$ and $\xi'$ there lie $[\frac{1}{2}\xi'] - [\frac{1}{2}\xi^0]$ even integers and $[\frac{1}{2}\xi' + \frac{1}{2}] - [\frac{1}{2}\xi^0 + \frac{1}{2}]$ odd integers, and so on, that, considering only the particular value of $\eta$,

$$\begin{aligned}
(\pm 1)\Sigma\Theta\frac{fM}{m} &= \{[\tfrac{1}{2}\xi'] - [\tfrac{1}{2}\xi^0] - [\tfrac{1}{2}\xi' + \tfrac{1}{2}] + [\tfrac{1}{2}\xi^0 + \tfrac{1}{2}]\} \\
&\qquad . \Theta\frac{(\xi^0 + \eta i)M}{m} \\
&\quad + \{[\tfrac{1}{2}\xi''] - [\tfrac{1}{2}\xi'] - [\tfrac{1}{2}\xi'' + \tfrac{1}{2}] + [\tfrac{1}{2}\xi' + \tfrac{1}{2}]\} \\
&\qquad . \{\Theta\frac{(\xi^0 + \eta i)M}{m} + \delta'\} \\
&\quad + \{[\tfrac{1}{2}\xi'''] - [\tfrac{1}{2}\xi''] - [\tfrac{1}{2}\xi''' + \tfrac{1}{2}] + [\tfrac{1}{2}\xi'' + \tfrac{1}{2}]\} \\
&\qquad . \{\Theta\frac{(\xi^0 + \eta i)M}{m} + \delta' + \delta''\} \\
&\quad + \text{etc.} \\
&\quad + \{[\tfrac{1}{2}\xi^{00}] - [\tfrac{1}{2}\xi^n] - [\tfrac{1}{2}\xi^{00} + \tfrac{1}{2}] + [\tfrac{1}{2}\xi^n + \tfrac{1}{2}]\} \\
&\qquad . \{\Theta\frac{(\xi^0 + \eta i)M}{m} + \delta' + \delta'' + \ldots + \delta^n\}
\end{aligned}$$

$$\begin{aligned}
= \; & -([\tfrac{1}{2}\xi^0] - [\tfrac{1}{2}\xi^0 + \tfrac{1}{2}]) \,.\, \Theta\frac{(\xi^0 + \eta i)M}{m} \\
& -([\tfrac{1}{2}\xi'] - [\tfrac{1}{2}\xi' + \tfrac{1}{2}]) \,.\, \delta' \\
& -([\tfrac{1}{2}\xi''] - [\tfrac{1}{2}\xi'' + \tfrac{1}{2}]) \,.\, \delta'' \\
& -\text{etc.} \\
& -([\tfrac{1}{2}\xi^n] - [\tfrac{1}{2}\xi^n + \tfrac{1}{2}]) \,.\, \delta^n \\
& +([\tfrac{1}{2}\xi^{00}] - [\tfrac{1}{2}\xi^{00} + \tfrac{1}{2}]) \,.\, \Theta\frac{(\xi^{00} + \eta i)M}{m}
\end{aligned}$$

(where the upper sign applies for even $\eta$, the lower for odd.)

The numbers $\delta', \delta'', \delta'''$, and so on, can have no values other than $+1$ and $-1$. The value $+1$ is obtained by $\delta'$ when, denoting the values of $X, Y$ belonging to $\xi'$ by $X', Y'$,

| $\frac{M}{m}$ in quadrant 1. | $\frac{M}{m}$ in quadrant 2. |
|---|---|
| $X'$ even integer and $[Y]$ odd | $X'$ even integer $[Y]$ even |
| $Y'$ even integer and $[X]$ even | $Y'$ even integer $[X]$ even |

| $\frac{M}{m}$ in quadrant 3. | $\frac{M}{m}$ in quadrant 4. |
|---|---|
| $X'$ even integer and $[Y]$ even | $X'$ even integer $[Y]$ odd |
| $Y'$ even integer and $[X]$ odd | $Y'$ even integer $[X]$ odd |

Whenever one of these conditions changes to its opposite, $\delta = -1$; whenever both change, $\delta = +1$ remains.

**11.**

The following formulas provide a more convenient overview of these calculations:

$$\begin{aligned} \text{here } m &= a + bi, & aa + bb &= d \\ M &= A + Bi, & AA + BB &= D \\ \frac{dM}{m} &= \alpha + \beta i, & \alpha &= aA + bB, \quad \beta = aB - bA \\ \frac{\xi + i\eta}{m} &= x + iy, & M(x + iy) &= X + iY \end{aligned}$$

If $\eta$ and $X$ are given, then

$$\begin{aligned} &1. \quad \xi = \tfrac{\beta\eta}{\alpha} + \tfrac{dX}{\alpha} \\ &2. \quad Y = \tfrac{D\eta}{\alpha} + \tfrac{\beta X}{\alpha} \end{aligned}$$

If $\eta$ and $Y$ are given, then

$$\begin{aligned} &3. \quad \xi = -\tfrac{\alpha\eta}{\beta} + \tfrac{dY}{\beta} \\ &4. \quad X = -\tfrac{D\eta}{\beta} + \tfrac{\alpha Y}{\beta} \end{aligned}$$

If $\eta$ and $x$ are given, then

$$\begin{aligned} &5. \quad \xi = -\tfrac{b\eta}{a} + \tfrac{dx}{a} \\ &6. \quad X = -\tfrac{B\eta}{a} + \tfrac{\alpha x}{a} \\ &7. \quad Y = \tfrac{A\eta}{a} + \tfrac{\beta x}{a} \end{aligned}$$

If $\eta$ and $y$ are given, then

$$8. \quad \xi = \frac{a\eta}{b} - \frac{dy}{b}$$
$$9. \quad X = \frac{A\eta}{b} - \frac{\alpha y}{b}$$
$$10. \quad Y = \frac{B\eta}{b} - \frac{\beta y}{b}$$

**12.**

The rule of Article 10 can now be expressed as follows. Giving $\eta$ a particular value, we have

$$\Sigma \pm \Theta \frac{fM}{m} = k^0 \Theta(X^0 + Y^0 i) - k^{00} \Theta(X^{00} + Y^{00} i) + \Sigma k$$

Here $k^0 = 0$ when $[\xi^0]$ is even; $k^0 = +1$ when $[\xi^0]$ is odd and $\eta$ even; $k^0 = -1$ when $[\xi^0]$ is odd and $\eta$ odd. $k^{00}$ is determined in the same way by $[\xi^{00}]$ and $\eta$. Finally, $\Sigma k$ is a sum of as many numbers as there are, between $\xi = \xi^0$ and $\xi = \xi^{00}$, integer values of $X$ or $Y$; in each case $k = 0$ when the corresponding $[\xi]$ is even, and $= \pm 1$ when $[\xi]$ is odd. The sign is determined as follows. If $X$ is an integer, then $k = 1$ when all the following hold:

| | |
|---|---|
| $\eta$ | even |
| $X$ | even |
| $[Y]$ | even |
| $\frac{M}{m}$ | in the second or third quadrant, i.e. $\alpha$ negative |

If one or three of these conditions are absent, then $k = -1$; if two or all four are absent, $k = 1$ remains. If, on the other hand, $Y$ is an integer, then $k = 1$ when all, two, or none of the 4 conditions

| | |
|---|---|
| $\eta$ | even |
| $Y$ | even |
| $[X]$ | even |
| $\frac{M}{m}$ | in the first or second quadrant, i.e. $\beta$ positive |

are satisfied.

**13.**

We must now consider separately the cases in which $\xi^0$ or $\xi^{00}$ (or $X^0, Y^0, X^{00}, Y^{00}$) is an integer. There are four cases to distinguish here, taking $a$ and $A$ to be odd.

I. If $m$ lies in the first quadrant, then for $x = \frac{1}{2}, y = 0$, $\eta = \frac{1}{2}b$ is an integer. Then $Y^{00} = \frac{1}{2}B$ is an integer, and $\Theta(X^{00} + Y^{00} i)$ will be $= \Theta(\frac{1}{2}A + \frac{1}{2}Bi)$ only when $\beta$ is negative. When $\beta$ is positive, $\Theta(\frac{1}{2}A + (\frac{1}{2}B - 1)i)$ is to be taken instead.

II. If $m$ lies in the second quadrant, then for $x = 0, y = 0$ one has $\eta = 0$. For this value of $\eta$, $X^{00} = 0, Y^{00} = 0$. This gives

$$\Theta(X^{00} + Y^{00}i) = \begin{array}{ll} 2, & \text{according as } \frac{M}{m} \text{ lies in quadrant 1.} \\ 3, & 2. \\ 0 & 3. \\ 1 & 4., \text{ and } k^{00} = 1 \end{array}$$

III. If $m$ lies in the third quadrant, then for $x = \frac{1}{2}, y = 0$, $\eta = \frac{1}{2}b$ is an integer, for which $X^0 + Y^0 i = \frac{1}{2}A + \frac{1}{2}Bi$. Then put

$$\Theta(X^0 + Y^0 i) = \Theta(\tfrac{1}{2}A + (\tfrac{1}{2}B - 1)i)$$

whenever $\beta$ is negative.

IV. If $m$ lies in the fourth quadrant, then for $\eta = 0$ put $\Theta(X^0 + Y^0 i = 0, 1, 2, 3$, according as $\frac{M}{m}$ lies in quadrant 1. 2. 3. 4.; $k^0 = 0$.

**14.**

The preceding investigations now yield the following determination of the decident.

Collect all values of $x$ and $y$ that lie *within* the bounds 0 and $\frac{1}{2}$ and for which either $\eta$ and $X$ or $\eta$ and $Y$ are integers. For each $x + iy$, use the rules of Article 12 to determine the value of $k$.

Further, collect all values on the boundaries, that is, where either $x = 0$ or $\frac{1}{2}$ while $y$ lies *between* 0 and $\frac{1}{2}$, or $y = 0$ or $= \frac{1}{2}$ while $x$ lies *between* 0 and $\frac{1}{2}$, such that $\eta$ is an integer and $[\xi]$ odd. Determine the corresponding $l$ as follows. Let $\Theta M(x + yi) = \pm\theta$, with the upper sign for even, the lower for odd $\eta$

| | then for $m$ in | | | |
|---|---|---|---|---|
| for | quadrant 1. | quadrant 2. | quadrant 3. | quadrant 4. |
| $y = 0$ | $l = -\theta$ | $l = -\theta$ | $l = +\theta$ | $l = +\theta$ |
| $x = \frac{1}{2}$ | $l = -\theta$ | $l = +\theta$ | $l = +\theta$ | $l = -\theta$ |
| $y = \frac{1}{2}$ | $l = +\theta$ | $l = +\theta$ | $l = -\theta$ | $l = -\theta$ |
| $x = 0$ | $l = +\theta$ | $l = -\theta$ | $l = -\theta$ | $l = +\theta$ |

More briefly,

$$l = \pm\theta,$$

| | |
|---|---|
| the sign is the same as that of $a$ | if $x = 0$ |
| the opposite | if $x = \frac{1}{2}$ |
| the same as that of $b$ | if $y = \frac{1}{2}$ |
| the opposite | if $y = 0$ |

To $\Sigma k + \Sigma l$ one must then add

$$\left.\begin{array}{l}\text{if } m \text{ lies in the second quadrant: } 2, 1, 0, 3 \\ \text{if } m \text{ lies in the fourth quadrant: } 0,\end{array}\right\} \text{ according as } \frac{M}{m} \text{ lies in quadrant } \begin{array}{l}1.\ 2.\ 3.\ 4.\end{array}$$

if $m$ lies in the first quadrant and $\frac{1}{2}(a-1)$ is odd,

$$\begin{array}{ll}\Theta(\frac{1}{2}A + (\frac{1}{2}B - 1)i) & \text{if } \beta \text{ positive} \\ \Theta(\frac{1}{2}A + \frac{1}{2}Bi) & \text{if } \beta \text{ negative}\end{array}$$

if $m$ lies in the third quadrant and $\frac{1}{2}(a-1)$ is odd,

$$\begin{array}{ll}-\Theta(\frac{1}{2}A + \frac{1}{2}Bi) & \text{if } \beta \text{ positive} \\ -\Theta(\frac{1}{2}A + (\frac{1}{2}B - 1)i) & \text{if } \beta \text{ negative.}\end{array}$$

---

## [IV.]

### 1.

Biquadratic residue? $m = a + bi$; $aa + bb = d$

Modulus $M = A + Bi$, $AA + BB = D$

$$\frac{mD}{M} = \mu \qquad \mu = \alpha + \beta i, \quad \alpha = aA + bB, \quad \beta = Ab - Ba$$

$$\xi + \eta i = \pi; \quad \pi m = x + yi = p; \quad \pi M = X + Yi = P$$

Relations

$$\begin{array}{lll} x = a\xi - b\eta & d\xi = ax + by & D\xi = AX + BY \\ y = b\xi + a\eta & d\eta = -bx + ay & D\eta = -BX + AY \\ X = A\xi - B\eta & dX = \alpha x + \beta y & Dx = \alpha X - \beta Y \\ Y = B\xi + A\eta & dY = -\beta x + \alpha y & Dy = \beta X + \alpha Y \end{array}$$

$$\begin{aligned} \beta\xi &= -Bx + bX = Ay - aY \\ \beta\eta &= -Ax + aX = -By + bY \\ \alpha\xi &= Ax + aY = By + aX \\ \alpha\eta &= -Bx + aY = Ay - bX \end{aligned}$$

Those $\pi$ for which $\xi$ and $\eta$ lie between 0 and $\frac{1}{2}$ shall be denoted by $\pi^0$, and the corresponding $p$ and $P$ by $p^0$ and $P^0$; those $\pi$ for which $\eta = 0$ and $\xi$ lies between 0 and $\frac{1}{2}$, by $\pi'$; those for which $\xi = \frac{1}{2}$ and $\eta$ lies between 0 and $\frac{1}{2}$, by $\pi''$; those $\pi$ for which $\eta = \frac{1}{2}$ and $\xi$ lies between 0 and $\frac{1}{2}$, by $\pi'''$; finally, those for which $\xi = 0$ and $\eta$ lies between 0 and $\frac{1}{2}$, by $\pi''''$.

The decident of $\frac{m}{M}$ is found as follows:

Collect all *integer* $P^0$; the corresponding $x^0$ and $y^0$ will therefore be fractional. Let the respective intensors of $p^0$ be $t^0$, that is, the numbers $0, 1.\ 2, 3$, according as

| | | | | |
|---|---|---|---|---|
| $[x^0]$ | even, | odd, | odd, | even |
| $[y^0]$ | even, | even, | odd, | odd |

The required decident is then $= \Sigma \pm t^0$, where the upper sign is to be taken for even $P^0$, the lower for odd.

This is the *first* method.

**2.**

We now wish to arrange the individual $P^0$ according to the values of $Y^0$. Restricting ourselves to the case in which $a, b, A, B$ are positive, the smallest value is given by $Y^0 \ldots +1$, the largest by $\frac{1}{2}(A+B-1)$. For each particular value of $Y^0$, the values of $X^0$ must lie between certain bounds, namely

I. when $A-B$ is positive

| if | between | and |
|---|---|---|
| $Y < \frac{1}{2}B$ | $-\frac{BY^0}{A}$ | $\frac{AY^0}{B}$ |
| $Y = \frac{1}{2}B$ | $-\frac{BB}{2A}$ | $\frac{1}{2}A$ |
| $Y > \frac{1}{2}B$ and $< \frac{1}{2}A$ | $-\frac{BY^0}{A}$ | $-\frac{BY^0}{A} + \frac{D}{2A}$ |
| $Y > \frac{1}{2}A$ | $\frac{AY^0}{B} - \frac{D}{2B^0}$ | $-\frac{BY^0}{A} + \frac{D}{2A}$ |

II. When $A-B$ is negative,

| if | between | and |
|---|---|---|
| $Y < \frac{1}{2}A$ | $-\frac{BY^0}{A}$ | $\frac{AY^0}{B}$ |
| $Y > \frac{1}{2}A$ and $< \frac{1}{2}B$ | $\frac{AY^0}{B} - \frac{D}{2B}$ | $\frac{AY^0}{B}$ |
| $Y = \frac{1}{2}B$ | $\frac{AB-D}{2B}$ | $\frac{1}{2}A$ |
| $Y > \frac{1}{2}B$ | $\frac{AY^0}{B} - \frac{D}{2B}$ | $-\frac{BY^0}{A} + \frac{D}{2A}$ |

At the lower bounds either $\xi = 0$ or $\eta = \frac{1}{2}$; at the upper bounds, on the other hand, either $\eta = 0$ or $\xi = \frac{1}{2}$. It is easy to prove that the bounds of $x$ are never integers.

**3.**

We now wish to express the partial sums for each particular $Y^0$ in another way. At the bounds $p$ will have particular values, which may be denoted by $p^*, p^{**}$, and while $X$ changes continuously from one bound to the other, $p$ will pass continuously from $p^*$ to $p^{**}$. But the integer contained in $[p]$ changes here in jumps, with either the real or the imaginary part always changing by one unit. The changes occur at the values of $X$

$$X',\ X'',\ X'''\ldots X^\mu$$

which are already ordered by size and to which the values of $p$

$$p',\ p''.\ p'''\ldots$$

correspond.

The last $X^\mu$ can also be identical with $X^{**}$ when $\beta$ is positive, or $X'$ identical with $X^*$, and so on.

The $x$ are increasing here; thus, when $x^{**}$ is an integer, it is counted as $x^\mu$.

The $y$ are increasing for positive $\beta$; then $y^{**}$ is fully included in the count;

decreasing for negative $\beta$; then $y^*$ is included in the count.

Let the intensors of $p^*, p^{**}$ be $\lambda^*$ and $\lambda^{**}$

$$\begin{array}{rrl} \text{the intensor from } p' & \text{up to } p''\ldots & \lambda^* + \delta' \\ p'' & \text{up to } p'''\ldots & \lambda^* + \delta' + \delta'' \\ p^\mu & \text{up to } p^{**}\ldots & \lambda^* + \delta' + \delta''\ldots + \delta^\mu = \lambda^{**} \end{array}$$

Then the partial sum, insofar as $Y^0$ is even, becomes

$$\begin{aligned} &= \lambda^*(g-h) + (\lambda^* + \delta')(g'-h') \\ &\quad + (\lambda^* + \delta' + \delta'')(g''-h'') + \text{etc.} + \lambda^{**}(g^\mu - h^\mu) \end{aligned}$$

where $g$ denotes the number of even $X^0$ from $X^*$ to $X'$, and $h$ that of the odd ones.

**4.**

This formula can also be expressed as follows:

$$\begin{aligned}
&\lambda^*\{[\tfrac{1}{2}X^* - \tfrac{1}{2}] - [\tfrac{1}{2}X^*]\}\\
&+\delta'\{[\tfrac{1}{2}X' - \tfrac{1}{2}] - [\tfrac{1}{2}X']\}\\
&+\delta''\{[\tfrac{1}{2}X'' - \tfrac{1}{2}] - [\tfrac{1}{2}X'']\}\\
&+\text{etc.}\\
&+\delta^\mu\{[\tfrac{1}{2}X^\mu - \tfrac{1}{2}] - [\tfrac{1}{2}X^\mu]\}\\
&-\lambda^{**}\{[\tfrac{1}{2}X^{**} - \tfrac{1}{2}] - [\tfrac{1}{2}X^{**}]\}
\end{aligned}$$

or by

$$\lambda^*\varepsilon^* + \delta'\varepsilon' + \delta''\varepsilon'' + \text{etc.} + \delta^\mu\varepsilon^\mu - \lambda^{**}\varepsilon^{**}$$

where in general $\varepsilon = 0$ if $[X]$ odd
and $= -1$ if $[X]$ is even and $Y$ even
$+1$ if $[X]$ even and $Y$ odd.

For $\delta$, on the other hand, the values are

| | | $\beta$ | |
|---|---|---|---|
| | | positive | negative |
| if $x$ an even integer, | $[y]$ even | $-1$ | $-1$ |
| | $[y]$ odd | $+1$ | $+1$ |
| $x$ an odd integer, | $[y]$ even | $+1$ | $+1$ |
| | $[y]$ odd | $-1$ | $-1$ |
| $y$ an even integer, | $[x]$ even | $-3$ | $+3$ |
| | $[x]$ odd | $-1$ | $+1$ |
| $y$ an odd integer, | $[x]$ even | $+3$ | $-3$ |
| | $[x]$ odd | $+1$ | $-1$ |

**5.**

From this we derive the following second method of determining the decident.

I. Collect all combinations of integer values of $Y$ and $x$ having the following properties:

1. that $\xi = \frac{Ax+bY}{\alpha}$ lies between 0 and $\frac{1}{2}$, with the second bound included;
2. that $\eta = \frac{-Bx+aY}{\alpha}$ lies between 0 and $\frac{1}{2}$, with the first bound included.

II. For these calculate

$$X = \frac{Dx + \beta Y}{\alpha}$$
$$y = \frac{\beta x + dY}{\alpha}$$

III. Omit all those for which $[X]$ is odd, and divide the remaining ones, for which $[X]$ is even, into two classes;

in the first class put those for which, simultaneously,

$Y$ even, $x$ even, $[y]$ even
or where one of these conditions holds;

in the second class, those for which two of these conditions, or none at all, hold,

or in I. where $[Y + x + y]$ even
II. where $[Y + x + y]$ odd

and call the excess of the number in the first class over that in the second $c$.

IV. Collect all combinations of integer values of $Y$ and $y$ having the following properties:

1. that $\xi = \frac{Ay-aY}{\beta}$ lies between 0 and $\frac{1}{2}$, with the first bound included when $\beta$ is negative, the second included when $\beta$ is positive;
2. that $\eta = \frac{-By+bY}{\beta}$ lies between 0 and $\frac{1}{2}$, with the first bound included for positive $\beta$, the second for negative.

[V.]

**[1.]**

$$\begin{array}{ll} \text{Modulus} & M = A + Bi \quad AA + BB = D \\ \text{Residue} & m = a + bi, \quad aa + bb = d \end{array}$$

$$\frac{mD}{M} = \mu = \alpha + \beta i = aA + bB + (Ab - Ba)i$$

$$\xi + \eta i = \pi, \quad \pi m = p = x + yi; \quad \pi M = P = X + Yi$$

$\omega$ is an undetermined infinitely small real positive quantity.

**[2.]**

*Preparation.*

I. Collect all $\pi$ for which

$\xi$ nonnegative and no greater than $\frac{1}{2}$
$\eta$ positive and less than $\frac{1}{2}$
Either $x$ or $y$ is an integer
Either $X$ or $Y$ is an integer

and, for each $\pi$, determine the quantity $\varepsilon$ by the following rule:

$$\begin{array}{ll} \text{Let } p^0 & \text{be the nearest integer divisible by } 1+i \text{ to } p \\ P^0 & \text{be the nearest integer divisible by } 1+i \text{ to } P \end{array}$$

and put $\varepsilon = \pm 1$, where the sign is always the same as the sign of the imaginary part of the quantity

$$\frac{p - p^0}{P - P^0}(\alpha - \beta i)$$

The following are the special rules. First class, $x$ and $X$ integers

$$\beta\xi = -Bx + bX$$
$$\beta\eta = -Ax + aX$$
$$\beta y = -\alpha x + dX$$
$$\beta Y = -Dx + \alpha X$$

$\varepsilon = -1$ when $\beta$ is positive, $x$ even, $[y]$ even, $X$ even, $[Y]$ even, or when only an odd number of these conditions holds.

$\varepsilon = +1$ when none or an even number of these conditions holds.

Second class, $y$ and $Y$ integers

$$\begin{aligned}\beta\xi &= +Ay - aY\\ \beta\eta &= -By + bY\\ \beta x &= +\alpha y - dY\\ \beta X &= +Dy - \alpha Y\end{aligned}$$

$\varepsilon = -1$ when $\beta$ is positive, $[x]$ even, $y$ even, $[X]$ even, $Y$ even, or when an odd number of these conditions holds.

$\varepsilon = +1$ when none or an even number holds.

Third class, $y$ and $X$ integers

$$\begin{aligned}\alpha\xi &= +By + aX\\ \alpha\eta &= +Ay - bX\\ \alpha x &= -\beta y + dX\\ \alpha Y &= +Dy - \beta X\end{aligned}$$

$\varepsilon = -1$ when $\alpha$ is positive, $[x], y, X, [Y]$ are all even, or when an odd number of these conditions holds.

$\varepsilon = +1$ when none or an even number holds.

Fourth class, $x$ and $Y$ integers

$$\begin{aligned}\alpha\xi &= +Ax + bY\\ \alpha\eta &= -Bx + aY\\ \alpha y &= +\beta x + dY\\ \alpha X &= +Dx + \beta Y\end{aligned}$$

$\varepsilon = +1$ when $\alpha$ is positive, $x, [y], [X], Y$ are all even, or when an odd number of these conditions holds.

$\varepsilon = -1$ when none or an even number of them holds.

**[3.]**

II. Collect all $\pi$ for which

$\xi$ positive and less than $\frac{1}{2}$
$\eta = \omega$
and either $X$ or $Y$ is an integer,

and put $\varepsilon = \pm 1$, taking the sign of the imaginary part of

$$\frac{M}{P - P^0}$$

Special rule: first class, $X$ an integer

$$\begin{aligned} A\xi &= +X + B\omega \\ Ax &= +aX - \beta\omega \\ Ay &= +bX - \alpha\omega \\ AY &= +BX + D\omega \end{aligned}$$

$\varepsilon = -1$ when $A$ is positive, $X$ and $[Y]$ are even, or when only one condition holds.

$\varepsilon = +1$ when none or two hold.

Second class, $Y$ an integer

$$\begin{aligned} B\xi &= +Y - A\omega \\ Bx &= +aY - \alpha\omega \\ By &= +bY - \beta\omega \\ BX &= +AY - D\omega \end{aligned}$$

$\varepsilon = +1$ when $B$ is positive, $[X]$ and $Y$ are even, or when only one condition holds.

$\varepsilon = -1$ when none or two hold.

**[4.]**

III. Collect all $\pi$ for which

$\xi$ and $\eta$ satisfy the same conditions as in II;

either $x$ or $y$ is an integer,

and put $\varepsilon = \pm 1$ with the sign of the imaginary part of

$$\frac{m}{p - p^0}$$

Special rules: first class, $x$ an integer

$$\begin{aligned} a\xi &= +x + b\omega \\ ay &= +bx + d\omega \\ aX &= +Ax + \beta\omega \\ aY &= +Bx + \alpha\omega \end{aligned}$$

$\varepsilon = -1$ when $a$ is positive, $x, [y]$ are both even, or when only one condition holds.

$\varepsilon = +1$ when none or two hold.

Second class, $y$ an integer

$$\begin{aligned} b\xi &= +y - a\omega \\ bx &= +ay - d\omega \\ bX &= +Ay - \alpha\omega \\ bY &= +By + \beta\omega \end{aligned}$$

$\varepsilon = +1$ when $b$ is positive, $[x]$ and $y$ are even, or when only one condition holds.

$\varepsilon = -1$ when none holds.

**[5.]**

IV. Collect all $\pi$ for which

$$\begin{aligned}&\xi = \tfrac{1}{2} + \tfrac{1}{2}\omega\\ &\eta \text{ positive and less than } \tfrac{1}{2}\\ &X \text{ or } Y \text{ is an integer,}\end{aligned}$$

and put $\varepsilon = \pm 1$ with the sign of the imaginary part of

$$\frac{iM}{P - P^0}$$

Special rules: first class, $X$ an integer,

$$\begin{aligned}2B\eta &= +A - 2X + A\omega\\ 2Bx &= -\beta + 2bX - \beta\omega\\ 2By &= +\alpha - 2aX + \alpha\omega\\ 2BY &= +D - 2AX + D\omega\end{aligned}$$

$\varepsilon = +1$ when $B$ is positive, $X, [Y]$ are even, or when one condition holds,
$\varepsilon = -1$ when none or two conditions hold.
Second class, $Y$ an integer

$$\begin{aligned}2A\eta &= -B + 2Y - B\omega\\ 2Ax &= +\alpha - 2bY + \alpha\omega\\ 2Ay &= +\beta + 2aY + \beta\omega\\ 2AX &= +D - 2BY + D\omega\end{aligned}$$

$\varepsilon = +1$ when $A$ is positive, $[X], Y$ are even, or when one holds;
$\varepsilon = -1$ when none or two conditions hold.

**[6.]**

V. Collect all $\pi$ for which

$$\xi, \eta \text{ satisfy the same conditions as in IV,}$$
$$\text{and where } x \text{ or } y \text{ is an integer,}$$

and put $\varepsilon = \pm 1$ with the sign of the imaginary part of

$$\frac{im}{p - p^0}$$

Special rules: first class, $x$ an integer

$$\begin{aligned} 2b\eta &= +a - 2x + a\omega \\ 2by &= +d - 2ax + d\omega \\ 2bX &= +\beta + 2Bx + \beta\omega \\ 2bY &= +\alpha - 2Ax + \alpha\omega \end{aligned}$$

$\varepsilon = +1$ when $b$ is positive, $x, [y]$ are even, or when one condition holds;
$\varepsilon = -1$ when none or two conditions hold.

Second class, $y$ an integer

$$\begin{aligned} 2a\eta &= -b + 2y - b\omega \\ 2ax &= +d - 2by + d\omega \\ 2aX &= +\alpha - 2By + \alpha\omega \\ 2aY &= -\beta + 2Ay - \beta\omega \end{aligned}$$

$\varepsilon = +1$ when $a$ is positive, $[x], y$ are even, or when one condition holds;
$\varepsilon = -1$ when none or two conditions hold.

**[7.]**

The first method now gives the following result:

$$\begin{aligned} 4 \text{ decident} = \text{I}\quad & \Sigma\varepsilon \text{ from all} \\ & -4\Sigma\varepsilon \text{ from those where } y \text{ integer } [x] \text{ even} \\ \text{II.}\quad & -\Sigma\varepsilon \text{ from all} \\ & +4\Sigma\varepsilon \text{ from those where } [x] \text{ even } [y] \text{ odd} \\ \text{IV.}\quad & +4\Sigma\varepsilon \text{ from those where not both } [x] \text{ and } [y] \text{ even} \\ & +Q \end{aligned}$$

Here

| for $A$ | $B$ | $Q =$ |
|---|---|---|
| $+$ | $+$ | $-\text{Intens.} +\mu\omega i + \text{Int.} \frac{1}{2}m - \mu\omega$ |
| $+$ | $-$ | $-\text{Intens.} +\mu\omega + \text{Int.} \frac{1}{2}mi - \mu\omega i$ |
| $-$ | $-$ | $-\text{Intens.} -\mu\omega i - \text{Int.} \frac{1}{2}m + \mu\omega$ |
| $-$ | $+$ | $-\text{Intens.} -\mu\omega - \text{Int.} \frac{1}{2}mi + \mu\omega i$ |

The following table represents this:

| $\alpha$ | $\beta$ | $A$ | $B$ | $A$ | $B$ | $A$ | $B$ | $A$ | $B$ |
|---|---|---|---|---|---|---|---|---|---|
| | | $+$ | $+$ | $+$ | $-$ | $-$ | $-$ | $-$ | $+$ |
| $+$ | $+$ | $+2$ | $0$ | $0$ | $+2$ | $-3$ | $-5$ | $-3$ | $-5$ |
| $+$ | $-$ | $0$ | $+2$ | $-2$ | $0$ | $-5$ | $-3$ | $-1$ | $-3$ |
| $-$ | $-$ | $-3$ | $-1$ | $-1$ | $+1$ | $-4$ | $-2$ | $0$ | $-2$ |
| $-$ | $+$ | $+1$ | $-1$ | $-1$ | $+1$ | $0$ | $-2$ | $-4$ | $-6$ |
| $\frac{m-1}{2}$ | | even, | odd | | | | | | |

The first part of

$$2Q \ldots -3 - (\alpha\beta AB) + (A\alpha) + (A\beta) + (B\alpha) - (B\beta)$$

The whole $2Q$ for

$$\begin{aligned} \frac{m-1}{2} \text{ odd}\quad & -3 + 4(A) - (B) - (\beta) + (\alpha A) + (\alpha B) \\ & + (\beta A) - (\beta B) - (\alpha\beta AB) \\ \frac{m-1}{2} \text{ even}\quad & -3 + 2(A) + (B) + (\beta) + (\alpha A) + (\alpha B) \\ & + (\beta A) - (\beta B) + 2(\beta AB) - (\alpha\beta AB) \end{aligned}$$

Example

$$\begin{array}{r|l} m = +3-2i & 13 \\ M = -1-6i & 37 \end{array} \qquad \begin{array}{r} (1+i)M + (-2+i)m = 1 \\ mM = -15-16i \end{array}$$

$$\mu = +9+20i$$

$$\begin{array}{r|r|r|r} +4+13i & +38+31i & +1 & -1 \\ +5+7i & +29+11i & 0 & +0 \\ +6+i & +20-9i & 0-i & -3 \\ +10+14i & +58+22i & +1 & +1 \\ +11+8i & +49+2i & +1 & -1 \\ +12+2i & +40-18i & +1-i & +2 \\ +16+15i & +78+13i & +2 & -0 \\ +17+9i & +69-7i & +1-i & +2 \\ +18+3i & +60-27i & +1-i & -2 \\ \hline & & \multicolumn{2}{r}{\text{decident} = -2} \end{array}$$

$$\begin{array}{ll} \text{I}\,. & +1 \quad 0 \\ \text{II}\,. & 0 \quad -4 \\ \text{IV}\,. & \quad 0 \\ Q = & -5 \\ \hline & -8 \end{array}$$

Good.

I

$$\begin{array}{r|r|r|r|r|r|r} x & X & 20\xi & 20\eta & 20y & 20Y & \varepsilon \\ +1 & 0 & 6 & 1 & -9 & -37 & -1 \\ & +1 & 4 & 4 & +4 & -28 & -1 \\ & +2 & 2 & 7 & +17 & -19 & -1 \\ +2 & +1 & 10 & 5 & -5 & -65 & -1 \\ & +2 & 8 & 8 & +8 & -56 & +1 \\ & & & & & & -3 \end{array}$$

$$\begin{array}{r|r|r|r|r|r|r} y & Y & 20\xi & 20\eta & 20x & 20X & \varepsilon \\ 0 & -1 & 3 & 2 & +13 & +9 & +1^* \\ & -2 & 6 & 4 & +26 & +18 & +1 \\ & -3 & 9 & 6 & +39 & +27 & +1 \\ +1 & -1 & 2 & 8 & +22 & +46 & +1 \\ & & & & & & +4 \\ & & & & & & (+1) \end{array}$$

$$\begin{array}{r|r|r|r|r|r|r} y & X & 9\xi & 9\eta & 9x & 9Y & \varepsilon \\ 0 & +1 & 3 & 2 & +13 & -20 & +1 \\ -2 & +2 & 0 & 3 & +6 & -3 & -1^* \\ & & & & & & 0(-1) \end{array}$$

| $x$ | $Y$ | $9\xi$ | $9\eta$ | $9y$ | $9X$ | $\varepsilon$ |
|---|---|---|---|---|---|---|
| $+1$ | $-1$ | $1$ | $3$ | $+7$ | $+17$ | $-1$ |
| $+2$ | $-3$ | $4$ | $3$ | $+1$ | $+14$ | $+1$ |
| | | | | | | $0$ |

I . . . $+1(0)$

II

| $Y$ | $6\xi$ | $6x$ | $6y$ | $6X$ | $\varepsilon$ |
|---|---|---|---|---|---|
| $-1$ | $+1$ | $+3+9\omega$ | $-2$ | $-1$ | $-1^*$ |
| $-2$ | $+2$ | $+6+9\omega$ | $-4$ | $-2$ | $+1$ |
| | | | | | $0(-1)$ |

III

| $x$ | $3\xi$ | $3y$ | $3X$ | $3Y$ | $\varepsilon$ |
|---|---|---|---|---|---|
| $+1$ | $1$ | $-2$ | $-1$ | $-6+9\omega$ | $-1$ |

IV

| $X$ | $12\eta$ | $12x$ | $12y$ | $12Y$ | $\varepsilon$ |
|---|---|---|---|---|---|
| $0$ | $+1$ | $+20+20\omega$ | $-9$ | $-37$ | $-1^*$ |
| $+1$ | $+3$ | $+24+20\omega$ | $-3$ | $-39$ | $+1^*$ |
| $+2$ | $+5$ | $+28+20\omega$ | $+3$ | $-41$ | $-1$ |
| | | | | | $-1(0)$ |

V

| $x$ | $4\eta$ | $4y$ | $4X$ | $4Y$ | $\varepsilon$ |
|---|---|---|---|---|---|
| $+2$ | $+1$ | $-1$ | $+4-20\omega$ | $-13$ | $+1$ |

| $y$ | $6\eta$ | $6x$ | $6X$ | $6Y$ | $\varepsilon$ |
|---|---|---|---|---|---|
| $0$ | $+2$ | $+13$ | $+9$ | $-20$ | $+1$ |

**[8.]**

We now wish to examine the result of I more closely. There are four combinations:
$1^0$, when $x$ and $X$ are integers. Here

$$\beta\xi = -Bx + bX$$
$$\beta\eta = -Ax + aX$$
$$\beta y = -\alpha x + dX$$
$$\beta Y = -Dx + \alpha X$$

Let $y^0$ and $Y^0$ be the absolutely least residues of $-\alpha x + dX$, $-Dx + \alpha X$ modulo $\beta$, and $\beta y = \beta y' + y^0$, $\beta Y = \beta Y' + Y^0$, and put

$$\begin{aligned} \beta u &= -By^0 + bY^0 \qquad & y^0 &= +au - bt \\ \beta t &= -Ay^0 + aY^0 \qquad & Y^0 &= +Au - Bt \end{aligned}$$

Then $t, u$ will be integers, namely

$$\begin{aligned} -u + ti &= M(x + y'i) - m(X + Y'i)\ ^* \\ i(t + ui) &= Mi(y' - y) - mi(Y' - Y) \end{aligned}$$

and then $\varepsilon = -1$ when
$t + u$ is even, $\beta, y^0, Y^0$ positive, or when two or none of the conditions holds; otherwise $\varepsilon = +1$.

We put

$$\begin{aligned} t + ui = +\theta &\quad \text{when } y^0 \text{ and } Y^0 \text{ both positive} \\ -\theta &\quad y^0 \text{ and } Y^0 \text{ both negative} \\ +\theta' &\quad y^0 \text{ positive } Y^0 \text{ negative} \\ -\theta' &\quad y^0 \text{ negative } Y^0 \text{ positive} \end{aligned}$$

| | | | |
|---|---|---|---|
| to each divisible by $1+i$ | $\theta$ | there then corresponds | $\varepsilon = -1$ |
| to each divisible by $1+i$ | $\theta'$ | | $\varepsilon = +1$ |
| to each not divisible by $1+i$ | $\theta$ | | $\varepsilon = +1$ |
| to each not divisible by $1+i$ | $\theta'$ | | $\varepsilon = -1$ |

provided that $\beta$ is positive.

For negative $\beta$ the rule is reversed.

| $x$ | $X$ | $y^0$ | $Y^0$ | $t + ui$ | $\theta$ | $\theta'$ | $\varepsilon$ |
|---|---|---|---|---|---|---|---|
| $+1$ | $0$ | $-9$ | $+3$ | $0 - 3i$ | | $0 + 3i$ | $-1$ |
| | $+1$ | $+4$ | $-8$ | $-1 + 2i$ | | $-1 + 2i$ | $-1$ |
| | $+2$ | $-3$ | $+1$ | $0 - i$ | | $0 + i$ | $-1$ |
| $+2$ | $+1$ | $-5$ | $-5$ | $-1 - i$ | $+1 + i$ | | $-1$ |
| | $+2$ | $+8$ | $+4$ | $+1 + 2i$ | $+1 + 2i$ | | $+1$ |

**[9.]**

$2^0$, when $y$ and $Y$ are integers. Here let $x', X'$ be the nearest integers to $x$ and $X$, and

$$x - x' = \frac{x^0}{\beta}, \quad X - X' = \frac{X^0}{\beta}$$

and put

$$\begin{aligned} \beta(t + ui) &= -Mx^0 + mX^0 = -Mp^0 + mP^0, \\ t + ui &= Mp' - mP' \end{aligned}$$

that is,

$$\beta t = -Ax^0 + aX^0 \qquad x^0 = -bt + au$$
$$\beta u = -Bx^0 + bX^0 \qquad X^0 = -Bt + Au$$

Then

$\varepsilon = -1$, when $\beta$ positive, $x^0$ positive, $X^0$ positive, $t+u$ even
etc.

We put

| | | *better* |
|---|---|---|
| $t+ui = +\theta$ | when $x^0$ and $X^0$ positive | $+\theta''$ |
| $= -\theta$ | when both negative | $-\theta''$ |
| $= +\theta'$ | when $x^0$ positive, $X^0$ negative | $-\theta$ |
| $= -\theta'$ | when $x^0$ negative, $X^0$ positive | $+\theta$ |

where the same rule for $\varepsilon$ will hold as above.

| $y$ | $Y$ | $x^0$ | $X^0$ | $t+ui$ | $\theta$ | $\theta'$ | $\varepsilon$ |
|---|---|---|---|---|---|---|---|
| 0 | $-1$ | $-7$ | $+9$ | $+1-3i$ | | $-1+3i$ | $+1$ |
| | $-2$ | $+6$ | $-2$ | $0+2i$ | | $0+2i$ | $+1$ |
| | $-3$ | $-1$ | $+7$ | $+1-i$ | | $-1+i$ | $+1$ |
| $+1$ | $-1$ | $+2$ | $+6$ | $+1$ | $+1$ | | $+1$ |

It can now be proved

1) That all $\theta$ obtained from (1) and from (2) are distinct from one another. Call their collection $\Theta$.

2) That all $\theta = T + Ui$ have the property that

$$-bT + aU$$
$$-BT + AU$$

are positive numbers less than $\frac{1}{2}\beta$.

3) That if $T, U$ are two integers satisfying the conditions just mentioned, $T + Ui$ is certainly in $\Theta$.

(How, then? It is based on the above equation *.)

4) The same holds in a similar way for $\theta'$, whose collection $\Theta'$ will consist of those numbers $T' + U'i$ for which

$$-bT' + aU'$$
$$-(-BT' + AU')$$

are positive numbers less than $\frac{1}{2}\beta$.

In our case

$$\begin{array}{r|r||r|r||r|r||r|r}
\theta & \varepsilon & \theta' & \varepsilon & \theta'' & \varepsilon & \theta''' & \varepsilon \\
+1 & +1 & 0+i & -1 & +2-i & +1 & +1 & -1 \\
+1+i & -1 & 0+2i & +1 & +3-i & -1 & +2 & +1 \\
+1+2i & +1 & 0+3i & -1 & & & & \\
 & & -1+i & +1 & & & & \\
 & & -1+2i & -1 & & & & \\
 & & -1+3i & +1 & & & &
\end{array}$$

Here

$$\begin{aligned}
\theta &= -_*M + {}_*m \\
\theta' &= -_*M - {}_*m \\
\theta'' &= +_*Mi + {}_*m \\
\theta''' &= +_*Mi - {}_*m
\end{aligned}$$

**[10.]**

$3^0$. $y$ and $X$ integers. Let $x', Y'$ be the nearest integers to $x$ and $Y$, and

$$x' + yi = p', \quad X + Y'i = P'; \quad p - p' = \frac{p^0}{\alpha}, \quad P - P' = \frac{P^0}{\alpha}$$

and put

$$\begin{aligned}
i(t+ui) = Mp' - mP' &= -\frac{Mp^0}{\alpha} + \frac{mP^0}{\alpha} \\
&= -\frac{Mx^0}{\alpha} + \frac{miY^0}{\alpha} \quad \text{that is} \\
\alpha t = -Bx^0 + aY^0 \quad &\text{then} \quad x^0 = -bt + au \\
\alpha u = +Ax^0 + bY^0 \quad &\qquad\quad Y^0 = +At + Bu
\end{aligned}$$

Then

$\varepsilon = -1$, when $\alpha$ positive, $x^0$ positive, $Y^0$ positive, $t+u$ even etc.

We put

| | | *better* |
|---|---|---|
| $t+ui = +\theta''$ | when $x^0$ positive, $Y^0$ positive | $+\theta'$ |
| $= +\theta'''$ | when $x^0$ positive, $Y^0$ negative | $-\theta'''$ |
| $-\theta''$ | when $x^0$ negative, $Y^0$ negative | $-\theta'$ |
| $-\theta'''$ | when $x^0$ negative, $Y^0$ positive | $+\theta'''$ |

Thus, for each $\theta'' \ldots \varepsilon = -1$
$\theta''' \ldots \varepsilon = +1$

provided that $\theta''$ or $\theta'''$ is divisible by $1+i$ and $\alpha$ is positive.

| $y$ | $X$ | $x^0$ | $X^0$ | $t+ui$ | $\theta''$ | $\theta'''$ | $\varepsilon$ |
|---|---|---|---|---|---|---|---|
| 0 | +1 | +4 | −2 | +2 | | +2 | +1 |
| +1 | +2 | −3 | −3 | $-3+i$ | $+3-i$ | | −1 |

**[11.]**

Class 4, $x$ and $Y$ integers. With assumptions similar to those in 3, put

$$-(t+ui) = Mp' - mP' = -\frac{Mp^0}{\alpha} + \frac{mP^0}{\alpha}$$
$$= -\frac{Miy^0}{\alpha} + \frac{mX^0}{\alpha}$$

$$\alpha t = -By^0 - aX^0 \qquad y^0 = -bt + au$$
$$\alpha u = +Ay^0 - bX^0 \qquad X^0 = -At - Bu$$

Then

$\varepsilon = +1$ when $\alpha$ positive, $y^0$ positive, $X^0$ positive, $t+u$ even etc.

We put

$$\begin{aligned} t+ui = +\theta'' &\quad \text{when } y^0 \text{ positive } X^0 \text{ negative} \\ +\theta''' &\quad \text{when } y^0 \text{ positive } X^0 \text{ positive} \\ -\theta'' &\quad \text{when } y^0 \text{ negative } X^0 \text{ positive} \\ -\theta''' &\quad \text{when } y^0 \text{ negative } X^0 \text{ negative} \end{aligned}$$

For $\varepsilon$ the rule then holds (as above in 3), provided that $\alpha$ is positive:

$$\varepsilon = -1 \text{ for each (divisibility by } 1+i\text{):} \begin{cases} \text{divisible} & \theta'' \\ \text{not divisible} & \theta''' \end{cases}$$

$$\varepsilon = +1 \text{ for each (divisibility by } 1-i\text{):} \begin{cases} \text{divisible} & \theta''' \\ \text{not divisible} & \theta'' \end{cases}$$

| $x$ | $Y$ | $y^0$ | $X^0$ | $t+ui$ | $\theta''$ | $\theta'''$ | $\varepsilon$ |
|---|---|---|---|---|---|---|---|
| $+1$ | $-1$ | $-2$ | $-1$ | $-1$ | | $+1$ | $-1$ |
| $+2$ | $-3$ | $+1$ | $-4$ | $+2-i$ | $+2-i$ | | $+1$ |

The collection of all $\theta''$ from 3 and 4, which we denote by $\Theta''$, therefore consists of all numbers $T+Ui$ for which

$$\left.\begin{aligned} -bT + aU \\ \text{and } +AT + BU \end{aligned}\right\} \text{ positive and less than } \frac{1}{2}\alpha$$

the collection of all $\theta''' \ldots$ $(\Theta''')$ of those for which

$$\left.\begin{aligned} -bT + aU \\ -AT - BU \end{aligned}\right\} \text{ positive and less than } \frac{1}{2}\alpha$$

**[12.]**

After the above improvement, the rule therefore reads as follows: Let

$\Theta$ contain all numbers $T+Ui$, where $+bT-aU$ positive $-BT+AU$ positive and $<\frac{1}{2}\beta$

$\Theta'$ contain all numbers $T+Ui$, where $-bT+aU$ positive $+AT+BU$ positive and $<\frac{1}{2}\alpha$


$\Theta''$ contain all numbers $T+Ui$, where $-bT+aU$ positive $-BT+AU$ positive and $<\frac{1}{2}\beta$

$\Theta'''$ contain all numbers $T+Ui$, where $+bT-aU$ positive $+AT+BU$ positive and $<\frac{1}{2}\alpha$

provided that $\beta$ or $\alpha$, respectively, is positive.

| For all divisible by | then | when |
|---|---|---|
| $1+i$ | $\varepsilon =$ | |
| $\theta$ | $+1$ | $\beta$ positive |
| $\theta'$ | $-1$ | $\alpha$ positive |
| $\theta''$ | $-1$ | $\beta$ positive |
| $\theta'''$ | $+1$ | $\alpha$ positive |

| $\Theta$ | | $\Theta'$ | | $\Theta''$ | | $\Theta'''$ | |
|---|---|---|---|---|---|---|---|
| $0-i$ | $-1$ | $+2-i$ | $+1$ | $+1$ | $+1$ | $-1$ | $-1$ |
| $0-2i$ | $+1$ | $+3-i$ | $-1$ | $+1+i$ | $-1$ | $-2$ | $+1$ |
| $0-3i$ | $-1$ | | | $+1+2i$ | $+1$ | | |
| $+1-i$ | $+1$ | | | | | | |
| $+1-2i$ | $-1$ | | | | | | |
| $+1-3i$ | $+1$ | | | | | | |




**[13.]**

This yields the following rule. Let the results of the instructions be

$$\text{II}\ldots\ G, \quad \text{III}\ldots\ g, \quad \text{IV}\ldots\ H, \quad \text{V}\ldots\ h$$

Then

$$\begin{aligned} 4\Theta &= R = 0 \\ 4\Theta' &= -g + G - h - H + R' \\ 4\Theta'' &= -2g + 2G + R'' \\ 4\Theta''' &= -g + G + h + H + R''' \end{aligned}$$

In our example

$$G = 0, \quad g = -1, \quad H = -1, \quad h = +2,$$
$$R' = 0, \quad R'' = +2, \quad R''' = -2$$
$$4\Theta = 0, \quad 4\Theta' = +1 - 1 + 0 = 0,$$
$$4\Theta'' = +2 + 2 = +4, \quad 4\Theta''' = +1 + 1 - 2 = 0$$

and the corrections $R, R'$ etc. are determined as follows:

$$R = \begin{cases} -(\beta) + \frac{1}{2}(\alpha\beta) + \frac{1}{2}(\alpha\beta abAB) \\ -\frac{1}{2}(b) - \frac{1}{2}(B) \\ +(\beta) - \frac{1}{2}(\alpha\beta) - \frac{1}{2}(\alpha\beta abAB) \\ +\frac{1}{2}(b) + \frac{1}{2}(B) \end{cases}$$

$$R' = \begin{cases} +(\alpha) - \frac{1}{2}(\alpha\beta) + \frac{1}{2}(\alpha\beta abAB) \\ -\frac{1}{2}(b) - \frac{1}{2}(A) \\ 0 \\ -\frac{1}{2}(a) + \frac{1}{2}(B) \end{cases}$$

$$R'' = \begin{cases} +(\beta) + \frac{1}{2}(\alpha\beta) + \frac{1}{2}(\alpha\beta abAB) \\ -\frac{1}{2}(b) + \frac{1}{2}(B) \\ +(\beta) + \frac{1}{2}(\alpha\beta) + \frac{1}{2}(\alpha\beta abAB) \\ -\frac{1}{2}(b) + \frac{1}{2}(B) \end{cases}$$

$$R''' = \begin{cases} -(\alpha) - \frac{1}{2}(\alpha\beta) + \frac{1}{2}(\alpha\beta abAB) \\ -\frac{1}{2}(b) + \frac{1}{2}(A) \\ 0 \\ +\frac{1}{2}(a) + \frac{1}{2}(B) \end{cases}$$

where the quantities in parentheses supply only the signs.

Thus

$$R + R' + R'' + R''' = 2(\alpha\beta abAB) - 2(b) + 2(B) + 2(\beta)$$

consequently

$$\begin{aligned} \Theta + \Theta' + \Theta'' + \Theta''' = & -g + G + \tfrac{1}{2}(\alpha\beta abAB) \\ & + \tfrac{1}{2}(\beta) - \tfrac{1}{2}(b) + \tfrac{1}{2}(B) \\ = & -g + G + S \end{aligned}$$

| $ab$ | | + | + | | − | + | | − | − | | + | − | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| $A$ | $B$ | $\alpha$ | $\beta$ | $S$ | $\alpha$ | $\beta$ | $S$ | $\alpha$ | $\beta$ | $S$ | $\alpha$ | $\beta$ | $S$ |
| + | + | + | + | +1 | − | + | +1 | − | − | +1 | + | − | +1 |
| | | + | − | −1 | + | + | 0 | − | + | +1 | − | − | 0 |
| − | + | + | − | 0 | + | + | +1 | − | + | +2 | − | − | +1 |
| | | − | − | −1 | + | − | −1 | + | + | +1 | − | + | +1 |
| − | − | − | − | −1 | + | − | −1 | + | + | +1 | − | + | +1 |
| | | − | + | −1 | − | − | −2 | + | − | −1 | + | + | 0 |
| + | − | − | + | 0 | − | − | −1 | + | − | 0 | + | + | +1 |
| | | + | + | −1 | − | + | −1 | − | − | −1 | + | − | −1 |

**[14.]**

The first rule now accordingly takes the following form:

$4\,\text{Dec.} =$ I. $-4\Sigma\varepsilon$ from those where $y$ integer, $[x]$ even
II. $+4\Sigma\varepsilon$ from those where $[x]$ even, $[y]$ odd
III. $-\Sigma\varepsilon$ from all
IV. $+4\Sigma\varepsilon$ from those where not both $[x]$ and $[y]$ even
$+Q+S$

In our example

| | |
|---|---|
| I.... | 0 |
| II.... | −4 |
| III | +1 |
| IV | 0 |
| $Q$ | −5 |
| $S$ | 0 |
| | −8 |

Now suppose that in III those for which $y$ is an integer, $[x]$ even, are specially marked. Then

$$\left.\begin{array}{l}\text{III. } \Sigma\varepsilon \text{ from all} \\ \quad -4\Sigma\varepsilon \text{ from the specially marked}\end{array}\right\} = \text{Intensor } (\tfrac{1}{2}-\omega)m - \text{Intens. } \omega m = -W$$

Here

| | $a$ | $b$ | | |
|---|---|---|---|---|
| | + | + | −3 | −1 |
| | − | + | −2 | 0 |
| $W=$ | − | − | +2 | 0 |
| | − | + | +3 | +1 |
| | | | $\frac{m-1}{2}$ even | $\frac{m-1}{2}$ odd |

Thus

$$\begin{aligned} 4\,\text{decident} = {} & \text{I. } -4\Sigma\varepsilon \quad y \text{ integer, } [x] \text{ even} \\ & \text{II. } +4\Sigma\varepsilon \quad [x] \text{ even, } [y] \text{ odd} \\ & \text{III. } -4\Sigma\varepsilon \quad y \text{ integer, } [x] \text{ even} \\ & \text{IV. } +4\Sigma\varepsilon \quad \text{all where not both } [x], [y] \text{ even} \\ & +Q+S+W \end{aligned}$$

Table for $\frac{1}{4}(Q+S+W)$

| $a$ | $b$ | $A$ | $B$ | $\alpha$ | $\beta$ | | | $a$ | $b$ | $A$ | $B$ | $\alpha$ | $\beta$ | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| + | + | + | + | + | + | 0 | 0 | − | − | + | + | − | − | 0 | 0 |
| | | | | + | − | −1 | 0 | | | | | − | + | +1 | 0 |
| | | − | + | + | − | −1 | −1 | | | − | + | − | + | 0 | −1 |
| | | | | − | − | −1 | −1 | | | | | + | + | 0 | −1 |
| | | − | − | − | − | −2 | −1 | | | − | − | + | + | 0 | −1 |
| | | | | − | + | −1 | −1 | | | | | + | − | −1 | −1 |
| | | + | − | − | + | −1 | 0 | | | + | − | + | − | 0 | 0 |
| | | | | + | + | −1 | 0 | | | | | − | − | 0 | 0 |
| − | + | + | + | − | + | 0 | 0 | + | − | + | + | + | − | +1 | +1 |
| | | | | + | + | 0 | 0 | | | | | − | − | 0 | 0 |
| | | − | + | + | + | −1 | −1 | | | − | + | − | − | +1 | 0 |
| | | | | + | − | −1 | −1 | | | | | − | + | 0 | −1 |
| | | − | − | + | − | −2 | −1 | | | − | − | − | + | +1 | 0 |
| | | | | − | − | −2 | −1 | | | | | + | + | 0 | −1 |
| | | + | − | − | − | −1 | 0 | | | + | − | + | + | +1 | +1 |
| | | | | − | + | −1 | 0 | | | | | + | − | 0 | 0 |

**[15.]**

The second method is as follows:

$$\begin{aligned}\text{decident} = \text{I. } &+\Sigma\varepsilon, \text{ where } Y \text{ integer, } [X] \text{ even}\\ &-4\Sigma\varepsilon, \text{ among these, where also } y \text{ integer, } [x] \text{ even}\\ \text{II. } &+\Sigma\lambda\varepsilon, \text{ where } Y \text{ integer, } [X] \text{ even;}\\ &\quad \lambda \text{ is the intensor of } p\\ \text{II. } &-\Sigma\lambda'\varepsilon, \text{ where } X \text{ integer, } [Y] \text{ odd}\\ &\quad \lambda' \text{ the intensor of } ip = 1,\ 2,\ 3,\ 0\\ &\qquad \text{when } \lambda = 0,\ 1,\ 2,\ 3\\ \text{IV. } &+\Sigma\lambda\varepsilon, \text{ where } Y \text{ integer, } [X] \text{ even,}\\ &\quad \lambda \text{ the intensor of } p\\ \text{IV. } &+\Sigma\lambda'\varepsilon, \text{ where } X \text{ integer, } [Y] \text{ even}\\ &\quad \lambda' \text{ the intensor of } im - ip = 0 \quad 3 \quad 2 \quad 1\\ &\qquad \text{when Int. } p = 0 \quad 1 \quad 2 \quad 3\\ &+q\end{aligned}$$

Here $q = 0$ when $\frac{M-1}{2}$ is odd, i.e. divisible only by $1+i$, not by 2, and not at the same time $AB + -$; otherwise,

| $A$ | $B$ | | $\alpha$ | $\beta$ | $\alpha$ | $\beta$ | $\alpha$ | $\beta$ | $\alpha$ | $\beta$ |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | + | + | − | + | − | − | + | − |
| + | + | Int. $\frac{1}{2}(m-\omega\mu)$ | +3 | +1 | +3 | +1 | +0 | +2 | +0 | +2 |
| − | + | 0 | 0 | | 0 | | 0 | | 0 | |
| − | − | −Int. $\frac{1}{2}(m+\omega\mu)$ | −0 | −2 | −0 | −2 | −3 | −1 | −3 | −1 |
| + | − | −Int. $\omega\mu$ | −0 | | −1 | | −2 | | −3 | |

Where two numbers are given, the first applies to even $\frac{m-1}{2}$, the other to odd.

In our example:

| I. | $y$ | $Y$ | $20x$ | $20Y$ | $\varepsilon$ | |
|---|---|---|---|---|---|---|
| | 0 | −1 | +13 | +9 | +1* | +3 |
| | 0 | −2 | +26 | +18 | +1 | −4 |
| | +1 | −1 | +22 | +46 | +1 | −1 |
| | | | | | +3(+1) | Dec. = −2 |

II. Absent.

| IV. | $X$ | $12x$ | $12y$ | $12Y$ | $\varepsilon$ | $\lambda$ | $\lambda'$ | $\lambda'\varepsilon$ |
|---|---|---|---|---|---|---|---|---|
| | 0 | $+20+20\omega$ | −9 | −37 | −1 | 2 | 2 | −2 |
| | +1 | $+24+20\omega$ | −3 | −39 | +1 | 3 | 1 | +1 |
| | +2 | $+28+20\omega$ | +3 | −41 | −1 | 0 | 0 | 0 |
| | | | | | | | | −1 |

The contributions from II combine into the following result:

$$\begin{aligned}
\text{II. } &-\Sigma\varepsilon, \text{ where } Y \text{ integer, } [X] \text{ even}\\
&+4\Sigma\varepsilon, \text{ from these same ones, when also } [x] \text{ even,}\\
&\quad [y] \text{ odd}\\
\text{II. } &+\Sigma\lambda'\varepsilon, \text{ where } Y \text{ integer, } [X] \text{ even}\\
&-\Sigma\lambda'\varepsilon, \text{ where } X \text{ integer, } [Y] \text{ odd,}\\
&\quad \lambda' \text{ the intensor of } ip
\end{aligned}$$

The last two parts in turn combine into

$$\begin{aligned}
\text{III. } &+\Sigma\varepsilon, \text{ where } [X] \text{ even, } [Y] \text{ odd}\\
&-4\Sigma\varepsilon, \text{ when also } x \text{ an integer, } [y] \text{ odd}\\
&+r+s
\end{aligned}$$

where $r = 0$ when $AB\ldots.\begin{cases} + & + \\ - & + \\ - & - \end{cases}$

and $r = \text{Int. } \omega mi$ when $AB\,..\,+\ -$

$$s = -\text{Int. } (\tfrac{1}{2}-\omega)mi, \quad \text{when } \frac{M-1}{2} \text{ even and } B \text{ positive,}$$
$$\text{in all other cases } = 0$$

In our example

$$\text{III. Drops out.} \quad r = 0,\ s = 0$$

The contributions from IV combine into the following result:

$$\begin{aligned}
\text{IV. } &+4\Sigma\varepsilon, \text{ where } Y \text{ integer, } [X] \text{ even and not both}\\
&\quad [x] \text{ even, } [y] \text{ even}\\
&-\Sigma\lambda'\varepsilon, \text{ where } Y \text{ integer, } [X] \text{ even}\\
&+\Sigma\lambda'\varepsilon, \text{ where } X \text{ integer, } [Y] \text{ even}\\
&\quad \text{where } \lambda' \text{ Int. of } im-ip
\end{aligned}$$

The last two parts in turn combine into

$$\begin{aligned}
\text{V. } &+\Sigma\varepsilon, \text{ where not both } [X] \text{ and } [Y] \text{ even}\\
&-4\Sigma\varepsilon, \text{ where also } x \text{ an integer, } [y] \text{ even}\\
&+w
\end{aligned}$$

Here $w = 0$ when $\frac{M-1}{2}$ is even and $A$ positive; in all other cases

$$= -\text{Intensor } (\tfrac{1}{2}i+\omega)m$$

In our example

V.

| $y$ | $6x$ | $6X$ | $6Y$ | $\varepsilon$ |
|---|---|---|---|---|
| 0 | +13 | +9 | −20 | +1 |

$$w = -2$$

Table for $q, r, s, w$ and their sum.

| $a$ | $b$ | $A$ | $B$ | $\alpha$ | $\beta$ | $\frac{M-1}{2}$ even, $\frac{m-1}{2}$ even | $\frac{M-1}{2}$ odd, $\frac{m-1}{2}$ even | $\frac{M-1}{2}$ even, $\frac{m-1}{2}$ odd | $\frac{M-1}{2}$ odd, $\frac{m-1}{2}$ odd |
|---|---|---|---|---|---|---|---|---|---|
| + | + | + | + | + | + | +3 0 0 0 +3 | 0 0 0 0 0 | +1 0 −2 0 −1 | 0 0 0 −2 −2 |
| | | | | | − | 0 0 0 0 0 | 0 0 0 0 0 | +2 0 −2 0 0 | 0 0 0 −2 −2 |
| | | | | − | + | 0 0 0 0 0 | 0 0 0 0 0 | 0 0 −2 −2 −4 | 0 0 0 −2 −2 |
| | | | | | − | 0 0 0 0 0 | 0 0 0 0 0 | 0 0 −2 −2 −4 | 0 0 0 −2 −2 |
| | | | − | − | − | −3 0 0 0 −3 | 0 0 0 0 0 | −1 0 0 −2 −3 | 0 0 0 −2 −2 |
| | | | | | + | 0 0 0 0 0 | 0 0 0 0 0 | −2 0 0 −2 −4 | 0 0 0 −2 −2 |
| | | | | + | − | −1 +1 0 0 0 | −1 +1 0 0 0 | −1 +1 0 0 0 | −1 +1 0 −2 −2 |
| | | | | | + | 0 +1 0 0 +1 | 0 +1 0 0 +1 | 0 +1 0 0 +1 | 0 +1 0 −2 −1 |
| | | | + | + | − | +3 0 0 0 +3 | 0 0 0 −1 −1 | +1 0 −2 0 −1 | 0 0 0 −3 −3 |
| | | | | | + | +3 0 0 0 +3 | 0 0 0 −1 −1 | +1 0 −2 0 −1 | 0 0 0 −3 −3 |
| | | | | − | + | 0 0 0 −1 −1 | 0 0 0 −1 −1 | 0 0 −2 −3 −5 | 0 0 0 −3 −3 |
| | | | | | − | 0 0 0 −1 −1 | 0 0 0 −1 −1 | 0 0 −2 −3 −5 | 0 0 0 −3 −3 |
| | | | − | − | + | −3 0 0 −1 −4 | 0 0 0 −1 −1 | −1 0 0 −3 −4 | 0 0 0 −3 −3 |
| | | | | | − | −3 0 0 −1 −4 | 0 0 0 −1 −1 | −1 0 0 −3 −4 | 0 0 0 −3 −3 |
| | | | | + | − | −2 +2 0 0 0 | −2 +2 0 −1 −1 | −2 +2 0 0 0 | −2 +2 0 −3 −3 |
| | | | | | + | −1 +2 0 0 +1 | −1 +2 0 −1 0 | −1 +2 0 0 +1 | −1 +2 0 −3 −2 |
| | | + | + | − | − | 0 0 −1 0 −1 | 0 0 0 −1 −1 | +2 0 −3 0 −1 | 0 0 0 −3 −3 |
| | | | | | + | +3 0 −1 0 +2 | 0 0 0 −1 −1 | +1 0 −3 0 −2 | 0 0 0 −3 −3 |
| | | | | − | + | 0 0 −1 −1 −2 | 0 0 0 −1 −1 | 0 0 −3 −3 −6 | 0 0 0 −3 −3 |
| | | | | | − | 0 0 −1 −1 −2 | 0 0 0 −1 −1 | 0 0 −3 −3 −6 | 0 0 0 −3 −3 |
| | | | | − | − | 0 0 0 −1 −1 | 0 0 0 −1 −1 | −2 0 0 −3 −5 | 0 0 0 −3 −3 |
| | | | | | + | −3 0 0 −1 −4 | 0 0 0 −1 −1 | −1 0 0 −3 −4 | 0 0 0 −3 −3 |
| | | | | + | − | −3 +3 0 0 0 | −3 +3 0 −1 −1 | −3 +3 0 0 0 | −3 +3 0 −3 −3 |
| | | | | | + | −2 +3 0 0 +1 | −2 +3 0 −1 0 | −2 +3 0 0 +1 | −2 +3 0 −3 −2 |
| | | | + | + | − | 0 0 −1 0 −1 | 0 0 0 0 0 | +2 0 −3 0 −1 | 0 0 0 −2 −2 |
| | | | | | + | 0 0 −1 0 −1 | 0 0 0 0 0 | +2 0 −3 0 −1 | 0 0 0 −2 −2 |
| | | | | − | − | 0 0 −1 0 −1 | 0 0 0 0 0 | 0 0 −3 −2 −5 | 0 0 0 −2 −2 |
| | | | | | + | 0 0 −1 0 −1 | 0 0 0 0 0 | 0 0 −3 −2 −5 | 0 0 0 −2 −2 |
| | | | − | − | − | 0 0 0 0 0 | 0 0 0 0 0 | −2 0 0 −2 −4 | 0 0 0 −2 −2 |
| | | | | | + | 0 0 0 0 0 | 0 0 0 0 0 | −2 0 0 −2 −4 | 0 0 0 −2 −2 |
| | | | | + | − | 0 0 0 0 0 | 0 0 0 0 0 | 0 0 0 0 0 | 0 0 0 −2 −2 |
| | | | | | + | −3 0 0 0 −3 | −3 0 0 0 −3 | −3 0 0 0 −3 | −3 0 0 −2 −5 |

**[16.]**

Consequently,

$$\text{Dec. } \frac{m}{M} - \text{Dec. } \frac{M}{m} =$$

I. $-4\Sigma\varepsilon$, where $y, Y$ integer, $[x], [X]$ even

II. $+4\Sigma\varepsilon$, $Y$ integer, $[X]$ even, $[x]$ even, $[y]$ odd

III. $-4\Sigma\varepsilon$, $x$ integer, $[X]$ even, $[Y]$ odd, $[y]$ odd

IV. $+4\Sigma\varepsilon$, $Y$ integer, $[X]$ even, and not both $[x], [y]$ even

V. $-4\Sigma\varepsilon$, $x$ integer, $[y]$ even and not both $[X], [Y]$ even

$+\psi$

Here $\psi$ is represented in the following table:

| $a$ | $b$ | $A$ | $B$ | $\alpha$ | $\beta$ | $\psi$ | $a$ | $b$ | $A$ | $B$ | $\alpha$ | $\beta$ | $\psi$ |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| + | + | + | + | + | + | +4 0 0 −2 | − | − | + | + | − | − | 0 0 0 −2 |
| | | + | + | + | − | 0 0 0 −2 | | | + | + | − | + | +4 0 0 −2 |
| | | − | + | + | − | 0 −4 0 −2 | | | − | + | − | + | 0 −4 0 −2 |
| | | − | + | − | − | 0 −4 0 −2 | | | − | + | + | + | 0 −4 0 −2 |
| | | − | − | − | − | −4 −4 0 −2 | | | − | − | + | + | 0 −4 0 −2 |
| | | − | − | − | + | 0 −4 0 −2 | | | − | − | + | − | −4 −4 0 −2 |
| | | + | − | − | + | 0 0 0 −2 | | | + | − | + | − | 0 0 0 −2 |
| | | + | − | + | + | 0 0 0 −2 | | | + | − | − | − | 0 0 0 −2 |
| − | + | + | + | − | + | +4 0 0 −2 | + | − | + | + | + | − | 0 0 0 −2 |
| | | + | + | + | + | +4 0 0 −2 | | | + | + | − | − | 0 0 0 −2 |
| | | − | + | + | + | 0 −4 0 −2 | | | − | + | − | − | 0 −4 0 −2 |
| | | − | + | + | − | 0 −4 0 −2 | | | − | + | − | + | 0 −4 0 −2 |
| | | − | − | + | − | −4 −4 0 −2 | | | − | − | − | + | 0 −4 0 −2 |
| | | − | − | − | − | −4 −4 0 −2 | | | − | − | + | + | 0 −4 0 −2 |
| | | + | − | − | − | 0 0 0 −2 | | | + | − | + | + | 0 0 0 −2 |
| | | + | − | − | + | 0 0 0 −2 | | | + | − | + | − | −4 −4 −4 −6 |

| | |
|---|---|
| Here the first two columns apply | to $\frac{1}{2}(M-1)$ even |
| the last two | to $\frac{1}{2}(M-1)$ odd |
| the first and third | to $\frac{1}{2}(m-1)$ even |
| the second and fourth | to $\frac{1}{2}(m-1)$ odd |

**[17.]**

The 128 cases distinguished in the above table in determining $\psi$ can be expressed much more briefly as follows:

$$\psi = k + l$$

$k = -4$ when $a, A, \alpha, b, B, \beta$ have, respectively, the signs $+\ +\ +\ -\ -\ -$; otherwise always

$$k = 0$$

| | | | |
|---|---|---|---|
| $\frac{1}{2}(M-1)$ even | $\frac{1}{2}(m-1)$ even | $l = +4$, | when $AB\beta$ positive |
| | | $-4$, | when $AB\beta$ negative |
| | | $0$, | in all other cases |
| $\frac{1}{2}(M-1)$ even | $\frac{1}{2}(m-1)$ odd | $l = -4$, | when $A$ negative |
| | | $0$, | when $A$ positive |
| $\frac{1}{2}(M-1)$ odd | $\frac{1}{2}(m-1)$ even | $l = 0$ | |
| $\frac{1}{2}(M-1)$ odd | $\frac{1}{2}(m-1)$ odd | $l = -2$ | |

It remains to be tried whether it is more advantageous to take $A$ and $a$ positive, but $m \equiv 1$, $M \equiv 1$ only modulo 2 (not modulo $2 + 2i$). The final result must be

$$\text{Dec. } \frac{m}{M} - \text{Dec. } \frac{M}{m} \equiv$$

| | $M \equiv$ | | | |
|---|---|---|---|---|
| $m \equiv$ | $1$ | $1+2i$ | $3$ | $3+2i$ |
| $1$ | 0 | 0 | 0 | 0 |
| $1+2i$ | 0 | 2 | 2 | 0 |
| $3$ | 0 | 2 | 0 | 2 |
| $3+2i$ | 0 | 0 | 2 | 2 |

Everything modulo 4.

[VI.]

# Theory of Biquadratic Residues.

---

**1.**

Let a sequence of complex integers $\varphi, \varphi', \varphi''$, and so on, satisfy two conditions: first, they are all pairwise incongruent modulo $\mu = \alpha + \beta i$, where $\alpha$ and $\beta$ denote real integers; second, every complex integer is congruent to one of them modulo $\mu$. Under this assumption, the collection of numbers $\varphi, \varphi', \varphi''$, and so on, is called a complete residue system for the modulus $\mu$. It has been proved that the number of its members equals the norm of $\mu$, that is, the number $\alpha\alpha + \beta\beta$, which will be denoted by $\nu$.

**2.**

Among the numbers $\varphi, \varphi', \varphi''$, and so on, one is divisible by $\mu$. If it is excluded and the collection of the others is denoted by $\chi$, then $\chi$ forms a complete system of residues not divisible by the modulus, their number being $= \nu - 1$. If we restrict the investigation to odd moduli, $\nu - 1$ is divisible by 4. Furthermore, $f, if, -f, -if$ will then be pairwise incongruent; hence the numbers in $\chi$ congruent, respectively, to $if, -f, -if$ are distinct from one another and from $f$. (Associated and composite numbers.)

This partitions $\chi$ into four groups or partial systems $\kappa, \kappa', \kappa'', \kappa'''$. Place any number from $\chi$, for example $\varphi$, in the group $\kappa$, and the three members of $\chi$ congruent to $i\varphi, -\varphi, -i\varphi$ in the groups $\kappa', \kappa'', \kappa'''$, in that order. After removing these four numbers from $\chi$, again place any one of the remaining numbers in $\kappa$, and the three that depend on it in the same way in $\kappa', \kappa'', \kappa'''$. Continue until the entire system $\chi$ has been distributed. The groups $\kappa, \kappa', \kappa'', \kappa'''$ will be called associated quarter systems. They clearly have the following properties:

1) Each quarter system consists of $\frac{1}{4}(\nu - 1) = \frac{1}{4}(\alpha\alpha + \beta\beta - 1)$ numbers.

2) The characteristic property of a quarter system is that none of its numbers, either itself or multiplied by $i, -1$ or $-i$, is congruent to another number in the same quarter system, whereas every number not divisible by $\mu$, either itself or multiplied by $i, -1$ or $-i$, occurs in it or is congruent to one of its members.

3) Just as multiplying the numbers in $\kappa$ by $i, -1$ and $-i$ gives, respectively, the numbers in $\kappa', \kappa'', \kappa'''$, or numbers congruent to them, multiplying the numbers in $\kappa'$ by those factors reproduces, respectively, the numbers

in $\kappa'', \kappa''', \kappa$; multiplying the numbers in $\kappa''$ similarly reproduces the numbers in $\kappa''', \kappa, \kappa'$; finally, multiplying the numbers in $\kappa'''$ reproduces $\kappa, \kappa', \kappa''$.

For brevity, this mutual dependence of the four quarter systems can be expressed as

$$\begin{aligned} \kappa' &\equiv i\kappa, \\ \kappa'' &\equiv -\kappa \equiv i\kappa', \\ \kappa''' &\equiv i\kappa'' \equiv -\kappa' \equiv -i\kappa. \end{aligned}$$

**3.**

If $m$ denotes a complex integer having no common divisor with $\mu$, and every number of a quarter system $\kappa$ is multiplied by $m$, the products, or any numbers congruent to them, themselves form a quarter system. Likewise, multiplying the numbers of the systems $\kappa', \kappa'', \kappa'''$ gives three further quarter systems associated with the first. The proof is easy to give. Denote these four new systems by $m\kappa, m\kappa', m\kappa'', m\kappa'''$, whether the products themselves or only numbers congruent to them are chosen. In the latter case, the choice can always be made from one of the systems $\kappa, \kappa', \kappa'', \kappa'''$. Thus the system $\chi$ has been partitioned into quarter systems in two different ways, at least generally speaking. Suppose that $m\kappa$ has in common

$$\begin{aligned} \text{with } \kappa \ldots\ldots &\quad \lambda \text{ numbers} \\ \kappa' \ldots\ldots &\quad \lambda' \text{ numbers} \\ \kappa'' \ldots\ldots &\quad \lambda'' \text{ numbers} \\ \kappa''' \ldots\ldots &\quad \lambda''' \text{ numbers} \end{aligned}$$

Then $\kappa'$ with $m\kappa'$, $\kappa''$ with $m\kappa''$, and $\kappa'''$ with $m\kappa'''$ will also have $\lambda$ members in common; $\kappa''$ with $m\kappa'$, $\kappa'''$ with $m\kappa''$, $\kappa$ with $m\kappa'''$, $\lambda'$ members, and so on. Let $\varepsilon$ be the least residue of $\lambda' + 2\lambda'' + 3\lambda'''$ modulo 4, or let $\varepsilon$ be one of the four numbers $0, 1, 2, 3$ according as $\lambda' + 2\lambda'' + 3\lambda'''$ has the form $4n, 4n+1, 4n+2, 4n+3$. I now claim that $\varepsilon$ is independent of the arrangement of the quarter system $\kappa$.

To make the proof easier, I use the following notation. Let $\Pi\psi$ denote the number $0, 1, 2, 3$ according as the number $\psi$, not divisible by $\mu$, occurs (itself or through representation by congruence) in the group $\kappa, \kappa', \kappa'', \kappa'''$. The following consequences are therefore immediate:

I. $\Pi(i\psi) \equiv 1 + \Pi\psi \pmod 4$.

II. The number $i^{-\Pi\psi} \,.\, \psi$ occurs, itself or through representation by congruence, in the group $\kappa$.

III. $\Sigma\Pi m\varphi \equiv \varepsilon, \pmod 4$, when the sum extends over all members $\varphi$ occurring in $\kappa$.

Now let $k$ be another quarter system, consisting of $f, f', f''$, and so on, while $\kappa$ consists of $\varphi, \varphi', \varphi''$, and so on. I assume, as is allowed because the *order* of the members of $\kappa$ is arbitrary, that $f$ is identical with or associated with $\varphi$, $f'$ with $\varphi'$, $f''$ with $\varphi''$, and so on. Let the quarter groups associated with $k$ be $k'(\equiv ik)$, $k''(\equiv -k)$, $k'''(\equiv -ik)$. Furthermore, let the characteristic P have the same meaning for the groups $k, k', k'', k'''$ as $\Pi$ has for $\kappa, \kappa', \kappa'', \kappa'''$, so that $\mathrm{P}\psi = 0, 1, 2, 3$ according as $\psi$ belongs to $k, k', k'', k'''$.

Thus, if the partition of $\chi$ into the quarter systems $k, k', k'', k'''$ is used instead of $\kappa, \kappa', \kappa'', \kappa'''$, the least residue of $\mathrm{P}mf + \mathrm{P}mf' + \mathrm{P}mf'' + \mathrm{P}mf'''$, and so on, and so on, or of $\Sigma\mathrm{P}mf$, modulo 4, takes the place of $\varepsilon$; and it remains to prove that $\Sigma\mathrm{P}mf - \Sigma\Pi m\varphi$ is divisible by 4.

We write this quantity as

$$\begin{array}{r} \mathrm{P}mf + \mathrm{P}mf' + \mathrm{P}mf'' + \mathrm{P}mf''' + \text{and so on} \\ -\mathrm{P}m\varphi - \mathrm{P}m\varphi' - \mathrm{P}m\varphi'' - \mathrm{P}m\varphi'''- \\ +\mathrm{P}m\varphi + \mathrm{P}m\varphi' + \mathrm{P}m\varphi'' + \mathrm{P}m\varphi'''+ \\ -\Pi m\varphi - \Pi m\varphi' - \Pi m\varphi'' - \Pi m\varphi'''- \end{array}$$

Since, by assumption, $f$ and $\varphi$ are congruent or associated, the same holds for $mf$ and $m\psi$, and

$$\left.\begin{array}{r} f \equiv i^{-\mathrm{P}\varphi}\varphi \\ i^{-\mathrm{P}mf}mf \equiv i^{-\mathrm{P}m\varphi}m\varphi \end{array}\right\} \text{ mod. } \mu$$

From this it follows easily that $\mathrm{P}mf - \mathrm{P}m\varphi \equiv -\mathrm{P}\varphi$ (mod. 4) and that the sum of the top two rows $\equiv -\Sigma\mathrm{P}\varphi$. Furthermore, since $\mathrm{P}m\varphi - \Pi m\varphi \equiv \mathrm{P}(i^{-\Pi m\varphi}m\varphi)$, $m\varphi \,.\, i^{-\Pi m\varphi}$ belongs to $\kappa$, and the collection of *all* $m\varphi \,.\, i^{-\Pi m\varphi}$, regardless of order, agrees with all $\varphi$, the sum of *all* $\mathrm{P}(m\varphi \,.\, i^{-\Pi m\varphi})$ will equal the sum of all $\mathrm{P}\varphi$. Thus the sum of the third and fourth rows $\equiv \Sigma\mathrm{P}\varphi$ (mod. 4), and hence the sum of all four rows $\equiv 0$ (mod. 4). Which was to be proved.

Since $\varepsilon$ thus depends only on $m$ and $\mu$, independently of the choice of quarter systems, we shall call $\varepsilon$ the character of the number $m$ in relation to the modulus $\mu$ and denote it by Ch. $m$ (mod. $\mu$). It is easy to see that this only generalizes, or includes, the definition given (Art. ...) for the case in which $\mu$ is prime.

**4.**

I now turn to *specific* arrangements of the quarter systems, and set the modulus, to be denoted by $m$, $= ea + ebi$, so that the positive integer $e$ denotes the greatest real divisor, or the greatest divisor common to the two components of $m$; thus $a, b$ are relatively prime. The quarter system with the simplest arrangement will be the one whose members $x + iy$ are such that $ax + by$ is positive and less than $\frac{1}{2}e(aa + bb)$, while $ay - bx$ is nonnegative and likewise less than $\frac{1}{2}e(aa + bb)$; the latter condition is deliberately expressed so as to include the cases in which $ay - bx = 0$. It is easy to see that there will be $\frac{1}{2}(e-1)$ such cases in all, namely

$$\begin{array}{llll} x & = a, & y & = b \\ x & = 2a, & y & = 2b \\ x & = 3a, & y & = 3b \\ & \text{and so on, up to} & & \\ x & = \frac{1}{2}(e-1)a, & y & = \frac{1}{2}(e-1)b \end{array}$$

and hence none at all if the components of $m$ have no common divisor. Let us call this quarter system $k$, and let $k', k'', k'''$ be those obtained by multiplying the numbers belonging to $k$ by $i, -1, -i$; or one may also put

$$k' = m + ik, \quad k'' = (1+i)m - k, \quad k''' = im - ik$$

This gives the following rule for deciding whether any given integer $x + iy$ not divisible by $m$ is congruent to a member of $k, k', k''$ or $k'''$. Namely, $2(ax + by)$ can be put in the form $Pe(aa + bb) + Q$, and $2(ay - bx)$ in the form $Re(aa + bb) + S$, with $P, Q, R, S$ real integers satisfying the following conditions:

if $x + iy$ is congruent to a number from

| so that | $k$ | $k'$ | $k''$ | $k'''$ |
|---|---|---|---|---|
| $P$ | even | odd | odd | even |
| $R$ | even | even | odd | odd |
| $Q$ | positive | positive | positive | nonnegative |
| $e(aa+bb) - Q$ | positive | nonnegative | positive | positive |
| $S$ | nonnegative | positive | positive | positive |
| $e(aa+bb) - S$ | positive | positive | nonnegative | positive |

The matter is easier to survey if the cases in which neither of the numbers $ax + by$, $ay - bx$ is divisible by $e(aa + bb)$ are distinguished from the rest.

I. In the first case, simply take for $P$ the (algebraically) smaller of the two integers between which $\frac{2(ax+by)}{e(aa+bb)}$ lies strictly, and likewise for $R$ the smaller of the two between which $\frac{2(ay-bx)}{e(aa+bb)}$ lies.

II. If $ax + by$ is divisible by $e(aa + bb)$, then $ay - bx$ will be divisible by $aa + bb$, but not by $e(aa + bb)$ (since otherwise $x + iy$ would be divisible by $ea + ebi$). If now $R$, that is, the integer immediately below $\frac{2(ay-bx)}{e(aa+bb)}$, is even, then $x + iy$ will be congruent modulo $ea + ebi$ to a number from $k'$, but to one from $k'''$ if $R$ is odd.

III. If $ay + bx$ is divisible by $e(aa + bb)$, but $ax + by$ is not, then $x + iy$ will be congruent to a number from $k$ or from $k''$, according as the integer algebraically immediately below $\frac{2(ax+by)}{e(aa+bb)}$ is even or odd.

**5.**

The following method for determining the character of a given integer $M = A + Bi$ relative to the odd modulus $m = ea + ebi$ that does not divide it is readily derived from the above.

For brevity we use the following notation. If $p$ denotes any nonintegral real number, let $[p]$ denote the integer that makes both $p - [p]$ and $1 + [p] - p$ positive. This definition therefore excludes applying the notation to integers. If the definition required neither $p - [p]$ nor $1 + [p] - p$ to be negative, the symbol $[p]$ would be ambiguous when $p$ is an integer. One could also, as was done in an earlier paper, require $1 + [p] - p$ to be positive and $p - [p]$ merely to be nonnegative. For our purpose it is somewhat more convenient to keep to the first definition.

Accordingly, the quarter system $k$ consists of all integers $f$ for which, on putting $\frac{2f}{m} = \xi + \eta i$, $\xi$ lies strictly between 0 and 1, while $\eta$ lies between 0 and 1, with 0 included; or

$$\begin{aligned} & [\xi] = 0, \quad [\eta] = 0 \\ \text{or} \quad & [\xi] = 0, \quad \eta = 0 \end{aligned}$$

If now, for every $f$, we put $\frac{2fM}{m} = \xi + i\eta$ and take

$\Psi f = 0$ if both $[\xi]$ even and either $[\eta]$ even or $\eta$ integer
1 $[\eta]$ even and either $[\xi]$ odd or $\xi$ integer
2 $[\xi]$ odd and either $[\eta]$ odd or $\eta$ integer
3 $[\eta]$ odd and either $[\xi]$ even or $\xi$ integer

or, in tabular form,

| | $[\eta]$ even | $[\eta]$ odd | $\eta$ integer |
|---|---|---|---|
| $[\xi]$ even | 0 | 3 | 0 |
| $[\xi]$ odd | 1 | 2 | 2 |
| $\xi$ integer | 1 | 3 | − . |

which may be denoted by $\nabla(\xi+i\eta)$, then the required character of the number $M$ relative to the modulus $m$ equals the least residue of $\Sigma\Psi f$ modulo 4.

**6.**

The rule given in the preceding article is general; for the case in which $M$ is odd, however, we shall give it another form. At the same time we shall assume that the real parts of $m$ and $M$ are odd, and hence their imaginary parts even.

To each number $f$ belonging to the quarter system $k$ we assign another number $g$, derived from $f$ as follows. On putting $\frac{2f}{m} = \xi + i\eta$, four cases are distinguished:

I. If $[2\xi] = 0$ and either $[2\eta] = 0$ or $\eta = 0$
II. If $[2\xi] = 1$ and either $[2\eta] = 0$ or $\eta = 0$
III. If $[2\xi] = 0$ and $[2\eta] = 1$
IV. If $[2\xi] = 1$ and $[2\eta] = 1$

In the first case put $g = 2f$, in the second $g = im - 2if$, in the third $g = m + 2if$, and in the fourth $g = (1+i)m - 2f$. It is easy to see that the collection of all $g$ forms a complete quarter system $l$; its defining property is that, on putting $\frac{g}{m} = \xi + i\eta$, the following conditions hold together:

| | | | |
|---|---|---|---|
| either | $[\xi] = 0,$ | $[\eta] = 0$ | |
| or | $\eta = 0,$ | $[\xi] = 0$ | and $g$ divisible by $1+i$ |
| or | $\xi = 0,$ | $[\eta] = 0$ | and $g$ not divisible by $1+i$ |

It follows that $l$ differs from $k$ only in that the numbers in $k$ for which $\eta = 0$ and which are not divisible by $1+i$, namely

$$a+bi, \quad 3(a+bi), \quad 5(a+bi) \ldots \frac{e-3}{2}(a+bi)$$
$$\text{or up to } \frac{e-1}{2}(a+bi)$$

according as $e$ has the form $4n+1$ or $4n+3$, are absent from $l$, while their products by $i$ occur in the latter collection instead. At the same time one sees that, for $e = 1$, that is, when $m$ is divisible by no real integer, $k$ and $l$ are entirely the same.

It now remains to derive $\Psi f$ directly from the $g$ corresponding to $f$. The result is that, for the above 4 cases,

I. $\Psi f = \nabla \frac{gM}{m}$

II. $\Psi f = \begin{cases} -\nabla \frac{gM}{m} & \text{if neither the real nor imaginary part of } \frac{gM}{m} \text{ is an integer} \\ 1 - \nabla \frac{gM}{m} & \text{if either is an integer} \end{cases}$

III. as II. IV. as I.

## Remarks.

Judging by the position of the manuscripts in a notebook, the fragments designated I and II in this printed edition date from 1811 or the time immediately following. Of the preceding attempts to prove the fundamental theorem for biquadratic residues by the methods applied here to the residue $1 + i$, a record survives that settles the special case of the residue $1 + 2i$ and starts from the determination of the biquadratic character appended as a note to Article 6 of Fragment III. The manuscripts also supplement the historical information published by Gauss in the notices of his arithmetical papers: the theorems included in Articles 15 to 20 of the *Theoria residuorum biquadrat.* were written down before the *Theoria motus corporum coel.* was worked out. The investigations of cubic residues mentioned in those notices were probably never worked up in full; what is recorded consists of proofs of the reciprocity theorems for two primes, one of them real, carried out with the methods provided by the paper *Disquisitionum circa aequationes puras ulterior evolutio.*

Fragments III to VI form separate booklets in the manuscripts. For the first three, the handwriting points to a date not far from that of Fragments I and II, while a considerably later date must be assumed for the last, No. VI.

[I.] Art. 10. The number of lattice points in $(z, z', z'+\zeta, z+\zeta)$ is determined by the theorem: if $a$ and $b$ are relatively prime, the straight line drawn from $\xi + \eta i$ to $\xi + \eta i + a + bi$ passes through one lattice point when the imaginary part of $(\xi + \eta i) \,.\, (-a + bi)$ is an integer.

[I.] Art. 17. The first transformation of the last $S$ in the expression for $\Delta S$ is obtained by dividing the region in question into three parts such that this $S$ becomes

$$\begin{aligned} &S\big(\tfrac{1}{2}m + \tfrac{1}{2} - \tfrac{1}{2}i,\ (\tfrac{1}{2} + \tfrac{1}{2}i)m,\\ &\qquad (\tfrac{1}{2} + \tfrac{1}{2}i)m + i,\ \tfrac{1}{2}m + \tfrac{1}{2} + \tfrac{1}{2}i_{(+)}\big)\\ &- S\big(\tfrac{1}{2}m,\ \tfrac{1}{2}m + \tfrac{1}{2} + \tfrac{1}{2}i_{(+)},\ \tfrac{1}{2}m + \tfrac{1}{2} - \tfrac{1}{2}i\big)\\ &- S\big((\tfrac{1}{2} + \tfrac{1}{2}i)m,\ \tfrac{1}{2}m,\ \tfrac{1}{2}m + \tfrac{1}{2} - \tfrac{1}{2}i\big) \end{aligned}$$

then counting the lattice points in the first part by means of the theorem in Article 10, and further noting that the second part contains no lattice point.

The second transformation is obtained by multiplying the quantities corresponding to the vertices of the third part by $i$ and increasing them by the integer $(1 - i)\frac{m-1}{2}$; finally, the third is obtained by comparing the resulting figure, according to the rule of Article 16, with the figure displaced from it by $\frac{-1}{1+i}$.

[I.] Art. [18.] An explanation of the first scheme is found in Articles 1 to 6 of the fragment written later and designated here by [II.].

[I.] Art. [18.] (2.) The geometric interpretation, with the aid of the two systems of lattice points

$$[-2iQ - \tfrac{1}{2}, \tfrac{1}{2}, iQ] = [-2iQ + \tfrac{1}{2}, \tfrac{1}{2}, iQ] = \mathrm{IV}^*$$

and

$$-[-2iQ + \tfrac{1}{2}i, -\tfrac{1}{2}i, iQ] = -[-2iQ - \tfrac{1}{2}i, -\tfrac{1}{2}i, iQ] = \mathrm{X}^*$$

gives the equations

$$\begin{aligned} \mathrm{IV} - \mathrm{IV}^* + [-2iQ, \tfrac{1}{2}i, -\tfrac{1}{2}] &= -\mathrm{X} + \mathrm{X}^* + [-iQ, \tfrac{1}{2}i, -\tfrac{1}{2}] \\ \mathrm{XIII} + \mathrm{IV}^* &= [-2iQ, 1, iQ] = -I(-2iQ) + I(-iQ) \\ \mathrm{V} + \mathrm{X}^* &= -[-2iQ, -i, iQ] = R(-2iQ) - R(-iQ) \end{aligned}$$

where, in general, $Rx$ and $Ix$ denote the greatest integers contained in the real part and in the coefficient of the imaginary part of $x$, respectively.

But $[-2iQ, \frac{1}{2}i, -\frac{1}{2}]$ is the number of lattice points in the square whose centre is at $-2iQ - \frac{1}{4} + \frac{1}{4}i$ and whose endpoints have the displacements $\frac{1}{2}$ and $\frac{1}{2}i$ between them.

$[-iQ, \frac{1}{2}i, -\frac{1}{2}]$ or $[-2iQ, -\frac{1}{2}i, \frac{1}{2}]$ is the number of lattice points in an equal square with centre $-2iQ + \frac{1}{4} - \frac{1}{4}i$.

[I.] Art. [18.] (3.) With the aid of the lattice points $[0, \frac{1}{2}i, -iQ] = -[-\frac{1}{2}i, -\frac{1}{2}i, -iQ] = \mathrm{I}^*$ one obtains

$$\begin{aligned} \mathrm{VII} - \mathrm{XII} &= \mathrm{I} - \mathrm{I}^* + [-iQ, \tfrac{1}{2}, \tfrac{1}{2}i] - [0, \tfrac{1}{2}, \tfrac{1}{2}i] \\ &= \mathrm{I} - \mathrm{I}^* + [-2iQ, \tfrac{1}{2}, \tfrac{1}{2}i] \\ \mathrm{II} - \mathrm{I}^* &= [0, -i, -iQ] = -R0 + R(-iQ) \\ \mathrm{IX} - \mathrm{I} &= [0, -1, -iQ] = I0 - I(-iQ) \\ \mathrm{VI} - \mathrm{XII} &= -[\tfrac{1}{2}i, -1, -iQ] = -I(\tfrac{1}{2}i) + I(-iQ + \tfrac{1}{2}i) \\ \mathrm{VIII} - \mathrm{XIV} &= -[\tfrac{1}{2}, -1 - i, \tfrac{1}{2}im_{(-)}] = \frac{a-1}{2} + \frac{b}{2} \end{aligned}$$

[II.] Art. 10. We have

$$\begin{aligned} \Upsilon P' &\equiv -\Upsilon(-iP' + \tfrac{1}{2}im), \\ \Upsilon P'' &\equiv -1 - \Upsilon(P'' - \tfrac{1}{2}im), \\ \Upsilon P''' &\equiv 1 + \Upsilon(-iP''') \pmod 4 \end{aligned}$$

and $-iP' + \frac{1}{2}im$, $P'' - \frac{1}{2}im$ are the lattice points in I and VI, respectively, increased by $\frac{1}{2}i$.

[III.] Art. 6. I have added to the present paper the rules given in the note for determining $\mathrm{Dec.}\frac{M}{m}$, taking them from another place in the manuscripts.

The first of these two rules, which, as is easy to see, agrees with the second, follows from that of Article 6, because

$$k \equiv f \,.\, i^{-n} \text{ (mod. } m), \quad n = \Theta\frac{2fM}{m}, \quad p = mm',$$
$$\left[\frac{2km'}{p}\right]^2 \equiv \Theta\frac{2km'}{p} \text{ (mod. } 2+2i)$$

holds.

[III.] Art. 8 contains in the manuscript an example for Article 7, namely the determination of the decident of $-1+2i$ for the modulus $-11+4i$.

[III.] Art. 10. For the remark 'to be expressed differently', compare Article 3 of the following fragment [IV].

[IV.] Articles 1. 2. 4 contain in the manuscript, besides what is printed here, the application to the two examples for $m = 5+8i$, $M = 9+4i$ and for $m = 9+4i$, $M = 5+8i$. [V.] Art. [7.] Here, as in Article 1 of the preceding fragment [IV], Dec.$\frac{m}{M}$ denotes the value of

$$\Sigma(-1)^X \Sigma(-1)^Y \text{Int.}\, p$$

where the summation extends over all integers $X$ and $Y$ for which the corresponding $\xi$ and $\eta$ lie within the bounds 0 and $\frac{1}{2}$.

The text states that the formulas for the decident in Articles 7 and 15 were found in two distinct ways. To keep these explanations from becoming too long, however, they are derived here from a common source.

For any fixed integer value of $X$, let $Y^*$ be the smaller and $Y^{**}$ the larger of the two values of $Y$ corresponding to the boundary values of $\xi$, $\eta$. Let the values of $p$ corresponding to $Y^*$ and $Y^{**}$ be $p^*$ and $p^{**}$. Like $Y^*$ and $Y^{**}$, they cannot be equal to one another, since the sum $\Sigma$ does not extend over the boundary values of $\xi$ and $\eta$.

Sum over all values of $Y$ occurring for the same $X$, in the same way as in the preceding two papers [III] and [IV]. Put $[\frac{1}{2}Y'' - \frac{1}{2}] - [\frac{1}{2}Y' - \frac{1}{2}]$ for the number of odd integers between $Y'$ and $Y''$, and add the intensors referring to the bounds $\xi = 0$ and $= \frac{1}{2}$ twice, with opposite signs. This gives for $\Sigma(-1)^Y \text{Int.}\, p$ the expression consisting of seven parts

$$-\Sigma[-\text{Int.}(p - \mu\omega i) + \text{Int.}(p + \mu\omega i)]$$
$$\text{taking all } p \text{ for which } [Y] \text{ even,}$$
$$x \text{ or } y \text{ integer, incl. } \xi = 0 \text{ and } \tfrac{1}{2}, \quad \text{excl. } \eta = 0 \text{ and } \tfrac{1}{2}$$

$$\left.\begin{array}{l} -\text{Int.}(p^* - \mu\omega i) \text{ if } [Y^*] \text{ even} \\ +\text{Int.}(p^{**} + \mu\omega i) \text{ if } [Y^{**}] \text{ even} \end{array}\right\} \; \xi = 0 \text{ or } \tfrac{1}{2}, \quad 0 < \eta < \tfrac{1}{2}$$

$$\left.\begin{array}{l} -\text{Int.}(p^* + \mu\omega i) \text{ if } [Y^*] \text{ even} \\ +\text{Int.}(p^{**} - \mu\omega i) \text{ if } [Y^{**}] \text{ even} \end{array}\right\} \; 0 < \xi < \tfrac{1}{2}, \quad \eta = 0 \text{ or } \tfrac{1}{2}$$

$$\left.\begin{array}{l} -\text{Int.}(p^* + \mu\omega i) \text{ if } [Y^*] \text{ even} \\ +\text{Int.}(p^{**} - \mu\omega i) \text{ if } [Y^{**} - \omega] \text{ even} \end{array}\right\} \; \xi = 0 \text{ or } \tfrac{1}{2}, \quad \eta = 0 \text{ or } \tfrac{1}{2}$$

which, multiplied by $(-1)^X$ and summed over all integer values of $X$, gives the decident $\frac{m}{M}$.

In this way, the first part of the expression gives, from the $\varepsilon$ formed according to Article 2, Rule I,

$$\begin{array}{c} \Sigma\varepsilon \text{ where } X \text{ integer, } [Y] \text{ even} \\ -\Sigma\varepsilon \text{ where also } y \text{ integer, } [x] \text{ even} \end{array}$$

For the following parts one may put

$$\left.\begin{array}{l} -(B)\text{Int.}(p - B\mu\omega i) \text{ if } \xi = 0 \\ +(B)\text{Int.}(p + B\mu\omega i) \text{ if } \xi = \frac{1}{2} \end{array}\right\} \; X \text{ integer, } [Y] \text{ even, } 0 < \eta < \tfrac{1}{2}$$

$$\left.\begin{array}{l} -(A)\text{Int.}(p + A\mu\omega i) \text{ if } \eta = 0 \\ +(A)\text{Int.}(p - A\mu\omega i) \text{ if } \eta = \frac{1}{2} \end{array}\right\} \; X \text{ integer, } [Y] \text{ even, } 0 < \xi < \tfrac{1}{2}$$

$$\begin{array}{c} -\frac{1}{2}[(A') + (B')]\text{Int.}(p + A'\mu\omega i) \\ \text{if } X \text{ integer, } [Y + A'\omega] \text{ even, } \xi \text{ and } \eta = 0 \text{ or } \frac{1}{2} \end{array}$$

where $A' = +A$ or $-A$ when $\eta = 0$ or $\frac{1}{2}$, and $B' = +B$ or $-B$ when $\xi = 0$ or $\frac{1}{2}$; here, for example, the notation $(A) : +1$ or $-1$ is used according as $A$ is positive or negative.

Multiply by $(-1)^X$ and sum over $X$, making the following substitutions in these expressions, namely in the

$$\begin{array}{rl} \text{first} & p - B\mu\omega i,\ P - BD\omega i,\ \pi - AB\omega i - BB\omega,\ X,\ [Y] \\ & \text{respectively into } ip,\ iP,\ i\pi,\ -Y,\ [X] \\ \text{second} & p + B\mu\omega i,\ P + BD\omega i,\ \pi + AB\omega i + BB\omega,\ X,\ [Y] \\ & \ldots \quad p,\ P,\ \pi \quad X,\ [Y] \\ \text{third} & p + A\mu\omega i,\ P + AD\omega i,\ \pi + AA\omega i + AB\omega,\ X,\ [Y] \\ & \ldots \quad p,\ P,\ \pi \quad X,\ [Y] \\ \text{fourth} & p - A\mu\omega i,\ P - AD\omega i,\ \pi - AA\omega i - AB\omega,\ X,\ [Y] \\ & \ldots \text{ into } im - ip,\ iM - iP,\ i - i\pi,\ Y - B,\ A - 1 - [X] \end{array}$$

and denote the result from the fifth expression by $Q_1$. One then obtains

$$
\begin{aligned}
&- \Sigma(-1)^{Y}(B)\text{Int.}\, ip, \\
&\qquad \text{where } Y \text{ integer, } [X] \text{ even, } \eta = \omega,\ 0 < \xi < \tfrac{1}{2} \\
&+ \Sigma(-1)^{X}(B)\text{Int.}\, p, \\
&\qquad \text{where } X \text{ integer, } [Y] \text{ even, } \xi = \tfrac{1}{2} + \omega,\ 0 < \eta < \tfrac{1}{2} \\
&- \Sigma(-1)^{X}(A)\text{Int.}\, p, \\
&\qquad \text{where } X \text{ integer, } [Y] \text{ even, } \eta = \omega,\ 0 < \xi < \tfrac{1}{2} \\
&+ \Sigma(-1)^{Y}(A)\text{Int.}(im - ip), \\
&\qquad \text{where } Y \text{ integer, } [X] \text{ even, } \xi = \tfrac{1}{2} + \omega,\ 0 < \eta < \tfrac{1}{2} \\
&+ Q_1
\end{aligned}
$$

Examining the individual cases shows that, under the assumption $M \equiv 1 \pmod{2 + 2i}$,

$$
\begin{aligned}
Q_1 = \ & - \text{Int.}\, \mu\omega i \text{ if } M \text{ in quadrant 1} \\
& - \text{Int.}(\tfrac{1}{2}mi + \mu\omega i), \quad \text{if } M \text{ in quadrant 2 and } \frac{M-1}{2} \text{ even} \\
& + \text{Int.}(\tfrac{1}{2}mi - \mu\omega i), \quad \text{if } M \text{ in quadrant 4 and } \frac{M-1}{2} \text{ even}
\end{aligned}
$$

where a complex number is called even or odd according as it is divisible by 2 or not.

Thus, using the $\varepsilon$ specified in Rules II and IV, one obtains

$$
\begin{aligned}
\text{Dec.}\frac{m}{M} = \text{I}, \quad & \Sigma\varepsilon, \text{ where } X \text{ integer, } [Y] \text{ even} \\
& - 4\Sigma\varepsilon, \text{ where also } y \text{ integer, } [x] \text{ even} \\
\text{II}, \quad & - \Sigma\varepsilon\text{Int.}\, ip, \text{ where } Y \text{ integer, } [X] \text{ even} \\
\text{IV}, \quad & + \Sigma\varepsilon\text{Int.}\, p, \text{ where } X \text{ integer, } [Y] \text{ even} \\
\text{II}, \quad & + \Sigma\varepsilon\text{Int.}\, p, \text{ where } X \text{ integer, } [Y] \text{ even} \\
\text{IV}, \quad & + \Sigma\varepsilon\text{Int.}(im - ip), \text{ where } Y \text{ integer, } [X] \text{ even} \\
& + Q_1
\end{aligned}
$$

The expression for the decident takes another form if one first sums over $X$, expressing the number of odd integers between $X'$ and $X''$ by $[\frac{1}{2}X''+\frac{1}{2}]-[\frac{1}{2}X'+\frac{1}{2}]$, namely

$$\begin{aligned}
\text{Dec.}\frac{m}{M} = \text{I}, \quad & \Sigma\varepsilon, \text{ where } Y \text{ integer, } [X] \text{ odd} \\
& -4\Sigma\varepsilon, \text{ where also } y \text{ integer, } [x] \text{ even} \\
\text{II}, \quad & -\Sigma\varepsilon\text{Int.}\, ip, \text{ where } X \text{ integer, } [Y] \text{ even} \\
\text{IV}, \quad & +\Sigma\varepsilon\text{Int.}\, p, \text{ where } Y \text{ integer, } [X] \text{ odd} \\
\text{II}, \quad & +\Sigma\varepsilon\text{Int.}\, p, \text{ where } Y \text{ integer, } [X] \text{ odd} \\
\text{IV}, \quad & +\Sigma\varepsilon\text{Int.}(im-ip), \text{ where } X \text{ integer, } [Y] \text{ odd} \\
& +Q_2
\end{aligned}$$

$$\begin{aligned}
Q_2 = & -\text{Int.}(-\mu\omega), \quad \text{if } M \text{ in quadrant 2} \\
& +\text{Int.}(\tfrac{1}{2}m-\mu\omega), \quad \text{if } M \text{ in quadrant 1 and } \frac{M-1}{2} \text{ odd} \\
& -\text{Int.}(\tfrac{1}{2}m+\mu\omega), \quad \text{if } M \text{ in quadrant 3 and } \frac{M-1}{2} \text{ odd}
\end{aligned}$$

If one sums over $Y$ first, but uses the second method just applied for determining the number of odd integers between two values, one obtains

$$\begin{aligned}
\text{Dec.}\frac{m}{M} = \text{I}, \quad & \Sigma\varepsilon, \text{ where } X \text{ integer, } [Y] \text{ odd} \\
& -4\Sigma\varepsilon, \text{ where also } y \text{ integer, } [x] \text{ even} \\
\text{II}, \quad & -\Sigma\varepsilon\text{Int.}\, ip, \text{ where } Y \text{ integer, } [X] \text{ odd} \\
\text{IV}, \quad & +\Sigma\varepsilon\text{Int.}\, p, \text{ where } X \text{ integer, } [Y] \text{ odd} \\
\text{II}, \quad & +\Sigma\varepsilon\text{Int.}\, p, \text{ where } X \text{ integer, } [Y] \text{ odd} \\
\text{IV}, \quad & +\Sigma\varepsilon\text{Int.}(im-ip), \text{ where } Y \text{ integer, } [X] \text{ odd} \\
& +Q_3
\end{aligned}$$

$$\begin{aligned}
Q_3 = & -\text{Int.}(-\mu\omega i), \quad \text{if } M \text{ in quadrant 3} \\
& -\text{Int.}(\tfrac{1}{2}mi+\mu\omega i), \quad \text{if } M \text{ in quadrant 2 and } \frac{M-1}{2} \text{ odd} \\
& +\text{Int.}(\tfrac{1}{2}mi-\mu\omega i), \quad \text{if } M \text{ in quadrant 4 and } \frac{M-1}{2} \text{ odd}
\end{aligned}$$

If one also sums over $X$ first and uses the first method for expressing the number of odd integers between two values, one obtains the form for the

decident given in [V], Article 15, where the quantity $q$ can also be defined by the following equation:

$$\begin{aligned} q = & -\text{Int.}\,\mu\omega, \quad \text{if } M \text{ in quadrant 4} \\ & +\text{Int.}(\tfrac{1}{2}m - \mu\omega), \quad \text{if } M \text{ in quadrant 1 and } \frac{M-1}{2} \text{ even} \\ & -\text{Int.}(\tfrac{1}{2}m + \mu\omega), \quad \text{if } M \text{ in quadrant 3 and } \frac{M-1}{2} \text{ even} \end{aligned}$$

Combining these four expressions for the decident gives the result stated in [V], Article 7, since $\text{Int.}\,ip - \text{Int.}\,p$ equals 3 when $[x]$ is even and $[y]$ odd, and otherwise equals 1; furthermore, $\text{Int.}(im - ip) + \text{Int.}\,p$ equals 0 when $[x]$ and $[y]$ are even, and equals 4 in all other cases.

[V.] Art. [7.] In the first column, the first table for the example gives the values of $\frac{37.P}{M}$ or $37(\xi + \eta i)$ corresponding to each integer $P$, when $0 < \xi < \frac{1}{2}$, $0 < \eta < \frac{1}{2}$; in the second, $\frac{37.Pm}{M}$ or $37\,.\,p$; in the third, the greatest integer contained in $p$; and in the fourth, $\pm\text{Int.}\,p$, where the upper sign applies when $P$ is divisible by $1 + i$, and the lower when $P$ is not divisible by $1 + i$.

[V.] Art. [9.]. [12.] The improved notation for the $\theta$ is indicated only for the second and third classes, Articles 9. 10, but is also to be extended to the first and fourth, Articles 8. 11. Accordingly, a $\theta^\lambda = T + Ui$ has the index $\lambda, = 0, 1, 2$ or 3 for which the quantities $\varphi^0$ and $\Phi^0$, determined by the equations

$$\begin{aligned} & i^\lambda M = \mathfrak{A} + \mathfrak{B}i, \quad i^{-\lambda}\mu = \rho + \sigma i \\ & \sigma\varphi^0 = -\text{Coeff. Imag. } \theta^\lambda(a - bi) = +bT - aU \\ & \sigma\Phi^0 = +\text{Coeff. Imag. } \theta^\lambda(\mathfrak{A} - \mathfrak{B}i) = -\mathfrak{B}T + \mathfrak{A}U \end{aligned}$$

lie between 0 and $\frac{1}{2}$.

To prove, following the indications in Article 9 (3), that if $T$, $U$ are two real integers satisfying the conditions just stated, then $T + Ui$ also belongs to the collection $\theta^\lambda$ determined in one of the four combinations, Articles 8. 11, let $\varphi'$, $\Phi'$ denote the complex integers for which the equation

$$T + Ui = \varphi' i^\lambda M + \Phi' m$$

holds, and for which one of the four quantities $\pm\frac{\varphi' - \varphi^0}{m}$, $\pm i\frac{\varphi' - \varphi^0}{m}$ has its real part and the coefficient of its imaginary part between 0 and $\frac{1}{2}$ (*Theoria residuorum biquadr. artt.* 45, 46). As is readily seen by examining the sixteen individual cases in the four combinations, the quantity in question is then $\frac{p}{m}$, and the corresponding quantity among $\pm\frac{\Phi' - \Phi^0}{m}$, $\pm i\frac{\Phi' - \Phi^0}{M}$ is $\frac{P}{M}$, since $\frac{\Phi' - \Phi^0}{M} = -\frac{\varphi' - \varphi^0}{m} i^\lambda$.

This way of representing the quantities $\frac{p}{m}$ or $\frac{P}{M}$ also shows that the sum I, $\Sigma\varepsilon$, over all, equals $\Theta + \Theta' + \Theta'' + \Theta'''$, where $\Theta^\lambda$ denotes the sum of

the $\varepsilon$ which, for each lattice point $\theta$ inside the parallelogram $0,\ \frac{1}{2}m,\ \frac{1}{2}m + \frac{1}{2}i^\lambda M,\ \frac{1}{2}i^\lambda M$, are to be put

$= +1$ if $\theta$ is divisible by $1+i$ and Coeff. Imag. $\mu i^{-\lambda}$ is positive, or if neither condition holds; but

$= -1$ if only one holds.

[V.] Art. 13. The quantity $\Theta^\lambda$ can be determined either by the four different ways of summing used above for Dec.$\frac{m}{M}$, or, what is essentially the same, by the methods indicated in [II], Article 11. These are to be based on the four constructions obtained by joining the points whose $\theta$ is a multiple of $1+i$ to the points $\theta+1$, $\theta+i$, $\theta-1$ and $\theta-i$, respectively.

On the boundary of the parallelogram just mentioned, let a $\theta$ correspond to every point for which the real or imaginary part of $\theta$ is an integer. Denote by $\theta^0$ the nearest integer divisible by $1+i$ to $\theta$, and by $l$ the displacement from a point on the straight boundary segment containing $\theta$ to any following point on the same straight line; thus, for example, for this parallelogram, the quantities $m,\ Mi^\lambda,\ -m,\ -Mi^\lambda$, in order. If one puts

$$\varepsilon = \pm 1 \text{ with the sign of the imaginary part of } \frac{l}{\theta - \theta^0}$$

then combining the four results obtained in either way gives $4\Theta^\lambda = -\Sigma\varepsilon$.

One avoids determining the $\theta$ and $\varepsilon$ at the vertices separately by replacing this parallelogram with another whose boundaries are infinitely near those of the former and which does not include the two points 0 and $\frac{1}{2}m + \frac{1}{2}i^\lambda M$. The boundary of such a parallelogram is obtained by placing it on the positive sides of the lines

$$0 \ldots . \tfrac{1}{2}m, \quad \tfrac{1}{2}m + \tfrac{1}{2}i^\lambda M \ldots . \tfrac{1}{2}m,$$
$$\tfrac{1}{2}m + \tfrac{1}{2}i^\lambda M \ldots . \tfrac{1}{2}i^\lambda M, \quad 0 \ldots . \tfrac{1}{2}i^\lambda M$$

Assign the infinitely small positive quantities $\omega_1, \omega_2, \omega_3, \omega_4$ to the four resulting straight lines, in order. For the points $\theta$ on them one may then put

$$\begin{aligned}
\theta = p &= m(\xi + \omega_1 i),\\
-\theta i^\lambda + \tfrac{1}{2}mi^\lambda + \tfrac{1}{2}M = P &= M(\xi + \omega_2 i),\\
-\theta + \tfrac{1}{2}m + \tfrac{1}{2}Mi^\lambda = p &= m(\xi + \omega_3 i),\\
\theta i^{-\lambda} = P &= M(\xi + \omega_4 i) \quad \text{if } \lambda \text{ even}
\end{aligned}$$

$$\begin{aligned}
\theta = p &= m(\xi + \omega_1 i),\\
\theta i^{1-\lambda} - \tfrac{1}{2}mi^{1-\lambda} + \tfrac{1}{2}M = P &= M(\tfrac{1}{2} + \omega_2 + \eta i),\\
\theta i + \tfrac{1}{2}m - \tfrac{1}{2}Mi^{1+\lambda} = p &= m(\tfrac{1}{2} + \omega_3 + \eta i),\\
\theta i^{-\lambda} = P &= M(\xi + \omega_4 i) \quad \text{if } \lambda \text{ odd}
\end{aligned}$$

where, in part, $\xi$ and $\eta$ also extend beyond the range of real values from 0 to $\frac{1}{2}$ by infinitely small quantities, to close the figure.

Let $G$, $g$, $H$, $h$ denote the sums of the $\varepsilon$ formed according to Rules II, III, IV, V, respectively (in Articles 3 to 6). Let $G'$ or $G_{\prime}$, and $g'$ or $g_{\prime}$, include the $\varepsilon$ arising for the infinitely small values of $\xi$ that may occur in the second parallelogram, but otherwise formed according to Rules II and III, respectively. Further, let $G''$ or $G_{\prime\prime}$, and $g''$ or $g_{\prime\prime}$, similarly refer to the same rules, but to the infinitely small values of $\frac{1}{2} - \xi$; and finally let $H', h_{\prime}, H'', h_{\prime\prime}$ refer to Rules IV, V, IV, V and the infinitely small values $\frac{1}{2} - \eta, \frac{1}{2} - \eta, \eta, \eta$, respectively. Then

$$\begin{aligned} 4\Theta^{\lambda} = & -(g + g' + g'') - i^{\lambda}(G + G' + G'') \\ & + i^{\lambda}(g + g_{\prime} + g_{\prime\prime}) + (G + G_{\prime} + G_{\prime\prime}) \quad \text{if } \lambda \text{ even} \\ 4\Theta^{\lambda} = & -(g + g' + g'') - i^{1-\lambda}(H + H' + H'') \\ & - i^{\lambda-1}(h + h_{\prime} + h_{\prime\prime}) + (G + G_{\prime} + G_{\prime\prime}) \quad \text{if } \lambda \text{ odd} \end{aligned}$$

For the vertex $\theta$ of the parallelogram nearest to 0, let $\xi_1$ denote the corresponding value of the $\xi$ of the first side, and $\xi_4$ the corresponding value of the $\xi$ of the fourth side, so that

$$\theta = m(\xi_1 + \omega_1 i) = i^{\lambda} M(\xi_4 + \omega_4 i)$$

Then the $\xi$ corresponding to a point $p$ on the first side or its extension with real part equal to 0 is given by

$$(\text{Real. } p = 0), \quad \xi - \xi_1 = \sigma a \mathfrak{A} \omega_1 - \sigma \omega_4$$

where the positive factors of the infinitely small positive quantities have been replaced by unity, and $\sigma$, $\mathfrak{A}$ denote the real quantities determined by

$$\rho + \sigma i = i^{-\lambda}(\alpha + \beta i), \quad \mathfrak{A} + \mathfrak{B} i = i^{\lambda}(A + Bi)$$

This point $p$ lies on the first side itself if $\xi - \xi_1$ is positive; hence, taking $\omega_1$ infinitely small compared with $\omega_4$, if $\sigma$ is negative. The point $p^0$ nearest to $p$ whose representing number is divisible by $1 + i$ is 0; thus $\text{Imag.}\frac{m}{p - p^0}$ or $\text{Imag.}\frac{1}{\xi + \omega_1 i}$ has the minus sign. Therefore, for $\text{Real. } p = 0$, one obtains

$$\varepsilon = -1 \text{ if } (\sigma) = -1, \quad \varepsilon = 0 \text{ if } (\sigma) = +1, \quad \text{i.e. } \varepsilon = -\tfrac{1}{2} + \tfrac{1}{2}(\sigma)$$

and in the same way, for $\text{Imag. } p = 0$,

$$\xi - \xi_1 = \sigma b \mathfrak{B} \omega_1 - \sigma \omega_4, \quad \text{Imag.}\frac{m}{p - p^0} = \text{Imag.}\frac{1}{\xi + \omega_1 i},$$
$$\varepsilon = -\tfrac{1}{2} + \tfrac{1}{2}(\sigma) \quad \text{thus } g' = -1 + (\sigma)$$

For the fourth side, $P^0 = 0$, $\text{Imag.}\frac{M}{P-P^0} = \text{Imag.}\frac{1}{\xi+\omega_4 i}$,

$$\begin{aligned}
\text{thus for Real.}(i^\lambda P) = 0; \quad & \xi - \xi_4 = -\sigma a\mathfrak{A}\omega_4 + \sigma\omega_1, \\
& \varepsilon = -\tfrac{1}{2} + \tfrac{1}{2}(\sigma a\mathfrak{A}) \\
\text{and for Imag.}(i^\lambda P) = 0; \quad & \xi - \xi_4 = -\sigma b\mathfrak{B}\omega_4 + \sigma\omega_1, \\
& \varepsilon = -\tfrac{1}{2} + \tfrac{1}{2}(\sigma b\mathfrak{B})
\end{aligned}$$

and consequently $G_{\prime} = -1 + \frac{1}{2}(\sigma a\mathfrak{A}) + \frac{1}{2}(\sigma b\mathfrak{B})$, or, since $\rho = a\mathfrak{A} + b\mathfrak{B}$, $G_{\prime} = -1 + \frac{1}{2}(\rho\sigma) + \frac{1}{2}(\rho\sigma ab\mathfrak{A}\mathfrak{B})$.

The part $R_1^\lambda$ of $4\Theta^\lambda$ arising from the portion of the boundary infinitely near 0 is therefore

$$R_1^\lambda = +G_{\prime} - g' = -(\sigma) + \tfrac{1}{2}(\rho\sigma) + \tfrac{1}{2}(\rho\sigma ab\mathfrak{A}\mathfrak{B})$$

Similar considerations give, for the parts $R_2^\lambda$, $R_3^\lambda$, $R_4^\lambda$, which have the same relations to the points $\frac{1}{2}m$, $\frac{1}{2}m + \frac{1}{2}Mi^\lambda$, $\frac{1}{2}Mi^\lambda$, respectively, as $R_1^\lambda$ has to 0, when $\lambda$ is even,

$$\begin{aligned}
R_2^\lambda &= -g'' - i^\lambda G'' = -\tfrac{1}{2}(b) - \tfrac{1}{2}(\mathfrak{B}), \\
R_3^\lambda &= -i^\lambda G' + i^\lambda g_{\prime} = i^\lambda(\sigma) - \tfrac{1}{2}i^\lambda(\rho\sigma) - \tfrac{1}{2}i^\lambda(\rho\sigma ab\mathfrak{A}\mathfrak{B}) \\
R_4^\lambda &= +i^\lambda g_{\prime\prime} + G_{\prime\prime} = \tfrac{1}{2}i^\lambda(b) + \tfrac{1}{2}i^\lambda(\mathfrak{B})
\end{aligned}$$

and when $\lambda$ is odd,

$$\begin{aligned}
R_2^\lambda &= -g'' - i^{1-\lambda}H'' = -\tfrac{1}{2}(b) - \tfrac{1}{2}(\mathfrak{B}), \\
R_3^\lambda &= -i^{1-\lambda}H' - i^{\lambda-1}h_{\prime} = 0, \\
R_4^\lambda &= -i^{\lambda-1}h_{\prime\prime} + G_{\prime\prime} = \tfrac{1}{2}i^{\lambda+1}(a) + \tfrac{1}{2}i^{\lambda+1}(\mathfrak{A})
\end{aligned}$$

[V.] Art. [14.] The sums of the $\varepsilon$ formed according to Rule III are evaluated by means of the equation

$$\begin{aligned}
&\text{III,} \quad \Sigma\varepsilon \text{ from all } -4\Sigma\varepsilon \text{ from those where } y \text{ integer, } [x] \text{ even} \\
&= \text{III,} \quad \Sigma[-\text{Int.}(p - m\omega) + \text{Int.}(p + m\omega)]
\end{aligned}$$

which is readily verified from the definition of the $\varepsilon$. Here $p$ takes all values satisfying the conditions stated under Rule III. Except for the two intensors corresponding to the smallest ($\xi^*$) and the greatest admissible value ($\xi^{**}$) of $\xi$, respectively, which equal

$$\begin{gathered}
-\text{Int.}(p^* - m\omega) \text{ and } +\text{Int.}(p^{**} + m\omega) \text{ or} \\
-\text{Int.}(\omega m) \text{ and } +\text{Int.}(\tfrac{1}{2} - \omega)m
\end{gathered}$$

these intensors can always be paired as $+\text{Int.}(p' + m\omega)$ and $-\text{Int.}(p'' - m\omega)$, so that between $\xi'$ and $\xi''$, corresponding to $p'$ and $p''$, there is no value of

$\xi$ making the real or imaginary part of $p$ an integer. Thus the two intensors always cancel one another.

[V.] Art. [15.] We have

$$\begin{aligned}
\text{II}, \quad & + \Sigma\varepsilon \text{Int.}\, ip \text{ where } Y \text{ integer } [X] \text{ even}, \\
& - \Sigma\varepsilon \text{Int.}\, ip \text{ where } X \text{ integer } [Y] \text{ odd} \\
= \Sigma[ & -\text{Int.}\, i(p - m\omega) + \text{Int.}\, i(p + m\omega)] \\
& \text{for those } p, \text{ for which } x \text{ or } y \text{ integer, } [X] \text{ even} \\
& Y \text{ odd, } 0 < \xi < \tfrac{1}{2}, \quad \eta = \omega \\
& + \text{Int.}\, i(p^* - m\omega) \text{ if } [X^*] \text{ even } [Y^*] \text{ odd} \\
& - \text{Int.}\, i(p^{**} + m\omega) \text{ if } [X^{**}] \text{ even } [Y^{**}] \text{ odd}
\end{aligned}$$

as one readily verifies by performing the summation on the second side of the equation, by the method used in the preceding note, over each interval from $\xi_\prime$ to $\xi_{\prime\prime}$ so small that between $\xi_\prime$ and $\xi_{\prime\prime}$ there is no $\xi$ making the real or imaginary part of the corresponding $P$ an integer. Using the $\varepsilon$ formed according to Rule III gives the second side of this equation the form stated in Article 15.

[V.] Art. [15.] The sums of the $\lambda'\varepsilon$ formed according to Rule IV are transformed into the sums of the $\varepsilon$ from V by considerations of the same kind as in the last note, also using the equation

$$\begin{aligned}
\text{V}, \quad & \Sigma\varepsilon \text{ from all}, \ -4\Sigma\varepsilon \text{ where } x \text{ integer } [y] \text{ even}, \\
= & +\text{Int.}(\tfrac{1}{2}im + m\omega) - \text{Int.}(\tfrac{1}{2}im + \tfrac{1}{2} - \omega m)
\end{aligned}$$

which corresponds to the preceding evaluation of the sum of the $\varepsilon$ in Rule III.

[V.] Art. [17.] Define the auxiliary quantities $U$, $T$, $L$, $V$ by the equations

$$\begin{aligned}
U &= 1 - 2(B) - (AB) + (\alpha a) + (\beta b) + (\alpha b) - (\beta a) \\
&\quad + (\alpha\beta ab) - (\alpha\beta abAB) \quad \text{or} \\
U &= (1 + (A))(1 - (B))(1 - (a\beta) + (b\beta) + (\alpha\beta))
\end{aligned}$$

$$\begin{gathered}
\text{since } (\alpha a) + (\beta b) = (A) + (A\alpha\beta ab), \\
(\alpha b) - (\beta a) = (B) - (B\alpha\beta ab),
\end{gathered}$$

$$\begin{aligned}
T &= -2 + (a) + (b) - (\beta) - 2(B) - (\alpha ab) \quad \text{or} \\
T &= -2 + (a) + (b) - (\beta) - 2(B) - (aA) + (bB) - (\beta AB) \\
L &= (1 + (A))(1 + (B))(1 + (\beta)) - (1 - (A))(1 - (B))(1 - (\beta)) \\
V &= (1 + (A))(1 - (B))(-2(a) + 2(\beta) + (ab) + (\alpha\beta)) \quad \text{or} \\
V &= (1 + (A))(1 - (B))(-(a) + (b) + (\beta) + (ab) - (\beta ab) + (\alpha\beta))
\end{aligned}$$

$$\text{since } (a) + (b) = (\beta) + (\beta ab) \text{ if } A \text{ positive } B \text{ negative}$$

and denote by $W'$, $S'$, $Q'$ the quantities into which $W$, $S$, $Q$ in the expression for Dec.$\frac{m}{M}$ in Article 14 pass when $m$ and $M$, and hence $\alpha+\beta i$ and $\alpha-\beta i$, are interchanged. Then

$$\begin{aligned}
2W' &= -5(B)-(AB) \quad \text{if } \frac{M-1}{2} \text{ even}\\
2W' &= -(B)-(AB) \quad \text{if } \frac{M-1}{2} \text{ odd}\\
2S' &= -(\alpha\beta abAB)-(\beta)-(B)+(b)\\
2Q'+2S'+2W' &= 2T+U \quad \text{if } \frac{M-1}{2} \text{ even}\\
2Q'+2S'+2W' &= -4+4(a)+U \quad \text{if } \frac{M-1}{2} \text{ odd}\\
8\psi &= 8(q+r+s+w)-(2Q'+2S'+2W')
\end{aligned}$$

If one replaces $8(q+r+s+w)$ here by its value stated in Article 15, but writes it in the form

$$\begin{array}{llll}
2T+4L+V & \text{if } \frac{M-1}{2} \text{ even} & \frac{m-1}{2} \text{ even}\\
2T-16+16(A)+V & \text{if } \frac{M-1}{2} \text{ even} & \frac{m-1}{2} \text{ odd}\\
-4+4(a)+V & \text{if } \frac{M-1}{2} \text{ odd} & \frac{m-1}{2} \text{ even}\\
-20+4(a)+V & \text{if } \frac{M-1}{2} \text{ odd} & \frac{m-1}{2} \text{ odd}
\end{array}$$

and notes that

$$V-U=-\tfrac{1}{2}(1+(a))(1+(A))(1+(\alpha))(1-(b))(1-(B))(1-(\beta))$$

holds, one obtains for $\psi$ the method of determination given in Article 17.

[VI.] Art. 3. The incomplete reference may be taken to refer to Article 4 of Fragment III.

Schering.

# ON THE THEORY OF COMPLEX NUMBERS.

---

## [I.]

## NEW THEORY OF THE DECOMPOSITION OF CUBES.

I. We assume that the equation $x^3 + y^3 + z^3 = 0$ has a solution, namely $x = a,\ y = b,\ z = c$, where $a, b, c$ have no common divisor and are therefore also pairwise coprime. We set

$$\begin{aligned} b + c &= \alpha \\ c + a &= \beta \\ a + b &= \gamma \end{aligned}$$

where $\alpha, \beta, \gamma$ must also be pairwise coprime. For if $\alpha$ and $\beta$ had a common divisor, it would also divide $a^3$ and $b^3$, so $a$ and $b$ would have to have a common divisor too.

We now have

$$(\beta + \gamma - \alpha)^3 + (\gamma + \alpha - \beta)^3 + (\alpha + \beta - \gamma)^3 = 0$$

but the identity

$$\begin{aligned} &(\beta + \gamma - \alpha)^3 + (\gamma + \alpha - \beta)^3 + (\alpha + \beta - \gamma)^3 \\ &= (\alpha + \beta + \gamma)^3 - 24\alpha\beta\gamma \end{aligned}$$

holds. Consequently,

$$(\alpha + \beta + \gamma)^3 = 24\alpha\beta\gamma$$

If $\alpha, \beta, \gamma$ are real numbers, then $\alpha + \beta + \gamma$ will be divisible by 3, hence $\alpha+\beta+\gamma)^3$ by 27, and therefore $\alpha\beta\gamma$ by 9. Thus one of the numbers $\alpha, \beta, \gamma$, for example $\gamma$, must be divisible by 9, and so must $c^3$; consequently $c$ is divisible by 3.

If, however, $\alpha, \beta, \gamma$ are imaginary numbers, we conclude that $\alpha + \beta + \gamma$ must be divisible by $1 - \varepsilon$, hence $24\alpha\beta\gamma$ by $(1 - \varepsilon)^3$, and therefore $\alpha\beta\gamma$ by $1 - \varepsilon$. Thus one of the numbers $\alpha, \beta, \gamma$ is divisible by $1 - \varepsilon$, and consequently so is one of the numbers $a, b, c$.

II. In general, we have the identity

$$\begin{aligned} &(p + q + r)^3 + (p + q\varepsilon + r\varepsilon\varepsilon)^8 + (p + q\varepsilon\varepsilon + r\varepsilon)^3 \\ &= 27pqr + 3(p + q + r)(p + q\varepsilon + r\varepsilon\varepsilon)(p + q\varepsilon\varepsilon + r\varepsilon) \end{aligned}$$

Consequently, if $p+q+r=0$, then

$$(p+q\varepsilon+r\varepsilon\varepsilon)^3+(p+q\varepsilon\varepsilon+r\varepsilon)^3-27pqr=0$$

If $p,q,r$, are themselves cubes, namely $=a^3,b^3,c^3$ respectively, that is, if the equation $x^3+y^3+z^3=0$ has a solution, then, on setting

$$\begin{aligned} a^3+b^3\varepsilon+c^3\varepsilon\varepsilon &= a' \\ a^3+b^3\varepsilon\varepsilon+c^3\varepsilon &= b' \\ -3abc &= c' \end{aligned}$$

we also have $a'^3+b'^3+c'^3=0$. From this new solution a third can be derived in the same way, and so on. It is easy to see that if the first solution consists of real numbers, the third will too.

It should also be noted that if $a,b,c$ have no common factor, the same will hold for $a',b',c'$, apart from the factor $1-\varepsilon$. For

$$\begin{aligned} \frac{a'}{1-\varepsilon} &= -\varepsilon\varepsilon a^3+\varepsilon b^3=a^2-\varepsilon c^3=-b^3+\varepsilon\varepsilon c^3 \\ \frac{b'}{1-\varepsilon} &= a^3-\varepsilon b^3=-\varepsilon\varepsilon a^3+\varepsilon c^3=\varepsilon\varepsilon b^3-c^3 \\ \frac{c'}{1-\varepsilon} &= (\varepsilon\varepsilon-1)abc \end{aligned}$$

The first two numbers therefore have no factor in common with $a$, $b$, or $c$, and cannot be divisible by $1-\varepsilon$ either, unless $a,b,c$ are all divisible by $1-\varepsilon$; thus they also have no factor in common with the third.

III. But the reverse path is also open. We have seen that one of the quantities is divisible by $1-\varepsilon$; let it be $c$. Since we can substitute $a\varepsilon$ or $a\varepsilon\varepsilon$ for $a$, and likewise $b\varepsilon$ or $b\varepsilon\varepsilon$ for $b$, we may assume that $a$ is either $\equiv 1$ or $\equiv -1$. We shall assume the former, since in the other case $b\equiv 1$ would hold, and it would only need to be interchanged with $a$. Accordingly, we set

$$\begin{aligned} a &= 1+3\alpha \\ b &= -1+3\beta \end{aligned}$$

and

$$\begin{aligned} \frac{a\varepsilon+b\varepsilon\varepsilon}{\varepsilon-\varepsilon\varepsilon} &= 1+(\varepsilon\varepsilon-\varepsilon)(\alpha\varepsilon+\beta\varepsilon\varepsilon)=A \\ \frac{a\varepsilon\varepsilon+b\varepsilon}{\varepsilon-\varepsilon\varepsilon} &= -1+(\varepsilon\varepsilon-\varepsilon)(\alpha\varepsilon\varepsilon+\beta\varepsilon)=B \\ \frac{a+b}{\varepsilon-\varepsilon\varepsilon} &= (\varepsilon\varepsilon-\varepsilon)(\alpha+\beta)=C \end{aligned}$$

where $A+B+C=0$, and $ABC=\frac{a^3+b^3}{(\varepsilon-\varepsilon\varepsilon)^3}=\left(\frac{c}{\varepsilon\varepsilon-\varepsilon}\right)^3$

Since here

$$a = -\varepsilon A + \varepsilon\varepsilon B$$
$$b = \varepsilon\varepsilon A - \varepsilon B$$

$A$ and $B$ cannot have a common factor, because it would otherwise also be a common factor of $a$ and $b$. Since $A + B + C = 0$, $C$ consequently cannot have a factor in common with either $A$ or $B$. It follows easily that $A$ and $B$, and hence also $C$, are cubes. For $\left(\frac{c}{\varepsilon\varepsilon-\varepsilon}\right)^3$ will be divisible by $\varepsilon - \varepsilon\varepsilon$, hence also by $(\varepsilon - \varepsilon\varepsilon)^3$, or $\alpha + \beta$ by 3; therefore $A \equiv 1$, $B \equiv -1 \pmod 3$.

If we now set

$$A = a'^3$$
$$B = b'^3$$
$$C = c'^3$$

we have derived from the solution of the equation $x^3 + y^3 + z^3 = 0$

$$x = a$$
$$y = b$$
$$z = c$$

another solution,

$$x = a'$$
$$y = b'$$
$$z = c'$$

$$\text{where } a'^3 b'^3 c'^3 = \frac{c^3}{(\varepsilon\varepsilon - \varepsilon)^3}$$

so $c'$ will contain the factor $1 - \varepsilon$ one fewer time than $c$. But this is absurd if $c$ is divisible only by a fixed power of $1 - \varepsilon$, that is, if $c$ differs from 0. For by continuing these operations we would otherwise eventually reach a solution in which $z$ was not divisible by $1 - \varepsilon$ at all, contrary to (I).

---

We can take a similar path for powers with exponent 5. Namely, if $a^5 + b^5 + c^5 = 0$, we set $b + c = \alpha$, $c + a = \beta$, $a + b = \gamma$, and then

$$\begin{aligned} 0 &= (2a)^5 + (2b)^5 + (2c)^5 \\ &= (\beta + \gamma - \alpha)^5 + (\gamma + \alpha - \beta)^5 + (\alpha + \beta - \gamma)^5 \\ &= (\alpha + \beta + \gamma)^5 - 80\alpha\beta\gamma(\alpha\alpha + \beta\beta + \gamma\gamma) \end{aligned}$$

But $(\alpha+\beta+\gamma)^5 = 80\alpha\beta\gamma(\alpha\alpha+\beta\beta+\gamma\gamma)$ cannot hold unless one of the numbers $\alpha, \beta, \gamma$ is divisible by $1-\varepsilon$. For if none of them were divisible, both $\alpha+\beta+\gamma$ and $\alpha\alpha+\beta\beta+\gamma\gamma$ would have to be divisible by $1-\varepsilon$, and consequently so would $2(\alpha\alpha+\beta\beta+\gamma\gamma)+2(\alpha+\beta+\gamma)(\alpha+\beta-\gamma) = (2\alpha+\beta)^2+3\beta\beta$, which is impossible.

We can also present this as follows. If $a^5+b^5+c^5=0$, then

$$\begin{aligned} 4(a+b+c)^5 &= 5(b+c)(c+a)(a+b) \\ &\qquad [(a+2b+3c)^2+3(a+c)^2-8(a+b+c)c] \\ &= 5(b+c)(c+a)(a+b) \\ &\qquad [(b-c)^2+3(b+c)^2+4(a+b+c)a] \end{aligned}$$

$$4(a+b+c)^5+5abc[(b-c)^2+3(b+c)^2] = 5(a+b+c)\{\dots\}$$

The rest of the proof would be very similar to the preceding one.

But if we try the same procedure with seventh powers, we cannot prove that for a given solution

$$a^7+b^7+c^7=0$$

one of the quantities $a, b, c$ must be divisible by 7. For we obtain only

$$\begin{aligned} (\alpha+\beta+\gamma)^7 = 5\alpha\beta\gamma\{&3(\alpha^4+\beta^4+\gamma^4) \\ &+10(\alpha\alpha\beta\beta+\alpha\alpha\gamma\gamma+\beta\beta\gamma\gamma)\} \end{aligned}$$

which can hold without $\alpha, \beta, \gamma$ being divisible by $1-\varepsilon$.

Hopefully, however, this can in future be derived from the nature of the determinants and the units.

---

## [II.]

## DETERMINATION OF THE NEAREST INTEGER.

Let

$$\begin{aligned} \varepsilon^3 = 1, \quad m = a+b\varepsilon+c\varepsilon\varepsilon \\ 2a-b-c = A+\alpha \\ 2b-c-a = B+\beta \\ 2c-a-b = C+\gamma \end{aligned}$$

where $A, B, C$ are integers and $\alpha, \beta, \gamma$ are positive proper fractions. We then have

$$A+B+C+\alpha+\beta+\gamma = 0$$

so there are three cases to distinguish:

I. $\alpha + \beta + \gamma = 0$, hence $\alpha = 0$, $\beta = 0$, $\gamma = 0$

1, $A \equiv B \equiv C \pmod 3$. Here $m$ itself is an integer.

2, $A - B \equiv B - C \equiv C - A \equiv \pm 1 \pmod 3$. Here $m \pm \frac{\varepsilon - \varepsilon\varepsilon}{3} \cdot \varepsilon^n$ is an integer.

II. $\alpha + \beta + \gamma = 1$.

Here each of

$$A + B\varepsilon + C\varepsilon\varepsilon + 1$$
$$A + B\varepsilon + C\varepsilon\varepsilon + \varepsilon$$
$$A + B\varepsilon + C\varepsilon\varepsilon + \varepsilon\varepsilon$$

is divisible by $1 - \varepsilon$, and one of these numbers by 3. The quotient, or

$$m + \frac{\varepsilon^n - \alpha - \beta\varepsilon - \gamma\varepsilon\varepsilon}{3}$$

is the required integer.

III. $\alpha + \beta + \gamma = 2$

Here

$$A + B\varepsilon + C\varepsilon\varepsilon + \varepsilon + \varepsilon\varepsilon$$
$$A + B\varepsilon + C\varepsilon\varepsilon + \varepsilon\varepsilon + 1$$
$$A + B\varepsilon + C\varepsilon\varepsilon + 1 + \varepsilon$$

are divisible by $1 - \varepsilon$, and one of these numbers by 3. The quotient, or

$$m + \frac{\varepsilon^n(\varepsilon + \varepsilon\varepsilon) - \alpha - \beta\varepsilon - \gamma\varepsilon\varepsilon}{3}$$

is the required integer.

In all three cases the remainder has the form

$$x + y\varepsilon + z\varepsilon\varepsilon$$

so that $x, y, z$ are smaller than $\frac{1}{3}$ in absolute value, and $x + y + z = 0$. This necessarily gives

$$xx + yy + zz = 2xx - 2yz = 2yy - 2xz = 2zz - 2xy < \tfrac{2}{9}$$

because two of the three quantities $x, y, z$ must have the same sign. Consequently, the determinant of the remainder is

$$= \tfrac{3}{2}(xx + yy + zz) < \tfrac{1}{3} \qquad \text{Q. E. D.}$$

The nearest integer is found *more conveniently* as follows. Given $a + b\varepsilon + c\varepsilon\varepsilon = m$, set

$$b - a = C + \gamma$$
$$c - b = A + \alpha$$
$$a - c = B + \beta$$

where $A, B, C$ are the next smaller integers and $\alpha, \beta, \gamma$ are positive fractions. There are three cases to distinguish:

I. $\alpha + \beta + \gamma = 0$; then $m$ itself is an integer

II. $\alpha + \beta + \gamma = 1$; then the nearest integer is

$$\begin{array}{lll} B + (B+C)\varepsilon & & \text{if } \alpha \text{ is the largest fraction.} \\ \quad C\varepsilon + (A+C)\varepsilon\varepsilon & & \beta \\ A + B \quad . \quad +A\varepsilon\varepsilon & & \gamma \end{array}$$

II. $\alpha + \beta + \gamma = 2$; then the nearest integer is

$$\begin{array}{lll} B + 1 + (B+C+2)\varepsilon & & \text{if } \alpha \text{ is the smallest fraction.} \\ \quad (C+1)\varepsilon + (A+C+2)\varepsilon\varepsilon & & \beta \\ A + B + 2 \quad . \quad +(A+1)\varepsilon\varepsilon & & \gamma \end{array}$$

In II, 1, the remainder is $\beta + (\beta+\gamma)\varepsilon$, whose determinant is

$$\begin{aligned} &= \beta\beta + \beta\gamma + \gamma\gamma \\ &= \tfrac{1}{3} - \tfrac{1}{6}[(\alpha-\beta)(1+3\beta) + (\alpha-\gamma)(1+3\gamma)] \end{aligned}$$

*More simply still:*

Arrange the fractions $a - [a], b - [b], c - [c]$ by size, and call them $p, q, r$ in that order. If all three are equal, $m$ is an integer. If they are unequal, let $t$ be any fraction between

$$\begin{array}{llll} p \text{ and } q, & \text{according as} & q - p & \text{is largest} \\ q \text{ and } r & & r - q & \\ r \text{ and } 1 + p & & 1 + p - r & \end{array}$$

Then

$$[a - t] + [b - t]\varepsilon + [c - t]\varepsilon\varepsilon$$

is the nearest integer.

---

[III.]

Let $\varepsilon^5 = 1$

$$a + b\varepsilon + c\varepsilon\varepsilon + d\varepsilon^2 + e\varepsilon^4 = q'$$

$$a + b\varepsilon^{-1} + c\varepsilon^{-2} + d\varepsilon^{-3} + e\varepsilon^{-4} = q''''$$

$$(a-b)^2 + (b-c)^2 + (c-d)^2 + (d-e)^2 + (e-a)^2 = 2p'$$

$$(a-c)^2 + (b-d)^2 + (c-e)^2 + (d-a)^2 + (e-b)^2 = 2p''$$

$$q'q'''' = -p'\varepsilon - p''\varepsilon\varepsilon - p''\varepsilon^3 - p'\varepsilon^4 = P'$$

$$q''q''' = -p'\varepsilon\varepsilon - p''\varepsilon^4 - p''\varepsilon - p'\varepsilon^3 = P''$$

$$\text{Determinant} = P'P'' = -p'p' + 3p'p'' - p''p''$$
$$\text{Measure} = 2p' + 2p'' = 2P' + 2P''$$
$$= 5(aa + bb + cc + dd + ee) - (a + b + c + d + e)^2$$

Multiplication by $1 - \varepsilon$ gives the new measure $= 8p'$

$$\text{Maximum measure} = 2\left(\frac{\sin 72^\circ}{\sin 36^\circ} + \frac{\sin 36^\circ}{\sin 72^\circ}\right)\sqrt{D} = 4{,}472\sqrt{D}$$

*Modulus* $= 1 - \varepsilon$

$$1 - \varepsilon = x$$
$$\varepsilon = 1 - x$$
$$\varepsilon\varepsilon = 1 - 2x + xx$$
$$\varepsilon^3 = 1 - 3x + 3xx - x^3$$
$$\varepsilon^4 = 1 - 4x + 6xx - 4x^3 + x^4$$
$$= -4 + 6x - 4xx + x^3$$

Thus

$$\frac{1 - \varepsilon^n}{1 - \varepsilon} \equiv n \text{ mod. } (1 - \varepsilon)$$
$$\varepsilon^n \equiv 1 - nx \text{ mod. } (1 - \varepsilon)^2$$
$$\left(\frac{\varepsilon + \varepsilon^4}{\varepsilon\varepsilon + \varepsilon^3}\right)^n \equiv 1 + nxx \text{ mod. } (1 - \varepsilon)^3$$

Thus a number that is $\equiv 1$ mod. $(1 - \varepsilon)^3$ can be a unit *only* if it is also $\equiv 1$ (mod. 5).

---

[IV.]

## SOME REMARKS ON THE MEASURE OF NUMBERS.

Let $\varepsilon^n = 1$, with $n$ prime,

$$m = a + a'\varepsilon + a''\varepsilon\varepsilon + a'''\varepsilon^3 + \ldots + a^{(n-1)}\varepsilon^{n-1} = f\varepsilon$$
$$D = f\varepsilon \,.\, f\varepsilon\varepsilon \,.\, f\varepsilon^3 \ldots f\varepsilon^{n-1}$$
$$f\varepsilon \,.\, f\varepsilon^{n-1} = -b'(\varepsilon + \varepsilon^{n-1}) - b''(\varepsilon^2 + \varepsilon^{-2})$$
$$- b'''(\varepsilon^3 + \varepsilon^{-3}) \ldots$$

then

$$2b' = (a - a')^2 + (a' - a'')^2 + (a'' - a''')^2 + \text{etc.}$$
$$2b'' = (a - a'')^2 + (a' - a''')^2 + (a'' - a'''')^2 + \text{etc.}$$
$$\text{etc.}$$

Here $b', b'', b''' \ldots$ are therefore all positive quantities; they are called the *partial measures* of $m$, and their sum

$$b' + b'' + b''' + \text{etc.} = n(aa + a'a' + a''a'' + \ldots) - (a + a' + a'' + \text{etc.})^2$$

the *general measure*. If we set

$$f\varepsilon \,.\, f\varepsilon^{n-1} = c', \quad f\varepsilon\varepsilon \,.\, f\varepsilon^{n-2} = c'' \quad \text{etc.}$$

then

$$c' + c'' + c''' + \text{etc.} + c^{\frac{1}{2}(n-1)} = b' + b'' + b''' + \text{etc.} + b^{\frac{1}{2}(n-1)}$$

$$c'(\varepsilon + \varepsilon^{n-1}) + c''(\varepsilon\varepsilon + \varepsilon^{n-2}) + c'''(\varepsilon^3 + \varepsilon^{n-3}) + \text{etc.} = 2(b' + b'' + b''' + \text{etc.} + b^{\frac{1}{2}(n-1)}) - nb'$$

$$c'(2 - \varepsilon - \varepsilon^{n-1}) + c''(2 - \varepsilon\varepsilon - \varepsilon^{n-2}) + c'''(2 - \varepsilon^3 - \varepsilon^{n-3}) + \text{etc.} = nb'$$

$$b' > \frac{n-1}{2n}(nD)^{\frac{2}{n-1}}, \quad b' + b'' + b''' + \text{etc.} > \frac{n-1}{2} \,.\, D^{\frac{2}{n-1}}$$

If, in general,

$$f\varepsilon \,.\, f\varepsilon^{n-1} = A + A'\varepsilon + A''\varepsilon\varepsilon + A'''\varepsilon^3 + \ldots$$

then the general measure is $\triangle = -A - A' - A'' - \text{etc.} + nA$

Measure of $(1 + \varepsilon) f\varepsilon \ldots$

$$\triangle' = 4\triangle - 2n(A - A') = 4\triangle - 2nb'$$

If $a + a' + a'' + \ldots = 0$, then $\triangle = n(aa + a'a' + a''a'' + \text{etc.})$

and if $A + A' + A'' + \ldots = 0$, then $\triangle = nA$, $\triangle' = n(2A + 2A')$

Thus, if one of the coefficients $A', A''$, etc., is negative and greater in absolute value than $\frac{1}{2}A$, the measure can be reduced without changing the determinant.

---

[V.]

If it should be confirmed that every unit is composed solely of factors of the form

$$\frac{\varepsilon^{\alpha} - \varepsilon^{\beta}}{\varepsilon^{\gamma} - \varepsilon^{\delta}}$$

the following theorem would be proved:

*If $f(\varepsilon)$ is a unit, then*

$$\frac{f(\varepsilon)}{f(\varepsilon^{-1})} = \varepsilon^{n}$$

Even without *assuming that* theorem, the conclusion is easy to prove. Let

$$\frac{f\varepsilon}{f\varepsilon^{-1}} = F\varepsilon$$

then

$$F\varepsilon \,.\, F\varepsilon^{-1} = 1$$

from which the theory of measure readily gives

$$F\varepsilon = \pm\varepsilon^{n}$$

The lower sign is impossible, however, since otherwise $f\varepsilon$ would have to be divisible by $1 - \varepsilon$.

It is easy to prove that the determinant of a number different from 0 cannot be $= 0$. If the determinant is divisible by $m$, the number itself is divisible by $1 - \varepsilon$. Consequently, if the determinant is divisible by $m^{m-1}$, the number itself must be divisible by $m$. This is absurd: with determinant 0, the number could first be divided by $m$, without changing its determinant, until it was no longer divisible. The first assertion follows as follows. Let the given number be

$$a + b\varepsilon + c\varepsilon\varepsilon + \text{etc.} \equiv a + b + c\,.\,. \mod 1 - \varepsilon$$

so its determinant is $\equiv (a + b + c\,.\,.)^{m-1} \mod 1 - \varepsilon$.

---

[VI.]

Let $\varepsilon^{n} = 1$

$$\begin{aligned} f\varepsilon &= a + b\varepsilon + c\varepsilon\varepsilon + d\varepsilon^{3} + \text{etc.} \\ m &= \text{determinant of this number} \\ \frac{m}{f\varepsilon} &= f\varepsilon\varepsilon \,.\, f\varepsilon^{3} \ldots f\varepsilon^{n-1} \\ &= A + B\varepsilon + C\varepsilon\varepsilon + \text{etc.} = F\varepsilon \end{aligned}$$

The number $f\varepsilon$ corresponds to a root of the congruence $x^n \equiv 1$ (mod. $m$.) Let it be $r$. We have

$$\begin{aligned} nA &= F1 + F\varepsilon + F\varepsilon\varepsilon + \ldots \\ nB &= F1 + \varepsilon^{-1}F\varepsilon + \varepsilon^{-2}F\varepsilon\varepsilon + \ldots \\ nC &= F1 + \varepsilon^{-2}F\varepsilon + \varepsilon^{-4}F\varepsilon\varepsilon + \ldots \\ &\text{etc.} \end{aligned}$$

Thus, since $F\varepsilon\varepsilon, F\varepsilon^3, F\varepsilon^4$, etc., are divisible by $f\varepsilon$,

$$\begin{gathered} nA - F1 - \varepsilon(nB - F1) \\ nA - F1 - \varepsilon\varepsilon(nC - F1) \\ nA - F1 - \varepsilon^3(nD - F1) \\ \text{etc.} \end{gathered}$$

are all divisible by $f\varepsilon$, or also

$$\begin{gathered} n(A - B) - \varepsilon n(B - C) \\ n(B - C) - \varepsilon n(C - D) \\ \text{etc.} \end{gathered}$$

are divisible by $f\varepsilon$; consequently [if $f\varepsilon$ is not divisible by $1 - \varepsilon$, and $F\varepsilon$ is not divisible by a real integer]

$$\varepsilon \equiv \frac{A-B}{B-C} \equiv \frac{B-C}{C-D} \equiv \frac{C-D}{D-E} \text{ etc. (mod. } f\varepsilon)$$

---

## REMARKS.

The investigations collected here under the common heading On the Theory of Complex Numbers are scattered notes in the manuscript. They contain the essential steps of the proof of FERMAT's theorem for the third and fifth powers. In incomplete notes that are not printed here, numbers composed of cube roots of unity are studied by means of both the theory of binary quadratic forms and the division of the circle. When applying the latter, during the preparation of the paper *Disquisitionum circa aequationes puras ulterior evolutio*, the ternary cubic form into which $27\frac{x^n-1}{x-1}$ can be transformed for a prime $n \equiv 1 \bmod. 3$ was also set out, and the theory of composition of the related form $X^3 + mY^3 + mmZ^3 - 3mXYZ$ was developed at the same time.

The property of integers formed from cube roots of unity that is assumed in the investigations of Fragment [I], namely that each can be decomposed into prime factors in only one way, follows from EUCLID's procedure for finding the common divisors of two numbers, if one applies the theorem derived in [II] on the nearest integer to any given fraction.

That the same fundamental property holds for numbers composed of fifth roots of unity follows from this: the fractional remainder formed by a rule entirely analogous to that in [II], either for $m$ or else for $m$ multiplied by a suitable unit $E$, is such that subtracting it from the given number $mE$ produces an integer, and its determinant does not exceed unity. As indicated in [III], the units can be derived from the theory of binary quadratic forms of determinant 5, in conjunction with the decomposition of a real prime into four factors (for example, $11 = \text{Det. } (2 + \varepsilon)$), namely as products of powers of $\varepsilon$ and $1 + \varepsilon$.

SCHERING.

# TABLE

OF THE QUADRATIC CHARACTER

OF THE PRIMES FROM 2 TO 997 AS RESIDUES

WITH RESPECT

TO THE PRIMES FROM 3 TO 503 AS DIVISORS.

| | 2 | 3 | 5 | 7 | 11 | 13 | 17 | 19 | 23 | 29 | 31 | 37 | 41 | 43 | 47 | 53 | 59 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 3 | | − | | − | | − | | − | | | − | − | | − | | | | 3 |
| 5 | | | − | | − | | | − | | − | − | | − | | | | − | 5 |
| 7 | − | | | − | − | | | | − | − | | − | | − | | − | | 7 |
| 11 | | − | − | | − | | | | − | | − | − | | | − | − | − | 11 |
| 13 | | − | | | | − | − | | − | − | | | | − | | − | | 13 |
| 17 | − | | | | | − | − | − | | | | | | − | − | − | − | 17 |
| 19 | | | − | − | − | | − | − | − | | | | | − | − | | | 19 |
| 23 | − | − | | | | − | | | − | − | − | | − | | − | | − | 23 |
| 29 | | | − | − | | − | | | − | − | | | | | | − | − | 29 |
| 31 | − | | − | − | | | | − | | | − | | − | | − | | − | 31 |
| 37 | | − | | − | − | | | | | | | − | − | | − | − | | 37 |
| 41 | − | | − | | | | | | − | | − | − | − | − | | | − | 41 |
| 43 | | | | | − | − | − | | − | | − | | − | − | − | − | − | 43 |
| 47 | − | − | | − | | | − | | | | | − | | | − | − | − | 47 |
| 53 | | | | − | − | − | − | | | − | | − | | − | − | − | − | 53 |
| 59 | | − | − | − | | | − | − | | − | | | − | | | − | − | 59 |
| 61 | | − | − | | | − | | − | | | | | − | | − | | | 61 |
| 67 | | | | | | | − | − | − | − | | − | | | − | | − | 67 |
| 71 | − | − | − | | | | | − | | − | | − | | − | | | | 71 |
| 73 | − | − | | | | | | − | − | | | − | − | | | | | 73 |
| 79 | − | | − | | − | − | | − | − | | − | | | | | | | 79 |
| 83 | | − | | − | − | | − | | − | − | − | − | − | | | | − | 83 |
| 89 | − | | − | | − | | − | | | | | | | | − | − | | 89 |
| 97 | − | − | | | − | | | | | | − | | | − | − | − | | 97 |
| 101 | | | − | | | − | − | − | − | | − | − | | − | − | | | 101 |
| 103 | − | | | − | | − | − | − | − | − | | | − | | | | − | 103 |
| 107 | | − | | | − | − | | − | − | − | | − | − | | − | − | | 107 |
| 109 | | − | − | − | | | | | | − | − | | | − | | | | 109 |
| 113 | − | | | − | − | − | | | | | − | | − | | | − | | 113 |
| 127 | − | | | | − | − | − | − | | | − | − | − | | − | | | 127 |
| 131 | | − | − | − | − | − | | | | | | | − | − | | − | − | 131 |
| 137 | − | | | − | − | | − | − | | | | − | | | | | − | 137 |
| 139 | | | − | − | − | − | | | | − | − | − | − | | − | | | 139 |
| 149 | | | − | − | | | − | − | | − | − | − | | | − | − | | 149 |
| 151 | − | | − | | − | | − | − | | − | − | − | | − | − | | − | 151 |
| 157 | | − | | | − | − | − | − | | | − | − | | | − | | | 157 |
| 163 | | | | | | | | | | | | | − | − | − | − | | 163 |
| 167 | − | − | | − | − | | | − | | − | − | | | | − | | | 167 |
| 173 | | | | | | − | | | − | − | − | − | − | − | − | | | 173 |
| 179 | | − | − | | | − | − | − | | − | − | | | − | − | | − | 179 |
| 181 | | − | − | | − | − | | | | − | | − | | − | | | − | 181 |
| 191 | − | − | − | | | − | − | | − | | | | | − | | | − | 191 |
| 193 | − | − | | − | | | | | − | | − | | | − | | | − | 193 |
| 197 | | | | − | | | | − | − | − | | − | − | − | − | − | − | 197 |
| 199 | − | | − | − | | − | | | − | − | − | | | − | − | − | | 199 |
| 211 | | | − | | − | − | | − | | | | − | | − | − | − | − | 211 |
| 223 | − | | | − | | | − | − | | − | − | − | − | − | − | − | | 223 |
| 227 | | − | | − | − | | | − | − | − | | | | − | − | − | − | 227 |
| | 2 | 3 | 5 | 7 | 11 | 13 | 17 | 19 | 23 | 29 | 31 | 37 | 41 | 43 | 47 | 53 | 59 | |

| | 61 | 67 | 71 | 73 | 79 | 83 | 89 | 97 | 101 | 103 | 107 | 109 | 113 | 127 | 131 | 137 | 139 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 3 | − | − | | − | − | | | − | | − | | − | | − | | | − | 3 |
| 5 | − | | − | | − | | − | | − | | | − | | | − | | − | 5 |
| 7 | | − | − | | − | | | | | | − | − | − | − | | − | | 7 |
| 11 | | − | − | | | | − | − | | − | | | − | | | − | | 11 |
| 13 | − | | | | − | | | | − | − | − | | − | − | − | | − | 13 |
| 17 | | − | | | | − | − | | − | − | | | | − | | − | | 17 |
| 19 | − | | | − | | − | | | − | | | | | | − | − | − | 19 |
| 23 | | | − | − | | | | | − | | | | | − | − | | − | 23 |
| 29 | | − | − | | | − | | | | − | − | − | | | | | − | 29 |
| 31 | | − | − | | | | | − | − | − | − | − | − | | − | | | 31 |
| 37 | | − | − | − | | − | | | − | | − | | | − | | − | − | 37 |
| 41 | − | | | − | | − | | | | − | − | | − | − | − | | − | 41 |
| 43 | | − | | | − | − | | − | − | − | − | − | | − | | | − | 43 |
| 47 | − | | − | | − | − | − | − | − | − | | | | | − | | | 47 |
| 53 | | | | | | | − | − | | | − | | − | | − | | | 53 |
| 59 | | | − | | − | | | | | | − | | | − | | − | − | 59 |
| 61 | − | | | − | | − | | − | | − | − | − | − | − | − | − | | 61 |
| 67 | | − | − | − | | − | − | | | − | − | | | − | − | | | 67 |
| 71 | | | − | − | − | − | − | | − | − | − | − | | | − | | | 71 |
| 73 | − | − | − | − | − | | − | − | | | | − | | − | | − | | 73 |
| 79 | | − | | − | − | − | − | − | − | | | | | | − | | | 79 |
| 83 | − | | | | | − | | | | | | − | − | − | − | | | 83 |
| 89 | | − | − | − | − | | − | − | | | − | − | | | − | | − | 89 |
| 97 | − | | | − | − | | − | − | − | − | | − | − | | | | | 97 |
| 101 | | | − | | − | | | − | − | | − | | | | − | − | | 101 |
| 103 | − | | | | − | − | | − | | − | − | | | | − | − | − | 103 |
| 107 | − | | | | − | − | − | | − | | − | | | | | − | | 107 |
| 109 | − | | − | − | | − | − | − | | | | − | − | | − | − | | 109 |
| 113 | − | | | | | − | | − | | | | − | − | − | − | | − | 113 |
| 127 | − | | − | − | − | | | | | − | − | | − | − | − | | | 127 |
| 131 | − | | | | | | − | | − | | − | − | − | | − | | | 131 |
| 137 | − | | | − | | | | | − | − | − | − | | | | − | − | 137 |
| 139 | | − | − | | − | − | − | | | | − | | − | − | − | − | − | 139 |
| 149 | − | − | | − | | | | | | − | − | | − | − | | | | 149 |
| 151 | | | | | | | | − | | − | | | | − | | − | − | 151 |
| 157 | | − | − | | | | − | | − | | | − | − | − | | | | 157 |
| 163 | − | | − | | | − | | − | | | | | − | | − | | | 163 |
| 167 | − | | | | | | − | − | | | − | | | − | | − | | 167 |
| 173 | | − | | − | | − | − | | | | | − | − | | | − | − | 173 |
| 179 | − | − | | | | − | − | | − | | − | | | | | | − | 179 |
| 181 | | − | | − | − | | | | − | | | | | | | − | − | 181 |
| 191 | | − | | | − | | | − | | − | − | − | | | | | | 191 |
| 193 | | − | | | | − | | − | − | | − | − | | | − | − | − | 193 |
| 197 | − | | | | | − | | − | − | | − | − | | − | | − | | 197 |
| 199 | − | | | | − | | − | | | − | | | | | − | | − | 199 |
| 211 | | | − | − | − | − | | | − | − | − | − | − | | | − | − | 211 |
| 223 | | | | − | | − | − | | − | | | − | | − | − | | − | 223 |
| 227 | | | − | − | − | | − | − | − | − | | − | − | | − | | − | 227 |
| | 61 | 67 | 71 | 73 | 79 | 83 | 89 | 97 | 101 | 103 | 107 | 109 | 113 | 127 | 131 | 137 | 139 | |

| | 2 | 3 | 5 | 7 | 11 | 13 | 17 | 19 | 23 | 29 | 31 | 37 | 41 | 43 | 47 | 53 | 59 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 229 | | – | – | | – | | – | – | | | | – | | – | | – | | 229 |
| 233 | – | | | – | | – | | – | – | – | – | – | | | | | | 233 |
| 239 | – | – | – | | – | | – | | | – | – | | | | | | | 239 |
| 241 | – | – | – | | | | | | | – | | | – | | – | – | – | 241 |
| 251 | | – | – | – | | – | – | | – | | – | | – | | | | | 251 |
| 257 | – | | | | – | – | – | | – | – | – | | | | | | – | 257 |
| 263 | – | – | | | – | – | – | | – | | – | – | | – | | | | 263 |
| 269 | | | – | | – | – | | | – | | | – | – | – | – | – | | 269 |
| 271 | – | | – | – | – | | – | | | | – | – | – | | | – | | 271 |
| 277 | | – | | – | | – | | – | – | – | | | – | | – | | – | 277 |
| 281 | – | | – | – | | | – | | | – | – | | | – | | – | – | 281 |
| 283 | | | | – | – | – | | | – | – | | | – | | | | – | 283 |
| 293 | | | | | | | – | | | | – | – | | – | | – | – | 293 |
| 307 | | | | – | – | | – | – | | | | – | – | | | – | | 307 |
| 311 | – | – | – | – | | – | | | | | | | | | – | – | | 311 |
| 313 | – | – | | | – | – | | – | | – | | | | | | | | 313 |
| 317 | | | | – | – | | | | – | | – | – | | – | | – | – | 317 |
| 331 | | | – | | | | – | – | | | – | | | – | | – | | 331 |
| 337 | – | – | | – | | – | | | | | | – | – | – | – | | | 337 |
| 347 | | – | | | – | – | | | | – | – | | | – | | – | – | 347 |
| 349 | | – | – | | | | – | – | – | – | – | – | – | | | | | 349 |
| 353 | – | | | | – | | – | – | – | – | | | – | – | – | | | 353 |
| 359 | – | – | – | | – | | – | | – | | | – | – | | – | | | 359 |
| 367 | – | | | – | | – | | | – | | – | – | – | | – | – | – | 367 |
| 373 | | – | | – | | – | – | | | – | – | – | – | | | | – | 373 |
| 379 | | | – | | | | | – | – | | | – | – | | | | | 379 |
| 383 | – | – | | – | | | – | – | – | – | – | | | – | | | | 383 |
| 389 | | | – | – | – | – | – | – | | | | | – | | | | – | 389 |
| 397 | | – | | | – | | | – | – | – | – | – | | – | – | | | 397 |
| 401 | – | | – | – | – | | | | | – | | | – | – | – | | | 401 |
| 409 | – | – | – | | | | – | | – | | | | – | | | – | | 409 |
| 419 | | – | – | – | | – | | | – | – | | – | – | – | – | | – | 419 |
| 421 | | – | – | – | – | | – | | | | – | | | | | | | 421 |
| 431 | – | – | – | | – | | | – | – | – | | | – | | | – | – | 431 |
| 433 | – | – | | | – | – | – | | | | | – | – | | | – | – | 433 |
| 439 | – | | – | – | – | – | | – | | – | | | | | | – | | 439 |
| 443 | | – | | | | – | – | | | | | – | – | | – | | – | 443 |
| 449 | – | | – | – | – | | | | – | | | | – | | | – | – | 449 |
| 457 | – | – | | – | | | – | – | | – | | | | | – | | | 457 |
| 461 | | | – | | | | – | – | – | | | | – | – | | – | – | 461 |
| 463 | – | | | | | | – | | | – | – | | | – | – | | – | 463 |
| 467 | | – | | – | | – | – | | – | | | | – | – | – | – | – | 467 |
| 479 | – | – | – | – | – | | | | – | | | | | | | | | 479 |
| 487 | – | | | | | | | – | | – | – | | – | | | – | | 487 |
| 491 | | – | – | | – | – | – | | | | – | – | – | – | | | | 491 |
| 499 | | | – | | | | | | | – | – | | | – | – | | | 499 |
| 503 | – | – | | – | – | – | | | – | | | | | – | – | | – | 503 |
| | 2 | 3 | 5 | 7 | 11 | 13 | 17 | 19 | 23 | 29 | 31 | 37 | 41 | 43 | 47 | 53 | 59 | |

| | 61 | 67 | 71 | 73 | 79 | 83 | 89 | 97 | 101 | 103 | 107 | 109 | 113 | 127 | 131 | 137 | 139 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 229 | – | | – | | | – | | – | | – | | | | | | | | 229 |
| 233 | | | – | | | | – | | – | | – | – | – | | – | | | 233 |
| 239 | – | – | – | | | – | | | – | | | – | – | – | | | | 239 |
| 241 | – | – | | | – | – | | – | | | – | | – | | | | | 241 |
| 251 | | – | | – | – | – | – | | – | – | | | – | | – | | | 251 |
| 257 | – | – | | – | – | | – | | | | | | – | | | – | – | 257 |
| 263 | – | | | | | – | – | | | – | | – | | | | – | | 263 |
| 269 | – | – | | – | – | | – | – | | – | | | | – | – | | | 269 |
| 271 | – | – | | | – | – | – | | | – | | | | | | | – | 271 |
| 277 | | – | – | | – | – | – | | | | | | – | | – | | | 277 |
| 281 | | | | | – | | | | – | | | – | | | | – | | 281 |
| 283 | – | | – | – | | – | – | – | – | – | | | – | – | | – | | 283 |
| 293 | – | – | – | – | | – | | – | | | – | – | | | | – | | 293 |
| 307 | | | – | | – | – | – | – | – | – | – | – | – | – | | | | 307 |
| 311 | | – | | – | – | – | – | | | | – | – | – | – | | – | – | 311 |
| 313 | | | – | | – | – | | – | | – | – | | – | | | – | – | 313 |
| 317 | – | – | | – | – | – | – | | – | – | | | – | | – | | | 317 |
| 331 | | – | – | | – | – | – | | | – | | – | – | – | – | | – | 331 |
| 337 | | | | | – | | | | | – | – | | – | | – | – | | 337 |
| 347 | – | – | – | – | | – | – | | | | – | – | – | – | – | – | | 347 |
| 349 | | – | | – | | – | | | | | | – | | | | | – | 349 |
| 353 | – | | | – | | – | | – | | | | – | – | – | – | | | 353 |
| 359 | | | | – | – | | | | – | | – | | | – | – | | | 359 |
| 367 | – | – | | – | | – | – | | – | | – | | – | | | – | | 367 |
| 373 | | | – | – | | – | – | | – | – | – | – | | | | – | | 373 |
| 379 | – | – | | | – | – | | – | – | – | – | | | – | | – | – | 379 |
| 383 | | – | – | – | | | | | – | – | | | – | | | – | – | 383 |
| 389 | | – | | – | – | | | – | | | | | – | – | | – | | 389 |
| 397 | | – | | – | – | – | | – | | | – | | | – | – | – | | 397 |
| 401 | | | | – | | – | – | | | – | | – | – | | | | | 401 |
| 409 | | | – | | | – | – | | – | – | | – | | | – | – | – | 409 |
| 419 | | | | – | – | | | – | | | – | | | | – | – | – | 419 |
| 421 | | – | | | – | | | – | – | – | – | – | – | | – | | – | 421 |
| 431 | – | | | | | | | – | | | | – | | | | | – | 431 |
| 433 | | | | | – | | | | | | | – | | – | | – | – | 433 |
| 439 | – | | – | – | | – | | | | – | | – | – | – | | – | – | 439 |
| 443 | – | – | – | | – | | – | | | – | – | – | – | | – | – | – | 443 |
| 449 | – | – | | | | | – | – | – | | | | | – | | – | | 449 |
| 457 | | – | | – | – | | | | | | – | – | | – | – | | | 457 |
| 461 | – | – | | – | | | – | | | – | – | – | – | | | – | – | 461 |
| 463 | – | – | | – | – | | – | – | | – | | – | – | | – | | | 463 |
| 467 | | | – | | | – | – | – | | | | – | – | – | | – | – | 467 |
| 479 | – | | – | – | | | – | – | | – | – | – | | | – | – | – | 479 |
| 487 | – | – | – | – | | – | – | – | | – | – | | | – | | – | – | 487 |
| 491 | – | | – | | – | – | | – | – | | – | | | – | – | | – | 491 |
| 499 | | – | | – | | | | | – | – | – | – | | – | – | – | – | 499 |
| 503 | – | – | | – | – | – | | – | | | | | – | | – | | | 503 |
| | 61 | 67 | 71 | 73 | 79 | 83 | 89 | 97 | 101 | 103 | 107 | 109 | 113 | 127 | 131 | 137 | 139 | |

| | 149 | 151 | 157 | 163 | 167 | 173 | 179 | 181 | 191 | 193 | 197 | 199 | 211 | 223 | 227 | 229 | 233 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 3 | | – | – | – | | | | – | | – | | – | – | – | | – | | 3 |
| 5 | – | – | | | | | – | – | – | | | – | – | | | – | | 5 |
| 7 | – | – | | – | | | – | | – | – | – | | – | | | | – | 7 |
| 11 | | | – | – | | | – | – | – | | | – | | – | | – | | 11 |
| 13 | | | – | | | – | – | – | – | | | – | – | | | | – | 13 |
| 17 | – | – | – | | | | – | | – | | | | | – | | – | | 17 |
| 19 | – | | – | – | | | | | – | | – | – | | | | – | – | 19 |
| 23 | | – | | – | – | – | – | | | – | – | | – | – | | | – | 23 |
| 29 | – | – | | | – | – | – | – | | | – | – | | – | – | | – | 29 |
| 31 | – | | – | – | | – | | | – | – | | | – | | – | | – | 31 |
| 37 | – | – | – | | | – | | – | | | – | | – | – | | – | – | 37 |
| 41 | | | | – | | – | | | | | – | | | – | | | | 41 |
| 43 | | | | | – | – | | – | | – | – | | | | | – | | 43 |
| 47 | – | | – | | | – | | | – | | – | | | | | | | 47 |
| 53 | – | | | – | | | | | | | – | – | – | – | – | – | | 53 |
| 59 | | | | – | – | | | – | | – | – | – | | – | | | | 59 |
| 61 | – | | | – | – | | – | | | | – | – | | | | – | | 61 |
| 67 | – | – | – | – | – | – | | – | | – | | – | – | – | – | | | 67 |
| 71 | | – | – | | – | | – | | – | | | – | | – | | – | – | 71 |
| 73 | – | | | | | – | | – | | | | | – | – | – | | | 73 |
| 79 | | – | | – | – | | – | – | | | | | | – | | | | 79 |
| 83 | | – | | | – | – | | | – | – | – | – | | | – | – | | 83 |
| 89 | | | – | | – | – | – | | | | | – | | – | – | | – | 89 |
| 97 | | – | | – | – | | | | – | – | – | | | | – | – | | 97 |
| 101 | | | – | | | | – | – | | – | – | | – | – | – | | – | 101 |
| 103 | – | | | – | – | | – | | | | | | | – | | – | | 103 |
| 107 | – | – | | – | | | | | | – | – | – | | – | – | | – | 107 |
| 109 | | | – | | | – | | | – | – | – | | – | – | – | | – | 109 |
| 113 | – | | – | – | | – | | | | | | | – | | – | | – | 113 |
| 127 | – | | – | – | | | – | | – | | – | – | – | | – | | | 127 |
| 131 | | – | | | – | | – | | – | – | | | – | | | | – | 131 |
| 137 | | – | | | – | – | | – | | – | – | | – | | | | | 137 |
| 139 | | | | – | – | – | | – | – | – | | | | | | | | 139 |
| 149 | – | | | | | – | – | | – | | | | | | | – | | 149 |
| 151 | | – | | | – | – | | | – | – | | | | – | – | – | | 151 |
| 157 | | | – | | – | – | | | | – | – | – | | | | | – | 157 |
| 163 | | – | | – | – | – | – | | | | – | – | | – | – | | | 163 |
| 167 | | | – | | – | – | – | – | – | | | – | – | – | | – | – | 167 |
| 173 | – | – | – | – | – | – | – | | | | – | | – | | – | – | – | 173 |
| 179 | – | – | | | | – | – | | – | – | | – | | | – | | | 179 |
| 181 | | | | | – | | | – | | – | – | | | – | – | – | – | 181 |
| 191 | – | | | – | | | | | – | – | – | – | – | – | – | | | 191 |
| 193 | | – | – | | | | – | – | – | – | – | – | – | | | – | | 193 |
| 197 | | | – | – | | – | | – | – | – | – | | | – | | | – | 197 |
| 199 | | – | – | | | | | | | – | | – | | | – | | | 199 |
| 211 | | – | | – | | – | – | | | – | | – | – | | – | | | 211 |
| 223 | | | | | | | – | – | | | – | – | – | – | – | | | 223 |
| 227 | | | | | – | – | | – | | | | | | | – | | | 227 |
| | 149 | 151 | 157 | 163 | 167 | 173 | 179 | 181 | 191 | 193 | 197 | 199 | 211 | 223 | 227 | 229 | 233 | |

| | 239 | 241 | 251 | 257 | 263 | 269 | 271 | 277 | 281 | 283 | 293 | 307 | 311 | 313 | 317 | 331 | 337 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 3 | | – | | | | | – | – | | – | | – | | – | | – | – | 3 |
| 5 | – | – | – | | | – | – | | – | | | | – | | | – | | 5 |
| 7 | – | | | | – | | | – | – | | | | | | – | – | – | 7 |
| 11 | | | – | – | | – | | | | | | | – | – | – | – | | 11 |
| 13 | | | – | – | – | – | | – | | – | | | – | – | | | – | 13 |
| 17 | – | | – | – | – | | – | | – | | – | – | | | | – | | 17 |
| 19 | – | | – | | – | | – | – | | – | | | – | – | | | | 19 |
| 23 | – | | | – | | – | – | – | | | | – | – | | – | – | | 23 |
| 29 | – | – | | – | | | | – | – | – | | | | – | | | | 29 |
| 31 | | | | – | | | | | – | – | – | – | – | | – | | | 31 |
| 37 | | | | | – | – | – | | | | – | – | | | – | | – | 37 |
| 41 | | – | – | | | – | – | – | | – | | – | | | | | – | 41 |
| 43 | – | | – | | | – | – | | – | – | – | – | – | | – | | – | 43 |
| 47 | – | – | – | | – | – | – | – | | – | | – | | | | – | – | 47 |
| 53 | | – | | | | – | – | | – | | – | – | – | | – | – | | 53 |
| 59 | – | – | – | – | – | | – | – | – | | – | – | – | | – | – | | 59 |
| 61 | – | – | | – | – | – | – | | | – | – | | | | – | | | 61 |
| 67 | | – | | – | – | – | | – | | – | – | – | | | – | | | 67 |
| 71 | | | – | | – | | – | – | | | – | | – | – | | | | 71 |
| 73 | | | – | – | | – | | | | – | – | | – | | – | | | 73 |
| 79 | – | – | | – | – | – | | – | – | – | | | | – | – | | – | 79 |
| 83 | | – | | | | | | – | | | – | | | – | – | | | 83 |
| 89 | | | – | – | – | – | – | – | | – | | – | – | | – | – | | 89 |
| 97 | | – | | | | – | | | | – | – | – | | – | | | | 97 |
| 101 | – | | – | | | | | | – | – | | – | | | – | | | 101 |
| 103 | – | | | | | – | | | | | | | – | – | – | | – | 103 |
| 107 | – | – | – | | – | | – | | | – | – | | | – | | – | – | 107 |
| 109 | – | | | | – | | | | – | | – | – | – | | | – | | 109 |
| 113 | – | – | – | – | | | | – | | – | | – | – | – | – | – | – | 113 |
| 127 | | | – | | – | – | – | | | | | | | | | | | 127 |
| 131 | – | | | | – | – | – | – | | – | | – | – | | – | | – | 131 |
| 137 | | | | – | – | | | | – | – | – | | – | – | | | – | 137 |
| 139 | – | | – | – | – | | | | | – | | – | | – | | | | 139 |
| 149 | | | – | | – | – | | | – | | – | – | | | – | – | – | 149 |
| 151 | – | – | – | | | – | | | | | | – | – | – | | – | | 151 |
| 157 | – | | | – | – | | | – | – | – | | | – | – | – | – | | 157 |
| 163 | | | – | | – | | | | – | | | – | | – | | | | 163 |
| 167 | – | | – | | – | | | | – | – | – | | – | | – | | – | 167 |
| 173 | | | – | – | – | – | | | | | | | – | – | – | – | – | 173 |
| 179 | – | | | | | | | | | | | | | | – | | – | 179 |
| 181 | | – | – | | – | | | | – | – | | – | | – | – | | – | 181 |
| 191 | – | – | – | | – | – | – | – | – | – | – | | – | | | | | 191 |
| 193 | – | – | | – | | | | – | | | – | | – | | – | – | – | 193 |
| 197 | – | | – | – | | | | | | | – | | – | – | | – | | 197 |
| 199 | – | | – | – | – | – | – | | | | – | | – | – | | – | | 199 |
| 211 | | – | | – | – | – | | – | – | | – | – | – | | | – | – | 211 |
| 223 | – | – | – | – | | | | | – | – | | | | | – | | | 223 |
| 227 | – | | | – | – | | – | | | | | | – | | – | – | – | 227 |
| | 239 | 241 | 251 | 257 | 263 | 269 | 271 | 277 | 281 | 283 | 293 | 307 | 311 | 313 | 317 | 331 | 337 | |

| | 149 | 151 | 157 | 163 | 167 | 173 | 179 | 181 | 191 | 193 | 197 | 199 | 211 | 223 | 227 | 229 | 233 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 229 | – | – | | | – | – | | – | | – | | | | | | – | – | 229 |
| 233 | | | – | | – | – | | – | | | – | | | | | – | – | 233 |
| 239 | | | – | – | | | | | | – | – | | – | | | | | 239 |
| 241 | | – | | | | | | – | – | – | | | – | – | | – | – | 241 |
| 251 | – | | | | | – | – | – | | | – | | – | | – | | – | 251 |
| 257 | | | – | | | – | | | | – | – | – | – | – | – | | | 257 |
| 263 | – | – | – | | | – | – | – | | | | | | – | | | – | 263 |
| 269 | – | – | | | | – | | | – | | | – | – | | | | – | 269 |
| 271 | | – | | – | – | | – | | | | | | – | – | | – | – | 271 |
| 277 | | | – | | | | | | – | – | | | – | | | – | | 277 |
| 281 | – | | – | – | – | | | – | – | | | | – | – | | | | 281 |
| 283 | | – | – | – | | | – | – | | | | – | – | | – | | – | 283 |
| 293 | – | | | | – | | | | – | – | – | – | – | | | – | – | 293 |
| 307 | – | | | | – | | – | – | – | | | – | | – | – | – | – | 307 |
| 311 | | | – | – | | – | – | | | – | – | | | – | | – | | 311 |
| 313 | | – | – | – | | – | | – | | | – | – | | | | | | 313 |
| 317 | – | | – | | – | – | – | – | | – | | | | – | – | | | 317 |
| 331 | – | | – | – | – | – | – | | – | – | – | | | – | | | – | 331 |
| 337 | – | | | | – | – | – | – | | – | | | – | | – | – | – | 337 |
| 347 | – | | – | | – | – | | – | | | – | – | | | | – | | 347 |
| 349 | | – | – | | | | | – | – | | | | | – | – | | – | 349 |
| 353 | | | – | | – | | | – | – | | – | | | – | | | – | 353 |
| 359 | – | – | | | | – | | – | – | – | | | | | | – | – | 359 |
| 367 | – | – | | | | – | | – | – | | | – | – | | – | – | | 367 |
| 373 | | | | – | | | – | – | | | | | | | | – | | 373 |
| 379 | – | – | | | – | – | – | | – | – | – | | – | | – | | | 379 |
| 383 | – | | | | | – | | | | – | | | | – | – | – | | 383 |
| 389 | | | – | | | – | – | – | | – | | | – | – | | | | 389 |
| 397 | | – | | – | – | – | – | | – | | | | | | | – | | 397 |
| 401 | – | – | | | | – | – | – | | | – | | | – | | – | | 401 |
| 409 | | | | – | – | | – | | – | – | – | | | | – | – | | 409 |
| 419 | – | – | – | | | – | | | – | | – | – | | | | | | 419 |
| 421 | – | | | – | – | | | – | – | | | – | | | | – | | 421 |
| 431 | – | – | – | – | | – | – | | | | – | | | – | – | – | | 431 |
| 433 | | | | | – | | – | | – | | – | – | – | – | | – | – | 433 |
| 439 | | | | | – | | | | – | | | – | – | – | | – | | 439 |
| 443 | – | – | | – | – | | | – | – | | – | | | | – | – | – | 443 |
| 449 | | | | | – | | | – | – | – | – | – | | | – | – | | 449 |
| 457 | | – | – | – | | | | | – | | – | | | | – | – | – | 457 |
| 461 | | – | – | – | – | | | – | – | – | | – | | – | – | – | | 461 |
| 463 | – | | | – | – | – | – | – | | | | | – | | | – | | 463 |
| 467 | – | – | – | – | | – | – | | | – | | – | | – | – | – | – | 467 |
| 479 | | – | | – | – | – | | – | | – | – | | – | | | | – | 479 |
| 487 | | | – | | – | | | – | – | – | – | | | | | | | 487 |
| 491 | | | | – | | | – | – | | | – | – | | – | – | – | – | 491 |
| 499 | | – | | | – | | | – | | | – | – | – | | – | | – | 499 |
| 503 | | | | | | – | | | | | – | – | | – | | – | – | 503 |
| | 149 | 151 | 157 | 163 | 167 | 173 | 179 | 181 | 191 | 193 | 197 | 199 | 211 | 223 | 227 | 229 | 233 | |

| | 239 | 241 | 251 | 257 | 263 | 269 | 271 | 277 | 281 | 283 | 293 | 307 | 311 | 313 | 317 | 331 | 337 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 229 | | – | | | | | – | – | | | – | – | – | | | | – | 229 |
| 233 | | – | – | | – | – | – | | | – | – | – | | | | – | – | 233 |
| 239 | – | – | – | – | – | – | – | | | – | – | – | – | | | | – | 239 |
| 241 | – | – | – | – | | | – | – | – | | | | | – | | – | – | 241 |
| 251 | | – | – | | – | | – | | | | | | – | | – | – | – | 251 |
| 257 | – | – | | – | | | | | | – | – | – | | | – | | | 257 |
| 263 | | | | | – | – | – | | – | | | – | – | – | – | – | – | 263 |
| 269 | – | | | | – | – | | | | – | – | – | | – | | – | | 269 |
| 271 | | – | | | | | – | | – | | – | – | – | | | | | 271 |
| 277 | | – | | | | | | – | – | | – | – | – | – | – | | | 277 |
| 281 | | – | | | – | | – | – | – | – | | | | – | – | – | – | 281 |
| 283 | | | – | – | – | – | – | | – | – | – | – | – | | – | | – | 283 |
| 293 | – | | | – | | – | – | – | | – | – | – | | | – | – | | 293 |
| 307 | | | – | – | | – | | – | | | – | – | – | – | – | – | | 307 |
| 311 | | | | | | | | – | | | | | – | – | – | – | – | 311 |
| 313 | | – | | | – | – | | – | – | | | – | – | – | – | – | – | 313 |
| 317 | | | – | – | – | | | – | – | – | – | – | – | – | – | | | 317 |
| 331 | – | – | | | | – | – | | – | – | – | | | – | | – | – | 331 |
| 337 | – | – | – | | – | | | | – | – | | | – | – | | – | – | 337 |
| 347 | | – | – | | – | – | – | – | – | | – | | | | | | | 347 |
| 349 | – | – | | – | – | – | – | | – | – | – | – | | – | | | – | 349 |
| 353 | | | | | | – | – | – | – | – | – | – | – | | – | – | – | 353 |
| 359 | | – | | | | | | | – | – | | – | | | – | – | | 359 |
| 367 | | | – | | | | – | – | – | – | | | | – | | | | 367 |
| 373 | – | | – | – | | | | | | – | | – | | | | | – | 373 |
| 379 | – | | – | – | – | | – | – | – | | | – | – | – | | – | – | 379 |
| 383 | | | – | | – | | | – | | | – | | | – | – | – | | 383 |
| 389 | – | | | | | – | | – | | – | – | | – | – | | – | – | 389 |
| 397 | | | | – | | | – | – | – | | – | – | | | | | | 397 |
| 401 | – | – | | – | – | | | | | | | – | – | – | | – | – | 401 |
| 409 | – | | | | | | – | | – | | | – | – | – | – | | – | 409 |
| 419 | | | – | – | | – | | | – | | – | | | | | | – | 419 |
| 421 | – | – | | | | – | | – | – | – | | – | | – | – | | – | 421 |
| 431 | | | | | – | | | – | | – | | – | | | | | – | 431 |
| 433 | | – | | – | | – | – | – | | – | – | | | | | – | – | 433 |
| 439 | | | – | | | – | – | | – | – | | – | | | | – | | 439 |
| 443 | | | | | – | | – | | – | | – | – | – | | | – | | 443 |
| 449 | | | – | | – | – | – | | | | – | | | | | – | – | 449 |
| 457 | – | – | | – | | – | | | | – | | – | – | – | | – | | 457 |
| 461 | | | | | – | | | | – | | | | – | | – | – | | 461 |
| 463 | – | | – | | | | – | – | | – | – | | – | – | | | – | 463 |
| 467 | – | – | – | | | | | – | | – | | | | | – | – | | 467 |
| 479 | | – | – | – | | | – | – | | | – | – | | – | | – | | 479 |
| 487 | | – | – | | – | – | – | – | | | | – | – | – | | | – | 487 |
| 491 | – | – | | – | – | – | | – | | | – | | | | | – | | 491 |
| 499 | – | | – | – | – | | | – | – | | – | – | | | – | – | – | 499 |
| 503 | | | | – | – | | – | | – | – | – | | | | – | | | 503 |
| | 239 | 241 | 251 | 257 | 263 | 269 | 271 | 277 | 281 | 283 | 293 | 307 | 311 | 313 | 317 | 331 | 337 | |

| | 347 | 349 | 353 | 359 | 367 | 373 | 379 | 383 | 389 | 397 | 401 | 409 | 419 | 421 | 431 | 433 | 439 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 3 | | – | | | – | – | – | | | – | | – | | – | | – | – | 3 |
| 5 | | – | | – | | | – | | – | | – | – | – | – | – | | – | 5 |
| 7 | – | | | – | | – | – | | – | | – | | | – | – | | | 7 |
| 11 | | | – | | – | | – | – | – | – | – | | – | – | | – | | 11 |
| 13 | – | | | | – | – | | | – | | | | – | | | – | – | 13 |
| 17 | | – | – | – | | – | | – | – | | | – | | – | | – | | 17 |
| 19 | – | – | – | – | – | | | | – | – | | | – | | | | | 19 |
| 23 | – | – | – | | | | | | | – | | – | | | | | – | 23 |
| 29 | – | – | – | | | – | | – | | – | – | | – | | – | | – | 29 |
| 31 | | – | | – | | – | – | | | – | | | – | – | – | | – | 31 |
| 37 | | – | | – | – | – | – | | | – | | | – | | | – | | 37 |
| 41 | | – | – | – | – | – | – | | – | | – | – | – | | – | – | | 41 |
| 43 | | | – | – | – | | – | | | – | – | | | | – | | – | 43 |
| 47 | – | | – | | | | – | – | | – | – | | | | – | | – | 47 |
| 53 | – | | | | – | | | | | | | – | | | – | – | – | 53 |
| 59 | | | | – | | – | – | – | – | | | | | | | – | – | 59 |
| 61 | – | | – | | – | | – | | | | | | | | – | | – | 61 |
| 67 | | – | | – | | | | | – | – | | | – | – | – | | – | 67 |
| 71 | | | | – | – | – | – | | | | | – | – | | – | | | 71 |
| 73 | – | – | – | – | – | – | | – | – | – | – | | – | | | | – | 73 |
| 79 | – | | | | – | | | – | – | – | | | | – | – | – | – | 79 |
| 83 | | – | – | – | | – | | – | | – | – | – | – | | – | | | 83 |
| 89 | – | | | | – | – | | | | | – | – | | | | | | 89 |
| 97 | | | – | | | | – | | – | – | | | – | – | – | | | 97 |
| 101 | | | | – | – | – | – | – | | | | – | | – | | | | 101 |
| 103 | – | | | – | – | – | | | | | – | – | – | – | – | | | 103 |
| 107 | | | | | | – | | – | | – | | | | – | – | | – | 107 |
| 109 | – | – | – | | | – | | | | | – | – | | – | – | – | – | 109 |
| 113 | – | | – | | – | | | – | – | | – | | | – | | | – | 113 |
| 127 | | | – | | – | | | – | – | – | | | – | | – | – | | 127 |
| 131 | | | – | | – | | – | – | | – | | – | | – | – | | – | 131 |
| 137 | – | | | | – | – | – | – | – | – | | – | – | | | – | – | 137 |
| 139 | – | – | | – | – | | | | | | | – | | – | | – | | 139 |
| 149 | – | | | – | – | | – | – | | | – | | – | – | – | | | 149 |
| 151 | – | – | | | | | | – | | – | – | | | | | | – | 151 |
| 157 | – | – | – | | | | | | – | | | | – | | – | | | 157 |
| 163 | – | | | – | – | – | – | – | | – | | – | – | – | | | – | 163 |
| 167 | | | – | – | – | | | – | | – | | – | – | – | – | – | | 167 |
| 173 | – | | | – | – | | – | – | – | – | – | | – | | – | | | 173 |
| 179 | – | | | – | – | – | | – | – | – | – | – | – | | | – | – | 179 |
| 181 | – | – | – | – | – | – | | | – | | – | | | – | | | | 181 |
| 191 | – | – | – | | | | | – | | – | | – | | – | – | – | | 191 |
| 193 | | | | – | | | – | – | – | | | – | | | | | | 193 |
| 197 | – | | – | | | | – | | | | – | – | – | | – | – | | 197 |
| 199 | | | | – | | | – | – | | | | | | – | – | – | | 199 |
| 211 | – | | | – | | | | – | – | | | | – | | – | – | | 211 |
| 223 | – | – | – | – | – | | – | | – | | – | | – | | | – | | 223 |
| 227 | – | – | | – | | | | | | | | – | – | | | | – | 227 |
| | 347 | 349 | 353 | 359 | 367 | 373 | 379 | 383 | 389 | 397 | 401 | 409 | 419 | 421 | 431 | 433 | 439 | |

| | 443 | 449 | 457 | 461 | 463 | 467 | 479 | 487 | 491 | 499 | 503 | 509 | 521 | 523 | 541 | 547 | 557 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 3 | | | – | | – | | | – | | – | | | | – | – | – | | 3 |
| 5 | | – | | – | | | – | | – | – | | – | – | | – | | | 5 |
| 7 | – | – | – | | – | | | – | – | – | | | | | – | – | – | 7 |
| 11 | – | – | | | – | – | | – | | – | | – | – | | | | | 11 |
| 13 | – | | | | | – | | | – | | – | | – | – | | – | | 13 |
| 17 | – | | – | – | – | – | | | – | | | – | | – | | | – | 17 |
| 19 | – | | – | – | – | – | – | | – | – | – | | | | – | | – | 19 |
| 23 | – | – | | – | – | | | – | – | – | | – | | | – | – | | 23 |
| 29 | | | – | | – | | | – | | – | | – | – | – | | – | – | 29 |
| 31 | – | | | | | – | – | | | | – | | – | | – | – | | 31 |
| 37 | – | | | | | | | | – | | | – | – | | | | | 37 |
| 41 | – | – | | – | | – | | – | – | | | | | – | – | | | 41 |
| 43 | – | | | – | | | – | – | | | | – | | | – | – | – | 43 |
| 47 | | | – | | | | – | – | – | | | | – | – | – | | | 47 |
| 53 | | – | | – | | – | | – | | | | | – | – | – | – | | 53 |
| 59 | | – | | – | | | – | – | – | – | | | – | – | | – | – | 59 |
| 61 | – | – | | – | – | | – | – | – | | – | | | | | | | 61 |
| 67 | | – | – | – | | – | – | | – | | | – | | – | | | – | 67 |
| 71 | | | | | – | | | | | – | – | – | – | | | – | – | 71 |
| 73 | | | – | – | – | | – | – | | – | – | – | | – | | – | – | 73 |
| 79 | | | – | | | – | – | – | | – | | | | – | – | – | – | 79 |
| 83 | – | | | | | | – | | | – | | – | – | – | | – | – | 83 |
| 89 | – | – | | – | – | – | – | – | | | | – | | – | | | | 89 |
| 97 | | – | | – | – | – | – | – | – | | – | – | – | | | – | – | 97 |
| 101 | | – | | | | | | | – | – | | – | – | | – | | – | 101 |
| 103 | | | | – | | – | | | – | | – | – | | – | – | – | | 103 |
| 107 | | | – | – | – | – | | | | | – | – | | | | – | | 107 |
| 109 | – | | – | – | – | – | – | | | – | | – | | – | – | | – | 109 |
| 113 | – | | | – | – | – | | | | | – | – | – | | | – | – | 113 |
| 127 | – | – | – | | – | | – | | | | – | – | – | – | | | – | 127 |
| 131 | | | – | | | – | | – | | | | | | | | | – | 131 |
| 137 | – | – | | – | | – | – | – | | – | | – | | – | – | – | – | 137 |
| 139 | | | | – | – | | | | | | – | | | – | – | | – | 139 |
| 149 | – | | | | – | – | | | | | | | | – | | – | – | 149 |
| 151 | | | – | – | – | | | – | – | | – | | – | | – | – | | 151 |
| 157 | | | – | – | | – | | – | | | | | | – | | – | – | 157 |
| 163 | | | – | – | | | | – | | – | – | | | – | | – | | 163 |
| 167 | | – | | – | | – | | | – | | – | – | | – | | | – | 167 |
| 173 | | | | | – | – | – | | | | – | – | | – | – | | – | 173 |
| 179 | – | | | | | | – | – | | – | – | – | | | – | | – | 179 |
| 181 | – | – | | – | – | | – | – | – | – | | – | | – | | – | – | 181 |
| 191 | | – | – | – | – | – | – | | – | – | – | | | | | | | 191 |
| 193 | | – | | – | | – | – | – | | | | | | – | | – | | 193 |
| 197 | – | – | – | | | | – | – | – | – | – | | – | | | | – | 197 |
| 199 | – | – | | – | – | | – | – | | | | – | – | – | | | | 199 |
| 211 | – | | | | | – | | – | – | | – | – | – | – | – | – | | 211 |
| 223 | – | | | – | – | | – | – | | – | | – | | | | – | | 223 |
| 227 | | – | – | – | – | | – | – | | | – | | | – | – | | – | 227 |
| | 443 | 449 | 457 | 461 | 463 | 467 | 479 | 487 | 491 | 499 | 503 | 509 | 521 | 523 | 541 | 547 | 557 | |

| | 347 | 349 | 353 | 359 | 367 | 373 | 379 | 383 | 389 | 397 | 401 | 409 | 419 | 421 | 431 | 433 | 439 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 229 | – | | | – | – | – | | – | | – | – | – | | – | – | – | – | 229 |
| 233 | | – | – | – | | | | | | | | | | | | – | | 233 |
| 239 | – | – | | – | – | – | | – | – | | – | – | – | – | – | | – | 239 |
| 241 | – | – | | – | | | | | | | – | | | – | | – | | 241 |
| 251 | | | | – | | – | | | | | | | | | – | | | 251 |
| 257 | | – | | | | – | – | | | – | – | | – | | | – | | 257 |
| 263 | | – | | – | – | | | | | | – | | – | | | | – | 263 |
| 269 | – | – | – | | | | | | – | | | | – | – | | – | – | 269 |
| 271 | | – | – | – | | | | – | | – | | – | – | | – | – | | 271 |
| 277 | – | | – | | – | | – | – | – | – | | | | – | – | – | | 277 |
| 281 | – | – | – | – | – | | – | | | – | | – | – | – | | | – | 281 |
| 283 | – | – | – | | | – | – | – | – | | | | – | – | | – | | 283 |
| 293 | – | – | – | | | | | – | – | – | | | – | | | – | | 293 |
| 307 | – | – | – | | – | – | | – | | – | – | – | – | – | | | | 307 |
| 311 | – | | – | – | – | | | – | – | | – | – | – | | – | | – | 311 |
| 313 | | – | | | – | | – | – | – | | – | – | | – | | | | 313 |
| 317 | | | – | – | | | | – | | | | – | | – | | | | 317 |
| 331 | – | | – | | – | | | | – | | – | | – | | – | – | | 331 |
| 337 | | – | – | | | – | – | | – | | – | – | – | – | – | – | | 337 |
| 347 | – | | | – | | | | – | – | | | | | – | | | | 347 |
| 349 | | – | – | | | | | | | – | | – | – | | | | | 349 |
| 353 | | – | – | | | | | – | – | – | | | | – | – | | – | 353 |
| 359 | | | | – | – | | – | – | – | | | – | – | | – | – | – | 359 |
| 367 | – | | | | – | | | – | | – | | | – | | – | – | – | 367 |
| 373 | | | | | | – | | – | – | | – | – | – | – | | | – | 373 |
| 379 | – | | | | – | | – | – | | | – | – | | | | – | | 379 |
| 383 | | | – | | | – | | – | – | – | – | | – | – | – | – | – | 383 |
| 389 | – | | – | – | | – | | – | – | | | – | – | | – | – | | 389 |
| 397 | | – | – | | – | | | – | | – | – | – | | | – | – | – | 397 |
| 401 | | | | | | – | – | – | | – | – | – | – | – | | – | | 401 |
| 409 | | – | | – | | – | – | | – | – | – | – | – | – | | – | – | 409 |
| 419 | – | – | | | | – | – | | – | | – | – | – | | – | | – | 419 |
| 421 | – | | – | | | – | | – | | | – | – | | – | | – | | 421 |
| 431 | – | | – | | | | – | | – | – | | | | | – | – | – | 431 |
| 433 | | | | – | – | | – | – | – | – | – | – | | – | – | – | – | 433 |
| 439 | – | | – | | | – | – | | | – | | – | | | | – | – | 439 |
| 443 | – | – | | – | – | – | | – | – | | | – | – | | | | | 443 |
| 449 | – | – | | – | – | | – | | | | | – | | – | – | – | – | 449 |
| 457 | – | – | | – | | – | | | – | | – | – | – | – | | – | – | 457 |
| 461 | – | | | – | – | – | | | | – | | | | | – | | – | 461 |
| 463 | | | | – | – | – | | – | – | | | – | – | | | | – | 463 |
| 467 | | – | | | | – | | | – | | | | | – | | – | | 467 |
| 479 | | – | | | | – | | | | – | – | | | – | | | | 479 |
| 487 | | | | | | – | – | – | | – | – | | – | | – | – | – | 487 |
| 491 | | – | | | | | – | | – | | – | – | – | | | | | 491 |
| 499 | | | – | – | | | | | | – | – | – | | – | – | – | – | 499 |
| 503 | | | | | – | – | – | – | – | – | – | | | – | | – | | 503 |
| | 347 | 349 | 353 | 359 | 367 | 373 | 379 | 383 | 389 | 397 | 401 | 409 | 419 | 421 | 431 | 433 | 439 | |

| | 443 | 449 | 457 | 461 | 463 | 467 | 479 | 487 | 491 | 499 | 503 | 509 | 521 | 523 | 541 | 547 | 557 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 229 | – | – | – | – | – | – | | | – | | – | – | | | – | | – | 229 |
| 233 | – | | – | | | – | – | | – | – | – | | – | | | – | – | 233 |
| 239 | – | | – | | | | – | – | | | – | – | | – | | | | 239 |
| 241 | | | – | | | – | – | – | – | | | – | | – | – | | – | 241 |
| 251 | – | – | | | | | | | – | | – | – | | – | – | – | | 251 |
| 257 | | | – | | | | – | | – | – | – | | | – | | | | 257 |
| 263 | | – | | – | – | – | – | | | | | | – | | | | – | 263 |
| 269 | | – | – | | | | | – | – | | | | | | | – | | 269 |
| 271 | | – | | | | – | | | – | – | | – | – | – | | – | | 271 |
| 277 | | | | | – | – | – | – | – | – | | | | | – | – | – | 277 |
| 281 | – | | | – | | | | | | – | – | – | | – | | | – | 281 |
| 283 | – | | – | | | | – | – | – | – | | | – | – | | – | | 283 |
| 293 | – | – | | | – | | – | | – | – | – | – | – | | | – | | 293 |
| 307 | | | – | | – | – | | | – | | – | | | – | – | – | – | 307 |
| 311 | | | – | – | | – | – | | – | – | – | | – | – | | | | 311 |
| 313 | | | – | | – | | – | – | | | | – | – | – | – | – | | 313 |
| 317 | | | | – | | – | | | | – | – | | – | | | – | – | 317 |
| 331 | | – | – | – | – | | | – | | | – | | | | | – | | 331 |
| 337 | | – | | | – | | | – | | – | | – | | | | | – | 337 |
| 347 | | – | – | – | – | – | – | – | – | – | – | | | – | – | | | 347 |
| 349 | – | – | – | | | – | – | | – | | | | | | – | – | | 349 |
| 353 | | | | | | | | | | – | | – | – | | – | – | | 353 |
| 359 | | – | – | – | | – | – | – | – | | – | – | – | – | – | – | – | 359 |
| 367 | | – | | – | | – | – | – | – | – | | | | | – | | – | 367 |
| 373 | – | | – | – | – | – | – | – | | | – | | – | | | | – | 373 |
| 379 | – | – | | | – | – | – | | | – | | – | – | – | | | – | 379 |
| 383 | | | | | | – | – | | – | – | | – | – | | | | – | 383 |
| 389 | – | | – | | – | – | | | – | | – | – | | | | | – | 389 |
| 397 | | | | – | | | – | – | | – | – | | – | – | – | | – | 397 |
| 401 | | | – | | | | – | – | – | – | – | | | | – | – | – | 401 |
| 409 | – | – | – | | – | | | | – | – | | – | | | | – | | 409 |
| 419 | | | – | | | – | – | | | – | – | | – | | – | | | 419 |
| 421 | | – | – | | | – | – | | | – | – | | – | | | | | 421 |
| 431 | – | – | | – | – | – | – | | – | | – | | – | – | – | – | | 431 |
| 433 | | – | – | | | – | | – | | – | – | | – | | – | | | 433 |
| 439 | – | – | – | – | | – | – | | – | | – | – | | | – | | | 439 |
| 443 | – | | – | | | | – | | – | – | | – | | | | | – | 443 |
| 449 | | – | – | | – | – | | | | – | | | – | | – | – | | 449 |
| 457 | – | – | – | – | – | | | | – | – | | | – | | – | | – | 457 |
| 461 | | | – | – | | – | | – | – | | | | | – | – | | – | 461 |
| 463 | – | – | – | | – | – | – | | | – | | | – | – | – | – | – | 463 |
| 467 | – | – | | – | | – | – | | | | – | | | | – | | – | 467 |
| 479 | | | | | | | – | – | – | – | – | – | – | – | | | | 479 |
| 487 | – | | | – | – | – | | – | – | | – | | | – | | – | – | 487 |
| 491 | | | – | – | – | – | | | – | | – | | | | | – | | 491 |
| 499 | | – | – | | | – | | – | – | – | – | | – | – | | | | 499 |
| 503 | – | | | | – | | | | | | – | – | – | | | – | – | 503 |
| | 443 | 449 | 457 | 461 | 463 | 467 | 479 | 487 | 491 | 499 | 503 | 509 | 521 | 523 | 541 | 547 | 557 | |

| | 563 | 569 | 571 | 577 | 587 | 593 | 599 | 601 | 607 | 613 | 617 | 619 | 631 | 641 | 643 | 647 | 653 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 3 | | | – | – | | | | – | – | – | | – | – | | – | | | 3 |
| 5 | | – | – | | | | – | – | | | | – | – | – | | | | 5 |
| 7 | | – | – | | | | – | | | – | – | | – | – | | | – | 7 |
| 11 | | | | – | – | | – | | | | – | – | – | – | – | – | – | 11 |
| 13 | – | – | – | | | | – | – | – | | | | | – | | – | – | 13 |
| 17 | – | – | | – | – | – | – | | | – | | | – | | | – | | 17 |
| 19 | | | – | – | – | – | | | | – | – | – | – | | – | – | – | 19 |
| 23 | | | | – | – | – | – | – | – | | | | | | | – | – | 23 |
| 29 | | | – | | – | – | | | | – | | | – | | – | – | | 29 |
| 31 | – | | | – | | – | – | | – | | – | | | | | | – | 31 |
| 37 | | | – | | | – | – | – | | – | – | – | | – | | | | 37 |
| 41 | | – | | | | | – | | – | – | – | – | – | | | – | | 41 |
| 43 | – | – | | | | | – | | | – | – | – | | | – | | | 43 |
| 47 | | | – | | | | | – | | – | – | – | | | – | – | – | 47 |
| 53 | | | | – | – | – | – | | – | | | – | | | – | – | – | 53 |
| 59 | | | | – | | – | – | | – | | – | – | – | – | – | – | – | 59 |
| 61 | – | – | – | | | | | – | – | – | | – | | | | | | 61 |
| 67 | | – | – | | | | | – | – | – | – | – | | | – | | | 67 |
| 71 | | – | – | – | – | – | | | | – | – | | | – | – | – | | 71 |
| 73 | | | | | – | – | | | – | | | – | | – | | | – | 73 |
| 79 | – | – | – | | | – | – | | | | – | | | – | – | | – | 79 |
| 83 | – | | | | | – | | | – | | – | – | | | | | | 83 |
| 89 | | | | | – | | | – | – | – | | – | – | – | – | | | 89 |
| 97 | | | – | | | – | | | – | – | – | | – | | – | – | | 97 |
| 101 | – | – | | | – | – | | – | – | | | – | – | | – | | – | 101 |
| 103 | | | – | | – | | | | – | – | | – | – | – | – | – | | 103 |
| 107 | | – | – | – | – | | – | | | | | | | | – | | – | 107 |
| 109 | | | – | | | – | | | | | | – | | | | – | – | 109 |
| 113 | – | – | | | – | – | | – | | | – | | | | | – | – | 113 |
| 127 | | – | | – | – | | | | – | | | | | | – | | – | 127 |
| 131 | – | – | | – | – | | – | – | | – | | | – | – | | – | – | 131 |
| 137 | – | | | | – | | | | – | – | – | | | – | | – | – | 137 |
| 139 | – | – | | | – | – | | – | – | – | | – | | | | – | | 139 |
| 149 | – | | – | – | – | | | – | | – | | | – | – | – | | | 149 |
| 151 | – | – | – | – | | | | – | | – | | | | – | – | – | – | 151 |
| 157 | | | – | – | | – | | – | | | – | – | – | – | | – | – | 157 |
| 163 | – | | | – | | – | | | – | | | | | – | | – | – | 163 |
| 167 | – | | | – | | | – | – | | – | – | | – | | | | – | 167 |
| 173 | | | – | | | | | | – | | | – | | – | – | | | 173 |
| 179 | | | | | | – | | – | – | – | – | – | | | – | – | – | 179 |
| 181 | – | | | – | – | | – | | – | – | | | | | – | | | 181 |
| 191 | | | | – | | – | – | | – | – | | – | | – | | | – | 191 |
| 193 | – | | – | – | – | – | | | – | | | | | – | – | | | 193 |
| 197 | – | – | | | – | | | – | – | – | – | – | – | | | | – | 197 |
| 199 | – | | | | | | – | – | – | – | – | | | | – | – | – | 199 |
| 211 | | | | | | – | | – | – | | | | – | | | – | – | 211 |
| 223 | | | | – | | | – | | | | – | | | | – | – | | 223 |
| 227 | – | | | | – | – | | – | | – | | | – | | – | | | 227 |
| | 563 | 569 | 571 | 577 | 587 | 593 | 599 | 601 | 607 | 613 | 617 | 619 | 631 | 641 | 643 | 647 | 653 | |

| | 659 | 661 | 673 | 677 | 683 | 691 | 701 | 709 | 719 | 727 | 733 | 739 | 743 | 751 | 757 | 761 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 3 | | – | – | | | – | | – | | – | – | – | | – | – | | 3 |
| 5 | – | – | | | | – | – | – | – | | | – | | – | | – | 5 |
| 7 | – | | – | | – | | – | – | | | | – | – | – | – | | 7 |
| 11 | | – | | | – | – | | – | – | – | | | | – | – | | 11 |
| 13 | – | | – | – | | | – | | – | – | | | | – | – | | 13 |
| 17 | – | – | | | | | – | | | – | – | – | | | – | – | 17 |
| 19 | | | | | | – | – | – | – | – | – | – | | | – | – | 19 |
| 23 | | | – | | – | – | | | – | | | – | | | | – | 23 |
| 29 | | – | – | | – | – | – | – | – | | | | | | | – | 29 |
| 31 | – | – | | | – | – | – | | | – | – | | | – | | | 31 |
| 37 | – | | – | – | | – | | | – | | – | – | – | – | | – | 37 |
| 41 | | – | | – | | | – | | | | – | – | – | | | – | 41 |
| 43 | – | – | | | – | | – | – | – | | | | | | | | 43 |
| 47 | – | – | | | – | | | – | – | | – | – | | | | – | 47 |
| 53 | | – | – | | – | | | | | – | – | | – | – | – | | 53 |
| 59 | | – | | – | | | | – | | – | – | | – | | – | – | 59 |
| 61 | – | | | | – | – | | | – | – | – | | | – | – | | 61 |
| 67 | – | | | | | – | | – | – | | | | – | – | | – | 67 |
| 71 | – | | | – | | | | | – | | | – | | | | | 71 |
| 73 | – | – | – | | | | | | | – | – | – | | | – | | 73 |
| 79 | | | | – | – | | | | – | – | – | | – | – | – | – | 79 |
| 83 | – | | – | | | – | – | | | – | – | – | | – | – | | 83 |
| 89 | – | | – | | | – | – | | | | – | | | – | – | – | 89 |
| 97 | | – | – | – | – | – | – | | | – | – | | – | – | | | 97 |
| 101 | | | | – | – | – | – | | | – | | | – | | | – | 101 |
| 103 | – | | – | – | | | – | – | | | | – | | – | – | | 103 |
| 107 | | – | | – | – | – | | | | – | | | – | | | – | 107 |
| 109 | – | – | | | – | | | | | – | | | – | – | | | 109 |
| 113 | | | | – | | – | | – | – | – | | – | | | | – | 113 |
| 127 | | – | – | – | | | | – | – | | – | – | | | – | | 127 |
| 131 | – | | | | – | – | – | | – | | | – | | | – | | 131 |
| 137 | | | | – | – | | – | | – | | | | | | – | – | 137 |
| 139 | | | – | – | – | | – | | – | | – | – | | | | – | 139 |
| 149 | – | | | – | | – | | – | – | | | – | | – | | – | 149 |
| 151 | – | | – | | | | – | – | | – | | | – | | – | | 151 |
| 157 | – | – | | – | | | | – | | – | – | – | – | | | | 157 |
| 163 | | – | – | – | | – | – | – | | | – | – | – | | | | 163 |
| 167 | | | | – | | | – | | | | – | | – | | – | – | 167 |
| 173 | – | – | | – | – | – | – | | | – | – | – | – | | | | 173 |
| 179 | | – | | | – | | | – | – | | – | | – | | | – | 179 |
| 181 | – | | | | | – | | – | – | – | – | – | | – | – | – | 181 |
| 191 | – | | – | – | | – | – | – | | – | – | | – | | – | | 191 |
| 193 | | | | – | | – | | – | | | | | | – | | | 193 |
| 197 | | – | | | – | – | | | | – | – | – | | – | | | 197 |
| 199 | – | – | | – | – | – | – | – | – | – | | | | | – | | 199 |
| 211 | | | | – | | – | | – | | | – | | | | | | 211 |
| 223 | – | | – | – | – | | – | | – | – | – | | – | – | | | 223 |
| 227 | – | – | – | | | – | | – | | | | | – | – | – | | 227 |
| | 659 | 661 | 673 | 677 | 683 | 691 | 701 | 709 | 719 | 727 | 733 | 739 | 743 | 751 | 757 | 761 | |

| | 563 | 569 | 571 | 577 | 587 | 593 | 599 | 601 | 607 | 613 | 617 | 619 | 631 | 641 | 643 | 647 | 653 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 229 | | – | | | – | – | | | – | | – | – | – | – | – | | | 229 |
| 233 | | | – | | – | | – | – | – | | | | | – | – | – | – | 233 |
| 239 | – | – | – | – | – | | – | | | – | | | – | – | – | – | | 239 |
| 241 | – | – | | | | | | – | – | | – | | | – | – | | | 241 |
| 251 | | – | – | – | – | – | | | – | | – | – | | | | | | 251 |
| 257 | – | | – | | – | – | | | | – | | | – | | – | – | – | 257 |
| 263 | – | – | | – | – | | | – | – | | | – | – | | – | – | | 263 |
| 269 | – | | | | – | – | – | | | | – | – | – | – | – | | – | 269 |
| 271 | | | | – | – | | – | | | | | – | – | – | | | | 271 |
| 277 | – | | | – | | – | | – | | – | – | | | – | – | | | 277 |
| 281 | – | – | – | | – | – | | – | – | | | – | – | – | – | – | | 281 |
| 283 | – | | | – | | | | | – | | – | | | | – | – | | 283 |
| 293 | | – | – | – | – | | | – | – | | – | – | | – | – | – | – | 293 |
| 307 | – | – | – | | – | – | | – | | | | | – | | | | – | 307 |
| 311 | – | | – | | | – | – | | | | | | – | | – | – | | 311 |
| 313 | | – | | – | – | – | – | – | – | – | – | | | | | | – | 313 |
| 317 | | – | – | – | | | | | | | | – | | – | – | | | 317 |
| 331 | – | – | | | – | | – | – | – | | | | | | | – | | 331 |
| 337 | – | | – | | | – | – | | | | | – | – | | | – | – | 337 |
| 347 | | – | – | | | – | | | | | – | | – | | – | – | – | 347 |
| 349 | – | | | – | – | | | | – | – | – | | – | – | | – | – | 349 |
| 353 | | | – | | | – | | | – | – | | | | – | | | | 353 |
| 359 | – | | | | | | – | – | | – | | | – | – | | – | – | 359 |
| 367 | – | – | – | – | – | – | | – | – | | | – | – | – | | | | 367 |
| 373 | | – | – | – | | – | – | | | | | | – | | – | | – | 373 |
| 379 | | | | – | | | | – | | – | | | | – | | – | | 379 |
| 383 | | – | | | – | | – | | – | | | | – | – | – | | | 383 |
| 389 | | – | | | | | – | | – | | | – | | – | | | – | 389 |
| 397 | | – | | | – | – | – | | | | | | – | | – | – | – | 397 |
| 401 | – | | | – | – | | – | – | – | | | – | | | – | | – | 401 |
| 409 | – | – | – | | – | – | | – | | – | | | | | | | | 409 |
| 419 | – | | – | | | | – | | – | | – | | | | | | | 419 |
| 421 | – | | | | – | | – | – | | – | – | | | – | – | | – | 421 |
| 431 | – | – | | | | – | | | – | – | | | – | | – | – | | 431 |
| 433 | | – | | – | | | | | | | | | – | – | – | | | 433 |
| 439 | | – | | – | | – | – | – | | | | – | | | – | – | | 439 |
| 443 | – | – | | | – | | – | | – | – | – | | – | – | | – | | 443 |
| 449 | – | | – | – | | – | | | | – | | | | | – | – | – | 449 |
| 457 | | – | – | | – | – | | – | – | | | – | – | | | | – | 457 |
| 461 | – | | – | | – | – | – | | | | – | – | | – | | | | 461 |
| 463 | – | | | – | – | – | – | – | – | | – | – | – | – | | | – | 463 |
| 467 | | | | | – | | | – | | – | | – | – | – | | | – | 467 |
| 479 | – | – | – | – | – | | – | – | – | | – | – | | – | | – | | 479 |
| 487 | – | – | – | | – | – | | | – | | – | – | – | – | – | | – | 487 |
| 491 | | | – | | | | – | | | | – | | | | – | – | | 491 |
| 499 | – | – | | | – | | – | | | | – | – | – | – | – | | | 499 |
| 503 | | – | | | – | | – | – | – | | | | – | – | | – | – | 503 |
| | 563 | 569 | 571 | 577 | 587 | 593 | 599 | 601 | 607 | 613 | 617 | 619 | 631 | 641 | 643 | 647 | 653 | |

| | 659 | 661 | 673 | 677 | 683 | 691 | 701 | 709 | 719 | 727 | 733 | 739 | 743 | 751 | 757 | 761 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 229 | | – | – | | – | – | – | | | | – | | – | – | – | | 229 |
| 233 | | – | – | | – | – | – | | | – | | | | – | – | – | 233 |
| 239 | | – | | | | – | | | – | – | – | – | | – | – | – | 239 |
| 241 | – | | – | | – | – | | | – | – | – | – | – | | | | 241 |
| 251 | | | | – | – | – | | – | – | – | | – | – | – | – | | 251 |
| 257 | | | – | | – | | – | – | – | – | | – | | | | | 257 |
| 263 | – | | – | – | – | | | – | | | – | | | – | | – | 263 |
| 269 | – | | | | | | | | | – | | | – | – | | | 269 |
| 271 | | – | | | – | | | – | – | – | | | | | | – | 271 |
| 277 | | | | – | | | – | – | – | | | – | – | | – | – | 277 |
| 281 | | | – | | – | | | | – | – | | | – | | – | | 281 |
| 283 | – | – | | – | – | | – | – | | – | | | | – | | – | 283 |
| 293 | – | | – | – | – | | – | – | – | – | | – | | | | | 293 |
| 307 | | | | – | – | – | – | | – | – | – | | – | | | | 307 |
| 311 | | – | | | | | – | | | – | | – | – | | – | – | 311 |
| 313 | – | – | | | – | | – | – | | | – | – | – | | | | 313 |
| 317 | – | | – | – | – | – | – | | – | | – | – | | | – | | 317 |
| 331 | – | | | | – | | – | | | | – | – | – | – | – | | 331 |
| 337 | | – | – | – | – | | – | | | | | | | | | | 337 |
| 347 | | | – | – | | | | | – | – | – | | – | | | – | 347 |
| 349 | | – | – | | – | | – | | | – | | – | – | | | | 349 |
| 353 | – | | | – | – | – | | | | – | | | | | | | 353 |
| 359 | – | – | | | – | | | | – | – | – | | – | – | | | 359 |
| 367 | – | | | | | – | | | | | | | – | | – | | 367 |
| 373 | – | | – | | – | – | | – | – | | – | – | – | | | | 373 |
| 379 | – | | – | | – | | – | | | – | | | – | – | | | 379 |
| 383 | – | – | | – | – | | | | – | – | – | | | – | | – | 383 |
| 389 | – | – | | | – | – | – | – | – | | – | | – | | | – | 389 |
| 397 | | | – | | – | | – | – | – | – | | | | – | – | – | 397 |
| 401 | | | | – | | – | | – | – | | – | – | | – | – | – | 401 |
| 409 | – | | | | – | | | – | | – | – | | – | | | | 409 |
| 419 | – | | – | | – | | | | – | | | – | – | | | – | 419 |
| 421 | | – | – | – | | | | | | | – | – | – | | – | – | 421 |
| 431 | – | – | – | – | | | – | – | – | | – | | | – | – | – | 431 |
| 433 | | | | | | | | | – | – | – | – | – | – | – | – | 433 |
| 439 | – | – | – | | – | – | | | – | – | | | – | | | | 439 |
| 443 | | | | | | – | – | | | – | | | – | – | – | – | 443 |
| 449 | | – | – | – | | – | – | | | | | | | | – | – | 449 |
| 457 | | – | – | – | | | | – | – | | | – | – | – | – | – | 457 |
| 461 | – | | – | – | | | | – | – | – | – | | | – | – | | 461 |
| 463 | – | | | | – | – | | – | – | – | – | – | – | – | | – | 463 |
| 467 | – | | – | – | | | | | – | | – | – | – | – | | | 467 |
| 479 | – | | – | – | | | | – | – | | | | – | | – | | 479 |
| 487 | | | | | – | – | – | – | – | – | | | – | – | – | – | 487 |
| 491 | – | | – | | – | | – | – | | | | | | – | | | 491 |
| 499 | | | – | – | – | | | | | – | – | | | | – | | 499 |
| 503 | – | – | – | | | – | – | | – | – | | – | | | | – | 503 |
| | 659 | 661 | 673 | 677 | 683 | 691 | 701 | 709 | 719 | 727 | 733 | 739 | 743 | 751 | 757 | 761 | |

| | 769 | 773 | 787 | 797 | 809 | 811 | 821 | 823 | 827 | 829 | 839 | 853 | 857 | 859 | 863 | 877 | 881 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 3 | – | | – | | | – | | – | | – | | – | | – | | – | | 3 |
| 5 | – | | | | – | – | – | | | – | – | | | – | | | – | 5 |
| 7 | | | | | – | | – | – | – | | | | | | – | – | | 7 |
| 11 | | – | | – | | | | – | | – | – | | | – | – | | – | 11 |
| 13 | | | | – | – | | | – | | – | | | – | – | | | – | 13 |
| 17 | – | – | | – | | | | | | – | | | | – | – | | | 17 |
| 19 | – | | | | – | | – | – | | | | – | | – | | | – | 19 |
| 23 | | | | | – | – | – | – | | – | | – | – | – | – | – | | 23 |
| 29 | | | – | | | – | – | | | | | | – | | – | – | | 29 |
| 31 | – | | | | | – | | | | | – | – | – | | | – | | 31 |
| 37 | | – | – | | | – | – | – | | | – | | | | – | – | – | 37 |
| 41 | – | | – | – | | – | – | | | – | | – | – | – | – | – | – | 41 |
| 43 | – | | – | – | – | | – | – | – | | | – | – | | | – | – | 43 |
| 47 | – | – | | | | – | | – | – | | | – | | | – | | | 47 |
| 53 | | | | | | – | | – | | | – | | – | – | – | – | | 53 |
| 59 | | | – | | | | | | – | – | | – | | | | – | | 59 |
| 61 | | – | | – | – | | | | – | – | – | – | – | – | – | | – | 61 |
| 67 | | – | | – | | | – | – | – | – | – | – | | – | – | – | – | 67 |
| 71 | | | – | – | | – | – | | | – | – | – | – | | | – | – | 71 |
| 73 | | | – | – | – | – | – | | – | | – | – | – | | | – | | 73 |
| 79 | | – | – | | – | – | – | | | | – | | – | | – | – | | 79 |
| 83 | | – | – | | | – | | | | | – | – | – | – | – | | – | 83 |
| 89 | – | | | – | – | – | – | – | | | | | | | | | – | 89 |
| 97 | | – | – | | – | – | | – | | – | | | – | | | – | – | 97 |
| 101 | | | – | | – | | – | | – | – | – | – | – | | | | | 101 |
| 103 | | – | – | – | | | – | | | | – | – | – | | | | | 103 |
| 107 | | | | – | | – | | | | | – | | – | – | | | – | 107 |
| 109 | | | | – | – | – | | – | – | – | | | – | | – | – | – | 109 |
| 113 | – | – | – | | – | | – | – | – | | | – | | | – | | | 113 |
| 127 | | – | – | – | – | – | | – | | | | | | | | – | | 127 |
| 131 | – | | – | – | | – | – | | – | – | – | | | | – | – | | 131 |
| 137 | | – | | – | | – | – | – | | – | – | | | – | | | – | 137 |
| 139 | | – | | | | – | | | | | – | | | – | – | | – | 139 |
| 149 | – | – | – | | – | | – | | – | | | | – | – | – | – | | 149 |
| 151 | | – | – | – | | | | – | – | – | – | – | | | | | | 151 |
| 157 | – | – | | – | | | – | | – | – | | – | | | | | | 157 |
| 163 | | – | – | – | | | – | | | – | – | – | | | | – | | 163 |
| 167 | | | | | – | | | | | | – | – | – | – | – | – | | 167 |
| 173 | – | – | – | | – | – | | | – | – | | | | – | | | – | 173 |
| 179 | | – | | – | – | – | | – | | | | | – | | – | – | | 179 |
| 181 | – | – | | – | | – | | – | | | | – | – | – | – | | | 181 |
| 191 | – | – | – | | – | | | – | | – | – | | | | | | – | 191 |
| 193 | – | – | | – | | | – | | – | | – | – | – | | | | – | 193 |
| 197 | – | – | – | – | | – | – | | – | – | – | – | | | | | – | 197 |
| 199 | – | | | – | – | | – | | – | – | – | – | – | – | | – | | 199 |
| 211 | – | | – | | – | – | – | | – | – | | – | – | | – | | – | 211 |
| 223 | – | | | – | | | – | | | | | | – | | | | – | 223 |
| 227 | | – | | – | | | | | | | | – | – | | – | – | | 227 |
| | 769 | 773 | 787 | 797 | 809 | 811 | 821 | 823 | 827 | 829 | 839 | 853 | 857 | 859 | 863 | 877 | 881 | |

| | 883 | 887 | 907 | 911 | 919 | 929 | 937 | 941 | 947 | 953 | 967 | 971 | 977 | 983 | 991 | 997 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 3 | – | | – | | – | | – | | | | – | | | | – | – | 3 |
| 5 | | | | – | – | – | | – | | | | – | | | – | | 5 |
| 7 | – | | – | – | – | | | | – | – | – | | – | | – | | 7 |
| 11 | – | | – | – | | – | | | – | | | – | – | – | – | | 11 |
| 13 | – | – | – | – | – | | – | | | – | | – | | | – | – | 13 |
| 17 | – | | | | – | | – | | | – | – | – | – | | | | 17 |
| 19 | – | | | | – | – | – | | – | | – | | | | | – | 19 |
| 23 | – | – | | | | – | | | – | | – | | | | – | – | 23 |
| 29 | – | | | | – | – | – | – | | – | | | – | | – | | 29 |
| 31 | | – | – | | – | | – | | | | | – | – | | | – | 31 |
| 37 | | – | | | | – | – | – | | – | | – | | – | | | 37 |
| 41 | | | – | – | | | | – | – | – | | | | – | | | 41 |
| 43 | – | | – | | – | | | – | – | | – | – | – | | | | 43 |
| 47 | – | | – | – | | – | | – | – | | – | | – | | – | | 47 |
| 53 | | | – | – | | – | – | – | – | – | – | – | | – | – | – | 53 |
| 59 | – | | – | – | | | | | – | – | | – | | | | – | 59 |
| 61 | | | | – | – | – | – | | | | – | – | – | | – | | 61 |
| 67 | | – | – | – | | | | | – | – | – | – | – | | | – | 67 |
| 71 | | | | | | – | | – | – | – | | – | – | – | | – | 71 |
| 73 | | | | – | | | – | – | – | – | – | | | | | – | 73 |
| 79 | | – | – | – | – | | | – | | – | – | – | | | | – | 79 |
| 83 | | | – | – | | – | | – | | – | | | – | – | – | – | 83 |
| 89 | | | – | – | | – | – | | – | | | – | – | – | | – | 89 |
| 97 | | | | | | | – | | | | – | – | | | | – | 97 |
| 101 | | – | | | | – | | | | | – | | – | | – | – | 101 |
| 103 | – | – | – | | | – | | – | | – | | | – | – | – | | 103 |
| 107 | – | | | | | | – | – | | | – | | – | | | – | 107 |
| 109 | | – | – | | | | | | – | – | | | – | | | – | 109 |
| 113 | | | | – | – | – | | | | – | – | | | | – | | 113 |
| 127 | – | | – | – | – | | | – | | – | | – | – | – | | | 127 |
| 131 | | – | – | – | | – | – | | | – | | | – | | – | – | 131 |
| 137 | – | – | | | | – | – | – | | | – | | – | | – | – | 137 |
| 139 | – | | | – | | | | – | – | | | – | – | | | – | 139 |
| 149 | | – | | – | – | – | | – | – | | – | | | | | – | 149 |
| 151 | – | | – | – | | | – | | | – | | | | | – | – | 151 |
| 157 | | | – | – | | – | | – | | – | – | | – | | – | | 157 |
| 163 | | | | – | – | | | – | – | | – | – | | | | | 163 |
| 167 | – | | – | – | – | – | | | – | | – | | | | | – | 167 |
| 173 | | – | | | – | – | | | | – | | – | | – | – | – | 173 |
| 179 | | – | – | – | | | – | – | – | | | – | – | – | | | 179 |
| 181 | | | | | – | | | – | – | – | – | | | | | | 181 |
| 191 | | | | – | | | | – | | | – | – | | | – | | 191 |
| 193 | | | | – | – | – | – | – | – | – | – | – | – | – | | – | 193 |
| 197 | | | | | | | | | | | | | | | – | | 197 |
| 199 | | – | – | – | – | | | – | – | – | | – | | – | | – | 199 |
| 211 | | – | | | | | – | | – | – | – | | | – | | | 211 |
| 223 | | – | – | – | | – | | – | – | | | | | | | – | 223 |
| 227 | | | | – | – | – | – | – | | | – | – | – | – | | – | 227 |
| | 883 | 887 | 907 | 911 | 919 | 929 | 937 | 941 | 947 | 953 | 967 | 971 | 977 | 983 | 991 | 997 | |

| | 769 | 773 | 787 | 797 | 809 | 811 | 821 | 823 | 827 | 829 | 839 | 853 | 857 | 859 | 863 | 877 | 881 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 229 | – | | – | | | | – | | | | | | | – | – | | | 229 |
| 233 | | – | | – | – | – | | – | – | | | | | | | – | – | 233 |
| 239 | | | | – | | | | | – | | – | – | | – | | – | | 239 |
| 241 | | – | – | | | | – | – | | – | – | | – | | | – | – | 241 |
| 251 | – | – | | | | – | – | | – | | – | – | | – | – | – | | 251 |
| 257 | – | – | – | – | | | – | – | | – | – | | | – | – | | | 257 |
| 263 | – | | | – | | – | – | – | | | – | – | – | – | – | – | – | 263 |
| 269 | – | – | – | | | – | – | – | – | | | | | – | – | – | | 269 |
| 271 | | | – | | | | – | – | – | – | | – | – | | – | – | – | 271 |
| 277 | – | | | – | – | | – | | – | | | – | | – | | | | 277 |
| 281 | | – | – | | – | – | | – | – | – | – | – | – | – | – | – | | 281 |
| 283 | – | – | | | | | | – | – | – | | – | | – | | – | | 283 |
| 293 | | | | – | | – | – | – | | | – | – | – | | – | | | 293 |
| 307 | – | | | – | | | | – | | – | – | | | | | | | 307 |
| 311 | – | | | – | – | – | | – | | | | | – | – | | | | 311 |
| 313 | – | – | – | – | | – | | – | | | – | | | | – | | – | 313 |
| 317 | – | | | | – | | | | – | | – | | – | – | | | – | 317 |
| 331 | | – | – | | | – | | – | – | – | – | – | – | – | | – | – | 331 |
| 337 | – | | – | – | | – | – | – | | – | – | – | | | – | | | 337 |
| 347 | – | | – | | – | – | – | – | – | | | – | | | – | – | | 347 |
| 349 | | – | | | – | | – | – | | | | – | | | | | | 349 |
| 353 | | | – | – | | | | | – | | | | | – | – | – | | 353 |
| 359 | – | – | – | – | – | | | | | – | – | – | | – | | | | 359 |
| 367 | | | – | – | | | – | – | | – | – | | | | – | | | 367 |
| 373 | | – | – | – | – | | – | | – | – | – | – | – | | – | | | 373 |
| 379 | | | | – | – | | | | | | – | – | | – | – | – | | 379 |
| 383 | – | – | – | – | – | | – | – | | – | – | – | | – | | | | 383 |
| 389 | – | – | – | – | | | | – | – | | | | – | – | – | – | | 389 |
| 397 | – | | | – | | | – | – | – | – | | | | – | – | – | – | 397 |
| 401 | | – | | – | – | – | | | – | | | – | – | – | | | | 401 |
| 409 | – | – | | | – | | – | – | – | | | | | – | – | | | 409 |
| 419 | | | – | | | | – | | – | | – | – | | – | – | – | – | 419 |
| 421 | | | – | – | – | – | – | | – | | – | – | – | – | – | – | | 421 |
| 431 | – | – | | – | | – | | – | – | | | | | | – | – | – | 431 |
| 433 | | | – | | | | | | – | – | – | – | – | | – | – | | 433 |
| 439 | | – | | | | – | – | | | | – | | – | | – | | | 439 |
| 443 | | | – | | | | – | – | | – | | – | – | | – | | – | 443 |
| 449 | – | – | – | – | – | – | – | | | | – | – | – | – | – | | | 449 |
| 457 | | – | – | | | | | | – | | – | | – | – | – | | | 457 |
| 461 | – | | | – | – | – | | | – | – | | | | | – | – | – | 461 |
| 463 | – | | – | – | – | | – | | – | | – | | | | – | | – | 463 |
| 467 | | | | | – | | | – | – | – | | | – | | | – | | 467 |
| 479 | | – | – | | – | | | | | – | – | | – | | – | | | 479 |
| 487 | – | – | | | – | – | – | – | – | – | | | – | | | | – | 487 |
| 491 | | – | | | – | – | | – | | | | | | | – | – | | 491 |
| 499 | – | | | – | | | | – | – | | | | – | | – | | – | 499 |
| 503 | | | | – | | – | | | – | | – | – | – | | | | – | 503 |
| | 769 | 773 | 787 | 797 | 809 | 811 | 821 | 823 | 827 | 829 | 839 | 853 | 857 | 859 | 863 | 877 | 881 | |

| | 883 | 887 | 907 | 911 | 919 | 929 | 937 | 941 | 947 | 953 | 967 | 971 | 977 | 983 | 991 | 997 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 229 | – | | – | – | – | | | – | | – | – | – | – | | – | – | 229 |
| 233 | – | | – | | – | | | – | – | | | | | – | | | 233 |
| 239 | – | – | | | – | | – | | | | – | – | | – | | | 239 |
| 241 | – | – | | – | – | | – | | | | – | | | | – | | 241 |
| 251 | | | – | | | | | | – | | – | – | | | | | 251 |
| 257 | | – | – | – | | – | | | – | | – | – | | | | – | 257 |
| 263 | | – | | – | | – | – | | | | – | | | | | – | 263 |
| 269 | | – | – | | | | | | | | | – | – | – | | – | 269 |
| 271 | – | – | | – | – | | – | – | – | – | – | – | – | – | – | | 271 |
| 277 | – | | – | | – | | – | | – | – | – | | – | | – | | 277 |
| 281 | – | | – | – | | – | | – | | | – | – | | – | | | 281 |
| 283 | – | – | | – | – | | | – | | | | | | – | | | 283 |
| 293 | – | | | | – | | – | | – | | – | | | – | | | 293 |
| 307 | – | – | – | | – | | – | | – | | – | | | – | – | – | 307 |
| 311 | | – | | – | | | – | – | – | – | | | | – | | – | 311 |
| 313 | | – | – | | | | – | – | – | | | – | – | – | – | – | 313 |
| 317 | | – | – | – | | | | – | – | | – | | – | | – | | 317 |
| 331 | | – | – | | | | | – | | – | | | | | – | – | 331 |
| 337 | – | | – | – | | – | – | | – | | | | | – | | – | 337 |
| 347 | | | – | | – | – | – | | | – | | – | | – | – | | 347 |
| 349 | – | | | | | – | – | – | – | – | – | – | – | – | – | | 349 |
| 353 | – | – | – | | – | – | – | | | | – | – | – | – | – | | 353 |
| 359 | – | – | | – | | | – | | – | – | | – | | | – | | 359 |
| 367 | – | | – | | | – | | – | – | | | – | | | | | 367 |
| 373 | – | | | – | | | | | | | – | – | | | | – | 373 |
| 379 | – | – | – | | | – | – | | – | – | | – | – | – | | – | 379 |
| 383 | | – | | | – | | – | – | | | – | – | | – | – | | 383 |
| 389 | | | – | – | – | | – | | – | – | | – | | – | – | | 389 |
| 397 | | – | | | | | | – | | | – | | | | | | 397 |
| 401 | – | | | – | | | | | – | – | | – | – | – | | – | 401 |
| 409 | | – | – | | – | | | – | | – | | – | – | | | – | 409 |
| 419 | – | – | – | – | – | – | | | | – | – | | – | – | | | 419 |
| 421 | | – | | | – | | | – | – | | – | | – | | – | – | 421 |
| 431 | | – | – | – | – | | – | | | – | | – | – | – | | – | 431 |
| 433 | – | | – | | – | | | – | – | | | – | – | – | | | 433 |
| 439 | – | – | – | | | – | | – | – | | | – | – | | – | | 439 |
| 443 | | – | | – | | | – | – | – | – | – | | | | – | – | 443 |
| 449 | | – | – | | | | – | | – | – | | | | | – | – | 449 |
| 457 | | – | – | – | | | | – | | | | – | – | | | | 457 |
| 461 | – | | | | | | | – | – | | – | – | | – | | | 461 |
| 463 | | | – | | – | | | – | – | | | | | – | – | | 463 |
| 467 | | | | | – | – | – | – | – | | | | – | – | | – | 467 |
| 479 | | | – | – | – | – | | – | | | – | | | – | – | | 479 |
| 487 | | – | | – | | – | – | | – | | – | – | | – | | | 487 |
| 491 | | – | | | – | – | | | – | – | | | | – | – | – | 491 |
| 499 | – | | | – | – | | – | | | | | – | | – | – | | 499 |
| 503 | – | – | | | – | | | – | | – | | – | – | | – | | 503 |
| | 883 | 887 | 907 | 911 | 919 | 929 | 937 | 941 | 947 | 953 | 967 | 971 | 977 | 983 | 991 | 997 | |

# TABLE

FOR CONVERTING

COMMON FRACTIONS WITH DENOMINATORS FROM THE FIRST THOUSAND

INTO DECIMAL FRACTIONS.

3 (1)..6; (0)..3

7 (0)..428571

9 (1)..2; (2)..4; (3)..8; (4)..7; (5)..5; (0)..1

11 (1)..81; (2)..63; (3)..27; (4)..54; (0)..90

13 (1)..615384; (0)..769230

17 (0)..5882352941 176470

19 (0)..5263157894 73684210

23 (0)..4347826086 9565217391 30

27 (1)..740; (2)..481; (3)..962; (4)..925; (5)..851; (0)..370

29 (0)..3448275862 0689655172 41379310

31 (1)..4838709677 41935; (0)..3225806451 61290

37 (1)..351; (2)..756; (3)..783; (4)..918; (5)..594; (6)..972;
(7)..864; (8)..324; (9)..621; (10)..108; (11)..540; (0)..270

41 (1)..46341; (2)..78048; (3)..68292; (4)..09756; (5)..58536;
(6)..51219; (7)..07317; (0)..24390

43 (1)..5116279069 7674418604 6; (0)..2325581395 3488372093 0

47 (1)..2127659574 4680851063 8297872340 4255319148 936170

49 (0)..2040816326 5306122448 9795918367 3469387755 10

53 (1)..9056603773 584; (2)..5471698113 207; (3)..2264150943 396;
(0)..1886792452 830

59 (0)..1694915254 2472881355 9322033898 3050847457 6271186440
67796610

61 (0)..1639344262 2950819672 1311475409 8360655737 7049180327
8688524590

67 (1)..7910447761 1940298507 4626865671 641; (0)..1492537313
4328358208 9552238805 970

71 (1)..7323943661 9718309859 1549295774 64788; (0)..1408450704
2253521126 7605633802 81690

73 (1)..68493150; (2)..42465753; (3)..12328767; (4)..61643835;
(5)..08219178; (6)..41095890 (7)..05479452; (8)..27397260;
(0)..13698630

79 (1)..6708860759 493; (2)..4556962025 316; (3)..2151898734 177;
(4)..2405063291 139; (5)..9746835443 037; (0)..1265822784 810

81 (1)..358024691; (2)..938271604; (3)..320987654; (4)..530864197;
(5)..839506172; (0)..123456790

83 (1)..0240963855 4216867469 8795180722 8915662650 6;
(0)..1204819277 1084337349 3975903614 4578313253 0

89 (1)..3707865168 5393258426 9662921348 3146067415 7303;
(0)..1123595505 6179775280 8988764044 9438202247 1910

97 (0)..1030927835 0515463917 5257731958 7628865979 3814432989
6907216494 8453608247 4226804123 7113402061 855670

101 (1)..1980; (2)..3960; (3)..7920; (4)..5841; (5)..1683; (6)..3366;
(7)..6732; (8)..3465; (9)..6930; (10)..3861; (11)..7722;
(12)..5445; (13)..0891; (14)..1782; (15)..3564; (16)..7128;
(17)..4257; (18)..8514; (19)..7029; (20)..4059; (21)..8118;
(22)..6237; (23)..2475; (24)..4950; (0)..0990

103 (1)..5825242718 4466019417 4757281553 3980; (2)..4951456310
6796116504 8543689320 3883 (0)..0970873786 4077669902
9126213592 2330

107 (1)..8878504672 8971962616 8224299065 4205607476 6355140186 915
(0)..0934579439 2523364485 9813084112 1495327102 8037383177 570

109 (0)..0917431192 6605504587 1559633027 5229357798 1651376146
7889908256 8807339449 5412844036 6972477064 2201834862 38532110

113 (0)..0884955752 2123893805 3097345132 7433628318 5840707964
6017699115 0442477876 1061946902 6548672566 3716814159
2920353982 30

121 (1)..8925619834 7107438016 52; (2)..2396694214 8760330578 51;
(3)..3884297520 6611570247 93; (4)..5950413223 1404958677 68;
(0)..0826446280 9917355371 90;

127 (1)..3464566929 1338582677 1653543307 0866141732 28;
(2)..7244094488 1889763779 5275590551 1811023622 04;
(0)..0787401574 8031496062 9921259842 5196850393 70

131 (0)..0763358778 6259541984 7328244274 8091603053 4351145038
1679389312 9770992366 4122137404 5801526717 5572519083
9694656488 5496183206 1068702290

137 (1)..87591240; (2)..51094890; (3)..13138686; (4)..57664233;
(5)..91970802; (6)..03649635; (7)..43795620; (8)..25547445;
(9)..06569343; (10)..78832116; (11)..45985401; (12)..51824817;
(13)..21897810; (14)..62773722; (15)..53284671; (16)..39416058;
(0)..07299270

139 (1)..6187050359 7122302158 2733812949 6402877697 841726;
(2)..9208633093 5251798561 1510791366 9064748201 438848;
(0)..0719424460 4316546762 5899280575 5395683453 237410

149 (0)..0671140939 5973154362 4161073825 5033557046 9798657718
1208053691 2751677852 3489932885 9060402684 5637583892
6174496644 2953020134 2281879194 6308724832 21476510

151 (1)..5496688741 7218543046 3576158940 3973509933 7748344370
8609271523 1788079470 19867 (0)..0662251655 6291390728
4768211920 5298013245 0331125827 8145695364 2384105960 26490

157 (1)..1464968152 8662420382 1656050955 4140127388 5350318471
3375796178 3439490445 85987261 (0)..0636942675 1592356687
8980891719 7452229299 3630573248 4076433121 0191082802 54777070

163 (1)..2944785276 0736196319 0184049079 7546012269 9386503067
4846625766 8711656441 7177914110 4 (0)..0613496932 5153374233
1288343558 2822085889 5705521472 3926380368 0981595092
0245398773 0

167 (0)..0598802395 2095808383 2335329341 3173652694 6107784431
1377245508 9820359281 4371257485 0299401197 6047904191
6167664670 6586826347 3053892215 5688622754 4910179640
7185628742 514970

169 (1)..1065088757 3964497041 4201183431 9526627218 9349112426
0355029585 7988165680 47337278 (0)..0591715976 3313609467
4556213017 7514792899 4082840236 6863905325 4437869822 48520710

173 (1)..7398843930 6358381502 8901734104 0462427745 664;
(2)..6705202312 1387283236 9942196531 7919075144 508;
(3)..9826589595 3757225433 5260115606 9364161849 710;
(0)..0578034682 0809248554 9132947976 8786127167 630

179 (0)..0558659217 8770949720 6703910614 5251396648 0446927374
3016759776 5363128491 6201117318 4357541899 4413407821
2290502793 2960893854 7486033519 5530726256 9832402234
6368715083 7988826815 64245810

181 (0)..0552486187 8453038674 0331491712 7071823204 4198895027
6243093922 6519337016 5745856353 5911602209 9447513812
1546961325 9668508287 2928176795 5801104972 3756906077
3480662983 4254143646 4088397790

191 (1)..2198952879 5811518324 6073298429 3193717277 4869109947
6439790575 9162303664 9214659685 8638743455 49738;
(0)..0523560209 4240837696 3350785340 3141361256 5445026178
0104712041 8848167539 2670157068 0628272251 30890

193 (0)..0518134715 0259067357 5129533678 7564766839 3782383419
6891191709 8445595854 9222797927 4611398963 7305699481
8652849740 9326424870 4663212435 2331606217 6165803108
8082901554 4041450777 2020725388 6010362694 30

197 (1)..7055837563 4517766497 4619289340 1015228426 3959390862
9441624365 4822335025 3807106598 9847715736 04060913;
(0)..0507614213 1979695431 4720812182 7411167512 6903553299
4923857868 0203045685 2791878172 5888324873 09644670

199 (1) . . 3819095477 3869346733 6683417085 4271356783 9195979899 4974874371 8592964824 1206030150 7537688442 211055276; (0) . . 0502512562 8140703517 5879396984 9246231155 7788944723 6180904522 6130653266 3316582914 5728643216 080402010

211 (1) . . 3317535545 0236966824 6445497630; (2) . . 3222748815 1658767772 5118483412; (3) . . 2559241706 1611374407 5829383886; (4) . . 7914691943 1279620853 0805687203; (5) . . 5402843601 8957345971 5639810426; (6) . . 7819905213 2701421800 9478672985; (0) . . 0473933649 2890995260 6635071090

223 (0) . . 0448430493 2735426008 9686098654 7085201793 7219730941 7040358744 3946188340 8071748878 9237668161 4349775784 7533632286 9955156950 6726457399 1031390134 5291479820 6278026905 8295964125 5605381165 9192825112 1076233183 8565022421 5246636771 30

227 (1) . . 1806167400 8810572687 2246696035 2422907488 9867841409 6916299559 4713656387 6651982378 8546255506 6079295154 1850220264 317; (0) . . 0440528634 3612334801 7621145374 4493392070 4845814977 9735682819 3832599118 9427312775 3303964757 7092511013 2158590308 370

229 (0) . . 0436681222 7074235807 8602620087 3362445414 8471615720 5240174672 4890829694 3231441048 0349344978 1659388646 2882096069 8689956331 8777292576 4192139737 9912663755 4585152838 4279475982 5327510917 0305676855 8951965065 5021834061 1353711790 39301310

233 (0) . . 0429184549 3562231759 6566523605 1502145922 7467811158 7982832618 0257510729 6137339055 7959914163 0901287553 6480686695 2789799570 8154506437 7682403433 4763948497 8540772532 1888412017 1673819742 4892703862 6609442060 0858369098 7124463519 3133047210 30

239 (1) . . 4644351; (2) . . 2552301; (3) . . 9330543; (4) . . 6569037; (5) . . 9916317; (6) . . 7071129; (7) . . 7489539; (8) . . 2133891; (9) . . 4686192; (10) . . 4016736; (11) . . 0585774; (12) . . 0502092; (13) . . 7573221; (14) . . 5062761; (15) . . 7196652; (16) . . 1882845; (17) . . 5899581; (18) . . 6485355; (19) . . 6987447; (20) . . 4560669; (21) . . 9623430; (22) . . 6820083; (23) . . 8702928; (24) . . 4602510; (25) . . 1087866; (26) . . 8075313; (27) . . 2635983; (28) . . 2259414; (29) . . 9079497; (30) . . 7782426; (31) . . 2384937; (32) . . 3472803; (33) . . 1548117; (0) . . 0418410

241 (1)..5809128630 7053941908 7136929460; (2)..1327800829 8755186721 9917012448; (3)..8589211618 2572614107 8838174273; (4)..0248962655 6016597510 3734439834; (5)..3485477178 4232365145 2282157676; (6)..8796680497 9253112033 1950207468; (7)..3153526970 9543568464 7302904564; (0)..0414937759 3360995850 6224066390

243 (1)..6748971193 4156378600 8230452; (3)..8683127572 0164609053 4979423; (3)..4403292181 0699588477 3662551; (4)..6213991769 5473251028 8065843; (5)..3909465020 5761316872 4279835; (0)..0411522633 7448559670 7818930

251 (1)..4223107569 7211155378 4860557768 9243027888 4462151394; (2)..8764940239 0438247011 9521912350 5976095617 5298804780; (3)..2908366533 8645418326 6932270916 3346613545 8167330677; (4)..2828685258 9641434262 9482071713 1474103585 6573705179; (0)..0398406374 5019920318 7250996015 9362549800 7968127490

257 (0)..0389105058 3657587548 6381322957 1984435797 6653696498 0544747081 7120622568 0933852140 0778210116 7315175097 2762645914 3968871595 3307392996 1089494163 4241245136 1867704280 1556420233 4630350194 5525291828 7937743190 6614785992 2178988326 8482490272 3735408560 3112840466 926070

263 (0)..0380228136 8821292775 6653992395 4372623574 1444866920 1520912547 5285171102 6615969581 7490494296 5779467680 6083650190 1140684410 6463878326 9961977186 3117870722 4334600760 4562737642 5855513307 9847908745 2471482889 7338403041 8250950570 3422053231 9391634980 9885931558 9353612167 30

269 (0)..0371747211 8959107806 6914498141 2639405204 4609665427 5092936802 9739776951 6728624535 3159851301 1152416356 8773234200 7434944237 9182156133 8289962825 2788104089 2193308550 1858736059 4795539033 4572490706 3197026022 3048327137 5464684014 8698884758 3643122676 5799256505 5762081784 38661710

271 (1)..22140; (2)..32841; (3)..97047; (4)..82287; (5)..93726;
(6)..62361; (7)..74169; (8)..45018; (9)..70110; (10)..20664;
(11)..23985; (12)..43911; (13)..63468; (14)..80811; (15)..84870;
(16)..09225; (17)..55350; (18)..32103; (19)..92619; (20)..55719;
(21)..34317; (22)..05904; (23)..35424; (24)..12546; (25)..75276;
(26)..51660; (27)..09963; (28)..59778; (29)..58671; (30)..52029;
(31)..12177; (32)..73062; (33)..38376; (34)..30258; (35)..81549;
(36)..89298; (37)..35793; (38)..14760; (39)..88560; (40)..31365;
(41)..88191; (42)..29151; (43)..74907; (44)..49446; (45)..96678;
(46)..80073; (47)..80442; (48)..82656; (49)..95940; (50)..75645;
(51)..53874; (52)..23447; (53)..39483; (0)..03690

277 (1)..8880866425 9927797833 9350180505 4151624548 7364620938
6281588447 653429602 (2)..0469314079 4223826714 8014440433
2129963898 9169675090 2527075812 274368231 (3)..7545126353
7906137184 1155234657 0397111913 3574007220 2166064981
949458483 (0)..0361010830 3249097472 9241877256 3176895306
8592057761 7328519855 595667870

281 (1)..9217081850 5338078291 81494661; (2)..7722419928 8256227758
00711743; (3)..7010676156 5836298932 38434163; (4)..8576512455
5160142348 75444839; (5)..3131672597 8647686832 74021352;
(6)..9110320284 6975088967 97153024; (7)..1957295373 6654804270
46263345; (8)..5693950177 9359430604 98220640; (9)..7473309608
5409252669 03914590; (0)..0355871886 1209964412 81138790

283 (1)..1519434628 9752650176 6784452296 8197879858 6572438162
5441696113 0742049469 9646643109 5406360424 0282685512
3674911660 7773851590 1060070671 3780918727 9 (0)..0353356890
4593639575 9717314487 6325088339 2226148409 8939929328
6219081272 0848056537 1024734982 3321554770 3180212014
1342756183 7455830388 6925795053 0

289 (0)..0346020761 2456747404 8442906574 3944636678 2006920415
2249134948 0968858131 4878892733 5640138408 3044982698
9619377162 6297577854 6712802768 1660899653 9792387543
2525951557 0934256055 3633217993 0795847750 8650519031
1418685121 1072664359 8615916955 0173010380 6228373702
4221453287 1972318339 10

293 (1)..0375426621 1604095563 1399317406 1433447098 9761092150
1706484641 6382252559 7269624573 3788395904 4368600682
5938566552 9010238907 8498293515 3583617747 440273
(0)..0341296928 3276450511 9453924914 6757679180 8873720136
5187713310 5802047781 5699658703 0716723549 4880546075
0853242320 8191126279 8634812286 6894197952 218430

307 (1) . . 4951140065 1465798045 6026058631 9218241042 3452768729
6416938110 7491856677 5244299674 2671009771 9869706840
3908794788 2736156351 7915309446 2540716612 3778501628 664
(0) . . 0325732899 0228013029 3159609120 5211726384 3648208469
0553745928 3387622149 8371335504 8859934853 4201954397
3941368078 1758957654 7231270358 3061889250 8143322475 570

311 (1) . . 2958199356 9131832797 4276527331 1897106109 3247588424
4372990353 6977491961 4147909967 8456591639 8713826366
5594855305 4662379421 2218649517 6848874598 0707395498 39228
(0) . . 0321543408 3601286173 6334405144 6945337620 5787781350
4823151125 4019292604 5016077170 4180064308 6816720257
2347266881 0289389067 5241157556 2700964630 2250803858 52090

313 (0) . . 0319488817 8913738019 1693290734 8242811501 5974440894
5686900958 4664536741 2140575079 8722044728 4345047923
3226837060 7028753993 6102236421 7252396166 1341853035
1437699680 5111821086 2619808306 7092651757 1884984025
5591054313 0990415335 4632587859 4249201277 9552715654
9520766773 1629392971 2460063897 7635782747 6038338658
1469648562 30

317 (1) . . 2397476340 6940063091 4826498422 7129337539 4321766561
5141955835 9621451104 100946372 (2) . . 0220820189 2744479495
2681388012 6182965299 6845425867 5078864353 3123028391
167192429 (3) . . 5678233438 4858044164 0378548895 8990536277
6025236593 0599369085 1735015772 870662460 (0) . . 0315457413
2492113564 6687697160 8832807570 9779179810 7255520504
7318611987 381703470

331 (1) . . 1178247734 1389728096 6767371601 2084592145 0151057401
8126888217 5226586102 7190332326 2839879154 0785498489
4259818731 (2) . . 3595166163 1419939577 0392749244 7129909365
5589123867 0694864048 3383685800 6042296072 5075528700
9063444108 7613293051 (0) . . 0302114803 6253776435 0453172205
4380664652 5679758308 1570996978 8519637462 2356495468
2779466193 3534743202 4169184290

337 (0) . . 0296735905 0445103857 5667655786 3501483679 5252225519
2878338278 9317507418 3976261127 5964391691 3946587537
0919881305 6379821958 4569732937 6854599406 5281899109
7922848664 6884272997 0326409495 5489614243 3234421364
9851632047 4777448071 2166172106 8249258160 2373887240
3560830860 5341246290 8011869436 2017804154 3026706231
4540059347 1810089020 7715133531 157270

343. (0) . . 0291545189 5043731778 4256559766 7638483965 0145772594 7521865889 2128279883 3819241982 5072886297 3760932944 6064139941 6909620991 2536443148 6880466472 3032069970 8454810495 6268221574 3440233236 1516034985 4227405247 8134110787 1720116618 0758017492 7113702623 9067055393 5860058309 0379008746 3556851311 9533527696 7930

347 (1) . . 6023054755 0432276657 0605187319 8847262247 8386167146 9740634005 7636887608 0691642651 2968299711 8155619596 5417867435 1585014409 2219020172 9106628242 0749279538 9048991354 4668587896 253 (0) . . 0288184438 0403458213 2564841498 5590778097 9827089337 1757925072 0461095100 8645533141 2103746397 6945244956 7723342939 4812680115 2737752161 3832853025 9365994236 3112391930 8357348703 170

.349 (1) . . 3037249283 6676217765 0429799426 9340974212 0343839541 5472779369 6275071633 2378223495 7020057306 5902578796 5616045845 272206 (2) . . 8194842406 8767908309 4555873925 5014326647 5644699140 4011461318 0515759312 3209169054 4412607449 8567335243 5530085959 885386 (0) . . 0286532951 2893982808 0229226361 0315186246 4183381088 8252148997 1346704871 0601719197 7077363896 8481375358 1661891117 478510

353 (1) . . 7932011331 4447592067 9886685552 40; (2) . . 2096317280 4532577903 6827195467 42; (3) . . 8696883852 6912181303 1161473087 81; (4) . . 3512747875 3541076487 2521246458 92; (5) . . 8356940509 9150141643 0594900849 85; (6) . . 3994334277 6203966005 6657223796 03; (7) . . 1841359773 3711048158 6402266288 95; (8) . . 1558073654 3909348441 9263456090 65; (9) . . 3626062322 9461756373 9376770538 24; (10) . . 1529745042 4929178470 2549575070 82; (0) . . 0283286118 9801699716 7138810198 30;

359 (1) . . 3286908077 9944289693 5933147632 3119777158 7743732590 5292479108 6350974930 3621169916 4345403899 7214484679 6657381615 5988857938 7186629526 4623955431 7548746518 1058485821 7270194986 072423398 (0) . . 0278551532 0334261838 4401114206 1281337047 3537604456 8245125348 1894150417 8272980501 3927576601 6713091922 0055710306 4066852367 6880222841 2256267409 4707520891 3649025069 6378830083 565459610

361 (0)..0277008310 2493074792 2437673130 1939058171 7451523545
7063711911 3573407202 2160664819 9445983379 5013850415
5124653739 6121883656 5096952908 5872576177 2853185595
5678670360 1108033240 9972299168 9750692520 7756232686
9806094182 8254847645 4293628808 8642659279 7783933518
0055401662 0498614958 4487534626 0387811634 3490304709
1412742382 2714681440 4432132963 9889196675 90

367 (0)..0272479564 0326975476 8392370572 2070844686 6485013623
9782016348 7738419618 5286103542 2343324250 6811989100
8174386920 9809264305 1771117166 2125340599 4550408719
3460490463 2152588555 8583106267 0299727520 4359673024
5231607629 4277929155 3133514986 3760217983 6512261580
3814713896 4577656675 7493188010 8991825613 0790190735
6948228882 8337874659 4005449591 2806539509 5367847411
4441416893 732970

373 (1)..1983914209 1152815013 4048257372 6541554959 7855227882
0375335120 6434316353 8873994638 0697050938 3378016085
7908847184 9865951742 6273458445 0402144772 1179624664
8793565683 6461126005 3619302949 061662 (0)..0268096514
7453083109 9195710455 7640750670 2412868632 7077747989
2761394101 8766756032 1715817694 3699731903 4852546916
8900804289 5442359249 3297587131 3672922252 0107238605
8981233243 9678284182 305630

379 (0)..0263852242 7440633245 3825857519 7889182058 0474934036
9393139841 6886543535 6200527704 4854881266 4907651715
0395778364 1160949868 0738786279 6833773087 0712401055
4089709762 5329815303 4300791556 7282321899 7361477572
5593667546 1741424802 1108179419 5250659630 6068601583
1134564643 7794722955 1451187335 0923482849 6042216358
8390501319 2612137203 1662269129 2875989445 9102902374
6701846965 6992084432 71767810

383 (0)..0261096605 7441253263 7075718015 6657963446 4751958224
5430809399 4778067885 1174934725 8485639686 6840731070
4960835509 1383812010 4438642297 6501305483 0287206266
3185378590 0783289817 2323759791 1227154046 9973890339
4255874673 6292428198 4334203655 3524804177 5456919060
0522193211 4882506527 4151436031 3315926892 9503916449
0861618798 9556135770 2349869451 6971279373 3681462140
9921671018 2767624020 8877284595 30

389 (0)..0257069408 7403598971 7223650385 6041131105 3984575835
4755784061 6966580976 8637532133 6760925449 8714652956
2982005141 3881748071 9794344473 0077120822 6221079691
5167095115 6812339331 6195372750 6426735218 5089974293
0591259640 1028277634 9614395886 8894601542 4164524421
5938303341 9023136246 7866323907 4550128534 7043701799
4858611825 1928020565 5526992287 9177377892 0308483290
4884318766 0668380462 7249357326 47814910

397 (1)..3501259445 8438287153 6523929471 0327455919 3954659949
6221662468 5138539042 8211586901 7632241813 602015113;
(2)..5667506297 2292191435 7682619647 3551637279 5969773299
7481108312 3425692695 2141057934 5088161209 068010075;
(3)..3778337531 4861460957 1788413098 2367758186 3979848866
4987405541 5617128463 4760705289 6725440806 045340050;
(0)..0251889168 7657430730 4785894206 5491183879 0931989924
4332493702 7707808564 2317380352 6448362720 303022670

401 (1)..7381546134 6633416458 8528678304 2394014962 5935162094
7630922693 2668329177 0573566084 7880299251 8703241895
2618453865 3366583541 1471321695 7605985037 4064837905
2369077306 7331670822 9426433915 2119700748 1296758104
(0)..0249376558 6034912718 2044887780 5486284289 2768079800
4987531172 0698254364 0897755610 9725685785 5361596009
9750623441 3965087281 7955112219 4513715710 7231920199
5012468827 9301745635 9102244389 0274314214 4638403990

409 (1)..2542787286 0635696821 5158924205 3789731051 3447432762
8361858190 7090464547 6772616136 9193154034 2298288508
5574572127 1393643031 7848410757 9462102689 4865525672
3716381418 0929095354 5232273838 6308068459 6577017114 9144
(0)..0244498777 5061124694 3765281173 5941320293 3985330073
3496332518 3374083129 5843520782 3960880195 5990220048
8997555012 2249388753 0562347188 2640586797 0660146699
2665036674 8166259168 7041564792 1760391198 0440097799 5110

419 (0)..0238663484 4868735083 5322195704 0572792362 7684964200
4773269689 7374701670 6443914081 1455847255 3699284009
5465393794 7494033412 8878281622 9116945107 3985680190
9307875894 9880668257 7565632458 2338902147 9713603818
6157517899 7613365155 1312649164 6778042959 4272076372
3150357995 2267303102 6252983293 5560859188 5441527446
3007159904 5346062052 5059665871 1217183770 8830548926
0143198090 6921241050 1183317422 4343675417 6610978520
2863961813 84248210

421 (1)..2826603325 4156769596 1995249406 1757719714 .9643705463 1828978622 3277909738 7173396674 5843230403 8004750593 8242280285 0356294536 8171021377 6722090261 (2)..2636579572 4465558194 7743467933 4916864608 0760095011 8764845605 7007125890 7363420427 5534441805 2256532066 5083135391 9239904988 1235154394 2992874109 (0)..0237529691 2114014251 7814726840 8551068883 6104513064 1330166270 7838479809 9762470308 7885985748 2185273159 1448931116 3895486935 8669833729 2161520190

431 (1)..4872389791 1832946635 7308584686 7749419953 5962877030 1624129930 3944315545 2436194895 5916473317 8654292343 3874709976 7981438515 0812064965 1972197772 6218097447 7958236658 9327146171 6937354988 3990719257 5406032482 5986078886 31090 (0)..0232018561 4849187935 0348027842 2273781902 5522041763 3410672853 8283062645 0116009280 7424593967 5174013921 1136890951 2761020881 6705336426 9141531322 5058004640 3712296983 7587006960 5568445475 6380510440 8352668213 4570765661 25290

433 (0)..0230946882 2170900692 8406466512 7020785219 3995381062 3556581986 1431870669 7459584295 6120092378 7528868360 2771362586 6050808314 0877598152 4249422632 7944572748 2678983833 7182448036 9515011547 3441108545 0346420323 3256351039 2609699769 0531177829 0993071593 5334872979 2147806004 6189376443 4180138568 1293302540 4157043879 9076212471 1316397228 6374133949 1916859122 4018475750 5773672055 4272517321 0161662817 5519630484 9884526558 8914549653 5796766743 6489607390 30

439 (1)..4920273348 5193621867 8815489749 4305239179 9544419134 3963553530 7517084282 4601366742 5968109339 4077448747 1526195899 7722095671 9817767653 7585421412 3006833712 9840546697 0387243735 7630979498 8610478359 9088838268 7927107061 503416856 (0)..0227790432 8018223234 6241457858 7699316628 7015945330 2961275626 4236902050 1138952164 0091116173 1207289293 8496583143 5079726651 4806378132 1184510250 5694760820 0455580865 6036446469 2482915717 5398633257 4031890660 5922551252 847380410

443 (1) . . 4176072234 7629796839 7291196388 2618510158 0135440180 5869574492 0993227990 9706546275 3950338600 4514672686 2302483069 9774266365 6884875846 5011286681 7155756207 6749435665 9142212189 6162528216 7042889390 5191873589 1647855530 4740406320 5 (0) . . 0225733634 3115124153 4988713318 2844243792 3250564334 0857787810 3837471783 2957110609 4808126410 8352144469 5259593679 4582392776 5237020316 0270880361 1738148984 1986455981 9413092550 7900677200 9029345372 4604966139 9548532731 3769751693 0

449 (1) . . 7572383073 4966592427 6169265033 40; (2) . . 7461024498 8864142538 9755011135 85; (3) . . 3674832962 1380846325 1670378619 15; (4) . . 4944320712 6948775055 6792873051 22; (5) . . 8106904231 6258351893 0957683741 64; (6) . . 5634743875 2783964365 2561247216 03; (7) . . 1581291759 4654788418 7082405345 21; (8) . . 3763919821 8262806236 0801781737 19; (9) . . 7973273942 0935412026 7260579064 58; (10) . . 1091314031 1804008908 6859688195 99; (11) . . 7104677060 1336302895 3229398663 69; (12) . . 1559020044 5434298440 9799554565 70; (13) . . 3006681514 4766146993 3184855233 85; (0) . . 0222717149 2204899777 2828507795 10

457 (1) . . 7768052516 4113785557 9868708971 5536105032 8227571115 9737417943 1072210065 6455142231 9474835886 2144420131 2910284463 8949671772 4288840262 5820568927 7899343544 85 (2) . . 0765864332 6039387308 5339168490 1531728665 2078774617 0678336980 3063457330 4157549234 1356673960 6126914660 8315098468 2713347921 2253829321 6630196936 5426695842 45 (0) . . 0218818380 7439824945 2954048140 0437636761 4879649890 5908096280 0875273522 9759299781 1816192560 1750547045 9518599562 3632385120 3501094091 9037199124 7264770240 70

461 (0) . . 0216919739 6963123644 2516268980 .4772234273 3188720173 5357917570 4989154013 0151843817 7874186550 9761388286 3340563991 3232104121 4750542299 3492407809 1106290672 4511930585 6832971800 4338394793 9262472885 0325379609 5444685466 3774403470 7158351409 9783080260 3036876355 7483731019 5227765726 6811279826 4642082429 5010845986 9848156182 2125813449 0238611713 6659436008 6767895878 5249457700 6507592190 8893709327 5488069414 3167028199 5661605206 0737527114 9674620390 4555314533 6225596529 2841648590

463 (1) . . 7580993520 5183585313 1749460043 1965442764 5788336933
0453563714 9028077753 7796976241 9006479481 6414686825
0539956803 4557235421 1663066954 6436285097 1922246220 3023
(2) . . 9092872570 1943844492 4406047516 1987041036 7170626349
8920086393 0885529157 6673866090 7127429805 6155507559
3952483801 2958963282 9373650107 9913606911 4470842332 6133
(0) . . 0215982721 3822894168 4665226781 8574514038 8768898488
1209503239 7408207343 4125269978 4017278617 7105831533
4773218142 5485961123 1101511879 0496760259 1792656587 4730

| | | | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Divisor . . . . . | 3 | 9 | 11 | 13 | 27 | 31 | 37 | 41 | 43 | 53 | 67 | 71 | 73 | 79 | 81 | 83 | 89 | 101 |
| Primitive root . . | 2 | 2 | 2 | 6 | 2 | 17 | 5 | 6 | 28 | 26 | 12 | 62 | 5 | 29 | 11 | 50 | 30 | 2 |

| | | | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Divisor . . . . . | 103 | 107 | 121 | 127 | 137 | 139 | 151 | 157 | 163 | 169 | 173 | 191 | 197 | 199 | 211 | | | |
| Primitive root . . | 6 | 63 | 35 | 106 | 12 | 92 | 114 | 18 | 70 | 137 | 82 | 157 | 73 | 127 | 7 | | | |

| | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Divisor . . . . . | 227 | 239 | 241 | 243 | 251 | 271 | 277 | 281 | 283 | 293 | 307 | 311 | 317 | 331 | 347 |
| Primitive root . . | 163 | 35 | 14 | 65 | 111 | 6 | 80 | 54 | 259 | 89 | 138 | 258 | 71 | 37 | 125 |

| | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Divisor . . . . . | 349 | 353 | 359 | 373 | 397 | 401 | 409 | 421 | 431 | 439 | 443 | 449 | 457 | 463 | |
| Primitive root . . | 220 | 28 | 299 | 82 | 133 | 190 | 174 | 54 | 21 | 285 | 240 | 34 | 264 | 174 | |

467 2141327623 1263383297 6445396145 6102783725 9100642398
2869379014 9892933618 8436830835 1177730192 7194860813
7044967880 0856531049 2505353319 0578158458 2441113490
3640256959 3147751605 9957173447 5374732334 0471092077
0877944325 4817987152 0342612419 700<u>2141327</u> . . . . .

479 2087682672 2338204592 9018789144 0501043841 3361169102
2964509394 5720250521 9206680584 5511482254 6972860125
2609603340 2922755741 1273486430 0626304801 6701461377
8705636743 2150313152 4008350730 6889352818 3716075156
5762004175 3653444676 4091858037 578288100<u>2</u> . . . . .

487 2053388090 3490759753 5934291581 1088295687 8850102669
4045174537 9876796714 5790554414 7843942505 1334702258
7268993839 8357289527 7207392197 1252566735 1129363449
6919917864 4763860369 6098562628 3367556468 1724845995
8932238193 0184804928 1314163377 8234086242 2997946611
9096509240 2464065708 4188911704 3121149897 3305954825
4620123203 2854209445 5852156057 4948665297 7412731006
1601642710 4722792607 8028747433 2648870636 5503080082
1355236139 6303901437 3716632443 5318275154 0041067761
8069815195 0718685831 6221765913 757700<u>2053</u> . . .

491 2036659877 8004073319 7556008146 6395112016 2932790224
0325865580 4480651731 1608961303 4623217922 6069246435
8452138492 8716904276 9857433808 5539714867 6171079429
7352342158 8594704684 3177189409 3686354378 8187372708
7576374745 4175152749 4908350305 4989816700 6109979633
4012219959 2668024439 9185336048 8798370672 0977596741
3441955193 4826883910 3869653767 8207739307 5356415478
6150712830 9572301425 6619144602 8513238289 2057026476
5784114052 9531568228 1059063136 4562118126 2729124236
2525458248 4725050916 4969450101 8329938900 . . .

499 2004008016 0320641282 5651302605 2104208416 8336673346
6933867735 4709418837 6753507014 0280561122 2444889779
5591182364 7294589178 3567134268 5370741482 9659318637
2745490981 9639278557 1142284569 1382765531 0621242484
9699398797 5951903807 6152304609 2184368737 4749498997
9959919839 6793587174 3486973947 8957915831 6633266533
0661322645 2905811623 2464929859 7194388777 5551102204
4088176352 7054108216 4328657314 6292585170 3406813627
2545090180 3607214428 8577154308 6172344689 3787575150
3006012024 0480961923 8476953907 8156312625 2505010020 . . . .

503 1988071570 5765407554 6719681908 5487077534 7912524850
8946322067 5944333996 0238568588 4691848906 5606361829
0258449304 1749502982 1073558648 1113320079 5228628230
6163021868 7872763419 4831013916 5009940357 8528827037
7733598409 5427435387 6739562624 2544731610 3379721669
9801192842 9423459244 5328031809 1451292246 5208747514
9105367793 2405566600 3976143141 1530815109 3439363817
0974155069 5825049701 7892644135 1888667992 0477137176
9383697813 1212723658 0516898608 3499005964 2147117296
2226640159 0457256461 2326043737 5745526838 9662027833
00<u>19880715</u> . . . . .

509 1964636542 2396856581 5324165029 4695481335 9528487229
8624754420 4322200392 9273084479 3713163064 8330058939
0962671905 6974459724 9508840864 4400785854 6168958742
6326129666 0117878192 5343811394 8919449901 7681728880
1571709233 7917485265 2259332023 5756385068 7622789783
8899803536 3457760314 3418467583 4970530451 8664047151
2770137524 5579567779 9607072691 5520628683 6935166994
1060903732 8094302554 0275049115 9135559921 4145383104
1257367387 0333988212 1807465618 8605108055 0098231827
1119842829 0766208251 4734774066 7976424361 4931237721
021611001<u>9</u> . . . . .

521 1919385796 5451055662 1880998080 6142034548 9443378119
00191938 57 ....

523 1912045889 1013384321 2237093690 2485659655 8317399617
5908221797 3231357552 5812619502 8680688336 5200764818
3556405353 7284894837 4760994263 8623326959 8470363288
7189292543 0210325047 8011472275 3346080305 9273422562
1414913957 9349904397 7055449330 7839388145 3154875717
0172084130 01912045 88 .....

529 1890359168 2419659735 3497164461 2476370510 3969754253
3081285444 2344045368 6200378071 8336483931 9470699432
8922495274 1020793950 8506616257 0888468809 0737240075
6143667296 7863894139 8865784499 0548204158 7901701323
2514177693 7618147448 0151228733 4593572778 8279773156
8998109640 8317580340 2646502835 5387523629 4896030245
7466918714 5557655954 6313799621 9281663516 0680529300
5671077504 7258979206 0491493383 7429111531 1909262759
9243856332 7032136105 8601134215 5009451795 8412098298
6767485822 3062381852 5519848771 2665406427 2211720226
8431001890 ....

541 1848428835 4898336414 0480591497 2273567467 6524953789
2791127541 5896487985 2125693160 8133086876 1552680221
8114602587 8003696857 6709796672 8280961182 9944547134
9353049907 5785582255 0831792975 9704251386 3216266173
7523105360 4436229205 1756007393 7153419593 3456561922
3659889094 2698706099 8151571164 5101663385 9519408502
7726432532 3475046210 7208872458 4103512014 7874306839
1866913123 8447319778 1885397412 1996303142 3290203327
1719038817 0055452865 0646950092 4214417744 9168207024
0295748613 6783733826 2476894639 5563770794 8243992606
2846580406 6543438077 6340110905 7301293900 1848428835 ....

547 1828153564 8994515539 3053016453 3820840950 6398537477
1480804387 5685557586 8372943327 2394881170 01828153 56 ....

557 1795332136 4452423698 3842010771 9928186714 5421903052
0646319569 1202872531 4183123877 9174147217 2351885098
7432675044 8833034111 3105924596 0502692998 2046678635
5475763016 1579892280 0718132854 5780969479 3536804308
7971274685 8168761220 8258527827 6481149012 5673249551
1669658886 8940754039 4973070017 ....

563 1776198934 2806394316 1634103019 5381882770 8703374777
9751332149 2007104795 7371225577 2646536412 0781527531
0834813499 1119005328 5968028419 1829484902 3090586145
6483126110 1243339253 9964476021 3143872113 6767317939
6092362344 5825932504 4404973357 0159857904 0852575488
4547069271 7584369449 3783303730 01̲77619893̲ . . . . .

569 1757469244 2882249560 6326889279 4376098418 2776801405
9753954305 7996485061 5114235500 8787346221 4411247803
1634446397 1880492091 3884007029 8769771528 9982425307
5571177504 3936731107 2056239015 8172231985 9402460456
9420035149 3848857644 9912126537 7855887521 9683655536
0281195079 0861159929 7012302284 7100175746̲ . . . .

571 1751313485 1138353765 3239929947 4605954465 8493870402
8021015761 8213660245 1838879159 3695271453 5901926444
8336252189 1418563922 9422066549 9124343257 4430823117
3380035026 2697022767 0753064798 5989492119 0893169877
4080560420 3152364273 2049036777 5831873905 4290718038
5288966725 0437828371 2784588441 3309982486 8651488616
4623467600 7005253940 4553415061 2959719789 8423817863
3975481611 2084063047 2854640980 7355516637 4781085814
3607705779 3345008756 5674255691 7688266199 6497373029
7723292469 3520140105 0788091068 3012259194 3957968476
3572679509 6322241681 2609457092 8196147110 3327495621
7162872154 1155866900 1̲751313485 . . . .

577 1733102253 0329289428 0762564991 3344887348 3535528596
1871750433 2755632582 3223570190 6412478336 2218370883
8821490467 9376083188 9081455805 8925476603 1195840554
5927209705 3726169844 0207972270 3639514731 3691507798
9601386481 8024263431 5424610051 9930675909 8786828422
8769497400 3466204506 0658578856 1525129982 6689774696
7071057192 3743500866 5511265164 6447140381 2824956672
4436741767 7642980935 8752166377 8162911611 7850953206
2391681109 1854419410 7452339688 0415944540 7279029462
7383015597 9202772963 6048526863 0849220103 9861351819
7573656845 7538994800 6932409012 1317157712 3050259965
3379549393 4142114384 7487001̲733 . . . .

587 1703577512 7768313458 2623509369 1763202725 7240204429
3015332197 6149914821 1243611584 3270868824 5315161839
8637137989 7785349233 3901192504 2589437819 4207836456
5587734241 9080068143 1005110732 5383304940 3747870528
1090289608 1771720613 2879045996 5928449744 4633730834
7529812606 4735945485 5195911413 9693356047 7001703577 ....

593 1686340640 8094435075 8853288364 2495784148 3979763912
3102866779 0893760539 6290050590 2192242833 0522765598
6509274873 5244519392 9173693086 0033726812 8161888701
5177065767 2849915682 9679595278 2462057335 5817875210
7925801011 8043844856 6610455311 9730185497 4704890387
8583473861 7200674536 2563237774 0303541315 3456998313
6593591905 5649241146 7116357504 2158516020 2360876897
1332209106 2394603709 9494097807 7571669477 2344013490
7251264755 4806070826 3069139966 2731871838 1112984822
9342327150 0843170320 4047217537 9426644182 1247892074
1989881956 1551433389 5446880269 8145025295 1096121416
5261382799 3254637436 7622259696 4586846543 0016863406 ....

599 1669449081 8030050083 4724540901 5025041736 2270450751
2520868113 5225375626 0434056761 2687813021 7028380634
3906510851 4190317195 3255425709 5158597662 7712854757
9298831385 6427378964 9415692821 3689482470 7846410684
4741235392 3205342237 0617696160 2671118530 8848080133
5559265442 4040066777 9632721202 0033388981 6360601001 ....

601 1663893510 8153078202 9950083194 6755407653 9101497504
1597337770 3826955074 8752079866 8885191347 7537437603
9933444259 5673876871 8801996672 2129783693 8435940099
8336106489 1846921797 0049916805 3244592346 0898502495
8402662229 6173044925 1247920133 1114808652 2462562396
0066555740 4326123128 1198003327 7870216306 1564059900
1663893510 ...

607 1647446457 9901153212 5205930807 2487644151 5650741350
9060955518 9456342668 8632619439 8682042833 6079077429
9835255354 2009884678 7479406919 2751235584 8434925864
9093904448 1054365733 1136738056 0131795716 6392092257
0016474464 ....

613 1631321370 3099510603 5889070146 8189233278 9559543230
0163132137 ....

617 1620745542 9497568881 6855753646 6774716369 5299837925
4457050243 1118314424 6353322528 3630470016 ....

619 1615508885 2988691437 8029079159 9353796445 8804523424
8788368336 0258481421 6478190630 0484652665 5896607431
3408723747 9806138933 7641357027 4636510500 8077544426
4943457189 0145395799 6768982229 4022617124 3941841680
1292407108 2390953150 2423263327 9483037156 7043618739
9030694668 8206785137 3182552504 0387722132 4717235945
0726978998 3844911147 0113085621 9709208400 6462035541
1954765751 2116316639 7415185783 5218093699 5153473344
1033925686 5912762520 1938610662 3586429725 3634894991
9224555735 0565428109 8546042003 2310177705 9773828756
0581583198 7075928917 6090468497 5767366720 5169628432
9563812600 9693053311 7932148626 8174474959 6122778675
2827140549 2730210016 . . . .

631 1584786053 8827258320 1267828843 1061806656 1014263074
4849445324 8811410459 5879556259 9049128367 6703645007
9239302694 1362916006 3391442155 3090332805 0713153724
2472266244 0570522979 3977812995 2456418383 5182250396
1965134706 8145800316 9572107765 4516640253 5657686212
3613312202 8526148969 8890649762 2820919175 9112519809
8256735340 7290015847 . . . .

641 1560062402 4960998439 9375975039 0015600624 . .

643 1555209953 3437013996 8895800933 1259720062 2083981337
4805598755 8320373250 3888024883 3592534992 2239502332
8149300155 . . . .

647 1545595054 0958268933 5394126738 7944358578 0525502318
3925811437 4034003091 1901081916 5378670788 2534775888
7171561051 0046367851 6228748068 0061823802 1638330757
3415765069 5517774343 1221020092 7357032457 4961360123
6476043276 6615146831 5301391035 5486862442 0401854714
0649149922 7202472952 0865533230 2936630602 7820710973
7248840803 7094281298 2998454404 9459041731 0664605873
2612055641 4219474497 6816074188 5625965996 9088098918
0834621329 2117465224 1112828438 9489953632 1483771251
9319938176 1978361669 2426584234 9304482225 6568778979
9072642967 5425038639 8763523956 7233384853 1684698608
9644513137 5579598145 2859350850 0772797527 0479134466
7697063369 3972179289 0262751159 1962905718 7017001545 . . . .

653 1531393568 1470137825 4211322312 4042879019 9081163859
1117917304 7473200612 5574272588 0551301684 5329249617
1516079632 4655436447 1669218989 2802450229 7090352220
5206738131 6998468606 4318529862 1745788667 6875957120
9800918836 1408882082 6952526799 3874425727 4119448698
3154670750 3828483920 3675344563 5528330781 0107197549
7702909647 7794793261 868300153̲1̲ ....

659 1517450682 · 8528072837 6327769347 4962063732 9286798179
0591805766 3125948406 6767830045 5235204855 8421851289
8330804248 8619119878 6039453717 7541729893 7784522003
0349013657 0561456752 6555386949 9241274658 5735963581
1836115326 2518968133 5356600910 4704097116 8437025796
6616084977 2382397572 0789074355 0834597875 5690440060
6980273141 1229135053 1107738998 4825493171 4719271623
6722306525 0379362670 7132018209 4081942336 8740515933
2321699544 7647951441 5781487101 6691957511 3808801213
9605462822 4582701062 2154779969 6509863429 4385432473
4446130500 7587253414 2640364188 1638846737 4810318664
6433990895 2959028831 5629742033 3839150227 6176024279
2109256449 1654021244 3095599393 0197268588 7708649468
892261001̲5̲ ....

661 1512859304 0847201210 2874432677 7609682299 5461422087
7458396369 1376701966 7170953101 3615733736 7624810892
5869894099 8487140695 9152798789 7125567322 2390317700
4538577912 2541603630 8623298033 2829046898 6384266263
2375189107 4130105900 1̲5̲12859304 ....

673 1485884101 0401188707 2808320950 9658246656 7607726597
3254086181 2778603268 9450222882 6151560178 3060921248
1426448736 9985141158 9895988112 9271916790 4903417533
4323922734 0267459138 1872213967 3105497771 1738484398
2169390787 5185735512 6300148588 ....

677 1477104874 4460856720 8271787296 8980797636 6322008862
6292466765 1403249630 7237813884 7858197932 0531757754
8005908419 4977843426 8833087149 1875923190 5465288035
4505169867 0605612998 5228951255 5391432791 7282127031
0192023633 6779911373 7075332348 5967503692 7621861152
1418020679 4682422451 9940915805 0221565731 1669128508
1240768094 5347119645 4948301329 394387001̲4̲ ....

683 1464128843 3382137628 1112737920 9370424597 3645680819
9121522693 9970717423 1332357247 4377745241 5812591508
0527086383 6017569546 1200585651 5373352855 0512445095
1683748169 8389458272 3279648609 0775988286 9692532942
8989751098 0966325036 6032210834 5534407027 8184480234
2606149341 1420204978 0380673499 2679355783 3089311859
4436310395 3147877013 1771595900 4392386530 0146412884 ....

691 1447178002 8943560057 8871201157 7424023154 8480463096
9609261939 2185238784 3704775687 4095513748 1910274963
8205499276 4109985528 2199710564 3994211287 9884225759
7684515195 3690303907 3806078147 6121562952 2431259044
8625180897 2503617945 0072358900 1447178022 ....

701 1426533523 5378031383 7375178316 6904422253 9229671897
2895863052 7817403708 9871611982 8815977175 4636233951
4978601997 1469329529 2439372325 2496433666 1911554921
5406562054 2082738944 3651925820 2567760342 3680456490
7275320970 0427960057 0613409415 1212553495 0071326676
1768901569 1868758915 8345221112 6961483594 8644793152
6390870185 4493580599 1440798858 7731811697 5748930099
8573466476 4621968616 2624821683 3095577746 0770328102
7104136947 2182596291 0128388017 1184022824 5363766048
5021398002 8530670470 7560627674 7503566333 8088445078
4593437945 7917261055 6348074179 7432239657 6319543509
2724679029 9572039942 9386590584 8787446504 9928673323
8231098430 8131241084 1654778887 3038516405 1355206847
3609129814 5506419400 8559201141 2268188302 4251069900 ....

709 1410437235 5430183356 8406205923 8363892806 7700987306
0648801128 3497884344 1466854724 9647390691 1142454160
7898448519 0409026798 3074753173 4837799717 9125528913
9633286318 7588152327 2214386459 8025387870 2397743300
4231311706 6290550070 5218617771 5091678420 3102961918
1946403385 0493653032 4400564174 8942172073 3427362482
3695345557 1227080394 9224259520 4513399153 7376586741
8899858956 2764456981 6643159379 4076163610 7193229901
2693935119 8871650211 5655853314 5275035260 9308885754
5839210155 1480959097 3201692524 6826516220 0282087447
1086036671 3681241184 7672778561 3540197461 2129760225
6699576868 8293370944 9929478138 2228490832 1579689703
8081805359 6614950634 6967559943 5825105782 7926657263
7517630465 4442877291 9605077574 0479548660 0846262341
3258110014 ....

719 1390320584 1446453407 5104311543 8108484005 5632823365
7858136300 4172461752 4339360222 5312934631 4325452016
6898470097 3574408901 2517385257 3018080667 5938803894
2976356050 0695410292 0723226703 7552155771 9054242002
7816411682 8929068150 2086230876 2169680111 2656467315
7162726008 3449235048 6787204450 6258692628 6509040333
7969401947 1488178025 0347705146 0361613351 8776077885
9527121001 ....

727 1375515818 4319119669 8762035763 4112792297 1114167812
9298486932 5997248968 3631361760 6602475928 4731774415
4057771664 3741403026 1348005502 0632737276 4786795048
1430536451 1691884456 6712517193 9477303988 9958734525
4470426409 9037138927 0976616231 0866574965 6121045392
0220082530 9491059147 1801925722 1458046767 5378266850
0687757909 2159559834 9381017881 7056396148 5557083906
4649243466 2998624484 1815680880 3301237964 2365887207
7028885832 1870701513 0674002751 0316368638 2393397524
0715268225 5845942228 3356258596 9738651994 4979367262
7235213204 9518569463 5488308115 5433287482 8060522696
0110041265 4745529573 5900962861 0729023383 7689133425
0343878954 6079779917 4690508940 8528198074 2778541953
2324621733 1499312242 0907840440 1650618982 1182943603
8514442916 0935350756 5337001375 ....

729 1371742112 4828532235 9396433470 5075445816 1865569272
9766803840 8779149519 8902606310 0137174211

733 1364256480 2182810368 3492496589 3587994542 9740791268
7585266030 0136425648 ....

739 1353179972 9364005412 7198917456 0216508795 6698240866
0351826792 9634641407 3071718538 5656292286 8741542625
1691474966 1705006765 8998646820 0270635994 5872801082
5439783491 2043301759 1339648173 2070365358 5926928281
4614343707 7131258457 3748308525 0338294993 2341001353 ....

743 1345895020 1884253028 2637954239 5693135935 3970390309
5558546433 3781965006 7294751009 4212651413 1897711978
4656796769 8519515477 7927321668 9098250336 4737550471
0632570659 4885598923 2839838492 5975773889 6366083445
4912516823 6877523553 1628532974 4279946164 1991924629
8788694481 8304172274 5625841184 3876177658 1426648721
3997308209 9596231493 9434724091 5208613728 1292059219
3808882907 1332436069 9865410497 9811574697 1736204576
0430686406 4602960969 0444145356 6621803499 3270524899
0578734858 6810228802 1534320323 0148048452 2207267833
1090174966 3526244952 8936742934 0511440107 6716016150
7402422611 0363391655 4508748317 6312247644 6837146702
5572005383 5800807537 0121130551 8169582772 5437415881
5612382234 1857335127 8600269179 0040376850 6056527590
8479138627 1870794078 0619111709 2866756393 0013458950 ....

751 1331557922 7696404793 6085219707 0572569906 7909454061
2516644474 0346205059 9201065246 3382157123 8348868175
7656458055 9254327563 2490013315 ....

757 1321003963 0118890356 6710700132 ....

761 1314060446 7805519053 8764783180 0262812089 3561103810
7752956636 0052562417 8712220762 1550591327 2010512483
5742444152 4310118265 4402102496 7148488830 4862023653
0880420499 3429697766 0972404730 6176084099 8685939553
2194480946 1235216819 9737187910 6438896189 2247043363
9947437582 1287779237 8449408672 7989487516 4257555847
5689881734 5597897503 2851511169 5137976346 9119579500
6570302233 9027595269 3823915900 1314060446 ....

769 1300390117 0351105331 5994798439 5318595578 6736020806
2418725617 6853055916 7750325097 5292587776 3328998699
6098829648 8946684005 2015604681 4044213263 9791937581
2743823146 9440832249 6749024707 4122236671 0013003901 ....

773 1293661060 8020698576 9728331177 2315653298 8357050452
7813712807 2445019404 9159120310 4786545924 9676584734
7994825355 7567917205 6921096675 2910737386 8046571798
1888745148 7710219922 3803363518 7580853816 3001293661 ....

787 1270648030 4955527318 9326556543 8373570520 9656925031
7662007623 8881829733 1639135959 3392630241 4231257941
5501905972 0457433290 9783989834 8157560355 7814485387
5476493011 4358322744 5997458703 9390088945 3621346886
9123252858 9580686149 9364675984 7522236340 5336721728
0813214739 5171537484 1168996188 0559085133 4180432020
3303684879 2884371029 2249047013 9771283354 5108005082
5921219822 1092757306 2261753494 2820838627 700$\underline{12}$70648 ....

797 1254705144 2910915934 7553324968 6323713927 2271016311
1668757841 9071518193 2245922208 2810539523 2120451693
8519447929 7365119196 9887076537 0138017565 8720200752
8230865746 5495608531 9949811794 2283563362 6097867001 ....

809 1236093943 1396786155 7478368355 9950556242 2744128553
7700865265 7601977750 3090234857 8491965389 3695920889
9876390605 6860321384 4252163164 4004944375 7725587144
6229913473 4239802224 9690976514 2150803461 0630407911
00$\underline{12360939}$ ....

811 1233045622 6880394574 5992601726 2638717632 5524044389
6424167694 2046855733 6621454993 8347718865 5980271270
0369913686 8064118372 3797780517 8791615289 7657213316
8927250308 2614056720 0986436498 1504315659 6794081381
0110974106 0419235511 7139334155 3637484586 9297163995
0678175092 4784217016 0295930949 4451294697 9038224414
3033292231 8125770653 5141800246 6091245376 0789149198
5203452527 7435265104 8088779284 8335388409 3711467324
2909987669 5437731196 0542540073 9827373612 8236744759
5561035758 3230579531 4426633785 4500616522 8113440197
2872996300 8631319358 8162762022 1948212083 8471023427
8668310727 4969173859 4327990135 6350184956 8434032059
1861898890 2589395807 6448828606 6584463625 1541307028
3600493218 2490752157 8298397040 6905055487 0530209617
7558569667 0776818742 2934648581 9975339087 5462392108
5080147965 4747225647 3489519112 2071516646 1159062885
3267570900 123$\underline{304}5\underline{662}$ ....

821 1218026796 5895249695 4933008526 1875761266 7478684531
0596833130 3288672350 7917174177 8319123020 7064555420
2192448233 8611449451 8879415347 1376370280 1461632155
9074299634 5919610231 4250913520 0974421437 2716199756
3946406820 9500609013 3982947624 8477466504 2630937880
6333739342 2655298416 5651644336 1753958587 0889159561
5103532277 7101096224 1169305724 7259439707 6735688185
1400730816 0779537149 8172959805 1157125456 7600487210
7186358099 8781973203 4104750304 5066991473 8124238733
2521315468 9403166869 6711327649 2082825822 1680876979
2935444579 7807551766 1388550548 1120584652 8623629719
8538367844 0925700365 4080389768 5749086479 9025578562
7283800243 6053593179 0499390986 6017052375 1522533495
7369062119 3666260657 7344701583 4348355663 8246041412
9110840438 4896467722 2898903775 8830694275 2740560292
3264311814 8599269183 9220462850 1827040194 8842874543
2399512789 2813641900 1218026796 ....

823 1215066828 6755771567 4362089914 9453219927 0959902794
6537059538 2746051032 8068043742 4058323207 7764277035
2369380315 9173754556 5006075334 1433778857 8371810449
5747266099 6354799513 9732685297 6913730255 1640340218
7120291616 0388821385 1761846901 5795868772 7825030376
6707168894 2891859052 2478736330 4981773997 5698663426
4884568651 2758201701 0935601458 0801944106 9258809234
5078979343 8639125151 8833535844 4714459295 2612393681
6524908869 9878493317 1324422843 2563791008 5054678007
2904009720 5346294046 1725394896 7193195625 7594167679
2223572296 4763061968 4082624544 3499392466 5856622114
2162818955 0425273390 0364520048 6026731470 2308626974
4835965978 1287970838 3961117861 4823815309 8420413122
7217496962 3329283110 5710814094 7752126366 9501822600
2430133657 3511543134 8724179829 8906439854 1919805589
3074119076 5492102065 6136087484 8116646415 5528554070
4738760631 8347509113 0012150668 ....

827 1209189842 8053204353 0834340991 5356711003 6275695284
1596130592 5030229746 0701330108 8270858524 7883917775
0906892382 1039903264 8125755743 6517533252 7206771463
1197097944 3772672309 5525997581 6203143893 5912938331
3180169286 5779927448 6094316807 7388149939 5405078597
3397823458 2829504232 1644498186 2152357920 1934703748
4885126964 9334945586 4570737605 8041112454 6553808948
0048367593 7122128174 1233373639 6614268440 1451027811
3663845223 700$\underline{1209189}$ ....

829 1206272617 6115802171 2907117008 4439083232 8106151990
3498190591 0735826296 7430639324 4873341375 1507840772
0144752714 1133896260 5548854041 0132689987 9372738238
8419782870 9288299155 6091676718 9384800965 0180940892
6417370325 6936067551 2665862484 9215922798 5524728588
6610373944 5114595898 6731001206 ....

839 1191895113 2300357568 5339690107 2705601907 0321811680
5721096543 5041716328 9630512514 8986889153 7544696066
7461263408 8200238379 0226460071 5137067938 0214541120
3814064362 3361144219 3087008343 2657926102 5029797377
8307508939 2133492252 6817640047 6758045292 0143027413
5876042908 2240762812 8724672228 8438617401 6686531585
2205005959 4755661501 7878426698 4505363528 0095351609
0584028605 4827175208 5816448152 5625744934 4457687723
4803337306 3170441001 ....

841 1189060642 0927467300 8323424494 6492271105 8263971462
5445897740 7847800237 8121284185 4934601664 6848989298
4542211652 7942925089 1795481569 5600475624 2568370986
9203329369 7978596908 4423305588 5850178359 0963139120
0951248513 6741973840 6658739595 7193816884 6611177170
0356718192 6278240190 2497027348 3947681331 7479191438
7633769322 2354340071 3436385255 6480380499 4054696789
5362663495 8382877526 7538644470 8680142687 2770511296
0760998810 9393579072 5326991676 5755053507 7288941736
0285374554 1022592152 1997621878 7158145065 3983353151
0107015457 7883472057 0749108204 5184304399 5243757431
6290130796 6706302021 4030915576 6944114149 8216409036
8608799048 7514863258 0261593341 2604042806 1831153388
8228299643 2818073721 7598097502 9726516052 3186682520
8085612366 2306777645 6599286563 6147443519 6195005945
3032104637 3365041617 1224732461 3555291319 8573127229
4887039239 00$\underline{11890}\underline{606}$ ....

853 1172332942 5556858147 7139507620 1641266119 5779601406
7995310668 2297772567 4091441969 5193434935 5216881594
3728018757 3270808909 7303634232 1219226260 2579132473
6225087924 9706916764 3610785463 0715123094 9589683470
1055099648 300$\underline{1172}$33$\underline{2}$ ....

857 1166861143 5239206534 4224037339 5565927654 6091015169
1948658109 6849474912 4854142357 0595099183 1971995332
5554259043 1738623103 8506417736 2893815635 9393232205
3675612602 1003500583 4305717619 6032672112 0186697782
9638273045 5075845974 3290548424 7374562427 0711785297
5495915985 9976662777 1295215869 3115519253 2088681446
9078179696 6161026837 8063010501 7502917152 8588098016
3360560093 3488914819 1365227537 9229871645 2742123687
2812135355 8926487747 9579929988 3313885647 6079346557
7596266044 3407234539 0898483080 5134189031 5052508751
4585764294 0490081680 2800466744 4574095682 6137689614
9358226371 0618436406 0676779463 2438739789 9649941656
9428238039 6732788798 1330221703 6172695449 2415402567
0945157526 2543757292 8821470245 0408401400 2333722287
0478413068 8448074679 1131855309 2182030338 3897316219
3698949824 9708284714 1190198366 3943990665 1108518086
3477246207 7012835472 5787631271 8786464410 7351225204
200700$\underline{1166}$ ....

859 1164144353 8998835855 6461001$\underline{1164}$ ....

863 1158748551 5643105446 1181923522 5955967555 0405561993
0475086906 1413673232 9084588644 2641946697 5666280417
1494785631 5179606025 4924681344 1483198146 0023174971
0312862108 9223638470 4519119351 1008111239 8609501738
1228273464 6581691772 8852838933 9513325608 3429895712
6303592120 5098493626 8829663962 9200463499 4206257242
1784472769 4090382387 0220162224 7972190034 7624565469
2931633835 4577056778 6790266512 1668597914 2526071842
4101969872 5376593279 2584009269 9884125144 8435689455
3881807647 7404403244 4959443800 6952491309 3858632676
7091541135 5735805330 2433371958 2850521436 8482039397
4507531865 5851680185 3997682502 8968713789 1077636152
9548088064 8899188876 0139049826 1877172653 5341830822
7114716106 6048667439 1657010438 7369640787 9490150637
3117033603 7079953650 0579374275 7821552723 0590961761
2977983777 5202780996 5237543453 0706836616 4542294322
1320973348 7833140208 5747392815 7589803012 7462340672
0741599073 00$\underline{11587485}$ ....

877 1140250855 1881413911 0604332953 2497149372 8620296465
2223489167 6168757126 5678449258 8369441277 0809578107
1835803876 8529076396 8072976054 7320410490 3078677309
0079817559 8631698973 7742303306 7274800456 1003420752
5655644241 7331812998 8597491448 1185860889 3956670467
5028506271 3797035347 7765108323 8312428734 3215507411
6305587229 1904218928 1641961231 4709236031 9270239452
6795895096 9213226909 9201824401 3683010262 2576966932
7251995438 9965792474 3443557582 668187011 ....

881 1135073779 7956867196 3677639046 5380249716 2315550510
7832009080 5902383654 9375709421 1123723041 9977298524
4040862656 0726447219 0692395005 6753688989 7843359818
3881952326 9012485811 5777525539 1600454029 5119182746
8785471055 6186152099 8864926220 2043132803 6322360953
4619750283 7684449489 2167990919 4097616345 0624290578
8876276958 0022701475 5959137343 9273552780 9307604994
3246311010 2156640181 6118047673 0987514188 4222474460
8399545970 4880817253 1214528944 3813847900 1135073779 ....

883 1132502831 2570781426 9535673839 1845979614 9490373725
9343148357 8708946772 3669309173 2729331823 3295583238
9580973952 4348810872 0271800679 5016987542 4688561721
4043035107 5877689694 2242355605 8890147225 3680634201
5855039637 5990939977 3499433748 5843714609 2865232163
0804077010 1925754813 1370328425 8210645526 6138165345
4133635334 0883352208 3805209513 0237825594 5639864099
6602441506 2287655719 1392978482 4462061155 1528878822
1970554926 3873159682 8992072480 1812004530 0113250283 ....

887 1127395715 8962795941 3754227733 9346110484 7801578354
0022547914 3179255918 8275084554 6786922209 6956031567
0800450958 2863585118 3765501691 0935738444 1939120631
3416009019 1657271702 3675310033 8218714768 8838782412
6268320180 3833145434 0473506200 6764374295 3776775648
2525366403 6076662908 6809470124 0135287485 9075535512
9650507328 0721533258 1736189402 4802705749 7181510710
2593010146 5614430665 1634723788 0496054114 9943630214
2051860202 9312288613 3032694475 7609921082 2998872604
2841037204 0586245772 2660653889 5152198421 6459977452
0856820744 0811724915 4453213077 7903043968 4329199549
0417136414 8816234498 3089064261 5558060879 3686583990
9808342728 2976324689 9661781285 2311161217 5873731679
8196166854 5659526493 7993235625 7046223224 3517474633
5963923337 0913190529 8759864712 5140924464 4870349492
6719278466 7418263810 5975197294 2502818489 2897406989
8534385569 3348365276 2119503945 8850056369 7857948139
7970687711 3866967305 5242390078 9177001127 ....

907 1102535832 4145534729 8787210584 3439911797 1334068357
2216097023 1532524807 0562293274 5314222712 2381477398
0154355016 5380374862 1830209481 8081587651 5986769570
0110253583 ....

911 1097694840 8342480790 3402854006 5861690450 0548847420
4171240395 1701427003 2930845225 0274423710 2085620197
5850713501 6465422612 5137211855 1042810098 7925356750
8232711306 2568605927 5521405049 3962678375 4116355653
1284302963 7760702524 6981339187 7058177826 5642151481
8880351262 3490669593 8529088913 2821075740 9440175631
1745334796 9264544456 6410537870 4720087815 5872667398
4632272228 3205268935 2360043907 7936333699 2316136114
1602634467 6180021953 8968166849 6158068057 0801317233
8090010976 ....

919 1088139281 8280739934 7116430903 1556039173 0141458106
6376496191 5125136017 4102285092 4918389553 8628944504
8966267682 2633297062 0239390642 0021762785 6365614798
6942328618 0631120783 4602829162 1327529923 8302502720
3482045701 8498367791 0772578890 0979325353 6452665941
2404787812 8400435255 7127312295 9738846572 3612622415
6692056583 2426550598 4766050054 4069640914 0369967355
8215451577 8019586507 0729053318 8248095756 2568008705
1142546245 9194776931 4472252448 3133841131 6648531011
9695321001 ....

929 1076426264 8008611410 1184068891 2809472551 1302475780
4090419806 2432723358 4499461786 8675995694 2949407965
5543595263 7244348762 1097954790 0968783638 3207750269
1065662002 1528525296 0172228202 3681377825 6189451022
6049515608 1808396124 8654467168 9989235737 3519913885
8988159311 0871905274 4886975242 1959095801 9375672766
4155005382 1313240043 0570505920 3444564047 3627556512
3789020452 0990312163 6167922497 3089343379 9784714747
0398277717 9763186221 7438105489 7739504843 9181916038
7513455328 3100107642 ....

937 1067235859 1248665955 1760939167 5560298826 0405549626
4674493062 9669156883 6712913553 8954108858 0576307363
9274279615 7950907150 4802561366 0618996798 2924226254
0021344717 1824973319 1035218783 3511205976 5208110992
5293489861 2593383137 6734258271 0779082177 1611526147
2785485592 3159018143 0096051227 3212379935 9658484525
0800426894 3436499466 3820704375 6670224119 5304162219
8505869797 2251867662 7534685165 4215581643 5432230522
9455709711 8463180362 8601921024 5464247598 7193169690
5016008537 8868729989 3276414087 5133404482 3906083244
3970117395 9445037353 2550693703 3084311632 8708644610
4589114194 2369263607 2572038420 4909284951 9743863393
8100320170 7577374599 7865528281 7502668089 6478121664
8879402347 9188900747 0651013874 0661686232 6574172892
2091782283 8847385272 1451440768 4098185699 0394877267
8762006403 4151547491 9957310565 6350053361 7929562433
2977588046 9583778014 9413020277 4813233724 6531483457
8441835645 6776947705 4429028815 3681963713 9807897545
3575240128 0683030949 8399146211 3127001067 ....

941 1062699256 1105207226 3549415515 4091392136 0255047821
4665249734 3251859723 6981934112 6461211477 1519659936
2380446333 6875664187 0350690754 5164718384 6971307120
0850159404 8884165781 0839532412 3273113708 8204038257
1732199787 4601487778 9585547290 1168969181 7215727948
9904357066 9500531349 6280552603 6131774707 7577045696
0680127523 9107332624 8671625929 8618490967 0563230605
7385759829 9681190223 1668437832 0935175345 3772582359
1923485653 5600425079 7024442082 8905419766 2061636556
8544102019 1285866099 8937300743 8894792773 6450584484
5908607863 9744952178 5334750265 6748140276 3018065887
3538788522 8480340063 7619553666 3124335812 9649309245
4835281615 3028692879 9149840595 1115834218 9160467587
6726886291 1795961742 8267800212 5398512221 0414452709
8831030818 2784272051 0095642933 0499468650 3719447396
3868225292 2422954303 9319872476 0892667375 1328374070
1381509032 9436769394 2614240170 0318809776 8331562167
9064824654 6227417640 8076514346 4399574920 2975557917
1094580233 7938363443 1455897980 8714133900 1062699256 ....

947 1055966209 0813093980 9926082365 3643083421 3305174234
4244984160 5068637803 5902851108 7645195353 7486800422
3864836325 2375923970 4329461457 2333685322 0696937697
9936642027 4551214361 1404435058 0781414994 7201689545
9345300950 3695881731 7845828933 4741288278 7750791974
6568109820 4857444561 7740232312 5659978880 6758183738
1203801478 3526927138 3315733896 5153115100 3167898627
2439281942 9778247096 0929250263 9915522703 2734952481
5205913410 7708553326 2935586061 2460401267 1594508975
7127771911 2988384371 7001055966 ....

953 1049317943 3368310598 1112277019 9370409233 9979013641
1332633788 0377754459 6012591815 3200419727 1773347324
2392444910 8079748163 6935991605 4564533053 5152151101
7838405036 7261280167 8908709338 9296956977 9643231899
2654774396 6421825813 2214060860 4407135362 0146904512
0671563483 7355718782 7911857292 7597061909 7586568730
3252885624 3441762854 1448058761 8048268625 3934942287
5131164742 9171038824 .7639034627 4921301154 2497376705
1416579223 5047219307 4501573976 9150052465 8971668415
5299055613 8509968520 4616998950 6820566631 6894018887
7229800629 5907660020 9863588667 3662119622 2455403987
4081846799 5802728226 6526757607 5550891920 2518363064
0083945435 4669464847 8488982161 5949632738 7198321091
2906610703 0430220356 7681007345 2256033578 1741867785
9391395592 8646379853 0954879328 4365162644 2812172088
1427072402 9380902413 4312696747 1143756558 2371458551
9412381951 7313746065 0577124868 8352570828 9611752360
9653725078 6988457502 6232948583 4207764952 7806925498
4260230849 9475341028 3315844700 9443861490 0314795383
00<u>10493179</u> . . . .

961 1040582726 3267429760 6659729448 4911550468 2622268470
3433922996 8782518210 1977107180 0208116545 2653485952
1331945889 6982310093 6524453694 0686784599 3756503642
0395421436 0041623309 0530697190 4266389177 9396462018
7304890738 8137356919 8751300728 4079084287 2008324661
8106139438 0853277835 5879292403 7460978147 7627471383
9750260145 6815816857 4401664932 3621227887 6170655567
1175858480 7492195629 5525494276 7950052029 1363163371
4880332986 4724245577 5234131113 4235171696 1498439125
9105098855 3590<u>10405</u> . . . .

967 1034126163 3919338159 2554291623 5780765253 3609100310
2378490175 8014477766 2874870734 2295760082 7300930713
5470527404 3433298862 4612202688 7280248190 2792140641
1582213029 9896587383 6608066184 0744570837 6421923474
6639089968 9762150982 4198552223 3712512926 5770423991
7269966928 6452947259 5656670113 7538779731 1271975180
9720785935 8841778697 00<u>10341261</u> . . . .

971 1029866117 4047373841 4006179196 7044284243 0484037075
1802265705 4582904222 4510813594 2327497425 3347064881
5653964984 5520082389 2893923789 9073120494 3357363542
7394438722 9660144181 2564366632 3377960865 0875386199
7940267765 1905252317 1987641606 5911431513 9031925849
6395468589 0834191555 0978372811 5345005149 3305870236
8692070030 8959835221 4212152420 1853759011 3285272914
5211122554 0679711637 4871266735 3244078269 8249227600
4119464469 6189495365 6024716786 8177136972 1936148300
7209062821 8331616889 8043254376 9309989701 3388259526
2615859938 2080329557 1575695159 6292481977 3429454170
9577754891 8640576725 0257466529 3511843460 3501544799
1761071060 7621009268 7950566426 3645726055 6127703398
5581874356 3336766220 3913491246 1380020597 3223480947
4768280123 5839340885 6848609680 7415036045 3141091658
0844490216 2718846549 9485066941 2976313079 2996910401
6477857878 4757981462 4098867147 2708547888 7744593202
8836251287 3326467559 2173017507 7239958805 3553038105
0463439752 8321318228 6302780638 5169927909 3717816683
8311019567 4562306900 1029866117 ....

977 1023541453 4288638689 8669396110 5424769703 1729785056
2947799385 8751279426 8167860798 3623336745 1381780962
1289662231 3203684749 2323439099 2835209825 9979529178
9314227226 2026612077 7891504605 9365404298 8741044012
2824974411 4636642784 0327533265 0972364380 7574206755
3735926305 0153531218 0143295803 4800409416 5813715455
4759467758 4442169907 8812691914 0225179119 7543500511
7707267144 3193449334 6980552712 3848515864 8925281473
8996929375 6397134083 9303991811 6683725690 8904810644
8311156601 8423746161 7195496417 6049129989 7645854657
1136131013 3060388945 7523029682 7021494370 5220061412
4872057318 3213920163 7666325486 1821903787 1033776867
9631525076 7656090071 6479017400 2047082906 8577277379
7338792221 0849539406 3459570112 5895598771 7502558853
6335721596 7246673490 2763561924 2579324462 6407369498
4646878198 5670419651 9959058341 8628454452 4053224155
5783009211 8730808597 7482088024 5649948822 9273285568
0655066530 1944728761 5148413510 7471852610 0307062436
0286591606 9600818833 1627430910 9518935516 8884339815
7625383828 0450358239 5087001023 ....

983 1017293997 9654120040 6917599186 1648016276 7039674465
9206510681 5869786368 2604272634 7914547304 1709053916
5818921668 3621566632 7568667344 8626653102 7466937945
0661241098 6775178026 4496439471 0071210579 8575788402
8484231943 0315361159 3692777212 6144455747 7110885045
7782299084 4354018311 2919633774 1607324516 7853509664
2929806714 1403865717 1922685656 1546286876 9074262461
8514750762 9704984740 5900305188 1993896236 0122075279
7558494404 8830111902 3397761953 2044760935 9104781281
7904374364 1912512716 1749745676 5005086469 9898270600
2034587995 9308240081 3835198372 3296032553 4079348931
8413021363 1739572736 5208545269 5829094608 3418107833
1637843336 7243133265 5137334689 7253306205 4933875890
1322482197 3550556052 8992878942 0142421159 7151576805
6968463886 0630722278 7385554425 2288911495 4221770091
5564598168 8708036622 5839267548 3214649033 5707019328
5859613428 2807731434 3845371312 3092573753 8148524923
7029501525 9409969481 1800610376 3987792472 0244150559
5116988809 7660223804 6795523906 4089521871 8209562563
5808748728 3825025432 3499491353 00<u>10172939</u> ....

991 1009081735 6205852674 0665993945 5095862764 8839556004 0363269424
8234106962 6639757820 3834510595 3582240161 4530776992 9364278506
5590312815 3380423814 3289606458 1231079717 4571140262 3612512613
5216952573 1584258324 9243188698 9845610494 4500504540 8678102926
3370332996 9727547931 3824419778 0020181634 7124117053 4813319878
9101917255 2976791120 0807265388 4964682139 2532795156 4076690211
9071644803 2290615539 8587285570 1311806256 3067608476 2865792129
1624621594 3491422805 2472250252 2704339051 4631685166 4984863773
9656912209 88900<u>10090</u> ....

997 1003009027 0812437311 9358074222 6680040120 3610832497 4924774322
9689067201 6048144433 2998996990 9729187562 6880641925 7773319959
8796389167 5025075225 6770310932 7983951855 566700<u>1003</u> ....

# TABLE OF THE FREQUENCY OF PRIME NUMBERS.

| | | | | | | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | 168 | 51 | 89 | 101 | 81 | 151 | 85 | 201 | 77 | 251 | 71 | 301 | 85 | 351 | 74 | 401 | 70 | 451 | 92 |
| 2 | 135 | 52 | 97 | 102 | 93 | 152 | 90 | 202 | 87 | 252 | 88 | 302 | 83 | 352 | 80 | 402 | 71 | 452 | 76 |
| 3 | 127 | 53 | 89 | 103 | 87 | 153 | 88 | 203 | 78 | 253 | 78 | 303 | 72 | 353 | 82 | 403 | 76 | 453 | 63 |
| 4 | 120 | 54 | 92 | 104 | 80 | 154 | 77 | 204 | 78 | 254 | 81 | 304 | 84 | 354 | 79 | 404 | 75 | 454 | 72 |
| 5 | 119 | 55 | 90 | 105 | 91 | 155 | 84 | 205 | 77 | 255 | 76 | 305 | 88 | 355 | 87 | 405 | 70 | 455 | 74 |
| 6 | 114 | 56 | 93 | 106 | 82 | 156 | 85 | 206 | 85 | 256 | 87 | 306 | 80 | 356 | 79 | 406 | 83 | 456 | 82 |
| 7 | 117 | 57 | 99 | 107 | 92 | 157 | 76 | 207 | 83 | 257 | 72 | 307 | 82 | 357 | 67 | 407 | 67 | 457 | 73 |
| 8 | 107 | 58 | 91 | 108 | 76 | 158 | 88 | 208 | 87 | 258 | 78 | 308 | 73 | 358 | 80 | 408 | 81 | 458 | 77 |
| 9 | 110 | 59 | 90 | 109 | 91 | 159 | 87 | 209 | 85 | 259 | 86 | 309 | 76 | 359 | 83 | 409 | 79 | 459 | 75 |
| 10 | 112 | 60 | 94 | 110 | 88 | 160 | 85 | 210 | 88 | 260 | 76 | 310 | 80 | 360 | 71 | 410 | 82 | 460 | 68 |
| 11 | 106 | 61 | 88 | 111 | 83 | 161 | 85 | 211 | 84 | 261 | 77 | 311 | 79 | 361 | 68 | 411 | 73 | 461 | 77 |
| 12 | 103 | 62 | 87 | 112 | 84 | 162 | 84 | 212 | 86 | 262 | 73 | 312 | 69 | 362 | 79 | 412 | 81 | 462 | 69 |
| 13 | 109 | 63 | 88 | 113 | 81 | 163 | 81 | 213 | 69 | 263 | 79 | 313 | 86 | 363 | 76 | 413 | 74 | 463 | 74 |
| 14 | 105 | 64 | 93 | 114 | 88 | 164 | 83 | 214 | 81 | 264 | 84 | 314 | 86 | 364 | 84 | 414 | 69 | 464 | 77 |
| 15 | 102 | 65 | 80 | 115 | 82 | 165 | 77 | 215 | 86 | 265 | 80 | 315 | 76 | 265 | 77 | 415 | 90 | 465 | 85 |
| 16 | 108 | 66 | 98 | 116 | 93 | 166 | 80 | 216 | 74 | 266 | 78 | 316 | 77 | 366 | 77 | 416 | 80 | 466 | 74 |
| 17 | 98 | 67 | 84 | 117 | 81 | 167 | 81 | 217 | 76 | 267 | 87 | 317 | 84 | 367 | 85 | 417 | 67 | 467 | 69 |
| 18 | 104 | 68 | 99 | 118 | 90 | 168 | 83 | 218 | 80 | 268 | 94 | 318 | 84 | 368 | 79 | 418 | 82 | 468 | 83 |
| 19 | 94 | 69 | 80 | 119 | 79 | 169 | 73 | 219 | 84 | 269 | 75 | 319 | 81 | 369 | 72 | 419 | 85 | 469 | 85 |
| 20 | 102 | 70 | 81 | 120 | 87 | 170 | 87 | 220 | 91 | 270 | 78 | 320 | 86 | 370 | 68 | 420 | 75 | 470 | 72 |
| 21 | 98 | 71 | 98 | 121 | 88 | 171 | 87 | 221 | 78 | 271 | 84 | 321 | 79 | 371 | 70 | 421 | 75 | 471 | 87 |
| 22 | 104 | 72 | 95 | 122 | 86 | 172 | 81 | 222 | 80 | 272 | 78 | 322 | 80 | 372 | 76 | 422 | 73 | 472 | 78 |
| 23 | 100 | 73 | 90 | 123 | 88 | 173 | 89 | 223 | 81 | 273 | 83 | 323 | 81 | 373 | 81 | 423 | 77 | 473 | 73 |
| 24 | 104 | 74 | 83 | 124 | 88 | 174 | 79 | 224 | 80 | 274 | 71 | 324 | 71 | 374 | 73 | 424 | 83 | 474 | 78 |
| 25 | 94 | 75 | 92 | 125 | 83 | 175 | 83 | 225 | 83 | 275 | 80 | 325 | 87 | 375 | 82 | 425 | 81 | 475 | 80 |
| 26 | 98 | 76 | 91 | 126 | 84 | 176 | 75 | 226 | 84 | 276 | 83 | 326 | 85 | 376 | 85 | 426 | 74 | 476 | 86 |
| 27 | 101 | 77 | 83 | 127 | 83 | 177 | 95 | 227 | 76 | 277 | 83 | 327 | 73 | 377 | 80 | 427 | 71 | 477 | 75 |
| 28 | 94 | 78 | 95 | 128 | 86 | 178 | 73 | 228 | 80 | 278 | 74 | 328 | 86 | 378 | 71 | 428 | 78 | 478 | 69 |
| 29 | 98 | 79 | 84 | 129 | 89 | 179 | 89 | 229 | 89 | 279 | 81 | 329 | 73 | 379 | 77 | 429 | 71 | 479 | 85 |
| 30 | 92 | 80 | 91 | 130 | 83 | 180 | 94 | 230 | 88 | 280 | 73 | 330 | 81 | 380 | 83 | 430 | 89 | 480 | 71 |
| 31 | 95 | 81 | 88 | 131 | 85 | 181 | 71 | 231 | 84 | 281 | 87 | 331 | 80 | 381 | 72 | 431 | 76 | 481 | 77 |
| 32 | 92 | 82 | 92 | 132 | 83 | 182 | 79 | 232 | 78 | 282 | 85 | 332 | 82 | 382 | 76 | 432 | 79 | 482 | 78 |
| 33 | 106 | 83 | 89 | 133 | 87 | 183 | 91 | 233 | 76 | 283 | 77 | 333 | 72 | 383 | 74 | 433 | 84 | 483 | 82 |
| 34 | 100 | 84 | 84 | 134 | 82 | 184 | 79 | 234 | 71 | 284 | 72 | 334 | 80 | 384 | 81 | 434 | 80 | 484 | 75 |
| 35 | 94 | 85 | 87 | 135 | 80 | 185 | 83 | 235 | 87 | 285 | 90 | 335 | 77 | 385 | 78 | 435 | 85 | 485 | 65 |
| 36 | 92 | 86 | 85 | 136 | 89 | 186 | 91 | 236 | 73 | 286 | 77 | 336 | 77 | 386 | 80 | 436 | 82 | 486 | 63 |
| 37 | 99 | 87 | 88 | 137 | 96 | 187 | 79 | 237 | 76 | 287 | 71 | 337 | 84 | 387 | 78 | 437 | 73 | 487 | 82 |
| 38 | 94 | 88 | 93 | 138 | 80 | 188 | 87 | 238 | 73 | 288 | 71 | 338 | 80 | 388 | 69 | 438 | 70 | 488 | 78 |
| 39 | 90 | 89 | 76 | 139 | 85 | 189 | 80 | 239 | 87 | 289 | 85 | 339 | 77 | 389 | 75 | 439 | 75 | 489 | 83 |
| 40 | 96 | 90 | 94 | 140 | 84 | 190 | 88 | 240 | 79 | 290 | 84 | 340 | 68 | 390 | 84 | 440 | 75 | 490 | 78 |
| 41 | 88 | 91 | 89 | 141 | 87 | 191 | 75 | 241 | 80 | 291 | 84 | 341 | 84 | 391 | 81 | 441 | 79 | 491 | 78 |
| 42 | 101 | 92 | 85 | 142 | 87 | 192 | 81 | 242 | 91 | 292 | 77 | 342 | 77 | 392 | 79 | 442 | 72 | 492 | 76 |
| 43 | 102 | 93 | 97 | 143 | 82 | 193 | 89 | 243 | 76 | 293 | 78 | 343 | 77 | 393 | 86 | 443 | 85 | 493 | 67 |
| 44 | 85 | 94 | 86 | 144 | 77 | 194 | 84 | 244 | 77 | 294 | 68 | 344 | 80 | 394 | 87 | 444 | 88 | 494 | 82 |
| 45 | 96 | 95 | 87 | 145 | 79 | 195 | 74 | 245 | 78 | 295 | 85 | 345 | 80 | 395 | 75 | 445 | 82 | 495 | 80 |
| 46 | 86 | 96 | 95 | 146 | 85 | 196 | 85 | 246 | 80 | 296 | 75 | 346 | 76 | 396 | 72 | 446 | 68 | 496 | 87 |
| 47 | 90 | 97 | 84 | 147 | 84 | 197 | 76 | 247 | 84 | 297 | 82 | 347 | 80 | 397 | 75 | 447 | 68 | 497 | 68 |
| 48 | 95 | 98 | 82 | 148 | 83 | 198 | 87 | 248 | 79 | 298 | 73 | 348 | 82 | 398 | 75 | 448 | 73 | 498 | 81 |
| 49 | 89 | 99 | 87 | 149 | 83 | 199 | 96 | 249 | 88 | 299 | 73 | 349 | 77 | 399 | 82 | 449 | 70 | 499 | 72 |
| 50 | 98 | 100 | 87 | 150 | 91 | 200 | 77 | 250 | 80 | 300 | 78 | 350 | 82 | 400 | 81 | 450 | 80 | 500 | 81 |

| 501 78 | 551 79 | 601 75 | 651 61 | 701 75 | 751 68 | 801 85 | 851 70 | 901 74 | 951 76 |
|---|---|---|---|---|---|---|---|---|---|
| 502 74 | 552 75 | 602 73 | 652 74 | 702 71 | 752 85 | 802 66 | 852 77 | 902 73 | 952 70 |
| 503 67 | 553 71 | 603 83 | 653 85 | 703 81 | 753 73 | 803 70 | 853 74 | 903 70 | 953 78 |
| 504 76 | 554 80 | 604 76 | 654 69 | 704 71 | 754 71 | 804 69 | 854 66 | 904 63 | 954 65 |
| 505 76 | 555 77 | 605 73 | 655 78 | 705 87 | 755 83 | 805 78 | 855 71 | 905 81 | 955 73 |
| 506 83 | 556 61 | 606 74 | 656 73 | 706 68 | 756 70 | 806 79 | 856 73 | 906 70 | 956 76 |
| 507 76 | 557 88 | 607 72 | 657 71 | 707 82 | 757 66 | 807 68 | 857 78 | 907 80 | 957 58 |
| 508 71 | 558 68 | 608 78 | 658 70 | 708 74 | 758 68 | 808 70 | 858 76 | 908 80 | 958 69 |
| 509 76 | 559 74 | 609 78 | 659 79 | 709 77 | 759 79 | 809 69 | 859 69 | 909 79 | 959 77 |
| 510 75 | 560 77 | 610 80 | 660 73 | 710 77 | 760 77 | 810 78 | 860 71 | 910 82 | 960 69 |
| 511 72 | 561 86 | 611 73 | 661 83 | 711 78 | 761 77 | 811 78 | 861 77 | 911 62 | 961 68 |
| 512 82 | 562 61 | 612 71 | 662 70 | 712 76 | 762 80 | 812 72 | 862 74 | 912 81 | 962 88 |
| 513 70 | 563 83 | 613 76 | 663 74 | 713 72 | 763 68 | 813 69 | 863 83 | 913 71 | 963 71 |
| 514 77 | 564 67 | 614 79 | 664 77 | 714 73 | 764 79 | 814 72 | 864 60 | 914 54 | 964 74 |
| 515 81 | 565 77 | 615 71 | 665 77 | 715 66 | 765 72 | 815 78 | 865 80 | 915 73 | 965 74 |
| 516 66 | 566 78 | 616 75 | 666 77 | 716 83 | 766 82 | 816 69 | 866 80 | 916 70 | 966 70 |
| 517 85 | 567 72 | 617 85 | 667 73 | 717 69 | 767 78 | 817 75 | 867 68 | 917 72 | 967 73 |
| 518 83 | 568 72 | 618 81 | 668 73 | 718 65 | 768 68 | 818 75 | 868 78 | 918 79 | 968 66 |
| 519 76 | 569 71 | 619 67 | 669 66 | 719 67 | 769 77 | 819 62 | 869 80 | 919 75 | 969 75 |
| 520 78 | 570 80 | 620 73 | 670 74 | 720 74 | 770 74 | 820 83 | 870 73 | 920 71 | 970 73 |
| 521 73 | 571 85 | 621 77 | 671 75 | 721 78 | 771 77 | 821 75 | 871 79 | 921 72 | 971 76 |
| 522 83 | 572 72 | 622 70 | 672 76 | 722 77 | 772 75 | 822 72 | 872 58 | 922 72 | 972 78 |
| 523 79 | 573 85 | 623 74 | 673 76 | 723 73 | 773 70 | 823 84 | 873 76 | 923 72 | 973 74 |
| 524 69 | 574 72 | 624 75 | 674 77 | 724 86 | 774 76 | 824 78 | 874 65 | 924 81 | 974 63 |
| 525 77 | 575 70 | 625 68 | 675 69 | 725 75 | 775 72 | 825 71 | 875 75 | 925 76 | 975 85 |
| 526 79 | 576 77 | 626 69 | 676 75 | 726 69 | 776 67 | 826 81 | 876 80 | 926 80 | 976 70 |
| 527 84 | 577 78 | 627 70 | 677 74 | 727 76 | 777 70 | 827 78 | 877 75 | 927 74 | 977 66 |
| 528 72 | 578 77 | 628 70 | 678 63 | 728 75 | 778 76 | 828 69 | 878 67 | 928 63 | 978 60 |
| 529 70 | 579 76 | 629 71 | 679 82 | 729 76 | 779 81 | 829 69 | 879 68 | 929 70 | 979 80 |
| 530 78 | 580 77 | 630 75 | 680 83 | 730 75 | 780 71 | 830 68 | 880 75 | 930 80 | 980 65 |
| 531 80 | 581 73 | 631 67 | 681 75 | 731 69 | 781 70 | 831 76 | 881 80 | 931 69 | 981 67 |
| 532 68 | 582 79 | 632 81 | 682 78 | 732 76 | 782 82 | 832 79 | 882 69 | 932 69 | 982 75 |
| 533 79 | 583 73 | 633 77 | 683 66 | 733 71 | 783 68 | 833 82 | 883 72 | 933 76 | 983 70 |
| 534 74 | 584 78 | 634 70 | 684 78 | 734 75 | 784 74 | 834 68 | 884 73 | 934 68 | 984 70 |
| 535 72 | 585 72 | 635 82 | 685 72 | 735 74 | 785 75 | 835 67 | 885 69 | 935 81 | 985 74 |
| 536 71 | 586 81 | 636 78 | 686 74 | 736 79 | 786 77 | 836 73 | 886 77 | 936 68 | 986 76 |
| 537 87 | 587 79 | 637 73 | 687 74 | 737 69 | 787 70 | 837 71 | 887 76 | 937 71 | 987 76 |
| 538 67 | 588 87 | 638 74 | 688 82 | 738 78 | 788 73 | 838 64 | 888 71 | 938 71 | 988 63 |
| 539 78 | 589 73 | 639 59 | 689 74 | 739 70 | 789 80 | 839 80 | 889 77 | 939 72 | 989 71 |
| 540 71 | 590 68 | 640 72 | 690 79 | 740 81 | 790 68 | 840 69 | 890 68 | 940 68 | 990 72 |
| 541 73 | 591 71 | 641 77 | 691 60 | 741 67 | 791 78 | 841 70 | 891 68 | 941 74 | 991 71 |
| 542 77 | 592 67 | 642 71 | 692 79 | 742 74 | 792 71 | 842 69 | 892 80 | 942 79 | 992 79 |
| 543 78 | 593 80 | 643 68 | 693 77 | 743 73 | 793 82 | 843 83 | 893 69 | 943 72 | 993 65 |
| 544 81 | 594 77 | 644 70 | 694 73 | 744 67 | 794 71 | 844 68 | 894 72 | 944 76 | 994 68 |
| 545 68 | 595 78 | 645 86 | 695 76 | 745 64 | 795 73 | 845 78 | 895 74 | 945 73 | 995 78 |
| 546 68 | 596 77 | 646 75 | 696 77 | 746 67 | 796 79 | 846 70 | 896 80 | 946 66 | 996 69 |
| 547 73 | 597 79 | 647 74 | 697 73 | 747 76 | 797 77 | 847 69 | 897 64 | 947 72 | 997 69 |
| 548 76 | 598 73 | 648 79 | 698 79 | 748 71 | 798 72 | 848 77 | 898 75 | 948 66 | 998 83 |
| 549 77 | 599 72 | 649 73 | 699 62 | 749 76 | 799 71 | 849 75 | 899 76 | 949 67 | 999 74 |
| 550 78 | 600 73 | 650 84 | 700 72 | 750 72 | 800 81 | 850 68 | 900 61 | 950 75 | 1000 65 |

1000000 . . . 1100000

| | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | 1 | | | | | | | | | | 1 |
| 2 | | 1 | | | | 1 | | 1 | 1 | | 4 |
| 3 | | 4 | 2 | 2 | 3 | 1 | 2 | 3 | 3 | 1 | 21 |
| 4 | 2 | 8 | 5 | 4 | 3 | 6 | 9 | 4 | 5 | 8 | 54 |
| 5 | 11 | 10 | 8 | 18 | 12 | 10 | 10 | 12 | 15 | 8 | 114 |
| 6 | 14 | 14 | 18 | 21 | 16 | 22 | 19 | 15 | 17 | 15 | 171 |
| 7 | 26 | 17 | 23 | 23 | 24 | 24 | 17 | 22 | 20 | 21 | 217 |
| 8 | 19 | 19 | 21 | 7 | 14 | 15 | 20 | 17 | 15 | 17 | 164 |
| 9 | 11 | 13 | 9 | 13 | 14 | 14 | 12 | 13 | 11 | 16 | 126 |
| 10 | 8 | 6 | 8 | 5 | 9 | 5 | 5 | 9 | 7 | 9 | 71 |
| 11 | 6 | 6 | 4 | 6 | 3 | 1 | 3 | 1 | 4 | 5 | 39 |
| 12 | 1 | 1 | 2 | 1 | 1 | 1 | 2 | 2 | 1 | | 12 |
| 13 | 1 | 1 | | | 1 | | 1 | 1 | 1 | | 6 |
| 14 | | | | | | | | | | | |
| 15 | | | | | | | | | | | |
| 16 | | | | | | | | | | | |
| | 752 | 719 | 732 | 700 | 731 | 698 | 713 | 722 | 706 | 737 | 7210 |

$$\int \frac{\mathrm{d}x}{\log x} = 7212{,}99$$

1100000 . . . 1200000

| | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 0 | | | | | | | | | | | 0 |
| 1 | | | | | 1 | | | | | | 1 |
| 2 | | | 1 | | | | 1 | 1 | | 2 | 5 |
| 3 | 4 | 3 | 3 | 3 | 3 | 3 | 3 | 2 | | 1 | 25 |
| 4 | 5 | 6 | 7 | 5 | 9 | 4 | 4 | 5 | 6 | 6 | 57 |
| 5 | 8 | 13 | 10 | 12 | 11 | 11 | 9 | 12 | 12 | 9 | 107 |
| 6 | 14 | 20 | 17 | 20 | 17 | 17 | 18 | 16 | 17 | 14 | 170 |
| 7 | 21 | 19 | 22 | 19 | 21 | 21 | 20 | 18 | 29 | 27 | 217 |
| 8 | 22 | 13 | 10 | 12 | 18 | 20 | 17 | 19 | 9 | 20 | 160 |
| 9 | 13 | 14 | 16 | 17 | 7 | 11 | 16 | 14 | 12 | 11 | 131 |
| 10 | 9 | 6 | 10 | 10 | 8 | 9 | 6 | 8 | 3 | 8 | 77 |
| 11 | 1 | 4 | 2 | 2 | 2 | 1 | 4 | 5 | 9 | 2 | 32 |
| 12 | 1 | 1 | 1 | | 2 | 3 | 1 | | 2 | | 11 |
| 13 | 1 | 1 | 1 | | | | 1 | | 1 | | 5 |
| 14 | 1 | | | | 1 | | | | | | 2 |
| | 736 | 710 | 716 | 713 | 697 | 725 | 729 | 723 | 735 | 710 | 7194 |

$$\int \frac{\mathrm{d}x}{\log x} = 7165{,}911$$

1200000 . . . 1300000

| | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 0 | | | | | | | | | | | |
| 1 | | | | | | | | 1 | 1 | | 2 |
| 2 | 2 | | 2 | 1 | | | | | | 1 | 6 |
| 3 | 3 | 2 | 4 | 5 | 4 | 3 | 1 | 4 | 3 | 3 | 32 |
| 4 | 7 | 7 | 7 | 3 | 5 | 7 | 12 | 2 | 3 | 10 | 63 |
| 5 | 15 | 12 | 12 | 15 | 10 | 14 | 9 | 15 | 6 | 12 | 120 |
| 6 | 16 | 14 | 13 | 19 | 17 | 16 | 16 | 15 | 20 | 14 | 160 |
| 7 | 24 | 15 | 25 | 24 | 21 | 20 | 15 | 22 | 24 | 24 | 214 |
| 8 | 17 | 19 | 16 | 11 | 17 | 15 | 22 | 18 | 19 | 14 | 168 |
| 9 | 8 | 12 | 7 | 10 | 12 | 13 | 14 | 13 | 13 | 9 | 111 |
| 10 | 3 | 11 | 10 | 8 | 10 | 4 | 5 | 3 | 9 | 10 | 73 |
| 11 | 3 | 6 | 3 | 2 | 3 | 5 | 3 | 6 | 1 | 3 | 35 |
| 12 | 1 | 1 | 1 | | | 1 | 3 | 1 | 1 | | 9 |
| 13 | 1 | 1 | | 1 | | 2 | | | | | 5 |
| 14 | | | | 1 | | | | | | | 1 |
| 16 | | | | | 1 | | | | | | 1 |
| | 676 | 744 | 693 | 693 | 724 | 713 | 718 | 709 | 722 | 689 | 7081 |

$$\int \frac{\mathrm{d}x}{\log x} = 7123{,}35$$

1300000 . . . 1400000

| | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | | | | | | 1 | | | | | 1 |
| 2 | | 1 | 1 | | 1 | 1 | 1 | 2 | 1 | 1 | 9 |
| 3 | 3 | 1 | 1 | 3 | 2 | 2 | 2 | | 5 | | 19 |
| 4 | 3 | 10 | 7 | 11 | 6 | 8 | 7 | 6 | 6 | 5 | 69 |
| 5 | 17 | 13 | 11 | 11 | 15 | 12 | 8 | 8 | 14 | 10 | 119 |
| 6 | 15 | 14 | 17 | 14 | 20 | 18 | 23 | 17 | 16 | 19 | 173 |
| 7 | 22 | 18 | 26 | 18 | 21 | 16 | 16 | 28 | 19 | 23 | 207 |
| 8 | 14 | 22 | 14 | 16 | 14 | 13 | 21 | 17 | 15 | 15 | 161 |
| 9 | 17 | 11 | 12 | 14 | 12 | 9 | 12 | 7 | 13 | 13 | 120 |
| 10 | 5 | 6 | 2 | 11 | 5 | 15 | 5 | 8 | 6 | 7 | 70 |
| 11 | 4 | 2 | 7 | 1 | 1 | 3 | 3 | 6 | 3 | 3 | 33 |
| 12 | | 1 | 2 | 1 | 2 | 2 | 2 | 1 | 1 | 3 | 15 |
| 13 | | 1 | | | 1 | | | | 1 | | 3 |
| 14 | | | | | | | | | | 1 | 1 |
| | 709 | 702 | 713 | 705 | 692 | 713 | 709 | 723 | 695 | 737 | 7098 |

$$\int \frac{\mathrm{d}x}{\log x} = 7084{,}48$$

1400000 . . . 1500000

| | 141 | 142 | 143 | 144 | 145 | 146 | 147 | 148 | 149 | 150 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 0 | | | | | | | | | | | 0 |
| 1 | | | 1 | | 1 | 1 | 1 | | | 1 | 5 |
| 2 | | 1 | 1 | | 2 | | | | 2 | 1 | 7 |
| 3 | 3 | 3 | 0 | 2 | 1 | 2 | 2 | 4 | 2 | 0 | 19 |
| 4 | 8 | 8 | 8 | 4 | 6 | 7 | 9 | 9 | 5 | 8 | 72 |
| 5 | 17 | 9 | 7 | 14 | 13 | 11 | 14 | 15 | 16 | 13 | 129 |
| 6 | 21 | 23 | 20 | 18 | 20 | 19 | 11 | 16 | 16 | 19 | 183 |
| 7 | 17 | 23 | 18 | 18 | 13 | 24 | 18 | 11 | 15 | 22 | 179 |
| 8 | 12 | 16 | 28 | 17 | 24 | 14 | 17 | 18 | 23 | 14 | 183 |
| 9 | 12 | 11 | 4 | 15 | 6 | 10 | 13 | 12 | 7 | 8 | 98 |
| 10 | 7 | 2 | 7 | 7 | 9 | 7 | 8 | 9 | 8 | 9 | 73 |
| 11 | 2 | 2 | 2 | 4 | 2 | 3 | 6 | 5 | 4 | 4 | 34 |
| 12 | 1 | 2 | 3 | 1 | 2 | 2 | 1 | 1 | 2 | 1 | 16 |
| 13 | | | 1 | | 1 | | | | | | 2 |
| | 679 | 680 | 717 | 723 | 703 | 701 | 716 | 705 | 706 | 698 | 7028 |

$$\int \frac{\mathrm{d}x}{\log x} = 7048{,}78186$$

1500000 . . . 1600000

| | 151 | 152 | 153 | 154 | 155 | 156 | 157 | 158 | 159 | 160 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 0 | | | | | | | | | | | 0 |
| 1 | | | | | | 1 | 1 | | | | 2 |
| 2 | | | 1 | 1 | 2 | | 2 | | 1 | 3 | 10 |
| 3 | 2 | | 4 | 2 | 2 | 3 | 5 | 3 | 6 | 1. | 28 |
| 4 | 8 | 5 | 5 | 7 | 9 | 13 | 6 | 10 | 7 | 7 | 77 |
| 5 | 8 | 19 | 9 | 13 | 11 | 9 | 12 | 15 | 11 | 17 | 124 |
| 6 | 16 | 20 | 25 | 21 | 20 | 12 | 26 | 14 | 23 | 22 | 199 |
| 7 | 19 | 21 | 18 | 19 | 18 | 15 | 12 | 19 | 11 | 20 | 172 |
| 8 | 19 | 12 | 15 | 18 | 15 | 17 | 10 | 17 | 15 | 11 | 149 |
| 9 | 16 | 14 | 16 | 12 | 8 | 17 | 15 | 6 | 11 | 9 | 124 |
| 10 | 8 | 3 | 5 | 4 | 9 | 7 | 5 | 10 | 10 | 2 | 63 |
| 11 | | 5 | 2 | 2 | 3 | 3 | 3 | 2 | 4 | 5 | 29 |
| 12 | 4 | | | 1 | 3 | 1 | 1 | 4 | 1 | 2 | 17 |
| 13 | | 1 | | | | 2 | 2 | | | 1 | 6 |
| | 731 | 702 | 691 | 686 | 698 | 714 | 680 | 701 | 693 | 675 | 6971 |

$$\int \frac{\mathrm{d}x}{\log x} = 7015{,}78776$$

1600000 . . . 1700000

| | 161 | 162 | 163 | 164 | 165 | 166 | 167 | 168 | 169 | 170 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 0 | | | | | | | | 1 | | | 1 |
| 1 | | | | | | 2 | | | | | 2 |
| 2 | 1 | 3 | 1 | | 1 | 1 | 2 | | | 2 | 11 |
| 3 | 3 | 3 | 3 | 4 | 4 | 2 | 2 | 4 | 4 | | 29 |
| 4 | 7 | 4 | 9 | 7 | 7 | 10 | 4 | 4 | 10 | 6 | 68 |
| 5 | 10 | 11 | 8 | 12 | 11 | 11 | 13 | 12 | 18 | 14 | 120 |
| 6 | 18 | 22 | 15 | 19 | 15 | 11 | 14 | 19 | 10 | 16 | 159 |
| 7 | 22 | 15 | 14 | 21 | 25 | 18 | 22 | 24 | 24 | 18 | 203 |
| 8 | 14 | 23 | 25 | 15 | 17 | 16 | 17 | 15 | 13 | 19 | 174 |
| 9 | 8 | 12 | 18 | 12 | 12 | 21 | 12 | 8 | 14 | 13 | 130 |
| 10 | 7 | 2 | 5 | 8 | 4 | 6 | 11 | 7 | 4 | 9 | 63 |
| 11 | 7 | 3 | 1 | 1 | 3 | 1 | 3 | 2 | 2 | 3 | 26 |
| 12 | 2 | 1 | 1 | 1 | | | | 4 | | | 9 |
| 13 | 1 | | | | 1 | | | | 1 | | 3 |
| 14 | | 1 | | | | | | | | | 1 |
| 15 | | | | | | 1 | | | | | 1 |
| | 719 | 694 | 710 | 692 | 692 | 700 | 716 | 702 | 675 | 712 | 7012 |

$$\int \frac{\mathrm{d}x}{\log x} = 6985{,}13714$$

1700000 . . . 1800000

| | 171 | 172 | 173 | 174 | 175 | 176 | 177 | 178 | 179 | 180 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 0 | | | | | | | | | | | 0 |
| 1 | | | | 1 | | | | | | | 1 |
| 2 | 1 | | | | | 2 | 1 | | | 1 | 5 |
| 3 | 3 | 4 | | 3 | 3 | 4 | 3 | 5 | 3 | 2 | 30 |
| 4 | 7 | 9 | 6 | 6 | 5 | 8 | 6 | 6 | 10 | 7 | 70 |
| 5 | 13 | 15 | 19 | 16 | 12 | 15 | 21 | 13 | 13 | 15 | 152 |
| 6 | 17 | 16 | 22 | 22 | 20 | 14 | 15 | 13 | 18 | 17 | 174 |
| 7 | 23 | 21 | 22 | 15 | 22 | 19 | 19 | 21 | 17 | 15 | 194 |
| 8 | 11 | 16 | 11 | 15 | 16 | 15 | 13 | 18 | 19 | 13 | 147 |
| 9 | 18 | 11 | 8 | 11 | 15 | 10 | 12 | 14 | 10 | 15 | 124 |
| 10 | 3 | 1 | 8 | 7 | 2 | 9 | 6 | 9 | 5 | 11 | 61 |
| 11 | 1 | 3 | 3 | 1 | 3 | 4 | 4 | 1 | 4 | 2 | 26 |
| 12 | 2 | 3 | | 1 | 1 | | | | 1 | 2 | 10 |
| 13 | 1 | 1 | 1 | 2 | | | | | | | 5 |
| 14 | | | | | | | | | | | |
| 15 | | | | | | | | | | | 1 |
| | 695 | 685 | 691 | 689 | 706 | 684 | 679 | 700 | 689 | 713 | 6931 |

$$\int \frac{\mathrm{d}x}{\log x} = 6956{,}53562$$

1800000 . . . 1900000

| | 181 | 182 | 183 | 184 | 185 | 186 | 187 | 188 | 189 | 190 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 0 | | | | | | | | | | | |
| 1 | | | | | | | | | | | |
| 2 | 1 | 1 | 2 | 1 | | 2 | | | | 3 | 10 |
| 3 | 3 | 2 | 1 | 5 | 1 | 1 | 1 | 2 | 3 | 3 | 22 |
| 4 | 6 | 5 | 10 | 12 | 7 | 6 | 5 | 5 | 8 | 7 | 71 |
| 5 | 14 | 15 | 10 | 11 | 11 | 12 | 19 | 17 | 12 | 14 | 135 |
| 6 | 13 | 20 | 14 | 15 | 21 | 19 | 16 | 19 | 23 | 15 | 175 |
| 7 | 25 | 26 | 18 | 17 | 21 | 21 | 20 | 22 | 19 | 17 | 206 |
| 8 | 15 | 19 | 13 | 18 | 22 | 15 | 19 | 15 | 11 | 14 | 161 |
| 9 | 10 | 7 | 19 | 13 | 9 | 8 | 10 | 12 | 10 | 15 | 113 |
| 10 | 9 | 4 | 8 | 6 | 6 | 13 | 4 | 6 | 8 | 10 | 74 |
| 11 | 2 | | 4 | | | 3 | 5 | 2 | 5 | 2 | 23 |
| 12 | 2 | 1 | 1 | 2 | 2 | | 1 | | 1 | | 10 |
| 13 | | | | | | | | | | | |
| 14 | | | | | | | | | | | |
| | 704 | 672 | 718 | 674 | 700 | 707 | 703 | 689 | 697 | 691 | 6955 |

$$\int \frac{\mathrm{d}x}{\log x} = 6929{,}73917$$

1900000 . . . 2000000

| | 191 | 192 | 193 | 194 | 195 | 196 | 197 | 198 | 199 | 200 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 0 | | | | | | | | | | | |
| 1 | 1 | | | | | | | | | | 1 |
| 2 | | | 1 | | | 1 | 2 | | 1 | | 5 |
| 3 | 4 | 3 | 1 | 2 | 10 | 1 | 3 | 4 | 4 | 2 | 34 |
| 4 | 5 | 4 | 4 | 6 | 4 | 9 | 7 | 10 | 11 | 7 | 67 |
| 5 | 12 | 18 | 15 | 18 | 11 | 12 | 11 | 16 | 11 | 12 | 136 |
| 6 | 19 | 20 | 18 | 16 | 17 | 24 | 20 | 20 | 18 | 10 | 182 |
| 7 | 21 | 20 | 23 | 27 | 20 | 16 | 25 | 17 | 21 | 31 | 221 |
| 8 | 16 | 10 | 16 | 14 | 14 | 18 | 17 | 15 | 8 | 20 | 148 |
| 9 | 15 | 14 | 8 | 8 | 9 | 12 | 8 | 8 | 15 | 6 | 103 |
| 10 | 5 | 6 | 8 | 6 | 11 | 4 | 5 | 5 | 6 | 6 | 62 |
| 11 | 2 | 4 | 6 | 2 | 3 | 2 | 1 | 3 | 2 | 5 | 30 |
| 12 | | 1 | | 1 | | | | 2 | 1 | | 5 |
| 13 | | | | | 1 | | 1 | | 1 | | 3 |
| 14 | | | | | | 1 | | | 1 | 1 | 3 |
| | 689 | 697 | 711 | 683 | 692 | 685 | 673 | 670 | 688 | 714 | 6902 |

$$\int \frac{\mathrm{d}x}{\log x} = 6904{,}54424$$

2000000 . . . 2100000

| | 201 | 202 | 203 | 204 | 205 | 206 | 207 | 208 | 209 | 210 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 0 | | | | | | | | | | | |
| 1 | | | | | 2 | 1 | | | | | 3 |
| 2 | | | 2 | 2 | 1 | | | 2 | 2 | 1 | 10 |
| 3 | 3 | 3 | 5 | 2 | 2 | 3 | 4 | 5 | 2 | 3 | 32 |
| 4 | 7 | 8 | 9 | 4 | 8 | 5 | 6 | 7 | 9 | 6 | 69 |
| 5 | 13 | 10 | 9 | 15 | 13 | 10 | 12 | 13 | 9 | 15 | 119 |
| 6 | 25 | 20 | 13 | 16 | 26 | 23 | 25 | 14 | 17 | 18 | 197 |
| 7 | 10 | 22 | 23 | 25 | 13 | 23 | 18 | 23 | 25 | 22 | 204 |
| 8 | 13 | 17 | 15 | 22 | 12 | 13 | 12 | 16 | 18 | 19 | 157 |
| 9 | 16 | 15 | 11 | 4 | 11 | 12 | 11 | 14 | 11 | 10 | 115 |
| 10 | 10 | 2 | 8 | 6 | 11 | 7 | 7 | 3 | 3 | 6 | 63 |
| 11 | 2 | 1 | 5 | 3 | | 1 | 3 | 3 | 3 | | 21 |
| 12 | | 2 | | | 1 | 2 | 2 | | 1 | | 8 |
| 13 | 1 | | | 1 | | | | | | | 2 |
| | 705 | 691 | 693 | 690 | 671 | 696 | 694 | 674 | 686 | 674 | 6874 |

$$\int \frac{\mathrm{d}x}{\log x} = 6880{,}780$$

2100000 . . . 2200000

| | 211 | 212 | 213 | 214 | 215 | 216 | 217 | 218 | 219 | 220 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 0 | | | | | | | | | | | |
| 1 | | 1 | | | | | 1 | | | | 2 |
| 2 | 1 | 1 | 1 | | | 1 | 1 | 1 | 1 | 2 | 9 |
| 3 | 5 | 3 | 3 | 1 | 4 | | 2 | 3 | 2 | 4 | 27 |
| 4 | 7 | 7 | 2 | 13 | 5 | 3 | 9 | 6 | 9 | 8 | 69 |
| 5 | 12 | 14 | 20 | 16 | 16 | 12 | 17 | 13 | 12 | 14 | 146 |
| 6 | 12 | 20 | 17 | 14 | 16 | 25 | 16 | 23 | 21 | 19 | 183 |
| 7 | 19 | 14 | 18 | 23 | 26 | 22 | 18 | 22 | 22 | 17 | 201 |
| 8 | 22 | 21 | 20 | 20 | 12 | 16 | 20 | 10 | 12 | 15 | 168 |
| 9 | 12 | 10 | 8 | 7 | 9 | 10 | 7 | 13 | 16 | 17 | 109 |
| 10 | 6 | 5 | 6 | 2 | 7 | 7 | 8· | 5 | 2 | 4 | 52 |
| 11 | 3 | 2 | 3 | 1 | 4 | 2 | 1 | | 2 | | 18 |
| 12 | 1 | 1 | 1 | 1 | 1 | 2 | | 2 | | | 9 |
| 13 | | 1 | 1 | 1 | | | | | 1 | | 4 |
| 14 | | | | 1 | | | | 2 | | | 3 |
| | 699 | 683 | 697 | 673 | 693 | 712 | 666 | 691 | 679 | 664 | 6857 |

$$\int \frac{\mathrm{d}x}{\log x} = 6858{,}292$$

2200000 . . . 2300000

| | 221 | 222 | 223 | 224 | 225 | 226 | 227 | 228 | 229 | 230 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | | | | | 1 | | | | 1 | | 2 |
| 2 | | | 1 | 1 | | 2 | | | 3 | 2 | 9 |
| 3 | 5 | 2 | 4 | 7 | 2 | 1 | 2 | 2 | 2 | 2 | 29 |
| 4 | 8 | 9 | 5 | 5 | 10 | 7 | 7 | 7 | 6 | 9 | 73 |
| 5 | 12 | 24 | 16 | 13 | 11 | 10 | 14 | 14 | 15 | 9 | 138 |
| 6 | 17 | 18 | 12 | 18 | 18 | 20 | 16 | 17 | 21 | 22 | 179 |
| 7 | 19 | 17 | 25 | 21 | 18 | 20 | 23 | 25 | 19 | 18 | 205 |
| 8 | 12 | 12 | 19 | 15 | 18 | 24 | 13 | 17 | 16 | 22 | 168 |
| 9 | 14 | 9 | 6 | 9 | 19 | 11 | 16 | 9 | 13 | 7 | 113 |
| 10 | 7 | 6 | 6 | 6 | 1 | 3 | 7 | 5 | | 3 | 44 |
| 11 | 5 | 2 | 4 | 4 | 2 | 2 | 2 | 2 | 2 | 5 | 30 |
| 12 | 1 | 1 | 2 | 1 | | | | 2 | 2 | 1 | 10 |
| | 701 | 660 | 695 | 680 | 683 | 688 | 701 | 694 | 662 | 685 | 6849 |

$$\int \frac{\mathrm{d}x}{\log x} = 6836{,}977$$

2300000 . . . 2400000

| | 231 | 232 | 233 | 234 | 235 | 236 | 237 | 238 | 239 | 240 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | 1 | 1 | 1 | | 1 | | | | | | 4 |
| 2 | | | 1 | | 2 | 2 | 1 | 2 | 1 | 2 | 11 |
| 3 | 4 | 1 | 2 | 3 | 4 | 3 | 2 | 6 | 4 | 3 | 32 |
| 4 | 8 | 12 | 10 | 9 | 8 | 7 | 3 | 13 | 7 | 9 | 86 |
| 5 | 13 | 18 | 13 | 14 | 17 | 12 | 10 | 13 | 10 | 16 | 136 |
| 6 | 15 | 16 | 21 | 20 | 16 | 21 | 26 | 11 | 14 | 16 | 176 |
| 7 | 22 | 25 | 20 | 20 | 13 | 16 | 26 | 18 | 19 | 15 | 194 |
| 8 | 13 | 9 | 16 | 21 | 17 | 15 | 15 | 15 | 21 | 16 | 158 |
| 9 | 13 | 9 | 6 | 7 | 14 | 14 | 8 | 13 | 14 | 14 | 112 |
| 10 | 7 | 5 | 3 | 2 | 5 | 5 | 8 | 7 | 8 | 5 | 55 |
| 11 | 3 | 3 | 4 | 3 | 3 | 5 | 1 | 2 | 2 | 2 | 28 |
| 12 | 1 | | 3 | 1· | | | | | | 2 | 7 |
| 13 | | 1 | | | | | | | | | 1 |
| | 690 | 662 | 672 | 671 | 666 | 690 | 691 | 660 | 705 | 680 | 6787 |

$$\int \frac{\mathrm{d}x}{\log x} = 6816{,}706$$

2400000 . . . 2500000

| | 241 | 242 | 243 | 244 | 245 | 246 | 247 | 248 | 249 | 250 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | | | | | | | | | 1 | | 1 |
| 2 | 2 | | | | | | 1 | 1 | 2 | 3 | 9 |
| 3 | 4 | 6 | 4 | 4 | 1 | 5 | 3 | 5 | 3 | 2 | 37 |
| 4 | 12 | 8 | 7 | 7 | 9 | 7 | 10 | 8 | 6 | 4 | 78 |
| 5 | 13 | 14 | 17 | 19 | 15 | 12 | 18 | 11 | 11 | 17 | 147 |
| 6 | 18 | 16 | 21 | 20 | 18 | 22 | 18 | 18 | 20 | 22 | 193 |
| 7 | 21 | 16 | 19 | 20 | 17 | 17 | 21 | 22 | 19 | 17 | 189 |
| 8 | 10 | 16 | 14 | 14 | 21 | 17 | 10 | 17 | 16 | 16 | 151 |
| 9 | 8 | 13 | 9 | 11 | 7 | 12 | 6 | 11 | 12 | 13 | 102 |
| 10 | 9 | 6 | 5 | 4 | 6 | 4 | 8 | 6 | 7 | 3 | 58 |
| 11 | 2 | 4 | 3 | 1 | 4 | 1 | 4 | 1 | 2 | 1 | 23 |
| 12 | 1 | | 1 | | 1 | 1 | 1 | | | 2 | 7 |
| 13 | | 1 | | | 1 | 2 | | | 1 | | 5 |
| | 660 | 690 | 672 | 657 | 701 | 687 | 666 | 672 | 687 | 674 | 6766 |

$$\int \frac{\mathrm{d}x}{\log x} = 6797{,}394$$

2500000 . . . 2600000

| | 251 | 252 | 253 | 254 | 255 | 256 | 257 | 258 | 259 | 260 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | | 1 | 1 | | | | | | | 1 | 3 |
| 2 | 1 | 2 | | 1 | 1 | | | | | | 5 |
| 3 | 4 | | 6 | 6 | 2 | 2 | 1 | 2 | 8 | 4 | 35 |
| 4 | 7 | 18 | 9 | 7 | 8 | 6 | 10 | 7 | 8 | 8 | 88 |
| 5 | 10 | 11 | 7 | 15 | 15 | 23 | 16 | 16 | 10 | 13 | 136 |
| 6 | 22 | 14 | 20 | 21 | 20 | 15 | 18 | 19 | 20 | 25 | 194 |
| 7 | 24 | 17 | 12 | 20 | 22 | 22 | 23 | 15 | 13 | 12 | 180 |
| 8 | 18 | 15 | 20 | 9 | 16 | 18 | 16 | 20 | 19 | 19 | 170 |
| 9 | 8 | 10 | 13 | 7 | 9 | 7 | 6 | 12 | 6 | 10 | 88 |
| 10 | 2 | 5 | 8 | 10 | 5 | 4 | 7 | 4 | 9 | 4 | 58 |
| 11 | 1 | 5 | 3 | 2 | 2 | 1 | 2 | 3 | 5 | | 24 |
| 12 | 1 | 1 | 1 | 2 | | 2 | 1 | 2 | 1 | 2 | 13 |
| 13 | 2 | 1 | | | | | | | 1 | 2 | 6 |
| | 677 | 675 | 696 | 670 | 670 | 671 | 678 | 698 | 693 | 676 | 6804 |

$$\int \frac{\mathrm{d}x}{\log x} = 6778{,}960$$

2600000 . . . 2700000

| | 261 | 262 | 263 | 264 | 265 | 266 | 267 | 268 | 269 | 270 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 0 | | | | 1 | | | | | | | 1 |
| 1 | | | 2 | | | | | 2 | | | 4 |
| 2 | 1 | 2 | 1 | | 2 | | 1 | 2 | 1 | | 10 |
| 3 | 3 | 6 | 2 | 2 | 4 | | 3 | 3 | 3 | 2 | 28 |
| 4 | 9 | 6 | 6 | 12 | 3 | 7 | 7 | 8 | 6 | 7 | 71 |
| 5 | 11 | 15 | 14 | 11 | 13 | 17 | 22 | 16 | 18 | 21 | 158 |
| 6 | 26 | 17 | 14 | 18 | 23 | 19 | 20 | 24 | 19 | 15 | 195 |
| 7 | 23 | 11 | 27 | 20 | 21 | 21 | 21 | 14 | 22 | 21 | 201 |
| 8 | 14 | 23 | 16 | 13 | 15 | 16 | 12 | 8 | 12 | 13 | 142 |
| 9 | 9 | 10 | 13 | 16 | 10 | 10 | 5 | 10 | 5 | 8 | 96 |
| 10 | 3 | 6 | 5 | 4 | 2 | 8 | 2 | 5 | 8 | 10 | 53 |
| 11 | 1 | 1 | | 1 | 3 | 1 | 4 | 4 | 5 | 2 | 22 |
| 12 | | 3 | | 2 | 4 | 1 | 1 | 4 | 1 | 1 | 17 |
| 13 | | | | | | | 1 | | | | 1 |
| 14 | | | | | | | 1 | | | | 1 |
| | 653 | 681 | 672 | 680 | 689 | 695 | 660 | 665 | 681 | 686 | 6762 |

$$\int \frac{\mathrm{d}x}{\log x} = 6761{,}332$$

2700000 . . . 2800000

| | 271 | 272 | 273 | 274 | 275 | 276 | 277 | 278 | 279 | 280 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | | | | | | 1 | | | 1 | | 2 |
| 2 | 2 | | 2 | | | 2 | | | 1 | | 7 |
| 3 | 4 | 5 | 6 | 5 | 4 | 4 | 4 | 3 | 3 | 5 | 43 |
| 4 | 9 | 7 | 16 | 7 | 8 | 9 | 8 | 8 | 12 | 11 | 95 |
| 5 | 10 | 14 | 13 | 14 | 13 | 12 | 13 | 17 | 11 | 18 | 135 |
| 6 | 24 | 18 | 15 | 28 | 19 | 20 | 15 | 21 | 18 | 17 | 195 |
| 7 | 18 | 22 | 15 | 20 | 24 | 16 | 23 | 19 | 22 | 9 | 188 |
| 8 | 13 | 10 | 13 | 12 | 15 | 13 | 20 | 19 | 15 | 15 | 145 |
| 9 | 9 | 9 | 10 | 4 | 11 | 12 | 9 | 7 | 8 | 8 | 87 |
| 10 | 6 | 9 | 6 | 7 | 5 | 8 | 7 | 2 | 7 | 10 | 67 |
| 11 | 3 | 3 | 4 | 2 | | 2 | | 3 | 1 | 6 | 24 |
| 12 | 1 | 3 | | 1 | 1 | 1 | | | 1 | 1 | 9 |
| 13 | | | | | | | 1 | 1 | | | 2 |
| 14 | | | | | | | | | | | |
| 15 | | | | | | | | | | | |
| 16 | | | | | | | | | | | |
| 17 | 1 | | | | | | | | | | 1 |
| | 679 | 695 | 644 | 657 | 672 | 671 | 684 | 666 | 662 | 684 | 6714 |

$$\int \frac{\mathrm{d}x}{\log x} = 6744{,}430$$

2800000 . . . 2900000

| | 281 | 282 | 283 | 284 | 285 | 286 | 287 | 288 | 289 | 290 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | | | | | 1 | | 1 | | | | 2 |
| 2 | | 2 | 4 | | 5 | 2 | | 1 | 1 | | 15 |
| 3 | 2 | 4 | 4 | 3 | 4 | 3 | 1 | 3 | 2 | 4 | 30 |
| 4 | 9 | 7 | 6 | 9 | 7 | 8 | 10 | 11 | 10 | 8 | 85 |
| 5 | 14 | 7 | 14 | 14 | 7 | 16 | 19 | 16 | 18 | 15 | 140 |
| 6 | 18 | 17 | 20 | 13 | 23 | 18 | 16 | 18 | 15 | 21 | 179 |
| 7 | 24 | 22 | 21 | 20 | 20 | 27 | 22 | 23 | 21 | 22 | 222 |
| 8 | 13 | 18 | 9 | 20 | 13 | 12 | 9 | 12 | 12 | 14 | 132 |
| 9 | 10 | 17 | 12 | 12 | 8 | 6 | 14 | 9 | 10 | 11 | 109 |
| 10 | 7 | 4 | 6 | 3 | 5 | 7 | 5 | 5 | 8 | 3 | 53 |
| 11 | | 1 | 3 | 3 | 5 | | 1 | 2 | 1 | 2 | 18 |
| 12 | 2 | 1 | | | | 1 | 2 | | 2 | | 8 |
| 13 | | | 1 | 2 | 2 | | | | | | 5 |
| 14 | 1 | | | 1 | | | | | | | 2 |
| | 690 | 695 | 667 | 704 | 671 | 654 | 672 | 653 | 676 | 662 | 6744 |

$$\int \frac{\mathrm{d}x}{\log x} = 6728{,}220$$

2900000 . . . 3000000

| | 291 | 292 | 293 | 294 | 295 | 296 | 297 | 298 | 299 | 300 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | | 1 | | | | | | | 1 | | 2 |
| 2 | | 1 | 2 | 1 | 2 | 2 | | 1 | 1 | 3 | 13 |
| 3 | 3 | 3 | 5 | 2 | 8 | 6 | 4 | 4 | 5 | 4 | 44 |
| 4 | 7 | 7 | 6 | 6 | 7 | 9 | 6 | 6 | 6 | 4 | 64 |
| 5 | 20 | 11 | 14 | 18 | 12 | 15 | 17 | 19 | 16 | 11 | 153 |
| 6 | 17 | 21 | 22 | 18 | 18 | 16 | 11 | 26 | 21 | 17 | 187 |
| 7 | 19 | 30 | 18 | 22 | 22 | 25 | 27 | 13 | 15 | 23 | 214 |
| 8 | 14 | 11 | 12 | 12 | 17 | 11 | 13 | 11 | 14 | 19 | 134 |
| 9 | 10 | 9 | 12 | 13 | 9 | 6 | 12 | 12 | 9 | 11 | 103 |
| 10 | 6 | 4 | 5 | 6 | 3 | 6 | 8 | 7 | 9 | 4 | 58 |
| 11 | 2 | 1 | 3 | 1 | | 2 | 1 | | 1 | 4 | 15 |
| 12 | 2 | 1 | | 1 | 2 | 2 | | 1 | 2 | | 11 |
| 13 | | | 1 | | | | | | | | 1 |
| 14 | | | | | | | | | | | |
| 15 | | | | | | | 1 | | | | 1 |
| | 680 | 663 | 671 | 680 | 649 | 652 | 694 | 658 | 671 | 687 | 6705 |

$$\int \frac{\mathrm{d}x}{\log x} = 6712{,}64$$

1000000 . . . 2000000

| | 110 | 120 | 130 | 140 | 150 | 160 | 170 | 180 | 190 | 200 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 0 | | | | | | | 1 | | | | 1 |
| 1 | 1 | 1 | 2 | 1 | 5 | 2 | 2 | 1 | | 1 | 16 |
| 2 | 4 | 5 | 6 | 9 | 7 | 10 | 11 | 5 | 10 | 5 | 72 |
| 3 | 21 | 25 | 32 | 19 | 19 | 28 | 29 | 30 | 22 | 34 | 259 |
| 4 | 54 | 57 | 63 | 69 | 72 | 77 | 68 | 70 | 71 | 67 | 668 |
| 5 | 114 | 107 | 120 | 119 | 129 | 124 | 120 | 152 | 135 | 136 | 1256 |
| 6 | 171 | 170 | 160 | 173 | 183 | 199 | 159 | 174 | 175 | 182 | 1746 |
| 7 | 217 | 217 | 214 | 207 | 179 | 172 | 203 | 194 | 206 | 221 | 2030 |
| 8 | 164 | 160 | 168 | 161 | 183 | 149 | 174 | 147 | 161 | 148 | 1615 |
| 9 | 126 | 131 | 111 | 120 | 98 | 124 | 130 | 124 | 113 | 103 | 1180 |
| 10 | 71 | 77 | 73 | 70 | 73 | 63 | 63 | 61 | 74 | 62 | 687 |
| 11 | 39 | 32 | 35 | 33 | 34 | 29 | 26 | 26 | 23 | 30 | 307 |
| 12 | 12 | 11 | 9 | 15 | 16 | 17 | 9 | 10 | 10 | 5 | 114 |
| 13 | 6 | 5 | 5 | 3 | 2 | 6 | 3 | 5 | | 3 | 38 |
| 14 | | 2 | 1 | 1 | | | 1 | | | 3 | 8 |
| 15 | | | | | | | 1 | 1 | | | 2 |
| 16 | | | 1 | | | | | | | | 1 |
| | 7210 | 7194 | 7081 | 7098 | 7028 | 6971 | 7012 | 6931 | 6955 | 6902 | 70382 |

$$\int \frac{\mathrm{d}x}{\log x} = 70427{,}78$$

2000000 . . . 3000000

| | 210 | 220 | 230 | 240 | 250 | 260 | 270 | 280 | 290 | 300 | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 0 | | | | | | | 1 | | | | 1 |
| 1 | 3 | 2 | 2 | 4 | 1 | 3 | 4 | 2 | 2 | 2 | 25 |
| 2 | 10 | 9 | 9 | 11 | 9 | 5 | 10 | 7 | 15 | 13 | 98 |
| 3 | 32 | 27 | 29 | 32 | 37 | 35 | 28 | 43 | 30 | 44 | 337 |
| 4 | 69 | 69 | 73 | 86 | 78 | 88 | 71 | 95 | 85 | 64 | 778 |
| 5 | 119 | 146 | 138 | 136 | 147 | 136 | 158 | 135 | 140 | 153 | 1408 |
| 6 | 197 | 183 | 179 | 176 | 193 | 194 | 195 | 195 | 179 | 187 | 1878 |
| 7 | 204 | 201 | 205 | 194 | 189 | 180 | 201 | 188 | 222 | 214 | 1998 |
| 8 | 157 | 168 | 168 | 158 | 151 | 170 | 142 | 145 | 132 | 134 | 1525 |
| 9 | 115 | 109 | 113 | 112 | 102 | 88 | 96 | 87 | 109 | 103 | 1034 |
| 10 | 63 | 52 | 44 | 55 | 58 | 58 | 53 | 67 | 53 | 58 | 561 |
| 11 | 21 | 18 | 30 | 28 | 23 | 24 | 22 | 24 | 18 | 15 | 223 |
| 12 | 8 | 9 | 10 | 7 | 7 | 13 | 17 | 9 | 8 | 11 | 99 |
| 13 | 2 | 4 | | 1 | 5 | 6 | 1 | 2 | 5 | 1 | 27 |
| 14 | | 3 | | | | | 1 | | 2 | | 6 |
| 15 | | | | | | | | | | 1 | 1 |
| 16 | | | | | | | | | | | |
| 17 | | | | | | | | 1 | | | 1 |
| | 6874 | 6857 | 6849 | 6787 | 6766 | 6804 | 6762 | 6714 | 6744 | 6705 | 67862 |

The group of one hundred numbers at position 26379 contains no prime number

The group of one hundred numbers at position 27050 contains 17 prime numbers.

$$\int \frac{\mathrm{d}x}{\log x} = 67915{,}733$$

# GAUSS TO ENKE.

---

Highly esteemed friend!

— — The remarks on the frequency of prime numbers that you kindly sent me interested me in more than one way. They reminded me of my own work on this subject, whose beginnings lie far in the past, in 1792 or 1793, when I acquired LAMBERT's supplements to the logarithm tables. Even before I had begun the finer investigations of higher arithmetic, one of my first tasks was to turn my attention to the decreasing frequency of prime numbers. For this purpose I counted them in each group of a thousand and wrote the results on one of the attached blank leaves. I soon saw that, despite all its fluctuations, this frequency was on average approximately inversely proportional to the logarithm, so that the number of all primes below a given bound $n$ would be approximately expressed by the integral

$$\int \frac{\mathrm{d}n}{\log n}$$

with the natural logarithm understood. Later, when I learned of the list printed in VEGA's tables (of 1796), extending up to 400031, I extended my count, which confirmed that relationship. In 1811 the appearance of CHERNAC's cribrum gave me great pleasure, and (since I had no patience for a sustained count in sequence) I very often spent an otherwise free quarter of an hour counting a group of a thousand, here or there; eventually, however, I gave it up altogether without quite finishing the million. Only later did I make use of GOLDSCHMIDT's diligence, partly to fill the remaining gaps in the first million and partly to continue the count using BURCKHARDT's tables Thus the first three millions have been counted (for many years now) and compared with the value of the integral. Here I give only a short extract:

| Below | there are prime numbers | Integral $\int \frac{\mathrm{d}n}{\log n}$ | Deviation | Your formula | Deviation |
|---|---|---|---|---|---|
| 500000 | 41556 | 41606,4 | +50,4 | 41596,9 | +40,9 |
| 1000000 | 78501 | 79627,5 | +126,5 | 78672,7 | +171,7 |
| 1500000 | 114112 | 114263,1 | +151,1 | 114374,0 | +264,0 |
| 2000000 | 148883 | 149054,8 | +171,8 | 149233,0 | +350,0 |
| 2500000 | 183016 | 183245,0 | +229,0 | 183495,1 | +479,1 |
| 3000000 | 216745 | 216970,6 | +225,6 | 217308,5 | +563,5 |

I did not know that LEGENDRE had also concerned himself with this subject. Your letter prompted me to look in his *Théorie des Nombres*, and in the second edition I found some pages on it that I must have overlooked earlier (or forgotten since). LEGENDRE uses the formula

$$\frac{n}{\log n - A}$$

where $A$ is supposed to be a constant, for which he takes 1,08366. A quick calculation then gives me the deviations in the above cases as

$$\begin{gathered} -23{,}3 \\ +42{,}2 \\ +68{,}1 \\ +92{,}8 \\ +159{,}1 \\ +167{,}6 \end{gathered}$$

These differences are even smaller than those with the integral, but they seem to grow faster as $n$ increases, so it is quite possible that, if continued much further, they would exceed the latter. To bring the count and the formula into agreement, one would have to set, respectively, in place of $A = 1{,}08366$

$$\begin{gathered} 1{,}09040 \\ 1{,}07682 \\ 1{,}07582 \\ 1{,}07529 \\ 1{,}07179 \\ 1{,}07297 \end{gathered}$$

It seems that, as $n$ grows, the (average) value of $A$ decreases, but I venture no conjecture as to whether the limit as $n$ grows to infinity will be 1 or a quantity different from 1. I cannot say that there are grounds for expecting a very simple limiting value; on the other hand, the excess of $A$ over 1 could quite well be a quantity of order $\frac{1}{\log n}$. I would be inclined to believe that the differential of the function in question must be simpler than the function itself. Whereas I have assumed $\frac{\mathrm{d}n}{\log n}$ for it, LEGENDRE's formula would assume a differential function that would be approximately $\frac{\mathrm{d}n}{\log n-(A-1)}$. Your formula, moreover, could for a very large $n$ be regarded as agreeing with

$$\frac{n}{\log n - \frac{1}{2k}}$$

where $k$ is the modulus of BRIGGS's logarithms, and hence with LEGENDRE's formula if

$$A = \tfrac{1}{2k} = 1{,}1513 \text{ is set.}$$

Finally, I should mention that I have noticed a couple of differences between your counts and mine.

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| Between | 59000 | and | 60000 | you have | 95 | I | 94 |
| | 101000 | | 102000 | | 94 | | 93 |

The first difference may be due to the prime number 59023 appearing twice in LAMBERT's supplements. The group of a thousand from 101000—102000 teems with errors in LAMBERT's supplements: in my copy I have marked 7 numbers that are not prime and inserted 2 that were missing. Could you not persuade the young DASE to count the primes in the following millions from the tables at the Academy, which, I fear, the public is not meant to possess? In that case I should mention that, in the 2nd and 3rd millions, the count was made at my direction according to a special scheme that I had already used myself for part of the first million. The counts for each 100000 occupy one (small) octavo page in 10 columns, each referring to one group of ten thousand; there is also a column before them (on the left) and another after them on the right. As an example, here is one vertical column and the two additional columns from the interval $1000000 \ldots 1100000$ — — —

For an explanation, take the 1st vertical column, for example. The group of ten thousand from 1000000 to 1010000 contains 100 groups of a hundred; among these there is 1 containing only one prime; none at all with 2 or 3; 2 with 4 primes each; 11 with 5 each, and so on. Together they give $752 = 1 \,.\, 1 + 4 \,.\, 2 + 5 \,.\, 11 + 6 \,.\, 14 + \,.\,.$ The last column contains the aggregates from the 10 individual columns. The numbers 14. 15. 16 in the first vertical column are superfluous here, since no groups of a hundred contain so many primes; but they come into use on the following sheets. Finally, the 10 pages are again combined into 1, and thus cover the entire 2nd million.

But it is time to break off. — — — With heartfelt wishes for your well-being

Always yours

Göttingen, 24 December 1849. C. F. GAUSS.

---

# TABLE
## OF THE NUMBER OF CLASSES
## OF BINARY QUADRATIC FORMS.

Hundred 1.
G. I. . . .(17). .(61)
1. 1. 2. 3.
4. 7
3 11. 19. 23.
27. 31. 43.
67
5 47. 79
7 71
9 59. 83
G. II. .(58). .(280)
1 5. 6. 8.
9. 10. 12.
13. 15. 16.
18. 22. 25.
28. 37. 58.
2 14. 17. 20.
32. 34. 36.
39. 46. 49.
52. 55. 63.
64. 73. 82.
97. 100
3 26. 29. 35.
38. 44. 50.
51. 53. 54.
61. 75. 76.
81. 87. 91.
92. 99
4 41. 62. 68.
94. 95. 98
5 74. 86
6 89
G. IV. .(25). .(136)
1 21. 24. 30.
33. 40. 42.
45. 48. 57.
60. 70. 72.
78. 85. 88.
93
2 56. 65. 66.
69. 77. 80.
84. 90. 96

Total 233. . .477
Irreg. 0 Impr. 74

Hundred 2.
G. I. . . . .(11). .(101)
3 163
5 103. 127
7 151
9 107. 139. 199
11 167
13 191
15 131. 179
G. II. . . . .(46). .(406)
2 142. 148. 193
3 106. 108. 109.
115. 118. 121.
123. 124. 135.
147. 157. 162.
169. 172. 175.
187
4 111. 113. 128.
137. 158. 178.
183. 196
5 119. 122. 125.
143. 159. 166.
181. 188. 197
6 116. 155. 171
7 101. 134. 149.
173
8 146. 164
10 194
G. IV. . . .(39). .(356)
1 102. 112. 130.
133. 177. 190
2 114. 117. 126.
132. 136. 138.
141. 144. 145.
150. 153. 154.
156. 160. 180.
184. 192. 198
3 104. 110. 129.
140. 152. 170.
174. 176. 182.
186. 189. 195.
200
4 161. 185
G. VIII. . .(4). . .(32)
1 105. 120. 165.
168

Total 291. . .895
Irreg. 0

Hundred 3.
G. I. . . . . (9). . . . . (109)
7 223
9 211. 243(*3*). 283
11 271
13 263
15 227. 239
21 251
G. II. . . . . (42). . . . . (482)
3 202. 207. 214. 235. 247. 262. 267. 268. 277. 298
4 226. 256. 289. 292. 295
5 218. 229. 242. 250
6 203. 212. 219. 233. 241. 244. 259. 274. 275. 279. 291
7 215. 278. 284. 287
8 254. 257
9 236. 293
10 206. 281
11 269
12 299
G. IV. . . . . (43). . . . . (512)
1 232. 253
2 205. 208. 213. 217. 220. 225. 228. 238. 252. 258. 265. 282. 288
3 201. 204. 216. 222. 231. 234. 237. 245. 246. 249. 255. 261. 270. 286. 294. 297. 300
4 221. 224. 248. 260. 272. 276
5 209. 230. 266. 290. 296
G. VIII. . . . (6). . . . . (64)
1 210. 240. 273. 280.
2 264. 285

Total 313. . . . . 1167
Irreg. 1 Impr. 183.

Hundred 4.
G. I. . . . . (9). . . . . (113)
7 343
9 307(*3*). 331. 367. 379
15 347
17 383
19 311. 359
G. II. . . . . (40). . . . . (554)
3 358. 397
4 313. 337. 382. 388
5 303. 316. 317. 319. 346. 361. 373. 375. 394
6 302. 323. 324. 327. 334. 351. 355. 363. 387
7 338. 349. 391
8 353
9 332. 335. 339(*3*). 362
10 386. 398
11 326. 389
12 356. 371. 395
13 314
G. IV. . . . . (43). . . . . (608)
2 301. 310. 322. 328. 333. 340. 352. 372. 400
3 304. 309. 315. 318. 325. 342. 348. 364. 366. 368. 370. 378. 393. 396
4 305. 306. 308. 320. 350. 354. 369. 376. 377. 380. 384. 392. 399
5 321. 344. 365. 381.
6 329
7 341. 374
G. VIII. . . . (8). . . . . (88)
1 312. 330. 345. 357. 385
2 336. 360. 390

Total 325. . . . . 1363
Irreg. 2 Impr. 229

# NEGATIVE DETERMINANTS.

Hundred 5.
G. I. . . . .(10). .(174)
7 463. 487
9 499
15 439. 443
21 431. 467
25 479
27 419. 491
G. II. . . .(33). .(512)
3 403. 427
4 457. 466. 478
5 412. 415. 421. 422. 423
6 433. 436. 475. 484
7 447. 454
8 407. 409. 452. 471
9 411. 428. 451. 459(*). 486
10 401. 449. 482. 500
13 458
14 404
15 461
16 446
G. IV. . . .(49). .(760)
2 418. 438. 442. 445. 448. 498
3 405. 417. 424. 430. 432. 435. 450. 453. 460. 472. 473. 477. 483. 490. 492. 493. 496
4 402. 406. 410. 414. 441. 444. 468. 469. 481. 485. 495
5 413. 437. 455. 470. 474. 476. 488. 489
6 416. 425. 426. 434. 464. 497
7 494
G. VIII. . .(8). .(120)
1 408. 462
2 420. 429. 456. 465. 480
3 440

Total 336. . .1566
Irreg. 1

Hundred 6.
G. I. . . . . .(7). .(133)
9 547
15 523. 571
21 503. 587
25 599
27 563
G. II. . . .(40). .(724)
4 562. 577. 583
5 508. 538. 541
6 507. 526. 529. 543. 567
7 502. 511. 535
8 512. 514. 548. 559. 578
9 515. 519. 527. 531. 556. 557. 575. 586
11 551. 554. 591
12 539. 542. 579. 593
14 596
15 509. 524. 566
16 521. 569
G. IV. . . .(41). .(672)
2 505. 522. 532. 553. 568. 592. 598
3 513. 517. 533. 537. 540. 550. 555. 565. 588. 595. 597
4 501. 518. 544. 558. 564. 573. 574. 576(*2*). 580(*2*). 582. 589
5 534. 572. 590
6 516. 549. 594
7 506. 530. 536. 581
8. . . 545. 584
G. VIII. .(12). .(200)
1 520
2 504. 510. 525. 528. 552. 561. 570. 585. 600
3 546. 560

Total 347. . .1729
Irreg. 2

Hundred 7.
G. I. . . . .(8). . .(138)
9 643
13 607. 631
15 619. 683. 691
25 647
33 659
G. II. . . .(37). .(718)
3 652
5 613. 625. 694
6 603. 617. 622. 628. 655. 667. 673. 676. 687
7 604. 634. 639. 653
9 661. 675(*3*). 679
10 601
11 623. 662. 668
12 674. 695
13 698
14 641. 686. 692
16 611. 635. 671. 677. 699
17 614
18 626
G. IV. . . .(43). .(812)
2 658. 697
3 606. 610. 618. 627. 637. 648. 669. 670. 682. 685. 688. 700
4 612. 632. 640. 642. 646. 657. 663
5 615. 633. 636. 638. 649. 664. 666. 678. 681
6 602. 605. 608. 620. 621. 650. 651. 684
7 654
8 644. 656
9 629
10 689
G. VIII. .(12). .(216)
2 609. 616. 624. 630. 645. 660. 672. 690. 693.
3 665. 680. 696.

Total 350. .1884
Irreg. 1

Hundred 8.
G. I. . . . . .(6). . .(110)
13 727
15 739. 751. 787
21 743
31 719
G. II. . . . . .(39). . .(860)
4 772
5 709. 757
6 718. 723. 763. 775
7 703. 733. 778
8 799
9 707. 722. 729. 747. 771. 783. 796
10 711. 724. 769. 788
11 758. 767
12 706. 766
13 746. 764. 773
15 716. 779. 797
16 791
17 701
18 731. 755(*3*)
20 734. 761
21 794
G. IV. . . . . .(42). . .(792)
2 708. 742. 793
3 702. 715. 730. 748. 753. 762. 795
4 712. 717. 721. 732. 735. 736. 738. 745. 768. 784. 785. 786. 790
5 726. 737. 750. 752. 754. 774. 781
6 704. 713. 725. 756. 759. 782. 800
8 710. 740. 749. 789
10 776
G. VIII. . . . .(13). . .(264)
1 760
2 720. 765. 777. 792. 798
3 705. 714. 728. 741. 744. 780
4 770

Total 356. . .2026
Irreg. 1

Hundred 9.
G. I. . . . . (8). . . . (164)
9 823. 883
21 811. 827. 859. 863.
29 887
33 839
G. II. . . . . (34). . . . (750)
4 862
5 847. 853. 877
6 802. 898
7 807. 838. 841. 892.
8 895
9 835. 843. 844. 867. 886. 891(*3*)
10 878
11 829. 871. 879
13 842
14 818. 831
15 803. 815. 821. 851. 875
16 809. 857
20 881
21 899
22 866
G. IV. . . . (47). . . (1024)
3 808. 813. 814. 817. 826. 828. 837. 856
4 820(*2*). 832. 834. 850. 852. 855. 865. 868. 873. 882. 889. 900(*2*)
5 822. 830. 872. 874
6 801. 804. 810. 812. 819. 833. 848. 864. 876. 890. 894
7 806. 845. 849. 860. 893
8 846. 869. 884(*2*). 896
10 824. 836
11 854
G. VIII. . . (10). . . . (200)
2 805. 858. 870. 880. 897
3 816. 825. 861. 885. 888
G. XVI. . . . (1). . . . (16)
1 840

Total 360. . . . . 2154
Irreg. 4

Hundred 10.
G. I. . . . . (8). . . . (174)
9 907
11 967
15 947
17 991
19 919
27 983
31 911
45 971
G. II. . . . . (33). . . (810)
5 982
6 955
7 997
8 943. 958. 961
9 922. 931. 963. 972
10 916. 927. 937. 977
12 932. 939. 964. 979. 995. 999
13 934. 951. 998
15 908. 923. 956
16 953
18 914. 929. 959. 974(*3*)
20 926
23 941
G. IV. . . . (45). . . . (976)
2 928
3 913. 918. 925. 933. 940. 942. 949. 970. 973. 988
4 903. 904. 993. 938. 946. 975. 994
5 917. 921. 968. 1000
6 901. 905. 915. 948. 954. 976. 978. 980. 981. 985. 987. 996
7 902. 906. 909. 935. 962
8 992
9 944. 950. 989
11 965. 986
G. VIII. . . (14). . . . (312)
2 910. 912. 952. 957. 960
3 924. 930. 936. 945. 966. 969. 984. 990
5 920

Total 366. . . . . 2272
Irreg. 1

Hundred 11.
G. I. . . . . (7). . . . . (191)
9 1087
15 1051
19 1063
23 1039
35 1031
39 1019
51 1091
G. II. . . . . (35). . . . (880)
5 1093
6 1003. 1027. 1033. 1042
8 1024. 1047
9 1018. 1059. 1075(*3*). 1083. 1099
10 1006. 1009
11 1021. 1082. 1084
12 1043. 1058
13 1013. 1052. 1061. 1094.
15 1007. 1069
16 1028
17 1079
18 1011. 1055. 1067. 1097
21 1004. 1046
22 1049. 1076
G. IV. . . . . (44). . . . . (984)
2 1012
3 1030. 1038. 1048. 1068 1072. 1090
4 1002. 1015. 1017. 1023. 1054. 1057. 1060. 1078. 1081
5 1037. 1066. 1071. 1098
6 1026. 1035. 1036. 1044. 1053. 1062. 1073. 1077. 1089. 1096. 1100
7 1010. 1014. 1029. 1086. 1095
8 1016. 1022. 1025. 1074. 1088(*2*)
9 1041. 1070
11. . . 1034
G. VIII. . . . . (14). . . . (344)
2 1005. 1008. 1032. 1045. 1065. 1092
3 1020. 1050. 1080
4 1040. 1056. 1085
5 1001. 1064

Total 365. . . . . 2399
Irreg. 2

Hundred 12.
G. I. . . . . (6). . . . . (148)
15 1123
21 1163. 1171
23 1103
27 1187(*3*)
41 1151
G. II. . . . . (36). . . . (924)
6 1108. 1138. 1198
7 1117. 1183
8 1129. 1153. 1156. 1159
9 1107(*3*). 1132. 1135. 1142. 1147
10 1143
11 1111. 1114. 1126. 1167
12 1127. 1186. 1191. 1195
15 1115. 1174. 1175. 1179
16 1119
18 1172. 1193
19 1199
20 1124
23 1181
24 1139
25 1109
28 1154
G. IV. . . . . (40). . . . (1064)
3 1162. 1177. 1192
4 1149. 1150. 1152. 1168. 1178. 1180
5 1102. 1125. 1137. 1165. 1182. 1189
6 1131. 1134. 1141. 1145. 1158. 1164. 1188
7 1101. 1112. 1133. 1136. 1148. 1157. 1194
8 1146
9 1116. 1118. 1161. 1166
10 1184
11 1121. 1130
12 1106. 1169. 1196
G. VIII. . . . . (18). . . . . (408)
2 1105. 1110. 1113. 1120. 1122. 1128. 1170. 1185. 1197
3 1144. 1155. 1173. 1176 1200
4 1104. 1140
5 1160. 1190

Total 382. . . . . 2544
Irreg. 2

Hundred 13.
G. I. . . . . .(6). . . . . .(190)
23 1279
27 1231. 1291
33 1283
35 1223
45 1259(*3*)
G. II. . . . . .(38). . . . . .(986)
5 1213
6 1227. 1243. 1255. 1282. 1297
7 1237
8 1201. 1252
9 1203. 1207. 1215. 1219. 1228(*3*). 1267(*3*)
10 1261. 1263. 1268.
12 1202. 1234. 1299
13 1247
14 1294
15 1250
16 1217. 1249
17 1277
18 1251. 1262. 1289
19 1229. 1244
20 1214. 1271
21 1211. 1226. 1238
29 1286
G. IV. . . . . .(40). . . . .(1008)
3 1222. 1258. 1285
4 1204. 1225. 1233. 1246. 1278
5 1210. 1212. 1257. 1264. 1270. 1273. 1276. 1287
6 1208. 1216. 1236. 1242. 1269. 1275. 1292. 1293. 1296. 1300
7 1206
8 1220(*2*). 1239. 1241. 1253. 1266. 1280. 1298
9 1235. 1274. 1295
10 1205. 1284
13 1256
VIII. . . . .(16). . . . . .(416)
2 1240. 1248. 1288. 1290
3 1218. 1230. 1254. 1260. 1272. 1281
4 1221. 1224. 1232. 1245
5 1209. 1265

Total 370. . . . . .2600
Irreg. 4

Hundred 14.
G. I. . . . . .(7). . . . . .(191)
11 1303
15 1327
27 1367. 1399
33 1307. 1331
45 1319
G. II. . . . . .(32). . . . . .(846)
5 1318
6 1387
7 1372
8 1348
9 1306. 1315(*3*). 1323(*3*). 1324. 1347. 1363. 1366. 1369. 1373. 1383
10 1375
11 1354
12 1321. 1339. 1351
13 1381
14 1346. 1359
15 1388
17 1343
18 1355. 1371
19 1382
21 1322
22 1391
24 1379
25 1301
30 1361
G. IV. . . . . .(46). . . . . .(1340)
4 1312(*2*). 1332(*2*). 1345. 1357. 1393
5 1317. 1333. 1338. 1342. 1378. 1384. 1390. 1398
6 1308. 1313. 1336. 1337. 1350. 1358. 1362. 1377. 1395. 1397
7 1311. 1335. 1341. 1352. 1374. 1389. 1396
8 1314. 1334
9 1310. 1325. 1328. 1329. 1340. 1356(*3*)
10 1376
11 1304. 1370
12 1316. 1385. 1394
14 1349. 1364
G. VIII. . . . . .(13). . . . . .(328)
2 1302. 1353. 1360. 1380
3 1309. 1330. 1368. 1392
4 1305. 1344. 1386. 1400
5 1326
G. XVI. . . . . .(2). . . . . .(32)
1 1320. 1365

Total 391. . . . . .2737
Irreg. 5 Progr. 3192

Hundred 15.
G. I. . . . . .(10). . . . . .(308)
9 1423
21 1483
23 1447. 1471
33 1459
37 1487
39 1439. 1451. 1499
45 1427
G. II. . . . . .(26). . . . . .(746)
6 1411. 1467
7 1402. 1453
8 1438
9 1458. 1468
10 1444. 1486. 1489. 1492.
11 1429. 1493
15 1431. 1478
16 1412
17 1415. 1418
18 1409. 1433. 1475
19 1436
21 1403
26 1481
29 1466
30 1454
G. IV. . . . . .(49). . . . . .(1348)
3 1432. 1435. 1450
4 1408. 1417. 1422. 1462. 1465. 1474. 1477. 1498
5 1495. 1497. 1500
6 1404. 1405. 1407. 1413. 1420. 1437. 1442. 1443. 1452. 1457. 1472
7 1401. 1414. 1441. 1455. 1461. 1473. 1479
8 1426. 1434. 1446. 1463. 1476
9 1419. 1445. 1448. 1490. 1491
10 1460. 1494
11 1406
12 1421. 1424. 1484
14 1469
G. VIII. . . . .(15). . . . . .(424)
2 1428. 1488
3 1425. 1456. 1464. 1480. 1482. 1485
4 1410. 1416. 1430. 1440(*2*). 1449. 1470
7 1496

Total 378. . . . . .2826
Irreg. 1 Propr. 3282

Hundred 16.
G. I. . . . . .(9). . . . . .(299)
15 1567
19 1543
21 1523
27 1579
33 1531. 1583
49 1511
51 1559. 1571
G. II. . . . . .(24). . . . . .(656)
6 1507. 1555. 1588
7 1527. 1597
9 1516. 1519. 1549. 1563
10 1522
11 1503. 1591
12 1502. 1587
17 1532. 1546. 1594
18 1539(*3*)
19 1535
20 1553
22 1538. 1556
25 1514
27 1574
G. IV. . . . . .(53). . . . . .(1564)
3 1558. 1593
4 1510. 1513(*2*). 1528. 1537. 1552. 1578. 1582(*2*). 1600(*2*)
5 1534, 1542. 1570. 1573. 1576
6 1501. 1506. 1521. 1525. 1548. 1572. 1575. 1585
7 1557. 1562. 1564. 1565. 1569. 1577
8 1504. 1508. 1536. 1551. 1561. 1568(*2*). 1598
9 1515. 1541. 1547. 1566. 1599
10 1509. 1524. 1544. 1592
11 1586
12 1517. 1526. 1550. 1580. 1595
13 1529. 1589
G. VIII. . . . .(13). . . . . .(360)
2 1540
3 1512. 1518. 1530. 1533. 1545. 1554. 1584
4 1520. 1590(*2*). 1596
5 1505. 1581
G. XVI. . . . . .(1). . . . . .(32)
2 1560

Total 389. . . . . .2911
Irreg. 6 Propr. 3416

Hundred 17.
G. I. . . . . .(6). . . . . .(182)
17 1663
21 1627
27 1607
33 1699
39 1667
45 1619
G. II. . . . . .(37). . . .(1116)
6 1618
7 1642
8 1657
9 1603. 1621. 1675(*). 1683. 1687
10 1678. 1681. 1684
11 1639. 1654. 1693
12 1647. 1651
13 1669
14 1609. 1623. 1697
15 1611. 1622. 1643. 1682
16 1636
19 1637. 1671
20 1604
21 1613. 1658
22 1631. 1646. 1655
26 1679
27 1676. 1691
28 1601
G. IV. . . . . .(41). . . . . .(1312)
3 1612
4 1633. 1660. 1698
5 1626. 1648. 1660. 1662. 1688. 1692. 1695
6 1615. 1620. 1635. 1659. 1666. 1668. 1690. 1696
7 1606. 1614. 1630. 1670
8 1602. 1628. 1673
9 1644. 1674. 1689
10 1625. 1629. 1652
11 1641. 1686
12 1649. 1661. 1664. 1694
13 1685
14 1616
16 1634
G. VIII. . . .(15). . . . . .(368)
2 1605. 1632. 1645. 1653 1672. 1677
3 1617. 1638
4 1608. 1610. 1624. 1640. 1650. 1656. 1665
G. XVI. . . . . .(1). . . . . .(32)
2 1680

Total 380. . . . . .3010
Irreg. 1 Impr. 513

Hundred 18.
G. I. . . . . .(5). . . . . .(95)
15 1723. 1747
17 1783
21 1787
27 1759
G. II. . . . . .(35). . . . .(1182)
10 1714. 1753. 1774
12 1726. 1731. 1732. 1762. 1777. 1795
13 1719. 1735. 1741. 1789
14 1703. 1711. 1775
15 1707. 1756. 1772. 1779
17 1733
18 1727. 1763(*3*). 1791
19 1754
21 1709. 1715. 1724
23 1718
24 1751
25 1766. 1799
26 1721
29 1706
30 1739
G. IV. . . . .(41). . . . .(1260)
3 1708
4 1717. 1737. 1738. 1780. 1792
5 1702. 1750. 1758. 1761. 1765. 1773. 1786. 1798
6 1728. 1743. 1744. 1771. 1782. 1797
7 1757. 1788
8 1764(*2*). 1767
9 1701. 1712. 1713. 1730. 1734. 1755(*). 1793
10 1745. 1746. 1748. 1778. 1796
11 1742
13 1790
16 1769. 1784
17 1781
G. VIII. . . . .(18). . . . .(536)
2 1705. 1710. 1752. 1768
3 1720. 1722. 1729. 1740. 1800
4 1716. 1725. 1776. 1794
5 1749. 1770
6 1704. 1736. 1760
G. XVI. . . . .(1). . . . . .(32)
2 1785

Total 399. . . . . .3105
Irreg. 3 Impr. 525

Hundred 19.
G. I. . . . . .(7). . . . . .(263)
15 1867
19 1831
27 1879
43 1847
45 1823. 1871
69 1811
G. II. . . . . .(31). . . . . .(994)
6 1807. 1873
7 1852
8 1822. 1828
9 1843. 1863. 1882
10 1858
11 1849
12 1803
14 1801. 1838
15 1819. 1835. 1875. 1891. 1894
17 1877
18 1899
19 1861
20 1839
21 1851. 1859. 1868. 1883
23 1814
24 1895
28 1874
30 1844
36 1889
G. IV. . . . . .(45). . . . . .(1416)
4 1813. 1842. 1864. 1897
5 1810. 1857. 1887. 1893
6 1812. 1815. 1818. 1825. 1827. 1837. 1878. 1888. 1892. 1900
7 1816. 1846. 1855. 1898
8 1802. 1808. 1866. 1876. 1884
9 1804. 1809. 1821. 1834. 1836. 1853. 1862
10 1805. 1817. 1829. 1841. 1850. 1854
12 1856. 1865
13 1832
14 1826
16 1886(*2*)
G. VIII. . . . .(16). . . . . .(496)
2 1870. 1885
3 1830. 1833. 1840. 1890
4 1824. 1845. 1860. 1872
5 1806. 1820. 1869. 1880. 1881. 1896
G. XVI. . . . .(1). . . . . .(16)
1 1848

Total 393. . . . . .3185
Irreg. 1 Impr. 513

Hundred 20.
G. I. . . . . .(6). . . . . .(252)
21 1987
27 1999
33 1951.
39 1907
63 1931(*3*)
69 1979
G. II. . . . . .(33). . . . . .(1090)
7 1948
8 1983
9 1915. 1927. 1933. 1963. 1996
10 1906. 1975
11 1903. 1942
12 1939. 1982. 1993
14 1954
15 1923
16 ·1922. 1943
18 1913. 1966. 1967. 1971(*3*)
21 1901. 1959. 1973. 1997
22 1919
25 1916
26 1934
27 1964. 1994
28 1991
35 1949
G. IV. . . . . .(43). . . . . .(1356)
3 1978
4 1912. 1918(*2*). 1945. 1957
5 1930. 1962. 1969. 1981
6 1908. 1917. 1926. 1936. 1941. 1947. 1972. 1984. 1990
7 1909. 1929. 1935
8 1911. 1924. 1940. 1958. 1961
9 1902. 1944. 1955. 1977. 1998
10 1921. 1928. 1952. 1956. 1985. 2000
12 1982. 1986. 1988
13 1970
14 1910
17 1946
G. VIII. . . . .(18). . . . . .(584)
2 1992
3 1905. 1932. 1950. 1960. 1968. 1995
4 1920. 1938. 1953. 1974. 1980. 1989
5 1904. 1965,
6 1914. 1925
7 1976

Total 388. . . . . .3282
Irreg. 3 Impr. 556

Hundred 21.
G. I. . . . . .(8). . . . . .(284)
21 2011. 2083
27 2003
33 2027
35 2087
45 2039. 2063
57 2099
G. II. . . . . .(30). . . . . .(1054)
6 2017. 2062
8 2095
9 2023. 2038. 2047. 2053
12 2059. 2098
14 2007. 2018
15 2043. 2071. 2092
16 2048
17 2026. 2029
19 2031. 2069
20 2078
21 2012
22 2089
24 2019
25 2042
27 2051. 2075(*3*)
28 2066
30 2036. 2081
32 2084
G. IV. . . . . .(42). . . . . .(1376).
4 2020. 2077
5 2032. 2073. 2074
6 2022. 2025. 2028. 2035. 2050. 2052. 2067. 2068. 2082. 2086. 2096
7 2008. 2033. 2044. 2055. 2058. 2094
8 2004. 2005. 2034. 2041. 2056
9 2014. 2049. 2060. 2076. 2079. 2091
10 2057. 2061
12 2006. 2009. 2045
13 2015
15 2096
17 2021
18 2054
G. VIII. . . . . .(19). . . . . .(632)
2 2002. 2013. 2080. 2088
3 2037. 2065
4 2010. 2016. 2046. 2072. 2100.
5 2030. 2070. 2085. 2093
6 2001. 2064. 2090
7 2024
G. XVI. . . . . .(1). . . . . .(32)
2 2040

Total 404. . . . . .3378
Irreg. 1 Impr. 560

Hundred 2
G. I. . .(5). .(149)
13 2143
21 2179
27 2187(*3*)
39 2131
49 2111
G. II. . .(33).(1174)
8 2113. 2137
9 2122. 2167. 2188
10 2164
11 2182
12 2107. 2116
13 2102. 2197
14 2127
15 2151. 2191
16 2153. 2194
17 2103. 2119
18 2155. 2161. 2199
21 2123. 2138. 2147. 2171. 2183. 2186
24 2195
28 2129
30 2126. 2159
32 2174
39 2141
G. IV.(46).(1592)
5 2101. 2118. 2125. 2152. 2158. 2173. 2178
6 2104. 2115. 2132. 2139. 2148. 2157. 2163. 2172.
7 2110. 2140. 2146. 2149. 2165
8 2134. 2176(*2*) 2192
9 2106. 2108. 2117. 2124. 2133(*3*). 2175. 2181. 2198
10 2150. 2154. 2166. 2177
11 2135
12 2105. 2156. 2168. 2169. 2196
14 2114. 2144. 2162
15 2189
16 2180
G. VIII.(14).(424)
2 2128. 2170
3 2160. 2185. 2190. 2193. 2200
4 2112(*2*). 2130. 2142
5 2109. 2121. 2136
7 2120
G. XVI.(2). . .(80)
2 2145
3 2184

Total 399. . .3419
Irreg. 4 Impr. 585

Hundred 23.
G. I. . . .(7). .(217)
15 2203
21 2251
29 2287
33 2267
35 2239
39 2207
45 .2243(*3*)
G. II. .(29).(1084)
7 2293
9 2221. 2227. 2283
10 2281
11 2215. 2263
12 2209
13 2218
14 2258
15 2237. 2269. 2284. 2299
16 2206
17 2234
18 2228
20 2297
21 2213. 2259
22 2271
24 2273
27 2252. 2291
28 2279
29 2231
32 2276
36 2219(*3*)
39 2246
G. IV.(46).(1612)
4 2212. 2242. 2248. 2272
5 2202. 2230. 2247. 2290
6 2223. 2257. 2260. 2268. 2275. 2278
7 2217. 2238. 2270
8 2236. 2245. 2254(*2*). 2286. 2292. 2298
9 2214. 2221. 2235(*3*). 2241. 2253. 2266. 2295. 2300.
10 2249. 2250. 2255. 2282
11 2204. 2216
12 2211. 2225. 2229
13 2222
14 2274
15 2264. 2285
16 2201
19 2294
G. VIII.(17).(584)
2 2233. 2277
3 2205. 2220. 2262
4 2208. 2232. 2244. 2256. 2265. 2289. 2296

5 2226. 2288
6 2240
7 2210
9 2261
G. XVI.(1)..(32)
2 2280

Total 401...3529
Irreg. 4 Impr. 571

Hundred 24
G. I....(7)..(291)
15 2347
29 2311. 2383
39 2371
57 2339
59 2399
63 2351
G. II..(32).(1106)
7 2335
8 2302. 2308. 2377
11 2326. 2374
12 2307. 2323. 2395
13 2362. 2367
14 2359
15 2303. 2319. 2341. 2363
16 2386
17 2333. 2389. 2391
19 2381
20 2375
21 2342. 2348. 2357
24 2327. 2372
25 2396
27 2315(*3*)
30 2393
32 2306
33 2309
G. IV..(40).(1520)
4 2332. 2353. 2368
5 2398
6 2305. 2317. 2322. 2338. 2358. 2388
7 2314. 2382
8 2304. 2312. 2313. 2329(*2*). 2343. 2350. 2356
9 2318. 2344. 2349. 2355. 2361. 2387
11 2334. 2364.
12 2316. 2331. 2376. 2379(*2*). 2390
14 2324. 2354. 2384
15 2321. 2330. 2378
16 2336
18 2369

G. VIII.(20).(648)
2 2392
3 2325. 2346. 2352. 2370. 2373. 2380
4 2320. 2328. 2337. 2340. 2365. 2385(*2*). 2397. 2400(*2*).
5 2360. 2394.
6 2301. 2376
7 2345
G. XVI. .(1). .(32)
2 2310

Total 407. . .3597
Irreg. 5 Impr. 611

Hundred 25.
G. I. . .(5). .(217)
21 2467
33 2423
37 2447
57 2459
69 2411
G. II. .(35).(1250)
9 2403(*3*). 2437(*3*).
2443. 2458
10 2407. 2452. 2473. 2487. 2500
12 2419. 2468. 2479
13 2428
14 2401. 2446. 2455
15 2476
16 2434. 2462
17 2463
18 2417. 2491
19 2477
20 2402. 2404. 2498
21 2427
22 2439
27 2426
28 2495
30 2483
31 2471
33 2435
38 2441
39 2474
G. IV.(38).(1472)
4 2410
5 2422. 2433. 2494
6 2412. 2413. 2425. 2451. 2475.
2482. 2488. 2493
7 2416. 2431. 2438. 2497
8 2454
9 2430(*3*). 2461
10 2449
11 2421. 2489. 2492
12 2420(*2*). 2432. 2450.
2466. 2484. 2499
13 2453. 2469. 2481
14 2429. 2486
15 2406. 2444
17 2456
18 2414
G. VIII.(22).(720)
3 2418. 2424. 2440. 2457. 2472.
2485
4 2436. 2442. 2445. 2448(*2*).
2464. 2465. 2470. 2478. 2490.
2496
5 2405. 2409. 2415. 2460
6 2408. 2480

Total 403. .3659
Irreg. 5 Impr. 595

Hundred 26.
G. I. . .(7). .(301)
21 2503
33 2539
35 2543
41 2551
51 2531
57 .2591
63 2579
G. II. .(29).(1028)
8 2578
9 2515. 2557. 2563(*3*).
2566. 2572
10 2527
12 2587. 2593
13 2524
15 2523. 2575. 2518. 2599
16 2521
18 2511. 2547
19 2554
20 2559
21 2507. 2571
22 2567. 2594
25 2582. 2588
28 2564
32 2519. 2558
35 2549
G. IV.(45).(1752)
4 2533. 2542(*2*)
5 2577
6 2512. 2517. 2538. 2545. 2548.
2592. 2598
7 2502. 2509. 2535. 2536. 2556.
2583
8 2506. 2513. 2528. 2560. 2569.
2589
9 2522. 2555. 2581. 2595
10 2514. 2529. 2532. 2596
11 2570. 2573
12 2597
13 2510
14 2501. 2525. 2586
15 2526. 2534. 2537. 2540
16 2546. 2561
18 2504. 2516
G. VIII.(18).(648)
3 2508. 2530. 2550. 2553. 2562.
2590
4 2568. 2580. 2584
5 2505. 2541. 2544. 2552 2585

6 2565. 2574. 2600
8 2576
G. XVI.(1)..(32)
2 2520

Total 405..3761
Irreg. 2 Impr. 641

Hundred 27.
G. I. . .(7). . .(231)
15 2647. 2683
23 2671
39 2659
43 2663
45 2699(*3*)
51 2687
G. II. .(29).(1196)
10 2638
11 2623. 2662. 2677
12 2611. 2689. 2692
14 2612
15 2602. 2643
16 2617. 2633. 2657
18 2619(*3*). 2627. 2644
21 2693
23 2614. 2615
24 2631. 2654
26 2642
27 2675(*3*)
30 2603. 2606
31 2621
33 2636
39 2651
42 2609
G. IV.(46).(1776)
3 2608
4 2605
5 2641
6 2620. 2629. 2650. 2653. 2658.
2667. 2668. 2676. 2682. 2695.
2698
7 2607
8 2601. 2628(*2*). 2655.
2674. 2686
9 2626. 2634. 2635. 2637.
2646(*3*). 2673. 2700(*3*)
10 2656. 2678
11 2645. 2648. 2649. 2672
12 2681. 2691. 2696
13 2679
15 2630. 2684. 2690. 2694
16 2624(*3*). 2639. 2669
17 2666
18 2681
G. VIII.(17).(584)
2 2632
3 2613. 2622. 2680. 2685. 2697
4 2652. 2665. 2688
5 2610. 2618. 2625. 2664. 2670.
6 2604. 2660
7 2616

G. XVI. .(1). .(32)
2 2640

Total 401. . .3819
Irreg. 7 Impr. 625

Hundred 28.
G. I. . .(6). .(208)
21 2707 2767
33 2731
39 2791
41 2719
53 2711
G. II.(29).(1190)
9 2787. 2797
10 2722. 2743
12 2713
13 2762
14 2734. 2753
15 2727. 2732. 2764
18 2723. 2763(*3*). 2783
19 2738. 2746. 2799
20 2777
21 2749. 2779
22 2798
27 2747. 2759
29 2741
30 2708.
31 2735
39 2771
40 2729
41 2789
G. IV.(47).(1864)
4 2773. 2788
5 2776
6 2704. 2710. 2715. 2718. 2725. 2740. 2748. 2752. 2755. 2758. 2794
7 2703. 2761. 2766. 2782
8 2733. 2742. 2775. 2785
9 2716. 2770. 2778. 2781. 2795
10 2724. 2751
11 2757
12 2701. 2702(*2*). 2739. 2754 2780
13 2721. 2792
14 2744. 2768. 2774
15 2750. 2796
17 2726
18 2705. 2786
20 2714. 2756(*2*)
G. VIII.(16).(568)
2 2737
3 2728. 2800

4 2706. 2709. 2717. 2745. 2772.
2790. 2793
5 2712. 2769
6 2720. 2736. 2784
7 2765
G. XVI.(2)..(64)
2 2730. 2760

Total 412. .3894
Irreg. 3 Impr. 644

Hundred 29.
G. I. .(6). .(250)
25 2887
27 2803
33 2851
45 2843
57 2879
63 2819
G. II. .(32).(1298)
8 2878
10 2818. 2836. 2857
11 2815. 2863
12 2827
13 2809. 2823. 2839. 2854
15 2875. 2883. 2899
16 2833
18 2867. 2897
19 2858
20 2866
23 2837
24 2811
26 2807
27 2859. 2891(*3*)
30 2801. 2855
31 2876
33 2861
34 2804. 2831. 2894
36 2846
G. IV.(43).(1680)
4 2842. 2893
5 2830. 2853
6 2802. 2872. 2892
7 2812. 2881
8 2817. 2845. 2847: 2848(*2*).
2868. 2884
9 2806. 2828. 2835(*3*).
2862. 2887. 2888. 2890. 2895
10 2810. 2844. 2869. 2871. 2874.
2896
11 2841
12 2816. 2822. 2824. 2825. 2852.
2873. 2900.
13 2826.
15 2813. 2864. 2889
16 2882
21 2834
G. VIII.(17).(648)
2 2832
3 2808. 2860
4 2821. 2829. 2850(*2*).
2865. 2880. 2898
5 2814. 2838. 2877. 2886
6 2820
7 2870

8 2840. 2849
G. XVI.(2)..(96)
3 2805. 2856

Total 410..3972
Irreg. 4 Impr. 636

Hundred 30.
G. I..(6)..(322)
31 2927
33 2971
39 2963
59 2903
73 2999
87 2939
G. II..(33).(1266)
8 2962
9 2902. 2923. 2998
10 2983
11 2917. 2935
12 2947. 2953. 2995
13 2908
14 2932
15 2956. 2986
17 2918
18 2916. 2943. 2979
20 2942. 2959
21 2911. 2931. 2972
24 2974. 2991
26 2969
27 2951. 2957
30 2978. 2987
33 2906
35 2909
43 2966
G. IV.(37).(1588)
5 2965
6 2907. 2914. 2938. 2977. 2997
7 2913. 2930. 2941. 2973
8 2944. 2946. 2949. 2980(*2*)
9 2950. 2955. 2988. 2989
10 2919. 2929. 2948. 2975
11 2922. 2933. 2934. 2967
12 2993(*2*)
13 2901. 2984
16 2921. 2994. 2996.
18 2915. 2924
20 2936. 2954. 2981
G. VIII.(24).(888)
2 2968
3 2905. 2920. 2937. 2970. 2982. 2992
4 2928. 2940. 2952. 2958. 2985
5 2904. 2910. 2926. 2990. 3000

6 2912. 2925. 2961. 2964. 2976
8 2945. 2960

Total 412. .4064
Irreg. 2 Impr. 714

Hundred 43.
G. I. . .(7). .(425)
27 4243
45 4219
51 4231
63 4283
65 4271
69 4211
105 4259
G. II.(32).(1592)
12 4222. 4258. 4267. 4273
13 4207. 4282
14 4279
15 4204. 4227
17 4261
18 4201. 4291(*3*). 4297
19 4252
21 4203. 4253
22 4223
24 4239. 4287. 4295
26 4217. 4244
27 4262. 4299
30 4276
31 4247
39 4229
40 4286
42 4274
54 4226. 4241
56 4289
G. IV.(38).(1780)
6 4225. 4237. 4288
7 4210. 4213
8 4228. 4249
9 4236. 4246. 4255. 4263. 4285. 4293. 4294. 4300(*3*)
11 4215. 4238. 4281. 4298
12 4212(*2*). 4232. 4251. 4275(*2*). 4292
13 4202. 4206. 4208. 4234
14 4205
15 4235. 4250. 4266
17 4269
18 4220. 4265. 4268
19 4254
20 4214
G. VIII(22). .(976)
3 4218. 4257. 4272
4 4216. 4240. 4260. 4270. 4278
5 4230. 4233. 4242. 4264. 4284
6 4245. 4248. 4277
7 4209
8 4221. 4224. 4296
9 4280
10 4256

G. XVI.(1)..(48)
3 4290

Total 415....4821
Irreg. 4

Hundred 51.
G. I...(8)..(546)
25 5023
45 5003
57 5059
63 5011
69 5087
83 5039
87 5051
117 5099
G. II..(22).(1104)
11 5077. 5098
15 5047. 5062
16 5086
18 5007. 5041. 5042. 5063
21 5027. 5043. 5091
24 5095
25 5071
27 5067. 5078
30 5009
32 5079
39 5021
42 5006
45 5036(*3*)
58 5081
G. IV.(51).(2728)
6 5020. 5065. 5083
7 5038
8 5002. 5008. 5017. 5058. 5089
9 5013. 5014. 5035. 5050. 5055. 5068(*3*). 5097
10 5026. 5028. 5044. 5052. 5053. 5056. 5072. 5092
11 5093
12 5022. 5076
13 5029. 5090
14 5046. 5074
15 5019. 5030. 5094
16 5012. 5031(*2*)
18 5004. 5033. 5034. 5048. 5054. 5069(*3*). 5075(*3*)
19 5001. 5018
20 5057. 5084
22 5015
26 5024. 5045
30 5066
G. VIII(16).(784)
3 5032

4 5037
5 5061. 5080
6 5010. 5025. 5049. 5070. 5073.
5082. 5085. 5088. 5100
8 5064
9 5096
10 5060
G. XVI.(3).(128)
2 5005
3 5016. 5040

Total 424. . .5290
Irreg. 5

Hundred 61.
G. I. . .(7). .(353)
27 6007. 6043
45 6067(*3*). 6091(*3*)
57 6079
71 6047
81 6011(*3*)
G. II. .(22).(1440)
12 6073
15 6022
17 6037
18 6087
22 6082. 6092
24 6098
25 6031. 6053
27 6075(*3*)
28 6046
31 6038
33 6019
35 6029
39 6059
41 6023
42 6051
46 6002
48 6071
49 6044
58 6089
63 6074
G. IV.(53).(3012)
7 6013. 6028
8 6001(*2*). 6052(*2*)
9 6015. 6021(*3*). 6055. 6063. 6064. 6070. 6076. 6078. 6085(*3*)
10 6010. 6034. 6049. 6058. 6088.
11 6012
12 6003. 6004. 6008. 6025. 6027. 6057. 6066. 6077. 6084. 6099. 6100(*2*)
13 6033. 6094
14 6062
15 6017. 6039. 6050. 6081. 6093.
16 6016
18 6065. 6083(*3*)
19 6009
20 6036. 6054
21 6035. 6095
24 6014(*2*). 6068. 6086
25 6056
26 6005
28 6026
30 6041
G. VIII.(14).(768)
3 6097

4 6040. 6042
6 6018. 6024. 6030. 6048
8 6032. 6061. 6069. 6080
9 6060
10 6020. 6096
G. XVI.(4).(208)
3 6045. 6072. 6090
4 6006

Total 439. .5781
Irreg. 11 Impr. 933

Hundred 62.
G. I. . .(5). .(265)
33 6163
39 6199
41 6143
59 6151
93 6131
G. II. .(28).(1704)
11 6127
14 6103
15 6115. 6147
16 6178
18 6183. 6187. 6196
19 6172·
21 6133
22 6121
27 6122
29 6166
30 6139
33 6124. 6167
34 6113
35 6134. 6197
37 6173
40 6159
41 6119
42. 6155. 6158
45 6107
49 6191
53 6101
60 6179
G. IV.(44).(2568)
6 6157
7 6108. 6142. 6193
8 6112. 6148
9. 6130. 6154. 6175
10 6106. 6117. 6169. 6198
11 6182. 6190
12 6123(*2*). 6181
14 6145. 6174. 6185
15 6102. 6109. 6125. 6126. 6129. 6171
16 6144. 6189

17 6128. 6184
18 6111. 6135. 6156(*3*).
6164
19 6140
20 6114. 6137. 6161. 6176. 6186
22 6152. 6194
25 6170
28 6146
G. VIII(20).(1200)
4 6118
5 6136. 6153. 6162. 6168
6 6132. 6138. 6150. 6165. 6177.
6180. 6192. 6195
7 6141
9 6188
10 6110
12 6104. 6200
13 6116
15 6149
G. XVI(3)..(128)
2 6160
3 6105. 6120

Total 445...5865
Irreg. 2 Impr. 975

Hundred 63.
G. I. . .(7). .(447)
43 6247
45 6211
51 6203. 6271. 6287
77 6263
129 6299
G. II. .(28).(1678)
12 6238. 6295
15 6259. 6268
16 6217
18 6267(*3). 6283(*3*)
20 6223. 6241. 6274
23 6218. 6277
25 6207. 6250
27 6227(*3*)
29 6229
30 6212. 6219
33 6243
36 6242
38 6257
42 6275
45 6239. 6291(*3*)
51 6236. 6284
57 6269
63 6221(*3*)
G. IV.(39).(2428)
6 6220. 6262
7 6253
8 6202. 6208(*2*)
9 6235. 6252. 6297.
10 6246. 6292
11 6298
12 6255 6286(*2*)
13 6282
14 6228. 6245. 6260. 6276
15 6214. 6234. 6249
16 6231. 6233. 6244. 6272(*4*)
17 6294
18 6226(*3*). 6251
19 6281. 6289
20 6261
21 6266
22 6224
23 6278
24 6209. 6215
25 6206
27 6296
35 6254
G. VIII(23).(1176)
3 6232
4 6273
5 6205. 6213. 6258. 6280. 6285
6 6210. 6222. 6225. 6237. 6288.

6290. 6300
7 6248. 6256. 6265. 6279. 6293
8 6204
9 6264
10 6201
11 6230
G. XVI(3)..(176)
3 6216
4 6240. 6270

Total 451..5905
Irreg. 9

Hundred 91.
G. I...(6)..(386)
27 9067
35 9007
45 9043
63 9091
99 9011
117 9059
G. II..(26).(1960)
15 9013
18 9003(*3*). 9055
19 9034
21 9004. 9046
26 9079
27 9031. 9094
29 9098
30 9001. 9076
33 9068
36 9049. 9083
40 9023
42 9041
44 9047
45 9019(*3*)
46 9038
48 9092
54 9099(*3*)
57 9029
69 9014. 9071
80 9026
G. IV.(42).(2928)
8 9087. 9088
9 9022. 9037. 9073. 9097
10 9025. 9058
11 9052
12 9018. 9027. 9028
13 9082
14 9017. 9032. 9066
15 9033. 9051. 9063
17 9002. 9057. 9070
18 9012. 9015. 9069(*)..
9075(*3*)

19 9053. 9095
20 9039. 9054. 9062. 9081
21 9036
22 9084
27 9008. 9050. 9074
28 9089
30 9005. 9035. 9056. 9077

G. VIII(23).(1640)
5 9010
6 9040. 9042. 9045. 9072. 9085. 9100
7 9078. 9080. 9090. 9093
8 9016(*2*). 9024. 9060
9 9061. 9074
10 9020
11 9006
12 9021. 9065
14 9096
17 9086
18 9044

G. XVI(3)..(176)
3 9030. 9048
5 9009

Total 45..7090
Irreg. 6 Impr. 1122

Hundred 92.
G. I. . .(5). .(295)
51 9199
57 9103. 9127
63 9187
67 9151
G. II. .(30).(2208)
13 9157
14 9172
19 9133
20 9124. 9183
21 9115
23 9181
27 9109. 9123. 9167. 9175
28 9137
29 9148
30 9147
32 9111
34 9122. 9166
36 9143
39 9107. 9171
40 9188
47 9173
54 9134. 9155
56 9182
57 9179
60 9131
62 9119
63 9194
72 9161
G. IV.(36).(2652)
7 9178
8 9118(*2*). 9193
9 9132. 9139. 9162. 9163(*3*)
10 9190
12 9121. 9142. 9153. 9186
13 9117
15 9106. 9136. 9195. 9196
17 9158
18 9154
19 9146
20 9169
21 9126. 9197
22 9138. 9189
24 9113. 9159
25 9101. 9125
27 9164.
28 9116
30 9140
31 9191
32 9104(*2*)
33 9149
34 9110

G. VIII(26).(1752)
4 9108
5 9102. 9160
6 9112. 9130. 9145. 9150
7 9174
8 9135. 9144. 9156. 9168. 9184. 9192. 9198
9 9105. 9114. 9180. 9185. 9200
10 9128. 9152
11 9129. 9170
12 9141
19 9176
G. XVI(3)..(176)
3 9177
4 9120. 9165

Total 465...7083
Irreg. 3 Impr. 1207

Hundred 93.
G. I...(4)..(340)
33 9283
75 9227
93 9203
139 9239
G. II..(27).(2092)
13 9277
17 9223
18 9241. 9298
21 9235
25 9293
27 9211. 9247
29 9244
30 9271
31 9263
33 9267
35 9279
36 9259
37 9274
39 9242. 9286. 9287
40 9278
45 9251(*3*)
49 9221
54 9209
60 9257.
63 9206
66 9236
75 9299
80 9281
G. IV.(47).(3216)
7 9262
8 9202. 9208. 9232
10 9238. 9289
11 9237. 9258

12 9207. 9219. 9220(*2*).
9228. 9243(*2*). 9268.
9292. 9259. 9297
13 9249. 9255
14 9217. 9226. 9253
15 9212. 9250. 9276
16 9214(*2*). 9216.
9248(*2*). 9252
17 9254
18 9234
21 9201. 9229. 9261
22 9233. 9245
24 9275. 9291
25 9231. 9294
26 9218. 9290
27 9260(*3*)
29 9215
30 9284
32 9224
36 9266(*2*)
G. VIII(20).(1440)
6 9205. 9213. 9265. 9270. 9288.
9300
7 9256
8 9222. 9225. 9273. 9280(*2*).
9285
9 9272
10 9210
11 9230
12 9204. 9264
14 9269. 9296
15 9246
G. XVI.(1)..(48)
3 9282
G. XXXII(1).(64)
2 9240

Total 454...7200
Irreg. 9 Ingr. 1145

Hundred 94.
G. I. . .(6). . .(478)
41 9319
51 9343
55 9391
87 9323
97 9311
147 9371
G. II. .(27).(1894)
15 9307. 9388
18 9355
21 9397
23 9382
25 9375
26 9337
27 9349
28 9327
30 9346. 9358. 9363. 9364
32 9377
34 9326. 9332
36 9347. 9379. 9395
41 9302
46 9351
49 9335
51 9383
52 9359
56 9314
57 9356
69 9341
G. IV.(39).(2848)
7 9340. 9367
9 9342(*3*). 9370
11 9304. 9322
12 9378. 9387
13 9313. 9318
14 9353. 9394
15 9325. 9331. 9339
16 9301. 9316(*2*)
17 9303
18 9315(*3*). 9357. 9362. 9376. 9385(*3*). 9396. 9398
21 9334. 9368. 9392
22 9305. 9317.
23 9365
24 9308. 9399
27 9369. 9374.
30 9386
32 9344(*2)
33 9329. 9389
G. VIII(25).(1752)
4 9310. 9328. 9373
5 9333
6 9352

7 9321. 9361. 9381.
8 9312(*2*). 9348. 9372.
9393. 9394. 9400
9 9390
10 9330. 9338. 9366
11 9306. 9309. 9336
12 9324
14 9350. 9380
13 9320
G. XVI.(3)..(176)
3 9384
4 9345. 9360

Total 464...7148
Irreg. 6 Impr. 1210

Hundred 95.
G. I....(8)..(708)
33 9403
45 9463
75 9439
91 9431
101 9479
105 9419
123 9467
135 9491
G. II..(24).(1706)
16 9433
18 9475
19 9466. 9487
20 9442
21 9427
24 9406. 9409. 9423
30 9459
33 9421
34 9458
36 9451. 9497(*3*)
39 9484
40 9473
42 9428
45 9411. 9437
46 9407
51 9413
57 9461
63 9404
71 9446
G. IV.(41).(2988)
8 9412(*2*). 9457(*2*).
9472
9 9493
11 9402. 9447. 9496
12 9445. 9469. 9483. 9498
13 9415. 9418. 9448. 9454. 9468.
9478

14 9436
15 9443. 9452
17 9410
18 9444. 9477. 9482. 9495
20 9500
21 9414. 9481. 9489. 9499
22 9441
24 9422(*2*). 9426. 9474.
9488
28 9494
29 9449
30 9455
35 9470
36 9476
42 9434

G. VIII(25).(1744)
4 9430
5 9417
6 9408. 9432. 9438. 9465. 9492
7 9453. 9462
8 9424. 9460. 9485. 9486. 9490
9 9420. 9450(*3*). 9471
.10 9440
11 9429. 9456
12 9425. 9435. 9464
14 9416
16 9401

G. XVI.(2)..(112)
3 9480
4 9405

Total 452..7258
Irreg. 5

Hundred 96.
G. I. . . .(5). .(471)
39 9547
69 9511. 9587
129 9551
165 9539
G. II. .(28).(1964)
16 9508
17 9535
18 9523. 9583
20 9598
24 9502. 9507
25 9559
26 9543
27 9531(*3*). 9563.
9574(*3*)
29 9532
30 9571
33 9527
34 9556. 9586
38 9524. 9567
42 9518
43 9578
48 9566
49 9599
51 9575
59 9596
60 9572
61 9533
64 9521
G. IV.(38).(2700)
8 9538. 9562. 9577
10 9517. 9553. 9573
12 9505. 9522. 9526. 9550. 9580.
9595(*2*)
13 9514. 9549. 9557
14 9501. 9529. 9542
15 9582
16 9544
17 9564
18 9558
19 9565
20 9503(*2*). 9589. 9591.
9593
21 9515. 9561
24 9519. 9579
25 9530. 9584
27 9509
28 9536
30 9506
38 9569
40 9554
G. VIII(26).(1960)
4 9568

5 9592
6 9552. 9585. 9597
7 9528
8 9510(*2*). 9513(*2*).
9537. 9540(*2*). 9588.
9600(*2*).
9 9541. 9548. 9555(*3*)
10 9525. 9534. 9594
12 9504. 9516(*2*). 9560
13 9545. 9594
14 9581
15 9512. 9590
G. XVI.(3).(176)
3 9520. 9570
5 9576

Total 469. .7271
Irreg. 10

Hundred 97
G. I. . . .(5). .(333)
33 9643
57 9619
71 9679
77 9631
95 9623
G. II. .(29).(2108)
12 9667
19 9661
20 9697
21 9607. 9613
24 9601
26 9655
27 9627(*3*). 9663.
9683(*3*)
28 9604. 9634. 9649
30 9687. 9691
32 9662
33 9692
36 9668
42 9651
43 9647
44 9602
45 9626
49 9677
52 9689
55 9629
57 9659
63 9671
66 9611
69 9644
G. IV. .(41).(3108)
9 9612(*3*). 9615. 9622.
9678

10 9628
11 9670
12 9652. 9673(*2*). 9676.
9682. 9700
13 9637
14 9642. 9658. 9664
15 9639
16 9610
17 9606
18 9603. 9675(*3*).
9693(*3*).
19 9638. 9694
20 9608. 9616. 9650. 9653
21 9699
22 9654. 9684
23 9695
24 9641. 9666
25 9609
29 9617. 9674. 9698
30 9621
32 9665
33 9635
42 9686
G. VIII(22).(1600)
4 9640
5 9618. 9625. 9685. 9688
6 9648. 9696
7 9633. 9646
8 9645. 9669
9 9630. 9657
10 9620. 9636
11 9681
12 9605. 9632. 9680(*2*)
15 9624
16 9614
18 9656
G. XVI.(3)..(192)
3 9672
4 9690
5 9660

Total 451..7341
Irreg. 7

Hundred 98.
G. I. . .(6). .(524)
39 9739. 9787
89 9767
105 9743
119 9791
133 9719
G. II.(24).(1646)
17 9703
18 9748(*3*). 9783
19 9727
21 9733
22 9742
24 9763
25 9781
26 9769. 9778
27 9747(*3*). 9751
30 9755
33 9799
35 9754
37 9788
39 9707. 9771
43 9722
46 9721. 9759
48 9731
76 9764
81 9749
G. IV.(43).(3292)
7 9718
9 9772
10 9732. 9793
11 9738. 9753
12 9706. 9745. 9762
13 9712. 9713
14 9757. 9775. 9784
15 9766. 9777. 9795
16 9736. 9797
17 9711
18 9715. 9723. 9782
19 9773
20 9714. 9796
21 9708. 9710. 9770. 9774
22 9725. 9756
24 9728
26 9794
27 9704. 9726
28 9716. 9734
29 9761
30 9746
33 9740
36 9779
38 9701
G. VIII(24).(1752)
5 9717. 9730

6 9760. 9790
7 9724. 9752
8 9702(*2*). 9729. 9735.
9758. 9780. 9792(*2*)
9 9725. 9709. 9720(*3*)
10 9741. 9750. 9798
11 9737. 9786
12 9800
14 9789
15 9776. 9785
G. XVI(3)..(192)
4 9744. 9765. 9768

Total 466..7406
Irreg. 5

Hundred 99.
G. I...(8).(638)
49 9871
51 9883
63 9811(*3*). 9859
75 9887
91 9839
111 9803
135 9851
G. II..(22).(1700)
20 9892
21 9829. 9862. 9868
22 9847
24 9817. 9874
26 9838
28 9886. 9895
30 9857
34 9826
42 9827
45 9899(*3*)
48 9863
49 9818
51 9819
52 9833
57 9836
60 9875
70 9806
77 9866
G. IV.(43).(3164)
6 9823
7 9802
8 9865
9 9832. 9843. 9853. 9898
10 9808. 9837. 9850
11 9813
12 9872. 9835
13 9846
14 9820. 9878

15 9822. 9882
16 9807. 9814. 9848
17 9852. 9893. 9897
18 9801(*3*). 9844. 9873.
9891
19 9815. 9879
21 9855
22 9841
25 9830
26 9812. 9876
27 9831
28 9881(*2*)
31 9884
32 9809(*2*)
34 9824
37 9854
39 9896
41 9869

G. VIII(24).(1712)
5 9877
6 9804. 9810. 9828. 9867. 9888.
9900
7 9805. 9858. 9885
8 9816. 9856. 9860
9 9825(*3*). 9849(*3*).
9889
10 9821. 9834
11 9894
12 9861. 9864
13 9842
16 9845
17 9890

G. XVI(3)..(192)
3 9870
4 9880
5 9840

Total 464..7406
Irreg. 7

Hundred 100.
G. I. . .(4). .(228)
39 9967
45 9907(*3*)
69 9931
75 9923
G. II.(28).(2302)
16 9991
18 9934(*3*)
23 9927. 9973
25 9949
27 9963(*3*)
28 9903. 9943
30 9938. 9979
31 9901
32 9986
34 9998
38 9908
39 9914. 9987
42 9939. 9947
46 9983
47 9935. 9946
50 9902
60 9995
63 9971
65 9959
67 9941
76 9929
85 9974
G. IV.(46).(3248)
7 9937
9 9925. 9958. 9997
10 9913. 9942. 9948. 10000
11 9970
12 9916. 9922. 9952. 9955(*2*).
9972(*2*)
13 9993
14 9988
15 9904. 9921. 9969. 9978
16 9961. 9985
17 9919. 9965
18 9910. 9915. 9964
19 9951. 9957. 9977
20 9992
21 9909
22 9953. 9956. 9962. 9999
23 9917. 9994
25 9981
26 9924
27 9989
28 9926. 9980
30 9932. 9950
34 9911

G. VIII(18).(1328)
4 9982
5 9976
6 9928. 9940. 9990
7 9906
8 9930. 9975. 9984
10 9918
11 9968
12 9905. 9920. 9936. 9954. 9996
13 9966
14 9944
G. XVI(4)..(240)
3 9933
4 9912. 9945. 9960

Total 452..7346
Irreg. 5

Hundred 117.
G. I...(1)..(147)
147 11699
G. II..(35).(2896)
16 11617
18 11698
19 11677
20 11614
22 11668
26 11647
27 11643. 11683(*3*)
29 11663
30 11602. .11623. 11659
35 11686
36 12603. 11631. 11633. 11667.
11671. 11689. 11691. 11695
37 11642
42 11657
43 11626
45 11611
51 11621
54 11619
56 11678
59 11612
63 11675
67 11639
70 11636
73 11654
81 11651(*3*)
90 11681
G. IV..(35).(2672)
8 11650
9 11608. 11692
11 11638. 11653
12 11635. 11641. 11673
13 11632. 11674

14 11652. 11662
15 11607. 11694
16 11629. 11665
17 11672
18 11601. 11627. 11637. 11664
19 11687
21 11644
23 11618
24 11615
25 11693
26 11604. 11669. 11679
28 11646
29 11630
32 11666. 11684
33 11624
35 11606

G. VIII(27).(2248)
6 11610. 11620. 11628. 11656. 11680. 11697
8 11605. 11622. 11658. 11670. 11682
9 11613. 11655
10 11616. 11625. 11676
11 11688
12 11648. 11661. 11700(*2*)
13 11634. 11645. 11649
14 11690
18 11660. 11696
21 11609

G. XVI.(2).(128)
3 11685
5 11640

Total 459. . .8091
Irreg. 3 Impr. prim. 1339

Hundred 118.
G. I. . . .(5). .(319)
39 11743
41 11719
63 11731
81 11779(*3*)
95 11783
G. II. .(28).(2560)
18 11707
21 11767
22 11727. 11758
25 11734
29 11722
30 11755
31 11701. 11708
33 11702. 11763
36 11762
39 11747. 11787
40 11791
49 11789
50 11794
53 11751
54 11723
60 11711. 11771. 11777
61 11735. 11738
65 11759
73 11717
87 11756
98 .11714
G. IV.(40).(3100)
8 11797
9 11782
10 11785
11 11740. 11757
12 11733. 11737. 11788
14 11728. 11746. 11764
15 11710. 11773. 11793
16 11705. 11775. 11776. 11796
17 11761
18 11754. 11772
19 11706
20 11716. 11768
21 11724. 11799
22 11703. 11732. 11749
25 11709. 11769. 11780
27 11795(*3*). 11798.
28 11786
29 11721. 11741.
30 11774
34 11729
40 11744
G. VIII(23).(1744)
4 11713(*3*)
6 11715. 11718. 11748. 11752

7 11720. 11742. 11778
8 11712. 11725(*2*). 11753.
11770. 11784. 11790
9 11800
10 11792
11 11736
12 11739. 11745
15 11766
16 11780
18 11726. 11765
G. XVI.(4)..(320)
4 11760
5 11704. 11730
6 11781

Total 469...8043
Irreg. 4 Impr. prim. 1369

Hundred 119.
G. I....(7)..(505)
31 11863
39 11827
47 11887
61 11839
75 11867
113 11807
139 11831
G. II...(23).(1990)
21 11878
24 11806. 11812. 11854
27 11851(*3*). 11881
30 11875
33 11884
39 11852
40 11833. 11897
43 11821
45 11871. 11899(*3*)
49 11813. 11846
52 11876
54 11828. 11843(*.3*)
58 11855
70 11801
72 11819
75 11879
G. IV.(44).(3608)
8 11848
9 11803. 11818. 11893
10 11860. 11862
11 11815
12 11823
13 11838
14 11842. 11892
15 11824. 11853. 11883
16 11857

18 11809. 11811. 11835(*3*).
11873. 11874
19 11841
20 11822(*2*). 11836. 11847.
11858. 11866
21 11859. 11888. 11898
23 11829
24 11826. 11889. 11896
26 11834. 11894
27 11804. 11810. 11861
29 11882
35 11870
36 11849
37 11885
42 11891
43 11864

G. VIII(22).(1688)
4 11872
6 11817. 11845. 11869. 11895
7 11830. 11890
8 11805. 11808(*2*). 11877
9 11802. 11820. 11825
10 11850. 11900
11 11837. 11868
12 11844. 11886
16 11814. 11840
20 11816

G. XVI.(4)..(304)
2 11880
4 11832
6 11856. 11865

Total 469...8095
Irreg. 6 Impr. 1337

Hundred 120.
G. I. . .(7). .(547)
39 11923
45 11971(*3*)
81 11903. 11939(*3*)
83 11927
95 11959
123 11987
G. II. .(22).(1912)
20 11953
21 11962
24 11995
27 11907(*9*). 11911. 11967
30 11943. 11947
31 11983
33 11974. 11979
41 11941
48 11963
49 11933. 11999
50 11975
57 11915
66 11906
69 11909
71 11981
73 11996
80 11969
G. IV.(45).(3564)
9 11992
10 11922. 11932. 11938. 11958
11 11902. 11965
12 11905. 11908. 11917. 11929. 11950. 11980. 11988(*2*). 11998
13 11986
15 11944. 11955
17 11982
18 11916. 11956. 11991
19 11918. 11993. 11994
20 11978
21 11901. 11957
23 11989
24 11926. 11964. 11972(*2*)
26 11936. 11945
27 11919. 11930. 11942. 11961(*3*)
29 11951
30 11912. 11948
32 11966(*2*)
33 11931
40 11924
42 11954

G. VIII(22).(1832)
6 11914. 11937. 11968. 11977
7 11973
8 11920(*2*). 11940. 11946
9 11913. 11925. 11935. 11949. 11952
10 11997. 12000
12 11904. 11910
15 11934
16 11976. 11984
17 11990
21 11921
G. XVI.(4)..(288)
3 11928
4 11985
5 11960
6 11970

Total 471...8143
Irreg. 8 Impr. prim. 1361

Thousand I.
G. I...(93)..(1277)

| | | |
|---|---|---|
| 1 | 1. 2. 3. 4. 7 .. | 5 |
| 3 | 11. 19. 23. 27. 31. 43. 67. 163 .. | 8 |
| 5 | 47. 79. 103. 127 .. | 4 |
| 7 | 71. 151. 223. 343. 463. 487 .. | 6 |
| 9. | 59. 83. 107. 139. 199. 211. 243(*3*). 283. 307(*3*). 331. 367. 379. 499. 547. 643. 823. 883. 907 .... | 18 |
| 11 | 167. 271. 967 ..... | 3 |
| 13 | 191. 263. 607. 631. 727 ..... | 5 |
| 15 | 131. 179. 227. 239. 347. 439. 443. 523. 571. 619. 683. 691. 739. 751. 787. 947 .. | 16 |
| 17 | 383. 991 .. | 2 |
| 19 | 311. 359. 919 ..... | 3 |
| 21 | 251. 431. 467. 503. 587. 743. 811. 827. 859. 863 .. | 10 |
| 25 | 479. 599. 647 ..... | 3 |
| 27 | 419. 491. 563. 983 .. | 4 |
| 29 | 887 ..... | 1 |
| 31 | 719. 911 .. | 2 |
| 33 | 659. 839 .. | 2 |
| 45 | 971 ..... | 1 |

Irreg. 2 pr. 2130

G. II.....(402).....(6068)
1 5. 6. 8. 9. 10. 12. 13. 15. 16. 18. 22. 25. 28. 37. 58 .. 15

| | | |
|---|---|---|
| 2 | 14. 17. 20. 32. 34. 36. 39. 46. 49. 52. 55. 63. 64. 73. 82. 97. 100. 142. 148. 193 . . | 20 |
| 3 | 26. 29. 35. 38. 44. 50. 51. 53. 54. 61. 75. 76. 81. 87. 91. 92. 99. 106. 108. 109. 115. 118. 121. 123. 124. 135. 147. 157. 162. 169. 172. 175. 187. 202. 207. 214. 235. 247. 262. 267. 268. 277. 298. 358. 397. 403. 427. 541. 652 . . . . . | 49 |
| 4 | 41. 62. 68. 94. 95. 98. 111. 113. 128. 137. 158. 178. 183. 196. 226. 256. 289. 292. 295. 313. 337. 382. 388. 415. 457. 466. 478. 562. 577. 583. 772. 862 . . . . . | 32 |
| 5 | 74. 86. 119. 122. 125. 143. 159. 166. 181. 188. 197. 218. 229. 242. 250. 303. 316. 317. 319. 346. 361. 373. 375. 394. 412. 421. 422. 423. 508. 538. 613. 625. 694. 709. 757. 847. 853. 877. 982 . . . . . | 39 |
| 6 | 89. 116. 155. 171. 203. 212. 219. 233. 241. 244. 259. 274. 275. 279. 291. 302. 323. 324. 327. 334. 351. 355. 363. 387. 433. 436. 475. 484. 507. 526. 529. 543. 567. 603. 617. 622. 628. 655. 667. 673. 676. 687. 718. 723. 763. 775. 802. 898. 955 . . . . . | 49 |
| 7 | 101. 134. 149. 173. 215. 278. 284. 287. 338. 349. 391. 447. 454. 502. 511. 535. 604. 634. 639. 653. 703. 733. 778. 807. 838. 841. 892. 997 . . . . . | 28 |
| 8 | 146. 164. 254. 257. 353. 407. 409. 452. 471. 512. 514. 527. 548. 559. 578. 722. 799. 895. 943. 958. 961 . . . . . | 22 |
| 9 | 194. 236. 293. 332. 335. 339(*3*). 362. 411. 428. 451. 459. 486. 515. 519. 531. 556. 557. 575. 586. 661. 675(*3*). 679. 707. 729. 747. 771. 783. 796. 835. 843. 844. 867. 886. 891(*3*). 922. 931. 963. 972 . . . . . | 37 |
| 10 | 206. 281. 386. 398. 401. 449. 482. 500. 601. 711. 724. 769. 788. 878. 916. 927. 937. 977 . . . . . | 18 |
| 11 | 269. 326. 389. 551. 554. 591. 623. 662. 668. 758. 767. 829. 842. 871. 879 . | 15 |
| 12 | 299. 356. 371. 395. 539. 542. 579. 593. 674. 695. 706. 766. 932. 939. 964. 979. 995. 999 . . . . . | 18 |
| 13 | 314. 458. 698. 746. 764. 773. 934. 951. 998 . . . . . | 9 |
| 14 | 404. 596. 641. 686. 692. 818. 831 . . . . . | 7 |
| 15 | 461. 509. 524. 566. 611. 635. 671. 677. 699. 716. 779. 797. 803. 815. 821. 851. 875. 908. 923. 956 . | 20 |
| 16 | 446. 521. 569. 791. 809. 857. 953 . . . . . | 7 |
| 17 | 614. 701 . . . . . | 2 |
| 18 | 626. 731. 755(*3*). 914. 929. 959. 974(*3*) . . . | 7 |
| 20 | 734. 761. 881. 926 . . . | 4 |
| 21 | 794. 899 . . . . . | 2 |
| 22 | 866 . . . . . | 1 |
| 23 | 941 . . . . . | 1 |

Irreg. 5 all *3* pr. 7394

G. IV. . .(417). . . . .(6620)

| | | |
|---|---|---|
| 1 | 21. 24. 30. 33. 40. 42. 45. 48. 57. 60. 70. 72. 78. 85. 88. 93. 102. 112. 130. 133. 177. 190. 232. 253 . . . | 24 |
| 2 | 56. 65. 66. 69. 77. 80. 84. 90. 96. 114. 117. 126. 132. 136. 138. 141. 144. 145. 150. 153. 154. 156. 160. 180. 184. 192. 198. 205. 208. 213. 217. 220. 225. 228. 238. 252. 258. 265. 282. 288. 301. 310. 322. 328. 333. 340. 352. 372. 400. 418. 438. 442. 445. 448. 498. 505. 522. 553. 568. 592. 598. 658. 697. 708. 742. 793. 928 : . . . . . | 67 |
| 3 | 104. 110. 129. 140. 152. 170. 174. 176. 182. 186. 189. 195. 200. 201. 204. 216. 222. 231. 234. 237. 245. 246. 249. 255. 261. 270. 286. 294. 297. 300. 304. 309. 315. 318. 325. 342. 348. 364. 366. 368. 370. 378. 393. 396. 405. 417. 424. 430. 432. 435. 450. 453. 460. 472. 473. 477. 483. 490. 492. 493. 496. 513. 517. 533. 537. 540. 550. 555. 565. 588. 595. 597. 606. 610. 618. 627. 637. 648. 669. 670. 682. 685. 688. 700. 702. 715. 730. 748. 753. 762. 784. 795. 808. 813. 814. 817. 826. 827. 837. 856. 913. 918. 925. 933. 940. 942. 949. 970. 973. 988 . | 110 |
| 4 | 161. 185. 221. 224. 248·. 260. 272. 276. 305. 306. 308. 320. 350. 354. 369. 376. 377. 380. 384. 392. 399. 402. 406. 410. 414. 441. 444. 468. 469. 481. 485. 495. 501. 518. 532. 544. 558. 564. 573. 574. 576(*2*). 580(*2*). 582. 589. 612. 632 640. 642. 646. 657. 663. 712. 717. 721. 732. 735. 736. 738. 745. 768. 785. 786. 790. 820(*2*). 832. 834. 850. 852. 855. 865. 868. 873. 882. 889. 900(*2*). 903. 904. 933. 938. 946. 975. 994 . . . . . | 82 |
| 5 | 209. 230. 266. 290. 296. 321. 344. 365. 381. 413. 425. 437. 455. 470. 474. 476. 488. 489. 534. 572. 590. 608. 615. 629. 633. 636. 638. 649. 664. 666. 678. 681. 726. 737. 750. 752. 754. 774. 781. 822. 830. 872. 874. 917. 921. 968. 1000 . . . . . | 47 |
| 6 | 329. 416. 426. 434. 464. 497. 516. 549. 594. 602. 605. 620. 621. 650. 651. 684. 704. 713. 725. 756. 759. 782. 800. 801. 804. 810. 812. 819. 833. 848. 864. 876. 890. 894. 901. 905. 915. 948. 954. 976. 978. 980. 981. 985. 987. 996 . . . . . | 46 |
| 7 | 341. 374. 494. 506. 530. 536. 581. 654. 806. 845. 849. 860. 893. 902. 906. 909. 935. 962 . . . . . | 18 |
| 8 | 545. 584. 644. 656. 710. 740. 749. 789. 846. 869. 884(*2*). 896. 992 . . . | 13 |
| 9 | 944. 950. 989 . . . . . | 3 |
| 10 | 689. 776. 824. 836 . . . . | 4 |
| 11 | 854. 965. 986 . . . . . | 3 |

pr. 6904

G. VIII. . .(87). . . . .(1496)

| | | |
|---|---|---|
| 1 | 105. 120. 165. 168. 210. 240. 273. 280. 312. 330. 345. 357. 385. 408. 462. 520. 760 . . . . . | 17 |

| | | |
|---|---|---|
| 2 | 264. 285. 336. 360. 390. 420. 429. 456. 465. 480. 504. 510. 525. 528. 552. 561. 570. 585. 600. 609. 616. 624. 630. 645. 660. 672. 690. 693. 720. 765. 777. 792. 798. 805. 858. 870. 880. 897. 910. 912. 952. 957. 960 ..... | 43 |
| 3 | 440. 546. 560. 665. 680. 696. 705. 714. 728. 741. 744. 780. 816. 825. 861. 885. 888. 924. 930. 936. 945. 966. 969. 984. 990 . | 25 |
| 4 | 770 ..... | 1 |
| 5 | 920 ..... | 1 |

G. XVI. . . . .(1). . . . . (16)

| | | |
|---|---|---|
| 1 | 840 ..... | 1 |

Total number of all

$$\begin{aligned} \text{genera} &= 3277 \\ \text{classes } p.p.p &= 15467 \\ \sqrt{1} + \sqrt{2} + \ldots + \sqrt{1000} &= 21097{,}661 \\ \text{Quotient} &= 0{,}733 \end{aligned}$$

Irreg. 11, 5(*2*), 6(*3*)

Thousand III.

G. I. . . . . (64). . . . . (2470)

| | | |
|---|---|---|
| 13 | 2143 ..... | 1 |
| 15 | 2203. 2347. 2647. 2683 .. | 4 |
| 21 | 2011. 2083. 2179. 2251. 2467. 2503. 2707. 2767 .. | 8 |
| 23 | 2671 ..... | 1 |
| 25 | 2887 ..... | 1 |
| 27 | 2003. 2187*. 2803 .... | 3 |
| 29 | 2287. 2311. 2383 ..... | 3 |
| 31 | 2927 ..... | 1 |
| 33 | 2027. 2267. 2423. 2539. 2731. 2851. 2971 ..... | 7 |
| 35 | 2087. 2239. 2543 ..... | 3 |
| 37 | 2447 ..... | 1 |
| 39 | 2131. 2207. 2371. 2659. 2791. 2963 ..... | 6 |
| 41 | 2551. 2719 ..... | 2 |
| 43 | 2663 ..... | 1 |
| 45 | 2039. 2063. 2243*. 2699*. 2843 ..... | 5 |
| 49 | 2111 ..... | 1 |
| 51 | 2531. 2687 ..... | 2 |
| 53 | 2711 ..... | 1 |
| 57 | 2099. 2339. 2459. 2591. 2879 ..... | 5 |
| 59 | 2399. 2903 ..... | 2 |
| 63 | 2351. 2579. 2819 ..... | 3 |

| | | |
|---|---|---|
| 69 | 2411 . . . . . | 1 |
| 73 | 2999 . . . . . | 1 |
| 87 | 2939 . . . . . | 1 |

Irreg. 3

G. II. . . .(311). . . . .(11646)

| | | |
|---|---|---|
| 6 | 2017. 2062 . . . . . | 2 |
| 7 | 2293. 2335 . . . . . | 2 |
| 8 | 2095. 2113. 2137. 2302. 2308. 2377. 2578. 2878. 2962 . . . . . | 9 |
| 9 | 2023. 2038. 2047. 2053. 2122. 2167. 2188. 2221. 2227. 2283. 2403(*3*). 2437(*3*). 2443. 2458. 2515. 2557. 2563(*3*). 2566. 2572. 2787. 2797. 2902. 2923. 2998, . . . . | 24 |
| 10 | 2164. 2281. 2407. 2452. 2473. 2487. 2500. 2527. 2638. 2722. 2743. 2818. 2836. 2857. 2983 . . . . . | 15 |
| 11 | 2182. 2215. 2263. 2326. 2374. 2623. 2662. 2677. 2815. 2863. 2917. 2935 . . | 12 |
| 12 | 2059. 2098. 2107. 2116. 2209. 2307. 2323. 2395. 2419. 2468. 2479. 2587. 2593. 2611. 2689. 2692. 2713. 2827. 2947. 2953. 2995 . . . . . | 21 |
| 13 | 2102. 2197. 2218. 2362. 2367. 2428. 2524. 2762. 2809. 2823. 2839. 2854. 2908 . . . . . | 13 |
| 14 | 2007. 2018. 2127. 2258. 2359. 2401. 2446. 2455. 2612. 2734. 2753. 2932 . . | 12 |
| 15 | 2043. 2071. 2092. 2151. 2191. 2237. 2269. 2284. 2299. 2303. 2319. 2341. 2363. 2476. 2523. 2575. 2578. 2599. 2602. 2643. 2727. 2732. 2764. 2875. 2883. 2899. 2956. 2986 . . | 28 |
| 16 | 2048. 2153. 2194. 2206. 2386. 2434. 2462. 2521. 2617. 2633. 2657. 2833 . . | 12 |
| 17 | 2026. 2029. 2103. 2119. 2234. 2333. 2389. 2391. 2463. 2918 . . . . . | 10 |
| 18 | 2155. 2161. 2199. 2228. 2417. 2491. 2511. 2547. 2619(*3*). 2627. 2644. 2723. 2763(*3*). 2783. 2867. 2897. 2916. 2943. 2979 . . . . . | 19 |
| 19 | 2031. 2069. 2381. 2477. 2554. 2738. 2746. 2799. 2858 . . . . . | 9 |
| 20 | 2078. 2297. 2375. 2402. 2404. 2498. 2559. 2777. 2866. 2942. 2959 . . . . | 11 |
| 21 | 2012. 2123. 2138. 2147. 2171. 2183. 2186. 2213. 2259. 2342. 2348. 2357. 2427. 2507. 2571. 2693. 2749. 2779. 2911. 2931. 2972 . . . . . | 21 |
| 22 | 2089. 2271. 2439. 2567. 2594. 2798 . . . . . | 6 |
| 23 | 2614. 2615. 2837 . . . . | 3 |
| 24 | 2019. 2195. 2273. 2327. 2372. 2631. 2654. 2811. 2974. 2991 . . . . . | 10 |
| 25 | 2042. 2396. 2582. 2588 . . | 4 |
| 26 | 2642. 2807. 2969 . . . . | 3 |

| | | |
|---|---|---|
| 27 | 2051. 2075(*3*). 2252. 2291. 2315(*3*). 2426. 2675(*3*). 2747. 2759. 2859. 2891(*3*). 2951. 2957 ..... | 13 |
| 28 | 2066. 2129. 2279. 2495. 2564 ..... | 5 |
| 29 | 2231. 2741 ..... | 2 |
| 30 | 2036. 2081. 2126. 2159. 2393. 2483. 2603. 2606. 2708. 2801. 2855. 2978. 2987 ..... | 13 |
| 31 | 2471. 2621. 2735. 2876 .. | 4 |
| 32 | 2084. 2174. 2276. 2306. 2519. 2558 ..... | 6 |
| 33 | 2309. 2435. 2636. 2861. 2906 ..... | 5 |
| 34 | 2804. 2831. 2894 .... | 3 |
| 35 | 2549. 2909 ..... | 2 |
| 36 | 2219(*3*). 2846 .... | 2 |
| 38 | 2441 ..... | 1 |
| 39 | 2141. 2246. 2474. 2651. 2771 ..... | 5 |
| 40 | 2729 ..... | 1 |
| 41 | 2789 ..... | 1 |
| 42 | 2609 ..... | 1 |
| 43 | 2966 ..... | 1 |

Irreg. 10

G. IV...(430).....(16232)

| | | |
|---|---|---|
| 3 | 2608 ..... | 1 |
| 4 | 2020. 2077. 2212. 2242. 2248. 2272. 2332. 2353. 2368. 2410. 2533. 2542(*2*) 2605. 2773. 2788. 2842. 2893 ..... | 17 |
| 5 | 2032. 2073. 2074. 2101. 2118. 2125. 2152. 2158. 2173. 2178. 2202. 2230. 2247. 2290. 2398. 2422. 2433, 2494. 2577. 2641. 2776. 2830. 2853. 2965 .. | 24 |
| 6 | 2022. 2025. 2028. 2035. 2050. 2052. 2067. 2068. 2082. 2086. 2096. 2104. 2115. 2132. 2139. 2148. 2157. 2163. 2172. 2223. 2257. 2260. 2268. 2275. 2278. 2305. 2317. 2322. 2338. 2358. 2388. 2412. 2413. 2425. 2451. 2475. 2482. 2488. 2493. 2512. 2517. 2538. 2545. 2548. 2592. 2598. 2620. 2629. 2650. 2653. 2658. 2667. 2668. 2676. 2682. 2695. 2698. 2704. 2710. 2715. 2718. 2725. 2740. 2748. 2752. 2755. 2758. 2794. | |
| | 2802. 2872. 2892. 2907. 2914. 2938. 2977. 2997 .. | 76 |
| 7 | 2008. 2033. 2044. 2055. 2058. 2094. 2110. 2140. 2146. 2149. 2165. 2217. 2238. 2270. 2314. 2382. 2416. 2431. 2438. 2497. 2502. 2509. 2535. 2536. 2556. 2583. 2607. 2703. 2761. 2766. 2782. 2812. 2881. 2913. 2930.. 2941. 2973 ..... | 37 |
| 8 | 2004. 2005. 2034. 2041. 2056. 2134. 2176(*2*). 2192. 2236. 2245. 2254(*2*). 2286. 2292. 2298. 2304. 2312. 2313. 2329(*2*). 2343. 2350. 2356. 2454. 2506. 2513. 2528. 2560. 2569. 2589. 2601. 2628(*2*). 2655. 2674. 2686. 2733. 2742. 2775. 2785. 2817. 2845. 2847. 2848(*2*) 2868. 2884. 2944. 2946. 2949. 2980(*2*) ..... | 47 |

| | | |
|---|---|---|
| 9 | 2014. 2049. 2060. 2076. 2079. 2091. 2106. 2108. 2117. 2124. 2133(*3*). 2175. 2181. 2198. 2214. 2221. 2235(*3*). 2241. 2253. 2266. 2295. 2300. 2318. 2344. 2349. 2355. 2361. 2387. 2430(*3*). 2461. 2522 2555. 2581. 2595. 2626. 2634. 2635. 2637. 2646(*3*). 2673. 2700(*3*). 2716. 2770. 2778. 2781. 2795. 2806. 2828. 2835(*3*). 2862. 2887. 2888. 2890. 2895. 2950. 2955. 2988. 2989 . . | 58 |
| 10 | 2057. 2061. 2150. 2154. 2166. 2177. 2249. 2250. 2255. 2282. 2449. 2514. 2529. 2532. 2596. 2656. 2678. 2724. 2751. 2810. 2844. 2869. 2871. 2874. 2896. 2919. 2929. 2948. 2975 . . . . . | 29 |
| 11 | 2135. 2204. 2216. 2334. 2364. 2421. 2489. 2492. 2570. 2573. 2645. 2648. 2649. 2672. 2757. 2841. 2922. 2933. 3934. 2967 . . | 20 |
| 12 | 2006. 2009. 2045. 2105. 2156. 2168. 2169. 2196. 2211. 2225. 2229. 2316. 2331. 2366. 2379(*2*). 2390. 2420(*2*). 2432. 2450. 2466. 2484. 2499. 2597. 2661. 2691. 2696. 2701. 2702(*2*). 2739. 2754. 2780· 2816. 2822. 2824. 2825. 2852. 2873. 2900. 2993(*2*) . . . . . | 39 |
| 13 | 2015. 2222. 2453. 2469. 2481. 2510. 2679. 2721. 2792. 2826. 2901. 2984 . . | 12 |
| 14 | 2114. 2144. 2162. 2274. 2324. 2354. 2384. 2429. 2486. 2501. 2525. 2587. 2744. 2768. 2774 . . . . | 15 |
| 15 | 2096. 2189. 2264. 2285. 2321. 2330. 2378. 2406. 2444. 2526. 2534. 2537. 2540. 2630. 2684. 2690. 2694. 2750. 2796. 2813. 2864. 2889 . . . . . | 22 |
| 16 | 2180. 2201. 2336. 2546. 2561. 2624(*2*). 2639. 2669. 2882. 2921. 2994. 2996 . . . . . | 12 |
| 17 | 2021. 2456. 2666. 2726 . . | 4 |
| 18 | 2054. 2369. 2414. 2504. 2516. 2681. 2705. 2786. 2915. 2924 . . . . . | 10 |
| 19 | 2294 . . . . . | 1 |
| 20 | 2714. 2756(*2*). 2936. 2954. 2981 . . . . . | 5 |
| 21 | 2834 . . . . . | 1 |

Irreg. 13 (2). 6 (3). Sa..19

G. VIII. . .(184). . . .(6344)

| | | |
|---|---|---|
| 2 | 2002. 2013. 2080. 2088. 2128. 2170. 2233. 2277. 2392. 2632. 2737. 2832. 2068 . . . . . | 13 |
| 3 | 2037. 2065. 2160. 2185. 2190. 2193. 2200. 2205. 2220. 2262. 2325. 2346. 2352. 2370. 2373. 2380. 2418. 2424. 2440. 2457. 2472. 2485. 2508. 2530. 2550. 2553. 2562. 2590. 2613. 2622. 2680. 2685. 2697. 2728. 2800. 2808. 2860. 2905. 2920. 2937. 2970. 2982. 2992 . . . . | 43 |
| 4 | 2010. 2016. 2046. 2072. 2100. 2112(*2*). 2130. 2142. 2208. 2232. 2244. 2256. 2265. 2289. 2296. 2320. 2328. 2337. 2340. 2365. 2385(*2*). 2397. 2400(*2*). 2436. 2442. 2445. 2448(*2*). 2464. 2465. 2470. 2478. 2490. 2496. 2568. 2580. 2584. 2652. 2665. 2688. 2706. 2709. 2717. 2745. 2772. 2790. 2793. 2821. 2829. 2850(*2*). 2865. 2880. 2898. 2928. 2940. 2952. 2958. 2985 . . . . . | 57 |

| | | |
|---|---|---|
| 5 | 2030. 2070. 2085. 2093. 2109. 2121. 2136. 2226. 2288. 2360. 2394. 2405. 2409. 2415. 2460. 2505. 2541. 2544. 2552. 2585. 2610. 2618. 2625. 2664. 2670. 2712. 2769. 2814. 2838. 2877. 2886. 2904. 2910. 2926. 2990: 3000 . . | 36 |
| 6 | 2001. 2064. 2090. 2240. 2301. 2376. 2408. 2480. 2564. 2574. 2600. 2604. 2660. 2720. 2736. 2784. 2820. 2912. 2925. 2961. 2964. 2976 . . . . . | 22 |
| 7 | 2024. 2120. 2210. 2345. 2616. 2765. 2870 . . . . . | 7 |
| 8 | 2576. 2840. 2849. 2945. 2960 . . . . . | 5 |
| 9 | 2261 . . . . . | 1 |

Irreg. 5 (2)

G. XVI. . . .(11). . . . .(400)

| | | |
|---|---|---|
| 2 | 2040. 2145. 2280. 2310. 2520. 2640. 2730. 2760 . . | 8 |
| 3 | 2184. 2805. 2856 . . . . . | 3 |

Total of all

| | | | |
|---|---|---|---|
| genera *p.p.p* | = 4054 | expected | 4051,3 |
| classes *p.p.p* | = 37092 | . . | 37074,3 |
| . . impr. *p.p* | = 6182 | | |

Irreg. 18(*2*). 19(*3*). Sa= 37

Thousand X.

Genera I.

| | |
|---|---|
| 27 | 9067 |
| 33 | 9283. 9403. 9643 |
| 35 | 9007 |
| 39 | 9547. 9739. 9787. 9967 |
| 41 | 9319 |
| 45 | 9043. 9463. 9907(*3*) |
| 49 | 9871 |
| 51 | 9199. 9343. 9883 |
| 55 | 9391 |
| 57 | 9103. 9127. 9619' |
| 63 | 9091. 9187. 9811(*3*). 9859 |
| 67 | 9151 |
| 69 | 9511. 9587. 9931 |
| 71 | 9679 |
| 75 | 9227. 9439. 9887. 9923 |
| 77 | 9631 |
| 87 | 9323 |
| 89 | 9767 |
| 91 | 9431. 9839 |

| | |
|---|---|
| 93 | 9203 |
| 95 | 9623 |
| 97 | 9311 |
| 99 | 9011 |
| 101 | 9479 |
| 105 | 9419. 9743 |
| 111 | 9059. 9803 |
| 119 | 9791 |
| 123 | 9467 |
| 129 | 9551 |
| 133 | 9719 |
| 135 | 9491. 9851 |
| 139 | 9239 |
| 147 | 9371 |
| 165 | 9539 |

.57. . .4401.

Genera II.

| | | |
|---|---|---|
| 12 | 9667 . . . . . | 1 |
| 13 | 9157. 9277 . . . . . | 2 |
| 14 | 9172 . . . . . | 1 |
| 15 | 9013. 9307. 9388 . . . . | 3 |
| 16 | 9433. 9508. 9991 . . . . | 3 |
| 17 | 9223. 9535. 9703 . . . . | 3 |
| 18 | 9003(*3*). 9055. 9241. 9298. 9355. 9475. 9523. 9583. 9748(*3*). 9783. 9934(*3*) . . . . . | 11 |
| 19 | 9034. 9133. 9466. 9487. 9661. 9727 . . . . . | 6 |
| 20 | 9124. 9183. 9442. 9598. 9697. 9892 . . . . . | 6 |
| 21 | 9004. 9046. 9115. 9235. 9397. 9427. 9607. 9613. 9733. 9829. 9862. 9868 . . | 12 |
| 22 | 9742. 9847 . . . . . | 2 |
| 23 | 9181. 9382. 9927. 9973 . . | 4 |
| 24 | 9406. 9409. 9423. 9502. 9507. 9601. 9763. 9817. 9874 . . . . . | 9 |
| 25 | 9293. 9375. 9559. 9781. 9949 . . . . . | 5 |
| 26 | 9079. 9337. 9543. 9655. 9769. 9778. 9838 . . . . . | 7 |
| 27 | 9031. 9094. 9109. 9123. 9167. 9175. 9211. 9247. 9349. 9531(*3*). 9563. 9574(*3*). 9627(*3*). 9663. 9683(*3*). 9747(*3*). 9751. 9963(*3*) . . . . . | 18 |
| 28 | 9137. 9327. 9604. 9634. 9649. 9886. 9895. 9903. 9943 . . . . . | 9 |
| 29 | 9098. 9148. 9244. 9532 . . | 4 |
| 30 | 9001. 9076. 9147. 9271. 9346. 9358. 9363. 9364. 9459. 9571. 9687. 9691. 9755. 9857. 9938. 9979 . . | 16 |
| 31 | 9263. 9901 . . . . . | 2 |
| 32 | 9111. 9377. 9662. 9986 . . | 4 |
| 33 | 9068. 9267. 9421. 9527. 9692. 9799 . . . . . | 6 |
| 34 | 9122. 9166. 9326. 9332. 9458. 9556. 9586. 9826. 9998 . . . . . | 9 |

| | | |
|---|---|---|
| 35 | 9279. 9754 ..... | 2 |
| 36 | 9049. 9083. 9143. 9259. 9347. 9379. 9395. 9451. 9497(*3*). 9668 ..... | 10 |
| 37 | 9274. 9788 ..... | 2 |
| 38 | 9524. 9567. 9908 ..... | 3 |
| 39 | 9107. 9171. 9242. 9286. 9287. 9484. 9707. 9771. 9914. 9987 ..... | 10 |
| 40 | 9023. 9188. 9278. 9473 .. | 4 |
| 41 | 9302 ..... | 1 |
| 42 | 9041. 9428. 9518. 9651. 9827. 9939. 9947 ..... | 7 |
| 43 | 9578. 9647. 9722 ..... | 3 |
| 44 | 9047. 9602 ..... | 2 |
| 45 | 9019(*3*). 9251(*3*). 9411. 9437. 9626. 9899(*3*). | 6 |
| 46 | 9038. 9351. 9407. 9721. 9759. 9983 ..... | 6 |
| 47 | 9173. 9935. 9946 ..... | 3 |
| 48 | 9092. 9566. 9731. 9863 .. | 4 |
| 49 | 9221. 9335. 9599. 9677. 9818 ..... | 5 |
| 50 | 9902 ..... | 1 |
| 51 | 9383. 9413. 9575. 9819 .. | 4 |
| 52 | 9359. 9689. 9833 ..... | 3 |
| 54 | 9099(*3*). 9134. 9155. 9209 ..... | 4 |
| 55 | 9629 ..... | 1 |
| 56 | 9182. 9314 ..... | 2 |
| 57 | 9029. 9179. 9356. 9461. 9659. 9836 ..... | 6 |
| 59 | 9596 ..... | 1 |
| 60 | 9131. 9257. 9572. 9875. 9995 ..... | 5 |
| 61 | 9533 ..... | 1 |
| 62 | 9119 ..... | 1 |
| 63 | 9194. 9206. 9404. 9671. 9971 ..... | 5 |
| 64 | 9521 ..... | 1 |
| 65 | 9959 ..... | 1 |
| 66 | 9236. 9611 ..... | 2 |
| 67 | 9941 ..... | 1 |
| 69 | 9014. 9071. 9341. 9644 .. | 4 |
| 70 | 9806 ..... | 1 |
| 71 | 9446 ..... | 1 |
| 72 | 9161 ..... | 1 |
| 75 | 9299 ..... | 1 |
| 76 | 9764. 9929 ..... | 2 |
| 77 | 9866 ..... | 1 |
| 80 | 9026. 9281 ..... | 2 |
| 81 | 9749 ..... | 1 |
| 85 | 9974 ..... | 1 |

Total 265. .19580
Irreg. 14.

Genera IV.

| | | |
|---|---|---|
| 6 | 9823 . . . . . | 1 |
| 7 | 9178. 9262. 9340. 9367. 9718. 9802. 9937 . . . . . | 7 |
| 8 | 9087. 9088. 9118(*2*). 9193. 9202. 9208. 9232. 9412(*2*). 9457(*2*). 9472. 9538. 9562. 9577. 9865 . . . . . | 14 |
| 9 | 9022. 9037. 9073. 9097. 9132. 9139. 9162. 9163(*3*). 9342(*3*). 9370. 9493. 9612(*3*). 9615. 9622. 9678. 9772. 9832. 9843. 9853. 9898. 9925. 9958. 9997 . . . . . | 23 |
| 10 | 9025. 9058. 9190. 9238. 9289. 9517. 9553. 9573. 9628. 9732. 9793. 9808. 9837. 9850. 9913. 9942. 9948. 10000 . . . . . | 18 |
| 11 | 9052. 9237. 9258. 9304. 9322. 9402. 9447. 9496. 9670. 9738. 9753. 9813. 9970 . . . . . | 13 |
| 12 | 9018. 9027. 9028. 9121. 9142. 9153. 9186. 9207. 9219. 9220(*2*). 9228. 9243(*2*). 9268. 9292. 9295. 9297. 9378. 9387. 9445. 9469. 9483. 9498. 9505. 9522. 9526. 9550. 9580. 9595(*2*). 9652. 9673(*2*). 9676. 9682. 9700. 9706. 9745. 9762. 9835. 9872. 9916. 9922. 9952. 9955(*2*). 9972(*2*) | 43 |
| 13 | 9082. 9117. 9249. 9255. 9313. 9318. 9415. 9418. 9448. 9454. 9468. 9478. 9514. 9549. 9557. 9637. 9712. 9713. 9846. 9993 . . | 20 |
| 14 | 9017. 9032. 9066. 9217. 9226. 9253. 9353. 9354. 9436. 9501. 9529. 9542. 9642. 9658. 9664. 9757. 9775. 9784. 9820. 9878. 9988 . . . . . | 21 |
| 15 | 9033. 9051. 9063. 9106. 9136. 9195. 9196. 9212. 9250. 9276. 9325. 9331. 9339. 9443. 9452. 9582. 9639. 9766. 9777. 9795. 9822. 9882. 9904. 9921. 9969. 9978 . . . . . | 26 |
| 16 | 9214(*2*). 9216(*2*). 9248(*2*). 9252. 9301. 9316(*2*). 9544. 9610. 9736. 9797. 9807. 9814. 9848. 9961. 9985 . . . . . | 15 |
| 17 | 9002. 9057. 9070. 9158. 9254. 9303. 9410. 9564. 9606. 9711. 9852. 9893. 9897. 9919. 9965 . . . . . | 15 |
| 18 | 9012. 9015. 9069(*3*). 9075(*3*). 9154. 9234. 9315(*3*). 9357. 9362. 9376. 9385(*3*). 9396. 9398. 9444. 9477. 9482. 9495. 9558. 9603. 9675(*3*). 9693(*3*). 9715. 9723. 9782. 9801(*3*). 9844. 9873. 9891. 9910. 9915. 9964 . . . . . | 31 |
| 19 | 9053. 9095. 9146. 9565. 9638. 9694. 9773. 9815. 9879. 9951. 9957. 9977 . . | 12 |
| 20 | 9039. 9054. 9062. 9081. 9169. 9500. 9503(*2*). 9589. 9591. 9593. 9608. 9616. 9650. 9653. 9714. 9796. 9992 . . . . . | 17 |
| 21 | 9036. 9126. 9197. 9201. 9229. 9261. 9334. 9368. 9392. 9414. 9481. 9489. 9499. 9515. 9561. 9699. 9708. 9710. 9770. 9774. 9855. 9909 . . . . . | 22 |
| 22 | 9084. 9138. 9189. 9233. 9245. 9305. 9317. 9441. 9654. 9684. 9725. 9756. 9841. 9953. 9956. 9962. 9999 . . . . . | 17 |
| 23 | 9365. 9695. 9917. 9994 . . | 4 |
| 24 | 9113. 9159. 9275. 9291. 9308. 9399. 9422(*2*). 9426. 9474. 9488. 9519. 9579. 9641. 9666. 9728 . . | 15 |

| | | |
|---|---|---|
| 25 | 9101. 9125. 9231. 9294. 9530. 9584. 9609. 9830. 9981 ..... | 9 |
| 26 | 9218. 9290. 9794. 9812. 9876. 9924 ..... | 6 |
| 27 | 9038. 9050. 9074. 9164. 9260(*3*). 9369. 9374. 9509. 9704. 9726. 9831. 9989 ..... | 12 |
| 28 | 9089. 9116. 9494. 9536. 9716. 9734. 9881(*2*). 9926. 9980 ..... | 9 |
| 29 | 9215. 9449. 9617. 9674. 9698. 9761 ..... | 6 |
| 30 | 9005. 9035. 9056. 9077. 9140. 9284. 9386. 9455. 9506. 9621. 9746. 9932. 9950 ..... | 13 |
| 31 | 9191. 9884 ..... | 2 |
| 32 | 9104(*2*). 9224. 9344(*2*). 9665. 9809(*2*) ..... | 5 |
| 33 | 9149. 9329. 9389. 9635. 9740 ..... | 5 |
| 34 | 9110. 9824. 9911 .... | 3 |
| 35 | 9470 ..... | 1 |
| 36 | 9266(*2*). 9476. 9779 .. | 3 |
| 37 | 9854 ..... | 1 |
| 38 | 9569. 9701 ..... | 2 |
| 39 | 9896 ..... | 1 |
| 40 | 9554 ..... | 1 |
| 41 | 9869 ..... | 1 |
| 42 | 9434. 9686 ..... | 2 |

Total 416...30144
Irreg. 20(*2*). 11(*3*)

Genera VIII.

| | | |
|---|---|---|
| 4 | 9108. 9310. 9328. 9373. 9430. 9568. 9640. 9982. | |
| 5 | 9010. 9102. 9160. 9333. 9417. 9592. 9618. 9625. 9685. 9688. 9717. 9730. 9877. 9976 ..... | 14 |
| 6 | 9040. 9042. 9045. 9072. 9085. 9100. 9112. 9130. 9145. 9150. 9206. 9213. 9265. 9270. 9288. 9300. 9352. 9408. 9432. 9438. 9465. 9492. 9552. 9585. 9597. 9648. 9696. 9760. 9790. 9804. 9810. 9828. 9867. 9888. 9900. 9928. 9940. 9990 ..... | 38 |
| 7 | 9078. 9080. 9090. 9093. 9174. 9256. 9321. 9361. 9381. 9453. 9462. 9528. 9633. 9646. 9724. 9752. 9805. 9858. 9885. 9906 .. | 20 |
| 8 | 9016(*2*). 9024. 9060. 9135. 9144. 9156. 9168. 9184. 9192. 9198. 9222. 9225. 9273. 9280(*2*). 9285. 9312(*2*). 9348. 9372. 9393. 9394. 9400. 9424. 9460. 9485. 9486. 9490. 9510(*2*). 9513(*2*). 9537. 9540(*2*). 9588. 9600(*2*). 9645. 9669. 9702(*2*). 9729. 9735. 9758. 9780. 9792(*2*). 9816. 9856. 9860. 9930. 9975. 9984 ..... | 46 |
| 9 | 9061. 9064. 9105. 9114. 9180. 9185. 9200. 9272. 9390. 9420. 9450(*3*). 9471. 9541. 9548. 9555(*3*). 9630. 9657. 9705. 9709. 9720(*3*). 9825(*3*). 9849(*3*). 9889 ... | 23 |
| 10 | 9020. 9128. 9152. 9210. 9330. 9338. 9366. 9440. 9525. 9534. 9594. 9620. 9636. 9741. 9750. 9798. 9821. 9834. 9918 .... | 19 |
| 11 | 9006. 9129. 9170. 9230. 9306. 9309. 9336. 9429. 9456. 9681. 9737. 9786. 9894. 9968 ..... | 14 |

| | | |
|---|---|---|
| 12 | 9021. 9065. 9141. 9204. 9264. 9324. 9425. 9435. 9461. 9504. 9516(*2*). | |
| | 9560. 9605. 9632. 9680(*2*). 9900. 9861. 9864. 9905. 9920. 9936. 9954. 9996 .. | 23 |
| 13 | 9545. 9594. 9842. 9966 .. | 4 |
| 14 | 9096. 9269. 9296. 9350. 9380. 9416. 9581. 9789. 9944 ..... | 9 |
| 15 | 9246. 9320. 9512. 9590. 9624. 9776. 9785 .... | 7 |
| 16 | 9401. 9614. 9845 .... | 3 |
| 17 | 9086. 9890 ..... | 2 |
| 18 | 9044. 9656 ..... | 2 |
| 19 | 9176 ..... | 1 |
| Total 233..16680 | | |

Irreg. 11(*2*), 5(*3*)

Genera XVI.

| | | |
|---|---|---|
| 3 | 9030. 9048. 9177. 9281. 9384. 9480. 9520. 9570. 9672. 9870. 9933 .... | 11 |
| 4 | 9120. 9165. 9345. 9360. 9405. 9690. 9744. 9765. 9768. 9880. 9912. 9945. 9960 ..... | 13 |
| 5 | 9009. 9576. 9660. 9840 .. | 4 |
| 28...1680. | | |

Genera XXXII.

| | | |
|---|---|---|
| 2 | 9240 ..... | 1 |
| 1...64 | | |

Sum of all

| | |
|---|---|
| classes *p.p.p.* | 72549 |
| expected | 72572 |
| $\Sigma\gamma\sqrt{D}$ | 72775 |
| genera *p.p.p.* | 4595 |
| expected | 4594,9 |

Irreg. 31(*2*). 32(*3*)...63

Greatest quotient

1,729662 from 9434. IV, 42

least

0,2421048 from 9823. IV, 6

Number of classes

| | |
|---|---|
| less than half the root | 244 |
| less than the root greater than half | 566 |
| greater than the root | 199 |

Eight hundred negative determinants.
of the form $-(15n + 7)$.

G. I. . . . . (93). . . . . (2793)

| | | |
|---|---|---|
| 1 | 7 . . . . . | 1 |
| 3 | 67 . . . . . | 1 |
| 5 | 127 . . . . . | 1 |
| 7 | 487 . . . . . | 1 |
| 9 | 307(*3*). 367. 547. 907. 1087 . . . . . | 5 |
| 11 | 967 . . . . . | 1 |
| 13 | 607. 727 . . . . . | 2 |
| 15 | 787. 1327. 1567. 1747. 1867. 2347. 2647 . . . . . | 7 |
| 17 | 4447 . . . . . | 1 |
| 19 | 3607. 4327. 5527 . . . . . | 3 |
| 21 | 1627. 1987. 2467. 2707. 2767. 3067. 3187. 3907. 5107. 5647 . . . . . | 10 |
| 23 | 1447. 3847 . . . . . | 2 |
| 25 | 2887 . . . . . | 1 |
| 27 | 3307. 3547(*3*). 4027(*3*). 4987. 6007. 6427. 7027. 9067. 10627 . . . . . | 9 |
| 29 | 2287. 7207. 7687 . . . . . | 3 |
| 31 | 3727. 8647 . . . . . | 2 |
| 33 | 3967. 4567. 5167. 6547. 6967. 7867. 8167. 10987 . | 8 |
| 35 | 9007 . . . . . | 1 |
| 37 | 6367 . . . . . | 1 |
| 39 | 4507. 5347. 7507. 9547. 9787. 9967. 10567. 11467. 11587. 11827 . . . . . | 10 |
| 41 | 11047 . . . . . | 1 |
| 43 | 5407. 6247. 8527. 8887 . . | 4 |
| 45 | 5227(*3*). 5827. 6067(*3*). 6607. 8287. 8467. 8707. 9907(*3*). 10267 . . . . | 9 |
| 47 | 7927. 11887 . . . . . | 2 |
| 51 | 6907. 10687 . . . . . | 2 |
| 53 | 11287 . . . . . | 1 |
| 57 | 9127 . . . . . | 1 |
| 61 | 11527 . . . . . | 1 |
| 63 | 9187 . . . . . | 1 |
| 69 | 10867 . . . . . | 1 |

Irreg. 6

G. II. . . . .(343). . . .(10010)

| | | |
|---|---|---|
| 1 | 22. 37 . . . . . | 2 |
| 2 | 52. 82. 97. 142 . . . . | 4 |
| 3 | 157. 172. 187. 202. 247. 262. 277. 397. 427. 652 . . | 10 |
| 4 | 292. 337. 382. 457. 562. 577. 772. 862 . . . . . | 8 |
| 5 | 412. 757. 847. 877. 982 . . | 5 |
| 6 | 622. 667. 802. 1027. 1042. 1282. 1297. 1387. 1507. 1807. 2017. 2062 . . . . . | 12 |
| 7 | 502. 892. 997. 1117. 1237. 1372. 1402. 1597. 1642. 1852 . . . . . | 10 |
| 8 | 1252. 1657. 1822. 2137. 2302. 2377. 2962. 3217. 4687 . . . . . | 9 |
| 9 | 922. 1132. 1147. 1207. 1267(*3*). 1687. 1882. 1927. 2047. 2122. 2167. 2227. 2437(*3*). 2557. 2572. 2797. 2902. 3037. 3292. 3427. 3532. 5692 . . | 22 |
| 10 | 937. 1492. 1522. 2407. 2452. 2527. 2722. 2857. 3007. 3412. 3697. 4057. 4162. 4372. 4852 . . . . | 15 |
| 11 | 1942. 2182. 2662. 2677. 2917. 3637. 3802. 4957. 5077. 5212. 6127. 6637 . . | 12 |
| 12 | 1732. 1762. 1777. 2107. 2587. 2692. 2827. 2947. 3127. 3202. 3742. 3787. 4132. 4222. 4267. 4387. 4657. 4747. 4867. 4882. 5182. 5587. 5707. 5947. 7417. 7492. 7522. 7987. 8002. 9667 . . . . . | 30 |
| 13 | 2197. 2362. 3622. 3862. 4207. 4282. 5482. 6742. 6847. 6997. 8422. 8572. 8842. 9157. 9277. 10207. 11302 . . . . . | 17 |
| 14 | 2932. 3022. 3457. 3487. 5422. 5602. 5812. 5887. 6337. 8017. 8782. 9172 . . | 12 |
| 15 | 2092. 2602. 3142. 3517. 3667. 3877. 4087. 4357. 4492. 4627. 5047. 5062. 5437. 6022. 6442. 6667. 6727. 6892. 6922. 7087. 7162. 7387. 7477. 7627. 8227. 8677. 8812. 8947. 9307. 10147. 10732. 10957 | 32 |
| 16 | 2617. 3247. 4612. 4702. 6217. 7177. 8452. 8962. 10327. 10462. 11617 . . . | 11 |
| 17 | 4762. 4927. 5197. 5287. 5557. 6037. 6502. 6652. 6982. 7132. 7327. 7402. 7642. 7702. 8047. 8317. 10042. 11482 . . . . . | 18 |
| 18 | 4297. 5737. 5767. 5857(*3*). 6187. 6382. 6577. 6787. 7057. 7267(*3*). 7732. 7807. 8107(*3*). 8212. 8347. 8407. 8482. 8737. 8767. 10747(*3*). 10882. 11257. 11347. 11707 . . . | 24 |
| 19 | 4252. 4597. 4822. 5722. 6172. 9487. 9727. 10597. 10837. 11062. 11197. 11677 | 12 |
| 20 | 3442. 4177. 7012. 8542. 9442. 9697. 9892. 10162. 10177 . . . . . | 9 |
| 21 | 5242. 5932. 6487. 7147. 7447. 8182. 8332. 9397. 9427. 9607. 9862. 10012. 10102. 10342. 10927. 11002. 11227. 11317. 11767. 11962 | 20 |
| 22 | 6082. 6772. 7297. 7537. 7822. 9742. 9847. 11167 . | 8 |
| 23 | 6277. 6397. 7237. 7717. 8902. 9382. 10522. 10762 . | 8 |
| 24 | 7747. 7972. 8377. 9502. 9817. 10372. 10657. 10807. 10852. 10942. 11107. 11422. 11812 . . . . . | 13 |
| 25 | 10477 . . . . . | 1 |
| 26 | 7342. 7762. 9337. 11647 . | 4 |

| | | |
|---|---|---|
| 27 | 8587. 9247. 10357. 10447. 11212 ..... | 5 |
| 28 | 11497 ..... | 1 |
| 29 | 9532. 11722 ..... | 2 |
| 30 | 10027. 11182. 11602. 11947 | 4 |
| 32 | 11332 ..... | 1 |
| 34 | 10702 ..... | 1 |
| 35 | 11437 ..... | 1 |

Irreg. 6

G. IV....(310)....(9688)

| | | |
|---|---|---|
| 1 | 112. 232 ..... | 2 |
| 2 | 217. 322. 352. 442. 532. 592. 697. 742. 1012 ... | 9 |
| 3 | 472. 517. 637. 682. 817. 1072. 1162. 1177. 1192. 1222. 1432. 1612 .... | 12 |
| 4 | 712. 832. 1057. 1312(*2*). 1357. 1417. 1462. 1477. 1537. 1552. 1582(*2*). 1717. 1792. 1897. 1912. 1957. 2077. 2212. 2242. 2272. 2332. 2542(*2*). 2842. 3172(*2*). 3322. 3502(*2). 3712 .... | 27 |
| 5 | 1102. 1342. 1702. 2032: 2152. 2422. 3082. 3367. 3382. 3397. 3817. 3922. 4102. 4342. 5272. 5377 .. | 16 |
| 6 | 1837. 1972. 2257. 2317. 2482. 2512. 2752. 2872. 2977. 3052. 3232. 3337. 3577. 3592. 3652. 3892. 3937. 3997. 4012. 4117. 4192. 4237. 4417. 4432. 4477. 4537. 4552. 4642. 4717. 4792. 4942. 4972. 5317. 5362. 5467. 5497. 5842. 6157. 6262. 6307. 6352. 6472. 6682. 6862. 7282. 7837. 10432 .... | 47 |
| 7 | 2497. 2782. 2812. 3097. 3112. 3277. 3352. 3562. 3982. 4582. 4732. 5122. 5137. 5302. 5902. 5977. 6142. 6517. 7792. 9262. 9367. 9802. 9937 .... | 23 |
| 8 | 3262(*2*). 3682. 3757. 3772. 3832. 4402. 4672(*2*). 4777(*2*). 5002. 5017.. 5257. 5392. 5572. 5632. 5752. 5917. 5962. 6052(*2*). 6112. 6202. 6532. 6697. 6757. 6817. 7252. 7312. 7357. 7582(*2*). 7597. 8122. 8197. 8257. 8497. 8992(*2*). 9202. 9232. 9412(*2*). 9457(*2*). 9472. 9562. 9577. 10132(*2*). 10312. 10537. 10642. 11092. 11377. 11512. 11572. 11797 | 50 |
| 9 | 4042. 4072. 4147. 4837. 4897. 4912(*3*). 5452. 5542. 5677. 5782. 5872. 6322. 6412. 6457. 7102. 7117. 7372. 7852. 7942. 8062. 8092. 8152. 8242(*3*). 8752. 8827(*3*). 9022. 9037. 9097. 9622. 9772. 9832. 9997. 10222. 10252(*3*). 10717. 10822. 11407. 11692. 11782. 11992 | 40 |
| 10 | 4462. 4807. 5092. 5332. 6292. 6592. 6712. 6802. 6937. 7432. 7897. 8077. 8137. 8302. 8557. 8617. 8692. 8857. 8977. 9517. 10057. 10087. 10297. 10402. 10552. 10777. 10972. 11122. 11272. 11932 ..... | 30 |
| 11 | 5662. 7222. 7567. 7612. 7882. 8362. 8662. 8917. 9052. 9322. 10417. 10492. 11452. 11902 ..... | 14 |

| | | |
|---|---|---|
| 12 | 5617. 6562. 6877. 7042. 7552. 7957. 8392. 8722. 9142. 9292. 9652. 9682. 9922. 9952. 10237. 10387. 10612. 10672. 11017. 11542. 11737. 11917 ..... | 22 |
| 13 | 8872. 9082. 9637. 9712. 10117. 11632 ..... | 6 |
| 14 | 8032. 9217. 9757. 11662. 11842 ..... | 5 |
| 15 | 8797. 10072. 10507 ... | 3 |
| 16 | 10282. 10897. 11392(*2*). 11857 ..... | 4 |

Irreg. 19

G. VIII....(54)....(1856)

| | | |
|---|---|---|
| 2 | 952. 1672. 2002. 2392. 2632. 2737 ..... | 6 |
| 3 | 2992. 3157. 3952. 4522. 5032. 5797. 6097. 6232. 6832. 7912 ..... | 10 |
| 4 | 3472. 4312. 5152(*2*). 5992. 6622. 6952. 7072(*2*). 7672(*2*). 8437. 8512. 8932. 9982. 11152. 11872 | 14 |
| 5 | 5512. 7192. 7462. 7657. 7777. 8632. 9592. 9877. 11242. 11362. 11557 ... | 11 |
| 6 | 8272. 9112. 9352. 10192. 10582. 10792. 10912. 11032. 11077. 11137. 11752. 11977 | 12 |
| 7 | 8602 ..... | 1 |

Irreg. 3

All genera 2451 expected 2445,10
all classes 24347 expected 24358,82

$$\left.\begin{array}{l}\left.\begin{array}{ll}8n\ldots.. & 3068\\ 8n+4\,.. & 3010\end{array}\right\}6078\\ \left.\begin{array}{ll}8n+2\,.. & 3062\\ 8n+6\,.. & 3034\end{array}\right\}6096\end{array}\right\}12174$$

$$\left.\begin{array}{l}\left.\begin{array}{ll}8n+1\,.. & 3076\\ 8n+5\,.. & 3026\end{array}\right\}6102\\ \left.\begin{array}{ll}8n+3\,.. & 3033\\ 8n+7\,.. & 3038\end{array}\right\}6071\end{array}\right\}12173$$

| | | |
|---|---|---|
| $7n\ldots..$ | $115\,..$ | 3411 |
| $7n+1\,..$ | $115\,..$ | 3009 |
| $7n+2\,..$ | $114\,..$ | 2975 |
| $7n+4\,..$ | $114\,..$ | 2993 |
| $R\,7\ldots$ | $343\,..$ | 8977 |
| $7n+3\,..$ | $114\,..$ | 3979 |
| $7n+5\,..$ | $114\,..$ | 3998 |
| $7n+6\,..$ | $114\,..$ | 3982 |
| $N\,7\ldots$ | $342\,..$ | 11959 |

Impr. classes 4049
Proper together with improper
28396 expected 28418,62

Eight hundred negative determinants.
of the form $-(15n + 13)$

G. I. . . .(91). . . . .(2561)

| | | |
|---|---|---|
| 3 | 43. 163 . . . . . | 2 |
| 5 | 103 . . . . . | 1 |
| 7 | 223. 343. 463 . . . . . | 3 |
| 9 | 283. 643. 823. 883. 1423 . | 5 |
| 11 | 1303 . . . . . | 1 |
| 13 | 2143 . . . . . | 1 |
| 15 | 523. 1123. 1723. 2203. 2683 . . . . . | 5 |
| 17 | 1663. 1783 . . . . . | 2 |
| 19 | 1063. 1543. 3343. 3463 . . | 4 |
| 21 | 1483. 2083. 2503. 4603. 5923 . . . . . | 5 |
| 23 | 4783. 6703 . . . . . | 2 |
| 25 | 5023. 5503 . . . . . | 2 |
| 27 | 2803. 3163. 3643. 3943. 4243. 4363. 4483. 4723. 4903. 5443(*3*). 6043. 6763. 6883. 7723(*3*). 8563. 8803(*3*). 11383 . | 17 |
| 29 | 2383. 3583. 3823. 5743. 8863 . . . . . | 5 |
| 31 | 11863 . . . . . | 1 |
| 33 | 4423. 4663. 5623. 5683. 6163. 6343. 6823. 7603. 8443. 9283. 9403. 9643 . . | 12 |
| 35 | 11503 . . . . . | 1 |
| 39 | 4003. 7243. 7963. 11743. 11923 . . . . . | 5 |
| 41 | 10903 . . . . . | 1 |
| 43 | 8263 . . . . . | 1 |
| 45 | 5323. 5563(*3*). 9043. 9463. 10243. 10663. 10723(*3*). 11083 . . . | 8 |
| 51 | 8623. 9343. 9883. 10303. 11443 . . . . . | 5 |
| 57 | 8923. 9103 . . . . . | 2 |

G. II. . . .(340). . . .(10110)

| | | |
|---|---|---|
| 1. | 13. 28. 58 . . . . . | 3 |
| 2 | 73. 148. 193 . . . . . | 3 |
| 3 | 118. 268. 298. 358. 403 . | 5 |
| 4 | 178. 313. 388. 478. 583 . | 5 |
| 5 | 373. 508. 538. 613. 853. 1093. 1213. 1318 . . . . | 8 |
| 6 | 433. 628. 673. 718. 763. 898. 1003. 1033. 1108. 1138. 1198. 1243. 1588. 1618. 1873 . . . . . | 15 |
| 7 | 703. 733. 778. 838. 1183. 1453. 1948. 2293 . . . . | 8 |
| 8 | 943. 958. 1153. 1348. 1438. 1828. 2113. 2308. 2578. 2878 . . . . . | 10 |

| | | |
|---|---|---|
| 9 | 1018. 1228(*3*). 1363. 1468. 1603. 1843. 1933. 1963. 2023. 2038. 2053. 2188. 2443. 2458. 2563(*3*). 2923. 2998. 3238. 3523. 3628. 3733. 3763. 4348. 4678. 5413 . . . . . | 25 |
| 10 | 1678. 1753. 1858. 2473. 2638. 2743. 2818. 2983. 3028. 3103. 3118. 3508. 4153 . . . . . | 13 |
| 11 | 1693. 1903. 2263. 2623. 2863. 3418. 3703. 3868. 3958. 4918. 5098. 8023. 8143 . . . . . | 13 |
| 12 | 1993. 2098. 2323. 2593. 2713. 2953. 3283. 3313. 3403. 3433. 3778. 3883. 4063. 4258. 4273. 4513. 4843. 5188. 5233. 5758. 5938. 6073. 6238. 8068 . . | 24 |
| 13 | 2218. 2428. 2908. 3373. 3613. 3853. 3898. 4093. 4618. 4933. 5383. 5818. 5878. 6598. 7078. 8383. 8743. 10333. 10543 . . . . | 19 |
| 14 | 3673. 3988. 4078. 4183. 4468. 4948. 5218. 5263. 6103. 6388. 6658. 6673. 7438. 7753. 7858. 8233. 8353. 11113. 11278 . . . . | 19 |
| 15 | 2518. 3148. 3223. 4138. 4813. 5203. 5398. 5653. 5803. 6268. 6403. 6463. 6778. 6988. 7123. 7303. 7363. 7468. 7483. 8053. 8458. 8698. 9013. 9388. 10483. 10588 . . . . . | 26 |
| 16 | 2833. 4993. 6178. 6628. 7183. 7393. 8548. 8578. 9433. 9508. 10558 . . . . | 11 |
| 17 | 3253. 4303. 5308. 5578. 7333. 7558. 8038. 9223. 9703 . . . . . | 9 |
| 18 | 3043. 3793. 4963. 5458. 5998. 6283(*3*). 6418. 6553. 6583(*3*). 6793. 7543. 9298. 9523. 9583. 9748(*3*). 10003. 10018. 10228. 10603. 10753. 10798(*3*). 11698 . . . | 22 |
| 19 | 5158. 5638. 6373. 6733. 7573. 7933. 8983. 9133. 10198. 11173 . . . . . | 10 |
| 20 | 3748. 5143. 6223. 6718. 6898. 7063. 7423. 8098. 8158. 9598. 10063. 10513. 10993. 11428. 11953 . . . | 15 |
| 21 | 4198. 6133. 6508. 6523. 7213. 7318. 7948. 7978. 8518. 9613. 9733. 9868. 10093. 10123. 10363. 10378. 10453. 10828. 11023. 11068. 11263. 11323. 11878 . . . . | 23 |
| 22 | 5113. 5953. 5983. 7663. 7783. 7873. 7903. 7993. 8713. 10273. 10708. 11668. 11758 . . . . . | 13 |
| 23 | 5788. 8278. 8293. 8893. 9973. 10423. 10618 . . . . | 7 |
| 24 | 4798. 6943. 7003. 7108. 8083. 8503. 8683. 8818. 9763. 10843. 10963. 11143. 11203 11353. 11563 . . . . | 15 |
| 25 | 10783. 11548 . . . . . | 2 |
| 26 | 8788. 9778. 9838 . . . . . | 3 |
| 27 | 8203. 11683(*3*) . . . . . | 2 |
| 28 | 9943. 10078. 11038. 11593 . | 4 |
| 29 | 9148 . . . . . | 1 |
| 30 | 9358. 11623 . . . . . | 2 |
| 31 | 11983 . . . . . | 1 |
| 32 | 10183. 10468 . . . . . | 2 |
| 33 | 11338 . . . . . | 1 |
| 40 | 11833 . . . . . | 1 |

G. IV....(320)....(10088)

| | | |
|---|---|---|
| 1 | 88. 133. 253 ..... | 3 |
| 2 | 208. 238. 328. 418. 448. 553. 568. 598. 658. 793. 928 ..... | 11 |
| 3 | 493. 688. 748. 808. 913. 973. 988. 1048. 1258. 1558. 1708. 1978. 2608 .. | 13 |
| 4 | 868. 1078. 1168. 1393. 1408. 1498. 1513(*2*). 1528. 1633. 1738. 1813. 1918(*2*). 2248. 2353. 2368. 2533. 2773. 2788(*2*). 2893. 3088. 3193. 3298. 3448 ..... | 23 |
| 5 | 1273. 1333. 1378. 1573. 1648. 1798. 2158. 2173. 2398. 3133. 3178. 3388. 3928. 3973. 4558. 4873 .. | 16 |
| 6 | 1888. 2068. 2278. 2338. 2413. 2488. 2548. 2653. 2668. 2698. 2758. 2938. 3073. 3208. 3268. 3478. | |
| | 3598. 3658. 3718. 4018. 4033. 4108. 4123. 4168. 4288. 4393. 4438. 4453. 4708. 4753. 4768. 4978. 5083. 5128. 5173. 5293. 5338. 5518. 5548. 5713. 5728. 6028. 6433. 6493. 6913. 7018. 7093. 7228. 7813. 8248. 9823 ..... | 51 |
| 7 | 2008. 3538. 3568. 3838. 4213. 4333. 4498. 4543. 4573. 5038. 5488. 5608. 5833. 5863. 6013. 6193. 6253. 6538. 6568. 6973. 7198. 7408. 9178. 9718. 10858. 10873 ..... | 26 |
| 8 | 2848(*2*). 3013. 3058. 3328. 3358(*2*). 3493. 4228. 4318(*2*). 4633(*2*). 4738. 5008. 5248(*2*). 5428. 5533. 5668. 5908. 6148(*2*). 6208(*2*). 6868(*2*). 6958(*2*). 7033. 7168. 7288. 7588. 7648. 7828. 8308. 8338. 8638(*2*). 8653. 8878(*2*). 9088. 9118(*2*). 9193. 9208. 9538. 10033. 10138. 11533. 11848 ..... | 40 |
| 9 | 3688. 4378. 4528. 4588. 4693. 4828. 5068(*3*). 5773. 6313. 6359. 6643. 6748. 7258. 7513. 7528. 7678. 7693(*3*). 7708. 7738. 7918. 8188. 8218. 8323(*3*). 8488. 8668. 8908. 8998. 9073. 9163(*3*). 9493. 9853. 9898. 9958. 10213. 10288. 10393. 10693. 11158. 11188. 11608. 11803. 11818. 11893 ..... | 43 |
| 10 | 4858. 5053. 5473. 5968. 6058. 6088. 6613. 7153. 7273. 7453. 7498. 7798. 8128. 8173. 8473. 8593. 8608. 8758. 8773. 8833. 8953. 9058. 9238. 9553. 9628. 9793. 9808. 9913. 10108. 10438. 10888. 10978. 11053. 11293. 11398. 11458. 11938 ..... | 37 |
| 11 | 6298. 7618. 7768. 8413. 8728. 10648. 11638. 11653 | 8 |
| 12 | 5353. 5893. 6838. 6928. 7048. 7348. 7633. 7843. 8428. 9028. 9268. 9673(*2*). 10048. 10258. 10348. 10768. 11008. 11218. 11308. 11488. 11578. 11788. 11908. 11998 | 24 |
| 13 | 8938. 9313. 9418. 9448. 9478. 10498. 10678. 10918. 11098. 11518 ..... | 10 |
| 14 | 6478. 7138. 9253. 9658. 9988. 10573. 10633. 10933. 11233. 11728 ..... | 10 |
| 15 | 8368. 10408. 10813. 11413. 11773 ..... | 5 |

G. VIII. . . .(48). . . .(1600)

| | | |
|---|---|---|
| 2 | 1288. 1768. 2128. 2233. 2968. 4048 . . . . . | 6 |
| 3 | 2728. 3553. 3913. 4648. 4888. 5278. 6448 . . . . . | 7 |
| 4 | 3808. 5593. 5698. 5848. 6118. 6328(*2*). 6688(*2*). 6853. 7378. 8398. 8848. 8968. 9328. 9373. 9568. 10168(*2*). 10528(*2*). 10948. 11368. 11713(*2*) | 20 |
| 5 | 4408. 5368. 6808. 9688. 10153. 10738 . . . . . | 6 |
| 6 | 7888. 8533. 9928. 10318. 11473. 11968 . . . . . | 6 |
| 7 | 8113. 11128. 11248 . . . . | 3 |

G. XVI. . . .(1). . . . . (32)

| | | |
|---|---|---|
| 2 | 8008 . . . . . | 1 |

Total G. 2451 Cl. 24391

Irreg. 37 impr. 4075

| | | | $15n+7$,13 | |
|---|---|---|---|---|
| $7n$ . . . . . | 115 . . | 3435 . . . | 230 . . | 6846 |
| $7n+1$ . . | 114 . . | 2988 . . . | 229 . . | 5997 |
| $7n+2$ . . | 114 . . . | 2987 . . . | 228 . . | 5962 |
| $7n+4$ . . | 114 . . | 2994 . . . | 228 . . | 5987 |
| $R\,7$ . . . | 342 . . | 8969 . . . | 685 . . | 17946 |
| $7n+3$ . . | 114 . . | 4005 . . . | 228 . . | 7984 |
| $7n+5$ . . | 114 . . | 3975 . . . | 228 . . | 7973 |
| $7n+6$ . . | 115 . . | 4007 . . . | 229 . . | 7989 |
| $N\,7$ . . . | 343 . . | 11987 . . . | 685 . . | 23946 |
| All . | 800. | 24391 . . | 1600 . . | 48738 |

Quot. min. 0,2349782 from 163,2608

max. 0,7354322 from 11833

Det. of the form

| | | | | | |
|---|---|---|---|---|---|
| $-(15n+13)$ . . | 68 | 305 | 271 | 128 | 27 |
| $-(15n+7)$ . . . | 69 | 316 | 264 | 130 | 21 |
| | 0,3 | 0,4 | 0,5 | 0,6 | |

[Det. in hundred 10000 of the form $15n$]

G. VIII. . . .(3). . . . .(1464)

| | |
|---|---|
| 48 | 999975 |
| 63 | 999945 |
| 72 | 999930 |

G. XVI. . . .(3). . . . .(2224)

| | |
|---|---|
| 39 | 999915 |
| *44 | 999900 |
| 56 | 999990(*2*) |

G. XXXII. .(1). . . . . .(576)
18 999960
104. . . . . .4264
Impr. 592

[Quotient $< \frac{1}{2} <$ $< 1 <$]

| | | | | | |
|---|---|---|---|---|---|
| in hundred | 11 | Det. | 24. | 1. | 56. . 19 |
| | 12 | | 28 | | 51 21 |
| | 13 | | 21 | | 58 21 |
| | 14 | | 27 | | 57 16 |
| | 15 | | 21 | | 62 17 |
| | 16 | | 22 | | 56 22 |
| | 17 | | 29 | | 52 19 |
| | 18 | | 25. | | 57 18 |
| | 19 | | 22 | | 64 14 |
| | 20 | | 21 | | 61 18 |
| | 21 | | 18 | | 62 20 |
| | 22 | | 18 | | 63 19 |
| | 23 | | 19 | | 61 20 |
| | 24 | | 27 | | 55 18 |
| | 25 | | 31 | | 51 18 |
| | 26 | | 26 | | 54 20 |
| | 27 | | 30 | | 51 19 |
| | 28 | | 27 | | 53 20 |
| | 29 | | 23 | | 57 20 |
| | 30 | | 24 | | 58 18 |
| | 117 | | 24 | | 56 20 |
| | 118 | | 20 | | 61 19 |
| | 119 | | 24 | | 61 15 |
| | 120 | | 29 | | 52 19 |

Negative determinants.

| in hundred | | Quotient | | |
|---|---|---|---|---|
| 1 | max. | 1,271998 | from det. | 89 |
| | min. | 0,2626128 | | 58 |
| 2 | | 1,435917 | | 194 |
| | | 0,2349782 | | 163 |

| | | |
|---|---|---|
| 22 | 1,685723 | 2141 |
| | 0,2808228 | 2143 |
| 23 | 1,645848 | 2246 |
| | 0,2923654 | 2293 |
| 24 | 1,479278 | 2369 |
| | 0,2897240 | 2335 |
| 27 | 1,6445315 | 2609 |
| | 0,2895883 | 2683 |
| 28 | 1,5527075 | 2789 |
| | 0,3030216 | 2783 |
| 29 | 1,5778996 | 2834 |
| | 0,2974718 | 2893 |
| 30 | 1,604748 | 2939 |
| | 0,2936893 | 2968 |
| 91 | 1,684117 | 9026 |
| | 0,2835515 | 9067 |
| 92 | 1,586777 | 9176 |
| | 0,2717044 | 9157 |
| 93 | 1,660820 | 9281 |
| | 0,2699414 | 9277 |
| 94 | 1,518533 | 9371 |
| | 0,2893063 | 9367 |
| 95 | 1,729662 | 9434 |
| | 0,3287980 | 9472 |
| 96 | 1,689400 | 9539 |
| | 0,3269906 | 9577 |
| 97 | 1,707014 | 9686 |
| | 0,2440986 | 9667 |
| 99 | 1,650848 | 9869 |
| | 0,2420048 | 9823 |
| 100 | 1,702214 | 9974 |
| | 0,2808862 | 9937 |
| 117 | 1,6654535 | 11681 |
| | 0,2964744 | 11650 |
| 118 | 1,810938 | 11714 |
| | 0,294621 | 11797 |
| 119 | 1,579112 | 11864 |
| | 0,2846194 | 11863 |
| 120 | 1,5326965 | 11921 |
| | 0,3287433 | 11992 |

Positive determinants.

Hundred 1.
Square determinants excluded: 10.

G. I. . . .(12)
1 2. 5. 13. 17. 29. 41. 53. 61. 73. 89. 97
3 37
G. II. . . .(51)
1 3. 6. 7. 8. 10. 11. 12. 14. 18. 19. 20. 21. 22. 23. 26. 27. 28. 31. 32. 33. 38. 43. 44. 45. 46. 47. 50. 52. 54. 57. 58. 59. 62. 65. 67. 68. 69. 71. 74. 76. 77. 83. 85. 86. 92. 93. 94. 98
2 34. 82
3 79
G. IV. . .(27)
1 15. 24. 30. 35. 39. 40. 42. 48. 51. 55. 56. 60. 63. 66. 70. 72. 75. 78. 80. 84. 87. 88. 90. 91. 95. 96. 99

Hundred 2.
Excluded: 4.

G. I. . . .(11)
1 109. 113. 125. 137. 149. 157. 173. 181. 193
3 101. 197
G. II. . . .(41)
1 103. 106. 107. 108. 116. 117. 118. 122. 124. 127. 128. 129. 131. 133. 134. 139. 142. 151. 153. 158. 161. 162. 163. 164. 166. 167. 172. 174. 177. 179. 185. 188. 191. 199
2 145. 146. 178. 194
3 141. 148. 189
G. IV. . .(40)
1 102. 104. 105. 110. 111. 112. 114. 115. 119. 123. 126. 130. 132. 135. 136. 138. 140. 143. 147. 152. 154. 155. 156. 159. 160. 165. 170. 171. 175. 176. 180. 182. 183. 184. 186. 187. 190. 192. 198. 200
G. VIII. . .(4)
1 120. 150. 168. 195

Hundred 3.

G. I
1 233. 241. 277. 281. 293
3 229. 257. 269
G. II
1 201. 202. 206. 211. 212. 213. 214. 217. 218. 229. 236. 237. 239. 242. 243. 244. 245. 249. 250. 251. 253. 261. 262. 263. 265. 268. 271. 278. 283. 284. 292. 297. 298
2 205. 221. 274

3 223. 226. 254. 291
G. IV
1 203. 204. 207. 215. 216. 222. 228. 230. 232. 234. 238. 246. 247. 248. 252. 258. 259. 260. 266. 267. 270. 272. 273. 275. 276. 279. 282. 285. 286. 287. 290. 294. 295. 296. 299. 300
2 219. 220. 224. 288
3 235
G. VIII
1 210. 231. 240. 255. 264. 280

Hundred 9.

G. I. . . . .(7)
1 809. 821. 853. 857. 881
3 829. 877
G. II. . . .(32)
1 801. 811. 823. 827. 833. 838. 844. 845. 849. 859. 862. 863. 865. 869. 873. 878. 883. 886. 887. 889. 893
2 802. 818. 866
3 813. 837. 839. 842. 892
5 817
6 898
14 (841)
G. IV. . . .(52)
1 803. 804. 805. 806. 807. 808. 810. 814. 815. 822. 824. 825. 826. 830. 831. 832. 834. 835. 836. 843. 846. 847. 848. 850. 851. 852. 854. 856. 860. 861. 864. 867. 868. 871. 872. 875. 879. 882. 885
2 812. 820. 828. 876. 884. 890. 891. 896. 897
3 874. 894. 895. 899
G. VIII. . .(8)
1 816. 819. 855. 858. 888
2 870. 880. (900)
G. XVI. . .(1)
1 840

Hundred 10.

G. I. . . .(6). . . . .(8)
1 929. 937. 941. 953. 977
3 997
G. II. . .(38). . . .(130)
1 907. 908. 911. 913. 917. 919. 921. 922. 926. 932. 947. 949. 956. 958. 964. 965. 967. 971. 972. 974. 981. 983. 989. 991. 998
2 914
3 905. 909. 916. 925. 933. 934. 973. 985. 993
5 982
6 901

[15 961]
G. IV....(40)...(224)
1 902. 918. 923. 927. 928. 931. 938. 942. 944. 945. 946. 948. 950. 951. 954. 955. 957. 962. 968. 969. 970. 976. 978. 980. 986. 988. 995. 996. 999. 1000
2 939. 943. 959. 963. 979. 992
3 906. 940
4 904. 994
G. VIII...(16)...(144)
1 903. 912. 915. 920. 924. 930. 935. 936. 952. 966. 975. 984. 987. 990
2 910. 960

Total 370

G. I
1 313. 317
3 349. 373. 389. 397. 557. 677. 701. 709. 733. 757. 761
5 401
7 577
G. II
1 301. 302. 307. 309. 311. 314.
2 305
3 316. 321. 325. 326
5 727
G. IV
1 303. 304. 308. 310. 318. 319. 320. 327
2 306. 322. 323
G. VIII
1 312. 315

# TABLE
# FOR
# CIRCLE DIVISION.

| | | | |
|---|---|---|---|
| 2 | 5 | 3 | 5 |
| 4 | 17 | 5 | 13 |
| 6 | 37 | 7 | 5 . 5 |
| 8 | 5 . 13 | 9 | 41 |
| 10 | 101 | 11 | 61 |
| 12 | 5 . 29 | 13 | 5 . 17 |
| 14 | 197 | 15 | 113 |
| 17 | 5 . 29 | 18 | 5 . 5 . 13 |
| 19 | 181 | 21 | 13 . 17 |
| 22 | 5 . 97 | 23 | 5 . 53 |
| 27 | 5 . 73 | 28 | 5 . 157 |
| 30 | 17 . 53 | 31 | 13 . 37 |
| 32 | 5 . 5 . 41 | 33 | 5 . 109 |
| 34 | 13 . 89 | 37 | 5 . 137 |
| 38 | 5 . 17 . 17 | 41 | 29 . 29 |
| 43 | 5 . 5 . 37 | 44 | 13 . 149 |
| 46 | 29 . 73 | 47 | 5 . 13 . 17 |
| 50 | 41 . 61 | 55 | 17 . 89 |
| 57 | 5 . 5 . 5 . 13 | 68 | 5 . 5 . 5 . 37 |
| 70 | 13 . 13 . 29 | 72 | 5 . 17 . 61 |
| 73 | 5 . 13 . 41 | 75 | 29 . 97 |
| 76 | 53 . 109 | 80 | 37 . 173 |
| 81 | 17 . 193 | 83 | 5 . 13 . 53 |
| 91 | 41 . 101 | 93 | 5 . 5 . 173 |
| 98 | 5 . 17 . 113 | 99 | 13 . 13 . 29 |
| 100 | 73 . 137 | 105 | 37 . 149 |
| 111 | 61 . 101 | 112 | 5 . 13 . 193 |
| 117 | 5 . 37 . 37 | 119 | 73 . 97 |
| 123 | 5 . 17 . 89 | 128 | 5 . 29 . 113 |
| 129 | 53 . 157 | 132 | 5 . 5 . 17 . 41 |
| 133 | 5 . 29 . 61 | 142 | 5 . 37 . 109 |
| 157 | 5 . 5 . 17 . 29 | 162 | 5 . 29 . 181 |
| 172 | 5 . 61 . 97 | 173 | 5 . 41 . 73 |
| 174 | 13 . 17 . 137 | 182 | 5 . 5 . 5 . 5 . 53 |
| 183 | 5 . 17 . 197 | 185 | 109 . 157 |
| 191 | 17 . 29 . 37 | 192 | 5 . 73 . 101 |
| 193 | 5 . 5 . 5 . 149 | 200 | 13 . 17 . 181 |
| 211 | 113 . 197 | 212 | 5 . 89 . 101 |
| 216 | 13 . 37 . 97 | 233 | 5 . 61 . 89 |
| 237 | 5 . 41 . 137 | 239 | 13 . 13 . 13 . 13 |
| 242 | 5 . 13 . 17 . 53 | 251 | 17 . 17 . 109 |
| 253 | 5 . 37 . 173 | 255 | 13 . 41 . 61 |

| | | | |
|---|---|---|---|
| 265 | 13 . 37 . 73 | 268 | 5 . 5 . 13 . 13 . 17 |
| 278 | 5 . 13 . 29 . 41 | 293 | 5 . 5 . 17 . 101 |
| 294 | 13 . 61 . 109 | 302 | 5 . 17 . 29 37 |
| 307 | 5 . 5 . 5 . 13 . 29 | 313 | 5 . 97 . 101 |
| 319 | 17 . 41 . 73 | 327 | 5 . 17 . 17 . 37 |
| 342 | 5 . 149 . 157 | 343 | 5 . 5 . 13 . 181 |
| 360 | 29 . 41 . 109 | 378 | 5 . 17 . 41 . 41 |
| 394 | 29 . 53 . 101 | 401 | 37 . 41 . 53 |
| 403 | 5 . 109 . 149 | 408 | 5 . 13 . 13 . 197 |
| 411 | 13 . 73 . 89 | 437 | 5 . 13 . 13 . 113 |
| 438 | 5 . 17 . 37 . 61 | 443 | 5 . 5 . 5 . 5 . 157 |
| 447 | 5 . 13 . 29 . 53 | 463 | 5 . 13 . 17 . 97 |
| 467 | 5 . 113 . 193 | 499 | 13 . 61 . 157 |
| 500 | 53 . 53 . 89 | 507 | 5 . 5 . 53 . 97 |
| 512 | 5 . 13 . 37 . 109 | 515 | 13 . 101 . 101 |
| 524 | 37 . 41 . 181 | 538 | 5 . 13 . 61 . 73 |
| 557 | 5 . 5 . 5 . 17 . 73 | 560 | 53 . 61 . 97 |
| 568 | 5 . 5 . 5 . 29 . 89 | 577 | 5 . 13 . 13 . 197 |
| 599 | 17 . 61 . 173 | 606 | 13 . 13 . 41 . 53 |
| 616 | 13 . 17 . 17 . 101 | 621 | 29 . 61 . 109 |
| 657 | 5 . 5 . 89 . 97 | 660 | 37 . 61 . 193 |
| 682 | 5 . 5 . 5 . 61 . 61 | 684 | 13 . 17 . 29 . 73 |
| 693 | 5 . 5 . 5 . 17 . 113 | 697 | 5 . 13 . 37 . 101 |
| 701 | 17 . 97 . 149 | 743 | 5 . 5 . 61 . 181 |
| 746 | 13 . 13 . 37 . 89 | 757 | 5 . 5 . 73 . 157 |
| 772 | 5 . 13 . 53 . 173 | 776 | 73 . 73 . 113 |
| 785 | 13 . 137 . 173 | 798 | 5 . 13 . 97 . 101 |
| 818 | 5 . 5 . 5 . 53 . 101 | 829 | 17 . 17 . 29 . 41 |
| 853 | 5 . 13 . 29 . 193 | 882 | 5 . 5 . 29 . 29 . 37 |
| 905 | 13 . 17 . 17 . 109 | 919 | 37 . 101 . 113 |
| 922 | 5 . 17 . 73 . 137 | 924 | 53 . 89 . 181 |
| 931 | 13 . 17 . 37 . 53 | 945 | 29 . 89 . 173 |
| 948 | 5 . 17 . 97 . 109 | 993 | 5 . 5 . 13 . 37 . 41 |
| 999 | 17 . 149 . 197 | 1032 | 5 . 5 . 13 . 29 . 113 |
| 1057 | 5 . 5 . 5 . 41 . 109 | 1067 | 5 . 17 . 37 . 181 |
| 1068 | 5 . 5 . 5 . 5 . 5 . 5 . 73 | 1087 | 5 . 13 . 61 . 149 |
| 1118 | 5 . 5 . 17 . 17 . 173 | 1123 | 5 . 13 . 89 . 109 |
| 1143 | 5 . 5 . 17 . 29 . 53 | 1148 | 5 . 29 . 61 . 149 |
| 1196 | 53 . 137 . 197 | 1228 | 5 . 17 . 113 . 157 |
| 1239 | 41 . 97 . 193 | 1270 | 61 . 137 . 193 |
| 1303 | 5 . 41 . 41 . 101 | 1341 | 73 . 109 . 113 |
| 1385 | 41 . 149 . 157 | 1393 | 5 . 5 . 197 . 197 |
| 1407 | 5 . 5 . 17 . 17 . 137 | 1432 | 5 . 5 . 5 . 5 . 17 . 193 |
| 1433 | 5 . 29 . 73 . 97 | 1467 | 5 . 29 . 41 . 181 |
| 1477 | 5 . 13 . 97 . 173 | 1560 | 17 . 37 . 53 . 73 |
| 1567 | 5 . 41 . 53 . 113 | 1568 | 5 . 5 . 5 . 13 . 17 . 89 |
| 1597 | 5 . 37 . 61 . 113 | 1607 | 5 . 5 . 13 . 29 . 137 |
| 1636 | 17 . 29 . 61 . 89 | 1744 | 137 . 149 . 149 |

| | | | |
|---|---|---|---|
| 1772 | 5 . 17 . 17 . 41 . 53 | 1818 | 5 . 5 . 5 . 137 . 193 |
| 1823 | 5 . 17 . 113 . 173 | 1832 | 5 . 5 . 17 . 53 . 149 |
| 1893 | 5 . 5 . 13 . 37 . 149 | 1918 | 5 . 5 . 37 . 41 . 97 |
| 1929 | 13 . 13 . 101 . 109 | 1955 | 13 . 29 . 37 . 137 |
| 1984 | 13 . 29 . 53 . 197 | 2010 | 13 . 17 . 101 . 181 |
| 2013 | 5 . 29 . 89 . 157 | 2018 | 5 . 5 . 29 . 41 . 137 |
| 2042 | 5 . 29 . 149 . 193 | 2059 | 13 . 41 . 41 . 97 |
| 2153 | 5 . 13 . 181 197 | 2163 | 5 . 13 . 17 . 29 . 73 |
| 2191 | 89 . 149 . 181 | 2309 | 13 . 53 . 53 . 73 |
| 2350 | 17 . 17 . 97 . 197 | 2428 | 5 . 41 . 149 . 193 |
| 2436 | 13 . 13 . 13 . 37 . 73 | 2515 | 101 . 173 . 181 |
| 2540 | 13 . 29 . 109 . 157 | 2547 | 5 . 37 . 89 . 197 |
| 2621 | 13 . 37 . 37 . 193 | 2673 | 5 . 13 . 17 . 53 . 61 |
| 2697 | 5 41 . 113 . 157 | 2738 | 5 . 13 . 29 . 41 . 97 |
| 2801 | 17 . 29 . 73 . 109 | 2818 | 5 . 5 . 5 . 17 . 37 . 101 |
| 2917 | 5 . 13 . 29 . 37 . 61 | 2943 | 5 . 5 . 5 . 5 . 13 . 13 . 41 |
| 3039 | 17 . 61 . 61 . 73 | 3112 | 5 . 13 . 13 . 73 . 157 |
| 3141 | 13 . 13 . 17 . 17 . 101 | 3149 | 17 . 29 . 89 . 113 |
| 3166 | 17 . 41 . 73 . 197 | 3207 | 5 . 5 . 29 . 41 . 173 |
| 3323 | 5 . 13 . 29 . 29 . 101 | 3362 | 5 . 13 . 17 . 53 . 193 |
| 3405 | 29 . 29 . 61 . 113 | 3458 | 5 . 73 . 181 . 181 |
| 3521 | 29 . 37 . 53 . 109 | 3532 | 5 . 5 . 17 . 149 . 197 |
| 3583 | 5 . 13 . 17 . 37 . 157 | 3740 | 41 . 41 . 53 . 157 |
| 3782 | 5 . 5 . 29 . 109 . 181 | 3793 | 5 . 5 . 53 . 61 . 89 |
| 3957 | 5 . 5 . 13 . 13 . 17 . 109 | 4193 | 5 . 5 . 5 . 5 . 5 . 29 . 97 |
| 4217 | 5 . 13 . 29 . 53 . 89 | 4232 | 5 . 5 . 41 . 101 . 173 |
| 4246 | 13 . 17 . 29 . 29 . 97 | 4327 | 5 . 89 . 109 . 193 |
| 4484 | 17 . 89 . 97 . 137 | 4535 | 17 . 53 . 101 . 113 |
| 4545 | 13 . 37 . 109 . 197 | 4581 | 13 . 53 . 97 . 157 |
| 4594 | 13 . 17 . 29 . 37 . 89 | 4662 | 5 . 13 . 13 . 17 . 17 . 89 |
| 4747 | 5 . 17 . 41 . 53 . 61 | 4906 | 13 . 53 . 181 . 193 |
| 4937 | 5 . 73 . 173 . 193 | 4952 | 5 . 37 . 41 . 53 . 61 |
| 5052 | 5 . 13 . 41 . 61 . 157 | 5087 | 5 . 17 . 29 . 29 . 181 |
| 5257 | 5 . 5 . 13 . 17 . 41 . 61 | 5283 | 5 . 13 . 17 . 73 . 173 |
| 5357 | 5 . 5 . 61 . 97 . 97 | 5443 | 5 . 5 . 5 . 5 . 137 . 173 |
| 5507 | 5 . 5 . 13 . 13 . 37 . 97 | 5648 | 5 . 17 . 53 . 73 . 97 |
| 5667 | 5 . 29 . 37 . 41 . 73 | 5701 | 29 . 53 . 97 . 109 |
| 5767 | 5 . 13 . 17 . 101 . 149 | 5928 | 5 . 29 . 29 . 61 . 137 |
| 5962 | 5 . 13 . 29 . 109 . 173 | 6065 | 17 . 53 . 137 . 149 |
| 6107 | 5 . 5 . 17 . 17 . 29 89 | 6118 | 5 . 5 . 13 . 41 . 53 . 53 |
| 6252 | 5 . 17 . 29 . 101 . 157 | 6481 | 17 . 37 . 173 . 193 |
| 6682 | 5 . 5 . 5 . 29 . 109 . 113 | 6898 | 5 . 13 . 17 . 17 . 17 . 149 |
| 6908 | 5 . 13 . 73 . 89 113 | 6943 | 5 . 5 . 5 . 29 . 61 . 109 |
| 6962 | 5 . 37 . 37 . 73 . 97 | 7093 | 5 . 5 . 13 . 17 . 29 . 157 |
| 7161 | 17 . 101 . 109 . 137 | 7443 | 5 . 5 . 5 . 37 . 53 . 113 |
| 7697 | 5 . 17 . 29 . 61 . 197 | 7782 | 5 . 5 . 13 . 17 . 97 . 113 |
| 8224 | 13 . 17 . 29 . 61 . 173 | 8307 | 5 . 5 . 5 . 5 . 5 . 61 . 181 |
| 8368 | 5 . 5 . 17 . 37 . 61 . 73 | | |

| | |
|---|---|
| 8393 | 5 . 5 . 13 . 29 . 37 . 101 |
| 8457 | 5 . 5 . 53 . 137 . 197 |
| 8578 | 5 . 37 . 41 . 89 . 109 |
| 9133 | 5 . 17 . 37 . 89 . 149 |
| 9152 | 5 . 29 . 41 . 73 . 193 |
| 9193 | 5 . 5 . 5 . 5 . 17 . 41 . 97 |
| 9298 | 5 . 41 . 53 . 73 . 109 |
| 9431 | 17 . 97 . 149 . 181 |
| 9466 | 29 . 37 . 37 . 37 . 61 |
| 9667 | 5 . 13 . 41 . 89 . 197 |
| 9703 | 5 . 13 . 13 . 17 . 29 . 113 |
| 9762 | 5 . 17 . 37 . 157 . 193 |
| 9872 | 5 . 13 . 13 . 29 . 41 . 97 |
| 9901 | 13 . 13 . 29 . 73 . 137 |
| 10298 | 5 . 17 . 61 . 113 . 181 |
| 10312 | 5 . 29 . 53 . 101 . 137 |
| 10833 | 5 . 17 . 41 . 113 . 149 |
| 11018 | 5 . 5 . 157 . 157 . 197 |
| 11471 | 13 . 17 . 41 . 53 . 137 |
| 11981 | 13 . 17 . 41 . 89 . 89 |
| 12332 | 5 . 5 . 13 . 41 . 101 . 113 |
| 12433 | 5 . 13 . 61 . 101 . 193 |
| 12882 | 5 . 5 . 17 . 37 . 61 . 173 |
| 12943 | 5 . 5 . 5 . 5 . 13 . 13 . 13 . 61 |
| 13043 | 5 . 5 . 17 . 17 . 61 . 193 |
| 13068 | 5 . 5 . 5 . 53 . 149 . 173 |
| 13241 | 29 . 101 . 173 . 173 |
| 13252 | 5 . 13 . 13 . 37 . 41 . 137 |
| 13545 | 17 . 17 . 53 . 53 . 113 |
| 13918 | 5 . 5 . 13 . 37 . 89 . 181 |
| 14140 | 17 . 29 . 37 . 97 . 113 |
| 14318 | 5 . 5 . 5 . 13 . 17 . 41 . 181 |
| 14573 | 5 . 17 . 73 . 109 . 157 |
| 14646 | 13 . 37 . 41 . 73 . 149 |
| 14773 | 5 . 13 . 13 . 29 . 61 . 73 |
| 14942 | 5 . 13 . 13 . 37 . 37 . 193 |
| 14958 | 5 . 13 . 101 . 173 . 197 |
| 15075 | 13 . 17 . 53 . 89 . 109 |
| 16513 | 5 . 29 . 53 . 113 . 157 |
| 16928 | 5 . 17 . 109 . 157 . 197 |
| 17191 | 13 . 17 . 61 . 97 . 113 |
| 17557 | 5 . 5 . 5 . 17 . 29 . 41 . 61 |
| 17766 | 13 . 37 . 73 . 89 . 101 |
| 17923 | 5 . 61 . 61 . 89 . 97 |
| 18258 | 5 . 29 . 97 . 137 . 173 |
| 18432 | 5 . 5 . 5 . 17 . 29 . 37 . 149 |
| 18543 | 5 . 5 . 13 . 17 . 29 . 29 . 37 |
| 19123 | 5 . 29 . 37 . 173 . 197 |
| 19283 | 5 . 37 . 37 . 157 . 173 |
| 19326 | 13 . 29 . 61 . 109 . 149 |
| 19534 | 13 . 13 . 13 . 29 . 53 . 113 |
| 19653 | 5 . 37 . 61 . 109 . 157 |
| 19703 | 5 . 13 . 13 . 29 . 89 . 89 |
| 19902 | 5 . 73 . 89 . 89 . 137 |
| 19911 | 13 . 17 . 29 . 157 . 197 |
| 20080 | 13 . 29 . 61 . 89 . 197 |
| 20457 | 5 . 5 . 13 . 29 . 149 . 149 |
| 21124 | 29 . 41 . 53 . 73 . 97 |
| 21705 | 13 . 17 . 61 . 101 . 173 |
| 21907 | 5 . 5 . 29 . 29 . 101 . 113 |
| 22008 | 5 . 41 . 109 . 149 . 157 |
| 22157 | 5 . 5 . 13 . 37 . 137 . 149 |
| 22231 | 29 . 37 . 41 . 41 . 137 |
| 24263 | 5 . 13 . 17 . 41 . 73 . 89 |
| 24331 | 13 . 17 . 89 . 101 . 149 |
| 24778 | 5 . 29 . 149 . 157 . 181 |
| 24816 | 17 . 17 . 61 . 181 . 193 |
| 25462 | 5 . 13 . 17 . 37 . 101 . 157 |
| 25523 | 5 . 53 . 73 . 113 . 149 |
| 25683 | 5 . 13 . 17 . 17 . 97 . 181 |
| 25793 | 5 . 5 . 17 . 29 . 137 . 197 |
| 25943 | 5 . 5 . 5 . 13 . 29 . 37 . 193 |
| 26018 | 5 . 5 . 13 . 97 . 109 . 197 |
| 27493 | 5 . 5 . 17 . 17 . 17 . 17 . 181 |
| 28205 | 13 . 29 . 73 . 97 . 149 |
| 28322 | 5 . 13 . 13 . 13 . 13 . 41 . 137 |
| 28862 | 5 . 17 . 17 . 53 . 73 . 149 |
| 29757 | 5 . 5 . 41 . 61 . 73 . 97 |
| 30027 | 5 . 29 . 89 . 181 . 193 |
| 30103 | 5 . 13 . 17 . 41 . 73 . 137 |
| 30383 | 5 . 17 . 17 . 37 . 89 . 97 |
| 31752 | 5 . 17 . 17 . 37 . 109 . 173 |
| 32258 | 5 . 13 . 37 . 41 . 61 . 173 |
| 32406 | 17 . 17 . 17 . 37 . 53 . 109 |
| 32807 | 5 . 5 . 5 . 13 . 61 . 61 . 89 |
| 32885 | 13 . 13 . 109 . 149 . 197 |
| 32973 | 5 . 13 . 37 . 37 . 41 . 149 |
| 33307 | 5 . 5 . 5 . 5 . 17 . 53 . 197 |
| 34208 | 5 . 13 . 13 . 17 . 29 . 53 . 53 |
| 34367 | 5 . 13 . 37 . 41 . 53 . 113 |
| 35857 | 5 . 5 . 17 . 61 . 137 . 181 |
| 36673 | 5 . 17 . 29 . 37 . 73 . 101 |
| 37057 | 5 . 5 . 5 . 5 . 73 . 101 . 149 |
| 37448 | 5 . 13 . 13 . 53 . 173 . 181 |
| 37770 | 13 . 17 . 29 . 41 . 61 . 89 |
| 38326 | 29 . 37 . 41 . 173 . 193 |

| | | | |
|---|---|---|---|
| 38807 | 5 . 5 . 5 . 17 . 37 . 61 . 157 | 39082 | 5 . 5 . 41 . 73 . 137 . 149 |
| 39307 | 5 . 5 . 5 . 13 . 13 . 13 . 29 97 | 39818 | 5 . 5 . 5 . 5 . 17 . 37 . 37 . 109 |
| 40188 | 5 . 13 . 37 . 61 . 101 . 109 | 40515 | 17 . 53 . 61 . 109 . 137 |
| 40568 | 5 . 5 . 5 . 13 . 53 . 97 . 197 | 41187 | 5 . 17 . 37 . 37 . 37 . 197 |
| 41319 | 13 . 41 . 101 . 101 . 157 | 41688 | 5 . 17 . 41 . 53 . 97 . 97 |
| 42658 | 5 . 13 . 13 . 97 . 149 . 149 | 42932 | 5 . 5 . 5 . 29 . 29 . 89 . 197 |
| 43633 | 5 . 13 . 29 . 41 . 109 . 113 | 43932 | 5 . 5 . 5 . 5 . 13 . 17 . 89 . 157 |
| 44179 | 13 . 13 . 13 . 17 . 17 . 29 . 53 | 44507 | 5 . 5 . 13 . 113 . 149 . 181 |
| 44733 | 5 . 89 . 101 . 113 . 197 | 45050 | 13 . 41 . 109 . 181 . 193 |
| 45068 | 5 . 5 . 5 . 41 . 61 . 73 . 89 | 46444 | 13 . 41 . 149 . 157 . 173 |
| 46617 | 5 . 53 . 137 . 173 . 173 | 47403 | 5 . 13 . 29 . 37 . 89 . 181 |
| 47783 | 5 . 13 . 17 . 53 . 101 . 193 | 48187 | 5 . 97 . 101 . 137 . 173 |
| 48737 | 5 . 29 . 29 . 53 . 73 . 73 | 49083 | 5 . 13 . 17 . 73 . 109 . 137 |
| 50052 | 5 . 17 . 41 . 41 . 89 . 197 | 51115 | 17 . 17 . 17 . 29 . 53 . 173 |
| 51387 | 5 . 17 . 37 . 53 . 89 . 89 | 51412 | 5 . 17 . 61 . 61 . 61 . 137 |
| 51917 | 5 . 13 . 37 . 53 . 97 . 109 | 52571 | 37 . 41 . 61 . 109 . 137 |
| 54193 | 5 . 5 . 5 . 5 . 5 . 41 . 73 . 157 | 54358 | 5 . 13 . 29 . 73 . 109 . 197 |
| 54507 | 5 . 5 . 37 . 53 . 157 . 193 | 57532 | 5 . 5 . 17 . 41 . 41 . 41 . 113 |
| 66347 | 5 . 13 . 13 . 17 . 37 . 41 . 101 | 67333 | 5 . 17 . 53 . 61 73 . 113 |
| 67852 | 5 . 13 . 37 . 89 . 137 . 157 | 68463 | 5 . 13 . 17 . 113 . 137 . 137 |
| 71564 | 37 . 61 . 97 . 149 . 157 | 71700 | 13 . 29 . 37 . 41 . 89 . 101 |
| 72662 | 5 . 13 . 17 . 29 . 37 . 61 . 73 | 74043 | 5 . 5 . 13 . 37 . 37 . 61 . 101 |
| 75382 | 5 . 5 . 13 . 17 . 73 . 73 . 193 | 78629 | 13 . 17 . 41 . 41 . 53 . 157 |
| 80593 | 5 . 5 . 17 . 17 . 17 . 137 . 193 | 80802 | 5 . 37 . 41 . 53 . 109 . 149 |
| 81141 | 13 . 61 . 137 . 157 . 193 | 81749 | 13 . 17 . 17 . 53 . 97 . 173 |
| 83071 | 37 . 61 . 89 . 89 . 193 | 83247 | 5 . 13 . 13 . 13 . 29 . 73 . 149 |
| 84141 | 13 . 29 . 29 . 41 . 53 . 149 | 85353 | 5 . 13 . 17 . 37 . 41 . 41 . 53 |
| 86143 | 5 . 5 . 13 . 17 . 61 . 101 . 109 | 88668 | 5 . 5 . 13 . 17 . 73 . 101 . 193 |
| 88699 | 29 . 53 . 89 . 149 . 193 | 88733 | 5 . 13 . 41 . 97 . 97 . 157 |
| 88868 | 5 . 5 . 29 . 37 . 37 . 73 . 109 | 89361 | 29 . 37 . 137 . 157 . 173 |
| 89471 | 13 . 13 . 41 . 41 . 73 . 193 | 90212 | 5 . 13 . 37 . 89 . 193 . 197 |
| 90657 | 5 . 5 . 13 . 17 . 61 . 89 . 137 | 93020 | 13 . 13 . 17 . 17 . 29 . 41 . 149 |
| 93197 | 5 . 37 . 53 . 53 . 61 . 137 | 99557 | 5 . 5 . 5 . 5 . 41 . 41 . 53 . 89 |
| 99893 | 5 . 5 . 29 . 181 . 193 . 197 | 101343 | 5 . 5 . 13 . 29 . 41 . 97 . 137 |
| 102163 | 5 . 37 . 41 . 41 . 97 . 173 | 104818 | 5 . 5 . 5 . 5 . 17 . 29 181 . 197 |
| 106242 | 5 . 53 . 53 . 73 . 101 . 109 | 109637 | 5 . 13 . 17 . 29 . 37 . 37 . 137 |
| 112595 | 17 . 29 . 41 . 53 . 61 . 97 | 112782 | 5 . 5 . 17 . 37 . 41 . 109 . 181 |
| 114669 | 17 . 37 . 53 . 53 . 61 . 61 | 117251 | 41 . 97 . 101 . 109 . 157 |
| 117307 | 5 . 5 . 5 . 13 . 149 . 157 . 181 | 117372 | 5 . 13 . 17 . 17 . 53 . 101 . 137 |
| 128482 | 5 . 5 . 17 . 29 . 89 . 101 . 149 | 129553 | 5 . 13 . 13 . 17 . 61 . 61 . 157 |
| 133749 | 13 . 13 . 37 . 53 . 137 . 197 | 136293 | 5 . 5 . 17 . 41 . 53 . 89 . 113 |
| 136404 | 13 . 17 . 29 . 97 . 173 . 173 | 137717 | 5 . 13 . 53 . 89 . 157 . 197 |
| 137883 | 5 . 13 . 17 . 29 . 29 . 53 . 193 | 141743 | 5 . 5 . 89 . 149 . 157 . 193 |
| 143382 | 5 . 5 . 13 . 13 . 17 . 17 . 113 . 149 | 145046 | 13 . 29 . 37 . 101 . 109 . 137 |
| 145231 | 13 . 37 . 37 . 41 . 97 . 149 | 148158 | 5 . 53 . 61 . 61 . 113 . 197 |
| 148582 | 5 . 5 . 13 . 53 . 73 . 97 . 181 | 150522 | 5 . 13 . 17 . 29 . 37 . 97 . 197 |
| 155317 | 5 . 41 . 61 . 73 . 73 . 181 | 157308 | 5 . 13 . 29 . 29 . 41 . 61 . 181 |
| 157318 | 5 . 5 . 5 . 5 . 5 . 5 . 13 . 37 37 . 89 | 159772 | 5 . 37 . 53 . 101 . 149 . 173 |

| | |
|---|---|
| 160590 | 29 . 29 . 29 . 89 . 109 . 109 |
| 162014 | 13 . 17 . 73 . 89 . 101 . 181 |
| 174118 | 5 . 5 . 17 . 41 . 89 . 113 . 173 |
| 180107 | 5 . 5 . 13 . 29 . 97 . 113 . 157 |
| 181743 | 5 . 5 . 17 . 17 . 73 . 173 . 181 |
| 161832 | 5 . 5 . 13 . 13 . 29 . 37 . 53 . 109 |
| 173932 | 5 . 5 . 5 . 5 . 5 . 13 . 73 . 101 . 101 |
| 177144 | 17 . 29 . 73 . 89 97 . 101 |
| 181343 | 5 . 5 . 17 . 37 . 53 . 109 . 181 |

| | |
|---|---|
| 184133 | 5 . 29 . 73 . 101 . 101 . 157 |
| 189782 | 5 . 5 . 13 . 61 . 89 . 137 . 149 |
| 190393 | 5 . 5 . 13 . 13 . 137 . 173 . 181 |
| 191407 | 5 . 5 . 13 . 13 . 17 . 37 . 61 . 113 |
| 191807 | 5 . 5 . 5 . 13 . 17 . 41 . 109 . 149 |
| 194708 | 5 . 5 . 29 . 41 . 73 . 101 . 173 |
| 201106 | 17 . 17 . 61 . 89 . 149 . 173 |
| 208048 | 5 . 53 . 89 . 109 . 113 . 149 |
| 210195 | 61 . 113 . 137 . 149 . 157 |
| 210943 | 5 . 5 . 5 . 13 . 37 . 37 . 73 . 137 |
| 211765 | 13 . 17 . 53 . 89 . 137 . 157 |
| 216676 | 13 . 29 . 41 . 97 . 173 . 181 |
| 219602 | 5 . 17 . 17 . 53 . 53 . 109 . 109 |
| 221382 | 5 . 5 . 13 . 73 . 101 . 113 . 181 |
| 228068 | 5 . 5 . 5 . 17 . 29 . 61 . 101 . 137 |
| 232643 | 5 . 5 . 13 . 13 . 13 . 41 . 61 . 197 |
| 236151 | 17 . 17 . 41 . 89 . 137 . 193 |
| 247643 | 5 . 5 . 17 . 29 . 73 . 173 . 197 |
| 249501 | 13 . 53 . 53 . 61 . 89 . 157 |
| 251103 | 5 . 13 . 17 . 17 . 89 . 109 . 173 |
| 256638 | 5 . 13 . 29 . 37 . 61 . 113 . 137 |
| 260359 | 13 . 13 . 17 . 41 . 53 . 61 . 89 |
| 262433 | 5 . 17 . 29 . 29 . 53 . 61 . 149 |
| 263317 | 5 . 17 . 17 . 17 . 89 . 101 . 157 |
| 263557 | 5 . 5 . 5 . 13 . 37 . 41 . 73 . 193 |
| 265842 | 5 . 13 . 17 . 17 . 101 . 193 . 193 |
| 267657 | 5 . 5 . 41 . 41 . 61 . 89 . 157 |
| 281897 | 5 . 13 . 29 . 37 . 37 . 89 . 173 |
| 286018 | 5 . 5 . 13 . 13 . 53 . 53 . 61 . 113 |
| 287228 | 5 . 17 . 17 . 29 . 73 . 149 . 181 |
| 289038 | 5 . 17 . 17 . 37 . 89 . 97 . 181 |
| 292362 | 5 . 13 . 17 . 41 . 61 . 157 . 197 |
| 298307 | 5 . 5 . 5 . 5 . 41 . 53 . 181 . 181 |
| 307939 | 13 . 29 . 61 . 101 . 137 . 149 |
| 309070 | 13 . 29 . 101 . 113 . 149 . 149 |
| 320078 | 5 . 13 . 41 . 61 . 73 . 89 . 97 |
| 322392 | 5 . 13 . 17 . 41 . 89 . 149 . 173 |
| 330182 | 5 . 5 . 5 . 5 . 5 . 13 . 29 . 37 . 41 . 61 |
| 331068 | 5 . 5 . 5 . 5 . 53 . 101 . 181 . 181 |
| 383807 | 5 . 5 . 5 . 13 . 37 . 73 . 97 . 173 |
| 385692 | 5 . 13 . 17 . 61 . 89 . 137 . 181 |

| | |
|---|---|
| 389163 | 5 . 13 29 . 41 . 89 . 101 . 109 |
| 390112 | 5 . 13 . 17 . 17 . 17 . 17 . 17 . 17 . 97 |
| 403639 | 29 . 29 . 37 . 97 . 137 . 197 |
| 409557 | 5 . 5 . 5 . 5 . 13 . 13 . 73 . 73 . 149 |
| 411787 | 5 . 17 . 17 . 53 . 97 . 101 . 113 |
| 418048 | 5 . 97 . 97 . 109 . 173 . 197 |
| 444753 | 5 . 53 . 109 . 113 . 157 . 193 |
| 447342 | 5 . 17 . 29 . 41 . 97 . 137 . 149 |
| 464307 | 5 . 5 . 5 . 29 . 37 . 73 . 101 . 109 |
| 465525 | 13 . 13 . 29 . 89 . 97 . 113 . 197 |
| 465694 | 13 . 13 . 17 . 29 . 73 . 181 . 197 |
| 478707 | 5 . 5 . 13 . 13 . 13 . 17 . 41 . 41 . 73 |
| 485298 | 5 . 13 . 13 13 . 13 . 29 . 29 . 37 . 53 |
| 494607 | 5 . 5 . 29 . 89 . 101 . 137 . 137 |
| 500150 | 41 . 61 . 73 . 73 . 137 . 137 |
| 508929 | 13 . 13 . 37 . 53 . 53 . 73 . 101 |
| 518734 | 13 . 17 . 37 . 37 . 53 . 97 . 173 |
| 520463 | 5 . 13 . 41 . 61 . 73 . 101 . 113 |
| 534568 | 5 . 5 . 5 . 13 . 89 . 89 . 149 . 149 |
| 538275 | 17 . 61 . 73 . 97 . 109 . 181 |
| 548630 | 37 . 41 . 89 . 109 . 113 . 181 |
| 566793 | 5 . 5 . 17 . 29 . 29 . 41 . 97 . 113 |
| 567923 | 5 . 13 . 17 . 17 . 17 . 41 . 109 . 113 |
| 571459 | 13 . 13 . 13 . 13 . 17 . 37 . 61 . 149 |
| 586455 | 29 . 41 . 73 . 89 . 113 . 197 |
| 606325 | 13 . 37 . 97 . 137 . 149 . 193 |
| 607533 | 5 . 17 . 29 . 73 . 97 . 97 . 109 |
| 617427 | 5 . 13 . 13 . 17 . 29 . 53 . 89 . 97 |
| 623888 | 5 . 13 . 37 . 41 . 113 . 181 . 193 |
| 627391 | 41 . 41 . 53 . 113 . 113 . 173 |
| 662843 | 5 . 5 . 17 . 113 . 137 . 173 . 193 |
| 672717 | 5 . 89 . 97 . 157 . 173 . 193 |
| 683982 | 5 . 5 . 17 . 29 . 61 . 61 . 101 . 101 |
| 700107 | 5 . 5 . 13 . 17 . 41 . 53 . 137 . 149 |
| 703175 | 13 . 17 . 17 . 29 . 97 . 149 . 157 |
| 704683 | 5 . 13 . 29 . 61 . 97 . 113 . 197 |
| 721068 | 5 . 5 . 5 . 5 . 17 . 29 . 109 . 113 . 137 |
| 780262 | 5 . 17 . 17 . 29 . 37 . 41 . 61 . 157 |
| 783568 | 5 . 5 . 5 . 5 . 17 . 29 . 101 . 109 . 181 |
| 791532 | 5 . 5 . 53 . 89 . 149 . 181 . 197 |
| 793921 | 17 . 17 . 17 . 29 . 73 . 157 . 193 |
| 812447 | 5 . 29 . 29 . 41 . 89 . 137 . 157 |
| 832902 | 5 . 13 . 13 . 13 . 17 . 109 . 173 . 197 |
| 848871 | 29 . 53 . 73 . 113 . 157 . 181 |
| 899168 | 5 . 5 . 17 . 37 . 53 . 73 . 97 . 137 |
| 907567 | 5 . 29 . 37 . 41 . 89 . 109 . 193 |
| 911111 | 17 . 41 . 101 . 173 . 173 . 197 |
| 936513 | 5 . 37 . 37 . 41 89 . 97 . 181 |

| | |
|---|---|
| 1000193 | 5 . 5 . 5 . 29 . 53 . 101 . 149 . 173 |
| 1010027 | 5 . 13 . 61 . 89 . 89 . 109 . 149 |
| 1024240 | 37 . 61 . 109 . 157 . 157 . 173 |
| 1031675 | 13 . 13 . 17 . 53 . 73 . 113 . 157 . 197 |
| 1049433 | 5 . 13 . 61 . 89 . 89 . 89 . 197 |
| 1059193 | 5 . 5 . 5 . 5 . 13 . 13 . 13 . 37 . 61 . 181 |
| 1067157 | 5 . 5 . 41 . 113 . 157 . 173 . 181 |
| 1068182 | 5 . 5 . 5 . 17 . 17 . 41 . 61 . 73 . 173 |
| 1083493 | 5 . 5 . 13 . 61 . 61 . 61 . 73 . 109 |
| 1089593 | 5 . 5 . 61 . 89 . 149 . 149 . 197 |
| 1131527 | 5 . 41 . 53 . 53 . 73 . 97 . 157 |
| 1139557 | 5 . 5 . 5 . 5 . 5 . 5 . 5 . 17 . 37 . 73 . 181 |
| 1143007 | 5 . 5 . 13 . 41 . 61 . 73 . 101 . 109 |
| 1197943 | 5 . 5 . 5 . 5 . 5 . 17 . 17 . 37 . 109 . 197 |
| 1264557 | 5 . 5 . 5 . 5 . 5 . 13 . 13 . 29 . 53 . 197 |
| 1306143 | 5 . 5 . 29 . 37 . 37 . 61 . 73 . 193 |
| 1351742 | 5 . 17 . 53 . 109 . 137 . 157 . 173 |
| 1373307 | 5 . 5 . 5 . 5 . 17 . 29 . 101 . 157 . 193 |
| 1387203 | 5 . 29 . 41 . 41 . 113 . 181 . 193 |
| 1402232 | 5 . 5 . 17 . 37 . 41 . 113 . 137 . 197 |
| 1413443 | 5 . 5 . 13 . 29 . 37 . 41 . 89 . 157 |
| 1477034 | 37 . 37 . 41 . 53 . 53 . 101 . 137 |
| 1518057 | 5 . 5 . 5 . 13 . 13 . 41 . 61 . 113 . 193 |
| 1528649 | 13 . 37 . 53 . 61 . 61 . 109 . 113 |
| 1615463 | 5 . 13 . 17 . 37 . 53 . 73 . 73 . 113 |
| 1618855 | 17 . 29 . 37 . 53 . 89 . 97 . 157 |
| 1635786 | 29 . 29 . 41 . 89 . 89 . 97 . 101 |
| 1664957 | 5 . 5 . 13 . 37 . 73 . 89 . 113 . 157 |
| 1750507 | 5 . 5 . 13 . 53 . 53 . 89 . 109 . 173 |
| 1766693 | 5 . 5 . 5 . 5 . 5 . 5 . 29 . 97 . 157 . 173 |
| 1824257 | 5 . 5 . 17 . 29 . 97 . 109 . 113 . 113 |
| 1909461 | 13 . 13 . 17 . 17 . 41 . 53 . 89 . 193 |
| 1954207 | 5 . 5 . 13 . 61 . 61 . 89 . 113 . 157 |
| 1984933 | 5 . 17 . 37 . 37 . 37 . 53 . 89 . 97 |
| 2036069 | 17 . 41 . 41 . 61 . 61 . 101 . 193 |
| 2050706 | 13 . 17 . 17 . 17 . 41 . 53 . 157 . 193 |
| 2052057 | 5 . 5 . 5 . 5 . 5 . 5 . 5 . 17 . 29 . 73 . 97 . 193 |
| 2126007 | 5 . 5 . 17 . 29 . 29 . 181 . 181 . 193 |
| 2277387 | 5 . 13 . 29 . 29 . 53 . 89 . 89 . 113 |
| 2298668 | 5 . 5 . 13 . 13 . 17 . 17 . 29 . 37 . 37 . 109 |
| 2343692 | 5 . 17 . 41 . 41 . 61 . 73 . 89 . 97 |
| 2353918 | 5 . 5 . 13 . 17 . 29 . 29 . 61 . 113 . 173 |
| 2379723 | 5 . 13 . 29 . 37 . 53 . 53 . 97 . 149 |
| 2457057 | 5 . 5 . 5 . 5 . 13 . 17 . 41 . 53 . 89 . 113 |
| 2471717 | 5 . 37 . 41 . 109 . 137 . 149 . 181 |
| 2475918 | 5 . 5 . 17 . 53 . 61 . 157 . 157 . 181 |
| 2478328 | 5 . 13 . 29 . 37 . 89 . 97 . 101 . 101 |
| 2484968 | 5 . 5 . 13 . 61 . 97 . 113 . 157 . 181 |

| | |
|---|---|
| 2680168 | 5 . 5 . 29 . 61 . 73 . 109 . 137 . 149 |
| 2733307 | 5 . 5 . 5 . 5 . 5 . 5 . 13 . 13 . 13 . 17 . 37 . 173 |
| 2809305 | 13 . 17 . 29 . 37 . 37 . 61 . 73 . 101 |
| 2923783 | 5 . 13 . 17 . 37 . 41 . 109 . 149 . 157 |
| 2959007 | 5 . 5 . 17 . 37 . 97 . 101 . 157 . 181 |
| 3014557 | 5 . 5 . 5 . 5 . 5 . 5 . 5 . 5 . 41 . 53 . 53 . 101 |
| 3025001 | 13 . 13 . 17 . 17 . 61 . 73 . 109 . 193 |
| 3136570 | 13 . 13 . 29 . 37 . 53 . 61 . 97 . 173 |
| 3139557 | 5 . 5 . 5 . 5 . 5 . 5 . 13 . 29 . 73 . 73 . 157 |
| 3272693 | 5 . 5 . 5 . 13 . 37 . 41 . 101 . 137 . 157 |
| 3370437 | 5 . 13 . 13 . 13 . 13 . 41 . 73 . 97 . 137 |
| 3449051 | 13 . 13 . 13 . 53 . 61 . 89 . 97 . 97 |
| 3637197 | 5 . 13 . 17 . 29 . 61 . 89 . 193 . 197 |
| 3800438 | 5 . 13 . 29 . 29 . 97 . 101 . 149 . 181 |
| 3801448 | 5 . 29 . 37 . 53 . 61 . 73 . 101 . 113 |
| 3815076 | 13 . 13 . 17 . 37 . 53 . 109 . 137 . 173 |
| 3894873 | 5 . 13 . 13 . 37 . 89 . 101 . 137 . 197 |
| 3911450 | 29 . 29 . 41 . 97 . 137 . 173 . 193 |
| 3931663 | 5 . 13 . 29 . 37 . 41 . 109 . 137 . 181 |
| 4000300 | 13 . 13 . 13 . 17 . 17 . 73 . 137 . 181 . 181 |
| 4079486 | 13 . 17 . 17 . 53 . 61 . 73 . 137 . 137 |
| 4218932 | 5 . 5 . 5 . 5 . 29 . 41 . 41 . 61 . 61 . 157 |
| 4466678 | 5 . 13 . 17 . 73 . 97 . 109 . 149 . 157 |
| 4650839 | 17 . 17 . 89 . 113 . 137 . 157 . 173 |
| 4697282 | 5 . 5 . 13 . 113 . 113 . 137 . 197 . 197 |
| 4751232 | 5 . 5 . 13 . 17 . 29 . 73 . 97 . 101 . 197 |
| 4773557 | 5 . 5 . 5 . 29 . 29 . 41 . 113 . 149 . 157 |
| 5033696 | 13 . 17 . 37 . 37 . 89 . 89 . 97 . 109 |

| | |
|---|---|
| 5982670 | 13 . 13 . 13 . 17 . 41 . 53 . 53 . 53 . 157 |
| 6151956 | 13 . 17 . 29 . 29 . 73 . 97 . 113 . 149 |
| 6208047 | 5 . 17 . 17 . 17 . 17 . 29 . 41 . 41 . 197 . 197 |
| 6225244 | 29 . 37 . 41 . 53 . 53 . 53 . 61 . 97 |
| 6315768 | 5 . 5 . 17 . 17 . 53 . 61 . 73 . 149 . 157 |
| 6356150. | 13 . 29 . 37 . 37 . 61 . 61 . 109 . 193 |
| 6367252 | 5 . 13 . 17 . 29 . 29 . 61 . 73 . 97 . 101 |
| 6656382 | 5 . 5 . 13 . 29 . 41 . 41 . 137 . 137 . 149 |
| 6817837 | 5 . 17 . 17 . 53 . 61 . 149 . 173 . 193 |
| 6829610 | 13 . 17 . 17 . 53 . 61 . 101 . 193 . 197 |
| 6981694 | 13 . 41 . 97 . 137 . 181 . 193 . 197 |
| 7138478 | 5 . 29 . 37 . 41 . 73 . 89 . 181 . 197 |
| 7620661 | 5 . 17 . 37 . 73 . 101 . 101 . 137 . 181 |
| 7691443 | 5 . 5 . 5 . 37 . 53 . 97 . 101 . 109 . 113 |
| 8082212 | 5 . 13 . 17 . 17 . 37 . 53 . 97 . 101 . 181 |
| 8571779 | 13 . 13 . 29 . 41 . 73 . 101 . 137 . 181 |
| 8809432 | 5 . 5 . 5 . 13 . 89 . 101 . 149 . 181 . 197 |
| 9407318 | 5 . 5 . 5 . 5 . 5 . 5 . 37 . 41 . 53 . 73 . 193 |

| | |
|---|---|
| 9548768 | 5 . 5 . 13 . 13 . 17 . 41 . 53 . 61 . 61 . 157 |
| 9614382 | 5 . 5 . 29 . 37 . 53 . 61 . 61 . 101 . 173 |
| 9639557 | 5 . 5 . 5 . 5 . 5 . 5 . 13 . 17 . 17 . 53 . 109 . 137 |
| 9689961 | 13 . 29 . 29 . 37 . 61 . 113 . 113 . 149 |
| 10328193 | 5 . 5 . 5 . 13 . 17 . 29 . 53 . 53 . 137 . 173 |
| 10669731 | 17 . 89 . 97 . 101 . 101 . 193 . 197 |
| 11131086 | 13 . 13 . 17 . 17 . 37 . 61 . 73 . 89 . 173 |
| 12477035 | 17 . 17 . 17 . 29 . 29 . 29 . 37 . 97 . 181 |
| 12514913 | 5 . 13 . 41 . 53 . 137 . 149 . 157 . 173 |
| 12750353 | 5 . 13 . 17 . 17 . 41 . 61 . 73 . 137 . 173 |
| 14698000 | 13 . 13 . 17 . 17 . 29 . 61 . 97 . 149 . 173 |
| 15165443 | 5 . 5 . 5 . 5 . 37 . 53 . 61 . 97 . 101 . 157 |
| 15986082 | 5 . 5 . 13 . 17 . 109 . 109 . 137 . 157 . 181 |
| 16317267 | 5 . 13 . 17 . 17 . 61 . 61 . 101 . 109 . 173 |
| 18378313 | 5 . 13 . 13 . 17 . 37 . 61 . 137 . 193 . 197 |
| 18975991 | 13 . 17 . 17 . 17 . 53 . 61 . 89 . 97 . 101 |
| 20198495 | 13 . 17 . 41 . 89 . 101 . 101 . 137 . 181 |
| 22866693 | 5 . 5 . 5 . 5 . 41 . 61 . 73 . 101 . 113 . 197 |
| 23747457 | 5 . 5 . 17 . 17 . 17 . 17 . 17 . 17 . 37 . 73 . 173 |
| 24208144 | 29 . 29 . 29 . 37 . 37 . 53 . 61 . 61 . 89 |
| 24280807 | 5 . 5 . 5 . 5 . 13 . 13 . 17 . 53 . 109 . 157 . 181 |
| 24310918 | 5 . 5 . 13 . 13 . 37 . 41 . 53 . 89 . 113 . 173 |
| 31011557 | 5 . 5 . 5 . 13 . 17 . 61 . 97 . 109 . 137 . 197 |
| 32944452 | 5 . 13 . 13 . 29 . 29 . 41 . 53 . 53 . 89 . 149 |
| 34436768 | 5 . 5 . 17 . 61 . 97 . 101 . 137 . 173 . 197 |
| 34602875 | 13 . 17 . 17 . 29 . 37 . 53 . 113 . 137 . 181 |
| 45500682 | 5 . 5 . 5 . 37 . 53 . 53 . 61 . 89 . 149 . 197 |
| 53365057 | 5 . 5 . 5 . 13 . 37 . 89 . 97 . 101 . 157 . 173 |
| 58305593 | 5 . 5 . 13 . 17 . 37 . 37 . 101 . 109 . 137 . 149 |
| 75505943 | 5 . 5 . 5 . 37 . 37 . 53 . 89 . 137 . 149 . 173 |
| 95665578 | 5 . 13 . 37 . 41 . 73 . 181 . 181 . 197 . 197 |
| 111530944 | 13 . 13 . 13 . 13 . 13 . 17 . 37 . 37 . 53 . 157 . 173 |
| 121042733 | 5 . 17 . 41 . 73 . 97 . 97 . 101 . 157 . 193 |
| 160007778 | 5 . 13 . 13 . 17 . 17 . 29 . 29 . 73 . 73 . 149 . 157 |
| 167207057 | 5 . 5 . 5 . 5 . 17 . 17 . 17 . 29 . 73 . 109 . 109 . 181 |
| 168623905 | 13 . 13 13 . 13 . 17 . 29 . 29 . 37 . 89 . 97 . 109 |
| 185507821 | 13 . 13 . 17 . 29 . 29 . 53 . 61 . 101 . 113 . 193 |
| 193788912 | 5 . 13 . 17 . 17 . 37 . 37 . 37 . 53 . 73 . 101 . 101 |
| 201229582 | 5 . 5 . 13 . 13 . 17 . 17 . 17 . 17 . 17 . 53 . 97 . 101 |
| 211823957 | 5 . 5 . 17 . 17 . 53 . 101 . 137 . 149 . 157 . 181 |
| 284862638 | 5 . 13 . 17 . 17 . 17 . 17 . 29 . 29 . 41 . 41 . 97 . 109 |
| 299252491 | 13 . 29 . 37 . 97 . 109 . 109 . 113 . 157 |
| 317742693 | 5 . 5 . 5 . 13 . 29 . 41 . 73 . 89 . 137 . 149 . 197 |
| 327012132 | 5 . 5 . 13 . 17 . 17 . 29 . 89 . 109 . 149 . 157 . 173 |
| 599832943 | 5 . 5 . 5 . 5 . 13 . 17 . 29 . 37 . 37 . 41 . 73 97 . 113 |
| 830426722 | 5 . 13 . 13 . 61 . 97 . 149 . 157 . 173 . 173 . 197 |
| 1112115023 | 5 . 17 . 17 . 61 . 73 . 113 . 157 . 173 . 173 . 181 |
| 1282794079 | 13 . 17 . 29 . 29 . 73 . 89 . 97 . 113 . 181 . 197 . 197 |

| | |
|---|---|
| 2189376182 | 5 . 5 . 5 . 17 . 17 . 29 . 29 . 53 . 61 . 61 . 89 . 89 . 101 |
| 2971354082 | 5 . 5 . 13 . 17 . 29 . 41 . 53 . 53 . 113 . 149 . 157 . 181 |
| 3955080927 | 5 . 13 . 17 . 17 . 17 . 17 . 53 . 53 . 61 . 61 . 101 . 149 . 173 . 197 |
| 8193535810 | 13 . 13 . 29 . 29 . 61 . 109 . 109 . 137 . 157 . 157 . 193 |
| 14033378718 | 5 . 5 . 13 . 13 . 17 . 17 . 61 . 61 . 61 . 61 . 73 . 73 . 157 . 181 |

| | |
|---|---|
| 5 | 2. 3. 7 |
| 13 | 5. 8. 18. 57. 239 |
| 17 | 4. 13. 21. 38. 47. 268 |
| 29 | 12. 17. 41. 70. 99. 157. 307 |
| 37 | 6. 31. 43. 68. 117. 191. 302. 327. 882. 18543* |
| 41 | 9. 32. 73. 132. 278. 378. 829. 993. 2943 |
| 53 | 23. 30. 83. 182. 242. 401. 447. 606. 931. 1143*. 1772. 6118. 34208. 44179. 85353. 485298 |
| 61 | 11. 50. 72. 133. 255. 438. 682. 2673. 2917. 4747*. 4952. 5257. 9466. 12943. 17557. 114669. 330182 |
| 73 | 27. 46. 173. 265. 319. 538. 557. 684. 1068. 1560*. 2163. 2309. 2436. 3039. 5667. 8368. 14773. 48737. 72662. 478707* |
| 89 | 34. 55. 123. 233. 411. 500. 568. 746. 1568. 1636*. 3793. 4217. 4594. 4662. 6107. 11981. 19703. 24263. 32807. 37770*. 45068. 51387. 99557. 157318. 260359. 24208144 |
| 97 | 22. 75. 119. 172. 216. 463. 507. 560. 657. 1433*. 1918. 2059. 2738. 4193. 4246. 5357. 5507. 5648. 6962. 9193*. 9872. 17923. 21124. 29757. 30383. 39307. 41688. 112595. 320078. 390112*. 617427. 1984933. 2343692. 3449051. 6225244 |
| 101 | 10. 91. 111. 192. 212. 293. 313. 394. 515. 616*. 697. 798. 818. 1303. 2818. 3141. 3323. 8393. 17766. 36673*. 66347. 71700. 74043. 173932. 177144. 508929. 683982. 1635786. 2478328. 2809305*. 3014557. 6367252. 18975991. 193788912. 201229582. 2189376182 |
| 109 | 33. 76. 142. 251. 294. 360. 512. 621. 905. 948*. 1057. 1123. 1929. 2801. 3521. 3957. 5701. 6943. 8578. 9298*. 15075. 32406. 39818. 40188. 51917. 86143. 88868. 106242. 160590. 161832*. 219602. 389163. 464307. 607533. 1083493. 1143007. 2298668. 5033696. 168623905. 284862638* |
| 113 | 15. 98. 128. 437. 693. 776. 919. 1032. 1341. 1567*. 1597. 3149. 3405. 4535. 6682. 6908. 7443. 7782. 9703. 12332*. 13545. 14140. 17191. 19534. 21907. 34367. 43633. 57532. 67333. 136293*. 191407. 286018. 411787. 520463. 566793. 567923. 1528649. 1615463. 1824257. 2277387*. 2457057. 3801448. 7691443. 599832943 |
| 137 | 37. 100. 174. 237. 922. 1407. 1607. 1955. 2018. 4484*. 5928. 7161. 9901. 10312. 11471. 13252. 19902. 22231. 28322. 30103*. 40517. 49083. 51412. 52571. 68463. 90657. 93197. 101343. 109637. 117372*. 145046. 210943. 228068. 256638. 494607. 500150. 721068. 899168. 1477034. 3370434*. 4079486. 9639557 |

| | |
|---|---|
| 149 | 44. 105. 193. 403. 701. 1087. 1148. 1744. 1832. 1893*. 5767. 6065. 6898. 9133. 10833. 14646. 18432. 19326. 20457. 22157*. 24331. 25523. 28205. 28862. 32973. 37057. 39082. 42658. 80802. 83247*. 84141. 93020. 128482. 143382. 145231. 189782. 191807. 208048. 262433. 307939*. 309070. 409557. 447342. 534568. 571459. 700107. 1010027. 2379723. 2680168. 6151956*. 6656382. 9689961. 32944452. 58305593 |
| 157 | 28. 129. 185. 342. 443. 499. 757. 1228. 1385. 2013*. 2540. 2697. 3112. 3583. 3740. 4581. 5052. 6252. 7093. 14573*. 16513. 19653. 22008. 25462. 38807. 41319. 43932. 54193. 67852. 71564*. 78629. 88733. 117251. 129553. 180107. 184133. 210195. 211765. 249503. 263317*. 267657. 703175. 780262. 812447. 1131527. 1413443. 1618855. 1664957. 1954207. 2923783*. 3139557. 3272693. 4218932. 4466678. 4773557. 5982670. 6315768. 9548768. 15165443. 160007778*. 299252491 |
| 173 | 80. 93. 253. 599. 772. 785. 945. 1118. 1477. 1823*. 3207. 4232. 5283. 5443. 5962. 8224. 12882. 13068. 13241. 18258*. 19283. 21705. 31752. 32258. 46444. 46617. 48187. 51115. 81749. 89361*. 102163. 136404. 159772. 174118. 194718. 201106. 251103. 281897. 322392. 383807*. 518734. 627391. 1000193. 1024240. 1068182. 1351742. 1750507. 1766693. 2353918. 2733307*. 3136570. 3815076. 4650839. 9614382. 10328193. 11131086. 12514913. 12750353. 14698000. 16317267*. 23747457. 24310918. 53365057. 75505943. 111530944. 327012132 |
| 181 | 19. 162. 200. 343. 524. 743. 924. 1067. 1467. 2010*. 2191. 2515. 3458. 3782. 5087. 8307. 9431. 10298. 13918. 14318*. 24778. 25683. 27493. 35857. 37448. 44507. 47403. 112782. 117307. 148582*. 155317. 157308. 162014. 181343. 181743. 190393. 216676. 221382. 287228. 289038*. 298307. 331068. 385692. 538275. 548630. 783568. 848871. 936513. 1059193. 1067157*. 1139557. 2471717. 2475918. 2484968. 2959007. 3800438. 3931663. 4000300. 7620662. 8082212*. 8571779. 12477035. 15986092. 20198495. 24280807. 34602875. 167207057. 211823957. 1112115023. 2971354082*. 14033378718 |
| 193 | 81. 112. 467. 660. 853. 1239. 1270. 1432. 1818. 2042*. 2428. 2621. 3362. 4327. 4906. 4937. 6481. 9152. 9762. 12433*. 13043. 14942. 24816. 25943. 30027. 28326. 45050. 47783. 54507. 75382*. 80593. 81141. 83071. 88668. 88699. 89471. 137883. 141743. 236151. 263557*. 265842. 444753. 606385. 623888. 662843. 672717. 793921. 907567. 1306143. 1373307*. 1387203. 1518057. 1909461. 2036069. 2050706. 2052057. 2126007. 3025001. 3911450. 6356150*. 6817837. 9407318. 121042733. 185507821. 8193535810 |
| 197 | 14. 183. 211. 408. 577. 999. 1196. 1393. 1984. 2153*. 2350. 2547. 3166. 3532. 4545. 7697. 8457. 9667. 11018. 14958*. 16928. 19123. 19911. 20080. 25793. 26018. 32885. 33307. 40568. 41187*. 42932. 44733. 50052. 54358. 90212. 99893. 104818. 133749. 137719. 148158*. 150522. 232643. 247643. 292362. 403639. 418048. 465525. 465694. 586455. 704683*. 791532. 832902. 911111. 1031675. 1049433. 1089593. 1197943. 1264557. 1402232. 3637197*. 3894873. 4697282. 4751232. 6208047. 6829610. 6981694. 7138478. 8809432. 10669731. 18378313*. 22866693. 31011557. 34436768. 45500682. 95665578. 317742693. 830426722. 1282794079. 3955080927 |

**Factorable $aa + 4$.**

| | | | |
|---|---|---|---|
| 1 | 5 | 3 | 13 |
| 5 | 29 | 7 | 53 |
| 9 | 5 . 17 | 11 | 5 . 5 . 5 |
| 13 | 173 | 19 | 5 . 73 |
| 21 | 5 . 89 | 23 | 13 . 41 |
| 25 | 17 . 37 | 29 | 5 . 13 . 13 |
| 31 | 5 . 193 | 39 | 5 . 5 . 61 |
| 43 | 17 . 109 | 49 | 5 . 13 . 37 |
| 53 | 29 . 97 | 59 | 5 . 17 . 41 |
| 61 | 5 . 5 . 149 | 63 | 29 . 137 |
| 81 | 5 . 13 . 101 | 83 | 61 . 113 |
| 99 | 5 . 37 . 53 | 101 | 5 . 13 . 157 |
| 111 | 5 . 5 . 17 . 29 | 121 | 5 . 29 . 101 |
| 127 | 13 . 17 . 73 | 141 | 5 . 41 . 97 |
| 143 | 113 . 181 | 161 | 5 . 5 . 17 . 61 |
| 169 | 5 . 29 . 197 | 179 | 5 . 13 . 17 . 29 |
| 199 | 5 . 89 . 89 | 205 | 13 . 53 . 61 |
| 211 | 5 . 5 . 13 . 137 | 213 | 17 . 17 . 157 |
| 219 | 5 . 53 . 181 | 237 | 13 . 29 . 149 |
| 247 | 17 . 37 . 97 | 261 | 5 . 5 . 5 . 5 . 109 |
| 283 | 13 . 61 . 101 | 309 | 5 . 13 . 13 . 113 |
| 311 | 5 . 5 . 53 . 73 | 335 | 13 . 89 . 97 |
| 359 | 5 . 149 . 173 | 393 | 13 . 109 . 109 |
| 417 | 17 . 53 . 193 | 419 | 5 . 13 . 37 . 73 |
| 469 | 5 . 29 . 37 . 41 | 485 | 17 . 101 . 137 |
| 527 | 29 . 61 . 157 | 535 | 17 . 113 . 149 |
| 611 | 5 . 5 . 109 . 137 | 679 | 5 . 13 . 41 . 173 |
| 691 | 5 . 29 . 37 . 89 | 705 | 13 . 13 . 17 . 173 |
| 749 | 5 . 29 . 53 . 73 | 759 | 5 . 29 . 29 . 137 |
| 761 | 5 . 5 . 5 . 41 . 113 | 829 | 5 . 13 . 97 . 109 |
| 841 | 5 . 17 . 53 . 157 | 943 | 17 . 17 . 17 . 181 |
| 961 | 5 . 5 . 17 . 41 . 53 | 1011 | 5 . 5 . 5 . 13 . 17 . 37 |
| 1043 | 13 . 13 . 41 . 157 | 1047 | 89 . 109 . 113 |
| 1089 | 5 . 5 . 13 . 41 . 89 | 1131 | 5 . 17 . 101 . 149 |
| 1159 | 5 . 37 . 53 . 137 | 1305 | 97 . 97 . 181 |
| 1351 | 5 . 17 . 109 . 197 | 1371 | 5 . 41 . 53 . 173 |
| 1381 | 5 . 13 . 13 . 37 . 61 | 1499 | 5 . 41 . 97 . 113 |
| 1581 | 5 . 41 . 89 . 137 | 1745 | 13 . 29 . 41 . 197 |
| 1801 | 5 . 37 . 89 . 197 | 1899 | 5 . 37 . 101 . 193 |
| 2025 | 13 . 29 . 73 . 149 | 2343 | 13 . 37 . 101 . 113 |
| 2355 | 17 . 41 . 73 . 109 | 2441 | 5 . 13 . 29 . 29 . 109 |

| | |
|---|---|
| 2677 | 17 . 17 . 137 . 181 |
| 3039 | 5 . 5 . 13 . 157 . 181 |
| 3339 | 5 . 5 . 41 . 73 . 149 |

| | |
|---|---|
| 3351 | 5 . 13 . 13 . 97 . 137 |
| 3377 | 13 . 61 . 73 . 197 |
| 3717 | 29 . 53 . 89 . 101 |
| 3749 | 5 . 17 . 37 . 41 . 109 |
| 4021 | 5 . 17 . 37 . 53 . 97 |
| 4123 | 17 . 29 . 29 . 29 . 41 |
| 4215 | 13 . 73 . 97 . 113 |
| 4761 | 5 . 5 . 5 . 13 . 13 . 29 . 37 |
| 4989 | 5 . 5 . 5 . 13 . 17 . 17 . 53 |
| 5041 | 5 . 13 . 13 . 17 . 29 . 61 |
| 5567 | 13 . 17 . 17 . 73 . 113 |
| 5573 | 29 . 61 . 97 . 181 |
| 5717 | 13 . 13 . 41 . 53 . 89 |
| 5821 | 5 . 13 . 37 . 73 . 193 |
| 6061 | 5 . 5 . 13 . 17 . 61 . 109 |
| 6261 | 5 . 5 . 5 . 53 . 61 . 97 |
| 6989 | 5 . 5 . 5 . 53 . 73 . 101 |
| 7319 | 5 . 17 . 73 . 89 . 97 |
| 7745 | 13 . 13 . 37 . 53 . 181 |
| 8049 | 5 . 17 . 53 . 73 . 197 |
| 8579 | 5 . 29 . 53 . 61 . 157 |
| 8879 | 5 . 29 . 41 . 89 . 149 |
| 9801 | 5 . 17 . 73 . 113 . 137 |
| 9817 | 17 . 37 . 37 . 41 . 101 |
| 9947 | 73 . 89 . 97 . 157 |
| 10039 | 5 . 5 . 13 . 17 . 17 . 29 . 37 |
| 12383 | 37 . 109 . 193 . 197 |
| 12605 | 17 . 37 . 41 . 61 . 101 |
| 12815 | 13 . 13 . 41 . 137 . 173 |
| 13251 | 5 . 17 . 101 . 113 . 181 |
| 13489 | 5 . 5 . 5 . 5 . 5 . 5 . 5 . 17 . 137 |
| 13507 | 17 . 17 . 41 . 89 . 173 |
| 14261 | 5 . 5 . 5 . 89 . 101 . 181 |
| 14901 | 5 . 13 . 13 . 13 . 17 . 29 . 41 |
| 16041 | 5 . 73 . 89 . 89 . 89 |
| 20511 | 5 . 5 . 5 . 13 . 17 . 97 . 157 |
| 20769 | 5 . 29 . 37 . 37 . 41 . 53 |
| 20875 | 13 . 29 . 53 . 113 . 193 |
| 21139 | 5 . 5 . 17 . 37 . 157 . 181 |
| 21161 | 5 . 5 . 13 . 89 . 113 . 137 |
| 21189 | 5 . 5 . 37 . 61 . 73 . 109 |
| 22805 | 13 . 17 . 89 . 137 . 193 |
| 23311 | 5 . 5 . 29 . 41 . 101 . 181 |
| 23901 | 5 . 29 . 137 . 149 . 193 |
| 23915 | 13 . 29 . 73 . 113 . 173 |
| 25689 | 5 . 5 . 29 . 41 . 149 . 149 |
| 27355 | 13 . 37 . 53 . 149 . 197 |
| 27411 | 5 . 5 . 41 . 61 . 61 . 197 |

| | |
|---|---|
| 27429 | 5 . 17 . 29 . 37 . 73 . 113 |
| 27611 | 5 . 5 . 41 . 61 . 89 . 137 |
| 29169 | 5 . 13 . 29 . 41 . 101 . 109 |
| 29691 | 5 . 17 . 17 . 29 . 109 . 193 |
| 29929 | 5 . 13 . 17 . 61 . 97 . 137 |
| 31351 | 5 . 37 . 149 . 181 . 197 |
| 32003 | 13 . 17 . 17 . 41 . 61 . 109 |
| 32139 | 5 . 5 . 13 . 17 . 73 . 197 |
| 32239 | 5 . 5 . 5 . 5 . 5 . 37 . 89 . 101 |
| 37579 | 5 . 17 . 29 . 41 . 89 . 157 |
| 44301 | 5 . 13 . 37 . 53 . 89 . 173. |
| 47389 | 5 . 5 . 53 . 97 . 101 . 173 |
| 47761 | 5 . 5 . 5 . 5 . 17 . 17 . 73 . 173 |
| 47829 | 5 . 17 . 37 . 41 . 113 . 157 |
| 49813 | 13 . 53 . 101 . 181 . 197 |
| 57989 | 5 . 5 . 5 . 53 . 53 . 61 . 157 |
| 63911 | 5 . 5 . 13 . 13 . 17 . 29 . 37 . 53 |
| 66361 | 5 . 5 . 41 . 113 . 193 . 197 |
| 79011 | 5 . 5 . 5 . 5 . 13 . 89 . 89 . 97 |
| 79871 | 5 . 29 . 37 . 61 . 101 . 193 |
| 81487 | 13 . 13 . 73 . 73 . 73 . 101 |
| 81669 | 5 . 13 . 29 . 113 . 173 . 181 |
| 86487 | 17 . 17 . 41 . 41 . 89 . 173 |
| 91587 | 17 . 29 . 29 . 37 . 101 . 157 |
| 95963 | 13 . 13 . 41 . 89 . 109 . 137 |
| 99011 | 5 . 5 . 5 . 5 . 5 . 13 . 29 . 53 . 157 |
| 99407 | 17 . 29 . 37 . 41 . 73 . 181 |
| 108111 | 5 . 5 . 13 . 17 . 97 . 113 . 193 |
| 110211 | 5 . 5 . 13 . 37 . 73 . 101 . 137 |
| 114611 | 5 . 5 . 13 . 13 . 89 . 181 . 193 |
| 115983 | 13 . 17 . 37 . 61 . 149 . 181 |
| 117281 | 5 . 29 . 61 . 89 . 101 . 173 |
| 128359 | 5 . 13 . 17 . 29 . 53 . 89 . 109 |
| 139701 | 5 . 13 . 53 . 149 . 193 . 197 |
| 140489 | 5 . 5 . 5 . 41 . 113 . 173 . 197 |
| 140871 | 5 . 13 . 17 . 37 . 61 . 73 . 109 |
| 142047 | 29 . 29 . 41 . 53 . 61 . 181 |
| 148939 | 5 . 5 . 29 . 29 . 73 . 97 . 149 |
| 183739 | 5 . 5 . 5 . 13 . 29 41 . 101 . 173 |
| 191279 | 5 . 13 . 13 . 29 . 73 . 113 . 181 |
| 203091 | 5 . 17 . 17 . 41 . 61 . 101 . 113 |
| 205111 | 5 . 5 . 13 . 73 . 97 . 101 . 181 |
| 206707 | 29 . 37 . 53 . 61 . 109 . 113 |
| 207171 | 5 . 13 . 17 . 29 . 89 . 101 . 149 |
| 211221 | 5 . 13 . 13 . 37 . 61 . 149 . 157 |
| 228179 | 5 . 13 . 13 . 61 . 73 . 101 . 137 |
| 234333 | 37 . 37 . 37 . 41 . 137 . 193 |
| 234881 | 5 . 13 . 13 . 17 . 17 . 17 . 97 . 137 |

| | |
|---|---|
| 241511 | 5 . 5 . 5 . 5 . 13 . 17 . 37 . 101 . 113 |
| 244299 | 5 . 13 . 17 . 37 . 97 . 101 . 149 |
| 245293 | 109 . 113 . 137 . 181 . 197 |
| 247699 | 5 . 13 . 17 . 41 . 61 . 149 . 149 |
| 257665 | 13 . 17 . 101 . 109 . 157 . 173 |
| 263489 | 5 . 5 . 5 . 5 . 5 . 5 . 29 . 37 . 41 . 101 |
| 269459 | 5 . 17 . 37 . 37 . 53 . 61 . 193 |
| 289589 | 5 . 5 . 29 . 61 . 97 . 113 . 173 |
| 302111 | 5 . 5 . 41 . 61 . 97 . 101 . 149 |
| 306757 | 17 . 29 . 53 . 101 . 181 . 197 |
| 313489 | 5 . 5 . 5 . 5 . 5 . 5 . 17 . 17 . 17 . 37 . 173 |
| 317039 | 5 . 5 . 53 . 61 . 89 . 89 . 157 |
| 326957 | 37 . 37 . 41 . 101 . 109 . 173 |
| 349835 | 17 . 41 . 73 . 97 . 137 . 181 |
| 355989 | 5 . 5 . 5 . 5 . 13 . 17 . 37 . 137 . 181 |
| 387921 | 5 . 61 . 101 . 137 . 181 . 197 |
| 396783 | 13 . 13 . 29 . 61 . 61 . 89 . 97 |
| 408489 | 5 . 5 . 5 . 5 . 13 . 29 . 73 . 89 . 109 |
| 466489 | 5 . 5 . 5 . 13 . 17 . 29 . 61 . 61 . 73 |
| 563235 | 13 . 17 . 41 . 61 . 61 . 97 . 97 |
| 567629 | 5 . 13 . 37 . 73 . 109 . 113 . 149 |
| 582997 | 37 . 41 . 73 . 113 . 157 . 173 |
| 588489 | 5 . 5 . 5 . 5 5 . 61 . 89 . 137 . 149 |
| 628261 | 5 . 5 . 5 . 13 . 17 . 29 . 41 . 61 . 197 |
| 634205 | 37 . 53 . 73 . 109 . 149 . 173 |
| 637855 | 13 . 37 . 41 . 41 . 61 . 73 . 113 |
| 834267 | 17 . 17 . 17 . 29 . 137 . 181 . 197 |
| 840421 | 5 . 13 . 17 . 137 . 149 . 173 . 181 |
| 851929 | 5 . 17 . 53 . 73 . 89 . 137 . 181 |
| 922769 | 5 . 13 . 13 . 17 . 41 . 61 . 137 . 173 |
| 966391 | 5 . 13 . 17 . 29 29 . 37 . 157 . 173 |
| 1029353 | 61 . 61 . 89 . 109 . 149 . 197 |
| 1165689 | 5 . 5 . 29 . 41 . 41 . 53 . 109 . 193 |
| 1230349 | 5 . 13 . 13 . 17 . 17 . 29 . 37 . 53 . 109 |
| 1299241 | 5 . 13 . 13 . 37 . 89 . 89 . 173 . 197 |
| 1341429 | 5 . 29 . 61 . 73 . 89 . 173 . 181 |
| 1362611 | 5 . 5 . 13 37 . 89 . 89 . 101 . 193 |
| 1493911 | 5 . 5 . 13 . 29 . 101 . 109 . 137 . 157 |
| 1499001 | 5 . 13 . 13 . 17 . 73 . 73 . 149 . 197 |
| 1780489 | 5 . 5 . 5 . 29 . 37 . 41 . 53 . 73 . 149 |
| 1996199 | 5 . 13 . 13 . 17 . 17 . 37 . 53 53 . 157 |
| 2028211 | 5 . 5 . 13 . 17 . 41 . 41 . 53 . 61 . 137 |
| 2050005 | 17 . 29 . 61 . 73 . 89 . 137 . 157 |
| 2159739 | 5 . 5 . 5 . 5 . 13 . 17 . 37 . 41 . 113 . 197 |
| 3376311 | 5 . 5 . 13 . 13 . 13 . 17 . 17 . 61 . 61 . 193 |
| 3666653 | 13 . 13 . 17 . 41 . 53 . 97 . 149 . 149 |
| 3872099 | 5 . 13 . 13 . 17 . 37 . 41 . 41 . 97 . 173 |
| 4370811 | 5 . 5 . 13 . 17 . 53 . 61 . 89 . 197 |

| | |
|---|---|
| 4490249 | 5 . 13 . 29 . 61 . 73 . 89 . 137 . 197 |
| 4705711 | 5 . 5 . 13 . 17 . 17 . 17 . 17 . 41 . 101 . 197 |
| 5125339 | 5 . 5 . 17 . 17 . 29 . 37 . 97 . 181 . 193 |
| 5472411 | 5 . 5 . 17 . 29 . 41 . 41 . 89 . 109 . 149 |
| 6101547 | 13 . 13 . 13 . 17 . 29 . 37 . 61 . 97 . 157 |
| 6489011 | 5 . 5 . 5 . 5 . 17 . 37 . 41 . 89 . 149 . 197 |
| 8175989 | 5 . 5 . 5 . 5 . 5 . 13 . 17 . 37 . 97 . 149 . 181 |
| 8649761 | 5 . 5 . 5 . 13 . 17 . 37 . 61 . 101 . 109 . 109 |
| 8812979 | 5 . 17 . 17 . 17 . 29 . 73 . 89 . 97 . 173 |
| 9530277 | 13 . 13 . 17 . 17 . 17 . 73 . 89 . 113 . 149 |
| 10126399 | 5 . 13 . 17 . 29 . 29 . 53 . 89 . 149 . 157 |
| 10251621 | 5 . 13 . 17 . 17 29 . 61 . 101 . 173 . 181 |
| 10763489 | 5 . 5 . 5 . 5 . 5 . 5 . 61 . 61 . 101 . 109 . 181 |
| 10831321 | 5 . 17 . 29 . 29 . 41 . 61 . 73 . 89 . 101 |
| 11398611 | 5 . 5 . 13 . 17 . 53 . 97 . 137 . 173 . 193 |
| 11483821 | 5 . 29 . 53 . 61 . 101 . 113 . 157 . 157 |
| 15035789 | 5 . 5 . 37 . 41 . 41 . 53 . 101 . 157 . 173 |
| 17363031 | 5 . 13 . 13 . 41 . 73 . 89 . 89 . 101 . 149 |

| | |
|---|---|
| 23866411 | 5 . 5 . 13 . 17 . 17 . 17 . 61 . 157 . 193 . 193 |
| 25252451 | 5 . 13 . 29 . 37 . 41 . 97 . 97 . 137 . 173 |
| 31456571 | 5 . 13 . 17 . 41 . 97 . 101 . 109 . 113 . 181 |
| 34411159 | 5 . 13 . 37 . 41 . 61 . 73 . 89 . 157 . 193 |
| 35272357 | 13 . 13 . 13 . 13 . 17 . 29 . 61 . 97 . 101 . 137 |
| 35944451 | 5 . 13 . 17 . 17 . 17 . 29 . 73 . 89 109 . 197 |
| 61017271 | 13 . 13 . 13 . 17 . 29 . 37 . 61 . 97 . 157 |
| 107402539 | 5 . 5 . 13 . 29 . 37 . 61 . 109 . 149 . 173 . 193 |
| 143828743 | 29 . 29 . 37 . 37 . 53 . 97 . 113 . 157 . 197 |
| 148757489 | 5 . 5 . 5 . 13 . 13 . 17 . 37 . 41 . 41 . 61 . 109 . 149 |
| 150446761 | 5 . 5 . 5 . 13 . 17 . 37 . 53 . 113 . 137 . 137 . 197 |
| 322564791 | 5 . 13 . 17 . 29 . 29 . 29 . 37 . 53 . 101 . 101 . 193 |
| 657182319 | 5 . 17 . 17 . 61 . 97 . 97 . 113 . 149 . 157 . 197 |
| 1359685525 | 13 . 17 . 53 . 61 . 61 . 97 . 113 . 157 . 157 . 157 |
| 4949475989 | 5 . 5 . 5 . 5 . 5 . 13 . 13 . 29 . 29 . 37 . 37 . 41 . 61 . 89 . 181 |
| 28608252345 | 13 . 29 . 29 . 29 . 37 . 53 . 61 . 73 . 97 . 113 . 149 . 181 |
| 112899039159 | 5 . 13 . 13 . 17 . 17 . 17 . 17 . 29 . 37 . 41 . 61 . 73 . 73 . 73 . 173 |

| | |
|---|---|
| 5 | 1. 11 |
| 13 | 3. 29 |
| 17 | 9 |
| 29 | 5. 111. 179 |
| 37 | 25. 49. 1011. 4761. 10039 |
| 41 | 23. 59. 469. 4123. 14901 |
| 53 | 7. 99. 961. 4989. 20769. 63911 |
| 61 | 39. 61. 205. 1381. 5041 |

| | |
|---|---|
| 73 | 19. 127. 311. 419. 749. 466489 |
| 89 | 21. 199. 691. 1089. 5717. 16041 |
| 97 | 53. 141. 247. 335. 4021. 6261. 7319. 79011. 396783. 563235* |
| 101 | 81. 121. 283. 3717. 6989. 9817. 12605. 32239. 81487. 263489*. 10831321 |
| 109 | 43. 261. 393. 829. 2355. 2441. 3749. 6061. 21189. 29169*. 32003. 128359. 140871. 408489. 1230349. 8649761 |
| 113 | 83. 309. 761. 1047. 1499. 2343. 5567. 27429. 203091. 206707*. 244299. 637855 |
| 137 | 63. 211. 485. 611. 759. 1159. 1581. 3351. 9801. 13489*. 21161. 27611. 29929. 95963. 110211. 228179. 234881. 2028211. 35272357 |
| 149 | 61. 237. 535. 1131. 2025. 3339. 8879. 25689. 148939. 207171*. 244299. 247699. 302111. 567629. 588489. 1780489. 3666653. 5472411. 9530277. 17363031*. 148757489 |
| 157 | 101. 213. 527. 841. 1043. 8579. 9947. 20511. 37579. 47829*. 57989. 91587. 99011. 211221. 317039. 1493911. 1996199. 2050005. 6101547. 10126399*. 11483821. 61017271. 1359685525 |
| 173 | 13. 359. 679. 705. 1371. 12815. 13507. 23915. 44301. 47389*. 47761. 86487. 117281. 183739. 257665. 289589. 313489. 326957. 582997. 634205*. 922769. 966391. 3872099. 8812979. 15035789. 25252451. 112899039159 |
| 181 | 143. 219. 943. 1305. 2677. 3039. 5573. 7745. 13251. 14261*. 21139. 23311. 81669. 99407. 115983. 142047. 191279. 205111. 349835. 355989*. 840421. 851929. 1341429. 8175989. 10251621. 10763489. 31456571. 4949475989. 28608252345 |
| 193 | 31. 417. 1899. 4215. 5821. 20875. 22805. 23901. 29691. 79871*. 108111. 114611. 234333. 269459. 1165689. 1362611. 3376311. 5125339. 11398611. 23866411*. 34411159. 107402539. 322564791 |
| 197 | 169. 1351. 1745. 1801. 3377. 8049. 12383. 27355. 27411. 31351*. 32139. 49813. 66361. 139701. 140489. 245293. 306757. 387921. 628261. 834267*. 1029353. 1299241. 1499001. 2159739. 4370811. 4490249. 4705711. 6489011. 35944451. 143828743*. 150446761. 657182319 |

## **Factorable** $aa + 9$

| | | | |
|---|---|---|---|
| 1 | 5 | 2 | 13 |
| 4 | 5 . 5 | 5 | 17 |
| 7 | 29 | 8 | 73 |
| 10 | 109 | 11 | 5 . 13 |
| 13 | 89 | 14 | 5 . 41 |
| 16 | 5 . 53 | 17 | 149 |
| 19 | 5 . 37 | 22 | 17 . 29 |
| 26 | 5 . 137 | 28 | 13 . 61 |
| 29 | 5 . 5 . 17 | 31 | 5 . 97 |
| 37 | 13 . 53 | 41 | 5 . 13 . 13 |
| 46 | 5 . 5 . 5 . 17 | 50 | 13 . 193 |
| 55 | 37 . 41 | 56 | 5 . 17 . 37 |
| 65 | 29 . 73 | 67 | 13 . 173 |
| 68 | 41 . 113 | 71 | 5 . 5 . 101 |
| 73 | 17 . 157 | 76 | 5 . 13 . 89 |

| | | | |
|---|---|---|---|
| 79 | 5 . 5 . 5 . 5 . 5 | 80 | 13 . 17 . 29 |
| 89 | 5 . 13 . 61 | 94 | 5 . 29 . 61 |
| 106 | 5 . 13 . 173 | 109 | 5 . 29 . 41 |
| 119 | 5 . 13 . 109 | 124 | 5 . 17 . 181 |
| 128 | 13 . 13 . 97 | 131 | 5 . 17 . 101 |
| 143 | 53 . 193 | 154 | 5 . 5 . 13 . 73 |
| 155 | 61 . 197 | 158 | 13 . 17 . 113 |
| 163 | 97 . 137 | 166 | 5 . 37 . 149 |
| 167 | 13 . 29 . 37 | 175 | 17 . 17 . 53 |
| 181 | 5 . 29 . 113 | 191 | 5 . 41 . 89 |
| 196 | 5 . 5 . 29 . 53 | 211 | 5 . 61 . 73 |
| 232 | 13 . 41 . 101 | 239 | 5 . 29 . 197 |
| 241 | 5 . 37 . 157 | 254 | 5 . 5 . 29 . 89 |
| 271 | 5 . 5 . 13 . 113 | 281 | 5 . 53 . 149 |
| 284 | 5 . 13 . 17 . 73 | 301 | 5 . 13 . 17 . 41 |
| 314 | 5 . 13 . 37 . 41 | 352 | 17 . 37 . 197 |
| 379 | 5 . 5 . 13 . 13 . 17 | 413 | 17 . 29 . 173 |
| 419 | 5 . 97 . 181 | 430 | 17 . 73 . 149 |
| 436 | 5 . 193 . 197 | 437 | 17 . 41 . 137 |
| 446 | 5 . 5 . 73 . 109 | 454 | 5 . 5 . 5 . 17 . 97 |
| 464 | 5 . 17 . 17 . 149 | 521 | 5 . 5 . 61 . 89 |
| 529 | 5 . 5 . 29 . 193 | 535 | 13 . 101 . 109 |
| 544 | 5 . 13 . 29 . 157 | 547 | 41 . 41 . 89 |
| 574 | 5 . 13 . 37 . 137 | 610 | 37 . 89 . 113 |
| 629 | 5 . 5 . 41 . 193 | 704 | 5 . 5 . 5 . 5 . 13 . 61 |
| 722 | 37 . 73 . 193 | 746 | 5 . 5 . 113 . 197 |
| 796 | 5 . 5 . 5 . 37 . 137 | 811 | 5 . 17 . 53 . 73 |
| 821 | 5 . 5 . 13 . 17 . 61 | 848 | 29 . 137 . 181 |
| 869 | 5 . 13 . 37 . 157 | 943 | 37 . 61 . 197 |
| 971 | 5 . 5 . 109 . 173 | 991 | 5 . 17 . 53 . 109 |
| 1015 | 17 . 157 . 193 | 1042 | 13 . 17 . 17 . 17 . 17 |
| 1055 | 13 . 13, 37 . 89 | 1070 | 61 . 137 . 137 |
| 1081 | 5 . 13 . 89 . 101 | 1129 | 5 . 5 . 13 . 37 . 53 |

| | |
|---|---|
| 1144 | 5 . 17 . 89 . 173 |
| 1175 | 41 . 113 . 149 |
| 1259 | 5 . 13 . 89 . 137 |
| 1298 | 13 . 29 . 41 . 109 |
| 1309 | 5 . 53 . 53 . 61 |
| 1324 | 5 . 13 . 149 . 181 |
| 1421 | 5 . 5 . 5 . 41 . 197 |
| 1559 | 5 . 17 . 17 . 29 . 29 |
| 1618 | 97 . 137 . 197 |
| 1627 | 13 . 17 . 53 . 113 |
| 1646 | 5 . 5 . 29 . 37 . 101 |
| 1675 | 13 . 29 . 61 . 61 |
| 1687 | 73 . 101 . 193 |

| | |
|---|---|
| 1805 | 13 . 29 . 29 . 149 |
| 1831 | 5 . 13 . 17 . 37 . 41 |
| 1909 | 5 . 13 . 17 . 17 . 97 |
| 1979 | 5 . 5 . 29 . 37 . 73 |
| 2069 | 5 . 13 . 13 . 17 . 149 |
| 2182 | 13 . 29 . 73 . 173 |
| 2351 | 5 . 13 . 17 . 41 . 61 |
| 2528 | 17 . 41 . 53 . 173 |
| 2596 | 5 . 5 . 17 . 101 . 257 |
| 2719 | 5 . 13 . 29 . 37 . 53 |
| 2839 | 5 . 61 . 73 . 181 |
| 2953 | 13 . 17 . 109 . 181 |
| 3038 | 17 . 29 . 97 . 193 |
| 3089 | 5 . 17 . 37 . 37 . 41 |
| 3458 | 29 . 41 . 89 . 113 |
| 3496 | 5 . 5 . 37 . 73 . 181 |
| 3571 | 5 . 5 . 37 . 61 . 113 |
| 3677 | 13 . 13 . 13 . 17 . 181 |
| 4136 | 5 . 13 . 17 . 113 . 137 |
| 4171 | 5 . 5 . 5 . 13 . 53 . 101 |
| 4196 | 5 . 5 . 41 . 89 . 193 |
| 4237 | 37 . 41 . 61 . 97 |
| 4459 | 5 . 17 . 29 . 37 . 109 |
| 4489 | 5 . 53 . 193 . 197 |
| 4496 | 5 . 5 . 13 . 37 . 41 . 41 |
| 4565 | 13 41 . 113 . 173 |
| 4786 | 5 . 13 . 53 . 61 . 109 |
| 5029 | 5 . 5 . 13 . 13 . 41 . 73 |
| 5111 | 5 . 13 . 13 . 13 . 29 . 41 |
| 5125 | 37 . 37 . 53 . 181 |
| 5198 | 13 . 13 . 29 . 37 . 149 |
| 5401 | 5 . 17 . 29 . 61 . 97 |
| 5549 | 5 . 13 . 41 . 53 . 109 |
| 5579 | 5 . 5 . 5 . 13 . 61 . 157 |
| 5605 | 13 . 17 . 17 . 37 . 113 |
| 5921 | 5 . 5 . 5 . 17 . 73 . 113 |
| 6329 | 5 . 5 . 5 . 5 . 5 . 13 . 17 . 29 |
| 6346 | 5 . 5 . 13 . 17 . 37 . 197 |
| 6421 | 5 . 5 . 5 . 17 . 89 . 109 |
| 6494 | 5 . 37 . 37 . 61 . 101 |
| 6641 | 5 . 13 . 37 . 53 . 173 |
| 6821 | 5 . 5 . 53 . 97 . 181 |
| 7196 | 5 . 5 . 17 . 37 . 37 . 89 |
| 7271 | 5 . 5 . 17 . 37 . 41 . 41 |
| 7489 | 5 . 29 . 41 . 53 . 89 |
| 7616 | 5 . 13 . 53 . 113 . 149 |
| 7646 | 5 . 5 . 13 . 13 . 101 . 137 |
| 7729 | 5 . 5 . 97 . 109 . 113 |

| | |
|---|---|
| 7934 | 5 . 17 . 53 . 89 . 157 |
| 9650 | 29 . 113 . 157 . 181 |
| 10012 | 13 . 13 . 29 . 113 . 181 |
| 10154 | 5 . 5 . 17 . 41 . 61 . 97 |
| 10447 | 29 . 73 . 149 . 173 |
| 10736 | 5 . 13 . 97 . 101 . 181 |
| 11074 | 5 . 13 . 61 . 157 . 197 |
| 11671 | 5 . 5 . 5 . 41 . 97 . 137 |
| 12109 | 5 . 17 . 41 . 109 . 193 |
| 12191 | 5 . 37 . 41 . 97 . 101 |
| 13561 | 5 . 13 . 13 . 17 . 37 . 173 |
| 14029 | 5 . 5 . 13 . 29 . 53 . 197 |
| 15004 | 5 . 5 . 13 . 37 . 97 . 193 |
| 16096 | 5 . 5 . 13 . 13 . 13 . 53 . 89 |
| 16291 | 5 . 13 . 17 . 29 . 41 . 101 |
| 17029 | 5 . 5 . 17 . 41 . 53 . 157 |
| 17357 | 13 . 41 . 41 . 61 . 113 |
| 17668 | 17 . 29 . 37 . 109 . 157 |
| 18671 | 5 . 5 . 5 . 5 . 5 . 17 . 17 . 193 |
| 19504 | 5 . 5 . 17 . 89 . 89 . 113 |
| 20651 | 5 . 61 . 61 . 73 . 157 |
| 20813 | 17 . 37 . 53 . 73 . 89 |
| 22085 | 13 . 41 . 53 . 89 . 97 |
| 22367 | 17 . 17 . 37 . 149 . 157 |
| 22700 | 13 . 17 . 29 . 37 . 41 . 53 |
| 23425 | 29 . 29 . 41 . 73 . 109 |
| 23671 | 5 . 5 . 5 . 5 . 13 . 29 . 29 . 41 |
| 23879 | 5 . 5 . 13 . 61 . 73 . 197 |
| 24001 | 5 . 61 . 61 . 113 . 137 |
| 24311 | 5 . 53 . 61 . 101 . 181 |
| 25645 | 89 . 137 . 149 . 181 |
| 26141 | 5 . 13 . 17 . 37 . 61 . 137 |
| 27341 | 5 . 13 . 17 . 17 . 101 . 197 |
| 27731 | 5 . 13 . 29 . 37 . 37 . 149 |
| 27805 | 13 . 37 . 73 . 101 . 109 |
| 27844 | 5 . 13 . 13 . 13 . 13 . 61 . 89 |
| 28804 | 5 . 5 . 29 . 37 . 157 . 197 |
| 28973 | 17 . 41 . 73 . 73 . 113 |
| 30544 | 5 . 17 . 29 . 37 . 53 . 193 |
| 33629 | 5 . 5 . 13 . 89 . 113 . 173 |
| 34010 | 13 . 13 . 29 . 53 . 61 . 73 |
| 38608 | 13 . 41 . 137 . 137 . 149 |
| 39704 | 5 . 5 . 5 . 13 . 73 . 97 . 137 |
| 40030 | 17 . 41 . 97 . 137 . 173 |
| 40304 | 5 . 5 . 73 . 73 . 89 . 137 |
| 42173 | 29 . 29 . 89 . 109 . 109 |
| 42421 | 5 . 5 . 5 . 5 . 13 . 37 . 41 . 73 |
| 43864 | 5 . 13 . 37 . 89 . 89 . 101 |

| | |
|---|---|
| 47296 | 5 . 5 . 5 . 13 . 73 . 109 . 173 |
| 47335 | 13 . 29 . 89 . 173 . 193 |
| 48046 | 5 . 5 . 5 . 5 . 13 . 29 . 97 . 101 |
| 48829 | 5 . 5 . 5 . 5 . 17 . 29 . 53 . 73 |
| 49883 | 13 . 17 . 17 . 41 . 41 . 197 |
| 49924 | 5 . 17 . 41 . 73 . 97 . 101 |
| 54871 | 5 . 5 . 13 . 17 . 53 . 53 . 97 |
| 54926 | 5 . 37 . 41 . 41 . 89 . 109 |
| 55709 | 5 . 97 . 109 . 149 . 197 |
| 57701 | 5 . 37 . 41 . 41 . 53 . 101 |
| 57839 | 5 . 13 . 13 . 13 . 13 . 13 . 17 . 53 |
| 62402 | 13 . 13 . 17 . 89 . 97 . 157 |
| 66584 | 5 . 13 . 17 . 17 . 53 61 . 73 |
| 70171 | 5 . 5 . 5 . 17 . 37 . 173 . 181 |
| 71021 | 5 . 5 . 13 . 13 . 13 . 17 . 37 . 73 |
| 71276 | 5 . 17 . 61 . 89 . 101 . 109 |
| 71354 | 5 . 5 . 17 . 37 . 41 . 53 . 149 |
| 73972 | 13 . 17 . 29 . 53 . 89 . 181 |
| 78829 | 5 . 5 . 5 . 5 . 29 . 37 . 41 . 113 |
| 79051 | 5 . 13 . 13 . 37 . 37 . 37 . 73 |
| 84563 | 13 . 17 . 17 . 17 . 17 . 37 . 89 |
| 84818 | 17 . 29 . 29 . 61 . 73 . 113 |
| 86221 | 5 . 5 . 61 . 73 . 173 . 193 |
| 88411 | 5 . 13 . 37 . 73 . 113 . 197 |
| 95071 | 5 . 5 . 13 . 37 . 61 . 61 . 101 |
| 95188 | 13 . 13 . 13 . 17 . 41 . 61 . 97 |
| 105274 | 5 . 41 . 53 . 73 . 89 . 157 |
| 109279 | 5 . 5 . 29 . 37 . 41 . 61 . 89 |
| 109991 | 5 . 13 . 53 . 89 . 109 . 181 |
| 112171 | 5 . 5 . 5 . 17 . 109 . 157 . 173 |
| 114499 | 5 . 29 . 29 . 41 . 193 . 197 |
| 114896 | 5 . 5 . 13 . 41 . 61 . 109 . 149 |
| 115079 | 5 . 5 . 5 . 5 . 29 . 53 . 61 . 113 |
| 116881 | 5 . 13 . 53 . 73 . 157 . 173 |
| 127114 | 5 . 17 . 29 . 29 . 37 . 41 . 149 |
| 133523 | 17 . 29 . 41 . 53 . 53 . 157 |
| 134764 | 5 . 17 . 97 . 101 . 113 . 193 |
| 137659 | 5 . 13 . 37 . 137 . 149 . 193 |
| 141581 | 5 . 13 . 13 . 17 . 37 109 . 173 |
| 146794 | 5 . 13 . 41 41 . 53 . 61 . 61 |
| 147409 | 5 . 13 . 13 . 41 . 53 . 61 97 |
| 154679 | 5 . 5 . 29 . 37 . 41 . 73 . 149 |
| 154729 | 5 . 5 . 17 . 17 . 61 . 157 . 173 |
| 157454 | 5 . 5 . 5 . 13 . 37 . 41 . 89 . 113 |
| 167689 | 5 . 13 . 13 . 13 . 73 . 89 . 197 |
| 170107 | 13 . 17 . 17 . 29 . 37 . 37 . 97 |
| 174427 | 61 . 89 . 113 . 137 . 181 |
| 178336 | 5 . 13 . 13 . 41 . 61 . 101 . 149 |

| | |
|---|---|
| 178988 | 17 . 37 . 37 . 73 . 109 . 173 |
| 180416 | 5 . 13 . 17 . 29 . 89 . 101 . 113 |
| 190021 | 5 . 5 . 17 . 41 . 53 . 113 . 173 |
| 190541 | 5 . 17 . 41 . 137 . 193 . 197 |
| 193546 | 5 . 5 . 5 . 13 . 13 . 97 . 101 . 181 |
| 193829 | 5 . 5 . 5 . 5 . 5 . 29 . 89 . 137 |
| 219754 | 5 . 5 . 13 . 17 . 17 . 53 . 89 . 109 |
| 249871 | 5 . 5 . 13 . 17 . 17 . 29 . 73 . 157 |
| 250250 | 37 . 41 . 53 . 61 . 113 . 113 |
| 252328 | 13 . 41 . 89 . 97 . 101 . 137 |
| 263681 | 5 . 13 . 13 . 37 . 101 . 101 . 109 |
| 265256 | 5 . 17 . 29 . 41 . 61 . 101 . 113 |
| 280750 | 13 . 13 . 17 . 61 . 61 . 73 . 101 |
| 286904 | 5 . 5 . 17 . 29 . 29 . 41 . 41 . 137 |
| 293687 | 17 . 37 . 61 . 73 . 89 . 173 |
| 311921 | 5 . 5 . 5 . 17 . 53 . 61 . 73 . 97 |
| 313454 | 5 . 5 . 5 . 13 . 13 . 13 . 13 . 13 . 29 . 73 |
| 314257 | 17 . 17 . 73 . 109 . 109 . 197 |
| 316739 | 5 . 17 . 37 . 41 . 73 . 73 . 73 |
| 324952 | 29 . 37 . 41 . 89 . 149 . 181 |
| 341569 | 5 . 17 . 29 . 29 . 53 . 89 . 173 |
| 347543 | 17 . 29 . 41 . 113 . 137 . 193 |
| 356809 | 5 . 13 . 29 . 37 . 41 . 113 . 197 |
| 377804 | 5 . 5 . 13 . 109 . 113 . 181 . 197 |
| 386722 | 13 . 29 . 109 . 109 . 173 . 193 |
| 393079 | 5 . 5 . 5 . 13 . 17 . 137 . 137 . 149 |
| 415825 | 17 . 37 . 37 . 109 . 173 . 197 |
| 419246 | 5 . 5 . 29 . 53 . 137 . 173 193 |
| 423475 | 17 . 41 . 89 . 89 . 109 . 149 |
| 448280 | 41 . 41 . 41 . 89 . 181 . 181 |
| 496004 | 5 . 5 . 13 . 17 . 37 . 41 . 149 . 197 |
| 512579 | 5 . 5 . 5 . 5 . 5 . 13 . 17 . 37 . 53 . 97 |
| 520921 | 5 . 5 . 5 . 13 . 53 . 97 . 109 . 149 |
| 524704 | 5 . 5 . 5 . 13 . 13 . 29 . 41 . 97 . 113 |
| 528967 | 17 . 29 . 101 . 109 . 149 . 173 |
| 539996 | 5 . 5 . 13 . 13 . 37 . 109 . 109 . 157 |
| 541829 | 5 . 5 . 5 . 13 . 17 . 29 . 41 . 41 . 109 |
| 550985 | 37 . 41 . 61 . 101 . 109 . 149 |
| 554279 | 5 . 5 . 13 . 13 . 37 . 61 . 89 . 181 |
| 599510 | 13 . 17 . 29 . 29 . 109 . 113 . 157 |
| 693775 | 17 . 17 . 17 . 41 . 97 . 109 . 113 |
| 700061 | 5 . 13 . 41 . 53 . 89 . 101 . 193 |
| 713291 | 5 . 17 . 29 . 29 . 29 . 41 . 41 . 73 |
| 744421 | 5 . 5 . 5 . 13 37 . 149 . 157 . 197 |
| 745249 | 5 . 13 . 13 . 29 . 29 . 53 . 73 . 101 |
| 792113 | 29 . 61 . 89 . 101 . 109 . 181 |
| 847319 | 5 . 17 . 37 . 73 . 101 . 113 . 137 |
| 859379 | 5 . 5 . 17 . 29 . 29 . 53 . 101 . 193 |

| | |
|---|---|
| 895208 | 13 . 17 . 17 . 17 . 29 . 41 . 61 . 173 |
| 895861 | 5 . 17 . 29 . 89 . 97 . 109 . 173 |
| 937766 | 5 . 13 . 17 . 29 . 41 . 53 . 73 . 173 |
| 947329 | 5 . 5 . 5 . 37 . 73 . 89 . 109 . 137 |
| 970454 | 5 . 5 . 5 . 37 . 73 . 97 . 149 . 193 |
| 984934 | 5 . 13 . 17 . 29 . 29 . 61 . 109 . 157 |
| 987406 | 5 . 17 . 53 . 73 . 101 . 149 . 197 |
| 1196173 | 53 . 89 . 97 . 101 . 113 . 137 |
| 1202704 | 5 . 5 . 5 . 17 . 17 . 37 . 61 . 113 . 157 |
| 1256084 | 5 . 13 . 17 . 29 . 53 . 61 . 97 . 157 |
| 1297090 | 13 . 29 . 37 . 41 . 109 . 137 . 197 |
| 1460288 | 13 . 13 . 13 . 17 . 29 . 101 . 101 . 193 |
| 1717025 | 13 . 29 41 . 53 . 73 . 157 . 157 |
| 1799921 | 5 . 5 . 5 . 5 . 5 . 17 . 29 . 37 . 157 . 181 |
| 1800254 | 5 . 5 . 17 . 37 . 97 . 101 . 109 . 193 |
| 2153956 | 5 . 17 . 101 . 101 . 157 . 173 . 197 |
| 2253046 | 5 . 5 . 5 . 5 . 5 . 29 . 41 . 53 . 149 . 173 |
| 2347195 | 17 . 29 . 41 . 53 . 137 . 137 . 137 |
| 2362579 | 5 . 5 . 5 . 5 . 5 . 5 . 5 . 13 . 29 . 37 . 197 |
| 2382560 | 13 . 29 . 37 . 101 . 113 . 181 . 197 |
| 2454779 | 5 . 5 . 13 . 41 . 73 . 109 . 157 . 181 |
| 2473954 | 5 . 5 . 5 . 13 . 17 . 17 . 29 . 41 . 97 . 113 |
| 2579296 | 5 . 5 . 5 . 5 . 17 . 29 . 41 . 61 . 89 . 97 |
| 2710934 | 5 . 17 . 17 . 17 . 37 . 41 . 53 . 61 . 61 |
| 2867521 | 5 . 5 . 17 . 61 . 73 . 101 . 137 . 157 |
| 2960596 | 5 . 5 . 13 . 17 . 41 . 53 . 73 . 73 . 137 |
| 3045079 | 5 . 5 . 5 . 5 . 13 . 17 . 17 . 101 . 113 . 173 |
| 3287839 | 5 . 17 . 17 . 17 . 29 . 53 . 53 . 73 . 73 |
| 3386888 | 13 . 13 . 17 . 29 . 37 . 137 . 157 . 173 |
| 3569269 | 5 . 13 . 29 . 41 . 53 . 89 . 101 . 173 |
| 4046131 | 5 . 13 . 17 . 29 . 89 . 101 . 157 . 181 |
| 4546271 | 5 . 5 . 13 . 17 . 41 . 53 . 53 . 109 . 149 |
| 4699704 | 5 . 5 . 5 . 13 . 61 . 73 . 113 . 137 . 197 |
| 4889605 | 37 . 41 . 53 . 61 . 73 . 173 . 193 |
| 8026096 | 5 . 5 . 17 . 37 . 37 . 89 . 101 . 109 . 113 |
| 8182343 | 17 . 17 . 29 . 41 . 73 . 73 . 101 . 181 |
| 8931226 | 5 . 37 . 53 . 61 . 89 . 89 . 113 . 149 |
| 9237421 | 5 . 5 . 5 . 5 . 5 . 13 . 17 . 17 . 29 . 29 . 29 . 149 |
| 9250762 | 41 . 89 . 97 . 101 . 101 . 137 . 173 |
| 10419736 | 5 . 13 . 13 . 29 . 37 . 73 . 101 . 109 . 149 |
| 11077571 | 5 . 5 . 13 . 13 . 41 . 109 . 113 . 149 . 193 |
| 12519856 | 5 . 13 . 37 37 . 41 . 53 . 61 . 97 . 137 |
| 13237028 | 17 . 29 . 37 . 41 . 61 . 149 . 149 . 173 |
| 13382956 | 5 . 13 . 17 . 29 . 29 . 37 . 137 . 193 . 197 |
| 14937769 | 5 . 13 . 17 . 37 . 53 . 61 . 61 . 101 . 137 |
| 19912579 | 5 . 5 . 5 . 5 . 5 . 13 . 41 . 41 . 97 . 173 . 173 |
| 20620229 | 5 . 5 . 37 . 37 . 61 . 73 . 73 . 97 . 197 |
| 22181629 | 5 . 5 . 13 . 13 . 13 . 17 . 29 . 37 . 41 . 53 . 113 |

| | |
|---|---|
| 23504986 | 5 . 13 . 13 . 13 . 29 . 29 . 29 . 41 . 53 . 73 |
| 25674911 | 5 . 13 . 29 . 37 . 41 . 73 . 89 . 113 . 157 |
| 26999399 | 5 . 13 . 13 . 29 . 37 . 97 . 109 . 193 . 197 |
| 33399844 | 5 . 17 . 17 . 29 . 37 . 37 . 41 . 73 . 73 . 89 |
| 33753059 | 5 . 13 . 13 . 41 . 89 . 97 . 101 . 109 . 173 |
| 34618846 | 5 . 5 . 13 . 13 . 13 . 17 . 17 . 37 . 97 . 109 . 193 |
| 34792409 | 5 . 13 . 13 . 13 . 17 . 29 . 41 . 101 . 137 . 197 |
| 40103726 | 5 . 17 . 37 . 41 . 109 . 109 . 197 |
| 41494546 | 5 . 5 . 5 . 13 . 61 . 61 . 89 . 109 . 149 . 197 |
| 48279454 | 5 . 5 . 5 . 5 . 13 . 29 . 41 . 41 . 41 . 61 . 181 |
| 60740461 | 5 . 13 . 17 . 17 . 17 . 41 . 73 . 97 . 101 . 197 |
| 64370954 | 5 . 5 . 5 . 13 . 17 . 37 . 53 . 61 . 73 . 89 . 193 |
| 96499349 | 5 . 13 . 17 . 29 37 . 41 . 61 . 73 . 137 . 157 |
| 105742171 | 5 . 5 . 5 . 13 . 17 . 29 . 29 . 37 . 41 . 41 . 53 . 73 |
| 110518796 | 5 . 5 . 5 . 13 . 37 . 53 . 53 . 61 . 73 . 109 . 149 |
| 111009121 | 5 . 5 . 13 . 17 . 17 . 37 . 37 . 37 . 73 . 113 . 157 |
| 113737804 | 5 . 5 . 13 . 13 . 29 . 29 . 53 . 73 . 89 . 97 . 109 |
| 117290203 | 17 . 29 . 37 . 41 . 53 . 73 . 109 . 113 . 193 |
| 149574656 | 5 . 29 . 41 . 61 . 73 . 137 . 173 . 181 . 197 |
| 163030454 | 5 . 5 . 5 . 13 . 13 . 17 . 37 . 37 . 53 . 73 . 89 . 157 |
| 165242573 | 13 . 29 . 37 . 41 . 73 . 97 . 109 . 157 . 197 |
| 178643779 | 5 . 5 . 13 . 41 . 41 . 61 . 113 . 137 . 157 . 197 |
| 200760094 | 5 . 13 . 17 . 17 . 37 . 37 . 37 . 41 . 53 . 101 . 193 |
| 323643829 | 5 . 5 . 5 . 5 . 5 . 5 . 13 . 13 . 17 . 37 . 61 . 73 . 73 . 97 |
| 401580454 | 5 . 5 . 5 . 13 . 53 . 61 . 73 . 73 . 97 . 137 . 137 . 193 |
| 478666540 | 17 . 17 . 29 . 37 . 41 . 61 . 73 . 149 . 157 . 173 |
| 1411168679 | 5 . 5 . 13 . 17 . 17 . 89 . 113 . 157 . 173 . 197 . 197 |

| | |
|---|---|
| 5 | 1. 4. 79 |
| 13 | 2. 11. 41 |
| 17 | 5. 29. 46. 379. 1042 |
| 29 | 7. 22. 80. 1559. 6329 |
| 37 | 19. 56. 167 |
| 41 | 14. 55. 109. 301. 314. 1831. 3089. 4496. 5111. 7271*. 23671 |
| 53 | 16. 37. 175. 196. 1129. 2719. 22700. 57839 |
| 61 | 28. 89. 94. 704. 821. 1309. 1675. 2351. 146794. 2710934* |
| 73 | 8. 65. 154. 211. 284. 811. 1979. 5029. 34010. 42421*. 48829. 66584. 71021. 79051. 313454. 316739. 713291. 3287839. 23504986. 105742171* |
| 89 | 13. 76. 191. 254. 521. 547. 1055. 7196. 7489. 16096*. 20813. 27844. 84563. 109279. 33399844 |
| 97 | 31. 128. 454. 1909. 4237. 5401. 10154. 22085. 54871. 95188*. 147409. 170107. 311921. 512579. 2579296. 323643829 |
| 101 | 71. 131. 232. 1081. 1646. 4171. 6494. 12191. 16291. 43864*. 48046. 49924. 57701. 95071. 280750. 745249 |
| 109 | 10. 119. 446. 535. 991. 1298. 4459. 4786. 5549. 6421*. 23425. 27805. 42173. 54926. 71276. 219754. 263681. 541829. 113737804 |

| | |
|---|---|
| 113 | 68. 158. 181. 271. 610. 1627. 3458. 3571. 5605. 5921*. 7729. 17357. 19504. 28973 78829. 84818. 115079. 157454. 180416. 250250*. 265256. 524704. 693775. 2473954. 8026096. 22181629 |
| 137 | 26. 163. 437. 574. 796. 1070. 1259. 4136. 7646. 11671*. 24001. 26141. 39704. 40304. 193829. 252328. 286904. 847319. 947329. 1196173*. 2347195. 2960596. 12519856. 14937769 |
| 149 | 17. 166. 281. 430. 464. 1175. 1805. 2069. 5198. 7616*. 27731. 38608. 71354. 114896. 127114. 154679. 178336. 393079. 423475. 520921*. 550985. 4546271. 8931226. 9237421. 10419736. 110518796 |
| 157 | 73. 241. 544. 869. 2596. 5579. 7934. 17029. 17668. 20651*. 22367. 62402. 105274. 133523. 249871. 539996. 599510. 984934. 1202704. 1256084*. 1717025. 2867521. 25674911. 96499349. 111009121. 163030454 |
| 173 | 67. 106. 413. 971. 1144. 2182. 2528. 4565. 6641. 10447*. 13561. 33629. 40030. 47296. 112171. 116881. 141581. 154729. 178988. 190021*. 293687. 341569. 528967. 895208. 895861. 937766. 2253046. 3045079. 3386888. 3569269*. 9250762. 13237028. 19912579. 33753059. 478666540 |
| 181 | 124. 419. 848. 1324. 2839. 2953. 3496. 3677. 5125. 6821*. 9650. 10012. 10736. 24311. 25645. 70171. 73972. 109991. 174427. 193546*. 324952. 448280. 554279. 792113. 1799921. 2454779. 4046131. 8182343. 48279454 |
| 193 | 50. 143. 529. 629. 722. 1015. 1687. 3038. 4196. 12109*. 15004. 18671. 30544. 47335. 86221. 134764. 137659. 347543. 386722. 419246*. 700061. 859379. 970454. 1460288. 1800254. 4889605. 11077571. 34618846. 64370954. 117290203*. 200760094. 401580454 |
| 197 | 155. 239. 352. 436. 746. 943. 1421. 1618. 4489. 6346*. 11074. 14029. 23879. 27341. 28804. 49883. 55709. 88411. 114499. 167689*. 190541. 314257. 356809. 377804. 415825. 496004. 744421. 987406. 1297090. 2153956*. 2362579. 2382560. 4699704. 13382956. 26620229. 26999399. 34792409. 40103726. 41494546. 60740461*. 149574656. 165242573. 178643779. 1411168679 |

## **Factorable** $aa + 16$**.**

| | |
|---|---|
| 1 | 17 |
| 3 | 5 . 5 |
| 5 | 41 |
| 7 | 5 . 13 |
| 9 | 97 |
| 11 | 137 |
| 13 | 5 . 37 |
| 17 | 5 . 61 |
| 19 | 13 . 29 |
| 23 | 5 . 109 |
| 27 | 5 . 149 |
| 33 | 5 . 13 . 17 |
| 35 | 17 . 73 |

| | |
|---|---|
| 39 | 29 . 53 |
| 45 | 13 . 157 |
| 47 | 5 . 5 . 89 |
| 53 | 5 . 5 . 113 |
| 61 | 37 . 101 |
| 67 | 5 . 17 . 53 |
| 77 | 5 . 29 . 41 |
| 87 | 5 . 37 . 41 |
| 97 | 5 . 5 . 13 . 29 |
| 103 | 5 . 5 . 5 . 5 . 17 |
| 105 | 61 . 181 |
| 131 | 89 . 193 |
| 135 | 17 . 29 . 37 |
| 137 | 5 . 13 . 17 . 17 |
| 141 | 101 . 197 |
| 147 | 5 . 5 . 5 . 173 |
| 173 | 5 . 53 . 113 |
| 227 | 5 . 13 . 13 . 61 |
| 241 | 13 . 41 . 109 |
| 253 | 5 . 5 . 13 . 197 |
| 257 | 5 . 73 . 181 |
| 263 | 5 . 101 . 137 |
| 271 | 17 . 29 . 149 |
| 279 | 13 . 53 . 113 |
| 309 | 29 . 37 . 89 |
| 357 | 5 . 13 . 37 . 53 |
| 383 | 5 . 13 . 37 . 61 |
| 397 | 5 . 5 . 5 . 13 . 97 |
| 403 | 5 5 . 73 . 89 |
| 487 | 5 . 13 . 41 . 89 |
| 505 | 37 . 61 . 113 |
| 545 | 17 . 101 . 173 |
| 569 | 41 . 53 . 149 |
| 579 | 13 . 17 . 37 . 41 |
| 619 | 29 . 73 . 181 |
| 647 | 5 . 5 . 5 . 17 . 197 |
| 677 | 5 . 29 . 29 . 109 |
| 747 | 5 . 5 . 13 . 17 . 101 |
| 851 | 13 . 17 . 29 . 113 |
| 897 | 5 . 5 . 5 . 41 . 157 |
| 903 | 5 . 5 . 13 . 13 . 193 |
| 987 | 5 . 17 . 73 . 157 |
| 1021 | 13 . 17 . 53 . 89 |
| 1203 | 5 . 5 . 13 . 61 . 73 |
| 1237 | 5 . 29 . 61 . 173 |
| 1293 | 5 . 13 . 17 . 17 . 89 |
| 1353 | 5 . 5 . 5 . 5 . 29 . 101 |
| 1359 | 13 . 17 . 61 . 137 |

| | |
|---|---|
| 1463 | 5 . 13 . 13 . 17 . 149 |
| 1553 | 5 . 5 . 13 . 41 . 181 |
| 1717 | 5 . 41 . 73 . 197 |
| 1837 | 5 . 17 . 29 . 37 . 37 |
| 1929 | 137 . 157 . 173 |
| 2203 | 5 . 5 . 13 . 109 . 137 |
| 2223 | 5 . 29 . 173 . 197 |
| 2243 | 5 . 13 . 17 . 29 . 157 |
| 2301 | 29 . 41 . 61 . 73 |
| 2447 | 5 . 5 . 17 . 37 . 193 |
| 2455 | 29 . 37 . 41 . 137 |
| 2477 | 5 . 13 . 13 . 53 . 137 |
| 2593 | 5 . 13 . 13 . 73 . 109 |
| 2687 | 5 . 17 . 29 . 29 . 101 |
| 2823 | 5 . 17 . 29 . 53 . 61 |
| 2957 | 5 . 13 . 17 . 41 . 193 |
| 3095 | 17 . 37 . 97 . 157 |
| 3113 | 5 . 13 . 29 . 53 . 97 |
| 3153 | 5 . 5 . 13 . 13 . 13 . 181 |
| 3247 | 5 . 5 . 53 . 73 . 109 |
| 3293 | 5 . 101 . 109 . 197 |
| 3603 | 5 . 5 . 5 . 17 . 41 . 149 |
| 3607 | 5 . 13 . 13 . 89 . 173 |
| 3777 | 5 . 13 41 . 53 . 101 |
| 3847 | 5 . 5 . 29 . 137 . 149 |
| 4453 | 5 . 5 . 17 . 37 . 97 |
| 4497 | 5 . 5 . 61 . 89 . 149 |
| 4505 | 13 . 13 . 29 . 41 . 101 |
| 4601 | 29 . 37 . 109 . 181 |
| 4647 | 5 . 5 . 5 . 13 . 97 . 137 |
| 4853 | 5 5 . 5 . 29 . 73 . 89 |
| 4897 | 5 . 5 . 5 . 5 . 17 . 37 . 61 |
| 5337 | 5 . 13 . 17 . 149 . 173 |
| 5473 | 5 . 17 . 53 . 61 . 109 |
| 5635 | 13 . 13 . 13 . 97 . 149 |
| 5897 | 5 . 5 . 5 . 29 . 53 . 181 |
| 5921 | 13 . 89 . 157 . 193 |
| 6051 | 13 . 17 . 29 . 29 . 197 |
| 6081 | 37 . 53 . 109 . 173 |
| 6427 | 5 . 17 . 53 . 53 . 173 |
| 6605 | 61 . 73 . 97 . 101 |
| 6727 | 5 . 13 . 61 . 101 . 113 |
| 7345 | 17 . 89 . 181 . 197 |
| 7413 | 5 . 17 . 37 . 101 . 173 |
| 7547 | 5 . 5 . 13 . 13 . 13 . 17 . 61 |
| 7683 | 5 . 17 . 37 . 137 . 137 |
| 7703 | 5 . 5 . 13 . 41 . 61 . 73 |
| 7963 | 5 . 13 . 89 . 97 . 113 |

| | |
|---|---|
| 8141 | 73 . 89 . 101 . 101 |
| 8523 | 5 . 37 . 41 . 61 . 157 |
| 8747 | 5 . 5 . 101 . 157 . 193 |
| 9133 | 5 . 13 . 61 . 109 . 193 |
| 9353 | 5 . 5 . 5 . 13 . 13 . 41 . 101 |
| 10003 | 5 . 5 . 13 . 37 . 53 . 157 |
| 11967 | 5 . 13 . 17 . 29 . 41 . 109 |
| 12045 | 13 . 29 . 53 . 53 . 137 |
| 12257 | 5 . 29 . 53 . 113 . 173 |
| 12603 | 5 . 5 . 5 . 5 . 13 . 113 . 173 |
| 12667 | 5 . 37 . 73 . 109 . 109 |
| 13397 | 5 . 5 . 5 . 13 . 17 . 73 . 89 |
| 16897 | 5 . 5 . 5 . 17 . 29 . 41 . 113 |
| 17477 | 5 . 17 . 17 . 29 . 37 . 197 |
| 17635 | 13 . 41 . 53 . 101 . 109 |
| 17853 | 5 . 5 . 5 . 109 . 149 . 157 |
| 19991 | 17 . 29 . 61 . 97 . 137 |
| 20167 | 5 . 41 . 97 . 113 . 181 |
| 23677 | 5 . 53 . 97 . 113 . 193 |
| 24447 | 5 . 5 . 13 . 13 . 17 . 53 . 157 |
| 24785 | 13 . 13 . 17 . 29 . 73 . 101 |
| 25617 | 5 . 13 . 29 . 37 . 97 . 97 |
| 28581 | 13 . 53 . 73 . 109 . 149 |
| 29217 | 5 . 13 . 37 . 37 . 53 . 181 |
| 29853 | 5 . 5 . 5 . 17 . 41 . 53 . 193 |
| 36107 | 5 . 13 . 17 . 53 . 113 . 197 |
| 36823 | 5 . 13 . 17 . 41 . 173 . 173 |
| 37579 | 5 . 17 . 29 . 41 . 89 . 157 |
| 38863 | 5 . 13 . 17 . 17 . 37 . 41 . 53 |
| 39653 | 5 . 5 . 29 . 101 . 109 . 197 |
| 40853 | 5 . 5 . 5 . 13 . 61 . 113 . 149 |
| 41373 | 5 . 13 . 29 . 29 . 173 . 181 |
| 44269 | 17 . 53 . 61 . 181 . 197 |
| 44947 | 5 . 5 . 13 . 17 . 17 . 137 . 157 |

| | |
|---|---|
| 45793 | 5 . 13 . 37 . 89 . 97 . 101 |
| 52157 | 5 . 17 . 17 . 17 . 37 . 41 . 73 |
| 52379 | 37 . 73 . 89 . 101 . 113 |
| 52393 | 5 . 17 . 29 . 41 . 157 . 173 |
| 57323 | 5 . 13 . 17 29 41 . 41 . 61 |
| 57803 | 5 . 5 . 89 . 97 . 113 . 137 |
| 66333 | 5 . 13 . 17 . 17 . 29 . 41 . 197 |
| 67327 | 5 . 37 . 41 . 61 . 97 . 101 |
| 68215 | 37 . 61 . 101 . 137 . 149 |
| 69347 | 5 . 5 . 101 . 101 . 109 . 173 |
| 73467 | 5 . 29 . 41 . 89 . 101 . 101 |
| 74133 | 5 . 13 . 13 . 41 . 41 . 53 . 73 |

| | |
|---|---|
| 81413 | 5 . 13 . 29 . 29 . 29 . 37 . 113 |
| 82817 | 5 . 13 . 73 . 89 . 109 . 149 |
| 82893 | 5 . 17 . 37 . 73 . 173 . 173 |
| 103317 | 5 . 13 . 13 . 29 . 37 . 61 . 193 |
| 104293 | 5 . 13 . 61 . 101 . 157 . 173 |
| 113699 | 29 . 29 . 29 . 53 . 73 . 137 |
| 126497 | 5 . 5 . 13 . 53 . 61 . 97 . 157 |
| 130553 | 5 . 5 . 13 . 53 . 97 . 101 . 101 |
| 132143 | 5 . 29 . 53 . 193 . 193 |
| 139477 | 5 . 37 . 41 . 97 . 137 . 193 |
| 150897 | 5 . 5 . 5 . 13 . 29 . 61 . 89 . 89 |
| 154821 | 29 . 37 . 41 . 41 . 97 . 137 |
| 158373 | 5 . 13 . 17 . 37 . 61 . 89 . 113 |
| 158509 | 17 . 17 . 53 . 101 . 109 . 149 |
| 161399 | 17 . 17 . 53 . 89 . 97 . 197 |
| 162383 | 5 . 17 . 89 . 149 . 149 . 157 |
| 171293 | 5 . 17 . 29 . 41 . 41 . 73 . 97 |
| 172569 | 13 . 29 . 29 . 101 . 149 . 181 |
| 174727 | 5 . 13 . 17 . 37 . 53 . 73 . 193 |
| 232147 | 5 . 5 . 5 . 13 . 41 . 41 . 109 . 181 |
| 239387 | 5 . 53 . 97 . 109 . 113 . 181 |
| 240347 | 5 . 5 . 17 . 41 . 89 . 193 . 193 |
| 242897 | 5 . 5 . 5 . 17 . 17 . 97 . 113 . 149 |
| 251817 | 5 . 13 . 29 . 41 . 53 . 113 . 137 |
| 260033 | 5 . 13 . 13 . 17 . 29 . 29 . 29 . 193 |
| 260575 | 17 . 29 . 61 . 73 . 157 . 197 |
| 300527 | 5 . 13 . 17 . 37 . 113 . 113 . 173 |
| 374203 | 5 . 5 . 17 . 89 . 113 . 181 . 181 |
| 378671 | 13 . 13 . 37 . 41 . 53 . 61 . 173 |
| 434441 | 13 . 13 . 37 . 37 . 41 . 101 . 197 |
| 577603 | 5 . 5 . 5 . 5 . 29 . 29 . 41 . 113 . 137 |
| 648447 | 5 . 5 . 13 . 17 . 61 . 61 . 113 . 181 |
| 650103 | 5 . 5 . 5 . 5 . 29 . 37 . 73 . 89 . 97 |
| 658783 | 5 . 17 . 17 . 29 . 41 . 41 . 61 . 101 |
| 696353 | 5 . 5 . 5 . 5 17 . 17 . 37 . 37 . 37 . 53 |
| 748853 | 5 . 5 . 5 . 5 . 5 . 61 . 109 . 137 . 197 |
| 870487 | 5 . 13 . 61 . 73 . 97 . 137 . 197 |
| 873503 | 5 . 5 . 13 . 13 61 . 109 . 157 . 173 |
| 970497 | 5 5 . 17 . 37 . 53 . 73 . 113 . 137 |
| 1193679 | 13 29 . 41 . 41 . 101 113 . 197 |
| 1229533 | 5 . 13 . 13 . 53 . 53 . 53 . 61 . 197 |
| 1259837 | 5 . 13 . 13 . 13 . 17 . 29 . 37 . 89 . 89 |
| 1335487 | 5 . 17 . 89 . 97 . 113 . 137 . 157 |
| 1404163 | 5 . 13 . 13 . 37 . 41 . 97 . 101 . 157 |
| 1626475 | 13 . 29 . 41 . 73 . 109 . 137 . 157 |
| 2008103 | 5 . 5 . 5 . 13 . 41 . 53 . 61 . 97 . 193 |
| 2083893 | 5 . 13 . 17 . 53 . 73 . 89 . 101 . 113 |
| 2116091 | 17 . 29 . 37 . 101 . 113 . 137 . 157 |

| | |
|---|---|
| 2373167 | 5 . 17 . 29 . 37 . 41 . 53 . 157 . 181 |
| 2960653 | 5 . 5 . 13 . 13 . 17 . 17 . 17 . 101 . 113 |
| 3258603 | 5 . 5 . 5 . 37 . 41 . 61 . 61 . 101 . 149 |
| 3611583 | 5 . 17 . 29 . 37 . 61 . 109 . 137 . 157 |
| 3898603 | 5 . 5 . 5 . 13 . 13 . 37 . 41 . 73 . 73 . 89 |
| 4945505 | 13 . 13 . 17 . 17 . 17 . 29 . 89 . 101 . 113 |
| 5431603 | 5 . 5 . 5 . 17 . 17 . 29 . 41 . 73 . 97 . 97 |
| 8180243 | 5 . 13 . 29 . 29 . 37 . 37 . 41 . 53 . 53 . 113 . 193 |
| 8268383 | 5 . 13 . 13 . 29 . 41 . 73 . 73 . 113 . 113 |
| 9993613 | 5 . 13 . 29 . 37 . 53 . 53 . 61 . 61 . 137 |
| 10311423 | 5 . 41 . 41 . 61 . 109 . 113 . 113 . 149 |
| 15305803 | 5 . 5 . 13 . 37 . 53 . 101 . 109 . 173 . 193 |
| 16626883 | 5 . 17 . 73 . 101 . 109 . 149 . 157 . 173 |
| 17545053 | 5 . 5 . 13 . 17 . 17 . 41 . 53 . 101 . 109 . 137 |
| 17916571 | 17 . 37 . 37 . 61 . 73 . 109 . 157 . 181 |
| 18500917 | 5 . 13 . 29 . 41 . 41 . 61 . 89 . 101 . 197 |
| 19344643 | 5 . 13 . 13 . 29 . 41 . 73 . 73 . 113 . 113 |
| 20278927 | 5 . 13 . 17 . 29 . 53 . 73 . 113 . 149 . 197 |
| 22858302 | 5 . 5 . 17 . 17 . 17 . 17 . 17 . 29 . 53 . 61 . 157 |
| 38648107 | 5 . 17 . 17 . 29 . 97 . 109 . 109 . 157 . 197 |
| 40473647 | 5 . 5 . 5 . 5 . 13 . 37 . 41 . 73 . 97 . 137 . 137 |
| 46113113 | 5 . 13 . 13 . 17 . 29 . 37 . 53 . 73 . 181 . 197 |
| 1082687431 | 13 . 17 . 29 53 . 61 . 97 . 109 157 . 173 . 197 |
| 1254102921 | 13 . 13 . 17 . 17 . 41 . 53 . 61 . 97 . 101 . 137 . 181 |

| | |
|---|---|
| 5 | 3 |
| 13 | 7 |
| 17 | 1. 33. 103. 137 |
| 29 | 19. 97 |
| 37 | 13. 135. 1837 |
| 41 | 5. 77. 87. 579 |
| 53 | 39. 67. 357. 38863. 696353 |
| 61 | 17. 227. 383. 2823. 4897. 7547. 57323 |
| 73 | 35. 1203. 2301. 7703. 52157. 74133 |
| 89 | 47. 309. 403. 487. 1021. 1293. 4853. 13397. 150897. 1259837*. 3898603 |
| 97 | 9. 397. 3113. 4453. 25617. 171293. 650103. 5431603 |
| 101 | 61. 747. 1353. 2687. 3777. 4505. 6605. 8141. 9353. 24785*. 45793. 67327. 73467. 130553. 658783 |
| 109 | 23. 241. 677. 2593. 3247. 5473. 11967. 12667. 17635 |
| 113 | 53. 173. 279. 505. 851. 6727. 7963. 16897. 52379. 81413*. 158373. 2083893. 2960653. 4945505. 8268383. 19344643 |
| 137 | 11. 263. 1359. 2203. 2455. 2477. 4647. 7683. 12045. 19991*. 57803. 113699. 154821. 251817. 577603. 970497. 9993613. 17545053. 40473647 |
| 149 | 11. 271. 569. 1463. 3603. 3849. 4497. 5635. 28581. 40853*. 68215. 82817. 158509. 242897. 3258603. 10311423 |

| | |
|---|---|
| 157 | 45. 897. 987. 2243. 3095. 8523. 10003. 17853. 24447. 37579*. 44947. 126497. 162384. 1335487. 1404163. 1626475. 2116091. 3611583. 22858302 |
| 173 | 147. 545. 1237. 1929. 3607. 5337. 6081. 6427. 7413. 12257*. 12603. 36823. 52393. 69347. 82893. 104293. 300527. 378671. 873503. 16626883* |
| 181 | 105. 257. 619. 1553. 3153. 4601. 5897. 20167. 29217. 41373*. 172569. 232147. 239387. 374203. 648447. 2373167. 17916571. 1254102921 |
| 193 | 131. 903. 2447. 2957. 5921. 8747. 9133. 23677. 29853. 103317. 132143 139477. 174727. 240347. 260033. 2008103. 8180243 15305803 |
| 197 | 141. 253. 647. 1717. 2223. 3293. 6051. 7345. 17477. 36107*. 39653. 44269. 66333. 161399. 260575. 434441. 748853. 870487. 1193679. 1229533*. 18500917. 20278927. 38648107. 46113113. 1082687431 |

## **Factorable** $aa + 25$**.**

| | |
|---|---|
| 1 | 13 |
| 2 | 29 |
| 3 | 17 |
| 4 | 41 |
| 6 | 61 |
| 7 | 37 |
| 8 | 89 |
| 9 | 53 |
| 11 | 73 |
| 12 | 13 . 13 |
| 13 | 97 |
| 14 | 13 . 17 |
| 17 | 157 |
| 19 | 193 |
| 27 | 13 . 29 |
| 31 | 17 . 29 |
| 37 | 17 . 41 |
| 38 | 13 . 113 |
| 44 | 37 . 53 |
| 48 | 17 . 137 |
| 51 | 13 . 101 |
| 53 | 13 . 109 |
| 54 | 17 . 173 |
| 56 | 29 . 109 |
| 62 | 53 . 73 |
| 67 | 37 . 61 |
| 71 | 17 . 149 |
| 78 | 41 . 149 |
| 81 | 37 . 89 |
| 84 | 73 . 97 |
| 86 | 41 . 181 |

| | |
|---|---|
| 89 | 29 . 137 |
| 97 | 53 . 89. |
| 99 | 17 . 17 . 17 |
| 116 | 13 . 17 . 61 |
| 118 | 13 . 29 . 37 |
| 119 | 41 . 173 |
| 127 | 41 . 197 |
| 151 | 101 . 113 |
| 157 | 13 . 13 . 73 |
| 168 | 13 . 41 . 53 |
| 174 | 157 . 193 |
| 181 | 13 . 13 . 97 |
| 201 | 17 . 29 . 41 |
| 207 | 13 . 17 . 97 |
| 209 | 13 . 41 . 41 |
| 227 | 149 . 173 |
| 252 | 17 . 37 . 101 |
| 259 | 13 . 29 . 89 |
| 267 | 181 . 197 |
| 274 | 13 . 53 . 109 |
| 303 | 17 . 37 . 73 |
| 309 | 17 . 53 53 |
| 311 | 13 . 61 . 61 |
| 324 | 13 . 41 . 197 |
| 326 | 13 . 13 . 17 . 37 |
| 354 | 17 . 73 . 101 |
| 363 | 13 . 37 . 137 |
| 376 | 13 . 73 . 149 |
| 377 | 17 . 37 . 113 |
| 414 | 37 . 41 . 113 |
| 433 | 29 . 53 . 61 |
| 437 | 29 . 37 . 89 |
| 454 | 13 . 101 . 157 |
| 488 | 37 . 41 . 157 |
| 521 | 13 . 53 . 197 |
| 573 | 13 . 73 . 173 |
| 611 | 29 . 41 . 157 |
| 636 | 13 . 29 . 29 . 37 |
| 638 | 13 . 173 . 181 |
| 677 | 13 . 17 . 17 . 61 |
| 733 | 37 . 53 . 137 |
| 753 | 13 . 113 . 193 |
| 768 | 13 . 17 . 17 . 157 |
| 816 | 41 . 109 . 149 |
| 819 | 17 . 109 . 181 |
| 857 | 13 . 13 . 41 . 53 |
| 858 | 37 . 101 . 197 |
| 898 | 13 . 17 . 41 . 89 |

| | |
|---|---|
| 959 | 29 . 101 . 157 |
| 984 | 29 . 173 . 193 |
| 1092 | 61 . 113 . 173 |
| 1104 | 13 . 29 . 53 . 61 |
| 1177 | 37 . 97 . 193 |
| 1252 | 73 . 109 . 197 |
| 1364 | 13 . 13 . 101 . 109 |
| 1431 | 13 . 17 . 41 . 113 |
| 1442 | 13 . 17 . 97 . 97 |
| 1544 | 17 . 17 . 73 . 113 |
| 1561 | 13 . 17 . 37 . 149 |
| 1733 | 97 . 113 . 137 |
| 1767 | 13 . 29 . 41 . 101 |
| 1887 | 29 . 29 . 29 . 73 |
| 2128 | 17 . 41 . 73 . 89 |
| 2144 | 13 . 29 . 89 . 137 |
| 2341 | 13 . 41 . 53 . 97 |
| 2434 | 17 . 29 . 61 . 197 |
| 2751 | 17 . 41 . 61 . 89 |
| 2887 | 13 . 17 . 109 . 173 |
| 2989 | 13 . 17 . 17 . 29 . 41 |
| 3199 | 13 . 13 . 13 . 17 . 137 |
| 3323 | 37 . 37 . 37 . 109 |
| 3471 | 17 . 37 . 61 . 157 |
| 3522 | 13 . 17 . 37 . 37 . 41 |
| 3654 | 13 . 61 . 113 . 149 |
| 3686 | 17 . 41 . 101 . 193 |
| 3788 | 17 . 61 . 101 . 137. |
| 4219 | 17 . 41 . 113 . 113 |
| 4264 | 17 . 61 . 89 . 197 |
| 4458 | 13 . 89 . 89 . 193 |
| 4798 | 13 . 89 . 101 . 197 |
| 4814 | 17 . 17 . 17 . 53 . 89 |
| 5154 | 17 . 89 . 97 . 181 |
| 5251 | 13 . 13 . 29 . 29 . 97 |
| 5706 | 13 . 101 . 137 . 181 |
| 5927 | 13 . 13 . 37 53 . 53 |
| 6001 | 29 . 73 . 97 . 173 |
| 6157 | 17 . 53 . 109 . 193 |
| 6581 | 29 . 53 . 73 . 193 |
| 6616 | 13 . 17 . 37 53 . 101 |
| 7359 | 13 . 97 . 109 . 197 |
| 7676 | 53 . 73 . 97 . 157 |
| 7753 | 41 . 61 . 61 . 197 |
| 8147 | 29 . 37 . 157 . 197 |
| 8231 | 17 . 101 . 109 . 181 |
| 8776 | 13 . 13 . 37 . 109 113 |
| 9209 | 73 . 73 . 73 . 109 |

| | |
|---|---|
| 10049 | 41 . 89 . 101 . 137 |
| 10061 | 13 . 17 . 29 . 53 . 149 |
| 10501 | 37 . 73 . 137 . 149 |
| 12468 | 13 . 29 . 41 . 89 . 113 |
| 12526 | 17 . 17 . 29 . 97 . 193 |
| 13786 | 73 . 101 . 149 . 173 |
| 13787 | 89 97 . 101 . 109 |
| 14067 | 13 . 29 37 . 41 . 173 |
| 14756 | 13 . 37 . 41 . 61 . 181 |
| 15807 | 13 . 17 . 29 . 101 . 193 |
| 17057 | 13 . 13 . 53 . 109 . 149 |
| 18123 | 13 . 29 . 37 . 61 . 193 |
| 18771 | 13 . 13 . 13 17 . 53 . 89 |
| 18823 | 13 . 29 . 41 . 73 . 157 |
| 19553 | 13 . 17 . 17 . 17 . 41 . 73 |
| 19751 | 17 . 17 . 17 . 29 . 37 . 37 |
| 20502 | 13 . 53 . 61 . 73 . 137 |
| 21009 | 13 . 17 . 37 . 137 . 197 |
| 21319 | 13 . 37 . 37 . 113 . 113 |
| 21527 | 13 . 53 . 61 . 137 . 149 |
| 21644 | 13 . 13 . 17 . 41 . 41 . 97 |
| 23488 | 29 . 37 . 53 . 89 . 109 |
| 24101 | 13 . 29 . 61 . 73 . 173 |
| 24358 | 17 . 29 . 41 . 149 . 197 |
| 25401 | 13 . 17 . 97 . 101 . 149 |
| 26707 | 29 . 29 . 37 . 73 . 157 |
| 30467 | 17 . 41 . 41 . 109 . 149 |
| 31226 | 17 . 53 . 61 . 113 . 157 |
| 33381 | 29 . 37 . 53 . 97 . 101 |
| 38011 | 13 . 41 . 89 . 97 . 157 |
| 38134 | 17 . 41 . 97 . 137 . 157 |
| 40559 | 13 . 17 . 17 . 37 . 61 . 97 |
| 41037 | 29 . 41 . 73 . 89 . 109 |
| 41891 | 17 . 37 . 73 . 97 . 197 |
| 44407 | 13 . 17 . 17 37 . 41 . 173 |
| 48062 | 13 . 17 . 53 . 53 . 61 . 61 |
| 49943 | 17 . 37 . 73 . 157 . 173 |
| 50051 | 13 . 17 . 173 . 181 . 181 |
| 56913 | 13 . 17 . 37 . 37 . 53 . 101 |
| 60347 | 13 . 17 . 29 . 29 . 97 . 101 |
| 68626 | 13 . 13 17 . 53 . 157 . 197 |
| 85699 | 37 . 61 89 . 101 . 181 |
| 87989 | 17 . 53 . 113 . 193 . 197 |
| 93469 | 13 . 13 . 13 . 17 . 29 . 37 . 109 |
| 95473 | 13 . 13 . 13 . 73 . 157 . 181 |
| 101151 | 37 . 41 . 101 . 173 . 193 |
| 108871 | 17 . 53 173 . 193 . 197 |
| 121479 | 17 . 17 . 29 . 41 . 109 . 197 |

| | |
|---|---|
| 141777 | 13 . 17 . 37 . 73 . 113 . 149 |
| 144808 | 13 . 41 . 61 . 61 . 97 . 109 |
| 152762 | 13 . 41 . 41 . 97 . 101 . 109 |
| 152803 | 13 . 29 . 37 . 41 . 137 . 149 |
| 155187 | 73 . 97 . 101 . 113 149 |
| 160314 | 29 . 37 41 . 61 . 61 . 157 |
| 172561 | 13 . 13 . 13 . 13 . 37 . 73 . 193 |
| 183971 | 17 . 37 . 41 . 73 . 89 . 101 |
| 188618 | 13 . 17 . 17 29 53 . 61 . 101 |
| 214482 | 29 . 37 . 53 . 61 . 89 . 149 |
| 214631 | 13 . 29 . 41 . 73 . 137 . 149 |
| 234852 | 17 . 41 . 53 . 73 . 113 . 181 |
| 249014 | 13 . 73 . 73 . 89 . 89 . 113 |
| 257841 | 13 . 29 . 61 . 89 . 101 . 149 |
| 279007 | 13 . 13 . 17 . 29 . 29 . 89 . 181 |
| 329219 | 17 . 37 . 73 . 89 . 89 . 149 |
| 329848 | 13 . 17 . 17 . 29 . 37 . 137 . 197 |
| 382537 | 13 . 17 . 17 . 29 . 61 . 101 . 109 |
| 422419 | 17 . 17 . 17 . 41 . 53 . 61 . 137 |
| 458742 | 17 . 17 . 113 . 173 . 193 . 193 |
| 484041 | 13 . 29 . 37 . 37 . 61 . 61 . 61 |
| 546534 | 13 . 13 . 37 . 41 . 41 . 157 . 181 |
| 564812 | 13 . 37 . 37 . 41 . 53 . 73 . 113 |
| 735331 | 13 . 13 . 37 . 41 . 53 . 101 197 |
| 743781 | 13 . 13 . 13 . 13 . 17 . 37 . 89 . 173 |
| 867847 | 17 . 89 . 89 . 137 . 137 . 149 |
| 938003 | 13 . 29 . 53 . 61 . 61 . 61 . 97 |
| 1000154 | 13 . 13 . 29 . 29 . 37 . 37 53 . 97. |
| 1964806 | 13 . 13 . 17 . 29 . 41 . 73 . 113 . 137 |
| 2144583 | 13 . 41 . 53 . 61 . 73 . 101 . 181 |
| 3589859 | 17 . 17 . 73 . 109 . 113 . 137 . 181 |
| 3879591 | 13 . 37 . 53 . 89 . 113 . 149 . 197 |
| 5693622 | 13 . 13 . 13 . 53 . 97 . 101 . 157 . 181 |
| 6991009 | 13 17 . 17 . 37 . 53 . 113 . 149 . 197 |
| 7062082 | 13 . 29 . 29 . 41 . 53 . 73 . 149 193 |
| 8489259 | 13 . 17 . 29 . 29 . 29 . 41 . 41 . 41 . 97 |
| 8717008 | 13 . 13 17 . 17 . 37 . 37 . 89 . 113 . 113 |
| 9707868 | 13 . 29 . 29 . 37 . 101 . 113 . 137 . 149 |
| 10305788 | 13 . 17 . 37 . 53 101 . 109 . 113 . 197 |
| 17462342 | 13 . 37 . 61 . 61 . 89 . 89 . 137 157 |
| 38722306 | 13 . 13 . 29 . 37 . 61 . 89 . 89 . 109 . 157 |
| 48162204 | 13 . 17 . 37 . 37 . 41 . 53 . 101 . 181 . 193 |
| 60920523 | 13 . 17 . 17 . 53 . 61 . 61 . 101 . 137 . 181 |

| | |
|---|---|
| 63769026 | 17 . 29 . 37 . 41 . 53 . 89 . 101 . 101 . 113 |
| 111771087 | 13 . 17 . 37 . 61 . 61 . 89 . 113 . 137 . 149 |
| 141757784 | 13 . 17 . 53 . 61 . 89 . 113 . 137 . 137 . 149 |

| | |
|---|---|
| 172642653 | 13 . 17 . 29 . 73 . 89 . 89 . 137 . 149 . 197 |
| 190067607 | 73 . 89 . 97 . 101 . 101 . 109 . 149 . 173 |
| 308956283 | 13 . 29 . 29 . 37 . 41 . 53 . 61 . 73 . 89 . 137 |
| 569329071 | 13 . 13 . 29 . 37 . 41 . 97 . 101 . 109 . 137 . 149 |

| | |
|---|---|
| 13 | 1. 12 |
| 17 | 3. 14. 99 |
| 29 | 2. 27. 31 |
| 37 | 7. 118. 326. 636. 19751 |
| 41 | 4. 37. 201. 209. 2989. 3522 |
| 53 | 9. 44. 168. 309. 857. 5927 |
| 61 | 6. 67. 116. 311. 433. 677. 1104. 48062. 484041 |
| 73 | 11. 62. 157. 303. 1887. 19553 |
| 89 | 8. 81. 97. 259. 437. 898. 2128. 2751. 4814. 18771* |
| 97 | 13. 84. 181. 207. 1442. 2341. 5251. 21644. 40559. 938203*. 1000154. 8489259 |
| 101 | 51. 252. 354. 1767. 6616. 33381. 56913. 60347. 183971. 188618* |
| 109 | 53. 56. 274. 1364. 3323. 9209. 13787. 23488. 41037. 93469*. 144808. 152762. 382537 |
| 113 | 38. 151. 377. 414. 1431. 1544. 4219. 8776. 12468. 21319*. 249014. 564812. 8717008. 63769026 |
| 137 | 48. 89. 363. 733. 1733. 2144. 3199. 3788. 10049. 20502*. 422419. 1964806. 308956283 |
| 149 | 71. 78. 376. 816. 1561. 3654. 10061. 10501. 17057. 21527*. 25401. 30467. 14777. 152803. 155187. 214482. 214631. 257841. 329219. 867847*. 9707868. 111771087. 141757784. 569329071 |
| 157 | 17. 454. 488. 611. 768. 959. 3471. 7676. 18823. 26707*. 31226. 38011. 38134. 160314. 17462342. 38722306 |
| 173 | 54. 119. 227. 573. 1092. 2887. 6001. 13786. 14067. 24101*. 44407. 49943. 743781. 190067607 |
| 181 | 86. 638. 819. 5154. 5706. 8231. 14756. 50051. 85699. 95473*. 234852. 279007. 546534. 2144583. 3589859. 5693622. 60920523 |
| 193 | 19. 174. 753. 984. 1177. 3686. 4458. 6157. 6581. 12526*. 15807. 18123. 101151. 172561. 458742. 7062082. 48162204 |
| 197 | 127. 267. 324. 521. 858. 1252. 2434. 4264. 4798. 7359*. 7753. 8147. 21009. 24358. 41891. 68626. 87989. 108871. 121479. 329848*. 735331. 3879591. 6991009. 10305788. 172642653 |

**Factorable $aa + 36$.**

| | |
|---|---|
| 1 | 37 |
| 5 | 61 |
| 7 | 5 . 17 |
| 11 | 157 |
| 13 | 5 . 41 |
| 17 | 5 . 5 . 13 |
| 23 | 5 . 113 |
| 35 | 13 . 97 |
| 41 | 17 . 101 |
| 43 | 5 . 13 . 29 |
| 61 | 13 . 17 . 17 |
| 67 | 5 . 5 . 181 |
| 73 | 5 . 29 . 37 |
| 85 | 53 . 137 |
| 89 | 73 . 109 |
| 95 | 13 . 17 . 41 |
| 113 | 5 . 13 . 197 |
| 115 | 89 . 149 |
| 127 | 5 . 53 . 61 |
| 191 | 13 . 53 . 53 |
| 203 | 5 . 73 . 113 |
| 217 | 5 . 5 . 5 . 13 . 29 |
| 233 | 5 . 5 . 41 . 53 |
| 293 | 5 . 89 . 193 |
| 295 | 13 . 37 . 181 |
| 307 | 5 . 109 . 173 |
| 347 | 5 . 13 . 17 . 109 |
| 445 | 37 . 53 . 101 |
| 479 | 29 . 41 . 193 |
| 517 | 5 . 5 . 17 . 17 . 37 |
| 565 | 29 . 101 . 109 |
| 617 | 5 . 5 . 97 . 157 |
| 667 | 5 . 5 . 13 . 37 . 37 |
| 673 | 5 . 17 . 73 . 73 |
| 737 | 5 . 13 . 61 . 137 |
| 763 | 5 . 13 . 13 . 13 . 53 |
| 953 | 5 . 13 . 89 . 157 |
| 971 | 13 . 29 . 41 . 61 |
| 1183 | 5 . 5 . 17 . 37 . 89 |
| 1333 | 5 . 5 . 17 . 37 . 113 |
| 1463 | 5 . 41 . 53 . 197 |
| 1517 | 5 . 5 . 13 . 73 . 97 |
| 1673 | 5 . 13 . 17 . 17 . 149 |
| 1717 | 5 . 5 . 5 . 5 . 53 . 89 |
| 1873 | 5 . 41 . 109 . 157 |

| | |
|---|---|
| 2201 | 13 . 41 . 61 . 149 |
| 2251 | 17 . 17 . 89 . 197 |
| 2383 | 5 . 5 . 13 . 101 . 173 |
| 2557 | 5 . 13 . 17 . 61 . 97 |
| 2567 | 5 . 5 . 29 . 61 . 149 |
| 2963 | 5 . 89 . 109 . 181 |
| 3181 | 13 . 37 . 109 . 193 |
| 3553 | 5 . 13 . 29 . 37 . 181 |
| 3767 | 5 . 5 . 17 . 173 . 193 |
| 4031 | 41 . 61 . 73 . 89 |
| 4277 | 5 . 17 . 29 . 41 . 181 |
| 4883 | 5 . 5 . 37 . 149 . 173 |
| 5009 | 13 . 97 . 101 . 197 |
| 5321 | 13 . 13 . 29 . 53 . 109 |
| 5467 | 5 . 5 . 5 . 5 . 17 . 29 . 97 |
| 5495 | 13 . 13 . 29 . 61 . 101 |
| 5497 | 5 . 173 . 181 . 193 |
| 6217 | 5 . 5 . 5 . 37 . 61 . 137 |
| 6221 | 29 . 73 . 101 . 181 |
| 6655 | 29 . 41 . 193 . 193 |
| 6827 | 5 . 17 . 61 . 89 . 101 |
| 7547 | 5 . 37 . 37 . 53 . 157 |
| 7717 | 5 . 5 . 5 . 53 . 89 . 101 |
| 7813 | 5 . 17 . 61 . 61 . 193 |
| 7861 | 13 . 17 . 137 . 157 |
| 7919 | 37 . 97 . 101 . 173 |
| 8459 | 13 . 17 . 41 . 53 . 149 |
| 10261 | 13 . 17 . 53 . 89 . 101 |
| 10565 | 13 . 13 . 41 . 89 . 181 |
| 13763 | 5 . 13 . 17 . 37 . 41 . 113 |
| 13823 | 5 . 13 . 109 . 149 181 |
| 14543 | 5 . 13 . 29 . 29 . 53 . 73 |
| 15245 | 13 . 37 . 61 . 89 . 89 |
| 15733 | 5 . 5 . 29 . 29 . 61 . 193 |
| 17617 | 5 . 5 . 29 . 41 . 53 . 197 |
| 18659 | 13 . 17 . 97 . 109 . 149 |
| 22345 | 17 . 29 . 53 . 97 . 197 |
| 22481 | 13 . 17 . 17 . 17 . 41 . 193 |
| 22583 | 5 . 5 . 17 . 101 . 109 . 109 |
| 22733 | 5 . 5 . 13 . 13 . 13 . 97 . 97 |
| 22867 | 5 . 5 29 . 37 . 101 . 193 |
| 23753 | 5 . 37 . 113 . 137 . 197 |
| 29129 | 13 . 53 . 89 . 101 . 137 |
| 29995 | 13 . 13 . 13 . 13 . 17 . 17 . 109 |
| 30845 | 13 . 13 . 17 . 41 . 41 . 197 |
| 31885 | 13 . 17 29 . 41 . 53 . 73 |
| 32647 | 5 . 13 . 17 . 73 . 73 . 181 |
| 38893 | 5 . 73 . 109 . 193 . 197 |

| | |
|---|---|
| 38923 | 5 . 17 . 37 . 53 . 61 . 149 |
| 39347 | 5 . 13 . 41 . 53 . 97 . 113 |
| 42133 | 5 . 5 . 17 . 17 . 17 . 97 . 149 |
| 44327 | 5 . 29 . 29 . 37 . 73 . 173 |
| 48967 | 5 . 5 . 5 . 13 . 17 . 29 . 41 . 73 |

| | |
|---|---|
| 49517 | 5 . 5 . 29 . 113 . 173 . 173 |
| 54167 | 5 . 5 . 13 . 13 . 37 . 137 . 137 |
| 61117 | 5 . 5 . 13 . 29 . 61 . 73 . 89 |
| 62825 | 13 . 17 . 37 . 41 . 61 . 193 |
| 74603 | 5 . 13 . 17 . 29 . 29 . 53 . 113 |
| 87217 | 5 . 5 . 5 . 17 . 17 . 29 . 53 . 137 |
| 96227 | 5 . 17 . 53 . 109 . 109 . 173 |
| 125909 | 13 . 17 . 61 . 73 . 89 . 181 |
| 130613 | 5 . 41 . 53 . 73 . 137 . 157 |
| 141709 | 13 . 13 . 29 . 37 . 37 . 41 . 73 |
| 151163 | 5 . 29 . 61 . 109 . 137 . 173 |
| 161035 | 13 . 41 . 53 . 61 . 101 . 149 |
| 171655 | 89 . 97 . 109 . 173 . 181 |
| 174565 | 29 . 37 . 37 . 41 . 97 . 193 |
| 182743 | 5 . 17 . 29 . 29 . 29 . 89 . 181 |
| 189353 | 5 . 17 . 109 . 157 . 157 . 157 |
| 191203 | 5 . 53 . 53 . 73 . 181 . 197 |
| 206407 | 5 . 17 . 17 . 29 . 41 . 137 . 181 |
| 256693 | 5 . 17 . 17 . 29 . 61 . 149 . 173 |
| 387833 | 5 . 13 . 37 . 53 . 53 . 61 . 73 |
| 427795 | 13 . 17 . 37 . 37 . 53 . 101 . 113 |
| 429347 | 5 . 13 . 13 . 37 . 173 . 173 . 197 |
| 449921 | 13 . 29 . 37 . 41 . 41 . 89 . 97 |
| 533789 | 13 . 13 . 29 . 29 . 113 . 113 . 157 |
| 726029 | 17 . 17 . 29 . 29 . 101 . 109 . 197 |
| 837533 | 5 . 5 . 5 . 37 . 97 . 101 . 113 . 137 |
| 1097105 | 13 . 17 . 17 . 97 . 109 . 157 . 193 |
| 1396529 | 13 . 13 . 13 . 37 . 41 . 53 . 61 . 181 |
| 2390717 | 5 . 5 . 5 . 13 . 17 . 29 . 29 . 37 . 61 . 109 |
| 2525527 | 5 . 13 . 17 . 61 . 89 97 . 97 . 113 |
| 5318933 | 5 . 5 . 13 . 17 . 29 . 89 . 97 . 113 . 181 |
| 6920333 | 5 . 5 . 13 . 17 . 17 . 17 . 37 . 61 . 97 . 137 |
| 9439957 | 5 . 17 . 17 . 17 . 17 . 17 . 37 . 37 . 53 . 173 |
| 11776417 | 5 . 5 . 17 . 41 . 73 . 73 . 89 . 97 . 173 |
| 45435967 | 5 . 5 . 5 . 29 . 41 . 53 . 61 . 113 . 193 . 197 |
| 70145903 | 5 . 13 . 13 . 17 . 41 . 53 . 73 . 97 . 113 . 197 |
| 90115783 | 5 . 5 . 5 . 5 . 5 . 5 . 5 . 5 . 5 . 5 . 5 . 13 . 13 . 13 . 17 . 61 . 73 |
| 716295433 | 5 . 5 . 13 . 13 . 29 . 29 . 37 . 41 . 61 . 89 89 . 197 |
| 2009136133 | 5 . 5 . 13 . 17 . 41 . 61 . 73 . 89 . 97 . 113 . 181 . 197 |

| | |
|---|---|
| 5 | . |
| 13 | 17 |
| 17 | 7. 61 |
| 29 | 43. 217 |
| 37 | 1. 73. 517. 667 |
| 41 | 13. 95 |
| 53 | 191. 233. 763 |
| 61 | 5. 127. 971 |
| 73 | 673. 14543. 31885. 48967. 141709. 387833. 90115783 |
| 89 | 1183. 1717. 4031. 15245. 61117 |
| 97 | 35. 1517. 2557. 5467. 22733. 449921 |
| 101 | 41. 445. 5495. 6827. 7717. 10261 |
| 109 | 89. 347. 565. 5321. 22583. 29995. 2390717 |
| 113 | 23. 203. 1333. 13763. 39347. 74603. 427795. 2525527 |
| 137 | 85. 737. 6217. 29129. 54167. 87217. 837533. 6920333 |
| 149 | 115. 1673. 2201. 2567. 8459. 18659. 38923. 42133. 161035 |
| 157 | 11. 617. 953. 1873. 7547. 7861. 130613. 189353. 533789 |
| 173 | 307. 2383. 4883. 7919. 44327. 49517. 96227. 151163. 256693. 9439957*. 11776417 |
| 181 | 67. 295. 2963. 3553. 4277. 6221. 10565. 13823. 32647. 125909*. 171655. 182743. 206407. 1396529. 5318933 |
| 193 | 293. 479. 3181. 3767. 5497. 6655. 7813. 15733. 22481. 22867*. 62825. 174565. 1097105 |
| 197 | 113. 1463. 2251. 5009. 17617 22345. 23753. 30845. 38893. 191203*. 429347. 726029. 45435967. 70145903. 716295433. 2009136133 |

**Factorable $aa + 49$.**

| | |
|---|---|
| 1 | 5 . 5 |
| 2 | 53 |
| 3 | 29 |
| 4 | 5 . 13 |
| 5 | 37 |
| 6 | 5 . 17 |
| 8 | 113 |
| 9 | 5 . 13 |
| 10 | 149 |
| 11 | 5 . 17 |
| 12 | 193 |
| 13 | 109 |
| 15 | 137 |
| 16 | 5 . 61 |
| 17 | 13 . 13 |
| 19 | 5 . 41 |
| 22 | 13 . 41 |

| | |
|---|---|
| 23 | 17 . 17 |
| 24 | 5 . 5 . 5 . 5 |
| 26 | 5 . 5 . 29 |
| 29 | 5 . 89 |
| 30 | 13 . 73 |
| 31 | 5 . 101 |
| 32 | 29 . 37 |
| 39 | 5 . 157 |
| 40 | 17 . 97 |
| 41 | 5 . 173 |
| 43 | 13 . 73 |
| 45 | 17 . 61 |
| 48 | 13 . 181 |
| 51 | 5 . 5 . 53 |
| 55 | 29 . 53 |
| 57 | 17 . 97 |
| 60 | 41 . 89 |
| 61 | 5 . 13 . 29 |
| 69 | 5 . 13 . 37 |
| 74 | 5 . 5 . 13 . 17 |
| 79 | 5 . 17 . 37 |
| 96 | 5 . 17 . 109 |
| 99 | 5 . 5 . 197 |
| 101 | 5 . 5 . 5 . 41 |
| 103 | 73 . 73 |
| 104 | 5 . 41 . 53 |
| 106 | 5 . 37 . 61 |
| 108 | 13 . 17 . 53 |
| 113 | 13 . 17 . 29 |
| 116 | 5 . 37 . 73 |
| 118 | 89 . 157 |
| 121 | 5 . 13 . 113 |
| 122 | 109 . 137 |
| 132 | 101 . 173 |
| 137 | 97 . 97 |
| 139 | 5 . 13 . 149 |
| 142 | 17 . 29 . 41 |
| 149 | 5 . 5 . 5 . 89 |
| 152 | 13 . 13 . 137 |
| 159 | 5 . 17 . 149 |
| 171 | 5 . 29 . 101 |
| 176 | 5 . 5 . 17 . 73 |
| 181 | 5 . 17 . 193 |
| 186 | 5 . 13 . 13 . 41 |
| 199 | 5 . 5 . 13 . 61 |
| 205 | 109 . 193 |
| 214 | 5 . 53 . 173 |
| 227 | 17 . 37 . 41 |

| | |
|---|---|
| 229 | 5 . 29 . 181 |
| 234 | 5 . 97 . 113 |
| 249 | 5 . 5 . 17 . 73 |
| 251 | 5 . 5 . 13 . 97 |
| 264 | 5 . 13 . 29 . 37 |
| 289 | 5 . 61 . 137 |
| 295 | 13 . 17 . 197 |
| 296 | 5 . 89 . 197 |
| 314 | 5 . 109 . 181 |
| 316 | 5 . 13 . 29 . 53 |
| 321 | 5 . 13 . 13 . 61 |
| 331 | 5 . 97 . 113 |
| 334 | 5 . 13 . 17 . 101 |
| 347 | 13 . 41 . 113 |
| 351 | 5 . 5 . 5 . 17 . 29 |
| 373 | 13 . 53 . 101 |
| 374 | 5 . 5 . 29 . 193 |
| 449 | 5 . 5 . 37 . 109 |
| 474 | 5 . 5 . 89 . 101 |
| 533 | 17 . 61 . 137 |
| 550 | 13 . 17 . 37 . 37 |
| 554 | 5 . 29 . 29 . 73 |
| 555 | 13 . 17 . 17 . 41 |
| 589 | 5 . 13 . 17 . 157 |
| 591 | 5 . 181 . 193 |
| 594 | 5 . 13 . 61 . 89 |
| 601 | 5 . 5 . 5 . 5 . 17 . 17 |
| 606 | 5 . 17 . 29 . 149 |
| 634 | 5 . 37 . 41 . 53 |
| 641 | 5 . 13 . 29 . 109 |
| 667 | 13 . 109 . 157 |
| 676 | 5 . 5 . 101 . 181 |
| 691 | 5 . 17 . 53 . 53 |
| 719 | 5 . 13 . 41 . 97 |
| 776 | 5 . 5 . 13 . 17 . 109 |
| 799 | 5 . 5 . 113 . 113 |
| 809 | 5 . 29 . 37 . 61 |
| 824 | 5 . 5 . 157 . 173 |
| 839 | 5 . 17 . 41 . 101 |
| 844 | 5 . 17 . 17 . 17 . 29 |
| 899 | 5 . 5 . 5 . 53 . 61 |
| 906 | 5 . 13 . 73 . 173 |
| 919 | 5 . 13 . 73 . 89 |
| 979 | 5 . 13 . 73 . 101 |
| 992 | 13 . 17 . 61 . 73 |
| 1009 | 5 . 17 . 53 . 113 |
| 1031 | 5 . 13 . 13 . 17 . 37 |
| 1041 | 5 . 29 . 37 . 101 |

| | |
|---|---|
| 1111 | 5 . 17 . 53 . 137 |
| 1128 | 17 . 29 . 29 . 89 |
| 1179 | 5 . 13 . 17 . 17 37 |
| 1186 | 5 . 29 . 89 . 109 |
| 1221 | 5 . 29 . 53 . 97 |
| 1252 | 13 . 17 . 41 . 173 |
| 1364 | 5 . 37 . 89 . 113 |

| | |
|---|---|
| 1374 | 5 . 5 . 13 . 37 . 157 |
| 1395 | 13 . 29 . 29 . 89 |
| 1556 | 5 . 13 . 193 . 193 |
| 1592 | 17 . 29 . 53 . 97 |
| 1609 | 5 . 17 . 97 . 157 |
| 1621 | 5 . 13 . 17 . 29 . 41 |
| 1629 | 5 . 13 137 . 149 |
| 1649 | 5 . 5 . 5 . 73 . 149 |
| 1689 | 5 . 17 . 97 . 173 |
| 1751 | 5 . 5 . 13 53 . 89 |
| 1766 | 5 . 29 . 137 . 157 |
| 1929 | 5 . 37 89 . 113 |
| 1949 | 5 5 . 17 . 41 . 109 |
| 2069 | 5 . 41 . 53 . 197 |
| 2076 | 5 . 5 . 13 . 89 . 149 |
| 2151 | 5 5 . 37 . 41 . 61 |
| 2374 | 5 . 5 . 17 . 89 . 149 |
| 2381 | 5 . 29 . 113 . 173 |
| 2521 | 5 37 . 89 . 193 |
| 2578 | 13 . 17 . 17 . 29 . 61 |
| 2595 | 17 . 37 . 53 101 |
| 2607 | 17 . 29 . 61 . 113 |
| 2659 | 5 . 37 . 97 . 197 |
| 2817 | 13 . 37 . 73 . 113 |
| 2851 | 5 5 . 5 . 13 41 . 61 |
| 2930 | 17 . 41 . 109 . 113 |
| 2944 | 5 61 . 157 . 181 |
| 2999 | 5 . 5 . 13 . 101 . 137 |
| 3100 | 17 . 29 101 . 193 |
| 3156 | 5 . 17 . 17 . 61 . 113 |
| 3251 | 5 5 . 29 . 37 . 197 |
| 3501 | 5 . 5 . 13 . 109 . 173 |
| 3547 | 17 . 37 . 73 . 137 |
| 3709 | 5 . 13 . 29 . 41 . 89 |
| 3753 | 13 . 41 . 73 . 181 |
| 4034 | 5 . 13 . 29 . 89 . 97 |
| 4065 | 13 . 37 . 89 . 193 |
| 4211 | 5 . 97 . 101 . 181 |
| 4286 | 5 . 13 . 41 . 61 . 113 |

| | |
|---|---|
| 4324 | 5 . 5 . 17 . 29 . 37 . 41 |
| 4399 | 5 . 5 . 5 . 5 . 113 . 137 |
| 4556 | 5 . 29 37 . 53 . 73 |
| 4629 | 5 . 73 . 149 . 197 |
| 4630 | 17 . 37 . 173 197 |
| 4657 | 37 . 37 . 89 . 89 |
| 4715 | 13 . 13 . 17 . 53 . 73 |
| 4749 | 5 . 5 . 13 . 13 . 17 . 157 |
| 4754 | 5 . 13 . 17 . 113 . 181 |
| 4778 | 37 . 41 . 101 . 149 |
| 4918 | 13 . 13 . 13 . 101 . 109 |
| 4927 | 29 . 53 . 53 . 149 |
| 5024 | 5 . 5 . 5 . 5 . 41 . 197 |
| 5101 | 5 . 5 . 5 . 29 . 37 . 97 |
| 5191 | 5 . 13 . 17 . 89 . 137 |
| 5221 | 5 . 101 . 137 . 197 |
| 5226 | 5 . 5 . 5 . 41 . 73 . 73 |
| 5841 | 5 . 13 . 37 . 41 . 173 |
| 6008 | 41 . 41 . 109 . 197 |
| 6029 | 5 . 17 . 29 . 73 . 101 |
| 6309 | 5 . 13 . 53 . 53 . 109 |
| 6381 | 5 . 17 . 17 . 73 . 193 |
| 6574 | 5 . 5 . 13 13 . 53 . 193 |
| 6899 | 5 5 . 5 . 5 . 13 . 29 . 101 |
| 6998 | 13 . 17 . 37 . 53 . 113 |
| 7276 | 5 . 5 . 13 . 29 . 41 . 137 |
| 7316 | 5 . 17 . 53 . 109 . 109 |
| 7440 | 13 . 17 . 41 . 41 . 149 |
| 7914 | 5 . 29 . 61 . 73 97 |
| 8149 | 5 . 5 . 5 . 5 . 5 . 5 . 5 . 5 . 5 . 17 |
| 8219 | 5 . 37 . 41 . 61 . 73 |
| 8251 | 5 . 5 . 13 . 17 . 61 . 101 |
| 8515 | 37 . 89 . 101 . 109 |
| 8753 | 13 . 137 137 . 157 |
| 8919 | 5 . 17 . 41 . 101 . 113 |
| 9161 | 5 . 13 . 29 . 113 . 197 |
| 9301 | 5 . 5 . 73 . 137 . 173 |
| 9546 | 5 . 13 . 97 . 97 . 149 |
| 9616 | 5 . 13 . 13 . 17 . 41 . 157 |
| 9837 | 13 . 17 . 37 . 61 . 97 |
| 9993 | 13 149 149 . 173 |
| 10291 | 5 . 17 . 37 . 113 . 149 |
| 10630 | 13 . 13 . 61 . 97 . 113 |
| 10651 | 5 . 5 . 13 . 37 . 53 . 89 |
| 10727 | 29 . 97 . 113 . 181 |
| 10761 | 5 . 41 . 53 . 73 . 73 |
| 10887 | 41 . 89 . 109 . 149 |
| 11185 | 53 . 89 . 89 . 149 |

| | |
|---|---|
| 11632 | 29 . 149 . 173 . 181 |
| 12151 | 5 . 5 . 13 . 13 . 101 . 173 |
| 12489 | 5 . 13 . 13 . 17 . 61 . 89 |
| 13101 | 5 . 5 . 5 . 5 . 17 . 41 . 197 |
| 13329 | 5 . 13 . 73 . 97 . 113 |
| 14351 | 5 . 5 . 5 . 5 . 37 61 . 73 |
| 14656 | 5 . 41 . 61 . 89 . 193 |
| 14811 | 5 . 13 . 109 . 113 . 137 |
| 15356 | 5 41 . 61 . 109 . 173 |
| 15425 | 17 . 29 . 29 . 53 . 157 |
| 15661 | 5 . 13 . 61 . 157 . 197 |
| 16393 | 61 . 101 . 113 193 |
| 16446 | 5 . 29 . 109 . 109 157 |
| 17247 | 13 . 37 37 . 61 137 |
| 18099 | 5 . 5 . 17 . 29 . 97 . 137 |
| 18976 | 5 . 5 . 5 . 13 . 37 . 53 . 113 |
| 19743 | 13 . 17 . 41 . 137 . 157 |
| 20297 | 13 . 13 . 13 . 29 . 53 . 61 |
| 20999 | 5 . 13 . 29 . 149 . 157 |
| 21768 | 97 . 137 . 181 . 197 |
| 21834 | 5 . 17 . 29 . 41 . 53 . 89 |
| 22156 | 5 . 13 . 13 . 53 . 97 . 113 |
| 22569 | 5 . 41 . 41 . 157 . 193 |
| 25726 | 5 . 5 . 5 . 29 . 41 . 61 . 73. |
| 25931 | 5 . 13 . 17 . 41 . 41 . 181 |
| 26016 | 5 . 17 . 29 . 37 . 41 . 181 |
| 26337 | 113 . 113 . 157 . 173 |
| 27200 | 13 . 37 . 97 . 101 . 157 |
| 27564 | 5 . 13 . 13 . 73 . 109 . 113 |
| 27721 | 5 . 17 . 17 . 29 . 53 . 173 |
| 29254 | 5 . 13 . 13 . 53 . 97 . 197 |
| 29501 | 5 . 5 . 13 . 17 . 17 . 41 . 113 |
| 30179 | 5 . 61 . 73 . 113 . 181. |
| 30424 | 5 . 5 . 13 . 17 . 29 . 53 . 109 |
| 31274 | 5 . 5 . 5 . 5 . 13 . 17 . 73 . 97 |
| 32491 | 5 . 13 . 37 . 41 . 53 . 101 |
| 33258 | 13 . 17 . 17 . 37 . 73 . 109 |
| 34134 | 5 . 13 . 41 . 53 . 73 . 113 |
| 35303 | 17 . 37 . 61 . 109 . 149 |
| 35361 | 5 . 41 . 113 . 137 . 197 |
| 35524 | 5 . 5 . 5 . 17 . 17 . 181 . 193 |
| 36149 | 5 . 5 . 5 . 13 . 13 . 157 . 197 |
| 37409 | 5 . 61 . 89 . 149 . 173 |
| 37836 | 5 . 17 . 17 . 61 . 109 . 149 |
| 38601 | 5 . 5 . 5 . 13 . 17 . 149 . 181 |
| 39901 | 5 . 5 . 13 . 13 . 29 . 73 . 89 |
| 41801 | 5 . 5 . 41 . 61 . 89 . 157 |
| 41859 | 5 . 73 . 89 . 149 . 181 |

| | |
|---|---|
| 43416 | 5 . 13 . 13 . 13 . 29 . 61 . 97 |
| 44677 | 13 . 53 . 97 . 109 . 137 |
| 44976 | 5 . 5 . 5 . 5 . 5 13 . 17 . 29 . 101 |
| 46229 | 5 . 17 . 29 . 41 . 97 . 109 |
| 48317 | 13 . 13 . 29 . 37 . 41 . 157 |
| 49099 | 5 . 5 . 41 . 73 . 89 . 181 |
| 50632 | 17 . 73 . 101 . 113 . 181 |
| 51176 | 5 . 5 . 17 . 17 . 37 . 97 . 101 |
| 54274 | 5 . 5 . 5 . 37 . 53 . 61 . 197 |
| 55774 | 5 . 5 . 5 . 13 . 89 . 137 . 157 |
| 58881 | 5 . 13 . 53 . 61 . 73 . 113 |
| 64051 | 5 . 5 . 73 . 73 . 89 . 173 |
| 64644 | 5 . 29 . 37 . 61 . 113 . 113 |
| 67785 | 17 . 53 . 109 . 149 . 157 |
| 69983 | 13 . 13 . 17 . 61 . 89 . 157 |
| 72851 | 5 . 5 . 5 . 17 . 17 . 17 . 29 . 149 |
| 73043 | 13 . 17 . 37 . 41 . 73 . 109 |
| 73672 | 17 . 17 . 37 . 53 . 61 . 157 |
| 73721 | 5 . 29 . 29 53 . 89 137 |
| 76534 | 5 . 29 . 53 . 53 . 197 |
| 80841 | 5 . 17 . 61 . 73 . 89 . 97 |
| 82307 | 13 . 53 . 157 . 173 . 181 |
| 83430 | 13 . 17 . 29 . 37 . 149 197 |
| 87369 | 5 . 13 . 17 . 89 . 197 . 197 |
| 88213 | 37 . 41 . 97 . 137 . 193 |
| 88406 | 5 . 17 . 53 . 89 . 101 . 193 |
| 88989 | 5 . 13 . 17 . 17 . 41 . 53 . 97 |
| 89149 | 5 . 5 . 5 . 29 . 89 . 109 . 113 |
| 92049 | 5 . 5 . 13 . 17 . 29 . 137 . 193 |
| 102735 | 13 . 13 . 13 . 89 . 137 . 197 |
| 124029 | 5 . 13 . 13 . 37 . 37 . 61 . 109 |
| 131863 | 13 . 17 . 37 . 97 . 97 . 113 |
| 133149 | 5 . 5 . 5 . 5 . 5 . 41 101 . 137 |
| 138199 | 5 . 5 . 13 . 17 . 101 . 109 . 157 |
| 139551 | 5 . 5 . 13 . 29 . 53 . 101 . 193 |
| 141633 | 17 . 17 . 17 . 17 . 29 . 41 . 101 |
| 148439 | 5 . 37 . 41 . 73 101 . 197 |
| 150410 | 17 . 37 . 41 . 61 . 73 . 197 |
| 150681 | 5 . 29 . 53 . 97 . 97 . 157 |
| 154876 | 5 . 5 . 17 . 41 . 73 . 109 . 173 |
| 157995 | 29 . 37 . 41 . 53 . 53 . 101 |
| 160168 | 17 . 37 . 41 . 53 . 137 . 137 |
| 163609 | 5 . 13 . 13 . 37 . 41 . 53 . 137 |
| 166851 | 5 . 5 . 5 . 5 . 5 . 13 . 41 . 61 . 137 |
| 167124 | 5 . 5 . 13 . 13 . 29 . 37 . 61 . 101 |
| 174074 | 5 . 5 . 13 . 17 . 157 . 181 . 193 |
| 178149 | 5 . 5 . 5 . 5 . 17 . 89 . 97 . 173 |
| 179565 | 13 . 17 . 29 . 113 . 113 . 197 |

| | |
|---|---|
| 187249 | 5 . 5 . 17 . 17 . 109 . 113 . 197 |
| 189538 | 73 . 109 . 149 . 157 . 193 |
| 207814 | 5 . 13 . 17 . 17 . 97 . 137 . 173 |
| 213263 | 29 . 37 . 41 . 73 . 73 . 97 |
| 217351 | 5 . 5 . 5 . 13 . 13 . 17 . 17 . 53 . 73 |
| 231755 | 13 . 17 . 17 . 17 . 61 . 61 . 113 |
| 273694 | 5 . 17 . 37 . 41 . 53 . 97 . 113 |
| 281226 | 5 . 5 . 5 . 5 . 61 . 73 . 157 . 181 |
| 288901 | 5 . 5 . 29 . 37 . 53 . 149 . 197 |
| 299399 | 5 . 5 . 5 . 5 . 13 . 29 . 37 . 53 . 97 |
| 307519 | 5 . 13 . 17 . 17 . 29 . 29 . 41 . 73 |
| 343066 | 5 . 13 . 17 . 17 . 17 . 41 . 89 . 101 |
| 345094 | 5 . 13 17 . 37 . 113 . 149 . 173 |
| 361409 | 5 . 13 . 17 . 53 . 61 . 101 . 181 |
| 375967 | 5 . 17 . 37 . 53 . 61 . 149 |
| 401444 | 5 . 13 . 17 . 17 . 29 . 29 . 101 . 101 |
| 408628 | 41 . 53 . 89 . 89 . 89 . 109 |
| 415848 | 13 . 17 . 17 . 37 . 101 . 109 . 113 |
| 417317 | 13 . 41 . 61 . 113 . 137 . 173 |
| 428021 | 5 . 13 . 37 . 41 . 61 . 97 . 157 |
| 448976 | 5 . 5 . 5 . 17 . 61 . 89 . 101 . 173 |
| 462953 | 29 . 41 . 53 . 101 . 113 . 149 |
| 521044 | 5 . 13 . 13 . 17 . 17 . 73 . 97 . 157 |
| 527329 | 5 . 17 . 37 . 61 . 61 . 109 . 109 |
| 658576 | 5 . 5 . 13 . 13 . 13 . 13 . 53 . 73 . 157 |
| 689601 | 5 . 5 . 5 . 37 . 41 . 73 . 89 . 193 |
| 788493 | 13 . 41 . 109 . 157 . 173 . 197 |
| 935601 | 5 . 5 . 5 . 5 . 5 . 13 . 13 . 17 . 29 . 41 . 41 |
| 979976 | 5 . 5 . 5 . 5 . 17 . 17 . 137 . 197 197 |
| 1055864 | 5 . 13 . 17 . 29 . 37 . 53 . 113 . 157 |

| | |
|---|---|
| 1538221 | 5 . 13 . 13 . 29 . 53 . 61 . 109 . 137 |
| 1686759 | 5 . 13 . 29 . 29 . 41 . 41 . 113 . 137 |
| 2001229 | 5 . 13 . 17 . 29 . 41 . 53 . 149 . 193 |
| 2446492 | 13 . 41 . 73 . 89 . 101 . 109 . 157 |
| 3254151 | 5 . 5 . 13 . 17 . 29 . 41 . 61 . 73 . 181 |
| 3297075 | 37 . 41 . 53 . 53 . 73 . 101 . 173 |
| 3643774 | 5 . 5 . 5 . 13 . 17 . 41 . 109 . 137 . 157 |
| 4515359 | 5 . 13 . 13 . 13 . 17 . 17 . 113 . 157 . 181 |
| 5307581 | 5 . 17 . 37 . 53 . 73 . 73 . 101 . 157 |
| 5456999 | 5 . 5 . 41 . 73 . 73 . 101 . 137 . 197 |
| 7936717 | 13 . 29 . 37 . 41 . 41 . 41 . 181 . 181 |
| 8555207 | 13 . 17 . 41 . 53 61 . 73 . 109 . 157 |
| 12448726 | 5 . 5 5 . 5 . 13 . 13 . 13 . 41 . 89 . 157 . 197 |
| 21432319 | 5 . 17 . 37 . 73 . 149 . 173 . 197 . 197 |
| 40407039 | 5 . 17 . 37 . 41 . 53 . 89 . 97 . 101 . 137 |
| 41719774 | 5 . 5 . 5 . 17 . 17 . 29 . 29 . 41 . 41 . 173 . 197 |

118135085 | 17 17 . 37 . 41 . 61 . 101 . 109 . 137 . 173

| | |
|---|---|
| 5 | 1. 24 |
| 13 | 4. 9. 17 |
| 17 | 6. 11. 23. 74. 601. 8149 |
| 29 | 3. 26. 61. 113. 351. 844 |
| 37 | 5. 32. 69. 79. 264. 550. 1031. 1179 |
| 41 | 19. 22. 101. 142. 186. 227. 555. 1621. 4324. 935601* |
| 53 | 2. 51. 55. 104. 108. 316. 634. 691 |
| 61 | 16. 45. 106. 199. 321. 809. 899. 2151. 2578. 2851*. 20297 |
| 73 | 30. 43. 103. 116. 176. 249. 554. 992. 4556. 4715*. 5226. 8219. 10761. 14351. 25726. 217351. 307519 |
| 89 | 29. 60. 149. 594. 919. 1128. 1395. 1751. 3709. 4657*. 10651. 12489. 21834. 39901 |
| 97 | 40. 57. 137. 251. 719. 1221. 1592. 4034. 5101. 7914*. 9837. 31274. 43416. 80841. 88989. 213263. 299399 |
| 101 | 31. 171. 334. 373. 474. 839. 979. 1041. 2595. 6029*. 6899. 8251. 32491. 44976. 51176. 141633. 157995. 167124. 343066. 401444* |
| 109 | 13. 96. 449. 641. 776. 1186. 1949. 4918. 6309. 7316*. 8515. 30424. 33258. 46229. 73043. 124029. 408628. 527329 |
| 113 | 8. 121. 234. 331. 347. 799. 1009. 1364. 1929. 2607*. 2817. 2930. 3156. 4286. 6998. 8919. 10630. 13329. 18976. 22156. 27564*. 29501. 34134. 58881. 64644. 89149. 131863. 231755. 273694. 415848 |
| 137 | 15. 122. 152. 289. 533. 1111. 2999. 3547. 4399. 5191*. 7276. 14811. 17247. 18099. 44677. 73721. 133149. 160168. 163609. 166851*. 1538221. 1686759. 40407039 |
| 149 | 10. 139. 159. 606. 1629. 1649. 2076. 2374. 4778. 4927*. 7440. 9546. 10291. 10887. 11185. 35303. 37836. 72851. 375967. 462953* |
| 157 | 39. 118. 589. 667. 1374. 1609. 1766. 4749. 8753. 9616*. 15425. 16446. 19743. 20999. 27200. 41801. 48317. 55774. 67785. 69983*. 73672. 138199. 150681. 428021. 521044. 658576. 1055864. 2446492. 3643774. 5307581*. 8555207 |
| 173 | 41. 132. 214. 824. 906. 1252. 1689. 2381. 3501. 5841*. 9301. 9993. 12151. 15356. 26337. 27721. 37409. 64051. 154876. 178149*. 207814. 345094. 417317. 448976. 3297075. 118135085 |
| 181 | 48. 229. 314. 676. 2944. 3753. 4211. 4754. 10727. 11632*. 25931. 26016. 30179. 38601. 41859. 49099. 50632. 82307. 281226. 361409*. 3254151. 4515359. 7936717 |
| 193 | 12. 181. 205. 374. 591. 1556. 2521. 3100. 4065. 6381*. 6574. 14656. 16393. 22569. 35524. 88213. 88406. 92049. 139551*. 174074. 189538. 689601. 2001229 |
| 197 | 99. 295. 296. 2069. 2659. 3251. 4629. 4630. 5024. 5221*. 6008. 9161. 13101. 15661. 21768. 29254. 35361. 36149. 54274. 76534*. 83430. 87369. 102735. 148439. 150410. 179565. 187249. 288901. 788493. 979976*. 5456999. 12448726. 21432319. 41719774 |

**Factorable $aa + 64$.**

| | |
|---|---|
| 1 | 5 . 13 |
| 3 | 73 |
| 5 | 89 |
| 7 | 113 |
| 9 | 5 . 29 |
| 11 | 5 . 37 |
| 15 | 17 . 17 |
| 19 | 5 . 5 . 17 |
| 21 | 5 . 101 |
| 25 | 13 . 53 |
| 27 | 13 . 61 |
| 29 | 5 . 181 |
| 31 | 5 . 5 . 41 |
| 49 | 5 . 17 . 29 |
| 51 | 5 . 13 . 41 |
| 53 | 13 . 13 . 17 |
| 63 | 37 . 109 |
| 67 | 29 . 157 |
| 69 | 5, 5 . 193 |
| 79 | 5 . 13 . 97 |
| 81 | 5 . 5 . 5 . 53 |
| 85 | 37 . 197 |
| 95 | 61 . 149 |
| 115 | 97 . 137 |
| 121 | 5 . 17 . 173 |
| 131 | 5 . 5 . 13 . 53 |
| 149 | 5 . 61 . 73 |
| 155 | 13 . 17 . 109 |
| 159 | 5 . 37 . 137 |
| 181 | 5 . 5 . 13 . 101 |
| 183 | 13 . 29 . 89 |
| 219 | 5 . 5 . 17 . 113 |
| 223 | 17 . 29 . 101 |
| 233 | 5 . 37 . 113 |
| 281 | 5 . 5 . 29 . 109 |
| 309 | 5 . 97 . 197 |
| 339 | 5 . 13 . 29 . 61 |
| 359 | 5 . 17 . 37 . 41 |
| 381 | 5 . 5 . 37 . 157 |
| 389 | 5 . 13 . 17 . 137 |
| 391 | 5 . 13 . 13 . 181 |
| 393 | 17 . 61 . 149 |
| 433 | 37 . 37 . 137 |
| 441 | 5 . 13 . 41 . 73 |
| 455 | 29 . 37 . 193 |

| | |
|---|---|
| 461 | 5 . 17 . 41 . 61 |
| 467 | 13 . 97 . 173 |
| 529 | 5 . 17 . 37 . 89 |
| 571 | 5 . 13 . 29 . 173 |
| 581 | 5 . 5 . 5 . 37 . 73 |
| 661 | 5 . 17 . 53 . 97 |
| 703 | 13 . 193 . 197 |
| 707 | 41 . 89 . 137 |
| 717 | 53 . 89 . 109 |
| 729 | 5 . 13 . 13 . 17 . 37 |
| 831 | 5 . 5 . 5 . 5 . 5 . 13 . 17 |
| 873 | 53 . 73 . 197 |
| 879 | 5 . 29 . 73 . 73 |

| | |
|---|---|
| 911 | 5 . 13 . 113 . 113 |
| 937 | 13 . 17 . 29 . 137 |
| 989 | 5 . 13 . 101 . 149 |
| 1035 | 17 . 29 . 41 . 53 |
| 1097 | 41 . 149 . 197 |
| 1141 | 5 . 17 . 17 . 17 . 53 |
| 1169 | 5 . 5 . 5 . 13 . 29 . 29 |
| 1171 | 5 . 13 . 17 . 17 . 73 |
| 1191 | 5 . 53 . 53 . 101 |
| 1247 | 13 . 37 . 53 . 61 |
| 1343 | 29 . 37 . 41 . 41 |
| 1419 | 5 . 5 . 5 . 89 . 181 |
| 1491 | 5 . 37 . 61 . 197 |
| 1589 | 5 . 41 . 109 . 113 |
| 1609 | 5 . 41 . 73 . 173 |
| 1613 | 13 . 17 . 61 . 193 |
| 1637 | 13 13 . 101 . 157 |
| 1681 | 5 . 5 . 17 . 61 . 109 |
| 1691 | 5 . 13 . 29 . 37 . 41 |
| 1749 | 5 . 17 . 17 . 29 . 73 |
| 1839 | 5 . 37 . 101 . 181 |
| 1861 | 5 . 37 . 97 . 193 |
| 1999 | 5 . 41 . 101 . 193 |
| 2019 | 5 . 5 . 41 . 41 . 97 |
| 2041 | 5 . 73 . 101 . 113 |
| 2055 | 13 . 17 . 97 . 197 |
| 2081 | 5 . 5 . 5 . 5 . 13 . 13 41 |
| 2131 | 5 . 5 . 13 . 89 . 157 |
| 2201 | 5 . 53 . 101 . 181 |
| 2445 | 13 . 29 . 101 . 157 |
| 2479 | 5 . 73 . 113 . 149 |
| 2601 | 5 . 13 . 29 . 37 . 97 |
| 2625 | 5 . 13 . 53 . 73 . 137 |

| | |
|---|---|
| 2771 | 5 . 73 . 109 . 193 |
| 2989 | 5 . 13 . 13 . 97 . 109 |
| 3171 | 5 . 13 . 37 . 37 . 113 |
| 3199 | 5 . 13 . 29 . 61 . 89 |
| 3413 | 29 . 41 . 97 . 101 |
| 3721 | 5 . 17 . 29 . 41 . 137 |
| 4031 | 5 . 5 . 13 . 17 . 17 . 173 |
| 4061 | 5 . 17 . 17 . 101 . 113 |
| 4109 | 5 . 13 . 13 . 13 . 29 . 53 |
| 4315 | 13 41 . 181 . 193 |
| 4419 | 5 . 5 . 5 . 13 . 61 . 197 |
| 4541 | 5 . 17 . 41 . 61 . 97 |
| 4979 | 5 . 29 . 37 . 97 . 157 |
| 5097 | 13 . 61 . 181 . 181 |
| 5169 | 5 . 5 . 5 . 37 . 53 . 109 |
| 5381 | 5 . 5 . 13 . 41 . 41 . 53 |
| 5459 | 5 . 13 . 17 . 149 . 181 |
| 5761 | 5 . 17 . 37 . 61 . 173 |
| 5869 | 5 . 5 . 89 . 113 . 137 |
| 6081 | 5 . 5 . 5 . 29 . 101 . 101 |
| 7781 | 5 . 5 . 29 . 37 . 37 . 61 |
| 9039 | 5 . 29 . 37 . 97 . 157 |
| 9779 | 5 . 37 . 73 . 73 . 97 |
| 10519 | 5 . 37 . 37 . 53 . 61 |
| 10527 | 41 . 109 . 137 . 181 |
| 10831 | 5 . 5 . 5 . 5 . 17 . 61 . 181 |
| 11511 | 5 . 17 . 41 . 193 . 197 |
| 13331 | 5 . 5 . 5 . 5 . 5 . 29 . 37 . 53 |
| 13581 | 5 . 5 . 5 . 17 . 29 41 . 73 |
| 13889 | 5 . 41 . 89 . 97 . 109 |
| 13911 | 5 . 13 . 13 . 29 . 53 . 149 |
| 14451 | 5 . 29 . 73 . 109 . 181 |
| 16069 | 5 . 5 . 13 . 37 . 109 . 197 |
| 16985 | 17 . 29 . 53 . 61 . 181 |
| 17421 | 5 . 13 . 137 . 173 . 197 |
| 18511 | 5 . 13 . 17 . 17 . 17 . 29 . 37 |
| 18563 | 13 . 37 . 41 . 101 . 173 |
| 18685 | 17 . 29 . 73 . 89 . 109 |
| 24061 | 5 . 29 . 29 . 37 . 61 . 61 |
| 24261 | 5 . 17 . 17 . 37 . 101 . 109 |
| 27665 | 13 37 . 41 . 197 . 197 |
| 29019 | 5 . 5 . 37 . 53 . 89 . 193 |
| 29981 | 5 . 5 . 37 . 41 . 137 . 173 |
| 31669 | 5 . 5 . 5 . 5 . 13 . 17 . 53 . 137 |
| 58397 | 13 . 17 . 29 . 37 . 73 . 197 |
| 59279 | 5 . 13 . 53 . 73 . 89 . 157 |
| 88789 | 5 . 13 17 . 29 . 37 . 61 . 109 |
| 103481 | 5 . 5 . 13 . 13 . 17 . 29 . 53 . 97 |

| | |
|---|---|
| 132081 | 5 . 5 . 5 . 5 . 5 . 5 . 5 . 13 . 89 . 193 |
| 170529 | 5 . 17 . 29 . 41 . 53 . 61 . 89 |
| 213331 | 5 . 5 . 5 . 5 . 5 . 13 . 13 . 17 . 37 . 137 |
| 383229 | 5 . 17 . 17 . 61 . 89 . 97 . 193 |
| 728391 | 5 . 13 . 73 . 97 . 101 . 101 . 113 |
| 1934581 | 5 . 5 . 5 . 5 . 13 . 17 . 17 . 17 . 29 . 53 . 61 |
| 2446081 | 5 . 5 . 5 . 13 . 17 . 17 . 37 . 53 . 73 . 89 |
| 4056181 | 5 . 5 . 13 . 17 . 29 . 41 . 101 . 137 . 181 |
| 5106581 | 5 . 5 . 5 . 13 . 17 . 37 . 41 . 61 . 101 . 101 |
| 14836119 | 5 . 5 . 13 . 13 . 17 . 17 . 17 . 17 . 29 . 137 . 157 |

| | |
|---|---|
| 13 | 1 |
| 17 | 15. 19. 53. 831 |
| 29 | 9. 49. 1169 |
| 37 | 11. 729. 18511 |
| 41 | 31. 51. 359. 1343. 1691. 2081 |
| 53 | 25. 81. 131. 1035. 1141. 4109. 5381. 13331 |
| 61 | 27. 339. 461. 1247. 7781. 10519. 24061. 1934581 |
| 73 | 3. 149. 441. 581. 879. 1171. 1749. 13581 |
| 89 | 5. 183. 529. 3199. 170529. 2446081 |
| 97 | 79.661. 2019. 2601. 4541. 9779. 103481 |
| 101 | 21. 181. 223. 1191. 3413. 6081. 5106581 |
| 109 | 63. 155. 281. 717. 1681. 2989. 5169. 13889. 18685. 24261*. 88789 |
| 113 | 7. 219. 233. 911. 1589. 2041. 3171. 4061. 728391 |
| 137 | 115. 159. 389. 433. 707. 937. 2625. 3721. 5869. 31669*. 213331 |
| 149 | 95. 393. 989. 2479. 13911 |
| 157 | 67. 381. 1637. 2131. 2445. 4979. 9039. 59279. 1483619 |
| 173 | 121. 467. 571. 1609. 4031. 5761. 18563. 29981 |
| 181 | 29. 391. 1419. 1839. 2201. 5097. 5459. 10527. 10831. 14451*. 16985. 4056181 |
| 193 | 69. 455. 1613. 1861. 1999. 2771. 4315. 29019. 132081. 383229 |
| 197 | 85. 309. 703. 873. 1097. 1491. 2055. 4419. 11511. 16069*. 17421. 27665. 58397 |

**Factorable $aa + 81$.**

| | |
|---|---|
| 1 | 41 |
| 2 | 5 . 17 |
| 4 | 97 |
| 5 | 53 |
| 7 | 5 . 13 |
| 8 | 5 . 29 |
| 10 | 181 |
| 11 | 101 |

| | |
|---|---|
| 13 | 5 . 5 . 5 |
| 17 | 5 . 37 |
| 19 | 13 . 17 |
| 20 | 13 . 37 |
| 22 | 5 . 113 |
| 23 | 5 . 61 |
| 28 | 5 . 173 |
| 32 | 5 . 13 . 17 |
| 37 | 5 . 5 . 29 |
| 38 | 5 . 5 61 |
| 40 | 41 . 41 |
| 43 | 5 . 193 |
| 49 | 17 . 73 |
| 50 | 29 . 89 |
| 53 | 5 . 17 . 17 |
| 58 | 5 . 13 . 53 |
| 59 | 13 . 137 |
| 62 | 5 . 5 . 157 |
| 71 | 13 . 197 |
| 79 | 29 . 109 |
| 83 | 5 . 17 . 41 |
| 91 | 37 . 113 |
| 95 | 29 . 157 |
| 97 | 5 · 13 . 73 |
| 98 | 5 . 13 . 149 |
| 101 | 53 . 97 |
| 112 | 5 . 5 . 5 . 101 |
| 122 | 5 . 41 . 73 |
| 124 | 13 . 29 . 41 |
| 128 | 5 . 37 . 89 |
| 137 | 5 . 5 . 13 . 29 |
| 139 | 89 . 109 |
| 163 | 5 . 5 . 13 . 41 |
| 188 | 5 . 5 . 13 . 109 |
| 191 | 101 . 181 |
| 202 | 5 . 13 . 17 . 37 |
| 215 | 13 . 13 . 137 |
| 217 | 5 . 53 . 89 |
| 247 | 5 . 41 . 149 |
| 248 | 5 . 109 . 113 |
| 253 | 5 . 13 . 17 . 29 |
| 268 | 5 . 73 . 197 |
| 287 | 5 . 5 . 17 . 97 |
| 292 | 5 . 13 . 13 . 101 |
| 313 | 5 . 5 . 37 . 53 |
| 317 | 5 . 89 . 113 |
| 323 | 5 . 53 . 197 |
| 352 | 5 . 137 . 181 |

| | |
|---|---|
| 376 | 17 . 53 . 157 |
| 389 | 17 . 61 . 73 |
| 406 | 17 . 89 . 109 |
| 409 | 13 . 41 . 157 |
| 427 | 5 . 17 . 29 . 37 |
| 461 | 13 . 13 . 17 . 37 |
| 472 | 5 . 29 . 29 . 53 |
| 487 | 5 . 5 . 5 . 13 . 73 |
| 491 | 17 . 41 . 173 |
| 533 | 5 . 157 . 181 |
| 547 | 5 . 173 . 173 |
| 553 | 5 . 13 . 13 . 181 |
| 566 | 13 . 157 . 157 |
| 575 | 37 . 41 . 109 |
| 578 | 5 . 13 . 53 . 97 |
| 587 | 5 . 5 . 61 . 113 |
| 617 | 5 . 13 . 29 . 101 |
| 631 | 13 . 17 . 17 . 53 |
| 662 | 5 . 5 . 89 . 197 |
| 683 | 5 . 13 . 37 . 97 |
| 694 | 53 . 61 . 149 |
| 733 | 5 . 17 . 29 . 109 |
| 737 | 5 . 5 . 5 . 41 . 53 |
| 763 | 5 . 5 . 5 . 17 . 137 |
| 797 | 5 . 17 . 37 . 101 |
| 862 | 5 . 5 . 5 . 5 . 29 . 41 |
| 877 | 5 . 13 . 61 . 97 |
| 920 | 17 . 17 . 29 . 101 |
| 1018 | 5 . 17 . 89 . 137 |
| 1037 | 5 . 5 . 137 . 157 |
| 1060 | 13 . 13 . 61 . 109 |
| 1108 | 5 . 41 . 53 . 113 |
| 1118 | 5 . 53 . 53 . 89 |
| 1168 | 5 . 29 . 97 . 97 |
| 1201 | 37 . 101 . 193 |
| 1229 | 13 . 13 . 41 . 109 |
| 1243 | 5 . 17 . 61 . 149 |
| 1265 | 73 . 97 . 113 |
| 1277 | 5 . 17 . 53 . 181 |
| 1313 | 5 . 5 . 29 . 29 . 41 |
| 1436 | 13 . 41 . 53 . 73 |
| 1447 | 5 . 17 . 109 . 113 |
| 1463 | 5 . 5 . 13 . 37 . 89 |
| 1475 | 13 . 13 . 41 . 157 |
| 1487 | 5 . 5 . 5 . 5 . 29 . 61 |
| 1585 | 53 . 137 . 173 |
| 1639 | 41 . 181 . 181 |
| 1645 | 13 . 29 . 37 . 97 |

| | |
|---|---|
| 1685 | 17 . 37 . 37 . 61 |
| 1702 | 5 . 17 . 173 . 197 |
| 1703 | 5 . 29 . 73 . 137 |
| 1787 | 5 . 5 . 13 . 17 . 17 . 17 |
| 1807 | 5 . 53 . 61 . 101 |
| 2008 | 5 . 13 . 17 . 41 . 89 |
| 2009 | 13 . 29 . 53 . 101 |
| 2041 | 97 . 109 . 197 |
| 2051 | 29 . 29 . 41 . 61 |
| 2125 | 13 . 29 . 53 . 113 |
| 2138 | 5 . 5 . 5 . 13 . 29 . 97 |
| 2293 | 5 . 17 . 157 . 197 |
| 2312 | 5 . 5 . 29 . 73 . 101 |
| 2450 | 13 . 17 . 157 . 173 |
| 2531 | 17 . 29 . 73 . 89 |
| 2623 | 5 . 41 . 97 . 173 |
| 2705 | 17 . 29 . 41 . 181 |
| 2888 | 5 . 5 . 5 . 5 . 5 . 17 . 157 |
| 2906 | 13 . 37 . 97 . 181 |
| 3088 | 5 . 5 . 13 . 13 . 37 . 61 |
| 3238 | 5 . 5 . 41 . 53 . 193 |
| 3322 | 5 . 13 . 41 . 41 . 101 |
| 3347 | 5 . 13 . 17 . 37 . 137 |
| 3503 | 5 . 13 . 13 . 53 . 137 |
| 3517 | 5 . 13 . 17 . 29 . 193 |
| 3791 | 29 . 37 . 37 . 181 |
| 3988 | 5 . 5 . 29 . 97 . 173 |
| 4010 | 13 . 13 . 17 . 29 . 193 |
| 4112 | 5 . 5 . 5 . 17 . 73 . 109 |
| 4354 | 17 . 61 . 101 . 181 |
| 4388 | 5 . 5 . 5 . 13 . 17 . 17 . 41 |
| 4429 | 29 . 41 . 73 . 113 |
| 4648 | 5 . 13 . 29 . 73 . 157 |
| 4657 | 5 . 101 . 109 . 197 |
| 4837 | 5 . 5 . 41 . 101 . 113 |
| 5083 | 5 . 29 . 41 . 41 . 53 |
| 5162 | 5 . 5 . 61 . 101 . 173 |
| 5557 | 5 . 13 . 17 . 89 . 157 |
| 5747 | 5 . 109 . 157 . 193 |
| 5792 | 5 . 13 . 13 . 29 . 37 . 37 |
| 5833 | 5 . 17 . 17 . 61 . 193 |
| 6013 | 5 . 5 . 5 . 5 . 5 . 5 . 13 . 89 |
| 6233 | 5 . 13 . 37 . 41 . 197 |
| 6458 | 5 . 17 . 37 . 89 . 149 |
| 6532 | 5 . 5 . 37 . 113 . 157 |
| 6689 | 13 . 97 . 113 . 157 |
| 6883 | 5 . 13 . 13 . 17 . 17 . 97 |
| 7097 | 5 . 29 . 29 . 53 113 |

| | |
|---|---|
| 7160 | 53 . 61 . 101 . 157 |
| 7793 | 5 . 13 . 29 . 89 . 181 |
| 8158 | 5 . 13 . 13 . 17 . 41 . 113 |
| 8273 | 5 . 29 . 53 . 61 . 73 |
| 8638 | 5 . 5 . 5 . 13 . 17 . 37 . 73 |
| 9028 | 5 . 13 . 73 . 89 . 193 |
| 9101 | 41 . 73 . 101 . 137 |
| 9295 | 61 . 73 89 . 109 |
| 9562 | 5 . 5 . 13 . 29 . 89 . 109 |
| 9587 | 5 . 5 . 13 . 13 . 73 . 149 |
| 9607 | 5 . 17 . 29 . 97 . 193 |
| 10118 | 5 . 37 . 53 . 53 . 197 |
| 10577 | 5 . 29 . 41 . 97 . 97 |
| 11563 | 5 . 5 13 29 . 41 . 173 |
| 12143 | 5 . 29 . 29 . 89 . 197 |
| 12565 | 13 . 17 . 29 . 109 . 113 |
| 15233 | 5 . 29 . 73 . 97 . 113 |
| 16237 | 5 . 5 . 5 . 17 . 17 . 41 . 89 |
| 16522 | 5 . 17 . 29 . 37 . 41 . 73 |
| 16777 | 5 . 13 . 13 . 17 . 97 . 101 |
| 17888 | 5 . 5 . 5 5 . 37 . 101 . 137 |
| 17972 | 5 . 13 . 29 . 53 . 53 . 61 |
| 18887 | 5 . 5 . 29 . 37 . 61 . 109 |
| 18974 | 13 . 13 . 17 . 29 . 29 . 149 |
| 19558 | 5 . 13 . 13 . 41 . 61 . 181 |
| 20362 | 5 . 5 . 5 . 113 . 149 . 197 |
| 23368 | 5 . 13 . 13 . 53 . 89 . 137 |
| 24083 | 5 . 13 . 157 . 157 . 181 |
| 24499 | 13 . 17 . 61 . 113 . 197 |
| 24958 | 5 . 17 . 37 . 37 . 53 . 101 |
| 29153 | 5 . 13 . 17 . 29 . 89 . 149 |
| 29242 | 5 . 17 . 17 . 61 . 89 . 109 |
| 29765 | 17 . 37 . 41 . 89 . 193 |
| 30218 | 5 . 13 . 41 . 41 . 61 . 137 |
| 31487 | 5 . 5 . 5 . 5 . 5 . 41 . 53 . 73 |
| 31843 | 5 . 13 . 17 . 17 . 137 . 197 |
| 34763 | 5 . 5 . 5 . 5 . 17 . 29 . 37 . 53 |
| 35137 | 5 . 5 . 17 . 73 . 101 . 197 |
| 35783 | 5 . 13 . 17 . 17 . 173 . 197 |
| 37147 | 5 . 13 . 17 . 41 . 97 . 157 |
| 38201 | 13 . 17 . 17 . 29 . 37 . 181 |
| 44987 | 5 . 5 . 5 . 13 . 29 . 109 . 197 |
| 46963 | 5 . 5 . 13 . 29 . 29 . 53 . 73 |
| 56387 | 5 . 5 . 13 . 17 . 53 . 61 . 89 |
| 57037 | 5 . 5 . 13 . 17 . 37 . 73 . 109 |
| 60743 | 5 . 13 . 17 . 17 . 17 . 53 . 109 |
| 61337 | 5 . 5 . 41 . 109 . 113 . 149 |
| 69107 | 5 . 17 . 53 . 53 . 73 . 137 |

| | |
|---|---|
| 69244 | 13 . 13 . 29 . 37 . 137 . 193 |
| 79813 | 5 . 5 . 13 . 17 . 53 . 73 . 149 |
| 86528 | 5 . 17 . 29 . 97 . 173 . 181 |
| 87263 | 5 . 5 . 5 . 5 . 5 . 13 . 17 . 37 . 149 |
| 97577 | 5 . 29 . 61 . 73 . 73 . 101 |
| 98063 | 5 . 5 . 73 . 97 . 157 . 173 |
| 121933 | 5 . 13 41 . 97 . 149 . 193 |
| 132683 | 5 . 17 . 29 . 109 . 181 . 181 |
| 157723 | 5 . 13 . 13 . 29 . 53 . 61 . 157 |
| 168703 | 5 . 37 . 37 . 61 . 173 . 197 |
| 181508 | 5 . 41 . 41 . 101 . 197 . 197 |
| 195787 | 5 . 5 . 13 . 17 . 29 . 37 . 53 . 61 |
| 237322 | 5 . 13 . 13 . 13 . 17 . 17 . 113 . 157 |
| 269861 | 13 . 37 . 41 . 101 . 101 . 181 |
| 278297 | 5 . 5 . 13 . 37 . 97 . 113 . 113 |
| 297212 | 5 . 5 . 13 . 97 . 113 . 137 . 193 |
| 314387 | 5 . 5 . 41 . 41 . 73 . 89 . 181 |
| 327737 | 5 . 5 . 5 . 5 . 13 . 13 . 29 . 89 . 197 |

| | |
|---|---|
| 349487 | 5 . 5 . 5 . 29 . 29 . 53 . 97 . 113 |
| 474013 | 5 . 5 . 5 . 13 . 17 . 17 . 29 . 73 . 113. |
| 609161 | 13 . 73 . 73 . 113 . 137 . 173 |
| 647665 | 29 . 37 . 53 . 97 . 193 . 197 |
| 934862 | 5 . 5 . 5 . 13 . 13 . 17 . 17 . 37 . 53 . 73 |
| 1125533 | 5 . 13 . 41 . 89 . 101 . 137 . 193 |
| 1158413 | 5 . 5 . 29 . 37 . 41 . 61 . 73 . 137 |
| 1880912 | 5 . 5 . 13 . 17 . 37 . 53 . 53 . 61 . 101 |
| 2023513 | 5 . 5 . 5 . 5 . 37 . 41 . 97 . 113 . 197 |
| 2092285 | 17 . 17 . 37 . 37 . 37 . 41 . 41 . 89 |
| 2898587 | 5 . 5 . 17 . 29 . 73 . 137 . 173 . 197 |
| 3559861 | 13 . 37 . 61 . 89 . 109 . 113 . 197 |
| 4034153 | 5 . 13 . 13 . 17 . 29 . 41 . 53 . 89 . 101 |
| 4676188 | 5 . 5 . 17 . 37 . 37 . 37 . 89 . 101 . 113 |
| 4802183 | 5 . 73 . 97 . 109 . 113 . 137 . 193 |
| 4947916 | 17 . 17 . 37 . 53 . 61 . 73 . 89 . 109 |
| 6678737 | 5 . 5 . 5 . 17 . 17 . 29 . 41 . 53 . 97 . 101 |
| 9578563 | 5 . 5 . 13 . 17 . 17 . 17 . 29 . 61 . 109 . 149 |
| 34928797 | 5 . 13 . 13 . 17 . 29 . 37 . 53 . 53 . 73 . 193 |
| 59554033 | 5 . 13 . 13 . 13 . 37 . 61 . 73 . 89 . 101 . 109 |

| | |
|---|---|
| 5 | 13 |
| 13 | 7 |
| 17 | 2. 19. 32. 53. 1787 |
| 29 | 8. 37. 137. 253 |
| 37 | 17. 20. 202. 427. 461. 5792 |

| | |
|---|---|
| 41 | 1. 40. 83. 124. 163. 862. 1313. 4388 |
| 53 | 5. 58. 313. 472. 631. 737. 5083. 34763 |
| 61 | 23. 38. 1487. 1685. 2051. 3088. 17972. 195787 |
| 73 | 49. 97. 122. 389. 487. 1436. 8273. 8638. 16522. 31487. 46963. 934862 |
| 89 | 50. 128. 217. 1118. 1463. 2008. 2531. 6013. 16237. 56387. 2092285 |
| 97 | 4. 101. 287. 578. 683. 877. 1168. 1645. 2138. 6883. 10577 |
| 101 | 11. 112. 292. 617. 797. 920.1807. 2009. 2312. 3322. 16777. 24958. 97577. 1880912. 4034153. 6678737 |
| 109 | 79. 139. 188. 406. 575. 733. 1060. 1229. 4112. 9295. 9562. 18887. 29242. 57037. 60743. 4947916. 59554033 |
| 113 | 22. 91. 248. 317. 587. 1108. 1265. 1447. 2125. 4429. 4837. 7097. 8158. 12565. 15233. 278297. 349487. 474013. 4676188 |
| 137 | 59. 215. 763. 1018. 1703. 3347. 3503. 9101. 17888. 23368. 30218. 69107. 1158413 |
| 149 | 98. 247. 694. 1243. 6458. 9587. 18974. 29153. 61337. 79813. 87263. 9578563 |
| 157 | 62. 95. 376. 409. 566. 1037. 1475. 2888. 4648. 5557. 6532. 6689. 7160. 37147. 157723. 237322 |
| 173 | 28. 491. 547. 1585. 2450. 2623. 3988. 5162. 11563. 98063. 609161 |
| 181 | 10. 191. 352. 533. 553. 1277. 1639. 2705. 2906. 3791. 4354. 7793. 19558. 24083. 38201. 86528. 132683. 269861. 314387 |
| 193 | 43. 1201. 3238. 3517. 4010. 5747. 5833. 9028. 9607. 29765. 69244. 121933. 297212. 1125533. 4802183. 34928797 |
| 197 | 71. 268. 323. 662. 1702. 2041. 2293. 4657. 6233. 10118. 12143. 20362. 24499. 31843. 35137. 35783. 44987. 168703. 181508. 327737. 647665. 2023513. 2898587. 3559861 |

# A NEW QUADRATURE OF THE CIRCLE.

## I. Acotg .5 = (2) − (13).

| | | | | | | |
|---|---|---|---|---|---|---|
| 26684971 | 0210071265 | 7279572372 | 8848455203 | 4183360499 | 1499728341 | 833 |
| | 364 | 7220869565 | 2173913043 | 4782608695 | 6521739130 | 434 |
| | | 4977954329 | 5694145758 | 6618876941 | 4575866188 | 769 |
| | | (43) . . . . . | 204560302 | 8420465116 | 2790697674 | 418 |
| | | (47) . . . . . | 299441 | 4645858042 | 5531914893 | 617 |
| | | (51) . . . . . | 441 | 5293752324 | 0156862745 | 098 |
| | | (55) . . . . . | 6550690367 | 0843578181 | 818 | |
| | | (59) . . . . . | 9770521 | 2254817540 | 338 | |
| | | (63) . . . . . | 14640 | 2730743726 | 600 | |
| | | (67) . . . . . | 22 | 0259630730 | 860 | |
| | | (71) . . . . . | 332561019 | 920 | | |
| | | (75) . . . . . | 503719 | 091 | | |
| | | (79) . . . . . | 765 | 142 | | |
| | | (83) . . . . . | 1 | 165 | | |
| | | (87) . . . . . | 1 | | | |
| 26684971 | 0210071630 | 9478396268 | 6921374192 | 0505397169 | 7498644659 | 104 |
| 0,2000640569 | 5190437336 | 8654456654 | 4566544566 | 5445665445 | 6654456654 | 456 |
| | 7710131 | 0688316235 | 2941176470 | 5882352941 | 1764705882 | 352 |
| | | 185127900 | 6896551724 | 1379310344 | 8275862068 | 965 |
| | | 266673 | 0412205651 | 4831745270 | 7696610135 | 634 |
| | | | 7830238 | 1276333963 | 9002267573 | 696 |
| | | (53) . . . . . | 16 | 9947155749 | 8300377358 | 490 |
| | | (57) . . . . . | 252833663 | 2909752140 | 350 | |
| | | (61) . . . . . | 378007 | 0506907695 | 003 | |
| | | (65) . . . . . | 567 | 5921253449 | 092 | |
| | | (69) . . . . . | 8555011744 | 329 | | |
| | | (73) . . . . . | 12937990 | 364 | | |
| | | (77) . . . . . | 19625 | 419 | | |
| | | (81) . . . . . | 29 | 850 | | |
| | | (85) . . . . . | 45 | | | |
| 0,2000640569 | 5198147467 | 9528161463 | 4824308667 | 9015775954 | 9600162348 | 045 |
| 26684971 | 0210071630 | 9478396268 | 6921374192 | 0505397169 | 7498644659 | 104 |
| 0,1973955598 | 4988075837 | 0049765194 | 7902934475 | 8510378785 | 2101517688 | 941 |

## II. $\operatorname{Acotg} .70 = 2(2) - 2(13) - (29)$

| | | | | | | |
|---|---|---|---|---|---|---|
| 0,0142857142 | 8571428571 | 4285714285 | 7142857142 | 8571428571 | 4285714285 | 714 |
| 3) 29154 | 5189504373 | 1778425655 | 9766763848 | 3965014577 | 2594752186 | 588 |
| 5) 5 | 9499018266 | 1986077229 | 7257095257 | 9282441839 | 7096447908 | 609 |
| 7) 12142656 | 7890201240 | 2509644305 | 1546792335 | 0693284989 | 369 | |
| 9) 2478 | 0932222490 | 0490308090 | 6745213631 | 0887896588 | 773 | |
| 11) 5057333106 | 6306222511 | 8552396982 | 3736915897 | 263 | | |
| 13) 1032108 | 7972715555 | 6146643346 | 3229334064 | 468 | | |
| 15) 210 | 6344484227 | 6644111559 | 8665965170 | 217 | | |
| 17) 429866221 | 2709519206 | 4407891013 | 300 | | | |
| 19) 87727 | 8002593779 | 4298858753 | 268 | | | |
| 21) 17 | 9036327059 | 9549856909 | 949 | | | |
| 23) 36538025 | 9306030583 | 042 | | | | |
| 25) 7456 | 7399858373 | 588 | | | | |
| 27) 1 | 5217836705 | 790 | | | | |
| 29) 3105680 | 960 | | | | | |
| 31) 633 | 812 | | | | | |
| 33) 128 | | | | | | |
| 9718 | 1729834791 | 0592808551 | 9922254616 | 1321671525 | 7531584062 | 196 |
| 1734665 | 2555743034 | 3215663472 | 1649541762 | 1527612141 | 338 | |
| 459757555 | 1482383864 | 7141126998 | 3976083263 | 387 | | |
| 14 | 0422965615 | 1776274103 | 9911064344 | 681 | | |
| 4617 | 2526452304 | 1805203092 | 277 | | | |
| 1588609 | 8230696981 | 871 | | | | |
| 563623581 | 695 | | | | | |
| 20 | 447 | | | | | |
| 9718 | 1731569456 | 3608309155 | 5043272185 | 4416655304 | 3545867487 | 890 |
| 0,0142857142 | 8571428571 | 4285714285 | 7142857142 | 8571428571 | 4285714285 | 714 |
| 1 | 1899803653 | 2397215445 | 9451419051 | 5856488367 | 9419289581 | 721 |
| 275 | 3436913610 | 0054478676 | 7416134847 | 8987544065 | 419 | |
| 79392 | 9844055042 | 7395895642 | 0248410312 | 651 | | |
| 25286248 | 3100559953 | 3200464177 | 252 | | | |
| 8525539383 | 8073802709 | 997 | | | | |
| 298 | 2695994334 | 943 | | | | |
| 107092 | 446 | | | | | |
| 3 | | | | | | |
| 0,0142857144 | 0471232500 | 0119922734 | 6518096163 | 0866047064 | 6911326560 | 146 |
| 9718 | 1731569456 | 3608309155 | 5043272185 | 4416655304 | 3545867487 | 890 |
| 0,0142847425 | 8739663043 | 6511613579 | 1474823977 | 6449391760 | 3365459072 | 256 |

III. $\operatorname{Acotg} .99 = -(2) + 2(13) - (29).$

| | | | | | | |
|---|---|---|---|---|---|---|
| 0,0101010101 | 0101010101 | 0101010101 | 0101010101 | 0101010101 | 0101010101 | 010 |
| 3) 10306 | 1015212836 | 4555667892 | 0621375472 | 9212335579 | 0228548210 | 397 |
| 5) 1 | 0515357128 | 1335022514 | 8344109925 | 0559046254 | 0127571528 | 232 |
| 7) 1072886 | 1471414653 | 8633643925 | 1020361090 | 3228255032 | 295 | |
| 9) 109 | 4670081768 | 6617552662 | 3737477845 | 2290911973 | 781 | |
| 11) 111689631 | 8506950061 | 4898861083 | 3430495960 | 817 | | |
| 13) 11395 | 7383788077 | 7534424942 | 4633550708 | 689 | | |
| 15) 1 | 1627118027 | 5561425816 | 2373699984 | 767 | | |
| 17) 1186319 | 5620402145 | 2725474308 | 742 | | | |
| 19) 121 | 0406654464 | 0491044329 | 589 | | | |
| 21) 123498281 | 2431434654 | 048 | | | | |
| 23) 12600 | 5796595391 | 761 | | | | |
| 25) 1 | 2856422466 | 625 | | | | |
| 27) 1311745 | 991 | | | | | |
| 29) 133 | 837 | | | | | |
| 31) 13 | | | | | | |
| 3435 | 3671737612 | 1518555964 | 0207125157 | 6404111859 | 6776182736 | 799 |
| 153269 | 4495916379 | 1233377703 | 5860051584 | 3318322147 | 470 | |
| 10153602 | 8955177278 | 3172623734 | 8493681450 | 983 | | |
| 775141201 | 8370761721 | 0824913332 | 317 | | | |
| 6 | 3705613392 | 8446897069 | 978 | | | |
| 547 | 8512895451 | 815 | | | | |
| 48583 | 184 | | | | | |
| 3435 | 3671890881 | 6024625946 | 1170821347 | 7513162840 | 6372940772 | 546 |
| 0,0101010101 | 0101010101 | 0101010101 | 0101010101 | 0101010101 | 0101010101 | 010 |
| 2103071425 | 6267004502 | 9668821985 | 0111809250 | 8025514305 | 646 | |
| 12 | 1630099085 | 4068616962 | 4859719760 | 5810101330 | 420 | |
| 876 | 5952599082 | 9041109610 | 9587196208 | 360 | | |
| 69783 | 5036494243 | 8395616135 | 808 | | | |
| 5880870 | 5353877840 | 668 | | | | |
| 514256898 | 665 | | | | | |
| 4 | 615 | | | | | |
| 0,0101010101 | 2204081538 | 7998024565 | 9791117914 | 9156023837 | 7787572825 | 192 |
| 3435 | 3671890881 | 6024625946 | 1170821347 | 7513162840 | 6372940772 | 546 |
| 0,0101006665 | 8532190657 | 1973398619 | 8620296567 | 1642860997 | 1414632052 | 646 |

IV. $\operatorname{Acotg} .307 = -3(2) + 3(5) + (13) + (29).$

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| | 0,0032573289 | 9022801302 | 9315960912 | 0521172638 | 4364820846 | 9055374592 | 833 |
| | 3) 345 | 6088648397 | 3443000295 | 0769987174 | 1094169326 | 1556823471 | 597 |
| | 5) 36669764 | 6489336152 | 1087234559 | 3817877050 | 8899421286 | 416 | |
| | 7) 389 | 0732490417 | 2580304164 | 8671007839 | 6274855960 | 501 | |
| | 9) 41281419 | 3298311709 | 4522474201 | 4009658191 | 131 | | |
| | 11) 438 | 0037913381 | 6985798496 | 2620441698 | 672 | | |
| | 13) 46473043 | 8878046300 | 7405517516 | 808 | | | |
| | 15) 493 | 0879254719 | 3742187243 | 551 | | | |
| | 17) 52317576 | 3638805103 | 367 | | | | |
| | 19) 555 | 9995380734 | 067 | | | | |
| | 21) 58897127 | 616 | | | | | |
| | 23) 624 | 909 | | | | | |
| 0,0032 | 115 | 2029549465 | 7814333365 | 0256662391 | 3698056442 | 0518941157 | 199 |
| | 55 | 5818927202 | 4654329166 | 4095858262 | 8039265137 | 214 | |
| | 39 | 8185264852 | 8816890772 | 3874585608 | 970 | | |
| | 32 | 8725283647 | 9582812482 | 903 | | | |
| | 29 | 2157651617 | 582 | | | | |
| | 27 | 169 | | | | | |
| | 115 | 2029549521 | 3633260607 | 3096256443 | 5336089154 | 4173256031 | 037 |
| | 0,0032573289 | 9022801302 | 9315960912 | 0521172638 | 4364820846 | 9055374592 | 833 |
| | 7333952 | 9297867230 | 4217446911 | 8763575410 | 1779884257 | 283 | |
| | 4586824 | 3699812412 | 1613608244 | 6001073132 | 347 | | |
| | 3574849 | 5298311253 | 9031193655 | 139 | | | |
| | 3077504 | 4919929711 | 962 | | | | |
| | 2804625 | 124 | | | | | |
| | 0,0032573289 | 9030135255 | 8618414966 | 8442006812 | 0043393260 | 0790259974 | 688 |
| | 115 | 2029549521 | 3633260607 | 3096256443 | 5336089154 | 4173256031 | 037 |
| | 0,0032573174 | 7000585734 | 4985154359 | 5345750368 | 4707304105 | 6617003943 | 651 |

TABLE OF ARCS CORRESPONDING TO GIVEN PRIME TANGENTS
IN UNITS OF THE RADIUS, TO 110 DIGITS.

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| 2 | 0,7853981633 | 9744830961 | 5660845819 | 8757210492 | 9234984377 | 6455243736 | 1480769541 | |
| 5 | 0,4636476090 | 0080611621 | 4256231461 | 2144020285 | 3705428612 | 0263810933 | 0887201978 | 6 |
| 13 | 0,5880026035 | 4756755124 | 5611080625 | 0854276017 | 0724605592 | 4353726047 | 2072 | |
| 17 | 0,2449786631 | 2686415417 | 2082481211 | | | | | |
| 29 | 0,3805063771 | 1236486630 | 3587916810 | 4331044974 | 0571365810 | 0837576305 | 623 | |
| 37 | 0,1651486774 | 1462683827 | 9128289643 | 9435540983 | 8 | | | |

# ON CALCULATING COMMON LOGARITHMS.

Calculate the logarithms of

$$\log \frac{1025}{*} = a$$
$$\log \frac{1024^2}{*} = b$$
$$\log \frac{81^2}{*} = c$$
$$\log \frac{125^2}{*} = d$$
$$\log \frac{99^2}{*} = e$$

($*$ denotes a denominator that is 1 less than the numerator) then, as is easily checked:

$$\log 2 = \frac{14\frac{3}{4} + \frac{5}{4}a + 2b - \frac{3}{4}c - 2d + e}{49} = f$$

It is also easy to derive

$$\log 41 = a + 12f - 2 = g$$
$$\log 3 = \frac{1 + c + 4f + g}{8}$$

$\log 11 \,.\, 31$ and $\log \frac{11}{7}$ and thus also $\log 7 \,.\, 31$

---

| | | | |
|---|---|---|---|
| $\frac{17}{23 \,.\, 73}$ | $\frac{7}{23 \,.\, 73}$ | $\frac{1680^2}{*}$ | from this 73 |
| $\frac{7 \,.\, 73}{17}$ | $\frac{136000}{*}$ | from these $\frac{7}{23}$ | $\left[\text{and } \frac{7}{23},\ \frac{7}{17}\right]$ |
| | $\frac{511^2}{*}$ | | |
| $7 \,.\, 7 \,.\, 43$ | $\frac{512001}{*}$ | $\frac{7 \,.\, 7}{17}$ from this and $\frac{7}{17}$ | $\left[\text{and } \frac{7}{23}\right]$ |
| $17 \,.\, 43$ | $\frac{730^2}{*}$ | | |

gives 7, 17, 23, 43

---

| | | | |
|---|---|---|---|
| $a$, | $\frac{81}{8}$ | $\frac{81}{80}$ | |
| $b$, | $\frac{41}{4}$ | $\frac{6561}{6560}$ | |
| $c$, | 2 | $\frac{1025}{1024}$ | |
| $d$, | 7 | $\frac{2401}{2400}$ | |
| $e$, | 29 | $\frac{1681^2}{1680.1682}$ | |
| $f$, | 43 | $\frac{512001}{512000}$ | |
| $g$, | 23 . 73 | $\frac{1680^2}{1679.1681}$ | |
| $h$, | 17 | $\frac{136000}{135999}$ | |
| $i$, | 73 | $\frac{511^2}{510.512}$ | $\frac{730^2}{729.731}$ |
| $k$, | 13 | $\frac{729^2}{728.730}$ | |
| $l$, | 19 | $\frac{512^2}{511.513}$ | |
| $m$, | 47 | $\frac{2116^2}{2115.2117}$ | |
| $n$, | 61 | $\frac{2500^2}{2499.2501}$ | |
| $o$, | 31 | $\frac{17081^2}{17080.17082}$ | |
| $p$, | 11 | $\frac{1024^2}{1023.1025}$ | $\frac{2001^2}{2000.2002}$ |
| $q$, | 71 | $\frac{10935^2}{10934.10936}$ | |
| $r$, | 53 | $\frac{788800}{788799}$ | |
| $s$, | 67 | $\frac{274700}{274699}$ | $\frac{3751^2}{*}$ |
| $t$, | 37 | $\frac{1000000}{999999}$ | $\frac{1331^2}{1330.1332}$ |

| | | | |
|---|---|---|---|
| $u$, | 59 | $\frac{3481^2}{3480.3481}$ | |
| $v$, | 89 | $\frac{4095^2}{4094.4096}$ | |
| $w$, | 83 | $\frac{6889^2}{6888.6890}$ | |
| $x$, | 79 | $\frac{3879^2}{3880.3882}$ | |
| $y$, | 97 | $\frac{13871^2}{13870.13872}$ | $\frac{46656^2}{46655.46657}$ |

[How these fractions can be used to calculate the logarithms of the small primes listed beside them, using the very rapidly convergent series in still increasing powers of $\frac{1}{x}$ for $\log \frac{x}{x-1}$, follows directly from the example at the beginning.

For calculating the logarithms of the smallest primes 2, 3, 7 one can, moreover, obtain still more advantageous series by suitably combining these numbers of Gauss with those given by Huyghens (Hugenii, Opera varia, Lugduni 1724 p. 457) as follows:

| | |
|---|---|
| 2 | $1000$ |
| | $1024 = 2^{10}$ |
| 3 | $32805 = 3^8 \,.\, 5$ |
| | $32768 = 2^{15}$ |
| 7 | $2400 = 100 \,.\, 2^3 \,.\, 3$ |
| | $2401 = 7^4$ |
| 11 | $9800 = 100 \,.\, 2 \,.\, 7^2$ |
| | $9801 = 3^4 \,.\, 11^2$ |
| 13 | $123200 = 100 \,.\, 2^4 \,.\, 7 \,.\, 11$ |
| | $123201 = 3^6 \,.\, 13$ |
| 17 | $2600 = 100 \,.\, 2 \,.\, 13$ |
| | $2601 = 3^2 \,.\, 17^2$ |
| 19 | $28899 = 3^2 \,.\, 13^2 \,.\, 19$ |
| | $28900 = 100 \,.\, 17^2$ |
| 23 | $25920 = 10 \,.\, 2^5 \,.\, 3^4$ |
| | $25921 = 7^2 \,.\, 23^2$ |
| 29 | $613088 = 2^5 \,.\, 7^2 \,.\, 17 \,.\, 23$ |
| | $613089 = 3^6 \,.\, 29^2$ |

31 $116280 = 10 \,.\, 2^2 \,.\, 3^2 \,.\, 17 \,.\, 19.$
$116281 = 11^2 \,.\, 31$
37 $165648 = 2^4 \,.\, 3 \,.\, 7 \,.\, 17 \,.\, 29$
$165649 = 11^2 \,.\, 37$
41 $1413720 = 10 \,.\, 2^2 \,.\, 3^3 \,.\, 7 \,.\, 11 \,.\, 17$
$1413721 = 29^2 \,.\, 41^2$
43 $978120 = 10 \,.\, 2^2 \,.\, 3^2 \,.\, 11 \,.\, 13 \,.\, 19$
$978121 = 23^2 \,.\, 43^2$
47 $664848 = 2^5 \,.\, 3^6 \,.\, 7 \,.\, 13$
$664849 = 31^2 \,.\, 47^2$
53 $3059000 = 100 \,.\, 7 \,.\, 19 \,.\, 23$
$3059001 = 3^2 \,.\, 11^2 \,.\, 53^2$
59 $5851560 = 10 \,.\, 2^2 \,.\, 3 \,.\, 11^2 \,.\, 13 \,.\, 31$
$5851561 = 41^2 \,.\, 59^2$
61 $3575880 = 10 \,.\, 2^2 \,.\, 3^3 \,.\, 7 \,.\, 11 \,.\, 43$
$3575881 = 31^2 \,.\, 61^2$
67 $1620528 = 2^4 \,.\, 3 \,.\, 7^2 \,.\, 13$
$1620529 = 19^2 \,.\, 67^2$
71 $2016399 = 3 \,.\, 7^2 \,.\, 11 \,.\, 29 \,.\, 43$
$2016400 = 100 \,.\, 2^2 \,.\, 71^2$
73 $5116644 = 2^2 \,.\, 3^2 \,.\, 13^2 \,.\, 29^2$
$5116645 = 7 \,.\, 17 \,.\, 19 \,.\, 31 \,.\, 73$
79 $5997600 = 100 \,.\, 2^3 \,.\, 3^2 \,.\, 7^2 \,.\, 17$
$5997601 = 31^2 \,.\, 79^2$
83 $1164240 = 10 \,.\, 2^3 \,.\, 3^3 \,.\, 7^2 \,.\, 11$
$1164241 = 13^2 \,.\, 83^2$
89 $2859480 = 10 \,.\, 2^2 \,.\, 3^2 \,.\, 13^2 \,.\, 47$
$2859481 = 19^2 \,.\, 89^2.$
97 $1138488 = 2^3 \,.\, 3 \,.\, 13 \,.\, 41 \,.\, 89$
$1138489 = 11^2 \,.\, 97^2$

To calculate the logarithms of all primes less than 200 one can use to advantage the decompositions of $aa + 1$, $a + 2$, $.. aa + 81$ found in the tables for circle division, provided one restricts attention to the numbers $a$ that themselves have only primes less than 200 as divisors. The other $a$ can then be used to calculate the logarithms of the larger prime divisors occurring in them.]

---

# POSTHUMOUS PAPERS. FIRST TEN THOUSAND SQUARES.

| | 10 | 20 | 30 | 40 | 50 | 60 | 70 | 80 | 90 | |
|---|---|---|---|---|---|---|---|---|---|---|
| 00 | 1000 | 4000 | 9000 | 16000 | 25000 | 36000 | 49000 | 64000 | 81000 | 000 |
| 1 | 02 | 04 | 06 | 08 | 10 | 12 | 14 | 16 | 18 | 001 |
| 2 | 04 | 08 | 12 | 16 | 20 | 24 | 28 | 32 | 36 | 004 |
| 3 | 06 | 12 | 18 | 24 | 30 | 36 | 42 | 48 | 54 | 009 |
| 4 | 08 | 16 | 24 | 32 | 40 | 48 | 56 | 64 | 72 | 016 |
| 5 | 10 | 20 | 30 | 40 | 50 | 60 | 70 | 80 | 81090 | 025 |
| 6 | 12 | 24 | 36 | 48 | 60 | 72 | 84 | 64096 | 81108 | 036 |
| 7 | 14 | 28 | 42 | 56 | 70 | 84 | 49098 | 64112 | 26 | 049 |
| 8 | 16 | 32 | 48 | 64 | 80 | 36096 | 49112 | 28 | 44 | 064 |
| 09 | 18 | 36 | 54 | 72 | 25090 | 36108 | 26 | 44 | 62 | 081 |
| 10 | 20 | 40 | 60 | 80 | 25100 | 20 | 40 | 60 | 80 | 100 |
| 1 | 22 | 44 | 66 | 88 | 10 | 32 | 54 | 76 | 81198 | 121 |
| 2 | 24 | 48 | 72 | 16096 | 20 | 44 | 68 | 64192 | 81216 | 144 |
| 3 | 26 | 52 | 78 | 16104 | 30 | 56 | 82 | 64208 | 34 | 169 |
| 4 | 28 | 56 | 84 | 12 | 40 | 68 | 49196 | 24 | 52 | 196 |
| 5 | 30 | 60 | 90 | 20 | 50 | 80 | 49210 | 40 | 70 | 225 |
| 6 | 32 | 64 | 9096 | 28 | 60 | 36192 | 24 | 56 | 81288 | 256 |
| 7 | 34 | 68 | 9102 | 36 | 70 | 36204 | 38 | 72 | 81306 | 289 |
| 8 | 36 | 72 | 08 | 44 | 80 | 16 | 52 | 64288 | 24 | 324 |
| 19 | 38 | 76 | 14 | 52 | 25190 | 28 | 66 | 64304 | 42 | 361 |
| 20 | 40 | 80 | 20 | 60 | 25200 | 40 | 80 | 20 | 60 | 400 |
| 1 | 42 | 84 | 26 | 68 | 10 | 52 | 49294 | 36 | 78 | 441 |
| 2 | 44 | 88 | 32 | 76 | 20 | 64 | 49308 | 52 | 81396 | 484 |
| 3 | 46 | 92 | 38 | 84 | 30 | 76 | 22 | 68 | 81414 | 529 |
| 4 | 1048 | 4096 | 44 | 16192 | 40 | 36288 | 36 | 64384 | 32 | 576 |
| 5 | 50 | 4100 | 50 | 16200 | 50 | 36300 | 50 | 64400 | 50 | 625 |
| 6 | 52 | 04 | 56 | 08 | 60 | 12 | 64 | 16 | 68 | 676 |
| 7 | 54 | 08 | 62 | 16 | 70 | 24 | 78 | 32 | 81486 | 729 |
| 8 | 56 | 12 | 68 | 24 | 80 | 36 | 49392 | 48 | 81504 | 784 |
| 29 | 58 | 16 | 74 | 32 | 25290 | 48 | 49406 | 64 | 22 | 841 |
| 30 | 60 | 20 | 80 | 40 | 25300 | 60 | 20 | 80 | 40 | 900 |
| 1 | 62 | 24 | 86 | 48 | 10 | 72 | 34 | 64496 | 58 | 961 |
| 2 | 65 | 29 | 93 | 57 | 21 | 85 | 49 | 64513 | 77 | 024 |
| 3 | 67 | 33 | 9199 | 65 | 31 | 36397 | 63 | 29 | 81595 | 089 |
| 4 | 69 | 37 | 9205 | 73 | 41 | 36409 | 77 | 45 | 81613 | 156 |
| 5 | 71 | 41 | 11 | 81 | 51 | 21 | 49491 | 61 | 31 | 225 |
| 6 | 73 | 45 | 17 | 89 | 61 | 33 | 49505 | 77 | 49 | 296 |
| 7 | 75 | 49 | 23 | 16297 | 71 | 45 | 19 | 64593 | 67 | 369 |
| 8 | 77 | 53 | 29 | 16305 | 81 | 57 | 33 | 64609 | 81685 | 444 |
| 39 | 79 | 57 | 35 | 13 | 25391 | 69 | 47 | 25 | 81703 | 521 |
| 40 | 81 | 61 | 41 | 21 | 25401 | 81 | 61 | 41 | 21 | 600 |
| 1 | 83 | 65 | 47 | 29 | 11 | 36493 | 75 | 57 | 39 | 681 |
| 2 | 85 | 69 | 53 | 37 | 21 | 36505 | 49589 | 73 | 57 | 764 |
| 3 | 87 | 73 | 59 | 45 | 31 | 17 | 49603 | 64689 | 75 | 849 |
| 4 | 89 | 77 | 65 | 53 | 41 | 29 | 17 | 64705 | 81793 | 936 |
| 5 | 92 | 82 | 72 | 62 | 52 | 42 | 32 | 22 | 81812 | 025 |
| 6 | 94 | 86 | 78 | 70 | 62 | 54 | 46 | 38 | 30 | 116 |
| 7 | 96 | 90 | 84 | 78 | 72 | 66 | 60 | 54 | 48 | 209 |
| 8 | 1098 | 94 | 90 | 86 | 82 | 78 | 74 | 70 | 66 | 304 |
| 94 | 1100 | 4198 | 9296 | 16394 | 25492 | 36590 | 49688 | 64786 | 81884 | 401 |

| 11 | 21 | 31 | 41 | 51 | 61 | 71 | 81 | 91 | |
|---|---|---|---|---|---|---|---|---|---|
| 1210 | 4410 | 9610 | 16810 | 26010 | 37210 | 50410 | 65610 | 82810 | 000 |
| 12 | 14 | 16 | 18 | 20 | 22 | 24 | 26 | 28 | 201 |
| 14 | 18 | 22 | 26 | 30 | 34 | 38 | 42 | 46 | 404 |
| 16 | 22 | 28 | 34 | 40 | 46 | 52 | 58 | 64 | 609 |
| 18 | 26 | 34 | 42 | 50 | 58 | 66 | 74 | 82882 | 816 |
| 21 | 31 | 41 | 51 | 61 | 71 | 81 | 65691 | 82901 | 025 |
| 23 | 35 | 47 | 59 | 71 | 83 | 50495 | 65707 | 19 | 236 |
| 25 | 39 | 53 | 67 | 81 | 37295 | 50509 | 23 | 37 | 449 |
| 27 | 43 | 59 | 75 | 26091 | 37307 | 23 | 39 | 55 | 664 |
| 29 | 47 | 65 | 83 | 26101 | 19 | 37 | 55 | 73 | 881 |
| 32 | 52 | 72 | 16892 | 12 | 32 | 52 | 72 | 82992 | 100 |
| 34 | 56 | 78 | 16900 | 22 | 44 | 66 | 65788 | 83010 | 321 |
| 36 | 60 | 84 | 08 | 32 | 56 | 80 | 65804 | 28 | 544 |
| 38 | 64 | 90 | 16 | 42 | 68 | 50594 | 20 | 46 | 769 |
| 40 | 68 | 9696 | 24 | 52 | 80 | 50608 | 36 | 64 | 996 |
| 43 | 73 | 9703 | 33 | 63 | 37393 | 23 | 53 | 83083 | 225 |
| 45 | 77 | 09 | 41 | 73 | 37405 | 37 | 69 | 83101 | 456 |
| 47 | 81 | 15 | 49 | 83 | 17 | 51 | 65885 | 19 | 689 |
| 49 | 85 | 21 | 57 | 26193 | 29 | 65 | 65901 | 37 | 924 |
| 52 | 90 | 28 | 66 | 26204 | 42 | 80 | 18 | 56 | 161 |
| 1254 | 94 | 34 | 74 | 14 | 54 | 50694 | 34 | 74 | 400 |
| 56 | 4498 | 40 | 82 | 24 | 66 | 50708 | 50 | 83192 | 641 |
| 58 | 4502 | 46 | 90 | 34 | 78 | 22 | 66 | 83210 | 884 |
| 61 | 07 | 53 | 16999 | 45 | 37491 | 37 | 83 | 29 | 129 |
| 63 | 11 | 59 | 17007 | 55 | 37503 | 51 | 65999 | 47 | 376 |
| 65 | 15 | 65 | 15 | 65 | 15 | 65 | 66015 | 65 | 625 |
| 67 | 19 | 71 | 23 | 75 | 27 | 79 | 31 | 83283 | 876 |
| 70 | 24 | 78 | 32 | 86 | 40 | 50794 | 48 | 83302 | 129 |
| 72 | 28 | 84 | 40 | 26296 | 52 | 50808 | 64 | 20 | 384 |
| 74 | 32 | 90 | 48 | 26306 | 64 | 22 | 80 | 38 | 641 |
| 76 | 36 | 9796 | 56 | 16 | 76 | 36 | 66096 | 56 | 900 |
| 79 | 41 | 9803 | 65 | 27 | 37589 | 51 | 66113 | 75 | 161 |
| 81 | 45 | 09 | 73 | 37 | 37601 | 65 | 29 | 83393 | 424 |
| 83 | 49 | 15 | 81 | 47 | 13 | 79 | 45 | 83411 | 689 |
| 85 | 53 | 21 | 89 | 57 | 25 | 50893 | 61 | 29 | 956 |
| 88 | 58 | 28 | 17098 | 68 | 38 | 50908 | 78 | 48 | 225 |
| 90 | 62 | 34 | 17106 | 78 | 50 | 22 | 66194 | 66 | 496 |
| 93 | 66 | 40 | 14 | 88 | 62 | 36 | 66210 | 83484 | 769 |
| 95 | 71 | 47 | 23 | 26399 | 75 | 51 | 27 | 83503 | 044 |
| 1297 | 75 | 53 | 31 | 26409 | 87 | 65 | 43 | 21 | 321 |
| 1299 | 79 | 59 | 39 | 19 | 37699 | 79 | 59 | 39 | 600 |
| 1301 | 83 | 65 | 47 | 29 | 37711 | 50993 | 75 | 57 | 881 |
| 04 | 88 | 72 | 56 | 40 | 24 | 51008 | 66292 | 76 | 164 |
| 06 | 92 | 78 | 64 | 50 | 36 | 22 | 66308 | 83594 | 449 |
| 08 | 4596 | 84 | 72 | 60 | 48 | 36 | 24 | 83612 | 736 |
| 11 | 4601 | 91 | 81 | 71 | 61 | 51 | 41 | 31 | 025 |
| 13 | 05 | 9897 | 89 | 81 | 73 | 65 | 57 | 49 | 316 |
| 15 | 09 | 9903 | 17197 | 26491 | 85 | 79 | 73 | 67 | 609 |
| 17 | 13 | 09 | 17205 | 26501 | 37797 | 51093 | 66389 | 83685 | 904 |
| 1320 | 4618 | 9916 | 17214 | 26512 | 37810 | 51108 | 66406 | 83704 | 201 |

| | 10 | 20 | 30 | 40 | 50 | 60 | 70 | 80 | 90 | |
|---|---|---|---|---|---|---|---|---|---|---|
| 50 | 1102 | 4202 | 9302 | 16402 | 25502 | 36602 | 49702 | 64802 | 81902 | 500 |
| 1 | 04 | 06 | 08 | 10 | 12 | 14 | 16 | 18 | 20 | 601 |
| 2 | 06 | 10 | 14 | 18 | 22 | 26 | 30 | 34 | 38 | 704 |
| 3 | 08 | 14 | 20 | 26 | 32 | 38 | 44 | 50 | 56 | 809 |
| 4 | 10 | 18 | 26 | 34 | 42 | 50 | 58 | 66 | 74 | 916 |
| 5 | 13 | 23 | 33 | 43 | 53 | 63 | 73 | 83 | 81993 | 025 |
| 6 | 15 | 27 | 39 | 51 | 63 | 75 | 49787 | 64899 | 82011 | 136 |
| 7 | 17 | 31 | 45 | 59 | 73 | 87 | 49801 | 64915 | 29 | 249 |
| 8 | 19 | 35 | 51 | 67 | 83 | 36699 | 15 | 31 | 47 | 364 |
| 59 | 21 | 39 | 57 | 75 | 25593 | 36711 | 29 | 47 | 65 | 481 |
| 60 | 23 | 43 | 63 | 83 | 25603 | 23 | 43 | 63 | 82083 | 600 |
| 1 | 25 | 47 | 69 | 91 | 13 | 35 | 57 | 79 | 82101 | 721 |
| 2 | 27 | 51 | 75 | 16499 | 23 | 47 | 71 | 64995 | 19 | 844 |
| 3 | 29 | 55 | 81 | 16507 | 33 | 59 | 49885 | 65011 | 37 | 969 |
| 4 | 32 | 60 | 88 | 16 | 44 | 72 | 49900 | 28 | 56 | 096 |
| 5 | 34 | 64 | 9394 | 24 | 54 | 84 | 14 | 44 | 74 | 225 |
| 6 | 36 | 68 | 9400 | 32 | 64 | 36796 | 28 | 60 | 82192 | 356 |
| 7 | 38 | 72 | 06 | 40 | 74 | 36808 | 42 | 76 | 82210 | 489 |
| 8 | 40 | 76 | 12 | 48 | 84 | 20 | 56 | 65092 | 28 | 624 |
| 69 | 42 | 80 | 18 | 56 | 25694 | 32 | 70 | 65108 | 46 | 761 |
| 70 | 1144 | 84 | 24 | 64 | 25704 | 44 | 84 | 24 | 64 | 900 |
| 1 | 47 | 89 | 31 | 73 | 15 | 57 | 49999 | 41 | 82283 | 041 |
| 2 | 49 | 93 | 37 | 81 | 25 | 69 | 50013 | 57 | 82301 | 184 |
| 3 | 51 | 4297 | 43 | 89 | 35 | 81 | 27 | 73 | 19 | 329 |
| 4 | 53 | 4301 | 49 | 16597 | 45 | 36893 | 41 | 65189 | 37 | 476 |
| 5 | 55 | 05 | 55 | 16605 | 55 | 36905 | 55 | 65205 | 55 | 625 |
| 6 | 57 | 09 | 61 | 13 | 65 | 17 | 69 | 21 | 73 | 776 |
| 7 | 59 | 13 | 67 | 21 | 75 | 29 | 83 | 37 | 82391 | 929 |
| 8 | 62 | 18 | 74 | 30 | 86 | 42 | 50098 | 53 | 82409 | 084 |
| 79 | 64 | 22 | 80 | 38 | 25796 | 54 | 50112 | 70 | 28 | 241 |
| 80 | 66 | 26 | 86 | 46 | 25806 | 66 | 26 | 65286 | 46 | 400 |
| 1 | 68 | 30 | 92 | 54 | 16 | 78 | 40 | 65302 | 64 | 561 |
| 2 | 70 | 34 | 9498 | 62 | 26 | 36990 | 54 | 18 | 82482 | 724 |
| 3 | 72 | 38 | 9504 | 70 | 36 | 37002 | 68 | 34 | 82500 | 889 |
| 4 | 75 | 43 | 11 | 79 | 47 | 15 | 83 | 51 | 19 | 056 |
| 5 | 77 | 47 | 17 | 87 | 57 | 27 | 50197 | 67 | 37 | 225 |
| 6 | 79 | 51 | 23 | 16695 | 67 | 39 | 50211 | 83 | 55 | 396 |
| 7 | 81 | 55 | 29 | 16703 | 77 | 51 | 25 | 65399 | 73 | 569 |
| 8 | 83 | 59 | 35 | 11 | 87 | 63 | 39 | 65415 | 82591 | 744 |
| 89 | 85 | 63 | 41 | 19 | 25897 | 75 | 53 | 31 | 82609 | 921 |
| 90 | 88 | 68 | 48 | 28 | 25908 | 37088 | 68 | 48 | 28 | 100 |
| 1 | 90 | 72 | 54 | 36 | 18 | 37100 | 82 | 64 | 46 | 281 |
| 2 | 92 | 76 | 60 | 44 | 28 | 12 | 50296 | 80 | 64 | 464 |
| 3 | 94 | 80 | 66 | 52 | 38 | 24 | 50310 | 65496 | 82682 | 649 |
| 4 | 96 | 84 | 72 | 60 | 48 | 36 | 24 | 65512 | 82700 | 836 |
| 5 | 1199 | 89 | 79 | 69 | 59 | 49 | 39 | 29 | 19 | 025 |
| 6 | 1201 | 93 | 85 | 77 | 69 | 61 | 53 | 45 | 37 | 216 |
| 7 | 03 | 4397 | 91 | 85 | 79 | 73 | 67 | 61 | 55 | 409 |
| 8 | 05 | 4401 | 9597 | 16793 | 89 | 85 | 81 | 77 | 73 | 604 |
| 9 | 1207 | 4405 | 9603 | 16801 | 25999 | 37197 | 50395 | 65693 | 82791 | 801 |

| 11 | 21 | 31 | 41 | 51 | 61 | 71 | 81 | 91 | |
|---|---|---|---|---|---|---|---|---|---|
| 1322 | 4622 | 9922 | 17222 | 26522 | 37822 | 51122 | 66422 | 83722 | 500 |
| 24 | 26 | 28 | 30 | 32 | 34 | 36 | 38 | 40 | 801 |
| 27 | 31 | 35 | 39 | 43 | 47 | 51 | 55 | 59 | 104 |
| 29 | 35 | 41 | 47 | 53 | 59 | 65 | 71 | 77 | 409 |
| 31 | 39 | 47 | 55 | 63 | 71 | 79 | 66487 | 83795 | 716 |
| 34 | 44 | 54 | 64 | 74 | 84 | 51194 | 66504 | 83814 | 025 |
| 36 | 48 | 60 | 72 | 84 | 37896 | 51208 | 20 | 32 | 336 |
| 38 | 52 | 66 | 80 | 26594 | 37908 | 22 | 36 | 50 | 649 |
| 40 | 56 | 72 | 88 | 26604 | 20 | 36 | 52 | 68 | 964 |
| 43 | 61 | 79 | 17297 | 15 | 33 | 51 | 69 | 83887 | 281 |
| 45 | 65 | 85 | 17305 | 25 | 45 | 65 | 66585 | 83905 | 600 |
| 47 | 69 | 91 | 13 | 35 | 57 | 79 | 66601 | 23 | 921 |
| 50 | 74 | 9998 | 22 | 46 | 70 | 51294 | 18 | 42 | 244 |
| 52 | 78 | 10004 | 30 | 56 | 82 | 51308 | 34 | 60 | 569 |
| 54 | 82 | 10 | 38 | 66 | 37994 | 22 | 50 | 78 | 896 |
| 57 | 87 | 17 | 47 | 77 | 38007 | 37 | 67 | 83997 | 225 |
| 1359 | 91 | 23 | 55 | 87 | 19 | 51 | 83 | 84015 | 556 |
| 1361 | 4695 | 29 | 63 | 26697 | 31 | 65 | 66699 | | 889 |
| 64 | 4700 | 36 | 72 | 26708 | 44 | 80 | 66716 | | 224 |
| 66 | 04 | 42 | 80 | 18 | 56 | 51394 | 32 | | 561 |
| 68 | 08 | 48 | 88 | 28 | 68 | 51408 | | | 900 |
| 71 | 13 | 55 | 17397 | 39 | 81 | 23 | | | 241 |
| 73 | 17 | 61 | 17405 | 49 | 38093 | 37 | | | 584 |
| 75 | 21 | 67 | 13 | 59 | 38105 | 51 | | | 929 |
| 78 | 26 | 74 | 22 | 70 | 18 | 66 | | | 276 |
| 80 | 30 | 80 | 30 | 80 | 30 | 80 | | | 625 |
| 82 | 34 | 86 | 38 | 26790 | 42 | 51494 | | | 976 |
| 85 | 39 | 93 | 47 | 26801 | 55 | 51509 | | | 329 |
| 87 | 43 | 10099 | 55 | 11 | 67 | 23 | | | 684 |
| 90 | 48 | 10106 | 64 | 22 | 80 | 38 | | | 041 |
| 92 | 52 | 12 | 72 | 32 | 38192 | 52 | | | 400 |
| 94 | 56 | 18 | 80 | 42 | 38204 | 66 | | | 761 |
| 97 | 61 | 25 | 89 | 53 | 17 | 81 | | | 124 |
| 1399 | 65 | 31 | 17497 | 63 | 29 | 51595 | | | 489 |
| 1401 | 69 | 37 | 17505 | 73 | 41 | 51609 | | | 856 |
| 04 | 74 | 44 | 14 | 84 | 54 | 24 | | | 225 |
| 06 | 78 | 50 | 22 | 26894 | 66 | 38 | | | 596 |
| 08 | 82 | 56 | 30 | 26904 | 78 | 52 | | | 969 |
| 11 | 87 | 63 | 39 | 15 | 38291 | 67 | | | 344 |
| 13 | 91 | 69 | 47 | 25 | 38303 | 81 | | | 721 |
| 16 | 4796 | 76 | 56 | 36 | 16 | 51696 | | | 100 |
| 18 | 4800 | 82 | 64 | 46 | 28 | 51710 | | | 481 |
| 20 | 04 | 88 | 72 | 56 | 40 | 24 | | | 864 |
| 23 | 09 | 10195 | 81 | 67 | 53 | 39 | | | 249 |
| 25 | 13 | 10201 | 89 | 77 | 65 | 53 | | | 636 |
| 28 | 18 | 08 | 17598 | 88 | 78 | 68 | | | 025 |
| 30 | 22 | 14 | 17606 | 26998 | 38390 | 82 | | | 416 |
| 32 | 26 | 20 | 14 | 27008 | 38402 | 51796 | | | 809 |
| 35 | 31 | 27 | 23 | 19 | 15 | 51811 | | | 204 |
| 1437 | 4835 | 10233 | 17631 | 27029 | 38427 | 51825 | | | 601 |

# POSTHUMOUS PAPERS INDICES OF PRIMES IN THE HIGHER NUMBER DOMAIN.

| | $-1+2i$ | $-1-2i$ | $-3$ | $3+2i$ | $3-2i$ | $1+4i$ | $1-4i$ | $-5+2i$ | $-5-2i$ | $-1+6i$ | $-1-6i$ | $5+4i$ | $5-4i$ | $-7$ | $7+2i$ | $7-2i$ | $-5+6i$ | $-5-6i$ | $-3+8i$ | $-3-8i$ | $5+8i$ | $5-8i$ | $9+4i$ | $9-4i$ |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | $i$ | $i$ | $1-i$ | $-1-i$ | $+1-i$ | $+1-2i$ | $+1-2i$ | $+1+i$ | $-1+i$ | $+1+i$ | $-1+i$ | $+2-i$ | $+1-2i$ | $+1-2i$ | $-1+i$ | $1+i$ | $1-i$ | $-1-i$ | $1+i$ | $-1-i$ | $1-i$ | $1-i$ | $1+i$ | $1+4i$ |
| $i$ | 1 | 1 | 2 | 3 | 3 | 4 | 4 | 7 | 7 | 9 | 9 | 10 | 10 | 12 | 13 | 13 | 15 | 15 | 18 | 18 | 22 | 22 | 24 | 24 |
| $1+i$ | 2 | 3 | 3 | 7 | 4 | 7 | 5 | 1 | 22 | 1 | 28 | 32 | 2 | 46 | 40 | 1 | 16 | 31 | 1 | 37 | 23 | 23 | 1 | 7 |
| $1-i$ | 1 | 2 | 1 | 4 | 1 | 3 | 1 | 22 | 15 | 28 | 19 | 22 | 32 | 34 | 27 | 40 | 1 | 16 | 55 | 19 | 1 | 1 | 73 | 79 |
| $-1+2i$ | $*$ | 3 | 7 | 4 | 5 | 9 | 2 | 18 | 12 | 13 | 6 | 27 | 21 | 25 | 19 | 10 | 35 | 9 | 68 | 65 | 10 | 40 | 20 | 63 |
| $-1-2i$ | 1 | $*$ | 5 | 11 | 4 | 14 | 15 | 12 | 18 | 6 | 31 | 31 | 17 | 31 | 10 | 45 | 39 | 5 | 11 | 68 | 40 | 54 | 81 | 44 |
| $-3$ | 3 | 1 | $*$ | 2 | 2 | 3 | 5 | 17 | 3 | 16 | 16 | 25 | 35 | 16 | 21 | 47 | 12 | 12 | 6 | 42 | 51 | 29 | 14 | 50 |
| $3+2i$ | 2 | 1 | 6 | $*$ | 1 | 13 | 9 | 13 | 23 | 14 | 35 | 26 | 33 | 3 | 4 | 44 | 28 | 52 | 9 | 44 | 79 | 38 | 36 | 23 |
| $3-2i$ | 3 | 2 | 2 | 7 | $*$ | 15 | 11 | 9 | 27 | 17 | 14 | 3 | 6 | 21 | 44 | 4 | 52 | 28 | 44 | 27 | 82 | 57 | 89 | 60 |
| $1+4i$ | 1 | 2 | 3 | 9 | 7 | $*$ | 6 | 15 | 20 | 21 | 2 | 8 | 9 | 29 | 22 | 24 | 38 | 11 | 23 | 52 | 27 | 81 | 78 | 1 |
| $1-4i$ | 2 | 3 | 1 | 1 | 3 | 10 | $*$ | 20 | 1 | 2 | 3 | 19 | 8 | 11 | 24 | 22 | 41 | 38 | 52 | 5 | 15 | 5 | 79 | 18 |
| $-5+2i$ | 0 | 2 | 1 | 3 | 11 | 11 | 12 | $*$ | 25 | 5 | 28 | 29 | 24 | 30 | 51 | 41 | 19 | 36 | 71 | 66 | 8 | 75 | 91 | 58 |
| $-5-2i$ | 2 | 0 | 3 | 5 | 9 | 4 | 13 | 11 | $*$ | 28 | 23 | 24 | 39 | 18 | 15 | 25 | 36 | 49 | 30 | 53 | 53 | 8 | 70 | 85 |
| $-1+6i$ | 3 | 0 | 4 | 8 | 3 | 5 | 10 | 19 | 22 | $*$ | 29 | 33 | 5 | 22 | 27 | 33 | 34 | 59 | 39 | 67 | 65 | 12 | 39 | 32 |
| $-1-6i$ | 0 | 1 | 4 | 9 | 8 | 6 | 3 | 22 | 5 | 11 | $*$ | 15 | 3 | 10 | 7 | 1 | 29 | 34 | 49 | 21 | 12 | 87 | 32 | 9 |
| $5+4i$ | 3 | 3 | 5 | 6 | 11 | 4 | 3 | 5 | 16 | 25 | 15 | $*$ | 32 | 9 | 16 | 9 | 46 | 32 | 69 | 13 | 26 | 55 | 4 | 43 |
| $5-4i$ | 1 | 1 | 7 | 5 | 6 | 5 | 12 | 16 | 19 | 33 | 7 | 32 | $*$ | 15 | 35 | 16 | 32 | 46 | 31 | 15 | 33 | 70 | 37 | 28 |
| $-7$ | 1 | 3 | 4 | 7 | 1 | 1 | 7 | 10 | 10 | 10 | 10 | 1 | 11 | $*$ | 28 | 28 | 23 | 53 | 51 | 33 | 83 | 61 | 11 | 5 |
| $7+2i$ | 1 | 0 | 1 | 10 | 12 | 2 | 8 | 21 | 9 | 1 | 25 | 36 | 7 | 44 | $*$ | 43 | 55 | 26 | 37 | 10 | 87 | 85 | 27 | 89 |
| $7-2i$ | 0 | 3 | 3 | 2 | 10 | 8 | 14 | 23 | 7 | 7 | 19 | 37 | 36 | 20 | 17 | $*$ | 26 | 25 | 46 | 55 | 19 | 65 | 23 | 21 |
| $-5+6i$ | 1 | 1 | 0 | 6 | 10 | 10 | 1 | 25 | 6 | 4 | 27 | 2 | 4 | 43 | 37 | 8 | $*$ | 31 | 21 | 43 | 85 | 2 | 45 | 29 |
| $-5-6i$ | 3 | 3 | 0 | 10 | 6 | 7 | 6 | 6 | 11 | 9 | 4 | 4 | 22 | 13 | 8 | 11 | 1 | $*$ | 25 | 39 | 46 | 19 | 83 | 3 |
| $-3+8i$ | 0 | 3 | 6 | 5 | 4 | 15 | 18 | 7 | 26 | 3 | 33 | 9 | 31 | 23 | 1 | 6 | 33 | 13 | $*$ | 26 | 28 | 24 | 73 | 13 |
| $-3-8i$ | 1 | 0 | 2 | 4 | 11 | 8 | 19 | 26 | 21 | 15 | 21 | 21 | 19 | 17 | 6 | 27 | 43 | 3 | 62 | $*$ | 24 | 28 | 67 | 7 |
| $5+8i$ | 2 | 0 | 7 | 7 | 10 | 11 | 15 | 4 | 5 | 9 | 8 | 38 | 13 | 19 | 39 | 7 | 25 | 2 | 24 | 56 | $*$ | 30 | 77 | 79 |
| $5-8i$ | 0 | 2 | 5 | 10 | 1 | 9 | 13 | 19 | 4 | 8 | 27 | 23 | 18 | 37 | 33 | 13 | 2 | 55 | 56 | 24 | 74 | $*$ | 1 | 35 |
| $9+4i$ | 0 | 1 | 2 | 8 | 5 | 14 | 7 | 11 | 14 | 35 | 4 | 28 | 25 | 39 | 3 | 39 | 21 | 7 | 1 | 11 | 49 | 1 | $*$ | 90 |
| $9\quad 4i$ | 3 | 0 | 6 | 11 | 8 | 1 | 2 | 14 | 25 | 4 | 17 | 35 | 28 | 33 | 13 | 29 | 37 | 51 | 65 | 19 | 23 | 71 | 6 | $*$ |
| $-1+10i$ | 2 | 2 | 5 | 2 | 0 | 7 | 7 | 16 | 17 | 31 | 17 | 21 | 10 | 5 | 46 | 35 | 50 | 45 | 39 | 34 | 57 | 42 | 10 | 54 |
| $-1-10i$ | 2 | 2 | 7 | 0 | 2 | 1 | 1 | 3 | 16 | 35 | 13 | 30 | 31 | 35 | 9 | 46 | 15 | 50 | 70 | 21 | 86 | 79 | 42 | 22 |
| $3+10i$ | 1 | 3 | 2 | 0 | 9 | 6 | 5 | 14 | 24 | 23 | 24 | 17 | 18 | 38 | 49 | 50 | 54 | 56 | 14 | 2 | 69 | 51 | 65 | 31 |
| $3-10i$ | 1 | 3 | 6 | 3 | 0 | 3 | 10 | 24 | 14 | 24 | 5 | 38 | 27 | 26 | 50 | 23 | 56 | 54 | 38 | 50 | 29 | 3 | 49 | 47 |
| $-7+8i$ | 3 | 2 | 7 | 1 | 9 | 14 | 9 | 22 | 9 | 7 | 22 | 7 | 15 | 12 | 5 | 23 | 48 | 6 | 4 | 47 | 11 | 68 | 58 | 82 |
| $-7-8i$ | 2 | 1 | 5 | 3 | 7 | 15 | 2 | 23 | 22 | 22 | 25 | 5 | 37 | 36 | 49 | 31 | 6 | 48 | 29 | 4 | 68 | 77 | 46 | 70 |
| $-11$ | 2 | 2 | 0 | 11 | 5 | 5 | 3 | 1 | 15 | 24 | 24 | 37 | 7 | 40 | 12 | 12 | 15 | 45 | 61 | 7 | 16 | 16 | 94 | 34 |
| $-11+4i$ | 0 | 1 | 3 | 4 | 3 | 7 | 15 | 14 | 27 | 29 | 13 | 36 | 29 | 26 | 32 | 21 | 57 | 29 | 54 | 30 | 41 | 18 | 9 | 38 |
| $-11-4i$ | 3 | 0 | 1 | 9 | 4 | 9 | 1 | 13 | 14 | 31 | 11 | 39 | 36 | 38 | 47 | 32 | 59 | 27 | 66 | 18 | 62 | 63 | 26 | 39 |
| $7+10i$ | 3 | 1 | 3 | 3 | 4 | 12 | 0 | 3 | 1 | 26 | 10 | 23 | 20 | 4 | 6 | 12 | 44 | 22 | 29 | 31 | 47 | 14 | 12 | 75 |
| $7-10i$ | 3 | 1 | 1 | 4 | 9 | 0 | 4 | 15 | 17 | 10 | 26 | 20 | 13 | 28 | 12 | 6 | 22 | 44 | 13 | 47 | 58 | 25 | 69 | 84 |
| $11+6i$ | 2 | 3 | 4 | 11 | 7 | 0 | 12 | 17 | 4 | 20 | 12 | 7 | 14 | 17 | 18 | 49 | 8 | 29 | 43 | 18 | 13 | 11 | 75 | 20 |
| $11-6i$ | 1 | 2 | 4 | 1 | 5 | 4 | 0 | 4 | 3 | 12 | 20 | 34 | 37 | 23 | 23 | 18 | 59 | 8 | 54 | 25 | 77 | 35 | 44 | 69 |
| $-23+2i$ | 1 | 0 | 7 | 2 | 8 | 15 | 4 | 27 | 1 | 20 | 18 | 18 | 32 | 7 | 33 | 36 | 24 | 7 | 26 | 28 | 72 | 66 | 38 | 77 |
| $-23-2i$ | 0 | 3 | 5 | 8 | 2 | 12 | 9 | 15 | 13 | 18 | 20 | 32 | 38 | 1 | 36 | 7 | 37 | 24 | 28 | 62 | 22 | 72 | 35 | 26 |
| $-9+20i$ | 0 | 0 | 2 | 11 | 1 | 3 | 6 | 8 | 16 | 19 | 3 | 20 | 37 | 15 | 42 | 37 | 5 | 42 | 53 | 59 | 39 | 56 | 61 | 16 |
| $-9-10i$ | 0 | 0 | 6 | 7 | 5 | 10 | 5 | 16 | 8 | 21 | 1 | 7 | 20 | 9 | 11 | 42 | 42 | 35 | 41 | 71 | 56 | 17 | 16 | 19 |
| $-7+12i$ | 2 | 1 | 4 | 0 | 11 | 9 | 4 | 12 | 13 | 1 | 14 | 11 | 22 | 20 | 30 | 5 | 42 | 19 | 59 | 50 | 81 | 64 | 87 | 87 |

## AUXILIARY TABLE

FOR SOLVING THE INDETERMINATE EQUATION

$$A = fxx + gyy$$

BY THE METHOD OF EXCLUSION.

---

It is assumed that a prime $p$ has been chosen as the exclusion number, and that none of the numbers $A$, $f$, $g$ is divisible by it. The table is also restricted to the two cases in which the value of the expression $\frac{A}{f}(\text{mod}.p)$ is either a fixed quadratic residue (always 1), or a fixed quadratic nonresidue modulo $p$. Finally, the table has been provided only for the case in which $fg$ is a quadratic residue of $p$, and the opposite case has been omitted entirely. Section Six of the *Disquisitiones Arithmeticae* gives sufficient instruction on how to derive easily from the table what it does not contain directly.

*Examples.* Let the equation to be solved be $21680143 = xx + 78yy$

1) Exclusion number $= 5$ $\qquad fg = \text{N}, \quad \frac{A}{f} \equiv 3 \equiv 2 \,.\, 2^2$

From the table $1, 4$ for $fg = R$ and hence $0, 2, 3$ for $fg = \text{N}$, and for the above case $0, 1, 4$ or excluded $5n \pm 2$

2) Exclusion number $= 7$ $\qquad fg = R, \quad \frac{A}{f} \equiv 2 \equiv 1, 3^2$

From the table $0, 1, 2, 5, 6$ For the present case $0, 3, 6, 1, 4$ and excluded $7n \pm 2$

3) Exclusion number $= 11$ $\qquad fg = R, \quad \frac{A}{f} \equiv 1$

Thus we obtain $0, 1, 3, 5, 6, 8, 10$; excluded $11n \pm 2, 4$

4) Exclusion number $= 17$ $\qquad fg = \text{N}, \quad \frac{A}{f} \equiv 9 \equiv 1 \,.\, 3^1$

$$0 \,.\, 1 \,.\, 3 \,.\, 4 \,.\, 6$$
$$0 \,.\, 3 \,.\, 8 \,.\, 5 \,.\, 1$$
$$0, 2, 3, 4, 6, 7$$

Exclusion number $\quad \pm 1, 5, 8$

| $p$ | Value of $\frac{A}{f}(\text{mod}.p)$ | Numbers, taken positively or negatively, to which $x$ must be congruent modulo $p$. |
|---|---|---|
| 3 | 1 | 0, 1 |
| | 2 | 1 |
| 5 | 1 | 0, 1 |
| | 2 | 1 |
| 7 | 1 | 0, 1, 2 |
| | 6 | 2, 3 |
| 11 | 1 | 0, 1, 3, 5 |
| | 10 | 1, 3, 4 |
| 13 | 1 | 0, 1, 2, 6 |
| | 2 | 1, 4, 5 |
| 17 | 1 | 0, 1, 3, 4, 6 |
| | 3 | 1, 2, 4, 6 |
| 19 | 1 | 0, 1, 2, 3, 4, 7 |
| | 18 | 1, 3, 6, 7, 8 |
| 23 | 1 | 0, 1, 4, 8, 9, 10, 11 |
| | 22 | 1, 2, 3, 4, 6, 8 |
| 29 | 1 | 0, 1, 5, 6, 8, 9, 11, 13 |
| | 2 | 1, 3, 5, 6, 8, 13, 14 |
| 31 | 1 | 0, 1, 2, 4, 5, 7, 10, 11, 13 |
| | 30 | 4, 5, 6, 8, 11, 12, 13, 14 |
| 37 | 1 | 0, 1, 2, 7, 8, 10, 11, 14, 16, 18 |
| | 2 | 1, 3, 6, 7, 8, 14, 15, 17, 18 |
| 41 | 1 | 0, 1 . 3, 9, 12, 13, 14, 16, 17, 18, 19 |
| | 3 | 1, 2, 6, 7, 8, 9, 11, 12, 13, 17 |
| 43 | 1 | 0, 1, 2, 3, 7, 8, 9, 11, 13, 17, 18, 20 |
| | 42 | 1, 2, 5, 6, 7, 8, 9, 16, 17, 19, 21 |
| 47 | 1 | 0, 1, 4, 6, 9, 10, 11, 16, 18, 19, 20, 22, 23 |
| | 46 | 2, 3, 5, 9, 13, 15, 16, 18, 19, 21, 22, 23 |
| 53 | 1 | 0, 1, 4, 5, 8, 10, 12, 13, 14, 16, 19, 20, 21, 22 |
| | 2 | 1, 3, 7, 8, 11, 12, 15, 16, 18, 21, 24, 25, 26 |
| 59 | 1 | 0, 1, 3, 5, 10, 11, 13, 14, 15, 17, 19, 22, 23, 24, 25, 29 |
| | 58 | 1, 3, 6 |

| Exclusion number $p$ | $\frac{A}{f}$ | $\frac{f}{A}$ | $fgRp$ Allowed | | Excluded |
|---|---|---|---|---|---|
| 3 | 1 | 1 | 0 | 1 . 2 | |
| | 2 | 2 | 1 . 2 | | 0 |
| 5 | 1 | 1 | 0 | 1 . 4 | 2 . 3 |
| | 2 | 3 | 1 . 4 | | 0 . 2 . 3 |
| | 3 | 2 | 2 . 3 | | 0 . 1 . 4 |
| | 4 | 4 | 0 | 2 . 3 | 1 . 4 |
| 7 | 1 | 1 | 0 . 2 . 5 | 1 . 6 | 3 . 4 |
| | 2 | 4 | 0 . 1 . 6 | 3 . 4 | 2 . 5 |
| | 3 | 5 | 1 . 3 . 4 . 6 | | 0 . 2 . 5 |
| | 4 | 2 | 0 . 3 . 4 | 2 . 5 | 1 . 6 |
| | 5 | 3 | 1 . 2 . 5 . 6 | | 0 . 3 . 4 |
| | 6 | 6 | 2 . 3 . 4 . 5 | | 0 . 1 . 6 |
| 11 | 1 | 1 | 0 . 3 . 5 . 6 . 8 | 1 . 10 | 2 . 4 . 7 . 9 |
| | 2 | 6 | 1 . 2 . 3 . 8 . 9 . 10 | | 0 . 4 . 5 . 6 . 7 |
| | 3 | 4 | 0 . 3 . 4 . 7 . 8 | 5 . 6 | 1 . 2 . 9 . 10 |
| | 4 | 3 | 0 . 1 . 5 . 6 . 10 | 2 . 9 | 3 . 4 . 7 . 8 |
| | 5 | 9 | 0 . 1 . 2 . 9 . 10 | 4 . 7 | 3 . 5 . 6 . 8 |
| | 6 | 2 | 1 . 4 . 5 . 6 . 7 . 10 | | 0 . 2 . 3 . 8 . 9 |
| | 7 | 8 | 2 . 3 . 5 . 6 . 8 . 9 | | 0 . 1 . 4 . 7 . 10 |
| | 8 | 7 | 2 . 4 . 5 . 6 . 7 . 9 | | 0 . 1 . 3 . 8 . 10 |
| | 9 | 5 | 0 . 2 . 4 . 7 . 9 | 3 . 8 | 1 . 5 . 6 . 10 |
| | 10 | 10 | 1 . 3 . 4 . 7 . 8 . 10 | | 0 . 2 . 5 . 6 . 9 |
| 13 | 1 | 1 | 0 . 2 . 6 . 7 . 11 | 1 . 12 | 3 . 4 . 5 . 8 . 9 . 10 |
| | 2 | 7 | 1 . 4 . 5 . 8 . 9 . 12 | | 0 . 2 . 3 . 6 . 7 . 10 . 11 |
| | 3 | 9 | 0 . 2 . 5 . 8 . 11 | 4 . 9 | 1 . 3 . 6 . 7 . 10 . 12 |
| | 4 | 10 | 0 . 1 . 4 . 9 . 12 | 2 . 11 | 3 . 5 . 6 . 7 . 8 . 10 |
| | 5 | 8 | 1 . 2 . 3 . 10 . 11 . 12 | | 0 . 4 . 5 . 6 . 7 . 8 . 9 |
| | 6 | 11 | 3 . 4 . 6 . 7 . 9 . 10 | | 0 . 1 . 2 . 5 . 8 . 11 . 12 |
| | 7 | 2 | 2 . 4 . 6 . 7 . 9 . 11 | | 0 . 1 . 3 . 5 . 8 . 10 . 12 |
| | 8 | 5 | 2 . 3 . 5 . 8 . 10 . 11 | | 0 . 1 . 4 . 6 . 7 . 9 . 12 |
| | 9 | 3 | 0 . 5 . 6 . 7 . 8 | 3 . 10 | 1 . 2 . 4 . 9 . 11 . 12 |
| | 10 | 4 | 0 . 1 . 3 . 10 . 12 | 6 . 7 | 2 . 4 . 5 . 8 . 9 . 11 |
| | 11 | 6 | 1 . 5 . 6 . 7 . 8 . 12 | | 0 . 2 . 3 . 4 . 9 . 10 . 11 |
| | 12 | 12 | 0 . 3 . 4 . 9 . 10 | 5 . 8 | 1 . 2 . 6 . 7 . 11 . 12 |
| 17 | 1 | 1 | 0 . 3 . 4 . 6 . 11 . 13 . 14 | 1 . 16 | 2 . 5 . 7 . 8 . 9 . 10 . 12 . 15 |
| | 2 | 9 | 0 . 1 . 2 . 7 . 10 . 15 . 16 | 6 . 11 | 3 . 4 . 5 . 8 . 9 . 12 . 13 . 14 |
| | 3 | 6 | 1 . 2 . 4 . 6 . 11 . 13 . 15 . 16 | | 0 . 3 . 5 . 7 . 8 . 9 . 10 . 12 . 14 |
| | 4 | 13 | 0 . 5 . 6 . 8 . 9 . 11 . 12 | 2 . 15 | 1 . 3 . 4 . 7 . 10 . 13 . 14 . 16 |
| | 5 | 7 | 1 . 2 . 3 . 8 . 9 . 14 15 . 16 | | 0 . 4 . 5 . 6 . 7 . 10 . 11 . 12 . 13 |
| | 6. | 3 | 2 . 5 . 6 . 7 . 10 . 11 . 12 . 15 | | 0 . 1 . 3 . 4 . 8 . 9 . 13 . 14 . 16 |
| | 7 | 5 | 3 . 4 . 5 . 7 . 10 . 12 . 13 . 14 | | 0 . 1 . 2 . 6 . 8 . 9 . 11 . 15 . 16 |
| | 8 | 15 | 0 . 2 . 3 . 4 . 13 . 14 . 15 | 5 . 12 | 1 . 6 . 7 . 8 . 9 . 10 . 11 . 16 |
| | 9 | 2 | 0 . 1 . 5 . 8 9 . 12 . 16 | 3 . 14 | 2 . 4 . 6 . 7 . 10 . 11 . 13 . 15 |
| | 10 | 12 | 1 . 3 . 5 . 6 . 11 . 12 . 14 . 16 | | 0 . 2 . 4 . 7 . 8 . 9 . 10 . 13 . 15 |
| | 11 | 14 | 3 . 6 . 7 . 8 . 9 . 10 . 11 . 14 | | 0 . 1 . 2 . 4 . 5 . 12 . 13 . 15 . 16 |
| | 12 | 10 | 2 . 4 . 5 . 8 . 9 . 12 . 13 . 15 | | 0 . 1 . 3 . 6 . 7 . 10 . 11 . 14 . 16 |
| | 13 | 4 | 0 . 2 . 3 . 7 . 10 . 14 . 15 | 8 . 9 | 1 . 4 . 5 . 6 . 11 . 12 . 13 . 16 |
| | 14 | 11 | 1 . 4 . 7 . 8 . 9 . 10 . 13 . 16 | | 0 . 2 . 3 . 5 . 6 . 11 . 12 . 14 . 15 |
| | 15 | 8 | 0 . 4 . 6 . 8 . 9 . 11 . 13 | .7 . 10 | 1 . 2 . 3 . 5 . 12 . 14 . 15 . 16 |
| | 16 | 16 | 0 . 1 . 5 . 7 . 10 . 12 . 16 | 4 . 13 | 2 . 3 . 6 . 8 . 9 . 11 . 14 . 15 |
| $p$ | $\frac{A}{f}$ | $\frac{f}{A}$ | Excluded | $fgNp$ | Allowed |

## POSTHUMOUS PAPERS. AUXILIARY TABLE FOR SOLVING THE EQUATION

$$A = fxx + gyy.$$

| Exclusion number $p$ | $\frac{A}{f}$ | $\frac{f}{A}$ | $fg\,R\,p$ Allowed | | $fg\,R\,p$ Excluded |
|---|---|---|---|---|---|
| 19 | 1 | 1 | 0 2 . 3 . 4 . 7 . 12 . 15 . 16 . 17 | 1 . 18 | 5 . 6 . 8 9 . 10 . 11 . 13 . 14 |
| | 2 | 10 | 1 . 2 . 4 . 6 . 9 . 10 . 13 . 15 . 17 . 18 | | 0 . 3 . 5 . 7 . 8 . 11 . 12 . 14 . 16 |
| | 3 | 13 | 4 . 5 . 6 . 7 . 9 . 10 . 12 . 13 . 14 . 15 | | 0 . 1 . 2 . 3 . 8 . 11 . 16 . 17 . 18 |
| | 4 | 5 | 0 . 4 . 5 . 6 . 8 . 11 . 13 . 14 . 15 | 2 . 17 | 1 . 3 . 7 . 9 . 10 . 12 . 16 . 18 |
| | 5 | 4 | 0 . 1 . 2 . 6 . 8 . 11 . 13 . 17 . 18 | 9 . 10 | 3 4 . 5 . 7 . 12 . 14 . 15 . 16 |
| | 6 | 16 | 0 . 1 . 3 . 4 . 9 . 10 . 15 . 16 . 18 | 5 . 14 | 2 6 . 7 . 8 . 11 . 12 . 13 . 17 |
| | 7 | 11 | 0 . 1 . 3 . 5 . 6 . 13 . 14 . 16 . 18 | 8 . 11 | 2 . 4 . 7 . 9 . 10 . 12 . 13 . 15 |
| | 8 | 12 | 1 . 2 . 4 . 7 . 8 . 11 . 12 . 15 . 17 . 18 | | 0 . 3 . 5 . 6 9 . 10 . 13 14 . 16 |
| | 9 | 17 | 0 . 2 . 6 . 7 . 9 . 10 . 12 . 13 . 17 | 3 . 16 | 1 . 4 . 5 . 8 . 11 . 14 . 15 . 18 |
| | 10 | 2 | 1 . 2 . 3 . 5 . 9 . 10 . 14 . 16 . 17 . 18 | | 0 . 4 . 6 . 7 . 8 . 11 . 12 . 13 . 15 |
| | 11 | 7 | 0 . 2 5 . 8 . 9 . 10 . 11 . 14 . 17 | 7 . 12 | 1 . 3 . 4 6 . 13 . 15 . 16 . 18 |
| | 12 | 8 | 1 · 5 . 7 . 8 . 9 . 10 . 11 . 12 . 14 . 18 | | 0 . 2 . 3 . 4 . 6 . 13 . 15 . 16 . 17 |
| | 13 | 3 | 2 . 3 . 4 . 5 . 8 . 11 . 14 . 15 . 16 . 17 | | 0 . 1 . 6 . 7 . 9 . 10 . 12 . 13 . 18 |
| | 14 | 15 | 3 . 4 . 6 . 8 . 9 . 10 . 11 . 13 . 15 . 16 | | 0 . 1 . 2 5 . 7 . 12 . 14 . 17 . 18 |
| | 15 | 14 | 2 . 3 . 5 . 6 . 7 . 12 . 13 . 14 . 16 . 17 | | 0 . 1 . 4 . 8 . 9 . 10 . 11 . 15 . 18 |
| | 16 | 6 | 0 . 3 . 7 . 8 . 9 . 10 . 11 . 12 . 16 | 4 . 15 | 1 . 2 . 5 . 6 . 13 . 14 . 17 . 18 |
| | 17 | 9 | 0 . 1 . 4 . 5 . 6 . 13 . 14 . 15 . 18 | 6 . 13 | 2 . 3 . 8 . 9 . 10 . 11 . 16 . 17 |
| | 18 | 18 | 1 . 3 . 6 . 7 . 8 . 11 . 12 . 13 . 16 . 18 | | 0 . 2 . 4 . 5 . 9 . 10 . 14 . 15 . 17 |
| 23 | 1 | 1 | 0.4.8.9.10.11.12.13.14.15.19 | 1 . 22 | 2 . 3 . 5 . 6 . 7 . 16 . 17 . 18 . 20 . 21 |
| | 2 | 12 | 0 . 1 . 3 . 4 . 6 9 . 14 . 17 . 19 . 20 . 22 | 5 . 18 | 2 . 7 . 8 . 10 . 11 . 12 . 13 . 15 . 16 . 21 |
| | 3 | 8 | 0 . 1 . 5 . 6 . 8 . 10 . 13 . 15 . 17 . 18 . 22 | 7 . 16 | 2 . 3 . 4 . 9 . 11 . 12 . 14 . 19 . 20 . 21 |
| | 4 | 6 | 0 . 1 . 3 . 5 . 7 . 8 . 15 . 16 . 18 . 20 . 22 | 2 . 21 | 4 . 6 . 9 . 10 . 11 . 12 . 13 . 14 . 17 . 19 |
| | 5 | 14 | 1 . 2 . 4 . 5 . 7 . 9 . 14 . 16 . 18 . 19 . 21 . 22 | | 0 . 3 . 6 . 8 . 10 . 11 . 12 . 13 . 15 . 17 . 20 |
| | 6 | 4 | 0 . 2 . 4 . 5 . 6 . 7 . 16 . 17 . 18 . 19 . 21 | 11 . 12 | 1 . 3 . 8 . 9 . 10 . 13 . 14 . 15 . 20 . 22 |
| | 7 | 10 | 1 . 2 . 7 . 8 . 9 . 11 12 . 14 . 15 . 16 . 21 . 22 | | 0 . 3 4 . 5 . 6 . 10 . 13 . 17 . 18 . 19 . 20 |
| | 8 | 3 | 0 . 2 . 5 . 6 . 8 . 11 . 12 . 15 17 . 18 . 21 | 10 . 13 | 1 . 3 . 4 . 7 . 9 . 14 . 16 . 19 . 20 . 22 |
| | 9 | 18 | 0 . 1 . 4 7 . 10 . 11 . 12 . 13 . 16 . 19 . 22 | 3 . 20 | 2 . 5 . 6 . 8 . 9 . 14 . 15 . 17 . 18 . 21 |
| | 10 | 7 | 1 . 2 . 3 . 5 . 10 . 11 . 12 . 13 . 18 . 20 . 21 . 22 | | 0 . 4 . 6 . 7 . 8 . 9 . 14 . 15 . 16 . 17 . 19 |
| | 11 | 21 | 3 . 4 . 5 . 7 . 8 . 10 13 . 15 . 16 . 18 . 19 . 20 | | 0 . 1 . 2 . 6 . 9 . 11 . 12 . 14 . 17 . 21 . 22 |
| | 12 | 2 | 0 . 2 . 3 . 7 . 10 . 11 . 12 . 13 . 16 . 20 . 21 | 9 . 14 | 1 . 4 . 5 . 6 . 8 . 15 . 17 . 18 . 19 . 22 |
| | 13 | 16 | 0 . 1 . 2 . 3 . 8 . 9 . 14 . 15 . 20 . 21 . 22 | 6 . 17 | 4 . 5 . 7 . 10 . 11 . 12 . 13 . 16 . 18 . 19 |
| | 14 | 5 | 1 . 5 . 6 . 9 . 10 11 . 12 . 13 14 . 17 . 18 . 22 | | 0 . 2 . 3 . 4 . 7 . 8 . 15 . 16 . 19 . 20 . 21 |
| | 15 | 20 | 3 . 5 . 6 . 7 . 9 . 11 . 12 . 14 . 16 . 17 . 18 . 20 | | 0 . 1 . 2 . 4 . 8 . 10 . 13 15 . 19 . 21 . 22 |
| | 16 | 13 | 0 . 2 . 6 . 7 . 9 . 10 . 13 . 14 . 16 . 17 21 | 4 . 19 | 1 . 3 . 5 . 8 . 11 . 12 . 15 . 18 . 20 . 22 |
| | 17 | 19 | 1 . 2 . 3 . 4 . 6 . 10 . 13 . 17 . 19 . 20 . 21 . 22 | | 0 . 5 . 7 . 8 . 9 . 11 . 12 . 14 . 15 . 16 . 18 |
| | 18 | 9 | 0 . 3 . 4 . 5 . 9 . 11 . 12 . 14 . 18 . 19 . 20 | 8 . 15 | 1 . 2 . 6 . 7 . 10 . 13 . 16 . 17 . 21 . 22 |
| | 19 | 17 | 1 . 4 . 6 . 7 . 8 . 11 . 12 . 15 . 16 . 17 . 19 . 22 | | 0 . 2 . 3 . 5 . 9 . 10 . 13 . 14 . 18 . 20 . 21 |
| | 20 | 15 | 2 . 4 . 5 . 8 . 9 . 10 . 13 . 14 . 15 . 18 19 . 21 | | 0 . 1 . 3 . 6 . 7 . 11 . 12 . 16 . 17 . 20 . 22 |
| | 21 | 11 | 3 . 6 . 7 . 8 . 9 . 10 . 13 . 14 . 15 . 16 . 17 . 20 | | 0 . 1 . 2 . 4 . 5 . 11 . 12 . 18 . 19 . 21 . 22 |
| | 22 | 22 | 2 . 3 . 4 . 6 8 . 11 . 12 . 15 . 17 . 19 . 20 . 21 | | 0 . 1 . 5 . 7 . 9 . 10 . 13 . 14 . 16 . 18 . 22 |

| $p$ | $\frac{A}{f}$ | $\frac{f}{A}$ | $fg\,N\,p$ Excluded | | $fg\,N\,p$ Allowed |
|---|---|---|---|---|---|

## SECTION EIGHT.

### A FULLER CONSIDERATION OF CERTAIN INVESTIGATIONS CONCERNING CIRCLE DIVISION.

---

**367.**

What we set out in the latter part of Section Seven, from Article 355 onward, certainly gives substantial examples of the great fertility of the theory of circle division, and of the remarkable connection linking this subject with various *arithmetic* investigations. There, however, the severe limits of space and time allowed us only to touch lightly on this field, which rewards our efforts with a richer harvest the further we advance into it. We therefore intend to take up here again one or another of the questions begun there and discuss them more fully: readers will certainly see with great wonder that the solution of several arithmetic problems, which anyone would have expected to be worlds apart from this subject, rests on this foundation.

**368.**

The investigation begun in Article 356 offers an especially fruitful subject. There we divided the collection of roots of the equation $x^n - 1 = 0$ (excluding unity) into two classes and showed how to determine the sum in each class. These sums were $= -\frac{1}{2} + \frac{1}{2}\sqrt{n}$ and $-\frac{1}{2} - \frac{1}{2}\sqrt{n}$ when $n$ has the form $4n+1$, or $= -\frac{1}{2} + \frac{1}{2}\sqrt{-n}$ and $-\frac{1}{2} - \frac{1}{2}\sqrt{-n}$ when $n$ has the form $4n+3$. Yet there we had not only restricted ourselves to the case where $n$ is a *prime number*; we had also, what was of much greater importance still, left the *sign* of the radical quantity undetermined, or rather neglected to support this determination, indicated in a few words, with a solid proof. We must therefore make good these deficiencies before anything else.

**.369.**

Now let $n$ be any positive integer, and $R$ a root of the equation $x^n - 1. = 0$ none of whose powers preceding the power with exponent $n$ equals unity (see Article 359, II.). As in Section VII, let us denote by $[\lambda]$ the power $R^\lambda$, so that $[0] = 1$, $[1]$, $[2]$, $[3] \ldots . [n-1]$ represent all the roots of the equation $x^n - 1 = 0$. Further, let us denote the sum

$$[0] + [1] + [4] + [9] + \ldots . + [(n-1)^2] \quad \text{by } \Sigma[\mathfrak{Q}]$$

and more generally

$$[0] + [\lambda] + [4\lambda] + [9\lambda] \ldots . + [\lambda(n-1)^2] \quad \text{by } \Sigma[\mathfrak{Q}\lambda]$$

so that $\mathfrak{Q}$ denotes, without further specification, the squares of the numbers $0, 1, 2, 3 \ldots . n-1$. Thus, just as in general $[\lambda] = [\mu]$ when $\lambda$, $\mu$ are any integers (positive or negative) congruent modulo $n$, we also have $\Sigma[\mathfrak{Q}\lambda] = \Sigma[\mathfrak{Q}\mu]$ when $\lambda \equiv \mu$. With these preparations we have the following.

**370.**

PROBLEM. *Determine the product of the two sums* $\Sigma[\mathfrak{Q}]$ *and* $\Sigma[-\mathfrak{Q}]$.

*Solution.* Since $nn \equiv 0$, $(n+1)^2 \equiv 1$, $(n+2)^2 \equiv 4$, etc. (mod.$n$), it is easy to see that $\Sigma[\mathfrak{Q}]$ becomes

$$\begin{aligned} &= [1] + [4] + [9] + [16] \ldots . + [nn] \\ &= [4] + [9] + [16] + [25] \ldots . + [(n+1)]^2 \\ &= [9] + [16] + [25] + [36] \ldots . + [(n+2)^2] \end{aligned}$$

etc. or in general

$$= [kk] + [(k+1)^2] + [(k+2)^2] + [(k+3)^2] \ldots . + [(n+k-1)^2]$$

Hence $[-kk] \times \Sigma[\mathfrak{Q}]$

$$= [0] + [2k+1] + [4k+4] + [6k+9] \ldots . + [(n-1)^2 + 2(n-1)k]$$

Hence $\Sigma[-\mathfrak{Q}] \times \Sigma[\mathfrak{Q}]$ expands into

$$\begin{aligned} &- [0] + [1] + [4] + [9] \ldots . + [(n-1)^2] \\ &+ [0] + [3] + [8] + [15] \ldots . + [nn-1] \\ &+ [0] + [5] + [12] + [21] \ldots . + [nn+2n-3] \\ &+ [0] + [7] + [16] + [27] \ldots . + [nn+4n-5] \\ &+ \text{etc.} \\ &+ [0] + [2n-1] + [4n] + [6n+3] \ldots . + [3nn-6n+3] \end{aligned}$$

Summing these parts *vertically* gives

$$\begin{aligned} n[0] + [1] \times \frac{1-[2n]}{1-[2]} + [4]\frac{1-[4n]}{1-[4]} + [9]\frac{1-[6n]}{1-[6]} \\ + \text{etc.} + [(n-1)^2] \times \frac{1-[2nn-2n]}{1-[2n-2]} \end{aligned}$$

In this expression all parts except the first vanish whenever $n$ is odd; for then all of $1-[2n]$, $1-[4n]$, $1-[6n]$, etc. become $= 0$, whereas none of the denominators $1-[2]$, $1-[4]$, $1-[6]$, $1-[8]$, etc. up to $1-[2n-2]$ does. When $n$ is even, however, one of the denominators is also $= 0$, namely $1-[n]$, to which the term $[\frac{1}{4}nn] \times \frac{1-[nn]}{1-[n]}$ corresponds; the sum of the parts from which

this term arose becomes $= n[\frac{1}{4}nn]$. Here again we must distinguish two cases. When $n$ is divisible by four, $\frac{1}{4}nn \equiv 0 \,(\mathrm{mod}.n)$ and therefore $[\frac{1}{4}nn] = 1$; but when $n$ is twice an odd number, $\frac{1}{4}nn \equiv \frac{1}{2}n \,(\mathrm{mod}.n)$ and therefore necessarily $[\frac{1}{4}nn] = -1$. We finally conclude that

1) for an odd value of $n$, the required product is $= n$
2) for a value divisible by four, the product is $= 2n$
3) for a value that is twice an odd number, it is $= 0$. Q. E. I.

**371.**

It is now worth considering the nature of the sum $\Sigma[\mathfrak{Q}]$ more closely.

I. Since we may replace the squares 0, 1, 4, 9, 16, etc. by their least residues modulo $n$, it is evident that if $M$ denotes, without further specification, the quadratic residues of the number $n$ from 0 to $n-1$, and $m$ the number of roots of the congruence $xx \equiv M(\mathrm{mod}.m)$, then $\Sigma[\mathfrak{Q}] = \Sigma m[M]$. We showed how to determine the number $m$ in Articles 104, 105.

II. If $n$ is an (odd) prime, then for $M = 0$ we have $m = 1$, whereas for every other value of $M$ we have $m = 2$. If $n$ is a power of an odd prime $= p^\nu$, then $m = 2$ for every value of $M$ not divisible by $p$ —

— — — — — — — — — — — — — — — — —

**372.**

If $n$ is an (odd) prime, the residues $m$ consist of zero, for which $M = 1$, and $\frac{1}{2}(n-1)$ other numbers, for which $M = 2$. Denoting these residues (excluding the residue 0), without further specification, by $\mu$, our progression is $= 1 + 2\Sigma r^\mu$. Further, if $\nu$ denotes, without further specification, all remaining numbers below $n$, whose number is likewise $\frac{1}{2}(n-1)$ and which comprise all quadratic nonresidues of $n$ below $n$, then evidently

$$1 + \Sigma r^\mu + \Sigma r^\nu = 1 + r + rr + r^3 \ldots + r^{n-1} = \frac{1 - r^n}{1 - r} = 0$$

Thus, setting the sum of our progression, that is $1 + 2\Sigma r^\mu = A$, we have $1 + 2\Sigma r^\nu = -A$, and also $\Sigma r^\mu - \Sigma r^\nu = A$.

By Article 356 we therefore have $A = \pm\sqrt{n}$ or $\pm\sqrt{-n}$, according as $n$ is $\equiv 1$ or $\equiv 3 \,(\mathrm{mod}.4)$. But this does not yet determine the sign of the root.

If, in our progression, which we shall denote by $\Pi$, we replace $r$ by another similar root of the equation $x^n - 1 = 0$, say $r' = r^\lambda$, let $\Pi'$ be the result.

**373.**

If $n$ is the square or a higher power of a prime, say $= p^\pi$, some residues $m$ consist of numbers not divisible by $p$; others are divisible by $pp$ but not by a higher power of $p$; others can be divided by $p^4$ but not. by $p^5$; and so on to those divisible by $p^{\pi-2}$ but not by $p^{\pi-1}$, or by $p^{\pi-1}$ but not by $p^\pi$, according as $\pi$ is even or odd. Finally there is the residue 0, the only one divisible by $p^\pi$ (compare Article 102). Now denote, without further specification, by $\mu$ the quadratic residues of the number $p$ below $p$, excluding zero (whose number is $= \frac{1}{2}(p-1)$). Those different classes of residues are then represented as follows. The first, which are not divisible by $p$, are represented by $\mu + kp$, where for $k$ we must substitute all integers from 0 to $p^{\pi-1} - 1$, so that the number of all residues contained in this form is $= \frac{1}{2}(p-1)p^{\pi-1}$; for each of these, $M = 2$. The sum of all terms in $\Pi$ corresponding to these residues is

$$= 2\Sigma r^{\mu+kp} = 2\Sigma r^\mu \,.\, \Sigma r^{kp}$$
$$= 2\Sigma r^\mu \,.\, \frac{r^{p^\pi} - 1}{r^p - 1} = 0$$

The second class of residues is represented by $\mu pp + kp^3$, where for $k$ we must substitute all integers from 0 to $p^{\pi-3} - 1$, so that the number of all residues contained in this form is $= \frac{1}{2}(p-1)p^{\pi-3}$; for each of them, however, $M = 2p$. The sum of the terms in $\Pi$ arising from these becomes

$$= 2p\Sigma r^{\mu pp+kp^3} = 2p\Sigma r^{\mu pp} \,.\, \Sigma r^{kp^3}$$
$$= 2p\Sigma r^{\mu pp} \,.\, \frac{r^{p^\pi} - 1}{r^{p^3} - 1} = 0$$

provided that $\pi > 3$. Similarly, the third, fourth, etc. class is represented by $\mu p^4 + kp^5$, $\mu p^6 + kp^7$, etc., where for $k$ we must take all integers from 0 to $p^{\pi-5} - 1$, $p^{\pi-7} - 1$, etc.; for these, $M = 2pp$, $M = 2p^3$, etc. The sum of the terms in $\Pi$ arising from the third, fourth, etc. class vanishes provided that $\pi > 5$, $\pi > 7$, etc.. respectively.

We conclude that when $\pi$ is even, in $\Pi$ only the terms corresponding to the residue 0 remain, and these are $= 1$; for these, however, $M = p^{\frac{1}{2}\pi}$, so that the sum of all terms in $\Pi$ is $= p^{\frac{1}{2}\pi}$.

---

## GAUSS TO DIRICHLET.

To Monsieur
Monsieur Lejeune Dirichlet in Paris.

I would have thanked you earlier for the memoir you so kindly sent me and for the great pleasure it gave me, had I not wished first to hear something of the outcome of my efforts in Berlin concerning your wishes and, I may add, my own. I am extremely pleased now to see, from a letter received from the Secretary of the Academy in Berlin, that we may hope that they will soon be willing to secure a suitable permanent position for you in your homeland.

The rarer this inclination is, the more pleased I am to find you so strongly drawn to the part of mathematics that has always been my favourite study. I sincerely wish you circumstances in which you remain as far as possible master of your time and of the choice of your work. Immediately after the appearance of my Disquisitiones, other kinds of work, and later my circumstances, greatly hindered me from following my inclination as much as I would have wished. Instead of a second part of that work, which I formerly intended, I shall in all probability have to confine myself to presenting a memoir on an individual subject from time to time. The three memoirs of this kind that have so far appeared in the 16th volume of the local Commentationes and in the first and fourth volumes of the Commentationes recentiores, however, contain (apart from a portion of the second) none of the subjects I already had in mind in 1801 for the continuation, but new ones. My later work of this kind likewise concerns a new subject, namely the theory of biquadratic residues, which I intend to present in about three memoirs; the first of these will shortly be printed for the sixth volume of the Comment. rec., and the main material for the remainder, as well as for the similar theory of cubic residues, may be regarded as essentially settled, although little of it has yet been properly put on paper.

Please give my regards to Herr von Humboldt, if he is still in Paris, and excuse my not writing to him directly now by my concern that the letter might not reach him, since, as I hear, he intended to leave Paris.

With sincere esteem

Yours most devotedly

Göttingen, 13 September 1826. C. F. Gauss.

I offer you, my highly esteemed friend, my most sincere thanks for your kind letter and for kindly sending your two memoirs. I am pleased to see the growing interest now beginning to be taken in investigations of higher arithmetic. I very much liked the successful way in which you derive the second, concerning the biquadratic residuacity of the number 2, from the first.

Presumably the 6th volume of our Commentationes has now reached Breslau, so you will probably be familiar by now with my Commentatio prima on biquadratic residues: if an opportunity arises, I would also gladly send you a separate copy. I could have chosen among several methods of proving the theorem in it; but you will see why I preferred the method used there, chiefly because the classification of 2 for those moduli for which it is a quadratic nonresidue (under $B$ or $D$) must be regarded as an essential part of the theorem, to which most other methods of proof seem inapplicable.

I have had all the material for the entire investigation for 23 years, and the proofs of its main theorems (among which the theorem in the first memoir is *not* yet to be counted) for about 14 years — (although I wish and hope to be able to simplify something further in the latter, the proofs) —. I have planned the investigation for about 3 memoirs. I have already begun writing the second, and hope to finish it before long, unless the surveying duties recently assigned to me again cause some delay.

I had already also announced the concluding theorem $b \equiv \frac{1}{2}rr \,(\mathrm{mod}.p)$ in the local gel. Anzeigen three years ago and drawn attention to the remarkable knot still to be untied in connection with it; but I have not yet heard of anyone attempting it. A few days ago I did succeed with *one half*, and this discovery gave me all the more pleasure because it does not rest on induction at all — for I confess that I would not have expected precisely *this* connection — but a priori on combining other *very intricate* and interesting investigations, already 28 years old but still entirely unpublished, of which a faint indication is given in the closing note of the Disquis. Arithm. p. 668 [Gauss Werke vol. I. p. 466].

This is a sufficient criterion for the case where $p$ has the form $8n + 5$.

Let the number of classes formed by the binary forms in each of the two kinds for the determinant $-p$ be $= k$. The beginning of a table I constructed up to the determinant $-3000$ is in Disquis. Arithm. p. 520. [Art. 303.] It should also be noted that for a $p$ of the assumed form we always have $k = 2m+1$ when $m$ denotes the number of decompositions of $p$ into three positive squares (I say positive to exclude 0), as Legendre found by induction and

as was first proved in the Disquis. Arithm. from the theory of *ternary* forms. For example,

| for $p =$ | 5, | 13, | 29, | 37, | 53, | 61, | 101, | 109, | 149, | 157 | etc. |
|---|---|---|---|---|---|---|---|---|---|---|---|
| $k =$ | 1, | 1, | 3, | 1, | 3, | 3, | 7, | 3, | 7, | 3 | |
| $m =$ | 0, | 0, | 1, | 0, | 1, | 1, | 3, | 1, | 3, | 1 | |
| | | | 4 | — | 1 | 9 | 1 4 16 | 9 | 1 4 36 | 4 | |
| | | | 9 | | 16 | 16 | 36 16 36 | 36 | 4 64 64 | 9 | |
| | | | 16 | | 36 | 36 | 64 81 49 | 64 | 144 81 49 | 144 | |

With this assumption, the value of $b$ that is $\equiv \frac{1}{2}rr \,(\text{mod.}p)$ is always

$$\equiv 2k + a - 1 \equiv 4m + a + 1 \;(\text{mod.}8)$$

which completely determines the sign of $b$. Here are 22 examples, doubling the extent of the table given at the end of the memoir.

| $p$ | $k$ | $a$ | $b$ | $p$ | $k$ | $a$ | $b$ |
|---|---|---|---|---|---|---|---|
| 5 | 1 | +1 | +2 | 181 | 5 | +9 | +10 |
| 13 | 1 | −3 | −2 | 197 | 5 | +1 | −14 |
| 29 | 3 | +5 | +2 | 229 | 5 | −15 | +2 |
| 37 | 1 | +1 | −6 | 269 | 11 | +13 | +10 |
| 53 | 3 | −7 | −2 | 277 | 3 | +9 | +14 |
| 61 | 3 | +5 | −6 | 293 | 9 | +17 | +2 |
| 101 | 7 | +1 | −10 | 317 | 5 | −11 | +14 |
| 109 | 3 | −3 | +10 | 349 | 7 | +5 | +18 |
| 149 | 7 | −7 | −10 | 373 | 5 | −7 | +18 |
| 157 | 3 | −11 | −6 | 389 | 11 | +17 | −10 |
| 173 | 7 | +13 | +2 | 397 | 3 | −19 | −6 |

The rule can therefore also be expressed as follows; (always assuming $p \equiv 5 \;(\text{mod.}8)$)

$$\begin{aligned} \text{We have } b &\equiv a + 1 \;(\text{mod.}8), && \text{if } m \text{ is even} \\ b &\equiv a + 5 && \text{if } m \text{ is odd.} \end{aligned}$$

I do not yet venture a conjecture as to whether a still simpler criterion is possible for distinguishing the case of even $m$ from the odd case *in advance*, that is, without knowing the value of $m$ itself, since, as I remarked above, this connection is still entirely new.

For the case $p \equiv 1 \pmod{8}$ the above congruence $b \equiv 2k + a - 1 \pmod{8}$ remains correct, but it no longer determines the sign of $b$, since both the positive and the negative values of $b$ satisfy it. Here $k$ is always even, $= 2m$ (with $m$ understood just as above), or $= 2m + 2$ if $m$ denotes the number of decompositions of $p$ into 3 positive *unequal* squares, and $b \equiv 0 \pmod{4}$, or $b \equiv -b \pmod{8}$. I conjecture that the case $p \equiv 1 \pmod{8}$ or $b \equiv 0 \pmod{4}$ calls for deeper investigation, and perhaps again

$$\begin{aligned} b &\equiv 4 \pmod{8} \quad \text{easier than } b \equiv 0 \pmod{8} \\ b &\equiv 8 \pmod{16} \quad \text{easier than } b \equiv 0 \pmod{16} \end{aligned}$$

etc.

With the highest esteem I remain

Yours most devotedly in friendship

Göttingen, 30 May 1828. C. F. Gauss.

---

# REMARKS.

In this second volume of GAUSS's works I have included all his memoirs, essays and tables on higher arithmetic, insofar as they are not already covered by the seven sections of the *Disqu. Arithm.*: the five memoirs published in the '*Commentationes societatis regiae scientiarum Gottingensis*' (in quarto), the reviews of these works of his own and of some works by others published in the '*Göttingischen Gelehrten Anzeigen*' (in octavo) (unsigned by GAUSS, but confirmed as his through the records of the Göttingen University Library), and a selection from his manuscript remains.

In the second printing I have also added the tables 'Circuli quadratura nova', 'On Calculating Logarithms', 'Quadratorum myrias prima', 'Indices of Primes in the Higher Number Domain', 'Auxiliary Table for Solving the Indeterminate Equation $A = fxx + gyy$ by the Method of Exclusion', as well as 'Sectio octava', as far as it was written down, and finally two letters from GAUSS to DIRICHLET, as essential parts of the history of higher arithmetic.

To make the subjects in so extensive a volume easier to survey, the theorems are distinguished typographically in the same way. To make both the earlier editions and the present one easier to use, references to the Disq. Arithm. give article numbers in place of page numbers, and references to memoirs give their own titles in place of their places of publication. The note originally attached to Article 2 of the memoir '*Theorematis arithmetici demonstratio nova*', which contained a correction to Article 139 of the Disqu. Arithm., has been inserted there at the relevant passage. The note on page 91 is taken from a handwritten note. Apart from this, the present edition differs from the earlier ones only in the correction of some printing errors. My additions are identified by square brackets [ ].

The *Table of the Quadratic Character of Primes* is printed in the manner of Tabula II of the Disqu. Arithm., described in Article 99 and especially used (in Article 331) for factoring numbers. The handwriting of the manuscript entitled '*Quadratorum numeris primis divisorum residua lateralia*' most closely resembles that of the second part of the table for converting common fractions into decimal fractions. In place of the horizontal strokes indicating a quadratic residue, it contains small circles; those in adjacent horizontal or vertical rows are always connected by lines. While proofreading I noticed several errors. I then found 190 discrepancies in the characters in a single comparison with JACOBI's *Canon Arithmeticus*, and by determining them directly found agreement with those printed tables. The edition here has been corrected accordingly.

The first part of the *Table for Converting Common Fractions into Decimal Fractions* is arranged here similarly to Tabula III of the Disqu. Arithm. For

the primes and their powers $p^\pi$ between 3 and 463 it contains the mantissas (1), (2)..(0) of the decimal fractions of $\frac{10.r}{p^\pi}$, $\frac{10.rr}{p^\pi}$..$\frac{10}{p^\pi}$. Here $r$ denotes unity, so that (1) = (2) = ..(0), when 10 is a primitive root of $p^\pi$; otherwise $r$ denotes the smallest of those primitive roots of $p^\pi$ for which, as base, the index of 10 takes its smallest value. To make the table easier to use, I have added the values other than 1 of $r$ on page 420. The manuscript, which also does not yet give the distinguishing digits of the different periods, most closely resembles the Analysis residuorum in appearance and seems to precede in date the piece placed here as the second part of the whole table. This second part contains, for the primes and their power $p^\pi$ between 467 and 997, the mantissas of the decimal fractions of $\frac{100}{p^\pi}$. The manuscript gives the divisors in decreasing order and ends with the words: *Explicitus October* 11. 1795. In print I have added, for the divisor 191 in period (1), the 71st digit, and omitted, for the divisor 829, a number between the 151st and 152nd digits.

The first part of the *Table of the Frequency of Primes*, discussed by Gauss himself in a letter (page 444), giving the number of primes in each of the first 1000 thousand groups, exists in a manuscript by Gauss. His manuscript remains, however, do not contain the counts mentioned in the letter of the hundred groups within the first million that contain a specified number of primes. The other part of the table, for the second and third million, comes from a manuscript by Goldschmidt alone. Herr Meissel, by counting and by his formula, has found the following corrections to pages 436 and 437:

| Thousand | Gauss | True Value | Thousand | Gauss | True Value |
|---|---|---|---|---|---|
| 20 | 102 | 104 | 546 | 68 | 69 |
| 159 | 87 | 77 | 601 | 75 | 76 |
| 199 | 96 | 86 | 625 | 68 | 78 |
| 206 | 85 | 83 | 668 | 73 | 74 |
| 245 | 78 | 88 | 675 | 69 | 73 |
| 289 | 85 | 77 | 784 | 74 | 75 |
| 290 | 84 | 85 | 800 | 81 | 71 |
| 334 | 80 | 81 | 879 | 68 | 78 |
| 352 | 80 | 81 | 985 | 74 | 70 |
| 501 | 78 | 79 | | | |

The formula of Encke mentioned in the letter from Gauss to Encke seems to be the following

$$\frac{n}{\log n} \sqrt[2\log n]{10}$$

which Encke gives in a letter to Gauss of 4 December 1849.

The *Table of the Number of Classes of Binary Quadratic Forms* gives the number of genera and classes and the index of irregularity for negative determinants in hundred groups 1 to 30, 43, 51, 61, 62, 63, 91 to 100, 117 to 120;

also, in a separate compilation, for those of the 1st, 3rd and 10th thousand groups, for the first 800 of the forms $-(15n+7)$ and $-(15n+13)$, and for some very large determinants; and for positive determinants in the 1st, 2nd, 3rd, 9th and 10th hundred groups and some others. The manuscript consists of separate slips with tables arranged in various ways. For instance, the older ones use the word Ordo in place of Genus, as in the individual hundred groups except the 9th and 10th for positive determinants, and also in some preliminary compilations in thousand groups. To make the arrangement easier to survey, the notation of the Disqu. Arithm. has been used throughout here. For lack of space, the greatest and least quotients obtained by dividing the number of classes by the determinant, and the numbers of determinants for which the quotient falls within certain bounds, are not placed below the individual hundred groups, but collected at the end of the table for negative determinants. Some surviving notes seem to indicate that Gauss first calculated the classes for determinants belonging to the same hundred group and the same remainder on division by 15. The determinants in these divisions were then arranged by the number of genera and classes, and finally all those in the same hundred group were ordered in the way reproduced here. Several later corrections have been inserted into the tables of individual hundred groups, but not into the compilations in thousand groups. Only the two tables with determinants of the forms $-(15n+7)$ and $-(15n+13)$ contain dates, respectively '*Expl. In. Febr. 1801*' and '*Expl. 27 Febr. 1807.*'

Among the errors I have noticed in these tables are the following. To make them easier to check, I give the period numbers of the fundamental classes, for example 4 . 4 . 2 for the determinant $-11713$, and the numbers represented by forms of the respective fundamental classes, for example 31 . 37 . 2. As in my memoir in volume 14 of the Memoirs of the Royal Society of Sciences at Göttingen, fundamental classes here mean classes which, together with the classes in their periods, produce by composition every properly primitive class of the determinant once and only once.

The following entries have already been omitted:

| | | | |
|---|---|---|---|
| Hundred 9 | G. IV ... | 3., | $-827[21 :: 3]$ |
| 26 | IV | 14 | $-2587[24 :: 11]$ |
| 26 | VIII | 6 | $-2564[56 :: 3]$ |
| 91 | I | 111 | $-9059[117 :: 5]$ |
| 120 | IV | 32 | $-11956^{*}2^{*}[36 . 2 :: 11 . 49]$ |
| 1 | I | 2 | $+37[3 :: 3]$ |
| 2 | I | 2 | $+101[3 :: 4]$ |

and added:

| | | | |
|---|---|---|---|
| Hundred 9 | G. IV . . . | 3 . . | −828[6 . 2 :: 31 . 23] |
| 26 | IV | 14 | −2586[28 . 2 :: 7 . 2] |
| 26 | VIII | 6 | −2565[12 . 2 . 2 :: 7 . 2 . 5] |
| 91 | I | 117 | −9059[117 :: 5] |
| 120 | IV | 32 | −11966*2*[32 . 4 :: 5 . 83] |
| 1 | I | 3 | +37[3 :: 3] |
| 2 | I | 3 | +101[3 :: 4] |

In the table for Hundred 3 and the last one on page 476, which in the manuscript occur together with a record of Hundreds 1 and 2 differing in appearance from the one printed here, the twelve divisions are designated, instead of I. Ordo unicus. 1; I. O. 2; I. O. 3; I. O. 4; II. Ordines duo. 1. 1; II. O. 1. 2; II. O. 2. 2; II. O. 3. 3; III. Ordines quatuor. 1. 1. 1. 1; III. O. 1. 1. 2. 2; III. O. 2. 2. 2. 2; IV. Ordines octo 1. 1. 1. 1. 1. 1. 1. 1, in the manner otherwise used here, as G. I. 1; G. I. 3; G. I. 5; G. I. 7; G. II. 1; G. II. 2; G. II. 3; G. II. 5; G. IV. 1; G. IV. 2; G. IV. 3; G. VIII. 1. Calculation gives, for example, 269. I. 3 [3 :: 4]; 235. IV. 3 [6 . 2 :: 3 . 5]; 401. I. 5 [5 :: 9]; 577. I. 7 [7 :: 3]; 727. II. 5 [10 :: 3]. (Genera I in place of Genus I on page 469 is a printing error).

Because of printing errors, omit:

| | | | |
|---|---|---|---|
| Hundred 27. | G. IV . . . . | 16 . . . | −2624*3* |
| 93 | IV | 16 | −9216 |
| 118 | VIII | 4 | −11713*3* |

and add:

| | | | |
|---|---|---|---|
| Hundred 27. | G. IV . . . . | 16 . . . | −2624*2*[16 . 4 :: 3 . 16] |
| 93 | IV | 16 | −9216*2*[16 . 4 :: 5 . 9] |
| 118 | VIII | 4 | −11713*2*[4 . 4 . 2 :: 31 . 37 . 2] |

According to my calculation, also omit:

| | | | |
|---|---|---|---|
| Hundred 10. | G. II . . . . | 9 . . . | −972[6 . 3 :: 7 . 13] |
| 17 | IV | 4 | −1660[10 . 2 :: 11 . 5] |
| 20 | IV | 12 | −1982[24 :: 3] |
| 21 | IV | 6 | −2096[30 . 2 :: 3 . 4] |
| 23 | IV | 9 | −2221[18 :: 10] |
| 24 | IV | 12 | −2376[12 . 2 . 2 :: 5 . 8 . 8] |
| 29 | IV | 9 | −2887[25 :: 8] |
| 61 | IV | 7 | −6028[12 . 2 :: 13 . 4] |

| | | | |
|---|---|---|---|
| 96 | VIII | 13 | $-9594[20 . 2 . 2 :: 31 . 2 . 13]$ |
| 118 | IV | 25 | $-11780[16 . 4 . 2 :: 3 . 8 . 19]$ |
| 118 | VIII | 16 | $-11780[16 . 4 . 2 :: 3 . 8 . 19]$ |
| 119 | VIII | 16 | $-11840[24 . 2 . 2 :: 5 . 9 . 7]$ |
| Thousand I | G. II . . . | 3 . . . | $-541[10 :: 11]$ |
| I | II | 4 | $-415[10 :: 13]$ |
| I | II | 8 | $-527[18 :: 3]$ |
| I | II | 8 | $-722[18 :: 3]$ |
| I | II | 9 | $-194[20 :: 5]$ |
| I | II | 9 | $-459[6 . 3 :: 5 . 9]$ |
| I | II | 9 | $-972[6 . 3 :: 7 . 13]$ |
| I | II | 11 | $-842[26 :: 13]$ |
| I | IV | 3 | $-784[8 . 2 :: 5 . 4]$ |
| I | IV | 4 | $-532[4 . 2 :: 13 . 7]$ |
| I | IV | 5 | $-425[12 . 2 :: 3 . 17]$ |
| I | IV | 5 | $-608[12 . 2 :: 13 . 27]$ |
| I | IV | 5 | $-629[18 . 2 :: 5 . 2]$ |
| III | II | 15 | $-2578[16 :: 13]$ |
| X | I | 111 | $-9059[117 :: 5]$ |
| forms $-(15n + 13)$ | IV | 4 | $-2788^*2^*[8 . 2 :: 19 . 17]$ |

and add:

| | | | |
|---|---|---|---|
| Hundred 10. | G. II . . . . | 9 | $-972^*3^*[6 . 3 :: 7 . 13]$ |
| 17 | IV | 12 | $-1700[24 . 2 :: 3 . 17]$ |
| 20 | IV | 12 | $-1937[24 . 2 :: 7 . 2]$ |
| 21 | IV | 6 | $-2097[12 . 2 :: 47 . 2]$ |
| 23 | IV | 9 | $-2224[18 . 2 :: 5 . 16]$ |
| 24 | IV | 12 | $-2366[24 . 2 :: 3 . 2]$ |
| 29 | IV | 9 | $-2885[18 . 2 :: 3 . 5]$ |
| 61 | IV | 6 | $-6028[12 . 2 :: 13 . 4]$ |
| 96 | VIII | 13 | $-9546[26 . 2 . 2 :: 5 . 3 . 37]$ |
| 118 | IV | 25 | $-11750[50 . 2 :: 3 . 47]$ |
| 118 | VIII | 16 | $-11780^*2^*[16 . 4 . 2 :: 3 . 8 . 19]$ |
| 119 | VIII | 16 | $-11840^*2^*[16 . 4 . 2 :: 3 . 16 . 5]$ |
| Thousand I. | G. II . . . . | 5 . . . | $-415[10 :: 13]$ |
| I | II | 5 | $-541[10 :: 11]$ |
| I | II | 9 | $-459^*3^*[6 . 3 :: 5 . 9]$ |
| I | II | 9 | $-527[18 :: 3]$ |
| I | II | 9 | $-722[18 :: 3]$ |
| I | II | 9 | $-972^*3^*[6 . 3 :: 7 . 13]$ |
| I | II | 10 | $-194[20 :: 5]$ |

| | | | |
|---|---|---|---|
| I | II | 13 | $-842[26 :: 13]$ |
| I | IV | 2 | $-532[4\,.\,2 :: 13\,.\,7]$ |
| I | IV | 4 | $-784[8\,.\,2 :: 5\,.\,4]$ |
| I | IV | 6 | $-425[12\,.\,2 :: 3\,.\,17]$ |
| I | IV | 6 | $-608[12\,.\,2 :: 13\,.\,27]$ |
| I | IV | 9 | $-629[18\,.\,2 :: 5\,.\,2]$ |
| III | II | 15 | $-2518[30 :: 19]$ |
| X | I | 117 | $-9059[117 :: 5]$ |
| forms $-(15n+13)$ | IV | 4 | $-2788[8\,.\,2 :: 19\,.\,17]$ |

The tables *for Circle Division* give, for 2452 numbers of the forms $aa+1$, $aa{+}4$, $aa{+}9$, $\ldots.\, aa{+}81$, all the odd prime divisors $p$ beside the corresponding $a$, in those cases where all the prime divisors are less than 200; only then are $aa+1$ and so on called factorable.

To make the tables easier to use, Gauss has constructed, for each table from which the complete factorizations of numbers of one of the particular forms can be determined, an auxiliary table giving beside each prime $p$ those numbers $a$ whose square, increased by 1 or 4 . . ., has $p$ as its greatest prime divisor.

The chief purpose of the tables is to facilitate the accurate calculation of arcs whose cotangents are given rational numbers. With their help, arcs for small cotangents can first be combined from arcs for large cotangents, reducing to a very small amount the calculations still needed for the series expressing arcs in terms of their cotangents. The developments towards this end in the manuscript remains are not extensive; the following is the one carried furthest. In it

$$\begin{array}{l}[2]\ [5]\ [13]\ [17]\ [29]\ [37]\ [41]\ [53]\ [61]\ldots[197]\\ (18)\ (57)\ (239)\ \left(\frac{79}{3}\right)\ldots\end{array}$$

denote the arcs with cotangents

$$1\quad 2\quad \frac{3}{2}\quad 4\quad \frac{5}{2}\quad 6\quad \frac{5}{4}\quad \frac{7}{2}\quad \frac{6}{5}\ldots 14\quad 18\quad 57\quad 239\quad \frac{79}{3}\ldots$$

Using the tables to factor $18+i$, $57+i$, $239+i$ into their complex prime factors gives

$$\begin{aligned}(18) &= 2[2]-2[5]-[13]\\ (57) &= -[2]+3[5]-[13]\\ (239) &= 3[2]\qquad -4[13]\end{aligned}$$

and from these

$$\begin{aligned}[2] &= 12(18)+8(57)-5(239)\\ [5] &= 7(18)+5(57)-3(239)\\ [13] &= 9(18)+6(57)-4(239)\end{aligned}$$

further, using the tables,

$$\begin{aligned}(268) &= -2[5] + 2[13] - [17] \quad \cdot \\ (38) &= -[5] \qquad\quad + 2[17]\end{aligned}$$

and from these, by eliminating [17] and substituting the values of [5], [13] obtained above,

$$(38) + 2(268) = (18) - (57) - (239)$$

Eliminating (18) then gives the new determination

$$\begin{aligned}[2] &= 12(38) + 20(57) + 7(239) + 24(268) \\ [5] &= 7(38) + 12(57) + 4(239) + 14(268) \\ [13] &= 9(38) + 15(57) + 5(239) + 18(268) \\ [17] &= 4(38) + 6(57) + 2(239) + 7(268)\end{aligned}$$

After successively using the cotangents 117, 327, 882, 18543, 307, 278, 378, 829, 993, 2943, 447, 606, 931, 1143, 1772, 6118, 34208, 44179, 85353, 485298, 17772, 9466, 330182, 5257, 114669, 12943, $[2][5]\ldots[61]$ are finally expressed in terms of (5257), (9466) ... (485298), and their coefficients are collected in the following columns:

| | 5257 | 9466 | 12943 | 34208 | 44179 | 85353 | 114669 | 330182 | 485298 |
|---|---|---|---|---|---|---|---|---|---|
| 2 | +2805 | −398 | +1950 | +1850 | +2021 | +2097 | +1484 | +1389 | +808 |
| 5 | +1656 | −235 | +1151 | +1092 | +1193 | +1238 | +876 | +820 | +477 |
| 13 | +2100 | −298 | +1460 | +1385 | +1513 | +1570 | +1111 | +1040 | +605 |
| 17 | +875 | −124 | +608 | +577 | +630 | +654 | +463 | +433 | +252 |
| 29 | +1359 | −193 | +945 | +896 | +979 | +1016 | +719 | +673 | +391 |
| 37 | +590 | −84 | +410 | +389 | +425 | +441 | +312 | +292 | +170 |
| 41 | +2410 | −342 | +1675 | +1589 | +1736 | +1802 | +1275 | +1193 | +694 |
| 53 | +994 | −141 | +691 | +655 | +716 | +743 | +526 | +492 | +286 |
| 61 | +2481 | −352 | +1725 | +1637 | +1788 | +1855 | +1313 | +1229 | +715 |

The correctness of these equations, which can be used to determine $[2][5]\ldots[61]$, is seen directly from the factorizations supplied by the tables above:

| | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| (5257) = | [2] | +2[5] | −[13] | +[17] | . | . | −[41] | . | −[61] |
| (9466) = | 2[2] | . | . | . | −[29] | −3[37] | . | . | −[61] |
| (12943) = | [2] | −4[5] | +3[13] | . | . | . | . | . | −[61] |
| (34208) = | 2[2] | −[5] | −2[13] | +[17] | +[29] | . | . | −2[53] | . |
| (44179) = | 3[2] | . | −3[13] | −2[17] | −[29] | . | . | +[53] | . |
| (85353) = | −[2] | −[5] | +[13] | −[17] | . | −[37] | +2[41] | −[53] | . |
| (114669) = | −3[2] | . | . | +[17] | . | +[37] | . | +2[53] | +2[61] |
| (330182) = | −4[2] | +5[5] | +[13] | . | +[29] | −[37] | −[41] | . | +[61] |
| (485298) = | −2[2] | −[5] | +4[13] | . | −2[29] | +[37] | . | +[53] | . |

Gauss collects in the following overview the methods used so far by calculators to determine $\frac{\pi}{4} = (1)$

| | |
|---|---|
| Machin | $(1) = 4(5) - (239)$ |
| | also Clausen |
| Euler | $= (2) + (3)$ |
| | (Euler to Goldbach 1746 May 28) |
| Vega | $= 5(7) + 2(\frac{79}{3})$ |
| | (Vega Thesaurus logar. p. 633) |
| Vega | $= 2(3) + (7)$ |
| | also Clausen (Astr. Nachr. vol. 25. p. 209) |
| Rutherford | $= 4(5) - (70) + (99)$ |
| | (Philos. Trans. 1841. p. 283) |
| Dase | $= (2) + (5) + (8)$ |
| | (Crelle Journal. vol. 27. p. 198) |
| Gauss. 1. | $= 12(18) + 8(57) - 5(239)$ |
| Gauss. 2. | $= 12(38) + 20(57) + 7(239) + 24(268)$ |

The first calculations for the tables date from the preparation of the *Disquiss. Arr.*; they were advanced especially in 1846 and 47. By 21 July 1847, 2283 factorizations had been entered in tables in the order reproduced here. The remaining 169 were calculated later, and I have included them in this printing, whose arrangement differs from the original only in using repeated factors instead of powers to make typesetting easier.

The manuscripts containing these last calculations seem to give the results in the form in which they were first found. The order in which the numbers $a$ appear suggests that the divisors of $aa + 1$ and so on were sought only for the smaller numbers, and that the larger ones were obtained from these by particular devices. Only the following rule is recorded, however: *From three numbers* $a$, $2a - n$, $2a + n$ *a fourth is found*

$$\frac{4a^3 - (nn-3)a}{nn+1}$$

*This is always an integer for* $n = 0$ *and* $n = 1$, *otherwise only*

$$\textit{for} \quad a \equiv 0 \quad \textit{and} \quad \equiv \pm\sqrt{-1} \quad \textit{mod}(nn+1) \quad \textit{when } n \textit{ is even}$$

$$\textit{and for} \quad a \equiv 0 \quad \textit{and} \quad \equiv \pm\sqrt{-1} \quad \textit{mod}\frac{nn+1}{2} \quad \textit{when } n \textit{ is odd}$$

| | | | |
|---|---|---|---|
| *Examples* | $a = 253,$ | $n = 6,$ | $1750507$ |
| | $a = 294,$ | $n = 11,$ | $832902$ |
| | $a = 119,$ | $n = 1,$ | $3370437$ |
| | $a = 57,$ | $n = 3,$ | $74043$ |
| | $a = 123,$ | $n = 9,$ | $90657$ |

The fourth number has no prime divisors other than those of the first three, and of these only the odd prime divisors belonging to the number $n$ are excluded.

The table '*Quadratorum myrias prima*' gives the thousands and hundreds in its heading row, the tens and units of the underlying number in its first vertical column, and, in the last vertical column of each individual table the three lowest digits of the square, and in the interior the four or five higher digits of the square.

The table '*Indices of Primes in the Higher Number Domain*' gives the modulus in each case in the top horizontal row, the base used in the second row, the residues in the first vertical column, and the indices in the interior. The stars $*$ denote zero residues.

The manuscript of the fragment '*Eighth Section. A Fuller Consideration of Certain Investigations Concerning Circle Division*' seems to date from the reworking of the 'Analysis residuorum' into the 'Disquisitiones arithmeticae'. The letters from GAUSS to DIRICHLET confirm the view that in higher arithmetic, too, GAUSS discovered considerably more than is found in his manuscript remains.

SCHERING.

# CONTENTS.

## GAUSS'S WORKS, VOLUME II. HIGHER ARITHMETIC.

*Papers.*


| | | |
|---|---|---|
| A New Proof of an Arithmetical Theorem . . . . . | 1808 Jan. . . | Page 1 |
| Summation of Certain Special Series . . . . . | 1808 Aug. . . | — 9 |
| New Proofs and Extensions of the Fundamental Theorem in the Theory of Quadratic Residues . . . . . | 1817 Feb. . | — 47 |
| Theory of Biquadratic Residues. First Paper . . . . . | 1825 Apr. . | — 65 |
| Theory of Biquadratic Residues. Second Paper . . . . . | 1831 Apr. . | — 93 |


*Reviews of His Own Writings.*


| | | |
|---|---|---|
| A New Proof of an Arithmetical Theorem . . . . . | 1808 May . . | — 151 |
| Summation of Certain Special Series . . . . . | 1808 Sept. . | — 155 |
| Of the Fundamental Theorem in the Theory of Residues, etc. . . . . | 1817 March . | — 159 |
| Theory of Biquadratic Residues. Paper I. . . . . | 1825 April . | — 165 |
| Theory of Biquadratic Residues. Paper II. . . . . | 1831 April . | — 169 |


*Reviews of Other Writings.*


| | | |
|---|---|---|
| [Dalberg] Investigations of the Arithmetical and Geometrical Irreducibility of Numbers and Their Powers . . . . . | 1809 March . | — 181 |
| Chernac. Arithmetical Sieve . . . . . | 1812 March . | — 181 |
| Burckhardt. Tables of Divisors . . . . . | 1814 Nov. 1816 Nov. 1817 Aug. . | — 183 |
| Erchinger. Construction of the Seventeen-Sided Polygon . . . . . | 1825 Dec. . | — 186 |
| Seeber. Investigations of the Properties of Positive Ternary Quadratic Forms . . . . . | 1831 July . . | — 188 |

*Manuscript Remains.*

Analysis of Residues:

Chapter Six. First Part. Solution of the Congruence $x^m - 1 \equiv 0$ Page 199
. . . . . .
Chapter Eight. General Investigations of Congruences . . . . . — 212
Further Development of the Investigations of Pure Equations — 243
. . . . . .
Proof of Some Theorems Concerning the Periods of Classes of Binary — 266
Quadratic Forms . . . . . .
On the Connection between the Number of Classes into Which Binary — 269
Quadratic Forms Are Distributed and Their Determinant. I. II . . X
. . . . . .
Geometrical Aspect of Ternary Forms . . . . . — 305
On the Theory of Biquadratic Residues. I. . . VI . . . . . — 313
On the Theory of Complex Numbers. I. . . VI . . . . . — 387
Table of the Quadratic Character of Primes . . . . . — 399
Table for Converting Common Fractions into Decimal Fractions — 411
. . . . . .
Table of the Frequency of Primes . . . . . — 435
Table of the Number of Classes of Binary Quadratic Forms . . . . . — 449
Table for Circle Division . . . . . — 477
A New Quadrature of the Circle . . . . . -. . . . . — . 497
On Calculating Logarithms . . . . . — 501
First Ten Thousand Squares . . . . . — 504
Indices of Primes in the Higher Number Domain . . . . . , . . . . . — 506
Auxiliary Table for Solving the Equation $A = fxx + gyy$ . . . . . — 509
Eighth Section. A Fuller Consideration of Certain Investigations — 510
Concerning Circle Division.
Letters from Gauss to Dirichlet . . . . . — 514

*Remarks* . . . . . — 519

---

GÖTTINGEN,
PRINTED BY THE DIETRICH UNIVERSITY PRINTING HOUSE.
W. FR. KAESTNER.